BIRLA CENTRAL LIBRARY
PILANI (Rajasthan)
Class No. R 583.03
Book No. J13 I. V-1
Accession No. 64798

INDEX KEWENSIS

AN ENUMERATION OF THE GENERA AND SPECIES

OF

FLOWERING PLANTS

FROM THE TIME OF LINNAEUS TO THE YEAR 1885 INCLUSIVE
TOGETHER WITH THEIR AUTHORS' NAMES, THE WORKS IN
WHICH THEY WERE FIRST PUBLISHED, THEIR NATIVE COUNTRIES
AND THEIR SYNONYMS

COMPILED AT THE EXPENSE OF
THE LATE CHARLES ROBERT DARWIN
UNDER THE DIRECTION OF JOSEPH D. HOOKER

BY

B. DAYDON JACKSON

VOLUME I

OXFORD
AT THE CLARENDON PRESS
M DCCC XCV

INDEX KEWENSIS

PLANTARUM PHANEROGAMARUM

NOMINA ET SYNONYMA OMNIUM GENERUM ET SPECIERUM

A LINNAEO USQUE AD ANNUM MDCCCLXXXV

COMPLECTENS

NOMINE RECEPTO AUCTORE PATRIA

UNICUIQUE PLANTAE SUBJECTIS

SUMPTIBUS BEATI CAROLI ROBERTI DARWIN

DUCTU ET CONSILIO JOSEPHI D. HOOKER

CONFECIT B. DAYDON JACKSON

TOMUS I

OXONII

E PRELO CLARENDONIANO

MDCCCXCV

OXFORD UNIVERSITY PRESS
AMEN HOUSE, E.C. 4
LONDON EDINBURGH GLASGOW NEW YORK
TORONTO MELBOURNE CAPE TOWN BOMBAY
CALCUTTA MADRAS
GEOFFREY CUMBERLEGE
PUBLISHER TO THE UNIVERSITY

First Edition 1895
Reprinted photographically in Great Britain
at the University Press, Oxford, 1946, by
John Johnson, Printer to the University

INDEX KEWENSIS

PLANTARUM PHANEROGAMARUM

NOMINA ET SYNONYMA OMNIUM GENERUM ET SPECIERUM

A LINNAEO USQUE AD ANNUM M.DCCC.LXXXV

COMPLECTENS

AA—ABELMOSCHUS.

AA, Reichb. f. Xen. Orchid. i. 18 (1858) = **Altensteinia**, H. B. & K. (Orchid.).
argyrolepis, Reichb. f. l. c. = Alt. argyrolepis.
palaeora, Reichb. f. l. c. = Alt. palaeoces.

AAMA, Hask. Cat. Hort. Bog. Alt. 171 (1844), sphalm. = ADAMIA, Wall. = **Dichroa**, Lour. (Saxifr.).

ABACOSA, Alef. in Bonplandia, ix. (1861) 102 = **Vicia**, Linn. (Legumin.).
americana, Alef. l. c. 104 = V. americana.
canescens, Alef. l. c. 103 = V. canescens.
dumetorum, Alef. l. c. 104 = V. dumetorum.
Kingii, Alef. l. c. 103 = V. Kingii.
micrantha, Alef. l. c. 104 = V. micrantha.
onobrychioides, Alef. l. c. 103 = V. onobrychioides.
pallida, Alef. l. c. = V. pallida.
patagonica, Alef. l. c. = V. patagonica.

ABALON, Adans. Fam. ii. 511 (err. typ. 512) (1763) = **Helonias**, Linn. (Liliac.).
albiflorum, Rafin. New Fl. Am. i. 35.—Am. bor.

ABALUM, Adans. Fam. 47 (1763) = prasc.

ABAMA, Adans. Fam. ii. 47 (1763) = **Narthecium**, Moehr. (Liliac.).
anthericoides, Steud. Nom. ed. I. 550, in syn. (1821) = N. ossifragum.
glutinosa, Rafin. New Fl. Am. i. 36 = Tofieldia glutinosa.
ossifraga, DC. Fl. Fr. iii. 171 = N. ossifragum.
pubens, Rafin. New Fl. Am. i. 35 = Tofieldia pubens.
pusilla, Rafin. l. c. 36.—Am. bor.
scabra, Rafin. l. c. 35; Fl. Tellur. 28 = Tofieldia pubens.

ABANDION, Adans. Fam. ii. 54 (1763) = **Bulbocodium**, Linn. (Liliac.).

ABAPHUS, Rafin. Fl. Tellur. iv. 19 (1836) = seq.

ABAPUS, Adans. Fam. ii. 57 (1763) = **Gethyllis**, Linn. (Amaryll.).
spiralis, Rafin. Fl. Tellur. iv. 19 = G. spiralis.

ABASICARPON, Andrz. ex Reichb. in Moesal. Handb. ed. II. 1150 (1858) = **Arabis**, Linn. (Crucif.).
patulum, Andrz. ex Reichb. l. c. = Arab. auriculata.

ABASOLOA, La Llave, in Nov. Veg. Desc. fasc. i. 11 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 362.
Tabasola, *La Llave*, *l. c.*—Mexic.

ABASTA, Ruiz & Pav. Prod. 78. t. 14 (1794). *XANTHACEAE*, Benth. & Hook. f. i. 799.
Grahibra, Mand. & Wedd. ex Benth. & Hook. f. i. 6.
Myrotrichum, Turcz. in Bull. Soc. Nat. Mosc. (1869), i. 254.
Ralbighia, Gardn. in Hook. Lond. Journ. Bot. iv. (1845). 97.

ABATIA :—
americana, *Eichl. in Mart. Fl. Bras.* xiii. 1. 510.—Bras.
brasiliensis, *Baill. Adansonia*, x. (1872) 245.—Bras.
parviflora, *Ruiz & Pav. Syst.* i. 136; *Prod.* t. 14.—Peruv.; N. Granat.
rugosa, *Ruiz & Pav. ll. cc.*—Peruv.
tomentosa, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 509.—Bras.
verbascifolia, H. B. & K. Nov. Gen. et Sp. v. 358 = parviflora.

ABAURIA, Becc. Malesia, i. (1877) 169. *LEGUMINOSAE*.
excelsa, *Becc. l. c.*—Malaya.

ABAZICARPUS, Andrz. ex DC. Syst. ii. 213 (1821) = **Arabis**, Linn. (Crucif.).

ABBEVILLEA, Berg, in Linnaea, xxvii. 425 (1856) = Campomanesia, Ruiz & Pav. (Myrtac.).
cerasoides, Berg, in Mart. Fl. Bras. xiv. 1. 436.—Bras.
chrysophylla, Berg, l. c. 437 = Psidium chrysophyllum.
eugenioides, Berg, l. c. 436.—Bras.
Penthiana, Berg, l. c. 433.—Bras.
Gardneriana, Berg, l. c. 436.—Bras.
Guaviroba, Berg, l. c. 432 = C. Guaviroba
Klotzschiana, Berg, l. c. 434.—Bras.
Langsdorffiana, Berg, l. c. 603.—Bras.
Martiana, Berg, l. c. 435.—Bras.
maschalantha, Berg, l. c. 432.—Bras.
neriifolia, Berg, l. c. 431.—Bras.
phaea, Berg, l. c. 435.—Bras.
plana, Berg, in Linnaea, xxvii. (1854) 427 (sphalm. = phaea).
punctulata, Berg, in Mart. Fl. Bras. xiv. 1. 432.—Bras.
recurvata, Berg, l. c. 438.—Bras.
Regeliana, Berg, l. c. 603.—Bras.
rufa, Berg, l. c. 437 = Psidium rufum.
rugosa, Berg, l. c.—Bras.
Schlechtendahliana, Berg, l. c. 435.—Bras.
Selloviana, Berg, l. c. 431.—Bras.
Widgreniana, Berg, in Linnaea, xxxi. (1861–62) 360. —Bras.

ABBOTIA, Rafin. New Fl. Am. i. 36 (1836) = Triglochin, Linn. (Naiad.).
filiformis, Rafin. l. c. = T. triandrum.
palustris, Rafin. l. c. 37 = T. palustre.
pusilla, Rafin. l. c. = T. palustre.

ABBOTTIA F. Muell. Fragm. ix. 121 (1875) *RUBIACEAE*.
singularis, *F. Muell. l. c.*—Austral.

ABELIA, R. Br. in Abel, Chin. App. 376 (1818) ii. 4. *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 4.
adenotricha, *Hance*, *in Journ. Bot.* ix. (1871) 193.—China.
biflora, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) VII. 153.—China.
chinensis, *R. Br. in Abel, Chin. App.* 376.—China.

ABELIA :—
coriacea, *Hemsl. Diagn. Pl. Nov.* 53.—Mexic.
corymbosa, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 606.—As. temp.
Davidii, Hance, in Seem. Journ. Bot. vi. (1868) 329 = biflora.
floribunda, *Decne. in Fl. des Serres*, Sér. I. ii. (Janv. 1846).—Mexic.
Hanceana, Mart. ex Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 216 = chinensis.
hirsuta, Walp. Rep. vi. 3 = speciosa.
rupestris, Lindl. in Journ. Hort. Soc. i. (1846) 63 = chinensis.
serrata, Sieb. & Zucc. Fl. Japon. i. 76. t. 84 = uniflora.
spathulata, *Sieb. & Zucc. l. c.* 77. t. 84.—Japon.
speciosa, *Decne. in Fl. des Serres*, ii. (1846) sub t. 5. —Mexic.
splendens, Hort. ex C. Koch, Dendrol. ii. 1. 20 = Lonicera caprifolioides.
trifiora, *R. Br. in Wall. Pl. As. Rar.* i. 14. t. 15.—Reg. Himal.
uniflora, *R. Br. in Wall. l. c.*—China.

ABELICEA, Reichb. Consp. 84 (1828); Baill. Hist. Pl. vi. 185 (1875) = **Zelkova**, Spach (Urtic.).
aquatica, Rafin. New Fl. Am. iii. 41 = Planera aquatica.
crenata, Baill. Hist. Fl. vi. 181 = Z. crenata.
cretica, Rafin. New Fl. Am. iii. 41 = Z. cretica.
Davidii, Buek, Ind. Gen. DC. Prod. iv. 1 = Hemipteles Davidii.
sibirica, Rafin. New Fl. Am. iii. 41 = Z. crenata.

ABELMOSCHUS, Medic. Malv. 45 (1787) = **Hibiscus**, Linn. (Malvac.).
achanicoides, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 196 = H. achanicoides.
aculeatus, Walp. Rep. i. 308 = H. surattensis, cannabinus.
acuminatus, C. Muell. in Walp. Ann. vi. 407.—Java.
albo-rubens, Benth. Fl. Austral. i. 209 = seq.
albo-ruber, F. Muell. Fragm. i. 67 = H. ficulneus.
angulosus, Wight & Arn. Prod. 53 = H. angulosus.
angustifolius, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 152 = H. angustifolius.
aquaticus, Walp. Rep. i. 311 = H. roseus.
ascendens, Walp. l. c. 310 = H. physaloides.
Baumia, Webb. Fragm. Fl. Aethiop. 46 = H. esculentus.
betulifolia, Wall. Cat. p. 87 et n. 1916 = H. Abelmoschus.
cancellatus, Wall. l. c. p. 87 et n. 1920 = H. cancellatus.
chinensis, Wall. l. c. p. 87 et n. 1916 = H. Abelmoschus.
ciliaris, Walp. Rep. ii. 308 = H. ciliaris.
cisplatinus, Walp. l. c. 309 = H. cisplatinus.
congener, Walp. l. c. 308 = H. cannabinus.
crinitus, Wall. Pl. As. Rar. i. 39 = H. cancellatus.
crunatus, Walp. Rep. i. 310 = H. crunatus.
cryptocarpus, Walp. l. c. = H. Abelmoschus.
cubensis, Walp. l. c. 309 = H. Abelmoschus.
cucurbitaceus, Walp. l. c. 310 = H. Abelmoschus.
decandrus, Medic. Malv. 46 = Pavonia praemorsa.
divaricatus, Walp. Rep. i. 309 = H. divaricatus.
esculentus, Moench, Meth. 617 = H. esculentus.

ABELMOSCHUS :—
floccineoides, Walp. Rep. i. 809 = H. floccineoides.
floccineus, Wight. & Arn. Prod. i. 53 = H. floccineus.
fusiformis, Wall. Cat. p. 87 et n. 1922 = H. cancellatus.
Genesii, Walp. Rep. i. 810 = H. illitifolius.
guineensis, Walp. l. c. = H. tiliaceus.
Hamabanus, Presl. Rel. Haenk. ii. 134 —Ins. Luzon.
laxiflorus, Walp. Rep. i. 810 = H. laxiflorus.
longifolius, Kostel. Allg. Med.-Pharm. Flora, v. 1859 = H. Abelmoschus.
moschatus, Bartl. in Linnæa, xiii. (1839) Litt. 81 = H. Bartlingianus.
magnifolius, Wall. Cat. p. 87 et n. 1919.—Ind. or. *Manihot*, Medic. Malv. 46 = H. Manihot.
mariannus, Presl, Rel. Rel. Haenk. ii. 134.—Ins. Marian.
moschatus, Medic. Malv. 46 = H. Abelmoschus.
multiformis, Wall. Cat. p. 87 et n. 1917.—Ind. or.
officinalis, Endl. ex Heynh. Nom. ii. 1 = H. esculentus.
palustris, Walp. Rep. i. 811 = H. Moscheutos.
pandurseformis, Hassk. in Tijdschr. Nederl. Ind. i. (1845) 184 = H. pandurseformis.
pentaphyllus, Voigt, Hort. Suburb. Calc. 119 = H. pentaphyllus.
persicifolius, Walp. Rep. i. 808 = H. persicifolius.
Pseudo-abelmoschus, Blume, Bijdr. 70 = H. Abelmoschus.
pungens, Wall. Cat. p. 87 et n. 1924 = H. pungens.
quinquelobus, Walp. Rep. i. 810 = H. quinquelobus.
racemosus, Wall. Cat. p. 87 et n. 1929 = H. cancellatus.
rhodopetalus, F. Muell. Fragm. ii. 113 = H. rhodopetalus.
roseus, Walp. Rep. i. 811 = H. Moscheutos.
rostellatus, Walp. l. c. 808 = H. rostellatus.
rugosus, Wight & Arn. Prod. i. 43 = H. rugosus.
splendens, Walp. Rep. i. 809 = H. splendens.
strictus, Voigt, Hort. Suburb. Calc. 119 = H. floccineus.
sublobatus, Presl, Rel. Haenk. ii. 134.—Ins. Pacif.
tetraphyllus, Wall. Cat. p. 87 et n. 1926 = H. tetraphyllus.
thrinodes, Voigt, Hort. Suburb. Calc. 120 = H. sulphureus, *Voigt.*
tulipæoflorus, Walp. Rep. v. 92 = H. tulipæoflorus.
venustus, Walp. l. c. i. 809 = H. mutabilis.
verrucosus, Walp. l. c. 808 = H. cannabinus.
Vrieslanus, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 184; xii. (1845) 88.—Malaya.
warreensis, Dalz. in Hook. Kew Journ. iii. (1851) 123 = H. tetraphyllus.
Zollingeri, C. Muell. in Walp. Ann. vi. 407.—Java.

ABENA, Neck. Elem. i. 296 (1790) = **Stachytarpheta**, Vahl (Verbenac.).

ABEREMOA, Aubl. Fl. Gui. i. 610, t. 245 (1775) = **Duguetia**, St. Hil. (Anonac.).
furfuracea, Baill. Hist. Pl. i. 204.—Bras.
guianensis, Aubl. Fl. Gui. i. 610.—Guiana.
lanceolata, Warm. in Kjoeb. Vidensk. Meddel. (1855) 146; (1873) 146.—Bras.
longifolia, Baill. Hist. Pl. i. 205 = D. longifolia.

ABERIA, Hochst. in Flora, xxvii. (1844) Bes. Beil. 2. *BIXINEÆ*, Benth. & Hook. f. i. 128.
ATHLESTE, Sond. in Linnæa, xxiii. (1850) 9, 10.
abyssinica, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 236.—Afr. trop.
caffra, *Harv. & Sond. Fl. Cap.* ii. *Add.* 584.—Afr. austr.
edulis, T. Anders. in Journ. Linn. Soc. vii. (1862) 67 = caffra.
Gardneri, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 235.—Zeylan.
longispina, *Harv. in Have. & Sond. Fl. Cap.* ii. 585.—Afr. austr.
macrocalyx, *Oliver, Fl. Trop. Afr.* i. 122.—Afr. trop.
mollis, *Oliver, l. c.*—Afr. trop.
tristis, *Sond. in Linnæa*, xxiii. (1850) 9.—Afr. austr.
verrucosa, *Hochst. in Flora*, xxvii. (1844) *Bes. Beil.* 2.—Abyss.
Zeyheri, *Sond. in Linnæa*, xxiii. (1850) 10.—Afr. austr.

ABESINA, Neck. Elem. i. 83 (1790) = **Verbesina**, Linn. (Compos.).

ABIES, D. Don, in Loud. Arb. Brit. iv. 2329 (1838) = **Picea**, Link (Conif.).

ABIES, [Tourn.] Linn. Fl. Lapp. 277 (1737); Link, in Linnæa, xv. (1841) 525. *CONIFERÆ*, Benth. & Hook. f. iii. 441.
PICEA, D. Don, in Loud. Arb. Brit. iv. 2329 (1838).
ajanensis, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 213 = Picea ajanensis.

ABIES :—
alba, Michx. Fl. Bor. Am. ii. 207 = Picea alba.
alba, Mall. Gard. Dict. ed. VIII. n. 1 (1768) = pectinata (1803).
Albertiana, A. Murr. in Proc. Hort. Soc. iii. (1863) 149 = Tsuga Mertensiana.
Alcockiana, Lindl. ex C. Koch, Dendrol. ii. 11. 245 = Picea Alcockiana.
Alcquesana, Gord. Pinet. Supp. 8 (“desc. falsa”) = pinsc.
amabilis, *Forb. Pin. Wob.* 126. t. 44.—Am. bor. occ.
amabilis, Parl. in DC. Prod. xvi. 11. 426, in parte = sublpina.
americana, Mill. Gard. Dict. ed. VIII. n. 6 = Tsuga canadensis.
Apollinis, Link, in Linnæa, xv. (1841) 528 = cephalonica.
Arangi, Siebold, in Verh. Batav. Gen. xii. (1830) 12 = Tsuga Sieboldii.
araucana, Poir. Encyc. Suppl vi. 524 = Araucaria excelsa.
arctica, A. Cunn. ex Gord. Pinet. 11 = Picea rubra.
arctica, A. Murr. in Journ. Bot. v. (1867) 258 = Picea alba.
argentea, De Chambr. Traité Prat. Arb. Résin. 17 = pectinata.
aromatica, Rafin. Atl. Journ. 119 = balsamea ?
atlantica, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 214 = Cedrus atlantica.
balsamea, Sm. ex Carr. Conif. ed. II. 229 = Cunninghamia sinensis.
bicolor, Maximo. in Bull. Acad. Pétersb. x. (1866) 488 = Picea Alcockiana.
bifida, Sieb. & Zucc Fl. Jap. ii 18 = firma.
bifolia, A. Murr. Syn. Var. Conif. 29 = lasiocarpa.
brachyphylla, Maximo. *in Bull. Acad. Pétersb.* x. (1866) 488.—Japon.
brevifolia, *Hook. & Arn. Bot. Beech. Voy.* 394.—Calif *Bridgesii*, Kellogg, in Proc. Calif. Acad. ii. (1863) 8 = Tsuga Mertensiana.
Brunoniana, Lindl. in Penny Cyclop. i. 80 = T. Brunoniana.
californica, Hort. ex Steud. Nom. ed. II. i. 1 = Pseudotsuga Douglasii.
canadensis, A. Murr. in Trans. Bot. Soc. Edinb. vi. (1860) 370 = magnifica.
canadensis, Michx. Fl. Bor. Am. ii. 206 = Tsuga canadensis.
canadensis, Mill. Gard. Dict. ed. VIII. n. 4 = Picea alba.
candicans, Fisch. ex Endl. Syn. Conif. 96 = pectinata.
cedroides, Griff. ex Carr. Conif. ed. I. 188 = Tsuga Brunoniana.
Cedrus, Poir. Encyc. vi. 510 = Cedrus Libani.
cephalonica, *Link, in Linnæa*, xv. (1841) 529.—Græcia.
chilotensis, Hort. ex Carr. Conif. ed. I. 222 = A. Pindrow.
chlorocarpa, Hort. ex Parl. in DC. Prod. xvi. 11. 425 = Webbiana.
chlorocarpa, Purk. ex Nym, Consp. 674 = Picea excelsa.
cilicica, *Carr. Conif.* ed. I. 229.—As. Min.
cinerea, Borkh. Forsth. i. 898 = Picea excelsa.
Clanbrassiana, Hort. ex Lond. Encyc. Pl. 804 = P. excelsa.
corrulea, Lodd. ex Forbes, Pin. Wob. 99 = P. alba.
corrulescens, Hort. ex C. Koch, Dendrol. ii. 11. 242 = P. rubra.
Columbaria, Desf. Tabl. ed. II. 247 = Araucaria imbricata.
concentata, Gord. Pinet. ed. II. 5 = Picea Engelmanni.
concolor, *Lindl. & Gord. in Journ. Hort. Soc.* v. (1850) 210.—Am. bor. occ.
coptrolepis, Poir. Encyc. vi. 524 = Libocedrus tetragona.
cursifolia, Salisb. in Trans. Linn. Soc. viii. (1807) 315 = Picea alba.
Deamara, Poir. Encyc. Suppl. v. 35 = Agathis loranthifolia.
Davidiana, *Franch. in Nouv. Mém. Mus. Par.* vi. (1884) 96.—As. or.
decidua, Wall.—Cf. C. Koch, Dendrol. ii. 11. 258 = Tsuga Brunoniana.
densa, Griff. Notul. iv. 19.—Cf. Webbiana.
denticulata, Poir. Encyc. vi. 520 = Picea nigra.
Deodara, Lindl. in Penny Cyclop. i. 84 = Cedrus Deodara.

ABIES :—
diversifolia, Maxim. in Bull. Acad. Pétersb. xii. (1868) 229 = Tsuga diversifolia.
Douglasii, Lindl. in Penny Cyclop. i. 82 = Pseudotsuga Douglasii.
Drummondi, Hort. ex Gord. Pinet. ed. I. 16 = pinsc.
dumosa, Loud. Arb. Brit. iv. 2326 = Tsuga Brunoniana.
Eichleri, Lasche, in Berl. Gartenz. i. (1882) 62 = Veitchii.
elegans, Sm. ex Knight, Syn. Conif. 36 = Picea excelsa.
Engelmanni, Parry, in Trans. Acad. St. Louis, ii. (1868) 212 = Picea Engelmanni.
eremita, Hort. Gall. ex C. Koch, Dendrol. ii. 11. 237 = Picea excelsa.
erythrocarpa, Purk. ex Nym. Consp. 674 = P. excelsa.
excelsa, Link, in Abh. Acad. Berl. (1827) 182 = pectinata.
excelsa, Poir. Encyc. vi. 518 = P. excelsa.
extrema, T. Fries, in Bot. Notiser (1867), 56 = P. excelsa.
falcata, *Rafin. Atl. Journ.* 120.—Am. bor.
fendomensis, Hort. ex Gord. Pinet. Suppl. 4 = P. excelsa.
Fischeriana, R. Neum. ex Parl. in DC. Prod. xvi. 11. 431 = homolepis.
firma, *Sieb. & Zucc. Fl. Jap.* ii. 15. t. 107.—Japon.
Fischeri, *Lindl. ex Loddi. Pl. Ross.* ii. 172.—Rossia.
Fortunei, A. Murr. Pin. Japan. 49 = Keteleeria Fortunei.
Fraseri, *Lindl. in Penny Cyclop.* i. 80.—Am. bor.
gigantea, Sm. ex Carr. Conif. ed. II. 227 = Picea excelsa.
glauca, Roezl, ex Gord. Pinet. 149 = religiosa.
glaucescens, Koezl, Cat. Gr. Conif. Mex. (1886–89) 5 = religiosa.
Glehni, F. Schmidt, Reisen Amurl. 176 = Picea Glehnii.
Gmelini, Rupr. in Beitr. Pfl. Russ. Reich. ii. 56 = Larix dahurica.
Gordoniana, Carr. Conif. ed. II. 298 = grandis.
grandis, *Lindl. in Penny Cyclop.* i. 80.—Am. bor. occ.
Gregoryana, Low, ex Gord. Pinet. Suppl. 4 = P. excelsa.
Griffithiana, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 214 = Larix Griffithii.
Hanburyana, Hort. ex Gord. Pinet. ed. II. 32, 421 = Tsuga Sieboldii ?
Harryana, McNab, in Proc. Roy. Irish Acad. (1876) 689 = homolepis.
heterophylla, C. Koch. in Linnæa, xxii. (1849) 295 = sibirica.
heterophylla, Rafin. Atl. Journ. 119 = Tsuga Mertensiana.
hirtella, Lindl. in Penny Cyclop. i. 81 = religiosa.
hispanica, De Chamb. Trait. Prat. Arb. Résin. 339 = A. Pinsapo.
holophylla, *Maxim. in Bull. Acad. Pétersb.* x. (1866) 487.—As. temp.
homolepis, *Sieb. & Zucc. Fl. Jap.* ii. 17.—Japon.
Hookeriana, A. Murr. in Edinb. N. Phil. Journ. N. S. i. (Jan.-Apr. 1855). 289 = Tsuga Hookeriana.
Hudsonia, Bosc, ex Carr. Conif. ed. I. 200 = Fraseri.
humilis, La Pyl. in Mém. Soc. Linn. Par. iv. (1826) 457 = Fraseri.
inverta, R. Sm. ex Gord. Pinet. Suppl. 4 = P. excelsa.
jesoënsis, Lindl. & Paxt. Flow. Gard. i. (1850–51) 42. f. 26 = Keteleeria Fortunei.
jesoensis, Sieb. & Zucc. Fl. Jap. ii. 19. t. 110 = Picea Alcockiana.
Kæmpferi, Lindl. in Penny Cyclop. i. 84 = Pseudolarix Kæmpferi.
kamtschatica, Rupr. in Beitr. Pfl. Russ. Reich. ii. 57 = Larix dahurica.
Khutrow, Lond. Encyc. Trees, 1089 = Picea Morinda.
Kotschyana, *Fenzl, ex Tchihat. Asie Min. Bot.* ii. 495.—As. Min.
lanceolata, Poir. Encyc. vi. 523 = Cunninghamia sinensis.
Larix, Poir. Encyc. Suppl. vi. 511 = Larix europæa.
lasiocarpa, *Hook. Fl. Bor. Am.* ii. 163.—Am. bor. occ.
lasiocarpa, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 210 = grandis, Lowiana.
laxa, C. Koch, Dendrol. ii. 11. 243 = Picea alba.
Lebdourii, Rupr. in Beitr. Pfl. Russ. Reich. ii. 56 = Larix dahurica.
leiolepis, Stev. ex Gord. Pinet. 152 = Nordmanniana.
Lemoniana, Booth, ex Gord. i. c. ed. I. 6 = Picea excelsa.
leptolepis, Sieb. & Zucc. Fl. Japon. 13. t. 108 = Larix japonica.
Lindleyana, Roezl, ex Carr. Conif. ed. II. 275 = religiosa.
Lowiana, *A. Murr. Syn. Var. Conif.* 27.—Am. bor. occ.

ABIES :—

Luscombeana, Lond. Arb. Brit. iv. 2225 = cephalonica.
magnifica, *A. Murr. in Proc. Roy. Hort. Soc.* iii. (1868) 218.—Calif.
Mariana, Mill. Gard. Dict. ed. VIII. n. 5 = Picea nigra.
Mariesii, *Mast. in Gard. Chron.* xli. (1879) 789.—As. or.
Maximowiczii, R. Nexm. Cat. (1868) ; ex Parl. in DC. Prod. xvi. ii. 481 = Picea Maximowiczii.
medioxima, Lawson, Pin. Brit. 159 = P. excelsa.
Menziesii, Lindl. in Penny Cyclop. i. 32 = Picea Menziesii.
Morhiana, Fisch. ex Parl. in DC. Prod. xvi. ii. 418 = Picea Menziesii.
Mertensiana, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 211 = Tsuga Mertensiana.
metensis, Hort. Par. ex Gord. Pinet. ed. II. 211 = pectinata.
microcarpa, Lindl. in Penny Cyclop. i. 33 = Larix microcarpa.
microphylla, *Rafin. Atl. Journ.* (1830) 110.—Am. bor.
microsperma, Lindl. in Gard. Chron. (1861) 22 = Picea ajanensis.
minuta, Knight, ex Gord. Pinet. ed. II. 8 = P. excelsa.
minor, Duham. ex Gord. i. c. ed. I. 148 = balsamea.
minor, Gilib. Exerc. Phytol. ii. 412 = pectinata.
minuta, Poir. Encyc. vi. 523 = P. rubra.
Momi, Siebold, in Verh. Batav. Gen. xii. (1830) 12 = firma.
montana, Nym. Syll. Suppl. 602 = P. excelsa.
Morinda, Lond. Encyc. Pl. 1294 = P. Morinda.
mucronata, Rafin. Atl. Journ. 120 = Pseudotsuga Douglasii.
mucronata, Rauch, ex Carr. Conif. ed. II. 329 = Picea excelsa.
nephrolepis, Maxim. in Bull. Acad. Pétersb. x. (1866) 486 = Veitchii.
nigra, Desf. Hist. Arbr. et Arb. ii. 580 = Picea nigra.
nigra, Durol, Harb. Baumz. ed. Pott, ii. 189 = P. nigra.
nigra, Engelm. in Am. Journ. Sc. xxxiii. (1888) 380 = Picea Engelmanni.
nobilis, *Lindl. in Penny Cyclop.* i. 30.—Am. bor. occ. Nordmanniana, *Spach, Hist. Vég. Phan.* xi. 418.—Reg. Caucas.
Novae-Angliae, C. Koch, Dendrol. ii. ti. 240 = Picea nigra.
numidica, *De Lannoy, ex Carr. in Rev. Hortic.* (1866) 106.—Afr. bor.
obliqua, Bong. ex Gord. Pinet. Supp. 10 = Pseudotsuga Douglasii.
obliquata, Rafin. ex Gord. i. c. = pracc.
obovata, Lond. Arb. Brit. iv. 2329 = Picea obovata.
Omorika, Nym. Consp. 673 = Picea Omorika.
orientalis, Poir. Encyc. vi. 518 = Picea orientalis.
panachaica, Heldr. ex Parl. in DC. Prod. xvi. ii. 422 = cephalonica.
Parsoniana, Hort. ex Mast. in Gard. Chron. (1880) i. 648 = Lowiana.
parvula, Knight, Syn. Conif. 36 = Picea excelsa.
Pattoni, Jeffrey, ex Gord. Pinet. Suppl. 6 = seq.
Pattoniana, Jeffrey, ex A. Murr. in Bot. Exped. Oregon. Circ. = Tsuga Hookeriana.
pectinata, *DC. Fl. Fr.* iii. 276.—Europ.
pectinata, Gilib. Exerc. Phytol. ii. 411 = P. excelsa.
pectinata, Poir. Encyc. vi. 528 = Picea rubra.
peloponnesiaca, Haage, ex Parl. in DC. Prod. xvi. ii. 422 = cephalonica.
pendula, Griff. ex Gord. Pinet. Suppl. 7 = Picea Morinda.
pendula, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 213 = Larix microcarpa.
Picea, Lindl. in Penny Cyclop. i. 29 = pectinata.
Picea, Mill. Gard. Dict. ed. VIII. n. 2 = Picea excelsa.
Pichta, Forbes, Pin. Wob. 113. t. 39 = sibirica.
Pindrow, *Spach, Hist. Vég. Phan.* xi. 428.—Reg. Himal. Pinsapo, *Boiss. in Bibl. Univ. Genève* (1838) ; *ex cj. Voy. Bot. Espagne*, ii. 584.—Hispan.
polita, Sieb. & Zucc. Fl. Jap. ii. 20. t. 111 = Picea polita.
Reginae-Amaliae, Heldr. in Regel, Gartenfl. ix. (1860) 312 = cephalonica.
religiosa, *Lindl. in Penny Cyclop.* i. 31.—Mexic.
Rinzi, Hort. Par. ex Gord. Pinet. Suppl. 50 = silicica.
Romanesia, Lond. ex Stend. Nom. ed. II. i. 1 = Pinus Laricio.
rubra, Petr. Encyc. vi. 520 = P. rubra.
sachalinensis, *Mast. in Gard. Chron.* (1879) ii. 588.—As. or.
sacra, *Franch. in Nouv. Arch. Mus. Par.* vii. (1882) 200.—As. or.
Schrenkiana, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 212 = Picea obovata.

ABIES :—

selinusia, Carr. in Fl. des Serres (1859), 69 = silicica.
sibirica, *Ledeb., Fl. Alt.* iv. 203.—Sibir.
sitchensis, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 212 = Picea Menziesii.
Smithiana, Lindl. in Penny Cyclop. i. 31 = P. Morinda.
spectabilis, Spach, Hist. Vég. Phan. xi. 422 = Webbiana.
spinulosa, Griff. Itin. 259, 265, 275 = Picea Morinda.
Standishiana, C. Koch, Dendrol. ii. ii. 257 = Tsuga Douglasii.
subalpina, Engelm. ex L. Ward, in Am. Natural. x. (1876) 555 = lasiocarpa.
subarctica, Nym. Consp. 674 = Picea excelsa.
sumatrana, Desf. Hort. Par. ed. III. 356 = Agathis lorantifolia.
taxifolia, Desf. i. c. = pectinata.
taxifolia, Nym. ex Gord. Pinet. 16 = Pseudotsuga Douglasii.
taxifolia, Jeffr. ex Gord. i. c. 18 = Tsuga Mertensiana.
taxifolia, Presl, Epim. Bot. 287 = Pseudotsuga Douglasii.
taxifolia, Rafin. New Fl. Am. i. 38.—Am. bor.
Tchugatskoi, Lawson, ex Gord. Pinet. Suppl. 50 = silicica.
Thunbergii, Lindl. in Penny Cyclop. i. 34 = Picea polita.
Tlapalcatsuda, Roezl, Cat. Gr. Conif. Mex. (1858–59) 5 = religiosa.
Torano, Siebold, in Verh. Bat. Gen. xii. (1830) 12 = Picea polita.
trigona, Rafin. Autik. Journ. (1820) 119 = Sequoia sempervirens!
Tschonoskyana, Regel, in Ind. Sem. Hort. Petrop. (1865) 32 = homolepis.
Tsuga, Sieb. & Zucc. Fl. Jap. ii. 14. t. 106 = Tsuga Sieboldii.
Veitchii, *Lindl. in Gard. Chron.* (1861) 23.—Japon.
venusta, C. Koch, Dendrol. ii. ii. 210 = bracteata.
viminalis, Wahlenb. Fl. Suec. 630 = Picea excelsa.
virescens, Hinterh. ex Nym. Consp. 674 = P. alba.
vulgaris, Poir. Encyc. vi. 514 = pectinata.
Webbiana, *Lindl. in Penny Cyclop.* i. 30.—Reg. Himal.
Williamsonii, Bridges, ex Gord. Pinet. Suppl. 12 = Tsuga Mertensiana.
Williamsonii, Newberry, in Williamson, in Pacif. Rail. Rep. vi. 63. t. 7 = Tsuga Hookeriana.

ABIGA, St. Lag. in Ann. Soc. Bot. Lyon, vii. 85 (1880) = **Ajuga**, Linn. (Labiat.).
cistifolia, St. Lag. i. c. 85, 119 = Ajuga Iva.

ABILDGAARDIA, Vahl, Enum. ii. 296 (1806)* = **Pimbristylis**, Vahl (Cyperac.).
aphylla, Vahl, Enum. ii. 348 = F. bahlenis.
Baorthryon, St. Hil. Voy. ii. 389 = F. bahlenis.
barbata, Beauv. Fl. Oware, ii. 47. t. 86. f. 1.—Afr. trop.
brevifolia, Stend. Syn. Pl. Cyp. 72 = F. setacea.
cinnamomiorum, Thw. Enum. Pl. Zeyl. 347 = F. cyperoides.
compressa, J. & C. Presl, Rel. Haenk. i. 179 = F. monostachya.
cyperoides, Nees & Meyen, in Wight, Contrib. 95 = F. cyperoides.
Engrostis, Nees & Meyen, l. c. = F. Engrostis.
fimbristyloides, F. Muell. Fragm. viii. 273 = F. disticha.
fulvescens, Thw. Enum. Pl Zeyl. 347 = F. fulvescens.
fusca, F. Muell. Fragm. viii. 273 = F. cyperoides.
fusca, Nees, in Wight, Contrib. 95.—Reg. Himal.
geminata, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 71 = F. bahlenis.
indica, Nees, in Linnaea, ix. (1835) 289 = F. monostachya.
javana, Nees, l. c. = F. monostachya.
javanica, Stend. Syn. Pl. Cyp. 73 = F. junces.
laevigata, Willd. ex Stend. Nom. ed. II. i. 1 = F. monostachya.
lanceolata, Schnm. Gain. Pl. 88 = F. tristachya.
monostachya, Vahl, Enum. ii. 296 = F. monostachya.
nervosa, J. & C. Presl, Rel. Haenk. i. 180 = F. schoenoides.
pauciflora, Kunth, Enum. Pl. ii. 249.—Ins. Molucc.
pilosa, Nees, in Linnaea, ix. (1835) 289.—Afr. trop.
polyrephala, Brongn. in Duperr. Voy. Bot. 176. t. 83 = Androtrichum polyrephalum.
polygama, Spreng. Nov. Prov. 45.—Bras.
polystachya, Spreng. Fl. Hal. Mant. 29 = Pharus angustifolius.
pubescens, J. & C. Presl, Reliq. Haenk. 180 = F. exilis.
Rottboelliana, Nees, ex Hook. & Arn. Bot. Beech. Voy. 373 = F. monostachya.
schoenoides, R. Br. Prod. 229 = F. squarrulosa.

ABILDGAARDIA :—
scirpoides, Nees, in Linnaea, ix (1835) 289 = F. bahlensis.
setacea, Griseb. Cat. Pl. Cub. 288.—Cuba.
ternata, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 71 = F. bahlensis.
tristachya, Vahl, Enum. ii. 297 = F. tristachya.
vaginata, R. Br. Prod. i. 229 = F. Brownii.

ABILDGARDIA, Reichb. Consp. 56 (1828) = pracc.

ABILGAARDIA, Poir. in Dict. Sc. Nat. i. Suppl 3 (1816) = pracc.

ABLANIA, Aubl. Pl. Gui. i. 585. t. 234 (1775) = **Sloanea**, Linn. (Tiliac.).
digitata, Spreng. Syst. iv. Cur Post. 210.—Bras.
guianensis, Aubl. Pl. Gui. i. 585. t. 234 = S. guianensis.
laurifolia, Pers. Syn. ii. 81 = S. guianensis.

ABOREA, Naud. in Rev. Hortic. (1862) 111. *CUCURBITACEAE*. Benth. & Hook. f. i. 824.
tenuifolia, *Naud. l. c.*—Bras.
viridiflora, *Naud. l. c.*—Am. austr.

ABOLA, Adans. Fam. ii. 31 (1763) = **Cinna**, Linn. (Gramin.).

ABOLA, Lindl. Fol. Orch. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 506.
radiata, *Lindl. l. c.*—N. Granat.

ABOLARIA, Neck. Elem. i. 105 (1790) = **Globularia**, Linn. (Selagin.).

ABOLBODA, Humb. & Bonpl. Pl. Aequin. ii. 109. t. 114 (1809). *XYRIDEAE*, Benth. & Hook. i. ii. 842.
CHLOERUM, Willd. ex Link, in Spreng. Jahrb. i. iii. 74 (1838).
POARCHON, Mart. ex Seub. in Mart. Fl. Bras. iii. i. 228 (1855).
Aubletii, *Kunth, Enum. Pl.* iv. 28.—Guian.
brasiliensis, *Kunth, l. c.* 26.—Bras.
grandis, *Griseb. in Linnaea*, xxi. (1848) 281.—Bras.
imberbis, *H. B. & K. Nov. Gen. et Sp.* i. 256.—Guian.
inermis, Link, ex Steud. Nom. ed. II. i. 2 = imberbis.
Poarchon, *Seub. in Mart. Fl. Bras.* iii. i. 228.—Bras.
Poeppigii, *Kunth, Enum. Pl.* iv. 27.—Bras.
pulchella, *Humb. & Bonpl. Pl. Aequin.* ii. 110. t. 114. —Venezuela.

ABROCHIS, Neck. Elem. iii. 130 (1790) = **Orchis**, Tourn.

ABROMA, Jacq. Hort. Vind. iii. t. 1 (1776). *STERCULIACEAE*, Benth. & Hook. f. i. 224.
ABROMA, Linn. f. Suppl. 241 (1781).
alata, *Blanco, Fl. Filip.* ed. I. 606.—Ins. Philipp.
angulata, Lam. Encyc. i. 126 = augusta.
angulosa, Poir. Encyc. Suppl. i. 32 = augusta.
augusta, *Linn. f. Suppl.* 241 (Ambroma).—As. trop.
communis, Blanco, Fl. Filip. ed. I. 606 = augusta.
denticulata, *Miq. Pl. Yungh.* 288.—Java.
ricngata, Lam. Encyc. i. 127 = augusta.
fastuosa, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 409.—As. trop. ; Austral.
fastuosa, Gaertn. Fruct. i. 307. t. 64 = augusta.
fastuosa, Jacq. Hort. Vind. iii. 8. t. 1 = augusta.
javanica, *Miq. Fl. Ind. Bat.* i. ii. 188.—Java.
Mariae, *Mart. in Denkschr. Bayer. Ges.* iii. (1837–43) 207.—Java.
mollis, *DC. Prod.* i. 485.—Ins. Molucc.
nitida, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 73.—Chili.
obliqua, *Presl, Rel. Haenk.* ii. 148.—Ins. Philipp.
tomentosa, Stend. Nom. ed. II. i. 2, sphalm.—Cf. Ambora tomentosa.
Wheleri, Retz. Obs. v. 27 = augusta.

ABRONIA, Juss. Gen. 448 (1789). *NYCTAGINEAE*, Benth. & Hook. f. iii. 7.
CYCLOPTERA, Nutt. ex A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 319.
TRICHATUS, L'Hérit. ex Willd. Sp. Pl. i. 807 (1799).
arenaria, *Menzies, ex Hook. Exot. Fl.* t. 193.—Calif.
californica, Raeusch.-Nom. ed. III. 191 = umbellata.
Crux-Maltae, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 71. f. 16.—Calif.
Cyclopters, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 319.—Texas.
fragrans, *Nutt. ex Hook. Kew Journ.* v. (1853) 261.—Calif.
glauca, *Menzies, ex Hook. Exot. Fl.* t. 194.—Calif.
gracilis, *Benth. Bot. Voy. Sulph.* 44.—Calif.
latifolia, *Eschsch. in Mém. Acad. Piterob.* v. (1826) 281.—Calif.

ABRONIA :—
maritima, *Nutt. ex S. Wats. Bot. Calif.* ii. 4.—Calif.
mellifera, *Dougl. in Hook. Bot. Mag.* t. 3879.—Calif.
Menziesii, *Dougl. ex Hook. Fl. Bor. Am.* ii. 135.—Calif.
micrantha, Cholsy, in DC. Prod. xiii. ii. 436 = A. Cyclopters.
nana, *S. Wats. in Proc. Am. Acad.* xiv. (1870) 294. —Am. bor. occ.
parviflora, *H. B. & K. Nov. Gen. et Sp.* ii. 216. t. 128. —N. Granat.
rosea, Hartweg, ex Lond. Hort. Brit. Suppl. iii. 479 = umbellata !
rotundifolia, Gaertn. f. Fruct. iii. t. 214 = umbellata.
speciosa, *Buchl. in Proc. Acad. Sc. Philad.* (1868) 7.—Am. bor. occ.
turbinata, *Torr. ex S. Wats. in Bot. King's Exped.* 286. t. 31.—Am. bor. occ.
umbellata, *Lam. Illustr.* i. 469. t. 105.—Calif.
vespertina, Dougl. ex Hook. Fl. Bor. Am. ii. 125 = mellifera.
villosa, *S. Wats. in Am. Naturalist*, vii. (1873) 302.—Am. bor.

ABROPHAËS, Rafin. Sylva Tellur. 99 (1838) = **Fothergilla**, Linn. f. (Hamamel.).
mirabilis, Rafin. l. c. = F. alnifolia.

ABROPHYLLUM, Hook. f. in Benth. Fl. Austral. ii. 437 (1864). *SAXIFRAGEAE*, Benth. & Hook. f. i. 547.
BRACHYNEMA, F. Muell. Fragm. iii. 90. (1862).
ornans, *Hook. f. l. c.*—Austral.

ABROTANELLA, Cass. in Dict. Sc. Nat. xxxvi. 27 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 431.
CRATELLA, Hook. f. Fl. Antarct. i. 23. t. 18 (1844).
SCLEROLEMA, Hook. f. in Lond. Journ. Bot. v. (1846) 444. t. 14.
TRIMEURON, Hook. f. Fl. Antarct. i. 23. t. 17 (1844).
emarginata, *Cass. in Dict. Sc. Nat.* xxxvi. 27 (1825).—Ins. Falkl.
forsteroides, *Hook. f. Handb. N. Zeal. Fl.* 139.—N. Zel.; Austral.
inconspicua, *Hook. f. l. c.* 140.—N. Zel.
linearifolia, *A. Gray, in Proc. Am. Acad.* v. (1861) 187.—Fuegia.
linearis, *Bergg. in Mimnekr. Fisiog. Sällsk. Lund* (1877), viii. p. 14.—N. Zel.
nivigena, *F. Muell. in Benth. Fl. Austral.* iii. 554.—Austral.
pusilla, *Hook. f. Handb. N. Zeal. Fl.* 139.—N. Zel.
rhynchocarpa, *Balf. f. in Phil. Trans.* clxviii. (1879) 359.—Ins. Rodr.
rosulata, *Hook. f. Handb. N. Zeal. Fl.* 139.—N. Zel.
scapigera, *F. Muell. ex Benth. Fl. Austral.* iii. 554.—Austral.
spathulata, *Hook. f. Handb. N. Zeal. Fl.* 139.—N. Zel.
submarginata, *A. Gray, in Proc. Am. Acad.* v. (1861) 187.—Regn. Magell.

ABROTANUM, [Tourn.] Linn. Syst. ed. I (1735) = **Artemisia**, Tourn. (Compos.).
alpestre, *amblyanum*, *brachylobium*, *congestum*, *incanescens*, *pauciflorum*, *pedunculare*, *platylobum*, *pulverulentum*, *rhodanicum*, *suave*, *virgatum*, *viridulum*, *serophilum*, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 69–72 = Art. Abrotanum.
frutescens, Gilib. Fl. Lituan. i. 173 = Art. campestris.

ABRUS, Linn. Hort. Cliff. 488 (1787). *LEGUMINOSAE*, Benth. & Hook. f. i. 587.
ZAGA, Rafin. Fl. Tellur. ii. 23 (1836).
acutifolius, Blume, ex Miq. Fl. Ind. Bat. i. 160 = pulchellus.
arboreus, Vell. Fl. Flum. 303; vii. t. 99 = Ormocarpum nitidum.
canescens, *Wehn. in Oliver, Fl. Trop. Afr.* ii. 175.—Afr. trop.
cantoniensis, *Hance, in Journ. Bot.* vi. (1868) 112.—China.
fruticulosus, *Wall. Cat.* n. 5830.—Ind. or.
laevigatus, E. Mey. Comm. i. 136 = pulchellus.
luscrius, *Vell. Fl. Flum.* 309; vii. t. 97.—Bras.
macrolestus, Noronha, in Verh. Batav. Gen. v. (1790), Art. iv. 7 = precatorius.
melanospermus, Hassk. Cat. Pl. Hort. Bog. Alt. 283 = pulchellus.
minor, Desv. in Ann. Sc. Nat. Sér. I. xi. (1826) 418 = precatorius.
mollis, *Hance, in Journ. Bot.* ix. (1871) 196.—China.
parciflorus, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 418 = precatorius.

ABUS :—
precatorius, *Linn. Syst. ed.* XII. 472.—Reg. trop.
pulchellus, *Wall. Cat.* n. 5819.—Ind. or.; Afr. austr.
squamulosus, E. Mey. Comm. i. 136 = precatorius.
tenuiflorus, *Benth. in Mart. Fl. Bras.* xv. i. 216.—Bras.

ABRYANTHEMUM, Neck. Elem. ii. 82 (1790) = **Mesembryanthemum**, Linn. (Ficoid.).

ABSINTHIUM, Tourn. ex Linn. Syst. ed. I (1735) = **Artemisia**, Tourn. (Compos.).
alpinum, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 240 = Art. lanata.
arborescens, Moench, Meth. 579 = Art. arborescens.
argentatum, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 272 = Art. argentea.
bipedale, Gilib. Fl. Lituan. i. 174 = Art. Absinthium.
borale, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 265 = Art. borealis.
campestre, Dulac, Fl. Hautes-Pyr. 502 = Art. campestris.
camphoratum, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 280 = Art. camphorata.
canariense, Besa. l. c. 229 = Art. canariensis.
candidum, Bieb. ex DC. Prod. vi. 125 = Art. frigida.
cernuum, Moench, Meth. 579 = Art. crithmifolia.
chinense, DC. Prod. vi. 118 sub n. 139 = Art. chinensis.
congestum, Lam. Fl. Fr. ii. 46 = Art. glacialis.
corymbosum, DC. Prod. vi. 122. sub n. 161 = Art. camphorata.
dentatum, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 265 = Art. rupestris.
fiscarianum, Fisch. ex Besa. l. c. 263 = Art. anethifolia.
frigidum, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 251 = Art. frigida.
glaciale, Lam. Illustr. iii. 273. t. 695 = Art. glacialis.
grandiflorum, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 282 = Art. grandiflora.
grandiflorum, Roehl. ex Stend. Nom. ed. II. i. 2 = Art. rupestris.
lagopchalium, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 238 = Art. lagocephala.
lanatum, Besa. l. c. 265 = Art. lanata.
Linnæi, Fl. Fr. ii. 46 = Art. Mutellina.
mollissimum, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 265 = Art. mollissima.
Mosæ, Besa. l. c. 238 = Art. Mosa.
Mutellina, Roehl. ex Stend. Nom. ed. II. i. 2 = Art. Mutellina.
nitens, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 235 = Art. sericea.
officinale, Brot. Fl. Lusit. i. 357 = Art. Absinthium.
pedunculare, Ster. in Mém. Soc. Nat. Mosc. iv. (1813) 63 = Art. splendens.
petrosum, Baumg. Enum. Stirp. Transs. iii. 90 = Art. Baumgartenii.
ponticum, Besa. Enum. Pl. Volh. 76 = Art. pontica.
rupestre, Schrank, Prim. Fl. Salish. 303 = Art. rupestris.
santonicum, Gueldenst. ex Ledeb. Fl. Ross. ii. 573 = Art. maritima.
saxatile, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 231 = Art. camphorata.
sericeum, Besa. l. c. 237 = Art. sericea.
Sieversianum, Besa. l. c. 259 = Art. Sieversiana.
spathulatum, Noronha, in Verh. Batav. Gen. v. (1790) Art. iv. 6 = Dichrocephala latifolia.
spicatum, Baumg. Enum. Stirp. Transs. iii. 90 = Art. spicata.
splendens, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 238 = Art. splendens.
tanacetifolium, Gaertn. Fruct. ii. 393 = Art. tanacetifolia.
trifidum, Schrank, Baier. Fl. ii. 362 = Art. austriaca.
virescens, Jusa. ex Besa. in Bull. Soc. Nat. Mosc. viii. (1835) 29 = Art. Jussiæana.
virgatum, Turcz. ex Ledeb. Fl. Ross. ii. 597 = Art. rupestris.
viride, Besa. in Bull. Soc. Nat. Mosc. i. (1829) 249 = Art. rupestris.
viridiflorum, Besa. l. c. 246 = Art. rupestris.
vulgare, Dulac, Fl. Hautes-Pyr. 502 = Art. vulgaris.
vulgare, Lam. Fl. Fr. ii. 45 = Art. Absinthium.

ABSYNTHIUM, Gaertn. Meyer & Scherb. Fl. Wett. iii. 196 (1801) = prec.

ABULFALI, Adans. Fam. ii. 290 (1763) = **Thymbra**, Linn. (Labiat.).

ABUMON, Adans. Fam. ii. 84 (1763) = **Agapanthus**, L'Hérit. (Liliac.).

ABUTA, [Barr.] Aubl. Pl. Gui. i. 618. t. 250 (1775). *MENISPERMACEAE*, Benth. & Hook. f. i. 35.
ANELARIA, Miers, in Ann. Nat. Hist. Ser. II. vii. (1851) 42.
BATSCHIA, Mutis, ex Thunb. in Nov. Act. Soc. Sc. Upsal. v. (1792) 120. t. 3.
TRICHOA, Pers. Syn. ii. 634 (1807).
acutifolia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 259.—Bras.
amara, Aubl. Pl. Gui. i. 620 = Aristolochia glaucescens.
barbata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 258.—Guiana.
candicans, *Rich. ex DC. Syst.* i. 548.—Guiana.
Candolleï, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 47.—Guiana.
concolor, Benth. in Journ. Linn. Soc. v. (1861) App. 49 = guianensis.
concolor, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 64. t. 188.—Am. austr.
Griebachii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 47.—Bras.
guianensis, *Eichl. in Flora*, xlvii. (1864) 390.—Guiana.
heterophylla, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 256.—Am. austr.
Imene, *Eichl. in Flora*, xlvii. (1864) 389.—Bras.
macrophylla, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 256.—Am. austr.
oblonga, *Miers, l. c.*—Am. austr.
oblongifolia, *Miers, l. c.*—Am. austr.
panurensis, *Eichl. in Flora*, xlvii. (1864) 390.—Bras.
Natophylla, Mart. ex Eichl. in Mart. Fl. Bras. xiii. i. 200 (= *Botryopsis platyphylla*).—Bras.
racemosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 48.—N. Granat.
rufescens, *Aubl. Pl. Gui.* i. 618. t. 250.—Guiana.
rufescens, DC. Syst. i. 542 = Candollei.
rufescens, Griseb. in Journ. Linn. Soc. iii. (1859) 109 = Griebachii.
scandens, DC. Syst. i. 542 in syn. = rufescens.
Seemanni, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 50.—Darien.
Selloana, *Eichl. in Flora*, xlvii. (1864) 389.—Bras.
spicata, *Triana & Planch. in Ann. Sc. Nat.* Sér. II. xvii. (1862) 119.—N. Granat.
tomentosa, Sagot, ex Benth. in Journ. Linn. Soc. v. (1861) Suppl. ii. 51 (Coccoloboe sp.).—Guian.
umbellata, Sagot, ex Benth. l. c. = Sciadotaenia cayennensis.

ABUTILAEA, F. Muell. in Linnæa, xxv. (1852) 379 = Abutilon, Tourn. (Malvac.).
cryptantha, F. Muell. l. c. = Abutilon Mitchelii.

ABUTILON, Tourn. ex Adans. Fam. ii. 398 (1763). *MALVACEAE*, Benth. & Hook. f. i. 204.
ABUTILARA, F. Muell. in Linnæa, xxv. (1852) 379.
BELOÉRA, Shuttlew. ex Benth. & Hook. f. Gen. i. (1862) 204.
BOGENHARDIA, Reichb. Nom. 200 (1841).
HERIBAUNTIA, Medic. Phil. Bot. i. 90 (1789).
LASS, Adans. Fam. ii. 400 (1763).
PRESTONIA, Soop. Introd. 281 (1777).
acerifolium, G. Don, Gen. Syst. i. 504 = Sida acerifolia.
affine, *G. Don, l. c.* 503.—Reg. Argent.
albescens, Miq. Fl. Jungh. 285 = crispum.
albidum, Sweet, Hort. Brit. ed. I. 54.—Ins. Canar.
albidum, *Walp. Rep.* i. 235.—Mexic.
albidum, Webb & Berth. Phyt. Canar. i. 89. t. 2 = fruticosum.
amboynense, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 200 = Sida Lindeniana.
americanum, Sweet, Hort. Brit. ed. I. 58 = indicum.
amplexifolium, *G. Don, Gen. Syst.* i. 509.—Mexic.
amplum, *Benth. Fl. Austral.* i. 200.—Austral.
Andersonianum, *Garcke, ex Anderss. Gulap. Our. Vegetat.* 280.—Ins. Galap.
Andicaudi, *Hassk. Diagn. Pl. Nov.* 94.—Mexic.
angulatum, Mast. in Oliver, Fl. Trop. Afr. i. 183 = intermedium.
anodoides, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 49.—Bras.
arborescens, Medic. Malv. 29 = indicum.
arborescum, *Sweet, Hort. Brit.* ed. I. 58.—Peruv.
Arnottianum, Walp. Rep. i. 234 = Sida Arnottiana.
asiaticum, *G. Don, Gen. Syst.* i. 508.—Reg. trop.
asiaticum, Guill. & Perr. Fl. Senegj. Tent. i. 87 = glaucum.
asiaticum, Sweet, Hort. Brit. ed. I. 58 = indicum.
atropurpureum, *G. Don, Gen. Syst.* i. 509.—Java.

ABUTILON :—

aurantiacum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 357.—Calif.

aureum, G. Don, in Sweet, Hort. Brit. ed. III. 80 = Sida aurea.

aurium, *Sweet, l. c.* ed. I. 58.—Geront. trop.

australe, *Presl, Bat. Bemerk.* 20 ; *Garcke, Ind. Sem. Hort. Berol.* (1861) 10.—Austral.

Avicennæ, *Gaertn. Fruct. ii.* 251. t. 135.—Geront. trop.

Bedfordianum, *A. St. Hil. in Ann. Sc. Nat.* Sér. II. xvIII. (1842) 48.—Bras.

Behrianum, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 13 = A. Avicennæ.

benedictum, *Bunb. in Proc. Linn. Soc.* i. (1841) 109.—Bras.

bidentatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 68. —Abyss. : Ind. or.

blandum, *Fenzl, Delect. Sem. Hort. Bot. Vindob.* (1858).—Mexic.

californicum, Benth. Bot. Voy. Sulph. 8 = A. Avicennæ.

calycinum, *Presl, Rel. Haenk.* ii. 115.—Peruv.

carneum, *A. St. Hil. Fl. Bras. Mer.* i. 205.—Bras.

ceratocarpum, *Hook. Bot. Misc.* iii. (1833) 154.—Chili.

Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 508.—Madag.

cinereum, *Griseb. in Goett. Abh.* xxiv. (1879) 45.—Reg. Argent.

circinnatum, *G. Don, Gen. Syst.* i. 502.—Bras.

cistifiorum, *G. Don, l. c.* 503.—Java.

coccineum, *Hort. ex Regel, Cat. Pl. Hort. Aksakov.* (1860) 3, nomen.—Hab. ?

confertiflorum, *A. Rich. Ess. Fl. Cub.* i. 158.—Cuba.

contractum, Sweet, Hort. Brit. ed. II. 64 = Sida contracta.

cornutum, Sweet, l. c. 65 = Sida cornuta.

crassiflolium, *G. Don, Gen. Syst.* i. 504.—Ind. occ.

criapum, *Sweet, Hort. Brit.* ed. I. 53.—Reg. trop.

cryptopetalum, *F. Muell. ex Benth. Fl. Austral.* i. 201. —Austral.

Cunninghamii, *Benth. l. c.* 205.—Austral.

cynosum, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 185.—Am. trop.

cysticarpum, Hance, in Walp. Ann. ii. 157 = indicum.

Darwini, *Hook. f. Bot. Mag.* t. 5917.—Bras.

decurpum, *Raffin. New Fl. Am.* i. 40.—Am. bor.

deflexum, *Moench, Meth.* 620.—Hab ?

densiflorum, *Walp. Rep.* i. 324.—Chili.

denticulatum, Webb, Fragm. Fl. Aethiop. 51 = fruticosum.

depauperatum, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 282.—Ins. Galap.

dianthum, *Presl, Reliq. Haenk.* ii. 114.—Peruv.

diplotrichum, F. Muell. in Linnæa, xxv. (1852) 380 = Fraseri.

disclusum, *Schlecht. in Linnæa*, 218.—Reg. Argent.

divaricatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 204.—Mexic.

domingense, *Turcz. l. c.* 205.—Ins. S. Doming.

elaeocarpoides, Webb, Fragm. Fl. Aethiop. 53 = ramosum.

elatum, *Griseb. Fl. Brit. W. Ind.* 79.—Ind. occ. ; Mexic.

elegans, *A. St. Hil. Fl. Bras. Mer.* i. 207.—Bras.

elegans, Walp. Rep. i. 325 = Sida elegans.

ellipticum, *Schlecht. in Linnæa*, xi. (1837) 868.—Mexic.

elongatum, Moench, Meth. Suppl. 205 = indicum.

erosum, *Schlecht. in Linnæa*, xi. (1837) 867.—Mexic.

esculentum, *A. St. Hil. Fl. Bras. Mer.* i. 240.—Bras.

exoletior, G. Don, Gen. Syst. i. 500 = Wissadula exoletior.

exonemum, *F. Muell. Fragm.* xi. 63.—Austral.

exstipulare, *G. Don, Gen. Syst.* i. 508.—Ins. Borbon.

falcatum, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvIII. (1842) 48.—Bras.

ferruginosum, H. B. & K. Nov. Gen. et Sp. v. 271 = Sida ferruginea.

Figarianum, Webb, Fragm. Fl. Aethiop. 52 = graveolens.

floribundum, *Schlecht. in Linnæa*, xi. (1837) 866.—Mexic.

foetidum, *Moench, Meth. Suppl.* 206.—Hab. ?

Fraseri, *Walp. Ann.* ii. 158.—Austral.

fruticosum, Medic. Malv. 29 = indicum.

fruticosum, *Guill. & Perr. Fl. Seneg. Tent.* i. 70.—As. et Afr. trop.

furfuraceum, *Miq. Fl. Ind. Bat.* i. ii. 144.—Java.

gossypiiformum, H. B. & K. Nov. Gen. et Sp. v. 274. t. 474 = Bastarnea.

gossypioides, *Bunb. Fl. Austroal.* i. 202.—Austral.

ABUTILON :—

giganteum, *Presl, Rel. Haenk.* ii. 116.—Am. trop.

glaucum, *G. Don, Gen. Syst.* i. 504.—Afr. trop.

glaucum, Sweet, Hort. Brit. ed. I. 54 = muticum.

glechomaefolium, *A. St. Hil. Fl. Bras. Mer.* i. 198. t. 41.—Bras.

globiflorum, *G. Don, Gen. Syst.* i. 502.—Maurit.

Goudotianum, *Triana & Planch. in Ann. Sc. Nat* Sér. IV. xvI. (1862) 184.—N. Granat.

grandiflorum, G. Don, Gen. Syst. i. 504 = indicum.

grandifolium, Sweet, Hort. Brit. ed. I. 53 = molle.

graveolens, Seem. Bot. Voy. Herald, 365 = indicum.

graveolens, *Wight & Arn. Prod.* 56.—Geront. trop.

Grevilleanum, *Walp. Rep.* i. 324.—Chili.

Guichenotianum, *Decne. in Herb. Timor. Descr.* 106 (1858).—Ins. Timor.

gynanthemum, *Griseb. in Goett. Abh.* xxiv. (1879) 47. —Reg. Argent.

Haenkeanum, *Presl, Rel. Haenk.* ii. 115.—Mexic.

halophilum, F. Muell. in Linnæa, xxv. (1852) 381 = Fraseri.

hernandioides, Sweet, Hort. Brit. ed. I. 53 = Sida hernandioides.

heterophyllum, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 187 = hirtum.

hirsutissimum, Moench, Meth. Suppl. 205 = indicum.

hirtum, *G. Don, Gen. Syst.* i. 503.—Reg. trop.

hirtum, Sweet, Hort. Brit. ed. I. 53 = graveolens.

holosericeum, *Scheele, in Linnæa*, xxi. (1847) 471.—Texas.

Hulseanum, *Torr. ex A. Gray, Pl. Fendl.* 28.—N. Mexic.

hypoleucum, *A. Gray, Pl. Wright.* i. 20.—Texas.

ibarreuse, *H. B. & K. Nov. Gen. et Sp.* v. 272.—Ecuador.

imberbe, G. Don, Gen. Syst. i. 503 = crispum.

inaequilaterum, *A. St. Hil. Fl. Bras. Mer.* i. 198. t. 40. —Bras.

incanum, *Sweet, Hort. Brit.* ed. I. 53.—Ins. Sandvic. ; Mexic.

indicum, *Sweet, l. c.* 54.—Geront. trop.

insigne, *Planch. in Van Houtte, Fl. des Serres*, Sér. I. vi. (1850-51) 41.—N. Granat.

integerrimum, *Hook. Bot. Mag.* t. 4360.—Mexic.

intermedium, *Hochst. ex Schweinf. Beitr. Fl. Aethiop.* 49.—Geront. trop.

Jacquinii, *G. Don, Gen. Syst.* i. 508.—Am bor.

Julianae, *Endl. Prod. Fl. Norf.* 185.—Ins. Norfolc.

Kotschyi, Hochst. in Webb, Fragm. Fl. Aethiop. 52 = hirtum.

lanatum, *Miq. in Linnaea*, xxii. (1849) 553.—Bras.

laxiflorum, Guill. & Perr. Fl. Seneg. Tent. i. 66 = Wissadula rostrata.

leiocarpum, *Griseb. Fl. Brit. W. Ind.* 79.—Ind. occ.

Lemmoni, *S. Wats. in Proc Am. Acad.* xx. (1885) 357.—Calif.

lepidum, F. Muell. Fragm. vi. 168 = Sida lepida.

Leschenaultianum, Sweet, Hort. Brit. ed. I. 58 = Wissadula Leschenaultiana.

Leucanthemum, *A. St. Hil. Fl. Bras. Mer.* i. 200.—Bras.

leucopetalum, *F. Muell. ex Benth. Fl. Austral.* i. 200. —Austral.

lignosum, G. Don, Gen. Syst. i. 501 = Jacquini.

longicuspe, *Hochst. ex A. Rich. Fl. Abyss.* i. 69.—Abyssin.

longiflobum, *F. Muell. Fragm.* ix. 130.—Austral.

lucianum, *Sweet, Hort. Brit.* ed. I. 53.—Ind. occ.

macranthum, *Peyr. in Linnæa*, xxx. (1859-60) 59.—Mexic.

macranthum, *A. St. Hil. Fl. Bras. Mer.* i. 208.—Bras.

macrocarpum, *Guill. & Perr. Fl. Seneg. Tent.* i. 69.—Afr. trop.

macrocarpum, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvIII. (1842) 47.—Bras.

macrophyllum, *St. Hil. & Naud. l. c.*—Bras.

macropodum, *Guill. & Perr. Fl. Seneg. Tent.* i. 69. t. 14.—Afr. trop.

macro, *F. Muell Fragm.* ix. 59.—Austral.

malachroides, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvIII. (1842) 49.—Bras.

marmoratum, Hort. ex Lem. in Illustr. Hortic. iii. (1856) t. 93 = Hibiscus marmoratus.

mauritianum, Medic. Malv. 28 = sec. *asiaticum*, Sweet, Hort. Brit. ed. I. 53 = indicum.

megapotamicum, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvIII. (1842) 49.—Am. trop.

melanocarpum, *St. Hil. & Naud. l. c.* 48.—Bras.

mendocinum, *Phil. Sert. Mend. Alt.* 6.—Chili.

Mentzelii, *Szem. Fl. Vit.* 15.—Ins. Sandvic.

mexicanum, *Presl, Rel. Haenk.* ii. 115.—Mexic.

ABUTILON :—

micranthum, G. Don, Gen. Syst. i. 502, sphalm. = macranthum.

micropetalum, *Benth. Fl. Australe.* i. 201.—Austral.

microphyllum, A. Rich. Fl. Abyss. i. 70 = fruticosum.

microspermum, *G. Don, Gen. Syst.* i. 501.—Hab. ? Mitchelii, *Benth. Fl. Austral.* i. 201.—Austral.

molle, *Sweet, Hort. Brit.* ed. II. 65.—Peruv.

mollicomum, *Sweet, l. c.* ed. I. 54.—Hab. ?

mollissimum, *Sweet, l. c.* 58.—Peruv.

montanum, *A. St. Hil. Fl. Bras. Mer.* i. 207.—Bras.

mucronulatum, A. Gray, in Proc. Am. Acad. v. (1862) 178 = Wissadula mucronulata.

muticum, *Sweet, Hort. Brit.* ed. II. 65.—Geront. trop.

muticum, Webb, Fragm. Fl. Aethiop. 51 = glaucum.

neligherrense, *Munro, in Wight, Ill.* t. 66.—Ind. or.

neurocarpum, *Miq. Fl. Sungk.* 335.—Java.

Newberryi, *S. Wats. in Proc. Am. Acad.* xi. (1876) 125.—Am. bor.

niveum, *Griseb. in Goett. Abh.* xix. (1874) 92.—Reg. Argent.

notolophium, *A. Gray, in Proc. Am. Acad.* v. (1875) 176.—Mexic.

nudiflorum, Sweet, Hort. Brit. ed. I. 58 = Wissadula nudiflora.

Nuttallii, *Torr. & Gray, Fl. N. Am.* i. 231.—Am. bor.

obtusum, *G. Don, Gen. Syst.* i. 501.—Hab. ?

occidentale, Medic. Malv. 29 = Gaya subtriloba.

Ochsenii, *Phil. in Linnaea*, xxviii. (1856) 613.—Chili.

orbiculatum, *G. Don, Gen. Syst.* i. 504.—China.

otocarpum, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 13.—Austral.

oxycarpum, *F. Muell. Fragm.* ii. 12.—Austral.

oxypetalum, *Planch & Linden, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 184.—N. Granat.

oxyphyllum, Presl, Bot. Bemerk. 20 = polyandrum.

paeoniaefiorum, *Walp. Rep.* v. 95.—Bras.

Palmeri, *A. Gray, in Proc. Am. Acad.* viii. (1870) 289. —Am. bor.

pannosum, Webb, Fragm. Fl. Aethiop. 52 = glaucum.

paranthemoides, *Griseb. in Goett. Abh.* xxiv. (1879) 46.—Reg. Argent

paranthemum, *Griseb. l. c.*—Reg. Argent.

Parishii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 357. —Arizona.

parviflorum, *A. St. Hil. Fl. Bras. Mer.* i. 201.—Bras.

parvulum, *A. Gray, Pl. Wright.* i. 21.—Texas.

patens, *A. St. Hil. Fl. Bras. Mer.* i. 200.—Bras.

paucifiorum, *A. St. Hil.* t. c. 208.—Bras.

pedunculare, *H. B. & K. Nov. Gen. et Sp.* v. 273.—Bras.

peraßne, Shuttl. ex Chapm. Fl. S. U. St. 58 = Jacquini.

periplocifolium, Sweet, Hort. Brit. ed. I. 53 = Wissadula rostrata.

pernolle, *Sweet, l. c.*—Florida ; Cuba.

petiolare, *H. B. & K. Nov. Gen. et Sp.* v. 273.—N. Granat.

pictum, *Walp. Rep.* i. 324.—Reg. Argent.

planiflorum, *C. Koch & Bouché, in C. Koch, Allg. Gartenz.* (1857) 98.—Mexic.

polyandrum, *G. Don, Gen. Syst.* i. 500.—Ind. or.

polyanthon, Sweet, Hort. Brit. ed. I. 53 = Sida polyantha.

populifolium, Sweet, l. c. = indicum.

pubescens, Moench, Meth. 620 = A. Avicennae.

pulchellum, Sweet, Hort. Brit. ed. I. 53 = sec. *pulchrum*, G. Don, Gen. Syst. i. 503 = Plagianthus pulchellus.

pyramidale, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 203.—N. Granat.

racemosum, *Schlecht. in Linnaea*, xi. (1837) 867.—Mexic.

ramiflorum, *A. St. Hil. Fl. Bras. Mer.* i. 199.—Bras.

ramosissimum, *Presl, Rel. Haenk.* ii. 118.—Mexic.

ramosum, G. Don, Gen. Syst. i. 501 = Sida cordifolia.

ramosum, *Guill. & Perr. Fl. Seneg. Tent.* i. 66.—Geront. trop.

reflexum, *Sweet, Hort. Brit.* ed. I. 53 ; *Presl, Rel. Haenk.* ii. 114.—Mexic.

Regnellii, *Miq. in Linnaea*, xxii. (1849) 554.—Bras.

rigidum, *G. Don, Gen. Syst.* i. 504.—Afr. trop.

rivulare, *A. St. Hil. Fl. Bras. Mer.* i. 202.—Bras.

rufescens, G. Don, Gen. Syst. i. 503 = rufiserve.

rufescens, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 202.—Mexic.

rufinerve, *A. St. Hil. Fl. Bras. Mer.* i. 205. t. 42.—Bras.

rufinervo, Seem. Bot. Voy. Herald, 63 = cymosum.

Sellowianum, *Regel, in Ann. Sc. Nat.* Sér. IV. xii. (1859) 379.—Bras.

septemlobum, *Miq. in Linnaea*, xxii. (1849) 554.—Bras.

sessiliflorum, *Presl, Rel. Haenk.* ii. 118.—Mexic.

ABUTILON :—
sidoides, Dalr. & Gibs. Bomb. Fl. 18 = ramosum.
sidoides, *Hemsl. Diagn. Pl. Nov.* 24.—Mexic.
Sonnerattianum, *Sweet, Hort. Brit.* ed. I. 54.—Afr. austr.
Sonorae, *A. Gray, Pl. Wright.* ii. 23.—Mexic.
spermannioides, Guill. & Perr. Fl. Seneg. Tent. i. 70 = ramosum.
speciosum, G. Don, Gen. Syst. i. 502 = Sida speciosa.
spicatum, H. B. & K. Nov. Gen. et Sp. v. 271 = Wissadula spicata.
stenopetalum, *Garcke, in Bot. Zeit.* viii. (1850) 683.—N. Granat.
stipulare, *Presl, Rel. Haenk.* ii. 114.—Luzon.
striatum, *Dicks. in Bot. Reg.* (1839) *Misc.* 89.—Bras. subviscosum, *Benth. Fl. Austral.* i. 202.—Austral.
sundaicum, *G. Don, Gen. Syst.* i. 500 (1831).—Java, terminale, *A. St. Hil. Fl. Bras. Mer.* i. 203.—Bras.
teamae, Torr. & Gray, Fl. N. Am. i. 231 = incanum.
Theophrasti, Medic. Malv. 38 = A. Avicennae.
Thurberi, *A. Gray, Pl. Nov. Thurb.* 307.—Texas. ; N. Mexic.
thyrsodendron, *Griseb. in Goett. Abh.* xxiv. (1879) 48. —Reg. Argent.
tiliaefolium, Sweet, Hort. Brit. ed. I. 53 = A. Avicennae.
timoriense, *G. Don, Gen. Syst.* i. 500 (1831).—Ins. Timor.
tomentosum, Wight & Arn. Prod. 56 = muticum.
tortuosum, Guill. & Perr. Fl. Seneg. Tent. i. 66 = graveolens.
trichodum, A. Rich. Ess. Fl. Cub. i. 155 = crispum.
trilobatum, *Hemsl. Diagn. Pl. Nov. Mex.* 24.—Mexic.
triquetrum, *Presl, Relig. Haenk.* ii. 115.—Mexic.
triquetrum, *Sweet, Hort. Brit.* ed. I. 53.—Ind. occ.
truncatum, *G. Don, Gen. Syst.* i. 503.—Ins. S. Doming. tubulosum, *Walp. Ann.* ii. 158.—Austral.
umbellatum, *Sweet, Hort. Brit.* ed. I. 53.—Ind. occ.
umbelliforme, G. Don, Gen. Syst. i. 501 = seq.
umbelliforme, *A. St. Hil. Fl. Bras. Mer.* i. 204.—Bras.
ureticarpum, Walp. Rep. i. 324 = cornucopiae.
velutinum, *G. Don, Gen. Syst.* i. 504.—Afr. trop.
velutinum, A. Gray, Illustr. Gen. Fl. Am. Bor. ii. 67. t. 135 = holosericeum.
venosum, *Lem. Hortic. Univ. Misc.* 1844 ; *ex Fl. des Serres*, ii. (Mars 1846) t. 3.—Mexic.
verbascoides, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 202.—Venezuela.
vesicarium, Sweet, Hort. Brit. ed. I. 54 = A. Avicennae.
vexillarium, E. Morr. Belg. Hortic. (1864) 259 = megapotamicum.
virens, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 48.—Bras.
virgatum, *Sweet, Hort. Brit.* ed. I. 53.—Hab. ?
vitifolium, *Presl, Rel. Haenk.* ii. 116.—Chili.
Wagenerianum, *Regel, Cat. Hort. Akademov.* (1860) 1, nomen.—Hab. ?
wissadifolium, *Griseb. in Goett. Abh.* xxiv. (1879) 47. —Reg. Argent.
Wrightii, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 169.—Texas.
zanzibaricum, *Bef. ex Mast. in Oliver, Fl. Trop. Afr.* i. 186.—Afr. trop.

ABUTUA, Lour. Fl. Cochinch. 680 (1790) = **Gnetum**, Linn.
africana, Lour. l. c. 681 (Quid ?).—Afr. or. *indica*, Lour. l. c. 680 = G. funiculare.

ACACALLIS, Lindl. Fol. Orchid. 1853. *ORCHIDEAE*, Benth. & Hook. f. iii. 544.
cyanea, *Lindl. l. c.*—Bras.

ACACIA, [Toum.] Linn. Fl. Zeyl. 217 (1737) ; Willd. Sp. Pl. iv. 1049 (1805). *LEGUMINOSAE*, Benth. & Hook. f. i. 594.
ARTHROSPHON, Hask. Retzia, i. 212 (1855).
CHITONANTHUS, Lehm. Pl. Preiss. ii. 368 (1847).
CUPARILLA, DERPAPHYLA, EBURNAX, ESCLERONA, GUMIFERA, HECATANDRA, POPONAX, SENEGALIA, ZIGMALOBA, Rafin. Sylva Tellur. 118–120 (1838).
FARNESIA, Gasp. Desc. Nuov. Gen. (1838).
HOOPESIA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 433, partim.
PHYLLODOCE, Link, Handb. ii. 132 (1831).
TETRACHEILOS, Lehm. Pl. Preiss. ii. 368 (1847).
VACHELLIA, Wight & Arn. Prod. 272 (1834).
abietina, Willd. Sp. Pl. iv. 1051 = linifolia.
abstergens, Steud. Nom. ed. II. i. 3 = concinna.
abyssinica, *Hochst. ex Benth. in Hook. Lond. Journ. Bot.* v. (1846) 97.—Abyssin.
ACACIA :—
acanthocarpa, Willd. Enum. Hort. Berol. 1057 = Mimosa acanthocarpa.
acanthocalaba, *F. Muell. Fragm.* iii. 127.—Austral.
acantholoba, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1089 = Mimosa acantholoba.
acanthophora, Steud. Nom. ed. II. i. 3 = Mimosa acanthocarpa.
acapulcensis, Kunth, Mim. 78. t. 24 = Lysiloma acapulcensis.
acatlensis, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 513.—Mexic.
acicularis, K. Br. in Ait. Hort. Kew. ed. II. v. 450 = juniperina.
acicularis, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 1056 = Farnesiana.
acinacea, *Lindl. in Mitch. Three Exped.* ii. 267.—Austral.
aciphylla, *Benth. in Linnaea*, xxvi. (1853–55) 627.—Austral.
Acradenia, F. Muell. in Journ. Linn. Soc. iii. (1859) 149 = umbellata.
acuifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 496.—Ins. Baham.
aculeaticarpa, Lag. Gen. et Sp. Nov. 16 = Mimosa acanthocarpa.
acuminata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 373.—Austral.
acuticarpa, Buch.-Ham. ex Wall. Cat. sub n. 5271 = Pithecolobium angulatum.
acutistipula, Mart. in Flora, xx. (1837) Beibl. ii. 107 = Mimosa acutistipula.
Adansonii, Guill. & Perr. Fl. Seneg. Tent. i. 249 = arabica.
adenantheroides, Mart. & Gal. in Bull. Acad. Brux. xii. (1845) 311 = Mimosa adenanthera.
adenocarpa, Walp. Rep. i. 924, sphalm. = seq.
adenopa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 206 = A. Farnesiana.
adenophylla, Spreng. Syst. iii. 140 = decurrens.
adenostylis, Fenzl, in Flora, xxvii. (1844) 312 = Dichrostachys nutans.
adnascens, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 517.—Bras.
adiantoides, Spreng. Syst. iii. 146 = Piptadenia laxa.
adnata, *F. Muell. in Melb. Chemist* (July 1882).—Austral.
adstringens, A. Cunn. in G. Don, Gen. Syst iii. 405 = falciformis.
adstringens, Mart. Reise Bras. i. 548—Stryphnodendron Barbatimam.
aduncа, A. Cunn. ex G. Don, Gen. Syst. iii. 406 = crassiuscula.
aegyptiaca, *Baill. Adansonia*, iv. (1863–64) 82.—Aegypt.
affinis, Sweet, Hort. Brit. ed. I. 102 = dealbata.
alata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 464.—Austral.
alba, Lamk, Hort. Rip. App. ii. 389 = Calliandra portoricensis.
alba, *De Vriese, in Mig. Fl. Sungh.* 270.—Malaya.
alba, Willd. Enum. Hort. Berol. Suppl. 68 = leucophloea.
albacora, Stend. Nom. ed. II. i. 3 = tortuosa.
albicans, Kunth, Mim. 87. t. 27 = Pithecolobium albicans.
albida, *Delile, Fl. Egypt.* 286. t. 52. f. 3.—Afr. bor.
albida, Lindl. Bot. Reg. t. 1317 = tortuosa.
alliacea, Buch.-Ham. in Wall. Cat. n. 5258 = caesia.
alpina, *F. Muell. Fragm.* iii. 129.—Austral.
amara, Willd. Sp. Pl. iv. 1074 = Albizzia amara.
amazonica, *Benth. in Trans. Linn. Soc.* xxx. (1875) 593.—Am. trop.
ambigua, *Hoffmgg. Vers. Pfl. Nachtr.* iii. 15.—Afr. trop.
ambigua, *Vogel, in Linnaea*, x. (1835–36) 600.—Am. trop.
ambivcarpa, R. Grah. in Wall. Cat. n. 5260 = pennata.
amblygona, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 392.—Austral.
amblyphylla, *F. Muell. in Wing, South Sc. Record* (July 1882).—Austral.
amentacea, *DC. Prod.* ii. 455.—Am. bor.
amentifera, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 141.—Austral.
amorna, Sieber, ex Walp. Rep. i. 897 = rubida.
amoena, *Wendl. Comm. Acac.* 16. t. 4.—Austral.
amythorthophylla, *Steud. ex A. Rich. Fl. Abyss.* i. 245. —Abyssin.
anceps, *DC. Mém. Lég.* xii. 446.—Austral.
anceps, Hook. Ic. Pl. t. 167 = complanata.
ancura, *F. Muell. in Linnaea*, xxvi. (1853–55) 627.—Austral.
Angico, Mart. Mat. Med. Bras. 53 = Piptadenia rigida.
angulata, Desv. Journ. Bot. iii. (1814) 88 = discolor.
ACACIA :—
angulosa, Bertol. Fl. Guatim. 42 = filicina.
angustifolia, Lodd. Bot. Cab. t. 768 = longifolia.
angustifolia, Mciisn. in Lehm. Pl. Preiss. i. 13 = triquetra.
angustifolia, Wendl. Comm. Acac. 34 = suaveolens.
angustifolia, DC. Prod. ii. 470 = Pithecolobium peregrinum.
angustistipua, Desf. Cat. Hort. Par. ed. III. 300 = Desmanthus virgatus.
anthelmintica, Balli. Hist. Pl. ii. 56 = Albizzia anthelmintica.
apiculata, Meisn. in Lehm. Pl. Preiss. i. 17 = subcaerulea.
arabica, *Willd. Sp. Pl.* iv. 1085.—Afr. ; As. trop.
Arav, Buch.-Ham. in Wall. Cat. n. 5240 = A. latisiliqua.
arborea, Benth. Fl. Hartw. 18 = Lysiloma microphylla.
arborea, Willd. Sp. Pl. iv. 1064 = Pithecolobium filicifolium.
Arouthos, Spreng. Syst. iii. 134 = juniperina.
arcuata, *Decne. Herb. Timor. Descr.* 133.—Ins. Timor.
arcuata, Sieber, ex Spreng. Syst. iii. 135 = melanoxylon.
arenosa, Willd. Sp. Pl. iv. 1054 (err. typ. 1060) = Mimosa arenosa.
argyrophylla, Hook. ex F. Muell. Pl. Vict. ii. 18 = brachybotrya.
arida, *A. Cunn. ex G. Don, Gen. Syst.* iii. 404.—Austral.
armata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 463.—Austral.
armata, Heyne, ex Wall. Cat. n. 9564 = A. Farnesiana.
armatoides, Hort. ex Walp. Rep. i. 886 = armata.
Aroma, Gill. in Hook. Bot. Misc. iii. (1833) 206 = macracantha.
aromatica, Poepp. ex Benth. in Trans. Linn. Soc. xxx. (1875) 502 = A. Farnesiana.
Arrophila, D. Don ! in Wall. Cat. n. 5257 = Albizzia stipulata !
Arrophula, D. Don. Prod. Fl. Nep. 247 = pennata.
Asak, Willd. Sp. Pl. iv. 1077 = hamulosa ?
asparagoides, *A. Cunn. in Field, N. S. Wales*, 348. —Austral.
aspera, *Lindl. in Mitch. Three Exped.* ii. 139.—Austral.
asperulacea, F. Muell. in Journ. Linn. Soc. iii. (1859) 128 = leptocephala.
aspidioides, G. F. W. Mey. Prim. Fl. Esseq. 165 = Pentaclethra filamentosa.
asplenioides, Nees, in Flora, iv. (1821) 803 = Calliandra merteniioides.
astringens, A. Cunn. ex G. Don, Gen. Syst. ii. 405 = penninervis.
atazacantha, *DC. Prod.* ii. 459.—Afr. trop.
ataxiphylla, *Benth. in Linnaea*, xxvi. (1853–55) 605. —Austral.
atomiphylla, Bunch. Trav. i. 341 = haematoxylon.
atramentaria, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 392.—Chili.
Auchori, Benth. l. c. 499 = nubica.
aulacocarpa, *A. Cunn. ex Benth. l. c.* 379.—Austral.
aurea, *Noronha, in Verh. Batav. Gen.* v. (1790) Art. iv. 5.—Java.
auriculaeformis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 377.—Austral.
auriculata, Buch.-Ham. ex Wall. Cat. sub n. 5237 = Albizzia stipulata.
auriculata, Cunn. ex Fourn. in Bull. Soc. Bot. Fr. xii. (1865) 400 = Albizzia auriculata.
aurita, Schlecht. in Linnaea, xii. (1838) 565 = Lysiloma aurita.
auronitens, *Lindl. Swan River App.* 15.—Austral.
Ausfeldi, Regel, Ind. Sem. Hort. Petrop. (1866) 106 ; Gartenfl. xvi. (1867) 225 = sapera.
axillaris, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 341.—Tasmania.
Azak, G. Don, Gen. Syst. ii. 413 = hamulosa ?
Bagueri, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 377.—Austral.
bahamensis, Griseb. Fl. Brit. W. Ind. 221 = Lysiloma latisiliqua.
balsamia, *Benth. in Trans. Linn. Soc.* xxx. (1875) 525.—Am. trop.
Balbisii, Spreng. Syst. iii. 140 = Pithecolobium fragrans.
Banoyrahaea, Bert. ex Colla, Hort. Rip. 1 ; DC. Prod. ii. 473 = Caesalpinia bijuga.
barbinervis, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 390.—Austral.
basaltica, F. Muell. & Benth. in Journ. Linn. Soc. iii. (1859) 146 = Albizzia basaltica.
Baueri, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 344.—Austral.
Baxteri, *Benth. l. c.* 387.—Austral.

ACACIA :—

Baxteri, Meisn. in Lehm. Pl. Preiss. i. 7 = congesta.
bella, Mart. ex Spreng. Syst. iii. 141 = Calliandra bella.
Benthamii, Meisn. in Lehm. Pl. Preiss. 11 = cochlearis.
Berlandieri, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 592.—Mexic.
Berneri, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 868. —Madag.
Berteriana, Balb. ex DC. Prod. ii. 470 = Pithecolobium fragrans.
Berteriana, *Spreng. Syst.* iii. 138.—Jamaica.
biceps, Willd. Sp. Pl. iv. 1075 = Leucaena glauca.
bidentata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 332.—Austral.
Bidwilli, *Benth. in Linnaea*, xxvi. (1853–55) 629.— Austral.
biflora, *R. Br. in Ait. Hort. Kew.* ed. II. v. 463.— Austral.
biflora, Pazi. Mag. ix. (1842) 221 = decipiens.
biglandulosa, Meisn. in Lehm. Pl. Preiss. ii. 205 = pentadenia.
Bilauna, Buch.-Ham. ex Wall. Cat. sub n. 5264 = odoratissima.
bimucronata, DC. Prod. ii. 469 = Mimosa sepiaria.
binervata, *DC. l. c.* 462.—Austral.
binervosa, DC. Mém. Lég. xii. 448 = bivenosa.
bivenosa, *DC. Prod.* ii. 465.—Austral.
Blomei, Ohlend. in Neue Allgem. Gartenz. (1845) 369 = saligna.
Boivini, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 868.— Madag.
bombycina, Benth. ex Paxt. Flow. Gard. ii. (1851–52). 101 = brachybotrya.
bonariensis, *Gill. in Hook. Bot. Misc.* iii. (1833) 207. —Reg. Argent.
Bonplandii, Gill. ex Hook. l. c. 205 = Mimosa Bonplandii.
bossiaoides, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 332.—Austral.
bossiaoides, Seem. Engel. Acac. t. 1 = glaucoptera.
botrycephala, Desf. Cat. Hort. Par. ed. III. 300 = discolor.
brachyacantha, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 1065 = Mimosa acanthocarpa.
brachybotrya, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 347.—Austral.
brachyloba, Willd. Sp. Pl. iv. 1071 = Desmanthus brachyloma.
brachyphylla, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853–55) 615.—Austral.
brachystachya, Benth. in Hook. Lond. Journ. Bot. i. (1842) 325 = incurva.
brachystachya, *Benth. Fl. Austral.* ii. 408.—Austral.
brasiliensis, *Spreng. Syst.* iii. 142.—Bras.
brevifolia, *Benth. Fl. Austral.* ii. 395.—Austral.
brevifolia, Lodd. Bot. Cab. t. 1235 = lunata.
brevipes, A. Cunn. in Bot. Mag. t. 358 = melanoxylon.
Browniana, Wendl. in Flora, ii. (1819) 139 = strigosa.
Brownii, Steud. Nom. ed. I. 2 = juniperina.
brunioides, *A. Cunn. ex G. Don, Gen. Syst.* ii. 404.— Austral.
Buchananiana, Voigt, Hort. Suburb. Calc. 293 = pinnata.
Burkei, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 86.—Afr. austr.
Burtiitii, *F. Muell. ex Benth. Fl. Austral.* ii. 400.— Austral.
Burmanniana, *DC. Prod.* ii. 461.—Ind. or.
buxifolia, *A. Cunn. in Field, N. S. Wales*, 344.—Austral.
Byneaeana, *Benth. in Linnaea*, xxvi. (1853–55) 614.— Austral.
caduca, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1089 = Mimosa caduca.
caesia, Roxb. ex Wall. Cat. n. 5253 A = pennata.
caesia, *Wight & Arn. Prod.* 278.—Ind. or.
caesia, Willd. Sp. Pl. iv. 1090, ex parte = A. Intsia.
caffra, *Willd. l. c.* 1078.—Afr. austr.
calamifolia, Sieber, ex Steud. Nom. ed. II. i. 4 = quadrilateralis.
calamifolia, *Sweet, ex Bot. Reg.* t. 839.—Austral.
Caleyi, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 347 = podalyriaefolia.
calicoma, *Meisn. in Hook. Lond. Journ. Bot.* ii. (1843) 104.—Afr. austr.
calligera, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 141 = Wickhami.
Calliotaenea, Schlecht. in Linnaea, xii. (1838) 568 = Calliandra grandiflora.
callosa, *Bertol. Pl. Nov. Hort. Bon.* i. 9.—Hab. ?

ACACIA :—

callosa, Spreng. Syst. iii. 138 = Piptadenia communis.
calycularis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 379.—Austral.
Campbellii, Arn in Nov. Act. Nat. Cur. xviii. i. (1836) 833 = planifrons.
campylacantha, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 242 = A. Suma.
campylophylla, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853–55) 605.—Austral.
Candolleana, Meisn. in Lehm. Pl. Preiss. ii. 206 = Drummondii.
canescens, R. Grah in Wall. Cat. n. 5256 = pennata.
canescens, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 314 = Mimosa depauperata.
capensis, Burch. Trav. i. 195 = horrida.
capensis, *Colla, in Mem. Acc. Torin.* xxxv. (1831) 175 —Afr. austr.
carracasana, Willd. Sp. Pl. iv. 1068 = Calliandra carassana.
carbonaria, Schlecht. in Linnaea, xii. (1838) 571 = villosa.
cardiophylla, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 345.—Austral.
Carringa, Buch.-Ham. in Wall. Cat. n. 5239 = Leucaena glauca.
castaicides, *Willd. Enum. Hort. Berol.* 1051.—Hab. ? *Catechu*, Brandis, For. Fl. 186 = A. Suma.
Catechu, Wight & Arn. Prod. 272 = A. Suma
Catechu, *Willd Sp. Pl.* iv. 1079.—Ind. or.
catechuoides, Benth. in Hook. Lond. Journ. Bot. i. (1842) 510 = A. Catechu.
caudata, DC. Prod. ii. 456 = Entada polystachya.
Cavenia, Bert. ex Bull. Férussac, xx. (1830) 108 = A. Farnesiana.
Cebil, *Griseb. in Goett. Abh.* xix. (1874) 136.—Reg. Argent.
cedrioides, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853–55) 615.—Austral.
celastrifolia, Benth. in Hook. Lond. Journ. Bot. i. (1842) 349 = myrtifolia.
centrophylla, *DC. Prod.* ii. 465.—Jamaica.
centrota, Spreng. ex DC. l. c. = praec.
Ceratonia, Willd. Sp. Pl. iv. 1091 = Mimosa Ceratonia.
chiococca, Zucc. in Abh. Akad. Muench. ii. (1832) 847 = filicina.
chordophylla, F. Muell. ex Benth. in Linnaea, xxvi (1853–55) 612 = rigens.
choriophylla, *Benth in Hook. Lond. Journ. Bot.* i. (1842) 495.—Ins. Baham.
chrysantha, *DC. Prod.* ii. 471.—Guiana.
chrysocarpa, Meisn. ind. Sem. Hort. Basil. (1842) 1 = spectabilis
chrysocoma, Miq. in Fl. Ind. Bat. i. 1. 6 = tomentosa.
Chundra, Willd. Sp. Pl. iv. 1078 = A. Catechu.
cibaria, *F. Muell. in Melb. Chemist* (July 1882).— Austral.
ciliata, K. Br. in Ait. Hort. Kew. ed. II. v. 465 = strigosa.
ciliata, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 1055.—Am. austr.
cinclinnata, *F. Muell. Fragm.* xi. 85.—Austral.
Cineraria, Willd. Sp. Pl. iv. 1057 = Prosopis spicigera.
cinerascens, Sieber, ex DC. Prod. ii. 454 = glaucescens.
cinerea, Spreng. Syst. iii. 143 = Dichrostachys cinerea.
Claussenii, Benth. in Hook. Lond. Journ. Bot. i. (1842) 514 = denudata.
clavata, Schlecht. in Linnaea, xx. (1847) 662 = montana.
clavigera, *E. Mey. Comm. Pl. Afr. Austr.* 168.—Afr. austr.
Coa, Walp. Ann. iv. 626 = Koa.
cochlearis, *Wendl. Comm. Acac.* 15.—Austral.
cochliacantha, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1081.—Am. austr.
cochliocarpa, *G. Don, Gen. Syst.* ii. 422.—Bras.
cochlocarpa, *Meisn. in Benth. Lond. Journ. Bot.* i. (1842) 514 = denudata.
clavata, Schlecht. in Linnaea, xx. (1847) 662 = montana.
clavigera, *E. Mey. Comm. Pl. Afr. Austr.* 168.—Afr. austr.
Coa, Walp. Ann. iv. 626 = Koa.
cochlearis, *Wendl. Comm. Acac.* 15.—Austral.
cochli

ACACIA :—

davisioides, A. Cunn. ex G. Don, Gen. Syst. ii. 402 = diffusa.

dealbata, A. Cunn. in Field, N. S. Wales, 345 = lunata.

dealbata, *Link, Enum. Hort. Berol.* 445.—Austral.

Decaisnea, Steud. Nom. ed. II. i. 4 = arcuata.

decipiens, *R. Br. in Ait. Hort. Kew.* ed. II. v. 468.—Ind. or. : Austral.

decora, *Reichb. Ic. Exot.* t. 199.—Austral.

decurrens, *Willd. Sp. Pl.* iv. 1073.—Austral.

decussata, Tenore, Cat. Ort. Nap. (1845) 77 = longifolia.

delibrata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 374.—Austral.

deltoidea, *A. Cunn. ex G. Don, Gen. Syst.* ii. 401.—Austral.

demostachya, Benth. Pl. Hartw. 12 = scapulcensis.

Dempsteri, *F. Muell. Fragm.* xi. 55.—Austral.

densa, Wall. Cat. n. 5202 = leucophloea.

densifolia, Benth. in Hook. Lond. Journ. Bot. i. (1842) 360 = aspera.

denticulosa, *F. Muell. Fragm.* x. 32.—Austral.

dentifera, *Benth. in Maund, Botanist*, iv. t. 179.—Austral.

denudata, Lehm. Del. Sem. Hort. Hamb. (1842) = pulchella.

depauperata, Mart. ex Steud. Nom. ed. II. i. 4 = Desmanthus virgatus.

dependens, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 372 = longifolia.

dermatophylla, *Benth. Fl. Austral.* ii. 346.—Austral.

dermatoxylon, Hartw. Pl. Hartw. 13 = Lysiloma acapulcensis.

detinens, *Burch. Trav.* i. 310.—Afr. austr.

dictyocarpa, Benth. in Linnæa, xxvi. (1853–55) 616 = brachybotrya.

dictyophleba, *F. Muell. Fragm.* iii. 128.—Austral.

Dietrichiana, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 140.—Austral.

diffusa, *Lindl. Bot. Reg.* t. 634.—Austral.

dilatata, *Benth. in Linnæa*, xxvi. (1853–55) 608.—Austral.

dilatata, *Steud. Nom.* ed. II. i. 4.—Austral.

dimidiata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 361.—Austral.

dimorpha, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 130 = binervata.

diptera, *Lindl. Swan River App.* 15.—Austral.

diptera, Willd. Enum. Hort. Berol. 1051 = Prosopis juliflora.

discolor, Whitley, ex Steud. Nom. ed. II. i 4 = prismatica.

discolor, *Willd. Sp. Pl.* iv. 1068.—Austral.

disstiflora, Benth. in Hook. Lond. Journ. Bot. i. (1842) 371 = longifolia.

dissoneura, *F. Muell. in Wing, South Sc. Record* (July 1883).—Austral.

distachya, Moç. & Sesse, ex DC. Prod. ii. 456 = Mimosa distachya.

divaricata, A. Cunn. ex Steud. Nom. ed. II. i. 4 = dilatata.

divaricata, Willd. Sp. Pl. iv. 1070 = Lysiloma Schiedeana.

divergens, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 351.—Austral.

diversifolia, Schlecht. in Linnæa, xii. (1838) 571 = Leucæna diversifolia.

dodonæifolia, *Willd. Enum. Hort. Berol. Suppl.* 68.—Austral.

dolabriformis, Colla, Hort. Ripul. 1 = decipiens.

dolabriformis, A. Cunn. in Hook. Ic. Pl. t. 169 = dimidiata.

dolabriformis, Wendl. Acac. 56 = Daviesia incrassata.

domingensis, Bert. ex DC. Prod. ii. 464 = Mimosa domingensis.

doratoxylon, *A. Cunn. in Field, N. S. Wales*, 345.—Austral.

doratoxylon, Meisn. in Lehm. Pl. Preiss. i. 19 = acuminata.

drepanocarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 137.—Austral.

Drummondii, *Benth. in Lindl. Swan River App.* 15.—Austral.

dumetorum, *DC. Prod.* ii. 458.—Bras.

dumosa, Wight & Arn. Prod. 276 = latronum.

dura, *Benth. in Linnæa*, xxvi. (1853–55) 672.—Austral.

Durandiana, Buckl. in Proc. Acad. Sc. Philad. 1861. (1862) 453 = Greggii.

dyaxophylla, Benth. in Hook. Lond. Journ. Bot. i. (1842) 346 = undulæfolia.

eburnea, *Willd. Sp. Pl.* iv. 1081.—Orient ; Ind. or.

ACACIA :—

echinula, DC. Prod. ii. 449 = juniperina.

Edgeworthii, *T. Anders. in Journ. Linn. Soc.* v. (1860) *Suppl.* 18.—Ind. or.

edulis, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 1066 = A. Farnesiana.

eglandulosa, DC. Mém. Lég. xii. 445 = cochlearis.

Ehrenbergiana, *Hayne, Arzneigew.* x. t. 29.—Afr. trop.

Ehrenbergii, T. Nees, Pl. Offic. t. 284 = præc.

Ehrenbergii, *Steud. Nom.* ed. II. i. 5.—Mexic.

elata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 358.—Austral.

elata, R. Grah. in Wall. Cat. n. 5233 = Albizzia procera.

elegans, Schlecht. in Linnæa, xii. (1838) 569 = filicina.

elephantina, Burch. Trav. ii. 236 = Elephantorrhiza Burchelli.

Elephantorrhiza, Burch. ex DC. Prod. ii. 457 = præc.

elliptica, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 347 = bivenosa.

elongata, *Lindley, ex DC. Prod.* ii. 451.—Austral.

emarginata, Wendl. Comm. Acac. 27 = stricta.

Emoryana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 522.—Am. bor.

Endlicheri, Meisn. ex Lehm. Pl. Preiss. i. 21 = strigosa.

ephedroides, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 370.—Austral.

ericæfolia, *Benth. l. c.* v. (1846) 98.—Afr. austr.

eriantha, *Desv. Journ. Bot.* iii. (1814) 68.—Ind. or. ?

ericæfolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 345.—Austral.

erinacea, *Benth. l. c.* 360.—Austral.

erioclada, *Benth. in Linnæa*, xxvi. (1853–55) 606.—Austral.

erioloba, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1215 = Edgeworthii.

eriolobæ, E. Mey. Comm. Pl. Afr. Austr. i. 171 = giraffæ.

eriophylla, Hort. ex Loud. Encyc. Fl. Suppl. ii. 1582 = viscidula.

erubescens, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 343.—Afr. trop.

erythrocalyx, Steud. ex A. Rich. Tent. Fl. Abyss. i. 243 = A. Suma.

erythrocephala, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 362 = aspera.

erythropus, Tenore, Ind. Sem. Hort. Nap. (1842) 12 = Riceana.

esculenta, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 114 = Leucæna pulverulenta.

esculenta, Moç. & Sesse, ex DC. Prod. ii. 470 = Leucæna esculenta.

Esterhazii, *Mackay, ex Regel, Cat. Pl. Hort. Absakov.* 2.—Austral.

estrophiolata, *F. Muell. in Wing, South Sc. Record* (July 1883).—Austral.

ethalia, *Schweinf. in Linnæa*, xxxv. (1867–68) 330. t. 7, 8.—Afr. trop.

excelsa, *Benth. in Mitch. Trop. Austral.* 295.—Austral.

explanata, Jungh. Reise, 260 ; ex Miq. Fl. Ind. Bat. i. 9 = heterophlœa.

exsudans, Lindl. in Mitch. Three Exped. ii. 214 = verniciflua.

extensa, *Lindl. Swan River App.* 15.—Austral.

fagonioides, Benth. in Hook. Lond. Journ. Bot. i. (1842) 357 = pulchella.

falcata, *Desf.* Tabl. ed. II. 207 = Prosopis juliflora.

falcata, Sieber, ex Steud. Nom. ed. II. i. 5 = plagiophylla.

falcata, *Willd. Sp. Pl.* iv. 1058.—Austral.

falcinella, Hoffmgg. Verz. Pfl. Nachtr. 125, 201 = procumbens.

falciformis, DC. Prod. ii. 462 = penninervia.

falciformis, Meisn. in Bot. Zeit. xiii. (1855) 11 = pycnantha.

falcinella, *Tausch, in Flora*, xix. (1836) 419.—Austral.

fallax, E. Mey. Comm. Pl. Afr. Austr. i. 169 = caffra.

farinosa, *Lindl. in Mitch. Three Exped.* ii. 146.—Austral.

Farnesiana, Wall. Cat. n. 5234 = pluriflora.

Farnesiana, *Willd. Sp. Pl.* iv. 1085.—Reg. trop.

fasciculata, R. Br. in Salt, Abyss. App. 65 = A. Lahai.

fasciculata, Guill. & Perr. Fl. Seneg. Tent. i. 253 = torulifera.

fasciculata, Kunth, Mim. 75. t. 28 = Mimosa fasciculata.

fasciculifera, *F. Muell. ex Benth. Fl. Austral.* ii. 341.—Austral.

ferox, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 97.—Afr. austr.

ferox, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 113.—Mexic.

ferruginea, *DC. Prod.* ii. 458.—Ind. or.

ACACIA :—

filifolia, *Willd. Sp. Pl.* iv. 1079.—Mexic.

filicina, Mart. in Flora, xx. (1837) Beibl. ii. 109 = Piptadenia nitida.

filifolia, Benth. in Hook. Lond. Journ. Bot. i. (1842) 339 = ephedroides.

fimbriata, A. Cunn. ex G. Don, Gen. Syst. ii. 406 = prominens.

Fistula, Schweinf. in Linnæa, xxxv. (1867–68) 344 = A. Seyal.

flaccida, *Spreng. Syst.* iii. 137.—Bras.

flava, Spreng. ex DC. Prod. ii. 469 = Piptadenia flava.

flavescens, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 381.—Austral.

flexicaulis, *Benth. l. c.* 505.—Texas.

flexifolia, *A. Cunn. ex Benth. l. c.* i. 359.—Austral.

flexuosa, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1082 = macracantha.

flexuosa, Lag. Gen. et Sp. Nov. 16 = Prosopis juliflora.

floribunda, Willd. Sp. Pl. iv. 1061 = longifolia.

fluviatilis, Spreng. Syst. v. Ind. 3 = polyphylla.

foetida, H. B. & K. Nov. Gen. et Sp. vi. 365 = Piptadenia foetida.

foliolosa, R. Grah. in Wall. Cat. n. 5241 = Albizzia myriophylla.

formosa, Kunth, Mim. 102. t. 33 = Calliandra formosa.

fragrans, Tenore, Cat. Ort. Nap. 1 = Calliandra caracasana.

Fraseri, Hook. Ic. Pl. t. 171 = podalyriæfolia.

frondosa, Willd. Sp. Pl. iv. 1074 = Leucæna glauca.

fruticosa, *Steud. Nom.* ed. II. i. 5.—Ind. or.

fruticosa, Lond. Hort. Brit. 409 = frutescens.

frutescens, Mart. Herb. Fl. Bras. 107 = Piptadenia latifolia.

frutescens, Voigt, Hort. Suburb. Calc. 261 = Mimosa fruticosa, *Roxb.*

fulgens, Charp. ex Fourn. in Bull. Soc. Bot. Fr. xix. (1865) 400 = Albizzia Charpentieri.

fulgens, Labill. Sert. Austr. Caled. 68. t. 67 = Albizzia fulgens.

furcata, Desv. Journ. Bot. iii. (1814) 67 = Prosopis juliflora.

furcata, *Gill. in Hook. Bot. Misc.* iii. (1838) 206.—Am. austr.

furcifera, Lindl. in Mitch. Three Exped. ii. 267 = armata.

furfuracea, G. Don, Gen. Syst. ii. 405 = lunata.

galioides, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 344.—Austral.

gemella, *Hoffmgg. Vers. Pfl. Nachtr.* ii. 40 ; iii. 16.—Hab. ?

genistifolia, Link, Enum. Hort. Berol. ii. 442 = diffusa.

genistoides, *A. Cunn. ex Benth. Fl. Austral.* ii. 330.—Austral.

Gerrardi, *Benth. in Trans. Linn. Soc.* xxx. (1875) 508. —Afr. austr.

gigantea, Noronha, in Verh. Batav. Gen. v. (1790) Art. iv. 7 = Farsia speciosa.

Gilberti, *Meisn. in Lehm. Pl. Preiss.* ii. 204.—Austral.

Gilesiana, *F. Muell. in Melb. Chemist* (July 1883).—Austral.

Gillisii, Steud. Nom. ed. II. i. 5 = farcata.

Giraffæ, Sieber, ex Steud. l. c. = A. Seyal.

giraffæ, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 236 = stenocarpa.

giraffæ, *Willd. Enum. Hort. Berol.* 1054.—Afr. austr.

glabrata, Schlecht. in Linnæa, xii. (1838) 569 = filicina.

gladiiformis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 354.—Austral.

glandulosa, Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 360 = Leucæna Forsteri.

glandulosa, Willd. Sp. Pl. iv. 1071 = Desmanthus brachylobus.

glauca, Moench, Meth. 466 = Leucæna glauca.

glaucescens, *Willd. Sp. Pl.* iv. 1052.—Austral.

glaucifolia, Baung. ex Lehm. Pl. Preiss. i. 14 in note = calcitrifolia.

glaucophylla, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 243.—Afr. trop.

glaucoptera, *Benth. in Linnæa*, xxvi. (1853–55) 604. —Austral.

glaucorrhoa, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 331.—Am. austr.

glutinosa, F. Muell. Fragm. iv. 6 = bidentata.

Goldiana, *Benth. Fl. Austral.* ii. 329.—Austral.

gonoclinoides, Mart. in Flora, xx. (1837) Beibl. ii. 109 = Piptadenia communis.

ACACIA:—
gonocarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 136.—Austral.
gonoclada, *F. Muell. l. c.* 140.—Austral.
gonophylla, *Benth. in Linnaea*, xxvi. (1853–55) 618. —Austral.
gracilis, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 311 = Calliandra formosa.
graminea, Lehm. Del. Sem. Hort. Hamb. (1842) = extensa.
grandiflora, Willd. Sp. Pl. iv. 1074 = Calliandra grandiflora.
grandis, Henfr. in Gard. Mag. Bot. iii. (1851) 177 = pulchella.
grandistipula, Benth. in Hook. Lond. Journ. Bot. i. (1842) 518 = paniculata.
grandistipula, *Benth. l. c.* 511.—Bras.
granulosa, Labill. Fl. N. Caled. 67. t. 66 = Albizzia granulosa.
grata, Willd. Enum. Hort. Berol. 1056 = Piptadenia macrocarpa.
gravesteni, A. Cunn. ex G. Don, Gen. Syst. ii. 404 = verniciflua.
Greggii, *A. Gray, Pl. Wright.* i. 65.—N. Mexico.
Gregorii, *F. Muell. Fragm.* iii. 47.—Austral.
Grevenna, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 864. —Madag.
Guachapele, *H. B. & K. Nov. Gen. et Sp.* vi. 281 = Lysiloma Guachapele.
guadalupensis, DC. Prod. ii. 464 = riparia.
guatemalensis, Stend. Nom. ed. II. 4. 5 = Albizzia stipulata.
guatemalensis, Hook. & Arn. Bot. Beech. Voy. 419 = Mimosa guatemalensis.
guayaquilensis, Desf. Tabl Hort. Par. ed. II. 208 = Mimosa guayaquilensis.
guianensis, Willd. Sp. Pl. iv. 1061 = Stryphnodendron guianense.
Guilandinæ, DC. Prod. ii. 465 = Mimosa obovata.
gummifera, Delile, Fl. Aeg. Illustr. 79 = spirocarpa.
gummifera, *Willd. Sp. Pl.* iv. 1056.—Afr. bor.
Gunnii, Benth. in Hook. Lond. Journ. Bot. i. (1842) 332 = vomeriformis.
guttata, *Hoffmgg. Verz. Pfl. Nachtr.* i. 201, 125.— Hab.?
guttulifera, *Hoffmgg. l. c.* iii. 16 = Martinicna.
gyrocarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 241 = albida.
Habbas, Link, Enum. Hort. Berol. ii. 444 = Mimosa asperа!
habbasioides, Boj. Hort. Maurit. 116 = concinna.
hadiensis, *DC. Prod.* ii. 472.—Arab.
haematomma, Bert. ex DC. Mém. Lég. xii. 448. t. 68 = Calliandra haematomma.
Haematoxylon, *Willd. Enum. Hort. Berol.* 1056.— Afr. austr.
hakeoides, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 354.—Austral.
Hamiltoni, Desv. in Ham. Prod. Fl. Ind. Occ. 59 = Calliandra portoricensis.
hamulosa, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 509.—Arab.
harpophylla, *F. Muell. ex Benth. Fl. Austral.* ii. 389. —Austral.
Hartwegi, Benth. Pl. Hartw. 13 = filicina.
Harveyi, *Benth. Fl. Austral.* ii. 368.—Austral.
hastulata, *Sm. in Rees, Cyclop.* xxxix. *Suppl.*— Austral.
Hayesii, *Benth. in Trans. Linn. Soc. xxx.* (1875) 524.— Am. trop.
hebecephala, A. Cunn. ex G. Don, Gen. Syst. ii. 403 = elongata.
hebeclada, *DC. Cat. Hort. Monsp.* 78; *Prod.* ii. 461.— Afr. austr.
hcoxophylla, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 243.—Afr. trop.
hemignostа, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 134.—Am. trop.
hemiteles, Benth. in Linnaea, xxvi. (1853–55) 619 = subnervulea.
Henne, Walp. Rep. v. 606 (sphalm. pro *A. Nemu*).
heteracantha, *Burch. Trav.* i. 398.—Afr. austr.
heterocarpa, Delile, Fl. Aeg. Illustr. 79 = Prosopis Stephaniana.
heterocilia, *Meisen. in Lehm. Pl. Preiss.* i. 18.— Austral.
heterocraera, *Benth. in Linnaea*, xxvi. (1853–55) 624. —Austral.

ACACIA:—
heterophylla, Hook. & Arn. Bot. Beech. Voy. 81 = A. Kea.
heterophylla, *Willd. Sp. Pl.* iv. 1054.—Ins. Mascar.
Hindsii, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 504.—Mexico.
hippuroides, *Steward, ex Benth. l. c.* 344.—Austral.
hirsuta, Schlecht. in Linnaea, xii. (1838) 572 = filicina.
hirta, *Boj. Hort. Maurit.* 115.—Ins. Maurit.
hirta, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 318 = Mimosa Galeottii.
hirta, Nutt. in Torr. & Gray, Fl. N. Am. i. 404 = filicina.
hirtella, *E. Mey. Comm. Pl. Afr. Austr.* 167.—Afr. austr.
hispidissima, DC. Prod. ii. 465 = pulchella.
hispidula, *Willd. Sp. Pl.* iv. 1054.—Austral.
Hoffmannseggii, *DC. Prod.* ii. 471.—Bras.
holocarpa, *Benth. Fl. Austral.* ii. 408.—Austral.
holosericea, *A. Cunn. ex G. Don, Gen. Syst.* ii. 407.— Austral.
homalocladn, *F. Muell. Fragm.* xi. 34.—Austral.
homalophylla, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 353.—Austral.
homomalla, Wendl. Acac. 49. t. 18 = glaucescens.
Hookeri, Meisn. in Lehm. Pl. Preiss. i. 11 = ericifolia.
Hookeriana, Zipp. ex Miq. Fl. Ind. Bat. i. 10 = concinna.
horrida, Spun. in Linnaea, xv. (1841) 199 = pennata.
horrida, *Willd. Sp. Pl.* iv. 1082.—Afr. austr.
horridula, *Meisen. in Lehm. Pl. Preiss.* i. 9.—Austral.
hostilis, Mart. Reise, i. 556 = Mimosa hostilis.
Houstoni, Willd. Sp. Pl. iv. 1062 = Calliandra Houstoni.
Huegelii, *Benth. in Enum. Pl. Hueg.* 42.—Austral.
Humboldtii, Desv. in Ham. Prod. Fl. Ind. Occ. 60 = macracantha.
humifusa, *A. Cunn. ex Benth. in Hook Lond. Journ. Bot.* i. (1842) 382.—Austral.
humilis, Schlecht. in Linnaea, xii. (1838) 567 = Calliandra humilis.
Hunteri, *Oliver, in Hook. Ic. Pl* t. 1850.—Arab.
hybrida, Lodd. Bot. Cab. t. 1842 = armata.
hypoglauca, *Hoffmgg. ex Heynh. Nomencl.* i. 8, nomen.—Afr. austr.
idiomorpha, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 325.—Austral.
ignorata, *Koch, in Allg. Garte.* (1858) 195.—Hab.?
imbricata, F. Muell. Fragm. i. 65 = lineata.
implexa, *Benth in Hook. Lond. Journ. Bot.* i. (1842) 505.—Austral.
impressa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 133.—Austral.
impressa, Lindl. in Bot. Reg. t. 1115 = penninervia.
incana, Spreng. Syst. iii. 127 = Mimosa incana.
incrassata, Hook. Ic. Pl. t. 870 = decipiens.
incurva, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 332.—Austral.
indica, Desv. Journ. Bot. i. (1814) 69 = Farnesiana.
ingrata, *Benth. Fl. Austral.* ii. 391.—Austral.
insignis, Hoffmgg. Verz. Pfl. 159 = Albizzia lophantha.
insignis, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 313 = filicina.
insularis, *A. Rich. Ess. Fl. Cub.* 464.—Ind. occ.
insulana, *Guill. in Ann. Sc. Nat.* Sér. II. vii. (1837) 360 = Leucaena Forsteri.
intermedia, A. Cunn. ex Hook. Bot. Mag. t. 3208 = longifolia.
intermedia, *Pasq. in Atti R. Acad. Nap.* ix. (1882) 7. —Hab.?
intertexta, Sieber, ex DC. Prod. ii. 464 = longifolia.
Intsia, Willd. Sp. Pl. iv. 1091 = caesia.
intsioides, DC. Prod. ii. 464 = praec.
inundata, Mart. Reise, i. 555 = Pithecolobium multiflorum.
inversata, Sieber, ex Spreng. Syst. iii. 141 = dealbata.
Isambergiana, Schimp. ex A. Rich. Tent. Fl. Abyss. i. 236 = Albizzia Julibrissin.
itaspkylla, F. Muell. ex Benth. in Linnaea, xxvi. (1853–55) 617 = neriifolia.
ixiophylla, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 354.—Austral.
Jacquemontii, *Benth. l. c.* 492.—Ind. or.
Jambertiana, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 863.—Madag.
javanica, *DC. Prod.* ii. 471.—Java.
Julibrissin, Willd. Sp. Pl. iv. 1065 = Albizzia Julibrissin.

ACACIA:—
Julibrissin, Sieber, ex Benth. in Trans. Linn. Soc. xxx. (1875) 529 = paniculata.
julifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 374.—Austral.
julifera, F. Muell. in Journ. Linn. Soc. iii. (1859) 139 = torulosa.
julifora, Willd. Sp. Pl. iv. 1076 = Prosopis juliflora.
juncifolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 341.—Austral.
juniperina, *Willd. Sp. Pl.* iv. 1049.—Austral.
Jupunba, Willd. l. c. 1087 = Pithecolobium trapezifolium.
Jurema, *Mart. Reise*, ii. 788; *Syst. Mat. Med. Fl. Bras.* 53.—Bras.
Kalkora, G. Don, in Lond. Hort. Brit. 409 = Albizzia Julibrissin.
Kantuffa, Bruce, ex Steud. Nom. ed. I. 2 = Mimosa Kantuffa.
Karroo, Hayne, Arzneigew. x. t. 38 = horrida.
Kempeana, *F. Muell. in Melb. Chemist* (July, 1882). —Austral.
Kirkii, *Oliver, Fl. Trop. Afr.* ii. 350.—Afr. trop.
Koa, *A. Gray, Bot. U. S. Expl. Exped.* i. 460.—Ins. Sandvic.
Kraussiana, *Meisen. ex Benth. in Hook. Lond. Journ Bot.* i. (1842) 515.—Afr. austr.
Kunthii, *Dekuk. Rivist. Nap.* i. 169; *ex Walp. Rep.* i 924.—N. Granat.
lacerans, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 511.—Bras.
lachnophylla, *F. Muell. in Wing, South. Sc. Record* (July, 1882).—Austral.
lacustris, Desf. Cat. Hort. Par. ed. III. 301 = Neptunia oleracea.
laeta, *R. Br. in Salt, Abyss. App.* 65.—Afr. bor.; Arab.
laevigata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1059 = Prosopis juliflora.
laevis, Hort. ex Steud. Nom. ed. II. i. 6 = Gleditschia triacanthos.
Lahai, *Steud. & Hochst. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 506.—Abyss.
Lambertiana, D. Don, in Bot. Reg. t. 721 = Calliandra Lambertiana.
lanata, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 315.—Mexico.
Langsdorffi, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 521.—Bras.
lanigera, *A. Cunn. in Field, N. S. Wales*, 345.— Austral.
lanuginosa, Hort. ex Regel, in Gartenfl. (1854) 155 = pulchella!
laricina, *Meisen. in Lehm. Pl. Preiss.* i. 6.—Austral.
lasiocarpa, Benth. in Enum. Pl. Hueg. 43 = pulchella.
lasiopetala, *Oliver, Fl. Trop. Afr.* ii. 346.—Afr. trop.
lasiophylla, *Benth. in Trans. Linn. Soc.* xxx. (1875) 527.—Am. trop.
latescens, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 386.—Austral.
latifolia, *Benth. l. c.* 382.—Austral.
latifolia, Desf. Tabl. Hort. Par. ed. II. 207 = A. melanoxylon.
latifrons, Hort. ex Scem. & Schmidt, in Flora, xxvii. (1844) 499 = elata.
latipes, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 334.—Austral.
latisiliqua, Willd. Sp. Pl. iv. 1067 = Lysiloma latisiliqua.
latispinosa, Desf. Cat. Hort. Par. ed. III. 299 = Mimosa latispinosa.
Latrobei, Meisen. in Lehm. Pl. Preiss. i. 10 = acinacea.
latronum, *Willd. Sp. Pl.* iv. 1077.—Ind. or.
laurifolia, *Willd. l. c.* 1058.—Ins. Pacif.
Lawsoni, *Ait. ex Steud. Nom. ed.* II. i. 6.—Austral.
laxa, *Willd. Sp. Pl.* iv. 1069 = Calliandra laxa.
laxiflora, DC. Prod. ii. 455 = Pithecolobium laxiflorum.
laxophylla, vide *laxophylla*.
Lebbek, Willd. Sp. Pl. iv. 1066 = Albizzia Lebbek.
lebbekoides, DC. Prod. ii. 467 = Albizzia lebbekoides.
Leichardtii, *Benth. Fl. Australiensis*. ii. 378.—Austral.
leiophylla, Benth. in Hook. Lond. Journ. Bot. i. (1842) 351 = Albizzia saligna.
leptiorllata, F. Muell. in Journ. Linn. Soc. iii. (1859) 147 = Farnesiana.
lenticularis, *Buch.-Ham. in Wall. Cat.* n. 5244.— Ind. or.

ACACIA:—
lenticifolia, *Desf. Tabl. Hort. Par.* ed. II. 208.—Mexico.
lenticifolia, *A. Rich. Ess. Fl. Cub.* 469.—Cuba.
lepidota, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 208 = Mimosa Bonplandii.
leprosa, *Sieber, ex DC. Prod.* ii. 450.—Austral.
leptocarpa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 376.—Austral.
leptocladia, *A. Cunn. ex Benth. l. c.* 385.—Austral.
leptonevra, *Benth. l. c.* 341.—Austral.
leptopetala, *Benth. in Linnæa*, xxvi. (1853–55) 619.—Austral.
leptophleba, F. Muell. in Journ. Linn. Soc. iii. (1859) 143 = salacocarpa.
leptophylla, DC. Cat. Hort. Monsp. 74 = Farnesiana.
leptophylla, F. Muell. Fragm. xix. 9 = A. Byneana.
leptosperma, *Sello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 265.—Ind. occ.
leptospermoides, *Benth. in Linnæa*, xxvi. (1853–55) 626.—Austral.
leptostachya, *Benth. Fl. Austral.* ii. 406.—Austral.
leucacantha, Bert ex Spreng Syst. iii. 144 = tortuosa.
leucacantha, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 376.—Afr. trop.
leucadendron, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 374 = glaucescens.
leucocephala, Link, Enum. Hort. Berol. ii. 444 = Leucæna glauca.
leucolobia, *Sweet, Hort. Brit.* ed. II. 166.—Austral.
leucophloea, *Willd. Sp. Pl.* iv. 1088.—As. trop.
leucophylla, Colvill, ex Sweet, Hort. Brit. ed. I. 101 = holosericea.
leucophylla, Lindl. in Mitch. Three Exped. ii. 18 = pendula.
ligulata, *Ait. ex Steud. Nom.* ed. II. i. 6.—Austral. —Cf. seq.
ligulata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 382 = salicina.
ligustrina, *Meisen. in Lehm. Pl. Preiss.* ii. 208.—Austral.
limbata, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 145.—Austral.
linarioides, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 371.—Austral.
Lindleyi, *Meisen. in Lehm. Pl. Preiss.* i. 14.—Austral.
linearis, Desv. in Ham. Prod. Fl. Ind. Occ. 591 = Calliadra portoricensis.
linearis, *Sims, Bot. Mag.* t. 2156.—Austral.
linearis, *A. Cunn. ex G. Don, Gen. Syst.* ii. 408.—Austral.
linifolia, *Benth. ex Schlecht. in Linnæa*, xxvi. (1853–55) 626.—Austral.
lingvatousana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 862.—Madag.
linifolia, *Willd. Sp. Pl.* iv. 1051.—Austral.
litakunensis, *Burch. Trav.* ii. 452.—Afr. austr.
littoralis, *A. Rich. Ess. Fl. Cub.* 467.—Cuba.
lomatocarpa, DC. Prod. ii. 467 = Albizzia stipulata.
longifolia, Pxzt. Mag. Bot. xii. (1846) 269 = dentifera.
longifolia, *Willd. Sp. Pl.* iv. 1052.—Austral.
longispicata, Benth. in Mitch. Journ. Trop. Austr. 298 = Cunninghamii.
longissima, Wendl. Comm. Acac. 45. t. 11 = linearis.
lophantha, Willd. Sp. Pl. iv. 1070 = Albizzia lophantha.
lophanthoides, DC. Prod. ii. 457 = villosa.
loxocarpa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 377.—Austral.
lozophylla, *Benth. in Linnæa*, xxvi. (1853–55) 622.—Austral.
lucana, *Boj. Hort. Maurit.* 116.—Madag.
lucida, Baill. Hist. Pl. ii. 61 = Albizzia lucida.
Ludwigii, Ohlend, Verz. (1844) 74; Walp. Ann. ii. 458 = myrtifolia.
Luehmanni, *F. Muell. Fragm.* xi. 216.—Austral.
lunata, Hort. Prag. ex Steud. Nom. ed. II. i. 6 = falcinella.
lunata, Lodd. Bot. Cab. t. 824 = myrtifolia.
lunata, *Sieber, ex DC. Prod.* ii. 452.—Austral.
lutea, Leaveuw. in Am. Journ. Sc. vii. (1824) 61 = Neptunia lutea.
lycopodiifolia, *A. Cunn. in Hook. Ic. Pl.* t. 172.—Austral.
lycopodioides, Desv. Journ. Bot. iii. (1814) 69 = Neptunia pinna.
lysiphlœa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 137.—Austral.

ACACIA:—
macocentra, Walp. Rep. i. 876. in syn.—sphalm.—Cf. malacocentra.
macracantha, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1080.—Am. austr.
macracanthoides, Bert. in DC. Prod. ii. 468 = macracantha.
macradenia, *Benth. in Mitch. Trop. Austral.* 360.—Austral.
macrocentra, Walp. Rep. iv. 200.—Cf. malacocentra.
macroloba, Willd. Sp. Pl. iv. 1054 = Pentaclethra filamentosa.
macrophylla, Bunge, Enum. Pl. Chin. Bor. 20 = Albizzia Lebbek.
macrostachya, *Reichenb. ex G. Don, Gen. Syst.* ii. 411. —Senegal.
Magdalenæ, Bert. ex DC. Prod. ii. 455 = Calliandra Magdalenæ.
Maitlandi, F. Muell. Fragm. iii. 46 - patens.
malacocentra, Mart. Herb Fl. Bras. 106 = Mimosa malacocentra.
malacophylla, *Benth. in A. Gray, Pl. Wright.* i. 64.—Mexico
malacophylla, Steud. ex A. Rich. Tent. Fl. Abyss. i. 235 = Albizzia pallida.
mammifera, *Schlecht. in Linnæa*, xii. (1838) 568.—Mexico.
Mangium, *Willd. Sp. Pl.* iv. 1053.—Ins. Moluc.
marginata, Buch.-Ham. in Wall. Cat. n. 5248 = Albizzia stipulata.
marginata, Steud. Nom. ed. I. 3 = Albizzia odoratissima.
marginata, R. Br. in Ait. Hort Kew. ed. II. iii. 462 = myrtifolia.
maritima, Benth. in Hook. Lond. Journ. Bot. i. (1842) 384 = discolor.
Martii, *Benth. l. c.* 510.—Bras.
martinicensis, Presl, Bot. Bemerk. 65 = paniculata, maschalocephala, *Griseb. Cat Pl. Cub.* 82.—Cuba.
mauroccna, *DC. Cat. Hort. Monsp.* 74.—Marocco.
media, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 316 = Calliandra angelina.
megaladena, Desv. Journ. Bot. iii. (1814) 69 = pennata.
megalantha, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 143.—Austral.
megaphylla, F. Muell. l. c. 117 = aneeps.
Meisneri, *Lehm. Del. Sem. Hort. Hamb.* (1842).—Austral.
melanoxera, *Benrling, in Vet. Akad. Handl. Stockh.* (1854) 53.—Panama.
melanochaetes, Zoll. in Nat. en Geneesk. Arch. Neel. Indie, iii. (1846) 80 = leucophloea.
melanoxylon, *R. Br. in Ait. Hort. Kew.* ed. II. v. 462. —Austral.
mellifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 397.—Abyss.
metrondorifiora, Schlecht. in Linnæa, xii. (1838) 567. = Calliandra Houstoni.
metrondorifolia, Hort. ex Steud. Nom. ed. II. i. 6 = dodonæifolia.
micracantha, A. Dietr. in Otto & Dietr. Allg. Gartenz. i. (1833) 93 = armata.
micracantha, Desv. Journ. Bot. iii. (1814) 69 = Pithecolobium parvifolium.
micradena, Benth. in Hook. Lond. Journ. Bot. i. (1842) 518 = leucocephala.
micrantha, *Benth. in Trans. Linn. Soc.* xxx. (1875) 596.—Mexico.
micrantha, Desv. in Ham. Prod. Fl. Ind. Occ. 59 = seq.
microcantha, G. Don, Gen. Syst. ii. 476 = Pithecolobium parvifolium.
microbotrya, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 353.—Austral.
microcarpa, *F. Muell. 2nd Gen. Report*, 7.—Austral.
microcephala, A. Rich. Ess. Fl. Cub. 469 = paniculata.
microcephala, R. Grah. in Wall. Cat. n. 5263 = leucophloea.
microcephala, Macfad. Fl. Jamaic. i. 216 = macracantha.
microneura, *Meisen. in Lehm. Pl. Preiss.* i. 19.—Austral.
microphylla, Roxb. ex Steud. Nom. ed. II. i. 6 = Albizzia myriophylla.
microphylla, Willd. Sp. Pl. iv. 1078 = Piptadenia peregrina.
Mearsii, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 523.—Bras.

ACACIA:—
Mikanii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 526.—Am. trop.
minutifolia, *F. Muell. Fragm.* viii. 248.—Austral.
Mirbeli, *Dehnh. Rivist. Nap.* i. 168; *ex Walp. Rep.* i. 922.—Austral.
mlsera, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 375.—Afr. trop.
Mitchelli, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 357.—Austral.
modesta, *Wall. Pl. As. Rar.* ii. 27. t. 130.—Ind. or.
monsta, Lindl. Bot. Reg. (1846) t. 67 = verticillata.
mollicula, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 313.—Mexico.
mollis, Wall. Pl. As. Rar. ii. 76. t. 177 = Albizzia Julibrissin.
mollissima, Willd. Enum. Hort. Berol. 1058 = decurrens.
monacantha, *Willd. l. c.* 1056.—Bras.
monliformis, *Grisseb. in Goett. Abh.* xix. (1874) 136.—Reg. Argent.
montana, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 360.—Austral.
montana, Jungh. in Nat. en Geneesk. Arch. Neerl. Indie, ii. (1845) 35 = Albizzia montana.
mozambicensis, Bolle, in Peters, Reise Mossamb. Bot. i. 15 = albida.
mucronata, Willd. Enum. Hort. Berol. Suppl. 68 = longifolia.
Muelleri, Benth. in Linnæa, xxvi. (1853–55) 608 = aneeps.
multiflora, H. B. & K. Nov. Gen. et Sp. vi. 377 = Pithecolobium multiflorum.
multiflora, Spreng. Syst. iii. 142 = polyantha.
multifoliolata, *Bertol. Misc. Bot.* xvi. 26.—Ins. S. Domingo.
multijuga, *Meisen. in Hook. Lond. Journ. Bot.* ii. (1848) 105.—Afr. austr.
multinervia, DC. Mém. Lég. xii. 445; Prod. ii. 250 = lanigera.
multiplicata, *Benth. Fl. Austral.* ii. 400.—Austral.
muricata, *Willd. Sp. Pl.* iv. 1058.—Ind. occ.
Murrayana, *F. Muell. ex Benth. Fl. Austral.* ii. 370.—Austral.
myradenia, Bert. ex Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 259 = Seriantlies myriadena.
myriobotrya, Meisan. in Lehm. Pl. Preiss. i. 15 = microbotrya.
myriophylla, R. Grah. in Wall. Cat. n. 5242 = Albizzia myriophylla.
myrtifolia, *Willd. Sp. Pl.* iv. 1054.—Austral.
natalitia, *E. Mey. Comm. Pl. Afr. Austr.* 167.—Afr. austr.
Nebouei, *Baill. Adansonia*, iv. (1863–64) 117.—Afr. trop.
Nellyrensa, R. Grah. in Wall. Cat. n. 5240 = amara.
nematophylla, F. Muell. ex Benth. in Linnæa, xxvi. (1853–55) 612 = calamifolia.
Nemu, Willd. Sp. Pl. iv. 1065 = Albizzia Julibrisin.
nerifolia, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 367.—Austral.
Nernstii, F. Muell. Fragm. iv. 3 = amblygona.
nervosa, *DC. Mém. Lég.* xii. 444.—Austral.
Neuaana, Buckl. ex Benth. in Trans. Linn. Soc. xxx. (1875) 592 sphalm. = Pithecolobium brevifolium.
neurocarpa, A. Cunn. in Hook. Ic. Pl. t. 168 = holosericea.
nigra, Cav. in C. Gay, Fl. Chil. ii. 253 = Calliandra chilensis.
nigrescens, *Oliver, Fl. Trop. Afr.* ii. 340.—Afr. trop.
nigricans, *R. Br. in Ait. Hort. Kew.* ed. II. v. 465.—Austral.
nilotica, Delile, Fl. Aegypt. Illustr. 79; Desf. Tabl. Hort. Par. ed. II. 208 = arabica.
Ninpe, H. B. & K. Nov. Gen. et Sp. vi. 382 = Piptadenia peregrina.
nitida, Willd. Sp. Pl. iv. 1086 = Pithecolobium umbellatum.
nitidula, *Benth. Fl. Austral.* ii. 361.—Austral.
nodifora, *Benth. in Linnæa*, xxvi. (1853–55) 621.—Austral.
nodosa, Willd. Sp. Pl. iv. 1016 = Cassia bacillaris.
notabilis, *F. Muell. Fragm.* i. 6.—Austral.
nubica, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 498.—Nubia.
nudiflora, *Willd. Sp. Pl.* iv. 1058.—Ind. occ.
Nuciana, Buckl. in Proc. Ac. Sc. Philad. '1861' (1862) 458 = Pithecolobium brevifolium.
nutans, Spreng. ex DC. Prod. ii. 429 = Mimosa leiocarpa] *nyssophylla*, F. Muell. Fragm. iv. 4 = colletioides.

ACACIA:—
obliqua, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 334.—Austral.
obliqua, Desv. Journ. Bot. iii. (1814) 67 = suaveolens.
obliquifolia, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 317 = Pithecolobium dulce.
obovata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 329.—Austral.
obscura, *A. DC. Not.* vi. *Pl. Rar. Jard. Gen.* 23. t. 3.—Austral.
obtusa, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1087 = macracantha.
obtusata. *Sieber, ex DC. Prod.* ii. 458.—Austral.
obtusifolia, A. Cunn. in Field, New South Wales, 345 = longifolia.
odorata, Desv. Journ. Bot. iii. (1814) 67 = suaveolens.
odoratissima, Buch.-Ham. ex Wall. Cat. n. 5233 = elata.
odoratissima, Willd. Sp. Pl. iv. 1068 = Albizzia odoratissima.
officinalis, *Dehnh. Rivist. Nap.* i. 170; *ex Walp. Rep.* i. 923.—Tasman.
Oldfieldii, F. Muell. Fragm. iv. 17 = acuminata.
oleaefolia, A. Cunn. ex G. Don. Gen. Syst ii. 405 = lunata.
oligoneura, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 139.—Austral.
oligophylla, *Hoffmgg. Verz. Pfl. Nachtr.* i. 201.—Hab.?
Oliveri, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 274.—Afr. trop.
omalophylla = homalophylla.
oncinocarpa, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 378.—Austral.
onchnophylla, *Benth. in Lindl. Swan Riv. App.* 15.—Austral
oraria, *F. Muell. Fragm.* xi. 66.—Austral.
ornithophora, Sweet, Pl. Austral. t. 94 = armata.
orthocarpa, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 136 = xylocarpa.
Oswaldi, *F. Muell. Pl. Vict.* ii. 27.—Austral.
Ouyrarema, *DC. Prod.* ii. 467.—Guiana.
ovoidea, Benth. in Hook. Lond. Journ. Bot. i. (1842) 339 = verticillata.
Oxycedrus, *Sieber, ex DC. Prod.* ii. 458.—Austral.
oxyclada, *F. Muell. ex Benth. Fl. Austral.* ii. 341.—Austral.
oxyphylla, R. Grah. in Wall. Cat. n. 5262 = caesia.
pachycarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 139.—Austral.
pachyloma, Mart. Herb. Fl. Bras. n. 1108 = Calliandra parviflora.
pallida, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 147.—Austral.
pallida, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1059 = Prosopis pallidora.
Palmeri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 350.—Am. bor. occ.
paludosa, Blume, ex Miq. Fl. Ind. Bat. i. 14 = pennata.
paniculiflora, Steud. in Flora, xxvi. ii. (1843) 760 = Mimosa myriadenia.
paniculata, *Willd. Sp. Pl.* iv. 1074.—Bras.
papulaeformis, Walp. Rep. i. 898 = buxifolia.
papyrocarpa, *Benth. Fl. Austral.* ii. 388.—Austral.
paradoxa, DC. Cat. Hort. Monsp. 74 = armata.
parvifolia, Hort. ex Steud. Nom. ed. II. i. 7 = guttata.
parvifolia, Willd. Sp. Pl. iv. 1086 = Pithecolobium parvifolium.
patens, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 120.—Austral.
patula, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 1055.—Am. austr.
pauciflora, *A. Rich. Ess. Fl. Cub.* 461.—Cuba.
Pauliksenyana, Ohlend. in Neue Allg. Gartenz. (1845) 369 = myrtifolia.
pectinata, Hoffmgg. Verz. Pfl. Nachtr. i. 201 = Hoffmannseggiana.
pectinata, H. B. & K. Nov. Gen. et Sp. vi. 382 = Parkia pectinata.
pedicellata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 392.—Bras.
pedunculata, Rozb. ex Steud. Nom. ed. II. i. 7 = Parkia biglandulosa ?
pedunculata, Willd. Sp. Pl. iv. 1084 = Farnesiana.
pellacantha, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 19 = macracantha.
pendula, *A. Cunn. ex G. Don, Gen. Syst.* ii. 404.—Austral.
penicilliferu, Lag. Gen. et Sp. Nov. 16 = filicina.

ACACIA:—
pennata, *Willd Sp. Pl.* iv. 1090.—As. et Afr. trop.
pennatula, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 390.—Mexico.
penninervis, *Sieber, ex DC. Prod.* ii. 452.—Austral.
pennivenia, *Bulf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 404.—Ins. Socotra.
pentadenia, *Lindl. in Bot. Reg.* t. 1521.—Austral.
pentaedra, Regel, Gartenfl. (1852) 238, t. 23 = extensa.
pentagona, Hook. f. in Hook. Nig. Fl. 331 = pennata.
pentaptera, Welw. Apont. 584 = pennata.
peregrina, Kunth, Mimos. Humb. 96. t. 30 = Piptadenia colubrina.
peregrina, Willd. Sp. Pl. iv. 1078 = Piptadenia peregrina.
Perrottetii, Steud. Nom ed. II. i. 7 = tortilis.
Petersiana, Sterl. Hort. Nymph. ex Steud. Nom. ed. II i. 7 = Prosopis Stephaniana.
personioides, *Ait. ex Steud. Nom.* ed. II. i. 7.—Austral.
Pervillei, *Benth. in Trans Linn. Soc.* xxx. (1875) 521.—Madag.
peruviana, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 1056.—Peruv.
Petersiana, *Bolle, in Peters, Reise Mossamb. Bot.* i. 4.—Mossamb.
petiolaris, Lehm. Novit Pl. Hort. Hamb. (1851) 7; Linnaea, xxv. (1852) 308 = pycnantha.
Pexce, *F. Muell. Fragm* iii. 121.—Austral.
Pharnesiana, Risso, Fl. Nice, 166 = Farnesiana.
philippinarum, Benth in Hook. Lond. Journ. Bot. i. (1842) 314 = concinna.
phlebocarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 119.—Austral.
phlebophylla, F. Muell 1st Gen. Rep. 19 = longifolia.
piauhiensis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 523.—Bras.
pilgera, A. Cunn. in Hook Bot. Mag sub t. 3394 = umbellifera.
pilosa, *Benth. in Linnaea*, xxvi. (1853–55) 607.—Austral.
pilosa, Bert. ex DC. Prod. ii. 455 = Calliandra haematomma ?
pimifolia, Benth. in Mitch. Trop. Austral. 342 = juncifolia.
pinetorum, Dals. & Gibs. Bomb. Fl 87 = pennata.
pinnata, Link, Enum. Hort. Berol. ii. 446 = tamarindifolia.
pityoides, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 185.—Austral.
plagiophylla, F. Muell. t. c. 181 = undulaefolia.
plagiophylla, Spreng. Syst. iii. 185 = falcata.
planifrons, *Wight & Arn. Prod.* 276.—Ind. or.
platycantha, *Schlecht. in Linnaea*, xii. (1838) 565.—Mexico.
platycarpa, F. Muell. in Journ. Linn. Soc. iii (1859) 145 = sericata.
platyloba, *Spreng. Syst.* iii. 144 —N. Granat.
platyloba, Bert. ex DC. Prod. ii. 472 = Pithecolobium scutiforum.
platyphylla, *Sweet, Hort. Brit.* ed. I. 102, nomen.—Austral.
platystyla, Lindl. in Bot. Reg. (1841) Misc. 3 = falcata.
plectocarpa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 375.—Austral.
plumosa, *Lowe, in Bot. Mag.* t. 3366.—Bras.
plurijuga, *Steud. Nom.* ed. II. i. 7.—As. trop.
podalyriaefolia, *A. Cunn. in G. Don, Gen. Syst.* ii. 405.—Austral.
polycephala, Willd. Sp. Pl. iv. 1079 = A. Catechu.
polyantha, Spreng ex Benth. in Trans. Linn. Soc. xxx. (1875) 591 = Pithecolobium multiflorum.
polyantha, Zipp. ex Span. in Linnaea, xv. (1841) 199 = juncifolia.
polybotrya, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 384.—Austral.
polycephala, DC. Prod. ii. 478 = concinna.
polycephala, R. Grah. in Wall. Cat. n. 5255 = pluricapitata.
polymorpha, *Pasp. Cat. Ort. Bot. Nap.* 3.—Austral.
polyphylla, Cles, in C. Gay, Fl. Chil. ii. 364 = Lysiloma polyphylla.
polyphylla, *DC. Cat. Hort. Monsp.* 74; *Prod.* ii. 469.—N. Granat.
polystachya, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 376.—Austral.
portoricensis, Willd. Sp. Pl. iv. 1069 = Calliandra portoricensis.
praecox, *Griseb. in Goett. Abh.* xix. 136.—Reg. Argent.

ACACIA:—
praelongata, *F. Muell. in Melb. Chemist*, iii. (Aug. 1888).—Austral.
praefolia, F. Muell. Fragm. i. 4 = sublanata.
praevisima, *F. Muell. 1st Gen. Report*, 12.—Austral.
Preissiana, Lehm. Del. Sem. Hort. Hamb. (1842) = obscura.
prenzana, Lowe, in Bot. Mag. t. 3408 = pennata.
prismatica, *Hoffmgg. Verz. Pfl.* 159.—Hab.?
procera, Willd. Sp. Pl. iv. 1083 = Albizzia procera.
prommens, *A. Cunn. ex G. Don, Gen. Syst.* ii. 406.—Austral.
prona, *Noronha, in Verh. Batav. Gen.* v. (1790) Art. iv 5, nomen.—Java.
propinqua, A. Rich. Ess. Fl. Cub. i. 466 = Albizzia Lebbek.
prosopoides, Moç. & Sesse, ex DC. Prod. ii. 460 = Piptadenia patens.
prosopona, *Schnyder, in Anal. Soc Cient. Argent.* lii. (1877) 159.—Reg. Argent.
prostrata, Lodd. Bot. Cab. t. 681 = diffusa.
prunescens, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 236, 298.—Burma.
pruinosa, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 388.—Austral.
prunifolia, C. Muell. in Walp. Ann. vii. 937, sphalm. —Cf. pravifolia.
pseudoarabica, *Blume, ex Miq. in Fl. Ind. Bat.* i. 8.—N Guinea.
pseudonitida, Miq. l. c. 12 = caesia.
pseudorichodes, *DC. Prod.* ii. 466.—Ins. S. Doming.
psilostachya, DC. l. c. 457 = Piptadenia psilostachya.
Psoralea, DC. l. c. 464 = Mimosa Psoralea.
pteridifolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 321.—Am. trop.
pteriqoidea, Seem. in Verh. Gartenb. (1846) 11 = microbotrya.
pterocarpa, F. Muell. in Journ. Linn. Soc. iii. (1859) 134 = excelsa.
pterocarpa, Steud. Nom. ed. I. 3 = Gagnebina tamaricina.
pterocladu, F. Muell. Fragm. iv. 3 = trigonophylla.
pteriphylla, Hoffmgg. Verz. Pfl. (1824) 209 = pennata.
pterygocarpa, Hochst. ex Benth. in Hook. Lond. Journ. Bot. v. (1846) 96 = subdita.
ptychophylla, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 142.—Austral.
pubescens, *R. Br. in Ait. Hort. Kew.* ed. II. v. 467.—Austral.
pubescens, *Schlecht. in Linnaea*, xii. (1838) 563.—Mexico.
pugioniformis, *Wendl. Comm. Acac.* 88. t. 9.—Austral.
pugioniformis, Wendl. in Flora, ii. (1819) 189 = juniperina.
pulchella, *R. Br. in Ait. Hort. Kew.* ed. II. v. 464.—Austral.
pulcherrima, Willd. Sp. Pl. iv. 1061 = Stryphnodendron floribundum.
pulverulenta, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 342 = calamifolia.
pulverulenta, Schlecht. in Linnaea, xii. (1838) 571 = Leucaena pulverulenta.
punctata, Desf. Cat. Hort. Par. ed. III. 300 = Neptunia plena.
punctata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1084 = macracantha.
pungens, Spreng. Syst. iii. 184 = juniperina.
purpurascens, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 277.—Afr. trop.
purpurea, *Bolle, in Peters, Reise Mossamb. Bot.* 6.—Afr. trop.
purpurea, Steud. Nom. ed. II. i. 7 = Inga tergemina.
pycnantha, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 351.—Austral.
pycnophylla, *Benth. Fl. Austral.* ii. 388.—Austral.
pycnostachya, *F. Muell. Pl. Vict.* ii. 33.—Austral
pyrifolia, *DC. Mém. Lég.* xii. 447.—Austral.
quadrangularis, Link, Enum. Hort. Berol. ii. 445 = Calliandra tetragona.
quadrigeminalulae, Mart. in Flora, xx. (1838); Beibl. ii. 110 = riparia.
quadrilateralis, DC. Prod. ii. 451 = pugioniformis.
quadrilateralis, Decne. Herb. Tim. 177 = juncifolia?
quadrimarginea, *F. Muell. Fragm.* x. 81.—Adstral.
quadrisulcata, *F. Muell. l. c.* iii. 127.—Austral.
Raddiana, Savi, Sopra Alc. Acac. Egiz. 1 = tortilis.
ramosissima, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 356 ex parte.—Austral.

ACACIA:—
rumenaisnina, Benth. in Hook. Lond. Journ. Bot. i. (1842) 356 cz parte = Harveyi.
reclinata, F. Muell. 1st Gen. Rep. 12 = leprosa.
recurva, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 519.—Bras.
rubicina, *Wawra & Peyr. Sert. Benguel.* 15.—Afr. trop.
reniformis, *Benth. in Hook. Ic. Pl.* t. 1165.—Mexico.
restiacea, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 328.—Australia.
reticulata, *Willd. Sp. Pl.* iv. 1056.—Afr. austr.
retinervis, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 379.—Austral.
retinodes, Schlecht. in Linnæa, xx. (1847) 664 = neriifolia.
retivenea, *F. Muell. Fragm.* iii. 128.—Austral.
retusa, Meisn. in Bot. Zeit. (1855) 10 = sphacelata.
revoluta, Kunth, Mimos. Humb. &c. t. 26 = Mimosa revoluta.
rhigiophylla, *F. Muell. ex Schlecht. Benth. in Linnæa*, xxvi. (1855–56) 611.—Austral.
rhodacantha, *Desf. Tabl. Hort. Par.* ed. II. 208.—Hab.!
rhombifolia, *Desf. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 687.—Hab.!
rhombifolia, *Poir. ex Steud. Nom.* ed. I. 8.—Ins. Trinit.
Ricana, *Hrnal. in Mazud. Botanist*, t. 135.—Tasman.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 459. t. 58. —Ins. Fiji.
rigens, *A. Cunn. ex G. Don, Gen. Syst.* ii. 408.—Austral.
rigidula, Benth. in Hook. Lond. Journ. Bot. i. (1842) 504 = amentacea.
riparia, Bert. ex Spreng. Syst. iii. 142 = polyphylla.
riparia, *H. B. & K. Nov. Gen. et Sp.* vi. 276.—N. Granat.
robusta, *Burch., Trav.* ii. 443.—Afr. austr.
Roemeriana, *Schenk. in Linnæa*, xxi. (1848) 456.—N. Mexico.
Rohriana, DC. Prod. ii. 457 = nudiflora.
rostellifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 366.—Austral.
rostelliformis, Seem. Ringef. Acac. t. 2 = microbotrya.
rostrata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1054 (err. typ. 1060) = Lysiloma rostrata.
rotundata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 591.—Am. trop.
rotundifolia, Hook. Bot. Mag. t. 4041 = obliqua.
Rovumæ, *Oliver, Fl. Trop. Afr.* ii. 353.—Afr. trop.
Roxburghii, Kostel. Allg. Med. pharm. Flora, iv. 1367 = Albuzzia myriophylla.
Roxburghii, Wight & Arn. Prod. i. 276 = planifrons et eburnea!
rubacensis, Mart. & Gal. in Bull. Acad. Brux. x. 11. (1843) 315 = Calliandra lasa.
rubida, *A. Cunn. in Field. New South Wales*, 344.—Austral.
rufa, Hamk. Cat. Pl. Hort. Bog. 290 = Albizzia rufa.
rugata, Buch.-Ham. in Wall. Cat. n. 5251 = concinna.
runciformis, A. Cunn. ex G. Don, Gen. Syst. ii. 404 = linosta.
rupestris, Stocks, ex Boiss. Fl. Orient. ii. 638 = A. Senegal.
rupicola, *F. Muell. ex Benth. in Linnæa*, xxvi. (1855–56) 610.—Austral.
ruscifolia, A. Cunn. ex G. Don, Gen. Syst. ii. 407 = verticillata.
rutæolia, *Ait. ex Steud. Nom.* ed. II. i. 8.—Austral.
rutæfolia, Link, Enum. Hort. Berol. ii. 444 = nigricans.
Sabeana, Buckl. in Proc. Ac. Sc. Philad. '1861' (1862) 463 = Leucæna retusa.
saccharata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 505 = albida.
salicina, *Lindl. in Mitch. Three Exped.* ii. 20.—Austral.
saligna, *Wendl. Comm. Acac.* 26.—Austral.
salisburgen, DC. Prod. ii. 456 = Prosopis julifora.
Saltiana, Steud. Nom. ed. II. i. 8 = A. Lahai.
saltuum, Jungh. in Bot. Zeit. iii. (1845) 703 = Albizzia mootana.
Sandra, Bedd. Fl. Sylv. t. 50 = A. Sundra.
sanguinea, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 245 = venosa.
sapindoides, A. Cunn. ex Sweet, Hort. Brit. ed. III. 198 = Pithecolobium pruinosum.
saponaria, Heyne, ex Benth. in Trans. Linn. Soc. xxx. (1875) 581 = concinna.
sarmentosa, Desv. Journ. Bot. iii. (1814) 70 = cæsia.
sarmentosa, Griseb. Fl. Brit. W. Ind. 221 = riparia.
Sasa, *Baill. Hist. Pl.* ii. 47.—Madag.

ACACIA:—
scabra, *Benth. in Linnæa*, xxvi. (1853–55) 605.—Austral.
scalpelliformis, *Meisn. in Lehm. Pl. Preiss.* ii. 200.—Austral.
scandens, Benth. in Hook. Lond. Journ. Bot. i. (1842) 517 = pluricoa.
scandens, Willd. Sp. Pl. iv. 1057 = Entada scandens.
scandens, Willd. Enum. Hort. Berol. 1057 = polyphylla.
scapuliformis, A. Cunn. in G. Don, Gen. Syst. ii. 405 = cultriformis.
schinoides, Benth. in Hook. Lond. Journ. Bot. i. (1842) 382 = prunoca.
Schottii, *Torr. in Bot. Mex. Bound.* 62.—Am. bor.
scirpifolia, *Meisn. in Bot. Zeit.* (1855) 10.—Austral.
sclerophylla, *Lindl. in Mitch. Three Exped.* ii. 130.—Austral.
scleroperma, *F. Muell. in Wing. South. Sc. Record*, (July, 1883).—Austral.
sclerosyla, *Juss. Fl. Antill.* i. t. 21.—Ind. occ.
secundiflora, Juss. ex DC. Prod. ii. 452 = Entada polystachya.
Seresia, Roxb. ex Steud. Nom. ed. II. i. 8 = Albizzia Lebbek.
selenicarpa, Desv. in Ham. Prod. Fl. Ind. Occ. 60 = albida.
Seliei, Spreng. Syst. iii. 137 = Calliandra brevipes.
semicordata, *Roxb. ex Steud. Nom.* ed. II. i. 8.—Ind. or.
semispinosa, Willd. ex Steud. l. c. = Mimosa semispinosa.
semiteretes, *Mart. ex Steud. l. c.*—Bras.
semiteretifolia, Knowl. & Weste. Fl. Cab. ii. (1838) 27 = verticillata.
Senegal, *Willd. Sp. Pl.* iv. 1077.—Geront. trop.
Sentu, *Mudl. 2nd Gen. Report*, 7 (1854).—Austral.
sericata, Ait. ex Steud. Nom. ed. II. i. 8 = seq. ?
sericata, *L. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 380.—Austral.
serica, Mart. & Gal. in Bull. Acad. Brux. x. 11. (1843) 311 = acuminata.
sericocephala, Fenzl, in Flora, xxvii. (1844) 312 = Albizzia amara.
sericophylla, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 122 = stenophylla.
Serra, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 519.—Bras.
serriformis, A. Cunn. in Bot. Mag. sub t. 3894 = undulatifolia.
sessiliceps, *F. Muell. in Melb. Chemist* (July, 1882). —Australia.
sessilis, Benth. in Hook. Lond. Journ. Bot. i. (1842) 336 = sphacelata.
setigera, Hook. Ic. Pl. t. 861 = undulacifolia.
setosa, *Dekmh. Rvoist. Nap.* i. 169; *ex Walp. Rep.* i. 924. —Hab.!
setosa, *Spreng. Syst.* iii. 137.—Bras.
setulifera, *Benth. in Linnæa*, xxvi. (1853–55) 625. —Australia.
Seyal, *Delile, Fl. Aegypte*, 266. t. 52. f. 2.—Aegypt.
Shuttleworthii, *Meisn. in Lehm. Pl. Preiss.* i. 7.—Australia.
siculiformis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 387.—Austral.
Sieberi, Tausch, ex Steud. Nom. ed. II. i. 8 = crassicarpa.
Sieberiana, *DC. Prod.* ii. 458.—Senegal.
Sieberiana, Scheele, in Linnæa, xxvii. (1853) 387 = discolor.
Sieberiana, Tausch, in Flora, xix. (1836) 420 = crassisiucula.
signata, *F. Muell. Fragm.* iv. 7.—Austral.
Siliquastrum, Lag. Gen. et Sp. Nov. 16 = Prosopis juliflora.
similis, *Zoll. & Morr. in Nat. en Genoesch. Arch. N. Ind.* iii. (1846) 81.—Malaya.
Sinuli, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 368.—Austral.
Smg, Galli. & Per. Test. Fl. Senag. i. 251 = Sieberiana.
Sirissa, Buch.-Ham. in Wall. Cat. n. 5246 = Albizzia Lebbek.
smilacifolia, Field. & Gardn. Sert. Fl. t. 3 = urophylla.
Smithiana, Roxb. ex Wall. Cat. n. 5237 = Albizzia stipulata.
Sobeana, *Buckl. Proc. Acad. Nat. Sc. Philad.* (1862) 468.—Am. bor.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1883) 513.—Ins. Socotra.

ACACIA:—
Solandri, *Benth. Fl. Austral.* ii. 406.—Austral.
somalensis, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 274. —Afr. trop.
somnians, Sterler, ex Steud. Nom. ed. II. i. 8 = Mimosa somnians.
Sophoræ, R. Br. in Ait. Hort. Kew. ed. II. v. 462 = longifolia.
spadicigera, *Cham. & Schlecht. in Linnæa*, v. (1830) 594.—Jamaica.
spathulata, *F. Muell. ex Benth. Fl. Austral.* ii. 356.—Austral.
spathulata, Tausch, in Flora, xix. (1836) 420 = longifolia.
speciosa, Hort. ex Steud. Nom. ed. II. i. 8 = Albizzia lophantha.
speciosa, Willd. Sp. Pl. iv. 1066 = Albizzia Lebbek.
spectabilis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 388.—Austral.
sphacelata, *Benth. l. c.* 338.—Austral.
sphærocephala, *Cham. & Schlecht. in Linnæa*, v. (1830) 594.—Mexico.
spinescens, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 393.—Austral.
Spini, *Balb. in Spin. Cat. Suppl.* (1823) 8.—Ins. Guadaloup.
spinosa, E. Mey. Comm. Pl. Afr. Austr. i. 170 = Dichrostachys nutans.
spinosissima, *Benth. ex Schlecht. in Linnæa*, xxvi. (1855–55) 621.—Austral.
spinulosa, *Sweet, ex Steud. Nom.* ed. II. i. 8.—Austral.
spirocarpa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 229. —Afr. trop.
spirorbis, *Labill. Sert. Austr. Caled.* 69. t. 69.—N. Caled.
spondylophylla, *F. Muell. Fragm.* viii. 248.—Austral.
Sprengelii, G. Don, Gen. Syst. ii. 410 = Berterians.
Sprengelii, Hook. & Arn. in Mook. Bot. Misc. iii. (1833) 300 = Mimosa incana.
squamata, *Lindl. Swan Riv. App.* 15.—Austral.
squamata, Morr. in Ann. Sec. Gand, lii. (1847) 209. t. 134 = restiacea.
squarrosa, Mart. ex Benth. in Hook. Lond. Journ. Bot. iii. (1844) 104 = Calliandra squarrosa.
stellata, *Willd. Sp. Pl.* iv. 1072.—Arab.
stenocarpa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 233.—Abyss.
stenophylla, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 386.—Austral.
stenoptera, *Benth. l. c.* 325.—Austral.
stenostachya, Desv. ex Ham. Prod. Fl. Ind. Occ. 59 = Piptadenia psilostachya.
Stephaniana, Willd. Sp. Pl. iv. 1088 = Prosopis Stephaniana.
stereophylla, *Meisn. in Lehm. Pl. Preiss.* ii. 208.—Austral.
stigmatophylla, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 327.—Austral.
stipellata, Schlecht. in Linnæa, xii. (1838) 574 = filicina.
stipulata, DC. Prod. ii. 469 = Albizzia stipulata.
stipuligera, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 144.—Austral.
stipulosa, *F. Muell. l. c.* 119.—Austral.
stolonifera, Burch. Trav. ii. 241 = hebeclada.
striata, *Kunth. & Bonpl. ex Willd. Sp. Pl.* iv. 1069.—Am. austr.
striatula, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 396.—Austral.
stricta, *Willd. Sp. Pl.* iv. 1052.—Austral.
strigosa, Lindl. in Mitch. Three Exped. ii. 135 = asperu.
strigosa, *Link, Enum. Hort. Berol.* ii. 444.—Austral.
strigosa, Spreng. Syst. iii. 137 = strigulosa.
strigulosa, *Steud. Nom.* ed. II. i. 8.—Peruv.
strombulifera, Willd. Sp. Pl. iv. 1085 = Prosopis strombulifera.
strongylophylla, *F. Muell. Fragm.* viii. 226.—Austral.
Stuartiana, F. Muell. ex Benth. in Linnæa, xxvi. (1855–56) 609 = aciculiformis.
suaveuala, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 863.—Madag.
suaveolens, *Willd. Sp. Pl.* iv. 1060.—Austral.
subalata, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 276.—Afr. trop.
subbiformis, Meisn. ex Lehm. Plant. Preiss. i. 16 = acuiliformis.
subcæsalpina, *Lindl. in Bot. Reg.* t. 1075.—Austral.

ACACIA:—
subdimidiata, Spitzerb. in Hoov. & De. Vr. Tijdschr. ix. (1843) 113 = Pithecolobium multiflorum.
suberosa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 409.—Austral.
subfalcata, Meisn. in Bot. Zeit. (1855) 11 = micro botrys.
subinermis, Berr. ex DC. Prod. ii. 462 = macracantha.
sublanata, *Benth. in Enum. Pl. Hueg.* 42.—Austral.
subporosa, *F. Muell. Pl. Vict.* ii. 24.—Austral.
subrhombea, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 864.—Madag.
subtenuata, *F. Muell. & Benth. in Journ. Linn. Soc.* iii. (1859) 124.—Austral.
subtilifolia, H. B. & K. Nov. Gen. et Sp. vi. 268 = Piptadenia subtilifolia.
subtilinervis, *F. Muell. Pl. Vict.* ii. 32.—Austral.
subtilis, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 48.—Austral.
subulata, *Bonpl. Jard. Malm.* 110. t. 45.—Austral.
sulcata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 460.—Austral.
sulcipes, Sieber, ex G. Don, Gen. Syst. ii. 419 = decurrens.
Suma, Buch.-Ham. ex Wall. Cat. sub n. 5227 = A. Sundra.
Suma, *Kurz, ex Brandis. For. Fl.* 187.—Ind. or.
Sundra, *DC. Prod.* ii. 458.—Ind. or.
Sundra, Roxb. in Wall. Cat. n. 5227 = A. Suma.
taitensis, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 278.—Afr. trop.
tamarindifolia, Grisch. Cat. Pl. Cub. 82 = Mimosa polyanclstra.
tamarindifolia, *Willd. Sp. Pl.* iv. 1092.—Am. austr.
tamarisciua, Willd. i. c. 1083 = Gagnebina tamariscina.
tazifolia, A. Cunn. in Field, New South Wales, 844 = trinervata.
taxifolia, Lodd. Bot. Cab. t. 1225 = Ricsena.
taxifolia, *Willd. Sp. Pl.* iv. 1050.—Cochinch.
Taylioriana, *F. Muell. in Wing, South. Sc. Record* (July, 1882).—Austral.
tenuriua, Mig. Fl. Ind. Bat. i. 14. t. 3 = pennata.
tenuiflora, Willd. Sp. Pl. iv. 1088 = Mimosa Cabrera.
tenuifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 87.—Austral.
tenuifolia, Willd. Sp. Pl. iv. 1091 = paniculata.
tenuissima, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 135 = xylocarpa.
teplirolea, A. Gray, Pl. Wright i. 65 = Berlandieri.
teretifolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 396.—Austral.
tetragona, Willd. Sp. Pl. iv. 1069 = Calliandra tetragona.
tetragonocarpa, *Meisn. in Lehm. Pl. Preiss.* i. 4.—Austral.
tetragonophylla, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 121.—Austral.
taxensis, Torr. & Gray, Fl. N. Am. i. 404 = filicina.
thegonocarpa, A. Cunn. ex Sweet, Hort. Brit. ed. III. 198, sphalm. = trigonocarpa.
Thibaudiana, DC. Prod. ii. 456 = Piptadenia moniliformis.
Thonningiana, F. Muell. Fragm. iv. 3 = Albizzia Thonningiana.
Tomentilla, Zipp. ex Span. in Linnaea, xv. (1841) 199 = pennata.
tomentosa, *Willd. Sp. Pl.* iv. 1087.—Ind. or.
torquata, Lag. Gen. et Sp. Nov. 16, n. 206 = Prosopis torquata.
torrilis, *Hayne, Arzneigew.* ix. t. 31.—Afr. trop.; Amb.
tortuosa, *Willd. Sp. Pl.* iv. 1088.—Am. austr.
torulosa, *Benth. in Journ. Linn. Soc.* iii. (1859) 139.—Austral.
transitiosa, *A. Cunn. in Hook. Ic. Pl.* t. 160.—Austral.
trapezoidea, DC. ex Steud. Nom. ed. II. i. 8 = dasiphylla.
triacantha, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 244 = glaucophylla.
triangularis, Benth. Enum. Pl. Hueg. 44 = bifora.
trichandra, Zucc. Pl. Nov. Fasc. ii. 41 = Leucaena diversifolia.
trichodes, Willd. Sp. Pl. iv. 1068 = Leucaena trichodes.
trichophylladen, Macfad. Fl. Jamaic. 230 = Piptadenia peregrina.
trigona, A. DC. Nat. vill. Fl. Rar. Jard. Gen. 20 = myrtifolia.
trigonocarpa, A. Cunn. ex Lond. Hort. Brit. 407 = longifolia.

ACACIA:—
trigonophylla, *Meisn. in Lehm. Pl. Preiss.* ii. 199.—Austral.
trinervata, *Sieber, ex DC. Prod.* ii. 451.—Austral.
trinervis, *Desv. Journ. Bot.* iii. (1814) 70.—Hab.?
trineura, *F. Muell. Pl. Vict.* ii. 26.—Austral.
triptera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 395.—Austral.
triptycha, *F. Muell. ex Benth. Fl. Austral.* ii. 387.—Austral.
triquetra, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 358.—Austral.
triquetra, Mart. Herb. Fl. Bras. 108 = Piptadenia stipulacea.
tristinosa, Stokes, Bot. Mat. Med. iii. 168 = A. Senegal.
trinervosa, F. Muell. Fragm. iv. 6 = heterocilia.
tristis, R. Grah. in Bot. Mag. t. 3420 = armata.
tristis, *Welw. ex Oliver, in Fl. Trop. Afr.* ii. 349.—Afr. trop.
truncata, Hort. ex Hoffmgg. Pfl. Verz. 84 = decipiens?
tuberosa, *Sterler, Hort. Nymph. ex Steud. Nom.* ed. II. i. 8.—Hab.?
tubulifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 520.—Peruv.
tucumanensis, *Grisch. in Goett. Abh.* xix. (1874) 135.—Am. austr.
tumida, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 144.—Austral.
Tumndra, Steud. Nom. ed. II. i. 8, sphalm. = Sundra.
ulicina, *Meisn. in Lehm. Pl. Preiss.* ii. 202.—Austral.
umbellata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot* i. (1842) 378.—Austral.
umbelliflora, Kunth, Mimos. Humb. 100. t. 18 = filicina.
umbraculata, Wight, in Wall. Cat. n. 5245 = latronum.
umbrosa, A. Cunn. ex G. Don, Gen. Syst. ii. 405 = bivona.
uncifera, *Benth. in Mitch. Trop. Austr* 241.—Austral.
uncinata, Lodd. Bot. Cab. t. 909 = undulaefolia.
uncinata, *Benth. in Linnaea*, xxvi. (1855-55) 613.—Austral.
uncinella, DC. ex Steud. Nom. ed. II. i. 8 = acanthocarpa.
uncinella, Desf. Cat. Hort. Par. ed. III. 299 = Mimosa uncinella.
undulata, Wild. Enum. Hort. Berol. Suppl. 68 = armata.
undulaefolia, *A. Cunn. ex G. Don, Gen. Syst.* ii. 404.—Austral.
ungulata, Desv. Journ. Bot. iii. (1814) 69 = Calliandra portoricensis.
unigiandulosa, Seem. & Schmidt, in Flora, xxvii. (1844) 495 = alata.
uropbylla, *Benth. ex Lindl. in Bot. Reg.* (1841), *Misc.* 24.—Austral.
vaga, *Willd. Sp. Pl.* iv. 1066.—Bras.
Valenzuelana, A. Rich. Ess. Fl. Cub. 462 = villosa.
variana, Benth. in Mitch. Journ. Trop. Austral. 182 = bivona.
velutina, Benth. in Hook. Lond. Journ. Bot. i. (1842) 519 = monacantha.
velutina, *Bertol. Pl. Nov. Hort. Bonon.* ii. 13. t. 4.—Ind. occ.
velutina, *DC. Prod.* ii. 459.—Bras.
venosa, *Hayne, ex Benth. in Hook. Lond. Journ. Bot.* v. (1846) 98.—Abyss.
venulosa, *Benth. l. c.* 366.—Austral.
venusta, *Regel & Korn. Ind. Sem. Hort. Petrop.* (1858.—Hab.?
vennuste, Willd. Enum. Hort. Berol. 1052 = Calliandra portoricensis.
vera, Willd. Sp. Pl. iv. 1085 = arabica.
Verek, Guill. & Perr. Tent. Fl. Senag. i. 245. t. 56 = A. Senegal.
verniciflua, *A. Cunn. in Field, New South Wales*, 844.—Austral.
verticillata, Steber, ex Benth. Fl. Austral. ii. 332 = juniperina.
vertillata, *Willd. Sp. Pl.* iv. 1049.—Austral.
verugera, *Schweinf. in Linnaea*, xxxv. (1867-68) 340. t. 9, 10.—Afr. trop.
vesperlina, Macfad. Fl. Jamaic. i. 218 = Calliandra portoricensis.
vestita, *Ker-Gawl. in Bot. Reg.* t. 698.—Austral.
viarum, *Allem. ex Tenore, Cat. Ort. Napol.* 77.—Bras.
viciodes, Forr. & Galln. Voy. Abyss. iii. 119 = mellifera.

ACACIA:—
Victoriae, Benth. in Mitch. Journ. Trop. Austral. 383 = senta.
villosa, *Willd. Sp. Pl.* iv. 1067.—Jamaica.
viminalis, *Ait. ex Steud. Nom* ed. II. i. 8.—Austral.
viminea, *Dietr. ex Steud. l. c.* 9.—Austral.
Vincentis, Grisch. Fl. Brit. W. Ind. 222 = Pithecolobium tortum.
Virchowiana, *Vatke & Hildeb. in Oest. Bot. Zeit.* xxx (1880) 375.—Afr. trop.
virescens, *DC. Cat. Hort. Monsp.* 74.—Am. austr.
virgata, Gaert. Fruct. ii. 317. t. 148 = Desmanthus virgatus.
virgata, Lodd. Bot. Cab. t. 1246 = verniciflua.
virgata, *Rafin. Fl. Ludov.* 136.—Am. bor
virgultosa, Vahl, ex Walp. Rep. i 929 = Inga virgultosa.
viridiflora, Kunth, Mimos. Humb. 81. t. 25 = Piptadenia subtilifolia.
viridiflora, Tweed. ex Benth. in Hook. Lond. Journ. Bot. iii. (1844) 98 = Calliandra viridiflora.
viridiramis, Burch. Trav. i. 300 = Xerocladia Zeyheri
virida, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 868.—Madag.
viscidula, Benth. in Mssnd, Botanist, iv. t. 179 = elongata.
viscidula, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 368.—Austral.
viscifua, F. Muell. Pl. Vict. ii. 24 = dodonaefolia.
Visco, *Lorents, ex Grisch. in Goett. Abh.* xxiv. (1879) 132.—Reg. Argent.
viscosa, Schrad. ex Wendl. Comm. Acac. 80. t. 7 = dodonaefolia.
Visite, *Grisch. in Goett. Abh.* xix. (1874) 135.—Reg. Argent.
vianeoides, *Colla, in Linnaea*, iv. (1829) *Lit.* 56.—Austral.
Vogeliana, Steud. Nom. ed. II. i. 9 = ambigua.
volubilis, *F. Muell. Fragm.* x. 98.—Austral.
vomeriformis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 392.—Austral
vulcanica, Korth. Hort. Bogor. ex Flora, xxx. (1847) 705 = Albizzia montana.
Wallichiana, DC. Prod. ii. 458 = A. Catechu.
Watsoniana, *F. Muell. in Benth. Fl. Austral.* ii. 374.—Austral.
Welwitschii, *Oliver, Fl. Trop. Afr.* ii. 341.—Afr. trop.
Westiana, DC. Prod. ii. 464 = riparia.
Whanii, *F. Muell. ex Benth. Fl. Austral.* ii. 386.—Austral.
Wichkani, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 379.—Austral.
Wightiana, R. Grah. in Wall. Cat. n. 5259 = Albizzia amara.
Wightii, *Baker, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 508.—Ind. or.
Wightii, Wight & Arn. Prod. 274 = Albizzia amara.
Wilkesiana, F. Muell. in Trans. Phil Soc. Victoria, i. (1855) 87 = calamifolia.
Willdenowiana, Wendl. Vers. K. Beiggart. (1845) 5 = diptera.
Wrightii, *Benth. ex A. Gray, in Pl Wright.* i. 64.—Texas.
xanthina, Benth. in Hook. Lond. Journ. Bot. i. (1842) 355 = bivenosa.
xanthophloea, *Benth. in Trans. Linn. Soc.* xxx. (1875) 511.—Afr. trop.
xiphocarpa, Hochst. ex Benth. in Hook. Lond. Journ. Bot. v. (1846) 98 = abyssinica.
xylocarpa, *A. Cunn. ex Benth. l. c.* i. (1842) 370.—Austral.
xylocarpa, Willd. Sp. Pl. iv. 1055 = Xylia dolabriformis.
Zygia, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 864.—Madag.
zygioides, *Baill. l. c.* 863.—Madag.

ACAENA, Linn. Mant. ii. 145 (1771).—*ROSACEAE*, Benth. & Hook. f. i. 622.
ÆKCISTRUM, Forst. Char. Gen. 8. t. 2. (1776).
LASIOCARPUS, Banks & Soland. ex Hook. f. Fl. Antarct. 264 (1847).
SPHÆRULA, Anders. ex Hook. f. l. c. 268 (1847).
aculeata, *Meyen, Reise.* i. 348.—Chili.
adscendens, *Vahl, Enum.* i. 297.—Am. austr.; Austral.
aginis, Hook. f. Fl. Antarct. 268. t. 96 = adscendens.

ACAENA:—
agrimonioides, *H. B. & K. Nov. Gen. et Sp.* vi. 281. —Mexico.
alpina, *Poepp. in Nov. Act. Nat. Cur.* xix. Suppl i. (1843) 326.—Chili.
andicola, Gill. ex Walp. Rep. ii. 48 = pinnatifida.
andina, *Phil. in Linnaea*, xxviii. (1856) 636.—Chili.
antarctica, *Hook. f. Fl. Antarct.* 269.—Reg. Magell.
argentea, Bert. ex Stend. Nom. ed. ii. 9 = Berteriana.
argentea, *Ruiz & Pav. Fl. Per.* i. 67. t. 103. f. b.—Chili.
Behriana, Schlecht. in Linnaea, xx. (1847) 660 = ovina.
Berteriana, *Bernh. ex Steud. Nom.* ed. ii. i. 9.—Chili.
Buchananii, *Hook. f. Handb. New Zeal Fl.* 57.—N. Zel.
Cadilla, *Hook. f. Fl. Antarct* 269.—Patagonia.
caespitosa, *Gill in Hook. Bot. Misc.* iii. 307 (1833).—Chili.
Calcitrapa, *Phil. in Linnaea*, xxxiii (1864–65) 67.—Chili.
canescens, *Phil. Fl. Atac.* 18.—Chili.
Closiana, *C. Gay, Fl. Chil.* ii 298.—Chili.
Coxi, *Phil. in Linnaea*, xxxiii. (1864–65) 62.—Chili.
cuneata, *Hook & Arn. in Hook. Bot. Misc.* iii. 307 (1833) = Reg Magell
cylindristachys, *Ruiz & Pav. Fl. Per.* i. 68 t. 104. f. 2.—Peruv.
decumbens, Menzies, in Hook. f. Fl. Antarct. 9 = Sanguisorbae
depressa, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 548. —N Zeland.
digitata, *Phil. in Anales Univ. Chil.* (1862) ii. 445.—Chili.
echinata, Lehm. Pl. Preiss, i. 95 = ovina.
elegans, *C. Gay, Fl. Chil.* ii. 297.—Chili.
elongata, *Linn. Mant.* ii. 200.—Mexico.
Espatoria, *Schlecht. in Linnaea*, ii. (1827) 80.—Uruguay.
exigua, *A. Gray, Bot. U. S. Expl. Exped.* i. 498.—Ins. Sandvic.
glaberrima, Phil. in Linnaea, xxviii. (1856) 668 = pumila.
glabra, *T. Buch. in Trans. N. Z. Inst.* iv. (1872) 226. t. 14.—N. Zel.
Grahamiana, Gill. ex Walp. Rep. ii. 48 = adscendens.
hirsuta, *Phil. in Anales Univ. Chil.* (1872) 712.—Chili.
Hectori, *R. Br. [ter.] in Trans. N. Z. Inst.* xvi. (1884) 389.—N. Zeland.
incisa, Lindl. Bot. Reg. t. 1371 = pinnatifida.
inermis, *Hook. f. Fl. N. Zel.* i. 54.—N. Zel.
integerrima, *Gill. in Hook. Bot. Misc.* iii. (1833) 306. —Chili.
Krauseri, Phil. in Anales Univ. Chil. (1862) 711 = laevigata.
laevigata, *Ait. Hort. Kew.* ed. II. i. 68.—Reg. Magell.
lappacea, *Ruiz & Pav. Fl. Per.* i. 66. t. 103. f. a.—Peruv.
latebrosa, *Ait. Hort. Kew.* ed. II. i. 67.—Afr. austr.
laxiflora, *Willd. ex Steud. Nom.* ed. II. i. 9.—Am. austr.
leptacantha, *Phil. in Linnaea*, xxxiii. (1864–65) 66.— Chili.
lucida, *Vahl, Enum.* i. 296.—Ins. Falkl.
lucida, Willd. ex Steud Nom. ed. II. i. 9 = pinnatifida.
macrocephala, *Poepp. Fragm. Syn.* 25.—Chili.
macrorhiza, *Hook. f. Fl. Antarct.* 266.—N. Granat.
macrostemon, *Hook. f in Hook. Lond. Journ. Bot.* vi. (1846) 476, 269.—Chili.
magellanica, *Vahl, Enum.* i. 297.—Reg. Magell.
microcephala, *Schlecht. in Linnaea*, xxviii. (1856) 463. —Chili.
microphylla, *Hook. f. Fl. N. Zel.* i. 55.—N. Zel.
montana, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1846) 276.—Tasman.
montana, *Phil. in Linnaea*, xxviii. (1856) 636.—Chili.
montevidensis, Hook. f. Fl. Antarct. 265 = A. Espatoria.
multifida, A. Dietr. Sp. Pl. i. 619 = trifida.
multifida, *Hook. f. Fl. Antarct.* 265.—Patagonia.
myriophylla, *Lindl. Bot. Reg. t.* 1371.—Reg. Argent.
Novae-Zelandiae, *T. Kirk, in Trans. N. Z. Inst.* iii. (1871) 177.—N. Zel.
ochreata, *Wedd. Chlor. And.* ii. 240.—Peruv.
oligacantha, *Phil. in Linnaea*, xxxiii. (1864–65) 66.— Chili.
ovalifolia, *Ruiz & Pav. Fl. Par.* i. 67. t. 103.—Am. austr.

ACAENA:—
ovalis, Pers. Syn. i. 142 = ovalifolia.
ovina, *A. Cunn. in Field, New South Wales*, 368.— Austral.
pectinata, *C. Presl, Epim. Bot.* 201.—Chili.
pinnatifida, Bert. ex Steud. Nom. ed. II. i. 9 = cuneata, myriophylla.
pinnatifida, Hort. ex Steud. l. c. = A. Berteriana.
pinnatifida, Lindl. Bot. Reg. t. 1371 = trifida.
pinnatifida, *Ruiz & Pav. Fl. Per.* i. 68. t. 104.—Chili.
Poeppigiana, *C. Gay, Fl. Chil* ii. 294.—Chili.
polycarpa, *Griseb. in Goett. Abh.* xix. (1874) 138.— Reg. Argent.
potentoides, *Wedd. ex Steud. Nom.* ed. II. i. 9.—Am. austr.
pumila, *Phil. in Linnaea*, xxxiii. (1864–65) 67.—Chili.
pumila, *Vahl, Enum.* i. 298.—Reg. Magell.
quinquefida, *Phil. in Linnaea*, xxvin. (1856) 636.—Chili.
Sanguisorbae, *Vahl, Enum.* i. 294.—Austral.
sarmentosa, Carmich. in Trans. Linn. Soc. xii. (1818) 509 = A. Sanguisorbae.
sericea, *Yacq. f. Ecl.* i. 81. t. 55.—Mexico.
splendens, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 306.—Chili.
stellaris, *Meyen, Reise*, i. 348.—Chili.
stricta, *Griseb. in Goett. Abh.* xix. (1874) 138.—Reg. Argent.
suaveolens, A. Gray, Bot. U. S. Expl. Exped. i. 748, sphalm. = Acacia suaveolens.
subincisa, *Wedd. Chlor And.* ii. 242.— Ecuador.
tridactyla, *Presl, Epim. Bot.* 201.—Mexico.
trifida, *Ruiz & Pav. Fl. Per.* i. 67. t. 104. f. c.—Calif.; Peru; Chili.
trifida, Steud. & Hochst. ex Steud. Nom. ed. II. i. 9 = Berteriana.
valdiviana, *Phil. in Linnaea*, xxxiii. (1864–65) 65.— Chili.
venulosa, *Griseb. in Goett. Abh.* vi. (1854) 118.—Reg. Magell.

ACAENOPS, Schrad. ex Steud. Nom. ed. II. 9 (1840) = DIPSACA, Linn.
pilosus, Fourr. in Ann. Soc. Linn. Lyon. N.S. xvi. (1868) 400 = D. pilosus.
vulgaris, Schrad. ex Steud. Nom. ed. II. i. 9 = D. pilosus.

ACAJOU, Toorn. ex Adans. Fam ii. 344 (1763) = seq.

ACAJUBA, Gaertn. Fruct. i. 192. t. 40. (1788) = **Anacardium**, Linn.
occidentalis, Gaert. l. c. = Anac. occidentale.

ACALYPHA, Linn. Coroll. Gen. 19 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii 311.
CALYPTROSPATHA, Klotzsch, ex Baill. Etudes Euph. 440 (1858).
CATURUS, Linn. ex Schreb. Gen. Pl. 677 (1791).
CUPAMENI, Adans. Fam. ii. 356 (1763).
GALURUS, Spreng. Anleit. ii. 71. 264 (1817).
GYMNALYPHA, Griseb. in Bonplandia, vi. (1858) 2.
LINOSTACHYS, Klotzsch, ex Schlecht. in Linnaea, xiii. (1847) 385.
ODONTEILEMA, Turcz. in Bull. Soc. Nat. Mose. xxi. 1. (1848) 587.
URTICASTRUM, Desnet. Schinesa Hort. Malab. 31 (1818).
abortiva, Hochst. ex Baill. Etud. Gén. Euphorb. 443 = indica.
accedens, *Muell. Arg. in Linnaea*, xxiv. (1855–56) 85.—Brasil.
acuminata, *Benth. in Hook. Kew Journ.* vi. (1854) 329. —Bras.
acuminata, Vahl, ex Baill. Adansonia, i. (1860) 267 = decidua.
acuta, Thunb. Fl. Cap. ed. Schult. 546 = Adenocline acuta.
adenophora, Griseb. in Goett. Nachr. (1865) 175 = membranacea.
adenostachya, *Muell. Arg. in Linnaea*, xxxiv. (1865) 91.—Mexico.
adenotricha, *A. Rich. Tent. Fl. Abyss.* ii. 246.—Abyss.
adscendens, Hormem. Hort. Hafn. Suppl. 108 = chamaedrifolia.
affinis, Klotzsch, in Nov. Act. Acad. Nat. Cur. xix. Suppl. i. (1843) 416 = stipulacea.
agrimonioides, D. Dietr. Syn. v. 277 err. typ. = anemioides.
alnifolia, Poir. Encyc. vi. 303 = Poiretii.

ACALYPHA:—
alnifolia, *Klein, ex Willd. Sp. Pl.* iv. 525.—Ind. or.
alnifolia, Wall. Cat. n. 7765 A = capitata.
alopecuroides, *Yacq. Ic. Rar.* iii. 19. t. 620; *Coll.* iii. 196. —Venezuela.
ambyodonta, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 265.—Bras.
amboynensis, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 233 = grandis.
amentacea, Roxb. Fl. Ind. iii. 676 = fruticosa.
andina, *Muell. Arg. in Flora*, lv. (1872) 26.—Reg. Andin.
anemioides, *H. B. & K. Nov. Gen. et Sp.* ii. 94.— Mexico.
angatensis, *Blanco, Fl. Filip.* ed. I. 750.—Ins Philipp
angolensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 325.— Afr. trop.
angustata, Sond. in Linnaea, xxiii. (1850) 115 = pedunculariis.
angustifolia, *Sw. Prod. Veg. Ind. Occ.* 99.—Ins. S. Doming
anisodonta, *Muell. Arg. in DC. Prod.* xv. ii. 819.—Ins. Pacif.
arborea, *Comm. ex Poir. Encyc.* vi. 305.—Ins. Masc.; St. Helen.
Arcuana, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 262.— Bras.
aristata, *H. B. & K. Nov. Gen. et Sp.* ii. 93.—N. Granat
arvensis, *Poepp. & Endl. Nov. Gen & Sp.* iii. 21.—Peruv.
australis, *Linn. Sp. Pl.* 1004.—Reg. trop.
Baillioniana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 44.—Afr. trop.
Baroni, *Baker, in Journ. Linn. Soc.* xx. (1884) 254.— Madag
benguelensis, *Muell. Arg in Journ. Bot.* ii. (1864) 325.—Afr. trop.
Berteroana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 33.—Porto-Rico.
betulaefolia, Sw. Prod. Veg. Ind. Occ. 99 = scabrosa.
betulina, Retz. Obs. Fasc. v. 30 = fruticosa.
betulina, Schwenf. Fl. quaedam Nilot. 18 = Cephalocroton cordofanus.
betulina, Sond. in Linnaea, xxiii. (1850) 116 = glabrata.
betuloides, Klotzsch, ex Baill. in Adansonia, v. (1864–65) 238 = variabilis.
betuloides, Pav. ex Klotzsch, in Seem. Bot. Herald, 101 = diversifolia.
Bilbergiana, Klotzsch, l. c. = diversifolia.
bipartita, *Muell. Arg. in Flora*, xlvii. (1864) 538.—Afr. trop.
bisetosa, *Spreng. Syst.* iii. 879.—Venezuel.
boehmerioides, *Miq. Fl. Ind. Bat. Suppl.* i. 459.—Ins. Pacif.
Boiviniana, *Baill. Adansonia*, i. (1860–61) 272.— Afr. trop.
boliviensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 162.—Bolivia.
Botteriana, *Muell. Arg. l. c.* 46.—Mexico.
brachiata, Kraus, in Flora, xxvii. (1845) 68, nomen = tenuis.
brachyandra, *Baill. in Adansonia*, v. (1864–65) 232. —Bras.
brachyclada, *Muell. Arg. in DC. Prod.* xv. ii. 862.— Peruv.
brachycarpa, *Hornem. Hort. Hafn.* ii. 909.—China.
bractesta, *Miq. Fl. Ind. Bat.* ii. 406.—Malaya.
brasiliensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 37.—Bras.
brevibracteata, *Muell. Arg. in DC. Prod.* xv. ii. 855.— Ins. Pacif.
brevicaulis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 13.—Mexico.
brevipes, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 248.— Bras.
brevipes, *Rafin. New Fl Am.* i. 44.—Am. bor.
Buchenavii, *Muell. Arg. in Bremen, Abh.* vii. (1880) 27.—Afr. trop.
bullata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 17. —Peruv.
californica, *Benth. Bot. Voy. Sulph.* 51.—Calif.
callosa, *Benth. Pl. Hartw.* 252.—N. Granat.
calycyfrancis, Wight, ex Wall. Cat. n. 7766 = brachystachya.
canana, *Muell. Arg. in Flora*, xlvii. (1864) 538. —N. Granat.
canescens, Wall. Cat. n. 7765 = indica.

ACALYPHA:—
capensis, D. Dietr, Syn. v. 878, sphalm. = corensis.
caperonioides, Baill. Adansonia, iii. (1862-63) 157 = peduncularis.
capillipes, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 40.—Austral.
capitata, Wall. Cat. n. 7788. A, B = fruticosa.
capitata, *Willd. Sp. Pl.* iv. 525.—Ind. or.
carolinians, Blanco, Fl. Filip. ed. I. 748 = indica.
carolinians, *Rll. Sketch.* ii. 645.—Am. bor.
carolinians, Walt. Fl. Carol. 238 = virginica.
carpinifolia, Poepp. ex Baill. Etud. Gén. Euphorb. 449 = leptostachya.
carpinifolia, *Porr. in Law. Encyc.* vi. 203. (1804).—Ins. S. Doming.
carthagenensis, *Jacq. Enum. Pl. Carib.* 32.—N. Granat.
Caturus, *Blume, Bijdr.* 629.—Java.
caudata, H. B. & K. Nov. Gen. et Sp. ii. 95 = macrostachya.
chamaedrifolia, Muell. Arg. in DC. Prod xv. ii. 879 = reptans.
Chiloombos, *Baill. Adansonia*, i. (1860–61) 269.—Ins. Comor.
chinensis, Roxb. Fl. Ind. iii. 677 = australis.
chortiandra, *Baill. Adansonia*, v. (1864–65) 285.—Bras.
ciliata, *Forsk. Fl. Aegypt. Arab.* 162.—Geront. trop.
ciliata, Wall. Cat. n. 7779 J = indica.
cincta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 20.—Mexico.
circinnata, A. Gray, ex Seem. List Vit. Pl. 11 = Wilkesiana.
Clausseni, *Baill. Adansonia*, v. (1864–65) 226.—Bras.
codonocalyx, *Baill. l. c.* i. (1860–61) 271.—Ins. Comor.
colorata, Spreng. Syst. iii. 879 = integrifolia.
Commersoniana, Baill. Adansonia, i. (1860–61) 267 = integrifolia.
communis, Muell. Arg. in Linnaea, xxxiv. (1865–66) 28 = variabilis.
conferta, Roxb. Fl. Ind. iii. 677 = brachystachya.
consimilis, *Muell. Arg. in DC. Prod.* xv. ii. 807.—Ins. Pacif.
conspicua, *Muell. Arg. l. c.* 892.—Mexico.
contermina, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 46.—Peruv.
corchorifolia, *A. Rich. in Sagra, Hist. Cuba*, iii. 208.—Ind. occ.
corchorifolia, Vahl, ex Baill. Etud. Gén. Euphorb. 443 = fallax.
corchorifolia, Willd. Sp. Pl. iv. 524 = reptans.
cordata, Drège, ex Baill. Adansonia, iii. (1863) 158 = Eckloni.
cordata, Thunb. Prod. Pl. Cap. 117 = decumbens.
cordifolia, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 186 = parvula.
cordifolia, *Griseb. in Goett. Abh.* xix. (1874) 97.—Reg. Argent.
cordifolia, Anders. Galap. Veg. 103 = parvula.
cordobensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 46.—Peruv.
corchorifolia, *A. Rich. in Sagra, Hist. Cuba*, iii. 208.—Ind. occ.
corchorifolia, Vahl, ex Baill. Etud. Gén. Euphorb. 443 = fallax.
corchorifolia, Willd. Sp. Pl. iv. 524 = reptans.
cordata, Drège, ex Baill. Adansonia, iii. (1863) 158 = Eckloni.
cordata, Thunb. Prod. Pl. Cap. 117 = decumbens.
cordifolia, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 186 = parvula.
cordifolia, *Griseb. in Goett. Abh.* xix. (1874) 97.—Reg. Argent.
cordifolia, Anders. Galap. Veg. 103 = parvula.
cordobensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 46.—Peruv.
corduvensis, *Muell. Arg. in Journ. Bot.* xii. (1874) 238.—Reg. Argent.
corensis, *Jacq. Enum. Pl. Carib.* 32.—Ind. occ.
crassa, Meisn. ex Krauss, in Flora, xxviii. (1845) 88 = peduncularis.
cremata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 245.—Afr. trop.
crenulata, Rafin. New Fl. Am. I. 44 = virginica.
cucullata, Poir. Encyc. Suppl. iv. 538 = macrostachya.
cuneata, *Poepp. & Endl. Nov. Gen.* iii. 32.—Am. austr.
Cunninghamii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 35.—Austral.
cuspidata, *Jacq. Hort. Schoenb.* ii. 68. t. 243.—Am. austr.
cydnidrifolia, Roxb. Fl. Ind. iii. 678 = Poiretii.
decidua, *Forsk. Fl. Aegypt. Arab.* 161.—Arab.
decumbens, *Thunb. Prod. Pl. Cap.* 117.—Afr. austr.
dentifera, Blume, Bijdr. 628 = hispida.
dentata, Schum. & Thonn. Beskr. Guin. Pl. 410 = Malloius oppositifolius.
denudata, *Muell. Arg. in DC. Prod.* xv. ii. 819.—Ins. Pacif.
depauperata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 190.—Mexico.
Deppeana, Schlecht. in Linnaea, vii. (1832) 585 = Schiedeana.
dictyoneura, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 16.—Peruv.
diffusa, Anders. Galap. Veget. 104 = parvula.
digyneia, *Rafin. Fl. Ludov.* 112.—Louisian.
digynostachya, *Baill. Adansonia*, v. (1864–65) 283.—Bras.
dimorpha, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 354.—Bras.
discolor, Boj. Hort. Maurit. 286 = integrifolia.
discolor, E. Mey Zwei Pfl. Docum. 129 (n. 20); in Flora, xxviii. (1845) 84 = decumbens.
distans, *Muell. Arg. in DC. Prod.* xv. ii. 820.—Cuba.
divaricata, Baill. in Adansonia, v. (1864–65) 284 = gracilis.
divaricata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 34.—Peruv.
divaricata, *Rafin. New. Fl. Am.* i. 44.—Am. bor.
diversifolia, *Jacq. Hort. Schoenb.* iii. 68. t. 244.—Am. austr.
domingensis, Spreng. Syst. Veg. iii. 880 = carpinifolia.
dumetorum, *Muell. Arg. in Journ. Bot.* ii. (1864) 334 —Afr. trop.
Duparquana, Baill. Adansonia, v. (1864–65) 229 = accedens.
echinata, *Rafin. New. Fl. Am.* i. 45.—Am. bor.
Eckloni, *Baill. Adansonia*, iii. (1862–63) 158.—Afr. austr.
elegantula, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 246 = brachystachya.
elliptica, *Sw. Prod. Veg. Ind Occ.* 99.—Jamaica.
emmenesis, *Baill. Adansonia*, i. (1860–61) 270.—Madag.
eremorum, *Muell. Arg. in Flora*, xlvii. (1864) 440.—Austral
erythrostachya, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 31.—N Granat.
estrellana, Baill. Adansonia, v. (1868–64) 287 = brasiliensis.
exaltata, *Baill. l. c.* ii. (1861–62) 225.—N. Caledn
fallax, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 48.—Ax. trop.
fasciculata, *Muell. Arg. l. c.* 81.—Ins. Mascar.
filiformis, Poir. Encyc. vi. 205 = arborea.
filiformis, Klotzsch, ex Schlecht. in Linnaea, xix. (1847) 285 = Schiechtendaliana.
fimbriata, Baill. Adansonia, i. (1860–61) 272 = Baillonia et crenata.
fimbriata, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 245.—Abyss.
fimbriata, Schum. & Thonn. Beskr. Guin. Pl. 409 = ciliata.
firmula, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 21.—Am. centr.
fisso, Herb. Madr. ex. Wall. Cat. n. 7786. B. = brachystachya.
flaccida, Hook. f. in Trans. Linn. Soc. xx. (1847) 186 = parvula.
flexuosa, Wight, ex Stend. Nom. ed. II. i. 9 = lanceolata.
Forsteriana, *Muell. Arg. in DC. Prod.* xv. ii. 807.—Ins. Pacif.
Fournieri, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 162.—Mexico.
Frederici, *Muell. Arg. in DC. Prod.* xv. ii. 828.—Mexico.
fruticosa, *Forsk. Fl. Aegypt. Arab.* 161.—Geront. trop.
fruticulosa, *Rafin. Fl. Ludov.* 112.—Louisiana.
fuscescens, *Muell. Arg. in DC. Prod.* xv. ii. 821.—Afr. trop.
gemina, Spreng. Syst. iii. 880 = australis.
glabrata, *Thunb. Prod. Pl. Cap.* 117.—Afr. austr.
glabrata, Vahl, ex A. Juss. Test. Euph. 105 = Thecacoris madagascariensis.
glandulifolia, Meisn. ex Krauss, in Flora, xxviii. (1845) 88 = peduncularis.
glandulosa, *Blanco, Fl. Filip.* ed. I. 749.—Ins. Philipp.
glandulosa, *Cav. in Anal. Hist. Nat. Madr.* ii. (1800) 141.—Mexico.
girchomanefolia, A. Rich. Fl. Cub. Fanerog. ii. 205 = reptans.
Gouldiana, Baill. Adansonia, i. (1860–61) 268 = arborea.
gracilens, A. Gray, Man. Bot. N. Unit. States, 390 = virginica.
gracilifera, Baill. Adansonia, i. (1860–61) 273 = integrifolia.
gracilis, *Spreng. Syst.* iv; *Cur. Post.* 315.—Bras.
grandidentata, *Muell. Arg. in DC. Prod.* xv. ii. 823.—Afr. austr.
grandifolia, Poir. Encyc. vi. 204 = Claoxylon crassifolium.
grandis, *Benth. in Hook . Lond. Journ. Bot.* ii. (184) 283 —Malaya.
Hartwegiana, Benth. ex Baill. Etud. Gén. Euphorb. 443 = diversifolia.
havanensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 49.—Cuba.
hederacea, *Torr. in Bot. Mex. Bound.* 200.—N. Mexico.
hernandifolia, *Sw. Prod. Veg Ind Occ.* 99.—Jamaica.
heterodonta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 12.—Am. trop.
hirsuta, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 248 = villicaulis.
hirsutissima, Willd. Sp. Pl. iv. 528 = macrostachya.
hirta, Cav. in Anal. Hist. Nat. ii. (1800) 141 = phleoides.
hirta, Spreng. Syst. iv; Cur. Post. 315 = variabilis.
hispida, Blume, Bijdr. 628 = boehmerioides.
hispida, *Burm. f. Fl. Ind.* 203. t. 61.—Ind. or.
hispida, Wall. Cat. n. 7780. pro parte = fallax.
hispida, Willd. Sp. Pl. iv. 532 = Poiretii.
Hochstetteriana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 39.—Afr. trop.
hologyna, *Baker, in Journ. Linn. Soc. xxi.* (1885) 441.—Madag.
Hystrix, Balb. ex Spreng. Syst. iii. 883 = aristata.
indica, *Linn. Sp. Pl.* 1003.—Ax. & Afr. trop.
indica, Vell. Fl. Flum. x. t. 64 = Poiretii.
infesta, *Poepp & Endl. Nov. Gen.* iii. 22.—Peruv.
inflexiana, Muell. Arg. in Linnaea, xxxiv. (1865–66) 22 = prace
insulana, *Muell. Arg. in Flora* (1864) 489.—Ins. Pacif.
integrifolia, *Willd. Sp. Pl.* iv. 530.—Ins. Mascar.
interrupta, Schlecht. in Linnaea, vii. (1832) 586 = Bernardin interrupta.
jamaicensis, *Rafin. New Fl. Am.* 46.—Jamaica.
japonica, Hoult. ex Steud. Nom. ed. I. 4 = Boehmeria spicata.
Jardini, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 36.—Ins. Pacif.
Karwinskii, *Muell. Arg. in Flora*, lv. (1872) 41.—Am. centr.
Klotzschiana, Muell. Arg. in DC. Prod. xv. ii. 857 = prunifolia.
Klotzschii, Baill. Adansonia, v. (1864–65) 231 = longifolia.
Kraussiana, Buching. ex Krauss, in Flora, xxvii. (1845) 84 = decumbens.
lacvifolia, *Muell. Arg. in DC. Prod.* xv. ii. 853.—Ins. Pacif.
laevigata, *Sw. Prod.* 99.—Jamaica.
laevigata, Willd. Sp. Pl iv. 527 = carpinifolia.
Lagascana, *Muell. Arg. in Flora*, lv. (1872) 27.—Am. centr.
lagoensis, *Muell Arg. in Mart Fl. Bras.* xi. ii. 367.—Bras.
lamnifolia, *Scheele, in Linnaea*, xxv. (1852) 587.—Afr. austr.
lanceolata, Wall. Cat. n. 7789 = australis.
lanceolata, *Willd. Sp. Pl.* iv. 524.—Ind. or.
Langiana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 159.— Mexico.
languida, E. Mey. & Sond. in Linnaea, xxiii. (1850) 116 = petiolaris.
laugnida, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 39.—Afr. austr.
lantanaefolia, Boj. Hort. Maurit. 286 = reticulata.
latifolia, *Muell. Arg. in DC. Prod.* xv. ii. 817.—Ins. Pacif.
latifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 19.—Ind. occ.
lecuensis, *Benth. in Hook. Nig. Fl.* 504 = Mareya micrantha.
Leptinei, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 14.—Ins. Pacif.
leptoclada, *Benth. in Bot. Voy. Sulph.* 164.—Mexico.
leptopoda, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 29.—Mexico.
leptorhachis, *Muell. Arg. l. c.* 7.—Cuba.
leptostachya, A. Rich. in Sagra, Hist. Cuba, iii. 205 = laxiflora.
lepostachya, H. B. & K. Nov. Gen et Sp. ii. 96 = diversifolia.
Liebmanni, Muell. Arg. in DC. Prod. xv. ii. 829 = seq.

ACALYPHA:—
Liebmanniana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 161.—Mexico.
Lindeniana, *Muell. Arg. in DC. Prod.* xv. ii. 827.—Mexico.
Lindheimeri, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 47.—Mexico.
Linostachys, Baill. Adansonia, v. (1864–65) 235 = villosa.
Livingstoniana, *Muell. Arg. in Flora*, xlvii. (1864) 440. —Afr. trop.
longestipularis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 51.—Mexico.
longifolia, Baill. Etud. Gen. Euph. 448 = cuneata.
longifolia, *Klotzsch, ex Baill. Adansonia*, v. (1864–65) 231.—Bras.
longispicata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 152.—Mexico.
Lyallii, *Baker, in Journ. Linn. Soc.* xx. (1884) 255 — Madag.
macrodonta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 51.—Peruv.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 96 = macrostachya.
macrosperma, Muell. Arg. in Linnaea, xxxiv. (1865–66) 18 = membranacea.
macrostachya, *Jacq. Hort. Schoenb.* ii. 68. t. 245.—Am. austr.
macrostachyoa, Poir. Encyc. vi. 208 = Poiretii.
macrostachyoides, *Muell. Arg. in DC. Prod.* xv. ii. 809. —Mexico.
major, Baill. Adansonia, v. (1864–65) 236 = Wedelliana.
malabarica, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 48.—Ind. or.
Mandoni, *Muell. Arg. l. c.* 162.—Bolivia.
Manniana, *Muell. Arg. in Flora*, xlvii. (1864) 441.—Afr. occ.
Meppe, Wild. Sp. Pl. iv. 526 = Macaranga Mappa.
marginata, *Spreng. Syst.* iii. 879.—Ind. or.
Martiana, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 359.—Bras.
melochiefolia, *Muell. Arg. in DC. Prod.* xv. ii. 821. —Mexico.
membranacea, *A. Rich. Fl. Cub. Fanerog.* ii. 204.—Cuba.
mexicana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 41.—Mexico.
micrantha, Benth. in Hook. Nig. Fl. 505 = Mareya micrantha.
microcephala, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 150.—Mexico.
microgyne, Poepp. & Endl. Nov. Gen iii. 21 = diversifolia.
microphylla, Klotzsch, in Seem. Bot. Voy. Herald, 278 = reptana.
microstachya, Benth. Pl. Hartw. 71 = mollis.
mollis, *H. B. & K. Nov. Gen. et. Sp.* ii. 94.—Mexico.
monococca, Engelm. ex A. Gray, Man. Bot. N. U. States, 390 = virginica.
monostachya, Benth. Pl. Hartw. 15 = anemioides.
monostachya, *Cav. in Anal. Hist. Nat. Madr.* ii. (1800) 188.—Mexico.
multicalis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 58.—Bras.
multipartita, Moench, Meth. Suppl. 122 = tomentosa.
muralis, *Zipp. ex Span. in Linnaea*, xv. (1841) 850, nomen.—Timor.
nana, Griseb. in Goett. Nachr. (1865) 176 = reptans.
nemorum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 36.—Austral.
neocaledonica, *Muell. Arg. in DC. Prod.* xv. ii. 812.—N. Caled.
neogranatensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 15.—N. Granat.
neomexicana, *Muell. Arg. l. c.* 19.—N. Mexico.
neptunica, *Muell. Arg. in Bremen Abh.* vii. (1880) 26. —Afr. trop.
nigritiana, *Muell. Arg. in Flora*, xlvii. (1864) 440.—Afr. trop.
nivea, Benth. in Bot. Voy. Sulph. 163. t. 53 = cuneata.
obscura, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 163. —Mexico.
obtusa, Thunb. Fl. Cap. ed. Schalt. 546 = Leidesia obtusa.
obtusata, Spreng. ex Steud. Nom. ed. II. i. 10. [Adenocline, sp.].—Afr. austr.
ocymoides, *H. B. & K. Nov. Gen. et. Sp.* ii. 93.—Mexico.
odorata, Steud. Nom. ed. II. ii. 696 sphalm. = integrifolia.

ACALYPHA:—
oliganthe, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 159.—Mexico.
oligodonta, *Muell. Arg. in DC. Prod.* xv. ii. 831.—Mexico.
ornata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 247.—Abyss.
ostryæfolia, Riddell, Syn. Fl. W. St. 83 = caroliniana.
ovalifolia, Baill. Adansonia, i. (1860–61) 269 = arborea.
oxyodonta, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 367. —Mexico.
pacifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 97.—Am. austr.
panamensis, Klotzsch, in Seem. Bot. Voy. Herald, 101 = diversifolia.
Pancheriana, *Baill. Adansonia*, ii. (1861–62) 225.—N. Caled.
pascuicola, *Miq. Fl. Ind. Bat.* i. ii. 406. (1859).—Geront. trop.
parvifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 161.—Mexico.
parvula, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 185. —Ins. Galap.
Pastoris, DC. ex Willd. Enum. Hort. Berol. 993 = Abyss.
Pastoris, Schrank, in Denkschr. Bayer. Ges. ii. (1822) 39 = phleoides.
patens, *Muell. Arg. in DC. Prod.* xv. ii. 847.—Afr. austr.
pauciflora, Hornem. Hort. Hafn. ii. 909 = australis.
Pavoniana, Muell. Arg. in Linnaea, xxxiv. (1865–66) 50 = arvensis.
Peckoltii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 365. —Bras.
peduncularis, E. Mey. Zwei Pfl. Docum. 186 n. 26 = Zeyheri.
peduncularis, *Meisn. ex Krauss, in Flora*, xxviii. (1845) 52.—Afr. austr.
pendula, Wright, ex Griseb. in Goett. Nachr. (1865) 176 = reptans.
pensilifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 25.—Ind. occ.
peruviana, *Muell. Arg. l. c.* 17.—Peruv.
Perrottetiana, Baill. Adansonia, i. (1860–61) 278 = arborea.
petiolaris, *Krauss, in Flora*, xxviii. (1845) 53.—Afr. austr.
petiolaris, Sond. in Linnaea, xxiii. (1850) 117 = Sonderiana.
phleoides, *Cav. in Anal. Hist. Nat. Madr.* ii. (1800) 189.—Mexico.
phleoides, Torr. Bot. Mex. Bound. Surv. 199 = Lindheimeri.
pilifera, Klotzsch, ex Baill. Adansonia, v. (1860–61) 227 = variabilis.
pilosa, *Cav. in Anal. Nat. Hist. Madr.* ii. (1800) 186.—Am. trop.; Ins. Philipp.
pinnata, Poir. Encyc. vi. 205 = Tragia pinnata.
platyphylla, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 6.—Peruv.
plicata, *Muell. Arg. in DC. Prod.* xv. ii. 858.—Bolivia.
Poiliana, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 360.—Bras.
Poiteni, *Spreng. Syst.* iii. 879.—Am.-trop.
polyorchis, *Muell. Arg. in Journ. Bot.* ii. (1864) 335.—Afr. trop.
polystachya, *Jacq. Hort. Schoenb.* iii. 64. t. 346.—Peruv.; Mexico.
popayanensis, H. B. & K. Nov. Gen. et Sp. vii. 173 = diversifolia.
portoricensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 22.—Ind. occ.
Pringlei, *S. Wats. in Proc. Am. Acad.* xx. (1885) 378. —Am. bor.
pruriefolia, H. B. & K. Nov. Gen. et Sp. ii. 92 = phleoides.
prurifolia, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 97.—Bras.
pruriens, *Nees & Mart. l. c.* 96.—Bras.
pullostachya, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 246.—Abyss.
pubiflora, *Baill. Adansonia*, i. (1860–61) 266.—Afr. trop.; Madag.
punctata, Meisn. ex Krauss, in Flora, xxvii. (1845) 53 = peduncularis.
purpurascens, *H. B. & K. Nov. Gen. et. Sp.* ii. 97.—Mexico.
pygmæa, A. Rich. Fl. Cub. Fanerog. ii. 205 = reptans.
racemosa, Heyne, in Wall. Cat. n. 7784 = paniculata.

ACALYPHA:—
radians, *Torr. in Bot. Mex. Bound.* 200.—N. Mexico.
radicans, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 33.—Bras.
Rætsia, *Baker, in Journ. Linn. Soc.* (1884) 254.—Madag.
reflexa, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 33. —Peruv.
remiformis, Hook. f. in Trans. Linn. Soc xx. (1847) 187 = parvula.
repanda, *Muell. Arg. in Flora*, xlvii. (1864) 439.—Ins. Pacif.
reptans, *Sw. Prod. Veg. Ind. Occ.* 99.—Ind. occ.
reticulata, Muell. Arg. in DC. Prod. xv. ii. 851 = arborea.
rhombifolia, Baill. Adansonia, v. (1864–65) 230 = Poiretii.
rhombifolia, *Schlecht. in Linnaea*, vii. (1832) 382.—Mexico.
rhomboides, *Raffin. New Fl. Am.* i. 45.—Am. bor.
Richardiana, *Baill. in Adansonia*, i. (1860–61) 268.—Madag.
Riedeliana, *Baill. l. c.* v. (1864–65) 231.—Bras.
rivularis, *Seem. List Vit. Pl.* 11.—Ins. Fiji.
rottleroides, *Baill. Adansonia*, i. (1860–61) 270.—Madag.
rotundifolia, Vahl, ex Baill. Etud. Gén. Euphorb. 442 = indica.
rubra, Roxb. in Beatson, Tracts St. Helena Append. 295 = arborea.
rubra, Wight, ex Wall. Cat. n. 7781 = ciliata.
rubra, Willd. Enum. Hort. Berol. 993 = phleoides.
Ruiziana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 16.—Peruv.
salicifolia, *Muell. Arg. in Flora*, xlvii. (1864) 438.—Peruv.
salviæfolia, *Baill. Etud. Gén. Euphorb.* 448, nomen.—Ins. Mascar. ?
samydæfolia, *Poepp. & Endl. Nov. Gen.* iii. 21. t. 264.—Peruv.
scabra, Vahl, ex Baill. Adansonia, i. (1860–61) 280 = Claoxylon parviflorum.
scabrosa, *Sw. Prod. Veg. Ind. Occ.* 100.—Jamaica.
scandens, *Benth. in Hook. Kew Journ.* vi. (1854) 329. —Am. trop.
Schiedeana, *Schlecht. in Linnaea*, vii. (1832) 384.—Mexico.
Schlechtendaliana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 6.—Mexico.
Schlumbergeri, *Muell. Arg. in DC. Prod.* xv. ii. 861.—Mexico.
Seemanni, Klotzsch, in Seem. Bot. Voy. Herald, 102 = macrostachya.
segetalis, *Muell. Arg. in Journ. Bot.* ii. (1864) 336. Afr. occ.
semiinuda, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 360.—Bras.
senensis, *Klotzsch. in Peters, Reise Mossamb. Bot.* 96. —Afr. or.
sentilis, *Baill. Adansonia*, v. (1864–65) 228.—Am. austr.
septemloba, *Muell. Arg. in Flora*, lv. (1872) 37.—Amer. centr.
serrata, Andrew, Galapag. Veget. 103 = parvula.
sessilis, Poir. Encyc. vi. 204 = australis.
sidræa, *A. Rich. Fl. Cub. Fanerog.* ii. 204.—Cuba.
silenifolia, *A. Rich. Tent. Fl. Abyss.* ii. 249.—Abyss.
sidæfolia, H. B. & K. Nov. Gen. et Sp. ii. 96 = macrostachya.
somalicum, Muell. Arg. in Bremen Abh. vii. (1880) 37 = segetalis.
Sondeziana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 9.—Afr. austr.
Spachiana, *Baill. Adansonia*, i. (1860–61) 272.—Madag.
spicata, Andrew, Galap. Vegt. 104 = parvula.
spicata, Forsk. Fl. Aegypt. Arab. 161 = indica.
spicifera, *Burm. f. Fl. Ind.* 208 (t. c. 208), t. 61.—Madag.
spicifera, Thou. ex Baill. Adansonia, i. (1860–61) 281 = Claoxylon Linostachys.
spiciflora, Poir. Encyc. vi. 206 = Claoxylon parviflorum.
spicigera, Klotzsch, in Seem. Bot. Voy. Herald, 101 = diversifolia.
spinescens, *Brash. in Hook. Ic. Pl.* t. 1292.—Malaya.
stenoloba, *Muell. Arg. in Flora*, lv. (1872) 41.—Peruv.
stipulacea, *Klotzsch, in Nov. Act. Acad. Nat. Cur.* xix. (1843) Suppl. I. 618.—Ins. Mexico.

ACALYPHA,—
stricta, Poepp. & Endl. Nov. Gen. iii. 21. t. 225.—Peruv.
strobilifera, Hook. f. in Trans. Linn. Soc. xx. (1847) 187 = parvula.
subconnexa, *Lag. Elench. Hort. Reg. Matr.* (1806); *Gen. et Sp. Nov.* 21.—Mexico.
subvillosa, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 341.—Bras.
supera, Forsk. Fl. Aegypt. Arab. 162 = brachystachya.
tarapotensis, *Muell. Arg. in DC. Prod.* xv. ii. 808.—Peruv.
tenuicaulis, Baill. Adansonia, v. (1864–65) 234 = multicaulis.
tenuifolia, *Muell. Arg. in DC. Prod.* xv. ii. 862.—Venezuela.
tenuiramea, *Muell. Arg. l. c.* 858.—Ins. Pacif.
tenuis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 30.—Afr. austr.
tiliaefolia, Poir. Encyc. vi. 208. (An Euphorbiacea !).—Ins. S. Doming.
tomentosa, *Blanco, Fl. Filip.* ed. I. 750.—Ins. Philipp.
tomentosa, Boj. Hort. Maurit. 285 = marginata.
tomentosa, *Sw. Prod. Veg. Ind. Occ.* 99.—Ins. S. Domingo.
trachyloba, *Muell. Arg. in Flora*, lv. (1872) 25.—Mexico.
tricholoba, *Muell. Arg. in Linnaea*, xxxiv. (1865–66). —Mexico.
tricolor, *Seem. Fl. Vit.* 225.—Ins. Pacif.
triloba, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 28. —Mexico.
tristis, Poepp. & Endl. Nov. Gen. et Sp. iii. 22 = macrostachya.
ulmifolia, Benth. Pl. Hartw. 259 = diversifolia.
unibracteata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 160.—Mexico.
urophylla, *Boiv. ex Baill. Adansonia*, i. (1860–61) 273.—Afr. trop.
urostachya, *Baill. l. c.* v. (1864–65) 229.—Bras.
urticaefolia, *Poir. Encyc.* vi. 206.—Am. austr.
urticoides, Klotzsch, ex Baill. Adansonia, v. (1864–65) 227 = communis.
vagans, *Cav. in Anal. Hist. Nat. Madr.* ii. (1800) 139.—Mexico.
Vahliana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 43.—Afr. trop.
variabilis, *Klotzsch, ex Baill. Adansonia*, v. (1864–65) 226.—Bras.
Vedeliana, Baill. Adansonia, ii. (1861–62) 224 = Macaranga Vedeliana.
velamea, *Baill. l. c.* v. (1864–65) 228 = Bras.
velutina, Hook. f. in Trans. Linn. Soc. xx. (1847) 186 = parvula.
venosa, Poir. Encyc. vi. 204 = Leptonema venosum.
ventita, Benth. in Bot. Voy. Sulph. 164 = cuspidata.
villicaulis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 248.—Abyss.
villosa, *Jacq. Enum. Pl. Carib.* 32.—Ind. occ.
villosa, Vahl, ex Baill. Etud. Gén. Euphorb. 442 = Poiretii.
virgata, Forst. f. Prod. 67 = Forsteriana.
virgata, *Linn. Pugill. Jamaic. Pl.* 28 (*Am. Acad.* v. 410).—Jamaica.
virgata, Vell. Fl. Flum. x. t. 68 = variabilis.
virgata, Thunb. Fl. Jap. 288 = australis.
virginica, *Linn. Sp. Pl.* 1003.—Am. bor.
virginica, Wall. Cat. n. 7779 G = fallax.
Wallichiana, Muell. Arg. in DC. Prod. xv. ii. 802 = acuminata.
Wallichii, Thw. Enum. Pl. Zeyl. 271 = paniculata.
Weddelliana, Baill. Adansonia, v. (1864–65) 232 = brasiliensis.
Welwitschiana, *Muell. Arg. in Journ. Bot.* ii. (1864) 324.—Afr. trop.
Wightiana, Muell. Arg. in Linnaea, xxxiv. (1865–66) 43 = lanceolata.
Wilkesiana, Muell. Arg. in DC. Prod. xv. ii. 817 = triodior.
wimbensis, *Muell. Arg. in Flora*, xlvii. (1864) 440.—Afr. trop.
Zeyheri, *Baill. Adansonia*, iii. (1862–63) 156.—Afr. austr.
zeylanica, *Raffn. New Fl. Amer.* i. 46, nomen.—Zeylan.
Zollingeri, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 46.—Sumbawa.

ACAMPE, Lindl. Fol. Orch. (1858).
ORCHIDEAE, Benth. & Hook. f. iii. 579.
Cephalotes, *Lindl. l. c.* 3.—Ind. or.
congener, *Lindl. l. c.* 3.—Ind. or.
dentata, *Lindl. l. c.* 3.—Ind. or.
excavata, *Lindl. l. c.* 3.—Ind. or.
Griffithii, *Reichb. f. in Flora*, lv. (1872) 277.—Ind. or.
intermedia, *Reichb. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 217.—Hab. ?
longifolia, *Lindl. Fol. Orchid. Acampe*, 1.—Ind. or.
multiflora, *Lindl. l. c.*—China.
pachyglossa, *Reichb. f. Otia Bot. Hamb.* 76; *et in Bot. Zeit.* xxxix. (1881) 449.—Afr. trop.
papillosa, *Lindl. Fol. Orch. Acampe*, 2.—Ind. or.
Kenschiana, *Reichb. f. Otia Bot. Hamb.* 77; *et in Bot. Zeit.* xxxix. (1881) 449.—Madag.
Wightiana, *Lindl. Fol. Orch. Acampe*, 2.—Ind. or.

ACAMPTOPAPPUS, A. Gray, in Proc. Am. Acad. viii. (1873) 634 = **Aplopappus**, Cass.—(Compos.).
Skockleyi, A. Gray, l. c. xvii. (1881–82) 208.—Am. bor.
sphaerocephalus, A. Gray, l. c. viii. (1873) 634.—Am. bor.

ACANOS, Adans. Fam. ii. 116 (1768) = **Onopordon**, Linn. (Compos.).
Spina, Scop. Fl. Carn. ed. II. ii. 132 = O. Acanthium.

ACANTACARYX, Arruda, ex Koster, Trav. Bras. 491 (1816) = **Caryocar**. Linn. (Ternstroem.).
pinguis, Arruda, ex Koster, l. c.—Bras.

ACANTHELLA, Hook. f. in Benth. & Hook. f. Gen. i. 748 (1867).—*MELASTOMACEAE*, Benth. & Hook. f. i. 746.
Spruceii, *Hook. f. l. c.*—Bras.

ACANTHINOPHYLLUM, Fr. Allem. in Rev. Brazil. i. (1858) 368 = **Sahagunia**, Liebm. (Urtic.).
strepitans, Fr. Allem. l. c. 369.—Bras.

ACANTHIUM, Hall. Enum. Stirp. Helv. ii. 674 (1742) = **Onopordon**, Linn. (Compos.).
Onopordon, Gueldenst. Reise Russl. ii 69 = O. Acanthium.

ACANTHOBOTRYA, Eckl. & Zeyh. Enum. 192 (1836) = **Lebeckia**, Thunb. (Legumin.).
angustifolia, Eckl. & Zeyh. l. c. 194 = L. multiflora.
armata, Eckl. & Zeyh. l. c. 193 = L. microphylla.
cinerea, Eckl. & Zeyh. l. c. 193 = L. microphylla.
decumbens, Eckl. & Zeyh. l. c. 193 = L. humilis.
distincta, Eckl. & Zeyh. l. c. 193 = L. distincta.
linearifolia, Benth. in Ann. Wien. Mus. ii. (1838) 142 = L. linearifolia.
microphylla, Benth. l. c. = L. microphylla.
pungens, Eckl. & Zeyh. Enum. 193 = L. pungens.
sericea, Eckl. & Zeyh. l. c. 193 = L. microphylla.
sessilifolia, Eckl. & Zeyh. l. c. 193 = L. marginata.

ACANTHOCARPEA, Klotzsch, in Peters, Reise Mossamb. Bot. 137. t. 24 (1861) = Ldmeum, Linn. (Ficoid.)
scabrida, Klotzsch, l. c. 139.—Mossamb.
sessilis, Klotzsch, l. c. 138. t. 24 = L. linifolium.

ACANTHOCARPUS, Lehm. Pl. Preiss. ii. 274 (1847). *JUNCEAE*, Benth. & Hook. f. iii. 664.
fimbriatus, *F. Muell. Census*, 119.—Austral.
Preissii, *Lehm. Pl. Preiss.* ii. 274.—Austral.
Serra, *F. Muell. Census*, 119.—Austral.

ACANTHOCARYA, Arruda, ex Endl. Gen. 1076 (1840) = **Caryocar**, Linn. (Ternstroem.).

ACANTHOCAULON, Klotzsch, in Endl. Gen. Suppl. iv. iii. 88 (1886) = **Platygyna**, Mercier (Euphorb.).
prunioni, Klotzsch (sphalm. pruricus), ex Endl. l. c. = P. urens.

ACANTHOCEPHALUS, Kar. & Klr. in Bull. Soc. Nat. Mosc. xv. (1842) 137. *COMPOSITAE*, Benth. & Hook. f. ii. 510.
HARPACHAENA, Bunge, Del. Sem. Hort. Dorp. (1845).
HARPOCARPUS, Endl. Gen. Suppl. iii. 10 (1843).
amplexifolius, *Kar. & Klr. in Bull. Soc. Nat. Mosc.* xv. (1842) 138.—Sibir. alsaic.
Besthamianus, *Regel, Pl. Nov. Fedsch.* 47.—Turkestan.

ACANTHOCHITON, Torr. in Sitgr. Rep. 170. t. 18 (1853). *AMARANTACEAE*, Benth. & Hook. f. iii. 29.
Wrightii, *Torr. l. c.*—N. Mexico.

ACANTHOCLADIUM, F. Muell. Fragm. ii. 155 (1861) = **Hallothrysum**, Wall. (Compos.).
Dockerii, F. Muell. l. c. 156 = H. Dockerii.

ACANTHOCLADUS, Klotzsch, ex Hassk. in Miq. Ann. Mus. Lugd. Bat. i. 184 (1864). *POLYGALEAE*, Benth. & Hook. f. i. 974.
albicans, *A. W. Benn. in Mart. Fl. Bras.* xiii. iii. 46. —Bras.
brasiliensis, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Lugd. Bat.* i. 186.—Bras.
microphylla, *Griseb. in Gott. Abh.* xxiv. (1879) 24.—Reg. Argent.

ACANTHODIUM, Delile, Fl. Egypte, 241. t. 33 (1812) = **Blepharis**, Juss. (Acanth.).
angustum, Nees, in DC. Prod. xi. 278.—Afr. austr.
capense, Nees, in Linnaea, xv. (1841) 361.—Afr. austr.
cardiifolium, Nees, in DC. Prod. xi. 278 = B. cardiifolia.
Delilii, Buek, Ind. DC. Prod. iii. 1 = B. edulis.
dispermum, E. Mey. ex Nees, in DC. Prod. xi. 278 = Acanthopsis disperma.
diversispinum, Nees, l. c. 275.—Afr. austr.
furcatum, Nees, l. c. 276.—Afr. austr.
glabrum, Nees, l. c. 278.—Afr. austr.
glaucum, Nees, l. c. 277.—Afr. austr.
grosum, Nees, l. c. 274.—Afr. austr.
grosum, Wight, Ic. t. 1855, 1856 = B. scindica.
hirtinervium, Nees, in DC. Prod. xi. 277.—Afr. austr.
hirtum, Hochst. ex Nees, l. c. 276 = B. scindica.
Hoffmannseggianum, Nees, l. c. xi. 277.—Afr. austr.
macrum, Nees, l. c. 276.—Afr. austr.
marginatum, Nees, l. c. 275.—Afr. austr.
plumulosum, E. Mey. ex Nees, l. c. 278 = Acanthopsis horrida.
procumbens, Nees, in Linnaea, xv. (1841) 362.—Afr. austr.
serrulatum, Nees, in DC. Prod. xi. 276.—Afr. austr.
sinuatum, Nees, in Linnaea, xv. (1841) 362.—Afr. austr.
spathulare, Nees, in DC. Prod. xi. 277.—Afr. austr.
spicatum, Acerbi, ex Nees, l. c. 67 = Nelsonia canescens.
spicatum, Delile, Fl. Egypte, 241. t. 33. f. 3 = B. edulis.
squarrosum, Nees, in DC. Prod. xi. 275.—Afr. austr.

ACANTHODUS, Rafin. Princ. Somiol. 26 (1814) = **Acanthus**, Tourn. (Acanthac.).

ACANTHOGLOSSUM, Blume, Bijdr. 881 (1825) = **Pholidota**, Lindl. (Orchid.).
nervosum, Blume, l. c. 382 = P. nervosa.

ACANTHOGONUM, Torr. Pacif. Rail. Rep. iv. 132 (1856) = **Chorisanthe**, R. Br. (Polygon.).
corrugatum, Torr. & Gray, l. c. v. 364 = C. corrugata.
rigidum, Torr. l. c. iv. 133 = C. rigida.

ACANTHOLEPIS, Less. in Linnaea, vi. (1831) 88. *COMPOSITAE*, Benth. & Hook. f. ii. 463.
orientalis, *Less. l. c.*—Ala. Min.; Persia.

ACANTHOLINUM, C. Koch, in Linnaea, xxi. (1848) 715, sphalm. = seq.

ACANTHOLIMON, Boiss. Diagn. Ser. I. vii. 89 (1846). *PLUMBAGINEAE*, Benth. & Hook. f. ii. 625.
acerosum, *Boiss. l. c.* Ser. I. vii. 90.—As. Min.
acmostegium, *Boiss. & Bukse, in Nouv. Mém. Mosc.* xii. (1860) 188.—Persia.
alatavicum, *Bunge, Gatt. Acanthol.* 40.—Sibir.
Alberti, *Regel, in Act. Hort. Petrop.* vi. (1879) 389.—Turkestan.
androsaceum, *Boiss. Diagn.* Ser. I. vii. 73.—Oriens.
araxanum, *Bunge, Gatt. Acanthol.* 38.—Persia.
arietalatum, *Bunge, l. c.* 90.—Persia.
armenum, *Boiss. & Huet, Diagn.* Ser. II. iv. 64.—Armenia.
aspadanum, *Bunge, Gatt. Acanthol.* 50.—Persia.
asyriacum, Boiss. Diagn. Ser. I. vii. 91 = venustum.

ACANTHOLIMON:—
atropatanum, *Bunge, Gatt. Acanthol.* 29.—Persia.
auganum, *Bunge, l. c.*—Aghlan.
avenaceum, *Bunge, l. c.* 25.—Persia.
Balansae, Boiss. ex Bunge, l. c. 36 = armenum.
baltanense, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 895.—As. Min.
Bodesanum, *Bunge, Gatt. Acanthol.* 42.—Persia.
brachyphyllum, *Boiss. in DC. Prod.* xii. 628.—Afghan.
brachystachyum, *Boiss. ex Bunge, Gatt. Acanthol.* 51.—Kurdistan.
bracteatum, *Boiss. Diagn.* Ser. I. vii. 70.—Armen. ; Persia.
brevicapum, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 840 = Kotschyi.
bromifolium, *Boiss. ex Bunge, Gatt. Acanthol.* 23.—Kurdistan.
cabulicum, *Boiss. in DC. Prod.* xii. 623.—Afghan.
caesareum, *Boiss. & Bal. Diagn.* Ser. II. iv. 66.—Cappadoc
calocephalum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1881) 77.—Afghan.
Calveri, *Boiss. Diagn.* Ser. II. iv. 65.—Armenia.
caryophyllaceum, *Boiss. Diagn.* Ser. I. vii. 78.—Kurdistan.
caryophyllaceum, Haussk. ex Bunge, Gatt. Acanthol. 37 = armenum.
catacnemum, Bunge, l. c. 48 = lycaonicum.
cephalotum, St. Lag. in Ann. Soc. Bot. Lyon. vii. (1880) 118 = A. Cephalotes.
Cephalotes, *Boiss. in DC. Prod.* xii. 622.—Afghan.
curviflorum, *Bunge, Gatt. Acanthol.* 57.—Persia.
cynosum, *Bunge, l. c.* 15.—Persia.
desertorum, *Regel, in Act. Hort. Petrop.* vi. (1879) 391.—Turkestan.
dispensoides, *Boiss. in DC. Prod.* xii. 624.—Afghan.
dustachyum, *Boiss. l. c.* 628.—Afghan.
Echinus, Bunge, Gatt. Acanthol. 46 = androsaceum.
erythraceum, *Bunge, in Act. Hort. Petrop.* iii. (1875) 99.—Turkestan.
eschkerense, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 844.—Persia.
fasciculare, *Boiss. in DC. Prod.* xii. 626.—Afghan.
Fausti, *Trautv. in Act. Hort. Petrop.* ii. (1873) 482.—Persia.
ferox, Boiss. Diagn. Ser. I. vii. 81 = A. Scorpius.
Fetschei, *Regel, Dec. Pl. Nov.* 8.—Turkest.
foetacearum, *Boiss. Diagn.* Ser. I. vii. 76.—Persia.
flexuosum, *Boiss. ex Bunge, Gatt. Acanthol.* 51.—Persia.
genistoides, *Boiss. Diagn.* Ser. I. vii. 76.—Persia.
glumaceum, *Boiss, l. c.* 75.—Armenia.
Griffithianum, *Boiss. in DC. Prod.* xii. 628.—Afghan.
gulistanum, *Bunge, Gatt. Acanthol.* 67.—Persia.
Haussknechti, Bunge, l. c. 87 = armenum.
herstense, *Bunge, l. c.* 55.—Afghan.
Hohenackeri, *Boiss. Diagn.* Ser. I. vii. 75.—Reg. Caucas.
Hohenackeri, Ledeb. Fl. Ross. iii. 470 = glumaceum.
horridum, *Bunge, Gatt. Acanthol.* 54.—Persia.
Huetii, *Boiss. Diagn.* Ser. II. iv. 65.—Armenia.
Hystrix, *Steqf, in Denkschr. Acad. Wien*, i. (1885) 71.—Persia.
iconicum, *Boiss & Heldr. ex Boiss. Fl. Orient.* iv. 839. —As. Min.
incomptum, Boiss. & Buhse, in Nouv. Mém. Mosc. xii. (1860) 183 = A. Scorpius.
Karelinii, *Bunge, Gatt. Acanthol.* 58.—Armenia.
kokandense, *Bunge, in Act. Hort. Petrop.* iii. (1875) 99.—Turkest.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 74.—Cilicia.
luridum, Bunge, Gatt. Acanthol. 52 = brachystachyum.
latifolium, *Boiss. Diagn.* Ser. II. iv. 61.—Kurdistan.
latifolium, Rupr. Ser. Thianach. 89 = Ruprechtii.
laxiflorum, Boiss. Fl. Orient. iv. 833 = venustum.
laxiflorum, *Boiss. ex Bunge, Gatt. Acanthol.* 38.—Syria.
leptostachyum, *Aitch. & Hemsl in Journ. Linn. Soc.* xviii. (1881) 76.—Afghan.
lepturoides, *Boiss. Diagn.* Ser. I. vii. 77 —Reg. Caucas.
leucacanthum, Boiss. l. c. 81 = A. Scorpius.
libanoticum, *Boiss. in DC. Prod.* xii. 630.—Syria.
Listoniae, Boiss. l. c. 631 = sacrosum.
longiflorum, *Boiss. Diagn.* Ser. II. iv. 62.—Afghan.
lycaonicum, *Boiss. & Heldr. ex Boiss. in DC. Prod.* xii. 637.—As. Min.

ACANTHOLIMON:—
lycopodioides, *Boiss. in DC. Prod.* xii. 632.—As. centr. ; Kashmir.
Maewalkianum, *Regel, in Act. Hort. Petrop.* vi. (1879) 391.—Samarkand.
melananthum, *Boiss. Diagn.* Ser. I. vii. 74.—Persia.
Munroanum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1881) 76.—Afghan.
oliganthum, *Boiss. Diagn.* Ser. I. vii. 77.—Persia.
Olivieri, *Boiss. l. c.* 80.—Persia.
parviflorum, *Regel, Pl. Nov Fedsch.* 73.—As. temp.
Peronii, *Boiss. Fl. Orient.* iv. 842.—Cilicia.
petraeum, *Boiss. ex Bunge, Gatt. Acanthol.* 28.—Kurdistan.
phrygium, Boiss. Diagn. Ser. I. vii. 79 = acerosum.
Pinardi, Boiss. l. c. = acerosum.
polystachyum, *Boiss. in DC. Prod.* xii. 624.—Afghan.
pterostegium, *Bunge, Gatt. Acanthol.* 15.—Persia.
puberulum, Boiss. & Bal. Diagn. Ser. II. iv. 62 = androsaceum.
quinquelobum, *Bunge, Gatt. Acanthol.* 56.—Persia.
restiaceum, *Bunge, l. c.* 22.—Persia.
roseum, *Boiss. Diagn.* Ser. I. vii. 71.—Persia ; Afghan.
rubellum, Boiss. ex Bunge, Gatt. Acanthol. 66 = roseum.
Ruprechti, *Bunge, Gatt. Acanthol.* 20.—Turkestan.
sabendicum, *Boiss. & Buhse, in Nouv. Mém. Mosc.* xii. (1860) 183.—Persia.
sarawschanicum, *Regel, Pl. Nov. Fedsch.* 73.—As. temp.
scabrellum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 845.—Persia.
schahrudicum, *Bunge, Gatt. Acanthol.* 65.—Persia.
schirazianum, *Boiss. Diagn.* Ser. I. vii. 72.—Persia.
scirpinum, *Bunge, Gatt. Acanthol.* 22.—Persia.
scorpioidrum, St. Lag. in Ann. Soc. Bot. Lyon. vii. (1880) 118 = A. Scorpius.
Scorpius, *Boiss. Diagn.* Ser. I. vii. 81.—Persia.
sengananse, *Bunge, Gatt. Acanthol.* 29.—Persia.
setiferum, *Bunge, l. c.* 68.—Turkestan.
spiculosum, Bunge, l. c. 17 = bracteatum.
Stocksii, *Boiss. Diagn.* Ser. II. iv. p. 63.—Beluchistan.
subulatum, *Boiss. in DC. Prod.* xii. 629.—Afghan.
Szovitsii, Boiss. & Buhse, in Nouv. Mém. Mosc. xii. (1860) 184 = Karelinii.
talagonicum, *Boiss. in DC. Prod.* xii. 625.—Persia.
tataricum, *Boiss. l. c.* 629.—Turkestan.
Tchihatcheffi, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. (1854) i. 34 = acerosum.
tenuiflorum, *Boiss Diagn.* Ser. I. vii. 78.—Trans Caucas.
tibeticum, Hook. f. & Thoms. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 479 = lycopodioides.
tomentellum, *Boiss. Diagn.* Ser. I. vii. 73.—Persia.
Tournefortii, *Boiss. l. c.* 74.—Armenia.
tragacanthinum, *Boiss. l. c.* 81.—Persia.
tragacanthinum, Griff. ex Bunge, Gatt. Acanthol. 64 = fasciculare.
tremestum, *Bunge, l. c.* 56.—Persia.
ulicinum, *Boiss. in DC. Prod.* xii. 627.—Syria.
venustum, *Boiss. Diagn.* Ser. I. vii. 80.—As. Min.
viscidulum, *Boiss. l. c.* 71.—Persia.
Wiedemanni, *Bunge, Gatt. Acanthol.* 45.—Anatolia.

ACANTHOLIPPIA, Griseb. in Goett. Abh. xix. (1874) 244 = *Lippia*, Linn. (Verben.).
hastulata, Griseb. l. c. xxiv. (1879) 279 = L. hastulata.
salsoloides, Griseb. l. c. xix. (1874) 244.—Reg. Argent.

ACANTHOLOMA, Gandich. ex Baill. Adansonia, vi. (1866) 281. t. 1 = **Psachystroma**, Klotzsch (Euphorb.).
spinosum, Baill. l. c. 287 = P. ilicifolium.

ACANTHOMINTHA, A. Gray, in Syn. Fl. N. Am. ii. 365 (1878). —*LABIATAE*, Benth. & Hook. f. ii. 1192.
ilicifolia, *A. Gray, l. c.*—Calif.
lanceolata, *Curran, in Bull. Calif. Acad.* i. (1884) 13.—Calif.

ACANTHONEMA, Hook. f. Bot. Mag. t. 5320 (1869). *GESNERACEAE*, Benth. & Hook. f. ii. 1091.
strigosum, *Hook. f. l. c.*—Afr. trop.

ACANTHONOTUS, Benth. in Hook. Nig. Fl. 293 (1849) = **Indigofera**, Linn. (Legum.).
echinatus, Benth. l. c. = I. echinata.

ACANTHONYCHIA, Rohrb. in Mart. Fl. Bras. xiv. ii. 249. t. 56 (1873) = **Pentaosana**, Bartl. (Illecebr.).
polycnemoides, Rohrb. l. c. 250 = P. polycnemoides.
ramarioinna, Rohrb. l. c. 249. t. 56 = P. polycnemoides.

ACANTHOPANAX, Miq. Ann. Mus. Lugd. Bat. i. 10 (1863). *ARALIACEAE*. Benth. & Hook. f. i. 930.
KALOPANAX, Miq. l. c. (1863).
PLECTRONIA, Lour. Fl. Cochinch. 169 (1790).
aculeatum, *Seem. in Journ. Bot.* v. (1867) 238.—As. trop.
asperatum, Franch. & Sav. Enum. Pl. Jap. ii. 378 = divaricatum.
divaricatum, *Seem. in Journ. Bot.* v. (1867) 239 — Japon.
innovans, *Franch. & Sav. Enum. Pl. Jap.* i. 198.—Japon.
japonicum, *Franch. & Sav. l. c.* ii. 377.—Japon.
ricinifolium, *Seem. in Journ. Bot.* vi. (1868) 140. —Japon.
sciadophylloides, *Franch. & Sav. Enum. Pl. Jap.* ii. 376.—Japon.
septem, Seem. in Journ. Bot. v. (1867) 239 = aculeatum.
sessiliflorum, *Seem. l. c.*—China.
spinosum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 10.—Japon.
trichodon, *Franch. & Sav. Enum. Pl. Jap.* ii. 377.—Japon.

ACANTHOPHIPPIUM, Blume, Bijdr. 553. t. 47 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 512.
bicolor, *Lindl. Bot. Reg.* xx. (1835) 1730.—Zeylan.
Curtisii, *Reichb. f. in Gard. Chron. N. S.* xv. (1881) i. 169.—Malaya.
javanicum, *Blume, Bijdr.* 354.—Java.
parviflorum, *Hassk. Cat. Bog. Alt.* 45.—Malaya.
ringiflorum, Griff. Notul. iii. 347 = sylhetense.
striatum, *Lindl. Bot. Reg.* (1838) *Misc.* 41.—Reg. Himal.
sylhetense, *Lindl. Gen. et Sp. Orch.* 177.—Reg. Himal.

ACANTHOPHOENIX, H. Wendl. in Fl. des Serres. Sér. II. vi. (1865–67) 181. t. 1706. *PALMEAE*, Benth. & Hook. f. iii. 898.
DRUDEANIA, H. Wendl. ex Seem. in Gard. Chron. (1870) 561.
crinita, *H. Wendl. l. c.*—Ins. Mascarens.
rubra, *H. Wendl. l. c.* sub t. 1706.—Ins. Mascarens.

ACANTHOPHYLLUM, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 87 = **Nanosauvia**, Jaus. (Compos.).
axillare, Hook. & Arn. l. c.(= Strongyloma glomeruloa).
roruulatum, Hook. & Arn. l. c. 48 = N. rosulata.

ACANTHOPHYLLUM, C. A. Mey. Verz. Pfl. Cauc. 210 (1831). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 145.
ALLOCHRUSA, Bunge, in Boiss. Fl. Orient. i. 559 (1867).
JORDANIA, Boiss. Diagn. Ser. I. viii. 93 (1849).
Alexandrae, *Smirn. ex Regel, in Act. Hort. Petrop.* v. (1877) 350.—Reg. Amur.
brucetanum, *Boiss. Diagn.* Ser. I. i. 43.—Persia.
Bungei, *Trautv. in Act. Hort. Petrop.* ii.(1873) 511.—Persia.
caespitosum, *Boiss. Diagn.* Ser. I. i. 43.—Persia.
crassifolium, *Boiss. l. c.*—Persia.
elatius, *Bunge, in Boiss. Fl. Orient.* i. 561.—Persia ; Turkest.
Fontanesii, Boiss. & Buhse, in Nouv. Mém. Mosc. xii. (1860) 89 = glandulosum.
Fontanesii, *Boiss. Diagn.* Ser. I. i. 41.—Persia.
glandulosum, *Bunge, ex Boiss. Fl. Orient.* i. 563.—Persia.
glaucum, Bunge, ex Boiss. l. c. 566 = sordidum.
gracile, *Bunge, ex Boiss. l. c.* 562.—Persia.
grandiflorum, *Stocks, in Hook. Kew Journ.* iv. (1852) 142.—Baluchist.
gypsophiloides, *Regel, Pl. Nov. Fedsch.* 15.—Turkest.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 349.—Turkest.
laxiflorum, Boiss. Diagn. Ser. II. i. 182 = Gypsophila ornata.
macrodon, *Edgew. in Hook. Fl. Brit. Ind.* i. 216.—Ind. or. ; Afghan.
microcephalum, *Boiss. Diagn.* Ser. I. i. 42.—Persia.
nitis, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. (1854) i. 35. t. 1.—As. Min.

ACANTHOPHYLLUM:—
mucronatum, *C. A. Mey. Verz. Pfl. Cauc.* 210.—Reg. Cauc.
mucronatum, Fenzl, in Ann. Wien. Mus. i. (1836) 37 = Tournefortii.
peniculatum, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) 1. 589.—Turkest.
pungens, Boiss. Fl. Orient. i. 661 = spinosum.
sordidum, *Bunge, ex Boiss. Fl. Orient.* i. 565.—Persia.
spinosum, Bunge, in Mém. Sav. étr. Pétersb. vii. (1851) 237 = elatum.
spinosum, *C. A. Mey. Verz. Pfl. Cauc.* 210.—Reg. Caucas.; Sibir.
squarrosum, *Boiss. Diagn.* Ser. II. i. 81.—Persia.
Stackianum, Boiss. l. c. = squarrosum.
Tournefortii, *Fenzl, in Ann. Wien. Mus.* ii. (1838) 310.—Armenia.
versicolor, *Pisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 31.—Persia.
verticillatum, *C. A. Mey. Verz. Pfl. Cauc.* 210.—Armenia.

ACANTHOPHYTON. Less. Syn. Comp. 128 (1832) = Olchorhum, Linn. (Compos.).
spinosum, Less. l. c. = C. spinosum.

ACANTHOPLEURA, C. Koch, in Bot. Zeit. vii. (1849) 408 = Polylophium, Boiss. (Umbell.).
involucrata, C. Koch, l. c. = P. orientale.

ACANTHOPRASIUM, Spenn. in T. Nees, Gen. Fl. Germ. Gamop. ii. (1843) = Ballota, Linn. (Labiat.).
frutescens, Spenn. l. c. = B. spinosa!

ACANTHOPSIS, Harv. in Hook. Lond. Journ. Bot. i. (1842) 28. *ACANTHACEAE*, Benth. & Hook. f. ii. 1089.
disperma, *Harv. l. c.*—Afr. austr.
horrida, *Nees, in DC. Prod.* xi. 278.—Afr. austr.

ACANTHORHIZA, H. Wendl. in Kerch. Palmiers, 230 (1878): Bot. Zeit. (1879) 147. *PALMAE*, Benth. & Hook. f. iii. 925.
aculeata, *H. Wendl. in Kerch. Palmiers*, 230; *Illust. Hortic.* xxvi. (1879) t. 367.—Mexico.
Chuco, *Drude, in Mart. Fl. Bras.* iii. 11. 554.—Bras.
Moolani, *Benth. & Hook. f. Gen.* iii. 925.—Mexico.
stenacanthа, H. Wendl. ex Linden, Cat. n. 87 (1871); Illust. Hortic. xxviii. (1881) 15 = aculeata.
Wallisi, *H. Wendl. ex Regel, in Gartenfl.* xxviii. (1879) 168. t. 977. (2.)—Hab.!
Warscewiczii, *H. Wendl. in Kerch. Palmiers*, 230.—Panama.

ACANTHOSICYOS, Welw. in Trans. Linn. Soc. xxvii. (1869) 80. *CUCURBITACEAE*, Benth. & Hook. f. i. 824.
horrida, *Welw. l. c.* 81.—Afr. trop.

ACANTHOSPERMA, Vell. Fl. Flum. viii. t. 152 (1827) = Acicarpha, Juss. (Calycer.).
littorale, Vell. l. c. = Acicarpha spathulata.

ACANTHOSPERMUM, Schrank, Pl. Rar. Hort. Monac. t. 53 (1819). *COMPOSITAE*, Benth. & Hook. f. ii. 349.
CENTROSPERMUM, H. B. & K. Nov. Gen. et Sp. iv. 270. t. 397 (1820).
ECHINODIUM, Poiteau, ex Cass. in Dict. Sc. Nat. lix. 235 (1829).
OXCYA, Vell. Fl. Flum. 344 (1825).
brasilium, *Schrank, Pl. Rar. Hort. Monac.* ii. 53.—Bras.
hirsutum, DC. Prod. v. 522 = brasilium.
hispidum, *DC. l. c.* 522.—Bras.
humile, *DC. l. c.* 522.—Ind. occ.
xanthioides, DC. l. c. 521 = brasilium.

ACANTHOSPORA, Spreng. Anleit. ii. 1. 255 (1817); Syst. ii. 25 (1830) = Tillandsia, Linn. (Bromeliac.).
conanthia, Spreng. Syst. ii. 25 = T. strobilantha.
juncea, Spreng. l. c. 25 = T. setacea.
vittata, D. Dietr. Syn. Pl. ii. 1056 = Cryptanthus bivittatus!

ACANTHOSTACHYS, Link, Klotzsch & Otto, Ic. Pl. Rar. t. 21. t. 9 (1841). *BROMELIACEAE*, Benth. & Hook. f. iii. 663.
strobilacea, *Link, Klotzsch, & Otto, l. c.* 22. t. 9.—Bras.

ACANTHOSTEMMA, Blume, Rumphia, iv. 29 (1848) = Hoya, R. Br. (Asclepiad.).
Hasseltii, Blume, l. c. = H. Hasseltii.
Kuhlii, Blume, l. c. = H. longifolia.
longifolium, Blume, l. c. = H. Rumphii.
pictum, Blume, l. c. = H. picta.
pruinosum, Blume, l. c. 30 = H. pruinosa.
puberum, Blume, l. c. 29 = H. rubera.
Rumphii, Blume, l. c. 29 = H. Rumphii.

ACANTHOSYRIS, Griseb. in Goett. Abh. xxiv. (1879) 151. *SANTALACEAE*, Benth. & Hook. f. iii. 234.
falcata, *Griseb. l. c.*—Reg. Argent.
spinescens, *Griseb. l. c.*—Reg. Argent.

ACANTHOTHECA, DC. Prod. vi. 78 (1837) = Dimorphotheca, Moench (Compos.).
dentata, DC. l. c. 74 = D. dentata.
integrifolia, DC. l. c. 74 = D. integrifolia.
pinnatisecta, DC. l. c. 73 = D. pinnata.

ACANTHOXANTHIUM, Fourr. in Ann. Soc. Linn. Lyon. N.S. xvii. (1869) 110 = Xanthium, Tourn. (Compos.).
spinosum, Fourr. l. c. 110 = X. spinosum.

ACANTHUS, Tourn. ex Linn. Syst. ed. I. (1735). *ACANTHACEAE*, Benth. & Hook. f. ii. 1090.
ACANTHODUS, Rafin. Princ. Somiol. 26 (1814).
CHEILOPSIS, Moq. in Ann. Sc. Nat. Sér. I. xxvii. (1832) 330.
DILIVARIA, Juss. Gen. 108 (1789).
ZONABLEPHIS, Rafin. Sylva Tellur. 148 (1838).
arboreus, *Forsk. Fl. Aegypt. Arab.* 116.—Aegypt.
Barteri, *Anders. in Journ. Linn. Soc.* vii. (1864) 87.—Afr. trop.
capensis, *Thunb. Prod. Pl. Cap.* 97.—Afr. austr.
cardiosecus, *Grif. Notul.* iv. 138.—Reg. Himal.
carduifolius, Linn. f. Suppl. 294 = Blepharis carduifolia.
Galeori, Burm. f. Fl. Ind. 139 = Blepharis boerhaviaefolia.
cristatus, Rottl. ex Nees, in DC. Prod. xi. 256 = Lepidagathis cristata.
Dellilei, Spreng. Syst. iii. 819 = Blepharis edulis.
Dioscoridis, *Linn. Cent.* ii. 28 (*Am. Acad. iv.* 321).—Syria.
Dolocarpha, Blanco, Fl. Filip. ed. I. 487 = ilicifolius.
ebracteatus, *Vahl, Symb. Bot.* iii. 75. t. 40.—Ax. trop.
edulis, Forsk. Fl. Aegypt. Arab. 115 = Blepharis edulis.
ferox, Linn. f. Suppl. 295 = Blepharis furcata.
glaber, E. Mey. ex Nees, in DC. Prod. xi. 278 = Blepharis carduifolia.
glaucus, E. Mey. ex Nees, l. c. 277 = Blepharis glauca.
glomeratus, Lam. Encyc. i. 23 = Blepharis glomerata.
hirsutus, *Boiss. Diagn.* Ser. I. iv. 86.—Orient.
hispanicus, Hort. ex Loud. Hort. Brit. 256 = niger.
humilis, Vahl, ex Nees, in DC. Prod. xi. 274 (= Acanthodium sinuatum.)
ilicifolius, *Linn. Sp. Pl.* 639 (err. typ. 939).—Ax. trop.; Austral.
ilicifolius, Lour. Fl. Cochinch. 375 = ebracteatus.
imbricatus, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) 217.—Ind. or.
integrifolius, E. Mey. ex Nees, in DC. Prod. xi. 276 = Blepharis furcata.
integrifolius, Thunb. Prod. Pl. Cap. 97 = Blepharis saxicaefolia.
Kirkii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 87. —Angola.
latifolius, Hort. ex Goese, in Gard. Chron. (1867) 856, 908 = mollis.
leucostachyus, *Wall. Cat.* n. 2512.—Ind. or.
longibracteatus, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) pt. 79.—Burma.
longifolius, *Poir. Encyc. Suppl.* i. 88.—Europ. austr.
longipetiolatus, *Kurz, For. Fl.* ii. 241.—Burma.
luxiflorus, Hort. cf. Gard. Chron. (1867) 908 = mollis.
macer, E. Mey. ex Nees, in DC. Prod. xi. 276 = Blepharis furcata.
maderaspatanus, Vahl, ex Nees, in DC. Prod. xi. 267 = Blepharis boerhaviaefolia.
maderaspatensis, Linn. Sp. Pl. 639 (err. typ. 939) = Blepharis mollugiuifolia.
mollis, Graf & Noë, ex Nees, in DC. Prod. xi. 270 = longifolius.
mollis, *Linn. Sp. Pl.* 639 (err. typ. 939).—Europ austr.

ACANTHUS:—
mollis, Riedel, ex Nees, in DC. Prod. xi. 271 = spinosus.
montanus, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 87.—Afr. trop.
mucronatus, Heyne, ex C. B. Clarke, in Hook. f. Fl. Br. Ind. iv. 517 = Lepidagathis spinosa.
niger, *Mill. Gard. Dict. ed.* VIII. n. 2.—Lusitan.
nitidus, *S. Moore, in Journ. Bot.* xviii. (1880) 232.—Afr. trop.
pectinatus, Willd. ex Nees, in DC. Prod. xi. 274 = Blepharis edulis.
polystachyus, *Delile, in Caill. Voy. au Méroé*, 2. t. 62. f. 2.—Afr. trop.
procumbens, Herb. Madr. ex Wall. Cat. sub n. 2513 = Blepharis boerhaviaefolia.
procumbens, Thunb. Prod. Pl. Cap. 97 = Blepharis glomerata.
pungens, Herb. Madr. ex Wall. Cat. n. 2498 = Lepidagathis pungens.
Raddei, Trautv. in Act. Hort. Petrop. iii. 11. (1875) 269 = Dioscoridis.
repandus, Nees, in DC. Prod. xi. 285, sphalm. = Anthacanthus repandus.
repens, Vahl, Symb. Bot. ii. 76 = Blepharis mollugiuifolia.
rubens, *Rafin. Fl. Ludov.* 86.—Am. bor.
spathularis, *E. Mey. ex Nees, in. DC. Prod.* xi. 277.—Am. bor.
spinosissimus, Desf. Tabl Hort. Par. 51 = spinosus.
spinosissimus, Host, Fl. Austr. ii. 218 = mollis.
spinosus, *Linn. Sp. Pl.* 639 (err. typ. 939).—Europ.
spinosus, Nees, in DC. Prod. xi 271 = hirsutus, syriacus
spinulosus, Host, Fl. Austr. ii. 217 = spinosus.
syriacus, *Boiss. Diagn.* Ser. I. xi. 185.—Syria.
tetragonus, *R. Br. Salt, Abyss. App.* 65; *Verm. Schr. ed. Nees*, iii. 249.—Abyss.
verus, *Wender. ex Steud. Nom.* ed. II. i. 11.—Hab.!
volubilis, *Wall. Pl. As. Rar.* ii. 66. t. 172.—Ind. or.

ACANTHYLLIS, Pomel, Nouv. Mat. Fl. Atl. 179 (1874) = Anthyllis, Riv. (Leguin.).
numidica, Pomel, l. c. = Anth. numidica.
tragacanthoides, Pomel, l. c. = Astragalus armatus.

ACARNA, All. Fl. Pedem. i. 158 (1785) = Atractylis, Linn. (Compos.).
caespitosa, Willd. Sp. Pl. iii. 1700 = Atractylis caespitosa.
cancellata, All. Fl. Pedem. i. 152 = Atractylis cancellata.
chinensis, Bunge, Enum. Pl. Chin. Bor. 36 = Atractylis ovata.
comosa, Spreng. Syst. iii. 380 = Atractylis comosa.
cyanoides, Bess. Enum. Pl. Volh. 32 = Jurinea cyanoides.
cyanoides, Wibel, Prim. Fl. Werth. 169 = Jurinea Pollichii.
gummifera, Willd. Sp. Pl. iii. 1699 = Carlina gummifera.
lanceа, Willd. l. c. 1700 = Atractylis ovata.
macrocephala, Willd. l. c. = Carlina Fontanesii.
macrophylla, Willd. l. c. = Carlina macrophylla.
mollis, Bess. Enum. Pl. Volh. 32 = Jurinea mollis.
ovata, Willd. Sp. Pl. iii. 1701 = Atractylis ovata.
rigida, Godet, ex DC. Prod. vi. 674 = Jurinea longifolia.
serratuloides, Spreng. Syst. iii. 379 = Atractylis serratuloides.

ACARNA, Hill, Veg. Syst. iv. 18 (1762) = Cnicus, Linn. (Compos.).
Casabonae, Hill, Hort. Kew. 59 = C. Casabonae.
leucographa, Hill, l. c. = Tyrimnus leucographus.
onopyxe, Hill, l. c. = Carduus Onopyxos.

ACARPHA, Griseb. in Goett. Abh. vi. (1858–55) 125 = Boopis, Juss. (Calycer.).
australis, Griseb. l. c. 126 = B. australis.

ACARPHAEA, Harv. & Gray, ex A. A. Gray, in Mem. Am. Acad. iv. (1849) 1. (Pl. Fendl.) 98 = Chaenactis, DC. (Compos.).
artemisiaefolia, Harv. & Gray, l. c. = C. artemisiaefolia.

ACASTE, Salisb. in Trans. Hort. Soc. i. (1812) 322 = Bubbiana, Ker-Gawl. (Irid.).
pulchra, Salisb. l. c. 323 = B. stricta.
venusta, Salisb. l. c. 323 = B. rubrocyanea.

ACCA, Berg, in Linnaea, xxvii. (1855) 138 = Feijdium, Linn. (Myrtac.).
dominguensis, Berg, l. c. 138.—Ind. occ.
peruviana, Berg, l. c. 139.—Peruv.

ACCIA. St. Hil. Morphol. 499 (1840) = **Flukenetia,** Linn. (Euphorb.).
scandens, St. Hill. l. c. (= Fragariopsis scandens).

ACCIPITRINA, Riv. ex Rupp. Fl. Jen. ed. Hall. 81 (1745) = **Sisymbrium,** Linn. (Crucif.).

ACCOROMBONA, Endl. Gen. Suppl. i. 1437 (1841) = **Galega,** Linn. (Legum.).
tricolor, Benth. ex Walp. Rep. i. 672 = G. officinalis.

ACEDILANTHUS, Benth. & Hook. f. Gen. iii. 885 (1881) sphalm. = seq.

ACELIDANTHUS, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 94. t. 28) (1847) = **Veratrum,** Linn. (Liliac.).
anticleoides, Trautv. & Mey. l. c. = V. Maackii.

ACENTRA, Phil. Sert. Mendoc. Alt. 8 (1871) *VIOLARIEAE.*
serrata, *Phil. l. c.*—Reg. Argent.

ACER, [Tourn.] Linn. Syst. ed. I. (1735). *SAPINDACEAE,* Benth. & Hook. f. i. 409.
EUACER, Opiz, Seznam, 49 (1852).
abchasicum, *Rupr. Fl. Cauc.* 979.—Reg. Caucas.
acuminatum, *Wall. ex D. Don, Prod. Fl. Nep.* 249. —Reg. Himal.
aetnense, Hort. ex C. Koch, Dendrol. 581 = platanoides.
affine, Hoffmgg. ex Walp. Rep. i. 410 = campestre.
affine, Opiz, in Flora, vii. (1824) Beil. i. 83; xii. (1829) 547 = campestre.
Amalise, *Orphan. ex Boiss. Fl. Orient.* i. 951.—Graecia.
amornum, Carr. in Rev. Hortic. (1867) 280 = palmatum.
argutum, *Maxim. ex Miq. in Arch. Nderl.* ii. (1867) 475.—Japon.
austriacum, Tratt. Arch. Gew. i. t. 86 = campestre.
barbatum, Hook. Fl. Bor. Am. i. 113 = glabrum.
barbatum, *Michx. Fl. Bor. Am.* ii. 252 = saccharinum.
barbinerve, *Maxim. ex Miq. in Arch. Nderl.* ii. (1867) 476.—Japon.
bohemicum, Presl, ex Opiz, Seznam, 9 = campestre.
Boscii, *Spach, Hist. Vég. Phan.* iii. 88.—Am. bor.?
Buergerianum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 86 = palmatum.
Buonipalia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 249 = oblongum.
caesium, *Wall. ex Brand. For. Fl.* iii. t. 21.—Reg. Himal.
californicum, D. Dietr. Syn. ii. 1288 = Negundo aceroides.
calliforniense, *Torr. & Gray, Fl. N. Am.* i. 250.—Calif.
Campbellii, *Hook. f. & Thoms., ex Brand. For. Fl.* 109.—Reg. Himal.
campestre, *Linn. Sp. Pl.* 1055.—Europ.; Orient.
canadense, Duham. Arb. i. t. 13 = pensylvanicum.
capillipes, *Maxim. ex Miq. in Arch. Nderl.* ii. (1867) 477.—Japon.
carolinianum, Walt. Fl. Carol. 251 = rubrum?
carpinifolium, *Sieb. & Zucc. in Abh. Abad. Muench.* iv. ii. (1845) 154.—Japon.
castaefolium, *Blume, Rumphia,* iii. 198. t. 167. B. f. 2. —Java.
caudatum, *Wall. Pl. As. Rar.* ii. 4.—Reg. Himal.
cinerascens, *Boiss Diagn.* Ser. I. vi. 29.—Persia.
circinnatum, *Pursh, Fl. Am. Sept.* i. 267.—Amer. bor.
circumlobatum, *Maxim. in Bull. Acad. Pétersb.* xii. (1867) 325.—Japon.
clasifolium, *C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 252.—Japon.
coccineum, Michx. f. Arb. Am. i. 208 = rubrum.
colchicum, Hartwiss, in Loud. Gard. Mag. (1840) 632 = pictum.
Collinsona, *Thumb. Diss. Acero,* 6.—Hab.?
collinsum, Tenore, Syll. 194 = campestre.
communatum, J. & C. Presl, Delic. Prag. 81 = monspessulanum.
cordifolium, Moench, Meth. 56 = tataricum.
coriaceum, Ledd. ex Spach, Hist. Vég. Phan. iii. 107 = polymorphum.
coriaceum, *Bosc, ex Tausch, in Flora,* xii. (1829) 550. —Hab.?
crataegifolium, *Sieb. & Zucc. in Abh. Abad. Muench.* iv. ii. (1845) 155.—Japon.

ACER :—
creticum, *Linn. Sp. Pl.* ed. II. 1497.—Oriens.
creticum, Schmidt, ex Tratt. Arch. Gew. i. t. 45 = parviflolium.
crispum, *Mœnch, Vers. Austr. Barume,* 2.—Hab.?
cultratum, Wall. Pl. As. Rar. ii. 4 = pictum.
dasycarpum, *Ehrh. Beitr.* iv. 94.—Am. bor.
Dedyle, Maxim. in Bull. Phys. Math. Acad. Sc. Pétersb. xv. (1856) 125 = spicatum.
dilatiolosum, *Blume, ex Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 251.—Japon.
discolor, *Maxim. in Bull. Acad. Sc. Pétersb.* xxvi. (1880) 486.—China.
dissectum, Thunb. Fl. Jap. 160 = palmatum.
distylum, *Sieb. & Zucc. in Abh. Abad. Muench.* iv. ii. (1845) 154.—Japon.
Ditrichii, Opiz, Seznam, 9 = A. Pseudo-Platanus.
Douglasii, Hook. Lond. Journ. Bot. vi. (1846) 77. t. 6 = glabrum.
Drummondii, Hook. & Arn. in l. c. i. (1834) 200 = rubrum.
eriocarpon, Opiz, in Flora, vii. (1824) i. Beil. 83 = campestre.
eriocarpum, Michx. Fl. Bor. Am. ii. 253 = dasycarpum. Fabri, *Hausc, in Journ. Bot.* xxii. (1884) 76.—China. Fieberi, *Opiz, Seznam,* 9.—Bohem.
floridanum, Hort. ex C. Koch, Dendrol. 542 = dasycarpum.
formosum, *Carr. in Rev. Hortic.* (1867) 300.—Japon.
fraxinifolium, Nutt. Gen. Am. i. 253 = Negundo aceroides.
fraxinifolium, *Rafin. New Fl. Am.* i. 48.—Am. bor.
Friederici-Guilelmi, Carr. in Rev. Hortic. (1867) 300 = palmatum.
Ginnala, Maxim. in Bull. Phys. Math. Acad. Sc. Pétersb. xv. (1856) 126 = tataricum.
glaucum, Hort. ex C. Koch, Dendrol. 543 = sanguineum.
glaucum, Marsh. Arbust. 2 = rubrum?
glabrum, *Torr. in Ann. Lyc. New York,* ii. (1828) 172. —Am. bor.
Granatense, Boiss. Elench. 25 = opulifolium.
grandidentata, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 249.—Am. bor.
Heldreichii, *Boiss. & Heldr. in Boiss. Diagn.* Ser. II. v. 71.—Graecia.
heterolobum, Opiz, Natural. ix. (1825) 114 = campestre.
heterophyllum, Willd. Berl. Baumz. 10. t. 1 = creticum.
hispanicum, Pourr. in Mém. Acad. Toul. iii. (1788) 305 = opulifolium.
Hookeri, *Miq. in Arch. Nderl.* ii. (1867) 471.—Reg. Himal.
humile, Saliab. Prod. 281 = creticum.
hybridum, Bosc, Dict. Agr. v. 251 = pensylvanicum.
hyrcanum, Hort. ex C. Koch, Dendrol. 543 = rubrum.
hyrcanum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 81.—Orient.
ibericum, Bieb. Fl. Taur. Cauc. ii. 447 = monspessulanum.
illyricum, Jacq. f. in Flora, xii. (1829) 551 = monspessulanum.
insigne, *Boiss. & Buhse, in Nouv. Mém. Mosc.* xii. (1860) 46.—Persia.
intermedium, Fauč. ex Nym. Consp. 135 = monspessulanum.
Isolobum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 204.—Burma.
italum, *Lauth, de Acere,* n. 8.—Europ.; Oriens.
japonicum, *Thunb. Fl. Jap.* 161.—Japon.
javanicum, Jungh. in Hoev. & De Vr. Tijdschr. viii. (1841) 301 = niveum.
javanicum, *Burm. f. Fl. Ind.* 221.—Java.
jucundum. *Carr. in Rev. Hortic.* (1867) 280.—Japon.
lacinistum, Borckh. ex Tratt. Arch. Gew. 85 = platanoides.
lactescens, Pers. Syn. i. 417 = platanoides.
lactum, *C. A. Mey. Verz. Pfl. Cauc.* 236.—Orient.
lævigatum, *Wall. Pl. As. Rar.* ii. 3. t. 104.—Reg. Himal.
laurifolium, D. Don, Prod. Fl. Nep. 249 = oblongum.
laurinum, *Hassk. in Hoev. & De Vr. Tijdschr.* x. (1843) 126 = niveum.
laurinum, Hassk. l. c. = niveum.
leiocarpon, Opiz, in Flora, vii. (1824) i. Beil. 82 = campestre.
leptoderum, Gass. ex Nym. Consp. 135 = opulifolium.

ACER :—
lobatum, Bosc, ex C. Koch, Dendrol. 519 = Boscii.
lobatum, *Rafin. New Fl. Am* i. 48.
Lobelii, Tenore, Fl. Nap. v. 29. t. 199 = platanoides.
macrocarpon, Opiz, in Flora, vii. (1824) i. Beil. 82 = campestre.
macrophyllum, *Pursh, Fl. Am. Sept.* i. 267.—Am. bor.
macropterum, Vis, in Mem. Ist. Venet. ix. (1860) 175; Suppl. 23 = Visianii.
majus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 635 = A. Pseudo-Platanus.
mandshuricum, Maxim. in Bull. Acad. Sc. Pétersb. xxi. (1868) 238.—Manchur.
marsicum, Guss. Fl. Nap. 875 = campestre.
Martini, Jord. Pugill. 52 = opulifolium.
marucum, Walp. Rep. i. 408 = campestre.
Maximowiczianum, *Miq. in Arch. Nderl.* ii. (1867) 478. —Japon.
mellicdorum, Opiz, Seznam, 9 = A. Pseudo-Platanus.
mexicanum, *A. Gray, in Proc. Am. Acad.* v. (1861) 176.—Mexico.
micranthum, *Sieb. & Zucc. in Abh. Abad. Muench.* iv. ii. (1845) 155.—Japon.
microcarpon, Mœnch, ex Opiz, Natural. vi. (1824) 48; in Flora, vii. (1824) i. 83 = campestre.
microphyllum, Opiz, in Flora, vii. (1824) i. Beil. 82; et Natural. vi. (1824) 48 = campestre.
molly, Opiz, l. c. 83 = campestre.
Mono, Maxim. in Bull. Phys. Math. Acad. Sc. Pétersb. xv. (1856) 126 = pictum.
monspessulanum, *Linn. Sp. Pl.* 1056.—Europ. austr.
montanum, Ait. Hort. Kew. ed. II. iii. 435 = spicatum.
montanum, Lam. Fl. Fr. ii. 553 = A. Pseudo-Platanus.
neapolitanum, Tenore, Fl. Nap. ii. 372. t. 100 = opulifolium.
neglectum, *Hoffmgg. ex Walp. Rep.* i. 410 = Hab.?
neglectum, *Lange, in Bot. Tidss.* xiii. (1882) 30.—An hybr.?
Negundo, Linn. Sp. Pl. 1056 = Negundo aceroides.
nigrum, Michx. f. Arb. Am. ii. 238. t. 16 = saccharinum.
nikoense, Maxim. in Bull. Acad. Sc. Pétersb. xii. (1868) 227 = Maximowiczianum.
nikoense, *Miq. in Arch Nderl.* ii. (1867) 478.—Japon.
niveum, *Blume, Rumphia,* iii. 193.—Java; Ind. or.
oblongum, *Wall. ex DC. Prod.* i. 598.—Reg. Himal.
obtusatum, Waldst. & Kit. ex Willd. Sp. Pl. iv. 984 — opulifolium.
obtusifolium, Sibth. & Sm. Fl. Græc. t. 361 = creticum.
Opalus, Mill. Gard. Dict. ed. VIII. n. 8 = opulifolium. Opizii, *Ortm. ex Opiz, Seznam,* 9.—Bohem.
opulifolium, Thnil. ex Loud. Hort. Brit. Supp. ii. 606 = A. Pseudo-Platanus.
opulifolium, *Vill. Hist. Pl. Dauph.* iv. 809 = Europ; Oriens.
orientale, Linn. Syst. ed. X. ii. 1300 = creticum.
ornatum, *Carr. in Rev. Hortic.* (1867) 300.—Japon.
orthopteron, Mass. ex Opiz, Natural. vi. (1824) 49; et Flora, vii. (1824) i. Beil. 83 = campestre.
pallidum, Ait. ex Steud. Nom. ed. II. i. 11 = dasycarpum.
palmatifidum, Tausch, ex Steud. l. c. = platanoides.
palmatisectum, Ortm. ex Celak. Prod. Fl. Boehm. 538 = campestre.
palmatum, Rafin. New Fl. Am. i. 48 = macrophyllum.
palmatum, Thunb. Fl. Jap. 162.—Japon.
palmifolium, Borckh. Fl. Kataeb. 107 = saccharinum.
parviflorum, Ekrh. Beitr. iv. 25 = spicatum.
parviflorum, *Franch. & Sav. Enum. Pl. Jap.* ii. 321. Japon.
parvifolium, *Tausch, in Flora,* xii. (1829) 545.—Europ. austr.
pectinatum, Wall. Pl. As. Rar. ii. 4 = caudatum.
pensylvanicum, Durol. Diss. 61 = spicatum.
pensylvanicum, *Linn. Sp. Pl.* 1055.—Am. bor.
pentapomicum, *Stewart, ex Brand. For. Fl.* 111.—Reg. Himal.
pictum, *Thunb. Fl. Jap.* 163.—Japon.
pilosum, *Maxim. in Bull. Acad. Sc. Pétersb.* xxvi. (1880) 486.—As. or.
pinnatum, *Lour., Fl. Cochinch.* ii. 649.—Cochinch.
platanifolium, *Grisb. Iter. Hung.* 300 (= villosum!).—Reg. Himal.
platanifolium, Steinz, Bot. Mat. Med. ii. 883 = platanoides.
platanoides, *Linn. Sp. Pl.* 1055.—Europ.; Oriens.
Platanus, Barm. f. Fl. Ind. 231 (= quid?).—Ind. or.
polycarpon, Opiz, Natural. vi. (1824) 49; in Flora, vii. (1824) i. Beil. 82 = campestre.

ACER :—
polymorphum, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1845) 158 = palmatum.
polymorphum, *Spach, Hist. Vég. Phan.* iii. 107.—Orient.
pracox, Opiz, Seznam, 9 = campestre.
procerum, Salisb. Prod. 389 = A. Pseudo-Platanus.
Pseudoplatanus, Falk, Beitr. ii. 279 = platanoides.
Pseudo-Platanus, *Linn. Sp. Pl.* 1054.—Europ.; Orient.
pubescens, *Prauch. in Ann. Sc. Nat. Sér.* VI. xv. (1883) 246.—Turkest.
purpurascens, *Franch. & Sav. Enum. Pl. Jap.* ii. 320. —Japon.
pycnanthum, *C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 260.—Japon.
quinquelobatum, Wagn. ex Opiz, in Flora, vi. 1. (1824) Bell. 83 = campestre.
quinquelobum, C. Koch, Dendrol. 540 = creticum.
quinquelobum, Gilib. Fl. Lituan. ii. 198 = A. Pseudo-Platanus.
rectangulum, Dulac, Fl. Hautes-Pyr. 242 = monspessulanum.
Reginæ-Amaliæ, Orphan. ex Boiss. Diagn. Ser. II. i. 106 = Amaliæ.
reticulatum, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 312.—Hongkong.
Reyyassi, Boiss. Diagn. Ser. I. ii. 5, 71 = hyrcanum.
robustum, Opiz, Seznam, 9 = campestre?
rotundifolium, Lam. Encyc. ii. 382 = Italum.
rotundum, Dulac, Fl. Hautes-Pyr. 242 = platanoides.
rubrum, Lam. Encyc. ii. 380 = dasycarpum.
rubrum, *Linn. Sp. Pl.* 1055.—Am. bor.
ruhrum, Wats. Dendrol. Brit. t. 169 = rubrum.
rufinerve, *Siob. & Zucc. in Abh. Akad. Muench.* iv. ii. (1845) 155.—Japon.
saccharinum, Linn. Sp. Pl. 1055 = dasycarpum.
saccharinum, *Wangenh. Nordamer. Holzarten* (1787) 96. t. ii. f. 96.—Am. bor.
saccharophorum, *C. Koch & Fint. Wochenschr.* (1859) 349.—Am. bor.
Saint, Hort. ex C. Koch, Dendrol. 541 = dasycarpum.
sanguineum, *Carr. in Rev. Hortic.* (1867) 380.—Japon.
sanguineum, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 176 = rubrum.
Schwedleri, Hort. ex C. Koch, Dendrol. 530 = platanoides.
Semenovii, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) t. 550.—As. temp.
sempervirens, Linn. Mant. i. 128 = heterophyllum.
sempervirens, *Tausch, in Sit.* (1829) 550.—Hab.? *septemlobum*, Thunb. Fl. Jap. 162 = palmatum.
serotinum, *Kit. in Linnæa*, xxxii. (1863) 552.—Europ.
sessilifolium, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1845) 158 = palmatum.
Sieboldianum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 87. —Japon.
sikkimense, *Miq in Arch. Nierl.* ii. (1867) 471.—Reg. Himal.
spicatum, *Lam. Encyc.* ii. 381.—Am. bor.
stachyophyllum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 604. —Reg. Himal.
stereulicaseum, Griff. Itin. Notes, 148 = pictum.
stereuliscum, Wall. Pl. As. Rar. ii. & t. 105 = villosum.
striatum, Du Roi, Harbk. Baumz. i. 8 = spicatum.
suberosum, Dum. Fl. Belg. Prod. 118 = campestre.
sylvestre, Wender. in Schr. Nat. Ges. Marb. ii. (1830) 350 = campestre.
syriacum, *Boiss. & Gaill Diagn.* Ser. II. v. 72.— Syria; Cyprus.
tataricum, *Linn. Sp. Pl.* 1054.—Orient.
tauricolum, Boiss. & Balansa, Diagn. Ser. II. v. 72 = hyrcanum.
Tauschianum, Opiz, Seznam, 9 = campestre.
tegmentosum, *Maxim. in Bull. Phys. Math. Acad. Sc. Pétersb.* xv. (1866) 135.—Reg. Amur.
tenuatum, *Moc. & Sesse, ex DC. Prod.* i. 596.—Mexico.
tenuerdonum, Dulac, Fl. Hautes-Pyr. 243 = opulifolium.
Thomasii, Miq. in Arch. Nérl. ii. (1867) 470 = villosum.
tomentosum, Kit. in Linnæa, xxxii. (1863) 552 = campestre.
tomentosum, Desf. Cat. Hort. Par. ed. III. 386 = ericarpum.
Trautvetteri, *Medw. ex Trautv. in Act. Hort. Petrop.* vii. (1880) 439 in obs.—Hab.?
trichnopetiolum, Stokes, Bot. Mat. Med. ii. 370 = pensylvanicum.
trifidum, *Hook. & Arn. Bot. Beech. Voy.* 174.—China.
triflorum, Thunb. Fl. Jap. 163 = Lindera triloba.
trilobatum, *Regel. New Fl. Am.* i. 48.—Am. bor.
trifolium, Duham, Arb. i. t. 10. f. 8 = monspessulanum.
ACHE :—
trilobatum, Lam. Encyc. ii. 389 = monspessulanum.
trilobum, Gilib. Fl. Lituan. ii. 198 = campestre.
trilobum, Moench, Meth. 66 = monspessulanum.
tripartitum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 248 = glabrum.
truncatum, Bunge, Enum. Pl. Chin. Bor. 10 = pictum.
uhurundurense, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 24) = spicatum.
Van-Volxemii, *Mast. in Gard. Chron.* vii. (1877) 72. —Reg. Caucas.
velutinum, Boiss. Diagn. Ser. I. vi. 28 = insigne.
villosum, J & C. Presl, Delic. Prag. 81 = A. Pseudo-Platanus.
villosum, *Wall. Pl. As. Rar* ii. 4.—Reg. Himal.
virens, *Thunb. Diss. Acere*, 3. (err. typ. 5).—Oriens.
virgatum, *Rafin. New Fl. Am.* i. 48.—Am. bor.
virginianum, Mill. ex Steud. Nom. ed. I. 5 = dasycarpum.
Visiani, *Nym. Consp.* 135.—Servia.
vitifolium, Opiz, ex Tausch, in Flora, xii. (1829) ii. 549 = platanoides.
Wendzelchii, Opiz, Seznam, 9 = campestre.

ACHÆNTHUS, Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1837) 349. t. 14. *BERBERIDEÆ*, Benth. & Hook. f. i. 44.
diphyllus, *Morr. & Decne. l. c.* 350.—Japon.
macrophyllus, *Blume, ex C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 258.—Japon.
sagittatus, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1845) 176 = Epimedium sagittatum.
triphyllus, *C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 258.—Japon.

ACHÆRAS, R. Br. in Ait. Hort. Kew. ed. II. v. 191 (1813). *SAPOTIDEÆ*, Benth. & Hook. f. iii. 621.
affinis, Boiss. Fl. Orient. v. 56.—As. Min.
alpina, Steud. Nom. ed. II. i. 12 = Herminium alpinum.
angustifolia, Lindl. in Wall. Cat. n. 7061 = longicrenis.
anthropomorpha, Steud. Nom. ed. II. i. 12 = anthropophora.
anthropophora, *R. Br. in Ait. Hort. Kew.* ed. II. v. 191. —Europ.
caprina, Lindl. Gen. et Sp. Orch. 282 = Orchis hircina.
densiflora, Boiss. Voy. Espagne, ii. 595 = Habenaria intacta.
Duguesei, Reichb f. fr. Fl. Germ. xiii-xiv. 171 = Orchis pyramidalis.
formosa, Lindl. Gen. et Sp. Orch. 282 = Orchis formosa.
hircina, Lindl. l. c. = Orchis hircina.
intacta, Reichb. f. lc. Fl. Germ. xiii-xiv. 2 = Habenaria intacta.
lancea, Steud. Nom. ed. II. i. 12 (= Ophryx lancea).
longibracteata, Reichb. f. lc. Fl. Germ. xiii et xiv. 3 = Orchis longibracteata.
longicrenis, *C. Wright, ex A. Gray, in Mem. Am. Acad. N. S.* v. (1858 9) 411.—Japon.
angustifolia, Decne. in DC. Prod. viii. 522 = Asclepias stenophylla.
asperula, Decne. l. c.—Mexico.
atropurpurea, Kellogg, in Proc. Calif. Acad. (1854–59) 65; repr. 69 = G. sordidiflus.
auriculata, Engelm. ex Torr. in Bot. Mex. Bound. 160. —Am. bor.
bifida, Rusby, ex A. Gray, in Proc. Am. Acad. xx. (1885) 296.—Arizona.
circinalis, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 332.—Mexico.
connivens, Decne. in DC. Prod. viii. 521 (= Anantherix connivens).
cordifolia, Benth. Pl. Hartw. 323 = G. cordifolius.
decumbens, Decne. in DC. Prod. viii. 522 = Asclepiodora decumbens.
Fraseri, Chapm. in Coulter, Bot. Gazette, iii. (1878) 12. —Florida.
gomphocarpoides, Benth. Pl. Hartw. 66 = Nephradenia nertiifolia.
gomphocarpoides, Decne. in Ann. Sc. Nat Sér. II. ix. (1838) 323.—Mexico.
humilis, Benth. Pl. Hartw. 291.—Mexico.

ACERATES :—
incarnata, Decne. in DC. Prod. viii. 523.—Am. bor.
lanceolata, Steud. Nom. ed. II. i. 12 = G. viridiflorus.
lanuginosa, Decne. in DC. Prod. viii. 523.—Am. bor.
latifolia, Torr. & Frém. in Frém. Rep. Exped. (1845) 317 = Asclepias cryptoceras.
longifolia, Ell. Sketch, i. 317 = Asclepias longifolius.
paniculata, Decne. in DC. Prod. viii. 521 = Asclepiodora viridis.
Schaffneri, *Fourn. in Ann. Sc. Nat. Sér.* VI. xiv. (1852) 386 —Mexico.
tomentosa, Torr. in Bot. Mex. Bound. 160 = G. tomentosus.
vinosa, *Fourn. in Ann. Sc. Nat. Sér.* VI. xiv. (1882) 387.—Mexico!
viridiflora, Ell. Sketch, 317 = G. viridiflorus.

ACERATIUM, DC. Prod. i. 519 (1824) = **Elæocarpus**, Linn. (Tiliac.).
Calomala, Blanco, Fl. Filip. ed. II. 306 = E. nitidus.
oppositifolium, DC. Prod. i. 519 = E. oppositifolius.

ACEROTIS, Rafin. New Fl. Am. i. 49 (1836) = **Asclepias**, Tourn.

ACETOSA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Rumex**, Linn. (Polygon.).
Acetosella, Mill. Gard. Dict. ed. VIII. n 2 = R. Acetosella.
aculeata, Thuz. in Mill. Dict. Suppl. i. 8 = R. bucephalophora.
agrestis, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
alpina, Mill. Gard. Dict. ed. VIII. n. 5 = R. montanus.
alpina, Moench, Meth. 357 = K. alpinus.
amplexicaulis, Rafin. New Fl. Am. iv. 53 = R. Acetosa.
angustata, Rafin. l. c. = R. Acetosa.
arifolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 64 = R. axifolius.
badensis, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
bipinnata, Chaz. in Mill. Dict. Suppl. i. 8 = R. pictus.
bucephalophora, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 145 = K. bucephalophorus.
digyna, Mill. Gard. Dict. ed. VIII. n. 4 = Oxyria digyna.
hastata, Moench, Meth. 357 = R. Acetosella.
hastifolia, Schur, Enum. Pl. Transs. 581 = R. Acetosa.
hastulata, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
heterophylla, Rafin. l. c. = R. maximus!
integrifolia, Rafin. l. c. = K. integrifolia.
intermedia, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 165 = R. intermedius.
Lunaria, Mill. Gard. Dict. ed. VIII. n. 8 = R. Lunaria.
luxurians, Chaz. in Mill. Dict. Suppl. i. 8 = R. luxurians.
magna, Gilib. Exercit. ii. 445 = R. Acetosa.
multifida, Chaz. in Mill. Dict. Suppl. i. 8 = R. multifidus.
officinalis, Gueldenst. ex Ledeb. Fl. Ross. iii. 510 = R. Acetosa.
oistoria, Rafin. New Fl. Am. iv. 53 = R. Acetosa.
parva, Gilib. Exercit. ii. 446 = R. Acetosella.
pratensis, Mill. Gard. Dict. ed. VIII. n. 1 = R. Acetosa.
repens, S. F. Gray, Nat. Arr. Brit. Pl. ii. 276 = R. Acetosella.
rosea, Mill. Gard. Dict. ed. VIII. n. 7 = R. roseus.
scutata, Mill. l. c. n. 8 = K. scutatus.
sterilis, Mill. l. c. n. 8 = R. Acetosella.
subalpina, Schur, Enum. Pl. Transs. 581 = R. Acetosa.
tingitana, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 145 = R. tingitanus.
tuberosa, Chaz. in Mill. Dict. Suppl. i. 8 = R. tuberosus.

ACETOSELLA, Rupp. Fl. Jen. ed. Hall. 127 (1745) = **Rumex**, Linn. (Polygon.).
vulgaris, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 145 = R. Acetosella.

ACHAEMENES, St. Lag. in Ann. Soc. Bot. Lyon. vii. (1880) 41 = **Achimenes**, P. Br. (Gesnerac.).

ACHAETA, Fourn. (Gram. Mex. 109) ex Benth. & Hook. f. Gen. iii. 1158 (1883) = **Deyeuxia**, Clar. (Gramin.).
gracilentula, Fourn. l. c. ex Hemsl. Biol. Centr. Am. Bot. iii. 558.—Mexico.
plumosa, Fourn. l. c. ex Hemsl. l. c. 563.—Mexico.

ACHAETOGERON, A. Gray, in Mem. Am. Acad. N.S. iv. (1849) (Pl. Fendl.) 72. *COMPOSITÆ*, Benth. & Hook. f. ii. 293.
Galeottii, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 119.—Mexico.
Seemannii, *A. Gray, ex Hemsl. l. c.* 120.—Mexico.

ACHAETOGERON :—
Wiklemi, *A. Gray, in Mem. Am. Acad. N. S.* iv. (1849) (*Pl. Fendl.*) 73.—Mexico.

ACHANIA, Sw. Prod. Veg. Ind. Occ. 109 (1788) = **Malvaviscus**, Dill. (Malvac.).
ciliata, Spreng. Syst. iii. 100 = M. ciliatus.
coccinea, Sallsb. Prod. 395 = M. arborea.
concinna, Spreng. Syst. iii. 100 = M. concinnus.
cordata, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 99 = M. Balbisii.
floridana, Rafin. New Fl. Am. i. 49 = Hibiscus floridanus!
Malvaviscus, Sw. Prod. Veg. Ind. Occ. 109 = M. arboreus.
mollis, Alt. Hort. Kew. ii. 459 = M. mollis.
pilosa, Sw. Prod. Veg. Ind. Occ. 109 = M. pilosus.
Poeppigii, Spreng. Syst. iii. 100 = Hibiscus Poeppigii.
stylosa, Schrank, in Flora, ii. (1819) 449.—Bras.
tomentosa, Sterler, ex Steud. Nom. ed. II. i. 12.—Hab.!

ACHARIA, Thunb. Prod. Pl. Cap. 14 (1794). *PASSIFLOREÆ*, Benth. & Hook. f. i. 814.
tragodes, *Thunb. l. c.*—Afr. austr.

ACHARITEA, Benth. in Benth. & Hook. f. Gen. ii. 1142 (1876). *VERBENACEÆ*, Benth. & Hook. tenuis, *Benth. l. c.*—Madag.

ACHARITERIUM, Bluff & Fingerh. Fl. Germ. ii. 845 (1838) = **Filago**, Linn. (Compos.).
arvense, Bluff & Fingerh. l. c. 846 = F. arvensis.

ACHASMA, Griff. Notul. iii. 426 (1851) = **Amomum**, Linn. (Scitam.).
macrochelios, Griff. l. c. 429.—Ind. or.
megalochelios, Griff. l. c. 426.—Ind. or.
microchelios, Griff. l. c. 427.—Ind. or.

ACHATOCARPUS, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 46. *AMARANTACEÆ*, Benth. & Hook. f. iii. 26.
nigricans, *Triana, l. c.* 46.—N. Granat.
praecox, *Griseb. in Goett. Abh.* xxiv. (1879) 32.—Reg. Argent.
spinulosus, *Griseb. l. c.* 32.—Reg. Argent.

ACHETARIA, Cham. & Schlecht. in Linnæa, ii. (1827) 566 = **Dyrichia**, Cham. (Scrophulariin.).
Sprengelii, Cham. & Schlecht. l. c. 557 = B. villosa.

ACHILLAEA, Linn. Phil. Bot. 29 (1764) = **Achillea**, Linn. (Compos.).

ACHILLEA, Linn. Syst. ed I (1735). *COMPOSITÆ*, Benth. & Hook. f. ii. 419.
AGERATOIDES, Ponted. ex Linn. Class. Pl. 527 (1747).
ALITUBUS, Dclac, Fl. Hautes-Pyr. 499 (1867).
ARTHROLEPIS, Boiss. Diagn. Ser. I. xi. 14 (1849).
CONFORATA, Cassalp. ex Fr. Herb. Linn. in Ann. Soc. Linn. Lyon. N. S. xvi. (1869) 91.
MILLEFOLIUM, Tourn. ex Adans. Fam. ii. 128 (1768).
PTARMICA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 174 (1745).
abrotanifolia, Hort. ex Steud. Nom. ed. II. i. 12 (= Pyrethrum myriophyllum).
abrotanifolia, Linn. Sp. Pl. 897 = Tanacetum millefolistum.
abrotanoides, *Vis. Fl. Dalmat.* ii. 81. t. 10. f. 1.—Europ.
abrotanifolia, [Chaix.] Man. Herb. 251 = Clavennæ.
acuminata, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Sibiria.
adultrina, Leresch, ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 167 = obscura.
aegyptiaca, S. G. Gmel. lt. ii. 198 = Gerberi.
aegyptiaca, Hort. ex Steud. Nom. ed. II. i. 12 = semipectinata.
aegyptiaca, Linn. Sp. Pl. 900 = Tournefortii.
aegyptiaca, Sm. Fl. Græc. ix. 71 = taygetea.
ageratifolia, *Benth. & Hook. f. Gen.* ii. 420.—Graecia.
Agerstum, *Linn. Sp. Pl.* 897.—Europ.
albicaulis, C. A. Mey. Verz. Pfl. Cauc. 76 = tenuifolia.
alkida, Willd. Enum. Hort. Berol. Suppl. 61 = tenuifolia.
aleppica, *DC. Prod.* vii. 296.—Orientia.
alpina, *Linn. Sp. Pl.* 899 = Europ.
ambigua, Boiss. Fl. Orient. iii. 295 = A. Millefolium.
ambigua, Pollini, Vieg. Lag. Garda, 117 = magna.

ACHILLEA :—
ambrosiaca, *Boiss. Fl. Orient.* iii. 276.—Thessal.
amoena, C. A. Mey. Vers. Pfl. Cauc. 76 = vermicularis.
anethifolia, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. (1865) I. 404 = A. Millefolium.
anglica, G. Don, in Loud. Hort. Brit. 355 = decolorans.
angustifolia, Salisb. Prod. 204 = alpina.
anthemoides, *Willd. Sp. Pl.* iii. 2100.—Hab.?
arabica, *Kotschy, in Sitzb. Acad. Wien. Math. Nat.* liii. Abth. i. (1866) 255.—Arab.
arborea, Hornem. Hort. Hafn. Suppl. 102 = Gonospermum elegans.
argentea, Sallsb. Prod. 204 = Clavennæ.
argentea, Lam. Encyc. i. 29 = Tanacetum argenteum.
armenorum, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 269.—Armen.
asperula, Ces. & Pass. ex Nym. Consp. 362 = Anthemis asperula.
asplenifolia, Leresch, ex Sch. Bip. in Oestr. Bot. Wochenschr. vi. (1856) 301 = Lereschel.
asplenifolia, *Vent. Jard. Cels*, t. 95.—Am. bor.!
atrata, *Linn. Sp. Pl.* 899.—Europ.
Auchel, *Boiss. Diagn.* Ser. I. vi. 87.—Persia.
aurea, Lam. Encyc. i. 26 = Chrysanthemum achilleæfolium.
auriculata, *Willd. Sp. Pl.* iii. 2208.—Orientia.
batkalensis, *Fisch. Hort. Gorenk.* ed. II. (1812) 41.—Sibir. batkal.
bandana, Buch.-Ham. in Wall. Cat. n. 3291 = Chrysanthelium indicum.
bannatica, Kit. ex DC. Prod. vi. 29 = crithmifolia.
Barbeyana, *Heldr. & Heimerl, in Denkschr. Acad. Wien*, xlviii. (1884) 163.—Graecia.
Bardana, Buch.-Ham. ex DC. Prod. v. 681. sphalm. = *bandana*.
Barrelieri, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Europ.
biaristata, Spreng. Syst. iv. Cur. post. 303 = Anacyclus clavatus.
bicolor, *Wender. in Flora*, xii. (1829) i. *Erg.* 25.—Europ.!
bipinnata, Linn. Sp. Pl. 900 (= Pyrethrum myriophyllum).
biserrata, *Bieb. Fl. Taur. Cauc.* ii. 334.—As. Min.
Bocconii, Koch, Ind. Sem. Hort. Erlang. (1834) = alpina.
Boissieri, *Haussk. ex Boiss. Fl. Orient.* iii. 272.—Persia.
Boissieriana, Heldr. ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 170 = Babeyana.
borealis, Bong. Veget. ins. Sitcha, 149 = A. Millefolium.
brachyphylla, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 273.—Mesopot.
bugloasis, Frivald. in Flora, xix. (1836) 483 = lingulata.
callichroa, *Boiss. Diagn.* Ser. I. vi. 86.—Persia.
camphorata, Gilib. Fl. Lituan. i. 177 = nobilis.
cantabrica, Rupr. ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 189 = sibirica.
Cancrini, *Gruner, in Bull. Soc. Nat. Mosc.* xli. (1868) 21. 417.—Tauria.
capillifolia, Poir. Encyc. Suppl. i. 102 = setacea.
capillata, Lapeyr. Hist. Abr. Pl. Pyr. 534 = chamaemelifolia.
capitata, Willd. Tract. Ach. 17 = Clavennæ.
cartilaginea, *Ledeb. Ind. Dorp. ex Boiss. Fl. Orient.* iii. 274.—Europ.; Orient.
chamaecyparissus, *Sch. Bip. in Flora*, xxxviii. (1855) 14.—Europ.
chamaecyparyssus, Reichb. f. Ic. Fl. Germ. xvi. 64. t. 1012 = prace.
chamaemelifolia, *Pourr. in Mém. Acad. Toul.* iii. (1788) 306.—Mont. Pyren.
chrysantha, Griseb. ex Boiss. Fl. Orient. iii. 295 = chrysocoma.
chrysocoma, *Frivald. in Flora*, xviii. (1835) 896.—Graecia.
cilicica, *Y. Gay, ex Tchihat. Asie Min. Bot.* ii. 270. — Min.
claudiopolitana, Janka, in Oestr. Bot. Wochenschr. (1854) 409 = impatiens.
claudiopolitanus, Wolff, ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 185 = impatiens.
Clavennæ, *Linn. Sp. Pl.* 898.—Europ.
Clusiana, *Tenoch, in Flora*, iv. (1821) ii. 551.—Europ.
Clusii, Sauter, in Flora, vii. (1824) i. 211 = Clusiana.
clypeolata, *Sibth. & Sm. Fl. Græc.* t. 898.—Orient.
coarctata, Poir. Encyc. Suppl. i. 94 = compacta.

ACHILLEA :—
collina, Becker, ex Reichb. Fl. Germ. Excurs. 850 = A. Millefolium.
collina, Schur, ex Nym. Consp. 367 = setacea.
communata, *Heimerl, in Flora*, lxvi. (1883) 385.—Europ.
compacta, Lam. Encyc. i. 27 = magna.
compacta, *Willd. Sp. Pl.* iii. 2206.—Orientia.
condensata, *Miq. Fl. Ind. Bat.* ii. 85.—Java.
conferta, *DC. Prod.* vi. 32.—Orientia.
coronopifolia, Willd. Tract. Ach. 18. n. 9. t. 1. f. 2 = tanacetifolia.
corymbifera, S. G. Gmel. lt. i. 187. t. 25. f. 1 = nobilis.
crassifolia, Hort. ex Steud. Nom. ed. II. i. 13 = A. Millefolium.
cretica, *Linn. Sp. Pl.* 899.—Oriens.
crispa, Lam. ex DC. Prod. vi 26 = asplenifolia.
cristata, Hort. ex DC. l. c. 25 = tanacetifolia.
cristata, Retz. Obs. Fasc. ii. 25 = falcata!
crustata, Willd. Tract. Ach. n. 4 et 8; Sp. Pl. iii. 2192 = alpina.
crithmifolia, Frivald. ex Hampe, in Flora, xx. (1837) i. 229 = micrantha.
crithmifolia, *Waldst. & Kit. Pl. Rar. Hung.* i. 68. t. 66.—Oriens.
crustata, *Schur, Enum. Pl. Transs.* 329.—Transylv.
cuneata, *Sch. Bip. in Linnæa*, xix. (1846) 813 = grandifolia.
cuneatiloba, *Boiss. & Buhse, in Nouv. Mém. Mosc.* xii. (1860) 117. t. 10. f. 2.—Persia.
cuneifolia, Lam. Encyc. i. 28 = A. Herba-rota.
cuspidata, Wall. Cat. n. 3230 = A. Millefolium.
cylindrica, Poir. Encyc. Suppl. i. 95 = nobilis.
damascena, DC. Prod. vii. 296 = falcata.
decipiens, Vest, in Flora, xii. (1829) i. Erg. 60 = nobilis.
decolorans, *Schrad. in Willd. Enum. Hort. Berol.* 913.—Europ.
decumbens, *Lam. Encyc.* i. 27.—Kamtschat.
dentato-serrata, *Heuffel, ex Heimerl, in Denkschr. Acad. Wien*, xlviii. (1884) 179.—Europ.
denticulata, Bess. ex Heimerl, l. c. 188 = sibirica.
dentifera, DC. Fl. Fr. Suppl. 485 = tanacetifolia.
dentifera, Reichb. Fl Germ. Exc 230 = A. Millefolium.
depressa, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. (1866) I. 405 = sibirica.
depressa, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 204.—Balkan.
distans, Waldst. & Kit. ex Willd. Sp. Pl. iii. 2207 = tanacetifolia.
diversifolia, Schur, Enum. Pl. Transs. 330 = crithmifolia.
dzimilensis, Boiss. Fl. Orient. iii. 259 = latiloba.
dracunculoides, Desf. Cat. Hort. Par. ed. III. 180 = A. Ptarmica.
Dracunculus, Linn. Sp. Pl. ed. Nov. ed. II. i. 18 = Artemisia Dracunculus!
dubiumlensis, C. Koch, in Linnæa, xxiv. (1851) 294 = latiloba.
dubia, *Fisch. ex Steud. Nom.* ed. II. i. 13.—Sibir.
Dumasiana x, *Vatke, in Oestr. Bot. Zeitschr.* xxii. (1872) 374.—Europ.
elegans, *Sch. Bip. in Flora*, xxxviii. (1855) 13.—Hispan.
erba-rota, 'Auct.' = A. Herba-rota.
eriadnica, Bertol. Fl. Ital. ix. 407 = nobilis.
erioclada, DC. Prod. vii. 296 = cretica.
eriophora, *DC. Prod.* vi. 31.—Persia.
Eupatorium, Bieb. Fl. Taur. Cauc. ii. 337 = filipendulina.
falcata, Lapeyr. Hist. Abr. Pl. Pyr. 534 = chamaemelifolia.
falcata, *Linn. Sp. Pl.* 897.—Orientia.
fasciculata, Burm. f. Fl. Cap. Prod. 27 = Eriocephalus umbellatus.
Feliciana x, *Bruegg. in Jahresb. Naturf. Ges. Graub. N. Fol.* xxiii.-xxiv. (1880) 105.—Europ.
filicifolia, Bieb. Fl. Taur. Cauc. ii. 336 = filipendulina.
fililoba, *Freyn, in Flora*, lxiv. (1881) 210.—Graecia.
filipendulina, *Lam. Encyc.* i. 27.—Orientia.
flabelifolia, Boiss. & Heldr. Diagn. Ser. I. xi. 37 = Tanacetum flabelliforme.
flabelliformis, Sibth. & Sm. Fl. Græc. ix. 73 = holosericea.
flava, Hort. ex Steud. Nom. ed. II. i. 13 = Chrysanthemum achilleæfolium.
floccosa, *Desf. Tab. Hort. Par.* 108 (nomen)—Orient.
formosa, *Sch. Bip. in Flora*, xxxviii. (1855) 13.—As. Min.
Frasili, *Sch. Bip. l. c.* xxv. (1862) *Beibl.* i. 159.—Graecia.

ACHILLEA :—

fragilis, Balb. ex DC. Prod. vi. 38 = A. Ptarmica.
fragrantissima, *Sch. Bip. in Flora*, xxxviii. (1855) 18. Orient; Aegypt.
frutescens, Willd. ex Ledeb. Fl. Ross. ii. 536 = Gerberi.
Genipi, Murr. App. Med. i. 168 = moschata.
Gerberi, Bieb. Fl. Taur. Cauc. i. 336 = nobilia.
Gerberi, *Willd. Sp. Pl.* iii. 2196.—Europ. ; Caucas.
glva, *Moench*, *Meth. Suppl.* 257.—Europ. ?
Glaveanae, (err. typ.) Schrank, Baier. Fl. ii. 409 = Clavennae
glomerata, Bieb. Fl. Taur. Cauc. iii. 585 = compacta.
gonlocephala, *Boiss. & Bal. in Boiss. Diagn.* Ser. II. vi. 98.—As. Min.
gracilis, Rafin. [' Fl. texensis, 12 '] Atl. Journ. 177 = A. Millefolium.
grandiflora, *Bieb. Fl. Taur. Cauc.* ii. 333.—As. Min.
grandifolia, *Friwald. in Flora*, xix. (1836), ii. 433.—Oriens.
grandifolia, Bieb. Fl. Taur. Cauc. ii. 444 = A. Ptarmica.
grandis, Fisch. ex Herd, in Bull. Soc. Nat. Mosc. (1865) i. 407 = A. Ptarmica.
grata, Fenzl, in Tchihat. Asie Min. Bot. ii. 261 = odorata.
Guszowii, Ces. & Pass. ex Nym. Consp. 362 = Anthemis abrotanifolia.
Haenbeana, Tausch, in Flora, iv. (1821), 567 = A. Millefolium.
Halleri, Crantz, Inst. i. 308 = strata.
Haussknechtiana, Aschers. in Festschr. Ges. Naturf. Fr. Berl. (1878) 243 = seq.
Haussknechtii, *Boiss. Fl. Orient.* iii. 284.—Persia.
helvetica, *Schleich. ex Gaud. Fl. Helv.* v. 368 = valesiaca ?
Herba-rota, *All. Fl. Pedem.* i. 180.—Europ. merid.
Herba-rotta, Vill. Prosp. Delph. 23 = prsec.
heterophylla, *Spreng, Nachr.* i. *Bot. Gart. Halle*, 9.—Sibir.
hispanica, Schrank, Fl. Rar. t. 70 = nobilia.
holosericea, *Sibth. & Sm. Fl. Grsec* t. 894.—Graecia.
hybrida, *Gaud. in Koch, Syn. Fl. Germ.* ed. II. 409.—Europ.
imbricata, Ancher, ex DC. Prod. vii. 296 = alcppica.
imbricata, Vent. Jard. Cels, t. 54 = membranacea.
impatiens, *Linn. Sp. Pl.* 898.—Sibir.
impunctata, Vest, in Flora, iii. (1820) 3, ? = moschata.
incisa, [*Clairv.*] *Man. Herb.* 251.—Europ
inodora, Linn. Sp. Pl. 900 = Lonas inodora.
intermedia, Schlich. Cat. Pl. (1821) 5 = A. Millefolium.
ircutiana, Sch. Bip. in Flora, xxxvii. (1855) 15 = A. Ptarmica.
Jaborneggi x, *Halacsy, in Oestr. Bot. Zeitschr.* xxvii. (1877) 45.—Europ.
Jacea, Griseb. Iter, ii. 264. ex Spicil. Fl. Rumel. ii. 216 = chrysocoma.
Jankæ, *Sch. Bip. ex Nym. Consp.* 366.—Europ.
japonica, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 125 = sibirica.
kellalensis, *Boiss. & Hausskn. in Boiss. Fl. Orient.* iii. 268.—Persia.
Kotschyi, Boiss. Diagn. Ser. II. iii. 19 = odorata.
Kraettliana x, *Bruegg. in Jahresb. Naturf. Ges. Graub. N. Fol.* xxii.-xxiv. (1880) 105, nomen.—Europ.
Laggeri, *Sch. Bip. in Oestr. Bot. Wochenschr.* vi. (1856) 300.—Europ.
lanata, Lam. Fl. Fr. iii. 640 = A. Millefolium.
lanata, *Spreng. Cat. Fl. Hal. ex Willd. Enum. Hort. Berol.* 915.—Hab. ?
lanulosa. *Nutt. in Journ. Acad. Philad.* vii. (1834) 86 = A. Millefolium.
latifolia, Salisb. Prod. 204 = macrophylla.
latiloba, *Ledeb. ex Nordm. in Bull. Acad. Sc. Pttersb.* ii. (1837) 213.—Caucas.
Ledebourii, Heimerl, in Flora, lxvi. (1883) 389 = impatiens.
lenensis, Turcz. ex DC. Prod. vii. 295 = inrsticana.
leptophylla, *Bieb. Fl. Taur. Cauc.* ii. 385.—Tauria.
leptophylla, C. Koch, ex Nym. Consp. 367 = crithmifolia.
Lerechei, *Sch. Bip. in Oestr. Bot. Wochenschr.* vi. (1856) 301.—Europ.
leucanthema, Pers. Syn. ii. 467 = A. Ptarmica.
ligustica, *All. Fl. Pedem.* i. 181. t. 52. f. 2.—Europ. ; Oriens.
ligustica, Vis. ex Nym. Consp. 367 = nobilia.
linearis, Stend. Nom. ed. II. i. 18 = A. Ptarmica.
linguiata, *Waldst. & Kit. Pl. Rar. Hung.* i. 2.—Europ. orient.
lissa, Scop. Delic. Insub. i. 6 = moschata.
lobata, *Forsk. Fl. Aegypt. Arab.* 218.—Aegypt.

ACHILLEA :—

lucida, Salisb. Prod. 204 = impatiens.
lycaonica, Boiss. & Heldr. Diagn. Ser. I. xi. 18 = conferta.
macrocephala, Rupr. Beitr. Pfl. Russ. Reiches, ii. (1845) 41 = A. Ptarmica.
macrophylla, Pill. & Mitterb. It. Posog. 111. t. 11 = Chrysanthemum macrophyllum.
macrophylla, *Linn. Sp. Pl.* 898.—Europ.
macrophylloides, Pand. ex Nym. Consp. 366 = A. grandifolia.
magna, Haenke, in Jiras. Beob. Riesengeb. 108 = A. Millefolium.
magna, All. Fl. Pedem. i. 184 = tanacetifolia.
magna, *Linn. Sp. Pl.* ed. II. 1267.—Europ.
magna, Sibth. & Sm. Fl. Graec. iii. 194 = ligustica.
major, *Heimerl, in Denkschr. Acad. Wien*, xlviii. (1884) 157.—Europ.
marginata, Turcz. ex Ledeb. Fl. Ross. ii. 532 = A. Millefolium.
maxima, Fischer, ex Heimerl, in Denkschr. Acad. Wien, xlviii (1884) = A. Ptarmica.
membranacea, *DC. Prod.* vi. 32.—Oriens.
Michelii, *Sch. Bip in Flora*, xxxviii. (1855) 14.—Europ.
micrantha, Bieb. Fl. Taur. Cauc. ii. 336 = pubescens.
micrantha, *Willd. Sp. Pl.* iii. 2209.—Soongar ; Oriens.
microloba, DC. Prod. vi. 32 = tereticifolia.
microphylla, Willd. Sp. Pl. iii. 2210 = odorata.
Millefolium, Ledeb. Fl. Alt. iv. 124 = setacea.
Millefolium, *Linn. Sp. Pl.* 899.—Europ. ; As. ; Am.
mollis, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 451, nomen.—Rossia.
mongolica, Fisch. ex Spreng. Nov. Prov. Hort. Hal. et Berol. (1813) i. 8 = alpina.
monocephala, *Boss. & Bal. Diagn.* Ser. I. v. 110.—Cilicia.
montana, *Schinch. ex DC. Prod.* vi. 21.—Europ.
monticola, Martr. Pl. Crit. du Tarn, 31 = A. Millefolium.
Morisiana, Reichb. Ic. Fl. Germ. xvi. 66. t. 1019 = A. Ptarmica.
moschata, *Jacq. Fl. Austriac.* v. 45.—Europ.
mucronata, Hort. ex Steud. Nom. ed. II. i. 13 = heterophylla.
mucronulata, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Europ.
multifida, Griseb. Spicil. Fl. Rumel. ii. 212 = strata.
multifida, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Graecia.
multifida, Sprun. ex Nym. Consp. 367 = setacea.
multiflora, *Hook. Fl. Bor. Am.* i. 318.—Am. bor.
multiplex, Remar. Fl. Orn. 78 = A. Ptarmica.
murcata, Ces. & Pass. ex Nym. Consp. 362 = Anthemis murcica.
muschensis, C. Koch, in Linnæa, xxv. (1851) 329 = vermicularis.
myriophylla, Willd. Enum. Hort. Berol. 916 = Tanacetum millefoliatum.
nana, *Lam.* Fl. Fr. ed. I. ii. 154 = A. Herba-rota.
nana, *Linn. Sp. Pl.* 899.—Europ.
Neumayeri, Fleidr. ex Heimerl, in Denkschr. Acad. Wien, xlviii (1884) 168 = Fraasii.
Neilreichii, Kern. in Oestr. Bot. Zeitschr. xxi. (1871) 148 = nobilis.
nitida, C. A. Mey. Syn. ii. 467 = A. Impatiens.
nivea, *Sprun. ex Nym. Consp.* 365.—Europ.
nobilis, *Linn. Sp. Pl.* 899.—Europ. ; Oriens.
nobilis, Koch. ex Nym. Consp. 367 = crithmifolia.
Novæ-Graecise, *Sch. Bip. in Flora* (1861), i. 158.—Enum.
obscura, *v. Nees, Ind. Sem. Hort. Bonn.* (1831) 1.—Europ.
occidentalis, Rafin. ex DC. Prod. vi. 24 = A. Millefolium.
ochroleuca, Ehrw. Skizze, 149 = A. Millefolium.
ochroleuca, Ehrh. Beitr. vii. 166 = nobilis.
ochroleuca, Waldst. & Kit. Pl. Rar. Hung. i. 33 = pectinata.
odorata, *Linn. Syst.* ed. X. 1225.—Europ. ; Oriens.
odorata, Fall. Reise, i. 279 = nobilis.
odorata, Wulf. in Jacq. Coll. i. 259 = setacea.
oligocephala, *DC. Prod.* vi. 32.—Mesopot.
orwiciensis, *Kit. & Kanitz, Relig. Kitaib.* 32, *in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 506.—Europ.
ossica, C. Koch, in Linnæa, xxiv. (1851) 328 = A. Millefolium.
oxydonta, *Boiss. Diagn.* Ser. I. xi 15.—Persia.

ACHILLEA :—

oxylepia, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 271.—Persia.
oxylobe, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Afr. austr.
pallescens, DC. Prod. vii. 298 = grandifolia.
pallescens, *Linn. Syst.* ed. X. 1225.—Hab. ?
pallida, Salisb. Prod. 204 = Tournefortii.
pannonica, Scheele, in Linnæa, xvii. (1885) 472 = Millefolium.
parthenifolia, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. (1865) i. 407 = A. Ptarmica.
parthenifolia, *Desf. ex Steud. Nom.* ed. II. i. 14, nomen.—Hab. ?
pascuifera, Lam. Encyc. i. 26. = Tanacetum Vahlii.
pectinata, Lam. Encyc. i. 28 = odorata.
pectinata, Willd. Sp. Pl. iii. 2197.—Hungar; Tauria.
Perosini, *Boiss. Fl. Orient.* iii 261.—As. Min.
pescedansifolia, Griseb. Spicil. Fl. Rumel. ii. 214 = grandifolia.
phrygia, *Boiss. & Bal. Diagn.* Ser. II vi 99.—As. Min.
pilosa, *Willd. Sp. Pl.* iii. 2196.—Hab. ?
pinnata, *Reichb. f. Ic. Fl. Germ.* xvi. 64; *Sch. Bip. in Flora*, xxxviii. (1855) 14.—Europ.
polyphylla, Schleich. ex Koch, Syn. Fl. Germ. ed. II. 411 = setacea.
pseudo-tanacetifolia, Wier. ex Reichb. Fl. Germ. xvi. 69 = magna.
pseudo-pectinata, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 179.—Turcia.
Ptarmica, *Linn. Sp. Pl.* 898.—Reg. temp. bor.
Ptarmica, Richards. in Frankl. 1st Journ. ed. II. App. 33 = multiflora.
ptarmicoides, Maxim. Prim. Fl. Amur. 154 = sibirica.
pubescens, Sibth. & Sm. Fl. Graec. t. 895 = micrantha.
pubescens, *Linn. Sp. Pl* 897.—Oriens.
pubescens, Willd Sp. Pl iii. 2206 = odorata.
pumila, Schur, Enum. Pl. Transs. 330 = tanacetifolia.
punctata, Moench, Meth. 603 = alpina.
punctata, Tenore, ex Fenzl, in Tchihat. Asie Min. Bot. ii. 264 = odorata.
pyrenaica, Sibth. ex DC. Fl. Fr. iv. 211 = A. Ptarmica.
ramosissima, Moench, Meth. Suppl. 256 = alpina ?
rectimarginata, Lapeyr. Hist. Abr. Pl. Pyr. 584 = chamaemelifolia.
Reichardtiana x, *G. Beck, in Sitzb. Zool. Bot. Ges. Wien*, xxxvii. (1878) 44.—Europ.
rigida, *Wallr. Beitr. Fl. Hercyn.* 381.—Europ.
rosea, Desf. Tabl. Hort. Par. ed. I. 105 ; ed. III. 181 = asplenifolia.
rosea, *Waldst. & Kit. Pl. Hung. Rar.*—Hungar.
rosmarinifolia, *Sch. Bip. in Flora*, xxxvii. (1855) 11.—Europ. occ.
rubra, Hildebr ex Steud. Nom. ed. II. i. 11 = dentifera.
rupestris, *Huter, ex Beck, in Denkschr. Acad. Wien*, xlviii (1882) 190.—Europ.
salicifolia, Bess. in DC. Prod. vi. 28 = A. Ptarmica.
salina, Schur, Enum. Pl. Transs. 328 = setacea.
sambucifolia, Desf. Cat. Hort. Par. ed. III. 180 = Chrysanthemum macrophyllum.
sambucina, Thonis, ex DC. Prod. vi. 58 = Chrysanthemum macrophyllum.
Santolina, Falk, Beitr. ii. 243 = Gerberi.
Santolina, C. A. Mey. Vers. Pfl. Cauc. 76 = tereticifolia.
Santolina, Hort. ex Willd. Sp. Pl. iii. 2200 = anthemoides.
Santolina, *Linn. Sp. Pl.* 896.—Orientes ; Afr. bor.
Santolina, Sibth. & Sm. Fl. Graec. ix. 70 = cretica.
Santolina, Wild. Tract. Ach. ex Steud. Nom. ed. I. 6 = tenuifolia.
santolinoides, *Leg. Gen. et Sp. Nov.* 80.—Hispan.
scabra, Host. Fl. Austr ii. 519 = A. Millefolium.
Schkuhrii, Spreng. ex Nym. Consp. 367 = nobilis.
Schurii, Sch. Bip. in Oestr. Bot. Wochenschr. vi. (1856) 300 = oxylobe.
Seidlii, J. & C. Presl, Fl. Cech. 172 = A. Millefolium.
semipectinata, Desf. Cat. Hort. Par. ed. III (1829) 408 = auriculata.
senecionifiora, *Vest, in Flora*, xii. (1829) *Erg.* i. 59 = dentifera.
serices, Janka, in Linnæa, xxx. (1859–60) 579 = compacta.
serica, Vis. ex Reichb. Fl. Germ. Excurs. ii. 228 = Clavennae.
serrata x, Retz. Obs. Fasc. ii. 25.—Sibir.
serrata, Sm. Engl. Bot. t. 2581 = decolorans.

ACHILLEA :—
serratifolia, Salisb. Prod. 204 = serrata.
serrulata, Hornem. Hort. Hafn. ii. 835 = A. Ptarmica.
setacea, *Waldst. & Kit. P. Rar. Hung.* i. 68. t. 66.—Europ. ; Orient.
setacea, Schwein. in Long. Exped. ii. 119 = A. Millefolium.
sibirica, *Ledeb. Ind. Sem. Hort. Dorpat*, (1811) ; *Fl. Ross.* ii. 528.—Reg. temp. bor.
sicula, Rafin. Prec. Déc. 40 ; et in Journ. Bot. ii. (1814) 273 = ligustica.
Siegfriedi ×, *Brugg. in Jahresb. Naturf. Ges. Graub. N. Fol.* xiii–xxiv. (1850) 58.—Europ.
silvatica, Auct. = sylvatica.
simplicicaulis, Boiss. Fl. Orient. iii. 259 = latilobа.
speciosa, Henckel, Adumbr. Fl. Hort. Hal. 1 = A. Ptarmica.
spinosa, *Schur, in Verh. Siebenb. Ver. Naturw.* x. (1859) 168.—Europ.
spinulifolia, *Fenzl, ex Tchihat. Asie Min. Bot.* ii. 268.—Taurus.
spinulosa, Schur, ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 185 = impatiens.
spithamea, *Cosa. & Dur. ex Cosa. in Ann. Sc. Nat.* Sér. IV. i. (1854) 239.—Algeria.
squarrosa, *Ait. Hort. Kew.* ed. I. iii. 242 (sp. dubia).—Hab. ?
squarrosa, Hassk. Cat. Pl. Hort. Bogor. 101 = sibirica.
stricta, Schleich. ex Koch, Syn. Fl. Germ. ed. T. 374 = magna.
subcartilagines, *Heimerl, in Denkschr. Acad. Wien*, xlviii. (1884) 185.—Europ.
subhirsuta, Gilib. Fl. Lituan. i. 217 = A. Millefolium.
sudetica, Opiz, in Hesperus (1818), 628, ex ej. Natural. n. vi. (1824) 49 = A. Millefolium.
sulphurea, *Boiss. Diagn.* Ser. I. xi. 17.—Syria.
sylvatica, Becker, Fl. Frankf. ii. 235 = A. Millefolium.
sylvatica, Tenore, Fl. Nap. iv. 126 = ligustica.
sylvestris, S. F. Gray, Nat. Arr. ii. 458 = A. Ptarmica.
Sermiferi, Fisch. & Mey. ex DC. Prod. vi. 26 = latilobа.
talagonica, *Boiss. Diagn.* Ser. I. xi. 17.—Persia.
tanacetifolia, *All. Fl. Pedem.* i. 183 = Europ. ; Orient.
tanacetifolia, Mill. Gard. Dict. ed. VIII. n. 7 = A. Millefolium !
taurica. Bieb. Fl. Taur. Cauc. i. 336 = leptophylla.
taygetea, *Boiss. & Heldr. Diag.* Ser. I. xi. 16.—Orient. tenuifolia, *Lam. Encyc.* i. 96.—Orient.
tenuifolia, Salisb. Prod. 204 = A. Millefolium.
tenuis, Schur, Enum. Pl. Transs. 329 = A. Millefolium.
teretifolia, *Willd. Sp. Pl.* iii. 2198.—Orient.
Thomasiana, *Hall. f. ex Hamilt, Bot. Mag.* 49.—Europ.
Thomasii, Thomas, ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 167 = prsec.
tomentosa, Frsas, ex Nym. Consp. 367 = holoserices.
tomentosa, Frivald. ex Nym. Consp. 368 = compacta.
tomentosa, *Linn. Sp. Pl.* 897.—Europ. ; Orient. ; As. bor.
tomentosa, Pall. Ind. Taur. ex Ledeb. Fl. Ross. ii. 536 = leptophylla.
tomentosa, Pursh, Fl. Am. Sept. 562 = A. Millefolium.
Tournefortii, *DC. Prod.* vi. 25.—Graecia.
Trautmanni ×, *Stein, in Oestr. Bot. Zeitschr.* xxviii. (1878) 249, nomen ; *cf. Heimerl, in Denkschr. Acad. Wien*, xlviii. (1882) 180.—Europ.
trichophylla, *Schrenk. Enum. Pl. Nov.* 48.—Soongaria.
trilinguulata, Sch. Bip. ex Boiss. Fl. Orient. iii. 271 = aleppica.
tyrolensis, Wender. in Flora, ix. (1826) 358 = Clavennae.
umbellata, *Sibth. & Sm. Prodr.* ii. 192.—Graecia.
valesiaca ×, *Suter, Fl. Helvet.* ii. 199.—Europ.
velutina, Desf. Cat. Hort. Par. ed. III. 180 = compacta.
vermicularis, *Trin. in Mém. Acad. Petersb.* vi. (1818) 494. t. 13.—Orient.
vermicularis, DC. Prod. vi. 23 = vermicularis.
verseonensis, *Kit. ex Kanitz, Reisp. Kiistk.* 93. *in Verh. Zool.-bot. Ges. Wien*, xiii. (1863) 506.—Europ.
virgata, Hort. ex DC. Prod. vi. 21 = tenuifolia.
virilis, *Reichb. f. in Fl. Germ.* xvi. t. 1018.—Europ.
viscosa, Lam. Fl. Fr. iii. 195 = A. Ageratum.
viscosa, *Sch. Bip. in Flora*, xxxviii. (1855) 14.—Europ.
Wilhelmsii, *C. Koch, in Linnaea*, xxiv. (1851) 838.—Orient.
Wolffii, Schur, ex Nym. Consp. 365 = impatiens.

ACHILLEOPSIS, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 9 = **Bulingia**, R. Br. (Sereul.).
densiflora, Turcz. l. c. 10 = R. densiflora.

ACHILLIOS, St. Lag. in Ann. Soc. Bot. Lyon. vii. (1880) 118 = **Achillea**, Linn. (Compos.).
millefoliatus, St. Lag. l. c. = A. Millefolium.

ACHIMENES, Vahl, Symb. Bot. ii. 71. (1791) = **Artemisia**, D. Don (Scroph.).
sesamioides, Vahl, l. c. = Art. sesamoides.

ACHIMENES, P. Br. Hist. Jamaic. 271. t. 20. f. 1. (1756). *GESNERACEAE*, Benth. & Hook. f. ii. 998.
CYRILLA, L'Hérit. Stirp. Nov. 147. t. 71 (1784).
DOLICHODIRA, Hanst. in Linnaea, xxvi. (1855) 205.
EUCODONIA, Hanst. l. c. 306.
GUTHNICKIA, Regel, in Flora, xxxii. (1849) 179.
HOUTTEA, Heyrh. Nom. ii. 306 (1846).
KORNICKIA, Regel, Ind. Sem. Hort. Petrop. (1857) 48 ; et Gartenfl. (1858) 309.
LOCHERIA, Regel, in Flora, xxxii. (1848) 251.
MANDULA, Decne. in Rev. Hortic. (1848) 468.
SCHEERIA, Seem. in Bot. Mag. t. 4743 (1853).
TREVIRANA, Willd. Enum. Hort Berol. 637 (1809).
amabilis, Decne. in Fl. des Serres, Sér. II. ii. (1857) 21. t. 1198 = Niegelia multiflora.
amoena, *Van Houtte, ex Hanst. in Linnaea*, xxvii. (1854) 718.—Mexico.
Andrieuxii, *DC. Prod.* vii. 536.—Mexico.
argyrostigma, Hook. Bot. Mag. t. 4175 = Koellikeria argyrostigma.
atrosanguinea, Lindl. in Journ. Hort. Soc. iii. (1848) 218 = fulgens.
autumnalis, *Hanst. in Linnaea*, xxvii. (1854) 784.—Guatemala.
candida, Lindl. in Journ. Hort. Soc. iii. (1848) 317 = Dicyrta candida.
cardinalis, *A. Distr. in Allg. Gartenz.* xv. (1847) 314.—Mexico.
coccinea, *Pers. Syn.* ii. 156.—Ind. occ.
cochinchinensis, Spreng. Syst. ii. 812 (= Diceros cochinchinensis).
cornifera, *DC. Prod.* vii. 536.—Am. austr.
cordata, Hort. ex Regel, in Flora, xxxii. (1849) 182 = grandiflora.
cornifera, Regel, in Bot. Mag. t. 719 = cornifera.
cupreata, Hook. Bot. Mag. t. 4812 = Episcia cupreata.
Diceros, Gmel. ex Steud. Nom. ed. I. 7. 278 (= Diceros cochinchinensis).
diversicolor, *Porpp. & Endl. Nov. Gen. et Sp.* iii. 9.—Peruv.
erinoides, *DC. Prod.* vii. 536.—Am. austr.
foliosa, *Morr. in Ann. Sc. Gand*, ii. (1846) 403. t. 91.—Mexico.
Georgeana, *Hort. ex Belg. Hortic.* x. (1860) 66. t. 5.—Hab. ?
Ghiesbrechtiana, Hort. ex Regel, in Flora, xxxi. (1849) 182 = grandiflora.
Ghiesbrechtii, Lindl. in Journ. Hort. Soc. v. (1850) 168 = grandiflora.
gloxinaeflora, Forkel, ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 318 = Gloxinia glabrata.
gracilis, Warcz. ex Regel, in Flora, xxxi. (1849) 180 = Tydaea candida.
grandiflora, *DC. Prod.* vii. 536.—Mexico.
Haageana, Otto, ex Regel, in Flora, xxxi. (1849) 182 = longiflora.
heterophylla, *DC. Prod.* vii. 536.—Bras.
hirsuta, *DC. l. c.*—Bras.
ichthyostoma, *Hanst. in Linnaea*, xxxiv. (1865–66) 434.—Bras.
ignescens, Lem. in Fl. des Serres, Sér I. iii. (1847) t. 248 = heterophylla.
incisa, *Klotzsch, ex Oerst. in Vidensk. Selsk. Shr.* V. v. (1861) 93.—Guatemala.
Jaurequiana, Warscz. ex Haage, in Fl. des Serres, Sér. I. v. (1849) t. 536 = longiflora.
Klotzschii, Paxt. Mag. Bot. xvi. (1849) 289 = longiflora.
Knightii, *Lem. in Fl. des Serres*, Sér. I. iv. (1848) t. 430.—Guatemala.
lanata, *Hanst. in Linnaea*, xxxiv. (1865–66) 434.—Mexico.
Liebmanni, *Lem. in Fl. des Serres*, Sér. I. iii. (1847) t. 262 ; *Misc.* 67 B.—Hab. ?
longiflora, *DC. Prod.* vii. 536.—Mexico.
maculata, Walp. Rep. ii. 719 = foliosa.
maculata, *Porpp. & Endl. Nov. Gen. et Sp.* iii. 8. t. 207 A.—Peru.
magnifica, Lindl. in Gard. Chron. (1856) 136 = Locheria magnifica.

ACHIMENES:—
Martensiana, Walp. Rep. ii. 719 = foliosa.
micrura, Lindl. in Journ. Hort. Soc. iii. (1848) 319 = Dicyrta Warscewiziana.
mollis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 9.—Chili.
multiflora, *Gardn. in Hook. Ic. Pl.* t. 498.—Bras.
nangelioides, *Van Houtte, Fl. des Serres*, Sér. II. viii. (1870) 117.—Hab. ?
ocellata, *Hook. Bot. Mag.* t. 4359.—N. Granat.
panamensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 475.—Panama.
patens, *Benth. Pl. Hartw.* 67.—Mexico.
pavoniana, *Oerst. in Vidensk. Selsk. Shr.* V. v. (1861) 91.—Costa Rica.
pedunculata, *Benth. Pl. Hartw.* 78.—Mexico.
picta, Benth. in Hook. Bot. Mag. t. 4126 = Tydaea picta.
purpurea, Pritzel, Ind. Ic. App. 3, sphalm. = seq.
pyropaea, Gordon, in Journ. Hort. Soc. ii. (1847) 268. t. 4. f. 1 = coccinea.
repens, Herb. Madr. ex Benth. in DC. Prod. x. 387 = Limnophila repens.
roseo, *Lindl. Bot. Reg.* (1841) 65.—Guatemala.
rupestris, Gardn. in Hook. Ic. Pl. t. 460.—Bras.
scabra, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 8.—Peruv.
Scherrii, *Hemsl. Biol. Centr.-Am. Bot.* ii. 476.—Mexico.
Shimeri, Gordon, in Journ. Hort. Soc ii. (1847) 293. t. 4. f. 2 = hirsuta.
superba, *Morits, ex Hanst. in Linnaea*, xxxiv. (1865–66) 436.—N. Granat.
tanella, *DC. Prod.* vii. 536.—Mexico.
tenuirima, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 9.—Peruv.
urticaefolia, *Poepp. & Endl. l. c.* 8. t. 207. B.—Peruv.
urticaefolia, Warcz. ex Regel, in Flora, xxxii. (1849) 181 = Dicyrta Warscewiziana.
Warscewiziana, *Regel, in Flora*, xxxii. (1849) 181.—Am. austr.

ACHIMUS, Poir. in Dict. Sc. Nat. v. 51 (1827) = **Strebilus**, Lour. (Urticac.).

ACHIRANTHES, P. Br. Hist. Jamaic. 180 (1756) = **Achyranthes**, Linn. (Amarantac.).

ACHIRIDA, Horan. Prod. Monog. Scitam. 18. t. 2. (1862) = **Canna**, Linn. (Scitam.).
iridiflora, Horan. l. c. = C. iridiflora.

ACHLAENA, Griseb. Cat. Pl. Cub. 228 (1866). *GRAMINEAE*, Benth. & Hook. f. iii. 1117.
piptostachya, *Griseb. l. c.* 229.—Cuba.

ACHLYS, DC. Syst. ii. 35 (1821). *BERBERIDEAE*, Benth. & Hook. f. i. 45.
japonica, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 51.—Japon.
triphylla, *DC. Syst.* ii. 35.—Am. bor.

ACHNATHERUM, Beauv. Agrost. 19. t. 6. f. 7 (1812) = **Stipa**, Linn. (Gramin.).
argentum, Beauv. l. c. 146 = S. Calamagrostis.
bromoides, Beauv. l. c. 146. 20 = S. Aristella.
Calamagrostis, Beauv. l. c. 146 = S. Calamagrostis.
capense, Beauv. l. c. 146 = Danthonia papillosa.
conspicuum, Beauv. l. c. 146. 20 = Arundo conspicua.
Hallerii, Beauv. l. c. 146 = S. Calamagrostis.
lanceolatum, Beauv. l. c. 146. 20 = Calamagrostis lanceolata.
multiaristum, Beauv. l. c. 146. 20 (= Urochne parviflora).
soboliferum, Beauv. l. c. 146. 20 = Muehlenbergia sobolifera.
speciosum, Sebast, in Nuov. Giorn. Bot. Ital. v. (1873) 317, nomen.—Europ.
tremiflorum, Beauv. Agrost. 146. 20 = Agrostis canina.

ACHNERIA, Beauv. Agrost. 72 (1812) = **Brizochaeta**, R. Br. (Gramin.).
brevifolia, Beauv. l. c. 73. 146 = E. mucronata.
capillaris, Beauv. l. c. 73. 146 = E. capillaris.
mucronata, Beauv. l. c. 73. 146 = E. mucronata.
obtusa, Beauv. l. c. 73. 146 = E. obtusa.

ACHNERIA, Munro, in Harv. Gen. Fl. Cap. ed. II. 449 (18 :8).
GRAMINEAE, Benth. & Hook. f. iii. 1168.

ACHNODON, Link, Hort. Berol. i. 65 (1827) = **Phleum**, Linn. (Gramin.).
arenarius, Link, i. c. = P. arenarium.
Bellardi, Link, i. c. = P. tenue.

ACHNODONTON, Beauv. Agrost. 24. t. 7. f. 5. (1812) = **Phleum**, Linn. (Gramin.).
arenarium, Trin. Gr. Unif. 166 = P. arenarium.
Bellardi, Beauv. Agrost. 25 = P. tenue.
tenuis, Beauv. l. c. = P. tenue.

ACHNOPHORA, F. Muell. in Trans. Roy. Soc. S. Austral. vi. (1883) 32. *COMPOSITAE.*
Tatei, *F. Muell. l. c.* 33.—Australia.

ACHRAS, Linn. Gen. ed. I. 365 (1737). *SAPOTACEAE*, Benth. & Hook. f. ii. 657.
SAPOTA, *Gaertn. Fruct.* ii. 108 t. 104 (1791).
arnhemica, F. Muell. ex Benth. Fl. Austral. iv. 280 = Sideroxylon arnhemicum.
australis, R. Br. Prod. 530 = S. australe.
Balata, Aubl. Pl. Gui. i. 308 = Mimusops Kauki.
brasiliensis, Willd. ex Steud. Nom. ed. II. i. 15 = Chrysophyllum brasiliense.
Brownlessiana, F. Muell. Fragm. vii. 111 = Sideroxylon Brownlessianum.
Cainito, Ruiz & Pav. Fl. Per. fri. 18 = Lucuma Cainito.
chartacea, F. Muell. ex Benth. Fl. Austral. iv. 281 = Sideroxylon chartaceum.
costata, Endl. Prod. Fl. Ins. Norf. 49 = S. costatum.
cotinifolia, F. Muell. in Nouv. Giorn. Bot. ital. (1871) 31 = Hormogyne cotinifolia.
cuneifolia, Poir. Encyc. vi. 534 = Sideroxylon obovatum!
dissecta, Linn. f. Suppl. 210; Forst. f. Plant. Esc. 48 = Mimusops Kauki.
elengioides, Bedd. Fl. Sylv. t. 285 = Sideroxylon tomentosum.
euphlebia, F. Muell. Fragm. vii. 110 = S. euphlebium.
ferruginea, Casar. Stirp. Bras. Dec. vii. 63 = A. Sapota.
glaucescens, Willd. ex Steud. Nom. ed. II. i. 15 = Lucuma Rivicoa.
Guapeda, Casar. Stirp. Bras. Dec. vii. 61 = L. laurifolia.
Howeana, F. Muell. Fragm. ix. 72 = Sideroxylon Howeanum.
laurifolia, Casar. Stirp. Bras. Dec. vii. 62 = Lucuma Casaretti.
laurifolea, F. Muell. ex Benth. Fl. Austral. iv. 282 = Sideroxylon Richardi.
Lucuma, Blanco, Fl. Filip. ed. I. 287 = Lucuma mammosa.
Lucuma, Ruiz & Pav. Fl. Per. fri. 17 = L. obovata.
mammosa, Bonpl. ex. Miq. in Mart. Fl. Bras. vii. 64 = L. Bonplandii.
mammosa, Linn. Sp. Pl. ed. II. 469 = L. mammosa.
mammosa, Sieber, ex A. DC. in DC. Prod. viii. 304 = Mimusops Kauki.
mammosa, Tuss. Fl. Antill. iii. t. 6 = Bassia Jusiaei.
montana, Poir. Encyc. vi. 533 = Diploön montana.
myrsinoides, A. Cunn. ex Benth. Fl. Austral. iv. 283 = Sideroxylon myrsinoides.
nigra, Poir. Encyc. vi. 532 = Diploön nigra.
obovata, F. Muell. ex Benth. Fl. Austral. iv. 283 = Sideroxylon Brownii.
octodecemfida, Stokes, Bot. Mat. Med. ii. 294 = Mimusops Kauki.
pallida, Poir. Encyc. vi. 533 = Sideroxylon pallidum.
pentagona, Poir. l. c. 533 = S. pentagonum.
Pohlmaniana, F. Muell. Fragm. v. 184 = Sideroxylon Pohlemannum.
retusa, Dennst. Schluss. Hort. Malab. 30 = Aglaia edulis.
retusa, Poir. Encyc. vi. 533 = Bumelia retusa.
rotundifolia, Poir. l. c. 534 = B. rotundifolia.
salicifolia, Linn. Sp. Pl. ed. II = Diploön salicifolia.
Sapota, *Linn. Sp. Pl.* ed. II. 470.—Am. austr.
setosa, *Schum. & Thonn. Beskr. Guin. Pl.* 176.—Afr. trop.
serrata, Poir. Encyc. Suppl. v. 86 = Prunus caroliniana.
sessiliflora, Poir. Encyc. vi. 531.—Ins. Borb.
tomentosa, Bedd. Forest. Man. Bot. 142 = Sideroxylon tomentosum.
venosa, Mart. Herb. Fl. Bras. 180 = Lucuma venosa.
verrucosa, Stokes, Bot. Mat. Med. ii. 294 = A. Sapota (?).
vitellina, Tuss. Fl. Antill. iv. 19 = Lucuma pauciflora.
xerocarpa, F. Muell. ex Benth. Fl. Austral. iv. 281 = Sideroxylon xerocarpum.
Zapota, Linn. Sp. Pl. ed. I. 192. 1190 = A. Sapota.
Zapotilla, Jacq. ex Nutt. Sylv. Am. iii. 90 = A. Sapota.

ACHROANTHES, Rafin. in Med. Repos. New York, v. (1808) 352 = **Microstylis**, Lindl. (Orchid.).
acuminata, Rafin. New Fl. Am. i. 59 = M. monophylla.
cilifolia, Rafin. l. c. = M. monophylla.
laxiflora, Rafin. l. c. 58 = M. ophioglossoides.
obtusifolia, Rafin. l. c. = M. ophioglossoides.
unifolia, Rafin. in Med. Repos. New York, v. (1808) 353 = M. ophioglossoides.

ACHROCHLOA, Griseb. Spicil. Fl. Rumel. ii. 446 (1845) = **Aircohlos**, Link (Gramin.).

ACHROMOLAENA, Cass. in Dict. Sc. Nat. lvi. 222 (1828) = **Cassinia**, K. Br. (Compos.).
viscosa, Cass. l. c. = C. quinqueseria.

ACHROSTACHYS, Benth. in Journ. Linn. Soc. xix. (1881) 134, err. typ. = **Athroostachys**, Benth. (Gramin.).

ACHRYSUM, A. Gray, in Hook. Kew Journ. iv. (1852) 228 = **Calocephalus**, R. Br. (Compos.).
glomeratum, A. Gray, l. c. 229 = C. multiflorus.

ACHUDEMIA, Blume, Mus. Bot. Lugd. Bat. ii. 57. t. 20. '1859' [1856]. *URTICACEAE*, Benth. & Hook. f. iii. 385.
japonica, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 241.—Japon.
javanica, *Blume, Mus. Bot. Lugd. Bat.* ii. 57. t. 20.—Java.

ACHYMUS, Vahl, ex Juss. in Dict. Sc. Nat. i. Suppl. 31 (1816) = **Bertolus**, Lam. (Urticac.).
pallidus, Boland. ex Blume, Mus. Bot. Lugd. Bat. ii. 79 = S. asper.

ACHYRACHAENA, Schau. Del. Sem. Hort. Vratisl. (1837) 3, ex Linnaea, xii. (1838) Litt. 87. *COMPOSITAE*, Benth. & Hook. f. ii. 396.
LEPIDOSTEPHANUS, Barth. Ind. Sem. Hort. Goett. (1837), ex Linnaea, xii. (1838) Litt. 82.
mollis, *Schau. l. c.* 3.— Am. bor. occ.

ACHYRANTHES, Linn. Gen. ed. I. 34 (1737). *AMARANTACEAE*, Benth. & Hook. f. iii. 35.
AMOGINNE, Rafin. New Fl. Am. iv. 44 (1836).
CADELARIA, Rafin. Fl Tellur. iii. 39 (1836).
CENTROSTACHYS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 497 (1835).
CODIVALIA, Rafin. Fl. Tellur. iii. 40 (1836).
ECLOTORIPA, Rafin. l. c. (1836).
STEIREMIS, Rafin. l. c. (1836).
abyssinica, *Nees, in Del. Sem. Hort. Vratisl.* (1850) 8. —Abyss.
alba, *Echl. & Zeyh. ex Moq. in DC. Prod.* xiii. ii. 311. —Afr. austr.
alopecuroides, Lam. Encyc. i. 548 = Aerva javanica.
alternifolia, Linn. Mant. i. 50 = Digera arvensis.
alternifolia, Linn. f. Suppl. 169 = Cyathula globosa.
alternifolia, Rumph. ex Wall. Cat. n. 6857 = Digera arvensis.
altissima, Jacq. Enum. Pl. Carib. 17 = Chamissoa altissima.
amaranthoides, Lam. Encyc. i. 548 = Digera arvensis.
angustifolia, *Benth. in Hook. Nig. Fl.* 492.—Afr. trop.
aquatica, *R. Br. Prod.* 417.—Ind. or.; Afr. trop.
arborescens, *R. Br. l. c.*—Ins. Norfolk.
argentea, Lam. Encyc. i. 545 = aspera.
aspera, Jacq. f. Ecl. 109. t. 74 = crispa.
aspera, Lam. Sp. Pl. 204.—Geront. trop.
aspera, Wall. Cat. n. 6924 H = bidentata.
atra, Pav. ex Moq. in DC. Prod. xiii. ii. 372 = Telanthera elongata.
atropurpurea, Lam. Encyc. i. 546 = Pupalia atropurpurea.
australis, *R. Br. Prod.* 417.—Austral.
avicularia, *E. Mey. ex Moq. in DC. Prod.* xiii. ii. 311.—Afr. austr.
axillaris, Willd. Enum. Hort. Berol. 270 = Alternanthera spicata.
baccata, Pav. ex Moq. in DC. Prod. xiii. ii. 350 = Chamissoa altissima.
bangalensis, *Lam. Encyc.* i. 459 (Sp. dub.).—Ind. or. bidentata, *Blume, Bijdr.* 545.—As. trop.
borbonica, *Willd. ex Roem. et Schult. Syst.* v. 549.—Ins. Borbon.
brachiata, Linn. Mant. 50 = Nothosaerva brachiata.
brasiliana, Thouin, ex Moq. in DC. Prod. xiii. ii. 360, 318 = Alternanthera procumbens.

ACHYRANTHES :—
calceolata, Russ. ex Wall. Cat. n. 6927 = Psilotrichum calceolatum.
Calesi, *Ibanta, in La Natural.* iii. (1879) 79. t. 8.—Am. centr.
canescens, *R. Br. Prod.* i. 417.—Austral.
capitata, *Forsk. Fl. Aegypt. Arab.* 48.—Arab.
capitata, Moq. in DC. Prod. xiii. ii. 310 = Involucrata.
capitulifora, *Bert. in Colla, Hort. Ripul.* 4. t. 18.—N. Granat.
carnosa, Wall. Cat. n. 6931 = Psilotrichum trichotomum
ciliata, Lam. Encyc. i. 549 = Digera arvensis.
conferta Ruiz, ex Seub. in Mart. Fl. Bras. v. i. 223 = Guillemines densa.
cordata, Hochst. in Flora, xxvii (1844) Beil. 6 = Psilostachys gnaphalobrya.
corymbosa, Heyne, ex Wall. Cat. n. 6918 = Celosia polygonoides.
corymbosa, Linn. Sp. Pl. 205 = Polycarpaea corymbosa.
crispa, *Desf. ex Poir. Encyc. Suppl.* ii. 10.—Am. austr.
cylindrica, Boj. ex Moq. in DC. Prod. xiii. ii. 378 = Cyathula cylindrica.
debilis, Poir. Encyc. Suppl. ii. 10 = Cyathula prostrata.
decumbens, Forsk. Fl. Aegypt. Arab. 47 = Celosia polygonoides.
diandra, *Roxb. Fl. Ind. ed. Carey*, ii. 504.—Zeylan.
dichotoma, *Linn. Mant.* 51.—Am. bor.
diffusa, Moench, Meth. 315 = Cyathula prostrata!
Digera, Poir. Encyc. Suppl. i. 2. 11 = Digera arvensis.
dioica, Pav. ex Moq. in DC. Prod. xiii. ii. 350 = Alternanthera flavescens.
echinata, Retz. Obs. Fasc. ii. 12 = Pupalia lappacea.
elliptici folia, Stokes, Bot. Mat. Med. i. 479 = aspera.
excelsa, Thou. ex Moq. in DC. Prod. xiii. ii. 315 = aspera.
fasciculata, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 179.—Afr. trop.
ferruginea, Roxb. Fl. Ind. ed. Carey, ii. 502 = Psilotrichum ferrugineum.
ficoidea, Lam. Encyc. i. 548 = Telanthera ficoidea.
filifolia, *Willd. ex Roem. et Schult. Syst.* v. 549.—Venezuela.
frumentacea, *Burm. f. Fl. Cap. Prod.* 7.—Afr. austr.
frutescens, Hort. ex Steud. Nom. ed. II. i. 15 = Digera arvensis.
fruticosa, Desf. Cat. Hort. Par. ed. III. 76 = aspera.
fruticosa, *Lam. Encyc.* i. 545.—Ind. or.
geminata, *Schum. & Thonn. Beskr. Guin. Pl.* 88.—Afr. trop.
geniculata, Pav. ex Moq. in DC. Prod. xiii. ii. 381 = Telanthera capitulifora.
glabella, *A. Gray, in Proc. Am. Acad.* vii. (1868) 300.—Ins. Sandv.
globosa, Pers. Syn. i. 259 = Cyathula globosa.
glauca, Heyne, ex Wall. Cat. n. 6927 = Psilotrichum calceolatum.
globulifera, Boj. ex Moq. in DC. Prod. xiii. ii. 329 = Cyathula globulifera.
grandifolia, *Moq. in DC. Prod.* xiii. ii. 318.—Java.
halimifolia, Lam. Encyc. i. 547 = Telanthera frutescens.
Heudelotii, Moq. in DC. Prod. xiii. ii. 310 = Pandiaka Heudelotii
hirtiflora, *A. Rich. Fl. Cub. Fanerog.* ii. 175.—Cuba.
hispida, Moq. in DC. Prod. xiii. ii. 312 = bidentata.
incana, *Fisch. ex Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 86.—Bras.
incana, Pav. ex Moq. in DC. Prod. xiii. ii. 365 = Telanthera frutescens.
incana, Roxb. Fl. Ind. i. 671; ed. Carey, ii. 495 = Aerva javanica.
indica, Mill. Gard. Dict. ed. VIII. n. 2 = aspera.
involucrata, Moq. in DC. Prod. xiii. ii. 310 = Pandiaka involucrata.
javanica, *Moq. l. c.* 312.—Java.
javanica, Pers. Syn. i. 259 = Aerva javanica.
lanata, Linn. Sp. Pl. 204 = Aerva lanata.
lanceolata, Klein, in Wall. Cat. n. 6928 = bidentata.
lanceolata, Zoll. & Mor. ex Moq. in DC. Prod. xiii. ii. 313 = javanica.
lappacea, Linn. Sp. Pl. 204 = Pupalia atropurpurea.
lappacea, Willd. Sp. Pl. i. 1193 = Pupalia lappacea.
leptostachya, E. Mey. in Hook. Lond. Journ. Bot. ii. (1843) 545 (err. typ. 445) = albs.
linarifolia, *Sw. in Vet. Akad. Handl. Stockh.* (1825) 48. —Ins. Mulucc.
Linkiana, Roem. & Schult. Syst. v. 545 = Chamissoa macrocarpa.

ACHYRANTHES:—
lupulina, Willd. ex Steud. Nom. ed. II. i. 15 = Telanthera lupulina.
mauritiana, *Moq. in DC. Prod.* xiii. II. 318.—Ins. Maurit.
mollis, Lepr. ex Seub. in Mart. Fl. Bras. v. 1. 230 = Cyathula prostrata.
mollis, *Schum. & Thonn. Beskr. Guin. Pl.* 137.—Afr. trop.
Monsoniae, Pers. Syn. i. 258 = Aerva Monsoniae.
montana, Klein, ex Wall. Cat. n. 6926 = Pallotrichum nudum.
Moorcroftiana, Hook. f. Fl. Brit. Ind. v. 8. lapsu, pro Axyrid = Exotia ceratoides.
mucronata, Herb. ex Moq. in DC. Prod. xiii. II. 248 = Allmania albida.
mucronata, Lam. Encyc. i. 547 = Alternanthera Achyrantha.
muricata, Linn. Sp. Pl. ed. II. 295 = Digera arvensis.
mutica, *A. Gray, ex H. Munn, in Proc. Am. Acad.* vii. (1867) 306.—Ins. Hawaii.
natans, Buch-Ham. ex Wall. Cat. n. 6982 = aquatica.
nervosa, *Roth, Nov. Pl. Sp.* 170.—Ind. or.
nitres, *Ait. Hort. Kew.* i. 296.—Ins. Canar.
nivea, Link, in Buch, Beschreib. Canar. 141 = Alternanthera Achyrantha.
nodiflora, Roxb. Fl. Ind. i. 678 = Allmania nodiflora.
nodosa, Bert. ex Moq. in DC. Prod. xiii. II. 252 = Alternanthera argentata.—Am. trop.
nodosa, *Vahl, in Schum. & Thonn. Beskr. Guin. Pl.* 139.—Afr. trop.
Novae-Brasiliae, Steud. Nom. ed. II. i. 15 = seq.
novae-brasilianae, Thouin, ex Roem. & Schult. Syst. v. 564 = Alternanthera procumbens.
nuda, Heyne, ex Wall. in Roxb. Fl. Ind. ed. Carey, ii. 507 = Pallotrichum nudum.
obcordata, Pav. ex Moq. in DC. Prod. xiii. II. 259 = Alternanthera Achyrantha.
obovatifolia, Stokes, Bot. Mat. Med. i. 480 = aspera.
obtusifolia, Lam. Encyc. i. 545 = aspera.
orbiculata, Heyne, in Roxb. Fl. Ind. ed. Carey, ii. 507 = Pupalia lappacea.
ovata, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 135.—Afr. trop.
paludosa, Bunb. in Proc. Linn. Soc. i. (1841) 109 = Telanthera philoxeroides.
paniculata, Forsk. Fl. Aegypt. Arab. 48 = Celosia candata.
papposa, Forsk. l. c. 48 = Exotia ceratoides.
patula, Linn. f. Suppl. 160 = Pupalia lappacea.
persica, Steud. Nom. ed. I. 8 = Aerva javanica.
picta, *Pauser. in Bot. Zeit.* (1864) 81.—Bras.
pilosa, Pav. ex Moq. in DC. Prod. xiii. II. 252 = Alternanthera argentata.
pilosiloides, Poit. ex Moq. l. c. 337 = Gossypianthus lanuginosus.
polygonata, Heyne, ex Wall. Cat. n. 6914 = Banalia thyrsiflora.
polygonoides, Heyne, ex Wall. l. c. n. 6926 = Pallotrichum nudum.
polygonoides, Lam. Encyc. i. 547 = Telanthera polygonoides.
polygonoides, Retz. Obs. Fasc. II. 12 = Digera arvensis.
polygonoides, R. Br. Prod. 416 = Telanthera polygonoides.
polystachya, *Forsk. Fl. Aegypt. Arab.* 48.—Arab.
porphyristachya, Wall. Cat. n. 6925 = aspera.
porrigens, Jacq. Hort. Schoenb. iii. 54 = Telanthera porrigens.
prostrata, Linn. Sp. Pl. ed. II. 296 = Cyathula prostrata.
prostrata, Willd. Sp. Pl. i. 1194 = Cyathula globosa.
pubescens, Roth, Nov. Pl. Sp. 171 = Aerva Monsonia.
pungens, Lam. Encyc. i. 546 = Aerva Monsonia.
purpurrea, Pav. ex Moq. in DC. Prod. xiii. II. 277 = Telanthera porrigens.
radicans, Cav. in Anal. Hist. Nat. Madr. iii. (1801) 97 = Alternanthera Achyrantha.
radicans, Heyne, in Roth, Nov. Pl. Sp. 170 = Aerva radicans.
repens, Ell. Sketch, i. 309 = Telanthera polygonoides.
repens, Heyne, in Roth, Nov. Pl. Sp. 167 = Cyathula prostrata.
repens, Linn. Sp. Pl. 205 = Alternanthera Achyrantha.
rivinaefolia, *Desf. Cat. Hort. Par.* ed. III. (1829) 390, 76.—Hab. ?
rosea, Spreng. Syst. iv. Cur. Post. 103 = Polycarpaea violacea.

ACHYRANTHES:—
rotundifolia, Klein, ex Wall. Cat. n. 6926 = Pupalia orbiculata.
rubrofusca, Wight, Ic. v. t. 1778 = aspera.
sanguinolenta, *Linn. Sp. Pl.* ed. II. 294.—Ind. or.
sarmentosa, Link, ex Seub. in Mart. Fl. Bras. v. 1. 241 = Chamissoa macrocarpa.
sarmentosa, *Vahl, Symb. Bot.* ii. 44.—Ind. or.
scandens, Hook. f. & Thoms. ex C. B. Clarke, in Journ. Linn. Soc. xv. (1877) 139 = Sclibanthus scandens.
scandens, Heyne, ex Moq. in DC. Prod. xiii. II. 248 = Iresine amaranatoides.
scandens, Roxb. Fl. Ind. i. 676.; ed. Wall. ii. 508 = Aerva scandens.
scandens, Thonn. ex Moq. in DC. Prod. xiii. II. 231 = Pupalia atropurpurea.
sequax, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 506 = Cyathula tomentosa.
sericea, *Heyne, ex Wall. Cat.* n. 6890.—Ind. or.
sericea, Roem. in Roxb. Fl. Ind. iii. 675; ed. Carey, ii. 507 = Pallotachya sericea.
serpyllifolia, Poir. Encyc. Suppl. ii. 11 = Telanthera serpyllifolia.
sessilis, Steud. Nom. ed. II. i. 16 = Alternanthera sessilis.
setacea, Roth, Nov. Pl. Sp. 168 = Aerva Monsonia.
sicula, Roth, Catalect. Bot. i. 39 = aspera.
spinosa, *Dorr. Hort. Dorion.* (1834–38).—Hab. ?
spinosa, Horsem. Hort. Hafn. i. 240 = Alternanthera spinosa.
splendens, *Mart. ex Moq. in DC. Prod.* xiii. II. 316.—Ins. Sandw.
stellata, Willd. Sp. Pl. i. 1195 = Polycarpaea stellata.
styracifolia, Lam. Encyc. i. 546 = Pupalia lappacea.
tenella, Pers. ex Moq. in DC. Prod. xiii. II. 241 = Iresine serpyllifolia.
tenuifolia, *Steud. Nom.* ed. II. i. 16 = aspera ?
tenuifolia, Willd. Sp. Pl. i. 1196 = Polycarpaea tenuifolia.
ternata, Host, ex Fisch. Mey. & Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 12 = Gomphrena decumbens.
Thomingii, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 137.—Afr. trop.
thyrsiflora, Heyne, ex Wall. Cat. n. 6914 = Banalia thyrsiflora.
tomentella, *Zipp. ex Span. in Linnaea*, xv. (1841) 345. —Malaya.
tomentosa, Roth, Nov. Pl. Sp. 167 = Cyathula tomentosa.
transvaa, Roxb. Fl. Ind. i. 678 = Alternanthera sessilis.
trichotoma, Perr. ex Moq. in DC. Prod. xiii. II. 322 = aquatica.
uncinulata, Willd. ex Roem. & Schult. Syst. v. 554 = Cyathula achyranthoides.
uncinulata, *Schrad. Ind. Sem. Hort. Goett.* (1838) 1.—Afr. austr.
velutina, *Hook. & Arn. Bot. Beech. Voy.* 68.—Ins. Society.
Verreauxii, Moq. in DC. Prod. xiii. II. 311 = alba.
Verschaffeltii, Lem. Illustr. Hortic. (1846) t. 409 = Iresine Herbstii.
verticillata, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1808) 26.—Afr. austr.
villosa, Blanco, Fl. Filip. ed. I. 189 = Alternanthera nodiflora.
villosa, Forsk. Fl. Aegypt. Arab. 48 = Aerva lanata.
violacea, Spreng. Syst. iv. Cur. Post. 102 = Polycarpaea violacea.
virgata, *[Desf.] Poir. Encyc. Suppl.* ii. 10.—Portorico.
virgata, Hort. ex Steud. Nom. ed. II. i. 16 = Celosia virgata.
virgata, Klein, ex Wall. Cat. n. 6915 = Banalia thyrsiflora.
Wightiana, Wall. Cat. n. 6938 = bidentata.

ACHYRASTRUM, Neck. Elem. i. 57 (1790) = **Hyoseris**, Linn. (Compos.).

ACHYROBACCHARIS, Sch. Bip. in Walp. Rep. ii. 952 (1843) = **Baccharis**, Linn. (Compos.).

ACHYROCLINE, Less. Syn. Comp. 332 (1889). *COMPOSITAE*, Benth. & Hook. f. ii. 306.
adoensis, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 438. —Abyss.
alata, *DC. Prod.* vi. 221.—Ecuad.
albicans, Griseb. in Goett. Abh. xxiv. (1879) 187 = satureioides.
argentina, *O. Hoffm. in Linnaea*, xliii. (1880–82) 185. —Reg. Argent.

ACHYROCLINE:—
batocana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 339.—Afr. trop.
bogotensis, *DC. Prod.* vi. 221.—N. Granat.
candicans, DC. l. c. = satureioides.
capitata, *Baker, in Mart. Fl. Bras.* vi. 111. 117 —Bras. caleioides, *DC. Prod.* vi. 221.—Peruv.
citrina, Griseb. in Goett. Abh. xxiv. (1879) 187 = satureioides.
coquimbensis, *Klatt, in Linnaea*, xlii. (1878–79) 112. —Chile.
disjuncta, *Hemsl. Bot. Voy. Challenger*, i. 11. 128.—Ins. Trinit.
flaccida, *DC. Prod.* vi. 220.—Bras.
flovescens, Griseb. in Goett. Abh. xix. (1874) 181 = alata.
glumacea, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 340.—Afr. trop.
hirta, *Klatt, in Linnaea*, xlii. (1878–79) 113.—N. Granat.
Hochstetteri, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 438.—Abyss.
latifolia, *Wedd. Chlor. And.* i. 148.—Boliv.
leptocephala, *DC. Prod.* vi. 220.—Madag.
lanuloides, *Vatke, in Linnaea*, xxxix. (1875) 489; *et in Oestr. Bot. Zeitschr.* (1875) 236.—Afr. trop.
madioides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 375.—Peruv.
mathioloefolia, DC. Prod. vi. 221 = satureioides.
mollis, Benth. Pl. Hartw. 207 = satureioides.
Moritziana, *Klatt, in Linnaea*, xlii. (1878–79) 112.—N. Granat.
pterocaula, DC. Prod. vi. 221 = alata.
rufescens, *DC. l. c.* 220.—N. Granat.
rugosa, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 19 = Stoechas tomentosa.
satureioides, *DC. Prod.* vi. 220.—Am. merid.
satureioides, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 424 = Stoechas Gardneri.
Schimperi, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 423.—Abyss.
sclerocladena, *Sch. Bip. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 341.—Afr. trop.
Steetzii, Vatke, in Oestr. Bot. Zeitschr. xxvii. (1877) 194 = batocana.
tomentosa, Schrank, in Denkschr. Akad. Muench. viii. (1821–22) 170 = Stoebe fusca.
Trianae, *Klatt, in Linnaea*, xlii. (1878–79) 113.—N. Granat.
Vargasiana, DC. Prod. vi. 220 = satureioides.
Vauthieriana, DC. l. c. = alata.
ventosa, *Klatt, in Linnaea*, xlii. (1878–79) 112.—Mexico.

ACHYROCOMA, Cass. in Dict. Sc Nat. v. 57 (1828) = **Vernonia**, Schreb. (Compos.).
lowentzon, Cass. l. c. xxvi. 21 = V. Achyrocoma.

ACHYROCOME, Schrank, in Denkschr. Akad. Muench. viii. (1821–22) 147 & 170 = **Elytropappus**, Sch. Bip. (Compos.).
ambigua, Schrank, l. c. = E. ambiguus.

ACHYRONIA, Royen, ex Linn. Gen. ed. II. 346 (1742) = **Aspalathus**, Linn. (Legumin.).

ACHYRONIA, Wendl. Obs. Bot. 39. t. 12. (1798) = **Priestleya**, DC. (Legum.).
arborea, Spreng. Syst. iii. 385, iv. Cur Post. 271, 387 = Carmichaelia australis.
villosa, Wendl. Obs. Bot. 39 = P. Thunbergii ?

ACHYROPAPPUS, Torr. & Gray, in Proc. Am. Acad. vii. (1868) 380. *ILLECEBRACEAE*, Benth. & Hook. f. iii. 15.
Cooperi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 331. —N. Mexico; Calif.
Palmeri, Hook. f. in Benth. & Hook. f. Gen. iii. 15 = Parryi.
Parryi, *Hemsl. Diag. Pl. Nov.* 36.—Mexico.

ACHYROPAPPUS, Bieb. ex Fisch. in DC. Prod. vi. 568 (1838) = **Trichoclepis**, DC. (Compos.).

ACHYROPAPPUS, H. B. & K. Nov. Gen. et Sp. iv. 257. t. 390 (1820) = **Schkuhria**, Roth (Compos.).
anthemoides, H. B. & K. l. c. = Bahia anthemoides.
meritimus, Poepp. ex DC. Prod. v. 687 = Bahia ambrosioides.

ACHYROPAPPUS :—
pedatus, Less. Syn. Comp. 289 = Florestina pedata.
schkuhrioides, *D. Don, ex Hook. & Arn. in Hook. Lond. Journ. Bot.* iii. (1840) 322.—Hab.?
schkuhrioides, Link & Otto, Abh. Neuer Gew. Gart. Berl. t. 80 = Schkuhria senecioides.
Woodhousei, A. Gray, in Proc. Am. Acad. vi. (1864-5) 546 = Bahia Woodhousei.

ACHYROPHORUS, Adans. Fam. ii. 112 (1763) = **Hypochoeris**, Linn. (Compos.).
acaulis, Remy, in C. Gay, Fl. Chil. iii. 448.—Chili.
albiflorus, *Sch. Bip. in Pollichia*, xvi-xvii. (1859) 52.—N. Granat.
alpicolus, Schur, Enum. Pl. Transs. 865 = H. maculata.
ambiguus, DC. Prod. vii. 93 = H. cretensis.
andinus, DC. l. c. 92 = H. andina.
angustissimus, Phil. in Linnæa, xxxiii. (1864-65) 127.—Chili.
asperioides, DC. Prod. vii. 94 = H. asperioides.
araucanus, Phil. in Linnæa, xxviii. (1856) 721.—Chili.
arenarius, DC. Prod. vii. 95 = H. arenaria.
auranticus, DC. l. c. 93 = H. grandiflora.
Balbisii, Hornem. Hort. Hafn. Suppl. 91 = H. Balbisii.
barbatus, Sch. Bip. in Pollichia, xvi-xvii. (1859) 48.—N. Granat.
Bipontinus, Sch. Bip. l. c. 54 = H. Gardneri.
Botsianus, Walp. Ann. ii. 956.—Creta.
brasiliensis, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 106 = H. brasiliensis.
brasiliensis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 128 = H. Gardneri.
brevicaulis, Phil. in Linnæa, xxxiii. (1864-65) 127.—Chili.
chilensis, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 104.—Chili.
chondrilloides, A. Gray, in Proc. Am. Acad. v. (1845) 145 = H. brasiliensis.
chrysanthus, DC. Prod. vii. 94 = H. chrysantha.
ciliatus, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 128 = H. grandiflorus.
clarionoides, Remy, in C. Gay, Fl. Chil. iii. 450.—Chili.
coronopifolius, Sch. Bip. in Flora, xxxviii. (1855) 123. —Reg. Magell.
cryptocephalus, Sch. Bip. in Lechl. Berb. Am. Austr. 57.—Peruv.
elatus, Wedd. Chlor. And. i. 223 = H. elata
discolor, Godr. in Mém. Acad. Stanislas, Sér. III. [xvii.] 1859 (1854), 407.—Hab.?
eriolaenus, Sch. Bip. in Lechl. Berb. Am. Austr. 57.—Peruv.
foliosus, Phil. Fl. Atac. 29.—Chili.
Gardneri, Sch. Bip. in Pollichia, xvi-xvii. (1859), 64 = H. Gardneri.
glabrescens, Steud. Nom. ed. II. i. 16 = H. arenaria.
glaucus, Phil. Fl. Atac. 29 = H. andina.
graminifolius, Sch. Bip. ex Walp. Rep. vi. 833 = tenuifolius.
grandiceratus, Phil. Fl. Atac. 29.—Chili.
grandiflorus, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 113.—Sibiria.
grandiflorus, Ledeb. Fl. Ross. ii. 777 = H. grandiflora.
helveticus, Hornem. Hort. Hafn. ii. 772 = H. uniflora.
helveticus, Less. Syn. Comp. 130 = H. helvetica.
hispidus, Hornem. Hort. Hafn. ii. 772 = H. hispida.
Hohenackeri, Sch. Bip. in Lechl. Berb. Am. Austr. 57. —Peruv.
Humboldtii, Sch. Bip. in Pollichia, xvi-xvii. (1859) 52. —N. Granat.
humilis, Phil. in Anales Univ. Chil. (1873) 482.—Chili.
incanus, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 100.—Am. austr.
Yunckeri, Sch. Bip. l. c. 96.—Chili.
Lechleri, Sch. Bip. in Lechl. Berb. Am. Austr. 54.—Chili.
Lessingii, Sch. Bip. l. c. 98.—Chili.
maculatus, Scop. Fl. Carn. ed. II. ii. 116 = H. maculata.
Meyerianus, Walp. Rep. vi. 836 = H. Meyeriana.
microcephalus, Sch. Bip. in Pollichia, xvi-xvii. (1859), 59 = H. brasiliensis.
microphyllus, Remy, in C. Gay, Fl. Chili. iii. 449.—Chili.
minimus, Hornem. Hort. Hafn. ii. 773 = H. minima.
montanus, Phil. in Linnæa, xxviii. (1856) 720.—Chili.
oderatus, Walp. Rep. vi. 836.—Chili.
ovalentus, DC. Prod. vii. 92 = H. cretensis.
paleatris, Phil. in Anales Univ. Chil. (1865) ii. 317 — Chili.

ACHYROPHORUS :—
petiolaris, DC. Prod. vii. 94 = H. petiolaris.
pinnatifidus, DC. l. c. 93 = H. pinnatifida.
praemorsus, Webb, ex Walp. Rep. vi. 829 = Webbii.
psychrophilus, Wedd. Chlor. And. i. 222.—Chili.
quitensis, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 130.—Ecuad.
radicatus, Scop. Fl. Carn. ed. II. ii. 117 = H. radicata.
ramosus, DC. Prod. vii. 92.—Hab.?
Robertia, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 137.—Italia.
roseus, Less. in Linnæa, v. (1830) 183. t. 2. f. 5 = Pinaropappus roseus.
sagittatus, Phil Sert. Mend. alt. 20.—Chili.
Scorzonerae, DC. Prod. vii. 94 = H. Scorzonerae.
Sellei, Sch. Bip. in Pollichia, xvi-xvii. (1859), 60 = H. brasiliensis.
sericoides, DC. Prod. vii. 93 = H. cretensis.
sessiliflorus, DC. l. c. 95 = H. sessiliflora.
setosus, Wedd. Chlor. And. i. 220.—Peruv.
sonchoides, DC. Prod. vii. 95 = H. sonchoides.
spathulatus, Remy, in C. Gay, Fl. Chil. iii. 447.—Chili.
spinulger, Phil. in Linnæa, xxviii. (1856) 721.—Chili.
stenocephala, A. Gray, ex Wedd. Chlor. And. i. 221.—Peruv.
taraxacifolius, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 101 = H. radicata.
taraxacoides, Walp. Rep. vi. 836.—Peruv.
tenerifolius, Remy, in C. Gay, Fl Chil. iii. 446.—Chili.
tenuifolius, DC. Prod. vii. 94 = H. tenuifolia.
tenuisectus, Sch. Bip. in Pollichia, xvi-xvii. (1859), 57 = H. variegata.
thrincoides, Remy, in C. Gay, Fl. Chil. iii. 445.—Chili
trichocephalus, Sch. Bip. in Pollichia, xvi-xvii. (1859) 57 = H. variegata.
uniflorus, F. W. Schmidt, Samml. Phy. Aufs. i. (1795) 294 = H. uniflora.
variegatus, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 119 = H. variegata.
Volchmanni, Phil. in Anales Univ. Chil. (1873) 482.—Chili.
Webbii, Sch. Bip in Nov. Act. Nat. Cur. xxi. i. (1845) 97.—Ins. Falk.

ACHYROPSIS, Benth. & Hook. f. Gen. iii. 36 (1880). *AMARANTACEAE*, Benth. & Hook. f. l. c. acicularis, *Benth. & Hook. f. l. c.*—Afr. austr.
alba, *Benth. & Hook. f. l. c.*—Afr. austr.
leptostachya, *Benth. & Hook. f. l. c.*—Afr. austr.

ACHYROSERIS, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 165 = **Soonsonera**, Linn. (Compos.).
macrospermum, Sch. Bip. l. c. = S. macrosperma.

ACHYROSPERMUM, Blume, Bijdr. 840 (1826). *LABIATAE*, Benth. & Hook. f. ii. 1208.
LAMPROSTACHYS, *Boj. ex Benth. in DC. Prod.* xii. 457 (1848).
SIPHOTORIS, *Boj. ex Benth. l. c.* (1848).
aethiopicum, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 56.—Afr. trop.
densiflorum, *Blume, Bijdr.* 841.—Ins. Malay.
frutcosum, *Benth. Lab. Gen. et Sp.* 643 & 745.—Madag.
philippinense, Benth. in DC. Prod. xii. 458 = densiflorum.
phlomoides, *Blume, Bijdr.* 841.—Java.
seychellarum, *Baker, Fl. Maurit.* 250.—Ins. Seychelles.
Wallichianum, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 573.—Reg. Himal.

ACHYROSTEPHUS, Kunze, ex Reichb. Consp. 212* (1828). (Compos.).

ACIA, Schreb. Gen. ii. 458 (1791) = **Couepia**, Aubl. (Rosac.).
amara, Willd. Sp. Pl. iii. 717 = C. guianensis.
dulcis, Willd. l. c. = C. guianensis.

ACIACHNE, Benth. in Hook. Ic. Pl. t. 1362 (1881). *GRAMINEAE*, Benth. & Hook. f. iii. 1143.
pulvinata, *Benth. l. c.*—N. Granat.

ACIANTHERA, Scheidw. in Otto & Dietr. Allgem. Gartenz. x. (1842) 292. = **Pleurothallis**, R. Br. (Orchid.).
punctata, Scheidw. l. c. = P. Acianthera.

ACIANTHUS, R. Br. Prod. 321 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 611.
stenalus, *Reichb. f. in Linnaea*, xli. (1877) 56 — N. Caled.
bifolius, R. Br. Prod. 322 = Chiloglottis diphylla.
caudatus, *R. Br. l. c.* 321.—Austral.
elegans, *Reichb. f. in Linnaea*, xli. (1877) 56.—N. Caled.
exsertus, *R. Br. Prod.* 321.—Austral.
fornicatus, *R. Br. l. c.*—Austral.
petiolatus, D. Don, Prod. Fl. Nep. 29 = Liparis nepalensis.
rivularis, *A. Cunn. in Hook. Comp. Bot. Mag.* ii. (1836) 876.—N. Zel.
Sinclairii, *Hook. f. Fl. N. Zeal.* i. 246.—N. Zel.
viridis, *Hook. f. Fl. Tasm.* ii. 272.—Austral.

ACIALYPTUS, A. Gray, Bot. U. S. Expl. Exped. t. 551. t. 67. (1854). *MYRTACEAE*, Benth. & Hook. f. i. 708.
Fallgarii, *F. Muell. Fragm.* viii. 15.—Ins. D. Howe.
myrtoides, *A. Gray, Bot. U. S. Expl. Exped.* i. 551. t. 67.—Ins. Fiji.
myrtoides, Seem. in Bonplandia, ix. (1861) 206 = Seemanni.
nitida, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 186; *et in Ann. Sc. Nat.* Sér. V. iii. (1865) 227.—N. Caled.
Seemanni, *A. Gray, in Bonplandia*, x. (1862) 85.—Ins. Fiji.

ACICARPA, Raddi, Agrost. Bras. 31 (1823) = **Panicum**, Linn. (Gramin.).
sacchariflora, Raddi, l. c. t. 1. f. 4 = P. leucophaeum.

ACICARPA, R. Br. in Trans. Linn. Soc. xii. (1818) 182 = seq.

ACICARPHA, Juss. in Ann. Mus. Par. ii. (1803) 347. t. 58. *CALYCERAEAE*, Benth. & Hook. f. ii. 162.
ACANTHOSPERMA, Vell. Fl. Flum. lc. viii. t. 152 (1827).
CRYPTOCARPHA, Cass. in Bull. Soc. Philom. (1817) 34.
ECHINOLEMA, Jacq. f. ex DC. Prod. v. 3 (1836).
SOMMEA, Bory, in. Ann. Gén. Sc. Phys. vi. (1820) 92. t. 87.
buplauroides, Less. in Linnæa, vi. (1831) 527 = Boopis buplauroides.
crassifolia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 402.—Bras.
lanata, *Lag. ex Pers.* Syn. ii. 488.—Mexico.
pinnatifida, *Miers, in Ann. & Mag. Nat. Hist.* Ser III. vi. (1860) 404.—Bras.
procumbens, *Less. in Linnaea*, vi. (1831) 527.—Bras.
runcinata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 404.—Bras.
spathulata, *R. Br. in Trans. Linn. Soc.* xii. (1818) 129.—Bras.
tribuloides, *Juss. in Ann. Mus. Par.* ii. (1803) 348. t. 58.—Reg. Argent.

ACICARPHAEA, Walp. Ann. ii. 877 (1852) = **Chaenactis**, DC. (Compos.).
artemisiaefolia, Walp. l. c. = C. artemisiaefolia.

ACICLINIUM, Torr. & Gray, Fl. N. Am. ii. 281 in adnot. (1841) = **Bigelowia**, DC. (Compos.).

ACIDANDRA, Mart. ex Spreng. Gen. i. 388 (1830) = **Zollernia**, Mart. (Legumin.).

ACIDANTHERA, Hochst. in Flora, xxvii. (1844) 25. *IRIDEAE*, Benth. & Hook. f. iii. 706.
SPHAEROSPORA, Klatt, in Linnaea, xxxii. (1863) 725.
HOUTTUYNIA, Houtt. Handl. Plantenk. xii. 448. t. 85 (1780).
aequinoctialis, *Baker, in Journ. Linn. Soc.* xvi. (1877) 160.—Afr. trop.
bicolor, *Hochst. in Flora*, xxvii. (1844) 25.—Abyss.
brachystachys, *Baker, in Journ. Bot.* xiv. (1876) 338.—Afr. trop.
brevicollis, *Baker, l. c.* 339.—Afr. trop.
exscapa, Baker, ex Bouché & Wittm. in Berl. Monatschr. Bef. Gartenb. xix. (1876) 15 = tubulosa.
flexuosa, *Baker, ex Bouché & Wittm. l. c.*—Afr.

ACIDANTHERA :—
graminifolia, *Baker, in Journ. Bot.* xiv. (1876) 338.—Afr. trop.
Huttoni, *Baker, l. c.* 339.—Afr. trop.
platypetala, *Baker, l. c.*—Afr. austr.
tubulosa, *Baker, in Journ. Linn. Soc.* xvi. (1877) 160.—Afr. trop.
unicolor, *Hockst. ex Baker, l. c.*—Abyss.

ACIDOCROTON, Griseb. Fl. Brit. W. Ind. 49 (1864).
EUPHORBIACEÆ, Benth. & Hook. f. iii. 391.
adelioides, *Griseb. l. c.*—Ind. occ.

ACIDOTON, Sw. Prod. Veg. Ind. Occ. 88 (1788).
EUPHORBIACEÆ, Benth. & Hook. f. iii. 326.
innocua, *Baill. Etud. Gén. Euphorb.* 402. t. 18. f. 10-11.—Ind. occ.
urens, Sw. *Prod. Veg. Ind. Occ.* 88.—Jamaica.

ACILEPIS, D. Don, Prod. Fl. Nep. 169 (1825) = Vernonia, Schreb. (Compos.).
cirsiifolia, Spreng. Syst. iii. 387 = Pacourina cirsiifolia.
lanata, Spreng. ex DC. Prod. vii. 26 = Dicoma tomentosa.
squarrosa, D. Don, Prod. Fl. Nep. 169 = V. teres.

ACINAX, Rafin. Fl. Tellur. iv. 58 (1836) = Costus, Linn. (Scitam.).

ACINETA, Lindl. Bot. Reg. (1843) Misc. 67. *ORCHIDEÆ*, Benth. & Hook. f. iii. 553.
NEIPPERGIA, Morr. in Ann. Soc. Agr. Gand, v. (1849) 375. t. 282.
Barkeri, *Lindl. in Bot. Reg.* (1843) *Misc.* 68.—Mexico.
chrysantha, *Lindl. & Paxt. Flow. Gard.* i. (1850-51) 81.—Am. centr.
cryptodonta, *Reichb. f. in Bonplandia*, ii. (1854) 92.—Am. centr.
dense, *Lindl. & Paxt. Flow. Gard.* i. 91 (1850-51).—Am. centr.
erythroxantha, *Reichb. f. in Bonplandia*, ii. (1854) 93.—Hab.?
fulva, Klotzsch, in Otto & Dietr. Allg. Gartenz. (1859) 146 = Humboldtii.
glauca, Linden, ex Lindl. & Paxt. Flow. Gard. i. (1850-51) 123 = Cynoches Pescatorei.
Hrubyana, *Reichb. f. in Gard. Chron.* N. S. xviii. (1882) 11. 109.—N. Granad.
Humboldtii, *Lindl. Bot. Reg.* (1843) *Misc.* 68.—Am. centr.
longicapa, Reichb. f. in Bot. Zeit. x. (1852) 706 = Lacaena bicolor.
Schilleriana, *Reichb. f. in Walp. Ann.* vi. 609.—Hab.?
Sella-turcica, *Reichb. f. in Bot. Zeit.* x. (1852) 705.—Am. centr.
sulcata, *Reichb. f. in Gard. Chron.* N. S. xi. (1879) 1. 653.—Ecuador?
superba, Reichb. in Walp. Ann. vi. 609 = Humboldtii.
Warszewiczii, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 145 = dense.

ACINODENDRUM, Linn. Gen. ed. I. 129 (1787) = Melastoma, Burm.!
aurea, Rafin. Sylva Tellur. 94.—Hab.?
cymosa, Rafin. l. c. = Amphiblema cymosum?
lanifera, Rafin. l. c.—Hab.?

ACINOLIS, Rafin. Sylva Tellur. 95 (1838) = Melastoma, Burm.
elliptica, Rafin. l. c. = M. ellipticum?

ACINOS, Rupp. Fl. Jen. ed. Hall. 288 (1745); Moench, Meth. 407 (1794) = Calamintha, Lam. (Labiat.).
arvensis, Host, Fragm. Flor. Austr. 1. (1827) 3. t. 2.—Europ.
graveolens, Link, Enum. Hort. Berol. ii. 117 = C. graveolens.
Herba-barona, G. Don, in Loud. Hort. Brit. 389 = Thymus Herba-Barona.
hirsutus, Host, Fl. Austr. ii. 181. (err. typ. 151) = C. alpina.
incanus, Griseb. Spicil. Fl. Rumel. ii. 128 = C. mariti-mus.
multiflorus, C. Koch, ex Boiss. Fl. Orient. iv. 579 = C. umbrosa.
patavinus, Pers. Syn. ii. 181. (err. typ. 151) = C. alpina.
purpurascens, Pers. l. c. = C. graveolens.
rotundifolius, Frivald. ex Walp. Rep. iii. 780 = C. alpina.
rotundifolius, Pers. Syn. ii. 181. (err. typ. 151) = C. Nepeta.
suaveolens, G. Don, ex Loud. Hort. Brit. 389 = C. graveolens.
thymoides, Moench, Meth. 407 = C. Acinos.
transsilvanica, Schur, Enum. Pl. Transs. 529 = C. Nepeta.
villosus, Pers. Syn. ii. 181. (err. typ. 151) = C. Acinos.
vulgaris, Pers. l. c. = C. Acinos.

ACINOTUM, Reichb. Handb. 260 (1837) = Mathiola, R. Br. (Crucif.).

ACIOA, Aubl. Pl. Gui. ii. 698. t. 280 (1775) = Couepia, Aubl. (Rosac.).
emarg, Nord. Nom. ed. I. 9 = C. guianensis.
Bellayana, Baill. Adansonia, vii. (1866-67) 324 = Griffonia Bellayana.
dulcis, Steud. Nom ed. I. 9 = C. guianensis.
guianensis, Aubl. Pl. Gui. ii. 698 = C. guianensis.
Icondere, Baill. Adansonia, vii. (1866-67) 323 = Griffonia Icondere.
Glaucon, Baill. l. c. 324 = Griffonia pallescens.
Smeathmanni, Baill. l. c. x. (1872) 244.—Afr. trop.

ACIOTIS, D. Don, in Mem. Wern. Soc. iv. (1823) 300.
MELASTOMACEÆ, Benth. & Hook. f. i. 744.
SPENNERA, Mart. in DC. Prod. iii. 115 (1828).
acuminifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Am. trop.
acuminifolia, Triana, l. c. = sphaeranthera, etc.
acutiflora, *Triana, l. c.* 52. pro parte.—Am. trop.
acutiflora, Triana, l. c. pro parte = laxa.
aequatorialis, *Cogn. in Mart. Fl. Bras.* xiv. iii. 464.—Bras.
amazonica, *Cogn. l. c.* 461.—Bras.
amœna, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Am. merid.
aquatica, *G. Don, in Loud. Hort. Brit.* 174.—Ins. St. Vincent.
brachyborria, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Bras.
caudiculata, *Triana, l. c.* 52.—Peruv.
circumscissa, *Triana, l. c.*—Bras.; Guiana.
dichotoma, *Cogn. in Mart. Fl. Bras.* xiv. iii. 460.—Guiana.
discolor, *D. Don, in Mem. Wern. Soc.* iv. (1823) 301.—Ins. St. Vincent.
dysophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Bras.; Peruv.
ferruginea, Triana, l. c. 51 = brachyborria.
fragilis, *Cogn. in Mart. Fl. Bras.* xiv. iii. 475.—Guiana.
herbacea, *Cogn. l. c.* 468.—Bras.
imberon, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Bras.; Guiana
latifolia, *Cogn. in Mart. Fl. Bras.* xiv. iii. 464.—Bras.
laxa, *Cogn. l. c.* 476.—Bras.
Loyanea, *Cogn. l. c.* 460.—Nicarag.
longifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1872) 52.—Bras.
Martiana, *Cogn. in Mart. Fl. Bras.* xiv. iii. 468.—Bras.
paludosa, *Triana, in Trans. Linn. Soc.* xxviii. (1872) 51.—Bras.

ACIOTIS:—
pendulifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Guiana.
polystachya, *Triana, l. c.* 52.—Bras.
purpurascens, *Triana, l. c.* 52. t. 3. f. 36.—Am. trop.
rostellata, *Triana, l. c.* 51.—Mexico.
rubricaulis, *Triana, l. c.* 52. t. 3. f. 35 b.—Bras.
Siebori, Triana, l. c. 52 = Nepsera aquatica.
silenifiora, *Triana, l. c.* 52.—Bras.
sphaeranthera, *Cogn. in Mart. Fl. Bras.* xiv. iii. 458.—Peruv.
uliginosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Venezuela.
viscosa, *Triana, l. c.* 52.—Am. trop.

ACIPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1846) i. 577 = Cambessedesia, DC. (Melast.).
lanatum, Turcz. l. c. 578 = Pyramia pityrophylla.
rugosum, Turcz. l. c. = C. rugosa.

ACIPHYLLA, Forst. Char. Gen. 125. t. 68 (1776).—
UMBELLIFERÆ, Benth. & Hook. f. i. 912.
ANISOTOME, Hook. f. Fl. Antarct. i. 16. t. 8, 9 (1845).
CALOCIADIUM, Endl. Gen. Suppl. iv. iii. 7 (1850).
GINGIDIUM, F. Muell. in Trans. Phil. Inst. Vict. i. (1855) 108.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 92.—N. Zel.
crenulata, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 336.—N. Zel.
Dobsoni, *Hook. f. Handb. N. Zeal. Fl.* 93.—N. Zel.
glacialis, *F. Muell. ex Benth. Fl. Austral.* iii. 375.—Austral.
Hectori, *Y. Buch, in Trans. N. Z. Inst.* xiv. (1882) 346.—N. Zel.
Lyallii, *Hook. f. Handb. N. Zeal. Fl.* 92.—N. Zel.
Monroi, *Hook. f. Fl. N. Zel.* ii. 330.—N. Zel.
montana, *Armst. in Trans. N. Z. Inst. iv.* (1872) 290.—N. Zel.
procumbens, *F. Muell. ex Benth. Fl. Austral.* iii. 375.—Austral.
simplicifolia, *F. Muell. ex Benth. l. c.*—Austral.
squarrosa, *Forst. Char. Gen.* 126.—N. Zel.
Traillii, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 371.—N. Zel.
Traversi, *F. Muell. ex Y. Buch, l. c.* vii. (1875) 385.—N. Zel.

ACIPHYLLAEA, A. Gray, in Mem. Am. Ac. N. S. iv. (1849) (Pl. Fendl.) 91 = Hymenatherum, Cass. (Compos.).
acerosa, A. Gray, l. c. = H. acerosum.

ACIPHYLLUM, Steud. Nom. ed. II. i. 17 (1840) = Chorizema, Labill. (Legum.).

ACIS, Salisb. Parad. Lond. sub t. 74 (1807) = Leucojum, Linn. (Amaryll.).
autumnalis, Salisb. l. c. = L. autumnale.
Broteri, Jord. & Fourr. Ic. Pl. Europ. 25 = L. trichophyllum.
grandiflora, Sweet, Brit. Fl. Gard. iii. sub t. 297 = L. trichophyllum.
hiemalis, M. Roem. Syn. Enant. 24 = L. hyemale.
hyemalis, Sweet, Brit. Fl. Gard. iii. sub t. 297 = prec.
longifolia, M. Roem. Syn. Enant. 25 = L. longifolium.
rosea, Sweet, Brit. Fl. Gard. t. 297 = L. hyemale.
spoerntha, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 51 = L. autumnale.
pulchella, Jord. & Fourr. l. c. = L. autumnale.
roseus, Sweet, ex Loud. Hort. Brit. Supp. iii. 482 = L. roseum.
trichophyllus, Sweet, Brit. Fl. Gard. iii. sub t. 297 = L. trichophyllum.

ACISANTHERA, P. Br. Hist. Jamaic. 217. t. 22. f. 1. (1756). *MELASTOMACEÆ*, Benth. & Hook. f. i. 739.
ANISOCENTRUM, Turcz. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 232.
DICRANANTHERA, Pavel, Symb. Bot. i. 75. t. 50 (1832).
EXODICILIS, Rafin. Sylva Tellur. 97 (1838).
MICCARPUS, Nand. in Ann. Sc. Nat. Sér. III. ii. (1844) 146.
NOTEROPHILA, Mart. Nov. Gen. et Sp. iii. 190. t. 254 (1829).
URANTHERA, Nand. in Ann. Sc. Nat Sér. III. ii. (1844) 146.

ACISANTHERA:—
adscendens, Wright, in Sauvalle, Cat. Fl. Cub. 53 = quadrata.
alata, *Cogn. in Mart. Fl. Bras.* xiv. III. 225.—Bras.
alsinaefolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 34. t. 2. f. 18 b.—Bras.
Beccabunga, Triana, l. c. 38 = brevifolia.
brevifolia, Cogn. in Mart. Fl. Bras. xiv. III. 216 = brevifolia. Botsiciana, *Cogn. l. c.* 209.—Bras.
brevifolia, *Griseb Fl. Brit. W. Ind.* 269.—Ind. occ.; Bras.
brevifolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 33 = trivalvis.
divaricata, *Cogn. in Mart. Fl. Bras.* xiv. III. 212.—Bras.
fluitans, *Cogn. l. c.* 214.—Bras.
Glaziovana, *Cogn. l. c.* 223.—Bras.
hedyotoides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 33.—Bras.
hedyotoides, Triana, l. c. pro parte = Salzmanni.
inundata, *Triana, l. c.*—Bras.
lanuginosa, *Triana, l. c.* t. 11.—Am. trop.
pellucida, *Wright, in Griseb. Cat. Pl. Cub.* 104.—Cuba.
pulchella, *Cogn. in Mart. Fl. Bras.* xiv. III. 211.—Bras.
punctatissima, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 34. t. 11. f. 18 c.—Bras.; N. Granat.
quadrata, *Yuss. ex Poir. Encyc. Suppl.* i. 111.—Jamaica.
recurva, *Griseb. in Goett. Abh.* ix. (1861) 58.—Bras.; Guiana.
recurva, Triana, *in Trans. Linn. Soc.* xxviii. (1871) 34. t. 2. f. 18 b.—Bras.
Salzmanni, *Cogn. in Mart. Fl. Bras.* xiv. III. 209.—Bras.
tetramera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 33.—Bras.
trivalvis, *Cogn. in Mart. Fl. Bras.* xiv. III. 217.—Bras.
varabilis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 34. t. 11. f. 18 g.—Bras.

ACISPERMUM, Neck. Elem. i. 34 (1790) = **Coreopsis**, Linn. (Compos.).

ACISTOMA, Zipp. ex Span. in Linnæa, xv. (1841) 202 = Woodfordia, Sm. (Lythrar.).
coccinea, Zipp. ex Span. l. c. = W. floribunda.

ACKAMA, A. Cunn. in Ann. Nat. Hist. ii. (1839) 858.
SAXIFRAGEÆ, Benth. & Hook. f. i. 653.
Muelleri, *Benth. Fl. Austral.* ii. 444.—Austral.
rosaefolia, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 858.—N. Zel.

ACLADODEA, Ruiz & Pav. Prod. 133. t. 29 (1794) = **Talisia**, Aubl. (Sapind.).
pinnata, Ruiz & Pav. Syst. Veg. 262 = T. Acladodes.

ACLEIA, DC. Prod. vi. 340 (1837) = **Senecio**, Linn. (Compos.).
belbeysia, DC. l. c. = S. belbeysius.

ACLEISANTHES, A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 259. *NYCTAGINEÆ*, Benth. & Hook. f. iii. 6.
PENTACROPHYTS, *A. Gray, l. c.* 261 (1858).
anisophylla, *A. Gray, l. c.*—Texas.
Berlandieri, *A. Gray, l. c.* 260.—Texas.
crassifolia, *A. Gray, l. c.*—Texas.
longiflora, *A. Gray, l. c.*—N. Mexico.
Wrightii, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 6.—Texas.

ACLINIA, Grisb. Fl. Brit. iii. 320 (1851) = **Dendrobium**, Sw. (Orchid.).

ACLISIA, E. Mey. in Presl, Rel. Haenk. i. 137. t. 25 (1827) = **Follilia**, Thunb. (Commel.).
cotobica, Hassk. Commel. Ind. 49 = P. macrophylla.
Cumingiana, Hassk. l. c. 47 = P. macrophylla.
elegans, Hassk. in Mig. Fl. Jungh. 49 = P. sorsogonensis.
floridu, Hort. Berol. (1837) ex Kunth, Enum. Pl. iv. 47 = Athyrocarpus pallidus.
Griffithii, Hassk. Commel. Ind. 46 = P. sorsogonensis.
indica, Wight, Ic. t. 2066 = P. sorsogonensis.
sorsogonensis, Kunth, Enum. Pl. iv. 74 = P. indica.
sorsogonensis, E. Mey. in Presl, Rel. Haenk. i. 138. t. 26 = P. sorsogonensis.

ACLISIA:—
guayaquilensis, *G. Don, Gen. Syst.* iv. 461.—Peruv.
subumbellata, C. B. Clarke, Commel. et Cyrt. Beng. t. 30 = P. subumbellata.
Thomsoni, C. B. Clarke, l. c. t. 31 = Aneilema Thomsoni.
Zollingeri, Hassk. Commel. Ind. 51 = P. Zollingeri.

ACMADENIA, Bartl. & Wendl. f. Diosm. 59. t. A (1824). *RUTACEÆ*, Benth. & Hook. f. i. 369.
intermedia, *Chem. in Linnæa*, v. (1830) 52.—Afr. austr.
lasiantha, *Sond. in Harv. & Sond. Fl. Cap.* i. 383.—Afr. austr.
casuarinoides, Turcz. in Bull Soc. Nat. Mosc. xxxi. (1858) 1. 339 = Macrostylis dubia.
cucullata, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 383.—Afr. austr.
densifolia, *Sond. l. c.*—Afr. austr.
flaccida, *Eckl. & Zeyh. Enum.* 105.—Afr. austr.
juniperina, *Bartl. & Wendl. f. Diosm.* 61.—Afr. austr.
laevigata, *Bartl. & Wendl. f. l. c.* 64.—Afr. austr.
laevigata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 383 = asiatifolia.
Mundiana, *Eckl. & Zeyh. Enum.* 105.—Afr. austr.
muricinodes, Eckl. & Zeyh. l. c. 106 = juniperina.
Niveni, *Sond. in Harv. & Sond. Fl. Cap.* i. 380.—Afr. austr.
obtusata, *Bartl. & Wendl. f. Diosm.* 68.—Afr. austr.
obtusata, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i. 376 = Diosma teretifolia.
pallopetala, *Sond. in Harv. & Sond. l. c.* i. 380.—Afr. austr.
pungens, *Bartl. & Wendl. f. Diosm.* 64.—Afr. austr.
pungens, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i. 381 = obtusata.
rosmarinifolia, *Bartl. in Linnæa*, xvii. (1844) 355.—Afr. austr.
strobilina, E. Mey. ex Harv. & Sond. Fl. Cap. i. 381 = tetragona.
tetragona, *Bartl. & Wendl. f. Diosm.* 65.—Afr. austr.
tetragona, Eckl. & Zeyh. Enum. 105 = laevigata.

ACMANTHERA, Griseb. in Mart. Fl. Bras. xii. 1. 28 (1858). *MALPIGHIACEÆ*, Benth. & Hook. f. i. 258.
latifolia, *Griseb. l. c.* 29.—Bras.

ACMELLA, Rich. in Pers. Syn. ii. 472 (1807) = **Spilanthes**, Jacq. (Compos.).
bifora, Spreng. Syst. iii. 591 = Wedelia bifora.
brachyglossa, Cass. in Dict. Sc. Nat. i. 268 = S. Acmella.
brasiliensis, Spreng. Syst. iii. 592 = Wedelia paludosa.
bupblthalmoides, Pers. Syn. iii. 478 = Heliopsis buphthalmoides.
caulorrhiza, Dellile, Cent. Fl. Afr. Méroé, 45. t. iii. f. 7 = S. Acmella.
ciliata, Cass. in Dict. Sc. Nat. xxiv. 331 = S. ciliata.
debilis, Cass. l. c. 331 = S. debilis.
fimbriata, Cass. l. c. = S. fimbriata.
flaviculis, Rafin. New Fl. Am. i. 52.—Am. bor.
Garcini, Spreng. Syst. iii. 591 = Anvillea Garcini.
globosa, Spreng. l. c. 592 = Zaluzania globosa.
hirta, Lag. Gen. et Sp. Nov. 30 = Jaegeria hirta.
intermedia, Rich. ex Pers. Syn. ii. 473 = S. intermedia.
lanceolata, Link, ex Spreng. Syst. iii. 444 = S. Pseudoacmella.
Linnæi, Cass. in Dict. Sc. Nat. xxiv. 330 = S. Acmella.
mauritiana, Rich. ex Pers. Syn. ii. 473 = S. Acmella.
Montana, Cass. in Dict. Sc. Nat. xxiv. 331 = S. Mutisii.
nudicaulis, Rafin. New Fl. Am. i. 52.—Am. bor.
Nuttaliana, Rafin. l. c. = S. repens.
occidentalis, Nutt. Gen. Am. ii. 171 = S. repens.
accidentalis, Rich. ex Pers. Syn. ii. 478 = S. Mutisii.
parvifolia, Rafin. New Fl. Am. i. 52.
repens, Rich. ex. Pers. Syn. ii. 478 = S. repens.
spilanthoides, Cass. Dict. Sc. Nat. xxiv. 329 = Wedelia carnosa.
tenella. Cass. in l. c. xxiv. 331 = S. tenella.
trifida, Link, Spreng. Syst. iii. 591 = Zaluzania trifoba.
uliginosa, Cass. in Dict. Sc. Nat. xxiv. 331 = S. Acmella.

ACMENA, DC. in Dict. Class. Hist. Nat. xi. 401 (1826) = **Eugenia**, Linn. (Myrtac.).
bracteolata, Walp. Rep. ii. 161 = E. bracteolata.
Chemponi, Benth. in Hook. Kew Journ. Bot. i. (1853) 118.—Hongkong.
chinensis, Planch. Hort. Donat. 84.—China.
claviflora, Walp. Rep. ii. 181 = E. claviflora.
elliptica, G. Don, ex Steud. Nom. ed. II. i. 17 = E. Smithii.

ACMENA:—
floribunda, DC. in Dict. Class. Hist. Nat. xi 401; et Prod. iii. 262 = E. Smithii.
Gerrardi, Harv. Gen. S. Afr. Pl. ed. II. 112.—Afr. austr.
grata, Walp. Rep. ii. 181 = E. grata.
inophylla, Wight, Ic. ii. t. 623 = E. inophylla.
Kingii, G. Don, Gen. Syst. ii. 851 = E. Smithii.
lanceolata, Thw. Enum. Pl. Zeyl. 119 = E. Wightiana.
leptantha, Walp. Rep. ii. 181 = E. leptantha.
oblata, Walp. l. c. = E. oblata.
parviflora, DC. Prod. iii. 262 = E. zeylanica.
Wightiana, Walp. Rep. ii. 181 = E. Wightiana.
zeylanica, Thw. Enum. Pl. Zeyl. 118 = E. zeylanica.

ACMISPON, Rafin. Atl. Journ. 144 (1832) = **Hosackia**, Dougl (Legumin.).
multiflorum, Rafin. New Fl. Am. i. 58 = sec. ? *sericeum*, Rafin. Atl. Journ. 145 = H. sericea.

ACMOSTIMA, Rafin. Sylva Tellur. 26 (1838) = **Pavetta**, Linn. (Rubiac.).
brevifolia, Rafin. l. c. = P. pentandra.
longifolium, Rafin. l. c. = P. barbata.

ACNADENA, Rafin. Sylva Tellur. 41 (1838) = **Cordia**, Linn. (Borrag.).
elliptica, Rafin. l. c. 41 = C. elliptica.

ACNIDA, Linn. Diss. Chen. 32 (1750) (Am. acad. iii. 19). *AMARANTACEÆ*, Benth. & Hook. f. iii. 29.
MONTELLIA, A. Gray, Man. Bot. N. U. States, ed. II. 389 (1856).
altissima, Michx. ex. Moq. in DC. Prod. xiii. II. 277 = rusocarpa.
altissima, Kidd. ex Moq. l. c. = tuberculata?
australis, *A. Gray, in Am. Natural.* x. (1876) 489.—Am. bor.
cannabina, Chapm. Fl. S. U. St. Suppl. 645 = australis.
cannabina, *Linn. Sp. Pl.* 1027.—Am. bor.
cuspidata, Bert. ex Spreng. Syst. iii. 908 = cannabina.
Elliott, *Rafin. New Fl. Am.* i. 54.—Am. bor.
floridana, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 376.—Am bor.
obtusifolia, *Rafin. New Fl. Am.* i. 54.—Am. bor.
oderata, *P. C. Dietr. ex Moq. in DC. Prod.* xiii. II. 278, nomen.—Hab.?
rhyssocarpa, Stend. Nom. ed. II. i. 17 = rusocarpa.
ruscocarpa, Willd. Sp. Pl. iv. 768 = rusocarpa.
ruscocarpa, Moq. in DC. Prod. xiii. II. 277 = tuberculata.
rusocarpa, *Michx. Fl. Bor. Am.* ii. 234. t. 50.—Am. bor.
salicifolia, *Rafin. in Am. Monthly Mag.* ii. (1817) 48.—Am. bor.
tamariscina, *A. Gray, in Am. Natural.* x. (1876) 489.—Am. bor.
tuberculata, *Moq. in DC. Prod.* xiii. II. 277.—Am. bor.

ACNISTUS, Schott, in Wien. Zeitschr. (1829) iv, 1180, ex Linnæa, vi. (1831) Litt. 54. *SOLANACEÆ*. Benth. & Hook. f. ii. 894.
CODOCHONIA, Dun. in DC. Prod. xiii. 1. 482 (1852).
LYCIOPLESIUM, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 320.
EPHAILA, HUANUCA, KOKABUS, PEDERLEA, PLICULA, TROZELIA, Rafin. Sylva Tellur. 54–55, 124 (1838).
aggregatus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 341.—Peruv.
arborescens, *Schiecht. in Linnæa*, vii. (1832) 67.—Mexico.
Benthami, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 342.—Mexico.
brevifloras, *Sendtn. in Mart. Fl. Bras.* x. 152.—Bras.
canaliferus, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 337 = Miersi.
canaliferus, *Schott, in Wien. Zeitschr.* (1829) iv. 1180.—Bras.
costrioides, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 343 = Lycium costroides.
confertiflorus, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 31.—Peruv.
ellipticus, *Hook. f. in Trans. Linn. Soc.* xx. (1850) 308.—Ins. Galap.
fasciculatus, *Benth. & Hook. f. Gen.* ii. 894.—Boliv.
floribundus, *G. Don, Gen. Syst.* iv. 461.—Peruv.
frutescens, *Bolle, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 299.—Porto Rico.
grandiflorus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 344.—Venezuela.

ACNISTUS:—
guayaquilensis, *G. Don, Gen. Syst.* iv. 461.—Peruv.
Mierii, *Dun. in DC. Prod.* xiii. i. 497.—Braz.
parviflorus, *Griseb. in Goett. Abh.* xix. (1874) 219.—Reg. Argent.
Plumieri, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 340.—Am. austr.
ramiflorus, *Miers, l. c.* 342.—Ind. occ.
sideroxyloides, *G. Don, Gen. Syst.* iv. 461.—N. Granat.
spatulatus, *G. Don, l. c.*—Peruv.
umbellatus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 342.—Peruv.
virgatus, *Griseb. in Gott. Abh.* xxiv. (1879) 247.—Reg. Argent.

ACONLORRHAPHE, H. Wendl. in Bot. Zeit. xxxvii. (1879) 145. *PALMAE.*

ACOIDIUM, Lindl. Bot. Reg. (1837) sub t. 1951 = Trichoceratrum, Poepp. & Endl. (Orchid.).
fuscum, Lindl. l. c. = T. fuscum.

ACOKANTHERA, G. Don, Gen. Syst. iv. 485 (1837). *APOCYNACEAE*, Benth. & Hook. f. ii. 696. Toxicophlaea, Harv. in Hook. Lond. Journ. Bot. i. (1842) 24.
Lamarkii, *G. Don, Gen. Syst.* iv. 485.—Afr. austr.
lycioides, *G. Don, l. c.*—Afr. austr.
pubescens, *G. Don, l. c.* 486.—Afr. austr.
spectabilis, *Hook. f. Bot. Mag.* t. 5359.—Afr. austr
venenata, *G. Don, Gen. Syst.* iv. 485.—Afr. austr.

ACOMA, Benth. Bot. Voy. Sulph. 29. t. 17 (1844) = Corecoxarpus, Benth. (Compos.).
dissectum, Benth. l. c. (= *Leptoxyne dissecta*).

ACOMIS, F. Muell. Fragm. ii. 89 (1860). *COMPOSITAE*, Benth. & Hook. f. ii. 317.
maura, *F. Muell. l. c.* iv. 14.—Austral.
rutidosea, *F. Muell. l. c.* ii. 89.—Austral.

ACONCEVIBIUM, Miq. Fl. Ind. Bat. i. xi. 869 (1859). *EUPHORBIACEAE.*
trinerve, *Miq. l. c.*—Moluccas.

ACONITUM, Tourn. ex Linn. Syst. ed. I. (1785). *RANUNCULACEAE*, Benth. & Hook. f. i. 9.
Anthora, Hall. in Rupp. Fl. Jen. ed. Hall. 290 (1745).
Cammarum, Fourr. in. Ann. Soc. Linn. Lyon. N. S. xvi. (1868) 397.
Lycoctonum, Fourr. l. c. 326.
Napellus, Rupp. Fl. Jen. ed. Hall. 275 (1745).
Nirbisia, G. Don, Gen. Syst. i. 68 (1831).
abbreviatum, Langsd. ex DC. Prod. i. 61 = Fischeri.
achranthum, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 285.—Sib. altaic.
acuminatum, Reichb. Uebers. Acon. 46 = paniculatum.
acutum, Reichb. l. c. 21 = A. Napellus.
aegobromum, Reichb. l. c. 89 = A. Lycoctonum.
alatum, Wender. in Linnaea, xv. (1841) List. 99.—A. Napellus.
altaicum, *Host, Fl. Austr.* ii. 76.—Europ.
albidum, Bernh. ex Reichb. Uebers. Acon. 31 = A. Napellus.
album, Ait. Hort. Kew. ii. 246 = variegatum.
allemum, Reichb. Uebers. Acon. 78. = A. Lycoctonum.
alpinum, Mill. Gard. Dict. ed. VIII. n. 7 = variegatum.
altigaleatum, Hayne, Arm. Gew. xii. t. 16 = variegatum.
altissimum, Mill. Gard. Dict. ed. VIII. n. 2 = A. Lycoctonum.
ambiguum, Reichb. Monog. Acon. t. 28 = A. Napellus.
amoenum, Reichb. Uebers. Acon. 33 = A. Napellus.
amplifrons, Reichb. l. c. 37 = A. Napellus.
angustifolium, Bernh. ex Reichb. l. c. 29 = A. Napellus.
Anthora, *Linn. Sp. Pl.* 533.—Europ.; Orient.; As. bor.
anthorum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Anthora.
anthoroidoum, DC. Syst. i. 366 = A. Anthora.
arctophorum, Reichb. Uebers. Acon. 71 = A. Lycoctonum.
arcuatum, *Maxim. Prim. Fl. Amur.* 27.—Reg. Amur.
Atios, Rorle. in Journ. As. Soc. Beng. i. (1832) 459 = heterophyllum.
atlanticum, *Coss. in Bull. Soc. Fr.* xxii. (1875) 58, nomen.—Afr. bor.
atrox, Walp. Rep. i. 57 = ferox.
australe, Reichb. Uebers. Acon. 71 = A. Lycoctonum.

ACONITUM:—
austriacum, Tratt. ex Reichb. Monog. Acon. '18' (t. 71) = A. Cammarum.
autumnale, Lindl. in Journ. Hortic. Soc. ii. (1847) 77 = Fischeri.
autumnale, Reichb. Monog. Acon. t. 17. f. 2 = A. Napellus.
baicalense, *Turcz. ex Steud. Nom.* ed. II. i. 18.—Sibir. baical.
barbatum, Patr. ex Pers. Syn. ii. 93 = A. Lycoctonum.
Baumgartenii, *Schur, in Verh. Siebenb. Ver. Naturw.* iii. (1852) 84.—Europ.
Bernhardianum, Reichb. Uebers. Acon. 84 = A. Napellus.
Bernhardianum, Wallr. Sched. Crit. i. 250. t. 2 = variegatum.
Besserianum, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1882) 58, nomen.—Podolia.
bicolor, Schult. Obs. Bot. 101 = A. Cammarum.
biflorum, *Fisch. ex DC. Syst.* i. 380.—Sibir. altaic.
borealo, Ser. ex Reichb. Uebers. Acon. 62 = A. Lycoctonum.
bracteosum, *Bercht. & Presl, Rostl.* i. *Ranunculac.* 197, nomen.—Europ.
Braunii, Reichb. Uebers. Acon. 29 = A. Napellus.
Breiterianum, Reichb. l. c. 54 = A. Cammarum.
bulbiferum, Reichb. l. c. 55 = variegatum.
caeruleum, Dulac, Fl. Hautes-Pyr. 223 = A. Napellus.
calibotryon, Reichb. Monog. Acon. 98. t. 16. f. 2 = A. Napellus.
Camarum, Schleich. Cat. (1821) 5 = paniculatum.
Cammarum, *Linn. Sp. Pl.* ed. II. 751.—Europ.
Cammarum, Jacq. Fl. Austr. v. t. 234 = variegatum.
camtschaticum, auct. = *kamtschaticum.*
Candollei, Reichb. ex Steud. Nom. ed. II. i. 18 = A. Anthora.
canescens, Schleich. ex Reichb. Fl. Germ. Excurs. 741 = A. Napellus.
Carmichaeli, Debeaux, in Act. Linn. Soc. Bord. xxxiii. (1879) 87 (Florule du Tientein, 61) = Fischeri.
cernuum, Baumg. ex Schur, Enum. Pl. Transs. 82 = A. Cammarum.
cernuum, Wulf. ex Koelle, Spicil. 17 = paniculatum.
Chamissonianum, Reichb. Uebers. Acon. 37 = A. Napellus.
chinense, Siebold, ex Fauc. Mag. v. (1838) 3, cum tab. = Fischeri.
chinense, *Sieb. ex Sieb. & Zucc. Fl. Jap. Fam. Nat.* 76 (76).—Japon.
ciliare, DC. Syst. i. 378 = villosum, volubile.
Clairvillianum ×, *Bruegg. in Vahrsch. Naturf. Ges. Graub.* N. F. xxiii.-xxiv. (1881) 93.—Europ.
Clusianum, Reichb. Monog. Acon. 13 = sec.
Clusii, Reichb. Uebers. Acon. 22 = A. Napellus.
columbianum, Nutt. in Torr. & Gray, Fl. N. Am. i. 84 = Fischeri.
commutatum, Reichb. Uebers. Acon. 86 = A. Napellus.
compactum, Reichb. l. c. 27 = A. Napellus.
conspersum, Reichb. l. c. 86 = A. Napellus.
cordatum, Rafin. New Fl. Am. i. 56.—Am. bor.
cordatum, Royle, Illustr. 56 = heterophyllum.
cynoctonum, Reichb. Uebers. Acon. 78 = A. Lycoctonum.
Decandollii, Reichb. l. c. 16 = A. Anthora.
decorum, Reichb. l. c. 51 = A. Cammarum.
delphinifolium, DC. Syst. i. 380 = A. Napellus.
delphinifolium, Hort. Prag. ex Steud. Nom. ed. II. i. 18 = A. Lycoctonum.
denudiflorum, Hoppe, ex Steud. Nom. ed. II. i. 18 = A. Napellus.
diosectum, D. Don, Prod. Fl. Nep. 197 = A. Napellus.
dissectum, Tausch, ex Reichb. Monog. Acon. t. 47 = A. Lycoctonum.
divergens, *Pand. Elem.* 18.—Europ.
divergens, *Rafin. New Fl. Am.* i. 56.—Am. bor.
dictum, Salisb. Prod. 376 = A. Napellus.
eminens, Koch, ex Reichb. Uebers. Acon. 35 = A. Napellus.
engadinense ×, *Bruegg. in Vahrsch. Naturf. Ges. Graub.* II. xxiii.-xxiv. (1880) 92.—Europ.
eriostemum, DC. Syst. i. 377 = A. Cammarum.
eulophum, Reichb. Uebers. Acon. 15 = A. Anthora.
exaltatum, Reichb. l. c. 94 = A. Napellus.
exaltatum, Bernh. ex Reichb. l. c. 51 = A. Cammarum.
excelsum, *Reichb. Monog. Acon.* t. 58.—Russia; Sibir.
excelsum, Turcz. Cat. Baikal. n. 70 = A. Lycoctonum.
ferox, *Wall. in Ser. Mus. Helv.* i. 160.—Reg. Himal.

ACONITUM:—
ferox, Wall. Cat. n. 4721 A (Fl. As. Rar. t. 41) = A. Napellus.
firmum, Reichb. Uebers. Acon. 20 = A. Napellus.
Fischeri, *Reichb. Monog. Acon.* t. 22.—As. bor.; Am bor. flaccidum, *Reichb. Uebers. Acon.* 39.—Sibir.
fontanum, Hoppe & Hornsch. ex Reichb. Monog. t. 38 in syn. = paniculatum.
foxwoodum, Presl, ex Reichb. l. c. t. 7 in syn. = variegatum.
foxwoodum, *Rafin. New Fl. Am.* i. 55.—Am. bor.
formosum, Reichb. Uebers. Acon. 26 = A. Napellus.
fornicatum, Gillib. Fl. Lituan. ii. 285 = A. Napellus.
Funckianum, Reichb. Monog. Acon. t. 66 = A. Napellus.
Funckii, Reichb. Uebers. Acon. 28 = A. Napellus.
galeatum, Steven, ex Steud. Nom. ed. II. i. 18 = orientale.
galeriflorum, Stokes, Bot. Mat. Med. iii. 216 = A. Lycoctonum.
galectonum, Reichb. Uebers. Acon. 67 = A. Lycoctonum.
gymnaliflorum, *Host, Fl. Austr.* ii. 76.—Europ.
gibbiferum, Reichb. Monog. Acon. t. 19 = paniculatum.
gibbosum, Ser. Mus. Helv. i. 141. t. 15. f. 14, 15 = variegatum.
gigantoum, Host, ex Steud. Nom. ed. II. i. 18 = A. Napellus.
gigantoum, Zumag. Fl. Pedem. ii. 845 = A. Lycoctonum?
glabrum, DC. Syst. i. 379 = variegatum.
Gmelini, Reichb. Uebers. Acon. 63 = A. Lycoctonum.
gracile, Reichb. l. c. 55 = variegatum.
grandiflorum, [Fisch.] Hort. Gorenk. ed. I. (1808) 77 = biflorum.
grandiflorum, Pall.; cf. Reichb. Uebers. Acon. 86 = A. Napellus.
guttatum, Turcz. ex Regel, in Bull. Soc. Nat. Mosc. (1861) ii. 114 = A. Napellus.
gymnandrum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 308.—China.
Halleri, Reichb. Uebers. Acon. 27 = A. Napellus.
hamatum, Hort. Berol. ex Steud. Nom. ed. II. i. 18 = variegatum.
hamatum, Reichb. in Cat. Hort. Berol. (1818); Uebers. Acon. 48 = A. Cammarum.
Hamiltonii, G. Don, Gen. Syst. i. 69 = A. Napellus.
hebegynum, DC. Syst. i. 376 = paniculatum.
heterophyllum, *Wall. Cat.* n. 4722.—Reg. Himal.
hiana, *Host, Fl. Austr.* ii. 74.—Europ.
hians, Reichb. Fl. Germ. Excurs. 740 = A. Napellus.
hispidum, DC. Syst. i. 367 = A. Lycoctonum.
Hoppeanum, Reichb. Monog. Acon. t. 65 = A. Napellus.
Hoppii, Reichb. Uebers. Acon. 24 = A. Napellus.
hortense, Hoppe, ex Reichb. Monog. Acon. t. 71 in syn. = A. Cammarum.
Hostianum, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 177 = A. Lycoctonum.
humile, Bernh. ex Reichb. Uebers. Acon. 34 = A. Napellus.
humile, Salisb. Prod. 375 = paniculatum.
illimitum, Reichb. Uebers. Acon. 64 = variegatum.
inclinatum, Sweet, Hort. Brit. ed. II. 9 = A. Anthora?
intermedium, DC. Syst. i. 374 = A. Cammarum.
intermedium, Gaud. ex Steud. Nom. ed. II. i. 18 = variegatum.
intermedium, Host, Fl. Austr. ii. 69 = A. Lycoctonum.
inuncturn, Koch, ex Reichb. Monog. Acon. '6. app.' (Index) = A. Napellus.
italicum, Tratt. ex Reichb. l. c. t. 7. in syn. = variegatum.
Jacquini, Reichb. Uebers. Acon. 17 = A. Anthora.
Jacquinianum, Host, Fl. Austr. ii. 68 = A. Lycoctonum.
japonicum, *Decne. in Ann. Hortic.* (1851) 178.—Japon.
japonicum, Hort. ex Steud. Nom. ed. II. i. 18 = variegatum.
japonicum, Thunb. Fl. Jap. 221 = uncinatum.
kamtschaticum, Pall. ex Reichb. Uebers. 39 = Fischeri.
Koehleri, Reichb. Uebers. Acon. 30 = A. Napellus.
Koellowianum, Reichb. l. c. 18 = A. Napellus.
Kunisechii, *Reichb. Monog. Acon.* t. 11.—As. bor. or.
lasciniosum, Schleich. ex Steud. Nom. ed. II. i. 18 = variegatum.
lasianum, Reichb. Uebers. Acon. 21 = A. Napellus.
lasve, Royle, Illustr. 56 = A. Lycoctonum.
lasvigatum, Schleich. ex Steud. Nom. ed. II. i. 18 = variegatum.
lagochilum, Reichb. Uebers. Acon. 71 = A. Lycoctonum.
Lamarckii, Reichb. Monog. Acon. t. 40 = A. Lycoctonum.
lasiostylum, Reichb. Uebers. Acon. 85 = variegatum.

ACONITUM:—
laxistomum, *Reichb. Monog. Acon.* t. 49.—Rosa.; Podol.
laxistomum, Spreng. Syst. ii. 620 = orientale.
laxiflorum, Schleich. Cat. (1821) 6 = A. Napellus.
laxum, Reichb. Illustr. Acon. t. 13 verso = A. Napellus.
leptanthum, *Reichb. Monog. Acon.* t. 64—Rossia.
lethale, Griff. Notul. iv. 782 = palmatum.
leucanthemum, Wender. in Linnæa, v. (1830) Litt. 53 = variegatum.
leucanthum, *Reichb. Uebers. Acon.* 55.—Hab.?
Lobelianum, *Host, Fl. Austr.* ii. 74.—Europ.
Lubanskyi, Reichb. Monog. Acon. t. 20 = Fischeri.
Lubarscii, Heynh. Nom. i. 18 = præc.
luparia, Reichb. Uebers. Acon. 74 = A. Lycoctonum.
lupicida, Reichb. l. c. 70 = A. Lycoctonum.
luridum, *Hook. f. & Thoms. Fl. Ind.* i. 55.—Reg. Himal.
luridum, Salisb. Prod. 375 = variegatum.
Lycoctonum, Bess. Enum. Pl. Volh. 22 = orientale.
Lycoctonum, *Linn. Sp. Pl.* 532.—Europ.; As. bor.
macranthum, Reichb. Uebers. Acon. 51 = variegatum.
macrophyllum, Hort. ex G. Don, in Loud. Hort. Brit. 338 = orientale.
macrorhynchum, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 53.—As. temp.
maximum, Pall. ex DC. Syst. i. 380; Prod. i. 61 = Fischeri.
medium, Schrad. Cat. Hort. Goett. ex Reichb. Monog. Acon. t. 71 in syn. = A. Cammarum.
melocranum, Reichb. Uebers. Acon. 65 = A. Lycoctonum.
Meyeri, Reichb. l. c. 33 = A. Napellus.
microphyllum, Gaud. ex Steud. Nom. ed. II. i. 19 = A. Napellus.
microstachyum, Reichb. Uebers. Acon. 86 = A. Napellus.
Mielichhoferi, Reichb. l. c. 29 = A. Napellus.
mixtum, Reichb. l. c. 61 = variegatum.
moldavicum, Hacq. ex Reichb. l. c. 67 = A. Lycoctonum.
molle, Reichb. l. c. 47 = paniculatum.
monanense, Schmidt, ex Reichb. Monog. Acon. t. 51 in obs. = A. Lycoctonum.
monogynum, Forsk. Fl. Aegypt. Arab. p. xxvii = Delphinium Aconiti.
multifidum, Koch, ex Reichb. Monog. Acon. t. 70 = A. Napellus.
multifidum, Royle, Illustr. 56 = A. Napellus.
myctonum, Reichb. Uebers. Acon. 58 = A. Lycoctonum.
nanum, Stephan, ex Ledeb. Fl. Ross. i. 70 = biflorum.
napelloides, Sw. ex Reichb. Monog. Acon. t. 69 in syn. = A. Napellus.
Napellus, *Linn. Sp. Pl.* 532.—Reg. bor. temp
Napellus, Pall. Reise, ii. 528 = villosum.
Napellus, S. G. Gmel. It. i. 8 = A. Lycoctonum.
Napellus, Thunb. Fl. Jap. 381 = Fischeri.
nasutum, Fisch. ex G. Don, Gen. Syst. i. 61 (1831) 58 = variegatum.
nasutum, Hook. Fl. Bor. Am. i. 26 = Fischeri.
neapolitanum, Tenore, Fl. Nap. iv. 327 = A. Lycoctonum.
nemorosum, Bleb. ex Reichb. Uebers. Acon. 17 = A. Anthora.
neomontanum, Baumg. Fl. Transs. ii. 100 = paniculatum.
neomontanum, Koelle, Spicil. 16 = A. Napellus.
neomontanum, Willd. Sp. Pl. ii. 1236 = A. Cammarum.
neubergense, DC. Syst. i. 379 = A. Napellus.
nitidum, Fisch. ex Steud. Nom. ed. II. i. 19 = barbatum.
nutans, Hort. Helmst. ex Steud. Nom. ed. II. i. 19 = A. Napellus.
Nuttallii, *Sweet, Hort. Brit. ed.* II. 584, nomen.—Am. bor.
obscurum, Reichb. ex S. Wats. Bibl. Ind. 2 = A. Napellus.
obtusiflorum, *Host, Fl. Austr.* ii. 72.—Europ.
ochotense, Reichb. Monog. Acon. t. 18.—Sibir. or.
ochranthum, C. A. Mey. in Ledeb. Fl. Alt. ii. 286 = A. Lycoctonum.
ochroleucum, Hort. ex Steud. Nom. ed. II. i. 19 = A. Lycoctonum.
ochroleucum, Saliseb. Prod. 375 = A. Anthora.
ochroleucum, Willd. Sp. Pl. ii. 1238 = orientale.
oligoanthemum, Kern. in Ber. Naturw. Ver. Innsbruck, i. (1870) 113 = A. Napellus.
oligocarpum, Reichb. Uebers. Acon. 24 = A. Napellus.
oregonense, *Rydb. New Fl. Am.* i. 57.—Oregon.
orientale, *Mill. Gard. Dict. ed.* VIII. n. 16.—Reg. Caucas.

ACONITUM:—
Ottonianum, Reichb. Uebers. Acon. 54 = A. Cammarum.
ovatum, Lindl. in Bot. Reg. (1840) Misc. 58 = heterophyllum!
Pallasii, Reichb. Uebers. Acon. 18 = A. Anthora.
pallidum, Nutt. ex Loud. Hort. Brit. Suppl. 483, nomen = Nuttallii.
pallidum, Reichb. Uebers. Acon. 65 = A. Lycoctonum.
palmatum, Reichb. l. c. 46 = A. Cammarum.
palmatum, *D. Don, Prod. Fl. Nep.* 196.—Reg. Himal.
paniculatum, *Lam. Fl. Fr.* iii. 646.—Europ.
paradoxum, Reichb. Monog. Acon. 76. t. 10. f. 8 = A. Napellus.
parviflorum, Reichb. Uebers. Acon. 46 = paniculatum.
parviflorum, *Host, Fl. Austr.* ii. 70.—Europ.
pauciflorum, Bertol. Misc. xxiv. p. 17. t. 6 = A. Napellus.
pauciflorum, Host, Fl. Austr. ii. 70 = A. Lycoctonum.
pernicissimum, Reichb. Uebers. Acon. 68 = A. Lycoctonum.
Phthora, Reichb. l. c. 71. = A. Lycoctonum.
plicatum, Koehler, ex Reichb. l. c. 29 = A. Napellus.
prodanum, *Reichb. l. c.* t. 33 = A. Napellus.
pubescens, Moench, Meth. 310 = A. Napellus.
pubiceps, *Rupr. Fl. Caucas.* 42.—Reg. Caucas.
pumilum, Schar, Enum. Pl. Transs. 33 = A. Napellus.
purpureum, *Host, Fl. Austr.* ii. 74.—Europ.
pygmæum, Steud. Nom. ed. I. 10 = A. Napellus.
pyramidale, Mill. Gard. Dict. ed. VIII. n. 6 = A. Napellus.
pyrenaicum, Hort. ex Steud. Nom. ed. II. i. 20 = leptanthum.
pyrenaicum, Koelle, Spicil. 24 = A. Lycoctonum.
pyrenaicum, Linn. Sp. Pl. 532 = A. Lycoctonum.
pyrenaicum, Pall. Reise, ii. 316 = A. Anthora.
Radenianum, *Regel, in Ind. Sem. Hort. Petrop.* (1861) 48; *et Fl. Uss.* 11.—Manchuria.
ranunculifolium, Reichb. Ic. Fl. Germ. iv. 22 = A. Lycoctonum.
ranunculoides, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 78.—Russia.
reclinatum, A. Gray, in Hook. Lond. Journ. Bot. ii. (1843) 116 = A. Napellus.
recognitum, Reichb. Uebers. Acon. 42 = volubile.
rectum, Bernh. ex Reichb. l. c. 69 = A. Lycoctonum.
reflexum, Reichb. l. c. 43 = paniculatum.
reichenbachianum, Reichb. l. c. 56 = variegatum.
rigidum, Reichb. l. c. 86 = A. Napellus.
rostratum, Bernh. Ind. Sem. Hort. Erf. (1818); Reichb. Uebers. Acon. 56 = variegatum.
rotundifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 139 = A. Napellus.
rubellum, Sweet, Hort. Brit. ed. I. 9 = A. Napellus.
rubicundum, Fisch. ex Steud. Nom. ed. II. i. 20 = A. Lycoctonum.
schallnense, *F. Schmidt, Reise Amurland.* 107.—Ins. Sachal.
scandens, Muhl. ex Reichb. Uebers. Acon. 38 = uncinatum.
scandens, *Regn. New Fl. Am.* i. 55.—Am. bor.
Schleicheri, Reichb Uebers. Acon. 85 = A. Napellus.
Scrukini, *Turcz. in Bull. Soc. Nat. Mosc.* xiii. (1840) 61.—Sibir.
semigaleatum, Pall. ex Reichb. Uebers. Acon. 38 = A. Napellus.
septentrionale, Koelle, Spicil. 23 = A. Lycoctonum.
sibiricum, Poir. Encyc. Suppl. i. 113 = barbatum.
simorrhi, Siebold, ex Lindl. & Paxt. Flow. Gard. i. 17 = Fischeri.
speciosum, Otto, ex DC. Prod. i. 60 = A. Cammarum.
spicatum, Donn, Hort. Cantab. ed. X. 209 = A. Napellus.
Sprengelii, Reichb. Uebers. Acon. 32 = A. Cammarum.
squarrosum, Linn. ex DC. Syst. i. 368 = barbatum.
squarrosum, Koch, ex Reichb. Uebers. Acon. 29 = A. Napellus.
Stoerkianum, Reichb. l. c. 49 = A. Cammarum.
strictissimum, Reichb. l. c. 69 = A. Lycoctonum.
strictum, Bernh. ex DC. Syst. i. 373 = A. Napellus.
strictum, Willd. ex Reichb. Uebers. Acon. 51 = A. Lycoctonum.
taurorivum, Reichb. Monog. Acon. 67. t. 12 = A. Napellus.
tauricum, Wulf. in Jacq. Coll. ii. 112 = A. Napellus.
Tauschianum, *Steud. Nom.* ed. II. i. 20.—Sibir.
tenuifolium, Reichb. Uebers. Acon. 24 = A. Napellus.
tenuifolium, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 58 = macrorhynchum.

ACONITUM:—
tenuissimum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 65 = *A.* Lycoctonum!
thelyphonum, Reichb. Uebers. Acon. 73 = A. Lycoctonum.
thrixphonum, Reichb. l. c. 73 = A. Lycoctonum.
Thunbergianum, Rupr. Sert. Tianschan. 38 = A. Napellus.
tortuosum, *Willd. Enum. Hort. Berol.* 576.—Sibir.
toxicarium, Salisb. Prod. 375 = A. Lycoctonum.
toxicum, Reichb. Uebers. Acon. 43 = paniculatum.
tragoctonum, Reichb. l. c. 78 = A. Lycoctonum.
transsilvanicum, Larch. ex Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 165 = A. Lycoctonum.
Trattinickii, *Bess. ex Reichb. Uebers. Acon.* 75, nomen. —Hab.!
trichocarpum, *Reichb. l. c.* 39.—Hab.!
triste, Fisch. ex Steud. Nom. ed. II. i. 20 = A. Lycoctonum.
truncatum, *Regn. New Fl. Am.* i. 55.—Am. bor.
tuberosum, *Host, Fl. Austr.* ii. 73.—Europ.
tuberosum, Patr. ex Reichb. Uebers. Acon. 16 = A. Anthora.
umbraticolum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 65 = A. Lycoctonum.
uncinatum, Hort. ex Steud. Nom. ed. II. i. 20 = variegatum.
uncinatum, *Linn. Sp. Pl.* ed. II. 750.—Am. bor.; Japon.
uncinatum, Sievers, ex Pall. Nord. Beitr. vii. 229 = biflorum.
variegatum, Hook. f. & Thoms. Fl. Ind. 56 = uncinatum.
variegatum, *Linn. Sp. Pl.* 532.—Europ.
venustum, Reichb. Uebers. Acon. 38 = A. Napellus.
versicolor, Reichb. l. c. 80 = A. Cammarum.
versicolor, Ster. ex Ledeb. Fl. Ross. i. 66 = A. Anthora.
villosum, Maxim. Ind. Fl. Pek. in Prim. Fl. Amur. 468 = Kusnezoffi.
villosum, *Reichb. Uebers. Acon.* 89.—Sibir.
virgatum, Reichb. l. c. 28 = A. Napellus.
virosum, D. Don, Prod. Fl. Nep. 196 = A. Napellus.
volubile, Moench, Meth. Suppl. 110 = variegatum.
volubile, *Koelle, Spicil.* 21.—Sibir. altaic.
volubile, Muhl. Cat. 54 = uncinatum.
vulgare, DC. Syst. i. 371 = A. Napellus.
vulparia, Reichb. Uebers. Acon. 70 = A. Lycoctonum.
Wildenowii, Reichb. l. c. 35 = A. Napellus.
Wilmeitanum, Delarb. Fl. Auv. ed. II. 499 = paniculatum.
Wulfenianum, Reichb. ex Steud. Nom. ed. II. i. 20 = A. Napellus.
zoctonum, Reichb. Uebers. Acon. 65 = A. Lycoctonum.

ACONOGONUM, Reichb. Consp. 162 (1828) = **Polygonum**, Linn.
alpinum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 64 = P. alpinum.

ACONTIAS, Schott, Melet. i. 19 (1832) = **Xanthosoma**, Schott (Arnld.).
conspurcatus, Schott, Prod. 198 = X. conspurcatum.
cubensis, Schott, in Bonplandia, vii. (1859) 168 = X. cubense.
divergifolius, Regel, Ind. Sem. Hort. Petrop. (1865) 44; Gartenfl. (1866) 68 = X. platylobum.
hastifolius, Schott, Melet. i. 19 = X. hastifolium.
helleborifolius, Schott, l. c. = X. helleborifolium.
Hoffmanni, Schott, Prod. 196 = X. Hoffmanni.
lucidus, G. Don, in Sweet, Hort. Brit. ed. III. 632 = Caladium lucidum.
pentaphyllus, Schott, Syn. 65 = X. pentaphyllum.
platylobus, Schott, in Ann. Wus. Lugd. Bat. i. 279 = X. platylobum.
Plumieri, Schott, Syn. i. 65 = X. Plumieri.
Riedelianus, Schott, Prod. 192 = X. Riedelianum.
striatipes, Schott, l. c. = Caladium striatipes.
variegatus, Kunth, Enum. Pl. iii. 45 = X. helleborifolium.
Wendlandii, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 276 = X. Hoffmanni.

ACOPHORUM, Gaudich. ex Steud. Nom. ed. II. 20. (1840), nomen. *GRAMINEÆ*!
taeniolepsum, *Gaudich, l. c.*—Hab.!

ACORIDIUM, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. I. (1843) 181 = **Ceratostylis**, Blume (Orchid.).
tenellum, Nees & Meyen, l. c. = C. gracilis.

ACORUS, Linn. Gen. ed. I. 104 (1737). *AROIDEAE*, Benth. & Hook. f. iii. 999.
americanus, Rafin. New Fl. Am. i. 57 = A. Calamus ?
angustifolius, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 284 = A. Calamus.
aromaticus, Gilib. Exercit. ii. 507 = A. Calamus.
Belangeri, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 284 = A. Calamus.
brasiliensium, Schott, ex Nees, in Mart. Fl. Bras. ii. t. 170 = Sclleria hirsta.
Calamus, *Linn. Sp. Pl.* 324.—Reg. bor. temp.
Calamus-aromaticus, [Clairv.] Man. Herb. 104 = A. Calamus.
Casta, Bertol. in Mem. Acad. Sci. Bolog. Scr. II. iv. (1864) 310 = A. Calamus.
cochinchinensis, *Schott, Meiet.* i. 22.—Cochinch.
Commersonii, Schott, in Ann. Mus. Lugd. Bot. i. 284 = A. Calamus.
communiatus, Schott, Prod. 578 = A. Calamus.
elatus, Salisb. Prod. 263 = A. Calamus.
europaeus, Dum. Fl. Belg. 162 = A. Calamus.
flexuosus, Rafin. 'New Fl. Tex. 29'; Atl. Journ. 178 = A. Calamus.
floridanus, Rafin. New Fl. Am. i. 57 = A. Calamus.
gramineus, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 474. —Japon.
Griffithii, Schott, in Oestr. Bot. Zeitschr. 1858) 351 = A. Calamus
humilis, Salisb. Prod. 263 = A. gramineus.
nilaghirensis, Schott, in Oestr. Bot. Zeitschr. (1859) 101 = A. Calamus.
odoratus, Lam. Fl. Fr. iii. 299 = A. Calamus.
Palmita, Lichtenst. Reise, ii. 256 = Prionium Palmita.
pusillus, Salisb. ex Verh. Bat. Genoot. xii. (1830) 2 = graminens.
spurius, Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 284 = A. Calamus.
Tatarinowii, Schott, in Oestr. Bot. Zeitschr. (1859) 101 = A. Calamus.
terrestris, Spreng. Syst. ii. 118 = A. Calamus.
triquetor, Turcz. ex Schott, Prod. 578 = A. Calamus.
undulatus, Stokes, Bot. Mat. Med. ii. 282 = A. Calamus.
verus, Rafin. Med. Fl. i. 26 = A. Calamus.

ACOSMIA, Benth. in Wall. Cat. n. 644 (1828) = Gypsophilia, Linn. (Caryophyll.).
rupestris, Benth. l. c. = G. cerastioides.

ACOSMIUM, Schott, in Spreng. Syst. iv. Cur. Post. 406 (1827) = **Sweetia**, Spreng. (Legumin.).
lenticelfolium, Schott, l. c. = S. lenticelfolia.
lenticelfolium, Vog. in Linnaea, xi. (1837) 395 = S. fruticosa.

ACOSMUS, Desv. in Desf. Cat. Hort. Par. ed. III. 238 (1829) = **Aspicarpa**, Lag. (Malpig.).
pruriens, Desv. l. c. = A. hirtella.

ACOSTA, Adans. Fam. ii. 117 (1765) = **Centaurea**, Linn. (Compos.).

ACOSTA, DC. Prod. v. 90 (1836) = **Spiracantha**, H. B. & K. (Compos.).
Rolandrae, DC. l. c. = S. cornifolia.

ACOSTA, Lour. Fl. Cochinch. 376 (1790) = **Agapetes**, G. Don (Vaccin.).
spicata, Lour. l. c. = A. Acosta.

ACOSTA, Ruiz & Pav. Prod. 1 (1794) = **Moutabea**, Aubl. (Polygal.).
aculeata, Ruiz & Pav. Fl. Per. i. 5 = M. Acosta.

ACOUROA, Aubl. Pl. Gui. ii. 758. t. 301 (1775) = **Monstaphyllum**, Rich. (Legumin.).
violacea, Aubl. l. c. (= *Geoffraea violacea*).

ACOURTIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 208 = **Perezia**, Lag. (Compos.).
formosa, DC. Prod. vii. 66 = P. fruticosa.
formosa, D. Don, in Trans. Linn. Soc. xvi. (1830) 208 = P. turbinata.
formosa, Hook. & Arn. Bot. Beech. Voy. 487 = P. rigida.
formosa, Sch. Bip. in Seem. Bot. Voy. Herald, t. 56 = P. patens.
hebeclada, DC. in Deless. Ic. Sel. iv. t. 96 = P. hebeclada.

ACOURTIA :—
macrocephala, Sch. Bip. in Seem. Bot. Voy. Herald, t. 55 = P. turbinata.
microcephala, DC Prod. vii. 66 = P. microcephala.
moschata, DC. l. c. = P. moschata.
rigida, DC. l. c. = P. rigida.
Seemannii, Sch. Bip. in Seem. Bot. Voy. Herald, t. 54 = P. Seemannii.
turbinata, DC. Prod. vii. 66 = P. turbinata.

ACRACHNE, Wight & Arn. ex Lindl. Introd. Nat. Syst. ed. II. 381 (1836) = **Elousina**, Gaert. (Gramin.).
eleusinoides, Wight & Arn. ex Steud. Nom. ed. II. i. 21 = E. verticillata.
verticillata, Wight & Arn.—Cf. Lindl. Introd. Nat. Syst. ed. II. 381 = E. verticillata.

ACRADENIA, Kipp. in Proc. Linn. Soc. ii. (1852) 201; ex Trans. Linn. Soc. xxi. (1855) 207. t. 23. *RUTACEAE*, Benth. & Hook. f. i. 393.
Frankliniae, *Kipp. ll. cc.*—Tasmania.
sieriodes, Kipp. in Proc. Linn. Soc. ii. (1852) 201 = Franklinae.

ACRAEA, Lindl. in Benth. Pl. Hartw. 155 (1845) = **Pterichis**, Lindl. (Orchid.).
Mendesii, Reichb. f. Xenia Orch. iii. 18.—Boliv.
multiflora, Lindl. Orchid. Linden. 26.—N. Granat.
parviflia, Lindl. in Benth. Pl. Hartw. 155.—Ecuador.
trilobe, Lindl. in Ann. Nat. Hist. xv. (1845) 386.—Peruv.
Wiegmani, Reichb. f. Otia Bot. Hamb. 53.—Bras.

ACRANDRA, Berg, in Linnaea, xxvii. (1854) 485; et in Mart. Fl. Bras. ix. 459 (1857) = **Campomanesia**, Ruiz & Pav. (Myrtac.).
laurifolia, Berg, in Mart. Fl. Bras. xiv. t. 459 = C. laurifolia.
Selloviana, Berg, l. c.—Bras.
verrucosa, Berg, l. c.—Bras.

ACRANTHERA, Arn. ex Meisn. Gen. 169 (115) (1838); et in Ann. Nat. Hist. iii. (1839) 20. *RUBIACEAE*, Benth. & Hook. f. ii. 84.
ARCHBOTRYPIS, R. Br. in Wall. Cat. n. 8898 (1847 ?).
anamalicca, *Bedd. Ic. Pl. Ind. Or.* i. 5.—Ind. or.
grandiflora, *Bedd. l. c.*—Ind. or.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 92.—Reg. Himal.
laxiflora (Wight), *Hook. f. l. c.*—Moluca.
tomentosa, *Hook. f. l. c.*—Reg. Himal.
uniflora, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 212 = Smeadea uniflora.
zeylanica, *Arn. in Ann. Nat. Hist.* iii. (1839) 21.—Zeylan.

ACRATHERUM, Link, Hort. Berol. i. 230 (1827) = **Arundinella**, Raddi (Gramin.).
fucatum, Nees, ex Steud. Syn. Pl. Gram. 114 = A. fucata.
miliaceum, Link, Hort. Berol. i. 230 = Arund. nepalensis.
pumilum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 414 = Arund. pumila.

ACRIDOCARPUS, Guill. & Perr. Tent. Fl. Seneg. i. 139 (1829). *MALPIGHIACEAE*, Benth. & Hook. f. i. 256.
ANOMALOPTERIS, G. Don, Gen. Syst. i. 647 (1831).
adenophorus, *A. Juss. in Arch. Mus. Par.* iii. (1843) 481.—Madag.
angolensis, *A. Juss. l. c.* 490.—Afr. trop.
argyrophyllus, *A. Juss. l. c.* 498.—Madag.
austro-caledonicus, *Baill. Adansonia*, xi. (1873-76) 248.—N. Caled.
Casaulilleti, A. Juss. in Arch. Mus. Par. iii. (1843) t. 15 = Smeathmannii.
chloropteras, *Oliver, Fl. Trop. Afr.* i. 270.—Afr. trop.
corymbosus, *Hook. f. in Hook. Ic. Pl.* t. 774.—Afr. trop.
cuneatus, *A. Juss. in Arch. Mus. Par.* iii. (1843) 489. —Madag.
galphimiaefolius, *A. Juss. l. c.* 491.—Afr. trop.
guianensis, A. Juss. l. c. t. 15 = Smeathmannii.
Himado, *S. Moore, in Journ. Bot.* xvili. (1880) 1.—Afr. trop.
longifolius, Hook. f. in Hook. Niger Fl. 244 = Smeathmannii.
natalitius, *A. Juss. in Arch. Mus. Par.* iii. (1843) 486. —Natal.

ACRIDOCARPUS :—
orientalis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 488. —Natal.
plagiopterus, *Guill. & Perr. Tent. Fl. Senog.* i. 138. t. 29.—Senegamb.
pruricus, *A. Juss. in Arch. Mus. Par.* iii. (1843) 492. —Natal.
Smeathmanni, *Guill. & Perr. Tent. Fl. Senog.* i. 124. —Afr. trop.
zanzibaricus, *A. Juss. in Arch. Mus. Par.* iii. (1843) 485. t. 15.—Zanzibar.

ACRILIA, Griseb. Fl. Brit. W. Ind. 129 (1864) = **Trichilia**, Linn. (Meliac.).
Sloanei, Griseb. l. c. = T. terminalis.

ACRIOPSIS, Reinw. ex Blume, Cat. Gew. Buitenz. 97 (1823); Blume, Bijdr. 376 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 586.
densiflora, *Lindl. in Bot. Reg.* (1847) sub t. 20.—Borneo.
Griffithii, Reichb. in Bonplandia, ii. (1854) 92 = javanica.
indica, *Wight, Ic.* v. t. 1748.—Malaya.
javanica, *Reinw. ex Blume, Cat. Gew. Buitenz.* 97.; *Blume, Bijdr.* 377.—Malaya.
picta, Lindl. in Bot. Reg. (1842) Misc. 69 = javanica.

ACRIULUS, Ridl. in Journ. Linn. Soc. xx. (1884) 336. *CYPERACEAE*.
griegiflolius, *Ridl. l. c.*—Madag.
madagascariensis, *Ridl. l. c.*—Madag.

ACROANTHES, Rafin. in Journ. Phys. lxxxix. 261 (1819) = **Microstylis**, Lindl. (Orchid.).

ACROCARPIDIUM, Miq. in Versl. en Meded. Kon. Nederl. Inst. '1842' (1843) 198. t. 2 = **Peperomia**, Ruiz & Pav. (Piperac.).
brevipes, Benth. Bot. Voy. Sulph. 166 = P. brevipes.
cordifolium, Miq. Syst. Pip. 56 = P. cordifolia.
exile, Miq. l. c. = P. exilis.
guayaquilense, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 411 = P. guayaquilensis.
Guildingianum, Miq. l. c. 412 = P. Guildingiana.
hispidulum, Miq. Syst. Pip. 54 = P. hispidula.
Yamamotoanum, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 412 = P. Jamesoniana.
majus, Miq. Syst. Pip. 60 = P. hederacea.
muricatum, Miq. l. c. 58 = P. metaquiloensis.
molle, Miq. in Versl. en Meded. Kon. Nederl. Inst. '1843' (1843) 199 = P. mollis.
nummulariifolium, Miq. l. c. 198. t. 2 = P. nummularifolia.
pulicare, Miq. l. c. 199 = P. scandens.
repens, Miq. l. c. = P. scandens.
rotundifolium, Miq. l. c. = P. circularis ?
scandens, Miq. Syst. Pip. 59 = P. scandens.
Sellowianum, Miq. l. c. 55 = P. hispidula.
serpens, Miq. in Versl. en. Meded. Kon. Nederl. Inst. '1842' (1843) 199 = Verhenilia serpens.
tenellum, Miq. l. c. = P. tenella.
urocarpum, Miq. Syst. Pip. 63 = P. urocarpa.
uroperanum, Schlecht. in Linnaea, xxvii. (1854) 486 = prace. !

ACROCARPUS, Nees, in Mart. Fl. Bras. ii. t. 158. t. 16, 17 (1842) = **Cryptangium**, Schrad. (Cyperac.).
campestris, Nees, l. c. 160 = C. campestre.
densiflolius, Nees, l. c. 159 = C. densiflolium.
humilis, Nees, l. c. 161 = C. humile.
junciformis, Nees, l. c. = C. junciforme.
minarum, Nees, l. c. 159 = C. minarum.
polyphyllus, Nees, l. c. 160 = C. polyphyllum.
stellatus, Nees, ex Boeck. in Linnaea, xxxvili. (1874) 422 = C. stellatum.

ACROCARPUS, Wight, ex Arn. in Mag. Zool. & Bot. ii. (1839) 547. *LEGUMINOSAE*, Benth. & Hook. f. i. 562.
acuminifolius, *Teysm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1866) 265.—Malaya.
fraxinifolius, *Wight & Arn. in Mag. Zool. & Bot.* ii. (1839) 547.—Ind. or.
gemmiflo, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 87.—Sumatra.

ACROCENTRON, Cass. in Dict. Sc. Nat. xlv. 37. (1826) = Centostrum, Linn. (Compos.).
Botteri, Webb, Iter Hisp. 33 = C. Botteri.

ACROCENTRON :—
collinum, Cass. in Dict. Sc. Nat. xlvii. 37 = C. collina.
granatense, Webb, Iter Hisp. 89 = C. granatensis.
ornatum, Webb, l. c. = C. ornata.
tenuifolium, Cass. in Dict. Sc. Nat. l. 254 = C. rupestris.

ACROCEPHALUS, Benth. in Bot. Reg. sub tt. 1282, 1300 (1829). *LABIATAE*, Benth. & Hook. f. ii. 1178. axillaris, *Benth. in DC. Prod.* xii. 48.—Ind. or.
Blumei, Benth. in Bot. Reg. sub t. 1300 = capitatus.
caeruleus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 135.—Afr. trop.
capitatus, *Benth. in Bot. Reg.* sub t. 1300.—Ind. or.; China.
cylindraceus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 135.—Afr. trop.
lilacinus, *Oliver, l. c.*—Afr. trop.
scariosus, Benth. in Bot. Reg. sub t. 1282 = capitatus.
verbenaceus, *Vatke, in Linnaea*, xl. (1876) 178 — Afr. trop.
villosus, *Benth. in Bot. Reg.* sub t. 1300; *Lab. Gen. et Sp.* 23.—Madag.

ACROCHAENE, Lindl. Fol. Orch. (1858). *ORCHIDEAE*, Benth. & Hook. f. iii. 507.
punctata, *Lindl. l. c.*—Ind. or
Rimanni, *Reichb. f. in Gard. Chron.* N. S. xvii. (1882) 1. 796.—As. trop.

ACROCLINIUM, A. Gray, in Hook. Kew Journ. iv. (1852) 270 = **Helipterum**, DC. (Compos.).
multicaule, A. Gray, l. c. 271 = H. chlorocephalum.
phyllocephalum, A. Gray, l. c. = Podotheca fuscescens.
roseum, Hook. Bot. Mag. t. 4801 = H. roseum.
rubellum, A. Gray, in Hook. Kew Journ. iv. (1852) 71 = H. rubellum.

ACROCOMIA, Mart. Hist. Nat Palm. ii. 66. tt. 56, 57 (1828). *PALMAE*, Benth. & Hook. f. iii. 943.
aculeata, Lodd. ex Mart. l. c. iii. 286 = sclerocarpa.
cubensis, Lodd. ex H. Wendl. Ind. Cycl. Pand. Cycad. 1 = lasiospatha.
fusiformis, Sweet, Hort. Brit. ed. 1. 432 = lasiospatha?
glaucophylla, *Drude, in Mart. Fl. Bras.* iii. ii. 392.—Bras.
globosa, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 286.—Ins. S. Vincent.
guianensis, *Lodd. ex G. Don, in Loud. Hort. Brit.* 382, nomen.—Guiana.
horrida, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 286.—Ins. Trinit.
intumescens, *Drude, in Mart. Fl. Bras.* iii. ii. 391.—Bras.
lasiospatha, *Mart. in Palm. Orbign.* 81.—Ind. occ.
lasiospatha, Wallace, Palms Amaz. 97. t. 37 = sclerocarpa.
mexicana, *Karw. ex Mart. Hist. Nat. Palm.* iii. 285. t. 138.—Mexico.
minor, Lodd. ex G. Don, in Loud. Hort. Brit. 382 = lasiospatha?
sclerocarpa, *Mart. Hist. Nat. Palm.* ii. 66. tt. 56, 57, 100. f. 5.—Am. trop.
sphaerocarpa, Desf. Cat. Hort. Par. ed. III. 80 = sclerocarpa.
tenuifrons, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 286.—Ins. S. Domingo.
Totai, *Mart. in Palm. Orbign.* 78. t. 9. f. 1.—Boliv.
viridens, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 47. —Nicaragua.
Zapotecis, Karw. ex H. Wendl. in Kerch. Palm. 230 = mexicana.

ACROCORION, Adans. Fam. ii. 57 (1763) = **Gelanthus**, Lim. (Amaryll.).

ACROCORYNE, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 314 = **Monostelma**, R. Br. (Asclep.).
cordiuse, Turcz. l. c. = M. balsamosum.

ACRODICLIDIUM, Nees, Lauxin. Disp. Progr. 18 (1838). *LAURINEAE*, Benth. & Hook. f. iii. 154.
TRIPLOSPERA, Rafin. Sylva Tellur. 134 (1838).
anacardioides, *Spruce, ex Meisn. in DC. Prod.* xv. t. 86.—Bras.
brasiliense, Nees, Syst. Laurin. 267 = Aydendron brasiliense.
Cassure, *Schomb. in Hook. Lond. Journ. Bot.* iii. (1844) 699.—Guiana.

ACRODICLIDIUM :—
chrysophyllum, *Meisn. in DC. Prod.* xv. i. 87.—Guiana.
dominicense, *Meisn. l. c.* 86.—Ins. Dominic.
Gardneri, *Meisn. in DC. Prod.* xv. i. 87.—Bras.
guianense, *Nees, Syst. Laurin.* 268.—Guiana.
ilu-uba, *Meisn. in DC. Prod.* xv. i. 86.—Bras.
Jamaicense, Griseb. Fl. Brit. W. Ind. i. 280, ex parte = Wrightii.
jamaicense, *Nees, Syst. Laurin.* 270.—Jamaic.
Kunthianum, Meisn. l. c. 269 = Aydendron aciphyllum.
oppositifolium, *Nees, in Linnaea*, xxi. (1848) 500.—Guiana.
salicifolium, *Griseb. Fl. Brit. W. Ind.* i. 280.—Ind. occ. sericeum, *Griseb. l. c.* 280.—Ins. Dominic.
Spruceі, *Meisn. in DC. Prod.* xv. i. 86.—Bras.
Wrightii, *Meisn. l. c.* 85.—Cuba.

ACRODRYON, Spreng. Syst. i. 865 (1825) = **Cephalanthus**, Linn. (Rubiac.).
angustifolius, Spreng. l. c. 886 = C. angustifolius.
orientalis, Spreng. l. c. 886 = C. orientalis.

ACROELYTRUM, Steud. in Flora, xxix. (1846) 20 = Lophatherum, Brongn. (Gramin.).
japonicum, Steud. l. c. 21 = L. elatum.

ACROGLOCHIN, Schrad. Cat. Hort. Goett. ex Schult. f Mant. i. 89, 227 (1822). *CHENOPODIACEAE*. Benth. & Hook. f. iii. 48.
BLITANTHUS, Reichb. Cat. Hort. Dresd. (1824).
LECANOCARPUS, Nees, Fl. Hort. Bonn. 4. t. 2. (1823).
chenopodioides, *Schrad. ex Schult. f. Mant.* i. 227.—Reg. Himal.; China.
persicarioides, Moq. in DC. Prod. xiii. ii. 354 = chenopodioides.
Schraderiana, Schult. f. ex Moq. l. c = chenopodioides

ACROGLYPHE, E. Mey. Zwei Pfl. Docum 100 (1843) = **Annesorhiza**, Cham. & Schlecht. (Umbell.).
flexuosa, E. Mey l. c. 100 = Ann. hirsuta + montana.
hispida, Sond. in Harv. & Sond. Fl. Cap. ii. 546 = Ann. villosa.
hispidula, E. Mey. Zwei Pfl. Docum. 74 = Ann. villosa.
runcinata, E. Mey. l. c. 100.—Afr. austr.

ACROLEPIS, Schrad. in Comm. Gotting. vii. (1832) 142 = **Stoinia**, Schrad. (Cyperac.).
ferruginea, Nees, ex Boeck. in Linnaea, xxxvii. (1871–73) 91.—Afr. austr.
ramosissima, Boeck. l. c. 89 = F. ramosissima.
trichodes, Schrad. in Comm. Gotting. vii. (1832) 142 = F. trichodes.

ACROLOBUS, Klotzsch, in Verh. Akad. Berl. (1856) 236. t. 3 = **Keisteria**, Linn. (Olacin.).
parvifolia, Klotzsch, l. c. 237. t. 3 = H. parvifolia.
Schomburgkiana, Klotzsch, l. c. 236. t. 3 = H. parvifolia.

ACROLASIA, Presl, Rel. Haenk. ii. 89. t. 55. (1831) = **Mentzelia**, Linn. (Loasac.).
bartonioides, Presl, l. c. 89 = M. albicaulis.
Salsoria, C. Gay, Fl. Chil. ii. 480.—Chili.
squalida, Hook. f. in Trans. Linn. Soc. xx. (1851) 222 = M. aspera?

ACROLOPHUS, Cass. in Dict. Sc. Nat. l. (1827) 253 = **Centaurea**, Linn. (Compos.).
maculosus, Cass. l. c. 253 = C. maculosa.
Mierghii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 92 = C. paniculata.
paniculatus, Cass. in Dict. Sc. Nat. l. 253 = C. paniculata.
rupicolus, Webb, ex Steud. Nom. ed. II. i. 23 = C. rupicola.
rigidulus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 97 = C. paniculata.
subalbidus, Fourr. l. c. = C. maculosa.
tenuisectus, Fourr. l. c. = C. maculosa.

ACRONEMA, Falcon. ex Edgew. in Trans. Linn. Soc. xx. (1851) 51 = **Pimpinella**, Linn. (Umbellif.).
tenerum, Edgew. l. c. = P. tenera.

ACRONIA, Presl, Rel. Haenk. i. 108 (1827); Symb. Bot. ii. 9 (1856) = **Cryptostylis**, R. Br. (Orchid.).
phalangifera, Presl, ll. cc. 104, 10.—Peruv.

ACRONODA, Hassk. Cat. Hort. Bogor. Alt. 208 (1844) = seq.

ACRONODIA, Blume, Bijdr. (1825) 128 = **Blasocarpus**, Linn. (Tiliac.).
punctata, Blume, l. c. = E. Acronodia.

ACRONOZUS, Steud. Nom. ed. II. i. 21 (1840) = praec.

ACRONYCHIA, Forst. Char. Gen. 53. t. 27 (1776) *RUTACEAE*, Benth. & Hook. f. i 302.
CUNTO, Adans. Fam. ii 446 (1763).
CYMINOSMA, Gaert. Fruct. i. 260. t. 53 (1788).
DORRIENIA, Dennst. Schluss. Hort. Malab. 31 (1818).
GELA, Lour. Fl. Cochinch. 222 (1790).
HUONIA, Montrous. in Mém. Acad. Lyon, x. (1860) 185.
JAMBOLANA, Adans. Fam. ii. 506 (1763).
JAMBOLIFERA, Linn. Dis. Dass. 15 (1747); Amoen Acad. i. 398 (1749).
KOELPINIA, Scop. Introd. 281, et Ind. (1777).
SELAS, Spreng. Syst. ii. 216 (1825).
acidula, *F. Muell. Fragm.* iv. 154.—Austral.
apiculata, *Miq. Fl. Ind. Bat Suppl.* 582.—Sumatra
arborea, *Blume, Bijdr.* 1144.—Java.
Baueri, *Schott, Fragm. Bot.* 5. t. 8.—Austral.
Cunninghamii, Hook. Bot. Mag. t. 3994 = Medicosma Cunninghamii.
Endlicheri, *Schott, Fragm. Bot.* 8. t. 9.—Ins. Norfolc.
Halmaheirae, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 245. —Malaya.
heterophylla, *A. Gray, Bot. U. S. Expl. Exped.* i. 338. t. 25.—Ins. Samoa.
Hillii, F. Muell. Fragm. i. 26 = Baueri.
imperforata, *F. Muell. l. c.*—Austral.
laevis, *Forst. Char. Gen* 54. t. 27.—Austral.
laurifolia, *Blume, Cat. Gew. Buitenz.* 27. 68; *Bijdr* 245.—As. trop.
laurina, F. Muell. 3rd Rep. Veg. Vict. 7; Fragm. i. 27 = laevis.
melicopoides, *F. Muell. Fragm.* v. 3.—Austral.
Minahassae, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 245 —Ins. Celebes.
oblongifolia, Endl. ex Heynh Nom. ii. 8 = laevis.
ovata, Endl. ex Heynh. l. c. = Baueri.
pedunculata, Miq. Fl. Ind. Bat. Suppl. 582 = laurifolia
petiolaris, *A. Gray, Bot. U. S. Expl. Exped.* i. 335. t. 33.—Ins. Fiji.
Porteri, *Hook. f. Fl. Brit. Ind.* i. 498.—Malaya.
retusa, *A. Gray, Bot. U. S. Expl. Exped.* i. 338. t. 34. —Ins. Samoa.
Richii, *A. Gray, l. c.* 336. t. 33.—Ins. Samoa.
simplicifolia, Steud. Nom. ed. II. i. 21 = Endlicheri tetrandra, *F. Muell. Fragm.* ix. 104.—Austral.
trifoliolata, *Zoll. & Mor. in Nat. en Genees. Arch Neerl. Ind.* ii. (1845) 585.—Java.
venitis, *F. Muell. Fragm.* iv. 155.—Austral.

ACROPERA, Lindl. Gen. et Sp. Orch. 172 (1838) = **Gongora**, Ruiz & Pav. (Orchid.).
armeniaca, Lindl. in Lindl. & Paxt. Flow. Gard. i. (1850–51) 94 = G. armeniaca.
atropurpurea, Hort. ex Heynh. Nom. iii. 8 = G. galeata.
Batemanni, Lindl. ex Reichb. f. in Gard. Chron. N. S. ii. (1874) 1. 322 = G. galeata.
citrina, Hort. ex Reichb. f. Xenia Orch. i. 51 = G. galeata.
cornuta, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 186 = G. armeniaca.
flavida, Klotzsch, l. c. xix. (1851) 317 = G. galeata.
fuscata. Hort. Belg. ex Heynh. Nom. iii. 8.—Mexico.
grandiflora, Hort. ex Pasq. Cat. Orto Bot. Napol. 8. nomen.—Hab.?
Loddigesii, Lindl. Gen. et Sp. Orch. 172 = G. galeata.
luteola, Hort. ex Heynh. Nom. ii. 8 = *fuscata*.
pallida, Hort. ex Heynh. l. c. = G. galeata.
purpurea, Hort. ex Heynh. l. c. = G. galeata.
sulphurea, Hort. ex Heynh. l. c. = *fuscata*.

ACROPETALUM, A. Juss. in Orb. Dict. xii. 617 (1840) = XEROPETALUM, Delile = **Dombeya**, Cav. (Stercul.).

ACROPHYLLUM, Benth. in Maund, Bot. ii. t. 95 (1840). *SAXIFRAGEAE*, Benth. & Hook. f. i. 652.
CALYCOMIS, D. Don, in Edinb. N. Phil. Journ. [ix.] (July—Oct. 1830) 93.

ACROPHYLLUM :—
venosum, *Benth. in Maund, Bot.* ii. t. 95.—Austral.
verticillatum, Hook. Bot. Mag. t. 4050 = venosum.

ACROPHYLLUM, E. Mey Zwei Pfl. Docum. 140 (1843) = **Kiggelaria**, DC. (Hixin.).
obtusatum, C. Muell. in Walp. Ann. vii. 283 = K. africana.
obtusum, E. Mey. Zwei Pfl. Docum. 140 = K. africana.

ACROPODIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 408 = **Aspalathus**, Linn. (Legumin.).
suffruticosum. Desv. l. c. = Asp. suffruticosa.

ACROPSELION, Spach, Hist. Vég. Phan. xiii. 168 (1846) = ACROSPELION, Bess = **Trisetum**, Pers. (Gramin.).

ACROPTILON, Endl. Ench. 247 (1841) = seq.

ACROPTILON, Cass. in Dict. Sc. Nat. I. 464 (1827) = **Centaurea**, Linn. (Compos.).
angustifolium, Cass. l. c. 466 = C. Picris.
obtusifolium, Cass. l. c 465 = C. Picris.
Picris, DC. Prod. vi. 662 = C. Picris.
repens, DC. l. c. 663 = C. Picris.
serratum, Cass. in Dict. Sc. Nat. I. 466 = C. Picris.
subdentatum, Cass. l. c. 465 = C. Picris.

ACROSANTHES, Eckl. & Zeyh. Enum. 328 (1837).
FICOIDEAE, Benth. & Hook. f. i. 855.
DIDASTR, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 472 (1862).
anceps, *Sond. in Harv. & Sond. l. c.*—Afr. austr.
angustifolia, *Eckl. & Zeyh. Enum.* 328.—Afr. austr.
decandra, *Fenzl, in Ann. Wien. Mus.* ii. (1838) 270. —Afr. austr.
fistulosa, *Eckl. & Zeyh. Enum.* 328.—Afr. austr.
humifusa, *Sond. in Harv. & Sond. Fl. Cap.* ii. 472.—Afr. austr.
teretifolia, *Eckl. & Zeyh. Enum.* 329.—Afr. austr.

ACROSPELION, Bess. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 59 = **Trisetum**, Pers. (Gramin.).

ACROSPIRA, Welw. ex Baker, in Trans. Linn. Soc. Bot. Ser. II. i. (1875) 255. t. 34. *LILIACEAE*, Benth. & Hook. f. iii. 787.
saphodeoloides, *Welw. l. c.*—Afr. trop.

ACROSSANTHES, Presl, Bot. Bemerk. 23 (1844) = **Vismia**, Vell. (Hypericin.).
Lhotskyanus, Presl, l. c. = V. decipiena.

ACROSTEMON, Klotzsch, in Linnaea, xii. (1838) 227 = **Grimschachia**, Klotzsch (Tetrac.).
equisetoides, Klotzsch, l. c. 228.—Afr. austr.
glandulosus, Rach, l. c. xxvi. (1858–55) 790.—Afr. austr.
hirsutus, Klotzsch, l. c. xii. (1838) 228.—Afr. austr.
incanus, Klotzsch, l. c. 228 = G. incana.
incurvus, Benth. in DC. Prod. vii. 702.—Afr. austr.

ACROTAPHROS, Stend. ex A. Rich. Tent. Fl. Abyss. i. 207. t. 38 [1847] = **Ormocarpum**, Beauv. (Legum.).
bibracteata, Stend. ex A. Rich. l. c. = O. bibracteatum.

ACROTICHE, Poir. in Dict. Sc. Nat. i. Suppl. 50 (1816) = **Acrotriche**, R. Br. (Epacrid.).

ACROTOMIA, Benth. in Endl. Gen. 937 (1838). *LABIATAE*, Benth. & Hook. f. ii. 1206.
hispida, *Benth. in DC. Prod.* xii. 436.—Afr. austr.
inflata, *Benth. l. c.*—Afr. austr.
pallescens, *Benth. l. c.*—Afr. austr.

ACROTREMA, Jack, in Malay Misc. i. (1820) Art. V. p. 36. *DILLENIACEAE*, Benth. & Hook. f. i. 13.
Arnottianum, *Wight, Illustr. Ind. Bot.* i. 9. t. 3.—Ind. or.
bullatum, Thw. Enum. Pl. Zeyl. 20 = uniforum.
costatum, *Jack, in Malay Misc.* i. (1820) Art. V. p. 36. —Ind. or.; Malaya.
costatum, Wall. Cat. n. 1117 B = Arnottianum.
dissectum, *Thw. in Hook. Kew Journ.* viii. (1856) 242. —Zeylan.

ACROTREMA :—
Gardneri, *Thw. Enum. Pl. Zeyl.* 2.—Zeylan.
intermedium, *Thw. l. c.*—Zeylan.
lanceolatum, *Hook. Ic. Pl.* sub t. 157.—Zeylan.
lyratum, *Thw. in Hook. Kew Journ.* viii. (1856) 242. —Zeylan.
sylvestre, *Thw. Enum. Pl. Zeyl.* 2 = uniforum.
Thwaitesii, *Hook. f. & Thoms. in Hook. Kew Journ.* viii. (1856) 241.—Zeylan.
uniforum, *Hook. Ic. Pl.* t. 157.—Zeylan.
Walkeri, *Wight, ex Thw. Enum. Pl. Zeyl.* 2.—Zeylan.
Wightianum, Wall. Cat. n. 3669 = costatum.

ACROTRICHE, R. Br. Prod. 547 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 614.
ERORELLIA, Regel, Gartenfl. i. (1852) 164. t. 18.
affinis, DC. Prod. vii. 757 = serrulata.
aggregata, *R. Br. Prod.* 547.—Austral.
aristata, Benth. in Enum. Pl. Hueg. 76 = Leucopogon cuspidatus.
cordata, R. Br. Prod. 548 = ovalifolia.
depressa, *R. Br. l. c.*—Austral.
divaricata, *R. Br. l. c.* 547.—Austral.
fasciculiflora, *Benth. Fl. Austral.* iv. 239.—Austral.
latifolia, A. Cunn. ex DC. Prod. vii. 757 = Leucopogon cordifolius.
Mangleszii, Sond. in Lehm. Pl. Preiss. i. 336 = ramiflora.
ovalifolia, *R. Br. Prod.* 548.—Austral.
patula, Hook. f. Fl. Tasm. i. 258 = serrulata.
paniia, *R. Br. Prod.* 547.—Austral.
prostrata, F. Muell. in Trans. Vict. Inst. (1855) 40 = serrulata.
ramiflora, *R. Br. Prod.* 547.—Austral.
ramiflora, Sond. in Linnaea, xxvi. (1853–55), 251 = fasciculiflora.
serrulata, *R. Br. Prod.* 547.—Austral.
subcordata, DC. Prod. vii. 757 = ovalifolia.

ACROXIS, Trin. ex Steud. Nom. ed. II. i. 22 (1840) = **Muehlenbergia**, Schreb. (Gramin.).

ACROZUS, Spreng. Syst. iv. Cur. Post. 145, 149 (1827) = **Rinsocarpus**, Linn. (Tiliac.).
punctatus, Spreng. l. c. = E. Acrocodia.

ACRYPHYLLUM, Lindl. Introd. Nat. Syst. ed. II. 156 (1836) err. typ. = ARCYPHYLLUM, Ell. = **Tephrosia**, Lour. (Legum.).

ACTAEA, [Tourn.] Linn. Syst. ed. I (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 9.
CHRISTOPHORIANA, Tourn. ex Moench, Meth. 276 (1794).
acuminata, Wall. Cat. n. 4726 = spicata.
alba, *Mill, Gard. Dict.* ed. VIII. n. 2.—Am. bor.
americana, Pursh, Fl. Am. Sept. 366 = alba.
arguta, Nutt. ex Torr. & Gray, Fl. N. Am. i. 35 = spicata.
aspera, Lour. Fl. Cochinch 382. = Tetracera sarmentosa.
brachypetala, var. a, DC. Syst. I. 535 = alba.
brachypetala, var. β, DC. l. c. = spicata.
Christophoriana, Gouan, Fl. Monsp. 152 = spicata.
Cimicifuga, Hook. Fl. Bor. Am. i. 26 = Cimicifuga elata.
Cimicifuga, Linn. Sp. Pl. 504 = Cimicifuga foetida.
cordifolia, DC. Syst. I. 535 = Cimicifuga cordifolia.
corymbosa, Stokes, Bot. Mat. Med. iii. 176 = spicata.
dahurica, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 = Cimicifuga foetida.
davurica, Franch. Pl. David. 28 = Cimicifuga davurica.
dioica, *Walt. Fl. Carol.* 152.—Carolina.
erythrocarpa, Fisch. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 20 (1835) = spicata.
frigida, Wall. Cat. n. 4725 = Cimicifuga foetida.
grandis, D. Dietr. Syn. Pl. iii. 233 = Trautvetteria palmata.
gyrostachya, Wender. Ind. Sem. Hort. Marb. (1840).—Cf. Linnaea, xv. (1841), Litt. 100 = Trautvetteria palmata.
japonica, Thunb. Fl. Jap. 231 = Cimicifuga japonica.
longipes, Spach, Hist. Vég. Phan. vii. 388 = spicata.
macrophylla, Baill. in Bull. Soc. Linn. Par. i. (1879) 224 = Anemopaeonia macrophylla.
macropoda, Turcz. ex Fisch. & Mey. 6de Ledeb. Fl. Ross. i. 72 = Cimicifuga foetida.
monogyna, Walt. Fl. Carol. 151 = Cimicifuga racemosa.
nigra, Gaert. Mey. & Scherb. Fl. Watt. ii. 365 = spicata.

ACTAEA :—
orthostachya, Wender. Ind. Sem. Hort. Marb. (1840). —Cf. Linnaea, xv. (1841) Litt. 100 = Cimicifuga racemosa.
pachypoda, Ell. Sketch, ii. 15 = alba.
palmata, DC. Syst. I. 535 = Trautvetteria palmata.
pentacarpa, Michx. ex Torr. & Gray, Fl. N Am. i. 39 = Cimicifuga americana.
pentagyna, Walt. Fl. Carol. 151 = Cimicifuga americana.
podocarpa, DC. Syst. i. 283 = Cimicifuga americana.
podocarpa, Schlecht. in Linnaea, vi. (1831) 588 = Cimicifuga foetida.
pteropoma, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 = Cimicifuga foetida.
racemosa, Gener. ex Steud. Nom. ed. II. i. 22 = Cimicifuga foetida.
racemosa, Gilib. Fl. Lituan. ii. 238 = spicata.
racemosa, Linn. Sp. Pl. 504 = Cimicifuga racemosa.
racemosa, Walt. ex Steud. Nom. ed. II. i. 22 = Trautvetteria palmata.
rubra, Bigel. Fl. Bost. ed. II. 211 = alba !
rubra, Willd. Enum. Hort. Berol. 561; Ledeb. Fl. Alt. ii. 275 = spicata.
simplex, Wormsk. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 = Cimicifuga foetida.
spicata, *Linn. Sp. Pl.* 504.—Reg. bor. temp.

ACTAEOGETON, Steud. Nom. ed. II. i. 22 (1840) = **Scirpus**, Linn. (Cyperac.).

ACTEGETON, Blume, Bijdr. 1143 (1826) = **Anima**, Lam. (Salvador).
sarmentosum, Blume, l. c. 1144 = A. sarmentosa.

ACTEGITON, Endl. Gen. 1089 (1840), err. typ. = praec.

ACTEPHILA, Blume, Bijdr. 581 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 269.
ANOMOSPERMUM, Dalz. in Hook. Kew Journ. iii. (1851) 298.
LITHOXYLON, Endl. Gen. 1122 (1840).
bantamensis, Miq. Fl. Ind. Bat. i. 11. 356 = excelsa.
excelsa, *Muell. Arg. in Linnaea*, xxxii. (1863) 78.—Ind. or.; Malaya.
grandifolia, *Baill. Adansonia*, vi. (1865–66) 330.—Austral.
javanica, Miq. Fl. Ind. Bat. i. 11. 356 = excelsa.
latifolia, *Benth. Fl. Austral.* vi. 80.—Austral.
major, *Muell. Arg. in Linnaea*, xxxii. (1863) 77.—Java.
Mooreana, *Baill. Adansonia*, vi. (1865–66) 330.—Austral.
neilgherrensis, Wight, Ic. t. 1910 = excelsa.
petiolaris, *Benth. Fl. Austral.* vi. 80.—Austral.
puberula, *Kurz, in Journ. As. Soc Beng.* xlii. (1873) ii. 236.—Ins. Andaman.
rectinervia, *Kurz, in Journ. Bot.* xiii. (1875) 329.—Ins. Nicobar.
sessilifolia, *Benth. Fl. Austral.* vi. 90.—Austral.
Thomsoni, Muell. Arg. in Linnaea, xxxiv. (1865) 65 = excelsa.
weylandica, Muell. Arg. l. c. 77 = excelsa.

ACTIMERIS, Raün. in Am. Monthly Mag. (1819) 195 = **Actinomeris**, Nutt. (Compos.).

ACTINANTHUS, Ehrenb. in Linnaea, iv. (1829) 393. *ACANTHACEAE*, Benth. & Hook. f. i. 908.
syriacus, *Ehrenb. l. c.*—As. mis.; Syria.

ACTINEA, Juss. in Ann. Mus. Par. ii. (1803) 425. t. 61 = Cephalophora, Cav. (Compos.).
acaulis, Spreng. Syst. iii. 574 = Actinella acaulis.
alternifolia, Spreng. l. c. = C. heterophylla.
biternata, Spreng. l. c. = Coreopsis biternata.
chrysanthemoides, H. B. & K. Nov. Gen. et Sp. iv. 298. t. 411 = Hymenonis chrysanthemoides.
communata, Spreng. Syst. iii. 574 = Calea pinnatifida.
heterophylla, Juss. in Ann. Mus. Par. ii. (1803) 425 = C. heterophylla.
integrifolia, H. B. & K. Nov. Gen. et Sp. iv. 297. t. 410 = Helenium integrifolium.
integrifolia, *Torr. in Ann. Lyc. New York*, ii. (1828) 213.—Am. bor.
ianata, Stend. Nom. ed. I. 15 = Bahia ianata.
ophotifolia, Spreng. Syst. iii. 574 = Blainvillea bahiensis.

ACTINELLA, Nutt. Gen. Am. ii. 178 (1818). *COMPOSITAE*, Benth. & Hook. f. ii. 414.
PHILOZORA, Buckl. in Proc. Ac. Sc. Philad. '1861.' (1862) 459.
PICRADENIA, Hook. Fl. Bor. Am. i. 317. t. 108 (1833).
acaulis, *Nutt. Gen. Am.* ii. 178.—Am. bor.
argentea, *A. Gray, in Mem. Am. Ac. iv.* (1849) (*Pl. Fendl.*) 100.—N. Mexico.
biennis, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 378.—Am. bor. occ.
Bigelovii, *A. Gray, Pl. Wright.* ii. 96, in adnot.—Mexico.
Brandegei, *T. C. Porter, ex A. Gray, in Proc. Am. Acad.* xiii. (1878) 378.—Am. bor. occ.
chrysanthemoides, *A. Gray, l. c.* xix. (1883) 82.—Mexico.
Cooperi, *A. Gray, l. c.* vii. (1868) 359.—Calif.
depressa, *Harv. & Gray, ex Gray, in Mem. Am. Acad.* iv. (1849) (*Pl. Fendl.*) 100.—Am. bor. occ.
glabra, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 379 = acaulis.
grandiflora, *Torr. & Gray, in Bost. Journ. Nat. Hist.* v. (1847) 110.—Am. bor. occ.
insignis, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xviii. (1883–83) 109.—Am. bor. occ.
lanata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 379 = acaulis.
lanata, Pursh, Fl. Am. Sept. ii. 560 = Eriophyllum caespitosum.
lanuginosa, Buckl. in Proc. Ac. Sc. Philad. (1862) 459 = scaposa.
leptoclada, *A. Gray, in Pacif. Rail Rep.* iv 107.—Am. bor. occ.
linearifolia, *Torr. & Gray, Fl. N. Am.* ii. 383.—Texas.
odorata, *A. Gray, in Mem. Am. Acad. iv.* (1849) (*Pl. Fendl.*) 101.—Am. bor. occ.
Palmeri, *A. Gray, in Proc. Am. Acad.* xix. (1883) 81.—Mexico.
Richardsonii, Nutt. in Trans. Am. Phil. Soc N. S. vii. (1841) 379 = biennis.
Rusbyi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 83.—N. Mexico.
scaposa, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 379.—Am. bor. occ
Torreyana, Nutt. l. c. = acaulis.
Vaseyi, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 213.—Am. bor. occ.

ACTINELLA, Pers. Syn. ii. 489 (1807) = **Cephalophora**, Cav. (Compos.).
heterophylla, Pers. l. c. = C. heterophylla.

ACTINIDIA, Lindl. Introd. Nat. Syst. ed. II. 439 (1836). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 184.
KALOMIKTA, Regel, in Bull. Phys. Math. Acad. Pétersb. xv. (1857) 219.
TROCHOSTIGMA, Sieb. & Zucc. in Abh. Akad. Wiss. Muench. iii. (1843) 726. t. 2. f. 2.
acuminata, *Budisch. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 167.—Manchur.
arguta, Franch. & Sav. Enum. Pl. Jap. i. 58 = callosa.
callosa, *Lindl. Introd. Nat. Syst.* ed. II. 439.—Reg. Himal.; Japon.; Manchur.
Championi, *Benth. Fl. Hongk.* 26.—China.
chinensis, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 308.—China.
cordifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 15.—Japon.
Davidii, *Franch. Pl. Davidian.* i. 57.—China.
eriantha, *Benth. in Journ. Linn. Soc.* v. (1861) 55.—Hab.?
fulvicoma, *Hance, in Journ. Bot.* xxiii. (1885) 321.—China.
Kolomikta, Maxim. Prim. Fl. Amur. 63 = callosa.
platyphylla, *A. Gray, in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 15.—Japon.
polygama, *Franch. & Sav. Enum. Pl. Jap.* i. 59.—Japon.
rufa, Franch. & Sav. l. c. 58 = callosa.
stipulata, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 55.—Reg. Himal.
sutchuensis, Budisch. ex Trautv. in Act. Hort. Petrop. viii. (1883) 167 = acuminata.
volubilis, *Franch. & Sav. Enum. Pl. Jap.* i. 59.—Japon.

ACTINOBOLE, Endl. Gen. Suppl. iii. 70 (1848) = Craspedolobium, A. Gray (Compos.).

ACTINOCARPUS, R. Br. Prod. 342 (1810) = Damasonium, Juss. (Alismac.).
australis, Spreng. in Esch & Gruber, Encyc. i. 848 = D. australe.
Damasonium, Sm. in Rees, Cycl. Suppl. n. 1 = D. stellatum.
europaeus, Spreng in Esch & Gruber, Encyc. i. (1818) 848 = D. stellatum.
mator, Bercht. & Presl, Rostl. i. Alismac. 8 = D. stellatum.
minor, R. Br. Prod. 343 = D. australe.

ACTINOCARYA, Benth in Benth. & Hook. f. Gen ii. 846 (1876). *BORAGINEAE*, Benth. & Hook. f. ii. 846.
tibetica, *Benth l. c.*—Reg. Himal.

ACTINOCHLOA, Willd. ex Beauv. Agrost. 41 (1812); et ex Roem. & Schult. Syst. ii. (1817) 22. 417 = Boutelloua, Lag (Gramin.).
barbata, Roem. & Schult. l. c. ii 420 = B. barbata.
bromoides, Roem. & Schult. l. c = B. bromoides
ciliata, Willd. ex Beauv. Agrost. 41 = Chloris ciliata.
gracilis, Willd. ex Roem. & Schult. Syst. ii. 418 = B. oligostachya.
hirsuta, Roem & Schult l. c. 149 = B. hirsuta.
humilis, Willd. ex Beauv. Agrost. 41 = B. prostrata
juncifolia, Roem. & Schult. Syst ii 420 = B. juncifolia.
ovata, Roem. & Schult. l. c. = B. ovata.
procumbens, Roem. & Schult. l. c. 417 = B. tenuis.
prostrata, Roem & Schult. l. c. 419 = B. prostrata.
scorpioides, Roem. & Schult. l. c. 420 = B. scorpioides
simplex, Roem. & Schult l. c. 418 = B. simplex.
tenuis, Willd. ex Beauv. Agrost 41 = B. tenuis.

ACTINOCLADUS, E. Mey. Ind. Sem. Hort Regiom. (1846) = **Capnophyllum**, Gaertn. (Umbellif.).
cinerascens, E. Mey. l. c. = C. africanum.

ACTINOCYCLUS, Klotzsch, in Monatsb. Akad Berl (1857) 14 = Pyrola, Tourn. (Ericac.).
secundus, Klotzsch, l. c. = P. secunda.

ACTINODAPHNE, Nees, in Wall. Pl. As. Rar. ii 68 (1831). *LAURINEAE*, Benth. & Hook. f. in 160
IOZOSTE, Nees, l. c. 68 (1831).
IOZOSMENE, Lindl. Introd. Nat. Syst. ed. II. 202 (1836).
acuminata, *Meisn. in DC. Prod.* xv. 1 211.—Japon
angustifolia, Hook. f. & Thoms. ex Meisn. in DC. Prod. xv. 1. 218 = Hookeri.
angustifolia, *Nees, in Wall. Pl. As. Rar.* iii 34.—Ind. or. ; Malaya.
arcolata, *Blume, Mus. Bot. Lugd Bat.* i. 344.—Java.
Blumii, Nees, Syst. Laurin. 598 = Litsea ferruginea.
borneensis, *Meisn. in DC. Prod.* xv 1. 218.—Borneo.
Candolleana, Meisn. l. c. 219 = speciosa.
celebica, *Miq. Fl. Ind. Bat.* i. 972.—Ins. Celebes.
chinensis, *Nees, Syst. Laurin.* 600.—China.
cochinchinensis, *Meisn. in DC. Prod.* xv. 1. 216 — Cochinch.
concolor, Blume, Mus. Bot. Lugd. Bat. i. 345 = sclerophera.
concolor, *Nees, in Wall. Pl. As. Rar.* iii. 31.—Ind. or.
confertiflora, *Meisn. in DC. Prod.* xv 1. 216.—Reg. Himal.
corymbosa, *Blume, Mus. Bot. Lugd. Bat.* i. 343.—Sumatra.
elegans, *Thw. Enum. Pl. Zeyl.* 256.—Zeylan.
furfuracea, *Blume, Mus. Bot. Lugd. Bat.* i. 345.—Malaya.
glabra, *Blume, l. c.* i. 344.—Java.
glauca, *Nees, Syst. Laurin.* 599.—Zeylan.
gomerata, *Nees, l. c.* 597.—Java.
gracilis, *Miq. Fl. Ind. Bat. Suppl.* 368.—Sumatra.
heterophylla, Blume, Mus. Bot. Lugd. Bat. i. 342 = angustifolia.
hirsuta, *Blume, l. c.* 345.—Java.
Hookeri, *Meisn. in DC. Prod.* xv. 1. 218.—Ind. or.
javanica, *Miq. Fl. Ind. Bat.* i. i. 969.—Java.
lanata, *Meisn. in DC. Prod.* xv. 1. 219.—Ind. or.
lancifolia, *Meisn. l. c.* 211.—Japon.
macrophylla, *Nees, Syst. Laurin.* 598.—Java.
macroptera, *Miq. Fl. Ind. Bat.* i. i. 970.—Sumatra.

ACTINODAPHNE:—
mollis, *Blume, Mus. Bot. Lugd. Bat.* i. 343.—Sumatra.
molochina, *Nees, Syst. Laurin.* 603.—Zeylan.
moluccana, *Blume, Mus. Bot. Lugd. Bat.* i. 344.—Ins. Molucc.
Moonii, Thw. Enum. Pl. Zeyl. 256 = molochina.
multiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 220.—Ins. Fiji.
obovata, *Blume, Mus. Bot. Lugd. Bat.* i. 342.—Ind. or.
pauciflora, *Blume, l. c.*—Borneo.
pedunculata, *Meisn. in DC. Prod.* xv. 1. 211.—China
procera, *Nees, Syst. Laurin.* 605.—Java.
pruinosa, *Nees, in Wall. Pl. As. Rar.* ii. 68.—Ind. or
pubescens, *Blume, Mus. Bot. Lugd. Bat.* i. 345.—Malaya.
quercina, *Blume, l. c.* 344.—Java.
reticulata, *Meisn. in DC. Prod.* xvi. 1. 212.—Ind. or.
rufescens, *Blume, Mus. Bot. Lugd. Bat.* i. 343.—Malaya.
Rumphii, *Blume, l. c.* 344.—Ins. Molucc
salicina, *Meisn. in DC. Prod.* xv. 1. 212.—Ind. or.
sclerophera, *Miq. Fl Ind. Bat.* i. i. 972.—Malaya.
sesquipedalis, *Hook f. & Thoms. ex Meisn. in DC Prod.* xv. 1. 216.—Ind. or.
Sieboldiana, *Miq. in Zoll. Syst Verz Ind. Archip.* 114, 116.—Japon.
sikkimensis, *Meisn in DC. Prod.* xvi. 1. 218.—Reg Himal.
speciosa, *Nees, Syst. Laurin.* 602.—Zeylan.
sphaerocarpa, *Nees, l. c.* 605.—Java
stenophylla, *Thw Enum. Pl. Zeyl.* 256.—Zeylan.
Thwaitesii, Meisn. in DC. Prod. xv. 1. 212 = stenophylla.

ACTINODIUM, Schan. ex Schlecht. in Linnaea, x (1835) 311. *MYRTACEAE*, Benth. & Hook. f. i. 696
TRIPHELIA, R. Br. ex Endl. in Enum. Pl. Hueg. 48 (1837).
Cunninghami, *Schau. ex Lindl. Introd. Nat. Syst.* ed. II. 440.—Austral.
prolifferum, Turcz. in Bull. Soc. Nat Mosc. xxii. (1849) 11. 17 = Cunninghami.

ACTINOLEMA, Fenzl, Pugill Pl. Nov. Syr. 16. t. 12 (1842) = **Astrantia**, Linn (Umbellif.).
eryngioides, Fenzl, l c.—Oriens.
macrolema, Boiss. Fl. Orient. ii 831.—Oriens.

ACTINOLEPIS, DC. Prod. v. 655 (1836). *COMPOSITAE*, Benth. & Hook. f. ii 399.
HYMENOXYS, Torr. & Gray, Fl. N. Am. ii. 380 (1841).
anthemoides, *A. Gray, in Bot. Calif.* i. 378.—Calif.
coronaria, *A. Gray, in Proc. Am. Acad.* ix. (1874) 197.—Calif.
lanosa, *A. Gray, l. c.*—Utah.
Lemmoni, *A. Gray, l. c.* xvi. (1881) 101.—Calif.
multicaulis, *DC. Prod.* v. 656.—Calif.
mutica, *A Gray, in Proc. Am. Acad.* ix. (1874) 197.—Calif.
nivea, *A. Gray, l c*—Calif.
tenella, *A. Gray, l. c.*—Calif.
Wallacei, *A. Gray, l. c.*—Utah.

ACTINOMERIS, Nutt. Gen. Am. ii. 181 (1818). *COMPOSITAE*, Benth. & Hook. f. ii. 379.
ACTIMERIS, Rafin. in Am. Monthly Mag. (1819) 195.
PTEROPHYTON, Cass. in Bull. Soc. Philom. (1818) 76.
RIDAN, Adans. Fam. ii. 180 (1763).
alata, Nutt. Gen. Am. ii. 181 = Verbesina occidentalis.
alba, *Torr. & Gray, Fl. N. Am.* ii. 385.—Am. bor.
alternifolia, DC. Prod v. 572 = squarrosa.
Elliott, *Rafin. Neo Fl. Am.* i. 61.—Am. bor.
helianthoides, *Nutt. Gen. Am.* ii. 181.—Am. bor.
heterophylla, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 6.—Am. bor.
longifolia, *A. Gray, in Smithson. Contrib.* v. (1853); *Pl. Wright.* ii. 89.—Arizona.
nudicaulis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 384.—Florida.
oppositifolia, DC. Prod. v. 575 = helianthoides.
oppositifolia, Fresen. Ind. Sem. Hort. Francof. (1836) = squarrosa.
OVATA, *Nutt. Gen. Am.* ii. 181.—Mexico.

ACTINOMERIS :—
pauciflora, *Nutt. in Am. Journ. Sc.* v. (1822) 301.—Florida.
pedunculosa, DC. Prod. v. 576 = Verbesina Capitaneja.
procera, Steud. Nom. ed. II. i. 28 = squarrosa.
squarrosa, *Nutt. Gen. Am.* ii. 181.—Am. bor. occ.
stricta, *Hemsl. Biol. Centr. Am. Bot.* ii. 186.—Mexico.
tetragona, *DC. Prod.* v. 576.—Mexico.
tetraptera, *DC. l. c.* 575.—Mexico.
Wrightii, *A. Gray, in Mem. Am. Acad.* iv. (1849) (*Pl. Fendl.*) 85.—N. Mexico.

ACTINOMORPHE, Miq. Comm. Phyt. 102 (1840) = **Heptapleurum**, Gaertn. (Araliac.).
humilis, Miq. l. c. = H. parasiticum.

ACTINOPAPPUS, Hook. f. ex A. Gray, in Hook. Kew Journ. iv. (1852) 226 = **Enydroeis**, DC. (Compos.).
Drummondii, A. Gray, l. c. = R. Pumilio.
perpusillus, Hook. f. ex A. Gray, l. c. = R. Pumilio.

ACTINOPHORA, Wall Cat. n. 1163 (1832) = **Schoutenia**, Korth (Tiliac.).
fragrans, Wall. l. c. = S ovata.

ACTINOPHYLLUM, Ruiz & Pav. Prod. 51. t. 8 (1794) = **Sciadophyllum**, P. Br. (Araliac.).
acuminatum, Pav. in Mem. Acad. Med. Madr. i. (1797) 196 = S. acuminatum.
angulatum, H. B. & K. Nov. Gen. et Sp. v. 92 = S. Humboldtianum.
angulatum, Pav. in Mem. Acad. Med. Madr. i. (1797) 194 = S. angulatum.
conicum, Pav. l. c. 195 = S. conicum.
digitatum, Wall. ex Steud. Nom. ed. II. i. 23 = S. digitatum.
divaricatum, Blume, Cat. Gew. Buitenz. 42 = S. divaricatum.
farinosum, Blume, l. c. 48 = S. tomentosum.
pedicellatum Pav. in Mem. Acad. Med. Madr. i. (1797) 195 = S. pedicellatum.
pentandrum, Pav. l. c. 197 = S. pentandrum.

ACTINORHYTIS, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 184. *PALMAE*, Benth. & Hook. f. iii. 889.
calapparia, *H. Wendl. & Drude, l. c.*—Malaya.

ACTINOSCHOENUS, Benth. in Hook. Ic. Pl. xiv. 33. t. 1346 (1881). *CYPERACEAE*, Benth. & Hook. f. iii. 1058.
ARTHROSTYLIS, Boeck. in Linnaea, xxxvii. (1873) 593.
chinensis, *Benth. l. c.*—Hongkong.
filiformis, *Benth. l. c.*—Zeylan.
Thouarsii, *Benth. l. c.*—Madag.

ACTINOSPERMUM, Ell. Sketch, ii. 446 (1824) = **Baldwinia**, Nutt. (Compos.).
angustifolium, Torr. & Gray, Fl. N. Am. ii. 389 = B. multiflora.
multiflorum, Ell. Sketch, ii. 441 = B. multiflora.

ACTINOSPORA, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 (1835) = **Cimicifuga**, Linn. (Ranunc.).
dahurica, Turcz. ex Fisch. & Mey. l. c. = C. davurica.
frigida, Fisch. & Mey. l. c. = C. frigida.
pterosperma, Walp. Ken. i. 60 = C. davurica.
Wallichii, Meissn. Gen. Comm. 4 = C. foetida.

ACTINOSTEMA, Lindl. Veg. Kingd. 281 (1846) err. typ. = **Actinostemon**, Mart. (Euphorb.).

ACTINOSTEMMA, Griff. Pl. Cantor, 24. t. 8 (1837). *CUCURBITACEAE*, Benth. & Hook. f. i. 838.
MITROSICYOS, Maxim. in Ann. Sc. Nat. Sér. IV. xiii. (1860) 95.
POMASTERION, Miq. Ann. Mus. Bot. Lugd. Bat. ii. (1865) 80.
japonicum, Miq. l. c. iii. 128 = racemosum.
lobatum, *Maxim. in Franch. & Sav. Enum. Pl. Jap.* i. 175.—Japon.
paniculatum, *Maxim. ex Cogn. in DC. Monog. Phan.* iii. 930.—China.
racemosum, *Maxim. ex Cogn. l. c.* 922.—China.

ACTINOSTEMMA :—
tenerum, *Griff. Pl. Cantor*, 25. t. 8.—Ind. or.
tenerum, Naud. in Ann. Sc. Nat. Sér. V. (1866) 39 = lobatum.

ACTINOSTEMON, Mart. ex Klotzsch, in Wiegm. Archiv, vii. (1841) 184. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 333.
DACTYLOSTEMON, Klotzsch, l. c. 181.
acuminatus, Baill. Etud. Gén. Euphorb. 532. t. 5 = concolor.
angustifolius, Klotzsch, ex Regel, Gartenfl. (1859) 363 = concolor.
caribaeus, *Griseb. in Goett. Abh.* vii. (1857) 168.—Bras.; Ind. occ.
concolor, Muell. Arg. in DC. Prod. xv. ii. 1193 = caribaeus.
cuneatus, Baill. Adansonia, v. (1864–65) 536.—Bras.
echinatus, *Muell. Arg. in Linnaea*, xxxii. (1863) 107.—Bras.
furcatus, Klotzsch, ex Baill. Etud. Gén. Euph. 532 = caribaeus.
grandifolius, Baill. l. c. = caribaeus.
imbricatus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 216.—Bras.
Klotzschianus, Baill. Adansonia, v. (1864–65) 534 (= *Dactylostemon grandifolius*).
lanceolatus, *Saldanha, in Baill. l. c.* viii. (1867–68) 363.—Bras.
lasiocarpoides, *Baill. l. c.* v. (1864–65) 534.—Bras.
lasiocarpus, *Baill. l. c.*—Bras.
macrocarpus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 597.—Bras.
marginatus. Klotzsch, ex Baill. Etud. Gén. Euph. 533 = caribaeus.
multiflorus, *Muell. Arg. in Linnaea*, xxxii. (1863) 111.—Bras.
oligandrus, *Baill. Adansonia*, v. (1864–65) 535.—Bras.
polymorphus, Muell. Arg. in Linnaea, xxxii. (1863) 108 = caribaeus.
sessilifolius, Klotzsch, Ind. Sem. Hort. Berol. App. (1851) 13 = caribaeus.
Sprengelii, Baill. Etud. Gén. Euph. 533 (= *Dactylostemon Klotzschii*).
trachycarpus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 591.—Bras.
verticillatus, *Baill. Adansonia*, v. (1864–65) 534.—Bras.

ACTINOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) 259 = **Seringia**, J. Gay (Sterculiac.).
lanceolatum, Turcz. l. c.—Austral.

ACTINOSTIGMA, Wallr. Annot. 560 (1858) = **Symphonia**, Linn. f. (Gutlif.).
speciosum, Wallr. l. c. = S. globulifera.

ACTINOSTROBUS, Miq. in Lehm. Pl. Preiss. i. 644 (1845). *CONIFERAE*, Benth. & Hook. f. iii. 425.
acuminatus, *Parl. Ind. Sem. Hort. Florent.* (1862) 25.—Austral.
pyramidalis, *Miq. in Lehm. Pl. Preiss.* i. 644.—Austral.

ACTINOTUS, Labill. Pl. Nov. Holl. i. 67. t. 93 (1804). *UMBELLIFERAE*, Benth. & Hook. f. i. 880, 1008.
ERIOCALIA, Sm. Exot. Bot. t. 78, 79 (1806).
HEMIPHURIS, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 469.
HOLOTOMUS, Endl. Gen. 766 (1839).
bellidioides, *Benth. Fl. Austral.* iii. 369.—Austral.
Gibbonii, *F. Muell. Fragm.* vi. 29.—Austral.
glomeratus, *Benth. Fl. Austral.* iii. 369.—Austral.
Helianthi, *Labill. Pl. Nov. Holl.* i. 67. t. 92.—Austral.
leptophlebia, *Benth. in Enum. Pl. Hueg.* 56.—Austral.
minor, *DC. Prod.* iv. 83.—Austral.
Novae-Zealandiae, *Petrie, in Trans. N. Z. Inst.* xiii. (1880) 294.—N. Zel.
omnifertilis, *F. Muell. ex Benth Fl. Austral.* iii. 368.—Austral.
rhomboideus, *Benth. Fl Austral.* iii. 368.—Austral.

ACTIPSIS, Rafin. Fl. Tellur. ii. 43 (1836) = **Solidago**, Linn. (Compos.).
glomerata, Rafin. l. c. = S. glomerata.
squarrosa, Rafin. l. c. = pauc. !

ACTISPERMUM, Rafin. New Fl. Am. i. 61 (1836) (= ACTINOSPERMUM, Ell.) = **Baldwinia**, Nutt. (Compos.).

ACUAN, Medic. Theod. 62 (1786) = **Desmanthus**. Willd. (Legum.).
arborescens, Medic. l. c.—Hab. ?
virgatum, Medic. l. c. = D. virgatus.

ACUBA, Link, Enum. Hort. Berol. ii. 386 (1822), sphalm. = **Aucuba**, Thunb. (Cornac.).

ACUBALUS, Neck. Elem. ii. 167 (1790) = **Cucubalus**, Tourn. (Caryophyll.).

ACUNA, Endl. Gen. 759 (1839) = seq.

ACUNNA, Ruiz & Pav. Prod. 69. t. 12 (1794) = **Befaria**, Mutis (Ericac.).
lanceolata, Cav. ex DC. Prod. vii. 731 = B. coarctata.
lanceolata, Ruiz & Pav. Syst. Veg. 124 = B. aestuans.
oblonga, Ruiz & Pav. l. c. 123 = B. oblonga.

ACURA, Hill, Veg. Syst. v. 52 (1769) (Compos.).

ACUROA, J. F. Gmel. Syst. 1083 (1791) = ACOUROA, Aubl. = **Scostaphyllum**, P. Br. (Legum.).

ACUSTON, Rafin. Sylva Tellur. 131 (1838) = **Farsetia**, Desv. (Crucif.).
clypeatum, Rafin. l. c. 131 = F. clypeata.
lunaroides, Rafin. l. c. = F. lunarioides.

ACYNOS, Pers. Syn. ii. 131 (1807) (err. typ. '151'). = ACINOS, Rupp. = **Calamintha**, Moench (Labiat.).

ACYNTHA, Medic. Theodora, 76 (1786) = **Sanseviera**, Thunb. (Haemodor.).
guineensis, Medic. l. c. = S. guineensis.

ADA, Lindl. Fol. Orch. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 566.
aurantiaca, *Lindl. l. c.*—N. Granat.

ADAMANTOGETON, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 165 (1842) = **Lagenocarpus**, Nees, (Cyper.).

ADAMBEA, Lam. Encyc. 89 (1783) = **Lagerstroemia**, Linn. (Lythrar.).
glabra, Lam. l. c. = L. Flos-reginae.
hirsuta, Lam. l. c. = L. hirsuta.

ADAMBOE, Rafin. Fl. Tellur. iv. 73 (1836) = **Argyreia**, Lour. (Convolv.).
bicolor, Rafin. l. c. = A. tiliaefolia.

ADAMIA, Wall. Tent. Fl. Nep. 46. t. 36 (1824) = **Diohroa**, Lour. (Saxifrag.).
chinensis, Gardn. & Champ. in Hook. Kew Journ. i. 311 (1849) = D. febrifuga.
cyanea, Wall. Tent. Fl. Nep. 46. t. 36 = D. febrifuga.
sylvatica, Meissn. in Walp. Rep. ii. 877 = D. febrifuga.
vercicolor, Fortune, in Lindl. Journ. Hort. Soc. i. (1846) 296 = D. febrifuga.

ADAMSIA, Fisch. ex Steud. Nom. ed. I. 866, 867, in syn. (1821) = **Genm**, Linn. (Rosac.).
anemonoides, Fisch. ex Steud. l. c. 866 = G. anemonoides.
glacialis, Fisch. ex Steud. l. c. 866 = G. glaciale.
reptans, Fisch. ex Steud. l. c. 867 = G reptens.
rotundifolia, Fisch. ex Steud. l. c. = G. rotundifolium.

ADAMSIA, Willd. in Mag. Ges. Naturf. Fr. Berl. ii. (1808) 16 = **Pueshkinia**, Adams (Liliac.).
scilloides, Willd. l. c. = P. scilloides.

ADANSONIA, Linn. Sp. Pl. 1190 (1753). *MALVACEAE*, Benth. & Hook. f. i. 209.
BAOBAB, Adans. Fam. ii. 399 (1763).
OPHELUS, Lour. Fl. Cochinch. 413 (1790).
Baobab, Linn. Sp. Pl. ed. II. 1680 = digitata.
Bechel, Gaert. Fruct. ii. 323. t. 138 = digitata.

ADANSONIA:—
digitata, *Linn. Sp. Pl.* 1190.—Afr. trop.; Ind. or.
Gregorii, *F. Muell. in Hook. Kew Journ.* ix. (1857) 14.—Austral.
integrifolia, Rafin. Sylva Tellur. 149 = A. Situla.
madagascariensis, *Baill. Adansonia*, xi. (1873-76) 251.—Madag.
microphylla, A. Rich. Fl. Cub. Fanerog. ii. 325, lapsu = Adelia microphylla.
pedunculosa, A. Rich. l. c. lapsu = Adelia pedunculosa.
Scutula, Steud. Nom. ed. II. i. 24, sphalm. = seq.
Situla, *Spreng. Syst.* iii. 124.—Afr. or.

ADAPHUS, Neck. Elem. ii. 240 (1790) = **Laurus**, Tourn.

ADATODA, Rafin. Fl. Tellur. iv. 60 (1836) = **Justicia**, Linn. (Acanth.).
arborea, Rafin. l. c. 61 = J. Adhatoda.
Betonica, Rafin. l. c. = J. Betonica.
repens, Rafin. l. c.—Hab. ?

ADELANTHUS, Endl. Gen. 1327. n. 6889; Suppl. ii. 90 (1841) = **Pyrenacantha**, Wight (Olacin.).
grandiflorus, Harv. ex Baill. Adansonia, x. (1872) 271 = P. grandiflora.
scandens, Endl. ex Baill. in DC. Prod. xvii. 18 = P. scandens.

ADELASTER, Lindl. in Proc. Roy. Hort. Soc. i. (1861) 568; Lem. in Illustr. Hortic. t. 930 = **Brunthemum** Linn. ? (Acanthac.).
albivenis, Lindl. ll. cc.—Hab. ?

ADELBERTIA, Meisan. Gen. Pl. 114 (1837) = **Meriania**, Sw. (Melast.).
calophylla, Meisan. l. c. (= *Davya calophylla*).—Bras.

ADELIA, P. Br. Nat. Hist. Jamaic. 361 (1756) = **Forestiera**, Poir. (Oleac.).
acuminata, Michx. Fl. Bor. Am. ii. 225 = F. acuminata.
cassinioides, Steud. Nom. ed. II. i. 24 = F. cassinioides.
ligustrina, Michx. Fl. Bor. Am. ii. 224 = F. ligustrina.
ovata, [Desf.] Hort. Par. ex Pers. Syn. ii. 635.—Am. bor.
porulosa, Michx. Fl. Bor. Am. ii. 224 = F. porulosa.
virgata, Poir. Encyc. Suppl. i. 132 = Securinega buxifolia.

ADELIA, Linn. Syst. ed. X. (1759) 1293. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 312.
RICINELLA, Muell. Arg. in Linnaea, xxxiv. (1865) 158.
Acidoton, Blanco, Fl. Filip. ed. I. 815 = Sumbavia rottleroides.
Acidoton, *Linn. Syst.* ed. X. 1293; *Pl. Jam. Pugill.* 29.—Jamaica; Cuba.
anomala, Jus. ex Steud. Nom. ed. II. i. 24 = ovata.
anomala, Poir. Encyc. Suppl. i. 132 = Erythrococca aculeata.
barbata, Blanco, Fl. Filip. ed. II. 561 = Mallotus ricinoides.
barbinervis, *Cham. & Schlecht. in Linnaea*, vi. (1831) 362.—Mexico.
Bernardia, Blanco, Fl. Filip. ed. I. 814 = Mallotus ricinoides.
Bernardia, Linn. Syst. ed. X. 1295 = Bernardia dichotoma.
caperoniaefolia, Baill. Adansonia, iv. (1864) 376 = Bernardia caperoniaefolia.
castaneicarpa, Roxb. Fl. Ind. iii. 848 = Chaetocarpus castanocarpus.
celastrinea, Baill. Adansonia, iv. (1864) 375 = Bernardia axillaris.
condiifolia, *Roxb. Fl. Ind.* iii. 849.—Ins. Moluce.
cuneata, Wall. Cat. n. 7964 = Homonoia retusa.
dodecandra, Seme, ex Cav. in Anal. Cienc. Nat. v. (1802) 264 = Bernardia interrupta.
ferruginea, Poir. ex Baill. Etud. Gén. Euph. 418 = Bernardia leprosa.
glandulosa, Blanco, Fl. Filip. ed. I. 814 = Alchornea javanensis.
grevilli, Seliah. Prod. 288 = A. Acidoton.
hirsutissima, Baill. Adansonia, iv. (1864) 372 = Bernardia pedunculosa.
Hookeliana, Baill. l. c. 378 = Bernardia axillaris.
javanica, *Mig. Fl. Ind. Bat.* i. ii. 568.—Java.

ADELIA:—
Martii, Spreng. Syst. iii. 147 = Bernardia axillaris.
microphylla, *A. Rich. Fl. Cub. Fanerog.* ii. 309.—Cuba.
monoica, Blanco, Fl. Filip. ed. II. 561 = Mallotus moluccanus.
myrtifolia, *Vent. ex Spreng. Syst.* iii. 147.—Hab. ?
nervifolia, Heyne, ex Roth, Nov. Pl. Sp. 375 = Homonoia riparia.
OSBECKIANA, *Hemsl. Biol. Centr. Am. Bot.* iii. 129.—Mexico.
papillaris, Blanco, Fl. Filip. ed. II. 562 = Mallotus Zollingeri.
pedunculosa, *A. Rich. Fl. Cub. Fanerog.* ii. 310.—Cuba.
pulchella, Baill. Adansonia, iv. (1864) 374 = Bernardia pulchella.
rennoas, Blanco, Fl. Filip. ed. II. 562 = Claoxylon Wallichianum.
retusa, Wight, Ic. t. 1869 = Homonoia retusa.
Ricinella, *Linn. Syst.* ed. X. 1298; *Pl. Jam. Pugill.* 29.—Ind. occ.
scabrida, Baill. Adansonia, iv. (1864) 378 = Bernardia axillaris.
scandens, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 350, nomen.—Malaya.
spartioides, Baill. Adansonia, v. (1866) 319 = Bernardia spartioides.
sylvestris, *Griseb. in Goett. Nachr.* (1865) 174.—Cuba.
tamandhana, Baill. Adansonia, iv. (1864) 374 = Bernardia tamandhana.
timoriana, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 350, nomen.—Ins. Timor.
trilobata, *Hemsl. Biol. Centr. Am. Bot.* iii. 130.—Panama.

ADELIOIDES, N. Br. ex Benth. Fl. Austral. i. 59 (1868) = seq.

ADELIOPSIS, Benth. in Benth. & Hook. f. Gen. i. 1. Add. 486 (1862). *MENISPERMACEAE*, Benth. & Hook. f. Gen. i. 1. Add. 486; i. 964.
ADELIOIDES, R. Br. ex Benth. Fl. Austral. i. 59 (1868).
decumbens, Benth. l. c.; et Fl. Austral. i. 59.—Austral.

ADELMANNIA, Reichb. Consp. 110 (1828) = **Borrichia**, Adans. (Compos.).

ADELOBOTRYS, DC. Prod. iii. 127 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 750.
SARMENTARIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 140.
SPHAXELLOPSIS, Steud. ex Naud. l. c. 137.
XERACINA, Rafin. Sylva Tellur. 98 (1838).
adscendens, *Triana, in Journ. Bot.* v. (1867) 210.—Am. trop.
barbata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 66.—Bras.
ciliata, *Triana, l. c.* 68.—Guiana.
fuscescens, *Triana, l. c.*—N. Granat.
laxiflora, *Triana, l. c.*—Guiana.
Lindeni, *Naud. in Belg. Hortic.* xv. (1865) 98.—Bras.
roraimifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 68.—Bras.
scandens, *DC. Prod.* iii. 127.—Guiana.
scandens, Macfad. Fl. Jamaic. ii. 99 = adscendens.

ADELODA, Rafin. Fl. Tellur. iv. 61 (1836) = **Justicia**, Linn. (Acanthac.).
integra, Rafin. l. c.—Hab. ?
serrata, Rafin. l. c. = J. serrata.

ADELONEMA, Schott, Prod. Aroid. 316 (1860) = **Homalonema** Schott (Aroid.).
erythropus, Schott, l. c. 317.—Bras.

ADELONENGA, Beca. in Ann. Jard. Buitenz. ii. (1885) 82. *PALMAE*, Benth. & Hook. f. iii. 885.
Geelvinkiana, *Becc. l. c.*—Malaya.
variabilis, *Becc. l. c.*—Malaya.

ADELOSA, Blume, Mus. Bot. Lugd. Bat. i. 176 (1850). *VERBENACEAE*, Benth. & Hook. f. ii. 1158.
microphylla, *Blume, l. c.*—Madag.

ADELOSTEMMA, Hook. f. in Hook. Ic. Pl. xv. (1883) 22. *ASCLEPIADEAE*.
gracillimum, *Hook. f. l. c.*—Burma.

ADELOSTIGMA, Steetz, in Peters, Reise Mossamb. Bot. 438 (1863). *COMPOSITAE*, Benth. & Hook. f. ii. 385.
athrixioides, *Steetz, l. c.* 430.—Afr. trop.
senegalensis, *Benth. in Hook. Ic. Pl.* t. 1144.—Afr. trop.

ADENACANTHUS, Nees, in Wall. Fl. As. Rar. iii. 75 & 126 (1832) = **Strobilanthes**, Blume (Acanthac.).
acuminatus, Nees, l. c. iii. 84 = S. acuminatus.
latifolia, Nees, in DC. Prod. xi. 197.—Hab. ?

ADENACHAENA, DC. Prod. vi. 49 (1837) = **Phymaspermum**, Less. (Compos.).
leptophylla, DC. l. c. = P. leptophyllum.
parvifolia, DC. l. c. = P. parvifolium.
pubescens, DC. l. c. 49.—Afr. austr.

ADENANDRA, Willd. Enum. Hort. Berol. 256. (1909). *RUTACEAE*, Benth. & Hook. f. i. 390.
GLANDULIFOLIA, Wendl. Collect. t. 83, 87 (1808).
HAENKEAEA, Usteri, Ann. Bot. vi. (1795) 117.
HAENKEA, F. W. Schmidt, Neue u. Seltne Pfl. 19 (1793); nomen prius.
OCENIA, F. G. Dietr. Lex. Gaert. Nachtr. v. 307 (1819).
OCKRA, F. G. Dietr. l. c. i. 195 (1815).
OCKENIA, Steud. Nom. ed. II. 24. (1840).
OKRA, Stend. l. c. ed. I. 562 (1821).
acuminata, Sweet, Hort. Brit. ed. I. 18 = amoena.
alba, *Korn. & Schult. Syst.* v. 451.—Afr. austr.
alternifolia, *Cham. & Schlecht. in Linnaea*, v. (1830) 52.—Afr. austr.
amoena, *Bartl. & Wendl. f. Diosm.* 80.—Afr. austr.
amoena, Link, Enum. Hort. Berol. i. 289 = fragrans.
Bartlingiana, Eckl. & Zeyh. Enum. 101 = viscida.
biseriata, *E. Mey. ex Bartl. & Wendl. f. Diosm.* 75.—Afr. austr.
brachypylla, *Schlecht. in Linnaea*, vi. (1831) 199.—Afr. austr.
calycina, *Sond. Nom.* ed. II. i. 24.—Afr. austr.
ciliata, *Sond. in Harv. & Sond. Fl. Cap.* i. 288.—Afr. austr.
cistoides, Eckl. & Zeyh. Enum. 100 = biseriata.
cordata, Link, Enum. Hort. Berol. i. 289 = Barosma serratifolia.
coriacea, Eckl. & Zeyh. Enum. 100 ex Sond. in Harv. & Sond. Fl. Cap. i. 289 = macrosepala.
coriacea, Bartl. & Wendl. f. Diosm. 75 = biseriata.
coriacea, *Lichtst. in Roem. & Schult. Syst.* v. 452.—Afr. austr.
crenata, Willd. ex Bartl. & Wendl. f. Diosm. 82 = umbellata.
cuspidata, Eckl. & Zeyh. Enum. 101 ex Sond. in Harv. & Sond. Fl. Cap. i. 285 = obtusata.
cuspidata, *E. Mey. ex Bartl. & Wendl. f. Diosm.* 87.—Afr. austr.
fragrans, *A. Gonsalt, in Rev. Hortic.* Sér. IV. i. (1852) 231.—Afr. austr.
fragrans, Eckl. & Zeyh. Enum. 99 = serpyllacea.
fragrans, *Roem. & Schult. Syst.* v. 451.—Afr. austr.
glandulosa, Lichtst. ex Roem. & Schult. Syst. v. 450 = umbellata.
gracilis, *Eckl. & Zeyh. Enum.* 101.—Afr. austr.
humilis, *Eckl. & Zeyh. l. c.* 99.—Afr. austr.
intermedia, Eckl. & Zeyh. l. c. 99 = marginata.
Kraussii, *Meisn. in Flora*, xxvii. (1844) 301.—Afr. austr.
laxiantha, *Sond. in Harv. & Sond. Fl. Cap.* i. 290.—Afr. austr.
linearis, A. Juss. in Mém. Mus. Par. xii. (1825) 471 = uniflora.
linifolia, *Bartl. in Linnaea*, xvii. (1843) 359.—Afr. austr.
macradenia, *Sond. in Harv. & Sond. Fl. Cap.* i. 289.—Afr. austr.
marginata, Eckl. & Zeyh. Enum. 99 = linifolia.
marginata, *Roem. & Schult. Syst.* v. 452.—Afr. austr.
mundiaefolia, *Eckl. & Zeyh. Enum.* 99.—Afr. austr.
muraltioides, *Eckl. & Zeyh. l. c.* 106.—Afr. austr.
obtusata, *Sond. in Harv. & Sond. Fl. Cap.* i. 285.—Afr. austr.
ovata, Link, Enum. Hort. Berol. i. 289 = amoena.

ADENANDRA:—
pubescens, *Send. in Harv. & Sond. Fl. Cap.* i. 385.—Afr. austr.
rosmarinifolia, A. Juss. in Mém. Mus. Par. xh. (1825) 471 = marginata.
rotundifolia, *Eckl. & Zeyh. Enum.* 101.—Afr. austr.
rugosa, Otto, ex Steud. Nom. ed. II. i. 24 = umbellata. Schmidtiana, *Steud. l c.*—Afr. austr.
serpyllacea, *Bartl. in Linnœa*, xvii. (1843) 859.—Afr. austr
serratifolia, Link, Enum. Hort. Berol. i. 239 = Barosma serratifolia.
speciosa, Link, l. c. = umbellata.
tetragona, Sweet, Hort. Suburb. 45 = Acmadenia tetragona.
trigona, *Eckl. & Zeyh. Enum.* 100.—Afr. austr.
umbellata, *Willd. Enum Hort. Berol.* 267.—Afr. austr.
uniflora, E. Mey. & Drège, in Harv. & Sond. Fl. Cap. i. 391 = serpyllacea.
uniflora, *Willd. Enum. Hort. Berol.* 256.—Afr. austr.
villosa, Eckl & Zeyh Enum p. 101 = pubescens.
villosa, Hort. ex Harv. & Sond. Fl. Cap. i. 388 = uniflora.
villosa, Lichtst. ex Roem. & Schult. Syst. v. 452 = cuspidata.
viscida, *Eckl. & Zeyh. Enum.* 101.—Afr. austr.
Wendlandiana, Eckl. & Zeyh. Enum. 100 = umbellata.

ADENANTHERA, Royen, ex Linn. Coroll. Gen. 7 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 590.
GONSII, Adans. Fam. ii. 318 (1763).
PONGELION, Adans. Fam. ii. 319 (1763).
abrosperma, *F. Muell. Progn.* v. 30.—Austral.
aculeata, Roxb. Hort. Beng. 90; Fl. Ind. ii. 37 = Prosopis spicigera.
bicolor, *Mem. Cat. Pl. Ceyl.* 84, *ex Thw. Enum. Pl. Zeyl* 98.—Zeylan.; Malaya.
Bonplandiana, H. B. & K. Nov. Gen. et Sp. vi. 311 = Entada polystachya.
chrysostachya, Benth. in Hook. Lond. Journ. Bot. iv. (1852) 348 = Piptadenia chrysostachya.
circinalis, DC. Prod. ii. 446 = Pithecolobium clypearia.
falcata, *Linn. Diss. Herb. Amb.* 14 (*Amoen. Acad.* iv. 124).—Ins. Molucc.
Falcataria, Linn. Sp. Pl. ed. II. 550 = falcata.
Gerrardi, *Schæff. in Flora*, lii. (1869) 307.—Malaya.
glauca, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 347, nomen.—Malaya.
Gogo, Blanco, Fl. Filip. ed. I. 358 = Entada scandens.
microsperma, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* xxvii. (1864) 58.—Java.
oudhensis, J. L. Stewart, ex Brand. For. Fl. Ind. 168 = Piptadenia oudhensis.
pavonina, *Linn. Sp. Pl.* 384.—As. trop.; Malaya.
polita, *Miq. Fl. Ind. Bat.* i. 47.—Malaya.
scandens, Forst. f. Prod. 33 = Entada scandens.
scandens, Vieill in Bull. Soc. Linn. Normand. ix. (1865) 240 = Entada Adenanthera.
tetraptera, Schum. & Thonn. Beskr. Guin. Pl. 213 = Tetrapleura Thonningii.
triphysa, Dennst. Schlüssn. Hort. Malab. 33 = Ailanthus malabarica.

ADENANTHES, Knight, Prot. 98. (1809) = seq.

ADENANTHOS, Labill. Pl. Nov. Holl. i. 28. t. 36 (1804). *PROTEACEAE*, Benth. & Hook. f. iii. 178.
apiculata, *R. Br. Prod. Suppl* 9.—Austral.
armata, Meissn. in DC. Prod. xiv. 313 = pungens.
barbigera, *Lindl. Swan Riv. App.* 36. n. 182.—Austral.
crenata, Willd. ex Spreng. Syst. i. 479 = cuneata.
cuneata, *Labill. Pl. Nov. Holl.* i. 28. t. 36.—Austral.
Cunninghamii, *Meissn. in Lehm. Pl. Preiss.* i. 513.—Austral.
Detmoldi, *F. Muell. Progn.* viii. 149.—Austral.
Dubosci, *F. Muell. l. c.* vi. 204.—Austral.
Drummondii, Meissn. in Lehm. Pl. Preiss. i. 514 = apiculata.
filifolia, *Benth. Fl. Austral.* v. 356.—Austral.
flabellifolia, (Salisb. in) Knight, Prot. 96 = cuneata.
flavidiflora, *F. Muell. Fragm.* i. 157.—Austral.
Forrestii, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 230.—Austral.
linearis, *Meissn. in DC. Prod.* xiv. 311.—Austral.

ADENANTHOS:—
Meisneri, *Lehm. Pl. Preiss.* i. 512; ii. 248.—Austral.
obovata, *Labill. Pl. Nov. Holl.* i. 29. t. 37.—Austral.
procumbens, Meissn. in Lehm. Pl. Preiss. i. 512; ii. 248 = apiculata.
pungens, *Meissn. l. c.* i. 515.—Austral.
sericea, *Labill. Pl. Nov. Holl.* i. 28. t. 38.—Austral.
terminalis, *R. Br. in Trans. Linn. Soc.* x. (1810) 152.—Austral.
velutina, *Meissn. in DC. Prod.* xiv. 312.—Austral.
venosa, *Meissn. l. c.*—Austral.

ADENANTHUS, Roem. & Schult. Syst. iii. 22, 397 (1818) = prasc.

ADENARIA, H. B. & K. Nov. Gen. et Sp. vi. 185. t. 549 (1823). *LYTHRARIEAE*, Benth. & Hook. f. i. 777.
floribunda, *H. B. & K. l. c.* 188. t. 549.—N. Granat.
grisseiodes, H. B. & K. l. c. 186 = floribunda.
lanceolata, *Beurling, in Vet. Akad. Handl. Stockh.* (1854) 124.—Panama.
parvifolia, Hook. Ic. Pl. t. 166 = floribunda.
purpurata, H. B. & K. Nov. Gen. et Sp. vi. 185 = floribunda.

ADENARIUM, Rafin. in Am. Monthly Mag. (1818) 266 = **Armeria**, Linn. (Caryophyl.).
marinum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 545 = Aren. peploides.
maritimum, Rafin. New Fl. Am. i. 62 = Aren. peploides.
peploides, Rafin. in Am. Monthly Mag. (1818) 266 = Aren. peploides.

ADENELBUTREOPHORA, Rodrig. Orch. Nov. ii. 170 (1881). *ORCHIDEAE*.
graminifolia, *Rodrig. l. c.* 171.—Bras.

ADENEMA, G. Don, Gen. Syst. iv. 201 (1837) = **Enicostema**, Blume (Gentian.).
hyssopifolium, G. Don, l. c. = E. littorale.

ADENESMA, Griseb. in DC. Prod. ix. 65 (1845) = prasc.

ADENIA, Forsk. Fl. Aegypt. Arab. 77 (1775) = **Modecca**, Linn. (Passif.).
venenata, Forsk. l. c.—Arabia.

ADENIA, Torr. Fl. New York, ii. t. 122 (1843) = **Pilea**, Lindl. (Urtic.).
pumila, Torr. l. c. = P. pumila.

ADENILEMA, Reichb. Nom. 159 (1841) = seq.

ADENILEMA, Blume, Bijdr. 1120 (1826) = **Notilia**, D. Don (Rosac.).
fallax, Blume, l. c. 1121 = N. thyrsiflora.

ADENILEMMA, Hassk. Cat. Hort. Bogor. Alt. 170 (1844) = prasc.

ADENIUM, Roem. & Schult. Syst. iv. 85 & 411 (1819). *APOCYNACEAE*, Benth. & Hook. f. ii. 722.
arboreum, *Ehrenb. Symb. Phys.* 4.—Afr. trop.
Honghel, *A. DC. in DC. Prod.* viii. 412.—Afr. trop.
Honghel, Lindl. Bot. Reg. xxxii. t. 54 = obesum.
multiflorum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 279.—Afr. trop.
namaquanum, Wyley, ex Harv. Thes. Cap. ii. 11. t. 117 = Pachypodium namaquanum.
obesum, *Roem. & Schult. Syst.* iv. 411.—Arab.; Afr. trop.
speciosum, *Fenzl, in Sitzb. Acad. Wien. Math. Nat.* li. *Abth.* ii. (1865) 140.—Afr. trop.

ADENOBASIUM, Presl, Symb. Bot. i. 29. t. 27 (1830) = **Sloanea**, Linn. (Tiliac.).
obtusifolium, Moric. Pl. Nouv. 52. t. 55 = S. obtusa.
salicifolium, Presl, Symb. Bot. i. 40. t. 27 = S. monosperma.

ADENOCALYMMA, Endl. Ench. 347 (1841) = seq.

ADENOCALYMNA, Mart. in Meisn. Gen. 300 (308) et in DC. Prod. ix. 199 (1840). *BIGNONIACEAE*, Benth & Hook. f. ii. 1086.
LEGUMINARIA, Bar. Monog. Bignon. 43 (1864).

ADENOCALYMNA:—
MEMORA, Miers. in Proc. Roy. Hort. Soc. iii. (1863) 165.
PACHYPTERA, DC. Prod. ix. (1840) 175.
acutissimum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 395.—Bras.
alliaceum, Miers, l. c. 394 = Bignonia alliacea.
brachybotrys, *DC. Prod.* ix. 203.—Guiana.
bractestum, *DC. l. c.* 200.—Bras.
bracteolatum, *DC. l. c.* 200.—Bolivia.
divaricatum, *Miers. in Ann. & Mag. Nat. Hist.* Ser. III vii. (1861) 390.—Bras.
comosum, *DC. Prod.* ix. 201.—Bras.
coriaceum, *DC. l. c.* 202.—Bras.
flavidum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III vii. (1861) 389.—Bras.
flavum, *Mart. ex DC. Prod.* ix. 202.—Bras.
floribundum, *DC. Prod.* ix. 201.—Bolivia.
grandifolium, *Mart. ex DC. l. c.* 199.—Bras.
Guillemini, *DC. l. c.* 202.—Bras.
inundatum, *Mart ex DC. l. c.* 201.—Bras.
japurense, Mart. ex DC. l. c. 203.—Bras.
laevigatum, Mart. ex DC. l. c. 200 = marginatum.
lanceolatum, *Miers, in Ann. & Mag. Nat. Hist.* Ser III. vii. (1861) 390.—Bras.
longeracemosum, *Mart. ex DC. Prod.* ix. 201.—Bras.
macranthum, Mart. ex DC. l. c. 208.—Bras.
macrophyllum, *DC. l. c.* 199.—Bras
macrostachyum, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 180.—Bras.
magnificum, *Mart ex DC. Prod.* ix. 202.—Bras.
marginatum, *DC. l c* 200.—Bras.
nitidum, *Mart. ex DC. l. c.*—Bras.
prasinum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 395.—Ins. Trop.
paniculatum, *Benth. ex Miers, l. c.* 393.—Bras.
plicatiflorum, *Mart. ex DC. Prod.* ix. 202.—Bras.
prehensile, *Miers, in Ann. & Mag. Nat. Hist.* Ser III. vii. (1861) 388.—Bras.
Salzmanni, *DC. Prod.* ix. 200.—Bras.
scabrusculum, *Mart. ex DC. l. c.* 201.—Bras.
scansile, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 388.—Bras.
separium, Mart. ex DC. Prod. ix. 200 = nitidum.
stridula, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 392.—Bras.
subsessiliflorum, *DC. Prod.* ix. 199.—Bras.
vernuciferum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 393.—Bras.

ADENOCALYX, Bert. ex H. B. & K. Nov. Gen. et Sp vi. 330 (1823) = COULTERIA, H. B. & K. = **Cassia-pinia**, Linn. (Legum.).
racemosus, Bert. ex H. B. & K. l. c. = C. mollis.
remotus, Steud. Nom. ed. II. i. 428, sphalm. = prasc.

ADENOCARPUM, D. Don, ex Hook. & Arn. in Hook. Lond. Journ. Bot. iii. (1841) 318 = ADENOSPERMUM, Hook. & Arn. = **Chrysanthallium**, Rich. (Compos.).
tuberculatum, D. Don, ex Hook. & Arn. l. c. = C. indicum.

ADENOCARPUS, DC. Fl. Fr. Suppl. 549 (1815). *LEGUMINOSAE*, Benth. & Hook. f. i. 481.
anagyrifolius, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 246; xxii. (1875) 57, nomen.—Marocc.
anagyrus, *Spreng. Syst.* iii. 226.—Ins. Canar.
anisochilus, *Boiss. Diagn.* Ser. II. ii. 5.—Hispan.
benguelensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 47.—Afr. trop.
Bivonii, *Presl, Fl. Sic.* i. p. xix. = intermedius.
Boissieri, Webb, Iter Hispan. 53 = decorticans.
cebennensis, Delile, Ind. Sem. Hort. Monsp. (1838) 1 = intermedius.
commutatus, Gus. Fl. Sic. Prod. ii. 875 = intermedius.
complicatus, J. Gay, ex Gren. & Godr. Fl. Fr. i. 364 = intermedius.
decorticans, *Boiss. in Bibl. Univ. Genèv.* (Feb. 1836). —Hispan.
divaricatus, Sweet, Hort. Brit. ed. I. 113 = intermedius.
foliolosus, *DC. Fl. Fr. Suppl.* 549.—Ins. Canar.
frankenioides, Choisy, in DC. Prod. ii. 256 = anagyrus.
graecus, Griseb. Spicil. Fl. Rumel. i. 30 = intermedius.
grandiflorus, *Boiss. in Bibl. Univ. Genèv.* (Feb. 1836). —Gallia.
hispanicus, *DC. Fl. Fr. Suppl.* 549.—Hispan.
intermedius, *DC. l. c.*—Europ.; Orient.

ADENOCARPUS :—
Mannli, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 189. —Afr. trop.
parviflorus, DC. Fl. Fr. Suppl. 549 = intermedius.
speciosus, *Pomel, Nouv. Mat. Fl. Atl.* 178 —Afr. bor.
telonensis, DC. Fl. Fr. Suppl. 549 = grandiflorus ?
umbellatus, *Cav. & Dur. in Ann. Sc. Nat.* Sér. IV. i. (1854) 223, nomen.—Afr. bor.
villosus, Boiss. Diagn. Ser I. ii. 14 = intermedius.
viscosus, Webb & Berth. Phyt. Canar. ii. 32 = anagyrus.
wolgensis, Spreng. Syst. iii. 226 = Calophaca wolgarica.

ADENOCAULON, Hook. Bot. Misc. i. 19. t. 15 (1830). *COMPOSITAE*, Benth. & Hook. f. ii. 344.
adhaerescens, Maxim. Prim. Fl. Amur. 152 = bicolor.
bicolor, *Hook. Bot. Misc.* i. (1830) 19. t. 15.—Am. bor.; Himal.; Am. bor.
chilense, *Less. in Linnaea*, vi. (1831) 107.—Chili.
himalaicum, Edgew. in Trans. Linn. Soc. xx. (1851) 64 = bicolor.
integrifolium, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 289 = bicolor.
Lechleri, *Sch. Bip. in Flora*, xxxviii. (1855) 118.—Reg. Magell.

ADENOCHAENA, DC. Prod. vi. 680 (1837) = ADENACANTHA, DC. = **Phymosperrnum**, Less. (Compos.).

ADENOCHETON, Fenzl, in Flora, xxvii. (1844) 312 = **Cocculus**, Linn. (Menisperm.).
phyllanthoides, Fenzl, l. c. = C. Leseba.

ADENOCHETUS, Baill. Etud. Gén. Euph. 658 (1858) = praec.

ADENOCHILUS, Hook. f. Fl. N. Zel. i. 246. t. 56 (1858). *ORCHIDEAE*, Benth. & Hook f. iii. 618.
gracilis, *Hook. f. l. c.*—N. Zel.
Nortoni, *Pfstg. Austral. Orch.* i. ii. 2.—Austral.

ADENOCHLAENA, Boiss. ex Baill. Etud. Gén. Euph. 472 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 307.
CENTROSTYLIS, Baill. Etud. Gén. Euph. 469 (1858).
SYMPHYLLIA, Baill. l. c. 472. t. 11 (1858).
SYMPHYLLIUM, Benth. in Journ. Linn. Soc. xvii. (1878) 228.
calycina, *Bedd. For. Man. Bot.* 207.—Ind. or.
indica, *Bedd. Fl. Sylv.* ii. t. 261.—Ind. or.
leucocephala, Baill. Etud. Gén. Euph. 473 = Cephalocroton leucocephalus.
silhetensis, *Benth. in Journ. Linn. Soc.* xvii. (1880) 228.—Ind. or.
zeylanica, *Thw. Enum. Pl. Zeyl.* 270.—Zeylan.

ADENOCLINE, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 59. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 310.
DIPLOSTYLIS, Sond. in Linnaea, xxiii. (1850) 118.
PARAUENOCLINE, Muell. Arg. in DC. Prod. xv. ii. 1141 (1866).
acuta, *Baill. Etud. Gén. Euph.* 457.—Afr. austr.
humilis, *Turcz. in Bull. Soc. Nat. Mosc.* xvi. (1843) 61.—Afr. austr.
Mercurialis, *Turcz. l. c.* 60.—Afr. austr.
ovalifolia, *Turcz. l. c.*—Afr. austr.
pauciflora, *Turcz. l. c.* 61.—Afr. austr.
serrata, *Turcz. l. c.* xxv. (1852) ii. 180.—Afr. austr.
sessiliflora, *Baill. Etud. Gén. Euph.* 457.—Afr. austr.
sessilifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xvi. (1843) 61.—Afr. austr.

ADENOCREPIS, Blume, Bijdr. 579 (1825) = **Baccaurea**, Lour. (Euphorb.).
javanica, Blume, l. c. = B. javanica.
lanceolata, Muell. Arg. in Linnaea, xxxii. (1863) 82 = B. lanceolata.
tetrandra, Baill. Etud. Gén. Euph. 601 = B. tetrandra.

ADENOCHAENA, Steud. Nom. ed. II. i. 25 (1840) sphalm. = **Adenachaena**, DC.

ADENOCYCLUS, Less. in Linnaea, iv. (1829) 387 = **Oliganthes**, Cass. (Compos.).
condensatus, Less. l. c. = O. condensata.

ADENODISCUS, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) ii. 504 = **Balotia**, A. Rich. (Tiliac.).
mexicanus, Turcz. l. c. = B. grewiaefolia.

ADENODUS, Lour. Fl. Cochinch. 294 (1790) = **Blaeocarpus**, Linn. (Tiliac.).
sylvestris, Lour. l. c. = E. sylvestris.

ADENOGRAMMA, Reichb. Iconogr. Exot. ii. 3. t. 109 (1827). *FICOIDEAE*, Benth. & Hook. f. i. 858.
STRUDELIA, Presl, Steud. Nov. Pl. Gen. (1829); et Symb. i. 3. t. 2 (1830).
diffusa, *Fenzl, in Ann. Wien. Mus.* ii. (1840) 275 (err. typ. 255).—Afr. austr.
galioides, *Fenzl, l. c.* 277.—Afr. austr.
lampocarpa, *E. Mey. ex Fenzl, in Ann. Wien. Mus.* ii. (1840) 276.—Afr. austr.
Mollugo, *Reichb. Iconogr. Exot.* ii. 3. t. 109.—Afr. austr.
oppositifolia, *Hassk in Flora*, xxxi. (1851) 754.—Java.
physocalyx, *Fenzl, in Ann. Wien. Mus.* ii. (1840) 276. —Afr. austr.
rigida, *Sond. in Harv. & Sond. Fl. Cap.* 149.—Afr. austr.
sylvatica, *Fenzl, in Ann. Wien. Mus.* ii. (1840) 275 (err. typ. 255).—Afr. austr.

ADENOGYNA, *Rafin. New Fl. Am.* i. 62 (1836) = **Saxifraga**, Linn.
sarmentosa, Rafin. l. c. 63 = S. sarmentosa.

ADENOGYNE, Klotzsch, in Wiegm. Archiv, vii. (1841) 138 = **Sebastiania**, Spreng. (Euphorb.).
angustifolia, Klotzsch, l. c. 184 = S. angustifolia.
brachyclada, Klotzsch, l. c. = S. brachyclada.
discolor, Klotzsch, l. c. = Gymnanthes marginata.
guianensis, Klotzsch, in Schomb. Fauna & Fl Guy. 1185 = S. guyanensis.
marginata, Klotzsch, in Wiegm. Archiv, vii. (1841) 184 = Gymnanthes marginata.
mucronata, Klotzsch, l. c. = S. Schottiana.
pachystachya, Klotzsch, l. c. = S. pachystachya.
retundifolia, Klotzsch, l. c. = S. brevifolia.
serrata, Klotzsch, l. c. = S. serrata.

ADENOGYNE, Rafin. Fl. Tellur. ii. 67 (1836) = **Saxifraga**, Linn.

ADENOGYNUM, Reichb. f. & Zoll. in Verh. Natuurk. Ver. Batavisch. Indie, i. (1856) 33; et in Linnaea, xxviii. (1856) 925 = **Chloradenia**, Baill. (Euphorb.) 312.
discolor, Reichb. f. & Zoll. ll. cc. = C. discolor.
odontophyllum, Miq. Fl. Ind. Bat. i. ii. 400.—Java.

ADENOGYRUS, Klotzsch, in App. Ind. Sem. Hort. Berol. (1854) 1; et in Walp. Ann. iv. 226 (1857) = **Scoliopia**, Schreb. (Bixin.).
Braunii, Klotzsch, ll. cc.—Hab. ?

ADENOLEPIS, Less. in Linnaea, vi. (1831) 510 = **Cosmos**, Cav. (Compos.).
calva, Sch. Bip. in Miq. Fl. Ned. Ind. ii. 79 = Cosmos sulphureus.
pulchella, Less. in Linnaea, vi. 510 (1831).—Ins. Sandvic.

ADENOLINUM, Reichb. Handb. (1837) 306 = **Linum**, Linn.
alpicola, Reichb. l. c. 307 = L. alpinum.
alpinum, Reichb. l. c.; et Ic. Fl. Germ. xvi. 65 = L. alpinum.
angustifolium, Reichb. l. c. 307 = L. angustifolium.
austriacum, Reichb. l. c. 307; et Ic. Fl. Germ. xvi. 66 = L. austriacum.
barbatulum, Reichb. Ic. Fl. Germ. xvi. 78 = L. austriacum.
barbulatum, Schur, Enum. Pl. Transs. 127 = L. barbulatum.
Leonii, Reichb. Ic. Fl. Germ. xvi. 65 = L. Leonii.
marginatum, Reichb. Handb. 307; et Ic. Fl. Germ. xvi. 78 = L. austriacum.
montanum, Reichb. Ic. Fl. Germ. xvi. 65 = L. montanum.
nervosum, Reichb. Handb. 307 = L. nervosum.
pallescens, Reichb. l. c. = L. pallescens.

ADENOLINUM :—
perenne, Reichb. Ic. Fl. Germ. xvi. 65 = L. perenne.
provinciale, Fourr. in Ann. Soc. Linn. Lyon, N.S. xvi. (1868) 250 = L. alpinum.
pyrenaicum, Reichb. Handb. 307 = L. alpinum.
sibiricum, Reichb. l. c. 307 = L. sibiricum.
squamulosum, Reichb. Ic. Fl. Germ. xvi. 66 = L. squamulosum.
Tommasinii, Reichb. l. c.—Europ.

ADENONCOS. Blume, Bijdr. 381 (1825) = **Sarcochilus**, R. Br. ? (Orchid.).
virens, *Blume, l. c.*—Java.

ADENONEMA, Bunge, in Mém. Sav. Étrang. Pétersb. ii. (1836) 548; *reimpr.* Verz. Suppl. Fl. Alt. 36 = **Stellaria**, Linn. (Caryophyll.).
petraeum, Bunge, ll. cc. = S. petraea.

ADENOON, Dalz. in Hook. Kew Journ. ii (1850) 344. *COMPOSITAE*, Benth. & Hook. f. ii. 226.
indicum, *Dals. l. c.*—Ind. or.

ADENOPA, Rafin. Fl. Tellur. iii. 37 (1836) = **Drosera**, Linn.
anglica, Rafin. l. c. = D. anglica.

ADENOPAPPUS, Benth. Pl. Hartw. 41 (1840). *COMPOSITAE*, Benth. & Hook f. ii. 411.
persaefolius, *Benth. l. c.*—Mexico.

ADENOPELTIS, Bert. in Bull. Férussac. xxi. (1830), nomen; et ex A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 24. *EUPHORBIACEAE*, Benth & Hook. f. iii. 338.
Collignaya, *Bert. l. c.*—Chili.

ADENOPETALUM, Klotzsch & Garcke, in Monatsb. Akad. Berl. 1859 (1860) 250; et Abh. Acad. Berl. (1860) 47 = **Euphorbia**, Linn.
boerhaaviifolium, Klotzsch & Garcke, ll. cc. 250, 47 = E. boerhaviifolia.
bracteatum, Klotzsch & Garcke, ll. cc. 48 = E. scabrella.
discolor, Klotzsch & Garcke. ll. cc. 49 = E. discolor.
elliptieum, Klotzsch & Garcke, in Monatsb. Akad. Berl. 1859 (1860) 250 = E. graminea.
elliptieum, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 50 = E. saccharata.
gramineum, Klotzsch & Garcke, in Monatsb. Akad Berl (1859) 250; et in Abh. Akad. Berl. (1860) 47 = E. graminea.
Hoffmanni, Klotzsch & Garcke, ll. cc. 250, 47 = E. graminea.
vrasuense, Klotzsch & Garcke, ll. cc. 250, 50 = E. graminea.
mexicanum, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 50 = E. graminea.
Orrstedii, Klotzsch & Garcke, in Monatsb. Akad. Berl. 1859) 250; et in Abh. Akad. Berl. (1860) 48 = E. graminea.
pictum, Klotzsch & Garcke, ll. cc. 250, 47.—Hab. ?
pubescens, Klotzsch & Garcke, ll. cc. 250, 49 = E. graminea.
sphaerorhizum, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 250.—Hab. ?
subsinuatum, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 250; et in Abh. Akad. Berl. (1860) 48 = E. graminea.

ADENOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxxi (1858) i. 416 = **Vitis**, Linn. (Ampel.).
palmatum, Turcz. l. c. 417.—Hab. ?

ADENOPHAEDRA, Muell. Arg. in Mart. Fl Bras. xi. ii. 385. t. 101 (1873). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 314.
megalophylla, *Muell. Arg. l. c.* 386.—Bras.

ADENOPHORA, Fisch. in Mém. Soc. Nat. Mosc. vi. (1823) 165. *CAMPANULACEAE*, Benth. & Hook. f. ii. 563.
FLOERKEA, Spreng. Anleit. ii. ii. 533 (1818).
communis, *Fisch. l. c.* 168.—Europ.
cordata, *Tausch, in Flora*, xxv. (1842) 288.—Hab. ?
coronata, A. DC. Mon. Camp. 363 = marsupiiflora.
coronopifolia, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 167.—As. bor.
crispata, Turcz. ex Ledeb. Fl. Ross. ii. 893 = coronata.

ADENOPHORA :—
denticulata, Fisch. in Mém. Soc. Nat. Mosc. vi. (1823) 167 = tricuspidata.
divaricata, *Franch. & Sav. Enum. Pl. Jap.* ii. 428.— Japon.
Fischeri, G. Don, in Loud. Hort. Brit. 75 = communis.
Gmelini, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 167.—Dahur.
intermedia, Sweet, Hort. Brit. ed. I. 249 = communis.
intermedia, Ledeb. Ind. Sem. Hort. Dorpat. (1824) 1 = coronata.
Isabellae, *Hemsl. in Journ. Bot.* xiv (1876) 207.— China.
Lamarkii, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 168.—Sibir.
latifolia, *Fisch. l. c.*—Sibir.
lilifolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 48 = communis.
lilifolia, Ledeb. Cat. Hort. Dorp. (1822); ex A. DC. Prod. vii. 492 = communis.
marsupiflora, *Fisch. in Mém Soc. Nat. Mosc.* vi. (1823) 167.—Dahur.
montana, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) ii. 486.—Dahur
nikoensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 428.— Japon.
perchiaefolia, G. Don, in Loud. Hort. Brit. 74 = latifolia.
periploceifolia, *A. DC. Mon. Camp.* 361.—Sibir.
Pichleri, Vis. Suppl. ex Nym. Consp. 479 = Campanula trichocalycina.
polymorpha, *Ledeb. Fl. Alt.* i. 246 in nota.—Rossia.
pomponiifolia, Fisch. in Mém. Soc. Nat. Mosc. vi. (1823) 167 = Gmelini.
Rabelaisiana, *G. Don, in Loud. Hort. Brit.* 75.— Sibir.
remotiflora, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 193. —Japon.
reticulata, *G. Don, in Loud. Hort. Brit. Suppl.* 471.— Sibir.
sinensis, *A. DC. Mon. Camp.* 354.—China.
stricta, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 192.—Japon.
stylosa, *Fisch in Mém. Soc. Nat. Mosc.* vi. (1823) 168.— As. bor.
stylosa, Reichb Iconogr. Exot. 32. t. 45 = communis.
suaveolens, Reichb. l. c. 22. t. 32 = communis.
tetraphylla, Fisch. in Mém. Soc. Nat. Mosc. vi. (1823) 169 = verticillata.
trachelioides, *Maxim. Prim. Fl. Amur.* 186.—China.
tricuspidata, A. DC. Mon. Camp. 355 = denticulata.
triphylla, *A. DC. l. c.* 355.—Japon.
verbenaefolia, *Bunge, ex Heynh. Nom.* ii. 9.—Japon.
verticillata, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 167.—Dahur.; Japon.

ADENOPHYLLUM, Pers. Syn. ii. 458 (1807). *COMPOSITAE*, Benth. & Hook. f. ii. 408.
DYSSODIA, Willd. Enum. Hort. Berol. 900 (1809).
LEBETINA, Cass. in Dict. Sc. Nat. xxv. 394 (1822).
SCHLECHTENDAHLIA, Willd. Sp. Pl. iii. 2125 (1803).
WILLEBRANDIA, Cav. Ic. Pl. i. 61. t. 89 (1791).
capillaceum, DC. Prod. v. 638 = coccineum.
coccineum, *Pers. Syn.* ii. 458. —Mexico.
Porophyllum, *Hemsl. Biol. Centr. Am. Bot.* ii. 218.— Mexico.
Wrightii, *A. Gray, Pl. Wright.* ii. 92.—Mexico.

ADENOPLEA, Radlk. in Bremen, Abh. viii. (1883) 406 = **Buddleia**, Houst. (Logan.).
baccata, Radlk. l. c. 407 = B. sinuata.
sinuata, Radlk. in Ber. Deutsch. Bot. Ges. ii. (1884) 261 = B. sinuata.

ADENOPLUSIA, Radlk. in Bremen Abh. viii. (1883) 461. *LOGANIACEAE.*
axillaris, *Radlk. l. c.* 462.—Madag.
Wildenowii, *Radlk. in Ber. Deutsch. Bot. Ges.* ii. (1884) 280.—Madag.

ADENOPODIA, C. Presl, Epimel. Bot. 206 (1849) = **Bintada**, Adans. (Legum.).
spicata, C. Presl, l. c. 207 = E. natalensis.

ADENOPUS, Benth. in Hook. Niger Fl. 372 (1849). *CUCURBITACEAE*, Benth. & Hook. f. i. 828.
abyssinicus, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 328. —Afr. trop.

ADENOPUS :—
brevifloras, *Benth. in Hook. Niger Fl.* 372.—Afr. trop.
Cienkowskii, Schweinf. Rel. Kotschy. 45. t. 30 = Peponia Cienkowskii.
eglandulosus, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 329. —Afr. trop.
longiflorus, *Benth. in Hook. Niger Fl.* 372.—Afr. trop.

ADENORHOPIUM, Reichb. Consp. 195 (1828) = ADENOROPIUM, Pohl, = **Jatropha**, Linn. (Euphorb.).

ADENORIMA, Rafin. Fl. Tellur. iv. 112 (1836) = **Euphorbia**, Linn.
punicea, Rafin. l. c. = E. punicea.

ADENOROPIUM, Pohl, Pl. Bras. Ic. i. 12. t. 9 (1827) = **Jatropha**, Linn. (Euphorb.).
Berterii, Pohl, l. c. 14 = J. Berterii.
divaricatum, Pohl, l. c. 14 = J. divaricata.
diversgens, Pohl, l. c. 15 = J. Pohliana.
elegans, Pohl, l. c. 15 = J. gossypifolia.
elliptic um, Pohl, l. c. 13. t. 9 = J. elliptica.
Pohlchii, Pohl, l. c. 15 = J. glandulosa.
glaucum, Pohl, l. c. 15 = J. glauca.
gossypifolium, Pohl, l. c. 16 = J. gossypifolia.
hernandiaefolium, Pohl, l. c. 14 = J. hernandiaefolia.
integerrimum, Pohl, l. c. 14 = J. integerrima.
Jacquini, Pohl, l. c. 15 = J. gossypifolia.
luxurians, Pohl, l. c. 16 = J. Pohliana.
Martiusii, Pohl, l. c. = J. Martiusii.
mollissimum, Pohl, l. c. 15 = J. Pohliana.
multifidum, Pohl, l. c. 16 = J. multifida.
mutabile, Pohl, l. c. 14 = J. mutabilis.
pandurae folium, Pohl, l. c. 14 = J. hastata.
peltatum, Pohl, l. c. 16 = J. peltata.
ribifolium, Pohl, l. c. 15 = J. gossypifolia.
Riedelianum, Pohl, l. c. 16 = J. Pohliana.
spinosum, Pohl, Pl. Bras. Ic. i. 15 = J. sphosa.
tomentosum, Pohl, l. c. 16 = J. tomentosa.
tripartitum, Pohl, l. c. 15 = Manihot tripartita.
variegatum, Pohl, l. c. 14 = J. variegata.
villosum, Pohl, l. c. 15 = J. Pohliana.

ADENOSACHNA, A. Juss. in Orb. Dict. xi. 255 (1849) = seq.

ADENOSACME, Wall. Cat. n. 6280-82 (1832). *RUBIACEAE*, Benth. & Hook. f. ii. 69.
LAWIA, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 14.
MYCETIA, Reinw. in Syll. Ratieb. ii. (1828) 9.
caudiflora, *Miq. Fl. Ind. Bat.* ii. 215.—Malaya.
fasciculata, *Miq. l. c.* 216.—Java.
holotricha, *Miq. l. c.* 217.—Sumatra.
lanceolata, *Miq. l. c.* 217.—Sumatra.
Lawii, *Hook. f. Fl. Brit. Ind.* iii. 96.—Ind. or.
longiflora, *Wall. Cat.* n. 6280.—Ind. or.
macrostachya, *Hook. f. Fl. Brit. Ind.* iii. 96.—Ind. or.
malayana, Wall. Cat. n. 6283 = longiflora.
nepalensis, Wall. l. c. n. 6281 = longiflora.
obovata, Miq. Fl. Ind. Bat. ii. 216 = fasciculata.
stipulata, *Hook. f. Fl. Brit. Ind.* iii. 95.—Ind. or.

ADENOSCILLA, Gren. & Godr. Fl. Fr. iii. 187 (1855) = **Scilla**, Linn. (Liliac.).
bifolia, Gren. & Godr. l. c. = S. bifolia.
vivalis, J. Gay, ex Baker, in Journ. Linn. Soc. xiii. (1873) 289 = S. bifolia.
unifolia, Texidor, in Rev. Prog. Cienc. xviii. (1868).— Ct. Bull. Soc. Bot. Fr. xvi. (1869); Bibl. 229 = S. bifolia ?

ADENOSELEN, Spach. Hist. Vég. Phan. x. 26 (1841) = ADENOSOLEN, DC, = **Maraemodes**, DC. (Compos.).

ADENOSEPALUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 259 = **Hypericum**, Linn.
hirsutum, Fourr. l. c. 258 = H. hirsutum.
montanum, Fourr. l. c. = H. montanum.
nummularium, Fourr. l. c. = H. nummularium.
pulchrum, Fourr. l. c. 257 = H. pulchrum.
tomentosum, Fourr. l. c. = H. tomentosum.

ADENOSMA, R. Br. Prod. 449 (1810). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 949.
ANISANTHERA, Griff. Notul. iv. 100 (1854).
PTEROSTIGMA, Benth. Scroph. Ind. 20 (1835).
SPATHESTIGMA, Hook. & Arn. Bot. Beech. Voy. 204 (1836).
Brownii, *Zipp. ex Span. in Linnaea*, xv. (1841) 332, nomen.—Malaya.
caeruleum, *R. Br. Prod.* 443.—As. trop.; Austral.
camphoratum, *Hook. f. Fl. Brit. Ind.* iv. 264.— Zeylan.
capitatum, *Benth. ex Hance, in Journ. Linn. Soc. xiii.* (1873) 114.—As. trop.
chenopodiifolia, *Sprang. Syst.* ii. 829.—Ins. Societ.
chenopodioides, *Zipp. ex Span. in Linnaea*, xv. (1841) 332, nomen.—Hab. ?
cuspidatum, Benth. in Wall. Cat. n. 8852 = Lindenbergia macrostachya.
fragrans, *Sprang. Syst.* ii. 829.—Ins. Societ.
grandiflorum, *Benth. ex Hance, in Journ. Linn. Soc. xiii.* (1874) 114.—China.
hirsutum, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 143.—Ins. Nicobar.
macrophyllum, *Benth. in Wall. Cat.* n. 8853.—Ind. or.
malabaricum, *Hook. f. Fl. Brit. Ind.* iv. 264.— Malabar.
microcephalum, *Hook. f. l. c.*—As. trop.
Mueller, *Benth. Fl. Austral.* iv. 485.—Austral.
ovatum, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 263.— Malaya.
subrepens, *Benth. ex Hook. f. l. c.*—Zeylan.
trifora, Seem. Fl. Vit. 284 = Limnophila Roxburghii.
villosum, Benth. in Wall. Cat. n. 8851 = caeruleum.

ADENOSMA, Nees, in Wall. Pl. As. Rar. iii. 75 (1832) = **Cardanthera**, Buch.-Ham. (Acanthac.).
affinis, Griff. Notul. iv. 133 = Strobilanthes Acrocephalus.
africana, T. Anders. in Journ. Linn. Soc. vii. (1864) 21 —Afr. trop.
balsamea, Spreng. Syst. iii. 829 = C. balsamica.
bipinata, Nees, in Wall. Pl. As. Rar. iii. 79 = C. avana.
elongata, Blume, Bijdr. 757 = Ebermaiera elongata.
Griffithii, T. Anders. in Journ. Linn. Soc. ix. (1867) 454 = C. Griffithii.
lanceolata, Blume, Bijdr. 757 = E. lanceolata.
pinnatifida, T. Anders. in Journ. Linn. Soc. ix. (1867) 455 = C. pinnatifida.
polysperma, Spreng. Syst. ii. 829 = Hygrophila Serpyllum.
spathulata, Blume, Bijdr. 737 = Ebermaiera glauca.
Thymus, Nees, in Wall. Pl. As. Rar. iii. 79 = C. balsamica.
trifora, Nees, l. c. = C. trifora.
Thwaitesii, T. Anders. in Thw. Enum. Pl. Zeyl. 224; et in Journ. Linn. Soc. ix. (1867) 455 = C. Thwaitesii.
uliginosa, Nees, in R. Br. Verm. Schrift. iii. 398 = C uliginosa.
verticillata, Nees, in DC. Prod. xi. 69 = C. verticillata.

ADENOSOLEN, DC. Prod vi. 186 (1837) = **Maraemodes**, DC. (Compos.).
tenuifolius, DC. l. c.—Afr. austr.

ADENOSPERMUM, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 318 = **Chrysanthellum**, Rich. (Compos.)
tuberculatum, Hook. & Arn. l. c. = C. indicum.

ADENOSTEGIA, Benth. in Lindl. Syst. Nat. ed. II. 445 (1836) = **Oncothyrsa**, Nutt. (Scrophul.).
rigida, Benth. l. c. = C. filifolius.

ADENOSTEMA, Desport. in Dict. Sc. Nat. i. 256 (1816) = seq.

ADENOSTEMMA, Forst. Char. Gen. 89. t. 45 (1776). *COMPOSITAE*, Benth. & Hook. f. ii. 289.
LAVENIA, Sw. Prod. Veg. Ind. Occ. 119 (1788).
angustifolium, Arn. in Nov. Act. Nat. Cur. xvii. i (1836) 347 = viscosum.
aquaticum, D. Don, Prod. Fl. Nep. 81 = viscosum.
Berterii, *DC. Prod.* v. 110.—Ins. fl. Danubg.
bifforum, Less. Syn. Comp. 128 = Wedelia biflora.
brasilianum, *Cass. in Dict. Sc. Nat.* xxv. 363.—Bras.
cotfrum, *DC. Prod.* v. 112.—Afr. austr.
decumbens, Don, ex Heynh. Nom. i. 76 = Berterii.
Dregei, *DC. Prod.* v. 111.—Afr. austr.

ADENOSTEMMA:—
elatum, D. Don, Prod. Fl. Nep. 181 = viscosum.
erectum, DC. Prod. v. 113 = viscosum.
fastigiatum, DC. l. c. 113 = viscosum.
glandulosum, Benth. ex Hook. f. Fl Brit. Ind. i. 342, sphalm. (= Odontostemma) = Arenaria Benthami.
glutinosum, *DC. Prod.* v. 113.—Ins. Sandw.
Gymnocoronis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 188 = Gymnocoronis spilanthoides.
hirsutum, *DC. Prod.* v. 113.—Java.
hirtiflorum, *Benth. Pl. Hartw.* 75.—Mexico.
lanceolatum, *Miq. Fl. Ind. Bat.* ii. 24.—Malaya.
latifolium, D. Don, Prod. Fl. Nep. 181 = viscosum.
leiocarpum, DC. in Wight, Contrib. 9 = viscosum.
macrophyllum, DC. Prod. v. 113 = viscosum.
madurense, DC. in Wight, Contrib. 9 = viscosum.
mauritianum, DC. Prod. v. 110 = viscosum.
microcephalum, DC. l. c. 111 = viscosum.
natalense, *DC. l. c.* 112.—Afr. austr.
nitidum, Spreng. Syst. ii. 357, lapsu = Gomortega nitida.
ovatum, *Miq. Fl. Ind. Bat.* ii. 25.—Malaya.
parviflorum, DC. Prod. v. 111 = viscosum.
Perrottetii, *DC. l. c.* 110.—Prom. Virid.
platyphyllum, *Cass. in Dict. Sc. Nat.* xxv. 362.—Peruv.
reticulatum, DC. in Wight, Contrib. 8 = viscosum.
rivale, Dalz. in Hook. Kew Journ. iii. (1851) 231 = viscosum.
Roylei, DC. Prod. v. 113 = viscosum.
rufescens, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 120, nomen.—Malaya.
rugosum, DC. in Deless. Ic. Sel. 4 t. 10 = viscosum.
scarealifolium, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 186 = suffruticosum.
Schimperi, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 382.—Abyss.
suffruticosum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 488.—Bras.
Swartzii, *Cass. in Dict. Sc. Nat.* xxv. 362.—Ind. occ.; Ecuador.
tinctorium, Cass. l. c. = viscosum.
triangulare, DC. Prod. v. 113 = viscosum.
viscosum, *Forst. Char. Gen.* 90.—Gerout. trop.

ADENOSTEMON, Spreng. Anleit. ii. 788 (1818) = seq

ADENOSTEMUM, Pers. Syn. i. 467 (1805) = **Gomortega**, Ruiz & Pav. (Laurin.).
nitidum, Bert. Merc. Chil. 557; ex Am. Journ. Sc. xix. (1831) 67 = Cryptocarya Peumus.
nitidum, Pers. Syn. i. 467 = G. nitida.

ADENOSTEPHANUS, Klotzsch, in Linnæa, xv. (1841) 51 = **Euplassa**, Salisb. (Proteac.).
austro-caledonicus, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 230.—N. Caled.
bahiensis, Meisn. in Mart. Fl. Bras. v. i. 94. t. 36.—Bras.
Blanchetii, Benth. Fl. Austral. v. 417.—Austral.
Engenensis, Meisn. in Mart. Fl. Bras. v. i. 95 = E. meridionalis.
incana, Klotzsch, in Linnæa, xv. (1841) 52.—Bras.
inæquatis, Endl. Gen. Suppl. iv. ii. 87.—Bras.
laxiflora, Meisn. in Mart. Fl. Bras. v. i. 94.—Bras.
nitida, Meisn. l. c.—Bras.
organensis, Endl. Gen. Suppl. iv. ii. 87.—Bras.
Salzmii, Klotzsch, in Linnæa, xv. (1841) 52.—Bras.

ADENOSTOMA, Blume, in Flora, viii. (1825) 680, nomen. *SCROPHULARINEÆ.*

ADENOSTOMA, Hook. & Arn. Bot. Beech. Voy. 189, et 338. t. 80 (1839). *ROSACEÆ*, Benth. & Hook. f. i. 613.
brevifolium, Nutt. ex Torr. & Gray, Fl. N. Am. i. 480 = fasciculatum.
fasciculatum, *Hook. & Arn. Bot. Beech. Voy.* 189, 338. t. 80.—Calif.
sparsifolium, *Torr. in Emory, Notes Mil. Reconnois.* 140; *Bot. Bound.* 68. t. 20.—Calif.; Mexico.

ADENOSTYLES, Cass. in Dict. Sc. Nat. i. Suppl. 59 (1816). *COMPOSITÆ*, Benth. & Hook. f. ii. 247.
ADENOSTYLIUM, Reichb. Ic. Fl. Germ. xvi. ii. t. 895 (1853).

ADENOSTYLES:—
albida, *Cass.* ' *Bull. Soc. Philom.* 1816;' *Dict. Sc. Nat.* i. *Suppl.* 60.—Europ.
alliariæ, Reichb. Fl. Germ. Excurs. 278 = albida.
Alliariæ, *Kern. in Oestr. Bot. Zeitschr.* xxi. (1871) 12. —Europ.
alpina, Bluff & Fing. Comp. Fl. Germ. ii. 329 = viridis.
alpina, *Kern. in Oestr. Bot. Zeitschr.* xxi. (1871) 12.—Europ.
australis, Nym. Consp. 396 = viridis.
candidissima, *Cass. in Dict. Sc. Nat.* i. *Suppl.* 60.—Europ.
chlora, *Reichb. f Ic. Fl. Germ.* xvi. 11.—Europ.
glabra, DC. Prod. v. 208 = viridis.
hirsuta, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = albida.
hybrida, DC. Prod. v. 204 = alpina, candidissima, leucophylla, viridis.
leucophylla, DC. l. c. vii 271 = orientalis.
leucophylla, Reichb. Fl. Germ. Excurs. 278 = candidissima.
macrophylla, Tenore, Syll. ex Nym. Consp. 396 = viridis.
Nardosmia, A. Gray, in Proc. Am. Acad. viii. (1878) 631 = Cacaliopsis Nardosmia.
nebrodensis, *Strobl, in Flora*, lxv. (1882) 196.—Sicil.
orientalis, *Boiss. Fl. Orient.* iii. 155.—Oriens.
Petasites, Bluff & Fing. Comp. Fl. Germ. ii. 329 = albida.
pontica, *C. Koch, in Linnæa*, xiii. (1839) 696.—As. Min.
pyrenaica, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 94.—Hispan.
viridis, *Cass. in Dict. Sc. Nat.* i. *Suppl.* 60.—Europ.

ADENOSYCLES, Benth. & Hook. f. Gen. iii. 599 (1883) err. typ. = seq.

ADENOSYTLIS, Blume, Bijdr. 414 (1825) = **Zeuxine**, Lindl. (Orchid.).
emarginata, Blume, l. c. = Z. sulcata.
integerrima, Blume, l. c. = Z. integerrima.

ADENOSYTLIUM, Reichb. f Ic. Fl. Germ. xvi. 2. t. 895 (1853) = **Adenostyles**, Cass. (Compos.).
dubium, Reichb. l. c. = Adenostyles dubia.
hybridum, Reichb. l. c. = Adenostyles viridis.

ADENOTHECA, Welw ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 255 = **Schizobasis**, Baker (Liliac.).
aphylla, Welw. ex Baker, l. c. = S. angolensis.

ADENOTRIAS, Jaub. & Spach, Illustr. Pl. Or. i. (1842) 76 = **Hypericum**, Linn.
Kotschyi, Jaub. & Spach, l. c. 77 (= *Triadenia Russeggeri*).—Syria.
phrygia, Jaub. & Spach, l. c. 76 = præc.

ADENOTRICHIA, Lindl. Bot. Reg. t. 1190 (1828) = **Sanecio**, Linn. (Compos.).
amplexicaulis, Lindl. l. c. = S. adenotrichius.
senecioides, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 22 = S. sinuatilobus.

ADENUM, G. Don, Gen. Syst. iv. (1837) 80 = **Adenium**, Roem. & Schult. (Apocyn.).

ADESMIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 94. *LEGUMINOSÆ*, Benth. & Hook. f. i. 517.
LOUDONIA, Bert. ex Hook. Bot. Misc. iii. (1833) 198.
PATAGONIUM, Schrank, in Munch. Denkschr. (1808) 93, nomen vetustius.
STREPTODESMA, A. Gray, in Proc. Am. Acad. iii. (1853) 9.
Aegiceras, *Phil. in Linnæa*, xxviii. (1856) 684.—Chili.
affinis, *Hook. f. Fl. Antarct.* 257.—Terr. Magell.
alpina, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
angulata, Hook. f. Fl. Antarct. ii. 257 = incana.
angustifolia, *Hook & Arn. Bot. Beech. Voy.* 19.—Chili.
aphylla, *Clos, in C. Gay, Fl. Chil.* ii. 177.—Chili.
aprica, *Phil. in Anales Univ. Chil.* (1872) 697.—Chili.
arachnipes, *Clos, in C. Gay, Fl. Chil.* ii. 158.—Chili.
arsucana, *Phil. in Linnæa*, xxviii. (1856) 683.—Chili.

ADESMIA:—
arborea, *Bert. Merc. Chil.* xii. 559; *cf. Bull. Féruss.* xx (1830) 108.—Chili.
arborea, Colla, in Mem. Acc. Torin. xxxvii. (1834) 58 = glutinosa.
argentea, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
argentea, *Meyen, Reise*, i. 402, *et in Nov. Act. Nat Cur.* xix. *Suppl.* i. (1843) 22.—Peruv.
argyrophylla, *Phil. in Linnæa*, xxviii. (1856) 631.—Chili.
Arnottiana, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
arvensis, *Phil. in Linnæa*, xxxiii. (1864–65) 55.—Chili.
aspera, *Gill. in Hook. Bot. Misc.* iii. (1833) 190.—Chili.
atacamensis, *Phil. Fl. Atac.* 16.—Chili.
axillaris, *Phil. in Linnæa*, xxviii. (1856) 683.—Chili.
balsamica, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 59. t. 10.—Chili.
balsamifera, *Hook & Arn. Bot. Beech. Voy.* 20.—Chili.
Berteroniana, Steud. Nom. ed. II. i. 27 = arborea.
bicolor, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Reg. Argent.
bijuga, *Phil. in Anales Univ. Chil.* (1884) reimpr. 8.—Chili.
boronioides, *Hook. f. Fl. Antarct.* 257.—Terr. Magell.
brachycarpa, *Phil. in Linnæa*, xxviii. (1856) 684.—Chil.
brasicata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 192.—Chili.
caespitosa, Gill. ex Hook. l. c. 191 = horrida.
calocarpa, *Phil. in Anales Univ. Chil.* (1865) ii. 834.—Chili.
calycina, Vog. in Linnæa, xii. (1838) 75 = punctata.
candida, *Hook. f. Fl. Antarct.* 257.—Patag
canescens, *Phil. in Anales Univ. Chil.* (1872) 708.—Chili.
capercarnica, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = horrida.
Capricornu, *Phil. in Anales Univ. Chil.* (1862) ii. 893; *et in Linnæa*, xxxiii. (1864–65) 58.—Chili.
Caragana, *Griseb. in Goett. Abh.* xix (1874) 119.—Reg. Argent.
chilitanensis, *Phil. in Anales Univ. Chil.* (1865) ii. 826.—Chili.
ciliata, *Vog. in Linnæa*, xii. (1838) 74.—Bras.
cinerea, *Clos, in C. Gay, Fl. Chil.* ii. 179.—Chili.
Closii, *Phil. in Anales Univ. Chil.* (1872) 700.—Chili.
colutеoides, *Gill. in Hook. Bot. Misc.* iii. (1833) 190.—Reg. Argent.
colutеoides, *Phil. in Anales Univ. Chil.* (1872) 788.—Chili.
compacta, *Phil. in Linnæa*, xxviii. (1856) 628.—Chili.
conferta, *Hook. & Arn. Bot. Beech. Voy.* 20.—Chili.
coronillоides, *Gill. in Hook. Bot. Misc.* iii. (1833) 190.—Chili.
corymbosa, *Clos, in C. Gay, Fl. Chil.* ii. 165.—Chili.
cuneata, *Meyen, Reise*, i. 402, *et in Nov. Act. Cur.* xix. *Suppl.* i. (1843) 22.—Chili.
curvifolia, *Clos, in C. Gay, Fl. Chil.* ii. 189.—Chili.
cytisoides, *Griseb. in Goett. Abh.* xix. (1874) 119.—Reg. Argent.
Davilæ, *Phil. in Anales Univ. Chil.* (1872) 699.—Chili.
decipiens, *Steud. Nom.* ed. II. i. 27.—Chili.
decumbens, *Phil. in Linnæa*, xxviii. (1856) 628.—Chili.
dendroides, *Phil. l. c.* 683.—Chili.
dentata, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Chili.
dentata, Gill. ex Hook. Bot. Misc. iii. (1833) 189 = Gilliesii.
denticulata, *Clos, in C. Gay, Fl. Chil.* ii. 191.—Chili.
denticulata, *Steud. Nom.* ed. II. i. 27.—Chili.
denudata, *Phil. in Linnæa*, xxxiii. (1864–65) 56.—Chili.
desertícola, *Phil. in Anales Univ. Chil.* (1865) ii. 844.—Chili.
dichotoma, *Clos, in C. Gay, Fl. Chil.* ii. 180.—Chili.
diffusa, *Bert. ex Steud. Nom.* ed. II. i. 27.—Chili.
Doniana, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
Echinus, *Presl, Symb. Bot.* ii. 14. t. 16.—Am. austr.
elata, *Clos, in C. Gay, Fl. Chil.* ii. 160.—Chili.
elegans, *Clos, l. c.* 186.—Chili.

ADESMIA :—
elongata, *Phil. in Linnaea*, xxvii. (1856) 633.—Chili.
emarginata, *Clos, in C. Gay, Fl. Chil.* ii. 187.—Chili.
eremophila, *Phil. Fl. Atac.* 15.—Chili.
erinacea, *Phil. l. c.* 16.—Chili.
exilis, *Clos, in C. Gay, Fl. Chil.* ii. 160.—Chili.
filicaulis, *Phil. in Anales Univ. Chil.* (1872) 702.—Chili.
filifolia, *Clos, in C. Gay, Fl. Chil.* ii. 157.—Chili.
filipes, *A. Gray, Bot. U. S. Expl. Exped.* i. 426.—Patag.
floribunda, *G. Don, Gen. Syst.* ii. 288.—Mexico.
frigida, *Phil. Fl. Atac.* 16.—Chili.
fruticulosa, *G. Don, Gen. Syst.* ii. 282.—Mexico.
Gayana, *Phil. in Linnaea*, xxviii. (1856) 684.—Am. austr.
genistoides, *Presl, Symb. Bot.* ii. 16.—Am. austr.
Germaini, *Phil. in Linnaea*, xxviii. (1856) 629.—Chili.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 188.—Chili.
glabriuscula, Vog. in Linnæa, x. (1836) 592 = radicans.
glandulosa, *Phil. in Anales Univ. Chil.* (1861) 47; *et in Linnæa*, xxxiii. (1864–65) 57.—Chili.
glauca, *Phil. in Linnaea*, xxvii. (1856) 682.—Chili.
glaucescens, *Phil in Anales Univ. Chil.* (1872) 707.—Chili.
glomerata, *Auct. ex Phil. f. Cat. Pl. Chil.* 49 (sp. dub.).—Chili.
glomerula, *Clos, in C. Gay, Fl. Chil.* ii. 184.—Chili.
glutinosa, *Hook. & Arn. Bot. Beech. Voy.* 19.—Chili.
gracilis, *Meyen, ex Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 25.—Chili.
grandiflora, *Gill. in Hook. Bot. Misc.* iii. (1833) 190.—Chili.
graveolens, *Phil. Fl. Atac.* 16.—Chili.
grisea, *Hook. f. Fl. Antarct.* 257.—Patag.
hirsuta, *Phil. in Anales Univ. Chil.* (1872) 706.—Chili.
hispidula, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Peruv.
Hookeriana, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
horrida, *Gill. in Hook. l. c.*—Chili.
humifusa, *Phil. in Anales Univ. Chil.* (1872) 698.—Chili.
humilis, *Phil. in Linnaea*, xxvii. (1856) 682.—Am. austr.
Hystrix, *Phil. Fl. Atac.* 16.—Chili.
incana, *Vog. in Linnæa*, xii. (1838) 76.—Bras.
inconspicua, *Phil. l. c.* xxviii. (1856) 681.—Chili.
inflexa, *Griseb. in Goett. Abh.* xix. (1874) 121.—Reg. Argent.
Kingii, *Phil. in Anales Univ. Chil.* (1872) 706.—Chili.
Lambertiana, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
lanata, *Hook. f. Fl. Antarct.* 256.—Terr. Magell.
Landbecki, *Phil. in Linnaea*, xxxiii. (1864–65) 59.—Chili.
latifolia, *Vog. l. c.* xii. (1838) 74.—Bras.
laxa, *Clos, in C. Gay, Fl. Chil.* ii. 162.—Chili.
leiocarpa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 189.—Chili.
leptacantha, *Phil. in Anales Univ. Chil.* (1872) 705.—Chili.
longipes, *Phil. in Linnaea*, xxviii. (1856) 680.—Am. austr.
longiseta, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Chili.
lotoides, *Hook. f. Fl. Antarct.* 225.—Terr. Magell.
Loudonis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 193.—Chili.
macrostachya, *Benth. in Mart. Fl. Bras.* xv. i. 55.—Bras.
Meyeniana, *Phil. in Linnaea*, xxviii. (1856) 682.—Am. austr.
Meyeriana, Walp. Rep. i. 729 = Aeschynomene micrantha.
micrantha, *Phil. Fl. Atac.* 15.—Chili.
microphylla, *Hook. & Arn. Bot. Beech. Voy.* 19. t. 9.—Chili.
mimosoides, *G. Don, Gen. Syst.* ii. 282.—Mexico.
miraflorensis, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 357.—Bolivia.
monosperma, *Clos, in C. Gay, Fl. Chil.* ii. 184.—Chili.
monosperma, *Steud. Nom.* ed. II. i. 27.—Chili.
montana, *Phil. in Linnaea*, xxviii. (1856) 629.—Am. austr.

ADESMIA :—
mucronata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 189.—Chili.
multicapsula, *Clos, in C. Gay, Fl. Chil.* ii. 190.—Chili.
muricata, Bert. ex Steud. Nom. ed. II. i. 27 = decipiens.
muricata, *DC. in Ann Sc. Nat.* Sér. I. iv. (1825) 95.—*muricata*, Gill. ex Hook. Bot. Misc. iii. (1833) 188 = retrofracta.
obcordata, *Clos, in C. Gay, Fl. Chil.* ii. 199.—Chili.
obovata, *Clos, l. c.*—Chili.
obscura, *Clos, l. c.* 188.—Chili.
odontophylla, *Phil. in Anales Univ. Chil.* (1865) 11. 844.—Chili.
oligophylla, *Phil. in Linnaea*, xxviii. (1856) 686.—Chili.
Ononcela, *Griseb. in Lechl. Berb. Am. Austr.* 55, nomen.—Chili.
oresigena, *Phil. in Anales Univ. Chil.* (1872) 698.—Chili.
oxallophylla, *Phil. l. c.* (1865) 11. 845.—Chili.
papposa, Bert. ex Steud Nom. ed. II. i. 27 = angustifolia.
papposa, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Chili.
parviflora, *Clos, in C. Gay, Fl. Chil.* ii. 158.—Chili.
parvifolia, *Phil. in Linnaea*, xxviii. (1856) 683.—Am. austr.
parvula, *Phil. in Anales Univ. Chil.* (1872) 697.—Chili.
pauciflora, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 25.—Chili.
Pearcei, *Phil. in Anales Univ. Chil.* (1872) 704.—Chili.
pedicellata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 191.—Chili.
pendula, Bert. ex Steud. Nom. ed. II. i. 27 = denticulata, diffusa.
pendula, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Reg. Argent.
phylledes, *Clos, in C. Gay, Fl. Chil.* ii. 186.—Chili.
pinifolia, *Gill. in Hook. Bot. Misc.* iii. (1833) 192.—Chili.
polycarpha, *Wedd. Chlor. And.* ii. 265.—Am. austr.
propinqua, *Clos, in C. Gay, Fl. Chil.* ii. 188.—Chili.
prostrata, *Clos, l. c.* 189.—Chili.
psoraleoides, *Phil. in Anales Univ. Chil.* (1865) 11. 845.—Chili.
psoraleoides, *Vog. in Linnæa*, xii. (1838) 80.—Bras.
pugionata, *Griseb. in Goett. Abh.* xix. (1874) 120.—Reg. Argent.
pulchra, *Phil. in Linnaea*, xxviii. (1856) 632.—Am. austr.
pumila, *Hook. f. Fl. Antarct.* 255.—Patag.
punctata, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Am. austr.
pungens, *Clos, in C. Gay, Fl. Chil.* ii. 197.—Chili.
pusilla, *Phil. Fl. Atac.* 15.—Chili.
quadrijuga, *Phil. in Anales Univ. Chil.* (1872) 705.—Chili.
racemosa, Steud. Nom. ed. II. i. 27 = Aeschynomene micrantha.
radicans, *Bert. ex Steud. l. c.*—Chili.
radicans, *Clos, in C. Gay, Fl. Chil.* ii. 178.—Chili.
radiciflora, *Clos, l. c.* 160.—Chili.
ramosissima, *Phil. in Linnaea*, xxxiii. (1864–65) 56.—Am. austr.
retrofracta, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 188.—Chili.
retusa, *Griseb. in Goett. Abh.* vi. (1855) 119.—Am. austr.
rupicola, *Wedd. Chlor. And.* ii. 266.—Am. austr.
Schickendantzii, *Griseb. in Goett. Abh.* xxiv. (1879) 104.—Reg. Argent.
Schneidert, *Phil. in Linnaea*, xxxiii. (1864–65) 60.—Am. austr.
sericea, *Griseb. in Lechl. Berb. Am. Austr.* 55, nomen.—Chili.
sessilifolia, *Phil. Fl. Atac.* 16.—Chili.
Simonsi, *Phil. in Anales Univ. Chil.* (1872) 701.—Chili.
Smithiana, Bert. ex Steud. Nom. ed. II. i. 27 = muricata.
Smithiana, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Am. austr.
spinosissima, *Meyen, Reise*, ii. 27; *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 24.—Chili.
stipulacea, *Clos, in C. Gay, Fl. Chil.* ii. 164.—Chili.
subterranea, *Clos, l. c.* 192.—Chili.
suffocata, *Hook. f. Fl. Antarct.* 256.—Terr. Magell.
tenella, *Hook. & Arn. Bot. Beech. Voy.* 19.—Chili.

ADESMIA :—
tomentosa, *Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 23.—Chili.
tribuloides, *Steud. Nom.* ed. II. i. 27.—Chili.
trifoliata, *Gill. in Hook. Bot. Misc.* iii. (1833) 194.—Chili.
trifoliata, *Phil. in Linnaea*, xxviii. (1856) 685.—Am. austr.
trijuga, *Gill. in Hook. Bot. Misc.* iii. (1833) 191.—Chili.
tristis, *Vog. in Linnæa*, xii. (1838) 77.—Bras.
ulicina, *Presl, Symb. Bot.* ii. 15.—Am. austr.
unpalliatensis, *Gill. in Hook. Bot. Misc.* iii. (1833) 192.—Chili.
Valdesia, *Clos, in C. Gay, Fl. Chil.* ii. 174.—Chili.
verrucosa, *Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 26.—Chili.
vesicaris, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 58.—Chili.
villosa, *Hook. f. Fl. Antarct.* 256.—Reg. Magell.
virgata, *Bert. ex Steud. Nom.* ed. II. i. 27.—Chili.
viscida, Bert. Merc. Chil. xii. 557; Colla, in Mem. Acc. Torin. xxxvii. (1834) 59 = mucronata.
viscosa, *Gill. in Hook. Bot. Misc.* iii. (1833) 192.—Chili.
Volckmanni, *Phil. in Linnaea*, xxviii. (1856) 619.—Am. austr.
Wallichiana, Gill. ex Hook. Bot. Misc. iii. (1833) 190 = Aeschynomene aspera.

ADHADOTA, Steud. Nom. ed. II. i. 27 (1840) err. typ. = seq.

ADHATODA, Nees, in DC. Prod. xi. 384 (1847). Genus minime retinendum, species fere omnes ad *Justiciam* referendae. Vide infra.

ADHATODA, Tourn. ex Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 393; Nees, in Wall. Pl. As. Rar. iii. 102 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1112.
CARIMA, Rafin. Fl. Tellur. iv. 62 (1836).
DUVERNOYA, E. Mey. in Drège, Zwei Pfl. Docum. 150 (1843), nomen.
acuminata, *Nees, in DC. Prod.* xi. 400.—Abyss.
adenostachya, *Nees, l. c.* 408.—Bras.
amaranthoides, Nees, l. c. 400 = ceylanica.
anagalloides, *Nees, l. c.* 408.—Afr. austr.
Anselliana, Nees, l. c. 403 = Dianthera Anselliana.
arenaria, Nees, in Wall. Pl. As. Rar. iii. 108 = Justicia trinervia.
argyrostachya, Nees, l. c. 102 = Justicia argyrostachya.
aspera, *Nees, in DC. Prod.* xi. 397.—Ins. Philipp.
aurantiaca, *Regel, Ind. Sem. Hort. Petrop.* (1857) 47.—Hab.!
auriculata, *Nees, in DC. Prod.* xi. 407.—Bras.
bahiensis, *Nees, in Mart. Fl. Bras.* ix. 148.—Bras.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* 540.—Ind. or.
Betonica, Nees, in Wall. Pl. As. Rar. iii. 108 = Justicia Betonica.
boerhaaviaefolia, *Nees, in DC. Prod.* xi. 409.—Madag.
bojeriana, *Nees, l. c.* 396.—Madag.
candicans, Nees, l. c. 396 = Dianthera candicans.
capensis, *Nees, l. c.*—Afr. austr.
carthagenensis, Nees, l. c. 409 = Beloperone violacea.
cheiranthifolia, *Nees, l. c.* 387.—Afr. austr.
ceylanica, Nees, l. c. 400 = Justicia ceylanica.
chinensis, *Benth. in Hook. Kew Journ.* v. (1853) 184.—Hongkong.
chrysotrichoma, *Nees, in DC. Prod.* xi. 408.—Bras.
ciliata, *Nees, l. c.* 396.—Bras.
congrua, *Nees, in Mart. Fl. Bras.* ix. 147.—Bras.
cuneata, *Nees, in DC. Prod.* xi. 392.—Afr. austr.
cuneifolia, *Nees, l. c.* 405.—Bras.
cydoniaefolia, *Nees, in Mart. Fl. Bras.* ix. 148.—Bras.
decussata, Nees, in DC. Prod. xi. 408 = Justicia decussata.
diffusa, Benth. in Hook. Niger Fl. 483 = Justicia insularis.
dioscophylla, *Nees, in DC. Prod.* xi. 394.—Afr. austr.
Dipteracanthus, Nees, l. c. 396 = Siphonoglossa pilosella.
divaricata, *Nees, l. c.* 391.—Afr. austr.
divergens, *Nees, l. c.* 406.—Am. merid.
diversiformis, Nees, l. c. 399 = Justicia glauca.
Ecatenbyrana, Nees, l. c. 404 = Beloperone Ecatenbyana.
fascista, *Nees, l. c.* 402.—Afr. austr.

ADHATODA:—
Felisbertiana, *Nees, in DC. Prod.* xi. 406.—Bras.
flava, *Nees, l. c.* 401.—Arab.
flexuosa, *Nees, in Mart. Fl. Bras.* ix. 148.—Bras.
formosissima, *Kotzsch, in Peters, Reise Mossamb. Bot.* 215.—Afr. trop.
furcata, Nees, in DC. Prod. xi. 398 = Justicia furcata.
genistiformis, *Nees, l. c.* 394.—Bras.
Gilliesii, *Nees, l. c.* 395.—Reg. Argent.
glandulosa, *Nees, l. c.* 405.—Bras.
glomerata, *Miq. Fl. Ind. Bat.* ii. 831.—Malaya
holosericea, *Nees, in Mart. Fl. Bras.* ix. 148.—Bras.
Hookeriana, Nees, in DC. Prod. xi. 403 = Justicia Hookeriana.
Hypericum, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 102.—Afr. trop.
hysopifolia, *Medic. in Act. Acad Theod. Palat vi. Phys.* (1790) 393 = Dicliptera ocymoides.
hysopifolia, *Nees, in DC. Prod.* xi. 392.—Afr. austr.
incana, *Nees, l. c.* 393.—Afr. austr.
juleptorifolia, Buchen, ex Steud. Nom. ed 1. 15 = Dicliptera ocymoides.
Kotschyi, *Nees, in DC. Prod.* xi. 397 —Afr trop.
leptantha, *Nees, l. c.* 390.—Afr. austr.
leptostachya, *Nees, l. c.* 403 —Arab.
Lindeniana, *Nees, l. c.* 405.—N Granat.
lithospermifolia, *Nees, l. c.* 398.—Peruv.
longiflora, Torr. in Bot. Mex. Bound 125 = Siphonoglossa longiflora.
longifolia, *Nees, in DC. Prod* xi 404.— Ins Mascar
lupulina, *Nees, l. c.* 385.—Afr. austr.
Magnolia, *Noronha, in Verh. Batav. Gen* v. (1790) ed. 1. Art. 4. 6.— Malaya.
major, *Nees, in DC. Prod.* xi. 397.—Abyss.
Martiana, *Nees, l. c.* 405.—Bras.
matammensis, *Schweinf in Verh. Zool. Bot. Ges Wien*, xviii. (1868) 674.—Afr. trop.
microphylla, *Klotzsch, in Peters, Reise Mossamb. Bot* 216.—Afr. trop.
minor, *Nees, in DC. Prod.* xi. 400 —Abyss.
mollissima, *Nees, l. c.* 391.—Afr. austr.
Moretiana, *Miq. Fl. Ind. Bat* ii. 830 —Malaya
Moritziana, *Nees, in DC. Prod.* xi. 388.—N. Granat
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 217.—Afr trop
natalensis, *Nees, in DC. Prod.* xi. 391.—Afr austr.
Neesiana, Ness, l. c. 397 = Justicia Neesiana.
nigerrhensis, Nees, in Wall. Pl. As. Rar. iii. 108 = Justicia nigerrhensis.
nigherrica, Nees, in DC. Prod. xi. 386 = Justicia nigerrhensis.
nuda, *Nees, l. c.* 402.—Arab.
odora, *Nees, l. c.* 399.—Arab.
Orbigniana, *Nees, l. c.* 404.—Bolivia.
orchiloides, *Nees, l. c.* 393.—Afr. austr.
organoides, *Nees, l. c.* 407.—Ind. or.
orixensis, Nees, l. c. 400 = Justicia glauca.
pacifica, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 164 = Justicia pacifica.
palustris, *Nees, in DC. Prod.* xi. 402.—Afr. trop.
paniculata, *Benth. in Hook. Niger Fl.* 482.—Afr. trop.
patula, *Nees, in DC. Prod.* xi. 393.—Afr. austr.
periplocifolia, *Nees, l. c.* 395.—Venezuela.
petiolaris, *Nees, l. c.* 402.—Afr. trop.
plicas, *Nees, l. c.* 405.—Bras.
plicata, *Nees, l. c.* 401.—Afr. trop.
protracta, *Nees, l. c.* 390.—Afr. austr.
pubescens, Moench, Meth. 481 = Gendarussa Adhatoda.
pygmæa, *Nees, in DC. Prod.* xi. 394.—Afr. austr.
quadrifaria, Nees, l. c. 398 = Justicia quadrifaria.
ramosissima, Nees, in Wall. Pl. As. Rar. iii. 108 = Justicia Betonica.
reflexiflora, *Nees, in DC. Prod.* xi. 398.—Ind. occ.
rostellaria, Nees, l. c. 397 = Justicia neglecta.
rostrata, Hochst. ex Oliver, in Trans. Linn. Soc. xxix. (1872) 131 = Asystasia Schimperi.
rostrata, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 104. —Afr. trop.
rotundifolia, *Nees, in DC. Prod.* xi. 391.—Afr. austr.
scabra, *Nees, l. c.* 407.—Venezuela.
Schimperiana, *Hochst. ex Nees, l. c.* 388.—Abyss.
sericea, Nees, fide Benth. & Hook. f. Gen. ii. 1115 = Jacobinia mexicana.
siparuensis, Benth. in Journ. Linn. Soc. ix. (1867) 518 = Rangia siparuensis.
sphæroperma, *Nees, in DC. Prod.* xi. 409.—Ind. occ.

ADHATODA:—
spicata, Burm. ex Steud. Nom. ed. I. 15, 274 = Rungia repens.
spicata, *Nees, in DC. Prod.* xi. 408.—Arab.
striata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 216. —Afr. trop.
suaveolens, *Nees, in DC. Prod.* xi. 401.—Arab.
subserrata, *Nees, l. c.* 409.—Ins. Philipp.
sulcata, *Nees, l. c.* 401.—Arab.
tetramera, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 301.—Porto-Rico.
thymifolia, *Nees, in DC. Prod.* xi. 392.—Afr. austr.
tranquebariensis, Nees, l. c. 399 = Justicus tranquebariensis.
trinervia, Nees, in Wall. Pl. As. Rar. iii. 108 = Justicia trinervia.
trisetiflora, *Teysm. & Binn in Tijdschr. Nederl. Ind.* xxix. (1867) 250.—Java.
tristis, *Nees, in DC. Prod.* xi. 404 —Afr. trop.
tubulosa, *Nees, l. c.* 392.—Afr. austr.
Tweediana, *Nees, l. c.* 395.—Bras.
umbrosa, *Nees, l. c.* 406.—Bras.
variegata, *Nees, l. c.* 385.—Afr. austr.
vasculosa, Nees, l. c. 407 = Justicia vasculosa.
Vasica, *Nees, in Wall. Pl. As. Rar.* iii. 103.— As. trop.
Velasquezii, *Nees, in DC Prod* xi. 409 —Guatemala.
ventricosa, Nees, l. c 407 = Justicua ventricosa.
verticillata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. 1 Art. iv. 6 —Java
wynaadensis, Nees, in DC. Prod. xi. 406 = Justicia wynaadensis.
Zollingeriana, Nees, l. c. xi. 398 = Justicia Zollengeriana.

ADHUNIA, Vell. Fl. Flum. 236 (1825) ; v. t. 118 (1827). —'Gen. dub.'
emarginata, Vell ll. cc.— Bras.

ADIANTHUM *truncatum*, Burm. f. Fl. Ind. 235. t. 66. f. 4 = Acacia decipiens.

ADIANTUM *truncatum*, Murr. Syst. Veg. 790 = Acacia decipiens.

ADICEA, Rafin. Cat. 13 (1824) = **Urtica**, Tourn.
glabra, Rafin. t. c. = U. glabra.
pumila, Rafin. ex Torr Fl. New York, ii. 223 = Pilea pumila.
trinervia, Rafin. ex Torr. l. c = Pilea pumila.

ADIKE, Rafin. New Fl. Am. i. 63 (1836) ; Fl. Tellur. iii. 48 (1836) = **Pilea**, Lindl. (Urtic.).
allophyla, Rafin. Fl. Tellur. iii. 48 = **P.** trianthemoides.
glaberrima, Rafin. New Fl. Am. i. 64 = P. smilacifolia.
lanceolata, Rafin. l. c. = P. lanceolata.
pumila, Rafin. l. c. 63 = P. pumila.
rhomboidea, Rafin. l. c. 64 = P. rhombea.

ADINA, Salisb. Parad Lond. t. 115 (1807). *RUBIACEÆ*, Benth. & Hook. f. ii. 30.
arahiodes, *Benth. & Hook f. Gen.* ii. 30.—Java.
cordifolia, *Benth. & Hook. f. l. c.*—Ind. or.
globiflora, *Salisb. Parad. Lond.* t. 115 (errore *globifera*). China.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 24.—Ind. or.
microcephala, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 40. —Afr. trop.
pedunculata, DC. Prod. iv. 349 = globiflora.
polyeephala, *Benth. Fl Hongk.* 146.—Ind. or.
racemosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 184.— China ; Japon.
rubella, *Hance, in Journ. Bot.* vi. (1868) 114.—China.
sessiliflora, *Benth. & Hook. f. Gen.* ii. 30.—Ind. or. trichotoma, *Benth. & Hook. f. l. c.*—Malaya.

ADINANDRA, Jack, in Malay Misc. ii. n. 7. p. 49 (1823). *TERNSTROEMIACEÆ*, Benth. & Hook. f. i. 192.
HAENKEA, Usteri, Ann. Bot. vi. (1795) 117.
SAROSANTHERA, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 103. t. 16 (1840 !).
acuminata, *Korth. l. c. ed. Temminck*, 109.—Sumatra.
cordata, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 114.
crenulata, *T. Anders. in Hook. f. Brit. Ind. Fl.* i. 268. —Ind. or.
cyrtopoda, Miq. Fl. Ind. Bat. Suppl. i. 478 = dumosa.

ADINANDRA:—
dasyantha, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 160 = integerrima.
dasyantha, *Korth. Verh. Nat. Gesch. Bot. ed. Temminck*. 108.—Borneo.
dumosa, *Jack, in Malay Misc.* ii. (1822) n. 7. p. 50. —Malaya ; China.
dumosa, Vidal, Sinops. Filip. t. 12 = integerrima.
excelsa, *Korth. Verh. Nat. Gesch. Bot. ed. Temminck*, 109.—Borneo.
glabra, Miq. Fl. Ind. Bat. Suppl. i. 479 = dumosa.
Griffithii, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 282.— Reg. Himal.
integerrima, *T. Anders. ex Dyer, l. c.*—Malaya.
Jackiana, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 106 = dumosa.
javanica, *Choisy, in Zoll. Syst. Vers. Ind. Archip.* 148, 148.—Java.
Lamponga, *Miq. Fl. Ind. Bat. Suppl.* i. 479.— Sumatra.
laxipetala, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 114.—Zeylan.
leiopetala, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 104.— Sumatra.
macrantha, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 421.—Java.
maculosa, *T. Anders. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 282.—Malaya.
Mannii, *Oliver, Fl. Trop. Afr.* i 170.—Afr. trop.
Millettii, *Benth. & Hook. f. ex Hance, in Journ. Bot.* xvi. (1878) 9.—China.
phlebophylla, *Hance, l. c.* xiv. (1876) 240.—Cambodia.
Sarosanthera, *Miq. Fl. Ind. Bat.* i. ii. 477.—Borneo.
stylosa, Miq. l. c Suppl. 478 = dumosa.
sylvestris, *Jack, in Malay Misc.* ii. (1822) n. 7 p. iii. —Sumatra.
trichocarpa, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 107 = dumosa.
villosa, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) —Burma.

ADIPE, Rafin. Fl. Tellur. ii. 101 (1836) = **Bifrenaria**, Lindl. (Orchid.) (1842).
fulva, Rafin. l. c. = B. racemosa.
racemosa, Rafin. l. c. = B. racemosa.

ADIPERA, Rafin. Sylva Tellur. 129 (1838) = **Cassia**, Tourn. (Legum.).
Herberti, Rafin. l. c. = C. Herbertiana.

ADISA, Steud. Nom. ed. II. i. 28 (1840), sphalm. = seq.

ADISCA, Blume, Bijdr. (1825) 609 = **Mallotus**, Lour. (Euphorb.).
acuminata, Blume, l. c. 610 = M. acuminatus.
albicans, Blume, l. c. 611 = Cephalocroton albicans.
floribunda, Blume, l. c. 610 = M. floribundus.
subfalcata, Zoll. in Linnæa, xxix. (1857–58) 464 = Macaranga subfalcata.
subpeltata, Blume, Bijdr. 610 = M. subpeltatus.
timoriana, Span. Ic. n. 26 ; et in Linnæa, xv. (1841) 948 = M. repandus.
Zippelii, Blume, Bijdr. 611 = M. ricinoides.

ADLERIA, Neck. Elem. ii. 642 (1790) = **Eperua**, Aubl. (Legum.).

ADLUMIA, Rafin. in Med. Repos. New York, v. (1808) 352. *PAPAVERACEÆ*, Benth. & Hook. f. i. 55.
BICUCULLA, Borckh. in Roem. Arch. i. ii. 46 (1797), nomen prius.
cirrhosa, *Rafin. l. c.*—Am. bor.

ADNARIA, Rafin. Fl. Ludov. 58 (1817). *VACCINIACEÆ.*
odorata, *Rafin. l. c.*—Am. bor.

ADNULA, Rafin. Fl. Tellur. ii. 87 (1836) = **Pelexia**, Poit. (Orchid.).
petiolaris, Rafin. l. c. = P. spiranthoides.

ADODENDRON, DC. Prod. vii. 725 = seq.

ADODENDRUM, Neck. Elem. i. 214 (1790) = **Rhododendron**, Reichb. (Ericac.).

ADOKETON, Rafin. Mus. N. Sc. 1 ; Nev. Fl. Am. i. 65 (1836) = **Telanthera**, R. Br. (Chenop.).
sessatile, Rafin. ll. cc. = T. frutescens.
uniflorum, Rafin. New Fl. Am. i. 65.—Am. bor.

ADOLIA, Lam. Encyc. i. (1783) 44 = **Boutia**, Comm. (sec. Rodl. Gen. Suppl. i. 1424) (Rhamnac.).
alba, Lam. l. c. 45 = S. indica.
rubra, Lam. l. c. = S. indica.

ADOLPHIA, Meisn. Gen. 70 (50) (1837). *RHAMNEAE*, Benth. & Hook. f. i. 384.
californica, *S. Wats. in Proc. Am. Acad.* xi. (1876) 126. —Calif.
infesta, *Meisn. Gen. Pl.* 70 (50).—Mexico.

ADONANTHE, Spach, Hist. Vég. Phan. vii. 227 (1839) = **Adonia**, Linn. (Ranunc.).
pyrenaica, Spach, l. c. 230 = Adonis pyrenaica.
sibirica, Spach, l. c. 229 = Adonis sibirica.
vernalis, Spach, l. c. 228 = Adonis vernalis.

ADONIASTRUM, Schur, in Verh. Naturf. Ver. Bruenn, xv. ii. (1877) 27 = **Adonis**, Dill. (Ranunc.).
vernale, Schur, l. c. = Adonis vernalis.

ADONIGERON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = **Senecio**, Linn. (Compos.).
adonidifolius, Fourr. l. c. = S. adonidifolius.

ADONIS, Dill. ex Linn. Syst. ed. I. (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 5.
ADONANTHE, Spach, Hist. Vég. Phan. vii. 227 (1839).
ADONIASTRUM, Schur, in Verh. Naturf. Ver. Bruenn, xv. ii. (1877) 27.
COSMARIUM, Dulac, Fl. Hautes-Pyr. 216 (1867).
aestivalis, Bieb. Fl. Taur. Cauc. iii. 278 = autumnalis.
aestivalis, *Linn. Sp. Pl.* ed. II. 771.—Europ ; Oriens.
aestivalis, Link, ex Webb & Berth. Phyt. Canar. i. 12 = microcarpa.
aethiopica, Thunb. Prod. Pl. Cap. 94 = Knowltonia gracilis.
alepplca, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 250.—Syria.
ambigua, Gaud. Fl. Helv. iii. 518 = aestivalis.
amurensis, *Regel & Radde, in Bull. Soc. Nat. Mosc.* (1861) ii. 25. t. 2. f. 15.—Manchur.
annua, Linn. Sp. Pl. 547 = aestivalis et autumnalis.
anomala, Wallr. Sched. Crit. 278 = flammea.
apennina, Linn. Sp. Pl. 548 = vernalis.
apennina, Pall. Reise, ii. 368 = wolgensis.
apennina, Schouw, ex DC. Prod. i. 25 = pyrenaica.
autumnalis, Bieb. Fl. Taur. Cauc. ii. 28 = aestivalis.
autumnalis, *Linn. Sp. Pl.* ed. II. 771.—Europ ; Oriens.
barbica, Cos. Not. Fl. Crit. ii. 25 = autumnalis.
caerulea, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 206.—China.
capensis, Linn. Sp. Pl. 548 = Knowltonia rigida.
capensis, Thunb. Prod. Pl. Cap. 94 = Knowltonia gracilis.
caucasica, Stev. in Bull. Soc. Nat. Mosc. xxi. (1848) ii. 271, 278 = flammea.
chaerophylla, Fisch. in DC. Prod. i. 25 = wolgensis.
Chrysocyathus, *Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind.* 15.—Reg. Himal.
citrina, DC. Syst. i. 228 = flammea.
citrina, Hoffm. Fl. Germ. i. 251 = aestivalis.
coriacea, Poir. Encyc. Suppl. i. 146 = Knowltonia rigida.
Cupaniana, Guss. Syn Fl. Sic. ii. 86 = microcarpa.
cyllenea, *Boiss. Heldr. & Orph. Diagn.* Sér. II. v. 5.—Graecia.
dasycfolia, Lam. Encyc. i. 46 = Knowltonia dasycfolia.
dcnvorica, Reichb. Ic. Crit. Cent. iv. 21. t. 351 = vernalis.
dentata, *Delile, Fl. Egypte*, 287.—Europ. ; Afr. bor.
dentata, Duby, ex Nym. Consp. 4 = aestivalis.
distorta, *Tenore, Fl. Nap.* iv. 327. t. 149.—Ital.
erioclypicla, *Boiss. Fl. Orient.* i. 17.—Armenia.
erosipetala, Stokes, Bot. Mat. Med. iii. 255 = autumnalis.
filia, Lam. f. Suppl. 271 = Knowltonia dasycfolia.
flammea, *Jacq. Fl. Austr.* iv. 29. t. 355.—Europ. ; Oriens.
flammea, Schleich. ex Steud. Nom. ed. II. i. 28 = aestivalis.
flava, Vill. Cat. Jard. Strasb. 247 = aestivalis.

ADONIS:—
fulgens, *Hochst. in Lorent, Wanderungen*, 338.—Ab. Mt.
gracilis, Poir. Encyc. Suppl. i. 147 = Knowltonia gracilis.
Hellebrus, Crantz, Stirp. Austr. ed. I. fasc. iii. 82 ; ed. II. 110 = vernalis.
hirsuta, Poir. Encyc. Suppl. i. 147 = Knowltonia hirsuta.
hybrida, Wolff, ex Nym. Consp. 4 = wolgensis.
Ingilica, Royle, Illustr. Himal. Pl. 53 = aestivalis.
intermedia, Webb & Berth. Phyt. Canar. i. 121 = microcarpa.
iresionea, Fisch. ex DC. Prod i. 25 = vernalis.
laserpitifolia, Poir. Encyc. Suppl. i. 147 = Knowltonia vesicatoria.
maculata, *Roth, Enum. Pl. Phan. Germ.* ii. 602.—Europ.
maculata, Wallr. Sched. Crit. 270 = aestivalis.
magellanica, Raeusch, Nom. ed. III. 162 = Hamadryas magellanica.
marginata, Bienert, ex Hook. f. Fl. Brit. Ind. i. 15 = scrobiculata.
Meschatiana, Andrz. ex Bess. Enum. Pl. Volh. 47 = wolgensis.
micrantha, DC. Syst. i. 222 = autumnalis.
microcarpa, *DC. l. c.*—Europ.
miniata, Jacq. Fl. Austr. iv. 28 = aestivalis.
palaestina, *Boiss. Diagn.* Ser. I. viii. 1.—Syria.
palustris, Raeusch, Nom. ed. III. 162 = Ranunculus sceleratus.
parviflora, *Fisch. ex DC. Prod.* i. 24.—Oriens.
parviflora, Janka, ex Nym. Consp. 4 = vernalis.
perramosa, Schur, Enum Pl. Transs. 6 = autumnalis.
persica, Boiss. Diagn. Ser. I. vi. 4 = dentata.
phoenicea, Bercht. et Presl, Rostl. i. Ranunculac. 42 = autumnalis.
pilosa, Raeusch, Nom. ed. III. 162 = Ranunculus pensylvanicus.
polymorpha, Zumag. Fl. Pedem. ii. 362 = aestivalis, autumnalis.
pratensis, Ledeb. Fl. Ross. i. 24 = vernalis.
Presliі, Tod. ex Nym. Consp. 4 = flammea.
pyrenaica, *DC. Fl. Fr.* v. 635.—Europ.
pyrenaica, Hook. f. & Thoms. Fl. Ind. 26 = A. Chrysocyathus.
riparia, *Rafin. New Fl. Am.* i. 66.—Am. bor.
scrobiculata, *Boiss. Diagn.* Ser. II. i. 6 —Afghan.
sibirica, *Patrin, ex Ledeb. Ind. Hort. Dorp. Suppl.* (1824) 1.—Sibir. altaic.
Sibthorpii, *Boiss. Orphan. & Heldr. in Bull. Congr. Bot. Pétersb.* 1869 (1870) 124.—Graecia.
squarrosa, *Stev. in Bull. Soc. Nat.-Mosc.* xxi. (1848) ii. 271, 272 = aestivalis.
vernalis, Asso, Syn. Stirp. Arag. 70 = dentata.
vernalis, *Linn. Sp. Pl.* 547.—Europ. ; As. bor.
vesicatoria, Linn. f. Suppl. 272 = Knowltonia vesicatoria.
villosa, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 1 = wolgensis.
wolgensis, *Stev. ex DC. Syst.* i. 545.—Europ.; Oriens; As. bor.

ADOPOGON, Neck. Elem. i. 55 (1790) = **Krigia**, Schreb. (Compos.).

ADORIUM, Rafin. Neogenyt. 3 (1825) = **Sesell**, Linn. (Umbellif.).
crassifolium, Rafin. l. c. = S. divaricatum.

ADOXA, Linn. Syst. ed. I. (1735). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 2.
MOSCHATELLA, Adans. Fam. ii. 248 (1763).
MOSCHATELLINA, [Tourn.] Hall. Enum. Stirp. Helv. i. 432 (1749).
inodora, Falc. ex Clarke, in Hook. f. Fl. Brit. Ind. iii. 2 = A. Moschatellina.
moschata, Dulac, Fl. Hautes-Pyr. 402 = A. Moschatellina.
Moschatellina, *Linn. Sp. Pl.* 367.—Europ. ; As. bor. ; Am. bor.
tuberosa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 498 = A. Moschatellina.

ADRASTAEA, DC. Syst. Veg. i. 424 (1818). *DILLENIACEAE*, Benth. & Hook. f. i. 15.
ADRASTEA, Spreng. Syst. ii. 360 (1825).
salicifolia, *DC. l. c.*—Austral.

ADRIANA, Gaudich. in Ann. Scr. Nat. Ser. I. v. (1825) 238. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 306.
TRACHYCARYON, Klotzsch, in Lehm. Pl. Preiss. i. 175 (1844).
acerifolia, Nelson, ex Mitch. Journ. Trop. Austral. 371 = glabrata.
Billardieri, Baill. Etud. Gén. Euph. Atl. 6. t. 2. f. 19 = quadripartita.
Gaudichaudi, Baill. Adansonia, vi. (1865–66) 312 = tomentosa.
glabrata, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 238.—Austral.
heterophylla, Hook. in Mitch. Journ. Trop. Austral. 371 = glabrata.
Hookeri, *Muell. Arg. in DC. Prod.* xv. ii. 893.—Austral.
Klotzschii, *Muell. Arg. l. c.* 892.—Austral.
quadripartita, *Gaudich. Bot. Voy. Freyc.* 439.—Austral.
tomentosa, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 238.—Austral.

ADROMISCHUS, Lem. Jard. Fleur. ii. (1852) Misc. 59 = **Cotyledon**, Tourn. (Crassul.).
caryophyllaceus, Lem. l. c. 60 = C. caryophyllaceus.
clavifolius, Lem. l. c. = C. clavifolia.
cristatus, Lem. l. c. = cristata.
cuneatus, Lem. l. c. = cuneata.
hemisphaericus, Lem. l. c. = C. hemisphaerica.
jasminiflorus, Lem. l. c.—C. jasminiflora.
maculatus, Lem. l. c. = C. maculata.
mammillaris, Lem. l. c. = C. mammillaris
mucronatus, Lem. l. c. = C. maculata.
rhombifolius, Lem. l. c. = C. rhombifolia.
robustus, Lem. l. c. = C. trifiora.

ADULPA, Endl. Gen. (1836) 119 (err. typ.) = seq.

ADULPA. Bosc, in Jaume St. Hil. Expos. Fam. Nat. i 65 (1805) = **Cyperus**, Linn.

ADVENTINA, Rafin. New Fl. Am. i. 67 (1836). *COMPOSITAE*.
ciliata, Rafin. l. c.—Am. bor.
parviflora, Rafin. l. c.—Am. bor.

ADYSETON, Adans. Fam. ii. 420 (1763) = **Alyssum**, Linn. (Crucif.).
alpestre, Sweet, Hort. Brit. ed. I. 467 = Alyss. alpestre.
altaicum, G. Don, in Sweet, l. c. ed. III. 80 = Alyss. Fischerianum.
argenteum, Sweet, l. c. ed. I. 467 = Alyss. argenteum.
atlanticum, G. Don, Gen. Syst. i. 178 = Alyss. atlanticum.
Bertolonii, Sweet, Hort. Brit. ed. I. 467 = Alyss. montanum.
calycinum, Scop. Fl. Carn. ed. II. ii. 18 = Alyss. calycinum.
campestre, Moench, Meth. Suppl. 87 = Alyss. campestre.
cuneifolium, Sweet, Hort. Brit. ed. I. 467 = Alyss. cuneifolium.
dentatum, G. Don, Gen. Syst. i. 179 = Camelina sativa.
diffusum, Sweet, Hort. Brit. ed. I. 467 = Alyss. montanum.
dubium, G. Don, Gen. Syst. i. 179 = Alyss. dubium.
edentulum, Link, Enum. Hort. Berol. ii. 157 = Alyss. gemonense.
Fischerianum, Sweet, Hort. Brit. ed. II. 25 = Alyss. canescens.
gemonense, Sweet, l. c. = Alyss. gemonense.
graphiodes, G. Don, Gen. Syst. i. 179 = Vesicaria graphiodes.
halimifolium, Link, Enum. Hort. Berol. ii. 157 = Alyss. halimifolium.
hyperboreum, G. Don, Gen. Syst. i. 179 = Alyss. hyperborreum.
lanigerum, G. Don, l. c. 178 = Alyss. lanigerum.
lenense, G. Don, l. c. = Alyss. lenense.
maritimum, Link, Enum. Hort. Berol. ii. 157 = Alyss. maritimum.
Marschallianum, Sweet, Hort. Brit. ed. II. 25 = Alyss. Marschallianum.
montanum, Scop. Fl. Carn. ed. II. ii. 14 = Alyss. Wulfenianum.
murale, Sweet, Hort. Brit. ed. I. 467 = Alyss. argenteum.
mutabile, Moench, Meth. 267 = Alyss. calycinum.

ADYSETON :—
nobrodense, Sweet, Hort. Brit. ed. II. 25 = Alysa. serpyllifolium.
obcordatum, G. Don, Gen. Syst. i. 179 = Alysa. obcordatum.
obtusifolium, Sweet, Hort. Brit. ed I. 467 = Alysa. alpestre.
orientale, Sweet, l. c. 467 = Alysa. orientale.
petiolodes, G. Don, Gen. Syst. i. 179 = Farsetia obovata.
repens, Sweet, Hort. Brit. ed. II. 25 = Alysa. alpestre.
saxranicum, G. Don, Gen. Syst. i. 178 = Alysa. alpestre.
saxatile, Sweet, Hort. Brit. ed. I. 467 = Alysa. saxatile.
serpyllifolium, Sweet, l. c. = Alysa. serpyllifolium.
sibiricum, G. Don, Gen. Syst. i. 179 = Alysa. saxatile.
spathuiatum, Sweet, Hort. Brit. ed. I. 467 = Alysa. spathulatum.
spinosum, Link, Enum. Hort. Berol. ii. 157 = Alysa. spinosum.
tortuosum, Sweet, Hort. Brit. ed. I. 467 = Alysa. alpestre.
verrale, Sweet, l. c. = Alysa. rostratum
Wulfenianum. Sweet, l. c. = Alysa. montanum.

ADYSETUM, Link, Enum. Hort. Berol. ii. 157 (1822) = prsec.

AECEOCLADES, Duchartre, in Orb Dict. ix. 170 (1848) err. typ. = OECEOCLADES, Lindl.

AECHMAEA, Brongn. in Ann. Sc Nat, Sér. II. xv. (1841) 371 = **Aechmea**, Ruiz & Pav. (Bromel.).

AECHMANDRA, Arn. in Hook. Lond. Journ. Bot iii. (1841) 274 = **Malothria**, Linn. (Cucurb.).
Blumeana, M. Roem. Syn. Pepon. 33 = M. Rumphiana.
conocarpa, Dalz. & Gibs. Bomb. Fl. 100 = Corallocarpus conocarpus.
deltoidea, Arn. in Hook. Lond. Journ. Bot. iii. (1841) 274 = M. zeyianica.
epigaea, Arn. l. c. = Corallocarpus epigeus
indica, C. B. Clarke, in Hook. f. Fl. Brit. Ind ii 626 = M. indica.
odorata, Hook. f. & Thoms. ex Benth. Fl. Hongkong, 124 = M. odorata.
rostrata, Arn. in Hook. Lond. Journ. Bot. iii. (1841) 274 = Kedrostis rostrata.
velutina, Dalz. & Gibs. in Bomb. Fl 200 = Corallocarpus velutina.
zeylanica, Thw. Enum. Pl. Zeyl. 125 = Cerasiocarpum zeylanicum.

AECHMANTHERA, Nees, in Wall. Pl. As. Rar. iii. 75, 87 (1837). *ACANTHACEAE*, Benth. & Hook f. ii. 1088.
gossypina, Nees, l. c. 87 = tomentosa.
leiosperma, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 429.—Ind. or.
tomentosa, *Nees, in Wall. Pl. As. Rar.* iii. 87.—Ind. or.
Wallichii, Nees, in DC. Prod. xi 170 = tomentosa.

AECHMEA, Ruiz & Pav. Prod. 47. t. 8 (1794). *BROMELIACEAE*, Benth. & Hook. f. iii. 668.
ANDROLEPIS, Brongn. ex Houllet, in Rev. Hortic. (1870) 12.
CANISTRUM, E. Morr. Belg. Hortic. (1873) 267. t. 15.
DIQUAMIA, Lem. Jard. Fleur. iii. (1851-3) t. 289.
ECHINOSTACHYS, Brongn. in Planch. Hort. Donat. 25 (1854).
ERONTAX, Rafin. Fl. Tellur. iv. 25 (1836).
HOHENBERGIA, Schult. f. Syst. vii. II. 71, 1251 (1830).
HOPLOPHYTUM, E. Morr. Belg. Hortic. (1868) 162.
LAMPROCOCCUS, Beer, Bromel. 108 (1857).
MACROCHORDION, De Vriese, in Jaarb. Nederl. Maat. Tuinb. (1858) 14.
PIRONNEAVA, Gaudich. Voy. Bonite, Bot. t. 63. (1839–52).
POTHUAVA, Gaudich. l. c. tt. 116, 117 (1839–52).
angustifolia, Poepp. & Endl. Nov. Gen. et Sp. ii. 48 = spicata.
aquilega, *Griseb. Fl. Brit. W. Ind.* 592.—Ind. occ.
augusta, *Baker, in Journ. Bot.* xvii. (1879) 162.—Am. trop.
aurantiaca, *Baker, l. c.* 235.—Am. trop.
Barlesi, *Baker, in Gard. Chron.* xx. (1883) 102.—Honduras.
bracteata, *Griseb. Fl. Brit. W. Ind.* 592.—Ind. occ.
brasiliensis, *Regel, Gartenfl.* (1885) 258.—Bras.
brasiocoides, *Baker, in Journ. Bot.* xx. (1882) 329.—Brit. Guiana.

AECHMEA:—
bromeliaefolia, *Baker, ex Benth. & Hook. f. Gen.* iii. 664.—Am. trop.
Burchellii, *Baker, in Journ. Bot.* xvii. (1879) 231.—Am. trop.
caerulea, E. Morr. Cat. (1871) 1 = seq.
caerulescens, *Hort. ex Baker, in Journ. Bot.* xvii. (1879) 227.—Am. trop.
calyculata, *Baker, in Journ. Bot.* xvii. (1879) 382.—Am. trop.
capitata, Baker, l. c. 167 = exsudans.
coelestis, *E. Morr. Fl. des Serres*, Sér. II. xi. (1875) t. t. 2146.—Am. trop.
columnaris, *André, in Illustr. Hortic.* xxv. (1878) 50.—N. Granat.
comata, *Baker, in Journ. Bot.* xvii. (1879) 234.—Am. trop.
contracta, *Baker, l. c.*—Am. trop.
corallina, *Brongn. ex Beer, Bromel.* 106.—Bras.
Cornui, Carr. in Rev. Hortic. (1885) 36 = nudicaulis.
Cumingii, *Baker, in Journ. Bot.* xvii. (1879) 227.—N. Granat.
cymoso-paniculata, *Baker, l. c.* 165.—Venezuela.
dactylina, *Baker, l. c.* 161.—Panama.
dichlamydes, *Baker, l. c.* 233.—Ins. Tobago.
discolor, Morr. in Ann Soc Hort. Gand, ii. (1846) 176. t. 66 = fulgens.
distans, *Griseb. Fl. Brit. W. Ind.* 592.—Ind. occ.
distans, Griseb. Pl. Cub. 253 = Wrightii.
distichantha, *Lem. Jard. Fleur.* iii. (1853) t. 269.—Bras.
exsudans, *E. Morr. ex C. H. Morr. in Belg. Hortic.* xxvii. (1878) 303.—Bras.
excavata, *Baker, in Journ. Bot.* xvii. (1879) 134.—Paraguay.
fasciata, *Baker, l. c.* 231.—Am. trop.
floribunda, *Mart. ex Schult. f. Syst.* vii. II. 1271.—Bras.
fulgens, *Brongn. in Ann. Sc. Nat. Sér.* II. xv. (1841) 371.—Guiana.
Furstenbergii, E. Morr. ex C. H. Morr. in Belg. Hortic. xxvii. (1878) 303 = Aechmea Furstenbergii.
Glaziovii, *Baker, in Journ. Bot.* xvii. (1879) 133.—Bras.
glomerata, *Hook. f. Bot. Mag.* t. 5668.—Bras.
Hookeri, *Lem. Illustr. Hortic.* xi. (1864) *Misc.* 42.—Bras.
Hystrix, *E Morr. ex C. H. Morr. in Belg. Hortic.* xxx. (1880) 90 & 243.—Guiana.
Jenmani, Baker, in Journ. Bot. xx. (1882) 329 = Mellinoni.
Lalindei, *Linden & Rodigas, in Illustr. Hortic.* xxx. (1883) t. 481.—N. Granat.
laxiflora, *Benth. in Bot. Voy. Sulph.* 173.—Mexico.
Legrelliana, *Baker, in Journ. Bot.* xvii. (1879) 236.—Am. merid.
Lindeni, *E. Morr. ex C. Koch, Wochenschr.* viii. (1865) 398.—Bras.
lingulata, *Baker, in Journ. Bot.* xvii. (1879) 164.—Jamaic.
Luddemanniana, Brongn. ex E. Morr. Cat. Bromel. (1871) 1 = caerulescens.
macrantha, Brongn. in Illustr. Hortic. (1880) 59 = Schiedeana.
Mariae-Reginae, *H. Wendl. in Hamb. Gartens.* xix. (1863) 23.—Am. centr.
martinicensis, *Baker, in Journ. Bot.* xvii. (1879) 132.—Ins. Martinic.
melanacantha, *De Vriese, in Tuinb. Fl.* i. (1855) 856.—Hab.?
Melinonii, *Hook. Bot. Mag.* t. 5225.—Am. trop.
Mertensii, *Schult. f. Syst.* vii. II. 1272.—Am. austr. mexicana, *Baker, in Journ. Bot.* xvii. (1879) 165.—Mexico.
Meyeri, Baker, 4. c. xviii. (1880) 16 = squillega.
microcantha, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 370 = Araeococcus micranthus.
mucronifora, *Hook. Bot. Mag.* t. 4832.—Guiana.
multiceps, *Baker, in Journ. Bot.* xviii. (1880) 49.—Am. trop.
Noettigii, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 117.—Bras.
nudicaulis, *Griseb. Fl. Brit. W. Ind.* 592.—Am. trop.
odora, *Baker, in Journ. Bot.* xvii. (1879) 226.—Am. trop.
organensis, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 116 = floribunda.
ornata, *Baker, in Journ. Bot.* xvii. (1879) 163.—Am. trop.
Ortgiesii, *Baker, l. c.* 236.—Am. trop.

AECHMEA:—
paniculata, *Ruiz & Pav. Fl. Per.* iii. 37.—Peruv.
paniculigera, *Griseb. Fl. Brit. W. Ind.* 593.—Ind. occ
parviflora, *Baker, in Journ. Bot.* xvii. (1879) 167.—Bras.
patentissima, *Baker, l. c.* 227.—Am. trop.
pectinata, *Baker, l. c.* 233.—Am. trop.
petropolitana, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 116.—Bras.
Pinellana, *Baker, in Journ. Bot.* xvii. (1879) 232.—Am. trop.
platynema, *Baker, l. c.* 166.—Bras.
polycephala, *Baker, l. c.* 164.—Ins. Jamaic.
pubescens, *Baker, l. c.* 185.—Nicarag.
pyramidalis, *Benth. in Bot. Voy. Sulph.* 178.—Mexico.
ramosa, *Mart. ex Schult. f. Syst.* vii. II. 1272.—Bras.
regularis, *Baker, in Journ. Bot.* xvii. (1879) 229.—Am. trop.
Schiedeana, *Schlecht. in Linnaea*, xviii. (1844) 487.—Mexico.
setigera, *Mart. ex Schult. f. Syst.* vii. II. 1273.—Bras.
spectabilis, *Brongn. ex Houllet, in Rev. Hortic.* (1875) 311.—Guatemala.
sphaerocephala, *Baker, in Journ. Bot.* xvii. (1879) 162.—Am. trop.
spicata, *Mart. ex Schult. f. Syst.* vii. II. 1273.—Bras.
suavcolens, *Knowl. & Westc. Flor. Cab.* iii. (1840) 177.—Bras.
subinermis, *Baker, in Journ. Bot.* xvii. (1879) 228.—Am. trop.
tillandsioides, *Baker, l. c.* 184.—Bras.
Veitchii, *Baker, in Bot. Mag.* t. 6329.—N. Granat.
viridis, *Baker, in Journ. Bot.* xvii. (1879) 235.—Am. austr.
vriesoides, Baker, l. c. 184 = tillandsioides.
Wrilbackea, Carr. in Rev. Hortic. (1871) 171 = seq
Weilbachii, *F. Didr. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 376.—Bras.
Wrightii, *Baker, in Journ. Bot.* xvii. (1879) 163.—Cuba.

AECHMOLEPIS, Decne. in DC. Prod. viii. (1844) 498 *ASCLEPIADEAE*, Benth. & Hook. f. ii. 741. rosmarini olia, *Decne. l. c.*—Afr. trop.

AECHMOPHORA, Spreng. ex Stend. Nom. ed. II. i. (1840) 29 = **Bromus**, Linn. (Gramin.).
arduensis, Spreng. ex Steud. l. c. = B. arduensis.

AEDEMONE, Kotschy, in Oestr. Bot. Zeitschr. viii. (1858) 116 = **Herminiera**, Guill. & Perr. (Legumin.).
mirabilis, Kotschy, l. c. = H. Elaphroxylon.

AEDMANNIA, Spach, in Orb. Dict. i. (1841) 139, err. typ. = OEDMANNIA, Thunb. (Legum.).

AEDULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. 4. i, nomen.

AEGIPHILA, Sw. Prod. Veg. Ind. Occ (1788) 31, err. typ. = **Aegiphila**, Jacq.

AEERIDIUM, Salisb. in Trans. Hort. Soc. i. (1812) 295 = **Aërides**, Lour. (Orchid.).
odorum, Salisb. l. c. = Aerides odoratum.

AEGELATIS, Roxb. Fl. Ind. ii. 111 (1832) err. typ. = **Aegialitis**, R. Br. (Plumbag.).

AEGIALEA, Klotzsch, in Walp. Ann. ii. (1852) 1118 = **Pieris**, D. Don (Ericac.).

AEGIALINA, Schult. Mant. ii. (1824) 13, 232 = **Koelezia**, Pers. (Gramin.).
tenuis, Schult. l. c. = K. villosa.

AEGIALINITES, Presl, Bot. Bemerk. (1844) 103 = **Aegialitis**, R. Br. (Plumbag.).
annulata, Presl, l. c. = Aegialitis annulata.
rotundifolia, Presl, l. c. = Aegialitis rotundifolia.

AEGIALITIS, R. Br. Prod. (1810) 426. *PLUMBAGINEAE*, Benth. & Hook. f. ii. 624.
AEGIALINITES, Presl, Bot. Bemerk. (1844) 103.
annulata, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 217 = rotundifolia.
annulata, *R. Br. Prod.* i. 426.—Austral.
rotundifolia, *Roxb. Fl. Ind.* ii. 111.—Ind. or.

AEGIALITIS, Trin. Fund. Agrost. 127. t. 9. (1820) = **Koeleria**, Pers. (Gramin.).
tenuis, Trin. l. c. = K. villosa.

AEGIALOPHILA, Boiss. & Heldr. Diagn. Ser. I. x. (1846) 105 = **Centaurea**, Linn. (Compos.).
cretica, Boiss. & Heldr. l. c. 106 = C. pumila.
humilis, Boiss. l. c. 105 = C. pumila.

AEGIANILITES, C. B. Clarke, in Hook. f. *Fl. Brit. Ind.* iii. 479 (1882), err. typ. = **Aegialitis**, K. Br.

AEGIATILIS, Griff. Not. iv. 207 (1854), err. typ. = praec.

AEGICERAS, Gaertn. Fruct. i. 216. t. 46 (1788). *MYRSINEAE*, Benth. & Hook. f. ii. 648.
MALASPINAEA, Presl, Reliq. Haenk. ii. 68. t. 61 (1830).
corniculatum, Blanco, Fl Filip. ed. 1. 79 = majus.
ferreum, Blume, Bijdr. 693 = majus
floridum, Roem. & Schult. Syst. iv. 512 = majus.
fragrans, Kon. in Kon. & Sims, Ann. Bot. i. (1805) 181 = majus.
majus, *Gaertn. Fruct.* i. 216 t. 46.—Geront. trop.
Malaspinaea, A. DC. Prod viii. 143 = majus.
minus, A. DC. l. c. 412 = majus.
minus, *Gaertn. Fruct.* i. 216. t 46.—Ind. or.
nigricans, A. Rich. Voy. Astrol. Bot. ii. 57. t. 21 = majus.
obcordatum, Steud. Nom. ed. II. i. 29 = majus.
obovatum, Blume, Bijdr. 693 = majus.

AEGICON, Adans. Fam. ii. 36 (1763) = **Trttioum**, Linn. (Gramin.).

AEGILOPS, Linn. Coroll. Gen. 20 (1737) = **Trttioum**, Linn. (Gramin.).
agropyroides, Godr. in Mém. Acad. Montp (Sect. Médic.) i. (1853) 456.—Syria.
aromatica, Walt. Fl. Carol. 249 = Ctenium americanum.
Aucheri, Boiss. Diagn. Ser. I. v. 74 = Agropyron Aucheri.
Augeri, Steud. Syn. Pl. Gram. 356 (err. typ.) = praec.
bicornis, Jaub. & Spach, Illustr. Pl. Or. iv. 11 = T. bicorne.
biuncialis, Vis. Fl. Dalm. 1.—Europ.
brachyathera, Pomel, Nouv. Mat. Fl. Atl. 389 — Afr.bor.
caudata, Bertol. Fl. Ital. vi. 622 = Agropyron Aucheri.
caudata, C. A. Mey. Verz. Pfl. Cauc. 26 = T. Aegylops.
caudata, Bab. Ellencho, 98 ; Griseb. Spicil. Fl. Rumel. ii. 435 = T. cylindricum.
caudata, Linn. Sp. Pl. 1051 = Agropyron villosum ?
ciliaris, Koen. ex Roem.& Schult. Syst. ii. 772.—Ind. or.
comosa, Sibth. & Sm. Fl. Graec. 75. t. 94.—Graecia.
cornuta, Steud. Syn. Pl. Gram. 356.—Graecia.
crassa, Boiss. Diagn. Ser. I. vii. 123.—Orient.
crithodium, Steud. Syn. Pl. Gram. 355 = T. monococcum.
cylindrica, C. A. Mey. Verz. Pfl. Cauc. 26 = T. Aegylops ?
cylindrica, Host, Gram. Austr. ii. 6. t. 7 = T. cylindricum.
cylindrica, Schur, Enum. Pl. Transs. 613 = T. ovatum.
cylindrica, Sibth. & Sm. Fl. Graec. t. 95 = T. caudatum.
divaricata, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 129 = T. ovatum.
echinata, Presl, Gram. Sic. 47 = T. triunciale.
Echinus, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 456.—Gallia ?
elongata, Lam. Fl. Fr. iii. 632 = T. trinnciale.
erigens, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 131 = T. ovatum.
erratica, Jord. & Fourr. l. c. 130 = T. ovatum.
exaltata, Linn. Mant. ii. App. 575 = Ophiurus corymbosus.
fuvoiatilis, Blanco, Fl. Filip. ed. 1. 47.—Ins. Philipp.
fragilis, Parl. Fl. Ital. i. 515 = T. fragile.
graniculata, Roth, Catalect. i. 121 = T. ovatum.
hordeiformis, Steud. Syn. Pl. Gram. 354.—Mesopotam.
Hystrix, Nutt. Gen. Am. i. 86.—Am. bor.
incurva, Linn. Sp. Pl. 1051 = Lepturus incurvatus.
incurvata, Linn. Sp. Pl. ed. II. 1490 = praec.
Kotschyi, Boiss. Diagn. Ser. I. vii. 129 = T. triunciale.
Lorentii, Steud. Syn. Pl. Gram. 354 = *Lorentii*.
loliacea, Jaub. & Spach, Illustr. Pl. Or. iv. 23 = *muticа*.

AEGILOPS:—
longearistata, Steud. Syn. Pl. Gram. 356.—Guiana.
Lorentii, Hochst. in Lorent, Wanderungen, 326 = T. ovatum.
macrochaeta, Shuttl. & Huet, ex Duval, in Bull. Soc. Bot. Fr. xvi. (1869) 384 = T. ovatum.
macrura, Jaub. & Spach, Illustr. Pl. Or. iv. 21 = Agropyron Aucheri.
microstachys, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 132 = T. ovatum.
muricata, Retz. Obs. ii. 27 = Ischaemum pectinatum.
mutica, Boiss. Diagn. Ser. I. v. 73.—Orient.
neglecta, Req. ex Bertol. Fl. Ital. i. 787 = T. triaristatum.
nigricans, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 128 = T. ovatum.
Notarisii, Clem. in Mem Acc. Torin. Ser. II. xvi. (1857) 335 = T. ovatum.
ovata, Linn. Sp. Pl. 1050 = T. ovatum.
ovata, Roth, in Usteri, Ann. iv. (1793) 41 = T. triaristatum.
parvula, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 131 = T. ovatum.
persica, Boiss. Diagn. Ser. I. vii. 129 = T. candatum.
platyathera, Jaub. & Spach, Illustr. Pl. Or. iv. 17.—Mesopotam.
procera, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 129 = T. ovatum.
pubiglumis, Jord. & Fourr. l. c. 131 = T. ovatum.
saccharina, Walt. Fl. Carol. 249 = Eleusine aegyptiaca.
sanguinea, Steud. Syn. Pl. Gram. 354.—Persia.
scula, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 129 = T. ovatum.
*spelta*e*formis*, Jord. in Ann. Sc. Nat. Sér. IV. iv. (1855) 313 = T. ovatum.
speltoides, Tausch, in Flora, xxxix. (1837) 109 = T. bicorne.
squarrosa, Willd. Sp. Pl. iv. 944 = T. venticosum.
squarrosa, Cav. Ic. t. 90. f. 2 = *platyathera*.
squarrosa, DC. ex Steud. Nom. ed. II. i. 29 = T. cylindricum.
squarrosa, Linn. Sp. Pl. 1051 = T. Aegylops.
subulata, Pomel, Nouv. Mat. Fl. Atl. 388.—Afr. bor
Tauschii, Coss. Notes Crit. 69 = T. Aegylops.
triaristata, Req. ex Bertol. Fl. Ital. i. 789 = T. triunciale.
triaristata, Willd. Sp. Pl. iv. 943 = T. triaristatum.
triquetrales, Jaub. & Spach, Illustr. Pl. Or. ii. 121. t. 200.—Asia.
tritcoides x, Req. ex Bertol. Fl. Ital. i. 788.—Europ.
truncatalis, Linn. Sp. Pl. 1051 = T. triunciale.
truncata, Ledeb. ex Trautv. in Act. Hort. Petrop. ix. (1884) 312, nomen.—Rossia.
uniaristata, Steud. Syn. Pl. Gram. 354.—Graecia.
unuaristata, Vis. Fl. Dalm. iii. 845.—Europ.
vagans, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 130 = T. ovatum.
ventricosa, Tausch, in Flora, xxxix. (1837) 108 = T. ventricosum.
virescens, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 130 = T. ovatum.

AEGINETIA, Cav. in Anal. Clenc. Nat. iii. (1801) 139 = **Bouvardia**, Salisb. (Rubiac.).
capitata, R. Grah. in Edinb. N. Phil. Journ. (Jan. 1839) 389 = Coccocipselum capituliflorum.
longiflora, Cav. in Anal. Hist. Nat. iii. (1801) 130 = Houstonia longiflora.
multiflora, Cav. l. c. = Bouvardia Cavanillesii.

AEGINETIA, Linn. Syst. ed. I. (1735). *OROBANCHACEAE*, Benth. & Hook. f. ii. 982.
CENTRONIA, Blume, Bijdr. 776 (1826).
CENTRONOTA, DC. ex Meisen. Gen. 303 (218) (1840); et Prod. ix. 286 (1845).
GASPARINIA, Endl. Gen. 1408 (1840).
TRONICENA, Steud. Nom. ed. II. ii. 721 (1841).
abbreviata, Buch.-Ham. ex Benth. Scroph. Ind. 55 = pedunculata.
aculis, Walp. Rep. iii. 481 = pedunculata.
Centronia, *Miq. Fl. Ind. Bat.* ii. 713.—Java.
coccinea, Juss. ex Walp. Rep. iii. 481 = Phelypaea coccinea.
Indica, *Linn. Sp. Pl.* 633 ; *Roxb. Pl. Corom.* i. 63. t. 91.—As. trop.
japonica, Steb. & Zucc. in Abh. Math. Akad. Muench. iv. iii. (1845) 341 = indica.
pedunculata, *Wall. Pl. As. Rar.* iii. 13. t. 219.—As. trop.

AEGIPHILA, Jacq. Stirp. Am. t. 16 (1763) ; Obs. Bot. ii. 3. t. 27 (1767). *VERBENACEAE*, Benth. & Hook. f. ii. 1151.
AMERINA, DC. in Meisen. Gen. 278 (186) (1840).
BRUECKIA, Klotzsch & Karst. Ausw. Gew. Venez. 21. t. 10 (1848).
MANABEA, Aubl. Pl. Gui. i. 61. t. 23, 25 (1775).
OMPHALOCOCCA, Willd. ex Roem. & Schult. Syst. iii. Mant. 10 (1827).
arborescens, *Vahl, Eclog.* i. 15. t. 10.—Am. austr.
aurea, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 213.—Cuba.
Bertersana, *Schau. in DC. Prod.* xi. 654.—N. Granat.
brachiata, *Schlecht. in Linnaea*, vi. (1831) 371.—Mexico.
brachiata, Vell. Fl. Flum 88 ; t. 93 = triantha.
cascelliformis, *Schau. in DC. Prod.* xi. 651.—Bras.
cestrifolia, *Gardn. in Hook. Lond. Journ. Bot* v. (1846) 184.—Bras.
cordata, *Poepp. ex DC. Prod* xi. 650.—Peruv
cornifolia, *Kunth, in Abh. Akad. Berl.* (1831) 215.—Bras.
cuspidata, *Mart. ex DC. Prod.* xi. 653.—Bras.
Deppeana, Steud. Nom. ed. II. i. 29 = brachiata.
diffusa, *Andr. Bot. Rep.* ix. t. 578.—Ind. occ.
dumosa, Salisb. Prod. 67 = martinicensis.
elata, *Sw. Prod. Veg Ind. Occ* 31.—Am. trop.
filipes, *Mart. & Schau. in DC. Prod.* xi. 652.—Bras.
flummensis, *Vell. Fl. Flum.* 38 ; t. 95.—Bras.
foetida, *Sw. Prod. Veg. Ind Occ.* 82.—Jamaic.
glabra, Lam. Illustr. t. 70. f. 1 = martinicensis.
glomerata, *Benth. Bot. Voy. Sulph.* 154.—Ecuador.
grandiflora, Hook. Bot. Mag. t. 4230 = Clerodendron grandiflorum.
grandifolia, Walp. Rep iv. 121 = magnifolia.
graveolens, *Mart. & Schau. in DC. Prod.* xi. 651.—Bras.
hirta, *Casar. Nov. Stirp. Bras. Decad.* 47.—Bras.
Humboldtii, *Schau. in DC. Prod.* xi. 652.—Venezuela.
incana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 218.—Cuba.
inflexa, Vell. Fl. Flum. 38 ; i. t. 96. (Rubiacea, sec. Schau.).—Bras.
integrifolia, *Jacq. Hist. Stirp. Am.* 15. t. 178. f. 7. —N. Granat.
laeta, *H. B. & K. Nov Gen. et Sp.* ii. 249.—N. Granat.
lacvigata, *Juss. in Ann. Mus. Par.* vii. (1806) 76.—Ind. occ.
laevis, *Poepp. ex Walp. Rep* iv. 110.—Bras.
laevis, *Willd. Sp. Pl.* i. 617.—Guiana.
lanuginosa, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 184 = obovata.
latifolia, Willd. ex Walp. Rep. iv. 120 = Mutisii.
laxiflora, *Benth. in Ann. Nat. Hist.* ii. (1839) 449.—Guiana.
Lhotskiana, *Cham. in Linnaea*, vii. (1832) 113.—Bras.
longifolia, *Turcz in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 218.—N. Granat.
Laschnati, *Schau. in DC. Prod.* xi. 651.—Bras.
lutea, Lam. Illustr. t. 70. f. 3 = laevis.
macrophylla, Desf. Cat. Hort. Paris. ed. III. 391 = magnifolia.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* ii. 251.—Am. austr.
macrophylla, Sieber, ex Schau. in DC. Prod. xi. 655 (= Rubiacea, sec. Schau.).—Ins. Trinit.
magnifolia, *Steud. Nom.* ed. II. i. 29.—Ind. occ.
Manabaea, Sw. Prod. Veg. Ind. Occ. i. 286 = laevis.
martinicensis, *Jacq. Obs. Bot.* ii. 3.—Ins. Martinic. mediterranea, *Vell. Fl. Flum.* 88 ; i. t. 94.—Bras.
membranacea, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 219.—Am. trop.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 250. t. 130.—Am. austr.
multiflora, *Ruiz & Pav. Fl. Per.* i. 50. t. 74.—Peruv.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* ii. 250. t. 131.—Am. austr.
Nuxia, Willd. Sp. Pl. i. 616 = Nuxia verticillata.
obducta, *Vell. Fl. Flum.* 89 ; i. t. 97.—Bras.
obovata, *Andr. Bot. Rep.* ix. 578.—Ind. occ.
olefera, Casar. Nov. Stirp. Bras. Decad. 47 = fluminensis.
peruviana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 219.—Peruv.
pubescens, Willd. ex Steud. Nom. ed. II. i. 29 = salutaris.
punctata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 219.—Jamaic.

AEGIPHILA:—
racemosa, Vell. Fl. Flum. 37 ; i. t. 36 = cupidista.
Riedeliana, *Schau. in DC. Prod.* xi. 649.—Bras.
salutaris, *H. B. & K. Nov. Gen. et Sp.* ii. 249.—Am. austr.
Selloviana, *Cham. in Linnaea*, vii. (1832) 111.—Bras.
serrata, Vell. Fl. Flum. 38 ; t. t. 92 = graveolens.
splendens, *Schau. in DC. Prod.* xi. 648.—Bras.
stipulata, Vell. Fl. Flum. 37 ; i. t. 90 (Rubiacea, sec. Schau.).—Bras.
straminea, *Hoffmgg. Vers Pfl. Nachtr.* iii. 18.—Bras.
tomentosa, *Cham. in Linnaea*, vii. (1832) 110.—Bras.
trianthe, *Schau. in DC. Prod.* xi. 650.—Bras.
trifida, *Sw. Prod. Veg. ind. Occ.* 32.—Jamaic.
umbellata, Vell. Fl. Flum. 37 ; i. t. 89. (Rubiacea, sec. Schau.).—Bras.
verrucosa, *Schau. in DC. Prod.* xi. 650.—N. Granat.
verticillata, Vell. Fl. Flum. 37 ; i. t. 91 = tomentosa.
viburnifolia, *Pers. in Ann. Mus. Par.* vii. (1806) 76. —Ins. Philipp.
villosa, *Vahl, Eclog.* i 16.—Guiana.
virgata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 220.—Bras.
vitelliniflora, *Klotzsch, ex Walp Rep.* iv. 128.—Hab.?

AEGIPHYLA, Stend. Nom. ed. II. i. 29 = **Aegiphila**, Jacq.

AEGLE, Correa, in Trans. Linn. Soc. v. (1800) 22. *RUTACEAE*, Benth. & Hook. f. i. 306.
BELOU, Adans Fam. ii. 408 (1763).
decandra, *F. Villar., in Blanco, Fl. Philipp* ed III. *Nov. App.* 38.—Ins Philipp.
Marmelos, *Correa, in Trans. Linn. Soc.* v. (1800) 233.—Ind. or.
sepiaria, DC. Prod. i. 538 = Citrus trifoliata.

AEGLE, Dulac, Fl. Hautes-Pyr. 43 (1867) = **Poidonia**, Koen. (Naiad.).
fragilis, Dulac, l. c. = P fragilis.

AEGOCHLOA, Benth. in Bot Reg. sub t. 1622 (1833) = **Gilia**, Ruiz & Pav. (Polemon.).
atractyloides, Benth. l. c. = G atractyloides.
cotulaefolia, Benth. l. c. = G cotulaefolia.
eryngioides, Benth. l. c. = G. Navaretia.
intertexta, Benth. l. c. = G. intertexta.
pubescens, Benth. l. c. = G. cotulaefolia.
pungens, Benth. l. c. = G. squarrosa.
Torreyi, G. Don, Gen. Syst. iv. 246 = G. pungens.

AEGOMARATHRUM, Steud. Nom. ed. II. i. 30 (1840) = **Cachrys**, Tourn. (Umbell.).
crispum, Steud. l. c. = C. crispa.
pterochlaena, Steud. l. c. = C. sicula.
siculum, Mor. ex Nym. Consp. 314 = Hippomarathrum Bocconi.

AEGONYCHON, S. F. Gray, Nat. Arr. Brit. Pl. ii. 354 (1821) = **Lithospermum**, Linn. (Borag.).
arvense, S. F. Gray, l. c. = L. arvense.
repens, S. F. Gray, l. c. = L. purpureo-caeruleum.

AEGOPICRON, Gieseke. Prael. 474 (1792) = AEGOPRICON, Linn. f. = **Maprounea**, Aubl. (Euphorb.).

AEGOPODION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = seq.

AEGOPODIUM, Knaut, ex Linn. Syst. ed. I. (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 893.
PODAGRARIA, Hall. Enum. Stirp. Helv. ii. 427 (1742).
alpestre, *Ledeb. Fl. Alt.* i. 354.—Sibir. altaic.
angelicaefolium, Salisb. Prod. 169 = Ae. Podagraria.
Carum, Wibel, Prim. Fl. Werth. 199 = Carum Carvi.
involucratum, Orphan. ex Boiss. Fl. Orient. ii. 923 = Physospermum aegopodioides.
latifolium, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 711 = Ae. Podagraria.
Podagraria, *Linn. Sp. Pl.* 265.—Europ.; As. bor.
ternatum, Gilib. Fl. Lituan. ii. 41 = Ae. Podagraria.

AEGOPOGON, Beauv. Agrost. 122. t. 22. f. 4. (1812) partim = **Amphipogon**, R. Br. (Gramin.).
avenaceus, Beauv. l. c. = Am. strictus.
debilis, Beauv. l. c. = Am. debilis.
lagurioides, Beauv. l. c. = Am. lagunoides.

AEGOPOGON:—
pusillus, Spreng. Syst. i. 270 = Chloris cruciata.
strictus, Beauv. Agrost. 122 = Am. strictus.
tenuinatus, Beauv. l. c. = Am. turbinatus.

AEGOPOGON, Beauv. Agrost. 122 (1812) partim ; Humb. & Bonpl in Willd. Sp. Pl. iv. 899 (1805). *GRAMINEAE*, Benth. & Hook. f. iii. 1121.
HYMENOTHECIUM, Lag. Gen. et Sp. Nov. 4 (1816).
SCHELLINGIA, Steud. in Flora, xxxii. (1850) 231. t. 1.
bryophilus, *Doell, in Mart. Fl. Bras.* ii. iii. 289.—Bras.
cenchroides, *Humb & Bonpl ex Willd. Sp. Pl.* iv. 899 —Venezuela.
gemmiflorus, *Humb. & Bonpl. Nov. Gen. et Sp.* iv. 133. t. 43.—Am trop.
multistrius, Steud. Syn. Pl. Gram. 146 = cenchroides.
quinquestrius, Koem & Schult. Syst. ii. 805 = cenchroides.
pusillus, Beauv. Agrost. 122 = geminiflorus.
rigidissimus, *Steud. Syn. Pl. Gram.* 146.—Texas.
setifer, Nees, in Linnaea, xix (1847) 691 = cenchroides.
submuticus, *Rupr. ex Trin. in Mém Acad. Pétersb Sét.* VI. v. ii (1840) 3.—Hab.?
tenellus, Trin. Gram. Unifl. 164 = geminiflorus.
trisetus, Koem. & Schult. Syst. ii. 805 = cenchroides.
unisetus, Koem. & Schult. l. c. = geminiflorus.

AEGOPORDON, Boiss. Diagn. Ser. 1. vi. (1845) 112 = **Jurinea**, Cass. (Compos.).
berardioides, Boiss. l. c. 118.—Persia.

AEGOPRICON, Linn. f. Suppl. 418 (1789) = seq.

AEGOPRICUM, Linn. Pl. Surinam. 16 (1775) , (Amoen. Acad. viii. 208 = **Maprounea**, Aubl. (Euphorb.).
brasiliense, Lam. l. c. = M. guianensis.

AEGOSERIS, Steud. Nom ed. II. i. (1840) 30 = **Crepis**, Linn. (Compos.).
leontodontoides, Fourr. in Ann. Soc Linn. Lyon, N. S. xvii. (1869) 103 = C. leontodontoides.
setosa, Fourr. l. c. = C. setosa.
suffreniana, Fourr. l. c. = C. suffreniana.

AEGOTOXICON, Molina, Sagg Chile, ed. II. 298 (1810) = **Aextoxicon**, Ruiz & Pav. (Euphorb.).

AEGOTOXICUM, Endl. Encb. 589 (1841) = praec.

AEGYLOPS, Honck. Syn i. 484 (1792) err. typ. = AEGILOPS, Linn.= **Tritticum**, Linn. (Gramin.).

AEIPHANES, Spreng. Anleit. ii. ii. 876 (1818) err. typ. = AIPHANES, Willd. = **Martinesia**, Ruiz & Pav. (Palm.).

AELUROPUS, Trin. Fund. Agrost. 148 (1820). *GRAMINEAE*, Benth. & Hook f. iii. 1193.
CALOTHECA, Spreng Anleit. ii i. 167 (1817) ; Syst. i. 347 (1825).
CHAMAEDACTYLUS, T. Nees, Gen. Fl. Germ. Monocot. i. n. 66 (1840?).
arabicus, *Steud. Nom.* ed II. i. 30.—Aegypt.; Abyss.
bombycinus, Figar. & De Not. in Mem. Acc. Torin. Ser II. xii. (1852) 258 = arabicus.
brevifolius, Nees, ex Steud. Nom. ed. II. i. 30 = pubescens.
concinnus, Figar. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 257 = laevis.
grandiflorus, *Munro, in Cat. Kew Distrib. Griffith*, &c. 97, nomen.—Afghan.
hirsutus, *Munro, l. c.* nomen.—Ind. or.
intermedius, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) ii. 292.—As. temp.
laevis, *Trin. Fund. Agrost.* 148.—Orosia.
litoralis, Parl. Fl. Ital. i. 461 = laevis.
mucronatus, Aschers. in Schweinf. Beitr. Fl. Aethiop. 297 = arabicus
nitiacus, *Steud. Nom.* ed. II. i. 30.—Aegypt.
pubescens, *Trin. ex Steud. Nom.* ed. II. i. 30.— Ind. or.
pungens, C. Koch, in Linnaea, xxi. (1848) 408 = laevis.
repens, Parl. Fl. Ital. i. 462 = villosus.
sinaicus, Figar. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 257 = laevis.
Smithii, *Steud. Nom.* ed. II. i 30.—Ins. Canar.
villosus, *Trin. Fund. Agrost.* 148.—Afr. bor.; Graecia.

AENOTHERA, Lam. Encyc. iv. 550 (1796), err. typ. = **Oenothera**, Linn. (Onagr.).

AEOLANTHUS, Mart. Amoen. Bot. Monac. 4. t. 2 (1825). *LABIATAE*, Benth. & Hook. f. ii. 1176.
ORCILANTHUS, E. Mey. Comm. Pl. Afr. Austr. ii. 280 (1837).
abyssinicus, *Hochst. ex Benth. in DC. Prod.* xii. 60.—Abyss.
ambustus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 138.—Afr. trop.
heliotropioides, *Oliver, l. c.* 137.—Afr. trop.
Livingstones, *C. Koch, Berl. Wochenschr.* iii. (1860) 188.—Afr. trop.
parvifolius, *Benth. in DC. Prod.* xii. 80.—Afr. austr.
pinnatifidus, *Hochst. ex Benth. l. c.*—Abyss.
pubescens, *Benth l. c.*—Afr trop.
repens, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 137. —Afr. trop.
suaveolens, G. Don, in Loud. Hort. Brit. 247 = suavis.
suavis, *Mart. Amoen. Bot. Monac.* 4. t. 2.—Bras. ; Afr.?
zanzibaricus, *S. Moore, in Journ. Bot.* xv. (1877) 68. —Afr. trop.

AEOLLANTHUS, Spreng. Syst. ii. 750 (1825) = praec.

AEONIA, Lindl. Bot. Reg. sub t. 817 (1824). *ORCHIDEAE*, Benth. & Hook. f. iii. 584.
BRACHYSTEPIS, Thou. ex Pritzel, Ind. Ic. 162 (1855).
ORONIA, Lindl Orch. Sced. 14, err. typ. (1826).
alata, *A. Rich. in Ann. Sc. Nat. Sér.* II. xvi. (1841) t 11.—Ind. or.
Auberti, *Lindl. Bot. Reg* sub t. 817.—Madag.
brachystachya, *Lindl. Gen. et Sp. Orch.* 245.—Madag
macrostachya, *Lindl. l. c.* 245.—Madag.
polystachya, *Benth. & Hook. f. Gen* iii. 584.— Madag.
roses, *Ridl. in Journ. Linn. Soc. xxi.* (1885) 496.— Madag.

AEONIUM, Webb. & Berth. Phyt. Canar. i. 184 (1836), t. 28 = **Sempervivum**, Linn. (Crassul.).
arboreum, Webb & Berth. l. c. 185 = S. arboreum.
balsamiferum, Webb & Berth l. c. 192 = S. balsamiferum.
barbatum, Webb & Berth. l. c. 188 = S. pilliferum.
Berthelotianum, Boiss. in Bonplandia, vii. (1859) 339
Bethencortianum, Webb, ex Bolle, l. c. 243.—Ins. Canar.
caespitosum, Webb & Berth. Phyt. Canar. i. 191 = S. caespitosum.
canariense, Webb & Berth. l. c. i. 196 = S. canariense.
Castello-Paivae, C. Bolle, in Bonplandia, vii. (1859) 240.—Ins. Canar.
ciliatum, Webb & Berth. Phyt. Canar. i. 195 = S. ciliatum.
cruentum, Webb & Berth. l. c. 186 = S. cruentum.
cuneatum, Webb & Berth. l. c. 197 = S. cuneatum.
decorum, Webb, ex C. Bolle, in Bonplandia, vii. (1859) 241.—Ins. Canar.
fruticosum, Webb & Berth. Phyt. Canar. i. 198 = S. frutescens.
glanduliosum, Webb & Berth. l. c. 185 = S. glandulosum.
glutinosum, Webb & Berth. l. c. 185 = S. glutinosum.
Goochiae, Webb. & Berth. l. c. 190 = S. Goochiae.
gorgoneum, J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 258.—Ins. Cap. Verd.
Haworthii, Webb & Berth. Phyt. Canar. i. 193 = S. Haworthii.
holochrysum, Webb & Berth. l. c. 194 = S. holochrysum.
laxum, Webb & Berth. l. c. i. 198 = S. laxum.
leucoblepharum, Webb, ex A. Rich. Tent. Fl. Abyss. i. 314 = S. chrysanthum.
Lindleyi, Webb & Berth. Phyt. Canar. i. 189 = S. villosum.
manriqueorum, C. Bolle, in Bonplandia, vii. (1859) 241.—Ins. Canar.
Meyerheimii, C. Bolle, l. c. 239.—Ins. Mader.
Saundersii, C. Bolle, l. c. 241.—Ins. Canar.
Smithii, Webb & Berth. Phyt. Canar. i. 187 = S. Smithii.
stripicladum, Webb & Berth. l. c. = S. strepsicladum.
subulaeforme, Webb & Berth. l. c. 185 = S. tabulaeforme.

AEONIUM :—
undulatum, Webb & Berth. Phyt. Canar. i. 197 = S. undulatum.
urbicum, Webb & Berth. l. c. 194 = S. urbicum.
vicastrum, Webb, ex C. Bolle, in Bonplandia, vii. (1859) 241.—Ins. Canar.
Webbii, C. Bolle, l. c. 238.—Ins. Canar.
Youngianum, Webb & Berth. Phyt. Canar. i. 197 = S. Youngianum.

AERA, Aschers. Fl. Brandenb. 830 (1864) = **Aira**, Linn. (Gramin.).

AERANGIS, Reichb. f. in Flora, xlviii. (1865) 190 = **Angraecum**, Thou. (Orchid.).
flabellifolia, Reichb. f. l. c. 191.—Afr. trop.
Kotschyi, Reichb. f. Otia Bot. Hamb. 78 = Angr. Kotschyi.

AERANTHES, Lindl. Bot. Reg. t. 817 (1824). *ORCHIDEAE*, iii. 576.
Arachnitis, *Lindl. Bot. Reg.* t. 817.—Ins. Mascar.
filiformis, Griseb. Fl. Brit. W. Ind. 625.—Ind. occ. grandiflorus, *Lindl. Bot. Reg.* t. 817.—Madag.
grandiflorus, Lindl. Gen. et Sp Orch. 248, pro parte = A. Arachnitis.
jamaicensis, Reichb. f. ex Griseb. Fl. Brit. W. Ind. 625 = Angraecum micranthum.
sesquipedalis, Lindl. Bot. Reg. sub t. 817 = Angraecum sesquipedale.

AERANTHUS, Bartl. Ord. 57 (1830) = praec.

AERANTHUS, Reichb. f. in Walp. Ann. vi. 899 (1861) = **Myrtaeidium**, Lindl. (Orchid.), &c.
aciculatus, Reichb. f. & Warm. in Otia Bot. Hamb. 91.—Bras.
arachnanthus, Reichb. f. in Flora, lxviii. (1885) 539.—Ins. Comor.
brachystachyus, Boj. Hort. Maurit. 314 = Aeonia brachystachya.
Calceolus, S. Moore, in Baker, Fl. Maurit. 358 = Angraecum Calceolus.
comorensis, Reichb. f. in Flora, lxviii. (1886) 540.—Ins. Comor.
Curnowianus, Reichb. f. in Gard. Chron. N. S. xix. (1883) 1. 306.—Madag.
dentiens, Reichb. f. in Flora, lxviii. (1885) 381.—Ins. Comor.
distichus, Reichb. f. in Walp. Ann. vi. 901 = Angraecum distichum.
erythrophlebius, Reichb. f. in Flora, xlviii. (1865) 190.—Afr. trop.
expansus, S. Moore, in Baker, Fl. Maurit. 351 = Angraecum expansum.
Fasciola, Reichb. f. in Walp. Ann. vi. 902 = Angraecum Fasciola.
filicornis, Reichb. f. l. c. 900 = M. filicorne.
fragrans, Reichb. f. l. c. 899 = Angraecum fragrans.
funalis, Reichb. f. l. c. 902 = Angraecum funale.
Gerrardi, Reichb. f. in Flora, l. (1867) 117.—Afr. austr.
Gladiator, Reichb. f. l. c. lxviii. (1885) 539.—Ins. Comor.
gladiflorus, Reichb. f. in Walp. Ann. vi. 900 = Angraecum gladiflolium.
gracilis, Reichb. f. in Flora, l. (1867) 117 = M. gracile.
Grandidieranus, Reichb. f. l. c. lxviii. (1885) 381.—Ins. Comor.
Guyonianus, Reichb. f. l. c. xlviii. (1865) 190 = Angraecum Guyonianum.
hippolulus, Reichb. f. in Linnaea, xli. (1877) 31.—N. Granat.
hymenanthus, Griseb. Cat. Fl. Cub. 264 = Dendrophylax hymenanthus.
intermedius, Reichb. f. & Warm. in Otia Bot. Hamb. 91.—Bras.
jamaicensis, Reichb. f. in Walp. Ann. vi. 901 = Angraecum micranthum.
Lanabergii, Reichb. f. l. c. = Angraecum Lanabergii.
Lehmanni, Reichb. f. Otia Bot. Hamb. 10.—Hab.?
Leonii, Reichb. in Flora, lxviii. (1885) 380 = seq.
Leonis, Reichb. f. in Gard. Chron. N. S. xxiii. (1885) 726 = Angraecum Humblotii.
Lindenii, Reichb. f. in Walp. Ann. vi. 902 = Angraecum Lindenii.
macrostachys, Reichb. f. l. c. 900 = Aeonia macrostachya.

AERANTHUS :—
Meirax, Reichb. f. in Flora, lxviii. (1885) 540.—Afr. austr.
micranthus, Reichb. f. in Walp. Ann. vi. 90 = Angraecum micranthum.
Monteverdii, Reichb. f. in Flora, xlviii. (1865) 279.—Hab.?
muscula, Reichb. f. l. c. 190.—Afr. trop.
negiectus, Reichb. f. & Warm. in Otia Bot. Hamb. 91.—Bras.
organensis, Reichb. f. in Walp. Ann. vi. 901 = Angraecum organense.
ornithorrhynchus, Reichb. f. l. c. 903 = Angraecum ornithorrhynchum.
packyrhynchus, Reichb. f. in Flora, xlviii. (1865) 279.—Ind. occ.
parahybunensis, Rodrig. Orch. Nov. ii. 245.—Bras.
pectinatus, Reichb. f. in Walp. Ann. vi. 900 = Angraecum pectinatum.
phalaenophorus, Reichb. f. in Flora, lxviii. (1885) 539.—Ins. Comor.
polystachyus, Reichb f. in Walp. Ann. vi. 901 = Aeonia polystachia.
porrectus, Reichb. f. in Flora, xlviii. (1865) 279.—Ind. occ.
pusillus, Reichb. f. l. c. l. (1867) 117 = M. pusillum.
rectus, S. Moore, in Baker, Fl. Maurit. 351 = Angraecum rectum.
rutilus, Reichb. f. in Flora, lxviii. (1885) 540.—Ins. Comor.
Sallei, Reichb. f. in Walp. Ann. vi. 902.—Ins. S. Doming.
Schiedei, Reichb. f. l. c. 901 = Campyloceutron Schiedei.
sesquipedalis, Reichb. in Flora, lxviii. (1885) 381 = Angraecum sesquipedale.
spathaceus, Griseb. Cat. Fl. Cub. 264.—Cuba.
Thouarsii, S. Moore, in Baker, Fl. Maurit. 35 = Angraecum filicorne.
trifurcus, Reichb. f. in Flora, lxviii. (1885) 540.—Ins. Comor.
volucris, Reichb. f. in Walp. Ann. vi. 900 = Aeonia Auberti.
xanthopollinius, Reichb. l. c. xlviii. (1865) 190.—Afr. trop.

AÉRIDES, Lour. Fl. Cochinch. 525 (1790). *ORCHIDEAE*. Benth. & Hook. f. iii. 576.
AERIDIUM, Salisb. in Trans. Hort. Soc. i. (1812) 295.
ORXERA, Raïn. Fl. Tellur. iv. 37 (1826).
acuminatum, *Lindl. Gen. et Sp. Orch.* 240.—Java.
affine, Wall. Cat. n. 7316 = multiflorum.
amplexicaule, Lindl. Gen. et Sp.Orch. 239 = Sarcochilus illaciinus.
ampullaceum, Roxb. Fl. Ind. iii. 476 = Sarcolabium ampullaceum.
anceps, Lindl. Gen. et Sp. Orch. 242 = Sarcochilus falcatus.
angustifolium, Lindl. l. c. 241 = Sarcochilus angustifolius.
approximulatum, Wall. Cat. n. 7315 = Sarcanthus appendiculatus.
Arachnites, Lindl. Gen. et Sp. Orch. 238 = Sarcochilus Arachnites.
Arachnites, Sw. in Schrad. Journ. ii. (1799) 238 = Arachnanthe moschifera.
Balfourianum, Reichb. f. in Gard. Chron. N. S. xxiv. (1885) ii. 196 = suaveolaneum.
Bernhardianum, *Reichb. f. l. c.* 650.—Borneo.
bicolor, Reinw. ex Lindl. Fol. Orchid. Vanda, 4 = Vanda furva.
Borssi, *Buch.-Ham. ex Sm. in Rees, Cyclop.* xxxix. n. 8.—Ind. or.
Brachensii, Hayah. Nom. ii. 11 = crispum.
Brookei, Batem. ex Lindl. in Bot. Reg. (1841) Misc. 55 = crispum.
calceolare, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 19.—Sumatra.
calceolare, Buch.-Ham. ex Sm. in Rees, Cyclop. xxxix. n. 11 = Saccolabium calceolare.
carnosum, Griff. Notul. iii. 345 = Doritis taeniifolia.
compressum, Lindl. Gen. et Sp. Orch. 240 = Dendrocolla compressa.
coriaceum, *Sw. in Schrad. Journ.* (1799) 234.—Madag.
corusatum, Roxb. Hort. Beng. 68; Lindl. Bot. Reg. t. 1486 = odoratum.

AÉRIDES :—
costatum, *' Weekh.' ex Steud. Nom.* ed. II. i. 30.—Ind. or.
crassifolium, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 146.—Burma.
crispum, *Lindl. in Wall. Cat.* n. 7319.—Malabar.
cylindricum, Hook. Bot. Mag. t. 4953 = A. vandarum.
cylindricum, *Lindl. in Wall. Cat.* n. 7317.—Ind. or.
decumbens, *Grisf. Notul.* iii. 365; *Ic. Pl. Asiat.* t. 390.—Burma.
dasypogon, *Sm. in Rees, Cyclop.* xxxix. n. 10.—Nepal.
dasypogon, Hort. ex Batem. in Hook. Bot. Mag. sub t. 5630 = Sarcanthus erinaceus.
difforme, Wall. ex Lindl. Gen. et Sp. Orch. 242 = Ornithochilus fuscus.
diurnum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 19.—Sumatra.
elongatum, *Blume, Bijdr.* 367.—Java.
emarginatum, Lindl. Gen. et Sp. Orch. 241 = Sarcochilus emarginatus.
Emericii, *Reichb. f. in Gard. Chron.* xviii. (1882) ii. 556.—Ins. Andam.
expansum, Reichb. f. l. c. 40 = falcatum.
falcatum, *Lindl. & Paxt. Fl. Gard.* ii. (1851) 142.—Burma.
Farmeri, *Boxall, ex Nowes, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 289.—Ins. Philipp.
Fenzlianum, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 263.—Ins. Philipp.
Fieldingi, *Lindl. ex E. Morr. in Belg. Hortic.* (1876) 288. t. 10.—Reg. Himal.
flavidum, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 101 = suaveolaneum.
Fleischeria, Sw. in Schrad. Journ. ii. (1799) 238 = Sarcochilus Arachnites.
guttatum, Roxb. Fl. Ind. iii. 471 = Rhynchostylis retusa.
Houlletianum, *Reichb. f. in Gard. Chron.* (1872) 1194.—Cochinch.
Hystrix, *Lindl. Gen. et Sp. Orch.* 242 = Sarcochilus Hystrix.
Hystrix, Lindl. in Journ. Linn. Soc. iii. (1859) 42 = Ornithochilus fuscus.
illustre, Reichb. f. in Gard. Chron. N. S. xviii. (1882) ii. 71 = maculosum.
inflexum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 294.—Malaya.
japonicum, *Reichb. f. in Hamb. Gartenz.* xix (1863) 210.—Japon.
jucundum, *Reichb. f. l. c.* xvi. (1860) 281.—Java.
Lacpedae, Hort. ex Reichb. f. in Otto et Dietr. Allg. Gartenz. xxiv. (1856) 219 = falcatum.
lasiopetala, *Willd. Sp. Pl.* iv. 120.—Ind. or.
latifolium, *Thw. Enum. Pl. Zeyl.* 439.—Zeylan.
Lawrenciae, *Reichb. f. in Gard. Chron.* xx. (1883) 460.—As. trop.
Leeanum, *Reichb. f. l. c.* xv. (1881) 656.—Hab.?
Leonei, *Reichb. f.*—Cf. *L'Orchidoph.* (1886) 201.—Burma.
lepidum, *Reichb. f. in Gard. Chron.* xix. (1883) 466.—Ind. or.
Lindleyanum, Wight, Ic. Pl. Ind. Or. 1677 = crispum.
Lobbii, Hort. ex Lem. Illustr. Hortic. xv. (1868) t. 550 = multiflorum.
Lobbii, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 293.—Burma.
McMorlandi, *Williams, Man. Orch.* 66; ed. VI. 105.—Ind. or.
macrostachyum, Spreng. Syst. iii. 718 = Aeonia macrostachya.
maculatum, Bech.-Ham. ex Sm. in Rees, Cyclop. xxxix. n. 9 = Vanda spathulata.
maculatum, *Lindl. Fragm.* 98.—Ins. Philipp.
maculosum, *Lindl. Bot. Reg.* xxxi. (1845) 58.—Ind. or.
margaritaceum, *Williams, Man. Orch.* 66; ed. VI. 106.—Ind. or.
marginatum, *Reichb. f. in Gard. Chron. N. S.* xxiii. (1885) i. 388.—Hab.?
matutinum, *Blume, Bijdr.* 366.—Java.
maximum, Willd. Sp. Pl. iv. 121 = Renanthera Arachnites.
Meadiii, Hort. ex S. Moor. in Belg. Hortic. (1876) 288 = falcatum.
minimum, Lindl. Gen. et Sp. Orch. 241 = Sarcochilus minimus.

AËRIDES :—
mittratum, Reichb. f. in Bot. Zeit. xxii. (1864) 415.—Ind. or.
multiflorum. Roxb. Fl. Ind. iii. 475.—Ind. or.
noble, Warn. Sel. Orch. Ser. I. t. 11 = suavissimum.
obtusum, Lindl. Gen. et Sp. Orch. 289 = Sarcochilus obtusus.
odoratum, Lour. Cochinch. 525.—China; Ind. or.
odoratum, Reinw. ex Blume, Ramphia, iv. 53 = suaveolens.
Ottgisianum, Reichb. f. in Gard. Chron. xxiii. (1885) 501.—Hab.?
pachyphyllum, Reichb. f. l. c. xiv. (1880) 231.—Burma.
pallidum, Blume, Rumphia, iv. 53.—Ins. Timor.
pallidum, Lindl. Gen. et Sp. Orch. 241 = Sarcochilus pallidus.
pallidum, Roxb. Fl. Ind. iii. 475 (= *Camarotis pallida*).
panniculatum, Ker-Gawl. Bot. Reg. fil. 220 = Vanda paniculata.
praemorsum, Willd. Sp. Pl. iv. 120.—Rhynchostylis retusa.
purpurascens, *Lindl. Gen. et Sp. Orch.* 240.—Java.
pusillum, Lindl. l. c. 241 = Sarcochilus pusillus.
quinquevalnera, *Lindl. Sert. Orch.* 20.—Malaya.
radiatum, *Roxb. Fl. Ind.* iii. 476.—Ind. or.
radicosum, *A. Rich. in Ann. Sc. Nat. Sér.* II. xv. (1841) 65. t. 1. f. c.—Ind. or.
Reichenbachii, Linden, in C. Koch & Fint. Wochenschr. (1858) 61 = suavissimum.
retusum, Sw. in Schrad. Journ. ii. (1799) 233 = Rhynchostylis retusa.
rigidum, *Buch.-Ham. ex Sm. in Rees, Cyclop.* xxxix. n. 12.—Ind. or.
Roebelenii, *Reichb. f. in Gard. Chron* xxi. (1884) 510.—Ins. Philipp.
Rohmsianum, Reichb. f. l. c. 206 = suavissimum.
roseum, Lodd. ex Paxt. Fl. Gard. t. 60 = multiflorum.
rostratum, Roxb. Fl. Ind. iii. 474 = Sarcochilus purpureus.
Ruckeri, *Hort. Donat. ex Lubbers, Cat. Pl. Rar. San Donato* (1880) 14.—Hab.?
Sanderianum, *Reichb. f. in Gard. Chron.* xxii. (1884) 184.—As. trop.
Schroderi, Hort. ex Reichb. f. in Allgem. Gartenz. (1855) 226 = maculosum.
sibbitanum, *Boxall, ex Nowes, in Blanco, Fl. Philipp.* ed. II. *Nov. App.* 289.—Ins. Philipp.; sphalm.? = Thibautianum.
spicatum, D. Don, Prod. Fl. Nep. 31 = Rhynchostylis retusa.
spurium, *Lindl. Gen. et Sp. Orch.* 241.—Java.
suaveolens, *Blume, Rumphia*, iv. 53.—Java.
suaveolens, *Roxb. Fl. Ind.* iii. 473.—Ind. or.
suavissimum, Lindl. in Journ. Hort. Soc. iv. (1849) 264. —Malacca.
subulatum, Lindl. Gen. et Sp. Orch. 241 = Sarcochilus subulatus.
Sulingi, Blume, Bijdr. 367 = Renanthera Sulingi.
taeniale, Lindl. Gen. et Sp. Orch. 239 = Dorttis taenialis.
teres, *Lindl. l. c.* 240.—Java.
tessellatum, Wight, in Wall. Cat. n. 7318 = Vanda Roxburghii.
testaceum, Lindl. Gen. et Sp. Orch. 238 = Vanda parviflora.
Teysmanni, *Miq. Choix Pl. Buitenz.* t. 18.—Malaya.
Thibautianum, *Reichb. f. in Gard. Chron.* (1866) 100.—Polynesia.
Thunbergii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 205.—Japon.
tinnoranum, *Miq. Fl. Ind. Bat.* iii. 695.—Ins. Timor.
trigonum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 177 = multiflorum.
umbellatum, Wall. Cat. n. 7137 = Saccolabium acutifolium.
uncinatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 294.—Java.
undulatum, Sm. in Rees, Cyclop. xxxix. n. 13 = Saccolabium praemorsum.
vandarum, *Reichb. f. in Gard. Chron.* (1867) 997.—Ind. or.
Veitchii, Hort. ex E. Morr. in Belg. Hortic. (1876) 291, xxvi. (1881) 198 = multiflorum.
virens, *Lindl. Bot. Reg.* (1843) *Misc.* 41.—Hab.?
viridiflorum, *Thw. Enum. Pl. Zeyl.* 426.—Zeylan.
Wightianum, Lindl. in Wall. Cat. n. 7320 = Vanda parviflora.

AËRIDES :—
Williamsii, Warn. Sel. Orch. Ser. I. t. 21 = Fieldingi Williamsianum, Hort. ex Gard. Chron. xxiii. (1885) 705 = suavissimum?
Zollingeri, *Reichb. f. in Bonplandia*, v (1857) 40.—Java.

AËRIDIUM, Pfeiff. Nom. i. 67 (1873) = AËRIDIUM, Salisb. = **Aërides**, Lour. (Orchid.).

AËROBION, Kaempf ex Spreng. Syst. iii. 679 (1826) = **Angraecum**, Thou. (Orchid.).
aphyllum, Spreng. l. c. 718 = Angr aphyllum.
Calceolus, Spreng. l. c. 717 = Angr. Calceolus.
carinatum, Spreng. l. c. 718 = Eulophia virens.
caulescens, Spreng. l. c. 717 = Mystacidium caulescens.
citratum, Spreng. l. c. 718 = Angr. citratum.
crassum, Spreng. l. c. 717 = Angr. crassum.
cucullatum, Spreng. l. c. = Angr cucullatum.
expansum, Spreng. l. c. 716 = Angr. expansum.
filicornu, Spreng. l. c. = Angr. filicornu.
fragrans, Spreng. l. c. = Angr. fragrans.
gladiifolium, Spreng. l. c. = Angr. gladiifolium.
gracile, Spreng. l. c. 717 = Angr. gracile.
implicatum, Spreng. l. c. 716 = Angr. implicatum.
inapertum, Spreng. l. c. 717 = Angr. inapertum.
maculatum, Spreng. l. c. 718 = Angr. maculatum.
multiflorum, Spreng. l. c. 717 = Angr multiflorum
palmiforme, Spreng. l. c. = Angr palmiforme.
parviflorum, Spreng. l. c. 718 = Angr. parviflorum.
pectinatum, Spreng. l. c. 717 = Angr. pectinatum.
rectum, Spreng. l. c. 716 = Angr. rectum.
recurvum, Spreng. l. c. = Angr recurvum
striatum, Spreng. l. c. 717 = Angr. striatum.
superbum, Spreng. l. c. 718 = Angr. superbum.
triquetrum, Spreng. l. c. 717 = Angr. triquetrum.

AËROKORION, Scop. Ann iv. Hist. Nat. 96 (1770) = AEROCORION, Adans. = **Galanthus**, Tourn. (Amaryll.).

AERUA, Juss. Gen. 88 (1789) = **Aerva**, Forsk. (Amarant.).

AERVA, Forsk Fl. Aegypt. Arab 170 (1775.) *AMARANTACEÆ* Benth. & Hook. f. iii. 84).
aegyptiaca, J. F. Gmel. Syst. ii. 1026 = tomentosa.
ambigua, *Moq. in DC. Prod.* xiii 11. 302.—*Afr. austr.*
bengalensis, Buch.-Ham ex Wall. Cat. sub n. 6911 = scandens.
Bovii, Edgew in Journ. Linn Soc. vi (1862) 206 = tomentosa.
brachiata, Mart. Beitr. Amarant. 88 = Nothosaerva brachiata.
Burchellii, Moq in DC Prod. xiii. 11. 302 = A. lanicura.
caudata, *Spr. Hort. Mauric.* 268.—Madag. ?
chenopodifolia, Boj. l. c. = Nothosaerva brachiata.
congesta, *Balf. f. ex Baker, Fl. Maurit.* 267.—Ins. Mascar.
elegans, *Moq. in DC. Prod.* xiii. 11. 303.—Arab.
floribunda, Wight, Ic. Pl. t. 1776 = lanata.
glabrata, *Hook. f. Fl. Brit. Ind.* iv. 728.—Ind. or.
lanata, Mart. in Nov. Act. Acad. Nat. Cur. xii. (1826) 291 = tomentosa.
javanica, Juss. in Ann. Mus. Par. xi. (1808) 181 = tomentosa.
lanata, *Juss. l. c.*—As. trop.; Arab.
leucura, *Moq. in DC. Prod.* xiii. 11. 302.—Afr. austr.
microphylla, *Moq. l. c.* 301.—' Mare rubrum.'
Monsonia, *Mart. Beitr. Amarant.* 88.—Ind. or.
nervosa, *Mart. l. c.*—Ind. or.
ovata, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 174 = tomentosa.
platyphylla, Ehrenb. ex Schweinf. l. c. = tomentosa.
pubescens, *Mart. Beitr. Amarant.* 88.—Ind. or.
radicans, Mart. l. c.—Ind. or.
revoluta, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 92.—Ins. Socotr.
sanguinea, Ker Moq. in DC. Prod. xiii. 11. 300 = sanguinolenta.
sanguinolenta, *Blume, Bijdr.* 547.—Java.
scandens, Gamble, Trees of Darjeeling = Stilbanthus scandens.
scandens, *Wall. Cat. n.* 6911.—As. et Afr. trop.
sericea, *Moq. in DC. Prod.* xiii. 11. 304.—Ins. Sandvic.
setacea, Mart. Beitr. Amarant. 88 = A. Monsonia.
suaveolens, Edgew. in Journ. Linn. Soc. vi. (1862) 207 = tomentosa.

AERVA :—
timorensis, *Moq. in DC. Prod* xiii. 11. 301.—Ins. Timor
timoriensis, Zipp. ex Span. in Linnaea, xv. (1841) 345 = sanguinolenta.
tomentosa, *Lam. Encyc.* i. 46.—As. et. Afr. trop.
tomentosa, Buch.-Ham. ex Wall. Cat. sub n. 6911 = scandens.
Tondala, Buch.-Ham. l. c. 6909 = lanata.
trichophylla, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop 174 = tomentosa.
velutina, Moq. in DC. Prod. xiii. 11 301 = Pupalia lappacea.
villosa, *Moq. l. c.* 304.—Ins. Maurit.
viridis, L. Mey. ex Moq. l. c. = lanata.
Wallichii, Moq. l. c. 300 = tomentosa.
Wightii, *Hook. f. Fl. Brit. Ind.* iv. 728.—Ind. or.

AESCHINANTHUS, Endl. Gen. 716 (1838) = **Aeschynanthus**, Jack (Gesneriac.).

AESCHINOMENE, Nocca in Usteri, Ann. Bot. v. (1798 16 = **Aeschynomene**, Linn. (Legum.).

AFSCHRION, Vell. Fl. Flum 58; i. t. 152 (1827) = **Piresana**, Lindl. (Simarub.).
crenata, Vell. il cc. = P. Vellozii.

AESCHYNANTHUS, Jack, in Trans. Linn. Soc. xiv. (1823) 42. *GESNERACEÆ*, Benth. & Hook. f. ii 1018.
RHEITHROPHYLLUM, Hassk. Cat. Hort. Bogor. Alt. 128 (1844)
TRICHOSPORUM, D. Don, in Edinb. Phil. Journ. vii. (1822) 84 (nomen prius).
acuminata, G. Don, Gen. Syst. iv. 667 = Chirita acuminata.
acuminata, *Wall. Cat. n.* 6897.—Ind. or.; China.
albida, A. DC. Prod. ix. 262 = purpurascens.
albida, *Steud. Nom.* ed. II. i. 32.—Java.
alba, Hort. ex Hanst. in Mart. Fl. Bras. viii. 418 = Codonanthe Devosiana.
amoena, *C. B. Clarke, in DC. Monog. Phan.* v. 47.—Java.
Andersoni, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 48. —Burma.
angustifolia, *Steud. Nom.* ed. II. i. 32.—Java.
apiculens, *Hance, in Journ. Bot.* xxi. (1883) 167.—China
arfakensis, *C. B. Clarke, in DC. Monog. Phan.* v. 86. —N. Guin.
atropurpurea, Van Houtte, Hort. Vanhout. Fasc. ii. 48 = purpurascens.
atrosanguinea, Van Houtte, Cat. (1851) ex C. B. Clarke, in DC. Monog. v 52.—Hab ?
Auckландiae, *Lem, Saraand.* 382, 386.—Borneo.
Auckландi, Hort. Low, ex Loud Hort. Brit. Supp. iii. 484 = pracc.
Beccarii, *C. B. Clarke, in DC. Monog. Phan.* v. 47.—Sumatra.
Blumei, Mor. ex C. B. Clarke, l. c. 48 = A. Lamponga.
Boschiana, De Vriese, in Morr. Ann. Soc. Agric. Bot. Gand, ii. (1846) 408 = A. Lamponga.
bracteata, Benth. Fl. Hongkong, 226 = acuminata.
bracteata, *Wall. Cat. n.* 794.—Ind. or.
brevicalyx, *Miq. Fl. Ind. Bat.* ii. 730.—Sumatra.
candida, *Henders. Cat.* (1851) *ex C. B. Clarke, in DC. Monog. Phan.* v. 52.—Hab.?
caudata, *C. B. Clarke, l. c.* 84.—Sumatra.
ceylanica, *Gard. in Calc. Journ. Nat. Hist.* vi. (1846) 474.—Zeylan.
ceylanica, Wight, Ic. t. 1347 = Perrottetii.
chinensis, Gard. & Champ. in Hook. Kew Journ. i. (1849) 238 = acuminata.
chiritoides, *C. B. Clarke, in DC. Monog. Phan.* v. 28. —Reg. Himal.
conferta, Griff. Priv. Journ. 68, nomen = Lepionotus confertus.
cordifolia, *Hook. Bot. Mag.* t. 5131.—Borneo.
cryptantha, *C. B. Clarke, in DC. Monog. Phan.* v. 50. —Borneo.
Curtisii, *C. B. Clarke, l. c.* 45.—Borneo.
discolor, T. Moore, in Paxt. Fl. Gard. iii. (1852-53) 53 = purpurascens.
elongata, *C. B. Clarke, in DC. Monog. Phan.* v. 45.—Borneo.
fulgens, *Wall. Cat. n.* 797.—Ind. or.
geminata, *Zoll. & Morr. in Nat. en Genees. Arch. Neerl. Ind.* ii. (1845) 574.—Java.

AESCHYNANTHUS :—
gracilis, Hort. ex Hanst. in Mart. Fl. Bras. viii. 418 = Codonanthe Devosiana.
gracilis, *Parish, ex C. B. Clarke, Comm. et Cyrt. Beng.* t. 48.—Ind. or.
grandiflora, J. Grah. Cat. Pl. Bomb. 146 = Perrottetii.
grandiflora, *Spreng. Syst.* iv. *Cur. Post.* 238.—Ind. or.
Griffithii, *R. Br. in Benn. Pl Jav. Rar.* 115.—Ind. or.
hians, *C. B. Clarke, in DC. Monog. Phan.* v. 51.—Borneo.
Hookeri, *C. B. Clarke, l. c.* 21.—Reg. Himal.
Horsfieldii, *R. Br. in Benn. Pl. Jav. Rar.* 116.—Java.
intermedia, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 414.—Malaya.
javanica, *Hook. Bot. Mag.* t. 4503.—Java.
Kingii, *C. B. Clarke, in DC. Monog. Phan.* v. 31.—Reg. Himal.
Lamponga, *Miq. Fl. Ind. Bat. Suppl.* 563.—Sumatra.
lepidosperma, *C. B. Clarke, in DC. Monog. Phan.* v. 35.—Borneo.
leptoclada, *C. B. Clarke, l. c.* 39.—N. Guin.
levipes, *C. B. Clarke, l. c.* 28.—Assam.
Lobbiana, *Hook. Bot. Mag.* t. 4260.—Java.
longicaulis, *Wall. Cat.* n. 388.—Ind. or.
longiflora, *Blume, Bijdr.* 766.—Malaya.
longiflora, Wall. Cat. n. 795 = superba.
macrocalyx, *C. B. Clarke, in DC. Monog. Phan.* v. 46.—Sumatra.
maculata, *Lindl. Bot. Reg.* (1841) t. 28.—Ind. or.
Mannii, *Kurz, ex C. B. Clarke, in DC. Monog. Phan.* v. 29.—Ind. or.
marmorata, *F. Moore, in Pant. Fl. Gard.* iii. (1852–53) 56.—Hab. ?
Masoniæ, *Kurz, ex C. B. Clarke, l. c.* 29—Ind. or.
micrantha, *C. B. Clarke, l. c.* 37.—Reg. Himal.
microphylla, *C. B. Clarke, l. c.* 49.—Ins. Celebes.
microtricha, *C. B. Clarke, l. c.* 51.—N. Guin.
minista, *Lindl. Bot. Reg.* N. S. xix. t. 61.—Java.
Motleyi, *C. B. Clarke, in DC. Monog. Phan.* v. 20.—Malaya.
Neesii, Zoll. & Moritzi, Syst. Verz. Zoll. 45 = pulchra.
obconica, *C. B. Clarke, in DC. Monog. Phan.* v. 50.—Malacca.
oblongifolia, G. Don, Gen. Syst. iv. 857 = Chirita acuminata.
obovata, *C. B. Clarke, in DC. Monog. Phan.* v. 47.—Java.
serratifica, C. B. Clarke, Comm. et Cyrt. Beng. t. 49 = Hookeri.
parasitica, Wall. Cat. n. 796 = grandiflora, pro parte.
parvifora, C. B. Clarke, Comm. et Cyrt. Beng. t. 47 = micrantha.
parviflora, *Spreng. Syst. iv. Cur. Post.* 238 = ramosissima.
parvifolia, *R. Br. in Benn. Pl. Jav.* 115.—Banca.
Paxtonii, Lindl. in Journ. Hort. Soc. iv. (1849) 79 = bracteata.
Peelii, Hook. f. & Thoms. Illustr. Himal. Pl. 17 = bracteata.
Perrottetii, *A. DC. in DC. Prod.* ix. 261.—Ind. or.
Perrottetii, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 475 = ceylanica.
philippinensis, *C. B. Clarke, in DC. Monog. Phan.* v. 39.—Ins. Philipp.
podocarpa, *C. B. Clarke, l. c.* 40.—N. Guin.
pulchella, Hort. ex Hanst. in Mart. Fl. Bras. viii. 418 = Codonanthe Devosiana.
pulchra, *G. Don, Gen. Syst.* iv. 856.—Malaya.
purpurascens, *Hassk. Cat. Pl. Hort. Bogor. Alt.* 154.—Java.
radicans, *Jack, in Trans. Linn. Soc.* xiv. (1823) 43.—Malaya.
radicans, Wall. Cat. n. 798 = Griffithii.
ramosissima, *Wall. Cat.* n. 799.—Ind. or.
Roxburghii, *Past. Bot. Dict. ex C. B. Clarke, in DC. Monog.* v. 52.—Hab. ?
rubiginosa, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 413.—Java.
siphonantha, *C. B. Clarke, in DC. Monog. Phan.* v. 45.—Borneo.
speciosa, *Hook. Bot. Mag.* t. 4820.—Java.
splendens, Lindl. & Paxt. Fl. Gard. iii. 169. t. 106 = speciosa.
splendida, *Lem. Jard. Fleur.* iii. (1858) t. 265.
superba, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 46.—Ind. or.
Sacokhoffii, Hort. Berol. ex C. B. Clarke, in DC. Monog. Phan. v. 48 = pulchra.
tetraquetra, *C. B. Clarke, l. c.* 36.—Sumatra.

AESCHYNANTHUS :—
Teysmanniana, *Miq. in Bot. Zeit.* vi. (1848) 509.—Java.
tricolor, *Hook. Bot. Mag.* t. 5081.—Java.
tubiflora, *C. B. Clarke, in DC. Monog. Phan.* v. 26.—N. Guin.
verticillata, *C. B. Clarke, l. c.* 40.—N. Guin.
viridiflora, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 34.—Ins. Celebes.
volubilis, *Jack, in Trans. Linn. Soc.* xiv. (1823) 42. t. 3. f. 3.—Sumatra.
Wallichii, *Benn. Pl. Jav. Rar.* 116.—Malaya.
Wilsoni, Walker, ex C. B. Clarke, in DC. Monog. v. 26 = ceylanica.
zebrina, Van Houtte [Cat. 1851] ex Walp. Rep. vi. 521 = marmorata.
zeylanica, Walp. Ann. iii. 95 = ceylanica.
Zollingeri, *C. B. Clarke, in DC. Monog. Phan.* v. 44.—Java.

AESCHYNOMENE, Linn. Gen. ed. I. 350 (1737).
LEGUMINOSAE, Benth. & Hook. f. i. 515.
GAJATI, Rumph. ex Adans. Fam. ii. 328 (1763).
MANTODDA, Adans. Fam. ii. 308 (1763).
MACROMISCUS, Turcz. in Bull. Soc. Nat. Mosc. (1846) ii. 507.
PATAGONIUM, E. Mey. Comm. Pl. Afr. Austr. 132 (1835).
ROCHEA, Scop. Introd. 296 (1777).
RUPPELLIA, A. Rich. Tent. Fl. Abyss. i. 208. t. 87 (1847).
abyssinica, Vatke, in Oest. Bot. Zeitschr. xxix. (1879) 324 = Ruppellii.
aegyptiaca, Steud. Nom. ed. II. i. 31 = Sesbania aegyptiaca.
acutangula, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 148.—Afr. trop.
americana, *Linn. Sp. Pl.* 713.—Ind. occ.; Am. trop.
aquatica, Steud. Nom. ed. II. i. 31 = aspera.
arborea, Linn. Sp. Pl. 713 = Desmodium umbellatum.
aristata, Jacq. Hort. Schoenb. i. 59 = Pictetia aristata.
aspera, *Linn. Sp. Pl.* 714.—Geront. trop.
aspera, Wall. Cat. n. 5666 = indica.
aspera, Muhl. ex Willd. Sp. Pl. iii. 1163 = hispida.
atropurpurea, *Span. in Linnaea*, xv. (1841) 192.—Ins. Timor.
bispinosa, Jacq. Ic. iii. 18 = Sesbania aculeata.
bracteosa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 150.—Afr. trop.
brasila, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 53 = Martiusii.
brasiliana, *DC. Prod.* ii. 322.—Bras.
brevifolia, *Pav. Encyc.* iv. 451.—Madag.
brevipes, *Benth. in Mart. Fl. Bras.* xv. 1. 66.—Bras.
Burmanni, *Raeusch. Nom.* ed. III. 214.—Ind. or.
cachemiriana, Cambess. in Jacquem. Voy. Bot. 40 = indica.
cannabina, Retz. Obs. Fasc. v. 26 = Sesbania aculeata.
cassioides, Desv. in Ham. Prod. Pl. Ind. Occ. 51 = Histrix.
cassioides, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 240 = Sesbania cassioides.
ciliata, Vog. in Linnaea, xii. (1838) 84 = hispida.
coccinea, Linn. f. Suppl. 330 = Sesbania grandiflora.
colutoides, Less. & A. Rich. Voy. Astrol. Bot. ii. 87. t. 32 = Ormocarpum sennoides.
conferta, *Benth. in Ann. Nat. Hist.* iii. (1839) 438.—Guian.
crepita, *Vacq. Fragm.* 37. t. 42. f. 2.—Venezuela.
cristata, *Vatke, in Oest. Bot. Zeitschr.* xxviii. (1878) 215.—Afr. trop.
debilis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 150.—Afr. trop.
decumbens, Zipp. ex Span. in Linnaea, xv. (1841) 198 = sericea.
densiflora, *Benth. in Hook. Lond Journ. Bot.* ii. (1840) 56.—Guian.
dentata, Lag. Gen. et Sp. Nov. 22 = Adesmia dentata.
diffusa, Roxb. ex Wight & Arn. Prod. 215 = Sesbania procumbens.
diffusa, Klein, ex Willd. Sp. Pl. iii. 1164 = indica.
dimidiata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 151.—Afr. trop.
divisa, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1826) 31 = americana.
Echinus, *Vog. in Linnaea*, xii. (1838) 92.—Bras.

AESCHYNOMENE :—
elegans, *Cham. & Schlecht. in Linnaea*, v. (1830) 583.—Mexico.
elongata, Salisb. Prod. 341 = Sesbania aegyptiaca.
erubescens, *E. Mey. Comm. Pl. Afr. Austr.* i. 133.—Afr. austr.
evenia, *Wright, in Sauv. Fl. Cub.* 27.—Cuba.
falcata, *DC. Prod.* ii. 322.—Bras.
fascicularis, *Cham. & Schlecht. in Linnaea*, v. (1830) 584.—Mexic.
filipes, *Basil. in Bull. Soc. Linn. Par.* i. (1884) 415.—Madag.
floss, *Mart. ex Benth. in Mart. Fl. Bras.* xv. 1. 61.—Bras.
fistulosa, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 259.—Porto Rico.
floribunda, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 186 = americana.
fluminensis, *Vell. Fl. Flum.* 310; vii. t. 119.—Bras.
frutescens, Poir. in Lam. Encyc. iv. 451 = Lespedeza capitata.
fruticulosa, Nees, ex G. Don, Gen. Syst. ii. 232 = Adesmia fruticulosa.
filgida, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 149.—Afr. trop.
fusa, Steud. Nom. ed. I. 17, sphalm. = seq.
fusca, *Desf. Tabl. Cat. Hort. Par.* ed. I. 198.—Hab. ?
gangetica, Poir. in Lam. Encyc. iv. 458 = Desmodium gangeticum.
glaberrima, Poir. Encyc. Suppl. iv. 76 = hispida ?
glabrescens, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 148.—Afr. trop.
glandulosa, Benth. in Miq. Fl. Jungh. 210 = javanica.
glandulosa, Poir. Encyc. Suppl. iv. 76 = americana.
gracilis, Miq. in Linnaea, xviii. (1844) 566 = paniculata.
gracilis, Vog. l. c. xii. (1838) 891 = falcata.
grandiflora, Linn. Sp. Pl. ed. II. 1050 = Sesbania grandiflora.
guayaquilensis, *G. Don, Gen. Syst.* ii. 284.—Ecuador.
hedysaroides, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 180.—Mexic.
herbacea, *Aubl. Pl. Gui.* ii. 775.—Guian.
heterophylla, *Lour. Fl. Cochinch.* 446.—Cochinch.
Heurckeana, *Baker, in Journ. Linn. Soc.* xx. (1884) 180.—Madag.
hirsuta, Vahl. Symma Pl. iv. 270 = Burmanni.
hirsuta, DC. Prod. ii. 322 = americana.
hirta, *Lam. Illustr.* t. 629, *ex ejusd. Encyc.* iv. 450.—Ind. or.
hispida, Roxb. ex Steud. Nom. ed. II. i. 31 = Robinia hispida.
hispida, *Willd. Sp. Pl.* iii. 1163.—Am. bor.
hispidula, *H. B. & K. Nov. Gen. et Sp.* vi. 580.—N. Granat.
hispidula, Lag. Gen. et Sp. Nov. 22. n. 297 = Adesmia hispidula.
Histrix, *Poir. Encyc. Suppl.* iv. 77.—Guian.
honesta, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1826) 32 = sensitiva.
incana, G. F. W. Mey. ex DC. Prod. ii. 332 = Desmodium incanum.
incana, *Vog. in Linnaea*, xii. (1838) 90.—Bras.
indica, Burm. f. Fl. Ind. 169 = Sesbania aegyptiaca.
indica, *Linn. Sp. Pl.* 713.—Ind. or.
indica, Wall. Cat. n. 5667 = aspera.
interrupta, *Benth. in Hook. Lond. Journ. Bot.* ii. (1840) 56.—Guian.
javanica, *Miq. Fl. Ind. Bat.* i. 275.—Java.
kashmiriana, Baker, in Hook. f. Fl. Brit. Ind. i. 151 = indica.
laevis, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 180.—Mexico.
Lagenaria, Lour. Fl. Cochinch. ii. 446 = aspera.
laevis, *Naronha, in Verh. Batav. Gen.* v. (1790) ed. I. art. iv. 5.—Malaya.
latifolia, Spreng. Syst. ii. 232 = Adesmia latifolia.
latistipulacea, Hill, ex Steud. Nom. ed. II. i. 31 = Poiretia latistipulacea.
laxiflora, *Boj. ex Baker, in Journ. Linn. Soc.* xx. (1884) 180.—Madag.
leptocladoya, *Benth. in Mart. Fl. Bras.* xv. 1. 63.—Am. austr.
litorella, *Vog. in Linnaea*, xii. (1838) 96.—Bras.
longifolia, Cav. Ic. iv. 6 = Pictetia longifolia.
longifolia, Orteg. Hort. Matr. Dec. 70 = Sesbania Cavanillesii.

AESCHYNOMENE:—
macropoda, DC. Prod. ii. 320 = indica.
maculata, Poir. in Lam. Encyc. iv. 459 = Desmodium gangeticum.
marginata, *Benth. in Mart. Fl. Bras.* xv. i. 66.—Am. trop.
Martii, *Benth. l. c.* 62.—Am. trop.
Martiusii, *Steud. Nom.* ed. II. i. 31.—Bras.
mazangayana, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 415.—Madag.
micrantha, *DC. Prod.* ii. 321.—Madag.
microphylla, *Desv. ex DC. l. c.* 325.—Bras.
mimosoides, Noes, ex G. Don, Gen. Syst. ii. 282 = Adesmia mimosoides.
mimulosa, Blume, ex Miq. Fl. Ind. Bat. i. 276 = javanica.
miniata, Orteg. Hort. Matr. Dec. 36 = Piscidia punicea.
mimosifolia, *Vatke, in Oest. Bot. Zeitschr.* xxix. (1879) 224.—Afr. trop.
mollicula, *H. B. & K. Nov. Gen. et Sp.* vi. 532.—Am. austr.
molucana, *Kostel. Allg. Med.-pharm. Flora*, iv. 1285.—Moluccs.
montana, Span. in Linnaea, xv. (1841) 192 = indica.
montevidensis, *Vog. in Linnaea*, xii. (1838) 88.—Reg. Argent.
mucronata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1840) 56.—Guian.
mucronulata, *Benth. l. c.* 56.—Guian.
multiflora, *Steud. Nom.* ed. II. i. 31.—Ind. or.
obovalis, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 415.—Madag.
oligantha, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 146.—Afr. trop.
oroboides, *Benth. in Mart. Fl. Bras.* xv. i. 64.—Am. austr.
paludosa, Roxb. Fl. Ind. iii. 338 = Sesbania aculeata
paniculata, *Willd. ex Vog. in Linnaea*, xii. (1838) 95.—Bras.
papposa, Lag. Gen. et Sp. Nov. 23 = Adesmia papposa.
papulosa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 146.—Afr. trop.
parviflora, *Michell, in Kjoeb. Vidensk. Meddel.* (1875) 66.—Bras.
patagonica, Hort. ex Steud. Nom. ed. II. i. 31 = Adesmia muricata.
patula, Poir. Encyc. Suppl. iv. 78 = sensitiva.
pauciflora, *Vog. in Linnaea*, xii. (1838) 93.—Bras.
paucifolia, *Vog. l. c.* 94.—Bras.
paucijuga, DC. Prod. ii. 321 = brasiliana.
picta, Cav. Ic. iv. 7 = Sesbania aegyptiaca.
pilosa, *Poir. in Lam. Encyc.* iv. 450.—Ind. or
platycarpa, *Benth. in Mart. Fl. Bras.* xv. i. 63.—Peruv.
platycarpa, Michx. Fl. Bor. Am. ii. 75 = Sesbania vesicaria.
pleurocervia, *DC. Prod.* ii. 321.—Ins. S. Doming.
plurisiticulata, *G. Don, Gen. Syst.* ii. 284.—Ecuador.
podocarpa, Vog. in Linnaea, xii. (1838) 89 = falcata.
procumbens, Roxb. Fl. Ind. iii. 337 = Sesbania procumbens.
prostrata, Poir. Encyc. Suppl. iv. 78 = viscidula.
pseudoacacia, Roxb. ex DC. Prod. ii. 261 = Robinia Pseudacacia.
pseudovisicosa, Blume, ex Miq. Fl. Ind. Bat. i. 276 = javanica.
pubernula, DC. Prod. ii. 321 = scoparia.
pubescens, *Poir. in Lam. Encyc.* iv. 450.—Ind. or.
pubescens, Vahl, ex DC. Prod. ii. 265 = Sesbania pubescens.
pudica, *F. G. Dietr. Lexik. Gaertn.* i. 186.—Aegypt.
pudica, *Zoll. & Morr. in Nat. en Genees. Arch. Neerl. Ind.* iii. (1846) 55, nomen.—Java.
puichella, *Planch. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 149.—Afr. trop.
pulchra, *Vatke, in Oest. Bot. Zeitschr.* xxix. (1879) 250.—Afr. trop.
pumila, Linn. Sp. Pl. ed. II. 1061 = indica.
punctata, Steud. Nom. ed. I. 17 = indica.
pygmaea, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 148.—Afr. trop.
quadrata, Schum. & Thonn. Beskr. Guin. Pl. 356 = indica.
racemosa, *Vog. in Linnaea*, xii. (1838) 92.—Bras.
ramosa, Poir. in Lam. Encyc. iv. 453 = Desmodium Scalpe.
rostrata, *Benth. in Mart. Fl. Bras.* xv. i. 60.—Bras.
Roxburghii, Spreng. Syst. iii. 228 = indica.
rudis, Benth. Pl. Hartw. 116 = hispidula.

AESCHYNOMENE:—
Ruppellii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 149.—Afr. trop.
scabra, *G. Don, Gen. Syst.* ii. 284.—Ecuador.
Schimperi, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 202.—Abyss.
Scoparia, *H. B. & K. Nov. Gen. et Sp.* vi. 529.—Peruv.
scoparia, *Splitgerb. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 107.—Bras.
Selloi, *Vog. in Linnaea*, xii. (1838) 82.—Bras.
sensitiva, Beauv. Fl. Owar. 89. t. 58 = indica.
sensitiva, *Sw. Fl. Ind. Occ.* iii. 1276.—Am. trop.
Sesban, Jacq. Collect. ii. 288 = Sesbania aculeata.
Sesban, Linn. Sp. Pl. 714 = Sesbania aegyptiaca.
silfolia, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 151.—Afr. trop.
spicata, Poir. in Lam. Encyc. iv. 452 = Desmodium spinulosum.
spinulosa, Roxb. Fl. Ind. iii. 338 = Sesbania aculeata.
submicosa, DC. Prod. ii. 321 = indica.
sulcata, H. B. & K. Nov. Gen. et Sp. vi. 530 = sensitiva.
surattensis, *Wight & Arn. Prod.* 219 = Sesbania aculeata?
Suymnata, Steud. Nom. ed. II. i. 31 = Sesbania aegyptiaca.
tecta, Vog. in Linnaea, xii. (1838) 87 = falcata.
tenuirama, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 150.—Afr. trop.
tenuis, *Griseb. Cat. Pl. Cub.* 72.—Cuba.
ternata, Spreng. ex DC. Prod. ii. 314 = Pictetia ternata.
timoriana, Span. Ic. 62; et in Linnaea, xv. (1841) 198 = sensitiva.
trachyloba, Miq. Fl. Ind. Bat. i. 276 = aspera.
tribuloides, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 414.—Madag.
trifora, Poir. in Lam. Encyc. iv. 451 = Desmodium triflorum.
tripartita, St. Hil. ex Steud. Nom. ed. II. i. 32 = Desmodium triquetrum.
trisperma, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1103.—Guian.
tuberculata, *Griseb. Cat. Pl. Cub.* 72.—Cuba.
uliginosa, Roxb. Fl. Ind. iii. 334 = Sesbania aculeata.
unifiora, *E. Mey. Comm. Pl. Afr. Austr.* i. 123.—Afr. austr.
verslcolor, *Wender. in Bot. Zeit.* i. (1843) 347.—Mexic.
villosa, *Poir. Encyc. Suppl.* iv. 76.—Ins. Porto-Rico.
violacea, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. *Art.* iv. 5.—Malaya.
virgata, Cav. Ic. iii. 47 = Coursetia virgata.
viscidula, *Michx. Fl. Bor. Am.* ii. 75.—Florida.
viscidula, Roxb. ex Willd. Enum. Hort. Berol. 776 = indica.
viscosa, *Vatke, in Bremen Abh.* vii. (1882) 247.—Afr. trop.
Warmingii, *Michell, in Kjoeb. Vidensk. Meddel.* (1875) 68.—Bras.

AESCULUS, Linn. Syst. ed. II. 32 (1740). *SAPINDACEÆ*, Benth. & Hook. f. i. 386.
Billia, Peyritsch, in Bot. Zeit. xvi. (1858) 153.
Calothyrsus, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 62.
Æsculus, Linn. Gen. ed. I. 109 (1737).
Hippocastanum, Tourn. ex Rupp. Fl. Jen. ed. Hall. 291. (1745).
Isypon, Rafin. Fl. Tellur. iv. 73 (1836).
Macrothyrsus, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 61.
Nemeropsis, Rafin. Alsog. Am. 68 (1838).
Oesculus, Neck. Elem. ii. 282 (1790).
Ozotis, Rafin. Alsog. Am. 71 (1838).
Pavia, Boerh. ex Mill. Gard. Dict. ed. VI. (1752).
Paviana, Rafin. Fl. Ludov. 87 (1817).
Putzeysia, Planch. & Linden, in Linden, Cat. n. 22 (1867).
alba, *Rafin. Alsog. Am.* 69.—Am. bor.
arguta, *Buchl. in Proc. Acad. Phil.* (1860) 448.—Am. bor.
asplenifolia, Hort. ex Loud. Encyc. Trees, 124 = A. Hippocastanum.
assamica, Griff. Notul. iv. 540 = punduana.
barbata, *Rafin. Alsog. Am.* 70.—Am. bor.
californica, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 251.—Calif.
carnea, Wats. Dendr. Brit. t. 121 = glabra.
Castanea, Gilib. Fl. Lituan. ii. 197 = A. Hippocastanum.

AESCULUS:—
chinensis, *Bunge, in Mém. Sav. étr. Pétersb.* ii. (1832) 84.—China.
discolor, Pursh, Fl. Am. Sept. i. 255 = flava.
disamilla, *Blume, Rumphia*, iii. 195.—Japon.
echinata, Muhl. Cat. Fl. Am. Sept. 38 = glabra.
flava, *Ait. Hort. Kew.* i. 494.—Am. bor.
glabra, *Willd. Enum. Hort. Berol.* 405.—Am. bor. Hippocastanum, *Linn. Sp. Pl.* 344.—Turc.
humilis, Lindl. Bot. Reg. t. 1018 = Æs. Pavia?
hybrida, *DC. Hort. Monsp.* (1818) 75 = flava.
indica, *Colebr. ex Wall. Cat.* n. 7181.—Reg. Himal.
intermedia, *Steud. Nom.* ed. II. i. 32.—Am. bor.
irvingata, *Rafin. Alsog. Am.* 70.—Am. bor.
livida, *Steud. Nom.* ed. II. i. 32.—Am. bor.
lutea, Wangh. in Schr. Berl. Ges. Naturf. Fr. viii. (1788) 138. t. 6 = flava.
Lyonii, Lond. Arb. Brit. i. 466 = A. Pavia.
macrostachya, Michx. Fl. Bor. Am. i. 220 = parviflora.
macrostachya, Pers. Syn. i. 408 = parviflora.
Mmmeigeri, Hort. ex C. Koch, Dendrol. i. 596 = A. Hippocastanum.
mexicana, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* i. 218.—Mexic.
muricata, *Rafin. Alsog. Am.* 68.—Am. bor.
neglecta, Lindl. Bot. Reg. t. 1009 = flava.
ochroleuca, *Rafin. Alsog. Am.* 68.—Am. bor.
odorata, *F. G. Dietr. Lexik. Gaertn. Nachtr.* i. 118.—Am. bor.
ohioensis, Michx. Arb. For. Am. Sept. iii. 242 = glabra.
pallida, Willd. Enum. Hort. Berol. 406 = glabra.
Parryi, *A. Gray, in Proc. Am. Acad. xvii.* (1881–82) 200.—Am. bor.
parviflora, *Walt. Fl. Carol.* 128.—Am. bor.
Pavia, *Linn. Sp. Pl.* 344.—Am. bor.
procera, Salisb. Prod. 380 = A. Hippocastanum.
punduana, *Wall. Cat.* n. 1189.—Reg. Himal.
rosea, Hort. ex Loud. Encyc. Trees, 127 = glabra.
rubella, Wender. Ind. Sem. Marb. (1858) 1. et 4 = glabra?
rubescens, Tausch, in Flora, xxi. (1838) 1. Beibl. 79 = A. Pavia?
rubicunda, Lodd. Bot. Cab. t. 1242 = glabra.
rubra, Pers. Syn. i. 408 = A. Pavia.
septenata, Stokes, Bot. Mat. Med. iii. 337 = A. Hippocastanum.
turbinata, *Blume, Rumphia*, iii. 195.—Japon.
verrucosa, *Rafin. Alsog. Am.* 69.—Am. bor.
versicolor, Wender. Ind. Sem. Hort. Marb. (1853) 4 = A. Pavia?
Watsoniana, D. Dietr. Syn. ii. 1225 = glabra.
Whitleyi, Hort. ex C. Koch, Dendrol. i. 511 = A. Pavia.

AETHEILEMA, R. Br. Prod. 478 (1810) = **Phaylopsis**, Willd. (Acanth.).

alopecuroides, Spreng. Syst. ii. 826 = Lepidagathis alopecuroides.
anisophyllum, E. Mey. ex Nees, in DC. Prod. xi. 262 = P. longifolia.
glutinosum, Steud. Nom. ed. II. i. 32 = P. parviflora.
Haenkei, Nees, in Mart. Fl. Bras. ix. 72 = Lepidagathis alopecuroides.
imbricatum, R. Br. Prod. i. 478 = P. longifolia.
javanicum, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 129.—Java.
longifolium, Spreng. Syst. Cur. Post. 236 = P. longifolia.
micranthum, Benth. in Hook. Nig. Fl. 480.—Afr. trop.
mucronatum, Griff. Notul. iv. 157 = P. parviflora.
parviflorum, Spreng. Syst. ii. 826 = P. parviflora.
Rothii, Steud. Nom. ed. II. i. 32 = P. parviflora.
reniforme, Nees, in Wall. Pl. As. Rar. iii. 94 = P. parviflora.
rigidum, Bartl. ex Nees, in DC. Prod. xi. 242 = Barleria solanifolia.
rupestre, Nees, l. c. 262.—Madag.

AETHEOLAENA, Cass. in Dict. Sc. Nat. xlviii. 450 (1827) = **Semecio**, Linn. (Compos.).
involucrata, Cass. l. c. 451 = S. involucratus.

AETHEONEMA, Reichb. Nom. 128 (1841) = **Geertnera**, Lam. (Logan.).

AETHEOPAPPUS, Cass. in Dict. Sc. Nat. i. 250 (1827) = **Centaurea**, Linn. (Compos.).
Balansæ, Boiss. Fl. Orient. iii. 604 = C. Balansæ.
pulcherrimus, Cass. in Dict. Sc. Nat. li. 52 = C. pulcherrima.

AETHEORHIZA, Cass. in Dict. Sc. Nat. xlviii. 426 (1827) = **Crepis**, Linn (Compos.).
bulbosa, Cass. l. c. = C. bulbosa.
montana, Willd. in Orst. Bot. Zeitschr. xxv. (1875) 110 = C. montana.

AETHEORRHIZA, Reichb. Consp 98 (1828) = praec.

AETHERIA, Endl Gen. 214 (1837) = **Hetaeria**, Blume (Orchid.).

AETHIONEMA, R. Br. in Ait. Hort. Kew ed. II. iv. 80 (1812). *CRUCIFERAE*, Benth. & Hook. f. i. 58.
Diastrophis, Fisch. & Mey. Ind Sem. Hort Petrop. ii. 35 (1835).
Disynoma, Rafin Fl. Tellur. iii 81 (1836).
Eirionema, Brongn. Enum. Gen. 92 (1843).
Eunomia, DC. Syst. ii. 555 (1821).
Oethionema, Knowl & Westc. Flor. Cab. i. (1837) 41.
arabicum, Andrz. ex DC. Syst ii. 560 = cappadocium.
armenum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 191.—Armen.
Balansae, Boiss. Diagn. Ser. II. vi. 20 (= *Crenularia glaucescens*).
banaticum, Janka, in Linnaea, xxx. (1859–60) 558 = saxatile.
Beyrichii, *Tausch, in Flora*, xix. (1836) 411.—Persia.
Bourgaei, *Boiss. Fl. Orient.* i. 344.—Syria.
Buxbaumii, DC. Syst. ii. 560 = cappadocicum.
caespitosum, Boiss. Fl. Orient. i. 343 = Iberidella caespitosa.
campylopterum, *Boiss. l. c.* 358 —Syria.
capitatum, *Boiss. & Balansa, Diagn.* Ser. II. v. 43.—As. Min.
cappadocium, *Spreng. Nov. Prov.* i. 2.—Orlens.
cardiophyllum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 43.—As. Min.
chlorsefolium, *Boiss. Fl. Orient.* i. 344.—As. Min.
clandestinum, *Delile, in Ann. Sc. Nat.* Sér II. vii (1837) 286.—Hab. ?
cordatum, *Boiss. Fl. Orient* i. 350.—Oriens.
cordifolium, *DC Syst.* ii. 561.—Orlens
coridifolium, Hort. ex Boiss Diagn. Ser. I. viii. 43 = pulchellum
cristatum, *DC Syst* ii. 560.—Orens.
diastrophis, *Bunge, Del Sem. Dorp.* (1841) 7.—Armen.
elongatum, *Boiss. Diagn.* Ser. I. vi. 17 —Persia
fimbriatum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 198.—Persia.
Fischeri, *Tausch, in Flora*, xix (1836) 410.—Sibir.
gracile, *DC. Syst.* ii. 559.—Carniol.
graecum, *Boiss. & Sprun. Diagn.* Ser I. vi. 16.—Graecia.
grandiflorum, *Boiss. & Hohen. l. c.* viii. 42 —Persia.
heterocarpum, *J. Gay, in Fisch. & Mey. Ind. Sem. Petrop.* iv. 1, nomen.—Syria.
heterophyllum, Boiss. Fl. Orient. i. 343 = Iberidella heterophylla.
iberideum, *Boiss. l. c.* i. 351.—Oriens.
laxum, *Boiss. Diagn.* Ser. II. v. 45.—Cilicia.
leptocarpum, *Delile, in Ann. Sc. Nat.* Sér. II. vii. (1837) 285.—Hab.?
lilgeosum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 348. —As. Min.
nembranaceum, *DC. Syst.* ii. 561.—Persia.
monospermum, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 80.—Hispan.
moricandianum, *Boiss. Fl. Orient.* i. 351.—As. Min.
oppositifolium, *Boiss. l. c.* 344.—Syria.
ovalifolium, *Boiss. l. c.* 351.—Oriens.
polygaloides, *DC. Syst.* ii. 562.—Ins. Scios.
polygaloides, Ledeb. Fl. Ross. i. 209 = armenum.
pulchellum, *Boiss. & Huet, Diagn.* Ser. II. v. 43.—Oriens.
pyrenaicum, *Boutigny, in Bull. Soc. Bot. Fr.* iv. (1857) 777.—Mont. Pyren.
rotundifolium, *Boiss. Fl. Orient.* i. 344.—Reg. Caucas.
rubescens, *Boiss. l. c.* 343.—As. Min.
sagittatum, Boiss. l. c. 342 = Iberidella sagittata.
salmasium, *Boiss. l. c.* 350.—Persia.
saxatile, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 80.—Europ. austr.
schistosum, *Boiss. & Kotschy, Diagn.* Ser. II. v. 42.—As. Min.
speciosum, *Boiss. & Huet, l. c.* 44.—Oriens.

AETHIONEMA:—
spicatum, *Post, in Hook. Ic. Pl.* xv. 62, t. 1478.—Syria.
stenopterum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 192.—Persia.
stylosum, *DC. Syst.* ii. 562.—Syria.
subulatum, *Boiss. Fl. Orient.* i. 349.—As. Min.
Szovitsii, *Boiss. l. c.* 346.—Persia.
tenue, Boiss. l. c. 348 = Iberdella tenuis.
theoilfolium, *Boiss. & Heldr. Diagn.* Ser. I. viii. 44.—As. Min.
Thomasianum, *Y. Gay, in Ann. Sc. Nat.* Sér. III. iv. (1845) 81.—Europ.
trinervium, Boiss. Fl. Orient. i. 342 = Iberidella trinervia

AETHIOPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii (1869) 184 = **Salvia**, Tourn. (Labiat.).
Sclarea, Fourr. l. c. = S. Sclarea.
vera, Fourr. l. c. = S. Aethiopis.

AETHONIA, D. Don, in Edinb. N. Phil. Journ. (Mar 1829) 309 = **Tolpis**, Adans. (Compos.).
filiformis, D. Don, l. c. = T. filiformis.
fruticosa, D. Don, l. c. = T. fruticosa.
lagopoda, D. Don, l. c. = T. lagopoda.

AETHUSA, Linn. Sp. Pl. 256 (1753). *UMBELLIFERAE*, Benth. & Hook. f. i. 907.
Cynapium, Kit. ex Rupp. Fl. Jen. ed. Hall. 284 (1745).
Ethusa, Linn. Syst. ed. I. (1735).
Ammi, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Ammi majus.
Bunium, Murr. Syst. Veg. ed. XIII. 286 = Carum Bunius.
capillacea, Nutt. ex DC. Prod. iv. 106 = Apiastrum angustifolium.
Cicuta, Neck. Delic. Gallo-Belg. i. 146 = A. Cynapium.
cynapioides, Bieb. Fl. Taur. Cauc. i. 227 = A. Cynapium.
Cynapium, *Linn. Sp. Pl.* 256.—Europ.; Orient.
cynica, Dulac, Fl. Hautes-Pyr. 358 = A. Cynapium.
divaricata, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Apium divaricatum.
elata, Friedl. ex Fisch. Cat. Hort. Gorenk. (1813) 45 = A. Cynapium.
fatua. Ait. Hort. Kew. i. 302 = Carum verticillatum.
leptophylla, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Apium leptophyllum.
Meum, Murr. Syst. Veg. ed. XIII. 286 = Meum athamanticum.
micrantha, Opiz, ex Nym. Consp. 297 = A. Cynapium.
montana, Lam. Fl. Fr. iii. 649 = Carum Bunus.
mutellina, Lam. l. c. 443 = Meum mutellina.
petroselinifolia, Gilib. Fl. Lituan. ii 25 = A. Cynapium.
pinnata, Nutt. ex DC. Prod. iv. 141 = Cynosciadium pinnatum.
segetalis, Boenn. Prod. Fl. Monast. 85 = A. Cynapium.
tenuifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 518 = A. Cynapium.
tenuifolium, Salisb. Prod. 166 = Meum athamanticum.
toxicaria, Salisb. l. c. = A. Cynapium.

AETIA, Adans., Fam. ii. 84 (1763) = **Combretum**, Linn.

AETOXICON, Ruiz & Pav. Prod. 131. t. 20 (1794). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 265.
Argotoxicon, Molina, Sagg. Chile. ed. II. 298 (1810).
punctatum, *Ruiz & Pav. Syst. Veg.* 260.—Chili.

AFARCA, Rafin. Sylva Tellur. 30 (1838) = **Sageretia**, Brongn. (Rhamn.).
parviflora, Rafin. l. c. = S. Michauxii.

AFFONSEA, A. St. Hil. Voy. Dism. i. 387 (1833). *LEGUMINOSAE*, Benth. & Hook. f. i. 599.
bullata, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 77. —Bras.
comosa, Benth. l. c. 76. t. 1 = jugiandifolia.
dassiflora, *Benth. l. c.*—Bras.
jugiandifolia, *A. St. Hil. Voy. Dism.* i. 388.—Bras.

AFZELIA, J. F. Gmel. Syst. 927 (1791) = **Swymeria**, Pursh (Scrophul.).
casuioides, J. F. Gmel. l. c. = S. tenuifolia.

AFZELIA, Sm. in Trans. Linn. Soc. iv. (1798) 221. *LEGUMINOSAE*, Benth. & Hook. f. i. 580.
Intsia, Thou. Gen. Nov. Madag. 22 (1806).
africana, *Sm. l. c.*—Afr. trop.
attenuata, *Klotzsch, in Peters, Reise Mossamb. Bot.* i. 20.—Afr. or. trop.
bijuga, *A. Gray, Bot. U. St. Expl. Exped.* i. 467.—Ins. Pacif.
bijuga, Spreng. Syst. iv. Cur. Post. 170 = Erioglossum cauliflorum.
bracteata, *Vog. ex Benth. in Hook. Ic. Pl.* t. 790.—Afr. trop.
cambodeensis, *Hance, in Journ. Bot.* xiv. (1876) 358.—Cambod.
coriacea, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 275.—Ind. or.
cuanzensis, Oliver, Fl. Trop. Afr. ii. 302 = quanzensis.
discolor, *Hort. ex Steud. Nom.* ed. II. i. 33.—Afr. trop.
grandis, Hort. ex Loud. Hort. Brit. 168 = Erythrophleum guineense.
madagascariensis, Baill. Hist. Pl. ii. 113 = bijuga.
palembanica, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 275.—Ind. or.
Pancovia, DC. Prod. ii. 507 = Erioglossum cauliforum.
Petersiana, Klotzsch, in Peters, Reise Mossamb. Bot. i. 19 = quanzensis.
quanzensis, *Welw. Apont.* 586.—Angola.
retusa, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) 11. 78.—Malacca.
rhomboidea, *Vidal, Phanerog. Cuming. Philipp.* 26, 110.—Ins. Philipp.
splendens, Hort. ex Loud. Hort. Brit. 168 = Carapa procera.

AGALINIS, Rafin. New Fl. Am. ii. 61 (1836) = **Gerardia**, Linn. (Scrophul.).
aphylla, Rafin. l. c. 65 = G. aphylla.
corymbosa, Rafin. l. c. 63 = G. Plukenetii.
fasciculata, Rafin. l. c.—G. purpurea.
filifolia, Rafin. l. c. 65 = G. filifolia.
longifolia, Rafin. l. c. 62 = G. aspera?
maritima, Rafin. l. c.—G. maritima.
microphylla, Rafin. l. c. 65.—Am. bor.
obtusifolia, Rafin. l. c. 64 = Aulaya obtusifolia?
palustris, Rafin. l. c. 62 = G. purpurea.
perennis, Rafin. l. c. 63 = G. linifolia.
Plukenetii, Rafin. l. c. = G. Plukenetii.
setacea, Rafin. l. c. 64 = G. setacea.
tenuifolia, Rafin. l. c. = G. tenuifolia.
virgata, Rafin. l. c. 62 = Seymeria virgata?

AGALLIS, Phil. in Linnaea, xxxiii. (1864–65) 12. *CRUCIFERAE*.
nuntana, *Phil. l. c.*—Chili.

AGALLOCHUM, Lam. Encyc. i. 48 (1783) = **Aquilaria**, Lam. (Thymel.).
officinarum, Lam. l. c. = Aquil. malaccensis.
prunisatinianum, Lam. l. c. 46.—As. or.
sylvestre, Lam. l. c. 49.—Ins. Timor.

AGALLOSTACHYS, Beer, Bromel. 35 (1857) = **Bromelia**, Linn.
anticacantha, Beer, l. c. 37 = B. fastuosa.
chrysantha, Beer, l. c. 35 = B. chrysantha.
Commeliniana, Beer, l. c. 39 = B. fastuosa.
fastuosa, Beer, l. c. 36 = B. fastuosa.
lacinisona, C. Koch, Wochenschr. ii. (1859) 140 = B. laciniosa.
Pinguin, Beer, Bromel. 36 = B. Pinguin.
sylvestris, Beer, l. c. 35 = B. sylvestris.
variegata, Beer, l. c. 38 = Billbergia speciosa.

AGALMA, Miq. Fl. Ind. Bat. i. 1. 751 (1855) = **Heptapleurum**, Gaert. (Araliac.).
aeculifolium, Seem. in Journ. Bot. ii. (1864) 296 = Brassaiopsis aculeata.
aromaticum, Seem. l. c.—Java.
elatum, Seem. l. c. = H. elatum.
glaucum, Seem. l. c. 299 = H. glaucum.
Griffithii, Seem. l. c. = H. elatum.
Horsfieldii, Seem. l. c. 298.—Java.
hoolanse, Seem. Revis. Hedenac. 108 = H. hoolanse.
lucuosum, Seem. in Journ. Bot. ii. (1864) 299 = H. octophyllum.
octophyllum, Seem. l. c. 296 = H. octophyllum.
racemosum, Seem. l. c. = H. racemosum.
reticulosum, Seem. l. c. 299.—Borneo.

AGALMA :—
rostratum, Seem. in Journ. Bot. ii. (1864) 298 = H. rostratum.
rugosum, Miq. Fl. Ind. Bat. i. 1. 752. t. 11.—Malaya.
simillimum, Miq. l. c.—Malaya.
tomentosum, Seem. in Journ. Bot. ii. (1864) 298 = H. impressum.
vitiense, Seem. Fl. Vit. 116.—Ins. Fiji.

AGALMA, Steud. Nom. ed. II. i. 33 (1840) = **Bonohua**, Tourn. (Compos.).

AGALMANTHUS, Hombr. & Jacquinot, ex Decne Voy. Astrol. et Zel. 78 (1845) = **Metroseideros**, Banks (Myrtac.).
umbellata, Hombr. & Jacquinot, ex Decne. l. c = M. lucida.

AGALMYLA, Blume, Bijdr. 766 (1826). *GESNERACEAE*, Benth. & Hook. f. ii. 1014
ORITHALIA, Blume, Fl. Jav. Pracf. 6 (1828).
ORYTHIA, 'Blume,' ex Endl. Gen. Suppl. i. 1408 (1841).
angustifolia, *Miq. Fl. Ind. Bat.* ii. 723.—Malaya.
asperifolia, Blume, Bijdr. 767 = Dichrotrichum asperifolium.
Beccarii, *C. B. Clarke*, *in DC. Monog. Phan.* v. 57.—Sumatra.
longistyla, Carr. in Rev. Hortic. (1878) 271 = staminea.
staminea, *Blume*, *Bijdr.* 767.—Java.
tuberculata, *Hook. f. Ic. Pl.* t. 897.—Borneo,

AGALOMA, Rafin. Fl. Tellur. iv. 116 (1836) = **Euphorbia**, Linn.
angustifolia, Rafin. l. c. 117 = E. angustifolia ?
corollata, Rafin. l. c. = E. corollata.
graminifolia, Rafin. l. c. = E. granulata.

AGANIPPEA, Mor. & Sesse, ex DC. Prod. vi. 3 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 358.
HELIOGENES, Benth. Pl. Hartw. 42 (1840).
bellidiflora, *Moç. & Sesse, ex DC. Prod.* vi. 3.—Mexic. dentata, *DC. l. c.*—Mexic.

AGANISIA, Lindl. in Bot. Reg. xxv. (1839) Misc. 45; xxvi. t. 32 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 544.
KORLLENSTEINIA, Reichb. f. in Bonplandia, ii. (1854) 17.
caerulea, *Reichb. f. in Gard. Chron.* N. S. vi. (1876) ii. 256.—Bras.
cyanea, Reichb. f. in Nov. Act. Nat. Cur. xxxv. ii. (1876) 13. t. 4 = Acacallis cyanea.
fimbriata, *Reichb. f. in Gard. Chron.* N. S. ii. (1874) ii. 459.—Guian.
lepida, *Linden & Reichb. f. in Nov. Act. Nat. Cur.* xxxv. ii. (1876) 15. t. 5.—Am. trop.
Oliveriana, *Reichb. f. in Gard. Chron.* N. S. ix (1878) i. 558.—Bras.
pulchella, *Lindl. in Bot. Reg.* (1839) *Misc.* 45; xxvi. (1840) t. 32.—Guian.

AGANON, Rafin. Sylva Tellur. 161 (1838) = **Calliocarpa**, Linn. (Verben.).
umbellata, Rafin. l. c. = C. arborea.

AGANOPE, Miq. Fl. Ind. Bat. i. 1. 151 (1855) = **Derris**, Lour. (Leguminh.).
floribunda, Miq. l. c. = D. thyrsiflora.
marginata, Miq. l. c. 152 = D. marginata.
microphylla, Miq. l. c.—Java.
submontana, Miq. l. c. Suppl. i. 399.—Sumatra.
sumatrana, Miq. l. c. 400.—Sumatra.

AGANOSMA, G. Don, Gen. Syst. iv. 77 (1837). *APOCYNACEAE*, Benth. & Hook. f. ii. 717.
GANOSMA, Decne. in DC. Prod. vii. 625 (1844), aphalm.
acuminata, G. Don, l. c. = marginata.
affinis, G. Don, l. c. = Ichnocarpus ovatifolius.
Blumei, *A. DC. in DC. Prod.* viii. 483.—Java.
Blumei, Wight, Ic. t. 1305 = caryophyllata.
calycina, *A. DC. in DC. Prod.* viii. 482.—Ind. or.
caryophyllata, *G. Don, Gen. Syst.* iv. 77.—Ind. or.
concanensis, Hook. Ic. Pl. v. t. 941 = Parsonsia spiralis.
conforta, G. Don, in Loud. Encyc. Pl. Suppl. ii. 1312 = cymosa.
cymosa, *G. Don*, *Gen. Syst.* iv. 77.—Ind. or.

AGANOSMA :—
Doniana, Wight, Ic. t. 1306 = cymosa.
elegans, G. Don, Gen. Syst. iv. 77 = cymosa.
Edithae, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 227 = Cryptolepis elegans.
euloba, Miq. Fl. Ind. Bat. ii. 447 = acuminata.
gracilis, *Hook. f. Fl. Brit. Ind.* iii. 664.—Reg. Himal.
laevigata, J. Grah. Cat. Bomb. Pl. 118 (= Parsonsiae sp.?) —Ind. or.
laevis, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 385 = Anodendron laeve.
macrocarpa, A. DC. in DC. Prod. viii 484 = marginata
marginata, *G. Don, Gen Syst.* iv. 77.—Ind. or.
Roxburghii, G. Don, l. c.—calycina.
velutina, *A. DC. in DC. Prod.* viii 484.—Malaya.
Wallichii, G. Don, Gen Syst. iv. 77 = calycina.

AGAPANTHUS, L'Hérit. Sert. Angl. 17 (1788) *LILIACEAE*, Benth. & Hook f. iii 798.
ABUMON, Adans. Fam. ii. 54 (1763).
MADHUIA, Dahl, Obs. Syst. Linn. 35 (1787).
africanus, Hoffmgg. Verz. Pfl 35 = umbellatus.
ensifolius, Willd. Sp. Pl. ii. 48 = Massonia ensifolia
medius, Lodd. ex Steud. Nom. ed. II. i. 33 = umbellatus.
minor, Lodd. Bot. Cab. t. 42 = umbellatus.
multiflorus, Willd. Enum. Hort. Berol. 358 = umbellatus.
praecox, Willd. l. c. = umbellatus
tuberosus, Linn. ex [DC. in] Red. Liliac. i. 6 = umbellatus.
umbellatus, *L'Hérit. Sert. Angl.* 18.—Afr. austr.
umbelliferus, Pers. Encyc. Suppl. i. 155 = umbellatus.
variegatus, Hort. ex Steud. Nom. ed. II. i. 33 = umbellatus.

AGAPATEA, Steud. in Bot. Zeit. xiv. (1856) 201 = **Distichia**, Nees & Meyen (Juncac.).
filamentosa, Buchen. in Bremen Abh. iv. (1874) 124 = Ind. filamentosa.
peruviana, Steud. in Bot. Zeit. xiv. (1856) 200 = D. muscoides.

AGAPETES, G. Don, Gen. Syst. iii. 862 (1834). *VACCINIACEAE*, Benth. & Hook. f. ii. 571.
CALLIULA, Klotzsch, in Linnaea, xxiv. (1851) 28.
PAPHIA, Seem. in Journ. Bot. ii. (1864) 77.
Acosta, Dun. in DC. Prod. viii. 556 = Vaccinium Acosta.
acuminata, D. Don, ex G. Don, Gen. Syst. iii. 862 = Corallobotrys acuminata.
amblyorudis, *Becc. Malesia*, i. (1878) 208.—Malaya.
angulata, *Benth. & Hook. f. Gen.* ii. 571.—Ind. or.
arborea, Dun. in DC. Prod. vii. 556 = Vaccinium Leschenaultii.
auriculata, *Benth. & Hook. f. Gen.* ii. 571.—Ind. or.
bracteata, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 448.—Ind. or.
buxifolia, *Nutt. ex Hook. f. Bot. Mag.* t. 5012 —Reg. Himal.
campanulata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 449.—Ind. or.
coriacea, G. Don, Gen Syst. iii. 863 = Vaccinium Korthalsii.
cuneifolia, G. Don, l. c. = Vaccinium cuneifolium.
discolor, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 448.—Reg. Himal.
elliptica, G. Don, Gen. Syst. iii. 862 = Vaccinium ellipticum.
floribunda, G. Don, l. c. = Vaccinium Teysmanni.
glabra, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 448.—Ind. or.
grandiflora, *Benth. & Hook f. Gen.* ii. 571.—Ind. or.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 449.—Reg. Himal.
hirsuta, Klotzsch, in Linnaea, xxiv. (1851) 38 = setigera.
laurifolia, G. Don, Gen Syst. iii. 862 = Vaccinium laurifolium.
linearifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 449.—Reg. Himal.
Lobbii, *C. B. Clarke*, l. c. 448.—Ind. or.
lonchnifiora, *D. Don, ex G. Don, Gen. Syst.* iii. 862.—Ind. or.
lucida, G. Don, l. c. 863 = Vaccinium luciduin.
macrantha, *Benth. & Hook f. Gen.* ii. 571.—Ind. or.
macrophylla, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 446.—Reg. Himal.
macrostemon, *C. B. Clarke*, *l. c.* 448.—Burma.
madagascariensis, *Dun. in DC. Prod.* viii. 555.—Madag.

AGAPETES :—
meliphagidum, *Becc. Malesia*, i. (1878) 208.—Malaya.
microphylla, *Yungh. in Nat. en Geneesk. Arch. Neerl Ind.* ii. (1845) 86.—Java.
minista, *Benth. & Hook. f. Gen.* ii. 571.—Reg. Himal.
mitraeicides, *Hook. f. ex C. B. Clarke, in Hook. f. Fl Brit. Ind.* iii. 447.—Reg. Himal.
myrtoidea, G Don, Gen. Syst. iii. 863 = Vaccinium myrtoides.
Myosmeles, *Becc. Malesia*, i. (1878) 209.—Malaya.
nana, *Benth. & Hook f. Gen.* ii. 571.—Reg. Himal.
Nuttallii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 445.—Reg. Himal.
obliqua, Benth. & Hook. f. Gen. ii. 571 = setigera
obovata, *Benth. & Hook. l. c* 572.—Reg Himal.
odontocera, Benth & Hook. f. l. c. 571 = variegata.
Parishii, *C. B Clarke, in Hook f. Fl. Brit. Ind* iii 445.—Ind. or.
pilifera, *Hook f ex C. B. Clarke, l. c.* 448.—Reg Himal
pulcherrima, Benth. & Hook. f Gen. ii. 571 = variegata.
ruca, *Yungh. in Nat. en Geneesk. Arch. Neerl. Ind.* ii. (1845) 85.—Java ?
salicifolia, *C. B. Clarke in Hook. f. Fl. Brit. Ind.* iii 445.—Reg. Himal.
salscifolia, Gardn. ex Thw. Enum. Pl. Zeyl. 170 = Vaccinium Leschenaultii.
saligna, *Benth. & Hook. f. Gen.* ii. 571.—Ind. or.
serrata. G. Don, Gen. Syst. iii. 862 = Vaccinium serratum.
setigera, *D. Don, ex G Don, l. c.*—Ind. or.
Sprengelii, G. Don, l. c. = Vaccinium Doniannm.
symplocifolia, D. Don, ex G Don, l. c. = Vaccinium Leschenaultii.
vaccinacea, Dun. in DC. Prod vii. 554 = Vaccinium serratum.
variegata, Dun. l. c. = setigera.
variegata, *D. Don, ex G. Don, Gen. Syst.* iii. 862.—Ind. or.
varingiaefolia, G. Don, l. c = Vaccinium varingiaefolium.
verticillata, D. Don, ex G. Don, l. c. = setigera.
vulgaris, Jungh. in Nat. en Geneesk. Arch. Neerl. Ind. ii. (1845) 84 = Vaccinium ellipticum
Wallichiana, Klotzsch, in Linnaea, xxiv. (1851) 38 = setigera.

AGARDHIA, Spreng. Syst. i. 4, 17 (1825) = **Qualea**, Aubl. (Vochys.).
cryptantha, Spreng. l. c. 17 = Q cryptantha.
grandiflora, Spreng l. c. = Q grandiflora ?

AGARISTA, DC. Prod. v 569 (1836) = **Coreopsis**, Linn. (Compos.).
calliopsidea, DC. l. c. = C calliopsidea.

AGARISTA, D. Don, in G. Don, Gen. Syst. iii. 537 (1834). *ERICACEAE*, Benth. & Hook. f. ii. 586.
AMECHANIA, DC. Prod. vii 578 (1838).
anasimonsans, *G. Don, l. c.* iii. 538.—N Granat
bracamorensis, *G. Don, l. c.* 537.—N. Granat.
buxifolia, G. Don, l. c = Agauria buxifolia.
chlorantha, *G. Don, l. c.* 538.—Bras.
eriophyla, *G. Don, l. c.*—Bras.
encalyptoides, *G. Don, l. c.* 537.—Bras.
ilicifolia, *G. Don, l. c.* 538.—Peruv.
Ilatiaiae, *Wawra, It. Princip. Coburg.* i. 78.—Bras.
multiflora, *G. Don, Gen. Syst.* iii. 537.—Bras.
pentifolia, *G. Don, l. c.* 538.—Bras.
pummularia, *G. Don, l. c* 537.—Bras.
oleifolia, *G. Don, l. c.* 538.—Bras.
Plutrix, *G. Don, l. c.*—Bras.
Pohlii, *G. Don, l. c.* 537.—Bras.
pulchella, *G. Don, l. c.* 538.—Bras.
pulchra, *G. Don, l. c.* 537.—Bras.
pyrifolia, G. Don, l. c. 538 = Agauria salicifolia.
salicifolia, G. Don, l. c. 537 = Agauria salicifolia.
serrulata, *G. Don, l. c.* 538.—Bras.
Sprengelii, *G. Don, l. c.* 537.—Bras.
subrotunda, *G. Don, l. c.*—Bras.

AGASSIZIA, Chav. Monog. Antirrh. 180. t 11 (1830) = **Galvezia**, Domb. (Scrophul.).
limensis, Chav. l. c. = G. limensis.

AGASSIZIA, Gray & Engelm. in Proc Am. Acad. i. (1847) 46 = **Gaillardia**, Fouger. (Compos.).
suavis, A. Gray & Engelm. l. c. 50 = G. simplex.

AGASSIZIA, Spach, Hist. Vég. Phan. iv. 347 (1835) = **Oenothera**, Linn. (Onagrar.).
cheiranthifolia, Spach, l. c. 348 = O. Agassizia.

AGASTA, Miers, in Trans. Linn. Soc. Ser. II. i. 59 (1878) = **Barringtonia**, Forst. (Myriac.).
asiatica, Miers, l. c. 61 = B. speciosa.
indica, Miers, l. c. 63 = B. speciosa.
splendida, Miers, l. c. 60 = B. speciosa.

AGASTACHYS, R. Br. in Trans. Linn. Soc. x. (1811) 188. *PROTEACEAE*, Benth. & Hook. f. iii. 176. odorata, *R. Br. l. c.*—Tasmania.

AGASTIANIS, Rafin. in New Fl. Am. iii. 85 (1836) = **Bophorus**, Linn. (Legum.).
secundiflora, Rafin. l. c. 86 = S. secundiflora.

AGASYLLIS, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1813) 259 = **Bilax**, Scop. (Umbellif.).
caucasica, Spreng. l. c. = S. caucasicum.
Galbanum, Spreng. l. c. = Peucedanum Galbanum.
latifolia, Bobs. Fl. Orient. ii. 980 = S. caucasicum.
salsa, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 259 = Peucedanum redivivum.

AGATEA, A. Gray, in Proc. Am. Acad. ii. (1852) 328 = **Agation**, Brongn. (Violar.).
violaris, A. Gray, l. c. 324 = Agation violare.

AGATEA, Rich, ex A. Gray, Bot. U. S. Expl. Exped. i. 609 (1854) = **Haplopetalum**, A. Gray (Rhisoph.).

AGATHAEA, Cass. in Bull. Soc. Philom. (1815) 176; Dict. Sc. Nat. i. Suppl. 77. t. 89 (1817) = **Felicia**, Cass. (Compos.).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 358 = F. Richardi.
amelioides, DC. Prod. v. 225 = Aster rotundifolius.
amoena, Sch. Bip. ex Walp. Rep. ii. 956 = Aster amoenus.
barbata, DC. Prod. v. 224 = Aster barbatus.
Bergeriana, DC. l. c. 223 = Aster Bergerianus.
brevifolia, DC. l. c. 222.—Afr. austr.
coelestis, Cass. Opusc. i. 8 = Aster rotundifolius.
corymbosa, Turcz. in Bull. Soc. Nat. Mosc. xxiv.(1851) ii. 60 = Gymnostephium corymbosum.
Gerstata, A. Rich. Tent. Fl. Abyss. i. 354 = F. Richardi.
diffusa, DC. Prod. v. 224 = Aster demissus.
Echlonis, Nees, Gen. et Sp. Ast. 362 = Aster Ecklonis.
elongata, Nees, l. c. 266 = Aster elongatus.
hirta, DC. Prod. v. 224 = Aster hirtus.
hispida, DC. l. c. = Aster elongatus.
Kraussii, Sch. Bip. ex Walp. Rep. ii. 95 = Aster Kraussii.
leiocarpa, DC. Prod. v. 225 = Aster leiocarpus.
linifolia, G. Don, in Loud. Hort. Brit. 349 = Enryops linifolius.
microphylla, Cass. in Bull. Soc. Philom. (1817) 183 = Aster aethiopicus.
natalensis, Sch. Bip. ex Walp. Rep. ii. 956 = Aster natalensis.
rotundifolia, Nees, Gen. et Sp. Ast. 258 = Aster rotundifolius.
scabrida, DC. Prod. v. 225 = Aster scabridus.
serrata, Nees, Gen. et Sp. Ast. 250 = Aster serratus.
spathulata, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. x. 48 = Amellus annuus.
spathulata, Kunze, Pugill. Pl. Ined. i. 9 = Amellus hispidus.
stricta, DC. Prod. v. 224 = F. adfinis.
strigosa, Nees, Gen. et Sp. Ast. 262 = Aster strigosus.
tenera, DC. Prod. v. 224 = Aster tener.
tricolor, Nees, Gen. et Sp. Ast. 267 = Aster tricolor.
Zeyheri, Nees, l. c. 260 = Aster linifolius.

AGATHELPIS, Cholay, in Mém. Soc. Phys. Genèv. ii. (1824) 85, 89. *SELAGINEAE*, Benth. & Hook. f. ii. 1129.
aduncа, *E. Mey. Comm. Pl. Afr. Austr.* i. 252.—Afr. austr.
angustifolia, *Choisy, in Mém. Soc. Phys. Genèv.* ii. (1824) 85.—Afr. austr.
brevifolia, *E. Mey. Comm. Pl. Afr. Austr.* i. 253.—Afr. austr.
mucronata, *E. Mey. l. c.*—Afr. austr.

AGATHELPIS :—
nitida, *E. Mey. Comm. Pl. Afr. Austr.* i. 252.—Afr. austr.
parvifolia, *Choisy, in Mém. Soc. Phys. Genèv.* ii. (1824) 85.—Afr. austr.

AGATHIS, Salisb. in Trans. Linn. Soc. viii. (1807) 311. t. 18. *CONIFERAE*, Benth. & Hook. f. iii. 426.
DAMMARA, [Rumph.] Lam. Encyc. ii. 259 (1786).
australis, *Steud. Nom.* ed. II. i. 84.—N. Zel.
Dammara, Rich. Conif 93. t. 19 = loranthifolia.
loranthifolia, *Salisb. in Trans. Linn. Soc.* viii. (1807) 312. t. 15.—Malaya.

AGATHISANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 294 (1861) = **Oldenlandia**, Plum. (Rubiac.).
Beyrei, Klotzsch, l. c. = O. Bojeri.
globosum, Klotzsch, l. c. = O. globosa.
Petersii, Klotzsch, l. c. 295 = O. Bojeri.

AGATHISANTHES, Blume, Bijdr. 645 (1825) = **Nyssa**, Linn. (Cornac.).
javanica, Blume, l. c. = N. sessiliflora.

AGATHOMERIS, Delaun. in Bon Jard. (1835) ex DC. Prod. vi. 157 = **Humea**, Sm. (Compos.).

AGATHOPHORA, Bunge (Anabas. Revis. 92); in Mém. Acad. Pétersb. Sér. VII. iv. (1862) n. 11 = **Halogeton**, C. A. Mey. (Chenopod.).
alopecuroides, Bunge, l. c. = H. alopecuroides.

AGATHOPHYLLUM, Jusa. Gen. 431 (1789) = **Ravensara**, Sonner. (Laurin.).
acuminatum, Willd. ex Meissn. in DC. Prod. xv. i. 110 = R. acuminata.
aromaticum, Willd. Sp. Pl. ii. 849 = R. aromatica.
cujabrae, Blume, Mus. Bot. Lugd. Bat. i. 340 (= *Mespilodaphne cujabraris*).
cymosum, Meissn. in DC. Prod. xv. i. 111 (= *Mespilodaphne cymosa*).
heteromorphum, Blume, Mus. Bot. Lugd. Bat. i. 340 (= *Mespilodaphne heteromorpha*).
Lindleyanum, Blume, l. c. (= *Mespilodaphne Lindleyana*).
marginatum, Blume, l. c. (= *Mespilodaphne marginata*).
Novianum, Blume, l. c. 339 (= *Mespilodaphne mauritiana*).
obtusatum, Blume, l. c. 340 (= *Mespilodaphne heteromorpha*).
Persoonianum, Blume, l. c. 341 (= *Mespilodaphne Persooniana*).
Rouwsara, Mirb. ex Steud. Nom. ed. II. 84 = R. aromatica.
retusum, Willd. ex Meissn. in DC. Prod. xv. i. 110 = R. retusa.

AGATHOPHYTUM, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 291. t. 10. c. = **Chenopodium**, Linn.
acuminatum, Schur, Enum. Pl. Transs. 879; Index = C. acuminatum.
Bonus-Henricus, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 291 = C. Bonus-Henricus.
crassifolium, Fuss, Fl. Transs. 553 = C. crassifolium.
glaucum, Fuss, l. c. = C. glaucum.
humifusum, Kitt. Taschenb. ed. II. 264 = C. crassifolium.
rubrum, Reichb. Fl. Sax. 321 = C. rubrum.

AGATHOSMA, Willd. Enum. Hort. Berol. 259 (1809). *RUTACEAE*, Benth. & Hook. f. i. 290.
BUCCO, Wendl. Coll. i. 9. etc. (1808).
DICHOSMA, DC. ex Loud. Hort. Brit. 85 (1830).
GYMNONYCHIUM, Bartl. in Linnaea, xvii. (1843) 554. t. 11.
HARTOGIA, Linn. Syst. ed. X. 939 (1759) (nomen prius).
acerosa, Eckl. & Zeyh. Enum. 114.—Afr. austr.
acuminata, Drège, ex Walp. Ann. I. 185 = punctata.
acuminata, Willd. Enum. Hort. Berol. 260 = imbricata.
acutiloba, *Sond. in Harv. & Sond. Fl. Cap.* i. 416.—Afr. austr.

AGATHOSMA :—
adenocaulis, *Eckl. & Zeyh. Enum.* 115.—Afr. austr.
affinis, *Sond. in Harv. & Sond. Fl. Cap.* i. 405.—Afr. austr.
alaris, *Cham. in Linnaea*, v. (1830) 58.—Afr. austr.
ambigua, *Sond. in Harv. & Sond. Fl. Cap.* i. 426.—Afr. austr.
ambigua, *Sweet, Hort. Brit.* ed. I. 89.—Afr. austr.
anomala, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 408.—Afr. austr.
apiculata, E. Mey. ex Sond. l. c. 405 = apiculata.
apiculata, E. Mey. in Bartl. & Wendl. f. Diosm. 176 = Barosma apiculata.
aristata, *Presl, Rel. Remerk.* 32.—Afr. austr.
asperifolia, *Eckl. & Zeyh. Enum.* 111.—Afr. austr.
aulonophila, Eckl. & Zeyh. l. c. 115 = commutata, virgata.
barbata, Spreng. Pugill. i. 20 = Macrostylis lanceolata.
barosmaefolia, *Eckl. & Zeyh. Enum.* 111.—Afr. austr.
barosmoides, *Sond. in Harv. & Sond. Fl. Cap.* i. 416.—Afr. austr.
Bathiana, Eckl. & Zeyh. Enum. 114 = A. Cerefolium, serpyllacea.
bernaliaefolia, Eckl. & Zeyh. l. c. 115 = virgata.
bifida, Bartl. & Wendl. f. Diosm. 152 = umbellata.
biophylla, Eckl. & Zeyh. Enum. 110 = hirta.
bisulca, *Bartl. & Wendl. f. Diosm.* 129.—Afr. austr.
blaerioides, *Cham. in Linnaea*, v. (1830) 56.—Afr. austr.
blaerioides, Eckl. & Zeyh. Enum. 113 = chortophila.
brevifolia, Lond. Hort. Brit. 86 = erecta.
brunioides, Lond. l. c.; G. Don, Gen. Syst. i. 787 = cuspidata.
capitata, *Sond. in Harv. & Sond. Fl. Cap.* i. 434.—Afr. austr.
cephalotes, E. Mey. ex Sond. l. c. 406 = globulifera.
Cerefolium, *Bartl. & Wendl. f. Diosm.* 139.—Afr. austr.
chortophila, *Eckl. & Zeyh. Enum.* 116.—Afr. austr.
ciliata, *Link, Enum. Hort. Berol.* i. 235.—Afr. austr.
clavata, Hoffmgg. Verz. Pfl. 36 = obtusa.
collina, *Eckl & Zeyh. Enum.* 110.—Afr. austr.
commutata, *Sond. in Harv. & Sond. Fl. Cap.* i. 412.—Afr. austr.
concinna, *Hoffmgg. Verz. Pfl.* 36.—Afr. austr.
cordata, Hoffmgg. l. c. = subcordata.
corymbosa, G. Don, Gen. Syst. i. 789 = villosa.
craspedota, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 415.—Afr. austr.
crassifolia, *Sond. l. c.* 438.—Afr. austr.
cuspidata, *Bartl. & Wendl. f. Diosm.* 132.—Afr. austr.
cyminodes, Eckl. & Zeyh. Enum. 117 = chortophila.
decumbens, *Eckl. & Zeyh. l. c.* 116.—Afr. austr.
dioica, Sweet, Hort. Brit. ed. I. 89 = Barosma lanceolata.
Dregeana, *Sond. in Harv. & Sond. Fl. Cap.* i. 410.—Afr. austr.
Eckloniana, *Schlecht. in Linnaea*, v. (1830) 207.—Afr. austr.
elata, *Sond. in Harv. & Sond. Fl. Cap.* i. 438.—Afr. austr.
elegans, *Cham. in Linnaea*, v. (1830) 54.—Afr. austr.
erecta, *Bartl. & Wendl. f. Diosm.* 135.—Afr. austr.
eriantha, Steud. Nom ed. II. i. 85 = Schlechtendalii.
ericoides, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.
fastigiata, *Eckl. & Zeyh. Enum.* 117.—Afr. austr.
filipetala, *Eckl. & Zeyh. l. c.* 112.—Afr. austr.
florida, *Sond. in Harv. & Sond. Fl. Cap.* i. 434.—Afr. austr.
fluvulenta, *Sond. l. c.* 437.—Afr. austr.
foetidissima, Hort. ex Steud. Nom. ed. II. i. 85 = Barosma foetidissima.
follosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 421.—Afr. austr.
fraudulenta, *Sond. l. c.* 415.—Afr. austr.
geminiflora, *Eckl. & Zeyh. Enum.* 115.—Afr. austr.
Gillivrayi, *Sond. in Harv. & Sond. Fl. Cap.* i. 427.—Afr. austr.
glabra, Eckl. & Zeyh. Enum. 116 = serpyllacea.
glabrata, *Bartl. & Wendl. f. Diosm.* 166.—Afr. austr.
glabrata, E. Mey. ex C. Mnall. in Walp. Ann. vii. 518 = crassifolia.
glandulosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 430.—Afr. austr.
glauca, Eckl. & Zeyh. Enum. 112 = hians.
globulifera, *Presl, Rel. Remerh.* 31.—Afr. austr.
goldmiedea, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.

AGATHOSMA :—
gonaquensis, *Eckl. & Zeyh. Enum.* 110.—Afr. austr.
gracilicaulis, *Sond. in Harv. & Sond. Fl. Cap.* i. 409. —Afr. austr.
graveolens, Meisn. in Flora, xxvii. (1844) 1. 302 = collina.
hirta, *Bartl. & Wendl. f. Diosm.* 188.—Afr. austr.
hirta, [Ker-Gawl.] in Bot. Reg. t. 369 = villosa.
hirtella, *Sond. in Harv. & Sond. Fl. Cap.* i. 420.—Afr. austr.
hispida, *Bartl. & Wendl. f. Diosm.* 182.—Afr. austr. Hookeri, *Sond. in Harv. & Sond. Fl. Cap.* i. 406.—Afr. austr.
humilis, *Sond. l. c.*—Afr. austr.
hybrida, Bartl. & Wendl. f. Diosm. 167 = rugosa.
hyponerva, Eckl. & Zeyh. Enum. 114 = serpyllacea.
imbricata, Drège, ex Walp. Ann. i. 155 = polyphylla.
imbricata, *Willd. Enum. Hort. Berol.* 259.—Afr. austr.
involucrata, *Eckl. & Zeyh. Enum.* 109.—Afr. austr.
Joubertiana, *Schlecht. in Linnaea*, vi. (1831) 207.— Afr. austr.
juncea, Hoffmgg. Verz. Pfl. 36 = imbricata.
juniperifolia, *Bartl. in Linnaea*, xvii. (1843) 376.— Afr. austr.
juniperina, *Sond. in Harv. & Sond. Fl. Cap.* i. 433.— Afr. austr.
lanceolata, Spreng. ex Bartl. & Wendl. f. Diosm. 156 = ciliata.
lancifolia, *Eckl. & Zeyh. Enum.* 109.—Afr. austr.
lanisphylla, G. Don, Gen. Syst. i. 788 = villosa.
latifolia, Loud. Hort. Brit. 85 = Barosma serratifolia.
latipetala, *Sond. in Harv. & Sond. Fl. Cap.* i. 422.— Afr. austr.
laxa, Bartl. & Wendl. f. Diosm. 162 = villosa.
lediformis, *Eckl. & Zeyh. Enum.* 109.—Afr. austr.
leptopermoides, Sond. in Harv. & Sond. Fl. Cap. i. 405 = arnstata.
linifolia, *Lichtst. ex Roem. & Schult. Syst.* v. 448 — Afr. austr.
lycopodioides, *Bartl. & Wendl. f. Diosm.* 148.—Afr. austr.
marifolia, *Eckl. & Zeyh. Enum.* 113.—Afr. austr.
Martiana, *Sond. in Harv. & Sond. Fl. Cap.* i. 416.— Afr. austr.
melalecoides, *Sond. l. c.* 431.—Afr. austr.
Meyeriana, *Presl, Bot. Bemerk* 32.—Afr. austr.
microphylla, *E. Mey. in Bartl. & Wendl. f. Diosm.* 173. —Afr. austr.
minus, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.
mollis, Bartl. & Wendl. f. Diosm. 168 = rugosa.
mollis, Eckl. & Zeyh. ex C. Muell. in Walp. Ann. vii. 514 = floridula.
montana, *Schlecht. in Linnaea*, vi. (1831) 207.—Afr. austr.
monticola, *Sond. in Harv. & Sond. Fl. Cap.* i. 409.— Afr. austr.
mucronulata, *Sond. l. c.* 417.—Afr. austr.
Mundtii, *Cham. in Linnaea*, v. (1830) 56.—Afr. austr.
myrsinites, G. Don, Gen. Syst. i. 789 = ciliata.
nigra, *Eckl. & Zeyh. Enum.* 117.—Afr. austr.
nigromontana, *Eckl. & Zeyh. l. c.* 111.—Afr. austr.
Niveni, *Sond. in Harv. & Sond. Fl. Cap.* i. 418.—Afr. austr.
obtusa, *Bartl. & Wendl. f. Diosm.* 169 = rugosa.
obtusata, G. Don, Gen. Syst. i. 789 = imbricata.
obtusifolia, *Presl, Bot. Bemerk.* 31.—Afr. austr.
orbiculata, *Bartl. & Wendl. f. Diosm.* 175.—Afr. austr.
Owanii, *Harv. & Sond. ex Harv. Thes. Cap.* ii. 42.— Afr. austr.
parviflora, *Bartl. & Wendl. f. Diosm.* 181.—Afr. austr.
patens, Hort. ex Barth. & Wendl. f. Diosm. 159 = A. Cerefolium.
patentissima, Eckl. & Zeyh. Enum. 114 = cuspidata.
patula, Eckl. & Zeyh. l. c. 116 = variabilis.
patula, *C. F. W. Mey. ex Bartl. & Wendl. f. Diosm.* 134.—Afr. austr.
pentachotoma, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 405.—Afr. austr.
perforata, *G. Don, Gen. Syst.* i. 789.—Afr. austr.
pilifera, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.
pinnifolia, *Sond. in Harv. & Sond. Fl. Cap.* i. 413.— Afr. austr.
platypetala, *Eckl. & Zeyh. Enum.* 116.—Afr. austr.

AGATHOSMA :—
polyphylla, *Presl, Bot. Bemerk.* 32.—Afr. austr.
prolifera, *Bartl. & Wendl. f. Diosm.* 185.—Afr. austr.
prolifera, Eckl. & Zeyh. Enum. 114 = bisulca.
propinqua, *Sond. in Harv. & Sond. Fl. Cap.* i. 423.— Afr. austr.
pubescens, G. Don, Gen. Syst. i. 788 = villosa.
pubescens, *Sond. in Harv. & Sond. Fl. Cap.* i. 412.— Afr. austr.
pubescens, *Willd. Enum. Hort. Berol.* 259.—Afr. austr.
pubigera, *Sond. in Harv. & Sond. Fl. Cap.* i. 414.— Afr. austr.
pulchella, Link, Enum. Hort. Berol. i. 238 = Barosma pulchella.
punctata, G. Don, Gen. Syst. i. 789 = ovata.
punctata, *Sond. in Harv. & Sond. Fl. Cap.* i. 417. —Afr. austr.
recurvifolia, *Sond. l. c.* 415.—Afr. austr.
reflexa, *Link, Enum. Hort. Berol.* i. 238.—Afr. austr.
robusta, *Eckl. & Zeyh. Enum.* 110.—Afr. austr.
rubra, *Willd. & Lichtst. ex Bartl. & Wendl. f. Diosm.* 178.—Afr. austr.
rufescens, G. Don, Gen. Syst. i. 789 = rubra.
rugosa, Link, Enum. Hort. Berol. i. 238 = pubescens.
sabulosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 406.— Afr. austr.
salina, Eckl. & Zeyh. Enum. 111 = asperifolia.
Schlechtendalii, *Sond. in Harv. & Sond. Fl. Cap.* i. 407.—Afr. austr.
Schlechtendahliana, Eckl. & Zeyh. Enum. 109 = Schlechtendalii.
sedifolia, *Schlecht. in Linnaea*, vi. (1831) 206 —Afr. austr.
serpyllacea, *Lichtst. ex Roem. & Schult. Syst.* v. 447. —Afr. austr.
serruriaefolia, Eckl. & Zeyh. Enum. 109 = bifida.
speciosa, Hoffmgg. Verz. Pfl. Nachtr. i. 128 = laxa.
spicata, *Lichtst. ex G. Don, Gen. Syst.* i. 788.—Afr. austr.
spinosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 417.— Afr. austr.
squamosa, *Bartl. & Wendl. f. Diosm.* 141.—Afr. austr.
stadensis, Eckl. & Zeyh. Enum. 113 = microphylla.
stenopetala, Steud. Nom. ed. II. i. 36 = filipetala.
suaveolens, Eckl. & Zeyh. Enum. 113 = A. Cerefolium.
subcordata, Hoffmgg. Verz. Pfl. 36, 163 = imbricata ?
tabularis, *Sond. in Harv. & Sond. Fl. Cap.* i. 403.— Afr. austr.
tenuis, *Sond. l. c.* 423.—Afr. austr.
tenuissima, Otto, ex Bartl. & Wendl. f. Diosm. 187 = ericoides.
teretifolia, G Don, Gen. Syst. i. 787 = Diosma teretifolia.
Thunbergiana, *Bartl. & Wendl. f. Diosm.* 150.—Afr. austr.
Thunbergiana, Schlecht. in Linnaea, vi. (1831) 204 = ambigua.
Thunbergiana, Sond. in Harv. & Sond. Fl. Cap. i. 425 = platypetala.
thymifolia, Hoffmgg. Verz. Pfl. 36 = acuminata.
thymifolia, *Schlecht. in Linnaea*, vi. (1831) 205.—Afr. austr.
thujoides, Sond. in Harv. & Sond. Fl. Cap. i. 437 = glabrata.
thyoides, G. Don, Gen. Syst. i. 787; Eckl. & Zeyh. Enum. 116 = glabrata.
trachyphylla, Eckl. & Zeyh. l. c. 110 = bispidata.
umbellata, Sond. in Harv. & Sond. Fl. Cap. i. 408 = bifida.
variabilis, *Sond. l. c.* 433.—Afr. austr.
Ventenatiana, *Bartl. & Wendl. f. Diosm.* 161.—Afr. austr.
vestita, Willd. ex Roem. & Schult. Syst. v. 447 = imbricata.
villosa, *Willd. Enum. Hort. Berol.* 259.—Afr. austr.
virgata, *Bartl. & Wendl. f. Diosm.* 139.—Afr. austr.
virgata, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 413 = commutata.
virgata, Spreng. ex Bartl. & Wendl. f. Diosm. 145 = imbricata.
Wendlandiana, G. Don, Gen. Syst. i. 788 = erecta.

AGATHOTES, D. Don, in Phil. Mag. viii. (1836) 77 = **Swertia**, Linn. (Gentian.).
alata, D. Don, ex G. Don, Gen. Syst. iv. 177 = S. alata.
Chirayta, D. Don, l. c. 163 = S. Chirata.
nervosa, G. Don, l. c. 177 = S. nervosa.

AGATHYRSUS, D. Don, in Edinb. N. Phil. Journ. (1838–39) 310 = **Lactuca**, Linn. et **Sonchus**, Tourn. (Compos.).
alpinus, D. Don, l. c. = L. alpina.
cacalaefolius, D. Don, l. c. (= *Mulgedium cacalaefolium*).
cauvulens, D. Don, l. c. = L. alpina.
cyaneus, D. Don, l. c. = L. hastata.
floridanus, D. Don, l. c. = L. leucophaea.
leucophaeus, Beck, Bot. 170 = L. leucophaea.
macrophyllus, Beck, l. c. = L. macrophylla.
Plumieri, D. Don, in Edinb. N. Phil. Journ. (1838–39) 310 = L. Plumieri.
prenanthoides, D. Don, l. c. = L. prenanthoides.
pulchellus, D. Don, l. c. = L. pulchella.
sibiricus, D. Don, l. c. = L. sibirica.
spicatus, D. Don, l. c.—Hab. ?
tataricus, D. Don, l. c. = L. tatarica.

AGATHYRUS, Rafin. New Fl. Am. iv. 84 (1836) = ENOXIS, Rafin. = **Lactuca**, Linn. ? (Compos.).

AGATI, Adans. Fam. ii. 326 (1763) = **Sesbania**, Pers. (Legum.).
cannabina, Desv. Journ. Bot. i. (1813) 190 = S. aculeata.
coccinea, Desv. l. c. t. 4 = S. grandiflora.
formosa, F. Muell. Fragm. ii. 88 = S. grandiflora.
grandiflora, Desv. Journ. Bot. i. (1813) 190. t. 4 = S. grandiflora.
tomentosa, Nutt. ex A. Gray, Bot. U. St. Expl. Exped. i. 469 = S. tomentosa.
virgata, Desv. Journ. Bot. i. (1813) 190 = Courretia virgata.

AGATION, Brongn. in Bull. Soc. Bot. Fr. viii. (1861) 79. *VIOLARIEAE*, Benth. & Hook. f. i. 118.
AGATEA, A. Gray, in Proc. Am. Acad. ii. (1852) 222. (nomen prius).
Pancheri, *Brongn. l. c.* 80.—N. Caled.
Vieillardii, *Brongn. l. c.*—N. Caled.
violare, *Brongn. l. c.*—Ins. Fiji.

AGATOPHYTON, Fourr in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 144 = AGATHOPHYTUM, Moq. = **Chenopodium**, Linn.

AGAURIA, Benth. & Hook. f. Gen. ii. 586 (1876). *ERICACEAE*, Benth. & Hook. f. ii. 586.
AGARISTA, D. Don, in G. Don, Gen. Syst. iii. 837 (1834) (quoad spp. Africanae).
buxifolia, *Baker, Fl. Maurit.* 187, *in obs.*—Ins. Borb.
polyphylla, *Baker, in Journ. Linn. Soc.* xx. (1884) 194.—Madag.
salicifolia, *Hook. f. ex Oliver, Fl. Trop. Afr.* iii. 488.— Afr. trop.

AGAVE, Linn. Hort. Ups. 87 (1748). *AMARYLLIDEAE*. Benth. & Hook. f. iii. 736.
ALLIBERTIA, Marion, in Rev. Hortic. Bouch. Rhone (Nov. 1882), ex Baker, in Gard. Chron. (1883) i. 176.
BONAPARTEA, Haw. Syn. Succ. 68 (1812).
LITTAEA, Tagliabue, in Bibl. Ital. i. 106 (1816).
MANFREDIA, Salisb. Gen. Pl. Fragm. 78 (1866).
albicans, *Jacobi, in Hamb. Gartens.* xxi. (1865) 256. —Mexic.
Alibertii, *Baker, in Gard. Chron.* N. S. xix. (1883) i. 176.—Mexic.?
aloides, *Jacobi, in Hamb. Gartens.* xxii. (1866) 265.— Hab. ?
aloiva, C. Koch, Wochenschr. iii. (1860) 37 = Sartorii.
altissima, Zuccag. Fl. Pedem. ii. 50 = americana.
americana, *Linn. Sp. Pl.* 323.—Am. trop.
amoena, Lem. ex Jacobi, in Hamb. Gartenz. xxi. (1865) 120 = A. Scolymus.
amurensis, Jacobi, l. c. xx. (1864) 548 = xylonacantha.
ananassoides, De Jonghe & Jacobi, Nachtr. i. 32 = Regeliana.
angustifolia, Haw. Syn. Succ. 72 = rigida.
angustifolia, Hort. Par. ex Steud. Nom. ed. II. i. 84 = Furcraea tuberosa.
angustissima, *Kegelin. in Trans. Acad. Sc. St. Louis*, iii. (1875) 306.—Mexic.
antillarum, Descourt. Antill. iv. 289. t. 284 = subsimplex.
applanata, *Lem. ex Jacobi, in Hamb. Gartens.* xx. (1864) 550.—Mexic.

AGAVE :—

aspera, Jacq. Enum. Vindob. App. 307 = Furcraea cubensis.

asperrima, *Jacobi*, *in Hamb. Gartenz.* xx. (1864) 561. —Texas.

atrovirens, *Karw. in Salm-Dyck*, *Hort.* (1834) 302.—Mexic.

attenuata, *Salm-Dyck*, *l. c.* 308.—Mexic.

auriculata, Hort. ex Baker, in Gard. Chron. (1877) II. 40 = A. Scolymus.

australis, Stunf. Nom. ed. I. 18 = Furcraea cubensis.

Baulan, *Perr. in Mem. Soc. Linn. Par.* iii. (1824) 97.—Java.

Beaucarnei, Lem. Illustr. Hortic. xi. (1864) Misc. 65 = Kerchovel.

Beauleriana, Jacobi, in Abh. Schles. Gesell. (1859) 150 = mexicana.

Bernhardi, *Jacobi*, *Nachtr.* i. 38.—Mexic.

Besseriana, De Smet, ex Jacobi, in Hamb. Gartenz. xxi. (1865) 155 = macroacantha.

Bonnetiana, Peacock, ex Baker, in Gard. Chron. (1877) II. 40 = ferox.

Botteri, *Baker*, *in Bot. Mag.* t. 6248.—Mexic.

Bouchei, *Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 217.—Mexic.

brachystachys, *Cav. Descr.* 453.—Mexic.

bracteosa, *S. Watz. ex Engelm. in Gard. Chron.* (1882) I. 776. I. 189.—Mexic ; Calif.

Brauniana, *Jacobi*, *in Hamb. Gartenz.* xxii. (1866) 216.—Mexic.

bromeliaefolia, *Salm-Dyck*, *Hort* (1834) 303.—Mexic

bulbifera, Bonpl. ex Steud. Nom. ed. II. i. 84 = mexicana.

bulbifera, *Salm-Dyck*, *Hort.* (1834) 8, 303.—Mexic.

caerulescens, Salm-Dyck, ex Jacobi, Monog. 38 = lophantha.

caespitosa, Tod. Hort. Bot. Panorm. (1877) 32. t. 8 = Sartorii.

californica, Jacobi, Nachtr. 47 = falcata.

Canartiana, Jacobi, in Abh. Schles. Gesell. (1869) 147 = atrovirens.

Candelabrum, Tod. Hort. Bot. Panorm. (1877) 66. t. 15 = rigida.

Cantula, Roxb. Fl. Ind. ii. 167 = vivipara.

carlhaea, *Hort. Kew. ex Baker*, *in Gard. Chron.* (1879) II. 683.—Ind. occ.

Celsiana, Jacobi, in Hamb Gartenz. xxi. (1865) 446 = Celsii.

Celsii, *Hook. Bot. Mag.* t. 4934.—Mexic.

chiapensis, Hort. Belg. ex Jacobi, in Hamb. Gartenz. xxii. (1866) 213 = polyacantha.

chloracantha, *Salm-Dyck*, *in Otto*, *Allg. Gartenz.* (1834) 50.—Mexic.

cinerascens, Jacobi, in Hamb. Gartenz. xx. (1864) 553 = applanata.

coarctata, Jacobi, in Abh. Schles. Gesell. (1869) 147 = atrovirens.

coccinea, *Roesl*, *ex Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 61.—Mexic.

cochlearis, *Jacobi*, *Nachtr.* ii. 67.—Mexic.

Cohniana, Jacobi, in Hamb. Gartenz. xxii. (1866) 221 = yuccaefolia.

Commelini, Salm-Dyck, Hort. (1834) 301 = Furcraea Commelyni.

concinna, *Baker*, *in Gard. Chron.* (1877) II. 137.—Mexic.

concinna, Hort. Angl. ex Baker, l. c. 717 = albicans.

conduplicata, *Jacobi & Bond. in Hamb. Gartenz.* xxi. (1865) 592.—Mexic.

Consideranti, Carr. in Rev. Hortic. (Nov. 1875) 429. f. 66 = A. Victoriæ-reginæ.

Corderoyi, Versch. ex Baker, in Gard. Chron. (1877) II. 396.—Mexic.

crenata, Jacobi, in Hamb. Gartenz. xxii. (1866) 176 = A. Scolymus.

cubensis, Jacq. Amer. 100 = Furcraea cubensis.

cucullata, *Lem. ex Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 124.—Mexico.

cuspidata, Hort. Saunders, ex Baker, in Gard. Chron. (1877) II. 137 = flaccida.

cyanophylla, Jacobi, in Hamb. Gartenz. xxii. (1866) 175 = mexicana.

dasylirioides, *Yacobi & Bouché*, *l. c.* xxi. (1865) 344.—Guatemal.

decaltata, Lem. ex Jacobi, l. c. 346 = dasylirioides.

Debaryana, Jacobi, in Abh. Schles. Gesell. (1869) 164 = prhinosa

Desmimeana, *Jacobi*, *l. c.* 153.—Mexic.

deflexispina, Jacobi, Nachtr. ii 72 = atrovirens.

AGAVE :—

Demesteriana, Hort. ex Jacobi, Monog. 218 = mirabiorensis.

De-Mesteriana, *Yacobi*, *in Hamb. Gartenz.* xxii. (1866) 167.—Mexic.

densiflora, *Hook. Bot. Mag.* t. 5006.—Mexic.

demisflora, Kegel, Gartenfl. (1858) 274. t. 410 = rupicola.

dentata, Hort. ex Baker, in Gard. Chron. (1877) II. 748 = prunosa.

Deserti, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 310.—Calif.

desmetiana, Hort. ex Baker, in Gard. Chron. (1877) I. 621 = horrida.

Desmettiana, Jacobi, in Hamb. Gartenz. xxii. (1866) 217 = miradorensis.

drimiæfolia, Hort. Petrop. ex Baker, in Gard. Chron. (1877) II. 808 = undulata.

Ehrenbergiana, Baker, l. c. 620 = Ehrenbergii.

Ehrenbergii, *Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 255.—Mexic.

echinoides, Jacobi, Nachtr. i. 48 = striata.

Ellemeetiana, *C. Koch*, *Wochenschr.* viii. (1865) 108.—Mexic.

elongata, Jacobi, in Hamb. Gartenz. xxi. (1865) 168 = rigida.

ensifera, Jacobi, Nachtr. 14 = univittata.

ensiformis, Hort. ex Baker, in Gard. Chron. (1877) II. 556 = striata.

excelsa, *Yacobi*, *in Hamb. Gartenz.* xxii. (1866) 215.—Honduras.

expansa, Jacobi, Nachtr. ii. 36 = americana.

falcata, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 304.—Mexic.

Feuziana, *Jacobi*, *in Hamb. Gartenz.* xxii. (1866) 170. Mexic.

ferox, *C. Koch*, *Wochenschr.* iii. (1860) 28.—Mexic.

filamentosa, Salm-Dyck, in Bonplandia, vii. (1859) 94 = filifera.

filifera, *Salm-Dyck*, *Hort.* (1834) 309.—Mexic.

flaccida, Haw. Syn. Succ. 73 = rubescens.

flaccida, *Yacobi*, *in Hamb. Gartenz.* xxii. (1866) 174.—Mexic.

flavescens, Salm-Dyck, Hort. Dyck. 8 = macroacantha.

flavovirens, *Yacobi*, *in Hamb. Gartenz.* xxii. (1866) 270. Mexic. ?

fortida, Linn. Sp. Pl. 323 = Furcraea gigantea.

fourcroydes, Lem. Illustr. Hortic. xi. (1864) Misc. 65 = rigida.

fragrantissima, *Yacq. Enum. Vindob. App.* 309.—Hab.?

Fuerstenbergii, Jacobi, Nachtr. ii. 72 = americana.

funifera, Lem. Illustr. Hortic. xi. (1864), Misc 65, sub t. 379 = Yucca funifera.

Funkiana, C. Koch & Bouché, in C. Koch, Wochenschr. iii. (1860) 47 = lophantha.

Galeottei, *Baker*, *in Gard. Chron.* (1877) II. 40.—Mexic.

geminiflora, *Ker-Gawl. in Brandt*, *Journ. Sc.* iii. (1817) no. 3. t. 1.—Peruv.

Ghiesbreghtii, *C. Koch*, *Wochenschr.* v. (1862) 83, 148.—Mexic.

gigantea, D. Dietr. Syn. Pl. ii. 1193 = Furcraea gigantea.

Gilbeyi, Hort. ex Gard. Chron. (1878) 1305 = horrida.

glabra, *Karw. in M. Korn. Syn. Ensat.* 293.—Mexic.

glaucescens, Hook. Bot. Mag. t. 5333 = attenuata.

glaucescens, *Otto*, *in M. Korn. Syn. Ensat.* 293.—Mexic.

Goeppertiana, *Yacobi*, *in Hamb. Gartenz.* xxii. (1866) 219.—Mexic.

gracilis, Jacobi, Nachtr. ii. 66 = atrovirens.

grandidentata, Hort. Belg. ex Jacobi, in Hamb. Gartenz. xxii. (1866) 114 = horrida.

granulosa, *Schnidw. in C. Koch*, *Wochenschr.* iv. (1861) 206.—Mexic.

Guedeneyri, *Houllet*, *in Rev. Hortic.* (1875) 466.—Mexic.

Gustaviana, Hort. Corderoy, ex Baker, in Gard. Chron. (1877) I. 8. viii. (1877) II. 201 = Maximiliana.

guttata, *Yacobi & Bouché*, *in Hamb. Gartenz.*, xxi. (1865) 561.—Mexic.

Haselloffii, *Yacobi*, *in Hamb. Gartenz.* xxii. (1866) 220.—Mexic.

Hawortkiana, M. Roem. Syn. Ensat. 290 = mexicana ?

Haynaldi, *Tod. Hort. Bot. Panorm.* I. 68. t. 24.—Mexic. ?

heteracantha, Hort. ex Jacobi, in Hamb. Gartenz. xxii. (1866) 62 = lophantha.

heteracantha, *Zucc. in Nov. Act. Nat. Cur.* xvi. (1833) II. 676.—Mexic.

AGAVE :—

hexapetala, *Yacq. Enum. Pl. Carib.* 12.—Ind. occ.

Hookeri, C. Koch, Wochenschr. viii. (1865) 104 = yuccaefolia.

Hookeri, *Yacobi*, *in Hamb. Gartenz.* xxii. (1866) 166.—Mexic.

Hookeri, Baker, in Bot. Mag. t. 6589 = Fenzliana.

horizontalis, *Yacobi*, *in Abh. Schles. Gesell.* (1869) 148, bis.—Mexic.

horrida, *Lem. ex Yacobi*, *in Hamb. Gartenz.* xx. (1864) 546.—Mexic.

Houlletii, Hort. Par. ex Jacobi, l. c. xxii. (1866) 368 = rigida.

Humboldtiana, *Yacobi*, *l. c.* 264.—Mexic.

humilis, M. Roem. Syn. Ensat. 285 = brachystachys.

hybrida, Versch. Cat. n. 12 (1865–69) = xylonacantha.

Hystrix, Hort. ex Baker, in Gard. Chron. (1877) II. 556 = striata.

inaequidens, *C. Koch*, *Wochenschr.* iii. (1860) 28.—Mexic.

Inghami, Versch. Cat. n. 16 (1871–72), 11 = Roezliana.

Ixtli, Karw. in Salm-Dyck, Hort. (1834) 304 = rigida.

ixtlioides, Hook. f. Bot. Mag. t. 5893 = rigida.

Yacobiana, Salm-Dyck. in Bonplandia, viii. (1860), 88 = atrovirens.

Yacquiniana, Schult. f. Syst. vii. 737 = lurida.

Karatto, *Mill. Gard. Dict.* ed. VIII. n. 6.—Mexic.

Karwinskii, Zucc. in Nov. Act. Nat. Cur. xvi. II. (1833) 675 = rigida.

Kellockii, Jacobi, in Abh. Schles. Gesell. (1869) 165 = prunosa.

Keratto, Steud. Nom. ed. II. i. 36 = Karatto.

Keratto, Salm-Dyck, Hort. Dyck. 304 = mexicana.

Keratto, Salm-Dyck, in Bonplandia, vii. (1859) 93 = Salmdyckii.

Kerchovei, *Lem. Illustr. Hortic.* xi. (1864) 64.—Mexic.

kewensis, *Yacobi*, *in Hamb. Gartenz.* xxii. (1866) 218. —Mexic.

Kochii, Jacobi, l. c. 117 = xylonacantha.

lamprochlora, Jacobi, in Abh. Schles. Gesell. (1869) 150 = Smithiana.

laticincta, *Versch. Cat.* n. 12 (1868) 2.—Mexic.

latifolia, *Karw. in M. Korn. Syn. Ensat.* 293.—Mexic.

latissima, Jacobi, in Hamb. Gartenz. xx. (1864) 551 = atrovirens.

Laurentiana, *Yacobi*, *l. c.* xxii. (1866) 266.—Mexic.

laxa, *Hort. Monac. ex Salm-Dyck*, *Hort.* (1834) 8.—Am. austr.

Lechegvillo, Torr. Bot. Mex. Bound. 213 = heteracantha.

Legrelliana, Jacobi, in Hamb. Gartenz. xxii. (1866) 266 = Ehrenbergii.?

Leguayana, Versch. Cat. n. 12 (1868) 2 = Ghiesbrechtii.

Lehmanni, Jacobi, in Abh. Schles. Gesell. (1869) 146 = atrovirens.

Lemaireï, *Hort. Y. Versch. ex Illustr. Hortic.* xi. (1864) *Misc.* 65.—Hab.?

lepida, D. Dietr. Syn. ii. 1192 = lurida.

Lindleyi, *Yacobi*, *in Abh. Schles. Gesell.* (1869) 152, bis.—Mexic.

linearis, Jacobi, in C. Koch, & Fint. Wochenschr. xii. (1869) 179 = macroacantha.

lophantha, *Schiede*, *in Linnaea*, iv. (1829) 552.—Mexic.

lurida, *Ait. Hort. Kew.* i. 473.—Mexic.

macracantha, *Tod. Hort. Bot. Panorm.* ii. 11. t. xxvii. —Mexic.?

macrocacantha, *Zucc. in Nov. Act. Acad. Nat. Cur.* xvi. II. (1833) 676.—Mexic.

maculata, *Regel*, *Ind. Sem. Hort. Petrop.* (1856) 16.—Texas.

maculosa, Hook. Bot. Mag. t. 5123 = maculata.

madagascariensis, Spreng. Syst. ii. 79 = Furcraea madagascariensis.

Magni, Desf. Tabl. ed. II. 83 = mexicana.

Maigretiana, Jacobi, in Hamb. Gartenz. xxii. (1866) 121 = horrida.

Maliterii, C. Koch, Wochenschr. v. (1862) 198 = radia.

Mangani, Desf. Tabl. Hort. Par. ed. II. 83 = mexicana.

marmorata, *Roezl*, *in Belg. Hortic.* (1883) 295.—Mexic.

Martiana, *C. Koch*, *Wochenschr.* iii. (1860) 26.—Mexic.

Maximiliana, *Hort. Saunders*, *ex Baker*, *in Gard. Chron.* (1877) II. 201.—Mexic.

AGAVE :—
megalacantha, *Hemsl. Diag. Pl. Nov. Mex.* 55.—Mexic.
melanacantha, *Lem. ex Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 215.—Mexic.
Mezeal, Ellem. ex C. Koch, Wochenschr. viii. (1865) 94 = A. Scolymus.
mexicana, *Lem. Encyc.* i. 59.—Mexic.
micracantha, *Salm-Dyck*, *in Bonplandia*, vii. (1859) 93.—Mexic.
Millori, Haw. Syn. Succ. 71 = americana.
miradorensis, *Jacobi*, *in Abh. Schles. Gesell.* (1869) 156.— Mexic.
mitis, *Mart. Del. Sem. Hort. Monac.* (1848) 4.—Mexic.
mitraeformis, Jacobi, in Abh. Schles. Gesell. (1859) 146 = atrovirens.
Mullmanni, *Jacobi*, *l. c.* (1870–71) 76.—Mexic.
Newberyi, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 309.—Arizona.
Nissoni, *Baker*, *in Gard. Chron.* (1877) i. 528.— Mexic.
Noachi, Jacobi, in Hamb. Gartenz. xxi. (1865) 221 = Sartorii.
oblongata, *Jacobi*, *in Abh. Schles. Gesell.* (1869) 155.— Mexic.
obscura, *Schiede*, *in Linnaea*, xviii. (1844) 413.— Mexic. ?
odorata, Pers. Syn. i. 380 = Furcraea cubensis.
Offoyana, *De Smet*, *ex Jacobi*, *in Hamb. Gartena.* xxi. (1865) 214.—Havana.
oligophylla, *Baker*, *in Gard. Chron* (1876) ii. 492.— Mexic. ?
ornata, Jacobi, in Hamb. Gartenz. xxi. (1865) 59 = americana.
Ottonis, Jacobi, l. c. xxii. (1866) 220 = atrovirens.
Ousselghemiana, Jacobi, in Abh. Schles. Gesell. (1869) 156 = albicans.
pallida, Salisb. Prod. 247 = virginica.
pallida, *Sartorius*, *ex Jacobi*, *in Hamb. Gartenz* xxi. (1865) 171.— Mexic.
Palmeri, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 313.—Arizona.
Parryi, *Engelm. l. c.* 311.—Arizona, Mexic.
parviflora, *Torr. Bot. Mex. Bound.* 214.—Mexic.
parvifolia, *Tod. Hort. Bot. Panorm.* i. (1877) 77.— Mexic.
Pascockii, *Croucher*, *in Gard. Chron.* (1873) 1400. t. 288.—Mexic.
pendula, Schnitz. in Zeits. Gartenb. Darmst. (1857) Jun. n. 21 = Sartorii.
perbella, Hort. ex Baker, in Gard Chron. (1877) 527, 528 = xylonacantha.
perlucida, Jacobi, in Abh. Schles. Gesell. (1869) 151 = Smithiana.
picta, Salm-Dyck, in Bonplandia, vii. (1859) 88 = americana.
polianthoides, *M. Roem. Syn. Ensat.* 286.—Mexic.
polyacantha, *Haw. Rev. Pl. Succ.* 35.—Mexic.
polyanthoides, Cham. & Schlecht. in Linnaea, vi. (1831) 55 = brachystachys.
polyanthoides, *Schiede*, *ex Schlecht. in Linnaea*, xviii. (1844) 413.—Mexic.
polyphylla, C. Koch, Wochenschr. iii. (1860) 38 = mexicana.
Poselgerii, Salm-Dyck, in Bonplandia, vii. (1859) 92 = heteracantha.
potatorum, *Zucc. in Nov. Act. Nat. Cur.* xvi. (1833) ii. 676.—Mexic.
prunosa, *Lem. ex Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 449.—Mexic.
pubescens, *Regel & Orig. in Gartenfl.* (1874) 227. t. 804.—Mexic.
pugioliformis, *Zucc. in Nov. Act. Nat. Cur.* xvi. (1833) ii. 676.—Mexic.
pulcherrima, Hort. ex C. Koch, Wochenschr. viii. (1865) 104 = Sartorii.
pulcherrima, *Otto*, *in M. Roem. Syn. Ensat.* 292.— Mexic.
pulchra, *Hort. Monac. in Salm-Dyck, Cat.* 8.—Am. auct.
pulverulenta, *Verach. Cat.* (1863).—Sierra Nevada.
punctata, *Hort. Berol. ex Kunth, Enum. Pl.* iii. 387.— Mexic.
quadrata, *Lem. Illustr. Hortic.* xi. (1864) *Misc.* 65.— Mexic.
ramosa, Moench, Meth. 642 = americana.
rarispina, Zucc. in Abh. Akad. Muench. iv. iii. (1846) 22 = striata.

AGAVE :—
Regeliana, Hort. ex Baker, in Gard. Chron. N. S. vii. (1877) i. 621 = horrida.
Regelii, *Jacobi*, *in Hamb. Gartenz.* xxii. (1866) 214.—Mexic.
regia, *Hort. Saunders*, *ex Baker*, *in Gard. Chron.* N. S. vii. (1877), ii. 620.—Mexic.
revoluta, *Klotzsch*, *in Otto & Dietr. Gartenz.* viii. (1840) 374.—Mexic.
rigida, *Mill. Gard. Dict.* ed. VIII. n. 8.—Mexic.
rigidissima, Jacobi, in C. Koch & Fint. Wochenschr. xii. (1869) 179 = Kerchovei.
Roezliana, *Baker*, *in Gard. Chron.* N. S. vii. (1877) i. 338. f. 79.—Mexic.
Rokanii, Hort. Belg. ex Jacobi, in Hamb. Gartenz. xx. (1864) 346 = Ghiesbrechtii.
rubescens, *Salm-Dyck*, *Hort.* (1834) 306.—Mexic.
rubrocincta, Jacobi, in Abh. Schles. Gesell. (1869) 153 = Sartorii.
rudis, *Lem. ex Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 216.—Mexic.
rufocincta, Jacobi, Nachtr. i. 88 = Sartorii.
Rumphii, Hassk. in Hoev. & De Vriese, Tydschr. x. (1843) 121 = vivipara.
rupicola, *Regel*, *Gartenfl.* vii. (1858) 312.—Mexic.
Salmdyckii, *Baker*, *in Gard. Chron.* N. S. viii. (1877) ii. 490.—Mexic.
Salmiana, Otto, in Otto & Dietr. Allg. Gartenz. (1842) 51 = atrovirens.
Saponaria, Lindl. Bot. Reg. (1838) Misc. 76 = brachystachys.
Sartorii, *C. Koch*, *Wochenschr.* iii. (1860) 37.—Mexic.
Saundersii, Hook. f. Bot. Mag. t. 5498 = A. Scolymus.
scabra, Salm-Dyck, in Bonplandia, vii. (1859) 89 = Wallisii.
schidigera, *Lem. in Illustr. Hortic.* viii. t. 280, verso (1860) ix. t. 330.—Mexic.
Schiechtendahlii, Jacobi, in Hamb. Gartenz. xx. (1864) 556 = atrovirens.
Schnittspahnii, Jacobi, l. c. xxi. (1865) 120 = A. Scolymus.
Schottii, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 305.—Mexic.
Scolymus, *Karw. in Salm-Dyck, Hort.* (1834) 307.— Mexic.
Seemanniana, *Jacobi*, *in Abh. Schles. Gesell.* (1869) 154.—Guatemal.
serrulata, *Karw. in Otto & Dietr. Allg. Gartenz.* (1842) 51.—Mexic.
sessiliflora, *Hemsl. Diag. Pl. Nov.* 55.—Mexic.
Shawii, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 314. t. 2–4.—Calif.
sisalana, Perrine, ex Engelm. l. c. (reimpr. 28) = rigida.
Smithiana, *Jacobi*, *in Hamb. Gartenz.* xxii. (1866) 203.—Mexic.
sobolifera, *Salm-Dyck*, *Hort.* (1834) 307, 309.—Ind. occ. ; Mexico.
spectabilis, Salisb. Prod. 247 = americana.
spectabilis, *Tod. Hort. Bot. Pan.* ii 4. t. 25.— Mexic. ?
spicata, *Cav. in Anal. Cienc. Nat.* v. (1802) 261 ; *Desc.* 164.—Cuba.
spicata, DC. Cat. Hort. Monsp. (1813) 74 = brachystachys.
spicata, Lour. Ind. Sem. (1826) ex Kunth, Enum. Pl. v. 681 = yuccaefolia.
spinosa, Steud. Nom. ed. II. i. 37 = Furcraea tuberosa.
splendens, *Jacobi*, *in Abh. Schles. Gesell.* (1870–71) 63.—Mexic.
stenophylla, Jacobi, in Hamb. Gartenz. xxii. (1866) 203 = vivipara.
striata, *Zucc. in Nov. Act. Nat. Cur.* xvi. ii. (1833) 678.—Mexic.
stricta, Salm Dyck, in Bonplandia, vii. (1859) 94 = striata.
subfalcata, Jacobi, in C. Koch & Fint. Wochenschr. xii. (1869) 179 = macracantha.
subinermis, *M. Roem. Syn. Ensat.* 289.—Bras.
sudburyensis, Hort. Pascock, ex Baker, in Gard. Chron. N. S. vii. (1877) ii. 137 = concinna.
Taylori *x*, *Hort. cf. Gard. Chron.* N. S. ii. (1874) 60 et vii. (1877) 621. f. 125.
Taylorsa x, Hort. Veitch. in Rev. Hortic. (1877) 36 = Taylori.
tehuacanensis, Karw. ex Otto in Otto & Dietr. Allg. Gartenz. (1842) 51 = atrovirens.

AGAVE :—
texasomuliana, Karw. ex M. Roem. Syn. Ensat. 292 = bromeliaefolia.
Theomenti, *Zuccagn. in Rem. Collect.* 138.—Mexic.
Thomsoniana, *Jacobi*, *in Hamb. Gartenz.* xxii. (1866) 203.—Mexic.
Toneliana, *Hort. Peacock*, *ex Baker*, *in Gard. Chron.* xv. (1881) i. 362.—Mexic. ?
triangularis, Jacobi, in C. Koch & Fint. Wochenschr. xii. (1869) 178 = horrida.
tuberosa, Mill. Gard. Dict. ed. VIII. n. 4 = Furcraea tuberosa.
uncinata, Jacobi, in Hamb. Gartenz. xxi. (1865) 165 = polyacantha.
undulata, *Klotzsch*, *in Otto & Dietr. Allg. Gartenz.* viii. (1840) 374.—Mexic.
univittata, *Haw. ex Salm-Dyck*, *Hort.* (1834) 308.— Mexic.
utahensis, *Engelm. in Bot. King, Exp.* 497.—Utah et Arizona.
Vandervinneni, *Lem. Illustr. Hortic.* xi. (1864), *Misc.* 64.—Mexic.
Vanderdonchii, Hort. ex Baker, in Gard. Chron. N. S. vii. (1877) i. 527 = xylonacantha.
variegata, Hort. ex Steud. Nom. ed. I. 18 = americana.
variegata, *Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 459.—Mexic.
Vera-Cruxx, Haw. Syn. Succ. 72 = seq.
Vera-cruz, Mill. Gard. Dict. ed. VIII. n. 7 = lurida.
Verschaffeltii, *Lem. ex Jacobi*, *in Hamb. Gartenz.* xxi. (1865) 122.—Mexic.
Victoriæ-Reginæ, *T. Moore*, *in Gard. Chron.* N. S. iv. (1875) ii. 485.—Mexic.
virginica, Hort. ex Baker, in Gard. Chron. N. S. xix. (1883) i. 176 = Allisertii.
virginica, *Linn. Sp. Pl.* 323.—Am. bor.
virginica, Mill. Gard. Dict. ed. VIII. n. 2 = americana.
viridissima, *Hort. Peacock*, *ex Baker*, *in Gard. Chron.* N. S. vii. (1877), ii. 137.—Mexic.
vivipara, Arruda, Diss. Pl. Bras. 23 = Furcraea agavephylla.
vivipara, *Linn. Sp. Pl.* 323.—Mexic.
Wallisii, *Jacobi*, *Nachtr.* ii. 78.—N. Granat.
Warelliana, *Hort. cf. Gard. Chron.* N. S. v. (1876), i. 603, *et Baker*, *l. c.* viii. (1877) ii. 264.—Mexic.
wiesenbergensis, *Wittm. in Berl. Gartenz.* (1885) 14.—Mexic.
Wislizeni, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 309.—Mexic.
xalapensis, Roezl, ex Jacobi, in Hamb. Gartenz. xxi. (1865) 61 = polyacantha.
xylacantha, Hort. cf. Gard. Chron. N. S. vii. (1877) i. 528 = xylonacantha.
xylonacantha, *Salm-Dyck*, *in Bonplandia*, vii. (1859) 92.—Mexico.
yuccaefolia, *DC. in Red. Lil.* t. 338, 329.—Mexic.
Yutii, Karw. ex G. Don, in Sweet, Hort. Brit. ed. III. 705 = rigida.
Zuccarinii, *Otto*, *in Allg. Gartenz.* x. (1842) 51, nomen. —Mexic. ?

AGDESTIS, Moç. & Sesse, ex DC. Syst. Veg. i. 542 (1818). *PHYTOLACCACEÆ*, Benth. & Hook. f. iii. 87.
clematidea, *Moç. & Sesse*, *l. c.*—Mexic.
tetrandra, De Not. Ind. Sem. Bot. Genuens. (1855) 39 = clematidea.

AGELAEA, Soland. ex Planch. in Linnaea, xxiii. (1850) 487. *CONNARACEÆ*, Benth. & Hook. f. i. 482.
HEMIANDRINA, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 171. t. 35.
ferruginea, Soland. ex Planch. l. c. 440 = Cnestis ferruginea.
glabriflora, *Hance*, *in Journ. Bot.* xiv. (1876) 287.— Cambodia.
Koneri, *O. Hoffm. & Hildebr. ex O. Hoffm. Sert. Madag.* 15.—Madag.
Lamarckii, *Planch. in Linnaea*, xxiii. (1850) 488.— Afr. trop.
nitida, *Soland. ex Planch. l. c.* 437.—Afr. trop.
oblique, *Baker*, *in Oliver*, *Fl. Trop. Afr.* i. 454.—Afr. trop.
pentagyna, Baill. in Bull. Soc. Linn. Par. i. (1882) 845 = Lamarckii.
pruriens, Soland. ex Planch. in Linnaea, xxiii. (1850) 440 = Cnestis corniculata.

AGELAEA :—
Thouarsiana, *Bull. Adansonia*, vii. (1866–67) 240.—Madag.
ventis, *Hook. f. Fl. Brit. Ind.* ii. 46.—Malaya.
villosa, *Soland. ex Planch. in Linnaea*, xxiii. (1850) 438.—Afr. trop.
Wallichii, *Hook. f. Fl. Brit. Ind.* ii. 47.—Malaya.

AGENIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447 (1836) = **Andropogon**, Linn. (Gramin.).
sulans, Nees, l. c. = A. Necsii.

AGENORA, D. Don, in Edinb. New Phil. Journ. (Mar. 1826) 210 = **Hypochoeris**, Linn. (Compos.).
laevigata, D. Don, l. c.—H. laevigata.

AGERATINA, Spach, Hist. Vég. Phan. x. 386 (1841) = **Eupatorium**, Linn. (Compos.).
agrestoides, Spach, l. c. = E. agrestoides.
aromatica, Spach, l. c. = E. aromaticum.
cordata, Spach, l. c. = E. aromaticum.

AGERATIOPSIS, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 246 (1873) = **Eupatorium**, Linn. (Compos.).
apiculata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 362 = E. apiculatum.

AGERATIUM, 'Adans.' ex Steud. Nom. ed. II. i. 87 (1841) = AGERATUM, Tourn. = **Brinus**, Linn. (Scrophul.).

AGERATIUM, Reichb. Consp. 209 (1828), err. typ = ACERATIUM, DC. = **Blascoacrpus**, Linn. (Tiliac.).

AGERATOIDES, Ponted. ex Linn. Class. Pl. 527 (1747) = **Achillea**, Linn. (Compos.).

AGERATUM, Linn. Gen. ed. I. 247 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 241.
CARLESTINA, Cass. Dict. Sc. Nat. vi. Suppl. 8. t. 93 (1817).
CARELIA, Adans. fl. 123 (1763).
COELESTINA, Cass. in Bull. Soc. Philom. (Jan. 1817) 10.
DECACHAETA, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 462.
OXYLOBUS, Moç. ex DC. Prod. v. 115 (1836).
adscendens, *Sch. Bip. ex Benth. & Hook. f. Gen.* ii. 243.—Mexic.
albidum, *Hemsl. Biol. Centr. Am. Bot.* ii. 81.—Mexic.
album, Steud. Nom. ed. I. 18 = conyzoides.
alternifolium, *Baker, in Mart. Fl. Bras.* vi. II. 195.—Bras.
altissimum, Linn. Sp. Pl. 839 = Eupatorium ageratoides.
angustifolium, Spreng. Syst. iii. 446 = Calea angustifolia.
aquaticum, Roxb. Hort. Beng. 61; et Fl. Ind. iii. 416 = Adenostemma viscosum.
arbutifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 149.—Mexic.
brachystephanum, *Regel, Gartenfl.* iii. (1854) 345. t. 108 f. c.—Caraccas.
caeruleum, Hort. ex Poir. Encyc. Suppl. i. 241 = conyzoides.
campuloclinioides, *Baker, in Mart. Fl. Bras.* vi. II. 196.—Bras.
ciliare, *Linn. Sp. Pl.* 839.—Ind. or. ?
ciliare, *Lour. Fl. Cochinch.* 484.—Cochinch.
coelestinum, Sims, Bot. Mag. t. 1730 = corymbosum.
corymbosum, Desf. Tabl. ed. I. 98 = conyzoides.
caeruleum, Sieber, ex Baker, in Mart. Fl. Bras. vi. II. 345 = Eupatorium macrophyllum.
confertum, *Baker, l. c.* 196.—Bras.
conspicuum, Hort. ex C. Koch & Fint. Wochenschr. i. (1858) 38 = Eupatorium glechonophyllum.
conyzoides, *Linn. Sp. Pl.* 839.—Keg. trop.
conyzoides, Sieber, ex Steud. Nom. ed. II. i. 37 = Eupatorium repandum.
cordifolium, Roxb. Fl. Ind. iii. 415 = conyzoides.
corymbosum, Benth. & Hook. f. Gen. ii. 242 = Agrianthus corymbosus.
corymbosum, *Zucc. ex Pers. Syn.* ii. 409.—Mexic.
dominguense, Spreng. Syst. iii. 446.—S. Doming.
echioides, *Hemsl. Biol. Centr. Am. Bot.* ii. 81.—Mexic.

AGERATUM :—
febrifugum, Sesse, ex DC. Prod. v. 104 = Piqueria trinervia.
glandulifolium, *Sch. Bip. ex Benth. & Hook. f. Gen.* ii. 242.—Mexic.
guianense, Aubl. Pl. Gui. ii. 800 = Eupatorium macrophyllum.
heterolepis, *Baker, in Mart. Fl. Bras.* vi. II. 198.—Bras.
hirsutum, Poir. Encyc. Suppl. i. 242 = seq.
hirtum, Lam. Encyc. i. 54 = conyzoides.
Houstonianum, *Mill. Gard. Dict.* ed. VIII. n. 2.—Mexic.
humile, Salisb. Prod. 168 = conyzoides.
iaocephalodes, *Hemsl. Biol. Centr. Am. Bot.* ii. 82.—Mexic.
Lassauxii, Carr. in Rev. Hortic. (1870–71) 90 = Conocliniun Lasaccuxii.
latifolium, Cav. Ic. iv. 33 = conyzoides.
laufolium, *Hemsl. Biol. Centr. Am. Bot.* ii. 82.—Guatemal.
linaare, Cav. Ic. III. 3. t 305 = Palafoxia linearis.
littorale, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 78.—Am. bor.
longifolium, *Baker, in Mart. Fl. Bras.* vi. II. 197.—Bras.
maritimum, H. B. & K. Nov. Gen. et Sp. iv. 150 = conyzoides.
matricaroides, *Less. Syn. Comp.* 155.—Cuba.
melissaefolium, *DC. Prod.* v. 109.—Bras.
mexicanum, Sims, Bot. Mag. t. 2524 = conyzoides.
mexicanum, Sweet, Brit. Fl. Gard. t. t. 89 = brachystephanum.
microcarpum, *Hemsl. Biol. Centr. Am. Bot.* ii. 82.—Costa-Rica.
microcephalum, *Hemsl. l. c.*—Mexic.
microphyllum, *Sch. Bip. in Seem. Bot. Voy. Herald*, 298.—Mexic.
muticum, *Griseb. Fl. Brit. W. Ind.* 358.—Jamaic.
nanum, Hort. ex Sch. Bip. in C. Koch & Fint. Wochenschr. i. (1858), Garten-Nachr. 26 = suffruticosum.
obtusifolium, Lam. Encyc. i. 54 = conyzoides.
odoratum, Vilm. Fl. Pl. Terre, ed. II. 42 = conyzoides.
palacatum, *Hemsl. Biol. Centr. Am. Bot.* ii. 83.—Mexic.
paniculatum, Hort. ex Steud. Nom. ed. II. i. 37 = Eupatorium paniculatum.
pedatum, Ortega, Hort. Matr. 38 = Florestina pedata.
petiolatum, *Hemsl. Biol. Centr. Am. Bot.* ii. 83.—Nicaragua.
Pohlianum, *Baker, in Mart. Fl. Bras.* vi. II. 197.—Bras.
punctatum, *Jacq. Hort. Schoenb.* iii. 28. t. 300.—Mexic.
punctatum, Ortega, Hort. Matr. 37 = Stevia Eupatoria.
purpureum, *Vitm. Summa Pl.* iv. 174.—Guian.
purpureum, Moç. & Sesse, ex DC. Prod. v. 122 = Stevia viscida.
quadriflorum, Blanco, Fl. Filip. ed. I. 624 = Elephantopus spicatus.
rubens, *Vic. Elench. Pl. Hort. Dinegro*, 9.—Hab. ?
salicifolium, *Hemsl. Biol. Centr. Am. Bot.* ii. 83.—Mexic.
scabrinodium, *Hemsl. l. c.*—Mexic.
scorpioideum, *Baker, in Mart. Fl. Bras.* vi. II. 197.—Am. trop.
sessilifolium, *Schau. in Linnaea*, xix. (1847) 715.—Mexic.
strictum, *Hemsl. Biol. Centr. Am. Bot.* ii. 83.—Mexic.
strictum, Sims, Bot. Mag. t. 2410 = Adenostemma viscosum.
suffruticosum, *Regel, Gartenfl.* iii. (1854) 389. t. 108.—Mexic.
tomentosum, *Hemsl. Biol. Centr. Am. Bot.* ii. 84.—Mexic.
viscosum, Ortega, Hort. Matr. 36 = Stevia lucida, salicifolia.
Wendlandii, *Hort. ex Vilm. Fl. Pl. Terre, Suppl.* 2.—Mexic.
Wrightii, *Torr. & Gray, in Proc. Am. Acad.* i. (1848) 46.—Texas.

AGERATUM, Tourn. ex Adans. Fam. ii. 210 (1763) = **Brinus**, Linn. (Scrophul.).

AGERELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139 = **Veronica**, Linn. (Scrophul.).
acinifolia, Fourr. l. c. = V. acinifolia.
armenia, Fourr. l. c. = V. arvensis.
verna, Fourr. l. c. = V. verna.

AGERIA, Adans. Fam. ii. 166 (1763) = **Ilex**, Linn.
acuminata, Rafin. Sylva Tellur. 46 = I. Maconcona.
Cassena, Rafin. l. c. 47 = I. Cassine.
graminata, Rafin. l. c. 48 = I. Cassine.
heterophylla, Rafin. l. c. = I. Dahoon.
mucronata, Rafin. l. c. 47.—Hab. ?
obovata, Rafin. l. c. 48 = I. Dahoon.
opaca, Rafin. l. c. 47 = I. opaca.
ovalis, Rafin. l. c. 48 = I. ovalifolia ?
palustris, Rafin. l. c. 47 = I. Dahoon.
retusa, Rafin. l. c. 46 = I. retusa.
uniflora, Rafin. l. c. 47 = I. uniflora.

AGGERANTHUS, Wight, Ic. v. 18. t. 1737 (1859) = **Elvia**, Lindl. (Orchid.).
marchantioides, Wight, l. c. = E. reticulata.

AGIALID, Adans. Fam. ii. 508 (1763) = **Balanites**, Delile (Simarub.).

AGIHALID, Alp. ex Juss. in Dict. Sc. Nat. i. 397 (1804) = prec.

AGINA, Neck. Elem. ii. 158 (1790) = **Sagina**, Linn. (Caryophyll.).

AGIRTA, Baill. Etud. Gén. Euph. 463 (1858) = **Tragia**, Linn. (Euphorb.).
Bolviniana, Baill. l. c. = T. Bolviniana.

AGLAEA, Steud. Nom. ed. I. 19 (1821) = **Malapsharsula**, Ker-Gawl. (Irid.).
graminea, Steud. l. c. = M. graminea.

AGLAIA, Allem. in Nov. Act. Nat. Cur. iv. (1770) 93 = **Cyperus**, Tourn. ?

AGLAIA, Lour. Fl. Cochinch. 173 (1790). *MELIACEAE*, Benth. & Hook. f. i. 354, 994.
AMERINA, Juss. Gen. 386 (1789).
ARGOPHILUM, Blanco, Fl. Filip. 186 (1837).
EUPHORA, Griff. Notul. iv. 547 (1854).
MILNEA, Roxb. Fl. Ind. ed. Wall. ii. 430 (1824).
NEMEDRA, Juss. in Mém. Mus. Par. xix. 228 (1830).
NYALELIA, Bennet. Schluess. Hort. Malab. 80 (1818).
SELBYA, M. Roem. Syn. Hesper. 89, 126 (1846).
acuminatissima, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* xvii. (1854) 42.—Malaya.
ancolana, Miq. Fl. Ind. Bat. Suppl. i. 506 = argentea.
andamanica, *Hiern, in Hook. Fl. Brit. Ind.* i. 555.—Ins. Andaman.
angustifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 55.—Sumatra.
apiocarpa, *Hiern, in Hook. Fl. Brit. Ind.* i. 555.—Zeylan.
argentea, *Blume, Bijdr.* 170.—Java.
aspera, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* xxvii. (1864) 42.—Java.
basiphylla, *A. Gray, Bot. U. St. Expl. Exped.* i. 287.—Ins. Fiji.
batjanica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 46.—Malaya.
chittagonga, Miq. l. c. 44 = Amoora chittagonga.
cordata, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 557.—Malacca.
coriacea, *Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 57.—Borneo.
crassinervia, *Kuro, ex Hiern, in Hook. f. Fl. Brit. Ind.* i. 556.—Tenasserim.
Cumingiana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 409 = Heamia Cumingiana.
decandra, Wall. Cat. n. 1276 = Amoora decandra.
denticulata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 410.—Luzon.
Dispanhorstii, *Miq. Fl. Ind. Bat. Suppl.* i. 507.—Sumatra.
Dookkoo, Griff. Notul. iv. 505 (= *Lansium domesticum*?).
edulis, *A. Gray, Bot. U. St. Expl. Exped.* i. 287.—Ind. or. ; Fiji.
edulis, Hiern, in Hook. Fl. Brit. Ind. iii. 556 = undulata, minutiflora.

AGLAIA :—
edulis, Seem. Fl. Vit. 37 = basiphylla.
elaeagnoides, *Benth. Fl. Austral.* i. 383.—Austral.
elliptica, Blume, Bijdr. 171 = Heamia elliptica.
eximia, Miq. Fl. Ind. Bat. Suppl. i. 506 = argentea.
Forstenii, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 46.—Amboina.
Ganggo, *Miq. Fl. Ind. Bat. Suppl.* i. 506.—Sumatra.
glabrata, Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. xxvii. (1864) 42 = Dysoxylum Halmaheirae.
glaziflora, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 555.—Malacca.
grandis, *Korth. in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 56.—Borneo.
grata, Wall. ex Voigt, Hort. Suburb. Calc. 138 = Roxburghiana.
Griffithii, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 146.—Malacca.
Halmaheirae, Miq. Ann. Mus. Bot. Lugd. Bat iv. 58 = Dysoxylum Halmaheirae.
hexandra, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 410.—Luzon.
hypoleuca, *Miq. Fl. Ind. Bat. Suppl.* i. 507 —Sumatra
inaequalis, Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. xxvii. (1864) 305 = Heamia elliptica.
khasiana, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 554.—Ind. or.
Korthalsii, *Miq. Ann. Mus. Bot. Lugd Bat.* iv. 42.—Sumatra.
latifolia, *Miq. l. c.*—Java.
laxiflora, *Miq. l. c.* 52.—Borneo.
lepidota, Miq. Fl. Ind. Bat. Suppl. i. 507 = Roxburghiana.
leptantha, *Miq. Ann. Mus. Bot. Lugd. Bat* iv 51.—Sumatra
littoralis, *Miq. l c.* 45.—Nov Guin.
Llanosiana, *C. DC. in DC. Monog Phan.* i. 621.—Ins. Philipp.
longifolia, *Teijsm. & Binn. in Tijdschr. Ned. Ind.* xxvii. (1864) 42.—Java.
macrobotrys, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 409.—Luzon.
macrophylla, Teijsm. & Binn. in Tijdschr. Ned. Ind. xxvii. (1864) 42 = Dysoxylum Halmaheirae.
madraporensis, Carey, ex Voigt, Hort. Suburb. Calc. 136 = Roxburghiana.
minutiflora, *Bedd. Fl. Ind. Or.* i. 44.—Ind. or.
mucronulata, *C. DC. in DC. Monog. Phan.* i. 601.—Java. multijuga, *Seem. Fl. Vit.* 37.—Ins. Fiji.
odorata, *Lour. Cochinch.* 173.—China.
odoratissima, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 218 = elaeagnoides.
odoratissima, Blume, Bijdr. 171 = Diepenhorstii.
oligantha, *C. DC. in DC. Monog. Phan.* i. 603.—Ins. Philipp.
oligocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 45.—Sumatra.
oligophylla, *Miq. Fl. Ind. Bat. Suppl.* i. 507.—Sumatra.
ovata, Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. xxvii. (1864) 43 = Heamia elliptica.
pachyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 57.—Sumatra.
palembanica, *Miq. Fl Ind. Bat. Suppl.* i. 507.—Borneo.
Pamattonis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 58.—Borneo.
paniculata, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 199.—Ind. or.
pedicellaris, *C. DC. in DC. Monog. Phan.* i. 607.—Ind. or.
pentaphylla, Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 48 = odorata.
perviridis, *Hiern, in Hook. Fl. Brit. Ind.* i. 556.—Ind. or.
pirifera, *Hance, in Journ. Bot.* xv. (1877) 331.—China.
polyantha, Bedd. Fl. Ind. Or. i. 44 = minutiflora.
polyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 56.—Java.
polystachya, Wall. in Roxb. Fl. Ind. ii. 429 = Amoora Rohituka.
pyenocarpa, *Miq. Fl. Ind. Bat. Suppl.* i. 196.—Sumatra.
pyramidata, *Hance, in Journ. Bot.* xv. (1877) 331.—China.
pyrrholepis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 47.—Java.
Reinwardtii, *Miq. l. c.* 51.—Celebes.
Roxburghiana, Kurz, in Journ. As. Soc. Beng. xliv. (1875) ii. 147 = undulata.

AGLAIA :—
Roxburghiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 41.—As. or.
rufa, *Miq. l. c.* 49.—Sumatra.
samoensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 336.—Ins. Navigat.
scripetala, Griff. Notul. iv. 505 = edulis.
sipamaea, Miq. Fl. Ind. Bat. Suppl. i. 506 = palembanica.
Spanoghei, Blume, ex Miq. l. c. 197 = Roxburghiana.
speciosa, *Blume, Bijdr.* 171.—Java.
speciosa, Teijsm. & Binn. Cat. Hort. Bogor. 211 = argentea.
spectabilis, Hort. Calc. ex C. DC. in DC. Monog. Phan. i. 595 = Amoora spectabilis.
subgrisea, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 54.—Malaya.
submonophylla, *Miq. l. c.* 40.—Borneo.
Sulingi, *Blume, Bijdr.* 170.—Java.
Sulingi, Hassk. Cat. Hort. Bogor. 220 = Diepenhorstii.
tenticulata, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 556.—Malaya.
Teysmanniana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 48.—Sumatra.
tomentosa, Teijsm. & Binn. in Tijdschr. Ned. Ind. xxvii. (1864) 43 = palembanica.
trichostemon, *C. DC. in DC. Monog. Phan.* i. 608.—Borneo.
Turczaninowii, *C. DC. l. c.* 628.—Ins. Philipp
undulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 44.—Ind. or.
Wallichii, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 555.—Ind. or.
Zippelii, *Miq. Ann Mus. Bot. Lugd. Bat.* iv. 54.—Nov. Guin.

AGLAIOPSIS, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 58 (1868–69) = **Haemia**, F. Muell. (Meliac.).
lanceifolia, Miq. l. c. 59 = H lanceifolia.
glaucescens, Miq. l. c. 58 = H. glaucescens.

AGLAIODENDRON, Remy, in Ann. Sc. Nat Sér. III. xii. (1879) 175 = **Plasia**, Ruiz & Pav. (Compos.).
cheiranthifolium, Remy, l. c. = P. cheiranthifolia.

AGLAODORUM, Schott, Gen. Aroid. t. 58 (1858) = **Aglaonema**, Schott (Aroid.).
Griffithii, Schott, l. c. = Aglaonema palustre.

AGLAONEMA, Schott, Melet. i. 20 (1829). *AROIDEAE*, Benth. & Hook. f. iii. 981.
AGLAORUM, Schott, Gen. Aroid. r. 58 (1858).
acutispathum, *N. E. Br. in Gard. Chron.* N. S. xxiv. (1885) ii. 89.—China.
commutatum, Schott, Syn. Aroid. 123 = Scindapsus Cuscuaria.
Cuscuaria, Miq. Fl. Ind. Bat iii. 217 = Scindapsus Cuscuaria.
fallax, Schott, ex Engl. in DC. Monog. ii. 439 = simplex.
gracile, Schott, in Ann. Mus. Lugd. Bat. i. 279 = pictum.
Hasskii, *Schott, in Bonplandia*, vii. (1859) 30.—Ins. Philipp.
Hookerianum, *Schott, l. c.*—Ind. or.
integrifolium, *Schott, Melet.* i. 20.—Malaya.
longicuspidatum, Schott, Prod. 304 = Schottianum.
maculatum, *Blume, Rumphia*, i. 154.—Ins. Philipp.
malaccense, Schott, in Bonplandia, vii. (1859) 30 = Schottianum.
marantifolium, *Blume, Rumphia*, i. 153. t. 66.—Malaya.
Mannii, *Hook. f. Bot. Mag.* t. 5760.—Afr. trop.
modestum, *Schott, ex Engl. in DC. Monog. Phan.* ii. 442.—Malaya.
nitidum, Kunth, Enum. Pl. iii. 56 = integrifolium.
oblongifolium, Kunth, l. c. 55 = marantifolium.
paludosum, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* xxv. (1863) 399.—Malaya.
pictum, *Kunth, Enum. Pl.* iii. 55.—Ind. or.; Malaya.
princeps, Kunth. l. c. = simplex.
propinquum, Schott, in Ann. Mus. Lugd. Bat. i. 280 = Schottianum.
pygmaeum, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 161 = Chamaecladon pygmaeum.
Schottianum, *Miq. Fl. Ind. Bat.* iii. 216.—Malaya.
simplex, *Blume, in Rumphia*, i. 153. t. 65.—Java.
subundulatum, *Zoll. Syst. Verz. Ind. Archip.* 76.—Celebes.

AGLITHEIS, Rafin. Fl. Tellur. ii. 17 (1836) = **Allium**, Linn. (Liliac.).
carinata, Rafin. l. c. 18 = All. carinatum.
tatarica, Rafin. l. c. = All. tataricum.
ursina, Rafin. l. c. = All. ursinum.

AGLOTOMA, Rafin. Fl. Tellur. ii. 44 (1836) = **Aster**, Linn. (Compos.).
multiflora, Rafin. l. c. = Aster multiflorus.

AGLYCIA, Willd. ex Stend. Nom. ed. II. i. 37 (1840) = **Erichloa**, Kunth (Gramin.).
coarctata, Willd. ex Steud. l. c. = E. polystachya.
distachya, Willd. ex Steud. l. c. = E. distachya.

AGNANTHUS, Vaill. ex Linn. Gen. ed. I. 366 (1737) = **Cornutia**, Linn. (Verben.).

AGNISTUS, G. Don, in Sweet, Hort. Brit. ed. III. 508 (1839). err. typ. = **Acnistus**, Schott (Solanac.).

AGNOSTUS, A. Cunn. in Lond. Hort. Brit. Suppl. i. 580 (1832) = **Stenocarpus**, R. Br. (Proteac.).
sinuatus, A. Cunn. l. c. = S. sinuatus.

AGNUS-CASTUS, [Tourn.] Carr. in Rev. Hortic. xlii. (1870–71) 418 = **Vitex**, Linn. (Verben.).
incisa, Carr. l. c. = V. Negundo.
Negundo, Carr. l. c. = V. Negundo.
robusta, Carr. l. c. = arg.]
vulgaris, Carr. l. c. 414 = V. Agnus-castus.

AGONANDRA, Miers, in Ann. Nat. Hist. Ser. II. viii. (1851) 179, nomen. *OLACINEAE*, Benth. & Hook f. i. 849.
brasiliensis, *Benth. & Hook. f. Gen.* i. 849.—Bras.
excelsa, *Griseb. in Gortt. Abh.* xxiv. (1879) 149.—Am. austr.

AGONIS, Lindl. Swan River App. 10 (1839). *MYRTACEAE*, Benth. & Hook. f. i. 708.
BILLOTTIA, R. Br. in Journ. Soc. Geogr. i. (1829) 19.
angustifolia, Schau. in Lehm. Pl. Preiss. i. 118 = linearifolia.
conspicua, Schau. l. c. = linearifolia.
flexuosa, *Lindl. Swan River App.* 10.—Austral.
floribunda, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 20.—Austral.
glabra, *Turcz. in Bull. Acad. Petersb.* x. (1852) 384. —Austral.
grandiflora, *Benth. Fl. Austral.* iii. 100.—Austral.
hypericifolia, *Schau. in Lehm. Pl. Preiss.* i. 117.—Austral.
juniperina, *Schau. l. c.* 118.—Austral.
linearifolia, *Lindl. Swan River App.* 10.—Austral.
marginata, *Schau. in Lehm. Pl. Preiss.* i. 117.—Austral.
obtusissima, *F. Muell. Fragm.* xi. 119.—Austral.
parviceps, *Schau. in Lehm. Pl. Preiss.* i. 119.—Austral.
Scortechniana, *F. Muell. Fragm.* xi. 118.—Austral.
spathulata, *Schau. in Lehm. Pl. Preiss.* i. 119.—Austral.
theaeformis, Schau. l. c. ii. 223 = hypericifolia.
undulata, *Benth. Fl. Austral.* iii. 100.—Austral.

AGONOLOBUS, Reichb. Nom. 188 (1841) = **Chirantthus**, Linn. (Crucif.).

AGONOMYRTUS, Schau. ex Reichb. Handb. 258 (1837) = **Leptospermum**, Forst. f (Myrtac.).

AGOPHYLLUM, Neck. Elem. ii. 227 (1790) = **Eygophyllum**, Linn.

AGORRHINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 126 = **Antirrhinum**, Tourn. (Scrophul.).
Orontium, Fourr. l. c. = Ant. Orontium.

AGOSERIS, Rafin. Fl. Ludov. 88 (1817) = **Troximon**, Nutt. (Compos.).
cuspidata, Steud. Nom. ed. II. i. 37 = T. marginatum.
glaucus, Steud. l. c. = T. glaucum.
parviflora, D. Dietr. Syn. Pl. iv. 1332 = T. parviflorum.
rosea, D. Dietr. l. c. = T. roseum.
taraxacifolia, D. Dietr. l. c. = T. taraxicifolium.

AGOSTANA, Bute, ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 236 (1821) = **Bupleurum**, Tourn. (Umbell.).
dioscorata, S. F. Gray, l. c. 237 = B. Odontites.
tenuissima, Bute, ex S. F. Gray, l. c. 236 = B. tenuissimum.

AGRAPHIS, Link, Handb. i. 166 (1829) = **Scilla**, Linn. (Liliac.).
campanulata, Link, l. c. = S. hispanica.
cernua, Reichb. Fl. Germ. Excurs. 105 = S. cernua.
nutans, Link, Handb. i. 166 = S. nutans.
patula, Reichb. Fl. Germ. Excurs. 106 = S. hispanica.
serotina, Heynh. Nom. ii. 14 = Dipcadi serotinum.

AGRAULUS, Beauv. Agrost. 5. t. 4 (1812) = **Agrostis**, Linn. (Gramin.).
alpinus, Beauv. l. c. = Agrostis rupestris.
anemagrostoides, Trin. in Spreng. N. Entdeck. ii. 56 = Agrostis effusa.
caninus, Beauv. Agrost. 5. t. 4, f. 7 = Agrostis canina.
flavescens, Panz. ex Steud. Nom. ed. I. 19, 840 = Festuca varia.
mexicanus, Presl, Reliq. Haenk. i. 236 = Agrostis tolucensis.
montanus, Panz. ex Steud. Nom. ed. I 19 = Agrostis alpina.
pallidus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 149 = Agrostis pallida.
setaceus, S. F. Gray, l. c. = Agrostis setacea.

AGRETTA, Eckl. Verz. 28 (1827) = **Tritonia**, Ker-Gawl. (Irid.).
crispa, Eckl. l. c. 24 = T. undulata.
grandiflora, Eckl. 23, nomen.—Afr. austr.
pallidiflorens, Eckl. l. c. 24.—Afr. austr.
pentandra, Eckl. l. c. 23 = T. scillaris.
rubusa, Steud. Nom. ed. II. i. 87.—Afr. austr.
stricta, Eckl. Verz. 23, nomen.—Afr. austr.

AGRIANTHUS, Mart. ex DC. Prod. v. 125 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 244.
campestris, *Mart. l. c.*—Bras.
corymbosus, *DC. l. c.* vii. 266.—Bras.
empetrifolius, *Mart. l. c.* v. 126.—Bras.

AGRICOLAEA, Schrank, in Denkschr. Akad. Muench. (1808) 98 = **Clerodendron**, Linn. (Verben.).
fragrans, Schrank, l. c. = C. fragrans.

AGRIMONIA, Tourn. ex Linn. Syst. ed. I. (1735). *ROSACEAE*, Benth. & Hook. f. i. 622.
AMONIA, Nestl. Pomet. 17 (1814).
AREMONIA, Neck. Elem. ii. 100 (1790).
STALLENANIA, Pollini, Hort. Veron. Pl. Nov. 10 (1816).
acutifolia, Dum. Fl. Belg. Prod. 97 = odorata.
adhaerens, Gillib. Fl. Lituan. ii. 185 = A. Eupatorium.
adscendens, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 211, nomen.—Kossia.
agrimonoides, *Linn. Sp. Pl.* 448.—Europ.
americana, Lucs, ex Wallr. Beitr. Bot. i. 43 = A. Eupatoria.
Blumei, *G. Don, Gen. Syst.* ii. 563.—Java; Japon.
bracteata, E. Mey. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 349 = A. Eupatoria.
bracteosa, E. Mey. Zwei Pfl. Docum. 51, 142, nomen = A. Eupatoria.
cagra, E. Mey. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 349 = A. Eupatoria.
canescens, Dum. Fl. Belg. Prod. 97 = odorata.
ceylonica, Wight, Ic. t. 934 = zeylanica.
conspicua, Tschern. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 346 = pilosa.
dahurica, Willd. ex Stev. ex DC. Prod. ii. 587 = pilosa.
decurrens, Schlecht. pat. ex Ledeb. Fl. Ross. ii. 82 = pracc.
decumbens, Linn. f. Suppl. 251 = Acaena latebrosa.
elata, Moench, Meth. 674 = repens.
elata, Salisb. Prod. 381 = A. Eupatoria.
Eupatoria, Michx. Fl. Am. Bor. i. 287 = parviflora.
Eupatoria, *Linn. Sp. Pl.* 448.—Reg. bor. temp.
Eupatorium, Thw. Enum. Pl. Zeyl. 109 = zeylanica.
eupatoroides, Gueldenst. ex Ledeb. Fl. Ross. ii. 81 = A. Eupatoria.
glabrata, Spreng. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 346 = A. Eupatoria.
Gochetiana, Andrz. ex C. A. Mey. l. c. = A. Eupatoria.

AGRIMONIA :—
graeca, *Boiss. & Heldr. ex Boiss. Fl. Orient.* ii. 728 = A. Eupatoria.
grandis, Andrz. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 343 = A. Eupatoria.
gryposepala, Wallr. Beitr. Bot. i. 49 = A. Eupatoria.
hirsuta, *Bong. ex C. A. Mey. in Bull. Sc. Acad. Pétersb.* x. (1842) 347.—Bras.
humilis, *Wallr. Beitr. Bot.* i. 87.—Dalmat.
incisa, *Torr. & Gray, Fl. N. Am.* i. 481.—Am. bor.
inermis, Pers. Syn. ii. 9 = agrimonioides.
javanica, Jungh. Java, i. 664, ex Miq. Fl. Ind. Bat. i. 1. 570 = Blumei.
lanata, Wall. Cat. n. 709 = A. Eupatoria.
leucantha, *Kunze, Del. Sem. Hort. Lips.* (1849).—Hab. ?
major, Urpensky, in Bull. Soc. Nat. Mosc. vii. (1834) 373 = odorata ?
microcarpa, Wallr. Beitr. Bot. i. 39 = A. Eupatoria.
minor, Mill. Gard. Dict. ed. VIII. n. 2 = A. Eupatoria.
nepalensis, D. Don, Prod. Fl. Nep. 229 = A. Eupatoria.
octandra, *Luce, Topogr. Nachr. Ins. Oesel*, 150.—Ins. Oesil.
odorata, *Mill. Gard. Dict.* ed. VIII. n. 3.—Europ.
officinalis, Lam. Encyc. i. 62 = A. Eupatoria.
officinarum, Lam. Fl. Fr. iii. 477 = A. Eupatoria.
parviflora, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 130. —Am. austr.
parviflora, Spreng. Syst. ii. 461 = A. Eupatoria.
pilosa, Ledeb. Ind. Hort Dorp. Suppl. (1822) 1 = A. Eupatoria.
platycarpa, Wallr. Beitr. Bot. i. 38 = A. Eupatoria.
procera, Wallr. Beitr. Fl. Hercyn. 303; in Linnaea, xiv. (1840) 573 = odorata.
pubescens, Wallr. Beitr. Bot. i. 45 = A. Eupatoria.
pumila, Muhl. ex Rafin. Fl. Ludov. 134, nomen. —Am. bor.
repens, Eckl & Zey. Enum. 264 = A. Eupatoria.
repens, *Linn Syst.* ed. X. 1046.—Hispan.; As. Min.
repens, Mutel, Fl. Fr. i. 342 = odorata.
robusta, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 211 = odorata ?
rostellata, Wallr. Beitr. Bot. i. 49 = A. Eupatoria.
serrifolia, Wallr. l. c. 40 = A. parviflora.
sessiliflora, Dum. Fl. Belg. Prod. 97 = odorata.
sororia, *Fisch, & Mey. ex C. A. Mey. in Bull. Sc. Acad. Pétersb.* x. (1842) 344.—Orient.
stipularia, Dum. Fl. Belg. Prod. 97 = odorata.
striata, Michx. Fl. Bor. Am. i. 287 = A. Eupatoria.
suaveolens, Hort. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 344 = odorata.
suaveolens, Blume, Bijdr. 1118 = Blumei.
suaveolens, Pursh, Fl. Am. Sept. i. 336 = parviflora.
sulcata, Dulac, Fl. Hautes-Pyr. 208 = A. Eupatoria.
viscidula, Bunge, in Mém. Sav. étr. St. Pétersb. ii. 100; (Enum. Pl. Chin. Bor. 26) = A. Eupatoria.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 577 = A. Eupatoria.
zeylanica, Moon, Cat. 37 = Alchemella indica.

AGRIODENDRON, Endl. Gen. 144 (1837) = **Aloë**, Linn. (Liliac.).

AGRIOPHYLLUM, Bieb. Fl. Taur. Cauc. iii. 6 (1819). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 84.
arenarium, *Bieb. l. c.*—Ax. bor.
arenarium, *Bunge, in Mém. Sav. étr. Pétersb.* vii. (1851) 456 (*Lehm. Rel.* 280).—As. temp.
gobicum, *Bunge, in Bull. Acad. St. Pétersb.* xxv. (1879) 855.—Sibir.
lateriflorum, *Moq. in DC. Prod.* xiii. ii. 139.—Armen.
latifolium, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 170.—Orient.
minus, *Fisch. & Mey. l. c.*—Turkest.
mongolicum, Hort. Aven. ex Moq. in DC. Prod. xiii. ii. 139 = arenarium.
pungens, Link, ex A. Dietr. Sp. Pl. i. 924 = Corispermum pungens.
squarrosum, Moq. in DC. Prod. xiii. ii. 139 = arenarium.
Tournefortii, Fisch. ex Moq. l. c. = lateriflorum.

AGRIPHYLLUM, Juss. Gen. 190 (1789) = **Berkheya**, Ehrh. (Compos.).
asteroides, Desf. Tabl. Hort. Par. ed. I. 109 = B. fruticosa.

AGRIPHYLLUM :—
ciliare, Desf. Cat. Hort. Par. ed. III. 186 = Collumia ciliaris.
echinatum, Desf. Tabl. Hort. Par. ed. I. 109 = B. cernua.
grandiflorum, Desf. Cat. Hort. Par. ed. II. 186 = B. grandiflora.
incanum, Desf. l. c. = B. fruticosa.
spinosum, Desf. ex Steud. Nom. ed. I. 19, 106 = B. fruticosa.

AGROCHARIS, Hochst. in Flora, xxvii. (1844) 1. 19 = **Caucalis**, Linn. (Umbell.).
gracilis, Hook. f. in Journ. Linn. Soc. vi. (1860) 9 = C. melanantha.
melanantha, Hochst. in Flora, xxvii. (1844) 1. 19 = C. melanantha.

AGROPHYLLUM, Neck. Elem. ii. 237 (1790) = **Zygophyllum**, Linn.

AGROPYRON, J. Gaert. in Nov. Comm. Petrop. xiv. (1770) i. 539. *GRAMINEAE* [Benth. & Hook. f. iii. 1209].
ANTHOSACHNE, Steud. Syn. Pl. Gram. 287 (1855).
BRACONOTIA, Godr. l. c. ed. I. iii. 191 (1844).
COSTIA, Willk. in Bot. Zeit. (1858) 377.
CREMOPYRUM, Schur, Enum. Pl. Transs. 807 (1866).
CRITHOPYRUM, Hort. Prag. ex Steud. Syn. Pl. Gram. 344 (1855).
ELYTRIGIA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190.
EREMOPYRUM, Jaub. & Spach, Illustr. Pl. Or. iv. 26 (1850–53).
HAYNALDIA, Schur, Enum. Pl. Transs. 807 (1866).
HETERANTHELIUM, Hochst. ex Janb. & Spach, Illustr. Pl. Or. iv. 24 (1850).
ROEGNERIA, C. Koch, in Linnaea, xxi. (1848) 418.
SECALIDIUM, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 91.
acutum, C. Koch, in Linnaea, xxi. (1848) 424 = repens.
acutum, *Roem. & Schult. Syst.* ii. 751.—Europ.
acutum, Reichb. ex Nym. Consp. 841 = pungens.
affine, Reichb. Ic. Fl. Germ. xi. 7 = acutum.
alpinum, Schur, Enum. Pl. Transs. 810.—Europ.
altissimum, Schur, l. c. 809 = repens.
angustifolium, Schult. Mant. ii. 413.—Sibir.
apricum, *Opiz, in Lotos*, iii. (1853) 63.—Europ.
aristatum, Bess. Enum. Pl. Volh. 41 = cristatum.
attenuatum, Roem. & Schult. Syst. ii. 751.—Ecuador.
Aucheri, *Boiss. Diagn. Ser.* I. v. 75.—Orient.
barbulatum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 91 = seq.
barbulatum, *Schur, Enum. Pl. Transs.* 808.—Europ.
biconne, Roem. & Schult. Syst. ii. 760 = Triticum biconne.
biflorum, *Roem. & Schult. l. c.*—Syria.
biforme, Flg. & De Not. in Mem. Torin. Ser. II. xii. (1853) 266 = panicum.
Bourgaei, *Boiss. Fl. Orient.* v. 669.—Armen.
brachyphyllum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 668.—Persia.
brevianum, Beauv. Agrost. 102 (aphalm. pro laevissimo) = Triticum polonicum.
bromiforme, Schur, Enum. Pl. Transs. 808 = repens.
bulbosum, *Boiss. Diagn. Ser.* I. v. 75.—Persia.
caesium, J. & C. Presl, Delic. Prag. 213 = repens.
caespitosum, *C. Koch, in Linnaea*, xxi. (1848) 424.—Armen.
Caldesii, *Goran. in Nuov. Giorn. Bot. ital.* xii. (1880) 145. —Ital.
campestre, *Godr. & Gren. Fl. Fr.* iii. 607.—Europ.
caninum, *Beauv. Agrost.* 102.—Europ.
capillare, Beauv. l. c. = repens.
caudatum, Beauv. l. c. = villosum.
collinum, Opiz, Natural. x. (1828) 246; Seznam, 12 = repens.
concinnum, De Not. ex Parl. Fl. Ital. i. 498 = pungens.
condensatum, *Presl, Rel. Haenk.* i. 296.—Chili.
crinitum, *Lindl. Hort. Beverl.* i. 32.—Hab. ?
cristatum, *Y. Gaert. in Nov. Comm. Petrop.* xiv. (1770) i. 540.—Iberia; Tauria; Sibir.
cristatum, *Beauv. Agrost.* 102; *Roem. & Schult. Syst.* ii. 756.—Hab. ?
curvifolium, Lange, in Kjoeb. Vidensk. Meddel. (1860) 56 = scirpeum.
dasyanthum, *Ledeb. in Ind. Sem. Dorpat.* (1839) 3.—Kossia.

AGROPYRON :—
dasystachyum, *Scrib. in Torrey Bot. Club.* x. (1888) 76. —Am. bor.
densiflorum, *Beauv. Agrost.* 102.—Sibir.
desertorum, Schult. Mant. ii. 412 = sibiricum ?
distans, C. Koch, in Linnaea, xxi. (1848) 426 = Bourgaei.
distichum, *Beauv. Agrost.* 102.—Afr. austr.
divaricatum, *Boiss. & Bal. in Bull. Soc. Bot. Fr.* lv. (1857) 307.—Orient.
divergens, *Nees, ex Steud. Syn. Pl. Gram.* 347.—Am. bor.
dumetorum, *Trautv. in Act. Hort. Petrop.* ix. (1884) 329.—Europ.
elatum, Opiz, Seznam, 12 = cristatum.
elongatum, Host, ex Beauv. Agrost. 102 = rigidum.
elymogenes, Arndt, ex Nym. Consp. 840 = strictum.
farctum, Boiss. Fl. Orient. v. 665 in syn. = junceum filiforme, *Poir. ex Roem. & Schult. Syst.* ii. 760.—Ins. Borbon.
firmum, J. & C. Presl, Fl. Cech. 28 = repens.
geminatum, Schult. Mant. iii 655 = Eleusine indica.
gigantеum, Roem. & Schult. Syst. ii. 758 = rigidum.
glaucum, Blanco, ex Nym. Consp. 841 = repens.
glaucum, *Roem. & Schult. Syst.* ii. 752.—Europ.
Halleri, Reichb. Fl. Germ. Excurs. 20 = Festuca Poa.
hirsutum, Hort. Berol. ex Ledeb. Fl. Alt. i. 114 = cristatum.
hispanicum, Presl, Cyp. et Gram. Sic. 49 = Festuca Poa.
hordeaceum, Boiss. Diagn. Ser. I. xiii. 67 = orientale.
imbricatum, *Roem. & Schult. Syst.* ii. 757.—Rossia.
intermedium, Beauv. Agrost. 102 = glaucum.
intermedium, Reichb. Fl. Germ. Excurs 140 = glaucum, junceum, et obtusiflorum.
junceum, *Beauv. Agrost.* 102.—Europ.
Kotschyanum, Boiss. & Hohenh. Diagn. Ser. I. xiii. 69 = patulum.
laevissimum, Besser, Agrost. 146 = Triticum polonicum.
lasianthum, Boiss. Diagn Ser. I xiii. 68 = orientale.
laxum, ' Fr.' ex Willk. & Lange, Prod. i. 109 = acutum.
lasium, *Boiss. Fl. Orient.* v. 661.—As. Min.
ligusticum, *De Not. in Atti Ottav. Riun. Sc. Ital.* 602.—Ital.
ligusticum, Savign. in Flora, xxx. (1847) 569 = Triticum ligusticum.
littorale, *Dum. Obs. Gram. Belg.* 97.—Europ
loliiforme, Schur, Enum. Pl. Transs. 808 = repens.
longearistatum, *Boiss. Fl. Orient.* v. 660.—Orient.
longiglume, *Hack. ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 11.—Europ.
maritimum, Beauv. Agrost. 102 = Cutandia maritima.
mucronatum, Opiz, Seznam, 12 = glaucum.
multiflorum, Beauv. Agrost. 102 = repens.
muricatum, Ehrh. Skizze, 119 = cristatum.
muricatum, *Schult. Mant.* ii. 414.—Sibir.
nubigenum, *Nees, ex Steud. Syn. Pl. Gram.* 342.—Java.
obtusiflorum, *Roem. & Schult. Syst.* ii. 758 —Belaur.
obtusiusculum, *Lange, Handb.* ed. II. 48.—Dan.; Scan.
orientale, *Roem. & Schult. Syst.* ii. 757.—Orient.; *Afr.* bor.
panormitanum, *Parl. ex Boiss. Voy.* 680; *Pl. Rar. Sic. Fasc.* ii. 20.—Europ.; Orient.
patulum, *Trin. Fund.* 183.—Aegypt.
pauciflorum, *Schur, in Verh. Siebenb. Ver. Naturw.* x. (1859) 77.—Europ.
pectinatum, *Beauv. Agrost.* 102.—Austral.; N. Zel.
pectiniforme, *Roem. & Schult. Syst.* ii. 758.—Tauria.
peruvianum, *Roem. & Schult. l. c.* 761.—Peruv.
pilosum, *Presl, Relig. Haenk.* i. 267.—Mexic.
pilosum, Schur, Enum. Pl. Transs. 809 = barbulatum.
Pousolzii, Godr. & Gren. Fl. Fr. iii. 608 = glaucum.
prostratum, Beauv. Agrost. 102 = tritioncum.
pseudo-caninum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 91 = caninum.
pumilum, Beauv. Agrost. 102 = tritioeum.
pungens, Reichb. ex Nym. Consp. 840 = acutum.
pungens, *Roem. & Schult. Syst.* ii. 753.—Europ.
pycnanthum, Gren. & Godr. Fl. Fr. iii. 606 = littorale.
Raegneri, *Boiss. Fl. Orient.* v. 663.—Reg. Caucas.
repens, *Beauv. Agrost.* 104.—Reg. temp.
Richardsoni, *Schrad. in Linnaea*, xii. (1838) 467 in syn.—Am. bor.
rigidum, *Beauv. Agrost.* 102.—Europ.
Rothii, *Opiz, in Lotos*, iii. (1853) 84.—Europ.
Rouxii, *Gren. & Duval, in Mém. Soc. Emul. Doubs*, III. iv. (1880) 291.—Europ.
rupestre, *Schult. Mant.* ii. 409.—Hab. ?
Sardagnolii, *De Not. Prosp. Fl. Ligust.* 57.—Liguria.

AGROPYRON :—
scabrum, *Beauv. Agrost.* 102.—Austral.
scirpeum, *Presl, Gram. Sic.* 49.—Sicilia.
Scribneri, *Vasey, in Bull. Torrey, Bot. Club*, x. (1883) 128.—Am. bor.
secundum, *Presl, Rel. Haenk.* i. 266.—Chili.
semicostatum, *Nees, ex Steud. Syn. Pl. Gram.* 346.—Afghan.
sepium, Besav. Agrost. 103 = repens.
sibiricum, *Beauv. l. c.*—Europ. et : As. occ.
squarrosum, Link, Hort. Berol. i. 82 = patulum.
stipaefolium, Cacrn. Consp. Fl. Charcov. 70, nomen.
subulosum, Nees, ex Steud. Syn. Pl. Gram. 346 = Triticum striatum.
strictum, *Reichb. in Moesel. Handb.* ed. II. iii. 1812.—Europ.
strigosum, *Boiss. Fl. Orient.* v. 661.—Orient.
subulatum, *Roem. & Schult. Syst.* ii. 761.—Aleppo.
Tauri, *Boiss. & Bal. in Bull. Soc. Bot. Fr.* iv. (1857) 307.—As. Min.
teneruin, *Vasey, in Coult. Bot. Gaz.* x. (1885) 258.—Am. bor.
tenuiculum, Steud. Nom. ed II. i. 88 = Festuca Poa.
Tourneforti, *De Not. in Atti Ottav. Riun. Sc. Ital.* 602.—Oriena.
Tournefortii, Savign. in Flora, xxx. (1847) 569 = Asticteri.
triticeum, *Y. Gaertn. in Nov. Comm. Petrop.* xiv. (1770) i. 539.—Europ. ; As. bor.
truncatum, *Pauz, Fl. Transs.* 749.—Europ.
unilaterale, Benuv. Agrost. 102 = Festuca tenuiflora.
vaginans, *Beauv. Agrost.* 102.—Am. merid.
Vaillantianum, *Trautv. in Act. Hort. Petrop.* ix. (1884) 330.—Europ.
variegatum, *Roem. & Schult. Syst.* ii. 759.—Hab. ?
velutinum, *Nees, in Hook. Lond. Journ. Bot.* ii. (1843) 417.—Austral.
villosum, *Link, Hort. Berol.* i. 81.—Europ. ; Oriens.
violaceum, *Vasey, Gram. Un. St.* 45.—Am. bor.

AGROPYRUM, Roem. & Schult. Syst. ii. 750 (1817) = praec.

AGROSINAPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 329 = Brassica, Tourn. (Cruciff.).
arvensis, Fourr. l. c. = B. arvensis.

AGROSTANA, Hill, Veg. Syst. v. 32 (1772) = Bupleurum, Linn. (Umbell.).

AGROSTEMA, Linn. Syst. ed. I. (1735) = seq.

AGROSTEMMA, Linn. Gen. ed. I. 185 (1737) = Lychnis, Linn. (Caryophyll.).
apetala, G. Don, Gen. Syst. i. 416 = L. apetala.
alpina, Forbes, Hort. Woburn. 104 = L. alpina.
Banksia, Meerb. Fl. Sel. 28 = L. coronaria.
brachypetala, Sweet, Hort. Brit. ed. I. 44 = L. apetala.
Bungeana, D. Don, in Sweet, Fl. Gard. Ser. II. 317 = Bungeana.
Coeli-rosea, Linn. Sp. Pl. 436 = L. Coeli-rosa.
Coronaria, Linn. l. c. = L. Coronaria.
corona, G. Don, Gen. Syst. i. 416 = L. corsica.
declinis, G. Don, l. c. = L. diclinis.
decumbens, Desaoo, ex Loud. Hort. Brit. Suppl. i. 580 = L. Lucombeana.
dioica, Dufour, ex G. Don, Gen. Syst. i. 416 = L. diclinis.
fimbriata, G. Don, l. c. 417 = L. indica.
Flos-cuculi, G. Don, l. c. = L. Flos-cuculi.
Flos-jovis, Linn. Sp. Pl. 436 = L. Flos-jovis.
Flos-jovis, Pollich, Hist. Pl. Palat. i. 443 = L. coronaria.
fulgens, Doell, Rhein. Fl. 643 = L. fulgens.
Githago, Linn. Sp. Pl. ed. II. 635 = L. Githago.
Githago, Retz. Fl. Scand. 86 = praec.
gracilis, Boiss. Diagn. Ser. II. i. 80.—As. Min.
grandiflora, Doell, Rhein. Fl. 643 = L. coronaria.
hirsuta, Gilib. Fl. Lituan. ii. 173 = L. Githago.
hirsuta, Stokes, Bot. Mat. Med. ii. 559 = L. Githago.
inflata, G. Don, Gen. Syst. i. 417 = L. inflata.
involucrata, G. Don, l. c. 416 = L. apetala.
laeta, G. Don, l. c. = Silene laeta.
lusitanica, G. Don, l. c. 417 = L. Flos-cuculi.
nicaeensis, Pers. Syn. i. 519 = L. Githago.
nivalis, G. Don, Gen. Syst. i. 416 = L. nivalis.
parviflora, Cav. ex Steud. Nom. ed. I. 19 = Silene laeta.
perennans, Haw. ex. Loud. Hort. Brit. Suppl. i. 580 = L. Dianoni.
pulchra, G. Don, Gen. Syst. i. 417 = Silene laciniata.

AGROSTEMMA :—
pusilla, Desf. ex Steud. Nom. ed. II. i. 88 = L. Coeli-rosa.
pyrenaica, G. Don, Gen. Syst. i. 417 = L. pyrenaica.
sanatilis, G. Don, in Sweet, Hort. Brit. ed. III. 67 = L. sanatilis.
sibirica, G. Don, Gen. Syst. i. 416 = L. sibirica.
suecica, Maund. ex Steud. Nom. ed. II. i. 88 = L. alpina.
sylvestris, G. Don, Gen. Syst. i. 416 = L. dioica.
trifiora, G. Don, l. c. 417 = L. trifiora.
variegata, G. Don, l. c. 416 = L. variegata.

AGROSTICULA, Raddi, Agrost. Bras. 83. t. 1. f. 2 (1823) = Sporobolus, R. Br. (Gram.).
muralis, Raddi, l. c. = S. minutiflorus.

AGROSTIS, Linn. Syst. ed. I. (1735) ; Gen. ed. I. 19 (1737). *GRAMINEAE.* Benth. & Hook. f. iii. 1149.
AGRAULUS, Besuv. Agrost. 5. t. 4 (1812).
AVENA, Hall. ex Scop. Introd. 74 (1777).
DRCANDOLIA, Beat.]'l. Malac.-et-Loire, 28 (1809).
NORONEMA, Rafin. Neogen. 4 (1832).
PERRILEMA, Presl, Reliq. Haenk. i. 288. t. 37 (1830).
TRICHODIUM, Michx. Fl. Bor. Am. i. 41. t. 8 (1803).
VILFA, Adans. Fam. ii. 495 (1763).
abahanensis, Less. ex Trin. in Mém. Acad. Péterab. Sér. VI. vi. 11. (1845) 325 = Michauxii.
aberrans, *Steud. Syn. Pl. Gram.* 422.—Reg. Magell.
actinoclada, F. Muell. Fragm. vi. 84 = Sporobolus actinocladus.
aculeata, Scop. Fl. Carn. ed. II. i. 62 = Crypsis aculeata.
acutiflora, Steud. Nom. ed. I. 19 = Deyeuxia acutiflora.
adscendens, Lange, in Kjoeb. Vidensk. Meddel (1860) 88 = verticillata.
aemula, K. Br. Prod. 172 = Deyeuxia Forsteri.
aenea, Spreng. Syst. Cur. Post. 837 = Sporobolus aeneus.
aeneus, *Trin. in Mém. Acad. Péterab.* Sér. VI. vi. 11. (1845) 332.—Sitcha.
aequata, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 418 = Deyeuxia aequata.
aequivalvis, *Trin. in Mém. Acad. Péterab.* Sér. VI. vi. 11. (1846) 362.—Am. bor. occ.
affinis, Kunth, Enum. Pl. i. 228 = Deyeuxia.
affinis, Schult. Mant. ii. 195 = Sporobolus Muehlenbergii.
africana, Poir. Encyc. Suppl. i. 354 = Sporobolus capensis.
airaeformis, *Steud. Syn. Pl. Gram.* 173.—Chili.
airoides, *Torr. ex Spreng. Syst. Cur. Post.* 31 = Sporobolus airoides.
alba, Kunze, in Flora, xxix. (1846) 678 = Reuterii.
alba, *Linn. Sp. Pl.* 63.—Europ. ; As.
albicans, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863), 91 = exarata.
albicans, Presl, ex Steud. Nom. ed. I. 19.—Hab. ?
alfieda, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 344.—Sarepta ; Rossia.
algida, (Soland. in) Phipps, Voy. 200 = Phippsia algida.
alopecuroides, Lam. Illustr. n. 812 = Polypogon monspeliensis.
alopecuroides, Link, ex Kunth, Enum. Pl. i. 228 = effusa.
alopecuroides, Vahl, ex Kunth, Enum. Pl. i. 86 = Panicum Myurus.
alpestris, Laest. ex Nym. Consp. 802 = rubra.
alpicola, *Hochst. in Flora*, xxxviii. (1855) 284.—Ab. or.
alpina, Leyss. Fl. Halen. 16 = canina.
alpina, Willd. Sp. Pl. i. 365 = rupestris.
alpina, Savi [Fl. Pisana, i. 81 ?], ex Steud. Nom. ed. I. 19 = vulgaris.
alpina, *Scop. Fl. Carn.* ed. II. i. 60.—Europ.
alpina, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. i. 71 = setacea.
altissima, *Tuckerm. in Am. Journ. Sc.* xiv. (1843) 44 = elata.
ambigua, Roem. & Schult. Syst. ii. 354 = alba.
anatolica, C. Koch, in Linnaea, xxi. (1848) 379 = verticillata.
andina, *Hook. in Anales Univ. Chil.* (1878) 561.—Chili.
Anemagrostis, Syme, Eng. Bot. ed. III. xi. 43 = Apera Spica-venti, et interrupta.
anemagrostoides, Trin. in Spreng. Neue Entdeck. ii. 56 = effusa.
anomala, Willd. Sp. Pl. i. 370 = elata.
antarctica, *Hook. f. Fl. Antarct.* 374. t. 132 —Reg. Antarct.
Antoniana, *Griseb. in Goett. Abh.* xxiv. (1879) 298.—Reg. Argent.
apetala, Besv. ex Trin. in Mém Acad. Péterab. Sér. VI. vi 11. (1845) 327 = Muehlenbergia diffusa.

AGROSTIS :—

sphaeae, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1846) 346.—N. Zembl.

aquatica, *Buckl. in Proc. Ac. Sc. Philad.* '1862.' (1863) 90.—Am. bor.

aquatica, Pourr. in Mém. Acad. Toul. iii. (1788) 306 = verticillata.

arachnoidea, Poir. Encyc. Suppl. i. 249 = Cinna arundinacea.

arachnoides, *Ell. Sketch*, i. 124.—Am. bor.

arenaria, Gouan, Illustr. 8 = Sporobolus pungens

arenaria, Schur, in Oestr. Bot. Zeitschr ix. (1859) 51 = amina.

armosa, Schar, Enum. Pl. Transs. 736 = canina.

argentea, Lam. Fl. Fr. iii 570 = Stipa Calamagrostis.

aristata, Sinel ex Steud. Nom. ed II. i. 39 = alba.

aristulata, *C. Muell. in Bot. Zeit.* xiv. (1856) 349.—Ins. Falkland.

articulata, Brot. Fl. Lusit. i 73 = Chaeturus fasciculatus.

arundinacea, Linn Sp. Pl. 61 = Deyeuxia sylvatica.

arundinacea, *Presl, in Rel. Haenk.* i. 238.—Peruv.

arundinacea, Vill. Hist. Pl. Dauph. ii. 79 = Calamagrostis varia.

arvensis, *Phil. in Linnaea*, xxix. (1857–58) 87.—Chili.

aspera, Michx Fl. Am Bor. i. 58 = Sporobolus asper.

aspera, Willd. ex Steud. Nom. ed. II. i. 39 = Muehlenbergia stipoides.

aspertifolia, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. 11. (1846) 317 = exarata.

asperiglume, *Steud. Syn. Pl. Gram.* 422.—Chili.

asperula, *Phil. in Linnaea*, xxix. (1857–58) 87.—Chili.

atropurpurea, Hort. Hafn. ex Steud. Nom. ed. II. i. 39 = vulgaris.

atrovirens, Roem. & Schult. Syst. ii. 361 = Sporobolus atrovirens.

auchlandica, Hook. f. Fl. Antarct i. 96 = Deschampsia caespitosa.

australis, *Lam. Encyc.* i. 57.—Europ.

australis, Linn. Mant. i. 30 = Gastridium australe.

avenacea, J. F. Gmel. Syst. i. 171 = Deyeuxia Forsteri.

avenoides, *Hook. f. Handb. New Zeal. Fl.* 330.—N. Zel.

barbata, Buckl. in Proc. Acad. Sc. Philad. '1862.' (1863) 334 = Sporobolus ramulosus.

barbata, Banks & Soland. ex Hook. f. Fl. Nov. Zel. i. 293 = Dichelachne crinita.

barbata, Pers. Syn. i. 75 = Sporobolus virginicus.

Bergiana, *Trin. Gram. Unifl.* 203.—Afr. austr.

bervica, Hort. Madr. ex Steud. Nom. ed. II. i. 39 (= *Urachne parviflora*).

Berlandieri, '*Pourn. Mex. Pl. Lu. Gram.* 96,' *ex Hemsl. Biol. Centr. Am. Bot.* iii. 550.—Mexic.

berroviana, Tuss. ex Kunth, Enum. Pl. i. 250 = Cynodon Dactylon.

Berteroiana, Spreng. ex Steud. Nom. ed. II. 39, 143 = Arundinella martinicensis.

Biebersteiniana, Claus, Beitr. Russ. Reich. viii. 264 = trichoclada.

Billardieri, R. Br. Prod. 171 = Deyeuxia Billardieri.

biseta, Spreng. ex Steud. Nom. ed. II. i. 39, 143 = Arundinella indica.

Boecheleri, Seubert, ex Steud. Syn. Pl. Gram. 218 (= *Gymnopogon distichophyllus*).

borbonica, *Steud. l. c.* 170.—Ins. Borbon.

borealis, Hartm. Skand. Fl. ed. IV. 23 = canina.

Bourgaei, '*Pourn. Mex. Pl. Lu. Gram.* 96,' *ex Hemsl. Biol. Centr. Am. Bot.* iii. 550.—Mexic.

brachystachys, *Steud. Syn. Pl. Gram.* 422.—Reg. Magell.

brachyphylla, Steud. l. c. 175 = Polypogon brachyphyllus.

brachyphylla, Willd. ex Steud. Nom. ed. II. i. 39 = Sporobolus atrovirens.

brachystachys, *Schult. Mant.* iii. 569.—Sicilia.

brasiliensis, Spreng. Nov. Prov. 45 = Reimaria acuta.

brevifolia, *Nutt. Gen. Am.* i. 44.—Am. bor.

breviglumis, *F. Muell. Census*, 133.—Austral.

brevis, Knapp, Gram. Brit. 116 = alba?

bromidioides, *Griseb. in Gortt. Abh.* xxiv. (1879) 293. —Reg. Argent.

bromoides, Linn. Mant. i. 30 = Stipa Aristella.

bryoides, *Dum. Fl. Belg.* 152.—Belgium.

byzantina, *Boiss. Diagn.* Ser. I. xiii. 46.—Byzantium.

caerulescens, Poir. Encyc. Suppl. i. 258 (= *Urachne coerulescens*).

caerulescens, vide etiam coerulescens, DC.

caespitosa, Gaudich. in Freyc. Voy. Bot. 497 (= *Vilfa Gaudichaudii*).

AGROSTIS :—

caespitosa, Hoppe, ex Steud. Nom. ed. II. i. 39 = repestris.

caespitosa, Presl, Rel. Haenk. i. 237 = Presliі.

caespitosa, Torr. in Ann. Lyc. N. York, i. (1824) 152 = Torreyi.

caespitosa, Willd. ex Steud. Nom. ed. II. i. 39 = Muehlenbergia quadridentata.

caespitosa, Salisb. Prod. 25 = Deschampsia caespitosa.

Calamagrostis, Georgi, Beach. Russ. Reich. iii. iv. 687 = Calamagrostis lanceolata.

Calamagrostis, Linn. Syst. ed. X. 872 = Stipa Calamagrostis.

Calamagrostis, Suter, Fl. Helv. i. 36 = Calamagrostis epigeios.

calamagrostoides, Regel, Ind. Sem. Hort. Petrop.(1865) 38 = Calamagrostis agrostoides.

californica, Trin. Agrost. ii. 113 = exarata !

calycina, Willd. ex Steud. Nom. ed. II. i. 39 ; ii. 530 = Schismus marginatus.

campestris, *Phil. in Linnaea*, xxix. (1857–58) 87.—Chili.

campyla, *Tuckerm. in Am. Journ. Sc.* Ser. II. vi. (1848) 231.—Am. bor.

canariensis, *Parl. in Webb & Berth. Phyt. Canar.* iii 391.—Ins. Canar.

Candolleі, Parl. Fl. Palerm. i. 68 = alba.

canescens, *Griseb. in Goett. Abh.* xix. (1874) 253.—Reg. Argent.

canescens, Salisb Prod. 25 = Corynephorus canescens.

canina, *Linn. Sp. Pl. ed.* 2.—Amphig.

canina, Ucria, Hort. Reg. Panorm. 87 = pallida.

capensis, *Lam. Encyc.* i. 58.—Afr. austr.

capensis, Steud. in Flora, xii. (1829) 467 = Bergiana.

capensis, Willd. Sp. Pl. i. 372 = Sporobolus capensis.

capillaris, All. Fl. Pedem. ii. 283 = canina.

capillaris, Huds. Fl. Angl. ed. II. 27 = alba.

capillaris, Boiss. & Lange, ex Anderss. in Walp. Ann. vi. 981 = truncatula.

capillaris, Desf. Fl. Atlant. i. 69 = Reuterii.

capillaris, Bieb. Fl. Taur. Cauc. i. 55 = trichoclada.

capillaris, *Linn. Sp. Pl. ed. Huds. Fl. Angl.* 27.—Europ.

capillaris, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. i. 55 = trichoclada.

capillaris, Pourr. ex Nym. Consp. 801 = olivetorum.

capillaris, Thore, Chlor. 26 = elegans

capillaris, Vill. Hist Pl. Dauph. ii. 78 = vulgaris.

cernua, De Not. ex Parl. Fl. Ital. i. 189 = alpina.

Carmichaeli, *Roem. & Schult. Mant.* iii. 571.—Ins. Tristan d'Acugn.

caryophylla, Salisb. Prod. 25 = Aira caryophyllea.

castriana, *Boiss. & Reut. Diagn. Hisp.* 26.—Hispan ; Afr. bor.

chamaecalamus, *Trin. in Mém. Acad Pétersb.* Sér. VI. vi. 11. (1846) 365.—Peruv.

Chamissonis, *Trin. l. c.* 374.—Ins. Paschalis.

chilensis, *Kunze, ex Trin. l. c.* 389.—Chili.

Chinantlae, '*Pourn. Mex. Pl. Lu. Gram.* 96,' *ex Hemsl. Biol. Centr. Am. Bot.* iii. 550.—Mexic.

chinensis, Koen. ex Steud. Nom. ed. II. i. 39 = Leptochloa tenerrima.

chonotica, *Phil. in Linnaea*, xxix. (1857–58) 87.—Chili.

ciliaris, Steud. Syn. Pl. Gram. 116 = Arundinella anomala.

ciliata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1846) 373.—Ind. or.

ciliata, Pall. ex Ledeb. Fl. Ross. iv. 404 = Pappophorum boreale.

ciliata, Thunb. Fl. Jap. 49 = Festuca Thunbergii.

ciliata, Willd. ex Steud. Nom. ed. II. i. 39 = Muehlenbergia ciliata.

cimicina, Poir. Encyc. Suppl. i. 258 = Panicum cimicinum.

Cinna, Lam. ex Roem. & Schult. Syst. ii. 373 = Cinna arundinacea.

Cinna, Retz. Obs. Fasc. v. 18 = Muehlenbergia mexicana.

clandestina, Spreng. Fl. Hal. Mant. 32 = Sporobolus asper.

clausa, *Phil. in Anales Univ. Chil.* (1873) 562.—Chili.

clausa, *Steud. Syn. Pl. Gram.* 169.—Mesopot.

clavata, *Trin. in Spreng. Neue Entdeck.* ii. 55.—Kamtschat.

coarctata, Ehrh. ex Hoffm. Fl. Germ. ed. II. i. 87 = alba.

coerulescens, DC. Fl. Fr. Suppl. 250 = Piptatherum coerulescens.

coguata, *Steud. Syn. Pl. Gram.* 421.—Reg. Magell.

AGROSTIS :—

colorata, Willd. ex Steud. Nom. ed. II. i. 40 = Muehlenbergia elegans.

comosa, Poir. Encyc. Suppl. i. 257 (= *Urachne parviflora*).

complanata, Alt. Hort. Kew. i. 96 = Chloris petraea.

composita, Poir Encyc. Suppl. i. 254.—Carolina.

compressa, Willd. in Roem. & Ust. Mag. iv. (1790) xi. 39 = alba.

compressa, *Poir. Encyc. Suppl.* i. 258.—Carolina.

compressa, *Doell, in Mart. Fl. Bras* ii. iii 27.—Bras.

compressa, Torr. Cat. Pl. N. York, 91 = Sporobolus compressus.

concinna, Tuckerm. in Journ. Am. Sc. Ser. II. xlv. (1843) 43 = canina.

condensata, Willd. ex Steud. Nom. ed. II. i. 40 = verticillata.

conferta, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. 312 = nana.

conferta, Willd. ex Steud. Nom. ed. II. i. 40 = alba.

conspicua, Roem. & Schult. Syst. ii. 364 = Arundo conspicua.

contracta, F. Muell. ex Hook. f. Fl. Tasm. ii. 126 = Deyeuxia scabra.

Cornucopiae, *Sm. ex Fraser, Monog.* 2.—Am. bor.

Cornucopiae, Lam. ex Steud. Nom. ed. II. i. 40 = laxiflora.

coromandeliana, Roxb. Fl. Ind. ed. Wall. i. 318 (= *Vilfa Roxburghiana*).—Ind. or.

coromandelina, Retz. Obs. Fasc. iv. 19 = Sporobolus coromandelinus.

corralensis, *Phil. in Linnaea*, xxxiii. (1864–65) 286.—Chili.

cremenica, Bess. ex Steud. Nom. ed. II. i. 39, 40 = alba.

crinita, Moench, Meth. 178 = Polypogon monspeliensis.

crinita, *R. Br. Prod.* i. 170.—Austral.

crinita, Rich. ex. Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 4 = Dichelachne Forsteriana.

crucata, Linn. Pl. Jam. Pugill. ii. 7 ; Syst. ed. X. 873 = Chloris cruciata.

cryptandra, Torr in Ann. Lyc. N. York, i. (1824) 151 = Sporobolus cryptandrus.

cuspidata, Willd. ex Steud. Nom. ed. II. i. 40 = Calamagrostis tolucensis.

cylindrica, *R. Br. Prod.* i. 171.—Austral.

dalmatica, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 357.—Europ.

debilis, Spreng. Syst. i. 249 = Deyeuxia Forsteri.

debilis, Spreng. Syst. i. 262 = Muehlenbergia debilis.

decipiens, K. Br. Prod. i. 175 = Deyeuxia scabra.

decipiens, Salzm. ex Steud. Syn. Pl. Gram. 101 = Eriochloa polystachya.

decumbens, Host, Ic. Gram. iv. t. 54 = alba.

decumbens, *Link, Hort. Berol.* i. 80.—Mexic.

delicatula, *Pourr. ex Kunth, Enum. Pl.* i. 219.—Lusitan.

delicatula, *Steud in Lechl. Berb. Am. Austr.* 55.—Peruv.

Delislei, *Hemsl. in Bot. Challeng.* i. 11. 269.—Ins. Amsterdam.

denso, Bieb. Fl. Taur. Cauc. i. 56 = verticillata.

densa, *F. Muell. Census*, 138.—Australia.

Desvauxi, *Phil. in Linnaea*, xxxiii. (1864–65) 288.—Chili.

diandra, Hornem. in Syll. Ratisb. ii. (1824) 198 = *diandra*.

diandra, Retz. Obs. Fasc. v. 19 = Sporobolus diander.

dianthera, *Schult. Mant.* ii. 571.—Ind. or.

diaphora, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. 11. (1846) 368 = Deyeuxia quadriseta.

difficilis, *Hemsl. Bot. Challeng.* i. 11. 268. t. 48. —Ins. Amsterdam.

diffusa, Banks & Soland. ex Hook. f. Fl. New Zeal. i. 298 = Deyeuxia Billardieri.

diffusa, Bess. Prim. Fl. Galic. i. 68 = verticillata.

diffusa, Host, Ic. Gram. iv. t. 55 = alba.

diffusa, Muhl. Desc. Gram. 64 = Muehlenbergia sylvatica.

digitata, Lam. Encyc. i. 59 = Paspalum digitatum.

dispar, Hort. Par. ex Steud. Nom. ed. II. i. 40 (= *Urachne parviflora*).

dispar, Michx. Fl. Am. Bor. i. 52 = stolata ?

distans, Kunze, Del. Sem. Hort. Lips. (1847) = Deyeuxia scabra.

distichа, Schweig. ex Steud. Nom. ed. II. i. 39, 40 = alba.

distichophylla, *Phil. Fl. Atac.* 54.—Chili.

distichophylla, *Roem. & Schult. Syst. ii.* 364.—Austral.

AGROSTIS :—
divaricata, *Griseb. in Goett. Nachr.* (1868) 81.—Ind. or.
divaricata, Hoffm. Fl. Germ. ed. II. i. 37 = vulgaris.
divaricata, Hort. ex Steud. Nom. ed. II. i. 40 = Chaeturus fasciculatus.
disparicata, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 713 = alba.
domingensis, Schult. Mant. iii. 570 = Sporobolus domingensis.
Dregeana, *Steud. Syn. Pl. Gram.* 173.—Afr. austr.
dubia, DC. Fl. Fr. iii. 21 = vulgaris.
dubia, Leers, Fl. Herborn. 21 = canina.
dubia, Auct. ex Steud. Nom. ed. II. i. 40 = Chaeturus fasciculatus.
dulcis, Hort. Par. ex Poir. Encyc. Suppl i. 250 = alba.
dulcis, Sibth. ex Kunth, Enum. Pl. i. 218 = verticillata.
dura, Willd. ex Steud. Nom. ed. II. i. 40 = Sporobolus virginicus!
Duriaei, Boiss. & Reut. ex Nym. Consp. 802 = truncatula.
Ecklonis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 364.—Afr. austr.
effusa, Lam. Encyc. i. 59 = Milium effusum.
effusa, *Spreng. Syst.* i. 260.—Europ.
elata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 317.—Am. bor.
elatior, *Steud. Syn. Pl. Gram* 174.—N. Zel.
elegans, Salisb. Prod. 25 = A. Cornucopiae.
elegans, Roem. & Schult. Syst. ii. 362 = Humboldtiana.
elegans, *Thore, ex Lossel. in Journ. Bot.* ii. (1849) 207. t. 5.—Europ.
Elliotiana, Schult. Mant. ii. 372 = arachnoides.
elongata, Lam. Illustr. i. 162 = Sporobolus indicus.
elongata, Roth, Nov. Sp. 41 = Sporobolus orientalis.
eminens, *Griseb. in Goett. Abh.* xix. (1874) 254.—Reg. Argent.
erecta, Spreng. Syst. i. 264 = Muehlenbergia aristata.
exarata, *Trin. Gram. Unisf.* 208; *Ic. Gram.* iii. t. 27.—Am. bor.
exasperata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 352.—Chili.
exigua, *Thurb. in S. Wats. Bot. Calif.* ii. 275.—Calif.
exilis, Loisel. in Mem. Soc. Linn. Par. vi. (1827) 399 = elegans.
expansa, Poir. ex Steud. Nom. ed. II. i. 40 = Cinna arundinacea.
falklandica, *Hook. f. Fl. Antarct.* 373.—Ins. Falkland.
fasciculoris, Sincl. Hort. Gram. Wob. 154 = canina.
fasciculata, *Roem. & Schult. Syst* ii. 362.—Ecuador.
fertilis, Steud. Syn. Pl. Gram. 170 = Sporobolus elongatus.
festucoides, Muhl. ex Kunth, Enum. Pl. i. 200 = Muehlenbergia glomerata.
festucoides, Vill. Prosp. 16 = alpina.
filifolia, *Link, in Schrad. Journ.* ii. (1799) 313.—Lusitan.
filiformis, Koen. ex Kunth, Enum. Pl. i. 261 = Cynodon elongatus.
filiformis, Spreng. Fl. Hal. Mant. 32 = Deyeuxia Forsteri!
filiformis, Vill. Hist. Pl. Dauph. ii. 78 = alpina.
filiformis, Willd. Enum. Hort. Berol. 95 = Muehlenbergia mexicana.
fimbellata, *Salzm. ex Steud. Syn. Pl. Gram.* 53 = Setaria sulcata!
flava, O. F. Muell. Fl. Dan. t. 751 = alba.
flaccescens, Host, Ic. Gram. iv. 52 = alpina.
flavida, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 48 = alba.
flavidula, *Steud. Syn. Pl. Gram.* 421.—Reg. Magellan.
foliata, *Hook. f. Fl. Antarct.* 95.—N. Granata.
foliosa, Hort. ex Kunth, Enum. Pl. i. 207 = Muehlenbergia mexicana.
foliosa, *Vasey, in Bull. Torrey Bot. Club*, x. (1883) 68.—Am. bor.
Fonckii, *Phil. in Linnaea*, xxxiii. (1864–65) 986.—Chili.
Forsteri, Rich. ex Roem. & Schult. Syst. ii. 359 = Deyeuxia Forsteri.
frigida, F. Muell. ex Benth. Fl. Austral. vii. 583 = Deyeuxia frigida.
frondosa, *Guss. Fl. Sic. Prod.* i. 58.—Sicilia.
frondosa, Poir. Encyc. Suppl. i. 253 = Muehlenbergia mexicana.
frondosa, Tenore, ex Spreng. Pugill. ii. 5 = vulgaris.
fulva, *Griseb. in Goett. Abh.* xxiv. (1879) 294.—Reg. Argent.
gaditana, *Nym. Syll.* 408.—Lusitan.
gallaeica, Hort. ex Steud. Nom. ed. I. 20; ed. II. i. 40 = Alopecia agrostiden.

AGROSTIS :—
Gayana, *E. Desv. in C. Gay, Fl. Chil.* vi. 313.—Chili.
gelida, F. Muell. in Trans. Vict. Inst. (1855) 43 = Muelleri.
gelida, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 348.—Peruv.
geminata, *Trin. Gram. Unisf.* 207.—Usalasch.
geniculata, Lam. Fl. Fr. iii. 572 = canina.
Ghiesbreghtii, *Fourn. Mex. Pl. En. Gram.* 97, *ex Hemsl. Biol. Centr. Am.* iii. 551.—Mexic.
gigantea, Roth, Tent. Fl. Germ. i. 81 = alba.
glabra, *Kunth, Enum. Pl.* i. 226.—Chili.
glabra, Presl, ex Trin. in Mém. Acad. Pétersb. Sér. VI. vi. 11. (1845) 342 = stricta.
glabrata, Spreng. Syst. i. 262 = Muehlenbergia glabrata.
glauca, Muhl. Descr. Gram. 76 = Deyeuxia Nuttaliana.
glauca, *Steud. Syn. Pl. Gram.* 176.—Mexic.
glaucescens, Opiz, ex Steud. Nom. ed. II. i. 40 = parvula.
glaucescens, Spreng. Syst. i. 258 = sicula.
glaucinea, Bast. Fl. Maine-et-Loire, Suppl. 25 = vulgaris.
globosa, Poir. Encyc. Suppl. i. 257 = Milium globosum.
glomerata, *Kunth, Enum. Pl.* i. 219.—Peruv.
glutinosa, Fisch. ex Kunth, l. c. 198 = Melinis minutiflora.
gracilis, Kit. ex Schult. Fl. Austr. ed. II. i. 180; et in Linnaea, xxxii. (1863) 300 = alba.
gracilis, Willd. ex Steud. Nom. ed. II. i. 40 = Muehlenbergia gracilis.
gracilis, Salisb. Prod. 25 = Koeleria cristata.
graeca, Sibth. Fl. Graec. t. 66 (= *Urachne parviflora*).
grandis, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. 11. (1845) 316 = exarata.
graveolentica, *Steud. Syn. Pl. Gram.* 175.—Greenland.
Gunniana, *F. Muell. Census*, 183.—Austral.
gynostyla, *Steud. Syn. Pl. Gram* 170.—Afr. austr.
Hegetschweileri × *Bruggeana*, *in Jahresb. Naturf. Ges. Graub.* ii. xxv. (1880) 60.—Helvet.
Helenae, *Steud. Syn. Pl. Gram.* 171.—Ins. S. Helena.
herbacea, Steud. l. c. 224 = Aira paradoxa.
Hildebrandii, Thurb. ex Boland. in Trans. Calif. Agr. Soc. (1884) 186 = aequivalvis.
hirsuta, *Linn. f. Suppl.* 108.—Teneriff.
hirtella, *Hochst. ex Steud. Syn. Pl. Gram.* 178.—Abyssin.
hispanica, Boiss. & Reut. Pugill. 120 = castellana.
hispida, Brot. ex Nym. Consp. 802 = truncatula.
hispida, Willd. Sp. Pl. i. 370 = vulgaris.
holciformis, Bleb. Fl. Taur. Cauc. i. 54 = Oryzopsis holciformis.
Hornungiana, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 46 = vulgaris.
Humboldtiana, *Steud. Nom.* ed. II. i. 40 — Quito.
humifusa, Roem. & Schult. Syst. ii. 361 = Sporobolus humifusus.
humilis, *Vasey, in Bull. Torr. Club*, x. (1883) 21.—Am. bor.
hybrida, Gand. Agrost. i. 66 = canina.
hygrometrica, Nees, in Mart. Fl. Bras. ii. 404 (= *Brachydium hygrometricum*).
hyperborea, Laest. ex Nym. Consp. 802 = rubra.
implicata, Spreng. Syst. i. 262 = Muehlenbergia implicata.
inaequiglumis, *Griseb. in Goett. Nachr.* (1868) 80.—Ind. or.
inconspicua, *Kunze, ex C. Gay, Fl. Chil.* vi. 315.—Chili.
incurvata, Scop. Fl. Carn. ed. II. i. 62 = Lepturus incurvus.
indica, Forsk. Fl. Aegypt. Arab. 104. n. 66 = Sporobolus coromandelinus.
indica, Linn. Sp. Pl. 63 = Sporobolus indicus.
infirma, *Steud. in Mg. Pl. Yungh.* 342.—Java.
insignita, Schur, Enum. Fl. Transs. 732 = alba.
intermedia, Balb. Elench. 35 = vulgaris.
interrupta, Linn. Syst. ed. x. 872 = Apera interrupta.
intricata, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 413 = scabra.
involuta, Muhl. Descr. Gram. 73 = Sporobolus asper.
involuta, *Petit. Encyc. Suppl.* i. 252.—Marocc.
jamaicensis, Poir. l. c. 255 = Digitaria filiformis.
Jamesoniana, *Steud. Syn. Pl. Gram.* 168 = Am. merid.
japonica, *Steud. l. c.* 171.—Japon.
javanica, Berm. ex Kunth, Enum. Pl. i. 213 = Ischaemum muticum.
juncea, Lam. Encyc. i. 60 = Sporobolus tremulus.
juncea, Michx. Fl. Am. Bor. i. 62 = Sporobolus junceus.

AGROSTIS :—
junciformis, Willd. ex Steud. Nom. ed. II. i. 41 = Deyeuxia juncifornis.
Jurzeí, *Link, in Schrad. Journ.* ii. (1799) 312.—Lusitan.; Hispan.
Kitaibelii, Schult. Fl. Austr. ed. II. i. 178 = alba.
koelerioides, *E. Desv. in C. Gay, Fl. Chil.* t. vi. 317.—Chili.
Labillardieri, Roem. & Schult. Syst. ii. 360 = Deyeuxia Billardieri.
lachnantha, *Nees, in Ind. Sem. Hort. Bot. Vratisl.* (1834).—Afr. austr.
Laersii, J. F. Gmel. Syst. ii. 169 = canina.
lanata, H. B. & K. Nov. Gen. et Sp. i. 186 = Epicampes lanata.
lanceolata, Roem. & Schult. Syst. ii. 364 = Calamagrostis lanceolata.
Langei, Nym. Consp. 801 = vulgaris.
lanigera, *Desv. Opusc.* 61.—Bras.
lasianiha, *Phil. in Linnaea*, xxix. (1857–58) 37.—Chili.
laterifl ra, Michx. Fl. Bor. Am. i. 53 = Muehlenbergia mexicana.
latifolia, Trevir. in Goeppert, Beschr. Bot. Gaert. in Breslau, 93 = Cinna arundinacea.
laxa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 147 = vulgaris.
laxa, Schreb. ex Pursh, Fl. Am. Sept. i. 61 = scabra.
laxiflora, *Poir. Encyc. Suppl.* i. 256.—Am. bor.
laxiflora, Richards. Frankl. Voy. App. 3 = scabra.
lazica, *Balansa, in Bull. Soc. Bot. Fr.* xxi. (1874) 12.—Orient.
Lechleri, *Steud. Syn. Pl. Gram.* 493.—Chili.
Leersii, J. F. Gmel. Syst. 1495 = canina.
lendigera, Poir. Encycl. Suppl. i. 259 = Gastridium australe.
lenta, Att. Hort. Kew. i. 96 = Panicum filiforme.
leptos, *Steud. Syn. Pl. Gram.* 169.—Louisiana.
leptostachys, *Hook. f. Fl. Antarct.* 94.—Reg. Antarct.
leptotricha, *E. Desv. in C. Gay, Fl. Chil.* vi. 316.—Chili.
Lessoniana, Steud. Nom. ed. II. i. 41 = Arundo conspicua.
ligulata, *Steud. Syn. Pl. Gram.* 178.—Austral.
limonias, *Phil. in Linnaea*, xxxiii. (1864–65) 287.—Chili.
limosa, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 82; et Oestr. Bot. Zeitschr. ix. (1859) 48 = alba.
Lu deniana, *Steud. Syn. Pl. Gram.* 171.—Colombia.
linearis, Retz. Obs. Fasc. iv. 19 = Cynodon Dactylon.
lithuanica, Bess. ex Roem. & Schult. Mant. 568, in add. = vulgaris.
littigua, *Steud. Syn. Pl. Gram.* 162.—Lusitan.
littoralis, Lam. Illustr. 161 = Sporobolus virginicus.
littoralis, With. Arr. Brit. Pl. ed. III. ii. 129. t. 28 = Polypogon littoralis.
lobata, R. Br. Prod. i. 171 = Deyeuxia quadriseta.
longiaristata, Herb. Ross. ex Kunth, Enum. i. 178 = Stipa splendens.
longiflora, Willd. ex Steud. Nom. ed. II. i. 41 = Cinna arundinacea.
longifolia, *Torr. ex A. Gray, Man. Bot. N. U. St.* ed. I. 575 = Sporobolus asper.
longiseta, Hort. ex Steud. Nom. ed. I. 21, 645 = Chaeturus fasciculatus.
lusitanica, Steud. Syn. Pl. Gram. 169 = castellana.
lutescens, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 299 = Arundo conspicua.
lutosa, Poir. Encycl. Suppl. i. 240 = Polypogon littoralis.
Lyallii, *Hook. f. Fl. N. Zeal.* 307.—N. Zel.
magellanica, *Lam. Illustr.* i. 160.—Reg. Magellan.
malaccensis, Willd. ex Kunth, Enum. Pl. i. 118 = Isachne pulchella.
maritima, Lam. Encyc. i. 61; Illustr. i. 161 = alba.
maritima, Wuth. Arr. Brit. Pl. ed. III. ii. 132 = Elymus arenarius!
Mattrella, Linn. Mant. iii. 185 = Zoysia pungens.
maxima, Roxb. Fl. Ind. i. 319 = Thyssanolaena Agrostis.
media, *Carmich. in Trans. Linn. Soc.* xii. (1819) 504. —Ins. Trist. da Cunha.
melanosperma, Lam. Encyc. i. 58 (= *Urachne Linnaei*).
melanteca, Trin. in Mém. Acad. St. Pétersb. ii. (1832) 170 = setacea.
Mertensii, *Trin. in Linnaea*, x. (1835) 302 = Chili; Unalaschk.
mexicana, *Kunth, Enum. Pl.* i. 233.—Mexic.
mexicana, Linn. Mant. 31 = Muehlenbergia mexicana.

AGROSTIS :—

mexicana, Pers. Syn. i. 74 = Deyeuxia canadensis.
Meyeni, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1846) 212.—Chili.
Michaelis, *Stud. Syn. Pl. Gram.* 171.—Ins. Azor.
Michauxii, Trin. Gram. Unif. 206 = scabra.
Michauxii, Zoograp. in Koem. Collect. 128 = Chaeturus fasciculatus.
micrantha, *Stend. Syn. Pl. Gram.* 170.—Nepal.
microcarpa, Hort. ex Steud. Nom ed. II. i. 164; ii 41 = Muehlenbergia debilis.
microphylla, Stend. Syn Pl. Gram. 164 = exarata.
microsperma, Lag. Hench. 2 = Muehlenbergia debilis.
militacea, Linn. Sp. Pl. 61 = Piptatherum miliaceum.
Milium comosum, Poir. Encyc. Suppl. i. 257 = Piptatherum miliaceum.
minima, Linn. Sp. Pl. 63 = Mibora verna.
minima, *Lour. Fl. Cochinch.* 50.—Cochinch.
minor, F. Muell. ex Benth. Fl. Austral. vii. 582 = Deyeuxia minor.
minutiflora, Desf. ex Steud. Nom. ed. II. i. 41 = Sporobolus minutiflorus.
minutiflora, Steud. Syn. Pl. Gram. 170 (= *Vilfa indica*).
minutissima, Steud. l. c. 171 = Sporobolus ramiculosus.
monandra, Homem. Hort. Hafn. i. 73 (= *Urachne parviflora*).
monostachya, Poir. Encyc. Suppl. i. 256 = Panicum amplexicaule.
montana, Krock. Fl. Siles. i. 110 = alpina.
montana, R. Br. Prod. i. 171 = Deyeuxia montana.
montevidensis, *Spreng. ex Nees, Agrost. Bras.* 408 = Montevid.
montevid., Roem. & Schult. Syst. ii. 359 = Deyeuxia montana.
Montis-aurei, Delarbre, Fl Averg. 8, ex Poir. Encyc. Suppl. i. 246 = Piptatherum multiflorum.
mucronata, *Presl, Rel. Haenk.* i. 388.—Am. bor. occ.
mucronata, Spreng. Syst. i. 262 = Muehlenbergia mucronata.
Muelleri, *Benth. Fl. Austral.* vii. 576.—Austral.
Muelleri, Presl, Bot. Bemerk. 120 = verticillata.
multiculis, *Hook. f Fl. Antarct.* 95.—Ins. Campbell.
multiculis, Kit. in Linnaea, xxxii. (1863) 306 = alba.
Munroana, Aitch. & Hemsl. Journ. Linn. Soc. xix. (1881) 192 = Calamagrostis Munroana.
muricata, *Kunth, Enum. Pl.* i. 213.—Peruv.
muscosa, *T. Kirk, in Trans. N. Z. Inst.* xiii. (1881) 335.—N. Zel.
Mustaphae, *Steud. Syn. Pl. Gram.* 169.—Afr. bor.
mutabilis, Knapp, Gram. Brit. 28 = alba.
mutabilis, Sibth. Fl. Oxon. 36 = setacea.
Myuros, Lam. ex Besuv. Agrost. 40 = Panicum Myurus.
nana, Delarbre, Fl. Averg. ed. II. 664 = pumila.
nana, *Kunth, Enum. Pl.* i. 266.—Peruv.
nardifolia, *Griseb. in Goett. Abh.* xix. (1874) 252.—Reg. Argent.
nebulosa, *Boiss. & Reut. Diagn.* 96.—Hispan.
nebulosa, Bourg. ex Reut. & Lange, in Bull. Soc. Bot. Fr. xi. (1864) 47 = delicatula.
Nessii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 361.—Afr. austr.
neglecta, Schult. Fl. Austr. ed. II. i. 179 = alba.
nemoralis, *Phil. in Linnaea*, xxx. (1859–60) 205.—Chili.
neogaea, *Steud. Syn. Pl. Gram.* 171.—N. Zembl.
nervosa, *Nees, ex Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 338.—Ind. or.
nevadensis, *Boiss. Elench.* 87.—Hispan.
nigra, With. Arr. Brit. Pl. ed. III. ii. 181 = vulgaris.
nigricans, *Salem. ex Steud. Nom.* ed. II. i. 41.—Bahia.
nigricans, Poir. Encyc. Suppl. i. 259 = Milium nigricans.
nitens, Guss. Fl. Sic. Prod. i. 50 = Triplachne nitens.
nivalis, F. Muell. in Trans. Vict. Inst. (1855) 43 = Deyeuxia nivalis.
nivea, Sincl. Hort. Gram. Woburn. 156 = alba ?
nobilis, De Not. ex Parl. Fl. Ital. i. 126 = canina.
nootkaensis, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. ii. (1845) 326 = nutkaensis.
Novae-Hollandiae, Beauv. Agrost. 148 = Deyeuxia Forsteri.
noveboracensis, Spreng. Syst. i. 260 = Epicampes lanata.
nutans, Poir. Encyc. Suppl. i. 255 = Ichnanthus pallens.
nutans, Spreng. Syst. i. 263 = Muehlenbergia gracilis.
nutkaensis, *Kunth, Enum. Pl.* i. 222.—Am. bor. occ.
oblique, Auct. ex Trin. Gram. Unifl. 196 = alba.

AGROSTIS :—

oblique, *Kit. in Linnaea*, xxxii. (1863) 306.—Austria.
obtusata, Auct. ex Steud. Nom. ed. II. i. 41 = Estonia obtusata.
odorata, *Lour. Cochinch.* 64.—Cochinch.
oligantha, Koem. & Schult. Syst. ii. 372 = pauciflora.
oliveriensis, *Gren. & Godr. Fl. Fr.* iii. 488.—Gallia.
orcophila, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 323.—Am. bor.
orientalis, Nees, Agrost. Bras. 393 = Sporobolus tenacissimus.
ovrorunis, Scenus, Reise Ist. (1805) 50 (= *Urachne parviflora*).
ovata, Forst. f. Prod. 8 = Echinopogon ovatus.
ovarensis, Roem & Schult Syst. ii. Mant. 199 = Sporobolus pyramidalis
pallens, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 328.—Am. bor.
pallida, *DC. Fl. Fr. Suppl.* 251.—Europ.
pallida, Schk. Handb. 3; Nachtr. 1 = canina.
pallida, With. Arr. Brit. Pl. ed. III. ii. 198. t. 22 = alba.
palustris, Huds. Fl. Angl. 37 = alba.
panicea, Lam. Encyc. i. 56; Illustr. 811 = Gastridium australe.
panicea, Ait. Hort. Kew. i. 94 = Polypogon monspeliensis.
paniculata, Lam. Illustr. 162 = Panicum Lamarckii.
paniculata, Hort. Madrit. ex Steud. Nom. ed. II. i. 41 = Muehlenbergiae sp. sec. Trin.
papposa, Schk. Handb. ii. 534 = Stipa Calamagrostis.
paradisiaca, Steud. Syn. Pl. Gram. 168 = chilensis.
paradoxa, Linn. Sp. Pl. 63 (= *Urachne Linnaei*).
parasitica, R. Br. in Ross, Voy. ed II. ii. 192 = Colpodium latifolium.
parviflora, R. Br. Prod. 170 = scabra.
parvula, Schult. Fl. Austr. ed. II. i. 189 = Sporobolus minutus.
patagonica, *Phil. in Linnaea*, xxix. (1857–58) 87.—Patagon.
patens, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 322.—Chili.
patula, Gaud. Agrost. Helv. i. 80 = alba.
pauciflora, *Pursh, Fl. Am. Sept.* i. 63.—Am. bor.
pauciflora, Schrad. Fl. Germ. 3 = alba.
paucinodis, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 242. —Reg. Magell.
pennata, *Tuckerm. in Am. Journ. Sc.* Ser. II. xlv. (1843) 44.—Am. bor.
pernambucensis, Spreng. Syst. i. 258 = Panicum rivulare.
peruviana, Spreng. Syst. i. 262 = Muehlenbergia Clomena.
phleoides, *E. Desv. in C. Gay, Fl. Chil.* vi. 320.—Chili.
phleoides, Willd. ex Steud. Nom. ed. II. i. 41 = Calamagrostis coarctata.
Pickeringii, *Tuckerm. in Am. Journ. Sc.* Ser. II. xlv. (1868) 3.—Am. bor.
pilosa, Schleich. ex Gand. Agrost. Helv. i. 75 = Calamagrostis tenella.
pilosa, *Less. in Rich. Voy. Astrol.* 134. t. 28.—N. Zel.
pilosa, Retz. Obs. Fasc. vi. 22 = Cynodon pilosus.
pilosula, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 372.—Ind. or.
pinifolia, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 255 = alba.
planifolia, *C. Koch, in Linnaea*, xxi. (1848) 380.—Armen.
plebeia, R. Br. Prod. i. 173 = Deyeuxia plebeia.
plicata, *Lour. Fl. Cochinch.* 64.—China.
plumosa, Tenore, Prod. i. 59 = Panicum Teneriffae.
poaeformis, Willd. ex Steud. Nom. ed. II. i. 42 = Poa subcaniflora.
polymorpha, Huds. Fl. Angl. ed. II. i. 31 = alba, vulgaris.
Polypogon, Salem. ex Steud. Syn. Pl. Gram. 178 = Muehlenbergia brasiliensis.
polystachya, Boxe, ex Steud. Nom. ed. II. i. 43 = canina.
Pourretii, Spreng. Syst. i. 260 = Avena Pourretii.
Pourretii, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 290 = effusa.
praecox, Salisb. Prod. 24 = Aira praecox.
Preslii, *Kunth, Enum. Pl.* i. 235.—Peruv.
procera, A. Rich. Voy. Astrol. 125 = Arundo conspicua.
procera, *Retz. Obs.* iv. 19.—Malabar.
procumbens, Roem. & Schult. Syst. ii. 371 = alba.

AGROSTIS :—

provincialis, G. Mey. Korresp. Bl. Naturf. Ver. Riga, xxvi. 56 (quid ?).—Romia.
prostrata, *Hook. f. Fl. Antarct.* 378.—Ins. Falkland.
pseudo-arundinacea, Schleich. ex Gaud. Agrost. Helv. i. 92 = Calamagrostis varia.
pubens, Willd. ex Steud. Nom. ed. II. i. 42 = Epicampes lanata.
pubescens, S. F. Gray, Nat. Arr. Brit. Pl. ii. 148 = alba.
pubescens, H. B. & K. Nov. Gen. et Sp. i. 136 = Epicampes pubescens.
pulchella, *Guss. Fl. Sic. Prod.* i. 58.—Sicilia.
pulchella, Knuth, Gram. i. 70 = Humboldtiana.
pulchella, Loisel. ex Trin. Gram. Unifl. 206 = elegans.
pulchella, Roth, ex Roem. & Schult. Syst ii. 367 = Sporobolus pulchellus.
pumila, Linn. Mant. 31 = vulgaris.
pumila, *Steud. Syn. Pl. Gram.* 171.—Ins. Adsocns.
punctata, Lam. Encyc. i. 58. n. 17 = Eriochloa polystachya.
pungens, Muhl. Desc. Gram. 72 = Zoysia pungens.
pungens, Pall. 'Ind. Taur.' ex Georgi, Beschr. iii. iv. 688 = Aeluropus littoralis.
pungens, Schreb. Graes. ii. 46 = Sporobolus pungens.
pungens, Pursh, Fl. Am. Sept. 64 = Sporobolus virginicus.
purpurascens, *Berl. ex Steud. Nom.* ed. II. i. 42.—S. Doming.
purpurascens, Sw. Prod. Veg. Ind Occ. 25 = Sporobolus purpurascens.
purpurea, Gaud. Agrost. Helv. i. 70 = Apera Spica-venti.
pusilla, Dum. Obs. Gram. Belg. 139 = canina.
pyramidalis, *Rich. ex Steud. Syn. Pl. Gram.* 158 (= *Vilfa Richardi*).—Ind. occ.
pyramidata, Lam. Illustr. i. 161 = Sporobolus africus.
pyrenaea, Timb. in Mem. Acad. Toul. Sér. IV. vi. (1856) 97 = alpina.
pyrenaica, Pourr. l. c. iii. (1788) 806 = repestris.
quadridentata, Spreng. Syst. i. 263 = Muehlenbergia quadridentata.
quadrifida, Labill. Pl. Nov. Holl. i. 20 = Pentapogon Billardieri.
quadrinata, R. Br. Prod. i. 171 = Deyeuxia quadriseta.
quitensis, Spreng. Syst. i. 262 = Muehlenbergia elegans.
quitensis, Willd. ex Steud. Nom. ed. II. i. 42 = Muehlenbergia calamagrostides.
racemosa, Michx. Fl. Bor. Am. i. 53 = Muehlenbergia glomerata.
radiata, Lam. Jl. Jam. Pugill. 7; Syst. ed. X. 878 = Chloris radiata.
ramosa, Poir. Encyc. Suppl. i. 257 = Eriochloa annulata.
ramosa, Roem. & Schult. Syst. ii. 364 = Calamagrostis lanceolata.
ramulosa, Carm. in Trans. Linn. Soc. xii. (1818) 504 = Carmichaeli.
ramulosa, Roem. & Schult. Syst. ii. 361 = Sporobolus ramulosus.
rara, R. Br. Prod. i. 171 = Dichelachne sciures.
rara, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 3 = Dichelachne sciurea.
Ravennae, Beauv. Agrost. 148 = kriathus Ravennae.
refracta, Moench, Meth. Suppl. 60 = verticillata.
Reinwardtii, *Miq. Fl. Ind. Bat.* iii. 750.—Malays.
repens, Sincl. Hort. Gram. Woburn. 148 = alba.
retroflracta, Willd. Enum. Hort. Berol. 94 = Deyeuxia Forsteri.
Reuterii, *Boiss. Voy. Espagne*, ii. 645.—Hispan.
Reygeri, [Honak.] Verz. Aller Gew. Teutschl. 204 = alpina ?
rigescens, *Presl, in Rel. Haenk.* i. 337.—Peruv.
rigida, Lamon, ex A. Rich. Voy. Astrol. 124 = Dichelachne rigida.
rigida, Spreng. Syst. i. 262 = Muehlenbergia rigida.
rigida, Willd. ex Steud. Nom. ed. II. i. 42 = Calamagrostis recta.
rigidula, *Steud. Syn. Pl. Gram.* 171.—Java.
rigidifolia, Poir. Encyc. Suppl. i. 257 = Panicum rigidifolium.
rivularis, Brot. Fl. Lusit. i. 76 = verticillata.
rosea, *Griseb. in Goett. Abh.* xix. (1874) 253.—Reg. Argent.
Roylana, *Trin. Agrost.* ii. 126.—Ind. or.
rubicunda, Boxe, ex Desv. Opusc. 55 = Cinna arundinacea.
rubra, Auct. ex Steud. Nom. ed. II. i. 42 = canina.

AGROSTIS :—
rubra, Hnds. Fl. Angl. 26 = Gastridium australe.
rubra, *Linn. Sp. Pl.* 92.—Europ.
rubra, Wahlenb. ex Nym. Consp. 861 = Deyeuxia scabra.
rudis, Roem. & Schult. Syst. ii. 360 = Deyeuxia scabra.
rupestris, *All. Fl. Pedem.* ii. 237.—Europ.; Am. bor.
rupestris, Willd. ex Kunth, Gram. i. 70 = alpina.
Ruprechtii, *Boiss. Fl. Orient.* v. 517.—Caucas.
salina, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 366 = alba.
salmantica, *Kunth, Gram.* i. 71.—Hispan.
scabertima, *Hassk. Pl. Jav. Rar.* 26.—Java.
scabra, R. Br. Prod. 172 = Deyeuxia scabra.
scabra, *Tuckerm. in Am. Journ. Sc.* Ser. II. xlv. (1845) 44.—Am. bor.
scabra, *Willd. Sp. Pl.* i. 370.—Am. bor.
scabriglumis, Boiss. & Reut. Pugill. 126 = alba.
scabriuscula, Bockl. in Proc. As. Sc. Philad. '1862.' (1863) 90 = scabra.
Schaffneri, *Fourn. Mex. Pl. En. Gram.* 94, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 551.—Mexic.
Schiedeana, *Fourn. l. c.* 96, *ex Hemsl. l. c.*—Mexic.
Schiedeana, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1846) 327.—Mexic.
Schimperiana, *Hochst. ex Steud. Syn. Pl. Gram.* 170.—Abyss.
Schlechteri, Jord. & Verlot, in F. Schultz, Arch. Fl. Fr. et Allem. 389, 347 = alpina.
Schottii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1846) 356.—Gibraltar.
Schultesii, *Kunth, Enum. Pl.* i. 228.—Pannonia.
Schweinitzii, Trin. in Mém. Acad. Pétersb. Sér. VI vi. (1846) 311 = elata.
scirpea, R. Br. Prod. i. 171 = Deyeuxia sclurea.
scoparia, Koen. ex Steud. Syn. Pl. Gram. 115 = Panicum semialatum.
Scouleri, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1846) 329.—Am. bor. occ.
semibarbata, Trin. l. c. 378 = Deyeuxia Forsteri.
sepium, Jacq. Enum. Stirp. Vindob. 12 (= *Urachne parviflora*).
serica, Banks & Soland. ex Hook. f. Fl. Nov. Zel. i. 299 = Arundo conspicua.
sericea, Ell. Sketch, i. 135 = Muehlenbergia capillaris.
serotina, Linn. Mant. 30 = Diplachne serotina.
serotina, Torr. Fl. Un. St. i. 88 = Sporobolus serotinus.
sesquiflora, *B. Desv. in C. Gay, Fl. Chil.* vi. 318.—Chili.
setacea, *Curt. Fl. Lond.* vi. t. 12.—Europ.
setacea, Vill. Hist. Pl. Dauph. ii. 76 = rupestris.
setifolia, Brot. Fl. Lusit. i. 74 = setacea.
setifolia, Willd. ex Steud. Nom. ed. II. i. 42 = tenuifolia.
setifolia, *Hook. f. Handb. New Zeal. Fl.* 329.—N. Zel.
setifolia, *Fourn. Mex. Pl. En. Gram.* 97 ; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 551.—Mexic.
setosa, Mubl. Descr. Gram. 68 = Muehlenbergia glomerata.
setosa, Spreng. Syst. i. 262 = Muehlenbergia debilis.
Shkuhrpfii, Hort. ex Steud. Nom. ed. II. i. 42 = verticillata.
sicula, Guss. ex Steud. Nom. ed. II. i. 42 = Gastridium muticum.
sicula, *Kunth, Rev. Gram.* i. 71.—Sicilia.
signata, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 47 = alba.
simensis, *Hochst. ex Steud. Syn. Pl. Gram.* 178.—Abyss.
simulans, *Hemsl. in Hook. Ic. Pl.* t. 1455.—Ins. Sta. Helena.
sinaica, Boiss. Diagn. Ser. I. xiii. 46 = alba.
sobolifera, Muhl. ex Willd. Enum. Hort. Berol. 95 = Muehlenbergia sobolifera.
Solandri, F. Muell. Veg. Chath. Isl. 80 = Deyeuxia Forsteri.
spicaeformis, Linn. f. Suppl. 108 = Perotis latifolia.
spicata, Thunb. Prod. Pl. Cap. 19 = Sporobolus capensis.
spicata, Vahl, Symb. i. 9 = Sporobolus spicatus.
Spica-venti, Linn. Sp. Pl. ed. I. 61 = Apera Spica-venti.
Sporobolus, Spreng. Nov. Prov. 46 = Sporobolus Spenglerii.
spuria, Roem. & Schult. Syst. ii. 846 = Calamagrostis lanceolata.
stellata, Willd. Sp. Pl. i. 376 = Cynodon Dactylon.
stipata, Koel. Gram. 77 = Stipa Calamagrostis.
stipoides, Spreng. Syst. i. 263 = Muehlenbergia stipoides.
stolonifera, Host, Gram. iv. t. 56 = alba.

AGROSTIS :—
stolonifera, Lecrs, Fl. Herb. t. 4. f. 6 = vulgaris.
stolonifera, *Linn. Sp. Pl.* 62.—Europ.
stolonifera, Linn. ex Sm. Fl. Brit. i. 80 = verticillata.
stoloniasus, Bess. in Roem. & Schult. Mant. 567, in add. = alba.
straminea, Hartm. Gram. Scand. 4 = alba.
stricta, *Buese, in Miq. Pl. Jungh.* 341.—Java.
stricta, *De Not. ex Parl. Fl. Ital.* i. 187.—Italia.
stricta, J. F. Gmel. Syst. 170 = alba.
stricta, Sincl. Hort. Gram. Woburn. 150 = canina.
stricta, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1846) 342.—Chili.
stricta, *Willd. Sp. Pl.* i. 366.—Am. bor.
suaveolens, Blytt, in Mag. for Naturvidensk. (1837); sec. Sommerfelt, in Skand. Akad. Handl. (1837) 25 = Coss. a. grandiacea.
subaristata, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 192.—Afghan.
subulata, *Hook f. Fl. Antarct.* 95 —Ins. Campbell.
sylvatica, Georgi, Beschr. Russ. Reich. iii. 687 = Calamagrostis sylvatica.
sylvatica, Hnds. Fl. Angl. 28 = alba.
sylvatica, Koel. Gram. 92 = vulgaris.
sylvatica, Torr. Fl. N. Amer. i. 87 = Muehlenbergia Willdenowii.
tabularis, Steud. Nom. ed. II. i. 42 = Bergiana.
tacubayensis, *Fourn. Mex. Pl. En. Gram.* 95, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 551.—Mexic.
tarda, *Dvorák, in Flora*, lx. (1877) 278.—Tyrolia.
tenacissima, Linn. f. Suppl. 107 = Sporobolus indicus.
tenella, Hoffm. Fl. Germ. ed. II. i. 36 = vulgaris.
tenella, Poir. Encyc. Suppl. i. 257 = Airopsis globosa.
tenella, Willd. ex Steud. Nom. ed. II. i. 42 = Muehlenbergia tenuifolia.
tenella, Schlech. ex Gaud. Agrost. Helv. i. 75 = Calamagrostis tenella.
tenerrima, Trin. Gram. Unifl. 205 = elegans.
Tenori, Pito, ex Lond. Hort. Brit. 26.—Italia ?
tenuiflora, Willd. Sp Pl. i. 364 = Muehlenbergia Willdenowii.
tenuiflora, Steud. Syn. Fl. Gram. 163 = perennans.
tenuifolia, Curt. Brit. Gr. 42 = canina.
tenuis, Bast. ex Roem. & Schult. Syst. ii. 277 = canina.
tenuis, Sibth. Fl. Oxon. 36 = vulgaris.
tenuis, *Vasey, in Bull. Torr. Club*, x. (1883) 21.—Calif.
tenuissima, Spreng. Syst. i. 258 = Sporobolus indicus.
tenuissima, Vest, ex Steud. Nom. ed. II. i. 42 = Polypogon monspeliensis.
Thunbergii, *Steud. Syn. Pl. Gram.* 163.—Japon.
thyrsoides, Boec, ex Steud. Nom. ed. II. i. 42 (= *Vilfa fulvescens*).
Tillieri, Willd. ex Steud. l. c. = Catabrosa aquatica.
tolucensis, *H. B. & K. Nov. Gen. et Sp.* i. 135.—Mexic.
tolucensis, Willd. ex Steud. Syn. Pl. Gram. 164 = virescens.
Torreyana, Schult. Mant. ii. 208 = compressa.
Torreyi, *Kunth, Enum. Pl.* i. 226.—Am. bor.
transilvanica, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 50 = canina.
tremula, Willd. Sp. Pl. i. 372 = Sporobolus tremulus.
triaristata, Knapp, Gram. Brit. 23 = Polypogon monspeliensis.
trichanthe, *Schrank, in Regensb. Bot. Denkschr.* ii. (1818) 4.—Labrador.
trichoclada, *Griseb. in Ledeb. Fl. Ross.* iv. 439.—Orima.
trichoides, *Roem. & Schult. Syst.* ii. 861.—Peruv.
trichopoda, Ell. Sketch, i. 136 = Muehlenbergia trichopodioides.
trichotoma, Auct. ex Steud. Nom. ed. II. i. 43 = alba.
trichotoma, *Nees, ex Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 336.—Monte Video.
trichotoma, *Kit. in Linnaea*, xxxii. (1863) 306.—Austria.
tricuspidata, Hack. in Oestr. Bot. Zeitschr. xxvii. (1877) 46 = canina.
trifora, Hort. Hafn. ex Steud. Nom. ed. II. i. 43 = Chaeturus fasciculatus.
trinervata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1846) 347.—Caucas.
tristoides, *Steud. Syn. Pl. Gram.* 172.—Chili.
trupica, *Beauv. Fl. Oware*, ii. 37.—Afr. trop.

AGROSTIS :—
truncatula, *Parl. Fl. Ital.* i. 185.—Europ.
uliginosa, *Phil. in Anales Univ. Chil.* (1866) ii. 328.—Chili.
umbellata, *Colla, in Mem. Acc. Torin.* xxxix. (1836) 98.—Chili.
umbellulata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi (1846) 370.—Afr. austr.
umbrosa, *Fers. ex Trin. l. c.* 345.—Gallia.
umbrosa, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 47 = vulgaris.
vaginaeflora, Torr. in A. Gray, Man. 576 = Sporobolus vaginaeflora.
vaginata, *Steud. Syn. Pl. Gram.* 173.—Ins. Sandvic.
valentina, *Roem. & Schult. Syst.* iii. 348.—Hispan.
valvata, *Steud. Syn. Pl. Gram.* 171.—Japon.
varia, Host, Gram. iv. t. 57 ; Fl. Austr. i. 100 = alba.
variana, Thuil. Fl. Par. ii. 35 = canina.
varians, *Trin. in Mem. Acad. Pétersb.* Sér. VI. vi. (1846) 314.—Am. bor.; Chili.
velutina, Poir. ex Steud. Nom. ed. II. i. 43 = Paspalum velutinum.
ventosa, Dulac, Fl. Hautes-Pyr. 74 = Apera Spica-venti.
ventricosa, Gouan, Hort. Monsp. 89. 547 = Gastridium australe.
venusta, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1846) 340.—Austral.
vernalis, Hort. Dorp. ex Trin. Gram. Unifl. 198 = alba.
vernalis, Poir. Encyc. Suppl. i. 259 = Milium vernale.
versicolor, Hoppe, ex Steud. Nom. ed. II. i. 43 = alpina.
versicolor, Hort. Hafn. ex Trin. Gram. Unif. 201, 203 = vulgaris.
versicolor, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 293 = Catabrosa versicolor.
verticillata, Lam. Encyc. i. 59 ; Illustr. n. 831 = Andropogon squarrosus.
verticillata, Thuil. Fl. Par. ii. 36 = vulgaris.
verticillata, *Vill. Prosp.* 16.—Europ.
vesuta, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 401.—Abyss.
Vidali, *Phil. in Anales Univ. Chil.* (1878) 561.—Chili.
Villarsii, Poir. Encyc. Suppl. i. 261 = verticillata.
villosa, A. Spreng. Syst. Suppl. 5 = Sporobolus ciliatus.
villosa, Poir. ex Steud. Nom. ed. II. i. 43 = Tricholoena insularis.
villosa, Kelchb. ex Steud. l. c. = Sporobolus villosus.
villosa, Vill. Hist. Pl. Dauph. i. 378 = Calamagrostis sylvatica.
vimealis, [Honck.] Verz. aller Gew. Teutschl. 205 = castellana.
vimealis, With. Arr. Brit. Pl. ed. III. ii. 127 = canina.
vimealis, Schreb. Spicil. 47 = alba.
violacea, *Phil. in Anales Univ. Chil.* (1873) 560.—Chili.
violacea, Thuil. Fl. Par. ii. 35 = vulgaris.
violaceo-purpurea, Gilib. Exercit. ii. 525 = canina.
virescens, *H. B. & K. Nov. Gen. et Sp.* i. 135.—Calif.; Mexic.
virescens, Rupr. in Bull. Acad. Brux. ix. (1842) 229 = tolucensis.
virgata, Willd. ex Steud. Nom. ed. II. i. 43 = Leptochloa tenerrima.
virginica, Forsk. Fl. Aegypt. Arab. 20 = Sporobolus spicatus.
virginica, Linn. Sp. Pl. 62 = Sporobolus virginicus.
virginica, Muhl. ex Steud. Nom. ed. II. i. 43 = Sporobolus vaginaeflorus.
viridis, Gouan, Hort. Monsp. 546 = Sporobolus pungens.
viridis, *Rafin. in Am. Monthly Mag.* (1811) 856, nomen.—Am. bor.
Virletii, *Fourn. Mex. Pl. En. Gram.* 96, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 551.—Mexic.
viridgaris, Bivona, ex Roem. & Schult. Syst. ii. 347 in nota = alba.
vulgaris, *With. Arr. Brit. Pl.* ed. III. ii. 132.—Reg. temp.
Wallichiana, *Steud. Syn. Pl. Gram.* 174.—Reg. Himal.
Wiluitzkii, Steud. l. c. 188 = Reuteri.
Wightii, *Nees, ex Steud. l. c.*—Ind. or.
Youngii, *Hook. f. Handb. New Zeal. Fl.* 330.—N. Zel.
Zenkeri, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1846) 363.—Ind. or.
zeylanica, *Klein, ex Steud. Syn. Pl. Gram* 88.—Zeylan.

AGROSTISTACHYS, Dalz. in Hook. Kew Journ. ii. (1850) 41. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 302.
SARCOLINIUM, Wight, Ic. v. 24. t. 1887, 1888, (1852).
africana, *Muell. Arg. in Flora*, xlvi. (1864) 534.—Afr. trop.
Gandichandii, *Muell. Arg. in Linnaea*, xxxix. (1865–66) 144.—Ind. or.
Hookeri, *Benth. & Hook. f. Gen.* iii. 303.—Ceylan.
indica, *Dals in Hook. Kew Journ. Bot.* ii. (1850) 41. —Ind. or.
longifolia, *Benth. ex Hook f. Fl. Brit. Ind.* v. 407.—Canara.
longifolia, Kurz, For. Fl. Brit. Burma, ii. 377 = Gandichandii.

AGROSTOCRINUM, F. Muell. Fragm is. 94 (1860). *LILIACEAE*, Benth. & Hook. f. iii. 792.
stypandroides, *F. Muell. l. c.* 95.—Austral.

AGROSTOMIA, Cerv in La Naturaleza (1870) 345 = **Panicum**, Linn. (Gramin.).
aristata, Cerv. l. c.—Mexic.
barbata, Cerv. l. c. 346.—Mexic.
mutica, Cerv. l. c. 345.—Mexic.

AGROSTOPHYLLUM, Blume, Bijdr. 368. t. 53 (1825). *ORCHIDEAE*, Benth. & Hook. f iii. 517.
DIPLOCONCHIUM, Schau in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 428. t. 12.
callosum, *Reichb. f. in Seem. Fl. Vit.* 296.—Reg. Himal.
Hasseltii, *Reichb. f. in Bonplandia*, v. (1857) 41; *et in Walp. Ann.* vi. 909.—Ind. or.
javanicum, *Blume, Bijdr.* 369. t. 53.—Java.
khasyanum, *Griff. in Calc. Journ. Nat Hist.* iv. (1844) 376.—Reg. Himal.
longifolium, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
megalurum, *Reichb. f in Seem. Fl. Vit.* 296.—Ins. Samoa.
planicaule, Reichb. f. in Walp. Ann. vi. 909 = khasyanum.

AGUAVA, Rafin. Sylva Tellur. 107 (1838) = **Myrcia**, DC. (Myrtac.).
guianensis, Rafin. l. c. = M. guianensis.
tomentosa, Rafin. l. c. = M. tomentosa.

AGYLLA, Phil. in Anales Univ. Chil. (1865) 1. 648 = **Cladium**, P. Br (Cyperac.).
ensifolia, Phil. l. c. = C. scirpoideum.

AGYLOPHORA, Neck. Elem. i. 145 (1790) = **Uncaria**, Schreb. (Rubiac.).

AGYNAIA, Hassk. in Flora, xxv. (1842); Beibl. 40 = seq.

AGYNEIA, Linn. Mant. ii. 161 (1771); Vent. Jard. Cels, t. 92 (1800). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 271.
DIPLOMORPHA, Griff. Notul. iv. 479 (1854).
affinis, Kurz, ex Teijsm. & Binn. in Tijdschr. Ned. Ind. xxvii. (1864) 118 = bacciformis.
bacciformis, *A. Juss. Tent. Euph.* 24.—Ind. or.; Malay.
Berberi, Spreng. Syst. iii. 19 = Phyllanthus grandifolius.
ciliata. Wall. Cat. n. 7892 = Trigonostemon semperflorens.
coccinea, [Buch.-Ham. in] Symes, Embassy, ed. II. iii. 317 = Glochidion coccineum.
flexuosa, Heyne, ex Wall. Cat. n. 7368 = Glochidion zeylanicum.
glomerulata, *Miq. Fl. Ind. Bat. Suppl.* 447.—Sumatra.
hirsuta, *Miq. l. c.* i. 11. 368.—Malaya.
impubera, *Miq. l. c.* 367.—Malaya.
impubera, Wall. Cat. n. 7869 = Glochidion multiloculare.
impubes, Herb. Madr.! Wall. Cat. n. 7990 partim = Glochidion littorale.
impubes, Linn. Mant. ii. 296 = Phyllanthus puberus.
impubes, Vent. Jard. Cels, 28. t. 28 = bacciformis.
lanceolata, *F. G. Dietr. Lexik. Gaerte.* i. 186.—Austral.!
latifolia, Moon, Cat. 65 = Aporosa latifolia.

AGYNEIA:—
multiflora, Hassk. Cat. Hort. Bog. Alt. 240 (= Phyllanthi, sp.).—Malaya.
multilocularis, Rottl. ex Willd. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 206 = Glochidion multiloculare.
obliqua, Willd. Sp. Pl. iv. 568 = Glochidion zeylanicum.
ovata, Poir. Encyc. Suppl. i. 243; Miq. Fl. Ind. Bat. i. ii. 367 = Sauropus silvicus.
phyllanthoides, Spreng. Syst. iii. 19 = bacciformis.
pinnata, Miq. Fl. Ind. Bat. i. ii. 368 = Glochidion obscurum.
pubera, Linn. Mant. ii. 296 = Phyllanthus puberus.
sinica, Miq. Fl. Ind. Bat. i. ii. 368 = Phyllanthus puberus.
tenera, Zoll. & Mor. ex Miq. Fl. Ind. Bat. i. ii. 365 = Andrachne australis.
tetrandra, Buch.-Ham. in Trans Linn. Soc. xv. (1826) 125 = Phyllanthus Hamiltonianus.
tetrandra, Wall. Cat. n. 7852 = Trigstemon Hookerianus.

AHOUAI, Tourn. ex Adans. Fam. ii. 171 (1763) = **Thevetia**, Linn. (Apocyn.).

AICHRYSON, Webb & Berth. Phyt. Canar. i. 180. t. 27 (1840) = **Sempervivum**, Linn. (Crassul.).
Berkreacurtianum, C. Bolle, in Bonplandia, vii. (1859) 248.—Ins. Canar.
Bollei, Webb, ex Bolle, l. c.—Ins. Canar.
dichotomum, Webb & Berth. Phyt. Canar. i. 181 = S. dichotomum.
pachycaulon, C. Bolle, in Bonplandia, vii. (1859) 244.—Ins. Canar.
palmense, Webb, ex C. Bolle, l. c. 243.—Ins. Canar.
Pariatorei, C. Bolle, l. c. 244.—Ins. Canar.
parviflorum, C. Bolle, l. c. 243.—Ins. Canar.
porphyrogennetos, C. Bolle, l. c.—Ins. Canar.
pulchellum, C. A. Mey. ex Ftisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 55.—Hab.?
punctatum, Webb & Berth. Phyt. Canar. i. 182 = S. punctatum.
pygmaeum, Webb & Berth. l. c. 184.—Ins. Canar.
radicescens, Webb & Berth. l. c. 183 = S. villosum.
sedifolium, Webb, ex C. Bolle, in Bonplandia, vii. (1859) 242.—Ins. Canar.
tortuosum, Webb & Berth. Phyt. Canar. i. 184 = S. tortuosum.
villosum, Webb & Berth. l. c. 181 = S. villosum.

AIDELUS, Spreng. Syst. iv. Cur. Post. 17 (1827) = **Veronica**, Tourn. (Scrophal.).
micranthus, Spreng. l. c.—Nepal.

AIDIA, Lour. Fl. Cochinch. 143 (1790). Gen. haud recogn. DC. Benth. & Hook. f. ii. 39.
cochinchinensis, Lour. l. c.—Cochinch.

AIGIROS, Rafin. Alsog. Am. 42 (1838) = **Populus**, Tourn. (Salic.).
heterophylos, Rafin. l. c. = P. heterophylla.

AIKINIA, R. Br. in Wall. Pl. As. Rar. iii. 65. t. 288 (1832) = **Epishamia**, Blume (Gesnerac.).
Brunonis, Wall. Pl. As. Rar. l. c. 66. t. 288 = E. Brunonis.
carnosa, G. Don, Gen. Syst. iv. 665 = E. carnosum.
Horsfieldii, R. Br. in Wall. Pl. As. Rar. iii. 66 = E. difforme.

AIKINIA, Salisb. ex A. DC. Monog. Camp. 185, 141 (1830) = **Wahlenbergia**, Schrad. (Campan.).
hederacea, Salisb. ex A. DC. l. c. = W. hederacea.

AIKINIA, Wall. Pl. As. Rar. iii. t. 278. p. 46 (1832) = **Ratesburgia**, Kunth (Gramin.).
elegans, Wall. l. c. = R. pulcherrima.

AILANTHUS, Desf. in Mém. Acad. Sc. Par. 1786. (1789) 265. t. 8. *SIMARUBEAE*, Benth. & Hook. f. i. 309.
excelsa, *Roxb. Pl. Corom.* i. t. 28.—Ind. or.
flavescens, Carr. in Rev. Hortic. (1865) 366 = Cedrela sinensis.
glandulosa, *Desf. in Mém. Acad. Sc. Par.* 1786. (1789) 265. t. 8.—China.
gracilis, Salisb. Prod. 171 = Brucea sumatrana.

AILANTHUS:—
imberbiflora, *F. Muell. Fragm.* iii. 42; xix. 44.—Austral.
integrifolia, Lam. ex Steud. Nom. ed. II. i. 43 = moluccana.
japonica, Hort. ex C. Koch, Dendrol. 569 = glandulosa.
malabarica, *DC. Prod.* ii. 69.—Malabar.
moluccana, *DC. l. c.*—Ins. Moluc.
Pongelion, J. F. Gmel. Syst. ii. 726 = glandulosa.
procera, Salisb. Prod. 171 = glandulosa.
punctata, F. Muell. Fragm. iii. 42 = Pentaceras australis.
rhodoptera, F. Muell. l. c. 43 = glandulosa.

AILANTUS, DC. Prod. ii. 68 (1825) = prsec.

AILUROSCHIA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 151 = **Astragalus**, Tourn. (Legum.).
amygdaliger, Stev. l. c. = Astrag. amygdaliger.
diffusa, Stev. l. c. = Astrag. dolichophyllus.
galactites, Stev. l. c. = Astrag. galactites.
glomeratus, Stev. l. c. = Astrag. glomeratus.
hypogaeus, Stev. l. c. = Astrag. hypogaeus.
rupifragus, Stev. l. c. 152 = Astrag. rupifragus.
scabriseta, Stev. l. c. 151 = Astrag. scabriseta.

AILLYA, De Vriese, in Nat. Verh. Haarl. x. (1854) 75 = **Goodenia**, Sm.
umbellata, De Vriese, l. c. = G. filiformis.

AIMORRA, Rafin. New Fl. Am. iv. 70 (1836). (Compos.).
acuminata, Rafin. l. c.—Am. bor.

AINSLIAEA, DC. Prod. vii. 18 (1838). *COMPOSITAE*, Benth. & Hook. f ii. 493.
DIASPANANTHUS, Miq. in Ann. Mus. Lugd. Bat. ii. 186 (1865).
acerifolia, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126.—Malaya.
affinis, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 187 = acerifolia.
angustifolia, *Hook. f. & Thoms. ex Clarke, in Journ. Linn. Soc.* xiv. (1875) 412.—Ind. or.
spiculata, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126.—Japon.
aptera, *DC. Prod.* vii. 14.—Reg. Himal.
aspertima, *Sch. Bip. in Pollichia*, xx-xxi. (1863) 407. —Nepal.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 318.—Burma.
cordifolia, *Franch. & Sav. Enum. Pl. Jap.* ii. 416.—Japon.
dissecta, *Franch. & Sav. l. c.* i. 264.—Japon.
fragrans, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 236.—Hongkong.
glumacea, Sch. Bip. in Pollichia, xx-xxi. (1863) 407; Klatt, in Sitzb. Muench. Akad. (1878) 97 = angustifolia.
latifolia, *Sch. Bip. in Pollichia*, xvii-xix. (1861) 190. —Nepal.
pteropoda, *DC. Prod.* vii. 14.—Reg. Himal.
silhetensis, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 411 = pteropoda.
uniflora, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126.—Java.
Walkeri, *Hook. f. Bot. Mag.* t. 6925.—Hongkong.

AINSWORTHIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 343 = **Tordylium**, Linn. (Umbell.).
Carmeli, Boiss. Diagn. Ser. I. x. 44.—Palestin.
cordata, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 343 = T. cordatum.
elegans, Boiss. & Bal. Diagn. Ser. II. v. 100.—Taurus.
trachycarpa, Boiss. l. c. Ser. I. x. 44.—Orient.

AIOLOTHECA, DC. Prod. v. 508 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 352.
perthamioides, *DC. l. c.*—Mexic.

AIOUEA, Aubl. Pl. Gui. i. 320. t. 120 (1775).
LAURINEAE, Benth. & Hook. f. iii. 158.
ARIVEA, Steud. Nom. ed. I. 35 (1821).
CELONANDRA, Neck. Elem. ii. 143 (1790).
DOUGLASSIA, Schreb. Gen. Pl. 600 (1791).
EHRHARDIA, Scop. fastrod. 127 (1777).
brasiliensis, *Meisn. in DC. Prod.* xv. i. 83.—Bras.

AIOUEA :—
densiflora, *Nees, in Linnaea*, xxi. (1848) 513.—Am. trop.
elliptica, *Meisn. in DC. Prod.* xv. i. 83.—Am. trop.
guianensis, Nees, Syst. Laurin. 863.—(Cf. Meisn. in DC. Prod. xv. i. 82) = sequ.?
guianensis, *Aubl. Pl. Gui.* t. 310.—Guiana.
guyanensis, Griseb. ex Meisn. in DC. Prod. xv. i. 83 = Aydendron riparium.
Malabonga, *Blanco, Fl. Filip* ed. I. 233.—Ins. Philipp.
saligna, *Meisn. in DC. Prod.* xv. i. 82.—Bras.
Schomburgkii, *Meisn. l. c.*—Am. trop.
tenella, Nees, in Linnaea, xxi. (1848) 512 = guianensis.
tenuiflora, Nees, ex Meisn. in DC. Prod. xv. i. 510 (= *Mespiloaphne indecora*).
trinervis, *Meisn. l. c.* 83.—Am. trop.

AIPHANES, Willd. in Mém. Acad. Berl. (1804) 32 = **Martinezia**, Ruiz & Pav (Palm.).
aculeata, Willd. l. c. = M. Aiphanes
caryotæfolia, H. Wendl. in Kerch. Palm. 230 = M. caryotæfolia.
corallina, H. Wendl. l. c. = M corallina.
elegans, H. Wendl. l. c. = M elegans.
erinacea, H. Wendl. l. c.—N Granat.
Lindeniana, H. Wendl. l. c. = M Lindeniana.
Pragn, H. B. & K. Nov. Gen. et Sp i. 303 = Euterpe Pragn.
truncata, H. Wendl. in Kerch. Palm. 230 = M truncata.

AIPYANTHUS, Stev. in Bull. Soc Nat. Mosc. (1851) i. 599 = **Macrotomia**, DC. (Horag.).
echioides, Stev. l. c. 600 = M. echioides.

AIRA, Linn. Fl. Lapp. 27 (1737); Gen. ed I. 335. n. 838 (1737). *GRAMINEÆ*, Benth & Hook. f. iii. 1156.
ANTINORIA, Parl. Fl. Palerm. i. 92 (1845).
ASPRIS, Adans. Fam. ii. 496 (1763).
FIORINIA, Parl. Fl. Ital. i. 232 (1848).
FUSSIA, Schur, Enum. Pl. Transs. 764 (1866).
LEPTOPHORA, Ehrh. Beitr. iv. 146 (1789).
MOLINERIA, Parl. Fl. Ital. i. 236 (1848).
PERIBALLIA, Trin. Fund. Agrost. 133 (1848).
PROINIRIA, Ehrh. Beitr. iv. 149 (1789).
acutiglumis, *Steud. Syn. Pl. Gram.* 221.—Chili.
aegilopoides, *Walt. Fl. Carol.* 78.—Carolina.
æstivosis, Guss. Fl. Sic. Syn. i. 151 = Corynephorus articulatus.
aggregata, Timer. ex Bor. Fl. Centr. Fr. ed. III. ii. 702 = caryophylles.
agrostidea, Guss. Fl. Sic. Prod. i. 61 = insularis.
agrostides, *Lousi. Notic.* 16.—Europ. amer.
alba, Vahl, in Roem. & Schult. Syst. ii. 688 = Deschampsia alba.
alba, Wulf. ex Kunth, Gram. 110 = Scleria elongata.
alpina, Lilljebl. Utkast Sv. Fl. ed. I. 49; ed. II. 41 = Hierochloe alpina.
alpina, Linn. Sp. Pl. 65 = Deschampsia alpina.
alpina, Roth, Tent. Fl. Germ. ii. i. 98 = Deschampsia cæspitosa.
alpina, Savi, Bot. Etrusc. i. 52 = Deschampsia juncea.
alpina, Vahl, in Fl. Dan. t. 963 = atropurpurea.
alpina, Sobol. ex Rupr. in Beitr. Pfl. Russ. Reich. iv. 37 = Deschampsia flexuosa.
altaica, Trin. in Bunge, Verz. Suppl. Fl. Alt. 8 = Poa soongarica.
altissima, Lam. Fl. Fr. iii. 581 = Deschampsia cæspitosa.
ambigua, Michx. Fl. Bor. Am. i. 61 = Deschampsia ambigua.
ambigua, De Not. in Ann. Sc. Nat. Sér. III. v. (1846) 366 = capillaris.
Andrzei, Aserus, ex Andrae, in Bot. Zeitschr. xiv. (1856) 206 = Deschampsia cæspitosa.
anomala, *Trin. in Linnaea*, x. (1836) 301.—Chili.
antarctica, Forst. f. Prod. 8 = Trisetum antarcticum.
antarctica, *Hook. fr. Fl.* i. 150.—Ins. Antarct.
aquatica, Guss. Fl. Sic. Prod. i. 61 = Catabrosa ochroleuca.
aquatica, Linn. Sp. Pl. 64 = Catabrosa aquatica.
aquatica, Huds. Fl. Angl. ed. II. 34, pro parte = Glyceria distans.
arctica, Spreng. Syst. Cur. Post. 88 = Deschampsia cæspitosa.

AIRA :—
arenaria, *F. W. Schmidt, in Neu. Abh. Boehm. Ges.* i. (1791) 11.—Europ.
argentea, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 3.—Ins. Mader.
aristulata, *Torr. Fl. U. St.* 182.—Am. bor.
articulata, Desf. Fl Atlant. i. 70 = Corynephorus articulatus.
arundinacea, Lilljebl. Utkast Sv. Fl. ed. I. 49 = Graphephorum arundinaceum.
arundinacea, Linn. Sp. Pl. 64 = Poa tatarica.
atropurpurea, *Wahlenb. Fl. Lapp.* 37.—Europ ; Am. bor.
atroviren, Thuil. Fl. Par. ed. II. i. 37 = Mollinia caerulea.
aurea, Steud. in Flora, xii. (1829) 470 = Airopsis aurea.
australis, Raoul, Choix, 12 = Deschampsia cæspitosa.
australis, Steud. Syn. Pl. Gram. 220 = Deschampsia australis.
avenacea, Spreng. Syst. i. 278 = Eriachne avenacea.
avenacea, Willd ex Steud. Nom. ed. II. i. 44 = Trisetum palustre.
baetica, Trin. in Bull Sc Acad. Pétersb i. (1836) 67 = Deschampsia baetica.
benghalensis, J. F. Gmel Syst. i. 174 = Arundo bengalensis.
Berteroniana, *Steud. Syn. Pl. Gram.* 220.—Chili.
Bieberatensis, Steud. l. c. 219 = Deschampsia cæspitosa.
bottnica, Wahlenb. Fl. Lapp 36. t. 4 = Deschampsia cæspitosa.
brasiliensis, Raddi, Agrost. Bras. 36 = Eragrostis airoides.
brasiliensis, Spreng. Syst. i. 278 = Arundinella brasiliensis.
breviaristata, Gilib. Exercit. ii. 528 = Deschampsia cæspitosa
brevisulmis, Loisel. Fl. Gall. i. 57 = Corynephorus canescens.
brevisetosa, Bieb. Fl. Taur. Cauc. iii. 63 = Deschampsia cæspitosa.
brevifolia, *Torrh, Fl. Am Sept.* i. 76.—Missouri.
brigantica, Chaix, in Vill. Hist. Pl. Dauph. ii. 378 = Glyceria distans.
bulbosa, Krock. Fl. Sil. Suppl. i. 131 = Koeleria cristata ?
caerulea, Linn Sp. Pl 63 = Molinia caerulea.
cæspitosa, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 350, pro parte = Festuca furgians.
cæspitosa, Linn Sp. Pl. 64 = Deschampsia cæspitosa.
cæspitosa, Muhl. Desc. Gram. 55 = Deschampsia ambigua.
calycina, *Steud. Syn. Pl. Gram* 220.—Ecuv.
canescens, All. Fl. Pedem. ii. 244 = Corynephorus articulatus.
canescens, Host, Gram. iv. t. 36 = caryophylles.
canescens, Linn. Sp. Pl. 65 = Corynephorus canescens.
capensis, Linn. f. Suppl. 108 = Ehrharta calycina.
capensis, Steud. in Flora, xii. (1829) 468 = Koeleria capensis.
capensis, Spreng. Syst. ii. 114 = Ehrharta villosa.
capillacea, Hort. Madr. ex Wall. Cat. sub n. 3804 = Leptochloa filiformis.
capillacea, *Lam. Illustr.* i. 177.—Am. bor.
capillaris, Guss. Fl. Sic. Prod. i. 64 = Avena corymbosa.
capillaris, *Host, Gram.* iv. 20. t. 35.—Europ. ; Oriens.
capillaris, Lag. in Varied. Cienc. ii. iv. (1805) 39 = Lagascea, Airopsis agrostidea.
capillaris, Salisb. ex Steud. Nom. ed. II. i. 44 = caryophylles.
capillaris, Savi, Fl. Pis. i. 86 = Deschampsia juncea.
caryophyllacea, Lears, Fl. Herborn. ed. II. 24 = Deschampsia flexuosa.
caryophyllea, Bertol. [ex Stend. ; an Fl. Ital. i. 455] = capillaris.
caryophyllea, *Linn. Sp. Pl.* 66.—Europ. ; Oriens.
catabrosoides, *Steud. Syn. Pl. Gram.* 222.—Caucas.
cespitosa, Muhl. Desc. Gram. 55 = Deschampsia ambigua.
Chamaemespilus, *Host, Fl. Austr.* ii. 8.—Europ.
chinensis, *Retz. Obs.* iii. 10.—China.
ciliaris, *Kit. in Linnaea*, xxvii. (1855) 810.—Austria.
ciliaris, Auct. ex Steud. Nom. ed. I. 28, 487 = Koeleria villosa.
ciliata, Spreng. Pugill. i. 7 = Eragrostis minutiflora.
ciliata, Spreng. Syst. i. 278 = Eriachne ciliata.
clavata, Gilib. Exercit. ii. 526 = Corynephorus canescens.
clavata, Willd. ex Steud. Nom. ed. II i. 44 = Corynephorus articulatus.
collina, Dum. Obs. Gram. Belg. 121 = Deschampsia flexuosa.
compressa, *Rafin. in Am. Monthly Mag.* (1811) 350. nomen.—Am. bor.
controversa, *Steud Syn. Pl. Gram.* 224.—Am. bor.
complanata, Steud. Nom. ed. II. i. 44 = Chlorispetræa.
corsica, Jord. Pugill. 353 = sicula.
corsica, Tausch, in Flora, xx. (1837) 102 = Deschampsia flexuosa.
corymbosa, Willd. ex Steud. Nom. ed. II. i. 44 = caryophylles.
corymbosa, Fauché & Chaub. Fl. Pelop. 5 n. 120. t. 7 [1 = intermedia.
cristata, Linn Sp. Pl. 63 = Koeleria cristata.
Cupaniana, Guss Fl. Sic. Syn. i. 145 = caryophylles.
cuprana, Fuss, Fl Transs. 722 = Deschampsia flexuosa.
curta, Jord. ex Bor Fl. Centr. Fr. ed. III. ii. 701 in nota = caryophylles.
dactyloides, Rochel, Pl. Rar. Ban. 32 f. 3 = Koeleria glauca.
danthonioides, Trin in Mém. Acad. Pétersb. Sér. VI. i (1830) 57 = Deschampsia calycina.
discolor, Thuil. Fl. Par. i. 39 = setacea.
distichophylla, Spreng. Syst. i. 278 = Panicum subulatum.
divaricata, Pourr. in Mém. Acad. Toul. iii. (1788) 307 = caryophylles.
durinscula, *Pour. Encyc. Suppl.* ii. 76.—Hab. ?
Edouardi, Reut. ex Jouve, in Bull. Soc. Bot. Fr. xii. (1865) 83 = Avena corymbosa.
effusa, Spreng. Syst i. 278 = Eriachne pallescens.
elatior, 'Hoffmgg.' ex Steud. Nom. ed. II. i. 44 = Melica Bauhini ?
elatior, *Steud. Syn. Pl. Gram.* 423.—Reg. Magellan.
elegans, Willd. ex Gaud. Agrost. Helv. i. 130, 355 = capillaris.
elegantes, 'Vill.' ex Steud. Nom. ed. II. i. 44 = Airopsis agrostides.
elegantissima, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 85 = capillaris.
elodes, Bingn. Fasc. Pl. Foroj. 10; cf. Flora (1840), 104 = Poa nemoralis.
elongata, Hook. Fl. Bor. Am. ii. 248 = Deschampsia elongata.
elongata, Salzm. ex Steud Nom. ed. II. i. 44 = Koeleria phleoides.
eufloruosa, Syme, Engl. Bot. ed. III. xi. 67 = Deschampsia flexuosa.
fascicularis, Steud. Syn. Pl. Gram. 219 = Corynephorus fasciculatus.
festucoides, Vill. Prosp. 16; Hist. Pl. Dauph. ii. 86 = Deschampsia alpina.
filiformis, *Korn. ex Roxb. Fl. Ind.* i. 328.—Ind. or.
flammea, Buchos, ex Honck. Syn. Pl. Germ. i. 521 = Deschampsia flexuosa.
flexilis, Link. ex Gieseke, Prael. 147 = Deschampsia flexuosa.
flexuosa, Linn. Sp. Pl. 65 = Deschampsia flexuosa.
Gayana, *Steud. Syn. Pl. Gram.* 220.—Chili.
gigantea, Schrenk, ex Ledeb. Fl. Ross. iv. 446 = Muehlenbergia alpestris.
gigantea, *Steud. Syn. Pl. Gram.* 224.—Ind. occ. ?
glabra, *Brot. Fl. Lusit.* 91.—Lusit.
glabrata, Brot. l. c. 92 = lendigera.
glauca, Hartm. ex Steud. Nom. ed. II. i. 44 = Deschampsia cæspitosa.
glauca, Rochel, Pl. Rar. Banat. 32 = Koeleria dactyloides.
glauca, Spreng. Nachtr. Bot. Gart. Halle, i. 10 = Dactylis glauca.
globosa, Thore, in Journ. Bot. Paris, i. (1808) 197 = Airopsis globosa.
glomerata, G. Don, ex S. F. Gray, Nat. Arr. Brit. Pl. iii. 137 = Deschampsia cæspitosa.
Gmelinii, [Honck.] Very. aller Gew. Teutschl. 212 = praecox ?
gracilis, Trin. ex Steud. Nom. ed. II. i. 44 = Koeleria cristata.
grandiflora, Bertol. Fl. Ital. i. 436 = Koeleria setacea.
grandiflora, Steud. Syn. Pl. Gram. 220 = Deschampsia grandiflora.
grandis, Nym. Consp. 807 = Deschampsia cæspitosa.

AIRA :—
Halleri, [Honck.] Vers. aller Gew. Teutschl. 212 = Trisetum rigidum.
Hartmaniana, Nym. Consp. 807 = Deschampsia caespitosa.
helodes. Steud. Nom. ed. II. i. 44 = Poa nemoralis.
hirsuta, Hall. f. ex Schleich. Cat. Nov. 5 = Koeleria hirsuta.
hispida, Spreng. Syst. i. 276 = Erischne capillaris.
holciformis, Steud. Syn. Pl. Gram. 221 = Deschampsia holciformis.
Holcus-lanatus, Vill. Hist. Pl. Dauph. ii. 87 = Holcus lanatus.
Holcus-mollis, Vill. l. c. 88 = Holcus mollis.
Hostii, *Steud. Syn. Pl. Gram.* 221.—Bohem.
humilis, Bieb. Fl. Taur. Cauc. i. 57 = Catabrosa humilis.
hybrida, Gaud. Agrost. Helv. i. 124 = Corynephorus articulatus.
incompleta, Boss. ex Steud. Nom. ed. II. i. 44 = Paspalum debile.
indica, Linn. Sp. Pl. ed. II. 94 = Panicum conglomeratum.
inflexa, Loisel. Fl. Gall. i. 56. t. 22 = Avena corymbosa.
insularis, *Boiss. Fl. Orient.* v. 628.—Europ. ; Orient.
Intermedia, *Guss. Fl. Sic. Prod. Suppl.* i. 105.—Europ.
interrupta, Rottl. ex Steud. Syn. Pl. Gram. 56 = Panicum canaliculatum.
involucrata, Cav. Ic. i. 88. t. 44 = Deschampsia involucrata.
ischaemoides, Koen. ex Trin. in Mém. Acad. Péterab. Sér. VI. iii. (1834) 228 = Panicum antipodum.
juncea, Vill. Prosp. 16 = Deschampsia juncea.
juncifolia, Wulf. ex Kunth, Enum. i. 322 = Scaleria tenuifolia.
Kingii, Hook. f. Fl. Antarct. ii. 376. t. 136 = Deschampsia caespitosa.
koeleroides, Peyr. in Linnaea, xxx. (1859-60) 5 = Deschampsia koeleroides.
Koenigii, J. F. Gmel. Syst. i. 174 = Eragrostis minutiflora.
la hrardnica, *Steud. Syn. Pl. Gram.* 220.—Labrador.
laevigata, Sm. Eng. Bot. t. 2102 = Deschampsia alpina.
laevis, Brot. Fl. Lusit. i. 90 = Deschampsia laevis.
laevis, Spreng. Syst. i. 278 = Erischne glauca.
Lagascæ, *Kunth, Gram.* i. 99.—Hispan. ; Corsic.
latifolia, Hook. Fl. Bor. Am. ii. 248 = Deschampsia latifolia.
latifolia, Steud. Syn. Pl. *Gram.* 219 = Deschampsia caespitosa.
latigluma, *Steud. l. c.* 221.—Abyss.
laxa, Kich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Orthoclada laxa.
Lerrii, Kunth, Enum. Pl. i. 289 = caryophyllea.
Leyei, Bor. in Soc. Angers, xxiv. (1855) ; et Fl. Centr. Fr. ed. III. ii. 700 = Deschampsia flexuosa.
lendigera, *Leg. in Varied. Cienc.* ii. iv. (1806) 39.—Hispan.
Lenzeri, Loisel. Fl. Gall. ed. II. i. 58 = caryophyllea.
Lenzii, Reichb. Fl. Germ. Excurs. 140* = prasc.
Lensei, Gren. & Godr. Fl. Fr. iii. 620 = prasc.
liburnica, Godr. ex Nym. Consp. 807 = Deschampsia caespitosa.
macrantha, Ledeb. in Mém. Acad. Pétersb. v. (1812) 315 = Koeleria cristata.
macranthera, *Steud. Syn. Pl. Gram.* 219.—Hispan.
macranthoidea, *Nees, ex Steud. l. c.* 221.—Madras.
magellanica, *Hook. f. Fl. Antarct.* ii. 376. t. 134.—Reg. Magellan.
magellanica, Lam. ex Beauv. Agrost. 65 = Hierochloë magellanica.
major, Syme, Engl. Bot. ed. III. xi. 68.—Deschampsia caespitosa, alpina.
media, Gouan, Illustr. 3 = Deschampsia caespitosa.
medii, Trin. in Mém. Acad. Péterab. Sér. VI. iv. (1841) Suppl. 57 = Deschampsia caespitosa.
meticodes, Michx. Fl. Bor. Am. i. 69 = Graphephorum meliicodes.
micrantha, *Steud. Syn. Pl. Gram.* 493.—Madag.
millacea, Vill. Hist. Pl. Dauph. ii. 81 = Poa millacea.
minuta, *Loef. It. Hisp.* 117.—Europ. ; Orient.
minuta, Loisel. Fl. Gall. i. 45 = Alropeis agrostides.
misurica, Spreng. Syst. ii. 578 = brevifolia.
mollis, Muhl. Descr. Gram. 89 = Estonia purpurascens.
mollis, Schreb. Spicil. 51 = Holcus mollis.
montana, Brot. ex Nym. Consp. 806 = Deschampsia stricta.

AIRA :—
montana, Desv. Obs. Angers, 55 = setacea.
montana, Koel. Gram. 131 = Deschampsia flexuosa.
montana, *Linn. Sp. Pl.* 65.—Europ.
mucronata, Spreng. Syst. i. 276 = Erischne brevifolia, mucronata.
multicaulis, Kunth, Enum. Pl. i. 292 = caryophyllea.
multiculmis, Dum. Obs. Gram. Belg. 121 = caryophyllea.
nana, Kunze, ex Steud. Nom. ed. II. i. 45 = praecox.
nana, Spreng. ex Steud. l. c. = caryophyllea.
nitida, Spreng. Mant. i. 32 = Estonia purpurascens.
nitida, *Steud. Syn. Pl. Gram.* 220.—Mexic.
nivea, *Host, Fl. Austr.* ii. 8.—Europ.
Notarisiana, Steud. Syn. Pl. Gram. 221 = capillaris.
nutans, [Web. in] Wigg. Prim. Fl. Holsat. 7 = Melica nutans.
obtusata, Michx. Fl. Bor. Am. i. 69 = Estonia obtusata.
oryzetorum, Spreng. Pugill. ii. 18 = Festuca arundinacea.
pallens, Muhl. Descr. Gram. 84 = Trisetum palustre.
pallens, *Spreng. Fl. Hal. Mant.* 36.—Hab. ?
pallescens, *Kit. in Roem. & Schult. Syst.* ii. 685.—Hungar.
paludosa, Roth, Tent. Fl. Germ. ii. i. 96 = Deschampsia flexuosa.
paludosa, Wibel, Prim. Fl. Werth. 118 = Deschampsia caespitosa.
palustris, Pollini, Cat. Ort. Veron. (1814) 8 = Festuca arundinacea.
panices, Willd. ex Steud. Nom. ed. II. i. 45 = Leptochloa filiformis.
paradoxa, *Steud. Syn. Pl. Gram.* 223.—Guinea.
parviflora, Thuil. Fl. Par. ed. II. i. 38 = Deschampsia caespitosa.
parvula, *Hook. f. Fl. Antarct.* ii. 377.—Fuegia.
patulipes, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 701 = caryophyllea.
Perszyniana, Spreng. in Mém. Acad. Péterab. (1810) 299. t. 7 = Estonia purpurascens.
petraea, Bellardi, ex Steud. Nom. ed. II. i. 45 ; ii. 361 = Sc. semoralis.
phleoides, Steud. Nom. ed. I. 24 = Koeleria phleoides.
pictiglume, *Steud. Syn. Pl. Gram.* 221.—Abyss.
pilosa, *Kit. in Linnaea*, xxii. (1849) 810.—Austria.
plesiantha, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 701 = caryophyllea.
poaeformis, Willd. ex Steud. Nom. ed. II. i. 45 = Poa depauperata.
praecox, *Linn. Sp. Pl.* 65.—Europ. ; Am. bor.
provincialis, Gennar. ex Nym. Consp. 814 = Avena corymbosa.
provincialis, Jord. ex Jouve, in Bull. Soc. Bot. Fr. xii. (1865) 54. t. 1 = capillaris.
pubescens, Vahl, Symb. iii. 9 = Koeleria villosa.
pulchella, Nocca & Balb. Fl. Ticin. i. 403. t. add. 1 = capillaris.
pulchella, Link, Hort. Berol. i. 180 = Avena corymbosa.
pulchella, Willd. Enum. Hort. Berol. 101 = lendigera.
pulchra, *Steud. Syn. Pl. Gram.* 220.—Chili.
pumila, Agardh, ex Roem. & Schult. Syst. ii. 678 = Sc. semoralis.
pumila, Parsh, Fl. Am. Sept. i. 76 = Poa annua.
pumila, Vill. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 372 = Deschampsia juncea.
purpurea, Muhl. Descr. Gram. 85 = Uralepsis aristulata.
purpurea, Walt. Fl. Carol. 78 = Triplasis purpurea.
purpureo-argentea, Gilib. Exercit. ii. 526 = caryophyllea.
pusilla, Schur, Enum. Pl. Transs. 754 = praecox.
rara, Spreng. Syst. i. 278 = Erischne rara.
refracta, Lag. in Varied. Cienc. ii. iv. (1806) 39 = Deschampsia refracta.
rigida, Dulac, Fl. Hautes-Pyr. 82 = caryophyllea.
Ruprechtii, Griseb. ex Hook. f. in Trans. Linn. Soc. xxiii. (1861) 346 = Avena Ruprechtii.
sabuletorum, Steud. Nom. ed. II. i. 45 = sabulonum.
sabulonum, *Labill. Sert. Austr. Caled.* 16. t. 21.—Nov. Caled.
saburralis, Jan, Elench. 8, ex Gaus. Fl. Sic. Syn. i. 148 = caryophyllea.
scaatthis, Balan. ex Trin. in Mém. Acad. Péterab. Sér. VI. i. (1831) 89 = capillaris.
scabra, *Kit. in Linnaea*, xxxii. (1863) 209.—Austr.
scabra, Hort. ex Kunth, Enum. Pl. i. 270 = Catabrosa viridula.

AIRA :—
scabra, Zuccagn. in Roem. Collect. 124 = Catabrosa scabra.
scabricostacea, Knapp, Gram. Brit. t. 22 = Deschampsia flexuosa.
seminudra, Walds. & Kit. in Schult. Oestr. Fl. i. 199 = Poa seminudra.
serotina, *Torr. sec. Trin. ex Steud. Nom.* ed. II. i. 45. —Am. bor.
setacea, *Huds. Fl Angl.* ed. I. 30 ; *Reis. Obs.* iii. 10.—Europ.
setacea, Pourr. ex Nym. Consp. 808 = Deschampsia juncea.
setiformis, *Host. sec. Trin. ex Steud. Nom.* ed. II. i. 45.—Hab. ?
sicula, Spreng. Syst. i. 277 = Holcus tenuis.
sicula, *Ind. Ind. Sem. Pan.* (1827) 38 ; *in Nuov. Giorn. Bot.* v. (1873) 156.—Sicilia.
spectabilis, *Steud. Syn. Pl. Gram.* 224.—Afr. austr.
spicaeformis, *Steud. l. c.* 424.—Reg. Magell.
spicata, Linn. Sp. Pl. 63 = Trisetum subspicatum.
spicata, Linn. l. c. 64 = Panicum inflcium.
splendida, Willd. ex Steud. Nom. ed. II. i. 45 = Deschampsia flexuosa.
squarrosa, Spreng. Syst. i. 276 = Erischne squarrosa.
stolonifera, Hausm. Fl. Tir. 890, sec. Nehlr. Nachtr. 25 = Deschampsia caespitosa.
stricta, Nym. Consp. 808 = Deschampsia stricta.
subaristata, Faye, ex Anderss. in Walp. Ann. vi. 992 = Deschampsia juncea.
subspicata, Linn. Syst.ed. X. 873 = Trisetum subspicatum.
subspicata, Raluv. in Gort. Fl. Belg. Foed. n. 74 = praecox.
subvirens, Lag. in Varied. Cienc. ii. iv. 39 = Deschampsia juncea.
superbiens, *Steud. Syn. Pl. Gram.* 424.—Reg. Magell.
subulorum, D. Dietr. Syn. i. 339 err. = sabulonum.
tenera, *Kit. in Schult. Oestr. Fl.* ed. II. i. 199.—Sisvonia ; Banatus.
Tenorei, Fuckel, ex Willk & Lange, Prod. i. 55 = Agrostis capillaris.
Tenorii, Guss. Fl. Sic. Prod. i. 62 = Avena corymbosa.
tenuifolia, *Steud. Syn. Pl. Gram.* 223.—Chili.
tenuissima, Spreng. Syst. i. 276 = Sporobolus minutiflorus.
Todari, Tenore, ex Tod. Ind. Sem. Hort. Panorm. (1859) (Linnaea, xxx. (1859-60) 758) = caryophyllea.
triaristata, *Clarke, Trav.* ii. 452.—Palaestina.
trichodes, Spreng. Syst. i. 276 = Agrostis trichodes.
trifera, Trivalt. in Flora, xix. (1836) 432 = Deschampsia caespitosa.
trifera, Ell. Sketch, i. 158 = Graphephorum Elliottii.
trifera, Roem. & Schult. Syst. ii. 680 = Deschampsia juncea.
trifera, Willd. ex Steud. Nom. ed. II. i. 45 = Corynephorus canescens.
trigitania, *Steud. Syn. Pl. Gram.* 223.—Texas.
truncata, Muhl. Descr. Gram. 83 = Estonia obtusata.
truncata, Salzm. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 56 = caryophyllea.
uliginosa, Weihe, ex Boenn. Prod. Fl. Monast. 25 = setacea.
uniaristata, *Cav. in Anales Cienc. Nat.* vi. (1803) 148.—Hab. ?
valesiaca, Colla, Herb. Pedem. t. 94. f. 1 = Koeleria setacea.
valesiaca, Bertol. Fl. Ital. i. 488 = prasc.
varia, Jacq. Hort. Vindob. 15 = Seleria caerulea.
variegata, *Steud. Syn. Pl. Gram.* 223. - Graecia.
versicolor, *Roem. & Schult. Syst.* ii. 679.—' Port. Egmont.'
ventita, *Steud. Syn. Pl. Gram.* 424.—Reg. Magell.
vicina, *Greff. Natal.* iii. 84.—Reg. Himal.
villosa, Gilib. Exercit. ii. 527 = montana.
villosa, Linn. f. Suppl. 109 = Elytrarta gigantea.
violacea, Gilib. Exercit. ii. 526 = Catabrosa aquatica.
violacea, *Steud. Syn. Pl. Gram.* 222.—Persia.
violacea, Willd. ex Steud. Nom. ed. II. i. 106 = Panicum antipodum.
vivipara, *Steud. Syn. Pl. Gram.* 222.—Groenland.
viridula, *Pers. Enchr. Suppl.* ii. 76.—Hab. ?
Wibeliana, *Steud. ex Pl. Zeas.* t. 2461, 2528.—Europ.
Wiblemii, Steud. Syn. Fl. Gram. 219 = Deschampsia caespitosa.

AIRIDIUM, Steud. Syn. Pl. Gram. 493 (1855) = Deschampsia, Beauv. (Gram.).
elegantulum, Steud. l. c. = D. setacea.

AIROCHLOA, Link, Hort. Berol. i. 126; ii. 276 (1827) = **Koeleria**, Pers. (Gram.).
albescens, Link, l. c. i. 128 = K. albescens.
capensis, Nees, in Mart. Fl. Bras. ii. 321 = K. capensis.
caudata, Link, in Linnaea, xvii. (1843) 405 = K. crassipes.
cristata, Link, Hort. Berol. i. 127 = K. cristata.
glauca, Link, ex Schur, Enum. Pl. Transs. 751 = K. glauca.
gracilis, Link, Hort. Berol. ii. 276 = K. cristata.
hirsuta, Link, l. c. i. 129 = K. hirsuta.
hispida, Link, l. c. = K. hispida.
permollis, Nees, ex Steud. Syn. Pl. Gram. 293 = K. permollis.
pubescens, Link, Hort. Berol. i. 128 = K. cristata.
uniflora, Hochst. in Flora, xxxviii. (1855) 580.—Afr. or.
valesiaca, Link, Hort. Berol. i. 127 = K. setacea.
villosa, Link, l. c. 128 = K. villosa.

AIROPSIS, Desv. Journ. Bot. i. (1808) 200, pro parte. *GRAMINEÆ*, Benth. & Hook. f. iii. 1156.
agrostidea, *DC. Fl. Fr. Suppl.* 963.—Europ. austr.
ampla, *Nees, ex Steud. Nom.* ed. II. i. 45.—Afr. austr.
aurea, *Nees, in Linnaea*, vii. (1832) 317.—Afr. austr.
brevifolia, Roem. & Schult. Syst. ii. 578 = Aira brevifolia.
Candollii, Desv. Journ. Bot. i. (1808) 200 = agrostidea.
capillaris, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 328 = Aira capillaris.
caryophyllea, Schur, l. c. = Aira caryophyllea.
Cupaniana, Lange, in Kjoeb. Vidensk. Meddel. (1860) 88 = caryophyllea.
globosa, *Desv. Journ. Bot.* i. (1808) 200.—Gall.; Hisp.
glabra, *Dum. Cours.* 54.—Ins. Borbon.
insularis, Nym. Syll. 411 = Aira insularis.
involucrata, Desv. Journ. Bot. i. (1808) 201 = Deschampsia involucrata.
jubata, *Griseb. in Goett. Abh.* xxiv. (1879) 992.—Reg. Argent.
lendigera, Lange, in Kjoeb. Vidensk. Meddel. (1860) 87 = Aira lendigera.
millegrana, Griseb. in Goett. Abh. xix. (1874) 262 = Eragrostis airoides.
minuta, Desv. Journ. Bot. i. (1808) 201 = Catabrosa minuta.
multiculmis, Lange, in Kjoeb. Vidensk. Meddel. (1860) 88 = Aira caryophyllea.
obtusata, Desv. Journ. Bot. i. (1808) 200 = Eatonia obtusata.
pallida, Nees, ex Steud. Nom. ed. II. i. 45 = Eriachne pallida.
peruviana, [Nees &] Meyen, Reise, ii. 484 = Diesanthelium supinum.
praecox, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 328 = Aira praecox.
pulchella, Tenore, Fl. Nap. iii. 56, 102 = Avena corymbosa.
Stendelii, Nees, in Linnaea, vii. (1832) 318 = Eriachne capensis.
tristis, Nees, ex Steud. Nom. ed. II. i. 45 = Eriachne tristis.
tuberculata, Nees, ex Steud. l. c. = Eriachne tuberculata.

AITCHISONIA, Hemsl. ex Aitch. in Journ. Linn. Soc. xix. (1882) 166. t. 14. *RUBIACEÆ*.
rosea, *Hemsl. l. c.*—Afghan.

AITHALES, Webb & Berth. Phyt. Canar. i. 178 (1838) = **Sedum**, Linn. (Crassul.).
caespitoses, Webb & Berth. l. c. 179, 180 = S. caespitosum.
rubens, Webb & Berth. l. c. 179 = S. rubens.

AITTARA, Endl. Gen. 254. (1837) = ATTARA.
Maregr. = **Hvodisa**, Forst. (Rutac.).

AITONIA, Trunb. in Phys. Sallsk. Handl. i. (1776) 106; Nov. Gen. 93. *SAPINDACEÆ*, Benth. & Hook. f. i. 411.
AYTONIA, Linn. f. Suppl. 49. 308 (1781).
capensis, *Thunb. Il. cc.*—Afr. austr.

AITOPSIS, Rafin. Fl. Tellur. iii. 91 (1836) = **Salvia**, Tourn. (Labiat.).
foliosa, Rafin. l. c. = S. foliosa.

AIZODRABA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 295 = **Draba**, Linn. (Crucif.).
aizoides, Fourr. l. c. = D. aizoides.
bengesiaca, Jord. & Fourr. l. c.—Gallia.
chrysantha, Jord. & Fourr. l. c.—Gallia.
ochroleuca, Jord. & Fourr. l. c. = D. algida.
savignea, Fourr. l. c. = D. aizoides.
trichocarpa, Jord. & Fourr. l. c.—Gallia.

AIZOON, Linn. Gen. ed. I. 161 (1737). *FICOIDEÆ*, Benth. & Hook. f. i. 854.
VESTINGIA, Heist. Syst. (1748); Fabric. Hort Helmst. 363 (1759).
angustifolium, D. Dietr. Syn. Pl. iii. 130 = Acrosanthes angustifolia.
argentinum, Eckl. & Zeyh. Enum. 325 = rigidum.
Casalcoronum, *Reichb. ex Spreng. Syst. Cur. Post.* 195.—Afr. austr.
canariense, Andr. Bot. Rep t. 201 = Sesuvium pedunculatum.
canariense, *Linn. Sp. Pl.* 488.—Ins. Canar.; Afr. bor.; Orient.
contaminatum, Eckl. & Zeyh. Enum. 326 = Galenia secunda.
crystallinum, Eckl. & Zeyh. l. c. 327 = Galenia crystallina.
elongatum, Eckl. & Zeyh. l. c. 326 = Galenia secunda.
fistulosum, D. Dietr. Syn. Pl. iii. 130 = Acrosanthes fistulosa.
frutescens, Eckl. & Zeyh. Enum. 327 = Galenia affinis.
fruticosum, Linn. f. Suppl. 261 = Galenia fruticosa.
galenoides, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 469.—Afr. austr.
glinoides, Eckl. & Zeyh. Enum. 325 = Galenia secunda.
glinoides, Linn. f. Suppl. 261.—Afr. austr.
Herniaria, Reichb. ex Steud. Nom. ed. II. i. 46 = Galenia hermiariaefolia.
hirsutum, Eckl. & Zeyh. Enum. 325 = glinoides.
hispanicum, *Linn. Sp. Pl.* 488.—Hispan.
lanceolatum, *Murr. Syst. Veg.* ed. XIII. 892.—Afr austr.
microphyllum, Bartl. in Linnaea, vii. (1832) 541 = Galenia hermiariaefolia.
montevidense, Spreng. ex Rohr, in Mart. Fl. Bras. xiv. ii. 170 = Sesuvium Portulacastrum.
mossamedense, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 554.—Afr. trop.
paniculatum, *Linn. Sp. Pl.* 488.—Afr. austr.
papulosum, Eckl. & Zeyh. Enum. 326 = Galenia papulosa.
perfoliatum, Linn. f. Suppl. 261 = Tetragonia Zeyheri.
procumbens, Cranz, Inst. i. 135 = canariense.
propinquum, Eckl. & Zeyh. Enum. 326 = Galenia secunda.
pubescens, Eckl. & Zeyh. l. c. = Galenia secunda.
quadrifidum, *F. Muell. Fragm.* ii. 148.—Austral.
rigidum, *Linn. f. Suppl.* 261.—Afr. austr.
sarmentosum, *Linn. f. l. c.* 260.—Afr. austr.
secundum, Linn. f. l. c. 261 = Galenia secunda.
sericeum, Fenzl, ex Harv. & Sond. Fl. Cap. ii. 470 = rigidum.
suaifolium, Moench, Meth. 698 = Hispanicum.
spathulatum, *Eckl. & Zeyh. Enum.* 326.—Afr. austr.
squamulosum, Eckl. & Zeyh. l. c. 326 = Galenia squamulosa.
stellatum, Lam. Encyc. iii. 418 = sarmentosum.
tortifolium, D. Dietr. Syn. Pl. iii. 130 = Acrosanthes tortifolia.
tomentosum, Lam. Encyc. iii. 418 = paniculatum.
virgatum, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 554.—Afr. trop.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 470.—Afr. austr.
zygophylloides, *F. Muell. Fragm.* viii. 229.—Austral.

AIZOUM, Linn. Syst. ed. I. (1755) = **Aizoon**, Linn. (Ficoid.).

AJAX, Salisb. in Trans. Hort. Soc. i. (1812) 343 = **Marcissus**, Linn. (Amaryll.).
(NOTA.—Cl. Baker *Ajaces* omnes ad *Narcissum Pseudo-Narcissum* ducit.)
abscissus, Haw. Rev. 116, 117 = N. abscissus.
albicans, Haw. Monog. 9 = N. Pseudo-Narcissus.
albus, Haw. Rev. 117 = N. moschatus.

AJAX:—
bicolor, Salisb. in Trans. Hort. Soc. i. (1812) 346 = N. Pseudo-Narcissus.
brevifolius, Haw. Monog. 2. n. 6 = N. Pseudo-Narcissus.
cambricus, Haw. l. c. 3 = N. Pseudo-Narcissus.
capax, M. Roem. Syn. Ensat. 201 = N. Pseudo-Narcissus.
cernuus, Haw. Monog. 2 = N. Pseudo-Narcissus.
cuneifolius, Salisb. in Trans. Hort. Soc. i. (1812) 343 = N. Pseudo-Narcissus.
cuneifolius, Haw. Rev. 48 = N. Pseudo-Narcissus.
cyclaminiformis, Haw. Monog. 2. n. 4 = N. Pseudo-Narcissus.
fenestralis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 191 = N. Pseudo-Narcissus.
festalis, Salisb. in Trans. Hort. Soc. i. (1812) 347 = N. Pseudo-Narcissus, serratus.
glaucus, M. Roem. Syn. Ensat. 199 = N. propinquus.
grandiflorus, Salisb. in Trans. Hort. Soc. i. (1812) 344 = N. major.
hexangularis, Herb. Amar. 305.—Hab.?
hispanicus, M. Roem. Syn. Ensat. 200 = N. Pseudo-Narcissus.
incomparabilis, Car. Prod. Fl. Tosc. 616 = N. incomparabilis.
lacinularis, Salisb. in Trans. Hort. Soc. i. (1812) 344 = N. propinquus.
lobularis, Haw. in Phil. Mag. (1830) 181 = N. Pseudo-Narcissus.
longiflorus, Salisb. in Trans. Hort. Soc. i. (1812) 349 = N. tortuosus.
lorifolius, Haw. Monog. 2. n. 7 = N. Pseudo-Narcissus.
lusitanicus, M. Roem. Syn. Ensat. 202 = N. triandrus.
luteus, Herb. Amar. 304, 314. t. 86. f. 83 = N. major.
major, Haw. Rev. 116 = N. major.
maximus, Haw. Monog. 3. n. 23 = N. Pseudo-Narcissus.
minimus, Haw. l. c. 1. n. 1 = N. Pseudo-Narcissus.
minor, Herb. Amar. 299, 315 = N. Pseudo-Narcissus.
moschatus, Haw. Monog. 2 = N. moschatus.
nanus, Haw. l. c. 3. n. 8 = N. Pseudo-Narcissus.
nobilis, Haw. Syn. 327 = N. Pseudo-Narcissus.
obvallaris, Haw. Rev. 120 = N. Pseudo-Narcissus.
odorus, Car. Prod. Fl. Tosc. 616 = N. odorus.
patulus, Salisb. in Trans. Hort. Soc. i. (1812) 348 = N. moschatus.
propinquus, Haw. Monog. iii. n. 22 = N. Pseudo-Narcissus.
Pseudo-narcissus, Haw. l. c. 2 = N. Pseudo-Narcissus.
pumilus, Haw. l. c. 1. n. 3 = N. Pseudo-Narcissus.
pygmaeus, M. Roem. Syn. Ensat. 201 = N. Pseudo-Narcissus.
radians, M. Roem. l. c. 198 = N. Pseudo-Narcissus.
reflexo, M. Roem. l. c. 202 = N. triandrus.
Rudbeckii, M. Roem. l. 201 = N. Pseudo-Narcissus.
rugilobus, Haw. Monog. 3 = N. Pseudo-Narcissus.
Sabiniamus, Herb. Amar. 306 = N. Pseudo-Narcissus.
serratus, Haw. Rev. 114 = N. serratus.
seungularis, M. Roem. Syn. Ensat. 201 = N. Pseudo-Narcissus.
spurius, Haw. Syn. 327 = N. Pseudo-Narcissus.
Telamonius, Haw. l. c. 326 = N. Pseudo-Narcissus.
tortuosus, Haw. Monog. 2 = N. Pseudo-Narcissus.
tubaeformis, Herb. Amar. 308 = N. Pseudo-Narcissus.

AJOVEA, Juss. Gen. 80 (1789) = **Aiouea**, Aubl. (Laurin.).
guianensis, Rafin. Sylva Tellur. 134 = Aiouea tenella.

AJUGA, Linn. Gen. ed. I. 167. n. 466 (1737). *LABIATÆ*, Benth. & Hook. f. ii. 1222.
ABIGA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 85.
BUGULA, Moench, Meth. 381 (1794).
CHAMARIPITYS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 230 (1745).
MOSCHARIA, Forsk. Fl. Aegypt. Arab. 158 (1775).
PHLEBOANTHE, Tausch, in Flora, xi. (1828) 332.
acaulis, *Brocchi, in Bibliot. Ital.* xxix. 83.—Europ.
adulterina, *Wallr. Beitr. Fl. Hercyn.* 231.—Europ.
africana, Pers. Syn. ii. 109 = Teucrium africanum.
alpestris, Dum. Fl. Belg. 43 = genevensis.
alpina, Fries, Mant. iii. 54 = reptans.
alpina, Vill. Hist. Pl. Dauph. ii. 347 = pyramidalis.
alpina, Lin. Mant. 80 = genevensis.
argyrea, *Steitz, in Denkschr. Acad. Wien* i. (1885) 103.—Lycia.
arenosa, Siev. ex Ledeb. Fl. Ross. ii. 149 = oblongata.

AJUGA :—
astolonosa, *Schur, Enum. Pl. Transs.* 545.—Europ.
australis, *R. Br. Prod.* 503.—Austral.
Barrelieri, Tenore, Fl. Nap. t. 340. f. 2 = reptans.
bombycina, *Boiss. Fl. Orient.* iv. 808.—As. Min.
brachystemon, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1883) 192.—Reg. Himal.
bracteosa, *Benth. in Wall. Pl. As. Rar.* i. 59.—Afr. trop.; Ind. or.; As. or.
caespitosa, Schleich. ex Steud. Nom. ed. II. i. 46 = pyramidalis.
capensis, Pers. Syn. ii. 109 = Teucrium africanum.
Chamaecistus, *Ging. ex Benth. Lab. Gen. et Sp.* 696. —Persia.
Chamaepytis, Gmel. Fl. Sic. Prod. ii. 66 = A. Chia.
Chamaepytis, *Schreb. Pl. Vert. Unilab.* 24.—Europ. Orient.
Chia, *Schreb. l. c.* 25 —Oriens.
ciliata, *Bunge, Enum. Pl. Chin. Bor.* 51.—Chin. bor.
comata, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 50. —Lycia.
controversa, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, vi. (1855) 186; xiv. (1863) 44.—Gallia.
crenata, Hochst. ex Benth. in DC. Prod. xii. 597 = bracteosa.
cryptostylon, Lagrèze-Fossat, ex Nym. Consp. 567 = genevensis.
cuneatifolia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 102.—Lycia.
decumbens, Tenore, Fl. Nap. Prod. 'p. xxxiii' = acaulis.
decumbens, *Thunb. Fl. Jap.* 248.—Japon.
decumbens, D. Don, Prod. 108 = macrosperma.
densiflora, Wall. Cat. n. 2081; Benth. in Wall. Pl. As. Rar. i. 606 = bracteosa.
densiflora, Tenore, Fl. Nap. t. 329. f. 2 = reptans.
depressa, Maxim. in Bull. Acad. Pétersb. xxvii. (1883) 192 = parviflora.
dictenarica, Benth. Lab. Gen. et Sp. 695 = australis.
distichа, Roxb. Fl. Ind. iii. 2 = Anisomeles malabarica.
euphrasioides, Boiss. Diagn. Ser. I. vii. 65 = A. Chamaecistus.
flaccida, *Baker, in Journ. Linn. Soc.* xx. (1884) 234.—Madag.
foliosa, Tratt. Arch. Gewaschk. i. 25 = genevensis.
fruticosa, Roxb. Fl. Ind. iii. 1 = Anisomeles malabarica.
furcata, Link, Enum. Hort. Berol. ii. 99 = Crainotome versicolor.
genevensis, *Linn. Sp. Pl.* 561.—Europ.; Orient.
genevensis, With. Arr. Brit. Pl. ed. III. iii. 516 = reptans.
geniculata, Maxim. in Bull. Acad. Pétersb. xxix. (1883) 189 = macrosperma.
glabra, Benth. Lab. Gen. et Sp. 700 = incvigata.
glabra, Presl, Fl. Sicul. Fragm. 26 = A. Chamaepitys.
glabrata, Benth. in Wall. Cat. n. 2041 = Anisomeles ovata.
grosse-serrata, *Franch. & Sav. Enum. Pl. Jap.* ii. 467. —Japon.
Hampeana × *A. Br. & Vatke, in Oestr. Bot. Zeitschr.* xxii. (1872) 357.—Europ.
humilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 114.— Japon.
hybrida × *Kerner, in Oestr. Bot. Zeitschr.* xxiv. (1874) 382.—Austria.
Hyacinthi, Wall. ex Benth. in Wall. Pl. As. Rar. i. 59, 60 = bracteosa.
incisa, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—Japon.
integrifolia, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 108.—Nepal.
intermedia, Boiss. & Orph. ex Boiss. Fl. Orient. iv. 809 = A. Chia.
interrupta, Dulac, Fl. Hautes-Pyr. 407 = genevensis.
Iva, *Schreb. Pl. Vert. Unilab.* 15.—Reg. Mediterr.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 114.— Japon.
laevigata, *Boiss. Fl. Orient.* iv. 804.—Oriens.
lanata, F. Mart. ex Stend. Nom. ed. II. i. 46 = genevensis.
latifolia, Host, Fl. Austr. ii. 119 = genevensis.
latifolia, Schur, Enum. Pl. Transs. 545 = astonoloza.
Laxmanni, *Benth. Lab. Gen. et Sp.* 697.—Oriens.
lobata, *D. Don, Prod. Fl. Nep.* 108.—Reg. Himal.
lucida, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, xxii. (1865) 9.—Corsica.
lupulina, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—China. bor.

AJUGA :—
lycia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 102. —Lycia.
macrosperma, *Wall. ex Benth. in Wall. Pl. As. Rar.* i. 58.—Ind. or.
mesopiana, Boiss. Diagn. Ser. I. vii. 62 = A. Chia.
mollissima, Wall. ex Steud. Nom. ed. II. i. 46 = Anisomeles ovata.
montana, Reichb. Fl. Germ. Exsurs. 314 = genevensis.
nacoza, Mill. ex Schreb. Pl. Vert. Unilab. 26 = A. Iva.
multiflora, *Bunge, Enum. Pl. Chin. Bor.* 51.—China bor.
nana, *Gilib. Fl. Lituan.* 223.—Europ.
Nanti, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, xiv. (1863) 44.—Gallia.
oblongata, *Bieb. Fl. Taur. Cauc. Suppl.* 388.—Caucas.
ophrydis, *Burch. ex Benth. in DC. Prod.* xii. 597.— Afr. austr.
orientalis, *Linn. Sp. Pl.* 561.—Oriens.
ovata, *Wall. Cat.* n. 2036.—Burma.
palaestina, Boiss. Diagn. Ser. I. xii. 62 = A. Chia.
parviflora, *Benth. in Wall. Pl. As. Rar.* i. 59.—Reg. Himal.
pseudoiva, Rob. Cast. ex DC. Fl. Fr. v. 395 = A. Iva.
pseudo-pyramidalis, *Schur, in Verh. Siebenb. Ver Naturw.* iv. (1853) 80.—Europ.
pumila, *Schur, Enum. Pl. Transs.* 545.—Europ.
pygmaea, *A. Gray, in Mem. Am. Acad.* N. S. VI. (1858-9) 402.—Japon.
pyramidalis, Bieb. Fl. Taur. Cauc. ii. 32 = genevensis.
pyramidalis, Huds. Fl. Ang. ed. I. 219 = reptans.
pyramidalis, *Linn. Sp. Pl.* 561.—Europ.
remota, Benth. in Wall. Pl. As. Rar. i. 59 = bracteosa.
repens, Gaeldenst. ex Ledeb. Fl. Ross. iii. 447 = reptans.
repens, Roxb. Fl. Ind. iii. 2 = macrosperma.
reptans, *Linn. Sp. Pl.* 561.—Europ.
robusta, *Baker, in Journ. Linn. Soc.* xx. (1884) 235.— Madag.
rotundifolia, *Willd. & Cat. ex Willd. in Linnaea*, xxx. (1859-60) 130.—Hispan.
rugosa, Host, Fl. Austr. ii. 120 = genevensis.
rupestris, Schleich. ex Steud. Nom. ed. II. i. 46 = pyramidalis.
salicifolia, *Schreb. Fl. Vert. Unilab.* 26.—As. Min.
salicifolia, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 265 = oblongata.
scoparia, Boiss. Diagn. Ser. I. vii. 63 = A. Chamaecistus.
sikkimensis, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 114 in syn. = macrosperma.
simata, *E. Br. Prod.* 503.—Austral.
stolonifera, *Jeanb. & Timb. in Bull. Sc. Phys. & Nat. Toulouse*, iii. (1875-76) 577.—Europ.
Tenorii, Presl, Fl. Sicul. Fragm. p. xxvi. = acaulis.
Thomasii, Maxim. in Bull. Acad. Pétersb. xxix. (1883) 189 = macrosperma.
tridentata, Benth. Lab. Gen. et Sp. 701 = australis.
tridactylites, Ging. ex Benth. l. c. 699 = A. Chia.
trifida, Dulac, Fl. Hautes-Pyr. 406 = A. Chamaepitys.
trifidifolia, Stokes, Bot. Mat. Med. iii. 288 = A. Chamaepitys.
vesiculifera, *Kurd. in Bull. Soc. Nat. Mosc.* xli. (1868) i. 394.—As. temp.
vestita, *Boiss. Diagn.* Ser. I. vii. 62.—Graecia.
virgata, Benth. Lab. Gen. et Sp. 700 = australis.
yesoensis, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* ii. 467.—Japon.

AJUVEA, Stend. Nom. ed. I. 25 (1821) = **Aiouea**, Aubl. (Lauria.).

AKANIA, Hook. f. in Benth. & Hook. f. Gen. i. 409 (1862). *SAPINDACEAE*, Benth. & Hook. f. i. 409. Hillii, *Hook. f. l. c.*—Austral.

AKEA, Stokes, Bot. Mat. Med. ii. 353 (1812) = **Blighia**, Koen. (Sapind.).
solitaria, Stokes, l. c. 354 = B. sapida.

AKEBIA, Decne. in Arch. Mus. Par. i. (1837) 195. t. 13. *BERBERIDEAE*, Benth. & Hook. f. i. 42.
clematifolia, *Sieb. & Zucc. Fl. Jap.* i. 146.—Japon.
lobata, *Decne. in Ann. Sc. Nat. Sér.* II. xii. (1839) 107. —Japon.
quercifolia, *Sieb. & Zucc. Fl. Jap.* i. 146.—Japon.
quinata, *Decne. in Ann. Sc. Nat. Sér.* II. xii. (1839) 107.—China.

AKEESIA, Tussac, Fl. Antill. i. 66 (1808) = **Blighia**, Koen. (Sapind.).
africana, Tussac, l. c. = B. sapida.

AKENTRA, Benj. in Linnaea, xx. (1847) 319 = **Utricularia**, Linn. (Lentib.).
inflata, Benj. l. c.—Surinam.

AKLEMA, Rafin. Fl. Tellur. iv. 114 (1836) = **Euphorbia**, Linn.
nudiflora, Rafin. l. c. = E. nudiflora.

AKYLOPSIS, Lehm. Ind. Sem. Hort. Hamb. (1850) 3; cf. Otto, Hamb. Gartz. vii. (1859) 549 = **Matricaria**, Linn. (Compos.).
suaveolens, Lehm. l. c. = M. discoides.

ALACOSPERMUM, Neck. Elem. i. 167 (1790) = **Cryptotaenia**, DC. (Umbell.).

ALAFIA, Thou. Gen. Nov. Madag. 11 (1809). *APOCYNACEAE*, Benth. & Hook. f. ii. 717.
BLASTOTROPHIS, Didrichs. in Kjoeb. Vidensk. Meddel. (1854) 192.
madagascariensis, Rich. ex A. DC. in DC. Prod. viii. 415 = Thouarsii.
pauciflora, *Radlk. in Bremen Abh.* viii. (1883) 403.— Afr.-trop.
Thouarsii, *Roem. Collect.* i. 203.—Madag.

ALAGOPHYLLA, Rafin. Fl. Tellur. ii. 33 (1836) = **Geonoma**, Linn.
dasyanthes, Rafin. l. c.—Hab.?

ALAGOPTERA, Mart. Gen. Comm. 267 (1842) = ALLAGOPTERA, Nees = **Diplothemium**, Mart. (Palm.).

ALALANTIA, Corr. in Ann. Mus. Hist. Nat. Par. vi. (1804) 388, *err. typ.* = **Atalantia**, Corr. (Rutac.).

ALAMANIA, La Llave & Lex. Nov. Veg. Descr. Fasc. ii. (Orch. Opusc.) 31. *ORCHIDEAE*, Benth. & Hook. f. ii. 526.
punices, *La Llave & Lex. l. c.*—Mexic.

ALAMANNIA, Lindl. Orch. Scel. 14 (1836) = praec.

ALANDINA, Neck. Elem. ii. 456 (1790) = **Moringa**, Juss.

ALANGIUM, Lam. Encyc. i. 174 (1788). *CORNACEAE*, Benth. & Hook. f. i. 949.
ANGOLAM, Adans. Fam. ii. 85 (1763).
KARA-ANGOLAM, Adans. Fam. ii. 84 (1763).
acuminatum, Wight, ex Stend. Nom. ed. II. i. 47 = Lamarckii.
begoniaefolium, Baill. Hist. Pl. vi. 270 = Marlea begoniaefolia.
cordifolium, *Zoll. & Mor. ex Zoll. Syst. Verz. Ind. Archip.* 68.—Malaya.
decapetalum, Lam. Encyc. i. 174 = Lamarckii.
ebracteum, Griff. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 742 = Marlea ebracea.
frutescens, *Zoll. & Mor. ex Zoll. Syst. Verz. Ind. Archip.* 68.—Malaya.
glandulosum, Thw. Enum. Fl. Zeyl. 133 = Lamarckii.
hexapetalum, Lam. Encyc. i. 174 = Lamarckii. Lamarckii, *Thw. Enum. Pl. Zeyl.* 133.—Geront. trop.
latifolium, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 741 = Lamarckii.
Mekillus, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 105. —Ins. Comor.
octopetalum, Blanco, Fl. Filip. ed. II. 210 = Lamarckii.
sundanum, Miq. Fl. Ind. Bat. i. 1. 774 = Lamarckii.
tomentosum, Lam. Encyc. i. 174 = Lamarckii.

ALANIA, Endl. Gen. Pl. 151 (1836). *LILIACEAE*, Benth. & Hook. f. iii. 797.
ALLANIA, Meisn. Gen. 401 (204) (1843).
Cunninghami, *Steud. Nom.* ed. II. i. 47.—Austral.
Endlicheri, Knuth, Enum. Pl. iv. 644 = Cunninghamii.

ALARCONIA, DC. Prod. v. 537 (1836) = **Wychia**, Nutt. (Compos.).
angustifolia, DC. l. c. = W. angustifolia.
helenioides, DC. l. c. = W. robusta.

ALATERNOIDES, Adans. Fam. ii. 304 (1763) = **Phylica**, Linn. (Rhamnac.).

ALATERNUS, [Tourn.] Mill. Gard. Dict. ed. VI. (1752) = **Rhamnus**, Linn.
alpinus, Moench, Meth 344 = R. alpinus.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 3 = R. Alaternus.
balearica, Duham. ex Steud. Nom. ed. I. 685 = R. Alaternus.
cordatus, Rafin. Sylva Tellur. 27.—Hab. ?
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = R. Alaternus.
glandulosus, Rafin. Sylva Tellur. 27 = R. glandulosus.
hispanicus, Hort. ex Steud. Nom. ed. I. 685 = R. Alaternus.
hybridus, Rafin. Sylva Tellur. 27 = R. hybridus.
integrifolius, Rafin. l. c. = R. integrifolius.
lanceolatus, Rafin. l. c. = R. lanceolatus.
latifolia, Mill. Gard. Dict. ed. VIII. n. 4 = R. Alaternus.
ovatus, Rafin. Sylva Tellur. 27.—Hab. ?
Phylica, Mill. Gard. Dict. ed. VIII. n. 1 = R. Alaternus.
prinoides, Rafin. Sylva Tellur. 27 = R. prinoides.
pumilus, Rafin. l. c. = R. pumilus.
rotundifolia, Hort. ex Steud. Nom. ed. I. 685 = R. Alaternus.
sempervirens, Roehl. ex Steud. Nom. ed. I. 493 = Lonicera sempervirens.
variegata, Hort. ex Steud. Nom. ed. I. 685 = R. Alaternus.

ALATHRAEA, Steud. Nom. ed. II. ii. 13, in syn. (1841) = seq.

ALATRAEA, Neck. Elem. i. 374 (1790) = **Phelypaea**, Tourn. (Orobanch.).

ALBERSIA, Kunth, Fl. Berol. ed. II. ii. 144 (1858) = **Amarantus**, Linn.
arenaria, Schur, Enum. Pl. Transs. 566 = Am. Blitum.
ascendens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 142 = Am. ascendens.
Blitum, Kunth, Fl. Berol. ed. II. ii. 144 = Am. Blitum.
caudata, Boiss. Fl. Orient. iv. 992 = Am. viridis.
deflexa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 142 = Am. deflexus.
gracilis, Webb & Berth. Phyt. Canar. iii. 237 = Am. gracilis.
lividus, Kunth, Fl. Berol. ed. II. ii. 144 = Am. Blitum.
oleracea, Kunth, l. c. = Am. Blitum.
polygama, Boiss. Fl. Orient. iv. 991 = Am. polygamus.
polygonoides, Kunth, Fl. Berol. ed. II. ii. 144 = Am. polygonoides.
polystachya, Kunth, l. c. = Am polystachyus.
prostrata, Kunth, l. c. = Am. deflexus.

ALBERTA, E. Mey. in Linnaea, xii. (1838) 258.
RUBIACEAE, Benth. & Hook. f. ii. 109.
laurifolia, *Baker, in Journ. Linn. Soc.* xviii. (1881) 271.—Madag.
magna, *E. Mey. in Linnaea*, xii. (1838) 258.—Afr. austr.
minor, *Baill. Adansonia*, xii. (1876-79) 247.—Madag.

ALBERTIA, Regel & Schmalh. in Act. Hort Petrop. v. (1877) 606. *UMBELLIFERAE.*
communata, *Regel & Schmalh. l. c.* 604.—Turkestan.
margaritifera, *Regel & Schmalh. l. c.* 605.—Turkestan.
palacea, *Regel & Schmalh. l. c.* 606.—Turkestan.

ALBERTINIA, Spreng. Neue Entdeck. ii. 133 (1821). *COMPOSITAE*, Benth. & Hook. f. ii. 227.
SYMBLOMERIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 384.
arborea, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 236 = Vanillosmopsis arborea.
bardanoides, Mart. in DC. Prod. v. 81 = Vernonia crotonoides.
bicolor, DC. Prod. v. 81 = Eremanthus bicolor.

ALBERTINIA :—
bicolor, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 236 = Eremanthus incanus.
brasiliensis, *Spreng. Neue Entdeck.* ii. 133.—Bras.
Candolleana, Gardn. in Hook Lond. Journ. Bot. v. (1846) 235 = Vanillosmopsis erythroppapa.
capitata, DC. Prod. v. 82 = Vanillosmopsis capitata.
Clausseni, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 15 = Vanillosmopsis erythroppapa.
crotonoides, DC. Prod. v. 81 = Vernonia crotonoides.
Elaeagnus, Mart ex DC. l. c. = Eremanthus Elaeagnus.
erythroppapa, *DC. in Deless. Ic. Sel.* iv. t. 5.—Bras.
erythroppapa, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 236 = Vanillosmopsis polycephala.
glomerulata, DC. Prod. v. 82 = Eremanthus glomerulatus.
gonocladus, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 144 = Piptolepis Martiana.
goyazensis, Gardn. in Hook Lond. Journ. Bot. vi. (1847) 425 = Eremanthus goyazensis.
incana, Less. in Linnaea, iv. (1829) 342 = Eremanthus incanus.
incanescens, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 15 = Vanillosmopsis erythroppapa.
mollis, Sch. Bip. ex Baker, l. c. 164 = Eremanthus mollis.
multiflora, Mart. in DC. Prod. v. 82 = Vanillosmopsis polycephala.
obtusata, Mart. l. c. 81 = Eremanthus glomerulatus.
Oleaster, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 145 = Vernonia Oleaster.
paludosa, DC. Prod. v. 81 = Eremanthus glomerulatus.
polycephala, DC. l. c. 82 = Vanillosmopsis polycephala.
rufescta, DC. l. c. 81 = Eremanthus glomerulatus.
salzigna, Mart. in DC. l. c. 82 = Vanillosmopsis polycephala.
stellata, *Gardn. in Hook. Lond. Journ Bot.* v. (1846) 235.—Bras.
verbascifolia, Mart. in DC. Prod. v. 81 = Vernonia crotonoides.

ALBERTINIA, DC. Prod. v. 89, pro parte (1836) = **Vanillosmopsis**, Sch. Bip. (Compos.).

ALBERTISIA, Becc. Malesia, i. (1877) 161. *MENISPERMEAE.*
papuana, *Becc. l. c.* 162 = N. Guin.

ALBIKIA, J. & C. Presl, Rel. Haenk. i. 134 et 34, 35 (1828) = **Hypolytrum**, Rich. (Cyperac.).
schoenoides, C. Presl, l. c. 135 = H. trinervium.
scirpoides, C. Presl, l. c. = H. latifolium.

ALBINA, Giseke, Prael. Linn. 199, 207 (1792) = **Alpinia**, Linn. (Scitam.).

ALBINEA, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zel. 87 (1853) = **Pleurophyllum**, Hook. f. (Compos.).
oresigeneea, Hombr. & Jacquinot, ex Decne. l. c. = P. oresigeneum.

ALBIZIA, Auct. = seq.

ALBIZZIA, Durazz. in Mag. Tosc. iii. iv. (1772) 11; F. Muell. in Journ. Bot. x. (1872) 8. *LEGUMINOSAE*, Benth. & Hook. f. i. 596.
BESPERA, A. Rich. Tent. Fl. Abyss. i. 253 (1847).
SERIANTHES, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 84 in nota.
SERICANDRA, Rafin. Sylva Tellur. 119 (1838).
ZYGIA, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 92.
acrodena, *Miq. Fl. Ind. Bat. i. Suppl.* 281.—Sumatra.
affinis, Fourn. in Ann. Sc. Nat. Sér. IV. xiv. (1860) 271 = amara.
altissima, Hook. f. in Hook. Niger Fl. 332 = Pithecolobium altissimum.
amara, *Boiv. in Encyc. six. Siècle*, ii. 84.—Ind. or. amoenissima, *F. Muell. Fragm.* viii. 165.—Austral.
angolensis, *Welw. in Oliver, Fl. Trop. Afr.* ii. 360.—Afr. trop.
anthelmintica, *Brongn. in Bull. Soc. Bot. Fr.* vii. (1860) 902.—Abyss.
auriculata, *Fourn.* in Ann. Sc. Nat. Sér. IV. xii. (1865) 400.—N. Caled.
basaltica, *Benth. Fl. Austral.* ii. 422.—Austral.

ALBIZZIA :—
Benthamiana, Blume, ex Miq. Fl. Ind. Bat. i. 80 = montana.
bigeminia, F. Muell. Sel. Pl. Indust. Cult. 264 = Pithecolobium bigeminum.
Boivini, *Fourn. in Ann. Sc. Nat. Sér. IV. xiv.* (1860) 378.—Madag.
brachypalyx, *Oliver, Fl. Trop. Afr.* ii. 361.—Afr. trop.
Browni, *Walp. Rep.* i. 926.—Afr. trop.
bubalina, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 129.—Ins. Nicob.
casuscuia, *Benth. Fl. Austral.* ii. 423.—Austral.
Championi, Benth. in Hook. Kew Journ. iv. (1852) 79 = Pithecolobium lucidum.
Charpentieri, *Fourn. in Bull. Soc. Bot. Fr.* xii. (1865) 400.—N. Caled.
clypearia, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 129.—Ins. Nicob.
conjugato-pinnata, *Vatke, in Oestr Bot. Zeitschr.* xxx. (1880) 278.—Afr. trop.
corniculata, F. Muell Sel. Pl. Indust. Cult. 264 = Pithecolobium subcoriaceum.
coriaria, *Welw. in Oliver, Fl. Trop. Afr.* ii. 360.—Afr. trop.
cyclosperma, Benth in Hook. Lond. Journ. Bot i. (1842) 527 = fulgens
dulcis, F. Muell. Sel. Pl. Indust. Cult. 12 = Pithecolobium dulce.
elata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = procera.
elegans, *Kurz, in Pegu Rep. App. B.* 47.—Burma.
elliptica, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 374.—Abyss.
fasciculata, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 129.—Ins. Nicob.
fastigiata, *Oliver, Fl. Trop. Afr.* ii. 361.—Afr. trop.
ferruginea, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 88.—Abyss.
fiorbunda, Fenzl, Reliq. Kotschy (1868). 5. t. 4 = anthelmintica.
Forbesi, *Benth. in Hook. Lond. Journ. Bot* iii. (1844) 92.—Afr. trop.
Fournieri, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 341 = Pithecolobium Fournieri.
fulgens, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844, 85.—Austral.; N. Caled.
glaberrima, *Benth. l. c.* 88.—Guinea.
glabrescens, *Oliver, Fl. Trop. Afr.* ii. 357.—Afr. trop.
glomeriflora, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 74.—Burma.
granulosa, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 85.—N. Caled.
Harveyi, *Fourn. in Bull. Soc. Bot. Fr.* xii. (1865) 399. —Afr. trop.
Henderonia, F. Muell. in Journ. Bot. x. (1872) 10 = Pithecolobium Hendersonii.
heterophylla, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 200.—Burma.
hypoleuca, Oliver, Fl. Trop. Afr. ii. 356 = Harveyi.
Isembergiana, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = A. Julibrissin.
Jaubertiana, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 381.—Madag.
Jiringa, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 200.—Burma.
Julibrissin, *Durazz. in Mag. Tosc.* iii. iv. (1772) 11.—As.; Afr. trop.
Junghuhniana, F. Muell. in Journ. Bot. x. (1872) 9 = Pithecolobium Junghuhnianum.
latifolia, Boiv. in Encyc. six. Siècle, ii. 83 = A. Lebbek
latistipula, F. Muell. Sel. Pl. Indust. Cult. 12 = Lyallona latistipula.
Lebbek, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 87.—Geront. trop.
lebbekioides, *Benth. l. c.* 89 = odoratissima.
lenticellosa, *Benth. in Trans. Linn. Soc. xxx.* (1875) 559.—N. Caled.
littoralis, *Teijsm. & Binn. in Tijdschr. Ned. Ind.* xxix. (1866) 259.—Moluce.
lomatocarpa, DC. Prod. ii. 467 = odoratissima.
lophantha, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 86.—Austral.
lucida, *Benth. l. c.*—Ind. or.; Malaya.
macrothyrsa, *Miq. Fl. Ind. Bat. i. Suppl.* 281.—Java.
malacophylla, Walp. Ann. ii. 467 = ferruginea.
marginata, Buch.-Ham. ex Wall. Cat. n. 5243 = stipulata.

ALBIZZIA :—
micrantha, Boiv. in Encyc. xix. Siècle, ii. 34 = odoratissima.
Milleti, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 89.—China.
mollis, Boiv. in Encyc. xix. Siècle, ii. 33 = A. Julibrissin.
moluccana, *Miq. Fl. Ind. Bat.* i. 26 —Ins. Molucc.
monilifera, *F. Muell. in Journ. Bot.* x. (1872) 10.—Austral.
mostana, *Benth. in Miq. Pl. Jungh.* 267.—Java.
mossambicensis, *Bolle, in Peters, Reise Mossamb. Bot.* i. 4.—Afr. trop.
myriophylla, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 90.—Ind. or.
nefasia, Walp. Ann. iii. 457 = Acacia abyssinica.
nollyvensa, K. Grab. in Wall. Cat. n. 6340 = amara.
Nemu, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = A. Julibrisin.
Neumanniana, *Hoynh. Nom.* ii. 17.—Austral.
obovata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 558. —N. Caled.
odoratissima, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 88.—Ind. or.
paivana, *Fourn. in Ann. Sc. Nat.* Sér. IV. xv. (1861) 172.—N. Caled.
pallida, *Fourn. l. c.* xiv. (1860) 375.—Abyss.
pallida, Harv. in Harv. & Sond. Fl. Cap. ii. 284 = Harveyi.
pedicellata, *Baker, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 568.—Malacca.
Petersiana, *Oliver, Fl. Trop. Afr.* ii. 362.—Afr. trop.
polyphylla, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 372.—Madag.
procera, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 89.—As. trop.; Austral.
pruinosa, F. Muell. in Journ. Bot. x. (1872) 9 = Pithecolobium pruinosum.
purpurascens, Blume, ex Miq. Fl. Ind. Bat. i. 1. 28 = stipulata.
purpurea, *Boiv. ex Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 376.—Ins. Comor.
Quartiniana, *Waip. Ann.* ii. 457.—Abyss.
ramiflora, *F. Muell. Fragm.* ix. 178.—Austral.
retusa, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 90.—Ins. Philipp.
rhombifolia, *Benth. l. c.* 87.—Senegal.
rivularis, Fourn. in Ann. Soc. Sc. Nat. Sér. IV. xv. (1861) 172 = filgens.
roxea, *Carr. in Rev. Hortic.* (1870-71) 490.—Ambor.
rostrata, Blume, Miq. Fl. Ind. Bat. i. 24 = Pithecolobium rostratum.
rotundata, *Blume, l. c.* 20.—N. Guinea.
rubiginosa, *Miq. l. c.* 27.—Java.
rufa, *Benth. in Hook. Lond. Journ. Bot.* v. (1844) 101.—As. trop.; Java.
salajerana, *Miq. Fl. Ind. Bat.* i. 1. 21.—Ins. Salager.
Saman, F. Muell. Sel. Pl. Indust. Cult. 13 = Pithecolobium Saman.
saponaria, *Blume, Miq. Fl. Ind. Bat.* i. 1. 19.—N. Guin.
Schimperiana, *Oliver, Fl. Trop. Afr.* ii. 359.—Afr. trop.
sericocephala, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 91 = amara.
Smithiana, Benth. l. c. i. (1842) 527 = stipulata.
speciosa, *Baker, l. c.* in A. Lebbek.
splendens, *Miq. Fl. Ind. Bat. Suppl.* 280.—Sumatra.
stipulata, *Boiv. in Encyc. xix. Siècle*, ii. 33.—Ind. or.
streptocarpa, *Fourn. in Bull. Soc. Bot. Fr.* xii. (1865) 359.—N. Caled.
subfalcata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 559.—N. Caled.
Sutherlandi, *F. Muell. Fragm.* vi. 22.—Austral.
tenerrima, De Vriese, in Miq. Fl. Jungh. 270 = Acacia pennata.
tengerensis, *Miq. Fl. Ind. Bat.* i. 25.—Java.
Teysmanni, *Kuru, in Journ. As. Soc. Beng.* xlv. (1876) ii. 299, 300.—Burma.
Thozetiana, *F. Muell. ex Benth. Fl. Austral.* ii. 422.—Austral.
tomentella, *Miq. Fl. Ind. Bat.* i. 1. 20.—Java.
Toona, F. Muell. in Journ. Bot. x. (1872) 10 = Pithecolobium grandiflorum.
umbrosa, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 86 = Calliandra umbrosa.
Vaillantii, F. Muell. Fragm. v. 9 = Archidendron Vaillantii.
versicolor, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 359.—Afr. trop.
viridis, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 377.—Madag.
Welwitschii, *Oliver, Fl. Trop. Afr.* ii. 362.—Afr. trop.
Wightii, Wight & Arn. Prod. 274 = amara.
Wightiana, K. Grah. in Wall. Cat. n. 6359 = amara.

ALBRADIA, D. Dietr. Syn. v. 280 (1852) = seq.

ALBRANDIA, Gandich. in Freyc. Voy. Bot. 509 (1826) = Streptium, Lour. (Urtic.).
Gaudichaudi, D. Dietr. Syn. Pl. v. 280 = S. asper !
orientalis, D. Dietr. l. c. = S. asper.
spinosa, D. Dietr. l. c. = Taxotrophis zeylanica.
tinnervosa, D. Dietr. l. c. = Taxotrophis zeylanica.

ALBUCA, Linn. Sp. Pl. ed. II. 438 (1762). *LILIACEÆ*, Benth. & Hook. f. iii. 809.
BRANCIONA, Salisb. Gen. Pl. Fragm. 86 (1866).
FALCONERA, Salisb. l. c. (1866).
NEMASTYLIS, Rafin. Fl. Tellur. iii. 59 (1836).
PALLASTHRA, Salisb. Gen. Pl. Fragm. 86 (1866).
VIRDIUS, Adans. Fam. ii. (19) (1763).
abyssinica, *Dryand. in Vet. Acad. Nya Handl. Stockh.* (1784) 297; *Jacq. Coll.* i. 55.—Afr. austr.
alba, Lam. Encyc i. 76 = altissima.
altissima, *Dryand. in Vet. Acad. Nya Handl. Stockh.* (1784) 293; *Jacq. Coll.* ii. 264.—Afr. austr.
angolensis, *Welw. Apont.* 591; *et in Saund. Ref. Bot.* i. 336.—Angola.
anthericoides, Sweet, Hort. Brit. ed. I. 417 = Ornithogalum suaveolens.
asurea, *Jacq. Coll.* iv. 302.—Afr. austr.
Bainesii, *Baker, in Journ. Linn. Soc.* xiii. (1873) 290. —Afr. trop.
caudata, *Jacq. Coll.* iv. 208.—Afr. austr.
chlorantha, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 351.—Angola.
coarctata, *Dryand. in Vet. Acad. Nya Handl. Stockh.* (1784) 295.—Afr. austr.
Cooperi, *Baker, in Journ. Bot.* xii. (1874) 366.—Afr. austr.
cornuta, [DC. in] Red. Liliac. i. t. 70 = altissima.
elastica, Salisb. Prod. 239 = abyssinica.
elata, Salisb. l. c. 238 = altissima.
Elwesi, Regel, in Act. Hort. Petrop. vi. (1879) 538 = Wakefieldii.
enuvaata, Ker-Gawl. in Bot. Mag. t. 871 = Urginea altissima.
fastigiata, *Dryand. in Vet. Acad. Nya Handl. Stockh.* (1784) 296.—Afr. austr.
fibrosa, *Baker, in Gard. Chron.* (1874) ii. 386.—Afr. austr.
filifolia, Ker-Gawl. Bot. Reg. vii. t. 557 = Urginea filifolia.
flaccida, Baker, in Saund. Bot. Ref. t. 336 = Cooperi.
flaccida, *Jacq. Coll.* iv. 201.—Afr. austr.
fragrans, *Jacq. Hort. Schœnb.* i. 44. t. 84.—Afr. austr.
fugax, Ker-Gawl. Bot. Reg. iv. 311 = Urginea fragrans.
galeata, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 351.—Angola.
Gardeni, Mook. Bot. Mag. t. 4842 = Speirantha convallarioides.
glandulosa, *Baker, in Gard. Chron.* (1875) i. 614.—Afr. austr.; China.
juncifolia, *Baker, l. c.* (1876) i. 534.—Afr. austr.
juncifolia, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 351 = monophylla.
longifolia, *Pisch. ex Steud. Nom.* ed. II. i. 47.—Afr. austr.
lutea, Lam. Encyc. i. 76 = major, minor.
major, *Linn. Sp. Pl.* ed. II. 438.—Afr. austr.
maxima, *Burm. f. Pl. Cap. Prod.* 9.—Afr. austr.
minor, Gledditsch, in Mem. Acad. Berl. (1796) 91 = Uropetalum serotinum.
minor, *Linn. Sp. Pl.* ed. II. 438.—Afr. austr.
monophylla, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 351.—Angola.
myogaloides, *Welw. ex Baker, l. c.* 350.—Angola.
Nelsoni, *N. E. Br. in Gard. Chron.* (1880) ii. 198.—Afr. austr.
nematodes, *Sweet, Hort. Brit.* ed. II. 526.—Afr. austr.
parviflora, *Donn, Hort. Cant.* ed. II. 41.—Afr. austr.
phyanden, Ker-Gawl. in Bot. Mag. t. 1046 = Urginea physodes.
polyphylla, *Baker, in Gard. Chron.* (1874) i. 471.—Afr. austr.
rigida, *Salisb. Prod.* 239.—Afr. austr.
setosa, *Jacq. Coll. Suppl.* 100. t. 14.—Afr. austr.
sordida, Salisb. Prod. 238 = major.
spiralis, *Linn. f. Suppl.* 196.—Afr. austr.
subspicata, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 351.—Angola.
tennifolia, *Baker, in Saund. Ref. Bot.* t. 335.—Afr. austr.
viridiflora, *Jacq. Coll. Suppl.* 98.—Afr. austr.
viscosa, *Linn. f. Suppl.* 196.—Afr. austr.
vittata, Ker-Gawl. in Bot. Mag. t. 1259 = Ornithogalum vittatum.
Wakefieldii, *Baker, in Bot. Mag.* t. 6429.—Afr. trop.
Yerburyi, *Ridl. in Journ. Bot.* xxii. (1884) 370.—Aden.

ALBUCEA, Reichb. Fl. Germ. Excurs. 109 (1830) = Ornithogalum, Linn. (Liliac.).
chlorantha, Reichb. Ic. Fl. Germ. t. 1089 = O. nutans.
nutans, Reichb. Fl. Germ. Excurs. 110 = O. nutans.

ALBUGA, Schreb. Gen. i. 221 (1789) *err. typ.* = **Albuca**, Linn. (Liliac.).

ALBUGOIDES, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 372 = Albuca, Linn. (Liliac.).
erectum, Medic. l. c.—Afr. austr.!

ALCAEA, Hill, Hort. Kew. 218 (1768); Burm. f. Fl. Ind. 149 (1768) = **Althæa**, Linn. (Malvac.).

ALCAMASPINORA, Nicronbs, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 6.—Quid !

ALCANNA, Gaertn. Fruct. ii. 338. t. 110 (1791) = **Lawsonia**, Linn. (Lythrar.).
spinosa, Gaertn. l. c. = L. alba.

ALCANNA, Orph. in Bull. Congr. Bot. Pétersb. 1869 (1870) 138 = **Alkanna**, Tausch (Borag.).

ALCEA (Tourn.) Linn. Syst. ed. I. (1735) = **Althæa**, Tourn. (Malvac.).
acaulis, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 351 = Alth. acaulis.
aegyptiaca, Boiss. Diagn. Ser. II. i. 103 = Alth. lavateræflora.
africana, Lour. Fl. Cochinch. 421 = Alth. africana.
apterocarpa, Boiss. Fl. Orient. i. 830 = Alth. apterocarpa.
Aucheri, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252 = Alth. Aucheri.
Calvertii, Boiss. Fl. Orient. i. 832 = Alth. Calvertii.
denudata, Boiss. l. c. 838.—Armenia.
digitata, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252 = Alth. lavateræflora.
ficifolia, Linn. Sp. Pl. 687 = Alth. ficifolia.
ficifolia, Nym. Consp. 127 = Alth. rosea.
glabrata, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252 = Alth. ficifolia.
Haussknechtii, Boiss. Fl. Orient. i. 830.—Syria.
Heldreichii, Boiss. l. c. 832 = Alth. Heldreichii.
Hohenackeri, Boiss. l. c. 838 = Alth. Hohenackeri.
indica, Burm. f. Fl. Ind. 149.—Java.
hurdica, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252, 253 = Alth. kurdica.
lasiocalycina, Boiss. Fl. Orient. i. 830.—Armenia.
lavateræflora, Boiss. l. c. 832 = Alth. lavateræflora.
litacina, Boiss. l. c. 831 = Alth. apterocarpa.
microchiton, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 254.—As. Min.
pallida, Bess. Enum. Pl. Volh. 26 = Alth. pallida.
palmata, Gilib. Fl. Lituan. i. 65 = Malva Alcea.
pinnatifida, Gilib. l. c. 66 = Malva moschata.
pontica, Jenka, in Oestr. Bot. Zeitschr. xxiii. (1873) 201.—Turcia
procumbens, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 326. t. 17.—Hab.?
remotiflora, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252 = Alth. remotiflora.
rhyticarpa, Trautv. ex Boiss. Fl. Orient. i. 837 = Alth. sulphurea.
rosea, Linn. Sp. Pl. 687 = Alth. rosea.
rosea, Falk, Beitr. ii. 221 = Alth. ficifolia.
rufescens, Boiss. Fl. Orient. i. 838 = Alth. rufescens.
rugosa, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 254 = Ficifolia.
schirasana, Alef. l. c. 252 = Alth. kurdica.
setosa, Alef. l. c. 252 = Alth. setosa.
striata, Alef. l. c. 252 = Alth. setosa.
sulphurea, Alef. l. c. = Alth. sulphurea.

ALCHEMILLA, Linn. Syst. ed. I. (1735). *ROSACEAE*, Benth. & Hook. f. i. 631.
APHANES, Linn. Gen. ed. I. 83 (1737).
PERCEPIER, Dill. ex Moench, Meth. 690 (1794).
abyssinica, *Fres. in Mus. Senkenb.* ii. (1837) 161.—Abyss.
achilleaefolia, Remy, in Ann. Sc. Nat. Sér. III. viii. (1847) 294, lines penultima = pinnata.
acutiloba, Stev. in Bull. Soc. Nat. Mosc xxix. (1856) ii. 178 = vulgaris.
alata, Ruiz & Pav. ex Steud. Nom. ed. II. i. 47 = pinnata.
algida x, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 64.—Europ.
alpestris, F. W. Schmidt, Fl. Boëm. iii. 88 = vulgaris.
alpina, Bieb. Fl. Taur. Cauc. i. 114 = sericea.
alpina, *Linn. Sp. Pl.* 123.—Reg. bor. temp.
ambigua, Jord. ex Nym. Consp. 286 = vulgaris.
Aphanes, Leers, Fl. Herborn. 64 = arvensis.
aphanoides, *Muiis, ex Linn. f. Suppl.* 122.—N. Granat.
argentea, Lam. Fl. Fr. iii. 309 = alpina.
arvensis, *Scop. Fl. Carn.* ed. II. i. 115.—Reg. bor. temp.
barbata, *Presl, Epim. Bot.* 199.—Peruv.
bifurcata, *Hils & Boj ex Baill. in Bull. Soc. Linn. Par.* i. (1882) 842.—Madag.
capensis, *Thunb. Prod. Pl. Cap.* 31.—Afr. austr.
ceylanica, Hook. f. Fl. Brit. Ind. ii. 361, sphalm. = indica.
ciliata, Willd. ex Steud. Nom. ed. II. i. 47, 48 = orbiculata.
conglomerata, F. W. Schmidt, Fl. Boem. iii. 89 = vulgaris.
conjuncta, Bab. in Mag. Nat. Hist. x. (1842) 25 = alpina.
cornucopioides, *Roem. & Schult. Syst.* iii. 471.—Hispan.
cryptantha, *Steud. ex A Rich. Tent. Fl. Abyss.* i. 259.—Abyss.
cuneata, Gaud. Sym. Fl. Helv. 117 = pentaphyllea.
cuneifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 432 = arvensis.
dendroides, *Zoll. in Nat. en Geneesk. Arch. Neerl. Indie*, i. (1844) 484.—Malaya.
diandra, Willd. ex Steud. Nom. ed. II. i. 48 = rupestris.
elongata, *Echl. & Zeyh. Enum.* 264—Afr. austr.
equisetiformis, *Trev. in Bot. Zeit.* vii. (1849) 215.—Venezuela.
erodiifolia, *Wall. Cat. Ind.* ii. 247.—N. Granat.
Eupatoria, *Michx. Fl. Bor. Am.* i. 287.—Am. bor.
fissa, Schum. ex Sturm, Deutschl. Fl. Heft 66 (1850) = pycnalica.
frigida, *Wedd. Chl. And.* ii. 245.—Boliv.
galioides, *Benth. Pl. Hartw.* 134.—N. Granat.
galioides, *Trev. in Bot. Zeit.* vii. (1849) 215.—Venezuela.
Geheebii, *Siegfr. ex Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 70.—Europ.
glaberrima, F. W. Schmidt, Fl. Boem. iii. 89 = vulgaris.
glabra, Saffr. in Flora, xix. (1836) 231 = vulgaris.
glabrata, Tausch, ex Steud. Nom. ed. II. i. 48 = vulgaris.
glaucescens, Wallr. in Linnaea, xiv. (1840) 549 = vulgaris.
Gunae, *Schwarinf. in Verh. Zool.-Bot. Ges. Wien*, xvii. (1866) 668.—Afr. trop.
helvetica x, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 64.—Europ.
heptaphylla, Schleich. ex Steud. Nom. ed. II. i. 48 = pentaphyllea
hirsuta, *H. B. & K. Nov. Gen. et Sp.* vi. 226.—Am. austr.
hybrida, F. W. Schmidt, Fl. Boëm. iii. 89 = vulgaris.
indica, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 3.—Mont. Nilghir. ; Zeylan.
intermedia, [Clairv.] Man. Herb. 48 = alpina.
latifolia, Salisb. Prod. 361 = vulgaris.
Lachenlana, *Griseb. in Goett. Abh.* xxiv. (1879) 124.—Reg. Argent.
lobata, Pall. Reise, ii. 508 = vulgaris.
lycopodioides, *Trev. in Bot. Zeit.* vii. (1849) 215.—Venezuela.
madagascariensis, *O. Hoffm. in Bremen Abh.* vii. (1883) 286.—Afr. trop.
Mandoniana, *Wedd. Chl. And. ii.* 246.—Boliv.
microcarpa, *Boiss. & Reut. Diagn. Hisp.* 11.—Hispan.
minor, Tausch, ex Host, Fl. Austr. i. 319 = pyrenaica.
minor, Huds. Fl. Angl. ed. I. 59 = vulgaris.
monanthos, Stokes, Bot. Mat. Med. i. 296 = arvensis.
monandra, *Jus. Prod. Veg. Ind. Occ.* 28.—Ind. occ.
montana, F. W. Schmidt, Fl. Boëm. iii. 88 = vulgaris.

ALCHEMILLA :—
multicaulis, Tausch, ex Steud. Nom. ed. II. i. 48 = vulgaris.
multifida, *Schoele, in Flora*, xxvi. (1843) 449.—Europ.
nivalis, *H. B. & K. Nov. Gen. et Sp.* vi. 224. t. 560.—Equador.
occidentalis, Nutt. in Torr. & Gray, Fl. N. Am. i. 432 = arvensis.
orbiculata, *Ruiz & Pav. Fl. Per.* i. 68.—Peruv.
palmata, *B. Mey. in Drège, Zwei Pfl. Docum.* 168, nomen.—Afr. austr.
palmata, Gilib. Exercit. ii. 429 = vulgaris.
palmatifida, Tausch, ex Reichb. Fl. Crit. i. 6 = pyrenaica.
paniculata, F. W. Schmidt, Fl. Boëm. iii. 89 = vulgaris.
pectinata, H. B. & K. Nov. Gen. et Sp. vi. 226 = orbiculata.
pedata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 260.—Abyss.
pentaphyllea, *Linn. Sp. Pl.* 123.—Europ.
pinnata, *Ruiz & Pav. Fl. Per.* i. 69.—Peruv.
polyleps, *Wedd. Chl. And.* ii. 246.—Am. austr.
potentilloides, Hils. & Boj. ex Baill. in Bull. Soc. Linn. Par. i. (1882) 848 = madagascariensis.
pratensis, F. W. Schmidt, Fl. Boëm. iii. 88 = vulgaris.
pubescens, Bieb. Fl. Taur. Cauc. i. 114 = vulgaris.
pubescens, Willd. Hort. Berol. t. 79 = pracc. ?
pusilla, *Pomel, Nouv. Mat. Fl. Atl.* 159.—Afr. bor.
pyrenaica, *Dufour, in Ann. Gén. Sc. Phys.* viii. (1821) 298.—Mont. Pyren.
repens, *Presl, Epim. Bot.* 199.—Peruv.
Rothii, *Oliver, Fl. Trop. Afr.* ii. 378.—Afr. trop.
rupestris, *H. B. & K. Nov. Gen. et Sp.* vi. 224.—Quito.
Rutenbergii, *O. Hoffm. in Bremen Abh.* vii. (1882) 336.—Afr. trop.
schnophylla, *Baker, in Journ. Linn. Soc.* xx. (1884) 137.—Madag.
sericata, Reichb. Pl. Crit. i. 6, in obs. = vulgaris.
sericea, *Willd. Enum. Hort. Berol.* 171, in nota.—Orient.
sibbaldiaefolia, *H. B. & K. Nov. Gen. et Sp.* vi. 226. t. 561.—Mexic.
stemmatophylla, *Wedd. Chl. And.* ii. 246. t. 75.—N. Granat.
subsericea, Rent. Compte-rend. Soc. Hallér. (1853–54) 20 = alpina.
sylvestris, F. W. Schmidt, Fl. Boëm. iii. 88 = vulgaris.
tenuicaulis, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 191.—Afr. trop.
trifida, Gilib. Exercit. ii. 430 = arvensis.
tripartita, *Ruiz & Pav. Fl. Per.* i. 68.—Peruv.
truncata, Tausch, ex Steud. Nom. ed. II. i. 48 = vulgaris.
velutina, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 354.—Am. bor.
venusta, *Cham. & Schlecht. in Linnaea*, v. (1830) 573.—Mexic.
villosa, *Tungh. in Nat. en Geneesk. Arch. Neerl. Indie*, ii. (1845) 46.—Java.
vulcanica, *Cham. & Schlecht. in Linnaea*, v. (1830) 572.—Mexic.
vulcanica, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie*, ii. (1845) 687.—Malaya.
vulgaris, *Linn. Sp. Pl.* 123.—Reg. bor. temp.
vulgaris, Wight, Ic. t. 929 = indica.

ALCHORNEA, Sw. Prod. Veg. Ind. Occ. 98 (1788). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 314.
APARISTHMIUM, Endl. Gen. 1119 (1841).
BLEKKERIA, Miq. Fl. Ind. Bat. i. ii. 407 (1859).
CLADODES, Lour. Fl. Cochinch. 574 (1790).
COELENOGYNE, J. Sm. in Trans. Linn. Soc. xviii. (1841) 509. t. 26.
DIDEROTIA, Baill. Adansonia, i. (1860–61) 274.
HERMIA, Humb. & Bonpl. in Willd. Sp. Pl. iv. 809 (1805).
LAUTEMBERGIA, Baill. Etud. Gén. Euph. 451 (1858).
ORFILEA, Baill. l. c. 452 (1858).
PALISYA, Baill. l. c. 503 (1858).
PSEUDOTERWIA, Miq. Fl. Ind. Bat. i. ii. 414 (1859).
SCHOUSBORA, Schum. Beskr. Guin. Pl. 449 (1827).
STIPELLARIA, Benth. in Hook. Kew Journ. vi. (1854) 9.
WETRIA, Baill. Etud. Gén. Euph. 409 (1858).

ALCHORNEA :—
acutifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 171.—Peruv.
Blumeana, *Muell. Arg. l. c.* 167.—Java.
castanesefolia, *A. Juss. Tent. Euph.* 42.—Fl. Orinoco.
columnaris, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 278.—Bras.
cordata, *Benth. in Hook. Niger Fl.* 507.—Afr. trop.
cordata, *Muell. Arg. in DC. Prod.* xv. ii. 901.—Am. trop.
cordifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 170.—Afr. trop.
cotinoides, *Muell. Arg. l. c.* 168.—Madag.
cuneata, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 122 = cuneifolia.
cuneifolia, *Muell. Arg. in DC. Prod.* xv. ii. 900.—Sumatra.
Davidi, *Franch. Pl. Davidian.* i. (1884) 264.—Mongolia.
discolor, *Poepp. & Endl. Nov. Gen.* iii. 19.—Bras.
Duparquetiana, *Baill. Adansonia*, xi. (1873–76) 175.—Afr. trop.
erythrosperma, *Klotzsch, ex Benth. in Hook. Kew Journ.* vi. (1854) 331.—Bras.
floribunda, *Muell. Arg. in Journ. Bot.* i. (1863) 336.—Afr. occ.
Gardneri, *Muell. Arg. in Flora*, xlvii. (1864) 485.—Bras.
glandulosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 18. t. 221.—Bras.
glandulosa, Poit. ex Baill. Etud. Gén. Euph. 446 = latifolia.
grandiflora, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 170.—Venezuela.
grandis, *Benth. Bot. Sulph. Voy.* 164.—Am. trop.
Hilariana, *Baill. Adansonia*, v. (1864–65) 240.—Bras.
hirtella, *Benth. in Hook. Niger Fl.* 507.—Afr. trop.
ilicifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 170.—Austral.
intermedia, *Klotzsch, ex Benth. in Hook. Kew Journ.* vi. (1854) 331.—Bras.
Iricurana, *Casar. Nov. Stirp. Bras. Dec.* 34.—Bras.
javensis, Muell. Arg. in Linnaea, xxxiv. (1865–66) 170 = rugosa.
jasminoides, Casar. Nov. Stirp. Bras. Dec. 15 = nemoralis.
latifolia, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 46 = cordata.
latifolia, *Sw. Prod. Veg. Ind. Occ.* 98.—Ind. occ.
macrophylla, Mart. Herb. Fl. Bras. 371 = cordata.
madagascariensis, *Muell. Arg. in DC. Prod.* xv. ii. 900.—Madag.
Martiana, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 275.—Bras.
megalophylla, *Muell. Arg. in Flora*, xlvii. (1864) 484.—N. Granat.
mollis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 168.—Ind. or.
multispicata, *Muell. Arg. l. c.* 168.—Madag.
nemoralis, *Mart. Herb. Fl. Bras.* 371.—Bras.
parvifiora, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 168.—Ins. Philipp.
parvifolia, *Klotzsch, ex Benth. in Hook. Kew Journ.* vi. (1854) 331.—Bras.
paniculata, Miq. in Linnaea, xxii. (1849) 797 = nemoralis.
platyphylla, *Muell. Arg. l. c.* xxxiv. (1865–66) 171.—Am. centr.
pulcherrima, Klotzsch, ex Benth. in Hook. Kew Journ. vi. (1854) 331 = nemoralis.
puberula, *Klotzsch, l. c.*—Bras.
pycnogyne, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 276.—Bras.
rotundifolia, Moric. ex Baill. Etud. Gén. Euph. 447 = nemoralis.
rufescens, *Franch. Pl. Davidian.* i. (1884) 265.—Mongol.
rugosa, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 170.—Ind. or. ; Malaya.
scandens, *Muell. Arg. l. c.* 170.—Cochinch.
Schomburgkii, Benth. in Hook. Kew Journ. vi. (1854) 330 = Gardneri.
Schomburgkii, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 46.—Am. austr.
sidaefolia, *Baill. Etud. Gén. Euph.* 447.—Bras.
similis, *Muell. Arg. in Flora*, xlvii. (1864) 484.—Mexic.

ALCHORNEA:—
subrotunda, Baill. Etud. Gén. Euph. 447 = glandulosa.
Thouetiana, *Baill. ex Benth. Fl. Austral.* vi. 187.—Austral.
tiliæfolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 168.—Ind. or.
trewioides, *Muell. Arg. l. c.*—China.
triplinervia, *Muell. Arg. in DC. Prod.* xv. ii. 909 = nemoralis.
villosa, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 168. —Malaya.
Zollingeri, Hassk. Cat. Hort. Bogor. Alt. 37; Retzia, 156 = villosa.

ALCHORNEOPSIS, Muell. Arg. in Linnæa, xxxiv. (1865–66) 156. *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 315.
floribunda, *Muell. Arg. l. c.*—Peruv.

ALCHYMILLA, Rupp. Fl. Jen. ed. Hall. 64 (1745) = **Alchemilla**, Linn. (Rosac.).

ALCINA, Cav. Ic. i. 10. t. 15 (1791) = **Melampodium**, Linn. (Compos.).
minor, Cass. in Dict. Sc. Nat. liv. 248 = M. sericeum.
ovatifolia, Jacq. Eclog. Pl. 73 = M. divaricatum.
perfoliata, Cav. Ic. i. 11. t. 15 = M. perfoliatum.

ALCINIA, H. B. & K. Nov. Gen. et Sp. iv. 274 (1820) = præc.

ALCIOPE, DC. Prod. v. 209 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 439.
CELMISIA, Cass. in Dict. Sc. Nat. vii. 356 (1817).
lanata, *DC. Prod.* v. 210.—Afr. austr.
tabularis, *DC. l. c.*—Afr. austr.

ALDAMA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 14 (1824) = **Solerocarpus**, Jacq. (Compos.).
dentata, La Llave, l. c. = S. dentatus.
dentata, Lass. in Linnæa, v. (1830) 154 = S. Schiedeanus.
unisierialis, A. Gray, Pl. Lindh. 228 = S. unisierialis.

ALDEA, Ruiz & Pav. Prod. 19 (1794) = **Phacelia**, Juss. (Hydrophyll.).
circinnata, Willd. Enum. Suppl. 9 = P. circinnata.
magellanica, Willd. ex Steud. Nom. ed. II. i. 48 = Hydrophyllum magellanicum.
peruviana, Ruiz & Pav. Fl. Per. ii. 8. t. 114 = P. peruviana.
pinnata, Ruiz & Pav. l. c. = P. peruviana.

ALDELASTER, C. Koch, in Belg. Hortic. (1861) 324 = ADELASTER, Veitch = **Eranthemum** ! Linn. (Acanth.).

ALDINA, Adans. Fam. ii. 328 (1763) = **Brya**, P. Br. (Legum.).

ALDINA, Endl. Gen. 1322 (1841). *LEGUMINOSÆ*, Benth. & Hook. f. i. 560.
ALIANIA, Benth. in Hook. Lond. Journ. Bot. ii. (1840) 91.
discolor, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 12.—Bras.
heterophylla, *Benth. l. c.* 13.—Bras.
insignis, *Endl. in Walp. Rep.* i. 843.—Guiana.
latifolia, *Benth. in Mart. Fl. Bras.* xv. ii. 13.—Bras.
macrophylla, *Spruce, ex Benth. l. c.* 13.—Bras.

ALDINIA, Scop. Introd. 178 (1777) = **Justicia**, Linn. (Acanth.).

ALDROVANDA, Monti, in Bonon. Sc. & Art. Inst. Comm. ii. 111. 404. t. 19 (1747); Linn. Dis. Chen. 39 (1751). *DROSERACEÆ*, Benth. & Hook. f. i. 655.
verticillata, Roxb. Fl. Ind. ii. 118 = vesiculosa.
vesiculosa, *Linn. Sp. Pl.* 281.—Europ.; Ind. or.

ALDUNATEA, Remy, in C. Gay, Fl. Chil. iii. 320. t. 38 (1849) = **Chaetanthera**, Ruiz & Pav. (Compos.).
chilensis, Remy, l. c. 322. t. 38. f. 1 (= *Oriastrum chilense*).

ALDUNATEA:—
gnaphalioides, Remy, in C. Gay, Fl. Chil. iii. 328 (= *Oriastrum gnaphalicides*).
lycopodioides, Remy, l. c. (= *Oriastrum gnaphalioides*).

ALECTORIDIA, A. Rich. Tent. Fl. Abyss. ii. 447. t. 99 (1851) = **Arthraxon**, Beauv. (Gramin.).
Quartiniana, A. Rich. l. c. 448 = Arth. ciliaris.

ALECTOROCTONUM, Schlecht. in Linnæa, xix. 259 (1847) = **Euphorbia**, Linn.
caracassanum, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 40 = E. caracassana.
cotinifolium, Schlecht. in Linnæa, xix. (1847) 259 = E. cotinifolia.
cotinoides, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 248 = E. cotinoides.
dilatatum, Klotzsch & Garcke, l. c. = E. dilatata.
nudiflorum, Klotzsch & Garcke, l. c. 40 = E. nudiflora.
ovatum, Schlecht. in Linnæa, xix. (1847) 262 = E. Schlechtendalii.
petiolare, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 248 = E. petiolaria.
Riedelianum, Klotzsch & Garcke, in Abh. l. c. (1860) 41 = E. cotinoides.
sanguinolentum, Klotzsch & Garcke, in Monatsb. l. c. (1859) 248 = E. haematodes.
scandens, Klotzsch & Garcke, l. c. = E. scandens.
Scotanum, Schlecht. in Linnæa, xix. (1847) 262 = E. Scotanum.
viride, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 248 = E. viridis.
Willdenowii, Klotzsch & Garcke, in Abh. l. c. (1860) 41 = E. cotinifolia.
Wrightii, Klotzsch & Garcke, in Monatsb. l. c. (1859) 248 = E. Wrightii.
Yavalquahuitl, Schlecht. in Linnæa, xix. (1847) 262 = E. Scotanum.

ALECTOROLOPHUS, Hall. Enum. Stirp. Helv. ii. 628 (1743); Moench, Meth. 440 (1794) = **Rhinanthus**, Linn. (Scrophul.).
alpinus, D. Dietr. Syn. Pl. iii. 553 = R. major.
alpinus, Reichb. f. Ic. Fl. Germ. xx. 66 = R. Crista-galli.
angustifolius, D. Dietr. Syn. Pl. iii. 553 = R. angustifolius.
buccalis, Heynh. Nom. ii. 18 = R. Crista-galli.
Crista-galli, Bieb. Fl. Taur. Cauc. ii. 68 = R. Crista-galli.
hirsutus, All. Fl. Pedem. i. 58 = R. Crista-galli.
major, Reichb. Ic. Fl. Crit. viii. 18. fig. 975; Ic. Fl. Germ. xx. 65 = R. major.
minor, Dum. Fl. Belg. 33 = R. minor.
minor, Reichb. Fl. Germ. Excurs. 858 = R. Crista-galli.
parviflorus, Wallr. Sched. Crit. 318 = R. minor.
pulchellus, Schum. ex Wimm. Fl. Schles. (1841) 277 = R. Crista-galli.
pulcher, Schum. ex Wimm. l. c. ed. III. (1857) 410 = R. Crista-galli.
ramosus, Schur, Enum. Pl. Transs. 580, Index = R. minor.
stenophyllus, Schur, l. c. = R. minor.
Trixago, Bieb. Fl. Taur. Cauc. ii. 68 = Bartsia Trixago.
villosus, Dum. Fl. Belg. 33 = R. Crista-galli.

ALECTRA, Thunb. Nov. Gen. 81 (1784). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 966.
GLOSSOSTYLIS, Cham. & Schlecht. in Linnæa, iii. (1828) 22.
HYMENOSPERMUM, Benth. in Wall. Cat. n. 3968 (1831).
STARBIA, Thou. Gen. Nov. Madag. 7 (1806).
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 118.—Abyss.
asperrima, *Benth. in DC. Prod.* x. 340.—Abyss.
brasiliensis, *Benth. l. c.* 339.—Bras.
capensis, *Thunb. Prod. Pl. Cap.* 97.—Afr. austr.
cordata, Benth. in DC. Prod. x. 339 = Indica.
hirsuta, *Klotzsch, in Peters, Reise Mossamb. Bot.* 221. —Afr. trop.
indica, *Benth. in DC. Prod.* x. 339.—Ind. or.
lurida, *Harv. Gen. S. Afr. Pl.* 250.—Afr. austr.

ALECTRA:—
major, *E. Mey. in Drège, Zwei Pfl. Docum.* 168.—Afr. austr.
melampyroides, *Benth. in DC. Prod.* x. 339.—Afr. austr.
minor, *E. Mey. in Drège, Zwei Pfl. Docum.* 168.—Afr. austr.
orobanchoides, *Benth. in DC. Prod.* x. 340.—Afr. austr.
parasitica, *A. Rich. Tent. Fl. Abyss.* ii. 117.—Abyss.
pedicularioides, *Baker, in Journ. Linn. Soc.* xx. (1884) 214.—Madag.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 116.—Afr. trop.
pumila, *Benth. in DC. Prod.* x. 340.—Afr. austr.
senegalensis, *Benth. l. c.* 339.—Afr. trop.
stricta, *Benth. l. c.* 338.—Bras.
Thomsoni, *Hook. f. Fl. Brit. Ind.* iv. 297.—Ind. or.
Vogelii, *Benth. in DC. Prod.* x. 339.—Afr. trop.

ALECTRYON, Gaertn. Fruct. i. 217. t. 46 (1788) *SAPINDACEÆ*, Benth. & Hook. f. i. 410.
cassecura, *DC. Prod.* i. 617.—Austral.
carinatum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 119.—N. Caled.
connatum, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 299, 307 = Nephelium connatum.
coriaceum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 118.—Austral.
excelsum, *Gaert. Fruct.* i. 216. t. 46.—N. Zel.
ferrugineum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 84.—Ins. Moluce.
glabrum, *Radlk. l. c.*—Ins. Timor.
laeve, *Radlk. l. c.* 119.—Austral.
semicinereum, *Radlk. l. c.* 118.—Austral.
serratum, *Radlk. l. c.*—Austral.
sphaerococcum, *Radlk. l. c.* 119.—Celebes.
subcinereum, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 534.—Austral.
subdentatum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 117.—Austral.
tomentosum, *Radlk. l. c.*—Austral.

ALEGRIA, Moç. & Sesse, ex DC. Prod. i. 516 (1824) = **Luehea**, Willd. (Tiliac.).
candida, Moç. & Sesse, l. c. 517 = L. platypetala..

ALEOME, Neck. Elem. iii. 68 (1790) = **Cleome**, Linn. (Cappar.).

ALEPIDEA, La Roche, Eryng. Hist. 19. t. 1 (1808). *UMBELLIFERÆ*, Benth. & Hook. f. i. 878.
anatymbica, Eckl. & Zeyh. Enum. 289.—Afr. austr.
ciliaris, *La Roche, Eryng. Hist.* 19. t. 1.—Afr. austr.
cordata, E. Mey. in Drège, Zwei Pfl. Docum. 163 = ciliaris.
ferrata, Steud. Nom. ed. II. i. 48, sphalm. = *serrata*.
longifolia, E. Mey. ex Steud. l. c. = anatymbica.
peduncularis, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 220.—Abyss.
serrata, Eckl. & Zeyh. Enum. 289 = ciliaris.
Woodii, *Oliver, in Hook. Ic. Pl.* t. 1452.—Afr. austr.

ALEPYRUM, B. Br. Prod. 252, 253 (1810) = **Centrolepis**, Labill.
monogynum, Hook. f. Fl. Tasm. ii. 77. t. 138 = C. monogyna.
Muelleri, Hook. f. l. c. 78 = C. glabra.
muscoides, Hook. f. l. c. 77 = C. muscoides.
nudicaule, R. Br. Prod. 253 = C. nuda.
pallidum, Hook. f. Fl. Nov. Zel. i. 268 = Gaimardia pallida.
polygynum, R. Br. Prod. 253 = C. polygyna.
Pumilio, R. Br. l. c. = C. polygyna.
viride, T. Kirk, in Journ. Linn. Soc. xix. (1884) 286.—N. Zel.

ALETHRIS, Linn. Ord. Nat. (1764) = seq.

ALETRIS, Linn. Dis. Chen. 15 (1751); (Amoen. Acad. iii. 11). *HAEMODORACEÆ*, Benth. & Hook. f. iii. 677.
STACHYOPOGON, Klotzsch, in Bot. Ergeb. Waldem. Reise, 49. t. 24 (1862).
alba, Michx. Fl. Bor. Am. i. 180 = farinosa.
arborea, Willd. Enum. Hort. Berol. 361 = Dracæna arborea.
aurea, *Walt. Fl. Carol.* 121.—Am. bor.

ALETRIS :—
bifolia, Burm. f. Fl. Cap. Prod. 10 = Lachenalia pendula.
capensis, Linn. Syst. ed. X. 985 = Veltheimia viridiflora.
chinensis, Lam. Encyc. i. 79 = Dracaena terminalis.
cochinchinensis, *Lour. Fl. Cochinch.* 304.—Cochinch.
draconoides, Roxb. ex Roem. & Schult. Syst. vii 347. —Ind. or.
farinosa, Ker-Gawl. in Bot. Mag. t. 1418 = aurea farinosa, *Linn. Sp. Pl.* 319.—Am. bor.
farinosa, Thunb. Fl. Jap 136 = japonica
fragrans, Linn. Sp. Pl. ed. II. 456 = Dracaena fragrans.
glauca, Ait. Hort. Kew. i. 468 = Veltheimia glauca
guineensis, Jacq. Hort. Vind. i. 63. t. 84 = Sansevieria guineensis.
hyacinthoides, Linn. Sp. Pl. ed. II. 456 = Sansevieria guineensis.
japonica, Houtt. Syst. xi. 486. t. 84. f. 2 = Funkia subcordata.
japonica, *Lamb. in Trans. Linn. Soc.* x. (1811) 407 — Japon.
japonica, Thunb. in Nov. Act. Ups. iii. (1780) 208 = Funkia ovata.
linguaeformis, *Burm f. Fl. Cap. Prod* 10.—Afr austr.
littoralis, Koen. ex Steud. Nom. ed. II. i. 49 = Urginia indica ?
pumila, Ait. Hort. Kew. i. 464 (= *Tritomanthe pumila*).
pumila, Donn, Hort. Cant. ed. II. 42 = Dracaena ovata.
punicea, Labill. Pl. Nov. Holl. i. 35. t. 111 = Blandfordia marginata.
sarmentosa, Andr. Bot. Rep. t. 54 (= *Trstomanthe media*).
Uvaria, Linn Mant 367 = Kniphofia aloides.
zeylanica, Mill. Gard. Dict. ed. VIII n. 4 = Sansevieria lanuginosa.

ALEURITES, Forst. Char. Gen. tu t 56 (1776). *EUPHORBIACEAE*, Benth. & Hook. f iii. 299
CAMIRIUM, Gaertn. Fruct ii 194 t 125 (1791).
DRYANDRA, Thunb. Fl. Jap. 18. t. 27 (1784).
ELAEOCOCCA, A. Juss. Euph. Tent. 38 t. 11 (1824).
TELOPEA, Soland. ex Baill. Etud. Gén. Euph. 345 (1858).
VERNICIA, Lour. Fl. Cochinch. 586 (1790).
Ambinux, Pers. Syn. ii. 579 = triloba.
communata, Geisel. Crot. Monogr. 32 = triloba.
cordata, *Steud. Nom.* ed. II. i. 49.—China; Japon.
cordifolia, Steud. l. c. = triloba.
japonica, Blume, in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 120 = cordata.
laccifera, Willd. Sp. Pl. iv. 590 = Croton aromaticus.
lanceolata, Hanco, Fl. Filip. ed. I. 757 = triloba.
lobata, Blanco, l. c. 756 = triloba.
moluccana, Willd. Sp. Pl. iv. 590 = triloba.
peltata, Geisel. Crot. Monogr. 81 = Mallotus peltatus.
pentaphylla, *Walt. Cat.* n. 7259.—Ind. or.
saponaria, *Blanco, Fl. Filip.* ed. II. 519.—Ins. Philipp.
triloba, *Forst. Char. Gen.* 112.—As. trop.; Ins Pacif.
triloba, Blanco, Fl. Filip. ed. I. 755 = triloba.
Vernicia, Hassk. in Flora, xxv. (1842); Beibl. ii. 40 = cordata.
verniciflua, Baill. Hist. Pl. v. 116 = praec.

ALEURITIA, Spach, Hist. Vég. Phan. ix. 360 (1840) = **Primula**, Linn.
cortusoides, Spach, l. c. 361 = P. cortusoides.
farinosa, Spach, l. c. = P. farinosa.
longiflora, Spach, l. c. 362 = P. longiflora.

ALEURODENDRON, Reinw. in Blume, Cat. Gew. Buitenz. 87 (1823), nomen; Syll. Ratisb. ii. (1828) 12 = **Melochia**, Linn. (Sterculiat.).
album, Reinw. ll. cc. = M. Alcurodendron.

ALEVIA, Baill. Etud. Gén. Euph. 508 (1858) = **Bernardia**, P. Br. (Euphorb.).
leptostachya, Baill. l. c. 509 = B. interrupta.

ALEXA, Moq. in DC. Prod. xiii. ii. 168 in nota (1849). *LEGUMINOSAE*, Benth. & Hook. f. i. 556.
ALEXANDRA, Schomb. in Hook. Lond. Journ. Bot. iv. (1845) 12.

ALEXANDRA, Bunge, in Linnaea, xvii. (1843) 120. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 67.
FERGOLATYN, Schenk, in Bull. Phys. Math. Acad. Pétersb. i. (1843) 361.
Lehmanni, *Bunge, in Linnaea* (1843), 130.—Scongar.

ALEXANDRA, Schomb. in Hook. Lond. Journ. Bot. iv. (1845), 12; et Barbac. (1845) 18 = **Alexa**, Moq. (Legum.).
Imperatricis, Schomb. ll. cc.—Guiana.

ALEXIA, Wight, Ic. t. 1293 (1850) = **Alyxia**, R. Br. (Apocyn.).

ALEXIS, Salisb. in Trans. Hort. Soc. i. (1812) 284 = **Amomum**, Linn. (Scitam.).
bifurca, Salisb. l. c. = A. exscapum.
grandiflora, Salisb. l. c. = A. grandiflorum.

ALEXITOXICON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 67 = **Vinostoxiourn**, Moench (Asclep.).

ALFONSIA, H. B. & K. Nov. Gen. et Sp. i. 306 (1815) = **Illaeis**, Jacq. (Palm.).
oleifera, H. B. & K. l. c. 307 = E. melanococca.

ALFREDIA, Cass. in Bull Soc. Philom. (1815) 175 = **Carduus**, Linn. (Compos.).
acanthelepis, Kar & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 304 = C. acanthelepis.
cernua, Cass. in Bull. Soc. Philom. (1815) 175 = C. cernuus.
japonica, Miq. Ann. Mus Bot. Lugd. Bat ii. 185 = Cnicus japonicus.
Karelini, Leieb. Fl. Ross. ii. 754 = C. acanthelepis.
nivea, Kar. & Kir. in Bull Soc. Nat. Mosc. xv. (1842) 295 = C. niveus
squarrosa, 'Kostel Select. Sem. Hort. Bot. Prag. (1838).'—Hab ?
stenolepis, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 452 = C. stenolepis.
suaveolens, Rupr. Sert Tianach. 56 = C. suaveolens.
tianschanica, Rupr. l. c.—As. centr.

ALGA, [Tourn.] Lam. Fl. Fr. iii. 539 (1778) = **Zostera**, Linn. (Naiad.)
marina, Lam l. c. = Z. marina.

ALGAROBIA, Benth. Pl. Hartw. 18 (1839) = **Prosopis**, Linn. (Legum.)
dulcis, Benth. l. c. = P. juliflora.
glandulosa, Torr. & Gray, Fl. N. Am. i. 399 = P. juliflora
juliflora, Heynh. Nom. ii. 18 = P. juliflora.

ALGERNONIA, Baill in Ann. Sc. Nat. Sér. IV. ix. 198 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 330.
TETRAPLANDRA, Baill. l. c. 200 (1858).
brasiliensis, *Baill. l. c.* 199.—Bras.
obovata, *Mmell. Arg. in Mart. Fl. Bras.* xi. ii. 586.—Bras.

ALGUELAGUEN, Feuill. ex Adans. Fam. ii. 505 (1763) = **Sphaeele**, Benth (Labiat.).

ALHAGI, Tourn. ex Adans. Fam. ii. 328 (1763); Desv. Journ. Bot. i. (1813) 120. t. 4. *LEGUMINOSAE*, Benth. & Hook. f. i 512.
MANNA, D. Don, Prod. Fl. Nep. 246 (1825).
camelorum, *Fisch. Hort. Gorenk.* ed. II. (1812) 72.—Orient. Ass. centr.
graecorum, *Boiss. Diagn.* Ser. I. ix. 114.—Graecia.
hardtuchorum, Boiss. & Hausak ex Boiss. Fl. Orient. i. 599 = maurorum.
kirghisorum, Schrenk, Enum. Pl. Nov. 84 = camelorum.
mannifera, Desv. Journ. Bot. i. (1813) 120 = maurorum.
mannifera, Janb. & Spach, Illustr. i. 401 = graecorum.
maurorum, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 397; *Desv. Journ.* i. (1813) 120. t. 4.—Orient.; Ind. or.; Syria.
nagusiensium, DC. Prod. ii. 352 = maurorum.
persarum, Boiss & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 76 = camelorum.
Pseudalhagi, Desv. Journ. Bot. i. (1813) 120 = camelorum.
turcorum, Boiss. Diagn. Ser. I. ix. 113 = camelorum.

ALHAGIA, Reichb. Nom. 158 (1841) = praec.

ALIBERTIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 234. t. 21. *RUBIACEAE*, Benth. & Hook. f. ii. 81.
CORDIERA, A. Rich. l. c. 222. t. 20 (1830).

ALIBERTIA :—
GARAPATICA, Karst. Fl. Colomb. i. 57. t. 26 (1858).
GARDENIOLA, Cham. in Linnaea, ix. (1834) 247.
GENIPELLA, Rich. ex DC. Prod. iv. 443 (1830).
MELANOPSIDIUM, Poit. ex A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 384.
SCEPERNOTHAMNUS, Cham. in Linnaea, ix. (1834) 248.
THIELEDOXA, Cham. l. c. 251 (1834).
edulis, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1830) 234. t. 21.—Guiana.
hexagyna, *Karst. Fl. Colomb.* i. 57.—N. Granat.
jacquinioides, *Griseb. Cat. Pl. Cub.* 123.—Cuba.
Melioana, *Hook. f. in Trans. Linn. Soc.* xxviii. (1872) 520.—Bras.
utilis, A. Rich. in Mém. Soc. Nat. Hist. Par. v. (1830) t. 21 = edulis.

ALIBREXIA, Miers. in Hook. Lond. Journ. Bot. iv. (1845) 506 = **Dolia**, Lindl. (Convolv.).
bryniifolia, Phil. in Linnaea, xxxiii. (1864-65) 209.—Chili.
Gayana, Gandich. Bot. Voy. Bonite, Atl. t. 105.—Chili ?
incana, Phil. ? t Atac. 45.—Chili.
revoluta, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 508 = D. revoluta.
rupicola, Miers, l. c. 506 = D. tomentosa.
tomentosa, Miers, l. c. 508 = D. tomentosa
villosa, Phil. Fl. Atac. 45.—Chili.

ALIBUM, Less. Syn. Comp. 152 (1832) = **Liabum**, Adans. (Compos.).
lsaboides, Less. l. c. = L. lyratum.

ALICABON, Rafin. Sylva Tellur. 56 (1838) = **Physalis**, Linn. (Solanac.).
barbadense, Rafin. l. c. 57 = P. barbadense.
somniferum, Rafin. l. c. 56 = P. somniferum.

ALICASTRUM, P. Br. Hist. Jamaic. 372 (1756) = **Broelmum**, Sw. (Urticac.).

ALICOSTA, Dulac, Fl. Hautes-Pyr. 331 (1867) = **Bartsia**, Linn. (Scrophul.).
alpina, Dulac, l. c. = B. alpina.
spicata, Dulac, l. c. = B. aspera.

ALICTERES, Neck. Elem. ii. 304 (1790) = **Heliotreae**, Linn. (Sterculiac.).

ALIFANA, Rafin. Sylva Tellur. 101 (1838) = **Brachyotum**, Triana (Melast.), etc.
canescens, Rafin. l. c. = Brachyotum canescens.
lutescens, Rafin. l. c. = Miconia lutescens.
montana, Rafin. l. c. = Conostegia montana.
striata, Rafin. l. c. = Brachyotum strigosum.

ALIFANUS, Pluk. ex Adans. Fam. ii. 284 (1763) = **Rhexia**, Linn. (Melast.).

ALINA, Adans. Fam. ii. 84 (1763).—Gen. dub.

ALIPENDULA, Neck. Elem. ii. 89 (1790) = **Spiraea**, Linn. (Rosac.).

ALIPSA, Hoffmgg. [Verz. Orchid. 20 (1840)] ex Linnaea, xvi. (1842) Litt. 228 = **Liparis**, Rich. (Orchid.).
foliosa, Hoffmgg. l. c. = L. reflexa ?

ALISETA, Rafin. New Fl. Am. ii. 22 (1836) = **Arnica** Linn. (Compos.).
plantaginea, Rafin. l. c. = A. alpina.

ALISMA, [Dill.] Linn. Syst. ed. I. (1735). *ALISMACEAE*, Benth. & Hook. f. iii. 1004.
BALDELLIA, Parl. Nov. Gen. Pl. Monocot. 57 (1854).
CALDESIA, Parl. Fl. Ital. iii. 598 (1858).
HELANTHIUM, Engelm. ex Benth. & Hook. f. Gen. iii. 1005 (1883).
NECTALISMA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 156.
acanthocarpum, *P. Muell. Fragm.* i. 28.—Austral.
alpestre, Cosa. in Bull. Soc. Bot. Fr. xi. (1864) 388 = Echinodorus alpestris.
Ancile, Mart. ex Steud. Nom. ed. II. i. 49, nomen = Sagittaria guyanensis.
Andrieuxii, Hook. & Arn. Bot. Beech. Voy. 311 = = Echinodorus ellipticus.

ALISMA :—
angustifolium, Gilib. Fl. Lituan. ii. 224 = Echinodorus ranunculoides.
angustifolium, Hoppe, Taschenb. (1797) 13 = A. Plantago.
apetalum, Buch.-Ham. ex Wall. Cat. n. 4996 = oligococcum.
aphylla, Steud. Nom. ed. II. i. 49 = præc.
arcuatum, Michell, in Bull. Soc. Bot. Fr. i. (1858) 312 = A. Plantago.
Berterii, Spreng. Syst. ii. 163 = Echinodorus rostratus.
Berteroanum, Balb. in Roem. & Schult. Syst. vii. 1605 = Echinodorus rostratus.
Bourgæi, Nym. Syll. 879 = Damasonium Bourgæi.
californicum, *Micheli*, *in A. DC. Monog. Phan.* iii. 34. —Calif.
calophyllum, Wall. Cat. n. 4997 = parnassifolium.
canaliculatum, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1857) 4 = A. Plantago.
cordifolium, Linn. Sp. Pl. 343 = Echinodorus macrophyllus, rostratus.
cordifolium, Thunb. Fl. Jap. 153 = A. Plantago.
cristatum, Wall. Cat. n. 4994 = Sagittaria guyanensis.
Damasonium, Desf. Fl. Atlant. i. 324 = Damasonium Bourgæi.
Damasonium, Linn. Sp. Pl. 343 = Damasonium stellatum.
Damasonium, Willd. Prod. Fl. Berol. 132 = parnassifolium.
difformifolium, *Steud. in Lechl. Berb. Am. Austr.* 58. —Chili.
diversifolium, Gilib. Fl. Lituan. ii. 228 = Elisma natans.
dubium, Willd. Prod. Fl. Berol. 132 = parnassifolium.
echinocarpum, Seub. in Mart. Fl. Bras. iii. 1. 105. t. 18. f. 3 = Sagittaria guyanensis.
ellipticum, Mart. in Roem. & Schult. Syst. vii. 1607 = Echinodorus ellipticus.
ensaandrum, Hochst. ex Michell, in DC. Monog. Phan. iii. 32 = Burmatia ensaandra.
flavum, Linn. Sp. Pl. 343 = Limnocharis flava.
flavum, Thunb. Fl. Jap. 158 = A. Plantago.
floribundum, Seub. in Mart. Fl. Bras. iii. 1. 109 = Echinodorus grandiflorus.
Geyeri, Torr. in Nicollet, Rep. Hydrog. Mississip. (1843) 169 = A. Plantago.
glandulosum, Thw. Enum. Pl. Zeyl. 332 = oligococcum.
gramineum, C. C. Gmel. Fl. Bad. iv. 256 = A. Plantago.
graminifolium, Ehrh. ex Ledeb. Fl. Ross. iv. 40 = A. Plantago.
grandiflorum, Cham. & Schlecht. in Linnæa, ii. (1827) 152 = Echinodorus grandiflorus.
Hamiltonianum, Wall. Cat. n. 4993 = Sagittaria guyanensis.
heterophyllum, Schur, Enum. Pl. Transs. 880, Index = A. Plantago.
humile, Rich. ex Kunth, Enum. Pl. iii. 154 = Echinodorus humilis.
intermedium, *Grsf. ex Voigt, Hort. Suburb. Calc.* 680. —Ind. or.
intermedium, Mart. in Roem. & Schult. Syst. vii. 1609 = Echinodorus Martii.
Kotschyi, Hochst. in Flora, xxvi. (1843) 492 = Limnophyton obtusifolium.
lanceolatum, With. Bot. Arr. Brit. Pl. ed. III. ii. 362 = A. Plantago.
Lappula, Buch.-Ham. ex Kunth, Enum. Pl. iii. 162 = Sagittaria Lappula.
latifolium, Gilib. Fl. Lituan. ii. 223 = A. Plantago.
Lozelii, Eichw. Skizze, 127 = A. Plantago.
macrophyllum, Kunth, Enum. Pl. iii. 151 = Echinodorus macrophyllus.
major, S. F. Gray, Nat. Arr. Brit. Pl. ii. 216 = A. Plantago.
minus, Spreng. Syst. ii. 163 = Damasonium australe.
natans, Linn. Sp. Pl. 343 = Elisma natans.
natans, Pollich, Hist. Pl. Palat. iii. 319 = A. Plantago.
nathburnense, Steud. Nom. ed. II. i. 49 = Echinodorus ellipticus.
nymphæifolium, *Griseb. Cat. Pl. Cub.* 218.—Cuba.
obtusifolium, Thw. Enum. Pl. Zeyl. 332 = Limnophyton obtusifolium.
oderatum, Rafin. Fl. Ludov. 17 = A. Plantago.
oligococcum, *F. Muell. Fragm.* i. 23.—Austral.
palæfolium, Kunth, Enum. Pl. iii. 153 = Echinodorus ellipticus.

ALISMA :—
paniculatum, Stokes, Bot. Mat. Med. ii. 384 = A. Plantago.
parnassifolium, Buch.-Ham. ex Wall. Cat. n. 4997 = calophyllum.
parnassifolium, *Bassi, ex Linn. Syst.* ed. XII. ii. 280.— Europ.; Ind. or.; Afr. trop.; Austral.
parviflorum, Pursh, Fl. Am. bept. i. 252 = A. Plantago.
plantagineum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Plantago.
Plantago, *Linn. Sp. Pl.* 342 (A. Plantago-aquatica). —Reg. temp. bor.; Austral.
polyspermum, Nym. Syll. 879 = Damasonium polyspermum.
pubescens, Buch.-Ham. ex Wall. Cat. n. 4995 = Sagittaria guyanensis.
pubescens, Mart. in Roem. & Schult. Syst. vii. 1607 = Echinodorus ellipticus.
ranunculoides, All. Fl. Pedem. i. 284 = A. Plantago.
ranunculoides, Linn. Sp. Pl. 343 = Echinodorus ranunculoides.
ranunculoides, Willd. Prod. Fl. Berol. 132 = Elisma natans.
reniforme, D. Don, Prod. Fl. Nep. 22 = parnassifolium.
repens, Lam. Encyc. ii. 515 = Echinodorus ranunculoides.
rostrum, Rafin. in Med. Repos. New York, v. (1808) 354 = A. Plantago.
rostratum, Nutt. in Trans. Am. Phil. Soc. v. (1837) 159 = Echinodorus rostratus.
Sagittaria, Stokes, Bot. Mat. Med. ii. 385 = Sagittaria sagittifolia.
sagittifolium, Willd. Sp. Pl. ii. 277 = Limnophyton obtusifolium.
Sprengelii, Rich. ex Kunth, Enum. Pl. iii. 154 = Echinodorus rostratus.
stellatum, Buch.-Ham. ex Wall. Cat. n. 4998 = Sagittaria guyanensis.
stellatum, Lam. Fl. Fr. iii. 77 = Damasonium stellatum.
subalatum, Mart. in Roem. & Schult. Syst. vii. 1609 = Echinodorus subalatus.
subcordatum, Rafin. in Med. Repos. New York, v. (1808) 362 = A. Plantago.
subulatum, Linn. Sp. Pl. 343 = Sagittaria natans.
subulatum, Hill, Veg. Syst. x. 33 = præc.?
taeniæfolium, *Steud. in Lechl. Berb. Am. Austr.* 58.— Chili.
tenellum, Mart. in Roem. & Schult. Syst. vii. 1600 = Echinodorus tenellus.
tribnerve, *Link, ex Steud. Nom.* ed. II. i. 49.—Lusitan.
triviale, Pursh, Fl. Am. Sept. i. 252 = A. Plantago.
verticillatum, Dulac, Fl. Hautes-Pyr. 45 = A. Plantago.
virgatum, Hook. & Arn. Bot. Beech. Voy. 311 = Echinodorus virgatus.

ALISMORCHIS, Thou. Orch. Afr. t. 35 (1822) = **Calanthe**, R. Br. (Orchid.).
Controvis, Steud. Nom. ed. II. i. 49 = C. sylvatica.

ALISSON, Vill. Prosp. 38 (1779) = **Alyssum**, Linn. (Crucif.).

ALISSUM, Neck. Delic. Gallo-Belg. i. 276 (1768) = præc.

ALITUBUS, Dulac, Fl. Hautes-Pyr. 499 (1867) = **Achillea**, Linn. (Compos.).
Millefolium, Dulac, l. c. 500 = Ach. Millefolium.
pyrenaica, Dulac, l. c. = Ach. Ptarmica.
tomentosa, Dulac, l. c. = Ach. tomentosa.

ALIX, Comm. ex DC. Prod. v. 318 (1836) = **Peladia**, Jacq. (Compos.).
salsifolia, Comm. ex DC. l. c. 319 = P. retusa.

ALKANNA, Tausch, in Flora, vii. (1824) 284.
BORAGINEÆ, Benth. & Hook. f. ii. 857.
BAPHORHIZA, *Link*, Handb. ii. 278 (1829).
CAMPTOCARPUS, C. Koch, in Linnæa, xvii. (1843) 304.
CAMPTOLOCARYUM, DC. Prod. x. 92 in syn. (1846).
alba, Boiss. Voy. Espagne, 747 = Nonea alba.
amplexicaulis, A. DC. in DC. Prod. x. 101 = Kotschyana.
arcolata, *Boiss. Diagn.* Ser. I. iv. 46.—As. Min.
Aucheriana, A. DC. in DC. Prod. x. 103 = syriaca, rostellata.
aucheriana, Boiss. Fl. Or. Dec. ii. ? = arcolata.
baccifera, DC. Prod. x. 98 = graca.
bicolor, Boiss. Fl. Or. Dec. ii. ? = Kotschyana.
brachycalon, Boiss. l. c. = Kotschyana.

ALKANNA :—
brassicum, *Boiss. Diagn.* Ser. I. xi. 118.—Persia.
callistachys, Heldr. ex Nym. Consp. 517 = graca.
cappadocica, Boiss. & Bal. Diagn. Ser. II. vi. 132 = megacarpa.
controversa, *A. DC. in DC. Prod.* x. 108.—Hab.? cordifolia, *C. Koch, in Linnæa*, xxii. (1849) 640.— Pontus.
Alagirensis, DC. Prod. x. 96 = Pinardi.
frigida, *Boiss. Diagn.* Ser. I. vi. 94.—Persia.
gallæa, *Boiss. l. c.* Ser. I. xi. 118.—Palæstin.
graca, *Boiss. & Sprun. l. c.* Ser. I. iv. 47.—Græcia.
hirsutissima, *A. DC. in DC. Prod.* x. 101.—Orient.
incana, *Boiss. Diagn.* Ser. I. iv. 46.—As. Min.
Kotschyana, *DC. Prod.* x. 98.—Orient.
Lehmanni, A. DC. in DC. l. c. 588 = tinctoria.
lutea, *A. DC. l. c.* 109.—Europ.
macrophylla, *Boiss. & Heldr. Diagn.* Ser. I. xi. 120.— As. Min.; Syria.
macrorhyncha, *Boiss. & Heldr. l. c.* 120.—Græcia.
Mathioli, Tausch, in Flora, vii. (1824) 284 = tinctoria.
megacarpa, *A. DC. in DC. Prod.* x. 101.—Orient.
noneiformis, Griseb. Spicil. Fl. Rumel. ii. 90 = A. Pulmonaria.
orientalis, *Boiss. Diagn.* Ser. I. iv. 46.—Orient.
Pinardi, *Boiss. l. c.* 48.—As. Min.
primuliflora, *Griseb. Spicil. Fl. Rumel.* ii. 89 — Thracia. Pulmonaria, *Griseb. l. c.* 90.—Macedon.
rostellata, *Boissi Diagn.* Ser. I. xi. 119.—Taurus.
Sartoriana, *Boiss. & Heldr. l. c.* Ser. II. iii. 134.— Græcia.
scardica, *Griseb. Spicil. Fl. Rumel.* ii. 91.—Macedon.
Sieberi, *A. DC. in DC. Prod.* x. 99.—Creta.
strigosa, *Boiss. & Hohenk. in Boiss. Diagn.* Ser. I. iv. 46.—Syria.
syriaca, *Boiss. & Hohenk. Pl. Or. Nov. Doc.* ii. 7.— Syria.
tinctoria, DC. Prod. x. 99 = Sieberi.
tinctoria, *Tausch, in Flora*, vii. (1824) 284.—Orient.
tomentosa, Griseb. Spicil. Fl. Rumel. ii. 88 in note = Onosma mollis.
tubulosa, *Boiss. Diagn.* Ser. I. iv. 48.—As. Min.
versicolor, *Orph. in Bull. Congr. Bot. Pétersb.* 1869 (1870) 138.—Græcia.
viscidula, *Boiss. Diagn.* Ser. I. xi. 118.—Cappadoc. Wagneri, *Bartl. ex Tchihat. Asie Min. Bot.* ii. 107, nomen.—As. Min.

ALKEKENGI, Tourn. ex Hall. Enum. Stirp. Helv. ii. 508 (1742) = **Physalis**, Linn. (Solanac.).
Jacum, Moench, Meth. 479 = P. viscosa.
officinarum, Moench. Meth. Suppl. 177 = P. Alkekengi.
procumbens, Moench, Meth. 478 = P. pubescens.
pubescens, Moench, l. c. = P. peruviana.
villosa, Moench, Meth. Suppl. 178 = P. pubescens.

ALKIBIAS, Rafin. Fl. Tellur. iv. 119 (1836) (Compos.).
hastata, Rafin. l. c. (= *Linozyris hastata*).
macrocnata, Rafin. l. c. = Iphiona imperforiata.
ovata, Rafin. l. c. = Chrysocoma ovata.

ALLAANTHUS, Thw. in Hook. Kew Journ. vi. (1854) 302. t. 9. *URTICACEÆ*, Benth. & Hook. f. iii. 361.
Jasonica, *Benth. & Hook. f. Gen. iii.* 361.—Ins. Phil. zeylanicus, *Thw. in Hook. Kew Journ.* vi. (1854) 302. —Zeylan.

ALLAOPHANIA, Thw. Enum. Pl. Zeyl. 147 (1859). *RUBIACEÆ*, Benth. & Hook. f. ii. 120.
Arnottii, *Hook. f. Fl. Brit. Ind.* iii. 48.—Zeylan.
decipiens, *Thw. Enum. Pl. Zeyl.* 147.—Zeylan.

ALLAGANTHERA, Mart. Hort. Erlang. 69 (1814) = **Alternanthera**, Forsk. (Amarant.).
Forskohlei, Mart. l. c. = Alt. sessilis.

ALLAGAS, Rafin. Fl. Tellur. iv. 58. (1836) = **Halenia**, Borkh. (Gentianac.).

ALLAGOPAPPUS, Cass. in Dict. Sc. Nat. lvi. 21. (1828). *COMPOSITÆ*, Benth. & Hook. f. ii. 384.
chrysocomoides, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 126 = Pulicaria vaviaeha.
dichotomus, Cass. in *Dict. Sc. Nat.* lvi. 23.—Ins. Canar.
viscosissimus, *C. Bolle, in Bonplandia*, vii. (1859) 208.—Ins. Canar.

ALLAGOPTERA, Nees, in Nenwied, Reise nach Bras. ii. 385 (1821) = **Diplothemium**, Mart. (Palm.).
pumila, Nees, l. c. = D. maritimum.

ALLAGOSPERMA, M. Roem. Syn. Pepon. 68 (1846) = **Oxyaponia**, Manso (Cucurb.).
Taymia, M. Roem. l. c. = C. Tayuya.

ALLAGOSTACHYUM, Nees, ex Steud. Nom. ed. II. i. 50 (1840) = **Poa**, Linn. (Gramin.).
laxiclolus, Nees, l. c. = Poa alternans.

ALLAMANDA, Linn. Mant. ii. 146. (1771). *APOCYNACEAE*, Benth. & Hook. f. ii. 690.
Orellia, Aubl. Pl. Gui. i. 270. t. 106 (1775).
angustifolia, *Pohl, Pl. Bras.* i. 73. t. 59.—Bras.
Aublettii, Pohl, l. c. 75 = cathartica.
Blanchetii, *A. DC. in DC. Prod.* viii. 319.—Bras.
brasiliensis, Schott, ex Pohl, Pl. Bras. i. 73 = Schottii.
cathartica, *Linn. Mant.* ii. 214.—Am. trop.
cathartica, Schrad. in Goett. Gelehrt. Anz. (1821) 707 = Schottii.
Donlana, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 11.—Bras.
grandiflora, Lam. Encyc. iv. 601 = cathartica.
Henderson, *Bull, ex Dombr. Hor. Mag.* v. (1866) t. 968.—Guiana.
latifolia, Presl, Bot. Bemerk. 103 = cathartica.
Linnei, Pohl, Pl. Bras. i. 74 = cathartica.
Martii, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 12.—Bras.
nerlifolia, *Hook. Bot. Mag.* t. 4594.—Hab. ?
nobilis, *T. Moore, in Gard. Chron.* (1868) 180, 918.—Bras.
oenotherasfolia, *Pohl, Pl. Bras.* i. 71. t. 57.—Bras.
parviflora, *Presl, Bot. Bemerk.* 103.—Ins Martin.
polyantha, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 12.—Bras.
puberula, *A. DC. in DC. Prod.* viii. 319.—Bras.
Schottii, Hook. Bot. Mag. t. 4851 = cathartica.
Schottii, *Pohl, Pl. Bras.* i. 73. t. 58.—Bras.
setulosa, *Mig. in Linnaea*, xviii. (1844) 748.—Surinam
thevetifolia, *Muell. Arg. in Linnaea*, xxx. (1859-60) 888.—Bras.
verrucosa, Gardn. ex Muell. Arg. in Mart. Fl. Bras. vi. i. 14 = puberula.
verticillata, *Desf. Tabl. Bot. Par.* ed. II. 93.—Hab. ?
violacea, Gardn. & Field, Sert. t. 41 = Blanchetii.
Wardleyana, Lebas, in Rev. Hortic. (1877) 20 = Henderson.

ALLANBLACKIA, Oliver, in Benth. & Hook. f. Gen. i. 980 (1867). *GUTTIFERAE*, Benth. & Hook. f. i. 980.
floribunda, Oliver, in Journ. Linn. Soc. x. (1869) 43.—Afr. trop.

ALLANIA, Benth. in Hook. Lond. Journ. Bot. ii. (1840) 91 = **Aldina**, Endl. (Legum.).
insignis, Benth. l. c.—Guiana.

ALLANIA, Meisn. Gen. 401 (304), (1842) = **Alania**, Endl. (LiHac.).

ALLANTOMA, Miers, in Trans. Linn. Soc. xxx. (1874) 291. *MYRTACEAE*.
salzocarpa, *Miers, l. c.* 296.—Bras.
Burchellana, *Miers, l. c.* 294.—Bras.
corbula, *Miers, l. c.* 297.—Bras.
cylindrica, *Miers, l. c.* 294.—Bras.
dictyocarpa, *Miers, l. c.* 297.—Bras.
fagifolia, *Miers, l. c.* 298.—Guiana.
lineata, *Miers, l. c.* 297.—Bras.
macrocarpa, *Miers, l. c.* 295.—Bras.
multiflora, *Miers, l. c.* 291.—Guiana.
scutellata, *Miers, l. c.* 296.—Bras.
subumosa, *Miers, l. c.* 292.—Guiana.
torulosa, *Miers, l. c.* 293.—Bras.

ALLARDIA, Decne. in Jacquem. Voy. Bot. 87. t. 95, 96 (1841 ?). *COMPOSITAE*, Benth. & Hook. f. ii. 494.
Waldheimia, Kze. & Klr. in Bull. Soc. Nat. Mosc. (1842) 136.
glabra, *Decne. l. c.* 88. t. 96.—Thet.
Hugelii, Sch. Bip. in Pollichia, xx-xxi. (1863) 442 = glabra.
Ilenne, *Klotzi*, *in Sitzb. Akad. Muench.* (1876) 88.—Thet.

ALLARDIA :—
lanata, Kurz, ex C. B. Clarke, Comp. Ind. 144 = tomentosa.
nivea, *Hook. f. & Thoms. ex C. B. Clarke, l. c.* 145.—Thet. occ.
Stoliczkai, *C. B. Clarke, Comp. Ind.* 146.—Thet. occ.
tomentosa, *Decne. in Jacquem. Voy. Bot.* 87. t. 95.—Thet. occ.
tridactylites, Sch. Bip. in Pollichia, xx-xxi. (1863) 442 = glabra.
vestita, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 145.—Reg. Himal.

ALLARDTIA. A. Dietr. in Otto & Dietr. Allg. Gartenz. xx. (1852) 249 = **Tillandsia**, Linn. (Bromel.).
cyanea, A. Dietr. l. c. = T. cyanea.
Potschii, Antoine, in Oestr. Bot. Zeitschr. xxviii. (1878) 56 = T. utriculata.

ALLASIA, Lour. Fl. Cochinch. 84 (1790) Inc. sed.; cf. Benth. & Hook. f. i. 820.
Jobnil, *Manso, Enum.* 86.—Bras.
Payos, *Lour. Fl. Cochinch.* 85.—Afr. or.

ALLAZIA, Manso, Enum. 86. (1836) = prec.

ALLELOTHECA, Steud. Syn. Pl. Gram. 117 (1855) = **Lophatherum**, Brongn. (Gramin.).
Urvillei, Steud. l. c. = L. gracile.

ALLEMANIA, Endl. Gen. 304 in nota (1887) = **Allmania**, R. Br. (Amarant.).

ALLENDEA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 10 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 457.
lanceolata, *La Llave, l. c.*—Mexic.

ALLETOTHECA, Benth. & Hook. f. iii. 1191 (1883) sphalm. = ALLELOTHECA, Steud. = **Lophatherum**, Brongn. (Gramin.).

ALLIARIA, Marsh. ex Repp Fl. Jen. ed. Hall. 75 (1745); DC. Syst. Veg. ii. 488 = **Sisymbrium**, Linn. (Crucif.).
brachycarpa, Bleb. Fl. Taur. Cauc. Suppl 445 = Erysimum brachycarpon.
Fuckii, Kupr. Fl. Canc. 85 = S. Alliaria.
Mathiolı, Kupr. l. c. = S. Alliaria.
officinalis, Andrz. ex DC. Syst. Veg. ii. 489 = S. Alliaria.

ALLIBERTIA, Marion, in Rev. Hort. Bouch. Rhone, Nov. (1869); ex Baker, in Gard. Chron. N. S. xix. (1883) 176 = **Agave**, Linn. (Amaryll.).
intermedia, Marion, ex Baker, l. c. = A. Allbertii.

ALLINUM, Neck. Elem. i. 179 (1790) = **Selinum**, Linn. (Umbell.).

ALLIONIA, Loefl. Iter, 181 (1758); Linn. Syst. ed. X. 890 (1759). *NYCTAGINEAE*, Benth. & Hook. f. iii. 4.
WEHELIA, Loefl. Iter, 180 (1758).
aggregata, Spreng. Syst. i. 884 = Oxybaphus aggregatus.
albida, Walt. Fl. Carol. 84 = Oxybaphus albidus.
Cervantesii, Steud. Nom. ed. II. i. 50 = Oxybaphus Cervantesii.
cuculiata, Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 55 = Oxybaphus cucullatus.
decumbens, Spreng. Syst. i. 884 = Oxybaphus decumbens.
hirsuta, Pursh, Fl. Am. Sept. ii. 728 = Oxybaphus hirsutus.
incarnata, *Linn. Syst.* ed. X. 890.—Am. bor.; Ind. occ.; Am. austr.
linearis, Pursh, l. c. = Oxybaphus angustifolius.
mollugo, Benth. Bot. Voy. Sulph. 44 = Incarnata.
mendocina, *Phil. Sert. Mendoc. Alt.* 41.—Chili.
nyctaginea, Michx. Fl. Bor. Am. i. 100 = Oxybaphus nyctagineus.
ovata, Pursh, Fl. Am. Sept. i. 97 = Oxybaphus floribundus.
violacea, Linn. Syst. ed. X. 890 = Oxybaphus violaceus.

ALLIUM, [Tourn.] Linn. Syst. ed. I. (1735). *LILIACEAE*, Benth. & Hook. f. iii. 802.
Anguinum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 160.

ALLIUM :—
Ascalonicum, Renault, Fl. Dép. Orne, 38 (1800).
Cepa, Tourn. ex Linn. Gen. ed. I. 108 (1737).
Codonoprasum, Reichb. Consp. 66 (1828).
Geroscon, Rafin. Cat. 14 (1824).
Moenchia, Medic. in Act. Acad. Palat. vi. Phys. (1790) 348.
Molium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159.
Moly, Boerh. ex Moench, Meth. 286 (1794).
Nectaroscordum, Lindl. Bot. Reg. t. 1313 (1836).
Ophioscordon, Wallr. Sched. Crit. 129 (1822).
Porrum, Tourn. ex Linn. Syst. ed. I. (1735).
Rhizirideum, Fourr. l. c. 160.
Saturnia, Maratti, Diss. Romul. et Sat. 18. t. 2. (1772).
Schoenoprasum, H. B. & K. Nov. Gen. et Sp. i. 227 (1815).
Scordon, Fourr. l. c. 160.
Trigonea, Parlat. in Ocebio, 161 (1839).
Aglitheis, Endotis, Getuonis, Gynodon, Kalabotis, Kepa, Kromdon, Loncostemon, Maliga, Panstemum, Plexistena, Stelmesus, Stemodoxis, Rafin. Fl. Tellur. ii. 17-21 (1836).
Perisuba, Praskoinon, Rafin. l. c. iv. 87, 89 (1836).
Bearnice, Briseis, Butomissa, Camarilla, Canidia, Hexonychia, Hylogeton, Julus, Molyza, Phyllodolon, Raphione, Schoenissa, Xylorhiza, Salisb. Gen. Pl. Fragm. 66-92 (1866).
achaium, *Boiss. & Orph. ex Boiss. Fl. Orient.* v. 259.—Graecia.
acre, *Jr. Gr. C. Presl, Delic. Prag.* 147.—Sicilia.
acuminatum, *Hook. Fl. Bor.-Am.* ii. 184. t. 196.—Am. bor.
acutangulum, Ambros. Fl. Tyr. Austr. i. 565 = fallax.
acutangulum, Reichb. Ic. Crit. viii. t. 734 = montanum.
acutangulum, Schrad. Sem. Hort. Goett. 1808 = angulosum.
acutiflorum, Bové, ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 192 = paniculatum.
acutiflorum, Loisel. in Desv. Journ. Bot. ii. (1809) 279 = multiflorum.
acutum, Spreng. Pugill. i. 28 = A. Schoenoprasum.
adscendens, Kunth, Enum. Pl. iv. 886 = A. Ampeloprasum.
aestivum, Tineo, Pl. Rar. Sicil. 89 = sphaerocephalum.
affine, Boiss. & Heldr. Diagn. Ser. II. iv. 114 = vineale.
affine, Ledeb. Fl. Ross. iv. 166 = margaritaceum.
africanum, F. G. Dietr. Gartenlex. Nachtr. i. 160 = odoratissimum.
agrum, Kunth, Enum. Pl. iv. 455 = Ornithogalum afrum.
Aitchisoni, *Boiss. Fl. Orient.* v. 248.—Afghan.
Akaka, *Cosn. ex Roem. & Schult. Syst.* vii. 1132.—Sibir.
Akaka, Regel, in Act. Hort. Petrop. iii. (1875) 243 = karataviense.
alaviense, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) i. 448.—Turkestan.
alatum, Schreb. ex Roth, in Roem. Arch. i. iii. (1798) 40 = cernuum.
Alberti, *Regel, in Act. Hort. Petrop.* v. (1877) 682.—Turkestan.
albescens, Gass. Enum. Pl. Inarim. 888 = A. Ampeloprasum.
albidum, Bieb. Fl. Taur. Cauc. iii. Suppl. 260 = flavescens.
albidum, *Fisch. Cat. Gorensk.* (1812) 10.—Orient; Sibir.
albidum, J. & C. Presl, Delic. Prag. 146 = paniculatum.
album, Hort. ex [Delar. in] Red. Lil. v. 360 = ollatum.
album, Sauti, Viaggi, 852. ? = neapolitanum.
album, Spreng. Syst. ii. 38 = pendulinum.
Alexeianum, *Regel, in Act. Hort. Petrop.* iii. (1875) 244.—Turkestan.
alibile, *A. Rich. Tent. Fl. Abyss.* ii. 330.—Abyss.
alpicolum, *Verl. in Bull. Soc. Stat. Iabre*, xiv. (1872) 207.—Gallia.
alpinum, Regel, in Act. Hort. Petrop. iii. ii. (1875) 181 = narcissiflorum.
alpinum, Hegetschw. Fl. Helv. 329 = A. Schoenoprasum.
altaicum, Pall. Reise, ii. 727 = fistulosum.
altissimum, *Regel, in Act. Hort. Petrop.* viii. (1884) 666.—Bokhara.
Amanesi, Bor. Fl. Cent. Fr. ed. III. ii. 681. in obs. = paniculatum.
ambiguum, DC. Fl. Fr. iii. 230 = ochroleucum.
ambiguum, Sibth. & Sm. Fl. Graec. t. 237 = roseum.
ambiguum, Tenore, Link, in Linnaea, ix. (1834) 189 = neapolitanum.

ALLIUM :—
amblyophyllum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 510.—Songaria.
amethystinum, Tausch, in Syll. Ratisb. ii. 256 = A. Ampeloprasum.
ammophilum, Heuff. in Flora, xxvii. (1845) 241 ; et Verh. Zool.-Bot. Ges. Wien, viii. (1858) 210 = angulosum.
amoenum, G. Don, in Mem. Wern. Soc. vi. (1827) 75 = roseum.
Ampeloprasum, Boiss. Voy. Espagne (ii. 615) = polyanthum, baeticum.
Ampeloprasum, *Linn. Sp. Pl.* 295.—Europ. ; Orient.
Ampeloprasum, Thunb. Prod. Pl. Cap. 68 = rotundum.
amphibolum, *Ledeb. Fl. Alt.* ii. 5. t. 857.—Sibir.
amplectens, Torr. in Pacif. Rail. Rep. iv. 148 = attenuifolium.
anceps, *Kellogg, in Proc. Calif. Ac.* ii. (1868) 109. t. 32.—Calif.
Andersonii, G. Don, in Mem. Wern. Soc. vi. (1827) 59 = senescens.
andicolum, *Regel, in Act. Hort. Petrop.* iii. (1875) 211. —Peruv.
angolense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 262.—Afr. trop.
angulare, *Pall. Reise*, iii. 293.—Sibir. balc.
angulatum, Pali. l. c. 108. = prace.
angulosum, All. Fl. Pedem. ii. 158 = albidum.
angulosum, DC. Fl. Fr. iii. 222 = fallax.
angulosum, Krock. Fl. Sil. i. 324 = acutangulum, fallax, senescens.
angulosum, *Linn. Sp. Pl.* 300.—Sibir.
angulosum, Lour. Fl. Cochinch. 303 = nilgiricum.
angulosum, Pursh, Fl. Am. Sept. i. 228 = reticulatum.
angustum, *G. Don, in Mem. Wern. Soc.* vi. (1827) 93. —Sibir.
anisopodium, Ledeb. Fl. Ross. iv. 183 = tenuissimum.
apeninum, Huet, ex Nym. Consp. 740 = paniculatum.
appendiculatum, Ram. ex Pers. Syn. i. 356 = suaveolens.
approximatum, Gren. & Godr. Fl.Fr. iii. 200 = sphaerocephalum.
arenarium, Linn. Sp. Pl. 297 = A. Scorodoprasum.
arenarium, Sadler, Fl. Pesth. ex Reichb. Fl. Germ. Excurs. 40 = sativum.
arenarium, [Wahlenb. in] Svensk. Bot. t. 691 = vineale.
arenarium, Thunb. [Fl. Jap. 132] = japonicum.
armenum, Boiss. & Kotschy, ex Boiss. Fl. Orient. v. 254 = flavum.
armerioides, *Boiss. Diagn.* Ser. I. vii. 116.—Assyria.
arvense, Guss.-Ind. Sem. (1826) = sphaerocephalum.
ascalonicum, *Linn. Amoen. Acad.* iv. 454.—Ubique cult.
ascalonium, Bory & Chaub. Fl. Pelop. 21 = gomphrenoides.
ascendens, Tenore, Syll. 164 = A. Ampeloprasum.
Aschersonianum, *Barbey, Herbor. au Levant*, 163. t. 4. —Oriens.
asperum, G. Don, in Mem. Wern. Soc. vi. (1827) 42 = carinatum.
atropurpureum, Waldst. & Kit. Ic. Pl. Rar. Hung. i. 16. t. 17 = stenopetalum.
atrorubens, *S. Wats. in Bot. King, Exp.* 352.—Calif.
atrosanguineum, *Ker. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 508.—As. med.
atroviolaceum, Boiss. Diagn. Ser. I. vii. 112 = A. Ampeloprasum.
atroviolaceum, Hornem. ex Steud. Nom. ed. II. i. 150 = sphaerocephalum.
attenuatum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 110. f. 38.—Calif.
attenuifolium, *Kellogg, l. c.* 110. f. 34.—Calif.
Auctori, *Boiss. Diagn.* Ser. I. vii. 116.—Persia.
aureum, Lam. Fl. Fr. iii. 260 = A. Moly.
auriculatum, *Kunth, Enum. Pl.* iv. 418.—Ind. or.
australe, Gmn. ex Kunth, l. c. 404 = paniculatum.
auwerum, Bunge, in Mém. Sav. Étr. Acad. Pétersb. vii. (1851) 510 ; Reliq. Lehm. 334. n. 1866 = arvicolatum.
auwerum, Ledeb. Fl. Alt. ii. 18 = caeruleum.
Babingtonii, Borrer, in Engl. Bot. t. 2906 = A. Scorodoprasum.
backhitaricum, Boiss. Fl. Orient. v. 585 = bakhtiaricum.
Backhousianum, *Regel, Gartenfl.* (1888) 218.—Reg. Himal.
baeticum, *Boiss. Diagn.* Ser. I. vii. 118.—Hispan.
balcanense, Willd. Enum. Hort. Berol. 350 = senescens.
Bakeri, *Regel, in Act. Hort. Petrop.* iii. (1875) 141.—China ; Japon.
bakhtiaricum, *Regel, l. c.* 232.—Persia.

ALLIUM :—
Balansae, *Boiss. Fl. Orient.* v. 268.—Pontus.
Banerianum, *Baker, in Gard. Chron.* (1874) ii. 181.—Aegypt. ; Turcia !
Baumannianum, C. Koch, in Linnaea, xxii. (1849) 241 = rotundum.
Beckerianum, Regel, Ind. Sem. Hort. Petrop. (1860) 30 = tataricum.
Bertolonii, De Not. Rep. Fl. Lig. 414 = A. Ampeloprasum.
bidentatum, Fisch. ex Regel, in Act. Hort. Petrop. iii. (1875) 157 = tenuissimum.
Bidwelliae, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 281.—Am. bor.
bifolium, *Willd. ex Schult. f. Syst. Veg.* vii. 1117 = decipiens.
Bigelovii, *S. Wats in Bot. King, Exped.* v. 487. t. 38. f. 8 & 9.—N. Mexic.
bisceptrum, *S. Wats. l. c.* 351 & 487. t. 37. f. 1–8.—Calif.
bodinieranum, Fisch. ex Regel, in Act. Hort. Petrop. iii. (1875) 157 = tenuissimum.
Bidwelliae, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 281.—Am. bor.
bifolium, *Willd. ex Schult. f. Syst. Veg.* vii. 1117 = decipiens.
Bigelovii, *S. Wats in Bot. King, Exped.* v. 487. t. 38. f. 8 & 9.—N. Mexic.
bisceptrum, *S. Wats. l. c.* 351 & 487. t. 37. f. 1–8.—Calif.
Andersonii, G. Don, in Mem. Wern. Soc. vi. (1827) 59 = senescens.
andicolum, *Regel, in Act. Hort. Petrop.* iii. (1875) 211. —Peruv.
angolense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 262.—Afr. trop.
angulare, *Pall. Reise*, iii. 293.—Sibir. balc.
angulatum, Pali. l. c. 108. = prace.
angulosum, All. Fl. Pedem. ii. 158 = albidum.
angulosum, DC. Fl. Fr. iii. 222 = fallax.
angulosum, Krock. Fl. Sil. i. 324 = acutangulum, fallax, senescens.
angulosum, *Linn. Sp. Pl.* 300.—Sibir.
angulosum, Lour. Fl. Cochinch. 303 = nilgiricum.
angulosum, Pursh, Fl. Am. Sept. i. 228 = reticulatum.
angustum, *G. Don, in Mem. Wern. Soc.* vi. (1827) 93. —Sibir.
anisopodium, Ledeb. Fl. Ross. iv. 183 = tenuissimum.
apeninum, Huet, ex Nym. Consp. 740 = paniculatum.
appendiculatum, Ram. ex Pers. Syn. i. 356 = suaveolens.
approximatum, Gren. & Godr. Fl.Fr. iii. 200 = sphaerocephalum.
arenarium, Linn. Sp. Pl. 297 = A. Scorodoprasum.
arenarium, Sadler, Fl. Pesth. ex Reichb. Fl. Germ. Excurs. 40 = sativum.
arenarium, [Wahlenb. in] Svensk. Bot. t. 691 = vineale.
arenarium, Thunb. [Fl. Jap. 132] = japonicum.
armenum, Boiss. & Kotschy, ex Boiss. Fl. Orient. v. 254 = flavum.
armerioides, *Boiss. Diagn.* Ser. I. vii. 116.—Assyria.
arvense, Guss.-Ind. Sem. (1826) = sphaerocephalum.
ascalonicum, *Linn. Amoen. Acad.* iv. 454.—Ubique cult.
ascalonium, Bory & Chaub. Fl. Pelop. 21 = gomphrenoides.
ascendens, Tenore, Syll. 164 = A. Ampeloprasum.
Aschersonianum, *Barbey, Herbor. au Levant*, 163. t. 4. —Oriens.
asperum, G. Don, in Mem. Wern. Soc. vi. (1827) 42 = carinatum.
atropurpureum, Waldst. & Kit. Ic. Pl. Rar. Hung. i. 16. t. 17 = stenopetalum.
atrorubens, *S. Wats. in Bot. King, Exp.* 352.—Calif.
atrosanguineum, *Ker. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 508.—As. med.
atroviolaceum, Boiss. Diagn. Ser. I. vii. 112 = A. Ampeloprasum.
atroviolaceum, Hornem. ex Steud. Nom. ed. II. i. 150 = sphaerocephalum.
attenuatum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 110. f. 38.—Calif.
attenuifolium, *Kellogg, l. c.* 110. f. 34.—Calif.
Auctori, *Boiss. Diagn.* Ser. I. vii. 116.—Persia.
aureum, Lam. Fl. Fr. iii. 260 = A. Moly.
auriculatum, *Kunth, Enum. Pl.* iv. 418.—Ind. or.
australe, Gmn. ex Kunth, l. c. 404 = paniculatum.
auwerum, Bunge, in Mém. Sav. Étr. Acad. Pétersb. vii. (1851) 510 ; Reliq. Lehm. 334. n. 1866 = arvicolatum.
auwerum, Ledeb. Fl. Alt. ii. 18 = caeruleum.
Babingtonii, Borrer, in Engl. Bot. t. 2906 = A. Scorodoprasum.
backhitaricum, Boiss. Fl. Orient. v. 585 = bakhtiaricum.
Backhousianum, *Regel, Gartenfl.* (1888) 218.—Reg. Himal.
baeticum, *Boiss. Diagn.* Ser. I. vii. 118.—Hispan.
balcanense, Willd. Enum. Hort. Berol. 350 = senescens.
Bakeri, *Regel, in Act. Hort. Petrop.* iii. (1875) 141.—China ; Japon.
bakhtiaricum, *Regel, l. c.* 232.—Persia.

ALLIUM :—
callimischon, Link, in Linnaea, ix. (1834) 140 = paniculatum.
callistemon, Webb, ex Regel, in Act. Hort. Petrop. iii. (1875) 189 = flavum.
calyptratum, *Boiss. Diagn.* Ser. I. xiii. 30.—Syria.
Cambessi, De Not. Rep. Fl. Lig. 412 = rotundum.
campanulaeflorum, *Gcy. ex Hook. Kew Journ.* vii. (1855) 375.—Am. bor.
campanulatum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 281.—Am. bor.
campestre, Schleich. ex Steud. Nom. ed. II. i. 51 = vineale.
canadense, Hort. ex Schult. f. Syst. vii. 1007 = Purshii.
canadense, Herbb. ex Schult. f. l. c. 1095 = mutabile.
canadense, *Linn. Sp. Pl.* 1295.—Am. bor.
canadense, Michx. Fl Bor. Am. i. 191 = striatum.
candidissimum, Cav. Lecc. 994 = neapolitanum.
candidum, *C. Presl, Fl. Sic.* 42.—Sicilia.
canescens, Hort. ex Schult. f. Syst. vii. 1067 = glaucum.
capillare, Cav. Icones, iii. 4. t. 208 = moschatum.
capillare, Desf. Tabl. ed. 1. 27 = pusillum.
capitellatum, *Boiss. Diagn.* Ser. I. vi. 118.—Persia ; Afghan.
cappadocicum, *Boiss. & Balansa, ex Boiss. Fl. Orient.* v. 241.—Cappadoc.
cardiostemon, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 48.—Persia.
carduchorum, C. Koch, in Linnaea, xxii. (1849) 237 = Aucheri.
caricifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 854 = Pallasi.
carinatum, *Regel, in Act. Hort. Petrop.* vi. (1879) 582. —Turkestan.
carnatum, Habl. Taur. 152 = flavum.
carinatum, *Linn. Sp. Pl.* ed. 2. 297.—Europ.
carinatum, Schult. Fl Stary. 68 = arenarium.
carinatum, Sm. Engl. Bot. t. 1658 = oleraceum.
Carmeli, *Boiss. Diagn.* Ser. I. xiii. 28.—Palaestin.
carneum, Tenore, Fl. Nap. 159 = roseum.
carneum, Hort. ex Schult. f. Syst. vii. 1028 = stellricum
carneum, Targ.-Tozz. Ist. Bot. ed. II. i. t. 6 = ambigium.
carneum, Willd. Enum. Hort. Berol. 359 = ascalonicum.
carolinianum, [*Delar. in*] *Red. Lil.* t. 101.—Carolina.
caspium, *Buch. Fl. Taur. Cauc.* i. 265 ; iii. 260.—Ross. austr. ; Turkest. ; Sibir.
cassium, *Boiss. Diagn.* Ser. I. xiii. 28.—Taurus ; Syria.
caucasicum, Bieb. Fl. Tau. Cauc. iii. 258 = globosum.
caucasus, Turra, ex Sacc. in Atti Ist. Venet. Ser. III. ix. (1863–64) 487 = globosum.
Cepa, Banks, ex Schult. f. Syst. vii. 1008 = proliferum.
Cepa, Falk, Beitr. ii. 167 = fistulosum.
Cepa, *Linn. Sp. Pl.* ed. I. 300.—Persia ; Beluchist.
Cepa, Willd. ex Ledeb. Fl. Ross. iv. 169 = galanthum.
cepaeforme, G. Don, in Mem. Wern. Soc. vi. (1827) 31 = A. Cepa.
cepaeum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113 = A. Cepa.
ceratophyllum, Bess. ex Schult. f. Syst. vii. 1039 = fistulosum.
cernuum, Hort. ex Schult. f. l. c. 1081 = prostratum.
cernuum, *Roth, in Roem. Archiv*, i. ii. 40.—Am. bor.
Chamaemoly, *Linn. Sp. Pl.* 301.—Reg. Mediterr.
Chamaemoly, Viviani, Fl. Lib. Spec. 19 = umbilicosum.
chamaespathum, *Boiss. Diagn.* Ser. I. vi. 118.—Graecia.
chilense, *Regel, in Act. Hort. Petrop.* iii. ii. (1875) 218.—Chili.
chinense, G. Don, in Mem. Wern. Soc. vi. (1827) 59 = Thunbergii.
chinense, Maxim. Prim. Fl. Amur. 284 = odorum.
chionanthe, Boiss. Fl. Orient. v. 272 = sebdanense.
chloroneurum, *Boiss. l. c.* 247.—Persia.
chlorotepalum, Boiss. Diagn. Ser. I. xiii. 33.—Syrium.
christianum, Lall. ex Regel, in Act. Hort. Petrop. iii. ii. (1875) 246 = decipiens.
chrysanthum, Schur, Enum. Pl. Transs. 673 = globosum.
Christii, *Yamba, in Linnaea*, xxx. (1859–60) 604.—Europ. austr.
chrysanthemum, *Boiss. & Reut. ex Boiss. Fl. Orient.* v. 280.—Assyria.
chrysanthum, *Regel, in Act. Hort. Petrop.* iii. (1875) 91. —China.
ciliare, [Delar. in] Red. Lil. vi. t. 311 = subhirsutum.

ALLIUM :—

ciliatum, C. Koch, in Linnaea, xix. (1847) 11 = sphaerocephalum.
ciliatum, Cyr. Pl. Rar. Neap. Fasc. ii. 18 = subhirsutum.
cilicicum, *Boiss. Diagn.* Ser. I. vii. 115.—As. Min.
circinnatum, *Siebor, Reise Kreta*, ii. 216.—Graecia.
cirrhosum, Raeusch. Nom. ed. III. 74 = roseum.
cirrhosum, Vand. Fasc. Pl. 11 = flavum.
clathratum, *Ledeb. Fl. Alt.* ii. 18.—Sibir. alt.
Clusianum, Bory, Fl. Pelop. 21 = circinnatum.
Clusianum, Retz. in Willd. Sp. Pl. ii. 79 = subvillosum.
colchicifolium, *Boiss. Diagn. Ser.* II. iv. 112.—Armenia.
collinum, *Dougl. ex S. Wats. in Proc. Am. Acad.* xiv. (1879) 28.—Am. bor. occ.
collinum, Guss. in Tenore, Syll. 169 = paniculatum.
coloratum, Dulac, Fl. Hautes-Pyr. 110 = A. Schoenoprasum.
coloratum, Spreng. Syst. ii. 38 = flavum.
commune, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 6; ed. II. 68 (1827) = A. Cepa.
commutatum, Guss. Enum. Pl. Inarim. 17 = rotundum.
compactum, Thuil. Fl. Par. 167 = vineale.
complanatum, Boreau, Not. xxxv. p. 12 = oleraceum.
condensatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 121.—Sibir. baical.
confertum, Fisch. Cat. (1821).—Cf. G. Don, in Mem. Wern. Soc. vi. (1827) 68 = deflexum, strictum, senescens.
confertum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 127 = roseum.
congestum, G. Don, in Mem. Wern. Soc. vi. (1827) 66 = Stellerianum.
consanguineum, *Kunth, Enum. Pl.* iv. 431.—Reg. Himal.
consimile, Jord. ex Gren. & Godr. Fl. Fr. iii. 208 = carinatum.
contortum, Stokes, Bot. Mat. Med. ii. 229 = A. Scorodoprasum.
controversum, Costa, Suplem. Catal. 62 = pyrenaicum.
controversum, Schrad. Cat. Goett. ex Willd. Enum. Hort. Berol. 358 = sativum.
convallariifolium, Pall. ex Ledeb. Fl Ross iv. 184 = A. Victorialis.
Copaleri, Tineo, Cat. Pl. Hort. Panorm. 18 = paniculatum.
corbariense, Timb. ex Nym. Consp. 737 = subhirsutum.
cornutum, *Clem. It. Dalm.* t. I. f. 1.—Dalmatia.
cornutum, Kit. in Linnaea, xxxii. (1863) 340 = macrocephalum.
Cowani, Lindl. Bot. Reg. t. 758 = neapolitanum.
Crameri, *Aschers. & Boiss. ex Boiss. Fl. Orient.* v. 279. —Aegypt.
crinitum, Tausch, in Syll. Ratisb. ii. 255 = sphaerocephalum.
cristatum, *Boiss. Fl. Orient.* v. 287.—Thracia.
Cristophi, *Trautv. in Act. Hort. Petrop.* ix. (1884) 268.—As. temp.
croceum, *Torr. in Bot. Mex. Bound.* 218.—Calif.
Cumaria, Buch.-Ham. ex Wall. Cat. n. 5072 = A. Cepa.
Cupani, Guss. Pl. Rar. 143 = moschatum.
Cupani, *Refin. Caratteri*, (1818) 86. n. 206.—Reg. Mediterr.
cupuliferum, *Regel, in Act. Hort. Petrop.* iii. (1875) 284.—Turkestan.
curtum, *Boiss. & Gaillard, Diagn.* Ser. II. iv. 116.—Syria ; Aegypt.
Cusickii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 228. —Am. bor. occ.
cyaneum, *Regel, in Act. Hort. Petrop.*. iii. (1875) 174. —China.
Cydni, Schott & Kotschy, ex Schott, in Oestr. Bot. Wochenbl. (1854) 217 = casium.
Cyrilli, *Tenore, Fl. Nap.* iii. 864.—Italia ; Graecia ; Tauria.
dansbiale, Reichb. ex Kunth, Enum. Pl. iv. 428 = senescens.
dansbiale, Spreng. Fl. Hal. Mant. 38 = angulosum.
darwasicum, *Regel, in Act. Hort. Petrop.* vii. (1884) 659.—Bokhara.
Decaisnei, Presl, Bot. Bemerk. 114 = flavum.
decipiens, *Fisch. Hort. Gorenk.* ed. II. (1812) 10.— Reg. Caucas ; Turkest. ; Sibir.
declinatum, Retzb. Ic. Fl. Hort. Germ. v. 46. t. 477 = Stellerianum.
deflexum, Fisch. ex Kunth, Enum. Pl. iv. 428 = Stellerianum.

ALLIUM :—

deflexum, Willd. Sp. Pl. ii. 65 = lineare.
delicatulum, *Siewers, ex Willd. in Roem. & Schult. Syst.* vii. 1138.—Sibir.
dentiforum, De Not. Rep. Fl. Lig. 418 = sphaerocephalum.
denticulatum, Kit. in Linnaea, xxxii. (1863) 342 = carinatum.
dentiferum, Webb & Berth. Phyt. Canar. iii. 345. t. 234 = paniculatum.
denudatum, [Delar. in] Red. Lil. t. 357 = albidum.
Derderianum, Regel, in Act. Hort. Petrop. ii. (1875) 242 = naemanthoides.
descendens, Koch, Syn. Fl. Germ. ed. I. 719 = vineale.
descendens, Linn. Sp. Pl. ed. I. 299 = sphaerocephalum.
descendens, Pall. Ind. Taur. ex Schult. f. Syst. vii. 1011 = rotundum.
descendens, Sabth. & Sm. Fl. Gr. iv. t. 315. t. 316 = margaritaceum.
Désfeülen, Bor. Not. Nov. (1853) 17 ; (in Bull. Soc. Indust. d'Angers).—Cf. Ann. Soc. Linn. Maine-et-Loire, i. (1857) 63 = sphaerocephalum.
deserti, Boiss. ex Regel, in Act. Hort. Petrop. iii. (1875) 76 = Bourgeri.
desertorum, *Forsk. Fl. Aegypt. Arab.* 79.—Aegypt.
diaphanum, Janka, in Linnaea, xxx. (1859–60) 605 = tauricum.
dictyoprasum, Ledeb. Fl. Ross. iv. 166 = dyctioprasum.
dilutum, *Stapf, in Denkschr. Acad. Wien*, i. (1885) 13.—Persia.
Dioscoridis, *Sibth. & Sm. Prod. Fl. Gr.* i. 222.—Europ. austr. ; Orient.
djimilense, *Boiss. ex Regel, in Act. Hort. Petrop.* iii. (1875) 108.—Pontus.
dolonense, *Regel, l. c.* 113.—Sibir.
Douglasii, *Hook Fl. Bor. Am.* ii. 184. t. 197.—Calif.
Douglasii, Wood, in Proc. Philad. Acad. (1868) 170 = unifolium, bisceptrum.
Drégeanum, Kunth, Enum. Pl. iv. 382 = A. Scorodoprasum.
Drummondii, Regel, in Act. Hort. Petrop. iii. (1875) 112 = nutabile.
Durieuanum, Regel, l. c. 54 = A. Ampeloprasum.
Durieuanum, Walp. Ann. Bot. i. 860 = prasc.
Durieuanum, J. Gay, in Ann. Sc. Nat. Sér. III. viii. (1847) 218 = A. Ampeloprasum.
dyctioprasum, *C. A. Mey. ex Kunth, Enum. Pl.* iv. 390. —Armenia.
effusum, Boiss. Fl. Orient. v. 257 = stamineum.
elatum, *Regel, in Act. Hort. Petrop.* viii. (1884) 665.—Bokhara.
Elmeri, Regel, l. c. v (1877) 266 = acuminatum ?
ellipticum, Wall. Cat n. 5069 = A. Victorialis.
eminens, Gren. ex Parl. Fl. Ital. ii. 567 = sphaerocephalum.
Erdelii, Baker, in Bot. Mag. t. 6496 = orientale.
Erdelii, *Zucc. in Abh. Math. Phys. Acad. Muench.* iii. (1848) 388. t. 2.—Palaestin.
erectum, G. Don, in Mem. Wern. Soc. vi. (1827) 18 = rotundum.
ericetorum, Thore, Chlor. Land. 123 = ochroleucum, bakeri.
eriophyllum, *Boiss. Diagn.* Ser. I. vii. 112.—Persia.
erubescens, C. Koch, in Linnaea, xxii. (1849) 242 = rudbeckium.
erythraeum, *Gruseb. Spicil. Fl. Rumel.* ii. 396.— Macedon.
erythraeum, Weiss. ex Boiss. Fl. Orient. v. 258 = Weissii.
esculentum, Salisb. Prod. 385 = A. Cepa.
eucomum, Link & Otto, Ic. Pl. Rar. Hort. Berol. i. 15. t. 8 = moschatum.
euschoenoprasum, Syme, Engl. Bot. ed. III. ix. 815 = A. Schoenoprasum.
exaltatum, Kar. & Kir. ex Ledeb. Fl. Ross. iv. 178 = obliquum.
exile, Boiss. & Orph. Diagn. Ser. II. iv. 118 = sipylum.
exsertum, Baker, in Journ. Bot. xii. (1874) 294 = Bakeri.
exsertum, *G. Don, in Mem. Wern. Soc.* vi. (1827) 17. —Rossia.
falcifolium, *Hook. & Arn. Bot. Beech. Voy.* 400.— Calif.
fallax, *Schult. f. Syst.* vii. 1072.—Europ. ; As. Min.
Fetschenkoanum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 82.—Turkestan.
Fetisowi, *Regel, l. c.* v. (1877) 681.—Turkestan.
filidens, *Regel, l. c.* iii. 11. (1876) 174.—Turkestan.

ALLIUM :—

filiforme, *Colla, Herb. Pedem.* t. 90. f. 3.—Italia.
fimbriatum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 283.—Calif.
Fischeri, Bess. in Roem. & Schult. Syst. vii. 1081 = Stellerianum.
Fischeri, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 161.—Sibir.
fistile, S. T. Gray, Nat. Arr. ii. 188 = A. Ascalonicum.
fistile, *Steud. Nom.* ed II. i. 51, nomen.—As. Min.
fistulosum, *Linn. Sp. Pl.* 301.—Sibir.
flavescens, Bess. Enum. Pl. Volh. 56 = albidum.
flavescens, Poepp. ex Kunth, Enum. Pl. iv. 459 = striatum.
flavidum, *Ledeb. Fl. Alt.* ii. 7.—Sibir.
flavum, Less ex Ledeb. Fl. Ross. iv. 179 = delicatulum.
flavum, *Linn. Sp. Pl.* 299 —Europ. ; Oriens.
flavum, Salisb. Prod. 235 = A. Moly.
flexifolium, Jord. in Gren. & Godr. Fl. Fr. iii. 208 = carinatum.
fleuum, Wallst. & Kit. Pl. Rar. Hung. ii. 207 = carinatum.
fleuosum, Host, Fl. Austr. i. 422 = carinatum.
fleuosum, Urville, in Mém. Soc. Linn. Paris, i. (1822) 294 = staticaeforme.
foetidum, Willd. Enum. Suppl. 17 = flexum, carinatum.
folosum, Clar. ex [DC. in] Red. Lil. t. 216 = A. Schoenoprasum.
Fontanesii, J. Gay, in Ann. Sc. Nat. Sér. III. viii. (1847) 209 = flavum.
fragrans, Boj. Hort. Maurit. 348 = Milla borbonica.
fragrans, Cyr. ex Tenore, Fl. Nap. iii. 864 = Cyrilli.
fragrans, Vent. Jard. Cels, 36 = Nothocordum fragrans.
frigidum, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 34.— Graecia.
Friedrichsenum, Kunze, in Linnaea, xvi. (1842) 811 = margaritaceum.
fuscum, Schrad. ex Hornem. Hort. Hafn. ii. 962 = pulchellum.
fuscum, Spreng. Syst. ii. 37 = longispathum.
fuscum, Waldst. & Kit. Pl. Rar. Hung. iii. 267. t. 241 = paniculatum.
Fussii, *Kerner, in Oestr. Bot. Zeitschr.* xxviii. (1878) 155.—Austria.
galanthum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 508.—Songaria.
gaditanum, *Perez-Lara, ex Willk. in Illustr. Fl. Hisp.* 81.—Morocco.
Gardnisia, Bertero, ex Steud. Nom. ed. II. i. 664 = Brodiaea violacea.
Gasparrini, Guss. in Enum. Pl. Inarim. 337. t. 16 = A. Ampeloprasum.
Geyi, *Boiss Fl. Orient.* v. 271.—Cilicia.
Geyeri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 227. —Calif.
Ghorasdi, De Not. Rep. Fl. Lig. 412 = sphaerocephalum.
giganteum, *Regel, Gartenfl.* (1883) 91.—Reg. Himal.
glandulosum, *Link & Otto, Ic. Narc.* i. 38. t. 17.— Mexic.
glaucum, [Desf.] Hort. Par. ex [DC. in] Red. Lil. iv. t. 214 = foliosum.
glaucum, Hort. Par. ex Ledeb. Fl. Ross. iv. 167 = A. Schoenoprasum.
glaucum, Schrad. ex Poir. Encyc. Suppl. i. 365 = senescens.
globosum, *Sieb. Fl. Taur. Cauc.* i. 263 ; iii. 257.— Europ. austr. ; Reg. Casc. ; Sibir.
globosum, [*DC. in*] *Red. Lil.* iii. t. 179.—Hab. ?
glumaceum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 260.—Catsonia.
gomphrenoides, *Boiss. & Heldr. Diagn.* Ser I. vii. 114.—Lycaonia.
gorumense, *Boiss. Fl. Orient.* v. 283.—Cilicia.
Gouanii, G. Don, in Mem. Wern. Soc. vi. (1827) 35 = neapolitanum.
Govanianum, *Wall. Cat.* n. 5071.—Reg. Himal.
gracile, [Dryand. in] Ait. Hort. Kew. i. 429 = Nothoscordum fragrans.
graecum, J. Gay, ex Boiss Fl. Oriens. v. 271 = Gayi.
graecum, Urville, in Mém. Soc. Linn. Par. i. (1822) 293 = trifoliatum.
graminaceum, C. Koch, in Linnaea, xxii. (1849) 239 = rotundum.
graminifolium, *Buch, in Abh. Acad. Berl.* xvii. (1816) 262.—Ins. Canar.
graminifolium, Loisel. Fl. Gall. i. 258 = subhirsutum.
graminifolium, Pers. Syn. i. 356 = ochroleucum.
grandiflorum, Lam. Encyc. i. 68 = narcissiflorum.

ALLIUM :—

Grasi, *Regel, in Act. Hort. Petrop.* iii. (1875) 125.—Japon.

Griffithianum, Boiss. Ic. Ser. II. iv. 117 = rubellum.

Grimmi, *Regel, Dec. Pl. Nov.* 10.—*Cf. Act. Hort. Petrop.* viii. (1883) 272.—Turkestan.

Guicciardii, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 238 = flavum.

guttatum, Stev. in Mém. Soc. Nat. Mosc. ii. (1809) 178. t. 11 = margaritaceum.

haematchoites, *Boiss. ex Regel, Act. Hort. Petrop.* iii. 11. (1875) 240.—Persia.

haematochites, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 257.—Calif.

Halleri, Bab. Man. Brit. Bot. ed. I. 305 = A. Scorodoprasum.

Halleri, Crants, Inst. i. 488 = lineare.

Halleri, G. Don, in Mem. Wern. Soc. vi. (1827) 15 = A. Ampeloprasum.

Heldreichii, *Boiss. Diagn.* Ser. II. iv. 116.—Graecia.

hierochuntinum, *Boiss. Fl. Orient.* v. 244.—Palaestin.

himalayense, Regel, in Act. Hort. Petrop. iii. 11. (1875) 118 = Stockmanum.

hirsutum, Lam. Fl. Fr. iii. 262 = rubbirostrum.

hirsutum, *Zucc. in Abh. Bayer. Akad.* iii. (1843) 229. t. 2.—Palaestin.

hirtifolium, *Boiss. Fl. Orient.* v. 281.—Persia.

hirtovaginatum, Kunth, Enum. Pl. iv. 413 = Cupani.

Hoeltzeri, *Regel, in Act. Hort. Petrop.* viii. (1884) 657; *Regel, Gartenfl.* (1884) 291.—Turkestan.

holmense, Willd. ex Kunth, Enum. Pl. iv. 384 = A. Ampeloprasum.

Hookeri, Thw. Enum. Pl. Zeyl. 339 = Wallichii.

humile, *Kunth, Enum. Pl.* iv. 442.—Ind. or.

hyalinum, *Curran, in Bull. Calif. Acad.* n. 8. (1885) 155.—Calif.

hymenale, Kafa. Fl. Ladov. 19 = fragrans.

hymenorrhizum, *Ledeb. Fl. Alt.* ii. 12.—Turkest.; Persia.

hymettium, *Boiss. & Heldr. Diagn.* Ser. II. iv. 120.—Graecia.

iliense, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) 1. 452.—Soongaria.

illyricum, Jacq. Collect. iii. 278 = roseum.

illyricum, Link & Otto, ex Kunth, Enum. Pl. iv. 423 = senescens.

inaquale, Janka, in Linnaea, xxx. (1859–60) 603 = maschatum.

incarnatum, Hornem. Hort. Hafn. i. 338 = roseum.

indicum, Fisch. ex Roem. & Schult. Syst. vii. 1138 = tataricum.

inodorum, Ait. Hort. Kew. i. 427 = Nothoscordum fragrans.

inodorum, Willd. Sp. Pl. ii. 76 = angulosum.

insubricum, Boiss. & Reut. Cat. Sem. Jard. Genèv. (1854).—Cf. Linnaea, xxix. (1857) 721 = narcissiflorum.

intermedium, DC. Fl. Fr. vi. 818 = tataricum, paniculatum.

intermedium, G. Don, in Mem. Wern. Soc. vi. (1827) 88 = oleraceum.

Jacquemontii, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 102.—Ind. or.

Jacquemontii, Kunth, Enum. Pl. iv. 399 = rubellum.

japonicum, *Steud. Nom.* ed. II. i. 50.—Japon.

japonicum, *Regel in Act. Hort. Petrop.* iii. 11. (1875) 132.—Japon.

jeholense, *Franch. in Nouv. Arch. Mus. Hist. Nat. Par.* Sér. II. vii. (1884) 113; *Fl. Davidian.* i. (1884) 306.—China.

jenisseiense, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 237.—Persia.

jesdianum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 217.—Persia.

juldusicolum, *Regel, in Act. Hort. Petrop.* vi. (1879) 532.—Turkestan.

junceum, Jacquem. ex Baker, in Journ. Bot. xii. (1874) 295 = Jacquemontii.

junceum, *Sibth. & Sm. Prod.* i. 326.—Cyprus.

juncifolium, *D. Don, ex Steud. Nom.* ed. II. i. 52.—Chili.

karaknense, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 76.—Turkestan.

karataviense, *Regel, l. c.* 248.—Turkestan.

Karelinianum, *Regel, l. c.* 84.—Turkestan.

kermesinum, Reichb. ic. Fl. Germ. x. 25 = globosum.

Kesselringi, *Regel, in Act. Hort. Petrop.* viii. (1883) 272.—Turkestan.

ALLIUM :—

Klotzschi, Regel, in Act. Hort. Petrop. iii. 11. (1875) 93 = lilacinum.

Kochii, Lange, ex Nym. Consp. 736 = vineale.

kokanicum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 104.—Turkestan.

Korolkowii, *Regel, l. c.* ii. (1878) 407.—Turkestan.

Kotschyi, *Boiss. Diagn.* Ser. I. vii. 117.—Persia.

Kunthii, *G. Don, in Mem. Wern. Soc. vi.* (1827) 82.—Mexic.

Kuschakewicsi, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 117.—Turkestan.

laceratum, *Boiss. & Not, Diagn.* Ser. II. iv. 112.—Persia.

lachnophyllum, *Payne, in Pal. Expl. Soc. Cat. Pl.* 125.—Palaestin.

lacteum, Sibth. & Sm. Prod. 326 = neapolitanum.

lacunosum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 281.—Calif.

lactescens, Benth. Pl. Hartw. 389 = Brodiaea lactea.

lactinum, Pollini, Cat. Veron. (1812) 7 = neapolitanum.

lactum, Salisb. Prod. 285 = A. Porrum.

Lallemanni, Regel & Kach, in Ind. Sem. Hort. Petrop. (1856) 24 = decipiens.

latifolium, Gilib. Exercit. ii. 470 = ursinum.

latifolium, Jaub. & Spach, Illustr. Pl. Orient. ii. 8. t. 103 = A. Akaka.

laxiflorum, Tausch, in Flora, xii. (1829) Erg. 46 = vineale.

laxum, G. Don, in Mem. Wern. Soc. vi. (1827) 68 = angulosum.

Ledebourianum, *Schult. f. Syst.* vii. 1029.—Rossia.

Lehmannianum, *Mevchl. ex Bunge, in Mém. Sav. Étr. Pétersb.* vii. (1851) 509; *Rel. Lehm.* 333.—Turkestan.

Lemmoni, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 258.—Calif.

lepidum, Kunth, Enum. Pl. iv. 408 = paniculatum.

lepidum, Ledeb. Ic. Fl. Ross. iv. 17 = Pallasii.

leptophyllum, Schur, Enum. Pl. Transs. 674 = senescens.

leptophyllum, Wall. Cat. n. 5078 = rubellum.

Lessingii, Kar. ex Ledeb. Fl. Ross. iv. 170 = sabulosum.

leucanthum, C. Koch, in Linnaea, xxii. (1849) 240 = A. Ampeloprasum.

leucocephalum, Turcz. ex Ledeb. Fl. Ross. iv. 179 = flavidum.

Libani, *Boiss. Diagn.* Ser. I. xiii. 26.—Syria.

ligusticum, De Not. Prosp. Fl. Lig. 55 = flavum.

lilacinum, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 50. —Reg. Himal.

lilacinum, Royle, ex Regel, in Act. Hort. Petrop. iii. (1875) 89 = rubens.

liliiflorum, Zeyh. Cat. Schwetz. (1818) 15 = neapolitanum.

lineare, *Linn. Sp. Pl.* 295.—Sibir.

linearе, Tenore, Fl. Nap. i. 154 = sphaerocephalum.

linearе, Willd. ex Kunth, Enum. Pl. iv. 419 = strictum.

littorosum, Bertol. Amoen. It. 241 = vineale.

littorosum, G. Don, in Mem. Wern. Soc. vi. (1827) 8 = scorodoprasum.

longibulbum, Delac. Fl. Hantes-Pyr. 110 = A. Victorialis.

longicaule, *Hornem. Hort. Hafn. Suppl.* 130.—Asia.

longicuspe, Hort. ex Schult. f. Syst. vii. 1067 = glaucum.

longicuspis, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 65.—Kokanda.

longifolium, *Sprong. Syst.* ii. 88.—Am. mer.; Mexic.

longifolium, Lindl. Bot. Reg. t. 1034 = glandulosum.

longipetiolatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113 = unifolium.

longipetalum, Bertol. in Nov. Comm. Acad. Bonon. v. (1842) 439 = rubellum.

longipetalum, [Delar. in] Red. Lil. t. 316 = paniculatum.

longistylum, Baker, in Journ. Bot. xii. (1874) 294 = maculiferum.

loratum, *Baker, l. c.* 290.—Tibet. occ.

lusitanicum, *Lam. Encyc.* i. 70.—Lusitan.

lusitanicum, Link, ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 78 = A. Scheenoprasum.

lusitanicum, [Delar. in] Red. Lil. t. 271 = angulosum.

lutetum, Hort. ex Distr. Lexic. Gart. ed. II. i. 265 = obliquum.

Macleanii, *Baker, in Bot. Mag.* t. 6707.—Afghan.

Macmahlanum, *Hort. ex Regel, Gartenfl.* (1878) 261. t. 77. f. 2–8.—Hab.?

ALLIUM :—

macranthum, *Baker, in Journ. Bot.* xii. (1874) 293.—Reg. Himal.

macrocephalum, *Kit. in Linnaea*, xxxii. (1863) 340.—Austria.

macrochaetum, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* v. 289.—Assyria.

macrorhizon, Regel, in Act. Hort. Petrop. iii. 11. (1875) 154.—Turkestan.

macrorrhizum, Boiss. Diagn. Ser. I. xiii. 32 = hymenorhizum.

macrospermum, Boiss. & Kotschy, ex Boiss. Fl. Orient. v. 282 = orientale.

macrostemon, *Bunge, in Mém. Sav. Étr. Acad. Pétersb.* ii. (1835) 139.—China bor.

macrostylum, *Regel, in Act. Hort. Petrop.* vi. (1879) 396.—Tianschan.

macrum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 253. —Am. bor.

maddisi, *S. Wats. l. c.* 228.—Am. bor.

magicum, Linn. Sp. Pl. ed. II. 424 = nigrum.

mayale, Cyr. Pl. Rar. Neap. Fasc. i. 28. t. 4 = roseum.

margaritaceum, Heldr. ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 201 = staticiforme.

margaritaceum, Moench, Meth. Suppl. 80 = A. Scorodoprasum.

margaritaceum, *Sibth. & Sm. Fl. Graec. Prod.* i. 324. —Europ.; Orient.

marginatum, *Janka, in Termesz. Füzetek*, viii. (1884) 29.—Europ. austr.

maritimum, Kafu. Cavan. Cartt. 86. n. 205 = pusillum.

maritimum, *Torr. in Rep. U. St. & Mex. Bound.* 217.—Calif.

Maximowiczi, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 158.—As. or.

medium, G. Don, in Mem. Wern. Soc. vi. (1827) 88 = triquetrum.

megalobulbon, *Regel, in Act. Hort. Petrop.* vi. (1879) 526.—Turkestan.

melananthemum, *Pant. Elem.* 64.—Europ. merid.

microcephalum, Tausch, in Flora, ii. (1819) 462 = strictum.

microcephalum, Willd. ex Kunth, Enum. Pl. iv. 425 = angulosum.

minutiflorum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 242.—Persia.

mobilense, Regel, l. c. = mutabile.

modestum, Boiss. Diagn. Ser. I. xiii. 33 = paniculatum.

magadorense, Willd. ex Kunth, Enum. Pl. iv. 384 = A. Ampeloprasum.

Moly, Georgi, Beschreib. Russ. Reich. iii. iv. 895 = decipiens.

Moly, Grisch. & Schur, ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 199 = globosum.

Moly, *Linn. Sp. Pl.* 301.—Europ.

Moly, Ucria, Hort. Panorm. 153 = subhirsutum.

monadelphum, *Turcz. ex Kunth, Enum. Pl.* iv. 398.—Sibir. baical.

mongolicum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 160.—Mongolia.

monterotianum, Pomr. ex Willk. & Lange, Prod. Fl. Hisp. i. 307 = carinatum.

monspessulanum, Gouan, Illustr. 24 = nigrum.

monspessulanum, Willd. Enum. Suppl. 16 = paniculatum.

montanum, Guss. Pl. Rar. 142 = tauriforum.

montanum, Reichb. Ic. Fl. Germ. x. t. 463. f. 1060 = flavum.

montanum, Tenore, Fl. Nap. iii. 367 = Cupani.

montanum, F. W. Schmidt, ex Schult. f. Syst. vii. 1072 = lusum.

montanum. Sibth. & Sm. Fl. Graec. t. 319 = paniculatum.

moschatum, *Linn. Sp. Pl.* ed. I. 299.—Europ.; Orientes.

moschatum, Moris, Stirp. Sard. Elench. i. 46 = parciflorum.

moschatum, Sintenis, ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 199 = globosum.

moschatum, Urville, in Mém. Soc. Linn. Par. i. (1822) 299 = Cupani.

multibulbosum, Jacq. Fl. Austr. i. 9 = nigrum.

multiflorum, DC. Fl. Fr. vi. 816 = polyanthum.

multiflorum, *Desf. Fl. Atlant.* i. 295.—Algeria.

multiflorum, Kunth, Enum. Pl. iv. 387 = rotundum.

multiflorum, Regel, in Act. Hort. Petrop. iii. (1875) 61, pro parte = subhirsutum.

Murrayanum, Regel, in Gartenfl. (1878) 260. t. 770. f. 1 = nanningtonum.

mutabile, *Michx. Fl. Am. Bor.* i. 196.—Georgia.

ALLIUM :—
myrianthum, Boiss. Diagn. Ser. I. v. 59 = A. Ampeloprasum.
Napus, Pall. ex Linnæum. Pl. iv. 395 = A. Cepa.
narcissiflorum, *Vill. Prosp.* 18.—Europ.
narcissifolium, Lam. Fl. Fr. iii. 303 = pedemontanum.
narcissifolium, Sonp. Fl. Cam. ed. II. i. 238 = senescens.
neapolitanum, *Cyr. Pl. Rar. Neap.* i. 18. t. 4.—Europ. ; Oriens.
nebrodense, Guss. Prod. Fl. Sic. i. 404 = flavum.
neglectum, Wender. in Schrift. Ges. Nat. Marb. ii. (1831) 211 = A. Scorodoprasum.
nemorale, Salisb. Prod. 285 = ursinum.
Nereidum, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 244.—China.
neriniflorum, *G. Don, ex Loud. Encyc. Pl. Suppl.* ii. 1362.—China.
nerinifolium, Baker, in Journ. Bot. xii. (1874) 290 = Thunbergii.
nevadense, *S. Wats. in Bot. King, Exp.* 351, 487. t. 38 f. 1–8.—Calif.
Nevii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 281.—Calif.
nigrum, All. Fl. Pedem. ii. 158. t. 95 f. 1 = narcissiflorum.
nigrum, Bieb. Fl. Taur. Cauc. i. 266 = decipiens.
nigrum, G. Don, in Mem. Wern. Soc. vi. (1827) 89 = multibulbosum.
nigrum, *Linn. Sp. Pl.* ed. II. 430.—Europ. ; Oriens.
nigrum, Stbth. & Sm. Fl. Græc. i. 328 = Bauerianum.
nipponicum, *Franch. & Sav. Enum. Pl. Jap.* ii. 527. —Japon.
nitens, Sauzé & Maill. Cat. Pl. Deux-Sèvres, 51 = vineale.
nitidulum, Fisch. ex Ledeb. Fl. Ross. iv. 171 = Pallasii.
Nutschmanni, Willd. ex Ledeb. l. c. 175 = flavum.
nivale, Jacquem. ex Regel, in Act. Hort. Petrop iii. 11. (1875) 178 = Govanianum.
niveum, Roth, Cat. Bot. fasc. ii. 35 = subhirsutum
Noeanum, *Reut. ex Regel, in Act. Hort. Petrop.* iii. 11. (1875) 285.—As. Min.
nodicaule, *Lehm. Cat. Hort. Hamb.* (1826).—Bras
nutans, Herbb. quorr. ex Schult. f. Syst. vii. 1088 = cernuum.
nutans, *Linn. Sp. Pl.* 299.—Sibir.
Nuttallii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 227.—Am. bor.
obliquum, *Linn. Sp. Pl.* 296.—Sibir.
obscurum, Bieb. ex Schult. f. Syst. vii. 1008 = A. Scorodoprasum.
obtusiflorum, [DC. in] Red. Lil. t. 118 = pusillum.
obtusifolium, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 51.—Reg. Himal.
occidentale, A. Gray, in Proc. Am. Acad. vii. (1868) 390 = attenuifolium.
ochroleucum, Reichb. Ic. Fl. Germ. x. t. 498. f. 1090 = globosum.
ochroleucum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 204. t. 186.—Europ.
odoratissimum, *Desf. Fl. Atlant.* i. 289. t. 88.—Afr. bor.
odorum, Kar. & Kir. in Bull. Soc. Nat. Mosc (1841) 854 = angulosum.
odorum, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 48 = ochroleucum.
odorum, *Linn. Mant.* 62.—Sibir.
odorum, Lour. Fl. Cochinch. 251 = Thunbergii.
odorum, Tenore, Fl. Nap. Prod. 21 = nigrum.
odorum, Thunb. Fl. Jap. 132 = Thunbergii.
oleraceum, Des Moul. Cat. Dord. 142, ex parte = denticulatum.
oleraceum, *Linn. Sp. Pl.* 299.—Europ.
oliganthum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 856 = A. Schœnoprasum.
Olivieri, *Boiss. Fl. Orient.* v. 284.—Mesopotamia.
olympicum, *Boiss. Diagn.* Ser. I. v. 58.—Thracia.
Ophioscorodon, G. Don, in Mem. Wern. Soc. vi. (1827) 2 = sativum.
Opizii, Wolfner, in Lotos, iv. (1854) 176 = triquetrum.
oreophiloides, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 114.—Turkestan.
oreophilum, *C. A. Mey. Verz. Pl. Cauc.* 37.—Cancas.
oreoprasum, *Schrenk, in Bull. Sc. Acad. St. Pétersb.* x. (1842) 854.—Turkestan.
odinsale, *Boiss. Diagn.* Ser. I. xiii. 25.—Oriens.
ornithogaloides, Wall. Fl. Carol. 121 = striatum.

ALLIUM :—
Ostrowskianum, *Regel, in Act. Hort. Petrop.* vii. (1880) 545.—Turkestan.
ovifforum, *Regel. l. c.* viii. (1884) 658.—Reg. Himal.
oxypetalum, G. Don, in Mem. Wern. Soc. vi. (1827) 87 = oleraceum.
Pallasii, *Murr in Comm. Gott.* vi. (1775) 92. t. 3. —Sibir.
pallens, Linn. Sp. Pl. ed. II. 427 = paniculatum.
pallens, Reichb. Ic. Fl. Germ. x t 484 f. 1062 = flavum.
pallens, *Sibth & Sm. Fl. Græc.* t. 317.—Græcia.
Palmeri, *S. Wats. Bot. in Rep. Geol. Expl. Fortieth Parallel*, 487. t. 27. f. 10, 11.—N. Mexico.
palustre, Pourr. ex Lag. Elench. 13 = A. Schœnoprasum.
palustre, Salisb. Prod. 285 = sibiricum.
paniculatum, All. Fl. Pedem. ii. 157 = flavum
paniculatum, Ker-Gawl. in Bot. Mag. t. 1432 = caucasicum.
paniculatum, Vill. Hist. Pl. Dauph. ii. 854 = intermedium.
paniculatum, *Linn. Syst.* ed. X. 978.—Europ ; Oriens.
paniculatum, Stbth. & Sm. Fl. Græc. iv. t. 318 = fuscum.
paniculatum, Viv. Fl. Lib. Spec. 19 = nigrum.
pannonicum, Gromow. ex Trautv. in Act. Hort. Petrop. ix. (1884) 278, nomen.—Europ
papillare, *Boiss. Diagn* Ser. I. xiii 27.—Palæstin.
paradoxum, *G. Don, in Mem. Wern. Soc* vi. (1827) 72.—Caucas. ; Persia.
parciflorum, *Vis. App. Fl. Cors. Prod.* 2.—Corsic.
Parishii, *S. Wats. in Proc Am. Acad.* xvii. (1881-82) 380.—Am. bor.
Parryi, *S. Wats. l. c.* xiv. (1879) 231.—Calif.
parviflorum, Desf. Fl. Atlant. i. 290 = paniculatum.
parviflorum, Desv. Fl. Alg. 96 = sphærocephalum
parviflorum, *Linn Sp. Pl.* ed. II. 427.—Europ.
parviflorum, Thuill. Fl. Paris, ed. II i 166 = oleraceum.
parvifolium, Cranz, Inst. i. 49 = parviflorum.
parvum, *Kellogg, in Proc. Calif Acad* iii. (1863) 54. f. 13.—Calif.
Paterniliæ, *Boiss. Diagn.* Ser. II. iv. 114.—Reg Caucas.
paucibulbosum, *Haw. ex Steud. Nom.* ed. II i. 58 — Hispan.
pauciflorum, Gren. & Godr. Fl. Fr. iii. 210 = parciflorum.
pauciflorum, Willd ex Ledeb. Fl. Ross. iv. 169 = A. Cepa.
pedemontanum, Willd. Sp. Pl. ii. 77 = narcissiflorum.
pendulinum, Tenore, Fl. Nap. i 168. t. 31 = triquetrum.
permixtum, Guss. Fl. Sic. Prod. i. in add. 8 = subhirsutum.
persicum, Fisch ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 43 = A. Scorodoprasum
petiolatum, Lam. Fl. Fr. iii. 261 = ursinum.
petræum, Kar. & Kir. in Bull. Soc. Nat. Mosc xv. (1842) 511 = globosum.
phaleratum, Heldr. & Sart. ex Heldr. in Atti Congr. Bot. Firenz, 1874 (1876) 233 = staticiforme.
phaneranthemum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 235.—Syria.
philistæum, Boiss. Diagn. Ser. I. xiii. 26 = Erdelii.
phrygium, *Boiss. in Bal. ex Boiss. Fl. Orient.* v. 256. —Phrygia.
phthoicum, *Boiss. & Heldr. ex Boiss. l. c.* 274.—Oriens.
pilosum, *Sibth. & Sm. Fl. Græc.* t. 321.—Archip. Græc.
pisidicum, Boiss. & Heldr. ex Boiss. Diagn. Ser. I. xiii. 24 = Cupani.
plantaginense, Willk. & Lange, Prod Fl. Hisp. i. 211 = A. Victorialis.
plantagineum, Lam. Fl. Fr. iii. 262 = A. Victorialis.
platycaule, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 234.—Calif.
platyspathum, *Schrenk, Enum.* 7.—Turkestan.
platystemon, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 514 = oreophilum.
pleianthum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 238.—Am. bor.
Plummerae, *S. Wats. l. c.* xviii. (1882–83) 195.— Mexic.
Poeppigii, *Kunth, Enum. Pl.* iv. 456.—Chili.
polyanthum, Schult. f. Syst. vii. 1016 = rotundum.
polyphyllum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 508.—Turkestan.
polyrhizum, *Turcz. ex Regel, in Act. Hort. i Petrop.* iii. (1875) 169.—Sibir. baical.

ALLIUM :—
porphyroprasum, Heldr. & Sart. ex Boiss. Fl. Orient v. 233 = rotundum.
porraceum, S. F. Gray, Nat. Arr. ii. 181 = A. Ampeloprasum.
Porrum, Georgi, Beschreib. Russ. Reich. iii. iv. 891 = obliquum.
Porrum, *Linn. Sp. Pl.* 295.—Europ.
Potanini, *Regel, in Act. Hort. Petrop.* vi. (1879) 295.— Mongol.
præcursum, Reichb. Ic. Fl. Germ. v. 17. t. 429 = paniculatum.
pratense, Schleich. ex Kunth, Enum. Pl. iv. 382 = carinatum.
Prestianum, Schult. f. Syst. vii. 1132 = rotundum.
procerum, *Trautv. in Act. Hort. Petrop.* viii. (1884) 668 : ix. (1884) 274.—Bokhara.
proliferum, Schrad. ex Willd. Enum. Hort. Berol. 358 = A. Cepa
prostratum, Maxim. Prim. Fl. Amur. 283 = Maximowiczii.
prostratum, Trev. Ind. Sem. Vratisl. (1821) = Stellerianum.
pruinatum, Link, in Spreng. Syst. ii. 36 = monspessulanum.
Przewalskianum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 164.—China occ.
Pseudo-Cepa, Schrenk, in Bull. Sc. Acad. Péterab. x. (1842) 356 - galanthum.
Pseudo-ochroleucum, Schur, Enum. Pl. Transs. 673 = globosum.
pulchellum, *G. Don, in Mem. Wern. Soc.* vi. (1827) 46.—Europ ; Oriens.
pulchrum, Clark, Trav. iii. 337 = nigrum.
punctulatum, Schlecht. in Linnæa, xix. (1847) 401 = A. Schœnoprasum.
purpureum, Salisb. Prod. 285 = senescens.
purpureum, Schur, Enum. Pl. Transs 672 = carinatum.
Purshii, G. Don, in Mem. Wern. Soc. vi. (1827) 10 = vineale.
pusillum, *Cyr. ex Tenore, Cat. Neap.* (1813) 4.—Ital.
pusillum, Presl, Delic. Prag. 147 = Cupani.
pustilum, Willd. Enum. Hort. Berol. Suppl. 17 = delicatulum.
pustulosum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 248.—Catsonia.
pylæum, De Not. in Linnæa, xviii. (1844) 157 = A. Ampeloprasum.
pyrenaicum, *Costa, Suplem. Catal.* 92 ; *Costa & Vayr. in Anal. Soc. Esp. Hist. Nat.* ix. (1880) 87.—Pyren.
Raddeanum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 155.—Siberia.
ramosum, Georgi, Beschr. Russ. Reich. iii. iv. 892 = odorum.
ramosum, Jacq. Misc. ii. 308 = obliquum.
ramosum, *Linn. Sp. Pl.* 296.—Sibiria.
ramosum, Pall. ex Ledeb. Fl. Ross. iv. 184 = tataricum.
reflexum, F. G. Dietr. Lexic. Gaert. ed. II. i. 285 = A. Schœnoprasum.
reflexum, *Boiss. & Reut. ex Boiss. Fl. Orient.* v. 281. —Mesopot.
Regelii, *Trautv. in Act. Hort. Petrop.* ix. (1884) 275. —Transcauc.
Renari, *Regel, l. c.* vi. (1879) 521.—Turkestan.
reticulatum, Benth. Pl. Hartw. 339 = attenuifolium.
reticulatum *Pras. ex G. Don, in Mem. Wern.* Soc. vi. (1827) 36.—Am. bor.
reticulatum, J. & C. Presl, Fl. Cech. 78 = strictum.
reticulatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Victorialis.
reticulatum, Wallr. Sched. Crit. 135 = angulosum ?
Reuterianum, *Boiss. Diagn.* Ser. I. v. 60.—As. Min.
riparium, Opiz, Natural. vi. (1824) 50 = sibiricum.
robustum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 853 = atropurpureum.
Rosenbachianum, *Regel, in Act. Hort. Petrop.* viii. (1884) 664.—Bokhara.
roseum, Krock. Fl. Siles. i. 316. t. 43 = sibiricum.
roseum, *Linn. Sp. Pl.* ed. I. 296.—Reg. Mediterr.
Rothii, *Zucc. in Abh. Bayer. Akad.* iii. (1843) 222. t. 2. f. 2.—Palæstin.
rotundifolium, Lamnitz. ex Steud. Nom. ed. II. i. 58 = rotundum.
rotundum, All. Fl. Pedem. ii. 155 = acutiflorum.
rotundum, *Linn. Sp. Pl.* ed. II 423.—Europ. ; As. Min.
rotundum, Wimm. & Grab. Enum. 56 = vineale.
Rouyanum, Kanth, Enum. Pl. iv. 454 = odorum

ALLIUM :—
rubellum, *Bieb. Fl. Taur. Cauc.* i. 264.—Europ.; Oriens; Sibir.
rubellum, Presl, Fl. Sic. 42 = rotundum.
rubellum, Regel, in Act. Hort. Petrop. iii. 11. (1875) 107 = Tachtajan
rubellum, Willd. ex Ledeb. Fl. Ross. iv. 179 = globosum.
rubens, *Baker, in Journ Bot.* xii. (1874) 293.—Reg. Himal.
rubens, Schrad. ex Willd. Enum. 360 = Stellerianum.
rubicundum, Nlven, ex G. Don, in Mem. Wern. Soc. vi. (1827) 16 = rotundum.
rubicundum, *Willd. ex Kunth, Enum. Pl.* iv. 489.—Hab.?
rubrovittatum, *Boiss. & Heldr. in Boiss. Diagn.* Ser. I. xiii. 29.—Creta.
rudbaricum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 215.—Persia.
rupestre, *Bieb. Fl. Taur. Cauc.* iii. 250.—Iberia; Taur.
rupestre, Fisch. ex Schrank, in Denkschr. Regensb. Bot. Ges. ii. (1822) 38 = nutatile.
rupestre, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 260 = paniculatum.
Ruprechti, *Bull. Fl. O ient.* v. 264.—Caucas.
ruthenicum, Steud. Nom. ed. 11. i. 54 = flavum.
sabulosum, *Fisch. ex Goebel, Reise*, ii. 311.—Reg. Casp.
sabulosum, *Stev. ex Ker, & Kir. in Bull. Soc. Nat. Mosc.* (1842) 518.—Persia; Turkestan.
sacculiferum, *Maxim. Prim. Fl. Amur.* 281.—Reg. Amur.
sairamense, *Regel, in Act. Hort. Petrop.* vi. (1879) 520.—Turkestan.
Salesovianum, Fisch. ex Regel, l. c. iii. 11. (1875) 162 = Fischeri.
Sanbornii, *Wood, in Proc. Ac. Nat. Sc. Philad.* (1868) 171.—Calif.
sapidissimum, Hedw. in Roem. Collect. 76 = fistulosum.
saravschanicum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 244.—Turkestan.
sardoum, Moris, Stirp. Sard. Elench. fasc. ii. 10 = sphaerocephalum.
sativum, *Linn. Sp. Pl.* 296.—Europ.
Savii, Parl. Fl. Ital. ii. 554 = paniculatum.
savranicum, Bess. Cat. Sem. Hort. Crem. Suppl. (1818) = globosum.
saxatile, Bieb. Fl. Taur. Cauc. i. 264 = globosum.
saxatile, Hohen. ex Boiss. Fl. Orient. v. 268 = lepidum.
saxatile, Pall. Reise, iii. 293 = fistulosum.
saxatile, Willd. ex Ledeb. Fl. Ross. iv. 181 = albidum.
scaberrimum, Serres, in Bull. Soc. Bot. Fr. iv. (1857) 439 = sphaerocephalum.
scabrellum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 215.—Persia.
scabriflorum, *Boiss. Diagn.* Ser. I. v. 60.—Cappadoc.
scabrinsparum, *Boiss. l. c.* xiii. 31.—Persia.
scabrum, Gilib. Exercit. ii. 469 = oleraceum.
scaposum, *Benth. in Pl. Hartw.* 26.—Mexic.; Texas.
scariosum, Jan, ex Schult. f. Syst. vii. 1071 = rotundum.
schergianum, *Boiss. Fl. Orient.* v. 251.—Syria.
Schmidtianum, *Tausch, in Flora*, xii. (1829) *Erg.* 45.—Bohemia.
schoenoprasoides, Fries, ex Nym. Consp. 741 = A. Schoenoprasum.
schoenoprasoides, *Regel, in Act. Hort. Petrop.* v. (1877) 630.—Reg. Himal.
Schoenoprasum, *Linn. Sp. Pl.* 301.—Reg. temp. bor.
Schoenoprasum, Regel & Tiling, Fl. Ajan. 128 = Ledebourianum.
Schrenkii, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 173.—Ind. or.
Schuberti, *Zucc. in Abh. Bayer. Akad.* iii. (1843) 284. t. 5. f. 1.—Oriens.
Schulleri, Stev. ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 60 = Aucheri.
scilloides, Dougl. ex S. Wats. in Proc. Am. Acad. xiv. (1879) 229 = collinum.
Scordoprasum, Aschers. Fl. Brandenb. 725 = seq.
Scordoprasum, *Linn. Sp. Pl.* 297.—Europ.; Reg. Cauc.; Syria.
scorsonerafolium, [*DC. in*] *Red. Lil.* ii. t. 93.—Europ. austr.
segetum, *Jan, ex Schult. f. Syst.* vii. 1090.—Sicilia.
Sellovianum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 117.—Bras.; Chili.
Semenovi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) 1. 449.—Reg. Himal.
semiretschenskianum, *Regel, in Act. Hort. Petrop.* v. (1877) 630.—As. centr.

ALLIUM :—
senescens, Ker-Gawl. in Bot. Mag. t. 1150 = glaucum.
senescens, Host, Fl. Austr. i. 420 = fallax.
senescens, *Linn. Sp. Pl.* 299.—Europ.; Sibir.
senescens, Seter, Fl. Helv. i. 198 = acutangulum.
senescens, Mlq. Ann. Bot. Lugd. Bot. iii. 818 = odorum.
senescens, Thunb. Fl. Jap. 128 = japonicum.
serbicum, Vis. & Pancˇ. in Mem. Ist. Venet. xii. (1864) 479 = moschatum.
serotinum, Lapeyr. Fl. Pyr. t. 84 = ochroleucum × suaveolens.
serotinum, Schleich. ex G. Don, in Mem. Wern. Soc. vi. (1827) 63 = senescens.
serrulatum, Boiss. Fl. Orient. v. 241 = atroviolaceum.
serratum, *S. Wats. Bot. King, Exped.* 487. t. 87. f. 4, 5.—Calif.
sessiliflorum, Fisch. ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 140 = senescens.
setaceum, Waldst. & Kit. Pl. Rar. Hung. i. 70. t. 68 = moschatum.
setifolium, *Schrenk, Enum.* 6.—Soongaria.
Seweerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) 458.—Turkestan.
sibiricum, Linn. Mant. 562 = A. Schoenoprasum.
Sibthorpianum, Schult. f. Syst. vii. 1057 = paniculatum.
sicheum, Ucria, in Roem. Arch. i. 68 = Dioscoridis.
Siebersianum, Schult. f. Syst. vii. 1099 = neapolitanum.
sikkimense, *Baker, in Journ. Bot.* xii. (1874) 292.—Reg. Himal.
sinaicum, *Boiss. Diagn.* Ser. I. xiii. 31.—Arabia.
sindjarense, *Boiss. & Hausskn. ex Regel, in Act. Hort. Petrop.* iii. 11. (1875) 121.—Mesopotamia.
sinicum, *Debeaux, in Verh. Baten. Gen.* v. (1790) ed. I. *Art.* iv. 6; ed. II. (1827) 68.—China.
sipyleum, *Boiss. Diagn.* Ser. I. v. 58.—Lydia; Ins. Chios.
sordidum, Schur, Enum. Pl. Transs. 673 = globosum.
spathaceum, Steud. ex A. Rich. Tent. Fl. Abyss. ii. 380 = subhirsutum.
speciosum, Cyr. Pl. Rar. Neap. Fasc. ii. 18 = nigrum.
spectabile, De Not. Fl. Lig. 414 = A. Ampeloprasum.
sphaerocephalum, Crome, ex Schlecht. Fl. Berol. ii. 249 = rotundum.
sphaerocephalum, *Linn. Sp. Pl.* 297.—Europ.; Orient.
spirale, Hort. Dorp. ex Kunth, Enum. Pl. iv. 432 = sparsum.
spiralis, Willd. Enum. Suppl. 17 = senescens, nutans.
splendens, Mlq. in Ann. Mus. Lugd. Bat. iii. 154 = Bakeri.
spoliatum, *Willd. ex Schult. f. Syst.* vii. 1025.—Sibiria.
spurium, *G. Don, in Mem. Wern. Soc.* vi. (1827) 59 = senescens.
spurium, Hort. ex Schult. f. Syst. vii. 1067 = glaucum.
stamineum, *Boiss. Diagn.* Ser. II. iv. 119.—Oriens.
stellerianum, *Sibth. & Sm. Fl. Graec.* iv. t. 320.—Graecia.
stellatum, *Pras. ex Ker-Gawl. in Bot. Mag.* t. 1576.—Am. bor.
Stellerianum, Bess. Cat. Hort. Crem. (1824) = albidum.
Stellerianum, *Willd. Sp. Pl.* ii. 82.—Sibiria.
Stellerianum, Ledeb. Fl. Ross. iv. 179 = flavidum.
stenopetalum, *Boiss. & Kotschy, ex Regel, in Act. Hort. Petrop.* iii. 11. (1875) 231.—Cilicia.
stenophyllum, Gmel. Sic. Syn. i. 254 = margaritaceum.
stenophyllum, Regel, ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) ii. 446 = dolomocarnse.
stenophyllum, *Schrenk, in Bull. Phys.-Math. Acad. Petrop.* iii. (1844) 210.—Soongaria.
stenophyllum, Wall. Cat. n. 5073 = Jacquemontii.
Stevenii, Willd. ex Ledeb. Fl. Ross. iv. 176 = globosum.
stipitatum, *Regel, in Act. Hort. Petrop.* vii. (1880) 546.—As. centr.
Stocksianum, *Boiss. Diagn.* Ser. II. iv. 117.—Beluchistan.
Stolleski, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 160.—Tibet. occ.
Strachevi, *Baker, in Journ. Bot.* xii. (1874) 292.—Reg. Himal.
stramineum, Boiss. & Rent. Diagn. 25 = A. Moly.
stramineum, *Regel, Gartenfl.* xxv. (1876) 355.—Manchuria.
stramineum, Schur, Enum. Pl. Transs. 674 = angulosum.

ALLIUM :—
striatellum, *Lindl. in Trans. Hort. Soc.* vi. (1851) 97.—Chili.
striatum, Jacq. Coll. Suppl. 51 = Nothoscordum striatum.
strictum, *Schrad. Hort. Gœtt.* t. 1.—Europ.; Oriens.
suaveolens, Gaud. Fl. Helv. ii. 491 = strictum.
suaveolens, *Jacq. Coll.* ii. 305.—Europ.
suaveolens, Duby, Bot. Gall. i. 470 = ochroleucum.
subalbidum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 128 = roseum.
subalpinum, Pall. Ind. Taur. ex Kunth, Enum. Pl. iv. 448 = decipiens.
subhirtiflorum, *Calla, in Mem. Acc. Torin.* xxxix. (1836) 18. t. 52.—Chili.
subhirsutum, Delile, Fl. Aegypt. ex Boiss. Fl. Orient. v. 274 = neapolitanum.
subhirsutum, Desf. Fl. Atlant. 286 = subvillosum.
subhirsutum, *Linn. Sp. Pl.* 295.—Europ.; Oriens.
subhirsutum, Reichb. Fl. Germ. Excurs. 113 = subvillosum.
subhirsutum, Sieber, ex Kunth, Enum. Pl. iv. 441 = neapolitanum.
subquadrifloriforum, *Boiss. Fl. Orient.* v. 284.—Armenia.
subtilissimum, *Ledeb. Ic. Fl. Alt.* ii. 22. t. 160.—Sibir. altaic.
subvillosum, Salzm. ex Schult. f. Syst. vii. 1104 = subhirsutum.
sulcatum, [DC. in] Red. Lil. t. 482 = neapolitanum.
Sukia, Buch.-Ham. in D. Don, Prod. Fl. Nep. 58 = fragrans.
Suworowi, *Regel, in Act. Hort. Petrop.* vii. (1880) 546.—As. centr.
Syenitis, G. Don, in Mem. Wern. Soc. vi. (1827) 19 = rotundum.
Synnotianum, Schult. f. Syst. vii. 1015 = praec.
systemanthum, C. Koch, in Linnaea, xxii. (1849) 288 = rubellum.
syriacum, Boiss. Fl. Orient. v. 282 = A. Ampeloprasum.
Szowitsi, Regel, in Act. Hort. Petrop. iii. 11. (1875) 171 = strictum.
szurlense, *Lerch. ex Ledeb. in Verh. Siebenb. Ver. Naturw.* xxxiv. (1884) 55.—Europ.
talassicum, *Regel, in Act. Hort. Petrop.* v. (1877) 638.—Turkestan.
tartaricum, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 288 = odorum.
tartaricum, Hort. ex Schult. f. Syst. vii. 1066 = tartaricum.
tartaricum, *Linn. f. Suppl.* 196.—Rossia; Sibiria; Persia.
tauricolum, Boiss. Diagn. Ser. II. iv. 118 = frigidum.
tauricum, Hort. ex Schult. f. Syst. vii. 1021 = flavum.
tauricum, Pall. ex G. Don, in Mem. Wern. Soc. vi. (1827) 50 = globosum.
tchefoense, *Debeaux, in Actes Linn. Soc. Bord.* xxxii. (1878) 25; *Fl. Tché-Fou*, 188.—China.
Tchihatschewii, *Boiss. in Tchihat. Asie Min. Bot.* ii. 251.—Armenia.
Temoli, Sprng. Pugill. i. 29; Syst. ii. 35 = roseum.
tenue, C. Koch, in Linnaea, xxii. (1849) 288 = rubellum.
tenue, Frivald. ex Griseb. Spicil. Fl. Rumel. ii. 396 = erythraeum.
tenue, G. Don, in Mem. Wern. Soc. vi. (1827) 34 = Pallasii.
tenuiflorum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 206.—Turkestan.
tenuiflorum, Tenore, Fl. Neap. i. 165. t. 30 = paniculatum.
tenuifolium, *Frivald. in Soll. Ratisch.* ii. 83.—Europ.
tenuifolium, Salisb. Prod. 385 = A. Schoenoprasum.
tenuifolium, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 76 = globosum.
tenuissimum, Habl. Taur. 162 = moschatum.
tenuissimum, *Linn. Sp. Pl.* 301.—Sibiria.
tenuiflolium, *Regel, in Act. Hort. Petrop.* v. (1877) 629.—Turkestan.
thessalum, Boiss. Fl. Orient. v. 282 = A. Ampeloprasum.
Thomsoni, *Baker, in Journ. Bot.* xii. (1874) 294.—Reg. Himal.
Thunbergii, *G. Don, in Mem. Wern. Soc.* vi. (1827) 34.—Japon; China.

ALLIUM :—
tianschanicum, Rupr. in Mém. Acad. Pétersb. Sér. III. xiv. (1869) 88. (Sert. Tianach. 88) = hymenorhizum.
Tilingi, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 124. —Calif.
Tinei, J. & C. Presl, Delic. Prag. 145 = subhirsutum.
Tolmiei, Baker, ex Coult. Rocky Mt. Bot. 349 = tribracteatum.
tortuosum, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 76.—Europ.
tortuosum, Willd. ex Schult. f. Syst. vii. 1064 = lineare.
Tournefortii, Boiss. Fl. Orient. v. 274 = roseum.
trachyanthum, Griseb. Spicil. Fl. Rumel. ii. 395 = rotundum.
trachyphyllum, Boiss. & Sprun. Diagn. Ser. I. vii. 114 = sphaerocephalum.
transilvanicum, *Lerch. ex Kanitz, in Verh. Siebenb. Ver. Naturw.* xxiv. (1884) 54.—Europ.
Trautvetteri*anum*, *Regel, in Act. Hort. Petrop.* viii. (1884) 661.—Bokhara.
tribracteatum, *Torr. in Pacif. Rail. Rep.* iv. 148.— Calif.
trichocnemis, *γ. Gay, in Ann. Sc. Nat. Sér.* III. viii. (1847) 310.—Algeria.
tricoccum, [*Solanz. in*] *Ait. Hort. Kew.* i. 428.—Am. bor.
tricoccum, Blanco, Fl. Filip. ed. I. 239 = uliginosum.
tricorne, Desv. ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 145 = angulosum.
tricorne, Hort. ex Poir. Encyc. Suppl. i. 270 = cernuum.
trifiorum, *Rafin. in Med. Repos. New York*, v. (1808) 362.—Pennsylv.
trifoliatum, *Cyr. Pl. Rar. Neap. Fasc.* ii. 11.—Europ.; Orient.
tripedale, Trautv. in Act. Hort. Petrop. ii. (1873) 485 = Dioscoridis.
triquetrum, Hort. ex Schrad. in Schult. f. Syst. vii. 1078 = angulosum.
triquetrum, *Linn. Sp. Pl.* 300.—Europ.; Afr. bor.
triquetrum, Lour. Fl. Cochinch. 202 = Thunbergii.
triquetrum, Sebast. & Maur. Prod. Fl. Rom. 126 = pendulinum.
tristis, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845). —Orient.
trinulcum, Hort. Genev. ex Schult. f. Syst. vii. 1067 = glaucum.
Tschonoskianum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 160.—Japon.
Tschtschjas, *Regel, l. c.* 107.—Turkestan.
tuberosum, *Rottl. ex Spreng. Syst. i.* 38.—Malabar.
tuberosum, Roxb. Hort. Beng. 34 = odorum.
tulipaefolium, Ledeb. Fl. Alt. ii. 9; Ic. t. 87 = decipiens.
turkestanicum, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 197.—Turkestan.
turtschicum, *Regel, l. c.* vi. (1879) 529.—Aa. centr.
tythocephalum, *Schult. f. Syst.* vii. 1183.—Sibiria.
uliginosum, G. Don, in Mem. Wern. Soc. vi. (1827) 60 = odorum.
uliginosum, Kanitz, in Linnaea, xxvii. (1863) 399 = angulosum.
uliginosum, Ledeb. Fl. Alt. ii. 16 = Ledebourianum.
umbellatum, Hall. ex Stend. Nom. ed. I. 29 = odorum.
umbilicatum, *Boiss. Diagn.* Ser. II. iv. 118.—Belochistan.
undulatum, Fisch. ex Stend. Nom. ed. I. 29 = caspicum.
undulatum, Schousb. ex Trev. All. Sp. 8. n. 9 = nutans.
unifolium, *Kellogg, in Proc. Calif. Acad. Nat. Sc.* ii. (1863) 113. f. 85.—Calif.
uratense, *Franch. in Nouv. Arch. Mus. Hist. Nat. Par.* Sér. II. vii. (1884) 119 (*Pl. Davidian.* i. 304).— Aa. or.
urceolatum, *Regel, Gartenfl.* (1878) 336; *et in Act. Hort. Petrop.* ii. (1873) 405.—Turkestan.
urceolatum, G. Don, ex Sweet, Hort. Brit. ed. I. 422 = ursinum.
ursinum, *Linn. Sp. Pl.* ed. I. 300.—Europ.; As. bor.
Ursulini, Heldr. & Sart. ex Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 238 = staticiformis.
vaginatum, Pall. ex Roem. Fl. Orient. v. 266 = Cupani.
vaginatum, Rent. in Comptes-rend. Soc. Hallér. (1854-56) 126 = flavum.
valdevirum, Nym. Consp. 749 = flavum.

ALLIUM :—
validum, *S. Wats. in Bot. King, Exp.* 250.—Am. bor.
variegatum, *Boiss. Diagn.* Ser. I. vii. 116.—Assyria.
venustiflorum, *Griseb. ex Tchihat. Asie Min. Bot.* ii. 555.—As. Min.
vernale, Tineo, in Guss. Fl. Sic. Prod. Suppl. i. 96 = subvillosum.
veronense, Linn. Syst. ed. X. 978 = paniculatum.
veronense, Spreng. Pugill. ii. 80 = sphaerocephalum.
verrucosum, G. Don, in Mem. Wern. Soc. vi. (1827) 20 = rotundum.
vernicolor, *New. ex Chaub, in Actes Linn. Soc. Bord.* xix. (1853) 50.—Europ.
verticillatum, *Regel, in Act. Hort. Petrop.* vi. (1875) 518.—Turkestan.
Victorialis, *Linn. Sp. Pl.* 295.—Europ.; Caucas.; Sibir.
Vincetoxicum, Pall. ex Ledeb. Fl. Ross. iv. 186 = ursinum.
vineale, Sch. Bip. ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 46 = sphaerocephalum.
vineale, *Linn. Sp. Pl.* 299.—Europ.
violaceum, Gilib. Exercit. ii. 470 = arenarium.
violaceum, Wall. ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 143 = Wallichii.
violaceum, Willd. Enum. Suppl. 17 = carinatum.
virescens, Guss. in Tenore, Syll. 164 = margaritaceum.
virescens, Lam. Fl. Fr. iii. 259 = oleraceum.
virens, Lam. Encyc. i. 67 = oleraceum.
viridi-album, Tineo, Cat. Pl. Hort. Panorm. (1827) 275 = sphaerocephalum.
viridulum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 855 = delicatulum.
viridulum, *Ledeb. Ic. Fl. Alt.* iii. t. 278.—Sibir. altaic.
viviparum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 852.—Sibiria.
volkynicum, Bess. Cat. Hort. Crem. Suppl. iii. 2 = strictum.
vulcanicum, Boiss. Diagn. Ser. I. xiii. 88 = rubellum.
Waldsteinianum, Schult. f. Syst. vii. 1011 = seq.
Waldsteini, G. Don, in Mem. Wern. Soc. vi. (1827) 17 = rotundum.
Wallichianum, Hort. Leichtl. ex Regel, in Act. Hort. Petrop. iii. 11. (1875) 282 = acuminatum.
Wallichianum, *Stend. Nom.* ed. II. i. 58.—Reg. Himal.
Wallichii, Kunth, Enum. Pl. iv. 443 = praec.
Webbii, Clem. in Mem. Acc. Sc. Torino, Ser. II. xvi. (1857) 337. t. 8 = flavum.
Weissii, *Boiss. Fl. Orient.* v. 268.—Ins. Cyclad.
Welwitschii, *Regel, in Act. Hort. Petrop.* iii. 11. (1875) 66.—Lusit.
Wesselowskyi, *Regel, l. c.* vi. (1879) 561.—Turkestan.
Wiedemannianum, *Regel, l. c.* iii. 11. (1875) 199.— Anatolia.
Wierzbicki, *Heufel, ex Steud. Nom.* ed. II. i. 55.— Hungar.
Willdenowii, Kunth, Enum. Pl. iv. 452 = delicatulum.
Wilsii, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 232 = A. Ampeloprasum.
Winklerianum, *Regel, in Act. Hort. Petrop.* viii. (1884) 661.—Bokhara.
xanthicum, Griseb. & Schrenk, in Wiegm. Archiv, xviii. (1852) i. 358 = globosum.
zebdanense, *Boiss. & Not, Diagn.* Ser. II. iv. 118.— Syria; Armenia.

ALYMANIA, R. Br. in Wall. Cat. n. 6890-92 (1832). *AMARANTACEAE*, Benth. & Hook. f. iii. 27.
albida, *R. Br. l. c.* n. 6891.—Ind. or.; Malaya.
aspera, Wight, Ic. t. 1770 = nodiflora.
dichotoma, Wight, l. c. t. 1771 = nodiflora.
saculenta, R. Br. in Wall. Cat. n. 6892 = nodiflora.
nodiflora, *R. Br. l. c.* n. 6890.—Ind. or.; Malaya.
procumbens, Wight, ex Hook. f. Fl. Brit. Ind. iv. 717 = nodiflora.

ALLOBIA, Rafin. Fl. Tellur. iv. 116 (1836) = Euphorbia, Linn.
portlandica, Rafin. l. c. = E. portlandica.

ALLOBIUM, Miers, in Ann. Nat. Hist. Ser. II. viii. (1851) 178, 179 = Phoradendron, Nutt. (Loranth.).

ALLOBORGIA, Stend. Nom. ed. II. i. 55 (1840) = seq.

ALLOBORGIA, Tratt. Fl. Austriac. ii. 71 (1799) = Paradisea, Mazz. (Liliac.).

ALLOCARPUS, H. B. & K. Nov. Gen. et Sp. iv. 291. t. 405 (1820) = Calea, Linn. (Compos.).
caracasanus, H. B. & K. l. c. 292.—Venezuela.
integrifolius, DC. Prod. v. 676 = C. integrifolia.
Lindenii, Sch. Bip. ex Wedd. Chlor. And. i. 74.—N. Granat.
sabaioides, Less. Syn. Comp. 342 = C. sabaioides.
scabrifolius, Hook. & Arn. Bot. Beech. Voy. 300 = C. scabrifolia.

ALLOCERATIUM, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 129, 135 = Diptychocarpus, Trautv. (Crucif.).
strictum, Hook. f. & Thoms. l. c. 135 = D. strictus.

ALLOCHLAMYS, Moq. in DC. Prod. xiii. ii. 468 (1849) = Pleuropetalum, Hook. f. (Amarant.).
Darwinii, Moq. l. c. = P. Darwinii.

ALLOCHRUSA, Bunge, in Boiss. Fl. Orient. i. 559 (1867) = Acanthophyllum, C. A. Mey. (Caryophyll.).
Bungei, Boiss. l. c. 560 = Ac. Bungei.
persica, Boiss. l. c.—Persia.
pulchella, Bunge, ex Boiss. l. c. = *persica*.
versicolor, Boiss. l. c. 559 = Ac. versicolor.

ALLODAPE, Endl. Gen. 749 (1839) = Lebetanthus, Endl. (Epacrid.).
americana, Endl. l. c. = L. americanus.

ALLODAPHNE, Steud. Nom. ed. II. i. 55 (1840) = praec.

ALLOEOSPERMUM, Spreng. Anleit. ii. 901 (1818) = ALLOISPERMUM, Willd. = Calea, Linn. (Compos.).

ALLOHEMIA, Rafin. Sylva Tellur. 125 (1838) = Loranthus, Linn.
pedunculata, Rafin. l. c. = L. pedunculatus.
purpurea, Rafin. l. c. = L. occidentalis.

ALLOIANTHERUS, Stend. Syn. Pl. Gram. 217 (1855) = seq.

ALLOIATHEROS, Ell. Sketch. i. 146 (1817) = Gymnopogon, Beauv. (Gramin.).
ambiguus, Ell. (cf. Sketch, i. 146) ex Kunth, Enum. i. 284 = G. racemus.
aristatus, Rafin. in Ser. Bull. Bot. i. (1830) 221 = G. racemosus.
lepturoides, Steud. Nom. ed. II. i. 55 = G. racemosus.

ALLOIOZONIUM, Kunze, in Linnaea, xvii. (1843) 572 = Cryptosetamina, R. Br. (Compos.).
arctioides, Kunze, l. c. = C. calendulaceum.

ALLOISPERMUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 139 = Calea, Linn. (Compos.).
divaricatum, Willd. l. c. (= *Alocarpus caracasanus*).

ALLOMIA, DC. Prod. vii. 310 (1838) = Alomia, H. B. & K. (Compos.).

ALLOMORPHIA, Blume, in Flora, xiv. (1831) 822. *MELASTOMACEAE*, Benth. & Hook. f. i. 752.
exigua, *Blume, l. c.* 828.—Ind. or.; Malaya.
Griffithii, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 74.—Malacca.
hispida, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 58.—Burma.
ovalifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 74.—Ins. Fiji.
pauciflora, Benth. in Hook. Lond. Journ. Bot. i. (1842) 435 = Oxyspora pauciflora!
umbellulata, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 74.—Burma.

ALLOPHYLLUS, Gleditsch, Syst. 89 (1764) = seq.

ALLOPHYLUS, Linn. Fl. Zeyl. 58 (1747); Dis. Linn. 14 (1747); Am. Acad. i. 398 (1749). *SAPINDACEAE*, Benth. & Hook. f. i. 396.
TOXICODENDRUM, Gaertn. Fruct. i. 207. t. 44 (1788). USUBIS, Burm. f. Fl. Ind. 89. t. 28. f. 1. (1768).
africanus, Beauv. Fl. Owar. ii. 54. t. 107 = serjanioides.
amboinensis, *Blume, Rumphia*, iii. 129.—Ambolna.
aporeticus, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 185.—Burma.
Blancoi, Blume, Rumphia, iii. 129 = A. Cobbe. Cambesedei, *Blume, l. c.*—Malaya.

ALLOPHYLUS :—
celebicus, *Blume, Rumphia*, iii. 128.—Ins. Celebes.
Cobbe, *Blume, l. c.* 131.— An. or.
Cominia, Sw. Prod. Veg. Ind. Occ. 62 = S. Cominia.
Commersoni, *Blume, Rumphia*, iii. 129.—Ins. Borbon.
dimorphus, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1878) 126.—Ins. Philipp.
fliger, *Radlk. l. c.*—Ins. Philipp.
fulvinervis, *Blume, Rumphia*, iii. 183.—Java.
integrifolius, *Blume, l. c.* 129.— Malaya.
javensis, *Blume, l. c.* 126.—Java.
lanatus, Roxb. Hort. Beng. 88 = A. Cobbe.
leptococcus, *Radlk. in Act Congr. Bot. Amst.* 1877 (1879) 136.—Ins. Key.
ligustrinus, *Blume, Rumphia*, iii. 126.—Sumatra.
litoralis, *Blume, l. c.* 124 —Java.
nigrescens, *Blume, l. c.* 130.—Madag.
ornithrophiodes, Roxb. ex Wall. Cat. n. 8055 = A. Cobbe.
Pervillei, *Blume, Rumphia*, iii. 128.—Madag.
pinnatus, Roxb. Ic. t. 1407 = Sapindus fruticosus.
racemosus, Sw. Prod. Veg Ind. Occ. 62 = S. occidentalis.
rhoaiphyllus, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 507.—Ins. Socotra.
rigidus, Sw. Prod. Veg. Ind. Occ. 62 = S. rigida.
rufescens, *Blume, Rumphia*, iii. 159.—Borneo.
rugosus, *Blume, l. c.* 138.—Java.
sahgnus, *Blume, l. c.* 129.—Madag.
serratus, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 185.—Burma.
sessilis, *Blume, Rumphia*, iii. 125.— Java.
sumatranus, *Blume, l. c.* 132.—Sumatra.
suvdanus, Miq. Fl Ind. Bat. I. ii. 575 = A. Cobbe.
ternatus, Lour. Fl. Cochinch. 282 = A. Cobbe.
ternatus, Roxb ex Steud. Nom. ed. II. i. 55 = S. zeylanicus.
timorensis, *Blume, Rumphia*, iii. 130.—Timor.
timorensis, Miq. Fl. Ind. Bat. I. i 575 = A. Cobbe.
zeylanicus, *Linn. Sp. Pl.* 848.—Himal.

ALLOPLECTUS, Mart. Nov. Gen. et Sp. iii. 58. t. 228 (1829). *GESNERACEAE*, Benth. & Hook. f. ii. 1008.
ANISOPLECTUS, CALANTHUS, CALOPLECTUS, CALYCOPLECTUS, CORYTOPLECTUS, ERYTHRANTHUS, PRIONOPLECTUS, SACCOPLECTUS, Oerst. in Danske Vid. Selsk. Skr. V. v. (1858) 117-121.
CRANTZIA, Scop. Introd. 178 (1777).
HEINTZIA, Karst. Ausw. Gew. Venez. 34. t. 11. (1848).
GLOSSALOMA, Hanst. in Linnaea, xxvi. (1853-55) 209.
LOPHIA, Desv. in Ham. Prod. Fl. Ind. Occ. 47 (1825).
LOPHALIX, Rafin. Sylva Tellur. 70 (1838).
MACROCHLAMYS, Decne. in Ann. Hortic. (1849) 248.
OROBANCHIA, Vand. Fl. Lusit. Spec. 41. f. 18, 19 (1788).
POLYSTHYRANIA, Hanst. ex Oerst. in Danske Vid. Selsk. Skr. V. v. (1858) 121.
PTERYGOLOMA, Hanst. in Linnaea, xxvi. (1853-1855) 211.
TYSSACIA, Reichb. ex Steud. Nom. ed. II. ii. 797 (1841).
VIREYA, Rafin. Specch. i. 194 (1814).
angustifolius, *DC. Prod.* vii. 546.—Bras.
bicolor, *Klotzsch, ex Hanst. in Linnaea*, xxxiv. (1865–66) 377.—Bras.
bracteatus, *Linden, Cat.* n. 5 (1850) 2.—Hab.?
calicerastus, *C. Don, Gen. Syst. iv.* 655.—Neg. Andin.
capitatus, *Hook. Bot. Mag.* t. 4452.—Am. austr.
chrysanthus, *Planch. & Linden, in Fl. des Serres*, Sér. I. viii. (1852–53) 211.—N. Granat.
circinnatus, *Mart. Nov. Gen. et Sp.* iii. 58. t. 228. f. 1. —Bras.
coccineus, *Mart. l. c.* 57.—Guiana.
concolor, Hook. Bot. Mag. t. 4371 = dichrous.
coriaceus, *Hanst. in Linnaea*, xxxiv. (1865–66) 368.—Costa Rica.
corymbosus, *Klotzsch, ex Hanst. l. c.* 371.—Peruv.
cristatus, *Mart. Nov. Gen. et Sp.* iii. 57.—Ind. occ.
dariensenis, *Seem. Bot. Voy. Herald*, 187.—Darien.
dichrous, *DC. Prod.* vii. 546.—Bras.
dichrous, Hook. Bot. Mag. t. 4216 = sparsiflorus.
dimidiatus, *Benth. Pl. Hartw.* 235.—N. Granat.
eriocalyx, Galeott. ex Hook. Bot. Mag. t. 4371 = dichrous.
Forsethii, *Hanst. in Linnaea*, xxxiv. (1865–66) 375. —Am. centr.

ALLOPLECTUS :—
glaber, *DC. Prod.* vii. 546.—Mexic.
hispidus, *Mart. Nov. Gen. et Sp.* iii. 57.—Am. austr.
ichthyoderma, *Hanst. in Linnaea*, xxxiv. (1865–66) 372.—Costa Rica.
Kunthii, *G. Don, Gen. Syst.* iv. 655.—N. Granat.
lanatus, *Seem. Bot. Voy. Herald*, 186.—Panama.
macrophyllus, *Hemsl. Biol. Centr. Am. Bot.* ii. 484. —Costa Rica.
melittifolius, Mart. ex Loud. Encyc. Pl. 1402 = Tussacia pulchella.
meridensis, *Klotzsch, ex Hanst. in Linnaea*, xxxiv. (1865–66) 376.—N. Granat.
mollifolius, *Hanst. l. c.* 361.—Costa Rica.
parviflorus, *Hemsl. Biol. Centr. Am. Bot.* ii. 484.— Costa Rica.
Patrisii, *DC. Prod.* vii. 545 —Guiana.
peltatus, *Oliver, in Bot Mag. t.* 6383.—Am. centr.
pendulus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 6. t. 205.—Peruv.
Pinelianus, *Lem. in Fl. des Serres*. Sér. I. ii.(Août 1846) t. v.—Bras.
repens, *Hook. Bot. Mag.* t. 4250.—Ind. occ.
sanguineus, *Mart. Nov. Gen. et Sp.* iii. 57.—Am. austr.
scabridus, Lem. Jard. Fleur. i. (1851) Misc. 51 = glaber.
Schlimii, *Planch. & Linden, in Fl. des Serres*, Sér. I. viii. (1852–53) 211. t. 837.—N. Granat.
Schottii, G. Don, Gen. Syst. iv. 655 = dichrous.
semicordatus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 5. —Peruv.
spathiflorus, *Mart. Nov. Gen. et Sp.* iii. 55.—Bras.
speciosus, Hort. ex Hook. Bot. Mag. t. 4452 = capitatus.
speciosus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 6.— Chili.
strigosus, *Hanst. in Linnaea*, xxxiv. (1865–66) 374.— Mexic.
tenuis, *Benth. Pl. Hartw.* 235.—Peruv.
tetragonus, *Hanst. in Linnaea*, xxxiv. (1865–66) 368. —Costa Rica.
tigrinus, *Hanst. l. c.* 366.—N. Granat.
villosus, *Hanst. in Mart. Fl. Bras.* viii. 407.—Bras.
vittatus, *André, in Illust. Hortic.* xvii. (1870) t. 13.— Peruv.
zanonioides, *Linden & André, in Illust. Hortic.* xix. (1872) 110.—N. Granat.

ALLOPLEIA, Rafin. Med. Fl. ii. 109 (1830) = **Veronica**, Tourn. (Scrophul.).
rotundifolia, Rafin. l. c. = V. rotundifolia.

ALLOPYTHION, Schott, Gen. Aroid. t. 24 (1858) = **Thomasnia**, Wall. (Aroid.).
Hookeri, Schott, l. c. = T. Hookeri.

ALLOSAMPELA, Rafin. Med. Fl. ii. 122 (1830) = **Vitis**, Linn. (Ampel.).
heterophylla, Rafin. l. c. = V. heterophylla.

ALLOSCHEMONE, Schott, Prod. Syst. Aroid. 358 (1856) = **Monstera**, Adans. (Aroid.).
Poeppigiana, Schott, l. c. = M. occidentalis.

ALLOSPERMA, Rafin. Fl Tellur. iv. 122 (1836) = **Commelina**, Linn.
tuberosa, Rafin. l. c. = C. tuberosa.

ALLOSTIS, Rafin. Sylva Tellur. 104 (1838) = **Baeckea**, Linn. (Myrtac.).

ALLOTROPSIS, C. Presl, Rel. Haenk. i. 348. t. 47 (1830). Gen. dub. *GRAMIN.* Benth. & Hook. f. iii. 1096.
distachya, *C. Presl, l. c.*—Calif.

ALLOTRIA, Rafin. Fl. Tellur. iii. 70 (1836) = **Commelina**, Linn.
scabra, Rafin. l. c. = C. scabra.

ALLOTROPA, A. Gray, in Pacif. Rail. Rep. Bot. iv. 81 (1855); et in Proc. Am. Acad. vii. (1868) 368. *MONOTROPEAE*, Benth. & Hook. f. ii. 607.
virgata, *Torr. & Gray, in Proc. Am. Acad. l. c.* —Calif.

ALLOUVA, Plum. ex Aubl. Fl. Gui. ii. 921 (1775) = **Maranta**, Linn. (Scitam.).

ALLOZYGIA, Nasd. in Ann. Sc. Nat. Sér. III. xv. (1851) 309. t. 15 = **Oxypora**, DC. (Melast.).
cernua, Nasd. l. c. = O. cernua.

ALLUGHAS, Steud. Nom. ed. I. 29 (1821) = ALUGHAS, Lindl. = **Alpinia**, Linn. (Scitam.).

ALMANA, Rafin. Fl. Tellur. iv. 68 (1836) = **Gloxinia**, Linn. (Gesner.).
hirsuta, Rafin. l. c. = G. hirsuta.

ALMEIDA, Cham. in Linnaea, v. (1830) 56 = seq.

ALMEIDEA, St. Hil. in Bull. Soc. Philom. (1828) 129. *RUTACEAE*, Benth. & Hook. f. i. 384.
ARUSA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 102. t. 19, 27, 29.
acuminata, *St. Hil. ex G. Don, Gen Syst.* i. 798.—Bras.
alba, *St. Hil. ex G. Don, l. c.*—Bras.
caerulea, *St. Hil. ex G. Don, l. c.*—Bras.
lanceolata, *St. Hil. Pl. Rem.* 144.—Bras.
lilacina, *St. Hil. in Bull. Soc. Philom.* (1828) 129.— Bras.
longifolia, *St. Hil. l. c.*—Bras.
longipes, *Presl, Bot. Bemerk.* 82.—Bras.
macropetala, *Fisch. & Mey. Sert. Petrop.* i. 10.—Bras.
myrtantha, *Cham. in Linnaea*, v. (1830) 56.—Bras.
rubra, *St. Hil. in Bull. Soc. Philom.* (1828) 129.—Bras.

ALMELOVEENIA, Dennst. Schluss. Hort. Malab. 32 (1818) = **Moullava**, Adans. (Inc. Sed.).
spinosa, Dennst. l. c.—Ind. or.

ALMIDEIA, Reichb. Nom. 197 (1841) = **Almeidea**, St. Hil. (Rutac.).

ALMYRA, Salisb. in Trans. Hort. Soc. i. (1812) 336 = **Pancratium**, Linn. (Amaryll.).
illyrica, Salisb. Gen. Pl. Fragm. 100 = P. illyricum.
parviflora, Salisb. l. c. = P. parviflorum.
stellaris, Salisb. in Trans. Hort. Soc. i. (1812) 336 = P. illyricum.

ALNASTER, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 200 = Alnus, Tourn. (Cupul.).
fruticosus, Ledeb. Fl. Ross. iii. 655 = Alnus viridis.
viridis, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 201 = Alnus viridis.

ALNOBETULA, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858); (Enum. Pl. Trans.) 68 = **Alnus**, Tourn. (Cupul.).
viridis, Schur, l. c. = Alnus viridis.

ALNUS, [Tourn.] Linn. Syst. ed. II. (1735); Fl. Lapp. 260 (1737). *CUPULIFERAE*, Benth. & Hook. f. iii. 404.
ALNASTER, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 200.
ALNOBETULA, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858); (Enum. Pl. Trans. 68. n. 614).
CLETHROPSIS, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 201.
SKINDOPSIS, Zumag. Fl. Pedem. i. 249 (1849).
acuminata, *H. B. & K. Nov. Gen. et Sp.* ii. 20.—Am. austr.
Alnobetula, C. Koch, Dendrol. ii. i. 625 = viridis.
alpina, Borckh. Handb. Forstb. i. 477 = viridis.
americana, Hort. ex C. Koch, Dendrol. ii. i. 636 = serrulata.
arguta, *Benth. Pl. Hartw.* 348.—Mexic.
argutu, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 205 = acuminata.
aurea, Hort. ex C. Koch, Dendrol. ii. i. 630 = glutinosa.
autumnalis, Hartig, ex Garcke, Fl. Deutschl. ed. IV. 305 = serrulata.
badensis, Lange, ex Regel, in DC. Prod. xvi. ii. 168 = glutinosa.
barbata, Fries, ex Nym. Consp. 671 = pubescens.
barbata, C. A. Mey. Verz. Pfl. Cauc. 42 = glutinosa.
brembana, Rota, Frosp. Prov. Bergamasco, 79 = viridis.
canadensis, Lodd. ex Loud. Hort. Brit. 878 = serrulata.
castanaefolia, *Mirb. in Mém. Mus. Par.* xiv. (1827) 468. t. 21.—Peruv.
cordifolia, Hartig, ex Regel, in DC. Prod. xvi. ii. 167 = glutinosa.
communis, Desf. Tabl. Hort. Par. ed. I. 318 = glutinosa.
cordata, Desf. l. c. ed. II. 344 = cordifolia.
cordifolia, *Tenore, Fl. Nap. Prod.* 54. t. 99.—Europ.; Orient.
corylifolia, *Kern. ex Dalla Torre, Atl. Alpenfl.* 189.— Europ.

ALNUS :—
crispa, Hort. ex Steud. Nom. ed. II. i. 55 = undulata.
crispa, *Pursh, Fl. Am. Sept.* ii. 181.—Am. bor.
cuneata, Geyer, ex Walp. Ann. i. 386 = Cercocarpus parvifolius.
denticulata, C. A. Mey. Verz. Pfl. Cauc. 43 = glutinosa.
dioica, Roxb. Fl. Ind. iii. 580 = Aporosa Roxburghii.
dubia, Req. ex Regel, in DC. Prod. xvi. ii. 187 = glutinosa.
elliptica, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 381 = glutinosa.
emarginata, Krock. ex Steud. Nom. ed. I. 29 = glutinosa.
februaria, Kuntze, Taschen. Fl. Leipz. 238 = glutinosa.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* ii. 17.—N. Granat.
firma, *Sieb. & Zucc. in Abh. Acad. Muench.* iv. iii. (1846) 220.—Japon.
fruticosa, Rupr. Beitr. Pfl. Russ. Reich. ii. 58 = viridis.
glauca, Michx. f. Arb. Am. iii. 220. t. 4. f. 2 = incana.
glutinosa, *Medic. Pfl. Anat.* 393.—Europ.; As. bor.
Harinoli, Siebold, in Verh. Gen. Bat. xii. (1830) 25 = japonica.
hirsuta, Turcz. in Bull. Soc. Nat. Mosc. (1838) 101; xxvii. (1854) i. 406 = incana.
hybrida, Reichb. Ic. Fl. Germ. xii. t. 680. f. 1292 = glutinosa, incana.
incana, *Medic. Pfl. Anat.* 393.—Reg. bor. temp.
incisa, Steud. Nom. ed. I. 29 = glutinosa.
integrifolia, Roxb. ex Steud. Nom. ed. II. i. 56 = Aporosa Roxburghii.
intermedia, Hort. ex C. Koch, Dendrol. ii. i. 681 = oblongata.
intermedia, Schrad. ex Regel, in DC. Prod. xvi. ii 189 = incana.
japonica, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1845) 230 = maritima.
japonica, Steud. Nom. ed. II. i. 55 = Betula alba.
jorullensis, Benth. Pl. Hartweg. 72 = acuminata.
jorullensus, *H. B. & K. Nov. Gen. et Sp.* ii. 20.—Mexic.
laciniata, Ehrh. Beitr. iii. 22 = glutinosa.
lanuginosa, Gilib. Exercit. ii. 402 = incana.
latifolia, Desf. Cat. Hort. Par. ed. III. 352 = serrulata.
Lindeni, *Regel, in Nouv. Mém. Soc. Nat. Moac.* xiii. (1861) 144.—As. temp.
lobata, Nym. Consp. 671 = incana.
longifolia, Bové, ex Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 346 = orientalis.
macrocarpa, Lodd. ex Loud. Hort. Brit. 378 = glutinosa.
macrocarpa, Req. ex Nym. Consp. 672 = cordifolia.
macrophylla, Desf. ex Steud. Nom. ed. I. 29; ed. II. i. 56 = serrulata.
maritima, *Nutt. Syle. Am. Suppl.* i. 84. t. 10 (1842).—Am. bor.
microphylla, *Avret, in Bull. Soc. Bot. Fr.* xxvi. (1879) *Bibl.* 89.—Europ.
Mirbellii, *Spach, in Ann. Sc. Nat. Sér.* II xv. (1841) 204.—Am. austr.
Michelianus, M. A. Curtis, ex A. Gray, in Am. Journ. Sc. xlii. (1842) 42 = crispa.
montana x, *Brugg. in Jahresb. Naturf. Ges. Graub.* II. xxiii.-xxiv. (1880) 115.—Europ.
Moriseana, Bertol. Fl. Ital. x. 168 = glutinosa.
nana, [*Claire.*] *Man. Herb.* 280.—Europ.
neapolitana, Savi, Tratt. Alb. Tosc. ed. II. ii. 21 = cordifolia.
nepalensis, *D. Don, Prod. Fl. Nep.* 58.—Reg. Himal.
nigra, Gilib. Exercit. ii. 401 = glutinosa.
nitens, C. Koch, in Linnæa, xxii. (1849) 324 = glutinosa.
nitida, *Endl. in Gen. Suppl.* iv. ii. 20.—Reg. Himal.
obcordata, C. A. Mey. ex Steud. Nom. ed. II. i. 56, sphalm. = cordifolia.
oblongata, Kotschy, ex Regel, in DC. Prod. xvi. ii. 188 = orientalis.
oblongata, Regel, Monog. 118. t. 7. f. 8–9 = maritima.
oblongata, *Willd. Sp. Pl.* iv. 335.—Europ. austr.
oblongifolia, *Torr. Bot. Mex. Bound.* 204.—N. Mexic.
oregana, Nutt. Sylv. i. 9 = rubra.
orientalis, *Decne. in Ann. Sc. Nat. Sér.* II. iv. (1835) 348.—Orient.
ovata, Lodd. Bot. Cab. t. 1141 = crispa.
oxyacanthifolia, Lodd. Cat. (1836) = glutinosa.
pinnata, Hort. ex C. Koch, Dendrol. ii. i. 688 = incana.
prunifolia, Hort. ex C. Koch, l. c. 680 = glutinosa.
pubescens, Tausch, in Flora, xvii. (1834) 520 = glutinosa x incana.
pumila, *Steud. Nom.* ed. I. 29.—Am. bor.

ALNUS :—
quercifolia, Willd. Berl. Baumz. 45 = glutinosa.
repens, Wormsk. Fl. Dan. t. 2736 = viridis.
rhombifolia, *Nutt. Sylva*, i. 83.—Calif.
rotundifolia, Bertol. Fl. Ital. x. 160 = cordifolia.
rotundifolia, Stokes, Bot. Mat. Med. iv. 369 = glutinosa.
rubra, *Bongard, in Mém. Acad. Petersb.* Sér. VI. ii. (1833) 162.—Am. bor. occ.
rubra, Desf. ex Steud. Nom. ed. I. 29 = serrulata.
rufescens, Liebm. ex Hemsl. Biol. Centr. Am. Bot. iii. 165 = acuminata.
rugosa, Hort. ex Tausch, in DC. Prod. xvi. ii. 188 = serrulata.
rugosa, Spreng. Syst. iii. 848 = acuminata.
serrulata, *Willd. Sp. Pl.* iv. 336.—Am bor.
sibirica, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. (1838) 101; xxvii. (1854) i. 406 = incana.
suaveolens, Mor. ex Nym. Consp. 671 = glutinosa.
suaveolens, *Req. in Ann. Sc. Nat.* v (1825) 381.—Europ.
subcordata, C. A. Mey. Verz. Pfl. Cauc. = cordifolia.
subrotunda, *Desf. Tabl. Hort. Par.* ed. II. 244.—Hab.?
tiliæfolia, Hort. ex C Koch, Dendrol. ii. i. 634 = subcordata.
tenuifolia, *Nutt. Sylva*, i. 10.—Am. bor.
tirolensis, Nym. Consp. 671 = incana.
tomentosa, Harug, ex Regel, in DC. Prod. xvi. ii. 185 = orientalis.
tristis, Wormsk. ex Regel, l. c. 189 = viridis.
undulata, *Willd. Sp. Pl.* iv. 336.—Canada.
Vilmoriana, *Hort. ex Lebas, in Rev. Hortic.* (1866) 446.—Hab.?
viridis, *DC. Fl. Fr.* iii. 304; *Cham. in Linnæa*, vi. (1831) 538.—Europ.; As. bor.; Am. bor.
viridis, A. Gray, Man. Bot. N. Un. St. (1848) 424 = crispa.
vulgaris, Pers. Syn. ii. 550 = glutinosa.

ALOASIA, Neck. Elem. iii. 289 (1790); Schott, in Schott & Endl. Melet. 18 (1832). *AROIDEAE*, Benth. & Hook. f. iii 975.
ENSOLENANTHE, Schott, in Bonplandia, ix. (1861) 303.
acuminata, *Schott, in Bonplandia*, vii. (1859) 28.—Ind. or.
alba, *Schott, in Oestr. Bot. Wochenbl* ii (1852) 59; (1854) 410.—Zeylan.
argyronevra, C. Koch, Berl. Wochenschr. (1861) 8 = Caladium Schomburgkii.
Becarii, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 300.—Malaya.
communata, Schott, in Oestr. Bot. Wochenbl. (1854) 409 = Colocasia odora.
cucullata, *Schott, in Schott & Endl. Melet.* 18.—Ind. or.
cuprea, *C. Koch, in Berl. Wochenschr.* iv. (1861) 141; vii. (1864) 90.—Borneo.
decipiens, *Schott, in Bonplandia*, vii. (1859) 28.—Ind. or.
denudata, *Engl. in DC. Monog. Phan.* ii. 507.—Singapore.
erythrosa, C. Koch, in Berl. Wochenschr. (1861) 8 = Colocasia Schomburgkii.
fallax, Schott, in Bonplandia, vii. (1859) 28 = navicularis.
fornicata, *Schott, in Oestr. Bot. Wochenbl.* (1854) 410.—Ind. or.
grandis, *Miquel, in Rev. Hortic.* (1868) 380.—Hab.?
heteroneura, C. Koch, in Berl. Wochenschr. (1861) 283 = Caladium Schomburgkii.
indica, *Schott, in Oestr. Bot. Wochenbl.* (1854) 410, pro parte.—Ind. or.; Malaya.
Jenningsii, Hort. Veitch. Cf. Gard. Chron. (1868) 26; in Illustr. Hortic. (1869) t. 585 = Colocasia affinis.
Johnstonii, Hort. in Gard. Chron. xiii. (1880) 759; in Illustr. Hortic. xxvii. (1880) t. 395 = Cyrtosperma Johnstonii.
Korthalsii, *Schott, in Ann. Mus. Lugd. Bat.* i. 124.—Borneo.
Liervalli, *Hort. ex Hérincq, L'Hortic. Franç.* (1869) 248.—Ins. Philip.
lobata, *Rafin. Fl. Tellur.* iii. 64.—Hab.?
longiloba, *Miq. Fl. Ind. Bat.* iii. 207.—Malaya.
Lowii, *Hook. Bot. Mag.* t. 5876.—Borneo.
macrorhiza, *Schott, in Schott & Endl. Melet.* 18.—As. trop.; Austral.; Ins. Pacific.
Marchalli, Engl. in DC. Monog. Phan. ii. 494 = Colocasia Marchalli.
montana, *Schott, in Oestr. Bot. Wochenbl.* (1854) 410.—Ind. or.; Malaya.

ALOCASIA :—
metallica, Hook. Bot. Mag. t. 5190 = cuprea.
metallica, Schott, in Oestr. Bot. Wochenbl. (1854) 410 = indica.
navicularis, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1855) App. 2.—Reg. Himal.
odora, *C. Koch, l. c.* (1854), *App.* 5.—Ind. or.; China.
pallida, C. Koch & Bouché, l. c. = alba.
patula, Rafin. Fl. Tellur. iii. 64 = Sauromatum pedatum.
plumbea, Van Houtte, in Fl. des Serres, Sér. II. xi. (1875) 93. t. 2206 = indica.
porphyroneura, Lem. in Illustr. Hortic. (1861) t. 297 = Caladium picturatum.
Portei, *Berc. & Engl. in DC. Monog. Phan.* ii. 645.—N Guin.
pubera, *Schott, Syn. Aroid.* 47.—Java.
Putreys, *N. E. Br. in Illustr. Hortic.* xxix. (1882) t. 489, & p. 22.—Sumatra.
rapiformis, *Schott, Prod Aroid.* 157.—Burma.
Regina, *N E Br.—Cf. Gard. Chron.* (1884) i. 711.—Borneo.
Roezlii, Hort. ex Gard. Chron. iii. (1875) 589 = Caladium marmoratum.
rugosa, Schott, in Oestr. Bot. Wochenbl. (1854) 410 = cucullata.
scabriuscula, *N. E. Br. in Gard. Chron.* (1879) ii. 296.—Borneo.
Sedeni x, Hort. Veitch.—Cf. Gard. Chron. (1869) 587, 615 = cuprea x Lowii.
sinuata, *N. E. Br. in Gard. Chron.* (1885) ii. 678.—Malaya.
Thibautiana, *Mast. l. c.* (1878) t. 827.—Borneo.
undipes, C. Koch & Bouché, in Ind. Sem. Hort. Berol. (1854), App. 5 = Xanthosoma Jacquini.
variegata, C. Koch & Bouché, l. c. = indica.
Veitchii, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 125 = Lowii.
virginica, Rafin. Fl. Tellur. iii. 64 = Peltandra virginica.
zebrina, *C. Koch & Veitch, in Berl. Wochenschr.* vi. (1868) 381.—Ins. Philipp.

ALOË, Tourn. ex Linn. Syst. ed. I. (1735). *LILIACEAE*, Benth. & Hook. f. iii. 776.
AGRIODENDRON, Endl. Gen. 144 (1837).
BOWIEA, Haw. in Phil. Mag. (1824) 399.
BUSIPIO, Rafin. Fl. Tellur. iii. 76 (1866).
CATEVALA, Medic. Theodora, 67 (1786).
KUMARA, Medic. l. c. 69. t. 4 (1786).
PACHIDENDRON, Haw. Rev. Pl. Succ. 85 (1821).
PITYAS, Salisb. Gen. Pl Fragm. 76 (1866).
RHIPIDODENDRON, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 164.
abyssinica, *Lam. Encyc.* i. 86.—Abyss.
acinacifolia, Jacq. Ecl. 49. t. 31 = Gasteria acinacifolia.
acuminata, Haw. Syn. Pl. Succ. 84 = A. humilis.
acuminata, Lam. Encyc. i. 90 = Gasteria verrucosa.
africana, *Mill. Gard. Dict.* ed. VIII. n. 4.—Afr. austr.
africana, Salm-Dyck, Cat. 31 = Salm-Dyckiana.
agavefolia, *Tod. Hort. Bot. Panorm.* i. 55. t. 28.—Afr. trop.
albicans, Haw. in Trans. Linn. Soc. vii. (1804) 18 = Haworthia albicans.
albispina, *Haw. l. c.* 22.—Afr. austr.
albocincta, Haw. Suppl. Pl. Succ. 48 = striata.
altilinea, Schult. f. Syst. vii. 638 = Haworthia altilinea.
americana, Cranz, Inst. I. 466 = Agave americana.
andongensis, *Baker, in Trans. Linn. Soc. Sct. II.* i. (1878) 268.—Afr. trop.
angolensis, *Baker, l. c.*—Afr. trop.
angulata, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 276 = Gasteria disticha.
angusta, *Schult. f. Syst.* vii. 710.—Afr. austr.
angustifolia, Salm-Dyck, Obs. 57 = Gasteria disticha.
anomala, Haw. in Trans. Linn. Soc. vii. (1804) 26 = Haworthia recurva.
arabica, *Lam. Encyc.* i. 91.—Arab.
arabica, Salm-Dyck, Cat. 27 = microstigma.
arachnoidea, Mill. Gard. Dict. ed. VIII. n. 17 = Haworthia arachnoidea.
arachnoides, Thunb. Diss. Aloë, 7 = Haworthia arachnoides.
arborea, Medic. Beobacht. (1782) 305 = arborescens.
arborescens, *Mill. Gard. Dict.* ed. VIII. n. 3.—Afr. austr.

ALOË :—
aristata, *Haw. in Phil. Mag.* (Oct. 1825) 280.—Afr. austr.
aristata, *Schult. f. Syst.* vii. 688.—Afr. austr.
aspera, Haw. in Trans. Linn. Soc. vii. (1804) 6 = Apicra aspera.
asperuscula, Schult. f. Syst. vii. 658 = Haworthia asperiuscula.
asperula, Schult. f. l. c. 635 = Haworthia asperula.
Atherstoni, *Baker, in Journ. Linn. Soc.* xviii. (1881) 170.—Afr. austr.
atrovirens, DC. Pl. Grass. t. 51 = Haworthia atrovirens.
attenuata, Haw. in Trans. Linn. Soc. vii. (1804) 11 = Haworthia attenuata.
Baineši, *Dyer, in Gard. Chron.* (1874) i. 567.—Afr. austr.
barbadensis, Mill. Gard. Dict. ed. VIII. n. 2 = vera.
Barberæ, Dyer, in Gard. Chron. (1874) i. 566 = Baineši.
Barteri, *Baker, in Journ. Linn. Soc.* xviii. (1881) 168.—Afr. trop.
Bayfieldii, Salm-Dyck, Monog. ii. § 29. f. 14 = Gasteria Bayfieldii.
bicarinata, Spreng. Syst. ii. 70; Schult. f. Syst. vii. 652 = Apicra bicarinata.
bicolor, Schult. f. l. c. 683 = Gasteria bicolor.
Bolusii, *Baker, in Journ. Linn. Soc.* xviii. (1881) 179.—Afr. austr.
Bowiea, *Schult. f. Syst.* vii. 704.—Afr. austr.
Bowieana, Schult. f. l. c. 662 = Gasteria picta.
brachyphylla, Salm-Dyck, Hort. 383 = Gasteria brevifolia.
Bradleyana, Jacq. Hort. Schoenbr. iv. 11, in nota ad t. 421 = Haworthia atrovirens.
brevifolia, *Mill. Gard. Dict.* ed. VIII. in errat.—Afr. austr.
brevifolia, Haw. in Trans. Linn. Soc. vii. (1804) 32 = distans.
brevioribus, Mill. Gard. Dict. ed. VIII. n. 8, sphalm. = brevifolia.
brevis, *Schult. f. Syst.* vii. 649.—Afr. austr.
bullulata, Jacq. Fragm. 72. t. 109 = Apicra pentagona.
cæsia, *Salm-Dyck, Cat.* 39.—Afr. austr.
Candelabrum, *Tod. Hort. Panorm.* (1876) 46.—Hab.?
candicans, Schult. f. Syst. vii. 681 = Gasteria candicans.
capitata, *Baker, in Journ. Linn. Soc.* xx. (1883) 272.—Madag.
carinata, Ker-Gawl. in Bot. Mag. t. 1381 = Gasteria glabra.
carinata, DC. Pl. Grass. t. 63 = Gasteria verrucosa.
carinata, Mill. Gard. Dict. ed. VIII. n. 21 = Gasteria carinata.
carolinians, Hill, Veg. Syst. xiii. 64 = Agave virginica.
chinensis, *Steud. Nom.* ed. II. i. 56; *Baker, in Bot. Mag.* t. 6801.—China.
chloracantha, Schult. f. Syst. vii. 641 = Haworthia chloracantha.
chlorolenca, *Baker, in Gard. Chron.* (1877) ii. 88.—Afr. austr.
ciliaris, *Haw. in Phil. Mag.* (Oct. 1825) 281.—Afr. austr.
clavipeta, Schult. f. Syst. vii. 646, 1712 = Haworthia attenuata.
clavifiora, *Burch. Trav.* i. 272.—Afr. austr.
coarctata, Schult. f. Syst. vii. 647 = Haworthia coarctata.
Commelyni, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 362 = mitriformis.
commutata, *Tod. Hort. Bot. Panorm.* i. 75. t. 18.—Hab.?
concinna, Schult. f. Syst. vii. 653 = Haworthia viscosa.
congesta, Salm-Dyck, Monog. i. § 2. f. 1 = Apicra congesta.
consobrina, *Salm-Dyck, l. c.* § 18. f. 2.—Afr. austr.
conspurcata, Salm-Dyck, Obs. 59 = Gasteria disticha.
constricta, *Baker, in Journ. Linn. Soc.* xviii. (1881) 168.—Afr. trop.
Cooperi, *Baker, in Gard. Chron.* (1874) i. 623.—Afr. austr.
cordifolia, Schult. f. Syst. vii. 658 = Haworthia cordifolia.
crassifolia, Schult. f. l. c. 664 = Gasteria nigricans.
crassiflora, *Baker, in Journ. Linn. Soc.* xviii. (1880) 162.—Afr. trop.
Croucheri, Hook. f. Bot. Mag. t. 5813 = Gasteria Croucheri.

ALOË :—
cryptopoda, *Baker, in Journ. Bot.* xxii. (1884) 52.—Afr. trop.
curta, *Spreng. Syst.* ii. 69.—Afr. austr.
cuspidata, Schult. f. Syst. vii. 639 = Haworthia cuspidata.
cylindracea, Lam. Encyc. i. 69 = Apicra spiralis, et Haworthia rigida.
cymbæfolia, Schrad. Neues Journ. ii. (1807) 17 = Haworthia cymbæfolia.
cymbæformis, Haw. in Trans. Linn. Soc. vii. (1804) 8 = Haworthia cymbæformis.
decipiens, Schult. f. Syst. vii. 674 = Gasteria decipiens.
deltoidea, Hook. f. Bot. Mag. t. 6071 = Apicra deltoidea.
deltoideodonta, *Baker, in Journ. Linn. Soc.* xx. (1883) 271.—Madag.
denticulata, Schult. f. Syst. vii. 639 = Haworthia denticulata.
depauperata, Stend. Nom. ed. I. 80 = pendens.
depressa, *Haw. in Trans. Linn. Soc.* vii. (1804) 16.—Afr. austr.
depressa, Salm-Dyck, ex Steud. Nom. ed. II. i. 57 = distans.
dichotoma, *Linn. f. Suppl.* 206.—Afr. austr.
dictyodes, Schult. f. Syst. vii. 663 = Gasteria retata.
distans, *Haw. Syn. Pl. Succ.* 78.—Afr. austr.
disticha, *Linn. Sp. Pl.* 321.—Afr. austr.
disticha, Mill. Gard. Dict. ed. VIII. n. 5 = saponaria.
distichia, Thunb. Diss. Aloë, n. 9 = Gasteria verrucosa.
dorsalis, Haw. in Trans. Linn. Soc. vii. (1804) 35 = linearis.
drepanophylla, *Baker, in Gard. Chron.* iii. (1875) 814.—Afr. austr.
echinata, Willd. Enum. Hort. Berol. 385 = humilis.
Ecklonis, *Salm-Dyck, Monog.* ii. § 21. f. 9.—Afr. austr.
elegans, *Tod. Inc. Sem. Panorm.* (1880) 37.—Hab.?
elongata, Murr. in Comm. Gott. ix. (1789) 191 = vera.
elongata, Salm-Dyck, Monog. § 29. f. 15 = Gasteria trigona.
emarginata, Schult. f. Syst. vii. 681 = Gasteria acinaciformis.
ensifolia, *Schult. f. l. c.* 649.—Afr. austr.
erecta, Salm-Dyck, Aloë, § 6. f. 7 = Haworthia margaritifera.
excavata, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 276 = Gasteria excavata.
expansa, Haw. in Trans. Linn. Soc. vii. (1804) 8 = Haworthia rigida.
falcata, *Baker, in Journ. Linn. Soc.* xviii. (1880) 161.—Afr. austr.
fasciata, Salm-Dyck, in Schult. f. Syst. vii. 1713 = Haworthia fasciata.
ferox, *Mill. Gard. Dict.* ed. VIII. n. 22.—Afr. austr.
flabelliformis, Salisb. Prod. 246 = plicatilis.
flava, Pers. Syn. i. 378 = vera.
floribunda, Haw. in Trans. Linn. Soc. vii. (1804) 22 = mitriformis.
foetida, Crantz, Inst. i. 466 = Agave americana.
foliolosa, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Apicra folliolosa.
formosa, *Schult. f. Syst.* vii. 662.—Afr. austr.
fragrantissima, Jacq. Enum. Stirp. Vind. App. 399 = Dracæna fragrans.
frutescens, Salm-Dyck, Cat. 30 = arborescens.
fruticosa, Lam. Encyc. i. 67 = arborescens.
gasteriodes, *Baker, in Journ. Linn. Soc.* xvii. (1880) 166.—Afr. austr.
glabra, Salm-Dyck, Obs. 48 = Gasteria glabra.
glabrata, Salm-Dyck, Hort. 225 = Haworthia glabrata.
Gilson, *Mill. Gard. Dict.* ed. VIII. n. 16.—Afr. austr.
gracilis, *Haw. in Phil. Mag.* (Oct. 1825) 280.—Afr. austr.
graeuis, *Schult. f. Syst.* vii. 650.—Afr. austr.
granata, Schult. f. Syst. vii. Aloë, § 6. f. 6 = Haworthia margaritifera.
grandidentata, *Salm-Dyck, Obs.* (1832) 329.—Afr. austr.
Greenii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 165.—Afr. austr.
guineensis, Jacq. Enum. Stirp. Vind. App. 306 = Sansevieria guineensis.
guttata, Salm-Dyck, Hort. 382 = Gasteria nigricans.
Hamburiana, Nand. in Rev. Hortic. (1878) 166 = striata.
Haynaldi, *Tod. Hort. Panorm.* i. 75. t. 24.—Hab.?
hebes, Schult. f. Syst. vii. 657 = Haworthia cymbiformis.

ALOË :—
herbacea, DC. Hort. Monsp. 76 = Haworthia reticulata.
herbacea, Mill. Gard. Dict. ed. VIII. n. 18 = Haworthia atrovirens.
heterocantha, *Baker, in Journ. Linn. Soc.* xviii. (1880) 161.—Hab.?
hexapetala, *Salm-Dyck, Cat.* 38.—Hab.?
horrida, Haw. in Trans. Linn. Soc. vii. (1804) 27 = ferox.
humilis, *Mill. Abr. Gard. Dict.* ed. VI. n. 10 (1771).—Afr. austr.
hyacinthoides, Linn. Sp. Pl. 321 = Sansevieria zeylanica.
hybrida, Salm-Dyck, Cat. 7 = Haworthia hybrida.
imbricata, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Apicra spiralis.
incurva, Haw. Syn. Pl. Succ. 85 = humilis.
indica, Royle, Illust. Pl. Himal. 390 = vera.
indurata, Schult. f. Syst. vii. 654.—Afr. austr.
inermis, *Forsk. Fl. Aegypt. Arab.* 74.—Arabia.
inermis, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 161 = heteracantha.
inignis ×, *N. E. Br. in Gard. Chron.* xxiv. (1885) 40. f. 8.—Hab.?
intermedia, Haw. in Trans. Linn. Soc. vii. (1804) 12 = Gasteria verrucosa.
Krausii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 169.—Afr. austr.
laetepunctata, Schult. f. Syst. vii. 676 = Gasteria laetepunctata.
lasteviroua, Link, Enum. Hort. Berol. i. 395 = Haworthia lasteviroua.
lasvigata, Schult. Syst. vii. 686 = Haworthia albicans.
laevis, *Salm-Dyck, Cat.* 16.—Afr. austr.
latifolia, *Haw. Syn. Pl. Succ.* 82.—Afr. austr.
leptocaulon, Boj. Hort. Maurit. 245 = oligophylla?
leptophylla, N. E. Br. ex Baker, in Journ. Linn. Soc. xviii. (1881) 165 = latifolia.
limpida, *Schult. f. Syst.* vii. 688.—Afr. austr.
linearis, *Haw. in Trans. Linn. Soc.* vii. (1804) 18.—Afr. austr.
Lingua, Ker-Gawl. in Bot. Mag. t. 1322 = Gasteria verrucosa.
Lingua, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 277 = sp. pluria.
Lingua, Thunb. Diss. Aloë, 8 = Gasteria disticha.
linguiformis, DC. Pl. Grass. t. 68 = Gasteria sulcata.
linguiformis, Lam. f. Suppl. 206 = plicatilis.
linguiformis, Mill. Gard. Dict. ed. VIII. n. 13 = Gasteria disticha.
linguiformis, *Medic. Beobacht.* (1782) 64.—Hab.?
linita, *Schult. f. Syst.* vii. 693.—Afr. austr.
littorlis, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 293.—Afr. trop.
littoralis, Koen. ex Baker, in Journ. Linn. Soc. xviii. (1880) 176 = vera.
lomatophylloides, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 22.—Ins. Mascar.
longiaristata, Schult. f. Syst. vii. 684 = aristata.
longifolia, Haw. Syn. Pl. Succ. 88 = Gasteria disticha.
longifolia, Lam. Encyc. i. 90 = Tritomanthe Uvaria.
longistyla, *Baker, in Journ. Linn. Soc.* xviii. (1880) 158.—Afr. austr.
Lynchii ×, *Baker, in Gard. Chron.* (1881) i. 266.
maculata, Nichols. Dict. Gard. i. 62 = humilis.
Macowani, *Baker, in Journ. Linn. Soc.* xviii. (1881) 170.—Afr. austr.
macra, Haw. Suppl. Pl. Succ. 65, 102 = Lomatophyllum macrum.
macracantha, *Baker, in Journ. Linn. Soc.* xviii. (1880) 167.—Afr. austr.
macrocarpa, *Tod. Hort. Bot. Panorm.* 36. t. 9.—Abyss.
macroclada, *Baker, l. c.* xx. (1883) 273.—Madag.
maculata, Ker-Gawl. in Bot. Mag. t. 765 = Gasteria pulchra.
maculata, Desf. Tabl. Hort. Par. ed. I. 85 = obscura.
maculata, Forsk. Fl. Aegypt. Arab. 78 = abyssinica.
maculata, Thunb. Diss. Aloë, 6 = Gasteria maculata.
maculosa, Lam. Encyc. i. 67 = obscura, saponaria.
margaritifera, Burm. f. Fl. Cap. Prod. 10 = Haworthia margaritifera.
marginata, DC. Pl. Grass. t. 81 = Lomatophyllum borbonicum.
marginata, Lam. Encyc. i. 90 = Haworthia albicans.
marginata, Willd. Enum. Hort. Berol. 368 = Lomatophyllum borbonicum.
marmorata, Hitchen, ex Steud. Nom. ed. II. i. 58 = formosa.

ALOË :—

micracantha, *Hew. Suppl. Pl. Succ.* 105.—Afr. austr.
microstigma, *Salm-Dyck, Monog.* ii. § 36. f. 4.—Afr. austr.
minor, Schult. f. Syst. vii. 648 = Haworthia margaritifera.
mirabilis, Haw. in Trans. Linn. Soc. vii. (1804) 9 = Haworthia mirabilis.
mitraeformis, DC. Pl. Grass. 99; Haw. Syn. Succ. 77 = diœna, nobilis.
mitraeformis, Willd. Enum. Hort. Berol. 385 = mitriformis.
mitriformis, *Mill. Gard. Dict.* ed. VIII. n. 1.—Afr. austr.
mollis, Schult. Syst. vii. 665 = Gasteria mollis.
mucronata, *Schult. f. l. c.* 686.—Afr. austr.
multifaria, *Schult. f. l. c.* 686.—Afr. austr.
muricata, Haw. in Trans. Linn. Soc. vii. (1804) 25 = ferox.
muricata, Schult. Obs. 70 = glauca.
mutica, *Schult. f. Syst.* vii. 686.— Afr. austr.
myriacantha, *Schult. f. l. c.* 704.—Afr. austr.
neglecta, *Tenore, in Atti Accad. Sc. Nap.* v. i. (1837) p. 367.—Hab.?
nigra, Schult. f. Syst. vii. 657 = Haworthia nigra.
nigricans, Haw. in Trans. Linn. Soc. vii. (1804) 18 = Gasteria nigricans.
nitens, *Baker, in Journ. Linn. Soc.* xviii. (1880) 170. —Afr. austr.
nitens, Schult. f. Syst. vii. 680 = Gasteria acinacifolia.
nitida, Salm-Dyck, Cat. 18 = Gasteria nitida.
nobilis, *Haw. Syn. Pl. Succ.* 78.—Afr. austr.
obliqua, DC. Pl. Grass. t. 91 = Gasteria pulchra.
obliqua, Haw. in Trans. Linn. Soc. vii. (1804) 18 = Gasteria maculata.
obliqua, Jacq. Hort. Schoenb. iv. 9 = Gasteria nigricans.
obscura, *Mill. Gard. Dict.* ed. VIII. n. 6.—Afr. austr.
obscura, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 276 = Gasteria excavata.
obscurevirens, *Martinati, ex Vis. Orto Bot. Padov.* 182.—Hab.?
obtusa, Schult. f. Syst. vii. 679 = Gasteria obtusa.
obtusifolia, Salm-Dyck, Obs. 62 = Gasteria obtusifolia.
officinalis, Forsk. Fl. Aegypt. Arab. 78 = vera.
oligophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 372.—Madag.
pallescens, Haw. Rev. Pl. Succ. 41 = serrulata.
pallida, Schult. f. Syst. vii. 641 = Haworthia pallida.
palmiformis, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 368.—Angola.
paniculata, Jacq. Fragm. 48. t. 66 = striata.
papillosa, Salm-Dyck, Cat. 7 = Haworthia papillosa.
parva, Schult. f. Syst. vii. 658 = Haworthia tessellata.
Peacockii, Baker, in Journ. Linn. Soc. xviii. (1880) 175 = abyssinica.
pellucens, Haw. Syn. Pl. Succ. 96 = Haworthia translucens.
pendens, *Forsk. Fl. Aegypt. Arab.* 74.—Arabia.
pentagona, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Apicra pentagona.
pentagona, *Jacq. Fragm.* 72. t. 111.—Afr. austr.
pentagona, Salm-Dyck, Monog. i. § 1. f. 4 = quinquangularis.
perrarum, Tod. Hort. Bot. Panorm. i. 81. t. 21 = abyssinica.
perfoliata, Linn. Sp. Pl. 319, 320 = sp. plurim.
perfoliata, Meyen, Reise, ii. 45 = arborescens.
Perryi, *Baker, in Journ. Linn. Soc.* xviii. (1880) 161. —Ins. Socotra.
persicina, Haw. in Trans. Linn. Soc. vii. (1804) 25 = Crassula caryophyllosa.
picta, Thunb. Diss. Aloë, 6 = obscura.
planifolia, Baker, in Saund. Refug. Bot. t. 162 = Gasteria planifolia.
planifolia, Schult. f. Syst. vii. 688 = Haworthia cymbiformis.
platylepis, *Baker, in Gard. Chron.* (1877) ii. 36.—Afr. austr.
platyphylla, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 364.—Angola.
plicatilis, *Mill. Gard. Dict.* ed. VIII. n. 7.—Afr. austr.
pluridens, *Haw. in Phil. Mag.* (Oct. 1824) 299.—Afr. austr.
pluripunctata, *Schult. f. Syst.* vii. 681.—Afr. austr.
postgenita, *Schult. f. l. c.* 1714.—Afr. austr.

ALOË :—

praemisia, *Baker, in Journ. Linn. Soc.* xviii. (1880) 166. —Afr. austr.
proliféra, Haw. in Trans. Linn. Soc. vii. (1804) 16 = brevifolia.
pseudoafricana, Salm-Dyck, Cat. 31 = africana.
pseudonigrolata, Salm-Dyck, l. c. 16 = Gasteria subcarinata.
pseudoferox, Salm-Dyck, l. c. 31 = ferox.
pseudonigricana, Salm-Dyck, l. c. 17 = Gasteria nigricans.
pseudorigida, Salm-Dyck, l. c. 9 & 41 = Haworthia subrigida.
pseudotortuosa, Salm-Dyck, l. c. 8 & 41 = Haworthia viscosa.
pulchra, Jacq. Hort. Schoenbr. iv. 10. t. 419 = Gasteria pulchra.
pumila, Linn. Sp. Pl. 322 = sp. plurim.
Pumilio, Jacq. Hort. Schoenb. iv. 11 = Haworthia reticulata.
punctata, Haw. in Trans. Linn. Soc. vii. (1804) 26 = variegata.
punctata, *Martinati, in Vis. Orto Bot. Padov.* 182.— Hab.?
purpurascens, *Haw. in Trans. Linn. Soc.* vii. (1804) 20.—Afr. austr.
purpurea, Lam. Encyc. i. 85 = Lomatophyllum borbonicum.
pusilla, *Schult. f. Syst.* vii. 674.—Hab.?
quinquangularis, *Schult. f. l. c.* 658.—Afr. austr.
racemosa, Lam. Encyc. i. 91 = Gasteria verrucosa.
Radula, Jacq. Hort. Schoenb. iv. 11. t. 422 = Haworthia Radula.
Radula, Ker-Gawl. in Bot. Mag. t. 1845 = Haworthia attenuata.
ramifera, *Schult. f. Syst.* vii. 686.—Afr. austr.
ramosa, Haw. in Trans. Linn. Soc. vii. (1804) 26 = purpurascens.
recurva, Haw. l. c. 10 = Haworthia recurva.
reflexa, Van Marum, ex Steud. Nom. ed. II. i. 59 = diœna.
Reinwardtii, Salm-Dyck, Obs. (1821) 87 = Haworthia Reinwardtii.
repens, Schult. f. Syst. vii. 674 = Gasteria repens.
reticulata, Haw. in Trans. Linn. Soc. vii. (1804) 9 = Haworthia reticulata.
retusa, Linn. Sp. Pl. 322 = Haworthia retusa.
retusifolia, Haw. ex Steud. Nom. ed. II. i. 59 = Gasteria obtusifolia.
rhodacantha, DC. Pl. Grass. 44 = glauca.
rhodacrata, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 163 = striata.
rigida, Ker-Gawl. in Bot. Mag. t. 1887 = Haworthia tortuosa.
rigida, DC. Pl. Grass. t. 62 = Haworthia rigida.
rigida, Salisb. Prod. 245 = Kniphofia aloides.
rigida, Jacq. Fragm. t. 108 = Haworthia subrigida.
rubens, DC. Pl. Grass. t. 15 = vera.
rufescincta, Haw. Suppl. Pl. Succ. 45 = Lomatophyllum rufocinctum.
rugosa, Salm-Dyck, in Schult. f. Syst. vii. 1712 = Haworthia rugosa.
Sabila, *Kunw. ex Steud. Nom.* ed. II. i. 59.—Mexic.
Sahundra, *Poj. Hort. Maurit.* 246.—Madag.
Salm-Dyckiana, *Schult. f. in Syst.* vii. 710.—Afr. austr.
saponaria, *Haw. in Trans. Linn. Soc.* vii. (1804) 17.— Afr. austr.
scabrivirens, Salm-Dyck, Hort. 382; Schult. f. Syst. vii. 1713 = Gasteria verrucosa.
scabra, Schult. f. Syst. vii. 644 = Haworthia scabra.
Schimperi, *Tod. Hort. Panorm.* (1876) 48.—Abyss.
Schweinfurthii, Regel, Gartenfl. (1879) 97 = Cooperi.
Schweinfurthii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 175.—Afr. trop.
semiglabrata, Schult. f. Syst. vii. 650 = Haworthia semiglabrata.
seminargentifera, Salm-Dyck, Cat. 6 = Haworthia margaritifera.
Serra, *DC. Pl. Grass.* t. 80.—Afr. austr.
Serra, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 282 = depressa.
serrulata, *Haw. in Trans. Linn. Soc.* vii. (1804) 18.— Afr. austr.
setosa, Schult. f. Syst. vii. 641 = Haworthia setata.
siguoides, *Baker, in Journ. Linn. Soc.* xvii. (1880) 177.—Afr. austr.
sinuata, Thunb. Diss. Aloë, 6 = succotrina.
sinuata, Willd. Sp. Pl. ii. 187 = purpurascens.

ALOË :—

succotrina, Schult. f. Syst. vii. 701 = seq.
succotrina, DC. Pl. Grass. 61. t. 86 = succotrina.
sordida, Schult. f. Syst. vii. 644 = Haworthia sordida.
speciosa, *Baker, in Journ. Linn. Soc.* xviii. (1880) 178. —Afr. austr.
spicata, *Linn. f. Suppl.* 205; *Haw. Syn. Succ.* 76.— Afr. austr.
spinulosa, Salm-Dyck, Obs. (1892) 4 = mitriformis.
spiralis, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Apicra pentagona.
spiralis, Linn. Sp. Pl. 323 = Apicra spiralis.
spirella, Salm-Dyck, Cat. 10 = Apicra pentagona.
squarrosa, *Baker, ex Bull. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 97.—Ins. Socotr.
stenophylla, Schult. f. Syst. vii. 641 = Haworthia angustifolia.
striata, *Haw. in Trans. Linn. Soc.* vii. (1804) 18.— Afr. austr.
striatula, *Haw. in Phil. Mag.* (Oct. 1825) 281.—Afr. austr.
strigata, *Schult. f. Syst.* vii. 676.—Afr. austr.
subalbicans, Salm-Dyck, Monog. i. § 6. f. 1 = Haworthia margaritifera.
subattenuata, Salm-Dyck, in Schult. f. Syst. vii. 1712 = Haworthia subattenuata.
subcarinata, Salm-Dyck, Obs. 51 = Gasteria subcarinata.
subrecta, Haw. in Trans. Linn. Soc. vii. (1804) 16 = humilis.
subfasciata, Salm-Dyck, in Schult. f. Syst. vii. 1712 = Haworthia subfasciata.
subferox, *Spreng. Syst.* ii. 73.—Afr. austr.
subnigricans, Spreng. i. c. 71 = Gasteria nigricans.
subrigida, Schult. f. Syst. vii. 654 = Haworthia subrigida.
subtortuosa, Schult. f. l. c. = Haworthia viscosa.
subtuberculata, Haw. in Phil. Mag. (Oct. 1825) 280 = humilis.
subulata, Salm-Dyck, in Schult. f. Syst. vii. 1712 = Haworthia subulata.
subverrucosa, Salm-Dyck, Obs. 67 = Gasteria subverrucosa.
succotrina, *Lam. Encyc.* i. 85.—Afr. austr.
sulcata, Salm-Dyck, Obs. 54 = Gasteria sulcata.
supralaevis, *Haw. in Trans. Linn. Soc.* vii. (1804) 22. —Afr. austr.
tenuifolia, *Lam. Encyc.* i. 87.—Afr. austr.
tenuior, *Haw. in Phil. Mag.* (Oct. 1825) 281.—Afr. austr.
tessellata, Schult. f. Syst. vii. 685 = Haworthia tessellata.
Thraskii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 180.—Afr. austr.
torquata, Schult. f. Syst. vii. 654 = Haworthia viscosa.
tortuosa, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Haworthia tortuosa.
translucens, [Soland. in] Ait. Hort. Kew. ed. II. ii. 300 = Haworthia translucens.
triangularis, Lam. Encyc. i. 89 = Haworthia viscosa.
trichotoma, Colla, Hort. Ripul. 13 = saponaria.
tricolor, *Baker, in Bot. Mag.* t. 6824.—Afr. austr.
tricolor, Haw. in Trans. Linn. Soc. vii. (1804) 25 = Haworthia venosa.
trigona, Salm-Dyck, Monog. ii. § 39. f. 18 = Gasteria obtusa.
tripetala, Medic. Beobacht. (1788) 55 = plicatilis.
triticina, Medic. Theod. 74 = Gasteria carinata.
tuberculata, Haw. in Trans. Linn. Soc. vii. (1804) 16 = humilis.
tuberculata, *Lag. Gen. et Sp. Nov.* 18.—Hab.?
turgida, Schult. f. Syst. vii. 685 = Haworthia turgida.
umbellata, DC. Pl. Grass. t. 98 = saponaria.
undata, Schult. f. Syst. vii. 677 = Gasteria subcarinata.
Unerid, Linn. Sp. Pl. 323 = Kniphofia aloides.
vacillans, Forsk. Fl. Aegypt. Arab. 74 = vera.
variegata, Linn. l. c. 74 = arabica.
variegata, *Linn. Sp. Pl.* 321.—Afr. austr.
venosa, Lam. Encyc. i. 89 = Haworthia venosa.
venusta, *Schult. f. Syst.* vii. 681.—Hab.?
vera, *Linn. Sp. Pl.* 320.—Reg. medit.
vera, Mill. Gard. Dict. ed. VIII. n. 15 = succotrina.
verrucosa, Mill. l. c. n. 20 = Gasteria verrucosa.
verrucula, Medic. Beobacht. (1788) 68 = pract.
virens, *Haw. in Trans. Linn. Soc.* vii. (1804) 17.—Afr. austr.
virescens, *Schult. f. Syst.* vii. 687.—Hab.?

ALOE:—
virginica, Crantz, Inst. i. 466 = Agave virginica.
viscosa, Linn Sp. Pl. 322 = Haworthia viscosa.
vittata, *Schult. f. Syst. Veg.* vii. 662 —Afr. austr.
vivipara, Crantz, Inst. i. 466 = Agave vivipara.
vulgaris, Lam. Encyc. i. 86 = vera.
xanthacantha, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 292 = mutriformis.
yuccaefolia, A. Gray, in Proc. Am. Acad. vii. (1868) 350 = Hesperoloe yuccaefolia.
zebrina, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 264.—Afr. trop.
Zeyheri, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 178 = Hainesii
Zeyheri, Salm-Dyck, Aloe, § 29. f. 3bis = Gasteria Zeyheri.
zeylanica, Jacq. Hort. Vind 3:10 = Sansevieria zeylanica.

ALORATHEROS, Endl. Gen. 96 (1836) = ALLOIATHEROS, Ell. = **Gymnopogon**, Beauv. † (Gramin.)

ALOEXYLUM, Lour. Fl. Cochinch. 267 (1790) = **Aquilaria**, Lam. (Thymel.).
Agallochum, Lour. l. c. = Aq. Agallochum.
verum, Lour. l. c. = Aq. Agallochum.

ALOIDES, Boerh. ex Linn. Gen. ed. II. 258 (1742) = **Straticles**, Linn. (Hydrocharr.).

ALOIZONIUM, Lindl. Veg. Kingd. 718 (1847) = ALLOIOZONIUM, Kunze = **Cryptostemma**, R. Br. (Compos.).

ALOITIS, Rafin. Fl. Tellur. iii. 21 (1836) = **Gentiana** Linn.
ancepe, Rafin. l. c. 22.—Hab.?
parviflora, Rafin l. c. 21 = G. quinqueflora.
quinqueflora, Rafin. l. c. 22 = G. quinqueflora.

ALOMIA, H. B. & K. Nov. Gen. et Sp. iv. 151, 312. t. 354 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 240. LYCAPSUS, Phil. in Bot. Zeit. (1870) 499. t. 8.
ORSINIA, Bertol. ex DC. Prod. v. 104 (1836).
ageratoides, *H. B. & K. Nov. Gen. et Sp.* iv. 151. t. 354.—Mexic.
alata, *Hemsl. Biol. Centr. Am. Bot.* ii. 79.—Mexic.
angustata, *Benth. ex Baker, in Mart. Fl. Bras.* vi ii. 190.—Bras.
Armani, *Baker, l. c.* 191.—Bras.
cinerea, *Benth. ex Baker, l. c.*—Bras.
fastigiata, *Benth. & Baker, l. c.* 192.—Bras.
foliosa, *Benth. & Hook. f. Gen.* ii. 240.—Bras.
myrtoidea, *Baker, in Mart. Fl. Bras.* vi. ii. 192.—Bras.
Pohlii, *Baker, l. c.* 190.—Bras.
polyphylla, *Baker, l. c.* 191.—Bras.
splanthoides, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 338 = Gymnocoronis splanthoides.

ALONA, Lindl. Bot. Reg. (1844) t. 46. *CONVOLVULACEAE*, Benth. & Hook. f. ii. 879.
RAYERIA, Gaudich. Voy. Bonite, t. 108 (1839–46).
baccata, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
balsamifua, *Gaudich. Voy. Bonite*, t. 107.—Hab.?
carnosa, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
coelestis, *Lindl. l. c.*—Chili.
floccricola, *Phil. Fl. Atac.* 44.—Chili.
dubia, *Phil. in Anales Univ Chil.* (1865) ii. 850.—Chili.
ericifolia, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 500.—Chili.
floribunda, *Phil. in Anales Univ. Chil.* (1878) 525.—Chili.
glandulosa, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
longifolia, *Lindl. l. c.*—Chili.
micrantha, *Phil. Fl. Atac.* 44.—Chili.
microphylla, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 501.—Chili.
mollis, *Phil. Fl. Atac.* 44.—Chili.
obtusa, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
pusilla, *Phil. Fl. Atac.* 45.—Chili.
revoluta, *Lindl. Bot. Reg.* (1844) sub t. 46.—Peruv.
rostrata, Lindl. l. c. = Osteocarpus rostratus.
sphaerophylla, *Phil. Fl. Atac.* 44.—Chili.
tomentosa, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
vernicosa, *Phil. Fl. Atac.* 44.—Chili.
xerophila, *Phil. l. c.*—Chili.

ALONSOA, Ruiz & Pav. Syst. Veg. 150 (1798). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 980.
HEMIMERIS, Pers. Syn. ii. 163 (1807) partim; H. B. & K. Nov. Gen. et Sp. ii. 276 (1817).

ALONSOA:—
HEMITOMUS, L'Hérit. ex Desf. Tabl. ed. I. 66 (1804).
SCHISTANTHE, Kunze, in Linnaea, xvi. (1842) Lit. 109.
acutifolia, *Ruiz & Pav. Syst. Veg.* 153.—Peruv.
caulialata, *Ruiz & Pav. l. c.* 152.—Peruv.
hirsuta, *Steud. Nom.* ed. II. i. 60.—Bras.
incisifolia, *Ruiz & Pav. Syst. Veg.* 154.—Peruv.
intermedia, Lodd. Bot. Cab. t. 1458 = linearis.
linearifolia, *Steud Nom.* ed. II. i. 60.—Peruv.
linearis, *Ruiz & Pav. Syst. Veg.* 154.—Peruv.
linifolia, *Kocsl, in Belg Hortic.* xxiv. (1874) 38.—Peruv.
Mathewsii, *Benth. in DC. Prod.* x 250.—Peruv.
Meliisii, G. Don, Gen. Syst. iv. 513 = caulialata.
myrtifolia, *Kocsi, in Belg. Hortic.* xxiv. (1874) 39.—Peruv.
parviflora, *G. Don, Gen. Syst.* iv. 513.—Venezuela.
procumbens, *Ruiz & Pav., Fl. Peruv.* 154.—Peruv.
quadrifolia, *G. Don, Gen. Syst.* iv. 513.—Mexic.
unilabiata, Ruiz & Pav. ex Steud. Nom. ed. I. 57, 31 = Diascia unilabiata.
urticaefolia, Steud. l. c. ed. II. i. 61 = incisifolia.
Warscewiczii, *Regel, in Gartenfl.* iii. (1854) 211. t 91.—Peruv.

ALONZOA, Brongn. Enum. Genr. 59 (1843) = prasc.

ALOPECIAS, Stev. in Bull. Soc. Nat. Mosc. iv. 266 (1832) = **Astragalus**, Tourn. (Legum.).
alopecuroides, Stev. l. c. = Astr. alopecuroides.
barbatus, Stev. l. c. 266 = Astr. barbatus.
chartaceus, Stev. l. c. xxix. (1856) ii. 148 = Astr. chartaceus.
Clausi, Stev. l. c. 148 = Astr. vulpinus.
dasyanthus, Stev. l. c. iv. (1832) 266 = Astr. dasyanthus.
dasycephalus, Stev. l. c. 266 = Astr. roseus.
emarginatus, Stev. l. c. 266 = Astr. emarginatus.
Kotschyanus, Stev. l. c. xxix. (1856) ii. 148 = Astr. Kotschyanus.
lanatus, Stev. l. c. iv. (1832) 266 = Astr. lanatus.
Libanotis, Stev. l. c. 265 = Astr. lanatus.
macrocephalus, Stev. l. c. 266 = Astr. macrocephalus.
narbonensis, Stev. l. c. 266 = Astr. narbonensis.
orbiculatus, Stev. l. c. xxix. (1856) ii. 148 = Astr. orbiculatus.
ponticus, Stev. l. c. iv. (1832) 266 = Astr. ponticus.
roseus, Stev. l. c. xxix. (1856) ii. 148 = Astr. roseus.
Sieversianus, Stev. l. c. 148 = Astr. Sieversianus.
sphaerocephalus, Stev. l. c. iv. (1832) 266 = Astr. macrocephalus.
vulpinus, Stev. l. c. 266 = Astr. vulpinus.

ALOPECURUS, Linn. Syst. ed. I. (1735); Gen. ed. I. 18 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1140.
COLOBACHNE, Beauv. Agrost. 22. t. 6 (1812).
TOZZETIA, Savi, in Mem. Soc. Ital. Sc. viii. (1798) 477.
aequalis, Sobol. Fl. Petrop. 16 = fulvus.
alpinus, *Desv. Opusc.* 55.—Hab.?
agrestis, *Linn. Sp. Pl.* ed. II. 89.—Europ.; Canc.
agrestis, Siebei, ex Kunth, Enum. Pl. i. 24 = creticus.
alpestris, Wahlenb. ex Steud. Nom. ed. II. i. 61 = pratensis.
alpinus, *Sm. Fl. Brit.* iii. 1086.—Reg. bor. et arct.; Chili.
altissimus, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 13 = arundinaceus.
angustifolius, Sibth. & Sm. Fl. Graec. t. 64 = vaginatus.
antarcticus, Gles. in Brewst. Edinb. Cycl. x. 494 = alpinus.
antarcticus, Lechler, ex Steud. Syn. Pl. Gram. 168 = Lechleri.
antarcticus, Vahl, Symb. Bot. ii. 18 = alpinus.
anthoxantoides, *Boiss. Diagn.* Ser. I. xiii. 42.—Orient.
aquaticus, Dum. Obs. Gram. Belg. 183 = nigricans.
aristatus, Gouan, Hort. Monsp. 87 = Polypogon monspeliensis.
aristulatus, *Michx. Fl. Bor. Am.* i. 48.—Europ.; Am. bor.
arundinaceus, *Poir. Encyc.* viii. 766.—Europ.; Am. bor.; Orient.
arvensis, Gronow. ex Trautv. in Act. Hort. Petrop. ix. (1884) 389 = pratensis.

ALOPECURUS:—
Auckeri, Balansa, in Bull. Soc. Bot. Fr. xxi. (1874) 11 = vaginatus.
Auckeri, *Boiss. Diagn.* Sér. I. xiii. 41.—Persia.
australis, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 413 = geniculatus.
bashkirensis, Turcz. ex Ledeb. Fl. Ross. iv. 462 = brachystachyus.
borealis, Trin. Fund. Agrost 58 = alpinus.
brachyglossus, Peterm. Fl. Lips. (1844) 329 = pratensis. brachystachyus, *Bieb. Fl. Taur. Cauc. Suppl.* 56.—Sibir.
brachystachyus, C. Koch, in Linnaea, xxi. (1848) 388 = vaginatus.
brachystylus, Peterm. Fl. Lips. (1844) 230 = pratensis.
bulbosus, With. Bot. Arr. Brit. Pl. ed. II. i. 59 = geniculatus.
bulbosus, Poir. Voy. Barb. ii. 94 = macrostachyos.
caesius, Schur, Enum. Pl. Transs. 727 in syn. = geniculatus.
caespitosus, *Trin. Sp. Gram.* [xxi. t. 241. *sec. Pritzel, Ic. Ind.*].—Am. bor.
candicans, Salzm. ex Steud. Nom. ed. II. i. 61 = pratensis.
capensis, Eckl. ex Steud. Syn. Pl. Gram. 394 = Koeleria Alopecuros.
capensis, *Thunb. Prod. Fl. Cap.* 19.—Afr. austr.
capillaris, *Raeusch. Nom.* ed. III. 20.—Europ.
capitatus, Lam. Illustr. i. 168 = Phleum Gerardi.
carolinianus, Spreng. Nachtr. i. Bot. Gart. Halle, 10 = Agrostis Cornucopiae.
carolinianus, *Walt. Fl. Carol.* 74.—Am. bor.
cassius, Boiss Diagn. Ser. I. xiii. 41 = vaginatus.
castellanus, *Boiss. & Reut. Diagn. Pl. Hispan.* 26 = brachystachyus.
caudatus, Thunb. in Nov. Act. Ups. iii. (1780) 208 = Brylkinia caudata.
caudicans, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 707 = pratensis.
ciliatus, All. Fl. Pedem. ii. 285 = Koeleria phleoides.
ciliatus, Vitm. Summa Pl. i. 178 = Gerardi.
coerulescens, *Steud. & Hochst. ex Steud. Nom.* ed. II. i. 60, nomen.—As. Min.
colobachnoides, Trin. Fund. Agrost. 58 = brachystachyus.
creticus, *Trin. in Spreng. N. Entdeck.* ii. 45.—As. Min.; Crete.
creticus, Willk. ex Willk. & Lange, Prod. Fl. Hispan. i. 41 = agrestis.
crypsoides, *Griseb. Spicil. Fl. Rumel.* ii. 466.—Macedonia.
dasyanthus, Trautv. in Act. Hort. Petrop. v. (1877) 486 = vaginatus.
diandrus, *Greif. Notul.* iii. 11.—Ind. or.
echinatus, Thunb. Prod. Fl. Cap. 19 = Lasiochloa adscendens.
effusus, Link, ex Kunth, Enum. Pl. i. 328 = Agrostis effusa.
elongatus, *Peterm. in Flora*, xxvii. (1844) 230.—Europ.
elongatus, Poir. Encyc. Suppl. v. 498 = Polypogo elongatus.
exaltatus, *Less. in Linnaea*, ix. (1834) 207.—Sibir. ural.
fallacinus, *Gren. Fl. Mass. Advon.* (1859) 21.—Hab.?
fasciculatus, Poir. Encyc. viii. 780 = Chaetturus fasciculatus.
foliosus, Clark, Trav. ii. 261 = vaginatus.
fulvus, Forsk. ex Steud. Nom. ed. II. i. 61 = Heleochloa alopecuroides.
fulvus, Sm. Engl. Bot. xxi. t. 1467 = geniculatus.
geniculatus, *Linn. Sp. Pl.* ed. I. 60.—Reg. bor. temp.
geniculatus, Lindh. ex Scheele, in Linnaea, xxii. (1849) 840 = Setaria polystachya.
geniculatus, Sibth. ex Steud. Nom. ed. II. i. 61 = Heleochloa alopecuroides.
Gerardi, *Vill. Fl. Delph.* 5; *Hist. Pl. Dauph.* ii. 66. t. 2.—Europ.; Orient.
Giesekeanus, Koern. & Schult. Mant. ii. 254 = alpinus.
glacialis, *C. Koch, in Linnaea*, xxi. (1848) 383.—Orient.
glaucus, *Less, l. c.* ix. (1834) 206.—Sibir. ural.
glomeratus, Poir. Encyc. Suppl. v. 495 = Muehlenbergia glomerata.
gracilis, Trautv. in Act. Hort. Petrop. v. (1877) 486 = glacialis.
gracilis, *Willd. ex Trin. Agrost.* i. 16.—Carolina.
granulatus, *Beauv. Agrost.* 4, nomen.—Hab.?

ALOPECURUS :—
kordeiformis, Linn. Sp. Pl. 60 = Pennisetum compressum.
kordeiformis, Willd. ex Steud. Nom. ed. II. i. 61 = Pennisetum orientale.
hybridus, *Wimm. Fl. Schles.* ed. III. (1857) 31.—Europ.
indicus, Murr. Syst. Veg. ed. XIII. 92 = Pennisetum Linnaei.
intermedius, *Hallier, in Bot. Zeit.* xxi. (1868) *App.* i. 6.—Ins. Heligol.
intermedius, Bock, ex Steud. Nom. ed. II. i. 61 = arundinaceus.
interruptus, Poir. Encyc. Suppl. v. 495 = Polypogon interruptus.
japónicus, *Steud. Syn. Pl. Gram.* 149.—Japon.
Konradii, Opiz, ex Mert. & Koch, in Roehl. Deutschl. Fl. i. 483 = geniculatus.
laguriformis, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 182 = brachystachyus.
laguroides, Balansa, in Bull. Soc. Bot. Fr. xxi. (1874) 11 = vaginatus.
lanatus, *Stèth. & Sm. Prod.* i. 48.—Orient.
lasiochlamys, Boiss. Fl. Orient. v. 490 = textilis.
lasiostachyus, *Link, Hort. Berol.* i. 71.—Hispan.
latifolius, *Cav. Descr.* 37.—Ins. Tobago.
Lechleri, *Steud. Syn. Pl. Gram.* 148.—Valdivia.
litoreus, All. Fl. Pedem. ii. 235 = Koeleria villosa.
longearistatus, *Maxim. Prim. Fl. Amur.* 327.—Manchuria.
longiradiatus, Regel, ex Trautv. in Act. Hort. Petrop. ix. (1884) 354 = longearistatus.
longistylus, *Opiz, in Lotos*, ii. (1852) 228.—Bohem.
macrostachyos, *Poir. Encyc.* viii. 775.—Afr. bor.
magellanicus, Lam. Illustr. i. 168 = antarcticus
malacostachyus, A. Gray, in Perry, Exped. Jap. ii. 328 = japonicus.
maritimus, Poir. Encyc. viii. 780 = Polypogon maritimus.
monspeliensis, Linn. Sp. Pl. 61 = Polypogon monspeliensis.
mucronatus, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 597 = arundinaceus.
myosuroides, Huds. Fl. Angl. 28 = agrestis.
nepalensis, *Trin. ex Steud. Syn. Pl. Gram.* 148.—Nepalia.
nigrescens, Jacq. Ecl. ii. t. 18 = arundinaceus.
nigricans, Hornem. Hort. Hafn. i. 68 = arundinaceus.
nigricans, Koch, Syn. Fl. Germ. ii. 896 = pratensis.
nigricans, Wichura, in Jahrb. Schles. Ges. (1845, 61 – hybridus.
obscurus, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 18 = pratensis.
otiporensis, *Griff. Notul.* iii. 94.—Afghan.
ovatus, Forst. Fl. Austral. 8 = alpinus ?
ovatus, Knapp, Gram. Brit. 15 = alpinus.
Pallasii, Trin. Sp. Gram. t. 2 = vaginatus.
pallidus, *Dum. Obs. Gram. Belg.* 133.—Belg.
paludosus, Beauv. ex Mert. & Koch, in Roehl. Deutschl. Fl. i. 481 = geniculatus.
paludosus, Crantz, Inst. i. 368 = Polypogon monspeliensis.
paniceus, Cav. Descr. 37 = Polypogon maritimus.
paniceus, O. Muell. Fl. Dan. t. 861 = geniculatus.
paniceus, Linn. Sp. Pl. ed. II. 90; Lam. Fl. Fr. iii. 560 = Polypogon monspeliensis.
paniculatus, *Gouldensk. ex Ledeb. Fl. Ross.* iv. 465.—Reg. Caucas.
pedalis, Bosc, ex Beauv. Agrost. 4 = gracilis.
pedatus, Steud. Nom. ed. II. i. 61 = prasc.
phalaroides, C. Koch, in Linnaea, xix. (1847) 5 = Gerardi.
phleiformis, Beauv. Agrost. 4 = villosus.
pontícus, C. Koch, in Linnaea, xxi. (1848) 382 = glacialis.
pratensis, Bourg. ex Lange, in Bull. Soc. Bot. Fr. xi. (1864) 47 = castellanus.
pratensis, *Linn. Sp. Pl.* 60.—Reg. bor. temp.
pratensis, Trin. Sp. Gram. t. 45 = arundinaceus.
prorsus, Mitt. ex Hook. f. Stend. Fl. ed. I. 428 = geniculatus.
purpurascens, Link, in Linnaea, xvii. (1843) 400 = agrestis.
ramosus, *Poir. Encyc.* viii. 776.—Carolina.
repens, Bieb. Fl. Taur. Cauc. iii. 54 = arundinaceus.
ruthenicus, Weinm. Fl. Dan. t. 2881; et Cat. Dorpat. (1819) 10 = arundinaceus.
saccatus, *Vasey, in Contr. Bot. Gov.* vi. (1861) 290.—Oregon.

ALOPECURUS :—
scaber, Opiz, Natural. n. ix. (1825) 166 = pratensis.
sericeus, Gaert. Fruct. i. 2. t. 1. f. 2 = pratensis.
setarioides, *Grem. in Mém. Soc. Emul. Doubs*, Sér. III. ii. (1858) 459.—Gallia.
sibiricus, Hort. Vind. ex Roem. & Schult. Syst. ii. 271 = arundinaceus.
subcristatus, Pursh, Fl. Am. Sept. i. 66 = geniculatus.
tenellus, Poir. Encyc. Suppl. v. 496 = Polypogon tenellus.
textilis, *Boiss. Diagn.* Ser. I. xiii. 40.—Oriens.
tonsus, Dum. Obs. Gram. Belg. 132 = agrestis.
transsilvanicus, Schur, Enum. Fl. Transs. 727 = brachystachyus.
trivialis, Seidl, ex Opiz, Natural. n. ix. (1825) 116 = pratensis.
turiconsis, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 111.—Europ.
tuscheticus, Trautv. in Act. Hort. Petrop. ix. (1884) 354 = gracilis.
typhoides, Burm. f. Fl. Ind. 27 = Penicillaria spicata.
uralensis, *Uspenski, Ekaterinenburg.* 24.—Sibir. mer.
utriculatus, [*Soland. MS*] *Russ. Aleppo*, ed. II. ii. 243. —Europ.; Oriens.
vaginatus, *Pall. in Nov. Act. Acad. Petrop.* x. (1792) 304, nomen.—Tauria; Oriens.
variegatus, *Steud. Syn. Pl. Gram.* 148.—Chili.
varius, Auct. ex Steud. Nom. ed. I. 32, 771 = Sesleria caerulea.
ventricosus, Huds. Fl. Angl. ed. II. i. 28 = Gastridium australe.
ventricosus, Pers. Syn. i. 80 = arundinaceus.
villosus, Gilib. Exercit. ii. 517 = pratensis.
villosus, *Poir. Encyc.* viii. 777.—Helvet.
Vlassovii, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. ii. (1845) 42 = brachystachyus.

ALOPHIA, Herb. in Bot. Mag. sub t. 3779 (1838) = **Herbertia**, Sweet (Irid.).
Drummondiana, Herb. l. c. = H. Drummondiana.
geminulata, Klatt, in Mart. Fl. Bras iii. i. 517.—Bras.
linearis, Klatt, in Linnaea, xxxi. (1861-62) 558 = H. lineata.
linearis, Warm. in Kjoeb. Vidensk. Meddel. (1872) 138.—Bras.
pulchella, Benth. & Hook. f. Gen. iii. 692 = H. pulchella, lineata.
Sellowiana, Klatt, in Linnaea, xxxi. (1861-62) 557.—Bras.

ALOPHIUM, Cass. in Dict. Sc. Nat. liv. 498 (1828) = **Centaurea**, Linn. (Compos.).
tenuifolium, Cass. l. c. = C. Alophium.

ALOPHOCHLOA, Endl. Gen. 100 (1836) *sphalm.* = LOPHOCHILOA, Reichb. = **Koeleria**, Pers. (Gram.).

ALOPHYLLUS, Linn. Sp. Pl. ed. II. 496 (1762) = ALLOPHYLUS, Linn. (1747) = **Schmiedelia**, Linn. (1764) (Sapind.).

ALOPICARPUS, Neck. Elem. ii. 369 (1790) = **Paris**, Rupp. (Liliac.).

ALORANTHUS, F. S. Voigt, in Rich. Anal. Frucht, 29 (1811) *sphalm.* = **Chloranthus**, Sw.

ALOSEMIS, Rafin. Sylva Tellur. 96 (1838) = **Malastoma**, Linn.
grandiflora, Rafin. l. c.—Hab. ?
villosa, Rafin. l. c.—Hab. ?
seylanica, Rafin. l. c.—Zeylan.

ALOYSIA, Ort. & Palav. ex L'Hérit. Stirp. i. 21 (1784) = **Lippia**, Linn. (Verben.).
attenuata, Walp. Rep. iv. 42.—Bras.
chamaedryfolia, Cham. in Linnaea, vii. (1832) 234 = L. chamaedryfolia
chamaedryoides, Steud. Nom. ed. II. i. 62 = L. chamaedryfolia.
citriodora, Ortog. ex Pers. Syn. ii. 139 = L. citriodora.
floribunda, Mart. & Gal. in Bull. Acad. Brux. xi. (1844) II. 290 = L. lycioides.
lycioides, Cham. in Linnaea, vii. (1832) 237 = L. lycioides.
polygalaefolia, Cham. l. c. 236 = L. polygalaefolia.
scordonioides, Cham. l. c. 234 = L. scordonioides.
urticoides, Cham. l. c. 238 = L. urticoides.
virgata, Juss. in Ann. Mus. Par. vii. (1806) 78 = L. urticoides.

ALPHANDIA, Baill. Adansonia, xi. (1873) 85. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 300.
furfuracea, *Baill. l. c.* 86.—N. Caled.
resinosa, *Baill. l. c.* 86.—N. Caled.

ALPHITONIA, Reissek, ex Endl. Gen. 1098 (1840). *RHAMNEAE*, Benth. & Hook. f. i. 381.
Emmenosperma, F. Muell. Fragm. iii. 63 = Emmenosperma alphitonoides.
erubescens, *Baill. Adansonia*, xi. (1874-76) 271.—N. Caled.
excelsa, *Reissek, ex Endl. Gen.* 1098.—Austral.
franguloides, *A. Gray, Bot. U. St. Expl. Exped.* i. 280. —Ins. Viti.
moluccana, *Tenjsm. & Binn. Cat. Hort. Bogor.* 221.—Molucca.
xerocarpus, *Baill. Adansonia*, xi. (1873-76) 270.—N. Caled.
zizyphoides, *A. Gray, Bot. U. St. Expl. Exped.* i. 278.—Hab. ?

ALPHONSEA, Hook. f. & Thoms. Fl. Ind. i. 152 (1855). *ANONACEAE*, Benth. & Hook. f. i. 29.
ceramicola, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 29.—Malaya.
elliptica, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 90.—Malacca.
javanica, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 340; *in Flora*, liii. (1870) 242.—Java.
lutea, *Hook. f. & Thoms. Fl. Ind.* i. 158.—Ind. or.
luteo, Teijsm. & Binn. ex Scheff. in Flora, liii. (1870) 242 = javanica.
madraspatana, *Bedd. Fl. Sylv.* i. t. 76.—Ind. or.
Maingayi, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 90.—Malacca.
sclerocarpa, *Thw. Enum. Pl. Zeyl.* 11.—Zeylan.
ventricosa, *Hook f. & Thoms. Fl. Ind.* i. 152.—Ind. or.; Malaya.
zeylanica, *Hook. f. & Thoms. l. c.* 153.—Ind. or.

ALPINIA, Linn. Gen. ed. I. 332 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 648.
ALBINA, Gieseke, Prael. Linn. 199, 207 (1792).
ALUGHAS, Linn. Fl. Zeyl. 207 (1747).
BANGLIUM, Buch.-Ham. ex Wall. Cat. sub n. 6579 (1832).
BOJERIA, Rafin. Fl. Telluv. fo. 50 (1836).
BUEKIA, Gieseke, Prael. Linn. 199, 204 (1792).
CATIMBIUM, Juss. Gen. 62 (1789).
CENOLOPHON, Blume, Enum. Pl. Jav. 60 (1830).
DONANTHES, Rafin. Fl. Telluv. iv. 51 (1836).
GALANGA, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1 Art. iv 2; ed. II. 65 (1827).
HELLENIA, Willd. Sp. Pl. i. 4 (1797).
HERITIERA, Retz Obs. vi. 17. t. 1 (1791).
KOLOWRATIA, Presl, Rel. Haenk. i. 118 t. 20 (1827).
LANGUAS, Koen. in Retz. Obs. iii. 64 (1783).
MARTENSIA, Gieseke, Prael. Linn. 199, 207 (1792).
MONOCYSTIS, Lindl. Introd. Nat. Syst. ed. II. 445 (1836).
ZERUMBET, Wendl. Sert. Hannov. fasc. iv. 8. t. 19 (1798).
acrostachya, Steud. Nom. ed. II. i. 62 (= Amomi sp.). —Java.
alba, D. Dietr. Syn. Pl. i. 12 = Elettaria alba.
alba, *Rosc. in Trans. Linn. Soc.* viii. (1807) 346.—Ind. or.
alata, *A. Dietr. Sp. Pl.* i. 45.—China.
Allaghas, *Rosc. in Trans. Linn. Soc.* viii. (1807) 346. —Ind. or.
Anstillarum, *Roem. & Schult. Syst. Veg* i. 20.—Ind. occ.
aquatica, *Rosc. in Trans. Linn. Soc.* viii. (1807) 346.—Ind. or.
arctiflora, *F. Muell. Fragm.* viii. 25.—Austral.
aromatica, Aubl. Pl. Gui. i. 3 = Renealmia domingensis.
auriculata, *Rosc. Scit. Pl.* t. 74.—Ins. Molucc.
Blumei, D. Dietr. Syn. Pl. i. 13 = Javanica.
Boia, *Seem. Fl. Vit.* 290.—Ins. Fiji.
bracteata, Rosc. in Trans. Linn. Soc. xi. (1815) 281 = alata.
bracteata, Roxb. Fl. Ind. i. 61 = Roxburghii.
brevilabris, *Presl, Rel. Haenk.* i. 110. t. 17.—Ins. Luzon.
brevicapsa, Poepp. & Endl. Nov. Gen. et Sp. ii. t. 135 = Renealmia brevicapsa.
caerulea, *Benth. Fl. Austral.* vi. 265.—Austral.
calcarata, *Rosc. in Trans. Linn. Soc.* viii. (1807) 347. —Ind. or.; China.

ALPINIA :—
capitellata, *Yack, in Malay. Misc.* ii. (1822) n. vii. 4. —Malaya.
Cardamomum, Roxb. in As. Res. xi. (1810) 356 = Elettaria Cardamomum.
Cardamon-medium, Roxb. Fl. Ind. i. 72 = Amomum costatum.
caribæa, Gaert. Fruct. t. 86. t. 12 = Renealmia racemosa.
carnea, *Griff. Notul.* iii. 420.—Ind. or.
cernua, *Sims, Bot. Mag.* t. 1900.—Ind. or.
chinensis, *Roxc. in Trans. Linn. Soc.* viii. (1807) 346.—China.
coccinea, D. Dietr. Syn. Pl. i. 12 = Amomum coccineum.
comosa, Jacq. Coll. iv. 112 = Costus comosus.
conchigera, *Griff. Notul.* iii. 424.—Ind. or.
costata, Roxb. Corom. Pl. iii. 48. t. 252 = Amomum costatum.
cristata, *Griff. Notul.* iii. 421.—Ind. or.
diffusa, *Roxc. in Trans. Linn. Soc.* xi. (1815) 361.—China.
elatior, *Yack, in Malay. Misc.* ii. (1822) n. vii. 2.—Malaya.
exalta, Ledd. ex Stend. Nom. ed. II. i. 62 = calcarata.
exaltata, Roem. & Schult. Syst. i. 21. add. 568 = Renealmia exaltata.
fasciculata, Steud. Nom. ed. II. i. 62 = Amomum fasciculatum.
ferruginea, Steud. Nom. ed. II. i. 62 = mollis.
fœtens, D. Dietr. Syn. Pl. i. 12 = Amomum fœtens.
Galanga, *Willd. Sp. Pl.* i. 12.—As. trop.
gigantea, *Blume, Enum. Pl. Jav.* 59.—Java.
gigantea, F. Vill. in Blanco, Fl. Filip. ed. III. Nov. App. 98 = gracilis.
globosa, *Horan. Prod. Scitam.* 84.—China.
gracilis, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 316.—Ins. Philipp.
Hasskarl, *Pugil, Symb.* 66. t. 43.—Chili.
Hamiltoniana, *Wall. Cat.* n. 6579.—Ind. or.
hemisphærica, D. Dietr. Syn. Pl. i. 13 = Elettaria hemisphærica.
hirsuta, *Horan. Prod. Scitam.* 84.—Cochinch.
humilis, Vell. Fl. Flum. 1; i. t. 4 (= Renealmiæ sp.). —Bras.
humilis, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 237.—Java.
involucrata, *Griff. Notul.* iii. 422.—Ind. or.
jamaicensis, Gaert. Fruct. i. 86. t. 12 = Renealmia jamaicensis.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 140.—Japon.
javanica, *Blume, Enum. Pl. Jav.* 59.—Java.
javanica, D. Dietr. Syn. Pl. i. 13 = *acrostachya*.
latifolia, Willd. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) 1. 180 = Calathea latifolia.
linguiformis, Roxb. Fl. Ind. i. 73 = Elettaria linguiformis.
macrantha, Poepp. & Endl. Nov. Gen. et Sp. t. 184 = Renealmia macrantha.
macrantha, *Schegf. in Ann. Yard. Buitens.* i. (1876) 56.—Malaya.
maculata, *Roxc. in Trans. Linn. Soc.* viii. (1807) 347.—Ind. or.
magnifica, Roxc. Scit. t. 75 = Amomum magnificum.
malaccensis, Presl, Rel. Haenk. i. 110 = Hasskarl.
malaccensis, *Roxc. in Trans. Linn. Soc.* viii. (1807) 345.—Ind. or. ; java.
media, Sprengl. Syst. i. 14 = Amomum costatum.
mexana, Horna. Prod. Scitam. 27 = Zingiber macrostachyum.
miner, D. Dietr. Syn. Pl. i. 12 = Elettaria minor.
minuta, D. Dietr. l. c. = Elettaria minuta.
missionis, *Wall. Cat.* n. 6589.—Ind. or.
mollis, *Presl, Rel. Haenk.* 110.—Ins. Luzon.
multicostata, Roem. & Schult. Syst. i. 21 = Renealmia domingensis.
mutica, *Roxb. in As. Res.* xi. (1810) 354.—Ind. or.
Nicesana, *Y. Grab. Cat. Bomb. Pl.* 307.—Ind. or.
Nimmonii, J. Grah. l. c. 206 = Zingiber Nimmonia.
notans, *Roxc. in Sm. Exot. Bot.* ii. 93.—Ind. or.
occidentalis, Sw. Prod. Veg. Ind. Occ. 11 = Renealmia jamaicensis.
officinarum, *Hance, in Journ. Linn. Soc.* xiii. (1873) 6.—China.
oxyphylla, *Miq. in Journ. Bot. Néerl.* i. (1861) 93.—China.
Paco-Secora, Jacq. Fragm. 54. n. 17. t. 76 = Renealmia Paco-Secora.
pallida, D. Dietr. Syn. Pl. i. 13 = Elettaria pallida.

ALPINIA :—
paludosa, D. Dietr. Syn. Pl. i. 13 = Elettaria paludosa.
papuana, *Schegf. in Ann. Yard. Buitens.* i. (1876) 56. —N. Guin.
parviflora, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 316.—Ins. Philipp.
penicillata, *Roxc. in Trans. Linn. Soc.* xi. (1814) 360.—China.
phœnicea, *Kamphœverner, ex Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 154, nomen.—Ins. Nicobar.
Pinings, *D. Dietr. Syn. Pl.* i. 12.—Java.
porrecta, *Wall. Cat.* n. 6576.—Ind. or.
Prealuna, Steud. Nom. ed. II. i. 62 = Renealmia racemosa.
pumila, *Hook f. Bot. Mag.* t. 6892.—China.
punicea, Roxb. Fl. Ind. i. 71 = Elettaria punicea.
pyramidata, *Blume, Enum. Pl. Jav.* 58.—Java.
racemigera, *F. Muell. Fragm.* viii. 27.—Austral.
racemosa, Linn. Sp. Pl. ed. I. 2 = Renealmia racemosa.
racemosa, Sw. Obs. 5 = A. Antillarum.
Rafflesiana, *Wall. Cat.* n. 6575.—Ind. or.
Renealmia, Sm. in Rees, Cyclop. Suppl. xxxix. n. 14 = Renealmia exaltata.
Rheedii, *Wight, Ic.* t. 1. 2026.—Malabar.
Roscoeana, Steud. Nom. ed. II. i. 63 = alata.
Roxburghii, *Sweet, Hort. Brit.* ed. I. 390.—Ind. or.
rubra, *D. Dietr. Syn. Pl.* i. 13.—Java.
Ruitziana, Steud. Nom. ed. II. i. 63 = Renealmia Ruiziana.
serrulata, *Willd. ex Steud. l. c.*—Bras.
sessilis, Koen. in Rets. Obs. fasc. iii. 64 = Kaempferia Galanga.
Sinsil, *Roxc. Observ. Boccodifol.* 5.—Hab. ?
solaris, D. Dietr. Syn. Pl. i. 13 = Elettaria solaris.
speciosa, D. Dietr. l. c. 13 = Elettaria speciosa.
spicata, Jacq. Enum. Pl. Carib. 11 = Costus spicatus.
spicata, Presl, Rel. Haenk. i. 109 = Renealmia racemosa.
spicata, *Roxb. in As. Res.* xi. (1810) 356.—Ind. or.
spicata, Jacq. Hort. Schoenb. i. 1 = Costus Pisonis.
stachyodes, *Hance, in Journ. Linn. Soc.* xiii. (1873) 126.—China.
striata, *Hort. Angl. ex Link, Enum.* i. 5.—Ind. or.
strobilifera, Poepp. & Endl. Nov. Gen. et Sp. t. 186 = Renealmia strobilifera.
thyrsoidea, Steud. Nom. ed. II. i. 63 = Renealmia thyrsoidea.
tomentosa, *D. Dietr. Syn. Pl.* i. 12.—Java.
tubulata, *[Roxc. in] Bot. Reg.* t. 777.—Ind. occ.
uviformis, *Horan. Prod. Scitam.* 85.—Malaya.
viridiflora, *Griff. Notul.* iii. 423.—Ind. or.
vitiensis, *Seem. Fl. Vit.* 290.—Ins. Fiji.
Walang, *D. Dietr. Syn. Pl.* i. 13.—Java.

ALSCHINGERA, Vis. Fl. Dalm. iii. 69 (1849) = Physospermum, Cusson (Umbell.).
actæfolia, Vis. sec. Tomm. fide Nym. Consp. 315 = P. verticillatum.
verticillata, Vis. Fl. Dalm. iii. 69 = P. verticillatum.

ALSEIS, Schott, in Spreng. Syst. iv. Cur. Post. 404 (1827). *RUBIACEÆ*, Benth. & Hook. f. ii. 88. Blackiana, *Hemsl. Diag. Fl. Nov.* 30.—Panama.
floribunda, *Schott, in Spreng. Syst. iv. Cur. Post.* 404. —Bras.

ALSENOSMIA, Endl. Ench. 284 (1841) sphalm. = Alsenuosmia, A. Cunn. (Caprifol.)

ALSEODAPHNE, Nees, in Wall. Pl. As. Rar. ii. 71 (1831). *LAURINEÆ*, [Benth. & Hook. f. iii. 187.] EUPHORBE, Blume, ex Meisn. in DC. Prod. xv. i. 56, 59 in syn. (1864).
NOTHAPHOEBE, Blume, Mus. Bot. Lugd. Bat. i. 328 (1851).
amara, Kostel. Allg. med.-pharm. Fl. ii. 484 = Nectandra cymbarum.
bancana, *Miq. Fl. Ind. Bat.* i. 915.—Banca.
breviflora, *Benth. Fl. Hongk.* 392.—China.
chinensis, *Champ. ex Meisn. in DC. Prod.* xv. i. 58.—Hongkong.
costalis, *Nees, in Wall. Pl. As. Rar.* ii. 72.—Ind. or.
Cujumary, Kostel. Allg. med.-pharm. Fl. ii. 484 = Acotea Cujumary.
cymbarum, Kostel. l. c. = Nectandra cymbarum.
excelsa, *Blume, Mus. Bot. Lugd. Bat.* i. 328.—Java.
grandis, Kurz, For. Fl. Burma, ii. 293 = Dehaasia cuneata.

ALSEODAPHNE :—
grandis, Nees, in Wall. Pl. As. Rar. ii. 72 = Phœbe opaca.
lucida, *Nees, l. c.*—Singapore.
opifera, Kostel. Allg. med.-pharm. Fl. ii. 484 = Ocotea opifera.
polyneura, *Miq. Fl. Ind. Bat.* i. 916.—Penang.
semicarpifolia, *Nees, in Wall. Pl. As. Rar.* ii. 72.—Ind. or.

ALSEUOSMIA, A. Cunn. in Ann. Nat. Hist. ii. (1839) 209. *CAPRIFOLIACEÆ*, Benth. & Hook. f. ii. 6. FAGOIDES, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. ii. (1839) 210.
atriplicifolia, A. Cunn. l. c. 210 = Banksii.
Banksii, *A. Cunn. l. c.* 209.—N. Zel.
bupleurifolia, *A. Cunn. ex Heynh. Nom.* ii. 27.—N. Zel.
Hookeria, *Colenso, Excurs. North Isl.* 84.—N. Zel.
Ilex, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 210.—N. Zel.
ligustrifolia, A. Cunn. l. c. 209 = linarifolia.
linarifolia, *A. Cunn. l. c.*—N. Zel.
macrophylla, *A. Cunn. l. c.* 210.—N. Zel.
palæriformis, A. Cunn. l. c. = Banksii.
pusilla, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 341.—N. Zel.
quercifolia, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 210.—N. Zel.

ALSINANTHE, Reichb. lc. Fl. Germ. v. (1841) 29. t. 209 = Arenaria, Linn. (Caryophyll.).
biflora, Reichb. l. c. 30. t. 209. f. 4935 = Ar. biflora.
stricta, Reichb. l. c. 29. t. 209. f. 4935 = Ar. stricta.
uliginosa, Reichb. l. c. t. 209. f. 4935 = Ar. stricta.

ALSINANTHEMOS, J. G. Gmel. Fl. Sib. iv. 46 (1769) = Trientalis, Lann. (Primul.).

ALSINANTHUS, Reichb. Handb. 298 (1837) = Arenaria, Linn. (Caryophyll.).
trinervius, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 347 = Ar. trinervis.

ALSINASTRUM, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 15 = Illatina, Linn.
calistricoides, Rupr. Fl. Ingr. 197 = E. triandra.
galitsiolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 15 = E. Alsinastrum.
gymospermum, Rupr. Fl. Ingr. 195 = E. Hydropiper.
orthospermum, Rupr. l. c. 196 = E. Hydropiper *triandrum*, Rupr. l. c. 197 = E. triandra.

ALSINE, Burm. Fl. Zeyl. 18. t. 8 (1737) = Mollugo, Linn. (Ficoid.).

ALSINE, Tourn. ex Linn. Syst. ed. I. (1735) ; Gen. ed. I. 138 (1737) = Arenaria, Linn., Stellaria, Linn. &c. (Caryophyll.).

ALSINE, Scop. Fl. Carn. ed. II. 294 (1772) ; Wahlenb. Fl. Lapp. 127 (1812) = Arenaria, Linn. (Caryophyll.).
aizoides, Boiss. Diagn. Ser. I. i. 47.—Kurdistan.
alpina, Willk. in Bot. Zeit. v. (1847) 248 = Spergularia rubra.
Alsinastrum, Crantz, Inst. ii. 405 = Elatine Alsinastrum.
anatolica, Boiss. Diagn. Ser. I. viii. 97 = Ar. setacea.
apetala, Kit. ex Nym. Consp. 111 = Stellaria media.
arabica, Fisch. Mey. & Ave-Lall. Ind. Sem. Petrop. xi. 49 = Ar. tenuifolia.
arctica, Fenzl, Verbreit. Alsin. in tab. ad p. 16 ; et 36 = Ar. arctica.
Arduinei, Fenzl, l. c. in tab. ad p. 57 = Ar. graminifolia.
arctioides, Mert. & Koch, in Roehl. Deutsch. Fl. iii. 277 = Ar. arctioides.
aristata, [Soland. in] Russ. Aleppo, ed. II. ii. 240.—Syria.
arvensis, Crantz, Inst. ii. 406 = Spergula arvensis.
arvatica, Guss. Fl. Sic. Syn. i. 506 = Ar. macrocarpa.
attica, Boiss. & Sprun. Diagn. Ser. I. v. 34 = Ar. serpa.
Aucheriana, Boiss. Fl. Orient. i. 676 = Ar. Aucheriana.
aurantica, Wahlenb. Fl. Lapp. 129 = Ar. rostrata.
auriculariana, Lam. Fl. Fr. iii. 46 = Stellaria media.
barbata, Stokes, Bot. Mat. Med. ii. 337 = Stellaria media.

ALSINE :—
Barrelieri, Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 57.—Gallia.
Bartolotti, Huet, ex Nym. Consp. 119 = Arenaria procumbens.
baticica, Bluff & Fingh. Comp. ed. II. ii. 11. 99 = Ar. banatica.
Bauhinorum, J. Gay, ex Lamotte, Prod. i. 144 = Ar. linifolia.
bavarica, Crantz, Inst. ii. 407 = Ar. bavarica.
Biebersteinii, Rupr. Fl. Cauc. 215.—Caucas.
biflora, Wahlenb. Fl. Lapp. 128 = Ar. scandinavica.
Billardieri, Boiss. Diagn. Ser. I. viii. 95 = Ar. Meyeri.
bipartita, Gilib. Fl. Lituan. ii. 150 = Stellaria media.
brachycarpa, Boiss. Diagn. Ser. II. vi. 87 = Ar. tenuifolia.
brachypetala, Opiz, Natural. xi. (1826) 406 = Stellaria media.
brevicaulis, Sternb. ex Nym. Consp. 119 = Ar. lanceolata.
brevifiora, Gilib. Fl. Lituan. ii. 150 = Ar tenuifolia.
brevifolia, Chapm. Fl. S. Us. St. 49 = Ar. brevifolia.
brevis, Boiss. Diagn. Ser. I. viii. 96.—Persia; Arabia.
Bufonia, Crantz, Inst. ii. 405 = Bufonia perennis.
caespitosa, Ehrh. ex Nym. Consp. 119 = Ar. verna.
campestris, Boiss. Voy. Espagne, ii. 723 (= *Minuartia campestris*).—Hispan.
campestris, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. montana.
canaliculata, Dulac, Fl. Hautes-Pyr. 246 = Ar Cherleria.
capillipes, Boiss. ex Steud. Nom. ed. II. i. 63 = Ar. capillipes.
cappadocica, Boiss. Diagn. Ser. II. i. 88 = A. setacea.
catarctarum, Janka, in Körlem. Magyar Tudom. Akad. xii. (1876) 165.—Europ.
caucasica, Boiss. Diagn. Ser. II. i. 87 = Ar. montana.
caucasica, Rupr. Fl. Caucas. 208.—Caucas.
cerastiifolia, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. cerastiifolia.
Cherleriana, Fenzl, l. c. 37 = Stellaria petraea.
Cherleriana, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = Ar. Cherleria.
Chowenausii, Reut. ex Nym. Consp. 119 = Ar. recurva.
ciliata, Crantz, Inst. ii. 408.—Italia.
condensata, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. condensata.
conferta, Jord. Pugill. 35 = Ar. tenuifolia.
confusa, Heldr. & Sart. ex Nym. Consp. 118 = Ar. trichocalycina.
corymbulosa, Boiss. & Bal. Diagn. Ser. II. vi. 87 = Ar. setacea.
corymbulosa, Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 57 = Ar. Barrelieri.
costata, Fenzl, Verbreit. Alsin. 37 = Ar. costata.
Cupaniana, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 19 = Stellaria media.
decandra, Dalla Torre, Atl. Alpenfl. 78.—Europ.
decipiens, Fenzl, Pugill. Fl. Nov. Syr. 15.—Syria.
Delastrei, Bor. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 12 = Ar. Barrelieri.
diandra, Guss. Fl. Sic. Syn. i. 501 = Ar. diandra.
dianthifolia, Boiss. Diagn. Ser. I. viii. 99.—As. Min.
dichotoma, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Minuartia dichotoma.
Dillenii, Vill. Fl. Dalphin. 80.—Gallia.
Drummondii, Fenzl, ex Torr. & Gray, Fl. N. Am. i. 675 = Stellaria Nuttallii.
Elatina, Crantz, Inst. ii. 405 = Elatine Hydropiper.
elegans, Fenzl, Verbreit. Alsin. tab. 18 = Ar. elegans.
elongata, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 19 = Stellaria media.
emarginata, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. emarginata.
erecta, Crantz, Inst. ii. 406 = Cerastium quaternellum.
erythrocepala, Boiss. Diagn. Ser. I. viii. 95.—As. Min.
eurytaenia, Boiss. l. c. Ser. II. vi. 85.—Graecia.
exilenia, Nym. Consp. 119 = Ar. procumbens.
Facchinii, Hausm. Fl. Tyr. 186.—Tyrol.
falcata, Griseb. Fl. Rumel. i. 200 = Ar. setacea.
fasciculata, Wahlenb. Fl. Lapp. 129 = Ar. fasciculata.
fastigiata, Bab. Man. ed. I. 51 = Ar. fastigiata.
formosa, Fenzl, in Flora, xxvi. (1843) 405.—Assyria.
Funkii, Jord. Pugill. 36 = Ar. fasciculata.
filiformis, Fenzl, ex Heynh. Nom. ii. 37 = Ar. picta.
Gerardi, Wahlenb. Fl. Carpat. 128 = Ar. verna.
Gimmelii, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. verna.

ALSINE :—
glabella, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 20 = Stellaria media.
glabra, A. Gray, Man. ed. II. 57 = Arenaria glabra.
glandulosa, Dulac, Fl. Hautes-Pyr. 246 = Ar. cerastiifolia.
glandulosa, Boiss. & Huet, Diagn. Ser. II. v. 61 = Ar. juniperina.
glandulosa, Mutel, ex Nym. Consp. 116 = Ar. linifolia.
globulosa, C. A. Mey. Verz. Pfl. Cauc. 219 = Ar. globulosa.
glomerata, Fenzl, Verbreit. Alsin. in tab. ad p. 46 = Ar. glomerata.
graminifolia, J. F. Gmel. Syst. 507 = Ar. Rosani.
grandiflora, Crantz, Inst. ii. 408 = Ar. grandiflora.
granuliflora, Fenzl, in Ledeb. Fl. Ross. i. 346 = Ar. setacea.
Griffithii, Boiss. Diagn. Ser. II. i. 89 = Ar. Griffithii.
grisebachii, Gren. & Godr. Fl. Fr. i. 252 = Ar. Villarsii.
groenlandica, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. groenlandica.
Gussonii, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 20 = Stellaria media.
Halophila, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 567.—Altai.
Heldreichiana, Boiss. & Heldr in Boiss. Fl. Orient. i. 677 = Ar. juniperina.
Helmii, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. Helmii.
hernarioides, Rion, ex Nym. Consp. 119 = Ar. arctioides.
heteromalla, Bieb. Fl Taur. Cauc. i. 350 = Ar. setacea
heterosperma, Guss. Fl. Sic. Syn. i. 501.—Sicil.
hirsuta, Fenzl, Verbreit. Alsin. in tab. 46 = Ar. recurva.
hirta, Hartm. Handb. Skand. Fl. ed. III. 104 = Ar. verna.
hispanica, Fenzl, Verbreit. Alsin. in tab. ad p. 46 = Queria hispanica.
hispida, Crantz, Inst. ii. 408 = Ar. hispida.
hybrida, Jord. Pugill. 35 = Ar. tenuifolia.
imbricata, C. A. Mey. Verz. Pfl. Cauc. 217 = Ar. imbricata.
inamoena, C. A. Mey. l. c. 218 = Ar. imbricata.
intermedia, Boiss. Fl. Orient. i. 685.—As. Min.
intricata, Martr. Pl. Crit. Tarn, 15.—Gallia.
irregularis, Vis. ex Boiss. Fl. Orient. i. 684 = Ar. globulosa.
Jacquini, Koch, Syn. Fl. Germ. ed. II. 125 = Ar. fasciculata.
Jacquiniana, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = Ar. fasciculata.
juniperina, Wahlenb. Fl. Lapp. 129 = Ar. juniperina.
Jurensi, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. Jurensi.
babylica, Pomel, Nouv. Mat. Fl. Atl. 827.—Afr. bor.
Kitaibelii, Nym. Consp. 116 = Ar. laricifolia.
lanceolata, Mert. & Koch, in Roehl. Deutsch. Fl. ed. III. ii. 275 = Ar. lanceolata.
lancifolia, Vis. Ic. Fl. Dalm. t. 84 = Ar. globulosa.
Langii, G. Reuss, Kvêt. Slov. (1853) 76 = Ar. laricifolia.
laricifolia, Crantz, Inst. ii. 407 = Ar. laricifolia.
laricifolia, Linn. ex Nym. Consp. 117 = Ar. setacea.
laricina, Crantz, Inst. ii. 408 = Spergula laricina.
laterifora, Crantz, l. c. 406 = Ar. laterifiora.
latifolia, Pum. Fl. Belg. 109.—Belg.
laxa, Jord. Pugill. 34 = Ar. tenuifolia.
leucocephala, Boiss. Diagn. Ser. I. i. 46.—As. Min.
libanotica, Boiss. l. c. viii. 98.—Libanus.
lineata, C. A. Mey. in Ledeb. Fl. Ross. = Ar. juniperina.
linifolia, Vis. Fl. Dalmat. iii. t. 178 = Ar. linifolia.
litnifolia, Borb. in Rad. Jugos. Akad. Zagrab, xxxvi. (1876) 166.—Europ. austr.
Lydia, Boiss. Diagn. Ser. II. i. 86 = Ar. tenuifolia.
macrocarpa, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. macrocarpa.
marginata, C. A. Mey. Verz. Pfl. Cauc. 217 = Spergularia media.
marginata, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Spergularia marginata.
marginata, Reichb. Fl. Germ. Excurs. 666. n. 3662 = Ar. media.
marginata, Schreb. Spicil. 21 = Spergularia pentandra.
marina, Webb & Berth. Phyt. Canar. i. 147 = Spergularia Bourgaeui.
marina, Mert. & Koch, in Roehl. Deutsch. Fl. ii. 293 = Spergularia media.

ALSINE :—
media, Linn. Sp. Pl. ed. I. 272 = Spergularia media.
medica, Vell. Fl. Flum. 127; iii. t. 106 = Drymaria cordata.
mediterranea, Gren. Fl. Massil. Advers. (1859) 8.—Gallia.
mexicana, Boiss. Diagn. Ser. I. i. 45 = Arenaria tenuifolia.
Meyeri, Boiss. l. c. Ser. I. viii. 96 = Arenaria globulosa.
mexicana, Bartl. ex Presl, Rel. Haenk. ii. 14 = Ar. mexicana.
Michauxii, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. stricta.
microcarpa, Fenzl, ex Torr. & Gray, Fl. N. Am. i. 674 = Ar. Pitcheri.
minuta, Rohrb. in Linnaea, xxxvii. (1871–72) 246.—Chili.
moehringia, Crantz, Inst. ii. 405 = Ar. muscosa.
moehringioides, Rohrb. in Linnaea, xxxvii. (1871–72) 247 (= *Hymenella moehringioides*).
mollugnea, Lag. Gen. et Sp. Nov. (1816) 18 = Ar. mollugnea.
montana, Crantz, Inst. ii. 406 = Ar. montana.
mucronata, Linn. Sp. Pl. ed. II. 889 = Ar. mucronata.
mucronata, Poir. Encyc. Suppl. iv. 310 = Ar. fasciculata.
multicaulis, Crantz, Inst. ii. 406 = Ar. multicaulis.
multinervis, Boiss. Fl. Orient. i. 683.—As. Min.
Munbyi, Boiss. Diagn. Ser. II. i. 85 = Ar. Munbyi.
muscorum, Adams, ex Ledeb. Fl. Ross. i. 349 = Ar. verna.
mutabilis, Lap. ex Nym. Consp. 118 = Ar rostrata.
nemorum, Schreb. Spicil. 20 = Stellaria nemorum.
nodosa, Crantz, Inst. ii. 408 = Sagina nodosa.
occulta, Kar. & Kir in Bull. Soc. Nat. Mosc. xv. (1842) 590 = Ar. polygonoides, artica.
octandra, Schur, Enum. Pl. Transs. 882, Index = Ar. arctioides.
Orphanides, Boiss. Diagn. Ser. II. i. 62 = Ar. tenuifolia.
pallida, Dum. Fl. Belg. 109; et in Bull. Soc. Bot. Belg. xii. (1873) 71.—Belg.
palustris, Kellogg, in Proc. Calif. Acad. iii. (1868) 61 = Ar. palustris.
paniculata, Fenzl, Verbreit. Alsin. 37 = Ar. paniculata.
parnassica, Boiss. & Sprun. Diagn. Ser. I. i. 46.—Graecia.
patula, A. Gray, Man. Bot. U. St. ed. II. 58 = Ar. patula.
pauciflora, Kit. ex Nym. Consp. 119 = Ar. verna.
pentandra, Crantz, Inst. ii. 408 = Spergularia pentandra.
peploides, Crantz, l. c. 406 = Ar. peploides.
Pestalozzae, Boiss. Diagn. Ser. I. viii. 99.—As. Min.
petraea, Jord. Obs. Pl. Crit. vii. (1849) 14 = Ar. rostrata.
pharnacoides, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. filiformis.
picta, Fenzl, l. c. = Ar. filiformis.
pilifera, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 588 = Spergula laricina.
pinifolia, Fenzl, Verbreit. Alsin. in tab. ad p. 46 = Ar. pinifolia.
Pitcheri, Wood, Cl.-Book, 260 = Ar. Pitcheri.
polycarpa, Crantz, Inst. ii. 405 = Polycarpon tetraphyllum.
Preslii, Reuss, Kvêt. Slov. (1853) 76 = Ar. graminifolia.
procumbens, Crantz, Inst. ii. 404 = Sagina procumbens.
procumbens, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. geniculata.
propinqua, Lange, Fl. Dan. t. 2908 = Ar. verna.
prostrata, Forsk. Fl. Aegypt. Arab. 207 = Polycarpea prostrata.
pubescens, Presl, ex Nym. Consp. 117 = Ar. tenuifolia.
pulvinaria, Boiss. Diagn. Ser. I. i. 48.—Lydia.
Pumilio, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. arctica.
purpurascens, Bubani, in Nuov. Ann. Sc. Nat. Bologn. (1843) 94.—Italia!
purpurea, Heynh. Nom. i. 38 = Spergularia purpurea.
radicans, Guss. Fl. Sic. Syn. i. 502 = Spergularia rubra.
ramosissima, Fenzl, ex Schur, Enum. Pl. Transs. 118 = Ar. fasciculata.
recurva, Wahlenb. Fl. Helv. 87 = Ar. recurva.
repens, Vell. Fl. Flum. 127; iii. t. 105 = Stellaria media.

ALSINE:—
rimarum, Boiss. & Bal. in Boiss. Fl. Orient. i. 678.—Cilicia.
Rosani, Guss. Fl. Sic. Syn. i. 498 = Arenaria Rosani.
rosea, Presl, ex Nym. Consp. 119 = Ar. procumbens.
Rossi, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. stricta.
rostrata, Fenzl, l. c. 46 = Ar. rostrata.
rotundifolia, Stokes, Bot. Mat. Med. ii. 537 = Drymaria cordata.
Royleï, Fenzl, ex Hook. f. Fl. Brit. Ind. i. 236 = Ar. follosa.
rubella, Schrenk, ex Ledeb. Fl. Ross. i. 349 = Ar. verna.
rubella, Wahlenb. Fl. Lapp. 128. t. 6 = Ar. verna.
rubra, Webb & Berth. Phyt. Canar. i. 148 = Spergularia canariensis.
rubra, C. A. Mey. Verz. Pfl. Cauc. 217 = Spergularia salsuginea.
rubra, Crantz, Inst. ii. 407 = Spergularia rubra.
rupestris, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. rupestris.
Sagina, Crantz, Inst. ii. 405 = Bartonia tenella.
saginoides, Crantz, l. c. 406 = Sagina Spergella.
saxatilis, Crantz, l. c. 407 = Ar. saxatilis.
saxatilis, Wahlenb. Fl. Helv. 87 = Ar. verna.
Saxifraga, Boiss. Diagn. Ser. I. i. 47 = Ar. Saxifraga.
scabra, Stokes, Bot. Mat. Med. ii. 538 = Stellaria Holostea.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 47. —Abyss.
scleranthoides, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 400.—Caucas.
scleranthoides, Boiss. Diagn. Ser. II. i. 88 = Ar. setacea.
sedoides, Froel. in Koch, Syn. Fl. Germ. ed. I. 114 = Ar. sedoides.
segetalis, Linn. Sp. Pl. 272 = Spergularia segetalis.
Segueiri, Turra, ex Sacc. in Atti Ist. Venet. Ser., III. ix. (1863–64) 486.—Italia.
serpyllifolia, Crantz, Inst. ii. 406 = Ar. serpyllifolia.
setacea, Mert. & Koch, in Roehl. Deutsch. Fl. iii. 286 = Ar. setacea.
sinaica, Boiss. Diagn. Ser. I. viii. 100 = Ar. picta.
Smithii, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. globulosa.
spinulifora, Fenzl, l. c. = Habrosia spinulifora.
squarrosa, Fenzl, ex A. Gray, Man. Bot. U. St. ed. II. 57 = Ar. squarrosa.
striata, Crantz, Inst. ii. 407 = Ar. striata.
striata, Gren. in Mém. Soc. Doubs, (1841) 33. t. 1. f. 1 = Ar. laricifolia.
stricta, Mert. & Koch, in Roehl. Deutsch. Fl. iii. 278 = Ar. stricta.
stricta, Wahlenb. Fl. Lapp. 127 = Ar. lapponica.
subtilis, Fenzl, ex Boiss. Fl. Orient. i. 687 = Ar. tenuifolia.
subulifolia, Guss. Fl. Sic. Syn. i. 503 = Ar. mucronata.
succulenta, Delile, Fl. Egypt. 211. t. 24 = Polycarpon succulentum.
taurica, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 319 = Ar. Villarsii.
Tchihatchewii, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 247.—Ar. Min.
tenella, Pers. Fl. Transs. 113 = Ar. austriaca.
tenuifolia, Crantz, Inst. ii. 407 = Ar. tenuifolia.
tenuissima, Pomel, Nouv. Mat. Fl. Atl. 237.—Afr. bor.
tetraquetra, Crantz, Inst. ii. 406 = Ar. tetraquetra.
thymifolia, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. thymifolia.
trichocalycina, Heldr. & Sart. Diagn. Ser. II. i. 87.—Ital.; Graecia.
trifolia, Baring-Gould, ex Bab. in Journ. Linn. Soc. xi. (1871) 298 = Ar. norvegica.
trinervia, Crantz, Inst. ii. 406 = Ar. trinervia.
uliginosa, Vill. Prosp. Dauph. 48 = Stellaria aquatica.
umbellata, Lam. Fl. Fr. iii. 45 = Holosteum umbellatum.
umbellifera, Boiss. Diagn. Ser. II. v. 61.—Cilicia.
unilateralis, Moench, Meth. 228 = Spergularia segetalis.
uralensis, Clerc, in Bull. Soc. Oural. iv. 106.—Ural.
vaginosa, Boiss. & Orphan. Diagn. Ser. II. vi. 636 = Ar. glomerata.
verna, Wahlenb. Fl. Lapp. 129; Bartl. Beitr. ii. 63 = Ar. verna.
verticillata, Noroscha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 6.—Javn.
Villarsii, Fenzl, in Ledeb. Fl. Ross. 350 = Ar. juniperina.

ALSINE:—
Villarsii, Mert. & Koch, in Roehl. Deutsch. Fl. iii. 282 = Arenaria Villarsii.
Villarsiana, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = praec.
villosa, Heynh. Nom. i. 39 = Ar. villosa.
viscidula, Thouin, ex Nym. Consp. 117 = Ar. tenuifolia.
viscosa, Schreb. Spicil. 30 = Ar. tenuifolia.
vulgaris, Moench, Meth. 228 = Stellaria media.

ALSINELLA, Dill. ex Linn. Gen. ed. I. 118 (1737) = **Sagina**, Linn. (Caryophyll.).

ALSINELLA, Hornem. Nom. 33 (1827) = **Spergularia**, J. & C. Presl (Caryophyll.).
media, Hornem. l. c. = S. marina.

ALSINELLA, [Sw.] Summa Veg. Scand. 17 (1814) pro parte; Benth. in Wall. Cat. n. 680 (1832) = **Stellaria**, Linn. (Caryophyll.).
biflora, Sw. l. c. = S. biflora.
Wallichiana, Benth. l. c. = S. media.

ALSINELLA, [Sw.] Summa Veg. Scand. 17 (1814) pro parte; S. F. Gray, Nat. Arr. Brit. Pl. ii. 655 (1821) = **Arenaria**, Linn. (Caryophyll.).
ciliata, S. F. Gray, l. c. = Ar. ciliata.
rubella, Sw. Summa Veg. Scand. 17 = Ar. verna.
stricta, Sw. l. c. = Ar. stricta.
trinervis, S. F. Gray, Nat. Arr. ii. 655 = Ar. trinervis.
viscosa, Hartm. Fl. Dan. t. 1764 = Ar. tenuifolia.
serpyllifolia, S. F. Gray, Nat. Arr. ii. 665 = Ar. serpyllifolia.

ALSINELLA, Moench, Meth. 223 (1794) = **Cerastium**, Linn. (Caryophyll.).
erecta, Moench, l. c. = C. quaternellum.

ALSINIDENDRON, H. Mann, in Proc. Bost. Soc. Nat. Hist. x. (1866) 311. *CARYOPHYLLEAE*, Benth. & Hook. f. i. 978.
trinerve, *H. Mann, l. c.* 312.—Isa. Sandvic.

ALSOBIA, Hanst. in Linnaea, xxvi. (1853) 207 = **Episcia**, Mart. (Gesnetrac.).
antirrhina, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 114.—Am. centr.
punctata, Hanst. in Linnaea, xxvi. (1853) 207 = E. punctata.

ALSOCYDIA, Mart. ex DC. Prod. ix. 149, &c. (1845) = **Bignonia**, Linn. &c.
acutifiora, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Lundia latifolia.
angustidens, Mart. ex DC. l. c. ix. 179 = Cuspidaria angustidens.
cordata, Mart. ex DC. l. c. 168 = Bign. cordifolia.
crateraphora, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Bign. crateraphora.
dichotoma, Mart. ex DC. l. c. ix. 150 = Bign. corymbifera.
erubescens, Mart. ex DC. l. c. ix. 157 = Bign. erubescens.
firmula, Mart. ex DC. l. c. 146 = Bign. brachypoda.
glandulosa, Mart. ex DC. l. c. 149, 164 = Bign. glandulosa, lanceolata.
glutinosa, Mart. ex DC. l. c. 162 = Bign. glutinosa.
hebantha, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Lundia hebantha.
hirsuta, Mart. ex DC. l. c. ix. 156 = Bign. trichoclada.
laxianthera, Mart. ex DC. l. c. 180 = Lundia cordata.
longidens, Mart. ex DC. l. c. 190 = Anemopaegma longidens.
nitidula, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Lundia nitidula.
notorophila, Mart. ex Buek, l. c. = Bign. notorophila.
tetragonocaulos, Mart. ex DC. l. c. ix. 169 = Bign. tetragonocaulos.
triplinervia, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Bign. triplinervia.
variabilis, Mart. ex DC. l. c. ix. 149 = Bign. varians.
villosa, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Bign. velutina.
virginalis, Mart. ex Buek, l. c. = Lundia virginalis.

ALSODEIA, Thou. Hist. Vég. Afr. ii. 55. t. 17, 18 (1807). *VIOLARIEAE*, Benth. & Hook. f. i. 113.
CERANTHERA, Beauv. Fl. Owar. et Ben. ii. 10 (1807).

ALSODEIA:—
CERATANTHERA, Beauv. Fl. Owar. et Ben. ii. 10, in note (1807).
CONOHORIA, Anbl. Fl. Gui. 289. t. 95 (1775).
CONORIA, Juss. Gen. 387 (1789).
CUSPA, Humb. Relat. i. 366 (1814) ex H. B. & K. Nov. Gen. et Sp. vii. 242.
DIORYKTANDRA, Hassk. Retzia, 125 (1858).
GONOHORIA, G. Don, Gen. Syst. i. 340 (1831).
JUEGENSIA, Spreng. Syst. iii. 50 (1826).
MEDUSA, Lour. Fl. Cochinch. 406 (1790).
MEDUSULA, Pers. Syn. ii. 216 (1807).
PASSALIA, Soland. ex R. Br. in Tuckey, Congo, 440 (1818).
PASSOURA, Anbl. Fl. Gui. [Suppl.] 21. t. 280 (1775).
PENTALOBA, Lour. Fl. Cochinch. 124 (1790).
PHYSIPHORA, Soland. ex R. Br. in Tuckey, Congo, 441 (1818).
PROSTHESIA, Blume, Bijdr. 866 (1826).
RIANA, Anbl. Fl. Gui. 287. t. 94 (1775).
RINOREA, Anbl. l. c. 285. t. 93 (1775).
VARECA, Roxb. Fl. Ind. i. 647 (1832).
albicaulis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 557.—N. Granat.
andina, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 366. —N. Granat.
angustifolia, *Thou. Hist. Vég. Afr.* ii. 57. t. 18. f. 2.—Madag.
arborea, *Thou. l. c.* 57. t. 17. f. 1.—Madag.
arillosa*flora*, *Welw. in Oliver, Fl. Trop. Afr.* i. 108.—Afr. trop.
Aucuparia, *Welw. l. c.* 100.—Afr. trop.
auriculata, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 319.—Madag.
bahiensis, *Moric. Pl. Nouv. Am.* 68. t. 46.—Bras.
bengalensis, *Wall. in Trans. Med. & Phys. Soc. Calc.* vii. (1836) 224.—Ind. or.
brachypetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 556.—Afr. trop.
brevipes, *Benth. in Hook. Lond. Journ. Bot.* iv. (1842) 106.—Guiana.
Browni, Korth. in Neder. Kruid. Arch. i. (1848) 361 = zeylanica.
caledonica, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 556.—N. Caled.
calophylla, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 311.—Madag.
calycina, *Tul. l. c.* 318.—Madag.
castanesefolia, *Spreng. Syst.* iv. *Cur. Post.* 99.—Bras.
castanoides, *Welw. in Oliver, Fl. Trop. Afr.* i. 110.—Afr. trop.
caudata, *Oliver, l. c.* 107.—Afr. trop.
cauliflora, *Oliver, l. c.* 111.—Afr. trop.
ceylanica, Arct. vide zeylanica.
chrysodasys, *Miq. Fl. Ind. Bat. Suppl.* i. 390.—Sumatra.
corylifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 559.—Java.
Cuspa, *Spreng. Syst.* iv. *Cur. Post.* 99.—Cumana.
cymulosa, *Miq. Fl. Ind. Bat. Suppl.* i. 116.—Java.
cymulosa, *Welw. in Oliver, Fl. Trop. Afr.* i. 108.—Afr. trop.
dasycoula, Miq. Fl. Ind. Bat. Suppl. i. 391 = mollis.
dasypyrsia, *Miq l. c.*—Sumatra.
decora, *Trim. in Journ. Bot.* xxiii. (1885) 208.—Zeylan.
deflexa, *Benth. Bot. Voy. Sulph.* 57.—Peruv.
dentata, *Beauv. Fl. Owar. et Ben.* ii. 11. t. 95.—Afr. trop.
distichа, *Zoll. in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 216.—Malaya.
echinocarpa, *Korth. in Neder. Kruid. Arch.* i. (1848) 360.—Sumatra.
ellipsica, *Oliver, Fl. Trop. Afr.* i. 108.—Afr. trop.
falcata, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. i. 366. —Bras.
fasciculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 559.—Ins. Philipp.
flavescens, *Spreng. Syst.* i. 806.—Guiana.
floribunda, Moric. Pl. Nouv. Am. 70. t. 47 = castaneaefolia.
glabra, *Burgersdyke, in Miq. Fl. Sumgh.* 122.—Sumatra.
Gossypium, *Tul. in Ann. Sc. Nat. Sér. III.* vii. (1847) 367.—N. Granat.
grandis, *Miq. Fl. Ind. Bat. Suppl.* i. 391.—Sumatra.
Griffithii, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 197. —Burma.

ALSODEIA:—
guianensis, *Eichl. in Mart. Fl. Bras.* xiii. t. 867.—Bras.
guineensis, *Spreng. Syst.* i. 806.—Afr. trop.
haplobotrys, *Hassk. in Verol. Kon. Akad. Wetensch.* iv. (1856) 187.—Java.
Horneri, *Korth. in Neder. Kruid. Arch.* i. (1848) 860. —Borneo.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. ii. 117.—Java.
Horsfieldii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 559.—Java.
ilicifolia, *Walp. in Trans. Linn. Soc.* xxvii. (1869) 12. —Afr. trop.
integrifolia, *Steud. Nom.* ed. II. i. 64.—Bras.
javanica, *Miq. Fl. Ind. Bat.* i. ii. 117.—Sumatra.
lanceolata, *Oudem. in Arch. Néerl.* ii. (1867) 196; *et in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 68.—Ind. or.
latifolia, *Thou. Hist. Vég. Afr.* ii. 57. t. 18. f. 3.— Madag.
laxiflora, Benth. in Hook. Lond. Journ. Bot. iv. (1842) 107 = guianensis.
leucocalax, *Tul. in Ann. Sc. Nat. Sér.* V. ix. (1868) 312.—Madag.
Lindeniana, Tul. l. c. Sér. III. vii. (1847) 364 = guianensis.
longiflora, *Oudem. in Arch. Néerl.* ii. (1867) 195; *et in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. (1867) 68 = Guiana.
longipes, *Tul. in Ann. Sc. Nat. Sér.* V. ix. (1868) 300. —Madag.
longiracemosa, *Kurz, in Journ. As. Soc. Beng.* xxxix (1870) ii. 68.—Ind. or.
macrocarpa, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. t 885.—Bras.
macrophylla, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 468. t. 19.—Ins. Timor.
maculata, *Tul. in Ann. Sc. Nat.* Sér V. ix. (1868) 316. —Madag.
Maingayi, *Hook. f. & Thoms. ex Fl. Brit. Ind.* i. 188. —Malacca.
marginata, *Triana & Planch. in Ann. Sc. Nat.* Sér IV. xvii. (1862) 127.—Nov. Granat.
Martini, *Turcz. in Bull. Soc. Nat Mosc.* xxxvi. (1863) i. 557.—Guin.
Maximiliana, *Eichl. in Mart. Fl. Bras.* xiii. t. 881.— Bras.
megapotamica, *Spreng. Syst.* iv. *Cur. Post.* 99.—Bras.
mollis, *Hook. f. & Thoms. ex Fl. Brit. Ind.* i. 188.— Tenasserim.
monticola, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 316.—Madag.
mutica, *Tul. l. c.* 318.—Madag.
nigricans, Boiv. ex Tul. l. c. 305 = pubescens.
obtusa, *Korth. in Neder. Kruid. Arch.* i. (1848) 859.— Borneo.
owarensis, *Spreng. Syst.* i. 806.—Afr. trop.
pallida, *Klotzsch, in Schomb. Fauna et Fl. Guy.* 1165. —Guiana.
paniculata, *Mart. Nov. Gen. et Sp.* i. 30.—Bras.
paradoxa, *Blume, ex Oudem. in Arch. Neerl.* ii. (1867) 204; *et in Miq. Ann. Mus. Lugd. Bat. Bot.* iii. 71. —Java.
pauciflora, *Thou. Hist. Vég. Afr.* ii. 57. t. 17.— Madag.
Perrini, *Spreng. Syst.* i. 807.—Am. austr.
Pervilli, *Oudem. in Arch. Néerl.* ii. (1867) 206; *et in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 72.—Madag.
physiphora, *Mart. Nov. Gen. et Sp.* i. 28.—Bras.
Piparea, Spreng. Syst. i. 807 = Casearia dentata.
prunifolia, *Spreng. l. c.*—Guiana.
pubescens, *Thou. Hist. Vég. Afr.* ii. 57. t. 18. f. 3.— Madag.
pulchra, *Benth. in Hook. Journ. Bot.* iv. (1842) 106. —Gutiana.
pyginiflora, *Oudem. in Arch. Néerl.* ii. (1867) 205; *et in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 71.— Madag.
racemosa, Hook. f. & Thoms. ex Fl. Brit. Ind. i. 187 = longiracemosa.
racemosa, *Mart. & Zucc. Nov. Gen.* i. 29. t. 20.— Bras.
Ragnelii, Walp. Ann. ii. 67 = Ionidium atropurpureum.
Rimosa, Spreng. Syst. i. 807 = paniculata.
Roxburghii, *Hook. f. & Thoms. ex Fl. Brit. Ind.* i. 188.—Ind. or.
rubra, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 315.— Madag.

ALSODEIA:—
rugosa, *Miq. Fl. Ind. Bat. Suppl.* i. 390.—Sumatra.
sclerocarpa, *Burgersdyke, in Miq. Pl. Fungd.* 122.— Sumatra.
scmiglyrata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 559.—Java.
sessilis, *Spreng. Syst.* i. 806.—Cochinch.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 402.—Ins. Socotra.
spinosa, *Bowdin, ex Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 307.—Madag.
Spruceі, *Eichl. in Mart. Fl. Bras.* xiii. t. 885.—Bras.
squamosa, *Boivin, ex Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 307.—Madag.
Storckii, *Seem. in Journ. Bot.* iii. (1864) 75.—Ins. Fiji.
subintegrifolia, *Beauv. Fl. Owar. et Ben.* ii. 11. t 66. —Afr. trop.
sylvatica, *Seem. Bot. Voy. Herald*, 79. t. 40.— Panama.
trigonocarpa, *Zoll. & Reichb in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 216.—Malaya.
ulmifolia, *Spreng. Syst.* i. 807.—N. Granat.
verticillata, *Baron, ex Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 317.—Madag.
virgata, Thw. ex Hook. f. Fl. Brit. Ind. i. 180 = Scyphilandra virgata.
viridiflora, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 310.—Madag.
Wallichiana, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 187.—Penang.
Welwitschii, *Oliver, Fl. Trop. Afr.* i. 110.—Afr. trop.
zeylanica, *Thw. Enum. Pl. Zeyl.* 21.—Ind. or.

ALSODEIOPSIS, Oliver, in Benth. & Hook. f. Gen. i. 996 (1867). *OLACINEAE*, Benth. & Hook. l. c.
Mazzi, *Oliver, in Journ. Linn. Soc.* x. (1869) 48.— Afr. trop.

ALSOLINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349 = Linum, Tourn. *maritimum*, Fourr. l. c. = L. maritimum.

ALSOMITRA, M Roem. Syn. Pepon. 117 partim (1846). *CUCURBITACEAE*, Benth. & Hook. f. i. 840.
angulata, M. Roem. l. c. 117 = Zanonia indica.
Beccariana, *Cogn. in DC. Monog. Phan.* iii. 932.—Ins. Kei.
brasiliensis, *Cogn. in Mart. Fl. Bras.* vi. iv. 115.— Bras.
capricornica, *F Muell. Fragm.* vii. 61.—Austral.
cissoides, M. Roem. Syn. Pepon. 136 = Gynostemma pedata.
clavigera, *M. Roem. l. c.*; *Benth. & Hook. f. Gen.* i. 840.—Ind. or.
heterosperma, M. Roem. l. c. = Gomphogyne heterosperma.
Hookeri, *F. Muell. Fragm.* vi. 188.—Austral.
laxa, M. Roem. Syn. Pepon. 118 = Gynostemma laxa.
macrocarpa, M. Roem. l. c. 117 = Zanonia macrocarpa.
pedatifolia, *Cogn. in Mart. Fl. Bras.* vi. iv. 116.— Bras.
sarcophylla, *M. Roem. Syn. Pepon.* 118.—Burma; Siam.
Schefferiana, *Cogn. in DC. Monog. Phan.* iii. 932.— Ins. Celebes.
Stephensiana, *Cogn. l. c.* 934.—Ins. Barnardi.
timorensis, *M. Roem. Syn. Pepon.* 117.—Ins. Timor.

ALSTONIA, Mutis, ex Linn. f. Suppl. 89 (1781) = Symploocos, Linn. (Styrac.).
ciliata, Benth. Pl. Hartw. 48 = S. ciliata.
theaeformis, Linn. f. l. c. 264 = S. Alstonia.

ALSTONIA, Scop. Introd. 198 (1777) = Landolphia, R. Br. (Apocyn.).

ALSTONIA, R. Br. in Mem. Wern. Soc. i. (1809) 75. *APOCYNACEAE*, Benth. & Hook. f. ii. 705.
BLABEROPUS, A. DC. Prod. viii. 410 (1844).
PALA, [Rheede] Jus. in Ann. Mus. Par. xv. (1825) 246.
acuminata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140.— Ins. Ceram.
angustifolia, *Wall. Cat.* n. 1650.—Malaya.
angustiloba, *Miq. Fl. Ind. Bat.* ii. 488.—Malaya.
Batissa, Blanco, Fl. Filip. ed. II. Suppl. 589 = macrophylla.
calophylla, *Miq. Fl. Ind. Bat.* ii. 489.—Java.

ALSTONIA:—
constricta, *F. Muell. Fragm.* i. 57.—Austral.
costata, *R. Br. in Mem. Wern. Soc.* i. (1809) 75.—Ins. Tahiti.
costata, Wall. Cat. n. 1649 = macrophylla.
costulata, Miq. Fl. Ind. Bat. Suppl. 556 = Dyera costulata.
cuneata, Wall. Cat. n. 1645 = spathulata.
Deplanchei, *Heurck & Muell. Arg. in Flora*, liii. (1870) 171.—N. Caled.
edulis, *G. Benn. in Journ. Bot.* v. (1867) 150.—N. Caled.
exima, *Miq. Fl. Ind. Bat. Suppl.* 555.—Ins. Banca.
grandifolia, *Miq. l. c.*—Sumatra.
Kurzii, *Hook. f. Fl. Brit. Ind.* iii. 643.—Ins. Andaman.
lanceolata, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 199.—N. Caled.
Legouxiae, *Heurck & Muell Arg. in Flora*, liii. (1870) 170.—N. Caled.
Lenormandii, *Heurck & Muell. Arg. l. c.* 172.—N. Caled.
lineata, *Benth Fl Austral.* iv. 314.—Austral.
longisaima, *F. Muell. Papuan Plants*, 91.—N. Guin.
lucida, D. Don, Prod. Fl. Nep. 131 = Trachelospermum fragrans.
macrophylla, *Wall. Cat.* n. 1648 (*Fl. Brit. Ind* iii. 643) —Malaya.
mollis, *Benth. Fl. Austral.* iv. 315.—Austral.
nerilifolia, *D. Don, Prod. Fl. Nep.* 131.—Reg. Himal.
oleandrifolia, Lodd. ex Loud. Hort. Brit. 67 = scholaris.
ophiocylindra, *F. Muell. Fragm.* i. 57.—Austral.
plumosa, *Labill. Sert. Austr. Caled.* 28. t. 32.—N. Caled
polyphylla, *Miq. Fl. Ind. Bat. Suppl.* 556.—Ins. Banca.
quaternata, *Heurck & Muell. Arg. in Flora*, liii. (1870) 170.—N. Caled.
Roeperi, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 201.—N. Caled.
scholaris, *R. Br. in Mem. Wern. Soc.* i. (1809) 75.— Geront. trop.
sences, *Blume, Bijdr.* 1038.—Malaya.
spatulata, *Blume, l. c.* 1037.—Malaya.
spectabilis, *R. Br. in Mem. Wern. Soc.* i. (1809) 75.— Ins. Timor.
spectabilis, Kurz, For. Fl. Burma, ii. 183 = Kurzii.
subsessilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140.— Amboina.
venenata, *R. Br. in Mem. Wern. Soc.* i. (1809) 77.—Ind. or.
verticillosa, *F. Muell. Fragm.* vi. 116.—Austral.
Vieillardii, *Heurck & Muell. Arg. in Flora*, liii. (1870) 171.—N. Caled.
villosa, *Blume, Bijdr.* 1038.—Java; Austral.
villosa, Seem. Fl. Vit. 161 = vitiensis.
vitiensis, *Seem. l. c.* 480.—Ins. Fiji.

ALSTROEMERIA, Linn. Diss. Alstroem. 8 (1762).
AMARYLLIDEAE, Benth. & Hook. f. iii. 786.
LIGTU, Adans. Fam. p. 90 (1763).
LILAVIA, Rafin. Fl. Tellur. iv. 36 (1836).
PRIOPETALON, Rafin. Fl. Tellur. iv. 34 (1836).
acutifolia, Link & Otto, Ic. Pl. Rar. Hort. Bot. Berol. 57. t. 29 = Bomarea acutifolia.
affinis, Mart. & Gal. in Bull. Acad. Brux. x. (1843) reimpr. 6 = Bomarea edulis.
altilinea, *Presl, Rel. Haenk.* i. 121. t. 22.—Chili.
amoena, Sallsb. Prod. 248 = A. Pelegrina.
anceps, *Ruiz & Pav. Fl. Per.* iii. 61.—Peruv.
andina, Phil. in Linnaea, xxix. (1857–58) 69 = Bomarea.
angustifolia, Herb. Amar. 96 = A. Ligtu.
araucana, *Phil. in Anales Univ. Chil.* (1873) 547.— Chili.
argento-vittata, *Lem. Illustr. Hortic.* iv. (1857), *Misc.* 98; vi. (1858) t. 192.—Bras.
aurantiaca, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 206.—Chili.
aurea, *R. Grah. in Edinb. Phil. Journ.* (1838) 181.— Chili.
aurea, Meyen, Reise, i. 311 = aurantiaca.
Banksiana, M. Roem. Syn. Ecast. 259 = pulchella.
Barklayana, *Hort. ex Loud. Hort. Brit. Suppl.* ii. 606. —Hab.?
Berterolana, *Penny, ex Loud. l. c.*—Chili.
bicolor, Hook. Exot. Fl. i. t. 95 = A. Ligtu.
bracteata, *Ruiz & Pav. Fl. Per.* iii. 61.—Peruv.
brasiliensis, *Spreng. Syst.* ii. 81.—Bras.
Brademeyeriana, Willd. ex Kunth, Enum. Pl. ii. 808 = Bomarea multiflora.

ALSTROEMERIA :—
Burchellii, *Baker, in Journ. Bot.* xv. (1877) 262.—Bras. Caldasiana, Hensl. Bot. Centr. Am. Bot. iii. 326 = seq.
Caldasii, H. B. & K. Nov. Gen. et Sp. i. 328 = Bomarea Caldasiana.
caryophyllacea, *Jacq. Schoenbr.* vi. 33. t. 495 — Bras.
chilensis, *Y. Cver. ex Loud. Hort. Brit. Suppl.* ii. 608, nomen.—Chili.
chilensis, *Lem. in Fl. des Serres*, Sér. I. i. (1845) 98.—Chili.
chilotensis, *Phil. in Linnaea*, xxix. (1857–58) 71.—Ins. Chiloe.
chorillensis, *Herb. in Bot. Reg.* (1843) *Misc.* 64.—Chili.
ciliata, *Poepp. Fragm. Syn.* 6.—Chili.
citrina, Phil. in Linnaea, xxxiii. (1864–65) 264 = versicolor.
coccinea, Ruiz & Pav. Fl. Per. iii. 62 = Bomarea coccinea.
concolor, Steud. in Lechl. Berb. Am. Austr. 58 = aurantiaca.
cordifolia, *Ruiz & Pav. Fl. Per.* iii. 64.—Peruv.
crispata, *Phil. in Linnaea*, xxix. (1857–58) 70.—Chili.
crocea, *Phil. l. c.* xxxii. (1864–65) 263.—Chili.
crocea, Ruiz & Pav. Fl. Per. iii. 61 = Bomarea crocea.
Cummingiana, Herb. Amar. 96, 396 = versicolor.
Cunea, *Veil. Fl. Flum.* iii. t. 121 ; *Seub. in Mart. Fl. Bras.* iii. 172.—Bras.
Cunha, Vell. l. c. 181 = præc.
Curtisiana, E. Mey. ex Presl, Rel. Haenk. ii. 122 = A. Ligtu.
Curtisiana, Schult. f. Syst. vii. 735 = caryophyllacea.
declinata, Poepp. & Endl. Nov. Gen. et. 44. t. 160 = Bomarea dispar.
densiflora, Herb. in Bot. Mag. t. 5581 = Bomarea tomentosa.
denticulata, *Ruiz & Pav. Fl. Per.* iii. 62.—Peruv.
Despreuxia, *Poep. Cat. Orto Bot. Nap.* 7.—Hab. ?
Dizai, *Phil. in Linnaea*, xxxiii. (1864) 261.—Chili.
Didierana, *Jaub. in Bull. Soc. Bot. Fr.* viii. (1861) 118. —Peruv.
distichifolia, *Ruiz & Pav. Fl. Per.* iii. 60,—Peruv.
distichophylla, Spreng. Syst. ii. 80 = Bomarea distichophylla,
dulcis, Hook. Bot. Misc. ii. (1831) 237. t. 95 = Bomarea glaucescens.
edulis, Andr. ex Poir. Encyc. Suppl. iv. 348 = Bomarea edulis.
Errembaultii x, *Hort. ex Lem. in Fl. des Serres*, Sér. I. iii. (1847) 269.
exserens, Meyen, Reise, i. 84 = A. Ligtu.
Feuillaeana, E. Mey. ex Presl, Rel. Haenk. ii. = A. Ligtu.
filipendula, Seub. in Mart. Fl. Bras. iii. t. 130 = caryophyllacea.
fimbriata, Ruiz & Pav. Fl. Per. iii. 63 = Bomarea fimbriata.
flava, *Phil. in Linnaea*, xxxiii. (1864–65) 263.—Chili.
floribunda, *H. B. & K. Nov. Gen. et Sp.* i. 323.— Ecuador.
Flos-Martini, [Ker-Gawl. in] Bot. Reg. t. 781 = A. Ligtu.
fluminensis, *M. Kuntn. Syn. Enast.* 265.—Bras.
foliosa, *Mart. in Schult. f. Syst.* vii. 740.—Bras.
formosa, Pers. Syn. i. 381 = seq.
formosissima, Ruiz & Pav. Fl. Per. iii. 64 = Bomarea formosissima.
Gardneri, *Baker, in Journ. Bot.* xv. (1877) 261.—Bras.
Gayana, *Phil. in Linnaea*, xxix. (1857–58) 71.—Chili.
glaucescens, H. B. & K. Nov. Gen. et Sp. i. 323 = Bomarea glaucescens.
gloriosa, Cham. & Schlecht. in Linnaea, vi. (1831) 51 = Bomarea edulis.
grandifolia, *H. B. & K. Nov. Gen. et Sp.* i. 325.—Bras.
haemantha, *Ruiz & Pav. Fl. Per.* iii. 60.—Chili.
Herbertiana, M. Roem. Syn. Enast. 250 = revoluta.
hirsuta, H. B. & K. Nov. Gen. et Sp. i. 325 = Bomarea hirsuta.
hirtella, H. B. & K. l. c. 324 = Bomarea edulis.
hirtella, *Phil. in Linnaea*, xxix. (1857–58) 70.—Chili.
Hookeri, Lodd. Bot. Cab. t. 1272 = A. Ligtu.
Hookeriana, *C. Gay, Fl. Chil.* v. 90.—Chili.
Hookeriana, Schult. f. Syst. vii. 732 = A. Ligtu.
inaequalis, *Phil. in Linnaea*, xxix. (1857–58) 67.—Chili.
inconspicua, *Phil. Anales Univ. Chil.* (1873) 546.— Chili.
Inodora, *Herb. Amar.* 90. t. 2. f. 1.—Bras.
insignis, *Annls, ex Hérincq. L'Hortic. Franç.* iv. (1854) 2.—Peruv.
Isabellana, *Herb. Amar.* 88, 396. t. 6. f. 4–6.—Bras.

ALSTROEMERIA :—
Jacobi, *Hort. ex Neumann, Ann. Fl. et Pomona*, (1884) t. 29.—Hab. ?
Jacquiniana, Lem. in Fl. des Serres, Sér. I. ii. (Dec. 1846) t. 182 = Bomarea Jacquiniana.
Kingii, Phil. in Anales Univ. Chil. (1873) 546 = versicolor.
Kunziana, *Klotzsch, ex Kunth, Enum. Pl.* v. 769.— Chili.
laxa, *Ruiz & Pav. Fl. Per.* iii. 64.—Peruv.
Ligtu, Cranz, Inst. i. 467 = A. Ligtu.
Ligtu, Curt. Bot. Mag. t. 125 = caryophyllacea.
Ligtu, *Linn. Dis. Alstroem.* 10.—Chili.
linearifolia, Lindl. Bot. Reg. (1834) t. 58 = chorillensis.
linearifolia, *Ruiz & Pav. Fl. Per.* iii. 60 t. 289.—Peruv.
linifolia, H. B. & K. Nov. Gen. et Sp. i. 322 = Bomarea linifolia.
longistaminea, *Mart. in Schult. f. Syst.* vii. 739.— Bras.
longistyla, *Schenk, in Mart. Fl. Bras.* iii. t. 178.— Bras.
Macrocona, *Herb. Amar.* 90.—Chili.
macrocarpa, *Ruiz & Pav. Fl. Per.* iii. 63.—Peruv.
magnifica, *Herb. in Bot. Reg.* (1843) *Misc.* 64.— Chili.
Meyeniana, *Y. C. Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* t. (1843) 440.—Chili.
minuta, Mart. & Gal. in Bull. Acad. Brux. x. (1843) repr. 6 = Bomarea edulis.
montana, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 168 = Ixiolirion montanum.
monticola, *Mart. in Schult. f. Syst.* vii. 739.—Bras.
multiflora, Linn. f. Suppl. 207 = Bomarea multiflora.
mutabilis, Kunze, ex Kunth, Enum. Pl. v. 777 = aurantiaca.
Neillsana, Herb. Amar. 94 = spathulata.
Neillii, Gill. ex Hook. in Bot. Mag. t. 3105 = præc.
nemorosa, Gardn. in Bot. Mag. t. 3958 = inodora.
nivalis, Meyen, Reise, i. 315 = versicolor.
nivalis, *Phil. in Linnaea*, xxix. (1857–58) 69.—Chili.
nubigena, *Phil. l. c.* 67.—Chili.
Ochagsvii, *Annal, ex Hérincq, L'Hortic. Franç.* iv. (1854) 2.—Chili.
oculata, Lodd. Bot. Cab. t. 1851 = Bomarea Salsilla.
odorata, *Knight & Perry, ex Loud. Hort. Brit. Suppl.* III. 486.—Hab. ?
Oreas, *Y. C. Schau. in Nov. Act. Nat. Cur. xix. Suppl.* t. (1843) 440.—Am. austr.
ovata, Cav. Ic. i. 54. t. 76 = Bomarea edulis.
pallens, *Phil. in Linnaea*, xxxiii. (1864–65) 265.— Chili.
pallida, R. Grah. in Edinb. N. Phil. Journ. (Jul.–Oct. 1829) 344 = A. Ligtu.
parvula, *Phil. in Linnaea*, xxxiii. (1864–65) 261.— Chili.
pauciflora, Hort. ex Lem. in Fl. des Serres, Sér. I. ii. (Dec. 1846) t. 183 = Bomarea Jacquiniana.
pauciflora, *H. B. & K. Nov. Gen. et Sp.* i. 324.—N. Granat.
pauperula, *Phil. Fl. Atac.* 51.—Chili.
Pelegrina, *Linn. Diss. Alstroem.* 10 ; *Amoen. Acad.* vi. 247.—Peruv. ; Chili.
Pelegrina, Vell. Fl. Flum. 181 ; tii. t. 119 = caryophyllacea.
peregrina, Ruiz & Pav. Fl. Per. iii. 61 = A. Pelegrina.
peruviana, Van-Houtte, Fl. des Serres, xxi. (1876) 51. t. 2175 = versicolor.
plantaginea, *Mart. in Schult. f. Syst.* vii. 737.—Bras.
platyphylla, *Baker, in Journ. Bot.* xv. (1877) 261.— Bras.
Pohliana, Seub. in Mart. Fl. Bras. iii. t. 172 = A. Cunea.
Presliana, Herb. Amar. 95. t. 1. f. 4 = A. Ligtu.
Presliana, Kunth, Enum. Pl. v. 773 = præc.
psittacina, Lehm. Cat. Hort. Hamb. (1836) *ex* Schult. f. Syst. vii. 739 = pulchella.
pubernula, Phil. in Linnæa, xxix. (1857–58) 68 = versicolor.
pulchella, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 48 = versicolor.
pulchella, *Linn. f. Suppl.* 206.—Am. austr.
pulchella, Sims, Bot. Mag. t. 2364 = haemantha.
pulchella, Salisb. Prod. 344 = A. Ligtu.
pulchra, *Sims, Bot. Mag.* t. 2421 = A. Ligtu.
purpurea, Ruiz & Pav. Fl. Per. iii. 60 = Bomarea purpurea.

ALSTROEMERIA :—
purpurea, Willd. ex Steud. Nom. ed. II. i. 61, 62 = Bomarea Caldasiana.
pygmæa, *Herb. Amar.* 100, 397. t. 8. f. 4–13.— Peruv.
quillotensis, Herb. l. c. 97. t. 3. f. 2 = A. Pelegrina.
reclinata, *Poepp. & Endl. Gen. et Sp.* ii. 44. t. 160.— Chili.
recumbens, Herb. Amar. 97. t. 3. f. 2 = versicolor.
reflexa, *M. Roem. Syn. Enast.* 252.—Chili.
revoluta, *Ruiz & Pav. Fl. Per.* iii. 59.—Chili.
Riedelliana, *Andot, in Rev. Hortic. Sér.* II. iv. (1846–48) 207.— Bras.
rosea, Hook. Exot. Fl. t. 181 = A. Ligtu.
roseo, *Phil. Sert. Mend. Alt.* 48.—Chili.
roseo, *Ruiz & Pav. Fl. Per.* iii. 61.—Peruv.
rosmarinifolia, Benth. Pl. Hartw. 258 = Bomarea linifolia.
rubra, *Morel, in Rev. Hortic.* Sér. IV. ii (1853) 81. f. 5.—Bras.
Salsilla, Linn. Diss. Alstroem. 10 ; Amoen. Acad. vi. 247 = Bomarea Salsilla.
Salsilla, Vell. Fl. Flum. 181 ; iii. t. 120 = Bomarea edulis.
salsilloides, *Mart. in Schult. f. Syst.* vii. 748.— Bras.
scabrella, *Baker, in Journ. Bot.* xv. (1877) 261.— Bras.
secundifolia, Ruiz & Pav. Fl. Per. iii. 60 = Bomarea secundifolia.
Sellowiana, *Seub. in Mart. Fl. Bras.* iii. t. 173.—Bras.
sepium, Schott, ex Seub. l. c. 168 = Bomarea edulis.
sericeifolia, J. C. Schau. in Nov. Act. Nat. Cur. xix. Suppl. t. (1843) 441 = spathulata.
setacea, Ruiz & Pav. Fl. Per. iii. 62 = Bomarea tomentosa.
Simsii, Spreng. Syst. ii. 80 = haemantha.
Simsii, Sweet, Hort. Brit. ed. I. 408 = A. Ligtu.
spathulata, *Presl, Rel. Haenk.* ii. 122. t. 22. f. 2.— Peruv.
stenopetala, *Phil. in Anales Univ. Chil.* (1873) 547.— Chili.
stenopetala, *Seub. in Mart. Fl. Bras.* iii. t. 174.— Bras.
subrosulacea, *Herb. Amar.* 397. t. 5. f. 1. 2.—Chili.
tenuifolia, Herb. l. c. 94. t. 1. f. 47 = versicolor.
tigrina, Phil. in Linnaea, xxix. (1857–58) 68 = versicolor.
tigrina, Seub. in Mart. Fl. Bras. iii. t. 174 = brasiliensis.
tomentosa, Ruiz & Pav. Fl. Per. iii. 62 = Bomarea tomentosa.
torta, H. B. & K. Nov. Gen. et Sp. i. 323 = Bomarea torta.
tricolor, Hook. Exot. Fl. t. 65 = A. Ligtu.
triflora, Griff. Notul. iii. 340 = Ixiolirion montanum.
umbellata, Meyen, Reise, i. 356 = spathulata.
uniflora, Matth. ex Kunth, Enum. Pl. v. 762 = Bomarea glaucescens.
venustula, *Phil. in Linnaea*, xxxiii. (1864–65) 260.— Chili.
versicolor, *Ruiz & Pav. Fl. Per.* iii. 59.—Chili.
violacea, *Knight & Perry, ex Loud. Hort. Brit. Suppl.* iii. 486.—Hab. ?
violacea, *Phil. Fl. Atac.* 51.—Chili.
viridiflora, *Warm. in Kjoeb. Videnskb. Meddel.* (1872) 118.—Bras.
xanthina, Phil. in Anales Univ. Chil. (1873) 546 = versicolor.
xanioides, *Baker, in Journ. Bot.* xv. (1877) 262.— Bras.

ALTENSTEINIA, H. B. & K. Nov. Gen. et Sp. i. 333. t. 78. *fig.* 1615. *ORCHIDEÆ*, Benth. & Hook. f. Gen. 592.
Aa, Reichb. f. Xen. Orchid. i. 18 (1858).
Myrosmodes, Reichb. f. l. c. i. 19. t. 8 (1858).
argyrolepis, *Benth. & Hook. f. (in Kirk Kew.).*—Round. calcarata, *Reichb. f. l. c.* iii. 19.—Peruv.
cerea, *Reichb. f. Otia Bot. Hamb.* 51.—Peruv.
fimbriata, *H. B. & K. Nov. Gen. et Sp.* i. 333. t. 78.— Round.
gymnandra, *Reichb. f. Xenia Orch.* iii. 18.—Boliv.
inaequalis, *Reichb. l. c.* 19.—Boliv.
Maackeni, *Reichb. f. l. c.*—Peruv.
meniniana, *Reichb. f. l. c.* 20.—Peruv.
Matthewii, *Reichb. f. l. c.* 19.—Peruv.
nubigena, *Reichb. f. l. c.* 18 in note.—Hab. ?

ALTENSTEINIA:—
palacosa, *Kunth, Syn.* i. 326.—Ecuad.
paludosa, *Reichb. f. Xenia Orch.* iii. 19.—Bolliv.
pilifera, *H. B. & K. Nov. Gen. et Sp.* i. 333. t. 73.—Ecuad.
rostrata, *Reichb. f. Xenia Orch.* iii. 18.—Ecuad.
Scaptrum, *Reichb. f. l. c.* i. 18.—Ecuad.
virescens, *Lindl. in Ann. & Mag. Nat. Hist.* xv. (1845) 986.—N. Granad.
Weddelliana, *Reichb. f. Xenia Orch.* iii. 19.—Peruv.

ALTERNANTHERA, Forsk. Fl. Aegypt. Arab. 28 (1775). *AMARANTACEAE*, Benth. & Hook. f. iii 39.
ALLAGANTHERA, Mart. Hort. Erlang. 69 (1814).
ILLECEBRUM, Spreng. Anleit. ii. i. 317 (1817); Syst. i. 818 (1825).
LITHOPHILA, Sw. Prod. Veg. Ind. Occ. 14 (1788); Fl. Ind. Occ. i. 47. t. 1 (1797).
LITOPHILA, Sw. Prod. l. c.
PITYRANTHUS, Mart. in Denks. Acad. Muench. v. (1817) 179.
acaulis, *Anderss. Galap. Öarnes Veget.* 164.—Ins. Galap.
Achyranthа, *R. Br. Prod.* i. 417.—Am. trop.
Achyranthes, Forsk. Fl. Aegypt. Arab. p. lxix = sessilis.
albida, *Griseb. in Goett. Abh.* xix. (1874) (*Pl. Lorent.* 36.—Reg. Argent.
amabilis, Hort. ex Lem. Illust. Hortic. xv. (1868) t. 558 = Telanthera ficoidea.
angustifolia, Moq. in DC. Prod. xiii. *11.* 324 = polycephala.
angustifolia, *R. Br. Prod.* i. 417.—Austral.
argentata, *Moq. in DC. Prod.* xiii. *11.* 352.—Porto Rico
aurata, *Moq. l. c.* 358.—Am. austr.
axillaris, D. Dietr. Syst. i. 986 in syn. = spinosa.
Bauer, *Moq. in DC. Prod.* xiii. *11.* 354.—Austral.
brachiata, Hort. ex Moq. l. c. 304 = Aerva brachiata.
canescens, *H. B. & K. Nov. Gen. et Sp.* ii. 204.—Cumana.
canescens, Moq. in DC. Prod. xiii. *11.* 350 = Iresine canescens.
carcasana, *H. B. & K. Nov. Gen. et Sp.* ii. 205.—Venezuela.
caribaea, *Moq. in DC. Prod.* xiii. *11.* 354.—Ind. occ.
celosioides, *Moq. l. c.* 360.—Ins. Philipp.
ciliata, Poepp. ex Seub. in Mart. Fl. Bras. v. i. 183 = A. Achyrantha.
decipiens, *Benth. Fl. Austral.* v. 251.—Austral.
denticulata, R. Br. Prod. i. 417 = nodiflora.
denticulata, Wall. Cat. n. 6922 = sessilis.
dubia, Hort. Paris. ex Moq. in DC. Prod. xiii. *11.* 361 = sessilis!
dubia, H. B. & K. Nov. Gen. et Sp. ii. 209 = Trommsdorffia canescens.
echinata, Hort. Bonn. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 48 = arrecta.
echinata, Sm. in Rees, Cyclop. Suppl. n. 10 = A. Achyrantha.
erecta, Reichb. in DC. Prod. iii. 375 = Polycarpaea linearifolia.
ficoidea, R. Br. Prod. 417; Roem. & Schult. Syst. v. 555 = Telanthera ficoidea.
ficoidea, Gmelin. Fl. Brit. W. Ind. 67 = Telanthera maritima.
ficoidea, Beauv. Fl. Owar. ii. 66. f. 1 = denticulata.
ficoidea, Reichb. ex Moq. in DC. Prod. xiii. *11.* 357 = sessilis.
flavescens, H. B. & K. Nov. Gen. et Sp. ii. 207 = Telanthera flavescens.
flavescens, *Moq. in DC. Prod.* xiii. *11.* 350.—N. Granad.
frutescens, *R. Br. ex Spreng. Syst.* i. 819 in syn.—Am. austr.
glabra, Moq. in DC. Prod. xiii. *11.* 361 = denticulata?
gomphrenoides, *H. B. & K. Nov. Gen. et Sp.* ii. 208.—Peruv.
hemicircles, *Beurling, in Vet. Akad. Handl. Stockh.* (1764) 148.—Panama.
iresinoides, H. B. & K. Nov. Gen. et Sp. ii. 209 = Gomphrena frutescidea.
lanuginosa, Moq. in DC. Prod. xiii. *11.* 359 = Cladothrix lanuginosa.
lappulacea, Schlecht. ex Moq. l. c. 329 = Cyathula globulifera.
latifolia, Moq. l. c. 361 = Iresine latifolia.
lepidophylla, *Benth. Fl. Austral.* v. 251.—Austral.
leucantha, *Moq. in DC. Prod.* xiii. *11.* 357.—Jamaic.

ALTERNANTHERA:—
litteralis, Beauv. ex Moq. in DC. Prod. xiii. *11.* 365 = maritima.
linearis, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 107.—Porto Rico.
longipes, *Benth. Fl. Austral.* v. 259.—Austral.
lupulina, *H. B. & K. Nov. Gen. et Sp.* ii. 206.—Peruv.
maritima, *D. Dietr. Syn. Pl.* i. 867.—Bras.
maritima, St. Hil. Voy. Brés. ii. 43 = Telanthera maritima.
mexicana, *Moq. in DC. Prod.* xiii. *11.* 352.—Mexico ! muscoides. *Benth. & Hook. f. Gen.* iii. 39.—Ind. occ.
nana, F. Bauer, ex Moq. in DC. Prod. xiii. *11.* 341 = Iresine Brownii.
nana, *R. Br. Prod.* i. 417.—Austral.
nigricepe, Hook. Bot. Misc. ii. (1831) 287 = Telanthera elegans.
nigripes, Steud. Nom. ed. II. i. 65. (aphalm.) = praec.
nodiflora, *Griseb. in Goett. Abh.* xxiv. (1879) 86.—Reg. Argent.
nodiflora, *R. Br. Prod.* i. 417.—Geront. trop.
paniculata, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 108.—Porto Rico.
paniculata, *H. B. & K. Nov. Gen. et Sp.* ii. 208.—Cumana.
paronychioides, Hort. ex Regel, Gartenfl. (1869) 102 = Telanthera Bettzichiana.
paronychioides, *St. Hil. Voy. Brés.* ii. ii. 43.—Am. austr.
parviflora, *Moq. in DC. Prod.* xiii *11.* 360.—Mexic.!
philoxeroides, *Griseb. in Goett. Abh.* xxiv. (1879) 86.—Reg. Argent.
pilosa, *Moq. in DC. Prod.* xiii. *11.* 357.—Bras.
polycephala, *Benth. Fl. Austral.* v. 251.—Austral.
polygonoides, R. Br. Prod. 417 = Telanthera polygonoides.
praelonga, St. Hil. Voy. Brés. ii. ii. 409 = Telanthera praelongа.
procumbens, *Roem. & Schult. Syst.* v. 564.—Bras.
prostrata, D. Don, Prod. Fl. Nep. 76 = sessilis.
puberula, D. Dietr. Syn. Pl. i. 866 = Telanthera pubrula.
pubescens, Hort. Prag. ex Moq. in DC. Prod. xiii. *11.* 303 = Aerva lanata.
pulchella, *H. B. & K. Nov. Gen. et Sp.* ii. 205.—Bras.
pulverulenta, *Moq. in DC. Prod.* xiii. *11.* 351.—Peruv.
pungens, *H. B. & K. Nov Gen. et Sp.* ii. 206.—Bras.
purpurea, Hort. ex Pynaert, in Rev. Hortic. Belge, v. (1879) 25 = Telanthera versicolor.
radicata, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 262.—Ins. Galapag.
repens, Steud. Nom. ed. I. 84 = sessilis.
repens, Stead. l. c. ed. II. i. 65 = A. Achyrantha.
Richardi, Moq. in DC. Prod. xiii. *11.* 353 = Iresine interrupta.
rufa, D. Dietr. Syn. Pl. i. 866 = Telanthera rufa.
sericea, *H. B. & K. Nov. Gen. et Sp.* ii. 207.—Ecuad.
sessilis, Lem Illustr. Hortic. (1855) t. 447 = Telanthera amoena.
sessilis, *R. Br. Prod.* i. 417.—Reg. trop.
sibirica, *Steud. Nom* ed. II. i. 66.—Sibir.
spathulata, Lem. Illustr. Hortic. xii. (1865) t. 445 = Telanthera ficoidea.
spinosa, *Roem. & Schult. Syst.* v. 555.—Am. trop.
striguta, *Vanek in Hor. & De Vriese, Tydschr.* v. (1888–89) 259.—Malaya !
subcaposa, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 262.—Ins. Galap.
subfruticosa, *Terr. in Emory, Mex. Bound. Surv.* 181. —N. Mexico.
tenella, Colla, in Mem. Acc. Torin. xxxiii. (1829) 181. t. 8 = Telanthera polygonoides.
tenella, *Moq. in DC. Prod.* xiii. *11.* 355.—Hab.?
tomentella, *Seub. in Mart. Fl. Bras.* v. i. 186.—Bras.
trianoche, Steud. Nom. ed. I. 83, 84 = Telanthera ficoidea.
triandra, Lam. Encyc. i. 95 = sessilis.
truxillensis, *H. B. & K. Nov. Gen. et Sp.* ii. 206.—Peruv.
versicolor, Hort. ex Regel, Gartenfl. (1869) 101 = Telanthera versicolor.
villifora, *Schenk, in Linnaea*, xxii. (1849) 149.—Texas.
villosa, D. Dietr Syn. Pl. i. 867 = Brandesia villosa.
villosa, H. B. & K. Nov. Gen. et Sp. ii. 208 = Telanthera villosa.
villosa, *Sm. in Rees, Cycl. Suppl.* n. 9.—Hab.?
virgata, Spreng. ex Moq. in DC. Prod. xiii. *11.* 364 = Telanthera polygonoides.

ALTERNASEMINA, Silva Manso, Enum. 35 (1836) = Cayaponia, Mart. (Cucurb.).
Tayuia, Silva Manso, l. c. = C. Tayuya.

ALTHAEA, [Tourn.] Linn. Syst. ed. I (1735). *MALVACEAE*, Benth. & Hook. f. i. 200.
ALCEA, [Tourn.] Linn. Gen. ed. I. 306 (1737).
FERBERIA, Scop. Introd. 321 (1777).
acaulis, *Cav. Diss.* ii. t. 27. f. 3 (p. 93).—Syria.
aegyptiaca, Boiss. Diagn. Ser. II. i. 108 = lavateraefiora.
africana, *Linn. ex G. Don, Gen. Syst.* i. 467.—Afr. or.
ambigua, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 260 = Lavatera thuringiaca.
apterocarpa, *Fenzl, in Del. Sem. Hort. Vind.* (1858) 4.—Orient.
arborea, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 260 = Lavatera arborea.
armeniaca, *Tenore, Ind. Sem. Hort. Neap.* (1837) 1. —Oriens.
atropurpurea, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva atropurpurea.
Aucheri, *Boiss. Diagn.* Ser. I. vi. 28.—Persia; Afghan.
barbonica, *DC. Prod.* i. 438 (= Pavonia ?).—Ins. Borbon.
borealis, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva borealis.
Burchellii, DC. Prod. i. 438 = Pavonia Columella.
Calverti, *Boiss. Diagn.* Ser. II. v. 66.—Armenia.
cannabina, *Linn. Sp. Pl.* 686.—Europ.; Oriens.
caribaea, Sims, Bot. Mag. t. 1016 = rosea.
chinensis, Wall. Cat. n. 3039 = rosea.
coromandelina, Cav. Diss. ii 93 = rosea.
corymbosa, Sw. Prod. Veg. Ind. Occ. 101 = Pavonia paniculata.
cretica, Weinm. in Syll. Ratisb. ii. (1828) 171 = rosea.
crispa, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva crispa.
digitata, Boiss. Diagn. Ser. I. viii. 106 = lavateraefiora.
ficifolia, *Cav. Diss.* ii. 92. t. 26. f. 2.—Sibir.
ficifolia, Sibth. & Sm. Fl. Graec. i. 363 = rosea.
flavo-virens, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat Mosc.* xii. (1860) 44.—Sibir. baical.
flexuosa, Sims, Bot. Mag. t. 899 = rosea.
Frolovviana, *Fisch. ex G. Don, Gen. Syst.* i. 467.—Sibir.
frutex, Hort. ex Mill Gard. Dict. ed. VIII = Hibiscus syriacus.
garipensis, E. Mey. Zwei Pfl. Docum. 168 = Ludwigii.
garipina, E. Mey. ex Boiss. Fl. Orient. i. 824 = praec.
glabriflora, *Ledeb. ex Sweet, Hort. Brit.* ed. II. 60.—Sibir.
Godroni, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva ambigua.
grandiflora, Scop. Del. Insub. fi. 33 = Hibiscus roseus.
Heldreichii, *Boiss. Diagn.* Ser. II. i. 108.—Thessal.
hirsuta, *Linn. Sp. Pl.* 687.—Europ.; Oriens.
hirsuta, Sieber, ex Steud. Nom. ed. II. i. 66 = Malva crecta.
hispida, Moench, Meth. 612 = hirsuta.
Hohenackeri, *Boiss. & Huet, Diagn.* Ser. II. v. 67.—Oriens.
indica, Steud. Nom. ed. II. i. 66.—Java.
Irolowiana, Steud. l. c. sphalm. = Frolovviana.
hirrndenensis, Boiss. Diagn. Ser. II. v. 67 = lavateraefiora.
Kotschyi, Boiss. l. c. Ser. II. i. 103 = cannabina.
Kraussiana, Buching. ex Krauss, in Flora, xxviii. (1844) 298 = Pavonia Krausiana.
lasiica, *Schlecht. in Linnaea*, xvii. (1843) 127.—Oriens.
laevis, Moench, Meth. Suppl. 199 = Ludwigii.
lavateraefiora, *DC. Prod.* i. 437.—Orient.
lazistana, *A. DC. ex Schlecht. in Bot. Zeit.* vi. (1848) 341.—Hab.!
leucantha, Fisch. ex Loud. Hort. Brit. 287 = multiflora.
litticina, Boiss. & Kotschy, ex Boiss. Fl. Orient. i. 821 = apterocarpa.
longiflora, Boiss. & Reut. Diagn. Pl. Hispan. 9 = hirsuta.
Ludwigii, *Linn. Mant.* 98.—Oriens; Afr. bor. et austr.
marrotica, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva marrotica.
mauritiana, Alef. l. c. 260 = Malva mauritiana.
moenanthe, Link, in Linnaea, ix. (1834) 586 = rosea.
mollissima, *Kunze, l. c.* xx. (1847) 51, 404.—Mexic.
mucronifera, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva parviflora.
multiflora, Reichb. ex Regel, Ind. Sem. Hort. Petrop. (1867) 36 = armeniaca.

ALTHAEA :—
mutica, J. Gay, ex Ball, in Journ. Linn. Soc. xvi. (1878) 376 = hirsuta.
narbonensis, Jacq. Coll. ii. 275 = cannabina.
narbonnensis, *Pourr. ex Cav. Diss.* ii. 94. t. 29. f. 2.—Europ.
nicaeensis, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva nicaeensis.
nudiflora, *Lindl. in Trans. Hort. Soc.* vii. (1830) 251. —Sibir. altaic.
officinalis, *Linn. Sp. Pl.* ed. I. 686.—Europ.; Oriens.
pallida, *Wald. & Kit. Pl. Rar. Hung.* t. 47.—Hungar.; Turc.
parviflora, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva parviflora.
plebeia, Schult. ex Steud. Nom. ed. II. i. 66 = Lavatera plebeia.
pulchra, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 112. —Ind. or.
racemosa, Sw. Prod. Veg. Ind. Occ. 102 = Pavonia racemosa.
remotiflora, *Boiss. & Heldr. Diagn.* Ser. I. viii. 108. —As. Min.
rhyticarpa, Trautv. in Bull. Soc. Nat. Mosc. (1860) i. 454 = sulphurea.
rosea, *Cav. Diss.* ii. 91. t. 29. f. 3.—Orient.
rosea, Hohen. ex Boiss. Diagn. Ser. I. v. 67 = Hohenackeri.
rufescens, *Boiss. Diagn.* Ser. II. i. 102.—Syria.
setosa, *Boiss. l. c.* Ser. I. viii. 107.—Syria.
Sieberi, *Hort. ex G. Don, in Loud. Hort. Brit.* 486.—Sicilia.
sinensis, Blanco, Fl. Filip. ed. I. 552 = rosea.
sinensis, *Cav. Diss.* ii. t. 29. f. 3.—China.
speciosa, *Vis. Cat. Orto Bot. Padov.* 182.—Hab. ?
spicata, Spreng. Syst. iii. 108 = Hibiscus spicatus.
striata, *DC. Prod.* i. 437.—Arab.
sublobata, Stokes, Bot. Mat. Med. iii. 530 = officinalis.
sulphurea, *Boiss. & Hohen. Diagn.* Ser. I. viii. 108.—Persia; Afghan.; Soongar.
sylvestris, *Brignoli, ex Tenore, Syll.* 540.—Italia.
tabrisiana, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 44 = ficifolia.
taurinensis, C. A. Mey. Verz. Pfl. Cauc. 207 = armeniaca.
taurinensis, DC. Prod. i. 436 = officinalis.
thuringiaca, Kuntze, Taschen-Fl. Leipz. 212 = Lavatera thuringiaca.
Tournefortiana, *Tausch, in Flora*, xii. (1829) 9,—Creta.
ulmifolia, Pritzel, Ic. Ind. 50 = Sida glomerata.
verticillata, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva verticillata.
vulgaris, Alef. l. c. 260 = Malva vulgaris.

ALTHEA, Crantz, Stirp. Austr. 144 (1769) = praec.

ALTHENIA, Petit, in Ann. Sc. Observ. i. (1829) 451. *NAIADACEAE*, Benth. & Hook. f. iii. 916.
Barrandouii, Duval-Jouve, in Bull. Soc. Bot. Fr. xix. (1873) p. lxxxii; Compt.-rend. lxxv. (1872) 95 = filiformis.
filiformis, *Petit, in Ann. Sc. Observ.* i. (1829) 451.—Gallia.
setacea, Delile, ex Petit, l. c. = filiformis.

ALTHERIA, Thon. Gen. Nov. Madag. 19 (1809) = **Melochia**, Linn. (Sterculi.).
madagascariensis, DC. Prod. i. 493.—Madag.

ALTHINGIA, Steud. Nom. ed. II. i. 66 (1840) = seq.

ALTINGIA, [G. Don, in] Loud. Hort. Brit. 408 (1830) = **Arauoria**, Juss. (Conif.).
Cunninghamii, [G. Don,] l. c. = Ar. Cunninghamii.
excelsa, [G. Don,] l. c. = Ar. excelsa.

ALTINGIA, Noronha, in Verh. Bat. Gen. v. (1785) Art. ii. 9 ; ed. II. 41. *HAMAMELIDEAE*, Benth. & Hook. f. i. 669.
SEDGWICKIA, Griff. in As. Res. xix. (1836) 98. t. 15.
chinensis, Oliver, ex Hance, in Journ. Linn. Soc. xiii. (1873) 103.—China.
excelsa, Noronha, *in Verh. Bat. Gen.* v. ed. II. 41 (1827).—Malaya.

ALTORA, Adans. Fam. ii. 356 (1763) = **Cluytia**, Linn. (Euphorb.).

ALUGHAS, Linn. Fl. Zeyl. 207 (1747) = **Alpinia**, Linn. (Scitam.).

ALVARADOA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 100. *SAPINDACEAE*, Benth. & Hook. f. i. 411.
amorphoides, *Liebm. l. c.* 101.—Mexic.
arborescens, *Griseb. Cat. Pl. Cub.* 50.—Cuba.
jamaicensis, *Benth. Pl. Hartw.* 344.—Jamaica.
mexicana, *Liebm. ex Benth. l. c.*—Mexic.

ALVARDIA, Fenzl, in Flora, xxvii. (1844) 312 = **Pseudocadanum**, Linn. (Umbellif.).
arborescens, Fenzl, l. c. = P. fraxinifolium.

ALVAREZIA, Pav. ex Nees, in DC. Prod. xi. 465 (1847) = **Riechum**, R. Br. (Acanth.).
distichа, Pav. ex Nees, l. c. 246 = Lophostachys Alvarezia.
parviflora, Pav. ex Nees, l. c. 466 = B. Linnaei.

ALVESIA, Welw. Apont. 587. n. 47 (1858) = **Bauhinia**, Linn. (Legum.).
bauhinioides, Welw. l. c. = B. tomentosa.

ALVESIA, Welw. in Trans. Linn. Soc. xxvii. (1869) 55. t. 19. *LABIATAE*, Benth. & Hook. f. ii. 1176.
rosmarinifolia, *Welw. l. c.*—Afr. trop.

ALVISIA, Lindl. Fol. Orch. (1859) = **Eria**, Lindl. (Orchid.).
tenuis, Lindl. l. c. = E. articulata.

ALYCIA, Willd. ex Steud. Nom. ed. II. i. 66 (1840) = **Eriochlon**, H. B. & K. (Gramin.).
coarctata, Willd. ex Steud. l. c. = E. polystachya.
distachya, Willd. ex Steud. l. c. = E. polystachya.

ALYMERIA, D. Dietr. Syn. i. in Ind. p. xiii (1839) = **Alymeria**, Mart. = **Polycarpaea**, Lam. (Caryoph.).

ALYMNIA, Neck. Elem. i. 81 (1790) = **Polymnia**, Linn. (Compos.).

ALYOGYNE, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 12 = **Fugosia**, Juss. (Malvac.).
hakeifolia, Alef. l. c. = F. hakeaefolia.

ALYPUM, Fisch. Hort. Gorenk. ed. II. (1812) 19 = **Globularia**, Tourn. (Selagin.).
monspeliensium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 143.—G. Alypum.
salicifolium, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 19 = G. Alypum.
Solandri, Hort. ex Steud. Nom. ed. I. 373 = G. Alypum.

ALYSICARPUS, Neck. Elem. iii. 15 (1790) ; Desv. Journ. Bot i. (1813) 120. t. 4. *LEGUMINOSAE*, Benth. & Hook. f. i. 522.
FABRICIA, Scop. Introd. 307 (1777).
HALILIA, J. St. Hil. in Bull. Soc. Philom. (Dec. 1818) 192.
HEGETSCHWEILERA, Heer & Regel, Cat. Sem. Hort. Turic. (1849) ; ex Bot. Zeit. i. (1843) 47.
MYSICARPUS, Webb, in Hook. Niger Fl. 122 (1849) lapsu.
belgaumensis, *Wight, ic. t.* 92.—Ind. or.
bupleurifolius, Wall. Cat. n. 5761 = rugosus.
bupleurifolius, *DC. Prod.* ii. 352.—Ind. or.
cylindriceus, Desv. in Ann. Linn. Soc. Par. iv. (1826) 301 = vaginalis.
diversifolius, Wall. Cat. n. 5773 = vaginalis.
ferrugineus, *Hochst. & Steud. ex A. Rich Tent. Fl. Abyss.* i. 207.—Abyss.
glaber, E. Mey. Comm. Pl. Afr. Austr. 125 = rugosus.
glumaceus, *DC. Prod.* ii. 353.—Arab.
glumaceus, Wall. Cat. n. 5764 = rugosus.
gracilis, Edgew. in Journ. As. Soc. Beng. xxi. (1852) 170 = bupleurifolius.
gramineus, *Zoll. & Mor. in Nat. en Genees. Arch. Neerl. Ind.* iii. (1846) 89, nomen.—Malaya.
hamosus, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 171.—Ind. or.
Harnieri, Scheinf. Reliq. Kotschy, 34. t. 19 = vaginalis.
heterophyllus, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 158 = vaginalis.

ALYSICARPUS :—
Heyneanus, Wight & Arn. Prod. i. 234 = rugosus.
Hochstetteri, A. Rich. Tent. Fl. Abyss. i. 209 = rugosus.
longifolius, *Wight & Arn. Prod.* i. 233.—Ind. or.
ludens, Wall. Cat. n. 5762 = bupleurifolius, rugosus.
lupulinus, Stocks, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 159 = rugosus.
monilifer, *DC. Prod.* ii. 353.—Ind. or. ; Afr. trop.
nummularifolius, DC. l. c. = vaginalis.
nummularioides, Zipp. ex Span. in Linnaea, xv. (1841) 194 = scaber.
nummularius, Ball. in Bull. Soc. Linn. Par. i. (1884) 482 = vaginalis.
obovatus, Edgew. in Journ. As. Soc. Beng. xxi. (1852) 169 = rugosus.
paniculatus, *Bove. ex Baill. in Bull. Soc. Linn. Par.* i. (1884) 482.—Madag.
parviflorus, Dalz. in Hook. Kew Journ. iii. (1851) 211 = Desmodium parviflorum.
pilifera, Wall. Cat. n. 5766 = rugosus.
porrectus, *Welw ex Baker, in Oliver, Fl. Trop. Afr.* ii. 171.—Afr. trop.
pubescens, *Law, in Wight, Ic.* i. t. 250.—Ind. or.
pyilfer, Steud. Nom. ed. II. i. 66 = rugosus.
quadrangularis, Edgew. ex Baker, in Hook. f Fl. Brit. Ind. ii. 159 = tetragonolobus.
Quartinianus, A. Rich. Tent. Fl. Abyss. i. 208 = rugosus.
racemosus, Benth. in Linnaea, xxiv. (1851) 642 = belgaumensis.
rotundifolius, Dalz. ex Hook. f. Fl. Brit. Ind. ii. 172 = Desmodium rotundifolium.
rotundifolius, Wight, ex Hook. f. l. c. 157 = hamosus.
Rukharna, Buch.-Ham. ex Wall. Cat. n. 5771 = vaginalis, monilifera.
rugosus, *DC. Prod.* ii. 353.—Reg. trop.
rupicola, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 170.—Ind. or.
scaber, *Span. in Linnaea*, xv. (1841) 194.—Malaya.
scarious, K. Grah. in Wall. Cat. n. 5768 = rugosus.
spicatus, Royle, ex Hook. f. Fl. Brit. Ind. ii. 159 = longifolius.
styracifolius, DC. Prod. ii. 353 = rugosus.
tetragonolobus, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 169.—Ind. or.
timorensis, *Span. in Linnaea*, xv. (1841) 194.—Ins. Timor.
trifoliatus, Stocks, ex Hook. f. Fl. Brit. Ind. ii. 158 = vaginalis.
vaginalis, *DC. Prod.* ii. 353.—Ind. or.
vaginalis, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 170 = monilifer.
vaginalis, Wall. Cat. n. 5763 = longifolius, rugosus.
varius, Wall. l. c. n. 5768 = vaginalis.
Wallichii, Wight & Arn. Prod. i. 234 = rugosus.
Zeyheri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 280.—Afr. austr.

ALYSSOIDES, Tourn. ex Adans. Fam. ii. 419 (1763) = **Vesicaria**, Lam. (Crucif.).
creticum, Medic. Phil. Bot. i. 189 = Alyssum creticum.
gemonense, Medic. i. c. = Alyssum gemonense.
leucofolium, Medic. Pflanzeng. 62 = V. graeca.
sinuatum, Medic. Phil. Bot. i. 189 = Alyssum sinuatum.
tomentosum, Moench, Meth. 264 = Alyssum creticum.
utriculatum, Medic. Phil. Bot. 189 = V. graeca.

ALYSSOPSIS, Boiss. Diagn. Ser. I. vi. 14 (1842). *CRUCIFERAE*, Benth. & Hook. f. i. 74.
deflexa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 58.—Persia.
Drummondii, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 291 = Sisymbrium brachypodum.
Kotschyi, *Boiss. Diagn.* Ser. I. vi. 14.—Persia.
sagittata, Boiss. Fl. Orient. i. 182 = deflexum.

ALYSSOPSIS, Reichb. Nom. 181. n. 6961 (1841) = **Vesicaria**, Lam. (Crucif.).

ALYSSON, Crantz, Stirp. Austr. ed. II. 15 (1769) = seq.

ALYSSUM, Tourn. ex Linn. Syst. ed. I. (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 78.
ADENETA, Adans. Fam. ii. 420 (1763).
ANODONTEA, G. Don, Gen. Syst. i. 180 (1831).
AURINIA, Desv. Journ. Bot. iii. (1814) 163.
BERTEROA, DC. Syst. ii. 290 (1821).

ALYSSUM:—
Gluck, Lindl. Syn. Brit. Fl. 26 (1826).
Konig, Adans. Fam. ii. 420 (1763).
Konig, A. R. in Denh. Clapp. & Oud. Voy. App. 214 (1826).
Lobularia, Desv. Journ. Bot. iii. (1814) 162.
Meniocus, Desv. l. c. 173.
Morchia, Roth, Tent. Fl. Germ. i. 278 (1788).
Octadenia, R. Br. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 41 (1836).
Odontarrhena, C. A. Mey. in Ledeb. Fl. Alt. iii. 58 (1831).
Onodontea, G. Don, Gen. Syst. i. 148 (1831), sphalm.
Ptilonema, C. A. Mey. in Ledeb. Fl. Alt. iii. 50 (1831).
Ptilotrichum, C. A. Mey. l. c. 64 (1831).
Schwerckia, Andrz. ex DC. Syst. ii. 300 (1821).
Stevena, Andrz. ex DC. Syst. ii. 291 (1821).
affine, Tenore, Syll. 315 = orientale.
alcides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 158.—Armenia.
albescens, Schlecht. ex DC. Syst. ii. 309, et Prod. i. 162 = montanum.
algeriense, *Pomel, Nouv. Mat. Fl. Atl.* 232.—Afr. bor.
alpestre, Bieb. Fl. Taur. Cauc. ii. 102, Suppl. 482 = Marschallianum.
alpestre, *Linn. Mant.* 92.—Europ.
alpestre, Wulf. ex Nym. Consp. 56 = montanum.
alpestre, Salis, in Flora, xvii. (1877) ii. Beibl. 77 = Robertianum.
alpestre, Sibth. & Sm. Fl. Græc. t. 628 = argenteum.
alpinum, Scop. Fl. Carn ed II. ii. 9 = montanum.
alsinefolium, Host, Fl. Austr. ii. 248 = calycinum.
altaicum, C. A. Mey. in Ledeb. Fl. Alt. iii. 55 = Fischerianum.
alyssoides, Linn. Syst. ed. X. 1130 = calycinum.
andinum, *Rupr. Fl. Caucas.* 103.—Reg. Caucas.
arl(ch)ense, Wormsk. ex Vahl, in Fl. Dan. t. 1530 = Vesicaria arctica.
arvense, Lobel, Fl. Gallica, ed. I. 401 = montanum.
arvense, *Pers*., *Fl. Sic.* i. 56.—Sicilia.
argenteum, *Vitm. Summ.* iv. 30.—Europ.; Orient.
argyrophyllum, *Schott & Kotschy, in Oestr. Bot. Wochenbl.* vii. (1857) 229.—Asia Min.
armenum, *Boiss. Fl. Orient.* i. 378.—Armenia.
aruaticum, Jord. Diag. i. 200 = calycinum.
atlanticum, Desf. Fl. Atlant. ii. 71. t. 149 = montanum.
Auckeri, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 156 = foliosum.
aurantiacum, *Boiss. Fl. Orient.* i. 376.—Lycia.
auresum, *Boiss. l. c.* 386.—Orient.
Bertholonii, Desv. Journ. Bot. iii. (1814) 173 & 185 = argenteum.
Besseri, *Stev. ex DC. Syst.* ii. 300 = podolicum.
brugesiacum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 10 = montanum.
bidentatum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 108 = Draba brachycarpa.
Billmekii, *Willd. in Willd. & Lange, Prod.* iii. 830.—Hispan.
brachystachyum, Bieb. Fl. Taur. Cauc. iii. 484 = umbellatum.
bracteatum, Boiss. Diagn. Ser. I. xiii. 77 sphalm. = Acanthophyllum bracteatum.
bracteatum, *Boiss. & Buhse, in Nouv. Mem. Soc. Nat. Mosc.* xii. (1860) 18.—Persia; Armenia.
bramifolium, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 11 = montanum.
brigantiacum, Jord. & Fourr. l. c. 12 = montanum.
Bungei, *Boiss. Fl. Orient.* i. 374.—Persia.
calicinum, Neck. Delic. Gallo-Belg. i. 277 = calycinum.
calliebroeum, Boiss. Diagn. Ser. II. v. 84 = constellatum.
calycinum, Hasenel. ex Nym. Consp. 56 = campestre.
calycinum, *Lam. & Dc. Fl. fr.* ed. II. 808.—Europ.; Orient.
calycinum, Pall. ex DC. Syst. ii. 316 = dasycarpum.
calycocarpum, *Rupr. Fl. Cauc.* 103, in textu.—Tauria.
campestre, Hoffm. Fl. Deutschl. ed. II. ii. 43 = calycinum.
campestre, *Linn. Syst.* ed. X. 1130.—Europ.; Orient.
campestre, Marz. ex Poll. Fl. Ver. ii. 862 = montanum.
camariense, Del. ex Webb & Berth. Phyt. Canar. i. 91 = sphæroc.
camdidum, Schlecht. ex DC. Syst. ii. 308 = Marschallianum.
canescens, *DC. Syst.* ii. 323.—Sibir.; Reg. Himal.

ALYSSUM:—
canium, *Boiss. & Heldr. Diagn.* Ser. I. viii. 34.—Syria.
castellanum, Bourg. ex Nym. Consp. 57 = serpyllifolium.
cedrorum, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 169 = hirsutum.
Cephalotes, *Boiss. Diagn.* Ser. II. i. 84.—Caria.
chalcidicum, *Yanka, in Oestr. Bot. Zeitschr.* xxii. (1872) 176.—Turcia.
cheiranthifolium, Willd. Sp. Pl. iii. 468 = Farsetia clypeata.
cheirifolium, Berger. ex Steud. Nom. ed. I. 34 = saxatile.
chrysanthum, Boiss. Fl. Orient. i. 271 = argenteum.
ciliatum, Lam. Fl. Fr. ii. 479 = Draba aizoides.
ciliicum, *Boiss. & Bal. Diagn.* Ser. II. v. 84.—Cilicia.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 236.—Afr. bor.
clypeatum, Linn. Sp. Pl. ed. I. 651 = Farsetia clypeata.
cochlearioides, Roth, Nov. Pl. Sp. 322 = Cochlearia flava.
cochleatum, *Cos. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 11.—Afr. bor.
collinum, Brot. Phyt. Lusit. ii. 209. t. 180 = campestre.
collinum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 9 = montanum.
compactum, *De Not. ex Ces. Ic. Stirp. Ital.* fasc. iii. 15. t. 5.—Sicilia.
condensatum, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* i. 268.—Ins. Min.; Syria.
confertum, Hort. Berol. ex Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 155 = strictum.
constellatum, *Boiss. l. c.* Sér. IV. ii. (1854) 244.—Cilicia.
contemptum, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 177 = Szowitzianum.
cordatum, Spreng. ex Steud. Nom. ed. II. i. 67 = obcordatum.
cornutum, Steph. ex Ledeb. Fl. Ross. i. 139 = rostratum.
corsicum, *Duby, Bot. Gall.* ed. II. i. 84.—Corsica.
corsicum, Rob. ex Gren. & Godr. Fl. Fr. i. 117 = Robertianum.
corymbosum, *Boiss. Fl. Orient.* i. 265.—Turc.
corymbulosum, Boiss. l. c. 282 = umbellatum.
crenulatum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 33.—Syria.
cretaceum, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 109.—Sibir.
creticum, *Linn. Sp. Pl.* 661.—Creta.
cryptopetalum, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 142; *Lehm. Rel.* (1847) 142 = marginatum.
cuneifolium, Tenore, Fl. Nap. Prod. xxxvii. App.; Cat Neap. (1815) 58 = montanum.
dagestanicum, *Rupr. Fl. Caucas.* 104.—Caucas.
dahuricum, Schlecht. ex DC. Syst. ii. 715 = canescens.
damascenum, *Boiss. & Gaill. Diagn.* Ser. II. vi. 18.—Syria.
dasycarpum, *Steph. in Willd. Sp. Pl.* iii. 469.—Orient; Turkestan. Afghan.
dauricum, Schlecht. in DC. Syst. ii. 322, & Prod. i. 164 = canescens.
dawuricum, Willd. ex Ledeb. Fl. Ross. i. 138 = Fischerianum.
decoloratum, *Pomel, Nouv. Mat. Fl. Atl.* 236.—Afr. bor.
decumbens, Herbich, Stirp. Rar. Bucov. 47 = repens.
deltoideum, Linn. Sp. Pl. ed. II. 908 = Aubrietia deltoidea.
densiflorum, Desf. Cor. Tourn. 65. t. 48 = strictum.
dentatum, Nutt. Gen. Am. ii. 63 = Draba ramosissima.
denticulum, Willd. Enum. Hort. Berol. 671 = Camelina sativa.
dicranoides, Boiss. & Huet, Diagn. Ser. II. v. 80 = Draba rigida.
diffusum, Jord. & Fourr. ex Nym. Consp. 57 = argenteum.
diffusum, Tenore, App. Hort. Neap. (1815) 58 = montanum.
Draba, Bieb. ex Willd. Enum. Hort. Berol. 672 = linifolium.
Draba, Hort. ex Webb & Berth. Phyt. Canar. i. 91 = lybicum.
Draba, Willd. Enum. Hort. Berol. 672 = hirsutum.
dubium, Bory, in Ann. Gen. Soc. Phys. iii. (1820) 4 = Lagascæ.
edentulum, Waldst. & Kit. Pl. Rar. Hung. i. 95. t. 92 = gemonense.

ALYSSUM:—
elatum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 33.—As. Min.
emarginatum, *Zahl. ex Vis. Fl. Dalm.* iii. 117.—Dalmat.
erigena, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 9 = montanum.
eriocarpum, Pourr. ex Steud. Nom. ed. II. i. 67 = Clypeola eriocarpa.
eriophorum, Willd. Enum. Hort. Berol. 671 = Clypeola eriophora.
eriophyllum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 373.—As. Min.
erosulum, Genn. & Pestal. ex Clem. in Mem. Acc. Sc. Toria. Ser. II. xvi. (1857) 246 = saffruticosa.
eruaticum, Jord. Diag. i. 200 = calycinum.
filifolium, *Wahlenb. in Isis*, xxi. 998.—As. Min.
Fischeri, *Jord. Obs. Pl. Crit.* i. (1846) 10 = montanum.
floribundum, *Boiss. Diagn.* Ser. II. v. 83.—Cilicia.
foliosum, *Bory & Chaub. Nouv. Fl. Pélop.* 42 n. 1080.—Græcia; As. Min.
fruticulosum, Steud. Nom. ed. I. 34 = Vesicaria fruticulosa.
fulvescens, *Sibth. & Sm. Prod. Fl. Græc.* ii. 13.—Græcia; As. Min.
gemonense, *Linn. Mant.* 92.—Europ.
gemonense, Roth, Tent. Fl. Germ. iii. 85 (err. glemonense) = saxatile.
globosum, Steud. Nom. ed. I. 34 = Vesicaria globosa.
glomeratum, *Burch. ex DC. Syst.* ii. 316.—Afr. austr.
Gmelini, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 8 = montanum.
graphalodes, DC. Syst. ii. 311 = Vesicaria graphalodes.
gracile, *Reyn. in Am. Monthly Mag.* (1811) 356, nomen.—Hab.!
granatense, Boiss. & Reut. Pugill. Pl. Nov. (1852) 9 = calycinum.
granatense, Nym. Consp. 57 = serpyllifolium.
halimifolium, DC. Fl. Fr. v. 594 = pyrenaicum.
halimifolium, Lap. Hist. Abr. Pl. Pyr. 371 = macrocarpum.
halimifolium, Linn. Sp. Pl. ed. I. 650 = maritimum.
halimifolium, Pall. Reise, iii. 161 = lenense.
halimifolium, *Willd. Sp. Pl.* iii. 460.—Europ.
Helderi, *Kruch. Fl. Sil. Suppl.* ii. 168.—Europ.
Hausknechtii, *Boiss. Fl. Orient.* i. 269.—As. Min.
Heinzi, *Ullep. in Oestr. Bot. Zeitschr.* xxxv. (1885) 307.—Europ.
helianthemifolium, *Timb. & Yeanb. in Rev. Bot. Soc. Fr.* iii. (1884–85) 105.—Gallia.
heterophyllum, Reis & Pav. ex DC. Syst. ii. 342 = Lepidium perfoliatum.
heterotrichum, *Boiss. Diagn.* Ser. I. vi. 15.—Persia.
hirsutum, Bess. ex Nym. Consp. 56 = rostratum.
hirsutum, *Bieb. Fl. Taur. Cauc.* ii. 106.—Orient.
hirsutum, Hoben. Enum. Ellasebathpol. 248 = campestre.
homalocarpum, *Boiss. Fl. Orient.* i. 285.—Arabia.
horebicum, Boiss. in Ann. Sc. Nat. Ser. II. xvii. (1842) 156 = homalocarpum.
Huetii, *Boiss. Fl. Orient.* i. 387.—As. Min.
hyperboreum, *Linn. Sp. Pl.* 651.—Am. bor.
Idaeum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 35.—Creta.
illyricum, Willd. Enum. Suppl. 44 = linifolium.
imbricatum, Schlecht. ex DC. Syst. ii. 308, & Prod. i. 161 = canescens.
incanum, *Linn. Sp. Pl.* 650.—Europ.; Sibir.
Jonthlaspi, [Clairv.] Man. Herb. 218 = Clypeola Jonthlaspi.
Kotschyanum, Boiss. Fl. Orient. i. 270 = mesamphanum.
Lagascæ, *Yanka, in Termész. Füzetek*, vii. (1883) 114.—Europ.
lanigerum, *DC. Syst.* ii. 308.—Persia.
lanuginosum, Pourr. ex DC. Syst. ii. 327 = Clypeola eriophora.
Lapeyrousianum, *Jord. Obs. Pl. Crit.* i. (1846) 8 = Mont. Pyren.
latifolium, *Vis. Fl. Dalm.* iii. 118.—Dalmat.
leiocarpum, *Pomel, Nouv. Mat. Fl. Atl.* 235.—Afr. bor.
lenense, *Adams, in Mem. Soc. Nat. Mosc.* v. (1817) 110.—Sibir.
lepidotum, *Boiss. Diagn.* Ser. I. i. 78.—As. Min.
Lescuri, *A. Gray, Man. Bot. U. St.* ed. V. 72.—Am. bor.

ALYSSUM :—
lucasdeum, *Guss. Pl. Rar.* 268.—Ital.
libyrum, *Cass. in Bull. Soc. Bot. Fr.* iv. (1857) 467.—Afr. bor.
linearifolium, Lag. Gen. et Sp. Nov. 18 = linifolium.
linifolium, *Steph. ex Willd. Sp. Pl.* iii. 467.—Europ.; Oriens.
litorale, *Rupr. Fl. Cauc.* 103. in textu.—Littora Maris Balt.
longicaule, *Boiss. in Bibl. Genèv.* (1888) 407.—Hispan.
ludovicianum, Nutt. Gen. Am. ii. 63 = Vesicaria ludoviciana.
lunarioides, Spreng ex DC. Syst. ii. 292 = mutabile.
lunarioides, Willd. Sp. Pl. iii. 461 = Farsetia lunarioides.
lusitanicum, Brot. ex Nym. Consp. 56 = campestre.
luteolum, *Pomel, Nouv. Mat. Fl. Atl.* 334.—Afr. bor.
macrocalyx, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 12.—Afr. bor.
macrocarpum, *DC. Syst.* ii. 321.—Gall. austr.
macropodum, *Boiss. & Bal. Diagn.* Ser. II. vi. 18.—Cappadoc.
macrostylum, *Boiss. & Huet, ex Boiss. Fl. Orient.* i. 279.—As. Min.
magnatum, *Steud. Nom.* ed. II. i. 68 (*ex Boiss. in Ann. Sc. Nat.* xvii. (1842) 157).—Arab.; Persia; Afghan.
margitatum, *Timb. & Jeanb. in Rev. Bot. Soc. Fr.* iii. (1884–85) 106.—Gallia.
maritimum, *Lam. Encyc.* i. 98.—Europ.
Marschallianum, *Andrz. ex DC. Syst.* ii. 308.—Caucas.
masmenense, *Boiss. Diagn.* Ser. II. v. 86.—Cappadoc.
medium, Host, Fl. Austr. ii. 244 = gemonense.
menlocoides. *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 158.—Mesopot.
mesopotamicum, Fenzl, in Tchichat. Asie Min. Bot. i. 311 = argenteum.
Meyerianum, Rupr. Fl. Cauc. 100, in textu = tortuosum.
micranthum, Fisch. & Mey. in Linnæa, x. (1836) 83 = campestre.
microcarpon, Vis. Fl. Dalmat. iii. 115 = corymbosum.
microcarpum, Nellr. ex Nym. Consp. 55 = gemonense.
microcephalum, Boiss. Diagn. Ser. I. xiii. 77, sphalm. = Acanthophyllum microcephalum.
micropetalum, Fisch. in DC. Syst. ii. 318 = campestre.
micropetalum, *Kit. in Linnæa*, xxxii. (1863) 497.—Europ.
microphyllum, *Steud. Nom.* ed. II. i. 68.—Sibir. alt.
minimum, Linn. Sp. Pl. 651 = maritimum.
minimum, Pall. Reise, ii. 741 = linifolium.
minimum, *Willd. Sp. Pl.* iii. 464.—Europ.; Oriens; Reg. Himal.
minutiflorum, Boiss. Diagn. Ser. I. i. 73 = alpestre.
minutulum, Schleich. ex DC. Syst. ii. 307 = alpestre.
minutum, Patr. ex DC. l. c. 322 = canescens.
minutum, *Schlechl. ex DC. l. c.* 316.—Sibir.
modestum, Boiss. Diagn. Ser. II. vi. 17 = minutum.
montanum, Brot. Fl. Lusit. ii. 558 = campestre.
montanum, *Linn. Sp. Pl.* 650.—Europ.; Oriens.
montanum, Pall. Reise, i. 353 = Fischerianum.
montanum, Patr. ex DC. Syst. ii. 322 = canescens.
mouradicum, *Boiss. & Bal. Diagn.* Ser. II. vi. 16.—As. Min.
Muelleri, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 16.—Armenia; Persia.
murale, Bieb. Fl. Taur. Cauc. ii. 108 = alpestre.
murale, Waldst. & Kit. Pl. Rar. Hung. i. 5. t. 6 = argenteum.
murcicum, Jord. ex Nym. Consp. 57 = serpyllifolium.
muricatum, *Kit. in Linnæa*, xxxii. (1863) 498.—Europ.
mutabile, *Vent. Hard. Cels*, 55.—Ital.
myagroides, Afl. Fl. Fedem. i. 241 = Cæsalpinia saxatilis.
Myagrum, Wibel, Prim. Fl. Werth. 204 = Camelina sativa.
nanum, *Pomel, Nouv. Mat. Fl. Atl.* 336.—Afr. bor.
nebrodense, Tineo, Pugill. Pl. Sic. 13 = serpyllifolium.
novum, Kit. Ind. Pesth, t. 6 = alpestre.
numidicum, *Pomel, Nouv. Mat. Fl. Atl.* 337.—Afr. bor.
obcordatum, *DC. Syst.* ii. 324.—Hab.?
obliquum, Sibth. & Sm. Fl. Græc. t. 638 = mutabile.
obovatum, Turcz. in Bull. Soc. Nat. Mosc. (1837) i. 57 = sibiricum.
obtusifolium, Stev. ex DC. Syst. ii. 305 = alpestre.

ALYSSUM :—
ochroleucum, Boiss. Diagn. Ser. II. v. 36 = montanum.
Oederi, Durand, Fl. Bourg. i. 161 = Vesicaria utriculata.
orbiculare, *Timb. & Jeanb. in Rev. Bot. Soc. Fr.* iii. (1884–85) 106.—Europ.
orbiculatum, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 385.—Græcia.
orientale, *Ard. Animad. Alt.* 32.—Græcia; As. Min.
oropilum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 11 = montanum.
ovirense, *A. Kern. Sched. Fl. Exsicc.* ii. 99.—Austria.
oxyacarpum, *Boiss. & Bal. Diagn.* Ser. II. v. 85.—Cappadoc.
paniculatum, Desf. Choix, 67 = Vesicaria paniculata.
paniculatum, Willd. Enum. Hort. Berol. 671 = Neslia paniculata.
parviflorum, Bieb. Fl. Taur. Cauc. iii. 434 = campestre.
parviflorum, Hochst. & Steud. ex Steud. Nom. ed. II. i. 68 = marginatum.
parviflorum, Schloss. ex Nym. Consp. 57 = calycinum.
parvulum, *Schrad. in Linnæa*, x. (1836) *Lit.* 69.—Hab. ?
patulum, *Pomel, Nouv. Mat. Fl. Atl.* 338.—Afr. bor.
pedemontanum, *Rupr. Fl. Cauc.* 102, *in textu*.—Europ.
peltaroides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 158.—As. Min.
persicum, *Boiss. l. c.* 162.—Persia.
Peruzianum, J. Gay, ex Grem. & Godr. Fl. Fr. i. 118 = Lapeyrousianum.
petalodes, DC. Syst. ii. 312 = Farsetia obovata.
petraeum, Andrz. ex DC. l. c. 309 = saxatile.
petraeum, Ard. Animad. Alt. 30 = gemonense.
petrophilum, Heldr. ex Nym. Consp. 56 = montanum.
Peyrousianum, Willk. & Lange, Prod. Fl. Hisp. iii. 835 = Lapeyrousianum.
phymatocarpum, Schlosser & Vukot. Syll. 186 = calycinum.
plocospermum, Fenzl, in Russeg. Reise, Illustr. 54 = aureum.
podolicum, *Bess. Cat. Hort. Crem.* (1816) 8.—Europ.; As. Min.; Sibir.
polyodon, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 17 = campestre.
porphyreticum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 11 = montanum.
prascox, *Boiss. Fl. Orient.* i. 275.—As. Min.
psammeum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 9 = montanum.
psilocarpum, Boiss. Voy. Bot. Espagne, App. ii. 718 = calycinum.
pyrenaicum, *Lapeyr. Hist. Abr. Pl. Pyr.* 371. *Suppl.* 91.—Mont. Pyren.
ramosissimum, *Web. in Giorn. Bot. Ital.* ii. (1846) 215.—Græcia.
repens, Baumg. Enum. Transs. ii. 237 = montanum.
rhodanense, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 10 = montanum.
Robertianum, *Bernard, ex Gren. & Godr. Fl. Fr.* i. 117.—Corsica.
Rochelii, Andrz. ex Reichb. Ic. Pl. Crit. xi. p. 8. n. 4973 b = montanum.
rostratum, *Stev. in Mém. Acad. Pétersb.* iii. (1809–10) 304. f. 1.—Græcia; As. Min.
rostratum, Wierzb. ex Nym. Consp. 56 = montanum.
rudorale, Jord. Diagn. i. 198 = calycinum.
rupestre, *Host, Fl. Nop.* t. 66.—Europ.
rupestre, Willd. Enum. Hort. Berol. 672 = Graellsia saxatilis.
sabulosum, Jord. Diag. i. 199 = calycinum.
samariferum, *Boiss. & Heusch. ex Boiss. Fl. Orient.* i. 272.—As. Min.
samolifolium, Desf. Choix, 66 = Thlaspi samolifolium.
sativum, Scop. Fl. Carn. ed. II. ii. 9 = Camelina sativa.
savranicum, Andrz. ex Bess. Enum. Pl. Volh. 82 = alpestre.
saxatile, Bory & Chaub. Nouv. Fl. Pélop. 61 = orientale.
saxatile, Crantz, Stirp. Austr. 16 = gemonense.
saxatile, *Linn. Sp. Pl.* 650.—Europ.
saxatile, Sibth. & Sm. Prod. Fl. Græc. ii. 11 = orientale.
scabrum, Weinm. Cat. Hort. Dorpat. (1816) 11 = umbellatum.

ALYSSUM :—
Schlosseri, Heuff. ex Schlosser & Vukot. Syll. 186 = calycinum.
schirwanicum, Rupr. Fl. Cauc. 100 = tortuosum.
Scovitzianum, Fisch. & Mey. in Linnæa, xii. (1838) sphalm. 181 = Scovitzianum.
scutigerum, *Dur. in Expl. Sc. Algér.* t. 79. f. 4.—Afr. bor.
semidoxandra, *Schrenk, Boiss. Fl.* ii. 189.—Europ.
serpyllifolium, Bieb. Fl. Taur. Cauc. ii. 308 = tortuosum.
sergyllifolium, Desf. Fl. Atlant. ii. 70 = alpestre.
sibiricum, Fisch. ex DC. Syst. ii. 311 = Fischerianum.
sibiricum, Trevir. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 151 = dasycarpum.
sibiricum, *Willd. Sp. Pl.* iii. 465.—Sibir.
siculum, Jord. Diag. i. 202 = calycinum.
simplex, Rudolphi, in Schrad. Journ. ii. (1799) 290 = campestre.
sinuatum, *Linn. Sp. Pl.* 651.—Europ.; As. Min.
smyrnaeum, *C. A. Mey. in Bull. Sc. Acad. Pétersb.* vii. (1840) 182.—Smyrna.
spathulatum, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 110 = alpestre.
spathulatum, *Steph. in Willd. Sp. Pl.* iii. 465.—Sibir.
speciosum, *Pomel, Nouv. Mat. Fl. Atl.* 337.—Afr. bor.
sphacioticum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 35. —Creta.
spinosum, *Linn. Sp. Pl.* 650.—Europ.
Sprunerі, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 12 = montanum.
strictum, C. A. Mey. Verz. Pfl. Cauc. 127 = Szowitzianum.
strictum, *Willd. Sp. Pl.* iii. 464.—As. Min.
strigosum, [Soland. in] Russell, Aleppo, ed. II. ii. 257 = hirsutum.
strigulosum, *Ann, Fl. Iber.* vi. 571.—Hispan.
styiare, *Boiss. & Bal. ex Boiss. Fl. Orient.* i. 287.—As. Min.
styriacum, Jord. & Fourr. Brev. Pl. Nov fasc. ii. 7 = montanum.
subalpestre, Schur, Enum. Pl. Transs. 61 = argenteum.
subalpinum, Fall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. ii. 108 = tortuosum.
sublineare, Jord. Diagn. i. 201 = calycinum.
submolle, Boiss. & Heldr. ex Nym. Consp. 56 = foliosum.
suffrutescens, *Boiss. Fl. Orient.* i. 275.—Syria.
superbum, L'Hérit. ex DC. Syst. ii. 288 = Farsetia suffruticosa.
Scovitzianum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 51.—Oriens.
tenuifolium, Bieb. Fl. Taur. Cauc. Suppl. 435 = canescens.
tenuifolium, *Steph. in Willd. Sp. Pl.* iii. 460.—Ross.
tetrospermum, Bertol. in Nov. Comm. Acad. Bonon. vi. (1844) 226 = meniocoides.
tetrastemon, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 158.—As. Min.; Syria.
tortuosum, *Rupr. Fl. Caucas.* 98.—Caucas.
tortuosum, Waldst. & Kit. Pl. Rar. Hung. i. 94. t. 91 = alpestre.
transsylvanicum, Schur, Enum. Pl. Transs. 63 = montanum.
trichostachyum, *Rupr. Fl. Caucas.* 101.—Prov. Caucas.
triquetrum, Fortenschl. ex DC. Syst. ii. 290 = Farsetia triquetra.
trochocarpum, Fenzl, in Tchihat. Asie Min. Bot. i. 297 = argenteum.
tuberculatum, Kit. in Linnæa, xxxii. (1863) 498 = muriculatum.
turkestanicum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 5.—Turkest.
umbellatum, *Desv. Journ. Bot.* iii. (1814) 178 & 184. —As. Min.; Syria.
utriculatum, Linn. Mant. i. 93 = Vesicaria utriculata.
utriculatum, Sibth. & Sm. Fl. Græc. t. 637 = Vesicaria Barrelieri.
vagum, Jord. Diag. i. 196 = calycinum.
vernale, Kit. ex Horænn. Hort. Hafn. 691 = montanum.
Vesicaria, Linn. Sp. Pl. ed. I. 651 = Vesicaria utriculata.
vesicarioides, Andrz. ex Steud. Nom. ed. II. i. 68 = gemonense.
Wierzbicki, Heuff. in Flora, xviii. (1835) i. 249 = rostratum.

ALYSSUM:—
Williamsonii, De Roem. ex Willk. in Linnaea, xxv. (1852) 8 = grenatense.
Wulfenianum, Willd. Enum. Hort. Berol. Suppl. 44 = montanum.
xanthocarpum, Boiss. in Ann. Sc. Nat. Sér. xvii. (1842) 154 = macrocalyx.
xerophilum, Jord. & Fourr. Rev. Pl. Nov. fasc. ii. 10 = montanum.

ALYXIA, Banks, ex R. Br. Prod. 469 (1810). *APOCYNACEAE*, Benth. & Hook. f. ii. 697.
GYMPOGOON, Forst. Char. Gen. 35. t. 18 (1776), nomen prius.
actinophylla, A. Cunn. in Bot. Mag. sub t. 3813 = verticillosa.
affinis, *Hewrch & Muell. Arg. in Hewrch, Obs. Bot.* 198.—N. Caled.
aromatica, Reinw. ex A. DC. in DC. Prod. viii. 346 = stellata.
bracteolosa, *Rich, ex A. Gray, in Proc. Am. Acad.* v. (1863) 333.—Ins. Fiji.
brevifora, *Hewrch & Muell. Arg. in Hewrch, Obs. Bot.* 198.—N. Caled.
buxifolia, *R. Br. Prod.* 470.—Austral.
calophylla, Wall. Cat. n. 1607 = Winchia calophylla.
capitellata, Benth. in Enum. Pl. Hueg. 51 = buxifolia.
ceylanica, *Wight, Ic. Pl.* t. 1293.—Zeylan.
coriacea, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 541.—Ind. or.
cuspidata, Wall. ex A. DC. in DC. Prod. viii. 351 = Melodinus orientalis.
daphnoides, *A. Cunn. in Hook. Bot. Mag.* t. 3813.—Ins. Norfolc.
diapharocarpa, *Hewrch & Muell. Arg. in Flora*, li. (1870) 166.—N. Caled.
erythrocarpa, *Vatke, in Bremen Abh.* ix. (1885) 124.—Ins. Mascar.
fascicularis, *Benth. & Hook. f. Gen.* ii. 698, *in nota ad Hunteriam.*—Reg. Himal.
Forsteri, *A. Cunn. ex Loud. Hort. Brit. Suppl.* i. 580 (1830) = A. Gynopogon.
glaucescens, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 642; G. Don, Gen. Syst. iv. 97 = Winchia calophylla.
glaucophylla, *Hewrch & Muell. Arg. in Flora*, liii. (1870) 170.—N. Caled.
gracilis, *R. Muell. Fragm.* viii. 46.—Ins. D. Howe.
lucida, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 540.—Malaya.
madagascariensis, *A. DC. in DC. Prod.* viii. 345.—Madag.
obovata, *Seem. Fl. Vit.* 156, *in nota.*—Ins. Pinet.
obtusifolia, *R. Br. Prod.* 470.—Austral.
odorata, *Wall. Cat. n.* 1606.—Tenasserim.
olivaeformis, *Gaudich. in Freyc. Voy. Bot.* 451.—Ins. Sandvic.
pilosa, *Miq. Fl. Ind. Bat.* ii. 408.—Java.
podocarpa, *Hewrch & Muell. Arg. in Hewrch, Obs. Bot.* 197.—N. Caled.
pugioniformis, A. Cunn. in Hook. Bot. Mag. sub t. 3813 = ruscifolia.
pumila, *Hook. f. Fl. Brit. Ind.* iii. 635.—Malacca.
quinata, *Miq. Fl. Ind. Bat.* ii. 407.—Sumatra.
Reinwardti, Blume, Cat. Gew. Buit. 48 = stellata.
Richardsonii, Sweet, Hort. Brit. ed. I. 272 = ruscifolia.
ruscifolia, *R. Br. Prod.* 470.—Austral.
scandens, *Roem. & Schult. Syst.* iv. 440.—Ins. Societ.
sinensis, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 324.—Hongkong.
Spanoghena, *Miq. Fl. Ind. Bat.* ii. 409.—Ins. Timor.
spicata, *R. Br. Prod.* 470.—Austral.
squamulosa, *C. Moore & F. Muell. in F. Muell. Fragm.* viii. 47.—Ins. D. Howe.
stellata, *Roem. & Schult. Syst.* iv. 489.—As. trop.; Malaya; Ins. Pacif.
sulcata, Hook. & Arn. Bot. Beech. Voy. 90 = olivaeformis.
tetragona, R. Br. Prod. 470 = spicata.
Thouarsii, *F. Muell. Fragm.* x. 108.—Austral.
thyrsiflora, *Benth. Fl. Austral.* iv. 309.—Austral.
Tiascarnii, *Monitr. in Mém. Acad. Lyon*, x. (1860) 333.—N. Caled.
Torresiana, *Gaudich. in Freyc. Voy. Bot.* 451.—Ins. Marian.
seylanica, Walp. Ann. iii. 33 = ceylanica.

ALZALIA, F. G. Dietr. Lexik. Gaertn. i. 299 (1802) = **Alastes**, Ruiz & Pav. (Celastr.).

ALZINIANA, F. G. Dietr. ex Pfeiff. Nom. i. 129 (1872) = prsec.

ALZATEA, Ruiz & Pav. Prod. 40. t. 7 (1794). *CELASTRINEAE*, Benth. & Hook. f. i. 362.
mexicana, F. G. Dietr. Lexik. Gaertn. i. 299 (1802) Nachtr. = verticillata.
verticillata, *Ruiz & Pav. Fl. Per.* iii. 20. t. 241.—Peruv.

AMADEA, Adans. Fam. ii. 230 (1763) = **Androsace**, Tourn. (Primul.).

AMAGRIS, Rafin. in Préc. Somiol. 27 (1814) & Am. Monthly Mag. (1819) 390 = **Calamagrostis**, Adans. (Gramin.).

AMAIOUA, Aubl. Pl. Gui. Suppl. 13. t. 375 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 81.
EHRENBERGIA, Spreng. Neue Entdeck. ii. 129 (1821) & Syst. ii. 13 (1825).
EHRENBERGIA, Willd. ex Schult. f. Syst. vii. 91 (1830).
affinis, *Miq. in Linnaea*, xviii. (1844) 618.—Surinam.
africana, Spreng. Syst. ii. 126 = Coffea sanguebarica.
Hostaliana, *A. Rich. in Mém. Soc. Nat. Par.* v. (1829) 251 † (*Mém. Rub.* 170).—Bras.
corymbosa, *H. B & K. Nov. Gen. et Sp.* iii. 419. t. 294.—Ins.
edulis, *Baill. Hist. Pl.* vii. 387.—Guiana.
eriopila, *Baill. l. c.*—Guiana.
fagifolia, *Desf. in Mém. Mus. Par.* vi. (1820) 14. t. 5.—Guiana.
grandifolia, *Miq. in Linnaea*, xviii. (1844) 291.—Guiana.
guianensis, *Aubl. Pl. Gui. Suppl.* 13. t. 375.—Guiana.
hirsuta, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 25. t. 230.—Bras.
intermedia, *Mart. in Schult. f. Syst.* vii. 90.—Bras.
Laureaster, *Mart. in Flora*, xxiv. (1841); *Beibl.* ii. 84.—Bras.
longifolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 26.—Bras.
peruviana, *Desf. in Mém. Mus. Par.* vi. (1820) 16. t. 4.—Peruv.
sacciferа, *Mart. in Schult. f. Syst.* vii. 91.—Bras.
surinamensis, *Steud. in Flora*, xxvi. (1843) 763.—Surinam.
utilis, Baill. in Bull. Soc. Linn. Par. i. (1879) 230 = Alibertia edulis.

AMALAGO, Rafin. Syll. Syst. Veliur. 84 (1838) = **Piper**, Linn.
antillanus, Rafin. l. c. = P. officinarum.
Madurari, Rafin. l. c. = prsec.

AMALIA, Hort. Hisp. ex Endl. Gen. 188 (1837) = **Tillandsia**, Linn. (Bromel.).
Aeria-incola, Hort. Hisp. ex Steud. Nom. ed. II. i. 69 = T. diantholdes.

AMALIA, Reichb. Nom. 52 (1841) = **Laelia**, Lindl. (Orchid.).

AMALIAS, Hoffmgg. Preis-Vers. Orch. ex Linnaea, xvi. (1843) Litz. 228 = Laelia, Lindl. (Orchid.).
ancops, Hoffmgg. l. c. = Laelia ancops.

AMALOBOTRYA, Kunth, ex Meisn. in DC. Prod. xiv. 185 (1856-57) = **Symmeria**, Benth. (Polygon.).
latifolia, Kunth, ex Meisn. l. c. = S. paniculata.

AMAMELIS, Lem. in Orbign. Dict. iv. 745 (1849) = **Hamamelis**, Gron.

AMANNIA, Blume, Bijdr. 1130 (1826) = **Ammannia**, Houst. (Lythrar.).

AMANOA, Aubl. Pl. Gui. i. 256. t. 101 (1775). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 268.
acuminata, Thw. Enum. Pl. Zeyl. 428 = Cleistanthus acuminatus.
Boiviniana, *Baill. Etud. Gén. Euph.* 582.—Madag.
bracteata, Rich. ex Baill. l. c. 581 = guianensis.
bracteosa, *Planch. in Hook. Ic. Pl.* t. 797.—Afr. trop.
brasiliensis, Baill. Etud. Gén. Euph. 581. n. 4 = Gonatogyne brasiliensis.
chartacea, Baill. l. c. 582 = Cleistanthus chartaceus.
collina, Baill. l. c. = Cleistanthus collinus.
Cunninghamii, Baill. Adansonia, vi. (1865-66) 335 = Cleistanthus Cunninghamii.
Dallachiana, Baill. l. c. = Cleistanthus Dallachyanus.
divaricata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 22. t. 226.—Am. austr.
fagínea, Baill. Adansonia, vi. (1865-66) 336 = Briedelia fagínea.
ferruginea, Thw. Enum. Pl. Zeyl. 230 = Cleistanthus ferrugineus.
grandiflora, *Muell. Arg. in Flora*, lv. (1872) 2.—Surinam.
guianensis, *Aubl. Pl. Gui.* t. 256. t. 101.—Guiana.
indica, Thw. Enum. Pl. Zeyl. 428 = Cleistanthus robustus.
indica, Wight, Ic. t. 1911 = Cleistanthus patulus.
Leichardtii, Baill. Adansonia, vi. (1865-66) 336 = Briedelia fagínea.
oblongifolia, *Muell. Arg. in Linnaea*, xxxii. (1863) 77.—Bras.
ovata, Baill. Adansonia, vi. (1865-66) 336 = Briedelia exaltata.
pallida, Thw. Enum. Pl. Zeyl. 280 = Cleistanthus pallidus.
patula, Thw. l. c. = Cleistanthus patulus.
purpurascens, *Poepp. ex Baill. Etud. Gén. Euph.* 598.—Bras.
racemosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 23.—Bras.
ramiflora, Poepp. & Endl. l. c. = Richeria grandis.
robusta, Thw. Enum. Pl. Zeyl. 428 = Cleistanthus robustus.
strobilacea, *Muell. Arg. in Flora*, xlvii. (1864) 515.—Afr. occ.
strobilantha, Planch. in Hook. Niger Fl. t. 47 = bracteosa.
tomentosa, Baill. Adansonia, vi. (1865-66) 336 = Briedelia tomentosa.

AMAPA, Steud. Nom. ed. I. 35 (1821) = **Carapa**, Aubl. (Meliac.).
guianensis, Steud. l. c. = C. guianensis.

AMARABOYA, Linden, ex Mast. in Gard. Chron. (1871) 944. *MELASTOMACEAE*.
amabilis, *Linden, ex Mast. l. c.*—N. Granat.
princeps, *Linden, ex Mast. l. c.*—N. Granat.
splendida, *Linden, ex Mast. l. c.*—N. Granat.

AMARACANTHUS, Steud. Nom. ed. II. i. 69 (1840) sphalm. = seq.

AMARACARPUS, Blume, Bijdr. 954 (1826). *RUBIACEAE*, Benth. & Hook. f. ii. 180.
microphyllus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 211.—Ins. Celebes.
pubescens, *Blume, Bijdr.* 954.—Java.

AMARACUS, Gleditsch, Syst. 189 (1764) = **Origanum**, Linn. (Labiat.).
cordifolius, Montbr. & Auch. ex Benth. in Ann. Sc. Nat. Sér. II. vi. (1836) 48 = O. cordifolium.
Dictamnus, Benth. Lab. Gen. et Sp. 333 = O. Dictamnus.
nipplus, Rafin. Fl. Tellur. iii. 84 = O. nipplus.
tomentosus, Moench, Meth. Suppl. 127 = O. Dictamnes.
Tournefortii, Benth. Lab. Gen. et Sp. 333 = O. Tournefortii.

AMARALIA, Welw. ex Benth. & Hook. f. Gen. ii. 90 (1873) = Rhetbrownia, G. Don (Rubiac.).
bignoniaeflora, Welw. l. c. ii. 91 = S. folíosa.
Doniana, Welw. l. c. = S. folíosa.

AMARANTESIA, Hort. ex Regel, Gartenfl. (1869) 102 = Telanthera, R. Br. (Amarant.).
brasiliensis, Hort. ex Regel, l. c. = T. Bettzichiana.

AMARANTHUS, Linn. Syst. ed. I. (1735); Gen. ed. 1. 286 (1737). *AMARANTACEAE*, Benth. & Hook. f. iii. 28.
ALBERSIA, Kunth, Fl. Berol. ed. II. 144 (1888).
AMBLOGYNA, Rafin. Fl. Tellur. iii. 42 (1836).
BAJAN, Adans. Fam. ii. 506 (1763).
BLITON, Adans. l. c. (1763).
BLITUM, Linn. Gen. ed. I. 20 (1737).
DIMERIANDRA, Rafin. Neogenyt. 2 (1825).
DIMERIANTHUS, Rafin. Fl. Tellur. iii. 41 (1836).
EUXOLUS, Rafin. l. c. 42 (1836).
GLOMERARIA, Cav. Descr. 319 (1837).
MENGEA, Schau. in Nov. Act. Cur. XIX. Suppl. I. (Pl. Meyn.) 405 (1843).
PENTRIUS, Rafin. Fl. Tellur. iii. 42 (1836).
PYXIDIUM, Moq. Tand. in DC. Prod. xiii. II. 262 (1849) (vox fructus desc. lapas progenere habita).
RORMERIA, Moench, Meth. 341 (1794).
SARRATIA, Moq. ex DC. Prod. xiii. II. 233, 268 (1849).
SCLEROPUS, Schrad. Ind. Sem. Hort. Goett. (1885).

Acroglochin, Spreng. Syst. I. 927 = Acroglochin chenopodioides.
adscendens, Auct. vide *ascendens*.
aeneus, *Bess. Cat. Hort. Volkyn.* (1816) 8.—Hab.?
arruedes, Hochst. & Steud. ex A. Rich. Tent. Fl. Abyss. ii. 214 = Aerva lanata.
albiflorus, *Moq. in DC. Prod.* xiii. II. 268.—Hab.?
albus, *Linn. Syst.* ed. X. 1268.—Am. bor (Europ. alien.).
albus, Robin, Voy. Louis. 374; ex Moq. in DC. Prod. xiii. II. 265 = leucanthus.
albus, Rodsch. ex F. G. Dietr. Lex. Gaertn. Nachtr. II. i. 196 = A. Blitum.
albus, Thunb. Fl. Cap. ed. Schult. i. 215 = Thunbergii.
Alopecurus, Hochst. ex A. Br. & Bouché, Ind. Sem. Hort. Berol. App. (1872) 1 = caudatus.
altissimus, Riddell, Syn. Fl. West. St. (1835) 41 = Acnida rusocarpa.
amboinicus, Buch.-Ham. in Wall. Cat. n. 6987 = gangeticus.
Anacardana, Hook. f. Fl. Brit. Ind. iv. 719 = seq.
Anardana, Buch.-Ham. in Wall. Cat. n. 6998 = paniculatus.
angustifolius, Bieb. ex Willd. Sp. Pl. iv. 381 = sylvestris.
angustifolius, Lam. Encyc. i. 115 = A. Blitum.
Arardhanus, Sweet, Hort. Brit. ed. III. 569 = paniculatus.
arctioides, Perr. ex Moq. in DC. Prod. xiii. II. 332 = Pupalia lappacea.
ascendens, *Horwem. Hort. Hafn. Suppl.* 107.—China.
ascendens, Loisel. Not. 141 = viridis.
atropurpureus, *Roxb. Fl. Ind.* iii. 608.—Ind. or.
atrosanguineus, Hort. Lugd. (1838) ex Moq. in DC. Prod. xiii. II. 266 = hypochondriacus?
aureus, Bess. Cat. Hort. Volkyn. (1816) 8 = flavus?
aureus, *F. G. Dietr. Lexik. Gaertn.* i. 310.—Nepal.
aureus, Hort. ex Moq. in DC. Prod. xiii. II. 259 = hybridus.
bahiensis, *Mart. Herb. Fl. Bras.* n. 969.—Bahia.
Bellardi, Hort. ex Moq. in DC. Prod. xiii. II. 259 = hybridus.
Berckholdi, Hort. ex Moq. l. c. 257 = paniculatus.
Berckholdi, Seidl, ex Opiz, Boehemis Gew. 164; Natural. i. (1838) [2] = A. Blitum.
Berckholdi, Hort. ex Moq. in DC. Prod. xiii. II. 259 = hybridus.
Bernhardi, Hort. ex Moq. l. c. 258 = flavus.
bicolor, Nocca, ex Willd. Sp. Pl. iv. 384 = melancholicus.
bitioides, *S. Wats. in Proc. Am. Acad.* xii. (1877) 273.—Am. bor.
blitoides, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Blitum.
Blitum, Auct. ex Moq. in DC. Prod. xiii. II. 267, 274 = viridis.
Blitum, *Linn. Sp. Pl.* ed. I. 990.—Reg. temp. et trop.
Blitum, Moq. in DC. Prod. xiii. II. 268 = sylvestris.
Blitum, Reichb. Ic. ex Steud. Nom. ed. II. i. 69 = polygonoides.
brasiliensis, *Moq. in DC. Prod.* xiii. II. 266.—Bras.
bullatus, Bess. ex Spreng. Syst. i. 928 = retroflexus.
californicus, *S. Wats. in Bot. Calif.* ii. 48.—Calif.
campestris, *Willd. Sp. Pl.* iv. 382.—Ind. or.
canariensis, *Bess. Cat. Hort. Volkyn.* (1816) 8.—Ins. Canar.

AMARANTHUS:—
capitatus, Cat. Hort. Turic. (1827–28).—Hab.?
Caracasw, Bess. Cat. Hort. Volkyn. (1816) 8 = A. Caracu.
caracasanus, *H. B. & K. Nov. Gen. et Sp.* ii. 195.—Venezuela.
Caracu, *Linn. ex* Steud. Nom. ed. II. i. 69 = A. Caracu.
Cararia, Bess. Cat. Hort. Volhyn. (1816) 8 = prsec.?
Cararu, Hort. Paris. ex Moq. in DC. Prod. xiii. II. 274 = caudatus.
Carsru, *Facq. ex Zuccag. in Rosm. Collect.* 158.—Hab.?
carneus, Hort. ex Moq. in DC. Prod. xiii. II. 257 = paniculatus.
Cathcarti, Hort. ex Moq. l. c. 259 = hybridus.
Caturus, *Roxb. in Herb. Made. sec. Wall. Cat.* n 6900.—Ind. or.
caudatus, *Linn. Sp. Pl.* 990.—Orient; Afr. trop.; Ind. or.
caulifiorus, Link, Enum. Hort. Berol. ii. 389 = Acroglochin chenopodioides.
celosioides, *H. B. & K. Nov. Gen. et Sp.* ii. 194.—N. Granat.
cernuus, Bess. in Cat. Hort. Turic. (1827–8) 'An Celosia?'
chlorostachys, Moq.-Tand. in DC. Prod. xiii. II. 259 = paniculatus.
chlorostachys, *Willd. Amarant.* 34. t. 10. f. 19.—Europ.; Orient; Ind. or.
chlorostachys, Willk. ex Willk. & Lange, Prod. Fl. Hisp. i. 275 = retroflexus.
circinnatus, Hort. Paris. ex Poir. Encyc. Suppl. i. 811 = lividus?
cocsius, *F. G. Dietr. ex Moq. in DC. Prod.* xiii. II. 266.—Hab.?
commutatus, *Kœrner, in Oestr. Bot. Zeitschr* XXV. (1875) 194.—Austr.
coracanus, Mart. Hort. Erlang. 197 = carcasanus.
crassipes, Schlecht. in Linnæa, vi. (1831) 757 = polygonoides.
cristatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 7 = Celosia cristata.
crocatus, Bess. Cat. Hort. Volhyn. (1816) 9.—Hab.?
cruentus, Linn. Syst. ed. X. 1269 = paniculatus.
cruentus, Willd. ex Roxb. Fl. Ind. iii. 610 = caudatus.
curvipilus, Spreng. Syst. i. 928 = retroflexus.
cuspidatus, Vis. Cat. Hort. Patav. (1841) = gangeticus.
deflexus, *Linn. Mant.* ii. 295.—Europ.
Delileï, Richter & Loret, in Bull. Soc. Bot. Fr. xiii. (1866) 316 = patulus.
diacanthus, *Rafin. Fl. Ludov.* 81.—Am. bor.
diandrus, Spreng. Neue Entdeck. iii. 20 = Acroglochin chenopodioides.
diffusus, Dulac, Fl. Hautes-Pyr. 174 = A. Blitum.
dioicus, Michx. ex Moq. in DC. Prod. xiii. II. 277 = Acnida rusocarpa.
divaricatus, *Andrz. ex Lindem. Prod. Fl. Cherson.* 185.—Cherson.
dubius, Mart. Hort. Erlang. (1814) 197 = tristis.
edulis, Michx. ex Moq. in DC. Prod. xiii. II. 277 = Acnida cannabina.
emarginatus, *Salem. ex Moq. l. c.* 274 = viridis.
emersus, *F. Muell. ex Benth. Fl. Austral.* v. 216.—Austral.
esculentus, Bess. Cat. Hort. Volhyn. (1816) 9; ex Moq. in DC. Prod. xiii. II. 266 = paniculatus.
farinosus, Roxb. ex Moq. l. c. = paniculatus.
fasciatus, Roxb. Fl. Ind. iii. 609 = viridis.
Fatine, Mart. Hort. Erlang. 197.—Hab.?
fimbriatus, *S. Wats. in Bot. Calif.* ii. 48.—Calif.
flavescens, Hort. ex Moq. in DC. Prod. xiii. II. 259 = hybridus.
flavus, *Linn. Syst.* ed. X. 1269.—Ind. or.
flexuosus, Hort. ex Moq. in DC. Prod. xiii. II. 260 = tristis.
floridus, Benth. Bot. Voy. Sulph. 158. t. 51 = acrianus.
frumentaceus, Buch.-Ham. in Roxb. Fl. Ind. iii. 599 = paniculatus.
frutescens, Hort. ex Moq. in DC. Prod. xiii. II. 248 = Iresine amarantoides.
gangeticus, *Linn. Syst.* ed. X. 1268.—Reg. trop.
gangeticus, Wall. Cat. n. 6896 = A. Blitum.
giganteus, Bess. Cat. Hort. Volhyn. (1816) 91; F. G. Dietr. Lexik. Gaertn. i. 313.—Hab.?
giganteus, L'Hérit. ex Moq. in DC. Prod. xiii. II. 277 = Acnida cannabina.
graecizans, Desf. Tabl. Hort. Par. ed. I. 49 = caudatus.
graecizans, Auct. Hesp. ex Cat. Fl. Comp. Madr. 879 = albus.
graecizans, Linn. Sp. Pl. 990 = sylvestris.

AMARANTHUS:—
Greggii, *S. Wats. in Proc. Am. Acad.* xii. (1877) 274.—Am. bor. occ.
grandesquamis, Hort. ex Moq. in DC. Prod. xiii. II. 257 = paniculatus.
hecticus, Willd. Amaranth. 28 = hybridus.
hierichuntinus, *Vis. in Atti. Ist. Ven. Sc.* Ser. III. iv. (1858–59) 189.—Syria.
Huttoni, Hort. Veitch. cf. Gard. Chron. (1872) 215 = Celosia Huttoni.
hybridus, *Linn. Sp. Pl.* 990.—Am. bor.
hybridus, Vell. Fl. Flum. x. t. 98 = flavus?
hypochondriacus, *Linn. Sp. Pl.* 991.—Am. bor. (Europ. alien.).
inamoenus, Willd. Sp. Pl. iv. 386 (1805) Amar. 14 = gangeticus.
incarnatus, Hort. ex Moq. in DC. Prod. xiii. II. 257 = paniculatus.
incomptus, Willd. Enum. Hort. Berol. 64 = gangeticus.
incrustatus, Gren. & Godr. Prosp. Fl. Fr. 8 = chlorostachys.
intermedius, Guss. ex Moq. in DC. Prod. xiii. II. 259 = chlorostachys.
interruptus, *R. Br. Prod.* 414.—Austral.
japonicus, Houtt. ex Steud. Nom. ed. I. 86 = gangeticus.
lacteus, Hort. Amstelod. (1814) 59 = Aerva sanguinolenta.
laetus, Willd. Amaranth. 28; Sp. Pl. iv. 391 (1805) = hybridus.
lanatus, Dum.-Cours. Bot. Cult. ed. I. i. 640 = Aerva lanata.
lanceifolius, Roxb. Hort. Beng. 67 = gangeticus.
lanceolatus, Roxb. Fl. Ind. iii. 607 = gangeticus.
lanciflorus, *Del. ex Moq. in DC. Prod.* xiii. II. 276 = interruptus.
lanifiorus, Comell. ex Poll. Fl. Veron. iii. 114 = patulus.
leptostachyus, *Benth. Fl. Austral.* v. 214.—Austral.
leucanthus, *Rafin. Fl. Ludov.* 82.—Am. bor.
leucocarpus, *S. Wats. in Proc. Am. Acad.* x. (1875) 347.—Arizona.
limatus, R. Br. Prod. 414 = interruptus.
littoralis, Bernh. in Hort. Tar. (1813) ex Moq. in DC. Prod. xiii. II. 274 = viridis.
littoralis, Horum. Hort. Hafn. (1815) 893 = albus.
lividus, Hort. Petrop. ex Hook. f. Fl. Brit. Ind. iv. 721 = A. Blitum.
lividus, *Linn. Sp. Pl.* 990.—Am. bor.
lividus, Roxb. Fl. Ind. iii. 605 = gangeticus.
macrocarpus, *Benth. Fl. Austral.* v. 216.—Austral.
macrocaulos, Poir. Encyc. Suppl. i. 814 = Acnida cannabina.
macrostachyus, Merat, ex Moq. in DC. Prod. xiii. II. 266 = hypochondriacus.
major, Salzm. ex Moq. in DC. Prod. xiii. II. 274.—Hab.?
mangostanus, Blanco, Fl. Filip. ed. I. 711 = melancholicus.
mangostanus, *Linn. Cent.* i. 32; *Am. Acad.* iv. 294.—Ind. or.
Mantagazianus, *Passer. in Hort. Parm.* (1864) 6.—Am. austr.
maximus, Mill. Gard. Dict. ed. VIII. n. 5 = caudatus.
melancholicus, Linn. Sp. Pl. 989 = gangeticus.
miamiensis, *Riddell, Syn. Fl. West. St.* (1835) 41.—Am. bor.
minor, S. F. Gray, Nat. Arr. Brit. Pl. ii. 939 = A. Blitum.
minutus, Leschen. ex Moq. in DC. Prod. xiii. II. 304 = Aerva brachiata.
Mitchelli, *Benth. Fl. Australiensis*. v. 214.—Austral.
monstrosus, Cat. Hort. Div. (1837); Hort. Tonelle, ex Moq. in DC. Prod. xiii. II. 256 = hypochondriacus.
montevidensis, Hort. ex Moq. l. c. 257 = paniculatus.
morosus, *Reichb Fl. Germ. Excurs.* 856.—Europ.
mucronatus, Hort. Petrop. ex Hook. f. Fl. Brit. Ind. iv. 720 = mangostanus.
mucronatus, Poir. Encyc. Suppl. i. 811 = A. Blitum.
muricatus, *Gill. ex Moq. in DC. Prod.* xiii. II. 276.—Hab.?
nagiachus, Hort. ex Moq. l. c. 259 = hybridus.
napalensis, Cat. Hort. Lugd. (1826); ex Moq. l. c. = hybridus.
obovatus, *S. Wats. in Proc. Am. Acad.* xii. (1877) 275.—Am. bor.
officinalis, Gromow. ex Trautv. in Act. Hort. Petrop. ix. (1884) 189 = seq.?
olereceus, *Linn. Sp. Pl.* 1406.—Egypt.; Ind. or.
olereceus, Rodsch. Enumerth. 24 = A. Blitum.
olereceus, Benth. Fl. Ind. iii. 603 = gangeticus.
oliteorius, Sam. Cat. Hort. Volhyn. (1816) 9 = olereceus?

AMARANTHUS :—
pachystachys, Retchb. ex Moq. in DC. Prod. xiii. 11. 265 = parviramis.
pallidiflorus, *F. Muell. Fragm.* i. 140.—Austral.
pallidus, Bieb. Fl. Taur. Cauc. ii. 399 = A. Blitum.
Palmeri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 274. —Calif.
paniculatus, *Linn. Sp. Pl.* ed. II. 1406.—Am. bor. (in Orient. alien.).
paniculatus, Wall. Cat. n. 6904 = caudatus.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 278 ; *Contrib.* 79.—Reg. Argent.
parviensis, *Schk. Handb.* iii. 249.—Hab. ?
pastulus, *Bertol. Comm. Neog.* 171. t. 12.—Am. bor. (in Europ. alien.).
pendulinus, Hort. ex Moq. in DC. Prod. xiii. 11. 255 = caudatus.
pendulus, Hort. ex Moq. l. c. = caudatus.
pericarioides, Hort. ex Poir. Encyc. Suppl. i. 311 = Acroglochin chenopodioides.
polychrus, Raensch. Nom. ed. III. 275 = Pellionia brevifolia ?
polyflagellus, *Sprang. in Hort. Argent.* (1884) *ex Moq. in DC. Prod.* xiii. 11. 266.—Hab. ?
polygamus, *Linn. Cent.* i. ; *Am. Acad.* iv. 294.—Afr. trop. ; Ind. or.
polygamus, Wall. Cat. n. 514 = A. Blitum.
polygamus, Thw. Enum. Pl. Zeyl. 247 = mangostanus.
polygamus, Roxb. Fl. Ind. iii. 607 = gangeticus.
polygonoides, *Linn. Pl. Jam. Pugill.* ii. 27.—Jamaic ; *Zeyl.*
polygonoides, Zolling. ex Moq. in DC. Prod. xiii. 11. 274 = A. Blitum.
polygonoides, Roxb. Fl. Ind. iii. 721 = polygamus.
polystachyus, Buch.-Ham. in Wall. Cat. n. 6901 = viridis.
polystachyus, *Willd. Sp. Pl.* iv. 885.—Ind. or.
Powellii, *S. Wats. in Proc. Am. Acad.* x. (1875) 347. —Am. bor
prostratus, Balb. Misc. Bot. 44. t. 10 = deflexus.
prostratus, Bast. Fl. Maine-et-Loire, 344 = A. Blitum.
pumilus, *Raffin. in Med. Repos. New York*, v. (1808) 360.—Am. bor.
purpureus, Hort. ex Moq. in DC. Prod. xiii. 11. 257 = paniculatus.
pyramidalis, Noronha, in Verh. Batav. Gen. v. (1790), ed. I. Art. iv. 7 = Celosia argentea.
quitensis, *H. B. & K. Nov. Gen. et Sp.* ii. 194.—Peruv.
recurvatus, Desf. Cat. Hort. Par. ed. III. (1829) 39 = retroflexus.
retroflexus, *Linn. Sp. Pl.* 991.—Am. bor. (in Europ. alien.).
rhombeus, R. Br. Prod. 414 = interruptus.
rigidus, Schult. ex Stend. Nom. ed. II. i. 70 = retroflexus.
rotundifolius, Herb. Par. ex Moq. in DC. Prod. xiii. 11. 261 = gangeticus.
rubescens, Hort. ex Moq. l. c. 257 = paniculatus.
rubricaulis, *Page ; Hort. Angl. ex Moq. in DC. Prod.* xiii. 11. 267, nomen.—Ind. or.
ruderalis, Koch, ex Moq. l. c. 274 = A. Blitum.
sallefolius, *Hort. Veitch, ex Gard. Chron.* (1871) 1550. —Ins. Philipp.
sanguineus, Linn. Sp. Pl. ed. II. 1407 = paniculatus.
sanguineus, *Vell. Fl. Flum.* x. t. 99.—Bras.
sanguinolentus, Schrad. ex Moq. in DC Prod. xiii. 11. 267 = paniculatus ?
scandens, *Linn. f. Suppl.* 419.—Am. austr.
scariosus, *Benth. Bot. Voy. Sulph.*158. t. 51.—Guatemala.
spathulatus, Desf. ex Moq. in DC. Prod. xiii. 11. 276 = linearis.
speciosus, Sims, Bot. Mag. t. 2227 = paniculatus.
spicatus, Lam. Fl. Fr. ii. 193 = retroflexus.
spicatus, Wirzen, Diss. Fl. Casan. 454 = paniculatus.
spicatus, Retchb. Fl. Germ. Excurs. 585 = patulus.
spinosus, *Linn. Sp. Pl.* 991.—Reg. trop.
spiralus, *Zipp. ex Span. in Linnæa*, xv. (1841) 845, nomen.—Malaya.
strictus, Tenore, Syll. 197 = retroflexus.
strictus, Willd. Amaranth. 27 = paniculatus.
sylvestris, Desf. Tabl. Hort. Par. ed. I. 44.—Europ. ; Orient.
tamariscus, Pluch. Hort. Hal. ex Moq. in DC. Prod. xiii. 11. 267 = seq. ?
tamariscinus, Retzh. Fl. Ind. iii. 602 = A. Blitum.
tamariscinus, *Willd. Sp. Pl.* iv. 951.—Ind. or.
tamariscinus, Wall. Cat. n. 6902 = polygamus.
tenuis, *Benth. Fl. Austral.* v. 216.—Austral.

AMARANTHUS :—
Thunbergii, *Moq. in DC. Prod.* xiii. 11. 262.—Afr. austr.
Timorensi, Jord. ex Moq. l. c. 259 = patulus.
Torreyi, *Benth. ex S. Wats. Bot. Calif.* ii. 42.—Am. bor.
tortuosus, *Hornem. Hort. Hafn. Suppl.* 107.—Ins. S. Cruc.
tricolor, Linn. Sp. Pl. 989 = gangeticus.
tristis, Linn. l. c. = gangeticus.
tristis, Wall. Cat. n. 6905 = mangostanus.
trivialis, Rota, in Giorn. Bot. Ital. ii. 11. (1846) 267 = patulus.
undulatus, Lindl. in Mitch. Trop. Austr. 102 = Moschkil ?
undulatus, R. Br. Prod. 414 = interruptus.
urceolatus, *Benth. in Bot. Voy. Sulph.* 158.—Guayaquil.
venulosus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 276.—Am. bor.
vernus, *Opiz, ex Moq. in DC. Prod.* xiii. 11. 267, nomen. —Europ.
verticillatus, Pers. ex Moq. in DC. Prod. xiii. 11. 270 = polygonoides.
violaceus, Hort. ex Moq. l. c. = paniculatus.
viridis, All. Fl. Pedem. ii. 119 = A. Blitum.
viridis, *Linn. Sp. Pl.* ed. II. 1405.—Reg. trop.
viridis, Pall. ex Stend. Nom. ed. II. i. 86 = pallidus.
viridis, Pollich, ex Moq. in DC. Prod. xiii. 11. 268 = paniculatus.
Wrightii, *S. Wats. in Proc. Am. Acad.* xii. (1877) 275.—Am. bor.
zaireus, *Hornem. ex Moq. in DC. Prod.* xiii. 11. 267.—Hab. ?

AMARANTUS, Linn. Syst. ed. x. 1268 (1759) = **Amaranthus**, Linn.

AMARELLA, Gilib. Fl. Lituan. i. 36 (1781) = **Gentiana**, Linn.
acuta, Raffin. Fl. Tellur. iii. 21 = G. Amarella.
brevicaulis, Raffin. l. c. = G. Amarella.
gracilis, Raffin. l. c. = G. Amarella.
quadrifida, Gilib. Fl. Lituan. i. 36 = G. campestris.
quinquefida, Gilib. l. c. = G. Amarella.
ruyana, Raffin. Fl. Tellur. iii. 21.—Hab. ?
vulgaris; Raffin. l. c. 20 = G. campestris.

AMARENUS, C. Presl, Symb. Bot. i. 46 (1830) = **Trifolium**, Linn. (Legum.).
agrarius, C. Presl, l. c. = T. agrarium.
asperus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362 = T. agrarium.
badius, C. Presl, l. c. = T. badium.
campestris, Fourr. l. c. = T. campestre.
filiformis, C. Presl, l. c. = T. filiforme.
flavus, C. Presl, l. c. = T. flavum.
micranthus, Fourr. l. c. = T. filiforme.
patens, C. Presl, l. c. = T. patens.
procumbens, C. Presl, l. c. = T. procumbens.
Schreberi, Fourr. l. c. = T. procumbens.
Sebastiani, C. Presl, l. c. = T. Sebastiani.
spadiceus, C. Presl, l. c. = T. spadiceum.
speciosus, C. Presl, l. c. = T. speciosum.

AMARIA, Mutis, ex Caldas, in Seman. Nuev. Gran. ii. (1810) 25 = **Bauhinia**, Linn. (Legumin.).
petiolata, Mutis, l. c. = B. petiolata.
sessilifolia, Mutis, l. c.—N. Granat.

AMARIDIUM, Hort. ex Lebbers, Cat. Pl. Rar. San Donato, 15 (1860) = **Camaridium**, Lindl. (Orchid.).

AMARORIA, A. Gray, Bot. U. St. Expl. Exped. i. 356. t. 40 (1854). *SIMARUBEAE*, Benth. & Hook. f. i. 314.
soulameoides, A. Gray, l. c.—Ins. Fiji.

AMARYLLIS, Linn. Syst. ed. I. (1735). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 727.
ATAMOSCO, Adans. Fam. ii. 57 (1763).
BELLADONNA, Sweet, Hort. Brit. ed. II. 506. (1830).
CALLICORE, Link, Handb. i. 193 (1829).
CALLIBOE, Endl. Gen. 176 (1887).
acuminata, Ker-Gawl. in Bot. Reg. t. 584 = Hippeastrum rutilum.
adorna, Ker-Gawl. l. c. t. 1125 = Hippeastrum advenum.
aethnensis, 'Raffin.' vide *stenanii*.
africana, Lam. Encyc. i. 194 = Lycoris aurea.
Agathæ, Steud. Nom. ed. II. i. 78 = H. reticulatum.
alba, Forsk. Fl. Aegypt. Arab. 208 = Crinum album.

AMARYLLIS :—
Alberti, Lem. Illustr. Hortic. (1866) t. 498 = Hippeastrum Reginae.
alta, Steud. Nom. ed. II. i. 78 = Hippeastrum splendens.
Andersonii, Steud. l. c. i. 71 = Zephyranthes Andersoni.
andicola, Poepp Fragm. Syn. 5 (= *Habranthus andicolus*).
angusta, Schult. f. Syst. vii. 807 = Hippeastrum bifidum.
Aramasco, Linn. Sp. Pl. ed. II. 420 (sphalm.) = Zephyranthes Atamasco.
arvensis, F. G. Dietr. Lexic. Gaertn. Nachtr. i. 204 = peruviana.
Atamasco, Linn. Sp. Pl. 299 = Zephyranthes Atamasco.
Atamasco, Blanco, Fl. Filip. ed. I. 254 = Hippeastrum miniatum.
atanasia, Cranta, Inst. i. 486 = Zephyranthes Atamasca.
aucta, Tratt. Thes. 45 = Nerine undulata.
aulica, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 353 ; et in Bot. Reg. t. 44 = Hippeastrum aulicum.
aurea, Ruiz & Pav. Fl. Per. iii. 56 (= *Pyrolirion aureum*).
australasica, Ker-Gawl. in Bot. Reg. t. 426 = Crinum flaccidum.
australis, Spreng. Syst. ii. 54 = praec.
Bagnoldi, D. Dietr. Syn. Pl. ii. 1177 = Hippeastrum Bagnoldi.
bahiensis, DC. Pl. Rar. Jard. Genève, 7me Notice, 18 = Hippeastrum bahiense.
Bangoldi, Steud. Nom. ed. II. i. 70, sphalm = *Bagnoldi*.
Bambusiana, Lindl. Bot. Reg. (1842), t. 11. & Append. 18 = Brunvigia Sloteriana.
Belladonna, *Linn. Sp. Pl.* 293.—Afr. austr.
Belladonna, E. Mey. ex Steud. Nom. ed. II. i. 70 = Hippeastrum equestre.
Berteri, Spreng. Syst. ii. 49 (= *Habranthus robustus*).
bicolor, Ruiz & Pav. Fl. Per. iii. 57 = Hippeastrum bicolor.
bifida, Spreng. Syst. Cur. Post. 183 = Hippeastrum bifidum.
bifolia, Aubl. Pl. Gui. i. 304 = Zephyranthes roseo.
bivegineta, Donn, Hort. Cantab. ed. VIII. 98 = Cyrtanthus pallidus.
blanda, Ker-Gawl. in Bot. Mag. t. 1450 = A. Belladonna.
brasiliensis, Andr. Bot. Rep. t. 358 = Hippeastrum equestre, Reginae.
brasiliensis, Tratt. Tabul. t. 325 = Hippeastrum rutilum.
Broussonetii, [DC. in] Red. Liliac. ii. 62 = Crinum yuccaeflorum.
bulbisperma, Burm. f. Fl. Cap. Prod. 9 = Crinum capense.
caerulea, *Griseb. in Gortt. Abh.* xxiv. (1879) 220.—Reg. Argent.
calyptrata, *Hoffmgg. in Linnaea*, xvi. (1842) *Litt.* 279 —Hab. ?
calyptrata, Ker-Gawl. Rev. 11 ; et in Journ. Sc. & Arts, ii. (1817) 358 = Hippeastrum calyptratum.
candida, Lindl. in Bot. Reg. t. 734 = Zephyranthes candida.
candida, Tratt. Tabul. t. 488 = Crinum giganteum.
capensis, Linn. Fl. Afr. 10 ? (Amoen. Acad. vi. 86) = Hypoxis stellatus.
capensis, Will. Gard. Dict. ed. VIII. n. 12 = Crinum longifolium.
carinata, Spreng. Syst. Cur. Post. 152 = Zephyranthes carinata.
carnea, Schult. f. Syst. vii. 799 = Zephyranthes roseo.
Carolinae, Sterler, Hort. Nymph. (1836) 64 = Hippeastrum reticulatum.
caspia, Wild. Sp. Pl. ii. 62 = Allium caspium.
chilensis, L'Hérit. Sert. Angl. 11 = Hippeastrum chilense.
chilensis, Ruiz & Pav. Fl. Per. iii. 56 = Hippeastrum nobilissimum.
chloroleuca, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 347. t. 8. f. 1 = Zephyranthes chloroleuca.
citrina, Linn. Sp. Pl. ed. II. 422 = Sophane ciliaris.
cinnamomea, L'Hérit. Sert. Angl. 15 = Hessea crispa.
citrina, Sibth. & Sm. Fl. Graec. iii. t. 311 = Sternbergia colchiciflora.
Clusiana, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 346 = Sternbergia colchiciflora.

AMARYLLIS :—

coccinea, Molina, Saggio Chil. ed. II. 284 = chilensis.
colchiciflora, Hort. Mosq. ex Schult. f. Syst. vii. 1135 = Bulbocodium versicolor.
colchiciflora, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 345 = Sternbergia colchiciflora.
concinna, *Morris, Fl. Conspic.* t. 44.—Hab.?
coracina, Steud. Nom. ed. II. i. 70 = seq.
coranica, Ker-Gawl. in Bot. Reg. ii. 139 = Ammocharis falcata.
corvinensis, Bury, Hexand. Pl. 9 = Hippeastrum organense.
cornuta, Ker-Gawl. in Bot. Mag. fol. 1430 = Nerine curvifolia.
crispa, Jacq. Hort. Schoenb. i. 37 = Hessea crispa.
crocata, Ker-Gawl. in Bot. Reg. t. 88 = Hippeastrum rutilum.
curvifolia, Jacq. Hort. Schoenb. i. 33 = Nerine curvifolia.
Cybister, Herb. ex Fl. des Serres, v. t. 455, 456 = Hippeastrum Cybister.
cylindracea, L'Hérit. Sert. Angl. 15 = Cyrtanthus angustifolius.
cyrtanthoides, Sims, Bot. Mag. t. 2399 = Hippeastrum bicolor.
demissa, Steud. Nom. ed. II. i. 71.—Chili.
despoupratæ, Poepp. Fragm. Syn. 4 = Zephyranthes despropeæa.
distichа, Sims, Bot. Mag. sub t. 1948 = Crinum yuccæflorum.
distichа, Linn. f. Suppl. 195 = Buphane disticha.
Drummondii, Steud. Nom. ed. II. i. 71 = Cooperia Drummondii.
Dryades, Vell. Fl. Flum. 130; iii. t. 117 = Griffinia Dryades.
dubia, Houtt. Plantenk. xii. 181. t. 84. f. 1 = Nerine sarniensis.
dubia, Lin. Dias. Pl. Scrin. n. 39; et Am. Acad. viii. 354 = Hippeastrum equestre.
elata, Jacq. Hort. Schoenb. i. 38 = Vallota purpurea.
elegans, Spreng. Pugill. ii. 59 = Hippeastrum solandræflora.
Entreriana, Hoffmgg. in Linnæa, xliii. (1880–82) 187 = Zephyranthes mesochlora.
equestris, Ait. Hort. Kew. i. 417 = Hippeastrum equestre.
etnensis, Rafin. Caratteri, 84 = Sternbergia colchiciflora.
exigua, Schousb. Vextr. Marokko, 150 = Tapeinanthus humillis.
falcata, L'Hérit. Sert. Angl. 13 = Ammocharis falcata.
falcata, Lee, ex Steud. Nom. ed. II. i. 73, 475 = Cyrtanthus angustifolius.
flaccida, Weinm. in Flora, iii. (1820) 608 = Crinum flaccidum.
flammea, Ruiz & Pav. Fl. Per. iii. 56 (= *Pyrolirion flammeum*).—Peruv.; Chili.
flava, Pav. ex Roem. & Schult. Syst. vii. 806 = Zephyranthes flava.
flavovirens, Schott, ex Steud. Nom. ed. II. i. 71.—Cf. *fulvovirens*.
flexuosa, Jacq. Hort. Schoenb. i. 35 = Nerine flexuosa.
Forbesii, Lindl. in Trans. Hort. Soc. vi (1826) 87 = Crinum Forbesianum.
formosa, Salisb. Prod. 284 = seq.
formosissima, Linn. Sp. Pl. 293 = Sprekelia formosissima.
Fothergillia, Andr. Bot. Rep. iii. 168 = Nerine curvifolia.
Fothergillii, Pohr. Encyc. Suppl. i. 980 = prec.
fragida, Ker-Gawl. in Bot. Reg. t. 226 = Hippeastrum rutilum.
fulvovirens, Schott; in Schreib. Nachr. Oesterr. Naturf. 129 = Hippeastrum calyptratum.
Gardneri, Seub. in Mart. Fl. Bras. iii. 149 = Hippeastrum organense.
gigantea, Ait. Hort. Kew. ed. II. ii. 226 = Crinum giganteum.
gigantea, Van Marum, in Verh. Maats. Haarl. iii. (1806) 345 = Brunsvigia Josephinæ.
glaucescens, Mart. in Schult. f. Syst. vii. 813 = Hippeastrum rutilum.
gracilifolia, Schult. f. l. c. 806 (= *Habranthus gracilifolius*).
gracilis, Spreng. Syst. Cur. Post. 133 = prec.
grandiflora, D. Dietr. Syn. Pl. ii. 1181 = Brunsvigia grandiflora.
grandiflora, Steud. Nom. ed. II. i. 73 = Sprekelia formosissima.

AMARYLLIS :—

guttata, Linn. Sp. Pl. 294 = Buphane ciliaris.
haemanthoides, Tratt. Gartenpfl. 36 = Haemanthus amarylloides.
humilis, Jacq. Hort. Schoenb. i. 36 = Nerine humilis.
humilis, Link, Enum. i. 313 = Cyrtanthus uniflorus.
hyacinthina, Ker-Gawl. in Bot. Reg. t. 168 = Griffinia hyacinthina.
ignea, Lindl. Bot. Reg. t. 800 = Hippeastrum bicolor.
ignescens, Regel, Gartenfl. xxiii. (1874) 150 = Hippeastrum ignescens.
illustris, Vell. Fl. Flum. 131; iii. t. 118 = Hippeastrum psittacinum.
insignis, Ker-Gawl. Bot. Reg. t. 579 = Crinum latifolium.
intermedia, Lindl. Bot. Reg. t. 1146 (= *Habranthus intermedius*).
intermedia, Spreng. Syst. iv. Cur. Post. 138 = Griffinia intermedia.
Jacquinii, Tratt. Gartenpfl. 43 = Nerine sarniensis.
Jagua, Thompa. Bot. Disp. t. 6 = Crinum giganteum.
Josephinæ [Del.] Red. Lil. vii. 870 = Brunsvigia Josephinæ.
Josephiniana, Herb. Amar. 278 = prec.
Karwinskii, Zucc. in Otto & Dietr. Allg. Gartz. ii. (1834) 345 = Sprekelia formosissima.
hermetiana, Lindl. in Bot. Reg. t. 1638 = Hippeastrum bifidum.
laterita, *A. Dietr. in Otto & Dietr. Allg. Gartz.* xviii. (1850) 55.—Guinea.
latiсoma, Ker-Gawl. in Bot. Reg. vi. 497 = Nerine laxida.
latifolia, Lam. Encyc. i. 124 = Crinum giganteum.
latifolia, L'Hérit. Sert. 14 = Crinum ornatum.
leonensis, *Van Houtte, Hort. Vanhoutt. fasc.* i. (1845) 5.—Afr. trop.
Leopoldii, Hort. Veitch. in Gard. Chron. (1870) 788 = Hippeastrum Leopoldii.
Lindleyana, Schult. f. Syst. vii. 802 = Zephyranthes carinata, Lindleyana.
linearifolia, Molina, Chils, ed. II. 284 = Hippeastrum chilense.
Linearis, Willd. Sp. Pl. ii. 58 = Crinum lineare.
lineata, Colla, Hort. Rip. 3 = Hippeastrum vittatum.
lineata, Lam. Encyc. i. 123 = Crinum zeylanicum.
littoralis, Salisb. Prod. 280 = Crinum latifolium.
lituata, *Reichb. Ic.* t. 82.—Hab.?
longa, Hort. ex Schult. f. Syst. vii. 854 = Crinum yuccæflorum.
longifolia, Steud. Nom. ed. II. i. 70, 71 = Hippeastrum solandræflorum.
longifolia, Linn. Sp. Pl. 293 = Crinum longifolium.
longipetala, *Lem. Illustr. Hortic.* xiii. (1866) *Misc.* 78. Hab.?
lorifolia, Steud. Nom. ed. II. i. 71 (= *Habranthus lorifolius*).—Am. austr.
Lorii, *Morr. in Ann. Soc. Agr. Gand*, ii. (1846) 141. t. 62.—Hab.?
lucida, D. Dietr. Syn. Pl. ii. 1181 = Nerine lucida.
lutea, Bieb. Fl. Taur. Cauc. iii. 255 = Sternbergia Fischeriana.
lutea, Linn. Sp. Pl. 292 = Sternbergia lutea.
lutea, Pav. ex Roem. & Schult. Syst. vii. 797.—*Cf. lutescens*.
lutescens, Herb. ex Kunth, Enum. Pl. v. 497 = Hippeastrum lutescens.
maculata, L'Hérit. Sert. Angl. 10 (= *Habranthus maculatus*).
maranensis, Ker-Gawl. in Bot. Reg. t. 719 = Hippeastrum stylosum.
marginata, Jacq. Hort. Schoenb. i. 34 = Nerine marginata.
mesochlora, Steud. Nom. ed. II. i. 71 = Zephyranthes mesochlora.
mexicana, Hort. ex Savi, Fl. Ital. iii. 29. t. 94 = Vallota purpurea.
miniata, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 346 = Hippeastrum rutilum.
miniata, Ruiz & Pav. Fl. Per. iii. 57 = Hippeastrum miniatum.
minima, Griseb. in Goett. Abh. xxiv. (1879) 321 = Zephyranthes minima.
minima, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 350; Rev. 8 = Zephyranthes verecunda.
minor, D. Dietr. Syn. Pl. ii. 1180 = Brunsvigia minor.
minuta, H. B. & K. Nov. Gen. et Sp. i. 278 = Zephyranthes verecunda.
mollissima, Ker-Gawl. in Journ. Sc. & Arts, iii. (1817) 109 = Crinum latifolium.

AMARYLLIS :—

montana, Labill. Ic. Pl. Syr. Dec. ii. 5 = Ixiolirion montanum.
multiflora, D. Dietr. Syn. Pl. ii. 1180 = Brunsvigia gigantea.
multiflora, Tratt. Gartenpfl. 37 = Haemanthus multiflorus.
narcissa, H. B. & K. Nov. Gen. et Sp. i. 278 = Zephyranthes tubispatha.
narcissa, Pohr. Encyc. Suppl. i. 321 = Brunsvigia striata.
nivea, Schult. f. Syst. vii. 799 = Zephyranthes candida.
nobilis, Salisb. Prod. 285 = Brunsvigia gigantea.
ochroleuca, Ker-Gawl. ex Schult. f. Syst. vii. 797 = Zephyranthes chlorolenca.
orientalis, Linn. Sp. Pl. 293 = Brunsvigia gigantea.
ornata, Ait. Hort. Kew. i. 418 = Crinum yuccæflorum.
ornata, Ker-Gawl. in Bot. Mag. t. 1171 = Crinum zeylanicum.
Oshoshii, Sabine, ex Desf. Cat. ed. III. 386 = Crinum Oshoshii.
ovigera, Schult. f. Syst. vii. 840 = Pellosanthes javanica!
pallida, [Del.] Red. Liliac. viii. 679 = A. Belladonna.
palmis, Willd. ex Schult. f. Syst. vii. 801 = Zephyranthes verecunda.
pardina, Hook. f. Bot. Mag. t. 5645 = Hippeastrum pardinum.
parvula, Seub. in Mart. Fl. Bras. iii. 145 = Zephyranthes minima.
pendula, Salisb. Prod. 281 = Cyrtanthus obliquus.
peruviana, Ker-Gawl. in Bot. Mag. sub t. 1089 (= *Pyrolirion peruvianum*).
phycelloides, Steud. Nom. ed. II. i. 72 = Hippeastrum phycelloides.
platypetala, Lindl. ex Bury, Hexand. Pl. 15 = Lycoris aurea.
praclara, Sweet, Hort. Brit. ed. II. 506 = Hippeastrum reticulatum.
Princeps, Vell. Fl. Flum. 130; iii. t. 115 = Hippeastrum reticulatum.
Principis, Salm-Dyck, in Nov. Act. Nat. Cur. x. (1821) 134 = Hippeastrum reticulatum.
procera, Duchart. in Bull. Soc. Bot. Fr. x. (1863) 75 = Hippeastrum proceræum.
procera, Salisb. Prod. 280 = Crinum erubescens.
propinqua, *Salisb. l. c.*—Panama.
psittacina, Ker-Gawl. in Bot. Reg. t. 190 = Hippeastrum psittacinum.
pudica, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 348 = A. Belladonna.
pulchella, Salisb. Prod. 229 = Zephyranthes Atamasco.
pulchella, Spreng. ex Kunth, Enum. Pl. ii. 683 = Hessea crispa.
pulchra, Salisb. Prod. 283 = Nerine sarniensis.
pulverulenta, Lodd. Bot. Cab. t. 434 = Hippeastrum rutilum.
pumila, Spin. Jard. Sébast. 28; ex Spreng. Neue Entdeck. i. 252 = Zephyranthes pumila.
Pumilio, Ait. Hort. Kew. i. 415 = Cyrtanthus uniflorus.
punicea, Lam. Encyc. i. 123 = Hippeastrum equestre.
purpurea, Ait. Hort. Kew. i. 417 = Vallota purpurea.
radiata, L'Hérit. Sert. Angl. 12 = Lycoris radiata.
Radula, Jacq. Hort. Schoenb. i. 35 = Brunsvigia Radula.
Rayneri, Hook. Bot. Mag. t. 5653 = Hippeastrum proceræum.
regalis, Salisb. Prod. 282 = A. Belladonna.
Reginæ, Linn. Spec. Nat. ed. X. ii. 977 = Hippeastrum Reginæ.
reticulata, L'Hérit. Sert. Angl. 12 = Hippeastrum reticulatum.
revoluta, L'Hérit. l. c. 13 = Crinum bisetosum.
riparia, Burch. ex Kunth, Enum. Pl. ii. 680 = Crinum longifolium.
robusta, A. Dietr. in Otto & Dietr. Allg. Gartz. xviii. (1850) 41 = Hippeastrum sulcatum.
rubens, Sweet, ex Steud. Nom. ed. II. i. 74 (= *Habrantus robustus*).
Ruiziі, Regel, Gartenfl. (1874) 290. f. 800 = Hippeastrum equestre.
rosea, Lam. Encyc. i. 123 = A. Belladonna.
rosea, Spreng. Syst. iv. Cur. Post. 138 = Zephyranthes rosea.
rotundifolia, Lam. Encyc. i. 124 = Eurycles sylvestris.
Rouyieri, Hort. ex Carr. in Rev. Hortic. (1869) 312 = Hippeastrum sulcatum.
rutila, Ker-Gawl. Bot. 26; in Bot. Reg. t. t. 28 = Hippeastrum rutilum.

AMARYLLIS:—
rutilans, Lodd. ex Steud. Nom. ed. II. i. 74 = Hippeastrum rutilum.
sarniensis, Linn. Sp. Pl. 293 = Nerine sarniensis.
scillifolia, Hort. Angl. ex Schult. f. Syst. vii. 854 = Crinum distichum.
semiherbata, Steud. Nom. ed. II. i. 72, sphalm. (*loco* Antherici) = Bulbine bulbosa.
Slateriana, Lindl. Bot. Reg. (1844) App. 76 = Brunsvigia Slateriana.
solandraeflora, Lindl. Coll. Bot. t. 11 = Hippeastrum solandraeflorum.
spathacea, Schult. f. Syst. vii. 808 (= *Habranthus spathaceus*).—Bonaria.
speciosa, L'Hérit. Sert. Angl. 12 = Vallota purpurea.
spectabilis, Andr. Bot. Rep. t. 390 = Crinum yuccaefloreum.
spectabilis, Lodd. Bot. Cab. t. 159 = Hippeastrum Reginae.
spiralis, L'Hérit. Sert. Angl. 10 = Hessea spiralis.
staminea, Sensb. in Mart. Fl. Bras. iii. 150 = Hippeastrum stylosum.
stellaris, Jacq. Hort. Schoenb. i. 27 = Hessea stellaris.
stilosa, Bury, Hexand. Pl. t. 33 = Hippeastrum stylosum.
striata, Jacq. Hort. Schoenb. i. 36 = Brunsvigia striata.
striata, Lam. Encyc. i. 135 = Hippeastrum rutilum.
striatula, Schult. f. Syst. vii. 801 = Zephyranthes verecunda.
stylosa, Sweet, Hort. Brit. ed. I. 408 = Hippeastrum stylosum.
subherbata, Schult. f. Syst. vii. 819 = Hippeastrum rutilum.
superba, Bury, Hexand. Pl. t. 31 = Hippeastrum vittatum.
Swertiana, Steud. Nom. ed. II. i. 72 = Hippeastrum adoenum.
Swertii, Herb. ex Sweet, Hort. Brit. ed. II. 506 = Hippeastrum reticulatum.
sylvatica, Mart. in Schult. f. Syst. vii. 807 (= *Habranthus sylvaticus*).
Symoni, *Punch, in Yourn. Hort. Prat. Scr.* II. iii. (1859) 349. t. 20.—Hab.?
tatarica, Georgi, Beschreib. Russ. Reich. iii. iv. 890 = Ixiolirion Ledebouri.
tatarica, Pall. Reise, iii. 727 = Ixiolirion tataricum.
tenella, Thunb. ex Steud. Nom. ed. II. i. 72 = Carpolyza spiralis.
toxicaria, D. Dietr. Syn. Pl. ii. 1181 = Brunsvigia toxicaria.
tubiflora, L'Hérit. Sert. Angl. 10 (= *Pyrolirion aureum*). —Peruv.
tubispatha, L'Hérit. l. c. 9 = Zephyranthes tubispatha.
Umbella, L'Hérit. l. c. 15 = Cyrtanthus obliquus.
undulata, Linn. Syst. ed. XII. 287 = Nerine undulata.
unguiculata, Mart. ex Schult. f. Syst. vii. 789 = Hippeastrum calyptratum.
vaginata, D. Dietr. Syn. Pl. ii. 1181 = Strumaria truncata.
valparadisiaca, Steud. Nom. ed. II. i. 188 (= *Habranthus pallidus*).—Chili.
variabilis, Jacq. Hort. Schoenb. iv. 14 = Crinum variabile.
venusta, Ker-Gawl. in Bot. Mag. t. 1090 = Nerine sarniensis.
verecunda, Salisb. Prod. 229 = Zephyranthes Atamasco.
verecunda, Schult. f. Syst. vii. 800 = Zephyranthes verecunda.
vernalis, Mill. Gard. Dict. ed. VIII. n. 10 = Hypoxis stellata.
Verrazzii, *Hort.—Cf. Journ. et Flore des Jardins*, ii. (1882) t. 119.—Hab.?
versicolor, Spreng. Syst. Cur. Post. 128 (= *Habranthus versicolor*).
vittata, L'Hérit. Sert. Angl. 13 = Hippeastrum vittatum.
viviparа, Lam. Encyc. i. 128 = Crinum defixum.
yuccaeoides, Thunbg. Bot. Displ. t. 12 = Crinum yuccaefloreum.
zeylanica, Linn. Sp. Pl. 293 = Crinum zeylanicum.

AMARYLLIS, Sweet, Hort. Brit. ed. II. 506 (1830) = **Hippeastrum**, Herb.

AMASONIA, Linn. f. Suppl. 48, 294 (1781). *VERBENACEAE*. Benth. & Hook. f. ii. 1147.
Dipnystema, Neck. Elem. i. 262 (1790).
Diplostemma, 'Neck.' ex DC. Prod. vii. 506. in nota (1838) lapsu.
Talgoalba, Astli. Pl. Gui. ii. 235, t. 292 (1776).

AMASONIA:—
angustifolia, *Mart. & Schau. in DC. Prod.* xi. 678.—Bras.
arborea, *H. B. & K. Nov. Gen. et Sp.* ii. 268.—Venezuela.
erecta, *Linn. f. Suppl.* 294.—Guiana.
hirta, *Benth. in Ann. Nat. Hist.* ii. (1839) 451.—Guiana.
integerrima, *Spreng. Syst.* ii. 765.—N. Granat.
lasiocaulos, *Mart. & Schau. in DC. Prod.* xi. 678.—Am. austr.
punices, *Vahl, Eclog.* ii. 51.—Ins. Trinit.
Talgalea, Anct. ex Steud. Nom. ed. II. i. 74 = erecta.
velutina, *Schau. in DC. Prod.* xi. 677.—Bras.

AMATHEA, Rafin. Fl. Tellur. iv. 65 (1836) = **Aphelandra**, R. Br. (Acanth.).
pulchra, Rafin. l. c. = Aph. pulcherrima.

AMATULA, Medic. Bot. Beobacht 38 (1782) ; Malv. 106 (1787) = **Lycopersicum**, Mill. (Solan.).
Acna, Medic. Malv. 106 = L. esculentum.
rubra, Medic. l. c. = L. esculentum.

AMAURIA, Benth. Bot. Voy. Sulph. 81 (1844). *COMPOSITAE*. Benth. & Hook. f. ii. 404.
dissecta, *A. Gray, Pl. Fendl.* 104.—N. Mexic.
rotundifolia, *Benth. Bot. Voy. Sulph.* 81.—Calif.

AMAXITIS, Adans. Fam. ii. 34 (1763) = **Dactylis**, Linn. (Gramin.).

AMBAIBA, Adans. l. c. 377 (1763) = **Cecropia** Loefl. (Urtic.).

AMBASSA, Steetz, in Peters, Reise Mossamb. Bot. 364 (1863) = **Vernonia**, Schreb. (Compos.).
Hochstetteri, Steetz, l. c. = V. Hochstetteri.

AMBELANIA, Aubl. Pl. Gui. i. 265. t. 104 (1775). *APOCYNACEAE*. Benth. & Hook. f. ii. 694.
Bentuca, Adans. Fam. ii. 168 (1763).
Kasalio, Denst. Schluss. Hort. Malab. 30 (1818).
Willughbeia, Scop. in Schreb. Gen. 162 (1789).
acida, *Aubl. Pl. Gui.* i. 265. t. 104.—Guiana.
cuneata, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 17.—Bras.
laxa, *Muell. Arg. l. c.*—Am. austr.
macrophylla, *Muell. Arg. l. c.* 18.—Bras.
quadrangularis, *Muell. Arg. l. c.*—Bras.
Sagoti, *Muell. Arg. in Linnaea*, xxx. (1859–60) 389. —Guiana.
tenuiflora, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 16. —Bras.

AMBERBOA, Less. Syn. Comp. 8 (1832) = **Volutarella**, Cass. etc. (Compos.).
abyssinica, A. Rich. Tent. Fl. Abyss. i. 453 = Centaurea Hochstetteri.
Bulbi, Sch. Bip. ex C. Bolle, in Bonplandia, vii. (1859) 297.—Ins. Canar.
carthamoides, DC. Prod. vi. 561 (= *Phaetappus carthamoides*).
crupinoides, DC. l. c. 559 = V. bicolor.
decurrens, DC. l. c. 560 = Centaurea decurrens.
discolor, Lojac. in Natural. Sicil. iv. (1886) 183.—Sicil.
eriocephala, DC. Prod. vi. 559 = V. eriolepis.
glauca, Less. Syn. Comp. 8 = Centaurea moschata ?
Goniocaulon, DC. Prod vi. 558 = Goniocaulon glabrum.
Hochstetteri, Buching. ex Oliver, Fl. Trop. Afr. iii. 438 = Centaurea Hochstetteri.
incanescens, DC. Prod. vi. 561 = Centaurea integrifolia.
indica, DC. l. c. 558 = Goniocaulon glabrum.
lanceifolia, DC. l. c. 561 = Centaurea lanceifolia.
leucantha, Coss. Ind. Fl. Maroc. 21.—Algeria.
Lippii, DC. Prod. vi. 559 = V. Lippii.
microcephala, Boiss. & Rent. Diagn. Ser. II. iii. 69.—Algeria.
moschata, Less. Syn. Comp. 8 = Centaurea moschata.
muricata, DC. Prod. vi. 559 = V. muricata.
odorata, DC. l. c. = Centaurea moschata.
Perralderiana, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 62, nomen.—Afr. bor.
phaeopappa, DC. Prod. vi. 560 = Centaurea leucoidea.
sinuata, DC. l. c. 559 = V. bicolor.
spectabilis, DC. l. c. 561 (= *Phaeopappus spectabilis*).
Wallichiana, DC. l. c. 558 = Centaurea Wallichiana.
xanthocephala, DC. l. c. 561 (= *Psephellus xanthocephalus*).

AMBERBOI, Adans. Fam. ii. 117 (1763) = AMBERBOA.

AMBINAX, Comm. ex Juss. Gen. 389 (1789) = **Aleurites**, Forst. (Euphorb.).

AMBLACHAENIUM, Turcz. ex DC. Prod. vii. 94 (1838) = **Erysochoeris**, Linn. (Compos.).
auranthacum, Turcz. ex DC. l. c. = H. grandiflora.

AMBLATUM, (Tourn.) G. Don, Gen. Syst. iv. 635 (1837) = ANBLATUM, Tourn. = **Lathraea**, Linn. (Orobanch.).

AMBLEIA, Spach, Hist. Vég. Phan. ix. 166 (1840), nomen = **Stachys**, Linn. (Labiat.).

AMBLIRION, Rafin. in Am. Monthly Mag. (1818) 265 = **Fritillaria**, Linn. (Liliac.).
album, Sweet, Hort. Brit. ed. II. 538 = F. alba.
camtschatcense, Sweet, l. c. = F. kamtschatcensis.
gracum, Rafin. Fl. Tellur. iii. 85 = F. graecum ?
lanceolatum, Sweet, l. c. = F. kamtschatcensis.
pudicum, Rafin. in Am. Monthly Mag. (1818) 265 = F. pudica.

AMBLOGYNA, Rafin. Fl. Tellur. iii. 42 (1836) = **Amaranthus**, Linn.
fimbriata, A. Gray, in Proc. Am. Acad. v. (1861) 167, 168 = Amar. fimbriatus.
polygonoides, Dalz. & Gibs. Bomb. Fl. 219 = Amar. polygamus
polygonoides, Rafin. Fl. Tellur. iii. 42 = Amar. polygonoides.
scariosa, Moq. in DC. Prod. xiii. ii. 269 ; A. Gray, in Proc. Am. Acad. v. (1861) 169 = Amar. scariosus.
squarrulosa, A. Gray, ; c.—Ins. Galap.
Torreyi, A. Gray, l. c. 167, 169 = Amar. Torreyi
urceolata, Moq. in DC. Prod. xiii. ii. 269 ; A. Gray, l. c. 168 = Amar. urceolatus.

AMBLOSTIMA, Rafin. Fl. Tellur. ii. 26 (1836) = **Schoenolirion**, Torr. (Liliac.).
albiflora, Rafin. l. c. = Sc. Elliottii.
crocea, Rafin. l. c. = Sc. croceum.
latifolia, Rafin. ex S. Wats. in Proc. Am. Acad. xiv. (1879) 242 = Sc. Elliottii.

AMBLOSTOMA, Schreb. in Otto & Dietr. Gartenz. vi. (1888) 388. *ORCHIDEAE*. Benth. & Hook. f. iii. 522.
cernuum, *Schreb. l. c.*—Bras.
densum, *Reichb. f. Xenia Orch.* iii. 22.—Peruv.
micranthum, *Reichb. f. in Saund. Refug. sub* t. 101. —Boliv.
tridactylum, Reichb. f. in Walp. Ann. vi. 485 = cernuum.

AMBLYACHYRUM, Hochst. ex Steud. Syn. Pl. Gram. 413 (1855) = **Apocopis**, Nees (Gramin.).
mangalorense, Hochst. ex Steud. l. c. ; et in Flora, xxxix. (1856) 26 = Apoc. Wightii.

AMBLYANTHERA, Blume, Mus. Bot. Lugd. Bat. i. 50 (1849) = **Osbeckia**, Linn. (Melast.).
parvifolia, Blume, l. c. = O. parvifolia.
truncata, Blume, l. c. = O. truncata.

AMBLYANTHERA, Muell. Arg. in Mart. Fl. Bras. vi. i. 141, t. 44, 46 (1869) = **Mandevilla**, Lindl. (Apocyn.).
acutiloba, Muell. Arg. in Linnaea, xxx. (1859–60) 426 = Echites acutiloba.
andina, Muell. Arg. l. c. 425 = Echites andina.
Andrieuxii, Muell. Arg. l. c. 422 = Mandevilla Andrieuxii.
angustifolia, Muell. Arg. l. c. 430 = Echites angustifolia.
antennacea, Muell. Arg. l. c. 448 = Echites antennacea.
Benthami, Muell. Arg. l. c. 451 = Echites Benthami.
bogotensis, Muell. Arg. l. c. 452 = Echites bogotensis.
brachyloba, Muell. Arg. l. c. 423 = Echites brachyloba.
Bridgesii, Muell. Arg. l. c. 430.—Boliv.
campestris, Muell. Arg. in Mart. Fl. Bras. vi. i. 149 = Echites campestris.
ciliata, Muell. Arg. l. c. 146 = Echites ciliata.
Clausenii, Miers, Apocyn. 8. Am. 274, t. 26.—Am. austr.
convolvulacea, Muell. Arg. in Linnaea, xxx. (1859–60.) 430 = M. convolvulacea.

AMBLYANTHERA :—
crassifolia, Muell. Arg. in Mart. Fl. Bras. vi. 1. 143 = Echites crassifolia.
cuyabensis, Muell. Arg. l. c. 145 = Echites cuyabensis.
cuyabensis, Muell. Arg. in Linnæa, xxx. (1859–60) 450 = prec.
Fendleri, Muell. Arg. l. c. 417.—Venezuela.
foliola, Muell. Arg. l. c. 427 = Mandevilla foliosa.
fluminensis, Muell. Arg. in Mart. Fl. Bras. vi. 1. 148 = Echites fluminensis.
funiformis, Muell. Arg. l. c. 144 = Echites funiformis.
guyanensis, Muell. Arg. in Linnæa, xxx. (1859–60) 448 = Echites guianensis.
hirsuta, Miers, Apocyn. S. Am 185 = M. hispida ?
hispida, Muell. Arg. in Mart. Fl. Bras. vi. 1. 147 = M. hispida.
Karwinskii, Muell. Arg. in Linnæa, xxx. (1859–60) 426 = M. Karwinskii.
laxicarpa, Muell. Arg. in Kjoeb. Vidensk. Meddel. (1866) 113 = M. hispida.
leptophylla, Muell. Arg. in Mart. Fl. Bras. vi. 1. 142 = Echites leptophylla.
macrophylla, Muell. Arg. in Linnæa, xxx. (1859–60) 445 = Echites macrophylla.
madida, Muell. Arg. in Mart. Fl. Bras. vi. 1. 150 = Echites madida.
membranacea, Muell. Arg. in Linnæa, xxx. (1859–60) 428, in nota = Echites membranacea.
mexicana, Muell. Arg. l. c. 424 = Echites mexicana.
microcalyx, Muell. Arg. l. c. 428 = Echites microcalyx.
Moritziana, Muell. Arg. l. c. 431.—N. Granat.
ovata, Miers, Apocyn. S. Am. 188.—Boliv.
oaxacana, Muell. Arg. in Linnæa, xxx. (1859–60) 447 = M. oaxacana.
palustris, Muell. Arg. in Mart. Fl. Bras. vi. 1. 145 = M. palustris.
pandurata, Muell. Arg. in Linnæa, xxx. (1859–60) 448 = Echites pandurata.
Pavonii, Muell. Arg. l. c. 450 = Echites Pavonii.
Priceuri, Muell. Arg. l. c. 448 = Echites Priceuri.
Schiimi, Muell. Arg. l. c. 419.—N. Granat.
Spruceana, Muell. Arg. in Mart. Fl. Bras. vi. 1. 143.—Bras.
suaveolens, Muell. Arg. in Linnæa, xxx. (1859–60) 447 = M. suaveolens.
subsessilis, Muell. Arg. l. c. = Echites subsessilis.
torosa, Muell. Arg. l. c. 446 = Echites torulosa.
tubiflora, Muell. Arg. l. c. 448 = Echites tubiflora.
versicolor, Muell. Arg. in Mart. Fl. Bras. vi. 1. 146 = Echites versicolor.

AMBLYANTHUS, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 83. t. 6. *MYRSINEAE*, Benth. & Hook. f. ii. 648.
glandulosus, *A. DC. l. c.*—Reg. Himal.

AMBLYCARPUM, Lem. in Orb. Dict. i. 335 (1841) = **Amblyocarpum**, Fisch. & Mey. (Compos.).

AMBLYCHLOA, Link, in Linnæa, xvii. (1843) 399 = **Solerochloa**, Besv. (Gramin.).
dura, Link, l. c. = S. dura.

AMBLYGLOTTIS, Blume, Bijdr. 369. t. 64 (1825) = **Calanthe**, R. Br. (Orchid.).
abbreviata, Blume, l. c. 371 = C. abbreviata.
angustifolia, Blume, l. c. 369 = C. angustifolia.
emarginata, Blume, l. c. 370 = C. emarginata.
flava, Blume, l. c. t. 64 = C. veratrifolia.
pilosa, De Vriese, ex Lindl. Fol. Orch. Calanthe, 11 = C. pilosa.
pulchra, Blume, Bijdr. 371 = C. carculigoides.
speciosa, Blume, l. c. 371 = C. speciosa.
veratrifolia, Blume, l. c. 370 = C. veratrifolia.

AMBLYGONUM, Reichb. Handb. 236 (1837) = **Polygonum**, Linn.

AMBLYLEPIS, Decne. in Orb. Dict. i. 337 (1841) = **Amblyolepis**.

AMBLYOCALYX, Benth. in Benth. & Hook. f. Gen. ii. 698 (1876). *APOCYNACEAE*, Benth. & Hook. f. ii. 693.
Beccarii, *Benth. in Hook. Ic. Pl.* t. 1179.—Borneo.

AMBLYOCARPUM, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 30 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 336.
inuloides, *Fisch. & Mey. l. c.*—Reg. Caspica.

AMBLYOGLOSSUM, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 310 = **Tylophora**, R. Br. (Asclep.).
brevipes, Turcz. l. c. 311 = T. brevipes.
longipes, Turcz. l. c.—Malaya.

AMBLYOLEPIS, DC. Prod. v. 667 (1836) = **Helenium**, Linn. (Compos.).
setigera, DC. l. c.—Mexic.

AMBLYOPAPPUS, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 231. *COMPOSITAE*, Benth. & Hook. f. ii. 406.
AROMIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 395.
INFANTEA, Remy, in C. Gay, Fl. Chil. vi. 257. t. 48 (1849).
mendocinus, *Phil. Sert. Mend. Alt.* 26.—Chili.
neo-mexicanus, *A. Gray, in Pacif. Rail. Rep.* iv. 106.—Mexic.
pusillus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 231.—Chili.

AMBLYOPELIS, Stend. Nom. ed. II. i. 75 (1840), sphalm. = AMBLYOLEPIS, DC.

AMBLYOPOGON, Fisch. & Mey. ex DC. Prod. vi. 561 (1837) = **Centaurea**, Tourn. (Compos.).
Balansae, Boiss. Fl. Orient. iii. 604, in syn. = C. Balansae.
concinnus, Boiss. & Huet, Diagn. Ser. II. iii. 65 = C. pulchellus.
sincanescens, Fisch. & Mey. ex DC. Prod. vi. 561 = C. integrifolia.
speciosus, Boiss. Fl. Orient. iii. 604, in syn. = C. pulcherrima.

AMBLYORHINUM, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 168 = **Phyllactis**, Pers. (Valerian.).
angustifolium, Turcz. l. c. 169.—Venezuela.
grandiflorum, Turcz. l. c. 168.—Venezuela.
phyllactoides, Turcz. l. c. 171.—Venezuela.
spicatum, Turcz. l. c. 170 = P. cordifolia.
triplinervum, Turcz. l. c.—Venezuela.

AMBLYSPERMA, Benth. in Enum Pl. Huegg. 67 (1837) = **Trichoolina**, Cass. (Compos.).
scapigerum, Benth. l. c. = T. scapigera.

AMBLYSTIGMA, Benth. in Benth. & Hook. f. Gen. ii. 745 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 748.
cionophorum, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 366.—Reg. Argent.
hypoleucum, *Benth. in Hook. Ic. Pl.* xii. 76. t. 1188.—Bolivia.
pedunculare, *Benth. l. c.* 77. sub t. 1188.—Bolivia.

AMBLYTES, Dulac, Fl. Hautes-Pyr. 80 (1867) = **Molinia**, Scop. (Gramin.).
caerulea, Dulac, l. c. = M. caerulea.

AMBORA, Juss. Gen. 401 (1789) = **Tambourissa**, Somm. (Monim.).
alternifolia, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 31. = T. alternifolia.
amplifolia, Tul. in Arch. Mus. Par. viii. (1855–56) 299 = T. amplifolia.
cymosa, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 31 (an Robiaceæ ?).—Hab. ?
elliptica, Tul. l. c. = T. elliptica.
Ficus, Tul. l. c. 30 = T. Ficus.
frutescens, Bory ex Tul. l. c. = T. purpurea.
grandifolia, R. Br. ex Baker, Fl. Maurit. 257 = T. amplifolia.
leptophylla, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 29 = T. leptophylla.
neglecta, Tul. in Arch. Mus. Par. viii. (1855–56) 282 = T. neglecta.
obovata, Tul. in Ann. Sc. Nat. Sér. IV. t. iii. (1855) 31 = T. obovata.
peltata, R. Br. ex Baker, Fl. Maurit. 258 = T. peltata.
purpurea, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 30 = T. purpurea.
quadrifida, Poir. Encyc. vii. 565 = T. quadrifida.
racemosa, R. Br. ex Baker, Fl. Maurit. 258 = T. quadrifida.
religiosa, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 30 = T. religiosa.

AMBORA :—
sempervirens, F. G. Dietr. Lexik. Gaertn. i. 342 = T. quadrifida.
Sieberi, Tul. in Arch. Mus. Par. viii. (1855–56) 304 = T. Sieberi.
Tambourissa, Lam. ex Steud. Nom. ed. I. 37, 535 = T. quadrifida.
Tambourissa, Tul. in Arch. Mus. Par. viii. (1855–56) 302 = T. Bolvinii.
tetragona, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 30 = T. tetragona.
tomentosa, Bory, Voy. I. 317. t. 18 = Monimia rotundifolia.
tomentosa, Sieber, ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 31 = T. Sieberi.
vestita, Tul. l. c. = T. vestita.

AMBORELLA, Baill. Hist. Pl. i. 328 in nota (1869); et Adansonia, x. 354 (1878). *MONIMIACEAE*, Benth. & Hook. f. iii. 143.
trichopoda, *Baill. l. c.*—N. Caled.

AMBRARIA, Cruse, Rub. Cap. 16. t. 1, f. 3, 4 (1825) = **Nenax**, Gaertn. (Rubiac.).
acerosa, Sond. in Harv. & Sond. Fl. Cap. 34 = N. acerosa.
glabra, Cruse, Rub. Cap. 16. t. 1 = N. acerosa.
hirta, Cruse, l. c.—Afr. austr.
microphylla, Sond. in Harv. & Sond. Fl. Cap. 34.—Afr. austr.

AMBRARIA, Heist. Syst. 11 (1748) = **Anthospermum**, Linn. (Rubiac.).
Heisteri, Willd. Sp. Pl. iv. 782 in syn. = Anth. aethiopicum.

AMBRINA, Spach, Hist. Vég. Phan. v. 295 (1836) = **Chenopodium**, Linn.
ambrosioides, Spach, l. c. 297 = C. ambrosioides.
anthelmintica, Spach, l. c. 298 = C. ambrosioides.
Botrys, Moq. Chen. Enum. 37 = C. Botrys.
carinata, Moq. l. c. 41 = C. carinatum.
chilensis, Spach, Hist. Vég. Phan. v. 297 = C. chilense.
denudata, Phil. in Linnæa, xxix. (1857–58) 87 = Roubieva multifida.
dissecta, Moq. Chen. Enum. 38 = C. bipinnatifidum.
foetida, Moq. l. c. 38 = C. Botrys.
graveolens, Moq. l. c. 41 = C. graveolens.
incisa, Moq. l. c. 36 = C. incisum.
obovata, Moq. l. c. 40 = C. obovatum.
pinnatiscta, Spach, Hist. Vég. Phan. v. 296 = Roubieva multifida.
Pumilio, Moq. Chen. Enum. 42 = C. Pumilio.
retusa, Moq. l. c. 40 = C. retusa.
spathulata, Moq. l. c. 39 = C. spathulatum.
tenuis, Moq. l. c. 42 = Roubieva multifida.
tomentosa, Moq. l. c. 37 = C. tomentosum.

AMBROMA, Linn. f. Suppl. 84, 341 (1781) = **Abroma**, Jacq. (Stercul.).
Augusta, Linn. f. l. c. 341 = Abr. fastuosum.

AMBROSIA, Basal. in Comm. Bonon. v. (1768) 82 = **Ambrosinia**, Linn. (Aroid.).

AMBROSIA, Linn. Gen. ed. I. 287. n. 718 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 354.
abrotanifolia, Michx. Fl. Bor. Am. ii. 138 = artemisiaefolia.
acanthicarpa, Hook. Fl. Bor. Am. i. 309 = Franseria Hookeriana.
aptera, *DC. Prod.* v. 537.—Am. bor.
arborescens, Lam. Encyc. I. 127 = Franseria artemisioides.
arborescens, *Mill. Gard. Dict.* ed. VIII. n. 5.—Peruv.
artemisiaefolia, *Meyen & Walp. in Nov. Act. Nat. Cur. xix. Suppl.* i. (1843) 268.—Peruv.
artemisiaefolia, Bess. Prim. Fl. Galic. ii. 290 = maritima.
artemisiifolia, *Linn. Sp. Pl.* 988.—Am. bor.
bidentata, *Michx. Fl. Bor. Am.* ii. 138.—Am. bor.; Ind. occ.
cheiranthifolia, *A. Gray, in Torr. Bot. Mex. Bound.* 87.—Mexic.
chilensis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1840) 211.—Chili.
confertiflora, DC. Prod. v. 536 = Franseria tenuifolia.
coronopifolia, Torr. & Gray, Fl. N. Am. ii. 471 = palmatifida.
crithmifolia, DC. Prod. v. 535 = hispida.

AMBROSIA :—
cumanensis, *H. B. & K. Nov. Gen. et Sp.* iv. 276.—Venezuela.
elata, Salisb. Prod. 175 = artemisiifolia.
elatior, Bess. Prim. Fl. Galic. ii. 389 = maritima.
elatior, Linn. Sp. Pl. 987 = artemisiifolia.
frutticosa, DC. Prod. v. 535 = Franseria tenuifolia.
frutticosa, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 247 = arborescens.
giandulosa, Scheele, in Linnæa, xxii. (1849) 157 = pallostachya.
heterophylla, Muhl. ex Willd. Sp. Pl. iv. 377 = artemisiifolia.
hispida, *Pursh, Fl. Am. Sept. Suppl.* 743.—Florida.
hispida, Torr. in Ann. Lyc. N. York, ii. (1828) 216 = pallostachya.
humilis, *Vell. Fl. Flum.* x. t. 35.—Bras.
incisa, F. G. Dietr. Lexik. Gaertn. i. 844 = artemisiifolia.
integrifolia, Muhl. ex Willd. Sp. Pl. iii. 375 = trifida.
Lindheimeriana, Scheele, in Linnæa, xxii. (1849) 156 = pallostachya.
longistylis, A. Gray, Pl. Fendler. 79 = Franseria tenuifolia.
longistylis, *mult. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 344.—Am. bor.
maritima, Ferrero, ex DC. Prod. v. 525 = crithmifolia.
maritima, *Linn. Sp. Pl.* 988.—Europ.; Orient; Afr. bor.
maritima, Vell. Fl. Flum. x. t. 36 = polystachya.
microcephala, *DC. Prod.* v. 527.—Guiana.
orobanchifera, *Meyen, Reise*, i. 444.—Peruv.
paniculata, Michx. Fl. Bor. Am. ii. 183 = artemisiifolia.
peruviana, DC. Prod. v. 526 = pallostachya.
peruviana, Willd. Sp. Pl. iv. 377.—Peruv.
Pitcheri, Torr. ex Hook. Comp. Bot. Mag. i. (1835) 99 = Iva ciliata.
polystachya, *DC. Prod.* v. 526.—Bras.
pallostachya, *DC. l. c.*—Am. bor.
pumila, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 217.—Am. bor.
psilila, Rafin. ex DC. Prod. v. 527 = artemisiifolia.
scabra, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1840) 311.—Bras.
senegalensis, DC. Prod. v. 525 = maritima.
simplicifolia, Rasnosh. Nom. ed. III. 374 = artemisiifolia.
tacorensis, *Meyen, Reise*, i. 456.—Peruv.
tenuifolia, Gren. & Godr. Fl. Fr. ii. 395 = Franseria tenuifolia.
tenuifolia, Spreng. Syst. iii. 851 = polystachya ?
tenuifolia, Nutt. Gen. Am. ii. 186 = Franseria discolor.
trifida, *Linn. Sp. Pl.* 987.—Am. bor.
umbellata, Moench, Meth. Suppl. 272 = maritima.
villosissima, *Forsk. Fl. Aegypt. Arab.* 160.—Arabia.

AMBROSINIA, Linn. Gen. ed. VI. 517 (1764) = *AROIDEAE*, Benth. & Hook. f. iii. 994.
AMBROSIA, Bassi, in Comm. Bonon. v. (1763) 82.
Bassii, *Linn. Gen.* ed. VI. 517.—Europ.
ciliaris, Spreng. Syst. iii. 771 = seq.
ciliata, Roxb. Fl. Corom. iii. 90 = Cryptocoryne ciliata.
maculata, Usteri, in Roem. Arch. i. 70 = A. Bassii.
nervosa, Lam. Encyc. i. 128 = A. Bassii.
reticulata, Tin. Cat. Hort. Panorm. 94, 176, sec. Guss. Fl. Sic. Syn. ii. 11. 896 = A. Bassii.
retrospiralis, Roxb. Hort. Beng. 65 = Cryptocoryne retrospiralis.
Rumphiana, Volgt, Hort. Suburb. Calc. 685 = Cryptocoryne undulocilaris.
spiralis, Roxb. Hort. Beng. 65 = Cryptocoryne spiralis.
undulocilaris, Roxb. Fl. Ind. iii. 498 = Cryptocoryne undulocilaris.
velutina, Blume, ex Hoyuh. Nom. ii. 82 = A. Bassii.

AMBULI, Adans. Fam. ii. 308 (1763) = seq.

AMBULIA, Lam. Encyc. i. 128 (1783) = **Limnophila**, R. Br. (Scrophul.).
alata, Rafin. Fl. Tellur. iv. 56 = Gratiola tetragona.
aromatica, Lam. Encyc. i. 126 = L. gratissima.
rigida, Rafin. Fl. Tellur. iv. 56 = Gratiola tetragona.

AMBUYA, Rafin. Fl. Tellur. iv. 68 (1836) = **Aristolochia**, Tourn.
lobiana, Rafin. l. c. = *Ar.* cymbifera.

AMEBIA, Regel, Fl. Nov. Fedsch. 58 (1882) err. typ. = **Arnebia**, Forsk. (Borragin.).

AMECARPUS, Benth. in Lindl. Veg. Kingd. 554 (1847) = **Indigofera**, Linn. (Legumin.).

AMECHANIA, DC. Prod. vii. 578 (1839) = **Agarista**, D. Don (Ericac.).
hispidula, DC. l. c. 579 (= *Leucothoë hispidula*).
subcaesarea, DC. l. c. (= *Leucothoë subcaesarea*).

AMELANCHIER, Medic. Phil. Bot. i. 185 (1789).
ROSACEAE, Benth. & Hook. f. i. 626.
ARONIA, Pers. Syn. ii. 39 (1807).
PERAPHYLLUM, Nutt. in Torr. & Gray, Fl. N. Am. i. 474 (1840).
XEROMALON, Rafin. New Fl. Am. iii. 11 (1836).
alnifolia, *Nutt. in Journ. Acad. Phil.* vii. (1834) 22.—Am. bor.
asiatica, Endl. in Walp. Rep. ii. 55 = canadensis.
Bartramiana, M. Roem. Syn. Rosifl. 145 = canadensis.
Botryapium, DC. Prod. ii. 632 = canadensis.
canadensis, *Medic. Gesch.* 79; *Torr. & Gray, Fl. N. Am.* i. 478.—Am. bor.; As. or.
chinensis, Hort. ex C. Koch, Dendrol. i. 186 = Sorbus arbutifolia.
cretica, DC. Prod. ii. 632 = vulgaris.
denticulata, C. Koch, Dendrol. i. 183 = Cotoneaster denticulata.
florida, Lindl. Bot. Reg. t. 1589 = alnifolia.
grandiflora, Dougl. ex M. Roem. Syn. Rosifl. 145 = canadensis.
integrifolia, Bolss. & Hohen. Diagn. Ser. I. iii. 8 = vulgaris.
intermedia, Spach, Hist. Vég. Phan. ii. 85 = canadensis ?
japonica, Hort. ex C. Koch, Dendrol. i. 179 = canadensis.
melanocarpa, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 186.—Hab. ?
oblongifolia, M. Roem. Syn. Rosifl. 147 = canadensis.
oligocarpa, M. Roem. l. c. 146 = canadensis.
orbicularis, Borck. ex Steud. Nom. ed. II. i. 76 = vulgaris.
ovalis, Medic. Gesch. 78 = canadensis.
parviflora, *Boiss. Diagn.* Ser. I. iii. 9.—As. Min.
parviflora, Hort. ex Loud. Arb. Brit. ii. 877 = alnifolia.
pinifera, Boiss. & Heldr. Diagn. Ser. I. x. 2 = parviflora.
ponila, Nutt. ex Torr. & Gray, Fl. N. Am. i. 474 = alnifolia.
racemosa, Lindl. in Bot. Reg. (1847) sub. t. 88 = Exochorda grandiflora.
rotundifolia, M. Roem. Syn. Rosifl. 146 = canadensis.
sanguinea, DC. Prod. ii. 633 = canadensis ?
spicata, C. Koch, Dendrol. i. 182 = canadensis ?
subcordicularis, Borkh. Forstbot. ii. 1362 = vulgaris.
tomentosa, Steud. Nom. ed. II. ii. 186 in syn. = vulgaris ?
vulgaris, *Moench, Meth.* 682.—Europ.; Orient.
Wangenheimiana, M. Roem. Syn. Rosifl. 146 = canadensis.

AMELETIA, DC. in Mém. Soc. Phys. Genèv. iii. 11. (1826) 82 = **Ammannia**, Houst. (Lythrar.).
acutidens, Miq. Fl. Ind. Bat. i. 1. 617 = Amman. peploides.
chandra, F. Muell. Fragm. iii. 108 = Amman. diandra.
elatinoides, Blume, Mus. Bot. Lugd. Bat. ii. 186 = Amman. elatinoides.
elongata, Blume, l. c. 185 = Amman. peploides.
floribunda, Wight, Illustr. Ind. Bot. 206 = Ammannia floribunda.
indica, DC. in Mém. Soc. Hist. Nat. Genèv. iii. 11. (1825) 2 & 82. t. 3 f. A. = Ammian. peploides.
polystachya, Wall. Cat. n. 2094 = Ammian. peploides.
rotundifolia, Wight, Ic. t. 958 = Ammian. rotundifolia.
subpinnata, Benth. in Hook. Kew Journ. iv. (1842) 81 = Ammian. rotundifolia.
tenella, Blume, Mus. Bot. Lugd. Bat. ii. 186 = Ammian. tenella.
tenuis, Wight, Ic. Pl. Ind. Or. t. 257 B = Ammian. tenuis.
uliginosa, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 261 = Ammian. peploides.

AMELIA, Aiefeld, in Linnæa, xxviii. (1856) 25 = **Pyrola**, Tourn. (Ericac.).

AMELIA :—
media, Alef. in Linnæa, xxviii. (1856) 80 = Pyrola media.
minor, Alef. l. c. 25 = P. minor.

AMELINA, C. B. Clarke, Commel. et Cyrt. Beng. 88. t. 26 (1874) = **Amblemma**, R. Br. (Commelin.).
Wallichii, C. B. Clarke, l. c. 89. t. 26 = Amblema aequinoctiale.

AMELLUS, P. Br. Hist. Jamaic. 317 (1756) = **Melanthera**, Rohr. (Compos.).

AMELLUS, Linn. Syst. ed. x. 1225 (1759). *COMPOSITAE*, Benth. & Hook. f. ii. 265.
HARNELIA, Walp. Rep. ii. 974 (1848).
KRAUSSIA, Sch. Bip. in Flora, xxvii. (1844) 672.
alternifolius, Roth, Catalecta, ii. 111 = strigosus.
asiaticus, *Cass. in Dict. Sc. Nat.* xxxvii. 489.—Afr. austr.
annuus, *Willd. Sp. Pl.* iii. 2215.—Afr. austr.
caroliniana, Walt. Fl. Carol. 218 = Eclipta alba.
collopodius, *DC. Prod.* v. 214.—Afr. austr.
diffusus, Forst. in Comm. Gott. ix. (1787) 89 = Chiliotrichum amelloideum.
divaricatus, Gater. Fl. Montaub. 147 = Aster Novi-Belgii.
floribundus, Willd. ex DC. Prod. v. 96 = Lisbum igniarium.
fasciculosus, DC. Prod. v. 214 = hispidus.
hispidus, *DC. l. c.*—Afr. austr.
Lychnitis, *Linn. Syst.* ed. X. 1255.—Afr. austr.
Lychnitis, E. Mey. ex Steud. Nom. ed. II. i. 76 = tenuifolius.
microglossus, *DC. Prod.* v. 215.—Afr. austr.
mutabilis, Gater. Fl. Montaub. 147 = Aster mutabilis.
nanus, *DC. Prod.* v. 215.—Afr. austr.
officinalis, Gater. Fl. Montaub. 147 = Aster Amellus.
pallidus, Salisb. Prod. 205 = A. Lychnitis.
pedunculatus, Orteg. ex Willd. Enum. Hort. Berol. 916 = Tridax procumbens.
rosmarinifolius, Poepp. ex DC. Prod. v. 216 = Chiliotrichum rosmarinifolium.
scabridus, *DC. l. c.* 214.—Afr. austr.
speciosus, Gater. Fl. Montaub. 146 = Aster chinensis.
spinulosus, *Pursh, Fl. Am. Sept.* ii. 564.—Am. bor.
strigosus, Less. in Linnæa, vi. (1831) 110 = annuus.
tenuifolius, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
tridactylus, *DC. Prod.* v. 215.—Afr. austr.
umbellatus, Linn. Fl. Jam. Pugill. 24 (Syst. ed. X. 1225) = Lisbum Browni.
villosus, Pursh, Fl. Am. Sept. ii. 564 = Chrysopsis villosa.

AMERIMNON, P. Br. Hist. Jamaic. 288 (1756); DC. Prod. ii. 491 (1825) = **Dalbergia**, Linn. (Legumin.).
afine, Spreng. Syst. iii. 199.—Am. austr.
Brownei, Sw. Fl. Ind. Occ. 1284 = Dalbergia Amerimnon.
Brownii, Jacq. Enum. Pl. Carib. 27.—Ind. occ.
Ebenus, Sw. Prod. Veg. Ind. Occ. 104 = Brya Ebenus.
fascicularum, Buch.-Ham. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 109 = Millettia fruticosa.
guianense, Klotzsch, in Schomb. Faun. et Fl. Guy. 1904.—Guiana.
horridum, Dennst. Schlüssl. Hort. Malab. 84 (quid ?).—Ind. or.
latifolium, Willd. Sp. Pl. iii. 909 = Lonchocarpus latifolius.
microcalla, Blanco, Fl. Filip. ed. I. 563 = Dalbergia lanceolaria.
obovatum, Buch.-Ham. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 246 = Derris thyrsiflora.
pallidum, Buch.-Ham. ex Baker, l. c. 106 = Millettia auriculata.
parviflorum, Spreng. Nova Enidack. ii. 259.—Bras.
pinnatum, Jacq. Sel. Stirp. Am. 200 = Dalbergia Amerimnon.
polystachyon, Spreng. Syst. Cur. Post. 263 = Eysenhardtia amorphoides.
pubescens, Willd. Sp. Pl. iii. 909 = Pterocarpus scandens.
scandens, Willd. l. c. = Pterocarpus scandens.
secundum, Buch.-Ham. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 247 = Derris secunda.
Sieberi, Reichb. ex DC. Prod. ii. 420 = Ecastaphyllum Brownei.
strigulosum, H. B. & K. Nov. Gen. et Sp. vi. 388.—Venezuela.

AMERINA, DC. in Meisn. Gen. 378 (1840) = **Asgiphila**, Jacq. (Verben.).
ternifolia, DC. Prod. ix. 518.—N. Granat.
tenuiflora, DC. l. c. = Asg. tomentosa ?
tripetala, DC. l. c.—Ait. nostr.

AMERINA, Norohna, in Verh. Batav. Gen. v. (1790) Art. iv. 1; ed. II. (1827) 64 = **Aglaia**, Lour. (Meliac.).

AMERINA, Rafin. Alsog. Amer. 15 (1838) = **Salix**, Tourn.
Baldwiniana, Rafin. l. c. 16.—Am. bor.
caroliniana, Rafin. l. c. 15 = S. nigra.
glandulosa, Rafin. l. c. = S. pentandra.
Houstoniana, Rafin. l. c. = S. nigra.
micrurica, Rafin. l. c. 16.—Am. bor.
nigra, Rafin. l. c. 15 = S. nigra.
tinctoria, Rafin. l. c. 15.—Am. bor.
tomentosa, Rafin. l. c. = S. Lapponum.

AMETHYSTEA, Linn. Amoen. Acad. i. 385 (1749). *LABIATAE*, Benth. & Hook. f. ii. 1220.
AMETHYSTINA, Mess. ex Hall. in Act. Upsal. 1742 (1748) 51. t. 4.
caerulea, *Linn. Sp. Pl.* 21.—Sibiria.
corymbosa, Pers. Syn. i. 24 = caerulea.
trifida, Hill, Veg. Syst. xvii. 44 = caerulea.

AMETHYSTINA, Mess. ex Hall. in Act. Upsal. 1742 (1748) 51. t. 4 = praec.

AMETRON, Rafin. Sylva Tellur. 160 (1838) = **Rubus**, Tourn. (Rosac.).
pedatum, Rafin. l. c. 161 = R. pedatus.
pyrifolium, Rafin. l. c. = R. pyrifolius.

AMHERSTIA, Wall. Pl. As. Rar. i. 1. t. 1, 2 (1880). *LEGUMINOSAE*, Benth. & Hook. f. i. 578.
nobilis, *Wall. l. c.*—Burma.

AMIANTANTHUS, Kunth, Enum. Pl. iv. 179 (1843) = AMIANTHIUM, A. Gray = **Zygadenus**, Michx. (Liliac.).

AMIANTHEMUM, Stend. Nom. ed. II. i. 76 (1840) = seq.

AMIANTHIUM, A. Gray, in Ann. Lyc. N. York, iv. (1848) 121 = **Zygadenus**, Michx. (Liliac.).
angustifolium, A. Gray, l. c. 194 = Z. angustifolius.
aspericaule, A. Gray, l. c. 126 = Melanthium aspericaule.
leimanthoides, A. Gray, l. c. 125 = Z. leimanthoides.
macrotos, Rafin. Fl. Tellur. iv. 27 = Z. muscaetoxicum.
muscaetoxicum, A. Gray, in Ann. Lyc. N. York, iv. (1848) 122 = Z. muscaetoxicum.
Nuttallii, A. Gray, l. c. 128 = Z. Nuttallii, paniculatus.

AMIANTHUM, Rafin. Fl. Te lur. iv. 27 (1836) = praec.

AMICIA, H. B. & K. Nov. Gen. et Sp. vi. 511. t. 600. (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 514.
ZYGOMERIS, Moç. & Sesse, ex DC. Prod. ii. 515 (1825).
glandulosa, *H. B. & K. l. c.*—N. Granat.
mediagnas, *Griseb. in Goett. Abh.* xxiv. (1879) 105. —Reg. Argent.
Zygomeris, *DC. Prod.* ii. 515.—Mexico.

AMICTONIS, Rafin. Sylva Tellur. 161 (1838) = **Calli-carpa**, Linn. (Verbenac.).
japonica, Rafin. l. c.=C. longifolia.

AMIDA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 390 = **Madia**, Molina (Compos.).
gracilis, Nutt. l. c. = M. glomerata.
hirsuta, Nutt. l. c. = M. glomerata.

AMIDENA, Adans. Fam. ii. 470 (1763) = **Arisaema**, Mart. (Aroid.).
japonica, Rafin. Fl. Tellur. iv. 15 = Arisaema japonicum.

AMIRIS, La Llave, in La Naturaleza, vii. Apénd. 71 (1888) = **Amyris**, Linn. (Rutac.).

AMIROLA, Pers. Syn. ii. 565 (1807) = **Llagunoa**, Ruiz & Pav. (Sapindac.).
glandulosa, Hook. & Arn. Bot. Beech. Voy. 13 = Ll. glandulosa.
mollis, DC. Prod. i. 616 = Ll. mollis.
nitida, Pers. Syn. ii. 565 = Ll. nitida.
parvifolia, Sessé. Nom. ed. II. i. 76 = Ll. nitida.
prunifolia, DC. Prod. i. 616 = Ll. prunifolia.

AMISCHOTOLYPE, Hassk. in Flora, xlvi. (1863) 391 = **Forrestia**, A. Rich. (Commelin.).
glabrata, Hassk. l. c. 392 = F. glabrata.
marginata, Hassk. l. c. = F. marginata.
mollissima, Hassk. l. c. = F. mollis.

AMMANELLA, Miq. Fl. Ind. Bat. I. i. 618 (1855) = **Ammannia**, Linn. (Lythrar.).
lanata, Miq. l. c. 619 = Ammannia octandra.

AMMANNIA, [Houst.] Linn. Gen. ed. I. 337 (1737). *LYTHRARIEAE*, Benth. & Hook. f. i. 776.
AMELETIA, DC. in Mém. Soc. Phys. Genèv. iii. 11. (1826) 89.
CORNELIA, Ard. Specim. ii. 9. t. 1 (1764).
CRYPTOTHECA, Blume, Bijdr. 1198 (1826).
NEXILIS, Rafin. New Fl. Am. iv. 9 (1836).
ORTEGIOIDES, Soland. ex DC. Prod. iii. 76 (1828).
NIMMOLIA, Wight, in Madras Journ. Sc. vii. (1837) 311. t. 20.
ROTALA, Linn. Mant. ii. 148 (1771).
SELLOWIA, Koth, ex Roem. & Schult. Syst. v. 407 (1819).
SUFFRENIA, Bellardi, in Act. Taur. vii. (1796) 445. t. 1.
TRITHECA, Miq. Fl. Ind. Bat. i. 614 (1855).
WINTERLIA, Spreng. Syst. i. 519 (1825).
acutidens, Miq. Fl. Ind. Bat. i. i. 617 = peploides.
aegyptiaca, Kotschy, ex Koehne, in Engl. Bot. Jahrb. i. (1881) 253, 254 = urceolata, apiculata.
aegyptiaca, Willd. Hort. Berol. t. 6 = salicifolia.
alata, *Stend. Nom. ed.* II. i. 76.—Reg. Caucas.
anagalloides, Sond. in Linnaea, xxiii. (1850) 40 = Nesaea anagalloides.
apiculata, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 254. —Am. trop.
arenaria, H. B. & K. Nov. Gen. et Sp. vi. 190 = senegalensis.
aspera, Guill. & Perr. Tent. Fl. Senog. i. 304 = Nesaea aspera.
attenuata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 278 = baccifera.
auriculata, Willd. Hort. Berol. i. 7. t. 7 = senegalensis.
australasica, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 41 = multiflora.
baccifera, Roth, Sp. Nov. 100 = leptopetala.
baccifera, *Linn. Sp. Pl.* 120.—Geront. calid.
baccifera, Pollini, Fl. Veron. i. 178 = salicifolia.
Berart, Spreng. Suppl. Fl. Maine-et-Loire, 89 = Lythrum nummularifolium.
borysthenica, DC. Prod. iii. 78 = Lythrum nummularifolium.
borysthenica, Kar. & Kir. in Bull. Soc. Nat. Mosc. xiv. (1841) 491 = Peplis alternifolia.
caspia, Hohen. Enum. Talysch. 133 = baccifera.
caspica, Bch. Fl. Taur. Cauc. ii. 467; Suppl. 111 = salicifolia.
catholica, Cham. & Schlecht. in Linnaea, ii. (1827) 576 = latifolia.
catholica, Hook. & Arn. ex Seem. Bot. Voy. Herald, 284 = latifolia.
coccinea, Pers. Syn. i. 147 = octandra.
coccinea, Roxb. Fl. Hort. Univ. (Havn.) Programm. Descr. (1779) 7 = latifolia.
cordata, *Wight & Arn. Prod.* 304.—Ind. or.
crassicaulis, Guill. & Perr. Tent. Fl. Senog. i. 303 = Nesaea polyantha.
crinipes, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 49 = Nesaea crinipes.
cryptantha, *Baker, in Journ. Linn. Soc.* xxi. (1884) 245.—Madag.
debilis, Ait. Hort. Kew. ed. I. i. 163 = baccifera.
densiflora, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 589 = salicifolia.
densiflora, Roth, Sp. Nov. 99 = pentandra.
dentata, *A. Gray, Pl. Wright.* ii. 55.—Am. bor.
dentelloides, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 76.—Bengal.
diandra, *F. Muell. Fragm.* iii. 106.—Austral.
diffusa, Willd. Enum. Hort. Berol. i. 167 = senegalensis.
dodecandra, DC. in Mém. Soc. Phys. Genèv. iii. ii. (1826) 89. t. 2 = Nesaea Candolleli.
elatinoides, *DC. l. c.* 92. t. 3. f. B.—Afr. trop.
elatinoides, Schimp. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 468 (= *Rotala stagnina*).
elongata, Blume, Mus. Bot. Lugd. Bat. ii. 185 = peploides.

AMMANNIA:—
filiformis, Baill. Hist. Pl. vi. 488 (= *Rotala filiformis*).
filiformis, DC. in Mém. Soc. Phys. Genèv. iii. ii. (1826) t. 79 = senegalensis.
fimbriata, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 569 = pentandra.
floribunda, *C. B. Clarke, l. c.* 567.—Ind. or.
floribunda, Guill. & Perr. Tent. Fl. Senog. i. 302 = senegalensis, multiflora.
glauca, Wall. Cat. n. 2100 = salicifolia.
gracilis, Guill. & Perr. Tent. Fl. Senog. i. 301 = senegalensis.
hastata, DC. Prod. iii. 78 = latifolia.
hexandra, Wall. Cat. n. 2108 = pentandra.
Heyneana, Wall. l. c. 2104 = pentandra.
Hildebrandtii, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 257.—Afr. trop.
humilis, *Michx. Fl. Bor. Am.* i. 99.—Am. bor.
hyssopica, Fisch. ex Stend. Nom. ed. II. i. 77 = auriculata.
illecebroides, Arn. ex Koehne, in Engl. Bot. Jahrb. i. (1881) 161 (= *Rotala decussata*).
indica, Lam. Illustr. i. 311. n. 1555 = baccifera.
japonica, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 261.—Japon.
lanceolata, *Heyne, in Wall. Cat.* n. 2106.—Ind. or.
latifolia, *Linn. Sp. Pl.* 119.—Am. bor.; Ind. occ.
latifolia, Wall. Cat. n. 2096 = peploides.
latifolia, Walp. Rep. ii. 102 = sanguinolenta.
leptopetala, *Blume, Mus. Bot. Lugd. Bat.* ii. 184.—Japon.
linguiata, Griseb. Cat. Pl. Cub. 106 = latifolia.
littorea, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 261.—Japon.
loandensis, Welw. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 480 = Nesaea loandensis.
longipes, *Wright, in Sauv. Fl. Cub.* 53.—Ind. occ.
lythrifolia, Salisb. Prod. 65 = latifolia.
madagascariensis, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 129 = multiflora.
microcarpa, DC. in Mém. Soc. Phys. Genev. iii. ii. (1826) 93 = multiflora ?
monoflora, Blanco, Fl. Filip. ed. I. 64 = latifolia ?
multiflora, *Roxb. Hort. Beng.* 11; *Fl. Ind. ed. Carey*, i. 447.—Ind. or.
nana, Roxb. Il. co. 11, 448 = peploides.
nana, Roxb. in Wall. Cat. n. 2105 = pentandra.
Nuttallii, A. Gray, Man. Bot. N. U. St. ed. IV. add. p. xcii = Didiplis linearis.
occidentalis, DC. Prod. iii. 78 = humilis.
octandra, *Linn. f. Suppl.* 187.—Ind. or.; Malaya.
octandra, Cham. & Schlecht. in Linnaea, ii. (1827) 576 = latifolia.
pallida, Lehm. Sem. Hort. Hamb (1823) 3; (Linnaea, iii. (1828) 9) = latifolia.
parviflora, DC. Prod. iii. 77 = multiflora.
passerinoides, Welw. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 480 = Nesaea passerinoides.
pentandra, *Roxb. Fl. Ind.* i. 448.—Geront. trop.
peploides, *Spreng. Syst.* i. 444.—Orient; Aa. trop.
pinnatifida, *Linn. f. Suppl.* 187.—Java.
polystachya, Wall. Cat. n. 2094 = peploides.
Portula, Baill. in Bull. Soc. Linn. Par. i. (1874) 87 = Peplis Portula.
Prieuriana, Guill. & Perr. Tent. Fl. Senog. i. 303 = senegalensis.
prostrata, Buch.-Ham. ex Blume, Mus. Bot. Lugd. Bat. ii. 182 = baccifera.
purpurea, Lam. Encyc. i. 181 = humilis.
pusilla, Sond. in Linnaea, xxiii. (1848) 40 = senegalensis.
pygmaea, *Kurz, in Journ. Bot.* v. (1867) 276.—Ind. or.
racemosa, Poir. Encyc. Suppl. i. 329 = arenaria.
racemosa, Roth, Catalect. fasc. iii. 25 = octandra.
ramosa, Hill, Veg. Syst. xi. 14 = humilis.
ramosior, Blanco, Fl. Filip. ed. II. 64 = peploides.
ramosior, Linn. Sp. Pl. 120 = humilis.
ramosior, Linn. Mant. ii. 228 = latifolia.
repens, Rottl. ex Mart. in Acad. Muensch. Phil. vi. (1820) 150, ex DC. Prod. iii. 80 = peploides.
retusa, Koehne, in Engl. Bot. Jahrb. i. (1881) 254.—Afr. trop.
Richiei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 566. —Ind. or.
robusta, Heer, Ind. Sem. Hort. Turic. (1845) = latifolia.
rosea, Poir. Encyc. Suppl. i. 329 (= *Rotala mexicana*).

AMMANNIA :—
Rotala. *F. Muell. Fragm.* iii. 108.—Geront. trop.
rotundifolia, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 230.—Ind. or. ; China.
rubra, Buch.-Ham. l. c. = pentandra.
sagittata, DC. Prod. iii. 80 = latifolia.
salicifolia, Hiern, in Oliver, Fl. Trop. Afr. ii. 476 = baccifera.
salicifolia, *Monti, in Comm. Bonon.* i. (1767) 113.—As. & Afr. trop.
salsuginosa, *Guill. & Perr. Tent. Fl. Senneg.* i. 302.—Afr. trop.
sanguinolenta, Cham. & Schlecht. in Linnæa, v. (1830) 538 = auriculata.
sanguinolenta, Heyne, ex Stend. Nom. ed. II. i. 77 = salicifolia.
sanguinolenta, Sw. Prod. Veg. Ind. Occ. 83 = latifolia.
sanguinolenta, Hook. & Arn. ex Seem. Bot. Voy. Herald, 284 = latifolia.
sarcophylla, Welw. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 479 = Nesæa sarcophylla.
senegalensis, *Lam. Illustr.* i. 312. n. 155. t. 77. f. 2.—Geront. trop.
simplicicaulis, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 54.—Ind. or.
styloza, Flach. et Mey. Ind. Sem. Hort. Petrop. vii. 41 = latifolia.
subrotunda, *Wall. ex Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 55.—Ind. or.
subspicata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 484 = rotundifolia.
subspicata, Hohen. ex Hook. f. Fl. Brit. Ind. ii. 568 = pentandra.
tenella, *Guill. & Perr. Tent. Fl. Senneg.* i. 297.—Afr. trop.
tennis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 567.—Ind. or.
texana, Scheele, in Linnæa, xxi. (1848) 588 = latifolia.
trifera, R. Br. ex Benth. Fl. Austral. iii. 397 = Nesæa lanceolata.
trifera, Wall. Cat. n. 6328 = Nesæa trifora.
undulata, C. A. Mey. Ind. Sem. Hort. Petrop. ix. 56 = senegalensis.
uroscalata, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 478.—Afr. trop.
verticillaris, Bellf. Hist. Pl. vi. 489 (= *Rotala verticillaris*).
verticillata, Boiss. Fl. Orient. ii. 745 = baccifera.
verticillata, Lam. Encyc. i. 131 = salicifolia.
verticillata, Wight & Arn. Prod. i. 304 = lanceolata.
vesicatoria, Roxb. Hort. Beng. 11 ; Fl. Ind. ed. Carey, i. 447 = baccifera.
viridis, Willd. ex Homem. Hort. Hafn. i. 146 = baccifera.
Wallichii, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 94 = Hydrolythrum Wallichii.
Wormskioldii, *Flach. et Mey. Ind. Sem. Hort. Petrop.* vii. 42.—Bras.
Wrightii, *A. Gray, Pl. Wright.* ii. 55.—Am. bor.

AMMANTHUS, Boiss. & Heldr. Diagn. Ser. I. xi. 18 (1849) = Chrysanthemum, Tourn. (Compos.).
agrestifolius, Boiss. & Heldr. ex Boiss. Fl. Orient. iii. 358 = C. maritimum.
filicoides, Boiss. & Heldr. ex Boiss. Diagn. Ser. I. xi. 19.—Ins. Creta.
maritimus, Boiss. & Heldr. l. c. = C. maritimum.

AMMI, [Tourn.] Linn. Syst. ed. I. (1735) ; Gen. ed. I. 79 (1737). *UMBELLIFERÆ*, Benth. & Hook. f. i. 889.
DASYSPERMUM, Neck. Elem. i. 276 (1790).
GOHORIA, Neck. l. c. 272 (1790).
VIGNALA, Riv. ex Guerin, Fruct. i. 92. t. 21 (1788).
acaule, Sprang. in Roem. & Schult. Syst. vi. 530 = Chamæsciadium acaule.
ammifolium, *Lam. Encyc.* i. 122.—Orient. ; Sibir.
ammithifolium, Ledeb. Fl. Ross. ii. 246 = Artedia squamata.
Bunkeri, Heldr. ex Hoffm. Gen. Umbell. p. xviii ; Icon. Geogr. 23 = majus.
Broussonetii, DC. Prod. iv. 113 = majus ?
capillaceum, *Michx. Fl. Bor. Am.* i. 164 = Discopleura capillacea.
cicutæfolium, Willd. ex Roem. & Schult. Syst. vi. 531 = majus.
cicutarium, Willd. ex Roem. & Schult. l. c.—Am. calid.
AMMI, —
copticum, Linn. Mant. 56 = Carum copticum.
costatum, Ell. Sketch Fl. Carol. i. 350 = Discopleura capillacea.
crinitum, *Guss. Pl. Rar.* 128. t. 25.—Calabr. ; Sicil.
dauciifolium, Buch.-Ham. in Wall. Cat. n. 7309 = Seseli dauciifolium.
dauciifolium, Scop. Fl. Carn. ed. II. i. 307 = Libanotis athamantoides.
daucoides, Salem. ex DC. Prod. iv. 113 = Ammiopsis daucoides.
dentatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Visnaga.
divaricatum, Michx. Fl. Bor. Am. i. 168 = Apium divaricatum.
glaucum, Salisb. Prod. 162 = majus.
fœniculaceum, Link, Enum. Hort. Berol. i. 266 = Ligusticum fœniculaceum.
fœniculifolium, Lag. ex Lond. Hort. Brit. 103 = prasc. ?
glaucifolium, Blanco, Fl. Filip. ed. I. 213 = Carum copticum.
glaucifolium, Linn. Sp. Pl. 245 = majus.
Huntii, *H. C. Wats. in Hook. Lond. Journ. Bot.* vi. (1847) 384.—Ins. Azor.
indicum, *Buch.-Ham. in Wall. Cat.* n. 7308.—Ind. or.
intermedium, DC. Prod. iv. 113 = majus.
majus, *Linn. Sp. Pl.* ed. I. 243.—Europ. ; Orient ; Afr. bor.
majus, Wall. Fl. Carol. 113 = Discopleura capillacea.
meoides, Pers. Syn. i. 308 = anethifolium.
minus, *Hort. Gotting.* (1976) 8.—Europ.
paucriadiatum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 272 = majus.
petroselinoides, Presl, ex DC. Prod. iv. 102 = Carum peregrinum.
procerum, *Lowe, Man. Fl. Mad.* 351.—Madera.
pumilum, *DC. Prod.* iv. 113.—Lusitan.
pyrenaicum, *Lapeyr. Hist. Abr. Pl. Pyr.* 145.—Pyren.
rubricaule, Hornem. Hort. Hafn. i. 273 = Discopleura capillacea.
sanctæ, *Torrey. ex Bess. in Flora*, xvii. (1834) i. ; *Beibl.* 13, nomen.—Rossia.
trachycarpum, Fisch. ex Boiss. Fl. Orient. ii. 855 = Aphanopleura trachysperma.
Visnaga, *Lam. Ill. Pr.* iii. 463.—Reg. Mediterr.

AMMIANTHUS, Spruce, ex Benth. & Hook. f. Gen. ii. 98 (1873) = Retinyphyllum, Humb. & Bonpl. (Rubiac.).

AMMIOPSIS, Boiss. Diagn. Ser. II. ii. 96 (1856). *UMBELLIFERÆ*, Benth. & Hook. f. i. 927.
daucoides, *Boiss. l. c.* 97.—Mauritan.

AMMIOS, Moench, Meth. 99 (1794) = Carum, Tourn. (Umbellif.).
maricosa, Moench, l. c. = C. copticum.

AMMOBIUM, R. Br. in Bot. Mag. t. 2459 (1824). *COMPOSITÆ*, Benth. & Hook. f. ii. 316.
alatum, *R. Br. l. c.*—Austral.
craspedioides, *Benth. Fl. Austral.* iii. 584.—Austral.
plantagineum, *G. Don, in Lond. Hort. Brit.* 353.—Austral.
spathulatum, Gaudich. Voy. Freyc. Bot. 467. t. 91 = alatum.

AMMOBROMA, Torr. in Ann. Lyc. N. York, viii. (1867) 51. t. 1. *LENNOACEÆ*, Benth. & Hook. f. ii. 692.
Sonoræ, *Torr. l. c.* 52.—N. Mexic.

AMMOCHARIS, Herb. App. 17 (1821). *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 737.
FALLENIA, Salisb. Gen. Pl. Fragm. 116 (1866).
coronica, Herb. App. 17 = falcata.
falcata, *Herb. l. c.*—Afr. austr.
longifolia, M. Roem. Amar. 62 = Crinum longifolium.
Staterianæ, Kunth, Enum. Pl. v. 613 = Brunsvigia Staterianæ.

AMMOCHLOA, Boiss. Diagn. Ser. I. xiii. 51 (1858). *GRAMINEÆ*. Benth. & Hook. f. iii. 1181.
CEPHALOCHLOA, Coss. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 229.
palaestina, Boiss. Diagn. Ser. I. xiii. 52 = subacualis.

AMMOCHLOA :—
pungens, *Boiss. Diagn.* Ser. I. xiii. 52.—Afr. bor.
subacualis, *Balansa, ex Coss. in Bull. Soc. Bot. Fr.* i. (1854) 217 ; iv. (1857) 899.—Syria ; Afr. bor.

AMMODAUCUS, Coss. & Dur. in Bull. Soc. Bot. Fr. vi. (1859) 932 = Daucus, Linn. (Umbellif.).
leucotrichus, *Coss. l. c. et* xxii. (1875) 60.—Algeria.

AMMODENDRON, Fisch. ex DC. Prod. ii. 522 (1825). *LEGUMINOSÆ*, Benth. & Hook. f. i. 554.
caspium, Eichw. Casp.-Casc. 7 = Eichwaldii.
Conollyi, *Bunge, ex Boiss. Fl. Orient.* ii. 627.—Turkest.
Eichwaldii, *Ledeb. & Mey. in Eichw. Casp.-Casc.* 37. t. 36.—Reg. Casp.
Karelini, Bunge, Fl. Rel. Lehm. 69 = Conollyi.
Karelinii, *Fisch. & Mey. in Ledeb. Fl. Ross.* i. 717.—Reg. Casp.
Lehmanni, *Bunge, ex Boiss. Fl. Orient.* ii. 626.—Turkest.
persicum, *Bunge, ex Boiss. l. c.* 627.—Persia.
Sieversii, *DC. Prod.* ii. 522.—Soongaria.
Zablotskii, *Fisch. & Mey. ex Kar. in Bull. Soc. Nat. Mosc.* (1839) 150, nomen.—Rossia.

AMMODENIA, Patrin. ex J. G. Gmel. Fl. Sibir. iv. 160 (1769) = Armaria, Linn. (Caryophyll.).
pepleides, Rupr. in Beitr. Pfl. Russ. Reich. ii. 25 = Ar. pepleides.

AMMODIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 391 = Chrysopsis, Nutt. (Compos.).
oregana, Nutt. l. c. = C. oregana.

AMMODYTES, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 253 = Astragalus, Tourn. (Legum.).
mollissimus, Stev. l. c. = Astr. mollissimus.

AMMOGETON, Schrad. Ind. Sem. Hort. Goett. (1838) 1 = Troximon, Nutt. (Compos.).
scorsoneræfolius, Schrad. l. c. = T. glaucum.

AMMOIDES, Adans. Fam. ii. 96 (1763) = Carum, Tourn. (Umbellif.).

AMMOLIRION, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 815 = Rhemarus, Bieb. (Liliac.).
stenophyllum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 218 = E. Bungei.
Stevani, Kar. & Kir. l. c. = E. tuberiensis.

AMMONALIA, Desv. ex Endl. Gen. 966, in syn. (1840) = Armaria, Linn. (Caryophyll.).

AMMOPHILA, Host. Gram. Austr. iv. 24. t. 41 (1809). *GRAMINEÆ*, Benth. & Hook. f. iii. 1158.
PSAMMA, Beauv. Agrost. 148. t. 6. f. 1. (1818).
arenaria, Link, Hort. Berol. i. 106 = arundinacea.
arundinacea, *Host, Gram. Austr.* iv. 24. t. 41.—Europ. ; Am. bor.
baltica, *Link, Hort. Berol.* i. 106 ; *Dum. Fl. Belg.* 152. —Europ.
brevipilis, *Benth. ex Vasey, Cat. Gram. U. St.* 52.—Am. bor.
Curtisii, *Vasey, in Torrey Bot. Club*, xi. (1884) 7.—Florida.
longifolia, *Vasey, Agric. Grasses U. St.* 71.—Am. bor.

AMMORRHIZA, Ehrh. Phytophyl. (1789) ; Beitr. iv. 146 = Carex, Linn.

AMMOSELINUM, Torr. & Gray, in Pacif. Rail. Rep. ii. 165 (1855). *UMBELLIFERÆ*, Benth. & Hook. f. i. 909.
Popei, *Torr. & Gray, l. c.*—Texas.

AMMOSERIS, Endl. Gen. 800 (1838) = Launæa, Cass. (Compos.).
asplenifolia, D. Dietr. Syn. Pl. iv. 1819 = L. asplenifolia.
bellidifolia, D. Dietr. l. c. 1820 = L. bellidifolia.
Desfontainii, D. Dietr. l. c.—Afr. austr.
resedaefolia, D. Dietr. l. c. 1819 = L. resedaefolia.
setosa, D. Dietr. l. c.—Ind. or.
surrentanum, D. Dietr. l. c. = L. spinatifida.
surrinamensis, D. Dietr. l. c.—Guiana.

AMMOSPERMA, Hook. f. in Benth. & Hook. f. Gen. i. 89 (1862). *CRUCIFERÆ*, Benth. & Hook f. i. 88.
cinerea, *Hook. f. t. c.*; *Baill. Hist. Pl.* iii. 278.—Afr. bor.

AMMOTHAMNUS, Bunge, Enum. Pl. Lehm. 67. t. 19 (1847). *LEGUMINOSÆ*, Benth. & Hook. f. i. 555.
glibosus, *Boiss. Fl. Orient.* ii. 628.—Assyria. Lehmanni, *Bunge, l. c.* 343.—Turkest.

AMMYRSINE, Pursh, Fl. Am. Sept. i. 301 (1814) = Leiophyllum, Hedw. f. (Ericac).
buxifolia, Pursh, l. c. = L. buxifolium.
Lyoni, Sweet, Hort. Brit. ed. II. 344 = L. buxifolium.
prostrata, Sweet, ex Steud. Nom. ed. II. i. 78 = L. buxifolium.
thymifolia, Forbes, Hort. Woburn. 93 = L. buxifolium.

AMNI, Brongn. Enum. Genr. 110 (1843) err. typ. = Ammi, Tourn.

AMOGETON, Neck. Elem. iii. 267 (1790) = Aponogeton, Thunb. (Naiad.).

AMOLEIACHYRIS, Sch. Bip. in Walp. Rep. ii. 951 (1843), sphalm. = AMPHIACHYRIS, Nutt. = Gutierrezia, Lag. (Compos.).

AMOMIS, Berg. in Linnæa, xxvii. (1854) 416 = Pimenta, Lindl. (Myrtac.).
acris, Berg, l. c. 417 = P. acris.
oblongata, Berg, l. c. 421 (= P. acris ?).—Ind. occ. : cult. ; Guiana.
Pimenta, Berg, l. c. 418 (= P. acris ?).—Ind. occ. ; Guiana.
pimentoides, Berg, l. c. 420 = P. acris.

AMOMOPHYLLUM, Engl. in Gard. Chron. (1877) i. 139 = Spathiphyllum, Schott (Aroid.).
floribundum, Engl. l. c. 140 = S. floribundum.
Patini, Engl. l. c. 139 = S. Patini.

AMOMUM, Linn. Mus. Clif. 26 (1736); Gen. ed. I. 330 (1737). *SCITAMINEÆ*, Benth. & Hook. f. iii. 644.
ACHARMA, Griff. Notul. iii. 436 (1851).
ALEXIS, Salisb. in Trans. Hort. Soc. i. (1812) 284.
CENOLOPHON, Horan. Prod. Monog. Scitam. 36 (1863).
DIRACODES, Blume, Enum. Pl. Jav. i. 55 (1830).
DONACODES, Blume, l. c. 54 (1830).
EULINGERA, Gieske, Præl. Linn. 199, 200 (1792).
GRANTHUS, Retzius in Syll. Retzius. ii. (1838) 5.
GROCALLIS, Horan. Prod. Monog. Scit. 33 (1863).
GREENWAYA, Gieske, Præl. Linn. 199, 206 (1792).
HORNSTEDTIA, Retz. Obs. vi. 18 (1972).
MAROGNA, Salisb. in Trans. Hort. Soc. i. (1812) 283.
MEISTERA, Gieske, Præl. Linn. 199, 205 (1792).
NICOLLIA, Horan. Prod. Monog. Scitam. 32. t. 1 (1863).
PALUDANA, Gieske, Præl. Linn. 199, 207 (1792).
PHAOMERIA, Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
STENOCHASMA, Griff. Notul. iii. 431 (1851).
TORYMENIS, Salisb. in Trans. Hort. Soc. i. (1812) 282.
WURMBAINIA, Gieske, Præl. Linn. 199, 206 (1792).
ZEDOARIA, Rafin. Fl. Tellur. iv. 53 (1836).
aculeatum, *Roxb. in As. Res.* xi. (1810) 344.—Malaya, acuminatum, *Thw. Enum. Pl. Zeyl.* 317.—Zeylan.
Afzelii, Rosc. in Trans. Linn. Soc. vii. (1807) 254 = A. Granum-paradisi.
Alpinia, Retth. in Coll. Soc. Med. Havn. ii. (1775) 246 t. 3 = Renealmia racemosa.
angustifolium, Hanb. in Journ. Linn. Soc. xiii. (1873) 155 = Danielli.
angustifolium, Salisb. Prod. 4 = Zingiber officinale.
angustifolium, *Sonner. Voy. Indes*, iii. 276.—Madag.
aquaticum, *Rœusch. Nom.* ed. III. 1.—Zeylan.
arborescum, Lour. Fl. Cochinch. 7 = Costus Loureiri.
aromaticum, Roxb. Hort. Beng. 1 ; Fl. Ind. i. 44.—Ind. or.
arundinaceum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 109.—Afr. trop.
Benthamianum, *Trim. in Journ. Bot.* xxiii. (1885) 265. —Zeylan.

AMOMUM :—
bifidum, Stokes, Bot. Comment. i. 168 = Alpinia Allughas.
biforum, *Yach, in Malay. Misc.* i. (1820) n. 1. p. 2.—Penang.
caniseicarpum, *Benth. & Hook. f. Gen.* iii. 644.—Ind. or.
capitatum, *Roxb. ex Link, Yahrb.* ii. (1811) 64.—Ind. or.
Cardamomum, Koen. in Retz. Obs. fasc. iii. 59 = seq.
Cardamom, *Linn. Sp. Pl.* —Ind. or. ; Malaya.
Cardamomum, Willd. Enum. Hort. Berol. Suppl. 1 = Alpinia striata.
cereum, *Hook. f. in Kew Journ. Bot.* iv. (1854) 296. —Afr. trop.
Cevuga, *Seem. Fl. Vit.* 291.—Ins. Fiji.
ciliatum, *Blume, Enum. Pl. Jav.* 49.—Java.
citratum, Pereira, Mat. Med. ii. 1137 = cereum.
Clusii, Hook. f. Bot. Mag. t. 5250 = Danielli.
Clusii, *Sm. in Rees, Cyclop.* xxxix. n. 4.—Afr. trop.
coccineum, *Benth. & Hook f. Gen.* iii. 644.—Java.
compactum, Roem. & Schult. Syst. i. 28 = A. Cardamon.
corynostachyum, *Wall. Pl. As. Rar.* i. 48 t. 58.—Ind. or.
costatum, *Benth. & Hook f. Gen.* iii. 635.—Ind. or.
Curcuma, Jacq. Hort. Vindob. iii. 5. t. 4 = Curcuma longa.
Dalzellii, *F. Muell. Fragm.* viii. 25.—Austral.
Danielli, *Hook. f. in Kew Journ.* iv. (1854) 129. t. 5. —Afr. trop.
dealbatum, *Roxb. Hort. Beng.* 1; *Fl. Ind.* i. 43.—Ind. or.
echinatum, Willd. Sp. Pl. i. 8.—Ind. or.
elatum, Salisb. Prod. 5 = A. Granum-paradisi.
Emal, Rœusch. Nom. ed. III. 1 = Elettaria Cardamomum.
exscapum, Sims, in Kon. & Sims, Ann. Bot. i. (1805) 244 = A. Granum-paradisi.
fasciculatum, *Benth. & Hook. f. Gen.* iii. 645.—Ind. or.
Fenzlii, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 154.—Ins. Nicobar.
floribundum, *Benth. & Hook. f. Gen.* iii. 644.—Zeylan.
foetens, *Benth. & Hook f. l. c.*—Java.
fulvicrys, *Thw. Enum. Pl. Zeyl.* 317.—Zeylan.
Galanga, Lour. Fl. Cochinch. 5 = Alpinia Galanga.
giganteum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 109.—Afr. trop.
glaberrimum, *Benth. & Hook. f. Gen.* iii. 644.—Malaya.
globosum, Lour. Fl. Cochinch. 4 = Alpinia globosa.
gracile, *Blume, Enum. Pl. Jav.* 49.—Java.
graminifolium, *Wall. Cat.* n. 6585.—Ind.-or.
graminifolium, *Thw. Enum. Pl. Zeyl.* 480.—Zeylan.
grandiflorum, Sm. Exot. Bot. ii. 108. t. 11 = A. Granum-paradisi.
Linn. Sp. Pl. 2.—Afr. trop.
heteranthum, *Blume, Enum. Pl. Jav.* 49.—Java.
hirsutum, Lam. Illustr. i. 134. t. 3 = Costus speciosus.
hirsutum, Lour. Fl. Cochinch. 5 = Alpinia hirsuta.
hypoleucum, *Thw. Enum. Pl. Zeyl.* 318.—Zeylan.
involucratum, *Benth. & Hook. f. Gen.* iii. 644.—Zeylan.
Koenigii, *Roem. & Schult. Syst.* i. 20.—Ind. or.
latifolium, *Afzel. Remad. Guin.* i. 5.—Afr. trop.
latifolium, Salisb. Prod. 4 = Curcuma Zedoaria.
Leconta, *Koen. ex Retz. Obs. fasc.* iii. 58.—Malacca.
limbatum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 110.—Afr. trop.
lingulifera, *Benth. & Hook f. Gen.* iii. 644.—Ind. or.
littorale, *Koen. ex Retz. Obs. fasc.* iii. 58.—Burma.
longicaspum, *Hook. f. in Journ. Bot.* vi. (1854) 296.—Afr. trop.
macrostemon, Sm. in Rees, Cyclop. xxxix. n. 3 = Clusii.
madagascariense, Lam. Illustr. t. 2. f. 1 = angustifolium.
magnificum, *Benth. & Hook. f. Gen.* iii. 644.—Ins. Maurit.
Mannii, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 110.—Afr. trop.
margaritatum, *Roxb. ex Sprang. Sekrad. & Link, Yahrb.* ii. (1811) 64.—Ind. or.
masticatorium, *Thw. Enum. Pl. Zeyl.* 317.—Zeylan.
maximum, *Roxb. in As. Res.* xi. (1810) 344.—Malaya.
medium, Lour. Fl. Cochinch. 4 = Hellenia alba.
Melegueta, *Rosc. Monog. Scitam.* t. 98.—Afr. trop.
Miaga, Thunb. Fl. Jap. 14 = Zingiber Miaga.
mioideum, Koen. ex Retz. Obs. fasc. iii. 61 = Zingiber Cassumunar.

AMOMUM :—
muricatum, *Budd. in Madr. Journ. Sc.* ser. III. i. (1864) 68.—Ind. or.
nemorale, *Benth. & Hook. f. Gen.* iii. 644.—Zeylan.
nemorosum, Boj. Hort. Maurit. 237 = Danielli.
nigrum, Rœusch. Nom. ed. III. 1 = Alpinia Allughas.
nutans, Roem. & Schult. Syst. i. Mant. 40 = Alpinia nutans.
palustre, *Afzel. Sierp. Med. Nov.* 9.—Afr. trop.
parviflorum, Presl, Rel. Haenk. ii. 112. t. 19 = Alpinia parviflora.
petiolatum, Lam. Encyc. i. 136 = Costus spicatus.
planatilobes, *Budd. in Madr. Journ. Sc.* ser. III. i. (1864) 68.—Ind. or.
pilosum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 110.—Afr. occ.
pterocarpum, *Thw. Enum. Pl. Zeyl.* 317.—Zeylan.
pulchellum, Thw. l. c. 318 = Cyphostigma pulchellum.
pyramidale, Lam. Encyc. i. 136 = Renealmia racemosa.
racemosum, Lam. Illustr. t. 2. f. 2 = A. Cardamon.
racemosum, Roxb. & Pav. Fl. Per. i. 2 = Renealmia Ruiziana.
Renealmia, Lam. Illustr. i. 5 = Renealmia exaltata.
repens, Sonner. Voy. Indes, iii. 272 = Alpinia Cardamomum
roscum, *Benth. & Hook. f. Gen.* iii. 644.—Malaya.
roseum, Roxb. Fl. Corom. ii. t. 196 = Zingiber roseum.
Rumphii, Sm. in Rees, Cyclop. xxxix. n. 15 = Elettaria muscea.
ruiseacea, *Benth. & Hook. f. Gen.* iii. 645.—Zeylan.
Sceptrum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 109.—Afr. trop.
Scottii, F. Muell. Fragm. viii. 24 = Elettaria Scottiana.
scyphiferum, *Korn. ex Retz. Obs. fasc.* iii. 58.—Malacca.
sericeum, Roxb. Hort. Beng. 1; Fl. Ind. i. 44 = dealbatum.
spirale, Hort. ex Steud. Nom. ed. II. i. 78 = Costus Pisonis.
spurium, J. F. Gmel. Syst. i. 6 = Zingiber Zerumbet.
spurium, *Roem. & Schult. Syst.* i. 30.—Ind. or.
striatum, Stokes, Bot. Comment. i. 169 = Renealmia sylvestris.
strobilaceum, *Sm. in Rees, Cyclop.* xxxix. n. 8.—Afr. trop.
subsericeum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 110.—Afr. trop.
subulatum, *Roxb. Pl. Corom.* t. 227 ; *Fl. Ind.* i. 43.—Reg. Himal.
sylvestre, Poir. Encyc. Suppl. v. 548 = Zingiber Zerumbet.
sylvestre, Sw. Prod. Veg. Ind. Occ. 11 = Renealmia sylvestris.
Taraca, Horan. Prod. 30 = Alpinia Allughas.
Taraeca, Roxb. ex Rosc. Monog. Scitam. in Obs. de Amomo = præc.
thyrsoideum, Ruiz & Pav. Fl. Per. i. 2. t. 2 = Renealmia thyrsoidea.
uliginosum, *Korn. ex Retz. Obs. fasc.* iii. 56.—Ind. or.
uncinatum, Stokes, Bot. Comment. i. 161 = A. Cardamon.
villosum, *Lour. Fl. Cochinch.* i. 4.—Ind. or. ; China.
vitellinum, *Lindl. in Journ. Hort. Soc.* iii. (1847) 245. —Ind. or. ?
xanthioides, *Wall. Cat.* n. 6567.—Burma.
xanthorhizos, Roxb. ex Steud. Nom. ed. II. i. 78 = Zingiber Cassumunar.
Zedoaria, Berg. Mat. Med. 41 = Curcuma Zedoaria.
Zerumbet, Koen. ex Retz. Obs. fasc. iii. 55 = Curcuma Zedoaria.
Zerumbet, Linn. Sp. Pl. 1.—Zingiber Zerumbet.
Zingiber, Blanco, Fl. Filip. 2 = Zingiber Blancoi.
Zingiber, Linn. Sp. Pl. 1 = Zingiber officinale.
Zinziber, Hill, Veg. Syst. xvi. 50 = præc.

AMONIA, Nestl. Mon. Potent. 17 (1816) = Agrimonia, Linn. (Rosac.).
agrimonioides, Steud. Nom. ed. I. 89, 19 = Agr. Eupatoria.

AMOORA, Roxb. Pl. Corom. iii. 54. t. 258 (1819). *MELIACEÆ*, Benth. & Hook. f. i. 384.
AMBERBOSIA, Rœmb. Hort. Beng. 67 (1814) ; Fl. Ind. ii. 213 (1832).
APHANAMIXIS, Blume, Bijdr. 166 (1826).
MONOCOMA, Griff. Notul. iv. 662 (1854).
NYEMQUA, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 18.

AMOORA:—
Oraoma, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 411.
Sphaerosacme, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 429 (1824).
amboinensis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 36.—Amboina.
Aphanamixis, *Schult. f. Syst.* vii. 1621.—Malaya.
auriculata, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 37 = cucullata.
Balsacsana, *C. DC. in DC. Monog. Phan.* i. 590.—N. Caled.
borneensis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 36.—Borneo.
casarana, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 560.—Ind. or.
Championii, Benth. & Hook. f. Gen. i. 835; Hiern, in Hook. f. l. c. 562 = Pseudocarapa Championii.
chittagonga, *Hiern, in Hook. f. l. c.* 559.—Ind. or.
cucullata, *Roxb. Pl. Corom.* iii. 54. t. 258.—Ind. or.
Cuninghana, *C. DC. in DC. Monog. Phan.* i. 580.—Ins. Philipp.
decandra, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 562.—Ind. or.
dysoxyloides, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 200.—Burma.
filiciformis *Wight, Illustr. Ind. Bot.* i. 147.—Ind. or.
Ganggo, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 128 = Aglaia Ganggo.
grandifolia, Walp. Rep. i. 429; C. DC. in DC. Monog. Phan. i. 581 = A. Aphanamixis.
Korthalsii, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 36.—Borneo.
lactescens, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 200.—Burma.
lanceolata, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 560.—Malacca.
Lawii, *Benth. & Hook. f. Gen.* i. 835.—Ind. or.
lasa, Wight & Arn. Prod. 119 = cucullata.
macrophylla, Nimmo, in J. Grah. Cat. Bomb. Pl. t. 31 = A. Rohituka.
Maingayi, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 562.—Malacca.
moulmeiniana, *C. DC. in DC. Monog. Phan.* i. 584.—Burma.
nitidula, *Benth. Fl. Austral.* i. 388.—Austral. Perrottetiana, *Steud. Nom.* ed. II. i. 89.—Ins. Philipp.
polystachya, Wight & Arn. Prod. i. 119 = A. Rohituka.
Rohituka, *Wight & Arn. l. c.*—Ind. or.
Roxburghiana, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 211 = Santiria laevigata.
rubescens, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 561.—Singapore.
rubiginosa, *Hiern, l. c.*—Malacca.
spectabilis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 37. Burma.
sumatrana, *Miq. l. c.* 35.—Sumatra.
Teysmanniana, Miq. Fl. Ind. Bat. Suppl. i. 508 = Aglaia Teysmanniana.
timoriensis, *Wight & Arn. ex Steud. Nom.* ed. II. i. 78.—Ins. Timor.
tomentosa, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 211 = Santiria tomentosa.
Vieillardi, *C. DC. in DC. Monog. Phan.* i. 591.—N. Caled.

AMOORIA, Walp. Rep. i. 640 (1843), sphalm. = AMORIA, C. Presl, = **Trifolium**, Tourn. (Legum.).

AMORDICA, Neck. Elem. i. 241 (1790) = **Momordica**, Tourn. (Cucurb.).

AMOREA, Moq. in Delile, Cat. Hort. Monsp. (1844) = Cyclolomea, Moq. (Chenopod.).
platyphylla, Delile, l. c. = C. platyphyllum.

AMOREBUXIA, Moq. & Sesse, ex DC. Prod. ii. 688 (1856). *BUXIHAE*, Benth. & Hook. f. i. 194.
Eurvanthe, Cham. & Schlecht. in Linnaea, v. (1830) 234.
australioides, *A. Gray, Pl. Wright.* i. 20.—Am. bor.
palmatifolia, Moq. & Sesse, ex DC. Prod. ii. 688 = Schiedeana?
Schiedeana, *Planch. in Hook. Lond. Journ. Bot.* vi. (1846) 146. t. 1.—Am. bor.
Wrightii, *A. Gray, Pl. Wright.* ii. 26.—Am. bor.

AMOREUXIA, Moq. in Mém. Soc. Hist. Nat. Monsp. (1826) = Cyclolomea, Moq. (Chenopod.).
platyphylla, Moq. l. c. = C. platyphyllum.

AMORGINE, Rafin. New Fl. Am. iv. 44 (1836) = **Achyranthes**, Linn. (Amaranth.).
albescens, Rafin. l. c.—Am. bor.

AMORIA, C. Presl, Symb. Bot. i. 45 (1830) = **Trifolium**, Tourn. (Legum.).
angulata, C. Presl, l. c. 47 = T. angulatum.
Biasolettiana, C. Presl, l. c. = T. repens.
caespitosa, C. Presl, l. c. = T. Thalii.
calycina, C. Presl, l. c. 48. t. 30 = T. calycinum.
carolinians, C. Presl, l. c. 47 = T. carolinianum.
elegans, C. Presl, l. c. = T. elegans.
hybrida, C. Presl, l. c. = T. hybridum.
isthmocarpa, C. Presl, l. c. 50 = T. Jamininanum.
macropoda, C. Presl, l. c. 51. t. 31 = T. macropodum.
Michelisana, C. Presl, l. c. 47 = T. Michelianum.
nigrescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362 = T. Michelianum.
obscurata, C. Presl, Symb. Bot. i. 47 = T. polymorphum.
pallescens, C. Presl, l. c. = T. pallescens.
parviflora, C. Presl, l. c. = T. parviflorum.
polymorpha, C. Presl, l. c. = T. polymorphum.
repens, C. Presl, l. c. = T. repens.
reflexa, C. Presl, l. c. = T. reflexum.
Thalii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362 = T. Thalii.
Vivianii, C. Presl, ex Steud. Nom. ed. II. i. 78 = T. caeruleum.

AMORPHA, Linn. Gen. ed. I. 229 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 492
californica, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 306.—Ins. Philipp.
canescens, *Nutt. in Fras. Cat.* (1813).—Am. bor.
carolinsana, Croom, in Am. Journ. Sc. xxv. (1834) 74 = fruticosa.
coerulea, Lodd. Cat. (1830) = fruticosa.
croceo-lanata, Wats. Dendrol. t. 139 = fruticosa?
cynostachya, M. A. Curt. ex Torr. & Gray, Fl. N. Am. i. 306 (cit. falso) = fruticosa.
elata, *Buchl. ex Schlecht. in Linnaea*, xxiv. (1851) 687.—Hab.?
elata, Hayne, Dendrol. Fl. 134 = fruticosa.
emarginata, Sweet, Hort. Brit. ed. I. 191. n. 466 = fruticosa.
fragrans, Sweet, Brit. Fl. Gard. iii. t. 242 = fruticosa.
fruticosa, *Linn. Sp. Pl.* 713.—Am. bor.
frutescens, Port. Mex. Bound. 53 = californica.
frutescens, Thunb. Fl. Jap. 278 = Cladrastis amurensis.
Gaertneri, Hort. ex C. Koch, Dendrol. i. 70 = fruticosa.
Gaertneri, Hort. ex C. Koch, l. c. Index, 701 = praec.
glabra, Desf. Tabl. Hort. Par. ed. I. 193 = fruticosa.
glandulosa, *Blanco, Fl. Filip.* ed. I. 555.—Ins. Philipp. herbacea, *Walt. Fl. Carol.* 179.—Am. bor.
herbacea, Schlecht. Ind. Sem. Hort. Hal. (1848) 8.—Cf. Linnaea, xxiv. (1851) 185 = fruticosa.
humilis, Tausch, in Flora, xxi. (1838) ii. 750 = fruticosa?
laevigata, *Nutt ex Torr. & Gray, Fl. N. Am.* i. 306.—Am. bor.
Lewisii, Lodd. ex Loud. Hort. Brit. Suppl. ii. 609 = fruticosa.
Lodwigii, Hort. ex C. Koch, Dendrol. i. 70 = fruticosa.
lutea, *Rafin. Fl. Ludov.* 105.—Am. bor.
microphylla, *Pursh, Fl. Am. Sept.* ii. 466.—Am. bor.
nana, Nutt. in Fras. Cat. (1813) = fruticosa?
nonperforata, Schk. Bot. Handb. ii. 333 = fruticosa.
ornata, Wender. Ind. Sem. Hort. Marb. (1855) = fruticosa.
pauciflora, *Torr. & Gray, Fl. N. Am.* i. 306.—Am. bor.
pedalis, *Blanco, Fl. Filip.* ed. I. 553.—Ins. Philipp.
pendula, Carr. in Rev. Hortic. (1868) 340 = fruticosa?
perforata, Schk. Bot. Handb. ii. 333 = fruticosa.
pubescens, Schlecht. in Linnaea, xxiv. (1851) 691 = fruticosa.
pubescens, Willd. Berl. Baumz. 17 = herbacea.
pumila, Michx. Fl. Bor. Am. ii. 64 = herbacea.
pumila, Schlecht. Ind. Sem. Hort. Hal. (1848) 8.—Cf. Linnaea, xxiv. (1851) 185 = fruticosa.
punctata, *Rafin. New. Fl. Am.* iii. 14.—Am. bor.
Rabilas, *Lex. in La Llave & Lex. Nov. Veg. Descr. fasc.* i. 23.—Mexic.
Remeriana, Scheele, in Linnaea, xxi. (1848) 461 = paniculata.

AMORPHA:—
tenuesemis, Schult. in Kunze, Del. Sem. Hort. Lips. (1848) = fruticosa.
texana, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 452 = laevigata.
tomentosa, *Rafin. Fl. Ludov.* 105.—Am. bor.

AMORPHOCALYX, Klotzsch, in Rich. Schomb. Faun. & Fl. Guian. 1104, nomen, (1848) (Legumin.). Roraimae, *Klotzsch, l. c.*—Guiana.

AMORPHOPHALLUS, Blume, in Batav. Diar. (1825); ex Decne. Herb. Tim. Descr. 88 (1835). *AROIDEAE*, Benth. & Hook. f. iii. 970.
Brachyspatha, Schott, Syn. Aroid. 85 (1856).
Candarum, Reichb. ex Schott, Melet. i. 17 (1832).
Conophallus, Schott, Syn. Aroid. 84 (1856).
Corynophallus, Schott, in Oestr. Bot. Wochenbl. (1857) 389.
Hansaila, Schott, in Oestr. Bot. Zeitschr. (1856) 82.
Hydrosme, Schott, in Oestr. Bot. Wochenbl. (1857) 389.
Kunda, Rafin. Fl. Tellur. ii. 82 (1836).
Protiniphallus, Hook. f. Bot. Mag. t. 6195 (1875).
Pythion, Mart. in Flora, xiv. (1831) 409.
Tapinophallus, Baill. Dict. Bot. i. Ic. sine desc. (1877).
Beccari, *Engl. in Engl. Bot. Jahrb.* i. (1881) 182.—Sumatra.
Blumei, Schott, in Ann. Mus. Lugd. Bat. i. 194 = gigantenus.
bulbifer, *Blume, Rumphia*, i. 148.—Ind. or.
campanulatus, *Blume, ex Decne. Herb. Timor.* 88.—As. trop.
Chatty, *André, Illustr. Hortic.* xix. (1879) 861.—Ind. or.
commutatus, *Engl. in DC. Monog. Phan.* ii. 819.—Ind. or.
consimilis, *Blume, Rumphia*, i. 149.—Afr. trop.
decurrens, *Kunth, Enum. Pl.* iii. 581.—Ins. Philipp.
difformis, Blume, Rumphia, i. 149 = Anchomanes difformis.
dubius, *Blume, l. c.* 142.—Ind. or.; Malaya.
Fontanesii, *Kunth, Enum. Pl.* iii. 35.—Afr. trop.
Gigas, *Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind.* xxiv. (1862) 329.—Sumatra.
giganteus, *Blume, Rumphia*, i. 144. t. 34.—Ind. or.; Malaya.
gracilis, *Engl. in Engl. Bot. Jahrb.* i. (1881) 183.—Sumatra.
hirsutus, *Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind.* xxiv. (1862) 332.—Java.
Imperialis, *Hort. cf. Gard. Chron.* (1883) i. 510.—Ins. Philipp.
Konjac, C. Koch, in Berl. Allg. Gartenz. (1858) 166 = Rivieri.
Lacourii, *Linden & André, in Illustr. Hortic.* xxv. (1878) t. 816.—Cochinch.
leonensis, *Lem. in Fl. des Serres*, Sér. I. ii. (1846) t. 161.—Afr. trop.
longistylus, *Lours, Rep. Andam.* 50, nomen.—Ins. Andaman.
lyratus, *Kunth, Enum. Pl.* iii. 34.—Ind. or.
margaritifer, Kunth, l. c. = Plesmonium margaritiferum.
mossambicensis, *Klotzsch, ex Garcke, in Peters, Reise Mossamb. Bot.* 509.—Afr. trop.
Mulleri, *Blume, Rumphia*, i. 148.—Java.
nivosus, Lem. Illustr. Hortic. xii. (1865) t. 424 = Dracontium asperum.
palmaeformis, Dur. ex Riv. in Act. Linn. Soc. Bord. xxviii. (1869) 15; Rivière, Ul. Amorph. 4, 5 = Rivieri.
papillosus, *Hort. ex Rafarin, in Rev. Hortic.* (1871) 476.—Hab.?
Peyri, *cf. Gard. Chron.* (1880) ii. 190.—Ins. Seychelles.
planeus, *Teijsm & Binn. in Nat. Tijdschr. Nederl. Ind.* xxiv. (1862) 331.—Sumatra.
pumetularus, *Blume, Rumphia*, i. 148.—Java.
Rivieri, *Dur. ex Rivière, in Act. Linn. Soc. Bord.* xxviii. (1869) 15.—Cochinch. ? Japon.
sativus, *Blume, Rumphia*, i. 146.—Ins. Molucc.
spectabilis, *Engl. in DC. Monog. Phan.* ii. 816.—Java.
sylvaticus, Kunth, Enum. Pl. iii. 34 = Synantherias silvaticus.
Titanum, *Becc. in Bull. Soc. Tosc. Ortic.* (1879) 66.—Sumatra.
tuberculiger, *Engl. in DC. Monog. Phan.* ii. 817.—Reg. Himal.
variabilis, *Blume, Rumphia*, i. 146. t. 35.—Java; Ins. Philipp.

AMORPHOPHALLUS:—
virosus, *N. E. Br. in Gard. Chron.* (1886) i. 759.—Siam. *Wallichii*, Regel, Gartenfl. (1861) 322 = Dracontium asperum.
sebrinus, *Hort.—Cf. Gard. Chron.* (1865) 485.—Hab.? *zeylanicus*, *Blume*, *Rumphia*, i. 148.—Java; Zeylan.

AMORPHOSPERMUM, F. Muell. Fragm. vii. 119 (1870) = Leucoma, Juss. (Sapotac.).
austilguum, F. Muell. l. c.—Austral.

AMOSA, Neck. Elem. ii. 459 (1790) = Inga, Plum. (Legum.).

AMOUREUXIA, C. Muell. in Walp. Ann. iv. 840 (1857), sphalm. = Amoreuxia, Moç. & Sesse (Bixin.).

AMPALIS, Boj. Hort. Maurit. 291 (1837). *URTICACEAE*, Benth. & Hook. f. iii. 365.
madagascariensis, *Boj. l. c.*—Madag.

AMPELANUS, Rafin. in Am. Monthly Mag. (1819) 192 = Ampelopsis, Nutt. (Asclep.).

AMPELOCERA, Klotzsch, in Linnaea, xx. (1847) 541. *URTICACEAE*, Benth. & Hook. f. iii. 358.
cubensis, *Griseb. Cat. Pl. Cub.* 57.—Ind. occ. Ruizii, *Klotzsch, in Linnaea*, xx. 543.—Peruv.

AMPELOCISSUS, Planch. in Journ. Vigne Am. (1884) 374 = Vitis, Linn. (Ampelid.).
abyssinica, Planch. in Journ. Vigne Am. (1885) 24 = V. abyssinica.
acetosa, Planch. l. c. 96.—Austral.
aculeata, Planch. l. c. 95 = V. aculeata.
angolensis, Planch. l. c. 48 = V. angolensis.
arcuata, Planch. l. c. = V. arcuata.
asarifolia, Planch. l. c. 39 = V. asarifolia.
Baheri, Planch. l. c. 80 = V. Schimperiana.
bombycina, Planch. l. c. 31 = V. bombycina.
cavicaulis, Planch. l. c. 82 = V. cavicaulis.
Chantinii, Planch. l. c. 37 = V. Chantinii.
cinnamochroa, Planch. l. c. 31.—Am. trop.
concinna, Planch. l. c. 48 = V. concinna.
diasecta, Planch. l. c. = V. diasecta.
divaricata, Planch. l. c. (1884) 375 = V. divaricata.
elephantina, Planch. l. c. 379.—Ins. Borbon.
Hermanni, Planch. l. c. 373.—Cochinch.
heracleifolia, Planch. l. c. (1885) 48 = V. heracleifolia.
indica, Planch. l. c. (1884) 376 = V. indica.
ipomoeaefolia, Planch. l. c. (1885) 26 = V. ipomoeaefolia.
latifolia, Planch. l. c. (1884) 374 = V. latifolia.
Lecardii, Planch. l. c. (1885) 29 = V. Lecardii.
leonensis, Planch. l. c. 30 = V. leonensis.
Lepervierii, Planch. l. c. 44 = V. multistriata.
mossambicensis, Planch. l. c. 49 = V. mossambicensis.
obtusata, Planch. l. c. 48 = V. obtusata.
platanifolia, Planch. l. c. 47 = V. platanifolia.
rugosa, Planch. l. c. (1884) 374 = V. lanata, indica.
salmonea, Planch. l. c. (1885) 31 = V. salmonea.
sarcocephala, Planch. l. c. 46 = V. sarcocephala.
Schimperiana, Planch. l. c. 38 = V. Schimperiana.
sikkimensis, Planch. l. c. (1884) 375 = V. sikkimensis.
tomentosa, Planch. l. c. 375 = V. tomentosa.

AMPELODAPENE, Meisen. in DC. Prod. xv. i. 81 (1864). *LAURINEAE*, Benth. & Hook. f. iii. 198.
arunciflora, *Meisen. l. c.*—Bras.
dasyantha, *Meisen. l. c.*—Bras.
macrophylla, *Meisen. l. c.*—Bras.

AMPELODESMA, Banw. Agrost. 78. t. 15. f. 11 (1819). *GRAMINEAE*, Benth. & Hook. f. iii. 1178.
australe, Brongn. in Duperr. Voy. Bot. 81 = Arundo pilosa.
bicolor, *Kunth, Gram.* i. 79.—Afr. trop.
effusum, *Steud. Syn. Pl. Gram.* 196.—Afr. bor.
festucoides, Steud. l. c. = tenax.
tenax, *Link, Hort. Berol* i. 136.—Reg. Mediterr.

AMPELODESMOS, Link, Hort. Berol. i. 136 (1827) = prsec.

AMPELODESMUS, Woods, in Trans. Linn. Soc. xviii. (1836) 24 = prsec.

AMPELOPLIS, Rafin. Sylva Teller. 88 (1838) = Rhamnus, Linn.
chinensis, Rafin. l. c. = R. theezans.

AMPELOPSIS, [Rich. in] Michx. Fl. Bor. Am. i. 159 (1803) ex parte = Vitis, Linn. (Ampelid.).
acerifolia, Bunge, Enum. Pl. Chin. Bor. 12 = V. serjaniaefolia.
africana, Steud. Nom. ed. II. i. 219 = V. africana.
bipinnata, Michx. Fl. Bor. Am. i. 160 = V. arborea.
Bodria, DC. Prod. i. 633 = V. africana.
brevipedunculata, Maxim. ex Trautv. in Act. Hort. Petrop. vii. (1883) 176.—As. or.
citrulloides, Lebas, in Rev. Hortic. (1875) 179.—Hab.?
cordata, Michx. Fl. Bor. Am. i. 159 = V. indivisa.
cordifolia, Rafin. Med. Fl. ii. 122 = prsec.
dissecta, Carr. in Rev. Hortic. (1868) 10 = V. serjaniaefolia.
hederacea, DC. Prod. i. 633 = V. hederacea.
heptaphylla, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 450 = V. hederacea.
heptaphylla, Roem. & Schult. Syst. v. 321 = V. heptaphylla.
heterophylla, Blume, Bijdr. 194 = V. Landuk.
heterophylla, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 196 = V. heterophylla.
himalayana, Royle, Illustr. Himal. Bot. 149 = V. himalayana.
hirsuta, Donn, Hort. Cantab. 166 = hederacea.
humulifolia, Bunge, Enum. Pl. Chin. Bor. 12 = V. heterophylla.
indica, Blume, Bijdr. 198 = V. indica.
japonica, Hort.—Cf. Gard. Chron. (1868) 1118. = V. inconstans.
latifolia, Tausch, in Flora, xxi. (1838) ii. 736; Kostel. Allg. Med.-Pharm. Fl. iv. 1198 = V. latifolia.
lucida, Carr. in Rev. Hortic. (1868) 39 = V. serjaniaefolia.
minima, Hort.—Cf. Gard. Chron. (1878) 1469 = V. inconstans.
napyforma, Carr. in Rev. Hortic. (1870–71) 16.—China.
neilgherrensis, Wight, Spicil. Neilgh. i. 32. t. 36; et Ic. t. 966 = V. himalayana.
palmiloba, Carr. in Rev. Hortic. (1867) 451.—China.
pinnata, Roem. & Schult. Syst. v. 322 = V. pinnata.
pubescens, Schlecht. in Linnaea, x. (1836) 251 = V. pubescens.
quinquefolia, Michx. Fl. Bor. Am. i. 159. t. 162 = V. hederacea.
Regeliana, Carr. Rev. Hortic. (1866) 440.—Japon!
sempervirens, Hort.—Cf. Gard. Chron. (1883) ii. 491 & 590 = V. striata.
serjaniaefolia, Bunge, Enum. Pl. Chin. Bor. 12 = V. serjaniaefolia.
ternata, DC. Prod. i. 633 = V. tomentosa.
tricuspidata, Sieb. & Zucc. in Abh. Akad. Muench. iv. (1846) 196 = V. inconstans.
triloba, Carr. in Rev. Hortic. (1868) 39 = V. serjaniaefolia.
triparitia, Carr. l. c. = V. serjaniaefolia.
tuberosa, Carr. l. c. (1870–71) 16 = V. serjaniaefolia.
Veitchii, Hort.—Cf. Gard. Chron. (1868) 814 = V. inconstans.

AMPELOSICYOS, (AMPELOSYCIOS), Thou. Hist. Vég. Isles Afr. suscr. 68. t. 22 (1807) = Telfairia, Hook. (Cucurb.).
scandens, Thou. l. c. = T. pedata.

AMPELYGONUM, Lindl. in Bot. Reg. (1838) Misc. 62 = Polygonum, Linn.
chinense, Lindl. l. c. = P. chinense.
tinctorium, Steud. Nom. ed. II. i. 79 = P. tinctorium.

AMPEREA, A. Juss. Euph. Tent. 25. t. 10 (1824) *EUPHORBIACEAE*, Benth. & Hook. f. ii. 956.
conferta, *Benth. Fl. Austral.* vi. 83.—Austral.
cuneifornis, F. Muell. ex Baill. Rend. Gén. Euph. 455 = spartioides.
ericoides, *A. Juss. Tent. Euph.* 112. t. 10.—Austral.
micrantha, *Benth. Fl. Austral.* vi. 83.—Austral.
protensa, *Nees, in Lehm. Pl. Preiss.* ii. 229.—Austral.
racemiformis, Klotzsch, in Lehm. l. c. i. 176 = ericoides.
spartioides, *Brongn. in Duperr. Voy. Bot.* 236. t. 49.—Austral.

AMPHERA:—
subnuda, *Nees, in Lehm. Pl. Preiss.* ii. 229.—Austral.
volubilis, *F. Muell. ex Benth. Fl. Austral.* vi. 83.—Austral.

AMPHANIA, Banks, ex DC. in Mém. Soc. Phys. Genèv. i. (1821) 408 = Turnstroemia, Linn.
integrifolia, Soland. ex DC. l. c. 410 = T. elliptica.

AMPHEREPHIS, H. B. & K. Nov. Gen. et Sp. iv. 81. t. 314, 315 (1820) = Centrostachys, Cass. (Compos.).
aristata, H. B. & K. l. c. 95 = C. punctatum.
indica, Wall. [Cat. n. 2956]; Less. in Linnaea, vi. (1831) 686 = C. reticulatum.
intermedia, Link, Abhdl. 5. t. 29 = C. intermedium.
mollis, Wall. Cat. n. 2957 = C. molle.
mutica, H. B. & K. Nov. Gen. et Sp. iv. 96, 315 = C. muticum.
pilosa, Cass. ex Steud. Nom. ed. II. i. 324, in syn., = C. muticum.
pulicaeria, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 4 = Oiospermum involucratum.
pulchella, Cass. ex Steud. Nom. ed. II. i. 79 = C. pulchellum.
senegalensis, Less. in Linnaea, vi. (1831) 687 = Heyderia truncata.
Wightiana, Wall. Cat. n. 2958 = Carpesium cernuum.

AMPHACHYRIS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 313. *COMPOSITAE*, Benth. & Hook. f. ii. 250.
AMPHIPAPPUS, Torr. & Gray, in Proc. Bost. Soc. i. (1844) 210.
dracunculoides, *Nutt. l. c.*—Am. bor.
Fremontii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 633.—Calif.

AMPHIANTHUS, Torr. in Ann. Lyc. N. York, iv. (1837) 92. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 956.
pusillus, *Torr. l. c.*—Am. bor.

AMPHIBECIS, Schrank, in Syll. Ratisb. i. (1824) 86 = Centratherum, Cass. (Compos.).
viridescens, Schrank, l. c. = C. intermedium.

AMPHIBLEMMA, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) t. 7; xv. (1851) 50. *MELASTOMACEAE*, Benth. & Hook. f. i. 754.
cymosum, *Naud. l. c. xv.* (1851) 51.—Afr. trop.
molle, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 456; *in Trans. Linn. Soc.* xxviii. (1871) 78.—Afr. trop.
setosum, *Hook. f. ii. cc.*—Afr. trop.

AMPHIBOLIS, C. Agardh, Sp. Alg. ii. i. 474 (1822). *NAIADACEAE*, Benth. & Hook. f. iii. 1015.
GRAUMULLERA, Reichb. Consp. 48 (1828).
antarctica, Aschers. in Linnaea, xxxv. (1867–68) 164 = Cymodocea antarctica.
Gardneri, *C. Agardh, Sp. Alg.* ii. i. 474.—Austral.
sosterae folia, C. Agardh, l. c. 475 = Cymodocea antarctica.

AMPHIBROMUS, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 430. *GRAMINEAE*, Benth. & Hook. f. iii. 1161.
flaitana, *T. Kirk, in Trans. N. Zeal. Inst.* xvi. (1884) 374. t. 28.—N. Zel.
Neesii, *Steud. Syn. Pl. Gram.* 298.—Austral.

AMPHICALA, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 411 = Calea, Linn. (Compos.).
fruticosa, Gardn. l. c. 413 = C. rotundifolia.
gentianoides, Gardn. l. c. = Gaiesoppapus gentianoides.

AMPHICALYX, Blume, Fl. Jav. Pracf. p. vii (1828) = Diplycosia Blume (Ericac.).

AMPHICARPAEA, Ell. in Journ. Acad. Philad. i. (1818) 272 (Amphicarpa). *LEGUMINOSAE*, Benth. & Hook. f. i. 589.
CRYPTOCOSTIS, Spreng. Syst. iii. 818 (1826).
FALCATA, J. F. Gmel. Syst. 1181 (1791).
LOBOMON, Rafin. New Fl. Am. i. 81 (1836).
NAVIA, Rafin. in Med. Repos. N. York, v. (1808) 353.
TETRODEA, Rafin. New Fl. Am. i. 81 (1836).
clinia, *Rafin. l. c.* 82.—Am. bor.

AMPHICARPÆA :—
comosa, G. Don, in Lond. Hort. Brit. 314 = monoica.
cuspidata, *Rafin. New Fl. Am.* i. 83.—Am. bor.
dolichoides, Rafin. l. c. = monoica.
Edgeworthii, *Benth. in Miq. Pl. Jungh.* 331.—Reg. Himal.
Elliotii, Rafin. New Fl. Am. i. 83 = monoica.
ferruginea, Benth. in Miq. Pl. Jungh. 331 = Shuteria ferruginea.
heterophylla, Rafin. New Fl. Am. i. 83 = monoica.
monoica, *Ell. in Journ. Acad. Philad.* i. (1818) 372 ; *& ex Nutt. Gen. Am.* ii. 115.—Am. bor.
Pluheri, *Torr. & Gray, Fl. N. Am.* i. 292.—Am. bor.
sarmentosa, Ell. ex Nutt. Gen. Am. ii. 114 = monoica.
trisperma, Baker, in Hook. f. Fl. Brit. Ind. ii. 181, in obs. = Edgeworthii.
villosa, Rafin. New Fl. Am. i. 83 = monoica.

AMPHICARPON, Rafin. in Am. Monthly Mag. (Jan. 1818) 175 = seq.

AMPHICARPUM, Kunth, Gram. i. 28 (1829). *GRAMINEÆ*, Benth. & Hook. f. iii. 1099.
floridanum, *Chapman*, *Fl. S. U. St.* 572.—Florida. Purshii, *Kunth*, *Gram.* i. 28.—Am. bor.

AMPHICOME, Royle, Illustr. Himal. Pl. 296. t. 72. f. 1 (1886). *BIGNONIACEÆ*, Benth. & Hook. f. ii. 1049.
argute, *Royle*, *l. c.*—Reg. Himal.
Emodi, *Royle*, *ex Lindl. Bot. Reg.* (1838) t. 19.—Reg. Himal.

AMPHIDERRIS, Spach, Hist. Vég. Phan. x. 402 (1841) = Ortiea, R. Br. (Proteac.).

AMPHIDETHS, Fourn. in Mart. Fl. Bras. vi. iv. 218 (1885). *ASCLEPIADEÆ*.
lacinistus, *Fourn. l. c.* 214.—Bras.
quinquedentatus, *Fourn. l. c.* 218.—Bras.

AMPHIDONAX, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836) = Arundo, Linn. (Gramin.).
bengalensis, Nees, l. c. = Arundo bengalensis.

AMPHIDONAX, Nees, ex Steud. Syn. Pl. Gram. 197 = Ecnkeria, Trin. (Gramin.).
bifaria, Nees, ex Steud. l. c.—Japon.
Heynii, Nees, ex Steud. l. c. = Z. elegans.
obtusiflora, Thw. Enum. Pl. Zeyl. 370 = Z. obtusiflora.
tenella, Wight & Arn. ex Steud. Syn. Pl. Gram. 197. —Ind. or.

AMPHIDOXA, DC. Prod. vi. 246 (1837). *COMPOSITÆ*, Benth. & Hook. f. ii. 300.
filagines, *Ficalho & Hiern*, *in Trans. Linn. Soc.* Ser. II. ii. (1881) 31.—Afr. trop.
gnaphalodes, *DC. l. c.*—Afr. austr.

AMPHIGENES, Janka, in Linnæa, xxx. (1859–60) 619 = Festuca, Linn. (Gramin.).
carpathica, Janka, l. c. = F. carpathica.
nutans, Janka, l. c. = F. nutans.

AMPHIGLOSSA, DC. Prod. vi. 258 (1837). *COMPOSITÆ*, Benth. & Hook. f. ii. 324.
alopecuroides, *Sch. Bip. in Pollichia*, xvii–xix. (1861) 183.—Ab. austr.
callanoides, *DC. Prod.* vi. 259.—Afr. austr.
corrudæfolia, *DC. l. c.* 258.—Afr. austr.
nitidula, *DC. l. c.* 259.—Afr. austr.
perotrichoides, DC. l. c. 258 = Pterothrix perotrichoides.
tomentosa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 276. —Afr. austr.
trifaria, DC. Prod. vi. 258 = Pterothrix spinescens.

AMPHIGLOTTIS, Salisb. in Trans. Hort. Soc. i. (1812) 294 = Epidendrum, Linn. (Orchid.).
luridæ, Salisb. l. c. = E. ancepe.
secundæ, Salisb. l. c. = E. secundum.

AMPHILOBIUM, Lond. Hort. Brit. 242 (1830), sphalm. = Amphilophium, H. B. & K. (Bignon.).

AMPHILOCHIA, Mart. Nov. Gen. et Sp. i. 127. t. 77 (1824) = Qualea, Aubl. (Vochysiac.).

AMPHILOCHIA :—
acuminulata, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 343 = Q. cryptantha.
cordata, Mart. Nov. Gen. et Sp. i. 128 = Q. cordata.
dichotoma, Mart. l. c. 128. t. 77 = Q. dichotoma.
Lundii, Warm. in Kjoeb. Vidensk. Meddel. (1867) 27 = Q. Lundii.
marginata, Miq. in Linnæa, xxii. (1849) 796 = Q. marginata.

AMPHILOPHIUM, Kunth, in Journ. Phys. lxxxvii. (1818) 451. *BIGNONIACEÆ*, Benth. & Hook. f. ii. 1039.
glandulisferum, Mart. ex DC. Prod. ix. 193 = Vauthierl.
macrophyllum, *H. B. & K. Nov. Gen. et Sp.* iii. 149. —N. Granat.
molle, *Cham. & Schlecht. in Linnæa*, v. (1830) 120. —Mexic.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* iii. 149. t. 219. —N. Granat.
paniculatum, *H. B. & K. l. c.*—Honduras.
pubescens, Spreng. Syst. ii. 886 = Bignonia fallax.
Vauthieri, *DC. Rev. Bignon.* (1838) ; *Prod.* ix. 193.—Bras.

AMPHINOMIA, DC. Prod. ii. 522 (1825) = Lotononis, Eckl. & Zeyh. (Legumin.).
decumbens, DC. l. c. = L. polycephalus.

AMPHIOLANTHUS, Griseb. Cat. Pl. Cub. 186 (1866) = Mitranthemum, Michx. (Scrophul.).
arenaroides, Griseb. l. c. 187.—Cuba.
bryoides, Griseb. l. c. 186.—Cuba.

AMPHION, Salisb. Gen. Pl. Fragm. 66 (1866) = Semele, Kunth (Liliac.).

AMPHIONE, Rafin. Fl. Tellur. iv. 79 (1836) = Ipomæa, Linn. (Convolv.).
asarifolia, Rafin. l. c. = I. asarifolia.
campodia, Rafin. l. c.—Hab. ?
cymosa, Rafin. l. c. = I. cymosa.
lobata, Rafin. l. c.—Hab. ?
splendens, Rafin. Sylva Tellur. 130 = Argyreia splendens.
venosa, Rafin. Fl. Tellur. iv. 79 = I. venosa.

AMPHIPAPPUS, Torr. & Gray, in Proc. Bost. Soc. i. (1844) 210 ; Bost. Journ. Nat. Hist. v. (1847) 107 = Amphiachyris, Nutt. (Compos.).
Fremontii, Torr. & Gray, l. c. 108 = Amphiachyris Fremontii.

AMPHIPLEIS, Rafin. Fl. Tellur. iii. 75 (1836) = Nicotiana, Linn. (Solanac.).
fetida, Rafin. l. c.—Hab. ?
quadrivalvis, Rafin. l. c. = N. quadrivalvis.

AMPHIPOGON, R. Br. Prod. 175 (1810). *GRAMINEÆ*, Benth. & Hook. f. iii. 1145.
Argopogon, Beauv. Agrost. 122. t. 22. f. 4 (1812), non Labill.
Gamelythrum, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 415.
Pentascraspedon, Steud. Syn. Pl. Gram. 151 (1855).
asperascens, R. Br. Prod. 175 (1810) = strictus.
Brownei, F. Muell. Fragm. viii. 201 = strictus.
caricinus, F. Muell. in Linnæa, xxv. (1852) 445 = strictus.
cygnorum, *Nees*, *in Lehm. Pl. Preiss.* ii. 100.—Austral.
debilis, *R. Br. Prod.* i. 175.—Austral.
gracilis, Nees, in Lehm. Pl. Preiss. ii. 101 = strictus.
leptoides, F. Muell. Fragm. viii. 201 = cygnorum.
laguroides, *R. Br. Prod.* i. 175.—Austral.
strictus, *R. Br. l. c.*—Austral.
turbinatus, *R. Br. l. c.*—Austral.

AMPHIREPHIS, Nees & Mart. in Nov. Act. Acad. Nat. Cur. xii. (1824) 4 = AMPHERREPHIS, H. B. & K. = Centratherum, Cass. (Compos.).

AMPHIRHAPIS, DC. Prod. v. 348 (1836) = Inula, Linn. ; Microglossa, DC. & Schlagg., Linn. (Compos.).
albescens, DC. Prod. v. 348 = M. albescens.
chinensis, *Sch. Bip. in Flora*, xxxv. (1852) 58 = Solidago Virga-aurea.

AMPHIRHAPIS :—
cuspidata, DC. Prod. v. 348 = Inula cuspidata.
heterotricha, DC. l. c. = I. empatorioides.
indica, ‘Less.’ ex Hook. f. Fl. Brit. Ind. iii. 327, in syn. = Centratherum reticulatum.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 172.—Japon.
leiocarpa, Benth in Hook. Lond. Journ. Bot. i. (1842) 488 = Solidago Virga-aurea.
mollis, ‘Wall.’ ex Hook. f. Fl. Brit. Ind. iii. 327, in syn. = Centratherum molle.
pedunculosa, DC. Prod. v. 344 = Aster asperulus.
pubescens, DC. l. c. 348 = Solidago Virga-aurea.
rubricaulis, DC. l. c. = I. rubricaulis.
Wightiana, Hook. f. Fl. Brit. Ind. iii. 301 = Carpesium abrotanoides.
zeylanica, DC. Prod. vii. 279 = M. zeylanica.
Zollingeriana, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 136, nomen.—Malaya.

AMPHIRRHOX, Spreng. Syst. Cur. Post. 51 (1827). *VIOLARIEÆ*, Benth. & Hook. f. i. 118.
Braddleya, Vell. Fl. Flum. ii. t. 140 (1827).
Spathularia, St. Hil. Pl. Rem. Brit. t. 28 (1824).
latifolia, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 376.—Bras.
longifolia, *Spreng. Syst. Cur. Post.* 51.—Bras.
surinamensis, *Eichl. in Mart. Fl. Bras.* xiii. i. 377.—Bras.

AMPHISCOPIA, Nees, in Wall. Pl. As. Rar. iii. 112 (1832) = Justicia, Linn. (Acanth.).
aequilabris, Nees, in DC. Prod. xi. 358.—Bras.
Beyrichii, Nees, in Mart. Fl. Bras. ix. 130 = J. pummularia.
ciliata, Moricand, ex Nees, l. c.—Bras.
ciliata, Nees, in DC. Prod. xi. 359.—Reg. Argent.
comosa, Nees, l. c. 360 = J. comosa.
cuneyfolia, Nees, in Mart. Fl. Bras. ix. 130 = J. cuneifolia.
inficiens, Nees, in DC. Prod. xi. 357 = J. inficiens.
lancifolia, Nees, l. c.—Peruv.
Martiana, Nees, in Mart. Fl. Bras. ix. 130.—Bras.
Middletoni, Nees, in DC. Prod. xi. 356.—Afr. trop.
montana, Nees, l. c. 359.—Bras.
Moricandiana, Nees, l. c. 357.—Bras.
Pohliana, Nees, l. c. 359.—Bras.
polystachya, Nees, l. c. 356 = J. polystachya.
retusa, Nees, l. c. 357 = J. retusa.
rosea, Nees, l. c. 358 = Eranthemum roseum ?
serpens, Nees, l. c. 357 = Hypoestes serpens ?
strobilacea, Nees, l. c. 358.—Bras.
venosa, Nees, l. c. 356 = J. crinata.

AMPHISCOPIUM, St. Lag. in Ann. Soc. Bot. Lyon, vii. 119 (1880) = præc.

AMPHISTELMA, Griseb. Fl. Brit. W. Ind. 417 (1864) = Vincetoxicum, Moench (Asclep.).
angulatum, Fourn. in Mart. Fl. Bras. vi. iv. 226.—Bras.
aphyllum, Fourn. l. c.—Bras.
ephedroides, Griseb. Cat. Pl. Cub. 174.—Cuba.
exsertum, Griseb. in Goett. Abh. xxiv. (1879) 229.—Reg. Argent.
ferrugineum, Fourn. in Mart. Fl. Bras. vi. iv. (1885) 224.—Bras.
filiforme, Griseb. Fl. Brit. W. Ind. 418.—Ind. occ.
graminifolium, Griseb. Cat. Pl. Cub. 174.—Cuba.
leptoclados, Griseb. Fl. Brit. W. Ind. 418 = V. leptoclados.
linearifolium, Griseb. Cat. Pl. Cub. 175.—Cuba.
melanthum, Fourn. in Mart. Fl. Bras. vi. iv. 226 = V. melanthum.
parviflorum, Fourn. l. c. 224.—Bras.
Riedelii, Fourn. l. c. 225.—Bras.
salzmannum, Wright, ex Griseb. Cat. Pl. Cub. 175.—Cuba.
Sellowianum, Fourn. in Mart. Fl. Bras. vi. iv. 224.—Bras.
streptolobium, Fourn. l. c. 225.—Bras.
tomentosum, Fourn. l. c. 226.—Bras.

AMPHITECNA, Miers, in Trans. Linn. Soc. xxvi. (1868) 168. *BIGNONIACEÆ*.
macrophylla, *Miers*, *in Baill. in Rev. Hortic.* (1868) 465.—Am. centr.
nigripes, *Baill. l. c.*—Am. trop. ?

AMPHITHALEA, Eckl. & Zeyh. Enum. 167 (1836).
LEGUMINOSAE, Benth. & Hook. f. i. 472.
CAPHIANTHA, Eckl. & Zeyh. Enum. 171 (1836).
EPISTEMUM, Walp. in Linnæa, xiii. (1839) 473.
IGENHOUSSIA, K. Mey. Comm. Pl. Afr. Austr. 20 (1856).
ciliaris, Eckl. & Zeyh. Enum. 169 = Coelidium ciliare.
caniflora, *Eckl. & Zeyh. l. c.* 167.—Afr. austr.
densa, *Eckl. & Zeyh. l. c.*—Afr. austr.
densiflora, Eckl. & Zeyh. l. c. 168 = ericæfolia.
ericæfolia, *Eckl. & Zeyh. l. c.* 169.—Afr. austr.
hilaris, Eckl. & Zeyh. l. c. = ericæfolia.
humilis, Eckl. & Zeyh. l. c. 168 = intermedia.
incurvifolia, Eckl. & Zeyh. l. c. = ericæfolia.
intermedia, *Eckl. & Zeyh. l. c.*—Afr. austr.
Krauseana, Meisn. in Hook. Lond. journ. Bot. ii. (1843) 65 = virgata.
micrantha, *Walp. in Linnæa*, xiii. (1839) 470.—Afr. austr.
multiflora, Eckl. & Zeyh. Enum. 166 = ericæfolia.
perplexa, Eckl. & Zeyh. l. c. 170 = Coelidium roseum.
phyllicoides, *Eckl. & Zeyh. l. c.*—Afr. austr.
spinosa, Steud. Nom. ed. II. i. 80 = Coelidium spinosum.
torbitis, Stend. l. c. = Coelidium Vogelii.
verticillata, Stend. l. c. = Priestleya elliptica.
violacea, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 451.—Afr. austr.
virgata, *Eckl. & Zeyh. Enum.* 169.—Afr. austr.
Vogelii, Walp. in Linnæa, xiii. (1839) 473 = ericæfolia.
Williamsonii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 28.—Afr. austr.

AMPHIZOMA, Miers, in Trans. Linn. Soc. xxviii. (1872) 422 = Balsocia, Linn. (Hippocrat.).
laxiflora, Miers, l. c. = S. laxiflora.
Martiana, Miers, l. c. 428 = S. Martiana.
tenuicula, Miers, l. c.—Bras.

AMPHOCHAETA, Anderss. in Vet. Akad. Hand. Stockh. (1853) 136 = Pennisetum, Pers. (Gramin.).
exaltata, Anderss. l. c. = P. exaltatum.

AMPHODUS, Lindl. Bot. Reg. t. 1101 (1827) = Kennedya, Vent. (Legum.).
ovatus, Lindl. l. c. = K. rubicunda.

AMPHORCHIS, Thou. Orch. Iles Afr. Tabl. Synopt. (1822) = Cynorchis, Thou. (Orchid.).
calcarata, Thou. l. c. t. 4 = C. squamosa.
colorata, Steud. Nom. ed. II. i. 80, sphalm. = præc.
inermis, Thou. Orch. Iles Afr. t. 5 = Arnottia inermis.
laxiflora, Blume, Mus. Bot. Lugd. Bat. ii. 190.—Malaya.
lilacina, Ridl. in Journ. Linn. Soc. xxi. (1885) 518.—Madag.
milarmis, Steud. Nom. ed. II. i. 80, sphalm. = inermis.
occidentalis, Lindl. in journ. Linn. Soc. vi. (1862) 189.—Afr. trop.

AMPHORICARPUS, Vieiani, Fl. Dalm. ii. 27. t. 10. f. 2 (1844). *COMPOSITAE*, Benth. & Hook. f. ii. 465.
Neumayeri, *Visiani, Fl. Dalm.* ii. 28. t. 10. f. 2.—Dalmat.

AMPHORICARPUS, Spruce, ex Miers, in Trans. Linn. Soc. xxx. (1874) 388 = Courataari, Aubl. (Myrtac.).
nanpensis, Spruce, ex Miers, l. c. = C. nanpensis.

AMPHYMENIUM, H. B. & K. Nov. Gen. et Sp. vi. 380 (1823) = Pterocarpus, Linn. (Legum.).
cordatum, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1905, nomen.—Guiana.
floribum, Benth. in Hook. Kew journ. ii. (1850) 284 = P. Rohrii.
latifolium, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1905, nomen.—Guiana.
lanciflorum, Benth. in Hook. Lond. journ. Bot. ii.(1843) 66.—Bras.
laxum, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1906, nomen.—Guiana.
longifolium, Benth. in Mart. Fl. Bras. xv. i. 209 = P. sacyclocalyx.

AMPHYMENIUM :—
mediterraneum, Benth. in Ann. Wien. Mus. ii. (1838) 96 = Martia parvifolia.
nitidum, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1906, nomen.—Guiana.
orbiculatum, Benth. in Ann. Wien. Mus. ii. (1838) 95 = P. orbiculatus.
pubescens, H. B. & K. Nov. Gen. et Sp. vi. 380 = P. Amphymenium.
reticulatum, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1905, nomen.—Guiana.
Rohrii, H. B. & K. Nov. Gen. et Sp. vi. 380 = P. Rohrii.
tenuifolium, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1905, nomen.—Guiana.
villosum, Benth. in Ann. Wien. Mus. ii. (1838) 95 = P. villosum.
violaceum, Walp. Rep. i. 798 = P. violaceus.

AMPOMELE, Rafin. Sylva Tellur. 160 (1838) = Rubus, Tourn. (Rosac.).
triphylla, Rafin. l. c. = R. trifidus.

AMSINCKIA, Lehm. Del. Sem. Hort. Hamb. (1831) 7. *BORAGINEAE*, Benth. & Hook. f. ii. 851.
BENTHAMIA, Lindl. Nat. Syst. ed. I. 241 (1830).
angustifolia, *Lehm. l. c.*—Chili.
basistaminea, *Cesati, in Atti Acc. Sc. Nap.* v. (1873) vii. 14.—Reg. Andin.
Douglasiana, A. DC. in DC. Prod. x. 118 = spectabilis.
echinata, *A. Gray, in Proc. Am. Acad.* x. (1874) 54.—Calif.
hamata, *Walp. in Nov. Act. Nat. Cur. xix. Suppl.* i. (1843) 371.—Reg. Argent.
intermedia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1836) 26.—Calif.
lycopsoides, *Lehm. Del. Sem. Hort. Hamb* (1831) 7.—Calif.
mexicana, *Mart. & Galeot. in Bull. Acad. Brux.* xi. (1844) reimpr. 28.—Mexic.
parviflora, Bernh. Del. Sem. Hort. Erf. (1838) = angustifolia.
spectabilis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1836) 26.—Calif.
tessellata, *A. Gray, in Proc. Am. Acad.* x. (1874) 54.—Calif.
Venicosa, *Hook. & Arn. Bot. Beech. Voy.* 370.—Calif.

AMSONIA, Walt. Fl. Carol. 98 (1788). *APOCYNACEAE*, Benth. & Hook. f. ii. 702.
angustifolia, *Michx. Fl. Bor. Am.* i. 121.—Am. bor.
brevifolia, *A. Gray, in Proc. Am. Acad.* xii. (1877) 64.—Utah; Arizona.
ciliata, Walt. Fl. Carol. 98 = angustifolia.
elliptica, *Roem. & Schult. Syst.* iv. 492.—Japon.
latifolia, Michx. Fl. Bor. Am. i. 121 = A. Tabernæmontana.
longiflora, *Torr. Bot. Mex. Bound.* 159.—N. Mexic.
orientalis, Decne. in Jacquem. Voy. Bot. 105 = Rhazya orientalis.
Palmeri, *A. Gray, in Proc. Am. Acad.* xii. (1877) 64.—Arizona.
salicifolia, Pursh, Fl. Am. Sept. i. 184 = A. Tabernæmontana.
Tabernæmontana, *Walt. Fl. Carol.* 98.—Am. bor.
tomentosa, *Torr. & Frém. in Frém. Rep.* 1848–44 (1845) 316.—Am. bor.
triuris, Sm. in Rees, Cycl. xxxv. ad calc. art. Tabernæmontanæ, n. 2 = A. Tabernæmontana.

AMSORA, Bartl. Ord. 536 (1830), sphalm. = Amoora, Roxb. (Meliac.).

AMURA, Schult. f. Syst. vii. p. 11, 75 (1829) = Amoora, Roxb. (Meliac.).

AMYDRIUM, Schott, in Mlq. Ann. Mus. Lugd. Bat. i. 127 (1863). *AROIDEAE*, Benth. & Hook. f. iii. 993.
humile, *Schott, l. c.*—Borneo.

AMYGDALOPHORA, Nock. Elem. ii. 70 (1790) = Prunus, Tourn. (Rosac.).

AMYGDALOPSIS, Carr. in Rev. Hortic. (1868) 91 = Prunus, Tourn. (Rosac.).
aburnea, M. Roem. Syn. Rosif. 16 = P. aburnea.
furcata, M. Roem. l. c. = P. aburnea.
horrida, M. Roem. l. c. 15 (= *Amygdalus horrida*).—Persia.

AMYGDALOPSIS :—
Lindleyi, Carr. in Rev. Hortic. (1862) 91; (1868) 197 = P. triloba.
lycioides, M. Roem. Syn. Rosif. 15 (= *Amygdalus lycioides*).—As. Min. ; Syria.
Scorpius, M. Roem. l. c. = P. aburnea.
virgata, Carr. in Rev. Hortic. (1873) 200 = P. virgata.

AMYGDALUS, [Tourn.] Linn. Syst. ed. I. (1735) = Prunus, Tourn. (Rosac.).
agrestis, Boiss. Diagn. Ser. I. x. 1.—Syria.
amara, Hayne, Arzn. iv. in note ad t. 39. f. 1.—P. Amygdalus.
amygdalina, Oken, ex M. Roem. Syn. Rosif. 8 = P. Amygdalus.
arabica, Olivier, Voy. t. 47.—Arabia.
argentea, Lam. Encyc. i. 103 = orientalis.
Armeniaca, Dum. Fl. Belg. 91 = P. Armeniaca.
Armeniacaria, Oken, ex M. Roem. Syn. Rosif. 10 = P. Amygdalus.
Balansæ, Boiss. Diagn. Ser. II. vi. 71.—Phrygia.
Bessariana, Schott, ex Schlecht. in Abh. Naturf. Ges. Halle, ii. (1854) 1 = P. nana.
Beissneri, Carr. in Rev. Hortic. (1875) 362.—As. Min.
Bovei, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 124.—Syria.
brahuica, Boiss. Fl. Orient. ii. 645.—Beluchist.
camelliæflora, Morr. in Belg. Hortic. viii. (1858) 97. t. 96 = P. Persica.
campaniflora, Carr. in Rev. Hortic. (1872) 129.—Hab. ?
campestris, Bess. Enum. Pl. Volh. 46 = P. nana.
caryophyllacea, Hort. ex C. Koch, Dendrol. i. 85 = P. Persica.
carsiana, Oken, ex M. Roem. Syn. Rosif. 9 = P. Amygdalus.
chinensis, Hort. ex C. Koch, Dendrol. i. 85 = P. Persica.
chinensis, Lavoy, ex C. Koch, l. c. 79 = P. nana.
cochinchinensis, Lour. Fl. Cochinch. 316.—Cochinch.
collinus, Wall. Cat. n. 738 = P. Persica.
communis, Bunge, Enum. Pl. Chin. Bor. 21 = P. Persica.
communis, Linn. Sp. Pl. ed. I. 473 = P. Amygdalus.
cordifolia, Roxb. Fl. Ind. ii. 500.—China.
dasyleptis, Miq. in Journ. Bot. Néerl. i. (1861) 192 = P. Persica.
dianthiflora, Hort. ex C. Koch, Dendrol. i. 85 = P. Persica.
discolor, M. Roem. Syn. Rosif. 12 = orientalis.
dulcis, Mill. Gard. Dict. ed. VIII. n. 2 = P. Amygdalus.
eburnea, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 128.—Persia.
elæagrifolia, Spach, l. c. 130.—Persia.
elata, Salisb. Prod. 356 = P. Amygdalus.
Fischeriana, Spach, in Ann. Sc. Nat. Sér II. xix. (1843) 128.—Hab. ?
fragilis, Hell. Fl. Witreb. i. 500 = P. Amygdalus.
frutescens, Steud. Nom. ed. II. i. 81 = (*A. fruticosa*).
fruticosa, Wender. in Bot. Zeit. (1845) 588. t. 12.—Hab. ?
furcatus, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 124 = P. aburnea.
Garrisenana, Schlecht. in Abh. Ges. Halle, ii. (1854) 19 = P. nana.
georgica, Desf. Hist. Arbr. ii. 321 = P. nana.
glandulosa, Hort. ex C. Koch, Dendrol. i. 118 = P microphylla.
glandulosa, Hook. fc. Pl. t. 988 = P. glandulosa.
glandulosa, Lavoy, ex C. Koch, Dendrol. i. 118 = P. japonica.
grandiflora, Hort. ex C. Koch, l. c. 81 = P. Amygdalus.
grandiflora, Hort. Gall. ex C. Koch, l. c. i. 90 = P. Persica.
graca, Salisb. Prod. 356 = P. Persica.
heterophylla, Hort. ex C. Koch, Dendrol. i. 81 = P. Amygdalus.
Hoeckiana, Schlecht. in Abh. Ges. Halle, (1854) t. 22.—Sibir.
horrida, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 121.—Persia.
humilis, Edgew. in Trans. Linn. Soc. xx. (1846) 44 = P. Jaquemontii.
hybrida, Diarb. ex M. Roem. Syn. Rosif. 11 = P. Amygdalus.

AMYGDALUS —
incana, Pall. Fl. Ross. i. 18. t. 7 = Prunus prostrata.
tapahanensis, Thouin, in Ann. Mus. Par. Ser. I. viii. (1826) 425 = P. Persica.
Kotschyi, Hohen. ex Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 117 = *elaeagnifolia*.
lavris, D. Dietr. Syn. Pl. iii. 42 = P. Persica.
Ledebouriana, Schlecht. in Abh. Ges. Halle, (1854) 1 = P. nana.
leiocarpa, Boiss. Diagn. Ser. I. vi. 52.—Persia.
lycioides, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 120.—As. Min.; Syria.
microphylla, H. B. & K. Nov. Gen. et Sp. vi. 245. t. 564 = P. microphylla.
minor, Duh. Arbr. Fruit. i. t. 7.—Hab.?
nana, Linn. Sp. Pl. 473.—Europ.; Reg. Casc.; Sibir.
Nuci-persica, Reichb. Fl. Germ. Excurs. 647 = P. Persica.
orientalis, Mill. Gard. Dict. ed. VIII. n. 4.—As. Min.; Syria.
Pallasiana, Schlecht. in Abh. Ges. Halle, (1854) 1 = P. nana.
Pallasii, Turcz. ex Steud. Nom. ed. II. i. 81 = P. pedunculata.
pedunculata, Bunge, Enum. Pl. Chin. Bor. 22 = P. triloba.
pedunculata, Pall. in Nov. Act. Petrop. vii. (1789) 855. t. 8, 9.—Sibir.
Persica, Linn. Sp. Pl. 473 = P. Persica.
persico-amygdala, Reichb. Fl. Germ. Excurs. 647 = P. Amygdalus.
persicoides, C. Koch, Dendrol. i. 84 = P. Persica.
pilosa, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 189.—China.
prostrata, Sweet, Hort. Brit. ed. II. 178 = P. prostrata.
pumila, Linn. Mant. i. 74 = P. japonica.
Reuteri, Boiss. & Bahse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 79 = *horrida*.
rosaeflora, Hort. ex C. Koch, Dendrol. i. 85 = P. Persica.
salicifolia, Boiss. & Bal. Diagn. Ser. II. vi. 671 = Webbii.
sativa, Mill. Gard. Dict. ed. VIII. n. 3 = P. Amygdalus.
scoparia, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 109.—Persia.
scorpius, Spach, l. c. 122 = P. eburnea.
serratifolia, Hort. ex C. Koch, Dendrol. i. 79 = P. nana.
sibirica, Lodd. ex Lond. Hort. Brit. 199.—Sibir.
sinensis, Hort. ex Steud. Nom. ed. II. i. 81 = *cochinchinensis*.
spartioides, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 108.—Syria; Persia.
spathulata, Boiss. Diagn. Ser. I. vi. 53 = P. eburnea.
spinosissima, Bunge, Rel. Lehm. 106.—Turkest.
Stochnana, Boiss. Diagn. Ser. II. ii. 45.—Beluchist.
tomentosa, Hort. ex C. Koch, Dendrol. i. 81 = P. Amygdalus.
tomentosula, Steud. Nom. Bot. ed. II. i. 81 = P. Amygdalus.
Webbii, Spach, in Ann. Sc. Nat. Ser. II. xix. (1843) 117.—Phrygia.

AMYRIS, P. Br. Hist. Jam. 208 (1756). *BURSERACEAE*, Benth. & Hook. f. i. 337.
ELEMI, Adans. Fam. ii. 342 (1763).
ELEMIFERA, Linn. Hort. Cliff. 486 (1787).
RITINOPHORA, Neck. Elem. ii. 229 (1790).
acuminata, Poir. Encyc. Suppl. ii. 136 = Icica acuminata.
acuminata, Roxb. Hort. Beng. 28; Fl. Ind. ii. 246 = Commiphora caudata.
Agallocha, Roxb. Hort. Beng. 28 (1814) = Commiphora Agallocha.
altissima, Willd. Sp. Pl. ii. 336 = Protium altissimum.
ambrosiaca, Linn. f. Suppl. 216 = Protium Icicariba.
ambrosiaca, Moç. & Sesse, in DC. Prod. ii. 77, 78 = Icica serrata.
ambrosiaca, Spreng. Syst. ii. 218, pro parte = Protium guianense.
ambrosiaca, Willd. Sp. ii. 335 = Protium heptaphyllum.
ambrosiaca, Vell. Fl. Flum. iv. t. 3 = Protium heptaphyllum?
amicta, Roxb. ex Steud. Nom. ed. II. i. 81 = Clausena heptaphylla.
amicta, Willd. Sp. Pl. ii. 337 = Clausena anisatum.
arborea, Spreng. Syst. iv. Cur. Post. 149 = Elaphrium arboreum.
armata, Spreng. Syst. ii. 219 = Zanthoxylon Thunbergii.
axillaris, *Griseb. Cat. Pl. Cub.* 46.—Cuba.
balsamifera, *Linn. Syst.* ed. X. 1000.—Jamaica.
bicolor, Willd. ex Engl. in DC. Monog. Phan. iv. 54 = Bursera bicolor.
bipinnata, Moç. & Sesse, ex DC. Prod. ii. 82 = Bursera bipinnata.
brachyboutya, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 614.—Bras.
brasiliensis, Spreng. Syst. ii. 217 = Protium brasiliense.
brasiliensis, Willd. ex Engl. in DC. Monog. Phan. iv. 64 = Protium heptaphyllum.
capensis, Spreng. Syst. ii. 218 (= *Fagarastrum capense*).
Carana, Humb. Relat. ii. 421 = Protium Carana.
caranifera, Willd. ex Engl. in DC. Monog. Phan. iv. 49 = Bursera graveolens.
Commiphora, Roxb. Fl. Ind. ii. (1824) 244 = Commiphora Agallocha.
Cataf, 'Forsk.' = *A. Kataf.*
copallifera, *Oliva, in La Naturaleza*, (1870) 40.—Mexico.
copallifera, Spreng. Syst. ii. 219 = Elaphrium copalliferum.
cymosa, Reichb. ex Steud. Nom. ed. II. i. 81 = sylvatica.
decandra, *Brotero, ex Prits. in Bot. Zeit.* i. (1843) 561.—Bras.
decandra, Willd. Sp. Pl. ii. 336 = Icica decandra.
dentata, Blume, Bijdr. 1160 = Clausena javanica.
dentata, Willd. Sp. Pl. ii. 337 = Clausena Willdenowii.
diatrypa, *Spreng. Neue Entdeck.* iii. 48; *Syst.* ii. 217. —In Deming.
disica, Hort. ex Steud. Nom. ed. II. i. 81 = Schinus dependens.
Elaphrium, Spreng. Syst. ii. 219 = Bursera glabra.
elegans, Willd. ex Engl. in DC. Monog. Phan. iv. 57 = Bursera excelsa.
elemifera, Koen. ex Blume, Mus. Bot. Lugd. Bat. i. 218 = Canarium zeylanicum.
elemifera, Linn. Syst. ed. X. 1000 = Plumieri.
ennoandra, Willd. Sp. Pl. ii. 335 = Protium decandrum.
fagaroides, Spreng. Syst. iv. Cur. Post. 148 = Bursera fagaroides.
floridana, *Ware, ex Nutt. in Am. Journ. Sc.* v. (1822) 28.—Florida.
Funckiana, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 475.—Venezuela.
fruticosa, *Roxb. Hort. Beng.* 28.—Ind. or.
gileadensis, Roxb. Hort. Beng. 28; Fl. Ind. ii. 246 = C. Berryi.
gileadensis, Linn. Diss. Opob.; Amoen. Acad. vii. 68 = Commiphora Opobalsamum.
graveolens, Buch.-Ham. ex Steud. Nom. ed. II. i. 81 = Clausena excavata.
graveolens, Spreng. Syst. iv. Cur. Post. 148 = Bursera graveolens.
guianensis, *Aubl. Pl. Gui.* i. 336.—Guiana.
guianensis, Willd. Sp. Pl. ii. 335 = Protium guianense.
heptaphylla, Roxb. Hort. Beng. 28; Fl. Ind. ii. 248 = Clausena heptaphylla.
heterophylla, Willd. Sp. Pl. ii. 335 = Protium Aracouchini.
hexandra, Ham. Prod. Fl. Ind. Occ. 84 = Dacryodes hexandra.
Hypelate, Robinson, ex Lunan, Hort. Jamaic. i. 149 = Hypelate trifoliata.
inaequalis, Spreng. Syst. ii. 218 = Clausena inaequalis.
jorullensis, Spreng. Syst. Cur. Post. 148 = Bursera jorullensis.
Kafaia, Forsk. Fl. Aegypt. Arab. 80 = an sequ.?
Kataf, Forsk. l. c. = Commiphora Kataf.
lanuginosa, Spreng. Syst. iv. Cur. Post. 149 = Bursera lanuginosa.
Linalol, La Llave, in La Naturaleza, vii. (1885) Apénd. 78 = seq.
Linanoe, La Llave, ex Schlecht. in Linnaea, xvii. (1843) 253 = Bursera Alcexylon.
linosta, *C. Wright, ex Griseb. in Mem. Am. Acad.* N. S. vii. (1861) 177.—Ind. occ.
Lunani, Spreng. Syst. ii. 217 = maritima.
Lyptata, Steud. Nom. ed. II. i. 81, sphalm. = Hypelate trifoliata.
maritima, *Vog. Enum. Pl. Carib.* 19.—Am. trop.
maritima, Llanos, in Mem Acad. Madr. Ser. III. ii. 111. (1856) 503.—Ins. Philipp.?
maritima, Moç. & Sesse, ex DC. Prod. ii. 82 = A. Tecomaca.
multijuga, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 476.—Bras.
nana, Roxb. Hort. Beng. 28 = Clausena Willdenowii.
Nonoo-Nolandiae, Hort. ex Steud. Nom. ed. II. i. 81 = Botryceras laurinum.
oleosa, Lam. Encyc. i. 362 = Canarium microcarpum.
Opobalsamum, Linn. Diss. Opob.; Amoen. Acad. vii. 68; Forsk. Fl. Aegypt. 76 = Commiphora Opobalsamum.
papyrifera, Calli. ex Del. Voy. Meroé, 99 = Boswellia papyrifera.
penicillata, Spreng. Syst. ii. 219 = Bursera penicillata.
pentaphylla, Roxb. Hort. Beng. 28; Fl. Ind. ii. 247 = Clausena pentaphylla.
pernambucensis, *Arruda, in Koster, Trav. Bras.* 500.—Bras.
Philippaea, Robinson, ex Lunan, Hort. Jam. i. 150 = Hypelate trifoliata.
pinnata, *H. B. & K. Nov. Gen. et Sp.* vii. 87.—Am. trop.
Plumieri, *DC. Prod.* ii. 81.—Ind. occ.
polygama, Cav. Ic. iii. 20 = Schinus dependens.
Protium, Linn. Mant. i. 65 = Protium javanicum.
pubescens, Willd. ex Schlecht. in Linnaea, xvii. (1842) 527 = Elaphrium pubescens.
punctata, Roxb. Hort. Beng. 28; Fl. Ind. ii. 251 = Clausena excavata.
Robinsonii, *DC. Prod.* ii. 82.—Jamaica.
rugosa, Willd. ex Engl. in DC. Monog. Phan. iv. 58 = Bursera jorullensis.
serratifolia, Kottl. ex A. W. Benn. in Hook. f. Fl. Brit Ind. i. 539 = Balsamodendron Roxburghii.
simplicifolia, Roxb. Fl. Ind. ii. 244 = Atalantia Roxburghiana.
subfruticosa, Roxb l. c. 250 = Clausena suffruticosa.
sumatrana, Roxb. Hort. Beng. 28; Fl. Ind. ii. 250 = Clausena excavata.
sylvatica, *Jacq. Am. Select.* 107.—Ind. or.
Tecomaca, *Moç. & Sesse, Fl. Mex. ined.* t. 195, *ex DC. Prod.* ii. 82.—Mexico.
terebunthifolia, *Tenore, Ind. Sem. Hort. Neap.* (1837) 1.—Bras.
thyrsiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 475.—Mexico.
tomentosa, Spreng. Syst. ii. 219 = Bursera tomentosa.
toxifera, *Linn. Syst.* ed. X. 1000.—Ind. occ
ventricosa, Llave, ex Schlecht. in Linnaea, xvii. (1843) 246 = Bursera fagaroides.
zeylanica, Retz. Obs. fasc. iv. 25 = Canarium zeylanicum.

AMYRSIA, Rafin. Sylva Tellur. 106 (1838) = **Eugenia**, Linn. (Myrtac.).
compressa, Rafin. l. c. = E. compressa.
discolor, Rafin. l. c. = E. discolor.
foliosa, Rafin. l. c. = E. foliosa.
microphyla, Rafin. l. c. (= *Myrtus microphylla*).

ANABAENA, A. Juss. Euph. Tent. 46. t. 15 (1824) = **Plukenetia**, Linn. (Euphorb.).
tamnoides, A. Juss. l. c. t. 15 = P. tamnoides.

ANABASIS, Linn. Pl. Camts. (1750) (Amoen. Acad. ii. 247). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 72.
BORITA, Adans. Fam. ii. 262 (1763).
BRACHYLEPIS, C. A. Mey. Fl. Alt. i. 370 (1829).
FREDOLIA, Coss. & Dur. ex Moç. & Coss. in Bull. Soc. Bot. Fr. ix. (1862) 301.
MICROLEPIS, Eichw. Casp. Casc. 2 (1831).
affinis, Bunge, Rel. Lehm. 480 = Arthrophytum subulifolium.
affinis, C. A. Mey. in Schrenk, Enum. Pl. Nov. Song. (1841) 19 = brevifolia.
alopecuroides, Moç. in DC. Prod. xiii. ii. 219 = Halogeton alopecuroides.
Ammodendron, C. A. Mey. in Ledeb. Fl. Alt. i. 375 = Haloxylon Ammodendron.
annua, *Bunge, Anab. Rev.* 46.—Persia.
aphylla, *Linn. Sp. Pl.* 238.—Orient; Sibir.
arbuscula, *Thunb. ex Steud. Nom.* ed. II. i. 81.—Hab.? aretioides, *Moç. & Coss. in Bull. Soc. Bot. Fr.* ix (1862) 301. t. 3.—Afr. bor.
articulata, Chomlette, ex Pomel, Nouv. Mat. Fl. Atl. 334 = prostrata.
articulata, *Moq. in DC. Prod.* xiii. ii. 212.—Afr. bor.; Syria.
brachiata, *Fisch. & Mey. ex Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 439.—Orient; Turkest.
brachiata, Kar. & Kir. l. c. = truncata.

ANABASIS:—
brachylepis, *D. Distr. Syn. Pl.* ii. 995.—Reg. Casp.
brevifolia, *C. A. Mey. in Ledeb. Fl. Alt.* i. 377.—Sibir. alt.
caespitosa, Steph. ex Moq. in DC. Prod. xiii. ii. 216 = Nanophyton caspicum.
cinerea, Moq. l. c. 214 = Seidlitzia florida.
clavata, S. G. Gmel. It. ii. 41 = Salsola clavifolia.
conjugata, Hoffm. Hort. Moscq. (1808) No. 184 = Petrosimonia brachiata.
crassa, Moq. Chenop. Enum. 166 = articulata.
cretacea, C. A. Mey. Enum. ex Bunge, Anab. Rev. 89 = brachiata.
cretacea, *Pall. Reise*, i. 498.—Sibir.
cretica, Vitm. Summ. Pl. ii. 104 = cretacea.
Echinus, Bieb. Beschr. Laend. Casp. 152 = Noaea spinosissima.
Ehrenbergii, *Schweinf. ex Boiss. Fl. Orient.* iv. 970, in nota.—Nubia; Arab.
florida, Bieb. Fl. Taur. Cauc. i. 190 = Seidlitzia florida.
foliata, Pall. ex Bunge, Anab. Rev. 47 = Salsola foliosa.
foliosa, Pall. Reise, i. 260 = Salsola clavifolia.
foliosa, Linn. Sp. Pl. 223 = Salsola foliosa.
glomerata, Bieb. in Mém. Soc. Nat. Mosc. i. (1808) 110 = Halogeton glomeratus.
Haussknechtii, *Bunge, ex Boiss. Fl. Orient.* iv. 969.—Persia.
heterophylla, Walp. Ann. i. 571 (aphalm. = seq.).
heteroptera, Jaub. & Spach, Illustr. ii. 45. t. 134 = Girgensohnia oppositiflora.
intermedia, Moq. in DC. Prod. xiii. ii. 214 = phyllophora ?
juncea, Bieb. ex Moq. l. c. 208 = Ofaiston monandrum.
Kareliniana, Fenzl, in Ledeb. Fl. Ross. iii. 825 = truncata.
lutea, Moq. in DC. Prod. xiii. ii. 215 = setifera.
macroptera, *Moq. l. c.* 218.—Affghanist.
monandra, Schrad. Neues Journ. iii. (1809) 89 = Ofaiston pauciflorum.
multiflora, Moq. in DC. Prod. xiii. ii. 212 = Haloxylon multiflorum.
oppositiflora, Bieb. in Mém. Soc. Nat. Mosc. i. (1808) 148 = Girgensohnia oppositiflora.
oppositifolia, Bieb. Beschr. Laend. Casp. App. 154 = Petrosimonia crassifolia.
oppositifolia, Schrad. Neues Journ. iii. (1809) 89 = Halogeton oppositifolius.
phyllophora, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 781.—Syria; Soongaria.
prostrata, *Pomel, Nouv. Mat. Fl. Alt.* 334.—Afr. bor.
ramosa, *Sievers, in Pall. N. Nord. Beitr.* vii. 304.—Sibiria.
setifera, *Moq. Chenop. Enum.* 164.—Aegypt.; Oriens.
Sieversii, Willd. in Roem. & Schult. Syst. vi. 293 = Nanophyton caspicum.
spheniolaxima, Linn. f. Suppl. 178 = Noaea spinosissima.
subulifolia, C. A. Mey. in Bull. Acad. Pétersb. Sér. IV. i. (1843) 260 = phyllophora.
tamariscifolia, Linn. Syst. ed. X. 949 = Salsola tamariscifolia.
tamariscifolia, Webb, Iter Hisp. 17 = Salsola Webbii.
tatarica, Pall. Illustr. 18. t. 8 = aphylla.
Tournefortii, Jaub. & Spach, Illustr. Pl. Or. ii. 48. t. 129 = Noaea Tournefortii.
triandra, Bieb. Beschr. Laend. Casp. App. 154 = Petrosimonia Volvox.
truncata, *Bunge, Anab. Rev.* 38.—Reg. Casp.

ANARATA, Willd. ex Roem. & Schult. Syst. iv. 60. 707 (1819). Gen. dub. *LOGAN.* Benth. & Hook. f. ii. 788.
SULZERIA, Roem. & Schult. l. c.
odonata, *Humb. ex Roem. & Schult. l. c.*—Am. austr.

ANACAMPSEROS, Linn. Syst. ed. I. (1735); Gen. ed. I. 136 (1737). *PORTULACAE*, Benth. & Hook. f. i. 127.
RUELINGIA, Ehrh. Beitr. iii. 182 (1788).
TALINUM, Adans. Fam. ii. 245 (1763).
TELEPHIASTRUM, Medic. Phil. Bot. i. 100 (1789).
angustifolia, *Sweet, Hort. Brit.* ed. I. 170.—Afr. austr.
arachnoides, *Sims, Bot. Mag.* t. 1368.—Afr. austr.
filamentosa, *Sims, l. c.* 1367.—Afr. austr.
fastermodia, Haw. ex G. Don, Gen. Syst. iii. 75 = A. Telephiastrum.

ANACAMPSEROS:—
lanceolata, *Sweet, Hort. Brit.* ed. I. 170.—Afr. austr.
lanigera, *Burch. Trav.* ii. 338.—Afr. austr.
papyracea, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 383.—Afr. austr.
polyphylla, *Sweet, Hort. Brit.* ed. I. 170.—Afr. austr.
quinaria, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 383.—Afr. austr.
rotundifolia, *Sweet, Hort. Brit.* ed. I. 170.—Afr. austr.
rubens, Sweet, l. c. 170 = arachnoides.
rufescens, Sweet, l. c. 170 = arachnoides.
Telephiastrum, *DC. Cat. Hort. Monsp.* 77.—Afr. austr.
trigona, *DC. Prod.* iii. 356.—Afr. austr.
tuberosa, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 383.—Afr. austr.
varians, Sweet, Hort. Brit. ed. I. 170 = A. Telephiastrum.

ANACAMPSEROS, [Tourn.] Adans. Fam. ii. 248 (1763); Haw. Syn. Pl. Succ. 111 (1812) = Sedum, Linn. (Crassul.).
Aizoon, Haw. Syn. Pl. Succ. 113 = S. Aizoon.
albicans, Haw. l. c. 111 = S. albicans.
alsida, Haw. ex DC. Prod. iii. 402 = S. Telephium ?
alpestris, Jord. & Fourr. Brev. Nov. Pl. fasc. i. 32 = S. maximum.
angustifolia, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 168.—Gallia.
aprica, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 19 = S. maximum.
argula, Haw. Rev. Pl. Succ. 25 = S. argutum.
asurgens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 24 = S. maximum.
aurigerana, Jord. & Fourr. l. c. 30 = S. Telephium.
brugnatia, Jord. & Fourr. l. c. 24 = S. Telephium.
Borderi, Jord. & Fourr. l. c. 30 = S. Telephium.
buxicola, Jord. & Fourr. l. c. 27 = S. Telephium.
caerulescens, Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, ii. (1874) 70 = S. caeruleum ?
Cepaea, Willd. Sp. Pl. ii. 762 = S. Cepaea.
cebennensis, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 15 = S. maximum.
chlorotica, Jord. & Fourr. l. c. 20 = S. maximum.
ciliaris, Haw. Syn. Pl. Succ. 113 = S. involucratum.
collina, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 23 = S. maximum.
conferta, Jord. & Fourr. Ic. Pl. Eur. i. 36 = S. Telephium.
convexa, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 29 = S. Telephium.
cornica, Jord. & Fourr. l. c. 17 = S. maximum.
delphinensis, Jord. & Fourr. l. c. 21 = S. maximum.
dentata, Haw. Rev. Pl. Succ. 26 = S. dentatum.
divaricata, Haw. Syn. Pl. Succ. 113 = S. divaricatum.
dumeticola, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 28 = S. Telephium.
erubescens, Jord. & Fourr. l. c. 18 = S. maximum.
formosa, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 168.—Gallia.
hybrida, Haw. Syn. Pl. Succ. 113 = S. hybridum.
fullitana, Jord. & Fourr. Ic. Pl. Eur. i. 36 = S. Telephium.
lagidiola, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 29 = S. Telephium.
linguaefolia, Hort. ex Lem. Illustr. Hortic. x. (1863) Misc. 77. Cotyledon linguaefolia.
livida, Haw. Rev. Pl. Succ. 25 = S. lividum.
lugdunensis, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 26 = S. Telephium.
maxima, Haw. Syn. Pl. Succ. ed. Germ. 121 = S. maximum.
Milleri, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 22 = S. maximum.
minor, Jord. & Fourr. l. c. 19 = S. maximum.
monticolarum, Jord. & Fourr. l. c. 26 = S. Telephium.
Neuteri, Jord. & Fourr. l. c. 27 = S. Telephium.
pachyphylla, Jord. & Fourr. l. c. 17 = S. maximum.
pauciones, Haw. Rev. Pl. Succ. 24.—Hab. ?
pectinata, Haw. l. c. 24 = S. pectinatum.
populifolia, Haw. Syn. Pl. Succ. 113 = S. populifolium.
Pourretii, Jeanb. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, iii. (1875-76) 256.—Gallia.
praecolax, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 25 = S. Telephium.
praeruptorum, Jord. & Fourr. l. c. 20 = S. maximum.
purpurea, Haw. ex DC. Prod. iii. 402 = S. Telephium.
pycnantha, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 26 = S. Telephium.

ANACAMPSEROS:—
pyramidalis, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 169.—Gallia.
racemosa, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 20 = S. maximum.
repens, Jord. & Fourr. l. c. 31 = S. Telephium.
rhodanensis, Jord. & Fourr. l. c. 24 = S. Telephium.
rubella, Jord. & Fourr. l. c. 30 = S. Telephium.
rupifraga, Jord. & Fourr. l. c. 29 = S. Telephium.
Saxifraga, Jord. & Fourr. l. c. 26 = S. Telephium.
semperviroa, Haw. Syn. Pl. Succ. 113 = S. Anacampseros.
serotina, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 22 = S. maximum.
silvatica, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 167.—Gallia.
spathulata, Hort. ex Steud. Nom. ed. II. i. 83 = S. spathulatum.
spectabilis, Jord. & Fourr. Ic. Pl. Eur. i. 37 = S. spectabile.
sparia, Haw. Rev. Pl. Succ. 25 = S. stoloniferum.
stellata, Haw. Syn. Pl. Succ. 114 = S. stellatum.
subalbida, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 27 = S. Telephium.
subrotunda, Jord. & Fourr. l. c. 19 = S. maximum.
subpyrenaica, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 169.—Gallia.
telephioides, Haw. Syn. Pl. Succ. 114 = S. telephioides.
ternata, Haw. l. c. = S. ternatum.
ternata, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 21 = S. maximum.
Thevenesi, Timb. in Mém. Acad. Toulouse, Sér. VII. vi. (1874) 646.—Gallia.
triphylla, Haw. Syn. Pl. Succ. 111 = S. Telephium.
Verloti, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 22 = S. maximum.
viridula, Jord. & Fourr. l. c. 26 = S. Telephium.
vogesiaca, Jord. & Fourr. l. c. 31 = S. Telephium.
vulgaris, Haw. Syn. Pl. Succ. 112 = S. Telephium.

ANACAMPTA, Miers. Apocyn. S. Amer. 64 (1878) = Tabernaemontana, Linn. (Apocyn.).
acutiflora, Miers, l. c. 66 = T. acutiflora.
angulata, Miers, l. c. 65 = T. angulata.
congesta, Miers, l. c. = T. rubrostriolata.
hirsuta, Miers, l. c. 67 = T. hirsuta.
longifolia, Miers, l. c. 66 = T. longifolia.
submollis, Miers, l. c. 67 = T. submollis.

ANACAMPTIS, Rich. in Mém. Mus. Par. iv. (1818) 47, 55 = Orchis, Linn.
Brancifortii, Lindl. Gen. et Sp. Orch. 275 = O. Brancifortii.
condensata, C. Koch, in Linnaea, xxii. (1849) 286, in syn. = O. pyramidalis.
Durandi, Bréb. Fl. Norm. ed. II. 258 = O. pyramidalis.
pyramidalis, Rich. in Mém. Mus. Par. iv. (1818) 55 = O. pyramidalis.
quadripunctata, Lindl. Gen. et Sp. Orch. 275 = O. Brancifortii.
trichocera, C. Koch, in Linnaea, xxii. (1849) 286 = O. quadripunctata.

ANACARDIA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = Anacardium, Linn.
rhinocarpa, St. Lag. l. c. = Anacardium Rhinocarpus.

ANACARDIUM, Linn. Syst. ed. I. (1735); Gen. ed. I. 139 (1737); Rottb. in Act. Hafn. ii. (1775) 286. *ANACARDIACEAE*, Benth. & Hook. f. i. 490.
ACAJOU, Tourn. ex Adans. Fam. ii. 344 (1763).
ACAJUBA, Gaertn. Fruct. i. 193. t. 40 (1788).
CASUVIUM, Lam. Encyc. i. 23 (1786).
MONODYNAMUS, Pohl, Pl. Bras. ii. 87. t. 144 (1831).
RHINOCARPUS, Bert. & Balb. ex Fl. B. & K. Nov. Gen. et Sp. vii. 5. t. 601 (1825).
bracteosum, *Burch. Rodr. in Rev. Eugenh.* (38 Jul. 1863) sepp. p. 5.—Bras.
curassicicolium, *St. Hil. in Ann. Sc. Nat. Sér. I.* xxiii. (1831) 273.—Bras.
dubium, Roxb. Hort. Beng. [66]; Fl. Ind. ii. 813 = Gluta Benghas.
excelsum, *Noronha, ex Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
giganteum, *Hancock, ex Engl. in Mart. Fl. Bras.* xii. ii. 410 (*errore Hancor.*).—Bras.

ANACARDIUM:—
giganteum, *Loud. ex Steud. Nom.* ed. II. i. 89, nomen. —Guinea.
humile, Engl. in Mart. Fl. Bras. xii. II. 411 = nanum.
humile, Mart. Reise, i. 814 = pumilum.
humile, *St. Hil. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 373.—Bras.
latifolium, Lam. Encyc. i. 189 = Semecarpus Anacardium.
longifolium, Lam. l. c. = Semecarpus Anacardium.
mediterraneum, *Vell. Fl. Flum.* 163; iv. t. 46.—Bras.
nanum, *St. Hil. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 372.—Bras.
occidentale, *Linn. Sp. Pl.* ed. I. 383.—Ind. occ.
officinale, Prits. Ind. Ic. 59 = Semecarpus Anacardium.
officinarum, Gaertn. Fruct. i. 192 = Semecarpus Anacardium.
orientale, Auct. ex Steud. Nom. ed. II. i. 82 = Semecarpus Anacardium.
pumilum, Walp. Rep. i. 555 = nanum.
Rhinocarpus, *DC. Prod.* ii. 62.—Am. austr.
solitarium, Stokes, Bot. Mat. Med. ii. 159 = Semecarpus Anacardium.
Spruceanum, *Benth. ex Engl. in Mart. Fl. Bras.* xii. II. 410.—Bras.
subcordatum, Presl, Bot. Bemerk. 40 = curatellæfolium.
subterraneum, Lisi, Climats, etc., Brésil, 604 = humile.

ANACHARIS, Rich. in Mém. Inst. ii. (1811) 62 = **Blodea**, Michx. (Hydroch.).
Alsinastrum, Bab. ex Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 74 = E. canadensis.
callitrichoides, Rich. in Mém. Inst. ii. (1811) 7. t. 2.—Bras.
canadensis, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 75 = E. canadensis.
chilensis, Planch. l. c.—Chili.
Hilariana, Casp. olim ex Casp. in Monatsb. Berl. Akad. (1857) 18 = E. Naias.
Matthewsii, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 78.—Peruv.
Nuttallii, Planch. l. c. 74 = E. canadensis.
pomeranica, Peterm. Deutschl. Fl. (1849) 529. t. 82. fig. 650 = E. canadensis.
Tripteris, Steud. Nom. ed. II. i. 82 (*sphalm.* = Anacis, Schrank) = Coreopsis Tripteris.

ANACHEILIUM, Hoffmgg. ex Linnæa, xvi. (1842) Lit. 229 = **Epidendrum**, Linn. (Orchid.).
cochleatum, Hoffmgg. l. c. = E. cochleatum.

ANACHYRIS, Nees, in Hook. Kew Journ. ii. (1850) 103 = **Paspalum**, Linn. (Gramin.).
paspaloides, Nees, l. c. = P. malacophyllum.

ANACHYRIUM, Steud. Syn. Pl. Gram. 33 (1855) = præc.

ANACIS, Schrank, in Denkschr. Acad. Muench. v. (1817) 5 = **Coreopsis**, Linn. (Compos.).
Tripteris, Schrank, l. c. = C. Tripteris.

ANACOLOSA, Blume, Mus. Bot. Lugd. Bat. i. 250. t. 46 (1850). *OLACINEÆ*, Benth. & Hook. f. i. 848.
crassipes, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 158.—Pegu.
dessiflora, *Bedd. in Madr. Journ.* Ser. III. i. (1864) 36.—Travancor.
frutescens, *Blume, Mus. Bot. Lugd. Bat.* i. 251.—Java.
Griffithii, *Mast. in Hook. f. Fl. Brit. Ind.* i. 580.—Burma.
heptandra, *Maingay, ex Mast. l. c.* 581.—Malacca.
lucoides, *Mast. l. c.* 580.—Reg. Himal.
Maingayi, *Mast. l. c.*—Singapore.
Parviflora, *Baill. Adansonia*, iii. (1862–63) 119.—Ins. Mascar.
pubsrula, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 207.—Ins. Andaman.

ANACOLUS, Griseb. in DC. Prod. ix. 70 (1845) = **Hookinia**, Gærtn. (Gentian.).
Gaudichaudii, Griseb. l. c. = H. montana.

ANACTORION, Raf. Fl. Tellur. iv. 34 (1836) = **Synaccia**, Sweet (Irid.).
bicolor, Raf. l. c. = S. bicolor.
galeatum, Raf. l. c. = S. galeata.

ANACTINIA, Remy, in C. Gay, Fl. Chil. iv. 8 (1849) = **Nardophyllum**, Hook. (Compos.).
Hookeri, Remy, l. c. = N. humile.

ANACTIS, Cass. in Dict. Sc. Nat. xlvii. 510 (1827) = **Atractylis**, Linn. (Compos.).
cæspitosa, Cass. l. c. 1. 56 = At. cæspitosa.
glabra, Rafin. Fl. Tellur. ii. 44 = Aster vimineus?
pilosa, Rafin. l. c. = Carthamus lanatus.
serratuloides, Cass. in Dict. Sc. Nat. l. 56 = At. serratuloides.

ANACYCLODON, Jungh. in Nat. en Geneesk. Arch. Néerl. Indie, ii. (1845) 49 = **Leucopogon**, R. Br. (Epacrid.).
pungens, Jungh. l. c. = L. javanicus.

ANACYCLUS, Linn. Gen. ed. I. 256 (1737). *COMPOSITÆ*, Benth. & Hook. f. ii. 419.
ARTHROLEPIS, Bolss. Diagn. Ser. I. xi. 14 (1849).
CYRTOLEPIS, Less. in Linnæa, vi. (1831) 166.
HIORTHIA, Neck. Elem. i. 97 (1790).
LEUCOCYCLUS, Bolss. Diagn. Ser. I. xi. 13 (1849).
alexandrinus, *Willd. Sp. Pl.* iii. 2173.—Afr. bor.
anomalus, J. Gay, ex Boiss. Fl. Orient. iii. 283 = Anthemis palæstina.
anthemoides, Lag. ex Spreng. Syst. iii. 497 = Anthemis abrotanifolia.
aristulatus, Link, ex Nym. Consp. 363 = clavatus.
aureus, Brot. Fl Lusit. i. 368 = radiatus.
aureus, Lam. ex DC. Fl. Fr. iv. 202 = Matricaria aurea.
aureus, Linn. Mant. ii. 287 = Anthemis aurea.
australis, Sieber, ex Spreng. Syst. iii. 497 = Cotula australis.
Barrelieri, Guss. Pl. Rar. 857 = Achillea Barrelieri.
Bivonæ, Pers. Syn. ii. 462 = radiatus.
Candolii, Nym. Syll. 8 = clavatus.
ciliatus, Trautv. in Bull. Soc. Nat. Mosc. xli. (1868) I. 461 = Anthemis ciliata.
clavatus, *Pers. Syn.* ii. 465.—Reg. Mediterr.
creticus, Linn. Sp. Pl. 892 = Anthemis cretica.
cystolepidioides, *Pomel, Nouv. Mat. Fl. Atl.* 54.—Afr. bor.
depressus, *Bail, in Journ. Bot.* xi. (1873) 365.—Marocco.
dissimilis, *Pomel, Nouv. Mat. Fl. Atl.* 53.—Afr. bor.
divaricatus, Cav. ex Steud. Nom. ed. I. 41 = clavatus.
formosus, *Fenzl, ex Boiss. Diag.* Ser. I. xi. 14.—Cilicia.
hirsutus, Lam. Fl. Fr. iii. 47 = valentinus.
inconstans, *Pomel, Nouv. Mat. Fl. Atl.* 52.—Afr. bor.
inflatus, *Lehm. ex Steud. Nom.* ed. II. i. 82.—Sibir.
lanuginosus, Moench, Meth. 581 = valentinus.
linearifolius, *Boiss. & Reut. Pugill. Pi. Hispan.* 57.—Hispan.
maroccanus, *Bail, in Journ. Bot.* xi. (1873) 365.—Marocco.
maurianicus, *Pomel, Nouv. Mat. Fl. Atl.* 55.—Afr. bor.
membranaceus, *Benth. & Hook. f. Gen.* iii. 419.—Syria.
mucronulatus, Guss. Pl. Rar. 856 = Achillea Barrelieri.
mucronulatus, Hort. ex Steud. Nom. ed. II. i. 82 = clavatus.
nigellæfolius, *Boiss. Diagn.* Ser. I. xi. 13.—Syria; Assyr.
officinarum, *Hayne, Arzneigew.* ix. t. 46.—Europ. austr.
orientalis, Linn. Sp. Pl. 892 = Anthemis montana.
pallidus, Guss. Fl. Sic. Syn. ii. 494 = radiatus.
pectinatus, Bory & Chaub. Fl. Pélop. 89. t. 30 = Anthemis tagetes.
pedunculatus, *Pers. Syn.* ii. 465.—Barbar.
prostratus, *Pomel, Nouv. Mat. Fl. Atl.* 53.—Afr. bor.
Pseudopyrethrum, *Achers. in Bonplandia*, vi. (1858) 198.—Hab.?
pubescens, Brot. ex Link, Handb. i. 768; Reichb. Fl. Exc. ii. 295 = clavatus.
pulcher, Bess. ex DC. Prod. vi. 16 = officinarum.
purpurascens, DC. Fl. Fr. v. 581 = radiatus.
Pyrethrum, Spreng. Syst. iii. 497 = Cotula Pyrethraria. Pyrethrum, *DC. Fl. Fr. Suppl.* 480.—Afr. bor.; Orient.
radiatus, *Loisel. Fl. Gall.* ed. I. 533; *Pers. Syn.* ii. 465.—Reg. Mediterr.
tomentosus, DC. Fl. Fr. Suppl. 481 = clavatus.
truncatus, J. Gay, in Boiss. Fl. Orient. iii. 283 = Anthemis palæstina.
valentinus, *Linn. Sp. Pl.* 892.—Reg. Mediterr.

ANACYLANTHUS, Steud. Nom. ed. II. i. 82 (1840), sphalm. = **Ancylanthos**, Desf. (Rubiac.).

ANADELPHIA, Hack. in Engl. Bot. Jahrb. vi. (1885) 240 = **Andropogon**, Linn. (Gramin.).
virgata, Hack. l. c. 241 = Andr. leptocomus.

ANADENDRUM, Schott, in Bonplandia, v. (1857) 45. *AROIDEÆ*, Benth. & Hook. f. iii. 991.
affine, *Schott, in Ann. Mus. Lugd. Bat.* i. 282.—Sumatra.
cordatum, *Schott, l. c.* 283.—Sumatra.
Lobbi, Schott, in Bonplandia, v. (1857) 45 = montanum.
marginatum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 116.—Ind. or.
medium, Schott, in Bonplandia, v. (1857) 45 = Epipremnum medium.
montanum, *Schott, l. c.*—Java.
punctulatum, Schott, Prod. Aroid. 393 = Monstera punctulata.
semiovestitum, Schott, in Ann. Mus. Lugd. Bat. i. 283 = affine.

ANADENDRON, Wight, Illustr. Ind. Bot. ii. 164 (1850) err. typ. = **Anodendron**, A. DC. (Apocyn.).

ANADEMIA, Agardh, Aphor. 236 (1826), sphalm. = seq.

ANADENIA, R. Br. in Trans. Linn. Soc. x. (1809) 166 = **Grevillea**, R. Br. (Proteac.).
anethifolia, A. Cunn. ex Benth. Fl. Austral. v. 483 = G. anethifolia.
Aquifolium, Lindl. Swan River App. 31 = G. monticola.
Caleyi, R. Br. Prot. Nov. 16 = G. ramosissima.
filifolia, Endl. in Ann. Wien. Mus. ii. (1838) 209 = G. didymobotrya.
flexuosa, Lindl. Swan River App. 31 = G. flexuosa.
gracilis, Lindl. l. c. = G. Synaphese.
haboides, Lindl. l. c. 80 = Hakea undulata.
sticifolia, R. Br. in Trans. Linn. Soc. x. (1809) 167 = G. ilicifolia.
integrifolia, Endl. in Ann. Wien. Mus. ii. (1838) 209 = G. integrifolia.
Manglesii, R. Grah. in Hook. Ic. Pl. t. 887 = Manglesia cuneata.
pulchella, R. Br. in Trans. Linn. Soc. x. (1809) 167 = G. pulchella.
Rœi, Endl. Gen. Suppl. iv. 11. 33 = G. didymobotrya.
tenuiflora, Lindl. Swan River App. 31 = G. tenuiflora.
trifida, R. Br. in Trans. Linn. Soc. x. (1809) 167 = G. trifida.
triternata, A. Cunn. ex Heynh. Nom. ii. 34 = G. triternata.

ANACTOCALYX, Triana, ex Benth. & Hook. f. Gen. i. 765 (1867). *MELASTOMACEÆ*, Benth. & Hook. f. i. 765.
bracteosa, *Triana, l. c. & in Trans. Linn. Soc.* xxviii. (1871) 97.—Venezuela.

ANAECTOCHILUS, Lindl. Orch. 498 (1840) = **Anoectochilus**, Blume (Orchid.).

ANAFRENIUM, Arn. in Hook. Lond. Journ. Bot. iii. (1841) 148 = ANAPHRENIUM, E. Mey. = **Rhus**, Linn. (Anacard.).

ANAGALLIDASTRUM, Mich. ex Adans. Fam. ii. 230 (1763) = **Centunculus**, Linn. (Primul.).

ANAGALLIDIUM, Griseb. Gen. et Sp. Gent. 211 (1856) = **Swertia**, Linn. (Gentian.).
dichotomum, Griseb. l. c. 212 = S. dichotoma.
tetrapetalum, Griseb. l. c. = S. tetrapetala.

ANAGALLIS, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 48 (1737). *PRIMULACEÆ*, Benth. & Hook. f. ii. 687.
EUPARKEA, Banks & Soland. in Gaertn. Fruct. i. 230. t. 50 (1788).
IRASEKIA, S. F. Gray, Nat. Arr. ii. 299 (1821).
JIRASEKIA, F. W. Schmidt, Neue u. Selt. Pfl. 49 (1793).
TIRASEKIA, G. Don, in Sweet, Hort. Brit. ed. III. 568 (1839).

ANAGALLIS :—
alba, Buch.-Ham. ex Steud. Nom. ed. II. i. 88.—(Qu?id ?) Ind. or.
alternifolia, *Cav. Ic.* vi. 8. t. 506.—Chili.
angustifolia, Sallab. Prod. 121 = linifolia.
aquatica, Erndl, ex Ledeb. Fl. Ross. iii. 31 = Samolus Valerandi.
arabica, *Duby, in DC. Prod.* viii. 60.—Arab.
arvensis, *Linn. Sp. Pl.* 148.—Europ.; As. temp.
caerulea, Lam. Fl. Fr. ed. 1. ii. 383 = arvensis.
capensis, *Linn. Sp. Pl.* 149.—Afr. austr.
carnea, Schrank, Baier. Fl. i. 461 = arvensis.
Centunculus, Afzel. Veget. Suecan. (1785) 1 = Centunculus minimus.
collina, Schousb. Marok. 64 = linifolia.
crassifolia, *Thore, Chlor. Land.* 69.—Hispan.; Marocco.
filiformis, *Cham. & Schlecht. in Linnaea*, i. (1826) 225.—Bras.
foemina, Mill. Gard. Dict. ed. VIII. n. 2 = arvensis.
fruticosa, Vent. Choix, 14 = linifolia.
Gmelini, *Steud. Nom.* ed. II. i. 88.—Camtsch.
grandiflora, Andr. Repos. t. 367 = linifolia.
Huttoni, *Harv. in Proc. Dub. Univ. Zool. & Bot. Assoc.* i. (1859) 141.—Afr. austr.
indica, Sweet, Fl. Gard. t. 132 = arvensis.
Jacquemontis, Duby, in DC. Prod. viii. 69 = arvensis.
latifolia, Linn. Sp. Pl. 148 = arvensis.
linifolia, *Linn. l. c.* ed. II. 212.—Europ.
Marryattae ×, Sweet, ex G. Don, in Sweet, Hort. Brit. ed. III. 668.
mas, Vill. Hist. Pl. Dauph. ii. 461 = arvensis.
Monelli, Bieb. Fl. Taur. Cauc. i. 143 = arvensis.
Monelli, Linn. Sp. Pl. 148 = linifolia.
multangularis, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 568 = Lysimachia pyrsmidalis.
nummularifolia, *Baker, in Journ. Linn. Soc.* xx. (1884) 196.—Madag.
orientalis, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. xi. 49 = arvensis.
ovalis, Reis & Pav. Fl. Per. ii. 8 = Centunculus pentandrus.
palustris, [Clairv.] Man. Herb. 59 = tenella.
parviflora, *Hoffmgg. & Link, Fl. Port.* i. 325. t. 64.—Lusitan.
parviflora, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 491 = arvensis.
pedunculata, Salzm. ex Duby, in DC. Prod. viii. 72 = Centunculus pentandrus.
peploides, *Baker, in Journ. Linn. Soc.* xx. (1884) 196.—Madag.
Philippsii, Hort. ex Duby, in DC. Prod. viii. 70—linifolia.
phoenicea, Scop. Fl. Carn. ed. II. i. 139 = arvensis.
pilosa, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 83 = Lysimachia japonica.
platyphylla, *Baudo, in Exped. Alg.* t. 67.—Algeria.
pulchella, Salisb. Prod. 120 = arvensis.
pumila, Sw. Prod. Veg. Ind. Occ. 40 = Centunculus pentandrus.
punctiflora, Stokes, Bot. Mat. Med. i. 305 = arvensis.
pusilla, Sallab. Prod. 121 = Centunculus minimus.
repens, DC. Fl. Fr. v. 381 = arvensis.
repens, *Pomel, Nouv. Mat. Fl. Atl.* 126.—Algeria.
rubricaulis, Boi. ex Duby, in DC. Prod. viii. 72(= *Micropyxis rubricaulis*).
serpens, *Hochst. ex A. DC. Prod.* viii. 568.—Abyss.
serpyllifolia, Dum. Fl. Belg. 29 = tenella.
sessilis, Salzm. ex Duby, in DC. Prod. viii. 72 = Centunculus pentandrus.
spicata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. 1. Art. iv. 6.—Malaya.
tenella, *Linn. Mant.* 335.—Europ.
tenuicaulis, *Baker, in Journ. Bot.* xx. (1882) 172.—Madag.
variicillata. All. Fl. Pedem. i. 87 = arvensis.
Webbiana, Penny, ex Loud. Gard. Mag. v. (1829) 471 = *Marryattae*.
Wellsiana ×, *Penny, ex Loud. Hort. Brit. Suppl.* i. 580.
Willmoreana, Don, ex Hook. Bot. Mag. t. 3380 = linifolia.

ANAGALLOIDES, Krock. Fl. Siles. ii. t. 398 (1790) = **Vandellia**, Linn. (Scrophul.).
procumbens, Krock. l. c. = V. erecta.

ANAGANTHOS, Hook. f. in Fl. N. Zeal. i. 236 (1854) = **Australina**, Gaudich. (Urticac.).

ANAGLYPHA, DC. Prod. v. 311 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 326.
OXYLAENA, Benth. in Hook. Ic. Pl. t. 1109 (1876).
acicularis, *Benth. l. c.* xii. 9. t. 1109.—Afr. austr.
aspera, *DC. Prod.* v. 311.—Afr. austr.

ANAGYRIS, Linn. Syst. ed. I. (1735); Gen. ed. I. 219 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 463.
barbata, R. Grah. in Wall. Cat. n. 5341 = Thermopsis barbata.
cretica, *Mill. Gard. Dict.* ed. VIII. n. 2.—Ins. Archip.
foetida, *Linn. Sp. Pl.* 374.—Reg. Mediterr.; Orient.
foetida, Lour. Fl. Cochinch. 260; ed. Willd. i. 318 = sinensis.
glauca, Hort. ex Loud. Hort. Brit. 162 = foetida.
indica, Hort. ex Steud. Nom ed. II. i. 88 = sinensis.
indica, Lindl. in Trans. Hort. Soc. vii. (1830) 245 = Piptanthus nepalensis.
inodora, *Lour. Fl. Cochinch.* 260.—Cochinch.
latifolia, *Brouss. ex Willd. Enum. Hort. Berol.* 489.—Teneriffa.
neapolitana, Tenore, Syll. 198 = foetida.
nepalensis, R. Grah. in Wall. Cat. n. 5340 = Piptanthus nepalensis.
sinensis, *Steud. ex DC. Prod.* ii. 98 (‘forte ad Ormosiam referend.’).—China.

ANAITIS, DC. Prod. v. 628 (1836) = **Sanvitalia**, Linn. (Compos.).
acapulcensis, DC. l. c. 629 = S. acapulcensis.

ANALECTIS, Juss. in Jaume St. Hil. Expos. ii. 362 (1805) = **Symphoremа**, Rozb. (Verben.).

ANALITON, Raf. Fl. Tellur. iii. 46 (1836) = **Rumex**, Tourn. (Polygon.).
bipinnatus, Raf. l. c. = R. lacerus.

ANALYRIUM, E. Mey. Zwei Pfl. Docum. 168 (1843), *nomen*; Presl, in Abh. Boehm. Ges. V. iii. (1845) 505 = **Peucedanum**, Koch (Umbell.).
millefolium, E. Mey. l. c. = P. millefolium.

ANAMENIA, Vent. Malm. 22. t. 22 (1803) = **Knowltonia**, Salisb. (Ranunc.).
capensis, Hoffmgg. Vers. Pfl. Nachtr. i. 204 = K. rigida.
coriacea, Vent. Malm. 22 = K. rigida.
dauciifolia, Vent. ex Cav. in Anal. Clenc. Nat. vi. (1808) 344 = K. tenuifolia.
gracilis, Vent. Malm. i. 22, obs. = K. gracilis.
hirsuta, Vent. ex Cav. in Anal. Clenc. Nat. vi. (1808) 344 = K. hirsuta.
laserpitiifolia, Vent. ex Cav. l. c. = K. vesicatoria.
vesicatoria, Dum.-Cours. Bot. Cult. iv. 488 = K. vesicatoria.

ANAMIRTA, Colebr. in Trans. Linn. Soc. xiii. (1822) 52. *MENISPERMACEAE*, Benth. & Hook. f. i. 35.
Bauerianа, Endl. Atakt. Bot. t. 29, 40 = A. Cocculus.
Cocculus, Wight & Arn. Prod. i. 446 = paniculata.
flaweccna, Miq. Fl. Ind. Bat. i. ii. 79 = A. Cocculus.
floreusana, C. Muell. in Walp. Ann. vii. 65, sphalm. = flavescens.
jucunda, *Miers, Contr. Bot.* iii. 55.—Java.
lemniscata, *Miers, l. c.* 54.—Java.
Lourieri, *Pierre, Fl. For. Coch. Chin.* t. 110.—Cambodia.
luctuosa, *Miers, Contr. Bot.* iii. 55.—Java.
paniculata, *Colebr. in Trans. Linn. Soc. xiii.* (1822) 66.—Ind. or.
populifiera, *Miers, Contr. Bot.* iii. 54.—Timor.
racemosa, Colebr. ex Steud. Nom. ed. II. i. 88 = Cocculus suberosus.
tenifera, Miers, Contr. Bot. iii. 53 = paniculata.

ANAMOMIS, Griseb. Fl. Brit. W. Ind. 240 (1864) = **Myrtus**, Linn.
acuminata, Griseb. l. c. in obs. = Eugenia acuminata.
ferruginea, Griseb. Cat. Fl. Cub. 91 = Eugenia ferruginea!
fragrans, Griseb. Fl. Brit. W. Ind. 240 = Myrcia fragrans!

ANAMOMIS :—
Guayabillo, Griseb. Cat. Pl. Cub. 90 = Eugenia Guayabillo.
punctata, Griseb. Fl. Brit. W. Ind. 240 = Myrcia Balbisiana.

ANAMORPHA, Karst. & Triana, in Linnaea, xxviii. (1856) 443 = **Malochia**, Linn. (Stercul.).
walkeriodes, Karst. & Triana l. c. = M. lupulina.

ANANAS, Gaertn. Fruct. i. 30. t. 11 (1788) = **Bromelia**, Linn.
Pinguin, Gaertn. l. c. = B. Pinguin.

ANANAS, Tourn. ex Linn. Syst. ed. I. (1735); Mill. Gard. Dict. ed. VI. (1759); Adans. Fam. ii. 67 (1763). *BROMELIACEAE*, Benth. & Hook. f. iii. 662.
ACANTHOSTACHYS, Link, Klotzsch & Otto, Ic. Pl. Rar. i. 21. t. 9. (1841).
ANANASSA, Lindl. Bot. Reg. sub t. 1068 & t. 1081 (1827).
arvensis, *Steud. Nom.* ed. II. i. 88.—Bras.
bractеatus, *Schult. f. Syst.* vii. 1286.—Bras.
debilis, Schult. f. l. c. 1287 = sativus.
glaber, Mill. Gard. Dict. ed. VIII. n. 3 (et corr.) = sativus.
lucidus, Mill. l. c. n. 4 = sativus.
macrodontes, *E. Morr. Belg. Hortic.* xxviii. (1878) 140. t. 4 et 5.—Bras.
Mordilona, Hort. Linden, ex E. Morr. in Belg. Hortic. xxix. (1879) 302 = sativus.
muricatus, Schult. f. Syst. vii. 1287 = bracteatus.
ovatus, Mill. Gard. Dict. ed. VIII. n. 1 = sativus.
Porteanus, Hort. Veitch, ex C. Koch, Wochenschr. xiv. (1871) 180 = sativus.
pyramidalis, Mill. Gard. Dict. ed. VIII. n. 2 = sativus.
Sagenaria, Schult. f. Syst. vii. 1286 = bracteatus.
sativus, *Schult. f. l. c.* 1283.—Am. trop.
semiarratus, Schult. f. l. c. 1286 = sativus.
serotinus, Mill. Gard. Dict. ed. VIII. n. 5 = sativus.
viridis, Mill. l. c. n. 6 = sativus.

ANANASSA, Lindl. Bot. Reg. sub t. 1068 et t. 1081 (1827) = **Ananas**, Tourn. (Bromel.).
bracamorensis, Linden, Cat. n. 98. (1876) 3 = Ananas sativus.
bracteata, Lindl. Bot. Reg. sub t. 1068 et t. 1081 = Ananas bractеatus.
debilis, Lindl. l. c. sub t. 1068 = Ananas sativus.
lucida, Lindl. l. c. = Ananas sativus.
Mordilona, Linden, Cat. n. 29. (1869) 2 = Ananas sativus.
muricata, D. Dietr. Syn. Pl. ii. 1088 = Ananas bractеatus.
Porteana, Carr. in Rev. Hortic. (1878) 140 = Ananas sativus.
Sagenaria, D. Dietr. Syn. Pl. ii. 1088 = Ananas bractеatus.
sativa, Lindl. Bot. Reg. sub t. 1068 = Ananas sativus.
semiarrata, D. Dietr. Syn. Pl. ii. 1088 = Ananas sativus.

ANANDRIA, Slegesb. ex Linn. Amoen. Acad. i. 243. t. 11 (1745) = **Gerbera**, Gronov. (Compos.).
Bellidiastrum, DC. Prod. vii. 40 = G. Anandria.
dimorpha, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 11. 68 = G. Anandria.
discoidea, Less. in Linnaea, v. (1830) 247. t. 3 = G. Anandria.
vadata, Less. l. c. 246. t. 4 = G. Anandria.

ANANTHERIX, Nutt. Gen. Am. i. 189 (1818) = **Asclepiodora**, A. Gray (Asclep.).
angustifolia, Rafin. Atl. Journ. 146.—Am. bor.
Bungei, G. Don, Gen. Syst. iv. 146 = Pycnostelma chinense.
cornuta, A. Gray, in Proc. Am. Acad. xii. (1877) 66 = Asc. viridis.
decumbens, Nutt. in Trans. Am. Phil. Soc. v. (1837) 203 = Asc. decumbens.
grandiflora, Rafin. in New Fl. Am. iv. 89.—Am. bor.
Nuttaliana, G. Don, Gen. Syst. iv. 146 = Asc. decumbens.
ovata, Rafin. Atl. Journ. 146.—Am. bor.
paniculata, Nutt. in Trans. Am. Phil. Soc. v. (1837) 203 = Asc. viridis.
pumilio, Nutt. l. c.—Am. bor.
Torreyana, G. Don, Gen. Syst. iv. 146 = Asc. viridis.
verrucosa, Rafin. New Fl. Am. iv. 60 = Asc. viridis.
viridis, Nutt. Gen. Am. i. 189 = Asc. viridis.

ANANTHOCYCLUS, Vaill. ex Linn. Syst. ed. I. (1785) = **Cotula**, Tourn. (Compos.).

ANANTHOPUS, Rafin. Fl. Ludov. 20 (1817) = **Commelina**, Linn.
clandestinus, Rafin. l. c. 21 = C. ignorata.
cordatus, Rafin. l. c. 22 = C. Rafinesquei.
undulatus, Rafin. Fl. Tellur. iii. 70 = C. undulata.

ANAPHALIS, DC. Prod. vi. 271 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 303.
MARGARIPES, DC. ex Steud. Nom. ed. II. ii. 101 (1841).
adnata, *DC. Prod.* vi. 274.—Reg. Himal.
alata, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 478.—As. or.
araneosa, *DC. Prod.* vi. 275.—Reg. Himal.
aristata, *DC. in Wight, Contrib.* 21.—Ind. or.
Beddomei, *Hook. f. Fl. Brit. Ind.* iii. 282.—Ind. or.
brevifolia, *DC. Prod.* vi. 278.—Zeylan.
Burmo, DC. l. c. 275 = araneosa.
chinensis, DC. l. c. 273 = Roylean.
cinnomomea, *C. B. Clarke, Comp. Ind.* 104.—Ind. or.; China; Japon.
contorta, *Hook. f. Fl. Brit. Ind.* iii. 284.—Reg. Himal.
cuneifolia, *Hook. f. l. c.* 280.—Reg. Himal.
cutchica, *C. B. Clarke, Comp. Ind.* 111.—Kutch.
elliptica, DC. Prod. vi. 274 = oblongs.
Falconeri, C. B. Clarke, Comp. Ind. 107 = contorta.
fruticosa, *Hook. f. Fl. Brit. Ind.* iii. 282.—Zeylan.
Griffithii, *Hook. f. l. c.* 280.—Reg. Himal.
Hancockii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 479.—China.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 283.—Reg. Himal.
japonica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 480.—Japon.
javanica, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip* 134.—Java.
lactea, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 479.—As. or.
leptophylla, *DC. Prod.* vi. 273.—Ind. or.
limaria, DC. l. c. 274 = marcescens ?
longifolia, DC. l. c.—Java.
marcescens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 286.—Ind. or.
margaritacea, *Benth. & Hook. f. Gen.* ii. 308.—Am. bor.
monocephala, DC. Prod. vi. 272 = nubigena.
mucronata, DC. l. c. = nubigena.
neelgerryana, *DC. l. c.*—Ind. or.
Notoniana, *DC. l. c.* 278.—Ind. or.
nubigena, *DC. l. c.* 272.—Reg. Himal.
oblonga, *DC. l. c.* 274.—Ind. or.
oligandra, *DC. l. c.* 276.—Madag.
polylepis, DC. l. c. 272 = Royleana.
pterocaulon, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 478.—As. or.
racemifera, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 310.—As. or.
Royleana, *DC. Prod.* vi. 272.—Reg. Himal.
Royleana, C. B. Clarke, Comp. Ind. 104 = Griffithii.
semidecurrens, C. B. Clarke, l. c. 109 = araneosa.
sinica, *Hance, in Journ. Bot.* xii. (1874) 261.—China.
Stoliczkai, *C. B. Clarke, Comp. Ind.* 108.—Tibet. occ.
subumbellata, *C. B. Clarke, l. c.* 104.—Reg. Himal.
tenella, DC. Prod. vi. 273 = contorta.
Thwaitesii, *C. B. Clarke, Comp. Ind.* 110.—Zeylan.
Timmua, D. Don, Prod. Fl. Nep. 174 = cinnamomea.
triplinervis, *Sims, ex C. B. Clarke, Comp. Ind.* 105.—Reg. Himal.
triplinervis, Benth. ex Hance, in Journ. Bot. xvi. (1878) 12 = Hemsleyii.
virgata, *Thoms. in C. B. Clarke, Comp. Ind.* 107.—Reg. Himal.
viscida, *DC. Prod.* vi. 274.—Java.
Wightiana, Thw. Enum. Pl. Zeyl. 166 = zeylanica.
Wightiana, *DC. Prod.* vi. 278.—Ind. or.
xylorhiza, *Sch. Bip. ex Hook. f. Fl. Brit. Ind.* iii. 281.—Reg. Himal.
yedoensis, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 480.—Japon.
zeylanica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 286.—Zeylan.
Zollingeri, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 84 = Antennaria acuatilis.
Zollingeriana, Sch. Bip. ex Zoll. Syst. Vers. Ind. Archip. 134 = praec.

ANAPHRAGMA, Stev. in Bull. Soc. Nat. Mosc. iv. 265 (1839) = **Astragalus**, Tourn. (Legum.).

ANAPHRENIUM, E. Mey. ex Endl. Gen. Suppl. i. 1425 (1841). *ANACARDIACEAE*, (Benth. & Hook. f. i. 419).
HERRIA, Meisn. Gen. Comm. 55 (1837).
OZOROA, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 91. t. 1.
ROXEMERIA, Thunb. Nov. Gen. Pl. ix. 130 (1798), pro parte.
abyssinicum, *Hochst. in Flora*, xxvii. (1844) 32.—Afr. trop.
argenteum, *E. Mey. Zwei Pfl. Docum.* 163.—Afr. austr.
concolor, *E. Mey. l. c.*—Afr. austr.
dispar, *E. Mey. l. c.*— Afr austr.
longifolium, Bernh. in Flora, xxvii. (1844) 1. 849 = Frotorhus longifolia.
mucronatum, *Bernh. l. c.*—Afr. austr.
mucronifolium, *Sond. in Harv. & Sond. Fl. Cap.* i. 522 = praec.
paniculosum *Engl. in DC. Monog. Phan.* iv. 858.—Afr. austr.
pulcherrimum, *Schweinf. Fl. Aethiop. Beitr.* i. 92.—Afr. trop.

ANAPHYLLUM, Schott, Gen. Aroid. t. 83 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 996.
Wightii, *Schott, l. c.*—Ind. or.

ANAPODOPHYLLUM, Tourn. ex Moench, Meth. 277 (1794) = **Podophyllum**, Linn. (Berberid.).
peltatum, Moench, l. c. = P. peltatum.

ANARMODIUM, Schott, in Bonplandia, ix. 368 (1861) = **Premnoulus**, Schott (Aroid.).
canarense, Schott, l. c. = D. canariensis.

ANARMOSA, Miers, ex Hook Bot. Misc. iii. (1833) 338 = **Tetilla**, DC. (Saxifrag.).
gracilis, Miers, ex Hook. l. c. = T. hydrocotylifolia.

ANARRHINUM, Desf. Fl. Atlant. ii. 51 (1800). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 933
CARDIOTHECA, Ehrenb. ex Steud. Nom. ed. II. i. 329 (1840).
SIMMULETA, Forsk. Fl. Aegypt. Arab. 115 (1775).
abyssinicum, *Vatke, & Spach, Illustr. Pl. Or.* v. 52. t. 447.—Abyss.
arabicum, *Vatke, & Spach, l. c.* v. 50. t. 446.—Arabia.
bellidifolium, *Desf. Fl. Atlant.* ii. 51; *Chav. Monog.* 175. t. 10.—Reg. Mediterr.
bellidifolium, Fenzl, ex Jaub. & Spach, Illustr. Pl. Or. v. 54 = orientale.
brevifolium, *Coss. & Kral. in Bull. Soc. Bot. Fr.* iv. (1857) 407.—Afr. bor.
canadense, Poir. Encyc. Suppl. i. 334, in obs. = Linaria canadensis.
corsicum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 41 = bellidifolium.
crassifolium, Willd. Sp. Pl. iii. 261 = Linaria origanifolia.
Duriminium, *Pers. Syn.* ii. 159 — Lusitania.
ecalcaratum, Fenzl, ex Jaub. & Spach, Illustr. Pl. Or. v. 54 = orientale.
fruticosum, *Desf. Fl. Atlant.* ii. 52.—Afr. bor.
fruticosum, Schimp. ex Benth. in DC. Prod. x. 289 = orientale.
hirsutum, Hoffingg. & Link, Fl. Port. 227. t. 33 = A. Duriminium.
incarnatum, Poir. Encyc. Suppl. iv. 18, in obs. = Linaria bipartita.
Linnaeanum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 41 = bellidifolium.
laxiflorum, *Boiss. Elench.* 71.—Hispan.
lusitanicum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 41 = bellidifolium.
murale, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 557 = A. Duriminium.
orientale, *Benth. in DC. Prod.* x. 289.—Oriena.
pedatum, *Desf. Fl. Atlant.* ii. 51. t. 141.—Afr. bor.
pedicellatum, *T. Anders. in Journ. Linn. Soc. v. Suppl.* i. (1860) 26.—Aden.
pubescens, *Fresen. in Mus. Senkenb.* i. 89.—Arab. petr.
pubescens, Hrst. ex Loud. Hort. Brit. 250 = A. Duriminium.
tenellum, *Willd. Sp. Pl.* iii. 262.—Hispan.
violaceum, Dulac, Fl. Hautes-Pyr. 378 = bellidifolium.

ANARTHRIA, R. Br. Prod. 248 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1080.
canaliculata, Nees, in Lehm. Pl. Preiss. ii. 62 = gracilis.
gracilis, *R. Br. Prod.* 249.—Austral
gracilis, Nees, ex Benth. Fl. Austral. vii. 216 = Lepyrodia Anarthria.
grandiflora, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 50 = gracilis.
humilis, Nees, l. c. = gracilis.
lachnocaules, Nees, in Lehm. Pl. Preiss. ii. 62 = gracilis.
laevis, *R. Br. Prod.* 249.—Austral.
pauciflora, R. Br. l. c. 249 = prolifera.
polyphylla, *Nees, in Lehm. Pl. Preiss.* ii. 63.—Austral.
prolifera, *R. Br. Prod.* 249.—Austral.
scabra, *R. Br. l. c.*—Austral.

ANARTHROPHYLLUM, Benth. in Benth. & Hook f. Gen. i. 478 (1865). *LEGUMINOSAE*, Benth & Hook. f. i. 478.
audiculum, *Phil. f. Cat. Pl. Chil* 52.—Chili.
Beigii, *Hieron. in Bot. Acad. Nac. Cordova*, iii. (1879) 340.—Reg. argent.
Cumingii, *Phil. f. Cat. Pl Chil.* 52.—Chili.
desideratum, *Benth. in Benth. & Hook. f. Gen.* i. 478. —Patagon.
elegans, *Phil. f. Cat. Pl. Chil.* 52.—Chili.
juniperinum, *Phil. f. l. c.*—Chili.
rigidum, *Hieron. in Bot. Acad. Nac. Cordova*, iii (1879) 340.—Reg. Argent.
umbellatum, *Phil. f. Cat. Pl. Chil.* 52.—Chili.

ANAKTHROSYNE, E. Mey. Comm. Pl. Afr. Austr. 124 (1835) = **Pseudarthria**, Wight & Arn. (Legum.).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 204 = Desmodium spirale
cordata, Klotzsch, in Peters, Reise Mossamb. Bot. 39. t. 7 = Desmodium latifolium.
densiflora, Klotzsch, l. c. 41 = P. Hookeri.
gracilis, Klotzsch, l. c. 40 = P. confertiflora.
robusta, E. Mey. Comm. Pl. Afr. Austr. 124 = P. Hookeri.

ANARTIA, Miers, Apocyn. S Amer. 79. (1878) = **Tabernaemontana**, Linn. (Apocyn.).
bogotensis, Miers, l. c. 82.—N. Granat.
flavescens, Miers, l. c. = Malouetia flavescens.
fragrans, Miers, l. c. = T. flavicans
glabrata, Miers, l. c. 81 = T. oblongifolia.
Meyeri, Miers, l. c. 80 = T. Meyeri.
recurva, Miers, l. c. = T. recurva.
Wulfschlaegelii, Miers, l. c. 81 = T. Wullschlaegelii.

ANASCHOVADI, Adans. Fam. ii. 122 (1763) = **Elephantopus**, Vaill. (Compos.).

ANASSER, Juss. Gen. 150 (1789) = **Geniostoma**, Forst. (Logan.).
Borbonum, J. F. Gmel. Syst. i. 303 = G. borbonicum.
borbonica, Lam. Illustr. ii. 40 = G. borbonicum.
moluccana, Lam. l. c. = Strychnos Pseudo-quina.
Rumphii, Span. in Linnaea, xv. (1841) 325 = Pittosporum timorense.

ANASSERA, Pers. Syn. i. 265 (1805) = praec.

ANASTATICA, Linn. Gen. ed. I. 191 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 71.
HIERICONTIS, Adans. Fam. ii. 421 (1763).
IERIKONTIS, Adans. l. c. 565 (1763).
hierochuntica, Crantz, Strip. Austr. ed. II. fasc. i. 7 = Euclidium syriacum.
hierochuntica, *Linn. Sp. Pl.* ed. II. 905.—Orient.
littoralis, Salisb. Prod. 286 = hierochuntica.
syriaca, Linn. Sp. Pl. ed. II. 895 = Euclidium syriacum.

ANASTRABE, E. Mey. ex Benth. in Hook. Comp. Bot. Mag. ii. (1836) 84. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 987.
integerrima, *E. Mey. ex Benth. l. c.*—Afr. austr.
serrulata, *E. Mey. ex Benth. l. c.*—Afr. austr.

ANASTRAPHIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 295. *COMPOSITAE*, Benth. & Hook. f. ii. 491.
lilaefolia, *D. Don, l. c.*—Am. trop.
intertexta, *Wright, ex Griseb. Cat. Pl. Cub.* 158.—Cuba.
mantonoia, *Wright, ex Griseb. l. c.*—Cuba.
microcephala, *Griseb. l. c.*—Cuba.

ANASTREPHIA, Decne. in Orb. Dict. i. 482 (1841) = **Anastraphia**, D. Don (Compos.).

ANASTROPHEA, Wedd. in DC. Prod. xvii. 78 (1873) = **Sphaerothylax**, Bisch. (Fodostem.).
abyssinica, Wedd. l. c. 78.—Abyss.

ANASTROPHUS, Schlecht. in Bot. Zeit. viii. (1850) 681 = **Paspalum**, Linn. (Gram.).
barbatus, *compressus*, *dissitiflorus*, *nematanthus*, *pectinatus*, *pellitus*, *platyculis*, *platyculmis*, *pulchellus*, *setifolius*, Schlecht, l. c. 682, 683, omnes Paspali homonymnae.

ANASYLLIS, E. Mey. Zwei Pfl. Docum. 16 (1848) = **Loxostylis**, Spreng. (Anacard.)
angustifolia, E. Mey. l. c.—L. alata.
latifolia, E. Mey. l. c.—L. alata.

ANATHALLIS, Barb. Rodrig. Orch. Nov. i. 23 (1877) = **Pleurothallis**, R. Br. (Orchid.).
densiflora, Rodrig. l. c. ii. 78.—Bras.
fasciculata, Rodrig. l. c. i. 28.—Bras.
hetaerpala, Rodrig. l. c. ii. 75.—Bras.
heterophylla, Rodrig. l. c. ii. 74.—Bras.
micrantha, Rodrig. l. c. ii. 76.—Bras.
nectarifera, Rodrig. l. c. ii. 74.—Bras.
ornaperma, Rodrig. l. c. ii. 78.—Bras.
parahybunensis, Rodrig. l. c. ii. 76.—Bras.
pulvinata, Rodrig. l. c. ii. 77.—Bras.
racemosa, Rodrig. l. c. i. 24.—Bras.
secunda, Rodrig. l. c. ii. 72.—Bras.

ANATHERIX, Steud. Nom. ed. I. 41 (1821) err. typ. = **ANANTHERIX**, Nutt. = **Asclepiodora**, A. Gray (Asclep.).

ANATHERUM, Beauv. Agrost. 182. t. 22. f. 10, 11 (1812) = **Andropogon**, Linn. (Gram.).
abortivum, Beauv. Agrost. 169 = Panicum abortivum.
Bertoroanum, Spreng. ex Roem. & Schult. Syst. ii. Mant. ii. 448 = Imperata exaltata.
bicorne, Beauv. Agrost. 138 = Androp. bicornis.
brasiliense, Spreng. ex Steud. Nom. ed. II. i. 84 = Androp. leucostachyus.
caudatum, Schult. Mant. ii. 445 = Imperata exaltata.
domingrense, Roem. & Schult. Syst. ii. 809 = Androp. leucostachyus.
glabrum, Schult. Mant. ii. 446 = Androp. intermedius.
holcoides, Nees, Agrost. Bras. 324 = Saccharum holcoides.
inerme, Griseb. Cat. Pl. Cub. 236 = Androp. spathiflorus.
macrurum, Griseb. Pl. Wright. Cub. ii. 584 = Androp. macrourus.
megapotamicum, Spreng. Syst. Cur. Post. 38 = Elionurus rostratus.
montanum, Roem. & Schult. Syst. Mant. ii. 445 = Androp. montanus.
muricatum, Beauv. Agrost. 150 = Androp. squarrosus.
muticum, Beauv. l. c.= Chloris petraea ?
nitidum, Spreng. Syst. i. 290 = Chrysopogon fascus.
parviflorum, Spreng. l. c. = Chrysopogon violaceus.
pedunculatum, Desv. Opusc. 70.—Ind. occ.
portoricense, Spreng. l. c. = Imperata exaltata.
refractum, Beauv. Agrost. 150 = Androp. refractus.
scoparium, Desv. Opusc. 70.—Afric.
setifolium, Spreng. Syst. i. 290 = Androp. setifolium.
spathiflorum, Griseb. Cat. Pl. Cub. 236 = Androp. spathiflorus.
squarrosum, Beauv. Agrost. 138, 150 = Panicum abortivum.
tripaccoides, Spreng. Syst. i. 290 = Elionurus tripsacoides.
virgatum, Desv. Opusc. 71 = Androp. spathiflorus.
virginicum, Spreng. Pugill. ii. 16 = Androp. virginicus.

ANATROPA, Ehrenb. in Linnaea, iv. (1829) 402 = **Tetradilella**, Stev. (Rutac.).
tenella, Ehrenb. l. c. 404 = T. salsa.

ANAUA, Miq. Fl. Ind. Bat. Suppl. 410 (1860) = **Hemicyclia**, Wight & Arn. (Euphorb.).
sumatrana, Miq. l. c. = H. sumatrana.

ANAUXANOPETALUM, Teijsm. & Binn. in Journ. Bot. Néerl. i. (1861) 268 = **Swintonia**, Griff. (Anacard.).
Schwenkii, Teijsm. & Binn. l. c.= S. Schwenkii.

ANAVINGA, Adans. Fam. ii. 448 (1763); Lam. Illustr. t. 356 = **Cassaria**, Jacq. (Samyd.).
conocladifolia, Rafin. Sylva Tellur. 149 = C. conocladifolia.
fuliginosa, Blanco, Fl. Filip. ed. I. 272 = C. fuliginosa.
ilicifolia, Rafin. Sylva Tellur. 149 = C. ilicifolia.
lanceolata, Lam. Encyc. i. 148 = C. tomentosa ?
ovata, Lam. l. c. = C. ovata.
parvifolia, Lam. Tabl. ii. 484 = C. parvifolia.
Samyda, Gaertn. f. Fruct. iii. 240. t. 234 = C. Samyda.

ANAXAGORAEA, Mart. Fl. Bras. xiii. i. 39 (1841) = seq.

ANAXAGOREA, St. Hil. in Bull. Soc. Philom. (1825) 91. *ANONACEAE*, Benth. & Hook. f. i. 25.
RINGALOCARPUS, Teijsm. & Binn. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 22 (1865).
acuminata, *St. Hil. l. c.*—Bras.
brevipes, *Benth. in Hook. Kew Journ.* v. (1853) 8.—Bras.
crassipetala, *Hemsl. Diagn. Pl. Nov.* 2.—Nicaragua.
fruticosa, *Schott, in Flora*, iii. (1869) 302.—Malaya ?
javanica, *Blume, Fl. Jav. Rumph.* 66.—Java.
luzonensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 27.—Ind. or.; Ins. Philipp.
Meyeriana, *Zoll. in Linnaea*, xxix. (1857–58) 323.—Java.
pbaeocarpa, *Mart. Fl. Bras.* xiii. i. 40.—Bras.
prinoides, *St. Hil. & A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 211.—Bras.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* i. 383.—Sumatra.
zeylanica, Hook. f. & Thoms. Fl. Ind. i. 144 = Insc-nensis.

ANAXETON, Gaertn. Fruct. ii. 406 (1791); Cass. in Dict. Sc. Nat. xxvi. (1823) 59; xxxiv. 87 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 314.
ARGYRANTHUS, Neck. Elem. i. 75 (1790).
arborescens, Less. Syn. Comp. 254 = arboreum.
arboreum, *Gaertn. Fruct.* ii. 407.—Afr. austr.
asperum, *DC. Prod.* vi. 268.—Afr. austr.
crispum, Gaertn. Fruct. ii. 407 = Helichrysum crispum.
floridum, Lam. Tabl. iii. 269 = arboreum.
foetidum, Gaertn. Fruct. ii. 406. t. 166 = Helichrysum foetidum.
hirsutum, *Less. Syn. Comp.* 256.—Afr. austr.
imbricatum, Gaertn. Fruct. ii. 407 = Helichrysum undifolium.
nycthemerum, *Less. Syn. Comp.* 256.—Afr. austr.
revolutum, *DC. Prod.* vi. 268.—Afr. austr.
septentrionalis, Vatke, in Oestr. Bot. Zeitschr. xxvii. (1877) 194 = Gnaphalium stenophyllum.
virgatum, *DC. Prod.* vi. 268.—Afr. austr.

ANAXETON, Schrank, in Denkschr. Acad. Munch. viii. (1824) 146 et 169 = **Heliptesum**, DC. (Compos.).
corymbosum, Schrank, l. c. = Heliptesum eximium.
racemosum, Schrank, l. c. = Helichrysum mucronatum.

ANBLATUM, Tourn. ex Adans. Fam. ii. 207 (1763) = **Lathraea**, Linn. (Orobanch.).
tartaricum, Fisch. Hort. Gorenk. ed. II. 19 = Phelipaea salsa.
Tournefortii, G. Don, Gen. Syst. iv. 685 = Lathraea Squamaria.

ANCANA, F. Muell. Fragm. v. 37. t. 35 (1865) = **Uxona**, Linn. f. (Anonac.).
sternopetala, F. Muell. l. c.= U. Ancana.

ANCANTHIA, Steud. Nom. ed. II. i. 85 (1840) err. typ. = seq.

ANCATHIA, DC. in Guill. Arch. Bot. ii. (1833) 381 = **Ontoxa**, Tourn. (Compos.).
ignaria, DC. in Deless. Ic. Sel. iv. t. 73.—C. ignatiaria.

ANCHISTRA, A. St. Hil. in Ann. Sc. Nat. Sér. I. ii. (1824) 383; in Mém. Mus. Par. xi. (1824) 464; Fl. Us. i. 19 (1824); Fl. Rom. Bras. 290 (1854). *VIOLARIEAE*, Benth. & Hook. f. i. 117.
ozalitata, *Bichl. in Mart. Fl. Bras.* xiii. i. 355.—Bras.
pyrifolia, G. Don, Gen. Syst. i. 340 = salutaris.
pyrifolia, *A. St. Hil. Fl. Rom. Bras.* 290.—Bras.
Roquefeuilliana, Spreng. Syst. iv. Cur. Post. 98 = salutaris.

ANCHIETA—
salutaris, *A. St. Hil. in Ann. Sc. Nat.* Sér. I. ii. (1824) 383.—Bras.
Salloviana, *Cham. & Schlecht. in Linnaea*, ii. (1827) 148.—Bras.

ANCHIONIUM, Reichb. Consp. 181. err. typ. = **Anchonium**, DC.

ANCHOMANES, Schott, in Oestr. Bot. Wochenbl. iii. (1853) 314. *AROIDEAE*, Benth. & Hook. f. iii. 978.
difformis, Engl. in DC. Monog. Phan. ii. 304 = Hookeri.
dubius, *Schott, Prod. Aroid.* 185.—Afr. occ.
Hookeri, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 314.—Afr. trop.

ANCHONIUM, DC. in Mém. Mus. Par. vii. (1821) 242. *CRUCIFERAE*, Benth. & Hook. f. i. 101.
Billardierii, *DC. l. c.*—Syria.
clichryifolium, Boiss. Fl. Orient. i. 240 = Tournefortii.
Tournefortii, *Boiss. in Ann. Sc. Nat. Sér.* II. xvii. (1842) 336.—Orient.

ANCHUSA, Linn. Syst. ed. I. (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 855.
BRUNNERA, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 582.
BUGLOSSOIDES, Tausch, ex Reichb. Fl. Germ. Excurs. 844 (1831).
BUGLOSSUM, Gaertn. Fruct. i. 322. t. 67 (1788).
CARVOLOPHA, Fisch. & Trautv. Ind. Sem. Hort. Petrop. iii. 51 (1837).
PENTAGLOTTIS, Tausch, in Flora, xii. (1829) 648.
STOMOTECHIUM, Lehm. Asperif. 395 (1818).
Adami, [Pieri] in Ionios Anthol. v. 182 = sempervirens.
aegyptiaca, DC. Prod. x. 48 = flava.
aegyptiaca, Maly, ex Nym. Consp. iii. 511 = Lycopsis variegata.
affinis, *R. Br. in Salt, Voy. Abyss. App.* 62.—Abyss.
africana, *Burm. f. Fl. Cap.* 4.—Afr. austr.
Agardhii, *Lehm. Asperif.* 341.—Sibir.
aggregata, *Lehm. l. c.* 319. t. 47.—Sicil.; Oriens.
alpestris, Stev. in Trans. Linn. Soc. xi. (1815) 409. t. 32 = Nonea alpestris.
altissima, *Desf. Cat. Hort. Par.* ed. III. 207.—Hab. ?
amoena, J. F. Gmel. Syst. ii. 218 = Italica.
amplexicaulis, Sibth. & Sm. Prod. Fl. Graec. i. 116 = hybrida.
angustifolia, Balb. Misc. Alt. 9 = undulata.
angustifolia, DC. Fl. Fr. ed. III. iii. 682 = officinalis.
angustifolia, Georgi, It. i. 200 (= *Stenosolenium saxatile*).
angustifolia, *Linn. Sp. Pl.* 133.—Europ. austr.
angustifolia, Willd. Enum. Hort. Berol. i. 178 = ochroleuca.
angustissima, Bourg. ex Nym. Consp. 510 = undulata.
angustissima, C. Koch, in Linnaea, xxi. (1849) 688 = ochroleuca.
annua, Pall. Ind. Taur. ex A. DC. in DC. Prod. x. 48 = stylosa.
arvalis, Reichb. Fl. Germ. Excurs. 843 = officinalis.
arvensis, Bieb. Fl. Taur. Cauc. i. 128 = Lycopsis arvensis.
arvensis, Tausch, in Flora, vii. (1824) 388 = officinalis.
aspera, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 248 = ochroleuca.
asperrima, Delile, Fl. Aeg. Illustr. 55 = Arnebia hispidissima.
atlantica, *Ball, in Journ. Bot.* xi. (1873) 373.—Marocco.
Auchert, *A. DC. in DC. Prod.* x. 49.—As. Min.
aurea, Mill. Gard. Dict. ed. VIII. n. 9 = italica.
aurea, Schur, Enum. Pl. Transs. 466 = ochroleuca.
Barrelieri, *Vitm. Summa Pl.* i. 338.—Europ.; As. Min.
Beauverdiana, Nym. Consp. iii. 510 = officinalis.
biceps, *Vent, in Flora*, iv. (1821) 146.—Ital.
brachiolata, Viv. Fl. Lib. Spec. 10. t. 4 = Alkanna tinctoria.
balbosa, [*Pieri*] *in Ionios Anthol.* v. 182.—Corcyra.
calocrus, *Boiss. Voy. Espagne*, ii. 681. t. 132.—Hispan.
campestris, Mahl. Cat. 29 = Lithospermum tenuiflorum.
canescens, Stev. ex Ledeb. Fl. Ross. iii. 128 = ochroleuca.

ANCHUSA :—
Capitisi, Moris, Stirp. Sard. Elench. ii. 6 = hybrida.
capensis, *Thunb. Prod. Fl. Cap.* 84.—Afr. austr.
Caustiana, Fenzl, in Frieder. Reise in N. Griech. 283 = Pulmonaria mollis.
cespitosa, *Lam. Encyc.* i. 504.—Creta.
ciliata, *Cham. in Linnaea*, iv. (1829) 440.—Bras.
ciliata, Schlecht. ex Ledeb. Fl. Ross. iii. 118 = ochroleuca.
clavata, Viv. ex DC. Prod. x. 52 = undulata.
coronensis, Schur, Enum. Pl. Transs. 466 = Barrelieri.
cretica, Mill. Gard. Dict. ed. VIII. n. 7 = Lycopsis variegata.
crispa, *Moris*, *Stirp. Sard. Elench.* iii. 9.—Sardinia.
crispa, Viviani, Fl. Cors. App. 1 = hybrida.
crispata, Presl, Fl. Sic. i. p. xxxiii = hybrida.
cruciata, Cham. in Linnaea, iv. (1829) 438 = Antiphytum cruciatum.
dasyantha, *Cham. l. c.* 437.—Bras.
deflexa, Lehm. Ind. Sem. Hort. Hamb. (1822) 17 = hispida.
divaricata, Murr. in Nov. Comm. Gotting. (1776) 25 = Nonea lutea.
Dregei, *A. DC. in DC. Prod.* x. 44.—Afr. austr.
dubia, Nocca, Hort. Ticin t. 3 = Nonea lutea.
echinata, Lam. Encyc. i. 505 = strigosa.
echioides, Bieb. Fl. Taur. Cauc. i. 123 = Macrotomia echioides.
exasperata, Clark, Trav. ed. III. i. 625 = ochroleuca.
flava, Forsk. Fl. Aegypt. Arab. 40 = aegyptiaca.
foliosa, Lehm. ex DC. Prod. x. 53 = Myosotis foliosa.
Gmelini, Ledeb. ex Spreng. Syst. i. 562 = ochroleuca.
granatensis, *Boiss. Voy. Espagne*, ii. 480.—Hispan.
granatensis, Lange, Pugill. 191 = hybrida.
hirta, Michx. ex DC. Prod. x. 58 = Lithospermum hirtum.
hispida, *Forsk. Fl. Aegypt. Arab.* 40.—Oriens; Turkest. : Ind. bor.
hispida, Pall. ex Ledeb. Fl. Ross. iii. 120 = stylosa.
hispidissima, Sieber, ex DC. Prod. x. 94 = Arnebia hispidissima.
hyalotricha, Boiss. Fl. Or. Nov. Dec. ii. 9 = Aucheri.
hybrida, Boiss. ex DC. Prod. x. 42 = granatensis.
hybrida, *Tenore*, *Fl. Nap.* i. 65. t. 11.—Ital. ; Oriens.
incana, Ledeb. Fl. Ross. iii. 117 = ochroleuca.
incarnata, Schrad. ex Steud. Nom. ed. I. 42 = officinalis.
intermedia, Schur, Enum. Pl. Transs. 466 = ochroleuca.
italica, *Retz. Obs.* i. 12.—Reg. Mediterr.
Kunthii, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 372 = Antiphytum Walpersii.
lanata, Linn. Syst. ed. X. 914 = Cynoglossum cheirifolium.
lanceolata, Willd. ex Steud. Nom. ed. II. i. 85 = Krynitzkia linifolia.
lateriflora, *Dum. Fl. Belg. Prod.* 41.—Belgia.
latifolia, Lahm. Sem. Hort. Hamb. (1821) 3 = Nonea rosea.
laxiflora, DC. Fl. Fr. iii. 681 = Borago laxiflora.
leiosperma, Bory, Fl. Pélop. t. 7. f. 3 = Nonea obtusifolia.
leptophylla, Koch, Syn. Fl. Germ. ed. II. ii. 573 = officinalis.
leptophylla, Roem. & Schult. Syst. iv. 90 = ochroleuca.
leucantha, Willd. ex DC. Prod. x. 122 = Antiphytum linifolium.
leucifolia, Lehm. Asperif. 249 = Echinospermum barbatum.
leucopsiola, Tardent, Ess. Hist. Nat. Besser. 57 = Nonea pulla.
limbata, *Boiss. Diagn.* Ser. I. xi. 99.—As. Min.
linearifolia, Hochst. in Lorent, Wanderungen, 384 = ochroleuca.
linearifolia, Urville, in Mém. Soc. Linn. Par. i. (1822) 276 = ochroleuca.
linifolia, Lehm. Asperif. 215 = Krynitzkia linifolia.
littorea, *Moris*, *in Atti Ottava Riun. Ital. Scienzi*, 866.—Sardin.
longifolia, *Lam. Encyc.* i. 504.—Italia.
Lorentii, *A. DC. in DC. Prod.* x. 58 = ochroleuca.
lucida, Vitm. Summa Pl. i. 257 = italica.
Lumnitzeri, *Wettstein*, *ex Stapf*, *in Denkschr. Acad. Wien*, i. (1885) 66.—Lycia.
lutea, Bieb. Fl. Taur. Cauc. i. 126 = Nonea lutea.
lutea, Cav. Ic. i. 90 = Myosotis versicolor.
Lycopsoides, Benth. ex Link, Enum. Hort. Berol. i. 166 = officinalis.

ANCHUSA :—
Lycopsoides, Retz. f. Pugill. 8 = Nonea lutea.
macrocarpa, *Boiss. & Hohen. Diagn.* Ser. I. iv. 42.—Kurdist.
macrophylla, *Desf. Fl. Atlant.* i. 157.—Marocco.
maculata, Hornem. ex Steud. Nom. ed. II. i. 85 = officinalis.
micrantha, Roem. & Schult. Syst. iv. 98 = aggregata.
microcalyx, *Vis. in Flora*, xii. (1829) 8.—Dalmat.
Milleri, *Willd. Enum. Hort. Berol.* i. 179.—Aegypt. ; Arab.
multifoides, Benth. in Royle, Illustr. Pl. Himal. t. 78. f. 1 = Mertensia multifoides.
myosotidiflora, *Lehm. Asperif.* 284.—Sibir. ; Caucas. neglecta, *A. DC. in DC. Prod.* x. 49.—Oriens.
nigricans, Brot. Fl. Lusit. i. 398 = Nonea nigricans.
nonnoides, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 85 = undulata.
oblique, *Vis. Ind. Sem. Hort. Patav.* (1836).—As. Min.
ochroleuca, Baumg. Fl. Transs. i. 120 = officinalis.
ochroleuca, *Bieb. Fl. Taur. Cauc.* i. 125.—Transylv. ; Oriens.
officinalis, Bieb. l. c. 124 = ochroleuca.
officinalis, Gouan, Hort. Monsp. 81 ; Lam. Encyc. i. 502 = italica.
officinalis, *Linn. Sp. Pl.* 133.—Europ.
officinalis, Thunb. Prod. 84 ; Fl. Cap. ed. Schult. 162 = riparia.
opposititolia, H. B. & K. Nov. Gen. et Sp. iii. 91 = Krynitzkia linifolia.
orientalis, Linn Sp. Pl. 133 = Alkanna orientalis.
orientalis, Retzius, f. Ic. Fl. Germ. xviii. 63 t. 1810 = Lycopsis orientalis.
orientalis, Pall. Reise, iii. 694 = Nonea lutea.
ovata, Lehm. Asperif. 222 = Lycopsis orientalis.
paniculata, Ait. Hort. Kew. i. 177 = italica.
parnassica, Boiss. & Orph. Diagn. Ser. II. iii. 134 = officinalis.
parviflora, Sibth. & Sm. Fl. Graec. ii. 57. t. 167 = aggregata.
parviflora, Willd. Sp. Pl. i. 759. (Sp. delenda ; cf. Boiss. Fl. Or. iii. 102).—As. Min.
pauciflora, Roem. & Schult. Syst. iv. 774 = Eritrichium rupestre.
perlata, Lam. Encyc. i. 504 ; Illustr. n. 1820 = Lycopsis variegata.
petiolata, Hook. Bot. Mag. t. 3858 = Cynoglossum petiolatum.
picta, Bieb. Fl. Taur. Cauc. i. 127 = Nonea picta.
Pinardi, Boiss. Fl. Or. Nov. Dec. ii. 8 = hybrida.
procera, Bess. ex Link, Enum. Hort. Berol. i. 166 = ochroleuca.
pubescens, *Kit. in Rochel, Pl. Bann.* 11.—Bannat.
pulla, Bieb. Fl. Taur. Cauc. i. 125 = Nonea pulla.
pustulata, *Schur, Enum. Pl. Transs.* 466.—Transylv.
purpurea, H. B. & K. Nov. Gen. et Sp. iii. 92 = Krynitzkia linifolia.
racemosa, DC. Prod. x. 50 = Mertensia racemosa.
rhizantha, Viv. Dec. n. 20 = Alkanna tinctoria.
riparia, *DC. Prod.* x. 48.—Afr. austr.
rosea, Bieb. Fl. Taur. Cauc. i. 125 = Nonea rosea.
rupestris, Roem. & Schult. Syst. iv. 101 = Eritrichium rupestre.
ryssoxperma, Gand. ex A. Rich. Tent. Fl. Abyss. ii. 88 = Milleri.
sabulosa, Stev. ex Ledeb. Fl. Ross. iii. 118 = ochroleuca.
saxatilis, Pall. Reise, iii. 718 (= *Stenosolenium saxatile*).
Sellowiana, *Cham. in Linnaea*, viii. (1833) 115.—Bras.
sempervirens, *Linn. Sp. Pl.* 134.—Europ.
serrata, Roem. & Schult. Syst. iv. 774 = Eritrichium sericeum.
silkimensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 186.—Reg. Himal.
spathulata, R. Br. ex Roem. & Schult. Syst. iv. 100 = Myosotis spathulata.
spicata, Lam. Illustr. n. 1810 = angustifolia.
spinacarpos, Forsk. Fl. Aegypt. Arab. 40 = Echinospermum spinocarpum.
Sprunerі, Boiss. Diagn. Ser. I. xi. 98 = stylosa.
stenchelifolia, Cham. in Linnaea, iv. (1829) 439 = Antiphytum stenchaefolium.
stricta, *Herbich*, *in Flora*, xxxvii. (1855) 641.—Europ.
strigosa, *Labill. Syr. Dec.* iii. 7. t. 4.—Syria ; Persia.

ANCHUSA :—
strigosa, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 246.—Syria.
stylosa, *Bieb. Fl. Taur. Cauc.* i. 128.—Graecia ; As. Min. ; Tauria.
subrepanda, *Kit. in Linnaea*, xxxii. (1863) 457.—Europ.
tenella, Hornem. Hort. Hafn. i. 176 = Bothriospermum tenellum.
tenuiflora, Fisch. ex Ledeb. Fl. Ross. iii. 142 = Eritrichium saxatile.
tetraquetra, Cham. in Linnaea, viii. (1833) 113 = Antiphytum tetraquetrum.
thessala, *Boiss. & Sprun. Diagn.* Ser. I. xi. 99.—Thessal.
tinctoria, Linn Sp. Pl. ed. II. i. 192 = Alkanna tinctoria.
tinctoria, Pall. Ind. Taur. ex Lehm. Asperif. 266 = Nonea pulla.
tinctorea, Sieber, ex Steud Nom. ed. II. i. 86 = Lycopsis orientalis.
tinctoria, Woodv. Med. Bot. t. 92 = officinalis.
tomentosa, Boiss. Diagn. Ser. II. iii. 133 = ochroleuca.
tuberculata, Forsk. Fl. Aegypt. Arab. 40 = Alkanna tinctoria.
tuberosa, *H. B. & K. Nov. Gen. et Sp.* iii. 92.—Amer. austr.
undulata, *Linn. Sp. Pl.* 138.—Reg. Mediterr.
undulata, Pall. Reise, iii. 682 = ochroleuca.
undulata, Sibth. & Sm. Fl. Graec. ii. t. 168 = hybrida.
undulata, Vahl, ex DC. Prod. x. 49 = granatensis.
variegata, Lehm. Asperif. 228 = Lycopsis variegata.
ventricosa, Sibth. & Sm. Fl. Graec. ii. 58. t. 169 = Nonea alba.
ventricosa, Viv. Fl. Lib. Spec. 10. t. 6. f. 1 = Nonea Viviani.
verrucosa, Lam. Encyc. i. 504 ; Illustr. 403. n. 1819 = aegyptiaca.
verrucosa, Lejeun. ex Reichb. Fl. Germ. Excurs. 342 = Lycopsis variegata.
versicolor, Schur, Enum. Pl. Transs. 466 = ochroleuca.
versicolor, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 254 = Nonea rosea.
villosa, Roem. & Schult. Syst. iv. 775 = Eritrichium villosum.
villosa, *Spreng. Syst.* i. 560.—Hab. ?
virginiana, *Linn. Sp. Pl.* 138 (= Lithospermum hirtum ?).—Amer. bor.
virginica, Linn. Syst. ed. XII. 145 = præc.
vulgaris, Dum. Fl. Belg. 43 = sempervirens.
zeylanica, Jacq. f. Eclog. i. 47. t. 29 = Bothriospermum tenellum.
zeylanica, Vahl, ex Hornem. *Hort. Hafn.* i. 176 = Cynoglossum denticulatum.

ANCHUSOPSIS, Bisch. Del. Sem. Hort. Heidelb. (1852) 8 = **Lindelofia**, Lehm. (Borag.).
longiflora, Bisch. l. c. = L. spectabilis.

ANCISTROCARPHUS, A. Gray, in Proc. Am. Acad. vii. (1868) 355 = **Micropus**, Linn. (Compos.).
filagineus, A. Gray, l. c. 356.—Calif.

ANCISTROCARPUS, H. B. & K. Nov. Gen. et Sp. ii. 186. t. 122. (1817) = **Microtea**, Sw. (Phytolacc.).
hexander, J. Gay, ex Meq. in DC. Prod. xiii. ii. 17, 18 = M. mayuranensis.
mayuranensis, H. B. & K. Nov. Gen. et Sp. ii. 186 = M. mayuranensis.
mayuranensis, Spreng. Neue Entdeck. i. 257 = M. Spreng eldii.
Schrankii, Ledeb. Ind. Sem. Hort. Dorp. (1822) = M. mayuranensis.

ANCISTROCARPUS, Oliver, in Journ. Linn. Soc. ix. (1867) 173. *TILIACEAE*, Benth. & Hook. f. i. 986.
hermiphonus, *Oliver*, *l. c.* 174.—Afr. trop.
dentiphonus, *Oliver*, *l. c.*—Afr. trop.

ANCISTROCARYA, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 448. *BORAGINEAE*, Benth. & Hook. f. ii. 861.
japonica, *Maxim. l. c.* 444.—Japon.

ANCISTROCLADUS, Wall. Cat. n. 1059 (1832). *DIPTEROCARPEAE*, Benth. & Hook. f. i. 191.
BIGAMEA Koenig, ex Endl. Gen. 1188 (1840).

ANCISTROCLADUS:—
WORMIA, Vahl, in Skrift. Nat. Selsk. Kjobenh. vi. (1810) 104.
attenuatus, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 300.—Burma.
extensus, *Wall. Cat.* n. 1052.—Burma.
Griffithii, *Planch. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 316.—Burma.
guineensis, *Oliver, Fl. Trop. Afr.* i. 175.—Afr. trop.
Heyneanus, *Wall. Cat.* n. 7262.—Ind. or.
pinangianus, *Wall.* l. c. n. 1054.—Malaya.
sagittatus, Wall. l. c. n. 1055 = Tetramerista glabra.
stelligerus, Wall. l. c. n. 1058 = Wallichii.
Vahlii, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 325. —Zeylan.
Wallichii, *Planch. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 319.—Ind. or.

ANCISTRODESMUS, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 309 = **Microlepis**, Miq. (Melast.).
oleaefolius, Naud. l. c. 308 = M. oleaefolia.

ANCISTROLOBUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 302. t. 6 = **Cratoxylon**, Blume (Hyper.).
brevipes, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 333 = C. polyanthum.
floribundus, Turcz. l. c. 332 = C. floribundum.
formosus, Zoll. & Mor. Syst. Verz. Zoll. 25 = C. formosum.
glaucescens, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 333 = C. arborescens.
lignitrunus, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 352 = C. polyanthum.
micradenius, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 332 = C. Blancoi.

ANCISTROPHORA, A. Gray, in Mem. Am. Acad. Ser. II. vi. (1859) 457 = **Verbesina**, Linn. (Compos.).
Wrightii, A. Gray, l. c. = V. Wrightii.

ANCISTROPHYLLUM, G. Mann & H. Wendl. in Kerch. Palm. 230 (1878). *PALMAE*, Benth. & Hook. f. iii. 937.
LACCOSPERMA, G. Mann & H. Wendl. l. c. 249 (1878 \, secundiflorum, *G. Mann & H. Wendl. l. c.*—Afr. trop.

ANCISTROSTIGMA, Fenzl, in Ann. Wien. Mus. ii. (1838) 398; Nov. Stirp. Dec. 85 (1838) = **Trianthema**, Linn. (Ficoid.).
cypsoleoides, Fenzl, l. c. = T. cypseleoides.

ANCISTRUM, Forst. Char. Gen. 3. t. 2 (1776) = **Acaena**, Linn. (Rosac.).
Acaena, Lag. Gen. et Sp. Nov. 7 = Acaena serices.
acerosa, DC. Prod. ii. 592, sphalm. = praec.
adscendens, Poir. Encyc. Suppl. i. 847 = Acaena adscendens.
anserinaefolium, Forst. Char. Gen. 4 = Acaena Sanguisorbae.
argentiosum, Poir. Encyc. Suppl. i. 846 = Acaena argentea.
barbatum, Lam. Illustr. i. 77. t. 22. f. 3 = Margyricarpus setosus.
cylindristachyum, Poir. Encyc. Suppl. i. 847 = Acaena cylindristachya.
decumbens, Gaertn. Fruct. i. 163. t. 32 = Acaena Sanguisorbae.
decumbens, Thunb. Prod. Pl. Cap. 6 = Acaena latebrosa.
dianthrum, Forst. f. Prod. 10 = Acaena Sanguisorbae.
elongatum, Poir. Encyc. Suppl. i. 845 = Acaena elongata.
humile, Pers. Syn. i. 80 = Acaena adscendens.
laevigatum, Lag. Gen. et Sp. Nov. 7 = Acaena laevigata.
lappaceum, Poir. Encyc. Suppl. i. 844 = Acaena lappacea.
latebrosum, Ait. Hort. Kew. i. 16 = Acaena latebrosa.
lucidum, Ait. l. c. 15 = Acaena lucida.
magellanicum, Lam. Illustr. i. 76. t. 22. f. 2 = Acaena adscendens, laevigata.
ovalifolium, Sweet, Hort. Brit. ed. I. 148 = Acaena ovalifolia.
pinnatifidum, Poir. Encyc. Suppl. i. 845 = Acaena pinnatifida.
pumilum, Poir. l. c. 847 = Acaena pumila.
sepium, Vent. Jard. Cels. 6. t. 6 = Acaena ovalifolia.
Sanguisorbae, Linn. f. Suppl. 89 = Acaena Sanguisorbae.

ANCISTRUM:—
sarmentosum, Thon. Fl. Tristan d'Acugna, 44 = Acaena Sanguisorbae.
trifidum, Poir. Encyc. Suppl. i. 846 = Acaena trifida.

ANCLYANTHUS, A. Juss. in Orb. Dict. xi. 254 (1849) = **Ancylanthos**, Desf.

ANCOBUMIA, Harv. ex Baker, in Hook. Ic. Pl. t. 1927 (1877). *LILIACEAE*, Benth. & Hook. f. iii. 806.
cuspidata, *Haro. ex Baker, l. c.*—Chili.

ANCYLANTHOS, Desf. in Mém. Mus. Par. iv. (1818) 5. t. 2. *RUBIACEAE*, Benth. & Hook. f. ii. 119.
Balansii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 160.—Afr. trop.
cinerascens, *Welw. ex Hiern, l. c.* 159.—Afr. trop.
cistifolius, *Welw. ex Hiern, l. c.*—Afr. trop.
fulgidus, *Welw. ex Hiern, l. c.*—Afr. trop.
Monteiroi, *Oliver, in Hook. Ic. Pl.* t. 1208.—Afr. trop.
rubiginosus, *Desf. l. c.*—Angola.

ANCYLOCALYX, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 186. t. 2 = **Pterocarpus**, Linn. (Legumin.).
acuminata, Tul. l. c. 187 = P. Ancylocalyx.

ANCYLOCLADUS, Wall. Pl. As. Rar. iii. 45 (1832) = **Willughbeia**, Roxb. (Apocyn.).

ANCYLOGYNE, Nees, in Mart. Fl. Bras. ix. 63. t. 7 (1847) = **Sanchesia**, Ruiz & Pav. (Acanth.).
capitata, Nees, in DC. Prod. xi. 222.—Peruv.
longiflora, Hook. f. Bot. Mag. t. 5588.—Ecuador.
macracnemis, Nees, in Mart Fl. Bras. ix. 64.—Am. trop.
munita, Nees, l. c.—Am. trop.
peruviana, Nees, in DC. Prod. xi. 222.—Peruv.

ANCYROSSEMON, Poepp. & Endl. Nov. Gen. et Sp. iii. t. 272 (1845) = seq.

ANCYROSTEMMA, Poepp. & Endl. Nov. Gen. et Sp. iii. 65 (1845) = **Selerochrix**, Presl (Lossac.).
micranthum, Poepp. & Endl. l. c. = S. fasciculata.

ANDA, A. Juss. [In Dict. Sc. Nat. ii. 218 (1816)]; Euph. Tent. 89. t. 12 (1824) = **Joannesia**, Vell. (Euphorb.).
brasiliensis, Raddi, in Mem. Mod., xviii.; Fis. (1820) 404 = J. Princeps.
Gomezii, A. Juss. Euph. Tent. 89. t. 12 = J. Princeps.
Yomezii, Mart. in Regensb. Arch. Gew. iv. (1830) 85 = praec.
Pisonis, Mart. ex Steud. Nom. ed. II. i. 86 = J. Princeps.

ANDACA, Rafin. Fl. Tellur. iii. 82 (1836) = **Lotus**, Tourn. (Legumin.).
arabica, Rafin. l. c. = L. arabicus.

ANDERSONIA, R. Br. Prod. 558 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 617.
ANTHOCERALA, DC. Prod. vii. 755 (1839).
HOMALOSTOMA, Stschegl. in Bull. Soc. Nat. Mosc. (1859) 1. 32.
SPHINCTEROSTOMA, Stschegl. l. c. 31.
aristata, *Lindl. Swan Riv. App.* 25.—Austral.
brachyantheta, *F. Muell. Fragm.* vi. 61.—Austral.
brachyotis, F. Muell. l. c. iv. 126 = parviflola.
brevifolia, *Sond. in Lehm. Pl. Preiss.* i. 332.—Austral.
coerulea, *R. Br. Prod.* 554.—Austral.
colonna, *F. Muell. Fragm.* vi. 62.—Austral.
depressa, *R. Br. l. c.* 555.—Austral.
depressa, A. Cunn. ex DC. Prod. vii. 767 = squarrosa.
Preissii, Sond. in Lehm. Pl. Preiss. i. 331 = sprengelioides.
gracilis, *DC. Prod.* vii. 767.—Austral.
grandiflora, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) 1. 31.—Austral.
heterophylla, *Sond. in Lehm. Pl. Preiss.* i. 333.—Austral.
homatostoma, *Benth. Fl. Austral.* iv. 366.—Austral.
involucrata, *Sond. in Lehm. Pl. Preiss.* i. 331.—Austral.

ANDERSONIA:—
lanuginosa, A. Cunn. ex DC. Prod. vii. 767 = depressa.
latiflora, *F. Muell. Fragm.* vi. 61.—Austral.
Lehmanniana, Sond. in Lehm. Pl. Preiss. i. 330 = sprengelioides.
macranthera, F. Muell. Fragm. iv. 124 = heterophylla.
macronema, F. Muell. l. c. viii. 51 = Sprengelia macronema.
micrantha, *R. Br. Prod.* 558.—Austral.
parviflola, *R. Br. l. c.*—Austral.
patens, Sond. in Lehm. Pl. Preiss. i. 331 = sprengelioides.
patrica, *F. Muell. Fragm.* vi. 79.—Austral.
pauciflora, *Sond. in Lehm. Pl. Preiss.* i. 332.—Austral.
prostrata, Sond. l. c. 333 = depressa.
pubescens, Sond. l. c. 331 = sprengelioides.
setifolia, *Benth. Fl. Austral.* iv. 252.—Austral.
spiraphylla, F. Muell. Fragm. vi. 62 = grandiflora.
sprengelioides, *R. Br. Prod.* 558.—Austral.
squarrosa, R. Br. Prod. 558 = depressa.
subsessilis, *Pancher. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 67.—N. Caled.
subulata, *Benth. Fl. Austral.* iv. 256.—Austral.
variegata, *Sond. in Lehm. Pl. Preiss.* i. 334.—Austral.

ANDERSONIA, Buch.-Ham. ex Wall. Cat. sub n. 4014 (1832) = **Anogeissus**, Wall (Combret.).
acuminata, Roxb. ex Wight & Arn. Prod. 316 = Anog. acuminata.
altissima, Roxb. ex Wight & Arn. l. c. = Anog. latifolia.
lanceolata, Buch.-Ham. ex Wall. l. c. = Anog. acuminata.

ANDERSONIA, Roxb. Hort. Beng. [87] (1814); Fl. Ind. ii. 212 (1832) = **Amoora**, Roxb. (Meliac.).
cucullata, Roxb. ll. cc. = Am. cucullata.
Rohituka, Roxb. ll. cc. 87, 213 = Am. Rohituka.

ANDERSONIA, Willd. ex Roem. & Schult. Syst. v. 21 (1819) = **Gaertnera**, Lam. (Logan.).
vaginata, Willd. ex Roem. & Schult. l. c. = G. vaginata.

ANDICUS, Vell. Fl. Flum. 80 (1825) = **Joannesia**, Vell. (Euphorb.).
pentaphyllus, Vell. l. c. = J. Princeps.

ANDIRA, Lam. Encyc. i. 171 (1788). *LEGUMINOSAE*, Benth. & Hook. f. i. 550.
LUMBRICIDIA, Vell. Fl. Flum. 205 (1825).
POLTOLORIUM, Presl, Bot. Bemerk. 68 (1844).
VOUACAPOUA, Aubl. Pl. Gui. Suppl. 2. t. 373 (1775).
acuminata, Benth. in Ann. Wien. Mus. ii. (1838) 109 = inermis.
Amazonum, *Mart. ex Benth. l. c.* 107.—Bras.
anthelmintica, *Benth. l. c.* 108.—Bras.
Araroba, *Aguiar, Mem. sobre a Araroba*, 31.—*Cf. Pharm. Journ.* Ser. III. x. (1879–80) 44.—Bras.
Aubletii, *Benth. in Ann. Wien. Mus.* ii. (1838) 108.—Guiana.
bracteosa, Benth. l. c. 107 = Amazonum.
cinerascens, Mart. Herb. Fl. Bras. n. 1156 = Tipuana macrocarpa.
cubensis, *Benth. in Journ. Linn. Soc.* iv. (1860) *Suppl.* 121.—Cuba.
cubensis, *Benth. in Mart. Fl. Bras.* xv. i. 295.—Bras.
excelsa, *H. B. & K. Nov. Gen. et Sp.* vi. 385.—Mexic.
fraxinifolia, *Benth. in Ann. Wien. Mus.* ii. (1838) 108. —Bras.
frondosa, *Mart. in Walp. Rep.* i. 803.—Bras.
gebenica, *Baill. in Adansonia*, vi. (1866) 219 (adnot.). —Afr. trop.
grandiflora, Guill. & Perr. Tent. Fl. Seneg. i. 264 = inermis.
Hostfeldii, Leschen. in Ann. Mus. Par. xvi. (1810) t. 24 = sp.
Horsfieldii, Leschen. l. c. 451 (err. Horsfeldii) = Eschreta Horsfieldii.
humilis, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 109.—Bras.
inermis, *H. B. & K. Nov. Gen. et Sp.* vi. 385.—Ind. occ. ; Bras.
laurifolia, *Benth. in Ann. Wien. Mus.* ii. (1838) 109. —Bras.

ANDIRA :—
microcarpa, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 179.—Cuba.
nitida, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 109.—Bras.
obtusa, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 121 = retusa.
ormosioides, Benth. in Ann. Wien. Mus. ii. (1838) 108 = anthelmintnica.
paniculata, *Benth. l. c.* 109.—Bras.
parrifolia, *Mart. ex Benth. l. c.* 108.—Bras.
pauciflora, *Benth. l. c.* 109.—Bras.
Pisonis, Auct. ex Steud. Nom. ed. II. i. 87 = A. Aubletii.
Pisonis, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 108.—Bras.
racemosa, Lam. Encyc. i. 17 = excelsa.
retusa, *H. B. & K. Nov. Gen. et Sp.* vi. 385.—Guiana, riparia, *H. B. & K. l. c.*—N. Granat.
rosea, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 105.—Bras.
sapindoides, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 123.—Ind. occ.
spinulosa, *Mart ex Benth. in Ann. Wien. Mus* ii. (1838) 108.—Bras.
stipulacea, *Benth. l. c.* 107.—Bra
vermifuga, *Mart. ex Benth. l. c.* 108.—Bras.

ANDISCUS, Vell. Fl. Flum. ii. t. 86 (1827) = **Andious**, Vell.

ANDOUINIA, Reichb. Consp. 160 (1828) err. typ. = **Audouinia**, Brongn. (Bruniac.).

ANDRACHNE, Linn. Syst. ed. I (1735) *EUPHORBIACEAE*, Benth. & Hook. f. iii. 270.
ARACHNE, Neck. Elem. ii. 848 (1790).
ERACLISSA, Forsk. Fl. Aegypt. Arab. 208 (1775)
LEPIDANTHUS, Nutt. in Trans. Am. Phil. Soc v. (1841) 175.
LEPTOPUS, Decne. in Jacquem. Voy. Bot 155. t. 156 (1844).
MASCHALANTHUS, Nutt. in Trans. Am. Phil. Soc v. (1837) 175.
PHYLLANTHIDIA, Didrichs. in Vidensk. Meddel. Kjoeb. (1857) 150.
TELEPHIOIDES, Tourn. ex Moench, Meth. Suppl. 310 (1802).
apetala, Roxb. ex Wall. Cat. n. 7956 = Bischofia javanica.
arboren, *Mill. Gard. Dict.* ed. VIII. *n* 3.— Campech aspera, *Spreng. Syst.* iii. 884.—Orient; Afr. trop.; Ind. or.
australis, *Zoll. & Mor. in Natur. en Genees Arch. Neerl. Indie*, ii. (1845) 17.—Java.
Cadiaca, Roxb. ex Wall. Cat. n. 7877 = Cleistanthus collinus.
capensis, *Baill. in Adansonia*, iii. (1862–63) 168.—Afr. austr.
chinensis, *Bunge, Enum. Pl. Chin. Bor.* 59.—China.
colchica, *Fisch. & Mey. ex Boiss. Fl. Orient.* iv. 1137. —Caucas.
cordifolia, *Muell. Arg. in DC. Prod.* xv. ii. 334.—Reg. Himal.
Decaisneana, Baill. Etud. Gén. Euphorb. 577 = cordifolia.
Decaisnei, *Benth. Fl. Austral.* vi. 88.—Austral.
Doonkyboica, Heyne, ex Wall. Cat. n. 7879 = Bridelia retusa.
Dregeana, *Baill. Adansonia*, iii. (1862–63) 164.—Afr. austr.
elliptica, *Roth, Nov. Sp.* 264.—Ind. or.
frutescens, Ehret, in Phil. Trans. lvii. (1768) 114. t. 6 = Arbutus Andrachne.
fruticosa, Decne. ex Muell. Arg. in DC. Prod. xv. ii. 235 = Decaisnei.
fruticosa, *Linn. Sp. Pl.* 1014.—China; N. Hisp.
fruticosa, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 287 = Phyllanthus subcrosus.
fruticulosa, *Botan. Disqu.* Ser. I. vii. 86.—Persia.
maroccana, *Ball, in Journ. Bot.* xiii. (1875) 205.—Marocco.
microphylla, *Baill. Etud. Gén. Euph.* 577.—Peruv.
obtusifolia, Muell. Arg. in DC. Prod. xv. ii. 510, in syn. = sq.
orbiculata, Roth, Nov. Sp. 264 = Cleistanthus collinus.
ovalis, *Muell. Arg. in Linnaea*, xxxii. (1863) 78.—Afr. austr.
ovata, Lam. ex Poir. Encyc. Suppl. i. 248, in syn. = Agyneia ovata.
phyllanthoides, Muell. Arg. in DC. Prod. xv. ii. 435, in syn. = Roemeriana.
Roemeriana, *Muell. Arg. l. c.* 234.—Amer. bor.
rotundifolia, Elchw. ex C. A. Mey. Casp. 18. t. 20 = telephioides.
telephioides, *Linn. Sp. Pl.* 1014.—Reg. Mediterr.; Orient; Ind. or
tenera, *Msq. Fl. Ind. Bat.* i. ii. 365 —Malaya.
trifoliata, Roxb. Hort. Beng. 70; Fl. Ind. iii. 728 = Bischofia javanica.

ANDRADEA, Allemão, Diss. Andrad. (1845). *NYCTAGINEAE*, Benth. & Hook. f. iii. 11.
floribunda, *Allemão, l. c.*—Bras.

ANDRASTIS, Rafin. ex Benth. in Ann. Wien. Mus. ii. (1838) 86 = **Virgilia**, Lam. (Legum.).

ANDREOSKIA, Boiss. Fl. Orient. i. 803 (1867) = **Andrzeiowskia**, Reichb. (Crucif.).

ANDREOSKIA, DC. Prod. i. 190 (1824) = **Dontostemon**, Andrz. (Crucif.).
crassifolia, Bunge, ex Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 271.—Mongolia.
dentata, Bunge, Enum. Pl. Chin. Bor. 6 = D. dentatus.
eglandulosa, DC. Prod i. 190 = D. eglandulosus.
integrifolia, DC. l. c. = D. integrifolius.
parvifora, Turcz. Cat. Pl. Baic. n. 155. sec. Bull. Soc. Nat. Mosc. xv. (1842) 289 = Malcolmia Ledebourii.
pectinata, DC. Prod. i. 190 = D. pectinatus.

ANDREUSIA, Dun. in DC. Prod. vii. 560 (1839) = **Hornemannia**, Vahl (Vaccin.).
guadaiupensis, Dun. i. c. = H. martinicensis.

ANDREUSIA, Vent. Jard. Malm. 108. t. 108 (1807) = **Myoporum**, Banks
debilis, Vent. l. c. sub t. 108 = M. debile.
glabra, Vent. l. c. t. 108 = M. ellipticum.
salicifolia, Hort. ex Steud. Nom. ed. II. i. 87 = Freylinia cestroides.
scabra, Zeyh. ex Steud. l. c. = M. parviflorum.

ANDREWSIA, Spreng. Anleit. ii. i. 474 (1817); Syst. i. 428 (1825) = **Bartonia**, Muhl. (Gentian.).
aulumnalis, Spreng. Syst. i. 162 = B. tenella.
paniculata, Bart. ex Steud. Nom. ed. II. i. 87 = B. tenella.
verna, Spreng. Syst. i. 162 = B. verna.

ANDRIALA, Dence. in Orb. Dict. i. 485 (1841) err. typ. = **Andryala**, Linn. (Compos.).

ANDRIAPETALUM, Pohl, Pl. Bras. Ic. i. 113 (1827) = **Tanopsis**, Salisb. (Proteac.).
cayennense, Klotzsch, ex Meisn. in Mart. Fl. Bras. v. i. 78.—Bras.
ferrugineum, Meisn. in DC. Prod. xiv. 845.—Peruv.
muelleriana, Schott, ex Endl. Suppl. iv. ii. 82.—Bras.
polysiachya, Schott, ex Meisn. in DC. Prod. xiv. 345. —N. Granat.
reticulatum, Pohl, Pl. Bras. i. 115. t. 92.—Bras.
rubescens, Benth. ex Meisn. in DC. Prod. xiv. 347 = *Spruci*.
rubescens, Pohl, Fl. Bras. i. 114. t. 91.—Bras.
sessilifolium, Benth. ex Meisn. in DC. Prod. xiv. 346 = cayennense.
sessilifolium, Klotzsch, in Linnaea, xv. (1841) 53 = P. hamelioefolia.
Spruci, Meisn. in DC. Prod. xiv. 347.—Bras.
suaveolens, Klotzsch, in Linnaea, xx. (1847) 472.—N. Granat.
Yolombo, Posado, in Bull. Soc. Bot. Fr. xviii. (1871) 374.—N. Granat.

ANDRIEUXIA, DC. Prod. v. 559 (1836) = **Heliopsis**, Pers. (Compos.).
mexicana, DC. l. c. et in Deless. Ic. Sel. iv. t. 31 = H. buphthalmoides.
quitensis, Benth. Pl. Hartw. 306.—Ecuador.

ANDROCENTRUM, Lem. in Fl. des Serres, Sér. I. iii. (1847) 249, verso. *ACANTHACEAE*, Benth. & Hook. f. ii. 1084.
multiflorum, *Lem. l. c.*—Mexico.

ANDROCERA, Nutt. Gen. Am. i. 129 (1818) = **Solanum**, Linn.
lobata, Nutt. l. c. = S. rostratum.

ANDROCHILUS, Liebm. in Bot. Notiser, (1844) 101 = **Liparis**, Rich. (Orchid.).
campestris, Liebm. l. c. 197.—Mexico.

ANDROCOMA, Nees, in Hook. Journ. Bot. ii. (1840) 396 = **Bolrpus**, Linn. (Cyperac.).
speciosa, Nees, l. c. 397 = S. giganteus.

ANDROCYMBIUM, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 21. t. 2. *LILIACEAE*, Benth. & Hook. f. iii. 822.
CYMSANTHES, Salisb. in Trans. Hort. Soc. i. (1812) 329.
ERYTHROSTICTUS, Schlecht. in Linnaea, i. (1826) 90.
PLEXINIUM, Rafin. Fl. Tellur. ii. 32 (1836).
Burchellu, *Baker, in Journ. Bot.* xii. (1874) 246.—Afr. austr.
Burkei, *Baker, l. c.*—Afr. austr.
circinatum, *Baker, in Journ. Linn. Soc.* xvii. (1880) 448.—Afr. austr.
cucullatum, *Eckl. Verz.* 7.—Afr. austr.
cuspidatum, *Baker, in Journ. Bot.* xii. (1874) 245.—Afr. austr.
Dregei, *Presl, Bot. Bemerk.* 116.—Afr. austr.
eucomodes, Sweet, Brit. Flow. Gard. t. 165 = leucanthum.
encomodes, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 21.—Afr austr.
leucanthum, *Willd. l. c.*—Afr. austr.
littorale, *Eckl. Verz.* 7.—Afr. austr.
longipes, *Baker, in Journ. Bot.* xii. (1874) 246.—Afr. austr.
melanthioides, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 21.—Afr. austr.
palaestinum, *Baker, in Journ. Linn. Soc.* xvii. (1880) 445.—Palaestin.
punctatum, Baker, in Journ. Bot. xii. (1874) 246 = leucanthum.
punctatum, *Baker, in Journ. Linn. Soc.* xvii. (1880) 446.—Afr. bor.
striatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 336. —Abyss.
suboiatum, *Baker, in Journ. Bot.* xii. (1874) 245.—Afr. austr.
volutare, *Burch. Trav.* i. 218.—Afr. austr.
volutum, Burch. ex Baker, in Journ. Bot. xii. (1874) 246 = volutare.

ANDROGLOSSA, Benth. in Hook. Kew Journ. iv. (1852) 42 = **Sabia**, Colebr.
reticulatum, Benth. l. c. = S. limoniacea.

ANDROGRAPHIS, Wall. Pl. As. Rar. iii. 77 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1099.
ERIANTHERA, Nees, in Wall. Pl. As. Rar. iii. 77 (1832).
affinis, *Nees, l. c.* iii. 116.—Ind. or.
alata, *Nees, in DC. Prod.* xi. 516.—Ind. or.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 506.—Ind. or.
ceylanica, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 313 = corchorotyra.
ceylanica, Wight, Ic. t. 1560 = viscosula.
echioides, *Nees, in Wall. Pl. As. Rar.* iii. 117.—Ind.or.
elongata, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 302.—Ind. or.
glandulosa, *Nees, in Wall. Pl. As. Rar.* iii. 115.—Ind. or.
gracilis, Nees, in DC. Prod. xi. 516 = Wightiana.
humifusa, Wall. ex Nees, l. c. 514 = serpyllifolia.
lineata, *Nees, in Wall. Pl. As. Rar.* iii. 116.—Ind. or.
lobelioides, *Wight, Ic.* t. 1557.—Ind. or.
macrobotrys, *Nees, in DC. Prod.* xi. 516.—Ind. or.
Neesiana, Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 504 = stellulata.
Neesiana, *Wight, Ic.* t. 1561.—Ind. or.
orbiculata, Wall. ex Nees, in DC. Prod. xi. 514 = serpyllifolia.
ovata, *Benth. & Hook. f. Gen.* ii. 1100.—Ind. or.
paniculata, *Nees, in Wall. Pl. As. Rar.* iii. 116.—Ind. or.
Rothii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 506. —Ind. or.
serpyllifolia, *Wight, Ic.* t. 1557.—Ind. or.

ANDROGRAPHIS:—
stellulata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 506.—Ind. or.
stenophylla, *C. B. Clarke, l. c.* 503.—Ind. or.
subpatula, *C. B. Clarke, l. c.* 502.—Ind. or.
tenuiflora, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 502.—Ind. or.
viscidula, Steud. Nom. ed. II. i. 87 = viscosula.
viscosula, *Nees, in Wall. Pl. As. Rar.* iii. 116.—Ind. or.
Wightiana, *Arn. ex Nees, in DC. Prod.* xi. 517.—Ind. or.
Wightiana, T. Anders. in Thw. Enum. Pl. Zeyl. 232 = macrobotrys.

ANDROGYNE, Griff. Notul. iii. 279 (1851) = **Panisea**, Lindl. (Orchid.).

ANDROLEPIS, Brongn. ex Hooll. in Rev. Hortic. xlii. (1870) 12 = **Aechmea**, Ruiz & Pav. (Bromel.).
Skinneri, Brongn. l. c.—Guatemala.

ANDROMACHIA, Humb. & Bonpl. Pl. Aequin. ii. 104. t. 112 (1809) = **Idabum**, Adans. (Compos.).
aculis, H. B. & K. Nov. Gen. et Sp. iv. 98 = L. acaule.
alternifolia, Kunze, ex Steud. Nom. ed. II. i. 87 = Aplopappus chrysanthemifolius.
Doppeana, Less. in Linnaea, vi. (1831) 401 = L. Deppeanum.
excelsa, Poepp. & Endl. Nov. Gen. et Sp. iii. 44.—Peruv.
grandiflora, H. B. & K. Nov. Gen. et Sp. iv. 99 = L. grandiflorum.
hieracioides, H. B. & K. l. c. 98 = L. hieracioides.
igniarum, Humb. & Bonpl. Pl. Eq. ii. 104. t. 112 = L. igniarium.
Jussieui, Cass. in Bull. Soc. Philom. (1817) 184 = L. Jussieui.
meiastomoides, H. B. & K. Nov. Gen. et Sp. iv. 101. t. 137 = L. melastomoides.
nubigenum, H. B. & K. l. c. 102 = L. nubigenum.
Poiteaui, Cass. in Bull. Soc. Philom. (1817) 184 = L. Brownei.
sessilifolia, H. B. & K. Nov. Gen. et Sp. iv. 102 = L. sessiliflorum.
solidaginea, H. B. & K. l. c. 100 = L. solidagineum.
verbascifolia, H. B. & K. l. c. = L. verbascifolium.
volubilis, H. B. & K. l. c. 102 = L. volubile.

ANDROMEDA, Linn. Syst. ed. I. (1735); Gen. ed. I. 128 (1737), pro parte. *ERICACEAE*, Benth. & Hook. f. ii. 567.
Xolisma, Raño. in Am. Monthly Mag. (1819) 198.
acuminata, Alt. Hort. Kew. ii. 70 = Leucothoë acuminata.
acuminata, Sm. Exot. Bot. ii. 59 = Leucothoë Catesbæi.
adenothrix, Miq. Ann. Mus. Bot. Lugd. Bat. i. 81 = Gaultheria adenothrix.
alata, Herb. Dombey, ex Dun. in DC. Prod. vii. 562 (= *Proclusia alata*).—Peruv.
ambigua, *Schrad. in Gott. Gel. Ana.* i. (1821) 710.—Hab.?
americana, Hort. ex DC. Prod. vii. 607 = A. Polifolia.
anastomosans, Murr., ex Linn. f. Suppl. 330 = Gaultheria anastomosans.
angustifolia, Pursh. Fl. Am. Sept. i. 291 = Cassandra calyculata.
arborea, Linn. Sp. Pl. ed. I. 394 = Oxydendrum arboreum.
axillaris, Lam. Encyc. i. 157 = Leucothoë axillaris.
axillaris, Michx. Fl. Bor. Am. i. 258 = Leucothoë Catesbæi.
baccata, Wangenh. Beitr. Am. 80. t. 89. f. 89 = Gaylussacia resinosa.
bracteata, H. B. & K. Nov. Gen. et Sp. iii. 258. t. 269 = Agarista bracteata.
bracteata, Cav. Ic. vi. 42 = Gaultheria bracteata, tomentosa.
bracteosa, DC. Prod. vii. 607 = Arctostaphylos tomentosa.
bryanthe, Linn. Mant. 228 = Bryanthus Gmelini.
buxifolia, Lam. Encyc. i. 159 = Agauria buxifolia.
carnulea, Linn. Sp. Pl. 393 = Phyllodoce taxifolia.
calyculata, Linn. l. c. 394 = Cassandra calyculata.
campanulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 81.—Japon.
canadensis, Cels, ex Lam. Encyc. i. 158 = A. Polifolia.

ANDROMEDA:—
candida, Vell. Fl. Flum. 175; iv. t. 98 = Gaultheria candida.
capricida, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 148 = Pieris ovalifolia.
cassinefolia, Vent. Jard. Cels, 60 = Zenobia speciosa.
Catesbæi, Walt. Fl. Carol. 137 = Leucothoë Catesbæi.
cerea, Linn. f. Suppl. 238 = Vaccinium cereum.
cernua, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 81.—Japon.
chinensis, Lodd. Bot. Cab. t. 1642 = Vaccinium bracteatum.
chlorantha, Cham. in Linnæa, viii. (1833) 508 = Leucothoë chlorantha.
ciliata, Nees, in Princ. Neuwied, Reise, i. 104 = Leucothoë ciliata.
ciliciaba, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 80 = Menziesia ciliciaba.
ciliosa, Boj. ex DC. Prod. vii. 608 = (*Leucothoë Bojeri*).—Madag.
coccinea, Nees, in Princ. Neuwied, Reise, i. 104 = Leucothoë coccinea.
coccinea, Schrad. in Goett. Gel. Anz. i. (1821) 709 = Gaylussacia Pseudo-vaccinium.
columellaris, Fisch. ex Herd. in Act. Hort. Petrop. i. (1871–72) 330 = Cassiope Redowskyi.
coriacea, Alt. Hort. Kew. ed. I. ii. 70 = Pieris nitida.
corisifolia, Thunb. & Blilh. Fl. Bras. Dec. 1. n. 1 = Leucothoë corisifolia.
crassifolia, Pohl, Pl. Bras. ii. 34 = Leucothoë crassifolia.
crespa, Hort. Paris. ex Poir. Encyc. Suppl. i. 856 = Cassandra calyculata.
cubensis, *Griseb. Cat. Pl. Cub.* 91.—Cuba.
cupressiformis, Wall. ex D. Don, in Mem. Wern. Soc. iii. (1823) 413 = Cassiope fastigata.
cupressina, Hook. Fl. Bor. Am. ii. 38 = Cassiope Mertensiana.
Dahoëcia, Linn. Syst. ed. XII. 800 = Dahoëcia polifolia.
Dahoecia, Pall. Fl. Ross. ii. 57 = Phyllodoce taxifolia.
dealbata, Lindl. Bot. Reg. t. 1020 = Zenobia speciosa.
drosericifolia, Linn. Mant. 229 = Erica glutinosa.
elegans, Meerb. Pl. Sel. Ic. Pict. t. 27 = Leucothoë racemosa.
elliptica, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 126 = Pieris ovalifolia.
empetrifolia, Lam. Encyc. i. 155 = Pernettya empetrifolia.
ericoides, Pall. Fl. Ross. i. ii. 56 = Cassiope ericoides.
eriophylla, Vell. in Vand. Enell. Fl. Lusit. et Bras. Sp. ed. Koem. 107 = Gaultheria ferruginea.
eucalyptoides, Cham. & Schlecht. in Linnæa, i. (1826) 512 = Leucothoë eucalyptoides.
fasciculata, Sw. Prod. Veg. Ind. Occ. 73 = Lyonia fasciculata.
fastigiata, Wall. in As. Res. xiii. (1820) 394 = Cassiope fastigiata.
ferruginea, Walt. Fl. Carol. 138 = Lyonia ferruginea.
flexuosa, Moon, Cat. ex Walp. Ann. i. 480 = Gaultheria fragrantissima.
floribunda, Pursh, Fl. Am. Sept. i. 293 = Pieris floribunda.
Fontanesiana, Steud. Nom. ed. II. i. 88 = Leucothoë axillaris.
formosa, Wall. in As. Res. xiii. (1820) 395; Wall. Cat. n. 761 = Pieris formosa.
formosa, Willd. Berl. Trav. N. & S. Car. Georg. ii. (1791) = Leucothoë acuminata.
frondosa, Pursh, Fl. Am. Sept. i. 295 = Lyonia paniculata.
glandulosa, *Wright, in Sauv. Fl. Cub.* 84.—Cuba.
glauca, Hort. ex C. Koch, Dendrol. ii. i. 118 = Zenobia speciosa.
glaucifolia, Wender. Ind. Sem. Hort. Marb. (1827) in App. n. 2 = A. Polifolia.
glaucophylla, Link, Enum. Hort. Berol. i. 394 = A. Polifolia.
globuliflora, Hort. ex DC. Prod. vii. 599 = Lyonia paniculata.
glomerata, Cav. Ic. vi. 43 = Gaultheria anastomosans.
grandiflora, Hort. ex Steud. Nom. ed. II. i. 88 = A. Polifolia.
grandiflora, Herb. Mus. Par. ex Dun. in DC. Prod. vii. 593 = Cavanillesmea grandiflorum.
grandiflora, Meerb. Pl. Sel. Ic. Pict. t. 25 = Pieris Mariana.
Hahniana, *Meerb. l. c.* t. 24.—Hab.?
Hahniana, Steud. Nom. ed. I. 44; ed. II. i. 88, sphalm. = Hahniana.

ANDROMEDA:—
hirsuta, Vell. Fl. Flum. 175; iv. t. 99 = Gaultheria ferruginea.
hirta, Wild. Enum. Hort. Berol. Suppl. 23 = Leucothoë hirta.
hispidula, Mart. ex Meisn. in Mart. Fl. Bras. vii. 165 = Leucothoë Martii.
humilis, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 326 = Pernettya pumila.
hypnoides, Linn. Sp. Pl. ed. I. 393 = Cassiope hypnoides.
ilicifolia, Pers. Syn. i. 481 = Leucothoë ilicifolia.
jamaicensis, Sw. Prod. Veg. Ind. Occ. 73 = Lyonia jamaicensis.
japonica, Naves, in Blanco, Fl. Filip. ed. III. 120 = Micromeluan tephrocarpum.
Japonica, Thunb. Fl. Jap. 181 = Pieris japonica.
kalmiæfolia, *Hoffmgg. Verz. Pfl. Nachtr.* i. 204.—Am. bor.
kataghirensis, Hook. Ic. Pl. t. 246 = Gaultheria fragrantissima.
lacustris, *Wright, in Sauv. Fl. Cub.* 85.—Cuba.
lanceolata, Desf. Cat. Hort. Par. ed. III. 186 = Leucothoë Catesbæi.
lanceolata, Wall. in As. Res. xiii. (1820) 390 = Pieris ovalifolia.
lanceolata, Vell. Fl. Flum. 175; iv. t. 97 = Leucothoë multiflora.
latifolia, *Wright, in Sauv. Fl. Cub.* 85.—Cuba.
laurina, Michx. Fl. Bor. Am. i. 258 = Leucothoë acuminata.
laxiflora, Mart. ex Meisn. in Mart. Fl. Bras. vii. 157 = Leucothoë laxiflora.
ledifolia, H. B. & K. Proleg. p. xvi = Arctostaphylos pollifolia.
ligustrina, Muhl. Cat. 44; Ell. Sketch, i. 490 = Lyonia paniculata.
litteralis, Steud. Nom. ed. II. i. 88 = Leucothoë littoralis.
longifolia, Pursh, Fl. Bor. Sept. i. 293 = Leucothoë Catesbæi.
lucida, Jacq. Coll. i. 95 = Leucothoë acuminata.
lucida, Lam. Encyc. i. 157 = Pieris nitida.
lycopodioides, Pall. Fl. Ross. i. ii. 55 = Cassiope lycopodioides.
marginata, Duham. Arb. ed. nov. i. 158. t. 40 = Pieris nitida.
mariana, Jacq. Ic. Rar. iii. t. 465 = Pieris nitida.
mariana, Linn. Sp. Pl. 393 = Pieris mariana.
Mertensiana, Bong. Veget. Ins. Sitcha, 152. t. 5 = Cassiope Mertensiana.
mexicana, *Steudl. in Biol. Centr. Am. Bot.* ii. 282.—Mexico.
montana, Buckl. in Am. Journ. Sc. xlv. (1843) 172 = Pieris floribunda.
montana, Salisb. Prod. 290 = Dahoëcia polifolia.
multiflora, Poir. Encycl. Pl. Bras. ii. 33. t. 122 = Leucothoë multiflora.
musciformis, Poir. Encyc. Suppl. i. 353 = Bryanthus Gmelini.
myrsinites, *Lam. Encyc.* i. 155.—Reg. Magell.
myrtifolia, Salisb. Prod. 290 = Pieris nitida.
myrtifolia, Willd. ex DC. Prod. vii. 608 = Agarista bracteomensis.
nana, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 47.—Japon.
nereifolia, Cham. & Schlecht. in Linnæa, i. (1826) 522 = Leucothoë nereifolia.
nitida, Bartr. ex Marsh. Arbust. (1785) 8 = Pieris nitida.
nitida, Vell. Fl. Flum. 174; iv. t. 94 = Leucothoë revoluta.
nummularia, Cham. & Schlecht. in Linnæa, i. (1826) 530 = Leucothoë nummularia.
obtusa, *Wright, in Sauv. Fl. Cub.* 84.—Cuba.
octandra, Sw. Prod. Veg. Ind. Occ. 73 = Lyonia octandra.
olacefolia, *Hoffmgg. Verz. Pfl. Nachtr.* i. 204.—Hab.?
oleifolia, Cham. in Linnæa, viii. (1833) 504 = Leucothoë oleifolia.
oleifolia, Hort. ex Steud. Nom. ed. II. i. 89 = A. Polifolia.
ovalifolia, Wall. in As. Res. xiii. (1820) 391; Cat. n. 758 = Pieris ovalifolia.
ovata, Soland. ex DC. Prod. vii. 596 = Zenobia speciosa.
paniculata, Lina. Sp. Pl. 394 = Leucothoë racemosa.
paniculata, Willd. Sp. Pl. ii. 612 = Lyonia paniculata.

ANDROMEDA:—
parabolica, Duham. Arb. ed. II. i. 191 = Lyonia parabolica.
perulata, Maig. in Ann. Mus. Bot. Lugd. Bat. i. 31 = Eukianthus japonicus.
phillyreaefolia, Hook. Ic. Pl. t. 122 = Pieris phillyreaefolia.
pilulifera, Hort. ex DC. Prod. vii. II. 599 = Lyonia paniculata.
Pieris, Cham. in Linnæa, viii. (1833) 508 = Leucothoe subrotunda.
Polifolia, *Linn. Sp. Pl.* 393.—Reg. bor. et arct.
polyantha, Stend. Nom. Bot. ed. II. i. 89 = Lyonia paniculata.
populifolia, Lam. Encyc. i. 159 = Leucothoë acuminata.
prostrata, Cav. Ic. vi. 48 = Ferentya microphylla.
pubescens, Poir. Encyc. Suppl. i. 355 = Lyonia rubiginosa.
pulchella, Cham. ex Meisn. in Mart. Fl. Bras. vii. 161 = Leucothoe pulchra.
pulchella, Cham. in Linnæa, viii. (1833) 509 = Leucothoe pulchella.
pulchella, Salisb. Prod. 389 = Pieris mariana.
pulchra, Vell. Fl. Flum. 174; iv. t. 96 = Gaultheria venusta.
pulchra, Cham. & Schlecht. in Linnæa, i. (1826) 521 = Leucothoe pulchra.
pulverulenta, Bartr. Trav. N. & S. Car. Georg. ii. 476 = Zenobia speciosa.
punctata, Dumb. ex Dun. in DC. Prod. vii. 564 = Thibaudia punctatifolia.
pyrifolia, Thou. Obs 14; Pers. Syn. i. 481 = Agauria salicifolia.
racemosa, Linn. Sp. Pl. 394 = Leucothoë racemosa.
racemosa, Walt. Fl. Carol. 138 = Lyonia paniculata.
recurva, Buckl. in Am. Journ. Sc. xlv. (1843) 172 = Leucothoe recurva.
Redowskii, Cham. & Schlecht. in Linnæa, i. (1826) 517 = Cassiope Redowskii.
reticulata, Walt. Fl. Carol. 137 = Leucothoe acuminata.
revoluta, Bouton, ex DC. Prod. vii. II. 603 = Leucothoe Bojeri.
revoluta, Hort. ex Steud. Nom. ed. II. i. 89 = A. Polifolia.
revoluta, Spreng. Neue Entdeck. ii. 131 = Leucothoë revoluta.
rhomboidalis, Veill. in Duham. Arb. ed. II. i. 192 = Gaultheria ferruginea.
rigida, Humb. Itin. i. 397 = Gaultheria rigida.
rigida, Pursh, Fl. Am. Sept. i. 292 = Lyonia ferruginea.
rosmarinifolia, Gilib. Fl. Lithuan. i. 3 = A. Polifolia.
rubiginosa, Pers. Syn. i. 481 = Lyonia rubiginosa.
rupestris, Forst. f. Prod. n. 186 = Gaultheria rupestris.
salicifolia, Lam. Encyc. i. 159 = Agauria salicifolia.
secunda, Moench, Meth. 457 = A. Polifolia.
Sellowii, Stend. Nom. ed. II. i. 89 = Leucothoë pulchra.
serrata, Sm. ex DC. Prod. vii. II. 609 = Leucothoe acuminata.
serrata, Vell. Fl. Flum. 174; iv. t. 96 = Gaultheria elliptica.
serratifolia, Hort. ex DC. Prod. vii. 602 = Zenobia speciosa.
serrulata, Cham. in Linnæa, viii. (1833) 506 = Leucothoë serrulata.
sinensis, Steud. Nom. ed. II. i. 89 = Vaccinium bracteatum.
speciosa, Michx. Fl. Bor. Am. i. 256 = Zenobia speciosa.
speciosa, Willd. ex Steud. Nom. ed. II. i. 89 = Leucothoë racemosa.
spicata, Wats. Dendrol. i. t. 36 = Leucothoë racemosa.
spinulosa, Pursh, Fl. Am. Sept. i. 293 = Leucothoë Catesbæi.
squamulosa, D. Don, Prod. Fl. Nep. 149 = Pieris ovalifolia.
Stelleriana, Pall. Fl. Ross. i. II. 58 = Cassiope Stelleriana.
subrotunda, Pohl, Pl. Bras. ii. 93. t. 191 = Leucothoë subrotunda.
subsessilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 32.—Japan.
symplocifolia, Wall. Cat. n. 1522 = Vaccinium Leschenaultii.

ANDROMEDA:—
taxifolia, Pall. Fl. Ross. ii. 64. t. 72. f. 2 = Phyllodoce taxifolia.
tetragona, Linn. Sp. Pl. 393 = Cassiope tetragona.
tomentosa, Dum.-Cours. Bot. Cult. ed. II. iii. 495 = Lyonia racemosa.
trinervis, Donb. ex Dun. in DC. Prod. vii. 563 (= *Proclesia bracteata*).—Peruv.
vaccinifolia, Hort. ex C. Koch, Dendrol. ii. t. 116 = Pieris floribunda.
venulosa, DC. Prod. vii. 607 = Arctostaphylos Hookeri.
villosa, Wall. Cat. n. 762 = Pieris villosa.
Walteri, Willd. Enum. Hort. Berol. 488 = Leucothoë Catesbæi.
Watsonana, Steud. Nom. ed. II. i. 89 = Lyonia parabolica.

ANDROMYCIA, A. Rich. Fl. Cub. Fanerog. ii. 282. t. 89 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 987.
cubensis, *A. Rich. l. c.*—Cuba.

ANDROPHORANTHUS, Karst. in C. Koch & Fint. Wochenschr. (1859) 5 = **Caperonia**, St. Hil. (Euphorb.).
glandulosus, Karst. l. c. 6 = C. palustris.

ANDROPHYLAX, Steud. Nom. ed. II. i. 89 (1840) = seq.

ANDROPHYLAX, Wendl. Beobacht. 38 (1798) = **Cocculus**, Linn. (Menisp.).
scandens, Wendl. l. c. = C. carolinus.

ANDROPHTHOË, Scheff. in Flora, lii. (1870) 249, err. typ. = DENDROPHTHOE, Mart. = **Loranthus**, Linn.

ANDROPHYSA, Moq. ex scipso in DC. Prod. xiii. II. 201 (1849) = **Halocharis**, Moq. (Chenop.).

ANDROPOGON, Linn. Sp. Pl. ed. I. 1045 (1753). *GRAMINEAE*, Benth. & Hook. f. iii. 1183.
AGENIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447 (1836).
ALLOJANTHENA, Steud. Syn. Pl. Gram. 217 (1855).
ALLOJATHEROS, Rafin. in Ser. Bull. Bot 221 (1830).
ANGELPHIA, Hack. in Engl. Bot. Jahrb. vi. (1885) 240.
ANATHERUM, Beauv. Agrost. 128 t. 22 (1812).
ARTHROSTACHYS, Desv. Opusc. 74. t. 6 (1831).
CYMBACHNE, Retz. Obs. vi. 36, obs. ad calc. (1791).
CYMBOPOGON, Spreng. Pugill. ii. 14 (1815).
DICHANTHIUM, Willem. in Ust. Ann. Bot. xviii. 11 (1796).
DICTOMIS, H. B. & K. Nov. Gen. et Sp. i. 193. t. 64 (1815).
DIMERATHEMON, Rafin. Neogenyt. 4 (1825).
DIPLASANTHUM, Desv. Opusc. 66 (1831).
DISCHANTHIUM, Kunth, Enum. Pl. i. 498 (1833).
ERIOPODIUM, Hochst. in Flora, xxix. (1846) 115, in obs.
EUKLASTAXON, Steud. in Flora, xxxiii. (1850) 1. 229; Syn. Pl. Gram. 412 (1854).
EUPOGON, Desv. Opusc. 67, in obs. (1831).
EXOTHECA, Anders. in Nov. Act. Upsal. ii. (1856) 253. t. 8.
GRAFA, Am. ex Steud. Nom. ed. II. i. 705 (1840).
GYMNANDROPOGON, [Munro] Duthie, in Atkins. Gaz. N. W. Ind. 638 (1878?), nomen.
GYMNANTHELIA, Anderss. in Schweinf. Beitr. Fl. Aeth. 299 (1867).
HETEROCHLOA, Desv. Opusc. 66. t. 5. f. 3 (1831).
HOMOTHERUM, Nees, in Hook. & Arn. Bot. Beech. Voy. 239 (1836).
HYPARRHENIA, Anderss. in Nov. Act. Upsal. ii. (1856) 231, nomen.
HYPOGYNIUM, Nees, Agrost. Bras. 364 (1829).
LENORMANDIA, Steud. in Flora, xxxiii. (1850) 229, nomen; Syn. Pl. Gram. 359 (1854).
LEPTOCERCIS, Trin. Fund. Agrost. 208 (1820).
MANDELORNA, Steud. Syn. Pl. Gram. 359 (1854).
OROPOGON, Neck. Elem. iii. 211 (1790).
PITHECURUS, Willd. ex Kunth, Enum. Pl. i. 490 (1833).
SCHIZACHYRIUM, Nees, Agrost. Bras. 331 (1829).
SCHIZOPOGON, Reichb. Consp. 53 (1828).
SPIROTHERUS, Rafin. in Ser. Bull. Bot. i. (1830) 221.

ANDROPOGON:—
VETIVERIA, Thou. ex Virey, in Journ. Phar. Sér. I. xvi. (1827) 499.
abyssinicus, *R. Br. ex Fres. in Mus Senkenb.* ii. (1837) 146.—Abyss.
aciculatus, Willd. Sp. Pl. iv. 906 = Chrysopogon aciculatus.
aciculatus, Retz. Obs. v. 22 = Chrysopogon aciculatus.
aciculatus, Steud. Nom. ed. II. i. 90 = Chrysopogon aciculatus.
adustus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 259 = Elionurus laniflorus.
aesthesus, Steud. Syn. Pl. Gram. 376 = Ischaemum ciliare
aethiopicus, Rupr. ex Steud. Syn. Pl. Gram. 372 = Cayaux.
afer, Bruce, Trav. v. App 47 (Pollinize sp. ?).—Afr. bor.
affinis, J. Presl, in Rel. Haenk. i. 348 = Sorghum halepense.
affinis, *R. Br. Prod.* i. 201.—Austral.
affinis, Spreng Nov. Prov. 5 = annulatus.
Agenium, Steud. Syn. Pl. Gram. 396 = Nees.
agrostoides, *Spreg. in Anal. Soc. Cient. Argent.* xvi. (1883) 186.—Reg. Argent.
agrostoides, Speg. Pl. Nov Am. Austr. Dec. ii. 27; ex Anal. Soc. Cient. Argent. xvi. (1883) 15 = Chrysopogon avenaceus.
albescens, Journ. Mex. Pl. Enum. Gram. 56; ex Hemsl. Biol. Centr Am. iii. 580 = Chrysopogon nutans.
albidus, *Wall. Cat.* n. 8821.—Ind. or.
Alectoridia, Steud. Syn. Pl. Gram. 383 = Arthraxon ciliaris.
Allionii, DC. Fl. Fr. iii. 97 = Heteropogon contortus.
alopecuroides, Linn Sp Pl. 1045 = Eriauthus saccharoides.
alternans, J. Presl, in Rel. Haenk. i. 342 = Chrysopogon violaceus.
altissimus, Colla, in Mem. Acc. Torin. xxxix. (1836) 28. t. 80 = saccharoides.
altissimus, *Hochst. ex A. Br. in Flora*, xxiv. (1841) 277. —Abyss
amaurus, Buese, in Miq. Pl. Jungh. 360 = Pollinia praemorsa.
ambiguus, Gennari, ex Barbey, Consp. Fl. Sard. 190 = Heteropogon hirtus.
ambiguus, Michx. Fl. Bor. Am. i. 58 = Gymnopogon racemosus.
ambiguus, Steud. Syn. Pl. Gram. 385 = exaltatus.
amethystinus, *Steud. l. c.* 371.—Afr. trop.
amplectens, *Nees, Fl. Afr. Austr.* 104.—Afr. austr.
amplexifolius, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 274 = Arthraxon ciliaris.
amplifloras, Steud. Syn. Pl. Gram. 388 = A. Nardus.
amplus, J. Presl, in Rel. Haenk. i. 339 = glaucescens.
angustatus, Steud. Syn. l. 370 = apricus.
angustifolius, H. B. & K. Nov. Gen. et Sp. i. 184 = Trachypogon angustifolius.
angustifolius, Parl. fl. Palerm. i. 269 = pertusus.
angustifolius, Sibth. & Sm. Prod. Fl. Græc. i. 47 = A. Ischaemum.
anias, *Llanos, Fragm.* 39.—Ins. Philipp.
annulatus, F. Muell. Fragm. viii. 123 = articens.
annulatus, *Forsk. Fl. Aegypt. Arab.* 173.—Afr. bor.; Ind. or.; Austral.
annulatus, F. Schmidt, Fl. Ins. Caprverd. 157 = A. Ischaemum.
annulatus, Schult. Mant. ii. 453 = foveolatus.
annuus, *Hack. in Flora*, lxviii. (1885) 137.—Afr. trop.
anomalus, *Steud. Syn. Pl. Gram.* 397.—Arabia.
anthephoroides, Steud. l. c. 375 = Ischaemum anthephoroides.
antihirtoides, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 468.—Abyss.
antihirtoides, Rupr. in Bull. Acad. Brux. ix. (1842) 345 = Ruprechtii.
appendiculatus, *Nees, Fl. Afr. Austr.* 105.—Afr. austr.
apricus, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 70. —Reg. trop.
arctatus, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 20.—Florida.
arcuatus, Steud. Syn. Pl. Gram. 374 = Ischaemum fasciculatum.
arenarius, *Hack. in Flora*, lxviii. (1885) 134.—Uruguay.

ANDROPOGON:—
argenteus, DC. Cat. Hort. Monsp. 77 = saccharoides.
argenteus, Ell. Sketch, 148 = argyreus.
argutus, Steud. Syn. Pl. Gram. 391 = Chrysopogon argutus.
argyreus, *Schult. Mant.* ii. 450.—Am. bor.
Ariani, Edgew. in Journ. Linn. Soc. vi. (1862) 208 = A. Iwarancsa.
aristatum, Poir. Encyc. Suppl. i. 585 = asperum.
aristidoides, Steud. Syn. Pl. Gram. 397 = Chrysopogon ciliolatus.
aristulatus, *Hochst. ex Steud. Syn. Pl. Gram.* 397 = Chrysopogon Wightianus.
aristulatus, Steud. Syn. Pl. Gram. 372 (Ischaemi sp. 1).—Ins. Borbon.
armillaris, *Willd. ex Steud. Nom.* ed. II. i. 90.—Ind. or.
Arnottianus, Steud. Syn. Pl. Gram. 375 = Ischaemum rugosum.
Arnottii, *Nees, ex Steud. l. c.* 380.—Ind. or.
aromaticus, Roxb. ex Schult. Mant. 442 = muricatus.
aromaticus, Sieber, ex Schult. l. c. 481, 388 = A. Schoenanthus.
arrhenobasis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 460 = Heteropogon arrhenobasis.
arundinaceus, Berg. Desc. Pl. Cap. 356 = Bergii.
arundinaceus, Scop. Fl. Carn. ed. II. ii. 274 = Sorghum halepense.
arundinaceus, Voigt, Hort. Suburb. Calc. 706 = Anthistiria ciliata.
asper, Kunth, Rev. Gram. i. 166 = Danthonia papillosa.
asperifolius, Hack. in Flora, lxviii. (1885) 140 = Zollingeri.
assimilis, Steud. Syn. Pl. Gram. 397 = Chrysopogon pictus.
asthenus, Steud. l. c. 376 = Ischaemum ciliare.
asthenostachys, Steud. l. c. 381 = Pollinia articulata.
asticlus, *Busse, in Mig. Fl. Sungh.* 360.—Sumatra.
attenuatus, *Bertol. in Mem. Acc. Sc. Bolog.* x. (1859) 84; *Misc. Bot.* xx. 10, t. 2.—Guatemala.
Auctori, Boiss. Diagn. Ser. I. v. 77 = Chrysopogon serrulatus.
aureo-fulvus, Steud. Syn. Pl. Gram. 373 = Pollinia Cumingii.
aureo-villosus, Steud. Syn. Pl. Gram. 380 = Pollinia aurea.
aureus, Bory, Voy. i. 367. t. 21 = Pollinia aurea.
australis, *Spreng. Syst.* i. 387.—Austral.; Timor.
avenaceus, Berg. Desc. Pl. Cap. 356 = Bergii.
avenaceus, H. B. & K. Nov. Gen. et Sp. i. 189 = Sorghum halepense.
avenaceus, Michx. Fl. Bor. Am. i. 58 = Chrysopogon avenaceus.
barbatus, *Linn. Pl. Jam. Pugill.* 30; *Mant.* ii. 302 = Chloris barbata.
barbigerus, Steud. Syn. Pl. Gram. 376 = Ischaemum aristatum.
barbinodis, Lag. Gen. et Sp. Nov. 3 = saccharoides.
Barteri, *Hack. in Flora*, lxviii. (1885) 124.—Afr. trop.
Bellardi, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 317 = Heteropogon contortus.
bellariensis, *Hack. in Flora*, lxviii. (1885) 123.—Ind. or.
Belvisii, Desv. Opusc. 67 = argyreus.
Benthamianus, Steud. Syn. Pl. Gram. 382 = condensatus.
Bergii, *Roem. & Schult. Syst.* ii. 813.—Afr. austr.
Berterronianus, Steud. & Hochst. ex Steud. Syn. Pl. Gram. 380 = saccharoides.
Besseri, Kunth, Rev. Gram. i. 166 = Sorghum vulgare.
bewshiensis, Steud. Syn. Pl. Gram. 367 = Heteropogon contortus.
biaristatus, Steud. l. c. 379 = Pollinia ciliata.
bicolor, Nees, Fl. Afr. Austr. 113 = dichroos.
bicolor, Roxb. Hort. Beng. 21; Fl. Ind. i. 272 = Sorghum vulgare.
bicornis, Benth. Pl. Hartw. 263 = condensatus.
bicornis, *Linn. Sp. Pl.* 1046.—Am. austr.
bicornis, Forsk. Fl. Aegypt. Arab. 178 = A. Schoenanthus.
bifidus, Steud. Syn. Pl. Gram. 378 = Ischaemum ciliare.
bifoveolatus, Steud. l. c. 380 = pertusus.
bilinguis, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 262 = Elionurus bilinguis.

ANDROPOGON:—
binatus, Retz. Obs. v. 21 = Spodiopogon pertusus.
binatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 255 = tenellus.
bipennatus, *Hack. in Flora*, lxviii. (1885) 142.—Afr. trop.; Madag.
bisquamulatus, Hochst. in Flora, xxvii. (1844) 245 = Gayanus.
Blashki, Retz. Obs. ii. 27 = annulatus.
Blumii, Steud. Syn. Pl. Gram. 378 = Spodiopogon Blumii.
Boivini, *Steud. l. c.* 481.—Ins. Nosibé.
bombycinus, *R. Br. Prod.* i. 205.—Austral.
Bourgaei, *Hack. in Flora*, lxviii. (1885) 134.—Mexico.
bourgeanus, Steud. Syn. Pl. Gram. 382 = Ischaemum digitatum.
brachyatherus, *Hochst. in Flora*, xxvii. (1844) 241.—Abyss.
brachystachyus, *Chapm. Fl. S. Un. St. Suppl.* 668.—Florida.
bracteatus, *Willd. Sp. Pl.* iv. 914.—Am. trop.
brevearistatus, Steud. Syn. Pl. Gram. 398 = Chrysopogon Wightianus.
brevifolius, *Sw. Prod. Veg. Ind. Occ.* 26.—Geront. trop.
brevus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 265 = lateralis.
Broughlarti, *Steud. Nom.* ed. II. i. 90.—Ins. Molucc.
Brownii, Kunth, Rev. Gram. i. 163 = Ischaemum laxum.
Byronis, Steud. Syn. Pl. Gram. 398 = Spodiopogon Byronis.
Cabanillesii, *Hack. in Flora*, lxviii. (1885) 133.—Am. bor.
caerulescens, Kunth, Enum. Pl. i. 504 = Chrysopogon violascens.
caespitosus, A. Rich. Tent. Fl. Abyss. ii. 451 = Elionurus argenteus.
caespitosus, Phil. in Linnaea, xxxiii. (1864-65) 275 = hirtifolius.
caffrorum, Kunth, Rev. Gram. i. 165 = Sorghum vulgare.
campestris, Kunth, Enum. Pl. i. 489 = tener.
campestris, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 277.—Bras.
camporum, Trin. ex Steud. Syn. I. 378 = campestris.
canaliculatus, *Schum. Beshriv. Guin. Pl.* 52.—Guinea.
candidus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 266 = Elionurus candidus.
canescens, Kunth, Enum. Pl. i. 487 = Trachypogon polymorphus.
canus, *Steud. Nom.* ed. II. i. 90.—Mexico.
capensis, Houtt. Plant.-Synst. xii. t. 93. f. 3 = Chloris petraea.
capillaris, Heyne, ex Steud. Syn. Pl. Gram. 116 = Aristida mutica.
capillaris, *Kunth, Rev. Gram.* i. 166.—Afr. austr.
capillifolius, Steud. Syn. Pl. Gram. 397 = Chrysopogon violascens.
carinosus, *Linn. Sp. Pl.* ed. II. 1480.—As. trop.; Ins. Mascar.
carinatus, *Nees, Agrost. Bras.* ii. 280.—Bras.
castratus, Griff. Notul. iii. 89 = Arthraxon nudo!
caucasicus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 286 = intermedius.
caudatus, Bieb. Tabl. Prov. Côte Occ. Mer Casp. 56 = Erianthus Ravennae.
ceresiaeformis, *Nees, Fl. Afr. Austr.* 109.—Afr. austr.
cerifrus, Hack. in Mart. Fl. Bras. ii. III. 281 = A. Nardus.
cernuus, Kunth, Enum. Pl. i. 501 = Sorghum halepense.
cernuus, Roxb. Fl. Ind. i. 270 = Sorghum vulgare.
chlorideiformis, Gandich. in Freyc. Voy. Bot. 412 = Dimeria pilosissima.
chlorostachys, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 260 = Elionurus rostratus.
chordatus, Steud. Syn. Pl. Gram. 398 = Spodiopogon Blumii.
chrysatherus, F. Muell. in Linnaea, xxv. (1852) 448 = sericeus.
chrysopus, Cass. in Bull. Soc. Bot. Fr. xxii. (1875) 66 = Auctori.
chrysostachyus, *Steud. Syn. Pl. Gram.* 377.—Abyss.
ciliatus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 260 = Elionurus latiflorus.

ANDROPOGON:—
ciliatus, Ell. Sketch, i. 144 = Chrysopogon avenaceus.
ciliatus, Thunb. Fl. Jap. 40 = Anthistiria barbata.
ciliolatus, Steud. Syn. Pl. Gram. 396 = Chrysopogon serrulatus.
cinctus, Steud. l. c. 398 = Chrysopogon violascens.
circinnatus, Hochst. & Steud. ex Steud. l. c. 387 = laniger.
cirratus, *Hack. in Flora*, lxviii. (1885) 119.—Am. bor.
citratus, Hort. ex DC. Cat. Hort. Monsp. 78 = A. Schoenanthus.
citreus, K. Br. Prod. i. 203 = Elionurus citreus.
citrodorus, Hort. ex Desf. Tabl. ed. II. 15 = A. Schoenanthus.
citronus, Hort. ex Steud. Nom. ed. II. i. 90 = A. Nardus.
clandestinus, Hale, ex Vasey, Gram. Un. St. 19 = Elliottii.
clandestinus, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Schoenanthus.
cornutus, Steud. l. c. 395 = Chrysopogon serrulatus.
cosinus, Nees, ex Hook. & Arn. Bot. Beech. Voy. 244 = A. Schoenanthus.
cognatus, Steud. Syn. Pl. Gram. 374 = schirensis.
coleotrichus, *Steud. l. c.* 396.—Abyss.
commutatus, Aitch. in Journ. Linn. Soc. xix. (1882) 191 = laniger.
commutatus, *Steud. Syn. Pl. Gram.* 387.—Abyss.
comorensis, Steud. l. c. 481 (= Ischaemi sp.).
comosus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 461 = coleotrichus.
comosus, Link, Hort. Berol. i. 289 = annulatus.
compactus, Brot. Fl. Lusit. i. 88 = Sorghum halepense.
condensatus, *H. B. & K. Nov. Gen. et Sp.* i. 188.—Amer. austr.
condensatus, Rupr. in Bull. Acad. Brux. ix. (1842) 344 = Chrysopogon Francavillanus.
condylotrichus, *Hochst. ex Steud. Syn. Pl. Gram.* 377.—Abyss.
confertiflorus, Steud. l. c. 385 = A. Nardus.
confinis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 461.—Abyss.
conjugatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 260 (Pollinise sp.).—Bengal.
connatus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 464 = A. Schoenanthus.
consanguineus, *Kunth, Enum. Pl.* i. 494.—Bras.; Uruguay.
consanguineus, Trin. in Linnaea, x. (1836) 291 = condensatus.
consimilis, Steud. Syn. Pl. Gram. 394 = serratus.
contortus, Linn. Sp. Pl. ii. 1045 = Heteropogon contortus.
controversus, Steud. Syn. Pl. Gram. 391 = Sorghum halepense.
cordatifolius, Steud. l. c. 376 = Ischaemum conjugatum.
cordofanus, Hochst. in Flora, xxvii. (1844) 245 = Gayanus.
Cornucopiae, *Hack. l. c.* lxviii. (1885) 126.—Afr. trop.
corollatus, Steud. Syn. Pl. Gram. 389 = Ischaemum corollatum.
cotulifer, Thunb. Fl. Jap. 41 = Pollinia cotulifera.
courtallumensis, Steud. Syn. Pl. Gram. 377 = Apocopis Wightii.
corassipes, Steud. l. c. 375 = Ischaemum Sieboldii.
crinitus, Thunb. Fl. Jap. 40 = Pogonatherum saccharoideum.
Crupina, Kunth, Rev. Gram. i. 166 = Sorghum halepense.
cryptatherus, Steud. Syn. Pl. Gram. 376 = Ischaemum australe.
cubensis, *Hack. in Flora*, lxviii. (1885) 131.—Cuba.
curtifoundulatus, Steud. Nom. ed. II. i. 90 = Eutriana cartipendula.
Curtisianus, Steud. Syn. Pl. Gram. 390 = virginicus.
cuspidatus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 455 = Arthraxon cuspidatus.
cymbarius, *Linn. Mant.* ii. 308.—Afr. trop.; Ind. or.
dactyloides, Steud. Syn. Pl. Gram. 381 = Trachypogon polymorphus.
debilis, Kunth, Enum. Pl. i. 488 = brevifolius.
decolorans, H. B. & K. Nov. Gen. et Sp. i. 190 = Sorghum halepense.
decussatus, Steud. Syn. Pl. Gram. 377 = koloostachys.
demissus, *Steud. l. c.* 386.—Ind. or.
densiflorus, Steud. l. c. 386 = A. Schoenanthus.

ANDROPOGON :—

dennus, Desv. in Ham. Prod. Fl. Ind. Occ. 8 = concinnatus.

depauperatus, *Kunth*, *Rev. Gram.* i. 166.—Bengal.

depressus, Steud. Syn. Pl. Gram. 373 = Spodiopogon depressus.

Dianae, Steud. l. c. 371 = leucostachyus.

diatherus, Steud. l. c. 378 = Ischaemum latifolium.

dichroanthus, Steud. l. c. 393 = serratus.

dichroos, *Steud. l. c.* 380.—Afr. austr.

digitatus, Hochst. ex Steud. l. c. 382 = Trachypogon dactyloides.

digitatus, *Vahl, ex Trin. Fund.* 186.—Hab. ?

diplandrus, *Hack. in Flora*, lxviii. (1885) 123.—Afr. trop.

dissitiflorus, Michx. Fl. Bor. Am. i. 57 = virginicus.

dissitiflorus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 267 = consanguineus.

dissolutus, Steud. Syn. Pl. Gram. 381 = Trachypogon polymorphus.

distachyos, *Linn. Sp. Pl.* 1046.—Reg. Mediterr.

distans, Nees, ex Steud. Syn. Pl. Gram. 387 = A. Nardus.

divaricatus, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. i. 99 = Aristida Humboldtiana.

divaricatus, *Linn. Sp. Pl.* 1045.—Virginia.

diversiflorus, Steud. in Zoll. Syst. Verz. 58 = Pollinia praemorsa.

domingensis, Fourn. Mex. Pl. Enum. Gram. 61; ex Hemsl. Biol. Centr. Am. iii. 530 = leucostachyus.

domingensis, Spreng. ex Steud. Nom. ed. II. i. 91 = Trachypogon polymorphus.

domingensis, Steud. l. c. 45 = leucostachyus.

Donianus, Benth. in Hook. Niger Fl. 570 = Elionurus platypus.

Dregeanus, Nees, Gram. Afr. Austr. 112 = Schimperi.

Drummondii, Steud. Syn. Pl. Gram. 393 = Sorghum vulgare.

dubitatus, Steud. l. c. 394 = Sorghum halepense.

dubius, C. Koch, in Linnaea, xxi. (1848) 443 = Sorghum halepense.

dubius, Kunth, Rev. Gram. i. 166 = Elionurus isostratus.

dulcis, *Burm. f. Fl. Ind.* 219 = Sorghum vulgare.

durus, Steud. Nom. ed. II. i. 91 = Erianthus strictus.

echinatus, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 182.—Ind. or.

echinatus, Heyne, in Steud. Nom. ed. II. i. 91 = Arthraxon lanceolatus.

echinulatus, Steud. Syn. Pl. Gram. 397 = Chrysopogon Gryllus.

elatus, Steud. l. c. 376 = Ischaemum aristatum.

elegans, J. Gay, ex Steud. l. c. 364 = Elionurus elegans.

elegantissimus, Steud. l. c. = Elionurus Royleanus.

Elliottii, *Chapm. Fl. S. Un. St.* 581.—Am. bor.

elongatus, Spreng. Syst. i. 287 = Chrysopogon elongatus.

elythrophyllus, *Steud. Syn. Pl. Gram.* 378.—Mexico.

emersus, Fourn. Mex. Pl. Enum. Gram. 58; ex Hemsl. Biol. Centr. Am. iii. 527 = saccharoides.

erianthoides, *F. Muell. Fragm.* x. 75.—Australi.

Erianthus, Link, Hort. Berol. i. 248 = Erianthus saccharoides.

eriophorus, Scheele, in Flora, xxvii. (1844) 51 = virginicus.

eriophorus, Willd. Sp. Pl. iv. 910 = laniger.

eriostachyus, J. Presl, in Rel. Haenk. i. 339 = ternatus.

erythrocoleos, *Steud. Syn. Pl. Gram.* 391.—Nepal.

Esenbeckii, Steud. l. c. 375 = Ischaemum aristatum.

eucnemis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 276.—Guinea.

eucomus, *Nees, Fl. Afr. Austr.* 104.—Afr. trop. et austr.

exaltatus, *R. Br. Prod.* i. 202.—Austral.

exaratus, *Hack. in Flora*, lxviii. (1885) 135.—Paraguay.

excavatus, Hochst. l. c. xxix. (1846) 115 = A. Schoenanthus.

exilis, *Hochst. l.c.* xxvii. (1844) 143.—Afr. trop.

exsertus, Steud. Syn. Pl. Gram. 366 = brevifolius.

falcatus, Steud. l. c. 389 = Ischaemum pectinatum.

familiaris, *Steud. l. c.* 385.—Afr. trop.

fasciculatus, *Roxb. Hort. Beng.* 83; *Fl. Ind.* i. 269.—Ind. or.

fascicularis, Thw. Enum. Pl. Zeyl. 487 = intermedius.

fasciculatus, Linn. Sp. Pl. 1047 = Chloris radiata.

fasciculatus, Steud. Syn. Pl. Gram. 383 = Ischaemum fasciculatum.

fastigiatus, *Sw Prod. Veg. Ind. Occ.* 26.—Reg. trop.

ANDROPOGON :—

fernaisi, Fourn. Mex. Pl. Enum. Gram. 62; ex Hemsl. Biol. Centr. Am. iii. 527 = hirtiflorus.

ferrugineus, Hochst. ex Steud. Syn. Pl. Gram. 377 = chrysostachyus.

festucoides, J. Presl, in Rel. Haenk. i. 340 = squarrosus.

filifolius, *Steud. Syn. Pl. Gram.* 374.—Afr. austr.

filiformis, Pers. Syn. i. 103 = caricoides.

filiformis, Roxb. Hort. Beng. 6; Fl. Ind. i. 260 = Dimeria ornithopoda.

filipendulus, *Hochst. in Flora*, xxix. (1846) 115.—Afr. trop.; Zeylan.; Austral.

finitimus, *Hochst, ex A. Rich. Tent. Fl. Abyss.* ii. 465. —Abyss.; Ins. Mascar.

firmandus, Steud. Syn. Pl. Gram. 370 = Pollinia praemorsa.

firmus, Kunth, Enum. Pl. i. 486 = Heteropogon firmus.

flabelliformis, *Reab. ex Spreng. Pugill* ii. 9.—Ind. or.

flaccidus, A. Rich. Tent. Fl. Abyss. ii. 462 = brevifolius.

flavescens, *F. Presl, in Rel. Haenk.* i. 330.—Peruv.

flexile, Pott. Encyc. Suppl. i. 583 = scoparius.

flexuosus, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Nardus.

floridus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 265 = brevifolius.

foliatus, Steud. Syn. Pl. Gram. 389 = A. Schoenanthus.

foveolatus, *Delile, Fl. Egypte*, 160. t. 8. f. 2.—Afr. et As. trop.

fragilis, R. Br. Prod. i. 202 = brevifolius.

fragilissimus, Hochst. ex Steud. Syn. Pl. Gram. 384 = exilis.

Francavillanus, Fourn. Pl. Mex. Enum. ii. 50; ex Hemsl. Biol. Centr. Am. iii. 530 = Chrysopogon avenaceus.

fulvibarbis, *Trin in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 287.—Afr. trop.

fulvescens, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 463 = rufus.

fuloispica, Steud. Syn. Pl. Gram. 379 = Pollinia Cumingii.

fulvus, *Spreng. Pugill.* ii. 8.—Bengal.

furcatus, Muhl. in Willd. Sp. Pl. iv. 919 = provincialis

fuscus, J. Presl, in Rel. Haenk. i. 342 = serratus.

Galeottii, *Fourn. Mex. Pl. Enum. Gram.* 56; *ex Hemsl. Biol. Centr. Am.* iii. 527 = Chrysopogon nutans.

gallicus, *Steud. Syn. Pl. Gram.* 371.—Ins. Mascar.

garipensis, Steud. l. c. 379 = annulatus.

Gayanus, *Kunth, Rev. Gram.* i. 168.—Senegal.

geniculatus, Steud. Syn. Pl. Gram. 378 = Ischaemum aristatum.

Gerardi, Vitm. Summa Pl. vi. 16 = provincialis.

gibbus, Steud. Syn.Fl.Gram. 376 = Ischaemum aristatum.

Gidarba, *Buch.-Ham. in Wall. Cat.* n. 8797 —Ind. or.

gigantéus, Fenzl, ex Steud. Syn. Pl. Gram. 386 = A. Iwarancusa.

giganteus, Hochst. in Flora, xxvii. (1844) 242 = A. Schoenanthus.

gigenteus, Tenore, Fl. Nap. v. 285 = Heteropogon hirtus.

glaber, Roxb. Hort. Beng. 7; Fl. Ind. i. 271 = intermedius.

glabratus, Steud. Nom. Bot. ed. II. i. 91 = Pollinia glabrata.

glabratus, Steud. Syn. Pl. Gram. 395 = Chrysopogon Gryllus.

glabriflorus, Hochst ex Steud. l. c. 374 = polystherus.

glabriusculus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 468.—Abyss.

glaucescens, Fourn. Pl. Mex. ii. 60; ex Hemsl. Biol. Centr. Am. iii. 527, partim = Bourgaei.

glaucescens, *H. B. & K. Nov. Gen. et Sp.* i. 186.—Am. austr.

glaucescens, Schlecht. in Linnaea, vi. (1831) 39 = virginicus.

glaucopsis, Ell. ex Steud. Nom. ed. II. i. 91 = macrourus.

glaucopsis, Steud. Syn. Pl. Gram. 397 = montanus.

glaucus, Muhl. Descr. Uber. 278 = macrourus.

glaucus, Retz. Obs. v. 20 = Apluda mutica.

glaucus, Torr. in Ann. Lyc. N. York, i. (1824) 157 = saccharoides.

Glaziovii, *Hack. in Mart. Fl. Bras.* ii. iii. 286.—Rio de Janeiro.

Goeringi, Steud. ex Flora, xxix. (1846) 22 = A. Nardus.

gracilipes, *Hack. in Flora*, lxviii. (1885) 120.—Paraguay.

ANDROPOGON :—

gracilis, J. Presl, in Rel. Haenk. i. 336 = tener.

gracilis, *Spreng Syst.* i. 284.—Ind. occ.

grandiflorus, *Hack. in Flora*, lxviii. (1885) 127.—Afr. trop.

grandis, Nees, ex Steud. Syn. Pl. Gram. 389 = A. Nardus.

Grayia, Steud. Nom. ed. II. i. 91 = pseudograya.

Griffithsiae, Steud. Syn. Pl. Gram. 375 = Ischaemum rugosum.

Grisebachii, Steud. l. c. 365 = Elionurus Royleanus.

Gryllus, Linn. Cent. Fl. ii. 32 = Chrysopogon Gryllus.

guianensis, Steud. Syn. Pl. Gram. 382 = Ischaemum guianense.

guineensis, Schum. Beskr. Guin. Pl. 51 = Gayanus.

Haenkei, J. Presl, in Rel. Haenk. i. 340 = intermedius.

halepensis, Brot. Fl. Lusit. i. 80 = Sorghum halepense.

Hallii, *Hack. in Sitzb. Acad. Wien*, lxxxix. (1884) 127. —Am. bor.

hamatulus, Hook. & Arn. Bot. Beech. Voy. 244 = A. Nardus

hamulatus, Nees, ex Steud Syn. Pl. Gram. 388 = A. Nardus.

hermaphroditus, Pourr. in Mém. Acad. Toul. iii. (1788) 308 = Stipa aristella.

heteranthus, *Steud. Syn. Pl. Gram.* 370.—Hab. ?

heteranthus, Trin. ex Steud. Nom. ed. II. i. 91 = bicornis.

heterolithus *Nees, Fl. Afr. Austr.* 115.—Ind. or.

hexastachyus, Steud. Syn. Pl. Gram. 380 = Pollinia argentea.

Hildebrandtii, *Hack. in Flora*, lxviii. (1885) 141.— Madag.

himalayensis, Steud. Syn. Pl. Gram. 377 = Apocopis Royleanus.

hirsutus, H. B. & K. Nov. Gen. et Sp. i. 186 = saccharoides.

hirtiflorus,*Kunth, Rev. Gram.* ii 560. t. 198.—Am. trop.

hirtifolius, *F. Presl, in Rel. Haenk* i. 338.—Mexico.

hirtsnoides, Kegel, in Act. Hort. Petrop. vii. (1880) 657 = Ischaemum hirtinodes.

hirtus, Linn. Sp. Pl. 1046 = Heteropogon hirtus.

hispidissimus, Steud. Syn Pl. Gram. 367 = Heteropogon contortus.

hispidus, Willd. Sp. Pl. iv. 908 = Arundinella brasiliensis.

Hochstetters, Steud. Syn. Pl. Gram. 384 = fastigiatus.

Hohenackeri, Hochst. ex Steud. l. c. 368 = Heteropogon contortus.

holcoides, Kunth, Rev. Gram. ii. 49. t. 160 = Saccharum holcoides.

Humboldtianus, Steud. Syn. Pl. Gram. 392 = Chrysopogon avenaceus.

humilis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 458.— Abyss.

humilis, Wright, ex Steud. Syn. Gram. i. 358 = demissus.

hypogynus, *Hack. in Mart. Fl. Bras.* ii. iii. 290.— Bras.

imberbis, *Hack. in Flora*, lxviii. (1885) 119.—Paraguay.

imberbis, Steud. Syn. Pl. Gram. 375 = Ischaemum aristatum.

importunus, *Steud. l. c.* 382.—China.

incertus, Steud. l. c. 375 = Ischaemum ciliare.

incompletus, J. Presl, in Rel. Haenk. i. 342 = Chrysopogon nutans.

increscens, Steud. Syn. Pl. Gram. 396 = Chrysopogon serrulatus

incurvatus, Koen. ex Trin. in Spreng. Neue Entdeck. ii. 91 = caricosus.

incurvatus, *Retz. Obs.* v. 21.—Ind. or.

inermis, Steud. Syn. Pl. Gram. 390 = spathiflorus.

infirmus, Steud. l. c. 389 = Ischaemum fragile.

inaequiglus,Anderss. in Schweiaf. Beitr. 1.1. Aethiop. 31C = Ischaemum laxum.

inaequiflus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 458 = pertusus.

insularis, Linn. Fl. Jam. Pugill. 30; Syst. ed. X. 1304 = Panicum leucophaeum.

Intermedius, *R. Br. Prod.* i. 202.—Orient; Afr. et As. trop.; Austral.

intonsus, Nees, Fl. Afr. Austr. 114 = lepidus.

intortum, Cranta, Inst. i. 398 = Sporobolus indicus.

intrans, *F. Muell. Census*, 120.—Austral.

inundatus, F. Muell. in Linnaea, xxv. (1852) 44 = intermedius.

involucratus, Koen. ex Steud. Nom. ed. II. i. 91 = Apluda aristata.

involutus, Steud. Syn. Pl. Gram. 373 = Spodiopogon angustifolius.

ANDROPOGON :—
Ischaemum, *Linn. Sp. Pl.* 1047.—Eur.; As. et Afr. trop. et temp.
Ischaemum, Thunb. Fl. Cap. i. 108 = appendiculatus.
ischyranthus, Steud. Syn. Pl. Gram. 367 = triticeus.
Iwarancusa, Bolss. Fl. Orient. v. 466 = A. Schoenanthus.
Iwarancusa, Nees, Fl. Afr. Austr. 117 = A. Nardus.
Iwarancusa, *Roxb. in Phil. Trans.* lxxx. (1790) 284.—Ind. or.; Afr. trop.
Jamesii, Torr. in Marc. Rep. (1852) 302 = saccharoides.
javanicus, *Mtq. Fl. Ind. Bat.* iii. 446.—Java.
javanicus, Steud. Syn. Pl. Gram. 396 = Chrysopogon aciculatus.
jubatus, Balansa, in Bull. Soc. Bot. Fr. xix. (1872) 322 = seriocus.
juncens, *Rich. ex Steud. Nom.* ed. II. i. 91.—Ins. S. Domingo.
junciformis, *Desv. in Ham. Prod. Fl. Ind. Occ.* 9.—Ins. S. Cruz.
Koenigii, Raeusch. Nom. ed. III. 293 = caricosus.
kolecotachys, *Steud. Syn. Pl. Gram.* 376.—Ins. Mascar.
korotrostachys, Trin. in Mém. Acad. Péterxb. Sér. VI. ii. (1833) 278 = Pollinia articulata.
lachnatherus, Benth. Fl. Austral. vii. 584 = filipendulus.
laguriformis, Griseb. in Goett. Abh. xxiv. (1879) 309 = saccharoides.
lagunoides, DC. Cat. Hort. Monsp. 78 = saccharoides.
lanatus, R. Br. Prod. i. 202 = exaltatus.
lancolatus, Baker, Fl. Maur. and Seych. 414 = Arthraxon cuspidatus.
lancolatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 262 = Arthraxon lancolatus.
lancifolius, Trin. in Mém. Acad. Péterxb. Sér. VI. ii. (1833) 271 = Arthraxon microphyllus.
lanigor, *Dref. Fl. Atland.* ii. 379.—Orient.; Ind. or.; Afr. hor.
lanigor, Munro, in Journ. Linn. Soc. xix. (1882) 409 = Spodiopogon lanigor.
lanuginosus, H. B. & K. Nov. Gen. et Sp. i. 187 = leucostachys.
lasiocolens, Steud. Syn. Pl. Gram. 383 = Arthraxon ciliaris.
lateralis, *Nees, Agrost. Bras.* ii. 320.—Am. trop.
latiflorus, Steud. Syn. Pl. Gram. 364 = Ellonurus latiflorus.
latifolius, Kunth, Rev. Gram. i. 166 = Centotheca lappacea.
latifolius, Spreng. Syst. i. 236 = Ischaemum latifolium.
laxiflorus, Steud. Syn. Pl. Gram. 380 = Trachypogon polymorphus.
laxus, Roxb. Hort. Beng. 7; Fl. Ind. ed. Carey, i. 271 = Sorghum halepense.
laxus, Willd. Sp. Pl. iv. 907 = serratus.
Lechleri, Steud. ex Hack. in Mart. Fl. Bras. ii. III. 297 = condensatus.
lepidus, *Nees, Fl. Afr. Austr.* 118.—Abyss.
leptanthus, Steud. Syn. Pl. Gram. 391 = intermedius.
leptoclados, *Hack. in Flora*, lxviii. (1885) 122.—Paraguay.
leptoconnus, *Trin. in Mém. Acad. Péterxb.* Sér. VI. ii. (1833) 264.—Sierra Leone.
leptophyllus, Trin. l. c. = tener.
leptus, Steud. Syn. Pl. Gram. 397 = tener.
leptostachyus, Benth. in Hook. Niger Fl. 571 = semiberbis.
Leschenaultianus, Decne. Herb. Timor. Descr. 29 = Pollinia Cumingii.
leucopogon, Nees, in Linnaea, xix. (1847) 694 = saccharoides.
leucostachys, *H. B. & K. Nov. Gen. et Sp.* i. 187.—Am. trop.
Lhotskyi, Steud. Syn. Pl. Gram. i. 384 = condensatus.
Hanatherus, Steud. l. c. 387 = Heteropogon insignis.
Lindmanni, *Hack. in Flora*, lxviii. (1885) 132.—Mexico.
Ligularis, Kunth, Enum. Pl. i. 497 = Trachypogon polymorphus.
Lindeni, Steud. Syn. Pl. Gram. 389 = apricus.
linearius, Steud. l. c. 389 = Ischaemum laxum.
lithophilus, *Trin. in Mém. Acad. Péterb.* Sér. VI. ii. (1833) 277.—Bras.
lividus, *Thw. Enum. Pl. Zeyl.* 367.—Ind. or.
lodiculatis, Steud. Syn. Pl. Gram. 375 = Ischaemum aristatum.
lolioides, Fourn. Mex. Pl. Enum. Gram. 62; ex Hemsl. Biol. Centr. Am. iii. 528 = scoparius.
longiberbis, *Hack. in Flora*, lxviii. (1885) 131.—Florida.

ANDROPOGON :—
longipes, *Hack. l. c.* 138.—Ind. or.
Loxiannus, Steud. Syn. Pl. Gram. 383 = virginicus.
lucer, Steud. l. c. 371.—' Mont. cerul.'
Lucrasi, Steud. l. c. 377 = Ischaemum ciliare.
macrogiossus, Steud. l. c. 368 = Trachypogon polymorphus.
macrolepis, *Hack. in Flora*, lxviii. (1885) 125.—Afr. trop.
macrostachyus, Anderss. in Schweinf. Beitr. Fl. Aethiop. 310 = Ischaemum laxum.
macrostachyus, Fourn. Mex. Pl. Enum. Gram. 60; ex Hemsl. Biol. Centr. Am. iii. 528 = Liebmanni.
macrothrix, Trin. in Mém. Acad. Péterxb. Sér. VI. ii. (1833) 270 = ternatus.
macrourus, *Michx. Fl. Bor. Am.* i. 56.—Am. bor.; Mexico.
madagascariensis, Hack. in Flora, lxviii. (1885) 136 = leucopogon.
malacophyllus, Hochst. ex Steud. Syn. Pl. Gram. 372 = Ischaemum ciliare.
malacostachyus, Fourn. Pl. Mex. Pl. Enum. Gram. 63; ex Hemsl. Biol. Centr. Am. iii. 528 = hirtiflorus.
malacostachyus, *Y. Presl, in Rel. Haenk.* i. 337.—Am. centr.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 282. —Fernando Po.
marginatus, Steud. in Flora, xii. (1829) 472 = A. Nardus.
Mariannus, Steud. Syn. Pl. Gram. 382 = Ischaemum digitatum.
maritimus, Chapm. Fl. S. Un. St. Suppl. 663 = scoparius.
Martini, Roxb. Hort. Beng. 7; Fl. Ind. i. 280 = A. Schoenanthus.
Martini, Thw. Enum. Pl. Zeyl. 367 = A. Nardus.
medius, Steud. Syn Pl. Gram. 382 = Ischaemum intermedium.
megapotamicus, Spreng. Syst. iv. Cur. Post. 33 = Trachypogon polymorphus.
melanocarpus, Ell. Sketch. i. 146 = Heteropogon acuminatus.
messanensis, Biv. ex Parl. Fl. Ital. i. 148 = Heteropogon contortus.
Meyerianus, Steud. Syn. Pl. Gram. 375 = Ischaemum aristatum.
micans, Steud. l. c. 382 = Arthraxon cuspidatus.
micranthus, Kunth, Rev. Gram. i. 165 = Chrysopogon violascens.
microphyllus Trin. in Mém Acad. Péterb. Sér. VI. ii. (1833) 275 = Arthraxon microphyllus.
microstachyus, *Desv. in Ham. Prod. Fl. Ind. Occ.* 8. —Ind. occ.
millaceus, *Roxb. Hort. Beng.* 7; *Fl. Ind.* i. 276.—Ind. or.
millefornis, Schult. Mant. ii. 448 = prsec
Minarum, Kunth, Enum. Pl. i. 507 = Chrysopogon stipoides.
minor, Steud. Syn. Pl. Gram. 377 = Ischaemum Urvillianum.
minutiflorus, Steud. l. c. 379 = Pollinia argentea.
mollicomus, Kunth, Rev. Gram. i. 365. t. 94 = caricosus.
mollis, Kunth, l. c. ii. 561. t. 195 = Trachypogon polymorphus.
monandrus, Roxb. Hort. Beng. 82; Fl. Ind. i. 264 = Pogonatherum saccharoideum.
monatherus, *A. Rich. Tent Fl. Abyss.* ii 462.—Abyss.
monostachys, Spreng. Pugill. ii. 9 = foveolatus.
monostachyus, Steud. Syn. Pl. Gram. 369 = Ischaemum aristatum.
montanus, Koen. ex Trin. in Spreng. Neue Entdeck. ii. 98 = Chrysopogon montanus.
montanus, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 138 = micranthes.
montanus, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 271 = Chrysopogon pictus.
monticolus, Schult. Mant. iii. 665 = Chrysopogon montanus.
Montufari, H. B. & K. Nov. Gen. et Sp. i. 184 = Trachypogon polymorphus.
Muehlenbergianus, Schult. Mant. ii. 455 = argyreus.
multicaulis, Steud. Syn. Pl. Gram. 388 = Arthraxon microphyllus.
multinervis, Hochst. ex Steud. l. c. 374 = polystachyus.
muricatus, Retz. Obs. iii. 48 = squarrosus.
murinus, Steud. Syn. Pl. Gram. 377 = Ischaemum murinum.
muticus, Linn. Sp. Pl. ed. II. 1482 = Chloris petraea ?
muticus, Steud. Syn. Pl. Gram. 374 = Ischaemum muticum.

ANDROPOGON :—
Myosurus, Hack. in Mart. Fl. Bras. ii. III. 300 = hirtiflorus.
Myosurus, *Y. Presl, in Rel. Haenk.* i. 337.—Mexico.
nardoides, Nees, Fl. Afr. Austr. 116 = A. Schoenanthus.
Nardus, *Linn. Sp. Pl.* 1046.—Reg. trop.
Nardus, Trin. in Spreng. Neue Entdeck. ii. 90 = A. Schoenanthus.
Nees, *Kunth, Enum. Pl.* i. 491.—Bras.
Neesii, Trin. in Mém. Acad. Péterxb. Sér. VI. ii. (1833) 263 = tener.
nemophilus, Steud. Syn. Pl. Gram. 377 = Ischaemum filiforme.
nervatus, Hochst. in Flora, xxvii. (1844) 243 = A. Schoenanthus.
nervosus, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1802) 218 = Ischaemum laxum.
Newtonii, *Hack. in Bol. Soc. Bruter.* iii. (1885) 137.—Angola.
nigor, Kunth, Rev. Gram. i. 164 = Sorghum vulgare.
nigritanus, Benth. in Hook. Niger Fl. 573 = squarrosus.
nitens, Schreb. ex Steud. Nom. ed. II. i. 92 = serratus.
nitidus, Kunth, Rev. Gram. i. 166 = serratus.
nodiculatus, Hochst. ex Steud. Syn. Pl. Gram. 396 = Chrysopogon Arnottianus.
noduliosus, *Hack. in Flora*, lxviii. (1885) 116.—Afr. trop.
notopogon, Steud. Syn. Pl. Gram. 373 = Spodiopogon angustifolius.
nudus, Steud. l. c. 383 = Arthraxon ciliaris.
Nuttallii, Chapm. Fl. S. Un. St. 580 = Elionurus tripsacoides.
nutans, *Linn. Mant.* ii. 303 = Anthistiria ciliata.
nutans, Linn. Sp. Pl. 1045 = Chrysopogon nutans.
nutans, Muhl. Descr. Uber. 2:4 = avenaceus.
obliquiberbis, *Hack. in Flora*, lxviii. (1885) 117.—N. Caledon.
obtusiflora, Poir. Encyc. Suppl. i. 583 = brevifolius.
obtusus, Nees, in Hook. & Arn. Bot. Beech. Voy. 243 = annulatus.
obvallatus, Steud. Syn. Pl. Gram. 378 = Spodiopogon angustifolius.
odoratus, Auct. ex Steud. Nom. ed. II. i. 92 = squarrosus.
oliganthus, *Hochst. ex Steud. Syn. Pl. Gram.* 368.—Ind. or.
oligostachyus, Chapm. Fl. S. Un. St. 581 = hirtiflorus.
Oliveri, Boiss. Diagn Ser. I. v. 76 = laniger.
orthos, Schult. Mant. ii. 455 = foveolatus.
pachnodes, Trin. in Mém. Acad. Péterxb. Sér. VI. ii. (1833) 264 = A. Schoenanthus.
paleaceus, Steud. Syn. Pl. Gram. 376 = Apocopis Royleanus.
pallidus, Kunth, Rev. Gram. i. 165 = Chrysopogon Gryllus.
panicus, Steud. Syn. Pl. Gram. i. 375 = Ischaemum intermedium.
paniculatus, Kunth, Enum. Pl. i. 494 = condensatus.
paniculatus, Lam. Fl. Fr. iii. 688 = Chrysopogon Gryllus.
papernellanus, Parl Fl. Ital. i. 140 = pertusus.
papillipes, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 460. —Abyss.
papillosus, *Hochst. ex A. Rich. l. c.* 457.—Afr. trop.
parviflorus, Roxb. Hort. Beng. 82; Fl. Ind. i. 277 = brevifolius.
parviglcis, Steud. Syn. Pl. Gram. 397 = Chrysopogon violascens.
patentiviliosus, Steud. l. c. 373 = Ischaemum ciliare.
patulus, Mœnch, Meth. 180 = A. Ischaemum.
pauciflorus, Salzm. ex Steud. Syn. Pl. Gram. 371 = virginicus.
pectinatus, *Steud. l. c.* 389 = Ischaemum pectinatum.
pedicellatus, Steud. l. c. 394 = serratus.
pedunculatus, *Kunth, Rev. Gram.* i. 166.—Ind. or.
pedunculatus, Baker, Fl. Maur. 444 = A. Alopecurus.
pendulus, Nees, ex Steud. Syn. Pl. Gram. 368 = A. Nardus.
peninsulae, Steud. l. c. 396 = Chrysopogon Arnottianus.
pennatus, Willd. ex Steud. Nom. ed. II. i. 92 = laxiginosus.
perforatus, Trin. ex Fourn. Mex. Pl. Enum. Gram. 59, ex Hemsl. Biol. Centr. Am. iii. 528 = saccharoides.

ANDROPOGON :—

perfoliatus, Nees & Meyen, ex Steud. Nom. ed. II. i. 92 = internodius.

pertusus, *Willd. Sp. Pl.* iv. 922.—Sicil. ; As. et Afr. trop. ; Austral.

petiolaris, Steud. Syn. Pl. Gram. 398 = Pollinia Lehmanni.

Petitianus, Balansa, in Bull. Soc. Bot. Fr. xix. (1872) 292 = obliquiberbis.

Petitianus, A. Rich. Tent. Fl. Abyss. ii. 451 = exilis.

pilifer, Steud. Syn. Pl. Gram. 378 = pilosus.

pilosus, *Klein, ex Willd. Sp. Pl.* iv. 920.—Ind. or.

pilosus, Sieber, ex Kunth, Enum. Pl. i. 514 = Ischaemum latifolium.

piptatherus, *Hack. in Mart. Fl. Bras.* ii. III. 293.—Bras.

plagiolepis, Hochst. ex Steud. Syn. Pl. Gram. 374 = polyatherus.

platypus, Trin. in Mém. Acad Pétersb. Sér. VI. ii. (1838) 361 = Elionurus platypus.

plumosus, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 918 = Trachypogon polymorphus.

podotrichus, Hochst. ex Steud. Syn. Pl. Gram. 384 = Heteropogon hirtus.

poecilotrichus, *Hack. in Bol. Soc. Broter.* iii. (1885) 188.—Afr. trop.

pogonanthus, Boiss. & Bal. in Bull. Soc. Bot. Fr. v. (1858) 168 = Spodiopogon pogonanthus.

Pohlianus, *Hack. in Mart. Fl. Bras.* ii. III. 304.—Bras.

polyatherus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 455.—Abyss.

polydactylon, Linn. Amoen. Acad. v. 412 ; Sp. Pl. ed. II. 1488 = Chloris polydactyla.

polymorphus, Steud. Syn. Pl. Gram. 375 = Ischaemum muticum.

polyneurus, Steud. l. c. 385 = A. Schoenanthus.

polypsychios, *Steud. l. c.* 380.—Zeylan.

polystachyus, Roxb. Hort. Beng. 82 ; Fl. Ind. i. 265 = Heteropogon contortus.

polystichus, Hochst. ex Steud. Syn. Pl. Gram. 369 = Heteropogon acuminatus.

Preslii, Kunth, Enum. Pl. i. 489 = tener.

Princeps, A. Rich. Tent. Fl. Abyss. ii. 470 = Thelepogon elegans.

prionodes, Steud. Syn. Pl. Gram. 383 = Arthraxon lanceolatus

procerus, F. Muell. Fragm. viii. 124 = filipendulus.

procerus, *R. Br. Prod.* i. 202.—Austral.

productus, Regel, in Bull. Acad. Pétersb. x. (1866) 378 = Pollinia tenuis.

propinquus, Kunth, Enum. Pl. i. 502 = Sorghum halepense.

prostratus, Nees, ex Steud. Syn. Pl. Gram. 384 = Anthistiria prostrata.

provincialis, Retz. Obs. iii. 48 = Chloris radiata.

provincialis, *Lam. Encyc.* i. 376.—Europ. ; Am. bor.

pruinosus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 464 = A. Iwarancusa.

pruinosus, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Schoenanthus.

pseudogryza, *Steud. l. c.* 365.—Zeylan. ; China ; Madag.

pseudohirtus, Steud. in Flora, xli. (1829) 471 = A. Nardus.

pseudoischaemum, *Nees, ex Steud. Syn. Pl. Gram.* 380.—Ind. or.

pubescens, [Dryand. in] Ait. Hort. Kew, ed. I. iii. 423 = Chloris ciliata.

pubescens, Vitman, in Flora, xii. (1829) 1. Erg. 3 = Heteropogon hirtus.

pubescens, Willd. ex Kunth, Enum. Pl. i. 498 = mollicomis.

pubiflorus, Fourn. Mex. Fl. Enum. Gram. 57 ; ex Hemsl. Biol. Centr. Am. iii. 529 = hirtiflorus.

pulchellus, *D. Don, in Hook. Niger Fl.* 571.—Afr. trop.

pumilus, *Roxb. Hort. Beng.* 7 ; *Fl. Ind.* i. 277.—Ind. or.

punctatus, Roxb. ll. cc. 92, 266 = intermedius.

purpurascens, Muhl. ex Willd. Sp. Pl. iv. 913 = scoparius.

purpureo-sericeus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 469.—Abyss.

pusillus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 283.—Afr. trop.

pusillus, *Lam. Syst. ed.* XIII. 758 = Anthistiria ciliata.

ANDROPOGON :—

Quartinianus, *A. Rich. Tent. Fl. Abyss.* ii. 469.—Abyss.

quinqueplumis, Hochst. ex A. Rich. l. c. 450 = Chrysopogon serrulatus.

radicans, Lehm. Ind. Hort. Hamb. (1826) ; ex Kunth, Enum. Pl. i. 499 = A. Ischaemum.

racemosus, Poir. ex Steud. Nom. ed. I. 412, in syn. = ramosus.

ramosus, Blanco, Fl. Filip. ed. I. 37 = Ischaemum ciliare.

ramosus, *Forsk. Fl. Aegypt. Arab.* 173.—Aegypt. ; Arab.

Ravennae, Linn. Sp. Pl. ed. II. 1481 = Erianthus Ravennae.

reconditus, Steud. Syn. Pl. Gram. 386 = Gayanus.

rectibaccha, *Fourn. Mex. Pl. Enum. Gram.* 61 ; *ex Hemsl. Biol. Centr Am.* iii. 529.—Mexico.

rectus, Steud. Syn. Pl. Gram. 389 = A. Nardus.

refractus, *R. Br. Prod.* i. 202.—Ins. Molucc. et Pacific. ; Austral.

relictus, Steud. Syn. Pl. Gram. 377 = Ischaemum muticum.

repens, Steud. l. c. 374 = Ischaemum muticum.

Retzii, Steud. l. c. 375 = Ischaemum ciliare.

rhizophorus, Steud. l. c. 381 = Spodiopogon albidus.

Riedelii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 363.—Am. austr.

Robertianus, Steud. Syn. Pl. Gram. 369 = Ischaemum laxum.

rostratus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1838) 361 = Elionurus rostratus.

rottboellioides, Steud. Syn. Pl. Gram. 382 = Rottboellia rottboelloides.

Roxburghianus, Schult. Mant. ii. 451 = Dimeria ornithopoda.

Roxburghii, Nees, ex Steud. Syn. Pl. Gram. 395 = A. Nardus.

Royleanus, Steud. l. c. 397 = Chrysopogon Gryllus.

rubens, Kunth, Rev. Gram. i. 164 = Sorghum vulgare.

rudis, Steud. Syn. Pl. Gram. 383 = Arthraxon rudis.

rufidulus, Steud. l. c. 392 = Chrysopogon avenaceus.

rufispica, *Steud. l. c.* 379.—Java.

rufus, *Kunth, Enum. Pl. i.* 497.—Afr. trop. ; Bras.

rufus, Kuth. ex Kunth, l. c. 478 = Saccharum cayennense.

rugosus, Steud. Syn. Pl. Gram. 376 = Ischaemum rugosum.

Ruprechtii, *Hack. in Flora*, lxvii. (1885) 126.—Afr. trop. ; Mexico.

saccharatus, Kunth, Enum. Pl. i. 502 = Sorghum vulgare.

saccharatus, Roxb. Hort. Beng. 7 ; Fl. Ind. i. 274 = Sorghum vulgare.

saccharoides, *Sw. Prod. Veg. Ind. Occ.* 26.—Am. temp. et trop.

sanguinarius, *Schreb. Beschr Grass.* ii. 48.—Afr. bor.

scabriusculus, Kunth, Enum. Pl. i. 504 = Chrysopogon nutans.

scabriflorus, *Rupr. ex Hack. in Mart. Fl. Bras.* ii. III. 289.—Bras. ; Paraguay.

scandens, Roxb. Hort. Beng. 7 ; Fl. Ind. i. 262 = annulatus.

Schaffneri, Griseb. ex Fourn. Mex. Fl. Enum. Gram. 57 ; ex Hemsl. Biol. Centr. Am. iii. 529 = hirtiflorus.

schangulensis, Rupr. ex Steud. Syn. Pl. Gram. 389 = Ischaemum laxum.

Schimperi, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 466.—Abyss.

schirensis, *Hochst. l. c.* 456.—Abyss.

Schlumbergeri, *Fourn. Mex. Pl. Enum. Gram.* 59, *ex Hemsl. Biol. Centr. Am.* iii. 529.—Mexico.

Schoenanthus, Benth. Fl. Austral. vii. 534 = A. Nardus.

Schoenanthus, *Linn. Sp. Pl.* 1046.—As. et Afr. trop. et subtrop.

Schoenanthus, Thunb. Fl. Cap. i. 108 ; Wall. Pl. As. Rar. i. 80 = A. Nardus.

Schottii, *Rupr. ex Hack. in Mart. Fl. Bras.* ii. III. 290.—Am. trop.

Schweinfurthii, *Hack. in Flora*, lxvii. (1885) 118.—Afr. trop.

scoparius, *Michx. Fl. Bor. Am.* i. 57.—Am. bor. ; Mexico.

scoparius, J. Presl, in Rel. Haenk. i. 338 = condensatus.

scrobiculatus, Kunth, Enum. Pl. i. 507 = Heteropogon acuminatus.

ANDROPOGON :—

secundus, Kunth, Enum. Pl. i. 487 = Trachypogon polymorphus.

secundus, Willd. ex Nees, Agrost. Bras. 364 = Heteropogon contortus.

segetumensis, Steud. Syn. Pl. Gram. 367 = Heteropogon acuminatus.

segetum, Steud. l. c. 376 = Ischaemum rugosum.

Sehima, Steud. l. c. 369 = Ischaemum laxum.

semiberbis, *Kunth, Enum. Pl.* i. 489.—Afr. et Am. trop.

semisagittatus, Steud. Syn. Pl. Gram. 376 = Ischaemum semisagittatum.

seneurensis, Hochst. in Flora, xxvii. (1844) 248 = A. Iwarancusa.

sericeus, *R. Br. Prod.* i. 201.—Malaya ; Austral.

serratus, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 290 = Chrysopogon violaceus.

serratus, Retz. Obs. v. 21 = caricosus.

serratus, *Thunb. Fl. Jap.* 41.—As. et Afr. trop. ; Austral.

serrulatus, Link, Enum. Hort. Berol. i. 241 = Arthraxon lanceolatus.

setifolius, *Kunth, Rev. Gram.* i. 166.—Afr. austr.

setosus, Griseb. Cat. Pl. Cub. 235 = Chrysopogon avenaceus.

sibiricus, Steud. Syn. Pl. Gram. 398 = Spodiopogon sibiricus.

similimus, Steud. l. c. 373 = Ischaemum ciliare.

simplex, *Schum. Beskr. Guin. Pl.* 49.—Afr. trop.

sinaicus, Del. Ind. Sem. Hort. Monsp. (1836) ; et in Ann. Sc. Nat. Sér. II. vii. (1837) 255 = Heteropogon hirtus.

Smithianus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 242.—Afr. trop.

Sorghum, Brot. Fl. Lusit. i. 88 = Sorghum vulgare.

spathaceus, Trin. Fund. Agrost. 186 = macrourus.

spathiflorus, *Kunth, Enum. Pl.* i. 496.—Bras.

speciosus, Hort. Goett. ex Steud. Nom. ed. II. i. 93 = saccharoides.

speciosus, Steud. Syn. Pl. Gram. 375 = Rottboellia speciosa.

spicatus, Steud. Nom. ed. II. i. 93 = Trachypogon polymorphus.

Sprengelii, Kunth, Rev. Gram. i. 166 = Chrysopogon montanus.

squamulatus, Hochst. in Flora, xxvii. (1844) 244 = Gayanus.

squarrosus, Koen. ex Kunth, Enum. Pl. i. 126 = Panicum abortivum et paludosum.

squarrosus, *Linn. f. Suppl.* 493.—Kap. trop.

stenophyllus, Roem. & Schult. Syst. ii. 819 = Trachypogon angustifolius.

stipaeformis, Steud. Syn. Pl. Gram. 377 = Dimeria ornithopoda.

stipoides, *Cham. & Schlecht. in Linnaea*, vi. (1831) 89 = Chrysopogon Francavillanus.

stipoides, H. B. & K. Nov. Gen. et Sp. i. 139 = Chrysopogon avenaceus.

stipoides, Kunth, Enum. Pl. i. 487 = Trachypogon polymorphus.

stipoides, Steud. Syn. Pl. Gram. 368 = Trachypogon polymorphus.

stipoides, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 363 = Chrysopogon avenaceus.

striatus, Klein, ex Willd. Sp. Pl. iv. 908 = Ischaemum laxum.

strictus, Host, Gram. ii. 2. t. 2 = Erianthus Hostii.

strictus, Roxb. Hort. Beng. 82 ; Fl. Ind. i. 266 = foveolatus.

subglabrescens, Steud. Syn. Pl. Gram. 393 = Sorghum vulgare.

submembranaceus, *Steud. l. c.* 381.—Ins. Mascar.

submuticus, Steud. l. c. 382 = Arthraxon ciliaris.

subsquarrus, Steud. l. c. 397 = montanus.

subtilis, Steud. l. c. 396 = Chrysopogon subtilis.

subulatus, J. Presl, in Rel. Haenk. i. 341 = Chrysopogon aciculatus.

sudanifloccidatus, Steud. Syn. Pl. Gram. 380 = pertusus.

supercilliatus, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 239.—Austral. ; Timor.

Tabina, Steud. in Lechl. Berb. Am. Austr. 56 = hirtiflorus.

tacazensis, Steud. Syn. Pl. Gram. 369 = Ischaemum laxum.

tahitensis, Hook. & Arn. Bot. Beech. Voy. 72 = refractus.

ANDROPOGON :—
Tamba, Hochst. ex Steud. Syn. Pl. Gram. 385 = lepidus.
tectorum, *Schum. & Thonn. Beskr. Guin. Pl.* 49.— Guinea.
tenellus, J. Presl, in Rel. Haenk. i. 335 = brevifolius.
tenellus, *Roxb. Hort. Beng.* 6; *Fl. Ind.* i. 259.— Bengal.
tener, *Kunth, Rev. Gram.* ii. 505. t. 197.—Am. bor. et austr.
tenuiculus, Steud. Syn. Pl. Gram. 371 = sericeus.
tenuifolius, Steud. l. c. 365 = Ellonurus argenteus.
tenuispedicellatus, Steud. l. c. 398 = australis ?
tenuiarchia, Fourn. Mex. Pl. Enum. Gram. 5b ; ex Hemsl. Biol. Centr. Am. iii. 529 = saccharoides.
tenuis, R. Br. Prod. i. 201 = Heteropogon contortus.
ternarius, Bertol. Bot. Misc. xi 14 = provincialis.
ternarius, Desv. Opusc. 88 = squarrosus.
ternarius, Michx. Fl. Bor. Am. i. 57 = provincialis ?
tematus, *Nees, Agrost. Bras.* ii. 326.—Am. austr.
tetrastachyus, Ell. Sketch, i. 100 = virginicus.
Thunbergii, Kunth, Rev. Gram. i. 166 = Danthonia livida.
thymioides, Steud. Syn. Pl. Gram. 365 = Ellonurus thymoides.
timorensis, Steud. l. c. 376 = Spodiopogon Blumii.
tomentellus, Steud. l. c. 371 = Guyanus ?
Tong-Dong, Steud. l. c. 378 = Ischaemum rugosum.
Torreyanus, Steud. Nom. ed. II. i. 93 = saccharoides.
tortus, Willd. ex Steud. Nom. ed. II. i. 92 = Trachypogon polymorphus.
trachypus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 280 = bracteatus.
trichozygus, *Baker, in Journ. Linn. Soc.* xx. (1884) 306.—Madag.
tridentatus, Hochst. in Flora, xxvii. (1844) 246 = Gayanus
tridentatus, Roxb. Hort. Beng. 82 ; Fl. Ind. i. 261 = Apocopis tridentata.
Trinii, Steud. Syn. Pl. Gram. 395 = Chrysopogon serrulatus.
trispaccoides, Trin. ex Steud. Nom. ed. II. i. 92 ; et Syn. Pl. Gram. 364 = Ellonurus trispaccoides.
trispicatus, Schult. Mant. ii. 452 = Pollinia argentea.
tristachyus, Roxb. Hort. Beng. 6 ; Fl. Ind. i. 261 = Pollinia argentea.
tristachyus, H. B. & K. Nov. Gen. et Sp. i. 187 = glaucescens.
triticeus, R. Br. Prod. i. 201 = Heteropogon insignis.
triticiformis, Steud. Syn. Pl. Gram. 376 = Ischaemum triticeum.
tropicus, Kunth, Enum. Pl. i. 503 = australis.
tropicus, Spreng. Syst. i. 287 = serratus.
truchmenorum, Walp. Ann. iii. 801 = Sorghum vulgare.
truncatus, Steud. l. c. 368 = Trachypogon polymorphus.
tumbuckianus, *Rendle, ex Kunth, Enum. Pl.* i. 507, *in syn.*—Afr. austr.
tumbuchannus, Steud. Nom. ed. II. i. 93 = prasc.
umbrosus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 467 = lepidus.
undatus, Jacq. Coll. iii. 287 = Oplismenus Jacquini.
undulatus, Pers. Syn. i. 105 = annulatus.
urceolatus, *Hack. in Flora*, lxviii. (1885) 115.—Afr. centr.
Urvilleanus, Steud. Syn. Fl. Gram. 376 = Ischaemum Urvilleanum.
usurum, Steud. l. c. 392 = Sorghum vulgare.
Vachellii, Nees. ex Hook. & Arn. Bot. Beech. Voy. 343 = intermedius.
vaginatus, Ell. Sketch, 148 = virginicus.
vaginatus, J. Presl, in Rel. Haenk. i. 336 = semiberbis.
velatus, Kunth, Enum. Pl. i. 488 = semiberbis.
venustus, *Thw. Enum. Pl. Zeyl.* 367.—Zeylan.
versicolor, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Schoenanthus.
verticillatus, Boj. Hort. Maurit. i. 57 = squarrosus.
verticillatus, Roxb. Hort. Beng. 82 ; Fl. Ind. i. 267 = Chrysopogon verticillatus.
verticillatus, Schum. Beskr. Guin. Pl. 50 = fulvibarbis.
verticilliflorus, Steud. Syn. Pl. Gram. 393 = Sorghum halepense ?
vestitus, *Steud. l. c.* 376.—Bras.
villiferus, Steud. l. c. 376 = Ischaemum australe.
villosulus, Steud. l. c. 397 = Chrysopogon violaceus.
villosus, Gilib. Exercit. ii. 522 = A. Ischaemum.

ANDROPOGON :—
villosus, Lam. Fl. Fr. iii. 634 = A. Ischaemum et provincialis.
villosus, Thunb. Prod. Pl. Cap. 20 = Pollinia villosa.
vimmosus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 268 = Pollinia imberbis.
violaceus, Heyne, ex Steud. Syn. Pl. Gram. 414 = Arthraxon ciliaris.
violascens, Nees, ex Steud. l. c. 396 = Chrysopogon violascens.
virens, Spreng. Syst. i. 287 = Arundinella brasiliensis.
virgatus, Desv. in Ham. Prod. Ind. Occ. 9 = spathiflorus.
virginicus, *Linn. Sp. Pl.* 1046.—Am. bor.
virginicus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 269 = leucostachyus.
Wightianus, Steud. Syn. Pl. Gram. 395 = Chrysopogon Wightianus.
Wrightii, *Hack. in Flora*, lxviii. (1885) 139.—Am. bor.
xanthoblephara, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 281.—Afr. trop.
Wrightii, *Munro, ex Wright, in Sauv. Fl. Cub.* 201. —Cuba.
Zeae, *Rosm. & Schult. Syst.* ii. 818.—N. Granat.
zeylanicus, Am. ex Benth. Fl. Hongk. 428 = A. pseudograya.
zeylanicus, Steud. Syn. Pl. Gram. 397 = Chrysopogon Arnottianus.
Zollingeri, *Steud. l. c.* 349.—Java.

ANDROSACE, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 52 (1737). *PRIMULACEÆ*, Benth. & Hook. f. ii. 632.
AMADEA, Adans. Fam. ii. 230 (1763).
ARETIA, Linn. Sp. Pl. ed. 1. 141 (1753).
acaulis, Hort. ex Duby, in DC. Prod. viii. 52 = septentrionalis.
acaulis, *Cirillo, ex Steud. Nom. ed. II. i. 93 = A. Chamaejasme.
affinis, Biroli, in Brugnat. Giorn. Fis. II. iii. (1820) 143. t. 6 = obtusifolia.
Aitoom, *Duby, in DC. Prod.* viii. 50.—Reg. Himal. ; China.
alba, *Duby, l. c.* 55.—Am. bor.
albana, *Stev. in Trans. Linn. Soc.* xi. (1815) 410. t. 34.—Ax. Min. ; Caucas.
alismoides, Hornem. Hort. Hafn. i. 185 = lactiflora.
alpina, *Lam. Fl. Fr.* iii. 642.—Europ.
altaica, *C. Koch, in Linnaea*, xxiii. (1850) 612.—Reg. altaic.
apennina, Huet, ex Nym. Consp 607 = alpina.
aquatica, [Clairv.] Man. Herb. 56 = Hottonia palustris.
arachnoidea, Schott, Nyman & Kotschy, Anal. Bot. 17 = villosa.
azeica, *Chow. in Linnaea*, i. (1826) 230.—Am. arct.
Aretia, D. Dietr. Syn. i. 632 = bryoides.
Aretia, Lap. Hist. Abr. Pl. Pyr. 91 = imbricata.
Aretia, *Vill. Hist. Pl. Dauph.* ii. 469.—Europ.
arctioides, Herr, ex Duby, in DC. Prod. viii. 51 = obtusifolia.
arctioides x, *Kerner, in Oestr. Bot. Zeitschr.* xxv. (1875) 159.—Europ.
argentea, Lap. Hist. Abr. Pl. Pyr. 92 = imbricata.
arizonica, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 231.—Am. bor.
armenica, *Duby, in DC. Prod.* viii. 52.—Ax. Min.
articulata, *Schur, Enum. Pl. Transs.* 551.—Europ.
baikalensis, Fisch. ex Ledeb. Fl. Ross. iii. 19 = lactiflora.
baikalensis, Schlecht. ex Roem. & Schult. Syst. iv. 736 = ochotensis.
bidentata, C. Koch, in Linnaea, xxiii. (1850) 610 = multiscapa.
Bocconi, Huet, ex Homem. Hort. Hafn. i. 184 = elongata.
brevifolia, Vill. Hist. Pl. Dauph. ii. 480 = septentrionalis.
brigantiaca, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 106 = carnea.
Bruegger i x, *Jaeggi, ex Brugg. in Jahresb. Naturf. Ges. Graub. Neue Fol.* xxii. (1879) p. xxix.—Helvet.
bryoides, *DC. Fl. Fr.* iii. 440.—Helvet.
campicos, *Lehm. ex Spreng. in Isis*, (1817) 1286. n 162. t. 9.—Persia.
cana, *Willd. ex Steud. Nom.* ed. II. i. 93.—Am. austr.
capitata, Willd. ex Roem. & Schult. Syst. iv. 736 = A. Chamaejasme.

ANDROSACE :—
carinata, Torr. in Ann. Lyc. Nat. Hist. New York, i. (1824) 80 = N. Chamaejasme.
carnea, *Linn. Sp. Pl.* 142.—Europ.
carnoulia, Duby, in DC. Prod. viii. 54 = saxifragaefolia.
Chaixi, *Gren. & Godr. Fl. Fr.* ii. 458.—Europ. ; Dahuria ?
Chamaejasme, Bieb. Fl. Taur. Cauc. iii 182 = villosa.
Chamaejasme, *Willd. Sp. Pl.* i. 790.—Europ. ; Ax. Min. ; Ax. bor.
Chamaejasme, Wulf. in Jacq. Coll. i. 194 = obtusifolia.
Charpentieri, *Heer, in Moritzi, Graub. Vers.* 113.— Helvet.
ciliata, *DC. Fl. Fr.* iii. 44 b.—Europ.
commutata, *Schlecht. in Bot. Zeit.* xiv. (1856) 521.— Ax. temp.
cordifolia, *Wall. in As. Res.* xiii. (1820) 371.—Reg. Himal.
coronopifolia, Andr. Bot. Repos. t. 647 = lactiflora.
Crotii, *Watt, in Journ. Linn. Soc.* xx. (1884) 17.— Reg. Himal.
cylindrica, *DC. Fl. Fr.* iii. 439.—Mont. Pyren.
dasyphylla, Bange, in Ledeb. Fl. Alt. i. 218 = villosa.
Diapensia, Vill. Hist. Pl. Dauph. ii. 472 = bryoides.
diapensioides, Lap. Hist. Abr. Pl. Pyr. 93 = pyrenaica.
elegans, Duby, in DC. Prod. viii. 55 = rotundifolia.
elongata, *Linn. Sp. Pl.* ed. II. *App.* 1668.—Europ. ; Caucas.
elongata, Richards. App. Frankl. 1st Voy. 733 = septentrionalis.
elongata, Pall. ex Ledeb. Fl. Ross. iii : 21 = filiformis.
Escheri x, *Brugg. in Jahresb Naturf. Ges. Graub. Neue Fol.* xxiii-xxiv. (1880) 100.—Helvet.
farinosa, Spreng. Pugill. ii. 37 = Primula farinosa.
fasciculata, Willd. ex Roem. & Schult. Syst. iv. 736 = septentrionalis.
filiformis, *Retz Obs.* ii. 10.—Persia ; As. bor. ; Am. bor.
foliosa, Duby, ex Deene. in Jacquem. Voy. Bot. 142. t. 146 = sarmentosa.
foliosa, Klatt, in Linnaea, xxxii. (1863) 208 = strigilosa.
Friesii *Traute. in Act. Hort. Petrop.* ix. (1884) 7.— N. Zemlaya.
frutescens, Lap. Hist. Abr. Pl. Pyr. 92 = cylindrica.
geranifolia, *Watt, in Journ. Linn. Soc.* xx. (1882) 17. —Reg. Himal.
glacialis, Hoppe, in Koch, Syn. ed. II. iii 670 = alpina.
globifera, *Duby, in DC. Prod.* viii. 48.—Reg. Himal.
globifera, Klatt, in Linnaea, xxxii. (1863) 292 = microphylla.
globifera, Schur, Enum. Pl. Transs. 551 = villosa.
Gmelini, *Gaert. Fruct.* i. 232.—Sibir.
Halleri, [Hoock.] Vers. Gew. Teutschl. 561 = carnea.
Halleri, *Linn. Sp. Pl.* 142.—Europ.
Hausmanni, *Leyb. in Flora*, xxxv. (1852) 401.—Austria.
hedraeantha, *Griseb. Spicil. Fl. Rumel.* ii. 3.—Thracia.
Heerii, Hegetschw. ex Hausm. Fl. Tirol. iii. 1474 = Hausmanni.
Heerii, Koch, Syn. Fl. Germ. ed. II. 670 = bryoides.
helvetica, All. Fl. Pedem. i. 91 = bryoides.
helvetica, L. Dufour, in Act. Soc. Linn. Bord. viii. (1836) 100 = pubescens.
Hookeriana, *Klatt, in Linnaea*, xxxii. (1863) 293.— Reg. Himal.
hybrida x, *Kerner, in Oestr. Bot. Zeitschr.* xxv. (1875) 159.—Europ.
imbricata, *Lam. Fl. Fr.* ii. 353.—Europ.
incana, Lam. Illustr. i. 432 = villosa.
incisa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 16 = rotundifolia.
intermedia, Ledeb. Fl. Ross. iii. 20 = armenica.
Jacquemontii, Duby, in DC. Prod. viii. 50 = villosa.
Lackenalli, C. C. Gmel. Fl. Bad. i. 2 = carnea.
lactea, *Linn. Sp. Pl.* 142.—Europ.
lactea, Vill. Hist. Pl. Dauph. ii. 476 = obtusifolia.
lactea, Pall. Reise, ii. 718 = septentrionalis.
lactiflora, [*Fisch.*] *Hort. Gorenk.* ed. I. (1808) 80.— Sibir.
lactiflora, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 738. n. 717 = septentrionalis.
Laggeri, *Huet, in Ann. Sc. Nat. Sér. III.* xix. (1853) 224.—Europ.
lanuginosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 15.— Reg. Himal.
Lehmanni, *Wall. Cat.* n. 617.—Reg. Himal.
Lehmanniana, Spreng. in Isis, (1817) 1289 = A. Chamaejasme.
Linnæi, R. Grah. in Edinb. N. Phil. Journ. (Apr.-June, 1829) 179 = septentrionalis.

ANDROSACE:—
longifolia, C. Koch, in Linnæa, xxiii. (1850) 610 = albana.
longifolia, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1892) 209.—Mongolia.
longiscapa, C. Koch, in Linnæa, xxiii. (1850) 611 = A. Chamæjasme.
longiseta, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 61 = obtusifolia.
lutea, Lam. Fl. Fr. ii. 288 = Douglasia Vitaliana.
macrantha, *Beiss. & Huet, Diagn.* Ser. II. iii. 119.—Armen.
macrocarpa, *Ledeb. ex Duby, in DC. Prod.* viii. 55, nomen.—Sibir.
Mathildæ, Levier, in Nuov. Giorn. Bot. ix. (1877) 43 = alpina.
maxima, *Linn. Sp. Pl.* 141.—Europ.; Orient; As. et Afr. bor.
microphylla, *Hook. f. Fl. Brit. Ind.* iii. 498.—Reg. Himal.; Chin.
mucronifolia, *Watt, in Journ. Linn. Soc.* xx. (1884) 17.—Reg. Himal.
multiflora, Lam. Fl. Fr. ii. 259 = septentrionalis.
multiflora, Stead. Nom. ed. II. i. 94 = imbricata.
multiscapa, *Duby, in DC. Prod.* viii. 51.—As. Min.; Syria.
muscoidea, *Duby, l. c.* 48.—Reg. Himal.
nana, Hornem. Hort. Hafn. i. 184 = elongata.
neglecta, *Clarke, in Bull. Soc. Nat. Mosc.* xlv. (1872) i. 430.—Mont. Ural.
obconata, Wall. ex Duby, in DC. Prod. viii. 35 = Primula floribunda.
obtusifolia, *All. Fl. Pedem.* i. 90. t. 46. f. 1.—Europ.
obtusifolia, Willd. ex Ledeb. Fl. Ross. iii. 18 = A. Chamæjasme.
occidentalis, *Pursh, Fl. Am. Sept.* i. 137.—Am. bor.
ochotensis, *Willd. ex Roem. & Schult. Syst.* iv. 786.—Sibir.
odoratissima, Schreb. Dec. Pl. Min. Cogn. 7. t. 4 = villosa.
olympica, Boiss. Diagn. Ser. I. iv. 87 = villosa.
orbicularis, Roem. & Schult. Syst. iv. 816, Ind. = Gmelini.
Pacheri, Leyb. in Flora, xxxvi. (1858) 585 = alpina.
parviflora, Jacq. ex Duby, in DC. Prod. viii. 54 = rotundifolia.
patens, Wright, ex A. Gray, in Mem. Am. Acad. N. S. vi. (1858-9) 401 = saxifragæfolia.
pauciflora, Vill. Hist. Pl. Dauph. ii. 477 = obtusifolia.
pedemontana, Reichb Ic. Fl. Germ. xvii. 48 = obtusifolia.
pedunculata, *[Clairv.] Man. Herb.* 57.—Europ.
penicillata, Schott, Nyman & Kotschy, Anal. Bot. 18 = villosa.
pennina, Gand. Fl. Helv. ii. 108 = alpina.
pleiocarpa, Walp. Ann. iii. 7 = seq.
pliocarpa, C. Koch, in Linnæa, xxiii. (1850) 611 = armenica.
primulina, Spreng. Syst. iv. Cur. Post. 56 = Primula pusilla.
primuloides, D. Don, Prod. 81 = Primula pusilla.
primuloides, Duby, in DC. Prod. viii. 51 = sarmentosa.
primuloides, Moench, Meth. Suppl. 152 = Primula cortusioides.
pubescens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 105 = carnea.
pubescens, DC. Fl. Fr. iii. 488 = alpina.
pyrenaica, *Lam. Illustr.* i. 439.—Pyren.
radiata, Lehm. in Roem. & Schult. Syst. iv. 164 = filiformis.
Reverchoni, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 105 = carnea.
rosea, Jord. & Fourr. l. c. 104 = carnea.
rotundifolia, *Hardw. in As. Res.* iv. (1795) 350.—Reg. Himal.; China.
rotundifolia, Lehm. ex Roem. & Schult. Syst. iv. 165 = Gmelini.
rotundifolia, Sm. Exot. Bot. ii. 107 = saxifragæfolia.
rupina, [Clairv.] Man. Herb. 57 = Douglasia Vitaliana.
sarmentosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 14.—Reg. Himal.
saxifragæfolia, *Bunge, Enum. Pl. Chin. Bor.* 58.—China.
Seiugo, *Hook. f. & Thoms. ex Watt, in Journ. Linn. Soc.* xx. (1884) 18.—Reg. Himal.
sempervivoides, *Jacquem. ex Duby, in DC. Prod.* viii. 50.—Reg. Himal.; Mongol.

ANDROSACE:—
septentrionalis, *Linn. Sp. Pl.* 142.—Europ.; Caucas.; As. et Am. bor.
spathulata, Cav. Ic. v. 56 = Samolus spathulatus.
spatulata, Steud. Nom. ed. II. i. 98 = prænc.
stricta, Hartm. ex Duby, in DC. Prod. viii. 44 = Primula stricta.
strigiosa, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 10.—China.
tenerrima, Schlecht. ex Duby, in DC. Prod. viii. 47 = imbricata.
trifora, *Adans. in Mém. Soc. Nat. Mosc.* v. (1817) 89. —Sibir.
valerianoides, Lehm. ex Spreng. in Isis, (1817) 162 = albana.
villosa, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. iii. 498 = microphylla.
villosa, Jacq. Fl. Austr. t. 352 = A. Chamæjasme.
villosa, *Linn. Sp. Pl.* 142.—Europ.; Orient; As. temp.
villosa, S. G. Gmel. It. ii. 196 = maxima.
Vitaliana, Lap. Hist. Abr. Pl. Pyr. 94 = Douglasia Vitaliana.
Wiedemanni, *Boiss. Fl. Orient.* iv. 16.—Anatol.
Wulfeniana, Sieber, ex Koch, Syn. Fl. Germ. ed. II. ii. 670 = hedreantha.

ANDROSAEMUM, [Tourn.] ex Adans. Fam. ii. 444 (1763) = **Hypericum**, Linn.
calycinum, Presl, ex Steud. Nom. ed. II. i. 94 = H. calycinum.
Canbyi, Britton, Cosm. ex Nym. Consp. 181.—Ins. Baleor.
fortidum, Spach, ex Willk. & Lange, Prod. iii. 589 = H. hircinum.
grandiflorum, Reichb Ic. Fl. Germ. xvi. 70.—Europ.
hircinum, Spach, Hist. Vég. Phan. v. 419 = H. hircinum.
officinale, All. Fl. Pedem. ii. 147 = H. Androsæmum.
parviflorum, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 361 = H. elatum.
pyramidale, Spach, Hist. Vég. Phan. v. 417 = H. elatum.
vulgare, Gaert. Fruct. I. 292. t. 59 = H. Androsæmum.
Webbianum, Spach, Hist. Vég. Phan. v. 418 = H. grandiflorum.
xylosteifolium, Spach, l. c. 420 = H. inodorum.

ANDROSCEPIA, Brongn. in Duperr. Voy. Coq. Bot. 77 (1829) = **Anthistiria**, Linn. (Gram.).
Anathera, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 249 = Anth. Anathera.
Barberi, Anderss. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 176 = Anth. Barberi.
gigantea, Brongn. in Duperr. Voy. Coq. Bot. 78 = Anth. gigantea.
tremula, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 247 = Anth. tremula.

ANDROSEMUM, Neck. Elem. ii. 895 (1790) = præc.

ANDROSTEMMA, Lindl. Swan Riv. App. 46 (1839) = **Conostylis**, R. Br. (Haemodor.).
juncewm, Lindl. l. c. = C. Androstemma.

ANDROSTEPHIUM, Torr. in Bot. Mex. Bound. 218 (1859). *LILIACEAE*, Benth. & Hook. f. iii. 801.
breviflorum, *S. Wats. in Amer. Natural.* vii. (1873) 308.—Utah.
violaceum, *Torr. in Bot. Mex. Bound.* 219.—Texas.

ANDROSTOMA, Hook. f. Fl. Antarct. i. 44. t. 20 (1845) = **Cyathodes**, Labill. (Orchid.).
empetrifolia, Hook. f. l. c. = C. empetrifolia.

ANDROSTYLIUM, Mlq. in Nat. Verh. Wet. Haarl. vii. (Stirp Surin.) 99 (1851) = **Clusia**, Linn. (Guttif.).
Fockeanum, Mlq. l. c. = C. Fockeana.

ANDROSYNE, Salisb. Gen. Pl. Fragm. 61 (1866). An **Dioscorea**, Linn. 'Genus dubium, prope Paridem.' —Cf. Benth. & Hook. f. Gen. iii. 762.
gracilis, Salisb. l. c.—Ins. Nicob.

ANDROTRICHUM, Brongn in Duperr. Voy. Coq. Bot. 177. t. 32 (1829). *CYPERACEAE*, Benth. & Hook. f. iii. 1046.
CONOSTYLIUM, Nees, in Linnæa, ix. (1834) 283.
polycephalum, *Brongn. l. c.*—Bras.

ANDROTROPIS, R. Br. ex Wall. Cat. n. 6898 (1847) = **Acranthera**, Arn. (Rubiac.).
tomentosa, R. Br. ex Wall. l. c. = Acranth. tomentosa.

ANDRYALA, Linn. Gen. ed. I. 238 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 517.
ERIOPHORUS, Vaill. ex DC. Fl. Fr. iv. 86 (1808).
FORNEUM, Adans. Fam. ii. 112 (1763).
ROTHIA, Schreb. Gen. Pl. 531 (1791).
VOLGTIA, Roth, in Roem. & Ust. Mag. iv. (x.) 17 (1790).
aestivalis, *Pomel, Nouv. Mat. Fl. Alt.* 259 (err. typ. oestivalis).—Afr. bor.
Agardhii, *Hausskn. ex DC. Prod.* vii. 244.—Hispan.
allochrus, *Hoffmgg. & Link, Fl. Port.* ii. 154.—Hispan.
arenaria, Boiss. & Reut. Pugill. Pl. Hispan. 71 = integrifolia.
aurea, Scop. Ann. ii. Hist. Nat. (1769) 58 = Crepis aurea.
Bourgeauii, Sch. Bip. ex Webb. & Berth. Phyt. Canar. ii. 417 = pinnatifida.
candidissima, Desf. Cat. Hort. Par. ed. III. 399 = varia.
chirantifolia, L'Hérit. Stirp. Nov. 35. t. 18 = varia.
cheirantifolia, Link, ex Steud. Nom. ed. II. i. 94 = pinnatifida.
chondrilloides, *Scop. Fl. Carn.* ed. II. ii. 115.—Europ.
coronopifolia, Hoffmgg. & Link, Fl. Port. ii. 155 = pinnatifida.
corymbosa, Brot. Fl. Lusit. i. 387 = nigricans.
corymbosa, Lam. Illustr. iii. t. 657 = sinuata.
cossyrensis, Guss. Fl. Sic. Syn. ii. 407 = dentata.
crithmifolia, *Ait. Hort. Kew.* iii. 129.—Ins. Canar.
dentata, *Sibth. & Sm. Fl. Graec.* t. 811.—Reg. Mediterr.
denudata, Soland. ex Lowe, Man. Fl. Mad. i. 568 = cheirantifolia.
diffusa, Jan. ex DC. Prod. vii. 246 = integrifolia.
dissecta, *Hoffmgg. & Link, Fl. Port.* ii. 158.—Hispan.
Ficalhesana, *Daveau, in Bolet Soc. Brot.* i. (1883) 43, 51.—Lusit.
floccosa, *Pomel, Nouv. Mat. Fl. Alt.* 3.—Afr. bor.
glandulosa, Lam. Encyc. i. 154 = varia.
gomerana, Sch. Bip. ex Webb & Berth. Phyt. Canar. ii. 417 = pinnatifida.
incana, DC. Fl. Fr. Suppl. 445 = ragusina.
integrifolia, *Linn. Sp. Pl.* 808.—Reg. Mediterr.
integrifolia, Varia, Hort. Par. Panborm. 886, ex Guss. Fl. Sic. Syn. ii. 404 = Hieracium crinitum.
laciniata, Lam. Encyc. i. 153 = ragusina.
lanata, Habl. Taur. 155 = Leontodon biscutellæfolius.
lanata, Linn. Cent. I. n. 57, in Amoen. Acad. iv. 286 = Hieracium andryaloides.
lanata, Vill. Hist. Pl. Dauph. iii. 65 = integrifolia.
lanata, Sibth. & Sm. Prod. ii. 141 = Hieracium pannosum.
laxiflora, *DC. Prod.* vii. 246.—Hispan.; Marocc.
lyrata, Pourr. in Mém. Acad. Toul. iii. (1788) 308 = ragusina.
macrocephala, Boiss. ex DC. Prod. vii. 244 = ragusina.
malacitana, Haenseler, ex Willk. in Willk. & Lange, Prod. ii. 272 = laxiflora.
mogadorensis, *Coss. ex Hook f. Bot. Mag.* t. 6010.—Marocc.
mollis, [Asso] Mant. Stirp. Arag. 175 = integrifolia.
nemausensis, Vill. Prosp. 37 = Pterotheca sancta.
nigricans, *Poir. Voy. Barb.* ii. 228.—Numidia.
oxidiculis, Lam. Encyc. i. 154 = Pterotheca sancta.
oestivalis, err. typ. *vide* aestivalis.
parviflora, Lam. Fl. Fr. ii. 117 = integrifolia.
pinnatifida, *Ait. Hort. Kew.* ed. I. iii. 139.—Ins. Canar.
pontana, Vill. Prosp. 37 = Crepis montana.
ragusina, *Linn. Sp. Pl.* ed. II. 1186.—Reg. Mediterr.
ramosissima, Boiss. Elench. 68 = ragusina.
Reboudiana, *Pomel, Nouv. Mat. Fl. Alt.* 259.—Afr. bor.
robusta, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) repr. 18.—Ins. Mader.
Rothia, *Pers. Syn.* ii. 378.—Hispan.
runcinata, Pers. l. c. = integrifolia.
sinuata, Linn. Sp. Pl. 808 = integrifolia.
sinuata, Reichb. Ic. Fl. Germ. xiv. 1436 = integrifolia.
sinuata, Willk. & Lange, Prod. ii. 271 = ragusina.
tenuifolia, *DC. Prod.* vii. 245.—Sicil.
tenuifolia, Tenore, Fl. Neap. Syll. App. v. 88 = dentata.
tomentosa, Salisb. Prod. 182 = ragusina.

ANDRYALA:—
tomentosa, Soop. Delic. Insub. ii. 12 = varia.
undulata, J. & C. Presl, Delic. Prag. 113 = integrifolia.
uniflora, Schrank, Baier Fl. ii. 346 = integrifolia.
varia, *Lowe, ex DC. Prod.* vii. 245.—Ins. Mader.
Webbiana, Sch. Bip. ex Webb & Berth Phyt. Canar. iii. 417 = pinnatifida.

ANDREJEWSKYA, Reichb. Pl. Crit. i. 15. t. 18 (1837). *CRUCIFERÆ*, Benth. & Hook. f. i. 70.
MACROCERATIUM, Reichb. Consp. 184 (1828).
Cardamine, *Reichb. l. c.*—Pontus; Syria.

ANECHITES, Griseb. Fl. Brit. W. Ind. 410 (1864) *APOCYNACEÆ*, Benth. & Hook f. ii. 707.
adglutinata, *Mors, Apocyn. S. Am.* 236.—Ind. occ.
Asperuginis, *Griseb. Fl Brit W. Ind* 410.—Cuba.
circinalis, *Mors, Apocyn. S. Am.* 236.—Ind. occ.
lappulacea, *Miers, l. c* 237.—Ins. S. Doming.
revoluta, *Miers, l. c.* 238.—Ins.
thomasiana, *Miers, l. c.* 237.—Ins. S. Thomae.

ANECIO, Neck. Elem. i 28 (1790) = **Senecio**, Linn. (Compos.).

ANECOCHILUS. Blume, Bijdr. 411. t. 15 (1825) = **Anoectochilus**, Blume (1838) (Orchid.).

ANECTOCHILUS, Blume, Orch. Archip. Ind. 44 (1858) = **Anoectochilus**, Blume (Orchid.).

ANELEMA, R. Br. Prod. 270 (1810). *COMMELINACEÆ*, Benth. & Hook. f. iii. 849.
AMELINA, C. B. Clarke, Commel. et Cyrt. Beng. 38. t. 16 (1874).
ANILEMA, Kunth, Enum. Pl. iv. 64 (1843).
APHYLAX, Salisb. in Trans Hort. Soc. i. (1812) 371.
BAUSCHIA, Scop. in Vidensk. Meddel. Kjoeb. 128 (1872).
DICHORISTEMUM, Wight, l. c. vi. 31. t. 2078 (1853).
DICTYOSPERMUM, Wight, l. c. 29. t. 2089 (1853).
DILASIA, Rafin. Fl. Tellur. iv. 122 (1836).
LAMPRODITHYROS, Hassk. in Peters, Mossamb. Reise, Bot. 239 (1863).
MURDANNIA, Royle, Illustr. Bot. Himal. 408. t. 95 (1839).
PELETOCARPUS, Hassk. in Flora, xlix. (1866) 212.
PRIONOTACHYS, Hassk. l. c.
RHOPALEPHORA, Hassk. ex Mohl & Schlecht. in Bot. Zeit. (1864) 58.
STREPTYLIS, Rafin. Fl. Tellur. iv. 122 (1836).
TALIPULIA, Rafin. l. c. ii. 17 (1836).
acuminatum, Benth. Fl. Austral. vii. 85 = laxum.
aculeatum, *R. Br. Prod.* 270.—Austral.
adhaerens, Kunth, Enum. Pl. iv. 72 = sequinoctiale.
sequinoctiale, *Kunth, l. c.*—Afr. trop.
affine, R. Br. Prod. 271 = gramineum.
africanum, Beauv. Fl. Owar. ii. 57. t. 93 = Floscopa africana.
anthericoides, R. Br. Prod. 271 = gramineum.
asperum, Buch.-Ham. ex Wall. Cat. sub n. 5204 = Floscopa scandens.
benlenense, *Kunth, Enum. Pl.* iv. 73.—Afr. trop.
biflorum, *R. Br. Prod.* 270.—Austral.
bracteolatum, Marr. in Flora, xxiv. (1841) 11. Beibl. ii. 59 = ovato-oblongum.
brasiliense, *C. B. Clarke, in DC. Monog. Phan.* iii. 225. —Bras.
caladrinioides, *F. Muell. Fragm.* ix. 191.—Austral.
canaliculatum, Dals. in Hook. Kew Journ. iii. (1851) 137 = spiratum.
compressum, Dals. l. c. 138 = nudiflorum.
conspicuum, *Kunth, Enum. Pl.* iv. 69.—Ind. or.
cordatum, Buch.-Ham. ex Wall. Cat. sub n. 5228 = Commelina herbacea.
crispatum, R. Br. Prod. 270 = Pollia crispata.
crocea, Griff. Notul. iii. 235 = ochraceum.
cymosum, Kunth, Enum. Pl. iv. 69 = Floscopa scandens.
debile, Wall. Cat. n. 5215 = nudiflorum.
densiflorum, Kunth, Enum. Pl. iv. 69 = Floscopa scandens.
dicandrum, Buch.-Ham. in Wall. Cat. n. 5211 = nudiflorum.
didymum, Buch.-Ham. l. c. n. 5202 = Commelina tuberosa.
didymum, Seem. Bot. Voy. Herald, 422 = Pollia sorzogonensis.

ANELEMA:—
dimorphum, *Dals. in Hook. Kew Journ. iii.* (1851) 138. —Ind. or.
dimorphum, Thw. Enum. Pl. Zeyl. 322 = zeylanicum.
divergens, *C. B. Clarke, Commel. et Cyrt. Beng.* 38. t. 16.—Ind. or.
diversifolium, Hassk. in Miq. Pl. Jungh. 142 = nudiflorum.
Dregeanum, *Kunth, Enum. Pl.* iv. 73.—Afr. austr.
Ehrenbergii, *C. B. Clarke, in DC. Monog. Phan.* iii. 220.—Afr. trop.
elatum, *Dals. in Hook. Kew Journ.* iii. (1851) 137. —Ind. or
elatum, Kunth, Enum. Pl. iv. 70 = herbaceum.
evolvifolium, Wight, Ic. Pl. Ind. Or. t. 2072 = gigianteum.
esculentum, *Wall. Cat.* n. 5208.—Ind. or.
esculentum, Wall. l. c. list. C. (nec A. B.) = sinicum.
filiforme, Buch.-Ham. Wall. l. c. n. 5209 = vaginatum.
filipes, Mart. in Flora, xxiv. (1841) 11. Beibl. ii. 59 = geniculatum.
floribundum, Hook & Arn. Bot. Beech. Voy. 311 = Leptorhoeo filiformis.
foliosum, Hassk in Miq. Pl. Jungh. 144 = nudiflorum.
Forskalei, *Kunth, Enum. Pl.* iv. 71.—Arab.
Gardneri, *Seub. in Mart. Fl. Bras.* iii. 259.—Bras.
giganteum, *R. Br. Prod.* 271.—Austral.
glaucum, *Thw. ex C. B. Clarke, in DC. Monog. Phan.* iii. 200.—Zeylan.
gracile, *C. B. Clarke, l. c* 228.—Afr. occ.
gramineum, *R. Br. Prod.* 270.—Austral.
Hamiltonianum, *Wall. Cat.* n. 5222.—Assam.
herbaceum, *Wall. l. c.* n. 5223.—Ind. or.: Malaya.
herbaceum, Wall. l. c. n 5233 = protensum.
hirtum, *J. Rich. Tent. Fl. Abyss.* ii 343.—Abyss.
hispidum, D. Don, Prod. 45 = Floscopa scandens.
Hookeri, *C. B. Clarke, Commel. et Cyrt. Beng.* 38. t. 17.—Ind. or.
japonicum, *Kunth, Enum. Pl.* iv. 70.—Japon.
Junghuniannum, Miq. Fl. Ind. Bat. iii. 528 = nudiflorum.
Keisak, *Hassk. Commel. Ind.* 32.—Japon.
Koenigii, *Wall. Cat* n. 5214.—Ind. or.
lanceolatum, Baker, ex C. B. Clarke, in DC. Monog. Phan. iii. 227 = Ab. concolor.
lanceolatum, *Benth. in Hook. Niger Fl.* 546.—Afr. trop.
lancifolium, Griff. Notul. iii. 236 = nudiflorum.
langiinum, *Wall. Cat.* n. 5231.—Ind. or.
latifolium, Wight, Ic. Pl. Ind. Or. t. 2073 = herbaceum.
laxum, R. Br. Prod. 270 = Commelina laxa.
laxiflorum, Benth. ex C. B. Clarke, in DC. Monog. Phan. iii. 226 = ovato-oblongum.
laxum, *R. Br. Prod.* 270.—Austral.
linoides, Kunth, Enum. Pl. iv. 69 = herbaceum.
longifolium, Hook. Exot. Fl. t. 204 = gigianteum.
longifolium, Wall. Cat. n. 5213 = divergens.
loriforme, Hassk. in Miq. Fl. Jungh. 143 = nudiflorum.
Loureirii, *Hance, in Seem. Journ. Bot.* vi. (1868) 260. —China.
macrophyllum, R. Br. Prod. 270 = Pollia macrophylla.
medicum, *R. Br. l. c.* 271.—China.
melanoticum, Hance, in Seem. Journ. Bot. vii. (1869) 167 = esculentum.
micranthum, *Kunth, Enum. Pl.* iv. 70.—Ind. or.
minutum, Kunth, l. c. 66 = nudiflorum.
monadelphum, *R. Br. ex Kunth, l. c.* 70.—Java.
montanum, Thw. Enum. Pl. Zeyl. 322 = zeylanicum.
montanum, *Wight, in Wall. Cat.* n. 5208.—Ind. or.
nanum, Kunth, Enum. Pl. iv. 65 = spiratum.
nudicaule, Kunth, l. c. 67 = nudiflorum.
nudiflorum, F. Muell. Fragm. viii. 62 = giganteum.
nudiflorum, Miq. ex C. B. Clarke, in DC. Monog. Phan. iii. 206, 215 = dimorphum, Koenigii.
nudiflorum, *R. Br. Prod.* 271.—Ind. or.
nudiflorum, Wall. Cat. n. 5234 = vaginatum.
Nummularia, Miq. Fl. Ind. Bat. 535 = spiratum.
ochraceum, *Dals. in Hook. Kew Journ.* iii. (1851) 135.—Ind. or.
oliganthum, Franch. & Sav. Enum. Pl. Jap. ii. 522 = Keisak.
ovalifolium, *Hook. f. ex C. B. Clarke, in DC. Monog. Phan.* iii. 212.—Ind. or.
ovatum, *Wall. Cat.* n. 5206.—Malaya.
ovato-oblongum, *Benth. Fl. Austr.* ii. 71. t. 104.—Afr. trop.
paniculatum, *Wall. Cat.* n. 5216.—Ind. or.
paniculatum, Wight, Ic. Pl. Ind. Or. t. 2075 = dimorphum.

ANELEMA:—
pauciflorum, Dals. in Hook. Kew Journ. iii. (1851) 136 = vaginatum.
pauciflorum, *Wight, Ic. Pl. Ind. Or.* t. 2077.—Ind. or.
pedunculosum, *C. B. Clarke, in DC. Monog. Phan.* iii. 228.—Afr. austr.
Peterzii, *C. B. Clarke, l. c.* 225.—Afr. trop.
pilosum, Wall. Cat. n. 5219 = Commelina pilosa.
poseoides, *Seub. in Mart. Fl. Bras.* iii. 259.—Bras.
protensum, *Wall. Cat.* n. 5218.—Ind. or.
protensum, Thw. Enum Pl. Zeyl. 322 = zeylanicum.
radicans, D. Don, Prod. Fl. Nep. 45 = nudiflorum.
ramiforme, Buch.-Ham. in Wall. Cat. n. 5205 = Pollia subumbellata.
rivulare, A. Rich. Tent. Fl. Abyss. ii. 342 = Floscopa rivularis.
scaberrimum, Kunth, Enum. Pl. iv. 69 = protensum.
scapiflorum, *Kostel. Allg. Med.-Pharm. Fl.* i. 137.—Ind. or.
scapiflorum, Hook. f. & Thoms. ex C. B. Clarke, in DC. Monog. Phan. iii. 208 = divergens.
Schomburgkianum, *Kunth, Enum. Pl.* iv. 66.—Bras.
Schweinfurthii, *C. B. Clarke, in DC. Monog. Phan.* iii. 227.—Afr. trop.
sclerocarpum, *F. Muell. Fragm.* viii. 61.—Austral.
secundum, Wight, Ic. Pl. Ind. Or. t. 2074 = sinicum.
secundiflorum, Kunth, Enum. Pl. iv. 69 = Pollia sorzogonensis.
semsroes, Dals. in Hook. Kew Journ. iii. (1851) 136 = paniculatum.
sepalosum, *C. B. Clarke, in DC. Monog. Phan.* iii. 202. —Afr. trop.
serotinum, Don, ex C. B. Clarke, l. c. 200 = scapiflorum.
siliculosum, F. Muell. Fragm. viii. 61 = acuminatum.
siliculosum, *R. Br. Prod.* 270.—Austral.
simplex, Kunth, Enum. Pl. iv. 72 = sinicum.
sinicum, Herb. Berol. ex C. B. Clarke, in DC. Monog. Phan. iii. 211 = nudiflorum.
sinicum, [*Lindl. in*] *Bot. Reg.* t. 659.—China.
spectabile, Kurz, in Journ. As. Soc. Beng. xl. (1871) 11. 77 = Loureirii.
spicatum, Wall. Cat. n. 5217 = Loureirii.
spiratum, *R. Br. Prod.* 271.—Ind. or.
Tacasseanum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 343 = sequinoctiale.
Tacasseanum, Baker, in Trans. Linn. Soc. xxix. (1872) 163 = lanceolatum.
terminale, Wight, Ic. Pl. Ind. Or. t. 2076 = nudiflorum.
Thomsoni, *C. B. Clarke, in Journ. Linn. Soc.* xv. (1877) 121.—Reg. Himal.
trichocalcar, Schau. in Nov. Act. Acad. Leop. Carol. xix. (1848) Suppl. i. 448 (Pl. Meyen.) = nudiflorum.
triquetrum, *Wall. Cat.* n. 5220.—Ind. or.
tuberosum, Buch.-Ham. in Wall. Cat. n. 5207 = scapiflorum.
umbrosum, Kunth, Enum. Pl. iv. 71 = ovato-oblongum.
vaginatum, *R. Br. Prod.* 271.—Ind. or.
versicolor, *Dals. in Hook. Kew Journ.* iii. (1851) 136. —Ind. or.
violaceum, F. Muell. ex C. B. Clarke, in DC. Monog. Phan. iii. 205 = giganteum.
vitiense, *Seem. Fl. Vit.* 314. t. 96.—Ins. Fiji.
Welwitschii, *C. B. Clarke, in DC. Monog. Phan.* iii. 229.—Angola.
zeylanicum, *C. B. Clarke, l. c.* 204.—Zeylan.

ANELASMA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42 = Abruta, Aubl. (Menisp.).
concolor, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 260 = Ab. concolor.
Gardnerianum, Miers, l. c. Ser. II. vii. (1851) 43 = Ab. concolor.
guianense, Miers, l. c. = Ab. concolor.
intermissum, Miers, l. c. Ser. III. xiv. (1864) 260.—Bras.
jamaicense, Miers, ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 50 = Pachygone domingensis.
laurifolium, Sagot, ex Benth. l. c. = Ab. concolor.
laurifolium, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 260.—Ins. S. Thomae.
Martianum, Miers, l. c. = Ab. concolor.
minutiflorum, Sagot, ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 50 = Pachygone domingensis.

ANELASMA:—
pallidum, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 260 = Ab. concolor.
Sellowianum, Miers, l. c.—Bras.
Spruceanum, Miers, ex Benth. in Hook. Kew Jour. iii. (1851) 116 = Ab. concolor.
strumarum, Miers, in Ann. & Mag. Nat. Hist. Ser. III xiv. (1864) 260.—Bras.
urophyllum, Miers, l. c.—Bras.

ANEMAGROSTIS, Trin. Fund. Agrost. 128 (1820) = **Apera**, Adans. (Gram.).
interrupta, Trin. l. c. 120 = Ap. interrupta.
Neesiana, Wight & Arn. ex Steud. Syn. Pl Gram 115 = Panicum semialatum.
Spica-venti, Trin. Fund. Agrost. 129 = Ap Spica-venti.
tenella, Wight, ex Steud. Syn. Pl. Gram. 115 = Aristidinella tenella.

ANEMANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 323 = **Anemone**, Tourn (Kanunc.)
nemorosus, Fourr. l. c. = Anemone nemorosa.
ranunculoides, Fourr. l. c. = Anemone ranunculoides.

ANEMARRHENA, Bunge, in Mém. Acad Pétersb ii (1881) 140 (Enum. Pl. Chin. Bor. 66). *LILIACEÆ*, Benth. & Hook. f. iii. 782
asphodeloides, *Bunge, it. cc.*—China.

ANEMIA, Nutt. in Ann. Nat. Hist. Ser. I. i. (1838) 136 = **Ruthtuynia**, Thunb. (Piperac.).
californica, Nutt. l. c. = H. californica.

ANEMIOPSIS, Endl. Gen. Suppl. i. 1374 (1841) = ANEMOPSIS, Hook. & Arn.

ANEMITIS, Rafin. Fl. Tellur. iii. 87 (1836) = **Phlomis**, Tourn. (Labiat.).
virgida, Rafin. l. c. = P. Herba-venti.

ANEMONANTHEA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 724 (1821) = **Anemone**, Linn. (Ranunc.).
apennina, S. F. Gray, l. c. 724 = Anemone apennina.
nemorosa, S. F. Gray, l. c. 725 = Anemone nemorosa.
ranunculoides, S. F. Gray, l. c. = Anemone ranunculoides.

ANEMONE, Linn. Syst. ed. I. (1735). *RANUNCULACEÆ*, Benth. & Hook. f. i. 4.
ANEMANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N S. xvi. (1868) 323.
ANEMONANTHEA, S. F. Gray, Nat. Arr. Brit. Pl. ii 724 (1821).
ANEMONELLA, Spach, Hist. Veg. Phan. vii. 240 (1839).
BARNEOUDIA, C. Gay, Fl. Chil. i. 29. t. 1 (1846).
FLAMMARA, Hill, Veg. Syst. xvi. 37 (1770).
HARTIANA, Rafin. Neogenyt. 2 (1825).
HEPATICA, Dill. ex Linn. Syst. ed. I (1735)
HOMALOCARPUS, Schur, Enum. Pl. Transyl. 3 (1866).
ISOPYRUM, Adans. Fam. ii. 460 (1763).
ORIBA, Adans. l. c. 459 (1763).
PSEUDANTHUS, Ehrh. Beitr. iv. 149 (1789).
PULSATILLA, (Tourn.) Linn. Syst. ed. I (1735).
SYNDESMON, Hoffmgg. in Flora, xv. (1832) Intell. 24.
acanthifolia, *Rich. Ind. Sem. Hort. Patav.* (1841).—Hab. ?
aconitifolia, Michx. Fl. Bor. Am. i. 320 = dichotoma.
aconitifolia, Turcz. in Bull. Soc. Nat. Mosc. (1840) 61 = sarcissifiora.
acutiloba, *Lour. Ranunc. Canad.* 30.—Am. bor.
acutipetala, Schleich. Cat. Fl. Helv. ed. III. 6 = A. Pulsatilla.
aequinoctialis, *Pupp. Fragm. Syn.* 27.—Peruv.
affinis, G. Don, Gen. Syst. i. 21 = pratensis.
alba, Gilib. Fl. Lituan. ii. 275 = nemorosa.
alba, Juss. in Ann. Mus. Par. iii. (1804) 248. t. 20. f. 1 = sylvestris.
alba, Goaty & Pons, in Bull. Soc. Bot. Fr. xxx. (1883) p. lxxix = coronaria.
albana, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 264. —Orient; Ab. bor.
alborosea, Gilib. Fl. Lituan. ii. 276 = vernalis.
alchemilloides, *E. Mey. ex Pritz. in Linnæa*, xv. (1841) 758.—Afr. austr.

ANEMONE:—
alpina, *Linn. Sp. Pl.* 539.—Europ. ; Caucas. ; Am. bor.
alpina, Scop. Fl. Carn. ed. II. ii. 384 = baldensis.
alsatica, *Fisch. ex Ledeb. Fl. Alt.* ii. 262.—Sibir. a'sai'.
ambigua, Turcz. ex Pritz. in Linnæa, xv. (1841) 601 = albana.
angulosa, *Lam. Encyc.* i. 169.—Europ. or.
anomala, *Rafin. Fl. Ludov* 82 (quid ?)—Am. bor.
antucensis, *Poepp. Fragm. Syn.* 27.—Chili
apennina, Auct. Orient. ex Boiss. Fl. Orient. i. 18 = blanda.
apennina, *Linn. Sp. Pl.* 541.—Europ.
apiifolia, Scop. Fl. Carn. ed. II. i. 385 = alpina.
arborea, Hort. ex Steud. Nom. ed. II. i. 95 = capensis.
arctica, Fisch. in Linnæa, vi. (1831) 574 = Richardsoni.
argentea, Raim. ex Wedd. Chlor. Andina, ii. 305 = Ranunculus Raimondii.
armena, *Boiss. Fl. Orient.* i. 10 —As. Min.
baicalensis, *Turcz. in Bull. Soc. Nat Mosc.* xv. (1842) 40.—As. temp.
baldensis, G. Don, Gen. Syst i. 18 = multifida.
baldensis, Lam. Encyc. i. 614 = alpina.
baldensis, *Linn. Mant.* i. 78.—Europ.
barbulata, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) ii. VII. 149.—China.
Bauhini, *Tausch, in Flora*, xxv. (1842) i. 69.—Europ.
Berlandieri, Pritz. in Linnæa, xv. (1841) 628 = decapetala.
bicolor, Poepp. ex Pritz l. c. 624 = decapetala.
bifora, *DC. Syst.* i. 201.—Orient ; Reg. Himal.
bilobata, Phil f Cat Pl. Vasc. Chil. 5 = decapetala.
blanda, *Schott & Kotschy, in Oestr. Bot. Wochenbl.* iv. (1854) 129.—Orient.
Bogenbardiana, *Prtz. in Linnæa*, xv. (1841) 591.—Europ.
borealis, Richards. in Frankl. 1st Journ. ed. II. App. 22 = parviflora.
Bungeana, *Pritz. in Linnæa*, xv. (1841) 603.—Sibiria.
Burseriana, Scop. Fl. Carn. ed. II. i. 385 = alpina.
caciesiana, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 4.—China.
caerulea, *DC. Syst.* i. 203.—Sibir.
caerulea, Lam. Fl. Fr. iii. 318 = appenina.
caffra, *Harv. Gen. S. Afr. Pl.* 9.—Afr austr.
Campanella, Fisch. ex Ledeb. Fl. Ross. i. 22 = albana.
campanifora, Stokes, Bot. Mat. Med. iii. 247 = pratensis.
canadensis, Linn Syst ed. XII. iii. App. 231 = dichotoma.
capensis, *Lam. Encyc.* i. 164.—Afr. austr.
caroliniana, Torr. & Gray, Fl. N. Am. i. 12, partim = heterophylla.
caroliniana, Walt. Fl. Carol. 156 = decapetala.
caucasica, Willd. ex Rupr. Fl. Caucas. 14 = blanda.
cernua, *Thunb. Fl. Jap.* 238.—Japon.
chilensis, Spreng. ex Eichl. in Mart. Fl. Bras. xiii. i. 152 = decapetala.
chinensis, *Bunge, Enum. Pl. Chin.* 2.—China.
cocinea, Lord. Diag. i. 57 = coronaria.
colina, Salisb. Prod 371 = A. Pulsatilla.
Commersoniana, DC. in Deless. Ic. i. 4 = multifida.
coronaria, *Linn. Sp. Pl.* 539.—Reg. Mediterr. ; Orient.
coronarioides, Hanry, Prod. Hist. Nat. Var, 142 = coronaria.
crassifolia, *Hook. Ic. Pl.* t. 257.—Tasmania.
cuneata, Schlecht. in Linnæa, vi. (1831) 674 = parviflora.
cunifolia, Juss. in Ann. Mus. Par. iii. (1804) 248. t. 21 = parviflora.
curta, Wall. Cat. n. 4090 = Clematis montana.
cyanea, Risso, Fl. Nice, 7 = coronaria.
cylindrica, *A. Gray, in Ann. Lyc. N. York*, iii. (1828–36) 221.—Am. bor.
cynosurus, Griff. Notul. iv. 735 = Ranunculus falcatus.
dacica, Freyn, ex Borb. in Just, Bot. Jahresb. (1874) 1076, *err. typ.* = Arabis dacica.
dahurica, *Fisch. ex DC. Prod.* i. 17.—As. temp.
debilis, *Fisch. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 274 (bis).—Sibir.
decapetala, J. F. Gmel. Syst. 871 = Geum rivale.
decapetala, *Arduini, Animad. alt.* 27.—Am. bor. et austr.
decaphilla, Krock, Fl. Sil. ii. t. 285 = Geum rivale.
deltoidea, *Dougl. in Hook. Fl. Bor. Amer.* i. 6. t. 8. f. A.—Am. bor. occ.
denticulata, *Hook. f. & Thoms. Fl. Ind.* i. 28.—Reg. Himal.

ANEMONE:—
dichotoma, *Linn. Sp. Pl.* 540.—As. bor. ; Am. bor.
discolor, Royle, Illustr Bot. Himal. 8 = obtusiloba.
dodecaphylla, Krock, Fl. Sil. ii. t. 285 = Geum rivale
Drummondii, *S. Wats. Bot. Calif.* ii. 424.—Calif.
dubia, Bell. App. Fl Pedem. 232 = narcissiflora.
dubia, Wall Cat. n. 4097 = rivularis
elegans, Decne. in Rev. Hortic. (1852) 41. cum ic. = japonica.
elongata, *D. Don, Prod Fl Nep* 194.—Reg Himal.
eranthoides, *Regel, in Act Hort. Petrop.* viii. (1884) 691.—As. temp.
exigua, *Maxim. in Bull. Acad. Petersb.* xxiii. (1877) 306.—China
Falconeri, *Thoms. in Hook. Ic. Pl.* t. 899.—Reg. Himal.
Fauconni, *Harv. Gen. S. Afr. Pl. ed.* II. 2, nomen.—Natal.
fasciculata, Linn Sp. Pl 542 = narcissiflora
fasciculata, Vahl, Symb. Bot. iii. 74 = umbellata.
Fischeriana, *DC. Prod.* i. 20.—Sibiria.
flaccida, *Fr. Schmidt, Reisen Amuri* 103.—Japon China.
flava, Gilib. Fl Lituan ii. 274 = ranunculoides.
flaviscens, Zucc in Flora, ix. (1826) i. 373 = patens.
flemmensis, Scop. Ann. ii. Hist. Nat. (1796) 54 = alpina.
formosa, *Clark, Trav.* ii 145.— As Min.
fragifera, Wulf in Jacq. Misc. ii. 55 = baldensis.
fulgens, *J. Gay, in DC Prod.* i. 18.—Reg. Mediterr
fumariaefolia, Juss. in Ann. Mus Par iii. (1804) 247 = triternata.
geranifolia, Wall. Cat. n. 4098 = rivularis.
gigantea, Raim. ex Wedd. Chlor. And ii. 304 = Ranunculus gigantea.
glacialis, [Clairv.] Man. Herb. 180 = Halleri.
Gmelinoviana, *Ural. in Linnæa*, xliii. (1880–82) 25.—Ross.
globosa, Nutt. ex Pritz in Linnæa, xv. (1841) 673 = multifida.
Gmelinsana, *Presl, ex Prtz. l. c.* 697.—Sibiria.
Gortschakowii, Kar & Kir. in Bull. Soc. Nat Mosc. xv (1842) 131 = biflora.
Gouaniana, Lindl. Bot. Reg. (1844) Misc. 45 = polyanthes
Gouaniana, Wall. Cat. n. 4088 = obtusiloba.
graecia, *Fr. Schmidt, Reisen Amurl. Ivi.*—Japon.
grandiflora, Hoppe, ex Reichb. Fl. Germ. Excurs. 732 = alpina.
graveana, Goaty & Pons, in Bull. Soc. Bot. Fr. xxx (1883) p. lxxviii = coronaria.
Gray, *Behr. ex Kellogg, in Bull. Calif. Acad.* n. 1. (1864) 5.—Calif.
Griffithii, *Hook. f & Thoms Fl. Ind.* i. 24.—Reg. Himal.
groenlandica, O. F. Muell. in Fl. Dan. t. 566 = Coptis trifolia.
Hackelii, Steud. Nom. ed. I. 48 = Halleri.
Halleri, *All. Fl. Pedem.* ii. 170.—Europ.
heldreichiana, Richter, ex Pritz. in Linnæa, xv. (1841) 608 = pratensis.
helleborifolia, *DC. Syst.* i. 211.—Am austr.
Hepatica, *Linn. Sp. Pl.* 538.—Europ. ; Orient ; Am. bor.
hepaticifolia, *Hook. Ic. Pl.* t. 1.—Chili.
heterophylla, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 12. —Am. bor.
hirsuta, Gilib. Fl. Lituan. ii. 276 = sylvestris.
hirsuta, Moench, Meth. Suppl. 105 = virginiana.
hispida, Wall. Cat. n. 4094 = rivularis.
hortensis, Linn. Sp. Pl. 540 = stellata.
hortensis, Thore, Chl. Land. 238 = fulgens.
Hudsoniana, Richards. in Frankl. 1st Journ. ed. II. App. 22 = multifida.
hybrida, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 74 = japonica.
hybrida, Mig. ex Pritz. in Linnæa, xv. (1841) 573 = Halleri.
incisa, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 1 = caerulea.
indeterminata, Banks, ex Walp. Rep. i. 28, sphalm. = parviflora.
integrifolia, *Humb. & Bonpl. ex DC. Syst.* i. 217.—Am. merid.
intermedia, G. Don, Gen. Syst. i. 21 = patens.
intermedia, Hoppe, ex Steud. Nom. ed. I. 48, 49 = montana.
interminata, Schult. Obs. 101 = A. Pulsatilla.

ANEMONE :—
intermedia, Winkl. ex Pritz. in Linnæa, xv. (1841) 665 = A. nemorosa.
irregularis, Lam. Encyc. i. 167 = dichotoma.
isopyroides, *Ynss. in Ann. Mus. Par.* iii. (1804) 249. —Sibir.
Jamesoni, *Hook. Ic. Pl.* t. 670.—Ecuador.
Jankæ, *F. Schultz, in Flora*, xxxix. (1856) 205.—Transayl.
japonica, *Sieb. & Zucc. Fl. Jap.* i. 15.—Japon.
lancifolia, Pursh, Fl. Am. Sept. ii. 386 = nemorosa.
lanigera, C. Gay, Fl. Chil. i. 22 = multifida.
latifolia, Bell. ex Re, in Mem. Acc. Sc. Torin. xxxiii. (1829) 283 = fulgens.
Laxmanni, Steud. Nom. ed. II. i. 96 = dichotoma.
lepida, Jord. Diag. i. 59 = hortensis
libanotica, *Fisch. ex Pritz. in Linnæa*, xv. (1841) 697.—Kamtschatka.
linearia, Schleich. ex Pritz. l. c. 661 = caerulea.
lobata, Pers. Syn. ii. 96 = palmata.
longipetala, Schleich. ex Steud. Nom. ed. I. 48, 49 = patens.
longiscapa, Wall. Cat. n. 4691 = polyanthes.
ludoviciana, Nutt. Gen. Am. ii. 20 = patens.
lutea, Lam. Fl. Fr. iii. 318 = ranunculoides.
macrorrhiza, Domb. ex Eichl. in Mart. Fl. Bras. xiii. i. 121 = decapetala.
malvifolia, Linn. Sp. Pl. 1197 = palmata.
mexicana, *H. B. & K. Nov. Gen. et Sp.* v. 33 — Mexic.
micrantha, Hort. ex Steud. Nom. ed. II. i. 96 = alpina.
micrantha, Klotzsch, in Bot. Ergeb. Waldem. Reise, 133 = obtusiloba.
millefoliata, Bertol. Amoen. 574 = alpina.
minima, DC. Syst. i. 206 = nemorosa.
minuta, *Laxm. ex Sievers, in Pall. Nov. Nord. Beitr.* vii. 183.—Sibir.
mollis, Wall. Cat. n. 4689 = obtusiloba.
montana, *Hoppe, ex Sturm, Fl. Deutschl. Heft* xc. n. 2.—Europ ; Caucas.
Mounardi, Hanry, ex Arduino, Fl. Alp. Marit. 12 = coronaria.
multifida, *Poir. Encyc. Suppl.* i. 864.—Am. bor. et austr.
myrrhidifolia, Vill. Prosp. 50 = alpina.
narcissiflora, *Linn. Sp. Pl.* 542.—Europ.; Orient; As. bor. ; Am. bor.
narcissiflora, Hook. & Arn. Bot. Beech. Voy. 121 = multifida.
nemorosa, *Linn. Sp. Pl.* 541.—Europ.; As. bor.; Am. bor.
nemorosa, Schang. in Fall. Neu. Nord. Beitr. vi. 34 = alsatica.
nemorosa-alba, Crantz, Stirp. Austr. ed. I. fasc. ii. 101 ; ed. II. 121 = nemorosa.
nemorosa-lutea, Crantz, l. c. = ranunculoides.
nikomas, *Maxim. in Bull. Acad. Pétersb.* xviii. (1875) 275.—Japon.
nobilis, Jord. Diag. i. 57 = coronaria.
Nuttalliana, DC. Syst. i. 198 = patens.
Nuttallii, Nutt. in Journ. Acad. Soc. Nat. Phil. v. (1826) 158 = patens.
obsoleta, Hort. ex Steud. Nom. ed. II. i. 96 = pratensis.
obtusiloba, *D. Don, Prod.* 194.—Reg. Himal.
obtusiloba, Lindl. Bot. Reg. xxx. (1844) t. 65 = scopsoa.
occidentalis, *S. Wats. in Proc. Am. Acad.* xi. (1876) 121. —Calif.
ochotensis, Fisch. Hort. Gorenk. ed. II. (1812) 47 = sylvestris.
ochroleuca *v. Bruegg. in Verhh. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 77.—Helvet.
octopetala, *Patrin, ex Pritz. in Linnæa*, xv. (1841) 697, nomen.—Hab.?
Ornanthe, Ucria, in Roem. Arch. i. 69 = coronaria.
palmata, *Linn. Sp. Pl.* 538.—Reg. Mediterr.
parviflora, *Michx. Fl. Bor. Am.* i. 319.—Am. bor.
patens, Hoppe, in Sturm, Deutschl. Fl. Heft xivl. 1 = Halleri.
patens, *Linn. Sp. Pl.* 538.—Europ. ; Am. bor.
Pavonina, *Boiss. Diagn.* Ser. II. i. 6 ; *et ex Lorentz & Levier, in Journ. Bot.* xvii. (1879) 196.—Iberia.
pavonina, Lam. Encyc. i. 166 = hortensis.
pedata, Rafin. in Med. Repos. New York, v. (1808) 361 = nemorosa.
pennsylvanica, Linn. Mant. 247 = dichotoma.
pentaphylla, Hook. ex Pritz. in Linnæa, xv. (1841) 657 = nemorosa.
pinnatifida, Dalac, Fl. Hautes-Pyr. 214 = Halleri.
pixidenia, Sims. Agr. Tosc. 293 (1801) = A. Pulsatilla.

ANEMONE :—
Pitonii *v. Glowacki, in Verh. Zool.-Bot. Ges. Wien*, xix. (1869) 901.—Europ.
polyanthes, *D. Don, Prod. Fl. Nep.* 194.—Reg. Himal.
praecox, Salisb. Prod. 371 = A. Hepatica.
praentabilis, Jord. Diagn. i. 58 = coronaria.
pratensis, C. Koch, in Linnæa, xv. (1841) 648 = montana.
pratensis, *Linn. Sp. Pl.* 539.—Europ.
pratensis, Pall. ex Pritz. in Linnæa, xv. (1841) 687 = sylvestris.
pratensis, Sibth. Fl. Ox. 169 ; With. Arr. Brit. Pl. ed. III. 498 = A. Pulsatilla.
propinqua, G. Don, Gen. Syst. i. 21 = pratensis.
pubescens, Domb. ex Steud. Nom. ed. I. 466, in syn. = Ranunculus Krapfia.
Pulsatilla, Bieb. Fl. Taur. Cauc. ii. 18 = montana.
Pulsatilla, *Linn. Sp. Pl.* 539.—Europ.
Pulsatilla, Sturm, Fl. Deutschl. i. Heft lxxxix. 19 = pratensis.
pusica, Sims. Agr. Tosc. 223 = A. Pulsatilla.
pusilla, DC. Syst. i. 197 = coronaria.
pusilla, Gaertn. ex Steud. Nom. ed. I. 49 = Geum anemonoides.
pygmaea, Hort. ex Vill. Fl. Pl. Terre, ed. III. 90 = spennina.
pyrenaica, Pall. ex Pritz. in Linnæa, xv. (1841) 640 = spennina.
quinquefolia, Linn. Sp. Pl. 541 = nemorosa.
Raddeana, *Regel, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) ii. 16.—Reg. Amur.
ranunculiflora, St. Lag in Ann. Soc. Bot. Lyon, vii. (1880) 119 = ranunculoides.
ranunculiformis, St. Lag. l. c. 67 = ranunculoides.
ranunculoides, *Linn. Sp. Pl.* 541.—Europ.; Caucas.; Sibir.
ranunculoides, Richards. in Frankl. 1st Journ. App. 12 = Richardsoni.
ranunculoidi-nemorosa, Kunze, in Fl. Dan. t. 2738 = nemorosa.
reflexa, Leiteb. ex Spreng. Syst. ii. 662 = caerulea.
reflexa, *Stepb. in Willd. Sp. Pl.* ii. 1282.—Sibir.
Regina, Nisso, Fl. Nice, 6 = coronaria?
Richardsoni, *Hook. Fl. Bor. Am.* i. 8. t. 4.—Am. arct.
rigida, *C. Gay, Fl. Chil.* i. 25.—Chili.
Rissoana, Jord. Diag. i. 58 = coronaria.
rivularis, *Buch.-Ham ex DC. Syst.* i. 211.—Ind. or.
rivularis, Wall. Cat. n. 4692 = elongata.
rosea, Hanry, ex Nym. Consp. 3 = coronaria.
Rossii, *S. Moore, in Journ. Linn. Soc.* xvii. (1880) 376 t. 16.—China.
rubra, Lam. Encyc. i. 168 = A. Pulsatilla.
rupestris, Jacquem. Bot. Voy. 5 = ropfcola.
rupestris, Wall. *Cat.* n. 4696.—Reg. Himal.
rupicola, *Camb. in Jacquem. Bot. Voy.* iv. 5.—Reg. Himal.
Salesovii, Fisch. ex Pritz. in Linnæa, xv. (1841) 655 = alsatica.
sanguinea, Pursh, ex Pritz. l. c. 673 = multifida.
scaposa, Edgew. in Trans. Linn. Soc. xx. (1851) 27 = polyanthes.
Sellowii, *Pritz. in Linnæa*, xv. (1841) 667.—Bras.
sibirica, *Linn. Sp. Pl.* 541.—Sibir.
silvestris, vide sylvestris.
slavica, *Reuss, Berichtt. Centr. Litt.* i. 62.—Europ.
sordida, Schar, Enum. Pl. Transs. 3 = sylvestris.
speciosa, Adams, ex Pritz. in Linnæa, xv. (1841) 685 = narcissiflora.
speciosa, *Fisch. ex Pl. Caucas.* 12.—Caucas.
sphenophylla, *Poepp. Fragm. Syn.* 27.—Chili.
stellata, Lam. Encyc. i. 166 = hortensis.
stolonifera, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 226.—Japon.
sulphurea, Fil. Fedem. ii. 170 = vernalis.
sulphurea, Linn. Mant. 78 = alpina.
sumatrana, *De Vriese, in Mig. Pl. Jungh.* 76.—Sumatra.
sylvestris, *Linn. Sp. Pl.* 540.—Europ. ; Caucas.
sylvestris, Vill. Hist. Pl. Dauph. iii. 726 = A. Pulsatilla.
tenella, Banks, ex Pritz. in Linnæa, xv. (1841) 683 = parviflora.
tenella, Pursh, Fl. Am. Sept. ii. 386 = decapetala.
tenuifolia, *DC. Syst.* i. 196.—Afr. austr.
tenuifolia, Schleich. Cat. Fl. Helv. ed. IV. 7 = Bogenhardiana.
tetrasepala, *Royle, Illustr. Bot. Himal.* 53.—Reg. Himal.
thalictroides, *Linn. Sp. Pl.* 542.—Am. bor.

ANEMONE :—
Thomsoni, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 397.—Afr. trop.
transylvanica, *Heuff. in Verh. Zool.-Bot. Ver. Wien*, (1858) 42.—Europ.
trifolia, DC. Prod. i. 20 = reflexa.
triloba, *Linn. Sp. Pl.* 540.—Europ. ; Am. bor.
trifoliata, Georgi, Beschr. Russ. Reich. iii. iv. 1055 = reflexa.
triloba, Stokes, Bot. Mat. Med. iii. 248 = A. Hepatica.
trilobata, Juss. in Ann. Mus. Par. iii. (1804) 247. t. 1 = decapetala.
trilobata, Pers. Syn. ii. 97 = parviflora.
tritensis, *Vahl, Symb. Bot.* iii. 74. t. 65.—Am. austr.
trullifolia, *Hook. f. & Thoms. Fl. Ind.* i. 22.—Reg. Himal.
Tschernaewi, *Regel, in Act. Hort. Petrop.* viii. (1884) 690.—As. temp.
udensis, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 6).—Manchur.
umbellata, Lam. Fl. Fr. iii. 322 ; Willd. Sp. Pl. ii. 1284 = narcissiflora.
umbrosa, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 261.—Sibir.
uniflora, Pursh, ex Steud. Nom. ed. II. i. 96 = Thalictrum anemonoides.
uralensis, Fisch. ex DC. Prod. i. 19 = caerulea, baldensis.
Vahlii, Hornem. Fl. Dan. t. 2176 = Richardsoni.
variata, Jord, ex Nym. Consp. 3 = hortensis.
Ventenata, Hanry, ex Arduino, Fl. Alp. Marit. 12 = coronaria.
vernalis, *Linn. Sp. Pl.* 538.—Europ.
verticolor, Jord. Pugill. (1852) 1 = coronaria.
verticolor, Salisb. Prod. 371 = hortensis.
villosa, Royle, Illustr. Bot. Himal. 53 = polyanthes.
virginiana, *Linn. Sp. Pl.* 540.—Am. bor.
vitifolia, *Buch.-Ham. ex DC. Syst.* i. 210.—Reg. Himal.
Wahlenbergii, *Szontagh, ex Nym. Consp.* 2.—Europ.
Wallichiana, Royle, Illustr. Bot. Himal. 52 = albana.
Walteri, *Pursh, Fl. Am. Sept.* ii. 387.—Am. bor.
Wightiana, *Hook. Ic. Pl.* ii. t. 176.—Ind. or. or. *Wightiana*, Wall. Cat. n. 4697 = rivularis.
Wolfgangiana, *Bess. ex Koch, in Roehl. Deutschl. Fl.* iv. 106.—Europ.

ANEMONELLA, Spach, Hist. Vég. Phan. vii. 339 (1839) = Anemone, Linn. (Rauncx.)
thalictroides, Spach, l. c. 240 = Anemone thalictroides.

ANEMONOPSIS, Prits. Ind. Ic. 71, sphalm. = ANEMOPSIS, Hook.

ANEMONOPSIS, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 181. *RANUNCULACEÆ*, Benth. & Hook. f. i. 8.
XAVERIA, Endl. Gen. Suppl. iv. 80 (1847).
macrophylla, *Sieb. & Zucc. l. c.* 182.—Japon.

ANEMONOSPERMUM, Commers. ex Steud. Nom. ed. II. i. 480, in syn. (1840) = Cryptostemma, K. Br. (Compos.).

ANEMOPAEGMA, Mart. ex DC. Prod. ix. 187 (1845). *BIGNONIACEÆ*, Benth. & Hook. f. ii. 1037.
acutifolium, *DC. Prod.* ix. 188.—Bras.
album, *Mart. ex DC. l. c.*—Bras.
citrinum, *Mart. ex DC. l. c.* 189.—Bras.
clematideum, *Grisob. in Gott. Abh.* xix. (1874) 292. —Reg. Argent.
floridum, *Mart. ex DC. Prod.* ix. 188.—Bras.
glaucum, *Mart. ex DC. l. c.*—Bras. ; Peruv.?
laeve, *DC. l. c.* 188.—Bras.
lanceasfolium, *DC. l. c.* 188.—Bras.
longidentatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = seq.
longidens, *DC. Prod.* ix. 190.—Bras.
mirandum, *Mart. ex DC. l. c.* 187.—Bras.
orbiculatum, *DC. l. c.* 190.—Panama.
Poeppigii, *DC. l. c.*—Am. austr.
prostratum, *DC. l. c.* 189.—Bras.
quinquedentatum, *Mart. ex DC. l. c.* 190.—Bras.
racemosum, *Mart. ex DC. l. c.* 189.—Bras.
scabriusculum, *Mart. ex DC. l. c.* 188.—Bras.
sessilifolium, Mart. ex DC. l. c. 187 = mirandum.
triplinervium, *Mart. ex DC. l. c.* 188.—Bras.
Vauguianum, *DC. l. c.* 190.—Bras.
velutinum, *Mart. ex DC. l. c.* 189.—Bras.

ANEMOPSIS, [Hook.] in Ann. Nat. Hist. i. (1836) 186, in note; Hook. & Arn. Bot. Beech. Voy. 390. t. 93 = **Houttuynia**, Thunb. (Piperac.).
Berlanderi, C. DC. in Linnaea, xxxvii. (1871–73) 333 — H. californica.
californica, Hook. & Arn. Bot. Beech. Voy. 390. t. 92 = H. californica.
Ludovici-Salvatoris, Willk. in Regel, Gartenfl. xxvi. (1877) 293.—Calif.

ANEPSA, Rafin. Fl. Tellur. ii. 81 (1836) = **Veratrum**, Tourn. (Liliac.).
carinata, *graminifolia*, *latifolia*, *spicata*, Rafin. l. c.—(Spp. ignotæ).

ANEPSIAS, Schott, Gen. Aroid. t. 73 (1858) = **Rhodospatha**, Poepp. (Aroid.).
Moritziana, Schott, l. c.—Venezuela.

ANERINCLEISTUS, Korth. in Verh. Nat. Gesch. ed. Temminck, 250. t. 66 (1842). *MELASTOMACEÆ*, Benth. & Hook. f. i. 753.
Griffithii, *Hook. f. ex Triana*, in *Trans. Linn. Soc.* xxviii. (1871) 75.—Ind. or.
Helferi, *Hook. f. ex Triana*, l. c.—Ind. or.
hirsutus, *Korth. in Verh. Nat. Gesch.* 250. t. 68.—Sumatra.

ANERMA, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 170 (1842) = **Solevia**, Berg. (Cyperac.).
hispidula, Schrad. ex Nees, l. c.—S. hispidula.

ANESLEA, Reichb. Consp. 46 (1828) = ANNESLEA, Andr. = **Buryala**, Salisb. (Nymph.).

ANESORHIZA, Endl. Gen. 778 (1839), sphalm. = **Annesorhiza**, Cham. & Schlecht. (Umbell.).

ANETHANTHUS, Hiern, ex Benth. & Hook. f. Gen. ii. 1025 (1876). *GESNERACEÆ*, Benth. & Hook. f. ii. 1092.
alatus, *Benth. & Hook. f. Gen.* ii. 1025.—Bras.
gracilis, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 98.—Bras.
parviflora, *Benth. & Hook. f. Gen.* ii. 1025.—Mexic.
villosa, *Benth & Hook. l. c.*—Bras.

ANETHUM, Tourn. ex Linn. Gen. ed. I. 82 (1737) = **Peucedanum**, Linn. (Umbell.).
arvense, Bess. Ind. Sem. Hort. Crem. (1820) = Carum Ridolfia.
arvense, Salisb. Prod. 168 = Peucedanum graveolens.
capense, Thunb. in Hoffm. Phytogr. Blaett. i. (1803) 26 = Carum capense.
chrysenm, *Boiss. & Heldr. Diagn.* Ser. I. x. 82.—Hispan.
cymbocarpum, DC. Prod. iv. 126 = Cymbocarpum anethoides.
dulce, DC. Cat. Hort. Monsp. 78 = Foeniculum vulgare.
erythraeum, DC. Prod. iv. 126 = Cymbocarpum erythraeum.
Foeniculum, Linn. Sp. Pl. 263 = Foeniculum vulgare.
Foeniculum, Thunb. Prod. Pl. Cap. 51 = Carum capense.
graveolens, Linn. Sp. Pl. 263 = Peucedanum graveolens.
graveolens, Ucria, Hort. Pan. 128 = Carum Ridolfia.
minus, Conan, Illustr. 20 = Foeniculum vulgare.
Panmori, Roxb. Fl. Ind. ii. 94 = Foeniculum vulgare.
Panmorium, Roxb. ex Flem. in As. Res. xi. (1810) 156 = præc.
Pastinaca, Wibel, Prim. Fl. Werth. 146 = Peucedanum Pastinaca.
pimpinellum, Ruiz & Pav. ex Urban, in Mart. Fl. Bras. xi. t. 849 = Aplum Ammi.
piperitum, Ucria, in Roem. Arch. i. 68 = Foeniculum piperitum.
pumilum, Hort. ex Steud. Nom. ed. II. i. 97 = Carum Ridolfia.
rupestre, Salisb. Prod. 168 = Foeniculum vulgare.
segetum, Linn. Mant. 219 = Carum Ridolfia.
Sowa, Roxb. ex Flem. in As. Res. xi. (1810) 156 = Peucedanum graveolens.
trifoliatum, Reichb. Hort. Beng. 86; Fl. Ind. ii. 96 = Pimpinella Heyneana.

ANETIA, Endl. Gen. 928 (1839) = **Byrsanthus**, Guill. (Sapind.).
Byrsanthus, Steud. Nom. ed. II. i. 97 = B. Brownii.

ANEULOPHUS, Benth. in Benth. & Hook. f. Gen. i. 944 (1869). *LINEÆ*.
CAUCANTHUS, Forsk. Fl. Aegypt. Arab. 91 (1775).
africanus, *Benth. l. c.*—Afr. trop.

ANEURISCUS, Presl, Symb. Bot. i. 73. t. 48 (1832) = **Symphonia**, Linn. (Guttif.).
Aubletii, Presl, l. c. 73 = S. globulifera.
caserana, Presl, l. c. t. 48 = S. globulifera.

ANGADENIA, Miers, Apocyn. S. Amer. 178 (1878). *APOCYNACEÆ*.
almadensis, *Miers, l. c.* 179.—Bras.
amazonica, Miers, l. c. 175 (= *Anisolobus amazonicus*). Berterii, *Miers, l. c.* 180.—Ins. S. Doming.
cognata, Miers, l. c. 176 (= *Anisolobus cognatus*).
coriacea, *Miers, l. c.* 177.—Guiana.
cubensis, *Miers, l. c.* 182.—Cuba.
Cururu, Miers, l. c. 178 (= *Anisolobus Cururu*).
elegans, *Miers, l. c.* 178.—Reg. Amazon.
elliptica, *Miers, l. c.* 180.—Bras.
geminata, Miers, l. c. 178 = Odontadenia geminata.
grandifolia, *Miers, l. c.* 175.—Reg. Amazon.
havanensis, *Miers, l. c.* 181.—Cuba.
hypoglauca, Miers, l. c. 179 = Odontadenia hypoglauca.
latifolia, *Miers, l. c.* 176.—Bras.
Lindeniana, *Miers, l. c.* 180.—Cuba.
majuscula, *Miers, l. c.* 174.—Reg. Amazon.
nitida, Miers, l. c. 177 = Odontadenia nitida.
pandurata, Miers, l. c. 182 = Echites pandurata.
Poeppigii, Miers, l. c. 179 = Odontadenia Poeppigii.
Pretenii, *Miers, l. c.* 182.—Guiana.
prunoides, *Miers, l. c.* 177.—Bras.
reticulata, *Miers, l. c.* 179.—Bras.
Sagraei, Miers, l. c. 181 = Echites Sagraei.
Spruceri, Miers, l. c. 178 (= *Anisolobus Spruceri*).
sylvestris, Miers, l. c. 174 = Odontadenia sylvestris.
Valenzuelana, Miers, l. c. 181 = Echites Valenzuelana.

ANGEJA, Vandelli, Fasc. Pl. 13 (1771) = Quiid ?
Vandellii, Steud. Nom. ed. II. i. 97.—Bras.

ANGELANDRA, Endl. Gen. Suppl. iii. 69 (1843) = **Engelmannia**, Torr. & Gray (Compos.).
pinnatifida, Endl. ex Walp. Rep. ii. 976 = E. pinnatifida.

ANGELANDRA, Endl. Gen. Suppl. iv. 131. 91 (1850) = **Croton**, Linn. (Euphorb.).
ellipticus, Baill. Etud. Gén. Euph. 379 = C. ellipticum.

ANGELESIA, Korth. in Ned. Kruidk. Arch. iii. (1856) 384 = **Trichocarya**, Miq. (Rosac.).
splendens, Korth. l. c. = T. splendens.

ANGELICA, [Riv.] Linn. Syst. ed. I (1735). *UMBELLIFERÆ*, Benth. & Hook. f. i. 916.
ANGELOPHYLLUM, Rupr. Rev. Umb. Kamtsch. 8 (1859).
CALLEACE, Fisch. in Hoffm. Gen. Umb. ed. II. 170 (1816).
EUSTYLIS, Hook. f. Fl. N. Zel. i. 79 (1853).
GINGIDIUM, Forst. Char. Gen. 43. t. 21 (1776).
GOMPHOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. (1841) 537.
OSTERICUM, Hoffm. Gen. Umb. 162 (1814).
alpina, Fisch. ex Besser, Enum. ed. I. 49 = sylvestris.
angustifolia, Hoffm. Gen. Umb. 162 = Peucedanum angustifolium.
anomala, Ave-Lall. in Ind. Sem. Hort. Petrop. ix. 57 = sylvestris.
aquilegifolia, Lam. Fl. Fr. iii. 452 = Siler trilobum.
Archangelica, Linn. Sp. Pl. 250 = Archangelica officinalis.
arguta, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 620.—Am. bor.
atropurpurea, Linn. Sp. Pl. ed. I. 251 = Archangelica atropurpurea.
bracteata, Roxb. in Beata. Tracts, App. 297 = Sium heleniasum.
Breweri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 348.—Calif.
Candollei, Wall. Cat. n. 889 = Selinum Candollii.
Carvifolia, Vill. Prosp. 28; Spreng. Umbellif. Prod. 16 = Selinum Carvifolia.
cheerophyllea, *Lotterm. ex Eysenh. Diss. Accur. Pl. Pruss.* 19.—Livon.
citriodora, *Hance, in Journ. Bot.* ix. (1871) 191.—China.

ANGELICA—
controversa, *Zuccag. Fl. Pedem.* ii. 11.—Italia.
Curtisii, *Buckl. in Am. Journ. Sc.* xiv. (1848) 178.—Am. bor.
dahurica, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 35.—As. or.
Dawsoni, *S. Wats. in Proc. Am. Acad.* xx. (1885) 369.—Am. bor.
decipiens, *Hook. f. Handb. New Zeal. Fl.* 98.—N. Zel.
Diefenbachii, *Benth. & Hook. f. Gen.* i. 916.—N. Zel.
diversicolor, Dulac, Fl. Hautes-Pyr. 845 = Rarulii.
dura, C. Koch, in Linnaea, xvi. (1842) 357 (= *Tommasinia purpuracea*).
ebulifolia, Lap. Hist. Abr. Pl. Pyr. 156 = sylvestris.
Fischeri, Spreng. in Roem. & Schult. Syst. vi. 605 = Silaus longifolius.
flavescens, Hoffm. Gen. Umb. 162 = sylvestris.
Fiorenti, *Franch. & Sav. ex Maxim. in Bull. Acad. Pétersb.* xix. (1874) 274.—Japon.
geniculata, *Hook. f. Handb. New Zeal. Fl.* 98.—N. Zel.
genuflexa, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 620.—Am. bor.
Gingidium, *Hook. f. Handb. New Zeal. Fl.* 97.—N. Zel.
glauca, *Edgew. in Trans. Linn. Soc.* xx. (1846) 53.—Reg. Himal.
Gmelini, Wormsk. ex DC. Prod. iv. 170 = Archangelica Gmelini.
gracilis, *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) 222.—China.
graveolens, Steud. Nom. ed. II. ii. 555 in syn. = Peucedanum graveolens.
grossecrenata, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 275.—China bor.
hakonensis, *Maxim l. c.* 277.—Japon.
heterocarpa, *Lloyd, in Bull. Soc. Bot. Fr.* vi. (1859) 709.—Hispan.
hirsuta, Muhl. Cat. 80 = Archangelica hirsuta.
inæqualis, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 194.—Japon.
integrifolia, *Walt. Fl. Carol.* 115.—Carolina.
intermedia, Schult. ex Steud. Nom. ed. I. 49 = Archangelica officinalis.
japonica, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858 & 59) 390.—Japon.
Kiusiana, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 264.—Japon.
laevigata, Fisch. Cat. Hort. Gorenk. (1812) 45 = Peucedanum latifolium.
lævis, *Y. Gray, in Ann. Sc. Nat.* Sér. II. vi. (1836) 221 (err. typ. 121).—Hispan.
lancifolia, *Spreng. Syst.* i. 916.—Hab. ?
leporina, *S. Wats. in Proc. Am. Acad.* xii. (1877) 252.—Am. bor.
Levisticum, All. Fl. Pedem. ii. 10 = Levisticum officinale.
linearifolia, *A. Gray, in Proc. Am. Acad.* vii. (1868) 347.—Calif.
litoralis, Fries, Fl. Halland. 51 = Archangelica officinalis.
lobata, *Walt. Fl. Carol.* 115.—Am. bor.
lucida, *Linn. Sp. Pl.* 251.—Canada.
Lyallii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 374.—Am. bor.
macrophylla, Schur, Enum. Pl. Transs. 262 = sylvestris.
maculata, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 72.—Ins. Kuril.
major, Gilib. Fl. Lituan. ii. 24 = Archangelica officinalis.
major, *Lag. Gen. et Sp. Nov.* 18.—Hispan.
Maximowiczii, *Benth. ex Maxim. in Bull. Acad. Pétersb.* xix. (1874) 274.—Sibir.
mexicana, *Vatke, App. Sem. Hort. Berol.* (1876) *App.* 2. n. 12.—Mexico.
minor, Gilib. Fl. Lituan. ii. 28 = sylvestris.
Miqueliana, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 276.—Japon.
mongolica, *Franch. in Nouv. Arch. Mus. Par.* Sér. II. vi. (1883) 21; *Pl. Davidian.* i. (1884) 141.—Mongol.
montana, Brot. Fl. Lusit. i. 496 = sylvestris.
nemorosa, Tenore, Fl. Nap. t. 221; Syll. 561 = sylvestris.
norvegica, *Nym. Syll. Suppl.* 98.—Norveg.
officinalis, Bernh. Syst. Verz. Erf. 170 = Peucedanum Ostruthium.

ANGELICA:—
officinalis, Moench, Meth. 81 = Archangelica officinalis.
pachycarpa, *Lange, Descr. Ic. Ill* 7. t 9.—Hispan.
pachyptera, Avé-Lall. Ind. Sem. Hort. Petrop ix 58 = sylvestris.
paludapifolia, Lam. Fl Fr iii. 451 = Levisticum officinale.
palustris, *Hoffm Gen. Umb.* 102—Europ
paniculata, Lam. Encyc. i. 172 = Trochiscanthes nodiflorus.
pinnata, *S. Wats in King, Rep.* v 126—Am bor.
polyclada, *Franch. in Bull. Soc Bot. Fr.* xxvi (1879) 86.—Japon.
polymorpha, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 185.—Japon.
pratensis, Beh. in Fisch. Hort. Gorenk. (1812), ex Bess. Enum. Pl Volh. 94 = palustris.
pratensis, J. & C. Presl, Fl. Cech. 61 = sylvestris.
pubescens, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 84.—Japon.
procera, Salisb. Prod 164 = Archangelica officinalis.
pyrenaea, Spreng. Umb. 62 = Schnuts pyrenaeum.
Razulii, *Gouan, Illustr.* 13. t. 6.—Pyren.
Razulii, All. Fl Pedem ii. 10 = sylvestris.
refracta, *F. Schmidt, Reisen Amurl.* 138.—Ins. Sachalin.
Royana, Benth. in Fisch. Hort. Gorenk. (1812), ex Bess. Enum. Pl Volh. 94 = palustris.
ruthenica, Schott, ex Ledeb. Fl. Ross. ii. 296 = sylvestris.
sachalinensis, *Maxim Prim. Fl Amur.* 127.—As. or.
sativa, Mill. Gard. Dict. ed. VIII. n. 1 = Archangelica officinalis.
saxatilis, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 739.—Sibir.
scabra, Pett, in Ann. Sc. Obs. i. 92. t. 8 = Xatardia scabra.
scotica, Lam. Encyc. i. 178 = Ligusticum scoticum.
Siebold, Miq. Ann. Mus. Bot Lugd. Bat. iii. 61 (fide Maxim. = A. pubescens, Kitagawa et daburica) — Japon.
songorica, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 590.—Kokania.
Strationana, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 164.—Afghan
sylvestris, *Linn. Sp. Pl.* 251.—Europ.; Orient; As. bor.
ternata, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 32.—As. temp.
tomentosa, *S. Wats. in Proc. Am. Acad.* xi. (1876) 141.—Calif.; Am. bor.
Tournefortiana, Cusson, ex Steud. Nom. ed. I. 50 = Razulii.
Traversii, Hook. f. Handb. N. Zeal. Fl. 739 = Aciphylla Traversii.
trilobata, Desf. ex Steud. Nom. ed. I. 464, 50 = Siler trilobum.
triquinata, Michx. Fl Bor. Am. i. 167 = Archangelica atropurpurea.
triquinata, Turcz. ex Trautv. & Mey. in Middend. Reise (Fl. Ochot. 44) = Physolophium saxatile.
ursina, *Regel, in Act. Hort. Petrop.* v. (1877) 590.—Kamtsch.
verticillaris, Linn. Mant. 217 = Peucedanum verticillare.
verticillata, *Geyer, ex Hook. Lond. Journ. Bot.* vi. (1847) 283.—Am. bor.
villosa, Lag. Gen. et Sp. Nov. 12 = sylvestris.
Wheeleri, *S. Wats. in Am. Natural.* vii. (1873) 301.—Amer. bor.

ANGELINA, Pohl, ex Tul. in Arch. Mus. Nat. Par. viii. (1855) 363 = **Siparuna**, Aubl. (Monim.).
divergentifolia, Pohl, ex Tul. l. c.—S. guianensis.

ANGELOCARPA, Rupr. in Mém. Acad. Sc. Pétersb. Sér. VII xiv. (1869) n. iv. (Sert. Tianschan.) p. 48. *UMBELLIFERAE.*
brevicaulis, *Rupr. l. c.*—As. centr.

ANGELONIA, Humb. & Bonpl. Pl. Aequin. ii. 92. t. 108 (1809). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 980.
Phylacanthus, Benth. Scroph. Ind. 1 (1835).
Physidium, Schrad. in Goett. Gel. Anz. i. (1821) 714.
Schelveria, Nees & Mart. in Flora, iv. (1821) 299.

ANGELONIA:—
Thylacantha, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1822) 45.
angustifolia, *Benth. in DC. Prod.* x. 254.—Mexic.
arguta, *Benth. l. c.* 252.—Bras.
biflora, *Benth. l. c.* 254.—Bras.
bisaccata, *Benth. l. c.* 252.—Bras.
Blanchetti, *Benth. l. c.* 253.—Bras.
campestris, *Nees & Mart. in Flora*, iv. (1821) i. 299. —Bras.
ciliata, Gardn. ex Hook. Bot. Mag. sub t. 3648 = cornigera.
ciliolata, Benth. in DC. Prod. x. 252 = Hookeriana.
Coffrana, Moric. ex Benth. in DC. Prod. x. 252 = procumbens.
cornigera, *Hook. Bot. Mag.* t. 3648.—Bras.
crassifolia, *Benth. in DC. Prod.* x. 252.—Bras.
erlostachys, *Benth. l. c.* 254.—Bras.
Gardneri, *Hook. Bot. Mag.* t. 3754.—Bras.
goyazensis, *Benth. in DC. Prod.* x. 253.—Bras.
grandiflora, *C. Morr. in Ann. Soc. Hort. Gand*, iii. (1847) 93. t. 119.—Bras.
hirta, *Cham. in Linnaea*, viii. (1833) 27.—Bras.
Hookeriana, *Gardn. ex Benth. in DC. Prod.* x. 251.—Bras.
integerrima, *Spreng. Syst. iv. Cur. Post.* 285.—Bras.
Lessingii, *Y. Kicks, in Bull. Acad. Belg.* vi. i. (1839) 309.—Bras.
linearifolia, Moric. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 242 = Blanchetti.
lobelioides, Mart. ex J. A. Schmidt, l. c. 246 = cornigera.
micrantha, *Benth. in DC. Prod.* x. 252.—Bras.
minor, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 33.—Bras.
pilosella, *Y. Kicks, in Bull. Acad. Belg.* vi. i. (1839) 309.—Cuba.
prætensis, *Gardn. ex Benth. in DC. Prod.* x. 253.—Bras.
procumbens, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1822) 45.—Bras.
pubescens, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 16.—Bras.
Humboldtiana, *Humb. & Bonpl. Pl. Aequin.* ii. 92. t. 108.—Am. austr.
serrata, *Benth. in DC. Prod.* x. 253.—Bras.
spectabilis, Moric. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 243 = eriostachys.
tomentosa, *Morc. ex Benth. in DC. Prod.* x. 254.—Bras.
verascoides, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 240 = procumbens.
viscosa, Mart. ex J. A. Schmidt, l. c. 244 = tomentosa.

ANGELOPHYLLUM, Rupr. in Beitr. Pfl. Russ. Reich. xi. (Rev. Umb. Kamtsch.) 8 = **Angelica**, Linn. (Umbell.).
daharicum, Rupr. l. c. 9 = Angelica sylvestris.
ursinum, Rupr. l. c. 8 = Angelica ursina.

ANGELOPOGON, Poepp. ex Poepp. & Endl. Gen. et (Sp. i. 1 (1835) = **Myrodendron**, Banks & Soland. Santal.).

ANGERVILLA, Neck. Elem. i. 351 (1790) = **Stemodia**, Linn. (Scrophul.).

ANGIANTHUS, Wendl. Coll. ii. 21. t. 48 (1809). *COMPOSITAE*, Benth. & Hook. f. ii. 519.
Cephalosorus, A. Gray, in Hook. Kew Journ. iii. (1851) 102.
Chrysocoryne, Endl. in Bot. Zeit. i. (1843) 457.
Crossolepsis, Benth. in Enum. Pl. Hueg. 61 (1837).
Cylindrosorus, Benth. l. c. 62 (1837).
Dithyrostegia, A. Gray, in Hook. Kew Journ. iii. (1851) 100.
Epitriche, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 74.
Eriocladium, Lindl. Swan Riv. App. 24 (1839).
Gamozygia, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 75.
Hyalochlaena, A. Gray, in Hook. Kew Journ. iii. (1851) 101.
Ogcerostylus, Cass. in Dict. Sc. Nat. xlix. 221 (1827).
Oxerostylus, Steud. Nom. ed. II. ii. 242 (1841), sphalm.

ANGIANTHUS:—
Phyllocalymma, Benth. in Enum. Pl. Hueg. 61 (1837).
Piptostemma, Turcz. in Bull. Soc. Nat Mosc. xxiv. (1851) i. 191.
Pleurophappus, F. Muell. in Trans Viet. Inst. (1856) 87.
Pogonolepis, Steetz, in Lehm. Pl. Preiss. i. 440 (1845).
Sloskebus, Labill. Pl. Nov. Holl. ii. 57. t. 209 (1806), nomen prius.
Skiarhophorus, DC. ex Lindl. Introd. Nat. Syst. ed. II. 260 (1836).
Stylocoryne, Spreng. Syst. iii. 356, 451 (1826).
amplexicaulis, *Benth. Fl. Austral.* iii. 568.—Austral.
brachypappus, *F. Muell. in Trans. Phil. Soc. Vict. i.* (1855) 44.—Austral.
codonopappus, *F. Muell. Fragm.* ix. 2.—Austral.
Cunninghamii, *Benth. Fl. Austral.* iii. 565.—Austral.
damiseus, *Benth. l. c.* 567.—Austral.
Drummondii, *Benth. l. c.* 566.—Austral.
eriocephalus, *Benth. l. c.* 567.—Austral.
flavescens, Steetz, in Lehm. Pl. Preiss. i. 456 = tomentosus.
globifer, *Benth. Fl. Austral.* iii. 567.—Austral.
humifusus, *Benth. l. c.* 568.—Austral.
microcephalus, *Benth. l. c.* 566.—Austral.
micropoides, *Benth. l. c.* 565.—Austral.
Milnei, *Benth. l. c.* 564.—Austral.
myosuroides, *Benth. l. c.* 563.—Austral.
phyllocephalus, *Benth. l. c.* 565.—Austral.
platycephalus, *Benth l. c.* 566.—Austral.
pleurappus, *Benth. l. c.* 563.—Austral.
plumiger, *Benth. l. c.* 565.—Austral.
Preissianus, *Benth. l. c.* 566.—Austral.
pusillus, *Benth. l. c.* 564.—Austral.
pygmaeus, *Benth. l. c.* 567.—Austral.
strictus, *Benth. l. c.* 568.—Austral.
tenellus, *Benth. l. c.* 564.—Austral.
tomentosus, *Wendl. Coll.* ii. 21. t. 48 (1809).—Austral.

ANGIOPETALUM, Reinw. in Syll. Ratisb. ii. (1828) 7. *MYRSINEAE.*
punctatum, *Reinw. l. c.*—Java.

ANGKALANTHUS, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 98. *ACANTHACEAE.*
oligophylla, *Balf. f. l. c.* 89.—Socotra.

ANGOLAEA, Wedd. in DC. Prod. xvii. 300 (1873). *PODOSTEMACEAE*, Benth. & Hook. f. iii. 111.
fluitans, *Wedd. l. c.*—Afr. trop.

ANGOLAM, Adans. Fam. ii. 85 (1763) = **Alangium**, Linn. (Cornac.).

ANGOLAMIA, Scop. Introd. 107 (1777) = prasc.

ANGOPHORA, Cav. Ic. iv. 21. t. 388 (1797). *MYRTACEAE*, Benth. & Hook. f. i. 707.
cordifolia, *Cav. l. c.*—Austral.
floribunda, Sweet, Hort. Brit. ed. II. 209 = Eugenia Smithii, Ventenatii.
intermedia, *DC. Prod.* iii. 229.—Austral.
lanceolata, *Cav. Ic.* iv. 22. t. 387.—Austral.
subvelutina, *F. Muell. Fragm.* i. 31.—Austral.
Woodsiana, *F. M. Bailey, in Proc. Linn. Soc. N. S. W.* vi. (1881) 148.—Austral.

ANGORCHIS, Nees, in R. Br. Verm. Schrift. ii. 438, in adnot. (1826) = **Angraecum**, Bory (Orchid.).

ANGOSTURA, Roem. & Schult. Syst. iv. 188 (1819) = **Galipea**, Aubl. (Rutac.).
Cuspare, Roem. & Schult. l. c. = G. trifoliata.

ANGOSTYLIS, Benth. in Hook. Kew Journ. vi. (1854) 238. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 327.
longifolia, *Benth. l. c.*—Brus.

ANGRAECUM, Bory, Voy. i. 360. t. 19 (1804). *ORCHIDEAE*, Benth. & Hook. f. iii. 503.
Aeranga, Reichb. f. in Flora, xlviii. (1865) 190.
Arborium, Rasmpf. ex Spreng. Syst. iii. 674 (1826).
Angorchis, Nees, in R. Br. Verm. Schrift. ii. 438 in adnot. (1826).

ANGRAECUM :—
Listrostachys, Reichb. f. in Bot. Zeit. x. (1852) 880.
Microcaelia, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 235 (1851).
Plectrelminthus, Rafin. Fl. Tellur. iv. 42 (1836)
alcicorne, *Reichb. f. Otia Bot. Hamb* 117.—Afr. trop.
aphyllum, *Thou. Orch. Afr.* t. 78.—Ins. Masc.
apiculatum, *Hook. Bot. Mag.* t. 4159.—Afr. trop.
arachnopus, *Reichb. f. in Bonplandia*, ii. (1854) 93.—Afr. trop.
arcuatum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 204.—Afr. austr.
arcuatum, Lindl. ex Hook. f. in Bot. Mag. t. 5589 = Chailluanum.
armeniacum, *Lindl. in Bot. Reg.* (1839) *Misc.* 67 — Afr. trop.
articulatum, *Reichb. f. in Gard. Chron.* (1872) 78.—Madag.
ashantense, *Lindl. in Bot. Reg.* (1843) *Misc.* 56.—Afr. trop.
bicaudatum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 205.—Afr. austr.
bilobum, *Lindl. Bot. Reg.* (1840) *Misc.* 69.—Afr. occ.
Bootoni, *Reichb. f. Otia Bot. Hamb* 117.—Ins Comor
bracteosum, *Balf f. & S Moore, in Journ Bot* xiv. (1876) 208.—Ins. Borbon
brevifolium, *Lindl. Bot. Reg.* (1840) sub t. 68 — Bras.
Brongniartianum, *Reichb. f. ex Linden, Pescatoria*, i. t. 16.—Ins. Mascar.
Burchellii, *Reichb. f. in Flora*, i. (1867) 117.—Afr. austr.
Calceolus, *Thou. Orch. Afr* t. 78.—Ins. Mascar.
camptoplectron, *Reichb. f. in Bonplandia*, iii. (1855) 220.—Hab. ?
capense, Lindl. Gen. et Sp. Orch. 248 = Mystacidium filicorne.
capitatum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 187.—Afr. trop.
carinatum, Kostel. Allg. Med.-pharm. Fl i. 250 = Eulophia virens.
carpophorum, *Thou. Orch. Afr.* t. 76 —Afr. trop.
caudatum, *Lindl. Bot. Reg.* t. 1844.—Afr trop.
caulescens, Thou. Orch. Afr. t. 75 = Mystacidium caulescens..
Chailluanum, *Hook f. Bot. Mag* t. 5589.—Afr. trop.
Chlocchistae, *Reichb. f. in Linnaea*, xx. (1847) 678.—Natal.
Christyanum, *Reichb. f. in Gard. Chron.* (1880) i. 208.—Hab. ?
citratum, *Thou. Orch. Afr.* t. 61.—Madag
clandestinum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 206.—Afr. trop.
clavigerum, *Rudl. in Journ. Linn. Soc.* xxi. (1885) 485.—Madag.
conchiferum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 205.—Afr. austr.
cornutum, *Reichb. f. in Flora*, lxviii. (1885) 588.—Ins. Comor.
Cowanii, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 484.—Madag.
crassum, *Thou. Orch. Afr.* tt. 70 et 71.—Madag.
cryptodon, *Reichb. f. in Gard. Chron.* (1883) i. 307.—Madag.
cucullatum, *Thou. Orch. Afr.* t. 48.—Ins. Mascar.
cuticiferum, *Reichb. f. in Flora*, lxviii. (1885) 588.—Ins. Comor.
descendens, *Reichb. f. in Gard. Chron.* (1882) i. 568.—Hab. ?
distichum, *Lindl. Bot. Reg.* t. 1781.—Afr. trop.
eburneum, Lindl. l. c. (1839) xviii. 1522 = superbum.
eburneum, *Bory, Voy.* i. 359. t. 19.—Ins. Borbon.
elatum, Thou. Orch. Afr. t. 79 = Cryptopus elatus.
Ellisii, *Reichb. f. in Flora*, lv. (1872) 278.—Madag.
expansum, *Thou. Orch. Afr.* t. 57.—Ins. Mascar.
falcatum, *Lindl. Coll. Bot.* 35.—Chins.
fasciola, *Lindl. Bot. Reg.* xxvi. (1840) sub t. 68.—Demerara.
fastuosum, *Reichb. f. in Gard. Chron.* (1881) ii. 748, 844.—Madag.
filicorne, *Thou. Orch. Afr.* t. 53.—Ins. Mascar ; Madag.
filiforme, *Lindl. Gen. et Sp. Orch.* 248.—Ind. occ.
floribundum, *Reichb. f. in Gard. Chron.* (1885) i. 787 ; *in Flora*, lxviii. (1885) 330.—Ins. Comor.
fragrans, *Thou. Orch. Afr.* t. 54.—Ins. Mascar.
funale, Lindl. in Gard. Chron. (1846) 135 = Dendrophylax funalis.

ANGRAECUM :—
fuscatum, *Reichb. f. in Gard. Chron.* (1882) ii. 488.—Madag.
Gallesandrae, *Reichb. f. in Flora*, xlviii. (1865) 189.—Afr. occ.
Gilpinae, *Reichb. f. & S. Moore, in Journ. Linn. Soc.* xvi. (1877) 206.—Madag.
gladiifolium, *Thou. Orch. Afr.* t. 58.—Ins. Mascar. ; Madag.
globulosum, Hochst. ex A. Rich. Tent. Fl. Abyss ii. 235 = Cuyonianum.
glomeratum, *Ridl. in Gard. Chron.* (1885) ii. 678.—Afr. trop
gracile, *Thou. Orch. Afr.* t. 77.—Ins. Mascar.
Grantii, Bateman, ex Baker, in Trans. Linn. Soc. xxix. (1876) 158 = Kotschyi.
Guyonianum, *Reichb. f. in Linnaea*, xxii. (1849) 865.—Abyss.
Hildebrandtii, *Reichb. f. in Gard. Chron.* (1878) i. 725.—Afr. trop.
Humblotii, *Reichb. f. in Flora*, lxviii. (1885) 381.—Ins Comor.
hyaloides, *Reichb. f. in Gard. Chron.* (1880) i. 264.—Madag.
Ichneumoneum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 136.—Afr. occ.
imbricatum, *Lindl. l. c.* 137.—Afr. occ.
implicatum, *Thou. Orch Afr.* t. 58.—Madag
inapertum, Thou l. c. t 59 = Mystacidium inapertum.
infundibulare, *Lindl. in Journ. Linn. Soc.* vi. (1862) 136.—Afr. occ.
jamaicense, *Reichb. f & Wullschl. in Walp. Ann.* vi. 901 — Jamaic.
japonicum, *Linden, Illustr. Hortic* xxviii (1881) 64.—Japan.
Kotschyi, *Reichb. f. in Gard. Chron.* (1880) ii. 456.—Afr trop
Lanzerpti, *Reichb. f. in Nederl. Kruidk. Arch* iv. (1859) 316.—Caracas.
Leonis, Hort.—Cf. Gard. Chron. (1885) ii. 80–81 = Aerantthus Leonis
lepidotum, *Reichb. f. l. c.* (1880) i. 806, in obs.—Hab.?
Lindenii, *Lindl. l. c.* (1846) 135.—Cuba.
luridum, *Lindl. ex Steud. Nom.* ed. II. i. 98.—Afr. trop.
maculatum, Lindl. Coll. 15 = Eulophia maculata.
maxillariodes, *Ridl. in Journ Linn. Soc.* xxi. (1885) 479.—Madag
melanthianum, *Reichb. f. Otia Bot Hamb.* 117.—Afr. trop.
micranthum, *Lindl. Bot. Reg.* xxi. t. 1772 —Afr. trop.
modestum, *Hook f. Bot. Mag* t. 6693.—Madag.
monodon, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 102.—Afr. occ.
monophyllum, A. Rich. in Mém. Soc. Hist. Nat. Par iv. (1828) 58 = Eulophia monophylla.
multiflorum, Thou. Orch. Afr. t. 74 = Mystacidium caulescens.
Mystacidii, *Reichb. f. in Linnaea*, xx. (1847) 677.—Natal.
odoratissimum, *Reichb. f. in Bonplandia*, iv. (1856) 326.—Hab. ?
organense, *Reichb. f. in Walp. Ann.* vi. 901.—Bras.
ornithorhynchum, *Lindl. Bot Reg.* xxvi. (1840) sub t 68.—Bras.
palmatum, Thou. Orch. Afr. t. 69 = palmiforme.
palmiforme, *Thou. l. c.* 68.—Ins. Mascar.
parviflorum, *Thou. l. c.* t. 60.—Ins. Mascar.
parviflorum, *Ayres, ex Baker, Fl. Maurit.* 357.—Ins. Mascar.
pectinatum, *Thou. Orch. Afr.* t. 51.—Ins Mascar.; Madag.
pellucidum, *Lindl. in Journ. Linn. Soc.* vi (1862) 184.—Afr. occ.
pertusum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 205.—Afr. occ.
Pescatorianum, *Lindl. in Journ. Hort. Soc.* iv. (1849) 263.—Ins. Borbon.
physophorum. *Reichb. f. Otia Bot. Hamb.* (1881) 77 et 449.—Ins. Mascar.
Poeppigii, *Reichb. f. in Linnaea*, xxii. (1849) 858.—Cuba.
polystachyum, *Lindl. Bot. Reg.* xxvi. (1840) sub t. 68.—Peruv.
polystachyon, *A. Rich. in Mém. Hist. Nat. Par.* iv. (1828) 56 = Aëonia polystachya.
pugioniforme, *Klotzsch, Ind. Sem. Hort. Berol. App.* (1851) 12.—Venezuela.

ANGRAECUM :—
pusillum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 206.—Afr. austr.
pygmaeum, *Linden, Illustr. Hortic.* xxviii. (1881) 64.—Japon.
ramosum, *Thou. Orch. Afr.* t. 59 —Ins. Mascar.
rectum, *Thou l. c.* 55.—Ins. Mascar.
recurvum, *Thou l. c.* t. 56.—Ins. Mascar.
rhipsalisocium, *Reichb. f. in Flora*, xlviii. (1865) 189.—Afr. occ.
Rohrii, *Reichb. f. Otia Bot Hamb.* 117.—Abyss.
rostellare, *Reichb f. in Flora*, lxviii. (1885) 380 ; *et in Gard. Chron.* (1885) i. 726 —Ins. Comor.
rostratum, *Ridl. in Journ. Linn Soc.* xxi (1885) 485.—Madag.
Rutenbergianum, *Kraenzl. in Bremen Abh.* vii. (1882) 257.—Afr. trop.
sacciferum, *Lindl. in Hook. Comp Bot. Mag.* ii. (1836) 205.—Afr. austr.
Schiedei, Reichb. f. in Linnaea, xxii. (1849) 857 = Campyloceutron Schiedei.
Schimperianum, *Reichb. f. in Walp. Ann.* iii. 573.—Abyss.
Scottianum, *Reichb f in Gard. Chron.* (1878) ii. 556.—Ins. Comor.
Sedens, Hort. ex Nichols. Dict. Gard. i. 79 = arcuatum.
Sellowii, *Reichb f. in Linnaea*, xxii. (1849) 857.—Bras.
sesquipedale, *Thou. Orch Afr.* tt. 66 et 67.—Madag.
spathilabrum, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 478.—Madag.
stratum, *Thou. Orch. Afr.* t. 72.—Ins. Mascar.
subulatum, *Lindl. in. Hook. Comp Bot. Mag.* ii. (1836) 206.—Afr occ.
superbum, *Thou. Orch. Afr.* tt. 62 et 64.—Madag
tenue, *Lindl. Bot. Reg* xxvi. (1840) sub t. 68.—Bras.
teretifolium, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 484 —Madag.
tridens, *Lindl. l. c* vi. (1862) 135.—Afr. trop.
trilentatum, *Herv. Thes. Cap.* ii. 6.—Afr. austr.
triquetrum, *Thou. Orch Afr.* t. 49.—Ins. Mascar.
vagans, *Lindl. in Journ. Linn. Soc.* vi. (1862) 135.—Afr. occ.
virens, Lindl. Bot. Reg. (1847) t. 19 = eburneum.
ve-istum, *Lindl. l. c.* (1848) *Misc.* 6.—Afr. trop.
Welgelii, *Reichb. f. in Linnaea*, xxii. (1849) 857.—Guiana
xylopus. *Reichb f. in Flora*, lxviii. (1885) 538.—Ins. Comor.
yuccaefolium, *Boj. Hort. Maurit.* 218.—Ins Maurit.
zeylanicum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 40.—Ind. or.

ANGUILLARIA, R. Br. Prod. 278 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 824.
australis, F. Muell. Fragm. vii. 74 = dioica.
biglandulosa, R. Br. Prodr 278 = dioica.
capensis, Spreng Syst. ii. 146 = Androcymbium leucanthum.
caricina, *Schult. f. Syst.* vii. 1539.—Ind or.
cochinchinenas, *Spreng. Syst.* ii. 147.—Cochinch.
densiflora, *Benth. Fl. Australe*, vii. 29.—Austral.
dioica, *R. Br. Prod.* 278.—Austral.
Heynoana, Wall. Cat. n. 5088 = Ipbigenia indica.
indica, R. Br. Prod. 278 = Ipbigenia indica.
monantha, Endl. in Lehm. Pl. Preiss. ii 45 = dioica.
pygmaea, Endl. l. c. = Wurmbea pygmaea.
racemosa, Schult. f. Syst. vii. 1539 = Ipbigenia indica.
tenella, Endl. in Lehm. Pl. Preiss. ii. 45 = Wurmbea tenella.
uniflora, R. Br. Prod. 278 = dioica.

ANGUILLARIA, Gaert. Fruct. i. 372 (1788) = **Ardisia**, Sw. (Myrsin.).
bahamensis, Gaertn. l. c. 373 = Ar. excelsa.
Barthesia, Lam. Illustr. ii. 100 = Ar. Barthesia.
chrysophylla, Lam. l. c. = Ar. serrulata.
coriacea, Poir. Encyc. vii. 686 = Ar. coriacea.
crenulata, Poir. l. c. 688 = Parathesis crenulata.
excelsa, Lam. Illustr. ii. 110 = Ar. excelsa.
Icacorea, Lam. l. c. 109 = Ar. acuminata.
lateriflora, Lam. l. c. = Ar. lateriflora.
latifolia, Steud. Nom. ed II. i. 98 = Ar. lanifolia.
laurifolia, Lam. Illustr. ii. 109 = Ar. lanifolia.
parasitica, Lam. l. c. = Cremmadenia parasitica.
pyramidalis, Cav. Ic. vi. 1 = Ar. pyramidalis.
serrata, Cav. l. c. 2 = Ar. Cavanillesii.
serrulata, Poir. Encyc. vii. 686 = Ar. serrulata.
solanacea, Poir. l. c. 688 = Ar. humilis.

ANGUILLARIA :—
***tinifolia*, Lam.** Illustr. i. 109 = Ardisia tinifolia.
***zeylanica*, Gaertn.** Fruct. i. 378. t. 77. f. 1 = Ar. Moonii.

ANGUINA, Mich. ex Linn. Syst. ed. I. (1735) = **Trichosanthes**, Linn. (Cucurb.).

ANGUINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. 160 (1869) = **Allium**, Tourn. (Liliac.).
***Victoriale*,** Fourr. l. c. = Allium Victorialis.

ANGULOA, Ruiz & Pav. Prod. Fl. Per. 118. t. 26 (1794). *ORCHIDEAE*, Benth. & Hook. f. iii. 542.
Clowesii, *Lindl. Bot. Reg.* (1844) *Misc.* 26.—Venezuela.
dubia, *Reichb. f. in Gard. Chron.* (1882) i. 764.—N. Granat.
eburnea, *Hort. ex Nichols. Dict. Gard.* i. 79.—N. Granat.
grandiflora, H. B. & K. Nov. Gen. et Sp. i. 245 = Stanhopea Bucephalus.
Hernandezii, Kunth, Syn. i. 339 = Stanhopea tigrina.
Hohenlohii, *Morr. in Belg. Hortic.* lii. (1858) 201. t. 21.—N. Granat.
luridа, Link, in Preuss. Gart. Ver. i. 289. t. 6 = Catasetum luridum.
maclа x, *Reichb. f. in Gard. Chron.* (1881) ii. 33.
parpurea, *Lindeu, in Illustr. Hortic.* xviii. (1881) t. 427.—Venezuela.
Ruckeri, *Lindl. Bot. Reg.* (1846) t. 41.—N. Granat.
squalida, *Poepp. & Endl. Nov. Gen. et Sp.* i. 74.—Peruv.
superba, H. B. & K. Nov. Gen. et Sp. i. 243. t. 93 = Acineta Humboldtii.
uniflora, *Ruiz & Pav. Syst.* 228.—Peruv.
virginalis, Hort.—Cf. Gard. Chron. (1851) 392 = uniflora.

ANGURIA, [Tourn.] Linn. Syst. ed. I. (1735). *CUCURBITACEAE*, Benth. & Hook. f. i. 833.
aculeolata, Schlecht. in Linnaea, xxiv. (1851) 748 = Wilbrandia a verticillata.
affinis, Schlecht. l. c. 760 = pedata.
Arrabidae, Schlecht. l. c. 786 = Guranía multiflora.
bignoniacea, Poepp. & Endl. Nov. Gen. et Sp. ii. 58 = triphylla, Guranía cissoides.
Boissieriana, *Cogn. in DC. Monog. Phan.* iii. 673.—Peruv.
capitata, Poepp. & Endl. Nov. Gen. et Sp. ii. 52 = Guranía capitata.
cissoides, Benth. in Hook. Kew Journ. ii. (1850) 242 = Guranía cissoides.
Citrullus, Mill. Gard. Dict. ed. VIII. add. = Citrullus vulgaris.
dentata, Schlecht. in Linnaea, xxiv. (1851) 713 = pedata.
dubia, Hook. & Arn. Bot. Beech. Voy. 292 = Cayaponia attenuata.
eriantha, Poepp. & Endl. Nov. Gen. et Sp. ii. 52. t. 169 = Guranía eriantha.
eriantha, Scem. Bot. Voy. Herald, 127 = Guranía Seemanniana.
Gaudichaudiana, Schlecht. in Linnaea, xxiv. (1851) 758 = Wilbrandia verticillata.
glomerata, Eggers, Fl. St. Croix & Virgin. Isl. 55 = Corallocarpus glomeratus.
grandiflora, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1877).—Bras.
guianensis, Klotzsch, in Schomb. Reis. Brit. Guian. iii. 988 = Guranía guianensis.
heterophylla, Willd. ex Schlecht. in Linnaea, xxiv. (1851) 772 = Guranía cissoides.
integrifolia, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 9.—Bras.
indica, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
Jacquiniana, Schlecht. in Linnaea, xxiv. (1851) 708 = trilobata.
Kegeliana, Schlecht. l. c. 776 = Guranía Kegeliana.
Knethiana, *Schlecht. l. c.* 762.—Bras.
lacviosa, Schlecht. l. c. 755 = Apodanthera laciniosa.
leptantha, Schlecht. l. c. 779 = Helmontia leptantha.
lobata, *Linn. Amoen. Acad.* viii. 268.—Guian.
longipedunculata, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1877).—Mexico.
Mahoyana, Leon. in Fl. des Serres, Sér. I. iii. (1847) 223 = Guranía Mahoyana.

ANGURIA :—
Meyeniana, Schlecht. in Linnaea, xxiv. (1851) 750 = Wilbrandia verticillata.
multifidа, Mig. in Linnaea, xviii. (1846) 366 = Guranía multifida.
obtusiloba, Schlecht. in Linnaea, xxiv. (1851) 744 = (vitis sp.).
Ottoniana, *Schlecht. l. c.* 758.—Cuba.
pallida, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1877) 10, 22.—Ecuador.
pedata, *Vaca. Enum. Pl. Carib.* 3.—Ins. S. Doming.
pedata, Willd. ex Cogn. in DC. Monog. Phan. iii. 838 = Cyclanthera leptostachya.
pedatiserta, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 10 = Cyclanthera pedata.
pedisecta, *Nees & Mart. l. c.* = Apodanthera pedisecta.
Plumieriana, *Schlecht. in Linnaea*, xxiv. (1851) 709. —Ins. Martinic.
Poeppigiana, Schlecht. l. c. 769 = Dieudonnaea rhizantha.
polyanthos, Klotzsch, in Schomb. Reis. Brit. Guian. iii. 988 = Guranía Klotzschiana, G. subumbellata.
polyphylla, Schlecht. in Linnaea, xxiv. (1851) 712 = pedata.
rhizantha, Poepp. & Endl. Nov. Gen. et Sp. ii. 52. t. 171 = Dieudonnaea rhizantha.
rosea, H. B. & K. Nov. Gen. et Sp. ii. 122 = trilobata. Schomburgkiana, *Schlecht. in Linnaea*, xxiv. (1851) 765.—Guiana.
Sellowiana, Schlecht. l. c. 746 = Guranía Sellowiana.
Sellowiana, Wawra, Bot. Maximil. Reise, i. 57. t. 52 = Guranía Wawraei.
sinuata, Benth. in Hook. Kew Journ. ii. (1850) 242 = Guranía sinuata.
speciosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 51. t. 169 = Guranía speciosa.
spinulosa, Poepp. & Endl. l. c. 52. t. 170 = Guranía spinulosa.
subumbellata, Mig. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Select.) 108 = Guranía subumbellata, *M. Roem. Syn. Pepon.* i. 26.—Bras.
trifoliata, Aubl. Fl. Guian. ii. 842 = Guranía cissoides.
trifoliata, *Linn. Sp. Pl.* ed. II. 1376.—Ins. S. Domin.
trifolvata, Vell. Fl. Flum. x. t. 2 = ternata.
trilobata, *Jacq. Enum. Pl. Carib.* 31.—N. Granat.
trilobata, Vell. Fl. Flum. x. t. 1 = Guranía multiflora.
triphylla, *Mig. in Linnaea*, xix. (1847) 186.—Guian.
umbrosa, *H. B. & K. Nov. Gen. et Sp.* ii. 121.—Mexic.
Wagneriana, Schlecht. in Linnaea, xxiv. (1851) 785 = Guranía Wagneriana.
Warmingiana, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1877) 21.—Bras.
Warmingii, *Cogn. l. c.*—Bras.
Warscewiciana, Hort. Belg. ex Cogn. l. c. 18 = seq.
Warscewiczii, *Hook. f. in Bot. Mag.* t. 5804.—Mexic.

ANGUSTINEA, A. Gray, Bot. U. St. Expl. Exped. i. 598, in syn. (1854) = AUGUSTINEA, St. Hil. & Naud. = **Miconia**, Ruiz & Pav. (Melast.).

ANHALONIUM, Lem. Cact. Nov. Gen. et Sp. 1. (1839) = **Mammillaria**, Haw. (Cact.).
arcolosum, Lem. Illustr. Hortic. vi. (1859) Misc. 85.—Mexic.
elongatum, Salm-Dyck, Cact. Hort. Dyck. (1848) 15 = M. elongata.
Engelmanni, Lem. Cact. 42 = M. fissurata.
fissuratum, Monv. Cat. ex Forst. Cact. Handb. ed. II. i. 332 = sulcatum.
fissuratum, Engelm. Cact. Mex. Bound. 76 = M. fissurata.
Ketchubeyi, Lem. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 5 = Mammillaria sulcata.
Ketschubeyanum, Lem. Cact. 41; Illustr. Hortic. xvi. (1869) Misc. 72.—Mexic.
prismaticum, Lem. Cact. Nov. Gen. et Sp. 1 = M. prismatica.
pulchelligerum, Lem. Illustr. Hortic. xvi. (1869) 72; Hort. Monv. i. 275 = Mammillaria elongata.

ANHALONIUM :—
retusum, Salm-Dyck, Cact. Hort. Dyck. (1845) 15 = Mammillaria prismatica.
sulcatum, Salm-Dyck, l. c. ed. II. 5, 78.—Mexic.
Williamsii, Lem. ex Forst. Handb. Cact. ed. II. i. 283 = Echinocactus Williamsii.

ANIA, Lindl. in Wall. Cat. n. 8740–41; et Gen. et Sp. Orch. 129 (1831) = **Tainia**, Blume (Orchid.).
angustifolia, Lindl. in Wall. Cat. n. 8740.—Ind. or.; China.
bicornis, Lindl. in Bot. Reg. (1843) Misc. 87 = T. bicornis.
latifolia, Lindl. in Wall. Cat. n. 8741 = T. latifolia.
latifolia, Wight, Ic. Pl. lii. 914 = T. bicornis?
maculata, Thw. Enum. Pl. Zeyl. 301 = (Chrysoglossi sp.).—Zeylan.

ANIBA, Aubl. Pl. Gui. i. 327 (1775) = **Ocotea**, Aubl. (Lauria.).
guianensis, Aubl. l. c. = O. commutata.

ANICTOCLEA, Nimmo, in J. Grah. Cat. Pl. Bomb. 253 (1839) = **Tetrameles**, R. Br. (Datisc.).
Grahamiana, Nimmo, l. c. = T. nudiflora.

ANIDRUM, Neck. Elem. i. 188 (1790) = **Bifora**, Hoffm. (Umbell.).

ANIGOSIA, Salisb. in Trans. Hort. Soc. i. (1812) 327 = **Anigozanthos**, Labill. (Haemodor.).
flavida, Salisb. l. c. = Anigozanthos flavida.

ANIGOZANTHOS, Labill. Voy. i. 409. t. 22 (1798). *HAEMODORACEAE*, Benth & Hook. f iii. 676.
ANIGOSIA, Salisb. in Trans. Hort. Soc. i. (1812) 327.
SCHWAEGRICHENIA, Spreng. Syst ii. 9b (1825).
bicolor, *Endl. in Lehm. Pl. Preiss.* ii 26.—Austral.
coccinea, Paxt. Bot. Mag. v. 271 = flavida.
flavescens, Stend. Nom. ed. II. i. 98 = flavida.
flavida, [*DC. in*] *Red. Lil.* t. 176.—Austral.
fuliginosa, Hook. Bot. Mag. t. 4291 = Macropodia fumosa.
grandiflora, Salisb. Parad. Lond. t. 97 = flavida.
humilis, *Lindl. Swan Riv. App.* 46. t. 6 B.—Austral.
Manglesii, *D. Don, in Sweet, Brit. Fl. Gard.* Ser. II. t. 265.—Austral.
Manglesii, Maund. Botanist, t. 67 = flavida.
minima, Lehm. Pl. Preiss. ii. 274 = Preissii.
Preissii, *Endl. in Lehm. l. c.* 26.—Austral.
pulcherrima, *Hook. Bot. Mag.* t. 4180.—Austral.
rufa, *Labill. Voy.* i. 409. t. 22.—Austral.
tyrianthina, Hook. Bot. Mag. t. 4507 = rufa.
viridis, *Endl. in Lehm. Pl. Preiss.* ii. 25.—Austral.

ANILEMA, Kunth, Enum. Pl. iv. 64 (1843) = **Aneilema**, R. Br. (Commel.).

ANIOTUM, Soland. ex Endl. Gen. 822 (1837) = **Inocarpus**, Forst. (Legum.).

ANISACANTHA, R. Br. Prod. 510 (1810). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 62.
bicornis, F. Muell. Fragm. viii. 14 = Sclerolaena bicornis.
bicuspis, *F. Muell. in Trans. Vict. Inst.* (1855) 183.—Austral.
Birchii, *F. Muell. Fragm.* viii. 168.—Austral.
brevicuspis, *F. Muell. l. c.* iv. 150.—Austral.
diacantha, Nees, in Lehm. Pl. Preiss. i. 685 = Sclerolaena diacantha.
divaricata, *R. Br. Prod.* 410.—Austral.
Drummondii, *Benth. Fl. Austral.* v. 198.—Austral.
echinopsila, *F. Muell. Fragm.* viii. 14.—Austral.
erinacea, *Moq. in DC. Prod.* xiii. ii. 122.—Austral.
glabra, *F. Muell. ex Benth. Fl. Austral.* v. 200.—Austral. Mount.
gracilicuspis, *F. Muell. Fragm.* ii. 170.—Austral.
inermis, A. Cunn. ex Moq. in DC. Prod. xiii. ii. 122 = muricata.
hamitropidon, F. Muell. in Trans. Vict. Inst. (1855) 133 = Sclerolaena diacantha.
lanicuspis, F. Muell. Fragm. ii. 170 = Sclerolaena lanicuspis.
muricata, *Moq. Chenop. Enum.* 84.—Australia.
quinqueaculeata, Muell. in Trans. Vict. Inst. (1855) 134 = muricata.
tricuspis, *F. Muell. l. c.* 188.—Austral.

ANISACANTHUS, Nees, in Linnæa, xvi. (1842) 207. *ACANTHACEÆ*, Benth. & Hook. f. ii. 1117.
Bierbaumia, Kostel. Ind. Fl. Hort. Prag. (1844) 21.
Greggii, *A. Gray, Synop. Fl. N. Am.* ii. 328.—Mexic.
juncea, *Hemsl. Biol. Centr. Am. Bot.* 522.—Mexic.
pumilus, *Nees, in DC. Prod.* xi. 445.—Mexic.
Thurberi, *A. Gray, Synop. Fl. N. Am.* ii. 328.—Am. bor.
virgularis, *Nees, in DC. Prod.* xi. 445.—Mexic.
Wrightii, *A. Gray, Synop. Fl. N. Am.* ii. 328.—Am. bor.

ANISACTIS, Dulac, Fl. Hautes-Pyr. 347 (1867) = **Carum**, Linn. (Umbell.).
segetalis, Dulac, l. c. = C. Petroselinum.

ANISADENIA, Wall. Cat. n. 1510 (1830). *LINEAE*, Benth. & Hook. f. i. 348.
bhazyana, Griff. Notul. . 534 = saxatilis.
pubescens, *Grif. l. c.* 535.—Reg. Himal.
saxatilis, *Wall. Cat.* n. 1510.—Reg. Himal.

ANISANDRA, Bartl. in Lehm. Pl. Preiss. i. 361 (1845) = **Microcorys**, R. Br. (Labiat.).
glabra, Bartl. l. c. = M. glabra.

ANISANDRA, Planch. ex Oliver, Fl. Trop. Afr. i. 348 in syn. (1868) = **Ptychopetalum**, Benth. (Olacin.)
nigrescens, Planch. l. c. = P. sarcpa.

ANISANTHA, C. Koch, in Linnæa, xxi. (1848) 394 = **Bromus**, Linn. (Gramm.).
pontica, C. Koch, l. c. = B. tectorum.

ANISANTHERA, Griff. Notul. iv. 100 (1854) = **Adenosma**, R. Br. (Scrophul.).

ANISANTHERA, Rafin. Fl. Tellur. iii. 80 (1836) = **Casonia**, Savi (Borag.).
ciliata, Rafin. l. c. = C. Rauwolfii.

ANISANTHERA, Rafin. Fl. Tellur. ii. 60 (1836) = **Crotalaria**, Tourn. (Legum.).
hastata, Rafin. l. c. 61 = C. verrucosa.
versicolor, Rafin. l. c. 60 = C. verrucosa.

ANISANTHUS, Sweet, Hort. Brit. ed. I. 396 (1827) = **Anthotyxa**, Linn. (Irid.).
abyssinicus, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 344 = Anthotyxa abyssinica.
caryophyllaceus, Klatt, l. c. = Watsonia humilis.
Cunonia, Sweet, Hort. Brit. ed. I. 397 = Anth. Cunonia.
huillensis, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 344.—Afr. trop.
quadrangularis, Sweet, Hort. Brit. ed. I. 397 = Anth. quadrangularis.
saccatus, Klatt, in Linnæa, xxxv. (1867–68) 800 = Anth. saccata
splendens, Sweet, Brit. Flow. Gard. Ser. II. i. t. 84 = Anth. cæfra.

ANISANTHUS, Willd. ex Roem. & Schult. Syst. v. p. xiv (1819) = **Symphoricarpos**, Juss. (Caprifol.).
microphyllus, Willd. ex Roem. & Schult. l. c. 258 = S. microphyllus.

ANISEIA, Choisy (Conv. Or. 99) in Mém. Soc. Genèv. vi. (1833) 481; viii. (1838) 65 = **Ipomœa**, Linn. (Convolv.).
Ayulii, G. Don, Gen. Syst. iv. 295 = Hewittia bicolor.
orifolia, Meisn. in Mart. Fl. Bras. vii. 319.—Bras.
sagra, Kellogg, in Proc. Calif. Acad. v. (1873) 58.—Calif.
barlerioides, Choisy, Conv. Or. 109 = I. barleriodes.
bifora, Choisy, l. c. 101 = I. bifora.
brachiata, Hook. Fl. Bar. Jav. 516 = Hewittia bicolor.
calycina, Choisy, Conv. Or. 100 = I. calycina.
calycinoides, Choisy, in DC. Prod. ix. 481 = I. cissoides.
carnea, Griseb. Fl. Brit. W. Ind. 471, *sphalm.* = cærnea.
cærnea, Moric. N. Fl. Am. t. 85.—Bras.
cordata, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 367.—Chili.
dissectifolia, Walp. l. c.—Chili.
ANISEIA :—
emarginata, Hask. Cat. Hort. Bog. Alt. 139 = Ipomœa martinicensis.
enantia, Choisy, Conv. Var. 144 = I. martinicensis.
fulvicaulis, Hochst. ex Choisy, in DC. Prod. ix. 481.—Abyss.
gracilima, Choisy, l. c. 480.—Bras.
hastata, Meisn. in Mart. Fl. Bras. vii. 319 = I. phylloneura.
heterantha, Choisy, in DC. Prod. ix. 480.—Bras.
heterophylla, Meisn. in Mart. Fl. Bras. vii. 321 = I. heterophylla.
martinicensis, Choisy, Conv. Var. 144 = I. martinicensis.
Medium, Choisy, Conv. Or. 100 = I. denticulata.
nitens, Choisy, Conv. Var. 145.—Bras.
salicifolia, Choisy, l. c. 144 = I. martinicensis.
tomentosa, Meisn. in Mart. Fl. Bras. vii. 321 = I. martinicensis.
uniflora, Choisy, Conv. Or. 101 = I. martinicensis.
Velloziana, Choisy, in DC. Prod. ix. 480.—Bras.

ANISEION, St. Lag. in Ann. Soc. Bot. Lyon, vii. 119 (1880) = præc.

ANISEPTA, Rafin. Cat. 18 (1824) = **Croton**, Linn. (Euphorb.).
gravoleens, Rafin. l. c.—Am. bor.

ANISOCALYX, Hance, in Walp. Ann. iii. 195 (1853) = **Herpetris**, Gaertn. (Scrophul.).
limnanthiflorus, Hance, l. c. = H. Monniera.

ANISOCARPUS, Nutt. in. Trans. Am. Phil. Soc. Ser. II. vii. (1841) 388 = **Madia**, Molina (Compos.).
Bolanderi, A. Gray, in Proc. Am. Acad. vii. (1868) 360 = M. Bolanderi.
madioides, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 388 = M. Nuttallii.

ANISOCENTRA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 598 = **Tropæolum**, Linn. (Geran.).
cardiopetala, Turcz. l. c. = T. speciosum.

ANISOCENTRUM, Turcz. in Bull. Soc. Nat. Mosc. (1862) i. 322 = **Acisanthera**, P. Br. (Melast.)
Gardneri, Turcz. l. c. 323 = Acc. punctatissima.

ANISOCHÆTA, DC. Prod. v. 109 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 493.
mixantoides, *DC. l. c.* 110.—Afr. austr.

ANISOCHILUS, Wall. Pl. As. Rar. ii. 18 (1831). *LABIATAE*, Benth. & Hook. f. ii. 1177.
adenophorus, *Dals. & Gibs. Bomb. Fl.* 206.—Ind. or.
albidus, Wight, Ic. iv. t. 1435 = dysophylloides.
carnosus, *Wall. Pl. As. Rar.* ii. 18.—Ind. or.
crassus, Benth. in DC. Prod. xii. 81 = carnosus.
decussatus, Dals. & Gibs. Bomb. Fl. 206 = eriocephalus.
dysophylloides, *Benth. in Wall. Pl. As. Rar.* ii. 19.—Ind. or.
eriocephalus, *Benth. in DC. Prod.* xii. 81.—Ind. or.
eunervis, *Miq. Fl. Ind. Bat.* ii. 957.—Java.
glaber, Schred. Ind. Sem. Hort. Gott. (1838) 1 = carnosus.
pallidus, *Wall. Pl. As. Rar.* ii. 18.—Ind. or.
paniculatus, *Benth. in DC. Prod.* xii. 82.—Ind. or.
planagineus, *Hook. f. Fl. Brit. Ind.* iv. 628.—Ind. or.
polystachys, *Benth. in Wall. Pl. As. Rar.* ii. 19.—Ind. or.
purpureus, Wight, Ic. iv. t. 1435 = dysophylloides.
robustus, *Hook. f. Fl. Brit. Ind.* iv. 629.—Ind. or.
rupestris, Wight, ex Hook. f. Fl. Brit. Ind. iv. 627 = carnosus.
scaber, *Benth. in DC. Prod.* xii. 81.—Ind. or.
sericeus, *Benth. l. c.* 82.—Ind. or.
sinensis, *Hance, in Journ. Bot.* xxiii. (1885) 227.—China.
suffruticosus, *Wight, Ic.* iv. t. 1437.—Ind. or.
suffruticosus, Thw. Enum. Fl. Zeyl. 238 = paniculatus !
verticillatus, *Hook. f. Fl. Brit. Ind.* iv. 629.—Ind. or.
Wightii, *Hook. f. l. c.* 628.—Ind. or.

ANISOCOMA, Torr. & Gray, in Bost. Journ. Nat. Hist. v. (1844) 111. t. 18. *COMPOSITAE*, Benth. & Hook. f. ii. 518.
Pterotheranus, Kellogg, in Proc. Calif. Acad. iii. (1868) 20. *t. 4.*
acaulis, *Torr. & Gray, l. c.*—Calif.

ANISODENS, Dulac, Fl. Hautes-Pyr. 465 (1867) = **Scabiosa**, Tourn. (Dipsac.).
arvensis, Dulac, l. c. = S. arvensis.
longifolius, Dulac, l. c. = S. sylvatica.
sylvaticus, Dulac, l. c. = S. sylvatica.

ANISODERIS, Cass. in Dict. Sc. Nat. xlviii. 429 (1827) = **Crepis**, Linn. (Compos.).
fœtida, Fisch. & Mey. Ind. Sem. Petrop. iv. 32 = C. fœtida.
rhœadifolia, Fisch. & Mey. l. c. = C. fœtida.
rubra, Cass. in Dict. Sc. Nat. xlviii. 429 = C. rubra.

ANISODONTEA, Presl, Bot. Bemerk. 18 (1844) (in Abh. Boehm. Ges. Fol. V. iii.) = **Sphæralcea**, St. Hil. (Malvac.)
Dregeana, Presl, l. c. = S. Dregeana.

ANISODUS, Link & Otto, Ic. Sel. Pl. 77. t. 35 (1824) = **Scopola**, Jacq. (Solanac.).
humsis, Hook. f. ex C. B. Clarke, in Fl. Brit. Ind. iv. 243 = Mandragora caulescens.
luridus, Link & Otto, Ic. Sel. Pl. 77 = S. lurida.
stramonifolius, G. Don, in Lond. Hort. Brit. 61 = S. lurida.

ANISOLEPIS, Steetz, in Lehm. Pl. Preiss. i. 446 (1845) = **Helipterum**, DC. (Compos.).
Pyrethrum, Steetz, l. c. 447 = H. Pyrethrum.

ANISOLOBUS, A. DC. Prod. viii. 395 (1844) = **Odontadenia**, Benth. (Apocyn.).
amazonicus, Muell. Arg. in Mart. Fl. Bras. vi. i. 114.—Bras.
anomalus, Heurck & Muell. Arg. Obs. Bot. 160.—Peruv.
cognatus, Muell. Arg. in Mart. Fl. Bras. vi. i. 113.—Bras.
Cururu, Muell. Arg. l. c. 112.—Bras.
distinctus, Miers, Apocyn. S. Am. 169.—Am. austr.
Focken, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Sel.) 159.—Guian.
gracilipes, Muell. Arg. in Mart. Fl. Bras. vi. i. 115.—Bras.
hebecarpus, Muell. Arg. l. c. 111.—Bras.
Hostmanni, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Sel.) 159.—Surinam.
Kappleri, Miq. in Linnæa, xvIii. (1844) 787 = *Cururu.*
oblongus, Miers, Apocyn. S. Am. 169.—Am. austr.
Perrottetii, A. DC. in DC. Prod. viii. 395.—Bras.
Pohlianus, Miers, Apocyn. S. Am. 171.—Am. austr.
pridiifolius, Miers, l. c. 172.—Am. austr.
pulcherrimus, Miers, l. c. 171.—Am. austr.
puncticulosus, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Sel.) 158.—Guian.
rubidulus, Miers, Apocyn. S. Am. 173.—Am. austr.
Salzmanni, A. DC. in DC. Prod. viii. 395.—Bras.
Spruceri, Muell. Arg. in Mart. Fl. Bras. vi. i. 114.—Bras.
Stadelmeyeri, Muell. Arg. l. c. 113.—Bras.
Zuccarinianus, Miers, Apocyn. S. Am. 171.—Am. austr.

ANISOLOTUS, Bernh. Sem. Hort. Erf. (1837); Linnæa, xii. (1838) Lit. 75 = **Hosæusia**, Dougl. (Legum.).
anthyllioides, Bernh. ll. oc. = H. subpinnata.
Poeppigiana, Bernh. ll. oc.—Chili.
Wrangeliana, Bernh. ll. oc. = H. subpinnata.

ANISOMALLON, Baill. Adansonia, xi. (1874) 197. *OLACINEAE.*
cluniæfolium, *Baill. l. c.*—N. Caled.

ANISOMELES, R. Br. Prod. 503 (1810). *LABIATAE*, Benth. & Hook. f. ii. 1207.
albiflora, *Miq. Fl. Ind. Bat.* ii. 970.—Malaya.
australis, *Spreng. Syst. Ivr. Cur. Post.* 236.—Austral.
candicans, *Benth. in Wall. Pl. As. Rar.* i. 89.—Ind. or.
cœnesta, *Vacq. Ecl. Pl.* 127.—Ind. or.
distichæ, Heyne, ex Roth, Nov. Pl. Sp. 254 = ovata.
furcata, Lond. Hort. Brit. 329 = Craniotome versicolor.
glabrata, Benth in Wall. Cat. n. 6041 = ovata.
Heyneana, *Benth. in Wall. Pl. As. Rar.* i. 89.—Ind. or.

ANISOMELES :—
inodora, *R. Br. Prod.* 508.—Austral.
intermedia, Wight, in Benth. Lab. Gen. et Sp. 703 = malabarica.
malabarica, *R. Br. ex Sims, Bot. Mag.* t 2071.—Ind or.; Ins. Masc.
mollissima, Wall. Cat. n. 2089 = ovata.
moschata, *R. Br. Prod.* 503.—Austral.
nepalensis, Spreng. Syst. ii. 706 = Craniotome versicolor.
ovata, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 364.—Ind. or.; Malaya.
salviæfolia, *R. Br. Prod.* 503.—Austral.

ANISOMERIA, D. Don, in Edinb. N. Phil. Journ. xiii. (1832) 228. *PHYTOLACCACEÆ*, Benth. & Hook. f. iii. 85.
coriacea, *D. Don, l. c.*—Chili.
drastica, *Meq. in DC. Prod.* xiii. II. 25.—Chili.
fruticosa, *Phil. in Linnæa*, xxix. (1857–8) 88.—Chili.
littoralis, *Meq. in DC. Prod.* xiii II. 25.—Chili.

ANISOMERIS, Presl, Symb. Bot ii. 5. t. 54 (1833) = **Chomelia**, Jacq. (Rubiac.).
diversifolia, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 211.—Bras.
spinosa, Presl, Symb. Bot. ii. 5. t. 54 = C. Anisomeris.

ANISOMETROS, Hassk in Flora, xx (1847) 602 = **Pimpinella**, Linn. (Umbell.).
alpina, Hassk. l. c. = P. javana.

ANISONEMA, A. Juss. Euph. Tent. 19. t. 4 (1824) = **Phyllanthus**, Linn. (Euphorb.).
dubium, Blume, Bijdr. 589 = P. reticulatus.
eglandulosum, Decne. Herb. Tim. Descr. 154 = P. reticulatus.
glaucinum, Miq. Fl. Ind. Bat. Suppl. 449.—Sumatr.
hypoleucum, Miq. l. c.—Sumatr.
intermedium, Decne. in Nouv. Ann. Mus. Par. iii. (1831) 452 = P. reticulatus.
jamaicense, Griseb. Fl. Brit. W. Ind. 716 = P. jamaicensis.
multiflorum, Wight, Ic. t. 1839 = P. reticulatus.
reticulatum, A. Juss. Euph. Tent. t. 4. f. 11 = P. reticulatus.
Zollingeri, Miq. Fl. Ind. Bat. i. II. 575 = P. reticulatus.

ANISOPAPPUS, Hook. & Arn. Bot Beech. Voy. 196 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 339.
africanus, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 369.—Afr. trop.
chinensis, *Hook. & Arn. Bot. Beech. Voy.* 196.—China.

ANISOPETALA, Walp. Rep. iv. 228 (1845) = **Pelargonium**, L'Hérit. (Geran.).
crispa, Walp. l. c. = P. crispum.
purpurascentia, Walp. l. c. = P. purpurascens.
Radulea, Walp. l. c. = P. Radula, etc.
tomentosa, Walp. l. c. = P. tomentosa.

ANISOPETALUM, Hook. Exot. Fl. t. 149 (1825) = **Bulbophyllum**, Thou. (Orchid.).
Careyanum, Hook. l. c. = B. Careyanum.

ANISOPHYLLEA, R. Br. ex Sabine, in Trans. Hort. Soc. v. (1824) 446. *RHIZOPHOREÆ*, Benth. & Hook. f. i. 685.
ANISOPHYLLUM, G. Don, ex Hook. Niger Fl. 342, 575 (1849).
TETRACARPÆA, Benth. in Journ. Linn. Soc. iii. (1859) 72, sphalm.
TETRACRYPTA, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 314.
Beccariana, *Baill. Adansonia*, xi. (1875) 311.—Borneo.
disticha, Baill. l. c. = trapezoidalis.
Gardichaudiana, *Baill. l. c.*—Am. trop.
grandis, *Benth. in Journ. Linn. Soc.* i. (1857) 79, in nota.—Penang.
grandifolia, G. Henslow, in Hook. f. Fl. Brit. Ind. ii. 443 = grandis.
Griffithii, *Oliver, in Linn. Trans. Soc.* xxiii. (1862) 460.—Malacca.
laurina, *R. Br. ex Sabine, in Trans. Hort. Soc.* v. (1824) 446.—Afr. trop.

ANISOPHYLLEA :—
rhomboidea, *Baill. Adansonia*, xi. 1875) 310.—Borneo.
trapezoidalis, *Baill. l. c.* lii. (1862) 4.—Singapur.
zeylanica, *Benth. in Hook. Niger Fl.* 343, 575.—Zeylan.

ANISOPHYLLUM, Boiv. ex Baill. Adansonia, i. (1860–61) 159 = Croton, Linn. (Euphorb.).
acutifolium, Boiv. ex Baill. l. c. = C. muricatus.
scutiligerum, Boiv. ex Baill. l. c. 154 = C. Payeri anus.

ANISOPHYLLUM, G. Don, ex Benth. in Hook. Niger Fl. 342, 575 (1849) = **Anisophyllea**, R. Br. (Rhizoph.).

ANISOPHYLLUM, Haw. Syn. Pl. Succ. 159 (1812) = **Euphorbia**, Linn.
aegyptiacum, Schweinf. Beitr. Fl. Aethiop. 84 = E. aegyptiaca.
affine, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 29 = E. Balbiani.
amoenum, Klotzsch & Garcke, l. c. 33 = E adenoptera.
arabicum, Schweinf. Beitr. Aethiop. 84 = E. arabica.
arenarium, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 24 = E. zygophylloides.
Ateta, Klotzsch & Garcke, l. c. 87 = E. Atoto.
bahiense, Klotzsch & Garcke, l. c. 38 = E. bahiensis.
Berterianum, Klotzsch & Garcke, l. c. 82 = E. Bertoniana.
Besseri, Klotzsch & Garcke, l. c. 30 = E. Besseri.
Burmannianum, Klotzsch & Garcke, l. c. 25 = E. thymifolia.
caecorum, Klotzsch & Garcke, l. c. 38 = E. caecorum.
californicum, Klotzsch & Garcke, l. c. 36 = E. Deppeana.
centunculoides, Klotzsch & Garcke, l. c. 38 = E. centunculoides.
Chamaesyce, Haw. Syn. Pl. Succ. 160 = E. Chamaesyce.
Chamissonis, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 22 = E. Chamissonis.
chersinlepis, Klotzsch & Garcke, l. c. 21 = E. cheirolepis.
congenerum, Klotzsch & Garcke, l. c. 22 = E. congenera.
convolvuloides, Klotzsch & Garcke, l. c. 34 = E. convolvuloides.
cordatum, Klotzsch & Garcke, l. c. 68 = E. cordata.
crassipes, Klotzsch & Garcke, l. c. = E. macropus.
densiflorum, Klotzsch & Garcke, l. c. 28 = E. adenoptera.
dentatum, Haw. Syn. Pl. Succ. 162 = E. dentata.
dioicum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 31 = E. dioica.
emarginatum, Klotzsch & Garcke, l. c. 24 = E. emarginata.
Fendleri, Klotzsch & Garcke, l. c. 26 = E. Fendleri.
flexuosum, Klotzsch & Garcke, l. c. 35 = E. flexuosa.
Forskalii, Klotzsch & Garcke, l. c. 25 = E. Forskalii.
Geyeri, Klotzsch & Garcke, l. c. 28 = E. Geyeri.
glabratum, Klotzsch & Garcke, l. c. 35 = E. glabrata.
glaucophyllum, Klotzsch & Garcke, l. c. = E. glaucophylla.
goliathum, Klotzsch & Garcke, l. c. 29 = E. goliana.
granulatum, Schweinf. Beitr. Fl. Aethiop. 84 = E. granulata.
Humboldtii, Haw. Syn. Pl. Succ. 160 = E. graminea.
humifusum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 21 = E. humifusa.
humistratum, Klotzsch & Garcke, l. c. 26 = E. humistrata.
hypericifolium, Haw. Syn. Pl. Succ. 161 = E. hypericifolia.
hyssopifolium, Haw. l. c. = E. hyssopifolia.
hyssopifolium, Boiss. in DC. Prod. xv. II. 28 = E. hypericifolia.
inaequale, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 27 = E. adenoptera.
inaequilaterum, Klotzsch & Garcke, l. c. 23 = E. inaequilatera.
indicum, Schweinf. Beitr. Fl. Aethiop. 84 = E. hypericifolia.
Ipecacuanha, Haw. Syn. Pl. Succ. 164 = E. Ipecacuanha.

ANISOPHYLLUM :—
laevigatum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 35 = E. laevigata.
lasiocarpum, Klotzsch & Garcke, l. c. = E. lasiocarpa.
leiospermum, Klotzsch & Garcke, l. c. 23 = E. obliqua.
leucanthum, Klotzsch & Garcke, l. c. 27 = E. leucantha.
Lindenianum, Klotzsch & Garcke, l. c. 87 = E. Lindeniana.
linearе, Klotzsch & Garcke, l. c. 38 = E. linearis.
macropus, Klotzsch & Garcke, l. c. 33 = E. macropus.
maculatum, Haw. Syn. Pl. Succ. 162 = E. maculata.
melanadenium, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 32 = E. melanadenia.
Meyenianum, Klotzsch & Garcke, l. c. 25 = E. Meyeniana.
mozambicrnse, Klotzsch & Garcke, l. c. 30 = E. mozambicensis.
multiforme, Klotzsch & Garcke, l. c. 37 = E. multiformis.
Mundtii, Klotzsch & Garcke, l. c. 25 = E. parvifolia.
Nagleri, Klotzsch & Garcke, l. c. 31 = E. Nagleri.
nanum, Klotzsch & Garcke, l. c. 32 = E. chamaerhodos.
nodosum, Klotzsch & Garcke, l. c. 23 = E. clusiaefolia.
novo-mexicanum, Klotzsch & Garcke, l. c. 31.—N. Mexic.
ocymoideum, Haw. Syn. Pl. Succ. 161 = E. ocymoides.
orbiculatum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 24 = E. orbiculata.
ovalifolium, Klotzsch & Garcke, l. c. 26 = E. ovalifolia.
Peplis, Haw. Syn. Pl. Succ. 159 = E. Peplis.
pilulifernm, Haw. l. c. 163 = E. pilulifera.
polycnemoides, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 34 = E. polycnemoides.
polygonifolium, Haw. Syn. Pl. Succ. 160 = polygonifolia.
prostratum, Haw. l. c. 163 = E. prostrata.
ramosissimum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 30 = E. Garckeana.
rhytispermum, Klotzsch & Garcke, l. c. 34 = E. rhytisperma.
rosaeum, Haw. Syn Pl. Succ. 163 = E. rosea.
Ruizianum, Klotzsch & Garcke, l. c. 37 = E. Ruiziana.
scordifolium, Klotzsch & Garcke, l. c. 35 = E. scordifolia.
Selloi, Klotzsch & Garcke, l. c. 32 = E. Selloi.
serrile, Klotzsch & Garcke, l. c. 28 = E. stictospora.
serpens, Klotzsch & Garcke, l. c. 23 = E. serpens.
setigerum, Klotzsch & Garcke, l. c. 29 = E. setigera.
tenuiforme, Klotzsch & Garcke, l. c. 26 = E. velligera.
tettense, Klotzsch & Garcke, l. c. 34 = E. tettensis.
thymifolium, Haw. Syn. Pl. Succ. 160 = E. thymifolia.
Vahlii, Klotzsch & Garcke, l. c. 36 = E. Vahlii.
velleriflorum, Klotzsch & Garcke, l. c. 28 = E. velleriflora.
virgatum, Klotzsch & Garcke, l. c. 36 = E. Hookeri.

ANISOPHYLLUM, Jacq. Select. Am. 288 (1763) (Quid?).
pinnatum, Jacq. l. c.—Ins. Baru.

ANISOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 120 (Gen. Centr. Am. 46) = **Alloplectus**, Mart. (Gesneracˆ.).
pendulus, Oerst. l. c. = Allop. pendulus.

ANISOPLEURA, Fenzl, in Flora, xxvi. (1843) 459 = **Prangos**, Lindl. (Umbell.).
crenata, Fenzl, l. c.—Mesopotam.

ANISOPOGON, R. Br. Prod. 176 (1810). *GRAMINEÆ*, Benth. & Hook. f. iii. 1162.
avenaceus, *R. Br. l. c.*—Austral.
capensis, *Nees, Gram. Afr. Austr.* 265.—Afr. austr.

ANISOPTERA, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 65. t. 6 (1839). *DIPTEROCARPEÆ*, Benth. & Hook. f. i. 192.
ANTHEROTRICHE, Turcz. in Bull. Soc. Nat. Mosc. (1846) II. 503.
bantamensis, Hassk. Retzia, i. 140 = Vatica bantamensis.
costata, *Korth. Verh. Nat. Gesch. Bot.* 65. t. 6.—Borneo.
glabra, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 61.—Burma.

ANISOPTERA :—
Guiso, *A. DC. in DC. Prod.* xvi. ii. 616.—Ins. Philipp.
lanceolata, Walp. Ann. i. 118 = thurifera.
Mengachapoi, A. DC. in DC. Prod. xvi. II. 616 = Vatica Mangachapoi.
marginata, *Korth. Verh. Nat. Gesch. Bot.* 66. t. 6 — Borneo.
melanoxylon, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 160 = Vatica melanoxylon.
oblonga, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 301.— Burma.
odorata, Kurz, in Flora, lv. (1872) 190 = Vatica grandiflora.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* i. 485.— Sumatr.
plagata, *Blume, Mus. Bot. Lugd. Bat.* ii. 42.—Ins. Philipp.
polyandra, *Blume, l. c.* t. 6.—N. Guin.
thurifera, *Blume, l. c.*—Ins. Philipp.

ANISORA, Rafin. Sylva Tellur 74 (1838) = **Helioteres**, Linn. (Stercul.).
angulata et *Murri*, Rafin. l. c.—Hab.?

ANISORAMPHUS, DC. Prod vii. 251 (1838) = **Crepis**, Linn. (Compos.).
hypochaeridea, DC. l. c.—Afr. austr.
hypochaeroides, Hook. f. in Journ. Linn. Soc. vii. (1864) 204 = C. Hookeriana.

ANISOSCIADIUM, DC. Coll. Mém. v. 63. t. 15 (1829); Prod. iv. 284 (1830) = **Echinophora**, Linn. (Umbell.).
orientale, DC. Prod. l. c.—Persia.

ANISOSPERMA, Manso, Enum. Subst. Bras. 38 (1836). *CUCURBITACEAE*, Benth. & Hook. f. i. 820.
Passiflora, *Manso, l. c.*—Bras.

ANISOSTACHYA, Nees, in DC. Prod. xi. 368, 730 (1847) = **Justicia**, Linn. (Acanth.).
Bojeri, Nees, in DC. Prod xi. 368, 730.—Madag.
Commersoni, T. Anders. in Journ. Linn. Soc. vii. (1864) 46 = J. haplostachya.
volutina, Nees, in DC. Prod. xi. 730.—Madag.

ANISOSTEMON, Tuscz. in Bull. Soc. Nat Mosc. xx. (1847) i. 152 = **Connarus**, Linn.
trifoliatus, Turcz. l. c. = C. polyanthus.

ANISOSTICHUS, Bur. Monog. Bign. 43. t. 6 (1864) = **Bignonia**, Tourn.
capreolata, Bur. l. c. = B. capreolata.

ANISOSTICTUS, Benth. & Hook. f. Gen. ii 1038 (1888), lapsu = ANISOSTICHUS, Bur. = Bignonia, Tourn.

ANISOTES, Nees, in DC. Prod. xi. 424 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1111.
CALASIAS, Rafin. Fl. Tellur. iv. 64 (1836), nomen prius.
diversifolius, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 88.—Ins. Socotr.
trinalicus, *Nees, in DC. Prod.* xi. 424.—Arab.

ANISOTES, Lindl. Introd. Nat. Syst. ed. II. 100, 441 (1836) = **Lythrum**, Linn.
anomalus, Lindl. l. c. 441 = L. anomalum.
Hilarianus, Melam. Gen. Comm. 84 = L. anomalum.

ANISOTICTE, Bartl. Ord. 294 (1830) = **Capparis**, Linn. (C. frondosa, Jacq.).

ANISOTOMA, Fensl, in Linnaea, xvii. (1843) 880. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 781.
ANISOTOMARIA, Presl, Bot. Bemerk. 103 (1844).
DECACERAS, Harv. Thes. Cap. ii. 9. t. 114 (1868).
LOPHOSTEPHUS, Harv. l. c. 113 (1868).
Arnottii, *Benth. & Hook. f. Gen.* ii. 781.—Afr. austr.
cordifolia, *Fensl, in Linnaea*, xvii. (1843) 881.—Afr. austr.

ANISOTOMARIA, Presl, Bot. Bemerk. 103 (1844) = **Anisotoma**, Fensl (Asclep.).
mellifa, Presl, l. c. = Anisotoma cordifolia.

ANISOTOME, Hook. f. Fl. Antarct. 16. t. 8, 9 (1844) = **Asciphyllia**, Forst. (Umbell.).
antipoda, Hook. f. Fl. Antarct. 17.—Ins. Antarct.
aromatica, Hook. f. Fl. N. Zeal. i. 89.—N. Zel.
geniculata, Hook. f. l. c. 90 = Angelica geniculata.
Gingidium, Hook. f. l. c. 89 = Angelica Gingidium.
intermedia, Hook. f. l. c. = Ligusticum intermedium.
latifolia, Hook. f. Bot. Antarct. 16 = Ligusticum latifolium.
Lyallii, Hook. f. Fl. N. Zeal. i. 88 = Ac. Lyallii.
rosaefolia, Hook. f. l. c. 90 = Angelica rosaefolia.

ANISTYLIS, Rafin. Neogenyt. 4 (1825) = **Liparis**, Rich. (Orchid.).
Convallaria, Rafin. l. c. = L. lillifolia
lutea, Rafin. l. c. = L. Loeselii.

ANISUM, Riv. ex Linn. Gen. ed. I. 83 (1737) = **Pimpinella**, Linn. (Umbell.).
caffrum, Eckl. & Zeyh. Enum. 241 = P. caffra.
officinale, DC. Prod. iv 122 = P. Anisum.
officinarum, Moench, Meth. 100 = P. Anisum.
staderae, Eckl. & Zeyh. Enum. 241 = P. staderaia.
vulgare, Gaertn. Fruct. i. 72 = P. Anisum.

ANKYLOBUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 148 = **Astragalus**, Tourn. (Legum.).
caspicus, Stev. l. c. = A. caspius.
contortuplicatus, Stev. l. c. = A. contortuplicatus.
cruciatus, Stev. l. c. = A. cruciatus.
hamosus, Stev l. c. = A. hamosus.
reticulatus, Stev. l. c. = A reticulatus.

ANKYROPETALUM, Fenzl, in Bot. Zeit. i. (1843) 393 = **Gypsophila**, Linn (Caryophyll.).
arsusianum, Kotschy, in Boiss. Fl. Orient i. 538.—Syria.
coelesyriacum, Boiss. Diagn. Ser. I. viii. 59.—Oriens.
gypsophiloides, Fenzl, in Bot. Zeit. i. (1843) 393.—Mesopot.
Reuteri, Boiss. & Haussk. in Boiss. Fl. Orient. i. 538.—Orient.

ANNESLEA, W. Hook. in Salisb. Parad. Lond. t. 64 (1807) = ANNESLIA, Salisb.

ANNESLEA, Roxb. ex Andr. Bot. Rep. t. 618 (1810); Fl. Ind. ii. 573 (1832) = **Euryale**, Salisb. (Nymph.)
amazonica, Presl, Bot. Bemerk 103 = Victoria regia.
brasiliana, Presl, l. c. = Victoria regia.
spinosa, Andr. Bot. Rep. t. 618 = E. ferox.

ANNESLEA, Wall. Pl. As. Rar. i. 5. t. 5 (1829). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 182.
CALLOSMIA, Presl, Bot. Bemerk. 103 (1844).
crassipes, *Hook. ex Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 121.—Malaya.
fragrans, *Wall. Pl. As. Rar.* i. 5. t. 5.—Burma.
monticola, *Kurz, in Journ. As. Soc Beng.* xlii. (1873) II. 59.—Burma.

ANNESLIA, = ANNESLEA, W. Hook. = **Callianthera**, Benth. (Legum.).

ANNESLIA, Salisb. Parad. Lond. t. 64 (1807) = **Calliandra**, Benth. (1840) (Legum.).
falcifolia, Salisb. l. c. in textu = C. Houstoni.
grandiflora, Salisb. l. c. t. 64 = C. Houstoni.
Houstoni, Sweet, Hort. Brit. ed. II. 168 = C. Houstoni.

ANNESORHIZA, Cham. & Schlecht. in Linnaea, i. (1826) 898. *UMBELLIFERAE*, Benth. & Hook. f. i. 918.
ACROGLYPHE, E. Mey. Zwei Pfl. Docum. 100 (1845).
GLIA, Sond. in Harv. & Sond. Fl. Cap. ii. 547 (1862).
STENOSEMIS, E. Mey. Zwei Pfl. Docum. 141, 148 (1845).
capensis *Cham. & Schlecht. l. c.* 899.—Afr. austr.
elata, *Eckl. & Zeyh. Enum.* 245.—Afr. austr.
filicalis, *Eckl. & Zeyh. l. c.*—Afr. austr.
hirsuta, *Eckl. & Zeyh. l. c.*—Afr. austr.
macrocarpa, *Eckl. & Zeyh. l. c.*—Afr. austr.
montana, *Eckl. & Zeyh. l. c.* 244.—Afr. austr.
spuria, *Eckl. & Zeyh. l. c.* 243.—Afr. austr.
villosa, *Sond. in Harv. & Sond. Fl. Cap.* ii. 546.—Afr. austr.

ANNONA, Linn. Hort. Cliff. 222 (1737); Forsk. Fl. Aegypt. Arab. 102 = **Anona**, Linn.

ANNULARIA, Hochst. in Flora, xxiv. (1841) 670 = **Vocanga**, Thou. (Apocyn.).
natalensis, Hochst. l. c. 671 = V. Dregei.

ANOCHEILE, Hoffmgg. ex Reichb. Nom. 235 (1841) = **Epidendrum**, Linn (Orchid.).

ANODA, Cav. Diss. i. 38. t. 10. f. 3 (1785). *MALVACEAE*, Benth. & Hook. f. i. 202.
CAVANILLEA, Medic. Malv. 19 (1787).
acerifolia, Cav. in Anal. Clenc. Nat. vi. (1803) 336; DC. Prod. i. 459 = hastata.
angustifolia, *Spreng. Nachtr. i. Bot. Gart. Halle*, 11.— Hab.?
arenariaeflora, Hort. ex Steud. Nom. ed. II. i. 100 = crenatiflora.
brachyantha, Reichb. Hort. Bot. t. 34 = triangularis.
crenatiflora, Ortog. Hort. Matr. Dec. 96 = parviflora?
cristata, Schlecht in Linnaea, xi. (1837) 210 = lavateroides.
Dilleniana, Cav. Diss. i. 40. t. 11. f. 1 = cristata.
fernandeziana, *Steud in Flora*, xxxix. (1856) 487.— Ins. Juan Fern.
hastata, *Cav. Diss.* i. 39 t. 10. f. 8.—Mexic.; Ind. occ
hirsuta, *Phil. in Linnaea*, xxviii. (1856) 675.—Chili.
incarnata, *H. B. & K. Nov. Gen. et Sp.* v. 266.— Mexic.
incarnata, Spreng. ex Steud. Nom. ed. II. i. 100 = triangularis.
lanceolata, *Hook. & Arn. Bot. Beech. Voy.* 411.— Mexic.
lavateroides, *Medic. Malv.* 19.—Mexic.
Ochsenii, Phil. in Linnaea, xxviii. (1856) 618 = Abutilon Ochsenii.
Ortegae, Spreng. Nachtr. i. Bot. Gart. Halle, 11 = crenatiflora.
ovata, Meyen, Reise, ii. 139 = Sida fallax.
parviflora, *Cav. Ic.* v. 19. t. 431.—Mexic.
pentaschista, *A. Gray, Pl. Wright.* ii. 22.—Nov. Mexic.
populifolia, *Phil. in Linnaea*, xxviii. (1856) 813.— Chili.
pubescens, *Schlecht. in Linnaea*, xi. (1837) 218.— Mexic.
punicea, *Lag. Gen. et Sp. Nov.* t. 21.—Mexic.
reticulata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 389.—Am. bor.
strictiflora, *Steud. in Flora*, xxxix. (1856) 487.— Chili.
triangularis, *DC. Prod.* i. 459.—Mexic.
triloba, Cav. Diss. i. 39 t. 10. f. 3 = lavateriolides.
Wrightii, *A. Gray, Pl. Wright.* ii. 22.—Nov. Mexic.

ANODENDRON, A. DC. Prod. viii. 448 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 719.
Candollesanum, *Wight, Ic.* iv. t. 1309.—Malaya.
coriaceum, *Miq. Fl. Ind. Bat.* ii. 485.—Java.
infiatum, *Hassk. in Flora*, xxvii. (1845) 269.—Java.
laeve, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* i. 215.—Japon.
moluccanum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140.—Malaya.
paniculatum, *A. DC. in DC. Prod.* viii. 444.—Ind. or.; Malaya.
pauciflorum, *Hook. f. Fl. Brit. Ind.* iii. 669.— Penang.
rhinocarpum, *Thw. Enum. Pl. Zeyl.* 194.—Zeylan.
rubescens, Teijsm. & Binn. Cat. Hort. Bogor. 127 = Candollesanum.
tenuiflorum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140. —Sumatr.

ANODIA, Hassk. in Flora, xxv. (1842) II. Beibl. 88, in nota = **Anoda**, Cav. (Malvac.).

ANODISCUS, Benth. in Benth. & Hook. f. Gen. ii. 998 (1876). *GESNERACEAE*, Benth. & Hook. f. ii. 998.
peruvianus, *Benth. in Hook. Ic. Pl.* t. 1199.—Peruv.

ANODONTEA, Sweet, Hort. Brit. ed. I. 467 (1827) = **Alyssum**, Tourn. (Crucif.).
arenaria, G. Don, Gen. Syst. i. 180 = Al. montanum?

ANODONTEA :—
Auckeri, D. Dietr. Syn. Pl. iii. 643 = Alyssum foliosum.
canescens, Sweet, Hort. Brit. ed II. 25 = Alyssum canescens.
dasycarpa, Sweet, l. c. = Alyssum dasycarpum.
edentula, Sweet, l. c. ed I. 467 = Alyssum gemonense.
halimifolia, Sweet, l. c. = Alyssum halimifolium.
macrocarpa, Sweet, l. c. = Alyssum macrocarpum.
marginata, D. Dietr. Syn. Pl. iii. 643 = Alyssum marginatum.
meniocoides, D. Dietr. l. c. = Alyssum meniocoides.
peltaroides, D. Dietr. l. c. = Alyssum peltaroides.
pyrenaica, G. Don, Gen. Syst. i. 180 = Alyssum pyrenaicum.
rupestris, Sweet, Hort Brit. ed. I. 467 = Alyssum rupestris.
spinosa, Sweet, l. c. = Alyssum spinosum.
tenuifolia, G. Don, Gen. Syst. i. 180 = Alyssum tenuifolium.
xanthocarpa, D. Dietr. Syn. Pl. iii. 643 = Alyssum macrostylum.

ANODOPETALUM, A. Cunn. ex Endl. Gen. 818 (1839). *SAXIFRAGEAE*. Benth. & Hook. f. i. 652. biglandulosum, *A. Cunn. ex Hook. f. Fl. Tasm.* i. 148. —Australia.
glandulosum, Baill. Hist. Pl. iii. 378 = biglandulosum.

ANOECTOCHILUS, Blume, Bijdr. 411. t. 15 (1825) (Anecochilus). *ORCHIDEAE*, Benth. & Hook. f. iii. 598.
alto-lineatus, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 141 ¬ brevilabris.
albo-marginatus, *Lour. Encyc. Pl. Suppl.* iii. 1539.—Cochinch.
argyroneurus, Hort. ex Loud. Hort. Brit. Suppl. iii. 608 = Physurus argenteus.
argyroneurus, *C. Koch & Lauche, in Berl. Allg. Gartenz.* xxv. (1857) 4.—Java.
aureus, Hort. ex C. Koch & Lauche, l. c. = setaceus.
brevilabris, *Lindl. Gen. et Sp. Orch.* 499.—Ind. or.
Bullenï, *Low, ex Morr. in Belg. Hortic.* xii. (1862) 10. —Borneo.
crispus, *Lindl. in Journ. Linn. Soc.* i. (1857) 180.—Ind. or.
Crotonæ *Reichb. f. in Hamb. Gartenz.* xvi. (1866) 494.—Hab.?
Dawsonianus, Low, ex Reichb. f. in Gard. Chron. (1868) 1088 = Haemaria discolor.
Dominï v.—*Cf. Burbridge, in Gard. Chron.* (1871) 44.
elatus, *Lindl. in Journ. Linn. Soc.* i. (1857) 178.—Ind. or.
elongatus, *Mig. Fl. Ind. Bat.* iii. 784.—Java.
flavescens, Blume, Bijdr. 419 = Odontochilus flavescens.
flavus, Benth. & Hook f. Gen. iii. 598 = Odontochilus flavus.
Frederici-Augusti, *Reichb. f. in Hamb. Gartenz.* xvi. (1866) 177, 494.—Hab.?
glaber, *Mig. Fl. Ind. Bat.* iii. 782.—Malaya.
gracilis, *Blume, Bijdr.* 418.—Malaya.
grandiflorus, Lindl. in Journ. Linn. Soc. i. (1857) 179 = Odontochilus grandiflorus.
Hasseltii, *Mig. Fl. Ind. Bat.* iii. 784.—Java.
intermedius, *Hort. ex Lond. Encyc. Pl. Suppl.* ii. 1492. —Ind. or.
intermedius, *Reichb. f. in Hamb. Gartenz.* xvi. (1866) 177.—Hab.?
Jaubertii, Gaudich. in Voy. Bonite, t. 100 = Odontochilus Jaubertii.
lanceolatus, Lindl. Gen. et Sp. Orch. 499 = Odontochilus lanceolatus.
lanceolatus, Hort. ex Blume, Coll. des Orch. i. 50 = Roxburghii.
Lobbianus, Planch. in Fl. des Serres, Sér. I. v. (1849) 519 = Roxburghii.
longiflorus, Reichb. f. in Seem. Fl. Vit. 294 = Odontochilus longiflorus.
Lowii, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1492 = Dossinia marmorata.
luteus, Lindl. in Journ. Linn. Soc. i. (1857) 179 = Odontochilus lanceolatus.
neglectus, *Blume, Orch. Arch.* 47 *in nota*.—Zeylan.?
occultus, *Mig. Fl. Ned. Ind.* iii. 784.—Java.
Ovdianus, Hort.—Cf. Gard. Chron. (1869) 1226 = Haemaria discolor.
Ortgiesi, *Hort. ex Van-Geert, Cat.* n. 70 (1873).—N. Ornat.
Nevilleanus, *Low, ex C. Morr. in Belg. Hortic.* xii. (1862) 10.—Hab.?

ANOECTOCHILUS :—
pictus, Hort. ex Loud. Hort. Brit. Suppl. iii. 608 = Physurus pictus.
pubescens, *Blume, Bijdr.* 412.—Java.
regalis, *Blume, Orch. Arch. Ind.* 46.—Zeylan.
regalis, Low, ex C. Morr. in Belg. Hortic. xii. (1862) 4 = setaceus.
Reinwardtii, *Blume, Orch. Arch. Ind.* 48.—Malaya.
Roxburghii, *Lindl. in Wall. Cat.* n. 7387.—Ind. or.
Ruckeri, *Low, ex C. Morr. in Belg. Hortic.* xii. (1862) 10.—Borneo.
Rumphii, *Low, ex C. Morr. l. c.* 5.—Malaya.
sandwicensis, *Lindl. Gen. et Sp. Orch.* 500.—Ind. Sandv.
setaceus, *Blume, Bijdr.* 413.—Java.
setaceus, Lindl. Bot. Reg. t. 2010 = regalis.
setaceo-pictus, *C. Koch & Lauche, in Berl. Allg. Gartenz.* xxv. (1857) 4.—Hab.?
spicatus, *Mig. Fl. Ind. Bat.* iii. 784.—Borneo.
stratus, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1492 = Zeuxine regia.
tuniformis, *Mig. Fl. Ind. Bat.* iii. 783.—Sumatra.
Veitchianus, Hort. ex Blume, Coll. des Orch. i. 119 = Macodes petola.
xanthophyllus, Planch. Fl. des Serres, Sér. I. v. (1849) 519 = Frederici-Augusti.

ANOGEISSUS, Wall. Cat. n. 6014 (1829) *COMBRETACEAE*, Benth. & Hook. f. i. 687.
ANDERSONIA, Buch.-Ham. ex Wall Cat. l. c. (1832).
acuminata, *Wall. Cat.* n. 6014.—Ind. or.
hirta, Wall. Cat. n. 6016 = acuminata.
latifolia, *Wall. Cat.* n. 6015.—Ind. or.
leiocarpa, *Guill. & Perr. Tent. Fl. Senag.* i. 280. t. 65. —Trop. or.
myrtifolia, Wall. Cat. n. 4017 = pendula.
pendula, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 171.—Ind. or.
philippinensis, *Heurck & Muell. Arg. Obs. Bot.* 209. —Burma.

ANOGRA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 339 = Osmothera, Linn. (Osagr.).
Douglasiana, Spach, l. c. 340 = O. albicaulis.
Nuttalliana, Spach, l. c. = O. albicaulis.
pinnatifida, Spach, l. c. 341 = O. albicaulis.

ANOGYNA, Nees, in Hook. Journ. Bot. ii. (1840) 395 = Lagenocarpus, Nees (Cyperac.).
tremula, Nees, l. c. = L. tremula.

ANOIGANTHUS, Baker, in Journ. Bot. xvi. (1878) 76. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 733.
breviflorus, *Baker, l. c.*—Natal.
luteus, Baker, l. c. 77 = breviflorus.

ANOMA, Lour. Fl. Cochinch. 278 (1790) = Moringa, Juss.
cochinchinensis, Lour. l. c. 280 (Cascalpineae sp. ?).
Moringa, Lour. l. c. 279 = M. pterygosperma.
Morunga, Lour. l. c. = M. pterygosperma.

ANOMALANTHUS, Klotzsch, in Linnaea, xli. (1838) 238 = Stimcochilus, Klotzsch (Eriac.).
discolor, Klotzsch, l. c. 239.—Afr. austr.
scoparius, Klotzsch, l. c.—Afr. austr.

ANOMALOPTERIS, G. Don, Gen. Syst. i. 647 (1831) = Acridocarpus, Guill. & Perr. (Malpig.).
longifolia, G. Don, l. c. = Ac. Smeathmanni.
obovata, G. Don, l. c. = Ac. plagiopterus.
spicata, G. Don, l. c. = Ac. Smeathmanni.

ANOMALOSTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 162. t. 38 (1861) = Cissonia, Linn. (Capparid.).
brevirostris, Klotzsch, l. c.—Afr. trop.

ANOMALOTIS, Steud. Syn. Pl. Glum. 198 (1855) = Trisetaria, Forsk. (Gramin.).
quinquesetis, Steud. l. c. = T. quinquesetis.

ANOMANTHODIA, Hook. f. in Benth. & Hook. Gen. ii. 87 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 87.
auriculata, *Hook. f. l. c.*—Malaya.

ANOMANTIA, Rafin. ex DC. Prod. v. 617 (1836) = Verbesina, Linn. (Compos.). (V. virginica, Linn.)

ANOMATHECA, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1806) 237 = Lapeirousia, Pourr. (Irid.).
angolensis, Baker, in Journ. Bot. xiv. (1876) 337.—Afr. austr.
calamifolia, Klatt, in Abh. Naturf. Ges. Halle, xii. (1869) 358.—Afr. austr.
cruenta, Lindl. Bot. Reg. t. 1869.—Afr. austr.
grandiflora, Baker, in Journ. Bot. xiv. (1876) 337.—Afr. austr.
juncea, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1806) 237 = L. juncea.
xanthospila, Ker-Gawl. ex Spreng. Syst. i. 154.—Afr. austr.

ANOMAZA, Lawa. ex Salisb. in Trans. Hort. Soc. i. (1812) 323 = Lapeirousia, Pourr. (Irid.).
excisa, Lawa. Cat. 2; ex Salisb. l. c. = L. juncea.

ANOMIANTHUS, Zoll. in Linnaea, xxix. (1857-58) 324. *ANONACEAE*, Benth. & Hook. f. i. 27. heterocarpus, *Zoll. l. c.*—ins. Bantam.

ANOMOCARPUS, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 351; Contrib. ii. 28 = Callosera, Cav.
axillaris, Miers, l. c. 353; 28.—Chili.
eryngioides, Miers, l. c. 353; 29 = C. eryngioides.
leucacthenus, Miers, l. c. 355; 31.—Chili.
pulvinatus, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 354 = C. pulvinata.
subsessiliflorus, Miers, l. c. 352; Contrib. ii. 29 = C. sessiliflora.
tenuifolius, Miers, l. c. 356; 33 = A. tenuifolia.
tenuis, Miers, l. c. 356; 32.—Chili.

ANOMOCHLOA, Brongn. in Ann. Sc. Nat. Sér. III. xvi. (1851) 368. t. 23. *GRAMINEAE*, Benth. & Hook. f. iii. 1111.
marantoidea, Pritz. Ind. Ic. 74 = marantoidea.
marantoides, *Brongn. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 369.—Brasa.

ANOMORRHEGMIA, Meisn. Gen. i. 303 (1840) = Stauranthera, Benth. (Gesner.).
Miquelii, Meisn. l. c. ii. 212 = S. scalcarata.

ANOMOSANTHES, Blume, Rumphia, iii. 151, in obs. (1847) = Hemigyrosa, Blume (Saplnd.).
deficiens, Blume, l. c. = H. deficiens.

ANOMOSPERMUM, Dalz. in Hook. Kew Journ. iii. (1851) 238 = Actephila, Blume (Euphorb.).
excelsum, Dalz. l. c. = Actephila excelsa.

ANOMOSPERMUM, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 39. *MENISPERMACEAE*, Benth. & Hook. f. i. 35.
ELIBARRENA, Miers, l. c. Ser. III. xiii. (1864) 124.
actiniflorum, *Griseb. Cat. Pl. Cub.* 4.—Cuba.
grandifolium, *Eichl. in Flora*, xlvii. (1864) 388.—Bras.
Hostmanni, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 103 = Schomburgkii.
japurense, *Eichl. in Flora*, xlvii. (1864) 388.—Bras.
lucidum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 103.—Guian.
mitidum, Miers, l. c. = Schomburgkii.
oblongatum, *Miers, l. c.*—Bras.
ovatum, *Miers, l. c.*—Bras.
reticulatum, *Eichl. in Flora*, xlvii. (1864) 388.—Bras.
Schomburgkii, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. vii. (1851) 109; Ser. III. xiv. (1864) 103.—Guian.

ANOMOSTEPHIUM, DC. Prod. v. 560 (1836) = Aspilia, Thou. (Compos.).
angustifolium, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 290 = Aspilia foliosa.
buphthalmoides, DC. in Deless. Ic. Sel. iv. t. 93.—Candolleap.
elliptieum, DC. Prod. v. 560 = Aspilia elliptica.
foliosum, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 290 = Aspilia foliosa.
oblongifolium, DC. Prod. v. 560 = Aspilia foliosa.
ovatifolium, DC. l. c. = Aspilia ovatifolia.

ANONA, Linn. Syst. ed. I. (1735); Gen. ed. I. 188 (1737). *ANONACEAE*, Benth. & Hook. f. I. 27.
ANNONA, Linn. Hort. Cliff. 329 (1737).
acutiflora, *Mart. Fl. Bras.* xiii. t. 10.—Bras.
aethiopica, Steud. Nom. ed. II. ii. 737 = Xylopia aethiopica.
africana, *Linn. Sp. Pl.* 537.—Hab.?
amara, *Raeusch. Nom.* ed. III. 161.—Guiana.
Ambotay, *Aubl. Pl. Gui.* t. 616. t. 249.—Am. trop.
amplexicaulis, *Lam. Encyc.* ii. 127.—Madag.; Ins. Mascar.
antiocquensis, *Linden, Cat. n.* 87 (1871).—N. Granat.
arenaria, Schum. & Thonn. Beskr. Guin. Pl. 257 = senegalensis.
aromatica, Boj. Hort. Maurit. 6 = Xylopia Richardi.
asiatica, Heyne, ex Wall. Cat. sub n. 6475 = Unona panosa.
asiatica, *Linn. Sp. Pl.* 537.
asiatica, Vahl, Symb. Bot. iii. 78 = Forskahlii.
atabapensis, *H. B. & K. Nov. Gen. et Sp.* v. 58.—Guian.
australis, *St. Hil. Fl. Bras. Mer.* i. 33.—Bras.
axilliflora, *DC. Prod.* i. 86.—Guian.
Barteri, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 477.—Afr. trop.
bibracteata, Hook. Ic. Pl. iv. t. 328 = Guatteria bibracteata.
Bonplandiana, H. B. & K. Nov. Gen. et Sp. v. 58 = muricata.
bullata, *A. Rich. Ess. Fl. Cub.* 31.—Cuba.
cacuna, *Warm. in Kjoeb. Vidensk. Meddel.* (1873) 155.—Bras.
cascarilloides, *Griseb. Cat. Pl. Cub.* 2.—Cuba.
cauliflora, *Mart. Fl. Bras.* xiii. t. 16.—Bras.
Cherimolia, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. trop.
cherimolioides, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 27.—N. Granat.
cheriqueusis, *Linden, Cat. n.* 87 (1871).—N. Granat.
chrysocarpa, Lepr. in Guill. & Perr. Tent. Fl. Seneg. i. 6 = palustris.
chrysopetala, *Steud. in Flora*, xxvi. (1843) 754.—Guian.
chrysophylla, *Boj. in Ann. Sc. Nat.* xx. (1880) 52.—Ins. Mascar.
cinerea, *Dunal, Monog. Anon.* 71. t. 8.—Ind. occ.
colorata, *Vell. Fl. Flum.* 239; v. t. 139.—Bras.
conica, *Ruiz & Pav. Fl. Per.* v. t. 490; *ex G. Don, Gen. Syst.* i. 88.—Peruv.
conophylla, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 27.—N. Granat.
coriacea, *Mart. Fl. Bras.* xiii. t. 6.—Bras.
cornifolia, *St. Hil. Fl. Bras. Mer.* i. 33.—Bras.
crassifolia, *Mart. Fl. Bras.* xiii. t. 6.—Bras.
crotonifolia, *Mart. l. c.* 16.—Bras.
densicoma, *Mart. l. c.* 9.—Bras.
depressa, *Baill. in Adansonia*, viii. (1868) 267.—Mexic.
diodon, *St. Hil. Fl. Bras. Mer.* i. 34.—Bras.
distincta, Raeusch. Nom. ed. III. 160 = Forskahlii.
distincta, Willem. in Ust. Ann. Bot. xviii. 42 = Guatteria Willemetiana.
dodecapetala, Lam. Encyc. ii. 127 = Talauma Plumieri.
dolabripetala, Raddi, in Mem. Mod. xviii. Fis. (1820) 394 = Rollinia longifolia.
echinata, *Dunal, Monog. Anon.* p. 68. t. 4.—Guian.
exalbida, Vell. Fl. Flum. 240; v. t. 131 = Rollinia exalbida.
excelsa, *H. B. & K. Nov. Gen. et Sp.* v. 59.—Mexic.
exsucca, Dunal, Monog. Anon. 77 = Rollinia exsucca.
fagifolia, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 135.—Bras.
fostida, *Mart. Fl. Bras.* xiii. t. 16.—Bras.
Forskahlii, *DC. Syst.* i. 472.—Aegypt.
furfuracea, *St. Hil. Fl. Bras. Mer.* i. 34.—Bras.
glabra, Forsk. Fl. Aegypt. Arab. 103 = Forskahlii.
glabra, *Linn. Sp. Pl.* 537.—Hab.?
glauca, *Schum. & Thonn. in Dan. Vid. Selsh. Afh.* iv. (1894) 81.—Guinea.
globiflora, *Schlecht. in Linnaea*, x. (1836) 235.—Mexic.
grandiflora, *Lam. Encyc.* ii. 126.—Ins. Mascar.; Madag.
grandiflora, Bart. Trav. 20 & 171. t. 2 = Asimina grandiflora.
grandiflora, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 131.—Bras.
haematantha, *Mig. in Linnaea*, xxii. (1849) 465.—Guian.
hemipetala, Linn. f. Suppl. 270 = Artabotrys odoratissimus.

ANONA:—
Hostmanni, *Steud. in Flora*, xxvi. (1843) ii. 754.—Guian.
Humboldtiana, H. B. & K. Nov. Gen. et Sp. v. 56 = seq.
Humboldtii, *Dunal, Monog. Anon.* 64. t. 3.—Cumana.
humilis, *Benth. in Hook. Kew Journ.* iii. (1851) 118.—Bras.
hypoglauca, *Mart. Fl. Bras.* xiii. t. 18.—Bras.
involucrata, *Baill. Adansonia*, viii. (1868) 265.—Mexic.
lahophylla, *Triana, & Planch. in Ann. Sc. Nat. Sér.* IV. xvii. (1862) 27.—N. Granat.
laevigata, *Mart. Fl. Bras.* xiii. t. 17.—Bras.
laevis, *H. B. & K. Nov. Gen. et Sp.* v. 60.—Guian.; Venezuela.
lanceolata, Willd. ex Steud. Nom. ed. II. i. 100 = Guatteria mayprunensis.
lasiocalyx, *Mart. Fl. Bras.* xiii. t. 16.—Bras.
laurifolia, *Dunal, Monog. Anon.* 65.—Ind. occ.
lepidota, *Msg. in Linnaea*, xix. (1847) 129.—Guian.
Liebmanniana, *Baill. Adansonia*, viii. (1868) 266.—Mexic.
longifolia, Aubl. Pl. Gui. i. 615. t. 248 = Duguetia longifolia.
madagascariensis, Pers. Syn. ii. 95 = grandiflora.
Manirote, *H. B. & K. Nov. Gen. et Sp.* v. 59.—Guian.
Mannii, *Oliver, in Hook. Ic. Pl.* t. 1010.—Afr. trop.
Marcgravii, *Mart. Fl. Bras.* xiii. t. 5.—Bras.
mexicana, *Ladd. ex G. Don, in Loud. Hort. Brit.* 227.—Mexic.
micrantha, *Bert. ex Spreng. Syst.* ii. 640.—Ins. S. Doming.
microcarpa, Jacq. Fragm. 40. t. 44 = Diospyros Carpilla.
microcarpa, *Ruiz & Pav. Fl. Peruv.* v. t. 487; *ex G. Don, Gen. Syst.* i. 90.—Peruv.
montana, *Macfad. Fl. Jamaic.* i. 7.—Jamaic.
monticola, *Mart. Fl. Bras.* xiii. t. 7.—Bras.
mucosa, Jacq. Obs. i. 16 = Rollinia mucosa.
muricata, *Linn. Sp. Pl.* 536.—Am. trop.
muscosa, Aubl. Pl. Gui. i. 613 = Rollinia mucosa.
Myristica, Gaertn. Fruct. iii. 194. t. 195 = Monodora Myristica.
nitida, *Mart. Fl. Bras.* xiii. t. 9.—Bras.
nitida, Willd. ex Steud. Nom. ed. II. i. 101 = Guatteria mayprunensis.
obovata, Willd. Sp. Pl. ii. 1269 = Asimina grandiflora.
obtusiflora, Tuss. Fl. Antill. t. 38 = Rollinia mucosa.
obtusifolia, DC. Prod. i. 84 = Rollinia mucosa.
paludosa, *Aubl. Pl. Gui.* i. 611. t. 246.—Guian.
palustris, *Linn. Sp. Pl.* ed. II. 757.—Am. et Afr. trop.
Pavoni, *G. Don, Gen. Syst.* i. 89.—Peruv.
peduncularis, *Steud. in Flora*, xxvi. (1843) ii. 754.—Guian.
pendula, Salish. Prod. 380 = Asimina triloba.
Perrottetii, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 197.—Senegal.
pervaviana, Humb. & Bonpl. ex Dunal, Monog. Anon. 67 = uliginosa.
phaeolacdes, *Mart. Fl. Bras.* xiii. t. 45.—Bras.
Pisonis, *Mart. l. c.* 5.—Bras.
polyphylla, *DC. Syst.* i. 499.—Afr. trop.
Porpetae, Boiv. ex Baill. in Bull. Soc. Linn. Par. i. (1882) 841 = senegalensis.
pubescens, Salish. Prod. 380 = A. Chermolia.
punctata, *Aubl. Pl. Gui.* i. 614. t. 247.—Guian.
punicifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 26.—N. Granat.
purpurea, *Moç. & Sesse, ex Dunal, Monog. Anon.* 64.—Mexic.
pygmaea, Bartr. Trav. 18. t. 1 = Asimina pygmaea.
pyxiflorula, *Boj. Hort. Maurit.* 6; *Baker, Fl. Maurit.* 8.—Ins. Mascar.
quindeusis, *H. B. & K. Nov. Gen. et Sp.* v. 60.—N. Granat.
reticulata, *Linn. Sp. Pl.* 537.—Am. trop.
reticulata, Sieber, ex A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 200 = Rollinia mucosa.
reticulosa, Vell. Fl. Flum. 239; v. t. 130 = Pisonia.
rhizantha, *Eichl. in Eichl. Jahrb. Berl. ii.* (1883) 292.—Bras.
rhombipetala, *Ruiz & Pav. Fl. Per.* v. t. 489; *ex G. Don, Gen. Syst.* i. 87.—Peruv.
riparia, *H. B. & K. Nov. Gen. et Sp.* v. 59.—Peruv.
rufa, *Pers., Rel. Haenk.* ii. 75.—Am. trop.
rufescens, Aubl. ex Steud. Nom. ed. I. 51 (lapsu pro Abuta rufescens?).—Guian.

ANONA:—
Salzmanni, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 197.—Bras.
senegalensis, *Pers. Syn.* ii. 95.—Afr. trop.; Guian.
sericea, *Dunal, Monog. Anon.* 69.—Guian.
sessiliflora, *Benth. in Hook. Kew Journ.* v. (1853) 8.—Bras.
silvestris, Vell. Fl. Flum. 239; v. t. 128 = Rollinia sylvatica.
sinensis, *Raeusch. Nom.* ed. III. 160.—China.
sphaerocarpa, *Splitg. in Horven & De Vriese, Tijdschr.* ix. (1842) 96.—Guian.
spinescens, *Mart. Fl. Bras.* xiii. t. 11.—Bras.
squamosa, Delile, Ill. Aegypt. 17 = Forskahlii.
squamosa, *Linn. Sp. Pl.* 537.—Ind. occ.
squamosa, Vell. Fl. Flum. 239; v. t. 127 = obtusiflora.
surinila, *Rosb. Hort. Beng.* 48, nomen.—Ind. or.
sylvatica, St. Hil. Fl. Bras. Mer. i. 32 = Rollinia sylvatica.
sylvestris, Auct.—Cf. *silvestris*.
tenuiflora, *Mart. Fl. Bras.* xiii. t. 10.—Bras.
tenuifolia, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 198.—Guian.
triloba, Linn. Sp. Pl. 537 = Asimina triloba.
tripetala, Ait. Hort. Kew. ii. 252 = A. Cherimolia.
tuberosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 7.—Malaya.
uliginosa, *H. B. & K. Nov. Gen. et Sp.* v. 56.—Peruv.
uncinata, Lam. Encyc. ii. 127 = Artabotrys odoratissima.
unuflora, Dunal, Monog. Anon. 76 = Duguetia Candollei.
velutina, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 132.—Bras.
vepertonum, *Mart. Fl. Bras.* xiii. t. 16.—Bras.
xestropetala, Spreng. Syst. ii. 641 = Rollinia longifolia.
xylopifolia, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvi. (1842) 132.—Bras.

ANONIS, Tourn. ex Scop. Fl. Carn. ed. II. ii. 54 (1772) = Ononis, Linn. (Legum.).
americana, Aubl. Pl. Gui. ii. 768 = Stylosanthes procumbens, viscosa.
arvensis, Lam. Fl. Fr. ii. 611 = O. arvensis.
fruticosa, Lam. l. c. 612 = O. fruticosa.
minutissima, Lam. l. c. 610 = O. minutissima.
Natrix, Scop. Fl. Carn. ed. II. ii. 55 = O. Natrix.
pinguis, Lam. Fl. Fr. ii. 609 = O. Natrix.
pusilla, Lam. l. c. 610 = O. reclinata?
reclinata, Lam. l. c. 611 = O. reclinata.
rotundifolia, Lam. l. c. 612 = O. rotundifolia.
scariosa, Moench, Meth. 157 = O. mitissima.
spinosa, Scop. Fl. Carn. ed. II. ii. 54 = O. spinosa.
variegata, Lam. Fl. Fr. ii. 608 = O. variegata.
viscosa, Lam. l. c. 609 = O. viscosa.

ANONYCHIUM, Schweinf. Reliq. Kotschy. 7. t. 7 (1868) = Prosopis, Linn. (Legum.).
lancolatum, Schweinf. l. c. = P. oblonga.

ANONYMOS, Walt. Fl. Carol. (1788) (passim pro omnibus dubiis).
aquatica, Walt. l. c. 87 = Limnanthemum trachyspermum.
aquatica, Walt. l. c. 230 = Planera aquatica.
aquatica, Walt. l. c. 109 = Saururus cernuus.
aquatica, Walt. l. c. 187 = Nesaea verticillata.
asteroides, Michx. ex DC. Prod. v. 301 = Boltonia asteroides.
bractеata, Walt. Fl. Carol. 181 = Zornia tetraphylla.
capitata, Walt. l. c. 69 = Burmannia capitata.
carolinensis, Walt. l. c. 60 = Elytraria virgata.
carolinensis, Walt. l. c. 91 = Lithospermum hirtum.
carolinensis, Walt. l. c. 164 = Verbena Aubletia.
carolinensis, Walt. l. c. 188 = Ruellia carolinensis.
carolinensis, Walt. l. c. 188 = Amphicarpaea monoica.
cassioides, Walt. l. c. 171 = Seymeria tenuifolia.
ciliata, Walt. l. c. 197 = Listra graminifolia?
corymbosa, Walt. l. c. 189.—Am. bor.
dioscoroides, Croom, in Am. Journ. Sc. xxix. (1836) 165 = Croomia pancifora.
erecta, Walt. Fl. Carol. 160 = Houstonia caerulea.
flamea, Walt. l. c. = Gerardia flava.
frondosa, Walt. l. c. 189.—Am. bor.

ANONYMOS :—
frutescens, Walt. Fl. Carol. 186 = Wistaria speciosa.
graminifolia, Walt. l. c. 197 = Liatris graminifolia ?
lignatrigua, Walt. l. c. 186.—Am. bor.
odoratissima, Walt. l. c. 198 = Trillisia odoratissima.
paniculata, Walt. l. c. = Trillisia paniculata.
paucifolia, Walt. l. c. 170 = Gerardia ?
petiolata, Walt. l. c. 108 = Mitreola petiolata.
pilosa, Walt. l. c. 197 = Liatris pycnostachya.
pinnatifida, S. Wats. Bibl. Ind. 347, lapse = seq.
pinnatus, Walt. Fl. Carol. 108 = Petalostemon corymbosus.
procumbens, Walt. l. c. 86 = Anotis rotundifolia.
pudica, Walt. l. c. 128 = Urvularia perfoliata.
purpurea, Walt. l. c. 170 = Gerardia purpurea.
quaternata, Walt. l. c. 246 = Gerardiae sp.
quinata, Walt. l. c. = Gerardiae sp.
ramosa, Walt. l. c. 108 = Liatris scariosa.
repens, Walt. l. c. 110 = Dichondra repens.
repens, Walt. l. c. 160 = Mitchella repens.
rotundifolia, Walt. l. c. 181 = Crotalaria ovalis.
sagittalis, Walt. l. c. = Crotalaria sagittalis.
sempervirens, Walt. l. c. 98 = Gelsemium sempervirens.
sessilifolia, Walt. l. c. 108 = Mitreola petiolata.
setacea, Walt. l. c. 170 = Gerardia setacea ?
tinctoria, Walt. l. c. 68 = Lachnanthes tinctoria.
umbrosa, Walt. l. c. 63 = Micranthemum orbiculatum.
unifiora, Walt. l. c. 198 = Carphephorus bellidifolius ?

ANOOSPERMA, Kunze, in Bot. Zeit. i. (1848) 661, *err. typ.* = Oncosperma, Blume (Palm.).

ANOPLANTHUS, Endl. Gen. 727 (1839) = Phelypaea, Tourn. et Aphyllon, Mitch. (Orobanch.).
Bieberstenii, Reut in DC. Prod. xi. 42 = P. coccinea.
coccinea, Walp. Rep. iii. 481 = P. coccinea.
comosus, Walp. l. c. 480 = Aphyllon comosum.
fasciculatus, Walp. l. c. = Aphyllon fasciculatum.
Tournefortii, Reut. in DC. Prod. xi. 42 = P. coccinea.
uniflorus, Endl. Ic. Gen. t. 72 = Aphyllon uniflorum.

ANOPLOCARVUM, Ledeb. Fl. Ross. iii. 154 (1847) = Brittsichtum, Schrad. (Borag.).
compressum, Ledeb. l. c.—Rossia.

ANOPLON, Reichb. Consp. 212 ? (1828) = Phelypaea, Tourn. et Aphyllon, Mitch. (Orobanch.).
Bieberstenii, C. A. Mey. Verz. Pfl. Cauc. 104 = Phelypaea coccinea.
bifurum, G. Don, Gen. Syst. iv. 634 = Aphyllon uniflorum.
fasciculatum, G. Don, l. c. 633 = Aphyllon fasciculatum.
Tournefortii, G. Don, l. c. = Phelypaea coccinea.

ANOPLOPHYTUM, Beer, in Flora, xxxvii. (1854) 846 = Tillandsia, Linn. (Bromel.).
atranthos, Beer, Bromel. 49 = T. dianthoidea.
amarum, E. Morr. ex C. Morr. in Belg. Hortic. xxxiii. (1883) 265. t. 17 = T. pulchra.
Benthamianum, Beer, Bromel. 268 = T. Benthamiana.
bicolor, Beer, l. c. 41 = T. stricta.
calothyrsum, Beer, l. c. 263 = T. calothyrsus.
dianthoideum, Beer, l. c. 41 = T. dianthoidea.
didistichum, E. Morr. ex C. Morr. in Belg. Hortic. xxxi. (1881) 164 = T. Lorentziana.
Duratii, Beer, Bromel. 42 = T. Duratii.
flexuosum, Beer, l. c. 43 = T. flexuosa.
graminiforum, E. Morr. ex C. Morr. in Belg. Hortic. xxix. (1879) 226, & xxx. 191. t. 111 = T. gesminiflora.
gulanense, Beer, Bromel. 44.—Guian.
incanum, E. Morr. ex C. Morr. in Belg. Hortic. xxxi. (1881) 209. t. 11 = T. Gardneri.
linare, Beer, Bromel. 42 = T. linearis.
longibracteatum, Beer, l. c. 264 = T. purpurea.
pulchellum, Beer, l. c. 41 = T. pulchra.
Rollizonii, E. Morr. ex C. Morr. in Belg. Hortic. xxix. (1879) 275.—Braz.
roseum, Beer, Bromel 40 = T. dianthoidea.
rubidum, Beer, l. c. = T. geminiflora.
setaceum, Beer, l. c. 265 = T. compressum ?
straminosum, C. Koch, in Ind. Sem. Hort. Berol. (1856) 7 = T. stramines.
Sprengelianum, Beer, Bromel. 266 = T. Sprengelianum.

ANOPLOPHYTUM :—
strictum, Beer, Bromel. 39 = Tillandsia stricta.
strobilanthum, Beer, l. c. 42 = T. strobilantha.
vestitum, Beer, l. c. 265 = T. Benthamiana.
violaceum, Beer, l. c. = T. violacea.
vittatum, Beer, l. c. 48 = T. vittata.

ANOPTERUS, Labill. Pl. Nov. Holl. i. 85. t. 112 (1804). *SAXIFRAGEAE*, Benth. & Hook. f. i. 648.
glandulosus, *Labill. l. c.* 86.—Tasman.
Macleyanus, *F. Muell. in Journ. Pharm. Soc. Vict.* (1858) 43.—Austral.

ANOSMIA, Bemh. in Linnaea, vii. (1832) 608 = Smyrnium, Linn. (Umbell.).
idaea, Bemh. l. c. 609 = S. apiifolium.

ANOSPORUM, Nees, in Linnaea, ix. (1835) 287 = Cyperus, Linn.
Cephalotes, Kunth, in J. rm. As. Soc. Beng. xxxix. (1870) ii. 84 = C. Cephalotes.
Colymbetes, Boeck. in Bot. Zeit. xxvii. (1869) 26 = C. Colymbetes.
cubensis, Boeck. l. c. = Scirpus cubensis.
monocephalum, Nees, in Linnaea, ix. (1835) 287 = C. Cephalotes.
nudicaule, Boeck. in Bot. Zeit. xxvii. (1869) 26 = C. nudicaulis.
pallidum, Boeck. in Linnaea, xxxvi. (1869–70) 412 = C. platystylis.

ANOTIS, DC. Prod. iv. 431 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 99.
affinis, Benth. & Hook. f. Gen. ii. 99 = Leschenaultiana.
boerhaaviolides, *Wight & Arn. in Bull. Acad. Pétersb.* xxix. (1883) 185.—China.
calycina, *Hook. f. Fl. Brit. Ind.* iii. 78.—Ind. or.
capitata, Korth. in Ned. Kruid. Arch. ii. (1851) 151 = Hedyotis stipulata ?
carnosa, *Benth. & Hook. f. Gen.* ii. 59.—Ind. or.
Cervantesii, DC. Prod. iv. 432 = Hedyotis Cervantesii.
ciliolata, G. Don, Gen. Syst. iii. 535 = Houstonia purpurea.
corrulea, G. Don, l. c. = Houstonia caerulea.
confusa, DC. Prod. iv. 431 = Mallostoma conferta.
decipiens, *Hook. f. Fl. Brit. Ind.* iii. 72.—Ind. or.
dentelloides, Steud. Nom. ed. II. i. 701 = Oldenlandia dentelloides.
ericoides, DC. Prod. iv. 438 = Hedyotis ericoides.
filiformis, DC. l. c. 431 = Mallostoma filiformis.
Finlaysoniana, *Benth. & Hook. f. Gen.* ii. 60.—Siam.
foetida, *Benth. & Hook. f. Gen.* ii. 59.—Ind. or.
gentianoides, Walp. Rep. iv. 280 = Houstonia caerulea.
gracilis, *Hook. f. Fl. Brit. Ind.* iii. 71.—Reg. Himal.
hypnoides, DC. Prod. iv. 432 = Hedyotis hypnoides.
ingrata, *Benth. & Hook. f. Gen.* ii. 59 ; *et Hook. f. Fl. Brit. Ind.* iii. 71.—Reg. Himal.
juniperifolia, DC. Prod. iv. 432 = Mallostoma juniperifolia.
lanceaefolia, Benth. & Hook. f. Gen. ii. 59 = lancifolia.
lanceolata, DC. Prod. iv. 433 = Houstonia purpurea.
lancifolia, *Hook. f. Fl. Brit. Ind.* iii. 78.—Ind. or.
laricifolia, DC. Prod. iv. 433 = Mallostoma laricifolia.
Lechleriana, Schlecht. in Linnaea, xxviii. (1856) 488 = Oldenlandia Lechleriana.
Leschenaultiana, *Benth. & Hook. f. Gen.* ii. 59 ; *et Hook. f. Fl. Brit. Ind.* iii. 72.—Ind. or.
longifora, Benth. Pl. Hartw. 28 = Houstonia triflora.
longifolia, G. Don, Gen. Syst. iii. 535 = Houstonia purpurea.
microphylla, DC. Prod. iv. 433 = Hedyotis microphylla.
mitrasemoides, *Benth. & Hook. f. Gen.* ii. 60.—Austral.
monosperma, *Benth. & Hook. f. l. c.* 59 ; *et Hook. f. Fl. Brit. Ind.* iii. 75.—Ind. or.
Montholoni, *Hook. f. Fl. Brit. Ind.* iii. 78.—Ind. or.
nummularia, *Benth. & Hook. f. Gen.* ii. 60 ; *et Hook. f. Fl. Brit. Ind.* iii. 75.—Zeylan.
nummulariformis, Benth. & Hook. f. l. c. = preac.
oxyphylla, *Hook. f. l. c.* 72.—Ind. or.

ANOTIS :—
pillifera, *Schlecht. in Linnaea*, xxviii. (1856) 487.—Peruv.
pterocpora, *Benth. & Hook. f. Gen.* ii. 60.—Austral.
purpurea, G. Don, Gen. Syst. iii. 535 = Houstonia purpurea.
quadricleolaria, *Benth. & Hook. f. Gen.* ii. 59 ; *et Hook. f. Fl. Brit. Ind.* iii. 74.—Ind. or.
Rheedii, *Benth. & Hook. f. l. c. ; et Hook. f. l. c.* 78.—Ind. or.
Richardiana, *Benth. & Hook. f. l. c. ; et Hook. f. l. c.* 75.—Ind. or.
Ritchiei, *Hook. f. Fl. Brit. Ind.* iii. 74.—Ind. or.
rotundifolia, *DC. Prod.* iv. 433.—Am. bor.
Salzmanni, DC. l. c.—Oldenlandia Salzmanni.
serpens, *DC. l. c.*—Am. austr.
serpyllifolia, G. Don, Gen. Syst. iii. 535 = Houstonia serpyllifolia.
setosa, DC. Prod iv. 432 = Mallostoma setosa.
tenella, G. Don, Gen. Syst. iii. 535 = Houstonia tenella.
Thwaitesiana, *Masm. in Bull. Acad. Pétersb.* xxix. (1883) 163.—China.
thymifolia, DC. Prod. iv. 432 = Mallostoma thymifolia.
urophylla, *Benth. & Hook. f. Gen.* ii. 59 ; *et Hook. f. Fl. Brit. Ind.* iii. 72.—Ind. or.
Wightiana, *Benth. & Hook. f. l. c.* 60 ; *et Hook. f. l. c.* 75.—Ind. or.

ANPLECTRUM, A. Gray, Bot. U. St. Expl. Exped. i. 597 (1854). *MELASTOMACEAE*, Benth. & Hook. f. i. 758.
Aplectrum, Blume, in Flora, xiv. (1831) 502.
annulatum, *Triana, in Trans. Linn Soc.* xxviii. (1871) 84.—Penang.
assamicum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 546.—Reg. Himal.
bashatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 84.—Burma.
biligulatum, *Triana, l. c.* 85.—Sumatr.
conbus, *Blume, Mus. Bot. Lugd. Bat.* i. 88.—Sumatr.
cyanocarpum, Triana, in Trans. Linn. Soc. xxviii (1871) 84 = glaucum.
divaricatum, *Triana, l. c.*—Ind. or.
glaucum, *Triana, l. c.*—Malaya.
Korthalsii, *Triana, l. c.* 85.—Borneo.
latifolium, *Triana, l. c.*—Borneo
ligulatum, *Triana, l. c.*—Java.
myrtiforme, Nand. in Ann. Sc. Nat. Sér. III. xv. (1851) 306 = Medinilla myrtiformis.
myrtifolium, Miq. Fl. Ind. Bat. i. t. 555 = Medinilla myrtiformis.
nodosum, *Blume, Mus. Bot. Lugd. Bat.* i. 87. Sumatr.
nodosum, *Blume, l. c.* 37 —Sumatr.
ovalifolium, A. Gray, Bot. U. St. Expl. Exped. i. 597 = Allomorphia ovalifolia.
pallens, *Blume, Mus. Bot. Lugd. Bat.* i. 88.—Malaya.
parviflorum, Benth. Fl. Hongk. 116 = Blastus cochinchinensis.
reformatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 85.—Borneo.
rostratum, *Blume, in Flora*, xiv. (1831) 502.—Malaya.
stipulare, *Blume, l. c.* 503.—Java.
viminale, *Blume, l. c.* 509.—Malaya.

ANQUETILIA, Decne. in Jacquem. Voy. Bot. 161. t. 161 (1848) = Skimmia, Thunb. (Rutac.).
Laureola, Decne. l. c. = S. Laureola.

ANREDERA, Juss. Gen. 84 (1789). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 78, 1219.
Baseana, Steud. Nom. ed. II. i. 199 (1840).
Claibisia, Benth. & Hook. f. Gen. iii. 78 (1880).
Clarisia, Abat, in Act. Soc. Med. Sev. x. 418 ; ex Spreng. Cur. i. 909 (1830).
Serbera, Presl, in Oken, Isis, xxi. (1828) 375.
Cumingii, Hassk. in Flora, xlviii. (1865) 401 = spicata.
scandens, Moq. in DC. Prod. xiii. ii. 230 = spicata.
spicata, *Pers. Syn.* i. 207.—Am. trop. ; Ins. Philipp.
vesicaria, Gaertn. f. Fruct. iii. 176. t. 218 = spicata.
ventricosa, Poir. Encyc. Suppl. i. 391 = spicata.

ANSELLIA, Lindl. Bot. Reg. xxx. (1844) sub t. 12. *ORCHIDEAE*, Benth. & Hook. f. iii. 587.
africana, *Lindl. l. c.*—Afr. trop.
gigantea, *Reichb. f. in Linnaea*, xx. (1847) 978.—Natal

ANSERINA, Dum. Fl. Belg. 21 (1827) = **Chenopodium**, Linn.
Bonus-Henricus, Dum. l. c. = C. Bonus-Henricus.

ANSONIA, Bert. ex Hemsl. Bot. Voy. Challenger, i. 111 53 (1884) = **Lactoris**, Phil. (Piperac.).
nodulosa, Bert. ex Hemsl. l. c. = L. fernandezia.

ANSONIA, Rafin. New Fl. Am. iv. (1836) 58 = **Amsonia**, Walt. (Apocyn.).
elliptica, Rafin. l. c.—Am. bor.
latifolia, Rafin. l. c. = Amsonia Tabernaemontana.
salicifolia, Rafin. l. c. = Amsonia Tabernaemontana.
tenuifolia, Rafin. l. c.—Am. bor.

ANSTRUTHERIA, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 344. t. 4 = **Welfia**, Spreng. (Rhizoph.).
africana, Benth. in Journ. Linn. Soc. iii. (1859) 78 = W. africana.
ceylanica, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 345 = W. ceylanica.
madagascariensis, Benth. in Journ. Linn. Soc. iii. (1859) 78 = W. madagascariensis.
zeylanica, Henslow, in Hook f. Fl. Brit. Ind. ii. 441 = W. ceylanica.

ANTACANTHUS, Rich. ex DC. Prod. iv. 484 (1830) = **Scoloscanthus**, Vahl (Rubiac.).

ANTAGONIA, Griseb. in Goett. Abh. xix. (1874) 144 = **Cayaponia**, Manso (Cucurb.).
citrullifolia, Griseb. l. c. 145 = C. citrullifolia.
Sandia, Griseb. in Lorentz, Veg. del Nordest. Prov Entre Rios, 69 = C. Sandia.

ANTAUREA, Neck. Elem. i. 70 (1790) = **Centaurea**, Linn. (Compos.).

ANTELAEA, Gaertn. Fruct. i. 277. t. 68 (1788). Genus inc. sed.
canescens, *Cels, ex Heynh. Nom.* ii. 38.—Hab ?
javanica, *Gaertn. Fruct.* i. 277.—Java.

ANTENNARIA, Gaertn. Fruct. ii. 410. t. 167 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 301.
DIEVANTHUS, Rafin. in Am. Monthly Mag. (1818) 268.
alpina, *Gaertn. l. c.*—Reg. bor.
andina, *A. Gray, in Proc. Am. Acad.* v. (1861) 138.—Am. austr.
arctioides, *A. Gray, l. c.*—Peruv.
argentea, *Benth. Pl. Hartweg.* 319.—Calif.
carpathica, *R. Br. in Trans. Linn. Soc.* xii. (1818) 123. —Reg. temp. bor.
chilensis, *Remy, in C. Gay, Fl. Chil.* iv. 285.—Chili.
cinnamomea, DC. Prod. vi. 270 = Anaphalis cinnamomea.
contorta, D. Don, in Bot. Reg. t. 605 = Anaphalis contorta.
dimorpha, *Torr. & Gray, Fl. N. Am.* ii. 431.—Am. bor.
dioica, *Gaertn. Fruct.* ii. 410. t. 167. f. 3.—Europ.; As. bor.
flagellaris, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 219.—Am. bor.
Geyeri, *A. Gray, Fl. Pendl.* 107.—Am. bor.
hyperborea, D. Don, in Engl. Bot. Suppl. t. 2640 = dioica.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 178 = Anaphalis margaritacea.
japonica, Sch. Bip. in Zoll. Syst. Vers. Ind. Archip. 196; Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 200 = prasc.?
javanica, *DC. Prod.* vi. 271.—Java.
laboradorica, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 406 = alpina.
leontopodina, DC. Prod. vi. 269 = Leontopodium albidum.
Leontopodium, Gaertn. Fruct. ii. 410 = Leontopodium alpinum.
leucophylla, Holl. & Reichh. in Flora, xlii. (1859) 382 = Helichrysum obconicum.
linearifolia, *Wedd. Chlor. And.* i. 150.—Bolivia.
lanuginosa, *Torr. & Gray, Fl. N. Am.* ii. 430.—Am. bor. occ.
magellanica, *Sch. Bip. in Flora*, xxxviii. (1855) 117.—Reg. Magell.
margaritacea, R. Br. in Trans. Linn. Soc. xii. (1817) 139 = Anaphalis margaritacea.

ANTENNARIA:—
Meredithae, F. Muell. in Papers Roy. Soc. Tasm. (1870) 15 = Leontopodium Meredithae.
microcephala, *A. Gray, in Proc. Am. Acad.* x. (1874) 74.—Calif.
monocephala, DC. Prod. vi. 269 = alpina.
monoica, *Wedd. Chlor. And.* i. 150.—N. Granat.
montana, S. F. Gray, Nat. Arr. ii. 458 = dioica.
muscoides, *Hook. f. & Thoms. ex C. B. Clarke, Comp Ind.* 100.—Reg. Himal.
nana, *Hook. f. & Thoms ex C. B. Clarke, l. c.*—Reg. Himal.
nivalis, Tenore, ex Nym. Consp. 383 = Leontopodium alpinum
nubigena, F. Muell. in Trans Phil. Soc. Vict. i. (1855) 45 = Raoulia catipes.
parvifolia, Nutt. in Trans. Am Phil. Soc. N. S. vii. (1841) 406 = dioica.
Planchoni, F. Muell. Census, 80 = Raoulia Planchoni.
plantaginifolia, *R. Br. in Trans. Linn Soc.* xii. (1817) 123.—Am. bor.
plantaginifolia, Hook. Fl. Bor. Am. i. 330 = plantaginea.
racemosa, *Hook. l. c.*—Am. bor.
rubicunda, C. Koch, in Linnaea, xvii. (1843) 49 = Helichrysum arenarium.
saxatilis, *DC. Prod.* vi. 271.—Java.
semidecurrens, DC. l. c. = Anaphalis araneosa.
sordida, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 194.—Java.
Steetziana, *Turcz. in Bull. Soc. Nat. Mosc.* (1857) i. Acid. p. xxxix.—Dahuria.
stenophylla, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 213.—Am. bor.
triplinervis, Sims, Bot. Mag. t. 2468 = Anaphalis triplinervis.
uniceps, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 106.—Austral.
villosissima, D. Don, Prod. Fl. Nep. 175 = Anaphalis araneosa.
viscida, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 194. —Malaya.

ANTENORON, Rafin. Fl. Ludov. 28 (1817) = **Polygonum**, Linn.
racemosum, Rafin. l. c. = P. virginianum.

ANTEPHORA, Steud. Syn. Pl. Gram. 111 (1855) = **Anthephora**, Schreb. (Gram.).

ANTERISCIUM, Meyen, Reise, i. 315 (1834) sphalm. = **Asterisicum**, Cham. & Schlecht. (Umbellif.).

ANTHACANTHA, Lem. Illustr. Hortic. (1858) Misc. 64 = **Euphorbia**, Linn.
Desmetiana, Lem. l. c. = E. hexagona.

ANTHACANTHUS, Nees, in DC. Prod. xi. 460 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1097.
CRATIOLA, Rafin. Fl. Tellur. iv. 64 (1836).
OPLONIA, Rafin. l. c.
aciculatus, *Nees, l. c.*—Jamaica.
arinatus, *Nees, l. c.*—Jamaica.
blepinoros, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 195.—Ind. occ.
cuneatus, *Nees, in DC. Prod.* xi. 461.—Jamaica.
cuneifolius, Nees, l. c. 482 = cuneatus.
emarginatus, *Nees, l. c.* 461.—Ins. S. Thom.
jamaicensis, *Griseb. Fl. Brit. W. Ind.* 457.—Jamaica.
lycioides, *Nees, in DC. Prod.* xi. 461.—Ind. occ.
microphyllus, *Nees, l. c.*—Ind. occ.
purpurascens, *Griseb. Cat. Pl. Cub.* 198.—Cuba.
repandus, *Nees, in DC. Prod.* xi. 462.—Ins. Tanna.
sinuatus, *Nees, l. c.*—Ins. Tanna.
spinosus, *Nees, l. c.* 460.—Am. austr.
Sprengelii, *Nees, l. c.* 463.—Ins. S. Doming.
tetrastichus, *Griseb. Cat. Pl. Cub.* 197.—Cuba.
vincoides, *Nees, in DC. Prod.* xi. 461.—Madag.

ANTHACANTHUS, Rich. ex DC. Prod. iv. 484 (1830) = **Scoloscanthus**, Vahl (Rubiac.).

ANTHACTINIA, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 139 = **Pentaphora**, Linn.
Horsfieldii, M. Roem. Syn. Pepon. 191 = P. Horsfieldii.
longipes, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 139 = P. longipes.

ANTHACTINIA:—
moluccana, M. Roem. Syn. Pepon. 190 = Passiflora moluccana.
nepalensis, M. Roem. l. c. = P. nepalensis.
penangiana, M. Roem. l. c. 192 = Modecca nicobarica.
singaporiana, M. Roem. l. c. = Modecca singaporiana.
sumatrana, M. Roem. l. c. 191 = P. sumatrana.
timorensis, M. Roem. l. c. = P. moluccana.
Walkeri, M. Roem. l. c. = P. suberosa.

ANTHADENIA, Lem. Hort. Vanhoutt. fasc. i. 4 (1845); Fl. des Serres, ii. (Avr. 1846) 10. t. 6 = **Besanum**, Linn. (Pedalin.).
sesamoides, Lem. ll. cc.—Afr. trop.

ANTHAERANTIA, Beauv. Agrost. 48. t. 10. f. 7 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1098.
ANTHENANTIA, R. Br. in Flind. Voy. ii. App. iii. 593 (1814).
AULAXANTHUS, Ell. Sketch Bot. Carol. i. 102 (1817)
AULAXIA, Nutt. Gen. Am. i. 47 (1818).
LEPTOCORYPHIUM, Nees, Agrost. Bras. 88 (1824).
lanata, *Benth. in Journ. Linn. Soc.* xix. (1882) 39.—Mexic.
rufa, *Schult. Mant.* ii. 258.—Am. bor.
villosa, *Beauv. Agrost.* 48. t. 10. f. 7.—Am. bor.

ANTHAERIUM, Schott, in Oestr. Bot. Zeitschr. viii (1858) 179 *err. typ.*—**Anthurium**, Schott (Aroid.).

ANTHALOGEA, Rafin. New Fl. Am. iv. 88 (1836) = **Polygala**, Linn.
polygama, Rafin. l. c. = P. rubella.

ANTHANEMA, Rafin. Fl Tellur. iv. 90 (1836) = **Cuscuta**, Linn. (Convolv.).
capitata, Rafin. l. c. = C. capitata.
paradoxa, Rafin. l. c. = C. glomerata.
verticillata, Rafin. l. c. = C. glomerata.

ANTHANOTIS, Rafin. Fl. Ludov. 52, 149 (1817) = **Asclepiodora**, A. Gray (Asclep.).
pedicellata, Rafin. l. c. 52 = Podostigma pubescens.
procumbens, Rafin. l. c. 149.—Am. bor.
viridis, Rafin. l. c. 52 = Asc. viridis.

ANTHEDOSORUS, A. Gray, in Hook. Kew Journ. iii. (1851) 178 = **Myrtocoophalus**, Benth. (Compos.).
gracilis, A. Gray, l. c. = M. gracilis.

ANTHEILEMA, Rafin. Fl. Tellur. iv. 66 (1836) = **Phaylopsis**, Willd. (Acanth.).
imbricata, Rafin. l. c. = P. longifolia.
paniculata, Rafin. l. c. = Ruellia paniculata.

ANTHEISCHIMA, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 137 t. 27 (1840) = **Gordonia**, Ellis (Ternstroem.).
excelsa, Korth. l. c. = G. excelsa.

ANTHELIA, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 127 (1863–64). *AROIDEAE*, Benth. & Hook. f. iii. 968
nobilis, *Schott, l. c.*—Celebes.

ANTHELIS, Rafin. Chl. Astn. 9 (1813) = **Helianthemum**, Tourn. (Cistin.).
albidus, Rafin. l. c.—Italia.
arenaria, Rafin. New Fl. Am. iii. 81.—Am. bor.
glaucus, Rafin. Chl. Astn. 9 = H. glaucum.
glutinosa, Rafin. l. c. = H. glutinosum.
laevipes, Rafin. l. c. = H. laevipes.
podanisia, Rafin. New Fl. Am. iii. 80.—Am. bor.

ANTHELMINTHICA, P. Br. Hist. Jamaic. 156 (1756) = **Spigelia**, Linn. (Logan.).

ANTHEMA, Medic. Malv. 42 (1787) = **Lavatera**, Tourn. (Malvac.).
arborea, Medic. l. c. = L. arborea.
cretica, Medic. l. c. = L. cretica.
scabra, Moench, Meth. 613 = L. cretica.
Tenoreana, Presl, Fl. Sicul. i. 181 = L. arctica.

ANTHEMIOPSIS, Boj. ex DC. Prod. v. 547, 548 (1836) = **Wedelia**, Jacq. (Compos.).
elongata, Boj. l. c. 548 = W. elongata.
macrophylla, Boj. l. c. 547.—Afr. trop.

ANTHEMIS, Mich. ex Linn. Syst. ed. I. (1735).
COMPOSITAE, Benth. & Hook. f. ii. 420.
CHAMAEMELUM, Tourn. ex Adans. Fam. ii. 128 (1763). CHAMOMILLA, Godr. Fl. Lorr. ii. 19 (1843). COTA, J. Gay, in Guss. Fl. Sic. Syn. ii. 866 (1844). LEPIDOPHORUM, Neck. Elem. i. 14 (1790). LEUCANTHEMUM, Adans. Fam. ii. 127 (1763). LYONNETIA, Cass. in Dict. Sc. Nat. i. 106 (1827). MARCELIA, Cass. l. c. xxxiv. 107 (1825). MARUTA, S. F. Gray, Nat. Arr. ii. 456 (1821). MATRICARIA, Tourn. ex Adans. Fam. ii. 127 (1763). ORMENIS, Cass. in Dict. Sc. Nat. xxix. 180 (1823). PYRETHRARIA, Webb, Itin. Hisp. 37 (1838). RHETHOLEPIS, Cass. in Bull. Soc. Bot. Fr. iii. 737 (1856)

abrotanifolia, *Guss. Fl. Sic. Syn.* ii. 490.—Italia; Lasitan.
abrotanifolia, Willk. in Bot. Zeit. v. (1847) 858 — montana.
abstrithifolia, *Boiss. & Sprun. Diagn.* Ser. I. iv. 8.—Caria.
abyssinica, J. Gay, ex A. Rich. Tent. Fl. Abyss. i. 418 = A. Cotula.
abyssinica, *Sch. Bip ex Schweinf. Beitr. Fl. Aethiop.* 142, nomen.—Abyss.
aciphylla, *Boiss. Diagn.* Ser. I. iv. 7.—As. Min.
adenolifolia, *Boiss. Fl. Orient.* iii. 317.—Armenia.
aetnensis, *Schouw, in Spreng. Syst.* iii. 596.—Mont. Aetna.
afra, Burm. f. Fl. Cap. Prod. 97 = Omites dentata.
afra, *Linn. Syst.* ed. X. 1293.—Afr. austr.
ageratifolia, Sibth. & Sm. Fl. Graec. ix. 68. t. 887 = Achillea ageratifolia.
agrestis, *Presl, Fl. Sic.* p. xxix, nomen.—Sicilia.
agrestis, Wallr. Sched. Crit. 484 = arvensis.
aizoides, Boiss. & Orph. ex Hook. f. Bot. Mag. t. 6117 = Achillea ageratifolia.
Aizoon, Griseb. Spicil. Fl. Rumel. ii. 210 = Achillea ageratifolia.
albida, *Boiss. Diagn.* Ser. I. iv. 10.—As. Min.
alpina, Gouan, Fl. Monsp. 370 = montana.
alpina, *Linn. Cent.* ii. 31. n. 195.—Europ.
alpina, Moric. ex Nym. Consp. 340 = carpatica.
altaica, *Fisch. Hort. Gorenk.* ed. II. (1812) 41.—Sibir. alt.
altissima, Boiss. Diagn. Ser. I. xi. 12 = coelopoda
altissima, Bell. in Mem. Acad. Torin. v. (1790-91) 246 = peregrina.
altissima, Guss. ex Nym. Consp. 359 = brachycentros.
altissima, *Linn. Sp. Pl.* 893.—Europ.; Orient.
americana, Linn. l. c. 895 = Chrysanthellum procumbens.
americana, Mutis, in Linn. f. Suppl. 378 = Spilanthes Mutisii.
ammophila, *Boiss. & Heldr. Diagn.* Ser. I. xi. 9.—Pamphyl.
Anacyclus, J. Gay, ex Boiss. Fl. Orient. iii. 299 = orctica.
anatolica, *Boiss. Diagn.* Ser. I. xi. 10.—As. Min.
anemonifolia, Boj. ex Steud. Nom. ed. II. i. 102 = Epallage anemonifolia.
anglica, Spreng. Syst. iii. 594 = arvensis.
apennina, *Tausch, in Syll. Ratisb.* ii. 250.—Europ.
apiifolia, R. Br. in Bot. Reg. t. 527 = parthenoides.
arabica, Linn. Sp. Pl. 896 = Cladanthus arabicus.
arabica, Vis. Fl. Lib. Spec. 56 = Cyrtolepis alexandrina.
arenicola, *Boiss. Fl. Orient.* iii. 306.—Cilicia.
argaea, Boiss. & Bal. Diagn. Ser. II. vi. 97 = ptarmicaeformis.
artemisiaefolia, Willd. Sp. Pl. iii. 2184 = Chrysanthemum sinense.
artemisiifolia, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 421 = pracc.?
arvensis, *Linn. Sp. Pl.* ed. I. 894.—Europ.; Orient.
arvensis, Pall. ex Bieb. Fl. Taur. Cauc. ii. 330 = altissima.
asperula, *Bertol. Fl. Ital.* ix. 368.—Corsica.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 289.—Afr. bor.
auct., *Schaur, ex Heinzerl. in Denkschr. Acad. Wien*, xivii. (1884) 187.—Europ. austr.
Auckeri, Boiss. Diagn. Ser. I. xiii. 77, sphalm. = Achillea Auckeri.
aurea, *Prot. Fl. Lusit.* i. 394.—Europ. austr.
aurea, DC. Prod. vi. 7 = nobilis.
aurea, Webb, ex Nym. Consp. 375 = Matricaria aurea.
auriculata, *Boiss. Diagn.* Ser. I. iv. 5.—As. Min.
australis, Sibth. & Sm. Fl. Graec. ix. 67 = cyparissias.
australis, Willd. Sp. Pl. iii. 2177 = incrassata.

ANTHEMIS:—
austriaca, *Jacq. Fl. Austr.* v. 22.—Europ.; Oriens.
Barrelieri, Tenore, Fl. Nap. ii. 345 = Achillea Barrelieri.
Baumgartneriana, Schur, Enum. Pl. Transs. 384 = montana.
Belidiastrum, Linn. Syst. ed. X. 1223 = Omites Belidiastrum.
biaristata, DC. Fl. Fr. iv. 204 = Anacyclus clavatus.
Biebersteiniana, *C. Koch, in Linnaea*, xxiv. (1851) 320. —Oriens.
Blanchenna, *Boiss. Fl. Orient.* iii. 292.—Syria.
Bollei x, Asch. ex Nym. Consp. ii. 368 = A. Cotula.
Bourgei, *Boiss. & Reut. Pugill. Pl. Hispan.* 56.—Hispan.
Boveana, *J. Gay, in Expl. Sc. Algér.* t. 60. f. 2.—Algeria.
Bowei, Willk. & Lange Prod. ii. 91 = pracc.
Brachmanni, *Boiss. & Heldr. in Boiss. Diagn.* Ser. I. vi. 84.—Graecia.
brachycentros, *J. Gay, in Koch, Syn. Fl. Germ.* ed. II. 414.—Graecia; As. Min.
brachyglossa, C. Koch, in Linnaea, xxiv. (1851) 318 = tinctoria.
bupthalmoides, Jacq. Hort. Schoenb. ii. 13 = Heliopsis buphthalmoides.
byzantina, C. Koch, in Linnaea, xxiv. (1851) 321 = montana.
caespitosa, Herbich, in Flora, xl. (1857) 509 = tenuifolia.
cairica, Vis. Fl. Quaed. Aegypt. 36. t. 6.—Aegypt.; Arab.
callichroa, Boiss. Diagn. Ser. I. xiii. 77, sphalm. = Achillea callichroa.
campylocarpa, Boiss. Fl. Orient. iii. 319 = tripolitana.
candicans, *Boiss. l. c.* 293.—Lycia.
candidissima, *Willd. ex Spreng. Syst.* iii. 593.—Oriens; Lycia.
canescens, *Bert. Fl. Lusit.* i. 395.—Lusit.
canescens, Tenore, ex Nym. Consp. 359 = brachycentros.
carnea, *Boiss. Diagn.* Ser. I. iv. 4.—As. Min.
carpatica, *Willd. Sp. Pl.* iii. 2179.—Europ.; As. Min. Cassia, *Boiss. Diagn.* Ser. I. xi. 10.—Graecia.
caucasica, Adam, in Weber & Mohr, Beitr. i. 71 = Chrysanthemum caucasicum.
caucasica, Henckel, in Dumersm. Pl. Select. H. Halens. 9 = tinctoria.
caucasica, Hornem. Hort. Hafn. 832 = Triumfetti.
caucasica, C. A. Mey. Ver. Pfl. Cauc. 75 = Iberica.
Chamomilla, Tenore, Syll. 440 = brachycentros.
Chamomilla, *Willd. Enum. Hort. Berol.* 910.—'Europ. austr.'
Chamomilla-romana, Crantz, Inst. i. 300 = nobilis.
chamomilloides, Steven, ex Bieb. Fl. Taur. Cauc. ii. 331 = candidissima.
chia, *Linn. Sp. Pl.* 894.—Graecia; As. Min.
chilensis, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 347.—Chili.
chronia, Nym. Syll. Suppl. 9 = montana.
chrysantha, *J. Gay, in Expl. Sc. Algér.* t. 60. f. 9.—Afr. bor.
chrysantha, Schur, Enum. Pl. Transs. i. 331 = tinctoria.
chrysocephala, Boiss. & Reut. Diagn. Ser. II. iii. 25 = montana.
chrysolenca, *J. Gay, ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 50.—Algeria.
ciliata, *Boiss. Fl. Orient.* iii. 312.—Reg. Caucas.
cineres, *Pauk. Elem.* 80.—Europ. austr.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 50.—Afr. bor.
clementa, Desf. Fl. Atlant. ii. 287 = Anacyclus clavatus.
clavata, *Guss. Fl. Sic. Syn.* ii. 865.—Sicilia.
coarctata, Sibth. & Sm. Fl. Graec. ix. 68 = tinctoria.
coelopoda, *Boiss. Diagn.* Ser. I. xi. 12.—As. Min.; Syria.
collina, Jord. Cat. Jard. Dijon, (1848) 16 = montana.
concinna, Guss. Fl. Sic. Syn. ii. 870 = intermedia.
confusa, *Pomel, Nouv. Mat. Fl. Atl.* 49.—Afr. bor.
coronopifolia, *Willd. Sp. Pl.* iii. 2176.—Hispan.
Cornucopiae, *Boiss. Diagn.* Ser. I. xi. 8.—Palaest.
corymbosa, Haenke, in Jacq. Coll. ii. 78 = Achillea stricta.
corymbulosa, *Boiss. & Hausk. in Boiss. Fl. Orient.* iii. 316.—Mesopot.
Cosentejana, Reichb. f. Ic. Fl. Germ. xvi. 68. t. 1000 = austriaca.
cossyrensis, Guss. Fl. Sic. Syn. ii. 870 = secundiramea.
Cota, Koch, Syn. Fl. Germ. 414 = brachycentros.
Cota, *Linn. Sp. Pl.* 893.—Europ. austr.
Cota, Savi, ex DC. Prod. vi. 11 = austriaca.

ANTHEMIS:—
Cota, Sibth. & Sm. Fl. Graec. t. 880 = mixta.
Cotula, Blanco, Fl. Filip. ed. I. 885 = Eclipta erecta.
Cotula, *Linn. Sp. Pl.* 894.—Europ.; Afr. bor.; Orient.
Cotula-foetida, Crantz, Inst. i. 300 = A. Cotula.
cotuloides, Rafin. ex DC. Prod. vi. 18 = A. Cotula.
crassifolia, Humb. ex Steud. Nom. ed. II. i. 102 = Iva cheiranthifolia.
crassipes, Boiss. Diagn. Ser. I. xi. 9 = hyalina.
cretica, *Linn. Sp. Pl.* 895.—Ag. Mediterr.
cronia, Boiss. & Heldr. Diagn. Ser. II. iii. 22 = montana.
Cupaniana, Tod. ex Nym. Consp. 360 = montana.
cypria, *Boiss. Fl. Orient.* iii. 300.—Ins. Cyprus.
cyrenaica, *Cass. in Bull. Soc. Bot. Fr.* xix. (1872) 82.—As. Min.
dalmatica, *Scheele, in Linnaea*, xviii. (1844) 464.—Europ. austr.
damascena, *Boiss. & Gaill. in Boiss. Fl. Orient.* iii. 315.—Syria.
dentata, Boj. ex DC. Prod. vi. 4 = Epallage dentata.
dentata, DC. Prod. vi. 291, in syn. = Omites anthemoides.
dentata, Moench, Meth. 606 = austriaca.
deserti, *Boiss. Fl. Orient.* iii. 305.—Arab.
diffusa, Salzm. ex DC. Prod. vi. 8 = arvensis.
discoidea, *Spreng. Nachtr. i. Bot. Gart. Halle*, 11.—Hab.?
discoides, *Willd. Sp. Pl.* iii. 2187 = tinctoria.
divergens, *Boiss. Fl. Orient.* iii. 300.—Cappadoc.
dregeana, Nest. ex Nym. Consp. 361 = maritima.
dubia, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) 11. 290 = arvensis.
Duriaei, Delile, Ind. Sem. Hort. Monsp. (1843) ex Flach. & Mey. Ind. Sem. Hort. Petrop. xi. 50 = chrysolenca.
eriocephala, *J. & C. Presl, Delic. Prag.* 100.—Sicilia.
eximia, *Boiss. Fl. Orient.* iii. 292.—Reg. Caucas.
fallax, Willd. Enum. Hort. Berol. Suppl. 60 = fuscata.
fimbriata, *Boiss. Fl. Orient.* iii. 306.—Cilicia.
fontida, Lam. Fl. Fr. ii. 166 = A. Cotula.
fruticosa, C. Sm. ex DC. Prod. vi. 14 = Gonospermum revolutum.
fruticosa, *Linn. Cent. Pl.* ii. 31. n. 196; *Amoen. Acad.* iv. 330.—Ad. trop.
fruticulosa, *Bieb. Fl. Taur. Cauc.* ii. 329.—Reg. Caucas.
fumariaefolia, *Boiss. Fl. Orient.* iii. 304.—Cappadoc.
fumariolides, *Hockst. in Lorent, Wanderungen*, 330.—Armen.
fungosa, *Boiss. & Hausk. in Boiss. Fl. Orient.* iii 318.—Persia.
fuscata, *Brot. Fl. Lusit.* i. 394.—Reg. Mediterr. occ.
Garcini, Burm. f. Fl. Ind. 183 = Anvillea Garcini.
Gayana, *Boiss. Fl. Orient.* iii. 318.—Persia; Beluchist.
Gemmellari, *Tineo, Pl. Rar. Sicil.* 47.—Mont. Etna
Gerardiana, Jord. Obs. Pl. Crit. vii. (1849) 31 = montana.
glabra, Jacquem. ex Decne. in Jacquem. Voy. Bot. iv. 88 = Allardia glabra.
globosa, Orteg. Hort. Matr. 46 = Zaluzania globosa.
graminensis, Boiss. Elench. 60 = arvensis.
grandiflora, Host, Fl. Austr. ii. 506 = carpatica.
grandiflora, Jacquem. Voy. Bot. iv. 87 = Allardia tomentosa.
grandiflora, *Des Moul. in Act. Linn. Soc. Bord.* xx. (1855) 556.—Gallia.
grandiflora, Ramatuelle, in Journ. Hist. Nat. Par. ii. (1792) 294 = Chrysanthemum sinense.
granulata, *Pomel, Nouv. Mat. Fl. Atl.* 290.—Afr. bor.
graveolens, Sturm, in Verh. Siebenb. Ver. Naturw. iii. (1852) (Sert. Fl. Transs.) 40 = carpatica.
graveolens, *Boiss. Fl. Orient.* iii. 295.—Lycia.
Guicciardeti, Haldr. & Sart. ex Boiss. Fl. Orient. iii. 308 = peregrina.
hallimifolia, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 284.—Algeria.
halophila, *Boiss. & Bal. in Boiss. Fl. Orient.* iii. 285. —Cilicia.
Haussknechtii, *Boiss. & Reut. i. c.* 310.—Syria; Mesopot.
Heynoldi, Janka, in Oestr. Bot. Wochenbl. vi. (1856) 1 = arvensis.
hebeonica, *Boiss. & Kotschy, Diagn.* Ser. II. v. 108.—Palaest.
hemisphaerica, Herbich, Strp. Rar. Bucov. 59 = macranthes.
hemisphaerica, *Boiss. Fl. Orient.* iii. 314.—Persia.

ANTHEMIS :—

heraclastica, Boiss. & Heldr. ex Nym. Consp. 362 = peregrina.

heterophylla, *Ball, in Journ. Linn. Soc.* xvi. (1878) 507.—Marocco.

heterophylla, *Willd. Britr. Fl. Hercyn.* 231.—Europ.

hirpina, Tenore, ex Guss. Fl. Sic. Syn. ii. 868 = montana.

hispanica, Zuccagni, in Roem. Collect. 155 = mixta.

hondoensis, Willd. ex Steud. Nom. ed. II. i. 102 = Wedelia acapulcensis.

hungarica, Hornem. Hort. Hafn. Suppl. 101 = austriaca.

hyalina, *DC. Prod.* vi. 4.—Syria; Mesopot.

hydruntina, *Groves, in Journ. Linn. Soc.* xxi (1885) 588.—Ital.

Iberica, *Bieb. Fl. Taur. Cauc.* ii. 328.—Reg. Caucas.

incisa, *Viviani, Fl. Ital.* 9.—Italia.

inconspicua, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxvii. (1865) i. 409 = Matricaria discoidea.

incrassata, Link, Enum. Hort. Berol. ii. 345 = Anacyclus clavatus.

incrassata, *Lois.* Not. 129 = arvensis.

indurata, *Delile, Fl. Egypte*, 274. t. 47. f. 3.—Aegypt.

inodora, Gilib. Fl. Lituan. i. 228 = arvensis.

intermedia, *Guss. Fl. Sic. Syn.* ii. 491.—Sicil.

Italica, *Linn. Syst.* ed. X. 1228.—Ital.

kahirica, Boiss. Fl. Orient. iii. 316 = calrica.

Kitaibelii, DC. Prod. vi. 7 = carpatica.

Kitaibelii, Sprengel Syst. iii. 592 = montana.

Kotschyana, *Boiss. Diagn.* Ser. II. iii. 28.—Cilicia.

lanuginosa, Moench, Meth. 606 = Anacyclus radiatus.

laxa, *Spreng. Fl. Hal. Mant.* 62.—Hab.?

leucanthifolia, Linn. Fl. Rar. Afr. 33 = Osmites anthemoides.

leucanthemifolia, *Boiss. & Blanch. Diagn.* Ser. II. iii. 80.—Syria.

leucanthemoides, *Boiss. l. c.* Ser. I. iv. 9.—Smyrna.

leucocephala, Schur, Enum. Pl. Transs. 334 = montana.

libanotica, DC. Prod. vi. 9 = A. chia.

Lindleyi, Nym. Syll. 7 = montana.

lithuanica, Bess. ex DC. Prod. vi. 18 = A. Cotula.

littoralis, Roj.-Clem. Exs. Vit. And. 807 = maritima.

lutescens, Carv. in La Lucr. et Lex. Nov. Veg. Desc. fasc. I. 30 = Zalusania angusta.

lyonnetioides, *Boiss. Fl. Orient.* iii. 286.—Syria.

macedonica, *Boiss. & Orph. Diagn.* Ser. II. vi. 97.—Macedon.

macrantha, *Heuff. in Flora*, xvi. (1833) 362.—Austria.

maritima, Urville, Enum. Pl. Orient. 114 = secundiramea.

maritima, *Linn. Sp. Pl.* 893.—Reg. Mediterr.

maritima, Pall. ex Bieb. Fl. Taur. Cauc. ii. 330 = ruthenica.

maritima, Sibth. & Sm. Fl. Graec. ix. 64 = tinctoria.

Marschalliana, Willd. Sp. Pl. iii. 2187 = Biebersteiniana.

Melampodia, Spreng. Syst. iii. 592 = seq.

melampodina, *Delile, Fl. Egypte*, 268. t. 45.—Aegypt.; Arab.

melanacme, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 315.—Mesopot.

melanolcena, Trautv. in Bull. Soc. Nat. Mosc. xli. (1868) 1. 461 = nigrescens.

membranacea, Labill. Ic. Pl. Syr. dec. iii. 14 = Achillea membranacea.

Micheliana, Bertol. Cat. Sem. Hort. Bonon. (1838); et Fl. Ital. ix. 264 = maritima.

Micheliana, Guss. Fl. Sic. Syn. ii. ii. 868, obs. = montana.

micrantha, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii 300.—Assyria.

microsperma, *Boiss. & Kotschy, Diagn.* Ser. II. v. 108.—Aegypt.; Arab.

millefolia, Lam. Sp. Pl. 896 = Chrysanthemum millefoliatum.

millefoliata, Pall. Reise Statthalt. Russ. Reich. i. 87, 139 = praec.

minuta, *Labm. ex Steud. Nom.* ed. II. i. 108.—Hab.?

mixta, ' H. S.' ex Nym. Consp. 361 = parnassia.

mixta, *Linn. Sp. Pl.* 894.—Reg. Mediterr.

monantha, Willd. Sp. Pl. iii. 2187 = tinctoria.

monilicostata, *Pomel, Nouv. Mat. Fl. Atl.* 41.—Algeria.

montana, Koch, Syn. Fl. Germ. ed. II. 415 = carpatica.

montana, *Ledeb. Fl. Ross.* ii. 522 = sterilis.

montana, *Linn. Sp. Pl.* ed. II. 1261.—Europ.; Orient.

ANTHEMIS :—

montana, Sibth. & Sm. Prod. Fl. Graec. ii. 190 = carpatica.

mucronulata, C. Koch, in Linnaea, xxiv. (1851) 819 = Chrysanthemum caucasicum.

mucronulata, Bertol. Amoen. 46 = Achillea Barrelieri.

mucronulata, Hort. ex Steud. Nom. ed. II. i. 108 = Anacyclus clavatus.

mucronulata, Reichb. Fl. Germ. Excurs. 227 = carpatica.

multicaulis, Boiss. & Heldr. in Boiss. Fl. Orient. iii. 301 = arvensis.

multicaulis, *Yanka, in Oestr. Bot. Wochenbl.* vi. (1856) 368.—Europ. austr.

muricata, *Guss. Fl. Sic. Syn.* ii. 490.—Sicil.

myagroides, Herb. Madr. ex DC. Prod. v. 551 = Guizotia oleifera.

Neilreichii, Ortm. in Flora, xxxv. (1852) 676 = rubrotinctoria.

nicaeensis, Willd. Sp. Pl. iii. 2182 = arvensis.

nigrescens, Vest, in Flora, iii. (1820) 8 = austriaca.

nigricans, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 265 = rigescens.

nobilis, *Linn. Sp. Pl.* 894.—Europ.

occidentalis, Willd. Sp. Pl. iii. 2185 = Spilanthes Mutisii.

odontostephana, *Boiss. Diagn.* Ser. I. vi. 85.—Persia; Ind. or.

odorata, Art. Hort. Kew. iii. 386 = Urvinia odorata.

odorata, Lam. Fl. Fr. ii. 168 = nobilis.

oppositifolia, Lam. Encyc. i. 576 = Spilanthes Mutisii.

ormenioides, Boiss. Diagn. Ser. I. iv. 5 = Wiedemannia.

ovatifolia, Orteg. Hort. Matr. 73 = Heliopsis buphthalmoides.

oxyloba, *Boiss. Fl. Orient.* iii. 289.—Cilicia.

oxyloba, Schur, Enum. Pl. Transs. 334, sphalm. = Achillea oxyloba.

parcelobola, Boiss. Fl. Orient. iii. 286 = Kotschyana.

palestina, *Reut. ex Boiss. l. c.* 288.—As. Min.; Syria.

pallescens, Heldr. ex Nym. Consp. iii. 359 = tinctoria.

pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. xi. 8.—As. Min.

parnassica, Nym. Syll. Suppl. 1 = tinctoria.

Parnassii, Boiss. & Heldr. ex Nym. Consp. 359 = tinctoria.

parnesia, *Boiss. & Heldr. in Boiss. Fl. Orient.* iii. 305. —Graecia.

parthenoides, *Benth. ex Hornem. Hort. Hafn.* ii. 833. —Hab.?

pauciloba, *Boiss. Diagn.* Ser. I. vi. 83.—Syria; Mesopot.

pectinata, *Boiss. & Reut. l. c.* Ser. I. iv. 6.—As. Min.

pedunculata, *Benth. & Hook. f. Gen.* ii. 422.—Af. bor.

pedunculata, Bory & Chaub. Fl. Pélop. 59 = Anacyclus clavatus.

pedunculata, Desf. Fl. Atl. ii. 288 = Anacyclus pedunculatus.

pentelica, Boiss. ex Nym. Consp. 360 = montana.

peregrina, Bieb. Fl. Taur. Cauc. ii. 331 = candidissima.

peregrina, *Linn. Syst. Nat.* ed. X. 1228.—Ital.; Graecia; As. Min.

persepolitana, *Boiss. Diagn.* Ser. I. xi. 7.—Persia.

persica, *Boiss. l. c.* Ser. I. vi. 85.—Persia.

Petasloxae, *Boiss. l. c.* Ser. I. xi. 19.—Lycia.

petraea, Tenore, Fl. Nap. ii. 246 = carpatica.

philistea, *Boiss. Fl. Orient.* iii. 302.—Palestin.

plaeiaulis, *Durieu, ex Munby, Fl. Algér.* 95, nomen.—Algeria.

platyglossa, C. Koch, in Linnaea, xxiv. (1851) 819 = rigescens.

plebeia, *Boiss. & Not, in Boiss. Fl. Orient.* iii. 318.—Kurdist.

pontiloelpis, Boiss. Diagn. Ser. II. iii. 24 = Kotschyana.

pontica, Urville, Enum. Pl. Orient. 114 = ruthenica.

pontica, Sibth. & Sm. Fl. Graec. ix. 66 = tricolor.

pontica, Willd. Sp. Pl. iii. 2184 = montana.

praecox, Link, ex Willd. l. c. 2183 = fuscata.

prolifera, Pers. Syn. ii. 467 = Cladanthus arabicus.

pseudo-astrata, Schur, Enum. Pl. Transs. 327 = tenuifolia.

pseudocota, Viviani, Fl. Dalm. ii. 73 = brachycentros.

pseudocotula, *Boiss. Diagn.* Ser. I. vi. 86.—Cyrena.

psorosperma, Tenore, Viag. Abruz. 136 = A. Cotula.

ANTHEMIS :—

ptarmicaefolia, Adam, in Weber & Mohr, Beitr. I. 71 = Achillea grandiflora.

ptarmiciformis, *C. Koch, in Linnaea*, xxiv. (1851) 819. —As. Min.

ptarygantha, C. Koch, l. c. xvii. (1843) 44 = candidissima.

pubescens, Willd. Sp. Pl. iii. 2177 = Anacyclus clavatus.

punctata, Guss. ex Nym. Consp. 360 = montana.

punctata, *Vahl, Symb. Bot.* ii. 91. t. 46.—Atlas; Sicilia.

purpurascens, Nym. Consp. 368 = Anacyclus radiatus.

pygmaea, Bertol. in Nov. Comm. Acad. Bonon. vi. (1848) 287 (Misc. Bot. ii. 23) = Matricaria praecox.

pygmaea, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 142. nomen.—Afr. trop.

pyrenaica, *Midgre. in Bull. Soc Bot. Fr.* xli. (1865) 897.—Mont. Pyren.

pyrenaica, Sch. Bip. in Oestr. Bot. Wochenbl. iv. (1854) 164 = austriaca.

pyrethriformis, Schur, Enum. Pl. Transs. 335 = carpatica.

Pyrethrum, Gouan, Fl. Monsp. 370 = montana.

Pyrethrum, Linn. Sp. Pl. 895 = Anacyclus Pyrethrum.

ramosa, Link, ex Spreng. Syst. iii. 592 = A. Cotula.

Raschеyana, *Boiss. Diagn.* Ser. I. xi. 8.—Syria.

reflectens, Reichb. f. Ic. Fl. Germ. xvi 61. t. 1006 = arvensis.

reflexa, Moench, Meth. 605 = altissima.

repanda, *Lamk. Sp. Pl.* 895.—Lusitan.

repens, Walt. Fl. Carol. 211 = Spilanthes repens.

Requienii, Sch. Bip. ex Nym. Consp. 361 = ruthenica.

retusa, Delile, Fl. Aeg. Illustr. 74 = cairica.

retusa, Link, ex Spreng. Syst. iii. 593 = ruthenica.

retusa, Tenore, ex Nym. Consp. 359 = brachycentros.

Reuteriana, Boiss. Diagn. Ser. I. iv. 8 = smyrnaea.

revoluta, Chr. Sm. ex Buch, in Berl. Acad. Abh. 1816-17 (1819) 370 = Gonospermum revolutum.

rhodensis, *Boiss. Diagn.* Ser. I. xi. 11.—Ins. Rhodus.

rigescens, *Willd. Hort. Berol.* i. t. 62.—Reg. Caucas.

rosea, Jacquem. Voy. Bot. iv. 88 = Allardia glabra.

rosea, *Sibth. & Sm Fl. Graec.* t. 887.—Cyprus.

rotata, *Boiss. Fl. Orient.* iii. 318. —Aegypt.; Arab.

Rudolphiana, Adam, in Weber & Mohr, Beitr. I. 72 = Biebersteiniana.

rupestris, Boj. ex Steud. Nom. ed. II. i. 108 = Epallage rupestris.

ruthenica, *Bieb. Fl. Taur. Cauc.* ii. 330. n. 1772.—Europ. austr. or. ; Reg. Caucas.

ruthenica, Sch. Bip. ex Schur, Enum. Pl. Transs. 332 = austriaca.

saboifolia, Boj. ex DC. Prod. v. 3 = Epallage salvifolia.

sanctolinoides, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 284.—Algeria.

sanatilis, Baumg ex Schur, Enum. Pl. Transs. 334 = Baumgarteniana.

sanatilis, Bieb. Fl. Taur. Cauc. iii. 551 = sterilis.

sanatilis, DC. ex Willd. Enum. Hort. Berol. 910 = montana.

scaposa, J. Gay, in Boiss. Fl. Orient. iii. 318 = Gayana.

scariosa, Boiss. Diag. Ser. I. xi. 7 = persepolitana.

scariosa, *DC. Prod.* vi. 4.—Syria; Mesopot.

scariosa, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 263.—Syria.

schizostephana, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 314.—Kurdist.

Schurii, Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 187 = Achillea oxyloba.

secundiramea, *Biv.ona, Sicul. Pl. Cent.* ii. 10.—Sicil.; Aegypt.

segetalis, Tenore, Fl. Neap. Syll. App. v. 54 = brachycentros.

seselifc, Salisb. Prod. 204 = Cladanthus arabicus.

Sibthorpii, *Griseb. Spicil. Fl. Rumel.* ii. 210.—Graecia.

sicula, Guss. Fl. Sic. Syn. ii. 866 = arvensis.

sinuata, Cerv. in La Llave et Lex. Nov. Veg. Desc. fasc. I. 30 = Zalusania triloba.

Sismondaana, *Clem. in Mem. Accad. Sc. Torino*, Ser. II. xvi. (1857) 297 (*Sert. Orient.* 61).—Byzant.

smyrnaea, *Boiss. Diagn.* Ser. I. iv. 7.—Smyrna.

sphacelata, Willd. in Bot. Zeit. v. (1847) 858 = Anacyclus clavatus.

ANTHEMIS :—
sphacelata, *Y. & C. Prael, Delic. Prag.* 108.—Sicilia.
Sprunneri, Boiss. & Heldr. Diagn. Ser. II. iii. 24 = montana.
sterilis, *Sieb. in Bull. Soc Nat. Mosc.* xxix. (1856) ii. 379.—Tauria.
stiparum, *Pomel, Nouv. Mat. Fl. Atl.* 48.—Afr. bor.
stipulacea, Moench, Meth. Suppl. 258 = Chrysanthemum sinense.
styriaca, Vest, in Syll. Ratisb. i. 12 = carpatica.
suaveolens, *Schmagr. ex Steud. Nom.* ed. II. i. 108.—Hab.!
subulicola, *Pomel, Nouv. Mat. Fl. Atl.* 49.—Afr. bor.
sudfruticosa, *Hornem. Hort. Hafn. Suppl.* 101.—Hab.!
sulphurea ×, Wall. ex Nym. Consp. 368 = A. Cotula.
Tatrae, *Kit. ex Schult. Fl. Austr.* ii. 534.—Europ.
taygetea, *Boiss. & Heldr. Diagn.* Ser. I. vi. 88.—Graecia.
tenella, *Host, Fl. Austr.* ii 507.—Europ.
tenuifolia, *Schur, in Verh Siebenb. Ver. Naturw.* ii. (1851) 171.—Transylvan.
tenuiloba, Boiss. Diagn. Ser. II iii. 25 = montana.
tenuiscta, *Ball, in Journ. Bot.* xi. (1873) 365.—Marocco.
tenuiscta, *Boiss. Fl. Orient.* iii. 201.—Armenia.
tenuiscta, *Pomel, Nouv. Mat. Fl. Atl.* 50.—Algeria.
tenuisphenca, *Herbich, Strp. Rar. Bucov.* 59.—Austria.
tigrensis, *Y. Gay, ex A. Rich. Tent. Fl. Abyss.* i. 418.—Abyss.
tinctoria, *Linn. Sp. Pl.* 896.—Europ. ; Orient.
tomentosa, Gouan, ex Boiss. Fl. Orient. iii. 321 = Anacyclus clavatus
tomentosa, *Linn. Sp Pl.* 893.—As. Min.
tomentosa, O. F. Muell. in Fl. Dan. xii. t. 685 = arvensis.
tomentosa, Linn. ex Boiss. Fl. Orient. iii. 291 = montana.
tomentosa, Willd. Sp. Pl iii. 2176 = peregrina.
tricolor, *Boiss. Fl. Orient.* iii. 288.—Cyprus.
triloba, Orteg. Hort. Matr. 79 = Zaluzania triloba
trinervia, *Raousch. Nom.* ed. III. 249.—Am. merid.
tripartita, *Boiss. & Blanch. Diagn.* Ser. II. iii. 22.—Syria.
Triumfetti, All. Fl. Pedem. i. 187 = austriaca.
Trotskiana, *Claus, in Beitr. Pflanz. Russ. Reich.* viii. 399.—Turkestan.
tuberiformis, Guss. Fl. Sic. Syn. ii. 870 = Anacyclus clavatus.
tuberculata, *Boiss. Elench.* 59. n. 107.—Hispan.; Afr. bor.
tubicina, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 319.—Mesopot.
ubensis, *Pomel, Nouv. Mat. Fl. Atl.* 289.—Afr. bor.
umbilicata, *Boiss. & Huet, Diagn.* Ser. II. iii. 24.—Armenia.
Urvillaeana, Sch. Bip. ex Nym Consp. ii. 362 = tomentosa.
Urvillaei, Nym. Syll. 7 = cretica.
valentina, Linn. Sp. Pl. 895 = Anacyclus radiatus.
viridis, Blanco, Fl. Filip. ed. II. 436 = Eclipta erecta.
Visianii, Weiss, ex Boiss. Fl. Orient. iii. 311 = chia.
vulgaris, Fl. Dan. ex Steud. Nom. ed. I. 58 = Matricaria inodora.
vulgaris, Linn. ex Steud. Nom. l. c. = Matricaria Chamomilla.
Wiedemanniana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1835) 27.—As. Min.

ANTHEPHORA, Schreb. Beschr. Graes. ii. 105. t. 44 (1779). *GRAMINEAE*, Benth. & Hook. f. ii. 1123.
ANTEPHORA, Steud. Syn. Fl. Gram. 111 (1855).
HYPUDARURUS, Reichb. Nom. 37 (1841).
abyssinica, A. Rich. Tent. Fl. Abyss. ii 389 = Hochstetteri.
axillifera, Steud. Syn. Fl. Gram. 111 = Buchloe dactyloides.
Betangeri, Steud. l. c. = Hilaria cenchroides.
cuspidata, *Anderss. Enum. Pl. Ins. Galap.* 141.—Ins. Galap.
elegans, Rupr. ex Steud. Syn. Fl. Gram. 111 = Hochstetteri.
elegans, *Schreb. Beschr. Graes.* ii. 105. t. 44.—Mexico.

ANTHEPHORA :—
Hochstetteri, *Nees, in Flora*, xxvii. (1844) 349.—Abyss.
Kotschyi, Hochst. ex Steud. Syn. Pl. Gram. 111 = Hochstetteri.
persica, *Boiss. Diagn.* Ser. I. xiii. 44.—Persia.
pubescens, Nees, Fl. Afr. Austr. i. 74 = Cenchrus pubescens.
punctulata, *Steud. Syn. Pl. Gram.* i. 111.—Mexico.
villosa, Spreng. Neue Entdeck. iii. 14 = elegans.

ANTHERICLIS, Rafin. in Am. Monthly Mag. (1819) 105 ; Journ. Phys. lxxxix. (1819) 261 = Tipularia, Nutt. (Orchid.).

ANTHERICUM, Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 788.
DILANTHES, Salisb. Gen. Pl. Fragm. 70 (1866).
HESPERANTHES, S. Wats. in Proc. Am. Acad. xiv. (1879) 241.
LILIAGO, Salisb. Gen. Pl. Fragm. 70 (1866).
PESSULARIA, Salisb. l. c. (1866).
PHALANGANTHUS, Schrank, ex Haw. Syn. Pl. Succ. 67 (1819).
PHALANGIUM, Jass. Gen. 52 (1789).
TRACHYANDRA, Kunth, Enum. Pl. iv. 573 (1843).
BLEPHANTHERA, ENDOGONA, LICINIA, OMITHALUM, TRACHINEMA, Rafin. Fl. Tellur. ii. 27, 59 ; iii. 57.
adenanthera, Forst. f. Prod. 24 = Dianella emifolia.
affine, *Baker, in Journ. Bot.* x. (1872) 138.—Afr. austr.
Aitoni, Baker, in Journ. Linn. Soc. xv. (1876) (= Bulbinella, Gen. Pl.) 394.—Afr. austr.
albucoides, Ait. Hort. Kew. i. 449 = Ornithogalum suaveolens.
aloifolium, Salisb. Prod. 250 = Bulbine aloides.
aloides, Linn. Sp. Pl. 311 = Bulbine aloides.
altissimum, Mill. Gard. Dict. ed. VIII. n. 8 = Bulbine longiscapa.
amoenum, Salisb. Prod. 250 = A. Liliago.
anceps, *Baker, in Journ. Linn. Soc.* xv. (1876) 305.—Afr. austr.
anceps, Soland. ex Baker, l. c. 368 = Tricoryne anceps.
andongense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 257.—Afr. trop.
angulicaule, *Baker, in Journ. Linn. Soc.* xv. (1876) 305.—Afr. austr.
angustifolium, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 332.—Abyss.
annuum, Linn. Sp. Pl. 311 = Bulbine annua.
annuum, Pourr. ex Willd. & Lange, Prod. Fl. Hisp. i. 308 = Asphodelus tenuifolius.
aphyllum, *Forsk. Fl. Aegypt. Arab.* 209.—Arabia.
arenarium, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 259.—Afr. trop.
asperatum, *Baker, in Journ. Bot.* x. (1872) 138.—Afr. austr.
asphodeloides, Ait. Hort. Kew. i. 450 = Bulbine longiscapa.
asphodeloides, Linn. Sp. Pl. 311 = Bulbine asphodeloides.
asphodelum, *Forsk. Fl. Aegypt. Arab.* 209.—Ins. Rhodus.
auratum, *Fisch. ex Trautv. in Act. Hort. Petrop.* ix. (1886) 281.—Soongaria.
autumnale, Scop. Fl. Carn. ed. II. i. 247 = Scilla autumnalis.
baeticum, Steud. Nom. ed. II. i. 108 = A. Liliago.
benguelense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 257.—Angola.
bicolor, Desf. Fl. Atl. i. 304 = Morsgagnia bicolor.
bifolium, Scop. Fl. Carn. ed. II. i. 247 = Scilla bifolia.
bipedunculatum, Jacq. Coll. Suppl. 88 = trifolium.
blepharophorum, Schult. f. Syst. vii. 461 = ciliatum.
boeticum, Boiss. Voy. Espagne, ii. 619. t. 172 = A. Liliago.
brasiliense, *Baker, in Journ. Linn. Soc.* xv. (1876) 306.—Bras.
Brehmeanum, *Baker, in Journ. Bot.* x. (1872) 139.—Afr. austr.
breviculc, *Baker, in Journ. Linn. Soc.* xv. (1876) 298.—Afr. austr.
brevifolium, *Thunb. Prod. Pl. Cap.* 68.—Afr. austr.
bulbosum, R. Br. Prod. 275 = Bulbine bulbosa.

ANTHERICUM :—
Burkei, Baker, in Journ. Bot. x. (1872) 140 = Bulbinella Burkei.
caeruleum, Ruiz & Pav. Fl. Per. iii. 67. t. 299 = Pasithea caerulea.
calyculatum, Georgi, It. i. 206 = Tofieldia cernua.
calyculatum, Linn. Sp. Pl. 311 = Tofieldia calyculata.
calyculatum, Pall. Reise, iii. 58 = Tofieldia palustris.
calyptrocarpum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 258.—Afr. trop.
Cameroni, *Baker, in Journ. Linn. Soc.* xv. (1876) 96, 314.—Afr. trop.
canaliculatum, *Ait. Hort. Kew.* i. 448.—Afr. austr.
capillare, Spreng. Syst. ii. 88 ; Schult. f. Syst. vii. 457 = Bulbinella capillaris.
carnosum, Baker, in Journ. Linn. Soc. xv. (1876) 296 (= Bulbinellae sp. Gen. Pl.).—Hab.!
Cauda-felis, Linn. f. Suppl. 202 = Bulbinella caudata.
caudatum, Thunb. Prod. Pl. Cap. 68 = Bulbinella caudata.
caulescens, *Baker, in Journ. Linn. Soc.* xv. (1876) 303.—Afr. trop.
cernuum, *Hornem. Hort. Hafn.* i. 382.—Hab.!
Chamaemoly, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 332 = Chlorophytum tetraphyllum.
ciliatum, *Linn. f. Suppl.* 202.—Afr. austr.
ciliatum, Schult. f. Syst. vii. 466 = Humboldtii.
ciliatum, *Spreng. l. c.* ii. 84.—Venezuela.
ciliolatum, Baker, in Journ. Bot. x. (1872) 137 = Bulbinella ciliolata.
cirrhatum, Forst. f. Prod. n. 148 = Arthropodium cirrhatum.
coarctatum, Ruiz & Pav. Fl. Per. iii. 67 = Excremis ramosa.
colubrinum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 256 = Dasystachys colubrina.
comosum, Thunb. Prod. Pl. Cap. 68 = Chlorophytum Sternbergianum.
contortum, *Linn. f. Suppl.* 202.—Afr. austr.
Cooperi, *Baker, in Journ. Linn. Soc.* xv. (1876) 304. —Natal.
corymbosum, *Baker, in Journ. Bot.* xv. (1877) 71.—Afr. trop.
costatum, *Andr. Bot. Rep.* t. 298.—Afr. austr.
crispum, Thunb. Prod. Pl. Cap. 68 = Chlorophytum crispum.
croceum, *Schult. f. Syst.* vii. 476.—Georgia.
dendroides, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 249 = Asphodeline prolifera.
distichum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 257.—Angola.
divaricatum, Jacq. Hort. Schoenb. iv. 7 = revolutum.
Dorattii, Hort. ex Baker, in Journ. Linn. Soc. xv. (1876) 301 = ramosum.
Dregeanum, *Baker, in Journ. Bot.* x. (1872) 139.—Afr. austr.
Drimiopsis, *Baker, in Journ. Linn. Soc.* xv. (1876) 301.—Afr. trop.
ecremorrhizum, *Ruiz & Pav. Fl. Per.* iii. 67.—Peruv.
echeandioides, *Baker, in Bot. Mag.* cxl. (1885) t. 6809. —Mexic.!
elatum, Ait. Hort. Kew. i. 448 = Chlorophytum elatum.
elongatum, Willd. Sp. Pl. 196 = Jacquinianum.
emarginatum, Banks & Soland. ex Hook. f. Fl. N. Zeal. i. 255 = Dianella intermedia.
emiforme, *Veil. Fl. Flum.* 148 ; ill. t. 162.—Bras.
eriostemon, Berger. Fl. Pyr. ii. 168 = Simethis bicolor.
esculentum, Spreng. Syst. ii. 84 = Camassia esculenta.
exwoisiatum, Jacq. Coll. Suppl. 89 = Urginea exwiata.
falcatum, *Linn. f. Suppl.* 202.—Afr. austr.
falcatum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 256 = Dasystachys falcata.
fasciculatum, *Baker, in Journ. Linn. Soc.* xv. (1876) 316.—Afr. austr.
fastigiatum, Stend. Nom. ed. I. 58 = Chlorophytum elatum.
fenzowum, Thunb. Prod. Pl. Cap. 68 = Bulbine favosa.
filifolium, Jacq. Coll. Suppl. 93 = Urginea filifolia.
filifolium, Thunb. Prod. Pl. Cap. 68 = Urginea nematodes.
filiferum, Ait. Hort. Kew. i. 451 = Bulbinella triquetra.

ANTHERICUM :—

filiforme, Thunb. Prod. Pl. Cap. 62 = Jacquinianum.

fimbriatum, Steud. Nom. ed. I. 35 = Dichopogon Sieberianus?

fimbriatum, Thunb. Prod. Pl. Cap. 63 = longifolium.

flagelliforme, Baker, in Journ. Bot. x. (1872) 140 = Schizobasis flagelliformis.

flavescens, *Schult. f. Syst.* vii. 1692.—Mexic.

flexifolium, Jacq. Coll. Suppl. 93 = Jacquinianum.

flexifolium, *Linn. f. Suppl.* 201.—Afr. austr.

floreoreum, R. Br. ex Benth. Fl. Austral. vii. 50 = Corynotheca dichotoma.

floribundum, Ait. Hort. Kew. i. 447 = Bulbinella latifolia.

fragrans, Jacq. Hort. Schoenb. i. 45 = Urginea fragrans.

frutescens, Linn. Sp. Pl. 310 = Bulbine frutescens.

fruticosum, Salisb. Prod. 250 = prasc.

fugax, Moris, Elench. i. 46 = Urginea fugax

Gerrardi, *Baker, in Journ. Bot.* x. (1872) 137.—Afr. austr.

glaucum, *Ruiz & Pav. Fl. Per.* iii. 68.—Peruv.

Gmelinianum, Schult. f. Syst. vii. 481 = Melanthium sibiricum.

gracile, Baker, in Journ. Bot. x. (1872) 137 = Bulbinella gracilis.

gracile, Salisb. Prod. 249 = ramosum.

gracillimum, *Regel, in Act. Hort. Petrop.* vi. (1879) 269.—Hab.?

graecum, Linn. Sp. Pl. ed. II. 444 = Lloydia graeca.

gramineum, *Vell. Fl. Flum.* 148 ; iii. t. 161.—Bras.

gramineum, *Baker, in Journ. Linn. Soc.* xv. (1876) 302.—Madag.

graminifolium, Willd. Sp. Pl. ii. 139 = undulatum.

grandiflorum, Salisb. Prod. 250 = Paradisea Liliastrum.

graminiforme, Dulac, Fl. Hautes-Pyr. 115 = Paradisea Liliastrum.

Grantii, *Baker, in Trans. Linn. Soc.* xxix. (1875) 160. t. 108 B.—Afr. trop.

graptophytum, *Baker, in Gard. Chron.* (1882) i. 460.—Ins. Socotra.

Heynei, Rottl. ex Baker, in Journ. Linn. Soc. xv. (1876) 328 = Chlorophytum Heyneanum.

hirsutum, *Thunb. Prod. Pl. Cap.* 63.—Afr. austr.

hispidum, *Linn. Sp. Pl.* ed. II. 446.—Afr. austr.

Hookeri, Colenso, ex Hook. f. Handb. New Zeal. Fl. 396 (= Bulbinella sp. Gen. Pl.).—N. Zel.

Humboldtii, *Hemsl. Bot. Centr. Am. Bot.* iii. 374.—Mexic.

humile, *Hockst. ex A. Rich. Tent. Fl. Abyss.* ii. 331.—Abyss.

hyacinthoides, Willd. ex Kunth, Enum. Pl. iv. 836 = Scilla indica.

inconspicuum, *Baker, in Journ. Bot.* xv. (1877) 71.—Afr. trop.

incurvum, Thunb. Prod. Pl. Cap. 62 = Bulbine rostrata.

intermedium, Willk. in Willk. & Lange, Prod. Fl. Hisp. i. 302 = A. Liliago.

intricatum, Baker, in Journ. Bot. x. (1872) 140 = Schizobasis intricata.

involucratum, *Baker, in Journ. Linn. Soc.* xv. (1876) 311.—Afr. austr.

Jacquinianum, *Schult. f. Syst.* vii. 462.—Afr. austr

japonicum, *Thunb. Fl. Jap.* 138.—China ; Japon.

lagopus, Thunb. Prod. Pl. Cap. 63 = Bulbine longiscapa.

latifolium, Linn. f. Suppl. 202 = Bulbinella latifolia.

latifolium, Schult. f. Syst. vii. 477 = simplex.

latifolium, Banks & Soland. ex Kunth, Enum. Pl. iv. 632 = Anthropodium chrysatum.

leptophyllum, *Baker, in Journ. Linn. Soc.* xv. (1876) 317.—Mexic.

Liliago, *Linn. Sp. Pl.* 310.—Europ. ; Afr. ber.

Liliastrum, Linn. l. c. ed. II. 446 = Paradisea Liliastrum.

limosum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 207.—Afr. trop.

longepaniculatum, *Steud. ex Schult. f. Syst.* vii. 457, 1692.—Afr. austr.

longifolium, A. Rich. Tent. Fl. Abyss. ii. 333 = Chlorophytum longifolium.

longifolium, *Jacq. Coll. Suppl.* 92.—Afr. austr.

longiflorum, *N. E. Br. in Gard. Chron.* (1882) ii. 297.—Zanz.

longiscapum, Jacq. Coll. Suppl. 84 = Bulbine longiscapa.

ANTHERICUM :—

longistylum, *Baker, in Journ. Linn. Soc.* xv. (1876) 305.—Afr. austr.

Macowani, *Baker, l. c.* 309.—Afr. austr.

macrophyllum, A. Rich. Tent. Fl. Abyss. ii. 334 = Chlorophytum macrophyllum.

Makoyanum, *Regel, Gartenfl.* (1880) 36.—Hab.?

marginatum, Thunb. Prod. Pl. Cap. 63 = Urginea marginata.

Matthaei, Vand. Fasc. Fl. 12 = Simethis bicolor.

microbyanthemoides, Baker, in Journ. Bot. x. (1872) 136 = Bulbine mesembryanthemoides.

millefloroum, [DC. in] Red. Lil. t. 58 = Anthropodium paniculatum.

minus, Steud. Nom. ed. I. 53 = Anthropodium minus.

molle, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 259.—Angola.

monophyllum, *Baker, in Journ. Bot.* xvi. (1878) 324.—Afr. trop.

muricatum, *Linn. f. Suppl.* 202.—Afr. austr.

nanum, *Baker, in Journ. Linn. Soc.* xv. (1876) 305.—Mexic.

nematodes, Schult. f. Syst. vii. 472 = Urginea nematodes.

nepalense, Spreng. Syst. iv. Cur. Post. 185 = Chlorophytum nepalense.

nidulans, *Baker, in Journ. Linn. Soc.* xv. (1876) 314.—Afr. trop.

Nimmonii, J. Grah. Cat. Pl. Bomb. 220 = Chlorophytum glaucum.

niveum, Spreng. Syst. ii. 84 ; Schult. f. in Roem. & Schult. Syst. vii. 480 = Chlorophytum tuberosum.

non-ramosum, Gillb. Exercit. ii. 465 = A. Liliago.

nubicum, *Baker, in Journ. Linn. Soc.* xv. (1876) 301.—Afr. trop.

Nuttallianum, *Schult. f. Syst* vii. 477.—Am. bor.

nutans, Jacq. Coll. Suppl. 86 = Bulbine nutans.

nutans, *Thunb. Prod. Pl. Cap.* 63.—Afr. austr.

Oatesii, *Baker, in Journ. Bot.* xvi. (1878) 324.—Afr. trop.

orchideum, *Weiw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 258.—Afr. trop.

ornithogaloides, Baker, in Journ. Linn. Soc. xv. (1876) 294 = Bulbinella ornithogaloides.

ornithogaloides, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 332 = Chlorophytum tuberosum.

ossifragum, Georgi, Beschreib. Russ. Reich. iii. iv. 907 = Tofieldia cernua.

ossifragum, Linn. Sp. Pl. 311 = Narthecium ossifragum.

palustre, Salisb. Prod. 250 = Narthecium ossifragum.

paniculatum, Andr. Bot. Rep. t. 395 = Arthropodium paniculatum.

parviflorum, *Schult. f. Syst.* vii. 459.—Afr. austr.

Parkeri, *Baker, in Journ. Bot.* xx. (1882) 269.—Madag.

parviflorum, Benth. Fl. Hongk. 373 = Chlorophytum laxum.

parvum, *N. E. Br. in Gard. Chron.* (1880) ii. 588.—Bras.

pauciflorum, Thunb. Prod. Pl. Cap. 63 = trifforum.

pendulum, Hornem. Hort. Hafn. i. 333 = Arthropodium paniculatum.

peronanthum, Baker, in Journ. Bot. x. (1872) 137 = Bulbinella peronata.

peruvianum, Willd. ex Kunth, Enum. Pl. iv. 596 = Hamboldtii.

physodes, Jacq. Coll. Suppl. 94 ; Ic. t. 418 = Urginea physodes.

pilosum, Jacq. l. c. 87 ; Ic. t. 416 = Ornithogalum hispidum.

planifolium, Linn. Mant. ii. 224 = Simethis bicolor.

pleiostachyum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 256 = Dasystachys pleiostachya.

plumosum, Ruiz & Pav. Fl. Per. iii. 68 = Bottionea thysanotoides.

pomeridianum, Ker-Gawl. in Bot. Reg. t. 564 = Chlorogalum pomeridianum.

praemorsum, Jacq. Coll. Suppl. 85 = Bulbine praemorsa.

Pseudo-asphodelus, Jacq. Enum. Vind. 59, 238 = Tofieldia calyculata.

pterocaulon, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 258.—Angola.

pubescens, *Baker, in Journ. Linn. Soc.* xv. (1876) 309.—Afr. austr.

ANTHERICUM :—

pubirhachis, *Baker, l. c.* 302.—Afr. trop.

pudicum, *Baker, l. c.* 308.—Afr. austr.

pugioniforme, Jacq Coll. Suppl. 83 = Bulbine pugioniformis.

pulchellum, *Baker, in Journ. Bot.* x. (1872) 140.—Afr. austr.

pusillum, Jacq. Coll. Suppl. 95 = Urginea pusilla.

pyrenicarpum, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 259.—Afr. trop.

Quamash, Stend. Nom. ed. I. 54 = Camassia esculenta.

ramosissimum, *Schult. f. Syst.* vii. 469.—Mexic.

ramosum, Steud. Nom. ed. I. 54 = Melaspharerula graminifolia.

ramosum, *Linn. Sp. Pl.* 310.—Europ.

reflexum, Cav. Ic. iii. 21 = Echeandia terniflora.

revolutum, *Linn. Sp. Pl.* 310.—Afr. austr.

rigidum, *Baker, in Journ. Bot.* x. (1872) 141.—Afr austr.

Rosii, *Hook. f. Handb. New Zeal. Fl.* 285 (= Bulbinellae sp. Gen. Pl.).—N. Zel.

rostratum, Jacq Coll. Suppl. 82 = Bulbine rostrata.

Saltii, *Baker, in Journ. Linn. Soc.* xv. (1876) 309.—Abyss.

sarcodum, Muell. ex Kunth, Enum. Pl. iv. 335 = Urginea fugax.

scabrellum, *Baker, in Journ. Linn. Soc.* xv. (1876) 318.—Mexic.

scabrum, *Linn. f. Suppl.* 202.—Afr. austr.

Schulteti, *Baker, in Journ. Bot.* x. (1872) 140.—Afr austr.

schlidiforum, *Eckl. ex Steud. Nom.* ed. II. i. 248, in syn. ; *et ex Baker, in Journ. Linn. Soc.* xv. (1876) 298.—Afr. austr.

schlioides, *Schult. f. Syst.* vii. 477.—Reg. Argent.

semibarbatum, Hook. Bot. Mag. t. 3129 = Bulbine bulbosa.

semibarbatum, R. Br. Prod. 275 = Bulbine semibarbata.

serotinum, Linn. Sp. Pl. ed. II. 444 = Lloydia alpina

sessiliflorum, Steud. Nom. ed. II. i. 104 = scilliforum.

setosum, Willd. ex Schult. f. Syst. vii. i. 478 = Bulbinella robusta.

silbricum, *Hort. Erf. ex Steud. Nom.* ed. II. i. 104.—Silar.

simplex, *Willd. ex Kunth, Enum. Pl.* iv. 596.—Reg. Colomb.

Skinneri, *Baker, in Journ. Linn. Soc.* xv. (1876) 318.—Am. trop.

spbaceatum, *Baker, l. c.* 303.—Afr. trop.

spirale, Linn. Mant. ii. 224 = Ketospermum spirale.

spiratum, Thunb. Prod. Pl. Cap. 62 = Urginea filifolia.

squamosum, Linn. f. Suppl. 202 = hispidum.

stenocarpum, *Baker, in Journ. Linn. Soc.* xv. (1876) 317.—Mexic.

Sternbergianum, Schult. f. Syst. vii. 1693 = Chlorophytum Sternbergianum.

strictum, Steud. ex Steud. Nom. ed. I. 54 = Dichopogon strictus.

subpetiolatum, *Baker, in Journ. Linn. Soc.* xv. (1876) 303.—Afr. trop.

subtriquetrum, Jacq. Coll. iii. 273 = Amianthium muscaetoxicum.

succulentum, Salisb. Prod. 250 = Bulbine asphodeloides.

sulphureum, Waldst. & Kit. Pl. Rar. Hung. i. 98 = Ornithogalum pyrenaicum.

superpositum, *Baker, in Journ. Bot.* xvi. (1878) 324.—Afr. trop.

tenellum, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 256.—Afr. trop.

thryoides, *Baker, in Journ. Bot.* x. (1872) 139.—Afr. austr.

Torreyi, *Baker, in Journ. Linn. Soc.* xv. (1877) 317.—Mexic.

trifforum, *Ait. Hort. Kew.* i. 448.—Afr. austr.

trinervium, Viviani, Fl. Lib. Spec. 21. t. 8. f. 2 = Lloydia alpina.

tripedale, *Baker, in Journ. Bot.* xx. (1882) 269.—Madag.

triquetrum, Linn. f. Suppl. 202 = Bulbinella triquetra.

tuberosum, Roxb. Fl. Corom. ii. 20. t. 186 = Chlorophytum tuberosum.

undulatum, *Jacq. Coll. Suppl.* 87.—Afr. austr.

undulatum, Thunb. Prod. Pl. Cap. 63 = hispidum.

uniflorum, *Roxb. Hort. Beng.* 37 ; *Fl. Ind.* ii. 149.—Ind. or.

ANTHERICUM :—
ustulatum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 258.—Afr. trop.
variegatum, Hort. ex Fl. Mag. N. S. t. 152 = Chlorophytum elatum.
vermiculare, Soland. ex Baker, in Journ. Linn. Soc. xv. (1876) 308 = Isogeneschaculatum.
vespertinum, Jacq. Hort. Schoenb. i. 44 = falcatum.
vestitum, *Baker, in Journ. Linn. Soc.* xv. (1876) 307. —Mexic.
villosum, Labill. Ic. Pl. Syr. Dec. v. 14. t. 9 = Gagea bohemica.
viscosum, *Baker, in Journ. Bot.* x. (1872) 141.—Afr. austr.
Williamsii, Hort. ex Gard. Chron. (1875) i. 490 = Chlorophytum elatum.
yedoënse, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* ii. 239.—Japon.
zanguebaricum, *Baker, in Journ. Linn. Soc.* xv. (1876) 302.—Afr. trop.
Zeyheri, *Baker, in Journ. Bot.* x (1872) 140.—Afr. austr.

ANTHEROCEPHALA, DC. Prod. vii. 755 (1839) = Andersonia, R. Br. (Epacrid.).
Drummondii, DC. l. c. = And. aristata.

ANTHEROCERAS, Bert. in Am. Journ. Sc. xix. (1831) 301 = Leucocoryne, Lindl. (Liliac.).
odorata, Bert. (Merc. Chil. 561) in Am. Journ. Sc. xix. (1831) 303.—Chili.
ornithogaloides, Bert. ll. cc.—Chili.

ANTHEROSPERMA, Forr. ex Stud. Nom. ed. II. i. 105 (1840), err. typ. = Atherosperma, Labill. (Monim.).

ANTHEROTOMA, Hook. f. in Benth. & Hook. f. Gen. i. 745 (1867). *MELASTOMACEAE*, Benth. & Hook. f. i. 746.
Abeilli, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 444.—Afr. trop.
Naudini, *Hook. f. in Benth. & Hook. f. l. c.*—Madag.

ANTHEROTRICHE, Turcz. in Bull. Soc. Nat. Mosc. (1846) ii. 506 = Anisopteris, Korth. (Dipterocarp.).
lanceolata, Turcz. l. c. 506 = Anisopt. thurifera.

ANTHERURA, Lour. Fl. Cochinch. 144 (1790). *RUBIACEAE*, Benth. & Hook. f. ii. 29.
rubra, *Lour. l. c.*—Cochinch.; Molucc.

ANTHERYLIUM, Rohr & Vahl, in Skr. Naturh. Selsk. Hafn. ii. (1792) i. 211. t. 8. *LYTHRARIEAE*, Benth. & Hook. f. i. 782.
floribundum, Spreng. Syst. ii. 474 = Adenaria floribunda.
grisleoides, Spreng. l. c. = Adenaria floribunda.
nudiflorum, *Hemsl. Diagn. Fl. Nov.* 18.—Mexic.
pauciflorum, Spreng. l. c. = Adenaria floribunda.
Rohrii, *Vahl, in Skr. Naturh. Selsk. Hafn.* ii. (1792) i. 211.—Ind. occ.

ANTHERYTA, Rafin. Sylva Tellur. 95 (1838) = Tetraxygia, Rich. (Melast.).
granulosa, Rafin. l. c. 96 = T. granulosa.

ANTHESTERIA, Spreng. Anleit. ii. 79 (1817) = Anthistiria, Linn. f. (Gramin.).

ANTHILLIS, Neck. Delic. Gallo-Belg. ii. 308 (1768) = Anthyllis, Linn. (Legum.).

ANTHIPSIMUS, Rafin. in Journ. Phys. lxxxix. (1819) 105 = Panicum, Linn. ! (Gramin.).
gonopodes, Rafin. l. c.—Am. bor.

ANTHIRRINUM, Moench, Meth. 521 (1794) = Antirrhinum, Tourn. (Scroph.).

ANTHISTIRIA, Linn. f. Nov. Gram. Gen. 35 (1779) (Amoen. Acad. x. 36). *GRAMINEAE*, Benth. & Hook. f. iii. 1136.
ANDROSCEPIA, Bronga. in Duperr. Voy. Coq. Bot. 77 (1826).
ARISTARIA, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 296.

ANTHISTIRIA :—
pilifera, Steud. Syn. Pl. Gram. 400 = arguens.
pilosa, Presl, Rel. Haenk. i. 346 = Cymbopogon Humboldtii.
polystachya, Roxb. Hort. Beng. 6; Fl. Ind. i. 248 = imberbis.
prostrata, *Wild. Sp. Pl.* iv. 901.—Ind. or.
pseudo-cymbaria, Steud. Syn. Pl. Gram. 399 = Cymbopogon elegans.
puberula, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 240 = imberbis.
punctata, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 448 = imberbis.
quinquuplex, Steud. Syn. Pl. Gram. 400 = Cymbopogon elegans.
ramosa, Thunb. ex Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 247 = Apluda gigantea.
reflexa, H. B. & K. Nov. Gen. et Sp. i. 191 = Cymbopogon Humboldtii.
scandens, Roxb. Hort. Beng. 6; Fl. Ind. i. 248 = ciliata.
setosa, Thunb. ex Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 238 = arguens.
subglabra, Buese, in Miq. Pl. Jungh. 368 = imberbis.
subsericea, Nees, ex Steud. Syn. Pl. Gram. 401 = gigantea.
syriaca, Boiss. Diagn. Ser. I. xiii. 72 = imberbis.
tortilis, Presl, Rel. Haenk. i. 347 = Andropogon Nardus.
tremula, *Nees, ex Steud. Syn. Pl. Gram.* 401.—Ind. or.
villosa, Lam. Illustr. t. 841. f. 3 = gigantea.
vulcanica, *Steud. Syn. Pl. Gram.* 400.—Ins. Santorin.
vulpina, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 245 = gigantea.
Wightii, Nees, ex Steud. Syn. Pl. Gram. 400 = prostrata.

ANTHOBOLUS, R. Br. Prod. 357 (1810). *SANTALACEAE*, Benth. & Hook. f. iii. 280.
exocarpoides, *F. Muell. Fragm.* ix. 8.—Austral.
filifolius, *R. Br. Prod.* 357.—Austral.
foveolatus, *F. Muell. Fragm.* i. 212.—Austral.
leptomeroides, *F. Muell. l. c.* 21.—Austral.
tripterus, A. Cunn. ex A. DC. in DC. Prod. xiv. 637 = filifolius.
triqueter, *R. Br. Prod.* 357.—Austral.

ANTHOCEPHALUS, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 237, excl. syn.; Korth. Verh. Nat. Gesch. 154. t. 48 (1840!). *RUBIACEAE*, Benth. & Hook. f. ii. 29.
CASSALIDIUM, A. Rich. l. c. 290.
Cadamba, Miq. Fl. Ind. Bat. ii. 185 = morindaefolius.
chinensis, Hassk. in Flora, xxvii. (1845) 281 = Cephalanthus orientalis.
chinensis, Walp. Rep. ii. 491 = Naucles purpurea.
indicus, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 285 = Naucles purpurea.
morindaefolius, *Korth. Verh. Nat. Gesch.* 154. t. 48; Obs. Nancl. Ind. 20.—Sumatr.; Ind. or.

ANTHOCERAS, Baker, in Journ. Linn. Soc. xi. 374 (1871) = ANTHEROCERAS, Bert. = Leucocoryne, Lindl. (Liliac.).

ANTHOCHAETES, A. Gray, in Hook. Kew Journ. iv. (1852) 225 = Toxanthus, Turcz. (Compos.).
Drummondii, A. Gray, l. c. = T. perpusillus.
Muelleri, Sond. in Linnaea, xxv. (1852) 486 = T. Muelleri.

ANTHOCERCIS, Labill. Nov. Holl. Pl. ii. 10. t. 104 (1806). *SOLANACEAE*, Benth. & Hook. f. ii. 913.
CYPHANTHERA, Miers, in Ann. Nat. Hist. Ser. II. xi. (1853) 376.
SARMIENTA, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 71.
albicans, *A. Cunn. in Field, New South Wales, App.* 385. t. 3.—Austral.
amblyanthe, F. Muell. Fragm. i. 170.—spuncoides.
angustifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 21.—Austral.
antennifera, *Endl. in Ann. Wien. Mus.* ii. (1836) 201.—Austral.

ANTHOCERCIS :—
arborea, *F. Muell. Fragm.* i. 219.—Austral.
Endesii, *F. Muell. l. c.* ii. 189.—Austral.
fasciculata, *F. Muell. l. c.* i. 120.—Austral.
genistoides, *Miers, Illustr. S. Am. Pl.* ii. *App.* 27. t. 83 D.—Austral.
globulifa, Miers, l. c. = viscosa.
gracilis, *Benth. in DC. Prod.* x. 192.—Austral.
Hopwoodii, *F. Muell. Fragm.* ii. 138.—Austral.
ilicifolia, Hook. Bot. Mag. sub t. 3961 = viscosa.
intricata, *F. Muell. Fragm.* i. 211.—Austral.
Leichhardtii, *F. Muell. l. c.* vi. 142.—Austral.
littorea, Endl. Iconogr. t. 68 = viscosa littorea, *Lehill. Pl. N. Holl.* ii. 19. t. 158.—Austral.
microphylla, *F. Muell. Fragm.* i. 179.—Austral.
myosotidea, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 20.—Austral.
Odgeril, *F. Muell. Fragm.* x. 19.—Austral.
racemosa, *F. Muell. l. c.* i. 211.—Austral.
scabrella, *Benth. in DC. Prod.* x. 192.—Austral.
spinescens, F. Muell. Fragm. i. 121 = genistoides.
tasmanica, *Hook. f. Fl. Tasm.* i. 289. t. 92.—Austral.
viscosa, *R. Br. Prod.* 448.—Austral.

ANTHOCEROS dissectus, Mont. ex Wedd. in DC. Prod. xvii. 69 (= *Neolacis dissecta*) = **Apinagia**, Tul. (Podost.).

ANTHOCHLAMYS, Fenzl, ex Endl. Gen. 300 (1837). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 88.
PELTISPERMUM, Moq. Chenop. Enum. 102 (1840). polygaloides, [*Fenzl*!] *Moq. l. c.* 102.—Reg. Cancas.

ANTHOCHORTUS, Endl. Gen. 121 (1837); Kunth, Enum. Pl. iii. 485 (1841) = ANTOCHORTUS, Nees = **Willdenowia**, Thunb. (Restiac.).

ANTHOCHYTRUM, Reichb. f. Ic. Fl. Germ. xix. 89. t. 1432 (1859) = **Crepis**, Vaill. (Compos.).
alpinum, Reichb. f. l. c. 89. t. 81 = C. alpina.

ANTHOCLEISTA, Afzel. ex R. Br. in Tuckey, Congo, App. 449 (1818). *LOGANIACEAE*, Benth. & Hook. f. ii. 795.
madagascariensis, *Baker, in Journ. Bot.* xx. (1882) 173.—Madag.
macrophylla, *G. Don, in Loud. Hort. Brit.* 471.—Afr. trop.
nobilis, *G. Don, Gen. Syst.* iv. 68.—Afr. trop.
scandens, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 16.—Afr. trop.
Vogelii, *Planch. in Hook. Ic. Pl.* t. 793.—Afr. trop.

ANTHOCOMA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie, ii. (1845) 569 = **Phlomis**, Linn.? (Labiat.).
flavescens, Zoll. & Mor. l. c. 570.—Malaya.

ANTHOCOMETES, Nees, in DC. Prod. xi. 311 (1847) = **Monotheium**, Hochst. (Acanth.).
aristatus, Nees, l. c. 313 = M. aristatum.

ANTHODENDRON, Reichb. in Moessl. Handb. i. p. xl. 244, 308 (1827) = **Rhododendron**, Linn. (Ericac.).
flavum, Reichb. l. c. 309 = R. flavum.
nudiflorum, Reichb. l. c. = R. nudiflorum.
ponticum, Reichb. Fl. Germ. Excurs. 416 = R. ponticum.
viscosum, Reichb. in Moessl. Handb. ed. II. i. 424 = R. viscosum.

ANTHODISCUS, Endl. Gen. 1091 in syn. (1841) = ANTHODON, Ruiz & Pav. = **Salacia**, Linn. (Hippocrat.).

ANTHODISCUS, G. F. W. Mey. Prim. Fl. Essequ. 198 (1818). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 183.
guianensis, *G. F. W. Mey. l. c.*—Guiana.
peruanus, *Baill. Adansonia*, x. (1872) 241.—Peruv.
trifoliatus, *G. F. W. Mey. Prim. Fl. Essequ.* (1818) 194.—Guiana.

ANTHODON, Ruiz & Pav. Fl. Per. i. 45. t. 74 (1798) = **Salacia**, Linn. (Hippocrat.).
aphloioides, Miers, ex Peyr. in Mart. Fl. Bras. xi. t. 127 = Hippocratea megacarpa.
crenatifolium, Mart. in Schult. Mant. i. 248 = S. crassifolia.

ANTHODON :—
decussatum, Ruiz & Pav. Fl. Per. i. 45. t. 74. f. 2 = Hippocratea decussata.
elliptica, Mart. in Schult. Mant. i. 248 = Salacia elliptica.
glomeratum, Mart. l. c. = S. glomerata.
grandiflorum, Benth. in Hook. Kew Journ. iv. (1852) 10 = S. grandiflora.
grandifolium, Mart. in Schult. Mant. i. 248 = S. grandifolia.
laevigatum, Steud. Nom. ed. II. i. 105 = S. laevigata.
laxiflorum, Benth. in Hook. Kew Journ. iv. (1852) 10 = S. laxiflora.
micranthum, Mart. in Schult. Mant. i. 248 = S. micrantha.
oblongifolium, Mart. l. c. = S. elliptica.
paniculatum, Mart. l. c. = S. paniculata.
Sellowianum, Schult. Mant. iii. Add. ii. 411.—Bras.
trinervoe, Schult. l. c.—Bras.
undulatum, Mart. in Schult. Mant. i. 248 = S. arbores.
verrucarum, Reichb. in Mem. Am. Acad. N. S. viii. (1861) 173.—Cuba.

ANTHODUS, Mart. ex Roem. & Schult. Syst. Mant. i. 253 (1822) = ANTHODON, Ruiz & Pav. = **Salacia**, Linn. (Hippocrat.).

ANTHOGONIUM, Wall. ex Lindl. Introd. Nat. Syst. ed. II. 341 (1836), nomen. *ORCHIDEAE*, Benth. & Hook. f. iii 515.
gracile, *Wall. Cat.* n. 7898.—Reg. Himal.
Griffithii, Reichb. f. in Bonplandia, ii. (1854) 90 = gracile.

ANTHOLOMA, Labill. Voy. Rech. Pérouse, ii. 235. t. 41 (1800). *TILIACEAE*, Benth. & Hook. f. i. 239.
Billardieri, *Virill. in Bull. Soc. Linn. Normand.* ix. (1865) 335.—N. Caled.
montana, *Labill. Voy. Rech. Pérouse*, 236. t. 41.—N. Caled.

ANTHOLYZA, Linn. Syst. ed. I. (1735); Gen ed. I. 10 (1737). *IRIDEAE*, Benth. & Hook. f. iii. 710.
ANISANTHUS, Sweet, Hort. Brit. ed. II. 500 (1830); Brit. Fl. Gard. Ser. II. t. 84 (1836).
CUNONIA, Mill. Gard. Dict. ed. VIII. (1768).
GLADIOLUS, Geertn. Fruct. i. 31. t. 11. f. 4. (1788).
HOMOGLOSSUM, Salisb. in Trans. Hort. Soc. i. (1812) 325.
PETAMENES, Salisb. l. c. 324.
abbreviata, Pers. Syn. i. 42 = quadrangularis.
abyssinica, *A. Brongn. in A. Sich. Tent. Fl. Abyss.* ii. 208.—Abyss.
aethiopica, *Linn. Syst.* ed. X. 868.—Afr. austr.
aletroides, Burm. f. Fl. Cap. Prod. 1 = Watsonia aletroides.
bicolor, Gasp. ex Tenore, Cat. Orto Nap. (1845) 78; fide E. Morr. in Belg. Hortic. ii. (1852) 145 = aethiopica.
brevifolia, Hort. ex Steud. Nom. ed. II. i. 105 = Gladiolus brevifolius.
brevifolia, Eckl. ex Ball, in Journ. Linn. Soc. xvi. (1878) 179 = caffra.
caffra, *Bunks, ex Ker-Gawl. in Konig & Sims, Ann. Bot.* i. (1805) 282.—Afr. austr.
caryophyllacea, Burm. f. Fl. Cap. Prod. 1 = Watsonia humilis.
caryophyllea, Vahl, Enum. Pl. ii. 129 = Watsonia humilis.
cepacea, *Linn. Mant.* ii. 176.—Afr. austr.
coccinea, Spreng. Neue Entdeck. i. 253 = A. Cunonia.
Cunonia, *Linn. Sp. Pl.* 87.—Afr. austr.
flammea, Klatt, in Linnaea, xxxv. (1867–68) 299 = Babiana fimbriata.
floribunda, Salisb. in Trans. Hort. Soc. i. (1812) 324 = aethiopica.
fucata, *Baker, in Journ. Linn. Soc. xvi.* (1878) 180.—Afr. austr.
fulgens, Andr. Bot. Rep. t. 122 = Watsonia Meriana.
hirsuta, Klatt, in Linnaea, xxxv. (1867–68) 279 = Gladiolus scriceo-villosus.
hirsuta, Lam. Encyc. ii. 211 = Gladiolus mollis.
hirsuta, Pers. Syn. i. 42 = Babiana Thunbergii.
huillensis, *Welw. ex Baker, in Trans. Linn. Soc. Ser. II.* i. (1878) 270.—Afr. trop.
hypogaea, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 248.—Afr. austr.

ANTHOLYZA :—
intermedia, *Baker, in Journ. Linn. Soc.* xvi. (18) 180, nomen.—Afr. austr.
lucidior, *Linn. f. Suppl.* 96.—Afr. austr.
lucidior, Vahl, Enum. Pl. ii. 122 = Watsonia plantaginea.
Ludwigii, Eckl. Verz. 84 = aethiopica!
marginata, Pagu, ex Steud. Nom. ed. II. i. 106 = Watsonia marginata.
Maura, Linn. Mant. ii. 175 = Witsenia Maura.
Meriana, Blanco, Fl. Filip. ed. I. 34 = Sisyrinchium palmifolium.
Meriana, Linn. Sp. Pl. ed. II. 54 = Watsonia Meriana.
Merianella, Curt. Bot. Mag. t. 441 = Watsonia aletroides.
Merianella, *Murr. Syst.* ed. XIII. 77.—Afr. austr.
Meriania, Linn. Syst. ed. X. 865 = Watsonia Meriana.
montana, [Ker-Gawl. in] Konig & Sims, Ann. Bot. i. (1805) 283 = Gladiolus montanus.
nemorosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1869) 246.—Afr. austr.
nervosa, *Thunb. Prod. Pl. Cap.* 7.—Afr. austr.
orchidiflora, Klatt, in Linnaea, xxxii. (1863) 733 = Gladiolus arenarius.
paniculata, *Klatt, in Linnaea*, xxxv. (1867–68) 279.—Afr. austr.
plicata, Linn. f. Suppl. 96 = Babiana Thunbergii.
praealta, [Del. in] Red. Liliac. vii. t. 387 = aethiopica.
quadrangularis, *Burm. f. Fl. Cap. Prod.* 1.—Afr. austr.
quinquenervia, *Schrank, Comm. Acad. Gott.* 56.—Afr. austr.
ramosa, Klatt, in Linnaea, xxxii. (1863) 734 = Gladiolus montanus.
revoluta, *Burm., f. Fl. Cap. Prod.* 1.—Afr. austr.
ringens, Andr. Bot. Rep. t. 32 = aethiopica.
ringens, Linn. Sp. Pl. 87 = Babiana ringens.
rupestris, Eckl. ex Baker, in Journ. Linn. Soc. xvi. (1877) 179 = caffra.
saccata, *Baker, in Journ. Linn. Soc.* xvi. (1877) 180.—Afr. austr.
spicata, Andr. Bot. Rep. t. 56 = Watsonia brevifolia.
spicata, *Brehm, ex Klatt, in Abh. Naturf. Ges. Halle*, xii. (1869) 245.—Afr. austr.
spicata, Mill. Gard. Dict. ed. VIII. n. 2 = Gladiolus Milleri.
splendens, *Steud. Nom.* ed. II. i. 106.—Afr. austr.
striata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1869) 246.—Afr. austr.
tubulosa, Andr. Bot. Rep. t. 174 = Watsonia aletroides.
vittigera, Sallsb. in Trans. Hort. Soc. i. (1812) 324 = aethiopica.

ANTHOMELES, M. Roem. Syn. (Rosiflorae) 102 & 140 (1847) = **Crataegus**, Linn. (Rosac.).
aestivalis, M. Roem. l. c. 141 = Cr. aestivalis.
Douglasii, M. Roem. l. c. 140 = Cr. Douglasii.
flava, M. Roem. l. c. 142 = Cr. flava.
flexispina, M. Roem. l. c. = Cr. parvifolia.
glandulosa, M. Roem. l. c. 141 = Cr. flava.
opaca, M. Roem. l. c. = Cr. aestivalis.
rotundifolia, M. Roem. l. c. 140 = Cr. coccinea.
sanguinea, M. Roem. l. c. = Cr. sanguinea.
stipulosa, M. Roem. l. c. 143 = Cr. stipulosa.
subserrata, M. Roem. l. c. = Cr. subserrata.
turbinata, M. Roem. l. c. = Cr. flava.

ANTHONOTA, Beauv. Fl. Owar. et Ben. i. 70. t. 42 (1805) = **Macrolobium**, Schreb. (Legum.).
crassifolia, Baill. Adansonia, vi. (1866–67) 179.—Afr. trop.
elliptica, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 430. —Am. merid.
macrophylla, Beauv. Fl. Owar. et Ben. i. 70. t. 42 = M. Palisoti.

ANTHOPHYLLUM, Stend. Syn. Pl. Cyp. 160 (1855) = **Scirpus**, Linn. (Cyper.).
Urvillei, Steud. l. c. = S. frondosus.

ANTHOPOGON, Neck. Elem. ii. 13 (1790) = **Gentiana**, Linn.
barbata, Rafin. Fl. Tellur. iii. 85 = G. barbata.
ciliata, Rafin. l. c. = G. ciliata.
crinita, Rafin. l. c. = G. crinita.
detonsa, Rafin. l. c. = G. serrata.
virgata, Rafin. l. c.—Am. bor.

ANTHOPOGON, Nutt. Gen. Am. i. 81 (1818) = **Gymnopogon**, Beauv. (Gram.).
ashcrvus, Rafin. New Fl. Am. iv. 91.—Am. bor.

ANTHOPOGON :—
brevifolium, Rafin. New Fl. Am. iv. 91 = Gymnopogon brevifolius.
incarnatum, Rafin. l. c. 90.—Am. bor.
lepturoides, Nutt. Gen. Am. I. 83 = G. racemosus.

ANTHOPTERUS, Hook. Ic. Pl. t. 948 (1840). *VACCINIACEAE*, Benth. & Hook. f. ii. 566.
calycinus, Spruce, ex Benth. & Hook. f. Gen. ii. 570 = Ceratostema calycinum.
longiflorus, Benth. ex Benth. & Hook. f. l. c. = Ceratostema longiflorum, *Benth. & Hook. f.*
racemosus, *Benth. Pl. Hartw.* 221.—Reg. Colomb.
racemosus, *Hook. Ic. Pl.* iii. t. 243.—Peruv.
Wardii, *Ball, in Hook. Ic. Pl.* t. 1465.—Reg. Colomb.

ANTHORA, Hall. in Rupp. Fl. Jen. 290 (1745); DC. Prod. i. 56 = Aconitum, Tourn. (Ranunc.).
saxatilis, Fourr. in Ann. Soc. Linn. Lyon, N. S xvi. (1868) 326 = Ac. Anthora.
versicolor, Stev. ex Ser in DC. Prod. i. 57 = Ac. Anthora.

ANTHOSACHNE, Steud. Syn. Pl. Gram. 287 (1855) = Agropyron, J. Gaertn. (Gramin.).
australasica, Steud. l. c. = Agrop. scabrum.

ANTOSCHMIDTIA, Steud. Syn. Pl. Gram. 199 (1855) = Schmidtia, Steud. (Gramin.).

ANTHOSCIADIUM, Fenzl, in Endl. Gen. Suppl. iv. 111. 9 (1850) = Selinum, Linn. (Umbell.).

ANTHOSPERMUM, Linn. Gen. ed. I. 202 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 140.
CHRYSOSPERMUM, Reichb. Consp. 166 (1828) †
aethiopicum, *Linn. Sp. Pl.* 1003.—Afr. austr.
ambrosiacum, Moench, Meth. Suppl. 269 = aethiopicum.
aromaticum, Salisb. Prod. 59 = aethiopicum.
asperuloides, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 11.—Fernando Po.
Bergianum, *Cruse, Rub. Cap.* 8.—Afr. austr.
Burkei, *Sond. in Harv. & Sond. Fl. Cap.* iii. 29.—Afr. austr.
calycophyllum, *Sond. l. c.* 31.—Afr. austr.
ciliare, *Linn. Sp. Pl.* ed. II. 1521.—Afr. austr.
ciliare, Thunb. ex Sond. in Harv. & Sond. Fl. Cap. iii. 30 = hirtum.
confertum, *Cruse, in Walp. Rep.* ii. 402.—Afr. austr.
Crocyllis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 32.—Afr. austr.
Dregei, *Sond. l. c.* 29.—Afr. austr.
Ecklonis, *Sond. l. c.* 33.—Afr. austr.
emirnense, *Baker, in Journ. Bot.* xx. (1882) 139.—Madag.
ferruginum, *Eckl. & Zeyh. Enum.* 266.—Afr. austr.
fostidum, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. iii. 30 = lanceolatum.
galioides, Reichb. ex Spreng. Syst. iv. Cur. Post. 338 = ciliare.
Galopina, Thunb. Prod. Pl. Cap. 32 = Galopina circaeoides.
hedyotideum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 20.—Afr. austr.
herbaceum, Linn. f. Suppl. 440 = lanceolatum.
hirsutum, DC. Prod. vi. 580 = hirtum.
hirtum, *Cruse, Rub. Cap.* 11.—Afr. austr.
hispidulum, *E. Mey. ex Harv. & Sond. Fl. Cap.* iii. 29.—Afr. austr.
lanceolatum, Sieber, ex Harv. & Sond. l. c. 30 = hirtum.
lanceolatum, *Thunb. Prod. Fl. Cap.* 32.—Afr. austr.
latifolium, E. Mey. ex Harv. & Sond. l. c. 30 = lanceolatum.
Lichtensteinii, *Cruse, Rub. Cap.* 9, 15.—Afr. austr.
muriculatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 345.—Abyss.
nudosum, E. Mey. ex Harv. & Sond. Fl. Cap. iii. 30 = lanceolatum.
pachyrhizum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 229.—Afr. trop.
pauciolatum, *Cruse, Rub. Cap.* 9, 15.—Afr. austr.
plicatum, Hilsenb. & Boj. ex Baker, in Journ. Linn. Soc. xx. (1884) 143 = Myosurandra maculata.
polyacanthum, *Baker, l. c.* 171.—Madag.

ANTHOSPERMUM :—
prostratum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 28.—Afr. austr.
pumilum, *Sond. l. c.* 31.—Afr. austr.
rigidum, *Eckl. & Zeyh. Enum.* 267.—Afr. austr.
rubicaeum, Reichb. ex Spreng. Syst. iv. Cur. Post. 338 = hirtum.
scabrum, Thunb. Prod. Pl. Cap. 32 = Carpacoce scabra.
spathulatum, Spreng. Neue Entdeck. iii. 45 = ciliare.
spermacoceum, Reichb. ex Spreng. Syst. iv. Cur. Post. 338 = Carpacoce Spermacoce.
thymoides, *Baker, in Journ. Linn. Soc.* xx. (1884) 171.—Madag.
tricostatum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 28.—Afr. austr.

ANTHOSTEMA, A. Juss. Euph. Tent. 56. t. 18 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 261.
madagascariense, *Baill. Étud. Gén. Euph.* 60, 544.—Madag.
senegalensis, *Juss. Euph. Tent.* 117.—Afr. trop.

ANTHOTIUM, R. Br. Prod. 582 (1810). *GOODENOVIEAE*, Benth. & Hook. f. ii. 587.
glabrum, Poir. in Dict. Sc. Nat. ii. Suppl. 80 = humile.
humile, *R. Br. Prod.* 582.—Austral.
rubriflorum, *F. Muell. ex Benth. Fl. Austral.* iv. 45.—Austral.

ANTHOTROCHE, Endl. Nov. Stirp. Dec. 6 (1839). *SOLANACEAE*, Benth. & Hook. f. ii. 912.
Blackii, *F. Muell. Fragm.* viii. 232.—Austral.
pannosa, *Endl. Nov. Stirp. Dec.* 7.—Austral.
Walcottii, *F. Muell. Fragm.* i. 128.—Austral.

ANTHOXANTHUM, Linn. Gen. ed. I. 18 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1188.
XANTHANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. 188 (1881).
XANTHOGRANTHOS, St. Lag. l. c. vii. (1880) 119; viii. 189 (1881).
aculeatum, Lian. f. Suppl. 89 = Crypsis aculeata.
alpinum, Schur, Enum. Pl. Transs. 735 = odoratum.
amarum, *Brot. Phyt. Lusit.* 11. t. 4. f. 1.—Lusitan.
angustifolium, Plan.-Gir. Ensaya Fl. Gallega, 398 = Puelii.
aristatum, *Boiss. Voy. Espagne*, 688.—Hispan.
aristatum, Bor. Fl. Centr. Fr. ed. II. 576 = Puelii.
asperum, Mann, ex Opiz, Natural. vii. (1824) 51 = odoratum.
avenaceum, Pourr. ex Lange, in Kjoeb. Vidensk. Meddel. (1860) 26 = amarum.
avenaceum, *Rets. Obs.* ii. 5.—Ind. or.
Carremanum, Parl. Pl. Nov. 87 = aristatum.
crinitum, Lion. f. Suppl. 90 = Dichelachne crinita.
gigantteum, Walt. Fl. Carol. 65 = Erianthus saccharoides.
glaucum, Vahl, ex Roem. & Schult. Syst. i. 288 = amarum.
gracile, *Bivon. Stirp. Rar. Sicil.* i. [13] t. 1. f. 2.—Reg. Mediterr.
indicum, Linn. Sp. Pl. 28 = Perotis latifolia.
lastianum, *Salem. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 709.—Marocco.
Lloydii, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 697 = aristatum.
myrtheuse, Tod. ex Nym. Consp. 790 = Puelii.
odoratum, *Linn. Sp. Pl.* 28.—Europ.; As. bor.; Afr. bor.
ovatum, *Lag. Elench.* 2. n. 15.—Hispan.; Marocco.
paniculatum, Linn. Sp. Pl. 28 = Festuca spadicea.
Puelii, *Lecoq & Lamotte, Cat. Pl. Auver.* 388.—Europ.
pulchellum, D. Dietr. Syn. Pl. i. 625, err. typ. 69 = pulchrovirinum.
pulcherrimum, *Lour. Fl. Cochinch.* 86.—Cochinch.
Sommieri-ianum, Ricci, in Nuovo Giorn. Bot. xiii. (1881) 135 = aristatum.
spiculaeforme, Asch. ex Steud. Nom. ed. II. i. 106 = Perotis latifolia.
villosum, Dum. Obs. Gram. Belg. 129 = odoratum.

ANTHRISCUS, Bernh. Syst. Verz. Erf. 112 (1800). *UMBELLIFERAE*, Benth. & Hook. f. i. 899.

ANTHRISCUS :—
aboriva, Jord. Obs. Pl. Crit. vii. 28 = alpestris.
africana, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 196.—Afr. occ.
alpestris, *Wimm. & Grab. Fl. Silesiae*, i. 289.—Europ.
alpina, Jord. Obs. Pl. Crit. vii. 29 = sylvestris.
anatolica, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 59 = nemorosa.
capensis, Spreng. in Roem & Schult. Syst. vi. 526 = Cerum hispidum.
Cascalia, Bieb. Fl. Taur. Cauc. i. 230 = vulgaris.
Cerefolium, *Hoffm. Gen. Umb.* 41, 47. t. 1.—Europ.; Oriens; As. bor.
chaerophyllus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Cerefolium.
Cicutaria, Duby, in DC. Bot. Gall. i. 239 = alpestris.
dubia, Kabath, Fl. Gleiw. 86 = alpestris.
elatior, Bess. Enum. Pl. Volh. 58 = sylvestris.
fumarioides, *Spreng. Umb. Prod.* 27.—Europ.
gracilis, Hampe, in Flora, xxv. (1842) i. 64 = tenerrima.
hetrosantha, Schur, Enum. Pl. Transs. 278 = alpestris.
humilis, Bess. Enum. Pl. Volh. 13 = alpestris.
intermedia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 31 = sylvestris.
Kotschyi, *Fenzl, ex Boiss. Diagn.* Ser. II. v. 102.—Orient.
laevigata, Griseb. Spicil. Fl. Rumel. i. 367 = sylvestris.
lasiopocarpa, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 59.—Syria. Cilic.
longirostris, Bertol. Fl. Ital. iii. 197 = A. Cerefolium.
macrocarpa, Boiss. & Heldr. Diagn. Ser. II. ii. 100 = nemorosa.
mollis, *Boiss. & Reut. i. c.* 99.—Afr. bor.
neglecta, Boiss. & Kent. ex Lange, Ind. Sem. Hann. (1854) 22; Losc. & Pard. Ser. Inconf. Arag. 46 = vulgaris.
nemorosa, Baker & S. Moore, in Journ. Linn. Soc. xvii. (1879) 285 = sylvestris.
nemorosa, *Spreng. Umb. Prod.* 27.—Europ.; Oriens.
nitens, *Bernh. in Linnaea*, xii. (1838) *Litt.* 76 = Ruthenia.
nitida, *Garcke, Fl. Deutschl.* ed. VII. 180.—Europ.
nodosa, Pers. Syn. i. 320 = Chaerophyllum nodosum.
odoratissima, *Gueldenst. Reisen durch Russl.* i. 190.—Rossia.
pilosa, Schur, Enum. Pl. Transs. 272 = sylvestris.
Prescottii, *Vesersem. in Beitr. Pfl. Russ. Reich.* ix. 84.—Rossia.
procera, Bess. Enum. Pl. Volh. 13 = sylvestris.
rivularis, Dolliner, ex Vukot. in Rad. Jugos. Akad Zagrab, xxxix. (1877) 200 = nemorosa.
rupicola, Godet, Fl. Jura, i. 300 = alpestris.
Ruprechti, *Boiss. Fl. Orient.* ii. 914.—Reg. Caucas.
sativa, Bess. Enum. Pl. Volh. 13 = A. Cerefolium.
scandens, Stend. Nom. ed. II. i. 107, sphalm. = scq *Scandix*; Bieb. Fl. Taur. Cauc. i. 231 = Chaerophyllum nodosum.
silesia, *DC. Prod.* iv. 228.—Sicil.
silvestris, Vis. ex Nym. Consp. 301 = fumarioides.
sylvestris, *Hoffm. Gen. Umb.* 40, 46. t. 1. f. 19.—Europ.; Reg. Caucas.; As. bor.
taurica, *Fisch. ex Lond. Hort. Brit.* 101.—Tauria.
tenerrima, *Boiss. & Sprun. in Ann. Sc. Nat.* Sér. III. ii. (1869) 60.—Graecia; As. Min.
torquata, Duby, in DC. Bot. Gall. i. 239 = alpestris.
trachysperma, Reichb. ex Nym. Consp. 301 = A. Cerefolium.
trichosperma, Fers. Syn. i. 320 = Caucalis trichosperma.
trichosperma, Spreng. in Roem. & Schult. Syst. vi. 525 = A. Cerefolium.
tuberosa, *Spreng. in Roem. & Schult. Syst.* vi. 527.—Hab.?
villosa, Boiss. Fl. Orient. ii. 911 = nemorosa.
vulgaris, *Bernh. Syst. Verz. Erf.* 108.—Europ.; Oriens.

ANTHROCEPHALUS, [Schlecht. in] Linnaea, xiv. (1840) *Litt.* 179, sphalm. = Anthocephalus, A. Rich. (Rubiac.).

ANTHROPODIUM, Steud. in Bot. Mag. Ind. (1818) 5, err. typ. = Arthropodium, R. Br. (Liliac.).

ANTHROSTYLIS, D. Dietr. Syn. Pl. i. 182, 927 (1839), err. typ. = Arthrostylis, R. Br. (Cyper.).

ANTHURIUM, Schott, in Wien. Zeitschr. f. Kunst, iii. 838 (1829). *AROIDEÆ*, Benth. & Hook. f. iii. 998.
PLEUROSPA, Rafin. Fl. Tellur. iv. 8 (1836).
STERESPANTHERA, Rafin. l. c. 13 (1836).

acaule, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 83. t. 294.—Peruv.

acaule, *Schott, Melet.* i. 23.—Ind. occ.

acuminatum, Schott, in Oestr. Bot. Wochenbl. v. (1855) 66 = gracile.

aduncum, *Schott, Prod. Aroid.* 478.—Bras.

aemulum, *Schott, in Bonplandia*, vii. (1859) 165.—Mexic.

seranthe, *Hort. ex Baker, in Saund. Ref.* t. 278.—Am. trop.

affine, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 82. —Bras.

agnatum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181. —Nicaragua.

albo-costatum, *Miq. Hort. Amstelod.* (1858).—Hab.?

alienatum, *Schott, Prod. Aroid.* 507.—Peruv.

alienigenum, Schott, ex Engl. in DC. Monog. Phan. ii. 130 = Olfersianum.

amoenum, *Kunth, Ind. Sem. Hort. Berol.* (1848) 11.—Peruv.

amplum, Kunth, Enum. Pl. iii. 77 = Hookeri.

andegavense ×, Hort. cf. Gard. Chron. (1883) t. 510.

Andersonii, *Schott, in Oestr. Bot. Wochenbl.* (1857) 295.—Ind. occ.

andicola, *Liebm. in Kjoeb. Vidensk. Meddel.* (1849) 22. —Mexic.

Andraeanum, *Linden, Illustr. Hortic.* (1877) 43. t. 271. —Reg. Colomb.

angustatum, *Kunth, Enum. Pl.* iii. 79.—Venezuela.

Appunianum, *Schott, Prod. Aroid.* 512.—Venezuela.

arisaefolium, *Regel, in Ind. Sem. Hort. Petrop.* (1869) 10; *Gartenfl.* (1870) 98. t. 648.—' N. Caled.'

assimile, Schott, in Oestr. Bot. Wochenbl. (1855) 82 = Harrisii.

attenuatum, Miq. Cat. Hort. Amstel. (1852) = acaule.

Aublettii, Kunth, Enum. Pl. iii. 81 = pentaphyllum.

augustinum, *C. Koch & Lauche, Ind. Sem. Hort. Berol.* (1855) 7 = trinervium.

bahiense, N. E. Br. ex Engl. in DC. Monog. Phan. ii. 638 = bellum.

Bakeri, *Hook. f. Bot. Mag.* t. 6261.—Am. centr.

Belangeri, Engl. in Engl. Bot. Jahrb. i. (1881) 480 = gracile.

bellum, *Schott, in Oestr. Bot. Zeitschr.* (1859) 100.—Bras.

Beyrichianum, Kunth, Enum. Pl. iii. 69 = Harrisii.

Binoti, Linden, in Regel, Gartenfl. (1872) t. 723 = bellum.

bogotense, *Schott, in Oestr. Bot. Wochenbl.* (1857) 302.—Reg. Colomb.

bombacifolium, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 182.—Costa Rica.

Bouchenum, *C. Koch, Ind. Sem Hort. Berol.* (1855) *App.* 7.—Venezuela.

brachycostatum, *Schott, Prod. Aroid.* 468.—Mexic.

brachypathum, C. Koch & Bouché, in Koch, Allg. Gartenz. (1857) 238 = trinerve.

Bredemeyeri, *Schott, in Oestr. Bot. Wochenbl.* vii.(1857) 269.—Bras.

brevicaspum, *Kunth, Enum. Pl.* iii. 78; *Poepp. & Endl. Nov. Gen. et Sp.* iii. 84.—Peruv.

Brownii, *Mast. in Gard. Chron.* (1876) ii. 744.—Reg. Colomb.

candidum, Bull, Cat. (1875) 3. t. 11 = brevirostre.

cardiophyllum, C. Koch & August, Ind. Sem. Hort. Berol. (1854) *App.* 9 = nymphaeifolium.

carinatum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 275. —Reg. Colomb.

cartilagineum, *Kunth, Enum. Pl.* iii. 79.—Venezuela.

caucanum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 274. —Reg. Colomb.

caudatum, Kunth, Enum. Pl. iii. 80 = undatum.

Chemisonis, Schott, in Oestr. Bot. Wochenbl. (1857) 269 = Gaudichaudianum.

chelanicrista ×, N. E. Br. in Gard. Chron. (1885) ii. 650.

clavigerum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 84. —Peruv.

cochlastum, Moritz, ex Schott, Prod. Aroid. 499 = nymphaeifolium.

colocasiaefolium, *Decaisn. in Rev. Hortic.* (1879) 452. —Am. trop.?

commutatum, *Schott, in Bonplandia*, x. (1862) 146. —Hab.?

comtum, *Schott. l. c.* 87.—Bras.

ANTHURIUM:—

concinnatum, Schott, Prod. Aroid. 522 = bogotense.

concinnum, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 301 = Guildingii.

consanguineum, Kunth, Enum. Pl. iii. 72 = Harrisii.

consimile, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 294.—Peruv.

consobrinum, *Schott, l. c.* v. (1855) 66.—Nicaragua.

contentum, Schott, ex Engl. in DC. Monog. Phan. ii. 129 = Harrisii.

corvilimbum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 84.—Peruv.

cordato-sagittatum, Schott, in Oestr. Bot. Zeitschr ix. (1859) 100 = trinervium.

cordatum, *C. Koch & H. Sello, Ind. Sem. Hort. Berol.* (1853) *App.* 5.—Venezuela.

cordatum, G. Don, in Sweet, Hort. Brit. ed. III. 638 = Pothos cordata.

cordatum, Schott, Prod. Aroid. 491 = Lhotskyanum.

cordifolium, Hort. ex Gard. Chron. (1866) 934 = grande.

cordifolium, *Kunth, Enum. Pl.* iii. 76.—Costa Rica.

coriaceum, *G. Don, in Sweet, Hort. Brit.* ed. III. 638. —Bras.

costatum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1853) *App* 6.—Venezuela.

crassifolium, *N. E. Br. in Gard. Chron.* (1883) t. 10. —Reg. Colomb.

crassinervium, Hort. ex Engl. in DC. Monog. Phan. ii. 188 = acaule.

crassinervium, *Schott, Melet.* i. 22.—Panama.

crenatum, Kunth, Enum. Pl. iii. 75 = acaule.

cristallinum, *Linden & André, in Lindrn, Cat.* (1873) n. 90. t. 126.—Peruv.

cucullatum, Koch & Sello, Ind. Sem. Hort. Berol. (1853) *App.* 6 = andicola.

cupreum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 276.—Reg. Colomb.

cuspidatum, *Mast. in Gard. Chron.* (1875) i. 426.—Reg. Colomb.

cuspidifolium, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.—Costa Rica.

cymatophyllum, *C. Koch & Sello, in C. Koch, Wochenschr.* xi. (1868) 276.—Bras.

Dechenii, André, in Illustr. Hortic. xxiv. (1877) t. 269 = Spathiphyllum cannaeforme.

decurrens, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 85. t. 293.—Peruv.

dewuldeanum, Engl. in Engl. Bot. Jahrb. vi. (1885) 280. —Reg. Colomb.

Devansayanum ×, *Hort.—Cf. Gard. Chron.* (1888) i. 810.

digitatum, *G. Don, in Sweet, Hort. Brit.* ed. III. 638. —Peruv.

dolosum, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179 = scandens.

Dombeyanum, *Brongn. ex Engl. in DC. Monog. Phan.* ii. 146.—Peruv.

dominicense, *Schott, in Oestr. Bot. Zeitschr.* (1858) 350.—Ind. occ.

egregium, Schott, Prod. Aroid. 475 = ellipticum.

elatum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1858) App. 6 = pentaphyllum.

elegans, *Engl. in Engl. Bot. Jahrb.* i. (1881) 482.—Ins. Martinica.

ellipticum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1858) App. 6.—Venezuela.

emarginatum, *Baker, in Saund. Refug.* t. 274.—Am. trop.

emblema, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 275.—Peruv.

erythropodum, Miq. Hort. Amstelod. (1858) 8 = Harrisii.

fallax, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 309 = Guildingii.

Fendleri, *Schott, Prod. Aroid.* 468.—Venezuela.

Ferrierense ×.—Cf. Gard. Chron. (1888) ii. 788.

fissum, *C. Koch, in Ender, Ind. Aroid.* 109.—Reg. Colomb.

flavescens, *Poepp. & Endl. Nov. Gen. et Sp* iii. 83.— Peruv.

flavidum, *N. E. Br. in Gard. Chron.* (1883) ii. 651.— Reg. Colomb.

flexile, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.— Costa Rica.

flexuosum, *Kunth, Enum. Pl.* iii. 82.—Venezuela.

floribundum, Linden & André, in Illustr. Hortic. (1874) 23. t. 159 = Spathiphyllum floribundum.

ANTHURIUM:—

Fontanesii, *Schott, in Bonplandia*, x. (1862) 347.—Venezuela.

formosum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181. —Costa Rica.

Fraseri, *Engl. in DC. Monog. Phan.* ii. 112.—Ecuador.

fraternum, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 293 = Lindenianum.

Friedrichsthalii, Schott, l. c. v. (1855) 65 = Rudgeanum.

fucatum, Schott, l. c. viii. (1858) 387 = nymphaeifolium.

funiferum, Klotzsch & Karst. ex Engl. in DC. Monog. Phan. ii. 111 = Bredemeyeri.

Galeottianum, *C. Koch, in Ind. Sem. Hort. Berol.* (1855) *App.* 5.—Bras.

Galeottii, *C. Koch, l. c.* (1854).—Mexic.

Gaudichaudianum, *Kunth, Enum. Pl.* iii. 74.—Bras.

Geitnerianum, *A. Regel, in Ind. Sem. Hort. Petrop.* (1866) 38.—Am. trop.

Ghiesbrechtii, Linden, ex Schott, Prod. Aroid. 538 = podophyllum.

gladiifolium, *Schott, in Journ. Bot.* i. (1868) 5.— Bras.

glaucescens, Kunth, Enum. Pl. iii. 73 = coriaceum.

glaucum, Schott, Melet. i. 22 = coriaceum.

Glaziovii, *Hook. f. Bot. Mag.* t. 6838.—Bras.

gracile, *Lindl. Bot. Reg.* t. 1835.—Guiana.

grande, *Hort. Bull, ex Gard. Chron.* (1865) 934.— Am. austr.

grandifolium, *Kunth, Enum. Pl.* iii. 77.—Reg. Colomb.

grosarum, Schott, in Oestr. Bot. Zeitschr. (1859) 101 = pentaphyllum.

Guildingii, *Schott, in Oestr Bot. Wochenbl.* (1857) 301.—Ind. occ.

Gustavi, *Regel, Gartenfl.* (1878) 324, nomen; (1882) 67.—Reg. Colomb.

gymnopus, *Griseb. Cat. Pl. Cub.* 219.—Cuba occ.

Hahnii, *Engl. in Engl. Bot. Jahrb.* i. (1881) 481.—Ins. Martinica.

Harrisii, *G. Don, in Sweet, Hort. Brit.* ed. III. 638.— Bras.

helleborifolium, Hort. ex Engl. in DC. Monog. Phan. ii. 206 = variabile.

helleborifolium, *Schott, in Bonplandia*, x. (1862) 148. —Hab.?

Hoffmanni, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181. —Costa Rica.

Hoffmannsegii, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 325.—Bras.

Holtonianum, *Schott, l. c.* 317.—Reg. Colomb.

Hookeri, *Kunth, Enum. Pl.* iii. 74.—Ind. occ.

Hookeri, Schott, Prod. Aroid. 472 = tetragonum.

Huegelii, Schott, in Oestr. Bot. Wochenbl. v. (1855) 83 = Hookeri.

Humboldtianum, Kunth, Enum. Pl. iii. 78 = rubrinervium.

Humboldtianum, Schott, Prod. Aroid. 524 = polyrrhizon.

humile, Schott, l. c. 449 = amoenum.

hybridum, *Hort. ex Engl. in DC. Monog. Phan.* ii. 191.—Hab.?

hygrophilum, *Engl. in Bot. Jahrb.* vi. (1885) 276.— Reg. Colomb.

ianthinopodium, Schott, ex Engl. in Mart. Fl. Bras. iii. 11. 89 = Harrisii.

illepidum, Schott, in Oestr. Bot. Zeitschr. (1859) 100 = longifolium.

imperiale, *Miq. ex Schott, Prod. Aroid.* 465.—Hab.?

inamoenum, Schott, l. c. 504 = nymphaeifolium.

inconditum, Schott, in Oestr. Bot. Zeitschr. (1858) 181 = gracile.

inconspicuum, *N. E. Br. in Gard. Chron.* (1885) t. 787. —Bras.

indecorum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 350. —Peruv.

indecorum, Schott, ex Engl. in DC. Monog. Phan. ii. 205 = variabile.

insigne, *Mast. in Gard. Chron.* (1878) i. 430.—Reg. Colomb.

intermedium, Kunth, Enum. Pl. iii. 76 = Harrisii.

Isertianum, Schott, Prod. Aroid. 498 = Guildingii.

Ylleki, Schott, in Bonplandia, x. (1862) 5 = Harrisii.

Kalbreyeri, *Hort. ex Gard. Chron.* (1881) ii. 117.—N Granad.!

Karwinskyi, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 101; *Prod. Aroid.* 553.—Am. cent.

ANTHURIUM:—
Kastelskii, *Schott, in Oestr. Bot. Zeitschr.* (1859) 100. —Chili.
Kellermanni, *Schubert, ex Kellerm. in Wien. Illustr. Gart. Zeit.* ix. (1884) 6.—Hab.?
Kunthianum, *Liebm. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 272.—Mexic.
Kunthii, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 84.—Pervv.
lactiflorum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 277. —Reg. Colomb.
lanceolatum, *Kunth, Enum. Pl.* iii. 71.—Am. austr.
lanceolatum, G. Don, in Sweet, Hort. Brit. ed. III. 634 = Willdenowii.
lancifolium, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180. —Costa Rica.
Langsdorffii, *Schott, Prod. Aroid.* 458.—Bras.
lapathifolium, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 309 = ochranthum
Lauchenaum, *C. Koch, in Allgem. Gartens.* (1857) 191.—Bras.
Lechlerianum, *Schott, Prod. Aroid.* 584.—Pervv.
leptostachyum, Schott, in Oestr Bot. Wochenbl. v. (1855) 66 = Olfersianum.
leucocarpum, Schott, l. c. (1857) 53 = scandens.
leuconeurum, *Lem. in Illustr. Hortic.* ix. (1862) t. 314. —Am. trop.
Lhotzkyanum, *Schott, Prod. Aroid.* 491.—Bras.
Libonianum, Lindl. & Regel, in Gartenfl. (1857) 291. t. 558 = Gaudichaudianum.
Liebmanni, *Schott, in Bonplandia*, vii. (1859) 165.— Mexic.
Lindenianum, Hort. ex Engl. in DC. Monog. Phan. ii. 176 = nymphaeifolium.
Lindenianum, *C. Koch & August, in C. Koch, Allg. Gartenz.* (1857) 234.—Reg. Colomb.
Lindigii, Hrincq, L'Hortic. Franç. (1866) 17; N. E. Br. in Gard. Chron. (1879) ii. 554 = Lindenianum.
lingulifolium, Hort. ex Engl. in DC. Monog. Phan. ii. 124 = Geitnerianum.
longifolium, Pr. t. ex Engl. l. c. 129 = Harrisii.
longifolium, *GriDon, in Sweet, Hort. Brit.* ed. III. 633; *Kunth, Enum. Pl.* iii. 69.—Bras.
longipaphum, *Schott, Prod. Aroid.* 520.—Costa Rica.
lucidum. *Kunth, Enum. Pl.* iii. 78.—Bras.
Luschnathianum, Kunth, l. c. = Olfersianum.
maclentum, Schott, in Bonplandia, vii. (1859) 165 = gracile.
Maclesii, *Schott, in Oestr. Bot. Zeitschr.* (1858) 350. —Pervv.
macrolobium x, *Hort. Bull, in Gard. Chron.* (1883) i. 406.—Hab.?
macrophyllum, G. Don, in Sweet, Hort. Brit. ed. III. 638 = Pothos macrophylla.
macrophyllum, Schott, in Oestr. Bot. Wochenbl. v. (1855) 53 = Selloum.
macrospadix, *Lem. Illustr. Hortic.* ix. (1862) *Misc.* 90. —Guiana.
magnificum, *Linden, in Belg. Hortic.* xv. (1865) 98.— Reg. Colomb.
magnificum, Hort. ex Gard. Chron. (1865) 934 = grande.
Malyi, *Maximil. (Archd. Austr.) ex Schott, in Bonplandia*, x. (1862) 292.—Bras.
mandiocanum, Schott, Prod. Aroid. 454 = Harrisii.
margaritaceum, Baker, in Saund. Refug. t. 380 = Hoffmanni.
Martianum, *C. Koch & Kolb, in C. Koch, Wochenschr.* xi. (1868) 276.—Guiana.
Martini, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 338.—Guiana.
martinicense, *Engl. in Engl. Bot. Jahrb.* i. (1881) 480. —Ins. Martinico.
Maximiliani, *Schott, in Bonplandia*, x. (1862) 5.—Bras.
membranuliferum, Schott & Ohlend. ex Engl. in DC. Monog. Phan. ii. 196 = podophyllum.
metallicum, *Linden, ex Schott, Prod. Aroid.* 506.— Hab.?
mexicanum, *Engl. in DC. Monog. Phan.* ii. 105.—Mexic.
mexicanum, Liebm. in Kjoeb. Videnak. Meddel. (1849) 21 = Schlechtendalii.
microphyllum, *G. Don, in Sweet, Hort. Brit.* ed. III. 633.—Bras.
microspadix, Schott, in Oestr. Bot. Zeitschr. (1858) 180 = porrectum.
microstachyum, *Kunth, Enum. Pl.* iii. 78.—Reg. Colomb.
Miquelianum, *C. Koch & August, Ind. Sem. Hort. Berol.* (1855) 5.—Bras.
moricandum, *Hemsl. Diag. Pl. Nov.* 36.—Guatemala.

ANTHURIUM:—
myosuroides, *Endl. ex Kunth, Enum. Pl.* iii. 72.—Reg. Colomb.
neglectum, Miq. in Cat. Hort. Amstel. (1858) = Hookeri.
nitidum, *Benth. Pl. Hartw.* 255.—Reg. Colomb.
nymphaefolium, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1858) *App.* 6.—Venezuela.
obliquum, *Schott, in Oestr. Bot. Zeitschr.* (1856) 357.—Venezuela.
oblongatum, *Schott, ex Engl. in Mart. Fl. Bras.* iii. 89 = Olfersianum.
obtusilolium, *G. Don, in Sweet, Hort. Brit.* ed. III. 633; *Kunth, Enum. Pl.* 79.—Barbadoes.
obtusilobum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181.—Costa Rica.
ochranthum, *C. Koch, Ind. Sem. Hort. Berol.* (1858) *App.* 6.—Am. trop.
Oerstedianum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.—Costa Rica.
Olfersianum, *Kunth, Enum. Pl.* iii. 79.—Bras.
ornatum, Schott, in Oestr. Bot. Wochenbl. (1857) 294 = nymphaeifolium.
Ottonianum, *Kunth, Enum. Pl.* iii. 83.—Bras.
oxybelium, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 310 = nitidum.
oxycarpum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 83.— Pervv.
padicifolium, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 273 = Bras.
pallens, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.— Costa Rica.
palmatifidum, *Van Houtte, ex Schott, Prod. Aroid.* 557.—Hab.?
palmatum, *G. Don, in Sweet, Hort. Brit.* ed. III. 633; *Kunth, Enum. Pl.* iii. 80.—Ins. Dominic.
pandurseforme, *Schott, Prod. Aroid.* 536.—Costa Rica.
panduriforme, *Mart. ex Schott, in Oestr. Bot. Wochenbl.* (1855) 273.—Bras.
panduratum, Schott, l. c. = pandurseforme.
parvispathum, *Hemsl. Biol. Cent. Am. Bot.* iii. 492. —Guatemala.
Patini, R. Hogg, in Gard. Chron. (1875) i. 524 = Spathiphyllum Patini.
pedatifolium, Regel & Lieden, Petrop. (1865) 44; et in Gartenfl. (1866) 66. t. 501 = pedato-radiatum.
pedato-radiatum, *Schott, in Bonplandia*, vii. (1859) 287.—Mexic.
pedatum, *Endl. ex Kunth, Enum. Pl.* iii. 79.—N. Granat.
peltatum, Poepp. & Endl. Nov. Gen. et Sp. iii. 83. t. 296 = Homalonema peltata.
pentaphyllum, *G. Don, in Sweet, Hort. Brit.* ed. III. 633; *Kunth, Enum. Pl.* iii. 81.—Bras.
podophyllum, *Kunth, l. c.* 80.—Mexic.
Poeppeanum, Kunth, l. c. 68 = gracile.
Poiteauanum, Schott, Prod. Aroid. 458 = gracile.
polyrrhizon, *C. Koch & August, Ind. Sem. Hort. Berol.* (1856) *App.* 7.—Venezuela.
polytomosum, Schott, in Bonplandia, vii. (1859) 287 = podophyllum.
popayanense, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 274. —Reg. Colomb.
porrectum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180. —Costa Rica.
protensum, *Schott, l. c.* 181.—Am. centr.
pseudopodophyllum, Schott, in Bonplandia, vii. (1859) 326 = podophyllum.
pulchellum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 278. —Reg. Colomb.
Purdieanum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 53.—Reg. Colomb.
quindiuense, Schott, l. c. 293 = fraternum.
quinquenervium, *Kunth, Enum. Pl.* iii. 76.—Reg. Colomb.
quinquenervinerum, Schott, ex Engl. in Mart. Fl. Bras. iii. ii. 100 = undatum.
radicosa, *C. Koch & A. Haage, Ind. Sem. Hort. Berol.* (1854) *App.* 10.—Bras.
recusatum, *Schott, Prod. Aroid.* 467.—Cuba.
reflexum, *Hort. Par. ex Regel, Gartenfl.* (1866) 269.— Am. trop.
reflexum, G. Don, in Sweet, Hort. Brit. ed. III. 633 = coriaceum.
regale, *Linden, in Belg. Hortic.* xvi. (1866) 200.—Am. trop.
Regnellianum, *Engl. in Mart. Fl. Bras.* iii. ii. 96.— Bras.

ANTHURIUM:—
repandum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 317. —Reg. Colomb.
reptans, *G. Don, in Sweet, Hort. Brit.* ed. III. 633.— Bras.
reticulatum, *Benth. Pl. Hartw.* 255.—Pervv.
rigidulum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180 —Costa Rica.
Roezlii, *Regel, in Gartenfl.* (1879) 331.—Reg. Colomb.
Rothschildianum x, *Hort.—Cf. Gard. Chron.* (1883) i. 510.
rubescens, *G. Don, in Sweet, Hort. Brit.* ed. III. 634.— Bras.
rubidum, Schott, ex Engl. in DC. Monog. Phan. ii. 120 = Harrisii.
rubricaule, Kunth, Enum. Pl. iii. 89 = Harrisii.
rubrinervium, *G. Don, in Sweet, Hort. Brit.* ed. III. 633. —Venezuela.
Rudgeanum, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 65.—Bras.
rugosum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 389. —Venezuela.
sagittaria, Linden, ex Schott, Prod. Aroid. 597 = pandurseforme.
sagittatum, G. Don, in Sweet, Hort. Brit. ed. III. 633 = rubrinervium.
Salvinise, *Hemsl. Diag. Pl. Nov.* 36.—Guatemala.
sanguineum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 279. —Reg. Colomb.
Saundersii, *Hook. f. Bot. Mag.* t. 6216.—Bras.
scandens, *Engl. in Mart. Fl. Bras.* iii. ii. 78.—Bras.
Scherzerianum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 53.—Guatemala.
Schlechtendalii, *Kunth, Enum. Pl.* iii. 75.—Mexic.
scolopendrinum, Kunth, l. c. 68 = gracile.
scolopendroides, Hort. ex Engl. in DC. Monog. Phan. ii. 129 = Harrisii.
Selloum, *C. Koch, Ind. Sem. Hort. Berol.* (1854) — Ins. S. Jan.
Sellowianum, *Kunth, Enum. Pl.* iii. 70.—Bras.
signatum, *C. Koch & L. Mathieu, Ind. Sem. Hort. Berol.* (1856) 5.—Reg. Colomb.
sinuatum, *Benth. ex Schott, in Oestr. Bot. Wochenbl.* (1857) 318.—Bras.
smilacifrons, C. Koch, Ind. Sem. Hort. Berol. (1856) 8 = undatum.
solitarium, Schott, Prod. Aroid. 478 = affine.
Sondervianum, Schott, in Bonplandia, vi. (1858) 372 = pentaphyllum.
sororium, *Schott, Prod. Aroid.* 592.—Peruv.
spathiphyllum, *N. E. Br. in Gard. Chron.* (1877) i. 652. —Hab.?
spectabile, *Hérincq, L'Hortic. Franç.* (1866) 17.—N. Granat.
spectabile, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181. —Costa Rica.
splendidum, *Hort. Bull, ex Illustr. Hortic.* xxx. (1883) t. 510. (*Cf. Gard. Chron.* (1883) i. 381).—Reg. Colomb.
stipitatum, *Benth. Pl. Hartw.* 255.—Reg. Colomb.
striatum, *C. Koch & L. Mathieu, Ind. Sem. Hort. Berol.* (1855) 7.—Pervv.
strictum, *N. E. Br. in DC. Monog. Phan.* ii. 488. —Paraguay.
subcordatum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181.—Guatemala.
subhastatum, *Schott, l. c.* 350.—Reg. Colomb.
subnigitatum, *Kunth, Enum. Pl.* iii. 79.—Venezuela.
subsignatum, *Schott, in Bonplandia*, iii. (1862) 365.— Costa Rica.
subtriangulare, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 279.—Reg. Colomb.
subtrilobum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 193.—Venezuela.
suzannense, *Mig. in Nat. Verh. Wet. Haarl.* vii. (1851) 211.—Surinam.
Swartzianum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 333.—Ind. occ.
tapinocarpum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.—Costa Rica.
tetragonum, *Hemsl. ex Schott, Prod. Aroid.* 476.—Hab.?
tolimanse, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 277. —Reg. Colomb.
trifidum, Oliver, in Hook. f. Bot. Mag. t. 6329 = signatum.
trilobum, Hort. Lindl. in Illustr. Hortic. xxiv. (1877) t. 235 = signatum.
trinerve, *Mig. in Linnaea*, xvii. (1843) 67.—Bras.
trinervium, *Kunth, Enum. Pl.* iii. 76.—Bras.

ANTHURIUM:—
triphyllum, *Brongn. ex Schott, Prod. Aroid.* 548.—Bolivia.
truncatulum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 275.—Reg. Colomb.
umbrosum, *Liebm. in Kjoeb. Videnskb. Meddel.* (1849) 91.—Mexic.
undatum, *Schott, Melet.* i. 22.—Bras.
undulatum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1854) App. 107 = Harrisii.
undulifolium, C. Koch, ex Ender, Ind. Aroid. 17 = undatum.
Urvilleanum, Schott, Prod. Aroid. 459 = Harrisii.
variabile, *Kunth, Enum. Pl.* iii. 31.—Bras.
varians, Miq. Hort. Amstel. (1853) = acaule.
Veitchenum, Hort. in Rev. Hortic. (1877) 37 = seq.
Veitchi, *Mast. in Gard. Chron.* (1876) 775.—Reg. Colomb.
Vellosianum, Schott, Prod. Aroid. 482 = Olfersianum.
venosum, *Griseb. Cat. Pl. Cub.* 219.—Cuba.
violaceum, Schott, Melet. i. 23 = scandens.
violascens, Schott, in Oestr. Bot. Wochenbl. (1854) 89 = costatum.
viride, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1855) 6 = Sellowianum.
virgosum, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 100.—Bras.
Wagenarianum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1858) App. 7.—Venezuela.
Wallisii, *Mast. in Gard. Chron.* (1875) 420.—Reg. Colomb.
Waluiewi, *Regel, in Act. Hort. Petrop.* vi. (1879) 290.—Venezuela.
Warocqueanum, *Moore, Florist & Pomol.* (1878) 101.—Reg. Colomb.
Warscewiczii, C. Koch, Ind. Sem. Hort. Berol. (1855) App. 4 ; et in Ender, Ind. Aroid. 17 = sinuatum.
Wendlandii, *Schott, in Oestr. Bot. Zeitschr.* (1856) 182.—Costa Rica.
Willdenowii, Kunth, Enum. Pl. iii. 71 = lanceolatum.
Williamsii, Hort. ex Engl. in DC. Monog. Phan. ii. 117 = Scherzerianum.

ANTHYLLIS, Riv. ex Linn. Syst. ed. I (1735). *LEGUMINOSÆ*, Benth. & Hook. f. i. 488.
ACANTHYLLIS, Pomel, Nouv. Mat. Fl. Atl. 179 (1874).
BARBA-JOVIS, Tourn. ex Adans. Fam. ii. 321 (1763).
CORNICINA, Boiss. Voy. Bot. Espagne, 162 (1843).
DORYCNOPSIS, Boiss. l. c. 163 (1843).
FALKLORA, Rafin. Sylva Tellur. 89 (1838).
PSYBANTHYLLIS, Boiss. Voy. Bot. Espagne, t. 162 (1843).
POGONITIS, Reichb. Handb. 236 (1837).
TRIPODION, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 346.
VULNERARIA, Tourn. ex Hall. Enum. Stirp. Helv. ii. 589 (1742).
ZENOPOGON, Link, Handb. ii. 481 (1831).
affinis, Britting. ex Koch, in Roehl. Deutschl, Fl. ed. III. v. 194 = A. Vulneraria.
alpestris, Kit. ex Schult. Oestr. Fl. ed. II. ii. 317 = A. Vulneraria.
alpina, G. Don, in Loud. Hort. Brit. 283 = A. Vulneraria.
argentea, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 408 = Cytisopsis dorycnifolia.
argyrea, Salisb. Prod. 328 = A. Barba-Jovis.
arundana, Boiss. & Reut. Pugill. Pl. Nov. (1852) 35 = A. Vulneraria.
Asplatshi, Ser. in DC. Prod. ii. 169 = A. Hermanniæ.
asphaltoides, Linn. Cent. Pl. ii. ; Amœn. Acad. iv. 296 = Aspalathus anthylloides.
aurea, *Welden. ex Host, Fl. Austr.* ii. 319.—Graecia ; Dalmat.
balearica, *Cosa. ex Maras & Virgin. Cat. Pl. Balear.* 75.—Ins. Balear.
Barba-Jovis, *Linn. Sp. Pl.* 720.—Reg. Mediterr.
bicephala, Gilib. Fl. Lituan. ii. 96 = A. Vulneraria.
bicolor, Bertol. Fl. Ital. vii. 401 = A. Vulneraria.
bifloscula, *Munby, Fl. Alger.* 76.—Algeria.
bifloura, [*Soland. in*] *Russ. Aloppo.* ed. II. ii. 259.—Syria.
calycicola, Schur, Enum. Pl. Transs. 150 = A. Vulneraria.
capensis, Lam. Encyc. i. 305 = Lebeckia cytisoides.
carpatica, *Pantoč. in Magyar Növénvt. Lapok*, vi. (1883) 168.—Europ.
cephalotes, *Vis. Fl. Dalmat.* iii. 178.—Dalmatia.
chilensis, Bert. ex Steud. Nom. ed. II. i. 107 = Adesmia Loudonia.
chilensis, Ser. in DC. Prod. ii. 171 = Hosackia subpinnata.
cicerifolia, *Pourr. ex Colm. in Anal. Soc. Hist. Nat.* i. (1870) 88.—Hispan.
collina, Salisb. Prod. 328 = A. Vulneraria.
cornicina, *Linn. Sp. Pl.* 719.—Hispan.
cornicina, Pall. ex Georgi, Beschr. Russ. Reich. Nachtr. 394 = Medicago cretacea.
cornicina, Poir. Voy. Barb. ii. 311 = hamosa.
cretica, Lam. Encyc. i. 304 = Ebenus cretica.
cusanata, Boch.-Ham. ex D. Don, Prod. Fl. Nep. 246 = Lespedesa eriocarpa.
cytisoides, *Linn. Sp. Pl.* 720.—Europ.
Dillenii, Schult. ex Steud. Nom. ed. I. 56 = A. Vulneraria.
Erinacea, Linn. Sp. Pl. 720 = Erinacea pungens.
Fama, Gouan, Herbor. 173, nomen = A. Vulneraria.
Genistae, *Dufour, ex DC. Prod.* ii. 169.—Hispan.
genistoides, Dufour, in Bull. Soc. Bot. Fr. vii. (1860) 324 = A. Genistae.
Gerardi, *Linn. Mant.* 100.—Reg. Mediterr. occ.
glaucescens, Kit. in Linnæa, xxxii. (1863) 613 = alpestris.
gracilis, Salisb. Prod. 332 = A. cytisoides.
hamosa, *Desf. Fl. Atlant.* ii. 153.—Hispan. ; Afr. bor. Hermanniae, *Linn. Sp. Pl.* 720.—Reg. Mediterr. ; Orient.
heterophylla, *Linn. l. c.* ed. II. 1013.—Hispan.
hispida, Boiss. & Reut. Pugill. Pl. Nov. (1852) 86 = A. Vulneraria.
Hystrix, Sweet, Hort. Brit. ed. I. 114, nomen = Erinacea pungens !
incisa, Willd. Sp. Pl. iii. 1017 = Cicer crvoides.
indica, *Lour. Fl. Cochinch.* 429.—Cochinch.
involucrata, Linn. Mant. 265 = Lotononis involucrata.
italica, Pert. ex Loud. Hort. Brit. Suppl. ii. 610 = A. Vulneraria.
Jacquini, *A. Kern. in Zeitschr. Ferdinand. Tirol.* III. xv. (1870) 387.—Europ.
leguminosa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 597 = A. Vulneraria.
Lemanniana, *Lowe, in Hook. Kew Journ.* viii. (1856) 291.—Ins. Madera.
linifolia, Linn. Mant. 265 = Aspalathus linearifolia.
lotoides, *Linn. Sp. Pl.* 720.—Hispan.
macrocarpa, Welp. Rep. i. 634 = A. Vulneraria.
macrocephala, Wender. in Linnæa, v. (1830) Lit. 54 = A. Vulneraria.
marsiana, Schweig. in Fl. Dan. t. 2738 = A. Vulneraria.
monocephalos, Gilib. Fl. Lituan. ii. 97 = A. Vulneraria.
montana, *Linn. Sp. Pl.* 719.—Europ.
numidica, *Cosa. & Durr. in Bull. Soc. Bot. Fr.* iv. (1857) 896, nomen.—Algeria.
onobrychioides, *Cav. Ic.* ii. 40. t. 150.—Hispan.
onoides, Burm. f. Fl. Cap. Prod. 21 = Ononis anthylloides.
pallida, Opiz, ex Nym. Consp. 164 = A. Vulneraria.
pentaphylla, *Steud. Nom.* ed. II. i. 107.—Sibir.
podocephala, *Boiss. Elench.* 84.—Hispan.
polycephala, *Desf. Fl. Atlant.* ii. 150. t. 195.—Atlas.
polycephala, Reichb. Fl. Germ. Excurs. 516 = aurea.
polyphylla, Kit. ex DC. Prod. ii. 170 = A. Vulneraria.
pseudocytisus, Welp. Rep. ii. 846 = Cytisopsis dorycnifolia.
pubescens, Stokes, Bot. Mat. Med. iii. 517 = A. Vulneraria.
quinquefiora, Linn. f. Suppl. 335 = Aspalathus araneosa, parviflora.
Ramburei, *Boiss. Elench.* 85.—Hispan.
rosea, Willk. ex Reichr. in Anal. Soc. Esp. Hist. Nat. viii. (1879) 53 = balearica.
rubicunda, Wender. ex Steud. Nom. ed. II. i. 107 = A. Vulneraria.
rubra, Gouan, Herbor. 172, nomen = A. Vulneraria.
rupestris, Cass. Notes Crit. 155.—Hispan.
rustica, Willd. Gard. Dict. ed. VIII. n. 8 = A. Vulneraria.
rusticana, Wender. in Linnæa, v. (1830) Lit. 54 = A. Vulneraria.
sanguinea, Schur, Enum. Pl. Transs. 150, in syn. = A. Vulneraria.
sericea, *Lag. Gen. et Sp. Nov.* 22.—Hispan.
sericea, Willd. Sp. Pl. iii. 1014 = Ebenus Sibthorpii.
splendens, Willd. l. c. 1018 = A. Barba-Jovis.
Spruneri, Heldr. ex Nym. Consp. 164 = A. Vulneraria.
tenuifolia, Roth, Tent. Fl. Germ. i. 332.—Europ.
tetraphylla, *Linn. Sp. Pl.* 719.—Reg. Mediterr.
Tournefortii, Schult. ex Steud. Nom. ed. I. 56 = A. Vulneraria.
tragacanthoides, Desf. Fl. Atlant. ii. 150 = Astragalus armatus.
tragacanthoides, Labill. Ic. Pl. Syr. Dec. ii. 16 = Astragalus colutecoides.
tricolor, *Vahot. in Rad. Yugos. Ahad. Zagrab*, xxxiv. (1876).—Europ.
variegata, Boiss. Diagn. Ser. II. v. 81 = A. Vulneraria.
vicifiora, *Linn. f. Suppl.* 335.—Afr. austr.
Vulneraria, *Linn. Sp. Pl.* 719.—Europ. ; Orient ; Abyss.
vulnerarioides, *Bonj. ex Reichb. Fl. Germ. Excurs.* 515.—Ital.
Wahlbiana, Webb, ex Walp. Rep. i. 68 = tejedensis.
Webbiana, Hook. Bot. Mag. t. 3284 = A. Vulneraria.
Weldemana, Reichb. Fl. Germ. Excurs. 515 = A. Vulneraria.

ANTIARIS, Leach. in Ann. Mus. Par. xvi. (1810) 478. t. 22. *URTICACEÆ*, Benth. & Hook. f. iii. 371.
ANTSCHAR, Horsf. in Verh. Bat. Gen. vii. (1814) 8.
IPO, Pers. Syn. ii. 566 (1807).
LEPURANDRA, J. Grah. Cat. Pl. Bomb. 193 (1839).
Bennettii, *Seem. in Bonplandia*, x. (1862) 8.—Ins. Fiji.
d'ebia, Span. ex Hook. Comp. Bot. Mag. i. (1885) 310 = toxicaria.
innoxia, Blume, Rumphia, 172. t. 54 = toxicaria.
macrophylla, *R. Br. in Fiind. Voy.* ii. 602. t. 5.—Austral.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* i. 424.—Sumatra.
rufa, *Miq. l. c.* 428.—Sumatra.
saccadora, Dals. in Hook. Kew Journ. iii. (1851) 282 = toxicaria.
toxicaria, *Lesch. in Ann. Mus. Par.* xvi. (1810) 478.—Ind. or. ; Malaya.
zeylanica, Seem. in Bonplandia, x. (1862) 4, in note ; et Fl. Vit. 255 = toxicaria.

ANTICHARIS, Endl. Nov. Stirp. Dec. 22 ; Gen. 682 (1839) = Leichhardtia, t. 98. *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 928.
DORATANTHERA, Benth. in Endl. Gen. 685 (1839).
MEIBARRHENA, R. Br. in Salt, Abyssin. App. 68 (1816).
arabica, *Endl. Nov. Stirp. Dec.* 23.—Arabia.
arabica, Hochst. ex Benth. in DC. Prod. x. 347 = linearis.
glandulosa, *Aschers. in Monatsb. Acad. Berl.* (1866) 880.—Aegypt. ; Arab. ; Ind. or.
linearis, *Hochst. ex Aschers. l. c.* 882.—Afr. trop. ; Arab. ; Ind. or.

ANTICHLOA, Steud. Nom. ed. II. i. 108 (1840) = ACTINOCHLOA, Willd. = Boutelous, Lag. (Gram.).

ANTICHORUS, Linn. Mant. i. 9 (1767) = Corchorus, Linn. (Tiliac.).
depressus, Linn. l. c. 64 = C. Antichorus.

ANTICLEA, Kunth, Enum. Pl. iv. 191 (1843) = Zygadenus, Michx. (Liliac.).
Fremonti, Torr. in Facif. Rail. Rep. iv. 144 = Z. Fremonti.
glauca, Kunth, Enum. Pl. iv. 192 = Z. glaucus.
mexicana, Kunth, l. c. 192 = Z. mexicanus.
Nuttallii, Torr. in Pacif. Rail. Rep. iv. 144 = Z. Nuttallii.
sibirica, Kunth, Enum. Pl. iv. 191 = Z. sibiricus.
volcanica, Baker, in Journ. Linn. Soc. xvii. (1880) 483 = Z. volcanicus.

ANTICORYNE, Turcz. in Bull. Math.-Phys. Acad. Petersb. x. (1852) 382 = Reachea, Linn. (Myrtac.).
dicomoides, Turcz. l. c. = R. ovalifolia.

ANTIDAPHNE, Poepp. & Endl. Nov. Gen. et Sp. ii. 70. t. 198 (1838). *LORANTHACEÆ*, Benth. & Hook. f. iii. 216.
viscoidea, *Poepp. & Endl. l. c.*—Peruv.

ANTIDESMA, Burm. ex Linn. Diss. de Anandr. 6 (1745); Amoen. Acad. i. 249 (1749). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 284.
BRETTAM, Adans. Fam. ii. 354 (1763).
MINUTALIA, Fenzl, in Flora, xxvii. (1844) 312.
RHYTIS, Lour. Fl. Cochinch. 660 (1790).
SYLAGO, Linn. Mant. i. 16 (1767).
acidum, Retz. Obs. v. 30 = A. Ghesaembilla.
acuminatum, *Wight, Ic.* t. 1991.—Ind. or.
Alexiteria, Gaertn. Fruct. i. 188. t. 39 = A. Menasu ?
Alexiteria, *Linn. Sp. Pl.* 1027.—Ind. or.
Alexiteria, Willd. Sp. Pl. iv. 762 = nigricans.
alexiterium, Presl, Epim. Bot. 234 = leptocladum.
alnifolium, *Hook. Ic. Pl.* t. 481.—Afr. austr.
amboinense, Miq. Ann. Mus. Bot. Lugd. Bat. i. 218 = stipularе.
attenuatum, Wall. Cat. n. 7286 = velutinosum.
auritum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 208.—Java.
barbatum, Presl, Epim. Bot. 233 = rostratum.
bicolor, Hassk. Cat. Hort. Bog. Alt. 81 = Excaecaria bicolor.
bifrons, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 229 = venosum.
Blumei, Tul. l. c. 211 = tetrandrum.
Bouvinianum, Baill. Adansonia, ii. (1861–62) 45 = venosum.
Boviniense, *Baker, Fl. Maurit.* 306.—Ins. Mascar.
Bunius, *Spreng. Syst.* i. 836.—Ind. or.; Malaya.
Bunius, Wall. Cat. n. 7282 A. = nigricans.
celebicum, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 218.—Ins. Celebes.
cilsatum, Presl, Epim. Bot. 234 = A. Bunius.
comptum, *Tul. in Ann. Sc. Nat.* Sér. xv. (1851) 190.—Ind. or.
cordifolium, Presl, Epim. Bot. 235 = A. Bunius.
coriaceum, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 204 = fallax.
Cumingii, *Muell. Arg. in DC. Prod.* xv. ii. 249.—Ins. Philipp.
cuspidatum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 67.—Ind. or.
Dallachyanum, *Baill. Adansonia*, vi. (1865–66) 337.—Austral.
diandrum, *Heyne, ex Roth, Nov. Pl. Spec.* 369.—Ind. or.
diandrum, *Spreng. Syst* i. 826; *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 196.—Ind. or.
Dispenhorsti, *Miq. Fl. Ind. Bat. Suppl.* 467.—Sumatra.
digitaliforme, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 191.—Ins. Philipp.
diversifolium, *Miq. Fl. Ind. Bat. Suppl.* 468.—Sumatra.
ereetre, *F. Muell. ex Benth. Fl. Austral.* vi. 87.—Austral.
erythrocarpum, *Muell. Arg. in DC. Prod.* xv. ii. 258.—Java.
erythroxyloides, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 208.—Madag.
excavatum, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 216.—Ins. Celebes.
fallax, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 68.—Malaya.
filiforme, Blume, Bijdr. 1194 = Bennettia filiformis.
flexuosum, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 225 = nigricans.
floribundum, Tul. l. c. 189 = A. Bunius.
frutescens, *Loch, in Malay. Misc.* ii. (1800) n. vii. 91.—Sumatra.
fruticosum, *Muell. Arg. in DC. Prod.* xv. ii. 259.—Cochinch.
fruticulosum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 287.—Pegu.
Ghesaembilla, *Gaertn. Fruct.* i. 189. t. 39.—Ind. or.; Malaya; Austral.; Afr. trop.
glabrum, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 188 = A. Bunius.
Grossularia, Rasusch. Nom. ed. III. 287 = Embelia Ribes.
guineensis, G. Don, ex Hook. Niger Fl. 515, nomen = Uapaca guineensis.
heterophyllum, *Blume, Bijdr.* 1123.—Java.
heterophyllum, Presl, Epim. Bot. 234 = leptocladum.
japonicum, *Steb. & Zucc. in Abh. Baier. Akad.* iv. iii. (1846) 213.—China; Japon.
lachnatum, *Muell. Arg. in Flora*, xlvii. (1864) 529.—Afr. occ.
lanceolatum, *Boj. Hort. Maurit.* 289.—Ins. Mascar.

ANTIDESMA:—
lanceolarium, Wall. Cat. n. 7284 = diandrum.
lanceolarium, Moritzi, Syst. Verz. Pfl. Zoll. 73 = lanceolatum.
lanceolatum, Dalz. & Gibs. Bomb. Fl. 237 = A. Menasu.
lanceolatum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 195.—Ind. or.
leptobotryum, *Muell. Arg. in Flora*, xlvii. (1864) 529.—Afr. trop.
leptocladum, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 199 = Moritzii.
leucopodum, *Miq. Fl. Ind. Bat. Suppl.* 466.—Sumatra.
littoreum, *Blume, Bijdr.* 1123.—Java.
Lobblanum, *Muell. Arg. in DC. Prod.* xv. ii. 254.—Java.
longifolium, *Boj. Hort. Maurit.* 289.—Ins. Mascar.
longrifolium, Decne. ex Baill. Adansonia, v. (1866) 349, nomen = Richeria grandis.
lunatum, Miq. Fl. Ind. Bat. Suppl. 467 = Aporosa lunata.
macrophyllum, *Wall. ex Voigt, Hort. Suburb. Calc.* 295, nomen.—Ind. or.
madagascariense, *Lam. Encyc.* i. 306.—Madag.
Mauritianum, Muell. Arg. in Flora, xlvii. (1864) 519 = Thecacoris Mauritiana.
mauritanicum, *Presl, Epim. Bot.* 232.—Burma.
membranaceum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 68.—Afr. occ.
Menasu, *Miq. ex Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 215.—Ind. or.
Menasu, Kurz, For. Fl. ii. 360 = martabanicum.
minus, *Blume, Bijdr.* 1123.—Java.
mucosa, Wall. Cat. n. 7283 = coriaceum.
molle, Wall. Cat. n. 7287; Muell. Arg. in Linnaea, xxxiv. (1865–66) 67 = velutinosum.
montanum, *Blume, Bijdr.* 1124.—Java.
montana, Thw. Enum. Pl. Zeyl. 289 = pyrifolium.
Moritzii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 67.—Ind. or.; Malaya.
natalense, Harv. Thes. Cap. ii. 45 = venosum.
nervosum, Wall. Cat. n. 7289 = Gronniera nervosa.
neurocarpum, *Miq. Fl. Ind. Bat. Suppl.* 466.—Sumatra.
nigricans, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 225.—Ind. or.
nitidum, Tul. l. c. 193 = leptocladum.
oblongum, *Muell. Arg. in DC. Prod.* xv. ii. 254.—Ind. or.
oblongifolium, *Blume, Bijdr.* 1125.—Java.
ovalifolium, *Zipp. ex Span. in Linnaea*, xv. (1841) 350.—Ins. Timor.
pacificum, *Muell. Arg. in DC. Prod.* xv. ii. 254.—Ins. Pacif.
palembanicum, *Miq. Fl. Ind. Bat. Suppl.* i. 465.—Sumatra.
paniculatum, Roxb. ex Willd. Sp. Pl. iv. 764 = A. Ghesaembilla.
parvifolium, *F. Muell. Fragm.* iv. 86.—Austral.
perusimie, *Kurz, in Journ. Bot.* xiii. (1875) 330.—Ins. Nicobar.
petiolare, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 207.—Madag.
platyphyllum, *H. Mann, in Proc. Am. Acad.* vii. (1867) 208.—Hawaii.
pleuricum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 213.—Ins. Philipp.
poberum, *Zipp. ex Span. in Linnaea*, xv. (1841) 350.—Ins. Timor.
pubescens, Moritzi, Verz. Pfl. Zolling. 73 = Moritzii.
pubescens, Roxb. Pl. Corom. ii. 35. t. 167 = A. Ghesaembilla.
punctiolatum, *Miq. Fl. Ind. Bat. Suppl.* 468.—Sumatra.
pyrifolium, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 68.—Zeylan.
refractum, Muell. Arg. in DC. Prod. xv. ii. 257 = acuminatum.
retusum, *Zipp. ex Span. in Linnaea*, xv. (1841) 350, nomen.—Ins. Timor.
rhamnoides, *Presrgn. ex Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 217.—Hab. ?
Ribes, Rasusch. Nom. ed. III. 287 = Embelia Ribes.
rostratum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 218.—Ins. Philipp.
rotatum, Muell. Arg. in DC. Prod. xv. ii. 256 = cuspidatum.
rotundifolium, *Boj. Hort. Maurit.* 289.—Ins. Mascar.

ANTIDESMA:—
Roxburghii, *Wall. Cat.* n. 7288; *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 234.—Ind. or.
rufescens, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 231 = venosum.
rugosum, *Wall. ex Voigt, Hort. Suburb. Calc.* 295, nomen.—Ind. or.
Rumphii, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 238.—Amboina.
salaccense, Zoll. & Moritzi, Syst. Verz. Zoll. 74 = tetrandrum.
salicifolium, *Miq. Fl. Ind. Bat. Suppl.* i. 467.—Malaya.
salicifolium, Presl, Epim. Bot. 232 = rostratum.
scandens, *Lour. Fl. Cochinch.* ii. 617.—China.
Schultzii, *Benth. Fl. Austral.* vi. 86.—Austral.
simile, Muell. Arg. in DC. Prod. xv. ii. 259 = acuminatum.
sinuatum, *Benth. Fl. Austral.* vi. 87.—Austral.
sphaerocarpum, *Muell. Arg. in DC. Prod.* xv. ii. 255.—Ins. Pacif.
spicatum, Blanco, Fl. Filip. ed. I. 794 = A. Ghesaembilla.
stenopetalum, Muell. Arg. in Flora, xlvii. (1864) 520 = Thecacoris stenopetala.
Stillago, *Poir. Encyc. Suppl.* i. 408.—Hab. ?
stipulare, *Blume, Bijdr.* 1126.—Ins. Moluce.
sylvestre, Lam. Encyc. i. 307 = A. Ghesaembilla et A. Bunius.
sylvestre, Wall. Cat. n. 7281 = diandrum.
tetrandrum, *Blume, Bijdr.* 1124.—Java.
Thwaitesianum, Muell. Arg. in DC. Prod. xv. ii. 263 = A. Bunius.
tomentosum, *Blume, Bijdr.* 1126.—Java.
tomentosum, Blume, ex Miq. Fl. Ind. Bat. i. ii. 427 = velutinosum.
tomentosum, Voigt, Hort. Suburb. Calc. 294; Wight, Ic. t. 767–768 = Roxburghii.
triplinervium, Spreng. Neue Entdeck. ii. 116 = Alchornea triplinervia.
Tulasneanum, *Baill. Etud. Gén. Euph.* 609.—Ins. Mascar. ?
velutinosum, *Blume, Bijdr.* 1128.—Burma; Malaya.
velutinosum, Miq. Fl. Ind. Bat. i. ii. 428 = molle.
velutinum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 228.—Burma.
venosum, *E. Mey. ex Tul. l. c.* 232.—Afr. austr.
vestitum, Presl, Epim. Bot. 232 = A. Ghesaembilla.
Vogelianum, *Muell. Arg. in Flora*, xlvii. (1864) 529.—Afr. trop.
Wallichianum, Presl, Epim. Bot. 235 = diandrum.
zeylanicum, *Lam. Encyc.* i. 207; *Buch.-Ham. in Wall. Cat.* n. 8572.—Zeylan.
zeylanicum, Presl, Epim. Bot. 234 = pleuricum.

ANTIGONA, Vell. Fl. Flum. 186; iv. t. 145 (1827) = **Cassaria**, Jacq. (Samyd.).
serrata, Vell. ll. cc. 187; t. 145 = C. lasiophylla.

ANTIGONON, Endl. Gen. 810 (1837). *POLYGONACEAE*, Benth. & Hook. f. iii. 108.
cinerascens, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 14.—Mexic.
cordatum, *Mart. & Gal. l. c.*—Mexic.
guatemalense, *Meisn. in DC. Prod.* xiv. 134.—Guatemala.
insigne, Mast. in Gard. Chron. (1877) i. 780 = guatemalense.
leptopus, *Hook. & Arn. Bot. Beech. Voy.* 308. t. 69.—Am. austr.
platypus, *Hook. & Arn. l. c.* 309.—Mexic.

ANTINORIA, Parl. Fl. Palerm. i. 92 (1845) = **Aira** Linn. (Gram.).
agrostidea, Parl. l. c. 95 = Aira agrostidea.
insularis, Parl. l. c. 94 = Aira insularis.

ANTIPHYLA, Rafin. New Fl. Am. iv. 16 (1836) = **Malaochia**, Dill. (Stercul.).
serrata, Rafin. l. c. = M. serrata.

ANTIPHYLLA, Haw. Enum. Saxif. 43 (1821) = **Saxifraga**, Linn.
arctica, Haw. l. c. 45 = S. nanscoides.
biflora, Haw. l. c. 44 = S. biflora.
carnosa, Haw. l. c. 43 = S. oppositifolia.
oppositifolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 296 = S. oppositifolia.
retusa, Haw. Enum. Saxif. 44 = S. retusa.
spathulata, Haw. l. c. = S. spathulata.

ANTIPHYTUM, DC. ex Meisn. Gen. 380 (1840)
PORAGINEAE, Benth. & Hook. f. ii. 859.
cruciatum, *DC. Prod.* x. 121.—Bras.
floribundum, A. Gray, in Proc. Am. Acad. x. (1874) 55 = Krynitzkia floribunda.
fruticosum, Moç. ex DC. Prod. x. 121 = mexicanum.
heliotropioides, A. DC. l. c. 122 = Krynitzkia heliotropioides.
linifolium, DC. l. c. = Krynitzkia linifolia.
mexicanum, *DC. l. c.*—Mexic.
Parryi, S. Wats. in Proc. Am. Acad. xviii. (1882-83) 122 = Krynitzkia Parryi.
staechadifolium, *DC. Prod.* x. 121.—Bras
tetrasquetrum, *DC. l. c.* 122.—Bras.
Walpersii, *A. DC. l. c.*—Peruv.

ANTIRRHOEA, Comm. ex Juss. Gen. 204 (1789) (Antirhea). *RUBIACEAE* [Benth. & Hook. f. ii. 100].
GUETTARDELLA, Champ. in Hook Kew Journ. iv. 197 (1852).
PITTONIOTIS, Griseb. in Bonplandia, vi. (1858) 8.
STENOSTOMUM, Gaertn. f. Fruct. iii 69 (1805).
STURMIA, Gaertn. l. c. t. 192 (1805).
aristata, *Benth. & Hook f. Gen.* ii. 100.—Ind. occ.
bifurcata, *Benth. & Hook. l. c.*—Ind occ
borbonica, J. F. Gmel. Syst. i. 344 = Quivisia Sieberi.
dichotoma, *Hemsl. Biol. Centr. Am. Bot* ii. 42.—Mexic.
disca, Bory, ex DC. Prod. iv. 459 = verticillata.
frangulacea, *DC. Prod.* iv. 459.—Ins. Mascar.
lucida, *Benth. & Hook. f. Gen.* ii 100.—Ind. occ
Lastoanna, Comm. ex DC. Prod. iv. 459 = borbonica.
philippinensis, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 313.—Ins. Philipp.
protracta, *Hemsl. Biol. Centr. Am Bot.* ii. 42.—Mexic.
rotundata, *Benth. & Hook. f. Gen.* ii. 100.—Cuba.
strigosa, *Kerth. in Neder. Kruid. Arch.* ii. 11. (1851) 217.—Sumatr.
tenuifolia, *F. Muell. ex Benth. Fl. Austral.* iii. 418.—Austral.
trichantha, *Hemsl. Biol. Centr. Am. Bot.* ii. 42.—Panama.
verticillata, DC. Prod. iv. 459 = borbonica.
viscosa, *Benth. & Hook. f. Gen.* ii. 100.—Ind. occ.

ANTIRRHINUM, Tourn. ex Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 934.
AGORRHINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 126.
ASARINA, Tourn. ex Mill. Gard. Dict. ed. VIII (1768).
GAMBELLIA, Nutt. in Journ. Acad. Phil. Ser. II. i. (1847) 149.
ORONTIUM, Pers. Syn. ii. 158 (1807).
SAIROCARPUS, Nutt. ex A. DC. in DC. Prod. x. 592 (1846).
acutangulum, Tenore, Fl. Nap. ii. 47. t. 59 = Linaria Cymbalaria.
Adamsii, Willd. ex Ledeb. Fl. Ross. iii. 307 = Linaria macroura.
aegyptiacum, *Forsk. Fl. Aegypt. Arab.* 119.—Arab.
aegyptiacum, Linn. Sp. Pl. ed. I. 613 = Linaria aegyptiaca.
aequitriloburn, Viv. Fl. Lib. Spec. 68; Fl. Cors. Sp. Nov. 10 = Linaria aequitriloba.
arrugineum, Gouan, Illustr. 88 = Linaria triatis.
albaceum, Medic. Beobachtt. (1782) 136 = Linaria aegyptiaca.
albifrons, Sibth. & Sm. Prod. i. 482 = Linaria albifrons.
alsum, Lam. Fl. Fr. ii. 345 = Linaria chalepensis.
alpinum, Linn. Sp. Pl. 615 = Linaria alpina.
alsinaefolium, Viv. Fl. Lib. Spec. 67; Fl. Cors. Sp. Nov. 10 = Linaria alsinaefolia.
ambiguum, *Lange, in Kjoeb. Vidensk. Meddel.* (1863) 27 (*Pugill.* 208).—Hispan.
amethysteum, Vent. in Lam. Encyc. iv. 353 = Linaria amethystea.
amethystinum, Steud. Nom. ed. II. i. 108 = Linaria amethystea.
angustifolium, Poir. Encyc. Suppl. iv. 27 = sicnlum.
angustifolium, Urv. in Mém. Soc. Linn. Par. i. (1822) 339 = tortuosum !
angustissimum, Loisel. Not. Suppl. 167 = Linaria Italica.
aparinoides, Willd. Sp. Pl. iii. 247 = Linaria stricta.
aphyllum, Thunb. Prod. Pl. Cap. 105 = Utricularia capensis.
aquaticum, *Lour. Fl. Cochinch.* ii. 384 (Utricularieae sp. ?).—Cochinch.

ANTIRRHINUM:—
arenarium, Poir. Encyc. Suppl. iv. 26 = Linaria arenaria.
arvense, Linn. Sp. Pl. 614 = Linaria arvensis.
asarifolium, Salisb. Prod. 98 = scq.
Asarina, *Linn. Sp. Pl.* 612.—Italia.
asarinum, Lam. Fl. Fr. ii. 348 = A. Orontium.
azungera, Blanco, ex Cavel. in Farl. Fl. Ital. vi. 656 = siculum.
auriculatum, Linn. Fl. Fr. ii. 339 = Linaria Elatine barbatum, *Thunb. Prod. Pl. Cap.* 105.—Afr. austr.
Barrelieri, *Bor. Graenes Rec. Yard. Bot. Angers* (1854) = Hispan.
Bauhinni, Gaud. Fl. Helv. iii. 154 = Linaria italica.
bellidifolium, Linn. Sp. Pl. 617 = Anarrhinum bellidifolium.
bicorne, Linn. Pl. Rar Afr 11 (Amoen Acad. vi. 88) = Nemesia bicornis.
bipartitum, Vent. Jard. Cels, t 82 = Linaria bipartita.
bipunctatum, Cav. Ic i t. 33. f. 1 = Linaria amethystea.
bipunctatum, Linn. Sp. Pl. ed. I. 614 = Linaria glauca.
bipunctatum, Thuill. Fl. Paris, ed. II. 311 = Linaria supina.
brevifiorum, Gilib. Fl. Lituan i. 187 = A. Orontium.
Brewerii, *A. Gray, in Proc. Am. Acad.* vii (1868) 374 —Calif.
Broussonetii, Poir. Encyc Suppl. iv 28 = Linaria multipunctata.
caesium, Lag. ex Pers. Syn. ii. 157 = Linaria tristis ?
calycinum, [*Soland. in*] *Russ. Alepp*o, ed. II. ii. 256.—Syria.
calycinum, Vent. in Lam. Encyc. iv. 365 = A. Orontium.
canadense, Linn. Sp. Pl 618 = Linaria canadensis.
capense, *Burm f. Fl. Cap. Prod.* 16.—Afr. austr.
capense, Thunb. Prod. Pl. Cap 105 = Nemesia inearia.
capitatum, C. & J. Presl, Delic. Prag 74 = sempervirens.
capitellatum, Vent in Lam. Encyc. iv. 353 = Linaria glauca.
caprarium, Cavel, in Parl. Fl. Ital. vi. 862 = Linaria capraria.
caucasicum, Willd. ex Ledeb Fl. Ross. iii. 205 = Linaria spuria.
chalepense, Linn. Sp. Pl. 617 = Linaria chalepensis.
Charidemi, *Lange, in Kjoeb. Vidensk. Meddel* (1881) 99.—Lusit.
chytrosperum, *A. Gray, in Proc. Am. Acad.* xii. (1877) 81.—Arizona.
cimbalaria, Neck. Delic. Gallo-Belg. *J.* 267 = Linaria Cymbalaria.
cirrhosum, Linn. Mant. ii. 249 = Linaria cirrhosa.
comintanum, Blanco, Fl Filip. ed. I. 502 = Hygrophila salicifolia.
commune, Lam. Fl. Fr. ii. 340 = Linaria vulgaris.
compositum, *Loyac. in Natural. Sicil.* ii. (1883) 297.—Sicil.
confertiflorum, Benth. in DC. Prod. x. 592 = Mohavea viscida.
corsicum, *Vitm. Summa*, iii. 495.—Corsica.
Coulterianum, *Benth. in DC. Prod.* x. 592.—Calif.
craniolare, Stokes, Bot. Mat. Med. iii. 383 = A. Orontium.
crassifolium, Cav. Ic. ii. 11 = Linaria origanifolia.
cyathiferum, *Benth. Bot. Voy. Sulph.* 40. t. 19.—Calif.
Cymbalaria, Gmelidenst. ex Ledeb. Fl. Ross. iii. 205 = Linaria Elatine.
Cymbalaria, Linn. Sp. Pl. ed. I. 612 = Linaria Cymbalaria.
Cymbalaria, Sieber, ex Benth. in DC. Prod. x. 267 = Linaria scutangula.
dalmaticum, Guaidenst. ex Ledeb. Fl. Ross. iii. 209 = Linaria genistifolia.
dalmaticum, Linn. Sp. Pl. ed. I. 616 = Linaria dalmatica.
delawarense, *Rosh. Hort. Beng.* 45, nomen.—Am. bor. ?
denticulatum, Poir. Encyc. Suppl. iv. 28 = Nemesia monomotapensis.

ANTIRRHINUM:—
dentatum, Vahl, Symb. Bot. ii. 66 = Linaria scariosa.
diffusum, Bernh ex Steud. Nom. ed. II. i. 108 = latifolium.
diffusum, Brot. Phyt. t. 141 = Linaria multicaulis.
diminutum, *Pomel, Nouv. Mat. Fl. Atl.* 100.—Algeria.
dubium, Vill. Hist. Pl. Dauph ii. 487 = Linaria supina
Duriminium, Brot. Fl. Lusit i. 197 = Anarrhinum Duriminium.
ecaudatum, *Lucé, Topog. Nachr. Orzel*, 218.—Ross.
Elatine, Linn Sp. Pl. 612 = Linaria Elatine.
elatinifolium, Wahlenb. in Isis, xxi. 989 = Linaria Elatine.
elatinoides, Tenore, Syll 48. f. 2 = Linaria Elatine.
elatinoides, Willd. Sp. Pl. iii. 235 = Linaria elatinoides
elegans, Pers. Syn. ii. 156 = Linaria amethystea.
elegans, Tenore, Vag. Abruz. 175 = A. Orontium.
filiforme, Poir. Encyc Suppl. iv. 27 = Linaria rubrifolia.
fiipes. *A. Gray, in Bot. Jnes, Report*, 19.—Colorado.
flavum, Poir. Voy. Barb. ii 191 = Linaria flava.
flexuosum, *Pomel, Nouv. Mat. Fl. Atl.* 100.—Algeria
flexuosum, Willd. Sp. Pl. iii. 250 = Linaria flexuosa.
fragrans, *Pall. Reise Statth. Russ. Reich.* i. 111, 130.—Reg Casp.
fruticans, Thunb. Prod. Pl Cap. 105 = Nemesia fruticans.
fruticosum, Willd Sp. Pl. iii. 237 = Linaria fruticosa.
fuqux, Boiss. Diagn. ser II iii. 160 = Linaria persica.
galliotdes, Vent. in Lam. Encyc. iv. 351 = Linaria striata.
genistaefolium, ' Vill ' ex Benth. in DC. Prod. x. 272 = Linaria italica.
genistifolium, Lapeyr. Hist. Abr. Pl. Pyr. 354 = Linaria vulgaris.
genistifolium, Linn. Sp. Pl 616 = Linaria genistifolia.
genistifolium, Vill. Hist. Pl. Dauph. ii. 440 = Linaria italica.
gibbosum, Wall Pl. As Rar. ii. 44 = A. Orontium.
glandulosum, Lejeune, Fl. Spa, 320 = Linaria vulgaris.
glandulosum, *Lindl. Bot. Reg.* t. 1893.—Calif.
glaucophyllum, Brot. Phyt. Lusit. 188 = Linaria glaucophylla.
glaucum, Lapeyr. Hist. Abr. Pl. Pyr. 352 = Linaria supina.
glaucum, Linn. Cent. I. n. 48; in Amoen. Ac. iv. 277 = Linaria glauca.
glaurum, Stocks, in Wight, Ic. iv. t. 1459 = Schweinfurthia sphaerocarpa.
glutinosum, Thore, Chl. Land. 285 = Linaria thymifolia.
glutinosum, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 82. —Hispan.
gracile, Pers. Syn. ii. 156 = Linaria Pelissertana.
gracile, Rent. Ind. Sem. Hort. Genev. (1853) 4; ex Lange, in Willk & Lange, Prod. Fl. Hisp. ii. 582 = A. Orontium.
gracum, Borv & Chaub. Fl. Pélop. 175. t. 21 = Linaria graeca.
grandiflorum, Stokes, Bot. Mat. Med. iii. 384 = majus.
gratissimum, Koxb. ex Hook. f. Fl. Brit. Ind. iv. 266 = Limnophila Roxburghii.
Haelava, Forsk. Fl. Aegypt. Arab. 111 = Linaria Haelava.
heleracum, Lam. Fl. Fr. ii. 338 = Linaria Cymbalaria.
hederaefolium, Salisb. Prod. 98 = Linaria Cymbalaria.
hepaticifolium, Poir. Encyc. Suppl. iv. 19 = Linaria hepaticaefolia.
heterophyllum, Schousb. Vextr. Marokko, 194; Willd. Sp. Pl. iii. 234 = Linaria heterophylla.
hexandrum, Forsk. Fl. Aegypt. Arab. 43 = Vandellia crustacea.
hirtum, Linn. Sp. Pl. ed. I. 616 = Linaria hirta.
hispanicum, *Chav. Monog.* 38.—Hispan.
Huetii, Rent. in Ann. Sc. Nat. Sér. IV. ii. (1854) 330 = sempervirens.
humile, Salisb. Prod. 98 = A. Orontium.
hybridum, *Benth. Cat. Pl. Pyren.* 60.—Mont. Pyren.
incarnatum, Vent. in Lam. Encyc. iv. 364 = Linaria bipartita.
indicum, Royle, ex Ball, in journ. Linn. Soc. xvi. (1878) 295 = A. Orontium.
intermedium, Debeaux, in Bull. Soc. Bot. Fr. xx. (1873) 183 = latifolium.
intermedium, Hort. ex Steud. Nom. ed. II. i. 108 = striatum.

ANTIRRHINUM:—
Intrictum, *Ball, in Journ. Bot.* xiii. (1875) 178.—Marocco.
tisalicum, Mill. Gard. Dict. ed. VIII. n. 5 = majus.
jamaicense, Fisch. ex Steud. Nom. ed. II. i. 108 = siculum.
junceum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 377.—Calif.
juncreum, Linn. Cent. i.; Amoen. Acad. iv. 277; Vent. in Lam. Encyc. iv. 362 = Linaria spartea.
juncreum, Pall. Reise, ii. 733 = Linaria Loeselii.
Kallogrii, *Greene, in Torrey Bot. Club*, x. (1883) 126.—Calif.
Klugii, *S. Wats. in Bot. King, Exp.* 215. t. 21.—Utah.
laevigatum, Herb. Banks, ex Benth. in DC. Prod. x. 255 = Dimeta parviflora.
lagopodioides, Linn. f. Suppl. 279 = Linaria saxatilis.
lanigerum, Willd. Sp. Pl. iii. 333 = Linaria lanigera.
latifolium, Bourg. ex Willk. & Lange, Prod. ii. 584 = hispanicum.
latifolium, *Mill. Gard. Dict.* ed. VIII. n. 4; *DC. Fl. Fr.* v. 411.—Europe.
latifolium, Willd. Sp. Pl. iii. 338 = Linaria latifolia.
laxiflorum, Willd. l. c. 342 = Linaria laxiflora.
lepaleum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 378.—Calif.
Linaria, Linn. Sp. Pl. 616 = Linaria vulgaris.
linarioides, Linn. l. c. ed. II. 853 = Linaria vulgaris.
linifolium, Linn. l. c. 858 = Linaria linifolia.
linifolium, Pall. Ind. Taur. 101 = Linaria macroura.
Linkianum, Boiss. & Reut. Diagn. Ser. II. iii. 160 = latifolium.
linogriseum, Brot. Phyt. 135 = Linaria bipartita.
littorale, Bernh. in DC. Fl. Fr. v. 410; Pair. Encyc. Suppl. iv. 20 = Linaria littoralis.
longiflorum, Thunb. Prod. Pl. Cap. 105 = Diascia Thunbergiana.
lusitanicum, Vent. in Lam. Encyc. iv. 361 = Linaria tristis.
lusitanicum, Link, ex Steud. Nom. ed. II. i. 109 = Linaria pedunculata.
macrocarpum, Soland. in Aft. Hort. Kew. ii. 335 = Nemesia chamaedrifolia.
macrourum, Bieb. Fl. Taur. Cauc. ii. 75 = Linaria macroura.
maculatum, Moq. & Sesse, ex DC. Prod. vii. 534 = Achimenes foliosa.
majus, *Linn. Sp. Pl.* 617.—Reg. Mediterr.
marginatum, Desf. in Act. Soc. Hist. Nat. Par. i. (1794) 84. t. 7 = Linaria tristis.
maritimum, Poir. Encyc. Suppl. iv. 23 = Linaria supina.
massilience, Poir. ex Steud. Nom. ed. II. i. 67, 483 = Linaria maritima.
maurandioides, *A. Gray, in Proc. Am. Acad.* vi. (1866) 376.—Texas.
medium, Hort. ex Steud. Nom. ed. II. i. 109 = striatum.
meionanthum, Brot. Phyt. t. 196 = sempervirens.
meonanthum, Hoffmgg. & Link, Fl. Port. i. 261. t. 51 = sempervirens.
micranthum, Cav. Ic. i. 51 = Linaria parviflora.
microcarpum, *Pomel, Nouv. Mat. Fl. Alt.* 99.—Algeria.
minus, Linn. Sp. Pl. 617 = Linaria viscida.
molle, Blanco, Fl. Filip. ed. I. 608 = Hygrophila undulata.
molle, Boiss. Voy. Espagne, 449 = hispanicum.
molle, *Linn. Sp. Pl.* 1168.—Mont. Pyren.
molle, St. Amans, Bouq. Pyren. n. 154 = sempervirens.
monspessulanum, Georgi, Beschr. Russ. Reich. iii. iv. 1136 = Linaria Loeselii.
monspessulanum, Linn. Sp. Pl. 616 = Linaria striata.
montevidense, Mart. ex Colla, in Mem. Acc. Torino, xxxv. (1831) 178 = majus.
multibractestum, Brot. ex Steud. Nom. ed. II. i. 109 = variabile.
multicaule, Linn. Sp. Pl. 615 = Linaria multicaulis.
multicaule, Tenore, Fl. Nap. Prod. 86 = Linaria stricta.
multipunctatum, Brot. Fl. Lusit. i. 196 = Linaria amethystea.
murale, Sallsb. Prod. 98 = majus.
muralis, Pers. Syn. ii. 155 = Linaria Cymbalaria.
myplocium, Spreng. Neue Entdeck. iii. 169 = Linaria triphylla.
nicaeense, Risso, Fl. Nice, 347 = latifolium.
Nevinianum, *A. Gray, in Coult. Bot. Gazette*, ix. (1884) 84.—Calif.

ANTIRRHINUM:—
Nuttallianum, *Benth. in DC. Prod.* x. 592.—Calif.
ochroleucum, Salisb. Prod. 98 = Linaria vulgaris.
ocimifolium, Poirr. ex Sprong. Syst. ii. 738 = Linaria ocimifolia.
odoratissimum, Gaudinet. ex Ledeb. Fl. Ross. iii. 308 = Linaria Loeselii.
odorum, Bieb. Fl. Taur. Cauc. ii. 75 = Linaria Loeselii.
oppositiflorum, Linn. Encyc. Suppl. iv. 23 = Linaria villosa.
orbiculare, *Medic. Beobachtt.* (1788) 242.—Hab. ?
orchidiflorum, [Desf.] ex Pers. Syn. ii. 155 = Linaria bipartita.
Orcuttianum, *A. Gray, in Coult. Bot. Gazette*, ix. (1884) 53.—Calif.
origanifolium, Gouan, Hort. Monsp. 301 = Linaria rubrifolia.
origanifolium, Linn. Sp. Pl. 615 = Linaria origanifolia.
Orontium, *Linn. Sp. Pl.* 617.—Europ. : As. Min.; Afr. bor.
ottomanum, *Janka, in Termész. Füzetek*, iv. (1880) 297.—Turcia.
Oyyris, Cyrill. Pl. Neap. Nov. Fasc. ii. 21 = Linaria chalepensis.
pallidiflorum, Lam. Fl. Fr. ii. 341 = Linaria genistifolia.
pallidum, Tenore, Fl. Nap. i. 328 = Linaria pilosa.
paphlagonicum, Boiss. Mant. 86 = Schweinfurthia sphaerocarpa.
parviflorum, Jacq. Coll. iv. 304 = Linaria arvensis.
parviflorum, Willd Sp. Pl. iii. 345 = Linaria parviflora.
patens, Thunb. Prod. Pl. Cap. 105 = Nemesia patens.
pauciflorum, Stend. Nom. ed. II. i. 109 = Linaria multicaulis.
pauciflorum, Poepp. ex Chav. Monog. 149 = Linaria canadensis.
pedunculatum, Linn. Sp. Pl. ed. II. 857 = Linaria pedunculata.
Pelisserianum, Linn. l. c. ed. I. 615 = Linaria Pelisseriana.
Pelisserianum, Mout.-Font. ex Chav. Monog. 155 = Linaria gracilis.
pilosum, Bory, Fl. Pélop. 89 = Linaria microcalyx.
pilosum, Jacq. Obs. Bot. ii. 28 = Linaria pilosa.
pinifolium, Poir. Voy. Barb. ii. 198 = Linaria reticulata.
pinnatum, Linn. f. Suppl. 280 = Nemesia pinnata.
polygalaefolium, Brot. Phyt. t. 138 = Linaria supina.
polygalifolium, Poir. Encyc. Suppl. iv. 21 = Linaria italica.
polygalaefolium, Willd. ex Steud. Nom. ed. II. i. 109 = Linaria tristis.
polygamiflorum, Poir. Encyc. Suppl. iv. 26 = Linaria amethystea.
porcinum, *Lour. Fl. Cochinch.* ii. 384.—Cochinch.
praecox, Brot. Phyt. t. 131 = Linaria spartea.
prostratum, Cyrill. ex Chav. Monog. 120 = Linaria reflexa.
ptarmicum, A. Rich. Tent. Fl. Abyss. ii. 115 = Schweinfurthia pterosperma.
pubescens, Pers. Syn. ii. 157 = Linaria arenaria.
pubescens, Tenore, Fl. Nap. Prod. 86 = Linaria pallida.
pumilum, Poir. ex Steud. Nom. ed. II. i. 109 = Linaria supina.
purpureum, Duiac. Fl. Hautes-Pyr. 374 = A. Orontium.
purpureum, Linn. Sp. Pl. ed. I. 613 = Linaria purpurea.
pygmaeum, *Gouan, Illustr.* 37.—Europ. austr.
pyramidale, Vent. in Lam. Encyc. iv. 360 = Linaria pyramidata.
pyramidatum, Benth. in DC. Prod. x. 272 = praec.
pyrenaicum, Ram. ex DC. Fl. Fr. iii. 581 = Linaria supina.
quadrifidum, *G. Shaw, Cimel. Phys.* 84. t. 44.—Hab. ?
quinquelobatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 110 = A. Asarina.
quitoense, Stokes, Bot. Mat. Med. iii. 385 = Linaria Cymbalaria.
ramosissimum, *Cav. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 384.—Algeria.
reflexum, Linn. Syst. ed. X. 1113 = Linaria reflexa.
repens, Linn. Sp. Pl. 614 = Linaria striata.
reticulatum, Sm. ic. Pict. (3) = Linaria reticulata.
Rhodium, Boiss. ex Benth. in DC. Prod. x. 291 = majus.
rhytidospermum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 27 = Linaria persica.
roemeriana, Seb. & Maur. ex Nym. Consp. 386 = tortuosum.
rosmarinifolium, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Malaya.

ANTIRRHINUM:—
rotundifolium, Gilib. Fl. Lituan. i. 138 = Linaria spuria.
rubrifolium, Poir. Encyc. Suppl. v. 737 = Linaria rubrifolia.
rupestre, Boiss. & Reut. Pugill. Pl. Nov. (1852) 82 = glutinosum.
rupestre, Bourg. ex Willk. & Lange, Prod. ii. 585 = hispanicum.
ruscinonense, Debeaux, in Bull. Soc. Bot. Fr. xx. (1873) 14 = Barrelieri.
rybidocarpum, G. Don, Gen. Syst. iv. 516 = Linaria persica.
rytidocarpum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 27 = Linaria persica.
sagittatum, Poir. Encyc. Suppl. iv. 19 = Linaria sagittata.
saphyrinum, Brot. Fl. Lusit. i. 197 = Linaria amethystea.
saphyrinum, Sieber, ex Stend. Nom. ed. II. i. 109 = Linaria Pelisseriana.
sanatile, Bonani, Nann. Prod. 69 = Linaria arenaria.
sanatile, Linn. Cent. Pl. i. 16 = Linaria saxatilis.
scabrum, Thunb. Prod. Pl. Cap. 105 = Nemesia chamaedrifolia.
scariosum, Vent. in Lam. Encyc. iv. 349 = Linaria scariosa.
sempervirens, *Lapeyr. Fl. Pyr.* i. 7. t. 4; *Hist. Abr. Pl. Pyr.* 854.—Mont. Pyren.
serpyllifolium, Poirr. ex Chav. Monog. 98 = Linaria villosa.
siculum, *Mill. Gard. Dict.* ed. VIII. n. 6; *Ucria, in Nouv. Arch.* i. 69.—Sicilia.
simplex, *Link, in Schrad. Neues Journ.* ii. (1799) 300. Gallia.
simplex, Willd. Sp. Pl. iii. 343 = Linaria arvensis.
sparteum, Linn. Sp. Pl. ed. I. 1197 = Linaria spartea.
sparteum, Viviani, Fl. Lib. Spec. 38 = Linaria viscosa.
spartum, Linn. Cent. i. 17, postea *sparteum*.
speciosum, Dunn, Hort. Lect. ed. IV. 139 = Linaria bipartita.
speciosum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 376.—Calif.
spinescens, Viviani, Fl. Lib. Spec. 32. t. 27. f. 2 = Linaria aegyptiaca.
spurium, Brot. ex Steud. Nom. ed. II. i. 109 = Linaria reticulata.
spurium, Linn. Sp. Pl. 618 = Linaria spuria.
striatum, Lam. Fl. Fr. ii. 343 = Linaria striata.
striatum, *Schrank, ex Steud. Nom.* ed. I. 57.—Europ.
strictum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 375.—Calif.
strictum, Sibth. & Sm. Fl. Graec. vi. 75. t. 594 = Linaria Sibthorpiana.
strumosum, Banks, ex Benth. in DC. Prod. x. 260 = Nemesia strumosa.
subalpinum, Brot. Fl. Lusit. i. 196 = Linaria alpina.
subcordatum, *A. Gray, in Proc. Am. Acad.* xx. (1885) 306.—Calif.
subsessile, *A. Gray, in Coult. Bot. Gazette*, ix. (1884) 58.—Calif.
tubulatum, Cyrill. ex Chav. Monog. 136 = Linaria Loeselii.
supinum, Brot. Fl. Lusit. i. 191 = Linaria reticulata.
supinum, Linn. Sp. Pl. 615 = Linaria supina.
supinum, Pall. Reise, ii. 733 = Linaria Loeselii.
supinum, Sibth. & Sm. Fl. Graec. vi. 75. t. 595 = Linaria quadrifolia.
tenuifolium, Cav. Ic. i. 61. t. 80. f. 1 = Linaria tenella.
tenue, Viviani, Fl. Lib. Spec. 38. t. 16. f. 5-6; Spreng. Syst. ii. 695 = Linaria viscosa.
Thuillieri, Poir. Encyc. Suppl. iv. 26 = Linaria supina.
thymifolium, Vahl, Symb. Bot. ii. 67 = Linaria thymifolia.
tortuosum, *Boc. ex Lam. Encyc.* iv. 365.—Reg. Mediterr. occ.
Tournefortii, Poir. Encyc. Suppl. iv. 24 = Linaria Tournefortii.
triandr, Hort. ex Chav. Monog. 116 = Linaria triphylla.
tricolor, Poirr. ex Chav. l. c. 172 = Linaria amethystea.
triornithophorum, Linn. Sp. Pl. 613 = Linaria triornithophora.
triphyllum, Cav. Ic. ii. 61. t. 179 = Linaria Cavanillesii.
triphyllum, Linn. Sp. Pl. 613 = Linaria triphylla.
tristis, Linn. l. c. = Linaria tristis.
uniflorum, Moç. & Sesse, ex DC. Prod. vii. 532 = Gemeria uniflora.

ANTIRRHINUM :—
unilabiatum, Linn. f. Suppl. 279 = Diascia unilabiata.
vagans, *A. Gray, in Proc. Am. Acad.* vii. (1868) 376. —Calif.
variabile, *Link, ex Steud. Nom.* ed. I. 57.—Lusit.
vermiculor, Linn. Sp. Pl. 866 = Linaria vermiculor.
vesillo-calyculatum, *Kellogg, in Proc. Calif. Acad.* I. ed. II. (1873) 7, 26.—Calif
villosum, Lap. Fl. Pyren. 259 = Linaria origanifolia.
villosum, Linn. Sp. Pl. ed. II. 852 = Linaria villosa.
violaceum, Salisb. Prod. 87 = Linaria Pelisseriana.
Virga, *A. Gray, in Proc. Am. Acad.* vii. (1868) 378 — Calif.
virgatum, Brot. Fl. Lusit. i. 198 = Linaria spartea.
virgatum, Poir. Voy. Barb. ii. 192 = Linaria virgata.
viscosum, Curt. Bot. Mag. t. 368 = Linaria hirta.
viscosum, Linn. Cent. ii. 21 = Linaria viscosa.

ANTISOLA, Rafin. Sylva Tellur. 97 (1838) = **Miconia**, Ruiz & Pav. (Melast.).
racemosa, Rafin. l. c. = M. racemosa.

ANTISTHIRIA, Pers. Syst. 127 (1797), err. typ. = **Anthistiria**, Linn. (Gramin.).

ANTISTROPHE, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 84. *MYRSINEAE*, Benth. & Hook. f. ii. 647.
oxyantha, *A. DC. l. c.*—Reg. Himal.
serratifolia, *Hook. f. in Benth. & Hook. f. Gen.* ii. 647. —Ind. or.

ANTITAXIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 37 ; Ser. III. xx. (1867) 18. *MENISPERMACEAE*, Benth. & Hook. *f.* Gen. i. 33, 959, iii. 256.
calocarpa, *Kurz, in Journ. Bot.* xiii. (1875) 324.—Ins. Nicob.
cauliflora, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xx. (1867) 14.—Java.
fasciculata, *Miers, l. c.*—Malaya.
longifolia, *Miers, l. c.*—Ins. Timor.
lucida, *Miers, l. c.*—Java.

ANTITHRIXIA, DC. Prod. vi. 277 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 329.
abyssinica, *Benth. & Hook. f. Gen.* ii. 329.—Afr. trop.
angustifolia, *Oliver & Hiern, in Oliver, Fl. Afr. Trop.* iii. 356.—Afr. trop.
flavicoma, *DC. Prod.* vi. 278.—Afr. austr.

ANTITRAGUS, Gaertn. Fruct. ii. 7. t. 80 (1791) = **Crypsis**, Ait. (Gram.).
aculeatus, Gaertn. l. c. = C. aculeata.

ANTIZOMA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 41 ; Ser. III. xvii. (1866) 366 = **Cissampelos**, Linn. (Menisp.).
angustifolia, Miers, ex Harv. & Sond. Fl. Cap. i. 11 = C. angustifolia.
Burchelliana, Miers, l. c. = C. calcarifera.
calcarifera, Miers, l. c. 13 = C. calcarifera.
Harveyana, Miers, l. c. = C. angustifolia.
lycioides, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 366.—Afr. austr.
Miersiana, Harv. in Harv. & Sond. Fl. Cap. i. 18.—Afr. austr.

ANTOCHORTUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 451 (1836) = **Willdenowia**, Thunb. (Restiac.).
Schlechtii, Nees, l. c. = W. striata !

ANTODON, Neck. Elem. i. 56 (1790) = **Leontodon**, *Linn.* (Compos.).

ANTOMMARCHIA, Colla, Hort. Ripul. App. ii. 345 (1826) = **Correa**, Sm. (Rutac.).
rubra, Colla, l. c. 25. t. 72 = C. speciosa.
speciosa, Schlecht. in Linnaea, iv. (1829) Litt. 56 = C. speciosa.

ANTONIA, R. Br. in Wall. Pl. As. Rar. iii. 65 (1832) = Rhynchoglossum, Blume (Gesner.).
obliqua, R. Br. l. c. = R. obliquum.

ANTONIA, Pohl, Pl. Bras. Ic. ii. 13. t. 169 (1831). *LOGANIACEAE*, Benth. & Hook. f. ii. 796.

ANTONIA :—
Griffithii, Wight, Illustr. Ind. Bot. ii. t. 156 b = Norisia malaccensis.
ovata, *Pohl, Pl. Bras.* ii. 13. t. 109.—Bras.
pilosa, Hook. Ic. Pl. t. 64 = ovata.
pubescens, Bong. in Mém. Acad. Pétersb. Sér. VI. iii. (1835) 2. t. 1 = ovata.

ANTONIANA, Tuss. Fl. Antill. 2. t. 16 (1818) ex Grieb. Fl. Brit. W. Ind. 338 = **Faramea**, Aubl. (Rubiac.).
laurifolia, Tuss. l. c. = F. odoratissima.

ANTOPETITIA, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 261 = **Ornithopus**, Linn. (Legum.).
abyssinica, A. Rich. l. c. 262 = O. coriandrinus.
cancellata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 210 = Helminthocarpon abyssinicum.

ANTOPHYLAX, Poir. in Dict. Sc. Nat. ii. Suppl. 95 (1816), err. typ. = ANDROPHYLAX, Wendl. = **Cocculus**, DC. (Menisp.).

ANTOSCHMIDTIA, Steud. Syn. Gram. 199 in syn. Schmidtiae (1855) = **Schmidtia**, Steud. (Gram.).
quinqueseta, Boiss. Fl. Orient. v. 559 = S. quinqueseta.

ANTRIBA, Rafin. Sylva Tellur. 126 (1838) = **Loranthus**, Linn.
budleoides, Rafin. l. c. = L. Scurrula.

ANTROLEPIS, Welw. Apont. 578 (1858) = **Ascolepis**, Nees (Cyper.).
anthemiflora, Welw. l. c. = A. anthemiflora.
ciata, Welw. l. c. = A. ciata.
leucantha, Welw. l. c. = A. protea.
santolina, Welw. l. c. = A. protea.
sulphurea, Welw. l. c. = A. protea.

ANTROSPERMUM, Sch. Bip. in Flora, xxvii. (1844) 778 = **Venidium**, Less. (Compos.).
floribundum, Passer. Ind. Sem. Parm. (1859).—Hab.!
Krausii, Sch. Bip. in Flora, xxvii. (1844) 778 = V. Krausii.

ANTSCHAR, Horsf. in Verh. Bat. Gen. vii. (1814) 8 = **Antiaris**, Lesch. (Urtic.).

ANTURA, Forsk. Fl. Aegypt. Arab. 68 (1775) = **Carissa**, Linn. (Apocyn.).
hadiensis, J. F. Gmel. Syst. 495 = C. edulis.

ANUBIAS, Schott, in Oestr. Bot. Wochenbl. (1857) 398. *AROIDEAE*, Benth. & Hook. f. iii. 975.
Afzelii, *Schott, l. c.* 399.—Afr. occ.
Barteri, *Schott, Prod. Aroid.* 159.—Afr. occ.
heterophylla, *Engl. in DC. Monog. Phan.* ii. 485.—Afr. trop.

ANURUS, Presl, in Weitenw. Beitr. ii. (1837) 24 = **Lathyrus**, Linn. (Legum.).
linifolius, Presl, l. c. = L. Nissolia.
Nissolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 366 = L. Nissolia.

ANVILLEA, DC. Prod. v. 487 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 338.
SYCODIUM, Pomel, Nouv. Fl. Atl. 39 (1874).
Garcini, *DC. Prod.* v. 487.—Aegypt. ; Persia.
radiata, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 471, *nomen.*—Algeria.

ANTOPHIA, Michx. Fl. Bor. Am. i. 119 (1803). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 16.
ANTHALIS, Rafin. New Fl. Am. iv. 40 (1836).
Argyrocoma, Michx. Fl. Bor. Am. i. 114 = Paronychia Argyrocoma.
Baldwinii, Torr. & Gray, Fl. N. Am. i. 172 = Paronychia Baldwinii.
ossadensis, *Ell. Sketch.* i. 307.—Am. bor.
capillacea, *DC. Prod.* iii. 369.—Am. bor.
conferta, *Rafin. Atl. Journ.* (1832) 16.—Am. bor.
dichotoma, *Michx. Fl. Bor. Am.* i. 113.—Am. bor.
divaricata, *Rafin. New Fl. Am.* iv. 43.—Am. bor.
fastigiata, *Rafin. Atl. Journ.* (1832) 16.—Am. bor.
Hermiaria, Pers. Syn. I. 261 = Illecebrum hermiaroides.
hermiarioides, Ell. Sketch. i. 306 = Hermiaria americana.
hermiarioides, Michx. Fl. Bor. Am. i. 113 = Paronychia hermiaroides.

ANYCHIA :—
lateralis, *Rafin. Atl. Journ.* (1832) 16.—Am. bor.
nudiflora, *Rafin. New Fl. Am.* iv. 43.—Am. bor.
polygonoides, *Rafin. Atl. Journ.* (1832) 16.—Am. bor.

AOPLA, Lindl. Bot. Reg. sub t. 1701 (1835) = **Habenaria**, Linn.
reniformis, Lindl. l. c. = H. reniformis.

AOTUS, Sm. in Kon. & Sims, Ann. Bot. i. (1805) 504. *LEGUMINOSAE*, Benth. & Hook. f. i. 470.
carinata, *Meissn. in Lehm. Pl. Preiss.* ii. 215.—Austral.
cordifolia, *Benth. in Enum. Pl. Hueg.* 96.—Austral.
cordifolia, Lindl. & Paxt. Fl. Gard. i. 76 = Vultenaea ternata.
dilwynioides, Meisan. in Lehm. Pl. Preiss. i. 60 = Dillwynia floribanda.
Drummondii, *T. Moore, in Gard. Comp.* (1852) 97.—Austral.
ericoides, G. Don, Gen. Syst. ii. 120 = villosa.
ferruginea, Labill. Pl. Nov. Holl. i. t. 182 = villosa.
genistoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 1. 268.—Austral.
gracilis, Loud. Hort. Brit. Suppl. ii. 610 = gracillima.
gracillima, *Meissn. in Lehm. Pl. Preiss.* i. 59.—Austral.
intermedia, Meisan. l. c. 60 = gracillimas.
lanigera, *A. Cunn. ex Benth. in Ann. Wien. Mus.* ii. (1838) 78.—Austral.
mollis, *Benth. in Mitch. Journ. Trop. Austral.* 236.—Austral.
passerinoides, *Meissn. in Lehm. Pl. Preiss.* i. 61.—Austral.
phyllicoides, *F. Muell. ex Benth. Fl. Austral.* ii. 93.—Austral.
Preissii, *Meissn. in Lehm. Pl. Preiss.* ii. 214.—Austral.
procumbens, Meisan. l. c. i. 60 = Preissii.
Tietkensii, *F. Muell. Fragm.* x. 88.—Austral.
villosa, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 504. —Austral.
virgata, *DC. Prod.* ii. 108.—Austral.
Wuenthii, Regel, in Bot. Zeit. ix. (1851) 596 = Dillwynia floribanda.

APABUTA, Grisch. Ges. Abhandl. 280 (1880), nomen. *MENISPERMACEAE.*

APACTIS, Thunb. Nov. Gen. 66 (1783) = **Roydsia**, Roxb. (Capparid.).
japonica, Thunb. l. c.—Japon.

APALANTHE, Planch. in Ann. & Mag. Nat. Hist. Ser. II. i. (1848) 87 = **Blodea**, Michx. (Hydroch.).
granatensis, Planch. l. c. = E. granatensis.
guyanensis, Planch. l. c. = E. guyanensis.
Schweinitzii, Planch. l. c. = E. canadensis.

APALANTUS, 'Adans. Fam. ii. p. (19)' fide Pfeiff. Nom. i. 925 (1870) = **Hapalanthus**, Jacq.

APALATOA, Aubl. Pl. Gui. i. 883. t. 347 (1775) = **Crudia**, Schreb. (Legumin.).
aromatica, Lam. Encyc. iv. 559 = C. aromatica.
bantamensis, Baill Hist. Pl. ii. 104.—Java.
guianensis, Lam. Encyc. i. 808 = C. Apalatoa.
spicata, Aubl. Pl. Gui. i. 888 = C. Apalatoa.

APALOCHLAMYS, Cass. in Dict. Sc. Nat. lvi. 238 (1828) = **Cassinia**, R. Br. (Compos.).
Billardierii, DC. Prod. vi. 157 = C. spectabilis.
Endlicheri, DC. l. c. = C. spectabilis.
Kerrii, DC. l. c. = C. spectabilis.

APALOPTERA, Nutt. ex A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 319 = **Abronia**, Juss. (Nyctag.).
annua, Nutt. ex Hook. Kew Journ. v. (1853) 261 = Ab. cyclopters.

APALUS, DC. Prod. v. 507 (1836) = **Blennosperma**, Less. (Compos.).
anthemifolius, DC. l. c. = B. chilense.

APAMA, Lam. Encyc. i. 91 (1783) = **Bragantia**, Lour. (Aristoloch.).
laurifolia, Rafin. Sylva Tellur. 13 = B. Wallichii.
oblignosa, Lam. Encyc. i. 91 = B. Wallichii.

APARGIA, Scop. Fl. Carn. ed. II. ii. 113 (1772) = **Leontodon**, Linn. (Compos.).
alpina, Willd. Sp. Pl. iii. 1547 = L. pyrenaica.
angustifolia, Schleich. ex DC. Prod. vii. 102 = L. tenuiflora.
annua, Viviani, Pl. Arg. 88. t. 6 = L. hispidulus.
arenaria, Morettl. ex DC. Prod. vii. 103 = L. Bernii.
aspera, Sprun. ex Nym. Consp. 468 = L. graecus.
aspera, Waldst. & Kit. Pl. Rar. Hung. ii. 114 = L. asperus.
auraniaca, Tenore, Syll. 892 = L. pyrenaicus.
aurea, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 282 (Hoppe, ex DC. Prod. vii. 107) = Crepis aurea.
autumnalis, Hoffm. Deutschl. Fl. ed. II. ii. 118 = L. autumnalis.
Bernii, Jan, ex DC. Prod. vii. 103 = L. Bernii.
borealis, Bongard, Veget. Ins. Sitcha, 146 = L. borealis.
Brumati, Schied. ex Steud. Nom. ed. II. i. 110 = L. hastilis.
bulbosa, Balb. in Mem. Acad. Tor. (1808) 294 = L. tuberosus.
canescens, Sieber, ex DC. Prod. vii. 103 = L. Bernii.
caucasica, Bieb. Fl. Taur. Cauc. ii. 247 = L. hastilis.
caucasica, Reichb. in Moesel. Handb. ed. II. ii. 1378 = L. asperus.
chillensis, H. B. & K. Nov. Gen. et Sp. iv. 3 = L. chilensis.
cichoracea, Tenore, Prod. 46; Fl. Nap. ii. 166 = L. cichoraceus.
ciliata, Presl, Fl. Sicul. p. xxxi = L. Presliі.
communis, Spenn. Fl. Friburg. ii. 558 = L. hispidus.
coronopifolia, Willd. Sp. Pl. iii. 1558 = Picris radicata.
crispa, Willd. l. c. 1551 = L. crispus.
crocea, Less. in Linnaea, ix. (1834) 152. t. 182.—Sibir. ural.
crocea, Pollin. Fl. Veron. ii. 566. t. 2. f. 3 = L. pyrenaicus.
dandalatis, Scop. Fl. Carn. ed. II. ii. 114 = L. hastilis.
dubia, Willd. Sp. Pl. iii. 1549 = L. hastilis.
erictorum, Richt. [Taschenb. ed. II. (1880)] = L. hastilis.
fasciculata, Bivona, Stirp. Rar. Sicil. ii. [11] t. 2 = L. cichoraceus.
Gouani, Schleich. Cat. Pl. Helv. ed. IV. 7 = L. hirtus.
guestphalica, Boenning. ex Reichb. in DC. Prod. vii. 102 = L. hastilis.
Haenkei, Sternb. ex Steud. Nom. ed. II. i. 110 = L. pyrenaicus.
hastilis, Host, Syn. 493 = L. hastilis.
hastilis, Viviani, Fl. Lib. Spec. 30. t. 26. f. 1 = L. Salzmannis.
heterophylla, Moench, Meth. 542 = L. hastilis.
hieracioides, Willd. Sp. Pl. iii. 1554 = Picris hieracioides.
hirta, Genera. ex Steud. Nom. ed. II. i. 110; ii. 32 = L. pyrenaicus.
hirta, Steud. l. c. i. 110 = L. hispidus.
hirta, Scop. Fl. Carn. ed. II. ii. 114 = L. hirtus.
hispanica, Bieb. Fl. Taur. Cauc. ii. 248 = L. asperus.
hispanica, Willd. Sp. Pl. iii. 1562 = L. hispanicus.
hispida, Bory, in Ann. Sc. Gén. Phys. iii. (1820) 5 = L. Boryi.
hispida, Hoffm. Deutschl. Fl. ed. II. ii. 118 = L. hispidus.
hispida, Willd. Sp. Pl. iii. 1553 = L. hastilis.
hyoseridifora, Saut. ex Schur, Enum. Pl. Transs. 359 = L. caucasicus.
hyoseridifolia, Less. Syn. Comp. 182 = Crepis hyoseridifolia.
hyoseroides, Sieber, in Spreng. Syst. iv. 564 = L. creticus.
hyssopoides, Vest, in Flora, iii. (1820) 7 = L. hirtus?
incana, Mart. Fl. Mosq. 135 = L. hastilis.
incana, Scop. Fl. Carn. ed. II. ii. 114 = L. incanus.
incrassata, Moench, Meth. Suppl. 220 = L. hispidus.
intermedia, Schult. ex Steud. Nom. ed. II. i. 110 = L. hastilis.
laevis, Moench, Meth. 541 = L. hirtus.
leida, Moench, l. c. 543 = L. hastilis.
lucida, Tenore, Cat. Hort. Neap. App. (1819) 71 = L. lucidus.
lyrata, Tenore, Fl. Nap. Prod. 46 = L. tuberosus.
macrorrhiza, Gaertn. ex Spreng. Syst. iii. 665 = L. macrorrhizus.
madia, Host, Fl. Austr. ii. 404 = L. pyrenaicus.
obtusa, Kit. in Linnaea, xxvii. (1853) 406.—Europ. austr.
pratensis, Link, Handb. i. 791 = L. saxanalis.
pyrenaica, Sternb. & Hoppe, ex Steud. Nom. ed. I. 56 = L. alpinus.

APARGIA:—
rigens, Mart. ex Reichb. in Moesel. Handb. ed. II. ii. 1377 = Picris hieracioides.
Rosani, Tenore, Fl. Nap. App. v. 26 = Leontodon Villarsii.
saxina, Gaert. Meyer & Scherb. Fl. Wett. iii. 146 = L. crispus.
salsa, Schur, Enum. Pl. Transs. 384, Index = Taraxacum palustre.
saxatilis, Tenore, Cat. Hort. Neap. (1819) 59 = L. crispus.
scabra, Nocca, ex Steud. Nom. ed. II. i. 110 = Picris sapienoides.
serotina, Jan, ex Steud. l. c. = Hypochoeris radicata.
setosa, Moench, Meth. Suppl. 219 = Crepis setosa.
sicula, Guss. Fl. Sic. Syn. ii. 399 = L. hastilis.
sinuata, Kit. in Linnaea, xxxii. (1863) 404.—Europ. austr.
strigosa, Bieb. Fl. Taur. Cauc. ii. 249 = L. asperrimus.
sudetica, Link, ex Spreng. Syst. iii. 664 = L. hispidus.
Taraxaci, Hornem. in Fl. Dan. t. 1523 = L. autumnalis.
Taraxaci, Willd. Sp. Pl. iii. 1550 = L. Taraxaci.
taraxacifolia, Viv. Fl. Lib. Spec. 50. t. 22. f. 1 = Picris taraxacifolia.
tenuiflora, Gaud. Fl. Helv. vi. 362 = L. alpinus.
tergestina, Hoppe, ex DC. Prod. vii. 103 = L. crispus.
tuberosa, Willd. Sp. Pl. iii. 1549 = L. tuberosus.
umbellata, Schrank, in Denkschr. Regensb. Bot. Ges. ii. (1818) 66 = Picris hieracioides.
variegata, Willd. Sp. Pl. iii. 1552 = Hypochoeris variegata.
varia, Salzm. in Delile, Sem. Hort. Monsp. (1836) = L. Salzmanni.
Villarsii, Willd. Sp. Pl. iii. 1558 = L. Villarsii.

APARGIDIUM, Torr. & Gray, Fl. N. Am. ii. 474 (1841) = **Leontodon**, Linn. (Compos.).
boreale, Torr. & Gray, l. c. = L. borealis.

APARINANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 397 = **Galium**, Tourn. (Rubiac.).
muralis, Fourr. l. c. = G. murale.
verticillatus, Fourr. l. c. = G. verticillatum.

APARINE, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Galium**, Tourn. (Rubiac.).
borealis, Hill, Herb. Brit. ii. 151 = G. boreale.
hispida, Moench, Meth. 640 = G. Aparine.
latifolia, Moench, l. c. = G. Cruciata.
micrantha, All. ex DC. Prod. iv. 610 = G. murale.
parisecnsis, Hill, Herb. Brit. ii. 150 = G. parisiense.
spuria, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 397 = G. spurium.
tenera, Fourr. l. c. = G. spurium.
tricornis, Fourr. l. c. = G. tricorne.
verrucosa, Moench, Meth. 640 = G. sacharatum.
viscidiflora, Rafin. Ann. Nat. (1820) 14.—Am. bor.
Vaillantii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 397 = G. spurium.
vulgaris, Hill, Herb. Brit. ii. 149 = G. Aparine.

APARINELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 397 = **Galium**, Tourn. (Rubiac.).
decipiens, Fourr. l. c. = G. parisiense.
dinoricola, Fourr. l. c. = G. divaricatum.
lilifolia, Fourr. l. c. = G. parisiense.
parisiensis, Fourr. l. c. = G. parisiense.
ruricola, Fourr. l. c. = G. parisiense.
setacea, Fourr. l. c. = G. setaceum.
tenuicaulis, Fourr. l. c. = G. divaricatum.

APARISTHMIUM, Endl. Gen. 1112 (1840) = **Alchornea**, Sw. (Euphorb.).
cordatum, Baill. Adansonia, v. (1865–66) 307 = Al. cordata.
javanense, Hask. Hort. Bogor. 235 = Al. rugosa.
Spruceanum, Baill. Adansonia, v. (1865–66) 307 = Al. cordata.
sumatranum, Reichb. f. & Zoll. in Verh. Nat. Ver. Nederl. Ind. i. (1856) 28 = Al. villosa.

APATANTHUS, Viv. Fl. Lib. Spec. 54. t. 7. f. 3 (1824) = **Hieracium**, Linn. (Compos.).
crinitus, Viv. l. c. = H. crinitum.

APATELIA, Trattin. (Terratrocea.). Syn. ii. 8 (1822) = Saxifraga, Willd. (Terratrocea.).

APATELIA:—
biserrata, DC. Prod. i. 526 = S. biserrata.
glabrata, DC. Prod. i. 526 = Saxifraga serrata.
lanceolata, DC. l. c. = S. lanceolata.
scabra, G. Don, Gen. Syst. i. 569 = S. excelsa.
tomentosa, G. Don, l. c. 568 = S. tomentosa.

APATEMONE, Schott, Gen. Aroid. t. 57 (1858). *AROIDEAE*, Benth. & Hook. f. tii. 984. Modeyana, *Schott, l. c.*—Borneo.

APATITIA, Desv. in Ham. Prod. Pl. Ind. Occ. 42 (1825) = **Belluzia**, Neck. (Melisc.).
blakeoides, Desv. l. c. = B. Aubletii.

APATURIA, Lindl. Gen. et Sp. Orch. 130 (1831) = **Pachystoma**, Blume (Orchid.).
chinensis, Lindl. l. c. 131 = P. chinense.
Lindleyana, Wight, Ic. Pl. Ind. v. 1662 = P. senile.
montana, Lindl. in Wall. Cat. n. 8786 = P. senile.
senilis, Lindl. l. c. n. 8789 = P. senile.
Smithiana, Lindl. l. c. n. 8787 = P. senile.

APEGIA, Neck Elem. i. 251 (1790) = **Cercopegia**, Linn. (Asclep.).

APEIBA, Aubl. Pl. Gui. i. 537. t. 213 ad 216 (1775). *TILIACEAE*, Benth. & Hook. f. i. 287.
AUBLETIA, Schreb. Gen. 858 (1789).
OXYTANDRIUM, Neck. Elem. ii. 255 (1790).
aspera, *Aubl. l. c.* t. 545. t. 216.—Guiana.
australis, A. Rich. Fl. Nov. Zel. t. 34 = Entelea arborescens.
Cimbalaria, *Arruda, in Koster, Trav. Bras.* 488.—Bras.
discolor, *G. Don, Gen. Syst.* i. 555.—Bras.
echinata, *Gaertn. Fruct.* ii. 189. t. 121.—Hab.?
emarginata, Lam. Encyc. i. 210 = Sloanea emarginata.
glabra, *Aubl. Pl. Gui.* i. 541. t. 214.—Guiana.
hirsuta, Lam. Encyc. i. 208.—A. Tiburbou.
hispida, Gaertn. Fruct. ii. 121 = A. Petonno.
hypoleuca, *Steud. in Flora*, xxvi. (1843) ii. 755.—Guiana.
laevis, *G. Don, Gen. Syst.* i. 555.—Guiana.
levis, Swartz, Prod. Veg. Ind. Occ. 88 = glabra.
levis, Aubl. Pl. Gui. t. 214 = glabra.
membranacea, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) *Suppl.* ii. 61.—Venezuel.
Petonmo, *Aubl. Pl. Gui.* i. 543. t. 215.—Guiana.
Tibourbou, *Aubl. l. c.* 538. t. 213.—Guiana; Venezuel.
ulmifolia, *H. B. & K. Nov. Gen. et Sp.* v. 347.—Venezuel.

APEMON, Rafin. Fl. Tellur. ii. 11 (1836) = **Datura**, Linn. (Solan.).
crassicaule, Rafin. l. c. = D. cerntocaula.

APENTOSTERA, Rafin. New Fl. Am. ii. 72 (1836) = **Penstemon**, Mitch. et Chalone, Linn. (Scrophul.).
secundiflora, Rafin. l. c. = P. Menziesii.
trifora, Rafin. l. c. 73 = Chelone nemorosa.

APELLA, Scop. Introd. 107 (1777) = APPELLA, Adans. = **Laurus**, Linn.

APENULA, Neck. Elem. i. 234 (1790) = **Specularia**, Heist. (Campan.).

APERA, Adans. Fam. ii. 495 (1763). *GRAMINEAE*, Benth. & Hook. f. iii. 1152.
AGROSTIS, Trin. Fund. Agrost. 128 (1820).
arundinacea, Hook. f. Fl. N. Zeal. 295. t. 67 (= Stipae sp.).—N. Zel.
aspera, Beauv. Agrost. 151 = Sporobolus asper.
crinita, Beauv. l. c. 81 = Dichelachne crinita.
effusa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 148 = A. Spica-venti.
interrupta, *Beauv.* Agrost. 81.—Rossia austr.; Orient.
palustris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 148 = Agrostis alba.
procumbens, Auct. ex Steud. Nom. ed. II. i. 112 = Agrostis alba, vulgaris.
purpurea, Beauv. Agrost. 81; et Reichb. Ic. Fl. Germ. i. 22 = A. Spica-venti.
Spica-venti, *Beauv. l. c.*—Europ.; Orient.
tenuiflora, Beauv. l. c. 151 = Muhlenbergia Willdenowii.

APERIPHRACTA, Nees, ex Meisn. in DC. Prod. xv. i. 111 (1864) = **Cocotea**, Aubl. (Lauria.).

APERIPHRACTA :—
cuneata, Nees, ex Meisn. in DC. Prod. xv. i. 119 (= *Oreodaphne cuneata*).—Peruv.
Martiana, Nees, l. c. 117 = Oocotea rufo-tomentosa.
Minarum, Nees, l. c. 140 = O. umbrosa.
velutina, Nees, l. c. 139 = O. velutina.

APERULA, Blume, Mus. Bot. Lugd. Bat. i. 365 (1851) = **Lindera**, Thunb. (Laurin.).
asiatica, Meisn. in DC. Prod. xvi. 240 = L. asiatica.
citriodora, Blume, Mus. Bot. Lugd. Bat. i. 366.—Japon.
confusa, Meisn. in DC. Prod. xv. i. 248.—Java.
cuspidata, Blume, Mus. Bot. Lugd. Bat. i. 367.—Ins. Celebes.
densiflora, Meisn. in DC. Prod. xv. i. 242.—Java.
insignis, Blume, in Mus. Bot. Lugd. Bat. i. 367.—Sumatr.
lanceolata, Blume, l. c.—Java.
lucida, Blume, l. c. 366.—Java.
macrophylla, Blume, l. c. 367.—Java.
Neesiana, Blume, l. c. 366 = L. Neesiana.
oxyphylla, Blume, l. c. = L. oxyphylla.
pipericarpa, Meisn. in DC. Prod. xv. i. 242.—Sumatr.
polyantha, Blume, Mus. Bot. Lugd. Bat. i. 366.—Java.
pubescia, Blume, l. c. 367.—Java.
reticulata, Blume, l. c. = L. reticulata.
salicifolia, Blume, l. c.—Java.

APERULA, Gled. in Mém. Acad. Berl. (1751) 127, sphalm. = **Asperula**, Linn (Rubiac.).

APETAHIA, Baill. in Bull. Soc. Linn. Par. i. (1882) 311. *LOBELIACEAE.*
ralatecnia, *Baill. l. c.*—Ins. Pacif.

APETALON, Wight, Ic. v. 22. t. 1758 (1852) = **Didymoplexis**, Griff. (Orchid.).
minutum, Wight, l. c. = D. pallens.

APHACA, [Tourn.] Linn. Gen. ed. I. 217 (1737) = **Lathyrus**, Linn (Legum.).
disperma, Alef. in Oest. Bot. Zeitschr. xi. (1861) 282.—Aegypt.
marmorata, Alef. l. c. = L. polyanthus.
pseudophaca, Alef. in Bonplandia, ix. (1861) 139 = L. pseudosphaca.
vulgaris, C. Presl, in Weitenw. Beitr. ii. (1837) 24 = L. Aphaca.

APHAENANDRA, Miq. Fl. Ind. Bat. ii. 341 (1856). *RUBIACEAE.*
sumatrana, *Miq. l. c.*—Sumatr.

APHAERENA, Miers, in Proc. Hort. Soc. iii. (1863) 294. *SAMYDACEAE*, Benth. & Hook. f. i. 799.
spicata, *Miers, l. c.*—Bras.

APHANACTIS, Wedd. Chlor. And. i. 142. t. 37 A (1855). *COMPOSITAE*, Benth. & Hook. f. ii. 360.
Jamesoniana, *Wedd. l. c.*—Ecuador.

APHANAMIXIS, Blume, Bijdr. 165 (1825) = **Amoora**, Roxb. (Meliac.).
Blumei, Span. Ic. 14 ; et fn Linnaea, xv. (1841) 182 = Am. timorensis.
grandiflora, Blume, Bijdr. 165 = Am. grandiflolia.
Perrottetiana, A. Juss. Mém. Mél. 108 = Am. Perrottetiana.
rubiginosa, Griff. ex C. DC. in DC. Monog. Phan. i. 585 = Am. rubiginosa.
timorensis, A. Juss. Mém. Mél. 107 = Am. timorensis.

APHANANTHE, Link, Enum. Hort. Berol. i. 388 (1821) = **Microtea**, Sw. (Phytolacc.).
celosioides, Link, l. c. = M. mayumensis.

APHANANTHE, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 365. *URTICACEAE*, Benth. & Hook. f. iii. 365.
HOMOIOCELTIS, Blume, Mus. Bot. Lugd. Bat. ii. 64. t. 24 (1855) (1856).
aspera, *Planch. in DC. Prod.* xvii. 208.—Japon.
cuspidata, Planch. l. c. 209 = Girounera reticulata.
phlegmaria, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 367.—Austral.
rectinervia, *Planch. in DC. Prod.* xvii. 208.—Austral.

APHANANTHEMUM, Steud. Nom. ed. II. i. 111 (1840) = **Helianthemum**, Tourn. (Cistin.).
salicifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 240 = H. salicifolium.

APHANES, Linn. Gen. ed. I. 88 (1737) = **Alchemilla**, Linn. (Rosac.).
alata, Steud. Nom. ed. I. 58 = Al. pinnata.
arvensis, Linn. Sp. Pl. ed. I. 123 = Al. arvensis.
cornucopioides, Lag. Gen. et Sp. Nov. 99. n. 7 = Al. cornucopioides.
monogyna, [Pleri] in Ionios Anthol. iii. (1834) 694 = Al. arvensis.
orbiculata, Pers. Syn. i. 150. = Al. orbiculata.
pinnata, Pers. l. c. = Al. pinnata.
triloba, Gilib. Exercit. ii. 430 = Al. arvensis.
tripartita, Pers. Syn. i. 150 = Al. tripartita.
vulgaris, Schur, Enum. Pl. Transs. 884 = Al. arvensis.

APHANIA, Blume, Bijdr. 236 (1825) = **Sapindus**, Linn.
bifoliata, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 238 = S. bifoliatus.
cuspidata, Radlk. l. c. = S. cuspidatus.
Dunura, Radlk. l. c. = S. Dunura.
longipes, Radlk. Sapind. Holland-Ind. Nachtr. 68.—Malaya.
micrantha, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 238 = S. microcarpus.
montana, Blume, Bijdr. 236 = S. montanus.
paucijuga, Radlk. Sapind. Holland-Ind. Nachtr. 68.—Ins.
rubra, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 238 = S. attenuatus.
spreculacea, Radlk. l. c. = S. abyssinicus.
sphaerococca, Radlk. in Act. Congr. Bot. Amst. 1877 (1879) 91.—N. Guinea.

APHANISMA, Nutt. ex Moq. in DC. Prod. xiii. ii. 54 (1849). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 50.
blitoides, *Nutt. ex Moq. l. c.*—Calif.

APHANOCALYX, Oliver, in Hook. Ic. Pl. xi. 53. t. 1066 (1870). *LEGUMINOSAE.*
cynometroides, *Oliver, l. c.*—Afr. occ.

APHANOCHILUS, Benth. in Bot. Reg. sub t. 1282 (1829) = **Blaholtzia**, Willd. (Labiat.).
blandus, Benth. l. c. = E. blandu.
eriostachyus, Benth. l. c. = E. eriostachya.
flavus, Benth. l. c. = E. flava.
foetens, Benth. l. c. = E. incisa.
incisus, Benth. l. c. = E. incisa.
panculatus, Benth. l. c. = E. incisa.
pilosus, Benth. l. c. = E. pilosa.
polystachyus, Benth. l. c. = E. polystachya.

APHANODON, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1853) 110 = **Henriettella**, Naud. (Melast.).

APHANOMYRTUS, Miq. Fl. Ind. Bat. i. i. 480 (1855). *LAURIN.* gen. dub. (= **Basckia**?).
rostrata, *Miq. l. c.* 481.—Malaya.

APHANOMYXIS, DC. Prod. vii. 11. 706, in nota (1839) = APHANOMIXIS, Blume = **Amoora**, Roxb. (Meliac.).

APHANOPAPPUS, Endl. Gen. Suppl. ii. 48 (1842) = Lipochaeta, DC. (Compos.).
Nuttallii, Walp. Rep. ii. 930 = L. micrantha.

APHANOPETALUM, Endl. Gen. 818 (1839). *SAXIFRAGEAE*, Benth. & Hook. f. i. 650.
PLATYPETELA, J. Drum. in Hook. Kew Journ. vii. (1855) 55.
occidentale, *P. Muell. Fragm.* i. 228.—Austral.
resinosum, *Endl. Nov. Stirp. Dec.* 85.—Austral.

APHANOPLEURA, Boiss. Fl. Orient. ii. 855 (1872). *UMBELLIFERAE.*
trachyperma, *Boiss. l. c.*—Armenia.

APHANOSTEMMA, St. Hil. Fl. Bras. Mer. i. 19 (1825) = **Remunculus**, Linn.
apiifolia, St. Hil. l. c. 18 = R. apiifolius.

APHANOSTEPHUS, DC. Prod. v. 310 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 262.
KERRIA, DC. Prod. v. 309 (1836), ex parte.
LEUCOPHIDIUM, DC. Prod. vi. 43 (1837).
arizonicus, A. Gray, in Proc. Am. Acad. xvi. (1881) 81 = humilis.
arkansanus, *A. Gray, Pl. Wright* i. 93.—Texas.
humilis, *A. Gray, l. c.*—Texas.
pilosus, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 457 = ramosissimus.
ramosissimus, *DC. Prod.* v. 310.—Texas.
ramosus, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 81. —Mexic.
Riddellii, Torr. & Gray, Fl. N. Am. ii. 189 = ramosissimus.

APHANTOCHAETA, A. Gray, in Pacif. Rail. Rep. iv. 99 (1856). *COMPOSITAE*, Benth. & Hook. f. ii. 251.
exilis, *A. Gray, l. c.* 100.—Calif.

APHARICA, Schlecht. in Linnaea, vi. (1831) Litt. 78 = APHANIA, Blume = **Sapindus**, Linn.

APHELANDRA, R. Br. Prod. 475 (1810). *ACANTHACEAE*, Benth. & Hook. f. ii. 1102.
AMATHEA, Raffin. Fl. Tellur. iv. 65 (1836).
HEMISANDRA, Scheidw. in Bull. Acad. Brux. ix. i. (1842) 22.
HEMITOME, Nees, in DC. Prod. xi. 295 (1847).
HYDROMESTUS, Schendw. in Otto & Dietr. Gartenz. x. (1842) 285.
LAGOCHILIUM, Nees, in Mart. Fl. Bras. ix. 85. t. 10 (1847).
POECILOCNEMIS, Mart. ex Nees, in DC. Prod. xi. 295 (1847).
STROBILORHACHIS, Link, Klotzsch & Otto, Ic. Pl. Rar. 117. t. 48 (1839).
SYNANDRA, Schrad. in Goett. Gel. Anz. i. (1831) 715.
acanthifolia, *Hook. Ic. Pl.* t. 113.—Peruv.
Acanthus, *Nees, in DC. Prod.* xi. 302.—Ecuador.
acutifolia, *Nees, l. c.* 299.—Ind. occ.
aeroviena, *N. E. Br. in Illustr. Hortic.* xxxi. (1884) t. 537.—Bras.
aurantiaca, *Lindl. Bot. Reg.* 31. t. 12.—Mexic.
bullata, *H. Wendl. in Hamb. Gartenb.* xix. (1863) 30. —Am. centr.
carduifolia, Hook. Ic. Pl. t. 718 = formosa.
chartacea, *T. Anders. in Journ. Bot.* ii. (1864) 289.—Peruv.
Chamissoniana, *Nees, in Mart. Fl. Bras.* ix. 90.—Bras.
Clava, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 174. —Bras.
cristata, *R. Br. Prod.* 475, in obs.—Ind. occ.
cristata, H. B. & K. Nov. Gen. et Sp. ii. 286 = pectinata.
cristata, Lindl. Bot. Reg. t. 1477 = tetragona.
Dehnhardtii, *Tenore, Ind. Sem. Hort. Neap.* (1829) 14.—Hab. !
Deppeana, *Cham. & Schlecht. in Linnaea*, v. (1830) 96.—Mexic.
Fascinator, *Linden & André, Illustr. Hortic.* xxi. (1874) t. 184.—Am. trop.
flava, *Nees, in DC. Prod.* xi. 297.—Reg. Colomb.
formosa, *Nees, l. c.* 301.—Peruv.
fulgens, *Decne. in Rev. Hortic.* Sér. III. i. (1847) 21.—Mexic.
fulgens, *Hort. ex Regel & Rach, Gartenfl.* (1858) 243 = tenuiflora.
glabrata, *Willd. ex Nees, in DC. Prod.* xi. 296.—Am. merid.
Ghiesbreghtiana, Hort. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 388 = Jacobinia Ghiesbreghtiana.
Haenkeana, *Nees, Pl. Haenk. ex DC. Prod.* xi. 298.—Am. centr.
Hartwegiana, *Nees, in Benth. Pl. Hartw.* 236.—Reg. Colomb.
Hieronymi, *Griseb. in Gottt. Abh.* xxiv. (1879) 260.—Reg. Argent.
hirsuta, Mart. ex Nees, in DC. Prod. xi. 417 = Beloperone hirsuta.
hydrometsa, *Hemsl. Biol. Centr. Am. Bot.* ii. 512.—Mexic.
ignea, Nees, ex Steud. Nom. ed. II. i. 111 = Stenandrium igneum.

APHELANDRA:—
illustrata.—*Cf. Gard. Chron.* v. (1876) 608.—Bras.
laterita, *Loud. Encyc. Pl.* 1539.—Guatemala.
Leopoldii, Van Houtte, in Fl. des Serres, ix. Sér. I. (1853) 58 = squarrosa.
Liboniana, *Linden, ex Hook. f. Bot. Mag.* 5468.—Am. trop.
longicaespitosa, Hort. ex Lindl. in Paxt. Fl. Gard. ii. 78 = Thyrsacanthus callistachys.
longiscapa, Hort. ex Lem. in Fl. des Serres, Sér. I. iii. (1847) 240 = Thyrsacanthus Lemairianus.
lutea, *Nees, in DC. Prod.* xi. 300.—Bras.
lyrata, *Nees, l. c.* 309.—Am. austr.
macrostachya, *Nees, in Mart. Fl. Bras.* ix. 88.—Bras.
Margaritae, *E. Morr. in Belg. Hortic.* xxxiii. (1883) 315. t. 19.—Bras.
marginata, *Nees & Morr. in Nov. Act. Nat. Cur.* xi. (1832) t. 53.—Bras.
mediaurita, *Cogn. & March. Pl. Ornem.* i. (1878) t. 80.—Bras.
melagrinoula, *Nees, in DC. Prod.* xi. 297.—Reg. Colomb.
micana, *Morite. ex Vathe, Ind. Sem. Hort. Berol.* (1876) *App.* 1.—Venezuela.
monticola, Mart. ex Nees, in DC. Prod. xi. 364 = Jacobinia monticola.
mucronata, *Nees, in DC. Prod.* xi. 301.—Peruv.
nemoralis, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 90.—Bras.
nervosa, *T. Anders. in Journ. Bot.* ii. (1864) 289.—Peruv.
nitens, *Hook. f. Bot. Mag.* t. 5741.—Reg. Colomb.
nitida, Nees & Mart. in Nov. Act. Nat. Cur. xi. i. (1852) 57 = Celosicoria nitida.
noblis, *Hort.—Cf. Gard. Chron.* (1867) 461.—Hab.?
nuda, *Nees, in Mart. Fl. Bras.* ix. 89.—Bras.
ocstachys, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 179.—Bras.
ornata, *T. Anders. in Journ. Bot.* ii. (1864) 289.—Bras.
pectinata, *Willd. ex Nees, in DC. Prod.* xi. 297.—Am. austr.
Portesna, *Moral, in Fl. des Serres*, Sér. I. x. (1854–55) 41. t. 984.—Bras.
pringlaica, *Hiern, in Kjoob. Vidensk. Meddel.* (1877–76) 78.—Am. centr.
pulcherrima, *H. B. & K. Nov. Gen. et Sp. Pl.* ii. 236.—Reg. Colomb.
pulcherrima, Willd. ex Nees, in DC. Prod. xi. 295 = tetragona.
pumila, *Nichols. Dict. Gard.* i. 91.—Bras.
punctata, *Hort.—Cf. Gard. Chron.* (1881) ii. 86; *Illustr. Hortic.* xxix. (1882) t. 457.—Am. merid.
querolfola, *Nees, in DC. Prod.* xi. 302.—Reg. Colomb.
rupaula, *Nees, l. c.* 798.—Peruv.
Roezlii, Carr. in Rev. Hortic. (1872) 100 = suranticea.
runchnata, *Klotzsch, ex Nees, in DC. Prod.* xi. 302.—Reg. Colomb.
Schiedeana, *Cham. & Schlecht. in Linnaea*, v. (1830) 95.—Mexic.
sciophila, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 91.—Bras.
Sinclairiana, *Nees, in Benth. Bot. Voy. Sulph.* 146. t. 47.—Am. centr.
Sintenisi, *Lindau, Cat.* n. 95 (1876); *Rev. Hortic.* (1876) 237.—Peruv.
squarrosa, *Nees, in Mart. Fl. Bras.* ix. 89.—Bras.
stephanophysa, *Nees, l. c.* 90.—Am. austr.
sulphurea, *Hook. f. Bot. Mag.* t. 5951.—Ecuador.
tarapotensis, Lindau, ex Rafarin, in Rev. Hortic. (1877) 86 = Sinitzini.
tmaniform, *Regel & Rach, Gartenfl.* (1858) 342.—Hab.?
tetragona, *Nees, in DC. Prod.* xi. 295.—Reg. Colomb.
tridentata, *Hemsl. Biol. Centr. Am. Bot.* ii. 513.—Costa Rica.
variegata, *Morel, Fl. des Serres*, Sér. I. x. (1854–55) 53. t. 981.—Hab.?
verticillata, *Nees, in DC. Prod.* xi. 281.—Mexic.

APHELANDROS, St. Lag. in Ann. Soc. Bot. Lyon, vii. 119 (1880) = **Aphelandra**, R. Br. (Acanth.).

APHELEXIS, D. Don, in Mem. Wern. Soc. v. (1826) 546 = **Helichrysum**, Tourn. (Compos.).
adhaerens, Boj. ex DC. Prod. vi. 217.—Madag.
Candollei, Boj. ex DC. l. c.—Madag.
ericoides, Sweet, Hort. Brit. ed. I. 238 = Helichrysum ericoides.

APHELEXIS:—
fasciculata, D. Don, in Mem. Wern. Soc. v. (1826) 546 = H. scaposides.
Hilfermia, D. Don, l. c. 547 = H. scaposides.
heterophylla, G. Don, in Sweet, Hort. Brit. ed. III. 376 = Helipterum heterophyllum.
humilis, D. Don, in Mem. Wern. Soc. v. (1826) 547 = H. humile.
hypnoides, DC. Prod. vi. 218.—Madag.
incana, G. Don, in Sweet, Hort. Brit. ed. III. 379 = Helipterum Incanum.
lycopodioides, Boj. in DC. Prod. vi. 217.—Madag.
macrantha, Pxtt. Fl. Mag. xv. 269 = H. humile.
Majorifolia, DC. Prod. vi. 217.—Madag.
scaposides, D. Don, in Mem. Wern. Soc. v. (1826) 546 = H. scaposides.

APHELIA, R. Br. Prod. 251 (1810). *CENTROLEPIDEAE*, Benth. & Hook. f. iii. 1026.
BRIZULA, Hieron. in Abh. Naturf. Ges. Halle, xii. (1873) 306, 308 (Centrolep. 92, 94).
Briaula, *F. Muell. Fragm.* v. 208.—Austral.
cyperoides, *R. Br. Prod.* i. 252.—Austral.
Drummondii, *Benth. Fl. Austral.* vii. 201.—Austral.
gracilis, *Sond. in Linnaea*, xxviii. (1855) 237.—Austral.
Gunnii, Hook. f. Fl. Tasm. ii. 75. t. 138 = gracilis.
monogyna, Hieron. in Abh. Naturf. Ges. Halle, xii. (1873) 308 (Centrolep. 94) = Centrolepis monogyna.
nuiana, *Hook. f. ex Benth. Fl. Austral.* vii. 200.—Austral.
Pumilio, *F. Muell. in Linnaea*, xxviii. (1855) 226.—Austral.

APHILLANTHES, Neck. Elem. iii. 199 (1790) = **Aphyllanthes**, Tourn. (Liliac.).

APHLOIA, Benn. Pl. Jav. Rar. 192 (1838) in adnot. *BIXINEAE*, Benth. & Hook. f. i. 136.
NEUMANNIA, A. Rich. Fl. Cub. 96, in adnot. (1845).
integrifolia, Benn. Pl. Jav. Rar. 192 in adnot. = mauritiana.
magnacarinata, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 278.—Madag.
mauritiana, *Baker, Fl. Maurit.* 12.—Ins. Mascar.
theaeformis, Benn. Pl. Jav. Rar. 192, in adnot. = mauritiana.

APHOMA, Rafin. Fl. Tellur. ii. 81 (1836) = **Iphigenia**, Kunth (Liliac.).
angustifora, Rafin. l. c. 82 = I. indica.
cuneata, Rafin. l. c. = I. indica.

APHOMONIX, Rafin. Fl. Tellur. ii. 85 (1836) = **Saxifraga**, Tourn.
hederacea, Rafin. l. c. = S. sarmentosa.

APHONINA, Neck. Elem. iii. 237 (1790) = **Pariana**, Aubl. (Gramin.).

APHORA, Neck. Elem. iii. 50 (1790) = **Virgilia**, Lam. (Legum.).

APHORA, Nutt. in Trans. Am. Phil. Soc. Ser. II. v. (1837) 174 = **Argithamnia**, P. Br. (Euphorb.).
Blodgettii, Torr. ex Chapm. Fl. S. U. St. 408.—Florida.
catamarinensis, Griseb. in Goett. Abh. xxiv. (1879) 58 = catamarinensis.
Drummondii, Engelm. & Gray, Pl. Lindh. i. 25.—Texas.
humulis, Engelm. & Gray, l. c. 54 = Arg. humilis.
humilis, Torr. Bot. Mex. Bound. 196 = Arg. neomexicana.
laevis, A. Gray, ex Torr. l. c. = Arg. laevis.
lanceolata, Engelm. & Gray, Pl. Lindh. i. 26 = Arg. lanceolata.
mercurialina, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 174 = Arg. mercurialina.
pilostemma, Torr. in Bot. Mex. Bound. 197 = Arg. mercurialina.
serrata, Torr. l. c. = Arg. serrata.

APHRAGMIA, Nees, in Lindl. Nat. Syst. ed. II. 444 (1836) = **Rualtia**, Lin. (Acanth.).
Hookeri, Nees, l. c. = R. chlamydia.
rotundifolia, Nees, ex Benth. Bot. Voy. Sulph. 146 = R. floribunda.

APHRAGMUS, Andrz. ex DC. Prod. i. 309 (1824) = **Bryx**, Sternb. & Hoppe (Crucif.).
Eschscholtsiana, Andrz. ex DC. l. c.—Ins. Aleutian.

APHYLAX, Saliab. in Trans. Hort. Soc. i. (1812) 271 = **Anellema**, R. Br. (Commelin.).
spiralis, Saliab. l. c. = An. spiratum.

APHYLLANTHES, Tourn. ex Linn. Gen. ed. I. 357 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 795.
juncea, Salisb. Parad. Lond. t. 9 = monspeliensis.
monspeliensis, *Linn. Sp. Pl.* 294.—Reg. Mediterr.

APHYLLEIA, Champ. in Calc. Journ. Nat. Hist. vii. (1847) 468 = **Gerbera**, Gronov. (Compos.).

APHYLLOCALUON, Lag. Amen. Nat. i. 88 (1811) = **Gerbera**, Gronov. (Compos.).

APHYLLOCLADUS, Wedd. Chlor. And. i. 11. t. 3 (1855) = **Hyalis**, D. Don (Compos.).
decurrentus, Hieron. Ic. Fl. Argent. 28. t. 5. f. 1.—Reg. Argent.
spartioides, Wedd. l. c.—Bolivia.

APHYLLON, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) 221. *OROBANCHACEAE*, Benth. & Hook. f. ii. 928.
ANOPLON, Reichb. Consp. 213* (1828).
GYMNOCAULIS, Nutt. ex Benth. & Hook. f. Gen. ii. 993 (1876).
LOXANTHES, Rafin. Neogenyt. 3 (1825).
MYZORRHIZA, Phil. in Linnaea, xxix. (1857–58) 86.
THALESIA, Rafin. in Am. Monthly Mag. (1818) 267.
californicum, *A. Gray, in Bot. Calif.* i. 584.—Am. bor. occ.
comosum, *A. Gray, l. c.*—Am. bor. occ.
Cooperi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 307.—Am. bor. occ.
Dugesii, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 152.—Mexic.
fasciculatum, *Torr. & Gray, in A. Gray, Man. Bot. U. St.* ed. I. 281.—Am. bor.
ludovicianum, *A. Gray, in Bot. Calif.* i. 585.—Am. bor. multiflorum, *A. Gray, l. c.*—N. Mexic.
pinetorum, *A. Gray, l. c.*—Am. bor. occ.
tuberosum, *A. Gray, l. c.*—Calif.
uniflorum, *A. Gray, Man. Bot. U. St.* ed. I. 390.—Am. bor.

APHYLLORCHIS, Blume, 'Bijdr.' t. 77 (1825)' (nobis latet). *ORCHIDEAE*, Benth. & Hook. f. iii. 606.
Hasseltii, *Blume, Mus. Bot. Lugd. Bat.* i. 31.—Java.
montana, *Reichb. f. in Linnaea*, xli. (1877) 57.—Ind. or.
pallida, *Blume, Bijdr.* t. 77; *Mus. Bot. Lugd. Bat.* i. 30.—Malaya.
Spicnlaea, *Reichb. f. in Linnaea*, xli. (1877) 58.—Borneo.

APHYTEIA, Linn. Dis. Aphyt. 7 (1776); (Amoen. Acad. viii. 310. t. 7) (1785) = **Hydnora**, Thunb. (Cytin.).
Ackarii, Steud. Nom. ed. II. i. 111, sphalm. (Cf. Thunb. Fl. Cap. ed. Schult. 499) = H. africana.
Hydnora, Linn. f. Diss. 16 = H. africana.
multiceps, Burch. Trav. i. 213, in nota = Cytinus dioicus.
triceps, Steud. Nom. ed. II. i. 111 = H. triceps.

APIASTRUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 643 (1840). *UMBELLIFERAE*, Benth. & Hook. f. i. 886.
angustifolium, *Nutt. l. c.* 644.—Calif.
latifolium, Nutt. l. c. = angustifolium.

APICRA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 167. *LILIACEAE*, Benth. & Hook. f. iii. 776.
APRICA, D. Dietr. Syn. ii. 1097 (1840).
CATEVALA, Medic. Theod. 67 (1786).
albicans, Willd. l. c. 271 = Haworthia albicans.
amoenula, Willd. l. c. 272 = Haworthia venosa.
arachnoidea, Willd. l. c. 166 = Haworthia arachnoidea.
aspera, *Willd. l. c.* 274.—Afr. austr.

APICRA :—
aspersula, Haw. ex Steud. Nom. ed. II. i. 111 = Haworthia aspersula.
atrovirens, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 168 = Haworthia atrovirens.
attenuata, Willd. l. c. 270 = Haworthia attenuata.
bicarinata, *Herw. Suppl.* 68.—Afr. austr.
bullulata, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 278 = pentagona.
congesta, *Baker*, in *Journ. Linn. Soc.* xviii. (1880) 218.—Afr. austr.
cymbaefolia, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 271 = Haworthia cymbiformis.
deltoidea, *Baker*, in *Journ. Linn. Soc.* xviii. (1880) 217.—Afr. austr.
expansa, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 272 = Haworthia rigida.
fasciata, Willd. l. c. = Haworthia fasciata.
folliolosa, *Willd. l. c.* 274.—Afr. austr.
granata, Willd. l. c. 269 = Haworthia margaritifera.
imbricata, Willd. l. c. 273 = spiralis.
margaritifera, Willd. l. c. 269 = Haworthia margaritifera.
maxima, Steud. Nom. ed. I. 59; ed. II. i. 112 = Haworthia margaritifera.
minor, Steud. ll. cc. = Haworthia margaritifera.
mirabilis, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 269 = Haworthia mirabilis.
nigra, Haw. in Phil. Mag. (1824) 802 = Haworthia nigra.
patula, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 272 = Haworthia rigida.
pentagona, *Willd. l. c.* 270.—Afr. austr.
pseudo-rigida, Haw. Suppl. 62 = Haworthia subrigida.
Pumilio, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 271 = Haworthia reticulata.
quinquangularis, Sweet, Hort. Brt. ed. II. 521 = Aloë quinquangularis.
Radula, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 270 = Haworthia Radula.
recurva, Willd. l. c. = Haworthia recurva.
reticulata, Willd. l. c. 272 = Haworthia reticulata.
retusa, Willd. l. c. 271 = Haworthia retusa.
rigida, Willd. l. c. 272 = Haworthia subrigida.
spiralis, *Baker*, in *Journ. Linn. Soc.* xviii. (1880) 217.—Afr. austr.
spiralis, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 273 = pentagona.
spirella, Willd. ex Haw. Suppl. Pl. Succ. 64 = pentagona.
tortuosa, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 274 = Haworthia viscosa.
translucens, Willd. l. c. 168 = Haworthia translucens
tricolor, Willd. l. c. 271 = Haworthia venosa.
viscosa, Willd. l. c. 274 = Haworthia viscosa.

APILIA, Rafin. Alsog. Am. 89 (1838) = **Fraxinus**, Tourn. (Oleac.).
laciniata, Rafin. l. c. = F. excelsior, var. †
macrophylla, Rafin. l. c. = F. macrophylla.
pendula, Rafin. l. c. = F. excelsior.

APINAGIA, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 97. *PODOSTOMACEAE*, Benth. & Hook. f. iii. 111.
BLANDOWIA, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 109. t. 4. f. 2, nomen prinsepe.
MONOSTYLIS, Tul. Podost. Monog. 201 (1852).
NEOLACIS, Wedd. in DC. Prod. xvii. 59 (1873).
divaricata, *Tul. & Wedd. ex Tul. l. c.*—Bras.
fucoides, *Tul. l. c.* 98.—Bras.
Gardneriana, *Tul. l. c.*—Bras.
membranacea, *Tul. l. c.* 99.—Bras.
Preissii, *Tul. in Arch. Mus. Par.* vi. (1852) 108.—Itilis.
psyllophora, *Tul. & Wedd. ex Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 97.—Bras.
pusilla, *Tul. l. c.* 99.—Guiana.
pygmæa, *Tul. l. c.*—Bras.
Riedelii, *Tul. l. c.* 98.—Bras.
rupptoides, *Tul. l. c.*—Reg. Colomb.
stricta, *Griseb. ex Wedd. in DC. Prod.* xvii. 64.—Peruv.
Willdenowii, *Tul. in Arch. Mus. Par.* vi. (1852) 106.—Peruv.; Chili.

APINELLA, Neck. Elem. i. 191 (1790) = **Trinia**, Hoffm. (Umbellif.).
APINUS, Neck. Elem. iii. 369 (1790) = **Pinus**, Tourn. (Conif.).

APIOCARPUS, Montrous. in Mém. Acad. Lyon, x. (1860) 180 = **Akania**, Hook. f. (Sapindac.).
Moguini, Montrous. l. c. 191.—N. Caled.

APIOPETALUM, Baill. Adansonia, xii. (1878) 133. *ARALIACEAE*.
glabratum, *Baill. l. c.* 134.—N. Caled.
velutinum, *Baill. l. c.*—N. Caled.

APIOS, Moench, Meth. 165 (1794). *LEGUMINOSAE*, Benth. & Hook. f. i. 532
BRADLEA, Adans. Fam. ii. 324 (1763).
CYSTOTROPIS, Wall. Pl. As. Rar. i. 49. t. 62 (1830).
americana, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 355.—Am. bor.
carnea, *Benth. ex Baker*, in *Hook. f. Fl. Brit. Ind.* ii. 185.—Reg. Himal.
chinensis, Spreng. Syst. iii. 310 = Wisteria chinensis.
Fortunei, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 206.—China; Japon.
frutescens, Pursh, Fl. Am. Sept. ii. 474 = Wisteria frutescens.
tuberosa, *Moench, Meth.* 165.—Am. bor.

APIOSPERMUM, Klotzsch, in Abh. Akad. Berl. (1853) 251, 1855 = **Platia**, Linn. (Arold.).
obcordatum, Klotzsch, l. c. = P. Stratiotes.

APIROPHORUM, Neck. Elem. ii. 72 (1790) = **Pyrus**, Tourn. (Rosac.).

APISTA, Blume, Bijdr. 296 (1825) = **Podoohilus**, Blume (Orchid.).
tenuis, Blume, l. c. = P. tenuis.

APIUM, [Tourn.] Linn. Syst. ed. I. (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 888.
CELERI, Adans. Fam. ii. 498 (1763).
CICLOSPERMUM, Lag. Amen. Acad. Madr. ii. 101 (1811).
CRITAMUS, Hoffm. Gen. Umb. 182 (1814).
CYCLOSPERMUM, DC. Prod. iv. 105, in syn. (1830).
HELODIUM, Dum. Fl. Belg. 77 (1827).
HELOSCIA, Dum. Anal. Fam. 34 (1829).
HELOSCIADIUM, Koch, in Nov. Act. Nat. Cur. xii. t. (1825) 125.
LEPTOCAULIS, Nutt. ex DC. Prod. iv. 107 (1830).
MACHAIRIA, Neck. Elem. i. 172 (1790).
SELINON, Adans. Fam. ii. p. † (23) * 603 (1763).
SIUMIS, Rafin. New Fl. Am. iv. 31 (1836).
SPERMOLEPIS, Rafin. Neogenyt. 2 (1825).
Ammi, Crantz, Stirp. Austr. ed. I. fasc. iii. 109; ed. II. 217 = Ammi majus.
Ammi, *Urban*, in *Mart. Fl. Bras.* xi. i. 341.—Bras.
Amomum, Stokes, Bot. Mat. Med. ii. 106 = Sium Amomum.
Anisum, Crantz, Cl. Umb. Emend. 101 = Pimpinella Anisum.
australe, Thou. Fl. Trist. d'Acugn. 43 = prostratum.
bipinnatum, Walt. Fl. Carol. 115 = Selinum canadense.
biternatum, Stokes, Bot. Mat. Med. ii. 150 = Aegopodium Podagraria.
Carsonii, *Benth. & Hook. f. Gen.* i. 888.—Am. bor.
Carvi, Crantz, Stirp. Austr. ed. I. fasc. iii. 110; ed. II. 218 = Carum Carvi.
Celleri, Gaertn. Fruct. i. t. 22 = graveolens.
chilense, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 358.—Chili.
Commersonii, *DC. Prod.* iv. 101.—Hab.?
crassipes, *Reichb. f. ic. Fl. Germ.* xxi. 9.—Hispan.
crispum, Mill. Gard. Dict. ed. VIII. n. 3 = Carum Petroselinum.
decumbens, Eckl. & Zeyh. Enum. 340 = graveolens.
divaricatum, *Benth. & Hook. f. ex S. Wats. Bibl. Ind. N. Am. Bot.* 413.—Am. bor.
dulce, Mill. Gard. Dict. ed. VIII. n. 5 = graveolens.
echinatum, *Benth. & Hook. f. ex S. Wats. Bibl. Ind. N. Am. Bot.* 413.—Am. bor.
filiforme, Hook. Ic. t. 819 = prostratum.
flexuosum, *Phil. in Linnæa*, xxxiii. (1864–65) 91.—Chili.
fractophyllum, *Hornem. Hort. Hafn. Suppl.* 128.—Hab.?
glaucescens, *H. B. & K. Nov. Gen. et Sp.* v. 17.—Reg. Colomb.
glaucum, Crantz, Cl. Umb. Emend. 100 = Trinia vulgaris.
gracile, *Benth. & Hook. f. Gen.* i. 888.—Chili.

APIUM :—
graveolens, Cham. in Linnæa, i. (1826) 888 = prostratum.
graveolens, *Linn. Sp. Pl.* 264.—Europ.; Orient.; Ind. bor. occ.; Calif
humile, *Benth. & Hook. f. Gen.* i. 888.—Afr. austr.
inundatum, *Reichb. f. ic. Fl. Germ.* xxi. 9.—Europ
involucratum, Roxb. ex Flem. in As. Res. xi. (1810, 157 = Carum Roxburghianum.
junceum, Stokes, Bot. Mat. Med. ii. 155 = Carum segetum.
lactum, Salisb. Prod. 169 = Carum Petroselinum.
latifolium, Mill. Gard. Dict. ed. VIII. n. 3 = Carum Petroselinum.
latifolium, Poir. Encyc. iv. 369 = Carum peregrinum.
leptophyllum, *F. Muell. ex Benth. Fl. Austral.* iii. 372.—Austral.
linearе, *Benth. & Hook. f. Gen.* i. 888.—Am. bor.
lobatum, Gilib. Fl. Lituan. ii. 41 = graveolens.
lusitanicum, Mill. Gard. Dict. ed. VIII. n. 7 = graveolens
maritimum, Salisb. Prod. 169 = graveolens.
montanum, *H. B. & K. Nov. Gen. et Sp.* v. 17.—Reg. Colomb.
nodiflorum, *Reichb. f. ic. Fl. Germ.* xxi. 10.—Europ.
patens, *S. Wats. Bibl. Ind. N. Am. Bot.* 413.—Am. bor.
peregrinum, Crantz, Cl. Umb. Emend. 101 = Carum peregrinum.
petræum, Crantz, l. c. 104 = Ammi majus.
Petroselinum, Cham. in Linnæa, i. (1826) 889 = prostratum.
Petroselinum, Linn. Sp. Pl. ed. I. 264 = Carum Petroselinum.
Popei, *A. Gray*, in *Proc. Am. Acad.* vii. (1868) 343.—Am. bor.
prostratum, *Labill. Relat. i.* 141; *Vent. Jard. Malm.* 81.—Austral.; Reg. Antarct.
pumilum, Crantz, Stirp. Austr. ed. I. fasc. iii. 109; ed. II. 218 = Trinia vulgaris.
ranunculifolium, *H. B. & K. Nov. Gen. et Sp.* v. 18. t. 421.—Reg. Colomb.
rapaceum, Mill. Gard. Dict ed VIII. n. 6 = graveolens.
repens, *Reichb. f. ic. Fl. Germ.* xxi. 10.—Europ.
romanum, Zuccagni, in Roem. Collect. 135 = Carum Petroselinum.
segetum, Dum. Fl. Belg. 71 = Carum segetum.
Sium, Crantz, Stirp. Austr. ed. I. fasc. iii. 106; ed. II. 216 = Sium erectum.
ternatum, Pall. ex Steud. Nom. ed. II. i. 112 = Archangelica Gmelini.
ternatum, Schlecht. ex Roem. & Schult. Syst. vi. 431 = Ligusticum scoticum.
ternatum, Siebold, in Verh. Batav. Gen. xii. (1830) 46 = Ligusticum scoticum.
trifoliatum, Wight & Arn. Prod. i. 368, sphalm. = Pimpinella Heyneana.
tuberosum, Steud. Nom. ed. I. 89 = Carum Petroselinum
Visnaga, Crantz Cl. Umb. Emend. 104 = Ammi Visnaga.
vulgare, Lam. Fl. Fr. iii. 444 = Carum Petroselinum.

APLACTIS, Rafin. Fl. Tellur. ii. 42 (1836) = **Solidago**, Tourn. (Compos.).
paniculata, Rafin. l. c. = S. pauciflosculosa.

APLARINA, Rafin. New Fl. Am. iv. 99 (1836) = **Euphorbia**, Tourn.
macrophylla, Rafin. l. c. 100 = E. thymifolia.
prostrata, Rafin. l. c. 99 = E. prostrata.

APLECTRA, Rafin. Cat. 13 (1824) = **Aplectrum**, Nutt (Orchid.).
elatior, Rafin. l. c. = Aplectrum hyemale.

APLECTROCAPNOS, Boiss. & Reut. Diagn. Ser. I. v. 79 (1844) = **Sarcocapnos**, DC. (Papav.).
baetica, Boiss. & Reut. l. c. = S. baetica.
integrifolia, Boiss. Diagn. Ser. II. i. 13 = S. baetica.

APLECTRUM, Torr. Comp. Fl. St. 222 (1826). *ORCHIDEAE*, Benth. & Hook. f. iii. 497.
APLECTRA, Rafin. Cat. 13 (1824).
hyemale, *Torr. l. c.*—Am. bor.

APLECTRUM, Blume, in Flora, xiv. (1831) 502 = **Aplectrum**, A. Gray (Melast.).
confine, Blume, Mus. Bot. Lugd. Bat. i. 86 = Aspl. confine.
myrtiformе, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 205 = Medinilla myrtiformis.
nodosum, Blume, Mus. Bot. Lugd. Bat. i. 37 = Aspl. nodosum.

APLECTRUM :—
ovalifolium, Nand. ex A. Gray, Bot. U. St. Expl. Exp. i. 597 = Allomorpha ovalifolia.
pallens, Blume, Mus. Bot. Lugd. Bat. i. 38 = Anplectrum pallens.
rostratum, Blume, in Flora, xiv. (1831) 502 = Anpl. rostratum.
stipulare, Blume, l. c. 503 = Anpl. stipulare.
viminale, Blume, l. c. 502 = Anpl. viminale.

APLEURA, Phil. in Linnæa, xxxiii. (1864–65) 89.
UMBELLIFERÆ, Benth. & Hook. f. i. 875.
nucamentacea, *Phil. l. c.* 90.—Chili.

APLEXIA, Rafin. Neogenyt. 4 (1825) = **Leersia**, Sw. (Gram.).
virgata, Rafin. in Ser. Bull. Bot. 220 = L. virginica.
virginica, Rafin. Neogenyt. 4 = L. virginica.

APLILIA, Rafin. New Fl. Am. iii. 98 (1836) = **Fraxinus**, Tourn. (Oleac.).
lacinista, Rafin. l. c. = F. excelsior.
macrophylla, Rafin. l. c. = F. excelsior.

APLINA, Rafin. Fl. Tellur. iv. 119 (1836) = **Stachelina**, Linn. (Compos.).
fuscata, Rafin. l. c. = S. uniflosculis.

APLOCA, Neck. Elem. i. 255 (1790) = **Periploca**, Tourn. (Asclep.).

APLOCARYA, Lindl. Bot. Reg. (1844) sub t. 46 = **Dolichos**, Lindl. (Convolv.).
divaricata, Lindl. l. c.—Chili.

APLOCHLAMIS, Steud. Nom. ed. II. i. 809, in syn. (1840) sphalm. = APALOCHLAMYS, Cass. = **Cassinia**, R. Br. (Compos.).

APLOLEIA, Rafin. Fl. Tellur. ii. 17 (1836) = **Tradescantia**, Linn. (Commelin.).
diffusa, Rafin. l. c.—Hab. ?

APLOLOPHIUM, Cham. in Linnæa, vii. (1832) 556. *BIGNONIACEÆ*. Benth. & Hook. f. ii. 1038.
HAPLOLOPHIUM, Endl. Gen. 712 (1839).
Blanchetti, *DC. Prod.* ix. 192.—Bras.
bracteatum, *Cham. in Linnæa*, vii. (1832) 556.—Bras. Landii, *DC. Prod.* ix. 192.—Bras.

APLOPAPPUS, Cass. in Dict. Sc. Nat. lvi. 168 (1828). *COMPOSITÆ*, Benth. & Hook. f. ii. 253.
ACAMPHOPAPPUS, A. Gray, in Proc. Am. Acad. viii. (1873) 634.
CHROILEMA, Bernh. Del. Sem. Hort. Erf. (1840).
CROILEMA, Heynh. Nom. ii. 171 (1846).
CROPTILON, Rafin. Fl. Tellur. ii. 47 (1836).
ERIOCARPUM, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 320.
HAPLOPAPPUS, Endl. Gen. 386 (1838).
HOMOPAPPUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 380.
HOOREBEKIA, Cornelius. in Mussch. Hort. Gand. 190 (1817), nomen princeps.
ISOPAPPUS, Torr. & Gray, Fl. N. Am. ii. 289 (1841).
PRIONOPSIS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 329.
PYRROCOMA, Hook. Fl. Bor. Am. i. 306. t. 107 (1838).
SIDERANTHUS, Fras. ex Sweet, Hort. Brit. ed. I. 227 (1826).
STENOTUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 384.
acaulis, *A. Gray, in Proc. Am. Acad.* vii. (1868) 353. —Calif.
acerosus, *Bert. ex Phil. in Linnæa*, xxvii. (1856) 724. —Chili.
acuminatus, DC. Prod. v. 348 = Leucopsis sericea.
alpigenus, Torr. & Gray, Fl. N. Am. ii. 241 = Aster pulchellus.
altenifolius, Steud. Nom. ed. II. i. 87, sphalm. = chrysanthemifolius.
anthylloides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 561.—Chili.
asparagioides, *A. Gray, in Proc. Am. Acad.* vii. (1868) 354.—Calif.
araucanus, *Phil. in Anal. Univ. Chil.* (1873) 491.— Chili.
arbuscides, *Remy, in C. Gay, Fl. Chil.* iv. 58.—Chili.
arenarius, *Benth. Bot. Voy. Sulph.* 34 (Asteris sp. ?).— Calif.
argenteus, *Steud. Nom.* ed. II. i. 112.—Chili.

APLOPAPPUS :—
armerioides, *A. Gray, Color. Exp. Surv. Bot.* (1860) 16.—N. Mexic.
aureus, *A. Gray, Pl. Fendl.* 76.—Texas.
baccharoides, Benth. Bot. Voy. Sulph. 34 = Solidago occidentalis.
Bayldonii, *Remy, in C. Gay, Fl. Chil.* iv. 49.—Chili.
belitidioides, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 113 = Erigeron pappochroma.
Berterii, *DC. Prod.* v. 346.—Chili.
blepharophyllus, A. Gray, Pl. Wright. i. 97 = Aster gymnocephalus.
Bloomeri, *A. Gray, in Proc. Am. Acad.* vi. (1865) 541.—Am. bor.
Brandegei, *A. Gray, Syn. Fl. N. Am.* i. 11. 192.—Am. bor.
buxifolioides, *Remy, in C. Gay, Fl. Chil.* iv. 49.—Chili.
caespitosus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 380.—Chili.
calendulaceus, *Griseb. in Gortt. Abh.* xxiv. (1879) 179. —Reg. Argent.
canescens, DC. Prod. v. 349 = Chrysopsis villosa.
carthamoides, *A. Gray, in Proc. Acad. Sc. Philad.* (1863) 65.—Am. bor.
cervinus, *Rothrock, in Wheeler, Geogr. Surv.* 40. t. 6.— Am. bor.
cervinus, S. Wats. in Am. Natural. vii. (1873) 301 = nanus.
Chamissonis, *DC. Prod.* v. 347.—Chili.
chrysanthemifolius, *DC. l. c.* 348.—Chili.
ciliatus, *DC. l. c.* 346.—Am. bor.
coronopifolius, *DC. l. c.* 347.—Chili.
Coulteri, Harv. & Gray, ex A. Gray, Pl. Fendl. 75 = spinulosus.
croceus, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 65.—Am. bor.
cuneatus, *A. Gray, in Proc. Am. Acad.* viii. (1873) 635. —Calif.
cuneifolius, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 380.—Chili.
decurrens, *Remy, in C. Gay, Fl. Chil.* iv. 58.—Chili.
denudatus, *Remy, l. c.* 53.—Chili.
diffusus, DC. Prod. v. 349 = Leucopsis diffusa.
Diplopappus, *Remy, in C. Gay, Fl. Chil.* iv. 53.—Chili.
discoideus, DC. Prod. v. 350 = Bigelovia veneta.
divaricatus, *A. Gray, in Pacif. Rail. Rep.* iv. 99.— Am. bor.
Doellianus, *Phil. in Anal. Univ. Chil.* (1873) 489.— Chili.
ericoides, *DC. Prod.* v. 347.—Calif.
filifolius, DC. Prod. vii. 280 = Erigeron filifolius ?
flexifer, Hook. & Arn. Bot. Beech. Voy. 351 = Townsendia florifera.
foliosus, *DC. Prod.* v. 346.—Chili.
Fremontii, *A. Gray, in Proc. Acad. Sc. Philad.* (1863) 65.—Am. bor.
glabratus, *Phil. in Linnæa*, xxvii. (1856) 727.—Chili.
glutinosus, Cass. ex DC. Prod. v. 347 = coronopifolius, pulchellus.
glutinosus, *DC. Prod.* v. 347.—Chili.
gracilis, *A. Gray, Pl. Fendl.* 76.—Texas.
graminifolius, Benth. Pl. Hartw. 65 (err. typ. 61) = Chrysopsis graminifolia.
Groenei, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 80. —Am. bor.
grindelioides, *DC. Prod.* v. 348.—Chili.
Gunnii, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 111 = Erigeron pappochroma.
gymnocephalus, DC. Prod. v. 346 = Aster gymnocephalus.
Harnhei, DC. l. c. 349 = Corethrogyne filaginifolia.
Hallii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 389. —Oregon.
hirtellus, *Phil. in Linnæa*, xxvii. (1856) 726.—Chili.
hirtus, *A. Gray, Syn. Fl. N. Am.* i. 11. 127.—Am. bor.
hispidulus, *DC. Prod.* v. 348.—Chili.
Hookerianus, *A. Gray, Syn. Fl. N. Am.* i. 11. 131.— Texas.
Hookerianus, *DC. Prod.* vii. 279.—Chili.
Hookerinnus, A. Gray, Pl. Lindh. i. 40 = divaricatus.
Hoorebekia, *DC. Prod.* v. 346.—Chili.
Howellii, *A. Gray, Syn. Fl. N. Am.* i. 11. 446.—Oregon.
hypoleucus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 177 = Bigelovia hypoleuca.
illecifolius, *Remy, in C. Gay, Fl. Chil.* iv. 55.—Chili.
illinitus, *Phil. in Linnæa*, xxvii. (1856) 727.—Chili.
integrifolius, *T. C. Porter, in A. Gray, in Proc. Am. Acad.* xvi. (1881) 79.—Am. bor.

APLOPAPPUS :—
inuloides, Torr. & Gray, Fl. N. Am. ii. 241 = uniflorus.
junceus, Greene, in Bull. Acad. Calif. i. (1884) 190 = spinulosus.
lanatus, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 215. —Ins. Galapag.
lanceolatus, *Torr. & Gray, Fl. N. Am.* ii. 241.—Calif.
lanuginosus, *A. Gray, Bot. U. St. Expl. Exped.* ii. 347. —Am. bor.
laricifolius, *A. Gray, Pl. Wright.* ii. 80.—N. Mexic.
Lastarriaeus, *Remy, in C. Gay, Fl. Chil.* iv. 45.—Chili.
linearifolius, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 457 = Baccharis texana ?.
linearifolius, *DC. Prod.* v. 347.—Calif.
Lyallii, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 64.—Am. bor.
Macraeanus, *A. Gray, in Proc. Am. Acad.* v. (1861) 132.—Chili.
macrocephalus, *DC. Prod.* v. 348.—Chili.
Macronemus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 542.—Am. bor.
marginalis, *Phil. in Linnæa*, xxxiii. (1864–65) 139.— Chili.
marginatus, *Griseb. Cat. Pl. Cub.* 149.—Cuba.
medicinalis, *Phil. in Linnæa*, xxvii. (1856) 729.— Chili.
Menziesii, Torr. & Gray, Fl. N. Am. ii. 241 = Bigelovia Menziesii.
Meyenii, *Walp. in Nov. Act. Cur.* xix. *Suppl.* i. (1843) 261.—Chili.
mollis, A. Gray, in Proc. Am. Acad. xvi. (1881) 80 = Greenei.
monactis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 1. —Calif.
mucronatus, *Hook. & Arn. Bot. Beech. Voy.* 147.—Chili.
multicaulis, *A. Gray, in Am. Natural.* vii. (1874) 213. —Am. bor.
nanus, *Eaton, in Bot. King, Exped.* 159.—Am. bor.
nevadensis, *Kellogg, in Proc. Calif. Acad.* iii. (1863) 9. —Calif.
Nuttallii, *Torr. & Gray, Fl. N. Am.* ii. 242.—Am. bor.
ochagavianus, *Phil. in Linnæa*, xxvii. (1856) 725.— Chili.
Orcutti, *A. Gray, in Proc. Am. Acad.* xx. (1885) 297.— Am. bor.
Palmeri, *A. Gray, l. c.* xi. (1876) 74.—Calif.
paniculatus, A. Gray, l. c. vii. (1868) 354 = racemosus.
paniculatus, *Phil. in Anal. Univ. Chil.* (1873) 490.— Chili.
pappochromus, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 111 = Erigeron pappochroma.
Parryi, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 289.—Am. bor.
parvifolius, *A. Gray, in Proc. Am. Acad.* v. (1861) 122.—Chili.
patagonicus, *Phil. in Linnæa*, xxvii. (1856) 726.— Chili.
pauciclematus, *Phil. in Anal. Univ. Chil.* (1873) 492.—Chili.
pectinatus, *Phil. in Linnæa*, xxxiii. (1864–65) 140.— Chili.
pedunculosus, *Remy, in C. Gay, Fl. Chil.* iv. 51.— Chili.
phyllocephalus, DC. Prod. v. 347 = rubiginosus.
pinifolius, *A. Gray, in Proc. Am. Acad.* viii. (1873) 636.—Calif.
pinnatifidus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 380.—Am. bor.
Poeppigianus, *A. Gray, in Proc. Am. Acad.* v. (1862) 121.—Chili.
polyphyllus, *Phil. in Linnæa*, xxx. (1859–60) 192.— Chili.
pumeliodes, *DC. Prod.* v. 346.—Chili.
pulchellus, *DC. l. c.* 347.—Chili.
pygmaeus, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 289.—Am. bor.
racemosus, *Torr. in Stkgrove. Rep.* 162.—Am. bor.
radianus, *Remy, in C. Gay, Fl. Chil.* iv. 45.—Chili.
ramulosus, DC. Prod. v. 350 = Baccharis ramulosa.
reflexus, *Phil. in Linnæa*, xxvii. (1856) 728.—Chili.
Remyanus, *Wedd. Chlor. And.* i. 210.—Chili.
Rengifoanus, *Remy, in C. Gay, Fl. Chil.* iv. 48.— Chili.
resinosus, A. Gray, Bot. U. St. Expl. Exped. ii. 348 = nanus.
reticulatus, *Phil. in Linnæa*, xxvii. (1856) 728.— Chili.

APLOPAPPUS :—
rigidus, *Phil. Fl. Atac.* 30.—Chili.
rubiginosus, *Torr. & Gray, Fl. N. Am.* ii. 240.—Am. bor.
scaposus, DC. Prod. v. 350 = Leucopsis scaposa.
scaposus, *Remy, in C. Gay, Fl. Chil.* iv. 50.—Chili.
scrobiculatus, *DC. Prod.* v. 348.—Chili.
sericeus, *DC. l. c.* 349.—Chili
sericeus, Phil. in Linnæa, xxviii. (1856) 724 = Poeppigianus.
sphærocephala, *Harv. & Gray, ex A. Gray, Pl. Fendl.* 76.—Calif.
spinulosus, *Phil. in Anal. Univ. Chil.* (1878) 490.—Chili.
spinulosus, *DC. Prod.* v. 347.—Am. bor.
squarrosus, *Hook. & Arn. Bot. Beech. Voy.* 146.—Calif.
stellatus, Hook. f. in Lond. Journ. Bot. vi. (1847) 112 = Erigeron pappochroma.
stelliger, *Remy, in C. Gay, Fl. Chil.* iv. 52.—Chili.
stenophyllus, *A. Gray, Bot. U. St. Expl. Exped.* ii. 347.—Calif.
stoloniferus, *DC. Prod.* v. 349.—Mexic.
suffruticosus, Eaton, in Bot. King, Exped. 159 = Watsonii.
suffruticosus, *A. Gray, in Proc. Am. Acad.* vi. (1866) 542.—Calif.
tanacetus, Hook. f. in Lond. Journ. Bot. vi. (1847) 110 = Erigeron pappochroma.
tenuicaulis, Eaton, in Bot. King, Exped. 160 = lanceolatus.
tortifolius, Torr. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 109 = Aster tortifolius.
uncinatus, *Phil. in Linnæa*, xxviii. (1856) 728.—Chili.
uniflorus, *Torr. & Gray, Fl. N. Am.* ii. 241.—Am. bor.
velutinus, *Remy, in C. Gay, Fl. Chil.* iv. 57.—Chili.
villosus, *Phil. in Linnæa*, xxviii. (1856) 725.—Chili.
virgatus, *Phil. in Anal. Univ. Chil.* (1873) 491.—Chili.
Watsonii, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 79.—Am. bor.
Whitneyi, *A. Gray, l. c.* vii. (1868) 353.—Calif.

APLOPHYLLUM, Cass in Dict. Sc. Nat. xxxiii. 472 (1824) = **Mutisia**. Linn. (Compos.).
decurrens, Cass. l. c. = M. decurrens.
ilicifolia, Cass. l. c. = M. ilicifolia.
linearifolium, Cass. l. c. = M. linearifolia.

APLOPHYLLUM, A. Juss. in Mém. Mus. Par. xii. (1825) 464 = **Ruta**, Linn.
acutifolium, G. Don, Gen. Syst. i. 780 = R. acutifolia.
arabicum, Boiss. Diagn. Ser. I. viii. 127 = R. tuberculata.
arbusculum, Franch. in Revoil, Comali, 38.—Afr. trop.
armenum, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 179 = R. suaveolens?
Besseri, Spach, l. c. = R. Besseri
Bieberstenii, Spach, l. c. 178 = R. Buxbaumii.
Blanchei, Boiss. Fl. Orient. i. 937.—Orient.
Boissierianum, Viv. & Panč. in Mem. Ist. Venet. xv. (1870) 14 = R. patavina.
Bourgæi, Boiss. Fl. Orient. i. 928.—Armenia.
Broussonetianum, Cosa. in Bull. Soc. Bot. Fr. xx. (1873) 244 ; xxii. (1875) 86, nomen.—Afr. bor.
Buhsei, Boiss. Fl. Orient. i. 926.—Persia.
Bungei, Trautv. in Act. Hort. Petrop. vlii. (1883) 188.—Turkestan.
Buxbaumii, G. Don, Gen. Syst. i. 780 = R. Buxbaumii.
Buxbaumii, Schur, Enum. Pl. Transs. 140 = R. suaveolens.
canaliculatum, Boiss. Diagn. Ser. I. viii. 128.—Persia.
Candollianum, Spach, ex Jaub. & Spach, Illustr. Pl. Or. t. 270 = R. glabra.
cappadocicum, Boiss. Fl. Orient. i. 963.—Cappadoc.
Chesneyanum, Boiss. Fl. Orient. i. 941.—Mesopot.
ciliatum, Griseb. Spicil. Fl. Rumel. i. 130 = R. suaveolens.
ciliicum, Boiss. Fl. Orient. i. 927.—Cilicia.
congestum, Jaub. & Spach, Illustr. Pl. Or. iii. 77 = R. suaveolens.
consanguineum, Bunge, ex Boiss. Fl. Orient. i. 941.—Persia.
cordatum, G. Don, Gen. Syst. i. 780 = R. cordata.
coronatum, Griseb. Spicil. Fl. Rumel. i. 129 = R. patavina.
corymbulosum, Boiss. Diagn. Ser. II. i. 116 = R. Buxbaumii.

APLOPHYLLUM :—
cremulatum, Boiss. Diagn. Ser. II. i. 117.—Beluchist.
dahuricum, Juss. in Mém. Mus. Par. xii. (1825) 464 = Ruta dahurica.
dauricum, Ledeb. Fl. Ross. i. 492 = præc.
dichotomum, G. Don, Gen. Syst. i. 780 = R. dichotoma.
dichotomum, Griseb. Spicil. Fl. Rumel. i. 131 = R. Buxbaumii.
elatum, Fisch. & Mey. ex Kar. in Bull. Soc. Nat. Mosc. (1889) 149.—Turcomania.
erythræum, Boiss. Diagn. Ser. II. i. 114.—Afghan.
flexifolium, Spach, ex Jaub. & Spach, Illustr. Pl. Or. t. 265.—Persia.
flexuosum, Boiss. Diagn. Ser. II. i. 118.—Afghan.
fruticulosum, G. Don, Gen. Syst. i. 780 = R. fruticulosa.
furfuraceum, Bunge, in Boiss. Fl. Orient. i. 934.—Persia.
glaberrimum, Bunge, l. c. 930.—Persia.
glabrum, G. Don, Gen. Syst. i. 780 = R. glabra.
Griffithianum, Boiss. Diagn. Ser. II. i. 115.—Afghan.
halepense, Spach, in Jaub. & Spach, Illustr. Pl. Or. t. 266 = R. Buxbaumii.
hirsutum, Regel & Schmalh. ex Regel, Pl. Nov. Fedsch. 17.—As. temp. or.
huzsanicum, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 176 = R. linifolia.
kispidulum, Bunge, in Boiss. Fl. Orient. i. 933.—Persia.
Kotschyi, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 186.—Persia.
Kowalenskyi, Stachég. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 467.—Reg. Caucas.
lasiantherum, Bunge, in Schrift. Naturf Ver. Riga, i. (1847) 206 ; et Pl. Lehm. 289.—Turkest.
latifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 898.—As. cent.
linearc, G. Don, Gen. Syst. i. 780 = R. linearis.
linifolium, G. Don, l. c. = R. linifolia.
linifolium, Reichb. ex Nym. Consp. 142 = R. patavina.
linifolium, Schur, Enum. Pl. Transs. 140 = R. suaveolens.
longifolium, Boiss. Diagn. Ser. I. viii. 127.—Palæstin.
mesopotamicum, Boiss. l. c. Ser. II. i. 116 = R. Buxbaumii.
myrtifolium, Boiss. l. c. Ser. I. i. 63.—As. Min.
obtusifolium, Ledeb. Fl. Ross. i. 490 = R. obtusifolia.
patavinum, G. Don, Gen. Syst. i. 780 = R. patavina.
pedicellatum, Bunge, ex Boiss. Fl. Orient. i. 925.—Persia.
perforatum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 897 = R. acutifolia.
pilosum, Franch. in Ann. Sc. Nat. Sér. VI. xv. (1883) 248, t. 18.—Turkest.
pilosum, Lescegl. ex Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 278.—Persia.
propinquum, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 189.—Mesopot.
ptilostylum, Spach, ex Jaub. & Spach, Illustr. Pl. Or. t. 267.—Persia.
pubescens, Boiss. Voy. Bot. Espagne, i. 371 = R. linifolia.
pubescens, G. Don, Gen. Syst. i. 780 = R. pubescens.
pumilum, Boiss. Diagn. Ser. I. i. 62.—As. Min.
Reuteri, Boiss. l. c. i. 87.—Caria.
robustum, Bunge, in Schrift. Naturf. Ver. Riga, i. (1847) 208, et Pl. Lehm. 289.—Persia.
rosmarinifolium, G. Don, Gen. Syst. i. 781 = R. rosmarinifolia.
Sieversii, Fisch. in Schrenk, Enum. Pl. Nov. 89 = R. acutifolia.
spathulatum, Reichb. ex Nym. Consp. 143 = R. Buxbaumii.
Stockeianum, Boiss. Diagn. Ser. II. i. 117.—Beluchist.
suaveolens, G. Don, Gen. Syst. i. 780 = R. suaveolens.
suaveolens, Sweet, Hort. Brit. ed. III. 126 = R. Buxbaumii.
subcordatum, Bunge, in Boiss. Fl. Orient. i. 932.—Persia.
sulphureum, Boiss. Diagn. Ser. I. v. 87.—As. Min.
sybroticum, Boiss. l. c. viii. 126.—Syria.
tauricum, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 178 = R. taurica.
Tchihatchewi, Boiss. Diagn. Ser. II. vi. 41 = *sulphureum*.
telephioides, Boiss. l. c. i. 929.—Cappadoc.
tenue, Boiss. l. c. 932.—Armenia.
thesioides, G. Don, Gen. Syst. i. 780 = R. thesioides.

APLOPHYLLUM :—
trichostylum, Bunge, in Boiss. Fl. Orient. i. 941.—Persia.
tuberculatum, Juss. in Mém. Mus. Par. xii. (1825) 528 t. 17 = R. tuberculata.
versicolor, Fisch. & Mey. in Bull. Phys. Math. Akad. Petersb. iii. (1846) 307 = *lazaretium*.
villosulum, Boiss. & Hausskn. in Boiss. Fl. Orient. i. 936.—Syria.
villosum, G. Don, Gen. Syst. i. 780 = R. villosa.
virgatum, Spach, ex Jaub. & Spach, Illustr. Pl. Or. t. 267.—Persia.
vulcanicum, Boiss. & Heldr. Diagn. Ser. I. viii. 126.—Lycaonia.

APLOSTELLIS, Thou. Orch. Iles Afr. t. 24 (1822) = **Pogonia**, Juss. (Orchid.).
ambigua, A. Rich. in Mém. Soc. Hist. Nat Par. iv. (1828) 36 = P. Thouarsii.
simplex, Thou. Orch. Isles Afr. t. 24 = P. Thouarsii.
truncata, Lindl. Gen. et Sp. Orch. 411.—Ins. Mascar.

APLOSTEMON, Rafin. in Journ. Phys. lxxxix. (1819) 105 = **Scirpus**, Linn. etc. (Cyper.).
atropurpureum, Rafin. l. c. = Elecharis atropurpurea.
bracteatum, Rafin. l. c. = S. bracteatus.
compressum, Rafin. l. c. = S. Caricis.
polytrichoides, Rafin. l. c. = Fimbristylis polytrichoides.
triquetrum, Rafin. l. c. = Fimbristylis argentea.

APLOSTYLIS, Rafin. Fl. Tellur. iv. 91 (1836) = **Cuscuta**, Linn. (Convolv.).
lupuliformis, Rafin. l. c. = C. lupuliformis.

APLOTAXIS, DC. in Guill. Archiv. Bot. ii. (1833) 330 = **Saussurea**, DC. (Compos.).
albescens, DC. Prod. vi. 540 = S. albescens.
andryaloides, DC. l. c. 549 = S. sorocephala.
auriculata, DC. l. c. 541 = S. hypoleuca.
australasica, F. Muell. Fragm. i. 36 = S. affinis.
Bungei, DC. Prod. vi. 539 = S. affinis.
Bungei, Less. ex Ledeb. Fl. Ross. ii. 758 = Leuzea salina.
caespitosa, DC. Prod. vi. 539 = S. taraxacifolia.
candicans, DC. l. c. 540 = S. candicans.
Candolleana, DC. l. c. 541 = S. Candolleana.
carthamoides, Ham. in DC. l. c. 540 = S. affinis.
circinales, DC. l. c. = Cnicus arvensis.
deltoides, DC. l. c. 541 = S. deltoides.
denticulata, DC. l. c. 539 = S. denticulata.
discolor, DC. l. c. 541 = S. pipathera.
fastuosa, Decne. in Jacquem. Voy. Bot. 97. t. 105 = S. denticulata.
folosa, Edgew. in Trans. Linn. Soc. xx. (1846) 77 = S. foliosa.
Frolovii, DC. Prod. vi. 538 = S. Frolovii.
graphalodes, Royle, Illustr. Bot. Himal. 251. t. 59. f. 1 = S. sorocephala.
gossypina, DC. Prod. vi. 541 = S. gossypiphora.
involucrata, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 389.—As. temp.
Jacea, Klotzsch, in Bot. Ergeb. Waldem. Reise, 79. t. 79 = S. Jacea.
Lappa, Decne. in Jacquem. Voy. Bot. 96. t. 104 = S. Lappa.
leontodontoides, DC. Coll. Mém. ix. 29. t. 10 (Prod. vi. 539) = S. Kunthiana.
multicaulis, DC. in Deless. Ic. Sel. iv. t. 68 = S. affinis.
nepalensis, DC. l. c. t. 67 = S. Eriostemon.
niwea, DC. Prod. vi. 541 = S. deltoidea.
obvallata, DC. in Deless. Ic. Sel. iv. t. 69 = S. obvallata.
pungens, DC. Prod. vi. 539 = Cnicus arvensis.
Royleı, DC. Prod. vi. 538 = S. Royleı.
sacrana, Edgew. in Trans. Linn. Soc. xx. (1846) 76 = S. candicans.
Simpsoniana, Field. & Gardn. Sert. Pl. t. 36 = S. andryaloides.
sorocephala, Schrenk, Enum. Pl. Nov. (1841) 43 = S. sorocephala.
taraxacifolia, DC. Prod. vi. 539 = S. taraxacifolia.
uniflora, DC. l. c. 539 = S. uniflora.

APLOTHECA, Mart. ex Cham. in Linnæa, v. (1830) 91 = **Froelichia**, Moench. (Amarant.).
interrupta, Mart. ex Cham. l. c. = F. interrupta.

APLUDA, Linn. Sp. Pl. ed. i. 82 (1753). *GRAMINEAE*, Benth. & Hook. f. iii. 1137.
CALAMINA, Beauv. Agrost. 128. t. 23. f. 1 (1812).
DICTOSIS, Beauv. l. c. 132. t. 23. f. 8 (1812).
aristata, Linn. Cent. ii. 7; Sp. Pl. ed. II. 1487 = mutica.
ciliata, *Anderss. in Oefver. Vet. Acad. Stockh.* (1855) 177.—Ind. or.
communis, Nees & Arn. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 194 = mutica.
Cumingii, Buese, ex De Vriese, Pl. Ind. Bat. Reinw. 105 = mutica.
digitata, Linn. f. Suppl. 434 = Polytoca bracteata.
distachya, Beauv. Agrost. 159, 151 = Andropogon distachyus.
geniculata, Roxb. Hort. Beng. 8; Fl. Ind. i. 327 = mutica.
gigantea, Spreng. Syst. i. 290 = mutica.
glauca, Schreb. Graes. ii. 99 = mutica.
Gryllus, Beauv. Agrost. 133 = mutica.
Gryllus, Presl, Cyp. & Gram. Sic. 55 = Chrysopogon Gryllus.
humilis, Kunth, Enum. Pl. i. 517 = mutica.
imberbis, Steud. Nom. ed. II. i. 251, sphalm. = Anthistiria imberbis.
inermis, Regel, in Act. Hort. Petrop. vii. (1880) 658 = mutica.
Kobilia, Buch.-Ham. ex Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 194 = mutica.
microstachya, Nees, l. c. 108 = mutica.
mucronata, Steud. Syn. Pl. Gram. 404 = mutica.
mutica, *Linn. Sp. Pl.* 82.—Aa. trop.; Polynes.; Austral.
pedicellata, Buese, ex De Vriese, Pl. Ind. Bat. Reinw. 105 = mutica.
rostrata, Arn. & Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 194 = mutica.
scabra, Anderss. in Oefver. Vet. Acad. Stockh. (1855) 179 = mutica.
scirpoides, Walt. Fl. Carol. 250 (pl. dubia).—Carolina.
villosa, Schreb. Gram. ii. 92 = mutica.
Zeugites, Linn. Syst. ed. X. 1305; Pl. Jam. Pugill. 30 = Zeugites americana.
Zeygites, Aubl. Pl. Gui. ii. 933 = Ichnanthus pallens.

APOBALLIS, Schott. in Oestr. Bot. Zeitschr. (1858) 318 = **Schizmatoglottis**, Zoll. (Aroid.).
neglecta, Schott, l. c. 317 = S. rupestris.

APOCHORIS, Duby, in DC. Prod. viii. 67 (1844). *PRIMULACEAE*, Benth. & Hook. f. ii. 685.
pentapetala, *Duby, l. c.*—China.

APOCOPIS, Nees, in Proc. Linn. Soc. i. (1841) 93. *GRAMINEAE*, Benth. & Hook. f. iii. 1128.
AMBLYACHYRUM, Hochst. ex Steud. Syn. Pl. Gram. 418 (1855).
Beckeriii, Thw. ex Trimen, Cat. Pl. Ceylon, 107 = Wightii.
Royleana, *Nees, in Proc. Linn. Soc.* i. (1841) 94.—Ind. or.
tridentata, *Benth. in Journ. Linn. Soc.* xix. (1882) 67.—Ind. or.
Wightii, *Nees, ex Steud. Syn. Pl. Gram.* 377.—Ind. or.; China.
Wrightii, Munro, in Proc. Am. Acad. iv. (1864–65) 363 = Wightii.

APOCOPSIS, Meisn. Gen. 429 (828) (1843) = **Apocopis**, Nees, Gram.).

APOCYNUM, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 65 (1737). *APOCYNACEAE*, Benth. & Hook. f. ii. 716.
QUIRIVELIA, Poir. Encyc. vi. 42 (1804).
Acouci, Aubl. Pl. Gui. i. 274 = Forsteronia Acouci.
africanum, *Lour. Fl. Cochinch.* i. 168 (Quid ?).—Zanzib.
agglomeratum, *Poir. Encyc. Suppl.* i. 407.—Ins. S. Doming.
alterniflorum, Lour. Fl. Cochinch. i. 168 = Strophanthus alterniflorus.
androsaemifolium, *Linn. Sp. Pl.* 213.—Am. bor.
bursiforium, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 5.—Malaya.
ca. ense, *Lam. Encyc.* i. 312.—Ins. Canar.
cannabinum, *Linn. Sp. Pl.* 213.—Am. bor.
citrefolium, Descourt. Fl. Med. Antill. iii. 180. t. 191 = Echites citrina.
compressum, Moench, Meth. 464 = venetum.
APOCYNUM:—
cordatum, Mill. Gard. Dict. ed. VIII. n. 10 = Rhabdadenia cordata.
cordatum, Thunb. Prod. Pl. Cap. 47 = Astephanus cordatus.
cordifolium, Tenore, ex Steud. Nom. ed. II. i. 118 = prasc.
cotunifolium, *Lam. Encyc.* i. 315.—Java.
crassifolium, Salisb. Prod. 149 = Ichnocarpus frutescens.
cuneatum, Willd. ex Roem. & Schult. Syst. vi. 706 = Metastes fimbriatum.
erectum, Vell. Fl. Flum. 128; iii. t. 87 = Barjonia racemosa.
filiforme, Thunb. Prod. Pl. Cap. 47 = Eustegia filiformis.
floristratum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 5 = Strophanthus dichotomus.
foetidum, Burm. f. Fl. Ind. 71 = Paederia foetida.
frutescens, Afzel. Kemed. Guin. Coll. iv. 28 = Ichnocarpus Afzelii.
frutescens, Linn. Sp. Pl. 213 = Ichnocarpus frutescens
hastatum, Thunb. Prod. Pl. Cap. 47 = Eustegia hastata.
Hendersonii, *Hook. f. in Hume & Henders. Lahore to Yarkand*, 327.—Thibet.
hypericifolium, Ait. Hort. Kew. ed. I. i. 304 = cannabinum.
indicum, Lam. Encyc. i. 314 = reticulatum.
Jasminiflorum, Lour. Fl. Cochinch. 167 = Tylophora ovata.
lanceolatum, Thunb. Prod. Pl. Cap. 47 = Astephanus lanceolatus.
littoreum, Linn. f. Suppl. 169 = Astephanus linearis.
maculatum, Descourt. Fl. Med. Antill. iii. 177. t. 190 = Echites maculata.
mucronatum, Linn. f. Suppl. i. 169 = Eustegia hastata.
mucronatum, Blanco, Fl. Filip. ed. I. 302 = Toxocarpus gracilis.
muscipulum, Moench, Meth. 464 = androsaemifolium.
nervosum, *Mill. Gard. Dict.* ed. VIII. n. 9.—Vera Crux.
obliquum, Mill. l. c. n. 8 = Echites umbellata.
odoratissimum, *Lour. ex Poir. In. Ind.* 83.—Malaya.
oriense, Rottler, ex Hook. f. Fl. Brit. Ind. iv. 6 = Cryptolepis elegans.
paniculatum, Lam. Encyc. i. 314 = Forsteronia Acouci.
pictum, *Schenk, in Bull. Phys. Math. Acad. Petersb.* ii. (1844) 116.—Soongaria.
piscatorium, Dougl. ex A. DC. in DC. Prod. viii. 439 = cannabinum.
platyanthum, Salzm. ex A. DC. l. c. 594 = Philibertia cynanchoides.
pubescens, R. Br. in Mem. Wern. Soc. i. (1809) 68 = cannabinum.
purpureum, *Tausch, in Flora*, xix. (1836) ii. 385.—Am. bor.
reclinatum, Salzm. ex Decne. in DC. Prod. viii. 602 = Blepharodon pallidum.
reticulatum, Herb. Madras, ex Wall. Cat. n. 8247 = Decalepis Hamiltonii.
reticulatum, *Linn. Sp. Pl.* 214.—Ind. or.
reticulatum, Lour. Fl. Cochinch. 208 = Parsonsia Helicandra.
salicifolium, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 406 = Gomphocarpus fruticosus.
salicifolium, *Willd. ex Roem. & Schult. Syst.* iv. 796.—Ins. S. Doming.
scandens, Mill. Gard. Dict. ed. VIII. n. 5 = Echites asperuginis.
sibiricum, Jacq. Hort. Vindob. iii. 37 = cannabinum.
sibiricum, Pall. ex Roem. & Schult. Syst. iv. 406 = venetum.
soleare, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 5.—Malaya.
speciosissimum, Mill. Gard. Dict. ed. VIII. n. 4 (= *Haemadictyon suberectum*).
stellatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 5.—Malaya.
stylospermum, *Noronha, l. c.*—Malaya.
syriacum, S. G. Gmel. Reise, ii. 198, 257 = Calotropis procera.
tiliaefolium, Lam. Encyc. i. 314 = Dregea volubilis.
trifforum, Linn. f. Suppl. 169 = Astephanus trifforus.
umbellatum, Aubl. Pl. Gui. i. 276 = Tbamadia umbellata.
umbellatum, Salzm. ex Decne. in DC. Prod. viii. 585 = Oxypetalum denticulosum.
venetum, A. DC. Prod. viii. 440 = cannabinum.
venetum, *Linn. Sp. Pl.* 213.—*Reg. Mediterr.*; Orient.; As. bor.; Ind. or.

APOCYNUM:—
villosum, *Mill. Gard. Dict.* ed. VIII. n. 11.—Vera Crux.
viminale, Bassi, in Comm. Bonon. (1755) ex Roem. & Schult. Syst. vi. 113 = Sarcostemma viminale.
vincaefolium, *Burm. f. Fl. Ind.* 71.—Ind. or.
volubile, Vell. Fl. Flum. 128; iii. t. 88 = Araujia calycina.

APODANTHERA, Arn. in Hook. Journ. Bot. iii. 274 (1841). *CUCURBITACEAE*, Benth. & Hook. f. i. 834.
argentea, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1878) 42.—Bras.
aspera, *Cogn. l. c.* 43.—Mexic.
bifora, *Cogn. l. c.*—Ecuador.
Harnieri, *Cogn. l. c.* 44.—Mexic.
Galeottii, *Cogn. l. c.* 45.—Mexic.
gracilis, *Benth. Bot. Voy. Sulph.* 99 = Melothria fluminensis.
lacviuscula, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1878) 12.—Bras.
Mandonii, *Cogn. l. c.* 41.—Bolivia.
Mathewsi, *Arn. in Hook. Journ. Bot.* iii. (1841) 274.—Peruv.
mucronata, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1878) 40.—Peruv.
pedisecta, *Cogn. l. c.* 17.—Bras.
scabra, *Cogn. in Bull. Acad. Belg. Sér.* II. xlix. (1880) 192.—Ecuador.
smilacifolia, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1878) 42.—Bras.
undulata, *A. Gray, Pl. Wright.* ii. 60.—N. Mexic.

APODANTHES, Poit. in Ann. Sc. Nat. Sér. I. iii. (1824) 422. t. 26. *CYTINACEAE*, Benth. & Hook. f. iii. 118.
FROSTIA, Bertero, ex Guill. in Ann. Sc. Nat. Sér. II. ii. (1834) 19, in nota.
PILOSTYLES, Guill. in Ann. Sc. Nat. Sér. II. ii. (1834) 21. t. 3.
SARNA, Karst. in Nov. Act. Nat. Cur. xxvi. (1858) 920. t. 65.
Berteri, *Gardn. in Hook. Jc. Pl.* t. 655.—Chili.
Blanchetti, *Gardn. l. c.* t. 655 b.—Bras.
Calliandrae, *Gardn. l. c.* t. 844.—Bras.
Cascarae, *Poit. in Ann. Sc. Nat.* Sér. I. iii. (1824) 422. t. 26.—Am. austr.
Flacourtiae, *Karst. in Linnaea*, xxviii. (1856) 413.—Venezuela.

APODOCEPHALA, Baker, in Journ. Linn. Soc. xvi. (1876) 417. *COMPOSITAE*.
pauciflora, *Baker, l. c.*—Madag.

APODOLIRION, Baker, in Journ. Bot. xvi. (1878) 74. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 722.
Bolusii, *Baker, l. c.* 75.—Afr. austr.
Buchanani, *Baker, l. c.*—Natal.
lancifolium, *Benth. & Hook. f. Gen.* iii. 722.—Afr. austr.
Mackenii, *Baker, in Journ. Bot.* xvi. (1878) 75.—Natal.

APODOSTACHYS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 857 = **Brotlia**, A. Juss. (Phytolac.).
densiflora, Turcz. l. c. = E. volubilis.

APODYNOMENE, E. Mey. Comm. Pl. Afr. Austr. iii. (1835) = **Euphrosia**, Pers. (Legum.).
asmula, E. Mey. l. c. 118 = T. asmula.
diffusa, E. Mey. l. c. = T. diffusa.
grandiflora, E. Mey. l. c. 113 = T. grandiflora.
macropoda, E. Mey. l. c. 113 = T. macropoda.
Meyeri, Presl, Bot. Bemerk. 57 = T. grandiflora.

APODYTES, E. Mey. ex Bernh. in Linnaea, xii. (1838) 126; et ex Am. in Hook. Journ. Bot. iii. (1841) 155. *OLACINEAE*, Benth. & Hook. f. i. 351.
NOTHAPODYTES, Blume, Mus. Bot. Lugd. Bat. i. 248 (1850).
RHAPHIOSTYLIS, Planch. ex Benth. in Hook. Niger Fl. 389 (1849).
acutifolia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 92 = dimidiata.
andamanica, *Kurz, Andam. Rep.* 33.—Ins. Andam.
Beddomei, *Mast. in Hook. f. Fl. Brit. Ind.* i. 688.—Ind. or.
bonincusis, *Hook. f. in Hook. Jc. Pl.* t. 778.—Afr. trop.

APODYTES :—
Benthamiana, *Wight, Ic.* t. 1158.—Ind. or.
brachycalyx, *F. Muell. Fragm.* ix. 149.—Austral.
dimidiata, *E. Mey. ex Bernh. in Linnaea*, xii. (1838) 136.—Afr. austr.
Gardneriana, Miers, in Ann. & Mag. Nat. Hist. Ser. II. ix. (1852) 389 = Benthamiana.
grandifolia, *Benth. & Hook. f. Gen.* i. 852.—Madag.
nauritiana, *Benth. & Hook. f. l. c.*— Ins. Mascar.

APOGANDRUM, Neck. Elem. i. 216 (1790) = **Erios**, Linn.

APOGETON, Schred. ex Steud. Nom. ed. II. i. 114 (1840) = **Aponogeton**, Linn. f. (Naiad.).

APOGON, Ell. Sketch, ii. 267 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 509.
LAXANON, Rafin. New Fl. Am ii. 16 (1836).
gracilis, *DC. Prod.* vii. 79.— Am. bor.
humilis, *Ell. Sketch*, ii. 267.—Am. bor.
humilis, Nutt. in Trans. Am. Phil. Soc. N. s. vii. (1841) 424 = gracilis
lyratum, Nutt. in Journ. Acad. Philad. vii. (1834) 71 = humilis.
Wrightii, *A. Gray, Syn. Fl. N. Am.* i. ii. 411.—Texas.

APOGONIA, Fourn. Gram Mex. 63 ex Benth & Hook. f. Gen. iii. 1130 (1883) = **Rottboellia**, Linn. (Gramin.).
glabrata, Fourn. ex Hemsl. Biol. Centr Am. Bot. iii. 521 = Andropogon imberbis.
ramosa, Fourn. ex Hemsl. l. c. = R. aurita.

APOLANESIA, Reichb. Nom. 154 (1841) = **Apoplanesia**, Presl (Legum.).

APOLEPSIS, Hassk. Cat. Hort. Bogor. Alt. 150 (1844) = **Lepidagathis**, Willd. (Acanth.).
dispar, Hassk. l. c.= L. repanda.
repanda, Nees, in DC. Prod. xi. 261 = L. repanda.

APOLLINARIS, Linn. Syst. ed. I. (1735) = **Nicotiana**, Linn. † (Solan.).

APOLLONIAS, Nees, Laurin. Expos. 10 (1833). *LAURINEAE*, Benth. & Hook. f. iii. 152.
Arnottii, *Nees, Syst Laurin.* 670.—Ind. or.
barbusana, A. Br. in Verh. Beforerd. Gartenb. Preuss. xxi. (1852) 10 = canariensis.
canariensis, *Nees, Syst. Laurin.* 696.—Ind. or.
seylanica, Thw. Enum. Pl. Zeyl. 253 = Ritschmiedia oppositifolia.

APOMAEA, Neck. Elem. ii. 24 (1790) = **Ipomoea**, Linn. (Convolv.).

APOMOEA, Stend. Nom. ed. II. i. 114 (1840) = prace.

APONOA, Rafin. Sylva Tellur. 84 (1838) = **Columnea**, Linn. (Gesner.).
repens, Rafin. l. c. = C. scandens?

APONOGETON, Linn. f. Suppl. 93 (1781). *NAIADACEAE*, Benth. & Hook. f. iii. 1018.
AMOGETON, Neck. Elem. iii. 267 (1790).
APOGETON, Schrad. ex Steud. Nom. ed. II. i. 114 (1840).
HYDROGETON, Pers. Syn. i. 400 (1806).
LIMNOGETON, Edgew. in Lindl. Veg. Kingd. 210 (1847).
OUVIRANDRA, Thou. in Roem. Collect. 196 (1809); Gen. Nov. Madag. 2 (1810).
SPATHIUM, Edgew. in Journ. As. Soc. Beng. (1842) 145.
abyssinicum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 351.—Abyss.
angustifolium, *Ait. Hort. Kew.* i. 495.—Afr. austr.
Bernerianum, *Hook. f. in Benth. & Hook. f. Gen.* iii. 1014.—Madag.
crinifolium, *Lehm. ex Steud. Nom.* ed. II. i. 114.—Afr. austr.
crispum, *Thunb. Nov. Gen.* 78.—Zeylan.
desertorum, *Zeyh. ex Steud. Nom.* ed. II. i. 114.—Afr. austr.
distachyon, Edgew. in Hook. Lond. Journ. Bot. iii. (1844) 402 = distachyum.

APONOGETON :—
distachyum, *Thunb. Nov. Gen.* 74.—Afr. austr.
echinatum, *Roxb. Hort. Beng.* [87]; *et Fl. Ind. ed. Carey*, ii. 210.—Ind. or.
elongatum, *F. Muell. ex Benth. Fl. Austral.* vii. 188. —Austral
fenestrale, *Benth in Benth. & Hook. f. Gen.* iii. 1014.—Madag.
Holubii, *Oliver, in Hook. Ic. Pl.* t. 1470.—Afr. trop.
junceum, *Lehm. ex Steud. Nom.* ed. II. i. 114.—Afr. austr.
leptostachyum, *E. Mey. ex Baker, in Trans. Linn. Soc.* xxix. (1875) 158.—Afr. austr.
lineare, Vahl, Symb. Hot. iii. 51 = monostachyon.
microphyllum, *Roxb. Hort. Beng.* 26; *Fl. Ind. ed. Carey*, ii. 211.—Ind. or.
monostachyon, Edgew. in Hook. Lond. Journ. Bot. iii. (1844) 402 = seq
monostachyon, *Linn. f. Suppl.* 214.—Ind. or ; China.
natalense, *Oliver, in Hook. Ic Pl.* t. 1471.—Afr. austr.
Rehmanni, *Oliver, l. c.* 1471 B.—Afr. trop.
spathaceum, *E. Mey. in Drège, Zwei Pfl. Docum.* 165, *nomen.*—Afr. austr.
quadrangulare, *Baker, in Journ. Linn. Soc.* xviii. (1881) 279.—Madag.
ulvaceum, *Baker, l. c.*—Madag.
undulatum, Roxb. Hort. Beng. 26; Fl. Ind. ed. Carey, ii. 211 = crispum.
vallisnerioides, *Baker, in Trans. Linn. Soc.* xxix. (1875) 158.—Afr. trop.
zosteraefolium, *Schrad. ex Steud. Nom.* ed. II. i. 114 (an lapsu = Potamogeton zosteraefolius? Schum.).—Am. bor.

APOPHRAGMA, Griseb Gen. et Sp. Gent. 168 (1839) = **Curtia**, Cham. & Schlecht. (Gentian.).
tenuifolium, Griseb. l. c.— Guiana.

APOPHYLLUM, F. Muell. in Hook. Kew Journ ix. (1857) 306. *CAPPARIDEAE*, Benth. & Hook. f. i. 109.
anomalum, *F. Muell. l. c.* 307.—Austral.

APOPLANESIA, Presl, Symb. Bot. i. 68. t. 41 (1831). *LEGUMINOSAE*, Benth. & Hook. f. i. 492.
paniculata, *Presl, l. c.*—Mexic.

APOPLEUMON, Rafin. Fl. Tellur. iv. 72 (1836) = **Ipomoea**, Linn. (Convolv.).
bignonioides, Rafin l. c. = I. bignoniodes.

APORANTHUS, Bromf. Fl. Vect. 117 (1856) = **Trigonella**, Linn. (Legum.).
Trifolistrum, Bromf. l. c. = T. ornithopodioides

APORETIA, Walp. Ann. ii. 217 (1851-52) = seq.

APORETICA, Forst. Char. Gen. 131. t. 66 (1776) = **Schmidelia**, Linn. (Sapind.).
gemella, Blanco, Fl. Filip. ed. ii. 203 = S. Cobbe.
gemella, DC. Prod. i. 610 = S. cochinchinensis.
penicillata, Blanco, Fl. Filip. ed. I. 291 = S Cobbe.
pinnata, Forst. f. Prod. n. 392 = S. Cobbe.
pinnata, Hook. & Arn. Bot. Beech. Voy. 61 = Guioa lentiscifolia.
ternata, Forst. Char. 132 = S. Cobbe.

APOROCACTUS, Lem. Illustr. Hortic. vii. (1860) Misc. 67 = **Cereus**, Linn. (Cact.).
Baumannii, Lem. i. c. = C. Baumanni.
colubrinus, Lem. l. c. = C. colubrinus.
flagelliformis, Lem. l. c. = C. flagelliformis.
flagriformis, Lem. Cact. 68 = C. flagriformis.

APOROSA, Blume, Bijdr. 514 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 382.
LEIOCARPUS, Blume, Bijdr. 581 (1826).
LEPTIOSTACHYS, Wall. Cat. n. 6816 (1832).
SCEPA, Lindl. Introd. Nat. Syst. ed. II. 441 (1836).
TETRACTINOSTIGMA, Hassk. Hort. Bogor. Descr. 55 (1858).
acuminata, *Thw. Enum. Pl. Zeyl.* 288.—Ind. or.
affinis, Baill. Etud. Gén. Euph. 645 = Lindleyana.
arborea, *Muell. Arg. in DC. Prod.* xv. ii. 470.—Java.
arborescens, *Muell. Arg. l. c.*—Java.
aurita, *Baill. Etud. Gén. Euph.* 645.—Ins. Philipp.
aurita, Miq. Fl. Ind. Bat. i. ii. 481 = microcalyx.

APOROSA :—
australiana, *F. Muell. Census*, 20.—Austral.
Cumingiana, Baill. Etud. Gén. Euph. 645 = microcalyx.
dioica, Muell. Arg. in DC. Prod. xv. ii. 472 = Roxburghii.
filifolia, *Baill. Adansonia*, xi. (1873-76) 177.—Malaya; Cochinch.
frutescens, *Blume, Bijdr.* 514.—Java.
frutescens, Benth. Fl. Hongk. 317 = microcalyx.
fruticosa, *Muell. Arg. in DC. Prod.* xv. ii. 475.—Java.
fusiformis, *Thw. Enum. Pl. Zeyl.* 288.—Zeylan.
glabriflora, *Kurz, in Journ. Bot.* xiii. (1875) 330.— Ins. Nicobar.
lanceolata, Kurz, For. Fl. Burma, ii. 368 = Blanchoniana.
lanceolata, *Thw. Enum. Pl. Zeyl.* 288.—Zeylan.
latifolia, *Thw. l. c.*—Zeylan.
leptostachya, Benth. Fl. Hongk. 317 = microcalyx.
Lindleyana, *Baill. Etud. Gén. Euph.* 645 (Lyndleyana).—Ind. or.
lunatum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 239.—Malaya.
macrophylla, *Muell. Arg. in DC. Prod.* xv. ii. 470.— Burma.
microcalyx, *Hassk. in Bull. Soc. Bot. France*, vi. (1859) 714.—China; Austral.; Malaya.
microstachya, *Muell. Arg. in DC. Prod.* xv. ii. 474.— Burma.
Miquelliana, *Muell. Arg. l. c.*—Sumatra.
oblonga, *Muell. Arg. in Linnaea*, xxxii. (1863) 78.— Burma.
Planchoniana, *Baill. Etud. Gén. Euph.* 645.—Burma.
quadrilocularis, *Muell. Arg. in DC. Prod.* xv. ii. 476. —Sumatra.
Roxburghii, *Baill. Etud. Gén. Euph.* 645.—Ind. or.
sphaerocarpa, Muell. Arg. in Flora, xlvii. (1864) 519 = Lindleyana.
tetrapleura, *Hance, in Journ. Bot.* xiv. (1876) 260.— Cambodia.
Thwaitesii, Baill. Etud. Gén. Euph. 645 = fusiformis.
Tulasneana, Baill. l. c. = microstachya.
villosa, *Baill. l. c.*—Burma; Cochinch.
villosula, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 239.—Burma.

APORRHIZA, Radlk. in Sitzb. Math. Phys. Acad. Munch. viii. (1878) 338. *SAPINDACEAE*.
paniculata, *Radlk. l. c.* 339.—Afr. trop.

APORUM, Blume, Bijdr. 334. t. 89 (1825) = **Dendrobium**, Sw. (Orchid.).
acinaciforme, Griff. in Calc. Journ. Nat. Hist. v. (1845) 870 = D. acinaciforme.
anceps, Lindl. in Wall. Cat. n. 2020 = D. Nathanielis.
anceps, Lindl. Gen. et Sp. Orch. 71 = D. anceps.
concinnum, Lindl. ex Reichb. f. in Trans. Linn. Soc. xxx. (1874) 149 = D. atropurpureum.
cuspidatum, Wall. ex Lindl. in Bot. Reg. (1841) Misc. 2 = D. Nathanielis.
incrassatum, Blume, Bijdr. 334 = D. incrassatum.
incrassatum, Brongn. ex Lindl. in Journ. Linn. Soc. iii. (1859) 60 = Epidendrum aporoides?
individuum, Blume, Bijdr. 334. t. 89 = D. individuum.
individuum, Lindl. Gen. et Sp. Orch. 70 = D. Leonis.
Jenkinsii, Griff. in Calc. Journ. Nat. Hist. v. (1845) 867 = D. parciflorum.
Leonis, Lindl. in Bot. Reg. (1840) Misc. 20 = D. Leonis.
lobatum, Blume, Bijdr. 334 = D. lobatum.
micranthum, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 375. t. 16. f. 7 = D. Serra.
Roxburghii, Griff. l. c. v. (1845) 870 = D. Roxburghii.
Serra, Lindl. in Wall. Cat. n. 2021 = D. Serra.
sinuatum, Lindl. in Bot. Reg. (1841), Misc. 1 = D. sinuatum.
subterre, Griff. in Calc. Journ. Nat. Hist. v. (1845) 870 = D. acerosum.

APOSERIS, Neck. Elem. i. 57 (1790) = **Hyoseris**, Linn. (Compos.).
foetida, Cass. in Dict. Sc. Nat. xlviii. 427 = H. lucida.
foetida, Less. Syn. Comp. 138 = H. foetida.
lucida, Less. l. c. = H. lucida.

APOSTASIA, Blume, Bijdr. 432. t. 5 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 525.
MESODACTYLIS, Wall. Pl. As. Rar. i. 74, in obs. (1830).

APOSTASIA :—
NEUMAYERA, Reichb. f. in Flora, lv. (1872) 278.
NIEMEYERA, F. Muell. Fragm. vi. 96 (1867).
Brunonis, Griff. Notul. iii. 343 = nuda.
Lobbii, *Reichb. f. in Flora*, lv. (1872) 278.—Borneo.
nuda, *R. Br. in Wall. Pl. As. Rar.* i. 76. t. 85.—Ind. or.; Malaya.
odorata, *Blume, Bijdr.* 423.—Java.
stylidioides, *Reichb. f. in Flora*, lv. (1872) 278.—Austral.
Wallichii, *R. Br. in Wall. Pl. As. Rar.* i. 75. t. 84.—Ind. or.; Malaya.

APOTERIUM, Blume, Bijdr. 218 (1825) = **Calophyllum**, Linn. (Guttif.).
Sulatri, Blume, l. c. = C. spectabile.

APOXYANTHERA, Hochst. in Flora, xxvi. (1843) 78 = **Raphionacme**, Harv. (Asclep.).
pubescens, Hochst. l. c. = R. pubescens.

APOZIA, Willd. ex Steud. Nom. ed. II. i. 114 (1840) = **Micromeria**, Benth. (Labiat.).
chamædryoides, Willd. ex Steud. l. c. = M. Brownei.
herniarioides, Willd. ex Steud. l. c. = Herpestis herniarioides.
serpyllacea, Willd. ex Steud. l. c. = M. Brownei.

APPELLA, Adans. Fam. ii. 84 (1763) = **Laurus**, Linn.

APPENDICULA, Blume, Bijdr. 297. t. 40 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 588.
CONCHOCHILUS, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1843) 146.
METACHILUM, Lindl. Gen. et Sp. Orch. 74 (1830).
SCOLIOCHILUS, Reichb. f. Xenia Orch. ii. 118, in obs. (1873).
alba, *Blume, Bijdr.* 299.—Java.
ancepe, *Blume, l. c.*—Java.
angustifolia, *Blume, l. c.* 301.—Java.
bifaria, *Lindl. ex Benth. in Hook. Kew Journ.* vii. (1855) 35.—Ind. or.; China.
bracteosa, *Reichb. f. in Seem. Fl. Vit.* 299.—Ins. Fiji.
buxifolia, *Blume, Bijdr.* 300.—Java.
callosa, *Blume, l. c.* 303.—Burma; Malaya.
carnosa, *Blume, l. c.* 300.—Java.
Chalmersiana, *F. Muell. in Wing. South. Sc. Record*, N. S. (1. May 1885).—N. Guin.
congeneri, *Blume, Bijdr.* 303.—Java.
convuta, *Blume, l. c.* 302.—Java.
crinita, *Blume, l. c.* 298.—Java.
elegans, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
graminifolia, *Teijsm. & Binn. in Tijdschr. Neder. Ind.* xxiv. (1862) 322.—Java.
Hasseltii, *Blume, Bijdr.* 304.—Burma.
Hasseltii, Wight, Ic. t. 1748 = Agrostophyllum khasyanum.
Kuhlii, *Reichb. f. Xenia Orch.* ii. 116.—Java.
Lewisii, *Griff. in Calc. Journ. Nat. Hist.* iv. (1844) 378.—Malaya.
longifolia, *Blume, Bijdr.* 304.—Java.
membranacea, *Teijsm. & Binn. in Kruid. Arch.* iii. (1855) 399.—Java.
micrantha, Lindl. in Ann. Nat. Hist. xv. (1845) 886.—Ins. Philipp.
monocera, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
muricata, *Teijsm. & Binn. in Tijdschr. Neder. Ind.* xxiv. (1862) 322.—Java.
pasuliora, *Blume, Bijdr.* 300.—Java.
pendula, *Blume, l. c.* 298.—Java.
penicillata, *Blume, Rumphia*, iv. 46.—N. Guin.
peruligera, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
purpurascens, *Blume, Bijdr.* 302.—Java.
ramosa, *Blume, l. c.* 299.—Java.
reduplicata, Par. & Reichb. f. Otia Bot. Hamb. 45 = bifaria.
reflexa, *Blume, Bijdr.* 301.—Java.
rhodiola, *Reichb. f. Xenia Orchid.* ii. 118.—Java.
sarcoia, *Mig. Fl. Ind. Bat.* iii. 708.—Java.
stipulata, Griff. Notul. iii. 356 = callosa.
teres, Griff. l. c. 359 = Ceratostylis teres.
tenuicaulis, *Zoll. in Nat. en Geneesk. Arch. Neerl. Indie*, i. (1844) 408.—Java.
torta, *Blume, Bijdr.* 303.—Malaya.

APPENDICULA :—
tricornis, *Reichb. f. in Bonplandia*, v. (1857) 41.—Hab.? undulata, *Blume, Bijdr.* 301.—Java.
Vieillardi, *Reichb. f. in Linnaea*, xli. (1877) 76.—N. Caled.
viridiflora, *Teijsm. & Binn. in Tijdschr. Neder. Ind.* xxiv. (1862) 321.—Java.
xyriophora, *Reichb. f. in Seem. Fl. Vit.* 299.—Malaya.

APPENDICULARIA, DC. Prod. iii. 114 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 741.
thymifolia, *DC. l. c.*—Guiana.

APPUNIA, Hook. f. in Benth. & Hook. f. Gen. ii. 120 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 120.
BELLYNKIA, Muell. Arg. in Flora, xlviii. (1875) 465.
angulata, *Hasll. Pl. Pl.* vii. 415.—Bras.
tenuiflora, *Benth. & Hook. f. Gen.* ii. 120.—Guiana.

APRADUS, Adans. Fam. ii. 102 (1763) = **Arotopus**, Linn. (Umbell.).

APRELLA, Steud. Nom. ed. II. i. 152 (1840), err. typ = ASPRELLA, Schreb. = **Leersia**, Sw. (Gram.).

APREVALIA, Baill. in Bull. Soc. Linn. Par. i. (1884) 498. *LEGUMINOSAE*.
floribunda, *Baill. l. c.*—Madag.

APTANDRA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 201. *OLACINEAE*, Benth. & Hook. f. i. 345.
Benthamiana, Miers, l. c. Ser. III. iv. (1859) 360 = Iliciomicoides.
Iliciomicoides, *Spruce, ex Miers, l. c.*—Bras.
Spruceana, *Miers, l. c.* Ser. II. viii. (1851) 202.—Bras.
tubicina, *Benth. ex Miers, l. c.* Ser. III. iv. (1859) 360.—Bras.

APTERIA, Nutt. in Journ. Acad. Sc. Philad. vii. (1834) 54. *BURMANNIACEAE*, Benth. & Hook. f. iii. 458.
NEMITIS, Rafin. Fl. Tellur. iv. 38 (1836).
STEMOPTERA, Miers, in Proc. Linn. Soc. i. (1840) 62.
hymenanthera, Miq. in Verh. Nat. Wet. Haarl. vii. (1851) 216. t. 64 = setacea.
illacina, Miers, in Trans. Linn. Soc. xviii. (1841) 546. t. 27 = setacea.
orobanchioides, Hook. Ic. Pl. t. 254 = Dictyostega orobanchioides.
setacea, *Nutt. in Journ. Acad. Sc. Philad.* vii. (1834) 54. t. 2.—Am. bor.

APTERANTHES, Mik. in Nov. Act. Nat. Cur. xvii. (1835) 593. t. 41 = Bougueria, Wight & Arn. (Asclep.).
cylindrica, Decne. in Ann. Sc. Nat. Sér. V. xiii. (1870-1) 406.—Abyssin.
Gustoniana, Mik. in Nov. Act. Nat. Cur. xvii. (1835) 594. t. 42 = B. Gusconiana.
tessellata, Decne. in Ann. Sc. Nat. Sér. V. xiii. (1870-1) 406 = Echidnopsis cereiformis.

APTEROCARYON, Opiz, in Lotos, v. (1855) 258 = Betula, Linn. (Cupulif.).
Michauxii, Opiz, l. c. 259 = B. Michauxii.

APTERON, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 300. *RHAMNEAE*.
lanceolatum, *Kurz, l. c.* 301.—Tenasserim.

APTEUXIS, Griff. Notul. iv. 672 (1854) = **Pternandra**, Jack (Melast.).
trinervis, Griff. l. c. = P. coerulescens.

APTOSIMUM, Burch. Trav. i. 219 (1824). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 997.
CHELOSTIGMA, Hochst. in Flora, xxiv. (1841) 272.
OELENDORFIA, Lehm. Ind. Sem. Hort. Hamb. (1835).
adstringens, *Burch. Trav.* i. 226.—Afr. austr.
depressum, *Burch. l. c.* i. 220.—Afr. austr.
eriocephalum, *E. Mey. ex Benth. in Bot. Reg.* sub t. 1883.—Afr. austr.
lodivianum, *Burch. Trav.* i. 219.—Afr. austr.
procumbens, Burch. ex Steud. Nom. ed. II. i. 114 = depressum.

APTOSIMUM :—
pumilum, *Benth. in DC. Prod.* x. 545.—Afr. austr.
tragacanthoides, *E. Mey. ex Benth. in Bot. Reg.* sub t. 1883.—Afr. austr.
viscosum, *Benth. l. c.*—Afr. austr.

APTOTHECA, Miers, Apocyn. S. Am. 150 (1878). *APOCYNACEAE*.
corylifolia, *Miers. l. c.*—Cuba.

APULEIA, Gaertn. Fruct. ii. 439. t. 171 (1791) = **Berkheya**, Ehrh. (Compos.).
asteroides, Gaertn. l. c. 440 = B. fruticosa.
atractyloides, Less. Syn. Comp. 64 (= *Stobæa atractyloides*).
fruticosa, Bartl. Sem. Goett. (1835).—Cf. Linnæa, xii. (1838) Lit. 80 = B. fruticosa.
fruticosa, Gaertn. Fruct. ii. 440 = B. grandiflora.
heterophylla, Less. Syn. Comp. 65 (= *Stobæa heterophylla*).
rigida, Gaertn. Fruct. ii. 439. t. 171 (= *Stobæa Gaertneri*).
Zeyheri, Less. Syn. Comp. 64 = B. Zeyheri.

APULEIA, Mart. Herb. Fl. Bras. 128 (1887). *LEGUMINOSAE*, Benth. & Hook. f. i. 574.
ZENKERIA, Arn. in Mag. Zool. & Bot. ii. (1838) 548.
molaris, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 177.—Bras.
praecox, *Mart. Herb. Fl. Bras.* 128.—Bras.

AQUARTIA, Jacq. Enum. Pl. Carib. i. 12 (1760) = Solanum, Linn.
aculeata, Jacq. l. c.—S. Aquartia.
microphylla, Lam. Illustr. i. 504. t. 82. f. 2 = S. microphyllum.
microphylla, Willd. Sp. Pl. i. 932.—Ins. S. Doming.
solanacea, Willd. ex Steud. Nom. ed. II. i. 114.—N. Granat.
tomentosa, Lam. Illustr. p. 1603. t. 82. f. 1 = S. Aquartia.

AQUIFOLIUM, Tourn. ex Hall. Enum. Stirp. Helv. ii. 407 (1742) = **Ilex**, Linn.
crocum, Rafin. Sylva Tellur. 44 = I. Aquifolium?
ferox, Mill. ex Rafin. l. c. = I. Aquifolium.
heterophyllum, Rafin. l. c. = I. Aquifolium.
Ilex, Scop. Fl. Carn. ed. II. i. 128 = I. Aquifolium.
japonicum, Rafin. Sylva Tellur. 44 = Berberis japonica.
lanceolatum, Rafin. l. c.—Hab.?
planifolium, Rafin. l. c.—Hab.?
spinosum, Lam. Fl. Fr. iii. 252 = I. Aquifolium.
undulatum, Rafin. Sylva Tellur. 44 = I. Aquifolium.
vulgare, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = I. Aquifolium.

AQUILARIA, Lam. Encyc. ii. 610 (1788). *THYMELAEACEAE*, Benth. & Hook. f. iii. 200.
AGALLOCHUM, Lam. Encyc. i. 47 (1783).
OPHIOSPERMUM, Reichb. Consp. 82 (1828).
OPHISPERMUM, Lour. Fl. Cochinch. 350 (1790).
Agallocha, *Roxb. Hort. Beng.* 38; *Fl. Ind.* ii. 422.—Ind. or.; Burma.
bancana, Miq. Fl. Ind. Bat. Suppl. 358 = Gonostylus Miquelianus.
chinensis, Sprang. Syst. ii. 356 = A. Ophispermum.
grandiflora, *Benth. Fl. Hongk.* 297.—China.
macrophylla, Miq. Fl. Ind. Bat. Suppl. 356 = Gonostylus Miquelianus.
malaccensis, Benth. in Hook. Kew Journ. v. (1853) 195 = grandiflora.
malaccensis, *Lam. Encyc.* i. 49.—Malaya.
microcarpa, *Baill. Adansonia*, xi. (1873-76) 304.—Borneo.
Ophispermum, *Poir. in Dict. Sc. Nat.* xviii. 161.—China.
ovata, Cav. Diss. vii. 377 = malaccensis.
pentandra, *Blanco, Fl. Filip.* ed. I. 378.—Ins. Philipp. secundaria, *DC. Prod.* ii. 56.—Ins. Molucc.

AQUILEGIA [Tourn.] Linn. Syst. ed. I. (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 8.
adsvea, *Regel, in Ind. Sem. Hort. Petrop.* (1856) 32.—Hab.?
aggericola, Jord. Diagn. i. 37 = vulgaris.
alpina, Dalla Torre, Ic. t. 49 = pyrenaica.
alpina, Cesati, It. i. 229 = glandulosa.
alpina, Lam. Encyc. i. 150 = pyrenaica.

AQUILEGIA :—
alpina, *Linn. Sp. Pl.* 533.—Europ.; Sibir.
Amaltea, *Heldr. in Boiss. Diagn.* Ser. II. i. 11.—Thessal.
anemonoides, Willd. in Ges. Nat. Fr. Berl. Mag. v. (1811) 401 t. 9. f. 6 = Isopyrum grandiflorum.
aragonensis, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 966.—Hispan.
arbascensis, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse*, ii. (1874) 449; (*Massive d'Arbas*, 58).—Europ.
arctica, Loud. Hort. Brit. 610 = formosa.
atrata, Koch, in Flora, xiii. (1830) 1. 118 = vulgaris.
atropurpurea, Miq. in Ann. Mus. Lugd. Bat. iii. 8 = Buergeriana.
atropurpurea, Willd. Enum. Hort. Berol. 577 = viridiflora.
aurea, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 174 = chrysantha.
Bauhini, Schott, in Verh. Bot. Zool. Ver. Wien, iii. (1853) 128 = Einseleana.
Bernardi, Gren. & Godr. Fl. Fr. i. 45 = vulgaris.
Bertoloni, *Schott, in Verh. Bot. Zool. Ver. Wien*, iii. (1853) 127.—Europ.
bicolor, Ehrh. Beitr. viii. 146 = sibirica.
blanda x, Lem. Illustr. Horticl. (1857) t. 146 = vulgaris.
brachyceras, Turcz. ex Fisch. & Mey. in Maund, Bot. Gard. n. 755 = leptoceras.
Brunni, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10, nomen.—Europ.
brevistyla, *Hook. Fl. Bor. Am.* i. 24.—Am. bor.
Buergeriana, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1846) 188.—Japon.
caerulea, *James, in Long, Exped. Rocky Mount.* ii. 345.—Am. bor. occ.
californica, *A. Gray, in Proc. Am. Acad.* vii. (1868) 328.—Am. bor. occ.
californica, Hartw. in Gard. Chron. (1854) 886 = formosa.
campylocentra, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10.—Europ.
canadensis, Hook. Fl. Bor. Am. i. 24 = formosa.
canadensis, *Linn. Sp. Pl.* 533.—Am. bor.
canadensis, Pall. Reise, iii. 280 = viridiflora.
caucasica, Ledeb. ex Rupr. Fl. Cauc. 32 = olympica.
chrysantha, *A. Gray, in Proc. Am. Acad.* viii. (1873) 621.—N. Mexic.
coerulea, Auct. = caerulea.
collina, Jord. Diag. i. 84 = vulgaris.
concolor, Fisch. ex Steud. Nom. ed. II. i. 115 = vulgaris.
confusa, Rota, Prosp. Fl. Bergamo, 99 = Einseleana.
corniculata, Vill. Cat. Hort. Strasb. 250 = vulgaris.
cornuta, Gilib. Fl. Lituan. ii. 286 = vulgaris.
corsica, Soleir. ex Nym. Consp. 18 = vulgaris.
cyclophylla, Jeaub. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, iii. (1875–76) 555; (Massif Laurenti, 189) = pyrenaica.
dakurica, Patr. in Deless. Ic. Sel. t. 49 = viridiflora.
dichroa, *Freyn, in Flora*, xliii. (1880) 96.—Europ.
dioica, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10, nomen.—Europ.
discolor, Steud. Nom. ed. II. i. 115 = glandulosa.
discolor, *Lerier & Lorsseke, in Yourn. Bot.* xvii. (1879) 197.—Hispan.
dumaticola, Jord. Diag. i. 86 = vulgaris.
Ebneri, Zimm. ex Nym. Consp. 18 = vulgaris.
ecalcarata, Hort. ex Steud. Nom. ed. II. i. 115 = vulgaris.
Einseleana, *Fr. Schultz, Arch. Fl. Fr. et Allem.* 135.—Europ.
elata, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 41 = vulgaris.
elegans, Pepe, ex Steud. Nom. ed. II. i. 115 = vulgaris.
elegans, Saliab. Prod. 374 = canadensis.
eximia, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10, nomen.—Europ.
exilens, Van Houtte, ex Planch. in Fl. des Serres, Sér. II. ii. (1857) 13. t. 1168 = formosa.
flabellata, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1846) 188.—Japon.
flava, Lam. Encyc. i. 149 = viridiflora.
flavescens, *S. Wats. in Bot. King, Exp.* 10.—Calif.
flavifloru, Tenore, in Am. Natural. i. (1868) 268 = canadensis.
formosa, *Fisch. in DC. Prod.* i. 30.—Kamtsch.
formosa, *Hook. f. Bot. Mag.* t. 6969 = Hookeri.

AQUILEGIA :—
fragrans, *Benth. in Maund, Botanist*, iv. t. 181.—Reg. Himal.
Guerneri, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10, nomen.—Europ.
Garnieriana, Sweet, Brit. Fl. Gard. Ser. II. v. t. 103 = sibirica.
Gebleri, *Bess. ex Turcz. in Flora*, xvii. (1834) i. *Beibl.* 5.—Europ.
glandulosa, *Fisch. Hort. Gorenk.* ed. II. (1812) 48; *et ex Link, Enum. Hort. Berol.* ii. 84.—Sibir. altaic.
glandulosa, Miq. in Ann. Mus. Lugd. Bat. iii. 8 = flabellata.
glauca, *Lindl. Bot. Reg.* xxvi. (1840) t. 46.—Reg. Himal.
glaucescens, Baker, in Gard. Chron. (1878) ii. 208 = vulgaris.
glaucophylla, Steud. in Flora, xxxix. (1856) 407 = vulgaris.
grandiflora, Patr. ex DC. Syst. i. 337 = alpina.
grandiflora, Schangin, in Pall. N. Nord. Beitr. vi. 49 = glandulosa.
grata, *Maly, ex Borbas, in Magyar Tudom. Akad.* xii n. vi. (1882) 8, nomen.—Europ.
Haenkeana, Koch, Syn. Fl. Germ. 28 = vulgaris.
Haynaldi, *Borbas, in Magyar Tudom. Akad* xii. n. vi. (1882) 9.—Europ.
hirsuta, *Tineo.—Cf. Bull. Soc. Bot. Fr.* xxix. (1882), *Bibl.* 47.—Gallia merid.
hispanica, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10.—Hispan.
Hookeri, *Borbas, in Termész. Füzetek*, viii. (1884) 311. —Asia bor.
Huteri, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 9.—Europ.
hybrida, Sims, in Bot. Mag. t. 1221 = vulgaris.
inversa, Mill. Gard. Dict. ed. VIII. n. 3 = vulgaris.
Jousii, *Perry, in Am. Natural* viii. (1874) 211.—Am. bor.
jucunda, Fisch. & Lallem. Ind. Sem. Hort. Petrop. vi. (1840) 28 = glandulosa.
jucunda, Fisch. & Mey. ex Ledeb. Fl. Ross. i. 736 = vulgaris.
kansuensis, Jacquem. ex Camb. in Jacquem. Voy. Bot. 7. t. 5 = Moorcroftiana.
Karelini, Baker, in Gard. Chron. (1878) ii. 76 = vulgaris.
Karelinna, *C. A. Mey. ex Trautv. in Act. Hort. Petrop.* vii. (1883) 68.—Hab. ?
Kitaibelii, *Schott, in Verh. Bot. Zool. Ver. Wien*, iii. (1853) 129.—Armenia.
lactiflora, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 374.—Sibir.
leptoceras, Fisch. & Mey. *Ind. Sem. Hort. Petrop.* iv. 28.—Sibir.
leptoceras, Nntt. in Journ. Ac. Philad. vii. (1834) 9 = caerulea.
longissepala, *Zimm. ex Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 9, nomen.—Europ.
longisima, *A. Gray, ex S. Wats. in. Proc. Am. Acad.* xvii. (1881–82) 317.—Am. bor.
lutea, *Lam. Encyc.* i. 149.—Hab. ?
lutescens, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10, nomen.—Europ.
macrantha, Hook. & Arn. Bot. Beech. Voy. 317. t. 72 = caerulea.
macrocentra, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10, nomen.—Europ.
Magnolii, Loret, in Loret & Barrand. Fl. Montp. ii. 796 = Kitaibelii.
mexicana, Hook. Bot. Mag. t. 3919 = Skinneri.
minuta, Steph. ex Ledeb. Fl. Ross. i. 53 = Isopyrum grandiflorum.
mollis, *Yeanb. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse*, iii (1875–76) 555.—Gallia.
montana, Sternb. in Regensb. Denkschr. ii. (1818) 60 = alpina.
Moorcroftiana, *Wall. Cat.* n. 4712.—Reg. Himal.
Moorcroftiana, Steud. Nom. ed. II. i. 115, sphalm. = praec.
nemorosa, *Jord. Diag.* i. 83.—Gallia.
nevadensis, *Boiss. & Reut. Cat. Hort. Genev.* (1854) 4; *et in Ann. Sc. Nat.* Sér. IV. ii. (1854) 380.—Hispan.
nigricans, Baumg. Enum. Stirp. Transs. ii. 104 = vulgaris.
nivalis, Falc. ex Baker, in Gard. Chron. (1878) ii. 76 = glauca.

AQUILEGIA :—
nivea, Baumg. ex Baker, in Gard. Chron. (1878) ii. 76 = vulgaris.
olympica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 360.—Orient.
orthantha, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10, nomen.—Europ.
Ottonis, *Orph. ex Boiss. Diagn.* Ser. II. i. 11.—Graecia.
oxysepala, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 10).—As. or.
parciflora, Schur, Enum Pl. Transs. 28 = vulgaris.
parviflora, *Ledeb. in Mem. Acad. Pétersb.* v. (1814) 544.—Sibir.
platysepala, Reichb. Ic. Fl. Germ. t. 4730 = vulgaris.
plena, Hort. ex Steud. Nom. ed. II. i. 115 = vulgaris.
praecox, Jord. Diag. i. 85 = vulgaris.
pubiflora, *Wall. Cat.* n. 4714.—Ind or.
pycnotricha, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10.—Europ.
pyrenaica, *DC. Fl. Fr. iv.* 912.—Pyren.; Appenin.
pyrenaica, Koch, Syn. Fl. Germ. ed. II. 24 = Einseleana.
pyrenaica, Reichb. Ic. Fl. Germ. t. 4732 = Bertoloni.
Reuteri, Boiss. Diagn. Ser. II. i. 10 = Bertoloni.
Reuteriana, Reichb. f. ex Nym. Consp. 18 = alpina.
ruscinonensis, *Timb. & Yeanb. in Bull. Soc. Sc. Nat. Toulouse*, iii. (1875–76) 554.—Gallia.
Schinneri, Polt. in Rev. Hortic. Sér. II. iv. (1845–46) 185 = Skinneri.
Schottii, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10.—Europ.
sibirica, Don, ex Steud. Nom. ed. II. i. 115 = vulgaris.
sibirica, *Lam. Encyc.* i. 150.—Sibir.
sibirica, Schur, ex Nym. Consp. 18 = glandulosa.
silvestris, Neck. Delic Gallo-belg i. 234 = vulgaris.
sincana, *Hayek. Nom. ill.* 41.—China.
Skinneri, *Hook. Bot. Mag.* t. 3919.—Mexic.
speciosa, DC. Syst. i. 338 = sibirica.
speciosa, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 185 = arbascensis.
spectabilis, Lem. in Illustr. Hortic. xi (1864) t. 403 = sibirica.
stellata, Hort. ex Steud Nom. ed. I. 61; ed. II. i. 115 = vulgaris.
stenopetala, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 8, nomen.—Europ.
Sternbergii, Reichb. Fl. Germ Excurs 749 = alpina.
subalpina, Bor. Fl. Centr. Fr. ed. III. ii. 24 = vulgaris.
subrapposa, Borbas, in Magyar Tudom. Akad. xii. n. vi. (1882) 18 = vulgaris.
sulphurea, *Zimm. ex Borbas, l. c.* 9.—Europ.
sylvestris, vide *silvestris*.
sylvestris, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 4 = vulgaris.
Szaboi, *Borbas, in Magyar Tudom. Akad.* xii. n. vi. (1882) 10, nomen.—Europ
thalictrifolia, Schott & Kotschy, in Verh. Bot. Zool. Ver. Wien, iii. (1853) 130 = Einseleana.
thalictrifolia, Schlecht. in Linnaea, vi. (1831) 581 = parv flora.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1. 52) 94; iv. (1853) 4 = vulgaris.
truncata, Fisch. & Mey. Ind. Sem. Hort. Petrop. (1843) Suppl. 8 = formosa.
variegata, Moench, Meth. 311 = canadensis.
versicolor, Saliab. Prod. 374 = vulgaris.
viridiflora, *Pall. in Act. Petrop.* (1779) 260. t. 11.—Sibir.
viscosa, Gouan, Fl. Monsp. 267 = Kitaibelii.
viscosa, Reichb. Fl. Germ. Excurs. 748 = Einseleana.
viscosa, Suffr. ex Reichb. l. c. 740 = pyrenaica.
volubilis, *Maack, ex Trautv. in Act. Hort. Petrop.* viii. (1883) 54.—Manchuria.
vulgaris, *Linn. Sp. Pl.* 533.—Europ.; Orient.
vulgaris, Richards. App. Frankl. Journ. ed. II. 21 = brevistyla.
vulgaris, Thunb. Fl. Jap. 232 = flabellata.
Wittmaniana, Stev. ex Fisch. & Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. xi. Suppl. 15 = olympica.

AQUILICIA, Linn. Mant. ii. 211 (1771) = **Loca**, Linn. (Ampel.).
Ottilis, Gaertn. Fruct. i. 275 = L. sambucina.
sambucina, Linn. Mant. 211 = L. sambucina.

ARABIDAEA.—Cf. Rev. Hortic. (1868) 154 = **Arrabidaea**, DC. (Bignon.).

ARABIDIUM, Spach, Hist. Vég. Phan. vi. 436 (1838) = **Arabis**, Linn. (Crucif.).
alpestre, Spach, l. c. 436 = Arabis alpina.
alpinum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi (1868) 382 = Arabis alpina.
saxatilolum, Fourr. l. c. = Arabis saxatilis.

ARABIDOPSIS, Schur, Enum. Pl. Transs. 55 (1866) = **Stegnobrium**, Linn. (Crucif.).
pinnatifida, Rupr. Fl. Caucas. 86 = Braya pinnatifida. *Thaliana*, Schur, Enum. Pl. Transs. 55 = S. Thalianum.

ARABIS, Adans. Fam. ii. 422 (1763) = **Iberis**, Dill. (Crucif.).

ARABIS, Linn. Gen. ed. I. 198 (1737). *CRUCIFERÆ*, Benth. & Hook. f. i. 69.
ARABICARPON, Andrz. ex Reichb. in Mocsal. Handb. ii. 1150 (1838).
ARABIDIUM, Spach, Hist. Vég. Phan. vi. 436 (1838).
ARABISA, Reichb. Handb. 260 (1841).
CAULOPSIS, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvi. (1868) 382.
LOMARPORA, DC. ex Steud. Nom. ed. II. ii. 65 (1841).
PILOSELLA, Kostel. Hort. Prag. 104 (1844).
PSILARABIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 382.
STEVENIA, Adams & Fisch. in Mém. Soc. Nat. Mosc. v. 84 (1890).
TURRITIS, Tourn. ex Linn. Syst. ed. I. (1737).
accedens, Jord. Diag. i. 115 = hirsuta.
aizoides, *Bonato, Cat. Pl. Hort. Patav. (ex. Steud. Nom. ed. II. i. 119).*—Hab.?
alaschanica, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 491.—Mongolia.
albida, *Stev. in Fisch. Cat. Hort. Gorenk.* 51.—Reg. Meditert.; Orient.
Alliaria, Benth. Syst. Vert. Erf. 195 = Sisymbrium Alliaria.
Allionii, *DC. Fl. Fr.* iv. 676.—Pedemont.
alpestris, *Schleich. ex Reichb. Ic. Fl. Germ.* ii. 18. n. 4338 b.—Helvetia.
alpina, Georgi, Beschr. Russ. Reich. iii. iv. 1157 = albida.
alpina, Krock. ex Steud. Nom. ed. II. i. 115 = sagittata.
alpina, *Linn. Sp. Pl.* 664.—Europ. ; Am. bor.
alpina, Lucé, Fl. Odl. 228 = Gerardi.
amara, Banks, ex DC. Syst. ii. 246 = Cardamine rhomboidea.
amœna, DC. Syst. ii. 231 = petræa.
amplexicaulis, *Edgew. in Trans. Linn. Soc.* xx. (1851) 81.—Reg. Himal.
anulicola, R. E. & K. Nov. Gen. et Sp. v. 81 = Sisymbrium hispidulum.
androsaces, *Fenzl, Pugill. Pl. Nov. Syr.* 18.—As. Min.
angustifolia, Lam. Encyc. i. 220 = Hutchinsia alpina.
apennina, *Tausch, in Syll. Ratisb.* ii. (1829) 244.—Italia.
arcuata, Dulac, Fl. Hautes-Pyr. 200 = Sisymbrium Thalianum.
arcuata, *A. Gray, in Proc. Am. Acad.* vi. 187.—Calif.
arcuata, Shuttlw. ex Godet, Fl. Jura, i. (1852) 88 = alpestris.
arenaria, *Schrank, Fl. Monac.* iii. 256.—Europ.
arenosa, Cham. & Schlecht. in Linnæa, i. (1826) 17 = kamtschatica.
arenosa, *Scop. Fl. Carn.* ed. II. ii. 82.—Europ.
arvensis, Benth. Syst. Vert. Erf. 195 = Conringia austriaca.
arvensis, *Edgew. in Trans. Linn. Soc.* xx. (1851) 82. —Ind. bor. occ.
aspera, All. Ped. Auct. 18. t. 9. f. 2 = auriculata.
attenuata, Royle, ex Hook. f. in Journ. Linn. Soc. vi. (1862) 148 = tibetica.
aulaticioides, *Boiss. Fl. Orient.* i. 176.—Cilicia.
Aucheri, *Boiss. in Ann. Sc. Nat. Sér.* II. xvii. (1842) 63.—As. Min. ; Syria.
auriculata, *Lam. Encyc.* i. 219.—Europ. ; Orient.
Balansæ, *Boiss. & Reut. Diagn.* Sér. II. v. 17.—Algeria.
Barbarea, Benth. Syst. Vert. Erf. 195 = Barbarea vulgaris.
bellidifolia, Crantz, Stirp. Austr. ed. I. fasc. i. 49; ed. II. 44 = pumila.
bellidifolia, Lucé, Fl. Carn. ed. II. ii. 81 = Cardamine alpina.

ARABIS:—
bellidioides, Lam. Encyc. i. 220 = Cardamine alpina.
bijuga, *Wall. in Journ. Linn. Soc.* xviii. (1881) 378. —Reg. Himal.
Billardieri, DC. Syst. ii. 218 = albida.
blepharophylla, *Hook. & Arn. Bot. Beech. Voy.* 321.—Calif.
Boissieri, Griseb. Spicil. Fl. Rumel. i. 248 = drabæformis.
borealis, Andrz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 33 = hirsuta.
Borysi, Boiss. Voy. Bot. Espagne, ii. 26. t. 5 = Braya pinnatifida.
brachycarpa, *Rupr. Fl. Cauc.* 78, *in textu.*—Reg. Caucas.
brachypoda, Boiss. Elench. 8 = parvula.
bracteata, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 291.—Bolivia.
brassicæformis, *Wallr. Sched. Crit.* 359.—Europ.
Brunfelii, *Lange, Consp. Fl. Groenl.* i. 50.—Groenl.
brevifolia, DC. Syst. ii. 218 = albida.
Breweri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 123. —Calif.
bryoides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 55.—Græcia.
bulbosa, Muhl. Cat. 68 ; Spreng. Fl. Hal. Mant. 46 = Cardamine rhomboidea.
bursifolia, Lam. Fl. Fr. ii. 511 = Sisymbrium bursifolium.
cadmea, *Boiss. Diagn.* Ser. I. viii. 21.—Caria.
caerulea, *Haenke, in Jacq. Collect.* ii. 56.—Europ.
caespitosa, *Schleich. Cat. Pl. Helv.* ed. IV. 8.—Helvet.
camtschatica, Willd. ex DC. Syst. ii. 242 = Stelleri.
canadensis, *Linn. Sp. Pl.* 665.—Am. bor.
canadensis, Mill. Gard. Dict. ed. VIII. n. 6 = Diplotaxis muralis.
cancasica, Brocchi, in Bibliot. Ital. xvix. 90 = alpina.
canescens, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 83.—Am. bor.
cantabrica, *Lresche & Lovier, in Journ. Bot.* xvii. (1879) 197.—Hispan.
capensis, Linn. Syst. ed. X. 1135 ; Burm. f. Prod. Fl Cap. 18 = Heliophila pusilla.
Cardamines, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 292.—Austral.
carduchorum, *Boiss. Fl. Orient.* i. 178.—Armenia.
carpetana, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 822.—Hispan.
caucasica, Willd. Enum. Hort. Berol. Suppl. 45 = albida.
cebennasia, *DC. Syst.* ii. 234.—Gallia.
cenisia, Reut. Cat. Jard. Bot. Genèv. (1853) 4 = alpestris.
chrysantha, Rottl. ex Wight, Illustr. i. sub t. 19 = Lepidium sativum.
ciliaris, Willd. Enum. Hort. Berol. 684 = pumila.
ciliata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 107.—Europ.
ciliata, Auct. ex Nym. Consp. i. 33 = alpestris.
ciliata, Tausch, Ind. Pl. Sudet. (sec. Steud. Nom. ed. II. i. 117) = hirsuta.
ciliicea, Boiss. & Bal. Diagn. Ser. II. v. 17 = ionocalyx.
Cinsiana, Schrank, Fl. Monac. ii. 125 = alpina.
coerulea, vide cærulea.
colchagensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 294.—Chili.
collina, *Tenore, Prod. Fl. Nap.* 39.—Italia.
collisparsa, Jord. Diagn. i. 116 = hirsuta.
colorata, *Tausch, in Flora*, xix. (1836) ii. 405.—Europ.
Commersonii, *DC. Syst.* ii. 238.—Reg. Argent.
conferta, Willd. ex Reichb. Fl. Germ. Excurs. 680 = hirsuta.
coringioides, *Ball, in Journ. Bot.* xi. (1873) 297.—Hispan.
constricta, Griseb. Spicil. Fl. Rumel. i. 249 = hirsuta.
contracta, *Celak. Prod. Fl. Bohem.* 155.—Bohem.
contracta, Loesn. Fl. Peterg. iii. 925 = hirsuta.
cordata, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 51 = laxa.
Cossæ, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 890.—Hispan.
Crantziana, Ehrh. Beitr. v. 177 = petræa.
cramifolia, *Ledeb. in Baier. Bot. Ges. Denkschr.* iii. (1841) 107.—Rossia.
crenocarpa, Boiss. & Bal. Diagn. Ser. II. v. 16 = laxa.
crepidifolpoda, Griseb. ex Pant. in Oestr. Bot. Zeitschr. xxiii. (1873) 288 = pseudoturritis.

ARABIS:—
cretica, Boiss. & Heldr. Diagn. Ser. I. viii. 20 = muralis.
crispata, Willd. Enum. Hort. Berol. 684 = alpina.
croatica, Schott, ex Nyman & Kotschy, Analect. Bot. 44 = neglecta.
cuneifolia, Huchst. ex A. Rich. Tent. Fl. Abyss. i. 17 = alpina.
curtisiliqua, DC. Syst. ii. 228 = hirsuta.
Cusickii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 363.—Am. bor.
dacsca, Heuff ex Grisob. & Schenk, in Wiegm. Archiv, xvii. (1851) i. 307 = Hallen.
dasycarpa, Andrz. ex DC. Syst. ii. 244 = auriculata.
declinata, Schrad. Ind. Sem. Hort. Goett. (1831) ; (ex Linnæa, viii. (1833) Litt. 22) = Holboellii.
decumbens, *Ball, in Journ. Bot.* xi. (1873) 297.—Marocco.
deflexa, *Boiss. Fl. Orient.* i. 175.—Lycia.
dentata, [Clairv]. Man. Herb. 228 = Braya pinnatifida
dentata, *Torr. & Gray, Fl. N. Am.* i. 80.—Am. bor.
docsca, Puss, in Verh. Siebenb. Ver. Naturw. xiii. (1862) 150, sphalm. = dacica.
Douglasii, Torr. in Torr. & Gray, Fl. N Am. i. 83, in syn. (cum cit. falsa) = Cardamine rotundifolia.
drabæformis, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 55.—Bithyn.
drabæformis, Griseb. Spicil. Fl. Rumel. i. 248 = bryoides.
Dregeana, *Harv. & Sond. Fl. Cap.* i. 22.—Afr. austr.
Drummondii, *A. Gray, in Proc. Am. Acad.* vi. 187.—Am. bor.
elongata, Salisb. Prod. 372 = A. Turrita.
ericarpa, Schur, Enum. Pl. Transs. 46 = A. Turrita.
eriopoda, *Turcz. in Bull. Soc. Nat Mosc.* (1849) 63. —Sibir. or.
erubescens, *Ball, in Journ. Bot.* xi. (1873) 297.—Marocco.
erysimoides, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv (1842) 145.—Soongaria.
Eschscholtziana, Andrz. ex Ledeb. Fl. Ross. i. 118 = hirsuta.
falcata, Michx. Fl. Am. Bor. i. 81 = canadensis.
falcata, *A. Rich. Tent. Fl. Abyss.* i. 17.—Abyss.
falklandica, Benth & Hook. f. Gen. i. 69 = macloviana.
farinacea, *Rupr Fl. Caucas.* 71.—Caucasus.
fastigiata, Hook. f. Fl. N. Zel. ii. 324 = Cardamine fastigiata
flagellosa, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 72.—Japon.
flavicoma, *Bunge, in Mém. Acad. Pétersb.* Sér. VI. vii. (1859) 582.—Armenia.
flexuosa, Hort. Berol. ex Reichb. Fl. Germ. Excurs. 679 = serpyllifolia.
floribunda, Schur, Enum. Pl. Transs. 44 = petræa.
foliosa, *Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1862) 160.—Reg. Himal.
fruticulosa, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 19.—Sibir. alt.
furcata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 362.—Am. bor.
Gerardi, *Bess. in Koch, Syn. Fl. Germ.* ed. I. 38.—Europ. ; Reg. Casc.
gigantea, Hook. Ic. Pl. t. 259 = Cardamine styloса.
glabra, Bernh. Syst. Verz. Erf. 195 = perfoliata.
glandulosa, *Kar & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 146.—Tibet. ; Soongar.
glareosa, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 106 ; ix. (1858) 59 = arenosa.
glastifolia, *Reichb. Ic. Fl. Germ.* xii. 13.—Europ.
Gmelinii, Willd. ex Ledeb. Fl. Ross. i. 131 = Parrya macrocarpa.
gracilescens, Jord. Diag. i. 119 = hirsuta.
gracilis, Willd. ex Ledeb. Fl. Ross. i. 186 = Sisymbrium multisiliquosum.
gracilis, Royle, ex Hook. f. & Thoms in Journ. Linn. Soc. v. (1861) 159 = Sisymbrium Wallichii.
grandiflora, Linn. Amœn. Acad. ii. 279 = Parrya macrocarpa.
grandiflora, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 148 = alpina.
Halleri, *Linn. Sp. Pl.* ed. II. 929.—Europ.
Halleri, Lucé, Fl. Odl. 228 = sagittata.
hastulata, Bert. ex DC. Prod. i. 150 = Cardamine resedifolia.
Heldreichii, DC. Syst. ii. 237 = Farsetia longisiliquata.
hesperidoides, A. Gray, Mem. Bot. U. St. ed. V. 66 = Thelypodium pinnatifidum.

ARABIS:—
heteromalla, Schrad. Ind. Sem. Hort. Goett. ex Linnaea, viii. (1833) Litt. 32 = Holboellii.
heterophylla, Forst. ex DC. Syst. ii. 265 = Cardamine hirsuta.
heterophylla, Nutt. ex Torr. & Gray, Fl. N. Am. i. 81 = laevigata.
heterophylla, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1858) 59.—Austria.
himalaica, *Edgew. in Trans. Linn. Soc.* xx. (1851) 31. —Reg. Himal.
hirsuta, DC. Fl. Fr. iv. 675 = ciliata.
hirsuta, *Scop. Fl. Carn.* ed. II. ii. 30.—Europ.; Orient; Am. bor.
hirsuta, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 142 = alpina.
hirta, Lam. Encyc. i. 220 = stricta.
hirtella, Jord. Diag. i. 120 = hirsuta.
hispida, Soland. in Ait. Hort. Kew. ed. I. ii. 400 = stricta.
hispida, Murr. Syst. Veg. ed. XIII. 501 (cf. Linn. f. Suppl. 298) - petraea.
hispida, Moench, Meth. 258 = pendula. Holboellii, *Hornem. Fl. Dan.* t. 1879.—Am. bor.
Holboelliana, Hort. Hafn. ex Trevir. in Flora, xvi. (1833) 194 = praec.
Hookeri, *Lange, Consp. Fl. Groenl.* i. 50.—Groenl.
Hornungiana, *Schur, Enum. Pl. Transs* 43.— Transylv.
Huetii, *Traute. in Act. Hort. Petrop.* ii. (1878) 496.— Reg. Caucas.
humilis, Schleich. ex DC. Syst. ii 224 = muralis.
hybrida, Rent. Suppl. Cat. Fl. Genèv. 8 = alpestris.
idaeanensis, Jord. Diag. i. 117 = hirsuta.
incana, Moench, Meth. 257 = alpina.
incana, *Roth, Catalect. Bot.* i. 79.—Europ.
incana, Willd. Enum. ex Steud. Nom. ed. II. i. 116 = muralis.
incana, Willd. Herb. ex Steud. l. c. = collina.
incanescens, Munro, Pl. Agra, 6, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 146 = Farsetia Hamiltonii.
incarnata, *Lamb. ex DC. Syst.* ii. 210.—Sibir.
integrifolia, *Lapeyr. Hist. Abr. Pl. Pyr.* 585. *Suppl.* 98. —Europ.
Intermedia, *Bruggg. in Yahresb. Naturf. Ges. Graub.* II. xxv. (1882) 92.—Europ.
intermedia, Hoppe, ex Steud. Nom. ed. II. i. 116 = bellidifolia.
ionocalyx, *Boiss. & Heldr. Diagn.* Ser. I. viii. 20.— As. Min.
japonica, *Regel, Gartenfl.* (1863) 308. t. 414.— Japon.
japonica, A. Gray, in Mem. Am. Acad. N. S. vi. (1858) 381 = Arabis Stelleri.
jugicola, Jord. Diag. i. 121 = bellidifolia.
kamtschatica, Auct. vide *camtschatica*, Willd.
kamtschatica, *Fisch. ex DC. Syst.* ii. 231.—Kamtschatca.
kamtschatica, Pall. ex Ledeb. Fl. Ross. i. 185 = Sisymbrium mollissimum.
kansuensrnsis, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 149 = amplexicaulis.
Kochii, Jord. Diag. i. 113 = Gerardi.
kokanika, *Regel, Pl. Nov. Fedsch.* 6.—Turkestan.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 382.—Turkestan.
laciea, Stev. ex Rupr. Fl. Cauc. 389 = mollis.
laevigata, *DC. Syst.* ii. 227.—Am. bor.
laevigata, Hook. Fl. Bor. Am. i. 48 = Drummondii!
lanata, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 846.—Peruv.
lasiolobs, *Link, Enum. Hort. Berol.* ii. 168.—Mexico.
lateriphrenaea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 150 = A. Turrita.
latifolia, *Dur. in Expl. Sc. Algér.* t. 72.—Algeria.
latifolia, Willd. ex Ledeb. Fl. Ross. i. 181 = Parrya Lanuginosa.
lazica, *Sibth. & Sm. Prod. Fl. Graec.* ii. 38.—As. Min.
leptocarpa, *Wall, Cat.* n. 4764 = Sisymbrium Wallichii.
leptocarpaea, Fisch. ex Reichb. Fl. Germ. Excurs. 680 = alpina.
Lerchenfeldiana, Schur, Enum. Pl. Transs. 42 = alpina.
lenigata, Poir. Encyc. Suppl. i. 411 = laevigata.
libanica, Schrad. Ind. Sem. Hort. Goett. vii. (1832) 42 = Holboellii.
longistiqua, *C. & J. Presl, Delic. Prag.* 16.—Sicilia.

ARABIS:—
longifolia, DC. Syst. ii. 219 = albida.
longirostris, *S. Wats. in Bot. King, Exp.* 17.— Calif.
longisiliqua, Wallr. Sched. Crit. 359 = A. Turrita.
lucens, Dulac, Fl. Hautes-Pyr. 302 = bellidifolia.
lucida, *Linn. f. Suppl.* 296.—Europ.
ludovicana, *C. A. Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. 60.—Am. bor.
lusitanica, *Boiss. Diagn.* Ser. II. i. 20.—Lusitan.
Lyallii, *S. Wats. in Proc. Am. Acad.* xi. (1876) 122. —Am. bor. occ.
lyrata, DC. Syst. ii. 244 = canadensis.
lyrata, *Linn. Sp. Pl.* 665.—Am. bor.
maccloviana, *Hook. Ic. Pl.* t. 498.—Ins. Falkland.
macrocarpa, Torr. Mex. Bound. Rep. 32 = perfoliata.
madonia, *C. & J. Presl, Delic. Prag.* 15.—Sicilia.
major, S. F. Gray, Nat. Arr. Brit. Pl. ii. 676 = A. Turrita.
Marschalliana, Hort. ex Steud. Nom. ed. II. i. 116 = hirsuta.
mexicana, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 319.—Mexico.
microphylla, Nutt. ex Torr. & Gray, Fl. N. Am. i. 82 = Drummondii!
minor, Hort. Berol. ex DC. Syst. ii. 226 = incana.
mollis, Rafin. in Am. Monthly Mag. ii. (1817) 43 = canadensis.
mollis, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 270. —Reg. Caucas.
montana, Bernh. ex DC. Syst. ii. 225 = stricta.
montana, Lam. Encyc. i. 219 = hirsuta
Montbretiana, *Boiss. in Ann. Sc. Nat.* Ser. II. xvii. (1842) 58.—Orient.
monticola, Jord. Diag. i. 107 = alpina.
moschata, Hort. ex Reichb. Fl. Germ. Excurs. 680 = incana.
muliensis, Bell. ex DC. Syst. ii. 238 = Halleri.
multiflora, *DC. Syst.* ii. 243.—Mont. Pyren.
multijuga, *Borb. in Linnaea*, xlii. (1877) 604.— Europ.
muralis, *Bertol. Rar. Ital. Pl. Decad.* ii. 37.— Graecia.
muralis, Salisb. Prod. 272 = Sisymbrium Thalianum.
muricola, Jord. Diag. i. 125 = muralis.
Nasturtium, [Clairv.] Man. Herb. 223 = Nasturtium officinale.
neglecta, *Schult. Oestr. Fl.* ii. 248.—Mont. Carpath.
nemoralis, Steud. Nom. ed. II. i. 117 = hirsuta.
nemorensis, C. A. Mey. Verz. Pfl. Cauc. 179 = Gerardi.
nemorensis, Reichb. Fl. Germ. Excurs. 681 = hirsuta.
nepetaefolia, *Boiss. Fl. Orient.* i. 177.—Persia.
nivalis, Guss. Pl. Rar. 389 = serpyllifolia.
nivalis, Spreng. Syst. ii. 898 = Macropodium nivale.
Nordmanniana, Rupr. Fl. Cauc. 79 = mollis.
nova, Vill. Prosp. 39 = saxatilis.
nuda, Schreng. ex Reise, in Ann. Sc. Nat. Sér. II. xvii. (1842) 54 = Sisymbrium nudum.
nudicaulis, DC. Syst. ii. 240 = Parrya macrocarpa.
nudicaulis, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 22. —Afr. austr.
nutans, Moench, Meth. 258 = pumila.
nutans, Sims, Bot. Mag. t. 2219 = Allionii.
oblongata, *Wender. Hort. Marb.* (1824) ex Steud. Nom. ed. II. i. 117.—Hab.!
obtusifolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 58 = alpina.
ochroleuca, *Boiss. & Heldr. Diagn.* Ser. II. i. 21.— Graecia.
ochroleuca, Lam. Encyc. i. 218 = A. Turrita.
odyneida, Sibth. ex DC. Syst. ii. 219 = albida.
ovata, Poir. Encyc. Suppl. v. 557 = hirsuta.
ovirensis, Wulf. in Jacq. Coll. i. 196 = Halleri.
ovalifolia, *Chem. & Schlecht. in Linnaea*, i. (1826) 17.—Ins. Sandvic.
oxyota, DC. Syst. ii. 236 = pendula.
pachyrhiza, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 144.—Soongaria.
pangiensis, *Watt in Journ. Linn. Soc.* xviii. (1881) 376.—Reg. Himal.
parviflora, Rafin. in Am. Monthly Mag. ii. (1817) 43 = Sisymbrium Thalianum.
parvula, *Dufour, in DC. Syst.* ii. 228.—Europ. austr.
patens, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 142 = amplexicaulis.
patens, *Sullivant in Am. Journ. Sc.* i. (1819) 42, 49.— Am. bor.

ARABIS:—
Patrimiana, DC. Syst. ii. 236 = pendula.
patula, *Host, Fl. Austr.* ii. 271.—Austria.
patula, Torr. Bot. Mex. Bound. 32 = Holboellii.
patula, Weinm. Cat. Hort. Dorp. (1810) 18 = auriculata.
pauciflora, Garcke, Fl. Deutschl. ed. IV. 22 = brassicaeformis.
pedemontana, Bolss. Diagn Ser. I. i. 69 = cebennensis
pellegrea, Fenzl, in Kotschy, Keise Cilic. Taur. 891 = ionocalyx
pendula, Lachen. ex Reichb. Fl. Germ. Excurs. 681 = A. Turrita.
pendula, *Linn. Sp. Pl.* ed. I. 665.—Sibir.; Am. bor.
pendula, Stell. ex DC. Syst. ii. 242 = Stelleri.
perfoliata, *Lam. Encyc.* i. 219.—Europ.; Sibir.; Reg. Himal.
permixta, Jord. Diag. i. 110 = Gerardi.
petiolaris, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 187 —Texas.
petiolata, Bieb. Fl. Taur. Cauc. ii. 126 = Sisymbrium Allaria.
petraea, *Lam. Encyc.* i. 221.—Reg. bor. temp.
petricola, Jord. Diag. i. 121 = hirsuta.
petrogena, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 141 = arenosa.
Plaszekii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 490.—Chusa.
pinnatifida, Lam. Encyc. i. 221 = Braya pinnatifida.
planisiliqua, Pers. ex Reichb. Ic. Fl. Germ. xii. 18 = Gerardi.
platyosperma, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 519.—Calif.
praecox, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 664 = procurrens.
procera, Jord. Diag. i. 114 = Gerardi.
procurrens, *Waldst. & Kit. Pl. Rar. Hung.* ii. 154. t. 144.—Hungar.; Banat.; Thracia.
propera, Jord. Diag. i. 118 = hirsuta.
propinqua, Jord. l. c. = hirsuta.
pseudoturrita, *Boiss. & Heldr. Diagn.* Ser. II. i. 20.— Graecia.
pterosperma, Edgew. in Trans. Linn. Soc. xx. (1851) 33 = alpina.
pubercula, Nutt. ex Torr. & Gray, Fl. N. Am. i. 82 = canescens.
pubescens, *Poir. Encyc. Suppl.* i. 413.—Algeria.
pubicalyx, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 72. —Japon.
pubigera, Jord. Diag. i. 115 = Gerardi.
pumila, *Jacq. Fl. Austr.* iii. 44.—Europ.
purpurascens, Presl, Fl. Sicul. i. 50 = longisiliqua.
purpurea, *Sibth. & Sm. Fl. Graec.* vii. t. 643.—Ins. Cyprus.
ramosa, Lam. Fl. Fr. ii. 510 = Sisymbrium Thalianum.
recta, Vill. Hist. Pl. Dauph. iii. 514 = auriculata.
Reichenbachii, Syme, Eng. Bot. ed. III. i. 168 = hirsuta.
repanda, *S. Wats. in Proc. Am. Acad.* xi. (1876) 122. —Calif.
reptans, Lam. Encyc. i. 222 = Draba carolinianal
resedifora, H. B. & K. Nov. Gen. et Sp. v. 81 = Nasturtium arabiforme.
resedifolia, Lam. Fl. Fr. ii. 511 = Cardamine resedifolia.
retrofracta, R. Grah. in Edinb. Phil. Journ. (1827) 7 —Holboellii.
Retsiana, Beurl. ex Nym. Consp. i. 33 = hirsuta.
rhaetica, *Bruggg. in Yahresb. Naturf. Ges. Graub.* II. xxv. (1882) 68.—Europ.
rhombodea, Pers. Syn. ii. 204 = Cardamine rhomboidea.
rigida, Dulac, Fl. Hautes-Pyr. 300 = brassicaeformis.
rigida, Wallr. Beitr. Fl. Hercyn. 229 = auriculata.
rigidula, Jord. Diag. i. 108 = Gerardi.
Rosmarinea, Andrz. ex DC. Syst. ii. 236 = parvula.
rosea, DC. Syst. ii. 215 = muralis.
rosella, Jord. Diag. i. 126 = muralis.
rotundifolia, Rafin. in Am. Monthly Mag. ii. (1817) 43 = Draba caroliniana.
rubicaulis, Jord. Diag. i. 118 = Gerardi.
rugosa, Moench, Meth. 259 = A. Turrita.
runcinata, Lam. Encyc. i. 223 = petraea.
runcinata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 319.—Am. bor.
rupestris, Nutt. ex Torr. & Gray, Fl. N. Am. i. 81 = hirsuta.
sabauda, *Bruggg. in Yahresb. Naturf. Ges. Graub.* II. xxv. (1882) 67.—Europ.

ARABIS:—
sagittata, DC. Fl. Fr. vi. 592 = hirsuta.
saxatilis, *All. Fl. Pedem.* i. 268.—Europ.
saxaticola, Jord. Diag. i. 106 = alpina.
saxicola, *Edgew. in Trans. Linn. Soc.* xx. (1846) 83.—Reg. Himal.
saxigena, Jord. Diag. i. 127 = muralis.
scabra, All. Fl. Pedem. i. 269 = serpyllifolia pumila.
scabra, Gilib. Fl. Lituan. ii. 61 = Sisymbrium Thalianum.
scaligera, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 54 = Sisymbrium nudum.
Schievereckiana, *Andrz. in DC. Prod.* i. 145.—Tyrol.
Scopolitana, *Boiss. ex Nym. Consp.* i. 85.—Europ.
segetalis, Schur, Enum. Pl. Transs. 45 = arenosa.
serpyllifolia, Hoppe, ex DC. Syst. ii. 241 = vochinensis.
serpyllifolia, *Vill. Prosp.* 39.—Europ.
serrata, *Franch. & Sav. Enum. Pl. Jap.* i. 38.—Japon.
sibirica, *Hort. ex Steud. Nom.* ed. II. i. 117.—Sibir.
sicula, *Huet, ex Rent. Cat. Grass.* (1857).—Sicilia.
sicula, Ster. ex Nym. Suppl. 23 = alliaca.
Sieberi, *Tausch, in Flora*, xix. (1836) ii. 405.—Austr.
silicicolus, Lam. Encyc. i. 221 = Parrya macrocarpa.
simplicifolia, Schur, Enum. Pl. Transs. 48 = petraea.
sinaica, Boiss. Diagn. Ser. I. viii. 21 = auriculata.
sinaica, *Steud. Nom.* ed. II. i. 117, nom.—Arab.
sinuata, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 64 — Sibir.
sisymbroides, Hook. ex Hook. f. in Trans. Linn. Soc. xxiii. (1861) 314 = petraea.
Sophia, Bernh. Syst. Verz. Erf. 196 = Sisymbrium Sophia.
Soyeri, *Reut. & Huet, ex Huet, in Ann. Sc. Nat.* Sér. III. xix. (1853) 251.—Mont. Pyren.
sparsiflora, Nutt. ex Torr. & Gray, Fl. N. Am. i. 81 = Holboellii.
spathulata, *DC. Syst.* ii. 227.—Am. austr.
spathulata, *Nuttt. ex Torr. & Gray, Fl. N. Am.* i. 81.—Am. bor.
Stellari, *DC. Syst.* ii. 242.—As. bor. or.
stellulata, Desv. & Berthol. in Desv. Journ. Bot. ii. (1813) 76 = pumila.
stenocarpa, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 4 = saxatilis.
stenopetala, *Willd. Enum. Hort. Berol. Suppl.* 46.—Hab. ?
Steveniana, *Rupr. Fl. Cauc.* 289.—Reg. Cauc.
stolonifera, Clairv. ex DC. Syst. ii. 233 = Halleri.
stolonifera, *Hornem. Hort. Hafn.* ii. 618.—Hab. ?
stricta, Coste, Amplac. 23 = Costeæ.
stricta, *Huds. Fl. Angl.* 292.—Europ.; Am. bor.
stricta, Sieber, ex Tausch, in Flora, xix. (1836) ii. 406 = alpestris.
subcordacea, Gren. ex Nym. Consp. 84 = bellidifolia.
subnivalis, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 58.—Europ.
subnitens, Jord. Diag. i. 124 = alpestris.
subpinnatifida, *S. Wats. in Proc. Am. Acad.* xx. (1885) 253.—Am. bor. occ.
sudetica, Tausch, in Flora, xix. (1836) ii. 407 = hirsuta.
suecica, Fries, Summa Veg. 30, 147 = Sisymbrium suecicum.
suffrutescens, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 362.—Am. bor. occ.
sulphurea, *Boiss. Fl. Orient.* i. 172.—Persia.
supina, Lam. Fl. Fr. ii. 513 = Sisymbrium supinum.
sylvestris, Scop. Fl. Carn. ed. II. ii. 33 = Brunica Cheiranthus.
taraxacifolia, T. Anders. in Hook. f. Fl. Brit. Ind. i. 136 = arvensis.
tatarica, Pall. Reise, ii. 676 = Parrya macrocarpa.
tenella, *Guss. & Tenore, Fl. Neap. Syll. App.* v. 20.—Lucania.
tenella, Host, Fl. Austr. ii. 378 = Halleri.
Tenorii, Huet, ex Nym. Consp. 84 = alliaca.
Thaliana, Cranz, Stirp. Austr. ed. II. 89 = petraea.
Thaliana, Linn. Sp. Pl. 665 = Sisymbrium Thalianum.
Thomsoni, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 145 = tibetica.
thyrsoidea, Sibth. & Sm. Fl. Græc. t. 642 = alliaca.
tibetica, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 148.—Reg. Himal.
tomophylla, Bieb. Fl. Taur. Cauc. iii. 448 = Sisymbrium tomophyllum.
transsilvanica, Schur, Enum. Pl. Transs. 48 = petraea.

ARABIS:—
trichopoda, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 68.—Rossia.
tuberosa, Mahl. ex Pers. Syn. ii. 204 = Cardamine rhomboidea.
Turczaninowii, *Ledeb. Fl. Ross.* i. 128.—Sibir.
Turrita, *Linn. Sp. Pl.* 665.—Europ.; Orient.
Turrita, [Clairv.] Man. Herb. 238 = A. Turrita.
umbrosa, Crantz, Stirp. Austr. ed. I. fasc. i. 48; ed. II. 61 = A. Turrita.
umbrosa, Turcz. ex Stend. Nom. ed. I. ii. 118 = petræa.
undulata, Link, Enum. Hort. Berol. ii. 161 = alpina.
uralensis, Willd. ex Ledeb. Fl. Ross. i. 130 = petræa.
vera, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 105.—Reg. Mediterr.
versuta, Jord. Diag. i. 123 = alpestris.
violacea, Moench, Meth. 259 = verna.
virescens, Jord. Diag. i. 109 = Gerardi.
virginica, Poir. Encyc. Suppl. i. 413 = Cardamine hirsuta.
viscosa, DC. Syst. ii. 216 = albida.
vochinensis, *Spreng. Pugill.* i. 46. n. 81.—Austria.
wolgensis, Willd. ex Ledeb. Fl. Ross. i. 135 = Sisymbrium toxophyllum.
yokoscensis, *Franch. & Sav. Enum. Pl. Jap.* i. 84.—Japon.
Zeyheriana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 292.—Afr. austr.

ARABISA, Reichb. Handb. 260 (1837) = **Arabis**, Linn. (Crucif.).

ARACHIDNA, Plum. ex Moench, Meth. 121 (1794) = **Arachis**, Linn. (Legum.).
hypogæa, Moench, l. c. 122 = Arachis hypogæa.
quadrifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 7.—Malaya.

ARACHIS, Linn. Syst. ed. I. (1735). *LEGUMINOSÆ*, Benth. & Hook. f. i. 518.
ARACHIDNA, Moench, Meth. 121 (1794).
MUNDUBI, Adans. Fam. ii. 323 (1763).
africana, Burm. f. Fl. Cap. Prod. 73 = Voandzeia subterranea.
africana, Lour. Fl. Cochinch. ii. 430 = hypogæa.
americana, Tenore, in Mem. Ist. Incorr. i. (1811) 48 = hypogæa.
aprica, Walt. Fl. Carol. 183 = Stylosanthes elatior.
asiatica, Lour. Fl. Cochinch. ii. 430 = hypogæa.
fruticosa, Retz. Obs. Fasc. v. 26 = Stylosanthes mucronata.
glabrata, *Benth. in Trans. Linn. Soc.* xviii. (1841) 159.—Bras.
hypogæa, *Linn. Sp. Pl.* 741.—Reg. trop. (cult.).
marginata, *Gardn. in Hook. Ic. Pl.* t. 500.—Bras.
prostrata, *Benth. in Trans. Linn. Soc.* xviii. (1841) 159.—Bras.
pusilla, *Benth. l. c.*—Bras.
tuberosa, *Benth. l. c.*—Bras.
villosa, *Benth. l. c.*—Am. austr.

ARACHNANTHE, Blume, Rumphia, iv. 55. t. 196 (1838). *ORCHIDEÆ*, Benth. & Hook. f. iii. 572.
ARACHNIS, Blume, Bijdr. 365 (1825).
ARACHNORHA, Breda, Orch. Kabl. & Hass. (1827).
ARAHNYNCHIUM, Lindl. in Paxt. Fl. Gard. i. 142 (1851).
EDERALDA, Reichb. f. Xenia Orch. ii. 88 (1874).
bilingvis, *Benth. in Journ. Linn. Soc.* xviii. (1881) 332.—Ind. or.
Cathcartii, *Benth. & Hook. f. Gen.* iii. 578.—Reg. Himal.
Lowei, *Benth. & Hook. f. l. c.*—Borneo.
moschifera, *Blume, Rumphia*, iv. 55.—Malaya.
Salingii, *Benth. in Journ. Linn. Soc.* xviii. (1881) 332. —Malaya.

ARACHNE, Neck. Elem. ii. 348 (1790) = **Andrachne**, Linn. (Euphorb.).

ARACHNIMORPHA, Desv. in Ham. Prod. Fl. Ind. Occ. 28 (1825) = **Rondeletia**, Linn. (Rubiac.).
incana, Ham. l. c. = R. incana.

ARACHNIS, Blume, Bijdr. 365 (1825) = **Arachnanthe**, Blume (Orchid.).
moschifera, Blume, l. c. = Arachnanthe moschifera.

ARACHNITES, F. W. Schmidt, Fl. Boëm. i. 74 (1793) = **Ophrys**, Linn. (Orchid.).
alpina, F. W. Schmidt, l. c. = Herminium alpinum.
anthropophora, F. W. Schmidt, l. c. 77 = Aceras anthropophora.
apifera, Hoffm. Deutschl. Fl. ed. II. ii. 180 = O. apifera.
atrata, Tod. Hort. Panor. ii. t. 28 = O. aranifera.
Benantii, Tod. l. c.—Hab. ?
Bertolonii, Tod. Orch. Sicul. 79 = O. Bertolonii.
Biancae, Tod. l. c. 83 = O. Biancae.
bombyliflora, Tod. l. c. 91 = O. bombyliflora.
fuciflora, F. W. Schmidt, Fl. Boëm. i. 76 = O. Arachnites.
fusca, Tod. l. c. 98 = O. fusca.
Insectifera, Tod. Nouv. Gen. 12 = O. Insectifera.
lunulata, Tod. Orch. Sicul. 77 = O. lunulata.
lutea, Tod. l. c. 96 = O. fusca.
Monorchis, Hoffm. Deutschl. Fl. ed. II. ii. 179 = Herminium Monorchis.
muscifera, F. W. Schmidt, Fl. Boëm. i. 75 = O. muscifera.
oxyrhynchus, Tod. Orch. Sicul. 81 = O. Arachnites.
pallida, Tod. l. c. 100 = O. pallida.
Speculum, Tod. l. c. 95 = O. Speculum.
tenthredinifera, Tod. l. c. 86 = O. tenthredinifera.

ARACHNITIS, Phil. in Bot. Zeit. 217 (1864). *BURMANNIACEÆ*, Benth. & Hook. f. iii. 460. unifora, *Phil. l. c.*—Chili.

ARACHNOPOGON, Berg. ex Haberl, Cat. Pesth.: ex Steud. Nom. ed. II. i. 118 (1840), nomen.
heterospermum, Berg, l. c.—Hab. ?

ARACHNOSPERMUM, Berg (sec. Steud. Nom. ed. II. i. 118) = **Hypochæris**, Linn. (Compos.).

ARACHNOSPERMUM, F. W. Schmidt, Samml. Phys. Aufs. i. 274 (1795) = **Scorzonera**, Linn. (Compos.).
lacinistum, F. W. Schmidt, l. c. = S. laciniata.
resedifolium, F. W. Schmidt, l. c. = prænc. ?

ARACHNOTHRIX, Walp. Ann. ii. 776 (1852) = seq.

ARACHNOTHRYX, Planch. in Fl. des Serres, Sér. I. v. (1849) sub t. 442 = **Rondeletia**, Linn. (Rubiac.).
buddleioides, Planch. l. c. = R. buddleoides.
discolor, Planch. l. c. = R. discolor.
elongata, Planch. l. c. = R. elongata.
hispidula, Griseb. Fl. Brit. W. Ind. 325.—Ind. occ.
laniflora, Planch. in Fl. des Serres, Sér. I. v. (1849) sub t. 442 = R. laniflora.
leucophylla, Planch. l. c. = R. leucophylla.
reflexa, Planch. l. c. = R. reflexa.
rosea, Lindex, Hort. Linden. (1859) 3.—N. Granat.

ARACHUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 360 = **Vicia**, Tourn. (Legum.).
viciodes, Medic. l. c. = V. bithynica.

ARACIUM, Neck. Elem. i. 49 (1790) = **Orepis**, Linn. (Compos.).
alpinum, Mönn. Ess. Hierac. 73 = C. alpina.
aureum, D. Dietr. Syn. Pl. iv. 1839 = C. aurea.
chondrilloides, D. Dietr. l. c. 1830 = C. jacquini.
cydoniæfolium, D. Dietr. l. c. = Hieracium cydoniæfolium.
Froelichianum, D. Dietr. l. c. = C. incarnata.
Gmelini, D. Dietr. l. c. = C. Gmelini.
incarnatum, D. Dietr. l. c. 1839 = C. incarnata.
longipetum, Miq. Fl. Ind. Bat. ii. 111 = C. laevigata.
lyratum, Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) 1. 214 = C. lyrata.
mauritianum, D. Dietr. Syn. Pl. iv. 1830 = C. mauritianum.
multicaule, D. Dietr. l. c. = C. multicaulis.
paludosum, Mönn. Ess. Hierac. 73 = C. paludosa.
Paucicii, Sch. Bip. in Pollichia, xxii–iv. (1866) 260.—Europ.
praemorsum, D. Dietr. Syn. Pl. iv. 1839 = C. praemorsa.
prenanthoides, D. Dietr. l. c. 1830 = Hieracium prenanthoides.
pubescens, C. Koch, in Linnæa, xxiii. (1850) 661.—An. Min. ?
pygmæum, Miq. Fl. Ind. Bat. ii. 111 = C. multicaule.
succisæfolium, D. Dietr. Syn. Pl. iv. 1839 = C. succisæfolia.

ARAEOCOCCUS, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 370. *BROMELIACEAE*, Benth. & Hook. f. iii. 664.
micranthus, *Brongn. l. c.*—Guiana.

ARAGUS, Steud. Nom. ed. II. i. 118 (1840) = ARAGALLUS, Neck. = **Astragalus**, Tourn. (Leguminos.).

ARALIA, Tourn. ex Linn. Syst. ed. I. (1735); Gen. ed. I. 88 (1737). *ARALIACEAE*, Benth. & Hook. f. i. 936. AURELIANA, Lafit. ex Catesb. Nat. Hist. Carol. App. 16 (1754).
BLAKWELLIA, Scop. Introd. 326 (1777).
DIMORPHANTHUS, Miq. Comm. Phyt. 95 (1840).
GYNAPTERIA, Spach, Hist. Vég. Phan. viii. 113 (1839).
MOTHERWELLIA, F. Muell. Fragm. vii. 107 (1870).
NALAGU, Adans. Fam. ii. 445 (1763).

abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 336 = Heptapleurum abyssinicum.
acerifolia, Willd. ex Roem. & Schult. Syst. vi. 699 = Oreopanax acerifolium.
aculeata, Ham. ex Seem. in Journ. Bot. vi. (1868) 136 = Brassaiopsis aculeata.
acutifolia, Willd. ex Roem. & Schult. Syst. vi. 697 = Brassaiopsis aculeata.
angularis, Willd. l. c. 698 = Oreopanax angulare.
arborea, Linn., Fl. Jam. Pugill. ii. 11; Syst. ed. X. 967 = Dendropanax arboreum.
arborea, Vell. Fl. Flum. 129; iii. t. 110 = Gilibertia brasiliensis.
argentata, H. B. & K. Nov. Gen. et Sp. v. 4 = Oreopanax argentatum.
aspera, *Linden, Cat.* n. 18. p. 7.—Hab.?
armata, *Seem. in Journ. Bot* vi. (1868) 134.—Ind. or.
aromatica, Blume, Bijdr. 571 (= *Agalma aromaticum*). —Java.
aviculariaefolia, H. B. & K. Nov. Gen. et Sp. v. 2. t. 2 = Oreopanax aviculariaefolium.
Bastardiana, Decne. Bot. Vénus, 24 = Panax Bastardianum.
bipinnata, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 83 = montana.
bipinnata, Blanco, Fl. Filip. ed. I. 222 = javanica.
bipinnatifida, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 732.—Reg. Himal.
brevifolia, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 74.—Mexic.
cachemirica, *Decne. in Jacquem. Voy. Bot.* 72. t. 81.—Reg. Himal.
calyculata, Zoll. & Moritzi, Syst. Verz. Zoll. 41 = Macropanax oreophilum.
camerana, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 94 = spinosa.
capitata, Jacq. Enum. Pl. Carib. 18 = Oreopanax capitatum.
capitulata, *Yungh. & De Vriese, in Ann. Sc. Nat.* Sér. III. vi. (1846) 116.—Sumatra.
catalpaefolia, Willd. ex Roem. & Schult. Syst. vi. 697 = Oreopanax capitatum.
cheirophylla, Spreng. Syst. i. 953 = Oreopanax cheirophyllum.
chinensis, Blume, Bijdr. 570 = dasyphylla.
chinensis, Linn. Sp. Pl. 378 = spinosa.
cissifolia, *Griff. ex Seem. in Journ. Bot.* vi. (1868) 134. —Reg. Himal.
cochleata, Lam. Encyc. i. 234 = Panax cochleatum.
cordata, *Thunb. Fl. Jap.* 197.—Japon.
crassifolia, Soland. ex A. Cunn. in Ann. Nat. Hist. ii. (1839) 314 = Pseudopanax crassifolium.
crassinervia, H. B. & K. Nov. Gen. et Sp. v. 5 = Oreopanax crassinervium.
cumanensis, H. B. & K. l. c. 3 = Oreopanax cumanense.
dasyphylla, *Miq. Fl. Ind. Bat.* i. t. 751.—Java.
Decaisneana, Hance, in Journ. Bot. iv. (1866) 172 = spinosa.
decomposita, *Reinw. ex De Vriese, Pl. Ind. Bat. Reinw.* 84.—Java.
digitata, Roxb. Hort. Beng. 23; Fl. Ind. ii. 107 = Heptapleurum venulosum.
digitata, Willd. ex Roem. & Schult. Syst. vi. 701 = Oreopanax xalapense.
discolor, H. B. & K. Nov. Gen. et Sp. v. 4 = Oreopanax discolor.
disperma, Blume, Bijdr. 572 = Macropanax oreophilum.

ARALIA :—
dubia, Spreng. Syst. iv. Cur. Post. 125 = Trevesia palmata.
Echinops, Cham. & Schlecht. in Linnaea, v. (1830) 409 = Oreopanax Echinops.
edulis, Sieb. & Zucc. Fl. Jap. i. 57. t. 25 = cordata.
elata, Seem. in Journ. Bot. vi. (1868) 134 = spinosa.
elegantissima, *Veitch, in Gard. Chron.* (1872) 782.—Hab.?
ernacea, Hook. in Brewst. Edinb. Journ. Sc. vi. (1827) 64 = fatsia horrida.
Faribuki, *Sieb. ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 16.—Japon.
farinosa, Delile, ex Ferr. & Galin. Voy. Abyss. iii. 135 = Panax pinnatum.
ferox, *Miq. Fl. Ind. Bat.* i. t. 750.—Malaya.
fragrans, Willd. ex Roem. & Schult. Syst. vi. 701 = Sciadophyllum ferrugineum.
fruticifolia, *Chr. Moore, ex Fourn. in Illustr. Hortic.* xxii. (1876) 72. t. 240.—Ins. Pacif.
Finlaysoniana, *Seem. in Journ. Bot.* vi. (1868) 134.—Ind. or.
floribunda, H. B. & K. Nov. Gen. et Sp. v. 6. t. 416 = Oreopanax floribundum.
folliolosa, *Seem. in Journ. Bot.* vi. (1868) 134.—Reg. Himal.
fragrans, G. Don. ex Loud. Hort. Brit. 112 = Heteropanax fragrans.
Gemma, *Linden.—Cf. Gard. Chron.* (1880) ii. 150.—N. Caled.
gracilipes, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 8.—Ambor.
Gheebrechtii, Versch. ex Morr. in Belg. Hortic. xiii. (1863) 8 = Oreopanax xalapense.
Ginseng, *Basill. Hist. Pl.* vii. 197.—China.
glomerulata, Blume, Bijdr. 573 = Brassaiopsis speciosa.
gracilima, Linden, in Rev. Hortic. (1877) 88 = Veitchii.
grandis, Miq. Comm. Phyt. 95 = elata.
Guillofyei, *Cogn. & March. Pl. Ornem.* ii. (1874). t. 58.—Ins. Pacif.
heptaphylla, Willd. ex Spreng. Syst. i. 952 = Sciadophyllum ferrugineum.
heterophylla, A. Cunn. ex Hook. Ic. Pl. t. 583-584 = Pseudopanax crassifolium.
heterophylla, Mutis, ex H. B. & K. Nov. Gen. et Sp. v. 1 = Oreopanax Mutisianum.
heterophylla, Willd. ex Roem. & Schult. Syst. vi. 698 = Oreopanax acerifolium.
hispida, *Vent. Jard. Cels*, t. 41; *Michx. Fl. Bor. Am.* i. 183.—Am. bor.
Humboldtiana, Roem. & Schult. Syst. vi. 699 = Oreopanax floribundum.
humilis, *Cav. Ic.* iv. 7. t. 318.—N. Mexic.
hypoleuca, Presl, Epim. Bot. 250 = spinosa.
incisa, Willd. ex Roem. & Schult. Syst. vi. 699 = Oreopanax floribundum.
japonica, Hort. ex Seem. in Journ. Bot. vi. (1868) 135 = Fatsia japonica.
japonica, Thunb. Fl. Jap. 128 = Fatsia japonica.
jatrophaefolia, H. B. & K. Nov. Gen. et Sp. v. 6 = Oreopanax jatrophaefolium.
javanica, *Miq. Fl. Ind. Bat.* i. t. 749.—Java.
Kleinii, Steud. Nom. ed. II. i. 118 = Miquelia Kleinii.
laciniata, C. Gay, Fl. Chil. iii. 151 (= *Cheirodendron lasciviense*).—Chili.
lappaefolia, *Raeusch. Nom.* ed. III. 88; *et Rets. Obs.* vi. 37.—Ind. or.
Lariboki, *Kriebe, ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 16.—Japon.
latifolia, Wight & Arn. Prod. 376 = Polyscias pinnata.
Lavrana, *C. Koch, Wochenschr.* vii. (1864) 369.—Ambor.
Lessoni, Hook. f. Fl. Nov. Zel. 96 = Pseudopanax Lessonii.
longifolia, *D. Dietr. Syn. Pl.* ii. 1085.—Java.
longifolia, Reinw. ex App. Hort. Spaarn. Berg. 2 = Brassaia littorea.
lucescens, Blume, Bijdr. 572 (= Heptapleuri sp.).—Java.
lucida, *Hassk. Cat. Hort. Bogor. Alt.* 164.—Malaya.
lutescens, Auct. ex Steud. Nom. ed. II. err. typ. = Inconcens.
Lyallii, *T. Kirk, in Trans. N. Z. Inst.* xvii. (1885) 222, 295 = (Pseudopanacis sp. ?).—N. Zel.
macrophylla, A. Cunn. in Sweet, Hort. Brit. Suppl. i. 601 = Maryta latifolia.
macrophylla, Lindl. in Bot. Reg. (1844) Misc. 78 = cachemirica.

ARALIA :—
malabarica, *Bedd. Fl. Sylv.* ii. 121.—Ind. or.
mandschurica, Seem. in Journ. Bot. vi. (1868) 134 = spinosa.
mandshurica, *Rupr. in Bull. Acad. Pétersb.* xv. (1856) 134.—As. or.
Maralia, Roem. & Schult. Syst. vi. 704 = Panax Maralia.
Maximowiczii, Van Houtte, Fl. des Serres, Sér. II. xx. (1874) 89. t. 2067 = Acanthopanax ricinifolium.
micans, Willd. ex Roem. & Schult. Syst. vi. 701 = Panax splendens.
Mitsde, Siebold, in Linnaea, xxvi. (1853-55) 754 = Dendropanax japonicum.
monogynia, Vell. Fl. Flum. 129; iii. t. 111 = Gilibertia monogyna.
montana, *Blume, Bijdr.* 570.—Java.
Moorei, F. Muell. Fragm. ii. 108 = Heptapleurum venulosum.
Motherwelli, *F. Muell. l. c.* vii. 107.—Austral.
Muehlenbergiana, Roem. & Schult. Syst. vi. 704 = hispida.
multiflora, Pohl, ex DC. Prod. iv. 263 = Oreopanax capitatum.
Mutisiana, H. B. & K. Nov. Gen. et Sp. v. 1 = Oreopanax Mutisianum.
nobilis, *Hort. Williams.—Cf. Gard. Chron.* (1878) ii. 56; (1880) i. 759.—Hab.?
nodosa, Blume, Bijdr. 573 = Polyscias nodosa.
nudicaulis, Blume, l. c. 570 = cordata.
nudicaulis, *Linn. Sp. Pl.* 274.—Am. bor.
nutans, *Franch. & Sav. Enum. Pl. Jap.* ii. 376.—Japon.
obtusifolia, H. B. & K. Nov. Gen. et Sp. v. 5 = Oreopanax obtusilobum.
occidentalis, Schlecht. ex Ledeb. Fl. Ross. ii. 875 = Fatsia horrida.
octonata, Stokes, Bot. Mat. Med. ii. 181 = Heptapleurum octophyllum.
octophylla, Blanco, Fl. Filip. ed. II. 158 = Heptapleurum rigidum.
octophylla, Lour. Fl. Cochinch. i. 187 = Heptapleurum octophyllum.
palmata, Lam. Encyc. i. 234 = Trevesia Zippeliana.
palmata, Lour. Fl. Cochinch. 187 = Brassaiopsis Hainla.
palmata, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 82 = Trevesia sundaica.
palmata, Willd. ex Roem. & Schult. Syst. vi. 699 = Oreopanax chelrophyllum
paniculata, Phil. in Linnaea, xxviii. (1856) 694 = Panax valdiviense.
papyrifera, Hook. Kew Journ. iv. (1852) 50 = Fatsia papyrifera.
parvifolia, *Panch. & Sebert, Not. Bois. Nouv. Calléd.* 204.—N. Caled.
pendula, Blanco, Fl. Filip. ed. I. 223 = Polyscias nodosa.
pennata, Poir. Encyc. Suppl. i. 418, sphalm. = Trevesia palmata.
pentaphylla, Thunb. Fl. Jap. 128 = Acanthopanax spinosum.
pergamacea, Blume, Bijdr. 573 = Heptapleurum pergamacum.
pinnata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 335.—Abyss.
pinnatifida, Jungh. & De Vriese, in Ann. Sc. Nat. Sér. III. vi. (1846) 115 = Araliidium pinnatifidum.
Planchoniana, Hance, in Journ. Bot. iv. (1866) 172 = Acanthopanax spinosum.
platanifolia, H. B. & K. Nov. Gen. et Sp. v. 6 = Oreopanax platanifolium.
polaris, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zel. 86 = Stilbocarpa polaris.
Polyscias, Spreng. ex Seem. in Journ. Bot. vi. (1868) 138 = Polyscias umbellata.
Pseudo-ginseng, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 731.—Reg. Himal.
pubescens, DC. Cat. Hort. Monsp. (1813) 80 = humilis?
quinduensis, H. B. & K. Nov. Gen. et Sp. v. 8 = Sciodaphyllum quinduense.
quinquefolia, *Decne. & Planch. in Rev. Hortic. Sér.* IV. iii. (1854) 105.—Am. bor.
racemosa, *Linn. Sp. Pl.* 378.—Am. bor.
ramiflora, Pohl, ex March. in Mart. Fl. Bras. xi. i. 250 = Gilibertia cuneata.
Regeliana, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 73.—Mexico.
Reinwardtiana, Seem. in Journ. Bot. vi. (1868) 136 = Trevesia sundaica.

ARALIA :—
reticulata, Willd. ex Roem. & Schult. Syst. vi. 699 = Oreopanax reticulata.
rigida, Blume, Bijdr. 874 = Heptapleurum rigidum.
rupsea, Blume, l. c. 879 (= Heptapleuri sp.).—Java.
sallaifolia, *Vent. ex Seem. in Journ. Bot.* vi. (1868). 188.—Hab.!
scabra, Presl, ex DC. Prod. iv. 258 = pubescens.
scandens, Poir. Encyc. Suppl. i. 419 = Brassiopsis Hainla.
Scheffera, Spreng. Pl. Pugill. i 28 = Scheffiera digitata.
Sciadophyllum, Sw. Prod. Veg. Ind. Occ. 55 = Sciodaphyllum Browne.
septemnervia, H. B. & K. Nov. Gen. et Sp. v. 3 = Oreopanax capitatum.
Steboldii, Hort. ex C. Koch, Dendrol. i. 678 = Fatsia japonica.
similima, Blume, Bijdr. 871 (= Heptapleuri sp.).—Java.
sonetensis, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 76.—Bolivia.
spectabilis, Hort. Bull.—Cf. Gard. Chron. (1876) i. 603 = filicifolia.
spinosa, *Linn. Sp. Pl.* 273.—Am. bor.
tarchonanthifolia, Willd. ex Roem. & Schult. Syst. vi. 698 = Oreopanax sylvenniæfolium.
tenuifolia, *Pancher, ex Baill. Adansonia*, x. (1871-72) 372.—N. Caled.
Thibautii, *Versch. ex Morr. in Belg. Hortic.* xiii. (1863). —Mexic.
Thomsonii, *Seem. in Journ. Bot.* vi. (1868) 134.—Reg. Himal.
trifolia, Banks & Soland. ex Walp. Ann. v. 84 = Pseudopanax crassifolium.
trifolia, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 105.—Am. bor.
trifoliata, Meyen, Reise, ii. 332 = Acanthopanax aculeatum.
trigyna, Gaudich. Voy. t. 98 (= Panacis sp.).—Hab.!
trphylla, Poir. Encyc. Suppl. i. 418 = trifolia.
tripinnata, Blanco, Fl. Filip. ed I. 223 = Panax fruticosum.
turbacensis, H. B. & K. Nov. Gen. et Sp. v. 3 = Oreopanax turbacense.
umbellata, Pav. ex Seem. in Journ. Bot. vi. (1868) 138 = Gilibertia umbellata.
umbellata, Pohl, ex March. in Mart. Fl. Bras. xi. t. 250 = Gilibertia cuneata.
umbelliflora, Lam. Encyc. i. 325 = Osmoxylon ambolnense.
umbraculifera, Roxb. Hort. Beng. 22; Fl. Ind. iii. 108 = Polyscias nodosa.
urticæfolia, *Blume, ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 9.—Hab.!
valdivientis, C. Gay, Fl. Chil. iii. 152 = Panax valdiviense.
Veitchii, *Hort. ex T. Moore, in Flor. & Pom.* (1874) 5. —Hab.!
vitiensis, A. Gray, Bot. U. St. Expl. Exped. i. 715 = Scheffiera vitiensis.
Wallichiana, *D. Dietr. Syn. Pl.* ii. 1035.—Ind. or.
xalapensis, H. B. & K. Nov. Gen. et Sp. v. 8. = Oreopanax xalapense.

ARALIDIUM, Miq. Fl. Ind. Bat. i. t. 762. t. 13 (1855). *ARALIACEÆ*, Benth. & Hook. f. i. 936.
dentatum, *Miq. Fl. Ind. Bat. Suppl.* 840.—Sumatra.
pinnatifidum, *Miq. l. c.* i. t. 763. t. 13.—Malacca.

ARALIOPSIS, Kurz, in Andam. Rep. 89 (1870) = **Brassaiopsis**, Decne. & Planch. (Araliac.).
andamanica, Kurz, l. c. = B. palmata.

ARAPABACA, Adans. Fam. ii. 225 (1763) = **Spigelia**, Linn. (Logan.).

ARARIBA, Mart. in Sitzb. Acad. Muench. (1860) 328 = **Hicksiella**, Willd. (Rubiac.).
alba, Mart. l. c. 324.—Bras.
rubra, Mart. l. c.—Bras.

ARBOCARPUS, Scheff. in Ann. Jard. Buitens. ii. (1885) 10. *ANONACEÆ*.
velutinus, *Scheff. l. c.*—Java.

ARASCHCOOLIA, Sch. Bip. Pl. Kotsch. Nub. Exs. n. 101 (1878); ex Benth. & Hook. f. Gen. ii. 387 = **Centauria**, Grisebach (Compos.).
acaulis, Sch. Bip. ex Oliver, Fl. Trop. Afr. iii. 368 = G. acaulis.

ARAUCARIA, Juss. Gen. Pl. 418 (1789). *CONIFERÆ*, Benth. & Hook. f. iii. 427.
ALTINGIA, Loud. Hort. Brit. 408 (1830).
COLUMBEA, Salisb. in Trans. Linn. Soc. viii. (1807) 317.
DOMBEYA, Lam. Encyc. ii. 301. t. 838 (1786).
EUTACTA, Link, in Linnæa. xv. (1841) 548.
EUTASSA, Salisb. in Trans. Linn. Soc. viii. (1807) 316.
QUADRIFARIA, Manetti, ex Gord. Pinet. Suppl. 14 (1859).
albopica, Hort. ex Carr. Conif. ed. II. 611 = excelsa.
araucana, C. Koch, Dendrol. ii. ii. 206 = imbricata.
Balansae, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. xiii. (1870-71) 351.—N. Caled.
Bibbiani, Hort. Ital. ex Gord. Pinet. 24 = brasiliana.
Bidwillii, *Hook. Lond. Journ. Bot.* ii. (1843) 503. t. 18, 19.—Australi.
brasiliana, *A. Rich. in Dict. Class. Hist. Nat.* i. 512.—Bras.
brasilieensis, Loud. Hort. Brit. 408 = præc.
chilensis, Mirb. in Mém. Mus. Par. xiii. (1825) 49 = imbricata.
columnaris, Hook. in Bot. Mag. t. 4685 = Cookii.
Cookii, *R. Br. ex D. Don, in Trans. Linn. Soc.* xviii. (1841) 164.—N. Caled.
Cunninghamii, *Sweet, Hort. Brit.* ed. II. 475.—Austral.
Dombeyi, A. Rich. Conif. 86. t. 20 = imbricata.
elegans, Hort. ex Carr. Conif. ed. I. 415 = brasiliana.
excelsa, *R. Br. in Ait. Hort. Kew.* ed. II. v. 412.—Austral.; ins. Norf.
glauca, Antoine, Conif. 105. t. 44 = Cunninghamii.
Goldieana, *Hort. ex T. Moore, Flor. & Pom.* (1877) 89 —N. Caled.
gracilis, Hort. ex Carr. Conif. ed. I. 415 = brasiliana.
imbricata, *Pav. in Mem. Acad. Med. Madr.* i. (1797) 199.—Chili.
intermedia, *R. Br. ex Virill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 54.—N. Caled.
lanceolata, Hort. ex Carr. Conif. ed. I. 171 = Cunninghamia sinensis.
Lindleyana, Van Houtte, ex Gord. Pinet. 24 = brasiliana.
montana, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. xiii. (1870-71) 356.—N. Caled.
muelleri, *Brongn. & Gris, l. c.* 362.—N. Caled.
Nidolfiana, Savi f. in Atti Firenz. (1841) 458 et 788; et Giorn. Bot. Ital. ii. (1846) fasc. i. 62 = brasiliana.
Rulei, *F. Muell. ex Lindl. in Gard. Chron.* (1861) 868. —N. Caled.
Saviana, *Parl. in Bull. Soc. Bot. Fr.* viii. (1861) 89.— Bolivia.
subulata, *Virill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 55.—N. Caled.

ARAUCASIA, Benth. & Hook. f. Gen. iii. 919 (1888), *nom.* = ARAUSIACA, Blume = **Oraxia**, Zipp. (Palm.).

ARAUJIA, Brot. in Trans. Linn. Soc. xii. (1818) 62. *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 761.
LAGENNIA, Fourn. in Mart. Fl. Bras. vi. iv. 298 (1885).
PHYSANTHRAGMA, Zucc. ex Reichb. Consp. 131 (1828).
PHYSANTHUS, Mart. Nov. Gen. et Sp. i. 53. t. 32 (1824).
SCHLECHTERIA, Mart. l. c. 55. t. 33 (1824).
albens, G. Don, Gen. Syst. iv. 149 = sericifera.
angustifolia, *Steud. Nom.* ed. II. i. 119.—Bras.
calycina, Decne. in DC. Prod. viii. 534 = sericifera.
fusca, Griseb. ex Lorentz, in Goett. Abh. xxiv. (1879) 282 = megapotamica.
hortorum, *Fourn. in Mart. Fl. Bras.* vi. iv. 298.—Bras.
megapotamica, *G. Don, Gen. Syst.* iv. 149.—Bras.
sericifera, *Brot. in Trans. Linn. Soc.* xii. (1818) 62.—Peruv.
subhastata, *Fourn. in Mart. Fl. Bras.* vi. iv. 298.—Bras.
undulata, Visiani, Hort. Patav. (1832) 134 = hortorum.

ARAUSIACA, Blume, Ramphia, in præf. p. 8. (1836) tt. 119, 132 = **Oraxia**, Zipp. (Palm.).
excelsa, Blume, l. c. = O. regalis.

ARBUTUS, [Tourn.] Linn. Syst. ed. I (1735). *ERICACEÆ*. Benth. & Hook. f. ii. 631.
UBEDO, Hoffingg. & Link, Fl. Port. i. 418 (1809).
acadiensis, Linn. Sp. Pl. 395 (planta obscura).—Am. bor.

ARBUTUS :—
acerba, Gilib. Fl. Lituan. i. 5 = Arctostaphylos Uva-ursi.
alpina, Linn. Sp. Pl. 395 = Arctostaphylos alpina.
Andrachne, *Linn. Syst.* ed. X. 1024. n. 468.—Reg. Mediterr. or.
andrachnoides x, *Link, Enum. Hort. Berol.* i. 395.— Hab.!
arguta, Zucc. ex Steud. Nom. ed. II. i. 120, 219 = Arctostaphylos arguta.
buxifolia, Stokes, Bot. Mat. Med. ii. 509 = Arctostaphylos Uva-ursi.
Callicarpa, Buch, Canar. 146 = canariensis.
canadensis, *Pair. ex Steud. Nom.* ed. I. 82.—Canada
canariensis, *Duham. Arb.* ed. II. i. 89.—Ins. Canar.
cassinifolia, *Hort. ex Steud. Nom.* ed. II. i. 119.— Ins. Madera.
coriacea, Blume, Cat. Gew. Buitenz. 67 = Vaccinium Korthalsii.
crispa, Hoffingg. Verz. Pfl. Nachtr. i (1824) 204 = A Unedo.
Croomii, Hort.—Cf. Gard. Chron. (1884) ii. 492 = A. Unedo.
densiflora, Benth. Pl. Hartw. 15 = glandulosa.
densiflora, *H. B. & K Nov. Gen. et Sp* iii 280. t 260 —Mexic.
discolor, Hook. Ic. Pl. t 721 = Arctostaphylos arguta
empetrifolia, Pers. Syn. i. 468 = Pernettya empetrifolia.
ferruginea, *Linn. f. Suppl.* 238.—Reg. Colomb.
filiformis, Lam. Encyc. i. 325 = Chiococca serpyllifolia.
floribunda, *Mart. & Gal. in Bull. Acad. Brux.* ix (1842) 584.—Mexic.
furiens, *Hook. Bot. Beech. Voy.* 33.—Chili.
glaberrima, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Malaya
glandulosa, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 583 —Mexic.
herpetica, Coleb. ex Roxb. Hort. Beng. [90]; Fl. Ind. ii. 412 = Pieris ovalifolia.
hybrida, Ker-Gawl. in Bot. Reg. t. 619 = andrachnoides.
integrifolia, Lam. Encyc. i. 327 = A. Andrachne.
integrifolia, Sieber, ex Klotzsch, in Linnæa, xxiv. (1851) 731.—Sieberi.
integrifolia, Sims, Bot. Mag. t. 2819 = A. Unedo.
intermedia, Heldr. ex Nym. Consp. 190 = A. Unedo.
lanceolata, Lam. Encyc. i. 327 = Prunus caroliniana.
laurifolia, Buch.-Ham. ex D. Don, Prod Fl. Nep. 151 = Gaultheria fragrantissima.
laurifolia, Linn. f. Suppl. 238. (Cf. A. Gray, Syn. Fl. N. Am. ii. 1. 396) = A. Unedo.
laurifolia, Lindl. Bot. Reg. (1839) t. 67 = xalapensis.
laurina, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 585.—Mexic.
leucocarpa, Poepp. & Kunze, ex DC. Prod. vii. 586 = Pernettya leucocarpa.
longifolia, Andr. Bot. Rep. x. 664 = canariensis.
lucida, Hort. ex Steud. Nom. ed. II. i. 119 = A. Andrachne.
macrophylla, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 584.—Mexic.
Madrono, Humb. Nouv. Espagne, ii. 454, ex H. B. & K. Nov. Gen. et Sp. iii. 282 = Arctostaphylos pungens.
Menziesii, *Pursh, Fl. Bor. Am.* i. 282.—Am. bor.; Mexic.
microphylla, Forst. f. in Comm. Gotting. ix. (1789) 82 = Pernettya microphylla.
microphylla, Hort.—Cf. Gard. Chron. (1884) ii. 49 = A. Unedo.
Milleri, *Sweet, Hort. Brit.* ed. II. 381.—Orlena.
mollis, *H. B. & K. Nov. Gen. et Sp.* iii. 281. t. 260.— Mexic.
mucronata, Linn. f. Suppl. 239 = Pernettya mucronata.
myrtifolia, Willd. ex Steud. Nom. ed. II. i. 120, 119 = Arctostaphylos pungens.
myrtilloides, Humb. ex Steud. Nom. ed. II. i. 119 = Vaccinium stamineum.
myrtilloides, Poepp. ex DC. Prod. vii. 598 = Pernettya myrtilloides.
nuda, Steud. Nom. ed. II. i. 119 = Arctostaphylos pungens.
obtusifolia, *Rafin. Fl. Ludov.* 55.—Am. bor.
officinalis, Boiss. Fl. Orient. iii. 967 = Arctostaphylos Uva-ursi.
ovata, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 583.—Mexic.
paniculata, *Mart. & Gal. l. c.* 585.—Mexic.

ARBUTUS:—
petiolaris, *H. B. & K. Nov. Gen. et Sp.* iii. 281.—Mexic.
phyllireæfolia, Pers. Syn. i. 483 = Pernettya phyllireæfolia.
pilosa, R. Grah. in Bot. Mag. t. 3177 = Pernettya pilosa.
procera, Dougl. ex Lindl. Bot. Reg. t. 1758 = Menziesii.
procera, Soland. ex DC. Prod. vii. 561 = canariensis.
procumbens, *Kluk, ex Bess. Fl. Galic.* ii. 355.—Galicia.
procumbens, Salisb. Prod. 289 = Arctostaphylos Uva-ursi.
prostrata, Cav. Ic. vi. 43. t. 562 = Pernettya parvifolia.
prunifolia, *Klotzsch, in Linnæa*, xxiv. (1851) 73.—Mexic.
pumila, Forst. f. in Comm. Gottng. ix. (1789) 32 = Pernettya empetrifolia.
punctata, *Hook. & Arn. Bot. Beech. Voy.* 33.—Calif.
pungens, Hook. & Arn. l. c. 144 = Arctostaphylos Hookeri.
rigida, Banks, ex Hook. f. Fl. Antarct. ii. 326 = Pernettya mucronata.
rigida, Bert. ex Colla, in Mem. Acc. Sc. Torin. xxxviii. (1835) 120 = Pernettya rigida.
rigida, Willd. ex Steud. Nom. ed. II. i. 120, 119 = Arctostaphylos pungens.
rubescens, Bertol. Fl. Gustimal. 20. t. 7 = Arctostaphylos rubescens.
salicifolia, *Cels. ex Hoffmgg. Vers. Pft. Nachtr.* i. (1824) 204.—Oriens.
serpyllifolia, Lam. Encyc. i. 225 = Gaultheria antipoda.
serratifolia, Lodd. Bot. Cab. t. 580 = A. Andrachne.
serratifolia, Salisb. Prod. 288 = A. Unedo.
sibirica, *Forbes, Hort. Woburn.* 95, nomen.—Sibir.
Sieberi, Klotzsch, in Linnæa, xxiv. (1851) 71 = A. Andrachne.
spinulosa, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 532.—Mexic.
texana, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 460 = Menziesii.
thymifolia, Alt. Hort. Kew. ii. 72 = Chiogenes serpyllifolia.
tomentosa, Pursh, Fl. Am. Sept. i. 282 = Arctostaphylos pungens.
turbinata, Pers. ex Reichb. Fl. Germ. Excurs. i. 416 = A. Unedo.
Unedo, *Linn. Sp. Pl.* 505.—Reg. Mediterr. ; Hibern.
Uva-ursi, Linn. l. c. = Arctostaphylos Uva-ursi.
varians, *Benth. Pl. Hartw.* 77.—Mexic.
vernalis, Poepp. & Endl. Nov. Gen. i. 26. t. 42 = Gaultheria vernalis.
villosa, Willd. ex Klotzsch, in Linnæa, xxiv. (1851) 73 = glandulosa.
xalapensis, Andr. ex Hook. lc. Pl. t. 27 = Arctostaphylos oaxacana.
xalapensis, *H. B. & K. Nov. Gen. et Sp.* iii. 279.—Mexic.
xanthocarpa, *Wangen. Beitr.* 80.—Am. bor.

ARCANGELISIA, Becc. Malesia, i. (1877) 145.
MENISPERMACEÆ.
inclyta, *Becc. l. c.* 147.—Java ; Celebes.
lemniscata, *Becc. l. c.*—N. Guin.

ARCAULA, Rafin. Alsog. Am. 80 (1838) = **Quarosa**, Tourn. (Cupul.).
spicata, Rafin. l. c. = Q. spicata.

ARCEUTHOBIUM, Bieb. Fl. Taur. Cauc. Suppl. 629 (1819). *LORANTHACEÆ*, Benth. & Hook. f. iii. 218.
RAZOUMOWSKIA, Hoffm. ex Bieb. l. c. (1819).
abietinum, Engelm. in Proc. Am. Acad. vili. (1873) 401 = occidentale.
americanum, Nutt. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 214 = A. Oxycedri.
campylopodum, *A. Gray, l. c.*—Am. bor.
capense, J. S. Presl, Rostl. ii. 38 = Viscum capense.
cryptopodum, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 214 = vaginatum.
divaricatum, *Engelm. in Wheeler, Rep.* vi. 253.—Am. bor.
Douglasii, *Engelm. l. c.*—Am. bor.

ARCEUTHOBIUM :—
gracile, *Engelm. in A. Gray, Pl. Fendl.* 59.—Am. bor.
minutum, *Engelm. in Bull. Torrey Bot. Club*, ii. (1871) 43, 47.—Am. bor.
occidentale, *Engelm. ex S. Wats. in Bot. Calif.* ii. 107.—Calif.
opuntioides, J. Presl, Rostl. ii. 28 = Dendrophthora opuntioides.
Oxycedri, *Bieb. Fl. Taur. Cauc. Suppl.* 629.—Europ. ; Am. bor.
robustum, *Engelm. in A. Gray, Pl. Fendl.* 59.—Am. bor.
vaginatum, *J. Presl, Rostl.* ii. 28.—Am. bor.

ARCEUTHOBIUM, Griseb. Fl. Brit. W. Ind. 314 (1864) = **Dendrophthora**, Eichl. (Loranth.).
Bonaniæ, Griseb. Cat. Pl. Cub. 121 = D. Bonaniæ.
cupressoides, Griseb. Fl. Brit. W. Ind. 315 = D. cupressoides.
domingense, Griseb. Cat. Pl. Cub. 121 = D. dominguensis.
Equisetum, Griseb. in Mem. Am Acad. N. S. viii. (1861) 192 = D. Equisetum.
glaucum, Griseb. Fl Brit W. Ind. 315 = D. glaucum.
gracile, Griseb l. c. = D. gracilis.
Mancinellæ, Wright, ex Griseb. Cat. Pl. Cub. 121 = D. Mancinellæ.

ARCEUTHOS, Ant & Kotschy, in Oestr. Bot. Wochenbl. (1854) 249 = **Juniperus**, Linn. (Conif.).
drupacea, Ant. & Kotschy, l. c. = J. drupacea.

ARCHANGELICA, Hoffm. Gen. Umb. ed. I. 161 (1814) ; ed. II. 166 (1816). *UMBELLIFERÆ*, Benth & Hook. f. i. 917.
COELOPLEURUM, Ledeb. Fl. Ross. ii. 861 (1844).
CZERNIAJEVIA, Turcz. in Bull. Soc. Nat. Mosc. (1858) 93.
atropurpurea, *Hoffm. Gen. Umb.* 162.—Am. bor.
brevicaulis, *Reichb. f. in Journ. Bot.* xiv. (1876) 46.—As. temp.
communis, *C. A. Mey. ex Rupr. in Beitr. Pft. Russ. Reich.* xi. 13.—Kamtschatka.
decurrens, Ledeb. Fl. Alt. i. 316 = officinalis.
dentata, *Chapm. in Torr. & Gray, Fl. N. Am.* i. 622.—Florida.
discocarpa, *Fries, in Bot. Notiser*, (1857) 50.—Europ.
Gmelini, *DC. Prod.* iv. 170.—Am. bor. ; As. bor. occ.
hirsuta, *Torr & Gray, Fl. N. Am.* i. 622.—Am. bor.
inæqualis, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* i. 138.—Japon.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 62.—Japon.
littoralis, *Agardh, ex DC. Prod.* iv. 170.—Europ. arct.
major, Lag. Gen. et Sp. Nov. 18 = littoralis !
norvegica, Rupr. in Beitr. Pft. Russ. Reich. ii. 37 = officinalis.
officinalis, *Hoffm. Gen. Umb.* 162.—Europ. ; Sibir. ; Reg. Himal.
officinalis, Hook. Fl. Bor. Am. i. 267 = Gmelini.
peregrina, Nutt. ex Torr. & Gray, Fl. N. Am. i. 622 = Gmelini.
Royleï, Lindl. in Royle, Illustr. Bot. Himal. 232 = officinalis.
sativa, Bess. Enum. 13 = officinalis.
slavica, Reuss, Kvet. Slov. (1853) 184 = officinalis.
spuria, Wahlenb. Fl. Casp. 84, in obs. = littoralis.

ARCHEMORA, DC. Prod. iv. 188 (1830) = **Peucedanum**, Linn. (Umbell.).
ambigua, DC. l. c. = P. ternatum.
denticulata, DC. l. c. = P. ternatum.
Fendleri, A. Gray, Pl. Fendl. 56.—Texas.
rigida, DC. Mém. Ombell. 52 ; Prod. iv. 188 = P. ternata.
ternata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 631 = P. ternatum.
tricuspidata, DC. Prod. iv. 188 = P. ternatum.

ARCHERIA, Hook. f. Fl. Tasm. i. 269. t. 80, 81. *EPACRIDEÆ*, Benth. & Hook. f. ii. 616.
ericoæpa, *Hook. f. l. c.* 269. t. 80 B.—Tasmania.
hirtella, *Hook. f. l. c.* t. 81.—Tasmania.
minor, Hook. f. l. c. 264 = serpyllifolia.

ARCHERIA :—
racemosa, *Hook. f. Handb. New Zeal. Fl.* 180.—N. Zel.
serpyllifolia, *Hook. f. Fl. Tasm. i.* 268. t. 80 B.—Tasmania.
Traversii, *Hook. f. Handb. New Zeal. Fl.* 180.—N. Zel.

ARCHIDENDRON, F. Muell. Fragm. v. 59 (1865).
LEGUMINOSÆ, Benth. & Hook. f. i. 1004.
Lucyi, *F. Muell. l. c.* vi. 201.—Austral.
Vaillantii, *F. Muell. l. c.* v. 60.—Austral.

ARCHIMEDEA, Leandr. Sacram. ex A. St. Hil. in Ann. Sc. Nat. Sér. II. vii. (1837) 33 = **Lophophytum**, Schott & Endl. (Balanoph.).
pyramidalis, Leandr. Sacram. l. c. = L. mirabile.

ARCHITAEA, Mart. Nov. Gen. et Sp. i. 116. t. 78 (1824) = **Archytaea**, Mart. (Ternstr.).

ARCHONTOPHOENIX, H. Wendl. & Drude, in Linnæa, xxxix. (1875) 182. *PALMAE*, Benth. & Hook. f. iii. 889.
Alexandræ, *H. Wendl. & Drude, l. c.* 212.—Austral.
Cunninghamii, *H. Wendl. & Drude, l. c.* 214.—Austral.
Veitchii, *H. Wendl. & Drude, l. c.*—Hab. ?

ARCHYTAEA, Mart. Nov. Gen. et Sp. i. 117 (1824). *TERNSTROEMIACEÆ*, Benth. & Hook. f. i. 188.
PLOIARIUM, Korth. Verh. Nat. Gesch. Bot. 185. t 25 (1842).
multiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 203.—Guiana.
sessilis, *Scheff. in Tydschr. Nederl. Ind.* xxxii. (1871) 406.—Malaya.
triflora, *Mart. Nov. Gen. et Sp.* i. 117.—Bras.
Vahlii, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 163.—Malaya.

ARCTAGROSTIS, Griseb. in Ledeb. Fl. Ross. iv. 434 (1858). *GRAMINEÆ*, Benth. & Hook. f. iii. 1150.
latifolia, *Griseb. l. c.*—Reg. Arct.

ARCTIO (et ARCTION), Lam. Encyc. i. 235 (1788) = **Berardia**, Vill. (Compos.).
lanuginosa, Lam. l. c. = B. subacaulis.

ARCTIODRACON, A. Gray, in Mem. Am. Acad. Ser. II. vi. (1858-9) 408 = **Lysichitum**, Schott (Aroid.).
camtschaticum, A. Gray, l. c. 409 = L. camtschatense.
japonicum, A. Gray, l. c. = L. camtschatense.

ARCTIUM, Lam. Fl. Fr. ii. 70 (1778) = **Berardia**, Vill. (Compos.).
lanuginosum, Lam. l. c. = B. subacaulis.

ARCTIUM, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 466.
LAPPA, Juss. Gen. 173 (1789).
adhærens, Gilib. Fl. Lituan. i. 182 = A. Lappa.
Bardana, Willd. Sp. Pl. iii. 1682 = A. Lappa.
Carduelis, Linn. Mant. i. 108 = Carduus arctioides.
conglomeratum, Schur, ex Nym. Syll. Suppl. 4 = minus.
eumenus, Syme, Engl. Bot. ed. III. v. 26 = minus.
grandiflorum, Desf. Cat. Hort. Par. ed. III. 154 = majus.
intermedium, Hab. in Ann. & Mag. Nat. Hist. Ser. II. xvii. (1856) 374 = majus.
intermedium, Lange, Fl. Dan. t. 2668 = nemorosum.
Lappa, Kalm, Fl. Fennic. n. 449 ; Sm. Engl. Bot. t. 1238 = minus.
Lappa, Willd. Sp. Pl. iii. 1681 = majus.
Lappa, *Linn. Sp. Pl.* 816 (= majus, minus, &c.).—Europ.
majus, *Bernh. Syst. Verz. Erf.* 154.—Europ.
minus, *Bernh. l. c.*—Europ.
montanum, Schweig. ex Steud. Nom. ed. II. i. 120 = minus.
nemorosum, *Lejeune, in Reichb. Ic. Fl. Germ.* xv. 81.—Europ.
Personata, Linn. Sp. Pl. 816 = Carduus Personata.

ARCTIUM :—
Personata, Pall. Reise, iii. 439 (plants ignota).—Davuria.
pubens, Bab. in Ann. & Mag. Nat. Hist. Ser. II. xvii. (1856) 376 = nemorosum.
ruderale, Salisb. Prod. 154 = A. Lappa.
tomentosum, Mill. Gard. Dict. ed. VIII. n. 3; Bernh. Syst. Verz. Erf. 154 = majus.

ARCTOCALYX, Fenzl, in Otto & Dietr. Allg. Gartenz. xvi. 305 (1848) = **Solenophora**, Benth. (Gesner.).
Endlicherianus, C. Heller, ex Fenzl, l. c. 307 = S. Endlicheriana.
insignis, Fenzl, l. c. 306 = S. insignis.

ARCTOCARPUS, Blanco, Fl. Filip. ed. I. 666 (1837) = **Artocarpus**, Forst. (Urtic.).

ARCTOGERON, DC. Prod. v. 260 (1836) = **Aster**, Linn. (Compos.).
graminifolium, DC. l. c. 261.—Sibir.

ARCTOMECON, Torr. & Frém. in Frém. Rep. 312. t. 2 (1845). *PAPAVERACEÆ*, Benth. & Hook. f. i. 52.
californicum, *Torr. & Frém. l. c.*—Calif.

ARCTOPHILA, Rupr. in Beitr. Pfl. Russ. Reich. ii. 62 (1845) = **Poa**, Linn. (Gram.).
fulva, Nym. Consp. 830 = P. fulva.
Laestadii, Rupr. in Beitr. Pfl. Russ. Reich. ii. 62 = P. Laestadii.
pendulina, Nym. Consp. 830 = P. Laestadii.

ARCTOPUS, Linn. Coroll. Gen. 21 (1737); Hort. Cliff. 495 (1737). *UMBELLIFERÆ*, Benth. & Hook. f. i. 879.
ÆPEADUS, Adans. Fam. ii. 102, 509 (1763).
Dregei, *Sond. in Harv. & Sond. Fl. Cap.* ii. 565.—Afr. austr.
echinatus, *Linn. Sp. Pl.* 1058.—Afr. austr.
monacanthus, *Cormich. ex Harv. & Sond. Fl. Cap.* ii. 565.—Afr. austr.

ARCTOSTAPHYLOS, Adans. Fam. ii. 165 (1763). *ERICACEÆ*, Benth. & Hook. f. ii. 581.
COMAROSTAPHYLIS, Zucc. in Abh. Akad. Muench. ii. (1837) 321.
DAPHNIDOSTAPHYLIS, Klotzsch, in Linnæa, xxiv. (1851) 78.
MAIRANIA, Neck. Elem. i. 219 (1790).
UVA-URSI, Tourn. ex Moench, Meth. 470 (1794).
XEROBOTRYS, Nutt. in Trans. Am. Phil. Ser. II. viii (1842) 267.
XYLOCOCCUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1842) 256.
acuta, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1842) 256 = Hookeri?
alpina, Payot.—Cf. Bull. Soc. Bot. Fr. xxix. (1882) Bibl. 43, lapsu = *angustifolia*.
alpina, *Spreng. Syst.* ii. 287.—Am. bor. et arct.
Andersonii, *A. Gray, in Proc. Am. Acad.* ix. (1876) 88.—Calif.
angustifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 278. —Mexic.
angustifolia, Payot, Fl. du Mont Blanc, Phan. 174 = A. Uva-ursi.
arbutoides, *Hemsl. Biol. Centr. Am. Bot.* ii. 278.—Guatemala.
arguta, *Zucc. in Linnæa*, xii. (1838) *Litt.* 86; *DC. Prod.* vii. 585.—Mexic.
attenuata, *Hemsl. Biol. Centr. Am. Bot.* ii. 278.—Mexic.
bicolor, *A. Gray, in Proc. Am. Acad.* vii. (1868) 366. —Calif.
Clevelandi, A. Gray, l. c. xii. (1877) 61 = bicolor.
cordifolia, Lindl. Bot. Reg. sub t. 1791 = tomentosa.
discolor, DC. Prod. vii. 585 = arguta.
diversifolia, *Parry, ex A. Gray, Syn. Fl. N. Am.* ii. 1. 397.—Calif.
glauca, *Lindl. Bot. Reg.* sub t. 1791.—Calif.
glauca, S. Wats. in Bot. King. Exped. 210 = pungens.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* iii. 278.— Mexic.
Hartwegiana, *Hemsl. Biol. Centr. Am. Bot.* ii. 278. —Mexic.
Hookeri, *G. Don, Gen. Syst.* iii. 896.—Calif.
latifolia, Walp. Rep. ii. 726 = arg.

ARCTOSTAPHYLOS :—
latifolia, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 536.—Mexic.
longifolia, *Benth. Pl. Hartw.* 44.—Mexic.
mucronata, *Hemsl. Biol. Centr. Am. Bot.* ii. 279.— Mexic.
mucronifera, *DC. Prod.* vii. 585.—Mexic.
nevadensis, *A. Gray, Syn. Fl. N. Am.* ii. 1. 37.—Calif.
nitida, Benth. Pl. Hartw. 66 = arguta.
pumilaaris, *A. Gray, in Proc. Am. Acad.* vii. (1868) 366.—Calif.
oaxacana, Benth. Pl. Hartw. 45 = Hartwegiana.
OREANDRA, *DC. Prod.* vii. 585.—Mexic.
officinalis, Wimm. & Grab. Fl. Sil. i. 391 = A. Uva-ursi.
oppositifolia, *Parry, in Proc. Davenport Acad. Nat. Sc.* iv. (1884) 37.—Calif.
polifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 277. t. 258. —Mexic.
polifolia, Torr. Bot. Mex. Bound. 108 = oppositifolia.
procera, Lindl. ex Pritz. Ind. Ic. 90, sphalm. = tomentosa.
procumbens, Patze, Mey. & Elk. Fl. Preuss. 188 = A. Uva-ursi.
pumila, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. viii. (1843) 266.—Calif.
pungens, A. Gray, in Bot. Calif. i. 458, partim = Hookeri.
pungens, *H. B. & K. Nov. Gen. et Sp.* iii. 278. t. 259. —Calif.; Mexic.
rubescens, *Hemsl. Biol. Centr. Am. Bot.* ii. 279.— Guatemala.
salicifolia, Parry, in Proc. Davenport Acad. iv. (1884) 37 = oppositifolia.
spinulosa, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 537 = arguta.
tomentosa, *Lindl. Bot. Reg.* t. 1791.—Calif.
Uva-ursi, *Spreng. Syst.* ii. 287.—Reg. bor. et arct.
Vvatchii, Kellogg, in Proc. Calif. Acad. ii. (1863) 19 = bicolor.

ARCTOTHECA, Wendl. Beobacht. 41; et Hort. Herrenh. 8. t. 6 (1796). *COMPOSITÆ*, Benth. & Hook. f. ii. 458.
grandiflora, Drège, ex Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 93 = Arctotis Dregei.
grandiflora, Schrad. in Ind. Sem. Hort. Goett. (1832) 1.—Capens.
hirta, *Schrad. ex Steud. Nom.* ed. II. i. 120.—Afr. austr.
repens, *Wendl. Beobacht.* 41; *et Hort. Herrenh.* 8. t. 6.—Afr. austr.

ARCTOTIS, Linn. Gen. ed. I. 261 (1737). *COMPOSITÆ*, Benth. & Hook. f. ii. 458.
ANEMOSPERMOS, Adans. Fam. ii. 127 (1763).
LYCOTIS, Hoffmgg. Verz. Pfl. Nachtr. ii. 99 (1826).
ODONTOPTERA, Cass. in Dict. Sc. Nat. xxxv. 396 (1825).
STŒGNOTUS, Cass. l. c. (1825).
abrotanifolia, Steud. Nom. ed. II. i. 120 = Ursinia abrotanifolia.
acaulis, E. Mey. ex DC. Prod. vi. 466 = canescens.
acaulis, Jacq. Hort. Schoenb. ii. 16. t. 158 = adpresa.
acaulis, *Linn. Sp. Pl.* ed. II. 1306.—Afr. austr.
adpressa, *DC. Prod.* vi. 468.—Afr. austr.
adpressa, Eckl. ex Harv. & Sond. Fl. Cap. iii. 465 = Haplocarpha lyrata.
æquis, *Jacq. f. Eclog.* i. t. 59.—Afr. austr.
amœna, *Jacq. Fragm.* 75.—Afr. austr.
amplexicaulis, *Less. Syn. Comp.* 25.—Afr. austr.
angustifolia, Jacq. Hort. Schoenb. ii. 21 = aspera.
angustifolia, *Linn. Sp. Pl.* 928.—Afr. austr.
anthemoides, Linn. Syst. ed. X. 1234 = Ursinia anthemoides.
anthemoides, Thunb. ex Harv. & Sond. Fl. Cap. iii. 453 = Ursinia chrysanthemoides.
arborescens, *Jacq. Hort. Schoenb.* ii. 22.—Afr. austr.
arborescens, Willd. ex Harv. & Sond. Fl. Cap. iii. 453 = aspera.
argentea, Alt. Hort. Kew. iii. 278 = linearis.
argentea, *Thunb. Diss. Arct.* 4 et 7.—Afr. austr.
aspera, *Linn. Sp. Pl.* 929.—Afr. austr.
asplenifolia, Burm. f. Fl. Cap. Prod. 28 = Gazania pinnata.
aurea, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
aurosa, Kze-Cwzl. in Bot. Reg. t. t. 35 = aspera.
auriculata, Jacq. Hort. Fl. Schoenb. 2. t. 166 = aspera.
bellidifolia, *Berg. Descr. Pl. Cap.* 216.—Afr. austr.

ARCTOTIS :—
bicolor, Willd. Enum. Hort. Berol. 936 = aspera.
breviscapa, Thunb. Diss. Arct. 13 = leptorhiza.
Calendula, Linn. Sp. Pl. 922 = Cryptostemma calendulaceum.
calendulacea, Linn. Syst. ed. XII. 578 = Cryptostemma calendulaceum.
calendulacea, Thunb. ex Harv. & Sond. Fl. Cap. iii. 468 = Venidium hirsutum.
campanulata, *DC. Prod.* vi. 485.—Afr. austr.
candida, *Thunb. Diss. Arct.* 14.—Afr. austr.
canescens, *DC. Prod.* vi. 466.—Afr. austr.
cardiifolia, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
carolinana, *Wall. Fl. Carol.* 216.—Carolina.
caulescens, Thunb. Diss. Arct. 11 = aspera.
cernua, Thunb. l. c. 17 = Ursinia anthemoides.
Cineraria, *Jacq. Hort. Schoenb.* ii. 24.—Afr. austr.
coronopifolia, Salisb. Prod. 210 = Ursinia crithmifolia.
corrugata, DC. Prod. vi. 496 = Cryptostemma calendulaceum.
crithmifolia, Harv. in Harv. & Sond. Fl. Cap. iii. 444 = Ursinia crithmifolia.
crithmoides, Berg. Descr. Pl. Cap. 226 = Ursinia crithmifolia.
crithmoides, Jacq. Hort. Schoenb. ii. 15 = Ursinia pilifera.
cuneata, DC. Prod. vi. 489 = stœchadifolia.
cuprea, Hort. ex Stend. Nom. ed. II. i. 120 = sanea.
cuprea, Jacq. Hort. Schoenb. ii. 25 = aspera.
decumbens, Jacq. l. c. iii. 68. t. 331 = angustifolia.
decumbens, Thunb. Diss. Arct. 5 = stœchadifolia.
decurrens, Jacq. Hort. Schoenb. ii. 20. t. 165 = angustifolia.
dentata, Linn. Pl. Afr. Rar. 25; Amœn. Acad. vi. 107 = Ursinia dentata.
dentata, Thunb. Diss. Arct. 17 (= *Sphenogyne discolor*). —Afr. austr.
denudata, Thunb. l. c. 14 = laevis.
diffusa, *Thunb. l. c.* 9.—Afr. austr.
Dregei, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 11. 93.—Afr. austr.
echinata, DC. Prod. vi. 486 = Landtia hirsuta.
elatior, *Jacq. Hort. Schoenb.* ii. 23.—Afr. austr.
elatior, Willd. ex DC. Prod. v. 488 = amplexicaulis.
elegans, Thunb. Diss. Arct. 14 = Chrysanthemum indicum.
elongata, *Thunb. l. c.* 9.—Afr. austr.
fastuosa, *Jacq. Hort. Schoenb.* ii. 20. t. 166.—Afr. austr.
flaccida, *Jacq. l. c.* 19.—Afr. austr.
flamtzea, *Jacq. f. Fragm.* 41.—Afr. austr.
fœnticulum, Jacq. Hort. Schoenb. ii. 15 = Ursinia fœniculacea.
formosa, Thunb. Diss. Arct. 11 = aspera.
gigantea, *A. Rich. Tent. Fl. Abyss.* i. 449.—Abyss.
glabrata, Jacq. Hort. Schoenb. ii. 25 = laevis.
glandulosa, *Thunb. Diss. Arct.* 8.—Afr. austr.
Gnaphalyla, Jacq. Hort. Schoenb. ii. 23 = candida.
glutinosa, Sims, Bot. Mag. t. 1848 = Dimorphotheca cuneata.
grandiflora, Ait. Hort. Kew. iii. 272 = aculis.
grandiflora, Jacq. Hort. Schoenb. iii. t. 278 = laevis.
grandis, Thunb. Diss. Arct. 8 = stœchadifolia.
hispidella, Juss. ex DC. Prod. vii. 258 = Hispidella hispanica.
humilis, Salisb. Prod. 210 = aculis.
hypochondriaca, Willd. Sp. Pl. iii. 2348 = Cryptostemma calendulaceum.
incisa, Thunb. Diss. Arct. 10 = aspera.
interrupta, Thunb. l. c. 12 = Arctothoca repens.
Jacobæa, Sch. Bip. in Flora, xxvii. (1844) 772 = aspera.
Kraussii, Sch. Bip. l. c. 771 = angustifolia.
lacinlata, *Lam. Encyc.* i. 387.—Afr. austr.
laevis, *Thunb. Diss. Arct.* 10.—Afr. austr.
lanata, Thunb. l. c. 11 = Haplocarpha Thunbergii.
lanosa, Thunb. ex DC. Prod. vi. 508 = Didelta tomentosum.
lanuginosa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 456. —Afr. austr.
Leichtlinlana, *Lynch, in Gard. Chron.* (1885) ii. 80. —Afr. austr.
leiocarpa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 451.— Afr. austr.
leptorhiza, *DC. Prod.* vi. 486.—Afr. austr.
lanuginosa, *Hoffmgg. Verz. Pfl.* 40.—Hab.?
leucanthemoides, *Jacq. Hort. Schoenb.* ii. 19. t. 154.— Afr. austr.

ARCTOTIS:—
linearis, *Thunb. Diss. Arct.* 7.—Afr. austr.
lyrata, Willd. Sp. Pl. iii. 2356 = aspera.
maculata, Jacq. Hort. Schoenb. iii. 67 = aspera.
Masoniana, Spreng. Syst. iv. Car. Post. 304 = laevis.
melanocyclo, Willd. Enum. Hort. Berol. 937 = aspera.
microcephala, Thunb. ex DC. Prod. vi. 493 = Venidium decurrens.
muricata, Thunb. Diss. Arct. 10 = bellidifolia.
nodosa, Thunb. l. c. 15 = Chrysanthemum nodosum.
nudicaulis, Thunb. l. c. 16 (= *Sphenogyne nudicaulis*).—Afr. austr.
odorata, Steud. Nom. ed. II. i. 121 = Ursinia odorata.
oocephala, *DC. Prod.* vi. 486.—Afr. austr.
palacea, Linn. Pl. Afr. Rar. 25; Amoen. Acad. vi. 107 = Ursinia crithmifolia.
palacea, Thunb. Diss. Arct. 16 = Ursinia palacea.
paniculata, Jacq. Hort. Schoenb. iii. 68 = bellidifolia.
pannosa, *Jacq. l. c.* iv. 21.—Afr. austr.
paradoxa, Alt. Hort. Kew. iii. 378 = Ursinia anthemoides.
paradoxa, Linn. Cent. ii. 32; Amoen. Acad. iv. 330 = Ursinia paradoxa.
paradoxa, Thunb. Diss. Arct. 17 = Ursinia chrysanthemoides.
pectinata, Thunb. l. c. 14 = Ursinia dentata.
petiolata, *Thunb. l. c.* 10.—Afr. austr.
pilifera, Thunb. l. c. 18 = Ursinia pilifera.
piligera, Berg. Desc. Pl. Cap. 325 = Ursinia dentata.
pinnata, Thunb. Diss. Arct. 16 = Ursinia filiformis.
pinnatifida, *Thunb. l. c.* 7.—Afr. austr.
plantaginea, Linn. Sp. Pl. ed. II. 1306 = Venidium semipapposum.
plantaginifolia, Salisb. Prod. 210 = Venidium semipapposum.
populifolia, Berg. Desc. Pl. Cap. 323 = Cryptostemma niveum.
prostrata, Salisb. Prod. 210 = Arctotheca repens.
punctata, Thunb. Diss. Arct. 15 (= *Sphenogyne punctata*).—Afr. austr.
purpurea, *Willd. Enum. ex Steud. Nom.* ed. II. i. 121.—Afr. austr.
pusilla, *DC. Prod.* vi. 489.—Afr. austr.
pygmaea, *A. Rich. Tent. Fl. Abyss.* i. 445.—Abyss.
radicans, Linn. Mant. ii. 479, 480 = angustifolia.
repens, Jacq. Hort. Schoenb. iii. 31 = Arctotheca repens.
reptans, *Jacq. l. c.* 89. t. 382.—Afr. austr.
revoluta, *Jacq. l. c.* iii. 24.—Afr. austr.
rigida, Burm. f. Fl. Cap. Prod. 28 = Gazania Pavonia.
rosea, Jacq. Hort. Schoenb. ii. 18 = stoechadifolia.
runcinata, *Hoffegg. Verz. Pfl.* 40.—Hab. ?
scabra, Thunb. Diss. Arct. 9 = Venidium semipapposum.
scapigera, Thunb. l. c. 12 = acaulis.
scariosa, Alt. Hort. Kew. iii. 374 = Ursinia scariosa.
sericea, Thunb. Diss. Arct. 18 (= *Sphenogyne sericea*).—Afr. austr.
serrata, Thunb. l. c. 8 = Ursinia serrata.
spathulata, Thunb. ex Spreng. Syst. iii. 626 = Venidium perfoliatum.
speciosa, Jacq. Hort. Schoenb. ii. 18 = acaulis.
speciosa, Salisb. Prod. 209 = Cryptostemma calendulaceum.
spinulosa, Jacq. Hort. Schoenb. ii. 21 = fastuosa.
squarrosa, Jacq. l. c. 25 = laevis.
staticifolia, Poir. Encyc. Suppl. i. 488 = Gazania subulata.
stoechadifolia, *Berg. Desc. Pl. Cap.* 324.—Afr. austr.
subfusca, Gaertn. Fruct. ii. 439 = Cryptostemma hypochondriacum.
sulphurea, Salisb. Prod. 210 = aspera.
superba, Linn. Cent. ii. 32; Sp. Pl. ed. II. 1305 = Cryptostemma calendulaceum.
tenuifolia, Linn. Mant. 288 = Ursinia tenuifolia.
tenuifolia, Poir. Encyc. Suppl. i. 489 = Dimorphotheca graminifolia.
tenuifolia, Salisb. Prod. 210 = Ursinia scariosa.
tomentosa, Thunb. ex DC. Prod. vi. 493 = Venidium semipapposum.
tricolor, Jacq. Hort. Schoenb. ii. 16 = acaulis.
tricolor, Willd. ex DC. Prod. vi. 486 = elongata.
trifida, Thunb. Diss. Arct. 13 (= *Sphenogyne trifida*).—Afr. austr.
trifurcata, Burm. f. Fl. Cap. Prod. 28 = Ursinia crithmifolia.
tristis, Linn. Sp. Pl. 923 = Cryptostemma calendulaceum.
undulata, Gaertn. Fruct. ii. 438 = lachnaea.
undulata, Jacq. Hort. Schoenb. ii. 16 = acaulis.

ARCTOTIS:—
undulata, Thunb. Diss. Arct. 18 = aspera.
venidioides, *DC. Prod.* vi. 489.—Afr. austr.
verbascifolia, *Harv. in Harv. & Sond. Fl. Cap.* iii. 452.—Afr. austr.
virgata, *Jacq. Hort. Schoenb.* iii. 33.—Afr. austr.
vitellina, *Hoffegg. Verz. Pfl.* 40.—Hab. ?

ARCYNOSPERMUM, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 191. *MALVACEAE* (gen. dub.), Benth. & Hook. f. Gen. i. 199.
nodiflorum, *Turcz. l. c.*—Mexic.

ARCYPHYLLUM, Ell. in Journ. Acad. Philad. i. (1818) 371 = **Rhynchosia**, Lour. (Legum.).
caribaeum, Ell. l. c. 372 = R. caribaea.
capitatum, Benth. in Linnaea, xxii. (1849) 525 = R. Clausseni.
diforme, Ell. in Journ. Acad. Phil. i. (1818) 372 = R. tomentosa.
erectum, Ell. l. c. = R. tomentosa.
leucophyllum, Benth. in Linnaea, xxii. (1849) 525 = R. leucophylla.
reticulatum, Ell. in Journ. Acad. Philad. i. (1818) 372 = R. reticulata.
rhombifolium, Ell. l. c. = R. minima.
simplicifolium, Ell. l. c. 371 = R. tomentosa.

ARCYTHOPHYLLUM, Willd. ex Schlecht. in Linnaea, xxviii. (1856) 491 = secq.

ARCYTOPHYLLUM, [Willd.] Roem. & Schult. Mant. iii. 5 (1827) = **Mallostoma**, Karst (Rubiac.).
blacrioides, [Willd. ex] Roem. & Schult. l. c. = M. cataeoasata.
nitidum, Schlecht. in Linnaea, xxviii. (1856) 492 = M. nitida.
setosum, Schlecht. l. c. = M. setosa.

ARDERNIA, Salisb. Gen. Pl. Fragm. 35 (1866) = **Ornithogalum**, Linn. (Liliac.).
fuscata, Salisb. l. c. = O. fuscatum.

ARDINGHELIA, Comm. ex A. Juss. Tent. Euph. 21 (1824) = **Phyllanthus**, Linn. (Euphorb.).

ARDISIA, Gaertn. Fruct. ii. 78. t. 94 (1791) = Cyathodes, Labill. (Epacrid.).
acerosa, Gaertn. l. c. = C. acerosa.
lanceolata, Gaertn. f. l. c. iii. 148. t. 210.—Austral.
montana, Gaertn. f. l. c. 149.—Austral.
procumbens, Gaertn. f. l. c. 149 = Melichrus rotatus.
rufscarpa, Gaertn. f. l. c. 148. t. 210.—Austral.

ARDISIA, Sw. Prod. Veg. Ind. Occ. & 48 (1788); Fl. Ind. Occ. i. 467. t. 10 (1797). *MYRSINEAE*, Benth. & Hook. f. ii. 645.
ANGUILLARIA, Gaertn. Fruct. i. 379 (1788).
BADULA, Juss. Gen. 490 (1789).
BLADHIA, Thunb. Nov. Gen. i. 6 (1781), *nomen prior.*
BARTHESIA, Comfin. ex A. DC. in Trans. Linn. Soc. xvii. (1837) 112.
CLIMCANDRA, Miq. Pl. Jungh. i. 199 (1852).
CALIZIOLA, Rafin. Sylva Tellur. 167 (1838).
ICACOREA, Aubl. Pl. Gui. ii. Suppl. i. t. 368 (1775).
KATONTHELA, Adans. Fam. ii. 159 (1763).
MILNEA, Rafin. Sylva Tellur. 167 (1838).
MONOPORUS, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 91.
PICKARINIA, Nutt. in Journ. Acad. Sc. Philad. vii. (1834) 95.
PURKINJIA, Presl, Symb. Bot. ii. 17. t. 64 (1833).
PYRGUS, Lour. Fl. Cochinch. i. 120 (1790).
STYLOGYNE, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 91.
WERDILA, Steud. Nom. ed. II. i. 138 (1840).
VEDELA, Adans. Fam. ii. 502 (1763).
acuminata, Spreng. Syst. i. 661 = laterifiora.
acuminata, *Willd. Sp. Pl.* i. 1062.—Ind. occ.
adenanthera, Miq. in Mart. Fl. Bras. x. 285 = Parathesis adenanthera.
affinis, Blume, ex A. DC. in DC. Prod. viii. 96 = Myrsine affinis.
alternata, A. DC. in Ann. Sc. Nat. Sér. II. ii. (1841) 296, sphalm. = oxyphylla.
ambigua, *Mart. Herb. Fl. Bras.* 868.—Bras.

ARDISIA:—
amboinensis, *Scheff. Comm. Myrsin.* 75.—Amboin.
Amherstiana, A. DC. in Trans. Linn. Soc. xvii. (1834) 120 = reflexa.
amplexicaulis, *Brid. Ic. Pl. Ind. Or.* i. 87.—Ind. or.
amoena, Blume, Bijdr. 685 = Blumii.
ancepe, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 280 = colorata.
andamanica, *Kurz, For. Fl. Burma*, ii. 108.—Malaya.
angustifolia, *A. DC. in DC. Prod.* viii. 134.—Ins. Philipp.
arborescens, Wall. Cat. n. 2389 = hamilia.
arguta, *H. B. & K. Nov. Gen. et Sp.* iii. 247.—Am. austr.
attenuata, Wall. Cat. n. 2386 = oxyphylla.
ovenia, Blume, Bijdr. 691 = Myrsine capitellata.
bahamensis, A. DC. in Trans. Linn. Soc. xvii. (1834) 128 = excelsa.
Barthesia, *Steud. Nom.* ed. I. 64.—Zeylan.
Bazaul, Roem. & Schult. Syst. iv. 517 = Embelia robusta.
bhotanica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 528.—Reg. Himal.
bipinnata, *Baker, in Journ. Linn. Soc. xx.* (1884) 201.—Madag.
Blumii, A. DC. in Trans. Linn. Soc. xvii. (1834) 117 = colorata.
Boivinii, A. DC. in DC. Prod. viii. 129 = lanceolata.
bonaccensis, *Scheff. Comm. Myrsin.* 79.—Borneo.
bractesta, *Baker, in Oliver, Fl. Trop. Afr.* iii. 495.—Afr. trop.
bracteosa, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 127.—Mexic.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 67.—Burma.
brasiliensis, *Spreng. Syst. i.* 662.—Bras.
brevifora, *A. DC. in DC. Prod.* viii. 122.—Peruv.
brevipedata, *F. Muell. Fragm.* vi. 163.—Austral.
bumelioides, *Gaertn. in Mem. Am. Acad.* N. S. viii. (1863) 516.—Cuba.
calothyrsa, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 271; *et in Flora*, lii. (1870) 254.—Sumatra.
calycosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 292.—Nicaragua.
canaliculata, *Ladd. Bot. Cab.* t. 1083.—Reg. Colomb.
capitata, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 331.—Ins. Fiji.
capollina, *Moq. & Sesse, ex A. DC. in Trans. Linn. Soc.* xvii. (1834) 119.—Mexic.
caribaea, *A. DC. ex Miq. in Mart. Fl. Bras.* x. 289, in adnot.—Ins. Dominica.
caulifora, *Mart. & Schult. Syst.* iv. 515.—Ins. Philipp.
celebica, *Scheff. Comm. Myrsin.* 83.—Ins. Celebes.
cestrifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 218.—Reg. Colomb.
chinensis, *Benth. Fl. Hongk.* 207.—China.
chrysophylloides, Miq. Fl. Ind. Bat. Suppl. 574 = tuberculata.
clusioides, *Griseb. Fl. Brit. W. Ind.* 396.—Ind. occ.
coriacea, *Yungh. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 304 ('104' err. typ.).—Java.
colorata, Link, Enum. Hort. Berol. i. 194 = paniculata.
colorata, *Roxb. Hort. Beng.* 16; *Fl. Ind.* i. 561.—Ind. or.
complanata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 280 = colorata.
compressa, *H. B. & K. Nov. Gen. et Sp.* iii. 245.—Reg. Colomb.
confinis, *Miq. Fl. Ind. Bat. Suppl.* 576.—Sumatra.
coriacea, *Lodd. Cat. Hort. Veg. Ind. Occ.* 48.—Ind. occ.
courtallensis, *Wight, Ic.* t. 1315.—Ind. or.
crenata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 518.—Malaya.
crenata, *Sims, in Bot. Mag.* t. 1950.—China.
crenata, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 538.—Malaya; China.
crenulata, Lodd. Bot. Cab. t. 2 = crenata.
crenulata, Vent. Choix, t. 5 = Parathesis crenulata.
crispa, A. DC. in Trans. Linn. Soc. xvii. (1834) 124 = crenata.
cubana, *A. DC. l. c.*—Cuba.
Cumingiana, *A. DC. in DC. Prod.* viii. 136.—Ins. Philipp.
cuspidata, *Benth. Bot. Voy. Sulph.* 128.—Ins. Cocos.

ARDISIA :—
cybianthoides, *Miq. Fl. Ind. Bat.* ii. 1028.—Malaya.
cymosa, *Baker, in Oliver, Fl. Trop. Afr.* iii. 495.—Afr. trop.
cymosa, *Blume, Bijdr.* 689.—Java.
decipiens, *A. DC. in DC. Prod.* viii. 125.—Peruv.
decus-montis, *Miq. Pl. Yungh.* 193.—Java.
demissa, *Miq. Fl. Ind. Bat.* ii. 1018.—Malaya.
densa, Miq. i. c. Suppl. 575 = crenata.
dentienlata, *Blume, Bijdr.* 691.—Java.
depressa, *C. B. Clarke, in Hook. fl. Brit. Ind.* iii. 522. —Reg. Himal.
disticha, A. DC. in DC. Prod. viii. 129 = Pimelandra disticha.
divaricata, *Willd. ex Roem. & Schult. Syst.* iv. 804.— Hab. ?
divergens, *Roxb. Hort Beng.* 85; *Fl. Ind.* i. 583.— Malaya.
divergens, Thw. Enum. Pl. Zeyl. 174 pro parte = Gardneri.
Doca, Buch.-Ham. ex A. DC. in DC. Prod. viii. 126 = paniculata.
Doma, Wall. ex A. DC. in Trans. Linn. Soc. xvii. (1834) 118 = humilis.
domingensis, *Willd. ex Roem. & Schult. Syst.* iv. 803. —Ins. S. Domingo.
dubia, *Scheff. Comm. Myrsin.* 67.—Malaya.
durifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 214.—N. Granat.
elegans, Andr. Bot. Repos. t. 623 = lentiginosa.
elliptica, Sedd. For. Man. 188 = humilis.
elliptica, *Thunb. Nov. Gen.* viii. 119.—Zeylan.
erythroxyloides, *Thw. ex Roem. & Schult. Syst.* iv. 804.—Madag.
escalloniοides, *Schlecht. & Cham. in Linnaea*, vi. (1831) 893.—Mexic.
esculenta, *Pav. ex A. DC. in Trans. Linn. Soc.* xvii. (1834) 128.—Am. austr.
eugeniaefolia, Kurz, For. Fl. Burma, ii. 111 = Pimelandra Griffithii.
eugeniaefolia, Wall. Cat. n. 2276 = Pimelandra eugeniaefolia.
excelsa, *Ait. Hort. Kew.* i. 361.—Ins. Baham.; Mader.
eximia, *Miq. Pl. Yungh.* 196.—Java.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* iii. 244.—N. Granat.
fertilis, *Miq. Fl. Ind. Bat. Suppl.* 576 —Sumatra.
floribunda, *Roem. & Schult. Syst.* iv. 804.—Madag.
floribunda, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 272 = neriifolia.
floribunda, *Willd. ex Steud. Nom.* ed. II. i. 129, *nomen.* —Madag.
foetida, *Willd. ex Roem & Schult. Syst.* iv. 803.— Am. austr.
formosana, Rolfe, in Journ. Bot. xx. (1882) 358 = Sieboldii.
Forstenii, *Scheff. Comm. Myrsin.* 61.—Malaya.
frangulaefolia, *Zipp. et Scheff. l. c.* 82.—Malaya.
fuliginosa, *Blume, Bijdr.* 692.—Java.
fusca, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 127. t. 2.—Costa Rica.
fusco-pilosa, *Baker, in Journ. Linn. Soc.* xx. (1884) 200.—Madag.
Gardneri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 521.—Zeylan.
glabra, A. DC. in Trans. Linn. Soc. xvii. (1834) 123 = japonica.
glabrata, Blume, Bijdr. 692 = villosa.
glandulosa, Blume, l. c. 690 = crenata.
glandulosa, Roxb. Hort. Beng. 16; Fl. Ind. i. 584 = Amblyanthus glandulosus.
glandulose-marginata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 128.—Costa Rica.
gracilis, *Miq. in Mart. Fl. Bras.* x. 387.—Bras.
grandiflora, Wall. Cat. n. 2272 = colorata.
grandifolia, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 128.—Burma.
grandis, *Sonn. in Bonplandia*, ix. (1861) 258; *Fl. Vit.* 150.—Ins. Fiji.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 528. —Reg. Himal.
guadalupensis, *Duchass. ex Griseb. in Goett. Abh.* vii. (1857) 387.—Ins. Dominic.
Hamiltonii, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 126.—Ind. occ.
Hasseltii, Blume, ex Scheff. Comm. Myrsin. 86 = lanceolata.

ARDISIA :—
Helleriana, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 86.—Burma.
hortorum, *Maxim. in Gartenfl.* (1865) 368. t. 491.— Japon.
Hostmanni, *Miq. in Mart. Fl. Bras.* x. 388.—Guian.
humilis, *Vahl, Symb. Bot.* iii. 40.—Ind. or.; Malaya.
Hymenandra, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 293 = Hymenandra Wallichii.
Icara, *Buch.-Ham. in Wall. Cat.* n. 2264.—Ind. or.; Burma.
Insularis, *Baker, Fl. Maurit.* 190.—Maurit.
involucrata, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 66.—Reg. Himal.
iraxensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 134.—Costa Rica.
jacquinioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 517.—Ind. occ.
jambolifolia, Miq. Fl. Sungh. 196 = crassa.
japonica, *Blume, Bijdr.* 690.—China; Japon.
javanica, *A. DC. in DC. Prod.* viii. 130.—Java.
Jungbuhniana, *Miq. Pl. Jungh* 192.—Sumatra.
Icenani, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 526.—Reg. Himal.
khasiana, *C. B. Clarke, l. c.* 527.—Reg Himal.
Korthaliana, Scheff. Comm. Myrsin. 57 = Pimelandra Wallichii.
Kurzii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 525. —Burma.
lacvigata, *Blume, Bijdr.* 690.—Java.
laevis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 125. —Costa Rica.
lamponga, *Miq. Fl. Ind. Bat. Suppl.* 577.—Sumatra.
lanceolata, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 583.— Malaya.
laterifolia, *Sw. Prod. Veg. Ind. Occ.* 48.—Ind. occ.
latifolia, Roem. & Schult. Syst. iv. 514 = laurifolia.
latifolia, Sieber, ex A. DC. in Trans. Linn. Soc. xvii. (1834) 113 = guadalupensis.
latipes, *Mart. Herb. Fl. Bras.* n. 570.—Bras.
laurifolia, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 116.—Ind. occ.
lewisiana, Ker-Gawl. in Bot. Reg. t. 533 = crenata.
lepidota, *H. B. & K. Nov. Gen. et Sp.* iii. 247.—N. Granat.
leptantha, *Miq. in Mart. Fl. Bras.* x. 386.—Bras.
Leschenaultii, A. DC. in DC. Prod. viii. 133 = lanceolata.
Lhotzkya, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 127.—Bras.
Liebmanni, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 129. t. 2.—Mexico.
Lindleyana, D. Dietr. Syst. i. 617 = crenata.
linearifolia, *Miq. Fl. Ind. Bat. Suppl.* 577.—Sumatra.
lineata, *Roem. & Schult. Syst.* iv. 804.—Am. austr.
littoralis, Andr. Bot. Rep. t. 630 = humilis.
Lobbiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 529.—Ind. or.
longifolia, *Mart. ex Miq. in Mart. Fl. Bras.* x. 390.— Bras.
longipes, *Baker, in Journ. Linn. Soc.* xxi. (1885) 421. —Madag.
lurida, *Blume, Bijdr.* 687.—Java.
luxonensis, *Presl, Rel. Haenk.* ii. 85.—Ins. Philipp.
macrocarpa, *Scheff. Comm. Myrsin.* 91.—Borneo.
macrocarpa, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 277. —Reg. Himal.
macrophylla, *Reinw. ex Blume, Bijdr.* 691.—Java.
macrophylla, Wall. Cat. n. 2290 = grandifolia.
macroscypha, *Baker, in Journ. Linn. Soc.* xxi. (1885) 420.—Madag.
maculata, *Polt. ex A. DC. in Trans. Linn. Soc.* xvii. (1834) 116.—Ins. S. Doming.
mamillata, *Hance, in Journ. Bot.* xxii. (1884) 290.— Japon.
marginata, *Blume, Bijdr.* 688.—Java.
Martinae, *Miq. in Mart. Fl. Bras.* x. 386.—Bras.
melanocarpa, Schlecht. in Linnaea, viii. (1833) 526 = Parathesia melanocticta.
membranacea, *Wall. Cat.* n. 2288; *ex A. DC. Prod.* viii. 134.—Reg. Himal.
metallica, *N. E. Br. in Illustr. Hortic.* xxviii. (1881) t. 421.—Sumatra.
micrantha, *H. B. & K. Nov. Gen. et Sp.* iii. 246.— Venezuela.
microphylla, *Roem. & Schult. Syst.* iv. 804.—Madag.
Miquelana, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 367; *et in Flora*, liii. (1870) 253.—Malaya.

ARDISIA :—
missionis, *Wall. Cat.* n. 6524; *et ex A. DC. Prod.* viii. 130.—Zeylan.
mollis, Blume, Bijdr. 689 = villosa.
Moonii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 528.—Zeylan.
mucronata, Blume, Bijdr. 685 = lanceolata
multiflora, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 518.—Ind. occ.
multiflora, Willd. ex Roem. & Schult. Syst. iv. 804 = revoluta.
myrtantha, *Baker, in Journ. Linn. Soc.* xxi. (1885) 419.—Madag.
myristicaefolia, *Blume, ex Scheff. Comm. Myrsin.* 68. —Malaya.
nana, Buch.-Ham. ex A. DC. in Trans. Linn. Soc. xvii. (1834) 118 = humilis.
neriifolia, A. DC. Prod. viii. 127, pro parte = andamanica.
neriifolia, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 225 = depressa.
neriifolia, *Wall. Cat.* n. 2278.—Reg. Himal.
nicaraguensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 128.—Nicaragua.
nigrescens, *Oerst. l. c.* 130. t. 2.—Mexic.
nigricans, *Miq. in Mart. Fl. Bras.* x. 389.—Bras.
nitidula, *Baker, in Journ. Linn. Soc.* xx. (1884) 200. —Madag.
nigro-punctata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 127. t. 2.—Costa Rica.
nutans, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 121. —Java.
obtonga, A. DC. l. c. = oxyphylla.
obovata, Blume, Bijdr. 688 = humilis.
obovata, Ham. Prod. Fl. Ind. Occ. 26 = Hamiltonii.
odontophylla, Lindl. Bot. Reg. t. 1892 = chinensis.
odontopbylla, *Wall. Cat.* n. 2270.—Reg. Himal.
oleracea, Buch.-Ham. ex A. DC. in Trans. Linn. Soc. xvii. (1834) 118 = humilis.
oligantha, *Baker, in Journ. Linn. Soc.* xxi. (1885 419.—Madag.
Olivers, *Mast. in Gard. Chron.* (1877) ii. 680.—Costa Rica.
Opegrapha, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 126.—Costa Rica.
orinocensis, *H. B. & K. Nov. Gen. et Sp.* iii. 244.— Fluv. Orinoco.
ovata, Thunb. Nov. Gen. viii. 119 = colorata.
oxyantha, Wall. Cat. n. 2275 = Antistrophe oxyantha.
oxyphylla, *Wall. l. c.* n. 2291; *A. DC. Prod.* viii. 130 —Burma; Malaya.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 575.—Sumatra.
paniculata, A. DC. in DC. Prod. viii. 139 = A. Samara.
panicnlata, *Roxb. Hort. Beng.* 16; *Fl. Ind.* i. 580.— Ind. or.
parasitica, Sw. Prod. Veg. Ind. Occ. 48 = Grammadenia parasitica.
parvifolia, *Humb. ex Willd. sec. Roem. & Schult. Syst.* iv. 804.—Peruv.
pauciflora, Benth. Fl. Hongk. 206 = pentagona.
pauciflora, *Heyne, ex Roxb. Fl. Ind. ed. Carey*, ii. 279. —Ind. or.
pavonina, Buch.-Ham. ex A. DC. Prod. viii. 136 = odontophylla.
pedunculata, Boj. ex Baker, in Journ. Linn. Soc. xx. (1884) 203 = Oncostemum pedicellatum.
pedunculosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 279. —Reg. Himal.
pellucida, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 130. t. 2.—Mexic.
pentagona, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 124.—Java.
pergamacea, Blume, ex A. DC. in DC. Prod. viii. 85 = Embelia pergamacea.
Perrottetiana, *A. DC. in Ann. Sc. Nat. Sér.* II. xvi. (1841) 95.—Ins. Philipp.
peruviana, *Miq. in Mart. Fl. Bras.* x. 390.—Peruv.
petiolaris, *Scheff. Comm. Myrsin.* 91.—Sumatra.
philippinensis, *A. DC. in DC. Prod.* viii. 138.—Ins. Philipp.
Pickeringia, *Torr. & Gray, ex DC. Prod.* viii. 124.— Florida.
plagiosperma, *Scheff. Comm. Myrsin.* 85.—Malaya.
polycarpa, Wall. Cat. n. 2285 = colorata.
polyphylla, *Wall. l. c.* n. 2293; *A. DC. Prod.* viii. 131.—Burma.
polyophala, Wight, Illustr. t. 145 = humilis.
polygama, *A. DC. in DC. Prod.* viii. 138.—Moluccas.

ARDISIA :—
polyneura, Miq. Fl. Jungh. 198 = colorata.
polysticta, Miq. Fl. Ind. Bat. Suppl. 576 = crenata.
porosa *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 522. —Malaya.
primulaefolia, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 234.—China.
procumbens, Gaertn. f. Fruct. iii. t. 210 = Melichrus rotatus.
propinqua, *H. B. & K. Nov. Gen. et Sp.* vii. 218.—Venezuela.
pseudo-Icacorea, *Miq. in Mart. Fl. Bras.* x. 288.—Bras.
pseudojambosa, *F. Muell. Fragm.* iv. 81.—Austral.
pterocaulis, *Miq. Fl. Sumatr.* 194.—Sumatra.
pteropoda, *Miq. Fl. Ind. Bat.* iii. 1095.—Malaya.
pubicalyx, *Miq. l. c. Suppl.* 576.—Sumatra.
pumila, Blume, Bijdr. 668 = Labisia pothoina.
punctata, Blume, l. c. 687 = natans.
punctata, Jack, in Roxb. Fl. Ind. ed. Carey, ii. 275 = divergens.
punctata, Lindl. Bot Reg. t. 827 = crenata.
punctulosa, D. Dietr. Syn. Pl. i. 615 = divergens.
purpurea, Reinw. in Blume, Bijdr. 684 = lanceolata.
pusilla, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 126. —Japon.
pyramidalis, *Pers. Syn.* i. 283 ; *A. DC. in Trans. Linn. Soc.* xvii. (1834) 123.—Ind. occ.
pyramidalis, Roth, Nov. Pl. Sp. 128 = colorata.
pyrgina, St. Lag. in Ann. Soc. Bot. Lyon, vii (1880) 119 = A. Pyrgus.
Pyrgus, *Reinw. & Schult. Syst.* iv. 518.—China.
pyrifolia, *Willd. ex Roem. & Schult. l. c.* 804.—Madag.
quinquangularis, *A. DC. in Ann. Sc. Nat. Sér.* II. xvi. (1841) 95.—Amazon.
quinquegona, Blume, Bijdr. 689 = pentagona.
racemosa, *Spreng. Syst.* i. 661.—Bras.
ramiflora, *Gerst. in Kjoob. Vidensk. Meddel.* (1861) 182.—Costa Rica.
reclinata, *Scheff. Comm. Myrsin.* 69.—Malaya.
reflexa, *Wall. Cat.* n. 2282 ; *A. DC. Prod.* viii. 132.—Burma.
revoluta, *H. B. & K. Nov. Gen. et Sp.* iii. 246.—Mexico.
rhomboidea, *Wight, Ic. Pl. t.* 1213.—Ind. or.
rhynchocarpa, *Scheff. Comm. Myrsin.* 68.—Malaya.
rhynchophylla, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 520.—Reg. Himal.
rigida, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 87.—Burma.
rostrata, Hask. in Flora, lx. (1868) 26 = humilis.
Rothii, A. DC. in Trans. Linn. Soc. xvii. (1834) 117 = colorata.
Roxburghiana, D. Dietr. Syn. Pl. i. 617 = Amblyanthus glandulosus.
rubiginosa, *Miq. Pl. Jungh.* 192.—Sumatra.
salicifolia, A. DC. in DC. Prod. viii. 129 = humilis.
Samara, *A. DC. ex Buek, Index*, iii. *pseud.* 8.—Ind. or.
sanguinolenta, Blume, Bijdr. 685 = colorata.
sanguinolenta, Wall Cat. n. 2287 = Wallichii.
scandens, *Blume, Bijdr.* 686.—Java.
Schomburgkiana, *Miq. in Mart. Fl. Bras.* x. 289.—Bras.
scrobiculata, *Seem. Fl. Vit.* 150.—Ins. Fiji.
semicrenata, Mart. Herb. Fl. Bras. 571 = acuminata.
semidentata, *Miq. Fl. Sumatr.* 191.—Java.
serrata, Pers. Syn. i. 288 = Cavanillesii.
serratifolia, Bedd. in Madr. Journ. Ser. III. i. (1864) 51 = Antistropha serratifolia.
serrulata, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 87 = A. Icara.
serrulata, *Sw. Prod. Veg. Ind. Occ.* 48.—Ind. occ.
sessilis, *Scheff. Comm. Myrsin.* 69.—Malaya.
Sieberi, *Baker, Fl. Maurit.* 190.—Mauriti.
Sieboldi, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 190. —Japon.
simensis, Hort. ex Steud. Nom. ed. II. i. 129 = A. Pyrgus.
solanacea, Roxb. Hort. Beng. 16 ; Fl. Ind. i. 580 = humilis.
Spanoghei, *Scheff. Comm. Myrsin.* 62.—Malaya.
speciosa, Blume, Bijdr. 684 = lanceolata.
sphenobasis, *Scheff. Comm. Myrsin.* 63.—Malaya.
spicata, Wall. Cat. n. 2278 = Labisia pothoina.
squamulosa, *Presl, Rel. Haenk.* ii. 68.—Ins. Luzon.
Stenckii, *Seem. Fl. Vit.* 150.—Ins. Fiji.
stylosa, *Miq. Fl. Jungh.* 196.—Java.
subcoriacea, *Miq. Fl. Ind. Bat.* ii. 1028.—Malaya.
subsessilis, *Miq. Fl. Jungh.* 195.—Sumatra.
subsilvestris, *Miq. Fl. Ind. Bat.* ii. 1027.—Malaya.

ARDISIA :—
surinamensis, *Miq. in Mart. Fl. Bras.* x. 288.—Guiana.
tawoyana, A. DC. in Trans. Linn. Soc. xvii. (1834) 125 = villosa.
synneura, *Scheff. Comm. Myrsin.* 80.—Malaya.
tenuiflora, Kollmann, ex A. DC. Prod. viii. 127 = pauciflora.
tenuifolia, *Blume, Bijdr.* 686.—Java.
tenuiramis, *Miq. Pl. Jungh.* 196.—Java.
tentaculata, *Scheff. Comm. Myrsin* 58.—Ins. Tenate
tetrandra, H. B. & K. Nov Gen. et Sp. iii. 243 = Cybianthus Humboldtii.
Teysmanniana, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 868 ; *et in Flora*, liii. (1870) 353.—Malaya.
thyrsiflora, D. Don, Prod. Fl. Nep. 148 = neriifolia.
tinifolia, *Sw. Prod. Veg. Ind. Occ.* 48.—Jamaica.
tomentosa, *Presl, Rel. Haenk.* ii. 66.—Ins. Luzon.
Tsjeriam-Cottam, Roem. & Schult. Syst. iv. 518 = Embelia robusta.
tuberculata, *Wall. Cat.* n. 2274.—Malaya.
turbacensis, *H. B. & K. Nov. Gen. et Sp.* iii. 245.—N. Granat.
umbellata, *Baker, in Journ. Linn Soc.* xxi. (1885) 420.—Madag.
umbellata, Roxb. Hort. Beng. 16 ; Fl. Ind. i. 582 = humilis.
umbrosa, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie*, ii. (1846) 575.—Java.
undulata, *C. B. Clarke, in Hook. f. Fl. Brit Ind.* iii. 524. —Reg. Himal.
verrucosa, *Presl, Rel. Haenk.* ii. 65.—Ins. Philipp.
vestita, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 274 = villosa.
villosa, *Roxb. Hort. Beng.* 85 ; *Fl. Ind* i. 582.—Malaya.
virens, *Kurz, For. Fl.* ii. 575.—Reg. Himal. ; China.
viridiflora, *Blume, ex A. DC. in DC. Prod.* viii. 88.—Malaya.
vitiensis, *Seem. Fl. Vit.* 150.—Ins. Fiji.
volubilis, Wall. ex Steud. Nom. ed. II. i. 122 = colorata.
Wallichii, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 123. —Burma.
Wallichii, Thw. Enum. Pl. Zeyl. 174 = Moonii.
Wightiana, Wall. Cat. n. 2280 = humilis.
xylosteoides, *Griseb. Fl. Brit. W. Ind.* 395.—Jamaica.
zeylanica, *C. B. Clarke, in Hook. Fl. Brit. Ind.* iii. 520. —Zeylan.
Zollingeri, A. DC. in DC. Prod. viii. 670 = colorata.

ARDISIANDRA, Hook. f. in Journ. Linn. Soc. vii. (1864) 205. t. 1 *PRIMULACEAE*, Benth. & Hook. f. ii. 683.
sibthorpioides, *Hook. f. l. c.*—Afr. trop. occ.

ARDUINA, Adans. Fam. ii. 499 (1768) = **Kunclmannia**, Scop. (Umbell.).

ARDUINA, Mill. Ic. ii. t. 300 (1759) ; Linn. Mant. i. 7 (1767) = **Clarisea**, Linn. (Apocyn.).
acuminata, E. Mey. Comm. Pl. Afr. Austr. 191 = C. acuminata.
bispinosa, Linn. Mant. i. 53 = C. bispinosa.
bispinosa, Lodd. Bot. Cab. t. 887 = C. ferox.
edulis, Spreng. Syst. i. 669 = C. edulis.
erythrocarpa, Eckl. in S. Afr. Journ. i. (1830) 372 = C. erythrocarpa.
ferox, E. Mey. Comm. Pl. Afr. Austr. 191 = C. ferox.
grandiflora, E. Mey. l. c. 190 = C. grandiflora.
haematocarpa, Eckl. in S. Afr. Journ. i. (1830) 372 = C. haematocarpa.
macrocarpa, Eckl. l. c. = C. macrocarpa.

ARECA, Linn. Sp. Pl. ed. I. 1189 (1753). *PALMAE*, Benth. & Hook. f. iii. 883.
ARECA, Linn. Fl. Zeyl. 186 (1747).
alba, Bory, Voy. i. 306 = Dictyosperma album.
Alicae, *W. Hill, ex F. Muell. in Gartenfl.* (1879) 199. —Austral.
angulosa, *Gaertn. Linn. Pralect.* 80.—Ind. or.
arundinacea, *Becc. Malesia*, i. (1877) 23.—Borneo.
augusta, Kurz, in Journ. Bot. xiii. (1875) 351 = Ptychosperma augusta.
aurea, Hort. in Fl. des Serres, Sér. II. vii. (1867–68) 43 = Dictyosperma aureum.
Bambe, *Arruda, in Kuster, Trav. Bras.* 490.—Pernambuc.

ARECA :—
Banksii, A. Cunn. ex Kunth, Enum. Pl. iii. 185 = Rhopalostylis Banksii.
Baueri, Hook. f. ex Lem. Illustr. Hortic. xv. (1868) 575 = Rhopalostylis Baueri.
borbonica, Hort. ex Kunth, Enum. Pl. iii. 186 = Dictyosperma album.
borracensis, *Becc. Malesia*, i. (1877) 22.—Borneo.
Calapparia, Blume, Rumphia, ii. t. 100. f. 2 = Actinophyta Calapparia.
Catechu, *Linn. Sp. Pl.* 1189.—Ind. or. ; Malaya.
Catheeu, Burm. f. Fl. Ind. 241 = A. Catechu.
cincta, Walp. Ann. v. 808 = Acanthophoenix cincta.
cocoides, Griff. in Calc. Journ. Nat. Hist. iv. (1845) 454 = Actinorhytis Calapparia.
communis, Zipp. ex Blume, Rumphia, ii. 78, 119. t. 123 = Drymophloeus communis.
concinna, *Thw. Enum. Pl. Zeyl.* 338.—Zeylan.
cornuta, *Giseke, Linn. Praelect.* 81.—Ind. or.
coronata, Blume, in Mart. Hist. Nat. Palm. 179 = Pinanga coronata.
costata, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 200 (= Pinangae sp.).—Ins. Nicobar.
crinita, Bory, Voy. i. 307 = Acanthophoenix crinita.
cuneifolia, Stokes, Bot. Mat. Med. ii. 818 = humilis.
curvata, Griff. Ic. Pl. As. t. 248 = Pinanga disticha.
curvata, Griff. Notul. iii. 164 = Pinanga paradoxa.
Dicksonii, Roxb. Hort Beng. [105] ; Fl. Ind. iii. 616 = Pinanga Dicksonii.
disticha, Roxb. Il. cc. 620 = Pinanga disticha.
elaeocarpa, Reinw. ex Kunth, Enum. Pl. iii. 195 = Drymophloeus oliviaeformis.
erythrocarpa, H. Wendl. ex Kerch. Palm. 231 = erythropoda.
erythropoda, Miq. in Journ. Bot. Néerl. i. (1861) 6 = Cyrtostachys Renda.
Fraseri, Gaertn. Fruct. i. 19. t 7. fig. 2 = A. Catechu.
furcata, *Becc. Malesia*, i. (1877) 23.—Borneo.
furfuracea, Hort. ex H. Wendl. in Kerch. Palm. 231 = Dictyosperma furfuraceum.
gigantea, Hort. l. c. 254 = Pinanga ternatensis.
glandiformis, *Lam. Encyc* i. 241.—Ins. Molucc.
globuliferu, Lam. Encyc. 895 = Pinanga Kuhlii.
gracilis, Roxb. Hort. Beng. 68 ; Fl. Ind. iii. 619 = Pinanga gracilis
gracilis, Thou. ex Kunth, Enum. Pl. iii. 188 = Dypsis pinnalifrons.
gracilis, Giseke, Linn. Praelect. 80 = Drymophloeus appendiculata.
haematocarpon, Griff. Notul. iii. 165 = Pinanga malaiana.
hexasticha, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 201 = Pinanga hexasticha.
himalayana, Griff. ex H. Wendl. in Kerch. Palm. 231 = A. Catechu.
horrida, Griff. in Calc. Journ. Nat. Hist. v (1845) 463 = Oncosperma horrida.
hortensis, Lour. Fl. Cochinch. 668, in obs. = A. Catechu.
humilis, Blanco, ex H. Wendl. in Kerch. Palm. 231 = triandra ?
humilis, Roxb. ex H. Wendl. l. c. = Pinanga disticha.
humilis, Willd. Sp. Pl. iv. 595 = Drymophloeus saxatilis.
jobiensis, *Becc. Malesia*, i. (1877) 21.—N. Guin.
lactea, Hort. Bogor. ex Miq. Fl. Ind. Bat. iii. 10 = Dictyosperma album.
lanciformis, *Giseke, Linn. Praelect.* 81.—Ind. or.
latsaeta, Scheff. in Tijdschr. Nederl. Ind. xxxii. (1871) 168 = Pinanga latsaeta.
laxa, *Buch.-Ham. in Mem. Wern. Soc.* v. (1824) 309. —Ins. Andam.
litoralis, Blume, Rumphia, ii. 123, in obs. = Drymophloeus communis.
lutescens, Bory, Voy. ii. 296 = Chrysalidocarpus lutescens.
macrocalyx, *Zipp. ex Blume, Rumphia*, ii. 75. t. 101, 160, 163 D.—N. Guinea.
madagascarica, *Stadtmann, ex Boj. Hort. Maurit.* 305. —Madag.
madagascariensis, *Mart. Hist. Nat. Palm.* 179.—Madag.
malaiana, Griff. in Calc. Journ. Nat. Hist. v. (1845) 457 = Pinanga malaiana.
minuta, *Scheff. in Ann. Jard. Buitens.* i. (1876) 146. —Borneo.
monostachya, Mart. Hist. Nat. Palm. 175. t. 1 5. f. 4 = Bacularia monostachya.
nagensis, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 453.—Ind. or.

ARECA :—
Nenga, Blume, in Mart. Hist. Nat. Palm. 179 = Nenga Wendlandiana.
Nibung, Griff. ex H. Wendl. in Kerch. Palm. 231 = Oncosperma filamentosa.
Nibung, Mart. Hist. Nat. Palm. 178. t. 150. f. 4 = Oncosperma horrida.
nigra, Gaertn, ex H. Wendl. in Kerch. Palm. 231 = A. Catechu.
nobilis, Hort.—Cf. Gard. Chron. (1868) 549 = Nephrosperma Van-Houtteana.
Normanbyi, *F. Muell. Fragm.* viii. 235.—Austral.
oleracea, Jacq. Am. Select. 278 = Oreodoxa oleracea.
olivaeformis, Gaertn, in Ord. Nat. Pl. 79 = Drymophloeus olivaeformis.
orinoquensis, Gaertn. Fruct. i. 20 = Punanga Kuhlii.
oviformis, *Gaertn. Fruct. i. 20 = Punanga Kuhlii.*
oviformis, *Gaertn. Fruct. i. 20 = Punanga Kuhlii.*
oxycarpa, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1868) n. v. 1.—Malaya.
paniculata, Scheff. ex H. Wendl. in Kerch. Palm. 231 = Mischophloeus paniculata.
paradoxa, Griff. in Calc. Journ. Nat. Hist. v. (1845) 468 = Pinanga paradoxa.
parva, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Malaya.
polystachys, *Miq. ex H. Wendl. in Kerch. Palm.* 232. —Malaya.
propria, Hort. Bogor. ex Miq. Fl. Ind. Bat. iii. 10 = Dictyosperma album.
procera, Zipp. ex Blume, Rumphia, ii. 95 = Kentia procera.
pumila, Blume, in Mart. Hist. Nat. Palm. 177. t. 153. fig. 1–3 = Nenga Wendlandiana.
punicea, Zipp. ex Blume, Rumphia, ii. 72. t. 122 = Drymophloeus paniceus.
purpurea, *Hort. in Illustr. Hortic.* xxiv. (1877) t. 298. —Maurit.
rubra, Bory, Voy. i. 306 = Acanthophoenix rubra.
rubra, Hort. ex H. Wendl. in Kerch. Palm. 235 = Dictyosperma rubrum.
sanguinea, Zipp. ex Blume, Rumphia, ii. 72 = Drymophloeus paniceus.
sapida, Soland. ex Forst. f. Pl. Esc. 66 = Rhopalostylis sapida.
sapidissima, Burm. f. Fl. Ind. 42 = Drymophloeus saxatilis.
spicata, Lam. Encyc. i. 241 = Calyptrocalyx spicatus.
spinosa, Hasselt & Kunth, Enum. Pl. iii. 185 = Oncosperma filamentosum.
sybestria, Lour. Fl. Cochinch. ii. 568 = glandiformis.
tenella, *Becc. Malesia*, i. (1877) 22.—Borneo.
tigillaria, Jack, in Malay. Misc. ii. (1820) n. vii. p. 98 = Oncosperma filamentosum.
triandra, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 617.—Ind. or.
umbracea, *Linn, Enum. Hort. Berol.* ii. 408, nomen.—Hab.?
vaginata, Gieseke, Linn. Praelect. 78 = Drymophloeus appendiculatus.
Verschaffelti, Hort. ex Lem. in Illustrat. Hortic. xiii. (1866) t. 462–3 = Hyophorbe Verschaffelti.
vestiaria, *Gieseke, Linn. Praelect.* 78.—Hab.?
Wallichiana, Mart. Hist. Nat. Palm. 178 = Iguanura Wallichiana.
Wendlandiana, Scheff. ex H. Wendl. in Kerch. Palm. 233 = Nenga Wendlandiana.

ARELINA, Neck. Elem. i. 88 (1790) = **Berkheya**, Ehrh. (Compos.).

AREMONIA, Neck. Elem. ii. 100 (1790) = **Agrimonia**, Linn. (Rosac.).
agrimonoides, DC. Prod. ii. 588 = Agrimonia Eupatoria.

ARENARIA, Adans. Fam. ii. 256 (1763) = **Spergula**, Linn. (Caryophyll.).

ARENARIA, Rupp. ex Linn. Syst. ed. I. (1745). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 149.
ADENARIUM, Rafin. in Ann. Monthly Mag. (1818) 266.
ALSINANTHUS, Reichb. Handb. 298 (1837).
ALSINE, Scop. Fl. Carn. ed. I. 294 (1772).
ALSINELLA, [Sw.] Summa Veg. Scand. 17 (1814), pro parte; S. F. Gray, Nat. Arr. Brit. Pl. ii. 658 (1821).

ARENARIA :—
AMMODENIA, Patrin, ex S. Gmel. Fl. Sibir. iv. 160 (1769).
HIGELORIA, (Bigelowia) Rafin. in Journ. Phys. lxxiii. (1819) 289.
BRACHYSTEMMA, D. Don, Prod. Fl. Nep. 216 (1938).
CHERLERIA, Hall. It. Helv. ii. n. 79. t. 1 (1740).
CHETROPIS, Rafin. Fl. Tellur. iii. 80 (1836).
DOLOPHRAGMA, Fenzl, in Ann. Wien. Mus. i. (1836) 63. t. 7.
EREMOGONE, Fenzl, Verhr. Alein. 18. t. ad p. 18 (1838).
EUPHALIA, Rupr. Fl. Caucas. 220 (1869).
FACCHINIA, Reichb. Ic. Fl. Germ. v. 29 (1841).
GOUFFEIA, Roblll. & Cast. in DC. Fl. Fr. v. 609 (1815).
GRENIERA, J. Gay, in Ann. Sc. Nat. Sér. III. iv. 27 (1845).
GYPSOPHYTUM, Adans. Fam. ii. 256 (1763).
HALIANTHUS, Fries, Fl. Halland. 76 (1817).
HONKENYA, Ehrh. Beitr. ii. 180 (1788).
LEPTOPHYLLUM, Ehrh. l. c. iv. 147 (1789).
LEPYRODICLIS, Fenzl, in Endl. Gen. 966 (1840).
MERCKIA, Fisch. ex Cham. & Schlecht. in Linnaea, i. (1826) 59.
MERKIA, Reichb. Handb. 298 (1837).
MINUARTIA, Linn. Sp. Pl. 89 (1753).
MOEHRINGIA, Linn. Syst. ed. II. 22 (1740).
NEUMAYERA, Reichb. Ic. Fl. Germ. v. 80 (1841).
ODONTOSTEMMA, Benth. in Wall. Cat. n. 6465 (1828); G. Don, Gen. Syst. i. 449 (1831).
PETTERA, Reichb. Ic. Fl. Germ. v. 38. t. 220 (1841); Nom. 205 (1841).
PLINTHANTHE, Reichb. Handb. 298 (1837).
RHODALSINE, J. Gay, in Ann. Sc. Nat. Sér. III. iv. 25 (1845).
SABULINA, Reichb. Fl. Germ. Excurs. 785 (1832).
SIEHERA, Hoppe, in Flora, ii. (1819) 24.
SOMERAURRA, Hoppe, l. c. 26 (1819).
STROPHIUM, Dulac, Fl. Hautes-Pyr. 247 (1867).
TEYTHEA, Reichb. Nom. 205 (1841).
WIERZBICKIA, Reichb. l. c. (1841).
WILHELMSIA, Reichb. Consp. 206 (1828).
XERALSINE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1868) 347.
abietina, J. & C. Presl, Delic. Prag. i. 63 = grandiflora.
aceross, Boiss. & Heldr. Diagn. Ser. I. viii. 103.—As. Min.
achalensis, *Griseb. in Goett. Abh.* xxiv. (1879) 26.—Reg. Argent.
aciculoris, Fisch. ex DC. Prod. i. 408 = juniperina.
aculeata, *S. Wats. in Bot. King, Exp.* 40.—Am. bor.
adenotrichia, Bong. ex Ledeb. Fl. Ross. i. 291 = Gypsophila petraea.
affinis, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 196.—Hab.?
africana, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 184. —Afr. trop.
agglomerata, Poir. Encyc. Suppl. v. 6 = glomerata.
aggregata, Boiss. Elench. 32, in obs. = erinacea.
aggregata, Loisel. Fl. Gall. i. 317 = tetraquetra.
aizoifolia, Fisch. ex DC. Prod. i. 402 = capillaris.
algarbiensis, *Welw. ex Willk. Ic. Pl. Hispan.* i. 98.—Lusitan.
Alpamarae, *A. Gray, Bot. U. St. Expl. Exped.* i. 116. —Peruv.
alpicola, Tenore, Fl. Nap. iv. 224 = graminifolia.
alpina, Porter, Fl. Colorad. 14 = obtusa.
alsinoides, Raddi, Brev. Observ. 13 = Stellaria uliginosa.
alsinoides, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 201 = laxiuscula.
alsinoides, Fisch. ex DC. Prod. i. 404 = arctica.
amabilis, Bory, in Ann. Gén. Sc. Phys. iii. (1820) 5 = tetraquetra.
amdloois, *Gill. in Hook. Bot. Misc.* iii. (1833) 148.—Chili.
andina, *Rehrb. in Linnaea*, xxxvii. (1871–72) 385.—Bolivia.
andreaeana, Fisch. ex DC. Prod. i. 402 = rigida.
apetala, Vill. Prosp. 43 = biflora.
aphanantha, *Wedd. in Ann. Sc. Nat. Sér. V.* i. (1864) 233.—Peruv.
arabica, Hort. Berol. ex Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop. xi. 49 = tenuifolia.
arctica, *Stev. ex DC. Prod.* i. 404.—Reg. bor. et arct.
Artedia, Vis. Shirp. Dalm. 8. t. 9. f. 2 = graminifolia.

ARENARIA :—
arcticoides, *Porterschi. ex S. Gay, in Ann. Sc. Nat.* Sér. I. iii. (1824) 84, nota.—Europ.
aristata, Rafin. Caratteri, n. 187 = setacea.
armeniaca, *Boiss. Diagn.* Ser. I. i. 46.—Armenia.
Armeriastrum, *Boiss. Elench.* 23.—Hispan.
armerina, *Bory, in Ann. Gén. Sc. Phys.* iii. (1820) *b.* —Hispan.
arvatica, Fenzl, Fl. Sicul. 168 = mucronata.
arvensis, Wallr. Sched. Crit. 200 = Spergula arvensis.
Aucheriana, *Boiss. Diagn.* Ser. I. i. 50.—Persia.
austriaca, All. Fl. Pedem. ii. 112. t. 84. f. 2 = Villarsii.
austriaca, *Jacq. Fl. Austr.* iii. 29. t. 270.—Europ.
baetica, Salem. ex Ball, in Journ. Linn. Soc. xvi. (1877) 366 = emarginata.
baicalensis, Steud. Nom. ed. II. i. 127 = uliginosa.
Balansae, *Boiss. Fl. Orient.* i. 700.—As. Min. ? Persia.
baldensis, Reichb. ex Nym. Consp. 118 = bavarica.
balearica, *Linn. Syst. Nat.* ed. XII. App. 230.—Ins. Balear.; Corsic.
banatica, *Heuff. in Flora*, xxi. (1838) 869.—Hungaria.
Barrelieri, Vill. Hist. Pl. Dauph. iv. 684 = tenuifolia.
Bartolloti, *Tineo, Pl. Rar. Sic. Pugill.* i. 10.—Sicilia.
bavarica, *Linn. Cent.* ii. 17; *Amoen. Acad.* iv. 315.—Europ.
bavarica, *Pall. Reise*, ii. 322.—Rossia.
Benthami, *Edgew. & Hook. f. in Hook. f. Fl. Brit. Ind.* i. 242.—Himal.
Benthamii, *Fenzl, ex Torr. & Gray, Fl. N. Am.* i. 675. —N. Mexic.
Bertcroana, Phil. in Linnaea, xxviii. (1856) 678 = Spergularia villosa?
Besserianeii, Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 202 = graminifolia.
biflora, Grieb. Spicil. Fl. Rumel. i. 208 = rotundifolia.
biflora, *Linn. Mant.* 74.—Europ.
bianica, *Fenzl, ex Rohrb. in Linnaea*, xxxvii. (1871–72) 345.—Peruv.
blepharophylla, *Boiss. Fl. Orient.* i. 693.—Armenia.
Bocconi, Soleir. ex Steud. Nom. ed. II. i. 193, 125 = Spergularia media.
bonariensis, *Gill. in Hook. Bot. Misc.* iii. (1833) 148. —Reg. Argent.
Bourgaeana, Cosa. ex Willk. in Linnaea, xl. (1876) 117 = incrassata.
Bourgaei, *Hemsl. Diagn. Pl. Nov.* 21.—Mexic.
brevicaulis, Sternb. in Spreng. Pugill. i. 81 = lanceolata.
brevifora, Gilib. Fl. Lituan. ii. 155 = serpyllifolia.
brevifolia, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 180. —Am. bor.
Brotherana, *Trautv. in Act. Hort. Petrop.* viii. (1883) 151.—Rossia.
bryoides, Fisch. ex DC. Prod. i. 404 = arctica.
bryoides, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 201.—Mexic.
buniifolia, Poir. Encyc. vi. 363 = laterifiora.
carruleecens, Rudolph, ex Ledeb. Fl. Ross. i. 291 = Gypsophila petraea.
caespitosa, Ehrh. Beitr. v. 177 = verna.
caespitosa, *Phil. in Linnaea*, xxviii. (1856) 673.—Chili.
caespitosa, Saliab. Prod. 299 = balearica.
caespitosa, J. Vahl, ex Nym. Consp. 121 = Sagina nivalis.
californica, *Brewer, in Bot. Calif.* i. 69.—Calif.
calycanthka, Ledeb. in Mém. Acad. Sc. Pétersb. v. (1813) 564 = bellidifolia.
calycina, Hasmael. ex Willk. & Lange, Prod. iii. 603 = Sagina subulata.
calycina, Hort. ex Ledeb. Fl. Ross. i. 348 = tenuifolia.
calycina, *Poir. Voy. Barb.* ii. 157.—Afr. bor.
calycolata, *Poir. Encyc. Suppl.* v. 7.—Hungaria?
campestris, All. Fl. Pedem. ii. 114 = Spergularia rubra.
canadensis, Pers. Syn. i. 504 = Spergularia rubra.
canescens, *Vahl, ex Hornem. Hort. Hafn.* i. 264.—Hab.?
cantabrica, Amo, Fl. Iber. vi. 140 = serpyllifolia.
capensis, *Steud. Nom.* ed. II. i. 124.—Afr. austr.
capillacea, All. Fl. Pedem. ii. 265 = grandiflora.
capillacea, Willd. ex Ledeb. Fl. Ross. i. 367 = capillaris.
capillaris, *Poir. Encyc.* vi. 350.—Sibir.; Am. bor.
capillipes, *Boiss. Voy. Bot. Espagne*, 98, t. 27.—Hispan.; Afr. bor.
capitata, *Lam. Fl. Fr.* iii. 39.—Hisp.; Lusit.
carsicifolia, *Boiss. in Tchihat. Asie Mint. Bot.* i. 385.—Armenia.
carvifolioides, Walt. Fl. Carol. 141 = squarrosa.
casuis, *Botss. Diagn.* Ser. I. viii. 103.—Syria.

ARENARIA :—

caucasica, Adams, ex Ledeb. Fl. Ross. i. 354 = pinifolia.
Cephalotes, *Rich. Fl. Taur. Cauc.* i. 346.—Taurie.
cerastiifolia, Spreng. Syn. v. 62 Ind. = cerastiifolia.
cerastiifolia, *Ramond, ex DC. Fl. Fr.* v. 783.—Pyren.
cerastioides, Pers. Syn. i. 502 = purpurascens.
cerastioides, Poir. Voy. Barb. ii. 166 = spathulata.
Cervianæ, *Cham. & Schlecht. in Linnaea*, i. (1826) 63.—Chili.
Cherleria, *Hook. f. Stud. Fl.* ed. 1. 60.—Europ.
Cherleriæ, Fisch. ex DC. Prod. i. 409 = Stellaria petræa.
cherleriifolia, G. Don, in Lond. Hort. Brit. 478 = hirta.
cherleroides, Vill. Prosp. 48 = lanceolata.
chilensis, *DC. in Mém. Soc. Phys. Genèv.* vi. (1833) 282.—Chili.
ciliaris, *Lour. Trat. Pl. Aragon.* i. 69, 79.—Hispan.
ciliata, *Linn. Sp. Pl.* 425.—Europ.; Am. bor.
ciliata, Sibth. & Sm. Fl. Græc. t. 428 = cretica.
ciliolata, *Edgew. & Hook. f. in Hook. f. Fl. Brit. Ind.* i. 240.—Reg. Himal.
cinerea, *DC. Fl. Fr.* v. 611.—Gallia.
clandestina, *Portenschl. Pl. Dalm.* 13.—Dalmatia.
coerulescens, Auct. vide cærulescens.
collina, Turcz. ex Ledeb. Fl. Ross. i. 349 = verna.
colorata, Turcz. ex Steud. Nom. ed. II. i. 124 = capillaris.
condensata, C. & J. Presl, Delic. Prag. i. 62 = recurva.
conferta, *Boiss. Diagn. Ser. I.* i. 51.—Græcia.
conferta, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 293. —Bolivia.
congesta, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 178.—Calif.
conica, *Boiss. Voy. Bot. Espagne*, ii. 98. t. 31.—Hispan.
conimbricensis, *Brot. Fl. Lusit.* ii. 200.—Lusitan.; Hispan.
conimbricensis, J. Gay, ex Gren. & Godr. Fl. Fr. i. 260 = controversa.
connata, *Rafin. Atl. Journ.* 151.—Am. bor.
controversa, *Boiss. Pugill. Pl. Nov. Hisp.* 100.—Hispan.; Gallia.
cornica, Steud. Nom. ed. II. i. 124 = balearica.
costata, *Bunge, ex Ledeb. Fl. Alt.* ii. 170.—Sibir.
cretica, *Spreng. Syst.* ii. 896.—Græcia; Cyprus; Creta.
cucubaloides, *Sm. Pl. Ic. Ined.* i. 17. t. 17.—Armenia.
dahurica, Fisch. ex DC. Prod. i. 402 = juncea.
dawurica, Spreng. Syst. ii. 895 = Stellaria petræa.
debilis, *Hook. f. Fl. Brit. Ind.* i. 242.—Reg. Himal.
debilis, Turcz. ex Ledeb. Fl. Ross. i. 348 = verna.
decussata, Salisb. Prod. 299 = tetraquetra.
decussata, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 219.—Mexic.
defixa, Desne. in Ann. Sc. Nat. Sér. II. iii. (1835) 277 = graveolens.
densa, Kit. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 200 = gracilis.
densissima, *Wall. Cat.* n. 6440.—Reg. Himal.
denticulata, *Phil. Fl. Atac.* 10.—Chili.
depauperata, *C. Gay, Fl. Chil.* i. 270.—Chili.
diandra, Guss. Fl. Sic. Prod. i. 515 = Spergularia diandra.
dianthoides, Hort. ex Ledeb. Fl. Ross. i. 365 = gypsophiloides.
dianthoides, *Sm. Pl. Ic. Ined.* i. 16. t. 16.—Armen.; Persia.
dichotoma, *Krock. Fl. Sil.* ii. i. 55.—Silesia.
dichotoma, Moench, Meth. 224 = tenuifolia.
dicranoides, *H. B. & K. Nov. Gen. et Sp.* vi. 84.—Mexic.
dilfusa, Ell. Sketch, i. 519 = lanuginosa.
dilfusa, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 201 = sopnioroma.
dimorpha, Adams, ex Ser. in DC. Prod. i. 405 = verna.
Donglasii, *Fenzl, ex Torr. & Gray, Fl. N. Am.* i. 674.—Calif.
dryppides, *Boiss. Diagn. Ser.* I. i. 49.—Aa. Min.
dubia, *Suter, Fl. Helv.* i. 266.—Helvet.
echinata, *Poir. Encyc.* vi. 377.—Europ. austr.
elegans, Cham. & Schlecht. in Linnæa, i. (1826) 57 = *Sirtica*.
elongata, Balbis, ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 200 = australis, grandiflora.
emarginata, *Brot. Fl. Lusit.* ii. 203.—Lusitan.; Marocco.
eriocarpa, *Boiss. Voy. Bot. Espagne*, ii. 102.—Hispan.
erythrosepala, *Boiss. Diagn. Ser.* I. viii. 68 = setacea.

ARENARIA :—

extensa, *Dufour, in Ann. Sc. Gén. Phys.* vii. (1820) 291.—Hispan.
fasciculata, *Forsk. Fl. Aegypt. Arab.* 211.—Arabia.
fasciculata, *Gouan, Illustr.* 30.—Europ.
fasciculata, Hohen. Enum. Fl. Elisabethpol. 232 = sclerantha.
fasciculata, Pall. Ind. Taur.; Bieb. Casp. App. 309 = glomerata.
fasciculata, Sibth. & Sm. Fl. Græc. v. 30 = globulosa.
fastigiata, *Phil. in Linnæa*, xxviii. (1856) 678.—Chili.
fastigiata, Sm. Eng. Bot. xx. t. 1744 = rostrata.
fastigiata, *Schur, Enum. Pl. Transs.* 112.—Transsylv.
Fendleri, *A. Gray, Pl. Fendl.* 13.—Am. bor.
festucoides, *Benth. in Royle, Illustr. Bot. Himal.* 81. t. 21. f. 3.—Reg. Himal.
filicaulis, Fenzl, in Griseb. Spicil. Fl. Rumel. i. 208 = graveolens.
filifolia, Bieb. Fl. Taur. Cauc. i. 344 = graminifolia.
filifolia, Forsk. Fl. Aegypt. Arab. 211 = setacea.
filiformis, Lehtl. Ic. Pl. Syr. dec. iv. 8 = picta.
fimbriata, Bieb. ex Ledeb. Fl. Ross. i. 352 = laricina.
fimbriata, Salzm. ex Rindberg, Monog. Lepigon. 22 = Spergularia fimbriata.
flaccida, [Clairv.] Man. Herb. 149 = Villarsii.
flaccida, Froel. ex Retchb. Fl. Germ. Excurs. 790 = lanceolata.
flaccida, Roxb. Hort. Beng. 84; Fl. Ind. ii. 447 = Spergula pentandra.
flaccida, Schleich. Cat. Pl. Helv. ed. III. 46 = verna.
flexuosa, *Raffin. Atl. Journ.* 151.—Am. bor.
floribunda, *C. Gay, Fl. Chil.* i. 269.—Chili.
foliosa, *Royle, ex Edgew & Hook. f. in Fl. Brit. Ind.* i. 240.—Reg. Himal.
formosa, Fisch. ex DC. Prod i. 402 = capillaris.
Franklinii, *Dougl. ex Hook. Fl. Bor. Am.* i. 101.—Am. bor.
frigida, *Rupr. Fl. Caucas.* 202.—Caucas.
frutescens, Kit. ex Schult. Oestr. Fl. ed. II. i. 667 = setacea.
gallodes, *Ser. in DC. Prod.* i. 410.—Hab.?
geniculata, *Poir. Voy. Barb.* ii. 166.—Reg. Mediterr.
Gerardi, Willd. Sp. Pl. ii. 729 = verna.
Gieseke, Homem. Fl. Dan. t. 1518 = verna.
glabra, Balb. Misc. 21 = Villarsii.
glabra, Ell. Sketch, i. 520 = Stellaria uniflora.
glabra, *Michx. Fl. Bor. Am.* i. 274.—Am. bor.
glabra, Torr. Fl. U. St. 446 = graveolens.
glabrata, Cham. & Schlecht. in Linnæa, i. (1826) 56 = verna.
glandulosa, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 240. —Reg. Himal.
glandulosa, *Jacq. Hort. Schœnb.* iii. 56.—Afr. austr.
globiflora, *Wall. Cat.* n. 6439.—Reg. Himal.
globosa, Spreng. Syst. ii. 400 = globulosa.
globulosa, *Labill. Ic. Pl. Syr. dec.* iv. 6.—Oriens.
glomerata, *Bieb. Fl. Taur. Cauc.* i. 350.—Taurie.
glutinosa, Bieb. l. c. 344 = Holosteum umbellatum.
glutinosa, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 247. t. 2 = Ledebouriana.
glutinosa, Willd. ex Ledeb. Fl. Ross. i. 358 = imbricata.
Gmelins, Bong. ex Ledeb. l. c. 351 = Villarsii.
Gmelini, Fisch. ex DC. Prod. i. 402 = Gypsophila Gmelini.
gorgonea, *T. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 278.—Ins. Cap. Verd.
gothica, *Fries, Mant.* ii. 33.—Europ.
Gouffeu, Pers. in Duch. Rev. Bot. i. (Avr. 1846) 450 = controversa.
Gouffeis, *Chaub. Fl. Pélop.* n. 701. t. 24. f. 2.—Gallo. occ.
gracilis, *Elt. ex Schult. Pl. Rar. Hung.* iii. 305.—Croat.; Dalmat.
graeca, Boiss. ex Nym. Consp. 114 = graveolens.
graminifolia, Auct. ex Steud. Nom. ed. II. i. 124 = grandiflora.
graminea, *C. A. Mey. Verz. Pfl. Cauc.* 220.—Reg. Caucas.
graminifolia, *Arduin. Animad. Alt.* 95.—Italia; Dalmat.
graminifolia, *Schrad. Hort. Gœtt.* i. 5.—Hungar.; Reg. Cauc.; Sibiria.
grandiflora, Cons. ex Willd. in Lianæa, xl. (1876) 117 = incrassata.
grandiflora, *Linn. Syst.* ed. X. 1084.—Europ.
grandiflora, Pall. Reise, iii. 24 = arctica.
grandis, H. B. & K. Nov. Gen. et Sp. vi. 80 = Spergularia grandis.

ARENARIA :—

graveolens, *Schreb. in Nov. Act. Nat. Cur.* iii. (1767) 478.—Græcia; As. Min.; Syria; Arab.
Griffithii, *Boiss. Fl. Orient.* i. 697.—Tibet, occ.; Afghan.
grisenensis, Thomas, Suppl. Cat. (1842) = Villarsii.
groenlandica, *Spreng. Syst.* ii. 402.—Am. arct.
Guicciardii, *Heldr. in Boiss. Diagn.* Ser. II. v. 60.—Græcia.
Guilelmi-Waldemarii, Klotzsch, in Bot. Ergeb Waldem. Reise, 149 = festucoides.
gypsophiloides, *Linn. Mant.* i. 71.—As. Min.; Persia.
gypsophiloides, Stev. ex Ledeb. Fl. Ross. i. 366 = cucubaloides.
gypsophiloides, Willd. ex Ledeb. l. c. 366 = juncea.
Hornbeana, Bartl. ex Presl, Rel. Haenk. ii. 15 = laterifiora.
Halleri, Raeusch. Nom. ed. III. 180 = tenuifolia.
Halleri, Vitm. Summa, iii. 109 = bifora.
halophila, Bunge, ex Ledeb. Fl. Alt. ii. 168 = Spergularia media.
Helmii, *Fisch. ex DC. Prod.* i. 404.—Sibir. ural.
herniariifolia, Desf. Fl. Atlant. i. 358 = geniculata.
heteromalla, Pers. Syn. i. 504 = setacea.
heterosperma, Guss. Fl. Sic. Syn. i. 501 = Spergularia media.
hirsuta, Bieb. Fl. Taur. Cauc. i. 349 = recurva.
hirta, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = cretica.
hirta, Worumsk. in Fl. Dan. t. 1646 = verna.
hispanica, Spreng. Syst. ii. 896 = spathulata.
hispida, *Linn. Sp. Pl.* 425.—Europ.
hispida, Salzm. ex Presl, Fl. Sicul. 104 = Salsmanni.
Holostea, *Bieb. Fl. Taur. Cauc.* i. 345.—Reg. Caucas.
Holostea, Besupr. ex Ser. in DC. Prod. i. 402 = rigida.
holosteoides, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 241. —Reg. Himal.
Hookeri, Nutt. ex Torr. & Gray, Fl. N. Am. i. 178 = Franklinii.
Howellii, *S. Watts. in Proc. Am. Acad.* xx. (1885) 354.—Oregon.
humifusa, Wahlenb. Fl. Lapp. 129 = ciliata.
Huteri, *Kern. in Oestr. Bot. Zeitschr.* xxii. (1872) 268.—Europ.
hybrida, Vill. Prosp. 48 = tenuifolia.
imbricata, Banks, ex Steud. Nom. ed. II. i. 124, 125 = glabra.
imbricata, *Bieb. Fl. Taur. Cauc.* i. 344.—Reg. Caucas.
imbricata, Lang. Hedr. in Anal. Clenc. Nat. v. (1802) 278 = tetraquetra.
imbricata, Rafin. in Med. Repos. N. York, v. (1808) 361 = squarrosa.
inamœna, Steud. Nom. ed. II. i. 125 = imbricata.
incrassata, *Lange, Pl. Nov. Hisp.* fasc. i. t. 4. f. 2.—Hispan.
intricata, Dufour, ex Ser. in DC. Prod. i. 410 = montana.
intricata, *Y. Gay, ex Willk. & Lange, Prod. Fl. Hisp.* iii. 617, in syn.—Hispan.
isaurica, *Boiss. Fl. Orient.* i. 695.—As. Min.
Jamesoniana, *Rohrb. in Linnæa*, xxxvii. (1871-72) 267.—Ecuador.
juncea, *Bieb. Fl. Taur. Cauc.* iii. 309.—Taurie.
juniperina, *Linn. Mant.* 72.—Oriens; Turkest.
juniperina, Pursh, Fl. Am. Sept. 319 = verna.
juniperina, Vill. Hist. Pl. Dauph. iii. 634 = grandiflora.
Juressi, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 212.—Hispan.
Jussiei, *St. Hil. Fl. Bras. Mer.* ii. 174.—Bras.
kansuensis, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 428.—China.
kashmirica, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 236.—Reg. Himal.
Koriniana, *Fisch. ex Ledeb. Fl. Ross.* i. 363.—Rossia.
Kotschyana, Fenzl, in Russegg. Reise, ii. 930 = tmolea.
lanceolata, *Alt. Fl. Pedem.* i. 114.—Europ.
lanuginosa, *Rohrb. in Mart. Fl. Bras.* xiv. 11. 274.—Am. bor. et austr.
lapponica, *Sprengl. Syst.* ii. 402.—Lappon.
laricifolia, Bieb. Fl. Taur. Cauc. i. 347 = pinifolia.
laricifolia, Lapeyr. [Abr. i. 955] ex Steud. Nom. ed. II. i. 124 = grandiflora.
laricifolia, *Linn. Sp. Pl.* 424.—Europ.
laricifolia, Pall. Ind. Taur. [51] ex Steud. Nom. ed. I. 68; ed. II. i. 127 = setacea.
laricifolia, Pursh, Fl. Am. Sept. i. 319 = arctica.
laricina, Cham. & Schlecht. in Linnæa, i. (1826) 57 = Spergula laricina.
laterifiora, *Linn. Sp. Pl.* 423.—Sibir.; Am. bor.

ARENARIA:—
laterifiora, Darl. Fl. Cestr. 277 = Stellaria borealis.
laxa, Fisch. ex Ser. in DC. Prod. i. 404 = grandiflora.
Laxmanni, Fisch. ex Ser. l. c. 402 = capillaris.
Ledebouriana, *Fenzl, in Russegg. Reise*, ii. 981.—Armenia.
leptoclada, Guss. Fl. Sic. Syn. ii. 924 = serpyllifolia.
leptoclados, *Boiss. Fl. Orient.* i. 701.—Europ.; Afr. bor.; Orient.
leptophylla, *Cham. & Schlecht. in Linnaea*, v. (1830) 238.—Mexic.
Lessertiana, *Fenzl, in Russegg. Reise*, ii. 982.—Persia.
lesurina, *Loret, Herb. Loc. et Prost.* 18.—Gall. mer.
libanotica, Kotschy, in Boiss. Fl. Orient. i. 699 = setacea.
Liebmanniana, *Rohrb. in Linnaea*, xxxvii. (1871–72) 274.—Hras.
ligerina, Lec. & Lamotte, Cat. Pl. Cent. Fr. 104 = lesurina.
lignosa, *Phil. Fl. Atac.* 10.—Chili.
linariifolia, Poir. Encyc. vi. 396 = montana.
linaria, *C. A. Mey. ex Ledeb. Fl. Ross.* i. 351.—Rossia.
linifolia, Jacq. Fl. Austr. v. 22 = verna.
linifolia, Linn. Sp. Pl. ed. II. 607 = laricifolia.
linifolia, Willd. ex Spreng. Syst. ii. 899 = grandiflora.
littoralis, *Phil. in Linnaea*, xxviii. (1856) 673.—Chili.
littoralis, Salisb. Prod. 299 = peploides.
Lloydii, Jord. Pugill. 37 = serpyllifolia.
longifolia, Dc. Prod. i. 402 = graminifolia.
longifolia, Bieb. Fl. Taur. Cauc. i. 345 = saxatilis.
Loscosii, *Texid. in Losc. Tvat. Pl. Aragon*, ii. 28.—Hispan.
lychnidea, *Bieb. Fl. Taur. Cauc.* i. 347.—Reg. Caucas.
lychnidea, Willd. ex Ledeb. Fl. Ross. i. 367 = capillaris.
lycopodioides, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1818) 212.—Mexic.
macradenia, *S. Watts. in Proc. Am. Acad.* xvii. (1881–82) 367.—Am. bor.
macrocalyx, *Fenzl, in Syll. Ratisb.* ii. (1828) 243.—Syria.
macrocarpa, Hornem. Hort. Hafn. i. 264 = canescens.
macrocarpa, Ser. Ser in DC. Prod. i. 404 = laricifolia.
macrocarpa, *Pursh, Fl. Am. Sept.* i. 216.—Am. bor.
macrophylla, *Hook. Fl. Bor.-Am.* i. 102.—Ae. or.; Am. bor. occ.
macrorrhiza, Guic. ex Willk. & Lange, Prod. Fl. Hisp. iii. 166 = Spergularia media.
macrorrhiza, Requien, ex Loisel. Nouv. Not. 22 = Spergularia macrorrhiza.
macrospala, *Boiss. Diagn.* Ser. I. i. 52.—Ae. Min.
macrothèca, Hornem. ex Cham. & Schlecht. in Linnaea, i. (1826) 58 = Spergularia macrotheca.
Mandoniana, *Weddl. in Ann. Sc. Nat.* Sér. V. i. (1864) 294.—Reg. And.
marginata, DC. Fl. Fr. v. 793 = Spergularia marginata.
marginata, *Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1818) 212.—Sibir. or.
marina, All. Fl. Pedem. ii. 114 = Spergularia media.
marina, Pall. Reise, iii. 608 = Spergularia marina.
marina, Roth, Tent. Fl. Germ. i. 189 = Spergularia salina.
maritima, Stend. Nom. ed. II. i. 125 = Spergularia media.
Marschlinsii, Koch, in Flora, xxiv. (1841) ii. 509 = serpyllifolia.
media, DC. Prod. i. 401 = Spergularia marina.
media, Linn. Sp. Pl. ed. II. 906 = Spergularia media.
media, Poll. ex Steud. Nom. ed. II. ii. 637, in syn. = Spergula pentandra.
mediterranea, Ledeb. ex Link, Enum. Hort. Berol. i. 431 = macronata.
melaudryoides, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 241.—Reg. Himal.
mexicana, *Steud. Nom.* ed. II. i. 125.—Mexic.
Meyeri, *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 236.—Orient.; Ind. bor. occ.
Meyeri, *Fenzl, ex Ledeb. Fl. Ross.* i. 368.—Aa. temp.
Michauxii, Hook. f. in Trans. Linn. Soc. xxiii. (1861) 287, 323 = stricta.
microphylla, *Phil. in Linnaea*, xxviii. (1856) 673.—Peruv.
minuta, *C. Gay, Fl. Chil.* i. 273.—Chili.
minuta, Lapeyr. Hist. Abr. Pl. Pyr. 255 = grandiflora.
modesta, *Dufour, in Ann. Sc. Gén. Phys.* vii. (1820) 291.—Hispan.
mollugines, *Ser. in DC. Prod.* i. 400.—Mexic.
mollugifolia, *Pourr. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1818) 201.—Hispan.
montana, *Linn. Cent.* i. 19; *Amoen. Acad.* iv. 272.—Gall.; Hisp.
montana, Sieber, ex Steud. Nom. ed. II. i. 125 = macrocalyx.
montícola, Buckl. in Proc. Acad. Sc. Philad. '1861.' (1862) 449 = Benthamii.
monticola, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 238.—Reg. Himal.
Moritzii, *Brueggr. in Zeitschr. Perdinand. Tirol.* III. ix. (1860) 117.—Europ.
mucronata, DC. Fl. Fr. iv. 791 = rostrata.
mucronata, *Linn. Sp. Pl.* 424.—Ital.; Gall.
mucronata, Sibth. & Sm. Fl. Græc. t. 294 = tenuifolia.
multicaulis, Linn. Amoen. Acad. i. 402 = ciliata.
multicaulis, *Phil. in Linnaea*, xxviii. (1856) 673.—Chili.
Munbyi, *Munby, Cat. Pl. Algér.* 6.—Algeria.
muralis, Sieber, ex Spreng. Syst. ii. 897 = graveolens.
musciformis, *Tviana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 150.—N. Granat.
musciformis, *Wall. Cat.* n. 6401.—Tibet.
muscipula, *Fenzl, Verhr. Akin.* anal. tab. iii.—Afr. bor.
muscoides, *H. B. & K. Nov. Gen. et Sp.* vi. 34.—Am. austr.
muscorum, Adams, sec. Fisch. ex DC. Prod. i. 407 = uliginosa.
muscorum, *Fisch. ex DC. l. c.* 409.—Kamtschatka.
muscosa, *Med. in Act. Acad. Theod. Palat* iii. *Phys.* (1778) 202. t. 12.—Europ.
mutabilis, Lapeyr. Hist. Abr. Pl. Pyr. 256 = fasciculata.
nana, *Boiss. & Heldr. Diagn.* Ser. I. viii. 103.—Creta.
nana, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1818) 201.—Cotyledes.
nardifolia, Ledeb. ex Hook. Fl. Bor. Am. i. 98 (1830) = capillaris.
neelgherrensis, Wight & Arn. Prod. i. 48.—Reg. Himal.; Mont. Nilgiri.
nemorosa, H. B. & K. Nov. Gen. et Sp. vi. 35 = lanuginosa.
nemorosa, Stend. Nom. ed. II. i. 126, 127 = trinervia.
nepalensis, *Spreng Syst. Cur. Post.* 181.—Nepal.
nervosa, Lam. Fl. Fr. iii. 36 = trinervia.
nevadensis, Boiss. Diagn. Ser. II. i. 90 = conferta.
nilghervensis, Fenzl, var. *a*, *γ*, ex Boiss. Fl. Orient. i. 700, in syn. = Balansæ.
nitida, *Rohrb. in Linnaea*, xxxvii. (1871–73) 249.—Peru.
nodosa, Bory & Chaub. Fl. Pélop. 28. t. 15 = juniperina.
nodosa, Wallr. Sched. Crit. 200, in obs. = Sagina nodosa.
norvegica, *Gunn. Fl. Norv.* ii. 1100. t. 9. f. 7–9; *ex Hornem. in Fl. Dan.* t. 1289.—Scandin.
nudicaulis, Bertol. Bot. Misc. ii. 222. t. 8 = picta.
obtusa, All. Fl. Pedem. ii. 114 = polygonoides.
obtusa, *Torr. in Ann. Lyc. New York*, ii. (1828) 170.—Am. bor.
obtusiflora, *Kunze, in Flora*, xxix. (1846) 632.—Hispan.
occulta, Fisch. ex DC. Prod. i. 408 = polygonoides, var. stricta.
oligosperma, *C. Gay, Fl. Chil.* i. 272.—Chili.
orbicularis, *Vis. Fl. Dalm.* iii. 180.—Europ.
orbicularis, *Royle, ex Hook. f. Fl. Brit. Ind.* i. 240.—Reg. Himal.
Orbignyana, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 293.—Bolivia.
orcophila, *Hook. f. Fl. Brit. Ind.* i. 238.—Tibet.
oxitoides, *Adams, ex Fisch. in DC. Prod.* i. 402.—Sibir.
oxyantha, *Sibth. & Sm. Fl. Græc. Prod.* i. 308.—Græcia; Creta.
palustris, *C. Gay, Fl. Chil.* i. 271.—Chili.
palustris, *S. Wats. Bot. Calif.* i. 70.—Calif.
pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. viii. 102.—As. Min.
panicolata, *Bunge, in Ledeb. Fl. Alt.* ii. 168.—Sibir. alt.
papillosa, Steud. Nom. ed. II. i. 126 = graveolens.
paradoxa, Bertl. ex Presl, Rel. Haenk. ii. 15 = lanuginosa.
parviflora, *Benth. Pl. Hartw.* 168.—N. Granat.
patagonica, *Phil. in Linnaea*, xxviii. (1856) 671.—Patagonia.
patula, Martr. Fl. Tarn, 107 = serpyllifolia.
patula, *Michx. Fl. Bor. Am.* i. 278.—Am. bor.
pauciflora, *Kit. in Linnaea*, xxxii. (1863) 510.—Hungar.
pedunculosa, *Weddl. in Ann. Sc. Nat. Sér.* V. i. (1864) 294.—Bolivia.
pendula, *Waldst. & Kit. Pl. Rar.* i. 90. t. 87.—Banat.; Thracia.
pensylvanica, Muhl. Ind. Fl. Lanc. 169 = laterifiora?
pentagyna, *Moç. & Sesse, ex DC. Prod.* i. 395.—Mexic.
pentandra, Dufour, in Ann. Sc. Gén. Phys. vii. (1820) 292 = tenuifolia.
pentandra, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 439.—Soongaria.
pentandra, *Turcz. in Flora*, xvii. (1834) i. *Beibl.* 8.—Sibir. baical.
pentandra, Waldr. Sched. Crit. 206 = Spergula pentandra.
peploides, Lapeyr. Hist. Abr. Pl. Pyr. 251 = Polycarpon peploides.
peploides, *Linn. Sp. Pl.* 423.—Reg. bor. temp. et arct.
persica, Boiss. Diagn. Ser. I. i. 49 = Lessertiana.
peruviana, Poepp. ex Rohrb. in Mart. Fl. Bras. xiv. 11. 275 = lanuginosa.
Peyritschii, *Rohrb. in Linnaea*, xxxvii. (1871–73) 264.—Mexic.
pharmacoides, Ser. in DC. Prod. i. 408 = picta.
physodes, *Fisch. ex DC. l. c.* 413.—Am. bor.
picta, *Sibth. & Sm. Fl. Græc. Prod.* i. 304.—Aa. Min.; Syria.
pilifera, Fisch. ex Ledeb. Fl. Ross. i. 352 = Spergula laricina.
pinifolia, *Biel. Fl. Taur. Cauc.* i. 348.—Reg. Caucas.
Plaisii, *Phil. in Linnaea*, xxxiii. (1864–65) 20.—Chili.
Pitcheri, *Nutt. ex Torr & Gray, Fl. N. Am.* i. 180.—Am. bor.
plantaginea, Mert. & Koch, in Roehl. Fl. Deutschl. iii. 278, in syn = trinervia.
Poeppigiana, *Rohrb. in Linnaea*, xxxvii. (1871–72) 265.—Peruv.
polycnemifolia, *Boiss. Diagn.* Ser. I. i. 48.—Armenia; Persia.
polygonoides, *Wulf. in Jacq. Coll.* i. 241. t. 15.—Europ.; As. bor.
polytrichoides, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 237.—Reg. Himal.
Pomeli, *Munby, in Bull. Soc. Bot. Fr.* xl. (1864) 45.—Algeria.
Ponæ, Reichb. Ic. Fl. Germ. ii. 26 = bavarica.
portulacea, Lam. Fl. Fr. iii. 35 = peploides.
Preslii, Nym. Consp. 118 = graminifolia.
procera, Spreng. ex Hornem. Hort. Hafn. 424 = graminifolia.
procumbens, Bove, ex Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 277 = graveolens.
procumbens, *Rupr. Fl. Cauc.* 202.—Caucas.
procumbens, Vahl, Symb. Bot. i. 30. t. 33 = geniculata.
propinqua, Richards. in Frankl. 1st Journ. ed. II. App. 17 = verna.
prostrata, Ser. in DC. Prod. i. 400 = Polycarpæa prostrata.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 438.—China.
pseudarmeriastrum, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 46.—Gallia.
pubescens, Presl, ex Steud. Nom. ed. II. i. 126 = rubiginea.
pubescens, *Redowski, ex Ledeb. Fl. Ross.* i. 349.—Rossia.
pubescens, Stend. Nom. ed. I. 66 = Spergularia media.
pubescens, Urville, in Mém. Soc. Linn. Par. i. (1822) 306 = graveolens.
pubigena, *Steud. Nom.* ed. II. i. 126.—Sicilia.
pulchella, Hoffm. Hort. Mosq. 4 = Gypsophila elegans.
pulchra, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1818) 213.—Armenia.
quitensis, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 238.—Reg. Himal.
Pumilio, R. Br. in Parry, 2nd Voy. App. 281 = section.
purpurea, *Clem. in Lag. Gen. et Sp. Nov.* 15.—Hispan.; Marocco.

ARENARIA :—

pungens, Steph. ex Ser. in DC. Prod. i. 408 = subulata.
pungens, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 179.—Am. bor. occ.
purpurascens, *Ramond, ex DC. Fl. Fr.* iv. 785.—Pyren.
purpurea, Pers. Syn. i. 504 = Spergularia purpurea.
purpurea, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1818) 211 = rubicunda.
Purshiana, Ser. in DC. Prod. i. 414 = Stellaria humifusa.
pusilla, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 367.—Am. bor.
pycnophylla, *Rohrb. in Linnaea*, xxxvii. (1873–72) 250.—Bolivia.
pyrenaica, Hort. ex Steud. Nom. ed. II. i. 126 = montana.
quadrivalvis, R. Br. in Parry, 1st Voy. App. 271 = verna.
querioides, Pourr. ex Willk. in Bot. Zeit. v. (1847) 289 = capitata.
racemosa, *Willk. l. c.*—Hispan.
Raddeana, Ser. in DC. Prod. i. 419 = Stellaria uliginosa.
radiana, *Benth. Pl. Hartw.* 163.—N. Granat.
radicans, Hort. ex Steud. Nom. ed. II. i. 126 = pendula.
radicans, Sprong. Syst. ii. 400 = Spergularia rubra.
Raftesquiana, Ser. in DC. Prod. i. 409 = squarrosa.
Ramondi, Poir. Encyc. Suppl. v. 8 = cerastifolia.
ramosissima, Bieb. ex Steud. Nom. ed. I. 67 = setacea.
ramosissima, *Willd. Enum. Hort. Berol. Suppl.* 24.—Hungar.
recurva, *All. Fl. Pedem.* ii. 119. t. 89. f. 3.—Europ. Oriens.
Redowskii, *Cham. & Schlecht. in Linnaea*, i. (1826) 56.—As. or.
repens, *Ser. in DC. Prod.* i. 410.—Hab.?
repinas, *Hemsl. Diag. Pl. Nov.* 22.—Mexic.
retusa, *Boiss. Voy. Bot. Espagne*, 99. t. 27.—Hispan.
rhodia, Boiss. Diagn. Ser. I. i. 52 = oxypetala.
rigida, *Bieb. Fl. Taur. Cauc.* i. 346.—Iauria.
rivularis, *Phil. Fl. Atac.* 10.—Chili.
Rohrbachiana, *Garcke, in Linnaea*, xxxvii. (1871–72) 811.—Mexic.
Rosani, *Tenore, Prod. Fl. Nap.* 26.—Ital.; Dalmat.
rosea, Presl, Fl. Sicul. 165 = geniculata.
rosea, Salzm. ex Ball, in journ. Linn. Soc. xvi. (1878) 365 = emarginata.
Rossii, R. Br. in App. Parry, Voy. 18 = stricta.
Royleï, Fenzl, ex Klotzsch, in Bot. Ergeb. Waldem. Reise, 149 = lesiocoides.
rostrata, Waldst. & Kit ex Willd. Enum. Hort. Berol. 481 = laricifolia.
rostrata, *Pers. Syn.* i. 504.—Europ. alp.
rotundifolia, *Bieb. Fl. Taur. Cauc.* i. 348.—Graecia; Reg. Cauc.; As. Min.
rubella, Sm. Engl. Bot. Suppl. t. 2588 = verna.
rubicunda, *Spreng. Syst.* ii. 399.—Sibir.
rubra, Linn. Sp. Pl. 423 = Spergularia rubra.
rupestris, *Labill. Pl. Syr. dec.* iv. 8. t. 4. f. 1.—Liban.
rupicola, Hook. f. Fl. Brit. Ind. i. 248 err. typ. = seq.
rupifraga, Fenzl, ex Ledeb. Fl. Ross. i. 780 = Thylacospermum rupifragum.
rusciifolia, *Poir. Encyc.* vi. 365.—Canada.
ruscifolia, Requien, in Guer. Vaucl. ed. II. 254 = cinerea.
sabulinea, *Grisch. in Ruszegg. Reise*, ii. 333.—Mesopot.
sajanensis, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1818) 200.—Sibir, alt.
salina, Ser. in DC. Prod. i. 401 = Spergularia marina.
salsuginea, Bunge, ex Ledeb. Fl. Alt. ii. 163 = Spergularia diandra.
Salzmanni, *Presl, Fl. Sicul.* 168.—Sicil.
sancta, *Yenkn., in Termless. Pflanzen*, vi. (1882) 165.—Europ. austr.
saponarioides, *Boiss. & Bal. Diagn.* Ser. II. vi. 35.—Thrygia.
saxatilis, Bout. ex Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 631 = obtusiflora.
saxatilis, Asch. ex Steud. Nom. ed. II. i. 127 = setacea.
saxatilis, C. C. Gmel. Fl. Bad. ii. 367 = verna.
saxatilis, *Linn. Sp. Pl.* 423.—Europ.; Sibir.
saxatilis, Vill. Fl. Delphin. 43 = rostrata.
Saxifraga, *Pritzeld. in Flora*, xix. (1836) ii. 434.—Rumelia.
saxosa, *A. Gray, Pl. Wright.* ii. 18.—N. Mexic.
scabra, *Poir. Encyc.* vi. 377.—Europ.
scandinavica, Sprong. Syst. ii. 399 = Stellaria biflora.
scarioua, *Boiss. in Tchihat. Asie Min. Bot.* i. 294.—Armenia.
scopulorum, *H. B. & K. Nov. Gen. et Sp.* vi. 21.—Peruv.
sedoides, *Pront. ex Hoch, Syn. Fl. Germ.* ed. I. 114.—Tyrol.

ARENARIA :—

segetalis, Lam. Fl. Fr. iii. 43 = Spergularia segetalis.
sericea, Ser. in DC. Prod. i. 414 = purpurea.
serpens, Flisch. ex Ser. l. c. 404 = stricta.
serpens, *H. B. & K. Nov. Gen. et Sp.* vi. 32.—Ecuador.
serpyllifolia, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 619 = nevadensis.
serpyllifolia, *Linn. Sp. Pl.* 423.—Europ.; As. temp.; Ind. or.
serpyllifolia, Guss. Fl. Sic. Syn. i. 495 = leptoclados.
serpyllioides, *C. Gay, Fl. Chil.* i. 371.—Chili.
setacea, Fisch. ex Ser. in DC. Prod. i. 409 = lychnides.
setacea, Adams, ex Ser. l. c. 408 = capillaris.
setacea, *Thuill. Fl. Par.* ed. II. 230.—Reg. Mediterr.; Orient.; Sibir.
setosa, Willd. ex Ledeb. Fl. Ross. i. 368 = capillaris.
sibirica, Pers. Syn. i. 504 = capillaris.
sipyles, Boiss. Diagn Ser. I. i. 51 = graveolens.
sitchensis, D. Dietr. Syn. Pl. ii. 1565 = peploides.
Smithii, Steud. Nom. ed. II. i. 127 = globulosa.
soratensis, *Kohrb. in Linnaea*, xxxvii. (1871–72) 266.—Bolivia.
sordida, *Chamb. Fl. Pfalz.* n. 699. t. 24. f. 1.—Graecia.
spathulata, *Desf. Fl. Atlant.* i. 338.—Hispan.; Afr. bor.
Spergula, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 293 = Spergularia media.
spergulioides, *Rafin. Atl. Journ.* 153.—Am. bor.
spergulioides, *Trevir. ex Steud. Nom.* ed. II. i. 127.—Chili.
sphaerocarpa, Martr. Fl. Tarn, 107 = patula.
sphaerocarpa, Tenore, Syll. Fl. Neap. iv. 222 = serpyllifolia.
spinulifera, Ser. in DC. Prod. i. 406 = Habrosia spinulifera.
squarrosa, *Micks. Fl. Bor. Am.* i. 278.—Am. bor.
stellarioides, Pers. Syn. i. 508 = austriaca.
stellarioides, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1818) 209.—Caucas.
stenocarpa, *Phil. Fl. Atac.* 10.—Chili.
stenopetala, Turcz Cat. Baikal. n. 357, in Bull. Soc. Nat. Mosc. (1838) 89 = arctica.
stenophylla, Ledeb. Ind. Hort. Dorp.; Fl. Alt. ii. 165 = graminifolia.
Stevensiana, *Boiss. Fl. Orient.* i. 692.—Armenia.
stolonifera, Vill. ex DC. Prod. i. 404 = grandiflora.
Strscheyi, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 240.—Thibet.
striata, Linn. Cent. ii. 27 = capillacea, cephalotes, laricifolia, recurva, saxatilis, verna.
striata, *Vill. Hist. Pl. Dauph.* iv. 680. t. 47.—Gallia.
stricta, *Micks. Fl. Bor. Am.* i. 274.—Am. bor.
stygia, Boiss. & Heldr. Diagn. Ser. II. i. 91 = cretica.
subulata, *Ser. in DC. Prod.* i. 408.—Sibir.
subulifolia, Presl, Fl. Sicul. 162 = mucronata.
succulenta, Ser. in DC. Prod. i. 400 = Polycarpon succulentum.
sulcata, Schrebt. in Ges. Naturf. Fr. Berl. Mag. vii. (1818) 212 = hirta.
Szowitsii, *Boiss. Fl. Orient.* i. 695.—Reg. Caucas.
tenella, *Kit. in Schult. Oestr. Fl.* ed. II. i. 662.—Hungar.
tenella, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 179.—Am. bor.
tenella, Vrcz. ex Steud. Nom. ed. II. i. 127 = stricta.
tenera, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 242.—Reg. Himal.
tenuifolia, *Schrad. ex Steud. Nom.* ed. II. i. 127.—Hab.?
tenuifolia, *Linn. Sp. Pl.* 424.—Europ.
tenuis, *C. Gay, in Walp. Ann.* vii. 800.—Hispan.
teres, Kit. ex Steud. Nom. ed. I. 67 = setacea.
teretifolia, *Phil. Fl. Atac.* 10.—Chili.
tetragyna, *Willd. ex Schlecht. Ges. Naturf. Fr. Berl. Mag.* vii. (1818) 201.—Am. austr.
tetraquetra, *Linn. Sp. Pl.* 428.—Gallia.
tetrasticha, *Boiss. Diagn.* Ser. I. i. 50.—Persia.
thesiifolia, *Presl. ex Reichb. Ic. Fl. Germ.* v. t. 219.—Alp. Algoviae.
thymifolia, Pursh, Fl. Am. Sept. i. 317 = Stellaria humifusa.
thymifolia, *Sibth. & Sm. Fl. Graec. Prod.* i. 304.—Creta.
tmolea, *Boiss. Diagn.* Ser. I. i. 50.—As. Min.
tomentosa, *Willd. in Linnaea*, xxv. (1852) 15.—Hispan.
tremula, *Boiss. Diagn.* Ser. I. viii. 101.—Syria.
triandra, Schrank, Hort. Monac. t. 91 = mucronata.

ARENARIA :—

trichocalycina, *Tenore & Guss. ex Boiss. Fl. Orient.* i. 681.—Graecia.
trichotoma, Boiss. Diagn. Ser. II. v. 90 = Ledebouriana.
trichotoma, *Royle, ex Hook. f. Fl. Brit. Ind.* i. 235.—Reg. Himal.
trifora, Linn. Mant. 240 = grandiflora.
trifora, Vill. Prosp. 43 = Villarsii.
trinervia, *Linn. Sp. Pl.* 423.—Europ.
tschuktschorum, *Regel, in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 284.—Sibir.
serranica, Sprang. in Erech.' sec. Steud. Nom. ed. II. i. 124, 127 = graminifolia.
uliginosa, Schleich. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1818) 207 = stricta.
umbellata, [Clairv.] Man. Herb. 149 = Holosteum umbellatum.
umbellata, Sibth. & Sm. Fl. Graec. v. 28 = Holosteum tenerrimum.
umbellata, *[Soland. in] Russell, Aleppo*, ed. II. ii. 252.—As. Min.
umbrosa, *Bunge, in Ledeb. Fl. Alt.* ii. 178.—Sibir, alt.
unifora, *Luce, Topog. Nachr. Oessl*, 141.—Ins. Oesil.
uniflora, Poir. Encyc. vi. 375 = recurva.
unilateralis, Steud. Nom. ed. II. i. 127 = segetalis.
uralensis, *Pall. ex Spreng. Syst. ii.* 396.—Sibir. ural.
varia, Dulac, Fl. Hautes-Pyr. 249 = serpyllifolia.
valentina, *Boiss. Diagn.* Ser. II. i. 90.—Europ.
verna, Georgi, Beschr. Russ. Reich. iii. iv. 979 = setacea.
verna, Hort. ex Vilm. F. ex Terre, ed. 1. 756 = Sagina subulata.
verna, *Linn. Mant.* 72.—Europ.; Am. bor.
verticillata, Willd. Sp. Pl. ii. 725 = Acanthophyllum verticillatum.
Villarsii, *Balb. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1818) 199.—Europ. med.
villosa, *Ledeb. in Mém. Acad. Pétersb.* v. (1812) 534.—Sibir. or.
villosa, Wulf. in Jacq. Coll. iv. 319 = Wulfenii.
violacea, *Ledeb. in Mém. Acad. Pétersb.* v. (1812) 533.—Sibir.
viscida. Loisel. Not. 58 = serpyllifolia.
viscidula, Thuill. Fl. Par. i. 219 = tenuifolia.
viscosa, Fisch. ex Ser. in DC. Prod. i. 408 = subulata.
viscosa, Pers. Syn. i. 504 = tenuifolia.
viscosa, Pourr. ex Webb & Berth. Phyt. Canar. i. 149 = procumbens.
vulcanorum, *Maxim. in Franch. & Sav. Enum. Pl Jap.* i. 52.—Japon.
vulgaris, Bernh. Syst. Verz. Erf. 259 = Stellaria media.
Wallichiana, Ser. in DC. Prod. i. 411 = serpyllifolia.
Wulfenii, *Steud. Nom.* ed. II. i. 128.—Alp. Carniol.

ARENBERGIA, Mart. & Gal. in Bull. Acad. Brux. xi. t. (1844) 378 = Bunitoma, Salisb. (Gentian.).
glauca, Mart. & Gal. l. c. 374 = E. exaltatum.

ARENGA, Labill. in Mém. Inst. Par. iv. (1801) 209.
PALMAE, Benth. & Hook. f. iii. 917.
GOMUTUS, Correa, in Ann. Mus. Par. ix. (1807) 288. t. 24. f. 1.
SAGUERUS, Adans. Fam. ii. 24 (1763).
Griffithii, Seem. ex H. Wendl. in Kerch. Palm. 282 = saccharifera.
javanica, *Hort. ex H. Wendl. l. c.*—Java.
obtusifolia, *Mart. Hist. Nat. Palm.* 191. t. 147, 148.—Ind. or.; Malaya.
saccharifera, *Labill in Mém. Inst. Par.* iv. (1801) 209.—Malaya.
Westerhoutii, Griff. in Calc. Journ. Nat. Hist. v. (1845) 474 = obtusifolia.
Wightii, *Griff. l. c.* 475.—Ind. or.

ARETHUSA, Gronov. Fl. Virg. 184 (1743). *ORCHIDEAE*, Benth. & Hook. f. iii. 514.
CRYBE, Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
EPIPACTIS, Adans. Fam. ii. 70 (1763).
ORCHIDIUM, Mitchell, in Act. Nat. Cur. vii. (1748) 216, App.
alaris, Rasbach. Nom. ed. III. 962 = Pterygodium catholicum.
alata, Poir. Encyc. Suppl. i. 443 = Pterostylis praecox.
benguinensis, Herb. Calc. ex Hemsl. in journ. Linn. Soc. xx. (1884) 311 = Didymoplexis pallens.
bifoliata, Linn. f. Suppl. 403 = Epipactis bonariensis.
bulbosa, *Linn. Sp. Pl.* 950.—Am. bor.

ARETHUSA :—
caparsis, Linn. Dis. Pl. Rar. Afr. 28 ; Amoen. Acad. vi. 111 = Disperis caparsis.
catenata, Sm. Exot. Bot. ii. 89 = Calochenia carnea.
ciliaris, Linn. f. Suppl. 405 = Bartholina pectinata.
cordata, Poir. Encyc. Suppl. i. 444 = Disperis cordata.
crispa, Thunb. Prod. Pl. Cap. 3 = Corycium crispum.
cucullata, Poir. Encyc. Suppl. i. 444 = Disperis cucullata.
divaricata, Linn. Sp. Pl. 951 = Pogonia divaricata.
ecristata, Griff. Notul. iii. 378 = Didymoplexis pallens.
foliosa, *Walt. Fl. Carol.* 222.—Am. bor.
gentianoides, Sw. Fl. Ind. Occ iii. 1436 = Pogonia gentianoides.
glutinosa, Blanco, Fl. Filip. ed. I. 641 = Geodorum semicristatum.
japonica, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858-9) 409.—Japon.
lutea, Gaudich. in Ann. Sc. Nat Sér. I. v. (1825) 101 = Chloraea Gaudichaudii.
medeoloides, Pursh, Fl. Am. Sept. ii. 591 = Pogonia verticillata.
ophioglossoides, Linn. Sp. Pl. 951 = Pogonia ophioglossoides.
parviflora, Michx. Fl. Am. Bor. ii. 160 = Pogonia pendula.
pendula, Willd. Sp. Pl. iv. 89 = Pogonia pendula.
petraea, *Pers. Syn.* ii. 512.—Afr. trop.
picta, Anders. in Trans. Soc. Arts, xxv. (1867) 199 = Spiranthes picta.
plicata, Andr. Bot. Rep. v. 321 = Pogonia plicata.
racemosa, Walt. Fl. Carol. 222 = Ponthieva glandulosa.
roseo, *Benth in Journ. Linn. Soc.* xviii. (1881) 348.—Mexic.
secunda, Thunb. Prod. Pl. Cap. 3 = Disperis secunda.
simplex, Thou. Orch. Afr. t. 24 = Pogonia Thouarii.
spicata, Walt. Fl. Carol. 222 = Aplectrum hyemale.
tigridiaefolia, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 14.—Mexic.
triantophorea, Sw. in Vet. Akad. Nya Handl. Stockh. (1800) 230 = Pogonia pendula.
verticillata, Muehl. ex Willd. Sp. Pl. iv. 81 = Pogonia verticillata.
villosa, Linn. f. Suppl. 405 = Disperis villosa.

ARETIA, Hall. Enum. Stirp. Helv. ii. 485 (1742) ; Linn. Sp. Pl. ed. I. 141 (1753) = **Androsace**, Linn. (Primul.).
alba, Rafin. Fl. Ludov. 34 = Andr. alba.
alpina, Linn. Sp. Pl. 141 = Andr. alpina.
alpina, Wulf. in Jacq. Fl. Austr. 36 = Andr. hedraeantha.
argentea, Gaertn. f. Fruct. iii. 107 = Andr. imbricata.
baldensis, Turra, Fl. Ital. Prod. 68.—Italia.
brevis, Hegetschw. Fl. Schw. 190 = Andr. alpina.
brutia, Nym. Consp. 607 = Andr. hedraeantha.
bryoides, Lotsel. Fl. Gall. ed. I. 111 = Andr. bryoides.
cana, Willd. ex Stend. Nom. ed. II. i. 129, 527 = Draba aretioides.
Charpentieri, Heer, in Hegetschw. Fl. Schw. 987 = Andr. imbricata.
ciliata, Lotsel. Fl. Gall. ed. I. 112 = Andr. alpina.
cylindrica, Lotsel. l. c. 111 = Andr. cylindrica.
glacialis, Schleich. Cat. Pl. Helv. ed. IV. 8 = Andr. alpina.
Halleri, Linn. Sp. Pl. ed. II. 204, in syn. = Andr. carnea.
Hausmanni, Nym. Consp. 607 = Andr. Hausmanni.
hedraeantha, Nym. l. c. = Andr. hedraeantha.
Henrii, Hegetschw. Fl. Schw. 188 = Andr. Hausmanni.
helvetica, Georgi, Beschreib. Russ. Reich. iii. tv. 760 = Andr. villosa.
helvetica, Murr. Syst. ed. XIII. 162 = Andr. bryoides.
hirtella, Nym. Consp. 607 = Andr. alpina.
longiflora, Fisch. ex A. DC. in DC. Prod. viii. 668 = Dionysia odora.
multiflora, Vand. Fasc. Fl. 8.—Lusit. ?
pennina, Murith, Bot. Val. 58 = Andr. alpina.
pubescens, Lotsel. Fl. Gall. ed. I. 111 = Andr. alpina.
pyrenaica, Loisel. l. c. = Andr. pyrenaica.
rubra, Schleich. ex Steud. Nom. ed. II. i. 128 = Andr. hedraeantha.
tomentosa, Schleich. Cat. Pl. Helv. ed. III. 8 = Andr. imbricata, bryoides.
Vandelli, Turra, Fl. Ital. Prod. 68.—Italia.

ARETIA :—
Vitaliana, Murr. Syst. ed. XIII. 162 = Douglasia Vitaliana.
Wulfeniana, Sieber, ex Nym. Consp. 607 = Andr. hedraeantha.

ARETIA, Link, Handb. ii. 411 (1829) = **Primula**, Linn.
auriculata, Link, l. c. = P. Auricula.
carniolica, Link, l. c. 413 = P. carniolica.
ciliata, Link, l. c. = P. ciliata.
crenata, Link, l. c. 411 = P. marginata †
Floerkeana, Link, l. c. 413 = P. Floerkeana.
glaucescens, Link, l. c. 413 = P. calycina.
glutinosa, Link, l. c. 413 = P. glutinosa.
integrifolia, Link, l. c. 413 = P. integrifolia.
minima, Link, l. c. 413 = P. minima.
Palinuri, Link, l. c. 411 = P. Palinuri.
spectabilis, Link, l. c. 412 = P. spectabilis.
truncata, Link, l. c. 413 = P. minima.
villosa, Link, l. c. 411 = P. villosa.

ARETIASTRUM, Spach, Hist. Vég. Phan. x. 304 (1841) = **Valeriana**, Linn.

ARGAN, Dryand. in Trans. Linn. Soc. ii. (1794) 225 = seq.

ARGANIA, Roem. & Schult. Syst. iv. 46 (1819). *SAPOTACEAE*, Benth. & Hook. f. ii. 655.
VERLANGIA, Neck. Elem. ii. 125 (1790) ; Rafin. Sylva Tellur. 38 (1838).
Sideroxylon, *Roem. & Schult. l. c.* 502.—Marocco.

ARGELASIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 357 = **Genista**, Tourn. (Legum.).
cornica, Fourr. l. c. = G. cornica.
Scorpius, Fourr. l. c. = G. Scorpius.

ARGELIA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 331. t. 13, G = **Solenostemma**, Hayne (Asclep.).
Delilii, Decne. l. c. = S. Argel.

ARGEMONE, Tourn. ex Linn. Syst. ed. I. (1735). *PAPAVERACEAE*, Benth. & Hook. f. i. 52.
ECHTRUS, Lour. Fl. Cochinch. 344 (1790).
alba, *Lestib. Bot. Belg.* ed. II. iii. 138 (= seq.).
alba, Rafin. Fl. Ludov. 36 = mexicana.
albiflora, Hornem. Hort. Hafn. 469 = mexicana.
armeniaca, Linn. Sp. Pl. 509 = Papaver armeniacum. Barclayana, *Penny, ex Loud. Gard. Mag.* vi. (1830) 116. —Mexic.
cambrica, Desf. in Dict. Sc. Nat. ii. 481 = Meconopsis cambrica.
fruticosa, *Thurber, ex A. Gray, in Mem. Am. Acad.* N. S. v. (1855) (*Pl. Nov. Thurb.*) 306.—Mexic.
Georgiana, Croom, in Am. Journ. Sc. Ser. I. xxv. (1834) 75 = mexicana.
grandiflora, Sweet, Brit. Flow. Gard. iii. t. 226 = mexicana.
hispida, *A. Gray, Pl. Fendl.* 5.—Am. bor.
Hunnemannia, Otto, in Otto & Dietr. Allg. Gartens. i. (1833) 398 = Hunnemannia fumariaefolia †
intermedia, Sweet, Hort. Brit. ed. II. 588 = Barclayana. mexicana, *Linn. Sp. Pl.* 508.—Am. bor.; Mexic.
mucronata, Dum.-Cours. ex Steud. Nom. ed. II. i. 128 = mexicana.
munita, Dur. & Hilg. in Journ. Acad. Nat. Sc. Philad. Ser. II. (1855) 87 = hispida.
ochroleuca, *Sweet, Brit. Fl. Gard.* iii. t. 242.—Mexic. platyceras, *Link & Otto, Ic.* ii. 48.—Mexic.
formosa, Linn. Sp. Pl. 509 = Papaver pyrenaicum. rosea, *Hook. Bot. Misc.* ii. (1831) 207.—Chili.
scrobilis, Stokes, Bot. Mat. Med. iii. 195 = mexicana.
spinosa, Moench, Meth. 237 = mexicana.
sulphurea, Sweet, ex Loud. Hort. Brit. 216 = ochroleuca.
versicolor, Salisb. Prod. 376 = mexicana.

ARGENOPE, Salisb. Gen. Pl. Frag. 102 (1866) = **Narcissus**, Linn. (Amaryll.).
serotina, Salisb. l. c. = N. serotinus.

ARGENTINA, Lam. Fl. Fr. iii. 118 (1778) = **Potentilla**, Tourn. (Rosac.).
rubra, Lam. l. c. 120 = P. palustris.
rupestris, Lam. l. c. = P. rupestris.
supina, Lam. l. c. 119 = P. supina.
vulgaris, Lam. l. c. = P. Anserina.

ARGITHAMNIA, Sw. Prod. Veg. Ind. Occ. 89 (1788), emend. in el. Fl. i. 885 (1797). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 303.
APHORA, Nutt. in Trans. Am. Phil. Soc. Ser. II. v. 174 (1837).
ARGOTHAMNIA, Spreng. Syst. iii. 847 (1826).
ARGYROTHAMNIA, Meell. Arg. in Linnaea, xxxiv. (1865) 65.
ARGYTHAMNIA, P. Br. Hist. Jamaic. 338 (1756).
CHIROPETALUM, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 21.
CHLOROCAULON, Klotzsch, in Kndl. Gen. Suppl. iv. iii. 89 (1847).
DESFONTAINA, Vell. Fl. Flum. 95 (1825).
DITAXIS, A. Juss. Euph. Tent. 27. t. 7 (1824).
PHILYRA, Klotzsch, in Wiegm. Archiv, vii. (1841) 191.
SZAOPHYTON, Benth. Bot. Voy. Sulph. 59 (1844).
STENONIA, Didrichs. in Kjoeb. Vidensk. Meddel. (1857) 146.
adenophora, *A. Gray, in Proc. Am. Acad.* viii. (1870) 294.—Mexic.
anisotricha, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 314.—Bras.
aphoroides, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 146.—Texas.
Berteroana, *Muell. Arg. l. c.* 151.—Chili.
boliviensis, *Muell. Arg. l. c.* 149.—Chili.
brasiliensis, *Muell. Arg. l. c.* 144.—Bras.
brevirames, *Muell. Arg. l. c.* 146.—Bolivia.
calycina, *Muell. Arg. in Flora*, lv. (1872) 10.—Ind. occ.
candicans, *Sw. Prod. Veg. Ind. Occ.* 89.—Ind. occ.
canescens, *Phil. f. Cat. Pl. Vasc. Chil.* 292.—Chili.
cantonensis, *Hance, in Journ. Bot.* xvi. (1878) 14.—Chins.
catamarcensis, *Hieron. in Bull. Acad. Nac. Cordova*, iv. (1881) 449.—Reg. Argent.
desertorum, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 310.—Bras.
diolca, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 145.—Bras.
fascicolata, *Muell. Arg. l. c.* 148.—Ind. occ.
Fendleri, *Muell. Arg. l. c.* 145.—Venezuela.
foliosa, *Muell. Arg. l. c.* 150.—Bras.
Gardneri, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 310.—Bras.
guatemalensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 145.—Mexic.
gymnadenia, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 316.—Bras.
herbacea, *Spreng. Syst.* iii. 848.—Am. bor.
heterantha, *Muell. Arg. in DC. Prod.* xv. ii. 735.—Mexic.
humilis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 147.—Mexic.
illinoiensis, *Muell. Arg. l. c.* 146.—Mexic.
laevis, *Muell. Arg. l. c.* 147.—Mexic.
lanceolata, *Muell. Arg. l. c.* 148.—Calif.
lanciflola, *Muell. Arg. l. c.* 146.—Ind. occ.
lineata, *Baill. Adansonia*, iv. (1863–64) 288.—Bras.
melochiaefiora, *Muell. Arg. in DC. Prod.* xv. ii. 736.—Bras.
mercurialina, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 148.—Texas.
mollis, *Baill. Adansonia*, iv. (1863–64) 288.—Bras.
montevidensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 147.—Reg. Argent.
neo-mexicana, *Muell. Arg. l. c.*—N. Mexic.
pascalioides, *Mirb. ex Steud. Nom.* ed. II. i. 128.—Am. austr.
Pavoniana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 149.—Peruv.
pilosissima, *Muell. Arg. l. c.* 148.—Texas.
quinquecuspidata, *Muell. Arg. l. c.* 150.—Peruv.
Ruiziana, *Muell. Arg. l. c.* 151.—Peruv.
Schiedeana, *Muell. Arg. l. c.* 150.—Mexic.
serica, *Griseb. Fl. Brit. W. Ind.* 44.—Ind. occ.
sericophylla, *A. Gray, in Bot. Calif.* ii. 70.—Calif.
serrata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 147.—Calif.
Sinensis, *Muell. Arg. l. c.* 148.—Sina.
spaciata, *Muell. Arg. l. c.* 148.—Peruv.
tricocca, *Muell. Arg. l. c.* 150.—Bras.
trisuspidata, *Muell. Arg. l. c.*—Calif.
tuberculata, *Muell. Arg. l. c.* 148.—China.

ARGOLASIA, Juss. Gen. 60 (1789) = **Lazaria**, Ait. (Haemodor.).
aurea, Paxt., Fl. Am. Sept. i. 224, in syn. = Lophiola aurea.
caperata, J. F. Gmel. Syst. ii. 560 = L. plumosa.
lanata, Lam. ex Poir. Encyc. Suppl. i. 448 = L. plumosa.
plumosa, Juss. Gen. ed. Usteri, 67 = L. plumosa.

ARGOPHILUM, Blanco, Fl. Filip. 186 (1837) = **Aglaia**, Lour. (Meliac.).
pinnatum, *Blanco, l. c.* 186 = Aglaia angustifolia.

ARGOPHYLLUM, Blanco, Fl. Filip. p. lxv (1837) = prec.

ARGOPHYLLUM, Forst. Char. Gen. 29. t. 15 (1776).
SAXIFRAGEAE, Benth. & Hook. f. i. 646.
elliptícum, *Labill. Sert. Austr. Caled.* 39. t. 41.—N. Caled.
Lejourdani, *F. Muell. Fragm.* iv. 33.—Austral.
nitídum, *Forst. Char. Gen.* 30.—N. Caled.
sericeum, Poir. Encyc. Suppl. i. 449 = nitidum.

ARGORIPS, Rafin. Alsog. Am. 13 (1838) = **Salix**, Tourn.
alba, Rafin. l. c. = S. alba.
cerulea, Rafin. l. c. = S. alba.

ARGOSTEMMA, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 324 (1824). *RUBIACEAE*, Benth. & Hook. f. ii. 54.
POMANGIUM, Reinw. in Syll. Ratisb. ii. (1828) 10.
acutum, Wall. ex Hook. f. Fl. Brit. Ind. iii. 43 = verticillatum.
angustifólium, *Miq. Fl. Ind. Bat.* ii. 168.—Malaya.
begoniaeforme, *Miq. l. c.* 848.—Malaya.
borragineum, *Blume, in DC. Prod.* iv. 417.—Java.
calycinum, R. Br. ex Benn. Pl. Jav. Rar. i. 97 = Neurocalyx Wightii.
coenosciadicum, *Suring. in Arch. Néerl.* v. (1870) 116.—Java.
connatum, Dals. in Hook. Kew Journ. iii. (1851) 345 = courtallense.
courtallense, *Arn. in Ann. Nat. Hist.* iii. (1839) 22.—Ind. or.
Elatostemma, *Hook. f. Fl. Brit. Ind.* iii. 45.—Malaya.
fulvum, *Zoll. & Moritzi, Syst. Verz. Zoll.* 61.—Java.
glabérrimum, Dals. in Hook. Kew Journ. iii. (1851) 345 = verticillatum.
humile, *Benn. Pl. Jav. Rar.* i. 94.—Reg. Himal.
inaequale, *Benn. l. c.* 96.—Sumatra.
inaequilaterum, *Benn. l. c.* 95.—Java.
Junghuhnii, *Miq. Fl. Ind. Bat.* ii. 164.—Malaya.
khasianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 43.—Reg. Himal.
Korthalsii, *Miq. Fl. Ind. Bat.* ii. 165.—Malaya.
Kurzii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 45.—Burma.
laeve, *Benn. Pl. Jav. Rar.* i. 96.—Java.
lasvigatum, *Benn. l. c.*—Java.
Lobbii, *Hook. f. Fl. Brit. Ind.* iii. 44.—Burma.
longifólium, *Benn. Pl. Jav. Rar.* i. 97.—Sumatra.
macrocarpum, *Miq. Fl. Ind. Bat.* ii. 849.—Malaya.
microcnide, *Miq. l. c. Suppl.* 589.—Sumatra.
montanum, *Blume, in DC. Prod.* iv. 418.—Java.
Neesianum, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) [Pl. *Meyen.*] 849.—Ins. Philipp.
neurocalyx, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 229.—Malaya.
oligantha, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 181.—Ins. Andam.
ophirense, *Maing. ex Hook. f. Fl. Brit. Ind.* iii. 45.—Malaya.
Parishii, *Hook. f. l. c.* 44.—Burma.
parvifólium, *Benn. Pl. Jav. Rar.* i. 96.—Malaya.
pascuiflorum, *Blume, in DC. Prod.* iv. 418.—Java.
pedunculosum, *Miq. Fl. Ind. Bat.* ii. 849.—Malaya.
pictum, *Korth. in Noder. Kruid. Arch.* ii. ii. (1851) 153.—Sumatra.
pictum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 327.—Penang.
pulchrum, *Korth. in Noder. Kruid. Arch.* ii. ii. (1851) 152.—Sumatra.
pumilum, *Benn. Pl. Jav. Rar.* i. 96.—Ind. or.; Afr. trop.
rostratum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 326.—Reg. Himal.
sarmentosum, *Wall.* i. c. 324.—Reg. Himal.
sonerilóides, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 210.—Ins. Andaman.
spinulosum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 46.—Singapore.
subinaequale, *Benn. Pl. Jav. Rar.* i. 96.—Java.
tavoyanum, *Wall. ex Benn. l. c.* 94.—Burma.
Teysmanianum, *Miq. Fl. Ind. Bat.* ii. 849.—Java.
timorense, *Benn. Pl. Jav. Rar.* i. 95.—Timor.
uniflorum, *Blume, in DC. Prod.* iv. 418.—Java.
unifolium, *Benn. Pl. Jav. Rar.* i. 94.—Penang.
unifolium, Kurz, in Journ. As. Soc. xlvi. (1877) ii. 131 = Kurzii.
verticillatum, Wall. Cat. n. 8394, partim = khasianum.
verticillatum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 325.—Ind. or.
Wallichii, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. 350 (1843) [Pl. Meyen.].—Ins. Philipp.

ARGOTHAMNIA, Spreng. Syst. iii. 847 (1826) = **Argithamnia**, Sw. (Euphorb.).

ARGUZIA, Anon. ex Steud. Nom. ed. I. 68 (1821); Stev. in Bull. Soc. Nat. Mosc. (1851) i. 558 = **Tournefortia**, Linn. (Borag.).
cymmerica, Stev. l. c. 560 = T. sibirica.
Mezerschmidia, Stev. l. c. = T. sibirica.
sarmentosa, Steud. Nom. ed. I. 68 = T. Argusia.
repens, Rafin. Sylva Tellur. 167.—Sibir.?
rosmarinifolia, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 559 = T. sibirica.

ARGYLIA, D. Don, in Edinb. Phil. Journ. ix. (1823) 260. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1043.
adscendens, *DC. Prod.* ix. 225.—Chili.
australis, *Phil. in Anal. Univ. Chil.* (1873) 512.—Chili.
Bustillosii, *Phil. in Linnaea*, xxix. (1857–58) 13.—Chili.
canescens, *D. Don, in Edinb. Phil. Journ.* (*Apr.-Yune* 1823) 258.—Chili.
chrysantha, *Phil. in Linnaea*, xxxiii. (1864–65) 180.—Chili.
cremophila, *Phil. l. c.*—Chili.
Feuillei, *DC. Prod.* ix. 225.—Chili.
geranioides, *DC. l. c.*—Chili.
glabriuscula, *Phil. in Linnaea*, xxxiii. (1864–65) 179.—Chili.
glutinosa, *Phil. Fl. Atac.* 36.—Chili.
huidobriana, *Clos, in C. Gay, Fl. Chil.* iv. 411.—Chili.
incana, *Phil. Fl. Atac.* 36.—Chili.
lutea, *Phil. in Anal. Univ. Chil.* (1873) 511.—Chili.
potentillaefolia, *DC. Prod.* ix. 225.—Chili.
puberula, *DC. l. c.*—Chili.
radiata, *D. Don, in Edinb. Phil. Journ.* ix. (1823) 260.—Peruv.
radiata, Endl. ic. n. 1112 = canescens.
tomentosa, *Phil. Fl. Atac.* 36.—Chili.
trifoliata, DC. Prod. ix. 227 = aspallatensis.
aspallatensis, *DC. l. c.* 226.—Chili.
villosa, *Phil. in Anal. Univ. Chil.* (1873) 512.—Chili.
viridis, *Phil. in Linnaea*, xxxiii. (1864–65) 181.—Chili.

ARGYRA, Noronha, ex Baill. Adansonia, i. (1860–61) 169 = **Croton**, Linn. (Euphorb.).
fasciculata, Noronha, ex Baill. l. c. = C. Noronhae.

ARGYRANTHEMUM, Webb, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 258 (1836–50) = **Chrysanthemum**, Linn. (Compos.).
anethifolium, Webb, l. c. 265–267 = C. anethifolium.
coronopifolium, Webb, l. c. 272 = C. coronopifolium.
crithmifolium, C. Bolle, Ind. Sem. Hort. Berol. App. (1861) 5 = C. anethifolium.
dissectum, Lowe, Man. Fl. Mad. i. 464 = C. dissectum.
flaviculaceum, Webb, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 268 = C. anethifolium.
frutescens, Sch. Bip. l. c. 264 = C. frutescens.
gracile, Sch. Bip. l. c. 261 = C. gracile.
haematomma, Lowe, Man. Fl. Mad. i. 463 = C. haematomma.
jacobaeifolium, Webb, in Webb & Berth. Phyt. Canar. ii. 251.—Ins. Canar.
ochroleucum, Webb, l. c. 259 = C. ochroleucum.
ARGYRANTHEMUM:—
pinnatifidum, Webb, in Webb & Berth. Phyt. Canar. ii. 269 = C. pinnatifidum.
Webbii, Sch. Bip. in Webb & Berth. l. c. = C. Webbii.

ARGYRANTHUS, Neck. Elem. i. 75 (1790) = **Heliopterum**, DC. (Compos.).

ARGYREIA, Lour. Fl. Cochinch. 134 (1790). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 869.
ADAMMIA, Rafin. Fl. Tellur. iv. 73 (1836).
ARGYRON, St. Lag. in Ann. Soc. Bot. Lyon, vi. (1880) 120.
SAMUDRA, Rafin. Fl. Tellur. iv. 72 (1836).
abyssínica, *Choisy, in DC. Prod.* ix. 382.—Abyss.
acuta, Benth. Fl. Hongk. 237, in syn. = Championi.
acuta, *Lour. Fl. Cochinch.* 135.—China; Ind. or.
aggregata, Arn. ex Choisy, Conv. Or. 45 = Lettsomia aggregata.
alata, *Montrous. in Mém. Acad. Lyon*, x. (1860) 286.—N. Caled.
alulata, Miq. Fl. Ind. Bat. ii. 587 = Ipomoea Turpethum.
ampla, Choisy, Conv. Or. 38 = Roxburghii.
arbórea, *Lour. Fl. Cochinch.* 135.—China.
argentata, *Miq. Fl. Ind. Bat.* ii. 588.—Malaya.
argentea, *Arn. ex Choisy, Convo. Or.* 36.—Ind. or.
barbigera, Choisy, l. c. 42 = Lettsomia barbigera.
Bona-nox, Sweet, Hort. Brit. ed. I. 289 = Rivea hypocrateriformis.
bractesta, *Choisy, Convo. Or.* 39.—Ind. or.
capitata, Arn. ex Choisy, l. c. 41 = Lettsomia strigosa.
capitata, Brand. For. Fl. 343 = Lettsomia Thomsoni.
capitata, Kurz, For. Fl. Burm. ii. 216 = Lettsomia pegunesis.
Championi, Benth. Fl. Hongk. 236 = Lettsomia Championi.
Cholsyana, *Regel & Korrn. in Ind. Sem. Hort. Petrop.* (1858) 40.—Hab.?
Cholsyana, *Wight, ex Hook. f. in Fl. Brit. Ind.* iv. 190.—Ind. or.
courtallensis, Wight, ex Choisy, in DC. Prod. ix. 329 = malabarica.
cuneata, *Ker-Gawl. in Bot. Reg.* t. 661.—Ind. or.
cymosa, *Sweet, Hort. Brit.* ed. I. 289.—Ind. or.; Malaya.
Daltoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 190.—Ind. or.
elliptica, Arn. ex Choisy, Conv. Or. 35 = Lettsomia elliptica.
festiva, Wall. Pl. As. Rar. 68. t. 76.—China.
floribunda, *Wall. ex Voigt, Hort. Suburb. Calc.* 352, *nomen.*—Ind. or.
fulgens, *Choisy, Convo. Or.* 33.—Ind. or.
glabra, *Choisy, ex Zoll. Syst. Verz. Ind. Archip.* 128, 180.—Malaya.
Gulcheenii, *Choisy, Convo. Or.* 41.—Malaya.
hancorniaefolia, Gardn. ex Thw. Enum. Pl. Zeyl. 210 = Lettsomia hancorniaefolia.
hirsuta, *Wight & Arn. in Nov. Act. Nat. Cur.* xviii. (1836) i. 856.—Ind. or.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 186.—Reg. Himal.
involucrata, *C. B. Clarke*, l. c. 187.—Ind. or.
lanceolata, *Choisy, Convo. Or.* 39.—Burma; Malaya.
laurifolia, Voigt, Hort. Suburb. Calc. 351 = Lettsomia elliptica.
Lawii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 190.—Ind. or.
Leschenaultii, *Choisy, Convo. Or.* 31.—Ind. or.
Leschenaultii, Thw. Enum. Pl. Zeyl. 209 = serices.
malabarica, *Arn. ex Choisy, Convo. Or.* 38.—Ind. or.; Malaya.
megapotamica, *Griseb. in Gottt. Abh.* xxiv. (1879) 263.—Reg. Argent.
mollis, *Choisy, Convo. Or.* 38.—Ind. or.; Malaya.
multiflora, Voigt, Hort. Suburb. Calc. 351 = Roxburghii.
nellygherya, *Choisy, Convo. Or.* 32.—Ind. or.
nervosa, *Bej. Hort. Mauril.* 234.—Ins. Mascar.
nitida, *Choisy, Convo. Or.* 40.—Ind. or.; Ins. Philip.
oblonga, *Benth. Bot. Voy. Sulph.* 138.—Mexic.
obtecta, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 186.—Burma; Malaya.
obtusifolia, Lour. Fl. Cochinch. 135 = obtecta.
ornata, Sweet, Hort. Brit. ed. I. 289 = Rivea ornata.

ARGYREIA:—
argyrensis, Choisy, in DC. Prod. ix. 334 = Lettsomia aggregata.
pallida, *Choisy*, *Conv. Or.* 34.—Burma.
pilosa, *Wight & Arn. in Nov. Act. Nat. Cur.* xviii. (1836) i. 356.—Ind. or.
pomacea, *Choisy*, *Conv. Or.* 32.—Ind. or.
populifolia, *Choisy*, *l. c.*—Zeylan.
populifolia, Choisy, in DC. Prod. ix. 339 = Hookeri. Reinwardtiana, *Miq. Fl. Ind. Bat.* ii. 589.—Malaya.
Roxburghii, *Arn. ex Choisy*, *Conv. Or.* 37.—Ind. or.
rubicunda, Choisy, l. c. 44 = Lettsomia rubicunda.
sericea, *Dals. & Gib. Bomb. Fl.* 169.—Ind. or.
sericea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = argentea.
setosa, Arn. ex Choisy, Conv. Or. 43 = Lettsomia setosa.
speciosa, *Roj. Hort. Maurit.* 224.—Maurit.
speciosa, *Sweet*, *Hort. Brit.* ed. I. 389.—Ind. or.; Malaya.
splendens, *Sweet*, *l. c.*—Ind. or.
tiliaefolia. *Wight*, *Ic.* iv. t. 13. t. 1358 (sphalm. *tiliaefolia*). Ind. or.
tomentosa, *Choisy*, *Conv. Or.* 44.—Burma.
uniflora, Sweet, Hort. Brit. ed. I. 389 = Rivea hypocrateriformis.
vernusta, Choisy, Conv. Or. 37 = argentea.
Wallichii, *Choisy*, *l. c.* 39.—Ind. or.
zeylanica, Kurz, For. Fl. Brit. Burma, ii. 215, excl. syn. = Lettsomia Kurzii.
zeylanica, Voigt, Hort. Suburb. Calc. 351 = pomacea.

ARGYRELLA, Nsud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 300. t. 6 = **Diastoïa**, Benth. (Melast.).
canescens, Harv. Gen. S. Afr. Fl. ed. II. 118 = D. incana.
incana, Nsud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 300. t. 6. f. 7 = D. incana.
phaeotricha, Nsud. l. c. = D. phaeotricha.

ARGYREON, St. Lag. in Ann. Soc. Bot. Lyon, vii. 120 (1880) = **Argyreia**, Lour. (Convolv.).

ARGYREXIAS, Rafin. Fl. Tellur. iv. 86 (1836) = **Bohlum**, Tourn. (Borag.).
candicans, Rafin, l. c. = E. candicans.

ARGYROCHAETA, Cav. Ic. iv. 54. t. 378 (1791) = **Parthenium**, Linn. (Compos.).
bipinnatifida, Cav. l. c. = P. Hysterophorum.

ARGYROCOMA, Rafin. New Fl. Am. iv. 41 (1836) = **Anychia**, Rich. (Illecebr.).
imbricata, Rafin. l. c. 43 = Anychia dichotoma.

ARGYROCOME, Gaertn. Fruct. ii. 410 (1791); Schrank, in Denkschr. Acad. Wiss. Muench. vii. (1824) 146 et 165 = **Helipterum**, DC. (Compos.).
apiculata, Cass. in Dict. Sc. Nat. xxxix. 40 = Helichrysum apiculatum.
bracteata, Cass. l. c. = Helichrysum rutilans!
cricoides, Poir. Tabl. Encyc. iii. 270 = Helichrysum cricoides.
ferruginea, Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 162 = Helip. speciosissimum.
imbricata, Sack, Ind. DC. Prod. ii. 15 = Metalasia unifora.
retorta, Cass. in Dict. Sc. Nat. xxxiv. 39 = Helichrysum stoloniferum.
retorta, Gaertn. Fruct. ii. t. 167 = Helichrysum retortum.
seminuda, Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 164 = Helip. speciosissimum.
semipapposum, Cass. in Dict. Sc. Nat. xxxix. 40 = Helichrysum semipapposum.
Staehelina, Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 165 = Helichrysum squamosum.
vermiculata, Poir. Tabl. Encyc. iii. 270 = Metalasia unifora.
vulnerata, Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 164 = H. canescens.

ARGYRODENDRON, Klotzsch, in Peters, Reise Mossamb. Bot. 100 (1861) = **Croton**, Linn. (Euphorb.).
bicolor, Klotzsch, l. c. 102 = C. pseudolinum.
croatum, Boiv. ex Baill. Adansonia, i. (1861) 152 = C. Bernieri.
Peterzii, Klotzsch, in Peters, Reise Mossamb. Bot. i. 101 (= Combreti sp.).—Afr. trop.

ARGYRODENDRON, F. Muell. Fragm. i. 2 (1858) = **Tarristia**, Blume (Stercul.).
trifoliolatum, F. Muell. l. c. = T. Argyrodendron.

ARGYROGLOTTIS, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 88 = **Hallchrysum**, Vaill. (Compos.).
turbinata, Turcz. l. c. 84 = H. Argyroglottis.

ARGYROLOBIUM, Eckl. & Zeyh. Enum. 184 (1835). *LEGUMINOSAE*, Benth. & Hook. f. i. 480.
CHAMARCTHUS, Vis. Fl. Dalm. 272. t. 55 (1850).
CHASMONE, E. Mey. Comm. Pl. Afr. Austr. 71 (1836).
DIOLOTUS, Tausch, in Flora, xxv. (1842) 284.
GAMOCHILUM, Walp in Linnaea, xiii. (1839) 509.
TEPHROTHAMNUS, Sweet, Hort. Brit. ed. II. 126 (1830), nomen prius.
TRICHASMA, Walp. in Linnaea, xiii. (1839) 510.
abyssinicum, *Jaub. & Spach*, *Illustr.* i. 115.—Abyss.
aequinoctiale, *Welw. ex Baker*, *in Oliver*, *Fl. Trop. Afr.* ii. 46.—Afr. trop.
Andrewsianum, *Steud. Nom.* ed. II. i. 129.—Afr. austr.
angustifolium, *Eckl. & Zeyh. Enum.* 188.—Afr. austr.
arabicum, *Jaub. & Spach*, *Illustr.* i. 115; *Boiss. Diagn.* Ser. I. ii. 18.—Arab.
argenteum, *Eckl. & Zeyh. Enum.* 184.—Afr. austr.
argentenm, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 464 = Linnaeanum.
ascendens, *Walp. in Linnaea*, xiii. (1839) 507.—Afr. austr.
baptisioides, *Walp. l. c.* 506.—Afr. austr.
barbatum, *Walp. Rep.* ii. 845.—Afr. austr.
biflorum, *Eckl & Zeyh. Enum.* 186.—Afr. austr.
Bottae, *Jaub. & Spach*, *Illustr.* i. 115.—Arab.
calycinum, Bell, in Journ. Linn. Soc. xvi. (1878) 896, in obs. = Linnaeanum.
calycinum, *Jaub. & Spach*, *Illustr.* i. 115.—Dalmat.; Reg. Caucas. Persia.
candicans, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
collinum, *Eckl. & Zeyh. l. c.*—Afr. austr.
conanthum, Vog. ex Steud. Nom. ed. II. i. 129 = baptisioides.
connatum, *Harv. in Harv. & Sond. Fl. Cap.* 72.—Afr. austr.
crassifolium, *Eckl. & Zeyh. Enum.* 184.—Afr. austr.
crinitum, *Walp.in Linnaea*, xiii. (1839) 506.—Afr austr.
crotalarioides, *Jaub. & Spach*, *Illustr.* i. 114, 116.—Syria.
dasycarpum, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 46 = A. Schimperianum.
divaricatum, Jaub. & Spach, Illustr. i. 116 = flaccidum.
dorycnioides, *Baker*, *in Oliver*, *Fl. Trop. Afr.* ii. 45.—Afr. trop.
emirnense, *Baker*, *in Journ. Linn. Soc.* xx. (1883) 125. —Madag.
faltax, Bail, in Journ. Bot. xi. (1873) 302 = Linnaeanum.
filiforme, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
flaccidum, *Jaub. & Spach*, *Illustr.* i. 115.—Ind. or. goodioides, *Walp. Rep.* ii. 845.—Afr. austr.
gracile, Fenzl, in Flora, xxvi. (1843) 1. 394 = crotalaroides.
grandiflorum, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 80.—Afr. bor.
incanum, Eckl. & Zeyh. Enum. 185 = crassifolium.
involucratum, *Harv. in Harv. & Sond. Fl. Cap.* 75.—Afr. austr.
Kotschyi, *Boiss. Diagn.* Ser. I. vi. 32.—Persia.
Krebsianum, *Presl*, *Bot. Bemerk.* 139.—Afr. austr.
lanceolatum, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
Linnaeanum, *Walp.in Linnaea*, xiii. (1839) 508.—Reg. Mediterr.
longifolium, *Walp. Rep.* ii. 844.—Afr. austr.
lotoides, *Bunge*, *ex Trautv. in Act. Hort. Petrop.* ii. (1873) 519.—Armenia.
lotoides, *Harv. in Harv. & Sond. Fl. Cap.* 586.—Afr. austr.
microphyllum, *Ball*, *in Journ. Bot.* xi. (1873) 302.—Marocco.
modestum, Hochst. in Flora, xxvii. (1844) 1. 100 = abyssinicum.
molle, *Eckl. & Zeyh. Enum.* 187.—Afr. austr.
oboordatum, *Steud. Nom.* ed. II. i. 130.—Afr. austr.
obovatum, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
obsoletum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 70.—Afr. austr.
ornithopodioides, Jaub. & Spach, Illustr. i. 116 = roseum.
patens, *Eckl. & Zeyh. Enum.* 185.—Afr. austr.

ARGYROLOBIUM:—
pauciflorum, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
pechuelianum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 346.—Madag.
petiolare, *Steud. Nom.* ed. II. i. 130.—Afr. austr.
Petitianum, *A. Rich. Tent. Fl. Abyss.* i. 168.—Abyss.
pilosum, *Harv. in Harv. & Sond. Fl. Cap.* 71.—Afr. austr.
polyphyllum, *Eckl. & Zeyh. Enum.* 184.—Afr. austr.
prostratum, *Boiss. Diagn.* Ser. I. ii. 18.—Assyria.
pumilum, *Eckl. & Zeyh. Enum.* 185.—Afr. austr.
ramosissimum, *Baker*, *in Oliver*, *Fl. Trop. Afr.* ii. 46. —Afr. trop.
remotum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 162. —Abyss.
roseum, *Jaub. & Spach*, *Illustr.* i. 116.—Persia; Beluchist.
rupestre, *Walp. in Linnaea*, xiii. (1839) 508.—Afr. austr.
Saharae, *Pomel*, *Nouv. Mat. Fl. Atl.* 177.—Algeria.
Sanderroni, *Harv. in Harv. & Sond. Fl. Cap.* ii. 594.—Afr. austr.
Schimperianum, *Hochst. in Flora*, xxvii. (1844) 1. 100. —Afr. trop.
sericeum, *Eckl. & Zeyh. Enum.* 184.—Afr. austr.
sessilifolium, *Jaubu, in Oestr. Bot. Zeitschr.* xxii. (1872) 175.—Turcia.
speciosum, *Eckl. & Zeyh. Enum.* 187.—Afr. austr.
splendens. *Walp. Rep.* ii. 845.—Afr. austr.
stenophyllum, *Boiss. Fl. Orient.* ii. 32.—Afghan.
stipulaceum, *Ball*, *in Journ. Bot.* xi. (1873) 302.—Marocco.
stipulaceum, *Eckl. & Zeyh. Enum.* 187.—Afr. austr.
strictum, *Steud. Nom.* ed. II. i. 130.—Afr. austr.
Sutherlandi, *Harv. in Harv. & Sond. Fl. Cap.* 594.—Afr. austr.
syriacum, Boiss. Diagn. Ser. I. ii. 18 = crotalarioides.
tenue, *Walp. Rep.* i. 502.—Afr. austr.
trigonelloides, *Jaub. & Spach*, *Illustr.* i. 116, 118.—Persia.
tuberosum, *Eckl. & Zeyh. Enum.* 188.—Afr. austr.
umbellatum, *Vog. ex Steud. Nom.* ed. II. i. 130.—Afr. austr.
uniforum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 72.—Afr. austr.
uniflorum, *Jaub. & Spach*, *Illustr.* i. 115.—Oriens.
velutinum, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
venustum, *Eckl. & Zeyh. l. c.* 185.—Afr. austr.
virgatum, *Baker*, *in Oliver*, *Fl. Trop. Afr.* ii. 46.—Afr. trop.

ARGYROPHANES, Schlecht. in Linnaea, xx. (1847) 596 = **Helichrysum**, Vaill. (Compos.).
Behrii, Schlecht. l. c. = H. Baxteri.

ARGYROPHYLLUM, Pohl, ex Baker, in Mart. Fl. Bras. vi. ii. 150 (1873) = **Soaresia**, Sch. Bip. (Compos.).
ovali-ellipticum, Pohl, ex Baker, l. c. = S. velutina.

ARGYROPHYTON, Hook. Comp. Bot. Mag. ii. (1836) 168 = **Argyroxiphium**, DC. (Compos.).
Douglasii, Hook. l. c. = Argyroxiph. sandwicense.

ARGYROPSIS, M. Roem. Syn. Ecnat. 125 (1847) = **Euphrantthes**, Horl. (Amaryll.).
candida, M. Roem. l. c. = Z. candida.

ARGYRORCHIS, Blume, Orch. Archip. Ind. 120. t. 31, 36 (1858). *ORCHIDAE*, Benth. & Hook. f. iii. 506.
javanica, *Blume*, *l. c.*—Java.

ARGYROTHAMNIA, Muell. Arg. in Linnaea, xxxiv. (1865) 144 = **Argithamnia**, Sw. (Euphorb.).

ARGYROXIPHIUM, DC. Prod. v. 668; et Mém. Comp. t. 9 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 393.
ARGYROPHYTON, Hook. Comp. Bot. Mag. ii. (1836) 168.
macrocephalum, *A. Gray*, *in Proc. Am. Acad.* ii. (1852) 190.—Ins. Sandvic.
sandwicense, *DC. Prod.* v. 668.—Ins. Sandvic.

ARGYTAMNIA, Duchesne, in Dict. Sc. Nat. iv. 86 (1804) = **Argithamnia**, Sw. (Euphorb.).

ARGYTHAMNIA, P. Br. Hist. Jamaic. 286 (1756, sphalm. = **Argithamnia**, Sw. (Euphorb.).

ARHYNCHIUM, Lindl. & Paxt. Flow. Gard. i. (1850–51) 142 = **Arachnanthe**, Blume (Orchid.).

ARIA, Jacq. f. Cat. Hort. Vindob. (1826) ex Host, Fl. Austr. ii. 7 (1831) = **Pyrus**, Linn. (Rosac.).
alnifolia, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 166 = P. alnifolia.
alpina, M. J. Roem. Syn. iii. Rosif. 124 = P. alpina.
ambigua, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 165.—Hab. ?
angusta, M. J. Roem. Syn. iii. Rosif. 129 = P. intermedia.
betulaefolia, M. J. Roem. l. c. 124 = P. betulaefolia.
Chamaemespilus, Host, Fl. Austr. ii. 8 = P. Chamaemespilus.
crenata, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 162 = P. crenata.
cuspidata, M. J. Roem. Syn. iii. Rosif. 125 = P. cuspidata.
Decaisneana, Lavall. Arbor. Segrez. 61.—Hab. ?
decipiens, M. J. Roem. Syn. iii. Rosif. 129 = P. rotundifolia.
edulis, Guimp. ex Lavall. Arbor. Segrez. 63 = *Decaisneana*.
edulis, M. J. Roem. Syn. iii. Rosif. 124 = P. edulia.
flabellifolia, M. J. Roem. l. c. 127 = Crataegus corymbosa.
graeca, M. J. Roem. l. c. = P. Aria.
Hostii, Carr. Rev. Hortic. (1877) 210 = P. Hostii.
Hostii, Jacq. f. Cat. Hort. Vindob. (1826) ex Host, Fl. Austr. ii. 8 = P. intermedia.
intermedia, Schur, Enum. Pl. Transs. 207 = P. intermedia.
japonica, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 164 = P. japonica.
kamaonensis, M. J. Roem. Syn. iii. Rosif. 130 = P. lanata.
humaonensis, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 162 = praec.
lanata, Decne. l. c. 163 = P. lanata.
latifolia, M. J. Roem. Syn. iii. Rosif. 128 = P. rotundifolia.
Mougeoti, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 378 = P. Aria.
nivea, Host, Fl. Austr. ii. 8 = P. Aria.
obtusata, M. J. Roem. Syn. iii. Rosif. 125 = P. obtusata.
rotundifolia, M. J. Roem. l. c. 129 = P. rotundifolia.
scandica, M. J. Roem. l. c. 127 = P. intermedia.
semilobata, M. J. Roem. l. c. 129 = P. rotundifolia.
Sorottii, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 165.—Hab. ?
tiliaefolia, Decne. l. c. 166 = P. alnifolia.
vestita, M. J. Roem. Syn. iii. Rosif. 125 = P. vestita.

ARIETINUM, Beck, Bot. N. and Midl. U. St. 352 (1838) = **Cypripedium**, Linn. (Orchid.).
americanum, Beck, l. c. = C. Arietinum.

ARILLARIA, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 70 = **Ormosia**, Jack (Legum.).
robusta, Kurz, l. c. 71 = O. robusta.

ARILLASTRUM, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. xiv. (1867) 258 = **Spermolepis**, Brongn. & Gris (Myrtac.).
gummiferum, Panch. l. c. = S. gummiferum.

ARINEMIA, Rafin. Sylva Tellur. 49 (1838) = **Ilex**, Tourn.
lanceolata, Rafin. l. c. = I. lanceolata.

ARIOCARPUS, Scheidw. in Bull. Acad. Brux. v. (1838) 491 = **Mammillaria**, Haw. (Cactac.).
retusus, Scheidw. l. c. 492. t. 1 = M. prismatica.

ARIODENDRON, Meisn. Gen. Comm. 301 (1836–43), sphalm. = AGRIODENDRON, Endl. = **Aloë**, Linn. (Liliac.).

ARIONA, Pers. Syn. i. 212 (1805) = **Arjona**, Comm. (Santal.).

ARIOPSIS, J. Grah. Cat. Bomb. Pl. 253 (1839).
AROIDEAE, Benth. & Hook. f. iii. 973.
peltata, *J. Grah. l. c.*—Exp. Himal.
protensiflora, N. E. Br. in Rep. Roy. Gard. Kew, (1877) 83 = peltata.

ARISACONTIS, Schott, in Bonplandia, v. (1857) 129. *AROIDEAE*? Benth. & Hook. f. iii. 963.
Chamisonis, *Schott, l. c.*—Ins. Radack.

ARISAEMA, Mart. in Flora, xiv. (1831) ii. 459.
AROIDEAE, Benth. & Hook. f. iii. 966.
AMIDENA, Rafin. Fl. Tellur. iv. 15 (1836).
abbreviatum, *Schott, in Oestr. Bot. Zeitschr.* (1857) 382.—Reg. Himal.
affine, Schott, in Bonplandia, vii. (1859) 27 = concinnum.
album, *N. E. Br. in Journ. Linn. Soc.* xviii. (1882) 247.—Siam.
alismatum, Schott, in Bonplandia, vii. (1859) 26 = concinnum.
amplissimum, *Blume, Rumphia*, i. 110.—Japon.
amurense, *Maxim. Prim. Fl. Amur.* 264.—Reg. Amur.
anguustum, *Franch. & Sav. Enum. Pl. Jap.* ii. 507.—Japon.
atrorubens, *Blume, Rumphia*, i. 97.—Am. bor.
Bucci, Blume, l. c. 104 = A. Dracontium.
Bottae, *Schott, Prod. Aroid.* 42.—Arabia.
brasilianum, Blume, Rumphia, i. 97 = atrorubens.
candatum, *Engl. in DC. Monog. Phan.* ii. 559.—Ind. or.
cochinchinense, Blume, Rumphia, i. 107 = Pinella Wawrae.
commutatum, Schott, in Bonplandia, vii. (1859) 26 = tortuosum.
concinnum, *Schott, l. c.* 27.—Reg Himal.
consanguineum, Schott, l. c. = crubescens.
comutum, *Schott, l. c.*—Reg. Himal.
costatum, *Mart. in Flora*, xiv. (1831) ii. 459.—Reg. Himal.
Camingii, *Schott, Syn. Aroid.* 28.—Ins Philipp.
curvatum, Hook. Bot. Mag. t. 5981 = tortuosum.
curvatum, *Kunth, Enum. Pl.* iii. 20.—Reg. Himal.
cuspidatum, *Engl. in DC. Monog. Phan.* ii. 586.—Penang.
decipiens, *Schott, in Oestr. Bot. Zeitschr.* (1857) 378.—Java.
dolosum, Schott, in Bonplandia, vii. (1859) 26 = interruedium.
Dracontium, *Schott, Meletem.* i. 17.—Am. bor.
echinatum, *Schott, l. c.*—Reg. Himal.
eminens, Schott, in Oestr. Bot. Zeitschr. (1857) 357 = speciosum.
enneaphyllum, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 352.—Abyss.
crubescens, *Schott, Meletem.* i. 17.—Reg. Himal.
exile, *Schott, in Bonplandia*, vii. (1859) 26.—Reg. Himal.
fallax, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 123 = filiforme.
filicaudatum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1882) 258.—Zeylan.
filiforme, *Blume, Rumphia*, i. 102. t. 28.—Java.
filiforme, Thw. Enum. Pl. Zeyl. 334 = neglectum.
fimbriatum, *Mast. in Gard. Chron.* (1884) ii. 680.—Ins. Philipp.
flavum, *Schott, Prod. Aroid.* 40.—Arabia.
Franchetianum, *Engl. Bot. Jahrb.* i. (1881) 487.—Tibet.
Griffithii, Schott, in Bonplandia, vii. (1859) 26 = Leschenaultii.
galeatum, *N. E. Br. in Gard. Chron.* (1879) ii. 102.—Reg. Himal.
gracile, Kunth, Enum. Pl. iii. 21 = Typhonium gracile.
Griffithii, *Schott, Syn. Aroid.* 26.—Reg. Himal.
hastatum, Blume, Rumphia, i. 98 = atrorubens.
heliboriifolium, Schott, Syn. Aroid. 29 = tortuosum.
heptaphyllum, *Blume, Rumphia*, i. 109.—Ind. or.
heterophyllum, Blume, l. c. 110 = Thunbergii.
Hookerianum, Schott, in Oestr. Bot. Wochenbl. (1857) 334 = Griffithii.
Hookeri, Schott, Gen. Aroid. f. 11–19 = Griffithii.
Huegelii, Schott, Syn. Aroid. 27 = Leschenaultii.
inclusum, *N. E. Br. in Journ. Linn. Soc.* xvii. (1881) 249.—Java.
intermedium, *Blume, Rumphia*, i. 102.—Reg. Himal.
Inquemenii, *Blume, l. c.* 95.—Reg. Himal.
japonicum, *Blume, l. c.* 108.—Japon.
laminatum, Benth. Fl. Hongk. 343 = penicillatum.
laminatum, *Blume, Rumphia*, i. 99. t. 27.—China.
latisectum, Blume, l. c. 110 = japonicum.
Leschenaultii, *Blume, l. c.* 98.—Ind. or.
lobatum, *Engl. Bot. Jahrb.* i. (1881) 487.—Tibet.
Lowerii, Blume, Rumphia, i. 108 = Pinella tuberifera.
macrospathum, *Benth. Pl. Hartw.* 62.—Mexic.

ARISAEMA :—
macrourum, Kunth, Enum. Pl iii. 644 = Pinellia tuberifera.
Mahoyanum, Kunth, in Ind. Sem. Hort. Berol. (1845) 9 = filiforme.
mirabile, Schott, in Oestr. Bot. Zeitschr. (1857) 866 = speciosum.
Murrayi, Hook. Bot. Mag. t. 4388 = Jacquemontii.
neglectum, *Schott, in Bonplandia*, vii. (1859) 26.—Zeylan.
nepenthoides, *Mart. in Schott, Meletem.* i. 17.—Reg. Himal.
ochraceum, *Schott, in Bonplandia*, vii. (1859) 27.—Reg. Himal.
ornatum, *Miq. Fl. Ind. Bat.* iii. 220.—Sumatra.
papillosum, Steud. ex Schott, Prod. Aroid. 46 = Leschenaultii.
penicillatum, *N. E. Br. in Journ. Linn. Soc.* xvii. (1881) 248. t. 5.—Hongkong
pentaphyllum, *Schott, Meletem.* i. 17.—Ind. or.; China
Pluhratzi, Blume, Rumphia, i. 110 = A. Dracontium.
praecox, De Vriese, ex C. Koch, Allg. Gartenz. (1857) 85 = ringens.
propinquum, Schott, in Oestr. Bot. Zeitschr. (1857) 338 = intermedium.
pulchrum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1881) 252. t. 6.—Ind. or.
pumilum, Blume, Rumphia, i. 107 = trilobatum.
Pythonium, Blume, l. c. 108 = Zomicarpa Pythonium.
quinatum, *Schott, Syn. Aroid.* 28.—Am. bor.
ringens, *Schott, Meletem.* i. 17.—Japon.
Roxburghii, Kunth, Enum. Pl. iii. 18 = cuspidatum.
Schimperianum, *Schott, in Bonplandia*, vii. (1859) 27.—Abyss.
serotinum, Miq. ex Franch. & Sav. Enum. Pl. Jap. ii. 4 = Sieroti um.
serratum, Schott, Meletem. i. 17 = japonicum.
Steholzii, De Vriese, ex C. Koch, Allg. Gartenz. (1857) 85 = ringens.
Stierotum. *Svboid, ex Regel, in Ind. Sem. Hort. Petrop.* (1866) 44.—Japon.
sikokianum, *Franch. & Sav. Enum. Pl. Jap.* ii. 507.—Japon.
speciosum, *Mart. in Schott, Meletem.* i. 17.—Reg. Himal.
Steudelii, *Schott, in Bonplandia*, vii. (1859) 26.—Ind. or.
stictopodum, Miq. Fl. Ind. Bat. iii. 219 = filiforme.
Strachevanum, Schott, in Oestr. Bot. Zeitschr. (1857) 338 = intermedium
Tatarinowii, *Schott, in Bonplandia*, vii. (1859) 27.—China.
Thunbergii, *Blume, Rumphia*, i. 105.—Japon.
tortuosum, *Schott, Meletem.* i. 17.—Reg. Himal.
tortuosum, Steud. ex Engl. in DC. Monog. Phan. ii. 544 = Steudelii.
tripartitum, *Engl. l. c.* 538.—Japon.
triphyllum, *Schott, Meletem.* i. 17.—Am. bor.
utile, *Hook. f. ex Engl. in DC. Monog. Phan.* ii. 587.—Reg. Himal.
verrucosum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 341.—Reg. Himal.
violaceotinctum, Schott, in Bonplandia, vii. (1859) 28 = crubescens.
Wightii, Hook. f. Bot. Mag. t. 5507 = neglectum.
Wightii, Schott, in Bonplandia, vii. (1859) 26 = Jacquemontii.
zebrinum, Hort. ex Nichola. Dict. Gard. i. 112 = triphyllum.

ARISARON, Adans. Fam. ii. 470 (1763) = **Arisarum**, Tourn. (Aroid.).

ARISARUM, Hall. in Rupp. Fl. Jen. ed. II. 251 (1745) = Calla, Rich. (Aroid.).

ARISARUM, [Tourn.] Targ. Toz. in Anal. Mus. Pis. Fir. ii. it. (1810) 67. *AROIDEAE*, Benth. & Hook. f. iii. 965.
ARISARON, Adans. Fam. ii. 470 (1763).
BALMISA, Lag. Gen. et Sp. Nov. 17 (1816).
HOMAIDA, Rafin. Fl. Tellur. iii. 68 (1836).
aspergillum, Dun. in Mém. Acad. Montp. (Sect. Sc.) i. (1847) 8 = simorrhinum.
austriale, Mich. in Guill. Arch. i. (1833) 90 = vulgare.
asoricum, Schott, in Oestr. Bot. Wochenbl. (1857) 190 = vulgare.
Balansae, Schott, l. c. = vulgare.
Clusii, Schott, Prod. Aroid. 23 = vulgare.

ARISARUM :—
crassifolium, Schott, in Bonplandia, ix. (1861) 369 = vulgare.
Forbesii, Schott, in Oestr. Bot. Wochenbl. (1857) 190 = vulgare.
hastatum, *Fomel, Nouv. Mat. Fl. Atl.* 290.—Algeria.
italicum, Rafin. Fl. Tellur. iii. 63 = Arum italicum.
Jacquini, Schott, Prod. Aroid. 22 = vulgare.
Libani, Schott, l. c. 21 = vulgare.
maculatum, Rafin. Fl. Tellur. iii. 63 = Arum maculatum.
oligocarpum, *Fomel, Nouv. Mat. Fl. Atl.* 290.—Algeria.
ovatum, Rafin. Fl. Tellur. iii. 63 = Lagenandra toricaria.
pictum, Rafin. l. c. = Arum pictum.
proboscideum, *Savi, Ossero. Div. Pisante.* 6.—Italia.
serpentinum, Rafin. Fl. Tellur. iii. 63 = Arisarum vulgare.
Sibthorpii, Schott, Prod. Aroid. 21 = vulgare.
simorrhinum, *Dur. in Duch. Rev. Bot.* i. (1845–46) 360.—Algeria.
subalpinum, Kotschy, ex Engl. in DC. Monog. Phan. ii. 563 = vulgare.
subsessile, Webb & Berth. Phyt. Canar. iii. 293 = vulgare.
tingitanum, Schott, Prod. Aroid. 22 = vulgare.
Vesingii, Schott, l. c. 20 = vulgare.
vulgare, *Targ. Toz. in Anal. Mus. Fis. Fir.* ii. 11. (1810) 67.—Reg. Mediterr.

ARISTAEA, A. Rich. in Orb. Dict. ii. 186 (1842) = **Aristea**, Soland. (Irid.).

ARISTARIA, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 296 = **Anthistiria**, Linn. (Gramin.).
barbata, Jungh. l. c. 297 = Anth. arguens.

ARISTEA, [Soland. in] Ait. Hort. Kew. i. 67 (1789); iii. 506. *IRIDEAE*, Benth. & Hook. f. iii. 701.
CLEANTHE, Salisb. in Trans. Hort. Soc. i. (1812) 213.
GENLISIA, Reichb. Consp. 60 (1828).
NIVENIA, Vent. Dec. Gen. Nov. 5 (1808).
WEDOWIA, Eckl. Verz. 16 (1827).
africana, *Hoffmgg. Verz. Pfl.* 40.—Afr. trop.
alata, *Baker, in Journ. Linn. Soc.* xxi. (1885) 405.—Afr. trop.
ancepa, *Eckl. ex Klatt, in Linnaea*, xxxiv. (1865–66) 560.—Afr. austr.
angolensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 270.—Afr. trop.
angustifolia, *Baker, in Journ. Linn. Soc.* xx. (1884) 369.—Madag.
bracteata, Pers Syn. i. 41 = capitata.
capitata, *Ker-Gawl. in Bot. Mag.* t. 605.—Afr. austr.
cladocarpa, *Baker, in Journ. Linn. Soc.* xx. (1884) 366.—Madag.
corniles, Vahl, Enum. ii. 124 = capitata.
compressa, *Buching. ex Baker, in Journ. Linn. Soc.* xvi. (1878) 131.—Natal.
cyanea, [*Soland. in*] *Ait. Hort. Kew.* i. 67.—Afr. austr.
dichotoma, Eckl. ex Klatt, in Linnaea, xxxiv. (1865–66) 561 = Eckloni.
dichotoma, *Ker-Gawl. Gen. Irid.* 12.—Afr. austr.
diffusa, *Eckl. Verz.* 16, nomen.—Afr. austr.
Ecklonii, *Baker, in Journ. Linn. Soc.* xvi. (1878) 112. —Afr. austr.
eriophora, Pers. Syn. i. 41 = cyanea.
fruticosa, *Pers. l. c.*—Afr. austr.
glauca, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 881.—Afr. austr.
intermedia, Eckl. Verz. 16 = dichotoma.
juncifolia, *Eckl. l. c.*; *Baker, in Journ. Bot.* xiv. (1876) 267.—Afr. austr.
Kitchingii, *Baker, in Journ. Linn. Soc.* xx. (1884) 369.—Madag.
laxera, Stokes, Bot. Comment. i. 224 = cyanea.
ingens, Hort. ex Steud. Nom. ed. II. i. 150 = melaleuca.
madagascariensis, *Baker, in Journ. Bot.* xiv. (1876) 267.—Madag.
major, Andr. Bot. Rep. t. 160 = capitata.
melaleuca, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 236.—Afr. austr.
pusilla, *Ker-Gawl. l. c.*—Afr. austr.
racemosa, *Baker, in Journ. Bot.* xiv. (1876) 267.—Afr. austr.
reptans, *A. Dietr. Sp. Pl.* ii. 516.—Afr. austr.
schlechteana, *Harv. ex Baker, in Journ. Bot.* xiv. (1876) 267.—Afr. austr.

ARISTEA :—
spathacea, Spreng. Syst. i. 156 = Bobartia spathacea.
spicata, Pers. Syn. i. 41 = capitata.
spiralis, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 236.—Afr. austr.
torulosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 883.—Afr. austr.
umbellata, *Sprong. Syst.* i. 156.—Afr. austr.
Wredowia, *Steud. Nom.* ed. II. i. 180.—Afr. austr.

ARISTEGA, Miers, Contrib. Bot. iii. 376 (1871). *MENISPERMACEAE.*
laevifolia, *Miers, l. c.* 376. t. 51.—Ind. or.

ARISTELLA, Bertol. Fl. Ital. i. 690 (1838) = **Stipa**, Linn. (Gram.).
bromoides, Bertol. l. c. = S. Aristella.
longiflora, Regel, in Act. Hort. Petrop. vii. (1880) 645.—As. temp.

ARISTIDA, Linn. Sp. Pl. ed. I. 82 (1753). *GRAMINEAE*, Benth. & Hook. f. iii. 1140.
ARTHRATHERUM, Beauv. Agrost. 32. t. 8. f. 8 (1812).
CHAETARIA, Beauv. l. c. 30. t. 8. f. 5, 6 (1812).
CURTOPOGON, Beauv. l. c. 32. t. 8. f. 7 (1812).
KIELBULL, Adans. Fam. ii. 31 (1763).
MOULINSIA, Rafin. in Sec. Bull. Bot. i. (1830) 221.
ORTACHNE, Nees, in Seem. Bot. Voy. Herald, 225 (1852–57).
SCHISTACHNE, Figar. & De Not. in Mem. Accad. Torin. Ser. II. xii. (1852) 252.
STIPAGROSTIS, Nees, in Linnaea, vii. 280 (1832).
STREPTACHNE, H. B. & K. Nov. Gen. et Sp. i. 124. t. 40 (1815).
TRIXOSTIS, Rafin. in Ser. Bull. Bot. i. (1830) 221.
abyssinica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 134 = Schimperiana.
acutiflora, *Trin. & Rupr. l. c.* 167.—Aegypt.; Arabia.
adoënsis, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 370. —Abyss.
Adscensionis, *Linn. Sp. Pl.* 82.—Reg. temp. et calid.
Adscensionis, Walt. Fl. Carol. 74 = oligantha.
Adscensionis, Sw. Obs. 40 = cognata.
aequiramea, *Schlecht. in Linnaea*, xxii. (1849) 845.— Texas.
aethiopica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 184 = Adscensionis.
affinis, Kunth, Rev. Gram. i. 61 = purpurascens.
americana, Linn. f. Pl. Jam. Pugill. 7; Syst. ed. X. 871 = virgata, bromoides, juncifolia.
amplissima, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 155.—Afr. trop.
Anzilliana, *Poir. Encyc. Suppl.* i. 451.—Ind. occ.
Antoniana, *Steud. in Lechl. Berb. Am. Austr.* 56. —Peruv.
arabica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 185 = Adscensionis.
arenaria, *Gaudich. in Freyc. Voy. Bot.* 407.—Austral. arundinacea, *Linn. Mant.* 146.—Ind. or.
asperata, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 68 = setifolia.
barbata, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 583.—Mexic.
barbicollis, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 152.—Abyss.
basiramea, *Vasey, in Coult. Bot. Gaz.* ix. (1884) 76.— Am. bor.
Behrsiana, *F. Muell. in Trans. Vict. Inst.* i. (1855) 44. —Austral.
Beyrichiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 104.—Am. centr.
bipartita, *Steud. Syn. Pl. Gram.* 140.—Afr. austr.
brachypoda, *Tausch, in Flora*, xix. (1836) 506.— Aegypt.; Arabia.
brachyptera, *Cass. & Sal. in Bull. Soc. Bot. Fr.* v. (1858) 169.—Afr. bor.
brevifolia, *Steud. Nom.* ed. II. i. 180.—Afr. austr.
bromoides, Boehl. in Proc. Ac. Sc. Philad. 1863 (1863) 92 = purpurea.
bromoides, *H. B. & K. Nov. Gen. et Sp.* i. 123.—Am. bor. occ.; Ecuador.
bromoides, Salzm. ex Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 118 = longiflola.
caffra, *Bory, Voy.* ii. 276.—Afr. trop.
calliveides, *Thurb. ex Boland. in Trans. Calif. Agr. Soc.* (1866) 194.—Calif.

ARISTIDA :—
calophylla, Boiss. Fl. Orient. v. 687 = paradisea.
calycina, *R. Br. Prod.* i. 172.—Austral.
canariensis, Willd. Enum. Hort. Berol. 99 = Adscensionis.
capensis, Savi, in Mem. Moden. (1807) 196 = plumosa.
capensis, *Thunb. Prod. Pl. Cap.* 19.—Afr. austr.
capillacea, Cav. Ic. v. 43. t 448. f. 1 = Cumingiana.
capillacea, *Lam. Illustr.* i. 156.—Ind. or.; Am. austr.
capillaris, Kunth, Enum. Pl. i. 188, in syn. = Cumingiana.
candelana, *Anderss. Enum. Pl. Ins. Galap.* 144.—Ins. Galap.
chaetophylla, *Steud. Syn. Pl. Gram.* 420.—Madag.
chapadensis, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68.—Bras.
chinensis, *Munro, in Proc. Am. Acad.* iv. (1864–65) 363.—China.
Chloris, Salzm. ex Doell, in Mart. Fl. Bras. ii. iii. 88 = Cynnopogon foliosus.
chrysopitis, *Steud. Nom.* ed. II. i. 181.—Arabia.
cilliata, *Desf. in Schrad. N. Journ.* iii. (1809) 255.— Afr. bor. austr.; Arabia.
ciliata, Steud. & Hochst. ex Steud. Nom. ed. II. i. 131 = hirtiglumis.
coarctata, H. B. & K. Nov. Gen. et Sp. i. 122 = setifolia.
coarctata, Lichtst. ex Kunth, Enum. Pl. i. 195 = congesta.
cornulescens, Desf. Fl. Atlant. i. 109 = Adscensionis.
cornulescens, Sieber, ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 88 = Sieberiana.
cornulescens, Hochst. ex Steud. Syn. Pl. Gram. 139 = adoënsis
cognata, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 157 = purpurascens, stricta.
compacta, *Anderss. Enum. Pl. Ins. Galap.* 145.—Ins. Galap.
complanata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 85.—Bras.
concinna, *Sond. ex F. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 140.—Afr. austr.
condensata, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 19. —Am. bor.
congesta, *Roem. & Schult. Syst.* ii. 401.—Afr. austr.
contorta, F. Muell. in Trans. Vict. Inst. i. (1858) 44 = arenaria.
crinita, Steud. Nom. ed. II. i. 181 = Stipa crinita.
crinita, *Y. Presl, in Rel. Haenk.* i. 238.—Peruv.
Cumingiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 141.—Ins. Philipp.
curtiseta, A. Gray, in Proc. Acad. Sc. Philad. 1862 (1863) 334 = purpurea.
curvata, *Nees, ex A. Rich. Tent. Fl. Abyss.* ii. 392.— Abyss.; Afr. austr.
curvifolia, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 583.—Mexic.
cyanantha, *Steud. Syn. Pl. Gram.* 141.—Afghan.
decorata, *Steud. l. c.* 421.—Palaestin.
deflexa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 393. —Abyss.
demissiplex, *Steud. Syn. Pl. Gram.* 139.—Afr. trop.
depressa, Retz. Obs. iv. 23 = Adscensionis.
depressa, Trin. Gram. Unifl. 178 = setacea.
desmantha, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 109.—Texas.
dichotoma, *Michx. Fl. Bor. Am.* i. 41.—Am. bor.
diffusa, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 68 = vestita.
dispersa, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 139.—Mexic.
divaricata, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 99.—Mexic.
divaricata; Jaeg. Ecolog. 7. t. 6 = coarctata.
divaricata, Lag. Elench. 5 = spadicea.
divulsa, *Anderss. Enum. Pl. Ins. Galap.* 148.—Ins. Galap.
dominguensis, Kunth, Rev. Gram. i. 62 = Andropogon hirtiflorus.
Drummondi, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 159.—Afr. austr.
Ehrenbergii, *Trin. & Rupr. Sérpic.* 126.—Arabia.
elatior, Cav. Ic. vi. 56. t. 559 = Adscensionis.
elegans, Nudge, Fl. Guian. ii. 1. 20 = capillacea.
Elliottiana, *Steud. Syn. Pl. Gram.* 139.—Carolina.
elliptica, *Kunth, Enum.* iv. 1. 339.—Peru.

ARISTIDA :—
fasciculata, *Torr. in Ann. Lyc. N. York*, i. (1824) 154. —Canada.
Fendleriana, *Steud. Syn. Pl. Gram.* 430.—N. Mexic.
festucoides, *Poir. Encyc. Suppl.* i. 458.—Afr. trop.
filipendula, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 38 = purpurea.
flaccida, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 117.—Bras.
flexuosa, *Fourn. Mex. Pl. Enum. Gram.* 77, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 538.—Mexic.
floccosa, Coss. & Dur. in Expl. Sc. Algér. i. 82 = plumosa.
floridana, *Vasey, Gram. U. St.* 21.—Am. bor.
foeniculoides, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 160, 158.—Ind. or.
Forskahlei, Trausch, in Flora, xix. (1836) 506 = plumosa.
Freycinetii, Steud. Nom. ed. II. i. 131 = Stipa crinita.
funiculata, *Trin. ex Steud. l. c.*—Nubia.
funiculata, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 159.—Afr. bor.; Arab.; Beluchist.
furcata, Pois. ex Roem. & Schult. Syst. ii. 711 = Bouteloua bromoides.
Gardneriana, Steud. Syn. Pl. Gram. 127 = setifolia.
geminata, Willd. ex Steud. Nom. ed. II. i. 131 = Gymnopogon foliosa.
geminiflora, *Fourn. Mex. Pl. Enum. Gram.* 77, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 538.—Mexic.
geminiflora, Steud. Syn. Pl. Gram. 144 = geminifolia.
geminifolia, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 169.—Afr. austr.
gesticulata, *Rudge, Am. Monthly Mag.* (1817), *ex Trin. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 179.—Am. bor.
Geyeriana, Steud. Syn. Pl. Gram. 138 = purpurascens.
gibbosa, *Kunth, Enum. Pl.* i. 189.—Bras.
gigantea, Linn. f. Suppl. 113 = Adscensionis.
glabernima, *Steud Syn. Pl. Gram.* 135.—Chili.
glauca, Steud. l. c. = longiramea.
gossypina, *Bosc, ex Beauv Agrost.* 30, nomen ; *Roem. & Schult. Syst.* ii. 891.—Carolina
gracilis, *Ell. Sketch.* i. 141.—Am. bor.
gracillima, *Oliver, in Trans Linn. Soc.* xxix. (1875) 178.—Afr. trop.
Griesbachiana, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 538.—Mexic.
guineensis, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 187.—Guinea.
gyrans, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 18.—Am. bor.
Heymanni, *Regel, in Act. Hort. Petrop.* vii. (1880) 649. —As. centr.
hirtigluma, *Steud. Nom.* ed. II. i. 131 ; *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 171.—Aegypt. ; Syria; Arabia.
hordeacea, *Hochst. ex Steud. Syn. Pl. Gram.* 142 = Steudeliana.
hordeacea, *Kunth, Rev. Gram.* ii. t. 178.—Afr. trop.
Humboldtiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 118.—Mexic.
humilis, *H. B. & K. Nov. Gen. et Sp.* i. 191.—Venezuela.
hygrometrica, *R. Br. Prod.* 174.—Austral.
Hystrix, *Linn. f. Suppl.* 113.—Afr. austr.
implexa, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68. —Bras.
interrupta, *Cav. Ic.* v. 45. t. 471. f. 3.—Mexic.
Jacquiniana, *Tausch, in Flora*, xix. (1836) 508.—Am. austr.
jubellata, *Kunth, Rev. Gram.* i. 62.—Mexico.
junciformis, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 143.—Afr. austr.
Karwinskiana, *Trin. & Rupr. l. c.* 181.—Mexic.
Ketachji, Hochst. ex Steud. Syn. Pl. Gram. 149 = funiculata.
Kunthiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 151.—Afr. trop.
laevis, *Kunth, Enum. Pl.* i. 192.—Bras.
Lamarckii, Steud. Nom. ed. I. 69 = amplissima.
lanata, Forsk. Fl. Aegypt. Arab. 25 = plumosa.
lanata, *Poir. Encyc. Suppl.* i. 458.—Am. bor.
lanosa, *Muhl. Descr. Uber.* 174.—Am. bor.
lanuginosa, Desv. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iv. 22. (1836) 46 = setacea.
laxa, *Cav. Ic.* v. 44. t. 470.—Ins. Philipp. ; Am. austr.

ARISTIDA :—
laxa, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 86 = flaccida.
laxa, Willd. ex Trin. & Rupr. l. c. vii. (1849) 130 = diffusa.
leiocalycina, *Trin. & Rupr. l. c.* 161.—Guinea.
leptopoda, *Benth. Fl. Australiensis*. vii. 562.—Austral.
Liebmannii, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 534.—Mexic.
longerodista, *Steud. Syn. Pl. Gram.* 140.—Afr. trop.
longespicula, *Poir. Encyc. Suppl.* i. 459.—Am. bor.
longifolia, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 84.—Bras.
longiramea, *Y. Presl, in Rel. Haenk.* i. 224.—Mexic.
longiseta, *Steud. Syn. Pl. Gram.* 420.—N. Mexic.
lutescens, *Steud. l. c.* 145.—Afr. austr.
laxoniensis, *Cav. Ic.* v. 45. t. 470.—Ins. Philipp.
macranthera, Hochst. ex Boiss. Fl. Orient. v. 693 = funiculata.
macrathera, A. Rich. Tent. Fl. Abyss. ii. 398 = praec.
macrochaeta, *Steud. Syn. Pl. Gram* 134.—Am. bor.
macrochloa, *Hochst. in Flora*, xxxviii. (1855) 200.—Afr. or.
maritima, *Steud. Syn. Pl. Gram.* 137.—Gundel.
mauritiana, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 392.—Abyss.
mauritiana, Kunth, Rev. Gram. i. 265. t. 44 = Adscensionis.
meccana, *Hochst. ex Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 152.—Arabia.
meggiana, *Spreng. Syst. Cur. Post.* 31.—Bras.
mendocina, *Phil. Serr. Mend. Ali.* 47.—Chili.
micropoda, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 107.—Am. bor.
minuta, Pers. ex Roem. & Schult. Syst. ii. 711 (= *Eutriana Ledebourii*).—Ins. S. Doming.
modatica, *Steud. Syn. Pl. Gram.* 139.—Abyss.
muehlenbeckioides, *Fourn. Mex. Pl. Enum. Gram.* 79, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 534.—Mexic.
murina, *Cav. Ic.* v. 44. t. 469.—Reg. Argent.
natalitia, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 150.—Afr. trop.
namaquensis, *Trin. & Rupr. l. c.* 174.—Afr. austr.
nana, *Steud. Nom.* ed. II. i. 131.—Chili.
Necsiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 113.—Bras.
nigrescens, *Y. Presl, in Rel. Haenk.* i. 238.—Mexic.
nutans, *Forsk. & Hempr. ex Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 185.—Arabia.
nutans, Hort. Berol. ex Steud. Nom. ed. II. i. 131 = Adscensionis.
obtusa, *Delile, Fl. Egypte*, 175. t. 13. f. 2.—Afr. bor. et austr. ; Arab.
oligantha, *Michx. Fl. Bor. Am.* i. 41.—Am. bor.
orinbensis, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 534.—Chili?
pallens, Nutt. Gen. Am. i. 57 = oligantha.
pallens, *Cav. Ic.* v. 45. t. 468.—Chili.
Palmeri, *Vasey, in Torrey Bot. Club*, x. (1883) 42.—Am. bor.
pallida, *Steud. Syn. Pl. Gram.* 143.—Nubia.
palustris, *Vasey, Cat. Gram. U. St.* 35.—Am. bor.
paniculata, *Forsk. Fl. Aegypt. Arab.* 25.—Arabia.
papposa, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 178.—Afr. austr.
paradisea, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) 11. 1219.—Persia; Afghan.; Aegypt.; Arab.
parodii, J. A. Schmidt, Betr. Fl. Cap. Verd. Ins. 140 = funiculata.
parviflora, Steud. Syn. Pl. Gram. 140 = vagans.
parviflora, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 26 = oligantha.
pennata, *Trin. in Mém. Acad. Pétersb.* vi. (1815) 488. —Persia; Turkestan.
perrensis, *Fourn. ex Trin. & Rupr. l. c.* Sér. VI. vii. (1849) 104.—Carolina.
pilgeana, Buroh. ex Spreng. Syst. i. 269 = capensis.
pilgeana, Buroh. ex Roem. & Schult. Syst. ii. Mant. 579 = capensis.
pilosa, *Labill. Sert. N. Caled.* 13. t. 17.—N. Caled.
plumosa, Desf. Fl. Atlant. i. 109 = ciliata.
plumosa, *Linn. Sp. Pl.* ed. II. 1658.—Reg. Mediterr. ; Persia ; Arab.
pogonopoda, *Boiss. Fl. Orient.* v. 496.—Beluchistan.
polyclinos, *Wats. in Trans. Linn. Soc.* xxvii. (1869) 80.—Afr. trop.
proxima, *Steud. Syn. Pl. Gram.* 145.—Afr. austr.

ARISTIDA :—
pseudo-hystrix, *Steud. Syn. Pl. Gram.* 142.—Afr. austr.
pumila, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 85 = Adscensionis.
pungens, *Desf. Fl. Atlant.* i. 109. t. 85.—Afr. bor. ; Turkestan ; Sibiria.
pungens, Sieber, ex Trin. Gram. Unifl. Diss. i. 181 = vulnerans.
pungens, Steud. Syn. Pl. Gram. 145 = scoparia.
purpurascens, *Poir. Encyc. Suppl.* i. 459.—Am. bor. ; Jamaic.
purpurea, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 145.—Am. bor.
pusilla, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 140.—Afr. austr.
quinqueseta, *Steud Syn. Pl. Gram.* 420.—Maurit.
racemosa, Muhl. Descr. Uber. 172 = purpurascens.
racemosa, *Spreng. Mant.* i. 35.—Ins. S. Helen.
Raddiana, *Savi, in Mem. Moden.* (1837) 198.—Aegypt.
ramosa, *R. Br. Prod.* i. 178.—Austral.
ramosissima, *Engelm. ex A. Gray, Man. Bot. U. St.* ed V. 618.—Am. bor.
recurvata, *H. B. & K. Nov. Gen. et Sp.* i. 123.—Am. austr
refracta, *Griseb. Cat. Pl. Cub.* 228.—Cuba.
repens, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831 87.—Ins. Galap.
Riedelians, *Trin & Rupr. l. c.* Sér. VI. vii. (1849) 113. —Panama.
rigescens, *Roem. & Schult. Syst.* ii. 400.—Ind. or.
rigida, *Cav. Ic.* v. 44. t. 469.—Ins. Philipp.
rigida, Roth, Nov. Pl. Sp. 42 = rigescens.
riparia, *Trin. in Bull. Acad. Pétersb.* i. (1836) 68. —Bras.
Roemeriana, *Scheele, in Linnaea*, xxii. (1849) 848.—Texas.
Roylcana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 160.—Ind. or.
rufescens, *Steud. Syn. Pl. Gram.* 421.—Ins. Mascar.
sabulosa, Kunth, Enum. Pl. i. 197 = setifolia.
Sanctae-Luciae, *Trin. Diss.* ii. 25.—Bras.
scabra, *Kunth, Rev. Gram.* i. 62.—Mexico.
Schaffneri, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 534.—Mexic.
Schiedlana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 120.—Mexic.
Schimperi, Hochst. & Steud ex Steud. Syn. Pl. Gram. 143 = ciliata.
Schweinfurthii, *Boiss. Fl. Orient.* v. 493.—Aegypt. ; Arab.
scoparia, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 176.—Aegypt.
secunda, Ledeb. ex Steud. Nom. ed. II. i. 132 (= *Eutriana Ledebourii*).—Ins. S. Doming.
sericea, Ehrenb. ex Boiss. Fl. Orient. v. 497 = paradisea.
setacea, Hort. ex Steud. Nom. ed. II. i. 132 = Adscensionis.
setacea, *Retz. Obs.* iv. 22.—Ind. or.
setacea, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 84 = Adscensionis.
setifolia, *H. B. & K. Nov. Gen. et Sp.* i. 123.—Mexic ; Bras.
Sieberiana, *Trin. in Spreng. Neue Entdech.* ii. 61.—Afr. bor. ; Creta ; Syria.
simplex, *Steud. Syn. Pl. Gram.* 420.—Madag.
simpliciflora, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 18. —Am. bor.
simplicissima, *Steud. Syn. Pl. Gram.* 139.—Ins. Borbon.
sorsogonensis, *Y. Presl, in Rel. Haenk.* i. 224.—Ins. Luzon.
spadicea, *H. B. & K. Nov. Gen. et Sp.* i. 122.—Am. austr.
spadicea, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. 11. (1836) 43 = tincta.
spiciformis, *Ell. Sketch.* i. 141.—Am. bor.
spicigera, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 136.—Arab.
squarrosa, *Trin. in Spreng. Neue Entdech.* ii. 62.—Am. bor.
Stanleiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 155.—Abyss.
stipacea, Ehrenb. & Hempr. ex Boiss. Fl. Orient. v. 498 = funiculata.
stipaeformis, Hochst. ex Steud. Syn. Pl. Gram. 143 = pallida.
stipiformis, *Lam. Illustr.* i. 187.—Afr. trop.
stipoides, *R. Br. Prod.* i. 174.—Austral.
stricta, *Michx. Fl. Bor. Am.* i. 41.—Am. bor. et austr.

ARISTIDA :—
stricta, Mnhl. Desc. Gram. 174 = spiciformis.
stricta, Auct. ex Steud. Nom. ed. II. i. 182 = virgata.
stricta, Swtnd. Syn. Fl. Gram. 188 = purpurascens.
subacualis, *Steud. Nom.* ed. II. i. 182.—Afr. austr.
subaequans, *Doell, in Mart. Fl. Bras.* ii. III. 19.—Bras.
subdiffora, *Steud. Syn. Pl. Gram.* 188.—Ins. Guadeloupe.
subrecurvata, *Beauv. Agrost.* 152, nomen.—Hab. ?
subspicata, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 125.—Ins. Galap.
Swartziana, Steud. Syn. Fl. Gram. 187 = purpurascens.
Tenerifiæ, *Steud. l. c.* 420.—Ins. Teneriffa.
tenuiflora, *Steud. l. c.* 182.—Afr. trop.
tenuis, *Hochst. in Flora*, xxxvIII. (1855) 200.—Abyss.
tenuis, *Kunth, Rev. Gram.* i. 62.—Cumana; Mexic.
temipes, *Cav. Ic.* v. 46.—Panama.
Thomingii, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 127.—Afr. trop.
tincta, *Trin. & Rupr. l. c.* 111.—Guiana.
torta, *Kunth, Enum. Pl.* i. 190.—Bras.
trichodes, *Nees, in Hook. Kew Journ.* ii. (1850) 101. —Ins. Philipp.
tripilis, Thunb. Fl. Cayl. Diss. 2, ex Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 147 = Hystrix.
tuberculosa, *Nutt. Gen. Am.* i. 57.—Am. bor.
umbilicaris, Willd. ex Steud. Nom. ed. II. i. 182 = Bouteloua aristioides.
uniplumis, *Lichtst. in Roem. & Schult. Syst.* ii. 401.—Afr. austr.
Urvillei, *Steud. Syn. Pl. Gram.* 141.—Ins. Falkland.
vagans, *Cav. Ic.* v. 45. t. 471.—Austral.
vesita, *Thunb. Prod. Pl. Cap.* 19.—Afr. austr.
virgata, *Trin. in Spreng. Neue Entdeck.* ii. 60.—Am. bor.
Virleti, *Fourn. Mex. Pl. Enum. Gram.* 76, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 535.—Mexic.
vulgaris, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 131 = Adscensionis.
vulnerans, Trin. & Rupr. l. c. 175 = pungens.
vulpioides, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 251.—China.
Zeyheri, *Steud. Nom.* ed. II. i. 182.—Afr. austr.
Zilteli, *Aschers. in Verh. Bot. Ver. Brand.* xxi. (1880) 70.—Aegypt.

ARISTIDIUM, Lindl. Veg. Kingd. 116 (1847) = Bouteloua, Lag. (Gramin.).

ARISTOMENIA, Vell. Fl. Flum. 845 (1826) viii. t. 84 = Bulbine, Mikan (Compos.).
fruticosa, Vell. l. c. 846; viii. t. 84 = S. chrysantha.

ARISTOLOCHIA, Tourn. ex Linn. Syst. ed. I. (1758). *ARISTOLOCHIACEÆ*, Benth. & Hook. f. iii. 193; 1220.
AMBUYA, Rafin. Fl. Tellur. iv. 88 (1836).
DICTYANTHES, Rafin. in Loud. Gard. Mag. viii. (1832) 347.
DIOLOMELIS, Rafin. Fl. Tellur. iv. 88 (1836).
EINOMIA, Rafin. Med. Fl. i. 62 (1828).
ENDODICA, Rafin. l. c. (1826).
GLOSSULA, Reichb. Handb. 173 (1837).
GUACO, Liebm. in Forhandl. Skandin. Naturf. 1844 (1847) 308.
HEXAPLECTRIS, Rafin. Fl. Tellur. iv. 97 (1836).
HOCQUARTIA, Dum. Comm. Bot. 30 (1829).
HOWARDIA, Klotzsch, in Monatsb. Acad. Berl. (1858) 807.
ISIPHIA, Rafin. Med. Fl. ii. 283 (1830).
ISOTREMA, Rafin. in Am. Monthly Mag. (1819) 196.
PISTOLOCHIA, Rafin. Fl. Tellur. iv. 98 (1836).
PLAGISTRA, Rafin. l. c. (1836).
PROPHISA, Rafin. l. c. 99. (1836).
PYREPHIS, Rafin. l. c. (1836).
SIPHISIA, Rafin. Med. Fl. i. 62 (1828).
TROPEXA, Rafin. Fl. Tellur. iv. 98 (1836).
abbreviata, Mart. ex Mast. in Mart. Fl. Bras. iv. II. 108 = cymbifera.
abyssinica, Klotzsch, in Monatsb. Akad. Berl. (1859) 598 = bracteata.
acuminata, *Lam. Encyc.* i. 254.—Ind. or.
acuminata, Roxb. Hort. Beng. 64; Fl. Ind. iii. 489 = Roxburghiana.
acutiflora, *Duch. in Ann. Sc. Nat. Sér.* IV. ii. (1854) 54.—Fl. Amazon.

ARISTOLOCHIA :—
æthiopica, *Welw. Apont.* 548.—Afr. trop.
albida, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 75. —Afr. occ.
altissima, *Desf. Fl. Atlant.* ii. 324. t. 249.—Reg. Mediterr. or.
anatolica, Boiss. Fl. Orient. iv. 1080 = maurorum.
angulicida, *Yacq. Enum. Pl. Carib.* 80.—Am. austr.
angueicida, Pav. ex Duch. in DC. Prod. xv. i. 448 = velutina.
anguicida, Sieber, ex Duch. l. c. 479 = acuminata.
angulata, *Boj. ex Duch. l. c.* 467.—Madag.
angulicauta, Michx. Fl. Bor. Am. ii. 258 = tomentosa.
angulosa, Wall. ex Duch. in DC. Prod. xv. i. 460 = Roxburghiana.
angustifolia, *Boj. Hort. Maurit.* 277.—Ins. Maurit.
angustifolia, *Cham. in Linnæa*, vii. (1832) 211.—Bras.
antibysterica, *Mart. ex Duch. in DC. Prod.* xv. i. 477. —Bras.
appendiculata, Vell Fl. Flum. ix. t. 98 = macroura.
arborescens, *Linn. Sp. Pl.* 960.—Am.
arcuata, *Mast. in Mart. Fl. Bras.* iv. II. 101.—Am. trop.
areniculata, *Hance, in Journ. Bot.* xiv. (1876) 261.—Cambodia.
argentina, *Griseb. in Goett. Abh.* xix. (1874) 166.—Reg. Argent.
arkasana, *Lodd. Cat.* ed. XIV. (1826) 37.—Am. bor.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 186.—Algeria.
attica, Orphan. ex Duch. in DC. Prod. xv. i. 487 = longa.
auriculata, Vell Fl. Flum. ix. t. 98 = macroura.
arborescens, *Linn. Sp. Pl.* 960.—Am.
arcuata, *Mast. in Mart. Fl. Bras.* iv. II. 101.—Am. trop.
areniculata, *Hance, in Journ. Bot.* xiv. (1876) 261.—Cambodia.
argentina, *Griseb. in Goett. Abh.* xix. (1874) 166.—Reg. Argent.
arkasana, *Lodd. Cat.* ed. XIV. (1826) 37.—Am. bor.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 186.—Algeria.
attica, Orphan. ex Duch. in DC. Prod. xv. i. 487 = longa.
Aucherii, Jaub. & Spach, Illustr. i. 174. t. 99 = maurorum.
aurantiaca, *Duch. in DC. Prod.* xv. i. 475.—Venezuela.
auricularia, *Boiss. Diagn.* Ser. I. v. 49.—As. Min.
aurita, *Duch. in DC. Prod.* xv. i. 482.—Madag.
bactica, *Linn. Sp. Pl.* 961.—Europ.
barbata, *Yacq. Coll.* iii. 221.—Venezuela.
Baueri, Duch. in DC. Prod. xv. i. 484 = Thonetii.
Bernieri, *Duch. l. c.* 462.—Madag.
bifora, Willd. ex Duch. l. c. 467 = maxima.
bilabiata, *Linn. Sp. Pl.* ed. II. 1361.—Ins. S. Doming.
Billardieri, *Jaub. & Spach, Illustr.* i. 175. t. 100.—Syria; Cilicia.
bilobata, *Linn. Sp. Pl.* 960.—Ins. S. Doming.
birostris, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 60.—Bras.
Bodamae, *Dingt. in Flora*, xlvi. (1863) 301.—Thracia.
Bompiandi, Tenore, Cat. Sem. Hort. Neap. (1842) 12 = fimbriata.
Bottae, Jaub. & Spach, Illustr. i. 173. t. 98 = maurorum.
brachyura, *Duch. in DC. Prod.* xv. i. 466.—Ins. S. Doming.
bracteata, *Retz. Obs.* v. 29.—Ind. or.
bracteolata, Lam. Encyc. i. 256 = bracteata.
bracteosa, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 57.—Mexic.
Braniciana, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 155, nomen.—Hab. ?
brasiliensis, *Mart. & Zucc. Nov. Gen. et Sp.* i. 77.—Bras.
brevipes, *Benth. Pl. Hartw.* 15.—Mexic.
Bridgesii, *Duch. in DC. Prod.* xv. i. 463.—Chili.
Brunsueri, Jaub. & Spach, Illustr. ii. 40. t. 129 = Olivieri.
Burchellii, *Mast. in Mart. Fl. Bras.* iv. II. 90.—Am. trop.
californica, *Torr. in Pacif. Rail. Rep.* iv. 136.—Calif.
campanaeformis, DC. ex Duch. in DC. Prod. xv. i. 468 = Holostylis reniformis.
caracasana, Spreng. Syst. iii. 753 = trilobata.
caudata, Herb. Berol. ex Duch. in DC. Prod. xv. i. 446 = brachyura.
caudata, Booth, ex Lindl. in Bot. Reg. t. 1453 = macroura.
caudata, *Yacq. Enum. Pl. Carib.* 80.—Ins. S. Doming.
caudata, *Ferrari, in Anal. Soc. Cient. Argent.* v. (1878) 155 (*Centrib.* 45).—Reg. Argent.
Chamisonis, *Duch. in DC. Prod.* xv. i. 462.—Bras.
chilensis, *Miers, Trav. Chil.* ii. 331.—Chili.
chiquitensis, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 49.—Bolivia.
ciliata, Hook. Bot. Mag. t. 3736 = fimbriata.
ciliosa, Benth. in Mazuc, Bot. t. 20 = fimbriata.

ARISTOLOCHIA :—
Clausenii, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 67.—Bras.
clavicornis, *Griseb. Cat. Pl. Cub.* 115.—Cuba.
Clematitis, *Linn. Sp. Pl.* 963.—Europ.; Reg. Caucas.; As. Min.
clypeata, *Linden & André, Illustr. Hortic.* xvii. (1870) t. 60.—N. Granat.
condupicata, Poit. ex Duch. in DC. Prod. xv. i. 460 = oblongata.
conferta, *Mill. Gard. Dict.* ed. VIII. n. 11.—Mexic.
constricta, *Griseb. in Goett. Abh.* vii. (1857) 235.—Reg. Argent.
contorta, *Bunge, Enum. Pl. Chin. Bor.* 58.—China bor.
cordiflora, *Mutis, ex H. B. & K. Nov. Gen. et Sp.* ii. 149.—N. Granat.
cordigera, *Willd. ex Klotzsch, in Monatsb. Akad. Berl.* (1859) 691.—Bras.
coriacea, *Rafin. Atl. Journ.* 146.—Hab. ?
cornuta, *Mast. in Mart. Fl. Bras.* iv. II. 97.—Am. trop.
costaricensis, *Duch. in DC. Prod.* xv. i. 450.—Costa Rica.
crenata, Ehrenb. ex Duch. l. c. 478 = bracteata.
cretica, *Lam. Encyc.* i. 225.—Creta.
cubensis, *Linden, Cat.* n. 11. *Suppl.* (1856) 12.—Cuba.
cucullata, Pohl, ex Duch. in DC. Prod. xv. i. 482 = Holostylis reniformis.
cyclochila, Duch. in Ann. Sc. Nat. Sér. IV. ii. (1854) 45 = passiflorafolia.
cymbifera, *Mart. & Zucc. Nov. Gen. et Sp.* i. 76. t. 49.—Bras.
cynaechifolia, *Mart. & Zucc. l. c.* 77. t. 51.—Bras.
debilis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 197.—Japon.
delitantha, *F. Muell. Progm.* vi. 179.—Austral.
delitoides, *B. & H. ex Nov. Gen. et Sp.* ii. 146. t. 112. —Bras.
dictyantha, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 40.—Venezuela.
Duchartres, André, Le Mouv. Hortic. (1867) 61; Mast. in Gard. Chron. (1868) 516 = Ruiziana.
durior, *Hill, Veg. Syst.* xxi. 57.—Am. bor.
ecaudata, Duch. in DC. Prod. xv. i. 446 = dictyantha.
egensis, Poepp. ex Duch. l. c. 465 = acutiflora.
Ehrenbergiana, *Cham. in Linnæa*, vii. (1832) 210.—Ins. S. Doming.
elegans, *Mast. in Gard. Chron.* (1885) ii. 801.—Bras.
elliptica, Duch. in Ann. Sc. Nat. Sér. IV. ii. (1854) 51 = tigrina.
emarginata, *Duch. in DC. Prod.* xv. i. 467.—Bras.
emarginata, Willd. ex Duch. l. c. 466 = summulariifolia.
erecta, *Linn. Sp. Pl.* ed. II. 1362.—Vera Crux.
erianthe, *Mart. & Zucc. Nov. Gen. et Sp.* i. 78. t. 53.—Bras.
Eschscholtzii, Ledeb. ex Duch. in DC. Prod. xv. i. 481 = A. Tegaia.
euryantha, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 41.—Bras.
fasciculata, Lam. Fl. Fr. iii. 387 = A. Pistolochia.
filipendula, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 70.—Bras.
fimbriata, *Cham. in Linnæa*, vii. (1832) 210. t. 6.—Bras.
flexuosa, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 36. Mexic.
floribunda, *Lem. Illustr. Hortic.* (1858) t. 566.—Bras.
fluminensis, Bunk. in DC. Prod. Index, iv. 38 = odora.
foetens, *Lindl. Bot. Reg.* t. 1254.—Ind. occ.
foetida, *H. B. & K. Nov. Gen. et Sp.* ii. 147. t. 114.—Mexic.
fontanesii, Boiss. & Reut. Pugill. Pl. Hispan. 108 = longa.
Fordiana, *Hemsl. in Journ. Bot.* xxiii. (1885) 266.—China.
fragrantissima, *Ruiz, in Mem. Bejuc. Estrell.* (1868) 46.—Peruv.
fruticosa, Marsh. Arb. Am. 24 = macrophylla.
galeata, *Mart. & Zucc. Nov. Gen. et Sp.* i. 78. t. 50.—Bras.
galeata, Moritzi, Syst. Verz. 70 = cymbifera.
Galeottii, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 44.—Mexic.

ARISTOLOCHIA :—

Gardneri, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 69.—Bras.

Gaudichaudii, *Duch. l. c.* 72.—Ins. Kawak.

geminiflora, H. B. & K. Nov. Gen. et Sp. ii. 148. t. 117 = maxima.

gibbosa, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 58.—Mexic.

Giberti, *Hook. Bot. Mag.* t. 5845.—Reg. Argent.

gigantea, Hook. l. c. 4321 = grandiflora.

gigantea, *Mart. & Zucc. Nov. Gen. et Sp.* i. 75. t. 48.—Bras.

Gigas, Lindl. in Bot. Reg. (1842) Misc. 61. t. 60 = grandiflora.

glandulosa, *Kicks, in Bull. Acad. Brux.* déc. vii. (1880) ii. 11.—Cuba.

glauca, Desf. Fl. Atlant. ii. t. 250 = baetica.

glauca, Salisb. Prod. 215 = subglauca.

glaucescens, *H. B. & K. Nov. Gen. et Sp.* ii. 147. t. 116.—N. Granat.

Glaziovii, *Mast. in Mart. Fl. Bras.* iv. ii. 90.—Am. trop.

Goldiana, *Hook. f. in Trans. Linn. Soc.* xxv. (1865) 185. t. 14.—Afr. trop.

Gouanii, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 66.—N. Granat.

gracilis, *Duch. in DC. Prod.* xv. i. 460.—Am. austr.

grandiflora, *Arruda, in Kuster, Trav. Bras.* 499.—Pernamb.

grandiflora, Gomez, Obs. Bot. Med. ii. (1803) 14. t. 3; et in Mem. Ac. Lisboa, iii. (1812) Mém. dos Corresp. 54 = cymbifera.

grandiflora, *Sw. Prod. Veg. Ind. Occ.* 126.—Jamaica.

grandifolia, Saliab. Prod. 215 = macrophylla.

Griffithii, *Hook. f. & Thoms. ex Duch. in DC. Prod.* xv. i. 437.—Reg. Himal.

hastata, H. B. & K. Nov. Gen. et Sp. ii. 148 = pentandra.

hastata, *Yach, in Malay. Misc.* ii. (1822) n. vii. 6.—Malaya.

hastata, Klotzsch, in Monatsb. Acad. Berl. (1859) 597 = Zollingeri.

hastata, Nutt. Gen. Am. ii. 200 = A. Serpentaria.

hians, *Willd. in Mém. Soc. Nat. Mosc.* ii. (1809) 100.—Venezuela.

Hilariana, *Duch. in DC. Prod.* xv. i. 451.—Bras.

hirsuta, Linn. Syst. ed. XII. 601 = hirta.

hirsuta, Muhl. Desc. Uber. 18 = tomentosa.

hirta, Buch. in DC. Prod. Index, iv. 38 = torta.

hirta, Georgi, Beschreib. Russ. Reich. iii. v. 1975 = poetica.

hirta, *Linn. Sp. Pl.* 961.—As Min.

hispida, *Pohl, ex Duch. in DC. Prod.* xv. i. 498.—Bras.

iberica, *Fisch. & Mey. ex Duch. in DC. Prod.* xv. i. 498.—Reg. Caucas.

imbricata, *Mast. in Journ. Linn. Soc.* xiv. (1875) 494.—Ins. Philipp.

incisa, *Duch. in DC. Prod.* xv. i. 490.—Anatolia.

indica, *Linn. Sp. Pl.* 960.—Ind. or.

infesta, Saliab. Prod. 215 = A. Clematitis.

inflata, *H. B. & K. Nov. Gen. et Sp.* ii. 145. t. 111.—N. Granat.

insignis, *Versch. ex Gard. Chron.* (1868) 349.—Bras.

Ipensi, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 153 (*Contrib.* 46).—Paraguay.

Jachii, Steud. Nom. ed. II. i. 141 = hastata.

japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 186.—Japon.

Kaempferi, *Willd. Sp. Pl.* iv. 152.—Japon.

Karwinskii, *Duch. in DC. Prod.* xv. i. 442.—Mexic.

Kotschyi, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 237 = brachysta.

labiata, Willd. in Mém. Soc. Nat. Mosc. ii. (1809) 101. t. 6 = brasiliensis.

labiosa, Kze-Gawl. in Bot. Reg. (1822) t. 689 = cymbifera.

lanceolata, Wight, Ic. v. t. 1856 = indica.

Leprieurii, *Duch. in DC. Prod.* xv. i. 451.—Guiana.

leuconeura, *Linden, in Belg. Hortic.* viii. (1858) 165.—Am. austr.

Lindeniana, *Duch. in DC. Prod.* xv. i. 453.—Cuba.

lineolifolia, *Wright, ex Griseb. Cat. Pl. Cub.* 115.—Cuba.

lineata, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 153 (*Contrib.* 47).—Reg. Argent.

ARISTOLOCHIA :—

longa, Georgi, Beschr. Russ. Reich. iii. v. 1974 = A. Clematitis.

longa, *Linn. Sp. Pl.* 961.—Reg. Mediterr.

longa, Thunb. Fl. Japon. 144 = debilis.

longa, Woodv. Med. Bot. ii. t. 107 = baetica.

longiflora, *Engelm. & A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 259.—Texas.

longifolia, *Champ. ex Benth. in Hook. Kew Journ.* vi. (1854) 116.—Hongkong.

longifolia, Roxb. Hort. Beng. [102]; Fl. Ind. iii. 490 = moluccana.

lutea, Desf. Choix, 18 t. 8 = pallida.

lutescens, *Duch. in DC. Prod.* xv. i. 461.—Bras.

macradenia, Hook. Bot. Mag. t. 4467 = glandulosa.

macrocarpa, *Duch. in DC. Prod.* xv. i. 497.—Afr. occ.

macroglossa, Jaub. & Spach, Illustr. ii. 88. t. 127 = Tournefortii.

macrophylla, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 68.—Guiana.

macroura, *Gomez*, Lam. Encyc. i. 252 = A. Sipho.

macrota, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 48.—Guiana.

macroura, *Gomes, in Mem. Ac. Lisboa*, iii. (1812) *Mem. dos Corresp.* 77.—Bras.

Mannii, *Hook. f. in Trans. Linn. Soc.* xxv. (1865) 186.—Afr. trop.

manupitiensis, *Schrad. in Gott. Gel. Ans.* i. (1821) 719.—Hab. ?

Mathewsii, *Duch. in DC. Prod.* xv. i. 497.—Peruv.

mauritiana, Pers. Syn. ii. 537 = brasiliensis.

maurorum, Klotzsch, in Monatsb. Acad. Berl. (1859) 598 = bracteata.

maxorum, *Linn. Sp. Pl.* ed. II. 1368.—As. Min.; Persia.

maxima, Cham. in Linnaea, vii. (1832) 208 = Chamissonis.

maxima, *Jacq. Enum. Pl. Carib.* 30.—Ind. occ.

mayorensis, Fisch. ex Duch. in DC. Prod. xv. i. 479 = indica.

melastoma, *Manso, ex Duch. l. c.* 461.—Bras.

mexicana, *Kostel. Allg. Med.-Pharm. Fl.* ii. 467.—Mexic.

mexicana, A. Dietr. Syn. v. 196 = maxima.

micrantha, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 55. t. 5.—Mexic.

microphylla, *Willd. Enum. Hort. Berol. Suppl.* 68.—Hab. ?

microstoma, *Boiss. & Sprun. Diagn.* Ser. I. v. 50.—Graecia.

microura, Pohl, ex Duch. in DC. Prod. xv. i. 496 = setosa.

mollissima, *Hance, in Journ. Bot.* xvii. (1879) 300.—China.

moluccana, *Duch. in DC. Prod.* xv. i. 488.—Ins. Molucc.

moschata, Wedd. ex Duch. l. c. 475 = odoratissima.

multiflora, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 78.—Ins. Comoro.

multinervis, *Pomel, Nouv. Mat. Fl. Atl.* 187.—Algeria.

nervosa, *Duch. in DC. Prod.* xv. i. 480.—Ind. or.

nummularifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 145.—Venezuela.

oblonga, Vell. Fl. Flum. ix. t. 99 = rumicifolia.

oblonga, *Yacq. Hort. Schoenb.* ii. 29.—Hab. ?

obtusata, *Sw. Prod. Veg. Ind. Occ.* 126.—Ind. occ.

odora, *Steud. Nom.* ed. II. i. 128.—Bras.

odoratissima, Benth. Pl. Hartw. 82 = gibbosa.

odoratissima, *Linn. Sp. Pl.* ed. II. 1362.—Jamaica.

odoratissima, Vell. Fl. Flum. ix. t. 97 = odora.

officinalis, T. Nees, Fl. Offic. t. 144 = A. Serpentaria.

Olivieri, *Coleg. ex Boiss. Diagn.* Ser. I. v. 50.—Mesopot.

orbiculata, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 59.—Bras.

orbiculata, Vell. Fl. Flum. ix. t. 96 = cymbifera.

ornithocephala, Hook. Bot. Mag. t. 4120 = brasiliensis.

Ottonis, Klotzsch, ex Duch. in DC. Prod. xv. i. 476 = pandurata.

ovalifolia, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 50.—Mexic.

pandisantha, *Boiss. Diagn.* Ser. I. xii. 104.—Syria.

pallida, Salisb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 556 = longa.

pallida, *Willd. Sp. Pl. iv.* 162.—Europ.; As. Min.

pandurata, *Yacq. Hort. Schoenb.* iv. 49. t. 497.—Venezuela.

ARISTOLOCHIA :—

pandurata, Wall. ex Duch. in DC. Prod. xv. i. 479 = indica.

pandurifformis, Willd. Sp. Pl. iv. 152 = pandurata.

pannosa, *Mast. in Journ. Linn. Soc.* xxiv. (1878) 498.—Peruv.

papillaris, *Mast. in Mart. Fl. Bras.* iv. ii. 100.—Am. trop.

paraguariensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 156 (*Contrib.* 48).—Paraguay.

parnaraibensis, *Duch. in DC. Prod.* xv. i. 496.—Bras.

pardina, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 47.—Mexic.

parviflora, *Griseb. in Gott. Abh.* xxiv. (1879) 148.—Reg. Argent.

parvifolia, *Sibth. & Sm. Fl. Graec. Prod.* ii. 222.—Syria; Cyprus; Graecia !

parvifolia, Sprun. ex Duch. in DC. Prod. xv. i. 485 = microstoma.

passiflorafolia, *A. Rich. Fl. Cub. Fanerog.* ii. 195.—Cuba.

paucinervis, *Pomel, Nouv. Mat. Fl. Atl.* 186.—Algeria.

Pavoniana, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 54.—Mexic.

Pearcei, *Mast. in Journ. Linn. Soc.* xiv. (1875) 493.—Peruv.

Pearcei, *Phil. in Ann. Univ. Chil.* i. (1861) 68; *Linn. Journ.* xxiii. (1885) 231.—Chili.

peltata, *Linn. Sp. Pl.* 960.—Ins. S. Doming.

peltata, Sw. Obs. 341 = reniformis.

pentandra, *Jacq. Enum. Pl. Carib.* 30.—Cuba.

Peterziana, *Klotzsch, in Monatsb. Akad. Berl.* (1859) 599.—Afr. trop.

picta, Karst. Ausw. Schoen. Pfl. Venez. ii. 84 = pandurata.

pilosa, *H. B. & K. Nov. Gen. et Sp.* ii. 146. t. 118.—Ecuador.

piperifolia, *Grif. Itin. Bot.* 19.—Reg. Himal.

Pistolochia, Aucher, ex Duch. in DC. Prod. xv. i. 491 = incisa.

Pistolochia, *Linn. Sp. Pl.* ed. II. 962.—Reg. Mediterr. occ.

platanifolia, *Duch. in DC. Prod.* xv. i. 487.—Mexic.

platyloba, *Garcke, in Linnaea*, xxii. (1849) 69.—Guiana.

podocarpa, *A. Bertol. in Nov. Comm. Acad. Sc. Bon.* iv. (1840) 487.—Guatem.

Pohliana, *Duch. in DC. Prod.* xv. i. 498.—Bras.

polyrrhiza, *Sprengl. Syst. iv. Cur. Post.* 310.—Am. bor.

pontica, *Lam. Encyc.* i. 255.—As. Min.

praevenosa, *F. Muell. Fragm.* ii. 166.—Austral.

promissa, *Mast. in Gard. Chron.* (1879) i. 494.—Afr. trop.

prostrata, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 68.—Bolivia.

pubera, F. Baner, ex Duch. in DC. Prod. xv. i. 484 = Thozetii.

pubera, *R. Br. Prod.* 349.—Austral.

puberula, F. Muell. ex Duch. in DC. Prod. xv. i. 484 = pubera.

pubescens, Hort. ex C. Koch, Dendrol. ii. i. 395 = tomentosa.

pubescens, *Page, ex Steud. Nom.* ed. II. i. 128.—Am. bor.

pubescens, *Willd. ex Duch. in DC. Prod.* xv. i. 450.—Bras.

punctata, Balbis, ex Duch. l. c. 446 = brachyura.

punctata, *Lam. Encyc.* i. 253.—Ins. S. Doming.

pusilla, Pohl, ex Duch. in DC. Prod. xv. i. 466 = Clausseni.

quadrifloris, *Gueldenst. Reisen Russl.* i. 421.—Reg. Caucas.

Raja, *Mart. & Zucc. Nov. Gen. et Sp.* i. 78.—Bras.

recurvilabra, *Hance, in Journ. Bot.* xi. (1873) 75.—China.

refracta, Burch. ex Mast. in Mart. Fl. Bras. iv. ii. 98 = Chamissonis.

reniformis, Vell. Fl. Flum. ix. t. 100 = A. Raja.

reniformis, *Willd. Sp. Pl.* iv. 153.—Ins. S. Doming.

repens, *Mill. Gard. Dict.* ed. VIII. n. 13.—Mexic.

reticulata, Holton, ex Duch. in DC. Prod. xv. i. 457 = maxima.

reticulata, *Nutt. in Trans. Am. Phil. Soc.* N.S. v. (1837) 163.—Am. bor.

reticulata, Sesm. Bot. Voy. Herald, 193 = Mathewsii.

rigida, *Duch. in DC. Prod.* xv. i. 498.—Afr. austr.

ringens, Link & Otto, in Fl. Sel. ii. 88. t. 18 = brasiliensis.

ARISTOLOCHIA :—
ringens, *Vahl, Symb. Bot.* iii. 99.—Ind. or.
rotunda, Bové, ex Duch. in DC. Prod. xv. i. 437 = longa.
rotunda, Georgi, Beschr. Russ. Reich. iii. v. 1374 = A. Clematitis.
rotunda, *Linn. Sp. Pl.* 963.—Reg. Mediterr.
rotunda, Mog. & Sesse, ex Duch. in DC. Prod. xv. i. 442 = bracteosa.
Roxburghiana, *Klotzsch, in Monatsb. Akad. Berl.* (1858) 596.—Ind. or.; Malaya.
rubromarginata, Fisch. ex Duch. in DC. Prod. xv. i. 451 = Leprieurii.
rufomarginata, Leprieur, ex Duch. l. c. = Leprieurii.
rugosa, Lam. Encyc. i. 252 = obtusata.
Ruiziana, *Duch. in DC. Prod.* xv. i. 476.—Peruv.
rumicifolia, *Mart. & Zucc. Nov. Gen. et Sp.* i. 79.—Bras.
rumicifolia, R. Schomb. ex Duch. in DC. Prod. xv. i. 447 = dictyantha.
Rumphii, *Kostel. Allg. Med.-Pharm. Fl.* ii. 465.—Ins. Banda.
saccata, Hook. f. & Thoms. ex Klotzsch, in Monatsb. Akad. Berl. (1859) 608 = seq.
saccata, *Wall. Pl. As. Rar.* ii. 2. t. 108.—Reg. Himal.
sagittata, *Muhl. ex Duch. in DC. Prod.* xv. i. 434 = A. Serpentaria.
scabrida, Boiss. ex Duch. in DC. Prod. xv. i. 494 = hirta.
scabridula, Boiss. Diagn. Ser. I. xii. 105 = pseudantha.
scandens, *Mill. Gard. Dict.* ed. VIII. n. 10.—Hispan.
Sellowiana, Duch. in DC. Prod. xv. i. 458 = triangularis.
sempervirens, *Forsk. Fl. Aegypt.-Arab.* 156.—Arabia.
sempervirens, *Linn. Sp. Pl.* 961.—Creta.
sempervirens, Orph. ex Nym. Consp. 645 = altissima.
sepicola, *Mast. in Mart. Fl. Bras.* iv. ii. 100.—Am. trop.
sericea, *Benth. Pl. Hartw.* 81.—Mexic.
sericea, *Bianco, Fl. Filip.* 288.—Ins. Philipp.
Serpentaria, *Linn. Sp. Pl.* 961.—Am. bor.
serpentaria, DC. ex Duch. in DC. Prod. xv. i. 480 = nervosa.
sessilifolia, *Duch. l. c.* 464.—Bras.
setigera, Klotzsch, in Monatsb. Acad. Berl. (1859) 617 = chilensis.
setosa, *Duch. in DC. Prod.* xv. i. 496.—Bras.
sicula, *Tineo, in Guss. Fl. Sic. Syn.* ii. 878.—Sicilia.
sinarum, *Lindl. in Gard. Chron.* (1839) 708.—China.
Sipho, *L'Hérit. Stirp. Nov.* 13.—Am. bor.
smilacina, *Duch. in DC. Prod.* xv. i. 459.—Bras.
somaliensis, *Oliver, in Hook. Ic. Pl.* t. 1573.—Afr. trop.
Soyauxiana, *Oliver, l. c.* t. 1410.—Afr. trop.
spathulata, *Duch. in DC. Prod.* xv. i. 448.—Cuba.
Sprucei, *Mast. in Mart. Fl. Bras.* iv. ii. 88.—Am. trop.
strictiflora, Duch. in DC. Prod. xv. i. 484 = pubera.
subalata, *S. Wats. in Proc. Am. Acad.* xx. (1885) 372.—Mexic.
subcordata, Willd. ex Duch. in DC. Prod. xv. i. 468 = deltoidea.
subglauca, Lam. Encyc. i. 257; Brot. Fl. Lusit. ii. 598 = baetica.
subsagittata, Blanco, Fl. Filip. ed. I. 288 = A. Tagala.
surinamensis, Miq. in Linnæa, xviii. (1844) 246 = macrura.
surinamensis, *Willd. Sp. Pl.* iv. 151.—Guiana.
Tagala, *Cham. in Linnæa*, vii. (1832) 907.—Ins. Luzon.
talisoana, *Hook. & Arn. Bot. Beech. Voy.* 309.—Mexic.
tamaulipila, *Duch. in DC. Prod.* xv. i. 448.—Bras.
tapetaricha, Lem. Illustr. Hortic. iii. (1856) Misc. 22 = N. macrorura.
tenera, *Pohl, ex Duch. in DC. Prod.* xv. i. 466.—Grenad.
theriaca, *Mart. ex Duch. l. c.* 455.—Bras.
Thouvii, *F. Muell. Fragm.* ii. 167.—Austral.
Thwaitesii, *Hook. Bot. Mag.* t. 4918.—China.
tigrina, *A. Rich. Fl. Cub. Fanerog.* ii. 194.—Cuba.
timoriensis, *Decne. Herb. Tim. Descr.* 41.—Ins. Timor.
tomentosa, *Sims, Bot. Mag.* t. 1369.—Am. bor.
torta, Willd. ex Duch. in DC. Prod. xv. i. 466 = indata.
Tournefortii, *Nouk. & Orph. Illustr.* ii. 38. t. 128.—As. Min.; Ins. Chios.
triactina, *Hook. f. in Trans. Linn. Soc.* xxv. (1865) 126.—Afr. trop.

ARISTOLOCHIA :—
Trianaei, *Duch. in DC. Prod.* xv. i. 462.—N. Granat.
triangularis, *Cham. in Linnæa*, vii. (1832) 209. t. 6.—Bras.
tricaudata, *Lem. Illustr. Hortic.* xii. (1865) *Misc.* 49; xiii. (1866) t. 493.—Mexic.
trichosema, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 190.—Cuba.
trifida, Lam. Encyc. i. 251 = trilobata.
trilobe, Salisb. Prod. 214 = trilobata.
trilobata, Lam. Encyc. i. 251 = surinamensis.
tribobata, Lindl. Bot. Reg. t. 1899 = macroura.
trilobata, *Linn. Sp. Pl.* 960.—Ind. occ.
tripteris, *Ruiz. Fl. Ludov.* 94.—Am. bor.
trulliiformis, *Mast. in Mart. Fl. Bras.* iv. ii. 101.—Am. trop.
truncata, *Field. & Gardn. Sert. Pl.* t. 44.—Peruv.
turbacensis, *H. B. & K. Nov. Gen. et Sp.* ii. 149.—Am. bor.
Ubdesana, *Duch. in DC. Prod.* xv. i. 465.—Mexic.
undata, Mœnch, Meth. 719 = sempervirens.
unguliifolia, *Mast. in Journ. Linn. Soc.* xiv. (1875) 494.—Borneo.
utriculata, Seem. Bot. Voy. Herald, 198 = Mathewsii.
valentina, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 39.—Mexic.
vanillofla, *Duch. l. c.* 65.—Mexic.
velutina, *Duch. l. c.* 89.—Mexic.
veraguensis, *Duch. in DC. Prod.* xv. i. 458.—Am. centr.
Wageneriana, *Schlecht. in Linnæa*, xxvi. (1853) 631.—N. Granat.
Warmingii, *Mast. in Mart. Fl. Bras.* iv. ii. 109.—Am. trop.
Weddellii, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 62.—Bras.
Westlandi, *Hemsl. in Journ. Bot.* xxiii. (1885) 386.—China.
Wrightii, Seem. Bot. Voy. Herald, 331. t. 72 = brevipes.
Zollingeri, *Miq. Fl. Ind. Bat.* i. 1066.—Java.

ARISTOTELA, Adans. Fam. ii. 125 (1763) = **Othonna**, Linn. (Compos.).

ARISTOTELA, J. F. Gmel. Syst. 751 (1791) = **Aristotelia**, L'Hérit. (Tiliac.).

ARISTOTELEA, Lour. Fl. Cochinch. 522 (1790) = **Spiranthes**, Rich. (Orchid.).
spicata, Lour. l. c. = S. australis.

ARISTOTELEA, Spreng. Syst. ii. 450 (1825) = **Aristotelia**, L'Hérit. (Tiliac.).

ARISTOTELIA, L'Héret. Stirp. Nov. 31. t. 16 (1784). *TILIACEAE*, Benth. & Hook. f. i. 289.
ERAUSAMA, Deless. ex Steud. Nom. ed. II. i. 192 (1840).
FRIESIA, DC. Prod. i. 520 (1824).
australasica, *F. Muell. Fragm.* ii. 79.—Austral.
Braithwaitei, *F. Muell. in Wing. South. Sc. Record*, i. (1881) 149.—N. Hebrid.
Colensoi, *Hook. f. Handb. New Zeal. Fl.* 82.—N. Zel.
excta, *J. Ducn. in Trans. N. Z. Inst.* iii. (1871) 209.—N. Zel.
fruticosa, *Hook. f. Fl. N. Zeal.* i. 84.—N. Zel.
glabra, *Mirbel, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868) 44; *in Contrib. Bot.* ii. 180.—Am. merid.
glandulosa, Ruiz & Pav. Syst Veg. 125 = A. Macqui.
lucida, Salisb. Prod. 345 = A. Macqui.
Macqui, *L'Hérit. Stirp. Nov.* 31. t. 16.—Chili.
megalosperma, *F. Muell. Fragm.* ix. 84.—Austral.
peduncularis, *Hook. f. Fl. Tasm.* i. 52.—Tasmania.
racemosa, *Hook. f. Fl. N. Zeal.* i. 83.—N. Zel.

ARIVELA, Rafin. Sylva Tellur. 110 (1838) = **Cleome**, Linn. (Capparid.).
viscosa, Rafin. l. c. = C. viscosa.

ARIVONA, Steud. Nom. ed. II. i. 180 (1840) = **Arjona**, Comm. (Santal.).

ARJONA, Comm. ex Cav. Ic. iv. 57. t. 388 (1797). *SANTALACEAE*, Benth. & Hook. f. ii. 220.
ARIVONA, Steud. Nom. ed. II. i. 180 (1840).
ARJOONA, Endl. Gen. 295 (1836).
andina, *Phil. in Anal. Univ. Chil.* ii. (1862) 405; *et in Linnæa*, xxxiii. (1864-65) 223.—Chili.

ARJONA :—
appressa, *Phil. in Linnæa*, xxxiii. (1864-65) 233.—Chili.
linearis, *Miers, in Journ. Linn. Soc.* xvii. (1880) 180.—Reg. Argent.
longifolia, *Phil. in Anal. Univ. Chil* ii. (1862) 405.—Chili.
minima, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881, 80.—Chili.
patagonica, *Hemsl. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél.* 23.—Patagon.
pungens, *Phil. in Anal. Univ. Chil.* (1873) 587.—Chili.
pusilla, *Hook. f. Fl. Antarct.* ii. 342.—Patagon.
rigida, *Miers, in Journ. Linn. Soc.* xvii. (1880) 182.—Reg. Argent.
ruscifolia, *Poepp. in Nov. Act.-Nat. Cur.* xix. *Suppl.* i. (1843) (*Pl. Meyen.*) 112.—Chili.
tuberosa, *Cav. Ic.* iv. 57. t. 383.—Am. merid.
tuberosa, Phil. in Linnæa, xxxiii. (1864-65) 231 = rigida.

AKKEZOSTIS, Rafin. *New Fl. Am.* iv. 100 (1836) = **Cayaponia**, Manso (Cucurb.).
quinquelobata, Rafin. l. c. = C. Boykinii.

ARKOPODA, Rafin. Fl. Tellur. iii. 78 (1836) = **Reseda**, Tourn.
Luteola, Rafin. l. c. = R. Luteola.

ARMANIA, Bert. ex DC. Prod. v. 576 (1836) = **Enoplia**, Adans. (Compos.).
fruticulosa, Bert. l. c.—Am. austr.

ARMENIACA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Prunus**, Tourn. (Rosac.).
amarella, Schur, Enum. Pl. Transs. 178 = P. amarella.
atropurpurea, Loisel. in Duham. ed. Nov. v. 172. t. 553 = P. dasycarpa.
berriccocsa, Steudel, Fl. Auv. ed. II. 326 = P. Armeniaca.
brigantiaca, Pers. Syn. ii. 36 = P. brigantiaca.
dasycarpa, Pers. l. c. = P. dasycarpa.
Davidiana, Carr. in Rev. Hortic. (1870) 287.—China.
duracina, Dierb. ex Steud. Nom. ed. II. i. 533 = P. amarella.
epirotica, Gaertn. Mey. & Scherb. Fl. Wett. ii. 167 = P. Armeniaca.
fusca, Turp. & Poit. in Nouv. Duham. Arb. Fruit. i. 123. t. 60 = P. dasycarpa.
longifolia, Desc. Fl. Antill. ii. 108 = P. occidentalis.
lucida, May, in Rev. Hortic. (1878) 179 = P. lucida.
Mume, De Vries, Tijdsch. Fl. i. (1855) t. 1. t. 1 = V. Mume.
nigra, Pers. Syn. ii. 36 = P. dasycarpa.
pedunculata, Hort. ex Loud. Hort. Brit. Suppl. ii. 610 = P. pedunculata.
persicifolia, Loisel. in Duham. ed. Nov. v. 172. t. 552 = P. dasycarpa.
sibirica, Pers. Syn. ii. 36 = P. sibirica.
verrucosa, May, in Rev. Hortic. (1878) 299.—As. Min.
trichocarpa, M. Roem. Syn. Rosifl. 34 = P. tomentosa.
vulgaris, Lam. Encyc. i. 2 = P. Armeniaca.

ARMENIASTRUM, Lem. Jard. Fleur. iv. (1854) Misc. 78 = **Hexadesma**, A. Rich. (Verben.).
spicatastrum, Lem. l. c. 78 = E. spicatum.

ARMERIA, Linn. Syst. ed. I. (1735); Willd. Enum. Hort. Berol. 333 (1809). *PLUMBAGINEAE*, Benth. & Hook. f. ii. 698.
POLYANTHEMUM, Medic. in Staatsw. Vorles. Churpf. Phys. Ök. Ges. i. (1791) 238.
alliacea, Ebel in Hoffmgg. in Hoffmgg. & Link, Fl. Port. i. 441 = plantaginea.
alliacea, Griseb. Spicil. Fl. Rumel. ii. 296 = cariensis.
alliacea, Mutel, Fl. Fr. iii. 66. f. 406 = plantaginea.
alliacea, *Boiss. & Schmidt. Syst.* vi. 773.—Reg. Mediterr. occ.
alliacea, Schenck. Verz. Marok. 166 = baetica.
alliacea, Webb, ex Bail, in Journ. Linn. Soc. xvi. (1878) 560 = mauritanica.
alliacea, *Boiss. Voy. Bot. Espagne*, ii. 854.—Hispan.
alliodes, Willk. ex Willk. & Lange, Prod. Fl. Hisp. ii. 396 = longearistata.
alpina, Ebel, Armeriae Gen. Diss. (1840) 28 = Halleri.
alpina, Trivalt. ex Griseb. Spicil. Fl. Rumel. ii. 296 = carinensis.

ARMERIA:—
alpina, Gaud. Fl. Sic. Prod. 878 = nebrodensis.
alpina, Tenore, Syll. 160, ex Boiss. in DC. Prod. xii. 668 = majellensis.
alpina, Turcz. in Bull. Soc. Nat. Mosc. (1838) 99; xxv. (1852) ii. 399 = sibirica.
alpina, *Willd. Enum. Hort. Berol.* 338.—Europ.
andicola, C. Gay, ex Boiss. in DC. Prod. xii. 682 = andina.
andina, *Poepp. ex Boiss. l. c.*—Chili.
andrzeoczo, Boiss. l. c. 679 = vulgaris.
arctica, *Wallr. Beitr.* i. 193.—Unalaska.
arcuata, Welw. ex Boiss. & Kent. Pugill. Pl. Nov. 101 = boetica.
arenaria, Schult. in Roem. & Schult. Syst. vi. 771 = maritima.
argyrocephala, *Wallr. Beitr.* i. 206.—Europ.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 134.—Algeria.
australis, *Boiss. Voy. Bot. Espagne*, ii. 526.—Hispan.
berlengensis, *Daveau, in Bol. Soc. Broter.* ii. (1884) 24.—Lusitan.
boetica, *Boiss. Voy. Bot. Extagne*, ii. 740.—Hispan.
Boissieriana, *Cass. Notes Crit.* 44.—Hispan.
Bourgeæi, Boiss. ex Nym. Consp. 615 = Durieæi.
brachyphylla, *Boiss. in DC. Prod.* xii. 682.—Chili.
brevifolia, Kunze, ex Boiss. l. c. = chilensis.
bupleuroides, Gren. & Godr. Fl. Fr. ii. 736 = plantaginea.
caespitosa, *Boiss. in DC. Prod.* xii. 679.—Hispan.; Lusitan.
campestris, Wallr. Beitr. i. 208 = maritima.
canescens, *Boiss. in DC. Prod.* xii. 686.—Dalmatia.
cantabrica, *Boiss. & Reut. ex Willk. & Lange, Prod. Fl. Hisp.* ii. 366.—Hispan.
carinsia, *Boiss. in DC. Prod.* xii. 677.—As. Min.
Cephalotes, Hook. Bot. Mag. t. 4128 = mauritanica.
Cephalotes, Hoffmgg. & Link, Fl. Port. i. 440 = latifolia.
Cephalotes, Schousb. Vext. Marok. 146 = plantaginea.
Cephalotus, Boiss. in DC. Prod. xii. 675 = pinifolia.
chilensis, *Boiss. l. c.* 681.—Chili.
Chouletiana, *Pomel, Nouv. Mat. Fl. Atl.* 135.—Algeria.
cinerea, *Welw. ex Boiss. & Reut. Pugill. Pl. Nov.* 101.—Lusitan.
curvifolia, *Colla, in Mem. Acc. Torin.* xxxix. (1835) 5.—Hab.?
curvifolia, Bert. in Merc. Chil. 568; et Bull. Férussac, xx. (1830) 108 = chilensis.
denticulata, *Bertol. Amoen. Ital.* 77.—Italia.
denticulata, Portensch. Fl. Dalm. t. 6 = canescens.
dianthoides, Homem. ex Spreng. ex Boiss. in DC. Prod. xii. 683 = plantaginea.
Durieæi, *Boiss. l. c.* 684.—Hispan.
duriuscula, Bab. in Ann. Nat. Hist. Ser. II. iii. (1849) 436 = pubescens.
ebracteata, *Pomel, Nouv. Mat. Fl. Atl.* 132.—Algeria.
elongata, *Hoffm. Deutschl. Fl.* i. 150.—Europ.
elongata, Koch, in Mert. & Koch, Roehl. Deutschl. Fl. ed. III. ii. 457 = maritima.
eriophila, *Willk. in Bol. Soc. Broter.* ii. (1884) 145.—Lusitan.
exiguosa, Wallr. Beitr. i. 197 = pubescens.
fasciculata, *Willd. Enum. Hort. Berol.* i. 334.—Hispan.; Sardin.
fibrosa, *Pomel, Nouv. Mat. Fl. Atl.* 133.—Algeria.
filicaulis, *Boiss. Voy. Bot. Espagne*, ii. 527. t. 154.—Hispan.
formosa, Hort. ex Vilm. Fl. Fl. Terre, ed. I. 848 = Statice Pseudo-armeria.
formosa, Heynh. Nom. ii. 45 = mauritanica.
fruticosa, *Ladd. Bot. Cab.* t. 1587.—Europ. austr.
gaditana, *Boiss. in DC. Prod.* xii. 675.—Hispan.; Marocco.
glauca, Wallr. Arm. 315 = boetica.
glaucescens, *Desf. Cat. Hort. Par.* ed. III. 79.—Hab.?
globosa, Link, ex Boiss. in DC. Prod. xii. 684 = latifolia.
gracilis, Tenore, Syll. 158 = nebrodensis.
graudifiora, Hort. Schwetz. ex Boiss. in DC. Prod. xii. 684 = latifolia.
Gussonei, *Boiss. in DC. Prod.* xii. 687.—Sicil.
Halleri, Wallr. Beitr. i. 194 = alpina.
heterophylla, Wallr. l. c. 198 = nebrodensis.
hirta, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 369 = pubigera.
hirta, Willd. Enum. Hort. Berol. 336 = boetica.
humilis, Link, Handb. ii. 262 = caespitosa.

ARMERIA:—
intermedia, Link, ex Boiss. in DC. Prod. xii. 681 = pubescens.
juncea, *Girard, in Ann. Sc. Nat. Sér. III.* ii. (1844) 324.—Gall. merid.
juncea, Wallr. Beitr. i. 212 = allioides.
junciformis, Ebel, Armeriae Gen. Diss. (1840) 30 = filicaulis.
juniperifolia, J. Gay, ex Boiss. in DC. Prod. xii. 684 = Durieæi.
juniperifolia, Koch, in Flora, vi. (1823) 711 = leucocephala.
juniperifolia, Willd. ex Hoffmgg. & Link, Fl. Port. i. 443 = caespitosa.
Kochii, Boiss. in DC. Prod. xii. 686 = leucocephala.
labradorica, *Wallr. Beitr.* i. 185.—Groenl.; Labrador.
lachnolepis, *Pomel, Nouv. Mat. Fl. Atl.* 134.—Algeria.
Langei, *Boiss. ex Lange, in Kjoeb. Vidensk. Meddel.* (1861) 69.—Hispan.
latifolia, Moris, Fl. Sard. Elench. fasc. iii. 10 = Morisii.
latifolia, *Willd. Enum. Hort. Berol.* 334.—Lusitan.
leucantha, Salzm. ex Boiss. Voy. Bot. Espagne, ii. 525 = allioides.
leucocephala, *Koch, in Flora*, vi (1823) 712.—Corsica.
littoralis, Boiss. in DC. Prod. xii. 676 = Welwitschii.
littoralis, *Hoffmgg. & Link, Fl. Port.* t. 1. 441.—Lusitan.
littoralis, Willd. Enum. Hort. Berol. 333 = boetica.
longearistata, *Boiss. & Reut. Pugill. Pl. Nov.* 103.—Hispan.
longearistata, Bourg. in Bull. Soc. Bot. Fr. ii. (1855) 648 = plantaginea.
longifolia, *Desf. Cat. Hort. Par.* ed. III. 79, nomen.—Hab.?
lusitanica, Link, ex Boiss. in DC. Prod. xii. 675 = pinifolia.
maclovana, *Cham. in Linnaea*, vi. (1831) 567.—Ins. Falkland.
macrophylla, *Boiss. & Reut. Pugill. Pl. Nov.* 100.—Hispan.
macropoda, *Boiss. in DC. Prod.* xii. 688.—Italia.
maculata, Poepp. ex Boiss. l. c. 682 = andina.
madrensis, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) 12.—Madera.
majellensis, *Boiss. in DC. Prod.* xii. 685.—Italia.
maritima, Girard, ex Boiss. l. c. 680 = macrocnemis.
maritima, *Willd. Enum. Hort. Berol.* 333.—Europ.
mauritanica, *Wallr. Beitr.* i. 217.—Afr. bor.
meridionalis, Poepp. ex Boiss. in DC. Prod. xii. 682 = chilensis.
montana, G. Don, ex Loud. Hort. Brit. 114 = maritima.
montana, Soleir. ex Wallr. Beitr. i. 196 = leucocephala.
montana, Wallr. l. c. 210 = plantaginea.
Morisii, *Boiss. in DC. Prod.* xii. 687.—Sicil.
Muelleri, Huet, in Ann. Sc. Nat. Sér. III. xix. (1853) 255 = alpina.
multiceps, Wallr. Beitr. i. 196 = leucocephala.
nebrodensis, *Guss. Fl. Sic. Syn.* i. 366.—Italia.
neglecta, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 324.—Lusitan.
Orphanidis, Boiss. Diagn. Ser. II. iv. 71 = majellensis.
pinifolia, Bourg. ex Nym. Consp. 614 = macrophylla.
pinifolia, *Hoffmgg. & Link, Fl. Port.* i. 437.—Lusitan.
planifolia, Nym. Consp. 616. t. 15 = pubescens.
plantaginea, Boiss. Voy. Bot. Espagne, 526 = boetica.
plantaginea, *Willd. Enum. Hort. Berol.* 334.—Europ.
plantaginea, Webb, It. Hisp. 18 = latifolia.
praecox, Jord. in Bull. Soc. Dauph. fasc. iii. 79 = plantaginea.
pseudarmeria, Brot. Fl. Lusit. i. 488 = plantaginea.
pseud.-Armeria, Murr. Syst. Veg. ed. XIV. 300 = latifolia.
pubescens, Heynh. ex Steud. Nom. ed. II. i. 183 = maritima.
pubescens, *Link, in Report. Nat. Cur. Berol.* i. 180.—Europ.
pubigera, Boiss. in DC. Prod. xii. 678 = maritima.
pulchrerrima, *Boiss. l. c.* 688.—Gallia.
pungens, *Hoffmgg. & Link, Fl. Port.* i. 439.—Lusitan.
purpurea, *Koch, in Flora*, vi. (1823) 710 = elongata.
rhenana, Gremli, Exc. Fl. Schw. ed. IV. 350 = maritima.
rigida, Wallr. Beitr. i. 199 = plantaginea.
rumelica, Boiss. in DC. Prod. xii. 677 = carinsia.
ruscinonensis, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 339.—Hispan.

ARMERIA:—
sabulosa, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 537 = plantaginea.
sanguinolenta, *Wallr. Beitr.* i. 207.—Am. bor.
sardoa, *Spreng. Syst. Cur. Post.* 127.—Sardin.; Corsic.
scabra, Kunze, ex Boiss. in DC. Prod. xii. 683 = andina.
scabra, *Pall. in Willd. ex Roem. & Schult. Syst.* vi. 776.—As. bor.
scabriuscula, Kunze, ex Boiss. in DC. Prod. xii. 683 = andina.
scorzonerafolia, Tenore, Syll. 160 = macropoda.
scorzonerafolia, Willd. Enum. Hort. Berol. 334 = plantaginea.
scorzonerifolia, Link, in Schrad. Journ. iii. (1800) 60 = latifolia.
scorzonerifolia, Balb. & Nocca, Fl. Ticin. t. 6 = plantaginea.
scorzonerifolia, Friv. ex Nym. Consp. 615 = rumelica.
setacea, Delile, ex Nym. l. c. 616 = juncea.
setigera, Reichb. Ic. Fl. Germ. v. 84 = plantaginea.
sibirica, *Turcz. ex Boiss. in DC. Prod.* xii. 678.—Sibiria.
sicula, Heldr. ex Boiss. l. c. 685 = nebrodensis.
simplex, *Pomel, Nouv. Mat. Fl. Atl.* 132.—Algeria.
Soleirolii, Duby, Bot. Gall. Suppl. 1032 = leucocephala.
spinulosa, *Boiss. in DC. Prod.* xii. 675.—Barbaria.
splendens, *Boiss. Voy. Bot. Espagne*, ii. 749.—Hispan.
stenophylla, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 323 = plantaginea.
strusia, Balb. ex Boiss. in DC. Prod. xii. 685 = sardoa.
tenuis, Wallr. Beitr. i. 184 = filicaulis.
thessala, Boiss. Diagn. Ser. II. iv. 70 = cariensis.
Thomasii, Nym. Consp. 615 = leucocephala.
tingitana, Boiss. & Reut. Pugill. Pl. Nov. 102 = caditana.
trigonodies, Hort. Berol. ex Ebel, Armeriae Gen. Diss. (1840) 34 = plantaginea.
undulata, Boiss. in DC. Prod. xii. 685 = argyrocephala.
velutina, *Welw. ex Boiss. & Reut. Pugill. Pl. Nov.* 100.—Lusitan.
vestita, Willk in Willk. & Lange, Prod. Fl. Hisp. ii. 366 = longearistata.
villosa, *Girard, in Ann. Sc. Nat. Sér. III.* ii. (1844) 323.—Hispan.
vulgaris, Moris, Elench. ex Nym. Consp. 615 = sardoa.
vulgaris, Vis. Fl. Dalmat. ii. 6 = canescens.
vulgaris, Willd. Enum. Hort Berol. 333 = elongata.
Welwitschii, *Boiss. in DC. Prod.* xii. 676.—Hispan.
Willkommii, *Henriques, in Bol. Soc. Broter.* iii. (1884) 199, 224.—Hispan.

ARMERIASTRUM, Lindl. Veg. Kingd. 641 (1847) = **Acantholimon**, Boiss. (Plumbag.).

ARMODORUM, Breda, Orch. Kuhl. & Hass. (1827) = **Arachnanthe**, Blume (Orchid.).
distichum, Breda, l. c. [t. 6] = Arachnanthe Sulingi.

ARMORACIA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 426 (1800); Bernh. Syst. Verz. Erf. '258' (1800) = Cochlearia, Linn. (Crucif.).
amphibia, Peterm. Fl. Lips. Excurs. 477 = Nasturtium amphibium.
aquatica, Kostel. Allg. Med.-Pharm. Fl. v. 1571 = Nasturtium amphibium.
austriaca, Wimm. Fl. Schles. (1841) 80 = Nasturtium austriacum.
lapathifolia, Gilib. Fl. Lituan. ii. 52 = C. Armoracia.
macrocarpa, Baumg. Enum. Stirp. Transsylv. ii. 340 = C. macrocarpa.
rustica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 8 = seq.
rusticana, Gaertn. Mey. & Scherb. Fl. Wett. ii. 426 = C. Armoracia.
sativa, Bernh. Syst. Verz. Erf. 191 = C. Armoracia.
waldsteiana, Phil. in Linnaea, xxviii. (1856) 671.—Chili.

ARNEBIA, Forsk. Fl. Aegypt. Arab. 62 (1775).
BORAGINEAE, Benth. & Hook. f. ii. 863.
DIOCLEA, Spreng. Syst. i. 802 (1825).
MESNGNIHA, Endl. Gen. 648 (1839).
STENOGLENIUM, Turcz. in Bull. Soc. Nat. Mosc. (1840) 288.

ARNEBIA :—
Strobila, G. Don, Gen. Syst. iv. 327 (1837).
Toxostigma, A. Rich. Tent. Fl. Abyss. ii. 86. t. 77 (1851).
Bunge, *Beiss. Fl. Orient.* iv. 215.—Persia.
calycina, Stev. in Bull. Soc. Nat. Mosc. (1851) 598 = cornuta.
Cephalotes, A. DC. in DC. Prod. x. 96 = Macrotomia Cephalotes.
cornuta, *Fisch. & Mey., Ind. Sem. Hort. Petrop.* i. (1835) 23.—Orient; Turkest.
decumbens, *Coss. & Kralik, in Bull. Soc. Bot. Fr.* iv. (1857) 898, 402.—Oriena.
densiflora, Ledeb. Fl. Ross. iii. 140 = Macrotomia Cephalotes.
echioides, A. DC. in DC. Prod. x. 96 = Macrotomia echioides.
endochroma, Altch. in Journ. Linn. Soc. xviii. (1882) 81 = Macrotomia perennis.
fimbriata, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 507.—As. centr.
fimbriopetala, *Stocks, in Hook. Kew Journ.* iii. (1851) 180.—Persia; Beluchist.
flavescens, Boiss. Diagn. Ser. I. xi. 117 = linearifolia.
Griffithii, *Boiss. l. c.* Ser. II. iii. 126.—Ind. bor. occ.; Afghan.
guttata, *Bunge, Ind. Sem. Hort. Dorpat.* (1840) 7.—Sibir.
hispidissima, *DC. Prod.* x. 94.—Afr. trop.; Arab.; Ind. bor. occ.
linearifolia, *DC. l. c.* 95.—Aegypt.; Syria; Persia.
longiflora, C. Koch, in Linnæa, xxii. (1849) 640 = Macrotomia echioides.
minima, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, i. (1885) 30.—Persia.
obovata, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 407 (*Lehm. Rel. Bot.* 232).—Turkest.; Ind. or.
Olgæ, *Regel, Pl. Nov. Fedtsch.* 58.—As. centr.
perennis, A. DC. in DC. Prod. x. 95 = Macrotomia perennis.
Severnowi, *Regel, Pl. Nov. Fedtsch.* 58.—As. centr.
tetrastigma, *Forsk. Fl. Aegypt. Arab.* 62.—Arab.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 176.—Tibet. occ.
tibetana, *Kurz, in Journ. As. Soc. Beng.* xlii. (1874) ii. 189.—Tibet. occ.
timotoria, *Forsk. Fl. Aegypt. Arab.* 63.—Aegypt.; Arab.
tinctoria, A. DC. in DC. Prod. x. 96 = Macrotomia perennis.
Vicianii, Coss. & Dur. ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 240 = cornuta.

ARNEDINA, Reichb. Nom. ‘n. 1984’ (1841) = **Arundina**, Blume (Orchid.).

ARNICA, Rupp. ex Linn. Syst. ed. I. (1735). *COMPOSITÆ*, Benth. & Hook. f. ii. 440.
alpina, *Olin & Ladan, Diss.* 11.—Europ.; Am. bor.
alpina, Salisb. Prod. 201 = Doronicum scorpioides.
alpina, Willd. ex Stend. Nom. ed. I. 70 = montana.
altaica, *Turcz. in Bull. S. c. Nat. Mosc.* xx. (1847) ii. 19.—Mont. Altai.
amplexicaulis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 408.—Am. bor. occ.
amplexicaulis, Wall. ex DC. Prod. vi. 314 = Senecio amplexicaulis.
angustifolia, A. Gray, in Am. Journ. Sc. Ser. II. xxxii. (1862) 238 = Parryi.
angustifolia, Turcz. ex Ledeb. Fl. Ross. ii. 622 = montana.
angustifolia, Vahl, Fl. Dan. t. 1594 = alpina.
auriculata, Wild. ex Ledeb. Fl. Ross. ii. 622 = Senecio caucasicus.
austriaca, Hoppe, ex Schur, Enum. Pl. Transs. 240 = Doronicum austriacum.
Bellidiastrum, All. Fl. Pedem. i. 205 = Aster Bellidiastrum.
brevicaulis, *Rafin. New Fl. Am.* ii. 23.—Am. bor.
caffra, *Linn. Syst.* ed. X. 1219.—Afr. austr.
Chamissonis, *Less. in Linnæa*, vi. (1831) 238.—Am. bor.; Unalaschk.
ciliata, Thunb. Fl. Jap. 318 = Hypochœris grandiflorus.
Claytoni, Pursh, Fl. Am. Sept. ii. 527 = undulatifolia.
Clusii, All. Fl. Pedem. i. 205. t. 17. f. 1 = Doronicum Clusii.

ARNICA :—
cordata, E. Mey. ex DC. Prod. vii. 16 = Gerbera tomentosa.
cordata, Thunb. Prod. Pl. Cap. 154 = Gerbera cordata.
cordata, Wulf. in Roem. Arch. iii. iii. 108 = Doronicum caucasicum.
corditolia, *Hook. Fl. Bor. Am.* i. 331.—Am. bor.
coronopifolia, Linn. Sp. Pl. 885 = Gerbera coronopifolia.
corsica, Loisel. Fl. Gall. 376. t. 20 = Doronicum corsicum.
crenata, Thunb. Prod. Pl. Cap. 154 = Mairia crenata.
crocea, Linn. Sp. Pl. 885 = Gerbera Burmanni.
discoides, *Benth. Pl. Hartw.* 319.—Calif.
Doronicum, Jacq. Fl. Austr. i. 57 = Senecio Doronicum.
Doronicum, Pursh, Fl. Am. Sept. ii. 528 = Senecio Pseudo-arnica.
floccosa, Bert. ex DC. Prod. vi. 357 = Senecio arachnoideus.
foliosa, *Nutt. in Trans. Am. Phil. Soc.* N.S. vii. (1841) 407.—Am. bor.
fulgens, Pursh, Fl. Am. Sept. ii. 527 = alpina.
Gerbera, Burm. f. Fl. Cap. Prod. 27 = Gerbera ferruginea.
Gerbera, Linn. Sp. Pl. 885 = Gerbera asplenifolia.
glacialis, Wulf. in Jacq. Coll. i. 280; et Ic. Rar. iii. 15. t. 286 = Doronicum glaciale.
grandis, Thunb. Prod. Pl. Cap. 154 = Alcipe lanata.
helvetica, G. Don, ex Loud. Hort. Brit. 351 = montana.
hirsuta, Forsk. Fl. Aegypt. Arab. 151 = Gerbera hirsuta.
incana, Pers. Syn. ii. 452 = Trichocline incana.
intermedia, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851), 308.—Sibir.
insuloides, Vahl, Symb. Bot. ii. 91 = Heterolepis decipiens.
japonica, Thunb. Fl. Jap. 319 = Senecio japonicus.
lanata, Thunb. Prod. Pl. Cap. 154 = Alcipe lanata.
lanceolata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 407 = Chamissonis.
lanigera, Hook. Prod. Fl. Nesp. (1811) 49 = Senecio arachnoideus.
latifolia, *Bongard, Vég. Isl. Sitch.* 39.—Am. bor. occ.
linifolia, Wall. Cat. n. 2385 = Senecio latifolius.
longifolia, *Eaton, in Bot. King, Exped.* 186.—Am. bor. occ.
macrophylla, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 408 = corditolia.
maritima, Linn. Sp. Pl. 884 = Senecio Pseudo-arnica.
Menziesii, Hook. Fl. Bor. Am. i. 331 = latifolia.
microphylla, Walp. Rep. ii. 653, err. typ. = *macrophylla*.
Mollii, *Schrank, Prim. Fl. Salisb.* 218.—Europ.
mollis, Hook. Fl. Bor. Am. i. 331 = Chamissonis.
montana, Hook. l. c. 330 (pro parte) = foliosa.
montana, *Linn. Sp. Pl.* 884.—Europ.; As. bor.
nevadensis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 55. —Calif.
nivea, Wall. Cat. n. 3148 = Gerbera nivea.
nudicaulis, *Ell. Sketch*, ii. 288.—Am. bor.
obtusifolia, *Less. in Linnæa*, vi. (1831) 236.—Unalaschk.
obtusifolia, Wall. ex DC. Prod. vi. 314 = Senecio amicoides.
operina, Benth. & Less. Fl. Nov. Zel. 231 = sp.
operina, Forst. f. Prod. 56 = Olearia operina.
palmata, Thunb. Fl. Jap. 319 = Senecio syneilesis.
Parryi, *A. Gray, in Am. Nat.* viii. (1874) 218.—Am. bor. occ.
parviflora, *A. Gray, in Proc. Am. Acad.* vii. (1868) 363. —Calif.
pavoniana, Pers. Syn. ii. 452 = Werneria rigida.
petiolata, *Schur, Enum. Pl. Transs.* 242.—Europ.
Pilosella, Burm. ex DC. Prod. vii. 17 (= an Arctotidea? —*Cf. DC. in loc.*).—Afr. austr.
piloselloides, Linn. Pl. Afr. Rar. 22. n. 78 (1760) = Gerbera pilosselloides.
plantaginea, *Pursh, Fl. Am. Sept.* ii. 527.—Am. arct.
planiseginifolia, Gilib. Fl. Lituan. i. 210 = montana.
pyrolæfolia, Pers. Syn. ii. 454 = Gerbera Burmanni.
rotundifolia, Willd. Sp. Pl. iii. 2110 = Bellium rotundifolium.
sachalinensis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 55.—Ins. Sachal.
scorpioides, Adams, ex Ledeb. Fl. Ross. ii. 622 = Senecio cannosicus.

ARNICA :—
scorpioides, *Vacq. Austr.* iv. 26. t. 349.—Europ.
scorpioides, Linn. Sp. Pl. 884 = Doronicum grandiflorum.
senecionides, Fenzl, ex Boiss. Fl. Orient. iii. 396 = Senecio cilicius.
serrata, Thunb. Fl. Cap. ed. Schult. 669 = Gerbera ferruginea.
sinuata, Thunb. Prod. Pl. Cap. 154 = Gerbera ferruginea.
spinulosa, Pers. Syn. ii. 452 = Alcipe tabularis.
styriaca, Vill. Prosp. 33 = Doronicum Clusii.
tabularis, Thunb. Prod. Pl. Cap. 154 = Alcipe tabularis.
tussilaginea, Burm. f. Fl. Ind. 182 = Senecio Kaempferi.
unalaschkensis, *Less. in Linnæa*, vi. (1831) 238.—Ins. Aleut.; Unalaschk.
viscosa, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 374. —Calif.
Wulfeniana, Pollini, in Giorn. Fis. Pav. ix. (1826) 185 = Doronicum Columnæ.

ARNOCORINUM, Endl. & Lehm. ex Lehm. Pl. Preiss. ii. 41 (1846). *LILIACEÆ*, Benth. & Hook. f. iii. 795.
Drummondii, *Endl. ex Lehm. l. c.*—Austral.
glabrum, *Baker, in Journ. Linn. Soc.* xvi. (1880) 416. —Austral.
Preissii, *Lehm. in Preiss.* ii. 42.—Austral.

ARNOGLOSSUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 202 (1821) = **Plantago**, Tourn.
alpinum, S. F. Gray, l. c. 294 = P. alpina.
graminum, S. F. Gray, l. c. 293 = P. maritima.
incanum, S. F. Gray, l. c. 292 = P. media.
lanceolatum, S. F. Gray, l. c. 293 = P. lanceolata.
maritimum, S. F. Gray, l. c. = P. maritima.
subulatum, S. F. Gray, l. c. 294 = P. Coronopus.

ARNOGLOSSUM, Rafin. Fl. Ludov. 64 (1817). *COMPOSITÆ*.
plantagineum, Rafin. l. c. 65 (quid?).—Am. bor.

ARNOLDIA, Blume, Bijdr. 868 (1826) = **Weinmannia**, Linn. (Saxifr.)
fraxinifolia, Blume, in Flora, xli. (1858) 254 = W. fraxinifolia, Hook. f.
heterophylla, Blume, Bijdr. 868 = W. Blumei, *A. Gray*.
pinnata, Blume, l. c. = W. Blumei, *Planch.*

ARNOLDIA, Cass. in Dict. Sc. Nat. xxx. 320 (1824) = Dimorphotheca, Moench (Compos.).
aurea, Cass. l. c. 321 = D. chrysanthemifolia.
chrysanthemifolia, Steud. Nom. ed. II. i. 256 = D chrysanthemifolia.
Tragus, Steud. Nom. ed. II. i. 256 = D. Tragus.
Ventenatiana, Presl, Repert. 192, in nota = D. chrysanthemifolia.
viscosa, Steud. Nom. ed. II. i. 256 = D. cuneata.

ARNOPOGON, Willd. Sp. Pl. iii. 1496 (1803) = **Uropetes**, Desf. (Compos.).
asper, Willd. l. c. 1497 = U. picroides.
caprosia, Willd. l. c. = U. picroides.
Dalechampii, Willd. l. c. 1496 = U. Dalechampii.
picroides, Willd. l. c. = U. picroides.

ARNOSERIS, Gaertn. Fruct. ii. 355. t. 157 (1791). *COMPOSITÆ*, Benth. & Hook. f. ii. 507.
foetida, Dum. Fl. Belg. 63 = Crepis foetida.
gracilis, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 301 = Apogon gracilis.
humilis, Sch. Bip. l. c. = Apogon humilis.
minima, Dum. Fl. Belg. 63 = pusilla.
pusilla, *Gaertn. Fruct.* ii. 355. t. 157.—Europ.

ARNOTTIA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 29. *ORCHIDEÆ*, Benth. & Hook. f. iii. 492.
isomeria, *S. Moore, in Baker, Fl. Maurit.* 330.—Ins. Mascar.
mauritiana, *A. Rich. in Mém. Soc. Hist. Nat. Par.* iv. (1828) 30.—Ins. Mascar.

AROMADENDRON, Blume, Bijdr. 10 (1825) = **Talauma**, Juss. (Magnol.).
elegans, Blume, l. c. 10 = T. elegans.
glaucum, Korth. ex Miq. Fl. Ind. Bat. i. 2. 16 = T. glauca.

AROMADENDRON, Andr. ex Steud. Nom. ed. II. i. 134 (1840) et AROMADENDRUM, Andr. ex Steud. l. c. 696, in syn. (1840) = **Eucalyptus**, L'Hérit. (Myrtac.).

AROMIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 395 = **Amblyopappus**, Hook. (Compos.). *tenuifolia*, Nutt. l. c. 396 = Amblyop. pusillus.

ARON, Adans. Fam. ii. 470 (1763).
dracunculum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = Dracunculus vulgaris.
colocasium, St. Lag. l. c. = Colocasia antiquorum.

ARONGANA, Choisy, in DC. Prod. i. 541 (1824) = **Harongs**, Thou. (Hyperic.).

ARONIA, Mitchell, in Act. Acad. Nat. Car. viii. (1748) 905, App. = **Orontium**, Linn. (Aroid.).

ARONIA, Medic. Phil. Bot. i. 140 (1789) = **Cotoneaster**, Lindl. (Rosac.).

ARONIA, Pers. Syn. ii. 39 (1807) = **Amelanchier**, Lindl. (Rosac.).
alnifolia, Nutt. Gen. Am. i. 306 = Amel. alnifolia.
Amelanchier, Dum. Fl. Belg. 92 = Amel. vulgaris.
arborea, Barton, Comp. Fl. Phil. i. 92 = Amel. canadensis.
arbutifolia, Medic. Phil. Bot. i. 140 = Pyrus arbutifolia.
aria-chamaemespilus, Reichb. Fl. Germ. Excurs. 630 = Pyrus Chamaemespilus.
asiatica, Sieb. & Zucc. Fl. Jap. i. 87. t. 42 = Amel. canadensis.
Botryapium, Pers. Syn. ii. 39 = Amel. canadensis.
Chamaemespilus, Pers. l. c. = Pyrus Chamaemespilus.
cordata, Rafin. Med. Fl. ii. 196 = Amel. canadensis.
cretica, Pers. l. c. 40 = Amel. vulgaris.
densiflora, Spach, Hist. Veg. Phan. ii. 88 = Pyrus arbutifolia.
depressa, Spach, l. c. 90 = Pyrus arbutifolia.
floribunda, Spach, l. c. 89 = Pyrus arbutifolia.
floribunda, Sweet, Hort. Brit. ed. I. 485 = Pyrus spectabilis.
glabrescens, Spach, Hist. Vég. Phan. ii. 89 = Pyrus arbutifolia.
grandiflora, Sweet, Hort. Brit. ed. II. 177 = Pyrus arbutifolia.
Hostii, Carr. in Rev. Hortic. (1878) 470 = Pyrus intermedia.
japonica, Hort. ex C. Koch, Dendrol. i. 179 = Pyrus Aronica.
latifolia, Riddell, Suppl. Cat. Fl. Ohio, 24 = Amel. canadensis ?
melanocarpa, Nutt. ex Ell. Sketch, i. 556 = Pyrus arbutifolia.
nivea, Neum. ex Tausch, in Flora, xxi. (1838) 714 = Amel. canadensis.
ovalis, Pers. Syn. ii. 39 = Amel. canadensis.
petraea, Wallr. Beitr. Fl. Hercyn. 206.—Europ.
prunoides, Neum. ex Tausch, in Flora, xxi. (1838) 715 = Amel. canadensis.
pubescens, Spach, Hist. Vég. Phan. ii. 90 = Pyrus arbutifolia.
pumila, M. Roem. Syn. iii. Rosifl. 161 = Pyrus arbutifolia.
pyrifolia, Lam. Encyc. i. 58 = Pyrus arbutifolia.
racemosa, Schur, Enum. Pl. Transs. 207 = Amel. ovalis.
rotundifolia, Pers. Syn. ii. 39 = Amel. vulgaris.
sanguinea, Nutt. Gen. Am. i. 306 = Amel. canadensis.
sorbifolia, Spach, Hist. Vég. Phan. ii. 87 = Pyrus aparia.
aparia, M. Roem. Syn. iii. Rosifl. 158 = Pyrus aparia.
subcordata, Rafin. ex DC. Prod. ii. 632 = Amel. canadensis.
taurandina, Wallr. Beitr. Fl. Hercyn. 206 = Amel. vulgaris ?
Tournefortiana, M. Roem. Syn. iii. Rosifl. 160 = Pyrus pinnatifida.

ARONICUM, Neck. Elem. i. 27 (1790) = **Doronicum**, Linn. (Compos.).
alpinum, DC. Prod. vi. 230 = D. alpicum.

ARONICUM:—
Bauhini, Reichb. f. Ic. Fl. Germ. xvi. 32 = D. hirsutum.
carpaticum, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 187 = D. grandiflorum.
Clusii, Koch, Syn. ed. I. 882 = D. hirsutum.
cordatum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 41 = D. cordatum.
corsicum, DC. Prod. vi. 219 = D. corsicum.
Doronicum, Reichb. Fl. Germ. Excurs. 238 = D. Clusii.
glaciale, Reichb. l. c. 234 = D. glaciale.
latifolium, Reichb. l. c. = D. grandiflorum.
scorpioides, DC. Prod. vi. 219 ; Koch, Syn. ed. I. 882 = D. grandiflorum.
scorpioideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = D. grandiflorum.
wiesnum, Freyn & Gautier, in Bull. Soc. Bot. Fr. xxviii. (1881) 50.—Europ.

AROSMA, Rafin. Fl. Tellur. iii. 66 (1836) = **Celadium**, Vent. (Aroid.).
acutifolia, Rafin. l. c. = C. bicolor !
obtusifolia, Rafin. l. c. = C. bicolor !

AROTON, Neck. Elem. ii. 336 (1790) = **Croton**, Linn. (Euphorb.).

AROUNA, Aubl. Pl. Gui. i. 16. t. 5 (1775) = **Dialium**, Linn. (Legumin.).
divaricatum, Willd. Sp. Pl. i. 556 = D. divaricatum.
guianensis, Aubl. Pl. Gui. i. 16 = D. divaricatum.

ARPITIUM, Neck. Elem. i. 168 (1790) = **Ldgustioum**, Linn. (Umbell.).
simplex, Sweet, Hort. Brit. ed. II. 591 = L. simplex.

ARPOPHYLLUM, Llave & Lex. Nov. Veg. Descr. fasc. ii. (Orch. Opusc.) 19 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 492.
alpinum, *Lindl. in Benth. Pl. Hartw.* 98 (1849).—Mexic.
cardinale, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 282.—Mexic.
giganteum, *Hartw. ex Lindl. in Ann. Nat. Hist.* iv. (1840) 384.—Mexico.
medium, *Reichb. f. Beitr. Orch. Centr. Am.* 89.—Mexico.
spicatum, *Llave & Lex. Nov. Veg. Descr.* fasc. ii. (*Orch. Opusc*). 20.—Mexico.
squarrosum, Hort. Dorat. ex Lubbers, Cat. Pl. Rar. San Donato (1880) 15 = spicatum.

ARRABIDAEA, DC. in Bibl. Univ. Genev. xvii. (1838) 126 (Rev. Bign. 10). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1028.
CHARMIA, Schott, in Spreng. Syst. Cur. Post. 409 (1827).
VASCONCELLIA, Mart. in Flora, xxiv. (1841) II. Beibl. 13 (Herb. Fl. Bras. 252).
acutiolia, *DC. Prod.* ix. 185.—Bras.
Agnus-castus, *DC. l. c.* 188.—Bras.
argentea, *Wawra, in Oestr. Bot. Zeitschr.* (1879) 216. —Bras.
Bianchetti, DC. Prod. ix. 186 = Bignonia Bianchetti.
candicans, *DC. l. c.* 185.—Guiana.
Chica, Verl. in Rev. Hortic. (1868) 154 = Bignonia Chica.
Clausseni, *DC. l. c.* 186.—Bras.
conjugata, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 46.—Bras.
cuneifolia, *DC. Prod.* ix. 188.—Bras.
cordifolia, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1158.—Guiana.
Cuscutarum, *DC. Prod.* ix. 185.—Bras.
fasciculata, *DC. l. c.*—Bras.
florida, *DC. l. c.* 184.—Bras.
obovata, *DC. l. c.* 185.—Bras.
Orbigeana, *DC. l. c.* 184.—Bolivia.
paniculata, *Seem. in Trans. Linn. Soc.* xxiii. (1862) 14. —Ins. Martinic.
platyphylla, *DC. Prod.* ix. 186.—Bras.
Plaei, *DC. l. c.* 184.—Venezuela.
pruinosa, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1158. —Guiana.
roseo, *DC. Prod.* ix. 188.—Bras.
rubricaulis, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 198.—Bras.
Salzmanni, *DC. Rev. Bign.* 10. (1838).—Bras.

ARRABIDAEA,—
Sege, DC. in Bibl. Univ. Genèv. xvii. (1838) 126 ; et Prod. ix. 188 = A. Agnus-castus.
Sieberi, *DC. l. c.* 186.—Bras.
subincana, DC. l. c. 184 = Bignonia subincana.
tuberculata, *DC. l. c.*—Bras.
virescens, *DC. l. c.*—Bras.

ARRABIDAEA, Steud. Nom. ed. II. i. 135 (1840) = Cormonema, Reissek (Rhamn.).
spinosa, Steud. l. c. = C. spinosum.

ARRACACHA, DC. in Bibl. Univ. Genèv. xi. (1829) 74 ; et Prod. iv. 243 (1830) = seq.

ARRACACIA, Bancr. in Jamaic. Journ. iv. (1836) 18. *UMBELLIFERAE*, Benth. & Hook. f. i. 884.
ARRACACHA, DC. in Bibl. Univ. Genev. xi. (1829) 74 ; et Prod. iv. 243 (1830).
BANCROFTIA, Billb. in Linnæa Samfund. Handl. i. 1859 (1838) 89.
DEWEYA, Torr. & Gray, Fl. N. Am. i. 641 (1840).
PENTACRYPTA, Lehm. in Linnæa, v. (1830) 380. t. 8.
VELARA, DC. Prod. iv. 230 (1830).
acuminata, *Benth. Pl. Hartw.* 187.—Mexico.
atropurpurea, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* i. 564.—Mexico.
decumbens, *Benth. & Hook. f. l. c.*—Mexico.
esculenta, DC. Prod. iv. 244 = xanthorrhiza.
glaucescens, *Benth. Pl. Hartw.* 187.—Mexico.
moschata, *DC. Prod.* iv. 244.—Am. merid.
tolucensis, *Hemsl. Biol. Centr. Am. Bot.* i. 564.—Mexico.
xanthorrhiza, *Bancr. in Jamaic. Journ.* iv. (1836) 18. —Jamaica.

ARRHENACHNE, Cass. in Dict. Sc. Nat. iii. 253 (1828) = **Bacchariss**, Linn. (Compos.).
juncea, Cass. l. c. = B. juncea.

ARRHENATHERUM, Beauv. Agrost. 55. t. 11. f. 5 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1181.
americanum, Beauv. l. c. 56 = Avena pensylvanica.
asperum, Opiz, in Lotos, iii. (1853) 66 = avenaceum.
avenaceum, *Beauv. Agrost.* 55. t. 11. f. 5.—Europ.
avenaceum, Boiss. Voy. Bot. Espagne, ii. 657 = erianthum.
biaristatum, Peterm. Fl. Lips. Excurs. 106 = avenaceum.
bulbosum, Kunze, ex Anderss. in Walp. Ann. vi. 1008 = erianthum.
bulbosum, Presl, Cyp. et Gram. Sic. 29 = avenaceum.
cechicum, Opiz, in Lotos, iii. (1853) 66 = avenaceum.
elatius, Beauv. Agrost. 56 = avenaceum.
erianthum, *Boiss. & Reut. Diag. Pl. Hispan.* 121. —Lusit. Alg.
exaveus, Opiz, in Lotos, iii. (1853) 65 = avenaceum.
kentuckense, *Torr. ex Steud. Nom.* ed. II. i. 136.—Am. bor.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 122.—Orient.
longifolium, Dulac, Fl. Hautes-Pyr. 78 = pallens.
palaestinum, Boiss. Diagn. Ser. I. xiii. 51 = avenaceum.
pallens, *Link, Hort. Berol.* i. 132.—Lusit.; Hisp.
pennsylvanicum, Torr. Fl. N. Am. i. 130 = Avena pensylvanica.
precatorium, Beauv. Agrost. 56 = avenaceum.
rivulare, *Opiz, in Lotos*, iii. (1853) 65.—Europ.
rupestre, *Dum. Obs. Gram. Belg.* 146.—Europ.
Thorei, Desm. Cat. Dord. (1849) = pallens.
vaudilianum, Opiz, in Lotos, iii. (1853) 66 = avenaceum.

ARRHOSTOXYLON, Mart. ex Nees, in Mart. Fl. Bras. ix. 57. t. 6 et 209 (1847) = seq.

ARRHOSTOXYLUM, Mart. ex Nees, in DC. Prod. xi. 209 (1847) = Riedlia, Linn. (Acanth.).
echinocalyx, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 131 = R. echinocalyx.
acutecuspidatum, Mart. ex Nees, in Mart. Fl. Bras. ix. 58 = R. acutecuspida.
affine, Nees, in Mart. Fl. Bras. ix. 59.—Bras.
album, Nees, l. c. 60 = Riedlia alba.
amplexicaule, Nees, l. c. 59.—Bras.
ciliatiflorum, Nees, in DC. Prod. xi. 210 = R. ciliatiflora.

ARRHOSTOXYLUM :—
coccineum, Nees, in Mart. Fl. Bras. ix. 63 = R. coccineus.
costatum, Nees, l. c. 61 = R. costatus.
curviflorum, Nees, l. c. 62 = R. curviflorus.
formosum, Nees, in DC. Prod. xi. 215 = R. formosus.
fulgidum, Nees, in Mart. Fl. Bras. ix. 59 = R. fulgidus.
glabrum, Nees, l. c. 58.—Bras.
Hoenkeanum, Nees, in DC. Prod. xi. 211.—Peruv.
Hortogianum, Nees, l. c. 209.—N. Granat.
inflatum, Nees, l. c. 215 = R. inflatus.
jussieuoides, Nees, l. c. 212 = R. jussieuoides.
laxum, Nees, in Mart. Fl. Bras. ix. 57.—Bras.
longiflorum, Nees, l. c. 62 = sq.
longifolium, Nees, in DC. Prod. xi. 214 = R. longifolia.
microphyllum, Nees, in Mart. Fl. Bras. ix. 61 = Calophanes microphylla ?
nitidum, Nees, l. c. 59.—Bras.
pedunculosum, Nees, in DC. Prod. xi. 211 = R. pedunculosa.
repens, Mart. Fl. Bras. ix. 60.—Bras.
roseum, Nees, l. c. 61.—Bras.
rubrum, Nees, in DC. Prod. xi. 209 = R. rubra.
silvaticola, Nees, in Mart. Fl. Bras. ix. 60 = R. formosa.
splendidulum, Nees, l. c.—Bras.
stemonacanthoides, Nees, ex Kjoeb. Vidensk. Meddel. (1854) 130 = K. stemonacanthoides.
subsessile, Nees, in Mart. Fl. Bras. ix. 58.—Bras.
terminale, Nees, in DC. Prod. xi. 214.—Peruv.
violaceum, Nees, l. c. 216 = R. violacea.
Willdenovianum, Nees, l. c. 209.—Ind. occ.

ARRHYNCHIUM, Lindl. in Paxt. Flow. Gard. i. 142 (1850 51) = **Arachnanthe**, Blume (Orchid.).
labrosum, Lindl. & Paxt. l. c. = Arachn. bilingvis.

ARROSTIA, Rafin. Caratteri, 75 (1810) = **Gypsophila**, Linn. (Caryophyll.).
altissima, Rafin. Fl. Tellur. ii. 54 = G. altissima.
dichotoma, Rafin. Caratteri, 75 = G. Arrostii.
paniculata, Rafin. Fl. Tellur. ii. 53 = G. paniculata.

ARROWSMITHIA, DC. Prod. vii. 254 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 329.
stypheloides, *DC. l. c.*—Afr. austr.

ARROZIA, Schrad. ex Kunth, Enum. Pl. i. 11 (1833) = **Luziola**, Juss. (Gramin.).
micrantha, Schrad. l. c. (= *Caryochloa brasiliensis*). —Bras.

ARRUDEA, A. St. Hil. & Camb. Fl. Bras. Mer. i. 318. t. 66 (1827) = **Cluzia**, Linn. (Guttif.).
bicolor, Benth. in Hook. Kew Journ. iii. (1851) 166 = C. bicolor.
clusioides, A. St. Hil. et Camb. Fl. Bras. Mer. i. 319.—Bras.
purpurea, Splitg. in Hoev. & De Vriese, Tijdschr. ix. (1842) 109.—Guiana.

ARSACE, Four. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 118 = **Erica**, Linn.
arborea, Four. l. c. = E. arborea.

ARSIS, Lour. Fl. Cochinch. 335 (1790) = **Grewia**, Linn. (Tiliac.).
rugosa, Lour. l. c. = G. Microcos.

ARTABOTRYS, R. Br. in Bot. Reg. t. 493 (1820). *ANONACEAE*, Benth. & Hook. f. i. 24.
ROPALOPETALUM, Griff. Notul. iv. 18 (1845).
Blumei, Benth. Fl. Hongk. 10 = honkongensis.
Blumei, *Hook. f. & Thoms. Fl. Ind.* i. 128.—Java.
brachypetalus, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 467.—Afr.
burmanicus, *DC. in Mém. Soc. Phys. Genèv.* v. (1832) 212.—Burma.
caudatus, *Wall. Cat.* n. 6417.—Reg. Himal.
crassifolius, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 54.—Burma ; Malaya.
Cuminginus, *Vidal, Phan. Cuming. Philipp.* 109.—Ins. Philipp.
densiflorus, *Wall. ex Voigt, Hort. Suburb. Calc.* 16, nomen.—Ind. or.
hamatus, Benth. in Hook. Kew Journ. iii. (1851) 257 = hookongensis.
ARTABOTRYS:—
hamatus, Blume, Fl. Jav. Anon. 60. t. 29 = odoratissimus.
Hildebrandtii, *O. Hoffm. Sert. Pl. Madag.* 3.—Madag.
hookongensis, *Hance, in Journ. Bot.* viii. (1870) 71. —Hongkong.
inodorus, *Zipp. in Ann. Mus. Bot. Lugd. Bat.* ii. 41. —N. Guinea.
intermedius, *Hasск. Cat. Pl. Hort. Bogor.* 176.—Java.
Korui, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 54.—Burma.
macrophyllus, *Hook. f. in Hook. Niger Fl.* 307.—Afr. trop.
madagascariensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 42.—Madag.
Maingayi, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 55.—Malacca.
malayanus, Griff Notul. iv. 713 = Xylopia ferruginea.
odoratissimus, Blume, Fl. Jav. Anon. 59 = hookongensis.
odoratissimus, *R. Br. in Bot. Reg.* t. 493.—Ind. or. ; Malaya.
parviflorus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 41.—Sumatr.
pleurocarpus, *Maing. ex Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 54.—Malacca.
polygynus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 41.—Borneo.
siamensis, *Miq. l. c.* 42.—Siam.
speciosus, *Kurz, ex Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 55.—Ins. Andam.
suaveolens, *Blume, Fl. Jav. Anon.* 62.—Ind. or. ; Malaya.
sumatranus, *Miq. Fl. Ind. Bat. Suppl.* i. 375.—Sumatr.
Thomsoni, *Oliver, Fl. Trop. Afr.* i. 29.—Afr. trop.
unguicalatus, *Wall. Cat.* n. 6415 F.—Ind. or.
uncatus, Bull. Hist. Pl. i. 282 = odoratissimus.
zeylanicus, *Hook. f. & Thoms. Fl. Ind.* i. 128.—Ind. or.

ARTANEMA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 224 (1829). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 958.
ACHIMENES, Vahl, Symb. Bot. fi. 71 (1791).
DICEROS, Pers. Syn. ii. 164 (1807).
angustifolium, *Benth. Scroph. Ind.* 39.—Singapore.
fimbriatum, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 224.—Austral.
sesamoides, *Benth. Scroph. Ind.* 39.—Ind. or.

ARTANTHE, Miq. Comm. Ph./t. 40. t. 7, 8 (1840) = **Piper**, Linn.
abutilifolia, Miq. Syst. Pip. 385 = P. abutilifolium.
abutiloides, Miq. l. c. 384 = P. abutiloides.
acutifolia, Miq. l. c. 428 = P. acutifolium.
adenandra, Miq. l. c. 515 = P. adenandrum.
adenophora, Miq. l. c. 514 = P. adenophorum.
adenophylla, Miq. in Linnaea, xx. (1847) 170 = P. colubrinum.
adunca, Miq. Comm. Phyt. 49. t. 4 = P. aduncum.
aequalis, Miq. Syst. Pip. 511 = P. aequale.
aequirophylla, Miq. l. c. 391 = P. cardiophyllum.
affinis, Miq. in Linnaea, xviii. (1844) 238 = P. affine.
alaria, Miq. Syst. Pip. 406 = P. alare.
albida, Miq. l. c. 415 = P. albidum.
alnoides, Miq. l. c. 462 = P. alnoides.
alpina, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 451 = P. alpinum.
alveolata, Miq. Syst. Pip. 406 = P. alveolatum.
amasonica, Miq. in Linnaea, xx. (1847) 146 = P. amasonicum.
ampla, Miq. Syst. Pip. 501 = P. Rohrii.
amplectens, Miq. in Linnaea, xx. (1847) 154 = P. amplectens.
amplicaule, Miq. Syst. Pip. 415 = P. amplexicaule.
angusta, Miq. l. c. 396 = P. angustum.
angustifolia, Miq. l. c. 398 = P. consanguineum.
anonaefolia, Miq. l. c. 504 = P. anonaefolium.
apiculata, Klotzsch, in Hook. Journ. Bot. iv. (1845) 321 = P. anonaefolium.
appendiculata, Benth. Fl. Hartw. 284 = P. appendiculatum.
aronata, Miq. Syst. Pip. 538 = P. Opizianum.
arvalis, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 454 = P. arvalense.
aspera, Miq. l. c. 466 = P. hirsutum.
ARTANTHE :—
asperifolia, Miq. Syst. Pip. 441 = Piper hirsutum.
asperiuscula, Miq. l. c. 464 = P. hirsutum.
auriculata, Miq. in Linnaea, xx. (1847) 167 = P. auriculatum.
aurita, Miq. Syst. Pip. 400 = P. auritum.
avellana, Miq. l. c. 478 = P. avellanum.
baccana, Miq. in Linnaea, xx. (1847) 166 = P. baccans.
bahiensis, Presl, Epim. Bot. 237 = P. bahiense.
barbata, Miq. Syst. Pip. 434 = P. barbatum.
Barclayana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 460 = P. hirsutum.
Bartlingiana, Miq. Syst. Pip. 510 = P. Bartlingianum.
Brecheyana, Miq. l. c. 408 = P. elliptieum.
begoniaefolia, Miq. l. c. = P. begoniaefolium.
Benthamiana, Miq. in Hook. Lond. journ. Bot. iv. (1845) 447 = P. Benthamianum.
berbisensis, Miq. Syst. Pip. 500 = P. Hostmannianum.
Blanchetii, Miq. l. c. 478 = P. Blanchetii.
Botrytis, Miq. l. c. 538 = P. Botrytis.
brachypoda, Benth. Bot. Voy. Sulph. 167. t. 54 = P. brachypodon.
brachystachya, Miq. ex C. DC. in DC. Prod. xvi. t. 296 = P. brachystachyum.
brasiliensis, Miq. in Linnaea, xx. (1847) 140.—Bras.
Bredemeyeri, Miq. Syst. Pip. 429 = P. Bredemeyeri.
brevispicata, Miq. l. c. 508 = P. brevispicatum.
Brongniartii, Miq. l. c. 486 = P. Brongniartii.
Brunoniana, Miq. l. c. 509 = P. lanceolatum.
caladiifolia, Miq. l. c. 387 = P. caladiifolium.
caldasiana, Miq. in Verh. Kon. Akad. Wetensch. II. v. (1871) 337 ; et in Arch. Neerl. vi. (1871) 176.—Bras.
calocoma, Miq. in Seem. Bot. Voy. Herald, 199 = P. calocoma.
Cambessedei, Miq. in Linnaea, xx. (1847) 161 = P. Cambessedeii.
caracasana, Miq. Syst. Pip. 398 = P. caracasanum.
cardiophylla, Presl, Epim. Bot. 225 = P. cardiophyllum.
caripanis, Miq. Syst. Pip. 522 = P. caripense.
casapinnis, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 468 = P. casapinense.
casinioides, Miq. Syst. Pip. 490 = P. casinioides.
catalpaefolia, Miq. in Verh. en Meded. Kon. Nederl. Inst. 1842 (1843) 197 = P. marginatum.
caudata, Miq. l. c. 196 = P. marginatum.
cearrasis, Miq. Syst. Pip. 456 = P. angustifolium.
celtidifolia, Miq. l. c. 452 = P. celtidifolium.
cernua, Presl, Epim. Bot. 225 = P. cernuum.
Chamissonis, Miq. Syst. Pip. 487 = P. Chamissonis.
chaviciodes, Miq. in Linnaea, xx. (1847) 135 = P. chaviciodes.
chelsvana, Miq. Syst. Pip. 533 = P. tomentosum.
chimonanthifolia, Miq. l. c. 466 = P. chimonanthifolium.
Churumaya, Miq. l. c. 481 = P. Churumaya.
cinnamomifolia, Miq. l. c. 419.—Peruv.
Claussenianum, Miq. in Linnaea, xx. (1847) 153 = P. Claussenianum.
coccoloboides, Miq. Syst. Pip. 397 = P. coccoloboides.
colubrina, Miq. l. c. 512 = P. colubrinum.
concinna, Miq. in Linnaea, xx. (1847) 151 = P. concinnum.
consanguinea, Miq. Syst. Pip. 398 = P. consanguineum.
controversa, Miq. l. c. 448 = P. hispidum.
cornifolia, Miq. l. c. 479 = P. cornifolium.
coruscans, Miq. l. c. 411 = P. coruscans.
corylifolia, Klotzsch, in Hook. Journ. Bot. iv. (1845) 321 = P. dilatatum.
corylifolia, Miq. Syst. Pip. 474 = P. dilatatum.
crassistipulata, Miq. l. c. 551 = P. crassistipulatum.
crenata, Miq. l. c. 429 = P. crocatum.
cumanensis, Miq. l. c. 484 = P. cumanense.
curvata, Miq. l. c. 528 = P. curvatum.
cuspidata, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 466 = P. pseudo-peruvianum.
dactyoura, Miq. l. c. 452 = P. dactyoura.
dactyoda, Miq. l. c. 453 = P. dactyoides.
decipiens, Miq. Syst. Pip. 462 = P. decipiens.
decurrens, Miq. l. c. 403 = P. tuberculatum.
decurrens, Four. in Illustr. Hortic. xxiii. (1876) t. 289 = P. decurrens.
demerarana, Miq. in Hook. Lond. Journ. Bot iv. (1845) 464 = P. demerarianum.
densiflora, Miq. in Linnaea, xx. (1847) 179 = P. densiflorum.
dimidiata, Presl, Epim. Bot. 225 = P. dimidiatum.
diospyrifolia, Miq. Syst. Pip. 477 = P. diospyrifolium.
diversyna, Presl, Epim. Bot. 238 = P. diversynum.
dubia, Miq. Syst. Pip. 555 = P. chilense.

ARTANTHE :—
Durvilleana, Miq. Syst. Pip. 507 = P. anonaefolium.
elongata, Miq. l. c. 484 = P. angustifolium.
encheoides, Miq. l. c. 412 = P. crassinervium.
Endlicheriana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 447 = P. formosum.
eriopoda, Miq. l. c. 459 = P. eriopodon.
eucalyptifolia, Miq. Syst. Pip. 505 = P. eucalyptifolium.
excavata, Miq. l. c. 584 = P. excavatum.
eximia, Miq. l. c. 393 = P. obliquum.
exserens, Miq. in Linnaea, xx. (1847) 148 = P. crassinervium.
flagellaris, Miq. Syst. Pip. 393 = P. nobile.
flexicaulis, Miq. l. c. 538 = P. flexicaule.
flexuosa, Miq. l. c. 454 = P. flexuosum.
formosa, Miq. l. c. 386 = P. formosum.
fuliginea, Miq. l. c. 455 = P. fuligineum.
fuscescens, Miq. in Linnaea, xx. (1847) 144 = P. estrellense.
Gabrieliana, Miq. Syst. Pip. 476 = P. Gabrielanum.
Galeottii, Miq. l. c. 451 = P. aduncum.
Gaudichaudiana, Miq. l. c. 469 = P. Gaudichaudianum.
geniculata, Griseb. Cat. Pl. Cub. 65 = P. rigidum.
geniculata, Miq. in Versl. en Medel. Kon. Nederl. Inst. '1842' (1843) 194 = P. geniculatum.
glabella, Miq. Syst. Pip. 518 = P. glabellum.
glabrata, Miq. l. c. 481 = P. cumanense.
glabrescens, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 461 = P. glabrescens.
gracilis, Miq. Syst. Pip. 446 = P. gracile.
grandifolia, Miq. l. c. 408 = P. grandifolium.
granulosa, Miq. l. c. 435 = P. unguiculatum.
guianensis, Klotzsch, in Hook. Journ. Bot. iv. (1842) 324 = P. guianense.
Guillemininiana, Miq. Syst. Pip. 409 = P. crassinervium.
Haenkeana, Miq. in Linnaea, xx. (1847) 145 = P. Paulianum.
halesiaefolia, Miq. Syst. Pip. 419 = P. halesiaefolium.
Hartwegiana, Benth. Pl. Hartw. 254 = P. Hartwegianum.
heterophylla, Miq. Syst. Pip. 502 = P. heterophyllum.
hexagyna, Miq. l. c. 419 = P. Betle.
Hilariana, Miq. l. c. 491 = P. Hilarianum.
hirsuta, Miq. l. c. 446 = P. hirsutum.
hirtula, Presl, Epim. Bot. 226 = P. hirtulum.
hispida, Miq. Comm. Phyt. 50. t. viii. f. A = P. asperifolia.
Hookeriana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 446 = P. Hookerianum.
Hostmanniana, Miq. l. c. 465 = P. Hostmannianum.
hymenophylla, Miq. Syst. Pip. 482 = P. Kegelii.
imperialis, Miq. in Scem. Bot. Voy. Herald, 198. t. 38 = P. imperiale.
insignis, Miq. Syst. Pip. 394 = P. insigne.
Jacquemontiana, Miq. l. c. 468 = P. citrifolium.
jalapensis, Miq. l. c. 444 = P. jalapense.
jamaicensis, Griseb. Fl. Brit. W. Ind. 170 = P. geniculatum.
japurensis, Miq. in Linnaea, xx. (1847) 168 = P. japurense.
jamiensis, Miq. Syst. Pip. 401 = P. javitense.
Jussiaeana, Miq. l. c. 420 = P. Jussiaeanum.
Karwinskiana, Miq. l. c. 476 = P. Karwinskianum.
Kegeliana, Miq. in Linnaea, xxii. (1849) 77 = P. Kegelianum.
Kunthiana, Miq. Syst. Pip. 453 = P. hirsutum.
lacunosa, Miq. l. c. 427 = P. lacunosum.
lanceafolia, Miq. l. c. 438 = P. lanceaefolium.
lanceata, Miq. l. c. 526 = P. lanciflolium.
lapathifolium, Miq. l. c. 403 = P. lapathifolium.
laurifolia, Miq. l. c. 530 = P. lauriflolium.
Lehmannsiana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 466 = P. salicariaefolium.
lentaginoides, Miq. Syst. Pip. 520 = P. rivinoides.
Leprieurii, Miq. l. c. 536.—Guiana.
lepturu, Miq. l. c. 532 = P. lepturum.
Lessertiana, Miq. l. c. 405 = P. lapathifolium.
leucophylla, Miq. l. c. 466 = P. verbaecifolium.
Lhotzkyana, Miq. l. c. 417 = P. Lhotzkyanum.
Lindeniana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 458 = P. Actanthe.
linoides, Miq. Syst. Pip. 488 = P. linoides.
linguifolia, Miq. l. c. 538 = P. linguaefolium.
Lucasiana, Miq. l. c. 563 = P. Lucasianum.
Luschnathiana, Miq. l. c. 494 = P. geniculatum.

ARTANTHE :—
macrophylla, Griseb. Fl. Brit. W. Ind. 171 = P. geniculatum.
macrophylla, Miq. Syst. Pip. 498 = P. macrophyllum.
macrorhyncha, Miq. in Linnaea, xx. (1847) 173 = P. macrorhynchon.
macrostachya, Presl, Epim. Bot. 227 = P. macrostachyum.
magnifica, Linden, ex Fourn. in Illustr. Hortic. xxiii. (1876), 230 = P. decurrens.
magnifica, Miq. Syst. Pip. 391 = P. obliquum.
malacophylla, Presl, Epim. Bot. 227 = P. malacophyllum.
marginata, Miq. Syst. Pip. 381 = P. marginatum.
Martiana, Miq. l. c. 422 = P. Martianum.
Martinicae, Miq. l. c. 413.—Ins. Martinica.
Mauritiana, Miq. in Linnaea, xx. (1847) 169 = P. juliflorum.
melanosticta, Miq. Syst. Pip. 404 = P. melanostictum.
mexicana, Miq. l. c. 458 = P. mexicanum.
Meyeniana, Klotzsch, ex Miq. l. c. 524 = P. lepturum.
Meyeri, Miq. l. c. 454 = P. divaricatum.
microstachya, Miq. l. c. 527 = P. microstachyon.
Miersiana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 449 = P. Miersianum.
Mikaniana, Miq. Syst. Pip. 388 = P. Parthenium.
mitis, Miq. l. c. 407 = P. mite.
modesta, Miq. l. c. 517 = P. modestum.
mollicoma, Miq. l. c. 488 = P. mollicomum.
mollis, Miq. in Linnaea, xx. (1847) 165 = P. molliplilosum.
Moritziana, Miq. l. c. xviii. (1844) 714 = P. Moritzianum.
nitida, Miq. Comm. Phyt. 47. t. vii = P. nitidum.
nitidula, Miq. Syst. Pip. 534 = P. nitidulum.
nubigena, Miq. l. c. 433 = P. crocatum.
obesa, Miq. l. c. 470 = P. amalago.
oblonga, Miq. l. c. 480 = P. oblongum.
oblongata, Miq. l. c. 536 = P. oblongatum.
oblongifolia, Klotzsch, in Hook. Journ. Bot. iv. (1842) 323 = P. oblongifolium.
obovata, Miq. Syst. Pip. 508 = P. colubrinum.
obtusa, Miq. l. c. 416 = P. fluminense.
obtusifolia, Miq. in Linnaea, xx. (1847) 142 = P. obumbratum.
Olfersiana, Klotzsch, in Hook. Journ. Bot. iv. (1842) 324 = P. hirsutum.
Opiisii, Miq. Syst. Pip. 440 = P. asperifolium.
opulifolia, Miq. in Linnaea, xx. (1847) 160 = P. Candebeadedesii.
paraense, Miq. l. c. 156 = P. paraënse.
Parkeriana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 46 = P. Parkerianum.
patens, Miq. Syst. Pip. 534 = P. nervosum.
patula, Miq. l. c. 554 = P. patens.
Pavoni, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 450 = P. Pavoni.
pedunculariis, Miq. Syst. Pip. 531 = P. pedunculare.
persicariaefolia, Miq. l. c. 499 = P. persicariaefolium.
peruviana, Miq. l. c. 482 = P. pseudo-peruvianum.
phthiotrichia, Miq. l. c. 527 = P. phthiotrichon.
phytolaccaefolia, Miq. l. c. 534 = P. phytolaccaefolium.
pilosiuscula, Miq. l. c. 476 = P. pilosiusculum.
plantaginea, Miq. l. c. 389 = P. plantagineum.
Poeppigii, Miq. l. c. 487 = P. pseudo-mollicomum.
polyneura, Miq. l. c. 504 = P. indecorum.
populifolia, Miq. l. c. 418 = P. populifolium.
pothifolia, Miq. l. c. 399 = P. pothifolium.
pothomorpha, Miq. l. c. 403 = P. Pothomorphe.
pruniflolia, Miq. l. c. 438 = P. pruniflolium.
pseudo-adversuvyra, Miq. l. c. 407 = P. pseudo-chromasya.
psilostachya, Miq. l. c. 448 = P. psilostachyum.
punctata, Miq. l. c. 572 = P. punctatum.
pyrifolia, Miq. in Linnaea, xx. (1847) 159 = P. Masterdianum.
Radula, Miq. l. c. 426 = P. Bredemeyeri.
ramulosa, Miq. ex C. DC. in DC. Prod. xvi. 1. 388 = P. Hostmannianum.
Regnellii, Miq. in Linnaea, xxii. (1849) 574 = P. Regnellii.
renifolia, Miq. in Mart. Fl. Bras. iv. 1. 221.—Am. trop.
reticulosa, Miq. Syst. Pip. 554 = P. reticulosum.
rhododendrifolia, Miq. l. c. 506 = P. anonaefolium.
rhynchostachya, Miq. l. c. 539 = P. rhynchostachyum.
richardiaefolia, Miq. l. c. 395 = P. riclardiaefolium.
riparia, Miq. l. c. 409 = P. grandifolium.
rivinoides, Miq. l. c. 519 = P. rivinoides.

ARTANTHE :—
Rudgeana, Miq. in Linnaea, xxii. (1849) 77 = P. Rudgeanum.
rudis, Miq. Syst. Pip. 467 = P. rude.
rugosa, Miq. l. c. 468 = P. Rohrii.
Rutaiana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 444 = P. pangoense.
rumicifolia, Miq. in Linnaea, xx. (1847) 141 = P. rumicifolium.
salicariaefolia, Miq. Syst. Pip. 468 = P. salicariaefolium.
salicifolia, Miq. l. c. 523 = P. salicifolium.
saliunca, Miq. l. c. 528 = P. saliclnum.
salvaefolia, Miq. l. c. 430 = P. salviaefolium.
sancta, Miq. in Linnaea, xviii. (1844) 714 = P. sanctum.
saururoides, Miq. Syst. Pip. 414 = P. saururoides.
scabra, Miq. l. c. 447 = P. hirsutum.
scabrata, Presl, Epim. Bot. 226.—Mexico.
scabrifolia, Miq. Syst. Pip. 461 = P. scabrifolium.
scabriuscula, Miq. l. c. 447 = P. scabriusculum.
Schachii, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 448 = P. Schachii.
Schiedeanidula, Miq. Syst. Pip. 386 = P. reticulanum.
Schomburghii, Klotzsch, in Hook. Journ. Bot. iv. (1842) 322.—Guiana.
Schottii, Miq. in Linnaea, xx. (1847) 157 = P. Schottii.
scopulorum, Miq. Syst. Pip. 396 = P. scopulorum.
scutata, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 445 = P. subauriculum.
secunda, Miq. Syst. Pip. 516 = P. secundum.
Seemannsiana, Miq. in Scem. Bot. Voy. Herald, 199. t. 39 = P. auritum.
Sellowiana, Miq. Syst. Pip. 416 = P. Sellowianum.
septuplinervia, Miq. in Scem. Bot Voy. Herald, 199. t. 40 = P. septuplinervium.
serovia, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 443 = P. serovium.
spectabilis, Miq. in Linnaea, xx. (1847) 188 = P. cernuum.
staminea, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 461 = P. geniculatum.
stellispia, Miq. in Linnaea, xx. (1847) 163 = P. stellipilum.
stupposa, Miq. Syst. Pip. 480 = P. crocatum.
subaequalis, Presl, Epim. Bot. 226 = P. subaequale.
subauriculata, Miq. in Linnaea, xx. (1847) 160 = P. subauriculatum.
superba, Miq. l. c. 187 = P. obliquum.
Swartziana, Miq. Syst. Pip. 490 = P. Swartzianum.
tectoniaefolium, Miq. l. c. 396 = P. tectoniaefolium.
tejucensis, Miq. l. c. 496 = P. tejucense.
tenuicuspis, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 457 = P. tenuicuspe.
tiliaefolia, Miq. Syst. Pip. 388 = P. tiliaefolium.
trichonerva, Miq. l. c. 483.—Guiana.
trichostachya, Miq. l. c. 485 = P. citrifolium.
tricuspis, Miq. in Scem. Bot. Voy. Herald, 200. t. 42.—Panama.
trinervis, Miq. l. c. 200. t. 41 = P. trinerve.
tuberculata, Miq. Syst. Pip. 497 = P. tuberculatum.
tumida, Miq. l. c. 465 = P. tumidum.
ulmifolia, Miq. l. c. 472 = P. dilatatum.
umbrosa, Miq. l. c. 470 = P. amalago.
variegata, Miq. l. c. 485 = P. variegatum.
Vellosiana, Miq. l. c. 436 = P. aduncum.
velutina, Miq. l. c. 471 = P. velutinum.
verbaecifolia, Miq. l. c. 431 = P. verbascifolium.
verrucosa, Griseb. Fl. Brit. W. Ind. 171 = P. geniculatum.
Warszewiczcana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 469 = P. Warszewiczcana.
Wydleriana, Miq. Syst. Pip. 487 = P. cuspidatum.
xestophylla, Miq. l. c. 491 = P. geniculatum.
xylopioides, Miq. l. c. 492 = P. xylosteoides.
xylosteoides, Miq. l. c. 422 = P. xylosteoides.

ARTEDIA, Linn. Gen. ed. 1. 71 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 927.
muricata, Linn. Sp. Pl. 342 = Daucus muricatus.
squamata, *Linn. l. c.*—Orient.
squamata, Pall. in Nov. Act. Petrop. x. (1797) 308 = Seniecia crithmifolius.

ARTEMISIA, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 435.
ABROTANUM, [Tourn.] Neck. Elem. i. 98 (1790).
ABSINTHIUM, Tourn. ex Linn. Syst. ed. I (1735).

ARTEMISIA:—
OLIGOSPORUS, Cass. in Bull. Soc. Philom. (1817) 88.
PICROTHAMNUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 417.

abrotanifolium, Salisb. Prod. 191 = A. Abrotanum.
abrotanifolium, *Siev. in Pall. N. Nord. Beitr.* vii. 309. —Russia.
abrotanoides, Jacq. ex DC. Prod. vi. 106 = vestita.
abrotanoides, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 399 = californica.
Abrotanum, Habl. Taur. 128 = maritima.
Abrotanum, *Linn. Sp. Pl.* 845.—Europ. ; As. temp.
Abrotanum, Savi, Mat. Med. t. 88 = camphorata.
Abrotanum, S. G. Gmel. It. i. 21 = procera.
absinthia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Absinthium.
Absinthii, Hayne, ex Hook. f. Fl. Brit. Ind. iii 329 = pallens.
absinthioides, Poir. Encyc. Suppl. i. 461 = Siversiana.
Absinthium, *Linn. Sp. Pl.* 846.—Europ.
abyssinica, *Sch. Bip. in A. Rich. Tent. Fl. Abyss.* i. 420. —Afr. trop.
acetica, Jacquem. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 86 = maritima.
achilleæfolia, Tenore, Fl. Nap. 190 = camphorata.
achilleoides, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 195 = Tanacetum achilleoides.
Adamsii, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 27.—As. bor. or.
aerenula, Steph. ex Bess. l. c. 22 = rutæfolia.
æthiopica, Linn. Sp. Pl. 845.—A. Herba-alba.
affinis, Hausk. in Flora, xxv. (1842) 11. Beibl. 22 = vulgaris.
affinis, *Spreng. Nosn.* ed. II. i. 186.—Sibir.
afra, *Spreng. Hort. Schoensk.* iv. 84. t. 467.—Afr. austr.
alba, Habl. Taur. 144 = austriaca.
alba, Pall. Reise, i. 462 = maritima.
alba, Terra, Fl. Ital. Prod. 67 = camphorata.
albida, Ledeb. Fl. Alt. iv. 85 = maritima.
albida, Willd. ex Ledeb. Fl. Ross. ii. 566 = inodora.
Alliondi, Nym. Consp. 377 = nana.
alpina, Pall. ex Willd. *Sp. Pl.* iii. 1834 = lanata.
alpina, DC. ex Steud. Nom. ed. II. i. 186 = campestris.
alpina, Sch. Bip. ex Willk. in Flora, xxxiv. (1851) 746 = lanata.
altaica, Desf. Cat. Hort. Paris. (1829) 160 = pontica.
altissima, Kark. ex DC. Prod. vi. 108 = procera.
ambigua, Linn. Sp. Pl. ed. II. 1190 = Trichogyne laricifolia.
ambigua, Thunb. Prod. Pl. Cap. 147 = Seriphium vermiculatum.
Ammaniana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 80.—Sibir.
amorpha, *Raffn. Fl. Ludov.* 69.—Louisiana.
amygdalina, *Decne. in Jacquem. Voy. Bot.* 92. t. 100.—Reg. Himal.
andicola, Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 296 = mendocina.
andromeda, Senm. Bot. Voy. Herald, 34. t. 6 = senjavinensis.
anethifolia, Fisch. ex DC. Prod. vi. 108 = procera.
anethifolia, *Weber, ex Stechm. Artem.* 39.—Sibir.
angelica, Hort. ex DC. Prod. vi. 108 = maritima.
angustifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 450 = A. Abrotanum.
annua, *Linn. Sp. Pl.* 847.—Orient; As. bor.
annua, Pall. Reise, ii. 523 = sacrorum.
annualis, *S. Moore, in Journ. Bot.* xiii. (1875) 227.—China.
anthriscifolia, *Presl, ex Steud. Nom.* ed. II. i. 186.—Hab ?
apetala, Hort. Pest. ex Stend. l. c. = vulgaris.
apiacea, *Hance, in Walp. Ann.* ii. 695.—China.
apiea, Gebl. ex DC. Prod. vi. 125 = frigida.
aquatica, *Lour. Fl. Cochinch.* ii. 490.—China.
aragonensis, Lam. Encyc. ii. 269 = A. Herba-alba.
arborescens, *Linn. Sp. Pl.* ed. II. 1188.—Reg. Mediterr.
arbuscula, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 398.—Am. bor. occ.
arctica, Bess. ex Hook. Fl. Bor. Am. i. 322 = Richardsoniana.
arctica, Less. in Linnæa, vi. (1831) 218 = norvegica.
arenaria, *DC. Prod.* vi. 94.—Reg. Caucas.
argentea, Buch, Beschr. Canar. 148, 164 = canariensis.
argentea, *L'Hérit. Sert. Angl.* 23.—Sicil. ; Madera.
argentea, Sch. & Maur. ex Willk. & Lange, Prod. Fl. Hisp. ii. 68 = arborescens.
argyrophylla, *Ledeb. Fl. Alt*, iv. 106.—Sibir.

ARTEMISIA:—
armeniaca, *Lam. Encyc.* i. 268.—Armenia.
armeniaca, Willd. ex Ledeb. Fl. Ross. ii. 592 = Tournefortiana.
Assoana, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 69 = lanata.
atragonensis, Auct. vide *aragonensis*.
Athanasia, Bess. in Bull. Soc. Nat. Mosc. vii. (1834) 24 = Tanacetum fruticulosum.
atlantica, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ix. (1862) 398.—Algeria.
atrata, Lam. Encyc. i. 258 = spicata.
atropurpurea, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. (1867) i. 281 = arctica.
Aucheri, *Boiss. Fl. Orient.* iii. 367.—Persia.
australis, Ehrh. ex DC. Prod. vi. 120 = biennis.
australis, *Less. in Linnæa*, vi. (1831) 252.—Ins. Sandvic.
austriaca, *Jacq. Fl. Austr.* i. 61.—Europ. or. ; Orient ; As. bor.
baicalensis, Willd. ex DC. Prod. vi. 107 = sacrorum.
Balsamita, Willd. Enum. Hort. Berol. Suppl. 57 = pontica.
barguzinensis, Spreng. Syst. iii. 493 = borealis.
Barrelieri, *Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 87. —Hispan.
barrelieri, Webb, ex Willk. & Lange, Prod. Fl. Hisp. ii. 70 = granatensis.
Baumgarteni, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 78.—Europ. austr. or.
Besseriana, Ledeb. Fl. Ross. ii. 590 = lagocephala.
Biasolettiana, Visiani, Ind. Sem. Hort. Patav. (1856) = camphorata.
biennis, *Willd. Phyt.* 11. n. 39.—Am. bor. ; Sibir. ; Reg. Himal.
bifora, *Spreng. Nachr. i. Bot. Gart. Halle*, 12.—Hab ?
Bigelovii, *A. Gray, in Pacif. Rail. Rep.* iv. 110.—Am. bor.
Billardieriana, Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 81 = A. Herba-alba.
blepharolepis, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 240 (*Lédm. Reliq. Bot.* 154).—As. centr.
Bocconi, All. Fl. Pedem. i. 169 = nana, spicata.
Bolanderi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 48.—Calif.
borealis, *Pall. Reise*, iii. 755.—Am. bor. et arct. ; Sibir.
Boschniakiana, DC. Prod. vi. 104 = maritima.
brachyloba, *Franch. Pl. Davidian.* i. 171.—Mongol.
brevifolia, Wall. Cat. n. 5206 = maritima.
caerulescens, *Linn. Sp. Pl.* 848.—Reg. Mediterr.
campicola, Bess. in Hook. Fl. Bor. Am. i. 324 = Richardsoniana.
campitosa, *Ledeb. Fl. Alt.* iv. 80.—Sibir. alt.
californica, *Less. in Linnæa*, vi. (1831) 253.—Calif.
Campbellii, *Hook. f. & Thoms. ex Clarke, Comp. Ind.* 164.—Reg. Himal.
campestris, Auct. Hisp. ex Willk. & Lange, Prod. Fl. Hisp. ii. 72 = glutinosa.
campestris, Ledeb. Fl. Alt. iv. 86 = commutata.
campestris, *Linn. Sp. Pl.* 846.—Europ. ; As. bor. ; Orient ; Ind. or.
campestris, Pursh, Fl. Am. Sept. ii. 522 = canadensis ?
campestris, Scop. ex Steud. Nom. ed. II. i. 186 = camphorata.
campestris, Turcz. ex DC. Prod. vi. 96 = Ledebouriana.
campicola, *Phil. Prosp.* 31.—Europ. austr.
camschatica, Schlecht. ex Ledeb. Fl. Ross. ii. 568 = borealis.
cana, *Pursh, Fl. Am. Sept.* 521.—Am. bor.
canadensis, *Michx. Fl. Bor. Am.* ii. 129.—Am. bor. ; As. bor. or.
canariensis, *Less. in Linnæa*, vi. (1831) 217.—Ins. Canar.
canescens, Willd. Sp. Pl. iii. 1843 = armeniaca.
cannabina, Jacquem. ex Bess. in DC. Prod. vi. 97 = desertorum.
capillaris, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 176 = scoparia.
capillaris, *Thunb. Fl. Jap.* 309.—Japon.
capillifolia, Fisch. ex DC. Prod. vi. 123 = subvicinosa.
capillifolia, Lam. Encyc. i. 266 = Estafortiana foeniculaceum.
carrifolia, Buch.-Ham. ex Roxb. Fl. Ind. iii. 428 = roxburghiana.
carvifolia, *Roxb. Hort. Beng.* 61.—Ind. or. ; China.
carvifolia, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 25 = parrhelia.
caucasica, *Willd. Sp. Pl.* iii. 1838.—Reg. Caucas.
caudata, *Michx. Fl. Bor. Am.* ii. 130.—Am. bor.

ARTEMISIA:—
centiflora, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 498.—Mongolia.
cephalantha, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xl. (1867) i. 227 = glomerata.
cernua, Dufour, ex Willk. & Lange, Prod. Fl. Hisp. ii. 71 = hispanica.
cernua, Nutt. Gen. Am. ii. 148 = dracunculoides.
cernuifolora, Dufour, ex Willk. & Lange, Prod. Fl. Hisp. ii. 71 = hispanica.
chamaemelifolia, *Vill. Prosp.* 32.—Gall. ; Reg. Caucas. ; Persia.
Chamissoniana, Bess. ex Hook. Fl. Bor. Am. i. 324 = norvegica.
chamomillifolia, Parrot, in Engelh. & Parrot, Reise Krym. Kauk. ii. 131 = chamaemelifolia ?
Chiajcna, *Kunze, ex Rich. Bot. Méd.* ii. 612 ; *ex Steud. Nom.* ed. II. i. 187.—Orient ?
chinensis, Bess. in Linnæa, xv. (1841) 92 = lagocephala.
chinensis, Burm. ex DC. Prod. vi. 125 = mollis.
chinensis, *Linn. Sp. Pl.* 849.—China ; Sibir.
chinensis, Pursh, Fl. Am. Sept. ii. 521 = Stelleriana.
chrysantha, Fisch. ex DC. Prod. vi. 125 = frigida.
chrysantha, Steph. ex Ledeb. Fl. Ross. ii. 597 = rupestris.
Chrysanthemum, Russ. ex DC. Prod. vi. 140 = Sphaeromorphaea Ruselliana.
Cina, *Berg. Dactyll.* iv. t. 29*.—Orient.
cinerea, Hort. Par. ex DC. Prod. vi. 123 = camphorata.
Clauscns, *Pomel, Nouv. Mat. Fl. Alt.* 55.—Algeria.
coarctata, J. H. Fors. in Linn. Inst. Skrift. (reimpr. 12) = vulgaris.
coarctata, Turcz. ex Ledeb. Fl. Ross. ii. 567 = campestris.
coarctescens, Fisch. ex Bess. = sericea.
coarctescens, Presc. ex Ledeb. Fl. Ross. ii. 564 = Ammaniana.
collina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 39 = campestris.
columbiensis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 398 = nana.
Columnae, Tenore, Ind. Sem. Hort. Neap. (1839) 14 = camphorata.
commutata, Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 70 = campestris.
compacta, Fisch. ex DC. Prod. vi. 102 = maritima.
Contra, Linn. Mant. 282 = Sieberi.
Contra, Willd. ex Spreng. Syst. iii. 494 = maritima.
controversa, Jan, ex DC. Prod. vi. 126 = Siversiana.
Copa, *Phil. Fl. Atac.* 33.—Chili.
corymbosa, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 74 = glomerata.
corymbosa, Lam. Encyc. i. 266 = camphorata.
crithmifolia, *Linn. Sp. Pl.* 846.—Lusit. ; Gall.
crithmifolia, Sievers, in Pall. N. Nord. Beitr. vii. 338 = glabella.
cuneifolia, DC. Prod. vi. 126 = parviflora.
cuneifolia, Scheele, in Linnæa, xxii. (1849) 162 = mexicana ?
curilensis, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 67 = Meyeriana.
curilensis, Spreng. Syst. iii. 489 = leontopodioides.
Cyna, Pall. Reise, iii. 690 = maritima.
Czekanowskiana, *Trautv. in Act. Hort. Petrop.* v. (1877) 78.—Sibir.
Darwini, *Vilm. Summa*, v. 19.—Dahuria.
Decaisnei, *F. W. Klatt, in Abh. Naturf. Ges. Halle*, xv. (1883) 239.—Ind. or.
Delilliana, Bess. Enum. Pl. Volh. Suppl. 39 = monosperma.
delphinensis, Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 19 = Mutellina.
densiflora, Viv. App. Alt. Corn. fl. 4. t. 3 = gallica.
denticta, Willd. Sp. Pl. iii. 1836 = rupestris.
depressa, Turcz. ex DC. Prod. vi. 107 = Adamsii.
desertorum, Bess. ex Hook. Fl. Bor. Am. i. 325 = canadensis.
desertorum, *Spreng. Syst.* iii. 490.—As. centr. ; Sibir. ; Reg. Himal.
dieuleri, *Dougl. ex DC. Prod.* vi. 108.—Am. bor. occ.
discolor, *Turcz. & Gray, in Pacif. Rail. Rep.* ii. 126 = tanacetoides.
Douglasiana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 66 = tridentata.
dracunculoides, *Pursh, Fl. Am. Sept.* 742.—Am. bor.
Dracunculus, *Linn. Sp. Pl.* 848.—Europ. or. ; Orient ; Reg. Himal.

ARTEMISIA:—
Dracunculus, Pursh, Fl. Am. Sept. ii. 521 = dracunculoides.
dubia, Wall. Cat. n. 3307 = vulgaris.
Echegaryi, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 42.—Reg. Argent.
elegans, Fisch. ex Ledeb. Fl. Ross. ii. 579 = procera.
elegans, Jan, ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 104 = glacialis.
elegans, Roxb. Hort. Beng. 61; Fl. Ind. iii. 421 = scoparia.
elegans, Salisb. Prod. 191 = arborescens.
eranthema, *Bunge*, *in Mém. Sav. Etr. Pétersb.* vii. (1851) 639 (*Lehm. Reliq. Bot.* 168).—Turkest.
eriantha, Tenore, Ind. Sem. Hort. Neap. (1830) 14 = Baumgarteni.
eriocarpa, *Bunge*, *in Mém. Sav. Etr. Pétersb.* vii. (1851) 636 (*Lehm. Reliq. Bot.* 160).— Persia; Turkestan.
eriophora, Schlecht. ex Ledeb. Fl. Ross. ii. 586 = vulgaris.
eriopoda, *Bunge*, *Enum. Chin. Bor.* 87.—Chin. bor.
Eschscholtziana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 24.—Ins. Sandvic.
exilis, Fisch. ex DC. Prod. vi. 119 = annua.
Falconeri, *C. B Clarke*, *in Hook. f. Fl. Brit. Ind.* iii. 328.—Tibet. occ.
fasciculata, *Bieb. Fl. Taur. Cauc.* ii. 293.—Reg. Caucas.; Persia.
fasciculata, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iii. 318 = Tanacetum artemisioides.
filagrandea, Steehm. Artem. 29 = vallesiaca.
filifolia, *Torr. in Ann. Lyc. N. York*, ii. (1828) 211.—Am. bor.
Fischeriana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 21 = californica.
forniculacea, Stev. ex DC. Prod. vi. 108 = procera.
fortida, Jacquem. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 96 = desertorum.
foliosa, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 397 = californica.
Fontanesiana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 52.—Hab.?
fragrans, Eichw. Casp. Cauc. 6 = Sieberi.
fragrans, Willd. Sp. Pl. iii. 1835 = maritima.
fraxinoides, *Greene*, *in Bull. Torrey Club*, x. (1883) 42.—Am. bor. occ.
frigida, Eichw. Casp. Cauc. 31 = austriaca.
frigida, *Willd. Sp. Pl.* iii. 1835.—Sibir.; Am. arct.
frutescens, *Pall. Reise Staath. Russ. Reich.* i. 94, 177. —Rossia.
fruticosa, Comm. ex DC. Prod. vi. 101 = Stoebe paniculata.
fruticosa, Willd. ex DC. l. c. 94 = arenaria.
furcata, Bieb. Fl. Taur. Cauc. iii. 567 = trifurcata?
gallica, Schur, Enum. Pl. Transs. 224 = maritima.
gallica, *Willd. Sp. Pl.* iii. 1824.—Gall.; Hisp.
gargunica, Tenore, ex DC. Prod. vi. 122 = camphorata.
Gayana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 35. —Hispan.
gelida, Ledeb. in Mém. Acad. Pétersb. v. (1805) 565 = borealis.
Gmelii, Steehm. Artem. 17 = spicata.
gilvescens, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 175.— Japon.
glabella, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 441.—Soongaria.
glabra, Jacquem. ex Bess. in DC. Prod. vi. 97 = desertorum.
glabrata, DC. in Wight, Contrib. 20 = parviflora.
glabrescens, Stokes, Bot. Mat. Med. iv. 188 = campestris.
glacialis, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 70 = granatensis.
glacialis, Vitm. Sagg. 44 = lanata.
glacialis, Herb. Mosc. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 67 = Tilesiana.
glacialis, *Linn. Sp. Pl.* ed. II. 1167.—Europ. austr. med.
glacialis, Wall. in Jacq. Coll. ii. 118 = Mutellina.
glacialis, Schang. in Pall. N. Nord. Beitr. vi. 93 = commutata.
glauca, Bieb. ex Ledeb. Fl. Ross. ii. 564 = Ammassiana.
glauca, *Pall. ex Willd. Sp. Pl.* iii. 1831.—Reg. Himal.; Sibir.
glutinosa, Chem. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 94 = norvegica.
ARTEMISIA:—
globularia, Ledeb. ex Maxim. in Bull. Acad. Pétersb. xvii. (1872) 485 = glomerata.
glomerata, Hook & Arn. Bot. Beech. Voy. 125 = senjavincana.
glomerata, *Ledeb. in Mém. Acad. Pétersb.* v. (1805) 564.—Am. occ. et As. or. arct.
glomerata, Sieber, ex Spreng. Syst. iii. 489 = Sieberi.
glomerulata, *Sieber*, *ex Spreng. Syst.* iii. 489 = Sieberi.
glutinosa, Boiss. ex Walp. Rep. vi. 214 = variabilis.
glutinosa, *J. Gay*, *in DC. Prod.* vi 95.—Gallia; Ital.
Gmelini, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc ii. (1834) 28 = sacrorum
Gmelini, *Steehm. Artem* 30.—Sibir
Gmelini, Willd. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 67 = Lagopus.
gnaphaloides, Bess. in DC. Prod. vi. 105 = mendozana.
gnaphaloides, Nutt. Gen. Am. ii. 143 = ludoviciana.
gongonum, *Webb*, *in Hook. Niger Fl.* 142.—Ins. Cap. Verd.
gracilis, L'Hérit. ex DC. Prod. vi. 100 = scoparia.
granatensis, *Boiss. ex DC. l. c.* 298.—Hispan.
grandiflora, *DC. l. c.* 122.—Sibir.
grandiflora, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xl. (1867) 1. 217 = pontica.
grandiflora, Hoffm Deutschl. Fl. ed. II. ii. 136 = tupestris.
grata, Heyne, ex Steud. Nom. ed. II. i. 137 = Roxburghiana?
grata, Wall. Cat. n. 3294; Comp. n. 404 = Roxburghiana.
Griffithiana, Boiss Diagn Ser. II. iii. 31 = macrocephala.
groenlandica, Wormsk. in Fl Dan. t 1585 = borealis.
Halimodendron, Ledeb in Hook. f. Fl. Brit. Ind. iii. 321 = salsoloides.
Haloderon, *Turcz. in DC. Prod.* vi. 94.—Dahur.; Mongol.
Haussknechtii, *Boiss. Fl Orient.* iii. 374.—Persia.
helenioides, Schleich. Cat. Pl. Helv. ed. III. 8 = nana.
hemisphaerica, Roxb. Hort. Beng. 61; Fl. Ind. iii. 422 = Cotula hemisphaerica.
Herba-alba, *Asso*, *Syn. Arag.* 117.—Hispan.; Marocc.; Orient.
herbacea, Clarke, DC. Prod. vi. 108 = procera.
herbacea, *Ehrh. in Willd. Sp. Pl.* iii. 1828.—Sibir.
herbacea, Hort. Wratisl. ex Bess. in Nouv. Mém Soc. Nat. Mosc. iii. (1834) 80 = procera.
hermaphrodita, Ledeb. ex Ledeb. Fl. Ross. ii. 586 = campestris.
heterophylla, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 74 = Richardsoniana?
heterophylla, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 400 = vulgaris.
Heyneana, Wall. Cat. n. 3295 = vulgaris.
hirsuta, Rottl. in Spreng. Syst. iii 490 = Cyathocline lyrata.
hispanica, Jacq. Coll. i. 98 = biennis.
hispanica, *Lam. Encyc.* i. 363.—Hispan.
hispanica, Steehm. ex Bess. in Bull. Nat. Soc. Mosc. ix. (1836) 114 = vulgaris.
holotricha, *Bieb. ex Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 46.—Ukraina.
holosericea, Boiss. & Hausskn. in Boiss. Fl. Orient. iii. 374 = Haussknechtii.
holosericea, Ledeb. Fl. Alt. iv. 65 = sericea.
Hookeriana, Bess. in Nouv. Mém. Nat. Soc. Mosc. iii. (1834) 89 = indoviciana.
humilis, Bieb. Casp. App. 38 = maritima.
humilis, Will. Gard. Dict. ed. VIII. n. 7 = A. Abrotanum.
humilis, Wulf. in Jacq. Coll. iv. 295 = camphorata.
hypoleuca, Edgew. in Trans. Linn. Soc. xx. (1846) 71 = Roxburghiana.
hyrcana, Spreng. Syst. iii. 484 = annua.
ibetica, *Boiss. ex Boiss. & Buhse*, *in Nouv. Mém. Soc. Nat. Mosc.* viii. (1860) 130.—Reg. Cauc.
ignatzia, Maxim. Prim. Fl. Amur. 161 = vulgaris.
incanescens, Jord. in Ann. Soc. Agric. Lyon; ex Gren. & Godr. Fl. Fr. ii. 197 = camphorata.
incompta, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 400 = discolor.
scoupicina, Spreng. Nachr. i. Bot. Gart. Halle, 12 = biennis.
inculta, Delile, Fl. Egypte, 264 = A. Herba-alba.
ARTEMISIA:—
inculta, Sieber, ex DC. Prod. vi. 95 = monosperma.
indica, Willd Sp. Pl. iii. 1846 = vulgaris.
inferna, S. G. Gmel. ex Ledeb. Fl. Ross. ii. 566 = campestris.
inodora, Bieb. Fl. Taur. Cauc. ii. 295 = campestris.
inodora, Hook. & Arn. Bot. Beech. Voy. 150 = dracunculoides.
inodora, Mill. Gard. Dict. ed. VIII. n. 16 = A. Absinthium.
inodora, Willd. Enum. Hort. Berol. 864 = A. Dracunculus
insipida, *Schur*, *Enum. Pl. Transs.* 222.—Transylv.
insipida, *Vill. Prosp.* 32.—Gallia.
integrifolia, *Falk, Beitr.* ii. 229.—Reg. Casp.
integrifolia, Linn. Sp. Pl. 848 = vulgaris.
integrifolia, Pursh, Fl. Am Sept. 520 = ludoviciana
integrifolia, Richards. App. Frankl. Journ. ed. II. 30 = Purshiana.
intermedia, Host, Fl. Austr. ii. 160 = camphorata
intricata, *Franch. Pl. Davidian.* i. 170.—Mongol.
involucrata, Turcz ex Ledeb. Fl. Ross. ii. 598 = frigida.
italica, Bieb. ex Ledeb. l. c. 566 = campestris.
italica, *Steehm. Artem.* 26.—Italia.
Jacquemontiana, Bess. in Bull. Soc. Nat. Mosc. (1835) 59 = desertorum.
Jacquini, Rasuach. Nom. ed. III. 387 = biennis
japonica, *Thunb. Fl. Jap.* 310.—Japon.
japonica, F. Schmidt, Fl. Sachal. 149 = Keiskeana.
jenisea, Willd. ex Ledeb. Fl. Ross. ii. 598 = seq.
jeniscensis, Willd. ex Spreng. Syst. iii. 486 = frigida.
judaica, *Linn. Fl. Palaest. in Amoen. Acad.* iv. 463; *et Mant.* 281.—Aegypt.; Arab.
judaica, Lour. Fl. Cochinch. 489 = Lourelri.
juncea, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 383.— Soongaria.
Jussiaea, *J. Gay*, *ex Bess. l. c.* viii. (1835) 28.— Hispan.
kamtschatica, Willd. ex Bess. l. c. 67 = leontopodioides.
Keiskeana, *Miq. in Ann. Mus Bot. Lugd. Bat.* ii. 176.—Japon.
Koelreuteriana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 16.—Hab.?
kotschyana, *Klatt*, *in Sitzb. Akad. Muench.* (1878) 91.— Ind. or.
Kotschuensis, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 80 = Chrysanthemum lupinatum.
Krascheninnikoviana, Bess. l. c. 41 = macrobotrys, latifolia.
Krukitzana, Bess. l. c. 22 = lagocephala.
kulbadica, *Boiss. & Buhse*, *in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 120.—Reg. Casp.
kurilensis, Walp. Rep. vi. 217 = A. Lagopus.
lacinifata, *Willd. Sp. Pl.* iii. 1848.—Reg. Himal.; Sibir.
lactiflora, Wall. ex DC. Prod. vi. 115 = vulgaris.
lagocephala, *Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 23.—Sibir.
lagopina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = seq.
Lagopus, *Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 69.—Sibir.
lanata, Koch, Syn. Fl. Germ. ed. II. 402 = nitida.
lanata, Lam. Encyc. i. 367 = variabilis.
lanata, *Willd. Sp. Pl.* iii. 1828.—Hisp.; Ital.; Taur. *lanuginosa*, Poir. Encyc. Suppl. i. 402 = lanata.
larvifolia, S. G. Gmel. ii. ii. 8 = maritima.
lasiocline, Boiss. Diagn. Ser. I. vi. 93 = splendens.
latifolia, Ledeb. in Mém. Acad. Pétersb. v. (1805) 569 = laciniata.
lavandulaefolia, DC. Prod. vi. 110 = vulgaris.
lavandulifolia, Salisb. Prod. 191 = caerulescens.
Ledebouriana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 37.—Sibir.
ledoicrnai, Rochel, ex Spreng. Fugill. ii. 77 = campestris.
Lehmaniana, *Bunge*, *in Mém. Sav. Etr. Pétersb.* vii. (1851) 240 (*Lehm. Reliq. Bot.* 164).—Afghan.; Turkest.
leontopodioides, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 68 = glomerata.
leptophylla, *D. Don*, *Prod. Fl. Nep.* 183 (sp. dubia).— Nepal.
leptostachya, DC. Prod. vi. 113 = vulgaris.
Lercheana, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 441 = maritima.

ARTEMISIA :—

Lercheana, Presc. ex Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 38 = arenaria.

Lerchiana, Weber. ex Stechm. Artem. 24 = fragrans.

Lerchii, Beeber, ex Georgi, Beschr. Russ. Reich. iii. IV. 1235 = maritima.

Lessingiana, *Bess. in Linnaea*, XV. (1841) 90.— Sibir.

leucanthemifolia, Desf. ex DC. Prod. vi. 98 = desertorum.

lencolea, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 106.—Soongor.

leucophylla, Turcz. ex C. B. Clarke, Comp Ind. 102 = vulgaris.

Leuinii, Torr. & Gray, Fl. N. Am. ii. 417 (nom. delend.).

Libanotis, Bieb. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 49 = macrobotrys.

Lindheimeriana, Scheele, in Linnaea, xxii. (1849) 168 = mexicana.

Lindleyana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 35.—Am. bor. occ.

linearis, Gueldenst. ex Ledeb. Fl. Ross. ii. 566 = salsoloides.

Lippii, Jan, ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 79 = A. Herba-alba.

lithophila, Turcz. ex DC. Prod. vi. 122 = lagocephala.

littoralis, Rctz. Obs. v. 28 = Epaltes australis.

Lobelii, All. Fl. Pedem. i. 166 = camphorata.

lobulifolia, *Boiss. Fl. Orient.* iii. 364.—Afghan.

longepedunculata, Rudolphi, ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 77 = norvegica.

longifolia, *Nutt. Gen. Am.* ii. 142.—Am. bor. occ.

Lourciri, *Kostel. Allg. Med. Pharm. Fl.* ii. 669.— China.

ludoviciana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 38 = discolor.

ludoviciana, *Nutt. Gen. Am.* ii. 143.—Am. bor. occ.

mâora, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 364. —Catsonia.

macrantha, Ledeb. in Mém. Acad. Pétersb. v. (1805) 578 = sacrorum.

macrobotrys, Ledeb. Fl Alt. iv. 78 = laciniata.

macrocephala, *Jacquem. ex Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 28.—Afghan.; Tibet.; As. centr.

macrophylla, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 41 = laciniata.

macrorhiza, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 73.—Sibir.

maderaspatana, Linn. Sp. Pl. 849 = Grangea maderaspatana.

magellanica, *Sch. Bip. in Flora*, xxxvii. (1855) 116.— Reg. Magell.

maracandica, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 846 (Lehm. Rel. Bot. 169) = fragrans.

maritima, *Linn. Sp. Pl.* 846.—Europ.; Reg. Caucas.; Sibir.

maritima, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 74 = gallica.

marocana, *Cosr. in Bull. Soc. Bot. Fr.* xxii. (1875) 61.—Marocco.

Marschalliana, Spreng. Syst. iii. 496 = campestris.

matricarioides, Less. in Linnaea, vi. (1831) 210 = Matricaria discoides.

megacephala, *Rupr. Sert. Tianach.* 52.—As. centr.

melanolepis, *Boiss. Diagn.* Ser. 1. xi. 97.—Persia.

mendozana, *DC. Prod.* vi. 105.—Am. austr.

Mertensiana, Wallr. Sched. Crit. 466. t. 6 = laciniata.

Messerschmidtiana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 37 = sacrorum.

Messerschmidtii, Weber, ex Stechm. Artem. 19 = maritima.

mexicana, *Willd. ex Spreng. Syst.* iii. 496.—Mexico.

Meyeriana, Bess. in Linnaea, xv. (1841) 96 = A. Lagopus.

Michauxiana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 71 = discolor.

Michauxii, Bess. ex Hook. Fl. Bor. Am. i. 324 = pracc.

millefoliata, Bieb. ex DC. Prod. vi. 107 = sacrorum.

minima, Linn. Sp. Pl. 849 = Centipeda orbicularis.

minor, *Jacquem. ex Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 22.—Tibet.

mollis, *Y. Gay, in DC. Prod.* vi. 125.—Hab. ?

mollissima, *D. Don, Prod. Fl. Nep.* 182 (sp. ignot.).— Nepal.

mollucana, *Roxb. Hort. Beng.* [101]; *Fl. Ind.* iii. 417.—Ins. Moluce.

ARTEMISIA :—

mongolica, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 53 = vulgaris.

monogyna, Waldst. & Kit. Fl. Rar. i. 177. t. 75 = fragrans.

monosperma, *Delile, Fl. Egypte*, 268.—Syria; Aegypt.; Arab.

monostachya, *Bunge, ex Maxim. Fl. Amur.* 482.— Mongolia.

monspeliensis, Nym. Consp. 377 = glutinosa.

montana, Schlecht. ex Ledeb. Fl. Ross. ii. 591 = Richardsoniana.

montevidensis, *Spreng. Syst.* iii. 487.—Reg. Argent.

monticola, C. Koch, in Linnaea, xxiv. (1851) 847 = cancasica.

Moorcroftiana, *Wall. Cat. Comp.* n. 4006.—Himal. bor. occ.

Moxa, *DC. Prod.* vi. 121.—China.

multicaulis, Lefeb. Fl. Alt. iv. 60 = anethifolia.

mutabilis, Salisb. Prod. 191 = A. Santonicum.

Mutellina, S. G. Gmel. It. i. 159 = splendens.

Mutellina, *Vill. Prosp.* 31.—Europ. med.

myriantha, Wall. Cat. n. 8297 = vulgaris.

nana, *Gaud. Fl. Helv.* v. 331.—Helvet.

narvaniana, Visiani, ex DC. Prod. vi. 108 = procera.

neapolitana, Tenore, Prod. Fl. Nap. App. v. 28 = variabilis.

neglecta, Spreng. in Willd. Enum. Hort. Berol. ii. 880 = desertorum.

nepalensis, *Nees, in Flora*, xiv. (1831) 290.—Ind. or.

nilotica, Linn. Cent. i. 27; Sp. Pl. ed. II. 1188 = Cotula scariosides.

nitens, Stev. ex Ledeb. Fl. Ross. ii. 595 = sericea.

nitida, *Bertol. Mant.* 58.—Italia.

nitrosa, *Weber, ex Stechm. Artem.* 24.—Sibir.

nivea, Redowsk. ex Willd. Enum. Hort. Berol. 868 = austriaca.

norica, Leyb. in Flora, xxxvii. (1854) 569 = nana.

norvegica, *Fries, Novit. Suec.* ed. I. (1817) 56.— Norveg.

nubigena, Wall. Cat. n. 8283 = Tanacetum nubigenum.

nutans, Fras. ex Pursh, Fl. Am. Sept. 742 = dracunculoides.

nutans, Willd. Sp. Pl. iii. 1831 = maritima.

Nuttalliana, Bess. ex Hook. Fl. Bor. Am. i. 326 = dracunculoides.

obtusiloba, *Ledeb. Fl. Alt.* iv. 68.—Sibir. alt.

occidentalis, Salzm. ex DC. Prod. vi. 96 = campestris.

odorstissima, *Desf. Fl. Atlant.* ii. 268.—Algeria.

officinalis, *Gueldenst. ex Ledeb. Fl. Ross.* ii. 600.— Rossia.

ohotensis, Steph. ex Ledeb. l. c. 597 = rupestris.

oliganthes, Méigev. in Bull. Soc. Bot. Fr. xviii. (1871) 86.—Mutellina.

Olivieriana, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 849 (Rel. Lehm. 150) = fragrans.

Olivieriana, J. Gay, ex Bess. in DC. Prod. vi. 101 = A. Herba-alba.

Ontina, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 236 = A. Herba-alba.

orientalis, Herb. Madr. ex DC. Prod. vi. 120 = pallens.

orientalis, Willd. Sp. Pl. iii. 1836 = austriaca.

pachystachya, DC. Prod. vi. 114 = pycnocephala.

pacifica, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 401 = canadensis.

Palassiana, *Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 61.—Rossia.

Pallasii, Spreng. Syst. iii. 494 = maritima.

Pallasii, Willd. ex Spreng. l. c. 493 = borealis.

pallens, *Willd. ex DC. Prod.* vi. 120.—Ind. or.

palitda, Salisb. Prod. 191 = pontica.

palmata, Lam. Encyc. i. 268 = caerulescens.

palmata, Lap. Hist. Abr. Pl. Pyr. 564 = gallica.

palmata, Steph. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 52 = ruteefolia.

Palmeri, *A. Gray, in Proc. Am. Acad.* xi. (1876) 79.— Calif.

palustris, *Linn. Sp. Pl.* 846.—Dahur.; Mongol.

paniculaeformis, DC. Prod. vi. 107 = vulgaris.

paniculata, *Lam. Encyc.* i. 265.—Ital.; Gall.

paniculata, Rottl. ex Wight, Contrib. 19 = Cystheolite lyrata.

paniculata, Roxb. Hort. Beng. 61; Fl. Ind. iii. 418 = vulgaris.

paniculata, Sch. Bip. ex Willk. in Flora, xxxiv. (1851) 747 = variabilis.

Parishii, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 230.—Calif.

Parryi, *A. Gray, l. c.* vii. (1868) 361.—N. Mexico.

parviflora, *Buch.-Ham. ex Roxb. Hort. Beng.* 61; *Fl. Ind.* iii. 450.—Ind. or.; Burma.

parviflora, Wight, Cat. n. 1460 = vulgaris.

pauciflora, Bieb. Fl. Taur. Cauc. ii. 290 = arenaria.

pauciflora, Georgi, Beschr. Russ. Reich. Nachtr. 306 = maritima.

pauciflora, Sprung. Syst. iii. 491 = desertorum.

pectinata, *Pall. Reise*, iii. 255.—Dahur.; Mongol.

pedatifida, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 399.—Am. bor.

pedemontana, Balb. ex Lotsel. Not. 126 = lanata.

pedunculariis, Bieb. Fl. Taur. Cauc. iii. 566 = splendens.

pedunculata, Stend. Nom. ed. II 72 = splendens.

pedunculosa, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 176.—Japon.

pendula, Salisb. Prod. 191 = A. Absinthium.

pendula, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 89 = fragrans.

persica, *Boiss. Diagn.* Ser. I. vi. 91.—Persia; Afghan.; As. centr.

petraea, Turcz. ex Ledeb. Fl. Ross. ii. 596 = frigida.

petraea, Jan, ex DC. Prod. vi. 118 = Baumgartenii.

peucedanifolia, Jusa. ex DC. Prod. vi. 99 = canadensis.

pinnatifida, Jacquem. ex DC. Prod. vi. 120 = Tournefortiana.

piperita, Pall. ex Ledeb. Fl. Alt. iv. 68 = scoparia.

plattensis, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 397 = filifolia.

plumosa, Fisch. ex Bess. in DC. Prod. vi. 119 = annua.

pontica, Berm. f. Prod. 26 = alba.

pontica, *Linn. Sp. Pl.* 847.—Europ.; Reg. Caucas.; Soongaria.

potentillaefolia, Fisch. ex Spreng. Syst. iii. 494 = armeniaca.

potentilloides, A. Gray, in Proc. Am. Acad. vi. (1864) 551 = Tanacetum potentilloides.

Prescottiana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 72.—Am. bor. occ.

procera, Lap. Hist. Abr. Pl. Pyr. 508 = variabilis.

procera, *Willd. Sp. Pl.* iii. 1818.—Europ. austr. or.; As. Min.; Sibir.

procumbens, Schrad. ex DC. Cat. Hort. Monsp. 80 = frigida.

pseudo-pontica, *Schur, Enum. Pl. Transs.* 323.— Europ.

pubescens, *Ledeb. in Mém. Acad. Pétersb.* v. (1805) 568.—Sibir. or.

pulchella, Gmel. ex DC. Prod. vi. 109 = maritima.

pumila, Hort. ex Link, Enum. Hort. Berol. ii. 316 = frigida.

pumila, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 640 = Lindleyana.

punctata, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 43 = latifolia, macrobotrys.

purpurascens, Jacquem. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 80 = Roxburghiana.

Purshiana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 59.—Am. bor.

pusilla, *Lucc, Topog. Nachr. Oesel*, 273.—Europ. bor.

pycnocephala, *DC. Prod.* vi. 99.—Sibir.

pycnorrhiza, *Ledeb. Fl. Alt.* iv. 70.—Sibir. alt.

pycnostachya, Torr. & Gray, Fl. N. Am. 423 = pycnocephala.

pyrrocacha, Viviani, Fl. Lib. Spec. 54. t. 18. f. 5 = hispanica.

quadripedalis, Gillib. Fl. Lituan. i. 174 = vulgaris.

quinquelobata, *Transit. in Bull. Soc. Nat. Mosc.* xxxix. (1866) i. 346.—As. temp.

racemifera, Steph. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 38 = sacrorum.

racemosa, *Ait. in Linnaea*, xxxii. (1863) 886.— Europ.

racemosa, Méigev. in Bull. Soc. Bot. Fr. xii. (1865) 361; xviii. (1871) 367 = spicata.

radicans, C. Sm. ex DC. Prod. vi. 103 = nutans.

ramosa, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 168.—Ins. Canar.

ramosa, Lag. ex Willk. & Lange, Prod. Fl. Hisp. ii. 71 = hispanica.

ARTEMISIA :—
Redowskyi, Ledeb. in Mém. Acad. Pétersb. v. (1806) 574 = A. Dracunculus.
repens, Pall. ex Willd. Sp. Pl. iii. 1840 = austriaca.
reptans, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 148 —Ins. Canar.
revoluta, Edgew. in Trans. Linn. Soc. xx. (1846) 72 = Roxburghiana.
rhodantha, *Rupr. Sert. Tianschi.* 52.—As. centr.
Richardsoniana, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 289 = norvegica.
Richardsoniana, *Hook. Fl. Bor. Am.* 323.—Am. arct.
rigida, *A. Gray, in Proc. Am. Acad.* xix. (1883) 49.—Am. bor. occ.
riparia, Buch.-Ham. ex DC. Prod. vi. 119 = carvifolia.
Rothrockii, *A. Gray, in Bot. Calif.* i. 618.—Calif.
Roxburghiana, *Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 87.—Reg. Himal.
Royleana, *DC. Prod.* vi. 115 (sp. dubia).—Ind. bor. occ.
rubella, Moench, Meth. 579 = caerulescens.
rubriflora, Turcz. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1836) 54 = vulgaris.
rudevalis, Salisb. Prod. 191 = vulgaris.
rupestris, All. Fl. Pedem. i. 169 = Mutellina.
rupestris, Aasø, Syn. Arag. 117 = lanata.
rupestris, Leg. & Kodr. in Anal. Cienc. v. (1802) 284 = granatensis.
rupestris, Lam. Encyc. i. 262 = splendens.
rupestris, *Linn. Sp. Pl.* 847 ; *It. Goth.* 265.—Oeland. ; Saxon. ; Sibir.
rupestris, O. F. Muell. in Fl. Dan. t. 801 = norvegica.
rupestris, Scop. Fl. Carn. ed. II. ii. 146 = camphorata.
rupestris, Vill. Hist. Pl. Dauph. ii. 246 = spicata.
ruscuca, Siev. ex DC. Prod. vi. 108 = rutaefolia.
rutaefolia, *Steph. ex Spreng. Syst.* iii. 488.—Sibir. or.
sabulosa, Siev. ex DC. Prod. vi. 108 = procera.
sabulosa, Willd. ex Ledeb. Fl. Ross. ii. 561 = arenaria.
sachalensis, Tilas. ex Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 48 = capillaris.
sacrorum, *Ledeb. in Mém. Acad. Pétersb.* v. (1805) 571.—Rossia ; Tibet. ; Sibir.
Saharæ, *Pomel, Nouv. Mat. Fl. Atl.* 56.—Algeria.
saligna, Tenore, Prod. Fl. Nap. Suppl. i. 63 = variabilis
salina, Bess. Enum. Pl. Volh. 76 = maritima.
salina, *Willd. Sp. Pl.* ii. 1834.—Europ.
salsa, F. Martini, Gesch. Nat. vii. 288 = nitrosa.
salsoloides, *Willd. Sp. Pl.* iii. 1833.—Reg. Caucas. ; Tibet. ; Sibir.
samamisica, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 50 = vulgaris.
santalinæfolia, Turcz. ex Bess. l. c. 87 = sacrorum.
Santonica, Lam. Encyc. i. 268 = caerulescens.
Santonica, Pursh, Fl. Am. Sept. 521 (sp. obscura).—Am. bor.
Santonica, Woodw. Med. Bot. t 128 = suaveolens.
Santonicum, *Linn. Sp. Pl.* 845.—As. bor.
Santonicum, Sievers, ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 113 = campestris.
saxatilis, Reichb. Fl. Germ. Excurs. 302 = incanescens.
saxatilis, Waldst. & Kit. in Willd. Sp. Pl. iii. 1830 = camphorata.
saxorum, F. Schmidt, Reisen Amurl. 50, *err. typ.* = sacrorum.
scariosa, Rich. ex Bess. in DC. Prod. vi. 126 = Sieversiana.
Schimperi, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 142, nomen.—Abyssin. trop.
Schlagintweitiana, *Klatt, in Sitzb. Acad. Muench.* (1878) 90.—Turkestan.
Schmidtiana, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 489.—Japon.
Schrenkiana, *Ledeb. Fl. Ross.* ii. 578.—Sibir.
scopaeformis, *Ledeb. l. c.*—Sibir.
scoparia, *Waldst. & Kit. Pl. Rar. Hung.* i. 66. t. 65.—Europ. ; Orient ; Ind. or.
scoparlorum, *A. Gray, in Proc. Ac. Sc. Philad.* 1863 (1864) 66.—Am. bor. occ.
selengensis, Turcz. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 50 = vulgaris.
semanineusis, Bess. l. c. 65 = senjavinensis.
Semenk, *Forsk. Fl. Aegypt. Arab.* 218.—Arabia.
Semenkeia, *Vacquem. ex Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 76.—Ind. or.
senjavinensis, *Bess. l. c.* 64.—Am. occ. et As. or. Arct.
sericea, *Weber, in Stechm. Artem.* 16. n. 5.—Sibir.
sericophylla, Rupr. in Beitr. Pfl. Russ. Reich. ii. 41 = borealis.
Seriphium, Pourr. ex Willd. & Lange, Prod. Fl. Hisp. ii 71 = hispanica.
Seriphium, Wallr. Sched. Crit. 458 = maritima.
serotina, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 240 (*Lehm. Reliq. Bot.* 165).—Turkestan.
serrata, *Nutt. Gen. Am.* ii. 142.—Am. bor.
sibirica, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 439 = Tanacetum sibiricum.
Sieberi, Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 80 = A. Herba-alba.
sipontina, Tenore, Fl. Nap. v. 215 = caerulescens
Siversiana, *Ehrh. ex Willd. Sp. Pl.* iii. 1845.—Rossia ; As. bor. ; Reg Himal.
sogdiana, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 243 (*Lehm. Reliq. Bot.* 167).—Turkestan.
soongarica, *Schrenk, Enum. Pl. Nov.* 49 (1871).—Soongaria ; Turkest.
spicata, *Wulf. ex Jacq. Fl. Austr.* v. 46.—Europ.
spicigera, C. Koch, in Linnæa, xxiv. (1851) 345 = fragrans.
spinescens, *Eaton, in Bot. King, Exp.* 180. t. 19.—Am. bor. occ.
spithamæa, Pursh, Fl. Am. Sept 522 = borealis.
splendens, *Willd. Sp. Pl.* iii. 1822.—As Min. ; Reg. Caucas.
Stechmanniana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 35 = chamaemelifolia.
Stelleri, Stev. ex Ledeb. Fl. Ross. ii. 568 = borealis.
Stelleriana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 79.—Am. bor. ; Graecia ; Kamtschatka.
sternutatoria, Roxb. Hort. Beng. 61 ; Fl. Ind. iii. 423 = Centipeda orbiculata.
Steveniana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 94 = glomerata.
Stewartii, C. B. Clarke, Comp. Ind. 168 = annua.
Strachеyi, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 104.—Tibet. occ.
stricta, *Edgew. in Trans. Linn. Soc.* xx. (1846) 78.—Reg. Himal.
stricta, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xl. (1867) 1. 204 = desertorum.
stricta, Heyne, ex DC. Prod. vi. 374 = Cyathocline lyrata.
suaveolens, Fisch. in DC. l. c. 110 = annua.
suaveolens, *Lam. Encyc.* i. 269.—Europ.
subcana, *Ford. Ind. Sem. Hort. Divion.* (1848) 18.—Gallia.
subcanescens, Boiss. Diagn. Ser. I. vi, 91 = persica.
subulata, Willd. ex Steud. Nom. ed. II. i. 140 = arenaria.
subulosa, Willd ex DC. Prod. vi. 94 = arenaria.
subvisceos, *Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 24. —Sibir.
succulenta, *Ledeb. Fl. Alt.* iv. 81.—Sibir.
supina, Stev. ex DC. Prod. vi. 112 = hololeuca.
sylvatica, *Maxim. Prim. Fl. Amur.* 161.—Mandchur.
sylvatica, Schlecht. ex Ledeb. Fl. Ross. ii. 581 = laciniata.
tanacetifolia, All. Fl. Pedem. i. 166 = spicata.
tanacetifolia, Georgi, It. i. 239 = sacrorum.
tanacetifolia, *Linn. Sp. Pl.* 848.—Sibir.
tanacetoides, Less. in Linnæa, vi. (1831) 210 = Matricaria multiflora.
tatarica, *Weber, ex Stechm. Artem.* 19.—Tataria.
taurica, *Bieb. Fl. Taur. Cauc.* ii. 291.—Taur.
taurica, Less. in Linnæa, x. (1835-36) 170 = Lessingiana.
taurina, Willd. Sp. Pl. iii. 1837 = fragrans.
tenella, *Weber, ex Stechm. Artem.* 26.—Hab. ?
tenuiflora, Jacquem. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 63 = Tanacetum tenuifolium.
tenuifolia, Adams, in DC. Prod. vi. 107 = Adamsii.
tenuifolia, Ehrh. ex Ledeb. Fl. Ross. ii. 567 = commutata.
tenuisecta, Moench, Meth. 571 = pontica.
tenuifolia, Willd. Sp. Pl. iii. 1819 = Eupatorium foeniculaceum.
tenuissima, Spreng. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 80 = procera.
Thunbergiana, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 432 = spicata.
thuscula, *Cav., in Anal. Cienc. Nat.* iv. (1801) 86.—Ins. Teneriffæ.
tibetica, Hook. f. Fl. Brit. Ind. iii. 329 = minor.
tiflisensis, Adams, ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 63 = fasciculata.
Tilesii, Ledeb. in Mém. Acad. Pétersb. v. (1805) 568 = vulgaris.
tomentella, *Trautv. in Bull. Soc. Nat. Mosc.* xxxix. (1866) 1. 351.—As. centr.
Tournefortiana, *Reichb. fc. Exot.* i. t. 5.—Orient ; Reg. Himal
transsilvanica, Schur, Enum. Pl Transs. 321 = Baumgarteni.
Trautvetteriana, *Bess. in Mém. Sav. Etr. Pétersb.* iv. (1846) 464. t. 5.—Orient ; Afghan.
trichophylla, Wall. Cat. n. 3305 = scoparia.
tridentata, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 398.—Am. bor. occ.
trifida, *Nutt. l. c.*—Am. bor. occ.
trifida, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 198 = Tanacetum trifidum.
trifurcata, Desf. Cat. Hort. Par. ed. III. (1829) 161 et 399 = Fontanesiana.
trifurcata, *Steph. ex Spreng. Syst.* iii. 488.—Soongaria.
Tmaiana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 67.—Sibir.
Tscherniviana, Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 33 = arenaria.
Turczaninowiana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 28.—Afghan.
turkestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 311.—Turkestan.
scrarnica, Bess. ex Nym. Consp. 378 = Hausknechtii.
umbelliformis, Lam. Encyc. i. 262 = Mutellina.
umbrosa, Turcz. ex DC. Prod. vi. 113 = sclengensis.
uralensis, *Spreng. ex Rem. in Flora*, xvii. (1834) 1. *Britt.* 16, nomen.—Sibir.
Vahliana, *Kostel. Allg. Med.-Pharm. Fl.* ii. 608.—Persia.
valesiaca, Lam. Encyc. i. 269 = A. Herba-alba.
vallesiaca, All. Fl. Pedem. i. 169 = seq.
vallesiana, *Lam. Encyc.* i. 269.—Helv. austr. ; Ital.
variabilis, *Tenore, Fl. Neap. Prod. App.* v. 28.—Neap. ; Hisp.
Veriatorum, Lamotte, in Mém. Assoc. Franç. Cong. Clerm.-Fer (1876) 511 = vulgaris.
vermiculata, Linn. Mant. 291.—Sinche artemisioides.
vermiculata, Schaug. ex DC. Prod. vi. 90 = borealis.
vestita, *Wall. Cat.* n. 3301.—Reg. Himal. , China.
Villarsii, Gren. & Godr. Fl. Fr. ii. 180 = Baumgarteni.
violacea, Desf. Tabl. ed. II. 112 = vulgaris.
violacea, Ledeb. in Mém. Acad. Pétersb. v. (1805) 567 = borealis.
virens, Moench, Meth. Suppl. 233 = vulgaris.
virgata, *Gueldenst. ex Ledeb. Fl. Ross.* ii. 600.—Ucrania.
virgata, Richards. in Frankl. Journ. App. 306 = frigida.
viridifolia, Ledeb. Ind. Sem. Hort. Dorpat. (1828) 2 = rupestris.
viridis, Willd. ex DC. Prod. vi. 124 = rupestris.
viscosa, DC. l. c. 95 = glutinosa.
vulgaris, Burm. f. Fl. Ind. 177 = indica.
vulgaris, *Linn. Sp. Pl.* 848.—Reg. temp. bor.
Wadei, Edgew. in Trans. Linn. Soc. xx. (1846) 72 = annua.
Wallichiana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 69 = Roxburghiana.
Willdenowii, Dietr. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 115 = camphorata.
Wrightii, *A Gray, in Proc. Am. Acad.* xix. (1883) 48. —Mexico.
Wulfenii, Schleich. Cat. Pl. Helv. ed. IV. 8 = Mutellina.
Zollingeriana, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126.—Malaya.

ARTENEMA, G. Don, in Sweet, Hort. Brit. ed. III. 516 (1839) = **Artanema**, D. Don (Scrophul.).

ARTHOSTEMA, Neck. Elem. ii. 280 (1790) = **Gnetum**, Linn.

ARTHRATHERUM, Beauv. Agrost. 32. t. 8. f. 8 (1812) = **Aristida**, Linn. (Gramin.).
arenarium, Nees, in Lehm. Pl. Preiss. ii. 98 = Aristida arenaria.
brachyatherum, Coss. & Dur. Expl. Algér Glum. 291 = Aristida acutiflora.

ARTHRATHERUM :—
brevifolium, Nees, Fl. Afr. Austr. i. 183 = Aristida brevifolia.
calogtilum, Jaub. & Spach, Illustr. iv. 54 = Aristida paradisea.
capense, Nees, Fl. Afr. Austr. 177 = Aristida capensis.
ciliatum, Nees, in Linnæa, vii. (1832) 267 = Aristida ciliata.
comosum, J. Gay, ex Kunth, Enum. Pl. i. 104 = Aristida stipiformis.
elatum, Boiss. & Buhse, Diag. Ser. II. iv. 128 = Aristida pogonopitila.
hirtigiume, Jaub. & Spach, Illustr. iv. 52 = Aristida hirtigluma.
hygrometricum, Beauv. Agrost. 33 = Aristida hygrometrica.
Hystrix, Nees, in Linnæa, vii. (1832) 267 = Aristida vestita.
Kotschyi, Hochst. ex Trin. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 159 = Aristida funiculata.
lutescens, Nees, Fl. Afr. Austr. i. 179 = Aristida lutescens.
magnpotamicum, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. vii. (1848) 124 = Aristida implexa.
nanuquense, Nees, Fl. Afr. Austr. i. 185 = Aristida nanuquensis.
obtusum, Nees, l. c. 179 = Aristida obtusa.
paniciformis, Boiss. & Buhse, Diag. Ser. II. iv. 127.—Persia.
plumosum, Nees, Fl. Afr. Austr. i. 182 = Aristida plumosa.
pogonopitilum, Jaub. & Spach, Illustr. iv. 56 = Aristida pogonopitila.
pungens, Beauv. Agrost. 33 = Aristida pungens.
rigidum, Schult. Mant. ii. 212 = Aristida rigescens.
Schimperi, Nees, Fl. Afr. Austr. 178 = Aristida ciliata.
stipoides, Beauv. Agrost. 33 = Aristida stipoides.
subacaule, Nees, Fl. Afr. Austr. 180 = Aristida subacaulis.
uniplume, Nees, l. c. 181 = Aristida uniplumis.
vestitum, Nees, l. c. 174 = Aristida vestita.
Zeyheri, Nees, l. c. 177 = Aristida Zeyheri.

ARTHRAXON, Beauv. Agrost. 111. t. 11. f. 6 (1812).
GRAMINEÆ, Benth. & Hook. f. iii. 1128.
ALECTORIDIA, A. Rich. Tent. Fl. Abyss. ii. 447. t. 99 (1851).
BATHRATHRUM, Nees, in Edinb. N. Phil. Journ. xviii. (1835) 150.
LASIOLYTRUM, Steud. in Flora, xxix. (1846) 18.
LUCAEA, Kunth, Rev. Gram. ii. 483. t. 159 (1830).
PLEUROLITUS, Trin. Fund. Agrost. 174 (1820).
PSILOPOGON, Hochst. ex Reichb. Nom. 26 (1841).
ciliaris, *Beauv. l. c.*—Orient; As. et Afr. trop.; Austral.
colonatus, Hochst. in Flora, xxxix. (1856) 188 = ciliaris.
cuspidatus, *Hochst. l. c.*—Ind. or.
echinatus, Hochst. l. c. = lanceolatus.
gracilis, Hochst. l. c. = ciliaris.
japonicum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 288 = ciliaris.
Junghuhnii, *Hochst. in Flora*, xxxix. (1856) 189.—Java.
lanceolatus, *Hochst. l. c.* 188.—Ind. or.; Abyss.; Japon.
lanceifolius, Hochst. l. c. = microphyllus.
Langsdorffianus, Hochst. l. c. = ciliaris.
lasiocolos, Hochst. l. c. 189 = ciliaris.
major, Hochst. l. c. 188 = ciliaris.
micans, *Hochst. l. c.*—Ind. or.
microphyllus, *Hochst. l. c.*—Ind. or.
minor, Hochst. l. c. = microphyllus
modest, Hochst. l. c. = ciliaris.
plumbeus, Hochst. l. c. 189 = ciliaris.
rudis, *Hochst. l. c.* 188 = cuspidatus.
Schimperi, Hochst. l. c. 189 = ciliaris.
Schmidtii, Hochst. l. c. = ciliaris.
serrulatus, Hochst. l. c. 188 = lanceolatus.
submuticus, Hochst. l. c. = ciliaris.
violaceus, Hochst. l. c. = ciliaris.

ARTHROCARPUM, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 510. *LEGUMINOSÆ*.
gracile, Balf. f. l. c. 511.—Ins. Socotra.

ARTHROCHILUS, F. Muell. Fragm. i. 42 (1858) = Dyakaea, Lindl. (Orchid.).
irritabilis, F. Muell. l. c. 43 = D. irritabilis.

ARTHROCHORTUS, Lowe, in Hook. Kew Journ. viii. (1856) 301 = Lolium, Linn. (Gram.).
loliaceus, Lowe, l. c.—Ins. Desertas (Madera).

ARTHROCLIANTHUS, Baill. Adansonia, ix. (1870) 296. *LEGUMINOSÆ*.
saxgulans, *Baill. l. c.* 297.—N. Caled.

ARTHROCNEMUM, Moq. Chenop. Enum. 111. (1840), partim. *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 65.
ambigunm, *Moq. Chenop. Enum.* 112.—Austral.
arbuscula, Moq. l. c. 113 = Salicornia arbuscula.
australasicum, F. Muell. ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876) 308 = Salicornia quinqueflora.
Belangerianum, Moq. Chenop. Enum. 112 = Halocnemum caspicum.
bidens, Nees, in Lehm. Pl. Preiss. i. 633 = Salicornia bidens.
caspicum, Moq. in DC. Prod. xiii. 11.150 = Halocnemum caspicum.
ciliolatum, *Bunge, in Linnæa*, xxviii. (1856) 578, nomen.—As. Min.?
corallioides, *Loac. & Pard. Ser. Imperf. Pl. Arag.* 90; ed. II. 351.—Hispan.
fruticosum, Moq. Chenop. Enum. 111 = Salicornia fruticosa.
fruticosum, Torr. in Stansbury, Rep. 394 = Spirostachys occidentalis.
glaucum, *Delile, Fl. Egypt.* 69; *Ung. Sternb. in Atti Congr. Bot. Firenze*, 1874 (1876) 283.—Græcia; Afr. trop.; Ind. or.
halocnemoides, Nees, in Lehm. Pl. Preiss. i. 632 = Salicornia arbuscula.
indicum, *Moq. Chenop. Enum.* 113.—Ind. or.; Afr. trop.
indicum, Thw. Enum. Pl. Zeyl. 246 = Salicornia brachiata.
macrostachyum, *Bunge, ex Ung. Sternb. Vers. Syst. Salicorn.* 67.—Reg. Mediterr.
macrostachyum, Moq. ex Delporte, in Ind. Sem. Hort. Bot. Taur. (1854) 35 = glaucum.
macrostachyum, Torr. Bot. Mex. Bound. Surv. 184 = Spirostachys occidentalis.
triandrum, F. Muell. Fragm. i. 139 = Salicornia robusta.

ARTHROLEPIS, Boiss. Diagn. Ser. I. xi. 14 (1849) = Achillea, Vaill. (Compos.).
imbricata, Boiss. l. c. 15 = Ach. membranacea.
membranacea, Boiss. l. c. = Ach. membranacea.

ARTHROLOBIUM, Reichb. Consp. 161 (1828) = ARTROLOBIUM, Desv. = Ornithopus, Linn. (Legum.).
obcrctastum, Auct. = O. ebractentus.

ARTHROLOBUS, Andrz. ex DC. Syst. ii. 480 (1821) = Rapistrum, Crantz (Crucif.).
perennis, Andrz. ex DC. l. c. 481 = R. perenne.
rugosus, Andrz. ex DC. l. c. 482 = R. rugosum.

ARTHROLOBUS, Stev. ex DC. Syst. ii. 579 (1821) = Sisrigma, DC. (Crucif.).
torulosus, Stev. ex DC. l. c. 580 = S. torulosum.

ARTHROMISCHUS, Thw. Enum. Pl. Zeyl. 47 (1858) = Paramignya, Wight (Rutac.).
armata, Thw. l. c. = P. armata.

ARTHROPHYLLUM, Blume, Bijdr. 878 (1826). *ARALIACEÆ*, Benth. & Hook. f. i. 944.
MORMOPHIS, Jack, ex Wall. Cat. n. 4981 (1831).
alternifolium, Malog. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 734 = planatum.
Blumeanum, Zoll. & Moritzi, Syst. Verz. Pfl. Zoll. 41 = diversifolium.
ceylanicum, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 27. —Ceylan.
dilatatum, *Miq. Fl. Ind. Bat.* i. 1. 768.—Java.
diversifolium, *Blume, Bijdr.* 879.—Malaya.
elliptienm, Blume, l. c. = diversifolium.
javanicum, Blume, l. c. = diversifolium.
ovalifolium, Jungh. & De Vriese, in Ann. Sc. Nat. Sér. III. vi. (1846) 117 = diversifolium.
planatum, *Malog. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 734.—Malaya.
reticulatum, *Blume, ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 27.—Malaya.
serratifolium, *Miq. Fl. Ind. Bat. Suppl.* 241.—Ins. Banca.

ARTHROPHYLLUM, Boj. Hort. Maurit. 221 (1837) = Phyllarthron, DC. (Bignon.).
comorense, Boj. l. c. = P. comorense.
madagascariense, Boj. l. c.—Madag.

ARTHROPHYTUM, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 211. *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 70.
Lehmannianum, *Bunge, Revis. Anab.* 46 = subcilifolium.
subcilifolium, *Schrenk, l. c.*—Turkestan.; Soongaria.

ARTHROPODIUM, R. Br. Prod. 276 (1810). *LILIACEÆ*, Benth. & Hook. f. iii. 790.
appendiculatum, *Asa-Lall. ex Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop.* xi. 59.—Austral.
caespitosum, Spreng. Syst. ii. 87 = Stypandra caespitosa.
candidum, *Raoul, Choix Pl. Nouv. Zél.* 14.—N. Zel.
capillipes, *Endl. in Lehm. Pl. Preiss.* ii. 84.—Austral.
cirratum, *R. Br. in Bot. Mag.* t. 2350.—N. Zel.
dianellicaulum, *F. Muell. Fragm.* x. 56.—Austral.
elongatum, DC. Cat. Hort. Monsp. 81, in obs. = paniculatum.
fimbriatum, R. Br. Prod. 276 = Dichopogon Sieberianus.
glaucum, Spreng. Syst. ii. 87 = Stypandra glauca.
hispidum, Spreng. l. c. = Anthericum hispidum.
imbricatum, Spreng. l. c. = Stypandra glauca.
laxum, Hook. f. Fl. Tasm. ii. 51 = Dichopogon strictus.
laxum, *Sieber, ex Schult. f. Syst.* vii. 441.—Austral.
Lindleyi, Kunth, Enum. Pl. iv. 621 = paniculatum.
minus, Lindl. Bot. Reg. t. 866 = paniculatum.
minus, *R. Br. Prod.* 276.—Austral.
muricatum, Spreng. Syst. ii. 87 = Anthericum muricatum.
neo-caledonicum, *Baker, in Journ. Linn. Soc.* xv. (1877) 253.—N. Caled.
paniculatum, *R. Br. Prod.* 276.—Austral.
pendulum, *DC. Cat. Hort. Monsp.* 80 = paniculatum.
Preissii, *Endl. in Lehm. Pl. Preiss.* ii. 85.—Austral.
scabrum, Spreng. Syst. ii. 87 = Agrostocrinum stipandroides.
strictum, R. Br. Prod. 276 = Dichopogon strictus.
umbellatum, Spreng. Syst. ii. 87 = Stypandra umbellata.

ARTHROPOGON, Nees, Agrost. Bras. 219 (1829). *GRAMINEÆ*, Benth. & Hook. f. iii. 1119.
stipiatus, *Hackel, in Sitzb. Acad. Wien*, lxxxiv. (1884) 125.—Cuba.
villosus, *Nees, Agrost. Bras.* 219.—Bras.

ARTHROSOLEN, C. A. Mey. in Bull. Phys-Math. Acad. Pétersb. i. (1843) 359. *THYMELÆACEÆ*, Benth. & Hook. f. iii. 193.
anthylloides, C. A. Mey. l. c. = Lasiosiphon anthylloides.
calocephalus, *C. A. Mey. l. c.*—Afr. austr.
chrysanthus, *Salma, ex Schweinf. Beitr. Fl. Aethiop.* 166.—Afr. trop.
glaucescens, *Oliver, in Journ. Linn. Soc.* xv. (1871) 96.—Afr. trop.
gymnostachys, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 359.—Afr. austr.
inconspicuus, *Meisn. in DC. Prod.* xiv. 560.—Afr. austr.
laxus, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 359.—Afr. austr.
ornatus, *Meisn. in DC. Prod.* xiv. 559.—Afr. austr.
pimelioides, *Meisn. l. c.* 560.—Natal.
polycephalus, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 360.—Afr. austr.
sericocephalus, *Meisn. in DC. Prod.* xiv. 561.—Afr. austr.
somalensis, *Franch. in Revoil, Conutl.* 62. t. vi.—Somali.
spicatus, *C. A. Mey. in Bull. Phys-Math. Acad. Pétersb.* i. (1843) 359.—Afr. austr.
subglobosus, Endl. Gen. Suppl. iv. 11. 62 = Wikstroemia subglobosa.

ARTHROSTEON, Hamb. Retsia, i. 212 (1866) = Acacia, Willd. (Legum.).
stipulatum, Hamb. l. c. = Acacia comosa.

ARTHROSTACHYA, Link, Hort. Berol. i. 251 (1827) = Gundelia, Boerh. (Comp.).
scaposa, Link, l. c. = G. gundelioides.

ARTHROSTACHYS, Desv. Opusc. 74 (1831) = **Andropogon**, Linn. (Gramin.).
gracilis, Desv. l. c.—Ind. or.

ARTHROSTEMMA, Ruiz & Pav. Fl. Per. iv. t. 336 (1802). *MELASTOMACEÆ*, Benth. & Hook. f. i. 740.
HETERONOMA, DC. Prod. iii. 122 (1828).
MELANIUM, Rich. ex DC. l. c. 155 (1828).
alatum, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 85.—Venezuela.
alpestre, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 357 = Pterolepis alpestris.
anguturense, DC. Prod. iii. 135 = Pterolepis glomerata.
Aublettii, DC. l. c. 137 = Comolia Aublettii.
Bonplandii, DC. l. c. 126 = Monochætum Bonplandii.
brachyandrum, Cham. in Linnæa, ix. (1834) 454 = Pterolepis glomerata, Tibouchina versicolor.
calcaratum, DC. Prod. iii. 138 = Monochætum calcaratum.
campanulare, DC. l. c. 136 = Brachyotum campanulare.
campanulare, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 85.—Am. trop.
capillatum, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1850) 361.—Ind. occ.
cataphractum, Naud. l. c. 360 = Pterolepis cataphracta.
caudulatum, Ruiz & Pav. Fl. Per. iv. t. 327 = Aciotis caudulata.
cerastifolium, DC. ex Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 122 = Tibouchina cerastifolia.
ciliatum, *Ruiz & Pav. Fl. Per.* iv. t. 326.—Peruv.
clidemioides, Miq. in Linnæa, xxii. (1849) 541 = Pterolepis clidemioides.
cubense, A. Rich. Ess. Fl. Cub. 550 = Chætolepis cubensis.
Deppeanum, Schlecht. & Cham. in Linnæa, v. (1830) 566 = Monochætum Deppeanum.
dicranantherum, DC. Prod. iii. 138 = Monochætum dicranantherum.
exiguum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 359 = Pterolepis exigua.
filiforme, Naud. l. c. 358 = Pterolepis filiformis.
floribundum, Schlecht. in Linnæa, xiii. (1839) 481 = Monochætum rivulare.
fragile, *Lindl. in Journ. Hort. Soc.* iii. (1848) 74.—Mexic.
glomeratum, Cham. in Linnæa, (1834) ix. 454 = Pterolepis glomerata.
Gaudichianum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 359 = Pterolepis Gaudotiana.
herbaceum, DC. Prod. iii. 137 = Tibouchina herbacea.
heterotrichum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 360 = Pterolepis filiformis.
hirsutissimum, DC. Prod. iii. 137 = Tibouchina herbacea.
hirsutissimum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 357 = Pterolepis parciflora.
ladanoides, DC. Prod. iii. 136 = Pterolepis pusilla.
lanceolatum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 186.—Cuba.
latifolium, *D. Don, in Mem. Wern. Soc.* iv. (1823) 299.—Guiana.
linarium, D. Don, l. c. = Monochætum linarium.
latescens, *DC. Prod.* iii. 136 = Brachyotum latescens.
macranthera, Miq. in Linnæa, xxii. (1849) 540 = Tibouchina sebastianopoltana.
Martinianum, DC. Prod. iii. 137 = Tibouchina sebastianopoltana.
multiflorum, DC. l. c. 138 = Monochætum multiflorum.
multiflorum, D. Don, in Mem. Wern. Soc. iv. (1823) 299 = Monochætum dicranantherum.
myrtoideum, DC. Prod. iii. 138 = Monochætum myrtoideum.
nitidum, R. Grah. in Edinb. N. Phil. Journ. xii. (1832) 186 = Tibouchina nitida.
nummularioides, DC. Prod. iii. 137 = Comolia summularioides.
paniculatum, D. Don, in Mem. Wern. Soc. iv. (1823) 299 = Oxyporn paniculata.
parciflorum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 357 = Pterolepis parciflora.
piloselloides, DC. Prod. iii. 136 = Castrasella piloselloides.
polygonoides, DC. l. c. 137 = Pterolepis polygonoides.
pusillum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 357 = Pterolepis pusilla.
pusillum, Naud. l. c. 361 = Pterolepis pusilla.

ARTHROSTEMA :—
quinquenerve, DC. Prod. iii. 136 = Brachyotum quinquenerve.
rosmarinifolium, DC. l. c. = Brachyotum rosmarinifolium.
Salzmannii, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 356 = Pterolepis Salzmannii.
Schiedeanum, Steud. Nom. ed. II. i. 141 (= *Pleroma longiflorum* !).—Mexic.
strygnoneve, Cham. in Linnæa, ix. (1834) 453 = Tibouchina nitida.
versicolor, DC. Prod. iii. 137 = Tibouchina versicolor.
villosum, DC. l. c. = Comolia villosa.
villosum, Mart. Herb. Fl. Bras. n. 499 = Comolia ovalifolia.
volubile, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 32.—N. Granat.
Weddellianum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 356 = Pterolepis Weddelliana.

ARTHROSTEMMA, DC. Prod. iii. 134 (1828) pro parte = **Brachyotum**, Triana (Melast.).

ARTHROSTEMMA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 355 = **Pterolepis**, Miq. (Melast.).

ARTHROSTYGMA, Steud. Nom. ed. II. i. 141 (1840) = **Petrophila**, R. Br. (Proteac.).

ARTHROSTYLIDIUM, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 117. *GRAMINEÆ*, Benth. & Hook. f. Gen. iii. 1208.
AULONEMIA, Goudot, in Ann. Sc. Nat. Sér. III. v. (1846) 75. t. 5.
Barchellii, *Munro*, *in Trans. Linn. Soc.* xxvi. (1868) 48.—Bras.
capillifolium, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 531.—Ind. occ.
cubense, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 118. t. 4. f. 13.—Cuba.
excelsum, *Griseb. Fl. Brit. W. Ind.* 529.—Ind. occ.
fimbriatum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 531.—Ind. occ.
Hasskarl, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 117. t. 3. f. 12.—Peruv.
leptophyllum, *Doell, in Mart. Fl. Bras.* ii. III. 175.—Bras.
longiflorum, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 41.—Venezuela.
maculatum, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 118. t. 5.—Peruv.
pubescens, *Rupr. l. c.* 119. t. 4. f. 14.—Ind. occ.
racemiflorum, *Steud. Syn. Pl. Gram.* 386.—Mexic.
Schomburgkii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 41.—Guiana.
Trinii, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 119. t. 4. f. 15.—Bras.

ARTHROSTYLIS, R. Br. Prod. 229 (1810). *CYPERACEÆ*, Benth. & Hook. f. iii. 1058.
aphylla, Boeck. in Linnæa, xxxvii. (1871–72) 524 = Actinoschœnus Thouarii.
aphylla, *R. Br. Prod.* 229.—Austral.
chinensis, Benth. Fl. Hongk. 397 = Actinoschœnus chinensis.
filiformis, Thw. Enum. Pl. Zeyl. 352 = Actinoschœnus filiformis.
Thouarii, Kunth, Enum. Pl. ii. 284 = Actinoschœnus Thouarii.

ARTHROSTYLIS, Boeck. in Linnæa, xxxvii. (1871–72) 523 = **Actinoschœnus**, Benth. (Cyperac.).
aphylla, Boeck. l. c. 524 = Act. Thouarii.
filiformis, Boeck. l. c. = Act. filiformis.

ARTHROTAXIS, Endl. Gen. Suppl. i. 1372 (1841) err. typ. = **Athrotaxis**, D. Don (Conif.).

ARTHROTHAMNUS, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251 = **Euphorbia**, Linn.
Berg'ii, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 63.—Afr. austr.
brachiatus, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251; et Abh. Akad. Berl. (1860) 62 = E. brachiata.
Burmanni, Klotzsch & Garcke, ll. cc. = E. Burmanni.
cyparissus, Klotzsch & Garcke, ll. cc. 251, 63 = E. cyparissias.

ARTHROTHAMNUS :—
dendriformis, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251; et Abh. Akad. Berl. (1860) 62.—Afr. austr.
Echlonii, Klotzsch & Garcke, ll. cc. 251, 63.—Afr. austr.
scopiformis, Klotzsch & Garcke, ll. cc. = E. scopiformis.
Tiruculli, Klotzsch & Garcke, ll. cc. 251, 62 = E. Tiruculli.

ARTHROTRICHUM, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 500; et Edinb. N. Phil. Journ. N. S. xvii. (Jan.–Apr. 1863) 324 = **Trichintuna**, R. Br. (Amarant.).
calostachyum, F. Muell. ll. cc. = T. calostachyum.

ARTHROZAMIA, Relchb. Consp. 40 (1828) = **Encephalartos**, Lehm. (Cycad.).

ARTOCARPUS, Forst. Char. Gen. 101. t. 51 (1776). *URTICACEÆ*, Benth. & Hook. f. iii. 376.
DURIO, Adans. Fam. ii. 399 (1763).
POLYPHEMA, Lour. Fl. Cochinch. 546 (1790).
RADERMACHIA, Thunb. in Handl. Vet. Akad. Stockh. (1778) 252.
RIMA, Sonner. Voy. Nouv. Guin. 99. t. 57 ad 60 (1776).
SITODIUM, Gaertn. Fruct. i. 344. t. 71, 72 (1791).
angustifolia, *Roxb. Hort. Beng.* 66, nomen.—Ins. Malay.
anisophylla, *Miq. Fl. Ind. Bat. Suppl.* 422.—Sumatra.
benghalensis, Roxb. ex Wall. Cat. sub n. 4655 = A. Lakoocha.
bijaria, Wall. ex Miq. Ann. Bot. Mus. Lugd. Bat. iii. 218 = Sloetia Sideroxylon.
biformis, *Miq. Fl. Ind. Bat. Suppl.* 419.—Sumatra.
Blumei, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 111.—Java.
brasiliensis, *Gomes*, *in Mem. Ac. Lisboa*, iii. (1812) *Mem. dos Corresp.* 84.—Bras.
calophylla, *Kurz*, *For. Fl.* ii. 431.—Tenasserim.
Camansi, *Blanco*, *Fl. Filip.* ed. I. 670.—Ins. Philipp.
canarana, *Miq. in Verh. Nederl. Inst.* III. v. (1852) (*Anal. Bot. Ind.*) 20.—Ind. or.
Cannoni, *Hort. Bull. ex Van Houtte*, *in Fl. des Serres*, xxi. (1875) 181.—Ins. Soc.
cerifera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 212.—Java.
Chama, Buch.-Ham. ex Wall. Cat. sub n. 4657 = A. Chaplasha.
Chaplasha, Spreng. Syst. iii. 804 = A. Polyphema.
Chaplasha, *Roxb. Hort. Beng.* 66; *et Fl. Ind.* iii. 525.—Ind. or.
communis, Forst. Char. Gen. 100 = incisa ?
Cumingiana, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 113.—Ins. Philipp.
cuspidata, Griff. Notul. iv. 400 = rigida.
Dadak, *Miq. Fl. Ind. Bat. Suppl.* 420.—Sumatra.
dasyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 212.—Ins. Celebes.
dimorphophylla, *Miq. Fl. Ind. Bat. Suppl.* 417.—Ins. Banca.
echinata, Roxb. Hort. Beng. 66; et Fl. Ind. iii. 527 = rigida.
elastica, *Reinw. in Blume*, *Bijdr.* 481.—Borneo.
elongata, Miq. Fl. Ind. Bat. Suppl. 419 = Sloetia Sideroxylon.
erythrocarpa, Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 213 = A. Dadak.
erythrocarpa, Teijsm. ex Miq. l. c. 212 = dasyphylla.
exculpta, *Hort.—Cf. Gard. Chron.* (1878) ii. 84.—Hab. ?
Finlaysoniana, *Wall. Cat.* n. 4662.—Malaya.
glanca, *Blume*, *Bijdr.* 483.—Java.
glaucescens, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 120.—Java.
Gomeziana, *Wall. ex Trec. l. c.* 113.—Ind. or.
heterophylla, Lam. Encyc. iii. 309 = integrifolia.
hirsuta, *Lam. Encyc.* iii. 311.—Ind. or.
hirsutissima, *Kurz*, *in Tijdschr. Nederl. Ind.* xxvii. (1864) 382.—Ins. Banca.
hypargyrea, *Hance*, *ex Benth. Fl. Hongk.* 325.—Hongkong.
incisifolia, Stokes, Bot. Mat. Med. iv. 331 = incisa.
incisa, *Linn. f. Suppl.* 411.—Malaya; Ins. Pacif.
integrifolia, *Linn. f. l. c.* 412.—Ind. or.; Malaya.
Jaca, Lam. Encyc. iii. 309 = integrifolia.
Kemando, *Miq. Fl. Ind. Bat. Suppl.* 418.—Sumatra.

ARTOCARPUS :—
Kurtz, *Zoll. Syst. Verz. Ind. Archip.* 89, 95.—Malaya.
lacnicata, *Hort.—Cf. Gard. Chron.* (1878) ii. 159; *et Rev. Hortic.* (1877) 38.—Ins. Pacif.
Lanceha, Roxb. Hort. Beng. 66, nomen = A. Lakoocha.
lasvis, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 134.—Java.
Lakoocha, *Roxb. Fl. Ind.* iii. 574.—Ind. or.; Malaya.
lamellosa, Blanco, Fl. Filip. ed. 1. 667 = nitida.
lanceæfolia, *Roxb. Hort. Beng.* [108]; *Fl. Ind.* iii. 537.—Malaya.
lanceolata, *Trèc. in Ann. Sc. Nat. Sér.* III. viii. (1847) 181.—Manilla.
Limpato, *Miq. Fl. Ind. Bat. Suppl.* 421.—Sumatra.
madagascariensis, *Bej. Hort. Maurit.* 290.—Madag.
mariannensis, *Trèc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 114.—Ins. Mariana.
maxima, Blanco, Fl. Filip. ed. 1. 669 = integrifolia.
mollis, *Miq. Fl. Ind. Bat Suppl.* 420.—Sumatra.
mollis, Wall. Cat. n. 4661 = A. Lakoocha.
Nanea, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 7.—Malaya.
nitida, *Trèc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 119.—Ins. Philipp.
nobilis, *Thw. Enum. Pl. Zeyl.* 282.—Zeylan.
odoratissima, *Blanco, Fl. Filip.* ed. 1. 671.—Ins. Philipp.
ovata, Blanco, l. c. 666 = Cumingiana.
ovata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 7.—Malaya.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 424.—Java.
parviflora, *Vesgt, in Syll. Ratisb.* ii. (1826) 58.—Hab.?
pedunculatia, *Kurz. in Journ. Bot.* xiii. (1875) 331.—Ins. Nicobar.
pestolaris, *Miq. Fl. Ind. Bat. Suppl.* 422.—Sumatra.
philippensis, *Lam. Encyc.* iii. 210.—Ins. Philipp.
pilosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 7.—Malaya.
pilosa, Reinw. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 211 = A. Polyphema.
Polyphema, *Pers. Syn.* ii. 531.—Malaya.
pomiformis, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 400 = Gomesiana.
Pongin, Dennst. Schlussel, 30 = Brousonetia integrifolia.
pubescens, *Blume, Bijdr.* 481.—Java.
pubescens, Moon, Cat. Pl. Ceyl. 81 = nobilis.
pubescens, Willd. Sp. Pl. 189 = integrifolia.
reticulata, Heyne, ex Wall. Cat. sub n. 4655 = A. Lakoocha.
reticulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 218.—Ins. Celebes.
Riedelii, *Miq. l. c.*—Ins. Celebes.
rigida, *Blume, Bijdr.* 482.—Burma; Malaya.
rigida, Wall. Cat. n. 4656 A = Gomeziana.
Rima, Blanco, Fl. Filip. ed. 1. 671 = incisa.
rufa, *Miq. Fl. Ind. Bat. Suppl.* 423.—Malaya.
rufescens, *Korz, For. Fl.* ii. 481.—Ind. or.
rufescens, *Miq. Fl. Ind. Bat. Suppl.* 430.—Malaya.
serrata, Roxb. Hort. Beng. 66, nomen.—Ind. or.
Tampang, Miq. Fl. Ind. Bat. Suppl. 421 = rufescens, *Miq.*
tephrophylla, *Miq. l. c.* 422.—Sumatra.
Teysmanni, *Miq. l. c.* 418.—Sumatra.
tiphyphila, *Miq. Fl. Ind. Bat.* i. 1. 289.—Malaya.
variana, *Miq. Fl. Ind. Bat. Suppl.* 417.—Sumatra.
venenosa, *Zoll. & Mor. in Nat. en Genees. Arch. Neerl. Indie*, ii. (1846) 213.—Java.
Vriesiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 212.—Ins. Celebes.
Zollingeriana, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 90, 93.—Malaya.

ARTROLOBIUM, Desv. Journ. Bot. i. (1813) 121. t. 4 = Coronilla, Linn., Ornithopus, Linn., etc. (Leguminosæ).
creticum, Desv. l. c. = Coronilla cretica.
durum, Desv. l. c. = Ornithopus durus.
ebracteatum, Desv. l. c. = Ornithopus ebracteatus.
micranthum, Benth. in Mart. Fl. Bras. xv. i. 51.—Am. trop.
repandum, DC. Prod. ii. 311 = Ornithopus repandus.
scorpioides, Desv. Journ. Bot. i. (1813) 121 = Coronilla scorpioides.
Spruchii, Kalen. in Bull. Soc. Nat. Mosc. xviii. (1845) i. 232.—Tauria.
tauricum, Kalen. l. c.—Tauria.

ARUANA, Barm. f. Ind. Alt. (sign. G, verso) (1769) = **Myristica**, Linn.
silvestris, Burm. f. l. c. = M. moschata.

ARUBA, Aubl. Pl. Gui. i. 393. t. 135 (1775) = **Simaba**, Aubl. (Simarub.).
guianensis, Aubl. l. c.—Guiana.

ARUBA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 172. t. 19, 27, 29 = **Almeidea**, St. Hil. (Rutac.).
accuminata, Nees & Mart. l. c. 175 = Almeidea acuminata.
alba, Nees & Mart. l. c. 174 = Almeidea alba.
corrulea, Nees & Mart. l. c. = Almeidea cærulea.
myriacantha, Nees & Mart. ex Stend. Nom. ed. II. i. 142 = Almeidea myriantha.

ARUM, [Tourn.] Linn. Syst. ed. I. (1735), partim. *AROIDEAE*, Benth. & Hook. f. Gen. iii. 967.
GYMNOSTIUM, Schott, in Oestr. Bot. Wochenbl. (1855) 17.
abyssinicum, A. Rich. Tent. Fl. Abyss. ii. 353 = Sauromatum abyssinicum.
aculeatum, Stend. Nom. ed. I. 78 = Caladium aculeatum.
alatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 4.—Malaya.
albinervium, Kotschy, ex Engl. in DC. Monog. Phan. ii. 590 = hygrophilum.
albispathum, Hort. ex Engl. l. c. 591 = italicum.
albispathum, Stev. ex Ledeb. Fl. Ross. iv. 9 = orientale.
alpinum, Schott & Kotschy, in Bot. Zeit. ix. (1851) 385 = maculatum.
alternans, Engl. in DC. Monog. Phan. ii. 572 = Philodendron crassinervium.
alternum, Vell. Fl. Flum. ix. t. 116 = Philodendron crassinervium.
amphibium, Vell. l. c. 112 = Philodendron amphibium.
aquaticum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 5.—Malaya.
angulatum, *Grif. Notul.* iii. 143.—Ind. or.
aphyllum, Hook. in W. Gray, Western Afr. 386. t. A = Amorphophalli sp.
arborescens, Linn. Sp. Pl. 967 = Montrichardia arborescens.
Arisarum, Linn. l. c. 966 = Arisarum vulgare.
Arisarum, Lour. Fl. Cochinch. 655 = Typhonium cochinchinense.
atrorubens, Alt. Hort. Kew. iii. 315 = Arisæma atrorubens.
atrorubens, Spreng. Syst. iii. 769 = Pinellia tuberifera.
auritum, Linn. Syst. ed. X. 1251; Pl. Jam. Pugill. 26 = Syngonium auritum.
auritum, Vell. Fl. Flum. ix. t. 113 = Syngonium Vellozianum.
balsamita, Buchoz, Dec. Pl. vii. 11 = pictum.
Bassevianum, Schott, in Oestr. Bot. Zeitschr. (1858) 349 = maculatum.
bertolfiium, *Mill. Gard. Dict.* ed. VIII. n. 12.—Am. bor.
bicolor, Ait. Hort. Kew. iii. 316 = Caladium bicolor.
bubalinum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 4.—Malaya.
bulbiferum, Roxb. Hort. Beng. 65; Sims, Bot. Mag. t. 2072 = Amorphophallus bulbifera.
bulbiferum, Salisb. Prod. 260 = Pinellia tuberifera.
bulbosum, Pers. ex Kunth, Enum. Pl. iii. 54 = Pinellia tuberifera?
byzantinum, Schott, in Oestr. Bot. Wochenbl. (1857) 213 = italicum.
cæruleo-punctatum, Zipp. ex Kunth, Enum. Pl. iii. 33 = Amorphophallus variabilis.
calyptratum, Salisb. Prod. 261 = Arisarum vulgare.
campanulatum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 509 = Amorphophallus campanulatus.
canariense, Webb & Berth. Phyt. Canar. iii. 203 = italicum.
cannæfolium, Linn. f. Pl. Suria. n. 133 = Spathiphyllum cannæforme?
carratracense, Haenel. in Bot. Zeit. iv. (1846) 315 = Biarum Bovei.
catawrium, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 4.—Malaya.
ciliatum, Hort. ex Lond. Encyc. Suppl. ii. 1306 = Cryptocoryne ciliata.

ARUM :—
ciliciense, Kotschy, ex Engl. in DC. Monog. Phan. ii. 559 = rupicolum.
clavatum, Desf. Cat. Hort. Par. ed. III. 385 = Sauromatum pedatum.
Colocasia, Linn. Sp. Pl. 965 = Colocasia antiquorum.
colocasioides, Desf. Cat. Hort. Par. ed. III. 7 et 385 = Colocasia antiquorum.
concinnatum, Schott, Ic. Aroid. t. 39 = italicum.
conophalloides, Kotschy, ex Schott, Prod. Aroid. 97 = orientale.
consobrinum, *Schott, l. c.* 87.—Reg. Cauc.
cordatum, Vell. Fl. Flum. ix. t. 111 = Philodendron cordatum.
cordifolium, Bory, Voy. i. 376; ii. 67 = Alocasia indica.
corsicum, Loisel. Fl. Gall. ii. 617 = pictum.
costatum, Wall. Tent. Fl. Nep. t. 19 = Arisæma costatum.
crassipes, Boiss. Diagn. Ser. I. xiii. 9 = Helicophyllum crassipes.
crenatum, Wight, in Hook. Bot. Misc. ii. (1831) 100 = Theriophonum crenatum.
creticum, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 9.—Creta.
crinitum, Ait. Hort. Kew. iii. 314 = Helicodiceros muscivorus.
crudele, Salisb. Prod. 261 = Dieffenbachia seguina.
cucullatum, Lour. Fl. Cochinch. ii. 586 = Alocasia cucullata.
curvatum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 506 = Arisæma curvatum.
cuspidatum, Blume, Cat. Gew. Buitenz. 101 = Typhonium cuspidatum.
cuspidatum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 506 = Arisæma cuspidatum.
cylindraceum, Gasp. in Guss. Fl. Sic. Syn. ii. ii. 597 = italicum.
cyprium, Schott, in Bonplandia, x. (1861) 368 = Dracontium.
decurrens, Blanco, Fl. Filip. ed. I. 656 = Amorphophallus decurrens.
detruncatum, *C. A. Mey. ex Schott, Prod. Aroid.* 8.—Pers. bor.
Dioscoridis, *Sibth. & Sm. Fl. Græc. Prod.* ii. 245.—Græc.; An. Min.
discolor, *Veigt, in Syll. Ratisb.* ii. 50.—Hab.!
dissectatum, Dalac, Fl. Hautes-Pyr. 46 = italicum.
divaricatum, Linn. Sp. Pl. 966 = Typhonium divaricatum.
diversifolium, Blume, Cat. Gew. Buitenz. 102 = Typhonium diversifolium.
Dracontium, Linn. Sp. Pl. 964 = Arisæma Dracontium
Dracontium, Lour. Fl. Cochinch. 583 = Pinellia Wawræ.
Dracontium, Thunb. Fl. Jap. 238 = Arisæma Thunbergii.
Dracontium, Vell. Fl. Flum. ix. t. 108 = Staurostigma Lanchnathianum.
Dracunculus, Linn. Sp. Pl. 964 = Dracunculus vulgaris.
Dracunculus, Thunb. Fl. Jap. 238 = Arisæma japonicum.
echinatum, Wall. Pl. As. Rar. ii. 30. t. 136 = Arisæma echinatum.
Ehrenbergii, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 356.—Syria.
elongatum, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) ii. 67 = orientale.
erubescens, Wall. Pl. As. Rar. ii. 30. t. 135 = Arisæma erubescens.
esculentum, Linn. Sp. Pl. 965 = Colocasia antiquorum.
esserteum, Salisb. Prod. 260 = A. Dracontium.
filiforme, Reinw. in Blume, Cat. Gew. Buitenz. 102 = Arisæma filiforme.
flagelliferum, *Grif. Notul.* iii. 144.—Ind. or.
flagelliforme, Roxb. Hort. Beng. 65; Lodd. Bot. Cab. t. 896 = Typhonium cuspidatum.
flavum, Forsk. Fl. Ægypt. Arab. 157 = Arisæma flavum.
fœtidum, Salisb. Prod. 261 = italicum.
foveolatum, Roth, Pl. Ind. Or. 365 = Pinellia tuberifera.
foveolatum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 501 = Alocasia fœtidum.
foveolatum, Wight, Ic. iii. t. 789 = Alocasia decipiens.
fragus, Salisb. Prod. 260 = esculentum.
gracile, Roxb. Hort. Beng. 65; Fl. Ind. iii. 506 = Typhonium gracile.
gracile, Unver. ex Schott, Prod. Aroid. 81 = maculatum.
graminiforme, Lam. Encyc. iii. 20 = Biarum graminiforme.

ARUM:—
graminеum, [Soland. in] Russ. Aleppo, ed. II. ii. 264 = Biarum Russellianum.
grandifolium, Jacq. Hort. Schoenbr. t. 189 = Philodendrum grandifolium.
gratum, Schott, Syn. Aroid. 11 = orientale.
Griffithii, *Schott*, l. c. 15.—Afghanist.
guttatum, Salisb. Prod. 260 = Dracunculus vulgaris.
guttatum, Wall. Pl. As. Rar. ii. 10. t. 115 = Sauromatum guttatum.
hederaceum, Jacq. Enum. Pl. Carib. 31 = Philodendron Hoffmannii.
hederaceum, Willd. Sp. Pl. iv. 486 = Philodendron hederaceum.
Heidrichii, Orph. ex Boiss. Fl. Orient. v. 40 = maculatum.
helleborifolium, Jacq. Coll. iii. 217 = Xanthosoma helleborifolium.
heterophyllum, Ascher, ex Blume, Rumphia, i. 121 = Helicophyllum Rauwolffii.
hygrophilum, *Boiss. Diagn.* Ser. I. xiii. 8.—Syria.
immaculatum, Reichb. Ic. Fl. Germ. vii. 6; Schott, Prod. Aroid. 94 = italicum.
incomptum, Schott, l. c. 89 = orientale.
incurvatum, Lam. Fl. Fr. iii. 588 = Arisarum vulgare.
indicum, Lour. Fl. Cochinch. ii. 596 = Colocasia indica.
integrifolium, Link, Enum. Hort. Berol. ii. 894 = Aglaonema integrifolium.
intermedium, Schur, ex Schott, Prod. Aroid. 91 = maculatum.
intortum, [Soland. in] Russ. Aleppo, ed. II. ii. 264 = Helicophyllum Rauwolffii.
italicum, *Mill. Gard. Dict.* ed. VIII. n. 2.—Europ. Jacquemontii, *Blume, Rumphia*, i. 118.—As. trop.
Korolkowi, *Regel, in Act. Hort. Petrop.* ii. (1873) 407. —Turkestan.
Kotschyi, *Boiss. & Hohen. Diagn.* Ser. I. xiii. 8.—Pers. bor.
laceratum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4 = seq. ?
lacerum, Jacq. Hort. Schoenb. iv. 35 = Philodendron lacerum.
lanceolatum, Vell. Fl. Flum. ix. t. 114 = Philodendron crassinervium.
Liebоldii, Schott, Prod. Aroid. 27 = Dioscoridis.
lingulatum, *Linn. Pl. Jam. Pugill.* 26; *Syst.* ed. X. 1351.—Jamaica.
liniferum, Arruda, in Koster, Trav. Bras. 483 = Montrichardia linifera.
lividum, Salisb. Prod. 260 = Colocasia antiquorum.
longicyrrhum, Schott, Syn. Aroid. 14 = hygrophilum.
longispathum, Reichb. Ic. Fl. Germ. vii. 5 = orientale.
lyratum, Roxb. Hort. Beng. [108]; Fl. Ind. iii. 508 = Amorphophallus lyratus.
macrorrhizum, Linn. Sp. Pl. 965 = Alocasia macrorrhiza.
macrourum, *Bunge, Enum. Pl. Chin. Bor.* 67.—China bor.
maculatum, All. Fl. Pedem. ii. 228 = italicum.
maculatum, Gueldenst. ex Ledeb. Fl. Ross. iv. 9 = orientale.
maculatum, *Linn. Sp. Pl.* 966.—Europ.
Malyi, Schott, Prod. Aroid. 98 = maculatum.
margaritiferum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 513 = Plesmonium margaritiferum.
maximum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 5.—Malaya.
maximum, Schott, Prod. Aroid. 86 = italicum.
melanopus, *Boiss. Fl. Orient.* v. 40.—Oriens.
minutum, Willd. Sp. Pl. iv. 484 = Theriophonum crenatum.
monsteroides, Roxb. Hort. Beng. 65; Fl. Ind. iii. 497 = Colocasia montana.
mucronatum, Lam. Encyc. iii. 12 = Alocasia macrorrhiza.
mucronotum, Spreng. Syst. iii. 769 = Xanthosoma mucronotum.
Mulleri, Zipp. ex Blume, Rumphia, i. 148 = Amorphophallus Mulleri.
Murreyi, J. Grah. Cat. Bomb. Pl. 229 = Arisaema Jacquemontii.
muscivorum, Linn. f. Suppl. 410 = Helicodiceros muscivorus.
nepenthoides, Wall. Tent. Fl. Nep. 26. t. 18 = Arisaema nepenthoides.
Nicholsii, Schott, Prod. Aroid. 86 = italicum.
nigrum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 212.—Europ.
nigrum, Vell. Fl. Flum. ix. t. 107 = Xanthosoma violaceum.
Nordmanni, Schott, Syn. Aroid. 12 = orientale.
novum, *Linn. f. Pl. Suria.* n. 192.—Guiana.
nunsidicum, Schott, Syn. Aroid. 10 = italicum.
nymphaefolium, Roxb. Hort. Beng. 66; Fl. Ind. iii. 495 = Colocasia antiquorum.
oblongum, Vell. Fl. Flum. ix. t. 115 = Philodendron oblongum.
obtusilobum, *Link, Enum. Hort. Berol.* ii. 898.—Hab.?
odoratum, Heynh. Nom. ii. 46 = seq.
odorum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 499 = Alocasia odora.
orientale, *Bieb. Fl. Taur. Cauc.* ii. 207.—Taur.; Caucas.
orientale, Vis. ex Nym. Consp. 754 = nigrum.
orixense, R. Br. Prod. 336 = Typhonium Brownii.
orixense, Roxb. ex Andr. Bot. Repos. t. 556 = Typhonium trilobatum.
ovatum, Linn. Sp. Pl. 967 = Lagenandra toxicaria.
palaestinum, *Boiss. Diagn.* Ser. I. xiii. 6.—Orient.
pedatum, Fisch. ex Spreng. Syst. iii. 769 = Sauromatum guttatum.
pedatum, Willd. Enum. Hort. Berol. Suppl. 54 = Sauromatum pedatum.
pellucidum, Fulchir, ex Kunth, Enum. Pl. iii. 42 = Caladium bicolor.
peltatum, Lam. Encyc. iii. 13 = Caladium esculentum.
pentaphyllum, Linn. Sp. Pl. 964 = Arisaema pentaphyllum.
pentaphyllum, Vell. Fl. Flum. ix. t. 104 = Xanthosoma pentaphyllum.
peregrinum, Linn. Sp. Pl. 966 = Alocasia macrorrhiza.
Petteri, Schott, Syn. Aroid. 12 = orientale.
philistaeum, *Kotschy, ex Schott, Prod. Aroid.* 79.—Palaestina.
phrygium, *Boiss. ex Tchihat. Asia Min. Bot.* ii. 648.—Asia min.
pictum, *Linn. f. Suppl.* 410.—Ins. balear.; Corsic.
pictum, Pett. Bot. Wegw. (1882); ex Schott, Syn. Aroid. 13 = orientale.
pinnatifidum, Jacq. Hort. Schoenb. iii. 31 = Philodendron pinnatifidum.
pinnatisectum, Vell. Fl. Flum. ix. t. 110 = Philodendron bipinnatifidum.
polymorphum, *Buchl. in Am. Journ. Sc.* xlv. (1843) 178.—Am. bor.
polyphyllum, *Link, Enum. Hort. Berol.* ii. 898.—Asia min.
ponticum, Schott, in Bonplandia, x. (1862) 148 = italicum.
probosciferum, Linn. Sp. Pl. 966 = Arisarum proboscideum.
ptychospermum, Zipp. ex Kunth, Enum. Pl. iii. 36 = Typhonium cuspidatum.
pulchrum, Salisb. Prod. 260 = Caladium bicolor.
pumilum, Kotschy, ex Engl. in DC. Monog. Phan. ii. 586 = philistaeum.
pumilum, Lam. Encyc. iii. 8 = Typhonium trilobatum.
punctatum, Desf. Cat. Hort. Par. ed. III. 385 = Alocasia indica.
punctulatum, Zipp. ex Kunth, Enum. Pl. iii. 34 = Amorphophallus bulbifer.
purpureum, *Thunb. Fl. Jap.* 20.—Japon.
pyrrosanum, Dufour, in Lap. Hist. Abr. Fl. Pyr. Suppl. 148 = maculatum.
Pythonium, Mart. Auswahl, 13 = Zomicarpa Pythonium.
quinarium, Nutt. Gen. Am. ii. 222 = Arisaema quinarium.
ramosum, *Link; Enum. Hort. Berol.* ii. 894.—Hab.?
rapiformе, Roxb. Hort. Beng. 65; Fl. Ind. iii. 497 = Alocasia rapiformis.
Rauwolffii, Blume, Rumphia, i. 122 = Helicophyllum Rauwolffii.
rauwolfii, Rodesch, 'in Bald. Diar. xxvIII. 32'; ex G. F. W. Mey, Esseq. 328 = Dieffenbachia Seguine.
ringens, *Thunb. in Trans. Linn. Soc.* ii. (1794) 337.—Japon.
Rumphii, Gaudich. in Freyc. Voy. Bot. 197. t. 34 = Amorphophallus campanulatus.
rupicola, *Boiss. Diagn.* Ser. I. xiii. 7.—Syria.
sagittaefolium, Hort. ex Steud. Nom. ed. II. i. 148 = Caladium bastifolium.

ARUM:—
sagittaefolium, Linn. Sp. Pl. 966 = Xanthosoma sagittaefolium.
sagittatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 4.—Malaya.
sagittifolium, Link, Enum. Hort. Berol. ii. 893 = Xanthosoma bastifolium.
sagittifolium, Rodsch. Observ. 30 = Uropsatha sagittifolia.
Saururus, Burger, ex Kunth, Enum. Pl. iii. 19 = Arisaema japonicum.
Seguine, Jacq. Enum. Pl. Carib. 31 = Dieffenbachia Seguine.
Seguinum, Linn. Sp. Pl. ed. II. 1371 = Dieffenbachia Seguine.
serratum, Thunb. in Trans. Linn. Soc. ii. (1794) 338 = Arisaema japonicum.
sessiliflorum, *Roxb. Hort. Beng.* 65; *Fl. Ind.* iii. 507. —Ind. or.
simense, Hort. ex Engl. in DC. Monog. Phan. ii. 571 —Sauromatum venosum.
speciosum, Wall. Tent. Fl. Nep. 29. t. 20 = Arisaema speciosum.
spectabile, Schott, in Oestr. Bot. Wochenbl. (1857) 175 = Dioscoridis.
spectabile, Zipp. ex Kunth, Enum. Pl. iii. 34 = Amorphophallus bulbifer
spiculatum, Blume, Rumphia, i. 121 = Helicophyllum crassipes.
spirale, Retz. Obs. i. 30 = Cryptocoryne spiralis.
spirale, Salisb. Prod. 259 = Helicodiceros crinitus.
stercorarium, Zipp. ex Kunth, Enum. Pl. iii. 33 = Amorphophallus variabilis.
strepiceros, *Sieber, ex Steud. Nom.* ed. II. i. 148.— Palaestina.
subulatum, Desf. Cat. Hort. Par. ed. III. 7 et 385 = Pinellia tuberifera.
sylvaticum, Roxb. Hort. Beng. [108]; Fl. Ind. iii. 511 = Synaiptorias silvaticus.
syriacum, Blume, Rumphia, i. 119 = Dioscoridis.
syriacum, Spreng. Syst. iii. 768 = Biarum Russellianum.
tenuifolium, Linn. Sp. Pl. 967 = Biarum tenuifolium.
ternatum, Thunb. Fl. Jap. 233 = Pinellia tuberifera.
tortuosum, Wall. Pl. As. Rar. ii. 10 = Arisarum tortuosum.
tragopsantinum, Schott, ex Engl. in DC. Monog. Phan. ii. 592 = italicum.
trifidum, Desf. Cat. Hort. Par. ed. III. 7 et 385 = Hydrosme Fontanesii.
trifoliatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 4.—Malaya.
trilobatum, Linn. Sp. Pl. 965 = Typhonium divaricatum.
trilobatum, Thunb. Fl. Jap. 234 = Pinellia tripartita.
tripartitum, Jacq. Hort. Schoenb. ii. 33. t. 190 = Philodendron tripartitum.
triphyllum, Hont. Plantenk. ii. 11. 184 = Pinellia tuberifera.
triphyllum, Linn. Sp. Pl. 965 = Arisaema atrorubens.
triphyllum, Thunb. Fl. Jap. 233 = Arisaema ringens.
variolatum, Schott, Prod. Aroid. 81 = Xanthosoma violaceum.
venosum, Ait. Hort. Kew. iii. 315 = Sauromatum guttatum.
venosum, Balb. ex Engl. in DC. Monog. Phan. ii. 487 = Xanthosoma helleborifolium.
ventricosum, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 25.— Bras.
vermiculatum, Vell. Fl. Flum. ix. t. 108 = Caladium bicolor.
vernale, Salisb. Prod. 261 = maculatum.
virginicum, Linn. Sp. Pl. 966 = Peltandra virginica.
viviparum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 496 = Remusatia vivipara.
virescens, *Stegl. in Denkschr. Acad. Wien*, i. (1885) 6. —Persia.
wittatum, Salisb. Prod. 260 = Arisaema atrorubens.
vulgare, Lam. Fl. Fr. iii. 587 = maculatum.
xanthorrhizon, Jacq. Hort. Schoenbr. ii. 32 = Xanthosoma sagittifolium.
Zeleborі, Schott, Prod. Aroid. 94 = maculatum.

ARUNA, Schreb. Gen. i. 26 (1789) = AROUNA, Aubl. = Diallium, Linn. (Legumin.).

ARUNCUS, Linn. Syst. ed. I. (1735); Gen. ed. I. 310 (1737) = **Spiræa**, Tourn. (Rosac.).
americanus, Raf. Sylva Tellur. 152 = S. Aruncus.
astilboides, Maxim. in Act. Hort. Petrop. vi. (1879) 171 = S. Aruncus.
sylvester, Kostal. Ind. Hort. Prag. 15 = S. Aruncus.
vulgaris, Raf. Sylva Tellur. 152 = S. Aruncus.

ARUNDINA, Blume, Bijdr. 401 (1825). *ORCHIDEÆ*, Benth. & Hook. f. iii. 521.
DILOCHIA, Lindl. Gen. et Sp. Orch. 88 (1830).
affinis, Griff. Notul. iii. 330 = chinensis.
bambusifolia, *Lindl. in Wall. Cat.* n. 3781.—Ind. or.; Malaya.
chinensis, *Blume, Bijdr.* 402.—Reg. Himal.; China.
densa, *Lindl. in Bot. Reg.* (1842) *Misc.* 25.—Singapore.
Meyeni, *Reichb. f. in Linnæa*, xxv. (1852) 237.—China.
minor, *Lindl. Gen. et Sp. Orch.* 126.—Zeylan.
pentandra, *Reichb. f. Xenia Orch.* ii. 11.—Java.
Philippi, Reichb. f. in Linnæa, xxv. (1852) 227 = chinensis.
pulchella, *Teijsm. & Binn. in Nederl. Kruid. Arch.* iii. (1856) 400.—China.
pulchra, *Miq. in Fourn. Bot. Néerl.* i. (1861) 90.—China.
speciosa, Blume, Bijdr. 401. t. 78 = densiflora.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 37.—Ins. Pacif.
Wallichii, *Reichb. f. Xenia Orch.* ii. 13. t. 105.—Malaya.

ARUNDINARIA, Michx. Fl. Bor. Am. i. 73 (1803). *GRAMINEÆ*, Benth. & Hook. f. iii. 1207.
LUDOLFIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 320.
MACHOMAX, Rafin. in Med. Repos. New York, v. (1808) 352.
MIEGIA, Pers. Syn. i. 101 (1805).
THAMNOCALAMUS, Munro, in Trans. Linn. Soc. xxvi. (1868) 33.
TRIGLOSSUM, Fisch. Cat. Jard. Gorenk. (1812).
acuminata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 33.—Mexic.
amplissima, *Nees, in Linnæa*, ix. (1834) 479.—Bras.
aristulata, *Doell, in Mart. Fl. Bras.* ii. iii. 165.—Bras.
attenuata, *Doell, l. c.* 170.—Bras.
bambusina, Trin. Fund. Agrost. 97 = macrosperma.
callosa, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 30.—Reg. Himal.
debilis, *Thw. Enum. Pl. Zeyl.* 375.—Zeylan.
decalvata, *Doell, in Mart. Fl. Bras.* ii. iii. 170.—Bras.
densiflora, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 32.—Zeylan.
distans, *Trin. in Mém. Acad. Pétersb. Sér.* VI. iii. 11. (1836) 221.—Bras.
elegans, *Kunth, in Fourn. As. Soc. Beng.* xiii. (1873) ii. 249.—Burma.
falcata, *Nees, in Linnæa*, ix. (1834) 478.—Reg. Himal.
fastuosa, *Hance, in Fourn. Bot.* xiv. (1876) 339.—China.
floribunda, *Thw. Enum. Pl. Zeyl.* 375.—Zeylan.
Fortunei, Fenzl, in Gard. Chron. (1876) ii. 773 = Simoni.
Fortunei, *Riviére, in Bull. Soc. Acclim.* Sér. III. v. (1878) 397.—Japon.
gigantea, *Nutt. Gen. Am.* i. 39.—Am. bor.
glaucescens, *Beauv. Agrost.* 144, 152.—Ind. or.; China!
Griffithiana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 30.—Reg. Himal.
Hindsii, *Munro, l. c.* 31.—Hongkong.
hirsuta, *Munro, l. c.* 30.—Reg. Himal.
hispida, Steud. Syn. Pl. Gram. 335 = Wightiana.
Hookeriana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 29.—Reg. Himal.
intermedia, *Munro, l. c.* 28.—Reg. Himal.
interrupta, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. 11. (1836) 220 = falcata.
japonica, *Sieb. & Zucc. ex Steud. Syn. Pl. Gram.* 334.—Japon.
khasiana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 28.—Reg. Himal.
kurilensis, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* viii. (1850) 121.—Ins. Kurile.

ARUNDINARIA:—
longifolia, *Fourn. Mus. Pl. Exsicc. Gram.* 131, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 586.—Mexic.
longiramea, Munro, in Trans. Linn. Soc. xxvi. (1868) 19 = simica.
macrosperma, *Michx. Fl. Bor. Am.* i. 74.—Am. bor.
macrostachya, *Nees, in Linnæa*, ix. (1834) 481.—Bras.
Maximowiczii, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 783 = Simoni.
Metake, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 284 = japonica.
microphylla, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 32.—Reg. Himal.
molinifromis, Hochst. ex Munro, l. c. 19 = Wightiana.
multiflora, *Doell, in Mart. Fl. Bras.* ii. iii. 166.—Bras.
parviflora, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. 11. (1835) 619 = Arthrostylidium Trinii.
pinifolia, Nees, in Mart. Fl. Bras. ii. 525 = Chusquea pinifolia.
procera, Wall. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 34 = spathiflora.
racemosa, *Munro, l. c.* 17.—Reg. Himal.
radiata, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 115. t. 15. f. 9.—Bras.
Schomburgkii, Benn. in Trans. Linn. Soc. xviii. (1841) 563 = Arthrostylidium Schomburgkii.
Simoni, *A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 774.—China.
simica, *Munro, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 235.—China.
spathiflora, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. 11. (1835) 617.—Reg. Himal.
stolonifera, Kurz, Cat. Hort. Bot. Calcut. 79 = Phyllostachys nigra.
subercata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 29.—Reg. Himal.
teres, *Mubl. Desc. Uber.* 191.—Am. bor.
tessellata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 31.—Afr. austr.
Tristani, *Munro, l. c.* 35.—N. Granat.
Trinii, Hack. in Mart. Fl. Bras. ii. iii. 174 = Arthrostylidium Trinii.
utilis, Cleghorn, in Journ. Agr. Soc. Ind. xiii. (1865) 388 = falcata.
verticillata, *Nees, in Mart. Fl. Bras.* ii. iii. 528.—Bras.
Walkeriana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 21.—Zeylan.
Wightiana, *Thw.* Enum. Pl. Zeyl. 444 = Walkeriana.
Wightii, Benth. Fl. Hongk. 484 = simica.

ARUNDINELLA, Raddi, Agrost. Bras. 27. t. 1 (1823). *GRAMINEÆ*, Benth. & Hook. f. iii. 1118.
ACRATHERUM, Link, Hort. Berol. i. 220 (1824).
BRANDTIA, Kunth, Rev. Gram. ii. 511. t. 170 (1830).
CALAMOCHLOË, Reichb. Consp. 52 (1828).
GOLDBACHIA, Trin. in Sprengl. Neue Entdeck. ii. 81 (1821).
HITCHCOCKIA, Trin. ex Kunth, Enum. Pl. i. 515 (1833).
THYSANACHNE, Presl, Thymen. Nov. Pl. Gen. (1829); et Rel. Haenk. i. 255 (1830).
Acratherum, *Nees, ex Steud. Nom.* ed. II. i. 14.—Ind. or.
agrostoides, Trin. Ic. xciii. t. 265 = holcoides.
anomala, *Steud. Syn. Pl. Gram.* 116.—Japon.
aristiseta, Rupr. in Bull. Acad. Petr. ix. (1842) ii. 242 = Deppeana.
avenacea, *Munro, ex Thw. Enum. Pl. Zeyl.* 362.—Zeylan.
brasiliensis, *Raddi, Agrost. Bras.* 27.—Bras.
ciliata, *Nees, ex Miq. in Verh. Nederl. Inst.* III. iv. (1851) 30.—Ind. or.
crinita, *Trin. in Linnæa*, x. (1836) 299.—Peruv.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 538.—Ind. occ.
Deppeana, *Nees, in Bonplandia*, iii. (1855) 84.—Mexic.
Schlechtendalii, Nees, Fl. Afr. Austr. 80 = nepalensis.
elegantula, *Hook. f. in Fourn. Linn. Soc.* vii. (1864) 193.—Afr. occ.
fasciculata, *Trin. Gram. Sp.* xxiii. t. 267.—Bras.
fuscata, *Nees, ex Buese, in Mig. Pl. Jungh.* 320.—Ind. or.
glabra, *Hook. & Arn. Bot. Beech. Voy.* 287.—China.

ARUNDINELLA:—
Goeringii, *Steud. Syn. Pl. Gram.* 118.—Japon.
hirsuta, Hochst. ex Steud. l. c. 115 = setifera.
himalensis, *Nees, ex Steud. l. c.*—Reg. Himal.
holcoides, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 71 = Brandtia holcoides.
indica, *Trin. ex Steud. Nom.* ed. II. i. 143.—Ind. or.
khasiana, *Nees, ex Steud. Syn. Pl. Gram.* 118.—Reg. Himal.
latifolia, Fourn. Mus. Pl. Exsicc. Gram. 54; ex Hemsl. Biol. Centr. Am. Bot. iii. 515 = Deppeana.
martinicensis, *Trin. Gram. Panic.* ii. 85.—Ins. Martinic.
mesophylla, *Nees, ex Steud. Syn. Pl. Gram.* 116.—Ind. or.
Metzii, *Hochst. ex Miq. in Verh. Nederl. Inst.* III. iv. (1851) 31.—Ind. or.
Mikani, Nees, Agrost. Bras. 465 = brasiliensis.
mutica, *Nees, ex Steud. Syn. Pl. Gram.* 116.—Ind. or.
nepalensis, *Trin. Sp. Gram.* t. 268.—Ind. or.
nervosa, Nees, ex Hook. & Arn. Bot. Beech. Voy. 287 = Pankeana semialata.
pallida, *Nees, Agrost. Bras.* 465.—Mexic.
peruviana, *Steud. Syn. Pl. Gram.* 118.—Peruv.
phragmitoides, *Griseb. Cat. Pl. Cub.* 234.—Cuba.
pilosa, Hochst. ex Miq. in Verh. Nederl. Inst. III. iv. (1851) 30 = ciliata.
pumila, *Steud. Syn. Pl. Gram.* 114.—Abyss.
purpurea, *Hochst. ex Steud. l. c.* 115.—Ind. or.
rigida, *Nees, Fl. Afr. Austr.* 80.—Afr. austr.
Schultzii, *Benth. Fl. Austral.* vii. 545.—Austral.
scoparia, Fourn. Mus. Pl. Exsicc. Gram. 56; ex Hemsl. Biol. Centr. Am. Bot. iii. 515 = Deppeana.
setifera, *Steud. Syn. Pl. Gram.* 116.—Ind. or.
setosa, *Trin. Gram. Panic.* 85.—Nepal.
stipoides, *Hack. in Sitzb. Acad. Wien*, lxxxix. (1884) 193.—Madag.
stricta, *Nees, in Hook. Kew Journ.* ii. (1850) 102.—Hongkong.
tenella, *Nees & Wight, ex Steud. Nom.* ed. II. i. 143.—Ind. or.
tricuspidula, *Buese, in Mig. Pl. Jungh.* 360.—Java.
trilocula, *Steud. ex Steud. Syn. Pl. Gram.* 116.—Zeylan.
Wallichii, *Nees, ex Steud. l. c.* 114.—Ind. or.
Zollingeri, *Steud. l. c.* 115.—Java.

ARUNDO, Besuv. Agrost. 60. t. 13. f. 2 (1812) = **Phragmites**, Trin. (Gram.).

ARUNDO, Tourn. ex Linn. Gen. ed. I. 19 (1737). *GRAMINEÆ*, Benth. & Hook. f. iii. 1179.
AMPHIDONAX, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836).
DONACIUM, Fries, Bot. Notiser, (1843) 181.
DONAX, Besuv. Agrost. 77 (1812), pro parte.
SCOLOCHLOA, Mert. & Koch, in Roehl. Deutschl. Fl. i. 538 (1823).
acutiflora, Pollini, Fl. Veron. i. 88 = Deyeuxia varia.
acutiflora, Schrad. Fl. Germ. i. 217 = Deyeuxia acutiflora.
ægyptia, Delile, Fl. Aeg. Illustr. 4 = A. Donax.
ægyptiaca, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 984 = pnoc.
agrostis, *Kth. in Linnæa*, xxxii. (1863) 307.—Europ.
agrostis, Lour. Fl. Cochinch. 57 = Bambusa agrestis.
Agrostis, Lag. ex Willk. & Lange, Prod. Fl. Hisp. i. 58 = Molinia cærulea.
Agrostis, Scop. Fl. Carn. ed. II. i. 88 = Deyeuxia acutiflora.
Agrostis, [Weber, in] Wigg. Prim. Fl. Holsat. 10 = Deyeuxia acutiflora!
amphidonax, Pursh, Fl. Am. Sept. i. 66 = Deyeuxia canadensis.
arundinacea, *Raddi, Obs. Madera*, 10.—Madera.
aticoides, *Petr. Enayc.* vi. 270.—Am. bor.
Aloysiocrux, Gandhi. ex Urv. in Mém. Soc. Linn. Par. iv. (1826) 602 = Festuca Aloysiocrux.
altaïcola, Hall. ex Roem. & Schult. Syst. iii. 688 = Deyeuxia Hallerana.
alpina, [Host.] Ver. Gew. Teutschl. 221; Schult. Oestr. Fl. ed. II. i. 189 = Calamagrostis tenella.
alpina, Vill. ex Kunth, Enum. Pl. i. 388 = Deyeuxia Hallerana.
altissima, Benth. Cat. Pl. Pyr. 62 = Phragmites communis.
Amphidonaxum, Cyrill. Pl. Rar. Neap. Fasc. ii. 30 = Amphidonax tenax.
arundinacea, Roem. & Schult. Syst. ii. 703 = Calamagrostis Halleriana.

ARUNDO :—

antarctica, Urv. in Mém. Soc. Linn. Par. iv. (1826) 809 = Festuca antarctica.
arborea, Mill. Gard. Dict. ed. VIII. n. 5 = Bambusa arundinacea.
arenaria, Gorter, Fl. Ingr. 18 = Deyeuxia neglecta.
arenaria, Linn. Sp. Pl. 81 = Ammophila arundinacea.
argentea, [*Claire.*] *Man. Herb.* 16.—Helvet.
australis, Cav. ex Roem. & Schult. Syst. ii. 511 = Phragmites australis.
australis, A. Rich. ex Steud. Nom. ed. II. i. 148, 250 = conspicua.
baltica, Flueg. ex Schrad. Fl. Germ. 228. t. 5. f. 4 = Ammophila baltica.
Bambos, Burm. ex Steud. Nom. ed. II. i. 149 = Bambusa agrestis.
Bambos, Linn. Sp. Pl. 81 = Bambusa arundinacea.
Bambu, Lour. Fl. Cochinch. 56 = prace.
bengalensis, Boj. Hort. Maurit. 268 = Phragmites communis.
bengalensis, *Retz. Obs.* v. 20.—Bengal.
bicolor, *Desf. Fl. Atlant.* i. 107 ; *Poir. Voy. Barb.* ii. 104.—Afr. bor.
Bieberstiniana, Roem. & Schult. Syst. ii. Mant. 385 = Calamagrostis littorea.
bifaria, *Retz. Obs.* iv. 23.—Ind. or.
bifora, Lam. Illustr. i. 196 = Ampelodesmos tenax.
Boivini, *Steud. Syn. Pl. Gram.* 428.—Madag.
brevipilis, Torr. Fl. U. St. 95 = Ammophila brevipilis.
Calamagrostis, Bieb. Fl. Taur. Cauc. i. 78 ; ii. 456 = Calamagrostis epigeios.
Calamagrostis, Hall. f. in Roem. Arch. i. ii. 10 = Deyeuxia Halleriana.
Calamagrostis, Leyss. Fl. Halens. 28 = Deyeuxia sylvatica.
Calamagrostis, Linn. Sp. Pl. 82 = Calamagrostis lanceolata, *Calamaria*, Saliseb. Prod. 24 = Calamagrostis lanceolata.
calycina, Willd. ex Steud. Nom. ed. II. i. 144 = nitida.
canadensis, Michx. Fl. Am. Bor. i. 73 = Deyeuxia canadensis.
canadensis, Steud. Nom. ed. II. i. 144 = Deyeuxia Nuttaliana.
canescens, [Weber, in] Wigg. Prim. Fl. Holsat. 10 = Calamagrostis epigeios, lanceolata.
canella, Molina, ex Steud. Syn. Pl. Gram. 336 = Chusquea Cumingii.
capillata, *Nom. ex Chatab. in Actes Linn. Soc. Bord.* xix (1853) 44.—Hab. ?
cillata, [*Claire.*] *Man. Herb.* 16.—Helvet.
cinnoides, Muhl. Desc. Uber. 187 = Deyeuxia Nuttaliana.
Clarionis, Loisel. Fl. Gall. 719 = Deyeuxia Halleriana.
coarctata, Poir. Encyc. Suppl. iv. 704 = Deyeuxia coarctata.
coarctata, Torr. Fl. U. St. 94 = Deyeuxia Nuttaliana.
collina, Tenore, Fl. Nap. t. 108 = Phlaeris.
colorata, Ait. Hort. Kew. i. 116 = Phalaris arundinacea.
confinis, Willd. Enum. Hort. Berol. 127 = Calamagrostis confinis.
conspicua, *Forst. f. Prod.* 9.—N. Zel.
contracta, Gillib. Exercit. ii. 542 = Calamagrostis lanceolata.
cupidata, Schult. Mant. iii. 504 = Calamagrostis cupidata.
cylindrica, J. Gay, ex Steud. Nom. ed. II. i. 144 =, Deyeuxia acutiflora.
diecca, Spreng. Syst. i. 361 = Gynerium argenteum.
dioica, *Linn. Fl. Cochinch.* 70.—Cochinch.
Donax, Georgi, Beschreib. Russ. Reich. iii. iv. 706 = Arundinaria inrilensis.
Donax, *Linn. Sp. Pl.* 81.—Reg. Mediterr. ; Orient.
effusa, C. C. Gmel. Fl. Bad. 202 = Calamagrostis littorea.
effusa, Poir. Encyc. Suppl. iv. 706 = Deyeuxia effusa.
Epanentiana, Roem. & Schult. Syst. ii. 511 = Phragmites Epanentiana.
epigeios, Fresh. ex Steud. Nom. ed. II. i. 144 = imperata arundinacea.
epigeios, Huds. Fl. Ang. ed. I. 48 ; ed. II. 54 = Calamagrostis lanceolata.
epigeios, Linn. Sp. Pl. 82 = Calamagrostis epigeios.
eriantha, Poir. Encyc. Suppl. iv. 705 = Deyeuxia eriantha.
exaltata, Saliseb. Prod. 24 = Bambusa arundinacea.
fasciata, Andd. Fl. Gat. f. 42 = Calamagrostis fasciata.

ARUNDO :—

fastuosa, Willd. ex Steud. Nom. ed. II. i. 144 = Gynerium saccharoides.
Fax, Lour. Fl. Cochinch. 58 = Melocanna humilis.
festucacea, Willd. Enum. Hort. Berol. 126 = Graphephorum arundinaceum.
festucoides, Desf. Fl. Atlant. i. 108 = Ampelodesmos tenax.
filiformis, *Hassk. in: Hoev. & De Vriese, Tijdschr.* x. (1843) 117.—Java.
flexa, Willd. ex Steud. Nom. ed. II. i. 144 = Deyeuxia canadensis.
flexuosa, Bronga. in Duperr. Voy. Bot. 29. t. 4 = Phragmites flexuosa.
flexuosa, [Clairv.] Man. Herb. 16 = Aira flexuosa.
fulvida, *Q. Buch. in Trans. N. Z. Inst.* vi. (1874) 242. —N. Zel.
Gayana, *E. Desv. in C. Gay, Fl. Chil.* vi. 326.—Valdivia.
gigantea, Walt. Fl. Carol. 81 = Arundinaria macrosperma.
glauca, Bieb. Fl. Taur. Cauc. i. 79 = Calamagrostis littorea.
glauca, Hornem. Hort. Hafn. 74 = Deyeuxia Nuttalliana.
graeca, Link, in Linnaea, ix. (1834) 136 = Phragmites communis.
gramineoides, Schrank, in Regenab. Denkschr. i. 11. (1818) 8 = Calamagrostis groenlandica.
Halleri, Willd. Prod. Fl. Berl. 60 = Stipa Calamagrostis.
Halleriana, Gand. Agrost. Helv. i. 97 = Deyeuxia Halleriana.
Henslowiana, *Hook. & Arn. Bot. Beech. Voy.* 248.—Chile.
Hostii, Roem. & Schult. Syst. ii. 508 = Deyeuxia varia.
humilis, Roem. & Schult. i. c. 510 = Calamagrostis lanceolata, tenella.
hyperborea, Trin. ex Steud. Syn. Pl. Gram. 336 (= *Dupontia psilanthantha*).
indica, *Neronha, in Verh. Batav. Gen.* v. (1799) ed. 1. Art. IV. 7.—Malaya.
intermedia, C. C. Gmel. Fl. Bad. i. 266 = Calamagrostis epigeios.
intacta, Saliseb. Fl. Aeg Illustr. 82 = Phragmites communis
junciformis, Poir. Encyc. Suppl. iv. 705 = Deyeuxia juncifolia.
Kakao, Steud. Syn. Fl. Gram. 194 = conspicua ?
Karka, *Retz. Obs.* iv. 21 ; *Roxb. Hort. Beng.* 6 ; *Fl. Ind.* i. 348.—Ind. or.
Kila, Spreng. ex Steud. Nom. ed. II. i. 144 = Gynerium argenteum.
laevis, Steud. Nom. ed. II. i. 144 = Pos laevis.
Langsdorffii, Link, Enum. Hort. Berol. i. 74 = Deyeuxia purpurea.
lapponica, Wahlenb. Fl. Lapp. 27. t. 1 = Deyeuxia lapponica.
latifolia, Saliseb. Prod. 24 = A. Donax.
laxa, Roem. & Schult. Syst. ii. 502 = Calamagrostis littorea.
Leersii, Gott. Syst. Vers. Oberlaus. 17 = Calamagrostis lanceolata.
Leersii, J. Koch, Hamdb. 49 = Stipa Calamagrostis.
levis, Poir. Encyc. Suppl. iv. 707 = Pos laevis.
ligulata, *Poir. i. c.* 706 = Deyeuxia ligulata.
lithuanica, Schult. Mant. ii. 502 = Calamagrostis lanceolata.
littorea, Schrad. Fl. Germ. 212 = Calamagrostis littorea.
littoralis, Steud. Nom. ed. II. i. 144 = Ammophila arundinacea.
madagascariensis, *Kunth, Rev. Gram.* i. 278. t. 48.—Madag.
maritima, *Agardh, in Flora*, vii. (1824) 561.—Scandin.
mauritanica, Poir. Voy. Barb. ii. 104 ; Desf. Fl. Atlant. i. 108 = Phlaeris.
maxima, Forsk. Fl. Aegypt. Arab. 24 = Phragmites communis.
maxima, Lour. Fl. Cochinch. 58 = Bambusa maxima.
microstachia, Lam. Illustr. i. 196 = Phlaeris.
minutiflores, *Brongn. in Duperr. Voy. Bot.* 29.—Ins. Moluce.
mitis, Lour. Fl. Cochinch. 57 = Bambusa mitis.
montana, [Clairv.] Man. Herb. 15 = Deyeuxia montana.
montana, Roem. & Schult. Syst. ii. 506 = Deyeuxia sylvatica.
multiglabra, Lour. Fl. Cochinch. 58 = Gigantochloa verticillata.

ARUNDO :—

Naga, Koen. ex Steud. Nom. ed. II. i. 144 = Phragmites australis.
Nayra, Koen. ex Steud. l. c. ii. 321 = prace.
neglecta, Ehrh. Beitr. vi. 84, 137 = Deyeuxia neglecta.
nemorosa, Schult. Oestr. Fl. ed. II. i. 187 = Deyeuxia varia.
nitida, *H. B. & K. Nov. Gen. et Sp.* i. 149. t. 688.—Ecuador.
occidentalis, *Sieber, ex Schult. Mant.* ii. 289.—Ins. Martin.
orientalis, Mill. Gard. Dict. ed. VIII. n. 6 = Saccharum fuscum.
pallens, *Muhl. ex Steud. Nom.* ed. II. i. 144.—Am. bor.
palustris, Saliseb. Prod. 24 = Phragmites communis.
penicillata, Labill. Nov. Holl. Pl. i. 26 = Danthonia racemosa.
Phragmites, Linn. Sp. Pl. 81 = Phragmites communis.
picta, Kunze, ex Steud. Nom. ed. II. i. 144 = Pos plicta.
pilosa, *Urv. in Mém. Soc. Linn. Par.* iv. (1826) 808.—Ins. Falkland.
pleatoria, *Lour. Fl. Cochinch.* 55.—Cochinch.
planifolia, Poir. Encyc. Suppl. iv. 707 = Deyeuxia planifolia.
plebeia, Poir. l. c. = Deyeuxia plebeia.
Pliniana, *Turra, Farset.* 11 ; *Fl. Ital. Prod.* 63.—Reg. Mediterr.
Plinii, Turra, Farset. 11 = Pliniana.
plumosa, Schult Mant. 504 = Stipa papposa.
pourformis, Labill. Nov. Holl. Pl. i. 27. t. 35 = Pos Billardieri.
poaeoides, Poir. Encyc. Suppl. iv. 705 = Pos subuniflora.
Pseudophragmites, Hall. f. in Roem. Arch. i. ii. 11 = Calamagrostis littorea.
Pseudophragmites, Lejeune, Fl. Spa, 64 = Phragmites communis.
Pseudophragmites, Schrad. Fl. Germ. 218 = Deyeuxia Halleriana.
Pseudophragmites, Wahlenb. Fl. Lapp. 28 = Calamagrostis phragmitoides.
pungens, Auct. ex Steud. Nom. ed. II. i. 144 = Phragmites communis.
purpurascens, Schult. Mant. iii. 508 = Deyeuxia purpurascens.
purpurea, Trin. in Spreng. Neue Entdeck. ii. 52 = Deyeuxia purpurea.
pygmaea, *Moretti, ex Pollini, in Brugnat. Giorn. Fis.* ix. (1816) 22.—Italia.
pygmaea, Spreng. Pugill. i. 9 = Trisetum hispanicum.
Quila, Molina, Sagg. Chil. 279 = Chusquea Fernandesiana.
Quila, Poir. Encyc. vi. 274 = Chusquea Quila.
quitensis, Poir. l. c. Suppl. iv. 704 = Muehlenbergia calamagrostides.
Ragamowskii, Hort.—Cf. Gard. Chron. (1876) ii. 547 = Bambusa tessellata.
ramosa, Schult. Oestr. Fl. ed. II. i. 185 = Calamagrostis lanceolata.
recta, Poir. Encyc. Suppl. iv. 706 = Deyeuxia recta.
Raynaudiana, *Kunth, Rev. Gram.* i. 276. t. 48.—Egypt.
rigida, Poir. Encyc. Suppl. iv. 705 = Deyeuxia rigida.
riparia, Balbi, Prod. 24 = Phalaris arundinacea.
riparia, Willd. ex Steud. Nom. ed. II. i. 144 = Calamagrostis lanceolata.
rivularis, Auct. ex Steud. Nom. ed. II. i. 144 = Phragmites communis.
Roshergeii, Kunth, Rev. Gram. i. 79 = A. Karka.
Rugi, *Molina, Sagg. Chil.* 279.—Chili.
Sabuli, *Kit. in Linnaea*, xxxii. (1863) 206.—Europ.
saccharoides, Poir. Encyc. Suppl. iv. 708 = Gynerium saccharoides.
sagittata, Pers. Syn. i. 102 = Gynerium saccharoides.
salina, Lam. Fl. Fr. iii. 616 = A. Donax.
Schleicheriana, Bess. Fl. Gallic. i. 69 = Calamagrostis lanceolata.
Schreberi, Schult. Mant. iii. 508 = Gastridium australe.
scriptoria, Linn. Amoen. Ac. iv. 150 = A. Donax.
sedenensis, Loisel. Fl. Gall. 719 = Deyeuxia varia.
Selloana, Schult. Mant. iii. 506 = Gynerium argenteum.
semiannularis, Labill. Nov. Holl. Pl. i. 26. t. 33 = Danthonia semiannularis.
semiverticula, Hort. Ticin. ex Steud. Nom. ed. II. i. 144 = Pliniana.
speciosa, Schrad. Fl. Germ. i. 219. t. 5. f. 3 = Stipa Calamagrostis.

ARUNDO :—
spicoidea, Schkr. ex Steud. Nom. ed. II. i. 144 = Deyeuxia scutiflora.
stricta, Gilib. Exercit. ii. 549 = Calamagrostis epigeios.
stricta, Hoff. Cat. Kansk. 7 = Deyeuxia neglecta.
stricta, Spreng. Neue Entdeck. i. 247 = Deyeuxia Nuttalliana.
stricta, Timm, in Siemss. Mekleub. Mag. ii. 235, ex Schrad. Fl. Germ. 215 = Deyeuxia stricta.
strigosa, Wahlenb. Fl. Lapp. 29. t. 2 = Deyeuxia strigosa.
subulata, J. Gay, ex Reichb. Fl. Germ. Excurs. 97 = Deyeuxia scutiflora.
sylvatica, Schrad. Fl. Germ. i. 218. t. 4. f. 7 = Deyeuxia sylvatica.
Tabacaria, Lour. Fl. Cochinch. 58 = Bambusa Tabacaria.
tecta, Bianco, Fl. Filip. ed. I. 48 = A. Karka.
tecta, Walt. Fl. Carol. 81 = Arundinaria tecta.
tenax, Vahl, Symb. Bot. ii. 25 = Ampelodesmos tenax.
tenella, Schrad. Fl. Germ. 230 = Calamagrostis tenella.
tenella, Spreng. Pugill. ii. 6 = Muehlenbergia tenuifolia.
tenuifolia, Pohr. Encyc. Suppl. iv. 704 = Muehlenbergia tenuifolia.
tolucensis, Pohr. l. c. 705 = Deyeuxia tolucensis.
triodioides, Trin. Sp. Gram. t. 351 = Schedonorus littoralis.
valdiviana. *Molina, Saggio Chil.* 279.—Chili.
Vallatoria, S. F. Gray, Nat. Arr. Brit. Pl. ii. 198 = Phragmites communis.
varia, Schrad. Fl. Germ. 216 = Deyeuxia scutiflora.
versicolor, Mill. Gard. Dict. ed. VIII. n. 8 = A. Donax.
villosa, Asch. ex Schult. Mant. iii. 387 = Calamagrostis sylvatica.
villosissima, *Forsk. ex Steud. Nom.* ed. II. i. 145.—Reg. Byzant.
viminalis, Schult. Mant. iii. 602 = Calamagrostis lanceolata.
viridescens, Steud. Syn. Pl. Gram. 192 = seq.
viridiflavescens, Poir. Encyc. vi. 271 = Deyeuxia viridiflavescens.
vulgaris, Lam. Fl. Fr. iii. 615 = Phragmites communis.
vulnerans, Gilib. Exercit. ii. 541 = Phragmites communis.
Webbiana, *Steud. Syn. Pl. Gram.* 194.—Afr. austr.
Wilhelmii, Lecheb. in Mém. Acad. Petersb. vi. (1814) 598 = Tristemon rigidum.
Zollingeri, *Busse, in Miq. Pl. Jungh.* 248.—Java.

ARUNGANA, Pers. Syn. ii. 91 (1807) = **Haronga**, Thon. (1809) (Hyperic.).
crenata, Pers. l. c. = H. crenata.
mollusca, Pers. l. c. = H. mollusca.
paniculata, Pers. l. c. = H. madagascariensis.

ARUNIA, Pers. Syn. 251 (1797), *sphalm.* = **Brunia**, Linn.

ARVERSIA, Cambess. in St. Hil. Fl. Bras. Mer. ii. 184. t. 112 (1829) = **Polycarpon**, Linn. (Caryophyll.).
depressa, Klotzsch, in Peters, Reise Mossamb. Bot. i. 140 = P. depressum.
frankenioides, Cambess. in St. Hil. Fl. Bras. Mer. ii. 184 = P. depressum.
Lorentzii, Walp. Rep. i. 268 = P. depressum.

ARVIELA, Salisb. Gen. Pl. Fragm. 135 (1866) = **Ephyranthes**, Herb. (Amaryll.).
tubispatha, Salisb. l. c. = Z. tubispatha.

ARYTERA, Blume, Rumphia, iii. 170 (1847) = **Bazonia**, DC. (Sapind.).
angustifolia, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 552.—Heb.!
arcuata, Radlk. l. c. 554.—N. Caled.
brachyphylla, Radlk. l. c. 553.—N. Guin.
Brackenridgei, Radlk. l. c. 555.—Ins. Fiji.
chartacea, Radlk. l. c. 558.—Heb.!
distylis, Radlk. l. c.—Austral.
divaricata, F. Muell. in Trans. Phil. Inst. Victoria, iii. (1859) 25 = Nephelium divaricatum.
fuscolata, F. Muell. l. c. = Nephelium fuvcolatum.
Gjellerupiana, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 67.—Sumatra.
Karung, Miq. Fl. Ind. Bat. Suppl. i. 510.—Sumatra.

ARYTERA :—
Leichhardtii, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 558.—Austral.
lepidota, Radlk. l. c. 555.—N. Caled.
littoralis, Blume, Rumphia, iii. 170.—Sumatra.
macrocarpa, Miq. Fl. Ind. Bat. Suppl. i. 510 = Trioma malaccensis.
microphylla, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 558.—Austral.
montana, Blume, Rumphia, iii. 171.—Malaya.
morotaca, Miq. Fl. Ind. Bat. Suppl. i. 510.—Sumatra.
oligolepis, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 555.—Ins. Navig.
O'Shanesiana, Radlk. l. c. 554.—Austral.
pachyphylla, Radlk. l. c.—N. Caled.
rufescens, Radlk. l. c. 559.—Ins. Philipp.
semiglauca, F. Muell. in Trans. Phil. Inst. Victoria, iii. (1859) 25 = Cupania semiglauca.
Silaka, Miq. Fl. Ind. Bat. Suppl. i. 510.—Sumatra.

ASACARA, Rafin. Neogenyt. 2 (1825) = **Gleditschia**, Linn. (Legum.).
aquatica, Rafin. l. c. = G. monosperma.

ASAEMIA, Harv. in Harv. & Sond. Fl. Cap. iii. 187 (1864), in syn. *COMPOSITAE*, Benth. & Hook. f. ii. 433.
axillaris, *Harv. l. c.*—Afr. austr.

ASAGRAEA, Baill. Adansonia, ix. (1870) 232. *LEGUMINOSAE*.
spinosa, *Baill. l. c.* 233.—Am. bor. occ.

ASAGRAEA, Lindl. Bot. Reg. t. 33 (1839) = **Schoenocaulon**, A. Gray (Liliac.).
caricoides, Ernst, in Journ. Bot. (1871) 91 = S. officinale.
caricifolia, Kunth, Enum. Pl. iv. 666 = S. officinale.
officinalis, Lindl. in Bot. Reg. (1839) t. 33 = S. officinale.
texanifolia, Kunth, Enum. Pl. iv. 700 = S. officinale.

ASAPHES, DC. Prod. ii. 90 (1825) = **Toddalia**, Juss. (Rutac.).
lanceolata, DC. l. c. = T. lanceolata.

ASAPHES, Spreng. Syst. iv. Cur. Post. 222 (1827) = **Morina**, Tourn. (Dipsac.).
nepalensis, Spreng. l. c. 232 = M. persica.

ASARCA, Poepp. ex Lindl. in Quart. Journ. R. Instit. i. (1827) 52 = **Chloraea**, Lindl. (Orchid.).
acutifora, Poepp. & Endl. Nov. Gen. et Sp. ii. 14. t. 120, A.—Chili.
acutifolia, Steud. Nom. ed. II. i. 145, sphalm. = praec.
alaris, Lindl. Gen. et Sp. Orch. 408 = C. alaris.
arauacana, Phil. in Linnaea, xxix. (1857–58) 56.—Chili.
aurantiaca, Lindl. in Hook. Journ. Bot. i. (1834) 4 = C. auranticea.
Bertori, Reichb. f. in Linnaea, xix. (1847) 377.—Chili.
bidentata, Poepp. & Endl. Nov. Gen. et Sp. ii. 15. t. 121.—Chili.
brachypetala, Phil. in Anal. Univ. Chil. (1865) ii. 392.—Chili.
chrysostachya, Phil. in Linnaea, xxix. (1857–58) 55.—Chili.
Commersonii, Hook. f. Fl. Antarct. ii. 351 = C. Commersonii.
glandulifera, Poepp. & Endl. Nov. Gen. et Sp. ii. 14. t. 120. B = C. volveria.
Kingii, Hook. f. Fl. Antarct. ii. 351.—Reg. Magell.
leucantha, Poepp. & Endl. Nov. Gen. et Sp. ii. 13. t. 119.—Chili.
maculata, Steud. Nom. ed. II. i. 145 = seq.
maculosa, Poepp. Fragm. Syn. Pl. Chil. 90 = C. guianea.
odoratissima, Poepp. & Endl. Nov. Gen. et Sp. ii. 13. t. 118.—Chili.
parviflora, Poepp. Fragm. Syn. Pl. Chil. 90.—Chili.
platypetala, Reichb. f. in Linnaea, xxii. (1849) 868.—Chili.
sinuata, Lindl. Gen. et Sp. Orch. 406 = C. longibracteata.
speciosa, Lindl. in Quart. Journ. R. Instit. i. (1827) 52 = C. Pavonii.

ASARCA :—
spectabilis, Phil. in Linnaea, xxix. (1857–58) 55 = Chlorsea spectabilis.
sulphurea, Phil. l. c. 57.—Chili.
verrucosa, A. Rich. ex C. Gay, Fl. Chil. v. 465. t. 66 = C. verrucosa.

ASARINA, Tourn. ex Mill. Gard. Dict. ed. VIII. (1768) = **Antirrhinum**, Tourn. (Scrophul.).
cordifolia, Moench, Meth. Suppl. 173 = Ant. Asarina.
erecta, Mill. Gard. Dict. ed. VIII. n. 2 = 'Penstemon,' Mitch.!—Am. bor.
Lobelii, Quer, Fl. Esp. iii. 115. t. 80 = Ant. Asarina.
procumbens, Mill. Gard. Dict. ed. VIII. n. 1 = Ant. Asarina.

ASARUM, (Tourn.) Linn. Syst. ed. I (1735). *ARISTOLOCHIACEAE*. Benth. & Hook. f. iii. 132.
HETEROTROPA, Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 314. t. 10.
HEXASTYLIS, Raf. Neogenyt. 3 (1825).
albivenium, *Regel, Gartenfl.* xiii. (1864) 295. t. 460.—Japon.
arifolium, *Michx. Fl. Bor. Am.* i. 270.—Am. bor.
Blumei, *Duch. in DC. Prod.* xv. i. 437.—Japon.
canadense, *Linn. Sp. Pl.* 442.—Am. bor.
caulescens, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 162.—Japon.
caroliniannm, Walt Fl. Carol. 143 = canadense.
caudatum, *Lindl. Bot. Reg.* sub t. 1399.—Calif.
caudigerum, *Hance, in Journ. Bot.* xix. (1881) 142.—China.
elegans, *Duch. in DC. Prod.* xv. i. 426.—Japon.
europeum, *Linn. Sp. Pl.* 442.—Europ.; As. Min.; Sibir.
furcatum, *Rafin. New Fl. Am.* ii. 90.—Am. bor.
grandiflorum, *Lodd. Cab. ex Loud. Hort. Brit.* 187.—Am. bor.
grandiflorum, *Klotzsch, in Monatsb. Acad. Berl.* (1859) 586.—Am. bor.
grandiflorum, Steud. Nom. ed. II. i. 145 = grandiflorum.
Hartwegii, *S. Wats. in Proc. Am. Acad.* x. (1875) 346.—Calif.
heterotropoides, F. Schmidt, Reisen Amurl. 171 = Siebolldii.
himalaycum, *Hook. f. & Thoms. ex Klotzsch, in Monatsb. Akad. Berl.* (1859) 585.—Reg. Himal.
Hookeri, Field. & Gerdt. Ann. i. 32 = canadense.
Hypocistis, Linn. Sp. Pl. 442 = Cytinus Hypocistis.
ibericum, Stev. ex Ledeb. Fl. Ross. iii. 553 = europaeum.
latifolium, Salisb. Prod. 244 = canadense.
Lemmoni, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 294.—Calif.
leucoclcyon, Miq. Ann. Mus. Bot. Lugd. Bat. ii: 135 = albivenium.
lucidum, Sallsb. Prod. 244 = europaeum.
maculosum, Stokes, Bot. Mat. Med. iii. 7 = virginicum.
medium, Rafin. New Fl. Am. ii. 90 = arifolium?
officinale, Moench, Meth. 292 = europaeum.
parviflorum, Regel, Ind. Sem. Hort. Petrop. (1866) 89 = elegans.
parvifolium, Rafin. New Fl. Am. ii. 90 = virginicum.
reniformo, Stokes, Bot. Mat. Med. iii. 5 = europaeum.
reniformifolium, Rafin. New Fl. Am. ii. 91 = virginicum.
Sieboldii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 134.—Japon.
Thunbergii, *A. Br. Ind. Sem. Hort. Berol.* (1861) *App.* 18.—Japon.
variegatum, *A. Br. & Bouché, l. c.* 19.—Japon.
villosum, Muhl. ex Desh. in DC. Prod. xv. i. 424 = canadense.
virginianum, Crantz, Inst. i. 284 = seq.
virginicum, *Linn. Sp. Pl.* 442.—Am. bor.
virginicum, Thunb. Fl. Jap. 180 = altivenium.
virginicum, Walt. Fl. Carol. 143 = arifolium.
viridiflorum, Regel, Ind. Sem. Hort. Petrop. (1859) 83 = variegatum.

ASCALEA, Hill, Veg. *Syst.* iv. 14 (1762) = **Cardtuse**, Linn. (Compos.).
lanceata, Hill, Hort. Kew. 60 = Cnicus lanceolatus.
natans, Hill, l. c. = Card. natans.

ASCALONICUM, *Renault*, Fl. Dép. Orne, 89 (1800) = **Allium**, Tourn. (Liliac.).
nutans, Renault, l. c. = All. Ascalonicum.

ASCALONICUM:—
Schoenoprasum, Renault, Fl. Dép. Orne, 33 (1800) = Allium Schoenoprasum.
Scorodoprasum, Renault, l. c. = All. Scorodoprasum.

ASCANICA, Crantz, Inst. ii. 313 (1766) = **Patagonula**, Linn. (Borag.).
patagonica, Crantz, l. c. = P. americana.

ASCARICIDA, Cass. in Dict. Sc. Nat. iii. Suppl. 38 (1816) = **Vernonia**, Schreb. (Compos.).
adolensis, Steets, in Peters, Reise Mossamb. Bot. 358 = V. adoënsis.
anthelmintica, Sweet, Hort. Brit. ed. I. 219 = V. anthelmintica.
Buchingeri, Steets, in Peters, Reise Mossamb. Bot. 359 = V. Buchingeri.
indica, Cass. in Dict. Sc. Nat. iii. Suppl. 38 = V. anthelmintica.
Kotschyana, Steets, in Peters, Reise Mossamb. Bot. 359 = V. Kotschyana.
mossambicensis, Steets, l. c. 358 = V. mossambicensis.
Richardi, Steets, l. c. = V. adoënsis.
triplinervia, G. Don, in Loud. Hort. Brit. 333 = Baccharis triplinervis.

ASCARIDIA, Reichb. Nom. 99 (1841) = prec.

ASCARINA, Forst. Char. Gen. 117. t. 59 (1776). *CHLORANTHACEÆ*, Benth. & Hook. f. iii. 134.
lanceolata, *Hook. f. in Journ. Linn. Soc.* i. (1856) 137.—Ins. Pacif.
lucida, *Hook. f. Fl. N. Zel.* i. 228.—N. Zel.
lurida, Solms, in DC. Prod. xvi. 1. 478 = rubricaulis.
polystachya, *Forst. Char. Gen.* 59, 117. t. 59.—Ins. Societ.
rubra, Poir. Encyc. ix. 475 = Myrica sapida.
rubricaulis, *Solms, in DC. Prod.* xvi. i. 478.—N. Caled.
serrata, Blume, Enum. Pl. Jav. i. 79 = Chloranthus brachystachyus.

ASCHAMIA, Salisb. Gen. Pl. Fragm. 184 (1866) = **Hippeastrum**, Herb. (Amaryll.).
equestris, Salisb. l. c. = H. equestre.
psittacina, Salisb. l. c. = H. psittacinum.
Reginæ, Salisb. l. c. = H. Reginæ.

ASCHENBORNIA, Schau. in Linnæa, xix. (1847) 716. *COMPOSITÆ*, Benth. & Hook. f. ii. 341.
heteropoda, *Schau. l. c.*—Mexic.

ASCHENFELDTIA, F. Muell. ex Meisn. in DC. Prod. xiv. 515 (1856) = **Pimelea**, Banks & Soland. (Thymel.).
pimeleoides, F. Muell. ex Meisn. l. c. = P. microcephalis.

ASCIADIUM, Griseb. Cat. Pl. Cub. 118 (1866). *UMBELLIFERÆ*.
coronopifolium, *Griseb. l. c.*—Cuba.

ASCIUM, Schreb. Gen. 358 (1789) = **Norantea**, Aubl. (Ternstr.).
adamantium, Steud. Nom. ed. II. i. 145 = N. adamantium.
anomalum, Spreng. Syst. iv. Cur. Post. 207 = N. anomala.
Aubletii, Spreng. Syst. ii. 599 = N. guianensis.
Berterii, Spreng. l. c. = N. Berterii.
goyasense, Steud. Nom. ed. II. i. 145 = N. goyasensis.
japurense, Steud. l. c. = N. japurensis.
Norantea, Raeusch. Nom. ed. III. 156 = N. guianensis.
paraënse, Steud. Nom. ed. II. i. 145 = N. paraënsis.
Salisii, Spreng. Syst. ii. 599 = N. cuneifolia, brasiliensis.
violaceum, Vahl, Ecl. Am. ii. 42. t. 20 = N. guianensis.

ASCLEA, Rafin. Sylva Tellur. 58 (1838) (= Lycium ? Linn.) (Solan.).
mexicana, Rafin. l. c. (= L. capsulare).—Mexic.

ASCLEPIAS, Linn. Gen. ed. I. 94 (1737). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 764.
SUBGEN. I. Rafin. New Fl. Am. i. 49 (1836).
SECT. I. Schlechter, Asclep. Afr. 1 (1895).
SUBSECT. I. Schlechter, l. c. 2.
abyssinica, Jacq. Fl. Austr. iii. 43. t. 280 = Gomphocarpus abyssinicus.
acerosa, Linn. Sp. Pl. 216 = Gomphocarpus acerosus.
adscendens, Roxb. Hort. Beng. 20; Fl. Ind. ii. 61 = Gomphocarpus adscendens.
albens, Wight, Contrib. 60 = Gomphocarpus albens.
amplexicaulis, Willd. Sp. Pl. i. 1256 = Gomphocarpus amplexicaulis.
angustissima, Linn. Sp. Pl. 214 = Gomphocarpus fruticosus.
arborescens, Linn. Sp. Pl. 215 = Gomphocarpus arborescens.
arenaria, Torr. in Bot. Mex. Bound. 162.—Am. bor.
asperula, Decne. in DC. Prod. viii. 579 = Gomphocarpus asperulus.
Baumii, Schlechter, Asclep. Afr. 2 (1895).
bicuspidata, Hook. & Arn. in Hook. Journ. Bot. i. 343 = Gomphocarpus bicuspidatus.
brachystephana, Engelm. ex Torr. in Bot. Mex. Bound. 162.—Am. bor. occ.
brevicornu, Schlechter, Asclep. Afr. 2 (1895).
Burchellii, Decne. in DC. Prod. viii. 579 = Gomphocarpus Burchellii.
cancellata, Burm. f. Fl. Cap. Prod. 7.—Afr. austr.
candida, Vell. Fl. Flum. 113; iii. t. 66.—Bras.
canescens, Willd. in Ges. Naturf. Fr. Nes. Schr. iii. (1801) 418 = Cynanchum glaucum.
canina, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 6.—Malaya.
carnosa, Blanco, Fl. Filip. ed. I. 208 = Hoya multiflora.
carnosa, Linn. f. Suppl. 170 = Hoya carnosa.
ciliata, Murr. ex Decne. in DC. Prod. viii. 572 = Xymalobium undulatum.
cinerea, Walt. Fl. Carol. 105.—Am. bor.
cirrosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 132.—Afr. trop.
citrifolia, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 295 = campestris.
citrifolia, Jacq. Coll. ii. 290 = variegata.
clausa, Jacq. Enum. Pl. Carib. 17 = Sarcostemma glaucum.
clausaefolia, *Horsf. ex Roem. & Schult. Syst.* vi. 82.—Java.
communis, Vell. Fl. Flum. 115; iii. t. 63 = Oxypetalum Banksii.
compressa, Moench, Meth. 717 = purpurascens.
connexa, Baldw. in Ell. Sketch, i. 320 = Asclepiodora viridis.
consanguinea, *Kunze, in Linnæa*, xx. (1847) 26.—Hab. ?
contorta, Forsk. Fl. Aegypt. Arab. p. cviii. n. 188 = Sarcostemma Forskohlianum.
convolvuloides, Heyne, ex Decne. in DC. Prod. viii. 572 = Holostemma Rheedianum.
convolvuloides, Willd. Sp. Pl. i. 1269 = Dæmia angolensis.
cordata, Burm. f. Fl. Ind. 72 = Pergularia minor.
cordata, Forsk. Fl. Aegypt. Arab. 49 = Dæmia cordata.
cordata, Walt. Fl. Carol. 105 = rubra ?
cordata, Vell. Fl. Flum. 116; iii. t. 56 = Roulinia fluminensis.
Cornuti, Decne. in DC. Prod. viii. 564 = syriaca.
Coulteri, *A. Gray, in Proc. Am. Acad.* xii. (1877) 71. —Mexic.
crassifolia, Hort. Par. ex Decne. in DC. Prod. viii. 572 = Gomphocarpus arborescens.
crassifolia, Linn. ex Decne. l. c. = Gomphocarpus fruticosus.
crispa, Berg. Descr. Pl. Cap. 75 = Gomphocarpus crispus.
cryptocoryna, *S. Wats. in Bot. King, Exp.* 268. t. 28.—Am. bor. occ.
cubensis, *Wender. in Bot. Zeit.* i. (1843) 830.—Cuba.
curassavica, Griseb. in Goett. Abh. xix. (1874) 189 = campestris.
curassavica, *Linn. Sp. Pl.* 215; *Willd. Sp. Pl.* i. 1266. —Am. austr.
curassavica, Lour. Fl. Cochinch. 170 = Toxocarpus Wightianus.
Curtisii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 85. —Am. bor.
Curupi, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 884.—Paraguay.
cutes, *Rafin. in Am. Monthly Mag.* (1817) 359.—Am. bor.
cynanchoides, *Humb. & Bonpl. ex Roem. & Schult. Syst.* vi. 86.—Venezuela.
Dermisi, Blanco, Fl. Filip. ed. I. 208 = Gymnema tingens.
Daemia, Forsk. Fl. Aegypt. Arab. 51 = Dæmia extensa.
dasypus, *Rafin. Att. Fourn.* 152.—Am. bor.
dasyurica, Willd. Sp. Pl. i. 1272 = Cynanchum roseum.
debilis, *Michx. Fl. Am. bor.* i. 116.—Am. bor.
decumbens, Lam. Sp. Pl. 216 = tuberosa.
densa, *Jacq. Enum. Pl. Carib.* 17.—Ind. occ.
dichotoma, Rottl. in Wall. Cat. n. 3300 = Tylophora tenuis.
Douglasii, Hook. Fl. Bor. Am. ii. 58. t. 97 = speciosa.
echinata, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iv. 35 = Marsdenia tenacissima.
echinata, Roxb. Hort. Beng. 20; Fl. Ind. ii. 44 = Dæmia extensa.

ASCLEPIAS:—
brevicornu, Schecle, in Linnæa, xxi. (1848) 755 = Asclepiodora decumbens.
Bridgesii, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 883.—Bolivia.
campestris, *Decne. in DC. Prod.* viii. 566.—Bras.
campestris, Vell. Fl. Flum. iii. t. 63 = Oxypetalum campestre.
cancellata, *Burm. f. Fl. Cap. Prod.* 7.—Afr. austr.
candida, *Vell. Fl. Flum.* 113; iii. t. 66.—Bras.
canescens, Willd. in Ges. Naturf. Fr. Nes. Schr. iii. (1801) 418 = Cynanchum glaucum.
canina, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Malaya.
carnosa, Blanco, Fl. Filip. ed. I. 208 = Hoya multiflora.
carnosa, Linn. f. Suppl. 170 = Hoya carnosa.
ciliata, Murr. ex Decne. in DC. Prod. viii. 572 = Xymalobium undulatum.
cinerea, *Walt. Fl. Carol.* 105.—Am. bor.
cirrosa, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 132.—Afr. trop.
citrifolia, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 295 = campestris.
citrifolia, Jacq. Coll. ii. 290 = variegata.
clausa, Jacq. Enum. Pl. Carib. 17 = Sarcostemma glaucum.
clausaefolia, *Horsf. ex Roem. & Schult. Syst.* vi. 82.—Java.
communis, Vell. Fl. Flum. 115; iii. t. 63 = Oxypetalum Banksii.
compressa, Moench, Meth. 717 = purpurascens.
connexa, Baldw. in Ell. Sketch, i. 320 = Asclepiodora viridis.
consanguinea, *Kunze, in Linnæa*, xx. (1847) 26.—Hab. ?
contorta, Forsk. Fl. Aegypt. Arab. p. cviii. n. 188 = Sarcostemma Forskohlianum.
convolvuloides, Heyne, ex Decne. in DC. Prod. viii. 572 = Holostemma Rheedianum.
convolvuloides, Willd. Sp. Pl. i. 1269 = Dæmia angolensis.
cordata, Burm. f. Fl. Ind. 72 = Pergularia minor.
cordata, Forsk. Fl. Aegypt. Arab. 49 = Dæmia cordata.
cordata, Walt. Fl. Carol. 105 = rubra ?
cordata, Vell. Fl. Flum. 116; iii. t. 56 = Roulinia fluminensis.
Cornuti, Decne. in DC. Prod. viii. 564 = syriaca.
Coulteri, *A. Gray, in Proc. Am. Acad.* xii. (1877) 71. —Mexic.
crassifolia, Hort. Par. ex Decne. in DC. Prod. viii. 572 = Gomphocarpus arborescens.
crassifolia, Linn. ex Decne. l. c. = Gomphocarpus fruticosus.
crispa, Berg. Descr. Pl. Cap. 75 = Gomphocarpus crispus.
cryptocoryna, *S. Wats. in Bot. King, Exp.* 268. t. 28.—Am. bor. occ.
cubensis, *Wender. in Bot. Zeit.* i. (1843) 830.—Cuba.
curassavica, Griseb. in Goett. Abh. xix. (1874) 189 = campestris.
curassavica, *Linn. Sp. Pl.* 215; *Willd. Sp. Pl.* i. 1266. —Am. austr.
curassavica, Lour. Fl. Cochinch. 170 = Toxocarpus Wightianus.
Curtisii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 85. —Am. bor.
Curupi, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 884.—Paraguay.
cutes, *Rafin. in Am. Monthly Mag.* (1817) 359.—Am. bor.
cynanchoides, *Humb. & Bonpl. ex Roem. & Schult. Syst.* vi. 86.—Venezuela.
Dermisi, Blanco, Fl. Filip. ed. I. 208 = Gymnema tingens.
Daemia, Forsk. Fl. Aegypt. Arab. 51 = Dæmia extensa.
dasypus, *Rafin. Att. Fourn.* 152.—Am. bor.
dasyurica, Willd. Sp. Pl. i. 1272 = Cynanchum roseum.
debilis, *Michx. Fl. Am. bor.* i. 116.—Am. bor.
decumbens, Lam. Sp. Pl. 216 = tuberosa.
densa, *Jacq. Enum. Pl. Carib.* 17.—Ind. occ.
dichotoma, Rottl. in Wall. Cat. n. 3300 = Tylophora tenuis.
Douglasii, Hook. Fl. Bor. Am. ii. 58. t. 97 = speciosa.
echinata, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iv. 35 = Marsdenia tenacissima.
echinata, Roxb. Hort. Beng. 20; Fl. Ind. ii. 44 = Dæmia extensa.

ASCLEPIAS:—
echinata, Decne. in DC. Prod. viii. 544 = Daemia angolensis.
ecornuta, Kellogg, ex A. Gray, Syn. Fl N. Am. ii. 1.100 = Gomphocarpus cordifolius.
elata, Benth. Pl. Hartw. 290 = glaucescens.
elegans, *Vell. Fl. Flum.* 116; Ill. t. 55.—Bras.
eriocarpa, *Benth. Pl. Hartw.* 323.—Calif.
eriocarpa, Torr. in Pacif. Rail. Rep. iv. 128 = vestita.
erosa, *Torr. in Bot. Mex. Bound.* 103.—Calif.
euphorbioefolia, *Engelm. in Proc. Am. Acad.* xvi. (1881) 104.—Mexic.
exaltata, Linn. Amoen. Acad. iii. 404 = syriaca.
exaltata, Link, ex Spreng. Syst. i. 848 = Asclepiodora viridis.
exaltata, Muhl. ex Bigel. Fl. Bost. 102 = phytolaccoides.
fascicularis, Decne. in DC. Prod. viii. 509 = mexicana.
fasciculata, Hemsl. Biol. Centr. Am. Bot. ii. 324, sphalm. = prasc.
Feayi, *Chapm. ex A. Gray, in Proc. Am. Acad.* xii. (1876) 72.—Am. bor. trop.
ferruginea, *Humb. & Bonpl. ex Roem. & Schult. Syst.* vi. 88.—Venezuela.
filiformis, Jacq. Enum. Pl. Carib. 17 = Sarcostemma Jacquini.
filiformis, Linn. f. Suppl. 169 (= *Lagarinthus tenuis*). —Ind. occ.; Afr. austr.
floridana, Lam. Encyc. i. 284 = Gomphocarpus longifolia.
foetida, Cav. Ic: ii. 45. t. 158 = Roulinia Jacquini.
Forskohlei, *Roem. & Schult. Syst.* vi. 85.—Arab.
fragrans, *Decne. in DC. Prod.* viii. 571.—Guatemala.
fragrans, *Raf. in Med. Repos. N. York*, v. (1808) 354. —Am. bor.
Frémontii, *Torr. in Pacif. Rail. Rep.* vi. 87.—Calif.
fruticosa, Linn. Sp. Pl. 216 = Gomphocarpus fruticosus.
fusca, Lour. Fl. Cochinch. 170 = Vincetoxicum medium.
fuscata, Willd. Enum. Hort. Berol. Suppl. 18 = pruse.
Galeottii, *Fourn. in Bull. Soc. Bot. Fr.* xiv. (1867) 250. —Mexic.
galioides, H. B. & K. Nov. Gen. et Sp. iii. 148 = verticillata.
geminata, Roxb. Hort. Beng. 20; Fl. Ind. ii. 45 = Gymnema sylvestre.
gigantea, Linn. Sp. Pl. 214 = Calotropis procera.
gigantea, Willd. Sp. Pl. i. 1264 = Calotropis gigantea.
glabra, Forsk. Fl. Aegypt. Arab. 51 = Daemia glabra.
glabra, Mill. Gard. Dict. ed. VIII. a. 12 = Gomphocarpus fruticosus.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* iii. 290. t. 287.—N. Mexic.
globosa, Stokes, Bot. Mat. Med. ii. 4 = syriaca.
gonocarpos, J. F. Gmel. Syst. 446 = Gonolobus carolinensis.
graminea, *Lam. Encyc.* i. 283.—Ind. or.
grandiflora, Linn. f. Suppl. 170 = Gomphocarpus grandiflorus.
grandiflora, *Fourn. in Ann. Sc. Nat. Sér.* VI. xiv. (1882) 379.—Mexic.
grandiflola, Bertol. in Mem. Acc. Sc. Bolog. iii. (1851) 189 (Misc. Bot. xii. 47) = syriaca.
Greenei, *Nutt. ex Lond. Hort. Brit. Suppl.* iii. 493. —Am. bor.
Hallii, *A. Gray, in Proc. Am. Acad.* xii. (1877) 69.— Colorado.
hastata, Bunge, in Mém. Sav. Etr. Acad. Péterab. ii. (1835) 117 = Cynanchum Bungei.
herbacea, Roxb. Hort. Beng. 21; Fl. Ind. ii. 50 = Calotropis Acia.
hispida, Vell. Fl. Flum. 115; Ill. t. 52 = Ditassa Gulliemiana.
humistrata, Walt. Fl. Carol. 105 = amplexicaulis.
hybrida, Michx. Fl. Am. Bor. i. 115 = variegata.
illinoënsis, Michx. ex Steud. Nom. ed. II. i. 146 = syriaca.
incarnata, *Linn. Sp. Pl.* 215.—Am. bor.
incarnata, Walt. Fl. Carol. 106 = Gomphocarpus longifolius.
involucrata, *Engelm. ex Torr. in Bot. Mex. Bound.* 103.—N. Mexic.
Jacquiniana, Spreng. Syst. iv. Cur. Post. 341 = Sarcostemma Jacquini.
Jamesii, *Torr. Bot. Mex. Bound.* 103.—Am. bor.
javanica, Burm. ex Decne. in DC. Prod. viii. 626 = Lepadenia reticulata.

ASCLEPIAS:—
Klaprothiana, *Roem. & Schult. Syst.* vi. 82.—Am. bor.
lactiflora, Linn. Sp. Pl. 216 = Gymnema lactiferum.
lanceolata, Ives, in Am. Journ. Sc. i. (1819) 252 = Gomphocarpus viridiflorus.
lanceolata, Walt. Fl. Carol. 105 = paupercula.
lancifolia, Steud. Nom. ed. I. 77 = rubra.
lanifiora, Forsk. Fl. Aegypt. Arab. 51 = Kanahia laniflora.
Langsdorffii, *Fourn. in Mart. Fl. Bras.* vi. iv. 208.— Bras.
lanuginosa, *H. B. & K. Nov. Gen. et Sp.* iii. 192.— Mexic.
lanuginosa, Nutt. Gen. Am. i. 168 (= *Acerates lanuginosa*).—Am. bor.
latifolia, *Raf. Atl. Journ.* 146.—Am. bor.
laurifolia, Michx. Fl. Bor. Am. i. 117 = rubra.
laurifolia, Roxb. Hort. Beng. 20; Fl. Ind. ii. 49 = Gymnema lucrifolius.
laxiflora, *Decne. in DC. Prod.* viii. 571.—Mexic.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 85.—Arizona.
leucophylla, Engelm. in Am. Natural. ix. (1875) 348 = cross.
Linaria, *Cav. Ic.* i. 42. t. 57.—Mexic.
Lindheimeri, Engelm. & A. Gray, Pl. Lindh. 250 = longicornu.
Lindleyi, Gill. ex Steud. Nom. ed. II. i. 145, 146 = variegata.
linearifolia, Pav. ex Decne. in DC. Prod. viii. 589 = Sarcostemma Domheyanum.
linearis, *Scheele, in Linnaea*, xxi. (1848) 758 = verticillata.
linifolia, H. B. & K. Nov. Gen. et Sp. iii. 190 = angustifolia, virgata ?
linifolia, *Lag. ex Spreng. Syst.* i. 847.—Mexic.
longicornu, *Kellogg, in Proc. Calif. Acad.* i. (1855 ?) 65.—Calif.
longicornu, *Benth. in Pl. Hartw.* 24.—Calif.
longifolia, Jacq. Enum. Pl. Carib. 17 (= *Schubertia longifolia*).—Ind. occ.
longifolia, *Michx. Fl. Bor. Am.* i. 116.—Am. bor.
longipetala, Scheele, in Linnaea, xxi. (1848) 757 = Asclepiodora viridis.
longistigma, Roxb. Hort. Beng. 85; Fl. Ind. ii. 46 = Toxocarpus Roxburghii.
lutea, *Raf. in Med. Repos. N. York*, v. (1808) 361.— Am. bor.
lutea, Mill. Gard. Dict. ed. VIII. a. 8 = Vincetoxicum intermedium.
macrantha, *Hochst. ex Decne. in DC. Prod.* viii. 558.— Abyss.
macrophylla, *Humb. & Bonpl. in Roem. & Schult. Syst.* vi. 86.—Cumana.
macrophylla, Nutt. in Journ. Acad. Philad. Ser. II. i. (1847) 180 = mexicana.
macrota, *Torr. Bot. Mex. Bound.* 104. t. 45.—N. Mexic.
margaritacea, *Hoffmgg. ex Roem. & Schult. Syst.* vi. 86. —Bras.
margaritata, Decne. in DC. Prod. viii. 566 = candida.
maritima, Jacq. Enum. Pl. Carib. 17 = Ibatia maritima.
maritima, *Pall. Reise*, iii. 557.—Rossia austr.
maritima, Rafin. ex Decne. in DC. Prod. viii. 567 = incarnata.
Meadii, *Torr. ex A. Gray, Man. Bot. U. St.* ed. II. Add. 704.—Am. bor.
media, Gilib. ex Steud. Nom. ed. II. i. 146 = Vincetoxicum medium.
megalotis, *Rafin. New Fl. Am.* iv. 61.—Am. bor.
melanta, *Decne. in DC. Prod.* viii. 570.—Mexic.
melleodora, St. Hil. Fl. Rem. Bras. 227 = nervosa.
mexicana, *Cav. Ic.* i. 42. t. 58 = Calif.; Mexic.
Michauxii, *Decne. in DC. Prod.* viii. 569.—Am. bor.
micrantha, *Roxb. Hort. Beng.* 21; *Fl. Ind.* ii. 50.— Ind. or.
microphylla, Heyne, ex Roth, Nov. Sp. Pl. 177 = acq.?
microphylla, Roxb. Hort. Beng. 85; Fl. Ind. ii. 55 = Pentatropis microphylla.
monantha, Vell. Fl. Flum. 115; Ill. t. 54 = Oxypetalum monanthum.
monilena, Roxb. Hort. Beng. 85; Fl. Ind. ii. 45 = Gymnema tingens.
montevidensis, *Spreng. ex Fourn. in Mart. Fl. Bras.* vi. iv. 202.—Reg. Argent.
mucronata, Thunb. Prod. Pl. Cap. 47 (= *Metaplocis mucronata*).—Afr. austr.

ASCLEPIAS:—
multicaulis, Vell. Fl. Flum. 117; Ill. t. 60 = Oxypetalum campestre.
multinervia, *Fourn. in Ann. Sc. Nat. Sér.* VI. xiv. (1882) 388.—Paraguay.
muricata, Schum. & Thonn. Beskr. Guin. Pl. 158 = Daemia angolensis.
neglecta, *Hemsl. Biol. Centr. Am. Bot.* ii. 325.— Mexic.
nervosa, *Decne. in DC. Prod.* viii. 566.—Bras.
nigra, Linn. Sp. Pl. 216 = Vincetoxicum nigrum.
nitida, Vell. Fl. Flum. 116; Ill. t. 59 = Oxypetalum umbellatum.
nivalis, J. F. Gmel. Syst. 444 = acq.
nivea, Forsk. Fl. Aegypt. Arab. 51 = Forskohlei.
nivea, *Linn. Sp. Pl.* 215.—Ind. occ.
nivea, Sims, Bot. Mag. t. 1181 = phytolaccoides.
nuda, *Thunb. & Schum. Beskr. Guin. Pl.* 155.—Afr. trop.
nummularia, *Torr. Bot. Mex. Bound.* 103. t. 45.— Am. bor.
nutans, Muhl. ex Steud. Nom. ed. II. i. 146 (= *Acerates viridiflorus*).—Am. bor.
Nuttalliana, A. Gray, Man. Bot. U. St. ed. II. 353, 704 = ovalifolia.
Nuttalliana, Torr. in Ann. Lyc. N. York, ii. (1828) 218 (= *Acerates lanuginosa*).—Am. bor.
nyctaginifolia, *A. Gray, in Proc. Am. Acad.* xii. (1877) 69.—Calif.
obovata, *Ell. Sketch*, 291.—Am. bor.
obtusifolia, Hort. Prag. ex Kunze, in Linnaea, xx. (1847) 27 = consanguinea.
obtusifolia, *Michx. Fl. Bor. Am.* i. 115.—Am. bor.
odoratissima, Roxb. Hort. Beng. 20; Fl. Ind. ii. 47 = Pergularia minor.
oenotheroides, Cham. & Schlecht. in Linnaea, v. (1830) 128 = longicornu.
otaroides, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 378.—Mexic.
ovalifolia, *Decne. in DC. Prod.* viii. 567.—Am. bor.
ovata, Aust. ex Steud. Nom. ed. II. i. 146 = Vincetoxicum officinale.
ovata, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1842) 268.—Mexic.
pallida, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 385.—Paraguay.
pallida, Roxb. Hort. Beng. 20; Fl. Ind. ii. 48 = Pergularia pallida.
pauciculatum, *Bunge, Enum. Pl. Chin. Bor.* 48.— China.
paradisica, Roxb. Hort. Beng. 20; Fl. Ind. ii. 42 = Hoya paradisica.
parviflora, Ait. Hort. Kew. i. 307 = perennis.
parviflora, Leconte, ex Decne. in DC. Prod. viii. 570 = verticillata.
parviflora, Roem. & Schult. Syst. vi. 89, sphalm. = parvifolia.
parvifolia, Decne. in DC. Prod. viii. 568 = perennis.
parvifolia, *Roem. & Schult. Syst.* iv. 801, ind.—Am. austr.
patula, Aust. ex Decne. in DC. Prod. viii. 536 = Calotropis procera.
pauperoula, *Michx. Fl. Bor. Am.* i. 116.—Am. bor.
pedicellata, Walt. Fl. Carol. 106 = Podostigma pubescens.
pellucida, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 381.—Mexic.
pendula, Roxb. Hort. Beng. 85; Fl. Ind. ii. 56 = Hoya pendula.
peregrina, Blanco, Fl. Filip. ed. I. 207 = Sarcolobus carinatus.
perennis, *Walt. Fl. Carol.* 107.—Am. bor.
periplocoefolia, Nutt. Gen. Am. i. 167 = rubra.
phytolaccoides, *Lyon. ex Pursh, Fl. Am. Sept.* i. 260. —Am. bor.
pilosa, Vell. Fl. Flum. 117; Ill. t. 62 = Oxypetalum foliosum.
pinifolia, Greene, in Torrey Bot. Club, viii. (1881) 5 = A. Linaria.
planifiora, Jacq. Enum. Pl. Carib. 17 = Gonolobus planiflorus.
plicata, *Heyne, ex Roth, Nov. Pl. Sp.* 178.—Ind. or.
polyantocyca, Walt. Fl. Carol. 107 = rubra ?
prasinula, *Benth. Pl. Hartw.* 23.—Mexic.
Primaps, *Bartl. Ind. Sem. Hort. Gott.* (1839); *ex Steetz, in Linnaea*, xx. (1847) 27.—Am. bor.
prostrata, Ait. Hort. Kew. i. 308 = Calotropis procera.
pseudurbana, *Raffn. Fl. Ludov.* 91.—Am. bor.

ASCLEPIAS:—
pseudosarsa, Roxb. Hort. Beng. 20; Fl. Ind. ii. 39 = Hemidesmus indicus.
pubescens, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 104.—Mexic.
pubescens, Linn. Mant. 215 = Gomphocarpus arborescens.
pubescens, Moench, Meth. 716 = syriaca.
pubigera, Dum. Fl. Belg. 52 = syriaca.
pulchella, Roxb. Hort. Beng. 21; Fl. Ind. ii. 54 = Raphistemma pulchellum.
pulchella, Salisb. Prod. 150 = perennis.
pulchra, Ehrh. ex Willd. Enum. Hort. Berol. 276 = incarnata.
pulchra, Vell. Fl. Flum. 116; ill. t. 58 = Oxypetalum pulchrum.
purpurascens, Georgi, Beschr. Russ. Reich. ill. iv. 818 = Vincetoxicum purpurascens.
purpurascens, *Linn. Sp. Pl.* 214.—Am. bor.
purpurascens, Walt. Fl. Carol. 105 = obtusifolia.
purpurea, Pall. Reise, ill. 260 (= *Cynoctonum roseum*). —Russia.
quadrifolia, *Jacq. Obs. Bot.* ii. 8.—Am. bor.
quinquedentata, *A. Gray, in Proc. Am. Acad.* xii. (1877) 71.—N. Mexic.
quinquedora, *Vell. Fl. Flum.* 118; ill. t. 68.—Bras.
racemosa, Jacq. Enum. Pl. Carib. 17 = Roulinia Jacquini.
racemosa, Roxb. Hort. Beng. 20; Fl. Ind. ii. 32 = Cosmostigma racemosum.
radiana, Forsk. Fl. Aegypt. Arab. 49 = Steinhelilia radiana.
ramosa, *Pourr. in Ann. Sc. Nat. Sér.* VI. xiv. (1882) 373.—Bolivia.
repanda, *Forsk. ex Steud. Nom.* ed. I. 76; ed. II. i. 146 = Gomphocarpus crispus.
revoluta, *Rafin. Fl. Ludov.* 51.—Am. bor.
rosea, *H. B. & K. Nov. Gen. et Sp.* iii. 189.—Mexic.
rosea, Roxb. Hort. Beng. 20; Fl. Ind. ii. 40 = Oxystelma esculentum.
rotundifolia, *Mill. Gard. Dict.* ed. VIII. n. 15.—Afr. austr.
rotundifolia, *Rafin. Atl. Journ.* 152.—Hab. ?
rubra, *Linn. Sp. Pl.* 217.—Am. bor.
rubra, Sievers, in Pall. N. Nord. Beitr. vii. 329 = Apocynum Venetum.
rubricaulis, *H. B. & K. Nov. Gen. et Sp.* iii. 148.—Mexic.
salicifolia, *Ledd. Bot. Cab.* t. 272.—Mexic.
scandens, Beauv. Fl. Owar. i. 92. t. 56 = Daemia saxolensis.
scandens, Mill. Gard. Dict. ed. VIII. n. 19 = Sarcostemma Swartzianum.
Schaffneri, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 108.—Mexic.
Selloana, *Pourr. in Mort. Fl. Bras.* vi. iv. 209.—Bras.
Serpentaria, *Rafin. Fl. Ludov.* 52.—Am. bor.
setosa, *Benth. Pl. Hartw.* 94.—Mexic.
setosa, Forsk. Fl. Aegypt. Arab. 51 = Gomphocarpus setosus.
sibirica, *Linn. Sp. Pl.* 217.—Sibir.
sibirica, Marr. in Comm. Gotting. (1779) 23 = Vincetoxicum sibiricum.
sinaïtica, *Hemsl. Bird. Centr. Am. Bot.* ii. 296.—Mexic.
singularis, *Vell. Fl. Flum.* 117; ill. t. 61.—Bras.
sinuosa, *Burm. f. Fl. Cap. Prod.* 7.—Afr. austr.
speciosa, *Torr. in Ann. Lyc. N. York*, ii. (1828) 218. —Am. bor. occ.
spinosa, *Vell. Fl. Flum.* 118; ill. t. 67.—Bras.
spiniflora, Beauv. Fl. Owar. i. 92. t. 56 = Daemia saxolensis.
spinosa, Forsk. Fl. Aegypt. Arab. 49 = Pentatropis spinosa.
stellata, Burm. ex Decne. in DC. Prod. viii. 573 = Hoya multiflora.
stenophylla, *A. Gray, in Proc. Am. Acad.* xii. (1876) 72.—Am. bor.
stipitacea, Forsk. Fl. Aegypt. Arab. 51 = Sarcostemma stipitaceum.
stomatosa, Leconte, ex Decne. in DC. Prod. viii. 569 = Matelea.
suberosa, Roxb. Hort. Beng. 35; Fl. Ind. ii. 36 = Leptadenia reticulata.
subulata, *Decne. in DC. Prod.* viii. 571.—Calif.
subulata, Forsk. Fl. Aegypt. Arab. p. xviii. n. 198. t. 7 = Caralluma subulata.
Sullivantii, *Engelm. ex A. Gray, Mem. Bot.* ed. I. 366. —Ohio, bor.
Swallenantii, Torr. Bot. Mex. Bound. 169 = glaucescens.
ASCLEPIAS:—
Suzuela, Roxb. Hort. Beng. 20; Fl. Ind. ii. 31 = Hoya imperialis.
syriaca, Blanco, Fl. Filip. ed. I. 204 = curassavica.
syriaca, *Linn. Sp. Pl.* ed. II. 214.—Am. bor.
tenacissima, Roxb. Pl. Corom. iii. 35. t. 240 = Marsdenia tenacissima.
tenuis, Herb. Heyne, ex Hook. f. Fl. Brit. Ind. iv. 41 = Tylophora cappariodifolia.
tenuissima, Roxb. Hort. Beng. 20; Fl. Ind. ii. 41 = Tylophora tenuissima.
tetropetala, Dennst. Schluss. Hort. Malab. 35 = Pentatropis microphylla.
tinctoria, Roxb. Hort. Beng. 20; Fl. Ind. ii. 48 = Marsdenia tinctoria.
tingens, Buch.-Ham. ex Roxb. Hort. Beng. 21; Fl. Ind. ii. 53 = Gymnema tingens.
tomentosa, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iv. 36 = Marsdenia tenacissima.
tomentosa, *Ell. Sketch*, i. 320.—Am. bor.
tomentosa, Thonn. ex Decne. in DC. Prod. viii. 638 = Leptadenia lanciflola.
tuberata, *Salisb. Prod.* 150 = Vincetoxicum officinale.
truncata, *Willd. Enum. Hort. Berol. Suppl.* 18.—Hab.?
truncata, Roxb. ex Decne. in DC. Prod. viii. 529 = Cynanchum alatum.
tuberosa, *Linn. Sp. Pl.* 217.—Am. bor.
tuberosa, Hort. ex Kunze, in Linnaea, xx. (1847) 27.—Hab.?
tuberosa, Roxb. Fl. Ind. ii. 38 = Leptadenia reticulata.
turicata, Roxb. Hort. Beng. 20; i. c. 35 = Cynanchum paniculiflorum.
tunicata, Hort. Calc. ex Hook. f. Fl. Brit. Ind. iv. 45 = Tylophora asthmatica.
umbellata, *Vell. Fl. Flum.* 118; ill. t. 66.—Bras.
uncialis, *Greene, in Coulter, Bot. Gas.* v. (1880) 64. —Am. bor. occ.
undulata, Jacq. Enum. Pl. Carib. 17 = Xysmalobium undulatum.
undulata, Linn. Sp. Pl. 214 = Pachycarpus crispus.
Vanilla, Rafin. in Am. Monthly Mag. iv. (1818) 39 = quadrifolia.
variegata, *Linn. Sp. Pl.* 215.—Am. bor.
verrucosa, Salisb. Prod. 150 = incarnata.
verticillata, *Linn. Sp. Pl.* 217.—Am. bor.
vestita, *Hook. & Arn. in Bot. Beech. Voy.* 363.—Calif.
vicinalis, Mart. Fl. Bras. vi. iv. 235 = Sarcostemma barbatum.
villosa, Balb. in Mem. Acc. Torin. vii. (1802–3) 366 = Leptadenia Balbisii.
villosa, Mill. Gard. Dict. ed. VIII. n. 14 = Gomphocarpus fruticosus.
villosa, *Rafin. in Med. Repos. N. York*, v. (1808) 353, nomen.—Am. bor.
viminalis, Steud. Nom. ed. II. i. 146 = Sarcostemma viminale.
viminalis, Sw. Prod. Veg. Ind. Occ. 58 = Philibertia viminalis.
Vincetoxicum, Linn. Sp. Pl. 216 = Vincetoxicum officinale.
virgata, Balb. in Cat. Hort. Taur. (1813) 14 = angustifolia.
viridiflora, Rafin. in Med. Repos. N. York, v. (1808) 360; Pursh, Fl. Am. Sept. i. 181 = Gomphocarpus viridiflorus.
viridiflora, Roxb. ex Decne. in DC. Prod. viii. 573 = Hoya viridiflora.
viridis, Walt. Fl. Carol. 107 = Asclepiodora viridis.
viridula, *Chapm. Fl. S. U. St.* 363.—Am. bor. trop.
Virletii, *Pourr. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 378.—Mexic.
volubilis, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iv. 63 = Leptadenia reticulata.
volubilis, Linn. f. Suppl. 170 = Dregea volubilis.
volubilis, Domb. ex Decne. in DC. Prod. viii. 539 = Sarcostemma Dombeyanum.
vomitoria, Koen. ex Hook. f. Fl. Brit. Ind. iv. 45 = Tylophora asthmatica.
Weddellii, *Pourr. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 381.—Bolivia.
Wrightii, Greene, in Proc. Am. Acad. xvi. (1881) 102 = syntrophilis.
xylacarcha, Benn. ex Decne. in DC. Prod. viii. 638 = Leptadenia reticulata.
ASCLEPIODORA, A. Gray, in Proc. Am. Acad. xii. (1876) 66. *ASCLEPIADEAE.*
ANANTHERIX, Nutt. Gen. Am. i. 169 (1818).
ANTHANOTIS, Rafin. Fl. Ludov. 52, 149 (1817), nomen prius.
asperula, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 369.—Mexic.
circinalis, Fourn. l. c. = Asclepias brachystephana.
decumbens, *A. Gray, in Proc. Am. Acad.* xii. (1876) 67.—Am. bor.
gomphocarpoides, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 369.—Mexic.
viridula, *A. Gray, in Proc. Am. Acad.* xii. (1876) 66.—Am. bor.

ASCOCHILUS, Blume, Fl. Jav. Pracf. p. vi. (1828) = **Geodorum**, Jack (Orchid.).

ASCOLEPIS, Nees, ex Steud. Syn. Pl. Cyp. 105 (1855). *CYPERACEAE*, Benth. & Hook. f. iii. 1054.
ANTROLEPIS, Welw. Apont. Photo-Geograph. in Annaes do Consel. Ultramarin. (1858) 578.
PLATYLEPIS, Kunth, Enum. Pl. ii. 289 (1835).
PTERACHNE, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 62, in nota (1842).
PTEROGYNE, Schrad. ex Nees, in Mart. l. c. (1842).
anthemiflora, *Welw. in Trans. Linn. Soc.* xxvii. (1868) 77.—Angola.
capensis, *Kunth. in Trans. Linn. Soc.* Ser. II. Bot. ii. (1884) 164.—Afr. austr. et trop.
cista, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 79.—Angola.
ericoscalides, *Nees, ex Steud. Syn. Pl. Cyp.* 105.—Abyss.
kyllingioides, Steud. l. c. = Lipocarpha microcephala.
prolata, *Welw. in Trans. Linn. Soc.* xxvii. (1868) 75.—Afr. trop.
pusilla, *Kidl. in Trans. Linn. Soc.* Ser. II. Bot. ii. (1884) 164.—Afr. occ.
speciosa, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 78. —Angola.
tenuior, Steud. Syn. Pl. Cyp. 105 = Scirpus squarrosus

ASCRA, Schott, in Spreng. Syst. iv. Cur. Post. 407 (1827) = **Banara**, Aubl. (Samyd.).
brasiliensis, Schott, l. c. = B. brasiliensis.

ASCYRUM, Linn. Gen. ed. I. 281 (1737). *HYPERICINEAE*, Benth. & Hook. f. i. 164.
HYPERICOIDES, Adans. Fam. ii. 448 (1763).
ISOPHYLLUM, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367.
amplericaule, *Michx. Fl. Bor. Am.* ii. 77.—Am. bor.
calycinum, Poir. Tabl. Encyc. iii. 199 = Hypericum calycinum.
coriaceum, Moench, Meth. 180 = Hypericum calycinum.
crucistatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69 = A. Crux-Andreae.
Crux-Andreae, Linn. Sp. Pl. 787 = Hypericum mutilum.
Crux-Andreae, *Linn. Sp. Pl.* ed. II. 1107.—Am. bor.; ind. occ.
cubense, *Griseb. Cat. Pl. Cub.* 40.—Cuba.
filicaule, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 262.—Reg. Himal.
grandiflorum, Moench, Meth. Suppl. 42 = Hypericum balearicum.
grandiflorum, Rafin. Fl. Ludov. 87 = stans.
helianthemifolium, Spach, Hist. Vég. Phan. v. 480 = A. Crux-Andreae.
humifusum, Labill. Nov. Holl. Pl. ii. 33. t. 175 = Hypericum japonicum.
hypericifolium, Sw. ex Rafin. Sylva Tellur. 116 = A. Hypericoides.
Hypericoides, *Linn. Sp. Pl.* 788 & ed. II. 1108 pro max. parte.—Ind. occ.
Hypericoides, Marsh. Arbust. 14 = stans.
grandiflorum, Rafin. Fl. Ludov. 87 = stans.
lancastrum, Labill. Nov. Holl. Pl. ii. 32. t. 174 = Hypericum gramineum.
linifolium, Spach, Hist. Vég. Phan. v. 459 = A. Crux-Andreae.
Michauxii, Spach, l. c. 460 = A. Crux-Andreae.
microsepalum, Torr. & Gray, Fl. N. Am. i. 157 = Hypericum microsepalum.
monogynum, Moench, Meth. Suppl. 42 = Hypericum monogynum.
montanum, Rafin. Sylva Tellur. 116 = A. Hypericoides.

ASCYRUM:—
multicaule, Michx. ex Willd. Sp. Pl. iii. 1472 = A. Crux-Andreae.
ramosissimum, Banks, ex Steud. Nom. ed. I. 77 = pumilum.
oblongifolium, Spach, Hist. Vég. Phan. v. 461 = A. Crux-Andreae.
pauciflorum, Nutt. Gen. Am. ii. 15 = pumilum.
Plumierii, Bertol. in Mem. Acc. Sc. Bologna. iv. (1853) 77 (Misc. Bot. xiii. 19) = A. Crux-Andreae.
pumilum, *Michx. Fl. Bor. Am.* ii. 77.—Am. bor.
sibiricum, Poir. Tabl. Encyc. iii. 900 = Hypericum Ascyrum.
simplex, Zeph. ex Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 389 = stans.
spathulatum, Spach, Hist. Vég. Phan. v. 462 = A. Crux-Andreae.
stans, *Michx. Fl. Bor. Am.* ii. 77.—Am. bor.
tetragonum, Moench, Meth. 130 = Hypericum Ascyron.
villosum, Linn. Sp. Pl. 788 = Hypericum setosum.

ASCYUM, Vahl, ex DC. Prod. i. 566 (1824) = ASCIUM, Schreb. = **Norantea**, Aubl. (Ternstr.).

ASEMEIA, Rafn. New Fl. Am. iv. 88 (1836) = Polygala, Linn.
alba, Rafn. l. c. = P. alba.
carnea, Rafn. l. c.—Am. bor.
leptopis, Rafn. l. c.—Am. bor.
rosea, Rafn. l. c. = P. Senega.

ASEMNANTHA, Hook. f. in Benth. & Hook. f. Gen. ii. 106 (1873). *RUBIACEÆ*, Benth. & Hook. f. ii. 106.
pubescens, *Hook. f. l. c.* 107.—Mexic.

ASEPHANANTHES, Bory, ex DC. Prod. iii. 328 (1828) = **Passiflora**, Linn.

ASICARIA, Neck. Elem. ii. 211 (1790) = **Polygonum**, Tourn.

ASIMIA, H. B. & K. Nov. Gen. et Sp. v. 61 (1821) = **Asimina**, Adans. (Anonac.).

ASIMINA, Adans. Fam. ii. 365 (1763). *ANONACEÆ*, Benth. & Hook. f. i. 24.
ORCHIDOCARPUM, Michx. Fl. Bor. Am. i. 329 (1808).
aurantiaca, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 25 = Orophea aurantiaca.
Blahni, *Griseb. Cat. Pl. Cub.* 3.—Cuba.
campechiana, *H. B. & K. Nov. Gen. et Sp.* v. 61.—Mexic.
campeniflora, Spach, Hist. Vég. Phan. vii. 538 = triloba.
conoidea, Spach, l. c. 530 = triloba?
cuneata, Shuttlw. ex Chapm. Fl. S. U. St. 15 = grandiflora.
glabra, Hort. ex C. Koch, Dendrol. i. 384 = triloba.
grandiflora, *Dun. Monog. Anon.* 84. t. 11.—Am. bor.
grandiflora, Hort. ex Spach, Hist. Vég. Phan. v. 530 = *conoidea*.
lanceolata, Hook. f. in Journ. Linn. Soc. i. (1857) 128, *err. typ.* = Ascarina lanceolata.
neglecta, *Griseb. Cat. Pl. Cub.* 3.—Cuba.
parviflora, *Dun. Monog. Anon.* 82. t. 9.—Am. bor.
pygmæa, *Dun. l. c.* 84. t. 10.—Am. bor.
reticulata, *Shuttlw. ex Chapm. Fl. S. U. St.* 15.—Florida.
rhomboifolia, *Griseb. Cat. Pl. Cub.* 3.—Ind. occ.
secundiflora, Shuttlw. ex Chapm. Fl. S. U. St. 15 = pygmæa.
triloba, *Dun. Monog. Anon.* 83.—Am. bor.

ASIPHONIA, Griff. in Trans. Linn. Soc. xix. (1845) 333. t. 37 = **Bragantia**, Lour. (Aristoloch.).
piperiformis, Griff. l. c. = B. corymbosa.

ASKIDIOSPERMA, Steud. in Flora, xxxiii. (1850) 229, nomen; et Syn. Pl. Cyp. ii. 267 (1855). *RESTIACEÆ*, Benth. & Hook. f. iii. 1092.
capitatum, *Steud. ll. cc.*—Afr. austr.

ASKOFAKE, Rafn. Fl. Tellur. iv. 106 (1836) = **Utricularia**, Linn. (Lentib.).
recurva, Rafn. l. c. = U. recurva.

ASKOLAME, Rafn. Fl. Tellur. ii. 11 (1836) = **Milla**, Cav. (Liliac.).
bifora, Rafn. l. c. 12 = *M. biflora*.

ASPALATHUS, Linn. Sp. Pl. 711 (1753). *LEGUMINOSÆ*, Benth. & Hook. f. i. 476.
ACROSCHINUM, Desv. in Ann. Sc. Nat. ix. (1826) 408.
CYPROCALYX, HETEROLATHUS, LAPASANTHUS, PACHYRAPHEA, PARASPALATHUS, PLAGIOSTIGMA, PSILOLEPUS, STREPTOSTEMA, et TRINCURIA, Presl, in Abh. Boehm. Ges. Wiss. Folge V. iii. 556-562 (Bot. Bemerk. 196-198 (1844).
DIALLOSPERMA, Rafn. Sylva Tellur. 69 (1838).
EXIOCYLAX, Neck. Elem. iii. 95 (1790).
NEFRAKIS, Rafn. Sylva Tellur. 69 (1838).
SARCOCALYX, Walp. in Linnæa, xiii. (1839) 479.
SARCOPHYLLUS, Thunb. in Schrad. Journ. i. (1799) 298.
SCALIGERA, Adans. Fam. ii. 373 (1763).
SEMETOR, Rafn. Sylva Tellur. 69 (1838).
bibetifera, E. Mey. Comm. Pl. Afr. Austr. 48 = sclophylla.
bibetina, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
scanthes, *Eckl. & Zeyh. Enum.* 215.—Afr. austr.
acanthophylla, Eckl. & Zeyh. l. c. 221 = Chamisaensis.
acicularis, *E. Mey. Comm. Pl. Afr. Austr.* 46.—Afr. austr.
aciphylla, *Harv. in Hore. & Sond. Fl. Cap.* ii. 129.—Afr. austr.
aculeata, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
aculeata, Willd. ex Walp. in Linnæa, xiii. (1839) 486 = tridentata.
acuminata, *Lam. Encyc.* i. 287.—Afr. austr.
acuminata, Thunb. Prod. Pl. Cap. 125 = ambigua.
adelphea, *Eckl. & Zeyh. Enum.* 219.—Afr. austr.
adscendens, Steud. Nom. ed. II. i. 147 = elongata.
aemula, *E. Mey. Comm. Pl. Afr. Austr.* 45.—Afr. austr.
affinis, Eckl. & Zeyh. Enum. 219 = microdon.
affinis, E. Mey. Comm. Pl. Afr. Austr. 60 = pinguis.
affinis, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
Agardhiana, *DC. Prod.* ii. 148.—Afr. austr.
albanensis, Eckl. & Zeyh. Enum. 211 = frankenioides.
albens, E. Mey. ex Benth. in Hook. Lond. Journ. Bot. vi. (1848) 648 = exilis.
albens, *Linn. Mant.* ii. 261; *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
albens, Willd. ex Walp. in Linnæa, xiii. (1839) 499 = aurea.
albiflora, Eckl. & Zeyh. Enum. 207 = thymifolia.
alopecuroides, E. Mey. Comm. Pl. Afr. Austr. 52 = Alopecurus.
Alopecurus, *Burch. ex Benth. in Hook. Lond. Journ. Bot.* vi. (1848) 628, (err. typ. 618).—Afr. austr.
alpina, Eckl. & Zeyh. Enum. 218 = rubro-fusca.
alternifolia, *Harv. in Hore. & Sond. Fl. Cap.* ii. 141.—Afr. austr.
alternifolia, *Sprang. Syst.* iii. 187.—Afr. austr.
ambigua, *DC. Prod.* ii. 148.—Afr. austr.
angustissima, *E. Mey. Comm. Pl. Afr. Austr.* 44.—Afr. austr.
anthylloides, Berg. Descr. Pl. Cap. 211 = lotoides.
anthylloides, *Linn. Sp. Pl.* ed. II. 1002.—Afr. austr.
appendiculata, E. Mey. in Linnæa, vii. (1832) 157 = ciliaris.
arachnoidea, Otto, ex Steud. Nom. ed. II. i. 147, 149 = setacea.
arachnoides, *Walp. in Linnæa*, xiii. (1839) 497.—Afr. austr.
arausica, *Linn. Sp. Pl.* 712.—Afr. austr.
arborea, *Lour. Fl. Cochinch.* 481.—Cochinch.
argentea, Eckl. & Zeyh. Enum. 208 = sericea.
argentea, *Linn. Sp. Pl.* 713.—Afr. austr.
argentea, J. F. Gmel. Syst. 1098 = sericea.
argyrea, *DC. Prod.* ii. 189.—Afr. austr.
aristata, *E. Mey. in Linnæa*, vii. (1832) 156.—Afr. austr.
armata, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
ascendens, E. Mey. Comm. Pl. Afr. Austr. 41 = elongata.
asparagoides, *Linn. f. Suppl.* 291.—Afr. austr.
astroides, Willd. ex Walp. in Linnæa, xiii. (1839) 505 = racemosa.
astroites, *Linn. Sp. Pl.* 711.—Afr. austr.
astroites, Thunb. Prod. Pl. Cap. 128 = racemosa.
aulonagnae, Eckl. & Zeyh. Enum. 216 = ciliaris.
batodes, Eckl. & Zeyh. l. c. 216 = sciophylla.
Benthamii, *Harv. in Hore. & Sond. Fl. Cap.* ii. 111.—Afr. austr.

ASPALATHUS:—
bicolor, Eckl. & Zeyh. Enum. 205 = retroflexa.
biflora, E. Mey. Comm. Pl. Afr. Austr. 64 = pedunculata.
bracteata, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
Burchellianus, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 693.—Afr. austr.
calcarata, *Harv. in Hore. & Sond. Fl. Cap.* ii. 116.—Afr. austr.
callosa, *Linn. Sp. Pl.* 713.—Afr. austr.
callosa, Sims, Bot. Mag. t. 2329 = araneosa.
canaliculata, E. Mey. Comm. Pl. Afr. Austr. 44 = stenophylla.
candicans, *Ait. Hort. Kew.* ed. II. iv. 264.—Afr. austr.
canescens, *Linn. Mant.* ii. 262.—Afr. austr.
capillaris, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 653.—Afr. austr.
capitata, *Linn. Fl. Afr. Rar.* 14; *Amœn. Acad.* vi. 92.—Afr. austr.
capitella, Burch. ex Benth. in Hook. Lond. Journ. Bot. vii. (1848) 597 = inops.
carnea, Eckl. & Zeyh. Enum. 1404 = thymifolia.
carnea, *Berg. Descr. Pl. Cap.* 206.—Afr. austr.
carnosa, Eckl. & Zeyh. Enum. 220 = pachyloba.
carnosa, Linn. Mant. 261 = arcuata.
Cephalotes, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 615 = Forbesii.
Cephalotes, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
cervicantha, Eckl. & Zeyh. Enum. 208 = spicata.
Chamisaensis, *Vog. in Linnæa*, x. (1835-36) 597.—Afr. austr.
Chenopoda, *Linn. Sp. Pl.* 711.—Afr. austr.
chortophila, Eckl. & Zeyh. Enum. 211 = frankenioides.
ciliaris, Sims, Bot. Mag. t. 2288 = A. Chenopoda.
ciliaris, *Linn. Mant.* 262.—Afr. austr.
cinerascens, *E. Mey. Comm. Pl. Afr. Austr.* 54.—Afr. austr.
cinerea, Thunb. Prod. Pl. Cap. 125 = cytisoides.
corrulescens, E. Mey. Comm. Pl. Afr. Austr. 54 = Lebeckia microphylla.
cognata, Presl, Bot. Bemerk. 196 = corymbosa.
collina, *Eckl. & Zeyh. Enum.* 220.—Afr. austr.
comosa, Eckl. & Zeyh. l. c. 209 = A. Alopecurus.
comosa, E. Mey. Comm. Pl. Afr. Austr. 56 = Benthamii.
comosa, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
comosa, Willd. ex Walp. in Linnæa, xiii. (1839) 490 = capitata.
conferta, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 596 = exigua.
corruscifolia, Berg.. Descr. Pl. Cap. 207 = genistoides, pungens.
corruscifolia, DC. Prod. ii. 189 = rigescens.
corymbosa, *E. Mey. in Linnæa*, vii. (1832) 159.—Afr. austr.
costulata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 641.—Afr. austr.
crassifolia, *Andr. Bot. Rep.* t. 351.—Afr. austr.
cretica, Linn. Sp. Pl. 713 = Anthyllis Hermanniae.
cuneata, D. Don, Prod. Fl. Nep. 246 = Lespedeza sericea.
cuneata, *E. Mey. Comm. Pl. Afr. Austr.* 37.—Afr. austr.
cymbaeformis, DC. Prod. ii. 140 = uniflora.
cytisoides, *Lam. Encyc.* i. 292.—Afr. austr.
dasyantha, *Eckl. & Zeyh. Enum.* 203.—Afr. austr.
decidualifolia, Eckl. & Zeyh. l. c. 210 = nigra.
densifolia, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 625.—Afr. austr.
diffusa, *Eckl. & Zeyh. Enum.* 208.—Afr. austr.
divaricata, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
divergens, E. Mey. Comm. Pl. Afr. Austr. 45 = prunc.
dimorpha, Willd. ex Walp. in Linnæa, xiii. (1839) 498 = microphylla
Dorycnium, Lam. Encyc. i. 290 = Dorycnium suffruticosum.
Dregeana, Walp. in Linnæa, xiii. (1839) 496 = incurva.
dubia, E. Mey. in Linnæa, vii. (1832) 157 = ciliaris.
dubia, Willd. ex Walp. in Linnæa, xiii. (1839) 504 = Amphithalea ericaefolia.
Ebenus, Linn. Syst. ed. X. 1158 = Brya Ebenus.
ebracteata, E. Mey. Comm. Pl. Afr. Austr. 51 = rigescens.
echinata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Hystrix.
Ecklonis, Vog. ex Walp. in Linnæa, xiii. (1839) 499 = ciliaris.

ASPALATHUS:—
elongata, *Eckl. & Zeyh. Enum.* 203.—Afr. austr.
elongata, E. Mey. Comm. Pl. Afr. Austr. 68 = incurva.
ericaefolia, Berg. Desc. Pl. Cap. 205 = unifora.
ericifolia, *Linn. Sp. Pl.* 711.—Afr. austr.
ericaefolia, Willd. ex Walp. in Linnaea, xiii. (1839) 495 = thymifolia.
ericoides, E. Mey. in Linnaea, vii. (1832) 160 = ericifolia.
erinacea, *Lam. Encyc. i.* 291.—Afr. austr.
eriophylla, *Walp. in Linnaea*, xiii. (1839) 499.—Afr. austr.
erythroides, *Eckl. & Zeyh. Enum.* 200.—Afr. austr.
exigua, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 597 = marginata.
exigua, *Eckl. & Zeyh. Enum.* 198.—Afr. austr.
exilis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 186.—Afr. austr.
falcata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 655.—Afr. austr.
ferox, *Harv. in Harv. & Sond. Fl. Cap.* ii. 188.—Afr. austr.
ferruginea, *Banks, ex Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 607.—Afr. austr.
filicaulis, *Eckl. & Zeyh. Enum.* 204.—Afr. austr.
filifolia, E. Mey. in Linnaea, vii. (1832) 158 = abietina.
flavescens, Presl, Bot. Bemerk. 126 = spinosa.
flexuosa, Thunb. Prod. Pl. Cap. 127 = mollis.
flexuosa, Willd. ex Walp. in Linnaea, xiii. (1839) 495 = ericifolia.
floribunda, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 598.—Afr. austr.
fornicata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 646.—Afr. austr.
frankenioides, *DC. Prod.* i. 189.—Afr. austr.
frankenioides, Eckl. & Zeyh. Enum. 207 = thymifolia.
frutescens, Vog. ex Walp. in Linnaea, xiii. (1839) 487 = pinea.
fusca, Thunb. Prod. Pl. Cap. 125; Fl. Cap. ed. Schult. 574 = callosa.
galeata, *E. Mey. Comm. Pl. Afr. Austr.* 49.—Afr. austr.
galioides, Berg. Desc. Pl. Cap. 210 = retroflexa.
galioides, *Linn. Mant.* ii. 260; *Thunb. Diss. Aspal.* 19.—Afr. austr.
galioides, Sieber, ex Walp. in Linnaea, xiii. (1839) 502 = divaricata.
galioides, Willd. ex Walp. l. c. 493, 509 = pedunculata.
garipensis, E. Mey. Comm. Pl. Afr. Austr. 44.—Afr. austr.
genistoides, Eckl. & Zeyh. Enum. 215 = laricifolia.
genistoides, *Linn. Mant.* 261.—Afr. austr.
genistoides, Thunb. Prod. Pl. Cap. 128 = pungens.
Gillii, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 637 = setacea.
glauca, Eckl. & Zeyh. Enum. 221 = obtusata.
globosa, *Andr. Bot. Rep.* t. 510.—Afr. austr.
globosa, Hort. ex Steud. Nom. ed. II. i. 748 = nigra.
globosa, Eckl. & Zeyh. Enum. 209 = spicata.
globulosa, E. Mey. in Linnaea, vii. (1832) 159 = nigra.
glomerata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 633.—Afr. austr.
glomerata, Linn. f. Suppl. 321 = capitata.
grandiflora, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 632.—Afr. austr.
heterophylla, *E. Mey. Comm. Pl. Afr. Austr.* 40.—Afr. austr.
heterophylla, Linn. f. Suppl. 321; Thunb. Prod. Pl. Cap. 126 = lotoides.
hirsuta, Eckl. & Zeyh. Enum. 212 = frankenioides.
hilaris, Eckl. & Zeyh. l. c. 214 = laricifolia.
hirta, *E. Mey. in Linnaea*, vii. (1832) 156.—Afr. austr.
hispida, Sieber, ex Presl, Bot. Bemerk. 131 = unifora.
hispida, Thunb. Prod. Pl. Cap. 127 = thymifolia.
holosericea, E. Mey. Comm. Pl. Afr. Austr. 37 = Buchenrodera holosericea.
horrida, Eckl. & Zeyh. Enum. 221 = spinosa.
Hystrix, Eckl. & Zeyh. l. c. 219 = Willdenowiana.
Hystrix, *Linn. f. Suppl.* 322.—Afr. austr.
incomitus, E. Mey. ex Benth. in Hook. Lond. Journ. Bot. vii. (1848) 648 = lacera.
incurva, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
incurva, Eckl. & Zeyh. Enum. 208 = cornuta.
incurva, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.

ASPALATHUS:—
incurvifolia, *Vog. ex Walp. in Linnaea*, xiii. (1839) 497.—Afr. austr.
indica, Linn. Sp. Pl. 712 = Indigofera aspalathoides.
iniquua, Eckl. & Zeyh. Enum. 219 = adelphea.
inops, *Eckl. & Zeyh. l. c.* 197.—Afr. austr.
intermedia, Eckl. & Zeyh. l. c. 211 = frankenioides.
involucrata, E. Mey. Comm. Pl. Afr. Austr. 38 = undulata.
involucrata, Pappe, ex Harv. & Sond. Fl. Cap. ii. 114 = nigra.
Jacobaea, *E. Mey. Comm. Pl. Afr. Austr.* 41.—Afr. austr.
Jonbertiana, Eckl. & Zeyh. Enum. 214 = canescens.
juniperina, Thunb. Prod. Pl. Cap. 126 = galioides.
kannatnaie, Eckl. & Zeyh. Enum. 207 = ericifolia.
Krauseana, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 69 = anthylloides.
lactea, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
lanata, *E. Mey. Comm. Pl. Afr. Austr.* 64.—Afr. austr.
lanceolata, *E. Mey. l. c.* 87.—Afr. austr.
laricifolia, *Berg. Desc. Pl. Cap.* 204.—Afr. austr.
laricifolia, Lam. Encyc. i. 287 = leptophylla.
laricina, DC. Prod. ii. 141 = laricifolia.
laricina, Steud. Nom. ed. II. i. 149 = tridentata.
laxata, *Linn. Mant.* 268.—Afr. austr.
lepida, *E. Mey. Comm. Pl. Afr. Austr.* 58.—Afr. austr.
leptocoma, Eckl. & Zeyh. Enum. 205 = divaricata.
leptophylla, *Eckl. & Zeyh. l. c.* 219.—Afr. austr.
leptothrix, Eckl. & Zeyh. l. c. 213 = frankenioides.
leucocephala, E. Mey. Comm. Pl. Afr. Austr. 41 = virgata.
leucophaea, *Harv. in Harv. & Sond. Fl. Cap.* ii. 112.—Afr. austr.
linearifolia, *DC. Prod.* ii. 142.—Afr. austr.
linifolia, E. Mey. in Linnaea, vii. (1832) 162 = linearifolia.
linifolia, Steud. in Flora, xiii. (1830) 543 = heterophylla.
longifolia, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 634.—Afr. austr.
longipes, *Harv. in Harv. & Sond. Fl. Cap.* ii. 141.—Afr. austr.
lotoides, E. Mey. Comm. Pl. Afr. Austr. 42 = Meyeri.
lotoides, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
lotoides, Willd. ex Walp. in Linnaea, xiii. (1839) 486, 508 = tridentata.
macrantha, *Harv. in Harv. & Sond. Fl. Cap.* ii. 117.—Afr. austr.
macrocarpa, *Eckl. & Zeyh. Enum.* 208.—Afr. austr.
macrocephala, *Steud. Nom.* ed. II. i. 149.—Afr. austr.
marginalis, *Eckl. & Zeyh. Enum.* 218.—Afr. austr.
marginata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 108.—Afr. austr.
melanoides, Eckl. & Zeyh. Enum. 210 = nigra.
Meyeri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 106.—Afr. austr.
Meyeriana, Eckl. & Zeyh. Enum. 216 = ciliaris.
micrantha, E. Mey. in Linnaea, vii. (1832) 161 = thymifolia.
microcarpa, *Burch. ex DC. Prod.* ii. 139.—Afr. austr.
microdon, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 642.—Afr. austr.
microphylla, DC. Prod. ii. 148 = divaricata.
microphylla, *Steud, in Flora*, xiii. (1830) ii. 545.—Afr. austr.
minutifolia, Vog. ex Walp. in Linnaea, xiii. (1839) 487 = nigra.
mollis, *Lam. Encyc.* i. 290.—Afr. austr.
mucronata, Linn. f. Suppl. 320 = Viborgia armata.
multiflora, Thunb. Prod. Pl. Cap. 128 = thymifolia.
multiflora, Sieber, ex Walp. in Linnaea, xiii. (1839) 495 = ericifolia.
Mundiana, Eckl. & Zeyh. Enum. 220 = verrucosa.
murallioides, Eckl. & Zeyh. l. c. 209 = mollis.
myrtillifolia, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 597.—Afr. austr.
nanthis, Eckl. & Zeyh. Enum. 213 = canescens.
nervosa, *E. Mey. Comm. Pl. Afr. Austr.* 62.—Afr. austr.
nigra, *Linn. Mant.* 262.—Afr. austr.
nigrescens, E. Mey. in Linnaea, vii. (1832) 159 = nigra.
nivea, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
nivea, Willd. ex Walp. in Linnaea, xiii. (1839) 486 = purpurea.
nodosa, *Vog. ex Walp. l. c.* 496.—Afr. austr.

ASPALATHUS:—
nudiflora, *Harv. in Harv. & Sond. Fl. Cap.* ii. 140.—Afr. austr.
obtusata, *Thunb. Prod. Pl. Cap.* 125.—Afr. austr.
opaca, *Eckl. & Zeyh. Enum.* 217.—Afr. austr.
orbicularis, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 595.—Afr. austr.
ornigera, Eckl. & Zeyh. Enum. 217 = ciliaris.
orientalis, *Linn. Mant.* 263.—Orient.
pachylobe, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 638.—Afr. austr.
pallens, Eckl. & Zeyh. Enum. 210 = nigra.
pallescens, *Eckl. & Zeyh. l. c.* 215.—Afr. austr.
papillosa, Eckl. & Zeyh. l. c. 217 = ciliaris.
Pappeana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 117.—Afr. austr.
parviflora, *Berg. Desc. Pl. Cap.* 208; *Thunb. Prod. Pl. Cap.* 122.—Afr. austr.
pedicellata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 135.—Afr. austr.
pedunculata, E. Mey. ex Walp. Ann. ii. 888 = bracteata, capillaris.
pedunculata, *L'Hérit. Sert. Angl.* t. 26.—Afr. austr.
persica, Burm. f. Fl. Ind. 155 = Lotus Garcini.
pilosa, Eckl. & Zeyh. Enum. 202 = tridentata.
pilosa, *Linn. Mant.* 263.—Afr. austr.
pilosa, Sieber, ex Walp. in Linnaea, xiii. (1839) 488 = arancosa.
pinea, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
pinguis, Eckl. & Zeyh. Enum. 220 = arida.
pinguis, E. Mey. Comm. Pl. Afr. Austr. 60 = affinis.
pinguis, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
pinnata, *Linn. Pl. Rar. Afr.* 14; *Amoen. Acad.* vi. 92.—Afr. austr.
Plukenetiana, Eckl. & Zeyh. Enum. 200 = rugosa.
polistes, Eckl. & Zeyh. l. c. 213 = frankenioides.
polyantha, *Walp. in Linnaea*, xiii. (1839) 485.—Afr. austr.
polycephala, *E. Mey. Comm. Pl. Afr. Austr.* 39.—Afr. austr.
Priori, *Harv. in Harv. & Sond. Fl. Cap.* ii. 127.—Afr. austr.
procumbens, *Eckl. & Zeyh. Enum.* 202.—Afr. austr.
procumbens, E. Mey. in Linnaea, vii. (1832) 162 = lotoides.
propinqua, *E. Mey. Comm. Pl. Afr. Austr.* 53.—Afr. austr.
prostrata, *Eckl. & Zeyh. Enum.* 206.—Afr. austr.
psoraleoides, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 598.—Afr. austr.
pulchella, E. Mey. Comm. Pl. Afr. Austr. 38 = Buchenrodera setacea.
pungens, *Thunb. Prod. Pl. Cap.* 129.—Afr. austr.
purpurascens, E. Mey. Comm. Pl. Afr. Austr. 44 = purpurea.
purpurea, *Eckl. & Zeyh. Enum.* 201.—Afr. austr.
quinquefolia, Linn. Pl. Rar. Afr. 14; Amoen. Acad. vi. 93 = lotoides.
quinquefolia, Thunb. Fl. Cap. 575 = virgata.
racemosa, *E. Mey. in Linnaea*, vii. (1832) 157.—Afr. austr.
ramulosa, E. Mey. l. c. 163 = scoparia.
recurva, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 649.—Afr. austr.
remota, *Eckl. & Zeyh. Enum.* 218.—Afr. austr.
retroflexa, Eckl. & Zeyh. l. c. 204 = abietina.
retroflexa, *Linn. Sp. Pl.* 712.—Afr. austr.
rigescens, *E. Mey. Comm. Pl. Afr. Austr.* 52.—Afr. austr.
rostrata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 631.—Afr. austr.
rubens, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
rubescens, Eckl. & Zeyh. Enum. 213 = adelphea.
rubrofusca, *Eckl. & Zeyh. l. c.* 216.—Afr. austr.
rugosa, *Thunb. Prod. Pl. Cap.* 125.—Afr. austr.
sanguinea, Eckl. & Zeyh. Enum. 213 = verniculata.
sanguinea, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
sarcantha, *Vog. ex Walp. in Linnaea*, xiii. (1839) 489.—Afr. austr.
sarcodes, *Vog. ex Walp. l. c.* 480.—Afr. austr.
scaphoides, Eckl. & Zeyh. Enum. 206 = unifora.
Scholliana, *Presl, Bot. Bemerk.* 129.—Afr. austr.
secundu, E. Mey. Comm. Pl. Afr. Austr. l. 67 = pungens.
securifolia, *Eckl. & Zeyh. Enum.* 198.—Afr. austr.
sericantha, E. Mey. Comm. Pl. Afr. Austr. 49 = laricifolia.
sericea, *Berg. Desc. Pl. Cap.* 213.—Afr. austr.

ASPALATHUS :—
setacea, Eckl. & Zeyh. Enum. 202 = Jacobæa.
setacea, Lam. Encyc. i. 288 = nivea.
setacea, *Linn. f. Suppl.* 321.—Afr. austr.
setacea, Thunb. Prod. Pl. Cap. 125 = Argyrolobium obsoletum.
setacea, *Eckl. & Zeyh. Enum.* 216.—Afr. austr.
Simsiana, Eckl. & Zeyh. l. c. 200 = arancosa.
spathulata, Eckl. & Zeyh. l. c. 198 = securifolia.
speciosa, *Stroud. in Flora*, xiii. (1830) 544.—Afr. austr.
spicata, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
spinescens, DC. Prod. ii. 138 = arida.
spinescens, Eckl. & Zeyh. Enum. 221 = divaricata.
spinescens, *Thunb. Prod. Pl. Cap.* 129.—Afr. austr.
spinifera, L'Hérit. ex Stroud. Non. ed. II. i. 107, 149 = Anthyllis Hermanniæ.
spinosa, *Linn. Sp. Pl.* ed. II. 1000.—Afr. austr.
squamosa, Thunb. Fl. Cap. ed. Schult. 581 = pedunculata.
squarrosa, Thunb. Prod. Pl. Cap. 128 = pedunculata.
stachyera, Eckl. & Zeyh. Enum. 202 = lotoides.
stenarthia, Eckl. & Zeyh. l. c. = argentea.
stellaris, *Eckl. & Zeyh. l. c.* 197.—Afr. austr.
stenophylla, *Eckl. & Zeyh. l. c.*—Afr. austr.
Stendelaina, Bp. ongn.) in Bull. Férus. xxvi. (1831) 278 = microphylla.
stricta, Steud. in Flora, xiii. (1830) ii. 548 = elongata.
suaveolens, *Eckl. & Zeyh. Enum.* 199.—Afr. austr.
subinermis, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 648.—Afr. austr.
subtingens, Eckl. & Zeyh. Enum. 212 = adelphea.
subulata, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
succulenta, E. Mey. in Linnæa, vii. (1832) 159 = verrucosa.
suffruticosa, *DC. Prod.* ii. 144.—Afr. austr.
sulcata, Vog ex Walp. in Linnæa, xiii. (1839) 490 = erythrodes.
tenuifolia, *DC. Prod.* ii. 148.—Afr. austr.
teres, *Eckl. & Zeyh. Enum.* 215.—Afr. austr.
thymifolia, Lam. Encyc. i. 288 = canescens.
thymifolia, *Linn. Sp. Pl.* 711.—Afr. austr.
thymifolia, Thunb. Prod. Pl. Cap. 127 = comosa.
tomentosa, E. Mey. Comm. Pl. Afr. Austr. 55 = frankenioides.
trichodes, E. Mey. l. c. 88 = Buchenroedera trichodes.
tridentata, Berg. ex Walp. in Linnæa, xiii. (1839) 482 = pilosa.
tridentata, Eckl. & Zeyh. Enum. 201 = ferruginea.
tridentata, *Linn. Sp. Pl.* 712.—Afr. austr.
trigona, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
triquetra, *Thunb. l. c.* 127.—Afr. austr.
truncata, *Eckl. & Zeyh. Enum.* 197.—Afr. austr.
tuberculata, *Walp. in Linnæa*, xiii. (1839) 497.—Afr. austr.
tylodes, Eckl. & Zeyh. Enum. 200 = callosa.
ulicina, *Eckl. & Zeyh. l. c.* 206.—Afr. austr.
undulata, *Eckl. & Zeyh. l. c.* 196.—Afr. austr.
uniflora, Eckl. & Zeyh. l. c. 219 = macrantha.
uniflora, *Linn. Sp. Pl.* 712.—Afr. austr.
vaginifera, Thunb. Diss. Aspal. 30 = Wildenowiana.
variana, Eckl. & Zeyh. Enum. 209 = ericifolia.
variegata, *Eckl. & Zeyh. l. c.* 201.—Afr. austr.
venosa, *E. Mey. Comm. Pl. Afr. Austr.* 89.—Afr. austr.
verticillata, *Lam. Encyc.* i. 288.—Afr. austr.
verrucosa, Eckl. & Zeyh. Enum. 220 = tuberculata.
verrucosa, *Linn. Sp. Pl.* 712.—Afr. austr.
verrucosa, Willd. ex Walp. in Linnæa, xiii. (1839) 486 = Wildenowiana.
versicolor, E. Mey. Comm. Pl. Afr. Austr. 48 = callosa.
villosa, *Thunb. Prod. Pl. Cap.* 125.—Afr. austr.
viminea, E. Mey. Comm. Pl. Afr. Austr. 35 = Buchenroedera viminea.
virgata, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
volvularia, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 7.—Malaya.
vulnerans, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 621 = hirta.
vulnerans, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
Wildenowiana, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 590.—Afr. austr.
Wurmbæana, *E. Mey. Comm. Pl. Afr. Austr.* 56.—Afr. austr.

ASPALATUS, St. Hil. in Dict. Sc. Nat. iii. 211 (1804) = Aspalathus, Linn. (Legum.).

ASPALTHIUM, Medic. Phil. Bot. i. 211 (1789) = ASPHALTHIUM, Medic. = **Psoralea**, Linn. (Legum.).

ASPARAGOPSIS, Kunth, in Abh. Akad. Berl. (1842) 35 = **Asparagus**, Tourn. (Liliac.).
abyssinica, Kunth, Enum. Pl. v. 101 = Asparagus racemosus.
acerosa, Kunth, l. c. 102 = Asparagus acerosus
adscendens, Kunth, l. c. = Asparagus adscendens.
aethiopica, Kunth, l. c. 99 = Asparagus aethiopicus.
alba, Kunth, l. c. 84 = Asparagus albus.
arborescens, Kunth, l. c. 79 = Asparagus arborescens.
ascendens, Baker, in Journ. Linn. Soc. xiv. (1875) 624 = Asparagus adscendens.
Brownei, Kunth, Enum. Pl. v. 108 = Asparagus racemosus.
consanguinea, Kunth, l. c. 76 = Asparagus consanguineus.
Curilla, Kunth, l. c. 102 = Asparagus Curillus.
Decaisnei, Kunth, l. c. 103 = Asparagus racemosus.
decumbens, Kunth, l. c. 77 = Asparagus crispus.
densiflora, Kunth, l. c. 96 = Asparagus sarmentosus.
denudata, Kunth, l. c. 82 = Asparagus denudatus.
Dregei, Kunth, l. c. 84 = Asparagus Thunbergianus.
flagellaris, Kunth, l. c. 103 = Asparagus flagellaris.
floribunda, Kunth, l. c. 93 = Asparagus racemosus.
grandiflora, Kunth, l. c. 90 = Asparagus umbellatus.
Hohenackeri, Kunth, l. c. 100 = Asparagus racemosus.
humilis, Hassk. in Versl. en Med. Akad. Wetensch. v. (1857) 93.—Japon.
javanica, Kunth, Enum. Pl. v. 100 = Asparagus racemosus.
juniperina, Kunth, l. c. 85 = Asparagus africanus.
Krebsiana, Kunth, l. c. 93 = Asparagus racemosus.
Lamarchii, Kunth, l. c. 87 = Asparagus africanus.
lancea, Kunth, l. c. 104 = Asparagus sarmentosus.
Lemei, Baker, in Journ. Linn. Soc. xiv. (1875) 611 = Asparagus umbellatus.
microrhaphis, Kunth, Enum. Pl. v. 83 = Asparagus microrhaphis.
minutiflora, Kunth, l. c. 89 = Asparagus minutiflorus.
Niveniana, Kunth, l. c. 88 = Asparagus africanus.
passerinoides, Kunth, l. c. 90 = Asparagus capensis.
retrofractus, Kunth, l. c. 86 = Asparagus retrofractus.
retrofracta, Schweinf. ex Baker, in Journ. Linn. Soc. xiv. (1875) 623 = Asparagus racemosus.
rubricaulis, Kunth, Enum. Pl. v. 87 = Asparagus rubricaulis.
sarmentosa, Dalz. & Gibs. Bomb. Fl. 246 = Asparagus racemosus.
sarmentosa, Kunth, Enum. Pl. v. 97 = Asparagus sarmentosus.
scandens, Kunth, l. c. 78 = Asparagus scandens.
Schlechtendalii, Kunth, l. c. 90 = Asparagus Thunbergianus.
scoparia, Kunth, l. c. 80 = Asparagus asiaticus.
setacea, Kunth, l. c. 82 = Asparagus declinatus.
sinica, Miq. in Journ. Bot. Néerl. i. (1861) 90 (= Asparagus lucidus ?).—Hongkong.
spinescens, Kunth, Enum. Pl. v. 95 = Asparagus stipulaceus.
stipulacea, Kunth, l. c. 91 = Asparagus stipulaceus.
suaveolens, Baker, in Journ. Linn. Soc. xiv. (1875) 617 = Asparagus suaveolens.
subpinnatifolius, Kunth, Enum. Pl. v. 94 = Asparagus racemosus.
Thunbergii, Kunth, l. c. 85 = Asparagus Thunbergianus.
triacantha, Kunth, l. c. 91 = Asparagus stipulaceus.
umbellata, Kunth, l. c. 79 = Asparagus umbellatus.
umbellulata, Kunth, l. c. 81 = Asparagus umbellulatus.
Willdenowii, Kunth, l. c. 86 = Asparagus asiaticus.
Zeyheri, Kunth, l. c. 92 = Asparagus stipulaceus.

ASPARAGUS, Tourn. ex Linn. Syst. ed. I. (1735); Gen. ed. I. 98 (1737). *LILIACEÆ*, Benth. & Hook. f. iii. 765.
ASPARAGOPSIS, Kunth, in Abh. Akad. Berl. (1842) 35.
ELIDE, Medic. Phil. Bot. ii. 71 (1791).
HECATRIA, Salisb. Gen. Pl. Fragm. 56 (1866).
MYRSIPHYLLUM, Willd. in Gen. Naturf. Fr. Berl. Mag. ii. (1808) 25.
abyssinicus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 313.—Abyssin.
aduncicaulis, Schweinf. ex Baker, in Journ. Linn. Soc. xiv. (1875) 616 = Schweinfurthii.

ASPARAGUS:—
acerosus, *Roxb. Hort. Beng.* 34; *Fl. Ind.* ii. 150.—Ind. or.; Burma.
acerosus, *Thunb. ex Schult. f. Syst.* vii. 837.—Afr. austr.
acutifolius, *Linn. Sp. Pl.* 314.—Reg. Mediterr.
adscendens, *Roxb. Hort. Beng.* 34; *Fl. Ind.* ii. 158.—Reg. Himal.; Afghan.
aethiopicus, *Linn. Mant.* i.—Afr. austr.
aetnensis, Tornab. in Atti Acc. Gioenia, Ser. II. xii. (1856) 345 = acutifolius.
africanus, *Lam. Encyc.* i. 295.—Afr. austr.
Agul, *Forsk. Fl. Aegypt. Arab.* p. lxv. *ex Schult. f. Syst.* vii. 837.—Arabia.
albus, *Linn. Sp. Pl.* 313.—Reg. Mediterr. occ.
albus, Thunb. Prod. Pl. Cap. 66 = Thunbergianus.
altilis, Anchera. Fl. Brandenb. 730 = officinalis.
altissimus, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 287.—Algeria.
amarus, DC. Cat. Hort. Monsp. 81 = scaber.
ambiguus, De Not. Rep. Fl. Lig. 401 = acutifolius.
angolensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 254.—Afr. trop.
aphyllus, *Linn. Sp. Pl.* 314.—Reg. Mediterr.
arborescens, *Willd. ex Schult. f. Syst.* vii. 837.—Ins. Teneriffa.
ascendens, Anot. vide adscendens.
asiaticus, *Linn. Sp. Pl.* 313.—Ind. or.; Afr. trop. et austr.
asiaticus, Wight, Ic. t. 2055 = subulatus.
benguellensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 255.—Angola.
brachyphyllus, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 78.—Reg. Daurih.; Soongaria; Afghan.
Bresleriauus, Schult. f. Syst. vii. 838 = maritimus.
brevifolius, *Boiss. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 602.—Lycia.
brevifolius, Tornab. in Atti Acc. Gioenia, Ser. II. xii. (1856) 359 = acutifolius.
Broussonetii, *Savy. Recog.* 3. t. 40.—Afr. bor.
Burchellii, *Baker, in Journ Linn. Soc.* xiv. (1875) 618.—Afr. austr.
Burkei, *Baker, l. c.* 607.—Afr. austr.
Buxbaumius, Gueldenst. ex Ledeb. Fl. Ross. iv. 199 = verticillatus.
capensis, *Linn. Sp. Pl.* 314.—Afr. austr.
capitatus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 607. —Ind. bor. occ.; Beluchist.
caspius, Hohen. Enum. Pl. Talusch. 24 = officinalis.
casplus, *Hort. Vind. ex Schult. f. Syst.* vii. 822.—Reg. Casp.
Chenayi, Baker, in Journ. Linn. Soc. xiv. (1875) 603 = filifolius.
collinus, Jan, ex Parl. Fl. Ital. iii. 14 = tenuifolius.
collinus, Schur, in Oestr. Bot. Zeitschr. x. (1860) 355 = officinalis.
communatus, Tenore, Fl. Nap. iii. 374 = acutifolius.
consanguineus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 615.—Afr. austr.
Cooperi, *Baker, in Gard. Chron.* (1874) i. 818.—Afr. austr.
Corradi, Scop. Fl. Carn. ed. II. i. 346 = acutifolius.
crispus, Boj. Hort. Maurit. 350 = umbellatulus.
crispus, *Lam. Encyc.* i. 295.—Afr. austr.
Curillus, *Buch.-Ham. ex Roxb.* 34; *Fl. Ind.* ii. 152.—Reg. Himal.
Curillus, Wall. Cat. n. 5155 E = officinus.
cuscutoides, *Durch. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 606.—Afr. austr.
davuricus, *Fisch. ex Link, Enum. Hort. Berol.* i. 248. —Davuria.
declinatus, Eckl. & Zeyh. ex Baker, in Journ. Linn. Soc. xiv. (1875) 615 = consanguineus.
declinatus, *Linn. Sp. Pl.* 313.—Afr. austr.
declinatus, Sieber, ex Baker, in Journ. Linn. Soc. xiv. (1875) 621 = retrofractus.
declinatus, Schousb. Vextr. Marok. 178 = foniculaceus.
decumbens, Jacq. Hort. Schœnb. i. 31 = crispus.
deflexus, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 254.—Angola.
densus, *Boland. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 613.—Afr. austr.
denudatus, *Baker, l. c.* 606.—Afr. austr.
dispersiflorus, Schlechtd. ex Kunth, Enum. Pl. v. 95 = juniperinus.
distichophyllus, Thunb. Prod. Pl. Cap. 66 = tetragonus.
distichophyllus, Brouss. ex Webb & Berth. Phyt. Canar. iii. 207 = umbellatus.

ASPARAGUS:—
divaricatus, Rottl. ex Baker, in Journ. Linn. Soc. xiv. (1875) 618 = asiaticus.
divaricatus, *Royle, Illustr. Bot. Himal.* i. 398.—Ind. or.
Draco, Linn. Sp. Pl. ed. II. 451 = Dracæna Draco.
Dregeanus, *Hort.—Cf. Gard. Chron.* (1877) ii. 598.—Afr. austr.
drepanophyllus, *Welw. ex. Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 254.—Afr. trop.
dubius, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 363 = racemosus.
dumosus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 606. —Sindh.
Eckloni, *Baker, l. c.* 616.—Afr. austr.
equisetifolius, Ledru, ex Webb & Berth. Phyt. Canar. iii. 328 = arborescens.
equisetoides, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 258.—Angola.
erectus, *Thunb. Prod. Pl. Cap.* 85.—Afr. austr.
esculentus, Salisb. Prod. 292 = officinalis.
euphraticus, Kunth, Enum. Pl v. 73 = filifolius.
exaltatus, C. Sm. & Buch, ex Webb & Berth. Phyt. Canar. iii. 326 = scoparius.
exuvialis, *Burch. Trav.* i. 432.—Afr. austr.
falcatus, Benth. in Hook. Lond. Journ. Bot. i. (1842) 493 = lucidus.
falsatus, *Linn. Sp. Pl.* 313.—Afr. trop. & austr.; Zeylan.
falcatus, Thunb. Fl. Jap. 189 = lucidus.
fasciculatus, R. Br. Prod. 281 = racemosus, acerosus.
fasciculatus, *Thunb. Pl. Cap.* ii. 329.—Afr. austr.
filicinus, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 49.— Reg. Himal.; Burma.
filifolius, *Kertol. Nov. Comm. Acad. Bonon.* v. (1842) 480 (*Misc. Bot.* i. 20).—Mesopot.
flagellaris, *Baker, in Journ. Linn. Soc.* xiv. (1875) 614.—Senegambia.
flexuosus, Thunb. Prod. Pl. Cap. 86 = crispus.
floribundus, Rottl. ex Baker, in Journ. Linn. Soc. xiv. (1875) 611 = Rottleri.
floriundus, *Royle, Illustr. Bot. Himal.* i. 398.—Ind. or.
foeniculaceus, *Lowe, in Journ. Linn. Soc.* v. (1864) 44. —Marocco.
fragilis, Burch. ex Baker, in Journ. Linn. Soc. xiv. (1875) 606 = denudatus.
Gaudichaudianus, *Kunth, Enum. Pl* v. 71.—Cochinch.
gibbus, Bunge, Enum. Pl. Chin. Bor. 65 = davuricus.
glycocarpus, Kunth, Enum. Pl. v. 61 = davuricus.
gonoclados, *Baker, in Journ. Linn. Soc.* xiv. (1875) 627.—Ind. or.
gracilis, *Royle, Illustr. Bot. Himal.* i. 393.—Reg. Himal.
gracilis, Salisb. Prod. 252 = asiaticus.
graminifolius, Linn. Sp. Pl. ed. II. 450 = Dracæna graminifolia.
grandiflorus, Willd. ex Webb & Berth. Phyt. Canar. iii. 327 = umbellatus.
Griffithii, *Baker, in Journ. Linn. Soc.* xiv. (1875) 604. —Afghan.
hedecarpus, Andr. ex Baker, l. c. 599 = caspius, officinalis.
horridus, Linn. f. Suppl. 208 = stipularis.
horridus, *Murr. Syst.* ed. XIII. 274.—Hab.?
hortensis, Mill. ex Baker, in Journ. Linn. Soc. xiv. (1875) 599 = officinalis.
inarimensis, Tornab. in Atti Acc. Gioenia, Ser. II. xii. (1856) 368 = acutifolius.
inderiensis, Blume, ex Ledeb. Fl. Alt. ii. 44 = maritimus.
insularis, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 246 = lucidus.
irregularis, *Baker, in Journ. Linn. Soc.* xiv. (1875) 630.—Afr. austr.
Jacquemonti, *Baker, l. c.* 613.—Ind. or.
Krausii, *Baker, l. c.* 626.—Afr. austr.
laxus, Salisb. Prod. 252 = æthiopicus.
laxiflorus, *Sieud. ex Baker*, in *Journ. Linn. Soc.* xiv. (1878) 253.—Ind. or.
laxus, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.
laxiculus, *Burch. Trav.* i. 537.—Afr. austr.
ligulatus, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
littoralis, Steu. in Bull. Soc. Nat. Mosc. xxx. (1857) ii. 98 = amboi.
longiflorus, *Prumph. Pl. Davidson.* i. 300.—As. or.
longifolius, *Hoch. ex Steud. Nom.* ed. II. i. 150.— Abyss.

ASPARAGUS:—
Lowei, *Kunth, Enum. Pl.* v. 78.—Madera.
Lownei, *Baker, in Journ. Linn. Soc.* xiv. (1875) 601. —Palæstin.
lucidus, *Lindl. Bot. Reg.* (1844) *Misc.* 29.—Japon.
Macowani, *Baker, in Journ. Linn. Soc.* xiv. (1875) 609.—Afr. austr.
madagascariensis, *Baker, l. c.* xvi. (1878) 206.—Madag.
marinus, Reichb. lc. Fl. Germ. t. 520 = scaber.
maritimus, Mill. Gard. Dict ed. VIII. n. 2 = scaber.
maritimus, *Pall. Reise*, ii. 339.—Rossia; Reg. Cauc.; Persia; Sibiria.
mauritianus, Lam. ex Desf. Tabl. ed. I. 20 = declinatus.
medeoloides, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.
micranthus, Lindl. Swan Riv. App. 58 = Corynotheca dichotoma.
micranthus, Sieb. & Zucc. ex Baker, in Journ. Linn Soc. xiv. (1875) 604 = schoberioides.
microphyllus, Burch. ex Baker, l. c. 619 = africanus.
microrrhabis, *Baker, l. c.* 612.—Afr. austr.
minutiflorus, *Baker, l. c.* 616.—Afr. austr.
mitis, *A. Rich. Tent. Fl. Abyss.* ii. 319.—Abyss.
monophyllus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 604.—Beloochist.
mozambicus, *Kunth, Enum. Pl.* v. 72.—Afr. trop.
multiflorus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 610.—Afr. austr.
neglectus, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 750 = trichophyllus.
Nelsoni, *Baker, in Journ. Linn. Soc.* xiv. (1875) 617. —Afr. austr.
nepalensis, *Baker, l. c.* 622.—Reg. Himal.
Nivenianus, *Schult. f. Syst.* vii. 831.—Afr. austr.
nodosus, *Soland. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 608.—Afr. austr.
officinalis, *Linn. Sp. Pl.* 313.—Europ.; Reg. Caucas.; Sibir.
oligoclonus, *Maxim. Prim. Fl. Amur.* 286.—Reg. Amur.
oligophyllus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 604.—As. temp.
orientalis, *Boss. Cat. Hort. Crem.* (1814).—Hab.?
oxycanthus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 625.—Afr. austr.
oxycarpus, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) ii. 92 = officinalis.
palæstinus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 602.—Palæstin.
pallidus, Salisb. Prod. 252 = albus.
paragus, Gueldenst. ex Ledeb. Fl. Ross. iv. 197 = officinalis.
parviflorus, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 180.—Sibir. boical.
Pastorianus, Webb & Berth. Phyt. Canar. iii. 329. t. 329 = albus.
Paull-Guilelmi, *Solms, in Schweinf. Beitr. Fl. Aethiop.* 208.—Afr. trop.
pectinatus, [DC. in] Red. Lil. t. 407 = scandens.
penduliflorus, Zipp. ex Span. in Linnæa, xv. (1841) 476 = dubius.
persicus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 608. —Persia.
Petersianus, *Kunth, Enum. Pl.* v. 72.—Afr. trop.?
Pettianus, *A. Rich. Tent. Fl. Abyss.* ii. 320.—Abyss.
phyllacanthus, Lam. Encyc. i. 296 = aphyllus.
pilosus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 610. —Afr. trop.
planiuscula, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
plocamoides, Webb, in Steud. Nom. ed. II. i. 150 = scoparius.
plumosus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 613. —Afr. austr.
polyphyllus, Stev. ex Ledeb. Fl. Ross. iv. 198 = officinalis.
prostratus, Dum. Fl. Belg. 188 = officinalis.
pullarus, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 255.—Angola.
puberulus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 616.—Afr. austr.
pubescens, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 254.—Angola.
purpurascens, Bieb. ex Baker, in Journ. Linn. Soc. xiv. (1875) 609 = trichophyllus.
racemosus, *Willd. Sp. Pl.* ii. 152.—Ind. or.; Afr. trop.; Austral.
ramosissimus, *Baker, in Gard. Chron.* (1874) ii. 6.— Afr. austr.
retrofractus, *Forsk. Fl. Aegypt. Arab.* 78.—Arabia.

ASPARAGUS:—
retrofractus, *Linn. Sp. Pl.* 313.—Afr. austr.
retrofractus, C. Sm. ex Webb & Berth. Phyt. Canar. iii. 326 = arborescens.
retrofractus, *Schousb. Vestr. Marok.* 174 (sp. dubia). —Marocco.
rivalis, *Burch. ex Schult. f. Syst.* vii. 837.—Afr. austr.
Rottleri, *Baker, in Journ. Linn. Soc.* xiv. (1875) 611. —Ind. or.
ruber, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
rubricaulis, *Baker, in Journ. Linn. Soc.* xiv. (1875) 616.—Ind. or.
rubicundus, Berg. Descr. Pl. Cap. 88 = africanus, capensis, stipularcus.
sarmentosus, Heyne, in Wall. Cat. n. 5154 = volubilis.
sarmentosus, *Linn. Sp. Pl.* 314.—Ind. or.
sativus, Mill. Gard. Dict. ed. VIII. Add. = officinalis.
scaber, *Brign. Pesc. Fl. Perojus.* 22.—Reg. Mediterr. et Caucas.
scaber, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 11 = umbellatus.
scaberulus, *A. Rich. Tent. Fl. Abyss.* ii. 320.—Abyss.
scandens, Gueldenst. ex Ledeb. Fl. Ross. iv. 199 = verticillatus.
scandens, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.
schoberioides, *Kunth, Enum. Pl.* v. 70.—Japon.
Schweinfurthii, *Baker, in Journ. Linn. Soc.* xiv. (1875) 616.—Afr. trop.
scoparius, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 11.—Marocco; Ins. Madera.
Sieboldii, *Maxim. Prim. Fl. Amur.* 387.—Japon.
simulans, *Baker, in Journ. Linn. Soc.* xiv. (1875) 609. —Madag.
sinensis, Nolszt, ex Baker, l. c. 607 = crispus.
Smithianus, *Webb, ex Steud. Nom.* ed. II. i. 150, nomen. —Ins. Teneriff.
spinescens, *Steud. in Schult. f. Syst.* vii. 834.—Afr. austr.
squarrosus, *Y. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 165.—Ins. Cap. Verd.
stachyoides, *Spreng. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 624.—Afr. austr.
stellatus, *Baker, l. c.* 612.—Afr. austr.
stipulaceus, *Lam. Encyc.* i. 296.—Afr. austr.
stipularis, *Forsk. Fl. Aegypt. Arab.* 72.—Reg. Mediterr.; Ins. Canar.
striatus, *Thunb. Prod. Pl. Cap.* 85.—Afr. austr.
suaveolens, *Burch. Trav.* i. 226.—Afr. austr.
subalatus, *Steud. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 614 (*err.* subalatus).—Ind. or.
subulatus, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.
sylvaticus, Burch. ex Baker, in Saund. Ref. Bot. i. 214 = virgatus.
sylvaticus, Waldst. & Kit. Pl. Rar. Hung. iii. 238 = tenuifolius.
Tauscheri, Fisch. ex Ledeb. Fl. Alt. ii. 44 = maritimus.
tenuifolius, Gilib. Exercit. ii. 459 = officinalis.
tenuifolius, *Lam. Encyc.* i. 294.—Europ. austr.
terminalis, Linn. Sp. Pl. ed. II. 450 = Cordyline terminalis.
tetragonus, Bresler, Diss. n. 15 = racemosus.
Thunbergianus, *Schult. f. Syst.* vii. 833.—Afr. austr.
triacanthus, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
tricarinatus, C. Koch, in Linnæa, xxii. (1849) 266 = verticillatus.
tricarinus [*Delile, in*] *Red. Lil.* t. 451.—Hab.?
trichophyllus, *Bunge, Enum. Pl. Chin. Bor.* 65.— Afghan.; As. med.; Chin. bor.
umbellatus, *Link, in Buch. Beschr. Canar. Ins.* 140.— Ins. Canar.
umbellatus, *Royle, Illustr. Bot. Himal.* i. 393.—Ind. or.
umbellatus, *Bressler, Diss.* n. 24.—Ins. Maurit.
umbricola, *Kit. ex Kanitz, Relig. Kitaib.* 121; (*in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 545.— Europ.
undulatus, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.
vaginellatus, *Bej. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 612.—Madag.
verticillaris, Lam. Encyc. i. 297 = verticillatus.
verticillatus, *Linn. Sp. Pl.* ed. II. 450.—Ross. austr.; Oriens.
virgatus, *Baker, in Saund. Ref. Bot.* t. 214.—Afr. austr.
volubilis, Georgi, Beschreib. Russ. Reich. Nachtr. 268 = verticillatus.
volubilis, Buch.-Ham. ex Wall. Cat. n. 5154 = adscendens.
volubilis, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.

ASPARAGUS:—
vulgaris, Gaudenst. ex Ledeb. Fl. Ross. iv. 197 = officinalis.
Wrightii, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 413.—Japon.
zanzibaricus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 614.—Afr. trop.

ASPARIA, Lindl. Gen. et Sp. Orch. 139 (1838). *ORCHIDEÆ*, Benth. & Hook. f. iii. 560.
TROPHIANTHUS, Scheidw. in Otto & Dietr. Allg. Gartenz. xii. (1844) 218.
Bikoriana, Reichb. f. in Walp. Ann. vi. 849 = seq.
Bikriana, *Reichb. f. in Bonplandia*, ii. (1854) 90.—Hab.†
epidendroides, *Lindl. in Hook. Journ. Bot.* i. (1834) 5.—Panama.
fragrans, Klotzsch, Ind. Sem. Hort. Berol. (1858) 12 = epidendroides.
lunata, *Lindl. Bot. Reg.* xxii. (1836) sub t. 1907.—Bras.
lyrata, [*Reichb. f. in*] *Bot. Zeit.* x. (1852) 721.—Bras.
papillonacea, *Reichb. f. in Gard. Chron.* (1876) ii. 100. —Costa Rica.
principissa, *Reichb. f. in Bot. Zeit.* x. (1852) 687.—Veraguas.
psittacina, *Reichb. f. in Gard. Chron.* x. (1878) ii. 684. —Ecuad.
variegata, *Lindl. Bot. Reg.* xxii. t. 1907.—Am. austr.

ASPASIA, E. Mey. ex Pfeiff. Nom. i. 295 (1873) = Stachys, Linn. (Labiat.).

ASPASIA, Salisb. Gen. Pl. Fragm. 84 (1866) = Ornithogalum, Linn. (Liliac.).
aurea, Salisb. l. c. = O. thyrsoides.
flavescens, Salisb. l. c. = O. thyrsoides.
lactea, Salisb. l. c. = O. lacteum.
revoluta, Salisb. l. c. = O. revolutum.
thyrsoides, Salisb. l. c. = O. thyrsoides.

ASPEGRENIA, Poepp. & Endl. Nov. Gen. et Sp. ii. 12. t. 118 (1837) = Octomeria, R. Br. (Orchid.).
aciculata, Poepp. & Endl. l. c. = O. aciculata.

ASPELINA, Cass. in Dict. Sc. Nat. xli. 166 (1826) = Senecio, Linn. (Compos.).
nivea, Cass. l. c. = S. Aspelina.

ASPERA, Columna, ex Moench, Meth. 641 (1794) = Galium, Tourn. (Rubiac.).
muralis, Lowe, Man. Fl. Mad. i. 369 = G. murale.
sudana, Moench, Meth. 641 = G. murale.

ASPERELLA, Humb. in Roem. & Ust. Mag. iii. vii. 5 (1790). *GRAMINEÆ*, Benth. & Hook. f. iii. 1207 (Asprella).
GYMNOSTICHUM, Schreb. Beschr. Gras. ii. 127. t. 47 (1810).
HYSTRIX, Moench, Meth. 294 (1794).
Hystrix, *Humb. l. c.*—Oriens.

ASPERUGO, [Tourn.] Linn. Syst. ed. I. (1735). *BORAGINEÆ*, Benth. & Hook. f. ii. 853.
aegyptiaca, Linn. Sp. Pl. ed. II. 196 = Anchusa flava.
alba, [Pier!] in Ionics Anthol. v. 192 = procumbens.
divaricata, Murr. in Nov. Comm. Gotting. (1776) 25. t. 2 = Nonea lutea.
procumbens, *Linn. Sp. Pl.* 138.—Europ.; Arab.; Orient; As. bor.
vulgaris, Dum.-Cours. Bot. Cult. ii. 167 = procumbens.
verrucosa, Boiss. Fl. Orient. iv. 159, in syn. = Anchusa flava.

ASPERULA, Linn. Syst. ed. I (1735). *RUBIACEÆ*, Benth. & Hook. f. ii. 150.
LEPTUNIS, Stev. in Bull. Soc. Nat. Mosc. xxix.(1856) ii. 396, in textu.
BLEHMACROSTEMMA, CHLOROSTEMMA, CYNANCHICA, et GALIOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 295.
acerosa, C. Koch, in Linnæa, xix. (1847) 31 = lilaciflora.
affinis, Boiss. & Huet, Diagn. Ser. II. ii. 110 = cynanchica.
algarica, Pers. Syn. i. 124 = hirsuta.
Allionii, Baumg. Enum. i. 80 = hexaphylla.
alpigena, Schur, Enum. Pl. Transs. 277 = cynanchica.
alpina, Bieb. Fl. Taur. Cauc. Suppl. 108 = cynanchica.
Aparine, *Sess. Fl. Calic.* i. 114.—Europ. austr. occ.; Orient; Sibiria.

ASPERULA:—
arcadiensis, *Sims, Bot. Mag.* t. 2146.—Graecia.
arvatscuk, Reut. Cat. Pl. Genève, ed. II. 101 = cynanchica.
aristata, Griseb. Spicil. Fl. Rumel. ii. 166 = longiflora.
aristata, *Linn. f. Suppl.* 130.—Europ.; Orient.
arundana, Ball, ex Nym. Consp. ii. 334 = pendula.
arvensis, *Linn. Sp. Pl.* 108.—Europ.; Orient.
aspera, *Boiss. Fl. Orient.* iii. 28.—Reg. Cauc.; Persia.
asperrima, *Boiss. Elench.* 56.—Hispan.
Athos, Boiss. Diagn. Ser. I. x. 69 = suberosa.
azurea, Janb. & Spach, Illustr. i. 151. t. 89 = orientalis.
Boissieri, Steud. ex Boiss. Voy. Bot. Espagne, 787 = effusa.
Botschantzewii, Stud. Nom. ed. II. i. 161 = praec.
brachyantha, *Boiss. Diagn.* Ser. II. vi. 66.—Persia.
brevifora, *Boiss. l. c.* Ser. I. x. 68.—Syria.
brevifolia, *Trent. Vard. Cat.*, t. 68.—As. Min.
brunnea, Boiss. Fl. Orient. iii. 32 = brevifolia.
bryoides, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 106.—Lycia.
calabrica, Linn. f. Suppl. 120 = Putoria calabrica.
canescens, Vis. Fl. Dalmat. iii. 11. t. 25 = aristata.
capitata, Kit. ex Reichb. f. lc. Fl. Germ. xvii. t. 1182 = hexaphylla.
cespitosa, Boiss. Fl. Orient. iii. 39 = lilaciflora.
chlorantha, *Boiss. & Heldr. Diagn.* Ser. II. vi. 90.—Graecia.
ciliaris, DC. Prod. iv. 582 = styloes.
ciliata, Moench, Meth. 484 = Putoria calabrica.
ciliata, *Rochel, Banat.* t. ix. f. 23; *Kegel, Ind. Sem. Horop.* (1854) 31.—Bannat.
collina, Salisb. 388 = cynanchica.
commutata, Presl, Fl. Sic. Prod. 62 = longiflora.
conferta, *Hook. f. in Lond. Journ. Bot.* vi. (1847) 468 [bis].—Austral.
consanguinea, Klotzsch, in Bot. Ergeb. Waldem. Reise, 87 = Galium trifforum.
crassifolia, Linn. Mant. 37 = tomentosa.
cretacea, Willd. ex Roem. & Schult. Syst. iii. 529 = cynanchica.
cruciata, *Vell. Fl. Flum.* 44; t. t. 114.—Bras.
cynoes, *Vell. l. c.* 44; t. t. 113.—Bras.
cynanchica, *Linn. Sp. Pl.* 104.—Europ.; As. Min.
Danlevskiana, *Besser, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 302.—Turkestan.
debilis, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 2 = humifusa.
deficiens, *Viviani, in Ann. Stor. Nat.* iv. 281.—Corsica.
densifolia, Vahl, ex Willd. & Lange, Prod. Fl. Hisp. ii. 301 = hirsuta.
digyna, Dufour, in Ann. Sc. Gén. Phyt. vii. (1820) 395 = stellata.
divaricata, *Bischoff, Del. Sem. Hort. Heidelb.* (1846) 4. —Hab.†
divergens, Boiss. Diagn. Ser. II. vi. 90 = graveolens.
dubia, Willd. ex Roem. & Schult. Syst. iii. Mant. 189 = arvensis.
effusa, *Boiss. Voy. Bot. Espagne*, 280.—Hispan.
elongata, *Ledeb. Enum. Pl. Nov.* 58.—Sonogaria.
eriocarpa, Boiss. Elench. 57 = asperrima.
fallax, Ehrh. ex Stend. Nom. ed. II. i. 161 = laevigata.
fascicularis, Boiss. Diagn. Ser. I. x. 69 = stricta.
flaccida, Tenore, Fl. Nap. t. 110 = longiflora.
fortida, Pers. Syn. i. 124 = Putoria calabrica.
fragiliflora, *Boiss. & Hausskn. in Boiss. Fl. Orient.* iii. 36.—Persia.
fragrantissima, *Armst. in Trans. N. Z. Inst.* xiv. (1882) 350.—N. Zel.
galioides, Mx. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 13; et viii. (1881) 161 = galioides.
galioides, *Bieb. Fl. Taur. Cauc.* i. 101.—Europ.; Reg. Cauc.
galioides, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 313 = Galium muricum.
galioides, Jacquem. Journ. ex Hook. f. Fl. Brit. Ind. iii. 205 = Galium trifforum.
geminiflora, *F. Muell. Progrn.* v. 147.—Austral.
glaberrima, Bunge, ex Nym. Consp. 333 = involucrata.
glabra, C. Koch, in Linnæa, xix. (1847) 31 = involucrata.
glauca, Bess. Enum. Pl. Volh. 7 = galioides.
glaucescens, *Griseb. Spicil. Fl. Rumel.* ii. 166.—Orbelus.
gracilis, *C. A. Mey. Verz. Pfl. Cauc.* 154.—Reg. Caucas.
graveolens, *Bieb. ex Ben. in Roem. & Schult. Syst.* iii. *Mant.* 276.—Tauria; As. Min.
Grunii, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 468 [bis].—Austral.

ASPERULA:—
Gussonii, *Boiss. Diagn.* Ser. I. x. 68.—Sicil.
Haussknechtii, *Boiss. Fl. Or. Nov. Dec.* i. 2.—Assyria.
hexaphylla, *All. Fl. Pedem.* i. 19. t. 77. f. 3.—Ital.; Hungar.
hexaphylla, Georgi, Beschr. Russ. Reich. iii. iv. 738 = crassa.
hirsuta, *Desf. Fl. Atlant.* i. 127.—Lusit.; Afr. bor.
hirta, *Ramond, in Bull. Soc. Philom.* (1800) 181. t. 9. —Mont. Pyren.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 87 = Galium trifforum.
humifusa, *Bess. Cat. Hort. Crem.* (1811) *Suppl.* iii. 4; *Rich. Fl. Taur. Cauc. Suppl.* 106.—Podol.; Orient.
incana, Bory & Chaub. Nouv. Fl. Pélop. 10 = taygetea.
inuna, *Sibth. & Sm. Prod. Fl. Graec.* i. 88.—Crete.
involucrata, *Wahlenb. in Isis*, xxi. (1828) 971.—As. Min.
Jordani, Perr. & Song. ex Nym. Consp. 334 = longiflora.
Kotschyana, *Boiss. Fl. Orient.* iii. 27.—Kurdistan.
laevigata, DC. Prod. iv. 582, partim = involucrata.
laevigata, *Linn. Mant.* 38.—Europ. austr.
laxiflora, *Boiss. in Bal. ex Boiss. Fl. Orient.* iii. 37.—Pontus.
libanotica, *Boiss. Diagn.* Ser. I. x. 69.—Syria.
lilaciflora, *Boiss. l. c.* iii. 38.—As. Min.
littoralis, *Sibth. & Sm. Prod. Fl. Graec.* i. 88.—Pontus.
longiflora, Nok, ex Nym. Consp. 333 = cynanchica.
longiflora, *Waldst. & Kit. Pl. Rar. Hung.* ii. 162. t. 150.—Europ.
longifolia, Sibth. & Sm. Fl. Graec. Prod. i. 87 = Galium longifolium.
luteu, *Sibth. & Sm. l. c.* 88.—Graecia.
Lycia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 109.—Lycia.
macrocalada, Huet, in Ann. Sc. Nat. Sér. III. xix. (1853) 255 = cynanchica.
macrorrhiza, Hoffmgg. & Link, Fl. Port. ii. 41. t. 84. = aristata.
maritima, Tenore, ex Nym. Consp. 334 = tomentosa.
matrisylva, Gilib. Exercit. Bot. Univ. Viln. 5 = odorata.
microphylla, *Boiss. Diagn.* Ser. I. xi. 34.—Persia.
minima, *Hook. f. in Lond. Journ. Bot.* vi. (1847) 464 [bis].—Tasmania.
minor, S. F. Gray, Nat. Arr. Brit. Pl. ii. 481 = cynanchica.
mollis, Boiss. Diagn. Ser. I. iii. 31 = arcadiensis.
mollugitoides, *Reichb. Fl. Germ. Excurs.* 206.—Reg. Caucas.
montana, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 151 = cynanchica.
macronata, *Otto, ex Steud. Nom.* ed. II. i. 151.—Hab.†
multiflora, Lapeyr. Hist. Abr. Pl. Pyr. 82 = cynanchica.
Mungieri, Boiss. & Heldr. Diagn. Ser. I. x. 61 = latifo.
muscos, *Boiss. & Heldr. l. c.* Ser. II. ii. 109.—Thessalia.
mutica, Boiss. Fl. Orient. iii. 43 = suavis.
neglecta, Moris, Elench. Sard. ii. 4 = pumila.
neglecta, *Guss. Fl. Ker.* 70. t. 13. f. 2.—Italia.
Nelirischii, *Bech, in Verh. Zool.-Bot. Ges. Wien*, xxxii. (1883) 183.—Europ.
nemoralis, Salisb. Prod. 88 = taurina.
nitens, Guss. Fl. Ker. 70. t. 13. f. 3 = neglecta.
nitida, *Sibth. & Sm. Fl. Graec. Prod.* i. 88.—As. Min.
odora, Salisb. Prod. 87 = odorata.
odorata, *Linn. Sp. Pl.* 108.—Europ.; Orient.
oligantha, F. Müell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 111, 123 = Rubia syrticola.
oppositifolia, *Regel & Schmalh. ex Regel, Pl. Nov. Pedsch.* 62.—As. centr.
orientalis, *Boiss. & Hausskn. Diagn.* Ser. I. iii. 30.—As. Min.; Reg. Caucas.
ovalifolia, Boiss. & Huet, Diagn. Ser. II. i. 113 = gracilis.
palustris, Tenore, Fl. Nap. 111 = tinctoria.
paniculata, Boiss. Elench. 56 = effusa.
paniculata, *Bunge, in Ledeb. Fl. Alt.* i. 140.—Sibir. alt.
papillosa, Kem. ex Nym. Consp. 333 = Galium aristatum.
papillosa, *Lange, in Kjoeb. Vidensk. Meddel.* 1112 [*Pugill.* ii. 121].—Hispan.
pendula, *Boiss. Elench.* 55.—Hispan.

ASPERULA :—
perpusilla, *Hook. Fl. Nov. Zel.* 114.—N. Zel.
Pestalozzæ, Boiss. Diagn. Ser. II. ii. 118 = graveolens.
Platygallium, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 284.—Manchur.
pontica, *Boiss. Fl. Orient.* iii. 89.—Pontus.
prostrata, *C. Koch, in Linnæa*, xxiv. (1851) 461.—Armenia.
pulvinaris, Nym. Consp. ii. 884 = listera.
pumila, *Moris, in Mem. Acc. Torin.* xxxviii. (1835) 96. —Sardinia.
pusilla, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 464 [bis]=Gunnii.
pyrenaïca, Bois. Diagn. Ser. I. x. 64 = hmnifusa
pyrenaica, Linn. Sp. Pl. 104 = cynanchica.
repens, Brot. Fl. Lusit. i. 152 = hirsuta.
rigida, *Sibth. & Sm. Fl. Græc. Prod.* i. 89.—Creta.
rivalis, Sibth. & Sm. l. c. 87 = A. Aparine.
rotundifolia, Linn. Mant. ii. 230 = læviguta.
rubeola, Gratel. in Bull. Soc. Nat. Hist. Bord. i. (1836) 311 = cynanchica.
rubeola, Lam. Fl. Fr. iii. 375 = cynanchica, tinctoria.
rubioides, *Schur, Enum. Pl. Transs.* 276.—Europ.
rumelica, Boiss. Diagn. Ser. II. ii. 113 = graveolens.
rupestris, Reichb. Fl. Germ. Excurs. 205 = scutellaris.
rupicola, Jord. Pugill. 76 = cynanchica.
sanasilia, Lam. Illustr. n. 189 = nitida, pyrenaica.
scabra, Link, Enum. Hort. Berol. i. 184 = tomentosa ?
scabra, Moris, Elench. Sard. ii. 4 = pumila.
scabra, J. & C. Presl, Delic. Prag. 124 = aristata.
scoparia, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 463 [bis].—Australasia.
scutellaris, *Vis. Sem. Rar. Hort. Patav.* (1836):—Albania.
setigerula, *Boiss. Diagn.* Ser. I. vi. 65.—Persia.
setosa, *Jaub. & Spach, Illustr.* i. 152. t. 82. B.—Syria ; Persia.
setulosa, Boiss. Diagn. Ser. I. x. 61 = litoralis.
sherardioides, *Jaub. & Spach, Illustr.* i. 153.—Persia.
sinaica, Delile, ex Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 168 = Galium sinaicum.
sparrerae, C. Koch, in Linnæa, xxiv. (1851) 461 = glomerata.
Stellana, *Vis. Fl. Dalmat.* iii. 11.—Dalmatia.
stricta, *Boiss. Diagn.* Ser. I. iii. 33.—Græcia ; As. Min.
strictissima, Schur, Enum. Pl. Transs. 277 = galioides.
styloza, *Boiss. Diagn.* Ser. I. iii. 31.—Persia.
suaveolens, Schrad. ex DC. Prod. iv. 583 = longiflora.
suavis, *Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop.* ix. 62.—Anatolia.
subalpina, Schur, Enum. Pl. Transs. 277 = cynanchica.
suberosa, Guss. Fl. Sic. Syn. i. 180 = Gussonii.
suberosa, *Sibth. & Sm. Fl. Græc. Prod.* i. 89.—Græcia.
subsimplex, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 468 [bis].—Tasmania.
subvolutina, DC. Prod. iv. 585 = Galium leiophyllum.
sufruticosa, *Boiss. & Heldr. Diagn.* Ser. II. ii. 111.—Græcia.
supina, Bieb. Fl. Taur. Cauc. i. 101 = cynanchica.
taurina, *Linn. Sp. Pl.* 108.—Europ. ; Oriens.
taygetea, *Boiss. & Heldr. Diagn.* Ser. I. x. 60.—Græcia.
Tenori, Wallr. ex Roem. & Schult. Syst. iii. Mant. 189, in adn. = tomentosa.
ternifiora, Jord. Pugill. 77 = cynanchica.
tenuifolia, *Boiss. Diagn.* Ser. I. iii. 32.—As. Min.
tenuissima, *C. Koch, in Linnæa*, xxiv. (1851) 462.—Oriens.
thessala, Boiss. & Heldr. Diagn. Ser. II. ii. 111 = longiflora.
tinctoria, *Linn. Sp. Pl.* 104.—Europ.
tomentosa, *Tenore, Prod. Fl. Neap.* p. xii.—Ins. Caprar.
Tournefortii, *Sieber, Reise Kreta*, i. 239 ; ii. 316.—Creta.
triandra, Ohlib. Fl. Lituan. i. 14 = tinctoria.
trichodes, *Y. Gay, in DC. Prod.* iv. 582.—Persia.
trinervia, Lam. Fl. Fr. iii. 376 = taurina.
tuberosa, Gaudat. ex Steud. Nom. ed. I. 80 ; ed. II. i. 182 = mollugoides.
tyrolica, Bess. Enum. Pl. Volh. 41 = galioides.
umbellata, Willd. ex DC. Prod. iv. 586 = hexaphylla.

ASPHALATHUS, Burm. f. Fl. Ind. 155 ; et Fl. Cap. Prod. 21 (1768) = **Aspalathus**, Linn. (Legum.).

ASPHALTHIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 390 = **Psoralea**, Linn. (Legum.).

ASPHALTHIUM :—
bituminosum, Medic. Phil. Bot. i. 211 = Psoralea bituminosa.
ellipticum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 381 = seq. ?
frutescens, Medic. l. c. 380 = P. bituminosa.
herbaceum, Medic. l. c. 381 = psoraloides.
nigrum, Medic. l. c. = præc. ?
palaestinum, Medic. l. c. = P. palæstina.

ASPHALTIUM, Fourr. in Ann. Soc. Lyon, N. S. xvi. (1868) 365 = ASPHALTHIUM, Medic. = **Psoralea**, Linn. (Legum.).

ASPHODELINE, Reichb. Fl. Germ. Excurs. 116 (1830). *LILIACEÆ*, Benth. & Hook. f. iii. 783.
Dorydium, Salisb. Gen. Pl. Fragm. 72 (1866).
Heroion, Rafin. Fl. Tellur. iv. 13 (1836).
ambigua, *Y. Gay, ex Baker, in Journ. Linn. Soc.* xv. (1877) 276.—Oriens.
Balansæ, *Y. Gay, ex Baker, l. c.* 277.—Cilicia.
brevicaulis, *Y. Gay, ex Baker, l. c.* 276.—Aa. Min. ; Mesopot.
capillaris, *Kunth, Enum. Pl.* iv. 562.—Hab ?
cretica, Boiss. ex Baker, in Journ. Linn. Soc. xv. (1877) 276 = brevicaulis.
cretica, Vis. Fl. Dalm. i. 152 = liburnica.
damascena, *Baker, in Journ. Linn. Soc.* xv. (1877) 276. —Syria.
globifera, *Y. Gay, ex Baker, l. c.* 274.—Syria.
isthmocarpa, *Y. Gay, ex Baker, l. c.* 278.—As. Min.
liburnica, Reichb. Fl. Germ. Excurs. 116.—Europ. austr. or.
lutea, *Reichb. l. c.*—Reg. Mediterr. et Caucas. ; As. Min.
Morisiana, Clem. in Mem. Acc. Torin. Sc. Nat. II. xvi. (1857) 326 = taurica.
parviflora, *Baker, in Journ. Linn. Soc.* xv. (1877) 276. —Byzantium.
prismatocarpa, *Y. Gay, ex Boiss. Fl. Orient.* v. 320.—Cappadoc.
prolifera, *Kunth, Enum. Pl.* iv. 563.—Armenia ; Reg. Cauc.
rigidiflora, *Baker, in Journ. Linn. Soc.* xv. (1877) 277. —As. Min.
Saovitzii, C. Koch, in Linnæa, xxii. (1849) 261 = tenuior.
taurica, *Kunth, Enum. Pl.* iv. 561.—Oriens.
tenuior, *Ledeb. Fl. Ross.* ii. 198.—Oriens ; Reg. Caucas. ; Persia.

ASPHODELOIDES, Moench, Meth. 634 (1794) = **Asphodelus**, Linn. (Liliac.).
ramosa, Moench, l. c. = Asphodelus ramosus.

ASPHODELOPSIS, Stend. ex Baker, in Journ. Linn. Soc. xv. (1876) 321 = Chlorophytum, Ker-Gawl.
arangadiensis, Steud. ex Baker, l. c. 332 = C. attenuatum.

ASPHODULUS, [Tourn.] Linn. Syst. ed. I (1735). *LILIACEÆ*, Benth. & Hook. f. iii. 782.
Asphodeloides, Moench, Meth. 634 (1794).
Claudonia, Pomel, Mat. Fl. Atl. 3 (1874).
Gethosyne, Salisb. Gen. Pl. Fragm. 72 (1866).
Ophioprason, Salisb. l. c. (1866).
Verinea, Pomel, Mat. Fl. Atl. 1 (1874).
abyssinicus, Hort. ex Schult. f. Syst. vii. 600 = Albuca abyssinica.
scanlis, *Desf. Fl. Atlant.* i. 302. t. 89.—Barbaria.
aestivus, Brignat. Stirp. Rar. i. 5 = Asphodelus liburnicus.
aestivus, Brot. Fl. Lusit. i. 525 = microcarpus.
aestivus, Reichb. Fl. Crit. i. 451 = tenuifolius.
affinis, Jord. Fl. Ital. ii. 396 = microcarpus.
africanus, Lam. Encyc. i. 300 = Albuca abyssinica.
africanus, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 730 = ramosus.
albus, Bolss. Voy. Bot. Espagne, 618 = cerasiferus.
albus, Mill. Gard. Dict. ed. VIII. n. 3 = ramosus.
albus, T. Ness, Gen. iv. t. 20 = fistulosus.
albus, *Willd. Sp. Pl.* ii. 133.—Europ. austr.
altaicus, Pall. in Act. Acad. Petrop. (1779) ii. 258. 10 = Eremurus altaicus.
ambiguus, *Jord. in Bull. Soc. Bot. Fr.* vii. (1860) 735. —Gallia.

ASPHODELUS :—
apicacarpus, Hoffmgg. ex Kunth, Enum. Pl. iv. 557 = aestivus.
appropinquatus, Gren. & Godr. Fl. Fr. iii. 238 = fistulosus.
Arrondeaui, Lloyd, Fl. Ouest Fr. ed. III 317 = albus.
asiaticus, *Haw. Syn. Pl. Succ.* 332.—Asia.
Audibertii, Requien, ex Schult. f. Syst. vii. 1695 = ramosus.
australis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 124 = ramosus.
bracteatus, Willd. ex Schult. f. Syst. vii. 484 = Asphodeline lutea.
brevicaulis, Bertol. in Nov. Comm. Acad. Bonon. v. (1842) 430 (Misc. Bot. i. 20) = Asphodeline brevicaulis.
canariensis, C. Sm. & Buch, in Abh. Acad. Berl. (1815–17) 262, 266 = intermedius.
capensis, Burm. f. Fl. Cap. Prod. 10 (= Cyanella ?).—Afr. austr.
capensis, Linn. Syst. ed. X. 982 = Chlorophytum elatum.
capillaris, [DC. in] Red. Lil. t. 380 = Asphodeline liburnica.
cerasiferus, *Y. Gay, in Bull. Soc. Bot. Fr.* iv. (1857) 610 ; *et in Ann. Sc. Nat.* Sér. IV. vii. (1857) 197.—Reg. Mediterr. occ.
Chamaemoly, Hochst. ex Baker, in Journ. Linn. Soc. xv. (1877) 328 = Chlorophytum tetraphyllum.
Chambertoni, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 736 = ramosus.
clavatus, Roxb. Hort. Beng. 24 ; Fl. Ind. ii. 148 = tenuifolius.
clavosus, Don, ex Steud. Nom. ed. II. i. 252, sphalm. = clavatus.
collinus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 123 = ramosus.
comosus, Houtt. Plantenk. xii. 386. t. 83 = Eucomis punctata.
comosus, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 737 = ramosus.
cornicus, Jord.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 739 = ramosus.
creticus, Lam. Encyc. i. 300 = Asphodeline liburnica.
crinifex, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 729 = albus.
damascenus, Boiss. Diagn. Ser. I. xiii. 22 = Asphodeline damascena.
delphinensis, Gren. & Godr. in Mém. Soc. Emul. Doubs, Sér. II. vi. (1855) 18 = albus.
Dioscuri, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 121 = ramosus.
fistulosus, *Linn. Sp. Pl.* 309.—Reg. Mediterr. ; Oriens.
fuscatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 128 = ramosus.
glaucus, Ascher, ex Boiss. Fl. Orient. v. 328 = Eremurus cappadocicus.
glaucescens, Jord.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 739 = ramosus.
gramineus, Mill. ex Steud. Nom. ed. I. 80, 278 ; ed. II. i. 152 = Melasphaerula graminea.
grandiflorus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 128 = ramosus.
inderiensis, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 257 = Eremurus inderiensis.
infestus, Parl. Fl. Ital. ii. 600 = microcarpus.
intermedius, Hornem. Hort. Hafn. Suppl. 131 = fistulosus.
latifolius, Breyn. ex Schult. f. Syst. vii. 486 = Wachendorfia paniculata.
liburnicus, Scop. Fl. Carn. ed. II. i. 345 = Asphodeline liburnica.
littoralis, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 729 = albus.
luteus, Linn. Sp. Pl. 309 = Asphodeline lutea.
macrocarpus, Parl. Fl. Ital. ii. 604 = albus.
major, Four. ex Willk. & Lange, Prod. Fl. Hisp. i. 504 = albus.
microanthus, Cosa. & Kralik, ex Cosa. in Bull. Soc. Bot. Fr. iv. (1857) 497 = viscidulus.
microcarpus, Reichb. Fl. Germ. Excurs. 117 ; Ic. Fl. Germ. ix. 30. n. 1116 = tenuifolius.
microcarpus, *Viviani, Fl. Cors.* 5.—Reg. Mediterr. ; Ins. Canar.
Moriseanus, *Parl. Fl. Ital.* ii. 596.—Italia.
neglectus, Schult. f. Syst. vii. 488 = albus.
nervosus, *Pomel, Nouv. Mat. Fl. Atl.* 385.—Afr. bor.
occidentalis, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 731 = albus.

ASPHODELUS:—
albiensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 122 = ramosus.
Cranensis, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 733 = ramosus.
parviflorus, Wight, Ic. t. 2062 = fistulosus.
pendulinus, *Cav. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 396, 497.—Algeria; Arab.
persicus, Fisch. & Mey. ex Boiss. Fl. Orient. v. 317 = Asphodeline tenuior.
persicus, Jaub. & Spach, Illustr. ii. 2. t. 109 = Eremurus persicus.
pratensis, Fourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 308 = microcarpus.
procerus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 124 = ramosus.
*prolifer*us, Bieb. Fl. Taur. Cauc. i. 280 = Asphodeline prolifera.
proterophyllus, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 737 = ramosus.
pyrenaicus, Jord. l. c. 732 = albus.
racemosus, Link, Enum. Hort. Berol. i. 238 = ramosus.
ramosus, Colm. ex Willk. & Lange, Prod. Fl. Hisp. i. 303 = microcarpus.
ramosus, Gouan, ex Willk. & Lange, l. c. 304 = cerasiferus.
ramosus, Lapeyr. Hist. Abr. Pl. Pyr. 188 = neglectus.
ramosus, *Linn. Sp. Pl.* ed. 1. 310.—Europ. austr.
ramosus, Linn. l. c. ed. II. 444 = microcarpus, cerasiferus.
ramosus, Murr. in Nov. Comm. Gotting. (1776) 37. t. 71 = albus.
ramosus, Sturm, Deutschl. Fl. Heft vi. t. 6 = fistulosus.
refractus, Boiss. Diagn. Ser. I. xiii. (1854) 23 = pendulinus.
regius, Hort. ex Heynh. Nom. ii. 48 = Eremurus spectabilis.
repens, *Pomel, Nouv. Mat. Fl. Atl.* 256.—Afr. bor.
rigidifolius, Boiss. & Heldr. Diagn. Ser. I. xiii. 22 = Asphodeline rigidifolia.
Rouxii, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 738 = ramosus.
serotinus, Sibth. ex Jacq. Eclog. 118 = Asphodeline liburnica.
sibiricus, Hort. ex Schult. f. Syst. vii. 483 = Asphodeline lutea.
sibiricus, Sievers, in Pall. N. Nord. Beitr. vii. 257 = Eremurus altaicus.
simplex, Pers. Syn. i. 367 = albus.
sphaerocarpus, Gren. & Godr. Fl. Fr. iii. 223 = albus.
spicatus, Desf. ex Baker, in Journ. Linn. Soc. xv. (1877) 371 = ramosus.
stenchadensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 122 = ramosus.
strictus, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
subalpinus, Gren. & Godr. Fl. Fr. iii. 224 = albus.
Sawitzii, C. Koch, in Linnæa, xxii. (1849) 261 = Asphodeline tenuior.
tardiflorus, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 736 = ramosus.
tauricus, Lodd. Bot. Cab. t. 1102 = Asphodeline tenuior.
tauricus, Pall. in Nov. Act. Petrop. x. 309; Bieb. Fl. Taur. Cauc. i. 279 = Asphodeline taurica.
tenuifolius, C. Koch, in Linnæa, xxii. (1849) 261 = Asphodeline tenuior.
tenuifolius, *Cav. in Anal. Cienc. Nat.* iii. (1801) 46.—Ind. or.
tenuior, Fisch. Cat. Hort. Gorenk. (1812) 9 = Asphodeline tenuior.
validus, Jord.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 740 = ramosus.
Villarsii, Verl. ex Jord. Billotia, 107, 109; et in Bull. Soc. Stat. Isère, Sér. II. iv. (1860) 151 = ramosus.
virgatus, Jord.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 738 = ramosus.
visciculus, *Boiss. Diagn.* Ser. I. vii. 136.—Afr. bor.; Arab.

ASPICARIA, D. Dietr. Syn. i. 2, 27 (1839) = seq.

ASPICARPA, Rich. in *Mém. Mus. Par.* ii. (1815) 398. t. 1; Lag. Gen. et Sp. Nov. 1 (1816). *MALPIGHIACEÆ*, Benth. & Hook. f. i. 361.
ACOSMIUM, Desv. in Desf. Cat. Hort. Par. (1829) 224.
Hartwegiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 506.—Mexic.

ASPICARPA:—
hirtella, *Rich. in Mém. Mus. Par.* ii. (1815) 399. t. 13.—Mexic.
hysopifolia, *A. Gray*, in Bost. Journ. Nat. Hist. vi. (1850) 167.—Texas.
longipes, *A. Gray, Pl. Wright.* i. 37.—N. Mexic.
sericea, *Griseb. in Goett. Abh.* xxiv. (1879) 68.—Reg. Argent.
urens, Lag. Gen. et Sp. Nov. i. 1. n. 5 = hirtella.

ASPIDALIS, Gaertn. Fruct. ii. t. 171 (1791) = **Berkheya**, Ehrh. (Compos.).
araneosa, Gaertn. l. c. = B. cernua.

ASPIDANDRA, Hassk. in Verh. en Med. Akad. Wetensch. iv. (1855) 141. *EUPHORBIACEÆ*? Gen. dub.'
fragrans, *Hassk. l. c.*—Java.

ASPIDANTHERA, Benth. in Hook. Journ. Bot. iii. (1841) 317 = **Perdinandusa**, Pohl (Rubiac.).
rudgeoides, Benth. l. c.—Bras.

ASPIDELIUM, Zollik. ex DC. Prod. vii. 150 (1838) = **Chondrilla**, Tourn. (Compos.).

ASPIDISTRA, Ker-Gawl. in Bot. Reg. t. 628 (1823). *LILIACEÆ*, Benth. & Hook. f. iii. 773.
MACROGYNE, Link & Otto, Ic. Pl. Sel. t. 31 (1828).
PLECTOGYNE, Link, in Otto & Dietr. Allg. Gartenz. (1836) 266.
PORPAX, Salisb. Gen. Pl. Fragm. 9 (1866).
elatior, *Blume, in Hoev. & De Vriese, Tijdschr.* i. (1834) 76. t. 4.—Japon.
lurida, *Ker-Gawl. in Bot. Reg.* t. 628.—China.
lurida, Sieber, ex Presl, in Flora, xii. (1829) 568 = Weschmoria hirsuta.
punctata, Lindl. in Bot. Reg. t. 977 = lurida.
variegata, Regel, in Flora, xxxiii. (1850) 354 = elatior.

ASPIDOCARPUS, Neck. Elem. ii. 123 (1790) = **Rhamnus**, Tourn.

ASPIDOCARYA, Hook. f. & Thoms. Fl. Ind. i. 180 (1855). *MENISPERMACEÆ*, Benth. & Hook. f. i. 33.
hirsuta, *Bucc. Malesia*, i. (1877) 136.—Malaya.
uvifera, *Hook. f. & Thoms. Fl. Ind.* i. 180.—Reg. or.

ASPIDOGLOSSUM, E. Mey. Comm. Pl. Afr. Austr. 200 (1837) = **Schizoglossum**, E. Mey. (Asclep.).
bifarium, E. Mey. l. c. 201.—Afr. austr.
fasciculare, E. Mey. l. c. 200.—Afr. austr.
heterophyllum, E. Mey. l. c.—Afr. austr.

ASPIDOPTERYS, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 366. *MALPIGHIACEÆ*, Benth. & Hook. f. i. 369.
canarensis, *Dalz. in Hook. Kew Journ.* iii. (1851) 37. —Ind. or.
concolor, *A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 366.—Burma.
cordata, *A. Juss. l. c.* 367.—Ind. or.
elliptica, *A. Juss. l. c.* 366.—Ind. or.
glabrescens, A. Juss. l. c. 367 = Roxburghiana.
glomerata, Wight, Ic. t. 1986 = canarensis.
Heiferiana, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) 11. 137.—Burma.
hirsuta, *A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 367.—Burma.
lanuginosa, A. Juss. l. c. = nutans.
nutans, A. Juss. l. c. = Wallichii.
nutans, *Hook. f. Fl. Brit. Ind.* i. 421.—Reg. Himal.
oxyphylla, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 367 = Roxburghiana.
retundifolia, A. Juss. l. c. = nutans.
Roxburghiana, *A. Juss. l. c.*—Ind. or.
tomentosa, *A. Juss. l. c.*—Java.
Wallichii, *Hook. f. Fl. Brit. Ind.* i. 421.—Reg. Himal.

ASPIDOSPERMA, Mart. & Zucc. Nov. Gen. et Sp. i. 27. t. 34 ad 36 (1824). *APOCYNACEÆ*, Benth. & Hook. f. ii. 703.
MACAGLIA, Rich. ex Vahl, in Skrivt. Nat. Selsk. Kiøbenh. vi. (1810) 107.
PELTOSTEMMUM, DC. in Bibl. Univ. Genèv. xvii. (1838) 153.

ASPIDOSPERMA:—
anonalum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 61.—Bras.
aquaticum, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 204.—Bras.
australe, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 88.—Bras.
bicolor, *Mart. Nov. Gen. et Sp.* i. 60.—Bras.
camporum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 49.—Bras.
condylocarpon, *Muell. Arg. l. c.* 55.—Bras.
cylindrocarpum, *Muell. Arg. l. c.* 54.—Bras.
dasycarpon, *A. DC. in DC. Prod.* viii. 396.—Bras.
decipiens, *Muell. Arg. in Linnæa*, xxx. (1859–60) 398. —Venezuela.
desmanthum, *Benth. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 59.—Bras.
discolor, *A. DC. in DC. Prod.* viii. 398.—Bras.
dispermum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 60.—Bras.
eburneum, *Allem. ex Saldanha, in Ann. Sc. Nat.* Sér. V. xix. (1874) 218.—Bras.
excelsum, *Benth. in Hook. Journ. Bot.* iii. (1841) 345.—Guiana.
Francisii, *A. DC. in DC. Prod.* viii. 398.—Guiana.
Gardneri, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 47.—Bras.
Gomezianum, *A. DC. in DC. Prod.* viii. 397.—Bras.
Hilarianum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 50.—Bras.
lagotoxe, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 102.—Bras.
latisiliquo, *A. DC. in DC. Prod.* viii. 676.—Guiana.
leucomelanum, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 105.—Bras.
Lhotskianum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 60. —Bras.
macrocarpon, *Mart. Nov. Gen. et Sp.* i. 59.—Bras.
macrophyllum, *Muell. Arg. in Linnæa*, xxx. (1859–60) 397.—Guiana.
Martii, *Silva Manso, in Mart. Herb. Fl. Bras.* 163.—Bras.
megalocarpon, *Muell. Arg. in Linnæa*, xxx. (1859–60) 400.—Mexic.
melanocalyx, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 52. —Bras.
mollis, Mart. Herb. Fl. Bras. 163 = pyrifolium.
multiflorum, *A. DC. in DC. Prod.* viii. 397.—Guiana.
nitidum, *Benth. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 59.—Bras.
noble, *Muell. Arg. l. c.* 44.—Bras.
oblongum, *A. DC. in DC. Prod.* viii. 399.—Guiana.
obscurum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 48.—Bras.
olivaceum, *Muell. Arg. l. c.* 57.—Bras.
pachypterum, *Muell. Arg. l. c.* 51.—Bras.
pallidiflorum, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 101.—Bras.
parvifolium, *A. DC. in DC. Prod.* viii. 398.—Bras.
platyphyllum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 46.—Bras.
polyneuron, *Muell. Arg. l. c.* 57.—Bras.
Pohlianum, *Muell. Arg. l. c.* 44.—Bras.
populifolium, *A. DC. in DC. Prod.* viii. 397.—Bras.
pyricollum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 58.—Bras.
pyrifolium, *Mart. Nov. Gen. et Sp.* i. 60.—Bras.
Quebracho-blanco, *Schlechtd. in Bot. Zeit.* xix. (1861) 137.—Arn. austr.
Quebracho-colorado, *Schlechtd. l. c.* 139.—Arn. austr.
ramiflorum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 55.—Bras.
reflectum, *Mart. Nov. Gen. et Sp.* i. 60.—Bras.
Riedelii, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 56.—Bras.
Sellowii, *Muell. Arg. l. c.*—Bras.
sessiliflorum, *Muell. Arg. in Linnæa*, xxx. (1859–60) 399.—Ins. Trinit.
Spruceanum, *Benth. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 62.—Bras.
subincanum, *Mart. Herb. Fl. Bras.* n. 162; *ex A. DC. in DC. Prod.* viii. 397.—Bras.
tomentosum, *Mart. Nov. Gen. et Sp.* i. 58. t. 34.—Bras.
Vauguoli, *A. DC. in DC. Prod.* viii. 399.—Venezuela.
velutinum, Fisch. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 46 = tomentosum.
verbenifolium, *Muell. Arg. l. c.* 46.—Bras.

ASPIDOSPERMA:—
venosum, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 106.—Bras.
verrucicaule, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 51.—Bras.
Warmingii, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 101.—Bras.

ASPIDOSTIGMA, Hochst. in Flora, xxvii. (1844) 18 = **Toddalia**, Juss. (Rutac.).
acuminatum, Hochst. l. c. = T. noblis.

ASPILIA, Thou. Gen. Nov. Madag. 12 (1806). *COMPOSITAE*, Benth. & Hook. f. ii. 371.
ANOMOSTEPHIUM, DC. Prod. v. 560 (1836).
CORONOCARPUS, Schum. & Thonn. Beskr. Guin. Pl. 893 (1827).
DIPTEROTHECA, Sch. Bip. in Flora, xxv. (1842) 434.
HARPEPHORA, Endl. Gen. Suppl. i. 1382 (1841).
WURTGENIA, Sch. Bip. in Flora, xxv. (1842) 435.
abyssinica, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 379.—Afr. trop.
abyssinica, *Vatke, in Linnaea*, xxxix. (1875) 495.—Abyss.
angustifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 380.—Afr. trop.
anomala, *Baker, in Mart. Fl. Bras.* vi. III. 204.—Bras.
aristata, *Griseb. in Goett. Abh.* xxiv. (1879) 191.—Reg. Argent.
asperrima, *Baker, in Mart. Fl. Bras.* vi. iii. 200.—Bras.
atimnata, *Baker, l. c.* 190.—Bras.
aurantiaca, *Griseb. in Goett. Abh.* xix. (1874) 188.—Reg. Argent.
Barozzi, *Baker, in Journ. Linn. Soc.* xx. (1884) 188.—Madag.
Bojeri, *DC. Prod.* v. 561.—Madag.
buphthalmiflora, *Griseb. in Goett. Abh.* xix. (1874) 186.—Reg. Argent.
Burchellii, *Baker, in Mart. Fl. Bras.* vi. III. 194.—Bras.
calendulacea, Griseb. in Goett. Abh. xxiv. (1879) 191 = buphthalmiflora.
Clausseniana, *Baker, in Mart. Fl. Bras.* vi. III. 202.—Bras.
ecliptaefolia, *Baker, l. c.* 197.—Bras.
elegans, *Benth. & Hook. f. Gen.* ii. 372.—Bras.
elliptica, *Baker, in Mart. Fl. Bras.* vi. III. 192.—Bras.
floribunda, *Baker, l. c.* 198.—Bras.
foliacea, *Baker, l. c.* 198.—Bras.
foliosa, *Benth. & Hook. f. Gen.* ii. 372.—Bras.
fruticosa, *Baker, in Mart. Fl. Bras.* vi. III. 204.—Bras. Guyana, *Benth. & Hook. f. Gen.* ii. 372.—Afr. trop.
glabra, *Benth. & Hook. f. l. c.*—Bras.
gracilis, *Baker, in Mart. Fl. Bras.* vi. III. 198.—Bras.
helianthoides, *Benth. & Hook. f. Gen.* ii. 372.—Afr. trop.
hirsuta, *Benth. & Hook. f. l. c.*—Bras.
hispidula, *Baker, in Mart. Fl. Bras.* vi. III. 201.—Bras.
Kotschyana, *Benth. & Hook. f. Gen.* ii. 372.—Afr. trop.
laevisiana, *Baker, in Mart. Fl. Bras.* vi. III. 202.—Bras.
latifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 379.—Afr. trop.
linearifolia, *Baker, in Mart. Fl. Bras.* vi. III. 192.—Bras.
linearifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 380.—Afr. trop.
Martii, *Baker, in Mart. Fl. Bras.* vi. III. 195.—Bras.
mexicana, *Benth. & Hook. f. Gen.* ii. 372.—Mexic.
multiflora, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 378.—Afr. trop.
oblonga, *Baker, in Mart. Fl. Bras.* vi. III. 198.—Bras.
ovalifolia, *Baker, l. c.* 199.—Bras.
paucisetosa, *Griseb. in Goett. Abh.* xxiv. (1879) 191.—Reg. Argent.
phyllostachya, *Baker, in Mart. Fl. Bras.* vi. III. 201.—Bras.
podophylla, *Baker, l. c.* 200.—Bras.
Pohlii, *Baker, l. c.* 201.—Bras.
Princeana, *Benth. & Hook. f. Gen.* ii. 372.—Afr. trop.
procumbens, *Baker, in Mart. Fl. Bras.* vi. III. 194.—Bras.
psadia, *Baker, l. c.* 193.—Bras.
rufens, *Baker, l. c.* 198.—Bras.
salicifolia, *Baker, l. c.* 202.—Bras.
Riedelii, *Baker, l. c.* 196.—Bras.

ASPILIA :—
rudis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 380.—Afr. trop.
Schimperi, *Oliver & Hiern, l. c.* 379.—Afr. trop.
serrulata, *Baker, in Mart. Fl. Bras.* vi. III. 204.—Bras.
setosa, *Griseb. in Goett. Abh.* xxiv. (1879) 192.—Reg. Argent.
silphioides, *Benth. & Hook. f. Gen.* ii. 372.—Bras.
Smithiana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 380.—Afr. trop.
squarrosa, *Baker, in Mart. Fl. Bras.* vi. III. 203.—Bras.
strigosa, *Benth. & Hook. f. Gen.* ii. 372.—Mexic.
subalpestris, *Baker, in Mart. Fl. Bras.* vi. III. 202.—Bras.
subpetiolata, *Baker, l. c.* 203.—Bras.
Thouarii, *DC. Prod.* v. 561.—Madag.
tomentosa, *Baker, in Mart. Fl. Bras.* vi. III. 199.—Bras.
Warmingii, *Baker, l. c.* 192.—Bras.
weddelianformis, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1877) 197.—Afr. trop.

ASPILOBIUM, Soland. ex A. Cunn. in Ann. Nat. Hist. (Sept. 1838) 45 = **Geniostoma**, Forst. (Logan.).

ASPILOTUM, S[oland.]ex Steud. Nom. ed. II. i. 669, 152 (1840) = **Geniostoma**, Forst. (Logan.).
laevigatum, S[oland.] ex Steud. l. c. = G. ligustrifolium.

ASPITTUM, Neck. ex Steud. Nom. ed. II. ii. 10 (1841) = **Laserpitium**, Linn. (Umbell.).

ASPLA, Reichb. Nom. 50 (1841) err. typ. = AOPLA, Lindl. = **Habenaria**, Linn. (Orchid.).

ASPRELLA, Host, Gram. Austr. iv. 17. t. 29 (1809) = **Paliurus**, Trin. (Gram.).
nardiformis, Host, l. c. = P. nardoides.

ASPRELLA, Willd. Enum. Hort. Berol. 132 (1809) = **Aspris**, Willd. (Humb. (Gram.).

ASPRELLA, Schreb. Gen. 45 (1789) = **Leersia**, Sw. (Gram.).
australis, Roem. & Schult. Syst. ii. 267 = L. hexandra.
brasiliensis, Koem.& Schult. Mant. 153 = L. brasiliensis.
digitata, Lam. Illustr. i. 187 = L. digitata.
echinea, Rafin. in Am. Monthly Mag. (1819) 190.—Am. bor.
flexuosa, Dulac, Fl. Hautes-Pyr. 71 = L. oryzoides.
hexandra, Roem. & Schult. Mant. 551 = L. hexandra.
Hystrix, Link, Enum. Hort. Berol. 100 = Asperella Hystrix.
imbricata, Roem. & Schult. Syst. ii. 268 = L. virginica.
lenticularis, Roem. & Schult. l. c. 267 = L. lenticularis.
major, Fres. ex Steud. Nom. ed. II. i. 150 = Asperella Hystrix.
mexicana, Roem. & Schult. Mant. 551 = L. hexandra.
monandra, Roem. & Schult. l. c. = L. monandra.
oryzoides, Lam. Illustr. i. 167 ; Beauv. Agrost. 9. t. 4. f. 9 = L. oryzoides.
ovata, Roem. & Schult. Syst. ii. 267 = L. lenticularis.
purpurea, Boj. Hort. Maunt. 276 = L. hexandra.
sibirica, Trauv. in Act. Hort. Petrop. v. (1877) 182.—Siberia.
virginica, Roem. & Schult. Syst. ii. 268 = L. virginica.

ASPRIS, Adans. Fam. ii. 496 (1763) = **Aira**, Linn. (Gramin.) (= Aira praecox).

ASSA, Houtt. Plantenk. v. 275 (1776) = **Tetracera**, Linn. (Dillen.).
exotica, F. G. Gmel. Syst. 539 = T. Assa.
indica, [Christm. & Panz. in] Houtt. Pflanz. Syst. iv. 40 = T. Assa.

ASSARACUS, Haw. Monog. Narcias. 4 ; in Sweet, Brit. Flow. Gard. iv. (1838) = **Narcissus**, Linn. (Amaryll.).
capax, Haw. l. c. = N. triandrus.
reflexus, Haw. l. c. = N. triandrus.

ASSONIA, Cav. Diss. iii. 120. t. 42 (1786) = **Dombeya**, Cav. (Stercul.).
Erythroxylon, Sims, Bot. Mag. sub t. 1000 = Trochetia Erythroxylon.
populnea, Cav. Diss. iii. 120. t. 42. f. 3 = D. populnea.
silvaticoides, DC. Prod. i. 498.—Ins. Borbon.

ASTARTEA, DC. in Dict. Class. xi. 400 (1826) ; Prod. ii. 210 (1828). *MYRTACEAE*, Benth. & Hook. f. i. 702.
ambigua, *F. Muell. Fragm.* ii. 32.—Austral.
aspera, Schau. in Linnaea, xvii. (1843) 242 = fasciularis.
clavulata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 338.—Austral.
corniculata, Schau. in Lehm. Pl. Preiss. i. 113 = fasciularis.
Endlicheriana, Schau. in Linnaea, xvii. (1843) 242 = fasciularis.
fasciularis, *DC. Prod.* iii. 210.—Austral.
glomerulosa, Schau. in Linnaea, xvii. (1843) 242 = fasciularis.
intratropica, *F. Muell. Fragm* i. 83.—Austral.
laricifolia, Schau. in Linnaea, xvii. (1843) 242 = fasciularis.
leptophylla, Schau. l. c. = fasciularis.
muricata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 334.—Austral.
scoparia, Schau. in Linnaea, xvii. (1843) 242 = fasciularis.

ASTELIA, Banks & Soland. ex R. Br. Prod. 291 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 781.
FUNCKIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 19.
HAMELINIA, A. Rich. Voy. Astrolob. ; Fl. Nouv. Zél. 158. t. 24 (1832).
albicans, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 252.—N. Zel.
alpina, *R. Br. Prod.* 291.—Austral.
Banksii, *A. Cunn. in Hook. Comp. Bot. Mag.* ii. (1836) 374.—N. Zel.
Cunninghamii, *Hook. f. Fl. N. Zel.* i. 259.—N. Zel.
fragrans, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 333.—N. Zel.
grandis, *Hook. f. ex T. Kirk, in Trans. N. Z. Inst.* i. (1868) 93.—N. Zel.
Hookeriana, T. Kirk, l. c. iv. (1872) 244 = Cunninghamii.
linearis, *Hook. f. Fl. Antarct.* 76.—Ins. Antarct.
Menziesiana, *Sm. in Rees, Cycl.* xxxix. *App.* ; *Schult. f. Syst.* vii. 1506.—Ins. Sandvic.
microperma, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 251.—N. Zel.
montana, *Seem. Fl. Vit.* 313.—Ins. Fiji.
nervosa, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 260.—N. Zel.
polyneuron, *Colenso, in Trans. N. Z. Inst.* xiv. (1882) 223.—N. Zel.
psychrocharis, F. Muell. in Hook. Kew Journ. viii. (1856) 332 = alpina.
pumila, *R. Br. Prod.* 291.—Terr. Magell.
Ricardi, Kunth, Enum. Pl. iii. 365 = Banksii.
sericea, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 261 = Banksii.
Solandri, *A. Cunn. in Hook. Comp. Bot. Mag.* ii. (1836) 374.—N. Zel.
spicata, *Colenso, in Trans. N. Z. Inst.* xiv. (1882) 335.—N. Zel.
stylosa, F. Muell. in Hook. f. Fl. Tasm. ii. 61 = Milligania stylosa.
trinervia, *T. Kirk, in Trans. N. Z. Inst.* i. (1868) 93.—N. Zel.
vernalvides, *Gaudich. in Freyc. Voy. Bot.* 490. t. 81.—Ins. Sandvic.
Waialealia, *Wawra, in Flora*, lviii. (1875) 243.—Ins. Sandvic.

ASTELMA, R. Br. in Bot. Reg. sub t. 532 (1820) = **Helichrysum**, Vaill. et **Helipterum**, DC. (Compos.).
canescens, D. Don, in Mem. Wern. Soc. v. (1824) 545 = Helipterum canescens.
crassifolium, D. Don, l. c. 543 = Helipterum ferrugineum.
eximium, R. Br. in Bot. Reg. sub t. 532 = Helipterum eximium.
fragrans, Sweet, Brit. Brit. ed. I. 222 = Helichrysum fragrans.
fruticans, Ker-Gawl. Bot. Reg. t. 726 = Helichrysum fruticans.
imbricatum, D. Don, in Mem. Wern. Soc. v. (1824) 544 = Helichrysum imbricatum.
linare, D. Don, l. c. = Helichrysum ericaefolium.
millefloroum, D. Don, l. c. 543 = Helipterum phlomoides.

ASTELMA;—
modestum, Sieber, ex Hook. Bot. Mag. t. 2710 = Helipterum gnaphaloides.
protensis, D. Don, in Mem. Wern. Soc. v. (1824) 548. —Afr. austr.
retortum, D. Don, ex Steud. Nom. ed. II. i. 152 = Helichrysum retortum.
spectantissimum, D. Don, in Mem. Soc. Wern. v. (1824) 544 = Helipterum speciosissimum.
spirale, G. Don, in Sweet, Hort. Brit. ed. III. 378 = Helipterum variegatum.
Stachelina, D. Don, in Mem. Wern. Soc. v. (1824) 543 = Helipterum speciosissimum.
variegatum, D. Don, l. c. 544 = Helipterum variegatum.
virgatum, G. Don, in Sweet, Hort. Brit. ed. III. 378 = Helipterum virgatum.

ASTEMMA, Less. Syn. Comp. 216 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 356.
dubium, *Less. l. c.*—Ecuador.

ASTEMON, Regel, Ind. Sem. Hort. Petrop. (1860) 38 = Bystropogon, L'Hérit. (Labiat.).
graveolens, Regel, l. c. (*an* Hyptis suaveolens ?).—Bolivia.

ASTEPHANANTHES, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 138 = Passiflora, Linn.
bilobata, Bory, l. c. = P. bilobata.

ASTEPHANUS, R. Br. in Mem. Wern. Soc. i. (1809) 54. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 747.
HARMAN, E. May. Comm. Pl. Afr. Austr. 233 (1887).
TYLODONTIA, Griseb. Cat. Pl. Cub. 175 (1866).
arenarius, *Decne. in DC. Prod.* viii. 507.—Madag.
badius, *E. May. Comm. Pl. Afr. Austr.* 294.—Afr. austr.
Berterii, *Spreng. Syst.* i. 855.—N. Granat.
cernuus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 343.—Ins. Pemba.
cordatus, *R. Br. in Mem. Wern. Soc.* i. (1809) 54.—Afr. austr.
cordifolius, *Phil. in Anal. Univ. Chil.* ii. (1862) 399; *et Linnaea*, xxxiii. (1864-65) 178.—Chili.
cubensis, *H. B. & K. Nov. Gen. et Sp.* iii. 206. t. 237. —Cuba.
Dregei, *D. Dietr. Syn. Pl.* ii. 909.—Afr. austr.
dumosus, D. Dietr. l. c. = Rhyssolobium dumosum.
fruticosus, *E. May. Comm. Pl. Afr. Austr.* 223.—Afr. austr.
fruticosus, Steud. Nom. ed. II. i. 153 = frutescens.
Gardneri, *Fourn. in Mart. Fl. Bras.* vi. iv. 199.—Bras.
geminiflorus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 342.—Chili.
lanceolatus, *R. Br. in Mem. Wern. Soc.* i. (1809) 54. —Afr. austr.
linearis, *R. Br. l. c.*—Afr. austr.
marginatus, *Decne. in DC. Prod.* viii. 508.—Afr. austr.
Masoni, *Korn. & Schuit. Syst.* vi. 124.—Afr. austr.
mitophorus, *Griseb. in Gott. Abh.* xix. (1874) 205.—Reg. Argent.
mucronatus, *Schlecht. in Linnaea*, viii. (1833) 518.—Mexic.
nigrescens, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1881) 367.—Mexic.
ovalifolius, *A. Rich. Fl Cub. Fanerog.* ii. 95.—Cuba.
ovatus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 342.—Madag.
pauciflorus, *E. May. Comm. Pl. Afr. Austr.* 224.—Afr. austr.
peruvianus, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 367.—Peruv.
recurvatus, *Klotzsch, in Peters, Reise Mossamb. Bot.* 274.—Afr. trop.
Schimperi, *Vatke, in Oestr. Bot. Zeitschr.* xxvi. (1876) 146.—Afr. trop.
streptocoryne, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 367.—Paraguay.
trifiorus, *R. Br. in Mem. Wern. Soc.* i. (1809) 54.—Afr. austr.
urceolatus, *Griseb. Cat. Pl. Cub.* 173.—Cuba.
utahensis, *Engelm. in Am. Natural.* ix. (1875) 549.—Am. bor. occ.
Zeyheri, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 314.—Afr. austr.

ASTER, Tourn. ex Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 271.
ACTIPSIS, AGLOTOMA, ANACTIS, DIPLACTIS, FIMBRISTIMA, GALACTELLA (GALASTELLA), KALIMARIS, LEIACHENIS, MESOLIGUS, TRIPOLIUM, VENATRIS, VIRGULUS, Rafin. Fl. Tellur. ii. 48–47 (1836).
ARCTOGERON, DC. Prod. v. 260 (1836).
BELLIDIASTRUM, Cass. in Dict. Sc. Nat. xxxvii. 458 (1825).
BIOTIA, DC. Prod. v. 264 (1836).
CALIMERIS, Nees, Gen. et Sp. Aster. 225 (1832).
CRINITA, Moench, Meth. 578 (1794).
CRINITARIA, Cass. in Dict. Sc. Nat. xxxvii. 475 (1825).
DIETERIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 300.
DIPLOPAPPUS, Cass. in Bull. Soc. Philom. (1817) 137; et in Dict. Sc. Nat. xiii. 308 (1819).
DOELLINGERIA, Nees, Gen. et Sp. Aster. 177 (1832).
EUCEPHALUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 298.
EURYBIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 464 (1821).
EXATHAMNUS, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 46.
GALATEA, Cass. in Bull. Soc. Philom. (1818) 165; et in Dict. Sc. Nat. xviii. 56 (1820).
HELEASTRUM, DC. Prod. v. 263 (1836).
HERSILEA, Klotzsch, Bot. Ergeb. Waldem. Reise Bot. 53 (1862).
HETEROCHAETA, DC. Prod. v. 289 (1836).
HOMOSTYLIUM, Nees, in Linnaea, xviii. (1844) 513.
KALIMERIS, Cass. in Dict. Sc. Nat. xxxvii. 464 (1825).
LINOSYRIS, Cass. l. c. 460, 476 (1825).
MACHAERANTHERA, Nees, Gen. et Sp. Aster. 224 (1832).
MARGARITA, Gaud. Fl. Helv. v. 335 (1829).
NOTICASTRUM, DC. Prod. v. 279 (1836).
PINARIA, Neck. Elem. i. 5 (1790).
PSYCHROGETON, Boiss. Fl. Orient. iii. 156 (1875).
RHINACTINA, Less. in Linnaea, vi. (1831) 119.
SYMPHYOTRICHUM, Nees, Gen. et Sp. Aster. 135 (1832).
TRIPOLIUM, Nees, l. c. 152 (1832).
TURCZANINOWIA, DC. Prod. v. 257 (1836).
VIRGAUREA, Rafin. ex DC. Prod. v. 248 (1836).
XYLORHIZA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 297.
abbreviatus, Nees, Syn. Aster. 16 = patulus.
abyssinicus, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 333 = Felicia Richardi.
acaulis, *Wedd. Chlor. And.* i. 189.—Peruv.
Acmella, Pall. Reise, iii. 320 = A. Acmella.
acris, S. G. Gmel. It. i. 173. t. 85. f. 2 = dracunculoides.
acris, Lapeyr. Hist. Abr. Pl. Pyr. 518 = salignus.
acris, Leyss. Fl. Halens. 216 = A. Amellus.
acris, *Linn. Sp. Pl.* ed. II. 1238.—Europ. austr.; Caucas.
acris, Willd. Sp. Pl. iii. 2023 = trinervis.
acuductus, Labill. Nov. Holl. Pl. ii. 52. t. 200 = Olearia ramulosa.
acuminatus, *Michx. Fl. Bor. Am.* ii. 109.—Am. bor.
acuminatus, Nees, Gen. et Sp. Aster. 60 = patulus.
adenolaenius, F. Muell. Fragm. v. 67 = Olearia adenolaenia.
adenophorus, F. Muell. l. c. 78 = Olearia adenophora.
adfinis, Less. Syn. Comp. 174 = Felicia affinis.
adnatus, *Nutt. in Journ. Acad. Philad.* vii. (1834) 89. —Am. bor.
adscendens, *Lindl. in DC. Prod.* v. 231.—Am. bor.
adscitivus, Willd. Enum. Hort. Berol. ii. 884 = Novi-Belgii.
aestivus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 208 = longifolius.
aestivus, A. Gray, Man. Bot. U. St. ed. V. 358 partim = junceus.
aestivus, Rothrock, in Wheeler, Rep. 264 = occidentalis.
aethiopicus, *Burm. f. Fl. Cap. Prod.* 27.—Afr. austr.
agrestoides, Turcz. in Bull. Soc. Nat. Mosc. (1847) n. vii. 194 = trinervius.
Altonii, *Forbes, Hort. Woburn.* 188.—Hab. ?
alatus, Aikin, in Eaton & J. Wright, Man. Bot. 146 = patens.
albescens, Wall. Cat. n. 2974 = Microglossa albescens.

ASTER, *continued*—
albus, Eaton & J. Wright, Man. Bot. 147 = ptarmicoides.
albus, Willd. ex Spreng. Syst. iii. 558 = A. Amellus.
alpicola, F. Muell. Fragm. v. 70 = Olearia alpicola.
alpinus, A. Gray, in Proc. Am. Acad. vii. (1872) 389 = pulchellus.
alpinus, Gueldenst. ex Ledeb. Fl. Ross. ii. 476 = A. Amellus.
alpinus, *Linn. Sp. Pl.* 872.—Europ.; Orient; Ins. bor.
altaicus, *Willd. Enum. Hort. Berol.* 881.—Persia; Reg. Himal.; Ins. bor.
altissimus, *Mill. Gard. Dict. ed.* VIII. n. 20.—Am. bor.
altissimus, Moench, Meth. 601 = Novae-Angliae.
alwartensis, Lodd. Cat. (1807) ex Sims, Bot. Mag. t. 2321 = Erigeron pulchellus.
alysoides, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 198.—Mongol.
amabilis, *Wender. ex Nees, Gen. et Sp. Aster.* 262.—Hab. ?
ambiguus, Bernh. ex Nees, l. c. 188 = macrophyllus.
amelloides, Hoffm. in Goett. Gel. Anz. ix. (1800) 1253 = A. Amellus.
Amellus, Aitch. in Journ. Linn. Soc. xix. (1882) 168 = pseudamellus.
Amellus, *Linn. Sp. Pl.* 873.—Europ.; As. Min.; Reg. Caucas.
amethystinus, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 294.—Am. bor.
Ammani, Lindl. ex DC. Prod. v. 229 = sibiricus.
amoenus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 84.—Afr. austr.
amorsus, Lam. Encyc. i. 305, 307 = punctatus.
amplexicaulis, Lam. l. c. 304 = Novae-Angliae.
amplexicaulis, Muhl. in Willd. Sp. Pl. iii. 2046 = laevis.
amplexicaulis, Michx. Fl. Bor. Am. ii. 114 = patens.
amplus, *Lindl. in DC. Prod.* v. 236.—Am. arct.
amygdalinus, Lam. Encyc. i. 305 = umbellatus.
Andersoni, *A. Gray, in Proc. Am. Acad.* vii. (1868) 353.—Calif.
Andersoni, *E. Forst. ex Forbes, Hort. Woburn.* 188.—Am. bor.
andinus, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 290.—Am. bor.
angustifolius, Jacq. Hort. Schoenb. iii. 63 = Felicia angustifolia.
angustifolius, Lindl. in Royle, Illustr. Bot. Himal. 251. t. 58 = altaicus.
angustissimus, Tausch, in Flora, xi. (1828) 487 = Heteropappus hispidus.
angustus, Nees, Syn. Aster. 26 = paniculatus.
angustus, Schur, Enum. Pl. Transs. 306 = Kayserianus.
angustus, *Torr. & Gray, Fl. N. Am.* ii. 162.—As. bor.; Am. bor.
annectens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 78.—Afr. austr.
annuus, Linn. Sp. Pl. 875 = Erigeron annuus.
anomalus, *Engelm. in Torr. & Gray, Fl. N. Am.* ii. 503.—Am. bor.
Ansif, *Mart. in Denkschr. Muench. Akad.* vi. (1830) 196.—Arabia.
antirrhinifolius, Saliab. Prod. 197 = Cymbalariae.
argenteus, *Ando, Syn.* 121. t. 8. f. 2; *Lam. Encyc.* i. 302.—Hispan.
argoensis, Bout. ex Willk. & Lange, Prod. Fl. Hisp. ii. 36 = Willkommii.
arduenae, *Forbes, Hort. Woburn.* 188.—Hab. ?
armatioides, *Eaton, ex A. Gray, in Proc. Am. Acad.* viii. (1878) 647.—Am. bor. occ.
argentinus, Michx. Fl. Bor. Am. ii. 111 = sericeus.
argophyllus, Labill. Nov. Holl. Pl. ii. 52. t. 201 = Olearia argophylla.
argumentosus, DC. Prod. v. 237 = flaccidus.
argutus, R. Br. ex Benth. Fl. Austral. iii. 488 = Olearia arguta.
argutus, Nees, Gen. et Sp. Aster. 60 = Novi-Belgii.
Arnottii, Nees, ex Torr. & Gray, Fl. N. Am. ii. 115 = patens.
arrogonensis, Asch. = argoensis.
artemisiaeflorus, Poir. Encyc. Suppl i. 500 = Tradescanti.
artemisioides, F. Muell. Fragm. v. 65 = Olearia tubuliflora.
ascendens, Hook. Fl. Bor. Am. ii. 8 = adscendens.
asper, Buch.-Ham. ex DC. Prod. v. 277 = asperrimus.
asper, *Nees, Gen. et Sp. Aster.* 84.—Am. bor.
asperrimus, *Less. in Linnaea*, v. (1830) 148.—Mexic.
asperrimus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 292 = grandiflorus.

ASTER :—
asperrimus, Wall. Cat. n. 2970 = trioevia.
asperuginosus, Eaton, in Bot. King. Exp. 149 = Erigeron asperuginosus.
asperulus, Torr. & Gray, Fl. N. Am. ii. 190 = undulatus.
asperulus, *Phil. in Linnaea*, xxxiii. (1864-65) 131.—Chili.
brevipes, *Benth. Fl. Hongk.* 175.—China.
britannicus, All. Fl. Pedem. i. 197 = Inula britannica.
brumalis, Nees, Gen. et Sp. Aster. 70 = Novi-Belgii.
Bubonium, Scop. Fl. Carn. ed. II. ii. 178 = Inula spiraeifolia.
bupleurifolius, Hort. Monsp. ex Nees, Gen. et Sp. Aster. 386 = laevis.
bupleuroides, *Forbes, Hort. Woburn.* 188.—Hab.?
Burchellii, Hook. f. Ic. Pl. sub t. 1056 = Commidendron Burchellii.
Burkei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 76.—Afr. austr.
cabulicus, Lindl. Bot. Reg. (1843) Misc. 62 = Microglossa albescens.
caespitosus, Hort. ex Lindl. Bot. Reg. xix. t. 1571 = Novi-Belgii.
caespitosus, *E. Forst. ex Forbes, Hort. Woburn.* 188.—Hab.?
caffrorum, Less. Syn. Comp. 179 = Felicia caffrorum.
calcareus, *F. Muell. Census*, 78.—Austral.
calendulaefolius, *Lozini. Herb. Lmnar.* iv. 274.—Hab.?
californicus, Less. in Linnaea, vi. (1831) 33 = Erigeron glaucus.
campestris, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 293.—Am. bor. occ.
camporum, *Gardn. in Hook. Lond. Journ. Bot.* vii (1848) 79.—Bras.
Candyi, Vasey, ex A. Gray, Syn. Fl. N. Am. i. 198 = foliaceus.
Candollii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 80.—Afr. austr.
canescens, *Pursh, Fl. Am. Sept.* ii. 547.—Am. bor. occ.
canus, *Waldst. & Kit. Pl. Rar. Hung.* i. 28.—Europ.
capensis, Less. Syn. Comp. 188 = rotundifolius.
Capsii, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 304.—Turkest.
caricifolius, H. B. & K. Nov. Gen. et Sp. iv. 92. t. 338 = pauciflorus.
carneus, Fries, ex Baker, in Gard. Chron (1884) II. 680 = paniculatus.
carneus, *Nees, Syn. Aster.* 26.—Am. bor.
carneus, Torr. & Gray, partim, ex Baker, in Gard. Chron. (1884) II. 680 = salicifolius.
carnosus, *A. Gray, ex Hemsl. in Biol. Centr. Am. Bot.* ii. 130.—Am. bor. occ.
carnosus, Gillib. Fl. Lituan. i. 214 = A. Tripolium.
carolinianus, *Walt. Fl. Carol.* 208.—Am. bor.
caryophyllus, Stead. Nom. ed. II. i. 154, sphalm. = Callistephus.
cassiniaeformis, *Maund. Bot. Gard.* vi. n. 572.—Russia.
Cassiniae, F. Muell. Fragm. v. 68 = Olearia Cassiniae.
castaneifolius, *Willd. & Costa, in Linnaea*, xxx. (1859-60) 106.—Hispan.
caulescens, Hook. Bot. Mag. lxxiii. (1847) App. 34 = Microglossa albescens.
caucasicus, *Willd. Sp. Pl.* iii. 2090.—Reg. Caucas.
Celmisia, F. Muell. Fragm. v. 84 = Celmisia longifolia.
chamaedrifolius, Lam. ex DC. Prod. vii. 13 = Printzia Herrii.
Chamissonis, *A. Gray, in Torr. Bot. U. St. Expl. Exped.* 541.—Calif.
Chatanani, *Torr. & Gray, Fl. N. Am.* ii. 161.—Am. bor.
Charieis, Less. Syn. Comp. 171 = strigosus.
chilensis, Nees, Gen. et Sp. Aster. 138 = Chamisconis.
chinensis, Linn. Sp. Pl. 877 = Callistephus hortensis.
chrysanthemifolius, Hort. ex Steud. Nom. ed. II. i. 154 = filifolius.
chrysanthemoides, Willd. ex Spreng. Syst. iii. 538 = tanacetifolius.
chrysocomifolius, Hort. ex Steud. Nom. ed. II. i. 154 = Nolletia chrysocomoides.
chrysocomoides, Desf. Atb. i. 283 = Nolletia chrysocomoides.
chrysocomoides, Turcz. ex DC. Prod. vii. 272 = alpinus.
chrysophyllus, A. Cunn. ex DC. l. c. v. 268 = Olearia chrysophylla.
ciliaris, Thunb. ex DC. l. c. 218 = Mairia taxifolia.
ciliatus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 295 = phyllolepis.
ciliatus, *Walt. Fl. Carol.* 209 (sp. dub.).—Am. bor.

ASTER :—
ciliatus, Muhl. in Willd. Sp. Pl. iii. 2037 = multiflorus.
ciliolatus, Lindl. in DC. Prod. v. 285 = Lindleyanus.
coerulescens, DC. l. c. 335 = salicifolius.
Collinii, Nutt. in Journ. Acad. Philad. vii. (1834) 82 = Sericocarpus tortifolius.
collinus, Saliab. Prod. 198 = A. Amellus.
coloradoënsis, *A. Gray, in Proc. Am. Acad.* xi. (1876) 76.—Colorado.
communis, *A. Gray, Syn. Fl. N. Am.* i. II. 185.—Am. bor. occ.
concinnus, Colla, Hort. Ripul. App. iii. 22. t. 13 = Novae-Angliae.
concinnus, Saliab. Prod. 197 = fruticosus.
concinnus, *Willd. Enum. Hort. Berol.* 884.—Am. bor.
concolor, *Linn. Sp. Pl.* 874.—Am. bor.
confertus, Desf. Cat. Hort. Par. ed. III. 401 = puniceus.
confertus, Nees, Syn. Aster. 21 = versicolor.
conocephalus, F. Muell. Fragm. v. 67 = Olearia conocephala.
consanguineus, Ledeb. Fl. Ross. ii. 473 = peregrinus.
conspicuus, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) 144 = tataricus.
conspicuus, *Lindl. in DC. Prod.* v. 280.—Am. bor. occ.
Conyzae, Griesael. Kleine Bot. Schrift. 122 = Inula vulgaris.
conyzoides, Desf. Cat. Hort. Par. ed. III. 175 = Felicia angustifolia.
conyzoides, Willd. Sp. Pl. iii. 2048 = Sericocarpus conyzoides.
cordifolius, Lam. Encyc. i. 307 = paniculatus.
cordifolius, *Linn. Sp. Pl.* 875.—Am. bor.
cordifolius, Michx. Fl. Bor. Am. ii. 112 = corymbosus.
cordifolius, Muhl. ex Willd. Sp. Pl. iii. 2035 = sagittifolius.
coriaceus, Forst. f. Prod. 56. n. 297 = Celmisia coriacea.
coriaceus, Hort. ex DC. Prod. v. 276 = fruticosus.
coridifolius, Michx. ex Willd. Sp. Pl. iii. 2028 = dumosus.
cornifolius, Muhl. ex Willd. l. c. 2039 = infirmus.
Cornuti, Mill. ex Nees, in DC. Prod. v. 284 = abbreviatus.
Cornuti, Wendl. ex Nees, Gen. et Sp. Aster. 58 = patulus.
corymbosus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 207. —Am. bor.
corymbosus, Moench, Meth. Suppl. 250 = Inula germanica.
crassifolius, *F. W. Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 326.—N. Granat.
crenatus, Less. Syn. Comp. 166 = Mairia crenata.
crenatus, Linn. Pl. Rar. Afr. 21 = Athrixia heterophylla.
crenifus, Thunb. Prod. Pl. Cap. 160 = Athrixia capensis.
crispus, Forsk. Fl. Aegypt. Arab. 150 = Pulicaria crispa.
crocatus, *Bertol. Fl. Guatimal.* 34.—Guatemala.
crocifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 89.—N. Granat.
cumeatus, A. Cunn. ex Steud. Nom. ed. II. i. 154 = Olearia stellulata.
Cunninghamii, *F. Muell. Census*, 78.—Austral.
Curtisii, *Torr. & Gray, Fl. N. Am.* ii. 110.—Am. bor.
Cusickii, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 99. —Am. bor. occ.
cyaneus, Hoffm. Phyt. Blaett. i. (1808) 71 = laevis.
cyanodiscus, F. Muell. Fragm. v. 82 = Olearia ramosissima.
cydoniaefolius, A. Cunn. ex DC. Prod. v. 267 = Olearia cydoniaefolia.
cylleneus, Orphan. ex Nym. Consp. 387 = alpinus.
Cymbalariae, *Ait. Hort. Kew.* ii. 197 ; *Thunb. Prod. Pl. Cap.* 160.—Afr. austr.
daharicus, *Benth. ex Baker, in Gard. Chron.* (1885) i. 208.—Sibir.
debilis, Boj. ex DC. Prod. vi. 276 = Athrixia debilis.
decorus, Desf. ex DC. l. c. v. 246 = laevis.
decumbens, *Baker, in Mart. Fl. Bras.* vi. III. 94.—Bras.
decurrens, A. Cunn. ex DC. Prod. v. 269 = Olearia decurrens.
decurrens, Moench, Meth. Suppl. 251 = Inula bifrons.
defluxus, Moench, l. c. 250 = laevis.
defluxus, *Tausch, in Flora*, xxii. (1839) 77.—Hab.? demissus, *Forbes, Hort. Woburn.* 188.—Hab.?

ASTER :—

demissa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 79. —Afr. austr.

dentatus, Andr. Bot. Rep. t. 61 = Olearia dentata.

dentatus, Thunb. Prod. Pl. Cap. 159 = Felicia fragilis.

denudatus, Nutt. in Trans. in Am. Phil. Soc. Ser. II. vii. (1841) 292 = adscendens.

depressus, Kth. in Linnæa, xxxii. (1863) 373 = A. Tripolium.

descriorum, *Tchertk. in Bull. Soc. Nat. Mosc.* vi. (1833) 18.—Reseda.

dichotomus, Ell. Sketch, ii. 366 = reticulatus.

diffusus, DC. Prod. v. 249, partim = vimineus.

diffusus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 205. —Am. bor.

dimorphophyllus, *Franch. & Sav. Enum. Pl. Jap.* i. 224.—Japon.

diplostephioides, *Benth. & Hook. f. Gen.* ii. 372.—Reg. Himal.

discoideus, Boiss. & Rent. Pugill. 54 = Willkommii.

discoidens, Ell. Sketch, ii. 358 = Solidago discoides.

discoidens, *Sond. ex Harv. & Sond. Fl. Cap.* iii. 76.—Afr. austr.

distortus, Turcz. ex Ayré-Lall. in Ind. Sem. Hort. Petrop. viii. 58. n. 846 = altaicus.

divaricatus, Lam. Encyc. i. 305 = acuminatus.

divaricatus, Linn. Sp. Pl. 873 = corymbosus.

divaricatus, *Nutt. in Torr. & Gray, Fl. N. Am.* ii. 168.—Am. bor. et austr.

divaricatus, Rafin. ex DC. Prod. v. 241 = divergens.

divaricatus, Spreng. Syst. iii. 539 = umbellatus.

divaricatus, Torr. & Gray, Fl. N. Am. ii. 168 = exilis.

divergens, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 205 = diffusus.

diversifolius, Michx. Fl. Bor. Am. ii. 118 = undulatus.

diversifolius, Spreng. Syst. iii. 533 = heterophyllus.

Donii, *E. Forst. ex Forbes, Hort. Woburn.* 188.—Hab. ?

Douglasii, Eaton, Bot. King, Exp. 141 = foliaceus.

Douglasii, *Lindl. in DC. Prod.* v. 289.—Am. bor. occ.

dracunculoides, Bess. Enum. Pl. Volh. 83 = acris.

dracunculoides, *Lam. Encyc.* i. 308.—Europ. austr.; As. bor.

dracunculoides, Willd. Sp. Pl. iii. 2050 = paniculatus.

Dregei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 74.—Afr. austr.

Drummondii, *Lindl. ex Hook. Comp. Bot. Mag.* i. (1836) 97.—Am. bor.

dumosus, DC. Prod. v. 241 = multiflorus.

dumosus, Hoffm. Phytogr. Blactt. i. (1808) 68 = ericoides.

dumosus, *Linn. Sp. Pl.* 873.—Am. bor.

dumosus, Nees, Syn. Aster. 31 = ericoides.

Durandii, Nutt. ex Durand, in Pacif. Rail. Rep. v. 8 = Menziesii.

dysenterica, Scop. Fl. Carn. ed. II. ii. 171 = Pulicaria dysenterica.

echinatus, Lour. Syn. Comp. 177 = Felicia echinata.

Ecklonis, *Less. i. c.* 170.—Afr. austr.

Ekrenbergii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 302.—Mexic.

elaeagnifolius, A. Cunn. ex DC. Prod. v. 267 = Olearia furfuracea.

elaeophilus, A. Cunn. l. c. 269 = Olearia heterophylla.

elatus, Bert. ex Steud. Nom. ed. II. i. 154 = corymbosus.

elegans, Hort. Par. ex A. Gray, Syn. Fl. N. Am. i. 11 154 = concinnus.

elegans, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 44.—Sikkim.

elegans, Nees, Syn. Aster. 30 = A. Amellus.

elegans, *Torr. & Gray, Fl. N. Am.* ii. 159.—Am. bor. occ.

elegans, Willd. Sp. Pl. iii. 2049 = spectabilis.

Elliottii, *Torr. & Gray, Fl. N. Am.* ii. 140.—Am. bor.

elliptieus, A. Cunn. ex DC. Prod. v. 271 = Olearia elliptica.

elodes, Torr. & Gray, Fl. N. Am. ii. 136 = Novi-Belgii.

elongatus, *Thunb. Prod. Pl. Cap.* 160.—Afr. austr.

elongifolius, A. Cunn. in Ann. Nat. Hist. ii. (1839) 197, sphalm. = *elaeagnifolius*.

eminens, Lindl. in Bot. Reg. t. 1614 = Novi-Belgii.

eminens, Nees, Gen. et Sp. Aster. 57 = junceus, paniculatus.

ASTER :—

eminens, Willd. Enum. Hort. Berol. 886 = salicifolius.

Engelmannii, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 382.—Am. bor. occ.

ensifolius, Scop. Fl. Carn. ed. II. ii. 174 = Inula ensifolia.

erectus, *Forbes, Hort. Woburn.* 188.—Hab. ?

eremophilus, *Bunge, Verz. Suppl. Fl. Alt.* 108.—Sibir. altaic.

ericaefolius, Forsk. Fl. Aegypt. Arab. 150 = Athrixia ericaefolia.

ericaefolius, *Rothr. in Coulter, Bot. Gas. ii.* (1877) 70. —Mexic.; Texas.

ericoides, Lam. Encyc. i. 304 = multiflorus.

ericoides, *Linn. Sp. Pl.* 875.—Am. bor.

erigeroides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 74. —Afr. austr.

erigeroides, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 48 = Erigeron fruticosus.

erubescens, Sieber, ex Benth. Fl. Austral. iii. 471 = Olearia myrsinoides.

erysigilifolius, *Torr. & Gray, Fl. N. Am.* ii. 502.—Am. bor.

espenbergensis, Nees, Gen. et Sp. Aster. 86 = sibiricus.

exasperatus, Link, Enum. Hort. Berol. ii. 238 = Olearia ramulosa.

exiguifolius, F. Muell. Fragm. v. 67 = Olearia exiguifolia.

exilifolius, F. Muell. l. c. 69 = Olearia exilifolia.

exilis, *Ell. Sketch*, ii. 344.—Am. bor.

exscapus, Richards. App. Frankl. Journ. ed. II. 32 = Townsendia sericea.

expansus, Nees, Syn. Aster. 28 = laevis.

Exsul, Lindl. Swan Riv. Fl. App. 24 = Olearia rudis.

falcatus, Eaton, in Bot. King, Exped. 140, ex parte = adscendens.

falcatus, *Lindl. in DC. Prod.* v. 241.—Am. arct.

fascicularis, E. Mey. ex DC. l. c. 290 = Felicia muricata.

fastigiatus, *Pisch. in Mém. Soc. Nat. Mosc.* iii. (1812) 74.—Sibir.

fastigiatus, Ledeb. Ic. Pl. Fl. Ross. t. 161 = Hauptii.

Fendleri, *A. Gray, Pl. Fendl.* 86.—Am. bor.

Fermeli, F. Muell. Fragm. v. 75 = Olearia Ferresii.

ferruginens, Edgew. in Trans. Linn. Soc. xx. (1846) 64 = Microglossa albescens.

ferruginens, Wendl. in Flora, ii. (1819) 676 = Olearia dentata.

ficoides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 76.—Afr. austr.

filaginifolius, Hook. & Arn. Bot. Beech. Voy. 146 = Corethrogyne filaginifolia.

filifolius, *Vent. Jard. Malm.* 83.—Afr. austr.

filifolius, Willd. ex Nees, Gen. et Sp. Aster. 210 = Felicia muricata.

filiformis, Eckl. ex DC. Prod. v. 218 = Matria taxifolia.

filiformis, Steud. Nom. ed. II. i. 154 = Felicia muricata.

flexuosus, Nutt. Gen. Am. ii. 154 = tenuifolius.

floribundus, *Nutt. i. c.* 158.—Am. bor.

floribundus, Willd. Sp. Pl. iii. 2048 = Novi-Belgii.

floriferus, F. Muell. Fragm. v. 93 = Olearia floribunda.

fluminensis, *Vell. Fl. Flum.* viii. t. 122.—Bras.

foliaceus, *Lindl. in DC. Prod.* v. 238.—Am. bor. occ.

foliaceus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 202 = vimineus.

foliaceus, Ell. Sketch, ii. 345 = dumosus.

foliosus, Pers. Syn. ii. 443 = dumosus.

formosissimus, *Hort. ex Heynh. Nom.* ii. 51. nomen; *Vilm. Fl. Pl. Terre*, ed. I. 98.—Hab. ?

Forsteri, *Borr. ex Forbes, Hort. Woburn.* 188.—Hab. ?

fragilis, Lindl. in Bot. Reg. t. 1527 = dumosus.

fragilis, Jacq. ex Steud. Nom. ed. I. 33 = linarifolius.

fragilis, Nees, Gen. et Sp. Aster. 101 = vimineus.

fragilis, *Schlecht. & Vahet. Syll. Fl. Croat.* 97.—Croatia.

fragilis, Willd. Sp. Pl. iii. 2051 = Tradescanti.

ASTER :—

Fremonti, *A. Gray, Syn. Fl. N. Am.* i. ii. 191.—Am. bor. occ.

frondosus, *Torr. & Gray, Fl. N. Am.* ii. 165.—Am. bor. occ.

frutetorum, Wimm. ex Nym. Consp. ii. 887 = adligus.

fruticosus, *Linn. Sp. Pl.* 873.—Afr. austr.

fruticulosus, Sims, Bot. Mag. t. 2356 = pilosiflorus.

fruticulosus, Burm. f. Prod. 27 = Felicia angustifolia.

fruticulosus, Krebs, ex DC. Prod. v. 276 = filifolius.

fruticulosus, Willd. Sp. Pl. iii. 2018, pro parte = obtusatus.

fruticulosus, Murr. Syst. Veg. ed. XIII. 638 = fruticosus.

fugax, Brot. Fl. Lusit. 385 = aragonensis.

furfuraceus, A. Rich. & Lesson, Fl. Nov. Zel. 216 = Olearia furfuracea.

Gardneri, *Baker, in Mart. Fl. Bras.* vi. iii. 95.—Bras.

Garibaldii, Bruegg. in Verh. Schweis. Naturf. Ges. Samaden, 1868 (1864) 229 = alpinus.

Gayanus, *DC. Prod.* v. 287.—Chili.

Gelderi, Bess. ex DC. l. c. 259 = altaicus.

Gerlachii, *Hance, in Journ. Bot.* xvii. (1880) 262.—China.

gigenteus, Hort. ex Vilm. Fl. Pl. Terre, ed I 998 = Vernonia noveboracensis.

Gilliesii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 49.—Chili.

glabellus, Nees, Syn. Aster. 31 = ericoides.

glaber, E. Mey. ex Nees, Gen. et Sp. Aster. 68 = vimineus.

glaber, *Mill. Gard. Dict. ed.* VIII. n. 17.—Am. bor.

glabriusculus, Torr. & Gray, Fl. N. Am. ii. 159 = xylorrhizus.

glacialis, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 291.—Am. bor.

glandulosus, Labill. Nov. Holl. Pl. ii. 50. t. 197 = Olearia glandulosa.

glaucescens, Wender. ex Nees, Gen. et Sp. Aster. 188 = laevis.

glaucus, Nees, Syn. Aster. 33 = laevis.

glaucus, *Torr. & Gray, Fl. N. Am.* ii. 159.—Am. bor. occ.

glomeratus, Bernh. ex DC. Prod. v. 265 = macrophyllus.

Glehni, *F. Schmidt, Reisen Amurl.* 146.—As. or.

glutescens, F. Muell. Fragm. v. 77 = Olearia glutinosa.

glutinosus, Cav. Ic. ii. 58 = Grindelia glutinosa.

glutinosus, Roxb. in Beats. App. 808 = Commidendron rugosum.

Gmelini, Ledeb. in Denkschr. Baier. Bot. Ges. iii. (1841) 77 = Senecio Gmelini.

Gmelini, Tausch, in Flora, xi. (1828) 486 = Heteropappus hispidus.

Gmelini, Tausch, l. c. xxii. (1839) i. 78 = altaicus.

gracilentus, Banks, ex DC. Prod. v. 237 = Chrysopsis villosa.

gracilentus, Torr. & Gray, Fl. N. Am. ii. 166 = virgatus.

gracilis, *Nutt. Gen. Am.* ii. 158.—Am. bor.

graminifolius, Pursh, Fl. Am. Sept. ii. 545 = Erigeron hyssopifolius.

grandiflorus, *Hoffm. Phyt. Blactt.* i. (1808) 65.—Hab. ?

grandiflorus, *Linn. Sp. Pl.* 877.—Am. bor.

grandiflorus, Nutt. Gen. Am. ii. 156 = paludosus.

gramineus, Nutt. in Journ. Ac. Nat. Sc. Phil. ii. (1821) 116 = oblongifolius.

gravis, F. Muell. Fragm. v. 93 = Olearia gravis.

Greenii, Nees, ex Torr. & Gray, Fl. N. Am. ii. 158 = nemoralis.

Greenii, Torr. & Gray, l. c. 184 = salicifolius.

grumosiflorus, Hook. f. Ic. Pl. t. 1056 = Commidendron spartioides.

gymnocephalus, *A. Gray, in Proc. Am. Acad.* xv. (1880) 93.—Texas; Mexic.

Hamiltoni, *Nees, Gen. et Sp. Aster.* 68.—Chili.

Hallii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 388 = Am. bor. occ.

Hauptii, *Ledeb. Fl. Alt.* iv. 100.—Sibir.

Haupti, Tausch. ex DC. Prod. v. 256 = dahuricus.

Haydeni, Porter, in Hayden, Geol. Rep. 1871 (1872) 448 = paludosus.

helenioides, DC. Prod. v. 242 = salicifolius.

Helenium, Scop. Fl. Carn. ed. II. ii. 171 = Inula Helenium.

heliophilus, F. Muell. Fragm. v. 66 = Olearia heliophila.

ASTER:—
Hersilei, *Klatt, Comp. Herb. Schlag. Hochas.* 368.—As. temp.
Herveyi, *A. Gray, Man. Bot. U. St. ed.* V. 280.—Am. bor.
hesperius, *A. Gray, Syn. Fl. N. Am.* i, II. 192.—Am. bor. occ.
heterochaeta, C. B. Clarke, Comp. Ind. 44 = Heterochaeta asteroides.
heteromallus, Wender. in Schrift. Marb. ii. (1831) VI. 54 = undulatus.
heterophyllus, Nees, Syn. Aster. 15 = paniculatus.
heterophyllus, Thunb. Prod. Pl. Cap. 160 = Athrixia heterophylla.
heterophyllus, Willd. Sp. Pl. iii. 55 = cordifolius.
heterophyllus, *Schloss. & Vukot. Syll. Fl. Croat.* 55.—Croatia.
Heynei, F. Muell. Fragm. v. 86 = Olearia xerophila.
hiemalis, Nees, Gen. et Sp. Aster. 77 = longifolius.
himalaicus, *C. B. Clarke, Comp. Ind.* 48.—Reg. Himal.
hirsuticaulis, Lindl. in DC. Prod. v. 242 = diffusus.
hirsutus, Host, Fl. Austr. ii. 485 = alpinus.
hirsutus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 80.—Afr. austr.
hirtellus, Lindl. in DC. Prod. v. 238 = sagittifolius.
hirtus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 82.—Afr. austr.
hirtus, *C. Koch, in Linnaea*, xxiii. (1850) 701.—Armenia.
hirtus, Scop. Fl. Carn. ed. II. ii. 173 = Inula hirta.
hirtus, Thunb. Prod. Pl. Cap. 159 = Felicia angustifolia.
hispidus, *Baker, in Saund. Ref. Bot.* t. 242.—Afr. austr.
hispidus, Lam. Encyc. i. 307 = punilceus.
hispidus, Thunb. Fl. Jap. 315 = Heteropappus hispidus.
holosericeus, Forst. f. Prod. 56. n. 296 = Celmisia holosericea.
homolepis, F. Muell. Fragm. v. 65 = Olearia homolepis.
Hookerii, *Forbes, Hort. Woburn.* 189.—Hab.!
horizontalis, Desf. Cat. Hort. Par. ed. III. 402 = diffusus.
Huegelii, F. Muell. Fragm. v. 76 = Olearia ciliata.
humilis, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 125 = Vittadinia australis.
humilis, Willd. Sp. Pl. iii. 2038 = infirmus.
hungaricus, Poir. Encyc. Suppl. i. 496 = saligna.
Hunnemanni, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.!
hybernus, *E. Forst. ex Forbes, l. c.*—Hab.!
hydrobadensis, Wall. Cat. n. 2973 = Pulicaria Wightiana.
hygrophilus, A. Cunn. ex DC. Prod. v. 269 = Olearia hygrophila.
hygrophilus, Berg. Desc. Pl. Cap. 387 = Felicia angustifolia.
hyssopifolius, Cav. Ic. t. 282 = sedifolius.
hyssopifolius, Linn. Mant. 114 = acris.
ibericus, Stev. ex Bieb. Fl. Taur. Cauc. ii. 311 = A. Amellus.
ignotus, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 11.—Ind. or.
ilicifolius, A. Cunn. ex DC. Prod. v. 268 = Olearia myrsinoides.
illinus, F. Muell. Fragm. v. 76 = Olearia elliptica.
imbricatus, *Linn. Pl. Rar. Afr.* 21.—Afr. austr.
imbricatus, Walp. Rep. ii. 574 = armarioides.
inaequalis, *Hill, Hort. Kew.* 16.—Am. bor.
incanus, *A. Gray, in Bot. Calif.* i. 332.—Calif.; Mexic.
incisus, *Pisch. in Mém. Soc. Nat. Mosc.* iii. (1819) 76.—Sibir.
inconspicuus, *Less. in Linnaea*, v. (1830) 143.—Mexic.
indicus, Heyne, in Wall. Cat. n. 2971 = Pulicaria Wightiana.
indicus, Linn. Sp. Pl. 876 = Boltonia indica.
infirmus, *Michx. Fl. Bor. Am.* ii. 109.—Am. bor.
infirmus, *Desf. Tabl. ed. II.* 122.—Hab.!
integrifolius, *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884). 224.—China.
integrifolius, *Nutt. in Trans. Am. Phil. Soc. Ser.* II. vii. (1841) 291.—Am. bor. occ.
interruptus, Turcz. ex DC. Prod. vii. 272 = flaccidus.
intybaceus, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 11 = tataricus.
involucratus, *Boiss. & Bal. ex Boiss. Fl. Orient.* iii. 189.—Reg. Caucas.

ASTER:—
inuloides, D. Don, Prod. Fl. Nep. 178 = Erigeron multiradiatus.
inuloides, Fisch. ex DC. Prod. v. 231 = sibiricus.
iodochrous, F. Muell. Fragm. v. 81 = Olearia iodochroa.
ircutianus, DC. Prod. v. 229 = tataricus.
Italicus, *Hill, Veg. Syst.* ii. 88.—Italia.
jamaicensis, *Less. in Linnaea*, v. (1830) 144.—Mexic.; Ind. occ.
japonicus, *Less. ex Nees, Gen. et Sp. Aster.* 33, 34.—Japon.
junceus, *Ait. Hort. Kew.* iii. 204.—Am. bor.
junceus, Nees, Syn. Aster. 26 = tenuifolius.
Kayserianus, *Schur, Enum. Pl. Transs.* 306.—Europ.
Kingii, *Eaton, Bot. King, Exp.* 141. t. 16.—Am. bor. occ.
Kraussii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 84.—Afr. austr.
Kumleini, Fries, ex A. Gray, Syn. Fl. N. Am. i. II. 179 = oblongifolius.
lacerus, Lindl. ex DC. Prod. v. 232 = sibiricus.
lacunarum, *Aitch. & Hemsl. in Journ. Linn. Soc* xix. (1882) 168.—Afghan.
laetus, Forst. ex Ledeb. Fl. Ross. ii. 479 = punctatus.
laevigatus, Hook. Bot. Mag. t. 2995 = laevis.
laevigatus, Lam. Encyc. i. 307 = Novi-Belgii.
laevigatus, Pursh, Fl. Am. Sept. ii. 553 = longifolius.
laevigatus, Willd. Sp. Pl. iii. 2046 = versicolor.
laevis, Lindl. in Bot. Reg. t. 1500 = versicolor.
laevis, *Linn. Sp. Pl.* 876.—Am. bor.
laevis, Vahl, ex DC. Prod. v. 283 = longifolius.
Lake, *C. B. Clarke, Comp. Ind.* 49.—Reg. Himal.
Lamarckianus, Nees, Gen. et Sp. Aster. 100 = paniculatus.
lanceolatus, Willd. Sp. Pl. iii. 2050 = paniculatus.
lanceolatus, Nutt. Gen. Am. ii. 158 = salicifolius.
lanuginosus, *Wendl. Beytr.* ii. 9.—Afr. austr.
lasiocarpus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 76.—Afr. austr.
lateralis, Hort. Bonn. ex Nees, Gen. et Sp. Aster. 74 = longifolius.
latifolius, Desf. Cat. Hort. Par. ed. III. 176 = macrophyllus.
latifolius, Banks, ex Steud. Nom. ed. II. i. 165 = acuminatus.
latifolius, Mill. Gard. Dict ed. VIII. n. 26 = trinervis?
lauterianus, *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) 224.—China.
laxus, *Willd. ex DC. Prod.* v. 274.—Mexic.
laxiflorus, Lindl. in Hook. Fl. Bor. Am. ii. 10 = junceus.
laxiflorus, Nees, Gen. et Sp. Aster. 69 = longifolius.
laxus, Torr. & Gray, Fl. N. Am. ii. 184 = Novi-Belgii.
laxus, Willd. Enum. Hort. Berol. ii. 857 = paniculatus.
ledifolius, A. Cunn. ex DC. Prod. v. 269 = Olearia ledifolia.
ledifolius, Pursh, Fl. Am. Sept. ii. 544 = nemoralis.
ledophyllus, A. Gray, in Proc. Am. Acad. xvi. (1880) 98 = Engelmanni.
leiocarpus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 84.—Afr. austr.
Lehmanni, F. Muell. Fragm. v. 66 = Olearia heleophila.
leiophyllus, *Franch. & Sav. Enum. Pl. Jap.* i. 223.—Japon.
leiophyllus, Schloss. & Vukot. Syll. Fl. Croat. 56 = saligneus.
Lemmoni, *A. Gray, Syn. Fl. N. Am.* i. II. 199.—Arizona.
lepidophyllus, Pers. Syn. ii. 442 = Olearia lepidophylla.
leptophyllus, *Desf. Tabl. ed. II.* 268; *DC. Prod.* v. 256.—Am. bor.
leptophyllus, *Fisch. Hort. Gorenk.* ed. II. (1812) 89, nomen.—Sibir.
leptophyllus, Hort. Par. ex DC. Prod. v. 242 = ericoides.
Lerchenfeldianus, *Schur, Enum. Pl. Transs.* 806.—Hungar.
leucanthemus, Desf. Cat. Hort. Par. ed. III. 401 = Tradescanti.
leucanthemus, *Rafin. in Med. Repos. N. York*, v. (1808) 359.—Am. bor.
leucrioides, Mnhl. ex DC. Prod. v. 230 = surculosus.
lilacinus, *Hoffegg. Verz. Pfl.* 41.—Hab.!
Lima, *Lindl. in DC. Prod.* v. 230.—Mexic.
linariifolius, *Linn. Sp. Pl.* 874.—Am. bor.
Lindeni, *Sch. Bip. in Seem. Bot. Voy. Herald*, 302.—Mexic.

ASTER:—
Lindleyanus, *Torr. & Gray, Fl. N. Am.* ii. 122.—Am. bor.
linifolius, Burm. f. Prod. 27 = Senecio rosmarinifolius.
linifolius, *Harv. in Harv. & Sond. Fl. Cap.* iii. 81.—Afr. austr.
linifolius, Linn. Sp. Pl. 874 = acris.
linifolius, Torr. & Gray, Fl. N. Am. ii. 162 = subulatus.
linoides, Hort. Gaybac. ex DC. Prod. v. 221 = Felicia caffrorum.
Linosyris, *Bernh. Syst. Verz. Erf.* 151.—Europ.
lyratus, Sims, Bot. Mag. t. 1509 = Olearia stellulata.
lithospermifolius, Desf. Tabl. ed. II. 268; DC. Prod. v. 259 = altaicus?
litoralis, *Phil. in Anal. Univ. Chil.* (1873) 486.—Chili.
livitus, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.!
longicaulis, Desf. ex DC. Prod. v. 253 = A. Tripolium.
longifolius, *Desf. Cat. Hort. Par.* ed. III. 400.—Hab.!
longifolius, A. Gray, Man. Bot. U. St. ed. V. 283 = Novi-Belgii.
longifolius, *Lam. Encyc.* i. 306.—Am. bor.
longifolius, Nees, Syn. Aster. 26 = laxiflolius.
longipes, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 78 = Leucopsis scaposa.
lucidus, Hort. Kew. ex Loud. Hort. Brit. 847 = laevis.
lucidus, Wender. in Flora, xii. (1829) i. Erg. 25 = paucens.
lucidus, *Moench, Meth. Suppl.* 252.—Hab.!
lusitanicus, Brot. Phyt. 68. i. 29 = aragonensis.
lutescens, Torr. & Gray, Fl. N. Am. ii. 160 = ptarmicoides.
luxurians, Spreng. Syst. iii. 538; Nees, Gen. et Sp. Aster. 88 = Novi-Belgii.
lyratus, Nees, Gen. et Sp. Aster. 188, in syn. = Olearia stellulata.
Maackii, *Regel, in Mém. Acad. Pétersb.* Sér. VII. iv. (1861) n. IV. 31.—Japon.
Macleai, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.!
macrophyllus, *Linn. Sp. Pl.* ed. II. ii. 1223.—Am. bor.
macrorrhinus, A. Cunn. ex DC. Prod. v. 266 = Vittadinia macrorrhiza.
macrorrhizus, *Thunb. Prod. Pl. Cap.* 159.—Afr. austr.
magellanicus, Lam. Illustr. iii. 259. t. 681 = Pereria lactucoides.
magellanicus, Spreng. Syst. iii. 536 = Chiliotrichium amelloideum.
magnificus, F. Muell. Fragm. v. 80 = Olearia magnifica.
magnistus, *H. B. & K. Nov. Gen. et Sp.* iv. 91.—N. Granat.
maritimus, Salisb. Prod. 198 = A. Tripolium.
Martii, *Baker, in Mart. Fl. Bras.* vi. III. 38.—Brasa.
marylandicus, Michx. Fl. Bor. Am. ii. 108 = Sericocarpus conyzoides.
maximus, Less. Syn. Comp. 182 = Erigeron maximus.
megalophyllus, F. Muell. Fragm. v. 70 = Olearia megalophylla.
megapotanicus, *Spreng. Syst.* iii. 536.—Bras.
Menziesii, *Lindl. in DC. Prod.* v. 248.—Am. bor. occ.
mesopotamicus, Steud. Nom. ed. I. 89 = megapotanicus.
mesopotamicus, *Spreng. Nachr.* i. *Bot. Gart. Halle*, 13.—Hab.!
mespilifolius, Less. Syn. Comp. 180 = Nidorella mespilifolia.
microcephalus, *Franch. & Sav. Enum. Pl. Jap.* i. 232.—Japon.
microphyllus, Labill. Nov. Holl. Pl. ii. 51 = Olearia lepidophylla.
microphyllus, Torr. ex Lindl. in DC. Prod. v. 244 = adnatus.
microphyllus, Vent. Jard. Malm. sub t. 58 = Olearia ramulosa.
microspermus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 72.—Afr. austr.
Milleri, *Forbes, Hort. Woburn.* 189.—Hab.!
mirabilis, *Torr. & Gray, Fl. N. Am.* ii. 165.—Am. bor.
miser, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 205 = Tradescanti.
miser, Lam. Encyc. i. 308 = virgatus.
miser, Linn. Sp. Pl. 877 (delend.—*Cf.* A. Gray, Syn. Fl. i. ii. 179).

ASTER :—
minor, Nutt. Gen. Am. ii. 158 = diffusus.
Mitchelii, *F. Muell. Fragm.* v. 78.—Austral.
modestus, *Lindl. in DC. Prod.* v. 281.—Am. bor. occ.
mollissimus, *Wall. Cat.* n. 2972.—Reg. Himal.
monanthus, *Vell. Fl. Flum.* viii. t. 121.—Bras.
mongolicus, *Franch. Pl. Davidian.* i. 161.—As. or.
monocephalus, Boiss. in Bal. ex Boiss. Fl. Orient. iii. 160 = caucasicus.
montanus, All. Fl. Pedem. i. 195 = Inula montana.
montanus, Nutt. Gen. Am. ii. 156 = sericeus.
montanus, R. Br. in Richards. App. Frankl. Journ. ed. I. 749 = sibiricus.
montevidensis, Griseb. in Goett. Abh. xix. (1874) 170 = Leucopis cliffusa.
monticolus, Lindl. ex DC. Prod. v. 287 = Erigeron monticolus.
Mooneyi, *F. Muell. Fragm.* viii. 144.—Austral.
morarensis, *H. B. & K. Nov. Gen. et Sp.* iv. 93.—Mexic.
Muehlenbergii, Tausch, in Flore, xxii. (1839) 77, in adnot. = amplexicaulis.
Muelleri, *F. Muell. Census*, 78.—Austral.
multicaule, Lindl. in DC. Prod. v. 287 = oblongifolius.
multiflorus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 116.—Am. bor.; Mexic.
multiflorus, Bertol. Misc. Bot. v. t. 5 = dumosus.
multiflorus, Nutt. Gen. Am. ii. 155 = vimineus.
multiradiatus, Wall. Cat. n. 2969 = Erigeron multiradiatus.
muricatus, F. Muell. Fragm. v. 66 = Olearia muricata.
muricatus, Thunb. Prod. Pl. Cap. 159 = Felicia muricata.
mutabilis, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 208 = Novi-Belgii.
mutabilis, *Vell. Fl. Flum.* viii. t. 122.—Bras.
mutabilis, *Linn. Sp. Pl.* 876 (sp. obscura).—Am. bor.
mutabilis, Willd. Sp. Pl. iii. 2045 = versicolor.
mutatus, Torr. & Gray, Fl. N. Am. ii. 149 = modestus.
myriantus, *Carr. in Rev. Hortic.* (1856) 439.—Hab.?
myrsinoides, Labill. Nov. Holl. Pl. ii. 58. t. 202 = Olearia myrsinoides.
myrtifolius, Willd. Enum. Hort. Berol. Suppl. 59 = uber.
nanasquanus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 79. —Afr. austr.
nanus, E. Mey. ex DC. Prod. v. 218 = Mairia microcephala.
natalensis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 81.—Afr. austr.
Neesii, Sch. Bip. in Flora, xxxix. (1856) 854 = Chamissonis.
nemoralis, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 198. —Am. bor.
Nerestii, F. Muell. Fragm. v. 81 = Olearia Nerestii.
nervosus, *Mill. Gard. Dict.* ed. VIII. n. 28.—Am. bor.
nitidulus, *DC. in Wight, Contrib.* 9.—Ind. or.
nivalis, Adam, in Weber & Mohr, Beitr. i. 67 = alpinus.
Nolanus, Sch. Bip. ex Nym. Consp. 387 = A. Amellus.
notocenicus, *Griseb. in Goett. Abh.* xxiv. (1879) 178.—Reg. Argent.
Novae-Angliae, *Linn. Sp. Pl.* 875.—Am. bor.
Novae-Scotiae, *Forbes, Hort. Woburn.* 159; *Steud. Nom.* ed. II. i. 156.—N. Scotia.
Novi-Belgii, *Linn. Sp. Pl.* 877.—Am. bor.
nudicaulis, Lam. Encyc. i. 308 = Lagenophora Commersonii.
nudiflorus, Nutt. Gen. Am. ii. 157 = A. Radula.
Nuttallii, Torr. & Gray, Fl. N. Am. ii. 126 = adscendens.
obcordatus, F. Muell. Fragm. v. 69 = Olearia obcordata.
obliquus, DC. Prod. v. 287 = carneus.
obliquus, Nees, Syn. Aster. 27 = salicifolius.
oblongifolius, *Nutt. Gen. Am.* ii. 156.—Am. bor.
obovatus, Ill. Sketch, ii. 368 = reticulatus.
obovatus, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 95.—Sibir. altaic.
obtusatus, *Thunb. Prod. Pl. Cap.* 159.—Afr. austr.
occidentalis, *Torr. & Gray, Fl. N. Am.* ii. 164.—Am. bor. occ.
ochraceus, H. B. & K. Nov. Gen. et Sp. iv. 84 = Tetramolopium ochraceum.
Oculus-Christi, Cav. ex Steud. Nom. ed. II. i. 156 = Inula Oculus-Christi.
odontophyllus, Wall. Cat. n. 2985 = Senecio linifolius.
odorus, *All. Fl. Pedem.* i. 197.—Italia.
officinalis, All. l. c. 194 = Inula Helenium.

ASTER :—
oligantheminus, *F. Muell. Census*, 78.—Austral.
onustus, Nees, Gen. et Sp. Aster. 122 = Novi-Belgi l.
oolantagineus, Riddell, Syn. Fl. W. States, 56 = azureus.
opposititolius, Linn. Amoen. Acad. v. 369 = Cymbalariae.
orarius, F. Muell. Fragm. v. 78 = Olearia glutinosa.
oreganus, *Torr. & Gray, Fl. N. Am.* ii. 168.—Am. bor. occ.
orientalis, S. G. Gmel. It. i. 155. t. 36. f. 1 = Inula britannica.
ornatus, Hort. Angl. ex Steud. Nom. ed. II. i. 156 = Herveyi.
Ottonis, *Ec. Forst. ex Forbes, Hort. Woburn.* 159.—Hab.?
pallens, Willd. Enum. Hort. Berol. Suppl. 58 = patulus.
pallidulus, B. C. Vog. in Trew, Pl. Rar. t. 28 = cordifolius.
Palmeri, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 209.—Texas.
paludosus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 201. —Am. bor.
palustris, Lam. Fl. Fr. ii. 148 = A. Tripolium.
panduratus, *Nees, ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 258.—China.
paniculatus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 207 = Lindleyanus.
paniculatus, *Lam. Encyc.* i. 306.—Am. bor.
paniculatus, *Mill. Gard. Dict.* ed. VIII. n. 24.—Hab.?
paniculatus, Muhl. in Ill. Sketch, ii. 365 = sagittifolius.
paniculatus, Nutt. Gen. Am. ii. 156 = undulatus.
paniculatus, Willd. Sp. Pl. iii. 2035 = cordifolius.
pannonicus, Jacq. Hort. Vindob. i. 3 = A. Tripolium.
pannosus, F. Muell. Fragm. v. 85 = Olearia pannosa.
Pappei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 79.—Afr. austr.
papposus, Willd. ex Spreng. Syst. iii. 525 = Printzia Bergii.
Parryi, *A. Gray, in Am. Natural.* viii. (1874) 212.—Am. bor. occ.
parviflorus, *A. Gray, in Bot. Calif.* i. 392.—Mexic.
parviflorus, Darlingt. Fl. Cestr. 442 = diffusus.
parviflorus, Nees, Syn. Aster. 29 = Tradescanti.
patens, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 201.—Am. bor.
patentissimus, Lindl. in DC. Prod. v. 282 = patens.
Patterson‖, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 272.—Colorado.
paulus, *Lam. Encyc.* i. 306.—Am. bor.
pauciedentatus, F. Muell. Fragm. v. 66 = Olearia paucidentata.
paucifloras, Lindl. ex DC. Prod. v. 281 = sibiricus.
pauciflorus, Martens, in Bull. Acad. Brux. viii. (1841) 57 = ericoides.
pauciflorus, *Nutt. Gen. Am.* ii. 154.—Am. bor.; Mexic.
pauciflorus, Wall. Cat. n. 2967 = asperulus.
pellitus, *H. B. & K. Nov. Gen. et Sp.* iv. 91.—N. Granat.
pendulus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 205 = diffusus.
pensilvanicus, Pohr. Encyc. Suppl. i. 498 = laevis.
peregrinus, Less. in Linnaea, vi. (1831) 128 = foliaceus.
peregrinus, *Pursh, Fl. Am. Sept.* ii. 556.—As. or.; Am. bor. arct.
perfoliatus, Less. Syn. Comp. 167 = Mairia perfoliata.
pernonioides A. Cunn. in DC. Prod. v. 267 = Olearia pernonioides.
perthensis, *DC. Prod.* v. 287.—Hungar.
petiolatus, *Haw. Thes. Cap.* ii. 35. t. 154.—Afr. austr.
philadelphicus, *Bart. Fl. Phil. Prod.* 81.—Am. bor.
phlogifolius, Muhl. in Willd. Sp. Pl. iii. 2084 = patens.
phlogoppapus, Labill. Nov. Holl. Pl. ii. 49. t. 195 = Olearia stellulata.
phlogotrichus, Spreng. Syst. iii. 525 = Olearia stellulata.
phyllicoides, H. B. & K. Nov. Gen. et Sp. iv. 96 = Tetramolopium phyllicoides.
phyllolepis, *Torr. & Gray, Pl. N. Am.* ii. 118.—Am. bor.
pichinchensis, Willd. ex Nees, Gen. et Sp. Aster. 208 = Tetramolopium tenerimum.
pilosus, Willd. Sp. Pl. iii. 2025 = ericoides.

ASTER :—
pimeloides, A. Cunn. in DC. Prod. v. 268 = Olearia pimeleoides.
pinifolius, F. Muell. Fragm. v. 71 = Olearia pinifolia.
pinifolius, Nees, Syn. Aster. 29 = polyphyllus.
pinnatus, Cav. ic. iii. 6. t. 213 = Dysodia pinnatoides.
plantagineus, Poepp. ex DC. Prod. v. 418 = Senecio plantagineus.
plantaginifolius, Nutt. ex Nees, Gen. et Sp. Aster. 209 = Sericocarpus conyzoides.
plumiflorus, *G. Don, in Lond. Hort. Brit.* 346.—Afr. austr.
Pohlii, *Baker, in Mart. Fl. Bras.* vi. iii. 23.—Bras.
pollifolius *Linn. Sp. Pl.* ed. II. 1234 = Printzia Bergii.
politus, Nees, Syn. Aster. 28 = laevis.
polyphyllus, Moench, Meth. Suppl. 353 = macrophyllus.
polyphyllus, Pursh, ex DC. Prod. v. 245 = tenuifolius.
polyphyllus, *Willd. ' Enum. Hort. Berol.* 888.—Am. bor.
Porteri, *A. Gray, in Proc. Am. Acad.* xvi. (1880) 99. —Am. bor. occ.
potosinus, *A. Gray, l. c.* xv. (1880) 32.—Mexic.
pradensis, *Phil. in Linnaea*, xxviii. (1856) 729.—Chili.
praealtus, Torr. & Gray, Fl. N. Am. ii. 134 = Novi-Belgii.
praealtus, Poir. Encyc. Suppl. i. 493 = salicifolius.
praecox, Lindl. in Hook. Fl. Bor. Am. ii. 9 = Lindleyanus.
praecox, Mill. Gard. Dict. ed. VIII. n. 19 = pyrenaeus.
praecox, Willd. Enum. Hort. Berol. Suppl. 58 = patulus.
pratensis, *Rafin. Fl. Ludov.* 67.—Am. bor.
Preissii, *F. Muell. Fragm.* v. 66 = Olearia helophila.
prenanthoides, *Muhl. in Willd. Sp. Pl.* iii. 2046.—Am. bor.
Prescottii, Lindl. in DC. Prod. v. 281 = sibiricus.
procumbens, *Mill. Gard. Dict.* ed. VIII. n. 32.—Mexic.
propinquus, A. Cunn. ex DC. Prod. v. 270 = Olearia ramulosa.
pseudamellus, Wender. in Schrift. Marb. ii. (18 1) 226 = amelloides.
pseudamellus, *Hook. f. Fl. Brit. Ind.* iii. 249.—Ind. or.
pseudoamellus, DC. Fl. Fr. Suppl. v. 469 = A. Amellus.
pseudodiffusus, *E. Forst. ex Forbes, Hort. Woburn.* 159.—Hab.?
pseudodumosus, *E. Forst. ex Forbes, l. c.*—Hab.?
ptarmicoides, *Torr. & Gray, Fl. N. Am.* ii. 160.—Am. bor.
pubescens, Homem. Hort. Hafn., Suppl. 93 = heterophyllus.
pubescens, Lam. Encyc. i. 307 = tataricus.
pubescens, Moench, Meth. 600 = Inula squarrosa.
pubescens, Nees, Syn. Aster. 16 = sagittifolius.
pulchellus, *Eaton, in Bot. King, Exp.* 143. t. 16.—Am. bor. occ.
pulchellus, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 255 = alpinus.
pulchellus, Willd. Sp. Pl. iii. 2019 = Erigeron pulchellus.
pulchellus, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xxxvii. (1865) 1. 376 = flaccidus.
pulcherrimus, Ledb. Bot. Reis. i. t. 6 = linarifolius.
Pulicaria, All. Fl. Pedem. i. 197 = Pulicaria vulgaris.
pulicarius, Scop. Fl. Carn. ed. II. ii. 172 = Pulicaria vulgaris.
puberulentus, Hoffmgg. Verz. Pfl. Nachtr. i. (1824) 165 = Olearia stellulata.
pumilus, Banks & Soland. ex Hook. f. Fl. N. Zeal. i. 136 = Vittadinia australis.
pumilus, Flisch. Cat. Hort. Gorenk. (1906) 62; ed II. (1912) 39 = altaicus.
punctatus, Lepoyr. Hist. Abr. Pl. Pyr. 512 = Jasione perennis.
punctatus, Waldst. & Kit. Pl. Rar. Hung. ii. t. 109 = acris.
punctulosus, Ell. Sketch, ii. 355 = Elliottii.
puniceus, *Linn. Sp. Pl.* 875.—Am. bor.
purpurascens, *Sch. Bip. in Seem. Bot. Voy. Herald*, 156.—Mexic.
purpuratus, Nees, Gen. et Sp. Aster. 118 = virgatus.
purpureo-caeruleseens, *Hill, Herb. Kew.* 17.—Am. bor.
purpureus, Gmelinist. ex Ledeb. Fl. Ross. ii. 477 = A. Amellus?
purpureus, *Sweet, ex Steud. Nom.* ed. I. 33.—Hab.?

ASTER:—
pusillus, *Hornem. Hort. Hafn.* ii. 812.—Hab.?
pygmæus, *Lindl. in DC. Prod.* v. 278.—Am. arct.
pyrenæus, *DC. Fl. Fr.* iv. 146.—Mont. Pyren.
pyrenaicus, Pourr. in Mém. Acad. Tool. iii. (1788) 808 = Erigeron uniflorus.
pyrrhopappus, *Botan. Pl. Orient.* iii. 158.—Afghan.
quercifolius, Less. ex Walp. Rep. vi. 138 = Erigeron quercifolius.
quercifolius, Sieber, ex DC. Prod. v. 272 = Olearia stellulata.
quitensis, *Willd. ex Spreng. Syst.* iii. 533.—Ecuador.
racemosus, *Ell. Sketch*, ii. 848.—Am. bor.
Radula, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 210.—Am. bor.
Radula, Less. in Linnæa, vi. (1831) 125 = Chamisscoia.
radulinus, *A. Gray, in Proc. Am. Acad.* viii. (1872) 388.—Am. bor. occ.
ramosissimus, Mill. Gard. Dict. ed. VIII. n. 21 = ericoides.
ramosus, *G. Don, in Loud. Hort. Brit.* 846.—Am. bor.
ramulosus, Labill. Nov. Holl. Pl. ii. 51 = Olearia ramulosa.
ramulosus, Lindl. in DC. Prod. v. 243 = falcatus.
ramulosus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 398 = adscendens.
ranunculoides, *Moench, Meth. Suppl.* 254.—Hab.?
recurvatus, Gmelin. ex Nees, Gen. et Sp. Aster. 97 = bellidiflorus.
recurvatus, Nees, l. c. 98 = tenuifolius.
recurvatus, Purah, Fl. Am. Sept. ii. 556 = Tradescanti.
recurvatus, Spreng. in Schrad. Journ. Bot. (1800) 195 = squarrulosus.
recurvatus, Willd. Sp. Pl. iii. 2047 = diffusus.
reflexus, Linn. Sp. Pl. 877 = Felicia reflexa.
regalis, Salisb. Prod. 198 = Callistephus hortensis.
Regnellii, *Baker, in Mart. Fl. Bras.* vi. 111. 21.—Bras.
repens, *H. B. & K. Nov. Gen. et Sp.* iv. 90. t. 332.—N. Granat.
reticulatus, *Pursh, Fl. Am. Sept.* ii. 548.—Am. bor.
retroflexus, A. Cunn. ex Hook. f. Fl. N. Zeal. i. 148 = Senecio glastifolius.
retroflexus, *Lindl. in DC. Prod.* v. 242.—Am. bor.
Reversii, Hort. ex Decne. in Rev. Hortic. Sér. IV. i. (1852) 181 = ericoides.
revolutus, *F. Muell. Census*, 78.—Austral.
rhomboideus, Lindl. ex DC. Prod. v. 280 = tataricus.
Richardsonii, Spreng. Syst. iii. 528 = sibiricus.
rigidulus, Desf. Tabl. ed. II. 122 = salicifolius.
rigidulus, Nees, Syn. Aster. 29 = leucanthemus, paitulus.
rigidus, Linn. Sp. Pl. 874 (ex parte) = linarifolius.
rigidus, Moench, Meth. Suppl. (1802) 5 = inula salicina.
riparius, *H. B. & K. Nov. Gen. et Sp.* iv. 92.—Arizona; Mexic.
riparius, Nees, Gen. et Sp. Aster. 280 = pesthiensis.
rivularis, *Cass. ex Loud. Hort. Brit.* 847.—Am. bor.
rivularis, Less. in Linnæa, v. (1830) 148 = Erigeron repens.
robustus, Hort. Bonn. ex Nees, Gen. et Sp. Aster. 282 = strictus.
robustus, Hort. ex Nees, l. c. 300 = salicifolius.
roscus, *Bish. ex Steu. in Mém. Soc. Nat. Mosc.* iii. (1812) 248.—Reg. Caucas.
rosmarinifolius, Desf. Cat. Hort. Par. ed. III. 401 = Novæ-Angliæ.
rosmarinifolius, A. Cunn. ex DC. Prod. v. 268 = Olearia rosmarinifolia.
rosmarinifolius, Spreng. Syst. iii. 526 = obtusatus.
rotundifolius, *Thunb. Prod. Pl. Cap.* 160.—Afr. austr.
Roxburghii, Hook. f. in Hook. Ic. Pl. xi. 45 = Conandendron robustum.
ruber, *Forbes, Hort. Woburn.* 189.—Hab.?
rubicundus, *Forbes, l. c.*—Hab.?
rubricaulis, Lam. Encyc. i. 306 = lævis.
rufibarbis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 76.—Afr. austr.
rugulosus, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 236.—Japon.
rupestris, H. B. & K. Nov. Gen. et Sp. iv. 94 = Tetramolopium rupestre.
rusanovdorus, A. Cann. ex DC. Prod. v. 269 = Olearia glandulosa.
sagittifolius, Ell. Sketch, ii. 303 = undulatus.

ASTER:—
sagittifolius, *Wedem. ex Willd. Sp. Pl.* iii. 2035.—Am. bor.
salicifolius, *Ait. Hort. Kew.* iii. 203; *Lam. Encyc.* i. 306.—Am. bor.
salicifolius, Nees, Syn. Aster. 26 = longifolius.
salicifolius, Richards. App. Frankl. Journ. ed. I. 748 = juncens.
salicifolius, Scholl. Fl. Barb. Suppl. 238 = paniculatus.
salicinus, Nees, Gen. et Sp. Aster. 171, in syn. = salignus.
salicinus, Seop. Fl. Carn. ed. II. ii. 179 = Inula salicina.
salignus, Willd. Sp. Pl. iii. 2040 = paniculatus.
salinus, Schrad. Hort. Gotting. 9 = A. Tripolium.
salsuginosus, Hook. Fl. Bor. Am. ii. 6, partim = peregrinus.
salsuginosus, Less. in Linnæa, vi. (1831) 124 = sibiricus.
salsuginosus, Richards. ex R. Br. in App. Frankl. Journ. ed. I. 749 = Erigeron salsuginosus.
sanguineus, Willd. ex Spreng. Syst. iii. 533 = paniculatus.
saturetæfolius, Banks, in Reliq. Houston. t. 19 = Pectis saturetoides.
Sayianus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 394 = modestus.
scaber, Ell. Sketch, ii. 368 = undulatus.
scaber, *Moench, Meth. Suppl.* 253.—Hab.?
scaber, *Thunb. Fl. Jap.* 218.—Japon.
scaberrimus, *Less. in Linnæa*, v. (1830) 148.—Mexic.
scaberulus, *Miq. in Journ. Bot. Néerl.* i. (1861) 100.—China.
scabridus, *E. Mey. ex DC. Prod.* v. 225.—Afr. austr.
scabridus, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 47 = trinervius.
scabris, Mill. Gard. Dict. ed. VIII. n. 16 = alpinus.
scabrosus, Bertol. Misc. Bot. vi. 29 = Senecio carpoticides.
scandens, Jacq. f. ex Spreng. Syst. iii. 524 = carolinianus.
scapigerus, *Ledeb. Fl. Ross.* ii. 472.—Sibiria.
scaposus, Klatt, in Sitzb. Akad. Muench. (1878) 84 = molliusculus.
scapuiensis, Nutt. in Linnæa, xxxii. (1863) 378 = A. Amelloides.
Schreberi, Nees, Syn. Aster. 16 = macrophyllus.
scoparius, DC. Prod. v. 245 = multiflorus.
scoparius, Nees, Syn. Aster. 28 = diffusus, Tradescanti.
scopulorum, *A. Gray, in Proc. Am. Acad.* xvi. (1880) 93.—Am. bor. occ.
scorpioides, Scop. Fl. Carn. ed. II. ii. 169 = Doronicum scorpioides.
scorzonerifolius, *Regel, Gartenfl.* (1878) 1; *et in Act. Hort. Petrop.* ii. (1878) 808.—Mexic.
secundiflorus, Desf. Tabl. ed. II. 268 = viminens.
secundiramosus, Zeyh. ex Sch. Bip. Tanacet. 20 = Matricaria nigellæfolia.
sedifolius, Linn. Sp. Pl. 874 = acris.
seiiper, Nees, in Nenwied, Reise Nord. Am. ii. 439 = diffusus.
sericeus, *Vent. Jard. Cels*, t. 38.—Am. bor.
serotinus, Willd. Sp. Pl. iii. 2049 = Novi-Belgii.
serratifolius, Schloss. & Vukot. Syll. Fl. Croat. 57 = paniculatus.
serratus, Lag. ex DC. Prod. v. 315 = Grindelia inuloides.
serratus, *Thunb. Prod. Pl. Cap.* 160.—Afr. austr.
serrulatus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 73.—Afr. austr.
sessilifolius, *Rich. ex Steud. Nom.* ed. II. i. 157.—Am. bor.
setosus, *Baker, in Mart. Fl. Bras.* vi. III. 22.—Bras.
shastensis, *A. Gray, in Bot. Calif.* ii. 323.—Calif.
Shortii, *Lindl. ex Hook. Fl. Bor. Am.* ii. 9; *Riddell, Syn. Pl. W. States*, 55.—Am. bor.
sibiricus, Lam. Encyc. i. 305 = pyrenæus.
sibiricus, *Linn. Sp. Pl.* 872.—Sibir.; Am. bor. occ.
sibiricus, Turcz. ex Torr. & Gray, Fl. N. Am. ii. 107 = montanus.
Siemssenii, F. Muell. Fragm. v. 71 = Olearia viscidula.
sikkimensis, *Hook. f. & Thoms. in Bot. Mag.* t. 4657.—Ind. or.
sibuifolius, *Turcz. in DC. Prod.* v. 297.—Sibiria.
simplex, Eaton, in Bot. Mag. Expod. 140 = oreganus.

ASTER:—
simplex, Willd. Enum. Hort. Berol. 887 = paniculatus.
simulans, *Harv. in Harv. & Sond. Fl. Cap.* iii. 73.—Afr. austr.
sinuatus, A. Cunn. ex DC. Prod. v. 272 = Olearia quercifolia.
Solandri, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.?
solidagineus, Michx. ex Willd. Sp. Pl. iii. 2024 = Sericocarpus solidagineus.
solidaginoides, Pers. Syn. ii. 448 = Sericocarpus solidagineus.
Sonderi, F. Muell. Fragm. v. 83 = Olearia grandiflora.
Sonoræ, A. Gray, Pl. Wright. ii. 76 = riparius.
sparsiflorus, Michx. Fl. Bor. Am. ii. 112 = dumosus.
sparsiflorus, Purah, Fl. Am. Sept. ii. 547 = tenuifolius.
spartioides, C. B. Clarke, Comp. Ind. 46 = altaicus.
spathulares, Brouss. ex DC. Prod. v. 315 = Grindelia inuloides.
spathulatus, Hort. ex Steud. Nom. ed. II. i. 157 = Inula salicina.
spathulatus, Lag. ex Nees, Gen. et Sp. Aster. 283 = Grindelia inuloides.
spathulatus, *Lindl. in DC. Prod.* v. 231.—Am. arct.
spathulifolius, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 216.—Japon.
speciosus, Hornem. Hort. Hafn. ii. 816 = spectabilis.
spectabilis, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 209.—Am. bor.
spectabilis, Hook. & Arn. Bot. Beech. Voy. 146 = Chammissonis.
spectabilis, Willd. Sp. Pl. iii. 2048 = Novi-Belgii.
spinosus, *Benth. Pl. Hartw.* 90.—Am. bor.; Mexic.
spinulosus, *Chapm. Fl. S. U. St.* 199.—Florida.
spurius, Willd. Sp. Pl. iii. 2032 = Novæ-Angliæ.
squarrosulus, Nees, Gen. et Sp. Aster. 84 = longifolius.
squarrosus, All. Fl. Pedem. i. 196 = Inula spiræifolia.
squarrosus, Banks, ex Steud. Nom. ed. II. i. 157 = linarifolius.
squarrosus, *Walt. Fl. Carol.* 209.—Carolina.
Steetzii, F. Muell. Fragm. v. 56 = Olearia stirlingea.
stellulatus, Labill. Nov. Holl. Pl. ii. 50. t. 196 = Olearia stellulata.
stenomeres, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 209.—Am. bor. occ.
stenophyllus, Lindl. in DC. Prod. v. 242 = salicifolius.
Strachevi, *Hook. f. Fl. Brit. Ind.* iii. 250.—Reg. Himal.
striatus, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 238.—China.
strictus, Poir. Encyc. Suppl. i. 498 = paniculatus.
strictus, Purah, Fl. Am. Sept. ii. 556 = A. Radula.
strictus, Wender. in Schriff. Marb. ii. (1839) vi. 54 = dracunculoides.
strigosus, *Lichtenst. ex Harv. & Sond. Fl. Cap.* iii. 88.—Afr. austr.
strigosus, Thunb. Prod. Fl. Cap. 159 = Amellus annuus.
Stuartii, F. Muell. Fragm. v. 76 = Olearia Stuartii.
subasper, Lindl. ex Hook. Comp. Bot. Mag. i. (1835) 97 = salicifolius.
subcymosus, Bernh. ex Nees, Gen. et Sp. Aster. 138 = macrophyllus.
subspicatus, Nees, l. c. 74 = Douglasii.
subulatus, Hort. ex Steud. Nom. ed. II. i. 157 = ericoides.
subulatus, *Michx. Fl. Bor. Am.* ii. 111.—Am. bor.
succulentus, Gilib. Exercit. i. 174 = A. Tripolium.
surculosus, *Michx. Fl. Bor. Am.* ii. 113.—Am. bor.
tamœcetifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 95.—Calif.; Mexic.
tardiflorus, *Linn. Sp. Pl.* ed. II. 1221.—Am. bor.
tardiflorus, Willd. Sp. Pl. iii. 2049 = Novi-Belgii.
tataricus, *Linn. f. Suppl.* 373.—Tataria.
tataricus, Turcz. in Bull. Soc. Nat. Mosc. (1838) 94 = sibiricus.
taxifolius, Linn. Fl. Alt. 23; Amœn. Acad. vi. 102 = Mœria taxifolia.
tenellus, Linn. ll. cc. = Felicia fragilis.
tenellus, *Thunb. Fl. Cap.* 686.—Afr. austr.
tenax, *Harv. in Harv. & Sond. Fl. Cap.* iii. 80.—Afr. austr.
tenuerimus, Less. in Linnæa, iv. (1829) 120 = Tetramolopium tenerrimum.

ASTER :—
tenue, *Kellogg, in Proc. Calif. Acad.* vii. (1877) 114. —Calif.
tenuicaulis, *Bart. Fl. Phil. Prod.* 81.—Am. bor.
tenuifolius, Ell. Sketch, ii. 347 = vimineus.
tenuifolius, *Linn. Sp. Pl.* 873.—Am. bor.
tenuifolius, Nees, Syn. Aster. 110, partim = polyphyllus.
tenuifolius, Willd. Sp. Pl. iii. 2032 = ericoides.
tenuis, *Less. ex Ledeb. Fl. Ross.* ii. 483.—Rossia.
teretifolius, F. Muell. Fragm. v. 77 = Olearia teretifolia.
Thomsoni, *C. B. Clarke, Comp. Ind.* 48.—Reg. Himal.
thyrsiflorus, Hoffm. ; Phyt. Blaett. i. (1808) 88 = Novi-Belgii.
tibeticus, *Hook. f. Fl. Brit. Ind.* iii. 251.—Tibet. occ.
Tilesii, Wlehstr. in Nya Handl. Stockh. (1822) 18 = peregrinus.
tinctorius, Wallr. Beitr. Fl. Hercyn. 271 ; et in Linnaea, xiv. (1840) 641 = A. Amellus.
tomentellus, Hook. & Arn. Bot. Beech. Voy. 146 = Corethrogyne filaginifolia.
tomentosus, Schrad. in Wendl. Sert. Hann. 8. t. 24 = Olearia dentata.
tortifolius, *A. Gray, in Proc. Am. Acad.* vii. (1868) 353.—Am. bor. occ.
tortifolius, Michx. Fl. Bor. Am. ii. 109 = Sericocarpus tortifolius.
Townshendii, Hook. f. Bot. Mag. t. 6430 = Bigelovii.
Tradescanti, Hoffm. Phyt. Blaett. i. (1808) 68. t. A. f. 2 = patulus.
Tradescanti, *Linn. Sp. Pl.* 876.—Am. bor.
Tradescanti, Michx. Fl. Bor. Am. ii. 118 = diffusus.
Tradescanti, Nees, Syn. Aster. 98 = paniculatus.
trianthus, *Vell. Fl. Flum.* viii. t. 120.—Bras.
tricephalus, *C. B. Clarke, Comp. Ind.* 43.—Reg. Himal.
tricolor, *Sperng. in Flora*, xiii. (1830) 1. *Beil.* 15, nomen.—Afr. austr.
trifurcatus, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 317 = Senecio trifurcatus.
trilineatus, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 91.—Mexic.
trinervifolius, Less. in Linnaea, ix. (1834) 188 = paniculatus.
trinervis, *Desf. Tabl.* ed. II. 122.—Gallia.
trinervis, Ruiz & Pav. ex DC. Prod. vii. 22 (= *Diastanzis trinervis*).—Peruv.
trinervius, Gilib. Fl. Lithuan. iii. 214 = A. Amellus.
trinervius, *Roxb. Hort. Beng.* 61 ; *Fl. Ind.* iii. 433.—Reg. Himal.
Tripolium, *Linn. Sp. Pl.* 872.—Europ. ; Afr. bor. ; Orient ; Aa. bor.
Tripolium, Wall. Fl. Cand. 209 = tenuifolius.
tubuliflorus, F. Muell. Fragm. v. 85 = Olearia tubuliflora.
turbinatus, *S. Moore, in Journ. Bot.* xvi. (1878) 182. —China.
turbinellus, *Lindl. ex Hook. Comp. Bot. Mag.* i. (1835) 98.—Am. bor.
Turczaninowii, F. Muell. Fragm. v. 67 = Olearia imbricata.
turkestanicus, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 308.—Turkestan.
umbellatus, *Mill. Gard. Dict.* ed. VIII. n. 22.—Am. bor.
unalaschkensis, Less. in Linnaea, vi. (1831) 122 = peregrinus.
unalaschkensis, Less. ex Bong. Veg. Isl. Sitcha, 148 = salsuginosus.
uncinatus, Steud. Nom. ed. II. i. 157 = Novi-Belgii.
undulatus, Ell. Sketch, ii. 361 = patens.
undulatus, *Linn. Sp. Pl.* 875.—Am. bor.
undulatus, Mœnch, Meth. 600 = Inula britannica.
undiflorus, Michx. Fl. Bor. Am. ii. 110 = nemoralis.
undiflorus, Mœnch, Meth. 602 = Novi-Belgii.
uropkyllus, Lindl. in DC. Prod. v. 233 = sagittifolius.
vagans, *E. Forst. ex Forbes, Hort. Woburn.* 190.—Hab. ?
Vaillantii, All. Fl. Pedem. i. 196 = Inula Vaillantii.
Vahlii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 49.—Chili.
vernicosus, F. Muell. Fragm. v. 67 = Olearia passerinoides.
versus, Linn. Sp. Pl. 876 = Erigeron radicatus.
vernicolor, Hort. ex Steud. Nom. ed. II. i. 157 = mutabilis.

ASTER :—
versicolor, *Willd. Sp. Pl.* iii. 2045.—Am. bor.
villosus, *Sch. Bip. in F. Schultz, Arch. Fl.* 180.—Europ.
villosus, Michx. Fl. Bor. Am. ii. 118 = ericoides.
villosus, Thunb. Prod. Pl. Cap. 159 = Felicia angustifolia.
vimineus, *Lam. Encyc.* i. 306.—Am. bor.
vimineus, Nees, Gen. et Sp. Aster. 68 = tardiflorus.
vimineus, Willd. Sp. Pl. iii. 2046 = virgatus.
virgatus, *Ell. Sketch.* ii. 358.—Am. bor.
virgatus, Mœnch, Meth. Suppl. 250 = tenuifolius.
virgatus, Nees, Syn. Aster. 27 = salicifolius.
virgineus, Nees, l. c. 22 = longifolius.
viscosus, Labill. Nov. Holl. Pl. ii. 58 = Olearia viscosa.
Walpersianus, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 359.—China.
Warmingii, *Baker, in Mart. Fl. Bras.* vi. iii. 28.—Bras.
Watsoni, *A. Gray, Syn. Fl. N. Am.* i. ii. 201.—Am. bor.
Willdenowii, *E. Forst. ex Forbes, Hort. Woburn.* 190. —Hab. ?
Willkommii, *Sch. Bip. ex Willk. in Flora*, xxxiv. (1851) 742.—Hispan.
Wolfii, Favrat, in Bull. Soc. Murith. vii-viii. (1879) 4 = alpinus.
Wrightii, *A. Gray, Pl. Wright.* ii. 75.—Mexic. ; Texas.
xerophilus, F. Muell. Fragm. v. 76 = Olearia xerophila.
xylorrhiza, *Torr. & Gray, Fl. N. Am.* ii. 158.—Am. bor. occ.
Zeyheri, Less. Syn. Comp. 177 = Felicia Zeyheri.

ASTERACANTHA, Nees, in Wall. Pl. As. Rar. iii. 75 (1832) = **Hygrophila**, R. Br. (Acanth.).
auriculata, Nees, in DC. Prod. xi. 246 = H. spinosa.
longifolia, Nees, in Wall. Pl. As. Rar. iii. 90 = H. spinosa.
macracantha, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 146 = H. spinosa.

ASTERANDRA. Klotzsch, in Wiegm. Archiv, vii. (1841) 1. = **Pyllanthus**, Linn. (Euphorb.).
cornifolia, Klotzsch, l. c. = P. grandifolius.
Sellowiana, Klotzsch, l. c. = P. grandifolius.

ASTERANTHA, Reichb. Handb. 190 (1837) = ASTERACANTHA, Nees, = **Hygrophila**, R. Br. (Acanth.).

ASTERANTHEMUM, Kunth, Enum. Pl. v. 151 (1850) = **Smilacina**, Desf. (Liliac.).
dahuricum, Kunth, l. c. 158 = S. dahurica.
trifoliatum, Kunth, l. c. = S. trifolia.
vulgare, Kunth, l. c. 152 = S. stellata.

ASTERANTHERA, Hanst. in Linnaea, xxvi. (1853) 211. *GESNERACEAE*, Benth. & Hook. f. ii. 1011. chilœnsis, *Hanst. l. c.*—Chiloe.
ovata, *Hanst. l. c.* xxxiv. (1865–66) 417.—Chili.

ASTERANTHOS, Desf. in Mém. Mus. Par. vi. (1820) 9. t. 3. *MYRTACEAE*, Benth. & Hook. f. i. 724. ASTERANTUS, Spreng. Syst. iii. 568 (1826).
brasiliensis, *Desf. l. c.*—Bras.

ASTERANTHUS, Endl. Gen. 209 (1837) = ASTRANTHUS, Lour. = **Homaltium**, Jacq. (Samyd.).

ASTERIAS, Borckh. in Roem. Arch. i. 1. (1796) 25 = **Gentiana**, Linn.
hybrida, G. Don, Gen. Syst. iv. 164 = G. lutea.
lutea, Borckh. in Roem. Arch. i. 1. (1796) 25 = *G. lutea*.

ASTERIASTIGMA, Bedd. Forest. Man. Bot. 236 (1873). *BIXINEAE*.
macrocarpa, *Bedd. l. c.*—Ind. or.

ASTERIDEA, Lindl. Swan Riv. App. 24 (1839) = **Athrixia**, Ker-Gawl. (Compos.).
gracilis, A. Gray, in Hook. Kew Journ. iv. (1852) 274 = Athr. gracilis.
multiceps, A. Gray, l. c. = Athr. multiceps.
puberulenta, Lindl. Swan Riv. App. 24 = Athr. australis.
stricta, A. Gray, in Hook. Kew Journ. iv. (1852) 275 = Athr. stricta.

ASTERIDIUM, Engelm. ex Walp. Rep. ii. 958 (1843) = **Chaetopappa**, DC. (Compos.).

ASTERINGA, E. Mey. ex DC. Prod. vi. 138 (1837) = **Pentasia**, Thunb. (Compos.).

ASTERIOIDES, Ponted. ex Linn. Class. Pl. 530 (1747) = **Buphthalmus**, Linn. (Compos.).

ASTERIPHOLIS, Ponted. ex Linn. Class. Pl. 530 (1747) = **Aster**, Linn. (Compos.).

ASTERISCIUM, Cham. & Schlecht. in Linnaea, i. (1826) 254. t. 5. *UMBELLIFERAE*, Benth. & Hook. f. i. 877.
BUSTILLOSIA, Clos, in C. Gay, Fl. Chil. iii. 106. t. 32 (1847).
CASSIDOCARPUS, Presl, ex DC. Prod. iv. 83 (1830).
DIPTERYGIA, Presl, l. c. 80 (1830).
EREMOCHARIS, Phil. Fl. Atacam. 25. t. 2 B (1860).
GYMNOPHYTON, Clos, in C. Gay, Fl. Chil. iii. 102. t. 32 (1847).
TRITARRHICUM, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) t. 369.
aemocarpon, *Clos, in C. Gay, Fl. Chil.* iii. 101.—Chili.
anethoides, *Turcz. in Bull. Soc. Nat. Mosc. xx.* (1847) t. 170.—Chili.
chilense, *Cham. & Schlecht. in Linnaea*, i. (1826) 254. t. 5.—Chili.
flexuosum, *Hemsl. Diag. Pl. Nov.* 16.—Mexic.
isatidicarpum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 349.—Chili.
oemocarpum, Walp. Ann. i. 978 = aemocarpon.
Poeppigii, DC. Prod. iv. 82 = chilense.
polycephalum, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 332.—Chili.
porcides, *Clos, in C. Gay, Fl. Chil.* iii. 100.—Chili.
ramosissimum, *Phil. in Linnaea*, xxxiii. (1864–65) 90. —Chili.
robustum, *Baill. Hist. Pl.* vii. 145.—Chili.
verrucosum, Meyen, Reise, i. 315 = chilense.

ASTERISCUS, Moench, Meth. 592 (1794) = **Odontospermum**, Neck. (Compos.).
aquaticus, Less. Syn. Comp. 210 = O. aquaticum.
aureus, Lange, Pugill. ii. 118.—Hispan.
brachiatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 60 = O. maritimum.
Daltoni, Walp. Ann. ii. 844 = O. Daltoni.
flaccidus, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 209 = Anvillea Garcini.
graveolens, Less. Syn. Comp. 210 = O. graveolens.
imbricatus, DC. Prod. v. 467 = O. imbricatum.
lanuginosus, Aucher, ex DC. l. c. vii. 287 = O. lanuginosum.
littoralis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 60 = O. maritimum.
maritimus, Less. Syn. Comp. 210 = O. maritimum.
mauritanicus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 61 = O. maritimum.
nanus, Nym. Consp. 391 = O. aquaticum.
odorus, DC. Prod. v. 486 = O. odorum.
pygmæus, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 471.—Afr. bor. ; Palæst. ; Beluchist.
Schimperi, Boiss. Fl. Orient. iii. 180.—Arab.
sericeus, DC. Prod. v. 486 = O. sericeum.
sessilis, Mœnch, Meth. 592 = O. aquaticum.
Smithii, Walp. Ann. ii. 844 = O. Smithii.
Vogelii, Walp. l. c. 845 = O. Vogelii.

ASTERISCUS, Tourn. ex Sch. Bip. in Webb, Phytogr. Canar. ii. 229 (1836–50) = **Pallenis**, Cass. (Compos.).
spinosus, Sch. Bip. l. c. 230 = P. spinosa.

ASTERISCUS, Reichb. Consp. 144 (1828) = **Asteriscidium**, Cham. (Umbell.).

ASTEROCARPUS, Eckl. & Zeyh. Enum. 122 (1835) = **Pterocelastrus**, Meisn. (Celastrin.).
arborescus, Eckl. & Zeyh. l. c. 123 = P. variabilis.
Burmanni, Eckl. & Zeyh. l. c. 123 = P. tricuspidatus.
Hirsutis, Eckl. & Zeyh. l. c. = P. Hirsutis.
nervosus, Eckl. & Zeyh. l. c. 123 = P. variabilis.
rostratus, Eckl. & Zeyh. l. c. = P. rostratus.
stenopterus, Eckl. & Zeyh. l. c. = P. stenopterus.
tetrapterus, Eckl. & Zeyh. l. c. = P. variabilis.
tricuspidatus, Eckl. & Zeyh. l. c. 124 = P. rostratus.
typicus, Eckl. & Zeyh. l. c. 123 = P. tricuspidatus.

ASTEROCARPUS, Reichb. Handb. 261 (1837) ; Gren. & Godr. · Fl. Fr. i. 190 = **Astrocarpus**, Neck. (Resed.).

ASTEROCEPHALUS, [Vaill.] Adans. Fam. ii. 162 (1763) = **Scabiosa**, Linn. (Dipsac.).
acutiflorus, Reichb. Fl. Germ. Excurs. 194 = S. maritima.
africanus, Spreng. Syst. i. 380 = S. cretica.
agrestis, Spreng. l. c. 381 = S. graminifolia.
altissimus, Spreng. l. c. = S. africana.
ambiguus, Reichb. Fl. Germ. Excurs. 194 = S. maritima.
amoenus, Spreng. Syst. i. 381 = S. amoena.
arenarius, Vis. Pl. Aeg. 6. t. 1 = S. arenaria.
argenteus, Spreng. Syst. i. 382 = S. ucranica.
atropurpureus, Reichb. Ic. Fl. Germ. xii. t. 1368 = S. maritima.
atropurpureus, Spreng. l. c. 381 = S. atropurpurea.
bannaticus, Spreng. Syst. i. 382 = S. banatica.
Biebersteinii, Sweet, Hort. Brit. ed. I. 205 = S. micrantha.
bipinnatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 58 = S. ochroleuca.
brachiatus, Reichb. Fl. Germ. Excurs. 194 = S. brachiata.
brevicaulis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 57 = S. rotata.
canescens, Lag. Gen. et Sp. Nov. 8, in obs.; Spreng. Syst. i. 380 = S. suaveolens.
caucasicus, Spreng. l. c. 381 = S. caucasica.
cochinchinensis, Spreng. l. c. 380 = S. cochinchinensis.
Columbaria, Wallr. Sched. Crit. 48 = S. Columbaria.
Columnae, Reichb. Fl. Germ. Excurs. 195 = S. holoserices.
crenatus, Spreng. Syst. i. 382 = S. crenata.
creticus, Spreng. l. c. 380 = S. cretica.
daucoides, Spreng. l. c. 383 = S. daucoides.
dentatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 57 = S. stellata.
dichotomus, Lag. Gen. et Sp. Nov. 8 = S. dichotoma.
diffusus, Spreng. Syst. iv. Cur. Post. 38 = S. diffusa.
drosaicolus, Lag. ex Steud. Nom. ed. I. 84 ; ed. II. i. 456 = S. sicula.
eburneus, Spreng. Syst. i. 382 = S. ucranica.
elegans, Lag. Gen. et Sp. Nov. 8 = S. caucasica.
flavescens, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 61 = S. ochroleuca.
graminifolius, Spreng. Syst. i. 380 = S. graminifolia.
graminifolius, Spreng. l. c. i. 383 = S. graminifolia.
grandiflorus, Sweet, Hort. Brit. ed. I. 205 = S. maritima.
Hladnikianus; Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 171 = S. lucida.
holoseriocus, Spreng. Syst. i. 381 = S. holoserices.
holoseriocus, Webb, Itin, 40 = S. aphacioides.
industrius, Spreng. Syst. i. 380 = S. africana.
integrifolius, Lag. ex Steud. Nom. ed. II. i. 456 = S. integrifolia.
intermedius, Lag. Gen. et Sp. Nov. 8 = S. lusitanica.
lacteus, Spreng. Syst. i. 383 = S. lactea.
legionensis, Spreng. l. c. 379 = S. legionensis.
limonifolius, Spreng. l. c. 380 = S. limonifolia.
lobatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 57 = S. rotata.
lucidus, Spreng. Syst. i. 380 = S. nitens.
lyratus, Spreng. l. c. 381 = S. sicula.
maritimus, Spreng. l. c. 381 = S. maritima.
Mauroni, Sweet, Hort. Brit. ed. I. 206 = S. nitens.
micranthus, Spreng. Syst. i. 381 = S. micrantha.
mollissimus, Spreng. l. c. 383 = S. Columbaria.
monspeliensis, Zumag. Fl. Pedem. i. 169 ; Jord. & Fourr. Ic. Fl. Europ. i. 47 = S. monspeliensis.
montanus, Schur, Enum. Pl. Transs. 399 = S. ochroleuca.
ochroleucus, Wallr. Sched. Crit. 50 ; Spreng. Syst. i. 383 = S. lucida.
palaestinus, Spreng. l. c. 380 = S. palaestina.
pauciflorus, Loud. Hort. Brit. Suppl. ii. 581 = S. graminifolia.
pectinatus, Lag. Gen. & Sp. Nov. 8 = S. pectinata.
pilosus, Lag. l. c. = S. uchroleuca.
polystemon, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 56 = S. rotata.
proliferus, Spreng. Syst. i. 380 = S. prolifera.
pseudo-lucidus, Schur, Enum. Pl. Transs. 399 = S. lucida.
pumilus, Spreng. Syst. i. 383 = S. Columbaria.

ASTEROCEPHALUS:—
pycnocephalus, Sweet, Hort. Brit. ed. I. 205 = S. Columbaria.
rotatus, Spreng. Syst. i. 381 = S. rotata.
rutaefolius, Sweet, Hort. Brit. ed. I. 205 = S. rutaefolia.
sanctilis, Spreng. Syst. i. 379 = S. sanctilis.
Scopolii, Reichb. Fl. Germ. Excurs. 195 = S. Scopolii.
sericeus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 59 = S. graminifolia.
setifer, Spreng. Syst. i. 381 = S. setifera.
siculus, Spreng. l. c. = S. sicula.
silenifolius, Spreng. l. c. 380 = S. brachiata.
simplex, Spreng. l. c. 383 = S. monspeliensis.
sphacioticus, Spreng. l. c. 382 = S. sphaciotica.
stellatus, Reichb. Fl. Germ. Excurs. 194 ; et Ic. Fl. Germ. xii. t. 1370 = S. stellata.
stellulatus, Spreng. Syst. i. 381 = S. monspelliensis.
strictus, Spreng. l. c. 380 = S. lucida.
suaveolens, Lag. Gen. & Sp. Nov. 8, in obs.; Wallr. Sched. Crit. 51 = S. suaveolens.
Succisa, Wallr. Sched. Crit. 52 = S. Succisa.
tomentosus, Spreng. Syst. i. 381 = S. Succisa.
trinervifolius, Vis. in Mem. Ist. Venet. xxi. (1880–82) 483 = S. trinervifolia.
ucranicus, Spreng. Syst i. 382 = S. ucranica.
ucrolatus, Spreng. l. c. = S. rutaefolia.
westinus, Facch. ex Reichb. Ic. Fl. Germ. xii. 20. n. 1373 = S. vestina.
wiresceus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 59 = S. graminifolia.
Wulfenii, Spreng. Syst. i. 381 = S. ochroleuca.
Wulfenii, Reichb. Fl. Germ. Excurs. 194, in syn. n 1156 = S. ucranica.

ASTEROCHAETE, Nees, in Linnaea, ix. (1834) 300. *CYPERACEAE*, Benth. & Hook. f. Gen. iii. 1062.
ingestilis, *Nees, l. c.*—Afr. austr.
arundinacea, *Kunth, Enum. Pl.* ii. 212.—N. Caled.
capitellata, *Nees, in Linnaea*, ix. (1834) 300 ; x. (1836) 194.—Afr. austr.
elongata, *Kunth, Enum. Pl.* ii. 212.—Ins. Maurit.
glomerata, *Nees, in Linnaea*, ix. (1834) 300 ; x. (1836) 194—Afr. austr.
Ludwigii, *Steckst. in Flora*, xxviii. (1845) 759, in nota. —Afr. austr.
nitens, *Kunth, Enum. Pl.* ii. 212.—Ins. Maurit.
tenuis, *Kunth, l. c.* 212.—Afr. austr.

ASTEROCHITON, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 188 = **Thomasia**, J. Gay (Sterculi.).
Zygonemus, Turcz. l. c. 139 = T. pygmaea.

ASTEROCHLAENA, Garcke, in Bot. Zeit. viii. (1850) 668 = **Pavonia**, Cav. (Malvac.).
cuspidata, Garcke, l. c. 668.—Peruv.

ASTEROCYTISUS, Schur, ex Fuss, Fl. Transs. 154 (1866) = **Genista**, Linn. (Legumin.).
radiatus, Schur, ex Fuss, l. c. = G. radiata.

ASTEROGEUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 294 (1821) = **Plantago**, Tourn.
lacinialum, S. F. Gray, l. c. = P. Coronopus.

ASTEROGYNE, H. Wendl. ex Benth. & Hook. f. Gen. iii. 914 (1883). *PALMAE*, Benth. & Hook. f. iii. 914.

ASTEROLASIA, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 9; et Hook. Kew Journ. viii. (1856) 94. *FUTACEAE*, Benth. & Hook. f. i. 394, 920.
PLEURANDROPSIS, Baill. Adansonia, x. (1872) 305.
UROCARPUS, J. Drumm. in Hook. Kew Journ. viii. (1856) 64.
buxifolia, *Benth. Fl. Austral.* i. 351.—Austral.
chorilaenoides, F. Muell. in Trans. Vict. Inst. i. (1855) 116 = Microcybe pauciflora.
cneoroides, *Benth. Fl. Austral.* i. 350.—Austral.
grandiflora, *Benth. l. c.* 352.—Austral.
mollis, *Benth. l. c.* 351.—Austral.
Muelleri, *Benth. l. c.* 350.—Austral.
pallida, *Benth. l. c.* 352.—Austral.
phebalioides, *Benth. l. c.*—Austral.
phebalioides, F. Muell. in Trans. Phil. Soc. Vict. i. (1854) 10 = pleurandroides.
pleurandroides, *F. Muell. ex Benth. Fl. Austral.* i. 351. —Austral.
squamuligera, *Benth. l. c.* 352.—Austral.

ASTEROLASIA:—
trymalioides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 10.—Austral.

ASTEROLINION, Bronga. Enum. Gen. 69 (1843) = seq.

ASTEROLINON, Hoffmgg. & Link, Fl. Port. i. 332 (1820). *PRIMULACEAE*, Benth. & Hook. f. ii. 686.
PELLETIERA, A. St. Hil. in Mém. Mus. Par. ix. (1822) 365.
adoense, *Kunze, in Linnaea*, xx. (1847) 37.—Abyss.
Linum-stellatum, Duby, in DC. Prod. viii. 68 = stellatum.
lysimachioidum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 64 = A. stellatum.
serpyllifolium, *Bali, in Journ. Linn. Soc.* xxi. (1884) 226.—Patagon.
stellatum, *Hoffmgg. & Link, Fl. Port.* i. 333.—Lusitan.
trinum, *Baudo, in Ann. Sc. Nat.* Sér. II. xx. (1843) 350. —Chili.

ASTEROMAEA, DC. Prod. v. 302 (1836) = seq.

ASTEROMOEA, Blume, Bijdr. 901 (1826) = **Boltonia**, L'Hérit. (Compos.).
indica, Blume, l. c. = B. indica.
pekinensis, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 225 = B. pekinensis.

ASTEROMYRTUS, Schau. in Linnaea, xvii. (1843) 242 = **Malaleuca**, Linn. (Myrtac.).
Gaertneri, Schau. l. c. 248 = M. angustifolia.

ASTEROPEA, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 79 = seq.

ASTEROPEIA, Thou. Hist. Vég. Afr. 51. t. 15. (1807). *SAMYDACEAE*, Benth. & Hook. f. i. 801.
amblyocarpa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 81.—Madag.
densiflora, *Baker, in Journ. Bot.* xx. (1882) 49.—Madag.
multiflora, *Thou. Hist. Vég. Afr.* 51. t. 15.—Madag.

ASTEROPHYLLUM, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 1077 (1829) = **Asperula**, Linn. (Rubiac.), &c.
Aparine, Schimp. & Spenn. l. c. = Galium Aparine.
Asperula, Schimp. & Spenn. l. c. = Asperula odorata.
galioides, Schimp. & Spenn. l. c. in obs. = Asperula galioides.
Galium, Schimp. & Spenn. l. c. = Gallum verum.
Scherardianum, Schimp. & Spenn. l. c. = Sherardia arvensis.
sylvaticum, Schimp. & Spenn. l. c. in obs. = Asperula odorata.
tinctorium, Schimp. & Spenn. l. c. = Asperula tinctoria.

ASTEROPSIS, Less. Syn. Comp. 188 (1832) = **Podocoma**, Cass. (Compos.).
macrocephala, Less. l. c.—Bras.

ASTEROPTERUS, Adans. Fam. ii. 124 (1763) ; Gaertn. Fruct. ii. 460. t. 173 (1791) = **Leysera**, Linn. (Compos.).
Calliconna, Gaertn. Fruct. ii. 460 = L. Calliconna.
cerulens, Rafin. Fl. Tellur. ii. 48 = an praec. ?

ASTEROPUS, Schult. Mant. iii. 15 (1827) = ASTROPUS, Spreng. = **Waltheria**, Linn. (Stercul.).

ASTEROSCHOENUS, Nees, in Mart. Fl. Bras. ii. t. 124 (1842) = **Synechoporus**, Vahl (Cyperac.).
cenangeniensis, Nees, l. c. 125 = R. cenangeniensis.
rigidus, Nees, l. c. 126 = R. rigida.
speciosus, Nees, l. c. 125 = R. speciosa.

ASTEROSPERMA, Less. Syn. Comp. 389 (1832) = **Felicia**, Cass. (Compos.).
chrysocomoides, Less. l. c.—Afr. austr.

ASTEROSTEMMA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 271. t. 10. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 775.
repandum, *Decne. l. c.*—Java.

ASTEROSTIGMA, Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 148; Schott, in Oestr. Bot. Wochenbl. (1852) 67 = **Staurostigma**, Scheidw. (Arnld.).
colubrinum, Schott, in Bonplandia, x. (1862) 86 = S. concinnum.
concinnum, Schott, in Oestr. Bot. Wochenbl. (1852) 67 = S. concinnum.
Langsdorffii, C. Koch, in Ind. Sem. Hort. Berol. (1854) 8 = S. concinnum.
lineolatum, Schott, in Bonplandia, x. (1862) 86 = S. concinnum.
Luschnathianum, Schott, Syn. Aroid. 126 = S. Luschnathianum.
Pavonii, Schott, Prod. Aroid. 339 = S. Pavoni.
Tweedianum, Schott, in Oestr. Bot. Zeitschr. (1859) 89 = S. Tweedianum.
Vellozianum, Schott, Syn. Aroid. 126 = S. Luschnathianum.
vermiculosum, Griseb. in Goett. Abh. xix. (1874) 247 = S. vermiculosum.
zebrina, Linden, in Belg. Hortic. xv. (1865) 98.—Bras.

ASTEROSTOMA, Blume, Mus. Bot. Lugd. Bat. i. 50 (1849) = **Cobeckia**, Linn. (Melast.).
aspera, Blume, l. c. = O. aspera.
decandra, Blume, l. c. = O. decandra.
decumbens, Blume, l. c. = O. decumbens.
Kleinii, Blume, l. c. = O. aspera.
multiflora, Blume, l. c. = O. multiflora.
octandra, Blume, l. c. = O. octandra.
repens, Blume, l. c. = Melastoma repens.
rotundifolia, Blume, l. c. = O. rotundifolia.
virgata, Blume, l. c. = O. virgata.
Walkeri, Blume, l. c. = O. Walkeri.
Wightiana, Blume, l. c. = O. octandra.

ASTEROTHRIX, Cass. in Dict. Sc. Nat. xlviii. 434 (1827) = **Leontodon**, Linn. (Compos.).
asperrima, Cass. l. c. = *L. asperrimus*.
crispa, Foerr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 100 = L. crispus.
hispanica, DC. Prod. vii. 127 = L. hispanicus.

ASTEROTRICHION, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. 19. t. 8 (1841) = **Plagianthus**, Forst. (Malvac.).
sidoides, Klotzsch, l. c. = P. sidoides.

ASTHENOCHLOA, Buese, in Miq. Pl. Jungh. 367 (1854). *GRAMINEAE*, Benth. & Hook. f. Gen. iii. 1096.
tenera, *Buese, l. c.* 368.—Java.

ASTHOTHECA, Miers, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 254 = **Clusia**, Linn. (Guttif.).

ASTIANTHUS, D. Don, in Edinb. Phil. Journ. ix. (1843) 262 = **Tecoma**, Juss. (Bignon.).
longifolius, D. Don, l. c. = T. viminalis.

ASTILBE, Buch. Ham. in D. Don, Prod. Fl. Nep. 210 (1825). *SAXIFRAGEAE*, Benth. & Hook. f. i. 634.
HOTEIA, Mort. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 316. t. 11.
Aruncus, Trev. in Bot. Zeit. xiii. (1855) 819 = Spiraea Aruncus.
barbata, Hort.—Cf. Gard. Chron. (1871) 547 = japonica.
chinensis, *Franch. & Sav. Enum. Pl. Jap.* i. 144.—China; Japon.
decandra, *D. Don, Prod. Fl. Nep.* 211.—Am. bor.
japonica, *A. Gray, in Hook. Lond. Journ. Bot.* ii. (1843) 134.—Japon.
odontophylla, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 96 = chinensis.
podophylla, Baill. Hist. Pl. iii. 332 = Rodgersia podophylla.
rivularis, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 210.—Reg. Himal.
rubra, *Hook. f. & Thoms. ex Hook. Bot. Mag.* t. 4959.—Reg. Himal.
speciosa, Jungh. Java, i. (1853) 250 = rivularis.
Stolitzkai, *Kurz, ex Serm. in Journ. Bot.* v. (1867) 240 (Spiraea sp. ?).—Himal. bor. occ.
Thunbergii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 96.—Japon.

ASTIRIA, Lindl. in Bot. Reg. (1844) Misc. 27. t. 49. *STERCULIACEAE*, Benth. & Hook. f. i. 231.
ROSEA, *Lindl. l. c.*—Manrit.

ASTOMA, DC. Coll. Mém. v. 71. t. 17 (1829); et Prod. iv. 249 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 885.
ASTOMAEA, Reichb. Handb. 218 (1837). sessiliflium, *DC. ll. cc.*—Syria.

ASTORGANTHUS, Endl. Cat. Hort. Vindob. ii. 196 (1843) = **Malloape**, Forst. (Rutac.).
Hurgelii, Endl. l. c. = M. simplex.

ASTRADELPHUS, Rémy, in Ann. Sc. Nat. Sér. III. xii. (1849) 185 = **Erigeron**, Linn. (Compos.).
calbucanus, Phil. in Anal. Univ. Chil. (1873) 485.—Chili.
chilensis, Rémy, in Ann. Sc. Nat. Sér. III. xii. (1849) 185 = E. chilensis.

ASTRAEA, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = **Croton**, Linn. (Euphorb.).
divaricatum, Klotzsch, l. c. = C. Klotzschii.
diversifolia, Klotzsch, l. c. = C. Klotzschii.
glandulifera, Klotzsch, ex Wawra, Bot. Ergeb. Maxim. Bras. ii. t. 42 = C. Klotzschii.
Jatropha, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. carnosus.
Klotzschii, Didr. in Kjoeh. Vidensk. Meddel. (1857) 137 = C. Klotzschii.
lobatum, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. lobatus.
Manihot, Klotzsch, l. c. = C. lobatus.
palmatum, Klotzsch, l. c. = C. lobatus.
Paulsna, Didr. in Kjoeh. Vidensk. Meddel. (1857) 138 = C. Paulsna.
pruniphila, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. Klotzschii.
Seemanni, Klotzsch, in Seem. Bot. Voy. Herald, 108 = C. lobatus.
tomentosum, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. comosus.

ASTRAEA, Schau. in Linnaea, xvii. (1843) 238 = **Thryptomene**, Endl. (Myrtac.).
saxicola, Schau. l. c. 239 = T. saxicola.

ASTRAGALOIDES, Adans. Fam. ii. 323 (1763) = **Astragalus**, Tourn. (Legum.).
Alopecurus, Moench, Meth. 168 = Astragalus alopecuroides.
alpina, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 375 = Astragalus alpinus.
australis, Medic. l. c. = Astragalus australis.
bicolor, Moench, Meth. Suppl. 87 = Astragalus uliginosus.
Cicera, Moench, Meth. 168 = Astragalus Cicer.
frigida, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 378 = Astragalus frigidus.
Glaux, Moench, Meth. 168 = Astragalus Glaux.
syphilitica, Moench, l. c. = Astragalus exscapus.

ASTRAGALUS, Tourn. ex Linn. Gen. ed. I. 215 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 536.
ALLUSCHIA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 151.
ALOPECIAS, Stev. l. c. 143 (1856).
AMMODYTES, Stev. l. c. iv. (1839) 263.
ANKYLOSUS, Stev. l. c. xxix. (1856) ii. 148.
ARAGALUS, Neck. Elem. iii. 12 (1790).
ARAGUS, Steud. Nom. ed. II. i. 118 (1840).
AULOSEMA, Walp. Rep. i. 694 (1842).
CARYOLOBIUM, Stev. in Bull. Soc. Nat. Mosc. iv. (1839) 266.
CHONDROCARPUS, Stev. l. c. 268 (1839).
CRACCINA, Stev. l. c. xxix. (1856) ii. 144.
CYMBICARPOS, Stev. l. c. iv. (1839) 266.
CYSTIUM, Stev. l. c. xxix. (1856) ii. 147.
DIDYMOPELTA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 469.
DIPELTA, Regel & Schmalh. l. c. 378 (1877).
DIPLOTHECA, Hochst. in Flora, xxix. (1846) 696.
EROPHACA, Boiss. Voy. Bot. Espagne, 176 (1839-45).
EULIUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 149.
EUPAEPIA, Stev. l. c. iv. (1839) 263.
ASTRAGALUS:—
FEIDANTHUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 148.
GLANDULA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 374.
GLOTTIS, Medic. l. c. 377 (1787).
GLYCYPHYLLA, Stev. in Bull. Soc. Nat. Mosc. iv. (1839) 266.
HAMAMA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364.
HAMOSA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 376.
HEDYPHYLLA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 142.
HOMALOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 350 (1840).
HYPOGLOTTIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364.
KENTROPHYTA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 358 (1840).
KIRCHNERIA, Opiz, in Lotos, viii. (1858) 65.
MACROSEMA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 146.
MEDYPHYLLA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 150.
MYOBROMA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 150.
ONIX, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 378.
PEDINA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 144.
PHACA, Roy. ex Linn. Coroll. 18 (1737).
PHILEMMOS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 146.
PHYSODRA, Rafin. Atl. Journ. 145 (1832).
PICHANA, Stev. in Bull. Soc. Nat. Mosc. iv. (1839) 263.
PODOCHERA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364.
PODOLOTUS, Royle, Illustr. Bot. Himal. 198 (1835).
PROSELIAS, Stev. in Bull. Soc. Nat. Mosc. iv. (1839) 263.
PSYCHIDIUM, Stev. l. c. 263 (1839).
RYSODIUM, Stev. l. c. (1839).
SEWERZOWIA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 380.
SOLENOTUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 148.
SPIESIA, Neck. Elem. iii. 13 (1790).
STELLA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 377.
THIUM, Steud. Nom. ed. I. 833 (1821).
TIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 373.
TRIQUETRA, Medic. l. c. 376 (1787).
XEROPHYRA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 150.
abakanensis, Less. ex Bunge, Astrag. ii. 23 = multicaulis.
abbreviatus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) ii. 348.—As. temp.
aboriginorum, *Richards. in Frankl. 1st Journ. App.* 746.—Am. bor. arct.
abyssinicus, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 198.—Abyss.
acanthostachys, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 236 = Roxanaeanus.
acaulis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 132.—Reg. Himal.
aciculatus, *Bunge, Astrag.* i. 90; ii. 161.—Cappadoc.
acinaciformis, *Boiss. Diagn.* Ser. I. ix. 73.—Arab.
acmonotrichus, *Fenzl, Pugill. Pl. Nov. Syr.* 5.—Cilicia.
acmophyllus, *Bunge, Astrag.* i. 98; ii. 166.—Cappadoc.
acmotrichus, Bunge, l. c. i. 109; ii. 169 = acmonotrichus.
acutiflorus, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 130 = Falconeri.
acutifolius, *Bunge, Astrag.* i. 130; ii. 237.—Persia.
acutifolius, DC. ex Steud. Nom. ed. I. 38 = A. Stella.
acutilobus, *Schischk. Cat. Pl. Helv.* ed. IV. 9.—Europ.
acutirostris, *S. Wats. in Proc. Am. Acad.* xx. (1885) 360.—Calif.
aduncus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
adenmacifolius, *Benth. ex Bunge, l. c.* i. 4; ii. 2.—Tibet. occ.
adpressus, Ehrenb. ex Walp. Ann. iv. 502 = guentheri.
adiandrum, *Schischk. Cat. Pl. Helv.* ed. IV. 9.—Europ.
adsurgens, *Pall. Astrag.* 19. t. 19. f. 1 (1800).—Sib.
adsurgens, Pall. ex Willd. Sp. iii. 1407 = Onobrychis.
aegiceras, Bunge, Astrag. i. 71; ii. 118 = Roxanaeanus.
aegobromus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
aegopodioides, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
aegyptiacus, *Lam. Encycl.* i. 284 (1783).—Egypt.
aemulans, *Karel. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) ii. 348.—As. temp.
aeneus, *Steph. ex Willd. Sp.* iii. 1407.—Sib.
aequicostatus, *Hook. f. & Thoms. ex Hook. Bot. Mag.* t. 4959.—Reg. Himal.
aequifoliolatus, *Boiss. Diagn.* Ser. I. ix. 73.—Arab.
aequivalvis, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
aerarius, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
aetnensis, *Presl, Del. Prag.* 124 (1822).—Sicil.
affinis, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
affinis, Hook. & Arn. ex Benth. in Hook. Lond. Journ. Bot. i. 113 = Hookeri.
africanus, Willd. Sp. iii. 1407 = Tragacantha.
agardhianus, *Fisch. & Mey. Ind. Sem. Petrop.* ii. 48 (1835).—Persia.
agathocles, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
agglutinatus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
agnicus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
agraphocalyx, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
agrestis, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
agrimoniifolius, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
agrimonioides, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
agrophilus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
agyrophorus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
ahmarensis, Schweinf. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 130 = Falconeri.
ajanensis, Fisch. ex Ledeb. Fl. Ross. i. 707 = Tilingii.
akmolinensis, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
alaicus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
alatavicus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albanus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albicaulis, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albicomus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albidus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albiflorus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albispinus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albocaudatus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albocephalus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albolanatus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
albovillosus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
alcedonius, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
alchidophilus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
alcyonius, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
alexandrinus, *Boiss. Diagn.* Ser. I. ix. 73.—Arab.
alexandrinus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 817.—Persia.

ASTRAGALUS:—

adsurgens, *Pall. Astrag.* 40. n. 44. t. 81.—Sibir.; Am. bor.

adsurgens, Torr. in Stansbury, Rep. 385 = lodanthus.

adsurgens, Willd. ex Steud. Nom. ed. II. i. 159 = virgatus.

aduuciformis, *Boiss. Diagn.* Ser. I. ii. 42.—As. Min.

aduncus, Bunge, Astrag. i. 102; ii. 179 = usturtensis.

aduncus, Bieb. Fl. Taur. Cauc. ii. 195 = Bungeanus.

aduncus, Hohen. Talysch. 107 = Cephalotes.

aduncus, *Willd. Sp. Pl* iii. 1389.—Reg. Caucas.

adustus, *Bunge, Astrag.* i. 89; ii. 134.—Kurdistan.

Aegiceras, *Willd. Enum. Hort. Berol. Suppl.* 52 = Ins. Chio.

aegobromus, *Boiss. & Hohen. Diagn.* Ser. I ix. 74.—Persia.

aegyptiacus, Mill. Gard. Dict. ed. VIII. n. 11 = annularis.

aegyptiacus, Spreng. Syst. iii. 298 = kahiricus.

aduropus, *Bunge, Astrag.* i. 89; ii. 157.—Reg. Caucas.; Persia.

aesicarus, *Pall. Reise*, iii. 751.—Sibir.

affghanus, *Boiss. Fl. Orient.* ii. 1093 (emend.).—Beluchist.

affinis, *Steud. Nom.* ed. II. i. 159.—Chili.

africanus, *Bunge, Astrag.* i. 61; ii. 104.—Algeria.

Agraniotii, *Orph. ex Boiss. Diagn.* Ser. II. ii. 29.—Laconia.

agrestis, Dougl. ex Hook. Fl. Bor. Am. i. 148 = hypoglottis.

aintabicus, *Boiss. Diagn.* Ser. I. ix. 86.—Syria; Mesopotam.

Aitchisoni, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 121.—Ind. bor. occ.

Ajfreidii, *Aitch. & Baker, in Journ. Linn. Soc.* xix. (1882) 157.—Afghanist.

ajubensis, *Bunge, Astrag.* i. 61; ii. 102.—Persia.

akkensis, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 57, nomen.—Afr. bor.

akmensis, *Bunge, Astrag.* i. 25; ii. 30.—Sibir. altaic.

alaschanus, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 31.—Mongolia.

alatavicus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 344.—Alatau.

albens, *Greene, in Bull. Calif. Acad.* i. ('1885') 156.—Calif.

Alberjilla, *Steud. Nom.* ed. II. i. 159.—Chili.

Alberti, *Bunge, in Act. Hort. Petrop.* vii. (1880) 375.—Turkestan.

albicans, *Bong. Vers. Pfl. Saisang-Nor*, 21. t. 2.—Soongaria.

albicaulis, *DC. Astrag.* 182. n. 59. t. 21.—Reg. Caucas.

albicaulis, Ledeb. Fl. Ross. i. 638 = medius.

albidus, Waldst. & Kit. Pl. Rar. Hung. i. 39. t. 40 = vesicarius.

alepplcus, *Boiss. Diagn.* Ser. I. ii. 58.—Syria; Mesopotam.

alexandrinus, *Boiss. l. c.* ix. 75.—Aegypt.; Arab.; Syria.

algarbiensis, *Coss. ex Bunge, Astrag.* i. 9; ii. 6.—Hispan.

alienus, *A. Gray, Bot. U. St. Expl. Exped.* i. 418.—Peruv.

alindanus, *Boiss. Diagn.* Ser. I. ii. 86.—Caria.

allochorus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 366.—Am. bor.

Alopecias, C. Koch, ex Bunge, Astrag. ii. 97 = maximus.

Alopecias, *Pall. Astrag.* 12. t. 9.—Persia; Turkestan.

alopecuroides, Ledeb. Fl. Ross. i. 633. n. 86 = maximus (ex parte).

alopecuroides, *Linn. Sp. Pl.* 755.—Sibiria; Helv.; Kamtschatca.

alopecuroides. Pall. Astrag. 9. t. 7 = vulpinus.

alopecuroideus. Lam. Fl. Fr. ii. 636 = narbonnensis.

alopecurus, *Pall. ex DC. Astrag.* 11. t. 8.—As. med.

Alpamarcas, *A. Gray, Bot. U. St. Expl. Exped.* i. 417.—Peruv.

alpinus, Ledeb. Fl. Ross. i. 601 = arcticus.

alpinus, *Linn. Sp. Pl.* i. 760.—Reg. bor. et arct.

alpinus, Vill. Hist. Pl. Dauph. i. 303 = A. Onobrychis.

alsaticus, *Bunge, Astrag.* i. 27; ii. 46.—Mont. Alsat.

altaicus, Pall. Astrag. 56 = Oxytropis brevirostra.

altaicoides, Kotschy, ex Walp. Ann. ii. 374 = elbrusensis.

altaicoides, *Lam. Encyc.* i. 317.—Armenia.

ASTRAGALUS:—

amalecitanus, *Boiss. Diagn.* Ser. I. ix. 46.—Palaestin.

amanus, *Boiss. Fl. Orient.* ii. 990.—Syria.

amarus, *Pall. Astrag.* 8. t. 6.—Reg. Casp.

amatus, *Clos, in C. Gay, Fl. Chil.* ii. 115.—Chili.

ambiguus, *Bunge, Astrag.* i. 86; ii. 150.—Syria.

ambigruus, Pall. Astrag. 54 = Oxytropis ambigua.

amblolepis, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 372.—Kurdistan.

Amherstianus, *Benth. in Royle, Illustr. Bot. Himal.* 199.—Reg. Himal.

ammocryptus, Boiss. Diagn. Ser. I. ix. 59. = tribuloides.

Ammodendron, *Bunge, Astrag.* i. 130; ii. 228.—As. med.

Ammodytes, *Pall. Voy. Ed. Gall. App.* n. 372. t. 88.—Sibir. or.

ammophilus, Bieb. ex Bess. in Flora, xv. (1832) ii. Beibl. 38 = virgatus.

ammophilus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 344.—Persia. Soongaria.

ammotrophus, *Bunge, Astrag.* i. 112; ii. 192.—Turkestan.

amoenus, *Fenzl, Pugill. Pl. Nov. Syr.* 4.—Cilicia.

amphiloxus, *Bunge, Astrag.* i. 90; ii. 161.—Anatolia.

amphoxys, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 366.—Am. bor.

ampullaticus, *S. Wats. in Am. Natural.* vii. (1873) 800.—Calif.

ampullatus, Pall. Reise, i. 750 = Oxytropis ampullata.

ampullifer, Griseb. Spicil. Fl. Rumel. i. 58 = physocalyx.

amygdaliger, Lesa. in Linnaea, ix. (1834) 176 = testiculatus.

amygdalinus, *Bunge, Astrag. Fedsch.* 238.—As. temp.

anacamptus, *Bunge, Astrag.* i. 110; ii. 190.—Persia.

anacardius, *Bunge, l. c.* i. 81; ii. 105.—Persia.

anatolicus, *Boiss. Diagn.* Ser. I. ii. 77.—Anatolia.

ancistrocarpus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 668.—Mesopotam.

ancistrumi, *Nouv. Mat. Fl. Atl.* 186.—Algeria.

ancocarpus, *Pomel, l. c.* 187.—Algeria.

ancylens, *Boiss. Fl. Orient.* ii. 640.—Cappadoc.

andalanicus, *Boiss. & Hausskn. ex Boiss. l. c.* 845.—Persia.

Andersonii, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 594.—Calif.

Andrusowi, *Bunge, Astrag.* i. 89; ii. 158.—Syria; Mesopotam.

andrachnesolius, *Fenzl, Pugill. Pl. Nov. Syr.* 4.—As. Min.

anemophilus, *Greene, in Bull. Calif. Acad.* i. (1885) 156.—Calif.

angustcuspus, Bunge, Astrag. i. 125; ii. 218 = strictus.

angarensis, *Turcz. ex Bunge, l. c.* i. 126; ii. 222.—Sibir. baical.

angorensis, Bunge, l. c. ii. 176 = leucocyaneus.

angulosus, *DC. Astrag.* 234. t. 45.—Syria.

angustatus, Boiss. Diagn. Ser. I. ix. 47 = ruscifolius.

angustatus, *Bunge, Astrag.* i. 125; ii. 220.—Persia.

angustiflorus, *C. Koch, in Linnaea*, xv. (1840) 720.—Armenia; Persia.

angustifolius, *Lam. Encyc.* i. 321.—Graecia; As. Min.

angustissimus, Bunge, Astrag. i. 125; ii. 222 = oxypetalus.

anisacanthus, *Boiss. Diagn.* Ser. I. ix. 108.—Afghan.

anisacanthus, Bunge, Astrag. ii. 138 = lutarius.

anisomerus, *Bunge, Astrag. Fedsch.* 223.—As. temp.

ankylotus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 37.—Iberg. Cauc.

ankylotus, Stachegl. in Bull. Soc. Nat. Mosc. (1854) i. 163 = commixtus.

annularis, *Forsk. Fl. Aegypt. Arab.* 139.—Aegypt.; Syria.

annuus, *DC. Astrag.* 127. n. 25.—Hab.?

anomalus, *Bunge, Astrag.* i. 85; ii. 62.—Tibet. occ.

anserinoides, *Boiss. Diagn.* Ser. I. ii. 76.—Persia.

antabicus, *Boiss. l. c.* ix. 86.—Anatolia.

anthylloides, *Lam. Encyc.* i. 320.—As. Min.

anthylloides, Pall. Astrag. 15 = Inflatus.

Antilibani, *Bunge, Astrag.* i. 55; ii. 90.—Syria.

Antiselli, *A. Gray, in Bot. Calif.* i. 192.—Calif.

antiochius, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 348.—Am. bor. occ.

Apollinem, *Boiss. Diagn.* Ser. II. ii. 27.—Graecia.

aprieus, *Bunge, Astrag.* i. 40; ii. 56.—Persia.

ASTRAGALUS:—

arabicus, Ehrenb. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 121 = prolixus.

arabicus, Ehrenb. ex Bunge, Astrag. i. 9; ii. 6 = fatinensis.

arabicus, *Kotschy, in Sitzb. Acad. Wien, Math.-Nat. lii. Abth.* i. (1866) 284.—Arabia.

arborescens, Bunge, Astrag. ii. 235 = paucijugus.

Arbuscula, *Pall. Astrag.* 19. n. 24. t. 17.—Sibir.

Arbuscula, Willd. ex Ledeb. Fl. Ross. i. 638 = fruticosus.

arcticus, *Bunge, Astrag.* i. 23; ii. 27.—Europ. arct.

arcticus, Spreng. Syst. iv. Cur. Post. 288 = Oxytropis uralensis.

arcticus, *Willd. Enum. Hort. Berol. Suppl.* 51.—Hab.?

arcuatus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 407.—As. Min.

arcuatus, Sieph. ex Ledeb. Fl. Ross. i. 649 = scaberrimus.

arenarius, Lapeyr. Hist. Abr. Pl. Pyr. 429 = baionensis.

arenarius, *Linn. Sp. Pl.* 759.—Dania; Suecia.

arenarius, O. F. Mssll. in Fl. Dan. t. 614 = danicus.

arenarius, Pall. Astrag. t. 84 = hypoglottis.

arenicola, *Pomel, Nouv. Mat. Fl. Atl.* 184.—Algeria.

arenosus, S. G. Gmel. ex Pall. Astrag. 35 = hyrcanus.

arequipensis, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 17.—Peruv.

argaeus, *Boiss. & Bal. Diagn.* Ser. II. vi. 53.—Cappadoc.

argenteus, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) 11. 28.—As. temp.

argentatus, Pall. Astrag. 60. t. 48 = Oxytropis argentata.

argenteus, Bertol. ex Vis. in Flora, xli. (1829) i. Erg. 18 = Muelleri.

argophyllus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 331 = glareosus.

argurieus, *Bunge, Astrag.* i. 103; ii. 181.—Mont. Ararat.

argutensis, *Bunge, l. c.* i. 98; ii. 170.—Armenia.

argyrophyllus, *Boiss. & Gaill. ex Boiss. Fl. Orient.* ii. 858.—Syria.

argyrostachyus, *Boiss. Diagn.* Ser. I. vi. 41.—Persia.

argyrothamnus, *Boiss. l. c.* ix. 89.—Syria.

aricus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 228.—Am. bor. occ.

aristatus, *L'Hérit. Stirp. Nov.* 170.—Europ. austr.

aristatus, Sieber, ex Steud. Nom. ed. II. i. 159 = angustifolius.

aristatus, Sibth. & Sm. Fl. Graec. vii. 25. t. 734 = Parnassi.

arizonicus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 398.—Arizona.

arkalycensis, *Bunge, Astrag.* i. 139; ii. 238.—As. temp.

arnatus, *Willd. Sp. Pl.* iii. 1330.—Numidia.

armenicus, *Boiss. Diagn.* Ser. I. ii. 86.—Armenia.

armeriastrum. Decne. ex Boiss. Fl. Orient. ii. 887 = cephalanthus.

arnacantha, *Bieb. Fl. Taur. Cauc.* ii. 205.—Reg. Caucas.

arnacreas, Bunge, Astrag. t. 12; ii. 14 = brachycreas.

aryllobus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 338.—As. temp.

arrectus, *A. Gray, in Proc. Am. Acad.* viii. (1870) 389.—Am bor.

Arthu-Schottii, A. Gray, l. c. vi. (1864) 209 = Coulteri.

artipes, *A. Gray, l. c.* xiii. (1878) 370.—Am. bor.

asaphes, *Bunge, Astrag. Fedsch.* 251.—As. temp.

ascocalyx, *Bunge, Astrag.* i. 137; ii. 234.—As. Min.

ascophorus, Fisch. ex Bunge, l. c. ii. 60 = brachystachys.

askius, *Bunge, l. c.* i. 120; ii. 208.—Persia.

aspadanus, *Bunge, l. c.* i. 63; ii. 107.—Persia.

asper, *Jacq. Ic. Rar.* i. 13. t. 152.—Europ. austr.; Reg. Caucas.

asperulus, Dufour, in Ann. Sc. Gén. vii. (1830) 295 = A. epiglottis.

Asterias, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 367 = cruciatus.

astragoides, *Pomel, Nouv. Mat. Fl. Atl.* 186.—Algeria.

atlanticus, *Ball, in Journ. Bot.* xi. (1873) 306.—Afr. bor. occ.

atropurpureus, *S. Wats. in Bot. King, Exp.* 69.—Am. bor. occ.

atropatanus, *Bunge, Astrag.* i. 83; ii. 138.—Persia.

ASTRAGALUS:—

atropatanus, Fisch. ex Bunge, Astrag. ii. 90 = pulchellus.
atropurpureus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 54.—As. Min.
atticus, Nym. Consp. 190 = Wulfenii.
Aucheri, *Boiss. Diagn.* Ser. I. ii. 46.—Armenia.
Auchervianus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 393. = pilosdes.
augustus, *Bunge, Astrag.* i. 86; ii. 45.—Afghanist.
aulacolobus, *Boiss. Diagn.* Ser. I. ix. 54.—Syria; Mesopotam.
aureus, *Willd. in Mém. Acad. Berl.* (1794–95) 29. t. 1. —Oriens.
Austinæ, *A. Gray, Bot. Calif.* i. 156.—Calif.
australis, *Lam. Fl. Fr.* ii. 587; *Bunge, Astrag.* i. 22; ii. 24.—Mont. Pyren.; Hispan.
austriacus, Delarb. Fl. Auv. ed. II. 466 = hamosus.
austriacus, *Linn. Sp. Pl.* 1070.—Europ.; Reg. Caucas.; Sibiria.
austriacus, Pall. ex Stend. Nom. ed. II. i. 159 = versicolor.
australiacus, Thore, ex DC. Prod. ii. 283 = baionnensis.
bachtiaricus, *Bunge, Astrag.* i. 38; ii. 53.—Persia.
bactrianus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 854.—Turkestan.
baeticus, *Linn. Sp. Pl.* ed. II. 1068.—Hispan.; Sicili.
baghensis, *Bunge, Astrag.* i. 80; ii. 135.—Persia.
balbut ensis, *Bunge, l. c.* i. 91; ii. 162.—Armenia.
balasalensis, *Bunge, l. c.* i. 139; ii. 238.—As. centr.
baicalensis, Pall. Astrag. 64. t. 52 = Oxytropis coerulea.
baicalia, Pall. Astrag. 93. t. 77. f. 1 = Oxytropis baicalia.
baionnensis, *Loisel. Fl. Gall.* ed. I. 474.—Gallia occ.
bakalinensis, *Bunge, in Mem. Sav Etr. Pétersb.* vii. (1851) 388 (*Pl. Lehm.* 92. t. 14).—Persia.
baksanis, *Bunge, Astrag.* i. 87; ii. 45.—Reg. Casp.; Ind. bor. occ.
Balansæ, *Boiss. Diagn.* Ser. II. vi. 59.—Cappadoc.
banaticus, Rochel, ex Nym. Consp. 189 = chlorocarpus.
Barbe-Aromis', Ehrenb. ex Bunge, Astrag. ii. 65 = Siberi.
Barbe-Jovis, *DC. Astrag.* 188.—As. Min.
Barbe-Jovis, Fenzl, ex Bunge, Astrag. ii. 149 = Fenzlii.
Barba-Mosis, *Ehrenb. ex Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 486.—Mont. Sinai.
barbatus, *Lam. Encyc.* i. 314.—Armenia.
barbatus, Sibth. & Sm. Fl. Græc. Prod. ii. 88 = Sibthorpianus.
Barrelieri, Dufour, in Ann. Sc. Gén. vii. (1826) 297 = incanus.
basianicus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 361.—Mesopotam.
bayonnensis, Auct. vide baionnensis.
Beckerianus, *Trautv. in Act. Hort. Petrop.* iv. (1876) 368.—Reg. Caucas.
Beckwithii, *Torr. & Gray, in Pacif. Rail. Rep.* iii. (1854) 120.—Am. bor. occ.
Bekes, Bertol. in Nov. Comm. Bonon. vi. (1844) 238 (Misc. Bot. ii. 19) = Russelii.
Belangerianus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) 484 = leiocladus.
Benthamianus, *Gill. in Hook. Bot. Misc.* iii. (1833) 187.—Chili.
Bergii, *Hieron. in Bol. Acad. Nac. Cordova*, iii. (1879) 243.—Reg. Argent.
Berteri, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 57. —Chili.
berythæus, *Boiss. & Blanche, Diagn.* Ser. II. ii. 33.—Syria.
berytius, *Bunge, Astrag.* i. 50; ii. 78.—Catasonia.
bethlehemiticus, *Boiss. Diagn.* Ser. I. ix. 55.—Palæstin.
bhotanensis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 126. —Reg. Himal.
bicarinatus, Stend. Nom. ed. II. i. 159 = Gueldenstædtia pauciflora.
bicolor, Desf. Cat. Hort. Par. ed. III. 478 = pictus.
bicolor, *Lam. Encyc.* i. 317.—Armenia.
bicolor, Stend. Nom. ed. II. i. 159 = nitidiformis.
bicristatus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 75.—Calif.
bicuspis, *Boiss. Fl. Orient.* 1094 (amend.).—As. Min.
bicuspis, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 406.—Tibet. occ.
bidentatus, *H. B. & K. Nov. Gen. et Sp.* vi. 403.—Am. calid.

ASTRAGALUS:—

bidentatus, Ehrenb. ex Bunge, Astrag. ii. 17 = hispidulus.
Bibersteinii, *Bunge, l. c.* i. 114; ii. 194.—Reg. Caucas.
Bienerti, *Bunge, l. c.* i. 80; ii. 135.—Persia.
bifidus, Turcz. ex Ledeb. Fl. Ross. i. 605 = multicaulis.
biflorus, Linn. Mant. 278 = Crotalaria biflora.
biflorus, Pall. Reise, iii. 306 = Gueldenstædtia pauciflora.
biflorus, *Vis. Fl. Lyb. Spec.* 44. t. 20.—Afr. bor.
Bigelovii, *A. Gray, Pl. Wright.* ii. 42.—Am. bor.; Mexic.
bingoellianus, Kotschy & Boiss. ex Boiss. Fl. Orient. ii. 857 = munchianus.
biovulatus, *Bunge, Astrag.* i. 11; ii. 11.—Persia.
bisarratus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = A. Biserrula.
Biserrula, *Bunge, Astrag.* i. 11; ii. 9.—Persia.
bisulcatus, *A. Gray, in Pacif. Rail. Rep.* xii. 49. t. 1. —Am. bor. occ.
bludshtanus, *Bunge, Astrag.* i. 76; ii. 129.—Belochist.
Bodenaus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 435.—Persia.
boeticus, *Linn. Sp. Pl.* 758.—Reg. Mediterr.
bohemicus, Mayer, in Abh. Boehm. Ges. iii. (1787) 314. t. 1 = virgatus.
Boissieri, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 394.—Hispan.
Boissieri, Bunge, Astrag. ii. 182, pro parte = creticus.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 337.—Calif.
bombycinus, *Boiss. Diagn.* Ser. I. ii. 50.—Aegypt.; Arab.; Persia.
Bonanni, J. & C. Presl, Delic. Prag. (1822) 42 = leucocephalus.
Bonannianus, Bunge, Astrag. ii. 242 = depressus.
bounacantinus, *Boiss. Fl. Orient.* ii. 870.—Lycia.
bonnoubilus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 99.—Persia.
Bourgæanus, *Coss. Notes Crit.* 160.—Hispan.
Bourgovii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 227.—Am. bor. occ.
brachybius, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 267 = hamosus.
brachybotrys, *Bunge, Del. Sem. Dorp.* (1839) 8; *et in Linnæa*, xiv. (1840) *Litt.* 116.—Sibir. alt.
brachycalyx, *Fisch. ex Boiss. Diagn.* Ser. I. ix. 90; *et in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 350.—Kurdist.
brachycarpus, *Bieb. Fl. Taur. Cauc.* ii. 201.—Reg. Caucas.
brachycentrus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 387.—Persia.
brachyceras, C. Koch, ex Bunge, Astrag. ii. 7 = tribuloides.
brachyceras, *Ledeb. Cat. Hort. Dorp.* (1823) 8.—Syria; Persia; Tauris.
brachycladus, *Boiss. Fl. Orient.* ii. 893.—Persia.
brachyclonus, *DC. Prod.* ii. 285.—Reg. Casp.
brachyodontus, *Boiss. Diagn.* Ser. I. ii. 46.—Persia.
brachyphyllus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 394 = condensatus.
brachypterus, Boiss. Diagn. Ser. I. ii. 69 = lagurus.
brachypterus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 397.—Oriens.
brachytropis, *Schrenk, Enum. Pl. Nov.* 79.—Soongaria.
brachystachys, *DC. Prod.* ii. 308.—Syria; Mesopotam.
brachytropis, *C. A. Mey. Verz. Pfl. Cauc.* 140.—Reg. Caucas.
Brachycarbidgei, *A. Gray, Bot. U. St. Expl. Exped.* i. 416.—Peruv.
bracteolatus, DC. Astrag. 208 = angustifolius.
brantsei, *Boiss. & Hoh, Diagn.* Ser. II. ii. 31.—Anatolia.
bractosus, C. A. Mey. ex Bunge, Astrag. ii. 185 = hyalolepis.
brachycarpus, Klotzsch, in Bot. Zeyeh. Waldem. Reise, 160 = melanostachys.
Brandegei, *Porter & Coult. Fl. Col.* 94.—Am. bor. occ.
brahuicus, *Bunge, Astrag.* i. 118, 206.—Belochist.
brasotensis, *Buckl. in Proc. Acad. Sc. Philad.* 1861 (1862) 452.—Texas.
brevicarinatus, DC. Astrag. 241. t. 49 = Gueldenstædtia pauciflora.
brevidorus, DC. l. c. 194. t. 31 = eriocephalus.
brevidorus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 386 = polyanthus.

ASTRAGALUS:—

brevidorus, Fenzl, ex Bunge, Astrag. ii. 149 = Fenzlii.
brevifolius, *Ledeb. Fl. Alt.* iii. 334.—Sibir. alt.
brevipes, *Bunge, Astrag.* i. 98; ii. 178.—Persia.
brevirostris, Pelt. Encyc. Suppl. i. 526 = Oxytropis brevirostris.
Breweri, *A. Gray, in Proc. Acad. Calif.* iii. (1868) 103. —Calif.
Brugueri, *Boiss. Diagn.* Ser. I. ix. 102.—Persia.
Bucorsa, *Willd. Enum. Hort. Berol. Suppl.* 51.—Ins. Maurit.
buchtormensis, *Pall. Astrag.* 176. t. 93.—Sibiria.
buchtormensis, Willd. ex Ledeb. Fl. Ross. i. 585 = Oxytropis Pallasii.
Buhsæanus, *Bunge, Astrag.* i. 120; ii. 207.—Persia.
Bulia, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 438 = cephalanthus.
bullarius, Pall. Reise, iii. 318 = Oxytropis mixotricha.
Bungeanus, *Boiss. Fl. Orient.* ii. 886.—Reg. Caucas.
Burseanus, *Benth. ex Harv. Thes. Cap.* i. 32. t. lxxxii. —Afr. austr.
Bustillonii, *Clos, in C. Gay, Fl. Chil.* ii. 117.—Chili.
byssantinus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) 379 = prusianus.
cabulicus, Boiss. Diagn. Ser. I. ix. 48 = Oxytropis cabulica.
cachinalensis, *Phil. Fl. Atac.* 15.—Chili.
cadmicus, *Boiss. Diagn.* Ser. I. ii. 44.—As. Min.
caelestis, *Boiss. Fl. Orient.* ii. 870.—Reg. Caucas.
caeruleus, Pall. Reise, iii. 298 = Oxytropis coerulea.
caeruleus, *Hort. Par. ex Bunge, Astrag.* i. 108; ii. 164.—Zeylan.
caespitosus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 230.—Am. bor. occ.
caespitosus, Pall. Astrag. 70. t. 57 = Oxytropis caespitosa.
calabricus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 402.—Calabria.
californicus, *Greene, in Bull. Calif. Acad.* n. 8 (1885) 157.—Calif.
callichrous, *Boiss. Diagn.* Ser. I. ix. 62.—Palæstin.
callilophus, *Bunge, Astrag.* i. 76; ii. 128.—Persia.
callistachys, *Bunge, in Bull. Soc. Nat. Mosc.* xxvi. (1853) 11. 463.—Persia.
calophyllus, *Boiss. & Heldr. Diagn.* Ser. II. ii. 26.—Anatolia.
calvescens, *Bunge, Astrag.* i. 42; ii. 65.—Persia.
calycinus, *Bieb. Fl. Taur. Cauc.* ii. 199.—Reg. Caucas.
calycosus, *Torr. ex S. Wats. in Bot. King, Exp.* 66.—Calif.
camelorum, *C. & W. Barbey, Herb. Levant*, 121. t. 8. —Aegypt.
campestris, *A. Gray, in Proc. Am. Acad.* vi. (1864) 229.—Am. bor. occ.
campestris, Linn. Sp. Pl. 761 = Oxytropis campestris.
camporum, *Benth. ex Bunge, Astrag.* ii. 3.—Afghanist.
camptocentrus, *Bunge, l. c.* i. 13; ii. 13.—Cappadocia.
campylanthus, *Boiss. Diagn.* Ser. I. vi. 42.—Persia.
campylorhychus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 32.—As. Min.; Persia; Soongaria.
campylosemus, *Boiss. Diagn.* Ser. I. ii. 32.—Cilicia.
campylotrichus, *Bunge, Astrag. Fedsch.* 207.—As. temp.
canadensis, *Linn. Sp. Pl.* 757.—Am. bor.
canalicularius, Willd. Enum. Hort. Berol. Suppl. 52 = scorpioides.
cancellatus, *Bunge, Astrag.* ii. 178.—Persia.
candicans, *Greene, in Bull. Calif. Acad.* n. 3 (1885) 156.—Calif.
candicinus, Pall. Astrag. 61. t. 49 = Oxytropis candicans.
candidissimus, *Ledeb. Fl. Alt.* iii. 309.—Sibir. alt.; Calif.
Candollianus, *Boiss. Diagn.* Ser. I. ii. 30.—Persia.
Candollianus, Royle, Illustr. Bot. Himal. 199 = Royleanus.
canescens, *Bunge, Astrag.* ii. 174.—Armenia.
canescens, DC. Astrag. 114. t. 18 = cinctoryloides.
canescens, Soland. ex Lowe, in Trans. Camb. Phil. Soc. iv. (1881) 34 = Solandri.
canus, *Ait. in Linnæa*, xxxii. (1863) 636.—Hungar.
cantabricus, Willd. Enum. Hort. Berol. Suppl. 53 = scorpioides.
cano-sior, *Bunge, Astrag.* i. 50; ii. 80.—Cappadoc.
canus, *Bunge, l. c.* ii. 195.—Persia.
canus, Willd. ex Ledeb. Fl. Ross. i. 606 = multicaulis.
capillipes, *Fisch. ex Bunge, Astrag.* i. 20; ii. 21.—Cabus.

ASTRAGALUS:—
capitatus, Lam. Fl. Fr. ii. 640 = A. Glanx.
capitatus, Linn. Sp. Pl. 755 = exaltatus.
Capito, *Boiss. & Hohen. Diagn.* Ser. I. ix. 40.—Persia.
cappadocicus, *Boiss. Diagn.* Ser. I. ii. 74.—As. Min.
caprinus, Guss. Fl. Sic. Syn. ii. 315 = Hystrii.
caprinus, *Linn. Sp. Pl.* ed. II. 1071.—Barbaria; Oriens.
caprinus, Pall. Reise, ii. 939 = dasyanthus.
Caraganae, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1836) 844.—Reg. Cauc.; Persia.
cardunchorum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 810.—Persia.
carinatus, *Boiss. Diagn.* Ser. I. ix. 56.—As. Min.
carinalis, Benth. ex Bunge, Astrag. i. 23; ii. 28 = himalayanus.
carnosus, Nutt. Gen. Am. ii. 100 = caryocarpus.
carnosus, Pursh, Fl. Am. Sept. ii. 740 = Sophora sericea.
carolinianus, Linn. Sp. Pl. 757 = canadensis.
caryocarpus, *Ker-Gawl. in Bot. Reg.* t. 176.—Am. bor.
caryocarpus, Torr. in Ann. Lyc. N. York, ii. (1828) 179 = plattensis.
caryolobus, *Bunge, Astrag.* i. 37.—Persia.
casapalensis, *Ball, in Journ. Linn. Soc.* xxii. (1885) 86.—And. Peruv.
Casei, *A. Gray, in Bot. Calif.* i. 154.—Calif.
caschmirensis, *Bunge, Astrag.* i. 30; ii. 84.—Kashmir.
caspicus, *Bieb. Fl. Taur. Cauc.* ii. 204.—Reg. Casp.; As. temp.
caspius, Hohen. ex Ledeb. Fl. Ross. i. 643 = praec.
castanaeformis, *S. Wats. in Proc. Am. Acad.* xx. (1885) 361.—Arizona.
castellanus, *Bunge, Astrag.* i. 9; ii. 5.—Hispan.
catacamptus, *Bunge, l. c.* ii. 191.—Persia.
catalinensis, Watt. in Journ. Acad. Sc. Phil. N. S. i. (1847) 159 = didymocarpus.
cataonicus, Boiss Fl. Orient. ii. 247 = affghanus.
catanonicus, *Bunge, Astrag.* i. 107; ii. 188.—Cataonia.
caucasicus, Bieb. Fl. Taur. Cauc. ii. 205 = Marschallianus.
caucasicus, *Pall. Astrag.* 2. t. 2.—Reg. Caucas.
caudatus, Pall. l. c. 62. t. 50 = Oxytropis songorica.
cauliculosus, *Boiss. Diagn.* Ser. II. ii. 25.—Armenia.
caucasomensis, *Phil. in Anal. Univ. Chil.* (1872) 691.—Chili.
Cedreti, *Boiss. Diagn.* Ser. I. ix. 39.—Mont. Libanus.
cephalanthus, *DC. Astrag.* 169. t. 88.—Persia.
cephalanthus, Hohen. ex Bunge, Astrag. ii. 117 = Hohenackeri.
cephalanthus, Kotschy, ex Bunge, l. c. ii. 120 = schirasicus.
cephalonicus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 424 = aristatus.
cephalonicus, *Presl, Bot. Bemerk.* 57.—Cephalonia.
Cephalotes, *Pall. Astrag.* 39. t. 34.—Reg. Caucas.
Cephalotes, [Soland. in] Russell, Aleppo, ed II. ii. 260 A. Andrachne.
cerasinus, *Baker, in Journ. Linn Soc.* xviii. (1878) 47. —Afghan.
cerasocrenus, *Bunge, Astrag.* i. 93; ii. 167.—Persia.
cerasoides, *Bieb. Fl. Taur. Cauc.* iii. 492.—Sibir. alt.
cerastoides, Bunge, Enum. Alt. 71 = senoceras.
Cerceini, Thomas, ex Steud. Nom. ed. II. i. 160 = lapponica.
chaboraicus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 487.—Mesopotam.
chastodes, *Bunge, in Mém. Sav. Etr. Acad. Pétersb.* vii. (1851) 272 (*Pl. Lehm.* 96).—Turkestan.
chastodes, Torr. ex A. Gray, in Proc. Am. Acad. vi. (1866) 194 = Spaldingii.
chastolobus, *Bunge, in Mém. Sav. Etr. Acad. Pétersb.* ii. (1835) 593.—Sibir. alt.
chastopodes, *Bunge, Astrag.* ii. 208.—Persia.
chalarasthus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 890.—Persia.
Chamaeleon, *A. Gray, in Ives, Rep. Bot.* 36.—Am. bor. occ.
Chardini, *Boiss. Diagn.* Ser. I. ii. 74.—Persia.
chartaceus, *Ledeb. Fl. Ross.* i. 694.—Persia.
chartostegius, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 810.—Orient; Persia.
Chesneyi, Bunge, Astrag. i. 118; ii. 202 = Wulfeni.
chilensis, *Lenn. f. Dresd.* t. 2.—China.
chlorochlorus, *Bunge, Astrag.* ii. 74.—Persia.
chlorosphaleus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 59.—As. Min.
chlorosphaleus, Kotschy, ex Bunge, Astrag. ii. 76 = dejectus.

ASTRAGALUS:—
chius, Boiss. & Orph. ex Boiss. Fl. Orient. ii. 870 = trojanus.
chlwensis, *Bunge, Astrag. Fedsch.* 247.—Khiva.
chloranthus, Pall. Astrag. 30. t. 25 = asper
chlorocarpus, Griseb. Spicil. Fl. Rumel. i. 50 = A. Onobrychis.
chlorocyaneus, *Boiss. & Reut. Diagn.* Ser. I. ix. 56.—Hispan.; Afr. bor. occ.
chlorodontus, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 39.—As. temp.
chlorophaeus, Bunge, Astrag. i. 50; ii. 79 = oxytropifolius.
chlorosphaerus, *Boiss. & Not, Diagn.* Ser. II. ii. 28.—Orient.
chlorostachys, Hook. f. & Thoms. ex Bunge, Astrag. ii. 98 = cariniis.
chlorostachys, *Lindl. in Trans. Hort. Soc.* vii. (1830) 249.—Reg. Himal.
chlorostegius, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 418.—Mesopotam.
choicus, Bunge, Astrag. i. 119; ii. 204 = latifolius.
chorassanicus, *Bunge, l. c.* i. 88; ii. 145.—Persia.
chorothianus, *Fisch. ex Bunge, l. c.* i. 52; ii. 86.—Reg. Caucas.
chorinensis, *Bunge, l. c.* i. 22; ii. 24.—Sibir.
Christianus, Aucher, ex Bunge, l. c. ii. 86 = diptherolobus.
Christianus, Heldr. ex Bunge, l. c. ii. 38 = plsidicus.
Christianus, *Linn. Sp. Pl.* 755.—As. Min.; Syria.
Christianus, Russell, ex Bunge, Astrag. ii. 36 = alepicus.
Christianus, Pinard, ex Bunge, l. c. ii. 88 = columnaris.
Christianus, Stev. in Pall. N. Nord. Beitr. vii. 293 = Sieversianus.
Christianus, Sibth. & Sm. Fl. Graec. Prod. ii. 85 = graecus.
Christianus, Vahl, Symb. Bot. i. 57 = tomentosus.
chromolepis, Boiss. & Hohen. Diagn. Ser. I. ix. 84 = aureus.
chrysanthus, *Boiss. & Hohen. l. c.* 78.—Persia.
chrysochlorus, Boiss. & Kotschy, in Boiss. l. c. Ser. II. v. 86 = pannosus.
chrysonotus, *Bunge, Astrag. Fedsch.* 310.—As. temp.
chrysophyllus, *Boss. Diagn.* Ser. I. ix. 38.—Syria.
chrysopterus, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 32.—China.
chrysostachys, *Boiss. Diagn.* Ser. I. ii. 69.—Persia.
chrysotrichus, *Boiss. l. c.* 75.—Persia.
chtonocephalus, *Boiss. & Bal. Diagn.* Ser. II. vi. 57. —Cappadoc.
Cicer, *Linn. Sp. Pl.* 757.—Europ. austr.; Reg. Caucas.; Sibiria.
cicerellus, *Boiss. & Bal. Diagn.* Ser. II. vi. 53.—Cappadoc.
cicerifolius, *Royle, ex Bunge, Astrag.* i. 44; ii. 70.—Reg. Himal.; Tibet. occ.
cicerisflius, Royle, ex Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 404 = scarlosus.
cicerinus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = A. Cicer.
ciliatus, *Boiss. Diagn.* Ser. I. ii. 59.—Cilicia.
ciliolatus, *Benth. ex Bunge, Astrag.* i. 27; ii. 33.—Reg. Himal.
cinereaceus, DC. ex Steud. Nom. ed. II. i. 160 = rupifragus, testiculatus.
cinereus, Boss. Enum. Pl. Volh. 29 = pallescens.
cinereus, DC. Astrag. 178 = reducnus.
cinereus, Turcz. Fl. Baic.-Dah. i. 223 = multicaulis.
cinereus, *Willd. Sp. Pl.* iii. 1215.—Armenia.
citrinus, *Bunge, Astrag.* i. 84; ii. 42.—Persia.
clandestinus, Roth, ex Spreng. Syst. iii. 300 = cymbiformis.
Clausii, C. A. Mey. in Goebel, Reise, ii. 265. t. 5 = vulpinus.
clavatus, *DC. Astrag.* 280. t. 44.—Armenia.
clavatus, Ledeb. Fl. Ross. i. 648 = brachycarpus.
Cleveland, *Greene, in Torrey Bot. Club.* ix. (1882) 121.—Am. bor.
Cloeianus, *Phil. in Anal. Univ. Chil.* (1872) 691.—Chili.
Clusii, *Boiss. Diagn.* Ser. I. ix. 101.—Hispan.
Clusii, *Pallini, ex Steud. Nom.* ed. I. 35; ed. II. i. 160.—Italia.
coarctatus, *Trautv. in Act. Hort. Petrop.* iv. (1876) 365.—Armenia.
cobrensis, *A. Gray, Pl. Wright.* ii. 48, in adnot.—Am. bor. occ.

ASTRAGALUS:—
cognatus, *C. A. Mey. in Schrenk, Enum. Pl. Nov.* 81. —As. centr.
collinus, *Boiss. Fl. Orient.* ii. 488.—As. Min.
collinus, *Dougl. ex Hook. Fl. Bor. Am.* 141, in syn.—Am. bor. occ.
columnaris, *Boiss. Diagn.* Ser. I. ix. 70.—As. Min.
coluteocarpus, *Boss. l. c.* 65.—Tibet. occ.; Afghanist.
coluteoides, *Willd. in Mém. Acad. Berl.* (1794-95) 27. —Syria.
commixtus, *Bunge, in Arb. Nat. Ver. Riga*, i. (1847) 246.—As. Min.; Persia; Turkest.
comosus, *Bunge, Astrag.* i. 56; ii. 93.—Persia.
compactus, Fisch ex Bunge, in Bull. Soc. Nat. Mosc. (1858) ii. 368 = zachlensis, strictifolius.
compactus, *Lam. Encyc.* i. 322.—Armenia.
compactus, C. A. Mey. Verz. Pfl. Cauc. 148 = Meyeri.
complanatus, *R. Br. ex Bunge, Astrag.* ii. 1.—Chin. bor.
complicatus, *Gill. in Hook. Bot. Misc.* iii. (1833) 187. —Chili.
compressus, *Ledeb. Fl. Alt.* iii. 304.—Sibir. alt.
concinnus, *Benth. ex Bunge, Astrag.* ii. 50.—Beluchist.
concretus, Benth. in Royle, Illustr. Bot. Himal. 199 = viciioides.
condensatus, *Ledeb. Fl. Ross.* i. 639.—Reg. Caucas.
condylocalyx, *Bertol. in Nov. Comm. Bonon.* vi. (1844) 281 (*Misc. Bot.* ii. 17).—Syria; Mesopotam.
confertiflorus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 368.—Am. bor.
confertus, Benth. ex Bunge, Astrag. ii. 27 = hippocrepidis.
confusus, *Bunge, l. c.* ii. 216.—Persia.
Congdoni, *S. Watt. in Proc. Am. Acad.* xx. (1885) 360.—Calif.
congestus, *Rakar, ex Aitch. in Journ. Linn. Soc.* xix. (1882) 158.—Ind. or.
conjunctus, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 371.—Am. bor.
consanguineus, *Dong. & C. A. Mey. Verz. Pfl. Saisang-Nor*, 22. t. 8.—China.
contortiplicatus, *Linn. Sp. Pl.* 758.—Europ. austr. occ.; Sibiria; Ind. or.
contortuplicatus, Sibth. & Sm. Fl. Graec. viii. 21. t. 729 = cruciatus.
Cooperi, *A. Gray, Man. Bot. U. St.* ed. II. 98.—Am. bor.
cordatus, *Bunge, Astrag.* i. 59; ii. 114.—Persia.
coriaceus, *Hemsl. Biol. Centr. Am. Bot.* i. 263.—Mexic.
corniculatus, *Bieb. Fl. Taur. Cauc.* iii. 492.—Besarab.; Podol.
corniculatus, Turcz. Cat. Fl. Baic. 256 = angarensis.
cornutus, *Bunge, Astrag.* i. 76; ii. 130.—Afghanist.
cornutus, Pall. Reise, i. 499 = vimineus.
Coronilla, *Bunge, Astrag.* ii. 10.—Persia.
corrugatus, *Bertol. Amon. Ital.* 88.—Aegypt.; Arab.; Pers.; Afghanist.
corydallinus, *Bunge, in Act. Hort. Petrop.* iii. (1875) 101.—Turkestan.
Cossonii, Bunge, Astrag. i. 118; ii. 192 = chlorocyaneus.
costatus, *Bunge, l. c.* i. 35; ii. 61.—Afghanist.
Coulteri, *Benth. Pl. Hartw.* 307.—Calif.
Cracca, *DC. Astrag.* 101. t. 9.—Peruv.
crassicarpus, Fras. Cat. ex DC. Prod. ii. 287 = caryocarpus.
crassicaulis, R. Grah. in Wall. Cat. n. 5932 = Caraguna crassicaula.
crassinervius, *Boss. Diagn.* Ser. II. v. 88.—Kurdistan.
crassipes, Fras. ex Steud. Nom. ed. II. i. 160 = caryocarpus.
crassispinus, *Bunge, Astrag.* i. 132; ii. 230.—Afghanist.
crassus, *Bunge, l. c.* i. 42; ii. 68.—Persia.
crenatus, Schult. Obs. 186 = corrugatus.
crenophilus, *Boiss. Diagn.* Ser. I. ix. 90.—Kurdistan.
cretaceus, *Boiss. l. c.* Ser. II. v. 84.—Syria.
cretaceus, Pall. in Nov. Act. Petrop. x. (1776) 56 = tauricus.
cretensis, Pall. Astrag. 5 = caspicus.
creticus, Boiss. Voy. Bot. Espagne, 181 = Boissieri.
creticus, DC. ex Boiss. Diagn. Ser. I. ix. 82 = cruentiflorus.
creticus, *Lam. Encyc.* i. 321.—Graecia.
creticus, Pinard, ex Bunge, Astrag. ii. 160 = ptilodes.
creticus, Willd. in Mém. Acad. Berl. (1794-95) 28 = echinoides.

ASTRAGALUS:—
creticus, Sibth. Fl. Grec. viii. 26. t. 736 = prusianus.
cristacanthus, Stev. ex Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1855) ii. 351 = strictatus.
crinitus, *Boiss. Diagn.* Ser. I. ii. 55.—Armenia.
cristatus, Gouan, Illustr. 50 = pentaglottis.
Crtschjahi, *Transv. in Act. Hort. Petrop.* viii. (1884) 278.—Rossia.
Crotalariae, A. Gray, in Proc. Am. Acad. vi. (1864) 216 = oocarpus.
cruciatus, *Link*, *Enum. Hort. Berol* ii. 256.—As. Min.; Syria.
cruciatus, Auct. ex Bunge, Astrag. ii. 8 = pseudostella.
cruciatus, Auct. quib. ex Boiss. Fl. Orient. ii. 226, in nota = polycalinus.
cruentiflorus, *Boiss. Diagn.* Ser. I. ix. 82.—Syria.
cruentus, Balb. Cat. Taur. App. i. 8 = corrugatus.
Cruikshankii, *Griseb. in Gortt. Abh.* xxiv. (1879) 103.—Reg. Argent.
cryptanthus, *Wedd. Chlor. Andi.* ii. 259.—Am. austr.
cryptocarpus, *DC. Astrag.* 137.—Armenia.
cucullaris, *Boiss. Diagn.* Ser. I. vi. 39.—Mesopotam.
cumanus, Beeb. ex Ledeb. Fl. Ross. i. 607 = lasioglottis.
cumanus, Willd. ex Steud. Nom. ed. II. i. 160, 159 = aduncus.
cuneifolius, *Bunge*, *Astrag.* i. 44; ii. 68.—Afghan.
curtipes, *A. Gray, in Proc. Calif. Acad.* iii. (1863) 103.—Am. bor. occ.
curvicarpus, *Boiss. Diagn.* Ser. I. ii. 67.—Persia.
curvirostris, *Boiss.* l. c. Ser. I. vi. 38.—Orient.
Cuscutae, *Bunge*, *Astrag.* i. 130; ii. 207.—Reg. Caucas.
Cusickii, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 270.—Am. bor.
cuspidatus, Bunge, Astrag. i. 117; ii. 198 = schizopterus.
cyaneus, *A. Gray, Pl. Fendl.* 34.—Am. bor. occ.
cygneus, Fisch. ex Bunge, Astrag. ii. 190 = subdijugus.
cylindraceus, *DC. Astrag.* 223.—Armenia.
cylindracrus, C. A. Mey. Verz. Pfl. Cauc. 143 = macrourus.
cylleneus, *Boiss. & Heldr. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 358.—Graecia.
cymbiformis, Willd. ex Brot. Phyt. 59 = cymbaecarpus.
cymbostegis, *Bunge*, *Astrag.* i. 85; ii. 14.—Catonia.
cyprius, *Boiss. Fl. Orient.* iii. 489.—As. Cyprus.
cyrenaicus, *Cosa. in Bull. Soc. Bot. Fr.* xiii. (1872) 30. —Cyren.
cyrtobasis, *Bunge*, *ex Boiss. Fl. Orient.* ii. 497. —Turkestan.
cyrtoides, A. Gray, in Proc. Am. Acad. vi. (1864) 201 = Gibbsii.
cystocalyx, *Ledeb. Fl. Ross.* i. 643.—Rossia.
cytisoides, *Bunge*, *Astrag.* i. 128; ii. 224.—Turkestan.
dactcus, *Hemfl. in Verh. Zool. Bot. Ges. Wien*, (1858) 98.—Europ.
dactylocarpus, *Boiss. Diagn.* Ser. I. ii. 60.—Mesopotam.
dasensis, *Boiss. l. c.* Ser. I. vi. 35.—Persia.
degenianus, Pall. Astrag. 88. t. 79 = Oxytropis prostrata.
daharicus, *DC. Prod.* ii. 285.—Dahuria; China.
daharicus, Patrin, ex DC. l. c. 277 = Oxytropis dissitiflora.
delgerianus, Willd. Sp. Pl. iii. 1201 = Oxytropis prostrata.
dalmaticus, Bunge, ex Nym. Consp. 190 = Soyeri.
Dalleianus, Bunge, Astrag. i. 44; ii. 69 = polycanthus.
damascenus, *Boiss. & Gaill. Diagn.* Ser. II. vi. 56.—Syria.
damascenus, DC. Prod. ii. 298 = colutecides.
danicus, *Retz. Obs.* iii. 41.—Europ.
Darumbium, *C. Gay, Fl. Chil.* ii. 112.—Chili.
dasyanthus, *Pall. Astrag.* 79. t. 65; *Retz*, iii. 749.—Hungar.; Sibir.; Reg. Cauc.
dasycephalus, Bens. ex Stev. in Bull. Soc. Nat. Mosc. (1852) 264 = roseus.
dasyglottis, Fisch. ex DC. Prod. ii. 282 = A. Hypoglottis.
dasyglottis, Pall. ex Steud. Nom. ed. II. i. 160 = pentaglottis.
dasyphyllus, Pall. Astrag. 91. t. 75 = Oxytropis lanata.
daucricus, C. Koch, in Linnaea, xv. (1841) 719 = Stevenianus.
daucricus, Auct. = daharicus.

ASTRAGALUS:—
dauricus, Ledeb. Ic. Fl. Ross. t. 315 = daharicus.
dauricus, Patrin, ex Ledeb. Fl. Ross. i. 597 = Oxytropis dissitiflora.
dealbatus, Ehrenb. ex Bunge, Astrag. ii. 164 = Rossianus.
decalbatus, Pall. Astrag. t. 29. f. 1 = albicaulis.
debilis, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 60.—Am. bor.
decumbiger, Bunge, Astrag. i. 43; ii. 67 = pallacanthus.
decipiens, Bunge, l. c. i. 107; ii. 188 = Sibthorpianus.
declinatus, Salisb. Prod. 337 = monspessulanus.
declinatus, *Willd. Sp. Pl.* iii. 1204.—Armenia.
decumbens, Boiss. Fl. Orient. ii. 620 = leucocyaneus.
decumbens, *A. Gray, in Proc. Am. Acad.* vi. (1864) 239.—Am. bor. occ.
decurrens, *Boiss. Diagn.* Ser. I. vi. 40.—Mesopotam.
defluxus, Pall. in Act. Acad. Petrop. ii. (1779) 268. t. 15 = Oxytropis deflexa.
delmacanthus, *Boiss Diagn.* Ser. I. ix. 76.—Syria.
delnensis, Bunge, Astrag. i. 55; ii. 91 = griseus.
demavendicus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 70.—Persia.
demirsus, Boiss. & Heldr. Diagn. Ser. I. ix. 50 = anacrus.
denudatum, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 325 = myriacanthus
denudatus, Stev. l. c. 328 = microcephalus.
depauperatus, *Ledeb. Fl. Alt.* i. 314.—Sibir. alt.
dependens, *Bunge*, *ex Maximow in Bull. Acad. Petersb.* xxvi. (1880) 471.—China occ.
depressus, Bertol. in Nov. Comm. Bonon. vi. (1844) 329 (Misc. ii. 332) = cruciatus.
depressus, *Linn. Cent. Pl.* ii. 29; *Amoen. Acad.* iv. 327.—Europ. austr.; Orient.
depressus, Pall. Reise, i. 239 = diffusus.
depressus, Sibth. & Sm. Fl. Graec. viii. 24. t. 733 = leucophaeus.
derbendicus, *Bunge*, *Astrag.* i. 40; ii. 55.—Persia.
desertorum, *Lehm. ex Spreng. Syst.* iii. 300.—Deserta Aden.
dianthus, *Bunge*, *Astrag. Fedsch.* 233.—As. centr.
diaphanus, Dougl. ex Hook. Fl. Bor. Am. i. 161 = lentiginosus.
dichopterus, Pall. Astrag. t. 39 = austriacus.
dictyocarpus, *Boiss. Diagn.* Ser. I. ii. 84.—Syria.
dictyoloxus, *C. A. Mey. ex Bunge, Astrag.* i. 60; ii. 98.—Persia.
dictyophysus, *Rend. ex Bunge*, l. c. i. 74; ii. 124.—Pontus.
dicyrtis, *Bunge*, *Astrag. Fedsch.* 235.—As. centr.
didymocarpus, *Hook. & Arn. Bot. Beech. Voy.* 354.—Calif.
didymophysus, *Bunge*, *Astrag.* i. 113; ii. 193.—Persia.
diffusus, Willd. Sp. Pl. iii. 1321 = dolichophyllus.
dilectus, *Phil. in Anal. Univ. Chil.* (1872) 690.—Chili.
dilutus, *Bunge*, *Del. Sem. Dorpat.* (1840) 7.—*Cf. Linnaea*, xv. (1841) *Litt.* 85.—Orient.
diopogon, *Bunge*, *Astrag.* i. 70; ii. 117.—Belochist.
diversicolor, Schott, ex Boiss. Fl. Orient. ii. 508 = Oxytropis dioritica.
dipelis, *Bunge*, *in Act. Hort. Petrop.* vii. (1880) 368. —Turkestan.
diphaces, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 349.—Am. bor. occ.
diphtherites, *Fenzl*, *in Flora*, xxvi. (1843) 396.—Mesopotam.
diphtherolobus, *Bunge*, *Astrag.* i. 81; ii. 36.—Syria.
diphyllus, Lam. Encyc. i. 330 = Oxytropis microphylla.
diphyses, *A. Gray, Pl. Fendl.* 34.—Am. bor. occ.
dipodurus, *Bunge*, *Astrag.* i. 88; ii. 156.—Syria.
dipsaceus, *Bunge*, *l. c.* i. 59; ii. 97.—Anatolia.
discolor, *Bunge*, *ex Maxim. in Bull. Acad. Petersb.* xxiv. (1878) 39.—Mongolia.
disperuus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 266.—Am. bor.
dissitiflorus, *Bunge*, *in Mém. Acad. Petersb. Sér. VI.* vii. (1850) 585.—As. temp.

ASTRAGALUS:—
distans, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 128 = macropterus.
distans, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 472.—Persia.
distans, *A. Gray, Bot. U. St. Expl. Exped.* i. 412.—Patagonia.
distortus, *Torr. & Gray, Fl. N. Am.* i. 338.—Am. bor.
divaricatus, *Bunge*, *Astrag.* i. 98; ii. 166.—Persia.
diversifolius, A. Gray, in Proc. Am. Acad. vi. (1864) 230 = junceus.
Doctii, *Phil. in Linnaea*, xxxiii. (1864–65) 48.—Am. austr.
dollichophyllus, *Pall. Astrag.* 84. n. 90. t. 68.—Reg. Caucas.
dolius, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 340. —Kurdistan.
Dombeyi, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 482.—Peruv.
Donianus, *DC. Prod.* ii. 283.—Nepal.
dorcocerus, *Bunge*, *Astrag.* i. 13; ii. 14.—Cilicia; Mesopot.
dorychnoides, Scop. Delic. Insub. ii. 104 = lusitanicus.
doryenioides, Dougl. ex G. Don, Gen. Syst. ii. 256 = succumbens.
Douglasii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 218.—Am. bor. occ.
drepanocarpus, Hochst. ex Bunge, Astrag. i. 14; ii. 16 = gyzensis.
drepanolobus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 75.—Am. bor. occ.
drepanophorus, *Griseb. in Gortt. Abh.* xxiv. (1879) 102.—Reg. Argent.
Drummondii, *Dougl. in Hook. Fl. Bor. Am.* i. 153.—Am. bor.
drupaceus, *Orphan. ex Boiss. Diagn.* Ser. II. ii. 32.—Graecia.
drusorum, *Boiss. Diagn.* Ser. I. ix. 78.—Syria.
dubius, DC. Astrag. 167. n. 115 = buchtormensis.
dyctiocarpus, Boiss. Diagn. Ser. I. ii. 84, sphalm. = dictyocarpus.
ebenoides, *Boiss. l. c.* Ser. I. ii. 70.—Persia.
echatanus, *Bunge*, *Astrag.* i. 71; ii. 118.—Persia.
Echegarayi, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 22.—Reg. Argent.
Echidna, *Bunge*, *Astrag.* i. 82; ii. 143.—Persia.
echinatus, Murr. Hort. Bot. Stirp. 222 = pentaglottis.
echinoides, L'Hérit. Stirp. Nov. 170 = angustifolius.
Echinops, *Ascher, ex Boiss. Diagn.* Ser. I. ii. 57.—Syria.
Echinus, Boiss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 280 = Boissieri.
Echinus, *DC. Astrag.* 197. t. 84.—Syria.
Echinus, C. A. Mey. Verz. Pfl. Cauc. 148 = caspius.
eclectus, Willd. Sp. Pl. iii. 1326 = angustifolius.
edulis, *Dur. ex Cosa. in Ann. Sc. Nat. Sér.* IV. i. (1854) 222.—Afr. bor.
effusus, *Bunge*, *Astrag.* i. 109; ii. 179.—Persia.
Ehrenbergii, *Bunge*, *l. c.* i. 61; ii. 102.—Syria.
elatus, *Boiss. & Bal. Diagn.* Ser. II. vi 57.—Cappadoc.
elbrusensis, *Boiss. l. c.* Ser. I. ix. 48.—Persia.
elegans, *Bunge*, *Astrag.* i. 55; ii. 89.—Persia.
Elliottii, D. Dietr. Syn. Pl. iv. 1080 = obcordatus.
ellipsoideus, Bong. & Mey. Suppl. Alt. 84, partim = kurtschumensis.
ellipsoideus, *Ledeb. Fl. Alt.* iii. 319.—Sibir. alt.
elmaleusis, *Bunge*, *Astrag.* i. 121; ii. 211.—Lycia.
elongatus, *Willd. Sp. Pl.* iii. 1214.—As. Min.
elymaiticus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 546.—Persia.
emarginatus, *DC. Astrag.* 150.—Orient.
emarginatus, *Ledeb. Ic. Fl. Syr. Dec.* i. 19.—Syria.
Emodi, *Steud. Nom.* ed. II. i. 161.—Reg. Himal.
enopius, *Trautv. in Act. Hort. Petrop.* iv. (1876) 264. —Reg. Caucas.
entomophyllus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 529.—Persia.
ephippium, *Pomel*, *Nouv. Mat. Fl. Atl.* 158.—Algeria.
epiglaucus, St. Leg. in Ann. Soc. Bot. Lyon, vii. (1880) 190 = epiglottis.
epiglottis, *Linn. Sp. Pl.* 759.—Reg. Mediterr.
epiglottoides, Willk. in Bot. Zeit. lii. (1845) 498 = A. Epiglottis.
epincos, *S. Wats. in Proc. Am. Acad.* x. (1875) 266. —Am. bor. occ.
eremophilus, *Boiss. Diagn.* Ser. I. ii. 54.—Aegypt.; Arab.; Belochist.

ASTRAGALUS:—
cremospartioides, *Regel, in Act. Hort. Petrop.* iii. (1878) 108.—Turkestan.
cremothamnus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 354.—Soongaria.
criacmus, *Boiss. Fl. Orient.* ii. 845.—Cappadoc.
crianthus, Fisch. ex Bobr. Fl. Orient. ii. 938 = crenophilus.
crianthus, Willd. Sp. Pl. iii. 1332 = gummifer.
crinecus, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 325.—Persia.
crinacus, Presl, Bot. Bemerk. 57 = angustifolius.
crioceltyr, *Bunge, Astrag.* i. 86; ii. 150.—Cilicia.
criocarpus, DC. Astrag. 287. t. 47 = mollis.
crioscarpus, *S. Wats. in Bot. King, Exp.* 71 & 440.—Am. bor. occ.
criocnemis, DC. Astrag. 188 = microcephalus.
criocephalus, Balansa, ex Bunge, Astrag. ii. 161 = acicularis.
criocephalus, Waldst. & Kit. Pl. Rar. Hung. i. 45. t. 46 = dasyanthus.
criocephalus, *Willd. Sp. Pl.* iii. 1338.—Armenia.
crioceras, *Fisch. & Mey. in Ledeb. Fl. Ross.* i. 626.—Ass. med.
criolobus, Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) 591 = depauperatus.
crioscorus, *Benth. ex Bunge, Astrag.* i. 33; ii. 40.—Afghan.
criophyllus, *Boiss. Diagn.* Ser. I. ix. 89.—As. Min.
criopodus, *Boiss. t. c.* Ser. I. ii. 48.—Persia.
crispoberus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 847.—Persia.
cristylus, *Boiss. & Haussk. ex Boiss. l. c.* 816.—Persia.
crocodes, Boiss. Diagn. Ser. I. ix. 58 = tribuloides.
crvoides, *Hook. & Arn. Bot. Beech. Voy.* 417.—Mexic.
crvoides, Turcz. in Bull. Soc. Nat. Mosc. (1838) 90 = miniatus.
erythrolepidius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = erythrolepis.
erythrolepis, *Boiss. Diagn.* Ser. I. ii. 65.—Persia.
erythrostemius, *Boiss. l. c.* 59.—Mesopotam.
eschkerensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 838.—Persia.
eubrychoides, *Boiss. Diagn.* Ser. I. ix. 41.—Cappadoc.
eucephalus, *Boiss l. c.* ii. 56.—Syria.
euoplus, *Trautv. in Act. Hort. Petrop.* iv. (1876) 364. —As. temp.
exaltatus, Bunge, Astrag. i. 25 = secundus.
eximius, *Bunge, Astrag. Fedsch.* 244.—As. centr.
expansus, Boiss. Diagn. Ser. I. ix. 48 = Kotschyanus.
exscapus, *Linn. Mant.* 275.—Europ.
exscapus, Auct. Transs.—Cf. Nym. Consp. 198 = pubiflorus.
fabaceus, DC. Prod. ii. 302 = striger.
fabaceus, *Bieb. Fl. Taur.-Cauc.* iii. 496.—Reg. Caucas.
fabaceus, Jan, ex Nym. Consp. 198 = Huetii.
falcatus, Desf. Fl. Atlant. ii. 188 = falciformis.
falcatus, *Lam. Encyc.* i. 310.—Reg. Caucas.; Sibir.
falciformis, *Desf. Emend. Atl. ex DC. Astrag.* 176.—Algeria.
falciniellus, *Boiss. Diagn.* Ser. I. ix. 68.—Aegypt.
Falconeri, *Bunge, Astrag.* i. 4; ii. 2.—Tibet. occ.
fallax, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 340 = mesolecus.
fallax, *S. Wats. in Proc. Am. Acad.* xx. (1885) 362.—Mexic.
farctus, *Bunge, in Mem. Sav. Etr. Pétersb.* vii. (1851) 376 (*Pl. Lehm.* 106).—Turkestan.
fasciculatus, *Bertol. in Nov. Comm. Bonon.* vi. (1844) 38.—Hab.?
fasciculifolius, *Boiss. Diagn.* Ser. I. ii. 78.—Persia.
fastidiosus, *Greene, in Bull. Calif. Acad.* n. 4 (1885) 186.—Calif.
fatmensis, *Hochst. ex Bunge, Astrag.* ii. 16.—Arabia; Beluchist.
Fendleri, *A. Gray, Pl. Wright.* ii. 44.—N. Mexic.
Fenalli, *Boiss. Diagn.* Ser. II. v. 89.—Cilicia.
ferrugineus, *C. Gay, Fl. Chil.* ii. 114.—Chili.
filaginens, Boiss. Diagn. Ser. I. ix. 69 = gossypinus.
filamentosus, *Bunge, Astrag.* i. 39; ii. 51.—Kurdist.
filicaulis, *Fisch. & Mey. Enum. Pl. Turc. et Pers.* 256. —Reg. Casp.; Turkest.; Afghan.
filifolius, A. Gray, in Pacif. Rail. Rep. iv. 80 = pictus.
filifolius, *Clos, in C. Gay, Fl. Chil.* ii. 111.—Chili.
filiformis, Pair. Encyc. Suppl. i. 538 = tanaitus.
filipes, *Torr. ex A. Gray, in Proc. Am. Acad.* vi. (1866) 226.—Am. bor. occ.
finitimus, *Bunge, Astrag.* i. 61; ii. 109.—Persia.

ASTRAGALUS:—
Fischeri, Buhse, in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 439 = macrocalyculus.
Fischerianus, Bunge, Astrag. i. 90; ii. 160 = pilodes.
flaccidus, *Bieb. Fl. Taur.* ii. 196. n. 1484.—Reg. Caucas.
flaccidus, Ledeb. Fl. Ross. i. 607 = onobrychioides.
flavescens, *Boiss. Diagn.* Ser. I. ii. 64.—As. Min.
flavus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 285.—Am. bor.
flexuosus, *Dougl. ex Hook. Fl. Bor. Am.* i. 141, in syn. —Am. bor.
flexus, *Fisch. in Bull. Phys.-Math. Acad. Pétersb.* iii. (1844) 307, in nota.—Persia; Beluchist.; Soongaria.
floccosus, *Boiss. Diagn.* Ser. I. ix. 61.—Persia.
floribundus, Pall. Astrag. 47. t. 37 = Oxytropis floribunda.
floridus, *Benth. ex Bunge, Astrag.* i. 24; ii. 28.—Reg. Himal.
floridus, Scop. Delic. Insub. ii. 108 = Christianus.
florulentus, *Boiss. & Haussk. Fl. Orient.* ii. 817.—Persia.
fodinarum, *Boiss. & Not, ex Bunge, Astrag.* i. 121; ii. 209.—Armenia.
foetidus, Vill. Prosp. 42 = Oxytropis foetida.
foliolosus, *Bunge, Astrag.* i. 74; ii. 125.—Anatolia.
follicularis, *Pall. Astrag.* 19. t. 14.—As. centr.
Fontanesii, *Coss. & Dur. ex Bunge, Astrag.* i. 75; ii. 126.—Algeria.
Forskahlei, *Boiss. Diagn.* Ser. I. ix. 101.—Aegypt.; Arab.; Syria.
fragiferus, *Bunge, Astrag.* i. 72; ii. 119.—Persia.
fragiformis, Willd. Sp. Pl. iii. 1261 = inflatus.
fragrans, *Willd. l. c.* 1284.—Orient.
Fraxinella, *Bunge, Astrag.* i. 39; ii. 52.—Persia.
frasinifolius, *DC. Astrag.* 142 n. 71.—Armenia.
Fremontii, Torr. in Pacif Rail. Rep. iv. 80 = malacus, lentiginosus.
Fresenii, *Decne. in Ann. Sc. Nat. Sér.* II. iii. (1835) 286.—Arab.
Frickii, *Bunge, Astrag.* i. 117; ii. 201.—Reg. Caucas.
frigidus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 219. —Am. bor.
Friesianus, Boiss. Diagn. Ser. I. ii. 78 = physocalyx.
frutescens, Hort. ex Kunth, Ind. Hort. Berol. (1845) 12 = Bongnaritia robiniodes.
fruticosus, Asso, Oryctogo. Arag. t. 2. f. 3 = Clusii.
fruticosus, Forsk. Fl. Aegypt. Arab. 139 = tomentosus.
fruticosus, Steud. Nom. ed. II. i. 161 = Hedysarum fruticosum.
fruticosus, *Pall. Astrag.* 21. n. 24. t. 19.—Sibir. or.
fruticosus, Ledeb. Fl. Alt. iii. 308 = vimineus.
gaeobotrys, *Boiss. & Bal. in Boiss., Diagn.* Ser. II. vi. 59.—Cappadoc.
Gaillardoti, *Boiss. Fl. Orient.* ii. 875.—Syria.
galactites, Ledeb. Fl. Ross. i. 656, ex parte = brevifolius.
galactites, *Pall. Astrag.* 85. t. 65.—Sibir.; Dahur.
galegiformis, Hohen. Enum. Pl. Elisabethpol. 250 = A. Glycyphyllos.
galegiformis, *Linn. Sp. Pl.* 756.—Reg. Caucas.
galegiformis, Pall. Reise, ii. 192 = uliginosus.
galegoides, Nutt. Gen. Am. ii. 100 = racemosus.
Garbancillo, *Cav. Ic.* i. 59, in textu.—Peruv.
Garbancillo, Pers. Syn. ii. 336 = prasc.
Garvansillo, Cav. Ic. i. t. 85 = Garbancillo.
Gebleri, *Fisch. ex Bong. Vers. Pfl. Saisang-Nor*, 34.—Soongaria.
geminans, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 830.—Persia.
geminiflorus, *Humb. & Bonpl. Pl. Aequin.* i. 126.—Ecuador.
genargenteus, Moris, Stirp. Sard. Elench. 11 = sirinicus.
genicularis, *Desf. Fl. Atlant.* ii. 186. t. 205.—Algeria.
genistoides, *Boiss. Diagn.* Ser. I. ix. 97.—Afghanist.
geobotrys, *Boiss. l. c.* Ser. II. vi. 62.—Phrygia.
geocyanus, *Boiss. Fl. Orient.* ii. 874.—Cappadoc.
georgicus, Boiss. Diagn. Ser. I. ii. 42 = Cephalotes.
Gerardi, Yanhe, in Termlxx. Füsteli. viii. (1884) 310. —Europ.
Gerardianus, R. Grah. in Wall. Cat. n. 5938 = Caragana Gerardiana.
gerensis, *Boiss. Diagn.* Ser. I. ix. 71.—Persia.
germesii, Boiss. & Buhse, ex Bunge, Astrag. ii. 64 = spinescens.
Germanii, *Phil. in Linnaea*, xxviii. (1856) 631.—Am. austr.

ASTRAGALUS:—
Geyeri, *A. Gray, in Proc. Am. Acad.* vi. (1864) 214. —Am. bor. occ.
ghilanicus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 329.—Persia.
gibberosus, Beriol. in Nov. Comm. Bonon. vi. (1844) 231 (Misc. Bot. ii. 17) = Chesneya rhytidosperma.
gibbosus, DC. Prod ii. 300 = Aminothamnus gibbosus.
Gibbsii, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 161. f. 50.—Calif.
giganteus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 370.—Am. bor.
gilensis, *Greene, in Torrey Bot. Club*, viii. (1881) 97. —N. Mexic.
Gillesii, *Phil. in Anal. Univ. Chil.* (1862) ii. 893.—Am. austr.
gilvus, *Boiss. Diagn.* Ser. I. ix. 71.—Caria.
glaber, DC. Astrag. 118. fragrans.
glaber, *Michx. Fl. Bor. Am* ii. 66.—Am. bor.
glaber, Lam. Encyc i. 325 = Oxytropis glabra.
glabrifolius, *Bunge, Astrag.* i. 93; ii. 167.—Afghanist.
glabriusculus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 204.—Am. bor. occ.
gladiatus, *Boiss. Diagn.* Ser. I. ii. 45.—As. Min.; Syria.
glanduliferus, *Debeaux, in Act. Linn. Soc. Bord.* xxxiii. (1879) 39.—China.
glareosus, *Dougl. ex Hook. Fl. Bor. Am.* i. 151.—Am. bor. occ.
glaucacanthos, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 481.—Persia.
glauciformis, *Pomel, Nouv. Mat. Fl. Atl.* 183.—Algeria.
glaucoides, S. G. Gmel. It i. 127. t. 22 = Coronilla varia.
glaucophyllus, *Bunge, Astrag.* i. 118; ii. 200.—Anatolia.
glaucus, Bieb. Fl. Taur. Cauc. ii. 203 = vesicarius.
Glaux, *Linn. Sp. Pl* 759.—Hispan.; Afr. bor. occ.
Glaux, Vall. Reise, ii. 464 = A. Hypoglottis.
glycyphillos, Neck. Delic. Gallo-Belg. ii. (1768) 312 = A. Glycyphyllos.
glycirrhiza, *Bunge, in Act. Hort. Petrop.* vii. (1880) 272.—As. centr.
globiflorus, *Boiss. Diagn.* Ser. I. ix. 81.—Persia.
globigerus, C. Koch, in Linnaea, xix. (1847) 64 = thraustus.
globosus, *Vahl, Symb. Bot* i 60.—Armenia.
glomeratus, *Ledeb. Fl. Alt.* iii. 327.—Sibir. alt.
glumaceus, *Boiss. Diagn.* Ser. I. ii. 60.—Persia.
glycyphyllodes, *DC. Prod.* ii. 292.—Reg. Caucas.
glycyphyllos, *Linn. Sp. Pl.* 758.—Europ.; Reg. Caucas.; Sibiria.
Gmelini, *Steud. Nomc. ed. II.* i. 161 = Laxmanni.
gomboforms, *Pomel, Nouv. Mat. Fl. Atl.* 187.—Algeria.
Gombo, *Coss. & Dur. ex Bunge, Astrag.* i. 42; ii 68. —Algeria.
Gomphobolium, *Benth. ex Bunge, l. c.* i. 89; ii. 50.—Beluchist.
gonistus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 330 = hypoglottis.
gossypinus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 376.—Kurdist.; Persia.
Govanianus, R. Grah. in Wall. Cat. n. 5925 = chlorostachys.
gracilentus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 223.—Am. bor. occ.
gracilipes, *Benth. ex Bunge, Astrag.* i. 15; ii. 17.—Tibet.
gracilis, *Nutt. Gen. Am.* ii. 100.—Am. bor.
gracilis, Stev. ex Ledeb. Fl. Ross. i. 609 = Steveníanus.
gracilis, Torr. in Ann. Lyc. N. York, ii. (1828) 179 = microlobus.
graecus, *Boiss. Diagn.* Ser. I. ii. 57.—Graecia.
Grahami, Steud. Nom. ed. II. i. 161 = vicioides.
Grahamianus, Fisch. in Bull. Soc. Nat. Mosc. (1853) ii. 406 = Jacquemontii.
Grahamianus, Royle, Illustr. Bot. Himal. 199. t. 36. f. 2 = polyacanthus
grammocalyx, *Boiss. & Hohen. Diagn.* Ser. I. ix. 52. —Persia.
granatensis, Lam. ex Steud. Nom. ed. I. 86; ed. II. i. 161 = A. Poterium.
granatensis, Lange, in Kjoeb. Vidensk. Meddel. (1865) 177 (Pugill. 273) = A. Glaux.
grandiflorus, *Bunge, Astrag. Fedsch.* 289.—As. centr.

ASTRAGALUS:—

grandiflorus, Linn. Sp. Pl. 761 = Hedysarum grandiflorum.

grandiflorus, Pall. Astrag. 57. t. 46 = Oxytropis grandiflora.

grandiflorus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 370.—Am. bor.

Grantii, *Bunge, Astrag.* i. 83; ii. 50.—Afghanist.

graveolens, *Buch.-Ham. in Wall. Cat.* n. 5929.—Reg. Himal.

Grayii, *Parry, ex S. Wats. in Am. Natural.* viii. (1874) 212.—Am. bor. occ.

Greenei, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 105. —N. Mexic.

Greggii, *S. Wats. l. c.* xvii. (1881–82) 348.—Am. bor. occ.

Griffithii, *Benth. ex Bunge, Astrag.* ii. 192.—Afghanist.

griseus, *Boiss. Diagn.* Ser. I. vi. 86.—Persia.

gryphus, *Cass. & Dur. ex Bunge, Astrag.* i. 12; ii. 18. —Algeria.

guatemalensis, *Hemsl. Biol. Centr. Am. Bot.* i. 264. Guatemala.

gudrunensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 462.—Assyria.

Guldenstaedtius, *Bunge, Astrag.* i. 119; ii. 203.— Persia.

gummifer, *Labill. in Journ. Phys.* (1790) 46.—Syria.

guttatus, [*Soland. in*] *Russell, Aleppo.* ed. II. ii. 260. —Aleppo.

gymnolobus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 426.—Armenia.

gymnopodus, *Boiss. Diagn.* Ser. I. ix. 66.—Afghanist.

gypsophilus, *Rouy, in Bull. Soc. Bot. Fr.* xxxi. (1884) 279, nomen.—Callia.

gyzensis, *Deisle, Fl. Egypt. Suppl.* 64.—Aft. bor.; Arab.; Persia; Soongaria.

Haarbachii, *Sprun. in Boiss. Diagn.* Ser. I. ii. 50.— Graecia.

haematocarpus, *Bunge, Astrag.* i. 118; ii. 201.—Reg. Caucas.

haleprensis, Fisch. ex Boiss. Fl. Orient. ii. 274 = aleppicus.

hallicasabus, *Lam. Encyc.* i. 220.—Armenia.

halicnacabus, Ledeb. Fl. Ross. i. 648 = mesites.

Halleri, All. Fl. Pedem. i. 348 = Oxytropis foetida.

Hallii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 224.— Am. bor. occ.

Halodendron, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 21.—As. temp.

hamatus, Lam. ex Willk. & Lange, Prod. Fl. Hisp. iii. 271 = A. Scalla.

hamosus, Balansa, ex Bunge, Astrag. ii. 17 = commixtus.

hamosus, Kotschy, ex Bunge, Astrag. ii. 14 = arnocerus.

hamosus, *Linn. Sp. Pl.* 758.—Reg. Mediterr.; Transcauc.; Ind. bor. occ.

hamosus, Pall. Reise, iii. 653 = reduncus, trimestris.

Hancockii, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 471.—China bor.

harpilobus, Boiss. Fl. Orient. ii. 284, in syn. = gyzensis.

Hartwegi, *Benth. Pl. Hartw.* 10. n. 58.—Am. bor. occ.

hasboyanus, *Boiss. Diagn.* Ser. I. ix. 77.—Syria.

Hausknechtii, *Bunge, Astrag.* i. 49; ii. 77.—Caucasia.

Haydenianus, *A. Gray, in Brandegee, Pl. S. W. Col.* 286.—Am. bor. occ.

hedysaroides, Linn. Sp. Pl. 756 = Hedysarum obscurum.

hedysaroides, Stevers, in Pall. N. Nord. Beitr. vii. 265 = A. Onobrychis.

hedysaroides, Willd. Sp. Pl. iii. 1264 = macrostachys.

heldreichii, Lerch. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = A. Cicer.

Heldreichii, *Boiss. Diagn.* Ser. I. ix. 48.—As. Min. bellenicus, *Boiss. Fl. Orient.* ii. 292.—Graecia.

Helmei, *Pomel, in Bonplandia*, viii. (1860) 56.— Mexic.

Helmii, *Fisch. ex DC. Prod.* ii. 301.—Mont. Ural.

helminthocarpus, Vill. Prosp. 48 = depressus.

hemiphaca, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 238.—Sibir.

Hemsleyi, *Aitch. & Baker, ex Aitch. in Journ. Linn. Soc.* xix. (1882) 158.—Afghanist.

Henderson, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 190.— Tibet. occ.

hercynus, *Bunge, Astrag.* i. 82; ii. 145.—Afghanist.

herrnianus, *Boiss. Diagn.* Ser. I. ix. 94.—Syria.

heterodontus, *Bunge, Astrag.* i. 86; ii. 56.—Persia.

ASTRAGALUS:—

Heydei, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 118.— Tibet, occ.

hians, Jacq. Coll. i. 87 = Oxytropis deflexa.

hilaris, *Bunge, Astrag.* i. 89; ii. 153.—Syria.

himalayanus, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 160.—Reg. Himal.

himalayensis, Hort. Par. ex Bunge, Astrag. ii. 93 = chlorostachys.

himalayensis, Jacquem. ex Heyrh. Nom. ii. 52 = seq.

himalensis, Jacquem. ex Baker, in Hook. Fl. Brit. Ind. ii. 128 = chlorostachys.

Hinkei, Sadl. ex Bunge, Astrag. ii. 217 = argenteus.

hippocrepidis, Benth. ex Bunge, l. c. i. 5; ii. 8.— Afghanist.

hippoglossus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = hypoglottis.

hirsutissimus, *DC. Astrag.* 119. t. 19.—Syria.

hirsutus, *Vahl, Symb. Bot.* i. 59.—As. Min.

hirtellus, Ledeb. Fl. Ross. i. 606 = vicisefolius.

hirticalyx, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* ii. 280.—Kurdistan.

hirtulus, Ledeb. Fl. Ross. i. 606 = vicisefolius.

hirtus, *Bunge, Astrag.* i. 84; ii. 57.—Persia.

hispanicus, *Coss. ex Bunge, l. c.* i. 135; ii. 231.— Hispan.

hispidulus, *DC. Astrag.* 132. t. 18.—Arabia.

hispidulus, Mlokz. ex Stend. Nom. ed. I. 86 = gummifer.

hispidus, Labill. Ic. Pl. Syr. Dec. i. 18 = barbatus.

Hoantii, *Franch. Pl. Davidian.* i. 66.—As. or.

Hohenackeri, *Boiss. Diagn.* Ser. I. ii. 70.—Reg. Caucas.

holargyreus, Bunge, in Mem. Sav. Etr. Pétersb. vii. (1851) 273 (Pl. Lehm. 97).—Schrenkianus.

hololeucus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 59.—Persia.

holopelius, *Bunge, Astrag.* i. 96; ii. 93.—Persia.

holopterus, *Turcz. ex Bunge, l. c.* i. 98; ii. 170.— Sibiria.

holosemius, *Bunge, l. c.* i. 55; ii. 89.—Persia.

Hookerianus, D. Dietr. Syn. Pl. iv. 1088 = erwoides.

Hookerianus, *A. Gray, Proc. Am. Acad.* vi. (1864) 215.—Am. bor. occ.

horsenaiatis, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 406 = Arbuscula.

Hornii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 398. —Am. bor. occ.

horridus, *Boiss. Diagn.* Ser. I. ii. 66.—Persia.

horridus, Willd. ex Bunge, Astrag. ii. 185 = arnacantha.

Hosackiae, *Greene, in Bull. Calif. Acad.* i. n. 8 (1885) 157.—Calif.

hosackioides, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 128.—Reg. Himal.

hostilis, *Boiss. Fl. Orient.* ii. 306.—Belachist.

Howellii, *A. Gray, in Proc. Am. Acad.* xv. (1880) 46. —Am. bor.

Huetii, *Bunge, Astrag.* i. 87; ii. 49.—Sicilia.

Humboldtii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 195.—Am. bor.

humifusus, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 119 = subumbellatus.

humifusus, Willd. Sp. Pl. iii. 1286 = vicisefolius.

humillis, *Bieb. Fl. Taur. Cauc.* ii. 208.—Reg. Caucas.

humilis, Meyer, ex Hook. Lond. Journ. Bot. vi. (1847) 211 = Shortianus.

humilis, Willd. ex Boiss. Diagn. Ser. I. ix. 83 = hamosus.

humillimus, *A. Gray, in Brandegee, Pl. S. W. Col.* 285.—Am. bor. occ.

humistratus, *A. Gray, Pl. Wright.* ii. 43.—Am. bor. occ.

hyalolepis, *Bunge, Astrag.* i. 105; ii. 185.—Reg. Caucas.

hybernus, Ehremb. ex Bunge, l. c. ii. 81 = hirsutissimus.

hybridus, S. G. Gmel. It. i. t. 18 = flaccidus.

hybridus, Willd. ex Ledeb. Fl. Ross. i. 628 = vesicarius.

hymenocalyx, *Boiss. Diagn.* Ser. I. ii. 55.—Persia.

hymenocarpus, Fisch. ex Bunge, Astrag. ii. 95 = chartaceus.

hymenochlaenus, Fisch. ex Bunge, l. c. i. 86; ii. 51 = angustifolius.

hymenocystis, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 449.—Persia.

hymenostegis, *Fisch. & Mey. l. c.* 448.—Persia.

ASTRAGALUS:—

hypoglaucus, *Ledeb. Fl. Ross. Alt.* v. 96; *Fl. Alt.* iii. 295.—Sibir. alt.

hypoglottis, Brot. Phytogr. t. 60 = pentaglottis.

hypoglottis, *Linn. Mant.* 274.—Reg. bor. temp.

hypoglottis, Desf. Fl. Atlant. ii. 188 = A. Glaux.

hypoglottis, DC. Astrag. 118 = danicus.

hypoglottoides, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 124.—Tibet. occ.

hypoleucus, *Schau. in Linnaea*, xx. (1847) 747.— Mexic.

hypoxylus, *S. Wats. in Proc. Am. Acad.* xviii. (1883–83) 193.—Am. bor. occ.

hypogeton, *Bunge, Astrag.* i. 89; ii. 140.—Persia.

hyrcanus, Kar. in Bull. Soc. Nat. Mosc. (1839) iii. = A. Ammodendron.

hyrcanus, *Pall. Astrag.* 35. t. 23.—Reg. Casp.

Hystrix, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 408.—Persia.

ibicinus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 293. —Persia.

ictericus, *Dingl. in Flora*, lxiv. (1881) 381.—Tiaracia.

ideus, *Bunge, Astrag.* i. 107; ii. 187.—Ins. Creta.

ilienais, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 20.—As. temp.

illyricus, [Bernh.] Sel. Sem. Hort. Erf. (1836).—Cf. Linnaea, xii. (1838) Lit. 74 = Wulfenii.

immersus, *Baker, ex Aitch. in Journ. Linn. Soc.* xviii. (1881) 45.—Afghanist.

impressus, Ehrenb. ex Bunge, Astrag. ii. 19 = peregrinus.

inarius, Pall. Astrag. 94. t. 77. f. 2 = Oxytropis inaria.

incanus, Chaub. ex Bunge, Astrag. ii. 202 = Chaubarii.

incanus, Gilib. Fl. Lithuan. ii. 119 = arenarius.

incanus, *Linn. Syst.* ed. X. 1175.—Gall. austr.

incanus, Pall. Astrag. iii. t. 84 = brachycarpus.

incanus, Sibth. & Sm. Fl. Graec. t. 782 = Sprunerl.

incanus, Siev. ex Ledeb. Fl. Ross. i. 592 = Oxytropis argentata.

incanus, Wulf. in Jacq. Coll. i. 230 = Wulfenii.

incertus, *Ledeb. Fl. Ross.* i. 647.—Reg. Caucas.; Armenia.

inconspicuus, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 126. —Reg. Himal.

incurvus, *Desf. Fl. Atlant.* ii. 182. t. 193.—Hispan.; Afr. bor. occ.

incurvus, Retzius. Fl. Germ. Exsurs. 519 = Wulfenii.

indersienis, *Claus, in Beitr. Pfl. Russ. Reich.* viii. 64. —Rossia.

indigoferoides, Horan. ex Bunge, Astrag. ii. 21 = capillipes.

inepsus, A. Gray, in Proc. Am. Acad. vi. (1864–65) 526 = lentiginosus.

infectus, *Boiss. Diagn.* Ser. I. ix. 95.—Afghanist.

infirmus, *Steud. Nom.* ed. II. i. 161.—Chili.

infistus, *DC. Astrag.* 90. n. 13.—Desert. Kirgh.

infistus, *Gill. ex Steud. Nom.* ed. II. i. 161.—Chili.

inflexus, *Dougl. ex G. Don, Gen. Syst.* iii. 256.—Am. bor. occ.

insularis, *Kellogg, in Bull. Calif. Acad.* n. i. (1884) 6. —Calif.

intarsensis, *Franch. in Ann. Sc. Nat. Sér. VI.* xv. (1883) 258.—Turkest.

intermedius, Boiss. Diagn. Ser. I. ii. 62 = commixtus.

intermedius, *Host, Fl. Austr.* ii. 361.—Europ.

intermedius, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 240.—Dahuria.

interruptus, Benth. ex Bunge, Astrag. ii. 10 = ammophilus.

iodanthus, *S. Wats. in Bot. King, Rep.* 70.—Am. bor. occ.

iodotropis, *Boiss. & Hohen. Diagn.* Ser. I. ix. 100.— Persia.

iranicus, Bunge, Astrag. i. 56; ii. 92 = mollis.

ischredensis, *Bunge, l. c.* i. 87; ii. 47.—Persia.

istenesis, Hort. ex Willd. Enum. Hort. Berol. 783 = falcatus, falciformis.

isopetalus, Boiss. Diagn. Ser. I. ix. 67 = lahtinus.

isphahanicus, *Boiss. l. c.* Ser. I. ii. 98.—Persia.

ispirensis, *Boiss. & Huet, l. c.* Ser. II. ii. 96.—Armenia.

Jacquemontii, Bunge, Astrag. i. 44; ii. 66 = polycarpus.

jacthicus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 286. —Persia.

Jenbertianus, *Reise, Fl. Orient.* ii. 24.—Phrygia.

Jenkinsianus, Bunge, Astrag. i. 90; ii. 160 = macrophthus.

ASTRAGALUS:—
japonus, *S. Wats. in Bot. King, Exp.* 73, 449.—Am. bor. occ.
jenisseensis, Less. ex Bunge, Astrag. ii. 188 = semibilocularis.
jendiamus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 61.—Persia.
Jessenii, *Bunge, Astrag.* i. 61; ii. 101.—Persia.
jodostachys, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 60.—Persia.
Johannis, *Boiss. Diagn.* Ser. I. vi. 41.—Persia.
jubatus, *Boiss. l. c.* ix. 104.—Persia; Afghan.
junceus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 380. —Am. bor. occ.
juncreus, Ledeb. in Spreng. Syst. iii. 397 = Stevenianus.
juvenalis, *Delile, Ind. Sem. Hort. Monsp.* (1836) 22.—Aegypt.
kadachoreusis, *Bunge, Astrag.* i. 108; ii. 182.—Reg. Caucas.
kahiricus, *DC. Prod.* ii. 292.—Aegypt.; Arab.; Persia; Afghan.
kaphernianus, Fisch. in Bull. Soc. Nat Mosc. xxvi. (1853) ii. 446 = persicus.
karabaghensis, *Bunge, Astrag.* i. 98; ii. 165.— Persia.
karakugensis, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 232.—Aa. Min.
karamanicus, Boiss. & Bal. Diagn. Ser. II. vi. 50 = orthodontus.
Karelini, Fisch. in Bull. Soc. Nat. Mosc. (1839) 152 = commixtus.
karputanus, *Boiss. Diagn.* Ser. II. ii. 26.—Armenia.
karsianus, *Bunge, Astrag.* ii. 115.—Armenia.
kaschmirensis, *Bunge*, l. c. ii. 246, Index = caschmirensis.
kellalensis, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 886.—Persia.
kentrodes, *Buhse, in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 412.—Persia.
kentrophyta, *A. Gray, in Proc. Acad. Philad.* (1862) 60.—Am. bor. occ.
kesseusis, *Bunge, Astrag.* i. 78; ii. 122.—Persia.
Kesslerі, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) i. 496.—Soongaria.
Keyserlingii, *Bunge, Astrag.* i. 83; ii. 139.—Persia.
khasianus, *Benth. ex Bunge, l. c.* i. 97; ii. 32.—Reg. Himal.
kirghisicus, Stschegl. in Bull. Soc. Nat. Mosc. xxvii. (1854) i. 161 = tribuloides.
kirvindicus, *Boiss. Diagn.* Ser. II. ii. 32.—Persia.
koburensis, *Bunge, Astrag.* i. 116; ii. 196.—Mongolia.
kohistanus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 354.—Turkest.
kohrudicus, *Bunge, Astrag.* i. 67; ii. 109.—Persia.
kokandensis, *Bunge, Astrag. Fedsch.* 216.—As. temp.
kongensis, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 190.—Him. bor. occ.
Korolkowii, *Bunge, Astrag. Fedsch.* 230.—As. temp.
koschakensis, *Boiss. Fl. Orient.* ii. 492.—Afghan.
Koschyanus, *Boiss. Diagn.* Ser. I. ii. 44.—Syria; Mesopotam.
Kotschyanus, Fisch. ex Bunge, Astrag. ii. 19 = tuberculosus.
Krauseanus, *Regel, in Act. Hort. Petrop.* iii. (1875) 104.—Turkestan.
kubistanus, *Bunge, Astrag.* i. 82; ii. 141.—Persia.
kuldshensis, *Bunge, in Act. Hort. Petrop.* vii. (1880) 374.—Turkestan.
kurramensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 46.—Afghan.
kurdicus, *Boiss. Diagn.* Ser. I. ix. 84.—Kurdistan; Syria.
kurumensis, *Bunge, in Act. Hort. Petrop.* vii. (1880) 370.—Afghan.
kurtschumensis, *Bunge, Astrag.* i. 129; ii. 226.— Soongaria.
labeadoricus, DC. Prod. ii. 287 = alpinus.
lacteus, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. ii. 31. —Graecia.
lacuiflorus, Ledeb. Fl. Alt. iii. 288 = testiculatus.
lactus, *Bunge, Astrag.* i. 84; ii. 41.—Afghan.
lagocephalus, *Fisch. & Mey. ex Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 197.—As. temp.
lagonyx, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 394.—Persia.
lagopodioides, Vahl, Symb. Bot. i. 64 = lagopoides.
lagopoides, Bunge, Astrag. i. 67; ii. 111 = unciformis.
lagopoides, *Lam. Encyc.* i. 329.—Armenia.
Lagowskii, *Trautv. in Bull. Phys.-Math. Acad. Pétersb.* xvi. (1858) 234.—Armenia.
lagouroides, *Pall. Reise*, iii. 750.—Sibir.
Lagurus, Eichw. Casp. Cauc. 681 = persicus.
Lagurus, Pall. Astrag. 16. t. 16 = lagouroides.
Lagurus, *Willd. in Mém. Acad. Berl.* (1794–95) 28.— Reg. Caucas.; Persia.
Lamarckii, *Boiss. Diagn.* Ser. I. ix. 89.—As. Min.
Lambertii, *Spreng. Syst.* iii. 908 = Oxytropis Lambertii.
lanatus, *Labill. Ic. Pl. Rar. Syr.* i. 21. t. 11.—Syria.
lanatus, Pall. Astrag. 108. t. 81 = Oxytropis Pallasii.
lanceolatus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 270.—Am. bor.
lanceolatus, *Bunge, Astrag.* i. 85; ii. 60.—Afghan.
lanceolatus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 246 = stipulatus.
lanigerus, *Desf. Fl. Atlant.* ii. 181. t. 202.—Afr. bor.
lanuginosus, Vis. Fl. Lib. Spec. 44 = alexandrinus.
lanuginosus, *Clos, in C. Gay, Fl. Chil.* ii. 121.—Chili.
lanuginosus, Desc. Fl. Pitt. Antill. Ind. viii. 53 = Sophora tomentosa.
lanuginosus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 409.—As. temp.
laricus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 81.— Persia.
lasianthus, C. A. Mey. in Bong. & Mey. Suppl. Fl. Alt. n. 92 = lasiopetalus.
lasiocaulos, Bunge, Astrag. i. 88; ii. 144 = pachyacanthus.
lasiocladus, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 185 = bicuspis.
lasiophyllus, *Dieb. Fl. Taur. Cauc.* iii. 486.—Reg. Caucas.
lasiopetalus, *Bunge, in Del. Sem. Dorp.* (1839).—*Cf. Linnaea* xiv. (1840) *Litt.* 117.—Soongaria.
lasiophyllus, *Ledeb. Fl. Ross.* ii. 227.—Orient.
lasiosemius, *Boiss. Diagn.* Ser. I. ix. 96.—Afghan.
lasiostylus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 367.—Turkestan.
latebracteatus, Boiss. in Tchihat. Asie Min. Bot. i. 54. = macrocholamys.
lateritius, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 886.—Kurdistan.
latifolius, DC. Astrag. 285 = Candolleanus.
latifolius, *Lam. Encyc.* i. 329.—Armen.; Persia.
laxiflorus, Boiss. Fl. Orient. ii. 413 = brachycodus.
laxiflorus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 390 = pallodothtis.
Laxmanni, Jacq. Hort. Vind. iii. 22. t. 37 = adsurgens.
Laxmanni, Nutt. Gen. Am. ii. 99 = striatus.
laxus, Boiss. & Hausskn. ex Boiss. Fl. Orient. ii. 269 = remotifolius.
Layneae, *Greene, in Bull. Calif. Acad.* i. n. 3. (1885) 156.—Calif.
Lehmannianus, *Bunge, in Arb. Nat. Ver. Riga*, i. (1847) 223.—Turkestan.
leiocalyx, Benth. ex Bunge, Astrag. i. 86; ii. 49 = lactus.
leiocarpus, Shuttlew. in Mag. Zool. & Bot. ii. (1838) 520 = crassipes.
leiocladus, *Boiss. Diagn.* Ser. I. ii. 62.—Persia.
leiolobus, *Bunge, Astrag.* i. 14; ii. 15.—Persia.
leiophyus, *Bunge, in Act. Hort. Petrop.* iii. (1875) 104.—Turkestan.
Lemmoni, *A. Gray, in Proc. Am. Acad.* viii. (1873) 698.—Am. bor. occ.
lentiformis, *A. Gray, in Bot. Calif.* i. 156.—Calif.
lentiginosus, *Dougl. ex Hook. Fl. Bor. Am.* i. 151.— Am. bor. occ.
leontinus, Wahlenb. Fl. Lapp. 191. t. 12. f. 4 = oroboides.
leontinus, *Wulf. in Jacq. Misc.* ii. 59.—Austr.; Helv.
leontopus, Bunge, Astrag. i. 89; ii. 157 = oleaefolius.
lepidanthus, *Boiss. Diagn.* Ser. I. ix. 86.—Syria.
leporinus, *Boiss. l. c.* Ser. II. ii. 30.—Armenia.
lepsensis, *Bunge, Astrag.* i. 26; ii. 29.—Soongaria.
leptacanthus, Benth. ex Bunge, l. c. i. 44; ii. 69 = leptocentrus.
leptacanthus, *Boiss. & Buhse, in Bull. Soc. Nat. Mosc.* (1853) ii. 423.—Persia.
leptaleus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 220. —Am. bor. occ.
leptocarpus, *Torr. & Gray, Fl. N. Am.* i. 334.— Texas.
leptocaulis, *Ledeb. Fl. Alt.* iii. 296.—Sibir. altaic.
leptocentrus, *Bunge, Astrag.* i. 44; ii. 69.—Reg. Himal.
leptochondron, Fisch. in Ball. Soc. Nat. Mosc. xxvi. (1853) ii. 345 = caspius.
leptodermus, *Bunge, Astrag. Fedsch.* 210.—As. centr.
leptophyllus, *Desf. Fl. Atlant.* ii. 188. t. 207.— Algeria.
leptophyllus, Nutt. in Journ. Acad. Philad. vii. (1834) 18 = stenophyllus.
leptophyllus, Pall. Reise, iii. 749 = Oxytropis leptophylla.
leptopus, *Benth. ex A. Gray, Pl. Wright.* ii. 43.— Mexic.
leptostachys, Pall. Astrag. 50. t. 40 = sulcatus.
leptostachys, Stev. in Willd. Sp. Pl. iii. 1265 = puberulus.
leptothalamus, Boiss. ex Bunge, Astrag. i. 135; ii. 232 = sec.
leptothamnus, *Boiss. Fl. Orient.* ii. 480.—Cilicia.
leptus, *Boiss. l. c.* 808.—Afghan.
lessertioides, *Benth. ex Bunge, Astrag.* i. 94; ii. 28.— Reg. Himal.
leucacanthus, *Boiss. Diagn.* Ser. I. ix. 98.—Aegypt.
leucanthus, Pall. Astrag. 59. t. 47 = Oxytropis leucantha.
leucocephalus, *R. Grah. in Wall. Cat.* n. 5923.—Reg. Himal.
leucocladus, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) 27.—As. temp.
leucocyaneus, *Griseb. Spicil. Fl. Rumel.* i. 50.—As. Min.
leucolachnus, *Boiss. Fl. Orient.* ii. 857.—Syria.
leucomallus, *Phil. in Linnaea*, xxx. (1859–60) 186.— Am. austr.
leuconeilas, *Bunge, Astrag.* i. 89; ii. 38.—Afghan.
leucophaeus, *Sm. in Trans. Linn. Soc.* i. (1791) 252. —Sicilia.
leucophyllus, Hook. Lond. Journ. Bot. vi. (1847) 211 = arrectus.
leucophyllus, *Torr. & Gray, Fl. N. Am.* i. 336.—Am. bor. occ.
leucophyllus, Willd. Sp. Pl. iii. 1281 = magnatilobus.
leucopsis, *Torr. Bot. Mex. Bound.* 56. t. 16.—Calif.
leucopilus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 896.—Kurdistan.
leucospermus, Bunge, in Arb. Nat. Ver. Riga, i. (1847) 250 = A. Alopecias.
libanoticus, Boiss. Diagn. Ser. II. v. 87 = A. Echinus.
Libanotis, Willd. Sp. Pl. iii. 1308 = lanatus.
lignosus, *DC. Astrag.* 175. t. 39.—Sibiria.
lilacinus, *Boiss. Diagn.* Ser. I. ix. 45.—Persia.
Lindheimeri, *Engelm. ex A. Gray, Pl. Wright.* i. 53. —Am. bor. occ.
linearifolius, Pers. Syn. ii. 336 = A. Onobrychis.
linearius, Willd. Sp. Pl. iii. 1302 = Oxytropis inaria.
linearius, *Lam. Encyc.* i. 314.—Armenia.
Listonіae, *Boiss. Diagn.* Ser. I. ix. 87.—Anatolia.
lithophilus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 344.—Soongar.
litoreus, *Phil. in Linnaea*, xxxiii. (1864–65) 47.— Chili.
litostachyus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 886.—Orient; Kurdistan.
lividus, Monb. ex Bunge, Astrag. ii. 19 = tuberculosus.
lobophorus, *Boiss. Diagn.* Ser. II. ii. 31.—Persia.
loochocarpus, *Torr. in Pacif. Rail. Rep.* iv. 80.—Am. bor. occ.
longebracteatus, Steud. Nom. ed. II. i. 109 = sanguinolentus.
longicaulis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 189.— Kashmir.
longicaulis, *Pomel, Nouv. Mat. Fl. Atl.* 185.— Algeria.
longicuspidatus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 190 = longicuspis.
longicuspis, *Bunge, Astrag.* i. 121; ii. 211.—Reg. Caucas.
longiflorus, Delile, Fl. Egypte, 256. t. 30 = kahiricus.
longiflorus, Gilib. Fl. Lithuan. ii. 111 = A. Onobrychis.
longiflorus, *Pall. Astrag.* 78. t. 60.—Reg. Casp.
longifolius, *Lam. Encyc.* i. 323.—Armenia.
longipes, Boiss. & Heldr. ex Boiss. Fl. Orient. ii. 480 = Apollinеus.

ASTRAGALUS:—
longipes, Steud. Nom. ed. II. i. 162 = rupifragus.
longipes, Ker. & Kir. in Bull. Soc. Nat. Mosc. (1841) 606 = macropterus.
longirostrus. Poir. Encyc. Suppl. i. 528 = Oxytropis longirostra.
longisetulus, *Bunge, Astrag.* i. 81 ; ii. 187.—Persia.
lotiflorus, *Hook. Fl. Bor. Am.* i. 152.—Am. bor. occ.
lotoides, Lam. Encyc. i. 816 = sinicus.
lunatus, Gillib. Fl. Lithuan. iii. 110 = glycyphyllos.
lunatus, *Pall. Astrag.* 44.—Persia.
lupulinus, *Pall. Reise*, iii. 750.—Siberia.
lusitanicus, *Lam. Encyc.* i. 812.—Reg. Mediterr. ; As. Min.
luteo-caeruleus, *Baker, in Journ. Linn. Soc.* xvii. (1878) 47.—Afghanist.
luxurians, *Bunge, Astrag.* i. 93 ; ii. 25.—Sibir.
Lyalli, *A. Gray, in Proc. Am. Acad.* vi. (1864) 195.—Am. bor. occ.
lycioides, *Boss. Diagn.* Ser. I. ii. 66.—Persia.
lycius, *Boiss. l. c.* ix. 45.—Lycia.
lydius, Boiss. l. c. ii. 48 = leucocyaneus.
lydius, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 393 = amphilocus.
macranthus, Willd. ex Ledeb. Fl. Ross. i. 650 = brachycarpus.
macrobotrys, *Bunge, Astrag.* i. 129 ; ii. 226.—Persia.
macrocarpus, *DC. Astrag.* 143. t. 93.—Syria.
macrocarpus, Pall. in Nov. Act. Petrop. x. (1776) 56 = subulatus.
macrocephalus, *Willd. Sp. Pl.* iii. 1360.—As. Min. ; Syria.
macrocras, C. A. Mey. ex Bong. Vers. Pfl. Saisang-Nor, 24 = macrolobus.
macrochlamys, *Boiss. ex Bunge, Astrag.* i. 49 ; ii. 77.—As. Min.
macrocladus, *Bunge, Astrag. Fedsch.* 293.—As. centr.
macrodon, *A. Gray, in Proc. Am. Acad.* vi. (1864) 216.—Am. bor. occ.
macrolacis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 63 = stenolepis.
macrolobus, *Bieb. Fl. Taur. Cauc.* iii. 498.—Sibir. altaic.
macrolobus, Bunge, Enum. Alt. 70 = macrocras.
macrolobus, Ledeb. Fl. Alt. iii. 307 = oriolobus, chaetolobus.
macronyx, *Bunge, Astrag. Fedsch.* 236.—As. centr.
macropelmatus, *Bunge, Astrag.* i. 86 ; ii. 43.—Persia.
macropetalus, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1844) 307.—As. temp.
macropodius, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 352 = aureus.
macropterus, *DC. Prod.* ii. 288.—Tibet. occ. ; Sibir.
macroptilus, *Boiss. ex Bunge, Astrag.* i. 90 ; ii. 160.—Phrygia.
macropus, Boiss. Diagn. Ser. II. vi. 51 = macrocsepus.
macropus, *Bunge, in Ark. Nat. Ver. Riga*, i. (1847) 238.—As. med.
macrorrhizus, *Cav. Ic.* ii. 28. t. 133.—Hispania.
macrorrhizus, D. Don, Prod. Fl. Nep. 245 = Donianus.
macrosepus, *Boiss. in Bunge, Astrag.* i. 49 ; ii. 77.—Cappadocia.
macrosemius, *Boiss. & Hohen. Diagn.* Ser. I. ix. 93.—Persia.
macrostachys, *DC. Astrag.* 141. t. 15.—Armenia.
macrotropis, *Bunge, Astrag.* i. 127 ; ii. 228.—Soongaria.
macrourus, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 546.—Reg. Caucas. ; Persia.
maculatus, *Bunge, Astrag.* i. 119 ; ii. 206.—Persia.
maculatus, Lam. Encyc. i. 816 = annularis.
Maddenianus, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 127.—Reg. Himal.
majellensis, Huet, ex Nym. Consp. 198 = australis.
malacensis, Salzm. ex Bunge, Astrag. ii. 12 = seamcus.
malacophyllus, *Benth. ex Bunge, l. c.* i. 86 ; ii. 61.—Reg. Himal.
malacophyllus, Hort. ex G. Don, in Loud. Hort. Brit. 307 = galegiformis.
malacotrichus, Fisch. & Kotschy, ex Bunge, Astrag. ii. 164 = Roussecanus.
malacus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 356.—Am. bor. occ.
marcoticus, *Delile, Fl. Egypt.* 257. t. 29. f. 3.—Aegypt.
maritimus, *Moris, Fl. Sard.* i. 238.—Sardinia.
maroniensis, *Dingl. in Flora*, xxiv. (1851) 382.—Thracia.

ASTRAGALUS:—
maroniticus, Boiss. & Balansa, Diagn. Ser. II. v. 58 = drusorum.
marschallianus, Lerch. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = dasyanthus.
Marschallianus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 326.—Orient ; Reg. Caucas.
marusiensis, Lerch. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = armarius.
masandranus, *Bunge, Astrag.* i. 114 ; ii. 193.—Persia.
massiliensis, *Jaub. & Spach*, i. 230.—Gallia.
Matthewsii, *S. Wats. in Proc. Am. Acad.* xviii. (1882-83) 192.—Am. bor.
mauritanicus, *Coss. in Bull. Soc. Bot. Fr.* iii. (1856) 678.—Algeria.
mauritanus, *Stev. in Bull. Soc. Nat. Mosc.* iv. (1832) 267.—Algeria.
Maximowiczii, *Trautv. in Act. Hort. Petrop.* ix. (1884) 379.—Rossia.
maximus, Lam. ex Steud. Nom. ed. II. i. 162 = A. Onobrychis.
maximus, *Willd. Sp. Pl.* iii. 1358.—Reg. Caucas. ; Armenia.
Maxwellii, Royle, Illustr. Bot. Himal. 199 = tibetanus ?
medicagineus, Boiss. Diagn. Ser. I. ix. 42 = Kotschyanus.
medius, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1843) 196.—Soongaria.
medullaris, Boiss. Diagn. Ser. I. ix. 66 = graveolens.
megacarpus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 215.—Am. bor. occ.
megalanthus, *DC. Astrag.* 131. t. 20.—Sibir.
megalocystis, *Bunge, l. c.* ii. 192.—Persia.
megalomerus, *Bunge, in Act. Hort. Petrop.* vii. (1880) 879.—As. centr.
megalotropis, *C. A. Mey. ex Bunge, Astrag.* i. 60 ; ii. 99.—Reg. Caucas.
melaleucus, *Bunge, Astrag.* ii. 100.—Persia.
melanocalyx, *Boiss. in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 59.—Persia.
melanocarpus, *Bunge, Astrag.* i. 51 ; ii. 81.—Orient.
melanocarpus, *Hook. Fl. Bor. Am.* i. 150.—Am. bor.
melanocarpus, Richards. App. Frankl. Journ. ed. II. 37 = missouriensis.
melanocephalus, *Boiss. Diagn.* Ser. I. ix. 50.—As. Min.
melanodon, *Boiss. l. c.* Ser. I. vi. 36.—Persia.
melanophrurius, *Boiss. Fl. Orient.* ii. 276.—Cappadoc.
melanostachys, *Benth. ex Bunge, Astrag.* i. 21 ; ii. 29.—Reg. Himal.
melanocalyx, Boiss. ex Bunge, l. c. ii. 188 = cataonicus.
melanotrichus, *Ledeb. Fl. Alt.* iii. 323.—Sibir. altaic.
melilotoides, *Pall. Voy.* t. 33 ; *Desc.* iii. 748.—Sibir. ; China.
melitus, *Boiss. Fl. Orient.* ii. 448.—Cappadocia.
melitenensis, *Boiss. Diagn.* Ser. I. ii. 45.—Cappadocia.
membranaceus, *Bunge, Astrag.* i. 25 ; ii. 30.—As. temp.
membranaceus, Moench, Meth. 167 = trimestris.
Menziesii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 217.—Calif.
Merklinii, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 70.—Persia.
meridionalis, *Bunge, Astrag.* i. 61 ; ii. 104.—Persia.
meschedensis, *Bunge, l. c.* i. 82 ; ii. 140.—Persia.
mestus, *Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 66.—Armenia.
mesogitanus, *Boiss. Diagn.* Ser. I. ii. 41.—As. Min.
mesoleios, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 63 = radkanensis.
mesoleios, *Boiss. & Hohen. Diagn.* Ser. I. ix. 91.—Persia.
mesotaenicus, *Boiss. l. c.* ii. 68.—Mesopotam.
mesopterus, *Griseb. Spicil. Fl. Rumel.* i. 49.—Macedon.
mexicanus, A. Gray, Fl. Lindh. 176 = plattensis.
mexicanus, *A. DC. Pl. Nov. Hort. Genev.* v. 17.—Am. bor.
Meyeri, *Boiss. Diagn.* Ser. I. ix. 86.—Armenia.
Michauxianus, *Boiss. l. c.* ii. 62.—Kurdistan.
micracme, *Boiss. & Noë, l. c.* Ser. II. v. 87.—Armenia.
micracistrus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 476.—Orient ; Assyria.
micranthellus, *Wedd. Chlor. And.* i. 262.—Peruv.
micranthus, *Desv. Journ. Bot.* iii. (1814) 76.—Hab. ?

ASTRAGALUS:—
micranthus, Nutt. in Journ. Acad. Philad. ii. (1821) 125 = Nuttallianus.
micranthus, Pall. ex Ledeb. Fl. Ross. i. 687 = strictellus.
microcarpus, *DC. Astrag.* 104.—Sibir.
microcephalus, *Willd. Sp. Pl.* iii. 1382.—As. Min.
microcystis, *Bunge, Astrag. Fedsch.* 306.—As. centr.
microcystis, *A. Gray, in Proc. Am. Acad.* vi. (1864) 220.—Am. bor. occ.
microdontus, *Baker, in Journ. Linn. Soc.* xviii. (1881) 46.—Afghan.
microlobus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 203.—Am. bor.
microphyllus, Bess. ex Steud. Nom. ed. II. i. 162 = A. Onobrychis.
microphyllus, Georgi, Beschr. Russ. Reich. Nachtr. 396 = adsurgens.
microphyllus, *Linn. Sp. Pl.* 757.—Sibir.
microphyllus, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 134 = cicerifolius.
microphyllus, Pall. Astrag. 92. t. 76 = Oxytropis microphylla.
microphyllus, Schrebl. & Martens, Fl. Wuertemb. 474 = A. Cicer.
microphysa, *Boiss. Diagn.* Ser. I. ii. 71.—Persia.
micropterus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 396.—Anatolia.
microchis, *Barbey, in Bull. Soc. Vaud. Sc. Nat.* xxi. (1886) 95.—Helvetia.
microthamnus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 393.—Assyria.
miniatus, *Bunge, Astrag.* i. 98 ; ii. 169.—Mongolia.
minimus, DC. Astrag. 64 = alpinus.
minimus, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 18.—Peruv.
minor, *Clos, in C. Gay, Fl. Chil.* ii. 107.—Chili.
minutissimus, *Wedd. Chlor. And.* ii. 257.—Peruv.
minutus, Boiss. Diagn. Ser. I. ix. 58 = tribuloides.
minutus, Buhse, ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xv. (1869) n. 7. 11 = blowitzius.
mirpourensis, Camb. in Jacquem. Voy. Bot. 39. t. 46 = Gouldenstaedtia mirpourensis.
miser, *Dougl. in Hook. Fl. Bor. Am.* i. 153.—Am. bor. occ.
missouriensis, *Nutt. Gen. Am.* ii. 99.—Am. bor.
Mitchellianus, *Beau. Diagn.* Ser. I. ix. 80.—Phrygia.
mixtus, Bieb. Cent. Pl. Rar. i. t. 26 = venicarnus.
modestus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 75.—Persia.
modestus, *Wedd. Chlor. And.* i. 263.—Am. austr.
Moellendorffi, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 472.—China bor.
mogollonicus, *Greene, in Torrey Bot. Club*, viii. (1881) 97.—Calif.
mohavensis, *S. Wats. in Proc. Am. Acad.* xx. (1885) 361.—Am. bor. occ.
moliscensis, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 367.—Am. bor. occ.
moldavicus, Steud. Nom. ed. I. 87 = A. Onobrychis.
mollis, *Bieb. Fl. Taur. Cauc.* iii. 495.—Reg. Caucas. Persia.
mollissimus, Presl, Bot. Bemerk. 57 = bethlemiticus.
mollissimus, *Torr. in Ann. Lyc. N. York*, ii. (1828) 178.—Am. bor. occ.
monadelphus, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 82.—China.
monanthus, *Boiss. Diagn.* Ser. I. ii. 78.—Persia.
mongholicus, *Bunge, Astrag.* i. 25 ; ii. 30.—Mongolia.
monophyllus, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 473.—Mongolia.
monspessulanus, Griseb. Spicil. Fl. Rumel. i. 60 = Wilfeni.
monspessulanus, *Linn. Sp. Pl.* 761.—Europ. austr. ; Reg. Cauc.
montanus, Bieb. Fl. Taur. Cauc. ii. 197 = Oxytropis cyanea.
montanus, Lapeyr. Hist. Abr. Pl. Pyr. 429 = Oxytropis pyrenaica.
montanus, Linn. Sp. Pl. 760 = Oxytropis montana.
montanus, Pall. Reise, iii. 199 = Oxytropis uralensis.
monstrosus, Brot. ex Steud. Nom. ed. I. 87 = macrorrhizus.
montanus, Jacq. Fl. Austr. 181 = alpinus.
monticola, *Phil. in Linnaea*, xxxiii. (1864-65) 47.—Chili.

ASTRAGALUS:—
Moorcroftianus, R. Grah. in Wall. Cat. n. 5933 = Sophora Moorcroftiana.
Mortoni, *Nutt. in Journ. Acad. Phil.* vii. (1834) 19. —Am. bor. occ.
mosulensis, *Bunge, Astrag.* i. 101; ii. 177.—Persia; Mesopotam.
muelleri, *Bunge, ex Boiss. Fl. Orient.* ii. 279.—Turkestan.
mucilaginosus, *Bunge, Astrag. Fedsch.* 223.—As centr.
mucronatus, *DC. Astrag.* 140. t. 25.—Armenia.
mucronifolius, *Boiss. Diagn.* Ser. I. ii 68.—Persia.
Muelleri, Stend. & Hochst. in Flora, x. (1827) 72 = A. Onobrychis.
mugodzharicus, Bunge, Astrag. i. 124; ii. 216 = mugosaricus.
mugosaricus, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 336.—Transural.
multiculis, *Ledeb. Fl. Alt.* iii. 295.—Sibir. altaic. et baical.
multicaulis, Nutt. ex Torr. & Gray, Fl. N. Am. i. 335 = pubentilissimus.
multiculis, Turcz. in Bull. Soc. Nat. Mosc. (1888) 90 = bifidus.
multiceps, *Wall. Cat.* n. 5237; *Bunge, Astrag.* ii. 69. —Reg. Himal.
multiflorus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 226.—Am. bor. occ.
multijugus, *DC. Prod.* ii. 301.—Mesopotam.
Munroi, *Benth. ex Bunge, Astrag.* i. 30; ii. 35.—Reg. Himal.; Kashgar.
muraricus, *Bunge, l. c.* i. 90; ii. 161.—Phrygia.
muricatus, Pall. Astrag. 89. t. 78 = Oxytropis muricata.
murinus, *Boiss. Diagn.* Ser. I. ii. 79.—Persia.
muschianus, *Kotschy & Boiss. ex Boiss. Fl. Orient.* ii. 357.—Armenia.
mutabilis, Bois. Fl. Orient. ii. 448 = paeclianthus.
mutabilis, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 125 = strictus.
myriacanthus, *Boiss. Diagn.* Ser. I. ix. 91.—Persia.
myriophyllus, *Bunge, Astrag. Fedsch.* 282.—As. centr. *myriophyllus*, Pall. Astrag. 87. t. 71 = Oxytropis myriophylla.
nanodes, *Bunge, in Act. Hort. Petrop.* vii. (1880) 377. —Turkest.
nanus, *DC. Astrag.* 114. t. 17.—Cilicia; Syria.
narbonensis, *Gouan, Illustr.* 49.—Gallia; Hispan.
narbonensis, Pall. Astrag. 13. t. 10 = posticus.
narbonensis, C. A. Mey. Verz. Pfl. Cauc. 148 = megalotropis.
nebrodensis, Presl, ex Steud. Nom. ed. II. i. 162 = Bolsiceri.
neglectus, *Fisch. ex Steud. l. c.*—Sibir.
nematodes, *Bunge, ex Boiss. Fl. Orient.* ii. 425.—Turkestan.
nervistipulus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 384.—Kurdistan.
neurocarpus, *Boiss. Diagn.* Ser. I. ii. 59.—Syria.
neurophyllus, *Franch. in Ann. Sc. Nat* Sér. VI. xv. (1883) 380.—Turkest.
nevadensis, *Boiss. Diagn.* Ser. I. ii. 68.—Hispan.
Newberryi, *A. Gray, in Proc. Am. Acad.* xii. (1877) 55.—Utah.
nicharensis, *Bunge, Astrag.* i. 30; ii. 50.—Beluchist.
nigrescens, A. Gray, in Am. Journ. Sc. Ser. H. xxxiii. (1862) 410 = multiflorus.
nigrescens, Nutt. in Journ. Acad. Philad. Ser. II. i. (1847) 153 = didymocarpus.
nigrescens, Pall. Astrag. 65. t. 58 = Oxytropis nigrescens.
nigrestriatus, C. Koch, in Linnaea, xv. (1841) 719 = ornithopodioides.
nitens, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 61 = viridis.
nitens, *Boiss. & Heldr. Diagn.* Ser. I. ix. 51.—As. Min.
nitens, *Host, Fl. Austr.* ii. 302.—Europ.
nitidus, Dougl. ex Hook. Fl. Bor. Am. i. 149 = adsurgens.
nivalis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 241.—Tibet. occ.; Soongaria.
nivosus, Coq. Monbr. ex Bunge, Astrag. ii. 189 = pelligeri.
Noeanensis, *Boiss. Diagn.* Ser. II. ii. 24.—Armenia.
nodiflorus, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) ii. 147.—Rossia.
Nortonii, *Nutt. in Journ. Acad. Phil.* vii. (1834) 19. —Am. bor.
norvegicus, *Grauer, Dec.* 13.—Norvegia.

ASTRAGALUS:—
nothoxys, *A. Gray, in Proc. Am. Acad.* vi. (1864) 232 —Am. bor. occ.
nubigenus, D. Don, Prod. Fl. Nep. 245 = Caragana crassicaulis.
nuciferus, *Bunge, Astrag. Fedsch.* 228.—Cappadoc.; Mesopotam.
nucleiferus, *Boiss. Diagn.* Ser. I. ii. 68.—Orient.
nudatus, Bunge, Astrag. i. 85; ii. 147 = kurdicus.
nudus, *Cloz, in C. Gay, Fl. Chil.* ii. 115 (1846).—Chili.
nudus, *S. Wats. in Bot. King, Exp.* 74, 448.—Am. bor. occ.
nummularioides, *Desf. Fl. Atlant. Emend. ex DC. Astrag.* 238.—Numidia.
nummularius, *DC. Astrag.* 221. t. 2. f. 2.—Ins. Creta.
nummularius, Desf. Fl. Atlant. ii. 189 = nummularioides.
nummularius, Lam. Encyc. i. 317 = pinetorum.
nummularius, C. A. Mey. ex Bunge, Astrag. ii. 57 = talyschensis.
nurensis, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 58.—Persia.
Nuttallianus, *DC. Prod.* ii. 289.—Am. bor. occ.
obcordatus, Boiss. Diagn. Ser. I. vi. 86 = sancardius
obcordatus, *Ell. Sketch*, ii. 227.—Am. bor.
oblongifolius, *Clos, in C. Gay, Fl. Chil.* ii. 109.—Chili.
obscurus, *S. Wats. in Bot. King, Exp.* 69, 437.—Am. bor. occ.
obtusifolius, *DC. Prod.* ii. 295.—Assyria.
obvallatus, Ehrenb. ex Bunge, Astrag. ii. 149 = crusiiformis.
oboviatus, Ehrenb. ex Bunge, l. c. ii. 168 = drusorum.
ochrostus, *Bunge, l. c.* i. 127; ii. 224.—Persia.
Ochrias, *Bunge, ex Maximow. in Bull. Acad. Pétersb.* xxiv. (1878) 88.—Mongolia.
ochrobius, *Bunge, Astrag.* i. 82; ii. 141.—Persia.
ochroleucus, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 87.—Algeria.
ochroleucus, Gilib. Fl. Lituan. ii. 111 = pilosus.
ocrochlorus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 83.—Persia.
odessanus, Bess. Enum. Pl. Volh. 80 = fruticosus.
odoratus, D. Dietr. Syn. Pl. iv. 1080 in syn. = obcordatus.
odoratus, *Lam. Encyc.* i. 311.—Reg. Caucas.; Persia; Soongaria.
oleifolius, *DC. Astrag.* 192.—As. Min.
olefolius, Trech. in Bull. Soc. Nat. Mosc. xxvi. (1853) 263 = sclerotropus.
Olgae, *Bunge, Astrag. Fedsch.* 228.—As. centr.
Olgini, Bunge, Astrag. i. 180; ii. 177 = mosulensis.
olygantus, Horan. ex Bunge, l. c. ii. 1 = complanatus.
oliganthus, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 237 = reticulatus.
oligophyllus, *Boiss. Diagn.* Ser. I. ii. 47.—Persia.
oligophyllus, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 197 = unijugus.
Olivieri, Klotzsch, ex Bunge, Astrag. i. 69; ii. 115 = mesopotamicus.
olopterus, DC. Prod. ii. 284 = austriacus.
olopterus, Turcz. ex Steud. Nom. ed. II. i. 162, 164 = vericolor.
olympicus, Bunge, Astrag. ii. 163 = balbutensis.
olympicus, Lam. ex Walp. Ann. iv. 514 = angustifolius.
olympicus, Pall. Astrag. 4. t. 4 = angustifolius.
oncocrotalus, *Bunge, Astrag.* ii. 11.—Persia.
onobrychioides, Bieb. Fl. Taur. Cauc. iii. 486 = A. Cephalotes.
onobrychioides, Bunge, Astrag. ii. 188 = Bungeanus.
Onobrychis, *Linn. Sp. Pl.* 760.—Europ. austr.; Orient.; Sibir.
Onobrychis, Poll. Hist. Pl. Palat. ii. 327 = hypoglottis.
onobrychis, *Gray, in Proc. Am. Acad.* vi. (1864) 218. —Am. bor. occ.
oocephalus, *Boiss. Diagn.* Ser. I. ii. 56.—Syria.
oophorus, *S. Wats. in Bot. King, Exp.* 73, 441.—Am. bor. occ.
ophiocarpus, *Benth. ex Boiss. Fl. Orient.* ii. 224.—Persia; Tibet. occ.
ophites, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 134 = cicerifolius.
orbicularis, Walp. Rep. i. 702 = seq.
orbicularis, *Ledeb. Fl. Ross. Alt.* i. 390; *Fl. Alt.* iii. 311.—Afghan.; Turkest.; Sibir. altaic.
Orbignyanus, *Wedd. Chlor. And.* ii. 260.—Bolivia.
Orcuttianus, *S. Wats. in Proc. Am. Acad.* xx. (1885) 361.—Calif.

ASTRAGALUS:—
oreades, *C. A. Mey. Verz. Pfl. Cauc.* 141.—Reg. Caucas.
oregonus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 335. —Am. bor. occ.
orientalis, Pers. ex DC. Prod. ii. 295 = macrocephalus.
ornithopodioides, *Lam. Encyc.* i. 312.—As. Min.; Persia.
orobioides, *Hornem. in Fl. Dan.* t. 1896.—Am. bor.
orthanthus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 195.—Mexic.
orthocarpus, *Boiss. Diagn.* Ser. I. ix. 68.—Afghan.
orthocarpus, Dougl. ex A. Gray, in Proc. Am. Acad. vi. (1864) 196 = canadensis.
orthodontus, Bunge, Astrag. i. 101; ii. 176 = karamaicus.
ortholobus, *Bunge, l. c.* i. 127; ii. 223.—Mont. Ural.
otiporensis, *Boiss. Fl. Orient.* ii. 271.—Afghan.
ourmiatensis, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 256.—Turkest.
ovalis, *Boiss. & Bal. Diagn.* Ser. II. vi. 52.—Cappadoc.
ovalicnsis, *Cloz, in C. Gay, Fl. Chil.* ii. 116.—Chili.
ovatus, *DC. Astrag.* 212. n. 107.—Armenia.
ovigerus, *Boiss. Diagn.* Ser. I. ii. 67.—Persia.
ovinus, *Boiss. l. c.* 76.—Armenia.
Owerini, *Bunge, Astrag.* i. 106; ii. 186.—Reg. Caucas.
oxyglottis, *Stev. in Bieb. Fl. Taur. Cauc.* ii. 192.—Reg. Caucas.; Persia; Turcoman.
oxyodon, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 125.—Tibet. occ.
oxypetalus, *Bunge, Astrag. Fedsch.* 278.—As. centr.
oxyphyllus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 375.—Syria.
oxyphyllus, Pall. Astrag. 90. t. 74 = Oxytropis oxyphylla.
oxyphysus, *A. Gray, in Proc. Calif. Acad.* iii. (1868) 108.—Am. bor. occ.
oxypterus, *Hemsl. Biol. Centr. Am. Bot.* i. 265.—Mexic.
oxysemus, *Bunge, Astrag.* i. 118; ii. 202.—Anatolia.
oxytropifolius, *Boiss. Diagn.* Ser. I. x. 37.—As. Min.
oxytropoides, Bunge, in Mém. Acad. Pétersb. Sér. VI. vii. (1850) 585; Sér. VII. xx. (1869) 185 = hemsus.
pachycaulinus, *Bunge, Astrag.* i. 93; ii. 148.—Persia.
pachycarpus, Torr. & Gray, Fl. N. Am. i. 332 = caryocarpus.
pachypus, *Greene, in Bull. Calif. Acad.* i. n. 3. (1885) 157.—Calif.
pachystachys, *Bunge, Astrag.* i. 80; ii. 134.—Persia.
paeclianthus, *Boiss. & Heldr. Diagn.* Ser. II. vi. 54.—Phrygia.
Pallasii, *Fisch. Hort. Gorensk.* (1812) 71; *et ex DC. Prod.* ii. 305 (1825).—Deserta Casp.
Pallasii, *Spreng. Fl. Hal. Mant.* (1807) 48.—Rossia.
pallescens, *Bieb. Fl. Taur. Cauc.* iii. 489.—Bessarab.; Podol.
pallidus, Boiss. Fl. Orient. ii. 372 = drusorum.
Palliseri, *A. Gray, in Proc. Am. Acad.* vi. (1864) 227. —Am. bor.
Palmeri, *A. Gray, l. c.* vii. (1868) 398.—Am. bor. occ.
pamphylicus, *Boiss. Fl. Orient.* ii. 289.—Cilicia.
Pancicii, *Hemfl. in Flora*, xxxvi. (1853) 621.—Europ.
pandustus, *Bunge, Astrag.* i. 89; ii. 95.—Anatolia.
pannonicus, Schult. Oestr. Fl. ed. II. ii. 385 = dasyanthus.
paucosus, *Fenzl, in Tchihat. Asie Min. Bot.* i. 2.—Cilicia.
paphlagonicus, Bunge, Astrag. i. 121; ii. 210 = atropurpureus.
paradoxus, *Bunge, l. c.* i. 64; ii. 108.—Persia.
parallopomenus, *Bunge, l. c.* i. 81; ii. 136.—Persia.
paralurgus, *Bunge, l. c.* i. 67; ii. 111.—Persia.
parapicticus, *Bunge, l. c.* i. 46; ii. 78.—Persia.
Parishii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 75. —Am. bor.
Parnassi, *Boiss. Diagn.* Ser. I. ix. 80.—Graecia; Orient.
Parnassi, Heldr. ex Nym. Consp. 192 = aristatus, cephalonicus.
parrowianus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 290.—Persia.
Parryi, C. Anders. Cat. Pl. Nev. 190 = malacus.
Parryi, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 410.—Am. bor. occ.
parviceps, *Bunge, Astrag.* i. 91; ii. 162.—As. Min.
parviflorus, Desf. ex Spreng. Syst. iii. 800, in syn. = racemosus.
parviflorus, Lam. Encyc. i. 310 = Oxytropis deflexa.

ASTRAGALUS:—

parviflorus, Nutt. ex A. Gray, in Proc. Am. Acad. vi. (1864) 202 = gracilis.

parvus, *Hemsl. Biol. Centr. Am. Bot.* i. 366.—Mexic.

Pastellianus, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 86 = vesicarius.

Patterson, *A. Gray, in Brandegee, Pl. S. W. Col.* 285. —Am. bor. occ.

pauciflorus, A. Gray, in Proc. Acad. Philad. (1863) 60 = leptaleus.

pauciflorus, Hook. in Lond. Journ. Bot. vi. (1849) 210 = alpinus.

pauciflorus, *Hook. Fl. Bor. Am.* i. 149.—Am. bor. occ.

pauciflorus, Pall. Astrag. 81. t. 66 = Gueldenstaedtia pauciflora.

paucijugus, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 196.—Turkestan.

paupar, *Bunge, Astrag. Fedsch.* 294.—As. centr.

pectinatus, *Dougl. ex Hook. Fl. Bor. Am.* i. 149.—Am. bor.

pectinatus, *Boiss. Diagn.* Ser. I. ii. 54.—Syria.

pedunculatus, *Royle, Illustr. Bot. Himal.* 199.—Tibet. occ.

Pehuenches, *Niederl. in Roca, Exped. Rio Negro*, ii. 306.—Patagonia.

pelliger, *Fenzl, Pugill. Pl. Nov. Syr.* 5.—Cilicia.

pellitus, *Bunge, Astrag.* i. 33; ii. 39.—Persia.

penduliflorus, Lam. Fl. Fr. ii. 608 = alpinus.

pendulius, DC. Astrag. 282 = elongatus.

penduliuz, Bunge, Astrag. ii. 211 = caelestis.

pennatus, *Bunge*, i. c. ii. 148.—Catacnia.

pentaglottis, *Linn. Mant.* 274.—Reg. Mediterr.

pentapetaloides, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 274 = Bexea.

peregrinus, Fisch. & Mey. ex Bunge, Astrag. ii. 84 = A. Caraganae.

peregrinus, *Vahl, Symb. Bot.* i. 57.—Aegypt.; Palaestin.

pestalozzae, *Boiss. & Heusch. ex Boiss. Fl. Orient.* ii. 861.—Persia.

peveranicus, C. A. Mey. ex Rupr. Fl. Bor. Ural. ii. 82. t. 11 = Halodi.

peyroullius, Bertol. in Nov. Comm. Bonon. vi. (1844) 230 = tribuloides.

phacophoidiformis, Boiss. Diagn. Ser. I. ix. 60 = ammophilus.

persicus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 8.—Reg. Casp.; Persia.

peruvianus, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 13.—Peruv.

Pestalozzae, Boiss. Diagn. Ser. I. ix. 41 = cadmicus.

petraeus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 333.—Mont. Alatav.

petropyimaia, *Bunge, Astrag.* i. 139; ii. 287.—Sibir.

phaeochrinus, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. 590 *; Verz. Suppl. Pl. Alt.* 90.—Sibir. altaic.

phlomoides, *Boiss. Diagn.* Ser. I. ii. 56.—Persia.

physaloides, *DC. Prod.* ii. 300.—Anatolia vel Syria.

physocalyx, *Fisch. in Bull. Sc. Acad. Pétersb.* ii. (1837) 74.—Thracia.

physocalyx, Kar. & Klr. in Bull. Soc. Nat. Mosc. (1841) 409 = cysticalyx.

physocarpus, Ledeb. Fl. Alt. iii. 325 = physodes.

physodes, *Linn. Sp. Pl.* 760.—Reg. Casp.

Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 415.—Peruv.

pictus, Boiss. & Gaill. Diagn. Ser. II. vi. 55 = condylicistus.

pictus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 214. —Am. bor. occ.

pictus, *Stend. Nom.* ed. II. i. 168.—Chili.

pimelioides, *Bunge, Astrag.* i. 88; ii. 52.—Persia.

pilosus, Linn. Sp. Pl. 756 = Oxytropis pilosa.

Pinardi, *Boiss. Diagn.* Ser. I. ix. 56.—As. Min.

pinetorum, *Boiss. l. c.* ii. 77.—As. Min.; Syria.

piptocephalus, *Boiss. & Heusch. ex Boiss. Fl. Orient.* ii. 876.—Persia.

pisidicus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 70.—Pisidia.

Pistaciae, Rost. ex DC. Prod. ii. 305 = elongatus.

plancus, *Cav. in C. Gay, Fl. Chil.* ii. 109.—Chili.

plataneus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 332. —Am. bor.

platyphyllus, Fisch. ex Bunge, Astrag. ii. 206 = cheiocus.

platyphyllus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 345.—Soongar.

platyrraphis, *Fisch. ex Bunge, Astrag.* i. 37; ii. 48.—Syria; Kurdist.; Mesopot.

platyosmeticus, *Bunge, Astrag.* i. 190; ii. 206.—Persia.

platytropis, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 526.—Am. bor. occ.

plebejus, *Boiss. Diagn.* Ser. I. vi. 27.—Persia.

plumosus, DC. ex Bunge, Astrag. ii. 148 = Lamarckii.

plumosus, Sieber, ex Bunge, l. c. ii. 59 = bethlemiticus.

plumosus, *Willd. Sp. Pl.* iii. 1388.—Galatia.

podocarpus, *C. A. Mey. Verz. Pfl. Casc.* 142.—Reg. Caucas.; Persia.

podocarpus, R. Grah. in Wall. Cat. n. 5980 = A. Emodi.

podocarpus, *Buch.-Ham. in Wall. l. c.*—Ind. or.

podolobus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 49.—Persia.

podosphaerus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 855.—Kurdistan.

poecilanthus, Bunge, Astrag. ii. 188 = pascilanthus.

polaris, *Benth. in Trans. Linn. Soc.* xxiii. (1861) 833. —Am. arct. occ.

polemitus, *Boiss. Fl. Orient.* ii. 806.—Belochist.

polemonieus, *Bunge, Astrag.* i. 125; ii. 283.—Anatolia.

pollotes, *Bunge, Astrag.* i. 116; ii. 197.—Sibir. altaic.

polonicus, *Royle, Illustr. Bot. Himal.* 199.—Reg. Himal.

polyacanthus, Wall. Cat. n. 5984 = Caragana polyacantha.

polycladius, *Boiss. Fl. Orient.* ii. 926, in nota.—Hispan.; Afr. bor.

polyanthus, *Bunge, Astrag.* i. 98; ii. 165.—Persia.

polycyanus, *Boiss. Diagn.* Ser. I. ix. 54.—Afghan.

polycephalus, Tenore, ex Bunge, Astrag. ii. 95 = ponticus.

polycnemus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 332.—Soongar.

Polygala, Pall. Astrag. n. 116. t. 83 = sanguinolentus.

polyphyllus, *Bunge, Astrag.* i. 35; ii. 59.—Reg. Caucas.

polyphyllus, Willd. Sp. Pl. iii. 1300 = Oxytropis microphylla.

pontieus, *Pall. Astrag.* 14. t. 11.—Reg. Caucas.; As. Min.

porphyracmus, *Boiss. Fl. Orient.* ii. 867.—Cappadoc.

porphyrobaphis, *Fisch. in Bull. Soc. Nat. Mosc.* (1858) ii. 423.—Persia.

porphyrolepis, Fisch. l. c. 878 = radkanensis.

porrectus, *S. Wats. in Bot. King, Exp.* 75, 411.—Am. bor. occ.

Potereus, DC. Astrag. 160 = Clusii.

Poterium, Pall. Astrag. i. t. 1-3 = arnacantha.

Poterium, *Vahl, Symb. Bot.* ii. 68.—Hispan.

praecox, Baumg. Enum. Stirp. Transs. ii. 362 = Wahlenii.

praetermissus, *Bull. in Journ. Bot.* xi. (1873) 306.—Marocco.

Presnii, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 222.—Am. bor. occ.

procerus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 851.—Persia.

procerus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 369. —Am. bor.

procumbens, *Hook & Arn. Bot. Beech. Voy.* 18.—Chili.

procumbens, Mill. Gard. Dict. ed. VIII. n. 18 = pentaglottis.

procumbens, *S. Wats. in Proc. Am. Acad.* xx. (1885) 361.—W. Mexic.

prolixus, *Sieber, ex Bunge, Astrag.* i. 9; ii. 6.—Afr. bor.; Ind. bor. occ.

prostratus, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. (1858) 90 = adsurgens.

prostratus, *Hook. & Arn. Bot. Beech. Voy.* 18.—Chili.

prostratus, Scop. Delic. Insub. ii. 106 = Oxytropis pauxilla.

pruinosus, Boiss. Diagn. Ser. I. ix. 38 = Boissieri.

Przewalskii, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxi. (1875) 85.—China.

pseudaustralis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. *Suppl.* 16.—Sibir. altaic.

pseudocalycinus, *Fisch. in Bull. Soc. Nat. Mosc.* (1858) ii. 847.—Anatolia.

pseudocicer, Opiz, in Flora, iv. (1821) 758 = A. Cicer.

pseudocorybchis, *Anders. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 228, nomen.—Hab.?

pseudopentaglottis, Boiss. & Noë, ex Boiss. Fl. Orient. ii. 886 = transcaucasicus.

pseudostella, *Delile, Fl. Egypt. Suppl.* 64.—Graecia; Aegypt.; Arab.

pseudotragacanthus, Pall. Astrag. 8. t. 8 = aureus.

pseudotragacanthus, Urv. in Mém. Linn. Soc. Par. i. (1822) 348 = thracicus.

psilacanthus, *Boiss. Diagn.* Ser. I. ix. 97.—Afghan.

psilocaulescens, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) 407 = ismaiensis.

psilocnus, *Bunge, Astrag.* i. 86; ii. 151.—Syria.

psilocentros, Fisch. in Bull. Soc. Nat. Mosc. (1858) ii. 405 = polycnanthus.

psilodontius, *Boiss. Diagn.* Ser. I. ix. 86.—Syria.

psilogliottis, Stev. in DC. Prod. ii. 288 = oxyglottis.

psilopterus, Bunge, Astrag. i. 45; ii. 71 = genistoides.

psilopus, Schrenk, in Bull. Sc. Acad. Pétersb. x. (1849) 364 = spartioides.

psilostylis, *Bunge, Astrag.* i. 37; ii. 152.—Persia.

psoraloides, Bunge, l. c. ii. 285 = asciculyx.

psoraloides, *Willd. Sp. Pl.* iii. 1300.—Cappadoc.

psoraloides, Lam. Encyc. i. 317; DC. Astrag. 145. t. 18 = canescens.

ptercocarpus, *S. Wats. in Bot. King, Exp.* 71.—Am. bor. occ.

pterocephalus, *Bunge, Astrag. Fedsch.* 262.—As. centr.

pterocystis, *DC. Prod.* ii. 294.—Austral.

ptilocephalus, *Baker, ex Aitch. in Journ. Linn. Soc.* xviii. (1881) 47.—Afghan.

ptilodes, *Boiss. Diagn.* Ser. I. ix. 81.—As. Min.

ptychophyllus, *Boiss. l. c.* vi. 43.—Persia.

pubentissimus, *Torr. & Gray, Fl. N. Am.* i. 693.—Am. bor. occ.

puberulus, Kar. & Klr. in Bull. Soc. Nat. Mosc. (1841) 405. n. 240 = macropterus.

puberulus, *Ledeb. Fl. Alt.* iii. 299.—Sibir. altaic.

pubescens, Schrank, Prim. Sel. S.=frigidus.

pubiflorus, *DC. Astrag.* 178.—Sibir.

pubigerus, Fisch. ex Trautv. in Act. Hort. Petrop. viii. (1883) 228 = pubiflorus.

pugioniferus, *Fisch. ex Bunge, Astrag.* i. 125; ii. 219. —Macedonia.

pugnaformis, L'Hérit. Stirp. Nov. 170 = longiflolus.

pugniformis, Willd. Sp. Pl. iii. 1324 = oleaefolius.

pulchellus, *Boiss. Diagn.* Ser. I. ii. 56.—Persia.

pulchellus, Sallab. Prod. 337 = danicus.

Pulsiferi, *A. Gray, in Proc. Am. Acad.* x. (1874) 69. —Am. bor. occ.

pulvinatus, *Bunge, Astrag.* i. 82; ii. 143.—Persia.

Pumilio, Pall. Astrag. 67. t. 55 = Oxytropis Pumilio.

punctatus, *Bunge, Astrag.* i. 130; ii. 206.—Persia.

pungens, Willd. Sp. Pl. iii. 1326 = sagustiflolus.

purpurascens, *Bunge, Astrag.* i. 35; ii. 49.—Afghan.

purpurascens, Jacq. f. ex Steud. Nom. ed. II. i. 168 = Onobrychis.

purpureus, Lam. Encyc. i. 314 = hypoglottis.

Purshii, *Dougl. ex G. Don, Gen. Syst.* ii. 271.—Am. bor. occ.

pusillus, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 19.—Peruv.

pycnocephalus, *Fisch. in Bull. Soc. Nat. Mosc.* (1858) ii. 871.—Kurdistan.

pycnocladus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 896.—Persia.

pycnolobus, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 594.—Sibir. altaic.

pycnophyllus, Kotschy, ex Bunge, Astrag. ii. 188 = mesolotus.

pycnophyllus, Stev. in Mém. Soc. Nat. Mosc. iv. (1814) 87 = microcephalus.

pycnorhizus, *Wall. Cat. n.* 5997.—Reg. Himal.

pycnostachys, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 527.—Am. bor. occ.

pygmaeus, DC. ex Bunge, Astrag. ii. 86 = nanus.

pygmaeus, Pall. Astrag. 66. t. 54 = Oxytropis nigrescens.

pyrenaicoides, Salzm. ex Bunge, Astrag. ii. 8 = epiglottis.

pyrrhotrichus, *Boiss. Diagn.* Ser. I. ix. 73.—Afghan.

quadrivalvatus, *Bunge, in Ank. Naturgf. Ver. Riga*, i. (1847) 245.—Persia; Afghan.; Turcoman.

quinquefoliolatus, *Bunge, Astrag.* i. 119; ii. 204.—Persia.

Quisqualis, *Bunge, Astrag. Fedsch.* 229.—As. centr.

racemosus, *Parsh, Fl. Am. Sept. ii. Suppl.* 740.—Am. bor.

racemulosus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 901.—Persia.

Raddeanus, Regel, Ind. Sem. Hort. Petrop. (1866) 40 = Sieckianus.

radiens, *Pomel, Nouv. Mat. Fl. Alt.* 135.—Algeria.

radiatus, Stend. Nom. ed. II. i. 169 = Sieberi.

ASTRAGALUS:—
radiatus, *Ehrenb. ex Bunge, Astrag.* i. 11; ii. 8.—Aegypt. Arab.
radiatus, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 367 = A. Stella.
radicans, *Hornem. Hort. Hafn.* ii. 708.—Hab.?
radicatus, Decne. in Ann. Sc. Nat. Sér. 11. iii. (1885) 268 = Sieberi.
radiciflorus, Steud. Nom. ed. II. i. 168 = testiculatus.
radicosus, *Bunge, Astrag.* i. 81; ii. 183.—Persia.
ramosissimus, *Soop. Delic. Insub.* ii. 105.—Sibir.
raphidontus, *Boiss. Diagn.* Ser. I. ix. 96.—Afghan.
rariflorus, *Ledeb. Fl. Alt.* iii. 313.—Soongaria.
Rattani, *A. Gray, in Proc. Am. Acad.* xix. (1883) 75. —Calif.
Rauwolfii, Vahl, Symb. Bot. i. 68 = Russelii.
Rebondi, *Cosz. ex Bunge, Astrag.* i. 87; ii. 49.—Afr. bor.
recognitus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 11. 452.—Persia.
recurvus, *Greene, in Bull. Calif. Acad.* i. n. 3. (1885) 155.—Calif.
reduncus, *Pall. Astrag.* 109. t. 92.—Reg. Caucas.
reflexistipulus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 44.—Japon.
reflexus, *Torr. & Gray, Fl. N. Am.* i. 334.—Am. bor. occ.
refractus, Boiss. & Buhse, in Boiss. Diagn. Ser. II. ix. 57 = Buhseanus.
refractus, *C. A. Mey. Verz. Pfl. Cauc.* 144.—Reg. Caucas.
Regelii, *Trautv. in Act. Hort. Petrop.* ii. (1873) 479. —Armenia.
Reinii, *Ball, in Journ. Linn. Soc.* xvi. (1878) 482.—Marocco.
remotiflorus, *Boiss. Diagn.* Ser. I. ii. 72.—Persia.
remotiflorus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 1095 (emend.).—Persia.
remotijugus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 74.—Persia.
reptans, *Willd. Hort. Berol.* ii. 88. t. 88.—Mexic.
resupinatus, Bieb. Fl. Taur. Cauc. iii. 493 = fragrans.
retamocarpus, *Boiss. & Hohen. ex Boiss. Diagn.* Ser. I. ix. 66.—Persia.
reticulatus, *Bieb. Fl. Taur. Cauc.* iii. 491.—Reg. Caucas.; Afghan.
retroflexus, Pall. Astrag. 88. t. 37 = Oxytropis deflexa.
retusus, DC. Astrag. 205. t. 36 = colutecoides.
retusus, Willd. in Mém. Acad. Berl. (1794–95) 24. t. 2 = augustifolius.
Reuterianus, *Boiss. Diagn.* Ser. I. ii. 71.—Persia.
reventus, *A. Gray, in Proc. Am. Acad.* xv. (1880) 40. —Am. bor.
rhacodes, *Bunge, Astrag. Fedsch.* 308.—As. temp.
rhizanthus, Boiss. Diagn. Ser. I. ii. 83 = carinesis.
rhizanthus, *Royle, Illustr. Bot. Himal.* 199.—Reg. Himal.
rhizocephalus, *Baker, ex Aitch. in Journ. Linn. Soc.* xviii. (1878) 46.—Ind. or.
rhodochrous, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 836.—Kurdistan.
rhodocarinus, *Boiss. & Hausskn. l. c.* 868.—Persia.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 414.—Peruv.
rigidulus, *Benth. ex Bunge, Astrag.* ii. 25.—Reg. Himal.
rigidus, Boiss. & Bal. Diagn. Ser. II. vi. 55 = melanocephalus.
Robbinsii, *A. Gray, Man. Bot. U. St.* ed. II. 98.—Am. bor.
Robbinsii, Matthew, Pl. Acad. 15 = alpinus.
robustus, *Bunge, Astrag.* i. 123; ii. 213.—Persia.
Rochelianus, Heuff. ex Grisb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 293 = A. Onobrychis.
roseus, *Ledeb. Fl. Alt.* iii. 320.—Sibir. altaic.
Rostani, Boiss. Diagn. Ser. I. ii. 45 = densifolius, chrysophyllus.
rostratus, *C. A. Mey. Verz. Pfl. Cauc.* 144.—Reg. Caucas.
rotundifolius, Presl, ex Steud. Nom. ed. II. i. 168 = glycyphyllus.
rotundifolius, Royle, Illustr. Bot. Himal. 199 = gruvcoicus.
rotundifolius, Willd. Sp. Pl. iii. 1317 = nummularioides.
Roussaeanus, *Boiss. Diagn.* Ser. I. x. 61.—Syria; Mesopotam.
Royleanus, Bunge, Astrag. i. 80; ii. 94 = Candollianus.
Royleanus, D. Don, Syn. Pl. iv. 1060 = strobiliferus, Royle.
rubriflorus, *Bunge, Astrag.* i. 67; ii. 100.—Persia.

ASTRAGALUS:—
rubrostriatus, *Bunge, Astrag.* i. 68; ii. 113.—Persia.
rudbaricus, Bunge, l. c. i. 128; ii. 215 = confusus.
rugosus, *Fisch. ex Bunge, l. c.* ii. 53.—Kurdistan.
rumelicus, Bunge, l. c. 81; ii. 137 = veluchensis.
runcinatus, *Boiss. Fl. Orient.* ii. 868.—Cappadoc.
rupicola, Bornm. ex Bunge, Astrag. ii. 19 = peregrinus.
rupifragus, *Pall. Astrag.* 86. t. 70.—Reg. Caucas.; Ural.
Ruprechti, *Bunge, Astrag.* i. 101; ii. 174.—Reg. Caucas.
Russelii, *Greene, in Bull. Calif. Acad.* i. n. 3 (1884) 8. —Calif.
rusciifolius, *Boiss. Diagn.* Ser. I. ii. 47.—Persia.
Russelii, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 260. t. 18.—Syria; Mesopotam.
rytidocarpus, *Ledeb. Fl. Alt.* iii. 316.—Sibir. altaic.
rytilobus, *Bunge, Astrag. Fedsch.* 309.—As. temp.
sabuletorum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 408 = arkalycensis.
sabuletorum, *Ledeb. Fl. Alt.* iii. 321.—Sibir. altaic.
sabulorum, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 368.—Am. bor.
saccatus, *Boiss. Diagn.* Ser. I. ix. 58.—Persia.
saccharensis, Ehrenb. ex Bunge, Astrag. ii. 16 = gyzensis.
saccocalyx, *Schrenk, Enum. Pl. Nov.* 88.—Soongaria.
sachalinensis, *Bunge, Astrag.* i. 28; ii. 36.—Ins. Sachalin.
saganlugensis, *Trautv. in Bull. Phys.-Math. Acad. Pétersb.* xvi. (1858) 252.—Armenia.
Saharae, *Pomel, Nouv. Mat. Fl. Atl.* 184.—Algeria.
Sahendi, *Buhse, in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 43.—Persia.
sakkarensis, Ehrenb. ex Boiss. Fl. Orient. ii. 934 = gyzensis.
salatavicus, *Bunge, Astrag.* i. 117; ii. 201.—Reg. Caucas.
salsuginosus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 341.—Soongaria.
salsula, *Soop. Delic. Insub.* ii. 104.—Sibir.
salsulatus, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 381 = masullicnsis.
samamensis, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 68.—Persia.
sanctus, *Boiss. Diagn.* Ser. I. ix. 47.—Syria.
sanguinolentus, *Bieb. Casp.* 190.—Reg. Caucas.
sarastagius, *Bunge, Astrag. Fedsch.* 309.—As. temp.
scaberrimus, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 91.—China bor.
scabrifolius, *Boiss. Diagn.* Ser. I. ii. 81.—Armenia.
scabrisetus, *Bong. Veg. Pfl. Saisang-Nor*, n. 89. t. 5.—Soongaria.
scaposus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 366.—Am. bor.
scaricosus, Benth. ex Bunge, Astrag. i. 44; ii. 71 = polyacanthus.
schahrudensis, *Bunge, l. c.* i. 61; ii. 108.—Persia.
Schanginianus, *Pall. Astrag.* 77. t. 78.—Sibiria.
Schelichowii, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 68.—Ochotsk.
Schimperi, *Boiss. Diagn.* Ser. I. ii. 83.—Arabia.
schisaicus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 457.—Persia.
schistaceus, *Bunge, Astrag.* i. 46; ii. 75.—Persia.
schistosus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 49.—Persia.
schizopterus, *Boiss. l. c.* ii. 83.—Cilicia; Syria.
Schmalhausenii, *Bunge, in Act. Hort. Petrop.* vii. (1880) 369.—Turkestan.
Schootianus, *Boiss. Diagn.* Ser. II. v. 59.—Cilicia; Cappadoc.
Schrenkianus, *Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 197.—Turkestan.
schumbicus, *Bunge, Astrag.* i. 76; ii. 139.—Persia.
sciadophorus, *Franch. Pl. Davidian.* i. 84.—As. or.
sciureus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 98.—Persia.
sclerocarpus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 228.—Am. bor. occ.
sclerocladus, *Bunge, Astrag.* i. 46; ii. 75.—Persia.
sclerocyphus, *Ledeb. Fl. Alt.* iii. 326; *Fl. Ross. Alt.* i. 266.—Sibir. altaic.
sclerocyclus, *Bunge, in Arb. Natucf. Ver. Riga*, i. (1847) 54.—Turkestan.
scoparioides, Ledeb. Fl. Ross. i. 615 = taurica.
scoparius, *Schrenk, Enum. Pl. Nov.* 82.—As. temp.
scopulorum, *Porter & Coult. Fl. Col.* 24.—Colorado.

ASTRAGALUS:—
scorpioides, *Pourr. ex Willd. Sp. Pl.* iii. 1280.—Hispan.
scorpius, *Boiss. Diagn.* Ser. I. ix. 108.—Persia.
scorpius, Buhse, ex Bunge, Astrag. ii. 127 = glaucacanthus.
Scorpiurus, Bunge, l. c. i. 5; ii. 8 = bicorpreipidis.
Scottianus, Stocks, ex Bunge, l. c. ii. 81 = medullaris.
scrobiculatus, Boiss. Fl. Orient. ii. 490 = gymnolobus.
secundus, *DC. Astrag.* 198. n. 55.—Sibir. austr.
secundus, Michx. Fl. Bor. Am. ii. 67 = alpinus.
segetalis, Boiss. Fl. Orient. ii. 277 = neurocarpus.
Segethi, *Phil. in Linnaea*, xxxiii. (1864–65) 47.—Chili.
sehabadensis, *Bunge, Astrag.* i. 68; ii. 113.—Persia.
Seidlitzii, *Bunge, l. c.* i. 40; ii. 56.—Persia.
Semenovii, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 23.—As. temp.
semiblocularis, *DC. Astrag.* 170.—Sibiria.
semibilocularis, *Fisch. ex Bunge, Astrag.* i. 103; ii. 183.—Sibir. altaic.
semipellitus, *Bunge, l. c.* i. 88; ii. 144.—Persia.
sempervirens, Lam. Encyc. i. 320 = aristatus.
sengnanesis, *Bunge, Astrag.* i. 82; ii. 142.—Persia.
serbicus, Panč ex Boiss. Fl. Orient. ii. 268 = glycyphylloides.
sericeus, DC. Astrag. 148 = globosus.
sericeus, Lam. Fl. Fr. ii. 645 = versicarius, A. Oxytropis, sordida.
sericeus, Willd. Enum. Hort. Berol. Suppl. 51 = hamosus.
sericoleucus, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 410.—Am. bor. occ.
sericophyllus, *Griseb. Spicil. Fl. Rumel.* i. 58.—Graecia.
sericostachyus, *Stocks, in Hook. Kew Journ.* iv. (1852) 146.—Beluchist.
serotinus, *A. Gray, in Pacif. Rail. Rep.* xii. 18, 51.—Am. bor. occ.
sesamicus, *Linn. Sp. Pl.* 759.—Reg. Mediterr.
sesamoeus, Pall. Astrag. 98 = oxyglottis.
sesamoides, *Boiss. Diagn.* Ser. I. ix. 59.—Afghan.
setbundites, Benth. in Royle, Illustr. Bot. Himal. 199 = trichocarpus.
sesquiflorus, *S. Wats. Proc. Am. Acad.* x. (1876) 346. —Utah.
setiferus, DC. Prod. ii. 296 = Cornulaca tragacanthoides.
setosus, Pall. Astrag. 55. t. 44 = Oxytropis setosa.
setulosus, *Boiss. & Bal. Diagn.* II. vi. 50.—Cappadoc.
Sewerzowii, Boiss. Fl. Orient. ii. 289 = seq.
Sewerzowii, *Bunge, Astrag.* i. 86; ii. 54.—Turkest.
seyanensis, *Bunge, l. c.* i. 87; ii. 154.—Syria.
Shortianus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 331. —Am. bor. occ.
Shortianus, S. Wats. Pl. Wheeler, 7 = cyaneus.
sibiricus, *Soop. Delic. Insub.* ii. 104.—Sibiria.
Sibthorpianus, *Boiss. Diagn.* Ser. I. ii. 84.—Bithynia.
siculus, *Bison. Giorn. Palerm.* n. 4.—Sicilia.
siculus, Guss. ex Nym. Consp. 199 = calabricus.
siculus, Schimp. ex Nym. l. c. = cephalonicus.
Sieberi, *DC. Prod.* ii. 295.—Aegypt.; Arab.
Sieversianus, *Pall. Astrag.* 15. t. 12.—Persia; Soongaria.
Sieversianus, Steph. ex Ledeb. Fl. Ross. i. 592 = Oxytropis argentea.
Sieversii, Coq. Monb. ex Bunge, Astrag. ii. 26 = diphtherolobus.
sigmoideus, *Bunge, l. c.* i. 128; ii. 214.—Anatolia.
sikkimensis, *Benth. ex Bunge, l. c.* i. 23; ii. 28.—Reg. Himal.
siliquosus, *Boiss. Diagn.* Ser. I. ii. 51.—Persia.
siliquosus, Boiss. l. c. vi. 40 = lobophorus.
silvicolus, Fritz. ex Ind. 132 = Oxytropis sylvatica.
simplicifolius, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 231.—Am. bor. occ.
sinaicus, Boiss. Diagn. Ser. I. ix. 87 = pseudostella.
singaricus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 868.—Mesopotam.
sinicus, *Linn. Mant.* i. 103.—China.
sinicus, Thunb. Fl. Jap. 290 = lotoides.
sirinicus, *Tenore, 'Prod. Fl. Neap. App.* v. 25 (1826)'; *Fl. Neap.* v. 182. t. 171.—Italia.
skyrodytes, *Bunge, Astrag.* i. 33; ii. 40.—Turkestan.
sobieus, *Bunge, l. c.* i. 126; ii. 220.—Persia.
sivasicus, *Bunge, l. c.* i. 93; ii. 166.—Armenia.
skythropos, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 81.—China.
soficus, *Bunge, Astrag.* i. 81; ii. 136.—Persia.
sogdianus, *Bunge, Astrag. Fedsch.* 268.—As. temp.

ASTRAGALUS :—
Solandri, *Lowe, in Hook. Kew Journ.* viii. (1856) 294. —Marocco; Madera.
somcheticus, C. Koch, in Linnæa, xv. (1841) 730 = sanguinolentus.
songaricus, Pall. Astrag. 68. t. 51 = Oxytropis songarica.
Sonorae, A. Gray, Pl. Wright. ii. 44 = arizonica.
sordidus, Willd. Sp. Pl. iii. 1313 = Oxytropis campestris.
Soyeri, *Buching. ex Bunge, Astrag.* i. 118; ii. 199.—Dalmat.
Spachianus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 67.—Persia.
Spaldingii, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 544.—Am. bor. occ.
sparsiflorus, *A. Gray, in Proc. Acad. Philad.* (1863) 60.—Am. bor. occ.
sparsus, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 367.—Arab.
spartioides, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 330.—Soongar.
speciosus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 67.—Persia.
speirocarpus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 225.—Am. bor. occ.
sphaeranthus, *Bous. Diagn.* Ser. I. vi. 37.—Persia.
sphaerocalyx, *Ledeb. Fl. Ross.* i. 647.—Mont. Ararat.
sphaerocarpus, *Clos, in C. Gay, Fl. Chil.* ii. 117.—Chili.
sphaerocarpus, *Decf. ex Steud. Nom.* ed. II. i. 168.—Chili.
sphaerocephalus, *Trautv. in Act. Hort. Petrop.* viii. (1883) 341.—Reg. Caucas.
sphaerocystis, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 273 (*Lehm. Rel. Bot.* (1847) 97).—As. centr.
sphaerophysa, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 388.—Soongar.
spicatus, Burm. ex Guill. & Perr. Tent. Fl. Seneg. 182 = Indigofera hirsuta.
spicatus, *Pall. Reise*, ii. 742.—Soongar.
spicatus, Nutt. in Torr. & Gray, Fl. N. Am. i. 336 = Mortoni.
spinellus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 815.—Persia.
spinescens, *Bunge, Astrag.* i. 49; ii. 54.—Persia.
spinosissimus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = A. Tragacantha.
spinosissimus, *Wall. Cat.* n. 5935.—Ind. or.
Sprunerі, *Boiss. Diagn.* Ser. I. ii. 79.—Graecia.
squalidus, *Boiss. l. c.* Ser. II. ii. 38.—As. Min.
squamulosus, Poir. Encyc. Suppl. i. 528 = Oxytropis squamulosa.
squarrosus, *Bunge, Astrag.* i. 129; ii. 236.—Persia.
Stankardi, Boiss. Fl. Orient. ii. 811 = Sahendi.
Stella, Bieb. Fl. Taur. Cauc. ii. 191 = cruciatus.
Stella, *Gouan, Illustr.* 50.—Gall. austr.; Afr. bor.
stellatus, Lam. Fl. Fr. ii. 641 = A. Stella.
stenanthus, *Bunge, Astrag. Fedsch.* 274.—As. temp.
stenocerus, *C. A. Mey. in Bong. Vers. Pfl. Saisang-Nor*, 94.—Soongar.
stenocystis, *Bunge, Astrag. Fedsch.* 301.—As. temp.
stenolepis, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 387.—Persia.
stenolobus, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 593 = ornatodes.
stenophyllus, *Torr. & Gray, Fl. N. Am.* i. 339.—Am. bor.
stenorrhachis, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) ii. 388 = andrachnaefolius.
stenosemius, *Boiss. Diagn.* Ser. II. ii. 27.—Armenia.
stenostegius, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 814.—Kurdistan.
Stevenianus, *DC. Prod.* ii. 285.—Reg. Caucas.; Persia.
Stewartii, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 139.—Ind. bor. occ.
stipitatus, *Benth. ex Boiss. Fl. Orient.* ii. 805.—Afghanist.
stipitatus, Sims, Bot. Mag. t. 2880 = stipulatus.
stipularis, Sims, ex D. Don, l. c. i. Index = seq.
stipulatus, *D. Don, ex Sims, l. c.* t. 2880.—Reg. Himal.
stipulatus, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 131 = graveolens.
Stocksii, *Benth. ex Bunge, Astrag.* i. 6; ii. 4.—Afghan.
stolenburgensis, Lerch. ex Schur, Enum. Pl. Transs. 164 = sriocephalus.
stramineus, *Boiss. Diagn.* Ser. II. v. 85.—Syria.
streptopus, *Greene, in Bull. Calif. Acad.* i. n. 3 (1885) 155.—Calif.
striatellus, *Bieb. Fl. Taur. Cauc.* ii. 189.—Reg. Caucas.
striatus, Nutt. in Torr. & Gray, Fl. N. Am. i. 330 = adsurgens.
strictifolius, *Boiss. Diagn.* Ser. I. ix. 78.—Kurdistan.; Persia.
strictispinus, *Boiss. Fl. Orient.* ii. 868.—Lycia.
strictsemus, Sievers, ex Pall. Astrag. 16 = follicularis.
strictus, *R. Grah. in Wall. Cat.* n. 5924.—Reg. Himal.
strictus, Siev. ex Fisch. in DC. Prod. ii. 301 = follicularis.
strigillosus, *Bunge, Astrag.* i. 102; ii. 178.—Anatolia.
strigulosus, *H. B. & K. Nov. Gen. et Sp.* vi. 494.—Mexic.
strobiliferus, *Lindl. in Bot. Reg.* (1849) *Misc.* 89.—Armenia.
strobilifrus, *Royle, Illustr. Bot. Himal.* 199.—Reg. Himal.
stromatodes, *Bunge, Astrag.* i. 85; ii. 148.—Syria.
subacaulis, Bunge, l. c. i. 101; ii. 178 = karputanus.
subalpinus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 69.—Persia.
subbellatus, *Bunge, Astrag. Fedsch.* 237.—Turkestan.
subbiflorus, Lag. ex DC. Prod. ii. 289 = scorploides.
subblyguus, *Ledeb. Fl. Ross.* i. 627.—Turkestan.
subcanescens, Ledeb. Fl. Ross. i. 644 = Ruprechti, hyalolepis.
subcaulescens, *Royle, Illustr. Bot. Himal.* 199.—Ind. or.
subcinereus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 366.—Am. bor.
subcompressus, *A. Gray, in Brandegee, Fl. S. W. Col.* 214.—Am. bor. occ.
suberosus, [Soland. in] Russell, Aleppo, ed. II. ii. 260 = tuberculosa.
subnitens, *Boiss. & Hohen. Diagn.* Ser. I. ix. 100.—Persia.
subsecundus, *Boiss. & Hohen. l. c.* 54.—Persia.
subsulatus, Desf. Fl. Atlant. ii. 186 = annularis.
subsulatus, Koch, in Flora, ex Steud. Nom. ed. II. i. 159, in syn. = argenteus.
subsulatus, *Pall. ex Bieb. Fl. Taur. Cauc.* ii. 193.—Reg. Caucas.; Tibet.; Afghan.; Sibir.
subsulfereus, *Boiss. & Kotschy, ex Bunge, Astrag.* i. 31; ii. 37.—Cilicia.
subsulfureus, DC. Astrag. 107 = subsulatus.
subumbellatus, Klotzsch, in Bot. Ergeb. Waldem. Reise, 159 = hippocrepicus.
succulentus, Richards. in Frankl. 1st Journ. App. 746 = eucyclocarpus.
succumbens, *Dougl. ex G. Don, Gen. Syst.* ii. 263.—Am. bor. occ.
succumbens, Torr. & Gray, in Pacif. Rail. Rep. ii. 168 = Parryi.
suffulcatus, *Bunge, Astrag.* i. 125; ii. 219.—Afghan.
suffruticosus, *DC. Astrag.* 82.—Sibir.
suffruticosus, Pall. Reise, i. 440 = fruticosus.
suidunensis, *Bunge, in Act. Hort. Petrop.* vii. (1880) 378.—Turkestan.
sulcatus, Lam. Fl. Fr. ii. 639 = austriacus.
sulcatus, *Linn. Sp. Pl.* 756.—Sibir.
sulfureus, *Bunge, Astrag.* i. 29; ii. 33.—Afghan.
sultanensis, Bunge, l. c. i. 17; ii. 19 = bombycinus.
superbus, *Bois. l. c.* i. 63; ii. 105.—Persia.
supinus, *C. A. Mey. ex Bunge, l. c.* i. 51; ii. 83.—Persia.
surgensis, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 402.—Mesopotam.
susianus, *Boiss. Diagn.* Ser. I. ii. 64.—Persia.
susianus, Kotschy, ex Bunge, Astrag. ii. 119 = acutus.
sylvaticus, Willd. Sp. Pl. iii. 1300 = Oxytropis sylvatica.
syloticolus, Pall. Astrag. 98 = Oxytropis sylvatica.
syriacus, *Linn. Sp. Pl.* 760.—Syria.
syriacus, Pall. ex Ledeb. Fl. Alt. iii. 296 = Laxmanni.
syriacus, Linn. Herb. ex Bunge, in Boiss. Fl. Orient. ii. 479 = viminens.
syrtochensis, *Bunge, Astrag.* i. 46; ii. 76.—Persia.
Szovitsii, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 378.—Armenia.

ASTRAGALUS :—
tabrisianus, *Bunge, in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 446.—Persia.
taeniolepis, Boiss. & Hohen. Diagn. Ser. I. ix. 99 = persicus.
talegoncus, *Boiss. & Hohen. l. c.* 87.—Persia.
talassaeus, *Boiss. & Bal. l. c.* Ser. II. vi. 58.—Cappadoc.
talyschensis, *Bunge, Astrag.* i. 40; ii. 57.—Reg. Caucas.
tanaiticus, *C. Koch, in Linnæa*, xxiv. (1851) 94.—Orient.
tarijensis, *Wedd. Chlor. And.* ii. 262.—Bolivia.
taschkendicus, *Bunge, Astrag. Fedsch.* 280.—As. temp.
tataricus, *Franch. Pl. Davidian.* i. 87.—Mongolia.
tauricolus, *Boiss. Diagn.* Ser. II. v. 86.—As. Min.
tauricus, Claus, in Goebel, Reise, ii. 269 = scopaeformis.
tauricus, *Pall. Astrag.* 48. t. 38.—Reg. Caucas.; Ural.; Soongar.
Tavernieri, *Boiss. & Gaill.* Ser. I. ii. 78.—Persia.
tegetarius, *S. Wats. in Bot. King, Exp.* 76, 444.—Am. bor. occ.
teheranicus, *Boiss. & Hohen. in Boiss. Diagn.* Ser. I. ix. 44.—Persia.
tenax, *Bunge, Astrag.* i. 97; ii. 110.—Persia.
tenellus, *Bunge, l. c.* i. 120; ii. 206.—Persia.
tenellus, Pursh, Fl. Am. Sept. ii. 478 = multiflorus.
tenellus, Willd. ex Steud. Nom. ed. II. i. 164 = strigulosus.
tener, *A. Gray, in Proc. Am. Acad.* vi. (1864) 206.—Calif.
tennesseensis, A. Gray, ex Chapm. Fl. S. U. St. 98 = plattensis.
tenuicaulis, *Benth. ex Bunge, Astrag.* i. 93; ii. 98.—Reg. Himal.
tenuifolius, *Desf. Fl. Atlant.* ii. 186.—Algeria.
tenuifolius, Georgi, It. i. 236 = versicolor.
tenuifolius, Linn. Sp. Pl. ed. II. 1065 = A. Onobrychis.
tenuirugis, *Boiss. Diagn.* Ser. I. ix. 61.—Arab.
tenuis, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 90 (*Cat. Baikal.* n. 354).—Sibir. baical.
tenuispinus, Bunge, Astrag. i. 44; ii. 70 = cicerifolius.
tephrodes, *A. Gray, Pl. Wright.* ii. 45.—Mexic.
tephrolobus, *Bunge, Astrag.* i. 127; ii. 222.—Sibir. altaic.
tephrostictes, *Boiss. Diagn.* Ser. I. ix. 69.—Afghan.
teres, Lam. Encyc. i. 324 = Oxytropis floribunda.
terminalis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 370.—Am. bor. occ.
testiculatus, *Pall. Astrag.* 82. t. 67.—Reg. Caucas.; Soongar.; Sibir.
tetragonocarpus, *Boiss. Diagn.* Ser. I. ii. 52.—Persia.
tetrapterus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 369.—Am. bor. occ.
tetrastichus, *Bunge, Astrag. Fedsch.* 279.—As. temp.
thessalus, Boiss. Diagn. Ser. I. ii. 80 = Sprunerі.
Thompsonae, *S. Wats. in Proc. Am. Acad.* x. (1875) 345.—Am. bor. occ.
Thomsonianus, Benth. ex Bunge, Astrag. ii. 324 = nivalis.
thracicus, *Griseb. Spicil. Pl. Rumel.* i. 55.—Thracia.
Thurberi, *A. Gray, in Mem. Am. Acad. Am.* N. S. v. (1855) 319.—Arizona.
tianschanicus, *Bunge, in Rupr. Sert. Tianschi.* 43.—As. temp.
tibetanus, *Benth. ex Bunge, Astrag.* i. 59; ii. 85.—Afghanist.
tigrensis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 193 = venosus.
Tigridis, *Boiss. Diagn.* Ser. I. ix. 55.—Mesopotam.
timuranus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 250.—Turkest.
tmoleus, *Boiss. Diagn.* Ser. I. ix. 79.—Lydia.
toensiolepis, Boiss. & Hohen. ex Boiss. l. c. 90 = persicus.
tokatensis, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 370.—Anatolia.
tomentosus, *Lam. Encyc.* i. 312.—Aegypt.
torrentum, *Bunge, Astrag.* i. 40; ii. 58.—Persia.
tortuosus, *DC. Prod.* ii. 299.—Mesopotam.
Torreyorii, *Boiss. Diagn.* Ser. I. ix. 99.—Cappadoc.
trachycarinatus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 381.—Persia.
trachyrrhinus, *Bunge, Astrag.* i. 101; ii. 178.—Armenia.
Tragacantha, Ucria, Hort. Panor. 318 = siculus.
Tragacantha, Habl. Taur. 112 = armacantha.

ASTRAGALUS:—
Tragacantha, *Linn. Sp. Pl.* 762.—Reg. Medit. or.
Tragacantha, Pall. *Astrag.* t. 4. f. 1. Pl. 9 = mamillensis.
Tragacantha, Vill. Hist. Pl. Dauph. iii. 470 = aristatus.
tragacanthoides, Hohen. Enum. Pl. Talysch. 8 = talyschensis.
tragacanthoides, Lam. ex Steud. Nom. ed. II. i. 164 = declinatus.
tragacanthoides, Linn. Sp. Pl. 762 = longiflorus.
tragacanthoides, Willd. Sp. Pl. iii. 1292 = lanigerus.
tragacanthoides, Pall. Reise, i. 148 = diffusus.
tragacanthus, Lam. Fl. Fr. ii. 642 = A. Tragacantha.
tragodes, Linn. Syst. ed. X. 1175 = longiflorus.
tranoxanus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 388.—Turkestan.
transsilvanicus, *Yanka, in Termész. Füzetek*, viii. (1884) 207.—Europ.
Trautvetteri, *Bunge, Astrag.* i. 89 ; ii. 54.—Soongaria.
triangularis, *DC. Prod.* ii. 291.—Hab. ?
tribuliflolius, *Benth. ex Bunge, Astrag.* i. 4 ; ii. 2.—Tibet. occ.
tribuliformis, Hochst. ex Boiss. Fl. Orient. iii. 225 = tribuloides.
tribuloides, *Delile, Fl. Aegypt. Illustr.* 70.—Afr. bor. ; Orient.
tribuloides, Kotschy, ex Bunge, Astrag. ii. 10 = ammophilus.
tricarinatus, *A. Gray, in Proc. Am. Acad.* xii. (1877) 56.—Am. bor.
trichocalyx, Nutt. ex Torr. & Gray, Fl. N. Am. i. 332 = mexicanus.
trichocalyx, *Trautv. in Act. Hort. Petrop.* iv. (1876) 362.—Reg. Caucas.
trichocarpus, *R. Grah. in Wall. Cat.* n. 5936.—Reg. Himal.
trichocarpus, Young, Fl. Texas, 228 = Nuttallianus.
tricholobus, *DC. Prod.* ii. 299.—Persia.
trichopodus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 218.—Am. bor. occ.
trichopterus, *Boiss. Fl. Orient.* ii. 292.—Syria.
trichostigma, *Bunge, Astrag.* i. 40 ; ii. 57.—Anatolia.
tricolor, *Bunge, i. c.* i. 139 ; ii. 237.—Soongar.
tridactylicus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 557.—Am. bor. occ.
trifloru, *A. Gray, Pl. Wright.* ii. 45.—Am. bor. occ.
trifoliatus, *Phil. in Linnaea*, xxviii. (1856) 681.—Chili.
trifoliolatus, *Boiss. Diagn.* Ser. I. ii. 48.—Syria.
trigonellioides, *Boiss. l. c.* 49.—Armenia.
trigonocarpus, *Bunge, Astrag.* i. 22 ; ii. 28.—As. centr.
trigonus, *DC. Astrag.* 186.—Aegypt. ; Arab.
trigonus, Sieber, ex DC. Prod. ii. 295 = Sieberi.
trimestris, Bieb. Fl. Taur. Cauc. ii. 190 = brachycerus.
trimestris, *Linn. Sp. Pl.* 761.—Aegypt.
trimestris, Pall. ex Ledeb. Fl. Ross. i. 668 = hamosus.
trimorphus, Viv. Fl. Lyb. Spec. 44. t. 10 = annularis.
triphyllus, Pall. Astrag. 68. t. 56. f. a = Oxytropis triphylla.
triphyllus, *Pursh, Fl. Am. Sept.* ii. 740.—Am. bor.
triquetrus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 267.—Am. bor.
tristachyus, *Bunge, Astrag.* i. 10 ; ii. 7.—Cilicia ; Mesopotam.
tristis, Nutt. in Torr. & Gray, Fl. N. Am. i. 336 = Mortoni.
troglodytus, *S. Wats. in Proc. Am. Acad.* xx. (1885) 362.—Arizona.
trojanus, *Stev. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 401.—Graecia ; As. Min. ; Orient.
tschanjensis, *Bunge, Astrag.* i. 23 ; ii. 24.—Sibir. altaic.
tuberculosus, *DC. Astrag.* 33. t. 22.—Syria ; Mesopotam.
tumidus, Bieb. Fl. Taur. Cauc. ii. 198 = fabaceus.
tumidus, DC. Astrag. 166 = Russelii.
tumidus, Willd. in Mém. Acad. Berl. (1794–95) 26 = Clusii.
tunetanus, *Willd. Sp. Pl.* iii. 1277.—Algeria.
turbinatus, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 381.—Persia ; Turkestan.
turcomanicus, *Bunge, Astrag.* i. 130 ; ii. 237.—Turcom.
Turczaninowii, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 242.—Turkestan.
turkestanus, *Bunge, ex Boiss. Fl. Orient.* ii. 278.—Turkestan.
tymphresteus, *Boiss. & Sprun. Diagn.* Ser. I. ii. 48. —Graecia.

ASTRAGALUS:—
tyrolensis, Sieber, ex Steud. Nom. ed. II. i. 164 = Oxytropis campestris.
ukrainicus, *Franch. Pl. Davidian.* i. 87.—Mongolia.
uliginosus, Bieb. Fl. Taur. Cauc. ii. 188 = glycyphylloides.
uliginosus, *Linn. Sp. Pl.* 757.—Sibiria.
umbellatus, *Bunge, Astrag.* i. 24 ; ii. 29.—N. Zemlaia.
umbellatus, Pers. Syn. ii. 335 = lotoides.
uncatus, *Linn. Sp. Pl.* ed. II. 1072.—Aleppo.
uncatus, Linn. Herb. ex Boiss. Fl. Orient. ii. 284 = trimestris.
uncinatus, *Bertol. Fl. Ital.* viii. 54.—Ins. Corsica.
uncinatus, Moench, Meth. 166 = baeticus.
uncinatus, *Pomel, Nouv. Mat. Fl. Atl.* 322.—Algeria.
uniflorus, *DC. Astrag.* 248. t. 50.—Peruv.
unifoliatus, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 231.—Turkestan.
unifultus, *L'Hérit. Stirp. Nov.* 168.—Peruv. ; Chili.
unijugus, *Bunge, Astrag.* i. 130 ; ii. 228.—Soongaria.
unilateralis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 404.—Sibir. altaic.
uralensis, Linn. Fl. Suec. 557 = Oxytropis campestris.
uralensis, Vahl, in Fl. Dan. t. 1041 = Oxytropis sordida.
uraniolimneus, *Boiss. Fl. Orient.* ii. 280.—Armenia.
urmiensis, *Bunge, Astrag.* i. 87 ; ii. 44.—Persia.
ursinus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 267. —Am. bor.
ustiortensus, *Bunge, Astrag.* i. 102 ; ii. 179.—Reg. Casp.
utahensis, *Torr. & Gray, in Pacif. Rail. Rep.* ii. (1854) 120.—Am. bor. occ.
utriger, Friv. ex Bunge, Astrag. ii. 108 = physocalyx.
utriger, Boiss. & Balse, ex Boiss. Fl. Orient. ii. 231 = macropelmatus.
utriger, *Pall. Astrag.* 75. t. 61.—Tauria.
utriger, C. A. Mey. Verz. Pfl. Cauc. 148 = bakuensis.
utriger, Willd. ex Ledeb. Fl. Ross. i. 654 = pubiflorus.
vaccarum, *A. Gray, Pl. Wright.* ii. 43.—Am. bor. occ.
vaginatus, *DC. Astrag.* 210. t. 37.—Cilicia.
vaginatus, *Pall. Astrag.* 46. t. 36.—Sibiria.
vaginatus, Richards. in Frankl. 1st Journ. App. 746 = Hallii.
valesianus, Thomas, ex Koch, Syn. Fl. Germ. ed. I. 182 = Oxytropis lapponica.
Vanillae, *Boiss. Diagn.* Ser. I. ii. 60.—Persia.
variabilis, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 33.—Mongolia.
variegatus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 289.—Turkest.
varius, S. G. Gmel. It. i. 116. t. 17 = virgatus.
Vaseyi, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 370.—Am. bor. occ.
vegetus, *Bunge, Astrag.* i. 103 ; ii. 181.—Persia.
vegetus, Sadl. Fl. Pest. i. 199 in nota = Muelleri.
veluchensis, *Boiss. Fl. Orient.* ii. 319.—Graecia.
velutinus, Sieber, ex Steud. Nom. ed. II. i. 164 = Oxytropis Halleri.
venosus, Aucher, ex Bunge, Astrag. ii. 217 = Aucheri.
venosus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 193. —Abyss.
venosus, Ehrenb. ex Bunge, Astrag. ii. 229 = angustifolius.
ventorum, *A. Gray, ex S. Wats. in Am. Natural.* vii. (1874) 115.—Am. bor. occ.
venulosus, *Boiss. Diagn.* Ser. I. ii. 75.—Cappadoc.
verus, *Georgi, Bemerk. Reise Russ. Reich.* i. 226.—Rossia.
verrucosus, *Moris, Stirp. Sard. Elench.* 13.—Sardinia.
versicolor, C. Koch, in Linnaea, xix. (1847) 65 = Sibthorpianus.
vesiculosus, *Pall. Astrag.* 45. t. 35.—Sibiria or.
verticillaris, *Bunge, Astrag.* i. 53 ; ii. 88.—Afghanist.
verticillaris, Linn. Mant. ii. 275 = Oxytropis myriophylla.
verticillaris, Georgi, Beschr. Russ. Reich. iii. iv. 1155 = Oxytropis verticillaris.
verticillaris, *Scop. Delic. Insub.* ii. 107.—Sibiria.
verus, DC. ex Bunge, Astrag. ii. 164 = Russeanus.
verus, *Olivier, Voy.* iii. t. 44.—Persia.
vesicarius, Lam. Fl. Fr. ii. 637 = A. Cicer.
vesicarius, *Linn. Sp. Pl.* 760.—Europ. austr. ; Tauria.
vesicarius, Pall. Reise, iii. App. 751 = Oxytropis caespitosa.
vesicarius, Siev. in Pall. N. Nord. Beitr. vii. 274 = inflatus.
vesiculosus, *Clos, in C. Gay, Fl. Chil.* ii. 190.—Chili.

ASTRAGALUS:—
vesiculosus, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. iii. 138 = mollis.
vestitus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 98.—Graecia.
vestitus, *S. Wats. Bibliog. Ind.* 202.—Calif.
vexillaris, *Boiss. Diagn.* Ser. I. ii. 49.—Mesopotam.
vicinesolius, *DC. Astrag.* 119.—Reg. Caucas. ; As. Min.
vicioides, *R. Grah. in Wall. Cat.* n. 5931.—Reg. Himal.
vicioides, Ledeb. Fl. Alt. iii. 301 = macropterus.
villosissimus, *Bunge, Astrag.* i. 129 ; ii. 225.—Turkestan.
villosus, Gueldenst. ex Ledeb. Fl. Ross. i. 654 = pubiflorus.
villosus, *Michx. Fl. Bor. Am.* ii. 67.—Am. bor.
vimineus, Pall. Astrag. 34. t. 21 = fruticosus.
virescens, Ait. Hort. Kew. iii. 78 = falcatus.
virgatus, *Pall. Astrag.* 20. t. 18.—Hung. ; Taur.
virgatus, Reichb. Fl. Germ. Excurs. 510, 958 = Muelleri.
vividis, *Bunge, Astrag.* i. 185 ; ii. 231.—Armenia.
viscosus, Chaix, ex Vill. Prosp. 42 = Oxytropis campestris.
Volckmanni, *Phil. in Anal. Univ. Chil.* (1861) i. 46. —Chili.
Vulnerariae, *DC. Prod* ii. 300.—As. Min.
vulnerarioides, Ait. Fl. Pedem. i. 343 = Anthyllis Vulneraria.
vulpinus, Ledeb. Fl. Alt. iii. 318 = lagocephalus.
vulpinus, *Willd. Sp. Pl.* iii. 1259.—Turkestan.
Wagneri, *Bartl. ex Bunge, Astrag.* ii. 125.—Persia.
Wardii, *A. Gray, in Proc. Am Acad.* xii. (1877) 55.—Utah.
wrioensis, *Boiss. & Kotschy, ex Bunge, Astrag.* i. 87 ; ii. 153.—Armenia.
Webberi, *A. Gray, in Bot. Calif.* i. 154.—Calif.
Webbianus, *R. Grah. in Wall. Cat.* n. 5936.—Reg. Himal.
Weidmannianus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) ii. 396.—Anatolia.
Whitneyi, *R. Gray, in Proc. Am. Acad.* vi. (1864–65) 536.—Calif.
Wilhelmsii, *Fisch. ex Ledeb. Fl. Ross.* i. 668 = resupinatus.
wingatanus, *S. Wats. in Proc. Am. Acad. xviii.* (1883–83) 192.—Am. bor. occ.
wolgensis, *Bunge, Astrag.* i. 26 ; ii. 47.—Rossia.
Wrightii, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 176.—Texas.
Wulfeni, *Koch, Syn. Fl. Germ.* ed. II. 207.—Dalmatia; Graecia.
xanthomelas, *Bunge, Astrag. Fedsch.* 314.—As. temp.
xanthotrichos, *Ledeb. Fl. Alt.* iii. 324.—Sibir. altaic.
xerophilus, *Ledeb. Fl. Ross.* i. 667.—Mont. Ararat.
Xiphidium, *Bunge, Astrag.* i. 25 ; ii. 213.—Reg. Caucas.
xiphocarpus, *Benth. ex Bunge, l. c.* i. 27 ; ii. 32.—Reg. Himal.
xylorrhizus, *Bunge, Astrag. Fedsch.* 267.—As. temp.
yuralicus, *Boiss. Fl. Orient.* ii. 296.—Cappadocia.
zacharensis, *Bunge, Astrag.* i. 22 ; ii. 27.—Syria.
zahnsiana, *Bunge, l. c.* i. 88 ; ii. 154.—Syria.
zagrosicus, *Boiss. & Haussk. ex Boiss Fl. Orient.* ii. 286.—Persia.
zanatanensis, *Benth. ex Bunge, Astrag.* i. 43 ; ii. 67.—Reg. Himal.
zerdaus, *Boiss. Diagn.* Ser. I. ii. 51.—Persia.
zeylanicus, Ledeb. ex Bunge, Astrag. ii. 184 = caeruleus.
Zohrabi, Bunge, l. c. i. 68 ; ii. 112 = lagopoides.

ASTRANTHIUM, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 312 = Bellis, Linn. (Compos.).
integrifolia, Nutt. l. c. = B. integrifolia.

ASTRANTHUS, Lour. Fl. Cochinch. 221 (1790) = Homalium, Jacq. (Samyd.).
cochinchinensis, Lour. l. c. 222 = H. tomentosum.
foetida, Wall. ex Mast. in Hook. f. Fl. Brit. Ind. ii. 598 = H. foetidum.

ASTRANTIA, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 879.
ACTINOLEMA, Fenzl, Pugill. Pl. Syr. 66. t. 19 (1849).
alpina, Star, ex Nym. Consp. 318 = gracilis.
alpina, [Chlivr.] Mor. Herb. 78 = minor.
bavarica, Schult. in Flora, xli. (1858) 121 = gracilis.

ASTRANTIA:—
Bieberstein, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 28.—Reg. Caucas.
candida, Mill. Gard. Dict. ed. VIII. n. 2 = major.
carinthiaca, Hoppe, ex Koch, Syn. Fl. Germ. ed. I. 281 = major.
carniolica, *Wulf. in Jacq. Fl. Austr.* v. 31.—Carinth.
carniolica, Koch, Syn. Fl. Germ. ed. I. 280 = gracilis.
caucasica, Willd. ex Spreng. Umb. 17 = major.
ciliaris, *Linn. f. Suppl.* 177.—Afr. austr.
croatica, *Tomm. ex Vukot. in Rad. Jugos. Akad. Zagrab*, xliv. (1878) 216.—Croatia.
Diapensia, Scop. Fl. Carn. ed. II. i. 186 = Sanicula europæa.
digitata, Moench, Meth. 94 = minor.
dondiaefolia, Schur, Enum. Pl. Transs. 245 = gracilis.
elatior, Frivald. in Flora, xix. (1836) 484 = major.
Epipactis, Scop. Fl. Carn. ed. II. i. 185 = Hacquetia Epipactis.
europæa, Linn. ex Rehm. Notiz. Veg. Gest. Schwarz. Merr. 63 = major!
gracilis, *Bartl. in Linnæa*, xv. (1841) *Litt.* 93.—Carniolia.
gracilis, Schur, in Oestr. Bot. Zeitschr. (1858) 22 = carniolica.
grandiflora, Poepp. ex DC. Prod. iv. 66 (= *Posoa corsacea*).—Reg. Andin.
helleborifolia, *Salisb. Prodr.* 159.—Reg. Caucas.
heterophylla, Willd. in Ges. Naturf. Fr. Berl. Neue Schr. iii. (1801) 419 = helleborifolia.
intermedia, Bieb. Fl. Taur. Cauc. Suppl. 194 = major.
major, Bieb. l. c. n. 509 = Bieberstein.
major, *Linn. Sp. Pl.* 236.—Reg. Caucas.
maxima, Pall. in Nov. Act. Petrop. vii. (1798) 357. t. 11 = helleborifolia.
microphylla, Schur, Enum. Pl. Transs. 245 = gracilis.
minor, *Linn. Sp. Pl.* 235.—Europ.
minor, Scop. Fl. Carn. ed. II. i. 187 = carniolica.
minor, Vitm. Summa Veg. ii. 194 = pauciflora.
montana, (Clairv.) Man. Herb. 78 = major.
neglecta, *C. Koch & Bouché, in App. Sem. Hort. Berol.* (1858) 22.—Europ.
nigra, Scop. Fl. Carn. ed. II. i. 188 = major.
pallida, Presl, ex Koch, Syn. Fl. Germ. ed. I. 280; ed. II. 309 = major.
pauciflora, *Bertol. in Desv. Journ. Bot.* ii. (1813) 76.—Italia.
rauscudifolia, *Reichb. f. Ic. Fl. Germ.* xxi. 3.—Europ.
sanicvioefolia, Salisb. Prod. 159 = major.
speciosa, Hort. ex Heynh. Nom. ii. 53 = helleborifolia.
transsilvanica, Schur, Enum. Pl. Transs. 245 = carniolica.
tridentata, *Parrot, Reise Krym Kauk.* ii. 130, nomen. —Reg. Caucas.
trifida, Hoffm. Umb. ed. II. 1. præf. 8 in nota = major.

ASTRAPAEA, Lindl. Coll. Bot. t. 14 (1822) = Dombeya, Cav. (Sterculi.).
floribunda, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 90 = D. Wallichii.
tiliæfolia, Sweet, Hort. Brit. ed. I. 58 = D. acutangula.
viscosa, Sweet, l. c. = D. cannabina.
Wallichii, Lindl. Coll. Bot. t. 14 = D. Wallichii.

ASTRELIA, F. Muell. [*Fragm.* x. 76 (1876), nomen] ex Benth. Fl. Austral. vii. 602 (1878). *GRAMINEÆ*, Benth. & Hook. f. iii. 1170.
etymologia, *F. Muell. ex F. M. Bailey, Grass. Queensl.* i. —Austral.
pectinata, *F. Muell. ex Benth. Fl. Austral.* vii. 602.—Austral.
tridentata, *F. Muell. l. c.*—Austral.

ASTREPHIA, Dufr. Hist. Valer. 50 (1811), pro parte. *VALERIANEÆ*, Benth. & Hook. f. ii. 155.
OLIGACOCE, Willd. ex DC. Prod. iv. 623 (1830).
carnosa, Dufr. l. c. β i = Valeriana carnosa.
chaerophylloides, *DC. Prod.* iv. 629.—Peruv.
chinensis, *Dupf. Hist. Valer.* 53.—China.
coarctata, Dufr. l. c. = Valeriana coarctata.
crispa, *Dufr. l. c.*—Chili.
hyalinorrhina, Dufr. l. c. 62 = Valeriana hyalinorrhina.
interrupta, Dufr. l. c. 41 = Valeriana interrupta.
laciniata, Dufr. l. c. 56 = chaerophylloides.
laxa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 364. —Chili.
lobata, *Hook. & Arn. l. c.*—Chili.

ASTREPHIA:—
mexicana, Hook. & Arn. Bot. Beech. Voy. Suppl. 481 = Phyllactis mexicana.
mexicana, Seem. Bot. Voy. Herald, 297 = Valeriana mexicana.
oblongifolia, Dufr. Hist. Valer. 60 = Valeriana oblongifolia.
polystachya, Dufr. l. c. 51 = Valerianopsis polystachya.
pratensis, Benth. Pl. Hartw. 39 = Phyllactis pratensis.
serrata, Dufr. Hist. Valer. 50 = Valeriana serrata.

ASTROCARPUS, Neck. Elem. ii. 248 (1790). *RESEDACEÆ*, Benth. & Hook. f. i. 111.
SESAMELLA, Reichb. Consp. 186 (1828).
abyssinicus, Hochst. ex Oliver, Fl. Trop. Afr. i. 103 = Cayluca abyssinica.
Clusii, *Y. Gay, in F. Schultz, Arch. Fl. Fr. et Allem.* 33.—Hispan.
cochlearifolius, *Nym. in Obser. Vet. Acad. Stockh.* xvIII. (1841) 191.—Lusitan.
interruptus, Bor. in Mém. Soc. Acad. Maine-et-Loire, viii. (1850) 7 = sesamoides.
minor, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 86 (*Pugill.* 295. t. 4. f. 1).—Europ.
purpurascens, Kaſn. Fl. Tellur. iii. 78 = Clusii.
sesamoides, Colm. ex Willk. & Lange, Prod. Fl. Hisp. iii. 899 = Clusii.
sesamoides, *DC. ex Duby, Bot. Gall.* i. 67.—Europ.
spathulæfolius, *Reveil, ex Bor. in Mém. Soc. Acad. Maine-et-Loire*, i. (1857) 17.—Gallia.
suffrutitus, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 83 (*Pugill.* 275. t. 4).—Hispan.

ASTROCARYUM, G. F. W. Mey. Prim. Fl. Esseq. 205 (1818). *PALMÆ*, Benth. & Hook. f. iii. 942.
TOXOPHORNIX, Schott, Nachr. K. K. Oester. Naturf. in Brasil. Ash. ii. (1832).
acanthopodium, *Barb.-Rodr. Enum. Palm. Nov.* 20.—Bras.
acaule, *Mart. Hist. Nat. Palm.* ii. 78. t. 24. 63. f. 5. —Bras.
aculeatum, Barb.-Rodr. Enum. Palm. Nov. 20 = Rodriguesii.
aculeatum, *G. F. W. Mey. Prim. Fl. Esseq.* 206.—Guiana.
aculeatum, Hort. ex H. Wendl. in Kerch. Palm. 280 = Acrocomia sclerocarpa.
aculeatum, Wallace, Palm Trees Amaz. 111. t. 43 = minus!
argenteum, *Hort. Bull.—Cf. Gard. Chron.* (1875) i. 491.—Hab. ?
aureum, *Griseb. & H. Wendl. in Griseb. Fl. Brit. W. Ind.* 521.—Ind. occ.
Aurora, De Vriese, in Jaarb. Maatsch. (1846) 12 = aculeatum.
Ayri, *Mart. Hist. Nat. Palm.* ii. 71. t. 59*.—Bras.
Borsigianum, C. Koch, ex Regel, Gartenfl. (1861) 29 = Syrrhopea grandifolia.
campestre, *Mart. Hist. Nat. Palm.* ii. 79. t. 63. f. 4. —Bras.
candescens, *Barb.-Rodr. Enum. Palm. Nov.* 22.—Bras.
Chichon, Linden, Illustr. Hortic. xxviii. (1881) 15 = mexicanum.
Ferdin, *Mart. in Palmet. Orb.* 64. t. 4. f. 1. 2.—Peruv.
farinosum, *Barb.-Rodr. Enum. Palm. Nov.* 21.—Bras.
ferrugineum, *Hort. ex H. Wendl. in Kerch. Palm.* 282. —Hab. ?
guianense, *Splitg. ex Mart. Hist. Nat. Palm.* iii. 233. —Guiana.
gynacanthum, *Mart. l. c.* ii. 71. t. 60, 61.—Bras.
gynacanthum, Wallace, Palm Trees Amaz. 108. t. 39 = A. Mumbaca!
Huaimi, *Mart. in Palmet. Orb.* 86. t. 19. fig. 3.—Bolivia.
humile, *Wallace, Palm Trees Amaz.* 115. t. 45.—Bras.
Jauari, *Mart. Hist. Nat. Palm.* ii. 76. t. 25 et 65. f. 1. —Bras.
fucuma, Linden, Illustr. Hortic. xxviii. (1881) 15, sphalm. = A. Tucuma.
Malybo, *Karst. in Linnæa*, xxvIII. (1856) 245.—N. Granat.
mexicanum, *Liebm. ex Mart. Hist. Nat. Palm.* iii. 233. —Mexic.

ASTROCARYUM:—
minus, *Trail, in Journ. Bot.* xv. (1877) 76.—Bras.
Munbaca, *Mart. Hist. Nat. Palm.* ii. 74.—Bras.
Murumuru, *Mart. l. c.* 70. t. 58, 59.—Bras.
Murumuru, *Wallace, Palm Trees Amaz.* 101. t. 38. —Bras.
panamanum, *Linden, Illustr. Hortic.* xxxvIII. (1881) 15. —N. Granat.
Paramaca, *Mart. in Palmet. Orb.* 88.—Guiana.
plicatum, *Drude, in Mart. Fl. Bras.* iii. ii. 375.—Guiana; Bras.
Princeps, *Barb.-Rodr. Enum. Palm. Nov.* 22.—Bras.
pygmæum, *Drude, in Mart. Fl. Bras.* iii. ii. 385.—Bras.
pumilum, *Hort. ex H. Wendl. in Kerch. Palm.* 282.—Hab. ?
Rodriguesii, *Trail, in Journ. Bot.* xv. (1877) 79.—Bras.
rostratum, *Hook. Bot. Mag.* t. 4778.—Bras.
sclerocarpum, Hort. ex H. Wendl. in Kerch. Palm. 282 = Acrocomia sclerocarpa.
sclerophyllum, *Drude, in Mart. Fl. Bras.* iii. ii. 377 —Bras.
segregatum, *Drude, l. c.* 382.—Bras.
tenuifolium, *Linden, Illustr. Hortic.* xxvIII. (1881) 15. —Bras.
Tucuma, *Mart. Hist. Nat. Palm.* ii. 77. t. 65. f. 2. —Bras.
tucumoides, *Drude, in Mart. Fl. Bras.* iii. ii. 381.—Bras.
vulgare, *Mart. Hist. Nat. Palm.* ii. 74. t. 62, 63.—Bras.
Warszewiczii, *Karst. in C. Koch & Fint. Wochenschr.* (1858) 297.—Am. austr.
Weddellii, *Drude, in Mart. Fl. Bras.* iii. ii. 383.—Bras.

ASTROCEPHALUS, Raſn. Fl. Tellur. iv. 94 (1836) = ASTROCEPHALUS, Vaill. = Scabiosa, Tourn. (Dipsac.).
ciliata, *integrifolia*, *rigida*, *sylvatica*, *tatarica*, *uralensis*, Raſn. l. c. = Scabiosa, ejusdem homonymus.

ASTROCOCCUS, Benth. in Hook. Kew Journ. vi. (1854) 327. *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 326.
coriaceus, *Baill. in Adansonia*, v. 305.—Bras.
cornutus, *Benth. in Hook. Kew Journ.* vi. (1854) 327. —Bras.

ASTROCOMA, Neck. Elem. i. 119 (1790) = Stenvia, Dahl (Brunlac.).

ASTRODENDRUM, Dennst. Schluss. Hort. Malab. 30 (1818) = Sterculia, Linn.
malabaricum, Dennst. l. c. = S. guttata.

ASTROGLOSSUS, Reichb. f. ex Benth. & Hook. f. Gen. iii. 588 (1883) = Trichocoros, H. B. & K. (Orchid.).

ASTROGYNE, Benth. Pl. Hartw. 14 (1839) = Croton, Linn. (Euphorb.).
crotonoides, Benth. l. c. = C. gracilis.

ASTROGYNE, Wall. ex Laws. in Hook. f. Fl. Brit. Ind. i. (1875) 629 = Stiphonodon, Griff. (Celastr.).
coriacea, Wall. ex Laws. l. c. = S. celastrineus.

ASTROLOBIUM, DC. Prod. ii. 311 (1825) = Ornithopus, Linn. et Coronilla, Linn. (Leguvin.).
cretheum, DC. l. c. 310 = C. cretica.
durum, DC. l. c. 311 = O. durus.
ebracteatum, DC. l. c. = O. ebracteatum.
repandum, DC. l. c. = C. repanda.
scorpioides, DC. l. c. = C. scorpioides.

ASTROLOMA, R. Br. Prod. 556 (1810). *EPACRIDEÆ*, Benth. & Hook. f. ii. 611.
MESOMELINA, Steudel. in Bull. Soc. Nat. Mosc. (1839) 1. 6.
PENTATAPHRUS, Schlecht. in Linnæa, xx. (1847) 618.
STENANTHERA, R. Br. Prod. 556 (1810).
STOMARRHENA, DC. Prod. viii. 700 (1844).
VENTENATIA, Cav. Ic. iv. 79 (1797).
VINTENATIA, Cav. l. c. 78 (1797).
Baxteri, *A. Cunn. ex DC. Prod.* viii. 700.—Austral.
Candollenaga, *Sond. in Lehm. Pl. Preiss.* i. 302.—Austral.
compactum, *R. Br. Prod.* 556.—Austral.

ASTROLOMA:—
conostephioides, *F. Muell. ex Benth. Fl. Austral.* iv. 158.—Austral.
cuneifolium, Sond. in Lehm. Pl. Preiss. i. 299 = compactum.
denticulatum, R. Br. Prod. 538 = humifusum.
dilatatum, Sond. in Lehm. Pl. Preiss. i. 298 = longiflorum.
dilcolor, Sond. l. c. = longiflorum.
divaricatum, *Sond. l. c.* 298.—Austral.
Drummondii, *Sond. l. c.*—Austral.
foliosum, *Sond. l. c.* 297.—Austral.
glaucescens, Sond. l. c. 298 = macrocalyx.
hirsutum, Stschegl. in Bull. Soc. Nat. Mosc. (1859) i. 7 = Drummondii.
humifusum, *R. Br. Prod.* 538.—Austral.
jumiperinum, F. Muell. Fragm. vi. 86 = microphyllum.
lasiocnemum, F. Muell. l. c. 40 = stomarrhena.
latifolium, Sond. in Lehm. Pl. Preiss. i. 302 = tectum.
longiflorum, *Sond. l. c.* 297.—Austral.
macrocalyx, *Sond. l. c.* 301.—Austral.
marginatum, Sond. l. c. 299 = divaricatum.
microcalyx, *Sond. l. c.* 298.—Austral.
microdonta, *F. Muell. ex Benth. Fl. Austral.* iv. 155. —Austral.
microphyllum, *Stschegl. in Bull. Soc. Nat. Mosc.* (1859) i. 7.—Austral.
pallidum, *R. Br. Prod.* 538.—Austral.
pallidum, Sond. in Linnaea, xxvi. (1853–55) 246 = humifusum.
pinifolium, *Benth. Fl. Austral.* iv. 159.—Austral.
prostratum, *R. Br. Prod.* 538.—Austral.
pungens, Stschegl. in Bull. Soc. Nat. Mosc. (1859) i. 8 = divaricatum.
splendens, Planch. in Fl. des Serres, Sér. I. x. (1854–55) 199. t. 1018 = divaricatum.
stomarrhena, *Sond. in Lehm. Pl. Preiss.* i. 301.—Austral.
tectum, *R. Br. Prod.* 538.—Austral.
xerophyllum, *Sond. in Lehm. Pl. Preiss.* i. 301.—Austral.

ASTRONIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. J. Art. IV. 1; Blume, Bijdr. 1080 (1826). *MELASTOMACEAE*, Benth. & Hook. f. i. 771.
ASTRONIDIUM, A. Gray, Bot. U. St. Expl. Exped. i. 583. t. 72 (1854).
LOMANODIA, Rafin. Sylva Tellur. 97 (1838).
NAUDINIA, Decne. ex Triana, in Seem. Fl. Vit. 86 (1866).
concolor, Zipp. ex Steud. Nom. ed. II. i. 165 = macrophylla.
confertiflora, *A. Gray, Bot. U. St. Expl. Exped.* i. 579. —Ins. Fiji.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 174.— Ins. Philipp.
Forsteri, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 258.—Ins. Tahiti.
fraterna, *A. Gray, Bot. U. St. Expl. Exped.* i. 576. t. 72.—Ins. Tahiti.
fraterna, Seem. Fl. Vit. 85, 86 = Forsteri.
glabra, Triana, in Trans. Linn. Soc. xxvii. (1871) 152 = Forsteri.
intermedia, Blume, Mus. Bot. Lugd. Bat. i. 9 = speciabilis.
macrophylla, *Blume, Bijdr.* 1080.—Malaya.
ovalifolia, *Naud. ex Decne. Bot. Voy. Vénus*, 27.—Ins. Tahiti.
papetaria, *Blume, Bijdr.* 967.—Ins. Moluce.
parviflora, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 152.—Ins. Fiji.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 577. t. 72.—Ins. Fiji.
pilosula, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 152.—Ins. Tahiti.
roberta, *Seem. Fl. Vit.* 86.—Ins. Fiji.
Rolfei, *Vidal, Phan. Cuming. Philipp.* 174.—Ins. Philipp.
samoënsis, *S. Moore, in Journ. Bot.* xvIII. (1880) 8.— Ins. Samoa.
sessilifolia, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 152.—Malaya.
spectabilis, *Blume, Bijdr.* 1080.—Java.
spectabilis, Zipp. ex Steud. Nom. ed. II. i. 165 = papetaria.
Storckii, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 152.—Ins. Fiji.

ASTRONIA:—
subcordata, *A. Gray, Bot. U. St. Expl. Exped.* i. 580. —Ins. Samoa.
tomentosa, *Seem. Fl. Vit.* 86.—Ins. Fiji.

ASTRONIDIUM, A. Gray, Bot. U. St. Expl. Exped. i. 581. t. 72 (1854) = **Astronia**, Blume (Melast.).
parviflorum, A. Gray, l. c. 582. t. 72 = Astronia parviflora.
Storckii, Seem. Fl. Vit. 87 = Astronia Storckii.

ASTRONIUM, Jacq. Enum. Pl. Carib. 10. *ANACARDIACEAE*, Benth. & Hook. f. i. 423.
Balansae, *Engl. in Bot. Jahrb.* i (1881) 45.—Paraguay.
Candollei, *Engl. l. c.*—Paraguay.
conchinum, *Schott, in Spreng. Syst. Cur. Post.* 404.— Bras.
fraxinifolium, *Schott, in Spreng. l. c.*—Bras.
gracile, *Engl. Bot. Jahrb. i.* (1881) 75.—Bras.
graveolens, *Jacq. Enum. Pl. Carib.* 33.—N. Granat.
insigne, March. Rev. Anac. 177 = Parishia insignis.
juglandifolium, Griseb. in Goett. Abh. xxiv. (1879) 94 = A. Urundeuva.
Libonianum, *T. Linden, Cat.* n. 17. (1862) 7.—Bras.
macrocalyx. *Engl. Bot. Jahrb.* i. (1881) 75.—Bras.
obliquum, *Griseb. Fl. Brit. W. Ind.* 176.—Ind. occ.
Planchonianum, *Engl. in Mart. Fl. Bras.* xii. ii. 399. —Am, trop.
Urundeuva, *Engl. Bot. Jahrb. i.* (1881) 45.—Bras.

ASTROPANAX, Seem. in Journ. Bot. iii. 176 (1865) = **Heptapleurum**, Gaertn. (Araliac.).
abyssinicum, Seem. l. c. 177 = H. abyssinicum.
Baikiei, Seem. l. c. = H. Baikiei.
Barteri, Seem l. c. = H. Barteri.
elatum, Seem. l. c. = H. elatum.
Mannii, Seem. l. c. = H. Mannii.

ASTROPETALUM, Griff. Notul. iv. 411. t. 565 (1854) = **Swintonia**, Griff. (Anacard.).

ASTROPHEA, Reichb. Consp. 182 (1828) = **Passiflora**, Linn.
emarginata, M. Roem. Syn. Pepon. 159 = P. emarginata.
glaberrima, Klotzsch, in Schomb. Fauna & Fl. Guy. 1155.—Guiana.
glauca, M. Roem. Syn. Pepon. 161 = P. arborea.
ovata, M. Roem. l. c. = P. ovata.

ASTROPHIA, Nutt. in Torr. & Gray, Fl. N. Am. i. 278 (1838) = **Lathyrus**, Linn. (Legum.).
littoralis, Nutt. l. c. = L. littoralis.

ASTROPHYLLUM, Torr. & Gray, in Bot. Pope Exped. (Pacif. Rail. Rep. ii.) 161 (1855). *RUTACEAE*, Benth. & Hook. f. i. 396.
dumosum, *Torr. & Gray, l. c.*—Am. bor. occ.

ASTROPHYTUM, Lem. Cact. Gen. Nov. Gen. et Sp. 3 (1839) = **Echinocactus**, Link & Otto (Cact.)
Asterias, Lem. Cact. 50 = E. Asterias.
myriostigma, Lem. Cact. Gen. Nov. et Sp. 4 = E. myriostigma.
prismaticum, Lem. Cact. 50 = E. myriostigma.

ASTROPUS, Spreng. Neue Entdeck. iii. 64 (1822) = **Waltheria**, Linn. (Sterc.).
tomentosus, Spreng. l. c. = W. Astropus.

ASTROSCHOENUS, Lindl. Veg. Kingd. 119 (1847) = **Asterochoenus**, Nees (Cyperac.).

ASTROSTEMMA, Benth. in Hook. Ic. Pl. xiv. 7. t. 1311 (1880). *ASCLEPIADEAE*.
spartioides, *Benth. l. c.*—Borneo.

ASTROTRICHA, DC. Coll. Mém. v. 29 t. 5 (1829); Prod. iv. 74 (1830). *ARALIACEAE*, Benth. & Hook. f. i. 937.
asperifolia, F. Muell. ex Klatt, in Linnaea, xxix. (1858) 709 = latifolia.
floccosa, *DC. Coll. Mém.* v. 30. t. 5.—Austral.
Hamptoni, *F. Muell. Fragm.* vi. 126.—Austral.
horsfeldes, *A. Cunn. ex Benth. Enum. Pl. Huegel.* 55. —Austral.
latifolia, *Benth. l. c.* 54.—Austral.

ASTROTRICHA:—
ledifolia, *DC. Coll. Mém.* v. 30. t. 6.—Austral.
linearis, *A. Cunn. ex Benth. Enum. Pl. Huegel.* 55.— Austral.
longifolia, *Benth. l. c.*—Austral.
pterocarpa, *Benth. Fl. Austral.* iii. 379.—Austral.

ASTYDAMIA, DC. Coll. Mém. v. 58 (1829); Prod. iv. 190 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 909.
canariensis, *DC. Prod.* iv. 190.—Ins. Teneriff.
latifolia, Baill. Hist. Pl. vii. 208 = canariensis.

ASTYLIS, Wight, Ic. t. 1992 (1853) = **Hemicyclia**, Wight & Arn. (Euphorb.).
venusta, Wight, l. c. = H. venusta.

ASTYLUS, Dulac, Fl. Hautes-Pyr. 188 (1867) = **Rutobinasia**, R. Br. (Crucif.).
alpinus, Dulac, l. c. = H. alpina.
petraea, Dulac, l. c. = H. petraea.

ASTYRIA, Lindl. Veg. Kingd. 864 (1847) = **Astiria**, Lindl. (Stercul.).

ASYNEUMA, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) 1. 385 = **Phyteuma**, Linn. (Campan.).
canescens, Griseb. & Schenk, l. c. = P. canescens.

ASYSTASIA, Blume, Bijdr. 796 (1826). *ACANTHACEAE*, Benth & Hook. f. ii. 1094.
DICENTRANTHERA, T. Anders. in Journ. Linn. Soc. vii. (1868) 52.
HENFREYEA, Lindl. Bot. Reg. (1847) t. 31.
INTRUSARIA, Rafin. Fl. Tellur. iv. 66 (1836).
MACKAYA, Harv. Thes. Cap. i. t. 13 (1859).
RAMUSIA, E. Mey. in Drège, Zwei Pfl. Docum. 215 (1843).
acuminata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 201.—Afr. trop.
africana, *Hort. Bogor. ex T. Anders. in Journ. Agri.-Hort. Soc. Ind.* N. S. i. (1858) 287.—Afr. trop.
atroviridis, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 526.—Reg. Himal.
Blumei, Nees, in DC. Prod. xi. 167 = intrusa.
Bojeriana, Nees, l. c. 166 = coromandeliana.
calycina, Benth. in Hook. Niger Fl. 478 = Strobilanthes Hehetus.
capensis, Nees, in Linnaea, xv. (1841) 356 = coromandeliana.
Charmian, *S. Moore, in Journ. Bot.* xviii. (1880) 38.— Afr. trop.
cheiranoides, *Nees, in Wall. Pl. As. Rar.* iii. 89.— Ind. or.
chinensis, *S. Moore, in Journ. Bot.* xiii. (1875) 228.— China.
comorensis, Boj. ex Nees, in DC. Prod. xi. 166 = coromandeliana.
coromandeliana, *Nees, in Wall. Pl As. Rar.* iii. 89.— Ind. or.; Malaya; Afr. trop. et austr.
coromandeliana, Wight, Ic. t. 1906 = crispata.
crispata, *Benth. in Linnaea*, xxiv. (1851) 647.— Ind. or.
denticulata, Nees, in Wall. Pl. As. Rar. iii. 89 = Strobilanthes denticulatus.
floribunda, *Klotzsch, in Peters, Reise Mossamb. Bot.* 200.—Afr. trop.
formosa, Hort. Calc. ex T. Anders. in Journ. Agri.-Hort. Soc. Ind. N. S. i. (1868) 270 = Ruellia formosa.
gangetica, T. Anders. in Thw. Enum. Pl. Zeyl. 235 = coromandeliana.
intrusa, *Blume, Bijdr.* 796.—Malaya.
intrusa, Nees, in DC. Prod. xi. 166 = coromandeliana.
Kunthiana, *Nees, in Wall. Pl. As. Rar.* iii. 89.— Penang.
lanceolata, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 524.—Burma.
Lawiana, *Dals. in Hook. Kew Journ.* iv. (1852) 344.— Ind. or.
macrocarpa, *Nees, in Wall. Pl. As. Rar.* iii. 89.—Reg. Himal.
multiflora, *Klotzsch, in Peters, Reise Mossamb. Bot.* 208.—Afr. trop.
myurensis, T. Anders. in Journ. Linn. Soc. ix. (1867) 524 = Lawiana.
Neesiana, *Nees, in Wall. Pl. As. Rar.* iii. 89.— Ind. or.; Burma.
nemorum, Nees, l. c. 90 = intrusa.
Panichanga, Nees, l. c. = Strobilanthes Panichanga.

ASYSTASIA :—
Parishii, T. Anders. in Journ. Linn. Soc. ix. (1867) 526 = Eranthemum Parishii.
paucifora, Nees, in Wall. Pl. As. Rar. iii. 90 = Codonacanthus pauciflorus.
plumbaginea, Nees, l. c. 89 = coromandeliana.
podostachys, *Klotzsch, in Peters, Reise Mossamb. Bot.* 199.—Afr. trop.
pubescens, *Klotzsch, l. c.* 202.—Afr. trop.
quaterna, *Nees, in DC. Prod.* xi. 166.—Afr. trop.
quercifolia, *Klotzsch, in Peters, Reise Mossamb. Bot.* 204.—Afr. trop.
rostrata, Solms, in Schweinf. Fl. Aethiop. 242 = Schimperi.
scabrida, *Klotzsch, in Peters, Reise Mossamb. Bot.* 202.—Afr. trop.
scandens, *Hook. Bot. Mag.* t. 4449.—Afr. trop.
Schimperi, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 40, 53.—Abyss.
subhastata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 200.—Afr. trop.
thyrsacanthus, T. Anders. in Journ. Linn. Soc. ix. (1867) 525 = Eranthemum indicum.
travancrica, *Bedd. Ic. Pl. Ind. Or.* i. 39.—Ind. or.
violacea, *Dalz. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 494.—Ind. or.
violacea, Dalz. in Hook. Kew Journ. ii. (1850) 139 = coromandeliana.
Vogeliana, *Benth. in Hook. Niger Fl.* 479.—Afr. trop.
Welwitschii, *S. Moore, in Journ. Bot.* xviii. (1880) 308. t. 213.—Afr. trop.

ATACCA, Lem. Jard. Fleur. ii. (1852) t. 186, 187 = seq.

ATACCIA, Presl, Rel. Haenk. 149 (1828) = **Tacca**, Forst.
aspera, Kunth, Enum. Pl. v. 464 = T. integrifolia.
cristata, Kunth, l. c. 466 = T. cristata.
integrifolia, Presl, Rel. Haenk. 149 = T. integrifolia.
laevis, Kunth, Enum. Pl. v. 466 = T. laevis.
lancaefolia, Kunth, l. c. 465 = T. lancaefolia.

ATADINUS, Rafin. Sylva Tellur. 30 (1838) = **Rhamnus**, Linn.
alpinus, Rafin. l. c. = R. alpinus.

ATAENIA, Endl. Gen. Suppl. v. 5 (1850) = ATENIA, Hook. & Arn. = **Carum**, Linn. (Umbell.).

ATALANTA, Nutt. Gen. Am. ii. 78 (1818) = **Cleome**, Linn. (Cappar.).
serrulata, Nutt. l. c. = C. serrulata.

ATALANTHUS, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 311 = **Sonchus**, Linn. (Compos.).
angustifolius, Pomel, Nouv. Mat. Fl. Atl. 8 (= *Zollikoferia angustifolia*).—Afr. bor.
arboreus, Sweet, Hort. Brit. ed. II. 273 = S. arboreus.
Candollianus, Pomel, Nouv. Mat. Fl. Atl. 8 (= *Zollikoferia Candolleana*).—Afr. bor.
divaricatus, Pomel, l. c. 7 = S. divaricatus.
longiloba, Pomel, l. c. (= *Zollikoferia longiloba*).—Afr. bor.
nudicaulis, Pomel, l. c. = Launaea nudicaulis.
pinnatus, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 311 = S. leptocephalus.
quercifolius, Pomel, l. c. 8 = S. quercifolius.
resedifolius, Pomel, l. c. (= *Zollikoferia resedifolia*).—Afr. bor.
spinosus, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 311 = Launaea spinosa.

ATALANTIA, Correa, in Ann. Mus. Par. vi. 883 (1805). *RUTACEAE*, Benth. & Hook. f. i. 505.
CHILOCALYX, Turcz. in Bull. Soc. Nat. Mosc. (1863) i. 585.
HELIA, M. Roem. Syn. Hesper. 42 (1846).
LAMPETIA, M. Roem. l. c. (1846).
MALNABERGAM, Adans. Fam. ii. 344 (1763).
MEROPE, M. Roem. Syn. Hesper. 44 (1846).
RISSOA, Arn. in Nov. Act. Nat. Cur. xvii. (1836) 934.
SCLEROTYLIS, Blume, Bijdr. 133 (1825).
SEVERINIA, Tenore, Ind. Sem. Hort. Neap. (1840).
bilocularis, Wall. Cat. n. 6356 = Limonia bilocularis.
buxifolia, *Oliver, in Benth. Fl. Hongk.* 51.—China.

ATALANTIA :—
buxifolia, Oliver, in Journ. Linn. Soc. v. Suppl. ii. (1861) 36 = Limonia bilocularis.
capitellata, Lindl. in Wall. Cat. n. 6355 = racemosa.
ceriscoides, Wall. l. c. n. 6354 = monophylla.
caudata, *Hook. f. Fl. Brit. Ind.* i. 513.—Reg. Himal.
ceylanica, *Oliver, in Journ. Linn. Soc. v. Suppl.* ii. (1861) 25.—Ind. or.
citrifolia, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 69.—Burma.
floribunda, Wight, Ic. iv. t. 1611 = monophylla.
glauca, *Benth. & Hook. f. Gen.* i. 806.—Austral.
Hindsii, *Oliver, in Journ. Linn. Soc. v. Suppl.* ii. (1861) 36.—China.
longispina, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 295.—Burma.
Loureiriana, *M. Roem. Syn. Hesper.* 37.—Cochinch.
macrophylla, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 186.—Ins. Nicobar.
missionis, *Oliver, in Journ. Linn. Soc. v. Suppl.* ii. (1861) 25.—Ind. or.
monophylla, Benth. in Hook. Lond. Journ. Bot. i. (1842) 488 = Hindsii.
monophylla, *DC. Prod.* i. 535.—Ind. or.
monophylla, Hook. & Arn. Bot. Beech. Voy. 172 = buxifolia.
nitida, *Oliver, in Journ. Linn. Soc. v. Suppl.* ii. (1861) 25.—Malaya.
parvifolia, M. Roem. Syn. Hesper. 37 = racemosa.
platystigma, Wight, Illustr. Ind. Bot. i. 108 = monophylla.
puberula, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 211.—Siam.
racemosa, *Wight & Arn. Prod.* 91; *et in Hook. Journ. Bot.* i. (1834) 64.—Ind. or.
recurva, *Benth. Fl. Austral.* i. 370.—Austral.
Roxburghiana, *Hook. f. Fl. Brit. Ind.* i. 513.—Malaya *Roxburgh*., Oliver, in Journ. Linn. Soc. v. Suppl. ii. 25 = caudata.
trimera, *Oliver, l. c.*—Malaya.
umbellata, M. Roem. Syn. Hesper. 37 = monophylla.

ATALAYA, Blume, Rumphia, iii. 186 (1847). *SAPINDACEAE*, Benth. & Hook. f. i. 401.
annularis, *Blume, Rumphia*, iii. 186.—Malaya.
australis, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 337 = Sapindus australis.
bijuga, Span. Ic. n. 19; in Linnaea, xv. (1841) 180 = Cupania salicifolia.
cochinchinensis, *Blume, Rumphia*, iii. 186.—Cochinch.
coriacea, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 326.—Ins. Dom. Howe.
hemiglauca, *F. Muell. ex Benth. Fl. Austral.* i. 468.—Austral.
multiflora, *Benth. l. c.*—Austral.
salicifolia, *Blume, Rumphia*, iii. 186.—Timor.; Austral.
variifolia, *F. Muell. ex Benth. Fl. Austral.* i. 468.—Linn.

ATAMASCO, Rafin. Neogenyt. 3 (1825) = **Amaryllis**, Linn.

ATAMISQUEA, Miers, Trav. Chili. ii. 599 (1826). *CAPPARIDACEAE*, Benth. & Hook. f. i. 109.
emarginata, *Miers, l. c.*—Chili.

ATAMOSCO, Adans. Fam. ii. 57 = **Amaryllis**, Linn.

ATAMOSKO, Adans. Fam. ii. 523 = prec.

ATASITES, Neck. Elem. i. 7 (1790) = **Gerbera**, Gronov. (Compos.).

ATAXIA, R. Br. Chlor. Melv. in App. Parry, Voy. 292 (1824) = **Hierochloë**, S. G. Gmel. (Gramin.).
Ecklonis, Nees, ex Trin. in Mém. Acad. Pétersb. v. (1840) 77 = H. Ecklonii.
Hookeri, Griseb. in Goett. Nachr. (1868) 77.—Ind. or.
Hornemanni, Kunth, Gram. i. 23; R. Br. in Benn. Pl. Jav. Rar. 8. t. 2.—Java.
mexicana, Rupr. in Bull. Acad. Brux. ix. (1842) 283 = H. mexicana.
tenuis, Trin. in Mém. Acad. Pétersb. v. (1840) 78. —Afr. austr.
Tongo, Nees, ex Trin. l. c. = H. Tongo.

ATE, Lindl. Gen. et Sp. Orch. 326 (1838) = **Habenaria**, Willd. (Orchid.).
acuminata, Thw. Enum. Pl. Zeyl. 300 = H. acuminata.
virens, Lindl. Gen. et Sp. Orch. 327 = H. barbata.

ATECOSA, Rafin. Fl. Tellur. iii. 46 (1836) = **Rumex**, Linn. (Polygon.).

ATELANDRA, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 289. t. 10. *MYRSINEAE*.
laurina, *Bello, l. c.*—Porto Rico.
obtusifolia, *Bello, l. c.*—Porto Rico.

ATELANDRA, Lindl. Swan Riv. App. 40. t. 5 (1839) = **Kennedya**, R. Br. (Labiat.).
incana, Lindl. l. c. = H. incana.
polystachya, Lindl. l. c. 40 = H. incana.

ATELEA, A. Rich. Pl. Vasc. Cub. 454 (1845) = **Acalea**, Moç. & Sessé (Legumin.).

ATELANTHERA, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 129. *CRUCIFERACEAE*, Benth. & Hook. f. i. 68.
perpusilla, *Hook. f. & Thoms. l. c.* 138.—Tibet. occ.

ATELEIA, Moç. & Sessé, ex DC. Mém. Lég. 395. t. 57 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 558.
apetala, *Grisb. Cat. Pl. Cub.* 80.—Cuba.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 180.—Cuba.
Glaziovecana, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 306.—Bras.
gummifera, *D. Dietr. Syn. Pl.* iv. 1219.—Ins. S. Doming.
microcarpa, *D. Dietr. l. c.*—Am. austr.
peltaria, *D. Dietr. l. c.*—Afr. austr.
pterocarpa, *Moç. & Sessé, ex DC. Lég. Mém.* 394.—Mexic.

ATELESTE, Sond. in Linnaea, xxiii. (1850) 9, 10 = **Aberia**, Hochst. (Bixin.).
tristis, Sond. l. c. 9 = Aberia tristis.
Zeyheri, Sond. l. c. 10 = Aberia Zeyheri.

ATELIANTHUS, Nutt. ex Benth. in DC. Prod. x. 455, in nota (1846) = **Synthyris**, Benth. (Scrophul.).
veronicoides, Nutt. ex Benth l. c. = S. plantaginea.

ATENIA, Hook. & Arn. Bot. Beech. Voy. 349 (1840) = **Carum**, Linn. (Umbell.).
Gairdneri, Hook. & Arn. l. c. = C. Gairdneri.

ATERAMNUS, P. Br. Hist. Jamaic. 339 (1756); 'nom. delend.' Benth. & Hook. f. Gen. iii. 256.

ATHALMUS, Neck. Elem. i. 20 (1790) = **Pallenis**, Cass. (Compos.).

ATHAMANTA, Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 900.
KILLINGA, Adans. Fam. ii. 498 (1763).
LIBANOTIS, Hall, Enum. Stirp. Helv. ii. 450 (1742).
PETRO-CARVI, Tausch, in Flora, xvii. (1834) 856.
TRIBULA, Hill, Veg. Syst. vi. 10 (1764).
TURBITH, Tausch, in Flora, xvii. (1834) 843.
acteliifolia, Wall. Cat. n. 566 = Pimpinella achilleifolia.
Ajowan, Wall. Cat. n. 572 = Carum copticum.
alaeis, Bieb. Fl. Taur. Cauc. i. 214 = Ligusticum alaeum.
alpina, Stokes, Bot. Mat. Med. ii. 79 = cretensis.
ambigua, *Rota. Obs.* iii. 98.—Hab. ?
anethifolia, Wall. Cat. n. 569 = Carum anethifolium.
annua, Linn. Sp. Pl. 245 = cretensis.
arctica, *Nym. Syll.* 155.—Europ.
asarea, *Vis. ex Reichb. Ic. Fl. Germ.* xxi. t. 1985.—Europ.
canescens, DC. Prod. iv. 155 (certe non Ath. ep. ex Boiss. Fl. Orient. ii. 971).—Oriens.
capensis, Burm. f. Fl. Cap. Prod. 7 = Ligusticum capense.
capillacea, Lam. Encyc. i. 325 = cretensis.
carvifolia, (Wah. in) Wigg. Prim. Fl. Hols. 27 = Selinum Carvifolia.
carvifolia, *Steph. ex Roem. & Schult. Syst.* vi. 496.—Russia.
carvifolia, Willd. ex Ledeb. Fl. Ross. ii. 272 = Seseli Hippomarathrum.
carvifolia, Willd. ex Ledeb. l. c. 214 = Pseudolaena balkanensis.
cenomica, Hort. Par. ex Flesh. & May. Ind. Sem. Hort. Petrop. ii. 39 = Seseli athamanticides.

ATHAMANTA:—
Cervaria, Linn. Syst. ed. X. 956 = Peucedanum Cervaria.
cervariaefolia, *DC. Prod.* iv. 156.—Ins. Teneriff.
cervariaefolia, Schrad. ex Spreng. Pugill. ii. 56 = Seseli buchoricoides.
cervariaefolia, Viv. ex Steud. Nom. ed. II. i. 166 = Laserpitium cretica.
chilliocladia, *Boiss. & Heldr. Diagn. Ser.* II. ii. 86.—Graecia.
chinensis, Linn. Sp. Pl. 245 = Selinum Monnieri.
coarctata, Ag. ex Nym. Consp. 296 = Seseli granatense.
collina, Salisb. Prod. 168 = Seseli Libanotis.
compacta, Ledeb. Fl. Alt. i. 327 = Schultzia compacta.
condensata, Linn. Sp. Pl. 1195 = Seseli condensatum.
condensata, Pall. ex Steud. Nom. ed. II. i. 166 = Pallasiana.
cretensis, Linn. *Sp. Pl.* 245.—Europ. austr.
cretensis, Schrank, Baier. Fl. i. 548 = Matthiola.
cretensis, Ucria, Hort. Panorm. 127 = sicula.
crinita, Ledeb. Fl. Alt. i. 326 = Schultzia crinita.
crithmifolia, Juss. ex DC. Prod. iv. 145 = Seseli gummiferum.
crithmoides, Lapeyr. Hist. Abr. Pl. Pyr. 148 = Seseli Monnieri.
daucifolia, Host, Fl. Austr. i. 362 = Seseli Monnieri.
daucoides, J. F. Gmel. Syst. 477 = Seseli Libanotis.
decussata, Gilib. Fl. Lituan. ii. 18 = Peucedanum Cervaria.
dense, *Boiss. & Orph. Diagn.* Ser. II. v. 98.—Graecia.
denudata, Fisch. ex Hornem. Hort. Hafn. Suppl. 32 (= *Cenolophium Fischeri*) —Rossia.
depressa, D. Don, Prod. Fl. Nep. 184 = Cortia Lindleyi.
diffusa, Wall. Cat. n. 570 = Seseli Libanotis.
divaricata, Gilib. Fl. Lituan. ii. 17 = Peucedanum Oreoselinum.
divaricifolia, Stokes, Bot. Mat. Med. ii. 78 = prasc.
flavescens, Vis. Orto Bot. Padov. (1842) 135 = Portenschlagia ramosissima.
flexuosa, Juss. ex DC. Prod. iv. 179 = Peucedanum palustre.
Galatia, Gmel. ex DC. Prod. iv. 230 = Pleurospermum Golaka.
Golaka, Hacq. Pl. Carn. t. 5 = Pleurospermum Golaka, gigantea, *D. Don, Prod. Fl. Nep.* 184 (Sp. dubia).—Nepal.
Haynaldi, Borb. & Uechtr. in Termész. Füzetek, i. (1877) 80. t. 6, 7 = Matthioli.
hungarica, Borb. ex Nym. Consp. 294 = Matthioli.
incana, Steph. ex Willd. Sp. Pl. i. 1402 = Seseli graveolens.
lasiantha, Willd. ex Roem. & Schult. Syst. vi. 494 = Pimpinella villosa.
lasiriflora, *Eckl. & Zeyh. Enum.* 248.—Afr. austr.
latifolia, Viv. Fl. Ital. 23 = Peucedanum Cervaria.
leucospermum, Poir. Encyc. Suppl. i. 533 = Seseli leucospermum.
Libanotis, Linn. Sp. Pl. 244 = Seseli Libanotis.
lucida, *Retz. Obs.* iii. 29.—Hab.?
macedonica, *Sprengel. in Roem. & Schult. Syst.* vi. 491. —Graecia; Afr. bor.
Mathioli, DC. Fl. Fr. iv. 818 = cretensis.
Matthioli, *Wulf. in Jacq. Coll.* i. 311.—Eur. austr. or.
maxima, Choisy, in DC. Prod. iv. 155 = cervariaefolia.
media, *Nym. Syll.* 154.—Europ.
Meum, Linn. Sp. Pl. 246 = Meum athamanticum.
Meum, Pall. Reise, ii. 93 = Schultzia crinita.
monstrosa, Steph. ex Roem. & Schult. Syst. Veg. vi. 495 = Schultzia compacta.
multiflora, Sibth. & Sm. Fl. Graec. iii. 69. t. 276 = Carum multiflorum.
multiflinoides, Lam. Encyc. i. 325 = cretensis.
Oreoselinum, Huds. Fl. Angl. 115 = Seseli Libanotis.
Oreoselinum, Linn. Sp. Pl. 244 = Peucedanum Oreoselinum.
Pallasiana, *Vest. ex Steud. Nom.* ed. II. i. 166.—Silibria.
panacifolia, Spreng. Sp. Umb. 140 = Magydaris panacina, tomentosa.
pastinaca, Savi, in Ust. Ann. Bot. xxi. 7 = Peucedanum palustre.
pubescens, Rets. Obs. iii. 23 = Seseli Libanotis.
pyrenaica, Jacq. Hort. Vindob. ii. 86 = Seseli Monnieri.
ramosissima, Link & Hoffingg. Fl. Port. 468. t. 106 = Seseli tortuosum.

ATHAMANTA:—
ramosissima, Portschl. in Roem. & Schult. Syst. vi. 496 = Portenschlagia ramosissima.
Reichenbachii, *Nym. Syll.* 154.—Europ.
rigida, Hornem. Hort. Hafn. 960 = Seseli buchoricoides.
rigida, *Roth, Enum. Pl. Phan. Germ.* i. 907.—Europ.
Riviniana, Host, ex Steud. Nom. ed. II. i. 166 = Seseli Libanotis.
Roxburghiana, Wall. Cat. n. 571 = Carum Roxburghianum.
rupestris, Reichb. Fl. Germ. Excurs. 470 = Matthioli.
rupestris, Vill. Hist. Pl. Dauph. ii. 648 = cretensis.
scabra, *Burm. f. Prod. Pl. Cap.* 7.—Afr. austr.
sibirica, Linn. Sp. Pl. 244 = Seseli sibiricum.
sicula, Ucria, Hort. Panor. 127 = Kundmannia sicula.
stricta, *Ledeb. ex Steud. Nom.* ed. II. i. 166.—Sibiria.
sylvestris, [Web. in] Wigg. Prim. Fl. Hols. 26 = Angelica sylvestris.
tenuifolia, Pall. ex Roem. & Schult. Syst. vi. 495 = Seseli tenuifolium.
tenuisecta, Willd. ex Spreng. in Roem. & Schult. l. c. = Peucedanum salinum.
teres, *D. Don, Prod. Fl. Nep.* 185 (Sp. dub.).—Nepal.
tortuosa, Spreng. Syst. i. 900 (= *Deverra tortuosa*).—Aegypt.
Turbith, Brot. Phytog. Lusit. 200. t. 169 = Seseli tortuosum.
verticillata, Portschl. Enum. Pl. Dalm. 14. t. 4 = Portenschlagia ramosissima.
verticillata, Sibth. & Sm. Fl. Graec. iii. 68. t. 275 = Carum graecum.

ATHAMUS, Neck. Elem. i. 85 (1790) = **Carlina**, Linn. (Compos.).

ATHANASIA, Linn. Sp. Pl. ed. II. 1180 (1762). *COMPOSITAE*, Benth. & Hook. f. ii. 416.
Bembicodium, Kunze, in Linnaea, xvi. (1842) 816.
Holophyllum, Less. Syn. Comp. 262 (1832).
Hymenolepis, Cass. in Bull. Soc. Philom. (1817) 188; et in Dict. Sc. Nat. xxii. 315 (1821).
Metagnanthus, Endl. Gen. 488 (1838).
Nebelia, Cass. in Dict. Sc. Nat. xxxiii. 59 (1824).
Pristocarpha, E. Mey. ex DC. Prod. vi. 86 (1837).
Saintmorysia, Endl. Gen. 486 (1836).
acerosa, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
annua, Linn. Sp. Pl. ed. II. 1182 = Lonas inodora.
aspera, *Thunb. Prod. Pl. Cap.* 145.—Afr. austr.
canescens, *Thunb. l. c.* 144.—Afr. austr.
capitata, Berg. Desc. Pl. Cap. 237 = scabra.
capitata, *Linn. Pl. Rar. Afr.* n. 50; *Amoen. Acad.* vi. 97.—Afr. austr.
cinerea, Linn. f. Suppl. 361 = tomentosa.
coronopifolia, *Harv. in Harv. & Sond. Fl. Cap.* iii. 191.—Natal.
crenata, *Linn. Sp. Pl.* 1180.—Afr. austr.
crithmifolia, *Linn. l. c.* 1181.—Afr. austr.
cuneifolia, *Linn. Encyc.* i. 327.—Afr. austr.
cuneiformis, *DC. Prod.* vi. 89.—Afr. austr.
cupressiformis, Comm. ex DC. Prod. v. 314 = Lepidophyllum cupressiforme.
dentata, *Linn. Sp. Pl.* ed. II. 1181.—Afr. austr.
dentata, Linn. f. Suppl. 361 = serrata.
dimorpha, *DC. Prod.* vi. 87.—Afr. austr.
Dregeana, *Harv. in Harv. & Sond. Fl. Cap.* iii. 191.—Afr. austr.
ebracteata, *E. Mey. ex DC. Prod.* vi. 87.—Afr. austr.
eriopoda, *DC. l. c.* 89.—Afr. austr.
fasciculata, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
filiformis, *Linn. f. Suppl.* 361.—Afr. austr.
fodinifera, Salisb. Prod. 189 = Pentzia flabelliformis.
fruticosa, E. Mey. ex DC. Prod. vi. 91 = microcephala.
fruticosa, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
geniculiflora, Linn. Mant. 464 = Relhania genistaefolia.
glabra, Thunb. Prod. Pl. Cap. 144 = capitata, microphylla.
glabrescens, *DC. Prod.* vi. 88.—Afr. austr.

ATHANASIA:—
glutinosa, E. Mey. ex DC. Prod. vi. 88 = Relhania genistaefolia.
graminifolia, Walt. Fl. Carol. 200 = Marshallia angustifolia.
hastata, Walt. l. c. 201 = Melanthera hastata.
hirsuta, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
imbricata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 196.—Afr. austr.
incisa, *Harv. l. c.* 190.—Afr. austr.
indica, Roxb. Hort. Beng. 61; Fl. Ind. iii. 417 = Goniocarion glabrum.
indivisa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 190.—Afr. austr.
juncea, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
laevigata, Linn. Pl. Afr. n. 50; Amoen. Acad. vi. 51 = aspera.
lanuginosa, Cav. Ic. i. 9 = canescens.
leucoclada, *Harv. in Harv. & Sond. Fl. Cap.* iii. 191.—Afr. austr.
linifolia, *Harv. l. c.* 198.—Afr. austr.
linifolia, Burm. f. Fl. Cap. Prod. 33 = longifolia.
longifolia, *Lam. Encyc.* i. 326.—Afr. austr.
maritima, Linn. Sp. Pl. ed. II. 1182 = Diotis maritima.
microcephala, D. Dietr. Syn. Pl. iv. 1401 = linifolia.
microphylla, *DC. Prod.* vi. 87.—Afr. austr.
Mundtii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 196.—Afr. austr.
obovata, Walt. Fl. Carol. 201 = Marshallia lanceolata.
pachycephala, *DC. Prod.* vi. 88.—Afr. austr.
palmatifida, *DC. l. c.* 89.—Afr. austr.
paniculata, Walt. Fl. Carol. 201 = Actinomeris alba.
parviflora, *Ait. Hort. Kew.* iii. 167.—Afr. austr.
parviflora, *Thunb. Prod. Pl. Cap.* 145.—Afr. austr.
pauciflora, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
pectinata, *Linn. f. Suppl.* 361.—Afr. austr.
pinnata, E. Mey. ex DC. Prod. vi. 88 = incisa.
pinnata, *Linn. f. Suppl.* 361.—Afr. austr.
pubescens, *Linn. Sp. Pl.* ed. II. 1182.—Afr. austr.
pubescens, Thunb. Prod. Pl. Cap. 144 = rugulosa.
pumila, Linn. f. Suppl. 362 = Relhania pectinulata.
punctata, Berg. Desc. Pl. Cap. 238 = capitata.
punctata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 190.—Afr. austr.
punctata, Thunb. Prod. Pl. Cap. 144 = capitata.
quinquedentata, *Thunb. Prod. Pl. Cap.* 145.—Afr. austr.
recurvifolia, Salisb. Prod. 189 = dentata.
rotundifolia, *DC. Prod.* vi. 87.—Afr. austr.
rugulosa, *E. Mey. ex DC. l. c.*—Afr. austr.
salicifolia, *Dietr. ex Steud. Nom.* ed. I. 91.—Afr. austr.
scabra, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
scaposa, *DC. Prod.* vi. 88.—Afr. austr.
schizoleps, *Harv. in Harv. & Sond. Fl. Cap.* iii. 190.—Afr. austr.
serrata, *Vitm. Summa Veg.* iv. 487.—Afr. austr.
sertulifera, *DC. Prod.* vi. 87.—Afr. austr.
sessiliflora, Linn. f. Suppl. 362 = Relhania sessiliflora.
spathulata, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
squarrosa, Linn. Pl. Rar. Afr. n. 52; Amoen. Acad. vi. 98 = Relhania squarrosa.
tanacetifolia, *Roth, Catalecta*, ii. 104.—Afr. austr.
tomentosa, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
tricuspis, Poir. Encyc. Suppl. i. 533 = trifurcata.
tridentata, Salisb. Prod. 189 = trifurcata.
trifurcata, *Linn. Sp. Pl.* ed. II. 1181.—Afr. austr.
trinervia, Walt. Fl. Carol. 201 = Marshallia latifolia.
unciformis, Linn. f. Suppl. 362 = Relhania cuneata.
velutina, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
virgata, *Jacq. Hort. Schoenb.* ii. t. 148.—Afr. austr.

ATHECIA, Gaertn. Fruct. i. 141 (1788) = **Forstera**, Linn. f. ? (Scyld.).
glabra, Gaertn. l. c. = F. sedifolia ?

ATHENAEA, Adans. Fam. ii. 121 (1763) = **Sparganophorus**, Vaill. (Compos.).

ATHENAEA, Schreb. Gen. 259 (1789) = **Cassaria**, Jacq. (Samyd.).

ATHENARA, Sendt. in Mart. Fl. Bras. x. 138 (1846). *SOLANACEAE*, Benth. & Hook. f. ii. 890.

ATHENAEA :—
LARNAX, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 87.
WITHERINGIA, Miers, l. c. iii. (1849) 145.
anonacea, *Sendt. in Mart. Fl. Bras.* x. 137. t. 18.—Bras.
hirsuta, *Sendt. l. c.* 137.—Bras.
Martiana, *Sendt. l. c.* 136.—Bras.
microantha, *Sendt. l. c.* 136.—Bras.
picta, *Sendt. l. c.* 134.—Bras.
popogoea, *Sendt. l. c.* 135.—Bras.
Pohliana, *Sendt. l. c.* 136.—Bras.
Schottiana, *Sendt. l. c.* 135.—Bras.
xalapensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 452.—Mexic.

ATHENANTHIA, Kunth, in Mém. Mus. Par. ii. (1815) 71 = **Anthesantia**, Beauv. (Gramin.).

ATHEOLAENA, Reichb. Consp. 106 (1828) = AETHEOLAENA, Cass. = **Senecio**, Linn. (Compos.).

ATHERANDRA, Decne. in DC. Prod. viii. 497 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 748.
acuminata, *Decne. l. c.*—Java.
acutifolia, *Decne. l. c.*—Malaya.
cuspidata, Blume, Mus. Bot. Lugd. Bat. i. 147 = acutifolia.
pubescens, Blume, l. c. = acutifolia.
Wallichii, Benth. & Hook. f. Gen. ii. 744 = Atherolepis Wallichii.

ATHERANTHERA, Mast. in Oliver, Fl. Trop. Afr. ii. 519 (1871); Trans. Linn. Soc. xxvii. (1871) 640. *PASSIFLOREAE*.
paniculata, *Mast. in Oliver, Fl. Trop. Afr.* ii. 519.—Afr. trop.
Welwitschii, Mast. in Trans. Linn. Soc. xxvii. (1871) 640 = paniculata.

ATHERNOTUS, Dulac, Fl. Hautes-Pyr. 74 (1867) = **Calamagrostis**, Adans. (Gramin.).
epigios, Dulac, l. c. = C. epigelos.
lanceolatus, Dulac, l. c. 75 = C. lanceolatus.
pyramidalis, Dulac, l. c. = Deyeuxia sylvatica.

ATHEROCEPHALA, DC. Prod. vii. 755 (1839) = **Andersonia**, R. Br. (Epacrid.).
Drummondii, DC. l. c. = Andersonia aristata.

ATHEROLEPSIS, Hook. f. in Hook. Ic. Pl. xv. 26. t. 1438 (1883). *ASCLEPIADEAE*.
Wallichii, *Hook. f. l. c.*—Burma.

ATHEROPHORA, Willd. ex Steud. Nom. ed. II. i. 167 (1840) = **Aegopogon**, Humb. & Bompl. (Gramin.).
secunda, Willd. l. c. = Aegopogon geminiflorus.

ATHEROPOGON, Muhl. ex Willd. Sp. Pl. iv. 937 (1805) = **Boutelouа**, Lag. (Gramin.).
acuminatus, Fourn. Mex. Pl. Fn. Gram. 139; ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
affinis, Fourn. ex Benth. & Hook. f. Gen. iii. 1169 = B. distichа!
americanus, Fourn. Mex. Pl. Fn. Gram. 140; ex Hemsl. Biol. Centr. Am. Bot. iii. 561 = B. bromoides.
Antillarum, Spreng. Syst. i. 294 = Aristida Antillarum.
aphudoides, Willd. Sp. Pl. iv. 937 = B. racemosa.
aristidoides, Roem. & Schult. Syst. ii. 415 = B. aristidoides.
bromoides, Roem. & Schult. l. c. = B. bromoides.
chondrosioides, Roem. & Schult. l. c. 416 = B. ovata.
curtipendulus, Fourn. Mex. Pl. Fn. Gram. 138; ex Hemsl. Biol. Centr. Am. Bot. iii. 563 = B. racemosa.
distichus, Spreng. Syst. i. 294 = B. distichа.
domingensis, Spreng. l. c. 293 (= *Eutriana juncea*). —Ins. S. Doming.
filiformis, Fourn. Mex. Pl. Fn. Gram. 140; ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
gracilis, Spreng. Syst. i. 293 = B. oligostachya.
hirtus, Spreng. l. c. = B. hirsuta.
humilis, Spreng. l. c. = B. prostrata.
juncifolius, Spreng. l. c. 294 = B. juncifolia.
modius, Fourn. Mex. Pl. Fn. Gram. 139; ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
oligostachyus, Nutt. Gen. Am. i. 78 = B. oligostachya.

ATHEROPOGON :—
Pentarrhaphis, Spreng. Syst. i. 294 (= *Eutriana Pentarrhaphis*).—Mexic.
polymorphus, Fourn. Mex. Pl. Fn. Gram. 141; ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
procumbens, Jacq. Eclog. Gram. 19 = B. tenuis.
radicosus, Fourn. Mex. Pl. Fn. Gram. 140; ex Hemsl. Biol. Centr. Am. Bot. iii. 561 = B. bromoides.
repens, Roem. & Schult. Syst. ii. 416 = B. bromoides.
scorpioides, Spreng. Syst. i. 293 = B. scorpioides.
stolonifer, Fourn. Mex. Pl. Fn. Gram. 140; ex Hemsl. Biol. Centr. Am. Bot. iii. 565.—Mexic.
Triaena, Spreng. Syst. i. 293 = B. Triathera.
villosus, Steud. Nom. ed. II. i. 167 (= *Eutriana villosa*).—Bras.

ATHEROSPERMA, Labill. Nov. Holl. Pl. ii. 74. t. 224 (1806). *MONIMIACEAE*, Benth. & Hook. f. iii. 144.
integrifolium, A. Cunn. ex A. DC. in DC. Prod. xvi. ii. 676 = moschatum.
microanthum, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 46 = Daphnandra micrantha.
moschatum, *Labill. Nov. Holl. Pl.* ii. 74. t. 224. —Austral.
Novae-Zelandiae, *Hook. f. Handb. New Zeal. Fl.* 240. —N. Zel.
repandum, *F. Muell. Fragm.* x. 105.—Austral.
Sassafras, A. Cunn. ex A. DC. in DC. Prod. xvi. ii. 676 = Doryphora Sassafras.
sempervirens, *Baill. Adansonia*, ix. (1868) 116; *Hist. Pl.* i. 232.—Chili.

ATHEROSTEMON, Blume, Mus. Bot. Lugd. Bat. i. 196 (1849). *ASCLEPIADEAE*.
javanicum, *Blume, l. c.*—Burma; Java.

ATHERSTONEA, Pappe, Sylva Cap. ed. II. 29 (1862) = Strychnos, Linn. (Loganiac.).
decussata, Pappe, l. c. = S. Atherstonei.

ATHERURUS, Blume, Rumphia, i. 186 (1885) = **Pinella**, Tenore (Aroid.).
ternatus, Blume, l. c. = P. tuberifera.
tripartitus, Blume, l. c. 187 = P. tripartita.

ATHESIANDRA, Miers, ex Benth. & Hook. f. Gen. i. 946 (1863) = **Pityohopetalum**, Benth. (Olacin.).
ancora, Miers, ex Oliver, Fl. Trop. Afr. i. 348 = P. ancepa.

ATHLIANTHUS, Endl. Gen. Suppl. ii. 68 (1842) = Justicia, Linn. (Acanth.).
spicciflorus, Nees, Linnaea, ix. 54.—Afr. trop.

ATHRIXIA, Ker-Gawl. in Bot. Reg. t. 681 (1823). *COMPOSITAE*, Benth. & Hook. f. ii. 338.
AUSTRALIA, Lindl. Swan Riv. App. 24 (1839).
CHRYSODISCUS, Steetz, in Lehm. Pl. Preiss. i. 360 (1845).
KLENZEA, Sch. Bip. ex Steud. Nom. ed. II. i. 847 (1840); et in Walp. Rep. ii. 973 (1843), partim.
TRICHOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 2. t. 83.
angustissima, *DC. Prod.* vi. 277.—Afr. austr.
australis, *Steetz, in Lehm. Pl. Preiss.* i. 482.—Austral.
capensis, *Ker-Gawl. in Bot. Reg.* t. 681.—Afr. austr.
Chatopoda, *F. Muell. Fragm.* x. 96.—Austral.
debilis, *DC. Prod.* vi. 276.—Madag.
elata, *Sond. in Linnaea*, xxiii. (1850) 67.—Afr. austr.
ericaefolia, *DC. Prod.* vi. 277.—Arabia.
fontana, *Mac Owan, in Journ. Linn. Soc.* xvii. (1881) 391.—Afr. austr.
Gerrardi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 292.—Afr. austr.
glandulosa, *Kunth, Ind. Sem. Hort. Berol.* (1848) 15. —Afr. austr.
gracilis, *Benth. Fl. Austral.* iii. 599.—Austral.
heterophylla, *Less. Syn. Comp.* 366.—Afr. austr.
multiceps, *Benth. Fl. Austral.* iii. 599.—Austral.
phylicoides, *DC. Prod.* vi. 277.—Afr. austr.
rosmarinifolia, Oliver & Hiern, in Oliver, Fl. Trop. Afr. iii. 355.—Afr. trop.
sessilifolia, *DC. Prod.* vi. 277.—Afr. austr.
stricta, *Benth. Fl. Austral.* iii. 600.—Austral.
tenella, *Benth. l. c.*—Austral.
tomentosa, Less. Syn. Comp. 364 = Pterothrix tomentosa.

ATHRODACTYLIS, Forst. Char. Gen. 149 (1776) = **Pandanus**, Linn.
spinosa, Forst. l. c. 150 = P. odoratissimus.

ATHROISMA, DC. in Guill. Arch. Bot. ii. (1833) 516. *COMPOSITAE*, Benth. & Hook. f. ii. 396.
laciniatum, *DC. l. c.* 517.—Ind. or.
viscida, Zoll. & Mor. in Nat. en Geneesk. Arch. Nerl. Indie, ii. (1845) 282 = laciniatum.

ATHROISMA, Griff. Notul. iv. 477 (1854) = **Trigonostemon**, Blume (Euphorb.).
dentatum, Griff. l. c. 478 = T. longifolius.
serratum, Griff. l. c. 477 = T. heteranthus.

ATHRONIA, Neck. Elem. i. 32 (1790) = **Spilanthes**, Jacq. (Compos.).

ATHROOSTACHYS, Benth. ex Benth. & Hook. f. Gen. iii. 1008 (1883). *GRAMINEAE*, Benth. & Hook. f. iii. 1208.
capitata, *Benth. l. c.* 1209.—Bras.

ATHROTAXIS, D. Don, in Trans. Linn. Soc. xviii. (1839) 171. t. 18, 14. *CONIFERAE*, Benth. & Hook. f. iii. 430.
ANTHROTAXIS; Endl. Gen. Suppl. i. 1372 (1841).
alpina, Van Houtte, ex Gord. Pinet. Suppl. 31 = selaginoides.
araucarioides, Brongn. ex Parl. in DC. Prod. xvi. ii. 496 = Dacrydium araucarioides.
cupressoides, *D. Don, in Trans. Linn. Soc.* xviii. (1839) 178. t. 18.—Tasmania.
Doniana, Henk. & Hochst. Nadelh. 221 = laxifolia.
Gunneana, *Hook. ex Carr. Conif.* ed. II. 207.—Tasmania.
imbricata, Hort. ex Carr. l. c. 204 = selaginoides.
imbricata, Maule, ex Gord. Pinet. Suppl. 16 = cupressoides.
Youcadan, Hort. ex Carr. Conif. ed. II. 55 = Juniperus gracilis.
laxifolia, *Hook. Ic. Pl.* t. 573.—Tasmania.
selaginoides, *D. Don, in Trans. Linn. Soc.* xviii. (1839) 172. t. 14.—Tasmania.
tetragona, Hook. Ic. Pl. t. 580 = Microcachrys tetragona.

ATHRUPHYLLUM, Lour. Fl. Cochinch. 120 (1790) = **Myrsine**, Linn.
linearе, Lour. l. c. = M. Athruphyllum.

ATHYMALUS, Neck. Elem. ii. 358 (1790) = **Euphorbia**, Linn.

ATHYROCARPUS, Schlecht. in Linnaea, xxvi. (1853–55) 454. *COMMELINACEAE*, Benth. & Hook. f. iii. 847.
PHAEOPHARRION, Hassk. in Flora, (1866) 212.
leiocarpus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 486.—*Mexic.*
pallidus, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 151 = Commelina pallida.
pallidus, *Schlecht. in Linnaea*, xxvi. (1853–55) 454.—Am. trop.
persicifolius, *Hemsl. Biol. Centr. Am. Bot.* iii. 486.—Panama.

ATHYRUS, Neck. Elem. iii. 89 (1790) = **Lathyrus**, Linn. (Legumin.).

ATHYSANUS, Greene, in Bull. Calif. Acad. i. (1885) 72. *CRUCIFERAE*.
pusillus, *Greene, l. c.*—Calif.

ATIMETA, Schott, Gen. Aroid. t. 71 (1858) = **Rodospatha**, Poepp. (Aroid.).
Blumenavea, Reichb. ex Peyr. Aroid. Maxim. 98. t. 19, 20.—Bras.
Martii, Schott, Prod. Aroid. 249 = R. oblongata.
Vidensiana, Schott, Gen. Aroid. t. 71 = R. oblongata.

ATIRSITA, Rafin. New Fl. Am. iv. 86 (1836) = **Bryantgium**, Towns. (Umbell.).
pumila, Rafin. l. c. 87 = R. gracile.

ATITARA, Mercur. ex Juss. in Dict. Sc. Nat. iii. 377 (1816) = **Brodiaea**, Forst. (Retac.).

ATKINSONIA, F. Muell. Fragm. v. 84 (1865) = **Loxanthus**, Linn.
ligustrina, *F. Muell. l. c.* v. 84; vi. 253 = L. spicatus.

ATLANTIA, Kurs, For. fl. Brit. Burma, i. 194 (1877) = **Atalantia**, Correa (Rutac.).

ATOCION, Adans. Fam. ii. 254 (1763) = **Silene**, Linn. (Caryophyll.).
Armeria, Pourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 844 = S. Armeria.

ATOLARIA, Neck. Elem. iii. 81 (1790) = **Crotalaria**, Linn. (Legum.).

ATOMOSTYLIS, Steud. in Flora, xxxiii. (1850) 229; Syn. Pl. Cyp. 215 (1855) = **Cyperus**, Linn.
cyperiformis, Steud. Syn. Pl. Cyp. 215 = C. nudicaulis.
flavescens, Steud. in Flora, xxxiii. (18..0) 229. = C. nudicaulis.

ATOSSA, Alef. in Bonplandia, ix. (1861) 100 = **Wiola**, Linn. (Legum.).
Chusi, Alef. l. c. 101 = V. oroboides.
sepium, Alef. l. c. = V. sepium.

ATRACTYLIA, Reichb. Nom. ‘n. 8416’ (1841) = seq.

ATRACTYLIS, Linn. Gen. ed. I. 246 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 405.
ACARNA, All. Fl. Pedem. i. 183 (1785).
ANACTIS, Cass. in Dict. Sc. Nat. xlvii. 510 (1827).
ATRACTYLODES, DC. Prod. vii. 48 (1838).
CHAMAELEON, Cass. in Dict. Sc. Nat. xlvii. 509 (1827).
CIRSELLIUM, Gaertn. Fruct. ii. 454. t. 168 (1791).
CROCODYLUS, ‘Adans.’ ex Steud. Nom. ed. II. i. 167, in syn. (1840).
SPADACTIS, Cass. in Dict. Sc. Nat. xlvii. 510 (1827).
acaulis, Pers. Syn. ii. 382, in syn. = gummifera.
amplexicaulis, Poir. Encyc. vi. 30 (= Stobaeae sp.).— Afr. austr.
angustifolia, Houtt. Plantenk. x. 518 = Berkheya lanceolata.
armena, Poir. Encyc. Suppl. iv. 637 (= *Onobroma armenum*).—Armen.; Persia.
carpitose, *Desf. Fl. Atlant.* ii. 254. t. 225.—Afr. bor. occ.
carpitosa, Viv. Fl. Lib. Spec. 52 = cancellata.
cancellata, *Linn. Sp. Pl.* 880.—Reg. Mediterr.; Orient.
canescens, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 519 = cancellata.
chinensis, DC. Prod. vi. 549 = ovata.
ciliaris, *Linn. Syst.* ed. X. 1202.—‘ Aethiopia.’
citrina, Coss. & Kralik, in Bull. Soc. Bot. Fr. iv. (1857) 471 = flava.
comosa, *Sieber, ex Cass. in Dict. Sc. Nat.* i. 58.—Syria.
cuneata, *Boiss. Fl. Orient.* iii. 454.—Afghanist.
diffusa, *Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 396, nomen.—Algeria.
echinata, *Pomel, Nouv. Mat. Fl. Atl.* 274.—Algeria.
flava, Coss. & Dur.—Cf. Bull. Soc. Bot. Fr. iv. (1857) 471 = citrina.
flava, *Desf. Fl. Atlant.* ii. 254. t. 222.—Afr. bor.; Arab.; Beluchist.
fruticosa, Linn. Sp. Pl. 839 = Berkheya fruticosa.
Fusus-agrestis, Gaertn. Fruct. ii. 381. t. 161 = Carthamus lanatus.
gummifera, *Linn. Sp. Pl.* 829.—Reg. Mediterr.
gummifera, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 519 = Carlina racemosa.
humilis, *Linn. Sp. Pl.* 829.—Europ. austr. occ.; Marocco.
lanata, Scop. Fl. Carn. ed. II. ii. 184 = Carthamus lanatus.
lanceos, Thunb. Fl. Jap. 306 = ovata.
leucocephala, Moench, Meth. 558 = Carthamus lanatus.
leucophaea, Gaertn. Fruct. ii. 381 = Carthamus leucocaulis.
macrocephala, *Desf. Fl. Atlant.* ii. 253.—Afr. bor. occ. macrophylla, *Desf. l. c.* 255.—Algeria.
mexicana, Linn. f. Suppl. 350 = Lycoseris mexicana.
microcephala, Coss. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 240 = serratuloides.
oppositifolia, Linn. Mant. ii. 477 = Berkheya grandiflora.
ovata, *Thunb. Fl. Jap.* 306.—Japon.
persica, Boiss. Diagn. Ser. I. v. 110 = Thevenotia persica.
phaeolepis, *Pomel, Nouv. Mat. Fl. Atl.* 273.—Algeria.
pilosa, Moench, Meth. 558 = Carthamus lanatus.

ATRACTYLIS:—
polycephala, *Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 461, nomen.—Algeria.
Pruniniana, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 250.—Ins. Canar.
prolifera, *Boiss. Diagn.* Ser. I. x. 96.—Palaestina.
purpurea, Linn. ex Sm. Ic. fasc. iii. 65 = seq.
purpurea, Linn. f. Suppl. 349 = Onoseris purpurea.
radiciflora, *DC. Prod.* vi. 551.—Gall. austr.
rigida, Poir. Encyc. Suppl. iv. 637 = Carthamus rigidus.
scabra, *Boiss. Diagn.* Ser. I. vi. 109.—Persia.
serrata, *Pomel, Nouv. Mat. Fl. Atl.* 20.—Afr. bor.
serratuloides, *Sieber, ex Cass. in Dict. Sc. Nat.* i. 58.—Algeria; Palaest.
speciosa, DC. in Deless. Ic. Sel. iv. t. 71 = comosa
triplinervia, Vahl, ex Steud. Nom. ed II. i. 168; ii. 83 = Lycoseris triplinervis.

ATRACTYLODES, DC. Prod. vii. 48 (1838) = **Atractylis**, Linn. (Compos.).
lanceos, DC. l. c. = Atractylis ovata.
lyrata, Sieb. & Zucc. in Abh Akad. Muench. iv. iii. (1846) 198 = Atractylis ovata.
ovata, DC. Prod. vii 48 = Atractylis ovata.

ATRAGENA, Linn. Syst. ed. I. (1735) = seq.

ATRAGENE, Linn. Diss. Dasc. 21 (1747); Amoen. Acad i. 182 = **Clematis**, Linn. (Ranunc.).
alpina, Linn. Sp. Pl. 542; Rev. Hortic. (1855) 323. t. 17. t. = C. alpina.
alpina, Pall. Fl. Ross. ii. 69 = C. verticillaris.
americana, Sims, Bot Mag. t 837 = C. verticillaris.
austriaca, Scop. Fl. Carn. ed. II. i. 367 = C. alpina.
austriaca, Host, in Linnaea, xi. (1837) 548, in obs = C. Wenderothii
balearica, Pers. Syn. ii. 98 = C. cirrhosa.
capensis, Burm. f. Fl. Cap. Prod. 16, pro parte = Anemone tenuifolia.
capensis, Linn. Sp. Pl. 543 = Anemone capensis.
ceylanica, Steud. Nom. ed. II. i. 168, vide *seylanica*.
cirrhosa, Pers. Syn. ii. 98 = C. cirrhosa.
Clematides, Crantz, Stirp. Austr. ed. I. fasc. ii. 111; ed. II. 127 = C. alpina.
columbiana, Nutt. in Journ. Acad. Phil. vii. (1834) 7 = C. verticillaris.
cordata, Wender. ex Steud. Nom. ed. II. i. 168 = C. Wenderothii.
florida, Pers. Syn. ii. 98 = C. florida.
indica, Desf. Tabl. ed. I. 193 = C. florida.
indica, Heyne, ex Steud. Nom. ed. II. i. 168, 379 = C. Gouriana.
japonica, Thunb. Fl. Jap. 238 = C. polypetala.
japonica, Wight, ex Steud. Nom. ed II. i. 168 = C. Gouriana.
lobata, Llanos, Fragm. 78 = Naravelia zeylanica.
macropetala, Ledeb. Fl. Alt. ii. 276 = C. macropetala.
occidentalis, Hornem. Hort. Hafn. ii. 820 = C. occidentalis.
ochotensis, Pall. Fl. Ross. i. 11. 69 = C. ochotensis.
patens, Decne. in Belg. Hortic. (1858) 16 = C. patens.
platysepala, Trautv. & C. A. Mey. in Middend. Reise, (Fl. Ochot. 5).—As. bor. occ.
polygama, Jacq. Am. Select. 171 = G. dominica.
polypetala, Thunb. Fl. Jap. 239 = Anemone japonica.
sibirica, Linn. Sp. ed. I. 543 = alpina.
speciosa, *Weinm. in Bull. Soc. Nat. Mosc.* (1850) i. 535.—Rossia.
tenuifolia, Linn. f. Suppl. 270 = Anemone tenuifolia.
tenuis, Thunb. Fl. Jap. 239 = Anemone tenuifolia.
trifernata, DC. Syst. i. 150 = C. tetternata.
Wenderothii, Schlecht. in Linnaea, xi. (1837) 650 = C. Wenderothii.
zeylanica, Blanco, Fl. Filip. ed. I. 461 = Naravelia zeylanica.
zeylanica, Linn. Sp. Pl. 543 = Naravelia zeylanica.

ATRAPHAXIS, Linn. Syst. ed. I (1735). *POLYGONACEAE*, Benth. & Hook. f. iii. 96.
PEDALIUM, Adans. Fam. ii. 277 (1763).
PERSICARIA, Neck. Elem. ii. 210 (1790).
TRAGOPYRUM, Bieb. Fl. Taur. Casc. iii. 284 (1819).
aghanica, *Meisn. in DC. Prod.* xiv. 76.—Afghanist.
angustifolia, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15, t. 115.—Armenia.
aphylla, *Pall. Reise*, i. 396.—Rossia.

ATRAPHAXIS:—
Aucheri, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15, 20. t. 114.—Persia.
Billardieri, *Jaub. & Spach, l. c.* 14, 17. t. 111.—Graecia; As. Min.; Syria.
buxifolia, *Jaub. & Spach, l. c.* 14.—Reg. Caucas.
Calcarii, Boiss. Diagn. Ser. II. iv. 76 = spinosa.
candida, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 1021.—Persia.
canescens, *Bunge, Ind. Sem. Hort. Dorpat.* (1839) 8. —As. bor.
compacta, *Ledeb. Fl. Alt.* ii. 55.—Turkestan.
crassifolia, Agardh, in Schult. f. Syst. vii. 1383 = spinosa.
cuneifolia, *Poir. Encyc. Suppl.* i. 585.—Am. bor.
dawurica, Jaub. & Spach, Illustr. Pl. Or. ii. 15 = Laxmanni.
decipiens, Jaub. & Spach, l. c. = Laxmanni.
dennaflora, C. Koch, in Linnaea, xxi. (1849) 212 = spinosa.
disaca, Bosc, ex Meisn. in DC. Prod. xiv. 80 = Polygonella parvifolia.
Fischeri, Jaub. & Spach, Illustr. Pl. Or. ii. 12 = spinosa.
frutescens, C Koch, Dendrol. ii. 1. 860 = Laxmanni.
graeca, Meisn. in DC. Prod. xiv. 77 = billardieri.
grandiflora, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15, 19 t. 118.—Armenia; Cappadoc.
inermis, Mill Gard. Dict. ed. VIII. n. 2 = undulata.
Karelini, Jaub. & Spach, Illustr. Pl. Or. ii. 12 = spinosa.
lactevirens, *Jaub & Spach, l. c.* 14.—Soongaria.
lancolata, *Bunge, in Mém. San. Etr. Pétersb.* vii (1851) 488 (*Lehm. Rel.* 807).—As. temp.
lanceolata, Meisn. in DC. Prod. xiv. 78 = Laxmanni.
Laxmanni, *Agardh, in Schult. f. Syst.* vii. 1382.—Sibir.
micrantha, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15.—Sibiria.
pungens, *Jaub. & Spach, l. c.* 14.—Soongaria.
pyrifolia, *Bunge, in Mém. San. Etr. Pétersb.* vii. (1851) 488 (*Lehm. Rel.* 807).—Turkestan.
replicata, Lam Encyc. i. 329 = spinosa.
sinaica, Fresen. in Mus. Senkenberg. i. 74; Jaub. & Spach, Illustr. Pl. Or. ii. 12 = spinosa.
spinosa, Eichw. Casp. Cauc. 7 = compacta.
spinosa, Laxm. ex Agardh, in Schult. f. Syst. vii. 1382 = crassifolia, Laxmanni.
spinosa, *Linn. Sp. Pl.* 475.—Orient.
suaedaefolia, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15.—Persia.
Tournefortii, *Jaub. & Spach, l. c.* 14, 18. t. 112.—Armenia.
undulata, Linn. Sp. Pl. 333 = Polygonum atraphaxoides.
undulata, *Willd. Sp. Pl.* ii. 249.—Afr. austr.
varnabitis, Jaub. & Spach, Illustr. Pl. Or. ii. 11, 13, 16. t. 110 = Billardieri.

ATRATEGIA, Bedd. ex Hook. f. Fl. Brit. Ind. i. 74 (1875) = **Gonothalamus**, Blume (Anonac.).
Thomsoni, Bedd. ex Hook. f. l. c. 75 = G. cardiopetalus.

ATREMA, DC. Coll. Mém. v. 71. t. 18 (1829); et Prod. iv. 259 (1830) = Bifora, Hoffm. (Umbellif.).
americanum, DC. Coll. Mém. v. 71. t. 18 = B. americana.
testiculatum, Miq. Fl. Ind. Bat. i. 1. 744 = B. testiculata.

ATRICHOSERIS, A. Gray, Syn. Fl. N. Am. i. ii. 410 (1884). *COMPOSITAE*.
platyphylla, *A. Gray, l. c.*—Am. bor. occ.

ATRIPLEX, [Tourn.] Linn. Syst. ed. I (1735); Gen. 317 (1737). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 58.
ENDOLEPIS, Torr. ex A. Gray, in Pacif. Rail. Rep. xii. 47. t. 8 (1860).
HAEIMUS, Wallr. Sched Crit. 117 (1822).
OBIONE, Gaertn. Fruct. ii. 198. t. 126 (1791).
PHYLLOTHECA, Nutt. ex Moq. in DC. Prod. xiii. ii. 98 (1849).
PTEROCARYA, Nutt. ex Moq. l. c. 106 (1849).
PTEROCHITON, Torr. in Frém. Rep. 818 (1846).
SCHIZOTHECA, Lindl. Veg. Kingd. 813 (1847).
THELEOPHYTON, Moq. in DC. Prod. xiii. ii. 115 (1849).

ATRIPLEX :—

abbreviata, Opiz, in Flora, vii. (1824) i. Beil. 81 = laciniata.
acanthocarpa, *S. Wats. in Proc. Am. Acad.* ix. (1874) 117.—Texas.
acuminata, Bieb. Fl. Taur. Cauc. iii. 641 = hortensis.
acuminata, Waldst. & Kit. Pl. Rar. Hung. ii. 107. t. 108 = nitens.
agrestis, Schur, Enum. Pl. Transs. 575 = patula.
alaskensis, *S. Wats. in Proc. Am. Acad.* ix. (1874) 108.—Am. bor. occ.
alba, Cranz, Inst. i. 206 = Chenopodium album.
alba, Salisb. Prod. 162 = Exomis striplicoides.
alba, Scop. Delic. Insub. ii. 16. t. 8 = roseа.
albicans, [Soland. in] Ait Hort. Kew. ed. I. iii. 430 = Exomis striplicoides.
albicans, Bess. Prim. Fl. Galic. i. 194 = roseа.
albicans, Willd. Sp. Pl. iv. 962 = laciniata.
Aldamar, *Grisb. Cat. Fl. Cub.* 292.—Cuba.
alexandrina, *Boiss Fl. Orient.* iv. 914.—Aegypt.
amaranticides, Gmel. ex Moq. in DC. Prod. xiii. ii. 116 = Axyris amarranthoides.
amblyostegia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 416.—As. centr.
ambrosioides, Cranz, Inst. i. 207 = Chenopodium ambrosioides.
angulata, *Benth. Fl. Austral.* v. 174.—Aus .l.
angustifolia, Sm. Fl. Brit. 1092 = patula.
anthelmintica, Cranz, Inst. i. 207 = Chenopodium ambrosoides.
arabicum, Ehrenb. ex Boiss. Fl. Orient. iv. 917 = farinosa.
arenaria, *Nutt. Gen. Am.* i. 198.—Am. bor.
arenaria, Tineo, Cat. Hort. Panorm. 276; Woods, in Bab. Man. ed. III. 271 = laciniata.
argentea, *Aschr. Am. Am.* i. 198; *ex S. Wats. Rev. Chenop.* 117.—Am. bor. occ.
argentea, Pall. ex Steud. Nom. ed. II. i. 168 = nitens.
argentea, Schrad. ex Moq. in DC. Prod. xiii. ii. 92 = roseа.
aristata, Cranz, Inst. i. 208 = Chenopodium aristatum.
Asoi, *Dufour, in Bull. Soc. Bot. Fr.* vii. (1860) 482. —Hispan.
astracanica, Balb. Hort. Taric. (1813) = laciniata.
astracanica, *Lag. Gen. et Sp. Nov.* (1816) 9.—As. temp.
astrachanica, Ledeb. Fl. Ross. iii. 719 = laciniata.
atacamensis, *Phil. Fl. Atac.* 48.—Chili.
Aucheri, Moq. Chenop. Enum. 51 = nitens.
australasica, Moq. l. c. 59 = patula.
axillaris, Tenore, Fl. Nap. Prod. i. 54 = roseа.
Babingtonii, Woods, in Bab. Man. ed. III. 270 = patula.
Barclayana, *D. Dietr. Syn. Pl.* v. 587.—Am. bor. occ.
Belangeri, *D. Dietr. l. c.* 585.—Persia; Afghanist.
bengbalensis, Lam. Encyc. i. 376 = hortensis.
Berlandieri, Moq. Chenop. Enum. 68 = canescens.
Besseriana, Roem. & Schult. Syst. vi. 284 = roseа.
Billardieri, *Hook. f. Fl. N. Zel.* i. 215.—Austral.; N. Zel.
Bocconi, Guss. Fl. Sic. Syn. ii. 588 = laciniata.
Bonus-Henricus, Cranz, Inst. i. 206 = Chenopodium Bonus-Henricus.
Botrys, Cranz, l. c. 207 = Chenopodium Botrys.
bracteosa, Trautv. in Act. Hort. Petrop. i. n. 1 (1870) 17 = dimorphostegia.
bracteosa, *S Wats. in Proc. Am. Acad.* ix. (1874) 115. —Calif.
Brewerі, *S. Wats. l. c.* 119.—Calif.
Bunburyana, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 274.—Austral.
californica, *Moq. in DC. Prod.* xiii. ii. 98.—Calif.
calotheca, *Fries, Novit. Fl. Suec. Mant.* iii. 164.— Suecia.
campanulata, *Benth. Fl. Austral.* v. 177.—Austral.
campestris, Koch & Zis, Cat. Pl. Palat. 24 = tatarica.
cana, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 306; *Moq. Chenop. Enum.* 61.—Sibiria.
candicans, *Link, ex Steud. Nom.* ed. II. i. 168.— Lusitan.
canescens, *Jones, Cat.* 178.—Am. bor. occ.
canescens, Nutt. Gen. Am. i. 197 = Nuttalii.
capensis, *Moq. Chenop. Enum.* 63.—Afr. austr.

ATRIPLEX :—

Chasicovii, *Bunge, ex Boiss. Fl. Orient.* iv. 913.— Persia.
chilensis, *Colla, in Mem. Adc. Torin.* xxxix. (1836) 7. t. 49.—Chili.
chrysellinus, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 279 = Billardieri.
cinerea, *Poir. Encyc. Suppl.* i. 471.—Austral.; N. Zel.
confertifolia, *S. Wats. in Proc. Am. Acad.* ix. (1874) 119.—Am. bor. occ.
coriacea, *Forsk. Fl. Aegypt. Arab.* 175.—Aegypt.
coriacea, Raddi, ex Moq. in DC. Prod. xiii. ii. 108 = glauca.
coronata, *S. Wats. in Proc. Am. Acad.* ix. (1874) 114. —Calif.
Coulteri, *D. Dietr. Syn. Pl.* v. 587.—Calif.
crassifolia, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 309.— Turkestan.; Ind. or.; Sibir.
crispa, *D. Dietr. Syn. Pl.* v. 586.—Persia.
cristata, Heyn, ex Moq. in DC. Prod. xiii. ii. 99; Koch. ex Moq. l. c. 109 = repens.
cristata, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 959.— N. Granat.
crystallina, Moq. l. c. 108, 116 = Billardieri.
crystallina, *Ehrenb. ex Jouss. Fl. Orient.* iv. 915.— Aegypt.
decumbens, Roem. & Schult. Syst. vi. 289 = prostrata.
decumbens, *S. Wats. in Proc. Am. Acad.* xii. (1877) 275.—Calif.
deltoidea, Bab. Man. ed. I. 253 = hastata.
denticulata, Moq. in DC. Prod. xiii. ii. 97 = semibaccata.
deserticola, *Phil. Fl. Atac.* 47.—Chili.
diffusa, Tenore, Prod. Fl. Nap. i. 57 = laciniata.
dimorphostegia, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 488.—Songar.
dioica, Rafin. in Am. Monthly Mag. (1818) 176 = Purshiana.
Drummondii, *Moq. in DC. Prod.* xiii. ii. 102.— Drummondii.
elachophylla, *F. Muell. Fragm.* vii. 8.—Austral.
elaeagnoides, Moq. Chenop. Enum. 66 = cinerea.
elongata, *D. Dietr. Syn. Pl.* v. 587.—Mexic.
elongata, Guss. Fl. Sic. Syn. ii. 592 = hastata.
Endolepis, *S. Wats. in Proc. Am. Acad.* ix. (1874) 110. —Am. bor.
erecta, Hook. Fl. Angl. ed. I. 376 = patula.
exilifolia, *F. Muell. Fragm.* vii. 9.—Austral.
expansa, *S. Wats. in Proc. Am. Acad.* ix. (1874) 116. —Am. bor. occ.
farinosa, Dum. Fl. Belg. 90 = laciniata.
farinosa, Moq. Chenop. Enum. 55 = Verreauxii.
farinosa, *Forsk. Fl. Aegypt. Arab.* 198.—Aegypt.; Arab.
fasciculata, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 377.—Am. bor.
fastuvalvis, *F. Muell. Fragm.* ix. 193.—Austral.
Flabellum, *Bunge, ex Boiss. Fl. Orient.* iv. 912.— Persia; Turkestan.
flavescens, Dum. Fl. Belg. 90 (Quid ?).—Belgium.
flexuosa, *Moq. Chenop. Enum.* 57.—Afr. austr.
foliolosa, Link, in Schrad. Journ. (1800) i. 54 = roseа.
foliolosa, *Phil. in Bot. Zeit.* xxviii. (1870) 500; *Anal. Univ. Chil.* (1875) 193.—Chili.
foliosa, Guss. ex Nym. Consp. 623 = roseа.
fruticosa, Nutt. ex Moq. in DC. Prod. xiii. ii. 112 = canescens.
Gardneri, D. Dietr. Syn. Fl. v. 587 = Gordoni.
gigantea, Poir. ex Moq. in DC. Prod. xiii. ii. 94 = hastata.
glauca, Brot. Fl. Lusit. i. 479 = candicans.
glauca, Cranta, Inst. i. 207 = Chenopodium glaucum.
glauca, Pall. ex Roem. & Schult. Syst. v. 277 = terrestricum.
glauca, Pall. Reise, i. 494 = cana.
glauca, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 645.—Lusitania.
glauca, *Linn. Cent.* i. 38.—Europ.
glomulifera, *Nees, in Lehm. Pl. Preiss.* i. 634.— Austral.
Gmelini, *C. A. Mey. in Bong. Veg. Sitch.* 42.—As. bor. or.
Gordoni, *Hook. Kew Journ.* v. (1853) 261.—Am. bor.
gracilis, Nutt. ex Moq. in DC. Prod. xiii. ii. 95 = hastata.
graeca, Sibth. & Sm. Fl. Graec. x. t. 963 = tatarica.
graeca, Willd. Sp. Pl. iv. 956 = hastata.

ATRIPLEX :—

Graeci, Tineo, Cat. Hort. Panorm. 277 = roseа.
Greggii, *S. Wats. in Proc. Am. Acad.* ix. (1874) 118. —Mexic.
Griffithii, *Moq. in DC. Prod.* xiii. ii. 102.—Persia; Afghanist.
halimoides, *Lindl. in Mitch. Three Exp.* i. 285.— Austral.
halimoides, Rafin. in Am. Monthly Mag. (1818) 176 = hastata.
halimoides, *Tineo, Cat. Hort. Panorm.* 277.—Hab.? *Halimus*, R. Br. Prod. 406 = cinerea.
Halimus, *Linn. Sp. Pl.* 1052.—Afr. bor. et austr.
Halimus, Nees, in Lehm. Fl. Preiss. i. 633 = cinerea, instides.
Halimus, Pall. Reise, ii. 477 = cana.
Halimus, Kitter, ex Moq. in DC. Prod. xiii. ii. 110 (= *Obione crispa*).—Ins. S. Doming.
hastaefolia, Salisb. Prod. 158 = hastata.
hastata, Geners. ex Steud. Nom. ed. II. i. 169 = laciniata.
hastata, Koch, Syn. Fl. Germ. ed. II. 702 = calotheca.
hastata, *Linn. Sp. Pl.* 1053.—Europ.; As. bor.
hastilifolia, Reichb. ex Steud. Nom. ed. II. i. 169 = patula.
Hermanniі, Willem. Phytogr. iii. 1992 = nitens.
heteranthа, Wight, Ic. v. t. 1787 = hortensis.
heterophylla, Nutt. ex Moq. in DC. Prod. xiii. ii. 112 = canescens.
heteroperma, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 448 (*Lehm. Rel.* 272).—As. centr.
hispanica, *S. G. Gmel. It.* iv. 40.—Rosa. austr.
holocarpa, *F. Muell. Fragm. Balb. Exp.* 19.—Austral.
hortensis, *Linn. Sp. Pl.* 1053.—Cosmopol. (cult.).
humilis, *F. Muell. Fragm.* iv. 48.—Austral.
hybrida, Cranta, Inst. i. 207 = Chenopodium hybridum!
hymenotheca, *S. Wats. in Proc. Am. Acad.* ix. (1874) 119.—Am. bor. occ.
hymenotheca, *Moq. in DC. Prod.* xiii. ii. 101.— Austral.
hypoleuca, Nees, in Lehm. Pl. Preiss. i. 633 = cinerea.
imbricata, *D. Dietr. Syn. Pl.* v. 586.—Bolivia.
incana, *F. G. Dietr. Vollst. Gaertn. Lexik. Nachtr.* i. 418.—Hispan.
incisa, Bieb. Fl. Taur. Cauc. iii. 641 = tatarica.
incisa, Hoben. ex Ledeb. Fl. Ross. iii. 720 = laciniata.
increasata, *F. Muell. Rep. Babb. Exp.* 30.—Austral.
inflata, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 75 = halimoides.
integrifolia, Krock. ex Steud. Nom. ed. II. i. 169 = triangularis.
instides, *Moq. Chenop. Enum.* 63.—Austral.
julacea, *S. Wats. in Proc. Am. Acad.* xx. (1885) 370.— Calif.
Katef, Ehrenb. ex Boiss. Fl. Orient. iv. 916 = A. Halimus.
Kentiji, Wall. Cat. n. 6951 = repens.
Kunthiana, *D. Dietr. Syn. Pl.* v. 586.—Am. austr.
lacera, Desf. Cat. Hort. Par. ed. III. 359 = calotheca.
laciniata, Linn. Cent. Pl. Pauab 126 = crassifolia.
laciniata, Bieb. Fl. Taur. Cauc. iii. 641 = roseа.
laciniata, *Linn. Sp. Pl.* 1053.—Europ.; Am. bor.
laciniata, Pursh, Fl. Am. Sept. i. 199 = Purshiana.
laciniata, Schkuhr, Handb. t. 249 = hastata.
lacvis, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 307.—Sibir.
Lampa, *Gill. ex Moq. in DC. Prod.* xiii. ii. 110.— Chili.
lanianthа, Boiss. Diagn. Ser. I. xii. 95 = tatarica.
latifolia, Wahlenb. Fl. Suec. ii. 690 = hastata.
Lehmanniana, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 451 = turcomanica.
lenticularis, *C. A. Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* (1838) 99, nomen.—As. temp.
lenticularis, Tausch, in Flora, xxiii. (1840) i. 16 = Tauschii.
lentiformis, *S. Wats. in Proc. Am. Acad.* ix. (1874) 118.—Am. bor. occ.
leptocarpa, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 74.—Australia.
leptoclada, Boiss. Diagn. Ser. II. iv. 74 = turcomanica.
leucoclada, *Boiss. Diagn. Ser.* I. xii. 95.—Aegypt.; Arab.
leucophylla, *D. Dietr. Syn. Pl.* v. 586.—Calif.
limbata, *Benth. Fl. Austral.* v. 178.—Austral.
limosa, *Schur, Enum. Pl. Transs.* 576.—Europ.
Lindleyi, Moq. in DC. Prod. xiii. ii. 100 = halimoides.

ATRIPLEX:—
linifolia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 958.—Mexic.
littoralis, *Linn. Sp. Pl.* 1064.—Reg. bor. temp.
longipes, Drejer & Fries, Summa Veg. Scand. 201 = hastata.
lucida, Desf. ex Mert. & Koch, in Roehl. Fl. Deutschl. ii. 311 = nitens.
lucida, Moench, Meth. Suppl. 121 = sibirica.
lunulata, Del. ex Moq. in DC. Prod. xiii. ii. 99 = mollis.
macrodira, Gass. Fl. Sic. Syn. ii. 592 = patula.
macrotheca, Hort. ex Moq. in DC. Prod. xiii. ii. 94 = calotheca.
marina, Linn. Mant. ii. 300 = littoralis.
maritima, Crantz, Inst. i. 208 = Suæda maritima.
maritima, Pall. Rosse, ii. 389 = littoralis.
mauritanica, *Boiss. & Reut. Pugill. Pl. Nov.* 106.—Hispan.
micrantha, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 788 = nitens.
micrantha, C. A. Mey. in Ledeb. Fl. Alt. iv. 308 = hastata.
micrantha, *Moq. Chenop. Enum.* 62.—Sibir.
microcarpa, *D. Dietr. Syn. Pl.* v. 586.—Calif.
microcarpa, *Benth. Fl. Austral.* v. 176.—Austral.
microcarpa, Waldst. & Kit. Pl. Rar. Hung. iii. 278. t. 250 = hastata.
microphylla, D. Dietr. Syn. Pl. v 536 = parvifolia, H. B. & K.
microphylla, Willd. Sp. Pl. iv. 958 = Chenopodium microphyllum.
microphylla, *Phil. Fl. Atac.* 48.—Chili.
microphylla, Schur, Enum. Pl. Transs. 578 = laciniata.
microsperma, Bab. Man. ed. I 258 = hastata.
microsperma, Waldst. & Kit. Pl. Rar. Hung. iii 278. t. 250 = hastata.
microtheca, Fries, Corp. Fl. Suec. 133 = hastata.
microtheca, Moq. in DC. Prod. xiii. ii. 91 = hortensis.
mollis, *Desf. Fl. Atlant.* ii. 391.—Algeria.
mollis, Uspenski, Jekaterinenburg. 106 = verrucifera.
Moneta, *Bunge, ex Boiss. Fl. Orient.* iv. 912.—Persia; Afghanist.
monilifera, *S. Wats. in Proc. Am. Acad.* ix (1874) 111.—Am. bor.
monoica, Moench, Meth. 353 = rosea.
montevidensis, *Spreng. Syst.* iii. 918.—Reg. Argent. Moquiniana, *Webb, ex Moq. in DC. Prod.* xiii. ii. 97.—Austral.
mucronata, Rafin. in Am. Monthly Mag. (1818) 176 = arenaria.
mucronata, *Phil. Fl. Atac.* 47.—Chili.
Muelleri, *Benth. Fl. Austral.* v. 175.—Austral.
multifida, Crantz, Inst. i. 207 = Roubieva multifida.
multifida, Desf. Tabl. ed. II. 51 = calotheca.
muralis, Crantz, Inst. i. 206 = Chenopodium murale.
muricata, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 959.—Am. austr.
nemorensis, Schur, Enum. Pl. Transs. 575 = patula.
nitens, *Schkuhr, Handb.* iii. (1803) 541.—Europ.; Orient.
nudicaule, *Boguslaw, in Erman, Archiv*, vi. 65.—Rossia.
nummularia, *Lindl. in Mitch. Journ. Trop. Austr.* (1848) 64.—Austral.
Nuttalli, *S. Wats. in Proc. Am. Acad* ix. (1874) 116.—Am. bor. occ.
oahuensis, *Meyen, Reise*, ii. 127.—Ins. Sandvic.
oblongifolia, Host, Fl. Austr. i. 320 = hastata.
oblongifolia, Waldst. & Kit. Pl. Rar. Hung. iii. 278. t. 211 = patula.
obovata, *Moq. Chenop. Enum.* 61.—Mexic.
obtusa, *Willd. Sp. Pl.* iv. 964.—Sibiria.
obtusata, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 42.—Hispan.
occidentalis, D. Dietr. Syn. Pl. v. 587 = canescens.
ocymifolia, Viv. Eg. 32 = coriacea.
odorata, Pers. Syn. i. 392 = Exomis albicans.
Olivieri, Moq. Chenop. Enum. 93 = tatarica.
oppositifolia, DC. Rapp. i. 13 = hastata.
oppositifolia, *Vill. Prosp.* 21.—Gallia.
oppositifolia, *S. Wats. in Proc. Am. Acad.* ix. (1874) 118.—Mexic.
orbicularis, *S. Wats. l. c.* xvii. (1882–82) 377.—Am. bor.
palaestina, *Boiss. Diagn.* Ser. I. xii. 96.—Syria; *Assyria*.

ATRIPLEX:—
Palmeri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 146.—Calif.
paludosa, *R. Br. Prod.* 406.—Austral.
paludosa, Nees, in Lehm. Pl. Preiss. i. 683 = Drummondii.
pamparum, *Griseb. in Gott. Abh.* xix. (1874) 69.—Reg. Argent.
Parishii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 377.—Am. bor.
Parryi, *S. Wats. l. c.* 378.—Am. bor.
parvifora, Host, Fl. Austr. i. 161 = seq.
parvifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 192.—Mexic.
parvifolia, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 16.—Marocco; Ins. Canar.
patagonica, *D. Dietr. Syn. Pl.* v. 586.—Patagonia.
patula, *Linn. Sp. Pl.* 1053.—Reg. temp. bor. et austr
patula, Sm. Fl. Brit. 1091; Engl. Fl. iv. 257 = hastata
pedunculata, *Linn. Cent. Pl.* i. 33.—Europ.; As. bor., Orient.
persica, *Boiss. Diagn.* Ser. I. xii. 96.—Persia.
peruviana, *Moq. in DC. Prod.* xiii ii. 102.—Peruv
Pharaonis, Hort. Avenion. ex Moq. l. c. 91 = microtheca.
philippica, *Weinm. in Syll. Ratisb.* i. (1828) 29.—Ins. Philipp.
phyllostegia, *S. Wats. in Proc. Am. Acad.* ix. (1874) 108.—Am. bor. occ.
Piquerez, Lag. Pl. Barcll. 67 = laciniata.
platyspatula, Guss. Fl. Sic. Syn. ii. 590 = hastata.
plebeja, *Cavmich. in Trans. Linn. Soc.* xii. (1819) 508.—Ins. Trist. d'Acunha.
polycarpa *S. Wats. in Proc. Am. Acad* ix. (1874) 117.—Calif.
polygama, *Sessé, in Lag. Gen. et Sp. Nov.* 12.—Mexic.
polymorpha, Cass. Germ. et Wedd. Cat. Pl. Par 108 = farinosa, patula.
polysperma, Crantz, Inst. i. 207 = Chenopodium polyspermum.
polysperma, Tenore, Prod. Fl. Nap. i. 57 = rosea.
pumilacoides, *Linn. Sp. Pl.* 1053.—Europ.; Syria; Afr. bor.
pumilacoides, Pall. Ind. Taur. 59 = verrucifera.
Powellii, *S. Wats. in Proc. Am. Acad.* ix. (1874) 114.—Arizona.
procumbens, Jundz. Fl. Lituan. 78 = patula.
procumbens, *Lesz. in Linnaea*, ix (1834) 208.—Sibir. alt.
prostrata, Bouch. Fl. Abbev. 76; Bab. Man. ed. I. 262 = hastata.
prostrata, *R. Br. Prod.* i. 406.—Austral.
prostrata, Moq. in DC. Prod. xiii. ii. 99 = cinerea.
prostrata, *Phil. in Anal. Univ. Chil.* (1873) 535.—Chili.
pruinosa, Sieber, ex Moq. in DC. Prod. xiii. ii. 93 = laciniata.
pruinosa, Sieber, ex Boiss. Fl. Orient. iv. 910 = tatarica.
Pseudo-nitens, *Schur, Enum. Pl. Transs.* 574.—Hungaria.
Pumilio, *R. Br. Prod.* 406.—Austral.
pungens, *Trautv. in Bull. Soc. Nat. Mosc.* xli. (1867) ii. 58.—Sibiria.
Purshiana, *Moq. Chenop. Enum.* 55.—Am. bor.
pusilla, *S. Wats. in Proc. Am. Acad.* ix. (1874) 110.—Am. bor. occ.
ramosissima, Nutt. ex Moq. in DC. Prod. xiii. ii. 111 = microsperma.
recta, Schur, Enum. Pl. Transs. 575 et Ind. = patula.
recurva, Lur. in Mém. Soc. Linn. Par. i. (1822) 284 = tatarica.
reniformis, R. Br. Prod. 406 = paludosa.
reniformis, F. Muell. Fragm. vii. 9 = stipitata.
repens, Allch. Cat. Pl. Panjab, 128 = Stocksii.
repens, *Roth, Nov. Pl. Sp.* 377.—Ind. or.
retusa, *C. Gay, Fl. Chil.* v. 240.—Chili.
rhagodioides, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 74.—Austral.
rosea, Bab. Man. Brit. ed. I. 253 = crassifolia.
rosea, *Linn. Sp. Pl.* ed. II. 1493.—Europ.; As. Min.; Afr. bor.
rosea, F. Muell. Fragm. vii. 9 = Muelleri.
rotundifolia, *Dombey, ex Moq. Chenop. Enum.* 70.—Peruv.
rotundifolia, Boiss. Fl. Orient. iv. 915 = crystallina.
rubrovirens, Hort. ex Moq. in DC. Prod. xiii. ii. 91 = hortensis.

ATRIPLEX:—
rubra, Crantz, Inst i. 206 = hortensis.
ruderalis, Wallr. Sched. Crit. 115 = hastata.
saccaria, *S. Wats. in Proc. Am. Acad.* ix. (1874) 112.—Utah.
Sachsi, Rostc. & Schum. Fl. Sedin. 401. t. 1 = hastata.
sagittata, Borckh. in Reinsch. Mag. 477 = nitens.
salicifolia, Lag. ex Moq. Chenop. Enum. 74 = polygama.
salicina, Pall. Reise. i. 495 = littoralis
salina, Sievers, in Pall. N. Nord. Beitr. vii. 327 = littoralis.
salina, Desf. in Journ. Bot. i. (1813) 48 = patula.
scoparia, Crantz, Inst. i. 208 = Kochia scoparia.
semibaccata, *R. Br. Prod.* 406.—Austral.
semibaccata, Moq. in DC. Prod. xiii. ii. 97 = spongiosa
semibracteata, Steud. Nom. ed. II. i. 169, sphalm. = semibaccata.
serotina, Crantz, Inst. i. 206 = Chenopodium serotinum.
serpyllifolia, *Bunge, in Act. Hort. Petrop.* v. (1877) 642.—Sibiria; Turkest.
serrata, Huds. Fl. Angl. ed. I. 377 = littoralis.
sibirica, *Linn. Sp. Pl.* ed. II. 1493.—Sibiria.
sicula, *Ucria, in Nov. Act. Acad. Panorm.* i. 70.—Sicilia.
sinuata, Hoffm. Deutschl. Fl. ed. II. ii. 277 = laciniata.
Smithii, Syme, Eng. Bot. ed III. viii. 32 = hastata.
spectabilis, Ehrh. ex Moq. in DC. Prod. xiii. ii. 91 = hortensis
spicata, Stokes, Bot. Mat. Med. ii. 24 = Anredera scandens?
spicata, *S. Wats. in Proc. Am. Acad.* ix. (1874) 108.—Calif.
spinacifolia, Stokes, Bot. Mat. Med. ii. 28 = hastata.
spinosa, *D. Dietr. Syn. Pl.* v. 586.—Am. centr.
spongiosa, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 74.—Austral.
stipitata, *Benth. Fl. Austral.* v. 168.—Austral.
stipitata, Westerl. in Linnaea, xl. (1876) 165 = calotheca.
Stocksii, *Boiss. Diagn.* Ser. II. iv. 78.—Scinde.
stylosa, *Viv. Fl. Eg.* 32.—Lybia.
subopposita, *Schur, in Verh. Siebenb. Ver. Naturw.* iv (1853) 66.—Europ.
Suckleyana, *S. Wats. in Proc. Am. Acad.* ix. (1874) 111.—Am. bor.
sulcata, Michx. ex Roem. & Schult. Syst. vi. 290 = littoralis.
tartarica, Crantz, Inst. i. 205 = tatarica.
tatarica, Auct. fid. Boiss. Fl. Orient. iv. 909 = patula.
tatarica, *Linn. Sp. Pl.* 1053.—As. centr.
tatarica, Schkuhr, Handb. t. 149 = oblongifolia.
Tauschii, *D. Dietr. Syn. Pl.* v. 588.—Hab. ?
tetrandra, Torr. ex Moq. in DC. Prod. xiii. ii. 93 = Parishiana.
texana, *S. Wats. in Proc. Am. Acad.* ix. (1874) 113.—Texas.
thunbergiaefolia, *Boiss. Fl. Orient.* iv. 911.—Babylonia.
Thunbergiana, Roem. & Schult. Syst. vi. 281 = laciniata.
Tornabeni, Tineo, in Guss. Fl. Sic. Syn. ii. 589 = laciniata.
Torreyana, *Rafin. Atl. Journ.* 146.—Am. bor.
Torreyi, *S. Wats. in Proc. Am. Acad.* ix. (1874) 118.—Am. bor. occ.
transiliunica, Schur, Enum. Pl. Transs. 577 = hastata.
triangularis, Willd. Sp. Pl. iv. 963 = hastata.
truncata, *A. Gray, in Proc. Am. Acad.* viii. (1873) 398.—Am. bor. occ.
turcomanica, *Fisch. & Mey. ex Karel. in Bull. Soc. Nat. Mosc.* (1839) ii. 169.—Persia.
turcomanica, Turcz. ex Ledeb. Fl. Ross. iii. 720 = laciniata.
undulata, *D. Dietr. Syn. Pl.* v. 586.—Patagon.
urbica, Crantz, Inst. i. 206 = Chenopodium urbicum.
velutinella, *F. Muell. Rep. Babb. Exp.* 30.—Austral.
venata, Moq. Chenop. Enum. 54 = microtheca.
venata, Willd. Sp. Pl. iv. 962 = tatarica.
Verreauxii, *Moq. in DC. Prod.* xiii. ii. 98.—Afr. austr.
verrucifera, *Bieb. Fl. Taur. Cauc.* ii. 441.—Reg. Caucas.; Persia.
verticillata, Lag. Gen. et Sp. Nov. 12 = rosea.
vesicaria, *Howard, ex Benth. Fl. Austral.* v. 179.—Austral.
virgata, Roth, Nov. Sp. Pl. 377 = hortensis.
virgata, Soop. Delic. Insub. ii. 14; Hoffm. Deutschl. Fl. ed. II. ii. 277 = patula.

ATRIPLEX :—
viridis, Crantz, Inst. t. 207 = Chenopodium album.
viridis, Hort. ex Moq. in DC. Prod. xiii. ii. 90 = nitens.
Vulvaria, Crantz, Inst. t. 207 = Chenopodium Vulvaria.
Wallii, *S. Wats. in Proc. Am. Acad.* ix. (1874) 112.—Am. bor.
Wrightii, *S. Wats. l. c.* 113.—N. Mexic.
zosterafolia, *S. Wats. l. c.* 109.—Am. bor.

ATROPA, Linn. Gen. ed. I. 49 (1737). *SOLANACEÆ*, Benth. & Hook. f. ii. 900.
BELLADONA, Adans. Fam. ii. 219 (1763).
acaulis, Stokes, Bot. Mat. Med. i. 396 = Mandragora officinarum.
acuminata, Royle, Illustr. Bot. Himal. 279 = A. Belladonna.
ambigua, Salisb. Prod. 132 = Scopola carniolica.
arborea, Willd. ex Dun. in DC. Prod. xiii. i. 497 = Acnistus arborescens.
arborescens, Linn. Cent. ii. 10 (Amœn. Acad. iv. 307) = Acnistus arborescens.
arborescens, Roem. & Schult. Syst. iv. 585 = Acnistus Plumieri.
arenaria, Willd. ex Roem. & Schult. Syst. iv. 586 = Nectouxia formosa.
aristata, Poir. Encyc. Suppl. i. 609 = Withania aristata.
aspera, Ruiz & Pav. Fl. Peruv. ii. 45 = Hebecladus asperus.
baccifera, *Willd. in Linnæa*, xxv. (1852) 50.—Hispan.
Belladonna, *Linn. Sp. Pl.* 181.—Europ.; Orient; Ind. or.
bicolor, Ruiz & Pav. Fl. Peruv. ii. 45 = Hebecladus bicolor.
biflora, Ruiz & Pav. l. c. = Hebecladus biflorus.
centoria, Pers. Syn. i. 219 = Saracha centoria.
daturæfolia, Thore, Chlor. Land. 74 = Nicandra physaloides.
dentata, Pers. Syn. i. 219 = Saracha dentata.
dependens, Mathews, ex Hook. Ic. Pl. t. 107 = Salpichroa dependens.
erecta, Hornem. Hort. Hafn. 957 = Hebecladus biflorus.
erecta, Zuccagni, in Roem. Collect. 129 = Saracha Zuccagniana.
flexuosa, Willd. ex Roem. & Schult. iv. 586; Dun. in DC. Prod. xiii. 465 = Hebecladus viridiflorus.
frutescens, Linn. Sp. Pl. 182 = Withania frutescens.
glandulosa, Hook. Bot. Misc. ii. (1831) 230 = Salpichroa glandulosum.
gymnosperma, Schmid. Ic. 67. t. 18 = Nolana prostrata.
herbacea, *Mill. Gard. Dict.* ed. VIII. n. 3.—Campechia.
hirsuta, Meyen, Reise, i. 466 = Salpichroa hirsutum.
hirtella, *Sprang. Syst.* i. 699.—Bras.
humifusa, Gouan, Hort. Monsp. 106 = Nolana prostrata.
humilis, Sal sb. Prod. 132 = Mandragora officinarum.
lethalis, Salisb. l. c. = A. Belladonna.
lutescens, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 241 = A. Belladonna.
Mandragora, Linn. Syst. ed. X. 988 = Mandragora officinarum.
Mandragora, Sibth. & Sm. Fl. Græc. iii. 26. t. 232 = Mandragora autumnalis.
origanifolia, Desf. Cat. Hort. Par. ed. III. 396 = Physalis curassavica, origanifolia.
physaloides, Linn. Sp. Pl. 181 = Nicandra physaloides.
physaloides, Georgi, Beschr. Russ. Reich. Nachtr. 261 = Scopola physaloides.
physaloides, Linn. Syst. ed. XII. 171 = *physaloides*.
picata, Roth, Catalecta, ii. 24 = Saracha procumbens.
procumbens, Cav. Ic. i. 53 = Saracha procumbens.
punctata, Pers. Syn. i. 219. = Saracha punctata.
ramosissima, Mathews, ex Dunal, in DC. Prod. xiii. i. 473 = Salpichroa ramosissimum.
rupicola, Walp. Rep. iii. 103, in syn. = seq.
rvcolnicis, F. G. Dietr. Vollst. Gaertn. Lexik. Nachtr. i. 422 = Hebecladus umbellatus.
rhomboidea, Gill. & Hook. in Hook. Bot. Misc. i. (1830) 126 = Salpichroa rhomboidum.
Rothii, Poir. Encyc. Suppl. i. 610 = Hebecladus umbellatus.
stderoxyloides, Willd. ex Roem. & Schult. Syst. iv. 586 = Acnistus arborescens.

ATROPA:—
solanacea, All. ex Steud. Nom. ed. II. i. 170 = Acnistus arborescens.
solanacea, Linn. Mant. ii. 205 = Solanum aggregatum.
spinosa, Meyen, Reise, i. 466 = Dunalia lycioides
umbellata, Ruiz & Pav. Fl. Peruv. ii. 44 = Hebecladus umbellatus.
villosa, Zuccagni, in Roem. Collect. 130 = Hebecladus spectabilis.
viridiflora, H. B. & K. Nov. Gen. et Sp. iii. 11 = Hebecladus viridiflorus.

ATROPIS, Rupr. Fl. Samoj. [Beitr. Fl. Russ. Reich. ii.] 64 (1845) = **Glyceria**, R. Br. (Gramin.).
angustata, Griseb. in Ledeb. Fl. Ross. iv. 390 = G. distans.
californica, Munro, ex A. Gray, in Proc. Acad. Sc. Philad. 1862 (1863) 286.—Am. bor. occ.
capillaris, Schur, Enum. Pl. Transs. 779 = G. distans.
carinata, Griseb. in Goett. Abh. xxiv. (1879) 291.—Reg. Argent.
convoluta, Griseb. in Ledeb. Fl. Ross. iv. 389 = G. distans.
distans, Rupr. Fl. Samoj. 64; Griseb. in Ledeb. Fl. Ross. iv. 388 = G. distans.
festucæformis, Schur, Enum. Pl. Transs. 780 = G. maritima.
intermedia, Schur, l. c. 779 = G. maritima.
magellanica, E. Desv. in C. Gay, Fl. Chil. vi. 394 = G. magellanica.
maritima, Griseb. in Ledeb. Fl. Ross. iv. 389 = G. distans.
pauciflora, Thurb. in S. Wats. Bot. Calif. ii. 510 = Poa pauciflora.
procumbens, Thurb. l. c. 509.—Am. bor.
Parade-maritima, Schur, Enum. Pl. Transs. 780 = G. maritima.
pumila, T. Kirk, in Gard. Chron. (1882) i. 261; Trans. N. Z. Inst. xiv. (1882) 379.—N. Zel.
scabrella, Thurb. in S. Wats. Bot. Calif. ii. 510 = Poa scabrella.
tenuiflora, Griseb. in Ledeb. Fl. Ross. iv. 389 = G. tenuifolia.
tenuifolia, Thurb. in S. Wats. Bot. Calif. ii. 510 = G. tenuifolia.
transeuntea, Schur, Enum. Pl. Transs. 889, Index = G. maritima.

ATRUTEGIA, *Bedd. in Madr. Journ.* Ser. III. i. (1864) 87. t. 1. *ANONACEÆ*, Benth. & Hook. f. Gen. i. 957.
wynadensis, *Bedd. l. c.*—Ind. or.

ATTALEA, H. B. & K. Nov. Gen. et Sp. i. 309 (1815). *PALMÆ*, Benth. & Hook. f. iii. 947.
acaulis, *Hort. ex H. Wendl. Cycl. Pand. Cycad.* 4.—Hab.?
agrestis, Barb.-Rodr. Enum. Palm. Nov. 42 = microcarpa.
amygdalina, *H. B. & K. Nov. Gen. et Sp.* i. 310. t. 95.—Am. austr.
bleoharospus, *Mart. in Palmet. Orb.* 116. t. 5. f. 2.—Bolivia.
butyrosa, Lodd. ex H. Wendl. in Kerch. Palm. 232 = humilis.
Cohune, Poepp. ex Mart. in Palmet. Orb. 119 = Scheelea Cephalotes.
Cohune, *Mart. l. c.* 121; *et Hist. Nat. Palm.* iii. 300. t. 167.—Honduras.
compta, *Mart. Hist. Nat. Palm.* ii. 137. t. 41, 75, 97.—Bras.
coronata, *Lodd. ex H. Wendl. Cycl. Pand. Cycad.* 5.—Hab.?
excelsa, *Mart. Hist. Nat. Palm.* ii. 135. t. 96. f. 3.—Bras.
exigua, *Drude, in Mart. Fl. Bras.* iii. ii. 439.—Bras.
funifera, *Mart. Hist. Nat. Palm.* ii. 136. t. 95, 96. f. 4.—Bras.
gomphocoeca, *Mart. l. c.* 201.—Am. austr.
grandis, *Hort. ex H. Wendl. in Kerch. Palm.* 233.—Hab.?
Humboldtiana, *Spruce, Palm. Amaz.* 168.—Am. trop.
humilis, *Mart. in Sprang. Syst.* ii. 694.—Bras.
limbata, *Soem. ex H. Wendl. in Kerch. Palm.* 233.—Hab.?
macrocarpa, *Linden, Illustr. Hortic.* xxviii. (1881) 15.—Bras.

ATTALEA:—
Magdalenae, *Linden, Illustr. Hortic.* xxviii. (1881) 15.—N. Granat.
Maraca, *Linden, l. c.*—N. Granat.
maracaibensis, *Mart. in Palmet. Orb.* 124.—Peruv.
Maripa, Mart. l. c. 123 = Maximiliana Maripa.
microcarpa, *Mart. l. c.* 125.—Bras.
monosperma, Barb.-Rodr. Enum. Palm. Nov. 42 = spectabilis.
nucifera, Karst. in Linnæa, xxviii. (1856) 255 = amygdalina.
phalerata, *Mart. in Sprang. Syst.* ii. 694.—Bras.
Piassaba, Barb.-Rodr. Enum. Palm. Nov. 42 = spectabilis.
Principis, Mart. in Palmet. Orb. 118. t. 4. f. 2 = Scheelea Principis.
purpurea, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—N. Granat.
paruénsis, *Linden, l. c.*—Fl. Amazon.
racemosa, Spruce, Palm. Amaz. 166 = Orbignia racemosa.
Rossii, *Lodd. ex Loud. Hort. Brit.* 387.—Bras.
rostrata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 50.—Costa Rica.
speciosa, *Mart. Hist. Nat. Palm.* ii. 138.—Bras.
spectabilis, *Mart. l. c.* 136. t. 96. f. 1, 2.—Bras.
spinosa, *Meyen, Reise*, i. 469.—Peruv.
Tissae, *Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Fl. Amazon.
tranativa, *Barb.-Rodr. Prot. App.* 49.—Bras.
venatorum, Mart. Hist. Nat. Palm. iii. 325 = Maximiliana venatorum.

ATTRACTILIS, Hall. ex Scop. Introd. (1777) 194 = **Atractylis**, Linn. (Compos.).

ATULANDRA, Rafin. Sylva Tellur. 81 (1838) = **Rhamnus**, Linn.
arragonensis, Rafin. l. c. = R. lycioides.
valentina, Rafin. l. c. = R. pumila.

ATUNA, Rafin. Sylva Tellur. 153 (1838) = **Heritiera**, Dryand. (Sterculi.).
racemosa, Rafin. l. c. = H. littoralis.

ATYLOSIA, Wight & Arn. Prod. 257 (1834). *LEGUMINOSÆ*, Benth. & Hook. f. i. 542.
CANTHAROSPERMUM, Wight & Arn. Prod. 255 (1834).
albicans, *Benth. in Miq. Pl. Junghh.* 248.—Ind. or.
barbata, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 216.—Ind. or.; Malaya.
calycina, *Kurz, in Journ. As. Soc. Beng.* xlii. (1874) ii. 156.—Ind. or.
candicans, Kurz, l. c. = Rhynchosia arvensis.
Candollei, *Wight & Arn. Prod.* i. 257.—Ind. or.
cinerea, *F. Muell. ex Benth. Fl. Austral.* ii. 264.—Austral.
circinnalis, Benth. in Miq. Pl. Junghh. 244 = Dunbaria circinnalis.
elongata, *Benth. l. c.* 243.—Ind. or.
geminiflora, *Dalz. in Journ. Linn. Soc.* xiii. (1873) 184.—Ind. or.
glandulosa, Dalz. l. c. = mollis.
goensis, Dalz. l. c. 186 = barbata.
grandiflora, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 214.—Reg. Himal.
grandiflolia, *F. Muell. ex Benth. Fl. Austral.* ii. 264.—Austral.
kulnensis, *Dalz. in Journ. Linn. Soc.* xiii. (1873) 186.—Ind. or.
\Lawii, *Dalz. l. c.*—Ind. or.
Lawii, Wight, ic. t. t. 93 = linosta.
linosta, *Wight & Arn. Prod.* i. 258.—Ind. or.
major, Wight & Arn. l. c. 257 = Candollei.
marmorata, *Banks, ex Benth. Fl. Austral.* ii. 263.—Austral.
mollis, *Benth. in Miq. Pl. Junghh.* 243.—Ind. or.; Malaya.
nivea, *Benth. l. c.*—Burma.
platycarpa, *Benth. l. c.*—Reg. Himal.
pluriflora, *F. Muell. ex Benth. Fl. Austral.* ii. 264.—Austral.
punctata, Dalz. in Journ. Linn. Soc. xiii. (1873) 186 = Dunbaria conspersa.
reticulata, *Benth. Fl. Austral.* ii. 263.—Austral.
rostrata, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 216.—Ind. or.
rugosa, *Wight & Arn. Prod.* i. 257.—Ind. or.

ATYLOSIA :—
scarabæoides, *Benth. in Miq. Pl. Jungh.* 242.—Austral.
sericea, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 218.—Ind. or.
subrhombea, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 51 = Dumbaria subrhombea.
villosa, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 214.—Reg. Himal.

ATYLUS, Salisb. Parad. Lond. sub t. 67 (1807) = **Petrophila**, R. Br. (Proteac.).
anemonifolia, Salisb. l. c. = Isopogon anemonifolius.
fucifolia, Salisb. l. c. = P. pulchella.

AUBENTONIA, Domb. ex Steud. Nom. ed. I. 93 (1821) = **Waltheria**, Linn. (Stercul.).
tomentosa, Domb. ex Steud. l. c. = W. ovata.

AUBERTIA, Bory, Voy. i. 356. t. 18 (1804) = **Zanthoxylum**, Linn.† (Rutac.).
Bajarnardia, Buch.-Ham. in Wall. Cat. sub n. 7118 = Z. Bajarnardia.
borbonica, Bory, Voy. i. 356. t. 18 (an Enodiae sp. !—Cf. Benth. & Hook. f. Gen. i. 296) = Z. Aubertia.
piperita, Buch.-Ham. ex Wall. Cat. sub n. 7117 = Z. Hamiltonianum.
Timber, Buch.-Ham. ex Wall. l. c. n. 7116 = Z. acanthopodium.
tomentosa, Bory, ex Steud. Nom. ed. II. i. 171 = Z. Aubertia.

AUBERTIA, Chapel. ex Baill. Adansonia, ii. (1861) 162, 166 = **Croton**, Linn. (Euphorb.).
argentea, Chapel. ex Baill. l. c. 162 = C. Noronhae.
glandulosa, Chapel. ex Baill. l. c. 166 = C. Chapelieri.

AUBION, Rafin. Sylva Tellur. 110 (1838) = **Cleome**, Linn. (Capparid.).
Chelidoni, Rafin. l. c. = C. Chelidonii.

AUBLETIA, Gaertn. Fruct. i. 379. t. 78. f. 2 (1788) = **Sonneratia**, Linn. f. (Lythrar.).
caseolaris, Gaertn. l. c. = S. acida.

AUBLETIA, Lour. Fl. Cochinch. 283 (1790) = **Paliurus**, Tourn. (Rhamnac.).
ramosissima, Lour. l. c. = P. ramosissimus.

AUBLETIA, Neck. Elem. i. 356 (1790) = **Ruellia**, Linn. (Acanth.).

AUBLETIA, Rich. in Pers. Syn. ii. 688 (1807) = **Monneria**, Linn. (Rutac.).
trifoliata, Pers. l. c. = M. trifolia.

AUBLETIA, Schreb. Gen. 358 (1789) = **Apeiba**, Aubl. (Tiliac.).
aspera, Willd. Sp. Pl. ii. 1156 = Apeiba aspera.
dicolor, Spreng. Neue Entdeck. ii. 166 = Apeiba dicolor.
laevis, Willd. Sp. Pl. ii. 1156 = Apeiba glabra.
Petoumaa, Willd. l. c. = Apeiba Petoumaa.
Tibourbou, Willd. l. c. = Apeiba Tibourbou.
Tiburbu, Sw. Fl. Ind. Occ. ii. 944 = prasc.

AUBRIETA, Adans. Fam. ii. 490 (1763) (Aubrieta); DC. in Mém. Mus. Par. vii. (1821) 393; Syst. ii. 293. *CRUCIFERAE*, Benth. & Hook. f. Gen. i. 73.
canescens, Boiss. Fl. Orient. i. 253 = deltoides.
Columnae, *Guss. Pl. Rar.* 266.—Ital.
croatica, *Schott, Nyman & Kotschy, Analect. Bot.* 47. —Croatia.
deltoides, *DC. Syst.* ii. 294.—Italia; Graecia; As. Min.
diffusa, Spreng. Syst. ii. 871 = Hellophila diffusa.
edentula, *Boiss. Fl. Orient.* i. 254.—Kurdistan.
erubescens, *Griseb. Spicil. Fl. Rumel.* i. 268.—Graecia.
floribunda, Spach, Hist. Vég. Phan. vi. 469 = deltoidea.
gracilis, *Sprun. ex Boiss. Diagn.* Ser. I. i. 74.—Graecia.
graeca, Griseb. Spicil. Fl. Rumel. i. 268 = deltoidea.
hesperidiflora, G. Don, ex Loud. Hort. Brit. 257 = deltoidea.
integrifolia, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 35 = deltoidea.
intermedia, *Heldr. & Orph. ex Boiss. Diagn.* Ser. II. i. 90.—Graecia.

AUBRIETIA :—
Kotschyi, *Boiss. & Hohen. Diagn.* Ser. I. viii. 93.—Persia.
italica, Boiss. Fl. Orient. i. 252, in nota = Columnae.
lepidioides, Spreng. Syst. ii. 871 = Heliophila diffusa.
libanotica, *Boiss. & Hohen. Diagn.* Ser. I. viii. 92.—Mont. Libanus.
olympica, *Boiss. Fl. Orient.* i. 251.—Bithynia.
parviflora, *Boiss. Diagn.* Ser. I. vi. 14.—Persia.
Pinardi, *Boiss. l. c.* Ser. II. i. 37.—As. Min.
purpurea, DC. Syst. ii. 294 = Arabis purpurea.

AUBRYA, Baill. Adansonia, ii. 265 (1861–62). *HUMIRIACEAE*, Benth. & Hook. f. i. 988.
gabonensis, *Baill. l. c.* 266.—Afr. trop.

AUCHERA, DC. Coll. Mém. ix. t. 11; Prod. vi. 557 (1837) = **Cousinia**, Cass. (Compos.).
floribunda, Regel, Cat. Pl. Hort. Aksakov. 13.—Hab.† *orientalis*, DC. Coll. Mém. ix. t. 11 = C. orientalis.

AUCKLANDIA, Falc. in Trans. Linn. Soc. xix. (1845) 28 = **Saussurea**, DC. (Compos.).
Costus, Falc. l. c. = S. Lappa.

AUCUBA, Thunb. Nov. Gen. 61 (1788). *CORNACEAE*, Benth. & Hook. f. i. 960.
EUBASIS, Salisb. Prod. 68 (1796).
chinensis, *Benth. Fl. Hongk.* 138.—China.
himalaica, *Hook. f. & Thoms. Illustr. Himal. Pl.* t. 12. —Reg. Himal.
japonica, *Thunb. Nov. Gen.* 61.—Japon.

AUCUBA, Cham. in Linnaea, v. (1830) 56, in syn. = ARUBA, Neck & Mart. = **Baputia**, Aubl. (Rutac.).

AUCUBAEPHYLLUM, Ahlburg, in Bot. Zeit. xxxvi. (1878) 113. *CORNACEAE*.
Ilookianae, *Ahlburg, l. c.*—Japon.

AUCUPARIA, Riv. ex Rupp. Fl. Jen. ed. Haller. 140 (1745) = **Pyrus**, Linn. (Rosac.).
pinnata, Pourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 178 = P. Aucuparia.
sylvestris, Medic. Gesch. 86 = P. Aucuparia.

AUDIBERTIA, Benth. in Bot. Reg. sub t. 1282 (1829) = **Mentha**, Linn. (Labiat.).
parviflora, Nym. Consp. 597 = M. Requieni.
pusilla, Benth. in Bot. Reg. sub t. 1282 = M. Requieni.

AUDIBERTIA, Benth. in Bot. Reg. t. 1469 (1831). *LABIATAE*, Benth. & Hook. f. ii. 1197.
capitata, *A. Gray, in Proc. Am. Acad.* vii. (1868) 387. —Calif.
Cleveland, *A. Gray, l. c.* x. (1874) 76.—Calif.
decurrens, Nutt. ex Benth. in DC. Prod. xii. 359 = grandiflora.
Dorrii, *Kellogg, in Proc. Calif. Acad.* ii. (1868) 190.—Calif.
grandiflora, *Benth. Lab. Gen. et Sp.* 312.—Calif.
humilis, *Benth. l. c.* 313.—Calif.
incana, *Benth. in Bot. Reg.* t. 1469.—Am. bor. occ.
stives, *Benth. Lab. Gen. et Sp.* 313.—Calif.
Palmeri, *A. Gray, in Bot. Calif.* i. 601.—Calif.
polystachya, *Benth. Lab. Gen. et Sp.* 314.—Calif.
spinulosa, Nutt. ex Benth. in DC. Prod. xii. 359 = stachyoides.
stachyoides, *Benth. Lab. Gen. et Sp.* 313.—Calif.
Vaseyi, *T. C. Porter, in Coult. Bot. Gazette*, vi. (1881) 207.—Calif.

AUDOUINIA, Brongn. in Ann. Sc. Nat. viii. (1826) 384. t. 88. *BRUNIACEAE*, Benth. & Hook. f. i. 673.
PAVINDA, Thunb. ex Bartl. Ord. 374 (1830).
capitata, *Brongn. l. c.*—Afr. austr.

AUGANTHUS, Link, Handb. i. 415 (1829) = **Primula**, Linn.
praecitens, Link, l. c. = P. sinensis.

AUGEA, Thunb. ex Retz. Obs. v. 2 (1789) = **Lanaria**, Ait. (Haemod.).
lanata, Thunb. ex Retz. l. c. = L. plumosa.

AUGEA, Thunb. Prod. Pl. Cap. 80 (1794); et Nov. Gen. 129 (1798). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 265.
capensis, *Thunb. Il. cc.* 80 et 129.—Afr. austr.

AUGIA, Lour. Fl. Cochinch. 837 (1790). 'Genus ignotum,' Benth. & Hook. f. Gen. i. 418.
sinensis, Lour. l. c. (= *Calophyllum Augia* ex Steud. Nom. ed. II. i. 171).—China.

AUGUSTA, Leandr. Sacram. in Denkschr. Akad. Muench. vii. (1819) 235. t. 14 = **Stiftia**, Mikan (Compos.).
grandiflora, Leandr. Sacram. l. c. = S. chrysantha.
parviflora, Leandr. Sacram. l. c. = S. parviflora.

AUGUSTA, Pohl, Pl. Bras. Ic. ii. t. 1. t. 101–105. *RUBIACEAE*, Benth. & Hook. f. ii. 51.
BONIFACIA, Manso, ex Steud. Nom. ed. II. i. 216 (1840).
SCHREIBERSIA, Pohl, in Flora, viii. (1825) 183.
attenuata, *Pohl, Pl. Bras. Ic.* ii. t. 104.—Bras.
glaucescens, *Pohl, l. c.* t. 105.—Bras.
lanceolata, *Pohl, l. c.* t. 101.—Bras.
oblongifolia, *Pohl, l. c.* t. 103.—Bras.
parviflora, *Pohl, l. c.* t. 102.—Bras.

AUGUSTEA, DC. Prod. iv. 404 (1830) = **Augusta**, Pohl (Rubiac.).

AUGUSTIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 124; et Abh. Akad. Berl. (1854) 80. t. 8. B. = **Begonia**, Linn.
caffra, Klotzsch, Il. cc. 124 & 81 = B. caffra.
Dregei, Klotzsch, Il. cc. 124 & 80 = B. Dregei.
natalensis, Klotzsch, Il. cc. 181 & 80 = B. natalensis.
suffruticosa, Klotzsch, Il. cc. 124 & 82 = B. suffruticosa.

AUGUSTINEA, Karst. in Linnaea, xxviii. (1856) 295 = **Bactris**, Jacq. (Palm.).
balanoidea, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 39 = B. balanoidea.
major, Karst. in Linnaea, xxviii. (1856) 295 = B. major.
ovata, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 38 = B. ovata.

AUGUSTINEA, St. Hil. & Nand. in Ann. Sc. Nat. Sér. III. ii. (1844) 145 = **Miconia**, Ruiz & Pav. (Melast.).
speciosa, St. Hil. & Nand. l. c. = M. speciosa.

AULACIA, Lour. Fl. Cochinch. 278 (1790) = **Clausena**, Burm. f. (Rutac.).
falcata, Lour. Fl. Cochinch. 278 (= *Cookia falcata*).—Cochinch.
falcifolia, Stokes, Bot. Mat. Med. ii. 482 = prasc.
punctata, Raeusch. Nom. ed. III. 119 = Cl. Wampi.

AULACIDIUM, Rich. ex DC. Prod. iii. 113 (1828) = **Macrocentrum**, Hook. f. (Melast.).
cristatum, Rich. ex DC. l. c. 113 = M. cristatum.
fasciculatum, Rich. ex DC. l. c. = M. fasciculatum.

AULACINTHUS, E. Mey. Comm. Pl. Afr. Austr. 155 (1835) = **Lotononis**, Eckl. & Zeyh. (Legum.).
gracilis, E. Mey. l. c. = L. gracilis.
rigidus, E. Mey. l. c. = L. rigidus.

AULACOCALYX, Hook. f. in Hook. Ic. Pl. t. 1126 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 109.
jasminiflora, *Hook. f. l. c.*—Afr. trop.

AULACOCARPUS, Berg, in Linnaea, xxvii. (' 1854 ') 345. *MYRTACEAE*, Benth. & Hook. f. i. 720.
crassifolius, *Berg, l. c.* 346.—Ins. Gorgonae quadrangularis, *Griseb. Fl. Brit. W. Ind.* 289.—Ind. occ.
Sellowianus, *Berg, in Linnaea*, xxvii. (' 1854 ') 346.—Bras.
Wrightii, *Griseb. Cat. Pl. Cub.* 90.—Cuba.

AULACODISCUS, Hook. f. in Benth. & Hook. f. Gen. ii. 71 (1873) (*non Ehrenb. quae Crypt.*). *RUBIACEAE*, Benth. & Hook. f. ii. 71.
premnoides, *Hook. f. l. c.*—Malaya.

AULACOPHYLLUM, Regel, Gartenfl. (1876) 141 = **Samia**, Linn. (Cycad.).
Lindeni, Regel, l. c. = Z. Lindeni.
montanum, Regel, l. c. = Z. montana.
Ortgiesi, Regel, l. c. = Z. Chigua.
Roezli, Regel, l. c. = Z. pseudo-parasitica.
Skinneri, Regel, l. c. 142 = Z. Skinneri.
Wallisi, Regel, l. c. = Z. Wallisii.

AULACORHYNCHUS, Ness, in Linnæa, ix. 305 (1834); ib. x. 198 (1835). *CYPERACEÆ*, Benth. & Hook. f. Gen. iii. 1048.
crinifolius, Steud. Nom. ed. II. i. 172 = seq.
crinifolius, *Ness, in Linnæa*, ix. 305; et x. 199.—Afr. austr.

AULACOSPERMUM, Ledeb. Fl. Alt. iv. 344 (1833) = Pleurospermum, Hoffm. (Umbell.).
amoenum, Ledeb. l. c. 235.—Mont. Altai.
cuneatum, Ledeb. l. c.—Mont. Altai.
multifidum, Meinsh. in Linnæa, xxx. (1859–60) 515. —Sibir.
simplex, Rupr. Sert. Tianach. 49 = P. stellatum.
tenuifolium, Meinsh. in Linnæa, xxx. (1859–60) 515. —Sibir.

AULACOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 149 = Rhynchothecа, Kuis & Pav. (Geran.).
inerme, Turcz. l. c. 150.—Reg. Andina.

AULAX, Berg. Descr. Pl. Cap. 83 (1767). *PROTEACEÆ*, Benth. & Hook. f. iii. 166.
cneorifolia, *Knight, Prot.* 18.—Afr. austr.
pinifolia, *Berg. Descr. Pl. Cap.* 83.—Afr. austr.
umbellata, R. Br. in Trans. Linn. Soc. x. (1810) 50 = cneorifolia.

AULAXANTHUS, Ell. Sketch, i. 102 (1817) = Anthaenantia, Beauv. (Gramin.).
ciliatus, Ell. l. c. = Anthaenantia villosa.
rufus, Ell. l. c. t. 6 = Anthaenantia rufa.

AULAXIA, Nutt. Gen. Am. i. 47 (1818) = Anthaenantia, Beauv. (Gramin.).
ciliata, Nutt. l. c. = Anthaenantia villosa.
rufa, Nutt. l. c. = Anthaenantia rufa.

AULAXIS, Haw. Enum. Saxif. 46 (1821) = Saxifraga, Linn.
micranthifolia, Haw. l. c. 47 = S. sroa.
nuda, Haw. l. c. = S. sroa.

AULAXIS, Steud. Nom. ed. II. i. 171 (1840) = AULAVIA, Nutt. = Anthaenantia, Beauv. (Gramin.).

AULAVA, Harv. Gen. Fl. Afr. Austr. 349 (1838) = Harveya, Hook. (Scrophul.).
capensis, Harv. l. c. 350 = H. capensis.
coccinea, Harv. Thes. Cap. i. 33.—Afr. austr.
grandiflora, Benth. in DC. Prod. x. 533.—Afr. austr.
obtusifolia, Benth. l. c. = H. obtusifolia.
pauciflora, Benth. l. c. 534.—Afr. austr.
purpurea, Benth. l. c. 533 = H. purpurea.
scarlatina, Benth. l. c. 534.—Afr. austr.
squamosa, Harv. Gen. Fl. Afr. Austr. 350 = H. squamosa.

AULEVA, D. Dietr. Syn. Pl. iii. 589 (1843) = præc.

AULICA, Rafin. Fl. Tellur. iv. 10 (1836) = Hippeastrum, Herb. (Amaryll.).
latifolia, Rafin. l. c. = H. sulicum.
platypetala, Rafin. l. c. = H. sulicum.
striata, Rafin. l. c. = H. sulicum.

AULIPHAS, Rafin. Sylva Tellur. 95 (1838) = Miconia, Ruiz & Pav. (Melast.).
ligustrina, Rafin. l. c. = M. ligustricides.

AULIZA, Salisb. in Trans. Hort. Soc. i. (1812) 294 = Epidendrum, Linn. (Orchid.).
ciliaris, Salisb. l. c. = E. ciliare.

AULOMYRCIA, Berg, in Linnæa, xxvii. (1854) 35 = Myrcia, DC. (Myrtac.).
Abrantes, Berg, l. c. xxx. (1859–60) 650.—Bras.
acutissima, Berg, l. c. 662.—Ind. occ.
aromatica, Berg, in Mart. Fl. Bras. xiv. i. 71.—Bras.
acutata, Berg, l. c. = M. acutata.
acutifolia, Berg, l. c. 80.—Bras.
Aethaea, Berg, l. c. 118.—Bras.
alagoiensis, Berg, l. c. 130 = M. alagoënsis.
alba-tomentosa, Berg, in Linnæa, xxvii. (1854) 62 = M. albo-tomentosa.
alliacea, Berg, l. c. 54.—Bras.
alternifolia, Berg, l. c. 72 = M. alternifolia.
amazonica, Berg, l. c. 41 = M. amazonica.

AULOMYRCIA :—
amethystina, Berg, in Mart. Fl. Bras. xiv. i. 108.—Bras.
andromedoides, Berg, in Linnæa, xxvii. (1854) 40 = Myrcia andromedoides.
andromedoides, Berg, l. c. xxx. (1859–60) 661.—Guiana.
angustifolia, Berg, in Mart. Fl. Bras. xiv. i. 135.—Bras.
anomala, Berg, l. c. 96 = M. anomala.
apiocarpa, Berg, l. c. 64.—Bras.
atropilosa, Berg, l. c. 101.—Bras.
atrovirens, Berg, l. c. 121.—Bras.
auco-lanata, Berg, in Linnæa, xxvii. (1854) 62.—Bras.
badia, Berg, in Mart. Fl. Bras. xiv. i. 147.—Bras.
bahiensis, Berg, l. c. 118.—Bras.
bella, Berg, l. c. 71 = M. bella.
bicarinata, Berg, l. c. 118.—Bras.
bicudensis, Berg, l. c. 857.—Bras.
bifurcata, Berg, l. c. 141.—Bras.
Billardiana, Berg, in Linnæa, xxvii. (1854) 51 = M. Billardiana.
binervinata, Berg, in Mart. Fl. Bras. xiv. i. 115.—Bras.
Blanchetiana, Berg, l. c. 65.—Bras.
bombycina, Berg, l. c. 95.—Bras.
Boryx, Berg, l. c. 116.—Bras.
bracteata, Berg, l. c. 554 = M. bracteata.
breviramis, Berg, l. c. 86.—Bras.
bullata, Berg, l. c. 96 = M. bullata.
buxifolia, Berg, l. c. 80 = M. buxifolia.
buxiusoa, Berg, in Linnæa, xxx. (1859–60) 664.—Guiana.
caerulescens, Berg, in Mart. Fl. Bras. xiv. i. 80.—Bras.
caesia, Berg, l. c. 58.—Bras.
calyptranthoides, Berg, l. c. 67.—Bras.
Camaraxana, Berg, l. c. 88 = M. Camaraxana.
Cambessedeana, Berg, in Linnæa, xxvii. (1854) 40 = M. Cambessedeana.
campestris, Berg, l. c. 74 = M. campestris.
capitata, Berg, in Mart. Fl. Bras. xiv. i. 554 = M. capitata.
capivarkyensis, Berg, in Linnæa, xxvii. (1854) 36 = M. capivarkyensis.
caracasana, Berg, l. c. 53 = M. caracasana.
cardiophylla, Berg, in Mart. Fl. Bras. xiv. i. 108 = M. cardiophylla.
casuinoides, Berg, in Linnæa, xxvii. (1854) 74 = M. casuinoides.
castrensis, Berg, in Mart. Fl. Bras. xiv. i. 111.—Bras.
chapadensis, Berg, l. c. 554.—Bras.
chilensis, Berg, in Linnæa, xxvii. (1854) 38 = M. chilensis.
chrysophylla, Berg, in Mart. Fl. Bras. xiv. i. 125.—Bras.
Clausseniana, Berg, l. c. 116.—Bras.
colina, Berg, in Linnæa, xxvii. (1854) 40.—Bras.
comosa, Berg, in Mart. Fl. Bras. xiv. i. 133.—Bras.
conduplicata, Berg, in Linnæa, xxvii. (1854) 76.—Guiana.
convisa, Berg, l. c. 56.—Bras.
coriacea, Berg, l. c. 70 = M. coriacea.
corymbifera, Berg, l. c. 74.—Bras.
corymbosa, Berg, l. c. 41 = M. corymbosa.
costata, Berg, in Mart. Fl. Bras. xiv. i. 79 = M. costata.
Coumeta, Berg, in Linnæa, xxvii. (1854) 60 = M. Coumeta.
crassifolia, Berg, l. c. 74 = M. crassifolia.
crassifolia, Berg, in Mart. Fl. Bras. xiv. i. 128.—Bras.
crenulata, Berg, l. c. 141.—Bras.
cuneata, Berg, in Linnæa, xxvii. (1854) 78.—Venezuela.
cupreo, Berg, in Mart. Fl. Bras. xiv. i. 77.—Bras.
curvifolioides, Berg, in Linnæa, xxvii. (1854) 55 = M. curvifolioides.
cymosa, Berg, in Mart. Fl. Bras. xiv. i. 562.—Bras.
daphnoides, Berg, in Linnæa, xxvii. (1854) 77 = M. daphnoides.
dealbata, Berg, l. c. 61 = M. dealbata.
debilis, Berg, l. c. 62 = M. debilis.
decrescens, Berg, in Mart. Fl. Bras. xiv. i. 126.—Bras.
densa, Berg, in Linnæa, xxvii. (1854) 37 = Calyptranthes densa.
desertorum, Berg, in Mart. Fl. Bras. xiv. i. 556.—Bras.
detergens, Berg, in Linnæa, xxvii. (1854) 46 = M. detergens.

AULOMYRCIA :—
diaphana, Berg, in Mart. Fl. Bras. xiv. i. 82.—Bras.
dichrome, Berg, in Linnæa, xxvii. (1854) 66.—Guiana.
dictyophlebs, Berg, in Mart. Fl. Bras. xiv. i. 560.—Bras.
dictyophylla, Berg, l. c. 72.—Bras.
dimorpha, Berg, l. c. 101.—Bras.
dinericata, Berg, in Linnæa, xxvii. (1854) 58 = M. divaricata.
Doniana, Berg, in Mart. Fl. Bras. xiv. i. 516 = M. Doniana.
dumosa, Berg, in Linnæa, xxx. (1859–60) 656.—Guiana.
edulis, Berg, l. c. 657.—Guiana.
egensis, Berg, in Mart. Fl. Bras. xiv. i. 99.—Bras.
elaeodendra, Berg, in Linnæa, xxvii. (1854) 76 = M. elaeodendra.
emarginata, Berg, in Mart. Fl. Bras. xiv. i. 134.—Bras.
enumerophylla, Berg, l. c. 98.—Bras.
eumorphoides, Berg, in Linnæa, xxvii. (1854) 79 = M. eximia.
fenestiana, Berg, l. c. 63 = M. Fenestiana.
ferruginea, Berg, in Mart. Fl. Bras. xiv. i. 562 = M. ferruginea.
fragilis, Berg, l. c. 117.—Bras.
Gardneriana, Berg, l. c. 129 = M. Gardneriana.
garopabensis, Berg, in Linnæa, xxvii. (1854) 47 = M. garopabensis.
Gaudichaudiana, Berg, in Mart. Fl. Bras. xiv. i. 88.—Bras.
gigantea, Berg, l. c. 548.—Bras.
glabra, Berg, l. c. 119.—Bras.
glandulosa, Berg, in Linnæa, xxvii. (1854) 79.—Bras.
glauca, Berg, l. c. 50 = M. glauca.
glaucescens, Berg, in Mart. Fl. Bras. xiv. i. 81.—Bras.
goyazensis, Berg, l. c. 85 = M. goyazensis.
grandiflora, Berg, l. c. 118.—Bras.
grandifolia, Berg, l. c. 97 = M. grandifolia.
hepatica, Berg, l. c. 122.—Bras.
hirtiflora, Berg, in Linnæa, xxvii. (1854) 40 = M. hirtiflora.
holosericea, Berg, in Mart. Fl. Bras. xiv. i. 99.—Bras.
Hostmanniana, Berg, in Linnæa, xxvii. (1854) 42.—Guiana.
hypericoides, Berg, l. c. 62 = M. hypericoides.
imbricata, Berg, l. c. 61 = M. imbricata.
insularis, Berg, in Mart. Fl. Bras. xiv. i. 96 = M. insularis.
intermedia, Berg, l. c. 107.—Bras.
Yacquiniana, Berg, in Linnæa, xxvii. (1854) 69 = M. coriacea.
jequitinhonhensis, Berg, in Mart. Fl. Bras. xiv. i. 137.—Bras.
Klotzschiana, Berg, l. c. 111 = M. Klotzschiana.
lacinioso, Berg, l. c. 545.—Bras.
lancea, Berg, l. c. 86.—Bras.
lanciflora, Berg, in Linnæa, xxx. (1859–60) 658.—Guiana.
langsdorffii, Berg, l. c. 102 = M. lanuginosa.
laricina, Berg, l. c. 61.—Bras.
Larrotiana, Berg, in Linnæa, xxvii. (1854) 58 = M. Larrotiana.
Laureola, Berg, in Mart. Fl. Bras. xiv. i. 110.—Bras.
laurifera, Berg, in Linnæa, xxvii. (1854) 64 = M. latrifera.
laurifolia, Berg, l. c. = M. laurifolia.
laxiflora, Berg, in Mart. Fl. Bras. xiv. i. 134 = M. laxiflora.
leptoclada, Berg, in Linnæa, xxvii. (1854) 60 = M. leptoclada.
Leucadendron, Berg, l. c. 64 = M. Leucadendron.
leucanthe, Berg, in Mart. Fl. Bras. xiv. i. 93.—Bras.
linearifolia, Berg, in Linnæa, xxvii. (1854) 35 = M. linearifolia.
linoeta, Berg, in Mart. Fl. Bras. xiv. i. 63.—Bras.
Lingua, Berg, in Linnæa, xxvii. (1854) 77.—Bras.
linguaeformis, Berg, in Mart. Fl. Bras. xiv. i. 125.—Bras.
longipes, Berg, in Linnæa, xxvii. (1854) 55.—Bras.
lucida, Berg, in Mart. Fl. Bras. xiv. i. 118.—Bras.
macrocarpa, Berg, in Linnæa, xxvii. (1854) 41 = M. macrocarpa.
macrophylla, Berg, in Mart. Fl. Bras. xiv. i. 99 = M. macrophylla.
Manaosi, Berg, l. c. 131.—Bras.

AULOMYRCIA :—
Maraguana, Berg, in Mart. Fl. Bras. xiv. 1. 120 = Myrcia Maraguana.
maritima, Berg. l. c. 553.—Bras.
Martiana, Berg, l. c. 133 = M. Martiana.
Matheuziana, Berg, in Linnæa, xxvii. (1854) 45.—Peruv.
micrantha, Berg, in Mart. Fl. Bras. xiv. 1. 516.—Bras.
microcarpa, Berg, l. c. 61 = M. microcarpa.
microphylla, Berg, in Linnæa, xxvii. (1854) 56.—Bras.
montana, Berg, l. c. 36 = M. montana.
mosigensis, Berg, l. c. = M. mosigensis.
multiflora, Berg, l. c. 47 = M. multiflora.
mutabilis, Berg, in Mart. Fl. Bras. xiv. 1. 70.—Bras.
myrtifolia, Berg, in Linnæa, xxvii. (1854) 36 = M. myrtifolia.
Neesiana, Berg, l. c. 36 = M. Neesiana.
nigrescens, Berg, l. c. 71 = M. nigrescens.
nigricans, Berg, in Mart. Fl. Bras. xiv. 1. 101.—Bras.
nigro-punctata, Berg, l. c. 116.—Bras.
nivea, Berg, in Linnæa, xxvii. (1854) 61 = M. nivea.
oblongata, Berg, l. c. 37 = M. oblongata.
obovata, Berg, l. c. 72.—Bras.
obscura, Berg, in Mart. Fl. Bras. xiv. 1. 132.—Bras.
obtecta, Berg, l. c. 117.—Bras.
obtusa, Berg, in Linnæa, xxvii. (1854) 66 = M. obtusa.
obtusata, Berg, l. c. 39.—Bras.
orthophylla, Berg, in Mart. Fl. Bras. xiv. 1. 549.—Bras.
Ottonis, Berg, in Linnæa, xxvii. (1854) 55.—Guiana.
ovalifolia, Berg, in Mart. Fl. Bras. xiv. 1. 81.—Bras.
ovalis, Berg, l. c. 107.—Bras.
pachyclada, Berg, l. c. 133.—Bras.
pallens, Berg, in Linnæa, xxvii. (1854) 73 = M. pallens.
pallida, Berg, in Mart. Fl. Bras. xiv. 1. 87.—Bras.
panicularis, Berg, l. c. 54.—Bras.
paniculata, Berg, in Linnæa, xxvii. (1854).50.—Bras.
paratensis, Berg, in Mart. Fl. Bras. xiv. 1. 76.—Bras.
parnahibensis, Berg, l. c. 135.—Bras.
pauciflora, Berg, l. c. 560 = M. pauciflora.
perforata, Berg, l. c. 88.—Bras.
pertusa, Berg, in Linnæa, xxvii. (1854) 47 = M. pertusa.
piauhiensis, Berg, in Mart. Fl. Bras. xiv. 1. 87.—Bras.
pilantha, Berg, l. c. 555.—Bras.
pinifolia, Berg, in Linnæa, xxvii. (1854) 35 = M. pinifolia.
pirarensis, Berg, l. c. 41.—Guiana.
platyclada, Berg, in Mart. Fl. Bras. xiv. 1. 545 = M. platyclada.
plumbea, Berg, l. c. 142.—Bras.
Poeppigiana, Berg, in Linnæa, xxvii. (1854) 78.—Bras.
Pohliana, Berg, in Mart. Fl. Bras. xiv. 1. 73.—Bras.
polyantha, Berg, l. c. 83 = M. Kunthiana.
polymorpha, Berg, in Linnæa, xxvii. (1854) 46.—Bras.
pruinosa, Berg, in Mart. Fl. Bras. xiv. 1. 114.—Bras.
prunifolia, Berg, in Linnæa, xxvii. (1854) 55 = M. prunifolia.
puberula, Berg, l. c. 54 = M. puberula.
pubiflora, Berg, l. c. 40 = M. pubiflora.
pulchra, Berg, in Mart. Fl. Bras. xiv. 1. 68.—Bras.
punctata, Berg, l. c. 546 = M. punctata.
pusilla, Berg, l. c. 140.—Bras.
pyrifolia, Berg, in Linnæa, xxvii. (1854) 44.—Bras.
racemosa, Berg, l. c. 82.—Bras.
ramuliflora, Berg, in Mart. Fl. Bras. xiv. 1. 64.—Bras.
ramulosa, Berg, in Linnæa, xxvii. (1854) 36 = M. ramulosa.
Regeliana, Berg, in Mart. Fl. Bras. xiv. 1. 557.—Bras.
reticulata, Berg, l. c. 108 = M. reticulata.
reticulosa, Berg, in Linnæa, xxvii. (1854) 63 = M. reticulosa.
retusa, Berg, in Mart. Fl. Bras. xiv. 1. 142.—Bras.
Richardiana, Berg, in Linnæa, xxx. (1859–60) 664.—Bras.
Riedeliana, Berg, in Mart. Fl. Bras. xiv. 1. 551.—Bras.
roraimensis, Berg, in Linnæa, xxvii. (1854) 68.—Guiana.
rordis, Berg, in Mart. Fl. Bras. xiv. 1. 559.—Bras.
rostrata, Berg, l. c. 544 = M. rostrata.
rosulana, Berg, in Linnæa, xxvii. (1854) 54.—Bras.
rotundifolia, Berg, in Mart. Fl. Bras. xiv. 1. 128.—Bras.

AULOMYRCIA :—
rubella, Berg, in Linnæa, xxvii. (1854) 40 = M. rubella.
rufa, Berg, in Mart. Fl. Bras. xiv. 1. 65.—Bras.
rufipes, Berg, in Linnæa, xxvii. (1854) 77 = M. rufipes.
rugosa, Berg, in Mart. Fl. Bras. xiv. 1. 110.—Bras.
Sagraea, Berg, in Linnæa, xxx. (1859–60) 655.—Cuba.
salicifolia, Berg, l. c. xxvii. (1854) 78 = M. salicifolia.
Salzmanni, Berg, in Mart. Fl. Bras. xiv. 1. 116.—Bras.
Schomburghiana, Berg, in Linnæa, xxvii. (1854) 75.—Bras.
Schrankiana, Berg, l. c. 79 = M. Schrankiana.
scrobiculata, Berg, in Mart. Fl. Bras. xiv. 1. 157 = M. scrobiculata.
Sonderiana, Berg, l. c. 110.—Bras.
spathulata, Berg, l. c. 91.—Bras.
sphærocarpa, Berg, in Linnæa, xxvii. (1854) 51 = M. sphærocarpa.
sphæroides, Berg, in Mart. Fl. Bras. xiv. 1. 68.—Bras.
Spruceana, Berg, l. c. 76.—Bras.
stenophylla, Berg, in Linnæa, xxx. (1859–60) 654.—Paraguay.
stictophylla, Berg, in Mart. Fl. Bras. xiv. 1. 67.—Bras.
stricta, Berg, l. c. 548.—Bras.
suaveolens, Berg, in Linnæa, xxvii. (1854) 78 = M. suaveolens.
subalpestris, Berg, l. c. 73 = M. subalpestris.
subcoriacea, Berg, in Mart. Fl. Bras. xiv. 1. 69.—Bras.
subcordata, Berg, in Linnæa, xxvii. (1854) 62 = M. subcordata.
subobliqua, Berg, l. c. 57.—Guiana.
subverticillaris, Berg, l. c. 78.—Bras.
sufruticosa, Berg, in Mart. Fl. Bras. xiv. 1. 136.—Bras.
supranitillaris, Berg, l. c. 69.—Bras.
surinamensis, Berg, in Linnæa, xxvii. (1854) 64 = M. surinamensis.
tenuiflora, Berg, in Mart. Fl. Bras. xiv. 1. 68.—Bras.
terrifolia, Berg, l. c. 134.—Bras.
thyrsiflora, Berg, in Linnæa, xxvii. (1854) 74.—Bras.
torta, Berg, l. c. 78 = M. torta.
tortuosa, Berg, in Mart. Fl. Bras. xiv. 1. 556.—Bras.
trichanthera Wawra, in Oestr. Bot. Zeitschr. xxix. (1879) 215.—Bras.
trifora, Berg, in Linnæa, xxvii. (1854) 79 = M. trifora.
trifolia, Berg, in Mart. Fl. Bras. xiv. 1. 107.—Bras.
undulata, Berg, l. c. 89.—Bras.
vaccinifolia, Berg, l. c. 140.—Bras.
variabilis, Berg, in Linnæa, xxvii. (1854) 69 = M. variabilis.
Vauthieriana, Berg, l. c. xxx. (1859–60) 655 = M. Vauthieriana.
velutina, Berg, in Mart. Fl. Bras. xiv. 1. 560.—Bras.
venulosa, Berg, in Linnæa, xxvii. (1854) 63 = M. venulosa.
vestita, Berg, l. c. 74 = M. vestita.
virgata, Berg, in Mart. Fl. Bras. xiv. 1. 136 = M. virgata.
Widgreniana, Berg, l. c. 70.—Bras.
Wullschlaegeliana, Berg, in Linnæa, xxvii. (1854) 48.—Guiana.

AULONEMIA, Gondot, in Ann. Sc. Nat. Sér. III. v. (1846) 76. t. 5 = **Arthrostylidium**, Rupr. (Gramin.).
Gaudichaudiana, Gondot, l. c.—N. Granat.

AULONIX, Rafin. Sylva Tellur. 25 (1838) = **Cytisus**, Linn. (Leguminos.).
biflorus, Rafin. l. c. = C. biflorus.

AULOSEMA, Walp. Rep. i. 694 (1842) = **Astragalus**, Linn. (Leguminos.).
eximium, Walp. l. c. = Astr. hypogeous.

AURANTIUM, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = Citrus, Linn. (Rutac.).
acre, Mill. l. c. ed. VIII. n. 1 = C. Aurantium.
decumanum, Mill. l. c. n. 4 = C. decumana.
humile, Mill. l. c. n. 5 = C. humilis.
myrtifolium, Descourt. Fl. Med. Antill. iii. 308. t. 219 = C. Aurantium.
orientale, Mill. Gard. Dict. ed. VIII. n. 3 = C. Aurantium.
silvestre, Prits. Ic. Ind. 137 = C. Aurantium.
sinense, Mill. Gard. Dict. ed. VIII. n. 2 = C. Aurantium.

AURELIA, Cass. Bull. Soc. Philom. (1815) 175; et in Dict. Sc. Nat. xxxvii. 468 (1816) = Grindelia, Willd. (Compos.).
amplexicaulis, Cass. l. c. = G. squarrosa.
discoidea, Cass. l. c. = G. glutinosa.

AURELIA :—
glutinosa, Cass.—Cf. Bull. Soc. Philom. (1815) 175 = Grindelia glutinosa.
squarrosa, Cass. ex Steud. Nom. ed. II. i. 171 = G. squarrosa.

AURELIA, J. Gay, in Ann. Sc. Nat. Sér. IV. x. (1858) 95 = **Narcissus**, Tourn. (Amaryll.).
Broussonetii, J. Gay, l. c. 96 = N. Broussonetii.

AURELIANA, Lafit. ex Catesb. Nat. Hist. Carol. App. 16 (1764) = **Aralia**, Linn.

AURELIANA, Sendt. in Mart. Fl. Bras. x. 188. t 19. (1846) = **Bassovia**, Aubl. (Solan.).
fasciculata, Sendt. l. c. 140 = B. fasciculata.
glomuliflora, Sendt. l. c. 185.—Bras.
lucida, Sendt. l. c. 189.—Bras.
tomentosa, Sendt. l. c. 140 = B. tomentosa.
velutina, Sendt. l. c. = B. velutina.

AUREOLARIA, Rafin. New Fl. Am. ii. 58 (1836) = Gerardia, Linn. (Scrophul.).
cuneifolia, Rafin. l. c. 59 = Herpestis pubescens ?
glauca, Rafin. l. c. 60 = G. quercifolia.
laevigata, Rafin. l. c. 59 = G. laevigata.
reticulata, Rafin. l. c.—Am. bor.
respatria, Rafin. l. c. 60 = G. repatria.
villosa, Rafin. l. c. 59 = G. flava.

AURICULA, Tourn. ex Spach, Hist. Vég. Phan. ix. 356 (1840) = Primula, Tourn.
crenata, Spach, l. c. 358 = P. marginata.
glutinosa, Spach, l. c. 359 = P. glutinosa.
hortensis, Spach, l. c. 357 = P. Auricula.
integrifolia, Spach, l. c. 359 = P. integrifolia.
minima, Spach, l. c. 360 = P. minima.
Palinuri, Spach, l. c. 357 = P. Palinuri.
villosa, Spach, l. c. 358 = P. minima.

AURILA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1, nomen (Quid ?).

AURINIA, Desv. Journ. Bot. iii. (1813) 162 = **Alyssum**, Linn. (Crucif.).
capensis, Desv. l. c. = Heliophila diffusa.
corymbosa, Griseb. Spicil. Fl. Rumel. i. 271 = Al. corymbosum.
edentula, Fisch. Mey. & Trautv. Ind. Sem. Hort. Petrop. iii. 4 = Al. gemonense.
gemonensis, Griseb. Spicil. Fl. Rumel. i. 272 = Al. gemonense.
media, Schur, Enum. Pl. Transs. 61 = Al. gemonense.
obovata, Desv. Journ. Bot. iii. (1813) 161 = Al. obovatum.
orientalis, Griseb. Spicil. Fl. Rumel. i. 272 = Al. orientale.
petraeum, Fuss, Fl. Transs. 64 = Al. gemonense.
saxatilis, Desv. Journ. Bot. iii. (1813) 162 = Al. saxatile.
saxatilis, Griseb. Spicil. Fl. Rumel. i. 272 = Al. orientale.
sinuata Griseb. l. c. 271 = Al. sinuatum.

AUROTA, Rafin. Fl. Tellur. iii. 61 (1836) = **Curculigo**, Gaertn. (Amaryll.).
latifolia, Rafin. l. c. = C. latifolia.

AURORA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1 = **Quisqualis**, Linn. (Combret.).

AUSTERIUM, Poit. ex DC. Prod. ii. 336 (1825) = **Rhynchosia**, Lour. (Leguminos.).

AUSTRALINA, Gandich. in Freyc. Voy. Bot. 505 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 304.
ANAGANTHOS, Hook. f. Fl. N. Zel. i. 226 (1854).
DIDYMOTOCA, E. Mey. Zwei Pfl. Docum. 178 (1843).
acuminata, *Wedd. in Ann. Sc. Nat. Sér.* IV. i. (1854) 219.—Ar. austr.
capensis, *Wedd. l. c.*—Ar. austr.
flaccida, Wedd. in Arch. Mus. Par. viii. (1855–56) 546. t. 20 C.—Schizoparias.
integrifolia, *Wedd. in Ann. Sc. Nat. Sér.* IV. i. (1854) 215.—Ar. austr.
Muelleri, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 545.—Austral.
Novae-Zelandiae, Hook. f. Fl. N. Zel. i. 226 = pusilla.

AUSTRALINA:—
pusilla, *Gaudich. in Freyc. Voy. Bot.* 505.—Austral. ; N. Zel.
Schlomeriana, *Wedd. in Ann. Sc. Nat. Sér.* IV. i. (1854) 213.—Abyss.
tasmanica, Hook. f. Fl. N. Zel. i. 236 = pusilla.

AUSTROBUXUS, Miq. Fl. Ind. Bat. Suppl. 444 (1860) = **Sarcococca**, Lindl. (Euphorb.).
nitidus, Miq. l. c.—Sumatra.

AUTOGENES, Rafin. Fl. Tellur. iv. 30 (1836) = **Narcissus**, Tourn. (Amaryll.).
angustifolius, Rafin. l. c. = N. poeticus.
biflorus, Rafin. l. c. = N. biflorus.
poeticus, Rafin. l. c. = N. poeticus.
tenuior, Rafin. l. c. = N. graecilis.

AUXEMMA, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 23. t. 5. *BORAGINEAE*, Benth. & Hook. f. ii. 840.
Gardneriana, *Miers, l. c.* 24. t. 5.—Bras.

AVELLANITA, Phil. in Linnaea, xxxiii. (1864–65) 257. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 289.
Bustillosi, *Phil. l. c.* 288.—Chili.

AVELLINIA, Parl. Fl. Nov. 59 (1842). *GRAMINEAE*, Benth. & Hook. f. iii. 1184.
Michelii, *Parl. l. c.*—Italia.
Savi, *Parl. l. c.* 61.—Italia.
scabriuscula, Nym. Consp. 815 = Koeleria scabriuscula.
tenuicula, *Nym. l. c.*—Hispan.

AVENA, Hall. ex Scop. Introd. 74 (1777) = **Agrostis**, Linn. (Gramin.).

AVENA, Linn. Syst. ed. I. (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1160.
ELICTOTRICHON, Bess. ex Andrz. Rys Botan. i. (1823) 9.
HELICOTRICHUM, Bess. in Reichb. Fl. Germ. Excurs. 140*. n. 869 (1832).
HEUFFELIA, Schur, Enum. Pl. Transs. 760 (1866).
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 415.—Abyss.
adsurgens, *Schur, Enum. Pl. Transs.* 762.—Hungar.
affinis, Bernh. ex Steud. Nom. ed. II. i. 171, 173 = sterilis.
agraria, Brot. Fl. Lusit. i. 105 = strigosa.
agraria-mutica, Brot. l. c. 106 = strigosa.
agraria-sesquitalera, Brot. l. c. = strigosa.
agropyroides, *Boiss. Diagn.* Ser. I. xiii. 50.—Graecia.
agrostidea, Fries, Mant. iii. 3 = Trisetum agrostideum.
agrostoides, Griseb. Spicil. Fl. Rumel. ii. 454 = Trisetum nyriauthum.
airoides, Koel. Gram. 208 = Trisetum subspicatum.
airoides, Pourr. ex Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 290 = Pourretii.
alba, *Vahl, Symb. Bot.* ii. 94.—Gall. austr. ; Afr. bor.
albinervis, *Boiss. Voy. Bot. Espagne*, ii. 656.—Hispan.
Alopecurus, Roth, Catalecta, iii. 20 = Trisetum Alopecurus.
alpestris, DC. Fl. Fr. vi. 960 = Trisetum pratense.
alpestris, Host, Gram. Austr. iii. 27. t. 39 = T. alpestre.
alpina, [Honck.] Vers. Gew. Teutschl. 302 = Aira montana !
alpina, Latour. Chlor. Lugd. 3 = Schenzeri.
alpina, Sm. in Trans. Linn. Soc. x. (1811) 335 = pratensis.
alpina, Trin. Fund. Agrost. 157 = Deschampsia alpina.
alta, Cav. ex Roem. & Schult. Syst. ii. 691 = strigosa.
ambigua, Schoenh. ex Nym. Consp. 810 = fatua.
amethystina, *DC. Fl. Fr.* iii. 56 ; vi. 960.—Gallia.
andropogoides, *Steud. in Flora*, xii. (1829) ii. 496.—Afr. austr.
angulica, Hort. ex Roem. & Schult. Syst. ii. 669 = sativa.
antarctica, Roem. & Schult. l. c. 676 = Trisetum antarcticum.
antarctica, Thunb. Prod. Fl. Cap. 22 = Danthonia antarctica.

AVENA:—
ardurensis, Lej. ex Steud. Nom. ed. II. i. 171 = strigosa.
arenaria, Spreng. Syst. i. 333 = Trisetum arenarium.
argaea, *Boiss. Fl. Orient.* v. 646.—Cappadoc.
argentiodeum, Schur, in Oestr. Bot. Zeitschr. x. (1860) 74 = Trisetum alpestre.
argentea, Willd. Enum. Hort. Berol. 125 = Trisetum argenteum.
argiuensis, *Steud. Syn. Pl. Gram.* 233.—Chili.
aristidoides, Steud. in Flora, xii. (1829) ii. 481. n. 23 = Danthonia Thunbergii.
aristoides, *Thunb. Prod. Pl. Cap.* 22.—Afr. austr.
arundinacea, Delile, Fl. Egypte, 171. t. 19 = Danthonia Forskohlei.
arvensis, Salisb. Prod. 28 = Bromus arvensis.
aspera, *Munro, ex Thw. Enum. Pl. Zeyl.* 372.—Zeylan.
atherantherea, Presl, Cyp. et Gram. Sic. 80 = barbata.
atropurpurea, Link, Hort. Berol. i. 119 = Aira atropurpurea.
aurata, All. Fl. Pedem. ii. 255 = setacea, Agrostis alpina.
australis, Parl. Fl. Ital. i. 285 = pratensis.
barbata, *Brot. Fl. Lusit.* i. 108.—Reg. Mediterr. ; As. Min. ; Arab.
barbata, *Pott, ex Link, in Schrad. Journ.* ii. (1799) 315.—Lusitan.
barregensis, Nym. Consp. 818 = Trisetum agrostideum.
Baumgartenii, *Steud. Syn. Pl. Gram.* 233.—Hungar.
Bellardi, *Colla, Herb. Pedem.* t. 93. f. 2.—Italia.
benghalensis, *Kunth, ex Steud. Nom.* ed. I. 94.—Ind. or.
Besseri, *Ledeb. Fl. Ross.* iv. 415.—Ross. med.
bifida, *Banon. Agrost.* 158, nomen.—Hab. !
biparitia, Link, Hort. Berol. i. 118 = Danthonia semiannularis.
Biasoi, Aschers. & Janka.—Cf. Aschers. & Kan. Cat. Coron. Serb. 9 = compressa.
borbonia, *Steud. Syn. Pl. Gram.* 234.—Ins. Borbon.
brevifolia, Host, Gram. iii. t. 40 = Trisetum distichophyllum.
brevifolia, Spreng. Syst. i. 334 = Trisetum rigidum.
brevis, *Roth, Tent. Fl. Germ.* i. 40.—Europ.
bromoides, *Gouan, Hort. Monsp.* 53.—Reg. Mediterr.
bromoides, Mert. & Koch, in Roehl. Fl. Deutschl. ed. III. i. 570 = pratensis.
Brownei, Spreng. Syst. i. 336 = Danthonia pallida.
Brubnasiana, *Gruner, in Bull. Soc. Nat. Mosc.* xli. (1867) ii. 458.—Reg. Casp.
bulbosa, Willd. in Ges. Naturf. Fr. Berl. Neue Schr. ii. (1799) 116. t. 5. f. 1 = Arrhenatherum avenaceum.
Burnoufii, Nym. Syll. 418 = Trisetum Burnoufii.
byzantina, C. Koch, in Linnaea, xxi. (1848) 392 = latifolia.
caespitosa, Kuntze, Taschen-Fl. Leipz. 45 = Deschampsia caespitosa.
calcicina, Vill. Hist. Pl. Dauph. ii. 148 = Danthonia provincialis.
callosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxix. (1856) i. 36.—Sibir.
calycina, Lam. Illustr. i. 900 = Danthonia calycina.
calycina, Willd. Sp. Pl. i. 458, in syn. = Danthonia provincialis.
canariensis, *Ness, ex Steud. Nom.* ed. II. i. 171.—Ins. Teneriff.
Candollei, Nym. Consp. 819 = Trisetum pratense.
canescens, [Weber, in] Wigg. Prim. Fl. Hols. 9 = Corynephorus canescens.
cantabrica, Lag. in Herrer. Agr. 141 = filifolia.
capensis, Burm. f. Fl. Cap. Prod. 3 = Anthistiria ciliata.
capensis, Linn. f. Suppl. 119 = Aristida capensis.
capensis, Steud. in Flora, xii. (1829) 481 = Danthonia papillosa.
capillacea, *Beauv. Agrost.* 153, nomen.—Hab. !
capillaris, *Mert. & Koch, in Roehl. Deutschl. Fl.* ed. III. i. 578.—Europ.
Carmeli, *Boiss. Diagn.* Ser. I. xiii. 50.—Syria.
carolinensis, Walt. Fl. Carol. 81 = Trisetum palustre.
carpatica, Host, Gram. Austr. iv. 18. t. 31 = Trisetum carpathicum.
caryophyllea, *Sibth. & Sm. Fl. Graec. Prod.* i. 67.—Archip. Graec.
caryophyllea, [Weber, in] Wigg. Prim. Fl. Hols. 10 = Aira caryophyllea.
Cassubitensis, Host, ex Roem. & Schult. Syst. ii. 691 = strigosa.

AVENA:—
Cavanillesii, Koch, Syn. Fl. Germ. ed. I. 797 = Trisetum Cavanillesii.
Cavanillesii, Lag. in Varied. Cienc. ii. iv. (1806) 89 = Stipa avenacea.
cernua, Kunth, Enum. Pl. i. 306 = Trisetum cernuum.
chinensis, Fisch. ex Roem. & Schult. Syst. ii. 669 = nuda.
chlorantha, Link, in Linnaea, xvii. (1843) 401 = Trisetum aureum.
ciliaris, Kit. ex Schult. Oestr. Fl. ed. II. i. 268 = Trisetum carpathicum.
cinerea, Auct. ex Roem. & Schult. Syst. ii. 669 = sativa.
clauda, *Dur. in Duch. Rev. Bot.* i. (1845–46) 360.—Gracia ; Aa. Min. ; Afr. bor.
coarctata, Desf. Cat. Hort. Par. ed. III. (1829) 22 = Gaudinia geminiflora.
colorata, Steud. in Flora, xii. (1829) ii. 481. n. 11 = Danthonia tortuosa.
compacta, *Boiss. & Heldr. Diagn.* Ser. I. vii. 122.—Gracia ; Servia.
compressa, *Heuff. in Flora*, xviii. (1835) 244.—Hungar.
condensata, Link, Enum. Hort. Berol. i. 89 = Trisetum aureum, *Tenore*.
convoluta, *Presl, Cyp. et Gram. Sic.* 31.—Europ. austr. or.
copaimbensis, Roem. & Schult. Syst. ii. 677 = Hierochloë utriculata.
cornensis, *Schur, Enum. Pl. Transs.* 763.—Transylv.
corymbosa, *Nym. Consp.* 814.—Corsica.
cristata, Auct. ex Roem. & Schult. Syst. ii. 758 = Agropyron cristatum.
Cupaniana, *Nym. Syll.* 414.—Ital. ; Creta.
cuspidata, Willd. ex Kunth, Enum. Pl. i. 289 = Calamagrostis cuspidata.
danuvii, Boiss. Diagn. Ser. I. vii. 123 = Trisetum rigidum.
decora, *Janka, in Termész. Füzetek*, viii. (1884) 24.—Europ.
depauperata, Sieber, ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 68 = Danthonia provincialis.
descrtorum, *Less. in Linnaea*, ix. (1834) 208.—Sibir.
desvauxioides, H. B. & K. Nov. Gen. et Sp. i. 147 = Deyeuxia triflora.
diepermis, Mill. Gard. Dict. ed. VIII. n. 1 = sativa.
distans, *Schur, in Oestr. Bot. Zeitschr.* xx. (1870) 22.—Europ.
distichea, Lam. Encyc. i. 330 = Trisetum rigidum.
distichophylla, Host, Gram. Austr. iii. 39. t. 58 = Trisetum argenteum.
distichophylla, Vill. Prosp. 16 ; Pl. Dauph. ii. 144 = Trisetum distichophyllum, rigidum.
dubia, Leers, Fl. Herborn. 41 = Gaudinia tenuis.
Dufourei, Nym. Consp. 813 = Trisetum Dufourei.
dura, Salisb. Prod. 29 = Festuca durinscula.
elata, *Forsk. Fl. Aegypt. Arab.* 19.—Constantinop.
elata, Salisb. Prod. 28 = Arrhenatherum avenaceum.
elatior, Linn. Sp. Pl. 79 = Arrhenatherum elatius.
elegans, Aschers. in Verh. Bot. Ver. Brand. iii.-iv. (1861–62) 279 = capillaris.
elephantina, Spreng. ex Steud. Nom. ed. II. i. 172 = andropogoides.
elephantina, Thunb. Prod. Pl. Cap. 23 = Danthonia elephantina.
elongata, H. B. & K. Nov. Gen. et Sp. i. 148 = Trisetum elongatum.
crianthe, *Dur. in Duch. Rev. Bot.* i. (1845–46) 360.—Afr. bor.
erianthe, *Hsch. in Oestr. Bot. Zeitschr.* xxvii. (1877) 192.—Hispan.
fallax, De Not. Ind. Sem. Hort. Reg. Genuens. 24 = sempervirens.
fallax, Pollini, Fl. Veron. iii. 778 = pubescens.
fallax, Roem. & Schult. Syst. ii. 692 = montana.
fallax, Tenore, Fl. Nap. iii. 98 = convoluta.
fatua, *Linn. Sp. Pl.* 80.—Europ. ; Orient ; As. bor.
fatua, Schreb. Beschr. Gras. 106. t. 18 = sterilis.
fatua, Vis. Fl. Lib. Spec. 6 = barbata.
fertilis, All. Auct. Pedem. 45 = Gaudinia tenuis.
festucaeformis, *Hochst. in Flora*, xxxviii. (1855) 278.—Afr. or.
filifolia, *Lag. Gen. & Sp. Nov.* 4.—Hispan.
filiformis, Forst. f. Prod. 9 = Deyeuxia Forsteri.
filiformis, Labill. Nov. Holl. Pl. i. 24. t. 31 = Deyeuxia filiformis.
flava, Hort. ex Roem. & Schult. Syst. ii. 669 = sativa.

AVENA:—
flavescens, Bont. ex Willk. & Lange, Prod. Fl. Hisp. i. 79 = Trisetum gallecicum.
flavescens, Linn. Sp. Pl. 80 = Trisetum pratense.
flexuosa, Mert. & Koch, in Roehl. Deutschl. Fl. ed. III. i. 570 = Deschampsia flexuosa.
flexuosa, *Schrank, in Denkschr. Reg. Bot. Ges.* ii. 7.—Labrador.
Forskoli, Vahl, Symb. Bot. ii. 35 = Danthonia Forskohlei.
Forsteri, Kunth, Rev. Gram. i. 104 = Deyeuxia Forsteri.
fragilis, Linn. Sp. Pl. 80 = Gaudinia fragilis.
Franzi, Orteg. [ex Spreng.] Bot. Gart. Halle, 14 = strigosa.
fusca, Ard. in Saggi Acc. Padov. ii. (1789) t. 4 = sativa.
fusca, Kit. ex Schult. Fl. Oestr. ed. II. i. 368 = Trisetum carpathicum.
fusca, *Schur, Enum. Pl. Transs.* 756.—Hungar.
fuscoflora, Schur, l. c. = fusca.
gallecica, Nym. Syll. Suppl. 71 = Trisetum gallecicum.
geminiflora, Kunth, Rev. Gram. 108, 299. t. 61 = Gaudinia geminiflora.
Georgiana, Roem. & Schult. Syst. ii. 669 = sativa.
georgica, Boiss. olim ex Fl. Orient. v. 540 = Ventenata macra.
georgica, Zuccag. in Roem. Collect. 126 = sativa.
gigantea, Sallsb. Prod. 28 = Bromus giganteus.
glabra, C. Koch, in Linnaea, xix. (1847) 5 = pubescens.
glabrata, Brot. ex Steud. Nom. ed II. i. 173 = corymbosa.
glacialis, Boiss. Elench. 87; Voy. Bot. Espagne, ii. 654 = Trisetum glaciale.
glauca, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 20 = Schenzeri.
glomerata, Steud. in Flora, xii. (1829) ii. 483 = Danthonia curvifolia.
glumosa, *Ell. Sketch.* ii.—Am. bor.
glumosa, Michx. Fl. Bor. Am. i. 72 = Danthonia spicata.
gracilis, Moris, Stirp. Sard. Elench. fasc. i. 50 = Trisetum Moriali.
Heidreichii, Parl. Fl. Ital. i. 275 = convoluta.
heteromalla, Moench, Meth. 196 = orientalis.
hexantha, Steud. in Flora, xii. (1829) ii. 487. n. 87 = Danthonia stricta.
hirsuta, Moench, Meth. Suppl. 64; Roth, Catalecta, iii. 19 = barbata.
hirta, *Schrad. in Goett. Gel. Ans.* (1821) iii. 2075.—Afr. austr.
hirtifolia, Boiss. Diagn. Ser. II. iv. 128 = pubescens.
hirtula, Lag. Gen. et Sp. Nov. 4 = barbata.
hispanica, Ard. in Saggi Acc. Padov. ii. (1789) 112 = sativa.
hispanica, Hort. ex Roem. & Schult. Syst. ii. 691 = strigosa.
hispanica, Lange, in Kjoeb. Vidensk. Meddel. (1860) 41 = Arrhenatherum erianthum.
hispida, Linn. f. Suppl. 111 = Tristachya leucothrix.
hispida, Nym. Syll. Suppl. 71 = Trisetum hispidum.
hybrida, Peterm. ex Reichb. Fl. Saxon. 17 = fatua.
Hoppeana, *Schaefe, in Flora*, xxvii. (1844) 57.—Ikeria.
Hostii, Boiss. & Rent. Pugill. 121 = Parlatorii.
Hugeniini, *De Not. ex Steud. Syn. Pl. Gram.* 425.—Italia.
hungarica, *Luce, Topogr. Nachr. Oesel*, 30.—Hungar.
hydrophila, F. Muell. ex Hook. f. Fl. Tasm. i. 121 = Amphibromus Neesii.
intermedia, Lindgren, in Bot. Notiser (1841), 151 = fatua.
intermedia, Guss. Fl. Sic. Prod. Suppl. 16 = corymbosa.
involucrata, Schrad. in Goett. Gel. Ans. (1821) iii. 2075 = Danthonia involucrata.
involuta, Presl, ex Steud. Nom. ed. II. i. 173 = semipervirens.
japonica, Steud. Syn. Pl. Gram. 231 = hirsuta.
Junghuhnii, *Buese, in Miq. Pl. Yungh.* 345.—Java.
Kotschyi, Spreng. Syst. i. 336 = Aristida arundinacea.
Kotschyi, *Steud. Syn. Pl. Gram.* 235.—Persia.
lachnantha, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 287.—Afr. trop.
lanuginosa, Nym. Syll. Suppl. 71 = Trisetum leconteum.
lanta, Sallsb. Prod. 28 = Brachypodium pinnatum.
lanuginosa, Schur, in Oestr. Bot. Zeitschr. x. (1860) 72 = pubescens.
lanuta, Hook. l. c. xxvii. (1877) 46 = albinervis

AVENA:—
lanata, Koel. Gram. 308 = Holcus lanatus.
lanata, Schrad. in Goett. Gel. Ans. (1821) iii. 2075 = Danthonia lanata.
lanuginosa, Gilib. Exercit. ii. 539 = fatua.
lasiantha, Link, in Linnaea, ix. (1834) 135 = Bromus intermedius.
latifolia, Host, Gram. Austr. iv. 19. t. 32 = planiculmis.
lendigera, Salsab. Prod. 28 = Gastridium australe.
leonina, Steud. in Flora, xii. (1829) ii. 484 = Danthonia setacea ?
leptostachys, *Hook. f. Fl. Antarct.* 378.—Reg. Magellan.
levis, Nym. Consp. 811 = albinervis.
lodunensis, Delastr. in 'Cat. Hort. Pict. (1835)' = sulcata.
Loeflingiana, Cav. Ic. i. t. 45. f. 1 = Trisetum Gaudinianum.
Loeflingiana, Geuns. ex Roem. & Schult. Syst. ii. 670 = Aira praecox.
Loeflingiana, Balb. Misc. Bot. i. 12 = Trisetum Cavanillesii.
Loeflingiana, Linn. Sp. Pl. 79 = Trisetum hispanicum.
Loeflingiana, Lam. Encyc. i. 352 = Trisetum paniceum.
longifolia, Req. in DC. & Duby, Bot. Gall. i. 614 = pratensis.
longifolia, Spreng. Syst. i. 356 = Danthonia longifolia.
longifolia, Thore, Prom. Golf. fasc. 92 = Arrhenatherum pallens.
longiglumis, *Dur. in Duch. Rev. Bot.* i. (1845-46) 359.—Afr. bor.
lucida, Bertol. Fl. Ital. i. 701 = pubescens.
ludoviciana, Dur. in Bull. Soc. Linn. Bord. xx. (1855) t. 1 = sterilis.
lupulina, Steud. in Flora, xii. (1829) ii. 486 = Danthonia lanata.
lupulina, Thunb. Prod. Pl. Cap. 28 = Danthonia lupulina.
lutea, *Linn. f. Suppl.* 112.—Ins. Martinic.
maculenta, Guss. Fl. Sic. Prod. Suppl. i. 29 = Koeleria maculenta.
macra, Hohen. Enum. Pl. Elisabethpol. 214 = rigida.
macra, Stev. in Bieb. Fl. Taur. Cauc. i. 77 = Ventenata macra.
macrocarpa, Steud. in Flora, xii. (1829) ii. 482 = Danthonia macrantha.
macrocarpa, Moench, Meth. 196 = sterilis.
macrochaeta, *Balansa & Dur. in Bull. Soc. Bot. Fr.* i. (1854) 318.—Algeria.
magellanica, *Beauv. Agrost.* 153, nomen.—Reg. Magellan.
Mandoniana, Cosa. & Bal. in Bull. Soc. Bot. Fr. xv. (1868) 185, nomen = marginata ?
marginata, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) 502.—Ins. Madera.
maxima, *Presl, Fl. Sicul.* p. xliv.—Sicilia.
mediolanensis, Balansa & De Not. in Comoll. Fl. Comens. i. 147 = Trisetum myrianthum.
Michelii, Guss. Fl. Sic. Syn 151 = Koeleria maculenta.
mollis, Koel. Gram. 300 = Holcus mollis.
mollis, Michx. Fl. Am. Bor. i. 72 = Trisetum subspicatum.
molli, Salisb. Prod. 28 = Bromus mollis.
montana, Brot. Fl. Lusit. i. 109 = Arrhenatherum Thorei.
montana, Roem. & Schult. Syst. ii. 674, in syn. = pratensis.
montana, *Vill. Fl. Dauph.* ii. 151.—Mont. Pyren.
montana, [Weber, in] Wigg. Prim. Fl. Holsat. 9 = Deschampsia flexuosa.
multiculmis, Dum. Agrost. 121 = Aira caryophyllea.
murralis, Sallsb. Prod. 28 = Festuca Myurus.
muricata, Spreng. Nene Entdeck. i. 247 = Aristida capensis, Danthonia papillosa.
mutica, *Kroch. Fl. Sil. Suppl.* i. 187.—Europ.
myriantha, Bertol. Fl. Ital. i. 723 = Trisetum myrianthum.
myoceuais, Spreng. Syst. i. 337.—Ind. or.
nana, Kunze, ex Steud. Nom. ed. II. i. 172, 45 = Aira praecox.
Neesii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 229.—Afr. trop.
neglecta, Bertol. Fl. Ital. i. 718 = Trisetum Gaudinianum.
neglecta, Savi, Fl. Pis. t. 1. f. 4 = Trisetum paniceum.
nemoralis, Sallsb. Prod. 28 = Bromus asper.
nervosa, Lam. Illustr. n. 1118 = strigosa.
nervosa, R. Br. Prod. i. 178 = Amphibromus Neesii.

AVENA:—
Neumeyeriana, Vis. Fl. Dalm. iii. 339 = compacta.
nigra, *Wallr. Beitr. Fl. Hercyn.* 174; *et in Linnaea*, xiv. (1840) 544.—Europ.
nitida, Desf. Fl. Atlant. i. 102 = Trisetum nitidum.
nodosa, Cullum, It. Scan. 261 = Arrhenatherum avenaceum.
Noeana, Nym. Syll. Suppl. 71 = Trisetum aureum, *Tenore*.
Notarisii, Parl. Fl. Ital. i. 279; ex Steud. Syn. Pl. Gram. 425 = sempervirens.
nuda, *Linn. Amoen. Acad.* iii. 401.—Europ. austr.
nuda, Thunb. Fl. Par. ed. II. i. 49 = brevis.
nutans, St. Lag. Rechb. Hist.—Cf. Bull. Soc. Bot. Fr. xxxi. (1884) Bibl. 201 = sterilis.
nutkaensis, *Y. Presl, in Rel. Haenk.* i. 254.—Sinus Nootka.
occidentalis, *Dur. Cat. Gr. Bord.* (1864) 25.—*Cf. Bull. Soc. Bot. Fr.* xii. (1865) *Bibl.* 78.—Ins. Ferro.
odorata, DC. Fl. Fr. iii. 42 = Hierochloë borealis.
odorata, Pers. Syst. i. 100 = Hierochloë borealis.
oligostachya, *Munro, ex Aitch. in Journ. Linn. Soc.* xviii. (1881) 108; xix. 193. t. 30.—Afghanist.
orientalis, *Schreb. Spicil.* 59.—Europ. austr.; Orient.
ovata, C. C. Gmel. ex Kunth, Enum. Pl. i. 295 = Trisetum ovatum.
ovina, Salisb. Prod. 22 = Festuca ovina.
palaestina, *Steud. Syn. Pl. Gram.* 426.—Palaestina.
pallens, Link, in Schrad. Journ. ii. (1799) 314 = Arrhenatherum pallens.
pallida, Sallsb. Prod. 24 = Holcus lanatus.
pallida, Thunb. Prod. Pl. Cap. 22 = Danthonia pallescens.
palustris, Michx. Fl. Bor. Am. i. 72 = Trisetum palustre.
pancsea, Hort. ex Roem. & Schult. Syst. ii. 625, 660 = Koeleria phleoides.
panicea, Lam. Illustr. i. 202 = Trisetum paniceum.
papillosa, Schrad. in Goett. Gel. Ans. (1821) iii. 2075 = Danthonia papillosa.
papillosa, Steud. in Flora, xii. (1829) ii. 484 = Danthonia scabra.
paradensis, *Kit. ex Kanitz, in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 548 (*Kelog. Kitaib.* 119).—Europ.
paradoxa, Willd. ex Kunth, Enum. Pl. i. 188 = Aristida dichotoma.
Parlatorii, *Woods, ex Dalla Torre, Atl. Alpenfl.* 228.—Italia.
parviflora, Desf. Fl. Atlant. i. 98 = Trisetum parviflorum.
parviflora, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 280 = Trisetum micranthum.
pauciflora, Auct. ex Roem. & Schult. Syst. ii. 660 = Trisetum paniceum.
pauperula, *Phil. in Linnaea*, xxix. (1857-58) 94.—Chili.
pendula, Gilib. Exercit. 539 = sativa.
penicillata, Willd. ex Steud. Nom. ed. II. i. 173; ii. 714 = Trisetum rigidum.
pennsylvanica, Muhl. Descr. Uber. 185 = Arrhenatherum kentuckense.
pensylvanica, Forsk. Fl. Aegypt. Arab. 28 = Danthonia Forskolei.
pensylvanica, *Linn. Sp. Pl.* 79.—Am. bor.
pensylvanica, Presl, Cyp. et Gram. Sic. 80 = maxima.
persica, *Steud. Syn. Pl. Gram.* 235.—Persia.
peruviana, Hort. ex Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 290 = Trisetum paniceum.
phleoides, Urv. in Mém. Soc. Linn. Par. iv. (1826) 601 = Trisetum phleoides.
pilosa, Anchez, ex Dur. in Duch. Rev. Bot. i. (1845-46) 360 = cincta.
pilosa, *Bieb. Fl. Taur. Cauc.* iii. 84.—Ad. Min.; Syria; Afr. bor.
pilosa, *Y. Presl, in Rel. Haenk.* i. 258.—Chili.
pilosa, *Scop. Fl. Carn.* ed. II. i. 95.—Carniolia.
planiculmis, *Schrad. Fl. Germ.* i. 381. t. 6. f. 2.—Europ.; As. bor.
planiculmis, Sm. Engl. Bot. t. 2141 = pratensis.
podolica, Pascal. ex Zuccagni, in Roem. Collect. 126 = sativa.
Pourretii, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 290. t. 8.—Hispan.
praecox, *Beauv. Agrost.* 89.—Lusitan.
praecox, Reichb. Fl. Germ. Excurs. 140 = pratensis.
pratensis, Boiss. Voy. Bot. Espagne, ii. 656 = bromoides.
pratensis, Brot. Fl. Lusit. ii. 110 = sulcata.

AVENA:—
pratensis, Gouan, ' Hort. 53 '; Kunth, Enum. Pl. i. 360 = pubescens.
pratensis, *Linn. Sp. Pl.* 80.—Europ.; As. bor.
pratensis, Sibth. & Sm. Fl. Græc. Prod. i. 67 = caryophyllea.
precatoria, Thuill. Fl. Par. ed. II. 58 = Arrhenatherum avenaceum.
Prenlli, *Kunth, Enum. Pl.* i. 304.—Chili.
provincialis, Nym. Syll. Suppl. 71 = capillaris.
provincialis, Poir. Encyc. Suppl. i. 541 = Danthonia provincialis.
pseudolucida, Schur, in Oestr. Bot. Zeitschr. x. (1860) 73 = pubescens.
pseudoviolacea, *Kern. ex Dalla Torre, Atl. Alpenfl.* 228.—Europ. alp.
pubescula, Guss. in Tenore, Prod. Fl. Nap. App. v. = Koeleria macilenta.
pubescens, *Huds. Fl. Angl.* 42.—Europ.; As. bor.
pulchella, Beauv. Agrost. 154, nomen = Aira lendigera.
pumila, Desf. Fl. Atlant. i. 108 = Trisetum pumilum
pumila, Urv. in Mém. Soc. Linn. Par. i. (1822) 267 = Trisetum aureum, *Ten.*
purpurascens, DC. Cat. Hort. Monsp. 82 = Trisetum pratense.
purpurascens, De Not. Ind. Sem. Genuen. (1847); Ann. Sc. Nat. Sér. III. ix. (1848) 323 = Trisetum alpestre.
purpurea, *Gueldenst. ex Ledeb. Fl. Ross.* iv. 419.—Rossia.
purpurea, Linn. f. Suppl. 119 = Danthonia purpurea.
pusilla, [Weber, in] Wigg. Prim. Fl. Hols. 9 = Aira præcox.
quadridentata, *Doell, in Mart. Fl. Bras.* ii. III. 100.—Brsz.
quadriseta, Labill. Nov. Holl. Pl. I. 25. t. 32 = Deyeuxia quadriseta.
quinquaseta, Steud. in Flora, xii. (1829) II. 485 = Trisetum Steudelii.
racemosa, *Thuill. Fl. Par.* ed. II. 59.—Gallia.
Radula, *Franch. & Sav. Enum. Pl. Jap.* ii. 174.—Japon.
redolens, Pers. Syn. i. 100 = Hierochloë redolens.
rigida, Bieb. Fl. Taur. Cauc. i. 77 = Trisetum rigidum.
rigida, *Steud. in Flora*, xii. (1829) II. 483.—Afr. austr.
Rotæa, De Not. ex Parl. Fl. Ital. i. 264 = Trisetum argenteum.
rubra, Zuccagni, in Roem. Collect. 126 = sativa.
rupestris, J. F. Gmel. Syst. *I* /*h* = Agrostis rupestris.
Ruprechtii, *Ledeb. Fl. Ross.* iv. 418.—Rossia acut.
sativa, *Linn. Sp. Pl.* 79.—Terr. temp. (cult.).
scabriuscula, Lag. in Varied. Cienc. ii. IV. (1805) 219 = Koeleria scabriuscula.
scabrivalvis, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 67.—Chili.
Schellians, *Hack. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 342.—Hab.?
Schenckeri, *All. Fl. Pedem.* ii. 255.—Europ.; Reg. Caucas.
secalina, Salisb. Prod. 22 = Bromus secalinus.
secunda, Salisb. l. c. = Arrhenatherum avenaceum.
sedenensis, DC. Fl. Fr. iii. 719 = montana.
segetalis, Bianca, ex Nym. Consp. 810 = barbata.
sempervirens, Bory & Chaub. Fl. Pélop. 7. n. 179 = Haldenskii.
sempervirens, Host, Gram. iii. 28. t. 41 = Pariatorii.
sempervirens, Bon. Enum. Pl. Vallo. 6 = Besseri.
sempervirens, Lapeyr. ex Benth. Cat. Pl. Pyr. 65 = montana.
sempervirens, Ledeb. Fl. Alt. i. 89 = desertorum.
sempervirens, *Vill. Prosp.* 17.—Europ.; As. bor.
sempervillosa, *Ledeb. Fl. Ross.* iv. 419.—As. or.
sesquitertia, Bieb. Fl. Taur. Cauc. i. 77 = Trisetum rigidum.
sesquitertia, Comol. Prod. 18 = lucida.
sesquitertia, Host, Syn. Fl. Austr. 60 = Trisetum alpestre.
sesquitertia, Hort. ex Steud. Nom. ed. I. 95 = barbata.
sesquitertia, Linn. Mant. i. 34 = pubescens.
sesquitertia, Tenore, ex Steud. Nom. ed. II. i. 178 = semperviress.
sesquitertia, Willd. Sp. Pl. i. 448 = Trisetum argenteum.
setacea, Bamg. ex Schur, Enum. Pl. Transs. 735 = Agrostis rupestris.
setacea, Bamg. l. c. 755 = Deschampsia flexuosa.
setacea, Parl. Fl. Ital. i. 138 = Pariatorii.

AVENA:—
setacea, Muhl. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 87 = Aristida dichotoma.
setacea, *Vill. Prosp.* 16.—Europ. austr.
setifolia, Brot. Fl. Lusit. 108 = Arrhenatherum pallens.
sibirica, Linn. Sp. Pl. 79 = Stipa sibirica.
Sibthorpii, Nym. Syll. 413 = bromoides.
sicula, Spreng. Syst. i. 336 = Trisetum asperum, *Tenore.*
Smithii, *Porter, ex A. Gray, Man. Bot. U. St.* ed. V. 640.—Am. bor.
spicæformis, Anct. ex Roem. & Schult. Syst. ii. 690 = Danthonia spicæformis.
spicata, All. ex Kunth, Enum. Pl. i. 312 = Triodia decumbens, Danthonia provincialis.
spicata, Ell. Sketch, i. 174 = Danthonia sericea.
spicata, Gilib. Exercit. 540 = pratensis.
spicata, Linn. Sp. Pl. 80 = Trisetum subspicatum.
spicata, Willd. Sp. Pl. i. 455, partim = Danthonia provincialis.
splendens, Boiss. Elench. 88 = albinervis.
splendens, Anct. Gall. ex Steud. Nom. ed. II. i. 178 = pratensis.
splendens, Guss. Fl. Sic. Prod. i. 126 = Trisetum pratense.
spurca, *Schrank, in Denkschr. Regensb. Ges.* i. II. (1818) 7.—Labrador.
sterilis, Delile, ex Boiss. Fl. Orient. v. 643 = fatua.
sterilis, *Linn. Sp. Pl.* ed. II. 118.—Reg. Mediterr.; Orient.
stipæformis, *Linn. Mant.* 34.—Afr. austr.
stipoides, Willd. ex Steud. Nom. ed. II. i. 178; ii. 146 = Cynosurus fimbriatus.
striata, Lam. Encyc. i. 332 = sempervirens.
striata, Trin. in Mém. Acad. Pétersb. Sér. VI. iv. II. (1838) 32 = calloza.
striata, *Michx. Fl. Bor. Am.* i. 73.—Am. bor.
striata, *Schousb. ex Willd. & Lange, Prod. Fl. Hisp.* i. 60.—Hispan.
striata, Vis. Fl. Dalm. i. 70 = convoluta.
stricta, Host, Fl. Austr. i. 127 = Danthonia provincialis.
strigosa, *Schreb. Spic.* 52.—Europ.; Reg. Casp.
strigosa, Vogler, Schediasma Bot. 32 = Gaudinia tenuis.
subalpigrostis, Hartm. Handb. ed. III. ex cj. ed. IV. 39 = Trisetum agrostideum.
subcylindrica, Ehrenb. ex Boiss. Fl. Orient. v. 686 = Trisetum glumaceum.
subspicata, [Clairv.] Man. Herb. 17 = Trisetum subspicatum.
subulata, Lam. Illustr. n. 1118 = setacea.
subulifolia, Schur, Enum. Pl. Transs. 768 = convoluta.
sulcata, *Y. Gay, in Delastre, Fl. Vienne*, 477. t. 4.—Hispan.
sylvatica, Salisb. Prod. 24 = Holcus mollis.
symphicarpa, Trin. ex Steud. Nom. ed. II. i. 178 = hirta.
syriaca, Boiss. & Bal. ex Boiss. Fl. Orient. v. 642 = sterilis.
tatarica, Ard. in Saggi Acc. Padov. ii. (1789) 101. t. 1 = orientalis.
tayyetana, *Steud. Syn. Pl. Gram.* 238.—Græcia.
Tenorii, Nym. Consp. 814 = corymbosa.
tenuiflora, *Bertol. ex Reichb. Ic. Fl. Germ.* i. 41. n. 1668. —Italia.
tennis, Moench, Meth. 195 = Ventenata avenacea
Thorei, DC. & Duby, Bot. Gall. i. 514 = Arrhenatherum pallens.
toluccnsis, H. B. & K. Nov. Gen. et Sp. i. 146 = Trisetum toluccnse.
triaristata, Vill. Prosp. 17 = Ventenata avenacea.
trichophylla, C. Koch, in Linnæa, xxi. (1848) 393 = fatua.
trichopodia, *Y. Presl, in Rel. Haenk.* i. 254.—Mexic.
trinia, Thunb. Prod. Pl. Cap. 23 = Danthonia Thunbergii.
trispcrma, Anct. ex Roem. & Schult. Syst. ii. 669 = sativa.
tricoides, *Lindl. in Mitch. Journ. Austral.* 265.—Austral.
truncata, Dalac, Fl. Hautes-Pyr. 38 = bromoides.
tuberosa, Gilib. Exercit. ii. 585 = Arrhenatherum avenaceum.
uniflora, *Parl. Pl. Nov.* 34.—Ins. Canar.
unilateralis, Bromm. ex Roem. & Schult. Syst. ii. 669 = orientalis.
valesiaca, Nym. Syll. 414 = Trisetum Cavanillesii.

AVENA:—
varia, Schur, in Oestr. Bot. Zeitschr. x. (1860) 75 = Trisetum carpaticum.
velutina, *Nym. Consp.* 813 = Trisetum velutinum.
ventricosa, *Balansa, in Bull. Soc. Bot. Fr.* i. (1854) 14 —Algeria.
verticillor, Bamg. ex Steud. Syn. Pl. Gram. 238 = Baumgartenii.
verticolor, Vill. Prosp. 17 = Scheuzeri.
verticolor, St. Am. Fl. Agen. 48 = sulcata.
ville, *Wallr. Beitr. Fl. Hercyn.* 178; *et in Linnæa*, xiv. (1840) 548.—German.
villosa, Bertol. Exc. de Re Herb. 6. n. 8 = Trisetum villosum.
villosa, Lag. ex Roem. & Schult. Syst. ii. 670 = hirsuta.
virescens, *Regel, in Act. Hort. Petrop.* vii. (1880) 695. —Turkestan.
viridis, H. B. & K. Nov. Gen. et Sp. i. 147 = Deyeuxia viridis, *Fourn.*
Wiesti, *Steud. Syn. Pl. Gram.* 231.—Aegypt.; Arab.; Persia.
Wilhelmsii, Spreng. Syst. i. 338 = Trisetum rigidum.

AVENELLA, Koch, ex Steud. Nom. ed. II. i. 173 (1840); Parl. Fl. Ital. i. 344 (1848) = Deschampsia, Beauv. (Gramin.).
cuprina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 86 = D. flexuosa.
flexuosa, Parl. Fl. Ital. i. 346 = D. flexuosa.
rubra, Jord. ex Nym. Consp. 806 = D. flexuosa.
uliginosa, Parl. Fl. Ital. i. 346 = D. uliginosa.

AVERRHOA, Linn. Syst. ed. I (1735). *GERANIACEÆ*, Benth. & Hook. f. i. 277.
CARAMBOLA, Adans. Fam. ii. 508 (1768).
acida, Linn. Sp. Pl. 439 = Phyllanthus Cicca.
acutangula, Stokes, Bot. Mat. Med. ii. 548 = A. Carambola.
Bilimbi, *Linn. Sp. Pl.* 438.—Ind. or.; China.
Carambola, *Linn. Sp. Pl.* 438.—Ind. or.; China.
frondosa, Salisb. Prod. 318 = Phyllanthus Cicca.
obtusangula, Stokes, Bot. Mat. Med. ii. 542 = A. Bilimbi.
pentandra, Blanco, Fl. Filip. ed. I. 392 = Connaropsis philippvica.
sinica, Hance, in Walp. Ann. ii. 241 = Rourea microphylla.

AVERRHOIDIUM, Baill. Adansonia, xi. 344 (1874). *SAPINDACEÆ*.
Gardnerianum, *Baill. l. c.*—Bras.

AVERSIA, G. Don, Gen. Syst. iii. 92 (1834) = ARVERSIA, Cambess. = **Polycarpon**, Linn. (Caryophyll.).

AVICHINNIA, Linn. Syst. ed. I (1735); Gen. ed. I. 37 (1737). *VERBENACEÆ*, Benth. & Hook. f. ii. 1160.
BONTIA, Linn. Gen. ed. I. 180 (1737).
DONATIA, Lœfl. Iter, 192 (1758).
HALODENDRUM, Thou. Gen. Nov. Madag. 8 (1806).
RACKA, J. F. Gmel. Syst. 345 (1791).
SCHUBA, Forsk. Fl. Aegypt. Arab. 37 (1775).
UPATA, Adans. Fam. ii. 201 (1763).
africana, Beauv. Fl. Owar. i. 80. t. 47 = officinalis.
alba, Blume, Bijdr. 831 = officinalis.
elliptica, Thunb. Fl. Bras. Dec. iii. 87 = officinalis.
floridana, Rafin. Atl. Journ. 148 = officinalis.
intermedia, Griff. in Trans. Linn. Soc. xx. (1846) 6 = officinalis.
Lamarckiana, Presl, Bot. Bemerk. 99 = officinalis.
Meyeri, Miq. in Linnæa, xviii. (1844) 382 = officinalis.
nitida, *Yacq. Enum. Pl. Carib.* 35.—Am. trop.; Afr. trop.
oblongifolia, Nutt. ex Chapm. Fl. S. U. St. 310 = officinalis.
obovata, Griff. Notul. iv. 389 = officinalis.
Ospata, Buch.-Ham. in Trans. Linn. Soc. xvii. (1837) 321 = officinalis.
officinalis, *Linn. Sp. Pl.* 110.—Littora tropica.
reniformis, Forst. f. Fl. Escul. 72 = officinalis.
tomentosa, Jacq. Enum. Pl. Carib. 25 = officinalis.

AVICEPS, Lindl. Gen. et Sp. Orch. 345 (1838) = Satyrium, Sw. (Orchid.).
pumila, Lindl. l. c. 346 = S. pumilum.

AVICULARIA, Stead. Nom. ed. II. i. 174 (1840) = **Polygonum**, Tourn.
indica, Didricha. in Bot. Notiser, (1850) 187 = P. plebeium.
maritima, Didricha. l. c. = P. maritimum.
vulgaris, Didricha. l. c. = P. aviculare.

AVIUNCULUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 367 = **Coronilla**, Linn. (Legum.).
scorpioides, Fourr. l. c. = C. scorpioides.

AVORNELA, Rafin. Sylva Tellur. 24 (1838) = **Genista**, Linn. (Legum.).
canariensis, Rafin. l. c. = G. canariensis.
candicans, Rafin. l. c. = Cytisus candicans.
linifolia, Rafin. l. c. = Cytisus linifolia.

AWAYUS, Rafin. Sylva Tellur. 151 (1838) = **Spiraea**, Linn. (Rosac.).
japonicus, Rafin. l. c. 151 = S. japonica.

AXANTHES, Blume, Bijdr. 1002 (1826) = **Urophyllum**, Wall. (Rubiac.).
arborea, Blume, l. c. 1003 = U. glabrum.
Blumeana, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 146 = U. Blumeanum.
cejanica, Wight, l. c. 146 = U. zeylanicum.
corymbosa, Blume, Bijdr. 1003 = U. corymbosum.
elliptica, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 144 = U. ellipticum.
enneandra, Wight, l. c. 144 = Anlacodiscus premnoides.
Griffithiana, Wight, l. c. 147 = U. Griffithianum.
hirsuta, Wight, l. c. 146 = U. hirsutum
longifolium, Wight, l. c. 145 = U. longifolium.
macrophylla, Blume, Bijdr. 1002 = U. macrophyllum.
philippensis, Cham. & Schlecht. ex Steud. Nom. ed. II. i. 174.—Ins. Luzon.
strigosa, Blume, Bijdr. 1003 = U. strigosum.
timoriensis, DC. Prod. iv. 441.—Ins. Timor.
tomentosa, Blume, ex DC. l. c. 440 = U. tomentosum.
zeylanica, Wight, Ic. t. 1163 = U. zeylanicum.

AXANTHOPSIS, Korth. in Nederl. Kruidk. Arch. ii. (1851) 195 = Urophyllum, Wall. (Rubiac.).
elongata, Korth. l. c. = U. elongatum.

AXENFELDIA, Baill. Etud. Gén. Euph. 419 (1858) = **Mallotus**, Lour. (Euphorb.).
intermedia, Baill. l. c. = M. muricatus.

AXIA, Lour. Fl. Cochinch. 85 (1790). *VALERIANACEAE*
cochinchinensis, *Lour. l. c.* 86.—Cochinch.

AXILLARIA, Rafin. in Am. Monthly Mag. (1818) 266 = Polygonatum, Adans. (Liliac.).

AXINAEA, Ruiz & Pav. Prod. 68 (1794). *MELASTOMACEAE*, Benth. & Hook. f. i. 749.
CLAETHRARIA, DC. Prod. iii. 103 (1828).
bracteata, Zipp. ex Blume, in Flora, (1831) 519 = Medinilla bracteata.
cramboda, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 69.—Peruv.
dependens, *Ruiz & Pav. Fl. Per.* iv. t. 411.—Peruv.
floribunda, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 69.—Ecuador.
glandulosa, *Ruiz & Pav. Fl. Per.* iv. t. 412.—Peruv.
grandifolia, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 69.—Venezuela.
lanceolata, *Ruiz & Pav. Fl. Per.* iv. 409; *Syst.* i. 122. —Peruv.
lepidota, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 69.—Ecuador.
macrophylla, *Triana*, *l. c.*—Ecuador.
Meriantea, *Triana*, *l. c.*—Venezuela.
muricata, D. Don, in Mem. Wern. Soc. iv. (1832) 321 = Miconia muricata.
purpurea, Ruiz & Pav. Fl. Per. iv. 410; Syst. i. 122 = Marianta radula.
Radula, Benth. Pl. Hartw. 180 = Marianta Radula.
reticophylla, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 68.—Ecuador.
ruginosa, *Triana*, *l. c.* 69.—N. Granat.
semifolia, *Triana*, *l. c.* 68.—Peruv.

AXINEA, Juss. in Dict. Sc. Nat. xxix. (1823) 508 = prec.

AXINANDRA, Thw. in Hook. Kew Journ. vi. (1854) 66. t. 1. C. *LYTHRARIEAE*, Benth. & Hook. f. i. 784.
alata, *Baill. Adansonia*, xii. (1876) 86.—Malaya.
Beccariana, *Baill. l. c.* 85.—Malaya.
coriacea, *Baill. l. c.*—Malaya.
Maingayi, *C. B. Clarke*, *in Hook. f. Fl. Brit. Ind.* ii. 581.—Malacca.
zeylanica, *Thw. in Hook. Kew Journ.* vi. (1854) 67.—Zeylan.

AXINANTHERA, Karst. in Linnaea, xxx. (1859–60) 157 = **Bellucia**, Neck. (Melast.).
Bellucia, Karst. l. c. = B. Axinanthera.
macrophylla, Karst. l. c. = B. superba!

AXINIPHYLLUM, Benth. in Hook. Ic. Pl. xii. 16. t. 1118 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 369.
corymbosum, *Benth. l. c.*—Mexic.
tomentosum, *Benth. l. c.* sub t. 1118.—Mexic.

AXIRIS, Linn. Diss. Chen. 43 (1751) = **Axyris**, Linn. (Chenop.).

AXOLOPHA, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = **Lavatera**, Linn. (Malvac.).
aegyptia, Alef. l. c. 259 = Malva aegyptia.
althaeoides, Alef. l. c. = Malva althaeoides.
hirsuta, Alef. l. c. = Althaea hirsuta.
Ludwigii, Alef. l. c. = Althaea Ludwigii.
maritima, Alef. l. c. = L. maritima.
Wigandi, Alef. l. c. = L. maritima.

AXOLUS, Rafin. Sylva Tellur. 61 (1838) = **Cephalanthus**, Linn. (Rubiac.).
angustifolius, Rafin. l. c. = C. angustifolius.

AXONOPUS, Beauv. Agrost. 12 (1812) = **Paspalum**, Linn. (Gramin.).
aureus, Beauv. l. c. = P. aureum.
cimicinus, Beauv. l. c. = Panicum cimicinum.
compressus, Beauv. l. c. = P. compressum.
corymbosus, Schult. Mant ii. 477 = Panicum corymbosum.
digitatus, Beauv. Agrost. 13 = P. digitatum.
fasciculatus, Beauv. l. c. = Panicum filiforme.
Poiretii, Roem. & Schult. Syst. ii. 818 = P. compressum.
savannarum, Schlecht. in Linnaea, xxvi. (1853–55) 153 = P. savannarum.

AXONOTECHIUM, Fenzl, in Ann. Wien. Mus. i. (1836) 884 = Orygia, Forsk. (Picrid.).
trianthemoides, Fenzl, l. c. 886 = O. decumbens.

AXYRIS, Linn. Syst. ed. VI. Add. (1748); Amoen. Acad. 24 (1751). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 56.
amaranthoides, *Linn. Sp. Pl.* 979.—Ass. bor.
ceratoides, Linn. l. c. = Eurotia ceratoides.
hybrida, Linn. l. c. 980 = amaranthoides.
Morcroftiana, R. Br. in Wall. Cat. n. 6950 = Eurotia ceratoides.
pentandra, *Vasq. Ann. Select.* 244.—Cuba.
prostrata, Linn. Sp. Pl. 980 = amaranthoides!
sphaerosperma, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. (1839) 46.—Mong. Altai.
trifaria, Moench, Meth. Suppl. 190 = amaranthoides.
villosa, *Meq. in DC. Prod.* xiii. ii. 117.—Afghan.

AYAPANA, Spach, Hist. Vég. Phan. x. 290 (1841) = Eupatorium, Linn. (Compos.).
officinalis, Spach, l. c. = E. Ayapana.

AYDENDRON, Nees, in Linnaea, viii. (1833) 86. *LAURINEAE*, Benth. & Hook. f. iii. 152.
ENGLICHERIA, Nees, in Linnaea, viii. (1833) 87.
GORPERTIA, Nees, Syst. Laurin. 365 (1836).
SCHAUERA, Nees, in Lindl. Nat. Syst. ed. II. 209 in adnot. (1836).
aciphyllum, *Nees*, *in Linnaea*, xxi. (1848) 496.—Guiana.
affinis, *Meisn. in DC. Prod.* xv. i. 93.—Bras.
amazonicum, *Meisn. l. c.* 89.—Bras.
argenteum, *Griseb. Fl. Brit. W. Ind.* 385.—Ind. occ.
brachteatum, *Nees*, *Syst. Laurin.* 256.—Ins. S. Vincent.

AYDENDRON:—
brasiliense, *Meisn. in Mart. Fl. Bras.* v. ii. 180.—Bras.
Cannella, *Meisn. in DC. Prod.* xv. i. 90.—Guiana.
cayennense, *Meisn. l. c.* 95.—Guiana.
citrifolium, Griseb. Fl. Brit. W. Ind. 284 = trinitatis.
citrifolium, *Nees, Syst. Laurin.* 257.—Bras.
cubense, *A. Rich. Fl. Cub. Famer.* ii. 187.—Cuba.
Cujumary, *Nees, Syst. Laurin.* 247.—Bras.
curvinanum, *Meisn. in DC. Prod.* xv. i. 100.—Guiana.
desertorum, *Nees, Syst. Laurin.* 260.—Bras.
firmulum, *Nees*, *in Linnaea*, viii. (1833) 86.—Bras.
floribundum, *Meisn. in DC. Prod.* xv. i. 88.—Bras.
Gardneri, *Meisn. l. c.* 89.—Bras.
glaucum, *Klotzsch & Karst. ex Meisn. l. c.* 90.—N. Granat.
goyazense, *Meisn. l. c.* 94.—Bras.
Hostmannianum, *Nees*, *in Linnaea*, xxi. (1848) 499.—Guiana.
intermedium, *Meisn. in DC. Prod.* xv. i. 89.—Bras.
laevigatum, *Meisn. l. c.* 88.—Bras.
Laurel, *Nees, Syst. Laurin.* 249.—Venezuela.
macrophyllum, *Meisn. in DC. Prod.* xv. i. 92.—N. Granat.
microbotrum, *Nees, Syst. Laurin.* 254.—Ins. Portorico.
Muca, *Nees*, *in Linnaea*, xxi. (1848) 498.—Peruv.
nitidum, *Meisn. in DC. Prod.* xv. i. 95.—Bras.
obtusilolium, *Nees, Syst. Laurin.* 258.—Am. austr.
pachyrrapum, *Meisn. in DC. Prod.* xv. i. 95.—Guiana.
pallidum, *Meisn. l. c.* 98.—Bras.
pauciflorum, *Meisn. l. c.* 89.—Bras.
parviflorum, *Meisn. l. c.* 90.—Bras.
pernaule, *Nees, Syst. Laurin.* 240.—Bras.
pisiiyense, *Meisn. in DC. Prod.* xv. i. 93.—Bras.
riparium, *Nees*, *in Linnaea*, xxi. (1848) 497.—Guiana.
robustum, *Klotzsch & Karst. ex Meisn. in DC. Prod.* xv. i. 91.—N. Granat.
salicifolium, *Nees, Syst. Laurin.* 252.—Ind. occ.; Guian.
Sellowii, *Meisn. in DC. Prod* xv. i. 88.—Bras.
sericeum, *Griseb. Fl. Brit. W. Ind.* 285.—Ind. occ.
suaveolens, *Nees*, *in Linnaea*, viii. (1833) 87.—Bras.
tenellum, *Meisn. in DC. Prod.* xv. i. 93.—Bras.
trinitatis, *Meisn. l. c.* 91.—Ins. Trinit.
verrucosum, *Nees, Syst. Laurin.* 259.—Bras.
verticillatum, *Meisn. in DC. Prod.* xv. i. 91.—Ins. S. Vincent.

AYENIA, Loefl. It. 199 (1758). *STERCULIACEAE*, Benth. & Hook. f. i. 295.
CYBIOTTIGNA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 155.
DAYENIA, Mill. Gard. Dict. ed. VII. (1759).
JUNGIA, Loefl. It. 199 (1758).
abutilifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 569 = magna.
acalyphifolia, *Griseb. in Gott. Abh.* xxiv. (1879) 50.—Reg. Argent.
angustifolia, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 84.—Bras.
cardiopetala, Seem. ex DC. Prod. i. 488 = magna.
cordifolia, *Seem. ex DC. l. c.*—Mexic.
cordobensis, *Hieron. Ic. Pl. Argent.* 50.—Reg. Argent.
cephalaefolia, *Griseb. Cat. Pl. Cub.* 29.—Cuba.
hirta, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 84.—Bras.
inermis, Medic. ex Steud. Nom. ed. II. i. 174 = pusilla.
laevigata, *Sw. Prod. Veg. Ind. Occ.* 97.—Ins. Jamaic.
linguiata, *Griseb. in Gott. Abh.* xxiv. (1879) 50.—Reg. Argent.
magna, *Linn. Syst. ed.* X. 1347.—Mexic.
mexicana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 569.—Mexic.
microphylla, *A. Gray*, *in Smithson. Contr.* iii. (1853) 24.—N. Mexic.
ovata, *Hemsl. Diag. Pl. Nov.* 4.—Mexic.
pusilla, *Linn. Syst. ed.* X. 1347.—Ins. Carib.; Peruv.
rotundifolia, *Hemsl. Diag. Pl. Nov.* 4.—Mexic.
sidaefolia, Hemsl. Biol. Centr. Am. Bot. i. 185 = mexicana.
sidaefolia, DC. Prod. i. 488 = tomentosa.

AYENIA:—
sidaefórmis, Loefl. It. 257 = tomentosa.
stipularis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 334.—N. Granat.
tomentosa, *Linn. Syst.* ed. X. 1247.—Venezuela.

AYLANTUS, Juss. Gen. 373 (1789) = **Allanthus**, Desf. (Simarub.).

AYLMERIA, Mart. Amaranth. 63; et in Nov. Act. Nat. Cur. xiii. (1826) 276 = **Polycarpaea**, Lam. (Caryophyll.).
rosea, Mart. ll. cc. = P. violacea.
violacea, Mart. ll. cc. = P. violacea.
Zollingeri, Fenzl, in Zoll. Syst. Verz. Ind. Archip. 141.—Malaya

AYPARIA, Rafin. Sylva Tellur. 154 (1838) = **Symplocos**, Linn. (Styrac.).
crenata, Rafin. l. c. (= *Dicalix cochinchinensis*, Lour.). —Cochinch.

AYTONIA, Linn. f. Suppl. 49, 308 (1781) = **Aitonia**, Thunb. (Sapind.).

AZADIRACHTA, A. Juss. in Mém. Mus. Par. xix. (1830) 220 (*Mél.* 68) (1830) = **Melia**, Linn.
indica, A. Juss. ll. cc. 221, 69 = Melia Azadirachta.

AZALEA, Gaertn. Fruct. i. 301. t. 63 (1788), pro parte = **Loiseleuria**, Desv. (Ericac.).

AZALEA, Linn. Syst. ed. I (1735) = **Rhododendron**, Linn. (Ericac.).
alba, Blume, ex DC. Prod. vii. 721 = R. album.
alba, Sweet, in Ann. Fromont, i. (1829) 104 = R. ledifolium.
amoena, Lindl. in Paxt. Flow. Gard. iii. (1852) t. 89 = R. indicum.
arborea, Bart. ex Steud. Nom. ed. II. i. 174 = R. arborescens.
arborea, Linn. Sp. Pl. 150 = R. ponticum.
arborescens, Pursh, Fl. Am. Sept. i. 158 = R. arborescens.
aurantiaca, Dietr. Darst. Vorz. Ziersfl. Heft i. 5. t. 1 = R. calendulaceum.
Beulii, Fortune, ex C. Koch, Dendrol. ii. 11. 175 = R. sublanceolatum.
bicolor, Pursh, Fl. Am. Sept. i. 158 = R. nudiflorum.
bullata, Forst. ex Steud. Nom. ed. II. i. 174 = Labetanthus americana.
Burgeri, Miq. Ann. Mus. Bot. Lugd. Bat. i. 84 = R. indicum.
calendulacea, Hook. & Arn. Bot. Beech. Voy. 361 = R. occidentale.
calendulacea, Michx. Fl. Bor. Am. i. 151 = R. calendulaceum.
californica, Torr. & Gray, ex Durand, in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 94 = R. californicum.
calycina, Linn. fard. Fl. iii. (1858) t. 257 = R. indicum.
canescens, Michx. Fl. Bor. Am. i. 150 = R. nudiflorum.
celebica, Blume, ex DC. Prod. viii. 735 = R. malayanum.
chinensis, Steud. Nom. ed. II. i. 175 = R. sinense.
crispiflora, Hook. Bot. Mag. t. 4726 = R. indicum.
crocea, Hoffmgg. Verz. Pfl. Nachtr. iii. 22 = R. calendulaceum!
dahurica, C. Koch, Dendrol. ii. 11. 181 = R. dauricum.
Danielsiana, Paxt. Mag. Bot. I. 196 = R. indicum.
daurumbens, DC. Prod. vii. 718 = R. indicum.
Farrerae, C. Koch, Dendrol. ii. 11. 178 = R. Farrerae.
ferruginea, Pall. Fl. Ross. ii. t. 70 = R. Anthopogon.
fimbriata, Dum.-Cours. Bot. Cult. ed. II. iii. 337.—Chin.
flammea, Prits. Ic. Ind. 131 = R. calendulaceum.
flava, Hoffmgg. Verz. Pfl. Nachtr. ii. 32 = R. calendulaceum!
fragrans, Adams, in Mém. Acad. Pétersb. ii. (1806) 329 = R. Anthopogon.
fragrans, Rafin. Ann. Nat. 13 = R. arborescens.
fulva, Michx. ex Lam. Journ. Hist. Nat. i. (1792) 410 = R. nudiflorum.
glandaensis, Hort. ex C. Koch, Dendrol. ii. 11. 185 = R. ponticum.
glauca, Lam. Illustr. 493. t. 110 = R. viscosum.
glauca, Mœnch. Pl. Sel. Ic. t. 6 = R. glaucum!

AZALEA:—
hispida, Pursh, Fl. Am. Sept. i. 154 = R. viscosum.
indica-alba, Lindl. in Bot. Reg. t. 811 (1824) = R. ledifolium.
indica, Linn. Sp. Pl. 150 = R. indicum.
japonica, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 406.—Japon.
javanica, Blume, ex DC. Prod. vii. 721 = R. javanicum.
Jodogawa, Hassk. Cat. Hort. Bog. Alt. 162 = R. Kaempferi.
lapponica, Linn. Sp. Pl. 151 = R. lapponicum.
lapponica, Pall. Fl. Ross. ii. 62. t. 70. f. 1 = R. Anthopogon.
ledifolia, Hook. Bot. Mag. t. 2901 = R. ledifolium.
lilliflora, Poit. in Rev. Hortic. (1859) 299 = R. ledifolium.
linariaefolia, Carr. l. c. (1872) 100 = seq.!
linearifolia, Hook. f. Bot. Mag. t. 5769 = R. linearifolium.
lucida, Meerb. Pl. Sel. Ic. 7 = R. lucidum.
Ludovicae, Herb. in Bot. Reg. xxxi. 60.—Hab.!
lugens, Dum.-Cours. Bot. Cult. ed. II. iii. 337.—Chin.
lutea, Linn. Sp. Pl. 150 = R. nudiflorum.
macrantha, Bunge, Enum. Pl. Chin. Bor. 41 = R. indicum.
malis, Blume, Bijdr. 853 = R. molle.
Mertensi, Sweet, ex Loud. Hort. Brit. Suppl. iii. 496 = R. calendulaceum.
Mertensiana, Hort. ex Spae, in Ann. Gand. ii. (1846) 325 = R. calendulaceum.
mucronata, Blume, Bijdr. 853 = R. ledifolium.
multiflora, Turcz. ex DC. Prod. vii. 715 = R. Anthopogon.
myrtifolia, Champ. in Hook. Bot. Mag. lxxvii. (1851) sub t. 4609 = R. ovatum.
nervosifolia, Fortune, ex Planch. in Rev. Hortic. Sér. IV. iii. (1854) 67 = R. ledifolium.
nitida, Pursh, Fl. Am. Sept. i. 158 = R. nitidum.
nivea, Dehnh. Cat. Pl. Hort. Camald. ed. II. 17.—Hab.!
nudiflora, Linn. Sp. Pl. ed. II. 214 = R. nudiflorum.
obtusa, Lindl. in Journ. Hort. Soc. i. (1846) 152 = R. indicum.
occidentalis, Torr. & Gray, ex Torr. in Pacif. Rail. Rep. iv. 116 = R. occidentale.
ovata, Lindl. in Journ. Hort. Soc. i. (1846) 149 = R. lepidotum.
pallida, Turcz. in Bull. Soc. Nat. Mosc. (1838) 96; xxi. (1848) ii. 498.—Mont. Baical.
periclymena, Pers. Syn. i. 318 = R. nudiflorum.
periclymenoides, Michx. Fl. Bor. Am. i. 151 = R. nudiflorum.
phaenicea, Hort. ex Planch. in Fl. des Serres, Sér. I. ix. (1853–54) 79 = R. puniceum.
pilosa, Michx. in Lam. Journ. Nat Hist. i. (1792) 410 = Menziesia globularis.
ponticа, Linn. Sp. Pl. 150 = R. ponticum.
procumbens, Linn. l. c. 151 = Loiseleuria procumbens.
prolifera, Poit. in Ann. Fromont, i. (1829) 104 = R. Kaempferi.
punctata, Lour. Fl. Cochinch. 118; ed. Willd. i. 189 = R. Loureirianum.
punicea, Sweet, in Ann. Fromont, i. (1829) 104 = R. sublanceolatum.
purpurea, Dum.-Cours. Bot. Cult. ed. II. iii. 337.—Ind. or.
quinquefolia, Lindl. Journ. Hort. Soc. iv. (1849) 291 = R. indicum.
Rawsonii, Paxt. Mag. Bot. ii. 121.—Hab.!
retusa, Blume, ex Steud. Nom. ed. II. i. 175 = R. retusum.
rosmarinifolia, Barm. f. Fl. Ind. 43 = R. ledifolium.
rubra, Meerb. Pl. Sel. Ic. 8.—Hab.!
scabra, Lodd. ex A. Gray, Syn. Fl. N. Am. ii. 11. 43 = R. viscosum.
serpyllifolia, A. Gray, in Perry, Jap. Exped. ii. 315 = R. serpyllifolium.
Seymouri, Lindl. in Bot. Reg. t. 1975 = R. Seymouri.
sinensis, Lodd. Bot. Cab. t. 885 = R. sinense.
speciosa, Willd. Enum. Hort. Berol. Suppl. 10 = R. speciosum.
squamata, Lindl. in Journ. Hort. Soc. i. (1846) 152 = R. Farrerae.
tomentosa, Duval, ex Steud. Nom. ed. I. 97 = Loiseleuria procumbens.
tubiflora, Blume, ex DC. Prod. vii. 725 = R. malayanum.

AZALEA:—
verticillata, Bart. ex Lodd. Cab. t. 1532.—Am. bor
virens, Dum.-Cours. Bot. Cult. ed. II. iii. 335 = R. viscosum.
viscosa, Linn. Sp. Pl. 151 = R. viscosum.
vittata, Dum.-Cours. Bot. Cult. ed. II. iii. 337 = R. vittatum.
vittato-punctata, Lem. Illustr. Horticr. (1854) t. 20 = prec.!

AZALTEA, Walp. Rep. i. 539 (1842) = **Alsatea**, Ruiz & Pav. (Celastr.).

AZAMARA, Hochst. ex Reichb. Nom. 145 (1841) = **Schmidelia**, Linn. (Sapind.).
trifoliata, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. 421 = S. africana.

AZANZA, Alef. in Bot. Zeit. xix. (1861) 298 = **Thespesia**, Corr. (Malvac.).
acuminata, Alef. l. c. 299 = T. populnea.
Lampas, Alef. l. c. 297 = T. Lampas.
Zollingeri, Alef. l. c. 298.—Java.

AZANZA, Moç. & Sessé, ex DC. Prod. i. 459 (1824) = **Hibiscus**, Linn. (Malvac.).
insignis, Moç. & Sessé, l. c. 454 = H. Azanza.

AZAOLA, Blanco, Fl. Filip. 402 (1837) = **Payena**, A. DC. (1844). (Sapot.).
Betis, Blanco, l. c. = P. Betis.
Leerii, Teijsm. & Binn. in Tijdschr. Nederl. Ind. vi. (1854) 116 = P. Leerii.

AZARA, Ruiz & Pav. Prod. 79. t. 36 (1799). *BIXINEAE*, Benth. & Hook. f. i. 127.
MYRTOPHYLLUM, Turcz. in Bull. Soc. Nat. Mosc. (1843) i. 604.
STLENIA, Bert. in Merc. Chil. (1829) ex Bull. Férussac. xx. (1830) 108.
alpina, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 20. t. 107. —Chili.
Berteroniana, *Steud. in Flora*, xxxix. (1856) 420.—Chili.
celastrina, *D. Don, in Edinb. N. Phil. Journ.* x. (Dec. 1830) 119.—Chili.
chilensis, *Hook. f. Fl. Antarct.* 248.—Chili.
dentata, *Ruiz & Pav. Syst.* 138.—Chili.
dubia, *Steud. in Flora*, xxxix. (1856) 418.—Chili.
Fernandeziana, C. Gay, Fl. Chil. i. 196 = serrata.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 144.—Chili.
hirtella, *Mey. in Linnaea*, xxv. (1852) 650.—Chili.
integrifolia, Bert. ex Steud. in Flora, xxxix. (1856) 420 = Berteroniana.
integrifolia, Miq. ex Steud. l. c. = Lechleriana.
integrifolia, *Ruiz & Pav. Syst.* 138.—Chilli.
intermedia, *C. Gay, Fl. Chil.* i. 194.—Chili.
lanceolata, *Hook. f. Fl. Antarct.* 248.—Chili.
Lechleriana, *Steud. in Flora*, xxxix. (1856) 420. — Chili.
Lilen, Bert. in Merc. Chil. (1829).—Cf. Bull. Férussac, ex. (1830) 108 = celastrina, spanifiora.
microphylla, *Hook. f. Fl. Antarct.* 244.—Chili.
salicifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 20.—Reg. Argent.
serrata, Bert. ex C. Muell. in Walp. Ann. vii. 926 = dubia.
serrata, *Ruiz & Pav. Prod.* 79. t. 36; *Syst.* 137.—Chili.
spanifiora, *Steud. Nom.* ed. II. i. 175.—Chili.
tomentosa, *Bert. ex Steud. in Flora*, xxxix. (1856) 419. —Chili.
umbellata, *Phil. in Linnaea*, xxviii. (1856) 610.—Chili.
umbellata, Presl, Rel. Haenk. ii. 92 = Laetia apetala.
Valdiviae, *Lechler, ex Steud. in Flora*, xxxix. (1856) 420.—Chili.

AZAROLUS, Borkh. Handb. Forstb. ii. 1224 (1803) = **Pyrus**, Linn. (Rosac.).
alpina, Borkh. l. c. 1234 = P. alpina.
arbutifolia, Borkh. l. c. 1236 = P. arbutifolia.
Aria, Borkh. l. c. 1230 = P. Aria.
Chamaemespilus, Borkh. l. c. 1237 = P. Chamaemespilus.
crataegifolia, Borkh. l. c. 1238 = Crataegus Azarolus.
heterophylla, Borkh. l. c. 1244.—Hab.?
hybrida, Borkh. l. c. 1239 = P. rotundifolia.
intermedia, Borkh. l. c. 1232 = P. intermedia.

AZAROLUS :—
maroccana, M. Roem. Syn. Rosifl. 138 = Crataegus Azarolus.
Oliveriana, M. Roem. l. c. = Crataegus pentagyna.
pinnatifida, Borkh. Handb. Forstb. ii. 1245 = Pyrus pinnatifida.
platyphylla, M. Roem. Syn. Rosifl. 138 = Crataegus pentagyna.
Pallasiana, Borkh. Handb. Forstb. ii. 1251 = P. Pollveria.
pubescens, M. Roem. Syn. Rosifl. 134 = Crataegus pubescens.
torminalis, Borkh. Handb. Forstb. ii. 1235 = P. torminalis.

AZEDARA, Rafin. Fl. Ludov. 135 (1817) = **Melia**, Linn.
speciosa, Rafin. l. c. = M. Azedarach.

AZEDARAC, Adans. Fam. ii. 342 (1763) = seq.

AZEDARACH, Tourn. ex Linn. Gen. ed. I. 127 (1737) = **Melia**, Linn.
Commelini, Medic. Bot. Beobacht. (1782) 164 = M. Azedarach.
deiteria, Medic. Malv. 116 = M. Azadir. ta.
edule, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 5 = Sandoricum indicum.
fraxinifolia, Moench, Meth. Suppl. 58 = M. Azadirachta.
nigrum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 5.—Malaya.
odoratum, Noronha, l. c. = M. Azedarach.
ramiflorum, Noronha, l. c. = Spicharis cauliflora.
racisum, Noronha, l. c.—Malaya.

AZEREDIA, Arruda, ex Allemão, Desemb. cum Ic. (1846). *BIXINEAE*, Benth. & Hook. f. i. 125.
pernambucana, *Allemão, l. c.*—Bras.

AZIMA, Lam. Encyc. i. 343 (1789). *SALVADORACEAE*, Benth. & Hook. f. ii. 681.
ACTEOTON, Blume, Bijdr. 1143 (1826).
KANDENA, Rafin. Sylva Tellur. 138 (1838).
MONETIA, L'Hérit. Stirp. Nov. i. t. 1 (1784).
angustifolia, *A. DC. in DC. Prod.* xvii. 29.—Madag.
diacantha, Lam. Encyc. i. 343 = Plectronia parviflora.
nova, *Blanco, Fl. Filip.* ed. I. 98.—Ins. Philipp.
nova, J. F. Gmel. Syst. 262 = tetracantha.
sarmentosa, *Benth. & Hook. f. Gen.* ii. 681.—Burma; Malaya.
sarmentosa, Vill. ex Vidal, Sinopa t. 65. f. C = tetracantha ?
scandens, Baill. Hist. Pl. vi. 13 = sarmentosa.
tetracantha, *Lam. Encyc.* i. 343.—Ind. or.

AZOPHORA, Neck. Elem. ii. 861 (1790) = **Rhinophora**, Linn.

AZORELLA, Lam. Encyc. i. 344 (1783). *UMBELLIFERAE*, Benth. & Hook. f. i. 874.
BOLAX, Comm. ex Juss. Gen. 236 (1789).
CHAMITIS, Banks, ex Gaertn. Fruct. i. 94. t. 22 (1788).
DICHOPETALUM, F. Muell. in Hook. Kew Journ. vii. (1855) 378. t. 11.
FRAGOSA, Ruiz & Pav. Prod. Fl. Per. 48. t. 34 (1794).
MICROSCIADIUM, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 468.
OSCHATZIA, Walp. Ann. i. 340 (1849).
PECTOPHYTUM, B. & K. Nov. Gen. et Sp. v. 28. t. 435 (1821).
POZOA, Lag. Gen. et Sp. Nov. 18 (1816).
apoda, *A. Gray, Bot. U. St. Expl. Exped.* i. 699.—Chili.
areticoides, *Willd. ex DC. Prod.* iv. 77.—Ecuador.
biloba, *Wedd. Chlor. And.* ii. 196.—Bolivia.
balsamo, *Clos, in C. Gay, Fl. Chil.* iii. 80.—Chili.
bryoides, *Phil. in Linnaea*, xxviii. (1856) 695.—Chili.
caespitosa, Cav. *Ic.* v. 57. t. 484.—Chili.
caespitosa, Vahl, Symb. Bot. iii. 48 = glebaria.
Chamitis, Pers. Syn. i. 308 = filamentosa.
cladorrhiza, *Pers. l. c.*—Peruv.
compacta, Labill. Nov. Holl. Pl. i. 75. t. 101 = Siebera compressa.
corymbosa, *Pers. Syn.* i. 308.—Peruv.
crassifolia, Pers. l. c. 308 = caespitosa.
crenata, *Pers. Syn.* 302.—Peruv.
cuneifolia, *F. Muell. ex Benth. Fl. Austral.* iii. 365.—Australia.
daucoides, Urv. in Mém. Soc. Linn. Par. iv. (1826) 618 = Oreomyrrhis andicola.
depauperata, *Phil. Fl. Atac.* 24.—Chili.
diapensioides, *A. Gray, Bot. U. St. Expl. Exped.* i. 702.—Peruv.
dichopetala, *Benth. Fl. Austral.* iii. 365.—Australia.
diversifolia, *Clos, in C. Gay, Fl. Chil.* iii. 86.—Chili.
filamentosa, *Lam. Encyc.* i. 344.—Terr. Magellan.
Gayana, *Phil. in Linnaea*, xxviii. (1856) 658.—Chili.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 347.—Chili.
glabra, *Wedd. Chlor. And.* 67.—Bolivia.
glebaria, *A. Gray, Bot. U. St. Expl. Exped.* i. 701.—Chili.
gummifera, Poir. Encyc. Suppl. i. 551 = caespitosa.
Hookeriana, Clos, in C. Gay, Fl. Chil. iii. 81 = caespitosa.
incisa, *Wedd. Chlor. And.* ii. 197.—Bolivia.
integrifolia, Raeusch. Nom. ed. III. 84 = filamentosa.
lanceolata, Labill. Nov. Holl. Pl. i. 74. t. 99 = Siebera Billardieri.
linearifolia, Cav. Ic. v. 57 = Siebera linearifolia.
lycopodioides, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 105. t. 3.—Reg. Magellan.
madreporica, *Clos, in C. Gay, Fl. Chil.* iii. 79.—Chili.
monantha, *Clos, l. c.*—Chili.
Muelleri, *Benth. Fl. Austral.* iii. 364.—Australia.
multifida, *Pers. Syn.* i. 308.—Peruv.
muscoides, *Phil. in Linnaea*, xxviii. (1856) 695.—Chili.
nervosa, *Phil. l. c.* 696.—Chili.
ovata, Labill. Nov. Holl. Pl. i. 74. t. 100 = Siebera Billardieri.
pedunculata, *Willd. ex DC. Prod.* iv. 78.—Ecuador.
pilgera, *Phil. in Linnaea*, xxviii. (1856) 652.—Chili.
pulvinata, *Wedd. Chlor. And.* ii. 194.—Ecuador.
Raunculus, *Urv. in Mém. Soc. Linn. Par.* iv. (1826) 618.—Reg. Magellan.
reniformis, Pers. Syn. i. 308.—Peruv.
Saxifraga, *Benth. Fl. Austral.* iii. 365.—Austral.
Selago, *Hook. f. Fl. Antarct.* 284.—Terr. Magell.; Ins. Kerguelen.
sessiliflora, *Phil. in Linnaea*, xxviii. (1856) 652.—Chili.
spinosa, *Pers. Syn.* i. 308.—Chili.
trincuspidata, Lam. Illustr. i. 189 = caespitosa.
trisaspidata, Poir. Encyc. Suppl. i. 551 = lycopodioides.
trifoliata, Walp. Ann. v. 69 = trifoliolata.
trifoliolata, *Hook. f. Fl. N. Zel.* i. (1853) t. 18.—N. Zel.
trifoliolata, *Clos, in C. Gay, Fl. Chil.* iii. 85. t. 30. f. 2.—Chili.
trifurcata, Hook. Ic. Pl. t. 539 = seq.
trifurcata, *Pers. Syn.* i. 308.—Reg. Magellan.
utricularia, *Griseb. in Gortt. Abh.* vi. (1854) 122.—Reg. Magellan.

AZURINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophul.).
aphylla, Fourr. l. c. 128 = V. aphylla.

B

BABACTES, DC. in Meisn. Gen. 302; Comm. 211 (1840); et Prod ix. 260 (1845) = **Chirita**, Buch.-Ham. (Gesnerac.).
oblongifolia, DC. l. c. = C. acuminata.

BABBAGIA, F. Muell. Rep. Babb. Exped. 21 (1858). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 61.
acroptera, *F. Muell. & Tate, in Trans. Roy. Soc. S. Austral.* vi. (1882) 108.—Australia.
dipteracoupra, *F. Muell. Rep. Babb. Exped.* 21.—Austral.
pentaptera, *F. Muell. & Tate, in Trans. Roy. Soc. S. Austral.* vi. (1882) 106.—Australia.
scleroptera, *F. Muell. in Wing, South Sc. Record*, (Nov. 1885).—Austral.

BABIANA, Ker-Gawl. in Bot. Mag. t. 676 (1809) *IRIDEAE*, Benth. & Hook. f. iii. 706.
ACAETE, Salisb. in Trans. Hort. Soc. i. (1812) 292.
angustifolia, *Sweet, Hort. Brit.* ed. I. 396; *Eckl. Vers.* 31.—Afr. austr.
atrocyanea, *Eckl. Vers.* 30.—Afr. austr.
atrodeltoides, *Eckl. l. c.*—Afr. austr.
Bainesii, *Baker, in Journ. Bot.* xiv. (1876) 335.—Afr. austr.
caesia, *Eckl. Vers.* 32.—Afr. austr.
caerulescens, *Eckl. l. c.*—Afr. austr.
cunefolia, *Baker, in Journ. Bot.* xiv. (1876) 335.—Afr. austr.
densiflora, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 348.—Afr. austr.
distichia, *Ker-Gawl. in Bot. Mag.* t. 926.—Afr. austr.
Dregei, *Baker, in Journ. Bot.* xiv. (1876) 336.—Afr. austr.
Ecklonii, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 348.—Afr. austr.
fimbriata, *Baker, in Journ. Linn. Soc.* xvi. (1878) 166.—Afr. austr.
flabelliformis, *Harv. ex Klatt, in Linnaea*, xxxv. (1867–68) 380.—Afr. austr.
flavocaesia, *Eckl. Vers.* 32.—Afr. austr.
fragrans, *Eckl. l. c.* 31.—Afr. austr.
Gladiola, *Spreng. ex Steud. Nom.* ed. II. i. 176.—Afr. austr.
hypogaea, *Burch. Trav.* ii. 589.—Afr. austr.
lilacina, *Eckl. Vers.* 31.—Afr. austr.
linearis, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 347.—Afr. austr.
maculata, *Klatt, l. c.* 349.—Afr. austr.
mucronata, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 284.—Afr. austr.
multiflora, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 351.—Afr. austr.
nana, *Spreng. Syst.* i. 156.—Afr. austr.
nervosa, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 284 = Antholyza nervosa.
obtusifolia, *Ker-Gawl. l. c.*—Ins. Mascar.
ochroleuca, *Sweet, Hort. Brit.* ed. I. 396.—Afr. austr.
parviflora, *Brehm, ex Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 350.—Afr. austr.
plicata, *Ker-Gawl. in Bot. Mag.* t. 576.—Afr. austr.
punctata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 348.—Afr. austr.
punices, *Eckl. Vers.* 32.—Afr. austr.
purpurea, Ker-Gawl. in Bot. Mag. sub t. 1019 = villosa.
pygmaea, *Spreng. ex Steud. Nom.* ed. II. i. 176.—Afr. austr.
pygmaea, Baker, in Journ. Linn. Soc. xvi. (1878) 165 = nana.
quadripartita, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 350.—Afr. austr.
reflexa, *Eckl. Vers.* 33.—Afr. austr.
reflexa, Ker-Gawl. Gen. Irid. 154 = plicata.
ringens, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 283.—Afr. austr.
rosea, *Eckl. Vers.* 32.—Afr. austr.
rubro-corulea, Pritz. Ind. Ic. 184 = seq.
rubrocyanea, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 284.—Afr. austr.
sambucina, *Ker-Gawl. l. c.*—Afr. austr.
scabra, *Brehm, ex Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 349.—Afr. austr.
secunda, *Ker-Gawl. Gen. Irid.* 154.—Afr. austr.
socotrana, *Hook. f. Bot. Mag.* t. 6585.—Ins. Socotr.
spathacea, *Ker-Gawl. in Bot. Mag.* t. 688.—Afr. austr.
Sprengelii, Baker, in Journ. Linn. Soc. xvi. (1878) 165 = pygmaea.
stellata, *Eckl. Vers.* 32.—Afr. austr.
stricta, *Ker-Gawl. in Bot. Mag.* t. 621, 687.—Afr. austr.
sulphurea, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 284.—Afr. austr.
tenuiflora, *Sweet, Hort. Brit.* ed. I. 396, nomen.—Afr. austr.
Thunbergii, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 283.—Afr. austr.
tubata, *Sweet, Hort. Brit.* ed. I. 396; *Eckl. Vers.* 33.—Afr. austr.
tubiflora, *Ker-Gawl. in Bot. Mag.* t. 680.—Afr. austr.
tubulosa, Ker-Gawl. Gen. Irid. 154 = tubiflora.
undulato-venosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 350.—Afr. austr.

BABIANA, —
villosa, *Ker-Gawl. in Bot. Mag.* t. 583.—Afr. austr.
villosula, *Ker-Gawl. ex Steud. Nom.* ed. II. i. 176.—Afr. austr.

BABINGTONIA, Lindl. in Bot. Reg. (1842) t. 10 = **Baeckea**, Linn. (Myrtac.).
Camphorosmae, Lindl. t. c. = Baeckea Camphorosmae.
corynophylla, F. Muell. Fragm. iv. 74 = Baeckea corynophylla.
crispiflora, F. Muell. l. c. = Baeckea crispiflora.
densifolia, F. Muell. l. c. = Baeckea densifolia.
gracilis, F. Muell. l. c. = Baeckea polyandra.
novo-anglica, F. Muell. l. c = Baeckea densifolia.
ovalifolia, F. Muell. l. c. = Baeckea ovalifolia.
pentagonantha, F. Muell. l.c. = Baeckea pentagonantha
pentandra, F. Muell. l. c. = Baeckea pentandra.
Preissiana, F. Muell. l. c. = Baeckea pentandra.
robusta, F. Muell. l. c. = Baeckea robusta.
subcuneata, F. Muell. l. c. = Baeckea subcuneata.
virgata, F. Muell. l. c. = Baeckea virgata.

BABIRON, Rafin. New Fl. Am. iv. 23 (1836) = **Daucus**, Tourn. (Umbell.), &c.
dichotomum, Rafin. l. c. 24 = Apium divaricatum.
divaricatum, Rafin. l. c. = Apium divaricatum.
pusillum, Rafin. l. c. 23 = D. pusillus.

BACA, Rafin. Princ. Somiol. 25 (1814) = **Boea**, Commers. (Gesneriac.).

BACASIA, Ruiz & Pav. Prod. Fl. Per. 105. t. 22 (1794) = **Barnadesia**, Mutis (Compos.).
corymbosa, Ruiz & Pav. Syst. Veg. 189 = Barn. corymbosa.
lanceolata, Ruiz & Pav. ex DC. Prod. vii. 8 = Barn. lanceolata.
spinosa, Ruiz & Pav. Syst. Veg. 188 = Chuquiraga spinosa.

BACCAUREA, Lour. Fl. Cochinch. 651 (1790). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 268.
ADENOCREPIS, Blume, Bijdr. 579 (1825).
CALYPTROON, Miq. Fl. Ind. Bat. Suppl. 471 (1860).
COCCOMELIA, Retzius. Syll. Ratisb. ii. (1832) 5.
HEDYCARPUS, Jack, in Trans. Linn. Soc. xiv. (1825) 118.
MICROSPALA, Miq. Fl. Ind. Bat. Suppl. 444 (1860).
PIERARDIA, Roxb. Hort. Beng. 28 (1814); Fl. Ind. ii. 254 (1832).
acuminata, *Muell. Arg. in DC. Prod.* xv. ii. 463.—Sumatra.
affinis, Muell. Arg. l. c. 459 = parviflora.
Barteri, *Muell. Arg. l. c.* 464.—Afr. trop. occ.
bornensis, *Muell. Arg. l. c.* 460.—Borneo.
bracteata, *Muell. Arg. l. c.* 466.—Malaya.
cauliflora, *Lour. Fl. Cochinch.* 661.—Cochinch.
costulata, *Muell. Arg. in DC. Prod.* xv. ii. 464.—Sumatra.
courtallensis, *Muell. Arg. l. c.* 459.—Ind. or.
dasystachya, *Muell. Arg. l. c.* 458.—Java.
deflexa, *Muell. Arg. l. c.* 463.—Java.
dulcis, *Muell. Arg. l. c.* 460.—Sumatra.
flaccida, *Muell. Arg. l. c.* 459.—Burma.
Griffoniana, *Muell. Arg. l. c.* 464.—Afr. occ.
javanica, *Muell. Arg. l. c.* 465.—Java.
lanceolata, *Muell. Arg. l. c.* 457.—Malaya.
macrocarpa, *Muell. Arg. l. c.* 459.—Malaya.
macrophylla, *Muell. Arg. l. c.* 464.—Malaya.
macrostachya, Wight, Ic. Pl. t. 1913 = sapida.
minutiflora, *Muell. Arg. in DC. Prod.* xv. ii. 463.—Java.
Motleyana, *Muell. Arg. l. c.* 461.—Malaya.
parviflora, *Muell. Arg. l. c.* 462.—Burma; Malaya.
prophyncha, Muell. Arg. l. c. 468 = sapida.
pubera, *Muell. Arg. l. c.* 458.—Sumatra.
pyrrhadasya, *Muell. Arg. l. c.* 462.—Sumatra.
racemosa, *Muell. Arg. l. c.* 461.—Java.
ramiflora, *Lour. Fl. Cochinch.* 661.—Cochinch.
sapida, Bedd. Fl. Sylv. t. 280 = courtallensis.
sapida, *Muell. Arg. in DC. Prod.* xv. ii. 459.—Ind. or.; Malaya.
Seemanni, *Muell. Arg. l. c.* 462.—Ins. Pacif.
stylaris, *Muell. Arg. l. c.* 458.—Ins. Pacif.
sumatrana, *Muell. Arg. l. c.* 466.—Sumatra.
sylvestris, *Lour. Fl. Cochinch.* 663.—Cochinch.
taitensis, *Muell. Arg. in DC. Prod.* xv. ii. 462.—Ins. Pacif.
tetrandra, *Muell. Arg. l. c.* 465.—Ins. Phillipp.
Wilkesiana, *Muell. Arg. l. c.* 461.—Ins. Pacif.

BACCHARIOIDES, Linn. Fl. Zeyl. 196 (1747) = seq.

BACCHARIS, Linn. Hort. Cliff. 404 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 386.
ACHYROBACCHARIS, Sch. Bip. in Walp. Rep. ii. 952 (1843).
ARRHENACHNE, Cass. in Dict. Sc. Nat. lii. 258 (1828).
MOLINA, Ruiz & Pav. Prod. Fl. Per. 111. t. 24 (1794).
PINGRAEA, Cass. in Dict. Sc. Nat. xli. 57 (1826).
POLYPAPPUS, Less. in Linnaea, iv. (1829) 314.
PTEROCLADIS, Lamb. ex G. Don, in Sweet, Hort. Brit ed. III. 849 (1839).
SERGILUS, Gaertn. Fruct. ii. 409. t. 174 (1791).
STEPHANANTHUS, Lehm. Ind. Sem. Hort. Hamb. (1826).—Cf. Flora, x. (1827) 96.
TURSENIA, Cass. in Dict. Sc. Nat. xxxvii. 480 (1825).
abiuthioides, Hook. & Arn. Bot. Beech. Voy. 57 = Tessaria abiuthioides.
acerosa, *Spreng. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 67.—Ecuador.
acutiloba, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 87 = incisa.
adnata, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 870 = Pluchea subdecurrens.
adscendens, *Pers. Syn.* ii. 424.—Peruv.
aegyptiaca, Forsk. ex DC. Prod. v. 450 = Pluchea Dioscoridis.
affinis, DC. l. c. 418 = subopposita.
affinis, Spreng. ex Baker, in Mart. Fl. Bras. vi. III. 94 = retusa.
Alamanni, DC. Prod. v. 402 = glutinosa.
alata, Hort. Berol. ex Lond. Hort. Brit. Suppl. ii. 612 = pedunculosa.
alaternoides, *H. B. & K. Nov. Gen. et Sp.* iv. 51.—Peruv.
alaternoides, Poepp. ex DC. Prod. v. 410 = Poeppigiana.
albida, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 41.—Reg. Argent.
alnifolia, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 264.—Peruv.
alpestris, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 122.—Bras.
alpestris, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 64 = rufescens.
alpina, *H. B. & K. Nov. Gen. et Sp.* iv. 48.—Peruv.
amplexicaulis, Wall. ex DC. Prod. v. 439 = Laggera flava.
amygdalina, Griseb. in Goett. Abh. xix. (1874) 175 = oxyodonta.
angulata, *Griseb. l. c.* xxiv. (1879) 180.—Reg. Argent.
angustifolia, Desf. Cat. Hort. Par. ed. III. 165 = B. Pingraea.
angustifolia, *Michx. Fl. Bor. Am.* ii. 125.—Am. bor.
anomala, *DC. Prod.* v. 408.—Bras.
aphylla. *DC. l. c.* 424.—Bras.
apiculata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 74 = trinervia.
arbores, *Linn. Mant.* 384.—Ins. Johann. (Afr. or.).
arborescens, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 428 = arborea.
arbutifolia, *Vahl, Symb. Bot.* iii. 97.—Peruv.
arctostaphyloides, *Baker, in Mart. Fl. Bras.* vi. III. 88.—Bras.
arenaria, *Baker, l. c.* 62.—Bras.
aretioides, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
aretioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 67.—Ecuador.
arguta, Gill. ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 35 = racemosa.
argyrea, Pers. Syn. ii. 435 = Psiadia trinervia.
artemisioides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 41.—Reg. Argent.
articulata, *Pers. Syn.* ii. 435.—Bras.
arunca, Schrank, ex DC. Prod. v. 381 = Conyza Gouani.
asperifolia, Benth. Fl. Hartw. 86 = Conyza asperifolia.
asperula, *Schau. in Linnaea*, xix. (1847) 734.—Mexic.
astoensis, *H. B. & K. Nov. Gen. et Sp.* iv. 67.—Ecuador.
asteroides, Colla, in Mem. Acc. Torin. xxxvii. (1835) 14. t. 25. f. 1 (= *Tripolium conspicuum* ?).—Chili.
attenuata, *D. Don, ex Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 34.—Uruguay.
auriculata, Wall. ex DC. Prod. v. 449 = Blumea hieracifolia.
avicenniaefolia, *DC. l. c.* 414.—Bras.

BACCHARIS—continued.
axillaris, *DC. Prod.* v. 407.—Bras.
axillaris, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 94 = halimifolia.
bahiensis, *Baker, l. c.* 88.—Bahia.
Bairdii, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 30.—Uruguay.
Baldwinii, Griseb. in Goett. Abh. xxiv. (1879) 181 = tridentata.
Balduinii, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 38 = rufescens.
balsamifera, *Benth. Pl. Hartw.* 202.—N. Granat.
balsamifera, Stokes, Bot. Mat. Med. iv. 204 = Blumea balsamifera.
banksiaefolia, Bert. in Merc. Chil. (1829).—Cf. Bull. Férussac, xx. (1830) 108 = Tessaria absinthioides.
berteroifolia, Walp. in Linnaea, xiv. (1840) 814 = brachylaenoides.
berberifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 67.—Ecuador.
Bezanillaeana, *Remy, in C. Gay, Fl. Chil.* iv. 98.—Chili.
bifrons, *Baker, in Mart. Fl. Bras.* vi. III. 54.—Bras.
Bigelovii, *A. Gray, in Torr. Bot. Mex. Bound.* 84.—Am. bor.; Mexic.
Blanchetiana, DC. Prod. vii. 281 = Moquinia flavescens.
bogotensis, *H. B. & K. Nov. Gen. et Sp.* iv. 61.—N. Granat.
brachylaenoides, *DC. Prod.* v. 421.—Bras.
brachyphylla, *A. Gray, Pl. Wright.* ii. 83.—Am. bor. occ.
bracteata, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 35 = dracunculifolia.
bracteolata, *DC. Prod.* v. 418.—Hab.?
brasiliana, Linn. Sp. Pl. ed. II. 1906 = Vernonia scabra.
brevifolia, *DC. Prod.* v. 409.—Bras.
brevisata, DC. l. c. 409 = serrulata.
bryoides, *Pers. Syn.* ii. 435.—Reg. Magellan.
buddleoides, *H. B. & K. Nov. Gen. et Sp.* iv. 64.—Peruv.
buplenroides, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 86 = ligustrina.
Burchellii, *Baker, in Mart. Fl. Bras.* vi. III. 44.—Bras.
buxifolia, *Pers. Syn.* ii. 424.—Peruv.
cacalioides, *Spreng. Nachr. i. Bot. Gart. Halle*, 14.—Hab.?
caerulescens, DC. Prod. v. 402 = glutinosa.
caespitosa, *Pers. Syn.* ii. 425.—Peruv.
callaetenioides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 265.—Chili.
callilepinos, *Griseb. in Goett. Abh.* xix. (1874) 177.—Reg. Argent.
calvescens, *DC. Prod.* v. 413.—Bras.
camporum, *DC. l. c.* 399.—Bras.
Candolleana, *Steud. Nom.* ed. II. i. 177.—Peruv.
canescens, *Spreng. Syst.* iii. 466.—Reg. Argent.
capitata, *H. B. & K. Nov. Gen. et Sp.* iv. 53.—Am. austr.
caprariaefolia, *DC. Prod.* v. 416.—Bras.
casinefolia, *DC. l. c.* 413.—Bras.
cassinoides, DC. l. c. = casinefolia.
cataphracta, *Spreng. Syst.* iii. 462.—Peruv.
Cebalotes, *DC. Prod.* v. 421.—Bras.
Chilco, *H. B. & K. Nov. Gen. et Sp.* iv. 56.—N. Granat.
chilensis, Wehnm. in Syll. Ratisb. ii. (1826) 21 = Podanthrus ovatifolius.
chiletoria, Sch. Bip. in Bonplandia, iv. (1856) 55 = B. Chilquilla.
Chilquilla, *DC. Prod.* v. 419.—Chili.
chimenia, Lour. Fl. Cochinch. ii. 494 = Blumea chimenia.
ciliata, *Gard. in Hook. Lond. Journ. Bot.* iv. (1845) 122.—Bras.
cinerea, *DC. Prod.* v. 400.—Bras.
cinnamomifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 65.—Ecuador.
Claussenii, *Baker, in Mart. Fl. Bras.* vi. III. 66.—Bras.
Claussenii, Sch. Bip. ex Baker, l. c. 88 = casinefolia.
cochinchinensis, *Spreng. Syst.* iii. 466.—Cochinch.
cognata, *DC. Prod.* v. 418.—Bras.
colletioides, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 75 = Lundii.
concava, *Pers. Syn.* ii. 425.—Peruv.; Chili.
confertifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 55.—Mexic.
confertifolia, Colla, in Mem. Acc. Torin. xxxvii. (1835) 13. t. 25. f. 2 = B. Chilquilla.

BACCHARIS:—
congesta, *DC. Prod.* v. 410.—Mexic.
consanguinea, DC. l. c. = pilularia.
conyzoides, *DC. l. c.* 423.—Bras.
coridifolia, *DC. l. c.* 423.—Bras.
Corydalis, DC. ex Sch. Bip. in Linnaea, xxx. (1859–60) 181 = casabifolia.
corymbosa, *Meyen, Reise*, i. 811.—Chili.
corymbosa, *Pers. Syn.* ii. 424.—Peruv.
crenulata, *Sprengel. Syst.* iii. 463.—Bras.
crispa, Spreng. l. c. 466 = genistelloides.
cultrata, *Baker, in Mart. Fl. Bras.* vi. III. 93.—Bras.
cuneata, DC. Prod. v. 408 = conferta.
cuneifolia, DC. l. c. 406 = magellanica.
cuneifolia, Mœnch, Meth. 574 = halimifolia.
cuneifolia, Wendl. ex Steud. Nom. ed. II. i. 177, 179 = Wendlandii.
curvifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 67 = rufescens.
cupressiformis, Pers. Syn. ii. 425 = Lepidophyllum cupressiforme.
cylindrica, *DC. Prod.* v. 426 = genistelloides.
daphnoides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 24 = casabifolia.
Darwinii, *Hook. & Arn. l. c.* 39.—Patagon.
deltoidea, *Baker, in Mart. Fl. Bras.* vi. III. 90.—Reg. Argent.
densiflora, *Wedd. Chlor. And.* i. 175.—Bolivia.
denticulata, *D. Don, in DC. Prod.* vii. 282.—Patagon.
dentifolia, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 66 = recurvata.
dentosa, Mart. ex Baker, l. c. 91 = subopposita.
depauperata, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 120 = serrulata.
dependens, *Pers. Syn.* ii. 424.—Peruv.
dioica, *Vahl, Symb. Bot.* iii. 98.—Ins. Carib.
Dioscoridis, Bové, ex DC. Prod. v. 382 = Conyza aegyptiaca.
Dioscoridis, Linn. Cent. Pl. i. 27; Amoen. Acad. iv. 289 = Pluchea hirsuta.
diptera, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 88 = articulata.
discolor, *Baker, l. c.* 48.—Bras.
discolor, Sch. Bip. ex Baker, l. c. 54 = B. Lychnophora.
distichea, Sch. Bip. ex Baker, l. c. 53 = calvescens.
divergens, *DC. Prod.* v. 400.—Mexic.
diversifolia, Spreng. ex Sch. Bip. in Walp. Rep. ii. 956 = Erigeron Sprengelii.
Doniana, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 26.—Bras.
Douglasii, *DC. Prod.* v. 400.—Calif.
dracunculifolia, *DC. l. c.* 421.—Bras.
effusa, *Grisb. in Gott. Abh.* xix. (1874) 177.—Reg. Argent.
elaeagnoides, *Steud. ex Sch. Bip. in Linnaea*, xxx. (1859–60) 181.—Bras.
elaeoides, *Remy, in C. Gay, Fl. Chil.* iv. 88.—Chili.
elegans, *H. B. & K. Nov. Gen. et Sp.* iv. 60. t. 324.—Mexic.
elliptica, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 33.—Bras.
emarginata, *Pers. Syn.* ii. 424.—Peruv.
Emoryi, *A. Gray, in Torr. Bot. Mex. Bound.* 88.—Am. bor. occ.
ericoides, Pers. Syn. ii. 425 = Lisbum ericoides.
erigeroides, *DC. Prod.* v. 415.—Bras.
ericcada, *DC. l. c.* 415.—Bras.
erioptera, Benth. in Ann. Nat. Hist. ii. (1839) 441 = Perocoulon virgatum.
eupatorioides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 29.—Chili.
farinosa, Spreng. in Roch & Grub. Allgem. Encyc. vi. (1831) 57 = glutinosa.
fasciculosa, *Kiatt, in Leopoldina*, xx. (1884) 91.—Mexic.
fastigiata, *Baker, in Mart. Fl. Bras.* vi. III. 39.—Bras.
ferruginea, *Pers. Syn.* ii. 425.—Peruv.
Feuillei, *DC. Prod.* v. 408.—Peruv.
filifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 39 = polyphylla.
fiabellata, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 28.—Chili.
flexuosa, *Baker, in Mart. Fl. Bras.* vi. III. 53.—Bras.
flexuosa, Mart. ex Baker, l. c. 53 = calvescens.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iv. 64.—N. Granat.
floribunda, *Phil. in Linnaea*, xxxiii. (1864–65) 145.—Chili.
foetida, Linn. Sp. Pl. 861 = Pluchea camphorata.
foliosa, *Gill. ex Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 30.—Chili.
frigida, *H. B. & K. Nov. Gen. et Sp.* iv. 49.—Peruv.
fuchsiaefolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 38 = argyrea.
fuliginosa, H. B. & K. Nov. Gen. et Sp. iv. 68 = Bigelovia faliginea.
fusca, *Turcz. in Bull. Soc. Mat. Mosc.* xxiv. (1851) II. 87.—Ecuador.
Gaudichaudiana, DC. Prod. v. 424 = articulata.
genistelloides, *Pers. Syn.* ii. 425.—Peruv.
genistelloides, Poepp. ex DC. Prod. v. 425 = sagittalis.
genistifolia, *DC. l. c.* 423.—Bras.
Gibertii, *Baker, in Mart. Fl. Bras.* vi. III. 52.—Bras.
Gillesii, *A. Gray, in Proc. Am. Acad.* v. (1861) 123.—Chili.
glabra, *Spreng. Syst.* iii. 466.—Reg. Argent.
glauca, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* (1843) 363.—Chili.
Glaziovii, *Baker, in Mart. Fl. Bras.* vi. III. 44.—Bras.
glomeruliflora, *Pers. Syn.* ii. 423.—Am. bor.; Ins. Bernud.
glutinosa, Hook. & Arn. Bot. Beech. Voy. 31 = Douglasii?
glutinosa, *Pers. Syn.* ii. 425.—Am. bor. et austr. occ.
gnaphalioides, *DC. Prod.* v. 415.—Bras.
gnaphalioides, Spreng. Syst. iii. 461 = Lucilia acutifolia.
gouldiflia, *H. B. & K. Nov. Gen. et Sp.* iv. 61.—Ecuador.
gracilis, *DC. Prod.* v. 423.—Bras.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* iv. 58.—Ecuador.
gratissima, Blume, ex DC. Prod. vi. 447 = Blumea appendiculata.
graveolens, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
grindeliaefolia, *Wedd. Chlor. And.* i. 177.—Peruv.
grisea, *Baker, in Mart. Fl. Bras.* vi. III. 65.—Bras.
Grisebachii, *Hieron. in Bot. Acad. Nac. Córdova*, iv. (1881) 86.—Reg. Argent.
guianensis, *Schomb. Fauna & Flora Guy.* 1135.—Guiana.
Haenkei, DC. Prod. v. 401 = Douglasii.
halimifolia, *Linn. Sp. Pl.* 860.—Am. bor.; Ind. occ.
halimifolia, Mœnch, Meth. Suppl. 235 = Conyza ivaefolia.
halimimorpha, *DC. Prod.* v. 411.—Bras.
hambatensis, *H. B. & K. Nov. Gen. et Sp.* iv. 62.—Ecuador.
Havardi, *A. Gray, Syn. Fl. N. Am.* i. II. 234.—Texas.
helichrysoides, *DC. Prod.* v. 415.—Bras.
hemiptilonodes, *Bush, Ind. Gen. DC. Prod.* i. p. v.—Bras.
heterophylla, *H. B. & K. Nov. Gen. et Sp.* iv. 62.—Mexic.
heterophylla, Spreng. Syst. iii. 464, pro parte = sebastianopolitana.
hieraciifolia, Lam. Encyc. i. 347 = Conyza Gouani.
hieraciifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 139.—Mexic.
hirta, *DC. Prod.* v. 405.—Bras.
hirtella, *DC. l. c.* 418.—Mexic.
Hookeriana, DC. Prod. v. 414 = Aplopappus Hookerianus.
humifusa, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 311 = Melalema humifusa.
humiflusa, *H. B. & K. Nov. Gen. et Sp.* iv. 48. t. 322.—Ecuador.
humilis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 93.—Bras.
Huydobriana, Remy, in C. Gay, Fl. Chil. iv. 90 = B. Phagnas
hypericifolia, *Baker, in Mart. Fl. Bras.* vi. III. 69.—Bras.
hyssopifolia, E. Mey. ex DC. Prod. v. 338 = Nidorella foliosa.
illicifolia, Lam. Encyc. i. 345 = Brachylaena dentata.
Imlana, *Benth. Pl. Hartw.* 302.—N. Granat.
ilinita, *DC. Prod.* v. 412.—Bras.
imanensis, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 31 = trinervis.
incarum, Wedd. Chlor. And. i. 170 = microphylla.
incisa, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 29.—Reg. Argent.
indica, Linn. Sp. Pl. 861 = Pluchea indica.
intermedia, *DC. Prod.* v. 411.—Chili.
intermixta, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 34.—Bras.
irasinoides, *H. B. & K. Nov. Gen. et Sp.* iv. 63.—N Granat.
ivaefolia, Blanco, Fl. Filip. ed. I. 627 = Conyza viscidula.
ivaefolia, Linn. Sp. Pl. 860 = Conyza ivaefolia.
juncea, *Desf. Cat. Hort Par.* ed. III. 168.—Bras.
juniciformis, *DC. Prod.* v. 426.—Bras.
lacrymans, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 94 = retusa.
laevis, *Spreng. Syst.* iii. 466.—Cochinch.
lanceolata, *H. B. & K. Nov. Gen. et Sp.* iv. 63.—Peruv.
lancifolia, *Less. in Linnaea*, ix. (1834) 266.—Mexic.
lanuginosa, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 32 = helichrysoides.
lateralis, *Baker, in Mart. Fl. Bras.* vi. III. 100.—Bras.
latifolia, *Pers. Syn.* ii. 424.—Peruv.
laxa, *Gardn. in Hook. Lond. Journ. Bot* iv. (1845) 121.—Bras.
Lechleri, *Sch. Bip. in Lechl. Berb. Am. Austr.* 53, nomen.—Chili.
ledifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 50.—Peruv.
lepidophylla, DC. Prod. v. 427 = Helichrysum becchaeroides.
leptocephala, DC. l. c. 413 = rufescens.
leptophylla, DC. l. c. 428 = rufescens.
leptosperm oides, DC. l. c. 421 = dracunculifolia.
Lessingiana, *DC. l. c.* 414.—Bras.
leucanthemifolia, Burm. f. ex DC. l. c. 387 = Conyza pinnatifolia.
leucolepis, Sch. Bip. in Linnaea, xxii. (1849) 570 = helichrysoides.
leucopappa, DC. Prod. v. 415 = helichrysoides.
ligustrina, *DC. l. c.* 421.—Bras.
limnophila, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 60 = ligustrina.
linearifolia, Pers. Syn. ii. 425 = Psiadia trinervia.
linearis, *Pers. l. c.*—Chili.
linearis, Poepp. ex Baker, in Mart. Fl. Bras. vi. III. 59 = serrulata.
lingulata, Kunze, in Linnaea, vi. (1831) 147 = rosmarinifolia.
linifolia, *DC. Prod.* v. 420.—Mexic.
lumifolia, Meyen, Reise, i. 811 = rosmarinifolia.
linifolia, *Phil. in Linnaea*, xxxiii. (1864–65) 145.—Chili.
longifolia, *DC. Prod.* v. 409.—Mexic.
longipes, *Kunze, ex DC. l. c.* 401.—Chili.
lorentzii, *Benth. Pl. Hartw.* 135.—Peruv.
lucida, Meyen, Reise, i. 460 = Vernonia phyllicaeformis.
Lundii, *DC. Prod.* v. 404.—Bras.
Lychnophora, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 35.—Bras.
lycioides, *Remy, in C. Gay, Fl. Chil.* iv. 99.—Chili.
lycopodioides, Pers. Syn. ii. 425 = Enotrix juniperifolia.
Macraei, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 32.—Chili.
macrantha, *H. B. & K. Nov. Gen. et Sp.* iv. 54.—Ecuador.
macrodonta, *DC. Prod.* v. 416.—Bras.
madagascariensis, Pers. Syn. ii. 423 = Psiadia madagascariensis.
magellanica, *Pers. l. c.* 425.—Reg. Magell.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 538, nomen.—Bolivia.
marcetiaefolia, *Benth. Pl. Hartw.* 209.—Reg. Granat.
marginalis, *DC. Prod.* v. 402.—Peruv.; Chili.
maritima, *Baker, in Mart. Fl. Bras.* vi. III. 73.—Bras.
maxima, *Baker, l. c.* 80.—Bras.
medullosa, DC. Prod. v. 406 = serrulata.
megacephala, D. Don, ex Baker, in Mart. Fl. Bras. vi. III. 79 = triplinervis.
megapotamica, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 33 = rufescens.
megapotamica, *Spreng. Syst.* iii. 461.—Bras.
melastomaefolia, Hook. & Arn. in Hook. Journ. Bot iii. (1841) 28 = oxyodonta.
mesoneura, DC. Prod. v. 413 = retusa.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 295.—Peruv.
microcephala, *DC. Prod.* v. 423.—Bras.
microdonta, *DC. l. c.* 416.—Bras.

BACCHARIS:—
microphylla, DC. Prod. v. 406 = Candolleana.
microphylla, *H. B. & K. Nov. Gen. et Sp.* iv. 58.—N. Granat.
micropoda, *Baker, in Mart. Fl. Bras.* vi. III. 88.—Bras.
micropteras, *Baker, l. c.* 42.—Bras.
milleflora, DC. Prod. v. 426 = genistelloides.
minutiflora, *Mart. ex Baker, in Mart. Fl. Bras.* vi. III. 69.—Bras.
mollis, *H. B. & K. Nov. Gen. et Sp.* iv. 58.—Ecuador.
monantha, Sch. Bip ex Baker, in Mart. Fl. Bras. vi. III. 69 = minutiflora.
montana, DC. Prod. v. 404 = macrodonta.
montana, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 78 = Schultzii.
montevidensis, Sch. Bip. ex Baker, l. c. 57 = cordifolia.
montevidensis, Spreng. Syst. iii. 460 = Vernonia nitidula.
mucronata, Hook. & Arn. Bot. Beech. Voy. 80 = Aplopappus Hookerianus.
mucronata, *H. B. & K. Nov. Gen. et Sp.* iv. 60.—Mexic.
Muelleri, *Baker, in Mart. Fl. Bras.* vi. III. 61.—Bras.
multiflora, *H. B. & K. Nov. Gen. et Sp.* iv. 59.—Mexic.
multisulcata, *Baker, in Mart. Fl. Bras.* vi. III. 45.—Bras.
myricaefolia, DC. Prod. v. 405 = semiserrata.
myriocephala, *Baker, in Mart. Fl. Bras.* vi. III. 93.—Bras.
myriocephala, DC. Prod. v. 426 = genistelloides.
myrsinites, *Pers. Syn.* ii. 424. n. 25.—Ins. S. Doming.
myrsinoides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 32 = brevifolia.
myrtilloides, *Griseb. in Goett. Abh.* xix. (1874) 176.—Reg. Argent.
nana, *D. Don, ex Baker, in Mart. Fl. Bras.* vi. III. 56.—Patagon.
Neaei, *DC. Prod.* v. 411.—Am. merid.
neriifolia, Linn. Sp. Pl. 860 = Brachylaena neriifolia.
nervosa, *DC. Prod.* v. 399.—Ind. occ.
nitida, *Pers. Syn.* ii. 425.—Peruv.
nitida, Wall. Cat. n. 3048 = Blumea chinensis.
nivalis, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Chili.
notoserglia, *Griseb. in Goett. Abh.* xxiv. (1879) 183.—Reg. Argent.
nuda, DC. Prod. v. 424 = aphylla.
oblongifolia, *Pers. Syn.* ii. 424; *Spreng. Syst.* iii. 460.—Chili.
obovata, *DC. Prod.* v. 408.—Peruv.
obovata, Hook. & Arn. Bot. Beech. Voy. 80 = umbelliformis.
obtusifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 51.—Peruv.
ochracea, *Spreng. Syst.* iii. 460.—Reg. Argent.
odorata, *H. B. & K. Nov. Gen. et Sp.* iv. 52.—N. Granat.
odorata, Spreng. in Ersch & Grub. Allgem. Encyc. vii. (1821) 27 = casinefolia.
olaefolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 86 = calvescens.
oligantha, DC. ex Thoms. in Journ. Linn. Soc. ix. (1867) 348 = Catamixia baccharoides.
oligophylla, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 33 = Pohlii.
opuntioides, *Mart. ex Baker, l. c.* 89.—Bras.
Orbignyana, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 237.—Chiquitos.
organensis, *Baker, in Mart. Fl. Bras.* vi. III. 74.—Bras.
orgyalis, *DC. Prod.* v. 416.—Bras.
ostenbaisii, *Sch. Bip. ex Hensl. Biol. Centr. Am. Bot.* ii. 180.—Mexic.
oriasbainia, Klatt, in Leopoldina, xx. (1884) 91 = resinosa.
orosacensis, *DC. Prod.* v. 422.—Peruv.
ovalis, Pers. Syn. ii. 424 = Pluchea ovalis.
ovata, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 33.—Chili.
ovata, Sieber, ex DC. Prod. v. 450 = Pluchea ovalis.
oxyodonta, *DC. l. c.* 404.—Bras.
oxyphylla, *DC. l. c.* 406.—Peruv.
paniculata, *DC. l. c.* 430.—Chili.
parviflora, *Less. in Linnaea*, v. (1830) 146.—Mexic.
parviflora, Pers. Syn. ii. 425 = marginalis.
parviflora, Poir. Encyc. Suppl. i. 636 = bracteolata.
parvifolia, *DC. Prod.* v. 424.—Peruv.

BACCHARIS:—
patagonica, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 29.—Reg. Magellan.
patens, *Baker, in Mart. Fl. Bras.* vi. III. 52.—Bras.
paucidentata, DC. Prod. v. 420 = rufescens.
paucidentata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 74 = dracunculifolia.
pauciflosculosa, *DC. Prod.* v. 418.—Bras.
pedalis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 64 = rufescens.
pedicularis, *DC. Prod.* v. 407.—Chili.
pentaptera, *DC. l. c.* 425.—Bras.
Pentlandii, *DC. l. c.* 416.—Bolivia.
pentziaefolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 96.—Bras.
petiolata, *DC. Prod.* v. 418.—Chili.
phagnaloides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 59 = microdonta.
phillippensis, *H. B. & K. Nov. Gen. et Sp.* iv. 55.—Ecuador.
phlycaefolia, *DC. l. c.* 415.—Bras.
phyliciaeformis, Meyen, Reise, ii. 31 = Vernonia phyliciaeformis.
phyllocides, *H. B. & K. Nov. Gen. et Sp.* iv. 49.—Peruv.
phyteumoides, *DC. Prod.* v. 425.—Bras.
pilularis, *DC. l. c.* 407.—Am. bor. occ.
Pingraea, *DC. Prod.* v. 490.—Chili.
Pingraea, Nutt. in Trans. Am. Phil. Soc. vii. (1841) 387 = glutinosa.
pinnatifida, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 227.—Patagon.
platensis, *Spreng. Syst.* iii. 465.—Reg. Argent.
platyptera, *DC. Prod.* v. 409.—Bras.
Plummerae, *A. Gray, in Proc. Am. Acad.* xv. (1880) 48.—Calif.
Poeppigiana, *DC. Prod.* v. 410.—Chili.
pollidoi, *Griseb. in Goett. Abh.* xix. (1874) 176.—Reg. Argent.
polyantha, *H. B. & K. Nov. Gen. et Sp.* iv. 64.—Peruv.; Mont. Vid.!
polyanthoides, Spreng. Syst. ii. 464 = helichrysoides.
polycephala, *Wedd. Chlor. And.* i. 173.—Bolivia.
polycephala, Sch. Bip. in Linnaea, xxx. (1859–60) 181 = vernonioides.
polygalaefolia, *H. B. & K. Nov. Gen. et Sp.* iv. 49.—Peruv.
polygona, *Baker, in Mart. Fl. Bras.* vi. III. 46.—Bras.
polylepis, Sch. Bip. ex Baker, l. c. 98 = subcapitata.
polyneura, Sch. Bip. ex Baker, l. c. 100 = reticularia.
polyphylla, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 87.—Bras.
polyphylla, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 69 = minutiflora.
polyptera, *DC. Prod.* v. 426.—Bras.
potosina, *A. Gray, in Proc. Am. Acad.* xv. (1880) 33.—Mexic.
prenanthoides, *Baker, in Mart. Fl. Bras.* vi. III. 84.—Bras.
prinoides, *H. B. & K. Nov. Gen. et Sp.* iv. 56.—Reg. Amazon.
proctorana, Baldw. in Hook. Journ. Bot. iii. (1841) 28 = Baldwinii.
prostrata, *Pers. Syn.* ii. 424.—Peruv.
pruniflora, *H. B. & K. Nov. Gen. et Sp.* iv. 63.—Peruv.
psiloclyx, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 48 = eriocladia.
ptarmicoides, DC. Prod. v. 419 = Bigelovii?
Pteroculon, Boj. ex DC. l. c. 455 = Pteroscalon Bojeri.
pteronioides, *DC. l. c.* 410.—N. Mexic.
puberula, *DC. l. c.* 401.—Bras.
pulchella, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 532, nomen.—Bolivia.
pulverulenta, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 237.—Peruv.
punctigera, DC. Prod. v. 404 = Lundii.
punctulata, DC. l. c. 405 = oxyodonta.
pyramidalis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 130.—Bras.
quadranguleris, Meyen, Reise, i. 460 = Lepidophyllum Meyeni.
quadrinervis, *Pers. Syn.* ii. 423.—Am. austr.
quinquenervia, *DC. Prod.* v. 399.—Bras.
quitensis, *H. B. & K. Nov. Gen. et Sp.* iv. 57.—Ecuador.

BACCHARIS:—
racemosa, *DC. Prod.* v. 401.—Chili.
radicans, DC. l. c. 416 = gnaphalioides.
Radlin, *Phil. in Linnaea*, xxxiii. (1864–65) 146.—Chili.
ramiflora, *A. Gray, in Proc. Am. Acad.* xv. (1880) 33.—Mexic.
ramosissima, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 84 = retusa.
ramulosa, A. Gray, Fl. Taarb. 301 = pteronioides.
rapunculoides, E. Mey. ex DC. Prod. v. 393 = Nidorella resedifolia.
recurvata, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 88.—Bras.
refracta, *Burch. ex Baker, in Mart. Fl. Bras.* vi. III. 64.—Bras.
Regnellii, *Sch. Bip. in Linnaea*, xxii. (1849) 571.—Bras.
resedifiora, Less. ex Baker, in Mart. Fl. Bras. vi. III. 58 = erigeroides.
retusa, *Steud. & Hochst. ex DC. Prod.* vii. 263.—Arab.
resinosa, Hook. & Arn. Bot. Beech. Voy. 81 = concava.
resinosa, *H. B. & K. Nov. Gen. et Sp.* iv. 52.—Am. austr.
retamoides, *Phil. Sert. Mend. Alt.* 22.—Chili.
reticularia, *DC. Prod.* v. 409.—Bras.
reticulata, Pers. Syn. ii. 425 = genistelloides.
retusa, *DC. Prod.* v. 412.—Bras.
revoluta, *H. B. & K. Nov. Gen. et Sp.* iv. 50.—Peruv.
rhetinodes, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 265.—Chili.
rhetinoides, *H. B. & K. Nov. Gen. et Sp.* iv. 66.—Peruv.; Bras.
rhomboidalis, *Remy, in C. Gay, Fl. Chil.* iv. 99.—Chili.
Riedelii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 82 = casinefolia.
rigida, Hook. & Arn. Bot. Beech. Voy. 57 = oblongifolia.
riparia, *H. B. & K. Nov. Gen. et Sp.* iv. 65.—Ecuador.
riparia, Poepp. ex DC. Prod. v. 418 = sessiliflora.
rivularis, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 88 = macrodonta.
Rohriana, Spreng. in Ersch & Grub. Allgem. Encyc. vii. (1821) 27 = dioica.
Rorainae, *Schomb. Fauna et Fl. Guy.* 1078, nomen.—Guiana.
rosmarinifolia, *Hook. & Arn. Bot. Beech. Voy.* 80.—Chili.
rotundifolia, *Spreng. Syst.* iii. 465.—Reg. Argent.
rufescens, *Spreng. l. c.* iii. 464.—Reg. Argent.
rufidula, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 81 = vernonioides.
rugosa, Willd. ex DC. Prod. v. 39 = Vernonia patens.
rupicola, *H. B. & K. Nov. Gen. et Sp. iv.* 52.—N. Granat.
sagittalis, *DC. Prod.* v. 425.—Chili.
salicifolia, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 387 = salicina.
salicifolia, *Pers. Syn.* ii. 425.—Peruv.
salicina, Rothr. in Wheeler, Rep. vi. 156 = Emoryi.
salicina, *Torr. & Gray, Fl. N. Am.* ii. 258.—Am. bor. occ.
saligna, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 60 = ligustrina.
saltensis, *Baker, l. c.* 98.—Bras.
Salvia, Lour. Fl. Cochinch. ii. 494 = Blumea balsamifera.
Salzmannii, *DC. Prod.* v. 409.—Bras.
sarothra, Phil. in Anal. Univ. Chil. i. (1861) 52; et in Linnaea, xxxiii. (1864–65) 147 = Heterothalamus spartioides.
sarothroides, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 211.—Calif.
scabra, *Pers. Syn.* ii. 424.—Ecuador.
scandens, *Less. in Linnaea*, v. (1830) 146.—Mexic.
scandens, *Pers. Syn.* ii. 424.—Peruv.
Schiedeana, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 33.—Bras.
Schlechtendahlii, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 50 = tarchonanthoides.
Schomburgkii, *Baker, l. c.* 99.—Guiana.
Schultzii, *Baker, l. c.* 78.—Bras.
scolopendra, *Hook. Ic. Pl.* i. t. 68.—Ecuador.
scoparia, *Sw. Fl. Ind. Occ.* iii. 1269.—Jamaica.
scoparioides, *Griseb. Cat. Pl. Cub.* 196.—Cuba.
scripta, *Griseb. in Goett. Abh.* xix. (1874) 178.—Reg. Argent.

BACCHARIS:—
sebastianopolitana, *Baker, in Mart. Fl. Bras.* vi. III. 66.—Bras.
Seemanni, *A. Gray, in Proc. Am. Acad.* xv. (1880) 33.—Mexic.
Sellot, *Baker, in Mart. Fl. Bras.* vi. III. 68.—Bras.
semiserrata, DC. Prod. v. 404. n. 38 = hemiptinodes.
semiserrata, DC. l. c. 419. n. 149 = Sternbergiana.
senegalensis, Pers. Syn. ii. 424 = Vernonia senegalensis.
sericula, Mart. in Flora, xxi. (1838) ii. Beibl. 61 = casuinefolia.
sergiloides, *A. Gray, in Pacif. Rail. Rep.* iv. 101.—Am. bor. occ.
serratifolia, *DC. Prod.* v. 408.—Mexic.
serratifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 59.—Mexic.; Ind. occ.
serrulata, *Pers. Syn.* ii. 423.—Bras.
sessiliflora, Michx. Fl. Bor. Am. ii. 135 = glomeruliflora.
sessiliflora, *Vahl, Symb. Bot.* iii. 97.—Bras.
sessiliifolia, *DC. Prod.* v. 418.—Chili.
sinensis, *Lour. Fl. Cochinch.* ii. 604.—Cochinch.
sinuata, *H. B. & K. Nov. Gen. et Sp.* iv. 58.—Peruv.
Solizzi, *Remy, in C. Gay, Fl. Chil.* iv. 98.—Chili.
Solisi, *Phil. in Anal. Univ. Chil.* (1856) ii. 830.—Chili.
sordescens, *DC. Prod.* v. 406.—Mexic.
spartea, *Benth. Bot. Voy. Sulph.* 114.—Am. centr.
spartioides, C. Gay, Fl. Chil. iv. 102 = Heterothalamus spartioides.
spathulata, *Schau. in Linnaea*, xix. (1847) 734.—Mexic.
spathulata, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 226.—Peruv.
speciosa, *DC. Prod.* v. 399.—Ins. Guadalup.
sphaerocephala, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 36.—Chili.
spiropoda, F. Muell. Fragm. i. 84 = Pluches baccharoides.
squarrosa, *Baker, in Mart. Fl. Bras.* vi. III. 50.—Bras.
squarrosa, *H. B. & K. Nov. Gen. et Sp.* iv. 57.—Mexic.
Steetzii, *Anderss. Galap. Oarnes Veg.* 177.—Ins. Galap.
stenocephala, *Baker, in Mart. Fl. Bras.* vi. III. 39.—Bras.
stenoptera, Sch. Bip. in Linnaea, xxxiv. (1865–66) 533 = subalata.
Sternbergiana, *Steud. Nom.* ed. II. i. 179.—Peruv.
striata, Kuiz & Pav. Syst. 207 = Fevillei.
Stroebeliana, *Ces. in Atti Soc. Acc. Sc. Nap.* v. (1873) VII. 12.—Europ.
stylosa, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 120.—Bras.
subalata, *Wedd. Chlor. And.* i. 174.—Bolivia.
subcapitata, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 85.—Bras.
subdentata, *DC. Prod.* v. 408.—Bras.
subporosa, *DC. l. c.* 413.—Bras.
subpenninervis, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 533, nomen.—Bolivia.
subscandens, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 76 = conyzoides.
subspathulata, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 87 = rufescens.
subulata, *D. Don, in Hook. Journ. Bot.* iii. (1841) 39.—Patagon.
sulcata, *DC. Prod.* v. 419.—Mexic.
synophala, Sch. Bip. in Bonplandia, iv. (1856) 54; et ex Baker, in Mart. Fl. Bras. vi. III. 99 = platypoda.
tarchonanthoides, *DC. Prod.* v. 414.—Bras.
teindalensis, *H. B. & K. Nov. Gen. et Sp.* iv. 417.—N. Granat.
tenella, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 42.—Patagon.
tenuifolia, *Linn. Sp. Pl.* 860.—' Afr.'
tenuifolia, DC. Prod. v. 433 = rufescens.
teucrifolia, Hook. & Arn. ex Baker, in Mart. Fl. Bras. vi. III. 77 = anomala.
texana, *A. Gray, Pl. Fendl.* 75.—Texas.
thesioides, *H. B. & K. Nov. Gen. et Sp.* iv. 61.—Mexic.; Arizona.
Thouesii, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 226.—Mexic.
thymifolia, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 38.—Reg. Argent.
thymifolia, Miers, in Proc. Hort. Soc. (1868) 349 = Wrightii.
thyoides, *Pers. Syn.* ii. 425.—Peruv.
Tola, *Phil. Fl. Atac.* 30.—Chili.
tomentosa, Steud. Nom. ed. II. i. 179 = Moquinia polymorpha.
tomentosa, *Pers. Syn.* ii. 424.—Peruv.
trichoclada, *DC. Prod.* v. 400.—Mexic.
trichotoma, *Klatt, in Leopoldina*, xx. (1884) 91.—Mexic.
tricuneata, Pers. Syn. ii. 424 = cuneata.
tridentata, Candich. Fl. Mal. 15 = magellanica.
tridentata, Poepp. ex DC. Prod. v. 411 = concava.
tridentata, *Vahl, Symb. Bot.* iii. 98.—Bras.
trifurcata, Trevir. in Nov. Act. Nat. Cur. xiii. (1826) 203 = Conyza stricta.
trimera, DC. Prod. v. 425 = genistelloides.
trinervis, *Pers. Syn.* ii. 423.—Bras.
triplinervis, DC. Prod. v. 404 = oxyodonta.
triplinervis, *Baker, in Mart. Fl. Bras.* vi. III. 79.—Bras.
tripteris, Mart. Mat. Med. Bras. 43 = genistelloides.
tripteris, Poepp. ex Baker, in Mart. Fl. Bras. vi. III. 42 = sagittalis.
truncata, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 82.—Bras.
truncata, *Phil. in Linnaea*, xxxiii. (1864) 148.—Chili.
tucumanensis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 34.—Reg. Argent.
Tweediei, *Hook. & Arn. l. c.* 33 = subopposita.
ulicina, *Hook. & Arn. l. c.* 38.—Patagon.
ulmifolia, *Burm. f. Fl. Cap. Prod.* 26 = Conyza incisa.
umbelliformis, *DC. Prod.* v. 410.—Chili.
uncinella, *DC. l. c.* 415.—Bras.
unifora, *Pers. Syn.* ii. 425.—Peruv.
Urvilleana, *Brongn. Bot. Voy. Coq.* t. 61.—Reg. Magell.
ustulata, *Benth. Pl. Hartw.* 135.—N. Granat.
vaccinioides, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 121.—Bras.
vaccinioides, *H. B. & K. Nov. Gen. et Sp.* iv. 50.—Mexic.
Vahlii, DC. Prod. v. 411 = dioica.
valdiviana, *Phil. in Linnaea*, xxviii. (1856) 788.—Chili.
varians, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 84 = rufescens.
Vauthieri, *DC. Prod.* v. 409.—Bras.
velutina, DC. l. c. 415 = ochraces.
veneta, H. B. & K. Nov. Gen. et Sp. iv. 68 = Bigelovia veneta.
venosa, *Pers. Syn.* ii. 425.—Peruv.
vernulosa, DC. Prod. v. 421 = brachylaenoides.
venusta, *H. B. & K. Nov. Gen. et Sp.* iv. 66.—N. Granat.
verbenaefolia, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 36 = hirta.
vernicosa, *Hook. & Arn. in Hook. l. c.* 28.—Uruguay.
vernonioides, *DC. Prod.* v. 422.—Bras.
villosa, *Vahl, Symb. Bot.* iii. 98.—Arabia.
viminea, *DC. Prod.* v. 400.—Calif.
vincaefolia, *Baker, in Mart. Fl. Bras.* vi. III. 83.—Reg. Argent.
virgata, Boj. ex DC. Prod. v. 385 = Conyza virgata.
viscosa, Hook. & Arn. Bot. Beech. Voy. 352 = Douglasii.
viscosa, Lam. Encyc. i. 345 = Psiadia integerrima.
viscosa, Walt. Fl. Carol. 202 = Pluches bifrons.
viscosa, DC. Prod. v. 412 = vernonioides.
Volckmanni, *Phil. in Linnaea*, xxxiii. (1864–65) 147.—Chili.
vulubilis, *H. B. & K. Nov. Gen. et Sp.* iv. 417.—Peruv.
vulneraria, *Baker, in Mart. Fl. Bras.* vi. III. 75.—Bras.
Wallenbergiana, D. Dietr. Syn. Fl. iv. 1447 = brachylaenoides.
Weddellii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 63 = rufescens.
Weirii, *Baker, l. c.* 67.—Bras.
Wendlandii, *Steud. Nom.* ed. II. i. 179.—Afr. austr.
Wrightii, *A. Gray, Pl. Wright.* i. 101.—Am. bor. occ.
Wrightii, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Seemanni.
xalapensis, H. B. & K. Nov. Gen. et Sp. iv. 56 = confecta.
xerophila, Mart. in Flora, xxiv. (1841) ii. Beibl. 11 = sessiliflora.
xiphophylla, *Baker, in Mart. Fl. Bras.* vi. III. 68.—Bras.

BACCHAROIDES, Moench, Meth. 578 (1794) = Vernonia, Schreb. (Compos.).
anthelmintica, Moench, l. c. = V. anthelmintica.

BACKHOUSIA, Hook. & Harv. in Bot. Mag. t. 4186 (1845). *MYRTACEÆ*, Benth. & Hook. f. i. 711.
angustifolia, *F. Muell. Fragm.* i. 79.—Austral.
BACKHOUSIA:—
citriodora, *F. Muell. Fragm.* i. 78.—Austral.
myrtifolia, *Hook. & Harv. in Bot. Mag.* t. 4183.—Austral.
riparia, Hook. & Harv. l. c. sub t. 4183 = myrtifolia.
sciadophora, *F. Muell. Fragm.* ii. 26, 171.—Austral.

BACONIA, DC. in Ann. Mus. Par. ix. (1807) 219 = **Pavetta**, Linn. (Rubiac.).
corymbosa, DC. l. c. = P. Baconia.
montana, Hook. f. in Journ. Linn. Soc. vii. (1864) 196 = P. Hookeriana.

BACOPA, Aubl. Pl. Gui. i. 128. t. 49 (1775). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 952.
aquatica, *Aubl. l. c.* 129. t. 49.—Guiana.
grandiflora, Mart. ex Benth. in Ann. Nat. Hist. ii. (1839) 448 = aquatica.

BACTRIS, Jacq. Stirp. Am. 271. t. 256 (1763). *PALMAE*, Benth. & Hook. f. iii. 941.
AUGUSTINEA, Karst. in Linnaea, xxvii. (1856) 895.
GUILIELMA, Mart. Palm. Fam. 21 (1824); Hist. Nat. Palm. ii. 31. t. 66, 67 (1824).
PYRENOGLYPHIS, Karst. Fl. Columb. ii. 141. t. 174 (1869).
acanthocarpa, *Mart. Hist. Nat. Palm.* ii. 92. t. 70, 71. f. 2.—Bras.
acanthocarpoides, Barb.-Rodr. Enum. Palm. Nov. 21 = acanthocarpa.
acanthocnemis, Mart. in Palmet. Orb. 67 = simplicifrons.
acanthophylla, *Mart. l. c.* 70.—Ins. Porto Rico.
acanthospatha, *Trail, ex Drude, in Mart. Fl. Bras.* iii. II. 354.—Bras.
actinocarpa, *Drude & Trail, ex Drude, in Mart. l. c.* 344.—Bras.
aculeifera, *Drude, l. c.* 352 in obs.—Bras.
acuminata, *Liebm. ex Mart. Hist. Nat. Palm.* iii. 221. Mexic.
arenaria, Barb.-Rodr. Enum. Palm. Nov. 29 = simplicifrons.
aristata, *Mart. Hist. Nat. Palm.* iii. 97. t. 78 a. f. 5.—Bras.
armata, Barb.-Rodr. Enum. Palm. Nov. 27 = chaetospatha.
arundinacea, *Trail, in Journ. Bot.* xv. (1877) 10.—Reg. Amazon.
Aubletiana, *Trail, l. c.* xiv. (1876) 372.—Guiana.
baculifera, *Karw. ex Mart. Hist. Nat. Palm.* iii. 222. —Mexic.
balanoides, *H. Wendl. in Kerch. Palm.* 238.—Costa Rica.
balanophora, *Spruce, in Journ. Linn. Soc.* xi. (1871) 146, 153.—Am. trop.
bicuspidata, *Spruce, l. c.* 146, 152.—Am. trop.
bidentata, *Spruce, l. c.* 146, 151.—Am. trop.
bifida, *Mart. Hist. Nat. Palm.* ii. 105. t. 73 c. —Bras.
bybda, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 44 = Oerstediana.
brevifolia, Spruce, in Journ. Linn. Soc. xi. (1871) 144, 147 = simplicifrons.
Brongniartii, *Mart. in Palmet. Orb.* 59. t. 7. f. 2.—Bolivia.
campestris, *Poepp. in Mart. Hist. Nat. Palm.* iii. 146. —Bras.
capillacea, *Trail, in Journ. Bot.* xv. (1877) 5.—Bras.
caracasana, *Lodd. ex Linden, Illustr. Hortic.* xxviii. (1881) 16.—Venezuela.
Carwalliana, *Linden, Illustr. Hortic.* xxviii. (1881) 16. —Bras.
caribaea, *Karst. in Linnaea*, xxviii. (1856) 408.—Ind. occ.
carolensis, Spruce, in Journ. Linn. Soc. xi. (1871) 145, 149 = simplicifrons.
caryotaefolia, *Mart. Hist. Nat. Palm.* ii. 106. t. 74. f. 3, 4.—Bras.
caryotifolia, *Drude, in Mart. Fl. Bras.* iii. II. 337.—Bras.
Catel, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Venezuela.
chaetophylla, *Mart. in Palmet. Orb.* 71.—Bras.
chaetorhachis, Mart. l. c. 61 = major.
chaetospatha, *Mart. Hist. Nat. Palm.* iii. 147.—Bras.
chloracantha, *Poepp. in Mart. l. c.* 145.—Bras.
ciliata, *Mart. l. c.* 95. t. 71. f. 8.—Peruv.
concinna, *Mart. l. c.* 99. t. 79. f. 4, 5, 6.—Bras.
confusa, Linden & H. Wendl. in Linnaea, xxviii. (1856) 847 = acanthospatha.

BACTRIS:—
Constantiae, *Barb.-Rodr. Enum. Palm. Nov.* 37.—Bras.
Coronilla, *Karst. in Linnaea*, xxvii. (1856) 407.—Venezuela.
Corusilla, Hort. ex H. Wendl. Cycl. Pand. Cycad. 5 = prec. ?
Cremgeriana, *Griseb. & Wendl. ex Griseb. Fl. Brit. W. Ind.* 520.—Ind. occ.
cucullata, *H. Wendl. in Kerch. Palm.* 238, nomen.—Hab. ?
Cuesco, *Cruez. ex Griseb. Fl. Brit. W. Ind.* 520.—Ind. occ.
Cusco, *Engl. in Linnaea*, xxxiii. (1864–65) 865.—N. Granat.
Cururna, Trail, in Journ. Bot. xv. (1877) 48 = socialis.
cuspidata, *Mart. Hist. Nat. Palm.* ii. 101. t. 78 B.—Bras.
Cuvaro, *Karst. in Linnaea*, xxviii. (1856) 406.—Venezuela.
cyagrodes, Barb.-Rodr. Enum. Palm. Nov. 33 = syagroides.
elatior, *Wallace, Palm Trees Amaz.* 81. t. xxx.—Bras.
elegans, *Hort. ex H. Wendl. in Kerch. Palm.* 233.—Hab. ?
elegans, *Trail, in Journ Bot.* xv. (1877) 3.—Bras.
ericrinus, Barb.-Rodr. Enum. Palm. Nov. 26 = bifida.
erosa, *Mart. Hist. Nat. Palm.* ii. 281.—Ind. occ.
eumorpha, *Trail, in Journ. Bot.* xv. (1877) 9. t. 184.—Bras.
exaltata, Barb.-Rodr. Enum. Palm. Nov. 32 = major ?
fauciun, *Mart. in Palmet. Orb.* 60. t. 6. f. 2.—Bolivia.
fissifrons, *Mart. Hist. Nat. Palm.* ii. 103. t. 78. b. f. 3, 4.—Bras.
flavispina, Hort. Belg. ex Heynh. Nom. ii. 57 = pallidispina.
floccosa, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 151 = cuspidata.
fusca, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 48.—Costa Rica.
Gasipaës, *H. B. & K. Nov. Gen. et Sp.* i. 302.—N. Granat.
Gastoniana, Trail, in Journ. Bot. xv. (1877) 48 = major.
geonomoides, *Drude, in Mart. Fl. Brit.* iii. 11. 255.—Bras.
glandulosa, *Oerst. L'Amér. Centr.* 15. t. 8. f. 2–15.—Bras.
glaucescens, *Drude, in Mart. Fl. Bras.* iii. 11. 245.—Bras.
Glazioviana, *Drude, l. c.* 248.—Bras.
globosa, Gaertn. Fruct. i. 22 = Acrocomia sclerocarpa.
gracilis, Barb.-Rodr. Enum. Palm. Nov. 37 = simplicifrons.
graminicaespa, *Barb.-Rodr. l. c.* 37.—Bras.
granatensis, *H. Wendl. ex Linden, Illustr. Hortic.* xxviii. (1881) 16.—N. Granat.
guineensis, Steud. Nom. ed. I. 96, 99 = minor.
hirta, *Mart. Hist. Nat. Palm.* ii. 105. t. 78. C. f. 3.—Bras.
horrida, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 41.—Nicaragua.
hylophila, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 152 = pectinata.
incommoda, *Trail, in Journ. Bot.* xv. (1877) 42.—Bras.
inermis, *Trail, ex Barb.-Rodr. Enum. Palm. Nov.* 30.—Bras.
infesta, *Mart. in Palmet. Orb.* 54. t. 7. f. 1.—Bolivia.
integrifolia, Wallace, Palm Trees Amaz. 91. t. xxxv = fissifrons.
interruptepinnata, *Barb.-Rodr. Enum. Palm. Nov.* 37.—Bras.
inundata, *Mart. in Palmet. Orb.* 58. t. 7. f. 3.—Bras.
jauaiusis, *Trail, in Journ. Bot.* xv. (1877) 40.—Bras.
leucanthia, *H. Wendl. in Linnaea*, xxviii. (1856) 345.—N. Granat.
Liboniana, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Bras.
linearifolia, Barb.-Rodr. Enum. Palm. Nov. 31 = pectinata.
littoralis, *Barb.-Rodr. l. c.* 36.—Bras.
longifrons, *Mart. Hist. Nat. Palm.* ii. 106.—Reg. Amazon.
longipes, *Poepp. in Mart. l. c.* 148. t. 78.—Bras.
Macanilla, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Venezuela.

BACTRIS:—
macracantha, *Mart. Hist. Nat. Palm.* ii. 95. t. 78.—Bras.
macracantha, *Mart. l. c.*—Bras.
macrocarpa, *Wallace, Palm Trees Amaz.* 85. t. xxxii.—Bras.
major, *Spruce, Stirp. Am.* 280. t. 171. f. 2.—N. Granat.
Maraja, *Mart. Hist. Nat. Palm.* ii. 93. t. 71. f. 1.—Bras.
Maraja-açu, *Barb.-Rodr. Enum. Palm. Nov.* 36.—Bras.
Maraja-y, Barb.-Rodr. l. c. ed. I. 29 = cuspidata.
martinezanefolia, Hort. ex H. Wendl. in Kerch. Palm. 234 = Martinezia corallina.
mexicana, *Mart. in Palmet. Orb.* 65.—Mexic.
microcarpa, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 153 = hylophila.
microspatha, Barb.-Rodr. Enum. Palm. Nov. 26 = cuspidata.
minas, *Miq. in Verh. Nat. Wet. Haarl.* vii. (1851) 207.—Guianas.
minima, Gaertn. Fruct. ii. 269. t. 189. f. 5 = Martinezia corallina.
minor, Gaertn. l. c. t. t. 9. f. 1 = Acrocomia lasiospatha ?
minor, *Spruce, Stirp. Am.* 279. t. 171. f. 1.—Bras.; N. Granat.
mitis, Mart. Hist. Nat. Palm. ii. 109 = cuspidata.
monticola, *Barb.-Rodr. Enum. Palm. Nov.* 34.—Bras.
negrensis, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 147 = simplicifrons.
nemorosa, *Barb.-Rodr. Enum. Palm. Nov.* 32.—Bras.
Neromanni, *Voigt, in Syll. Ratisb.* ii. (1828) 51.—Hab. ?
obovata, *H. Wendl. in Kerch. Palm.* 234.—Hab. ?
Oerstediana, *Trail, in Journ. Bot.* xv. (1877) 43.—Costa Rica.
oligocarpa, *Barb.-Rodr. Enum. Palm. Nov.* 28.—Bras.
ovata, *H. Wendl. in Kerch. Palm.* 234.—Nicaragua.
ovata, Stokes, Bot. Mat. Med. iv. 394 = major.
pallidispina, *Mart. in Palmet. Orb.* 62.—Guiana.
palustris, Barb.-Rodr. Enum. Palm. Nov. 36 = bidentula.
paraënsis, *Spitzg. ex H. Wendl. in Kerch. Palm.* 234.—Surinam.
pauciflora, *Barb.-Rodr. Enum. Palm. Nov.* 34.—Bras.
Pavoniana, *Mart. in Palmet. Orb.* 70.—Porto Rico.
pectinata, *Mart. Hist. Nat. Palm.* ii. 98. t. 60, 78. A. f. 1, 2.—Bras.
pilosa, *Mart. in Linnaea*, xxviii. (1856) 405.—N. Granat.
Piritu, *Wendl. ex H. Wendl. in Kerch. Palm.* 234.—N. Granat.
Piranga, *Trail, in Journ. Bot.* xv. (1877) 41.—Bras.
piscatorum, *Wedd. ex Drude, in Mart. Fl. Bras.* iii. 11. 254.—Bras.
Plumieriana, *Mart. in Palmet. Orb.* 64.—Ind. occ.
pnemorosa, *Poepp. ex Mart. l. c.* 56.—Peruv.
pulchra, *Trail, in Journ. Bot.* xv. (1877) 4. t. 184. f. 2.—Bras.
Puyamo, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Reg. Granat.
riparia, *Mart. Hist. Nat. Palm.* ii. 97. t. 71. f. 4.—Bras.
rivularis, *Barb.-Rodr. Enum. Palm. Nov.* 36.—Bras.
rotunda, Stokes, Bot. Mat. Med. iv. 394 = minor.
Seemanni, *Engl. in Linnaea* (1864–65), 867.—N. Granat.
sclophila, *Miq. in Verh. Nat. Wet. Haarl.* vii. (1851) 208.—Guianas.
setipinnata, Barb.-Rodr. Enum. Palm. Nov. 32 = hylophila.
setosa, *Mart. Hist. Nat. Palm.* ii. 94. t. 72. f. 1–3.—Bras.
setulosa, *Karst. in Linnaea*, xxviii. (1856) 408.—Venezuela.
simplicifrons, *Mart. Hist. Nat. Palm.* ii. 108. t. 78. C. f. 1–2.—Bras.
simplicifrons, Spruce, in Journ. Linn. Soc. xi. (1869) 146 = geonomoides.
socialis, Mart. in Palmet. Orb. 56. t. 14. f. 2 = major.
species, *Karst. in Linnaea*, xxviii. (1856) 409.—N. Granat.
sphaerocarpa, *Trail, in Journ. Bot.* xv. (1877) 8.—Amazon.
subglobosa, *H. Wendl. in Kerch. Palm.* 234.—Salvador.
syagroides, *Trail, in Journ. Bot.* xv. (1877) 76.—Bras.
sylvatica, *Barb.-Rodr. Enum. Palm. Nov.* 30.—Bras.

BACTRIS:—
tenera, *H. Wendl. in Kerch. Palm.* 234.—Hab. ?
tenuriua, *Mart. ex Drude, in Mart. Fl. Bras.* iii. 11. 253.—Bras.
tenuis, Wallace, Palm Trees Amaz. 87. t. xxxiii. = cuspidata.
tomentosa, *Mart. Hist. Nat. Palm.* ii. 100. t. 78. A. f. 3, 4.—Bras.
Trailiana, *Barb.-Rodr. Enum. Palm. Nov.* 27.—Bras.
trichospatha, *Trail, in Journ. Bot.* xv. (1877) 41, 42.—Bras.
turbinata, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 152 = pectinata.
turbinocarpa, *Barb.-Rodr. Enum. Palm. Nov.* 33.—Bras.
uaupensis, Spruce, in Journ. Linn. Soc. xi. (1871) 145, 150 = cuspidata.
umbraticola, *Barb.-Rodr. Enum. Palm. Nov.* 34.—Bras.
umbrosa, Barb.-Rodr. l. c. 29 = trichospatha.
utilis, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 418.—Costa Rica.
varinensis, *Hort. ex H. Wendl. in Kerch. Palm.* 234.—Hab. ?
villosa, *H. Wendl. ex Hemsl. Biol. Centr. Am. Bot.* iii. 418.—Costa Rica.
vulgaris, *Barb.-Rodr. Enum. Palm. Nov.* (1879) *App.* 43.—Bras.
xanthocarpa, Barb.-Rodr. l. c. 30 = simplicifrons ?

BACTYRILOBIUM, Willd. Enum. Hort. Berol. 439 (1809) = Cassia, Linn. (Legumin.).
biflorum, Willd. ex Steud. Nom. ed. II. i. 179 = C. bifloru.
ferrugineum, Schrad. in Goett. Gel. Anz. (1821) i. 713 = C. ferruginea.
Fistula, Willd. Enum. Hort. Berol. 440 = C. Fistula.
grande, Hornem. Hort. Hafn. i. 385 = C. grandis.
latifolium, Steud. Nom. ed. I. 41 = Anagyris latifolia.
molle, Schrad. in Goett. Gel. Anz. (1821) i. 718 = C. grandis.

BACULARIA, F. Muell. Fragm. vii. 108 (1870). *PALMAE*, Benth. & Hook. f. iii. 908.
LINOSPADIX, H. Wendl. & Drude, in Linnaea, xxxix. (1874) 177.
minor, *F. Muell. l. c.* xi. 58.—Austral.
monostachya, *F. Muell. l. c.* vii. 108; xi. 58.—Austral.

BADAMIA, Gaertn. Fruct. ii. 90. t. 97 (1791) = Terminalia, Linn. (Combret.).
Commersoni, Gaertn. l. c. = T. Catappa.

BADAROA, Bert. ex Steud. Nom. ed. II. i. 179 (1840) = seq.

BADEROA, Bert. ex Hook. Bot. Misc. iii. (1833) 324 = Bleyros, Linn. (Cucurb.).
armaria, Bert. ex Steud. Nom. ed. II. i. 179 = seq.
bryoniaefolia, Bert. ex Hook. l. c. = S. bryoniaefolius.

BADIERA, DC. Prod. i. 534 (1824). *POLYGALEAE*, Benth. & Hook. f. i. 127.
acuminata, *DC. l. c.* 535.—Mexic.
Berteriana, *Spreng. Syst.* iii. 178.—Ins. S. Doming.
divaricata, *DC. Prod.* i. 535.—Bras.
diversifolia, *DC. l. c.* 534.—Jamaica.
domingensis, *DC. l. c.*—Ins. S. Doming.
Pessaea, *DC. l. c.* 535.—Am. trop.

BADIERA, Hassk. Cat. Pl. Hort. Bogor. Alt. 227 (1844) = Polygala, Linn.
pulchra, Hassk. l. c. = P. venenosa.
venenosa, Hassk. l. c. = P. venenosa.

BADULA, Juss. Gen. 490 (1789) = Ardisia, Sw. (Myrsin.).
angustifolia, A. DC. in Trans. Linn. Soc. xvii. (1837) 114 = Embelia angustifolia.
arborea, Thou. ex A. DC. in DC. Prod. viii. 84 = Embelia arborea.
Barthesia, A. DC. in Trans. Linn. Soc. xvii. (1837) 113 = A. Barthesia.
Barthesia, Boj. Hort. Maurit. 196 = A. insularis.
borbonica, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 90.—Ins. Borbon.
caribaea, A. DC. l. c. 89 = A. caribaea.

BADULA:—
crassa, A. DC. in Trans. Linn. Soc. xvii. (1837) 114 = Ardisia insularis.
cylianthoides, A. DC. in Prod. viii. 107 = Cylianthus philippinensis.
discolor, Thou. ex A. DC. l. c. 89 = Oncostemum divaricatum.
ensolia, Thou. ex A. DC. l. c. 113 = (*Monoporus paludosus*).—Madag.
insularis, A. DC. in Trans. Linn. Soc. xvii. (1837) 113 = A. insularis.
iteoides, Benth. Pl. Hartw. 217.—N. Granat.
Lamarckiana, A. DC. in DC. Prod. viii. 110 = A. laurifolia.
lanceolata, Boj. ex A. DC. l. c. 90 = Oncostemum Bojeranum.
laterifiora, Griseb. in Goett. Abh. vii. (1857) 237 = A. laurifolia.
laurifolia, Boj. ex A. DC. in DC. Prod. viii. 108.—Madag.
Lhotskyana, A. DC. l. c. 111 = A. Lhotskyana.
Mamecillo, Schlecht. in Linnaea, xxvi. (1853–55) 671.—N. Granat.
micrantha, Boj. Hort. Maurit. 196 = A. insularis.
micrantha, A. DC. in Trans. Linn. Soc. xvii. (1837) 112.—Ins. Maurit.
multiflora, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 89 = A. insularis.
myrtifolia, Thou. ex A. DC. in DC. Prod. viii. 90 = Oncostemum myrtifolium.
nemorosa, Thou. ex A. DC. l. c. = Oncostemum nemorosum.
nigricans, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 89.—Bras.
ovalifolia, A. DC. in Trans. Linn. Soc. xvii. (1837) 114.—Ins. Maurit.
paludosa, Boj. ex A. DC. in DC. Prod. viii. 113 (= *Monoporus paludosus*).—Madag.
pauciflora, Boj. ex A. DC. l. c. 90 = Oncostemum pauciflorum.
peruviana, A. DC. l. c. 119 = A. nigricans.
philippinensis, A. DC. l. c. 108 = A. subcuneata.
reticulata, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 89 = A. insularis.
scandens, Boj. Hort. Maurit. 196 = Embelia micrantha.
Schomburgkiana, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 89.—Guiana.
Siebert, A. DC. in Trans. Linn. Soc. xvii. (1837) 113. t. 5 = A. guadalupensis.

BADUSA, A. Gray, in Proc. Am. Acad. iv. 308 (1860). *RUBIACEAE*, Benth. & Hook. f. ii. 42.
corymbiflora, *A. Gray, l. c.*—Ins. Pacif.

BADUSA Barthesia, Steud. Nom. ed. II. i. 98, in syn. (1840), sphalm. = BADULA, Juss. = Ardisia Barthesia.

BAEA, Comm. ex Juss. Gen. 121 (1789) = **Boea**, Comm. (Gesner.).

BAECKEA, Burm. f. Prod. Fl. Cap. 12 (1768) = **Brunia**, Linn.
africana, Burm. f. l. c. = Br. pinifolia.
cordata, Burm. f. l. c. = Br. racemosa.
lancifolia, Eckl. & Zeyh. Enum. 140 = Br. racemosa.
laurifolia, Sond. in Harv. & Sond. Fl. Cap. ii. 515, in syn. = Br. racemosa.
racemosa, Eckl. & Zeyh. Enum. 140 = Br. racemosa.
thymifiora, Harv. & Sond. Fl. Cap. ii. 604 = Br. pinifolia.
thymophora, Eckl. & Zeyh. Enum. 140 = Br. pinifolia.

BAECKEA, Linn. Sp. Pl. 856 (1753). *MYRTACEAE*, Benth. & Hook. f. i. 701.
ALLOUTIA, Raf. Sylva Tellur. 104 (1838).
BARINGTONIA, Lindl. in Bot. Reg. (1842) t. 10.
CAMPHOROMYRTUS, Schau. in Linnaea, xvii. (1843) 240.
CYATHOSTEMON, Turcz. in Bull. Phys.-Math. Acad. Petersb. x. (1852) 381.
EXICOMYRTUS, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 1. 254.
EUSYNOSTYLUS, Schau. in Linnaea, xvii. (1843) 239.
HARMOGIA, Schau. in Linnaea, xvii. (1843) 238.
ISERICARIA, Sm. in Trans. Linn. Soc. iii. (1797) 267.
JUNGIA, Gaertn. Fruct. i. 175. t. 35 (1788).
MOLLIA, J. F. Gmel. Syst. 490 (1791).
MURRINEA, Raf. Sylva Tellur. 104 (1838).
NEUHOFIA, Stokes, Bot. Mat. Med. i. 439 (1812).
OXYMYRRHINE, Schau. in Linnaea, xvii. (1843) 240.
RINZIA, Schau. l. c. xvii. 239 (1843).
SCHIDIOMYRTUS, Schau. l. c. 237 (1843).
TETRAPORA, Schau. l. c. 238 (1843).
TETRAPORIA, Miq. in Nederl. Kruidk. Arch iv. (1859) 160.
TJONGINA, Adans. Fam. ii. 234 (1763).
TRIPLARINA, Raf. Sylva Tellur. 104 (1838).
affinis, Endl. in Enum. Pl. Hueg. 51 = Astartea fascicularis.
affinis, Hook. Ic. Pl. t. 284 c = diffusa.
alpina, Lindl. in Mitch. Three Exped. ii. 178 = diffusa.
arbuscula, *R. Br. ex Benth. Fl. Austral.* iii. 79.—Austral.
astartioides, *Benth. l. c.* 80.—Austral.
Behrii, *F. Muell. Fragm.* iv. 68.—Austral.
Blaketii, *F. Muell. l. c.* viii. 181.—Austral.
brevifolia, *DC. Prod.* iii. 230.—Austral.
calycina, Lindl. in Mitch. Three Exped. ii. 190 = Thryptomene Mitchelliana.
camphorata, *R. Br. in Bot. Mag.* t. 2594.—Austral.
Camphorosmae, *Endl. in Enum. Pl. Hueg.* 51.—Austral.
Ascot.
capitata, *F. Muell. Census*, 54.—Austral.
carnulosa, Sieber, in Spreng. Syst. iv. Cur. Post. 149 = borealis.
chinensis, Gaertn. Fruct. i. 157 = frutescens.
ciliata, *F. Muell. Census*, 54.—Austral.
cochinchinensis, *Blume, Mus. Bot. Lugd. Bat.* i. 69.—Cochinch.
corymbulosa, *Benth. Fl. Austral.* iii. 87.—Austral.
corynophylla, *F. Muell. Fragm.* iv. 72.—Austral.
crassifolia, *Lindl. in Mitch. Three Exped.* ii. 115.—Austral.
crenatifolia, *F. Muell. Fragm.* iv. 70.—Austral.
crenulata, *DC. Prod.* iii. 230.—Austral.
crispiflora, *F. Muell. Fragm.* iv. 72.—Austral.
cryptandroides, *F. Muell. l. c.* x. 29.—Austral.
Cunninghamia, Schau. in Walp. Rep. ii. 920 = frutescens.
Cunninghamii, *Benth. Fl. Austral.* ii. 82.—Austral.
densifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 260.—Austral.
diffusa *Sieber, ex DC. Prod.* iii. 230.—Austral.
dimorphandra, *F. Muell. ex Benth. Fl. Austral.* iii. 74.—Austral.
dioemifolia, *Rudge, in Trans. Linn. Soc.* viii. (1807) 298. t. 18.—Austral.
discomoides, Sieber, ex DC. Prod. iii. 230 = crenulata.
Drummondii, *Benth. Fl. Austral.* iii. 75.—Austral.
ericaea, *F. Muell. Fragm.* i. 85.—Austral.
ericoides, *Bromfn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 184.—N. Caled.
ericoides, *Schlecht. in Bot. Zeit.* iv. (1846) 794.—Austral.
fasciculata, Sieber, in Spreng. Syst. iv. Cur. Post. 149 = densifolia.
floribunda, *Benth. Fl. Austral.* iii. 87.—Austral.
frutescens, *Linn. Sp. Pl.* 858.—Malaya; China.
fumana, *F. Muell. Fragm.* iv. 68.—Austral.
gracilis, *F. Cunn. in Field, New South Wales*, 349.—Austral.
grandiflora, *Benth. Fl. Austral.* iii. 89.—Austral.
Gunniana, *Schau. ex Walp. Rep.* ii. 920.—Austral.
involucrata, Endl. in Enum. Pl. Hueg. 170 = Scholtzia obovata.
laxiflora, *F. Muell. Census*, 54.—Austral.
leptocaulis, *Hook. f. in Hook. Ic. Pl.* t. 926.—Tasmania.
leptantha, *F. Muell. Census*, 54.—Austral.
leptocalyx, F. Muell. Fragm. i. 80 = Micromyrtus leptocalyx.
linifolia, *Rudge, in Trans. Linn. Soc.* viii. (1807) 297. Austral.
micrantha, DC. Prod. iii. 230 = Thryptomene baeckeacea.
micrantha, Hook. f. in Hook. Ic. Pl. t. 309 = Gunniana.
microphylla, A. Cunn. ex Schau. in Linnaea, xvii. (1843) 238 = Harmogia Cunninghamii.
microphylla, Sieber, ex Spreng. Syst. iv. Cur. Post. 149 = Micromyrtus microphylla.
multiclades, *Seem. in Journ. Bot.* ii. (1864) 74.—N. Caled.
obovata, DC. Prod. iii. 230 = Scholtzia obovata.
obtusifolia, *Bronga. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 185.—N. Caled.
ochropetala, *F. Muell. Fragm.* x. 29.—Austral.
oligandra, *F. Muell. Census*, 54.—Austral
oligomera, *Radlk. in Ber. Deutsch. Bot. Ges.* ii. (1884) 264.—Austral.
ovalifolia, *F. Muell. Fragm.* iv. 72.—Austral.
oxycoccoides, *Benth. Fl. Austral.* iii. 75.—Austral.
pachyphylla, *Benth. l. c.* 85.—Austral.
parviflora, *F. Muell. Census*, 54.—Austral.
parvula, DC. Prod. iii. 229 = virgata.
pentagonantha, *F. Muell. Fragm.* iv. 73.—Austral.
pentandra, *F. Muell. l. c.* ii. 81; iv. 72.—Austral.
phylicoides, A. Cunn. ex Schau. in Walp. Rep. ii. 921 = Kunzea peduncularis.
pinifolia, *DC. Prod.* iii. 229.—N. Caled.
platystemona, *Benth. Fl. Austral.* iii. 74.—Austral.
plicata, F. Muell. Fragm. i. 30 = Micromyrtus microphylla.
polyandra, *F. Muell. l. c.* iv. 72.—Austral.
polystemones, *F. Muell. l. c.* ii. 124.—Austral.
prostrata, Hook. f. Ic. Pl. t. 284 b = diffusa.
pulchella, *DC. Prod.* iii. 230.—Austral.
pygmaea, *R. Br. ex Benth. Fl. Austral.* iii. 86.—Austral.
ramosissima, *A. Cunn. in Field, New South Wales*, 348.—Austral.
robusta, *F. Muell. Fragm.* iv. 72.—Austral.
saxicola, *A. Cunn. ex Hook. Bot. Mag.* t. 3160.—Austral.
schollaraefolia, *Lehm. Pl. Preiss.* ii. 369.—Austral.
serpillifolia, *F. Muell. Fragm.* x. 30.—Austral.
sinensis, Gaertn. Fruct. i. 81 = frutescens.
spicata, A. Cunn. ex Schau. in Linnaea, xvii. (1843) 241 = Hypocalymma strictum.
spathulata, *F. Muell. Census*, 54.—Austral.
spinosa, *Sieber, ex Spreng. Syst. iv. Cur. Post.* 149.—Austral.
stenophylla, *F. Muell. Fragm.* i. 13.—Austral.
subcuneata, *F. Muell. l. c.* iv. 73.—Austral.
sumatrana, *Blume, Mus. Bot. Lugd. Bat.* 69.—Sumatra.
tarctiflora, *F. Muell. Census*, 54.—Austral.
tetragona, *F. Muell. ex Benth. Fl. Austral.* iii. 77.—Austral.
thymifolia, Hook. f. Ic. Pl. t. 284 a = diffusa.
trichophylla, Sieber, ex Spreng. Syst. iv. Cur. Post. 149 = linifolia.
uberiflora, *F. Muell. Census*, 53.—Austral.
umbellata, F. Muell. Fragm. iv. 69 = virgata.
umbelliflora, *F. Muell. Census*, 54.—Austral.
unicinella, *Benth. Fl. Austral.* iii. 84.—Austral.
utilis, F. Muell. ex Miq. in Neder. Kruidk. Arch. iv. (1859) 150 = Gunniana.
virgata, *Andr. Bot. Rep.* 598.—N. Caled.

BAEJICA, C. B. Clarke, in DC. Monog. Phan. v. 134 (1883) = **Boetica**, C. B. Clarke (Gesner.).

BAEOBOTRYS, Forst. Char. Gen. 21. t. 11 (1776) = **Maesa**, Forsk. (Myrsin.).
acuminata, Spreng. Syst. in Ind. v. 89 = M. ramentacea.
acuminata, Wall. Cat. n. 2231 = Gymnosporia acuminata.
argentea, Wall. l. c. n. 2216 = Maesa argentea.
Begotia, Buch.-Ham. ex A. DC. in DC. Prod. viii. 81 = M. macrophylla.
dubia, Wall. Cat. n. 2217 = M. dubia.
fragrans, Wall. l. c. n. 2202 = M. ramentacea.
fuloiserrie, Kollm. ex A. DC. in DC. Prod. viii. 82 = M. mollis.
glabra, Roxb. Hort. Beng. 16; Fl. Ind. i. 560 = M. ramentacea.
indica, Roxb. ll. cc. 16, 557 = M. indica.
indica, Wall. Cat. n. 2218. A & B = M. Chisia.
lanceolata, Blume, Bijdr. 863 = M. Blumii.
lanceolata, Spreng. Syst. iv. Cur. Post. 83 = M. ramentacea.
lanceolata, Vahl, Symb. Bot. i. 19 = M. lanceolata.
latifolia, Blume, Bijdr. 865 = M. indica.
lucida, Wall. Cat. n. 2223.—Ind. or.
lutea, Roxb. ex Steud. Nom. ed. II. i. 180 = M. lutea.
macrophylla, Wall. Cat. n. 2225 = M. macrophylla.
missionis, Wall. l. c. n. 6523 = M. ramentacea!

BAEOBOTRYS:—
mollis, Blume, Bijdr. 885 = Masea mollis.
mollissima, Blume, l. c. 886 = M. mollissima.
Morsha, Buch.-Ham. ex A. DC. in DC. Prod. viii. 80 = M. indica.
muscosa, Reinw. ex Blume, Bijdr. 886 = M. membranacea.
nemoralis, Forst. Char. Gen. 22 = M. nemoralis.
nemoralis, Roxb. Hort. Beng. 85; Fl. Ind. i. 559 = M. indica.
ovata, Wall. Cat. n. 3924 = M. ramentacea.
paniculata, Wall. l. c. n. 3920 = M. paniculata.
picta, Hochst. ex Walp. Ann. iii. 964 = M. lanceolata.
pubescens, Lodd. ex Lond. Hort. Brit. 71.—M. pubescens.
ramentacea, Roxb. Hort. Beng. 16; Fl. Ind. i. 558 = M. ramentacea.
rufescens, E. Mey. ex A. DC. in DC. Prod. viii. 81 = M. lanceolata.
tetrandra, Roxb. Hort. Beng. 85; Fl. Ind. i. 560 = M. tetrandra.
tomentosa, Buch.-Ham. ex Spreng. Syst. iv. Cur. Post. 88 = M. macrophylla.
virgata, Blume, Bijdr. 866 = M. indica.

BAEOMETRA, Salisb. in Trans. Hort. Soc. i. (1812) 330. *LILIACEAE*, Benth. & Hook. f. iii. 828.
JANIA, Schult. Syst. vii. 1. p. xcviii. 1658 (1830).
KOLBEA, Schlecht. in Linnaea, i. (1826) 80.
columellaris, *Salisb. l. c.*—Afr. austr.

BAEOTERPE, Salisb. Gen. Pl. Fragm. 17 (1866) = **Hyacinthus**, Linn. (Liliac.).
brevifolia, Salisb. l. c. 18 = H. brevifolius.
corymbosa, Salisb. l. c. = H. corymbosus.

BAEOTHRYON, Ehrh. Beitr. iv. 147 (1789) = **Scirpus**, Linn. (Cyperac.).
caespitosum, A. Dietr. Sp. Pl. ii. 89 = S. caespitosus.
confervoides, A. Dietr. l. c. 94 = S. confervoides.
cruinum, A. Dietr. l. c. = Ficinia filiformis.
festucoides, A. Dietr. l. c. = Heleocharis festucoides.
fistulosum, A. Dietr. l. c. = Heleocharis fistulosa.
flaweens, A. Dietr. l. c. 91 = Eleocharis ocreata.
Halleri, T. Nees, Gen. Germ. lic. ii. 17 = S. pauciflorus.
Hystrix, A. Dietr. Sp. Pl. ii. 92 = Hypodiscus aristatus.
nanum, A. Dietr. l. c. 91 = S. nanus.
pauciflorum, A. Dietr. l. c. 90 = S. pauciflorus.
pusillum, A. Dietr. l. c. 92 = S. pusillus.
pygmaeum, A. Dietr. l. c. = Eleocharis Chaetaria.
retroflexum, A. Dietr. l. c. 93 = Eleocharis Chaetaria.
tenuiculum, A. Dietr. l. c. = Eleocharis Rothiana.
trichodes, A. Dietr. l. c. = Eleocharis trichodes.
variegatum, A. Dietr. l. c. 93 = Eleocharis variegata.

BAERIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 29 (1835). *COMPOSITAE*, Benth. & Hook. f. ii. 399.
DICHAETA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 383.
PTILOMERIS, Nutt. in Am. Phil. Trans. N. S. vii. (1841) 381.
affinis, *A. Gray, in Proc. Am. Acad. xix.* (1883) 23.—Calif.
anthemoides, *A. Gray, l. c.* 23.—Calif.
cernua, *Greene, in Bull. Torrey Club*, x. (1883) 86.—Calif.
chrysostoma, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1835) 29.—Calif.
Cleveland, *A. Gray, in Proc. Am. Acad.* xix. (1883) 22.—Calif.
coronaria, *A. Gray, l. c.* 22.—Calif.
curta, *A. Gray, l. c.* 21.—Calif.
debilis, *Greene, ex A. Gray, Syn. Fl. N. Am.* i. ii. 326.—Calif.
Fremontii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 358; ix. (1874) 196.—Calif.
gracilis, *A. Gray, l. c.* ix. (1874) 196.—Calif.; Arizon.
hispidula, *A. Gray, Syn. Fl. N. Am.* i. ii. 325.—Calif.
macrantha, *A. Gray, in Proc. Am. Acad.* xix. (1883) 21.—Calif.
maritima, *A. Gray, l. c.* ix. (1874) 196; *Bot. Calif.* i. 376.—Calif.
mentzeliae, *A. Gray, in Proc. Am. Acad.* xix. (1883) 23.—Calif.
Palmeri, *A. Gray, in Bot. Calif.* i. 376.—Calif.

BAERIA:—
platycarpha, *A. Gray, in Proc. Am. Acad.* ix. (1874) 196.—Calif.
tenella, *A. Gray, l. c.* xix. (1883) 23.—Calif.
tenerrima, *A. Gray, l. c.* ix. (1874) 196.—Calif.
uliginosa, *A. Gray, l. c.*—Calif.

BAEUMERTA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 419 (1800) = **Nasturtium**, R. Br. (Crucif.).
Nasturtium, Gaertn. Mey. & Scherb. l. c. 467 = N. officinale.

BAGALATTA, Roxb. ex Reichb. Consp. 86 (1828) = **Tillaeora**, Colebr. (Menisperm.) (T. racemosa).

BAGASSA, Aubl. Pl. Gui. Suppl. 15. t. 376 (1775). *URTICACEAE*, Benth. & Hook. f. iii. 362.
LAUREA, Gaudich. Voy. Bonite, t. 88 (1836).
LOURIA, Kunth, in Linnaea, xiii. (1839) 502.
guianensis, *Aubl. l. c.*—Guiana.
Sagotiana, *Bur. ex Benth. & Hook. f. Gen.* iii. 362.—Guiana.

BAGNISIA, Becc. Malesia, i. 249. t. 19 (1878). *BURMANNIACEAE*, Benth. & Hook. f. iii. 482.
SARCOSIPHON, Blume, Mus. Bot. Lugd. Bat. i. 65. t. 18 (1849).
crocea, *Becc. l. c.*—Malaya.

BAHEL, Adans. Fam. ii. 210 (1763) = **Hygrophila**, R. Br. (Acanth.) (H. spinosa).

BAHIA, Lag. Gen. et Sp. Nov. 30 (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 402.
STYLESIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 377.
TRICHOPHYLLUM, Nutt. Gen. Am. ii. 166 (1818).
VILELTIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 402 (1873).
absinthifolia, *Benth. Pl. Hartw.* 18.—Texas; Mexic.
achilleoides, DC. Prod. v. 657 = Eriophyllum caespitosum.
ambigua, A. Gray, in Bot. Calif. i. 382 = Eriophyllum ambiguum.
ambrosioides, *Lag. Gen. et Sp. Nov.* 28, 30.—Chili.
angustifolia, DC. Prod. v. 656 = Calea angustifolia.
anthemoides, *A. Gray, in Proc. Am. Acad.* xv. (1880) 40.—Mexic.
arachnoidea, Fisch. & Avé-Lall. in Ind. Sem. Hort. Petrop. ix. 63 = Eriophyllum caespitosum.
artemisiaefolia, Less. in Linnaea, iv. (1829) 160 = Eriophyllum staechadifolium.
Bigelovii, *A. Gray, in Torr. Bot. Mex. Bound.* 96.—Texas.
biternata, *A. Gray, Pl. Wright.* ii. 95.—Texas.
chrysanthemoides, *A. Gray, in Proc. Am. Acad.* xix. (1883) 23.—Am. bor. occ.
confertiflora, DC. Prod. v. 657 = Eriophyllum confertiflorum.
cuneata, Kellogg, in Proc. Calif. Acad. v. (1873) 49 = Eriophyllum caespitosum.
dealbata, A. Gray, Pl. Fendl. 99 = absinthifolia.
Gilliesii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 23.—Reg. Argent.
gracilis, Hook. & Arn. Bot. Beech. Voy. 358 = Eriophyllum gracile.
integrifolia, DC. Prod. v. 656 = Eriophyllum caespitosum.
lanata, DC. l. c. 657 = Eriophyllum caespitosum.
latifolia, Benth. Bot. Voy. Sulph. 30 = Eriophyllum caespitosum.
latifolia, *Lindl. in Journ. Hort. Soc.* viii. (1853) 319.—Calif.
lemophylla, DC. Prod. v. 657 = Eriophyllum caespitosum.
leucophylla, Eaton, Bot. King, Exped. 17, pro parte = Eriophyllum Watsoni.
multiflora, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 373 = Eriophyllum caespitosum.
nec-mexicana, *A. Gray, in Proc. Am. Acad.* xix. (1883) 27.—N. Mexic.
nepetaefolia, A. Gray, l. c. v. (1861) 134 = sinuata.
nudicaulis, *A. Gray, l. c.* xix. (1883) 27.—Am. bor. occ.
obtusiflora, *A. Gray, l. c.*—Colorado.
oppositifolia, *DC. Prod.* v. 656.—Am. bor. occ.
Palmeri, *A. Gray, Syn. Fl. N. Am.* ii. 1. 402.—Calif.

BAHIA:—
parviflora, A. Gray, in Bot. Calif. i. 382 = Eriophyllum ambiguum.
pedata, *A. Gray, Pl. Wright.* i. 128.—Texas.
resinosa, DC. Prod. vi. 676 = Galinsoga resinosa.
rubella, A. Gray, in Torr. Bot. Mex. Bound. 96 = Eriophyllum Wallacei.
schkuhrioides, *A. Gray, in Proc. Am. Acad.* xix. (1883) 27.—Mexic.
sinuata, *Less. in Linnaea*, vi. (1831) 160.—Mexic.
staechadifolia, DC. Prod. v. 656 = Eriophyllum staechadifolium.
tenuifolia, *DC. l. c.* 657.—Calif.
trifida, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 374 = Eriophyllum confertiflorum.
trolliifolia, *DC. Prod.* v. 657.—Mexic.
Wallacei, A. Gray, in Pacif. Rail. Rep. iv. 105 = Eriospermum Wallacei.
Wallacei, A. Gray, in Proc. Nat. Hist. Bost. vii. (1859–61) 145 = Eriospermum ambiguum.
Woodhousii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 28.—N. Mexic.

BAHIOPSIS, Kellogg, in Proc. Calif. Acad. Nat. Soc. ii. (1863) 85 = **Viguiera**, H. B. & K. (Compos.).
lanata, Kellogg, l. c. = V. civea?

BAIKARA, Benth. in Benth. & Hook. f. Gen. i. 681 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
insigula, *Benth. in Trans. Linn. Soc.* xxv. (1866) 314. t. 41.—Afr. trop.
minor, *Oliver, Fl. Trop. Afr.* ii. 309.—Afr. trop.

BAILEYA, Harv. & Gray, ex Torr. in Emory, Notes Mil. Reconnois. 144 (1848). *COMPOSITAE*, Benth. & Hook. f. ii. 428.
multiradiata, *Harv. & Gray, ex Torr. l. c.*—Am. bor. occ.
pauciradiata, *Harv. & Gray, ex A. Gray, Pl. Fendl.* 105.—Calif.
pleniradiata, Harv. & Gray, ex A. Gray, l. c. = multiradiata.

BAILLIERIA, Aubl. Pl. Gui. ii. 804. t. 317 (1775) = Cilbedium, Linn. (Compos.).
aspera, Aubl. l. c. = C. surinamense.
Barbasco, H. B. & K. Nov. Gen. et Sp. iv. 259 = C. Barbasco.
caroliniana, Spreng. Syst. iii. 625 = Iva imbricata.
commelinoides, Less. Syn. Comp. 215 = Ichthyothere Cunabi.
cross, Poir. Encyc. Suppl. i. 563 = C. crossum.
nereifolia, H. B. & K. Nov. Gen. et Sp. iv. 259 = C. nereifolium.
pedunculosa, Poir. Encyc. Suppl. i. 563 = C. pedunculosum.
sylvestris, Aubl. Pl. Gui. ii. 807 = C. surinamense.
terebinthacea, Poir. Encyc. Suppl. i. 563 = C. terebinthaceum.

BAILLONIA, Boog. ex Baill. Adansonia, ii. (1861–62) 93. t. 7. *VERBENACEAE*, Benth. & Hook. f. ii. 1143.
DIOSTEA, Miers, in Trans. Linn. Soc. xxvii. (1870) 108. t. 28.
amabilis, *Boog. ex Baill. l. c.* 352.—Paraguay.
junces, *Benth. & Hook. f. Gen.* ii. 1144.—Chili.

BAIMO, Rafin. Fl. Tellur. iv. 27 (1836) = **Fritillaria**, Tourn. (Liliac.).
cirrhosa, Rafin. l. c. = F. japonica.

BAISSEA, A. DC. Prod. viii. 424 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 719.
CLEMORNIA, Wight, Ic. t. 1210, 1213 (1850).
acuminata, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 668.—Ceylan.
leonensis, *Benth. in Hook. Niger Fl.* 492.—Afr. trop.
malacocarpa, *Hook. f. Fl. Brit. Ind.* iii. 668.—Malacca.
multiflora, *A. DC. in Prod.* viii. 424.—Afr. trop.

BAITARIA, Ruiz & Pav. Prod. 62. t. 26 (1797) = **Calamdrinia**, H. B. & K. (Portulac.).
acaulis, *Ruiz & Pav. Fl. Per.* iv. t. 408; *Syst. Veg.* 111.—Peruv.

BAJAN, Adans. Fam. ii. 306 (1763) = **Amaranthus**, Linn.

BAKERIA, Seem. Journ. Bot. ii. (1864) 248 = **Flerandra**, A. Gray (Araliac.).
wilfordæ, Seem. l. c. = F. viticosa.

BALAKA, Becc. in Ann. Jard. Buitenz. ii. (1885) 91 in obs. *PALMAE.*
perbrevis, *Becc. l. c.*—Ins. Fiji.
Seemanni, *Becc. l. c.*—Ins. Fiji.

BALANAULAX, Rafin. Alsog. Am. 28 (1838) = Quercus, Tourn. (Cupul.).
malacca, Rafin. l. c. = Q. molucca.
Rumphi, Rafin. l. c. = Q. molucca ?

BALANGHAS, [Burm.] Rafin. Sylva Tellur. 72 (1838) = **Sterculia**, Linn.
rubiginosa, Rafin. l. c. = S. rubiginosa.
Telabo, Rafin. l. c. = S. Balanghas.

BALANGUE, Gaertn. Fruct. ii. 495. t. 180. f. 3 (1791). *JASMINEAE* ?
Gaertneri, *DC. Prod.* viii. 316.—Madag.

BALANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1 (Quid ?).

BALANITES, Delile, Fl. Egypte, 221. t. 28. f. 1. (1813) *SIMARUBEAE*, Benth. & Hook. f. i. 314.
AGIALID, Adans. Fam. ii. 508 (1763).
aegyptiaca, *Delile, l. c.*—Afr. bor. ; Arab. ; Palaestin.
aegyptica, Wall. Cat. n. 8855 = Roxburghii.
ferox, G. Don, Gen. Syst. i. 774 = aegyptiaca.
Roxburghii, *Planch. in Ann. Sc. Nat.* Sér. IV. iv. (1856) 258. t. 2.—Ind. or.

BALANOCARPUS, Bedd. Forest. Man. Bot. 236 [bis] (1873). *DIPTEROCARPEAE.*
ceros, *Bedd. l. c.* 237.—Ind. or.
utilis, *Bedd. l. c.*—Ind. or.

BALANOPHORA, Forst. Char. Gen. t. 50 (1776). *BALANOPHOREAE*, Benth. & Hook. f. iii. 285.
CYNOPSOLE, Endl. Gen. 74 (1838).
SARCOCORDYLIS, Wall. Cat. n. 7249 (1831 ?).
SCYNOPSOLE, Reichb. Handb. 164 (1837).
abbreviata, *Blume, Bijdr. Fl. Nederl. Ind.* i. 87.—Java.
affinis, Griff. in Proc. Linn. Soc. i. (1849) 220; Trans. Linn. Soc. xx. (1851) 94. t. 4 = dioica.
alsiacea, Jungh. in Nov. Act. Nat. Cur. xviii. Suppl. i. (1841) 205 = abbreviata.
alsiolata, Griff. in Proc. Linn. Soc. i. (1849) 220; Trans. Linn. Soc. xx. (1851) 94 = dioica.
Burmanniana, Griff. ii. cc. 219 et 93 = dioica.
capensis, Eckl. & Zeyh. ex Eichl. in DC. Prod. xvii. 125 = Mystropetalon Thomii.
dioica, *R. Br. in Wall. Cat.* n. 7245.—Ind. or. ; Burma.
dioica, Unger, in Ann. Wien. Mus. ii. (1838) t. 2. f. 1, 2 = elongata.
elongata, *Blume, Enum. Pl. Jav.* i. 87.—Java.
elongata, Schott & Endl. Melet. 18, partim = dioica.
fungosa, *Forst. Char. Gen.* 99. t. 50.—Austral. ; N. Hebrid.
gigantea, Wall. Cat. n. 7249 = globosa.
globosa, *Jungh. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 210.—Burma ; Java.
Harlandi, *Hook. f. in Trans. Linn. Soc.* xxii. (1859) 426.—Hongkong.
Hildebrandtii, *Reichb. f. in Journ. Bot.* xiv. (1876) 45. —Ins. Comoro.
indica, *Wall. Cat.* n. 7247.—Ind. or.
involucrata, *Hook. f. & Thoms. in Trans. Linn. Soc.* xxii. (1859) 20.—Reg. Himal.
Lowii, *Hook. f. l. c.* 426. t. 75 C.—Borneo.
maxima, Jungh. in Nov. Act. Nat. Cur. xviii. Suppl. i. (1841) 209. t. 1 = elongata.
pleta, Griff. in Proc. Linn. Soc. i. (1849) 220; Trans. Linn. Soc. xx. (1851) 94 = dioica.
picta, Miq. ex Eichl. in DC. Prod. xvii. 146 = indica.
polyandra, *Griff. in Trans. Linn. Soc.* xx. (1851) 94.—Reg. Himal.
rubicunda, *Becc. in Att. Soc. Ital. Sc. Nat. Milano*, xl. (1868) 197.—Borneo.
Thwaitesii, *Eichl. in DC. Prod.* xvii. 146, in note. —Ceylon.
typhina, Wall. Cat. n. 7246 = dioica.

BALANOPLIS, Rafin. Alsog. Amer. 29 (1838) = **Quercus**, Tourn. (Cupul.).
serrata, Rafin. l. c. 30 = Q. cuspidata.
tribuloides, Rafin. l. c. 29 = Castanopsis tribuloides.

BALANOPS, Baill. Adansonia, x. (1871) 117, 287. *BALANOPSEAE*, Benth. & Hook. f. iii. 341, 1223.
australiana, *F. Muell. Fragm.* x. 114.—Austral.
Balansae, *Baill. Adansonia*, x. (1872) 288.—N. Caled.
microcarpa, *Baill. l. c.* 340.—N. Caled.
oliviformis, *Baill. l. c.* 287.—N. Caled.
Pancheri, *Baill. l. c.* 118.—N. Caled.
Theophrasta, *Baill. l. c.* 288.—N. Caled.
Vieillardi, *Baill. l. c.* 118.—N. Caled.

BALANOPSIS, Rafin. Sylva Tellur. 134 (1838) = **Cocclea**, Arbl. (Laurin.).
acuta, Rafin. l. c. (= *Mespilodaphne cupularis*).
Cassia, Rafin l. c. = Litsaea zeylanica ?
cupulata, Rafin. l. c. (= *Mespilodaphne cupularis*).
oliptica, Rafin. l. c.—Hab. ?

BALANOPTERIS, Gaertn. Fruct. ii. 94. t. 98 (1791) = **Hartiera**, Ait. (Sterculiac.).
minor, Gaertn. l. c. t. 98. f. 2 = H. Fomes.
Tothila, Gaertn. l. c. p. 94. t. 99 = H. littoralis.

BALANOSTREBLUS, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 247. t. 19. *URTICACEAE*, Benth. & Hook. f. iii. 377.
ilicifolia, *Kurz, l. c.* 248.—Chittagong ; Burma ; Penang.

BALANSAEA, Boiss. & Rent. Pugill. Fl. Nov. 49 (1852) = **Chaerophyllum**, Linn. (Umbellif.).
Fontanesii, Boiss & Rent l. c. 50.—Hispan.
glaberrima, Lange, Pugill. iv. 238 = Daucus glaberrimus.

BALANTIUM, Desv. ex Ham. Prod. Fl. Ind Occ. 84 (1825) = **Parinarium**, Juss (Rutac.).
cordifolium, Desv. l. c.—Ind. occ. ?

BALARDIA, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 180. t. 111 (1829) = **Spergularia**, Presl (Caryophyll.).
platensis, Cambess l. c. = S. platensis.

BALAUSTION, Hook. Ic. Pl. t. 852 (1852). *MYRTACEAE*, Benth. & Hook. f. i. 702.
CHEYNIA, J. Drumm. in Hook. Kew Journ. vii. (1855) 24.
PUNICELLA, Turcz. in Bull. Acad. Pétersb. x. (1852) 332.
pulcherrimum, *Hook. Ic. Pl.* t. 852.—Austral.

BALBISIA, Cav. in Anal. Cienc. Nat. vii. (1804) 62. t. 46. *GERANIACEAE* Benth. & Hook. f. i. 276.
CRUCHSHANKSIA, Hook. & Arn. in Hook. Bot. Misc. ii. (1831) 211. t. 90.
LEDOCARPON, Desv. in Mém. Mus. Par. vi. (1818) 260.
Meyeniana, *Klotzsch, in Linnaea*, x. (1836) 492.—Pervv. peduncularis, *D. Don, in Edinb. N. Phil. Journ.* xi. (1831) 277.—Chili.
verticillata, *Cav. in Anal. Cienc. Nat.* vii. (1804) 62. t. 46.—Chili.

BALBISIA, DC. in Guill. Archiv. Bot. ii. (1838) 388 = **Rhetinodendron**, Meisn. (Compos.).
Berterii, DC. l. c. = R. Berterol.

BALBISIA, Wild. Sp. Pl. iii. 2214 (1803) = **Tridax**, Linn. (Compos.).
Caladonica, Spreng. Fl. Hal. Mant. 53 = Olearia viscosa.
canescens, Rich. ex Pers. Syn. ii. 470 = T. procumbens.
dimorpha, Cass. Opusc. iii. 91 ex DC. Prod. v. 679 = T. procumbens.
elongata, Willd. Sp. Pl. iii. 2214 = T. procumbens.
pedunculata, Hoffmgg. Verz. Pfl. (1824) 238 = T. procumbens.
suffruticosa, Dietr. ex DC. Prod. v. 267 = Olearia viscosa.

BALBOA, Liebm. in Kjoeb. Vidensk. Meddel. (1858) 106 = **Peghroeta**, Pers. (Legumin.).
diversifolia, Liebm. l. c. 107 = T. madrensis.

BALBOA, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 353. *GUTTIFERAE*, Benth. & Hook. f. i. 272.
membranacea, *Planch. & Triana, l. c.*—N. Granat.

BALDELLIA, Parl. Nuov. Gen. e Spec. 57 (1854) = **Alisma**, Linn.
ranunculoides, Parl. l. c. 58 = A. ranunculoides.

BALDINGERA, Dennst. Schluss. Hort Malab. 31 (1818) = **Fremna**, Linn. (Verben.).
glandulosa, Dennst. l. c. = F. longifolia.

BALDINGERA, Gaertn. Mey. & Scherb. Fl. Wett. i. 48 (1799) = **Phalaris**, Linn. (Gramin.).
arundinacea, Dum. Obs. Gram. Belg. 130 = P. arundinacea.
colorata, Gaertn. Mey. & Scherb. Fl. Wett. i. 96 = P arundinacea.

BALDINGERIA, Neck. Elem. i. 881 (1790) = **Cotula** Linn. (Compos.).

BALDINGERIA, F. W. Schmidt, Samml. Phys. Aufs i. 277 (1795) = **Leontodon**, Linn. (Compos.).
pygmaea, F. W. Schmidt, l. c. = L. hirtum ?
urens, F. W. Schmidt, l. c. = Hypochoeris cretensis

BALDWINIA, Nutt. Gen. Am. ii. 175 (1818) (Baldwina). *COMPOSITAE*, Benth. & Hook. f. ii 391.
ACTINOSPERMUM, Ell. Sketch. ii. 448 (1817).
ACTISPERMUM, Rafin. New Fl. Am. i. 61 (1836).
ENDOSMINA, Rafin. in Am. Monthly Mag. (1819) 195.
MNESITHEON, Rafin. Fl. Ludov 67 (1817).
bicolor, *Rafin. New Fl. Am.* iv. 73.—Am. bor.
lutea, *Rafin. l. c.*—Am. bor
multiflora, *Nutt. Gen. Am.* ii. 175.—Am. bor
uniflora, *Nutt. l. c.*—Am. bor.

BALDWINIA, Rafin. in Am. Monthly Mag (1818) 267 = **Passiflora**, Linn.
peltata, Rafin. l. c. = P. peltata.

BALENDASIA, Rafin. Fl. Tellur. iv. 105 (1836) = **Passerina**, Linn. (Thymel.).
ericoides, Rafin l. c. = P. ericoides.

BALENERDIA, Comm. ex Steud. Nom. ed. I. 99 (1821) = **Nanodea**, Banks (Santal.).

BALESSAM, Bruce, Trav. v. t. 25 (1790) = **Balsamodendron**, Kunth, sec. Endl. Gen. 1256 (Burserac.).

BALFOURIA, R. Br. in Mem. Wern. Soc. i. (1810) 70. = **Wrightia**, R. Br. (Apocyn.).
saligna, R. Br. Prod. 467.—W. saligna.

BALFOURODENDRON, Mello, ex Oliver, in Hook. Ic. Pl. xiii. 8. t. 1203–04 (1877). *RUTACEAE.*
eburneum, *Mello, l. c.*—Bras.

BALFURIA, Reichb. Consp. 184 (1828) = BALFOURIA, R. Br. = **Wrightia**, R. Br. (Apocyn.).

BALINGAYUM, Blanco, Fl. Filip. ed. I. 187 (1837) = **Erythropalum**, Blume (Olacin.).
decumbens, Blanco, l. c.—Ins. Philipp.

BALIOSPERMUM, Blume, Bijdr. 608 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 324.
angulare, Decne. ex Baill. Etud. Gén. Euph. 395 = axillare.
axillare, *Blume, Bijdr.* 604.—Ind. or. ; Malaya.
calycinum, *Muell. Arg. in Flora*, xlvii. (1864) 470.—Reg. Himal.
indicum, Decne. in Jacquem. Voy. Bot. 154. t. 155 = axillare.
micranthum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 215.—Reg. Himal.
montanum, Muell. Arg. in DC. Prod. xv. ii. 1135 = axillare.
Moritzianum, Baill. Etud. Gén. Euph. 395 = axillare.
polyandrum, Wight, Ic. t. 1885 = axillare.
reticulatum, *Kurz, in Flora*, lviii. (1875) 32.—Siam.
sinuatum, *Muell. Arg. in Flora*, xlvii. (1864) 470.—Reg. Himal.

BALLARDIA, Montr. in Mém. Acad. Lyon, x. (1860) 204 = **Clodesia**, Brongn. (Myrtac.).
elegans, Montr. l. c. 205.—N. Caled.

BALLARION, Rafin. in Am. Monthly Mag. (1818) 266 = **Stellaria**, Linn. (Caryophyll.).

BALLELA, Rafin. Fl. Tellur. iv. 70 (1836) = Campanula ? Linn.
repens, Rafin. l. c. = C. repens.

BALLEXERDA, Comm. ex A. DC. Prod. xiv. 675 (1857) = Nanodea, Banks (Santal.).

BALLIERIA, Juss. Gen. 188 (1789) = BAILLIERIA, Anbl. = Chiodalum, Linn. (Compos.).

BALLIMON, Rafin. New Fl. Am. iv. 28 (1836) = Daucus, Tourn. (Umbellif.).
maritimum, Rafin. l. c. = D. maritimus.
muricatulum, Rafin. l. c. = D. muricatus.

BALLOCHIA, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 86. *ACANTHACEAE.*
amoena, *Balf. f. l. c.* 87.—Ins. Socotr.
atro-virgata, *Balf. f. l. c.*—Ins. Socotr.
rotundifolia, *Balf. f. l. c.*—Ins. Socotr.

BALLOSPORUM, Salisb. Gen. Pl. Fragm. 142 (1866) = Gladiolus, Linn. (Irid.).
segetum, Salisb. l. c. = G. segetum.

BALLOTA, Linn. Gen. ed. I. 175 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1212.
ACANTHOPHASUM, Spach, in T. Nees, Gen. Fl. Germ. Gamop. i. (1843).
BERINGERIA, Neck. Elem. i. 312 (1790).
PSEUDODICTAMNUS, Tourn. ex Adans. Fam. ii. 191 (1763).
acetabulosa, *Benth. Lab. Gen. et Sp.* 595.—Graecia; As. Min.
africana, *Benth. l. c.* 594.—Afr. austr.
africana, Colm. Apuntes, 121 = hirsuta.
alba, *Linn. Fl. Suec. ed. II.* 308. n. 580.—Europ.
amplicata, Willd. ex Steud. Nom ed. II. i. 181 = nigra.
arabica, Hochst. & Steud. ex Steud. l. c. = Leucas urticaefolia.
aristata, Reichb. Fl. Germ. Excurs. 325 = nigra.
armena, Boiss. Diagn. Ser. II. iv. 54 = rotundifolia.
Aucheri, *Boiss. l. c.* Ser. I. v. 89.—Kurdist.; Persia.
borealis, Schweig. in Koenigsb. Arch. Naturw. i. (1812) 214 = alba.
bracteosa, Ball, in Journ. Linn. Soc. xvi. (1878) 629 = nigra.
bullata, *Pomel, Nouv. Mat. Fl. Atl.* 116.—Algeria.
cinerea, D. Don, Prod. Fl. Nep. 111 = Roylea elegans.
damascena, *Boiss. Diagn.* Ser. I. xii. 87.—Syria; Aegypt.
dictamnifolia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = pseudodictamnus.
distichoides, Linn. Mant. 88 = Anisomeles ovata.
distichoides, Rodech. Obs. 88 = Hyptis pectinata.
foetida, Lam. Fl. Fr. ii. 381 = nigra.
Forskahlei, *Benth. Lab. Gen. et Sp.* 599.—Arabia.
frutescens, Cos. ex Nym. Consp. 581 = spinosa.
glandulifera, Tausch, in Act. Hort. Petrop. iii. 11. (1875) 270 = rotundifolia.
Hildebrandtii, *Vatke & Kurtz, in Oestr. Bot. Zeitschr.* xxv. (1875) 95.—Afr. or.
hirsuta, *Benth. Lab. Gen. et Sp.* 595.—Hispan.; Afr. bor.
hirsuta, Schult. ex Steud. Nom. ed. II. i. 181 = nigra.
hirta, Hort. Berol. ex Steud. l. c. = nigra.
hispanica, *Benth. Lab. Gen. et Sp.* 597.—Europ. austr.
hispanica, Willk. & Lange, Prod. Fl. Hisp. ii. 446, in syn. = hirsuta.
integrifolia, *Benth. Lab. Gen. et Sp.* 599.—Ins. Cyprus.
italica, Benth. in DC. Prod. xii. 519 = rupestris.
lanata, Linn. Sp. Pl. 582 = Leonurus lanatus.
lanata, Willd. ex Ledeb. Fl. Ross. iii. 439 = Eremostachys laciniata.
lavendana, *Boiss. & Heldr. Diagn.* Ser. I. xii. 88.—As. Min.
limbata, Benth. Lab. Gen. et Sp. 748 = Otostegia limbata.
macrodonta, *Boiss. & Bal. Diagn.* Ser. II. iv. 54.—Cappadoc.
mauritanica, Steud. Nom. ed. II. i. 181 = seq.
mauritiana, Pers. Syn. ii. 136 = Anisomeles ovata.
microphylla, Benth. Lab. Gen. et Sp. 596 = Otostegia Schimperi.
mollissima, *Benth. l. c.* 595.—Hispan.
nigra, *Linn. Sp. Pl.* 583.—Europ.; Orient; Afr. bor.
obliqua, Benth. Lab. Gen. et Sp. 596 = saxatilis.
orbicularis, Lag. ex Willk. & Lange, Prod. Fl. Hisp. ii. 446 = hirsuta.

BALLOTA :—
orientalis, C. & J. Presl, Delic. Prag. 80 (Sp. dubia negligenda, Boiss.).—As. Min.
persica, Benth. Lab. Gen. et Sp. 595 = Otostegia macrophylla.
pilosa, Lour. Fl. Cochinch. 364 = Leucas Benthamiana.
pseudodictamnus, *Benth. Lab. Gen. et Sp.* 594.—Graecia; Creta.
rotundifolia, *C. Koch, in Linnaea*, xxi. (1848) 697.—Armenia.
roylecides, *Benth. in DC. Prod.* xii. 520.—Arab.
rubra, Schrad. ex Steud. Nom. ed. II. i. 182 = nigra.
ruderalis, Sw. in Palmstr. Sv. Bot. t. 389 (1809) = nigra.
rugosa, Benth. Lab. Gen. et Sp. 596 = saxatilis.
rupestris, *Vis. Fl. Dalm.* ii. 216.—Dalmat.
Russeliana, Benth. Lab. Gen. et Sp. 597 = saxatilis.
saxatilis, Guss. Fl. Sic. Syn. ii. 82 = rupestris.
saxatilis, *Sieber, ex Benth. Lab. Gen. et Sp.* 596.—Syria; As. Min.
Schimperi, Benth. in DC. Prod. xii. 519 = Otostegia Schimperi.
sepium, Paulet, in Pers. Syn. ii. 135 = nigra.
silvestris, Hoffmgg. & Link, ex Willk. & Lange, Prod. Fl. Hisp. ii. 446 = alba.
sordida, Salisb. Prod. 82 = nigra.
spinosa, *Link, Handb.* 476.—Gallia.
stachydiformis, *Hochst. in Flora*, xxvii. (1844) 100.—Abyss.
suaveolens, Linn. Syst. ed. X. 1100; Fl. Jam. Pugill. 15 = Hyptis suaveolens.
tomentosa, Rochsch. Obs. 85 = Hyptis pectinata.
undulata, *Benth. Lab. Gen. et Sp.* 595.—Syria; Arab.
urticaefolia, Ortim. ex Reichb. ic. Bot. viii. 30. t. 774 = nigra.
verticillata, *Noronha, in Verh. Batav. Gen. v.* (1790) ed. I. Art. iv. 8.—Malaya.
vulgaris, Hoffmgg. & Link, Fl. Port. t. 115 = nigra.

BALMEDA, Nocca, Hort. Ticin. 21 (1804) = Grewia, Linn. (Tiliac.).
corylifolia, Nocca, l. c. = G. corylifolia.

BALMISA, Lag. Gen. et Sp. Nov. 17 (1816) = Arisarum, Targ.-Tozz. (Aroid.).
vulgaris, Lag. l. c. = A. vulgare.

BALOGHIA, Endl. Prodr. Fl. Norf. 84 (1833). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 800.
STEIGERIA, Muell. Arg. in Linnaea, xxxiv. (1865) 15.
alternifolia, *Baill. Adansonia*, ii. (1861–62) 216.—N. Caled.
carunculata, *Baill. l. c.* 216.—N. Caled.
lucida, *Endl. Prod. Fl. Norf.* 84.—Ins. Norfolc.
Pancheri, *Baill. Adansonia*, ii. (1861–62) 214.—Austral.; N. Caled.

BALOSKION, Rafin. Fl. Tellur. iv. 32 (1836) = Rostic, Linn.
dichotomum, Rafin. l. c. = R. quadrifarius.

BALSAMAKIA, Lour. Fl. Cochinch. ii. 499 (1790) = Calophyllum, Linn. (Guttif.).
Inophyllum, Lour. l. c. 470 = C. Inophyllum.

BALSAMEA, Gled. in Berl. Ges. Naturf. Fr. Schr. iii. (1782) 287 = BALSAMODENDRUM, Kunth = Commiphora, Jacq. (sec. Engler) (Burser.).
abyssinica, Engl. in Bot. Jahrb. i. (1881) 41 = C. abyssinica.
africana, Baill. Hist. Pl. v. 295 = C. africana.
Agallocha, Baill. l. c. = C. Agallocha.
Berryi, Engl. in Bot. Jahrb. i. (1881) 41 = C. Berryi.
capensis, Engl. l. c. 43 = C. capensis.
caudata, Engl. l. c. = C. caudata.
edulis, Baill. in Bull. Soc. Linn. Par. i. (1877) 122 = C. edulis.
Harveyi, Engl. in Bot. Jahrb. i. (1881) 43 = C. Harveyi.
Hildebrandtii, Engl. l. c. = C. Hildebrandtii.
Kotschyi, Engl. l. c. 41 = C. africana.
madagascariensis, Engl. l. c. 42 = C. gnaidilobia.
marocanensis, Gled. in Berl. Ges. Naturf. Fr. Schr. iii. (1782) 137 = C. Opobalsamum.
mollis, Engl. in Bot. Jahrb. i. (1881) 42 = C. mollis.
Mukul, Baill. Hist. Pl. v. 295 = C. Mukul.
Myrrha, Baill. l. c. 294 = C. Myrrha.
pedunculata, Engl. in Bot. Jahrb. i. (1881) 43 = C. pedunculata.
pilosa, Engl. l. c. 41 = C. pilosa.

BALSAMEA :—
Roxburghii, Engl. in Bot. Jahrb. i. (1881) 41 = C. Agallocha.
Schimperi, Engl. l. c. = C. Schimperi.
Stocksiana, Engl. l. c. = C. Stocksiana.
zanzibarica, Baill. in Adansonia, xi. (1874) 180 = C. zanzibarica.

BALSAMIFLUA, Griff. Notul. iv. 382 (1854) = Populus, Linn. (Salicin.).
deltoides, Griff. l. c. = P. suphratica.

BALSAMINA, Tourn. ex Scop. Fl. Carn. ed. II. ii. 188 (1772); Gaertn. Fruct. ii. 151. t. 118 (1791) = Impatiens, Linn. (Geraniac.).
angustifolia, Blume, Bijdr. 239 = I. Balsamina.
bifida, Dc. Prod. i. 686 = I. flaccida.
capensis, Dc. l. c. = I. capensis.
carnea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 7.—Malaya.
chinensis, DC. Prod. i. 686 = I. chinensis.
coccinea, DC. l. c. 685 = I. Balsamina.
cochleata, DC. l. c. 686 = I. cochleata.
communis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 7.—Malaya.
comorensis, Boj. ex Baker, in Journ. Linn. Soc. xx. (1884) 114 = I. comorensis.
cornuta, DC. Prod. i. 686 = I. Balsamina.
cristata, Ser. Cat. Gr. Jard. Lyon, (1858) = I. scabrida ?
fasciculata, DC. Prod. i. 686 = I. chinensis.
foemina, Gaertn. Fruct. ii. 152 = I. Balsamina.
fruticosa, Lasch, ex Steud. Nom. ed. II. i. 182 = I. fruticosa.
fulva, Ser. Fl. Jard. iii. 455 = I. fulva.
glandulifera, Ser. Cat. Gr. Jard. Lyon, (1858) = I. glandulifera.
glandulifera, Lyall, ex Baker, in Journ. Linn. Soc. xx. (1884) 113 = I. Lyallii.
heterophylla, G. Don, Gen. Syst. i. 749 = I. chinensis.
hirsuta, Blume, Bijdr. 240 = I. hirsuta.
hortensis, Desp. in Dict. Sc. Nat. iii. 485 = I. Balsamina.
jamensis, Blume, Bijdr. 240 = I. javensis.
lacca, Medic. Malv. 71 = I. Balsamina.
latifolia, DC. Prod. i. 686 = I. latifolia.
Leschenaultii, DC. l. c. = I. Leschenaultii.
longicornis, Ser. Fl. Jard. iii. 452 = I. leptocorsa ?
lutea, Delarb. Fl. Auv. ed. II. 304 = I. Noli-tangere.
macrochila, Ser. Fl. Jard. iii. 454 = I. Roylei.
madagascariensis, DC. Prod. i. 686 = I. madagascariensis.
Masterviana, Paxt. Bot. Mag. vi. 75 = I. Masteriana.
micrantha, Blume, Bijdr. 240 = I. leptocorsa.
minor, DC. Prod. i. 686 = I. Kleinii.
minutiflora, Span. in Linnaea, xv. (1841) 185 = I. Balsamina.
mollis, G. Don, Gen. Syst. i. 749 = I. Balsamina.
mutilis, DC. Prod. i. 686 = I. mutila.
myosoroides, DC. l. c. = I. myosoroides.
Noli-tangere, Scop. Fl. Carn. ed. II. ii. 184 = I. Noli-tangere.
odorata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 208 = I. Balsamina.
oppositifolia, DC. Prod. i. 686 = I. oppositifolia.
parviflora, Ser. Cat. Gr. Jard. Lyon, (1853) = I. parviflora ?
puberula, G. Don, Gen. Syst. i. 749 = I. puberula.
racemosa, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 208 = I. Balsamina.
repens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 8.—Malaya.
roemerifolia, DC. Prod. i. 686 = I. oppositifolia.
Roylei, Ser. Fl. Jard. iii. 455. t. 7 = I. Roylei.
salicifolia, Boj. ex Baker, in Journ. Linn. Soc. xx. (1884) 113 = I. salicifolia.
scabriuscula, G. Don, Gen. Syst. i. 748 = I. scabriuscula.
setacea, Coleb. ex Hook. Exot. Fl. ii. t. 127 = I. chinensis.
Tila, DC. Prod. i. 686 = I. tripetala ?
tricornis, Ser. Fl. Jard. iii. 452 = I. scabrida.

BALSAMITA, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) t. 1 = Chrysanthemum, Linn. (Compos.).
agrostifolia, Desf. l. c. 2 = C. fuscullosum.
amana, DC. Fl. Fr. iv. 167 = Tanacetum amanum.
Audibertii, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 362 = Tanacetum Audibertii.
corymbosa, Salzm. in Flora, iv. (1821) 123 = C. fuscullosum.
dentata, Wall. Cat. n. 826 = Myriactis Wallichii.

BALSAMITA:—
flabelliformis, Pers. Syn. ii. 408 = Pentzia flabelliformis.
grandiflora, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) 1 (= *Plagius grandiflorus*).—Afr. bor.
major, Desf. l. c. 8 = C. Balsamita.
multifida, Roxas Clem. Essay. Vid, 292 = Tanacetum annuum.
suaveolens, Pers. Syn. ii. 408 = C. Balsamita.
tridentata, Delile, Fl. Egypte, 278 = Cotula tridentata.
virgata, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) 2 = C. discoidea.
vulgaris, Willd. Sp. Pl. iii. 1802 = Chrysanthemum Balsamita.

BALSAMOCARPON, Clos, in C. Gay, Fl. Chil. ii. 226. t. 90 (1846) = Casealpinia, Linn. (Legumin.).
brevifolium, Clos, l. c.—Chili.

BALSAMODENDRON, DC. Prod. ii. 76 (1825) = seq.

BALSAMODENDRUM, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 348 = Commiphora, Jacq. (Burser.).
abyssinicum, Engl. in Bot. Jahrb. i. (1881) 41 = C. abyssinica.
africanum, Arn. in Ann. Nat. Hist. iii. (1839) 87 = C. africana.
Agallocha, Wight & Arn. l. c. 86 = C. Agallocha.
Arnottianum, March. in Adansonia, vii. (1866–67) 266 = C. pubescens.
Berryanum, Volgt, Hort. Suburb. Calc. 149 = C. Berryi.
Berryi, Arn. in Ann. Nat. Hist. iii. (1839) 86 = C. Berryi.
caperae, Sond. in Harv. & Sond. Fl. Cap. i. 526 = C. capense.
caudatum, March. in Adansonia, vii. (1866–67) 266 = C. caudata.
Ehrenbergianum, O. Berg, in Bot. Zeit. xx. (1862) 168 = C. Opobalsamum.
gileadense, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 349 = C. Opobalsamum.
habessinica, O. Berg, in Bot. Zeit. xx. (1862) 161 = C. abyssinica.
Kataf, Kunth, l. c. = C. Kataf.
Kotschyi, O. Berg, in Bot. Zeit. xx. (1862) 162 = C. africanum.
madagascariense, March. in Adansonia, viii. (1867–68) 71. t. 6 = C. grandifolia.
molle, Oliver, Fl. Trop. Afr. i. 326 = C. mollis.
Mukul, Hook. ex Stocks, in Hook. Kew Journ. i. (1849) 259 = C. Mukul.
Myrrha, T. Nees, Pl. Officin. t. 182 = C. Myrrha.
Opobalsamum, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 348 = C. Opobalsamum.
parviflorum, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1883) 506.—Ins. Socotra.
pedunculatum, Kotschy & Peyr. Pl. Tinn. 11. t. 5; et Schweinf. Beitr. Fl. Aeth. 234 = C. pedunculata.
planiflorum, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 506.—Ins. Socotra.
Playfairii, Hook. f. ex Oliver, Fl. Trop. Afr. i. 326 = C. Myrrha.
pubescens, Stocks, in Journ. As. Soc. Bomb. ii. (1848) 395. t. 25 = C. Stocksiana.
Roxburghianum, March. ex Baill. Adansonia, vii. (1866–67) 266 = C. candata.
Roxburghii, Arn. in Ann. Nat. Hist. iii. (1839) 86 = C. Agallocha.
Roxburghii, Stocks, in Journ. As. Soc. Bomb. ii. (1848) 393 = C. Mukul.
Schimperi, O. Berg, in Bot. Zeit. xx. (1862) 162 = C. Schimperi.
socotranum, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1883) 505.—Ins. Socotr.
Wightii, Arn. in Ann. Nat. Hist. iii. (1839) 86 = C. Mukul.
zeylanicum, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 349 = Canarium zeylanicum.

BALSAMONA, Vand. Fasc. Pl. 15 (1771) = Cuphea, P. Br. (Lythrar.).
Pinto, Vand. l. c. = C. Balsamona.

BALSAMOPHLEOS, O. Berg, in Bot. Zeit. xx. (1862) 168 = Commiphora, Jacq. (Burser.).
Kataf, O. Berg, l. c. = C. Kataf.

BALSAMORHIZA, Hook. Fl. Bor. Am. i. 310 (1888).
COMPOSITAE, Benth. & Hook. f. ii. 306.
EPELETIA, Nutt. in Journ. Acad. Philad. vii. (1834) 87. t. 4.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 356.—Calif.
Careyana, *A. Gray, Pl. Fendl.* 81.—Am. bor. occ.
deltoides, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 351.—Am. bor. occ.
glabrata, Benth. Pl. Hartw. 317 = deltoides.
helianthoides, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 351 = sagittata.
hirsuta, *Nutt. l. c.* 349.—Am. bor. occ.
Hookeri, *Nutt. l. c.*—Am. bor. occ.
incana, Nutt. l. c. 350 = Hookeri.
macrophylla, *Nutt. l. c.*—Am. bor. occ.
sagittata, *Nutt. l. c.*—Am. bor. occ.
silphioides, Geyer, ex Hook. Lond. Journ. Bot. vi. (1847) 244 = Silphium laeve.
terebinthacea, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 349.—Am. bor. occ.

BALSAMUS, Stackh. De Liban. ii (1814) = Commiphora, Jacq. (Burser.).
Kafal, Stackh. l. c. 12 = C. abyssinica.
libanotus, Stackh. l. c. 11 = C. Opobalsamum?
mecanensis, Stackh l. c. 12 = C. Opobalsamum.
Theophrasti, Stackh. l. c. 11 = C. Opobalsamum.

BALTIMORA, Linn. Mant. 158 (1771). *COMPOSITAE*, Benth. & Hook. f. ii. 348.
FOVEARIA, Moench, Meth. Suppl. 243 (1802).
FOUGEROUXIA, Cass. in Dict. Sc. Nat. liv. 461 (1829).
NIEMUHRIA, Scop. Introd. 184 (1777).
SCOLOSPERMUM, Less. in Linnaea, v. (1830) 152.
TIMANTHRA, Schrad. Gen. Pl. (1806).
alata, Meerb. Afb. t. 23 = recta?
alba, Pers. Syn. ii. 429 = recta.
recta, *Linn. Mant.* 288.—Mexic.
scabra, Pohl, ex Baker, in Mart. Fl. Bras. vi. iii. 258 = Calea hymenoleptis.
scolospermum, *Steetz, in Seem. Bot. Voy. Herald*, 154. —Panama.
trinervata, Moench, Meth. 592 = recta.

BAMBOS, Retz. Obs. v. 24 (1789) = Bambusa, Schreb. (Gramin.).
arundinacea, Retz. Obs. v. 24 = Bambusa arundinacea.
Arunda, Soland. Fl. Ins. Ocean. Pacif. 217 = Schizostachyum glaucifolium.
quinquefora, Stokes, Bot. Mat. Med. ii. 286 = Bambusa arundinacea.
stricta, Roxb. Pl. Corom. i. 58. t. 80 (non text.) = Oxytenanthera Thwaitesii, et Dendrocalamus strictus.
verticillata, Poir. Encyc. viii. 708 = Gigantochloa verticillata.

BAMBUS, J. F. Gmel. Syst. 579 (1791) = seq.

BAMBUSA, Schreb. Gen. Pl. 236 (1789). *GRAMINEAE*, Benth. & Hook. f. iii. 1210.
BAMBOS, Retz. Obs. v. 24 (1789).
BAMBUS, J. F. Gmel. Syst. 579 (1791).
BERZEB, Kunth, Syn. i. 288 (1822).
ISCHUROCHLOA, Buese, in Miq. Pl. Jungh. 389 (1854).
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 439 = Oxytenanthera abyssinica.
affinis, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 93.—Burma.
agrestis, *Poir. Encyc.* viii. 704.—Ind. or.; Cochinch.
alpina, Bory, Voy. i. 310. t. 19 = Nastus borbonicus.
amahosana, *Lindl. in Penny Cyclop.* iii. 357 (1835). —Amboina.
amplexifolia, Schult. f. Syst. vii. 1348 = Gaudua amplexifolia.
andamanica, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 58.—Ins. Andam.
angulata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 94.—China.
Apua, *Schult. f. Syst.* vii. ii. 1353.—Java.
argenteo-striata, *Regel, Gartenfl.* (1865) 268.—Japon.
aristata, *Lodd. & Lindl. in Penny Cyc.* iii. (1836) 357. —Ind. or.
arundinacea, Hamb. & Bonpl. Pl. Aequin. i. 71 = Nastus borbonicus.
BAMBUSA:—
arundinacea, Ait. Hort. Kew. ed. II. ii. 316 = Thouarsii.
arundinacea, Wight, ex Steud. Nom. ed. II. i. 188 = orientalis.
arundinacea, *Willd Sp. Pl.* ii. 245.—Ind. or.
Arundo, Nees, in Linnaea, ix. (1834) 471 = arundinacea.
Arundo, Wight, ex Steud. Nom. ed. II. i. 188 = orientalis.
aspera, *Schult. f. Syst.* vii. ii. 1352.—Amboina.
ater, *Lindl. in Penny Cyc.* iii. (1835) 357.—Amboina.
attenuata, Thw. Enum. Pl. Zeyl. 375 = Teinostachyum attenuatum.
aurea, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 716 = Phyllostachys aurea.
aurea, *Siebold, ex Miq. Ann. Mus. Bot Lugd. Bat.* ii. 285.—Japon.
aureo-striata, *Regel, Gartenfl.* (1865) 262.—Japon.
aurriculata, Kurz, in Journ. As. Soc. Beng. xxxix (1870) ii. 86 = vulgaris.
baccifera, Roxb. Hort. Beng. 25; Fl. Ind. ii. 197 = Melocanna bambusoides.
Balcooa, *Roxb. l. c.* 25, 196 —Ind. or.
barbata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* i. (1835) 657 = Nastus barbatus.
Beecheyana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 108.—China.
bifolia, Siebold, ex Munro, l. c. 86 = Phyllostachys bambusoides.
Bitting, *Schult. f. Syst.* vii. ii. 1354.—Java.
Blancoi, *Steud. Syn. Pl. Gram.* 381.—Ins. Philipp.
Blumeana, Hook. & Arn. Bot. Beech. Voy. 254 = tuldoides.
Blumeana, *Schult. f. Syst.* vii. ii. 1348.—Java.
Brandisii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 109.—Burma.
brevifora, *Munro, l. c.* 96.—China.
caesa, Sieb. & Zucc. ex Munro, l. c. 89 = nana.
calostachya, *Kurz, in Journ. As. Soc. Beng* xlii. (1873) ii. 250.—Burma.
Cantori, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 111. —China.
capensis, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 144.—Afr. austr.
capitata, Trin. l. c. Sc. Nat. i. (1835) 626 = Gaudua capitata.
capitata, Wall. & Griff. in Wall. Cat. n. 8913 = Cephalostachyum capitatum.
capitata, *Willd. ex Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 48.—Madag.
China, *Franch. & Sav. Enum. Pl. Jap.* ii. 607.—Japon.
Chusque, Poir. Encyc. Suppl. v. 494 = Chusquea scandens.
cornuta, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 118. —Java.
crinita, Thomas, ex Munro, l. c. 157 = nutans.
critica, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 250.—Pegu.
dichotoma, Donn, Hort. Cantab. ed. IV. 76 = Phyllostachys nigra.
diffusa, Blanco, Fl. Filip. ed. I. 369 = Gaudua distorta.
distorta, *Nees, in Linnaea*, ix. (1834) 470.—Bras.
divaricatachya, Munro, in Trans. Linn. Soc. xxvi. (1868) 126 = Gigantochloa heterostachya.
dumetorum, Hance, ex Walp. Ann. iii. 781 = Schizostachyum dumetorum.
Duquilioi, Hort. ex Carr. in Rev. Hortic. (1869) 292 = Phyllostachys Quilioi.
edulis, *Carr. l. c.* (1866) 380.—China.
elegantissima, Hasak. Pl. Jav. 42 = Beesha elegantissima.
excelsa, Miq. Fl. Ind. Bat. iii. 418 = maxima.
falcata, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 98 = Arundinaria falcata.
Falconeri, *Munro, in Trans. Linn. Soc. xxvi.* (1868) 95.—Reg. Himal.
Fax, Poir. Encyc. viii. 704 = Melocanna humilis.
fera, *Miq. Fl. Ind. Bat.* iii. 418.—Amboina.
fetuloea, Royle, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 96 = Falconeri.
flagelliformis, Griff. ex Munro, l. c. 150 = Dendrocalamus flagelliformis.
flexuosa, *Munro, l. c.* 101.—China.
flexuosa, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 753 = Phyllostachys flexuosa.

BAMBUSA:—
floribunda, *Zoll. & Mor. ex Steud. Syn. Pl. Gram.* 330. —Java.
floribunda, Munro, in Trans. Linn. Soc. xxvi. (1868) 34 = Thamnocalamus Falconeri.
Fortunei, *Van Houtte, Fl. des Serres*, xv. (1863) t. 1535.—Japon.
fragilis, Spruce, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 78 = Guadua latifolia.
gigantea, Wall. Cat. Bot. Gard. Calc. 79 = Dendrocalamus giganteus.
glauca, Brune, ex Rupr. in Mém. Acad. Péterab. Sér. VI. v. (1839) 103 = Arundinaria glaucescens.
glauca, Lodd. ex Lindl. in Penny Cyclop. iii. 357 (1835) = nana.
glaucescens, Siebold, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 39 = nana.
glaucifolia, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 147 = Schizostachyum glaucifolium.
globiflora, *Griseb. in Goett. Nachr.* (1866) 79.—Ind. or.
glomerata, Royle, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 147 = Dendrocalamus strictus.
gracilis, *Hort. ex A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 682.—Hab.?
gracilis, Wall. Cat. n. 5033 = Oxytenanthera nigrociliata.
Griffithiana, *Munro, in Trans. Linn. Soc. xxvi.* (1868) 99.—Burma.
Guadua, Humb. & Bonpl. Pl. Aequin. 168. t. 20 = Guadua angustifolia.
Helferi, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 114. —Burma.
heterocycla, *Carr. in Rev. Hortic.* (1878) 354.—Hab.?
hexandra, Roxb. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 147 = Dendrocalamus strictus.
Hookeri, *A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 649.—Hab.?
Horsfieldii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 115.—Java.
humilis, Reichb. ex Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 140 = nana.
khasiana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 97. —Reg. Himal.
kumasasa, Zoll. ex Steud. Syn. Pl. Gram. 331 = Phyllostachys kumasaca.
latiflora, *Kurz, in Journ. As. Soc. Beng.* xli. (1873) ii. 250.—Ind. or.
latifolia, Humb. & Bonpl. Pl. Aequin. i. 68. t. 21 = Guadua latifolia.
levis, *Blanco, Fl. Filip.* ed. I. 372.—Ins. Philipp.
Linn, *Blanco, l. c.* 371.—Ins. Philipp.
lineata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 118. —Ins. Ternate.
longinodis, *Mig. Fl. Ind. Bat.* iii. 418.—Malaya.
longispatha, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 250.—Burma.
Læcoïae, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 115.—Ins. Philipp.
Lumampao, *Blanco, Fl. Filip.* ed. I. 372.—Ins. Philipp.
Mac-Clellandi, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 114.—Burma.
macra, Wall. ex Munro, l. c. 84 = Thamnocalamus spathiflorus.
macrocalmis, *A. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 634.—Hab.?
Macula, Buch.-Ham. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 91 = Tulda.
maculosacoriensis, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 681 = vulgaris.
marginata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 114.—Burma.
Masters*ii*, *Munro, l. c.* 118.—Ind. or.
maxima, Buch.-Ham. in Wall. Cat. n. 5089 = Dendrocalamus Hamiltonii.
maxima, *Poir. Encyc.* viii. 704.—Amboin.; China.
Maximowiczii, Hort. ex Munro, in Gard. Chron. (1876) ii. 774 = Arundinaria Fortunei.
Metake, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 284 = Arundinaria japonica.
microphylla, Griff. Journ. i. 249, 269 (1847) = Arundinaria microphylla.
nitis, Blanco, Fl. Filip. ed. I. 371 = Blancoi.
nitis, Hort. Paris. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 16 = Arundinaria japonica.
nitis, *Poir. Encyc.* viii. 704.—China.
nitis, Hort. ex Carr. in Rev. Hortic. (1866) 330 = edulis.

BAMBUSA:—
monogyna, *Blanco, Fl. Filip.* ed. I. 368.—Ins. Philipp.
monogyna, Griff. Notul. iii. 69 = Dendrocalamus Hamiltoni.
multiplex, *Raeusch. Nom.* ed. III. 108; *Schult. f. Syst.* vii. 11. 1350.—Cochinch.
nana, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 199.—Japon.
Nessiana, Arn. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 103 = arundinacea.
nigra, Lodd. Cat. (1833) ex Loud. Hort. Brit. 124 = Phyllostachys nigra.
nigricans, Hort. ex Steud. Nom. ed. II. i. 183 = Phyllostachys nigra.
nigro-ciliata, Bucse, in Miq. Pl. Jungh. 389 = Oxytenanthera nigro-ciliata.
nutans, *Wall. ex Munro, in Trans. Linn. Soc.* xxvi. (1868) 92.—Reg. Himal.
Oldhami, *Munro, l. c.* 109.—China.
orientalis, *Nees, in Linnaea*, ix. (1834) 475.—Ind. or.
pallida, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 97. —Ind. or.
paniculata, Willd. ex Steud. Nom. ed. II. i. 183 = Nastus borbonicus!
parviflora, Schult. f. Syst. vii. 11. 1350 = Arthrostylidium maculatum.
picta, *Lindl. in Penny Cyclop.* iii. 357 (1835).—Ins. Ceram.
picta, Sieb. & Zucc. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 111 = Fortunei.
polymorpha, *Munro, l. c.* 98.—Burma.
prava, *Lindl. in Penny Cyclop.* iii. 357 (1835).—Amboina.
pseudoarundinacea, Steud. Syn. Pl. Gram. 330 = Gigantochloa verticillata.
pubervula, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 285 = Phyllostachys nigra.
pubescens, *Doell, in Mart. Fl. Bras.* ii. iii. 189.—Bras.
pubescens, Lodd. ex Lindl. Penny Cyclop. iii. 357 (1835) = Dendrocalamus strictus.
pungens, Blanco, Fl. Filip. ed. I. 370 = arundinacea.
pygmæa, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 286.— Japon.
quadrangularis, *Fenzi, in Bull. Soc. Tosc. Ort.* v. (1880) 401.—Japon.
Quilioi, Hort. ex A & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 697 = Phyllostachys Quilioi.
Ragamowskii, Hort.—Cf. Gard. Chron. (1876) ii. 847 = viscellata.
regia, *Thoms. Cat. Hort. Bot. Calc.* 79; *ex Munro, in Trans. Linn. Soc.* xxvi. (1868) 116.—Ind. or.
reticulata, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 148 = Phyllostachys bambusoides.
Ritcheyi, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 118.—Ind. or.
Rumphiana, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 86.—Ind. or.
ruscifolia, Siebold, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 157 = Phyllostachys kumasaca.
scandens, Blume, ex Nees, in Flora, vii. (1824) 291 = Dendrocalamus Tjankorreh.
Schimperiana, Steud. Syn. Pl. Gram. 330 = Oxytenanthera abyssinica.
schizostachyoides, Kurz, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 134 = Melocanna Kurzii.
scriptoria, Dennst. Schlüssel. Hort. Malab. 21 = Melocanna bambusoides.
senanensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 606.— Japon.
siamensis, Kurz, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 116 = regia.
Siebori, Griseb. Fl. Brit. W. Ind. 598 = vulgaris.
Simoni, Carr. in Rev. Hortic. (1866) 330 = Arundinaria Simoni.
spinosa, Nees, in Linnaea, ix. (1834) 475 = arundinacea.
spinosa, Blume, ex Nees, in Flora, vii. (1825) 580 = Blumeana.
spinosa, *Roxb. Hort. Beng.* 25; *Buch.-Ham. in Trans. Linn. Soc.* xiii. (1822) 480.—Ind. or.
stellig, Kurz, ex Munro, in Trans. Linn. Soc. Bot. Lugd. Bat. ii. 285 = nana.
striata, *Lodd. ex Lindl. Penny Cyclop.* iii. 357 (1835). —China.
stricta, Roxb. Hort. Beng. 25; Fl. Ind. ii. 198 = Oxytenanthera Thwaitesii & Dendrocalamus strictus.
striduia, *Mem. Cat.* 26.—Zeylan.
sulfurea, Carr. in Rev. Hortic. (1878) 379 = Phyllostachys sulphurea.

BAMBUSA:—
surinamensis, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 60 = vulgaris.
tabacaria, *Poir. Encyc.* viii. 705.—Amboina; Java.
Tagoara, Nees, Agrost. Bras. 532 = Guadua Tagoara.
Teke, *Miq. Fl. Ind. Bat.* iii. 418.—Malaya; China.
tenuis, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 110. —Ins. Macass.
teres, *Buch.-Ham. in Wall. Cat.* n. 5026 B.—Ind. or.
tessellata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 110.—China.
textoria, *Blanco, Fl. Filip.* ed. I. 370.—Ins. Philipp.
Thouarsii, Kunth, Not. Gen. Bambusa, 8 = vulgaris.
trigyna, Roxb. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 91 = Tulda.
Tristi, Nees, in Linnaea, ix. (1834) 469 = Guadua Tristi.
Tulda, Benth. Fl. Hongkong. 434 = tuldoides.
Tulda, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 198.—Ind. or.; Burma.
tuldoides, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 98. —China.
variegata, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 285 = Fortunei.
vasaria, Buch.-Ham. in Wall. Cat. n. 5025 = Balcoos.
vasaria, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 122. —Malaya.
verticillata, Benth. Fl. Hongkong. 434 = Dendrocalamus latiflorus.
verticillata, Kortler, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 147 = Dendrocalamus strictus.
verticillata, Hook. & Arn. Bot. Beech. Voy. 254 = Beechcyana.
verticillata, Willd. Sp. Pl. ii. 245 = Gigantochloa verticillata.
villosula, *Kurz, For. Fl. Brit. Burma*, ii. 558.— Burma.
violacaens, Carr. in Rev. Hortic. (1869) 292 = Phyllostachys violascens.
virgata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1835) 234 = Guadua virgata.
viridi-glaucescens, Carr. in Rev. Hortic. (1869) 292 = nana.
viridi-striata, Regel, Ind. Sem. Hort. Petrop. (1866) 77 = striata.
vulgaris, Nees, Fl. Afr. Austr. 462 = Balcoos.
vulgaris, *Schrad. in Wendl. Coll. Pl.* ii. 26. t. 47. —Mexic.
Wightii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 111.—Ind. or.

BAMIA, [Linn.] R. Br. ex Wall. Pl. Asiat. Rar. i. 39 (1830) = Hibiscus, Tourn. (Malvac.).
Abelmoschus, R. Br. ex Wall. Cat. n. 1915 = H. Abelmoschus.
angulosa, Wall. Cat. n. 1927 = H. angulosus.
betulifolia, Wall. l. c. 1918 = H. Abelmoschus.
cancellata, Wall. l. c. 1930 = H. cancellatus.
chinensis, Wall. l. c. 1916 = H. Abelmoschus?
crinita, Wall. l. c. 1922 = H. cancellatus.
fruticosula, Wall. l. c. 1921 = H. cancellatus.
magnifica, Wall. l. c. 1919 = H. hostilis.
Manihot, Wall. l. c. 1926 = H. Manihot.
multifurcata, Wall. l. c. 1917 = H. Abelmoschus.
pungens, Wall. l. c. 1924 = H. pungens.
rugosa, Wall. l. c. 1923 = H. rugosus.
tetraphylla, Wall. l. c. 1925 = H. tetraphyllus.

BAMMIA, Rupp. Fl. Jen. ed. Hall. 29 (1745) = prasc.

BANALIA, Moq. in DC. Prod. offi. ii. 278 (1849).
AMARANTACEÆ, Benth. & Hook. f. iii. 37.
brasiliana, Moq. l. c. (= Chamissoea sp. ?).—Bras.
occidentalis, Moq. l. c. 279 = Nitrophila occidentalis.
thyrsiflora, *Moq. l. c.* 278.—Ind. or.

BANARA, Aubl. Fl. Gui. i. 547. t. 217 (1775).
SAMYDACEÆ, Benth. & Hook. f. i. 796.
ARCEA, Schott, in Spreng. Syst. iv. Cur. Post. 407 (1827).
BOCA, Vell. Fl. Flum. 329 (1829) | v. t. 118 (1827).
CHRISTIANNIA, Presl, Rel. Haenk. ii. 91. t. 47 (1836).
PINEDA, Ruiz & Pav. Prod. 76. t. 14 (1794).
XYLADENIUM, Desv. in Hamb. Prod. Fl. Ind. Occ. 43 (1825).
brasiliensis, *Benth. in Journ. Linn. Soc.* v. (1861). *Suppl.* ii. 93.—Bras.

BANARA:—
brevifolia, Blanco, Fl. Filip. ed. I. 426 = Scolopia dasyanthera.
dioica, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 94—Mexic.
domingensis, *Benth, l. c.*—Ins. S. Doming.
fagifolia, Vahl, Symb. Bot. iii. 65 = guianensis.
glaberrima, *Wright, ex Griseb. Cat. Pl. Cub.* 8.—Cuba.
glauca, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 93 = Kuhlia glauca.
grandiflora, *Spruce, ex Benth. l. c.* 93.—Peruv.
guianensis, *Aubl. Pl. Gui.* i. 548. t. 217.—Guiana; Bras.
ibaguensis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 291.—N. Granat.
incana, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 94.—Peruv.
ixiifora, *Benth. l. c.* 91.—Peruv.
mexicana, *A. Gray, in Proc. Am. Acad.* v. (1861) 174. —Mexic.
mollis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 290. —N. Granat.
nitida, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) *Suppl.* ii. 93.—Peruv.
parviflora, *Benth. l. c.* 91.—Bras.
pubescens, *Spruce, ex Benth. l. c.* 92.—Ecuador.
racemosa, *Blanco, Fl. Filip.* ed. I. 425.—Ins. Philipp.
reticulata, *Griseb. Cat. Pl. Cub.* 8.—Cuba.
Roxburghii, *Spreng. Syst.* ii. 479 (omnino dubia).—Ind. or.
tomentosa, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 240.—Bras.
ulmifolia, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 91 = Kuhlia ulmifolia.
Vellozii, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 331.—Bras.

BANCROFFTIA, Steud. Nom. ed. II. i. 188 (1840), = seq.

BANCROFTIA, Billh. in Linnæa Samfund. Handl. i. 1839 (1838) 39 = **Arrascaëta**, Bancr. (Umbell.).
moschata, Billh. l. c. = A. moschata.
xanthorrhiza, Billh. l. c. = A. xanthorrhiza.

BANCROFTIA, Macfad. Fl. Jamaic. i. 112 (1837) = **Tovaria**, Ruiz & Pav. (Cappalid.).
diffusa, Macfad. l. c. = T. pendula.

BANDEIRAEA, Welw. ex Benth. & Hook. f. Gen. i. 577 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
GRIFFONIA, Baill. in Adansonia, vi. (1865) 188. t. 9. simplicifolia, *Benth. in Trans. Linn. Soc.* xxv. (1866) 306, in nota.—Afr. trop.
spectos, *Welw. ex Benth. l. c.* 306. t. 40.—Afr. trop. tenuiflora, *Benth. l. c.* 307.—Afr. trop.

BANDERIA, Baill. Hist. Pl. ii. 120 (1870) = **Bandeiraea**, Welw. (Legum.).

BANDURA, Burm. Thes. Zeyl. 42. t. 17 (1737) = **Nepenthes**, Linn.

BANFFYA, Baumg. Enum. Stirp. Transs. i. 385 (1816) = **Gypsophila**, Linn. (Caryophyll.).
petræa, Baumg. l. c. = G. transylvanica.

BANGLIUM, Buch.-Ham. ex Wall. Cat. sub n. 6579 (1830) = **Alpinia**, Linn. (Scitam.).
sulphureum, Buch.-Ham. l. c. = A. Hamiltoniana.

BANIA, Becc. Malesia, i. (1877) 162. *MENISPERMACEAE*.
thyrsiflora, *Becc. l. c.*—N. Guinea.

BANISTERIA, Linn. Syst. ed. II. 32 (1740). *MALPIGHIACEAE*, Benth. & Hook. f. i. 357.
ANOMALOPTERYS, G. Don, Gen. Syst. i. 647 (1831). HOLOSTALON, Reichb. Nom. 207 (1841).
acanthocarpa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 394.—Ant. austr.
aculeata, *Mill. Gard. Dict.* ed. VIII. n. 6.—Mexic.
adenostalon, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 406.—Bras.
adenopoda, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 40. —Bras.
albicans, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 181.—Bras.
albida, Willd. ex Steud. Nom. ed. II. i. 188 = Gaudichaudia albida.
allophila, Sieber, ex A. Juss. in Arch. Mus. Par. iii. (1843) 356 = Brachypterys borealis.
amœna, Mart. ex A. Juss. l. c. 571 = Hiraea Gaudichaudiana.
anguiana, Vell. Fl. Flum. 191; iv. t. 160 = Stigmaphyllon affine.
anguiana, Linn. Sp. Pl. 427 = Stigmaphyllon anguiosum.
angustifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 38. —Bras.
anisandra, Griseb. in Linnæa, xiii. (1839) 190 = Clausenianæa.
anisandra, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 47. —Bras.
antifebrilis, *Ruiz, ex Griseb. in Linnæa*, xxii. (1849) 15.—Peruv.
appendiculata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 418.—Ins. S. Vincent.
appendiculata, Lamb. ex DC. Prod. i. 592 = Heteropteris appendiculata.
argentea, *Spreng. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 393.—Am. austr.
argyrophylla, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 45.—Bras.
atrosanguinea, *A. Juss. in Arch. Mus. Par.* iii. (1843) 389.—Bolivia.
aurea, Vell. Fl. Flum. 192; iv. t. 167 = Gardneriana.
auriculata, Cav. Diss. ix. 428. t. 255 = Stigmaphyllon auriculatum.
bauhiniæfolia, Linn. Sp. Pl. 427 = Hiptage Madablota.
Benthamiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 390.—Peruv.
bicolor, *Poepp. ex A. Juss. l. c.* 393.—N. Granat.
Billbergiana, *Beurling, in Vet. Akad. Handl. Stockh.* (1854) 118.—Panama.
Blanchetiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 430.—Am. austr.
bogotensis, Bonpl. ex Triana & Planch. Prod. Fl. N. Granat. i. 318 = Stigmaphyllon bogotense.
bonariensis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 157.—Reg. Argent.
brachiata, Linn. Sp. Pl. 426 = Heteropteris brachiata.
brachiata, Spreng. Syst. ii. 387 = Heteropteris platypetala.
brachiata, Willd. ex A. Juss. in Arch. Mus. Par. iii. (1843) 356 = Brachypterys borealis.
brachyptera, DC. Prod. i. 591 = Brachypterys borealis.
bracteata, DC. l. c. 590 = Tetrapteris inæqualis.
brevipes, Sesse, ex DC. Prod. i. 591 = Gaudichaudia brevipes.
Caapi, *Spruce, ex Griseb. in Mart. Fl. Bras.* xii. 111. 48.—Bras.
caerulea, Lam. Encyc. i. 367 = Heteropteris caerulea.
Calcitrapa, Desv. in Ham. Prod. Fl. Ind. Occ. 40 = Brachypterys borealis.
calocarpa, *Miq. in Linnæa*, xviii. (1844) 58.—Guiana.
campestris, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 36. —Bras.
chrysophylla, Lam. Encyc. i. 368 = Heteropteris chrysophylla.
ciliata, Lam. l. c. 369 = Stigmaphyllon ciliatum.
cinerascens, *Griseb. in Linnæa*, xxii. (1849) 18.—Guiana.
Clausseniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 422.—Bras.
coccinea, *Vell. Fl. Flum.* 191. iv. t. 161.—Bras.
connexa, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 406 = stellaris.
constricta, *Griseb. in Mart. Fl. Bras.* xii. 111. 46.—Bras.
convolvulifolia, Cav. Diss. ix. 428. t. 256 = Stigmaphyllon convolvulifolium.
cordata, Vell. Fl. Flum. 191; iv. t. 163 = rotundifolia.
cordistipula, *Mart. Herb. Fl. Bras.* 556 = Peixotos cordistipula.
coriacea, *G. Don, ex Griseb. in Mart. Fl. Bras.* xii. 111. 94.—Bras.
cornifolia, *Spreng. Syst.* ii. 388.—N. Granat.
corymbosa, *Griseb. in Mart. Fl. Bras.* xii. 111. 42, in note.—Bras.
corymbosa, Griseb. in Linnæa, xxii. (1849) 18 = cinerascens.
cristata, *Griseb. l. c.* 18.—Guiana.
crotonifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 36. —Bras.
cujabensis, Griseb. in Linnæa, xiii. (1839) 194 = Hiraea cujabensis.
deformis, Desv. in Ham. Prod. Fl. Ind. Occ. 40 = Stigmaphyllon anguiosum.
dichotoma, Linn. Sp. Pl. 427 = Stigmaphyllon rotundifolium.
dichotoma, DC. Prod. i. 588 = Stigmaphyllon convolvulifolium.
divaricata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 412. —Guiana.
diversifolia, Bert. ex DC. Prod. i. 588 = varifolia.
diversifolia, *H. B. & K. Nov. Gen. et Sp.* v. 159.—Cuba.
eglandulata, Vell. Fl. Flum. 190; iv. t. 155 = Tetrapteris crebriflora.
eglandulosa, DC. Prod. i. 590 = Heteropteris suberosa.
eglandulosa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 410.—Bras.
elegans, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xviii. (1862) 322.—N. Granat.
elliptica, H. B. & K. Nov. Gen. et Sp. v. 161 = Stigmaphyllon ellipticum.
emarginata, *Cav. Diss.* ix. 426. t. 249.—Am. austr.
erianthera, *A. Juss. in Arch. Mus. Par.* iii. (1843) 411.—Bras.
fagifolia, DC. Prod. i. 590 = Hiraea fagifolia.
ferruginea, *Cav. Diss.* ix. 424. t. 248.—Bras.; Guian.
ferruginea, Sesse, Bot. Voy. Herald, 92 partim = Tetrapteris Seemanni.
fimbriata, *Vell. Fl. Flum.* 192; iv. t. 165.—Bras.
fimbriata, Griseb. in Linnæa, xiii. (1839) 191 = nigrescens.
Fischeriana, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1858) 48.—Cuba.
floribunda, *DC. Prod.* i. 589.—Bras.
fruticosa, Vell. Fl. Flum. 192; iv. t. 164 = Heteropteris umbellata.
fulgens, DC. Prod. i. 590 = Stigmaphyllon puberum.
fulgens, G. F. W. Mey. Prim. Fl. Esseq. 181 = ferrugines.
fulgens, *Linn. Sp. Pl.* 427.—Ins. Guadalup.
fulgens, *Lam.* Encyc. i. 368 = Stigmaphyllon fulgens.
Gardneriana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 420.—Bras.
glandulosa, Pav. ex A. Juss. l. c. 478 = Heteropteris rubiginosa.
glauca, Desf. Cat. Hort. Par. ed. III. 406 = Stigmaphyllon ciliatum.
glaucophylla, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 319 = Verrucularia glaucophylla.
grata, *Griseb. in Mart. Fl. Bras.* xii. 111. 49.—Bras.
heterophylla, Willd. Sp. Pl. ii. 742 = Stigmaphyllon fulgens.
heterostyla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 408.—N. Granat.
hexandra, Vell. Fl. Flum. 188; iv. t. 149 = Schwantia elegans.
Humboldtiana, DC. Prod. i. 588 = Stigmaphyllon Humboldtianum.
hypericifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 42.—Bras.
intermedia, *A. Juss.* in Arch. Mus. Par. iii. (1843) 404 = stellaris.
jasminifolium, *A. Juss. l. c.* 409.—N. Granat.
Kraussiana, *Hochst. in Flora*, xxvii. (1844) 1. 296.—Afr. austr.
lacviifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 38. —Bras.
lanciifolia, West, ex Schlecht. in Linnæa, iv. (1829) 81 = Heteropteris laurifolia.
latifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 40.—Bras.
laurifolia, Linn. Sp. Pl. ed. II. 611 = Heteropteris laurifolia.
laurifolia, Rich. ex A. Juss. in Arch. Mus. Par. iii. (1843) 380 = Stigmaphyllon periplocifolium.
laurifolia, Sieber, ex A. Juss. l. c. 394 = Byrsonima spicata.
ledifolia, *H. B. & K. Nov. Gen. et Sp.* v. 159.—Cuba.
ledocarpa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 395. —Peruv.
leona, *Cav. Diss.* ix. 424 (sp. dubia).—Afr. trop.
leptocarpa, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 130.—Guiana.

BANISTERIA :—
lobulata, *E. Mey. in Nov. Act. Nat. Cur.* xii. ii. (1825) 818.—Guiana
longifolia, Sw. Prod. Veg. Ind. Occ. 75 = Heteropteris platyptera.
lucida, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 109.—Guiana.
lupuloides, Linn. Sp. Pl. 427 = Gouania domingensis.
lutea, *Ruiz, ex Griseb. in Linnæa*, xxii. (1849) 15.—Peruv.
macradena, *DC. Prod.* i. 590.—Guiana.
macrocarpa, Pers. Syn. i. 507 = Heteropteris platyptera.
macrophylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 418.—Bras.
macrostachya, Vell. Fl. Flum. 189 ; iv. t. 152 = Clausenianæ.
magnoliæfolia, Desv. ex Ham. Prod. Fl. Ind. Occ. 40 = Heteropteris platyptera.
maracabensis, A. Juss. in Arch. Mus. Par. iii. (1843) 418 = comifolia.
maritima, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Brachypteryx borealis.
Martiniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 413.—Guiana.
mediterranea, Vell. Fl. Flum. 191 ; iv. t. 162 = Janusia muricata.
megacarpa, Vell. l. c. 189 ; iv. t. 139 = Stigmaphyllon tomentosum.
megaphylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 396.—Bras.
membranifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 39.—Bras.
metallicolor, *A. Juss. l. c.* 46.—Bras.
microphylla, *Vacq. Obs.* iii. 7. t. 56.—Carolin.
Minarum, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 409 = schuxoptera.
monoptera, Vell. Fl. Flum. 189 ; iv. t. 158 = Heteropteris chrysophylla.
montana, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 37 = campestris.
mucronata, DC. Prod. i. 589 = Stigmaphyllon ellipticum.
multiciliata, Vell. Fl. Flum. 189 ; iv. t. 154 = Tetrapteris punctulata.
multiflora, Boj. ex A. Juss. in Arch. Mus. Par. iii. (1843) 424 = Sphedamnocarpus madagascariensis.
multiflora, DC. Prod. i. 589 = Heteropteris platyptera.
multifoliolata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 402.—Bras.
murienta, *Cav. Diss.* ix. 423. t. 248. f. 2.—Peruv.
mutabilis, Vell. Fl. Flum, 193 ; iv. t. 168 = Thryallis brachystachya.
nigrescens, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 44.—Bras.
nitens, Griseb. in Linnæa, xiii. (1839) 198 = argyrophylla.
nitida, Vell. Fl. Flum. 188 ; iv. t. 148 = Stigmaphyllon ciliatum.
nitida, Lam. Encyc. i. 369 = Heteropteris nitida.
nitrosiodora, *Griseb. in Goett. Abh. xxiv.* (1879) 65.—Reg. Argent.
nummifera, *A. Juss. in Arch. Mus. Par.* iii. (1843) 391.—Bras.
ochnæfolia, Mart. ex A. Juss. l. c. 464 = Heteropteris byrsonimæfolia.
orinocoensis, *H. B. & K. Nov. Gen. et Sp.* v. 162.—Venezuela.
ovata, Cav. Diss. ix. 429. t. 257. f. 1 = Brachypteryx borealis.
oxyclada, *A. Juss. in Arch. Mus. Par.* iii. (1843) 396.—Bras.
Pagnus, Stend. Nom. ed. II. i. 184 = Heteropteris syringæfolia.
palmata, *Cav. Diss.* ix. 430. t. 257. f. 2.—Am. austr.
paniculata, DC. Prod. i. 591 = Heteropteris Beecheyana.
parviflora, *A. Juss. in Arch. Mus. Par.* iii. (1843) 416.—Bras.
parvifolia, Vent. Choix, 51 = Heteropteris parvifolia.
patens, Griseb. in Linnæa, xiii. (1839) 200 = Heteropteris patens.
pauciflora, *H. B. & K. Nov. Gen. et Sp.* v. 159.—Cuba.
periplocæfolia, *Desf. Tabl.* ed. II. 160 = Stigmaphyllon retusum.
pilosa, H. B. & K. Nov. Gen. et Sp. v. 160 = Brachypteryx borealis.

BANISTERIA :—
platyptera, *Griseb. in Linnæa*, xxii. (1849) 17.—Venezuela.
podaliræa, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 892.—Cuba.
polita, *Miq. in Verh. Nat. Wet. Haarl.* vii. (1851) 80. t. 21.—Guiana.
præcox, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 133.—Bras.
Pragua, Vell. Fl. Flum. 190 ; iv. t. 138 = Heteropteris syringæfolia.
pruinosa, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 392.—Bras.
pubera, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Stigmaphyllon puberum.
pubiflora, DC. Prod. i. 591 = Heteropteris lanifolia.
pubipetala, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 41. t. 169.—Bras.
purpurea, Linn. Sp. Pl. 427 = Heteropteris purpurea.
Quapara, *Aubl. Pl. Gui.* i. 464. t. 186.—Guiana.
Riedeliana, *Regel, Ind. Sem. Hort. Petrop.* (1855) 17.—Cuba.
rigida, *A. Juss. in Arch. Mus. Par.* iii. (1843) 416.—Bras.
sagittata, *Cav. Diss.* ix. 430. t. 257.—Ins. S. Doming.
salicifolia, *DC. Prod.* i. 590.—Bras.
schizoptera, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 43.—Bras.
Schlimii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 584.—N. Granat.
Schomburgkiana, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 129.—Guiana.
schwanniodes, *Griseb. in Mart. Fl. Bras.* xii. iii. 54.—Bras.
scutellata, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 131.—Bras.
Sellowiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 419.—Bras.
septum, *Mart. ex A. Juss. l. c.* 414.—Bras.
sericea, Cav. Diss. ix. 429. t. 258 = Heteropteris sericea.
sericea, Dombey, ex A. Juss. in Arch. Mus. Par. iii. (1843) 394 = acanthocarpa.
sinemariensis, Aubl. Pl. Gui. i. 462. t. 185 = Stigmaphyllon puberum.
sinuata, *DC. Prod.* i. 588.—Guiana.
solitaria, Vell. Fl. Flum. 189 ; iv. t. 151 = Hiraea Salzmanniana.
Spixiana, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 361 = Heteropteris suberosa.
spiendens, DC. Prod. i. 588 = Stigmaphyllon fulgens.
Spruceana, *Griseb. in Mart. Fl. Bras.* xii. iii. 45.—Bras.
stellaris, *Griseb. in Linnæa*, xiii. (1839) 192.—Bras.
subcordata, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 326 = adenopoda.
suberosa, Willd. ex Stend. Nom. ed. II. i. 184 = Heteropteris suberosa.
tenuis, Lindl. in Bot. Reg. (1838) Misc. 72 = Heteropteris umbellata.
tenuifolia, *Vell. Fl. Flum.* 192 ; iv. t. 166.—Bras.
ternata, Vell. l. c. 190 ; iv. t. 159 = trifoliata.
ternata, Sesse, ex DC. Prod. i. 591 = Stigmaphyllon elliptinum.
tetraptera, Sonner. Voy. Indes, iii. 270 = Hiptage Madablota.
thyrsoidea, Griseb. in Linnæa, xiii. (1839) 199 = Heteropteris thyrsoidea.
tiliæfolia, H. B. & K. Nov. Gen. et Sp. v. 162 = Stigmaphyllon Humboldtianum.
tiliæfolia, *Vent. Choix*, 50. t. 50.—Java.
timoriensis, *DC. Prod.* i. 588.—Ins. Timor.
tomentosa, Desf. Tabl. ed. II. 160 = smaragnista.
tomentosa, Salzm. ex A. Juss. in Arch. Mus. Par. iii. (1843) 388 = Stigmaphyllon rotundifolium.
tomentosa, Schlecht. in Linnæa, x. (1835) 244 = Heteropteris Beecheyana.
torifera, Vell. Fl. Flum. 190 ; iv. t. 156 = crotonifolia.
trifoliata, *Stend. Nom.* ed. II. i. 184.—Bras.
tristelloceps, Boj. ex A. Juss. in Arch. Mus. Par. iii. (1843) 408 = Tristellateia Bojeriana.
umbellulata, Sieber, ex A. Juss. l. c. 396 = Brachypteryx borealis.
umbellulata, *DC. Prod.* i. 592.—Ins. S. Doming.
umbellulata, Sieber, ex A. Juss. in Arch. Mus. Par. iii. (1843) 42 = Byrsonima trinitensis.
unilatus, Vell. Fl. Flum. 190 ; iv. t. 157 = B. ferruginea.
ventenatoides, Lam. Encyc. i. 367 = Hiptage Madablota.

BANISTERIA :—
varia, Spreng. Syst. ii. 386 = varifolia.
varifolia, *DC. Prod.* i. 588.—N. Granat.
velutina, A. Juss. in Arch. Mus. Par. iii. (1843) 401 = crotonifolia.
verrucinervia, *Griseb. in Mart. Fl. Bras.* xii. iii. 398.—Bras.
verruculosa, *Sw. Ann. Bot. Posth.* 31.—Ind. occ.
vespertilionis, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 414 = pubipetala.
vitifolia, Sesse, ex DC. Prod. i. 588 = varifolia.
voinchita, Endl. ex Heynh. Nom. ii. 59 = Hiraea Simsiana.
zanzibarica, Boj. ex Steud. Nom. ed. II. i. 184 = Acridocarpus zanzibaricus.

BANTUM, Ces. ex Boiss. Fl. Orient. ii. 882 (1872) = Carum, Linn. (Umbell.).
ligusticifolium, Ces. ex Boiss. l. c. = C. multiflorum.

BANVOLIBA, Bowdich, Madeira, 258 (1825). *ACANTHACEÆ* ! (Genus omnino dub.).
violacea, *Bowdich, l. c.*—Madera.

BANKESIA, Bruce, Trav. ed. II. vii. 161 (1805) = **Brayera**, Kunth (Rosac.).

BANKSEA, Koen. in Retz. Obs. iii. 75 (1783) = **Costus**, Linn. (Scitamin.).
speciosa, Koen. l. c. = C. speciosus.

BANKSIA, Bruce, Trav. ed. II. vii. tt. 22, 23 = **Brayera**, Kunth (Rosac.).
abyssinica, 'Bruce', ex Steud. Nom. ed. II. i. 184 = Brayera anthelmintica.

BANKSIA, Domb. ex DC. Prod. iii. 88 (1828) = **Cuphea**, P. Br. (Lythrar.).
ciliata, Domb. ex DC. l. c. 88 = C. microphylla.
glutinosa, Domb. ex DC. l. c. = C. cordata.

BANKSIA, Forst. Char. Gen. 7. t. 4 (1776) = **Pimelea**, Banks (Thymel.).
Gnidia, Forst. l. c. 8 = P. Gnidia.
pilosa, Forst. f. in Nov. Act. Upsal. iii. (1780) 174 = P. pilosa.
prostrata, Forst. Char. Gen. 8 = P. prostrata.
tomentosa, Forst. l. c. = P. pilosa.

BANKSIA, Linn. f. Suppl. 15, 126 (1781). *PROTEACEÆ*, Benth. & Hook. f. iii. 184.
BANKSIINEÆ, R. Br. ex Steud. Nom. ed. II. i. 239 (1840).
aemula, *R. Br. in Trans. Linn. Soc.* x. (1811) 209.—Austral.
aemula, Sieber, ex Meisn. in DC. Prod. xiv. 461 = serrata.
aquifolia, Lindl. Swan Riv. App. 84 = ilicifolia.
asplenifolia, Salisb. Prod. 61 = integrifolia.
asplenifolia, Knight, Prot. 113 = marocscens.
attenuata, *R. Br. in Trans. Linn. Soc.* x. (1811) 209.—Austral.
australis, R. Br. l. c. 205 = marginata.
borbigera, Meisn. in Lehm. Pl. Preiss. ii. 264 = Goodii.
Banett, *R. Br. Prod. Nov.* 35.—Austral.
Baxteri, *R. Br. l. c.* 36.—Austral.
blechnifolia, F. Muell. Fragm. iv. 108 = repens.
Brownii, *Baxter, ex R. Br. Prod. Nov.* 37.—Austral.
Caleyi, *R. Br. l. c.* 35.—Austral.
Candolleana, *Meisn. in Hook. Kew Journ.* vii. (1855) 118.—Austral.
coccinea, *R. Br. in Trans. Linn. Soc.* x. (1811) 207.—Austral.
collina, *R. Br. l. c.* 204.—Austral.
compar, R. Br. l. c. 207 = integrifolia.
conchifera, Gaertn. Fruct. i. 221. t. 46 = serrata.
cunealfolia, Hofm. in Roem. & Schult. Syst. iii. Mant. 279 = integrifolia.
cunealfolia, *Wendl. f. ex Hoffmgg. Verz. Pfl. Nachtr.* ii. 64.—Austral.
Cunninghamii, Sieber, in Spreng. Syst. iv. Cur. Post. 67 = collina.
cylindrostachya, Lindl. Swan Riv. App. 84 = stenostela.
dactyloides, Gaertn. Fruct. i. 221. t. 47. f. 2 = Hakea dactyloides.
dentata, *Linn. f. Suppl.* 127.—Austral.
dentata, Wendl. Hort. Herrenh. t. 8 = serrata.
denticulata, Dum.-Cours. Bot. Cult. ed. II. vii. 196 = spinulosa.

BANKSIA:—
depressa, R. Br. in Trans. Linn. Soc. x. (1811) 205 = marginata.
dilleniaefolia, Knight, Prot. 118 = latifolia.
dryandroides, *Baxter, in Sweet, Fl. Austral.* t. 56.—Austral.
elatior, R. Br. in Trans. Linn. Soc. x. (1811) 209 = serrula.
elegans, *Meisn. in Hook. Kew Journ.* vii. (1855) 119.—Austral.
ericaefolia, *Linn. f. Suppl.* 127.—Austral.
fagifolia, Hoffm. ex Roem. & Schult. Syst. iii. Mant. 279 = latifolia.
ferrea, Vent. ex Spreng. Syst. i. 485 = marginata.
gibbosa, Sm. in White, Voy. 224. t. 23. f. 2 = Hakea gibbosa.
glauca, Cav. in Anal. Hist. Nat. i. (1799) 289 = integrifolia.
Goodii, *R. Br. Prot. Nov.* 86.—Austral.
grandidentata, Dum.-Cours. Bot. Cult. ed. II. vii. 108 = speciosa.
grandis, *Willd. Sp. Pl.* i. 535.—Austral.
Guentheri, *Regel, Ind. Sem. Hort. Petrop.* (1857) 37; *Gartenfl.* (1858) 80.—Austral.
Gunnii, Meisn. in Hook. Kew Journ. iv. (1852) 210 = marginata.
heterophylla, Hort. ex Meisn. in DC. Prod. xiv. 466 = Hakea wauvelensa.
Hookeri, Drumm. in Comp. Bot. Mag. (1848) 1 = Solandri.
Hookeriana, *Meisn. in Hook. Kew Journ.* vii. (1855) 119.—Austral.
Huegelii, *R. Br. ex Endl. Gen. Suppl.* iv. 11. 89.—Austral.
hypoleuca, Hoffmgg. Verz. Pfl. Nachtr. ii. 66 = marginata.
illicifolia, *R. Br. in Trans. Linn. Soc.* x. (1811) 211.—Austral. †
insularis, *R. Br. l. c.* 206 = marginata.
integerrima, *Dum.-Cours. Bot. Cult.* ed. II. ii. 421; vii. 108.—Austral. †
integrifolia, Labill. ex Meisn. in DC. Prod. xiv. 456 = marginata.
integrifolia, *Linn. f. Suppl.* 127.—Austral.
integrifolia, Schlecht. in Linnaea, xx. (1847) 584 = australis.
laevigata, *Meisn. in Hook. Kew Journ.* iv. (1852) 210.—Austral.
latifolia, *R. Br. in Trans. Linn. Soc.* x. (1811) 208.—Austral.
ledifolia, A. Cunn. ex Meisn. in DC. Prod. xiv. 454 = Cunninghamii.
Lemanniana, *Meisn. in Hook. Kew Journ.* iv. (1852) 210.—Austral.
Lindleyana, *Meisn. l. c.* vii. (1855) 120.—Austral.
littoralis, Lindl. in Bot. Reg. t. 1363 = collina.
littoralis, *R. Br. in Trans. Linn. Soc.* x. (1811) 204.—Austral.
longifolia, D. Dietr. Syn. i. 588 = latifolia.
longifolia, Hort. ex Stend. Nom. ed. II. i. 184 = oblongifolia.
longifolia, Doo, ex F. G. Dietr. Vollst. Gaertn. Lexik. ii. 150 = serrata.
macrophylla, Link, Handb. Hort. Berol. i. 116 = integrifolia.
marcescens, Bonpl. Jard. Malm. 116. t. 48 = marginata.
marcescens, Hook. Bot. Mag. t. 2608 = australis.
marcescens, *R. Br. in Trans. Linn. Soc.* x. (1811) 206.—Austral.
marginata, *Cav. in Anal. Hist. Nat.* i. (1799) 227.—Austral.
marginata, *R. Br. in Trans. Linn. Soc.* x. (1811) 204.—Austral.
media, *R. Br. Prot. Nov.* 85.—Austral.
Meisneri, *Lehm. Pl. Preiss.* i. 582.—Austral.
Menziesii, *R. Br. Prot. Nov.* 86.—Austral.
microstachya, Cav. in Anal. Hist. Nat. i. (1799) 234 = marginata.
microstylon, *Hort. Cantabr. ex F. G. Dietr. Vollst. Gaertn. Lexik.* ii. 156.—Austral.
mitis, Knight, Prot. 115 = serrata.
monosiliensis, *Gaertn. Fruct.* i. 221.—Austral.
nivea, Labill. Voy. i. 411. t. 24 = Dryandra nivea.
nutans, *R. Br. in Trans. Linn. Soc.* x. (1811) 208.—Austral.
oblongifolia, *Cav. Anal. Hist. Nat.* i. (1799) 235.—Austral.
oleaefolia, Lodd. Bot. Cab. t. 341 = marginata.

BANKSIA:—
occidentalis, *R. Br. in Trans. Linn. Soc.* x. (1811) 204.—Austral.
oleaefolia, Cav. in Anal. Hist. Nat. i. (1799) 228. t. 14 = integrifolia.
oleaefolia, Salisb. Prod. 51 = Hakea dactyloides
ornata, *F. Muell. ex Meisn. in Linnaea*, xxvi. (1853–55) 252.—Austral.
paludosa, R. Br. in Trans. Linn. Soc. x. (1811) 207 = linearis.
patula, R. Br. l. c. 206 = marginata.
petiolaris, *F. Muell. Fragm.* iv. 109.—Austral.
phylicaefolia, Salisb. Prod. 51 = ericaefolia.
pinifolia, Meisn. in Hook. Kew Journ. vii. (1855) 118 = sphaerocarpa.
pinifolia, Salisb. Prod. 51 = Hakea gibbosa.
pinnata, Hort. Cels, ex Roem. & Schult. Syst. iii. 444 = Hakea pinnata.
pinnatiserta, F. Muell. Fragm. vii. 58 = repens.
polycephalifolia, Knight, Prot. 118 = repens.
praemorsa, Andr. Bot. Rep. t. 258 = marcescens.
praemorsa, Hort. ex Dum.-Cours. Bot. Cult. vii. ed. II. 107 = marginata.
prionophylla, F. Muell. 1st Gen. Rep. 17 = collina.
prionotes, *Lindl. Swan River App.* 34.—Austral.
procumbens, Hort. ex Dum.-Cours. Bot. Cult. ed. II. vii. 107 = dentata.
prostrata, *R. Br. Prot. Nov.* 86.—Austral.
pubescens, *Donn, ex Roem. & Schult. Syst.* iii. 444.—Austral.
pulchella, *R. Br. in Trans. Linn. Soc.* x. (1811) 202.—Austral.
pyriformis, Gaertn. Fruct. i. 220. t. 47 = Xylomelum pyriforme.
quercifolia, *R. Br. in Trans. Linn. Soc.* x. (1811) 210.—Austral.
repens, *Labill. Voy.* i. 412. t. 23.—Austral.
reticulata, Hoffm. in Roem. & Schult. Syst. Mant. iii. 379 = integrifolia.
reticulata, *Wendl. f. ex Hoffmgg. Verz. Pfl. Nachtr.* ii. 8.—Austral.
Robur, Cav. in Anal. Hist. Nat. i. (1799) 226 = latifolia.
salicifolia, Cav. l. c. 230 = integrifolia.
Scoparium, *Meisn. in Hook. Kew Journ.* vii. (1855) 120.—Austral.
serraefolia, Knight, Prot. 112 = serrula.
serrata, *Linn. f. Suppl.* 136.—Austral.
serrata, Cav. Ic. vi. 37. t. 540 = serrula.
serratifolia, Salisb. Prod. 51 = serrula.
Solandri, *R. Br. Prot. Nov.* 86.—Austral.
speciosa, Lindl. in Bot. Reg. t. 1793 = Victoriae.
speciosa, *R. Br. in Trans. Linn. Soc.* x. (1811) 210.—Austral.
sphaerocarpa, *R. Br. l. c.* 208.—Austral.
spicata, Gaertn. Fruct. i. 221. t. 48 = integrifolia.
spinulosa, *Sm. New Holl. Pl.* i. 13. t. 4.—Austral.
tenuifolia, Salisb. Prod. 50 = Hakea scicularia.
teretifolia, Salisb. l. c. 51 = Hakea pugioniformis.
tomentosa, *F. Muell. in Hook. Kew Journ.* viii (1856) 227.—Austral.
tomentosa, Hort. ex Meisn. in DC. Prod. xiv. 454 = littoralis.
tricuspis, *Meisn. in Hook. Kew Journ.* vii. (1855) 118.—Austral.
trichantha, *Meisn. l. c.* 120.—Austral.
uncigera, Knight, Prot. 112 = latifolia.
undulata, Lindl. in Bot. Reg. t. 1816 = serrata.
verticillata, *R. Br. in Trans. Linn. Soc.* x. (1811) 207.—Australia.
Victoriae, *Meisn. in Hook. Kew Journ.* vii. (1855) 119.—Austral.

BANNISTERIA, Linn. Gen. ed. I. 377 (1737) = Banisteria, Linn. (Malpig.).

BANNISTERIOIDES, Linn. Fl. Zeyl. 192 (1747) = Pella, Gaertn. (Inc. sed.).

BAOBAB, Mill. Gard. Dict. ed. VI. 175 (1752) = Adansonia, Linn. (Malvac.).

BAPELA, Adol. ex Lodd. Bot. Cab. t. 867 (1825): *LEGUMINOSAE*, Benth. & Hook. f. i. 553.
BRACTEOLARIA, Hochst. in Flora, xxiv. (1841) II. 686.
CARPOLOBIA, G. Don, Gen. Syst. i. 370 (1831).
DELARIA, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 406.

BAPHIA:—
angolensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 249.—Afr. trop.
haematoxylon, Hochst. ex Hook. Niger Fl. 321 = nitida.
Hendelotiana, *Baill. Adansonia*, vi. (1866) 216.—Afr. trop.
Kirkii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 250.—Afr. trop.
laurifolia, Baill. Adansonia, vi. (1866) 213 = pubescens.
leptostemma, Baill. l. c. 214 = nitida.
maxima, *Baker, in Oliver, Fl. Trop. Afr.* ii. 250.—Afr. trop.
nitida, *Lodd. Bot. Cab.* t. 867.—Afr. trop.
pilosa, *Baill. Adansonia*, vi. (1866) 216.—Afr. trop.
polygalacea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 248.—Afr. trop.
pubescens, *Hook. f. in Hook. Niger Fl.* 320.—Afr. trop.
pyrifolia, Baill. in Bull. Soc. Linn. Par. i. (1885) 445 = polygalacea.
racemosa, *Hochst. in Flora*, xxiv. (1841) II. 688.—Afr. trop.
spathacea, *Hook. f. in Hook. Niger Fl.* 320.—Afr. trop.

BAPHIOPSIS, Benth. ex Baker, in Oliver, Fl. Trop. Afr. ii. 256 (1871). *LEGUMINOSAE*.
parviflora, *Benth. ex Baker, l. c.*—Afr. trop.

BAPHORHIZA, Link, Handb. i. 578 (1829) = Alkanna, Tausch (Borag.).
tinctoria, Link, l. c. = A. tinctoria.

BAPTISIA, Vent. Dec. Gen. Nov. 9 (1808). *LEGUMINOSAE*, Benth. & Hook. f. i. 466.
CROTALOPSIS, Michx. ex DC. Prod. ii. 100, in syn. (1825).
EAPLOSIA, LASINIA, PERICAULON, RIPASIA, Rafin. New Fl. Am. ii. 48–51 (1836).
alba, Hook. Bot. Mag. t. 1177 = leucantha.
alba, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 6.—Am. bor.
albiflora, *Rafin. New Fl. Am.* ii. 47.—Am. bor.
australis, *Sweet, Hort. Brit.* ed. II. 128.—Am. bor.
australis, Hort. ex Lehm. in Nov. Act. Nat. Cur. xiv. (1829) 808 = minor.
australis, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 6.—Am. bor.
bracteata, Muhl. ex Ell. Sketch, i. 469 = leucophaea.
caerulea, Eaton & Wright, Man. N. Am. Bot. 154 = australis.
calycosa, *Engelm. in Coult. Bot. Gaz.* iii. (1878) 66.—Florida.
confusa, Sweet, ex G. Don, Gen. Syst. ii. 113 = australis ?
exaltata, Sweet, Brit. Flow. Gard. i. t. 97 = australis.
fraxinifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 387, in obs. = Thermopsis fraxinifolia.
lanceolata, *Ell. Sketch*, i. 467.—Am. bor.
Lecontii, *Torr. & Gray, Fl. N. Am.* i. 396.—Florida.
leucantha, *Torr. & Gray, l. c.* 385.—Am. bor.
leucophaea, *Nutt. Gen. Am.* i. 283.—Am. bor.
lupinoides, *Burbidge, in Gard. Chron.* (1884) i. 827.—Am. bor. ?
megacarpa, *Chapm. ex Torr. & Gray, Fl. N. Am.* i. 396.—Florida.
microphylla, *Nutt. in Journ. Acad. Philad.* vii. (1834) 97.—Florida.
minor, Lehm. in Nov. Act. Nat. Cur. xiv. (1829) 808 = australis.
mollis, DC. Prod. ii. 100 = Thermopsis mollis.
mollis, Nutt. Gen. Am. i. 281 = Thermopsis fraxinifolia.
nepalensis, Hook. Exot. Fl. t. 131 = Piptanthus nepalensis.
perfoliata, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 5.—Am. bor.
retusa, *Rafin. New Fl. Am.* ii. 47.—Am. bor.
Serenae, *M. Curt. in Am. Journ. Sc.* Ser. II. vii. (1849) 406.—Am. bor.
simplicifolia, *Croom, in Am. Journ. Sc.* xxv. (1834) 74.—Florida.
sphaerocarpa, *Nutt. in Journ. Acad. Philad.* vii. (1834) 97.—Texas.
stipulacea, Ravenel, in Elliot Soc. Nat. Hist. i. (1856) t. 2 = microphylla.

BAPTISIA:—
sulphurea, *Engelm. in Coult. Bot. Gaz.* iii. (1878) 65. —Am. bor.
texana, Beckl. in Proc. Acad. Sc. Philad. '1861' (1862) 453 = australis?
tinctoria, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 6.—Am. bor.
undulata, Spreng. Syst. ii. 247 = lanceolata.
uniflora, *Nutt. Gen. Am.* i. 261, nomen—prsec.?
versicolor, Lodd. Bot. Cab. t. 1144 = *australis*?
versicolor, *Raf. New Fl. Am.* iv. 107, nomen.—Am. bor.
villosa, *Nutt. Gen. Am.* i. 281.—Carolina

BAPTISTONIA, Barb.-Rodr. Orch. Nov. i. 95 (1877). *ORCHIDEAE.*
echinata, *Barb.-Rodr. l. c.*—Bras.

BARANDA, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 502 = **Barringtonia**, Forst. (Myrtac.).
angustensis, Llanos, l. c. = B. macrostachya.

BARATRANTHUS, Miq. Fl. Ind. Bat. i. i. 684 (1855) = Loranthus, Linn.
axanthus, Miq. l. c.—Sumatra.

BARATTIA, A. Gray & Engelm. in Proc. Am. Acad. i. (1848) 48 = **Bnoella**, Adans. (Compos.).
calva, A. Gray & Engelm. l. c.—E. calva.

BARAULTIA, Spreng. Syst. ii. 818 (1825) = **Barraldeia**, Thou.

BARBACENIA, Vand. Fl. Lusit. & Bras. Spec. 21. t. 1 (1796). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 740.
CAMPDERIA, H. B. & K. Nov. Gen. et Sp. vii. 155 (1825).
PLEUROSTIMA, Raf. Fl. Tellur. ii. 97 (1836).
RADIA, A. Rich. in Kunth, Syn. Pl. Aequin. i. (1822).
VIENNA, Stend. ex Endl. Gen. 173 (1836).
Alexandrinæ, *Schnizl. ex Hook. Lond. Journ. Bot.* iv. (1845) 18.—Guiana.
bicolor, *Mart. Nov. Gen. et Sp.* i. 19. t. 13.—Bras.
brasiliensis, Willd. Sp. Pl. ii. 237 = Vandellii.
coccinea, *Mart. in Schult. f. Syst.* vii. 236.—Bras.
emilioides, *Mart. l. c.*—Bras.
exscapa, *Mart. Nov. Gen. et Sp.* i. 21. t. 14.—Bras.
flava, *Mart. ex Steud. Nom.* ed. II. i. 185.—Bras.
Gardneri, *Seub. in Mart. Fl. Bras. Ill.* i. 70. t. 8. f. 5. —Bras.
glauca, *Mart. in Schult. f. Syst.* vii. 238.—Bras
gracilis, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1881) 289 = Dasylirion acrotrichum.
ignea, *Mart. in Schult. f. Syst.* vii. 235.—Bras.
longiflora, *Mart. Nov. Gen. et Sp.* i. 19. t. 12.—Bras.
lanulosefolia, *Mart. in Schult. f. Syst.* vii. 235. —Bras.
macrantha, *Lem. Yard. Fl.* iv. (1854) t. 390.—Bras.
purpurea, *Hook. Bot. Mag.* t. 2777.—Bras.
Regnieri × , Hort. ex Moore & Ayres, Mag. Bot. ii. (1850) 309.—Bras.
rubro-virens, *Mart. Nov. Gen. et Sp.* i. 20. t. 14. f. 1.—Bras.
schidigera, *Lem. Yard. Fl.* ii. (1852) t. 198.—Bras.
squamata, *Lindl. in Bot. Reg.* (1848) *Misc.* 64.—Bras.
tomentosa, *Mart. Nov. Gen. et Sp.* i. 18. t. 11.—Bras.
tricolor, *Mart. l. c.* 18. t. 10.—Bras.
Vandellii, Schult. f. Syst. vii. 239 = glauca.
Vandellii, *Pohl, ex Seub. in Mart. Fl. Bras.* Ill. i. 72. —Bras.

BARBA-JOVIS, Adans. Fam. ii. 231 (1763) = **Anthyllis**, Linn. (Legumin.).
argyrophylla, Moench, Meth. 110 = A. Barba-Jovis.
linnæifolia, Moench, Meth. Suppl. 88 = A. Hermanniæ.
montana, Moench, l. c. 87 = A. montana.
tragacanthoides, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 347 = Astragalus armatus?

BARBAREA, Scop. Fl. Carn. ed. I. 529 (1760) = **Dentaria**, Linn. (Crucif.).

BARBAREA, R. Br. in Ait. Hort. Kew. ed. II. iv. 109 (1812). *CRUCIFERAE.* Benth. & Hook. f. i. 68. CAMPE, Dulac, Fl. Hautes-Pyr. 199 (1867).

BARBAREA:—
alpina, *C. Koch, in Linnæa*, xix. (1847) 819.—As. Min.
altaica, Andrz. ex Steud. Nom. ed. II. i. 185 = arcuata.
arcuata, Andrz. ex DC. Prod. i. 141 = taurica.
arcuata, *Reichb. in Flora*, v. (1822) 266.—Europ.; As. Min.; As. bor.
augustana, Boiss. Diagn. Ser. I. i. 69 = vulgaris.
australis, Hook. f. Fl. N. Zel. i. 14 = vulgaris.
australis, Jord. Diagn. i. 108 = præcox.
brachycarpa, *Boiss. Diagn.* Ser. I. v. 80.—As. Min.
bracteosa, *Guss. Fl. Sic. Prod.* ii. 257.—Sicilia.
brevicaulis, Jord. Diagn. i. 104 = rupicola.
brownifolia, Jord. l. c. 109 = præcox.
ciliciæ, Schott & Kotschy, ex Boiss. Fl. Orient. i. 185 = minor
ciliifera, *Steud. Nom.* ed. II. i. 185, nomen.—Græcia.
cretica, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 140.—Reg. Himal.
erysimoides, Schur, Enum. Pl. Transs. 41 = præcox.
heterophylla, *Willd. in Bot. Zeit.* v. (1847) 217.—Hispan.
hirsuta, Weihe, in Flora, xiii. (1830) 257 = vulgaris.
humilis, *Presl, Fl. Sic.* i. 45.—Sicilia.
iberica, DC. Syst. ii. 508 = arcuata.
Integrifolia, *DC. l. c.*—As. Min.
intermedia, Bor. Fl. Centr. Fr. ed. I. ii. 46 (1840) = vulgaris.
Kayserii, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 57 = vulgaris.
Linnæi, Spenn. Fl. Friburg. iii. 929 = præcox, vulgaris, taurica, &c.
longisiliqua, Jord. Diagn. i. 103 = præcox.
lyrata, Aschers. in Fl. Prov. Brandenb. i. 95 = vulgaris.
minor, *C. Koch, in Linnæa*, xix. (1847) 55.—As Min.; Syria.
oligosperma, *C. Koch, l. c.* 819.—Armenia.
orthoceras, *Ledeb. Hort. Dorp.* (1934).—Sibir. altaic.
palustris, Hegetschw. Fl. Schw. 654 = stricta.
parviflora, Fries, Novit ed. II. 207 = stricta.
patula, Fries, Nov. Mant. iii. 76 = præcox.
pecinata, *Presl, Nov. Mat. Fl. Sic.* 375.—Algeria.
planisiliqua, *C. A. Mey. in Middend. Reise (Fl. Ochot.* 14).—Sibir.
plantaginea, *DC. Syst.* ii. 208.—As. Min.; Persia.
præcox, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 109.—Europ.
prostrata, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1886) 346 = sicula.
pyrenaica, Jord. Diagn. i. 849 = vulgaris.
rivularis, Martr. Fl. Tarn. 44 = vulgaris.
rivularis, *Pearl. Elem.* 14.—Servia.
rupestris, Steud. Nom. ed. II. i. 185 = vulgaris.
rupicola, *Moris, Stirp. Sard. Elench.* i. 50.—Sardinia, sicula, Gren. & Godr. Fl. Fr. i. 92 = vulgaris.
sicula, *C. & P. Presl, Delic. Prag.* 17.—Ital.; As. Min.
stolonifera, *Pomel, Nouv. Mat. Fl. Atl.* 375.—Algeria.
stricta, Fellin. Fl. Vasc. Lapp. 6 = orthoceras.
stricta, *Andrz. in Bess. Enum. Pl. Volh.* 72.—Ross. med.
stricta, Willk. in Flora, xxxiv. (1851) 583 = vulgaris.
sylvestris, Jord. Diagn. i. 100 = vulgaris.
taurica, DC. Syst. ii. 207 = arcuata.
verna, Aschers. Fl. Prov. Brandenb. i. 96 = præcox.
vicina, Martr. Fl. Tarn, 45, in obs. = vulgaris.
vulgaris, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 109.—Europ.; As. temp.

BARBELLINA, Cass. in Dict. Sc. Nat. i. 440 (1827) = **Staehelina**, Linn. (Compos.).
sericea, Cass. l. c. = S. arborescens.

BARBERETTA, Harv. Gen. S. Afr. Pl. ed. II. 377 (1868). *HAEMODORACEAE*, Benth. & Hook. f. iii. 674.
aurea, *Harv. l. c.*—Afr. austr.

BARBERINA, Vell. Fl. Flum. 285 (1825); Ic. v. t. 117 (1827) = **Symplocos**, Linn. (Styrac.).
axillaris, Miers, in Journ. Linn. Soc. xvii. (1880) 294. —Ind. occ.
celastrinea, Miers, l. c. = S. celastrinea.
crenata, Miers, l. c.—Bras.
cubensis, Miers, l. c. = S. cubensis.
estrellensis, Miers, l. c. 293 = S. estrellensis.
hirsuta, Vell. Fl. Flum. 285; v. t. 117 = S. hirsuta.
lanceolata, Mart. Herb. Fl. Bras. n. 722 = S. lanceolata.
phaseoclades, Mart. l. c. 724 = S. phaseoclades.

BARBERINA:—
ramentacea, Miers, in Journ. Linn. Soc. xvii. (1880) 294 = S. ramentacea.
revoluta, Mart. Herb. Fl. Bras. n. 822 = S. revoluta.
rhamnifolia, Miers, in Journ. Linn. Soc. xvii. (1880) 293 = S. rhamnifolia.
Spruceana, Miers, l. c. 295.—Peruv.
tetrandra, Mart. Syst. Mat. Med. Bras. 49 = S. tetrandra.

BARBEUIA, Thou. Gen. Nov. Madag. 6 (1806). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 85. madagascariensis, *Steud. Nom.* ed. II. i. 186.—Madag.

BARBIERA, Spreng. Gen. ii. 587 (1831) = seq.

BARBIERIA, DC. Mém. Lég. 241. t. 29 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 495.
maynensis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 58.—Peruv.
pinnata, Baill. Hist. Pl. ii. 368 = polyphylla.
polyphylla, *DC. Mém. Lég.* vi. 249. t. 39.—Porto Rico.

BARBILUS, P. Br. Hist. Jamaic. 216 (1756) = **Trichilia**, Linn. (Meliac.).

BARBULA, Lour. Fl. Cochinch. 366 (1790) = **Caryopteris**, Bunge (Verben.).
sinensis, Lour. l. c. = C. incana.

BARBYLUS, Juss. Gen. 430 (1789) = **Trichilia**, Linn. (Melisc.).
Brownii, J. F. Gmel. Syst. 611 = T. hirta?
jamaicensis, DC. Prod. ii. 91 = T. hirta?

BARCELLA, Drude, in Engl. & Prantl, Bras. iii. 459. t. 106. (1881) = **Illaeis**, Jacq. (Palm.).
odora, Drude, l. c. = I. odora.

BARCENA, Dugès, in Rev. Cienc. Mexic. Dec. i. (1879) 8, cum tab.; et in La Naturaleza, iv. (1879) 281. *RHAMNEAE.*
guanajuatensis, *Dugès, ll. cc.*—Mexic.

BARCENIA, Dugès, in La Naturaleza, iv. (1879) 283 = prsec.

BARCKHAUSENIA, Menke, in Flora, xxvii. (1854) 723 = seq.

BARCKHAUSIA, DC. Fl. Fr. vi. 449 (1815) = BARKHAUSIA, Moench = **Crepis**, Linn. (Compos.).

BARCLAYA, Wall. in Trans. Linn. Soc. xv. (1827) 443. t. 18. *NYMPHAEACEAE*, Benth. & Hook. f. i. 47.
longifolia, *Wall. l. c.*—Burma.
Mottleyi, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 157. t. 21. f. 1–3.—Borneo.
oblonga, Wall. Cat. n. 599 = longifolia.

BARDANA, Hill, Veg. Syst. iv. 38 (1763) = **Arctium**, Linn. (Compos.).
Arctium, Hill, Hort. Kew. 64 = A. majus.
Lappa, Hill, Herb. Brit. i. 79 = A. majus.
minor, Hill, Hort. Kew. 64 = A. minus.

BARRERIA, Juss. in Dict. Sc. Nat. iv. 64 (1804) = BARRERIA, Scop. = **Poraquatiba**, Aubl. (Olacin.).

BARGEMONTIA, Gaudich. Voy. Bonite, t. 8. (1836–44?). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 880.
peruviana, *Gaudich. l. c.*—Peruv.

BARHAMIA, Klotzsch, in Seem. Bot. Voy. Herald, 104 (1853?) = **Croton**, Linn. (Euphorb.).
crassulosteris, Klotzsch, l. c. = C. populifolius.
hispida, Klotzsch, l. c. = C. populifolius.
macrostachya, Klotzsch, l. c. = C. populifolius.
multiplicata, Klotzsch, l. c. = C. urticæfolius.
ovalifolia, Klotzsch, l. c. = C. ovalifolius.
pauciserrata, Klotzsch, l. c. = C. populifolius.
urticæfolia, Klotzsch, l. c. = C. urticæfolius.

BARJONIA, Decne. in DC. Prod. viii. 512 (1844).
ASCLEPIADEÆ, Benth. & Hook. f. ii. 771.
chloræfolia, *Decne. l. c.*—Bras.
cymosa, *Fourn. in Mart. Fl. Bras.* vi. iv. 206.—Bras.
deltoidea, Decne. ex Fourn. l. c. 205 = racemosa.
linearis, *Decne. in DC. Prod.* viii. 512.—Bras.
obtusifolia, *Fourn. in Mart. Fl. Bras.* vi. iv. 206.—Bras.
racemosa, *Decne. in DC. Prod.* viii. 512.—Bras.
Warmingii, *Fourn. in Mart. Fl. Bras.* vi. iv. 206.—Bras.

BARKANIA, Ehrenb. [ex Reichb. Consp. 43 (1828)] in Abh. Acad. Berl. (1832) 429 = **Halophila**, Thou. (Hydroch.).
bullata, Ehrenb. in Abh. Acad. Berl. (1832) 429 = H. bullata.
punctata, Ehrenb. Symb. 5, ex Fritz. lc. Ind. Suppl. 43 = H. ovata.
stipulacea, Zanard. in Mem. Ist. Venet. vii. (1857) 235 = H. stipulacea.

BARKERIA, Knowles & Westc. Flor. Cab. ii. (1838) 7. t. 49 = **Epidendrum**, Linn. (Orchid.).
Barkeriola, Reichb. f. in Gard. Chron. (1884) ii. 616 = E. Barkeriola.
cycloteila, Reichb. f. l. c. (1880) i. 72 = E. cycloteilum.
elegans, Knowles & Westc. Flor. Cab. ii. (1838) 7 = E. elegans.
Lindleyana, Batem. ex Lindl. in Bot. Reg. (1842) Misc. 3 = E. Lindleyanum.
melanocaulon, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 22 = E. melanocaulon.
Skinneri, Paxt. Mag. Bot. xv. (1849) 1 = E. Skinneri.
spectabilis, Batem. in Lindl. Bot. Reg. (1842) Misc. 43 = E. spectabile.
Vanneriana, Reichb. f. in Gard. Chron. (1885) ii. 678 = E. Vannerianum.

BARKHAUSENIA. Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 69 = BORCKHAUSENIA, Gaert. Mey. & Scherb. = **Corydalis**, Vent. (Papav.).

BARKHAUSIA, Moench, Meth. 537 (1794) = **Crepis**, Linn. (Compos.).
aculeata, DC. Prod. vii. 159 = C. aculeata.
adenothrix, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 465 = C. Rueppellii.
albida, Cass. in Dict. Sc. Nat. xvi. 62 = C. albida.
alpina, DC. Fl. Fr. ed. III. iv. 41 = C. alpina.
amplexicaulis, Cos. & Dur. ex Cos. in Ann. Sc. Nat. Sér. IV. i. (1854) 282.—Algeria.
amplexifolia, Godr. in Mém. Acad. Montp. (Sect. Méd.) i. (1853) 487 = C. amplexifolia.
amygdalina, Lag. ex Stend. Nom. ed. II. i. 186 = C. foetida.
aparicioides, Spreng. Syst. iii. 652 = Chondrilla Petitdium.
arenaria, Pomel, Nouv. Mat. Fl. Atl. 4 = C. arenaria!
aspera, Reichb. in Mossel. Handb. ed. II. ii. 1405 = C. aspera.
auriculifolia, Reichb. l. c. 1410, in nota = C. incarnata!
Balbisiana, DC. Prod. vii. 155 = C. bursifolia.
balearica, Costa, Barcin. (1861) 6 = C. balearica.
ballotifolia, DC. Fl. Fr. v. 449 = C. ballotifolia.
Bivoniana, Reichb. in Mossel. Handb. ed. II. ii. 1408. —Europ.
bursifolia, Spreng. Syst. iii. 653 = C. bursifolia.
byzantina, DC. Prod. vii. 158 = C. foetida.
canspicua, Moris, Fl. Sard. 92 = C. canspicua.
Candolleï, Spreng. Syst. iii. 657 = C. foetida.
canescens, Spreng. l. c. 653, sub n. 16 = C. bursifolia.
carbonaria, Sch. Bip. in Mus. Senck. iii. 54 = C. carbonaria.
carolinioides, Nutt. Gen. Am. ii. 126 = Pyrrhopappus carolinianus.
cerinna, Reichb. in Mossel. Handb. ed. II. ii. 1406 = C. saxifraga.
chondrilloides, Bieb. & Fingerh. Comp. ii. 304 = C. foetidissima.
chondropsoides, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 864; (Lehm. Reliq. Bot. 208).—Turkest.
Clusiana, Pomel, Nouv. Mat. Fl. Atl. 4.—Algeria.

BARKHAUSIA:—
cornuta, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 27 = C. cornata.
cretica, Boiss. Diagn. Ser. I. xi. 53 = C. cretica.
decumbens, Gren. & Godr. Fl. Fr. ii. 332 = C. decumbens.
divaricata, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 26 = C. divaricata, *Lowe*.
divaricata, Schur, Enum. Pl. Transs. 378 = C. foetida.
dubia, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 27 —Ins. Madera.
echioides, Schur, Enum. Pl. Transs. 378 = C. foetida.
elata, Koch, in Linnæa, xxiii. (1850) 684 = C. alpina.
elegans, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 425 = C. elegans.
erucæfolia, Gren. & Godr. Fl. Fr. ii. 331 = C. crucæfolia.
fœtosa, DC. Prod. vii. 156 = C. nana.
floribunda, Pomel, Nouv. Mat. Fl. Atl. 260 = C. floribunda.
foetida, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 288 = C. foetida.
glandulosa, Pomel, Fl. Sic. p. xxxi = C. foetida.
gracilis, Dum. Fl. Belg. 61.—Belgium.
graveolens, Link, Enum. Hort. Berol. ii. 290 = C. foetida.
Hanseleri, Boiss. ex DC. Prod. vii. 158 = C. taraxacifolia.
heterocarpa, Boiss. Voy. Espagne, ii. 742 = C. taraxacifolia.
hirsutula, Spreng. Syst. iii. 652 = C. taraxacifolia.
hieracioides, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 27 = C. hieracioides.
hirsuta, Pomel, Nouv. Mat. Fl. Atl. 260 = C. hirsuta
hirta, DC. Koch, in Linnæa, xxiii. (1850) 688 = C. foetida.
hispida, Link, Enum. Hort. Berol. ii. 290 = C. setosa.
hyemalis, Bivona, Stirp. Rar. Sicil. i. [17] t. 2 = C. taraxacifolia.
interrupta, Reichb. in Mossel. Handb. ed. II. ii. 1407, in nota = C. Sieberi.
intybacea, Spreng. Syst. iii. 653 = C. taraxacifolia.
juvenalis, Delile, Ind. Sem. Hort. Monsp. (1856) 23 = C. juvenalis.
Kotschyana, Boiss. Diagn. Ser. I. vii. 13 = C. Kotschyana.
Kralikii, Pomel, Nouv. Mat. Fl. Atl. 5 = C. Kralikii.
lacinisata, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 26 = C. lacinisata.
leontodon, DC. Fl. Fr. ed. III. n. 2950 = C. leontodontoides.
leontodontoides, Spreng. Syst. iii. 653 = C. leontodontoides.
Lessingii, Hook. & Arn. Bot. Beech. Voy. i. 145 = Troximon aparsioides.
leucophæa, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 865; (Lehm. Reliq. Bot. 209).—As. centr.
leucorhoda, Reichb. in Mossel. Handb. ed. II. ii. 1409 = C. bicolor.
macrocephala, Willk. in Bot. Zeit. v. (1847) 860 = C. albida.
macrophylla, Spreng. Syst. iii. 652 = C. macrophylla.
Marschallii, C. A. Mey. Verz. Pfl. Casc. 57 = C. Marschallii.
melanocephala, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 865; (Lehm. Reliq. Bot. 209).—As. centr.
mucronata, Bertol. Fl. Ital. viii. 515 = C. alpestris.
muricata, Spreng. Syst. iii. 651 = C. aspera.
nana, DC. Prod. vii. 156 = C. nana.
nicæensis, Link, ex Spreng. Syst. iii. 653 = C. taraxacifolia.
nigricauna, Viv. ex Cos. in Bull. Soc. Bot. Fr. xii. (1865) 378 = C. nigricana.
nuda, Presl, ex Spreng. Syst. iii. 656 = C. leontodontoides.
nudicaulis, Viv. ex Cos. in Bull. Soc. Bot. Fr. xii. (1865) 378 = C. aspera.
numidica, Pomel, Nouv. Mat. Fl. Atl. 261 = C. numidica.
palustris, Besing. ex DC. Prod. vii. 170 = C. paludosa.
pinguis, Reichb. in Mossel. Handb. ed. II. ii. 1411 = C. foetida.
pterifolia, DC. Prod. vii. 157 = Pterotheca bifida.
pumilio, Reichb. in Mossel. Handb. ed. II. ii. 1409 = C. taraxacifolia.
pumilus, Dum. Fl. Belg. Prod. 61 = C. foetida.
purpurea, Bivona, Stirp. Rar. Mun. iv. 22 = Lagoseris purpurea.

BARKHAUSIA:—
radicata, Godr. in Mém. Acad. Montp. (Sect. Méd.). i. (1853) 486 = Hypochoeris radicata.
raphanifolia, Chev. ex Steud. Nom. ed. II. i. 187 = C taraxacifolia.
raphanifolia, Spreng. Syst. iii. 658 = C. raphanifolia.
recognita, DC. Prod. vii. 154 = C. recognita.
reflexa, C. & J. Presl, Delic. Prag. 111 = C. taraxacifolia.
repens, Spreng. Syst. iii. 652 —China.
rhoeadifolia, Bieb. Fl. Taur. Cauc. iii. 538 = C. foetida.
rodigiensis, Sch. Bip. ex Tchihat. Asie Min. Bot. ii. 374 = C. foetida.
Royleï, DC. Prod. vii. 157 = C. bifida.
rubra, Moench, Meth. 537 = C. rubra.
sardoa, Spreng. Syst. iv. Cur. Post. 304.—Sardin.
scapigera, Kunze, ex DC. Prod. vii. 152 (= *Macrorhynchus Pamphigii*).—Chili.
scariosa, Moench, Meth. 537 = C. alpina.
scariosa, Reichb. Fl. Germ. Exc. i. 256 = C. vesicaria.
Schimperi, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 466 = C. Schimperi.
Schultzii, Hochst. ex A. Rich. l. c. 464 = C. Schultzii.
senecioides, Spreng. Syst. iii. 652 = C. senecioides.
setosa, DC. Fl. Fr. ed. III. v. 44 = C. setosa.
spathulata, Spreng. Syst. iii. 651 = C. spathulata.
Sprengelii, Steud. Nom. ed. II. i. 187 = Troximon aparsioides.
stricta, Reichb. in Mossel. Handb. ed. II. ii. 1406 = C. neglecta.
Suffreniana, DC. Cat. Hort. Monsp. 88 = C. Suffreniana.
taraxacifolia, DC. Fl. Fr. iv. 48 = C. taraxacifolia.
taraxacifolia, Spreng. Syst. iii. 658 = C. taraxacoides.
taraxacoides, Reichb. in Mossel. Handb. ed. II. ii. 1406; DC. Prod. vii. 155 = C. taraxacoides.
tenella, Benth. in Hook. Lond. Journ. Bot. i. (1842) 488 = Lactuca versicolor.
tenerrima, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 464 = C. carbonaria.
tenerrima, Tenore, in Ann. St. Nat. Bolog. iv. 252 = C. leontodontoides.
tenuifolia, DC. Prod. vii. 155 = C. tenuifolia.
triangularis, C. Koch, in Linnæa, xxiii. (1850) 687 = C. foetida.
versicolor, Spreng. Syst. iii. 651 = Lactuca versicolor.
vesicaria, Spreng. l. c. 652 = C. vesicaria.
saxinthia, Margot. & Reut. Fl. Zanth. 62 = C. foetida

BARKHUSENIA, Hoppe, in Flora, xli. (1829) 512 = BARKHAUSIA, Moench = **Crepis**, Linn. (Compos.).

BARKLYA, F. Muell. in Journ. Linn. Soc. iii. (1859) 155; Fragm. i. 109 (1859). *LEGUMINOSÆ*, Benth. & Hook. f. i. 559.
syringifolia, *F. Muell. ll. cc.*— Austral.

BARLAEA, Reichb. f. in Linnæa, xli. (1877) 54 = **Habenaria**, Willd. (Orchid.).
calcarata, Reichb. f. l. c. = H. calcarata.

BARLERIA, Linn. Gen. ed. I. 669 (1787). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1091.
BARLERIACANTHUS, BARLERIANTHUS, BARLERIOPSIS, BARLERIOSIPHON, BARLERITES, DICRANACANTHUS, PRIONITIS et PSEUDOBARLERIA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 183–187.
BARLERIERA, J. F. Gmel. Syst. 959 (1791).
CRABBEA, Harv. Gen. Pl. Cap. ed. I. 376 (1838).
SOUBEYRANIA, Neck. Elem. i. 305 (1790).
WAHABIA, Fenzl, in Flora, xxvii. (1844) 312.
acanthoides, *Vahl, Symb. Bot.* i. 47.—Arab.; Aegypt.; Ind. bor. occ.
acanthophora, *Nees, in DC. Prod.* xi. 726.—Afr. trop.
aculeata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 85.—Ins. Socotr.
acuminata, Wight, Ic. t. 450 = tomentosa.
adolensis, *Deisle, in Rochet, Sec. Voy. Choa*, 341.—Afr. trop.
alata, *S. Moore, in Journ. Bot.* xviii. (1880) 266.—*Afr. trop.*
alba, Lodd. Bot. Cab. iv. t. 360 = cristata.
anomala, Spreng. Syst. ii. 838 = Bravaisia floribunda.
arabica, *Belang. ex Nees, in DC. Prod.* xi. 725.—Arabia.
argentea, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 96.—Ins. Socotr.

BARLERIA:—
Amottiana, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 311.—Zeylan.; Burma.
Aucheriana, Nees, in DC. Prod. xi. 284 = Hochstetteri.
auriculata, Schum. Beskr Guin. Pl. 386 = Hygrophila spinosa.
axillaris, Stend. Nom. ed. II. i. 187 = Lepidagathis trinervis ?
barbata, E. Mey. Zwei Pfl. Docum. 152 = Gueinzii.
Beddomei, T. Anders. ex Bedd. Ic. Ind. Or. i. 62 = Lawii.
bispinosa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 144 = B. Prionitis.
bispinosa, *Vahl, Symb. Bot.* i. 46.—Arab.
bispinosa, Nees, in Wall. Pl. As. Rar. iii. 94 = mysorensis.
bituminosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
Boivinii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 31.—Madag.
bractiata, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 490 = nitida.
brasiliensis, *Spreng. Neue Entdeck.* iii. 26.—Bras.
brevispina, K. Br. in Salt, Abyss. App. 64 = B. Prionitis.
Burchelliana, Nees, in DC. Prod. xi. 235 = macrostegia.
Burkeana, Sond. in Linnaea, xxiii. (1850) 92 = macrostegia.
buxifolia, *Linn. Sp. Pl.* 886.—Ind. or.
buxifolia, Roxb. Hort. Beng. 45; Fl. Ind. iii. 87 = obovata.
buxifolia, Wall. Cat. n. 2504 = mysorensis.
caerulea, Herb. Lamb. ex Nees, in DC. Prod. xi. 228 = cristata.
caerulea, Roxb. Hort. Beng. 45; Fl. Ind. iii. 89 = strigosa.
caerulea, Wall, Cat. n. 2509, partim = polytricha.
candida, *Nees, in DC. Prod.* xi. 240.—Orient.
capitata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 210.—Afr. trop.
cardiocalyx, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 105. —Afr. trop.
Carruthersiana, *S. Moore, in Journ. Bot.* xviii. (1880) 270.—Afr. trop.
chegoa, Herb. Madr. ex Wall. Cat. n. 2506 = cristata.
ciliata, E. Mey. ex Nees, in DC. Prod xi. 250 = Meyeriana.
ciliata, Roxb. Hort. Beng. 45; Fl. Ind. iii. 38 = cristata.
coccinea, Linn. Sp. Pl. 887 = Ruellia coccinea.
consanguinea, *Klotzsch, in Peters, Reise Mossamb. Bot.* 206.—Afr. trop.
conspicua, *Nees, in Wall. Pl. As. Rar.* iii. 92.—Ins. Penang.
cordifolia, Hochst. ex T. Anders. in Journ. Linn. Soc. vii. (1864) 29 = parviflora.
cornigera, Véry, ex Nees, in DC. Prod. xi. 248 = Hygrophila spinosa.
courtallica, *Nees, in Wight, Ic.* iv. t. 1529.—Ind. or.
cristata, Lam. Encyc. i. 380 = noctiflora.
cristata, *Linn. Sp. Pl.* 636.—Ind. or.; Burma.
cuspidata, *Heyne, ex Nees, in Wall. Pl. As. Rar.* iii. 93.—Ind. or.
cyanea, *S. Moore, in Journ. Bot.* xviii. (1880) 265.— Angola.
damarensis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 29.—Afr. austr. trop.
Decaisneana, *Nees, in DC. Prod.* xi. 230.—Madag.
diacantha, Hochst. ex Nees, l. c. 238 = B. Prionitis.
diandra, Heyne, ex Nees, in Wall. Pl. As. Rar. iii. 92 = polytricha ?
diandra, Hochst. & Steud. ex Nees, in DC. Prod. xi. 231 = Hochstetteri.
diandra, Schlecht. ex Nees, l. c. = obtusa.
dianthera, Herb. Madr. ex Wall. Cat. n. 2391 = Peristrophe montana.
dichotoma, Roxb. Hort. Beng. 45; Fl. Ind. iii. 39 = cristata.
diffusa, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 516 = Lepidagathis diffusa.
discolor, *Nees, in Benth. Bot. Voy. Sulph.* 146.— Mexic.
echinata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = B. Prionitis.
elata, Dals. in Hook. Kew Journ. iii. (1851) 227 = involucrata.
eranthemoides, R. Br. in Salt, Abyss. App. 64 = acanthoides.
fasciculata, Russ. ex Wall. Cat. n. 2500 = polytricha ?

BARLERIA:—
fissiflora, *Boj. ex Nees, in DC. Prod.* xi. 238.— Madag.
flava, *Jacq. Ecl.* 67. t. 46.—Arab.
gentianoides, Desf. ex Nees, in DC. Prod. xi. 224 = flava.
Gibsoni, *Dals. in Hook. Kew Journ.* ii. (1850) 339.— Ind. or.
glabra, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art iv. 8.—Malaya.
glabrata, Vahl, ex Nees, in DC. Prod. xi. 248 = Hygrophila spinosa.
glandulosa, Hochst. ex Nees, l. c. 243 = Lepidagathis glandulosa.
glauca, T. Anders. in Journ. Linn. Soc. ix. (1867) 493 = Gibsoni.
grandiflora, *Dals. in Hook. Kew Journ.* ii. (1850) 339. —Ind. or.
grandiflora, *R. Br. in Salt, Abyss. App.* 64 — Abyssin.
grandifolia, R. Br. ex T. Anders in Journ. Linn Soc. vii. (1864) 30, sphalm. = grandiflora.
grandis, Hochst. ex Nees, in DC. Prod. xi. 233 = grandiflora.
Grantii, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 127. —Afr. trop.
Guerkeana, *Sond. in Linnaea*, xxiii. (1850) 91.—Afr. austr.
halimoides, Nees, in DC. Prod. xi. 231 = Petalidium halimoides.
Hamiltoni, *Solms, in Schweinf. Beitr. Fl. Aethiop.* 106. —Afr. trop.
hexacantha, Bertol. in Ann. Stor. Nat. Bolog. fasc. ix. 440; Hiern, Fl. *Nov.* 3 (1898) = longifolia.
hexacantha, Moris, in Mem. Acc. Torin. xxxvi. (1888) 200. t. 7 = Hygrophila spinosa.
Hildebrandtii, *S. Moore, in Journ. Bot.* xv. (1877) 69. —Afr. trop.
hirsuta, Herb. Madr. ex Wall. Cat. n. 2503 = tomentosa.
hirsuta, Jacq. Obs. Bot. ii. 7 = Gonzalea spicata.
hirsuta, Nees, Wall. Pl. As. Rar. iii. 91; Cat. n. 7142 = polytricha.
Hochstetteri, *Nees, ex DC. Prod.* xi. 231.—Afr. bor.; Arab.; Scind.
hypocrateriformis, Hochst. ex T. Anders. in Journ. Linn. Soc. vii. (1863) 28 = B. Prionitis.
Hystrix, Linn. Mant. i. 89 = B. Prionitis.
ilicina, *E. Mey. ex T. Anders. in Journ. Linn. Soc* vii. (1864) 28.—Afr. austr.
inaequalis, Benth. in Flora, xxxii. (1849) 557 = Phaylopsis longifolia.
inaequalis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 149 = Phaylopsis longifolia.
indica, Linn. ex T. Anders. in Journ Linn. Soc. vii. (1864) 115 = cristata.
involucrata, *Nees, in Wall. Pl. As. Rar.* iii. 92.—Ind. or.
irritans, *Nees, in Linnaea*, xv. (1841) 859.—Afr. austr.
Kirkii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 30. —Afr. trop.
laciniata, Wall. Pl. As. Rar. iii. 91; Cat. n. 7144 = cristata.
lactea, Desf. ex Steud. Nom. ed. II. i. 187 = cristata.
lanciflolia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 28.—Afr. austr. trop.
Lawii, *T. Anders. l. c.* ix. (1867) 492.—Ind. or.
Lichtensteiniana, *Nees, in DC. Prod.* xi. 236.—Afr. austr.
limnophora, *S. Moore, in Journ. Linn. Soc.* xv. (1877) 95.—Ins. Celebes.
longiflora, *Linn. f. Suppl.* 290.—Ind. or.
longifolia, T. Anders. in Journ. Linn. Soc. ix. (1867) 491 = longiflora.
longifolia, Linn. Cent. Pl. ii. 22; Amoen. Acad. iv. 320 = Hygrophila spinosa.
lupulina, *Lindl. in Bot. Reg.* t. 1483.—Ind. or.; Ins. Maurit.
Mackenii, *Hook. f. Bot. Mag.* t. 5866.—Afr. austr.
Macrae, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 311.—Ind. or.
macracantha, R. Br. in Salt, Abyss. App. 64 = Hygrophila spinosa ?
macroloma, *Peyr. in Sitzb. Acad. Wien. Math.-Nat.* xxxvii. (1860) 576.—Afr. trop.
macrophylla, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 489 = strigosa.
macrostachya, Boj. Hort. Maurit. 260 = lupulina.
macrostegia, *Nees, in DC. Prod.* xi. 235.—Afr. austr.

BARLERIA:—
marginata, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 127.—Afr. trop.
Meyeriana, *Nees, in DC. Prod.* xi. 230.—Afr. austr.
micans, *Nees, in Benth. Bot. Voy. Sulph.* 146.— N. Granat.
mitis, *Kor-Gmel. in Bot. Reg.* t. 191.—Ind. or.
mollis, *R. Br. in Salt, Abyss. App.* 64.—Abyssin.
monostachya, *Benj. in Hook. Journ. Bot.* i. (1834) 356. —Ins. Maurit.
montana, *Herb. Madr. ex Nees, in Wall. Pl. As. Rar.* iii. 92.—Ind. or.
montana, Wight & Nees, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 487, pro parte = Gibsoni.
multiflora, *Hassk. Cat. Fl. Bogor. Alt.* 149.—Hab. ?
mysorensis, *Heyne, in Roth, Nov. Sp. Pl.* 313.— Ind. or.
napalensis, Nees, in Wall. Pl. As. Rar. iii. 91 = cristata.
nitida, *Jacq. ex Steud. Nom.* ed. II. i. 187.—Am. austr.
nitida, *Nees, in Wall. Pl. As. Rar.* iii. 91.—Ind. or.
noctiflora, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 145 = acanthoides.
noctiflora, *Linn. f. Suppl.* 290.—Ind. or.; Arab.
nuda, Nees, in Wall. Pl. As. Rar. iii. 92 = cristata.
nutans, *Nees, in DC. Prod.* xi. 227.—Zeylan.
obovata, Buch.-Ham. in Trans. Linn. Soc. xiv. (1825) 298 = mysorensis.
obtusa, *Nees, in Linnaea*, xv. (1841) 858.—Afr. austr.
odontacantha, Dum.-Cours. Bot. Cult. Cat. vii. 7 = flava.
opaca, *Nees, in DC. Prod.* xi. 230.—Afr. occ.
orbicularis, *Hochst. ex Anders, in Journ. Linn. Soc.* vii. (1864) 29.—Abyssin.
ovata, *E. Mey. ex Nees, in DC. Prod.* xi. 230.—Afr. austr.
paniculata, Herb. Madr. ex Nees, in Wall. Pl. As Rar. iii. 92 = tomentosa.
papillosa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 31.—Afr. austr.
parviflora, *R. Br. in Salt, Abyss. App.* 65.—Abyssin.
pentacantha, Arn. ex Nees, in DC. Prod. xi. 232 = involucrata et Arnottiana.
pilosa, *Heyne, ex Nees, in Wall. Pl. As. Rar.* iii. 93.— Ind. or.
polyneura, *S. Moore, in Journ. Bot.* xviii. (1880) 266. —Afr. trop.
polystachya, Hook. ex Nees, in DC. Prod. xi. 226 = strigosa.
polytricha, Boj. ex Nees, l. c. 236 (= *Polythrix stenandrium*).—Madag.
polytricha, *Wall. Pl. As. Rar.* i. 72. t. 82.—Ind. or.
prionoides, Hort. Monsp. ex Nees, in DC. Prod. xi. 229 = cristata.
Prionitis, *Linn. Sp. Pl.* 636.—As. et Afr. trop.
procumbens, *Lour. Fl. Cochinch.* 377.—China.
pubiflora, Benth. in Flora, xxxii. (1849) 558 = B. Prionitis.
pumila, Hochst. ex Nees, in DC. Prod. xi. 372 = Justicia simplex.
pungens, Heyne, ex Nees, in Wall. Pl. As. Rar. iii. 91 = polytricha ?
pungens, *Linn. f. Suppl.* 290.—Afr. austr.
pungens, Thunb. Prod. Pl. Cap. 98 = irritans.
pungens, Wall. Cat. n. 2494 = Lepidagathis pungens.
purpurea, Ledd. Bot. Cab. tr. t. 344 = macrostegia.
pyramidalis, Lam. ex Nees, in DC. Prod. 248 = Blechum Brownei.
quadriplexa, Stokes, Bot. Mat. Med. iii. 411 = B. Prionitis.
querimbensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 205.—Afr. trop.
racemosa, Herb. Madr. ex Wall. Cat. n. 2508 = tomentosa.
ramosa, Buch.-Ham. in Trans. Linn. Soc. xiv. (1825) 298 = noctiflora.
repens, *Nees, in DC. Prod.* xi. 230.—Afr. austr.
rhynchocarpa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 206.—Afr. trop.
Richardiana, *Nees, in DC. Prod.* xi. 235.—Madag.
rigida, Sprang. in Zeyh. ex Schlecht. in Linnaea, xiv. (1840) 204 = Pachypodium tomentosum.
rigida, *Willd. ex Nees, in DC. Prod.* xi. 242.— Afr. austr.
rubra, *Buch.-Ham. in Trans. Linn. Soc.* xiv. (1825) 296 (sp. dub.).—Ind. or.
ruellioides, Retzl. ex Nees, in DC. Prod. xi. 240 = noctiflora.

BARLERIA :—
ruellioides, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 30.—Afr. trop.
salicifolia, *S. Moore, in Journ. Bot.* xviii. (1880) 268. —Afr. trop.
saturnioides, *Griseb. Cat. Pl. Cub.* 195.—Cuba.
senegalensis, Nees, in DC. Prod. xi. 224 = flava.
serensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 209.—Afr. trop.
sepalosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 457.—Ind. or.
sericea, T. Anders. in Journ. Linn. Soc. vii. (1864) 115 = longiflora.
solanifolia, Linn. Sp. Pl. 637 = Barleriola solanifolia.
somalensis, *Franch. in Révoil, Somali*, 51.—Afr. or.
spicata, Roxb. Hort. Beng. 45 = B. Prionitis.
Spina-cylanica, Nees, in DC. Prod. xi. 242 = mysorensis.
spinicyma, *Nees, l. c.* 240.—Arabia.
spinosa, Hook. ex Nees, l. c. 247 = Hygrophila spinosa.
spinosa, Wall. Cat. n. 2497 = Lepidagathis spinosa.
spinulosa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 208.—Afr. trop.
squarrosa. *Klotzsch, l. c.* 207.—Afr. trop.
stellato-tomentosa, *S. Moore, in Journ. Bot.* xviii. (1880) 268.—Afr. trop.
stenophylla, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) 11. 95.—Burma.
stimulans, *E. Mey. ex Nees, in DC. Prod.* xi. 241.— Afr. austr.
Stocksii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 493.—Ind. or.
strigosa, *Willd. Sp. Pl.* iii. 379.—Ind. or.
terminalis, Nees, in DC. Prod. xi. 225 = strigosa.
tetracantha, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 85.—Ins. Socotr.
tomentosa, *Roth, Nov. Sp. Pl.* 314.—Ind. or.
tomentosa, Herb. Madr. ex Wall. Cat. n. 2503 = acuminata.
triacantha, Hochst. ex Nees, in DC. Prod. xi. 240 = acanthoides.
trinervia, Herb. Ham. ex Nees, l. c. 248 = Lepidagathis trinervis.
trispinosa, *Vahl, Symb. Bot.* i. 46.—Arab.
uiteragensis, Hochst. ex Nees, in DC. Prod. xi. 231 = obtusa.
unilateralis, Buch.-Ham. ex Wall. Cat. n. 2509 = polytricha.
ventricosa, *Hochst. ex Nees, in DC. Prod.* xi. 230.—Abyss.
wenulosa, Nees, l. c. 238 = cristata.
vestita, *T. Anders. in Thw. Enum. Pl. Zeyl.* 220.— Zeylan.
villosa, *S. Moore, in Journ. Bot.* xviii. (1880) 267.— Afr. trop.
violascens, *S. Moore, l. c.* 265.—Afr. trop.
Wolwitschii, *S. Moore, l. c.* 267.—Afr. trop.

BARLERIACANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136 = **Barleria**, Linn. (Acanth.).
acanthoides, Oerst. l. c. = Barleria acanthoides.
noctiflora, Oerst. l. c. = Barleria noctiflora.

BARLERIANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136 = **Barleria**, Linn. (Acanth.).
grandiflora, Oerst. l. c. = Barleria grandiflora.

BARLERIOLA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136. *ACANTHACEÆ*, Benth. & Hook. f ii. 1098.
solanifolia, *Oerst. l. c.*—Ins. S. Doming.

BARLERIOPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 133 = **Barleria**, Linn. (Acanth.).
discolor, Oerst. l. c. 134 = Barleria discolor.
glandulosa, Oerst. l. c. = Barleria micans.
micans, Oerst. l. c. = Barleria micans.

BARLERIOSIPHON, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136 = **Barleria**, Linn. (Acanth.).
longiflora, Oerst. l. c. = Barleria longiflora.

BARLERITES, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 137 = **Barleria**, Linn. (Acanth.).
Hochstetteri, Oerst. l. c. = Barleria Hochstetteri.

BARLIA, Parl. Nouv. Gen. Plant. Monocot. 5 (1858); Fl. Ital. iii. 445 (1858) = **Orchis**, Linn.
longibracteata, Parl. l. c. 6; Fl. Ital. iii. 447 = O. longibracteata.

BARNADESIA, Mutis, in Linn. f. Suppl. 55 (1781). *COMPOSITÆ*, Benth. & Hook. f. ii. 484.
BACAZIA, Ruiz & Pav. Prod. Fl. Per. 105. t. 22 (1794).
DIACANTHA, Less. in Linnaea, v. (1830) 243.
DOLICHOFFLYS, Cass. in Dict. Sc. Nat. lvi. 139 (1828).
FULCALDEA, Poir. Encyc. Suppl. v. 875 (1817).
PENTHEA, Spach, Hist. Vég. Phan. x. 9 (1841).
RHODACTINIA, Gardn. in Hook. Lond. Journ. vi. (1847) 449.
TURPINIA, Humb. & Bonpl. Pl. Aequin. i. 113. t. 38 (1808).
VOIGTIA, Spreng. Syst. iii. 367 (1826).
XENOPHONTA, Vell. Fl. Flum. 346 (1825).
arbores, *Sch. Bip. in* Linnaea, xxii. (1849) 569 = rosea.
berberoides, *Sch. Bip. in Lechl. Berb. Am. Austr.* 57. —Peruv.
corymbosa, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 278.—Peruv.
dianthiflora, Mart. ex Baker, in Mart. Fl. Bras vi. iii. 365 = rosea.
divaricata, *Griseb. in Gott. Abh.* xxiv. (1879) 209.— Reg. Argent.
Dombeyana, *Less. in* Linnaea, v. (1830) 246.—Peruv.
flavescens, *Meyen, Reise*, i 307.—Chili.
glochidiata, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 361 = syncantha.
hirsuta, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 527, nomen.—Bolivia.
lanata, *Meyen, Reise*, i. 347.—Chili.
lancolata, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 277.—Peruv.
laxa, D. Don, l. c. 280 = rosea.
media, *D. Don, l. c.* 276.—Peruv.
odorata, *Griseb. in Gott. Abh.* xxiv. (1879) 210.— Reg. Argent.
parviflora, *Spruce, ex Benth. & Hook. f. Gen.* ii. 485. —Ecuador.
polyscantha, *Wedd. Chlor. And.* i. 13.—Bolivia.
retinitis, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 277.—Peruv.
rosea, *Lindl. in Bot. Reg.* (1843) t. 39.—Am. austr.
spinosa, *Linn. f. Suppl.* 348.—Peruv.: N. Granat.
spinosa, Willd. ex DC. Prod. vii. 3 = arborea.
ulicina, Hook. & Arn. Bot. Beech. Voy. 29 = Chuquiraga ulicina.

BARNARDIA, Lindl. in Bot. Reg. t. 1029 (1826) = **Scilla**, Linn. (Liliac.).
indica, Wight, Ic. t. 2041 = S. indica.
japonica, Schult. f. Syst. vii. 555 = S. japonica.
scilloides, Lindl. in Bot. Reg. t. 1029 = S. chinensis.

BARNEOUDIA, C. Gay, Fl. Chil. i. 29. t. 1 (1845) = **Anemone**, Linn. (Ranunc.).
chilensis, C. Gay, l. c.—Chili.
Domeykoana, Leybold, in Anal. Univ. Chil. (1858) 159; et in Flora, xlii. (1859) 242.—Chili.
major, Phil. in Linnaea, xxviii. (1856) 609.—Chili.

BAROLA, Adans. Fam. ii. 344 (1763) = **Trichilia**, Linn. (Meliac.).

BAROLLAEA, Neck. Elem. ii. 322 (1790) = **Caryocar**, Linn. (Ternstroem.).

BAROMIA, Baker, in Journ. Bot. xx. (1882) 67. *NARCISSIDACEÆ*.
Tazetana, *Baker, l. c.*—Madag.

BAROSMA, Willd. Enum. Hort. Berol. 257 (1809). *RUTACEÆ*, Benth. & Hook. f. i. 290.
BARYOSMA, Roem. & Schult. Syst. v. 448 (1819).
BUCCO, Wendl. Collect. i. 18 (1808).
PHAMPHALIPSEA, Wendl. Collect. i. 15, 84 (1808).
acutata, *Sond. in Harv. & Sond. Fl. Cap.* i. 395.—Afr. austr.
alpina, *Eckl. & Zeyh. Enum.* 109.—Afr. austr.
angustifolia, *Bartl. & Wendl. f. Diosm.* 116.—Afr. austr.
apiculata, *Eckl. & Zeyh. Enum.* 104.—Afr. austr.
betulina, *Bartl. & Wendl. f. Diosm.* 103.—Afr. austr.
brunioides, Hoffmgg. Verz. Pfl. Nachtr. i. 128 = Agathosma cuspidata.
crenata, *Eckl. & Zeyh. Enum.* 103.—Afr. austr.

BAROSMA :—
crenata, *Sweet, Hort. Brit.* ed. I. 89.—Afr. austr.
crenulata, *Hook. Bot. Mag.* t. 3413.—Afr. austr.
dioica, *Bartl. & Wendl. f. Diosm.* 114.—Afr. austr.
Eckloniana, *Bartl. in Linnaea*, xvii. (1843) 363.—Afr. austr.
foetidissima, *Bartl. & Wendl. f. Diosm.* 118.—Afr austr.
gnidioides, *Eckl. & Zeyh. Enum.* 104.—Afr. austr.
graveolens, *G. Don, Gen. Syst.* i. 786.—Afr. austr
Kraussiana, *Meisn. in Flora*, xxvii. (1844) i. 303.— Afr. austr.
lanceolata, *Sond. in Harv. & Sond. Fl. Cap.* i. 397.— Afr. austr.
latifolia, *Roem. & Schult. Syst.* v. 449.—Afr. austr.
microcarpa, *Sond. in Harv. & Sond. Fl. Cap.* i. 398 — Afr. austr.
mucronata, *Meisn. in Flora*, xxvii. (1844) i. 303.— Afr. austr.
Niveni, *Sond. in Harv. & Sond. Fl. Cap.* i. 396.—Afr austr.
oblonga, *Bartl. & Wendl. f. Diosm.* 112.—Afr. austr
odorata, Willd. Enum. Hort. Berol. 257 = serratifolia.
orbicularis, Hort. ex Sweet, Hort. Brit. ed. I. 89 = betulina.
ovata, *Bartl. & Wendl. f. Diosm.* 109.—Afr. austr.
pauciflora, *Eckl. & Zeyh. Enum.* 103.—Afr. austr.
pulchella, *Bartl. & Wendl. f. Diosm.* 107.—Afr. austr.
pulchra, *Cham. in Linnaea*, v. (1830) 53.—Afr. austr.
pungens, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 397.— Afr. austr.
scoparia, *Eckl. & Zeyh. Enum.* 103.—Afr. austr.
serratifolia, *Willd. Enum. Hort. Berol.* 257.—Afr austr.
sertuliflora, *Delponte, in Mem. Acc. Torino*, Ser. II xiv. (1854) 408. t. 9.—Hab. ?
ternata, *Eckl. & Zeyh. Enum.* 108.—Afr. austr.
trichopodia, *Bartl. & Wendl. f. ex Juss. in Mém. Mus. Par.* xii. (1825) 474.—Afr. austr.
venusta, *Eckl. & Zeyh. Enum.* 102.—Afr. austr.

BARRALDEIA, Thou. Gen. Nov. Madag. 24 (1809) = **Carallia**, Roxb. (Rhizoph.).
madagascariensis, Spreng. Syst. ii. 318 (Baraultia).— Madag.

BARRATTIA, A. Gray, in Proc. Am. Acad. i. (1848) 48 = BARATTIA, A. Gray & Engelm. = **Emoclia**, Adans. (Compos.).

BARRELIERA, J. F. Gmel. Syst. 959 (1791) = **Barleria**, Linn. (Acanth.).

BARRERIA, Linn. Gen. ed. II. 131 (1742); (et BARRÈRA, Linn. Sp. Pl. ed. II. 893 (1762)). *asperata*, Linn. Sp. Pl. 874.—Afr. austr.

BARRERIA, Willd. Sp. Pl. i. 1145 (1798) = **Poraqueiba**, Aubl. (Olacinese).
guianensis, Racusch. Nom. ed. III. 259, nomen = P. guianensis.
theobromaefolia, Willd. Sp. Pl. i. 1145 = P. guianensis.

BARRINGTONIA, Forst. Char. Gen. 75. t. 38 (1776). *MYRTACEÆ*, Benth. & Hook. f. i. 720.
AGASTA, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. 59 (1875).
BARANDA, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 502.
BOTRYOROPIS, Presl, Epim. Bot. 220 (1849).
BUTONICA, Juss. Gen. 326 (1789).
DOXOMMA, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. (1875) 98.
MEGADENDRON, Miers, l. c. 109 (1875).
MENICHIRA, Sonner. Voy. N. Guin. 138 (1776).
METEORUS, Lour. Fl. Cochinch. 410 (1790).
MITRARIA, J. F. Gmel. Syst. 799 (1791).
STRAVADIUM, Juss. Gen. 326 (1789).
acuminata, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 206.—Borneo.
acutangula, *Gaertn. Fruct.* ii. 97. t. 101.—Ins. Moluce.
alata, Wall. Cat. n. 3833 = conoidea.
alba, *Blume, in Fl. des Serres*, Sér. I. vii. (1851-52) 28.—Ins. Moluce.
alba, Kostel. Allg. Med.-Pharm. Fl. iv. 1536 = racemosa
asiatica, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 181; xlvi. (1877) ii. 70 = speciosa.
angusta, *Kurz, l. c.* xlii. (1873) ii. 233.—Burma.
Butonica, *Forst. Char. Gen.* t. 38.—Ins. Pacif.

BARRINGTONIA :—
calyptrata. R. Br. ex Benth. Fl. Austral. iii. 388, in note = species ?
Careya, *F. Muell. Fragm.* v. 182.—Austral.
ceylanica, *Garän. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 508.—Ceylan.
cerviana, Kostel. Allg. Med.-Pharm. Fl. iv. 1586 = excelsa.
conoidea, *Griff. Notul.* iv. 656 ; *Ic. Pl. As.* t. 635, 636. f. 1.—Burma ; Malacca.
costata, *Miq. Fl. Ind. Bat.* i. 489.—Archip. Sunda.
cylindrostachya, Griff. Notul. iv. 655 = macrostachya.
edulis, *Seem. Fl. Vit.* 82.—Ins. Fiji.
elongata, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 206. —Java.
excelsa, *Blume, Bijdr.* 1097.—Malaya.
Helferi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 509. —Burma.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. 489.—Java.
incisa, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. (1876) 71. t. 14 = racemosa.
insignis, *Miq. Fl. Ind. Bat.* i. 488.—Malaya.
intermedia, *Vieill. in Bull. Soc. Linn. Normand.* x. (1886) 88.—N. Caled.
lasseniensis, *Vidal, Phan. Cuming. Philipp.* 13, 48, 118. —Ins. Philipp.
macrocarpa, *Hazsk. Cat. Hort. Bogor. Alt.* 268.—Java.
macrophylla, Miq. Fl. Ind. Bat. i. 491 = speciosa.
macrostachya, *Kurz, For. Fl. Brit. Burma*, i. 498 ; *et in Journ. As. Soc. Beng.* xlvi. (1877) 11. 71.—Burma.
Montrouzieri, *Vieill. in Bull. Soc. Linn. Normand.* x. (1886) 100.—N. Caled.
neocaledonica, *Vieill. l. c.* 99.—N. Caled.
nitida, *Miq. Fl. Ind. Bat.* i. 490.—Java.
pendula, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) 11. 71.—Burma.
pterocarpa, *Kurz, l. c.* xlii. (1873) 11. 284.—Burma.
pubescens, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. (1876) 83 = acutangula.
racemosa, *Roxb. Hort. Beng.* 52 ; *Fl. Ind.* ii. 684.— Ind. or. ; Malaya ; Polynes.
Retzwaittii, *Miq. Fl. Ind. Bat.* i. 488.—Java.
reticulata, *Miq. l. c.* 490.—Borneo.
rigida, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 510 = Symplocos rigida.
rubra, Blume, in Fl. des Serres, Sér. I. vii. (1851–52) 23 = racemosa.
samoënsis, *A. Gray, Bot. U. St. Expl. Exped.* i. 508.— Ins. Fiji.
sarcostachys, Miq. Fl. Ind. Bat. i. 490 = macrostachya.
Senegali, *Yardin, in Bull. Soc. Linn. Normand.* Sér. II. ix. (1875) 803.—N. Caled.
serrata, *Miq. Fl. Ind. Bat.* i. 488.—Java.
speciosa, Linn. f. Suppl. 212 = B. Butonica.
speciosa, Wall. Cat. n. 3633 B = racemosa.
spicata, *Blume, Bijdr.* 1097.—Java.
Strongdium, Blanco, Fl. Filip. ed. I. 558 = racemosa.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 313.—Sumatra.
timorensis, *Blume, in Ver. Houtte, Fl. des Serres*, Sér. I. vii. (1851–52) 23.—Ins. Timor.
Vriesei, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* ii. (1855) 308.—Java.

BARROËTIA. A. Gray, in Proc. Am. Acad. xv. (1880) 99. *COMPOSITÆ.*
Pavonii, *A. Gray, l. c.* xvii. (1881–82) 206.—Mexic.
setosa, *A. Gray, l. c.* xv. (1880) 29.—Mexic.
subuligera, *A. Gray, l. c.*—Mexic.

BARROTIA, Gaudich. Bot. Voy. Bonite, t. 13 (1852?) = Pandanus, Linn.
altissima, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 277 = F. Minde.
aragoënsis, Brongn. l. c. 278 = P. aragoënsis.
Balansæ, Brongn. l. c. 281 = P. Balansæ.
decumbens, Brongn. l. c. 285 = P. decumbens.
dicdon, Gaudich. Bot. Voy. Bonite, t. 13. f. 9–14 = P. furcatus.
Gaudichaudii, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 264 = P. tetrodon.
macrocarpa, Brongn. l. c. 279 = P. macrocarpus.
monadon, Gaudich. Bot. Voy. Bonite, t. 13. f. 15–24 = P. monadon.
Pancheri, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 288 = P. Pancheri.
sphærocephala, Brongn. l. c. 284 = P. sphærocephalus.
tetrodon, Gaudich. Bot. Voy. Bonite, t. 13. f. 1–8 = P. tetrodon.

BARROWIA, Decne. in DC. Prod. viii. 629 (1844). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 779.
Jasminiflora, *Decne. l. c.* 630.—Afr. austr.

BARTERIA, Hook. f. in Journ. Linn. Soc. v. (1861) 14. t. 3. *PASSIFLOREÆ*, Benth. & Hook. f. i. 813.
fistulosa, *Mast. in Oliver, Fl. Trop. Afr.* ii. 511.—Afr. trop.
nigritana, *Hook. f. in Journ. Linn. Soc.* v. (1861) 15. t. 3.—Afr. trop.

BARTHEA, Hook. f. in Benth. & Hook. f. Gen. i. 751 (1867). *MELASTOMACEÆ*, Benth. & Hook. f. i. c.
chinensis, *Hook. f. l. c.*—Hongkong.

BARTHESIA, Comm. ex A. DC. in Trans. Linn Soc. xvii. (1837) 112 = **Ardisia**, Sw. (Myrsin.).

BARTHOLINA, R. Br. in Ait. Hort. Kew. ed. II. v. 194 (1813). *ORCHIDEÆ*, Benth. & Hook. f. iii. 623.
LATHRISIA, Sw. Annot. Bot. 48 (1829).
Burmanniana, Ker-Gawl. in Brand. Journ. Sc. iv. (1818) 204. t. 5 = pectinata.
Ethelae, *Balus, in Journ. Linn. Soc.* xx. (1884) 472.— Afr. austr.
Lindleyana, Reichb. f. Otia Bot. Hamb. 119 = pectinata.
pectinata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 194.— Afr. austr.

BARTHOLLESIA, Manso, Enum. 45 (1836) = **Bertholletia**, Humb. & Bonpl. (Myrtac.).

BARTHRATHERUM, Anderss. in Walp. Ann. iii. 799 (1858) = BATHRATHERUM, Hochst. = **Arthraxon**, Beauv. (Gramin.).

BARTLETTIA, A. Gray, in Mem. Am. Acad. Ser. II. v. (1855) (Pl. Nov. Thurb.) 333. *COMPOSITÆ*, Benth. & Hook. f. ii. 246.
scaposa, *A. Gray, l. c.*—Mexic.

BARTLINGIA, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 373 = **Pultenæa**, Sm. (Legumin.).
obovata, Brongn. l. c. = P. obovata.

BARTLINGIA, F. Muell. ex Benth. Fl. Austral. vii. 63 (1878) = **Laxmannia**, R. Br. (Liliac.).
brachyphylla, F. Muell. Census, 119 = L. brachyphylla.
gracilis, F. Muell. Fragm. vii. 88 = L. gracilis.
grandiflora, F. Muell. Census, 118 = L. grandiflora.
minor, F. Muell. l. c. = L. ramosa.
ramosa, F. Muell. Fragm. vii. 88 = L. ramosa.
sessiliflora, F. Muell. in Papers Roy. Soc. Tasm. (1877) 116 = L. sessiliflora.
sessilis, F. Muell. Census, 119 = L. sessilis.
squarrosa, F. Muell. l. c. 118 = L. squarrosa.

BARTLINGIA, Reichb. in Flora, vii. (1824) 241 = **Plocama**, Ait. (Rubiac.).
scoparia, Reichb. l. c. = P. pendula.

BARTOLINA, Adans. Fam. ii. 124 (1763) = **Tridax**, Linn. (Compos.).

BARTSIA, Mühl. ex Willd. in Ges. Naturf. Freunde, Neue Schr. iii. (1801) 444. *GENTIANEÆ*, Benth. & Hook. f. ii. 813.
ANTHIA, Spreng. Syst. i. 428 (1825).
CENTAURELLA, Michx. Fl. Bor. Am. i. 97. t. 13 (1803).
CENTAURIUM, Pers. Syn. i. 187 (1805).
NUTTALLIA, Rafin. in Am. Monthly Mag. (1818) 175.
pubescens, Rafin. in Med. Repos. New York, v. (1808) 353 = verna ?
tenella, *Muhl. ex Willd. in Ges. Naturf. Freunde, Neue Schr.* iii. (1801) 445.—Am. bor.
uniflora, Rafin. in Med. Repos. New York, v. (1808) 353 = verna ?
verna, *Muhl. ex A. Gray, in Chapm. Fl. S. U. St.* 357. —Am. bor.

BARTONIA, Sims, Bot. Mag. t. 1487 (1804) = **Mentzelia**, Linn. (Loasac.).
albescens, Gill. & Arn. in Edinb. Geog. Journ. ii. (1831) 273.—Chili.
albicaulis, Dougl. ex Hook. Fl. Bor. Am. i. 222 = M. albicaulis.
aurea, Lindl. in Bot. Reg. t. 1691 = M. Lindleyi.

BARTONIA :—
decapetala, Sims, Bot. Mag. t. 1457 = Mentzelia ornata.
lasseniensis, Dougl. ex Hook. Fl. Bor. Am. i. 221. t. 69 = M. laevicaulis.
micrantha, Hook. & Arn. Bot. Beech. Voy. 343 = M. micrantha.
micrantha, Hook. in Lond. Journ. Bot. vi. (1847) 237 = M. dispersa.
multiflora, Nutt. in Journ. Acad. Sc. Phil. N. S. i. (1847) 130 = M. multiflora.
nuda, Pursh, Fl. Am. Sept. 328, 749 = M. nuda.
ornata, Hook. Lond. Journ. Bot. vi. (1847) 236 = M. laevicaulis.
ornata, Pursh, Fl. Am. Sept. 327 = M. ornata.
parviflora, Dougl. ex Hook. Fl. Bor. Am. i. 221 = M. laevicaulis.
pumila, Nutt. ex Torr. & Gray, Fl. N. Am. i. 505 = M. pumila.
sinuata, Presl, Rel. Haenk. ii. 33 = *albescens*.
Wrightii, Walp. Ann. ii. 656 = M. Wrightii.

BARTRAMIA, Linn. Dias. Dasa. 17 (1747) ; Fl. Zeyl. 77 (1747) ; Gaertn. Fruct. ii. 137. t. 111 (1791) = **Triumfetta**, Linn. (Tiliac.).
angulata, Lam. Ill. t. 430. f. 2 = T. rhomboidea.
crispifolia, Stokes, Bot. Mat. Med. iii. 14 = T. rhomboidea.
glandulosa, Lam. Ill. t. 400. f. 1 = T. rhomboidea.
indica, Linn. Sp. Pl. 889 = T. rhomboidea.
Lappago, Gaertn. Fruct. ii. 137. t. 111 = T. rhomboidea.
rhomboïdea, Stokes, Bot. Mat. Med. iii. 15 = T. rhomboidea.

BARTRAMIA, Salisb. Prod. 99 (1796) = **Penstemon**, Mitch. (Scrophular.).
concinna, Salisb. l. c. = P. pubescens.
pulchella, Salisb. l. c. = P. laevigata.

BARTSCHIA, Linn. Syst. ed. II. 26 (1740) = seq.

BARTSIA, Linn. Hort. Cliff. 325 (1737). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 977.
ALICOTA, Phil. Enum. Fl. Hautes-Pyr. 381 (1867).
BELLARDIA, All. Fl. Pedem. i. 61 (1785).
EUFRAGIA, Griseb. Spicil. Fl. Rumel. ii. 13 (1844).
LAGIOPERA, Hoffmgg. & Link, Fl. Port. i. 298 (1809).
ODONTITES, [Riv.] Hall. Enum. Stirp. Helv. ii. 627 (1742).
PARENTUCELLIA, Viv. Fl. Lib. Spec. 31. t. 21. f. 3 (1824).
STARHELINIA, Hall. Enum. Stirp. Helv. ii. 624 (1742).
TRIXAGO, Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 4.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 184 = B. Trixago.
acuminata, Pursh, Fl. Am. Sept. ii. 439 = Castilleja pallida.
alpina, *Linn. Sp. Pl.* 602.—Europ. ; Am. bor.
aspera, *Willd. & Lange, Fl. Hisp.* ii. 614, obs.—Hispan. ; Marocc.
bicolor, DC. Ic. Rar. t. 10 = B. Trixago.
blioba, *Wedd. Chlor. And.* ii. 123.—Peruv.
brevidora, *Benth. in DC. Prod.* x. 545.—Calif.
canescens, *Wedd. Chlor. And.* ii. 123.—Peruv.
capensis, Spreng. Syst. ii. 778 = B. Trixago.
carnea, *Griseb. Spicil. Fl. Rumel.* ii. 12.—Macedon.
chilensis, *Benth. in DC. Prod.* x. 547.—Chili.
cilleloides, *Wedd. Chlor. And.* ii. 123.—Bolivia.
coccinea, Linn. Sp. Pl. 602 = Castilleja coccinea.
crenoloba, *Wedd. Chlor. And.* ii. 124.—Bolivia.
decurva, *Hochst. ex Benth. in DC. Prod.* x. 545.— Abyssin.
densiflora, *Benth. l. c.* 546.—Peruv.
diffusa, *Benth. l. c.* 546.—Peruv.
elongata, *Wedd. Chlor. And.* ii. 127.—Am. merid.
euphrasioides, *Wedd. l. c.* 130.—Ecuador.
Figueiri, Lapeyr. Hist. Abr. Pl. Pyr. 545 = aspera.
filicoides, *Wedd. Chlor. And.* ii. 126.—Peruv.
Garidelii, Bell, in Journ. Linn. Soc. xvi. (1878) 602 = purpurea.
glauca, Pair. ex Steud. Nom. ed. II. i. 689 = Lagotis glauca.
gracilis, *Benth. in DC. Prod.* x. 548.—N. Granat.
granatensis, Bell, in Journ. Linn. Soc. xvi. (1878) 602 = purpurea.
grandiflora, Spreng. in Roch & Grah. Allgem. Encyc. vii. (1821) 491 = Castilleja scabiflora.
Gymnandra, Linn. f. Suppl. 270 = Lagotis glauca.
Gymnandra, Pursh, Fl. Am. Sept. ii. 430 = Synthyris rubra ?

BARTSIA :—
heterophylla, *Wedd. Chlor. And.* ii. 139.—N. Granat.
hispida, *Benth. in DC. Prod.* x. 547.—Peruv.
humilis, Lapeyr. Hist. Abr. Pl. Pyr. 344 = Euphrasia minima.
imbricata, Lapeyr. l. c. = Euphrasia officinalis.
inaequalis, *Benth. in DC. Prod.* x. 547.—Peruv. ; N. Granat.
integrifolia, *Wedd. Chlor. And.* ii. 131.—Peruv.
lanceolata, *Reichb. f. ic. Pl. Germ.* xx. 57. t. 107.—Europ.
lanifloru, *Benth. in DC. Prod.* x. 546.—N. Granat.
laricornata, Benth. l. c. 547 = santolinaefolia.
lattifolia, *Sibth. & Sm. Fl. Graec.* vi. 69. t. 586.—Reg. Mediterr. ; Reg. Casc. ; Persia.
laxiflora, *Benth. in DC. Prod.* x. 547.—N. Granat.
longiflora, *Hochst. ex Benth. l. c.* 545.—Abyss.
lutea, *Reichb. f. ic. Fl. Germ.* xx. 56.—Europ. ; Aa. Min.
marina, Gren. & Godr. Fl. Fr. ii. 611, in syn. = viscosa.
maritima, Pers. Syn. ii. 151 = B. Trixago.
melampyroides, *Benth. in DC. Prod.* x. 548.—Ecuador.
Meyniana, *Benth. l. c.* 546.—Peruv.
mollis, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. lviii. (1888). t. 397 = Veronica ciliata.
motica, *Benth. l. c.* 545.—Peruv.
Odontites, *Huds. Fl. Angl.* 268.—Europ. ; Reg. Mediterr.
orthocarpiflora, *Benth. in DC. Prod.* x. 545.—Peruv.
pallida, Linn. Sp. Pl. 602 = Castilleja pallida.
parviflora, Charpent. ex Benth. in DC. Prod. x. 544 = alpina.
parviflora, Thom. Cat. Pl. Suisses (1887) 6 = alpina.
parvifolia, *Benth in DC. Prod.* x. 545.—N. Granat.
patens, *Benth. l. c.* 546.—Peruv.
pedicularoides, *Benth. l. c.*—Ecuador.
peruviana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 400.—Peruv.
pumila, *Benth. in DC. Prod.* x. 546.—Peruv.
purpurea, *Ball, in Journ. Linn. Soc.* xvi. (1878) 602. —Hispan. ; Afr. bor.
purpurea, Dahy, Bot. Gall. 354 = lattifolia.
rhinanthoides, Hochst. ex Benth. in DC. Prod. x. 548 = B. Trixago.
rubricosa, Fall. Relse, iii. 84 = Castilleja pallida.
santolinaefolia, *Benth. in DC. Prod.* x. 547.—Mexic.
scabra, *Sprang. Syst.* ii. 778. = inaestr.
scordifolia, *Steud. Nom.* ed. II. i. 189.—Ins. Mader.
serotina, Bertol. Amoen. Ital. 83 = B. Odontites.
sibirica, Pall. Relse, ii. 123 = Castilleja pallida.
spicata, Ramond, in Bull. Soc. Philom. ii. (1799–1801) = aspera.
stricta, *Benth. in DC. Prod.* x. 547.—Ecuador.
subinclusa, *Benth. l. c.*—Peruv.
subinclusa, Griseb. ex Wedd. Chlor. And. ii. 126 = filiformis.
tenuifolia, Pursh, Fl. Am. Sept. ii. 439 = Orthocarpus tenuifolius.
trichophylla, *Wedd. Chlor. And.* ii. 132.—Peruv.
trifida, Sprang. Syst. ii. 772 = Bungea trifida.
Trixago, *Linn. Sp. Pl.* 602.—Europ. ; Afr. bor.
verna, *Reichb. f. ic. Fl. Germ.* xx. 57. t. 107.—Europ.
versicolor, Pers. Syn. ii. 151 = B. Trixago.
viscosa, *Linn. Sp. Pl.* 603.—Europ. ; Reg. Mediterr.

BARYA, Klotzsch, in Monatsb. Berl. Acad. (Marz 1854) 198 ; et in Abh. Akad. Berl. (1854) 22. t. 2, B = Begonia, Linn.
monadelpha, Klotzsch, ll. cc. 198, 23 = Begonia monadelpha.

BARYOSMA, Gaertn. Fruct. ii. 78. t. 98 (1791) = Dt. pteryx, Schreb. (Leguminos.).
oppositifolium, Pers. Syn. ii. 272 = D. oppositifolia.
Tonga, Gaertn. Fruct. ii. t. 98 = D. odorata.

BARYOSMA, Reem. & Schult. Syst. v. 448 (1819) = Bauconsa, Willd. (Rutac.).

BARYOSMA, Bunge, Del. Sem. Hort. Dorpat. (1839) 7 ; et in Linnaea, xiv. (1840) Lit. 117 = Haraeleum, Lisa. (Umbellif.).
villosum, Bunge, ll. cc. = H. villosum.

BARYXYLUM, Lour. Fl. Cochinch. 266 (1790). *LEGUMINOSAE* ?—Cf. Benth. & Hook. f. i. 464.
rubrum, *Lour. l. c.*—China.

BASALTOGETON, Salisb. Gen. Pl. Fragm. 27 (1866) = Scilla, Linn. (Liliac.).
peruvianum, Salisb. l. c. = S. peruviana.

BASANACANTHA, Hook. f. in Benth. & Hook. f. Gen. ii. 82 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
armata, *Hook. f. l. c.* 83.—Am. trop.
dioica, *Hook. f. l. c.*—N. Granat.
helocarpa, *Hook. f. l. c.*—Am. trop.
Humboldtiana, *Hook. f. l. c.*—Am. trop.
monantha, *Hook. f. l. c.*—Guatemala.
reticulata, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 98.—Mexic.
tetracantha, *Hook. f. in Benth. & Hook. f. Gen.* ii. 83. —Am. trop.

BASANANTHE, Peyr. in Bot. Zeit. xvii. (1859) 101 ; et in Sitzb. Acad. Wien. Math. Nat. xxxvii. (1860) 589. *PASSIFLOREAE*, Benth. & Hook. f. i. 812.
littoralis, Peyr. ll. cc.—Afr. trop.
Nummularia, *Wawr. in Trans. Linn. Soc.* xxvii. (1869) 28. t. 9.—Angola.

BASELA, Linn. Gen. ed. VI. (Vindob. reimpr.) 151 (1767) = seq.

BASELLA, [Rheede] Linn. Diss. Dass. 12 (1747) ; Amoen. Acad. i. 396. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 76.
GANDOLA, Rumph. ex Linn. Sp. Pl. ed. II. 890 (1763).
alba, Linn. Sp. Pl. 272 = rubra.
canariensis, Linn. Bach.-Hem. ex Wall. Cat. n. 6961 = rubra.
cordifolia, Lam. Encyc. i. 389 = rubra.
crassifolia, Salisb. Prod. 158 = rubra.
diffusa, Ruiz & Pav. ex Moq. in DC. Prod. xiii. ii. 226 (= *Tandonia diffusa*).—Peruv.
filiformis, Ruiz & Pav. ex Moq. l. c. 227 (= *Tandonia filiformis*).—Peruv.
Hookeriana, Moq. ex Moq. l. c. 225 = Tournonia Hookeriana.
japonica, Burm. f. Fl. Ind. 76. t. 39. f. 4 = rubra.
lucida, Linn. Syst. ed. X. 965. n. 548 = rubra.
marginata, H. B. & K. Nov. Gen. et Sp. ii. 189 (= *Tandonia marginata*).—Ecuador.
nigra, Lour. Fl. Cochinch. i. 158 = rubra.
obovata, H. B. & K. Nov. Gen. et Sp. ii. 189 (= *Tandonia obovata*).—N. Granat.
obtusifolia, Willd. ex Moq. in DC. Prod. xiii. ii. 227 (= *Tandonia obovata*).—N. Granat.
rosacea, Jacq. f. in Sprang. Syst. i. 960 = rubra.
rubra, *Linn. Sp. Pl.* 272.—Reg. trop.
trifida, Willd. in Roem. & Schult. Syst. vi. 695 (= *Tandonia marginata*).—Ecuador.
tuberosa, H. B. & K. Nov. Gen. et Sp. ii. 189 = Ullucus tuberosus.
vericaria, Lam. Illustr. t. 215. f. 1 = Anredera spicata.
volubilis, Salisb. Prod. 158 = rubra.

BASILAEA, Juss. ex Lam. Encyc. i. 383 (1783) ; et Gen. 52 (1789) = Rhizonia, *L'Hérit.* (1788) (Liliac.).
bifolia, Mirb. Hist. Nat. Pl. vii. (1804) 339 = Whitehaedia bifolia.
crocea, Lam. Encyc. i. 383 = E. regia.
nana, Mirb. Hist. Nat. Pl. vii. (1804) 339 = E. nana.
punctata, Mirb. l. c. ; Lam. Illustr. t. 289 = E. punctata.
regia, Mirb. l. c. 338 = E. regia.
undulata, Mirb. l. c. = E. undulata.

BASILEOPHYTA, F. Muell. First. Gen. Rep. 16 (1858) = Pleiadia, A. Cunn. (Gentianac.).
Friderici-Augusti, F. Muell. l. c. = F. australis.

BASILICUM, Moench, Meth. Suppl. 148 (1802) = Ocimum, Tourn. (Labiat.).
polystachion, Moench, l. c. = O. Basilicum.

BASILLAEA, R. Hedw. Gen. 289 (1809) = BASILAEA, Juss. = Bucomis, *L'Hérit.* (Liliac.).

BASILIMA, Raf. New Fl. Am. iii. 75 (1836) = Ocimum, Linn. (Rosac.).
pygmea, Raf. l. c. 76.—Am. bor.
scrobifolia, Raf. l. c. 75 = S. scrobifolia ?

BASISTEMON, Turcz. in Bull. Soc. Nat. Mosc. (1858) 214. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 1044.
bogotensis, *Turcz. l. c.*—N. Granat.
peruviana, *Benth. & Hook. f. Gen.* ii. 1245.—Peruv.

BASKERVILLA, Lindl. Gen. et Sp. Orch. 508 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 597.
assurgens, *Lindl. l. c.*—Peruv.

BASONCA, Rafin. Fl. Tellur. iv. 68 (1836) = **Martynia**, Linn. (Pedalin.).
longiflora, Rafin. l. c. = M. longiflora.

BASSELINIA, Vieill. in Bull. Soc. Linn. Normand. Sér. II. vi. 1872 (1873) 230. *PALMAE*.
Deplanchei, *Brongn. & Gris, l. c.* 232, nomen.—N. Caled.
gracilis, *Brongn. & Gris, l. c.* 231, nomen.—N. Caled.
kanaliensis, *Vieill. l. c.* 232, nomen.—N. Caled.
Lenormandi, *Vieill. l. c.* nomen.—N. Caled.
Pancheri, *Brongn. & Gris, l. c.* nomen.—N. Caled.

BASSIA, All. Misc. Taur. iii. 177. t. 4 (1766) = Chenolea, Thunb. (1781) (Chenopod.).
astrocarpha, F. Muell. Census, 140 = C. astrocarpa.
astrocarpa, F. Muell. Fragm. xii. 12 = C. astrocarpa.
bicornis, F. Muell. Census, 80 = Sclerolaena bicornis.
bicuspis, F. Muell. l. c. = Anisacantha bicuspis.
bifora, F. Muell. l. c. = Sclerolaena bifora.
Birchii, F. Muell. l. c. = Anisacantha Birchii.
brevicuspis, F. Muell. l. c. = Anisacantha brevicuspis.
carnosa, F. Muell. l. c. = C. carnosa.
Cornishiana, F. Muell. in Melb. Chem. & Drug. (1885). —Austral.
Dallachyana, F. Muell. Census, 80 = C. Dallachyana.
diacantha, F. Muell. l. c. = Sclerolaena diacantha.
divaricata, F. Muell. l. c. = Anisacantha divaricata.
Drummondii, F. Muell. l. c. = Anisacantha Drummondii.
echinopsila, F. Muell. l. c. = Anisacantha echinopsila.
enchylaenoides, F. Muell. l. c. = C. enchylaenoides.
Erahiensis, F. Muell. in Melb. Chem. & Drug. (Apr. 1885).—N. Guinea.
eurotioides, F. Muell. Census, 80 = C. eurotioides.
Forrestiana, F. Muell. Fragm. xii. 12 = C. Forrestiana.
glabra, F. Muell. Census, 80 = Anisacantha glabra.
hirsuta, Aschers. ex Nym. Consp. 625 = C. hirsuta.
lanicuspis, F. Muell. Census, 80 = Sclerolaena lanicuspis.
micrantha, F. Muell. l. c. = Enchylaena micrantha.
Muelleri, F. Muell. l. c. = C. Muelleri.
muricata, All. Misc. Taur. iii. t. 4 = C. hirsuta.
paradoxa, F. Muell. Census, 80 = Sclerolaena paradoxa.
quinquecuspis, F. Muell. l. c. = Anisacantha muricata.
salsuginosa, F. Muell. l. c. = Threlkeldia salsuginosa.
sclerolaenoides, F. Muell. l. c. = C. sclerolaenoides.
tricornis, F. Muell. l. c. = C. tricornis.
tridens, F. Muell. Fragm. xii. 13 = C. tridens.
uniflora, F. Muell. Census, 80 = Sclerolaena uniflora.

BASSIA, Koenig, ex Linn. Mant. ii. App. 555 (1771). *SAPOTACEAE*, Benth. & Hook. f. ii. 668.
DARTALUS, Thw. Enum. Pl. Zeyl. 176 (1860).
ILLIPE, F. Muell. Extra-trop. Pl. Am. ed. [5] (1884) 181.
MADHUCA, J. F. Gmel. Syst. 799 (1791).
albescens, Griseb. Cat. Pl. Cub. 164 (= Labourdonnaisiae sp.).—Cuba occ.
amicorum, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 337.—Isa. Pacif.
argentea, *De Vriese, Pl. Reinw.* 62.—Malacca.
Balem, *Miq. Fl. Ind. Bat. Suppl.* i. 583.—Sumatra.
Blanconi, *A. DC. in DC. Prod.* viii. 199.—Ins. Philipp.
butyracea, *Roxb. in As. Res.* viii. (1805) 477.—Ind. or.
caloneura, *Kurz, Andaman. Rep.* 141.—Ins. Andaman.
canaliculata, Bedd. Forest. Man. Bot. 141 = Dichopsis canaliculata.
Cocco, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 34.—N. Guinea.
cuneata, *Blume, Bijdr.* 675.—Malaya.
dubia, *Gaertn. Fruct.* ii. 104. t. 104.—Amboin.
elliptica, Dals. in Hook. Kew Journ. iii. (1851) 36 = Dichopsis elliptica.
elongata, *Miq. Fl. Ind. Bat. Suppl.* 582.—Sumatra.
falva, *Bedd. Forest. Man. Bot.* 140.—Ind. or.

BASSIA :—
galactoxyla, F. Muell. Fragm. vi. 27 = Lucuma galactoxylon.
grandiz, Bedd. Fl. Sylv. ii. t. 234 = Dichopsis grandi*.
hypoleuca, Miq. Fl. Ind. Bat. Suppl. 582 = Dichopsis obovata.
insignis, *Radlk. in Sitzb. Math.-Phys. Acad. Munch.* xii. (1882) 209.—Ind. or.
Jussaei, Griseb. Fl. Brit. W. Ind. 402 (sphalm.—*Cf.* Achras mammosa).—Ind. occ.
Korthalsii, *Pierre, ex Burck, in Ann. Jard. Buitenz.* v. (1885) 45.—Sumatra.
Krantzii, *Hance, in Journ. Bot.* xiv. (1876) 260.—Cambodia.
laevifolia, Bedd. Forest. Man. Bot. 141 = Dichopsis laevifolia.
lanceolata, Bedd. l. c.—Dichopsis lanceolata.
lanceolata, *A. DC. in DC. Prod.* viii. 199.—Ins. Philipp.
latifolia, *Roxb. Pl. Corom.* i. 20. t. 19.—Ind. or.
Lobbii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 546. —Burma.
longifolia, *Linn. Mant.* ii. *App.* 563.—Ind. or.
malabarica, *Bedd. Forest. Man. Bot.* 140.—Ind. or.
microphylla, *Hook. Ic. Pl.* i. t. 74.—Zeylan.
Moonii, *Bedd. Forest. Man. Bot.* 140.—Ind. or.
Mottleyana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 546.—Malaya.
neritifolia, *Bedd. Forest. Man. Bot.* 140.—Ind. or.
obovata, *Forst. f. Prod.* n. 200.—Ins. Tanna.
oleifera, *A. DC. in DC. Prod.* viii. 199.—Ins. Philipp.
pallida, *Burck, in Ann. Jard. Buitenz.* v. (1885) 44.—Sumatra.
Parkii, G. Don, Gen. Syst. iv. 86 = Butyrospermum Parkii.
parvifolia, A. DC. in DC. Prod. viii. 198 = microphylla.
pauciflora, Bedd. Forest. Man. Bot. 141 = Dichopsis pauciflora.
petiolaris, *Bedd. l. c.* 140.—Ind. or.
polyantha, Wall. Cat. n. 4166 = Dichopsis polyantha.
rubiginosa, Bedd. Forest. Man. Bot. 141 = Dichopsis rubiginosa.
serices, *Blume, Bijdr.* 674.—Malaya.
villosa, Wall. Cat. n. 4165 = latifolia.
Wightiana, Bedd. Forest. Man. Bot. 141 = Isonandra Wightiana.

BASSOVIA, Aubl. Pl. Gui. i. 217. t. 85 (1775). *SOLANACEAE*, Benth. & Hook. f. ii. 892.
AURELIANA, Scudd. in Mart. Fl. Bras. x. 188. t. 19 (1846).
WITHERINGIA, L'Hérit. Sert. Angl. i. f. 1 (1788).
brachypoda, *Dun. in DC. Prod.* xiii. 411.—Peruv.
cornuta, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 59.—Bras.
crassiflora, *Dun. in DC. Prod.* xiii. 409.—Bras.
fasciculata, *Dun. l. c.* 408.—Bras.
Gardneri, *Dun. l. c.* 409.—Bras.
glomuliflora, *Dun. l. c.* 405.—Bras.
hebepoda, *Dun. l. c.* 407 = lucida ?
laevis, *Dun. l. c.* 408.—Java.
leptopoda, *Dun. l. c.* 411.—Bras.
lucida, *Dun. l. c.* 408.—Guatemala.
pyrrastr, *Dun. l. c.* 408.—Bras.
Richardi, *Dun. l. c.* 408.—Guiana.
silvatica, Blanco, Fl. Filip. ed. II. 95 = Massa indica.
solanacea, *Benth. & Hook. f. Gen.* ii. 891.—Hab. ?
Spina-alba, *Griseb. in Gottt. Abhandl.* xxiv. (1879) 248.—Am. bor.
sylvatica, *Aubl. Pl. Gui.* i. 217.—Guiana.
tomentosa, *Dun. in DC. Prod.* xiii. 409.—Bras.
velutina, *Dun. l. c.* 410.—Bras.
Wallichii, Dun. l. c. 409 = Solanum crassipetalum.

BASTARDIA, H. B. & K. Nov. Gen. et Sp. v. 254. t. 472 (1821). *MALVACEAE*, Benth. & Hook. f. i. 203.
BOOMHARDIA, Reichb. Nom. 200 (1841).
angulata, Guill. & Perr. Tent. Fl. Senег. i. 65 = Abutilon intermedium.
aristata, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 300 = bivalvis.
bivalvis, *H. B. & K. Nov. Gen. et Sp.* v. 255, in nota.—Jamaica.
crispa, St. Hil. Fl. Bras. Mer. i. 194, in nota = Abutilon crispum.
foetida, Sweet, Hort. Brit. ed. II. 64 = Abutilon foetidum.
guayquilensis, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 300 = viscosa.
hirsutissima, Presl, Reliq. Haenk. ii. 112 = viscosa.

BASTARDIA :—
hirsutissima, Walp. Rep. i. 337 = praec.
nemoralis, St. Hil. Fl. Bras. Mer. i. 195 = Abutilon crispum.
parvifolia, H. B. & K. Nov. Gen. & Sp. v. 256 = viscosa.
spinifex, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 186 = bivalvis.
viscosa, *H. B. & K. Nov. Gen. et Sp.* v. 256.—Mexic.

BASTERA, J. F. Gmel. Syst. 1271 (1791) = BASTERIA, Houtt. = **Berkheya**, Ehrh. (Compos.).

BASTERIA, Mill. Gard. Dict. ed. VII (1759) = **Calycanthus**, Linn.
Calycanthus, Mill. l. c. ed VIII = C. floridus.

BASTERIA, Houtt. Nat. Hist. Pl. vi. 158 (1776) = **Berkheya**, Ehrh. (Compos.).
aculeata, Houtt. l. c. = Berkheya obovata.

BASTIA, Steud. Nomt. ed. II. i. 288, in syn. (1840), sphalm. = BUSTIA, Adans. = **Buphthalmum**, Linn. (Compos.).

BATANTHES, Rafin. Atl. Journ. 145 (1832) = **Gilia**, Ruiz & Pav. (Polemon.).
aggregata, Rafin. l. c. = G. aggregata.
longiflora, Rafin. l. c. = Collomia longiflora.
pungens, Rafin. l. c. = G. pungens.

BATATAS, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 434 ; (Conv Or. 52) = **Ipomoea**, Linn. (Convolv.).
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 64 = I. abyssinica.
acetosaefolia, Choisy, Conv. Rar. 124 = I. carnosa.
auriculata, Hochst. ex Choisy, in DC. Prod. ix. 350 = I. coriofanа.
betacea, Cavall. in Bot. Reg. (1839) Misc. 93 ; (1840) t. 56.—Guiana.
bignoniacaea, G. Don, Gen. Syst. iv. 261 = I. bignoniacea.
bombycina, Choisy, in DC. Prod. ix. 340 = I. bombycina.
bonariensis, Lindl. in Bot. Reg. (1838) Misc. 59 = I. fucifolia.
Canavillesii, G. Don, Gen. Syst. iv. 262 = I. Cavanillesii.
Choisyana, Wight, Ic. t. 491 = Argyreia Choisyana.
cissoides, Choisy, Conv. Or. 55 = I. cissoides.
crassicaulis, Benth. Bot. Voy. Sulph. 134.—Guayaquil.
edulis, Choisy, Conv Or. 58 = I. Batatas.
glaberrima, Hassk. Retzia, i. 65.—*Hab.* ?
glabra, Benth. in Hook. Lond. Journ. Bot. v. (1846) 362 = I. glabra.
glaucifolia, G. Don, Gen. Syst. iv. 262 = I. glaucifolia.
heterophylla, G. Don, l. c. 261 = I. heterophylla.
incurva, Benth. in Hook. Niger Fl. 464 = I. incurva ?
insignis, G. Don, in Sweet, Hort. Brit. ed. III. 482 = I. insignis.
Jalapa, Choisy, Conv. Rar. 125 = I. Jalapa.
littoralis, Choisy, l. c. 124 = I. carnosa.
Lourteiri, G. Don, Gen. Syst. iv. 262 = I. tuberosa.
macrorhiza, G. Don, l. c. 261 = I. macrorhizos.
maritima, Boj. Hort. Maurit. 225 = I. biloba.
multifora, Boj. ex Choisy, in DC. Prod. ix. 371 = I. staphylina.
paniculata, Choisy, Conv. Or. 54 = I. digitata.
Papirio, G. Don, Gen. Syst. iv. 261 = I. pubescens.
parvirasefolia, Choisy, Conv. Rar. 123 = I. parasitaefolia.
pentaphylla, Choisy, Conv. Or. 54 = I. pentaphylla.
pulchella, Boj. Hort. Maurit. 226 = I. pulchella.
quinquefolia, Choisy, Conv. Rar. 127 = I. quinquefolia.
senegalensis, G. Don, Gen. Syst. iv. 261 = I. palmata.
sinuata, Ker, Ic. Exsm. Pl. Ins. Inar. (1824) 210 = I. stolonifera.
subtriloba, G. Don, Gen. Syst. iv. 261 = I. pubescens.
ternata, G. Don, l. c. 262 = I. ternata.
tomentosa, Choisy, in DC. Prod. ix. 337 = I. tomentosa.
triloba, Choisy, Conv. Rar. 127 = I. triloba.
tuberosa, Boj. Hort. Maurit. 226 = I. tuberosa.
venosa, Boj. l. c. = I. cairica.
venosa, G. Don, Gen. Syst. iv. 261 = I. venosa.
villosa, Choisy, in DC. Prod. ix. 337 = I. villosa.
viridis, Voigt, Hort. Suburb. Calc. 354.—Ind. or. ?
Wallii, Morr. in Ann. Gand. ii. (1846) 285 = I. Wallii.
Wildenowii, G. Don, Gen. Syst. iv. 261 = I. heterophylla.
xanthorhiza, Boj. Hort. Maurit. 225 = I. Batatas.

BATEMANIA, Endl. Gen. 197 (1837) = seq.

BATEMANNIA, Lindl. in Bot. Reg. t. 1714 (1834). *ORCHIDEAE*, Benth. & Hook. f. iii. 546.
apiculata, *Reichb. f. in Linnaea*, xli. (1877) 109.—N. Granat.
armillata, *Reichb. f. in Gard. Chron.* (1878) i. 780.—Hab. ?
Beaumontii, *Reichb. f. in Walp. Ann.* iii. 544.—Bras.
Burtii, Endr. & Reichb. f. in Gard. Chron. (1878) 1099 = Zygopetalum Burtii.
Colleyi, *Lindl. Bot. Reg.* xx. (1885) t. 1714.—Guiana.
fimbriata, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 280.—N. Granat.
grandiflora, Reichb. f. l. c. iv. (1856) 328 = Zygopetalum grandiflorum.
Gustavi, *Reichb. f. in Linnaea*, xli. (1877) 108.—N Granat.
Lalindei, *Linden, Cat.* (1878) n. 90.—N. Granat.
lepida, *Reichb. f. in Gard. Chron.* (1878) i. 588.—Bras.
Meleagris, *Reichb. f. in Bonplandia*, iii. (1855) 217.—Bras.
Wallisii, *Reichb. f. in Gard. Chron.* (1880) i. 776.—N. Granat.

BATERIUM, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiii. (1864) 124 = **Haematocarpus**, Miers (1867) (Menisp.).
validum, Miers, l. c. = H. Thomsoni.

BATESIA, Spruce, in Benth. & Hook. f. Gen. i. 668 (1865). *LEGUMINOSAE*, Benth. & Hook. f. i. c. floribunda, *Spruce, ex Benth. in Trans. Linn. Soc.* xxv. (1866) 308. t. 37.—Bras. bor.

BATHYSA, Presl, in Abh. Boehm. Ges. Folge V. iii. (1845) 514 ; (Bot. Bemerk. 84). *RUBIACEAE* Benth. & Hook. f. ii. 49.
SCHOENLEINIA, Klotzsch, in Hayne, Arzneigew. xiv in nota ad t. 15 (1846).
VOIGTIA, Klotzsch, l. c. (1846)
stipulata, *Presl, l. c.*—Bras.

BATINDUM, Rafin. Sylva Tellur. 81 (1838) = **Oftia**, Adans. (Myopor.).
jasmineum, Rafin. l. c. 82 = O. africana.

BATIS, Linn. Syst. ed. X. 1380. n. 1172 (1759). *BATIDEAE*, Benth. & Hook. f. iii. 88.
americana, *Linn. Mant.* ii. 499.—Am. bor.
aurantiaca, Wall. ex Thw. Enum. Pl. Zeyl. 268 = Plecospermum spinosum.
californica, Torr. in Smithson. Contrib. vi. (1854) 8 = maritima.
fruticosa, Roxb. Hort. Beng. [71] ; Fl. Ind. iii. 768 — Cadrania javanensis et C. fruticosa.
hermaphrodita, *Blanco, Fl. Filip.* ed. I. 792.—Ins. Philipp.
maritima, *Linn. Syst.* ed. X. 1380.—Jamaica.
spinosa, Roxb. Hort. Beng. [71] ; Fl. Ind. iii. 762 = Plecospermum spinosum.
vermiculata, Hook. Fl. Bor. Am. ii. 128 = Sarcobatus vermiculatus.

BATOCARPUS, Karst. Fl. Colomb. ii. 67. t. 124 (1869?). *URTICACEAE*, Benth. & Hook. f. iii. 376.
orinocensis, *Karst. l. c.*—Venezuela.

BATOCVDIA, Mart. ex DC. Prod. ix. 148 (1845) = **Bignonia**, Linn.
caprolata, Mart. l. c. 146 = Bignonia caprolata.
cuneifolia, Mart. l. c. 157 = Bignonia cuneifolia.
exoleta, Mart. l. c. 145 = Bignonia exoleta.
Unguis, Mart. l. c. 146 = Bignonia Unguis-cati.

BATODENDRON, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1843) 261 = **Waoclinium**, Linn.
arboreum, Nutt. l. c. = V. arboreum.

BATRACHIUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 730 (1821) = **Ranunculus**, Tourn.
admirabile, Nyl. in Nyl. & Scal. Herb. Mus. Fenn. 35 = R. confervoides.
aquatile, Wimm. Fl. Schles. (1841), ii. 8 = R. aquatilis.
aspergillifolium, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 218 = R. trichophyllus.

BATRACHIUM :—
Backii, Wirtg. ex F. Schultz, Arch. Fl. Fr. et Allem. i. (1844) 293 = Ranunculus fluitans.
Baudotii, F. Schultz, l. c. 71 = R. Baudotii.
bipontinum, F. Schultz, ex Gren. & Godr. Fl. Fr. i. 24 = R. trichophyllus.
biternatum, Bercht. & Presl, Rostl. i. Ranunc. 49 = R. biternatus.
caespitosum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 722 = R. circinatus.
caespitosum, F. Schultz, Arch. Fl. Fr. et Allem. i. (1844) 71 = R. trichophyllus.
capillaceum, Bercht. & Presl, Rostl. i. Ranunc. 49 = R. trichophyllus.
carinatum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 28 = R. Baudotii.
circinatoides, Arv. in Soc. Ech. Dauph. iv. (1877) 105 = R. aquatilis ?
circinatum, Spach, Hist. Vég. Phan. vii. 201 = R. circinatus.
coenosum, Nym. in Bot. Notiser (1852), 100 = R. coenosus.
confervoides, Fries, in Bot. Notiser (1845), 121 = R. coenosus.
confusum, Garcke, Fl. Deutschl. ed. V. 8 = R. Petiveri.
confusum, Le Jolis, in Mém Soc. Sc. Nat. Cherb. vii. (1860) 257 = R. confusus.
dichotomum, Schmalh. ex Trautv. in Act. Hort. Petrop. ix. (1884) 363 = R. aquatilis.
divaricatum, Wimm. Fl. Schles. (1841) 10 = R. divaricatus.
diversifolium, Nym. Consp. 15 = R. diversifolius.
Drouetii, Nym. in Bot. Notiser (1852) 98 = R. Drouetii.
elongatum, F. Schultz, in Billotia, i. 113 = R. peltatus.
eradicatum, Lacst. ex Nym. Consp. 15 = R. confervoides.
flaccidum, Rupr. Fl. Cauc. 15 = R. aquatilis.
floribundum, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 216 = R. floribundus.
fluitans, Wimm. Fl. Schles. (1841) 9 = R. fluitans.
fluviatile, S. F. Gray, Nat. Arr. ii. 722 = R. fluviatilis.
Friesii, Nym. Consp. 15 = R. heterophyllus.
Godronii, Nym. l. c. 16 = R. diversifolius.
hederaceum, S. F. Gray, Nat. Arr. ii. 721 = R. hederaceus.
heterophyllum, S. F. Gray, l. c. = R. heterophyllus.
heterophyllum, Lange, Pugill. 251 = R. peltatus.
hololeucum, Garcke, Fl. Deutschl. ed. IV. 7 = R. hololeucus.
intermedium, Nym. Syll. 175 = R. intermedius.
Langei, F. Schultz, ex Nym. Consp. 16 = R. tripartitus.
leiospermum, Hartm. f. Skand. Fl. ed. XIII. 165 = R. hederaceus.
Lenormandii, F. Schultz, Arch. Fl. Fr. et Allem. i. (1844) 70 = R. Lenormandi.
longirostre, F. Schultz, l. c. 71 = R. longirostris.
lutarium, Kevel, in Act. Linn. Soc. Bord. xxv. (1864) 418. t. 4 = R. Lenormandi.
lutetum, Revel.—Cf. Bull. Soc. Bot. Fr. xii. (1865) Bibliog. 258 = R. heterophyllus.
lutulentum, Nym. Syll. Suppl. 29 = R. trichophyllus.
marinum, *b tus*, Nov. Mant. iii. 51 = R. Baudotii.
maximum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 28 = R. trichophyllus.
obtusiflorum, S. F. Gray, Nat. Arr. ii. 721 = R. tripartitus.
oleiocaes, [Bossch] Prod. Fl. Bat. 7 = R. hololeucus.
omoiophyllum, Nym. Consp. 16 = R. hederaceus.
pantothrix, S. F. Gray, Nat. Arr. ii. 723 = R. trichophyllus.
paucistaminum, F. Schultz, Arch. Fl. Fr. et Allem. (1844) 71 = R. trichophyllus.
pectinatum, Nym. Consp. 15 = R. trichophyllus.
peltatum, Bercht. & Presl, Rostl. i. Ranunc. 49 = R. peltatus.
penicillatum, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 216 = R. peltatus.
Petiveri, F. Schultz, Arch. Fl. Fr. et Allem. i. (1844) 71 = R. Petiveri.
pseudaquatilifolium, Dum. Fl. Belg. 127 = R. fluitans.
pseudofluitans, Nym. Consp. 16 = R. peltatus.
pumilum, Nym. in Bot. Notiser (1852), 98 = R. fluitans.
radians, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 215 = R. trichophyllus.

BATRACHIUM :—
radians, Revel, in Act. Linn. Soc. Bord. xix. (1853) 130 = Ranunculus radians.
rhipidyllum, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 215 = R. diversifolius.
rigidum, Dum. Fl. Belg. 127 = R. trichophyllus.
Rionii, Nym. in Bot. Notiser (1850) = R. scleratus.
tenellum, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 211 = R. tenellus.
trichophyllum, [Bosch] Prod. Fl. Bat. 7 ; Des Moul. in Act. Soc. Linn. Bord. xx. (1859) 456 = R. trichophyllus.
trinacrium, Nym. Consp. 16 = R. tripartitus.
tripartitum, S. F. Gray, Nat. Arr. ii. 721 = R. tripartitus.
triphyllos, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 214 = R. tripartitus.
truncatum, Dum. l. c. 215 = R. peltatus.
villosum, F. Schultz, ex Gren. & Godr. Fl. Fr. i. 24 = R. trichophyllus.

BATRATHERUM, Nees, in Edinb. N Phil. Journ. xviii. (1835) 180 = **Arthraxon**, Beauv. (Gram.).
cuspidatum, Hochst. in Flora, xxxix. (1856) 178 = A. cuspidatus.
echinatum, Nees, in Edinb. N. Phil. Journ. xviii. (1835) 181 = A. lanceolatus.
lanceolum, Nees, l. c. = A. lanceolatus.
micans, Nees, l. c. 182 = A. cuspidatus.
molle, Nees, l. c. 181 = A. microphyllus.
nudum, Nees, ex Steud. Syn. Pl. Gram. 388 = A. ciliaris.
plumbeum, [Munro, ex] Duthie, in Atkinson, Gazetteer, 640 = A. ciliaris
rude, Nees, ex Steud. Syn. Pl. Gram. 388 = A. rudis.
Schimperi, Nees, ex Hochst. in Flora, xxxix. (1856) 179 = A. microphyllus.
serrulatum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii 458 = A. lanceolatus.
submuticum, Nees, ex Steud. Syn. Pl Gram. 382 = A. ciliaris.

BATSCHIA, J. F. Gmel. Syst. ii. 815 (1791) = **Lithospermum**, Linn. (Boragin.).
albiflora, Raf. in New Fl. Am. iv. 19.—Am. bor.
canescens, Michx. Fl. Am. Bor. i. 130. t. 14 = L. canescens.
carolinianum, Roem. & Schult. Syst. iv. 52 = L. hirtum.
carolinianum, J. F. Gmel. Syst. i. 315 = L. hirtum.
conspicuum, R. Br. in Richards. App. Frankl. Journ. 782 = L. canescens.
decumbens, Nutt. Gen. Am. i. 114 = L. angustifolium.
distichum, G. Don, Gen. Syst. iv. 326 = L. distichum.
Gmelinii, Michx. Fl. Am. Bor. i. 130 = L. hirtum.
longiflora, Pursh, Fl. Am. Sept. i. 132 = L. angustifolium.
pilosa, G. Don, Gen. Syst. iv. 372 = L. pilosum.
sericea, Roem. & Schult. Syst. iv. 743 = L. canescens.
Torreyi, G. Don, Gen. Syst. iv. 372 = L. pilosum.

BATSCHIA, Moench, Meth. 567 (1794) = **Eupatorium**, Linn. (Compos.).
nivea, Moench, l. c. = E. ageratoides.

BATSCHIA, Mutis, ex Thunb. in Nov. Act. Ups. v. (1792) 130. t. 2 = **Abuta**, Aubl. (Menisp.).
conferta, Miers, in Seem. Bot. Voy. Herald, i. 76 = A. Seemanni.
conferta, DC. Prod. i. 108 = A. spicata.
racemosa, Thunb. in Nov. Act. Ups. v. (1792) 122 = A. racemosa.
spicata, Thunb. l. c. = A. spicata.

BATSCHIA, Vahl, Symb. Bot. iii. 89. t. 56 (1794) = **Humboldtia**, Vahl (Legum.).
laurifolia, Vahl, l. c. = H. laurifolia.

BATTATA, Hill, Hort. Kew. 148 (1768) = **Solanum**, Linn.
tuberosa, Hill, l. c. = S. tuberosum.

BAUCHRA, Fourn. ex Benth. in Journ. Linn. Soc. xix. (1881) 88. *GRAMINEAE*, Benth. & Hook. f. iii. 1140.
Karwinskii, *Fourn. Mex. Pl. Fn. Gram.* 87 ; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 550.—Mexic.

BAUDINIA, Lesch. ex DC. Prod. iii. 211 (1828) = **Calothamnus**, Labill. (Myrtac.).

BAUDINIA, Lesch. ex DC. Prod. vii. 511 (1839) = **Scaevola**, Linn. (Goodenov.).
humilis, Lesch. ex DC. l. c. 511 = S. striata.

BAUDOUINIA, Baill. Adansonia, vi. (1866) 193. t. 5. *LEGUMINOSAE*, Benth. & Hook f. i. 1003.
fluggeiformis, *Baill. l. c.* viii. (1868) 201.—Madag.
sollyaeformis, *Baill. l. c.* vi. (1866) 194.—Madag

BAUERA, Banks, ex Andr. Bot. Rep. t. 198 (1793). *SAXIFRAGEAE*, Benth. & Hook. f. i. 655.
Billardieri, D. Don, in Edinb. N. Phil. Journ. ix. (1830) 96 = rubioides.
capitata, *Ser. in DC. Prod.* iv. 13 —Austral.
galioides, Sieber, in Reichb. Ic. Exot. t. 77 = rubioides.
humilis, Sweet, Hort. Suburb. 124 = rubioides
juncea, Walp. Rep. ii. 875 = Tetrathecajuncea.
microphylla, Sieber, ex DC. Prod. iv. 13 = rubioides.
rubaefolia, Salisb. in Kon. & Sims, Ann. Bot. i. (1805) 514. t. 10 = rubioides.
rubioides, *Andr. Bot. Rep.* t. 198.—Australia.
sessiliflora, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 41.—Austral.

BAUHINIA, Linn. Gen. ed. 1. 126 (1787). *LEGUMINOSAE*, Benth. & Hook. f. i 575.
ALVESIA, Welw. Apont. 557. n 47 (1858).
AMAHIA, Mutis, ex Caldas, in Seman. Nuev. Gran. ii. (1810) 25.
BINARIA, Rafin. Sylva Tellur. 122 (1838).
CASSENIA, Rafin l. c. (1838)
CASPAREA, Ll. B. & K. Nov. Gen. et Sp. vi. 317 (1823).
CAULOTRETUS, Rich. ex Spreng. Syst. iv. Cur. Post. 406 (1827).
ELAYUNA, Rafin. Sylva Tellur. 145 (1838).
LACARA, Spreng. Neue Entdeck. iii. 56 (1822).
LASIOSEMA, Miq. Fl. Ind. Bat. i. 71 (1855).
LOCELLARIA, Welw. Apont. 588 (1858).
MANDARUS, Rafin. Sylva Tellur. 122 (1838).
MONOTELES, Rafin. l. c. (1838).
PAULETIA, Cav. Ic. v. 5. t. 466, 410 (1799).
PERLEBIA, Mart. Reise, i. 555 (1828).
PHANERA, Lour. Fl. Cochinch. 37 (1790)
PILOSTIGMA, Hochst. in Flora, xxix. (1846) 598.
SCHNELLA, Raddi, in Mem. Mod. xviii. Fis. (1820) 411.
TELESTRIA, Rafin. Sylva Tellur. 122 (1838).
abyssinica, A. Rich Tent Fl Abyss. i. 253 = reticulata.
acida, Reinw. in Flora, xxxi. (1848) 578 = malabarica.
aculeata, *Linn. Sp. Pl.* 374.—Am. austr.
aculeata, Vell. Fl. Flum. 171 ; iv. t. 83 = forficata.
acuminata, Bruce, Trav. Abyss. App. 167. t. 18 = B. Farea.
acuminata, *Linn. Sp. Pl.* 376.—Ind. or. ; Malaya ; China.
acuminata, Vell. Fl. Flum. 171 ; iv. t. 85.—Bras.
Aquensis, *Morsc. Pl. Nouv. Kar. Am.* 77. t. 51.—Bras.
Adansoniana, *Guill. & Perr. Tent. Fl. Senegi.* i. 265.—Afr. trop.
aegopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69, 120 = B. Pes-caprae.
affinis, *Vog. in Linnaea*, x. (1836) 594.—Bras.
Aimoata, Steud. Nom. ed. II. i. 101 (err. typ. 291), vide *Outmouti*.
alba, Buch.-Ham. ex Wall. Cat. sub n. 5796 = variegata.
albicans, *Vog. in Linnaea*, xiii. (1839) 304.—Bras.
americana, *Delaun. Herb. Amat.* t. 315.—Am. austr.
ampla, *Span. Ic.* 29 ; *in Linnaea*, xv. (1841) 202.—Timor.
anatomica, *Link, Enum. Hort. Berol.* i. 405.—Am. austr.
Andrieuxii, *Hemsl. Diag. Pl. Nov.* 48.—Mexic.
anguina, *Roxb. Hort. Beng.* 31 ; *Fl. Ind.* ii. 328.—Ind. or. ; Malaya.
anguicosa, *Vog. in Linnaea*, xiii. (1839) 312.—Bras.
armata, *Otto, Pl. Bras.* 25.—Bras.
articulata, Oliver, Fl. Trop. Afr. ii. 290, err. typ. = reticulata !
aurantiaca, *Boj. in Ann. Sc. Nat.* Sér. II. iv. (1835) 264.—Madag.
aurita, *Ait. Hort. Kew.* ii. 48.—Jamaica.

BAUHINIA:—
Bacurie, Buch.-Ham. ex Wall. Cat. sub n. 5793 = malabarica.
bahienais, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. (1836) 114 = Langsdorffiana.
Bauerriana, Steud. Nom. ed. II. i. 191 (err. typ. 291) = malabarica.
Benthami, Bedd. Forest. Man. Bot. 98; Ic. Pl. Ind. Or. i. 23 = phoenicea.
Benzoin, *Kotschy, ex Schweinf. Reliq. Kotsch.* (1868) 12. t. 11.—Afr. trop.
bicolor, *D. Dietr. Syn. Pl.* ii. 1476.—Bras.
bicuspidata, *Benth. in Mart. Fl. Bras.* xv. ii. 193.—Am. trop.
bidentata, *Yach, in Malay. Misc.* ii. (1822) n. vii. p. 76.—Malaya.
binata, Blanco, Fl. Filip. ed. I. 331 = Blancoi.
binata, Naves, ex Villar, in Blanco, Fl. Phil. ed. III. Nov. App. 72 = tomentosa.
Blancoi, *Benth. Fl. Austral.* ii. 396, in note; *Baker, in Hook. f. Fl. Brit. Ind.* ii. 278.—Siam; Ins. Philipp.
Bongardi, *Steud. Nom.* ed. II. i. 191 (err. typ. 291).—Bras.
Bowkeri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 396.—Afr. austr.
brachycarpa, *Wall. Cat.* n. 5786.—Burma.
brachystachya, Walp. Rep. i. 852 = cumanensis.
bracteata, *R. Grah. in Wall. Cat.* n. 5802.—Burma; Siam.
brasiliensis, *Vog. in Linnaea*, xiii. (1839) 302.—Bras.
Bredemeyeri, *Vog. l. c.*—Venezuela.
brevilobe, *Benth. in Mart. Fl. Bras.* xv. ii. 197.—Bras.
brevipes, Vog. in Linnaea, xiii. (1839) 307 = Bongardi.
Buchananii, Desv. in Ann. Sc. Nat. Sér. I. ix. (1839) 490 = diphylla.
Burchelli, *Benth. in Mart. Fl. Bras.* xv. ii. 186.—Bras.
Burkeana, Benth. ex Harv. & Sond. Fl. Cap. ii. 275 = esculenta.
Cancave, Buch.-Ham. ex Steud. Nom. ed. II. i. 191 (err. typ. 291) = variegata.
candicans, *Benth. in Mart. Fl. Bras.* xv. ii. 201.—Bras.
candida, Ait. Hort. Kew. ii. (1789) 49 = acuminata.
candida, Roxb. Hort. Beng. [81]; Fl. Ind. ii. 318 = variegata.
Carronii, *F. Muell. in Trans. Phil. Inst. Victoria*, iii. (1858) 49.—Austral.
castrata, *Blanco, Fl. Filip.* ed. I. 331.—Ins. Philipp.
castrata, *Hassk. in Flora*, xxv. (1842) *Beibl.* ii. 54.—Java.
Championi, *Benth. Fl. Hongkong*, 99.—Hongkong.
chelianthe, *D. Dietr. Syn. Pl.* ii. 1476.—Bras.
chinensis, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 43 = variegata.
chrysophylla, Vog. in Linnaea, xiii. (1839) 307 = rufa.
chrysophylla, Vahl, ex DC. Prod. ii. 517 = macrophylla.
chrysophylla, Vog. in Linnaea, xiii. (1839) 313 = splendens.
cinnamomea, *DC. Prod.* ii. 517.—Guiana.
cisoidea, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 287.—Afr. trop.
coccinea, *DC. Prod.* ii. 516.—Cochinch.
columbinensis, Vog. in Linnaea, xiii. (1839) 313 = cumanensis.
compacta, *Hort. ex Regel, Cat. Pl. Absahow.* 15.—Hab.?
confertiflora, *Benth. in Mart. Fl. Bras.* xv. ii. 206.—Bras.
cordata, Vog. in Linnaea, xi. (1837) 308 = rufa.
cordifolia, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 322 = emarginata.
corniculata, *Benth. in Mart. Fl. Bras.* xv. ii. 196.—Bras.
cornifolia, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 278.—Ins. Penang.
corymbadelphiana, DC. Mém. Leg. xiii. 486; Prod. ii. 515 = purpurea.
coronata, *Benth. in Mart. Fl. Bras.* xv. ii. 209.—Bras.
corymbosa, *Roxb. Hort. Beng.* [81]; *Fl. Ind.* ii. 322.—China.
cucullata, *Desv. Journ. Bot.* i. (1814) 74(= variegata ?).—Ind. or.
cumanensis, *H. B. & K. Nov. Gen. et Sp.* vi. 231.—Venezuela.

BAUHINIA:—
Cuminginna, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App* 72.—Ins. Philipp.
Cunninghamii, *Benth. Fl. Austral.* ii. 295.—Austral.
cupulata, *Benth. in Mart. Fl. Bras.* xv. ii. 188.—Bras.
curvula, *Benth. l. c.* 194.—Bras.
cuyabensis, *Steud. Nom.* ed. II. i. 191 (err. typ. 291).—Bras.
debilis, *Hassk. in Hoev. & De Vriew, Tijdschr.* x. (1843) 149.—Java.
dipetala, *Hemsl. Diag. Pl. Nov.* 48.—Mexic.
diphylla, *Buch.-Ham. in Symes, Embassy*, ed. II. iii. 311.—Burma; Malaya.
dipters, *Blume, ex Miq. in Verh. Nederl. Inst.* III. iii. (1850) 12.—Malaya.
divaricata, Hort. ex Steud. Nom. ed. II. i. 191 (err. typ. 291) = forficata.
divaricata, Lam. Encyc. i. 329 = Lamarckiana.
divaricata, *Linn. Sp. Pl.* 374.—Am. austr.
divergens, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 282.—Burma.
dodecandra, *D. Dietr. Syn. Pl.* ii. 1475.—Bras.
dubia, *G. Don, Gen. Syst.* ii. 463.—Bras.
dubia, Vog. in Linnaea, xiii. (1839) 314 = coronata.
dumosa, *Benth. in Mart. Fl. Bras.* xv. ii. 194.—Bras.
ebenina, *Karsch, Verh. Nat. Gesch. Bot.* (1859–42) 89.—Borneo.
emarginata, *Yach, in Malay. Misc.* ii. (1822) n. vii. p. 76.—Sumatra.
emarginata, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
emarginata, Wall. Cat. n. 5792 = retusa.
eriostachya, *Tenore, Ind. Sem. Hort. Neap.* (1842) 12. —Hab.?
escalsa, Blume, ex Korth. Verh. Nat. Gesch. Bot. 90 = ferruginea.
esculenta, *Burch. Trav.* ii. 589.—Afr. austr.
excisa, *Hemsl. Biol. Centr. Am. Bot.* i. 337.—Panama.
exinia, *Miq. Fl. Ind. Bat.* i. 1. 75.—Java.
Farez, *Desv. Journ. Bot.* i. (1814) 74.—Abyssin.
fascigiensis, *Kotschy, ex Schweinf. Reliq. Kotsch.* (1868) 14. t. 12.—Afr. trop.
ferruginea, D. Dietr. Syn. Pl. ii. 1476 = Bongardi.
ferruginea, *Roxb. Hort. Beng.* 90; *Fl. Ind.* ii. 331.—Malaya.
Finlaysoniana, *R. Grah. in Wall. Cat.* n. 5801.—Malaya.
flexuosa, *Meric. Pl. Nouv. Am.* 80. t. 53.—Bras.
florifera, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 480.—Bras.
forficata, *Link, Enum. Hort. Berol. i.* 404.—Bras.
foveolata, Dalz. in Journ. Linn. Soc. xiii. (1873) 188 = Lawii!
fulva, *Korth. Verh. Nat. Gesch. Bot.* (1839–42) 91.—Java.
furcata, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 429. —Am. bor.
fusconervis, *D. Dietr. Syn. Pl.* ii. 1476.—Bras.
Gardneri, *Benth. in Mart. Fl. Bras.* xv. ii. 186.—Bras.
garipensis, *E. Mey. Comm. Pl. Afr. Austr.* i. 162.—Afr. austr.
gemella, *Vog. in Linnaea*, xiii. (1839) 306.—Bras.
Gilesii, *F. Muell. & Bailey, in Wing, South Sc. Record*, ii. (1882) 151.—Austral.
glabra, *Vaup. Enum. Pl. Carib.* 20.—N. Granat.
glabrifolia, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 281.—Burma.
glandulosa, *DC. Prod.* ii. 518.—N. Barcel.
glauca, *Wall. Cat.* n. 5785; *Benth. Fl. Hongkong*, 99. —Burma; Malaya; China.
glaucescens, *DC. Prod.* ii. 516.—Venezuela.
Grahami, Steud. Nom. ed. II. i. 191 (err. typ. 291) = rufa.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 578.—Madag.
grandiflora, *Juss. in Poir. Encyc. Suppl.* i. 600.—Persa.
grandiflora, Blanco, Fl. Filip. ed. I. 333 = variegata.
grandiflora, *D. Dietr. Syn. Pl.* ii. 1475.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 377. t. 148.—Guiana.
heterandra, *Benth. in Mart. Fl. Bras.* xv. ii. 196.—Bras.
heterophylla, *Kunth, Mim.* i. 46.—Venezuela.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 107. —Afr. trop.

BAUHINIA:—
hirsuta, *Vog. in Linnaea*, xiii. (1839) 300.—Bras.
hirsuta, Weinm. in Ratisb. Syll. ii. (1828) 9 = acuminata.
hirta, *Steud. Nom.* ed. II. i. 191 (err. typ. 291).—Bras.
holophylla, *Steud. l. c.*; *D. Dietr. Syn. Pl.* ed. II. 1476.—Bras.
Hookeri, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 51.—Austral.
Hostmanniana, *Miq. in Verh. Nat. Wet. Haarl.* vii. (1851) 10.—Guiana.
Humbloteana, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 865.—Madag.
hymenaeaefolia, *Triana, ex Hemsl. Biol. Centr. Am. Bot.* i. 337.—Panama.
indica, *Lodd. ex Loud. Hort. Brit.* 165.—Ind. or.
inermis, Forsk. Fl. Aegypt. Arab. 85 = reticulata, tomentosa.
inermis, *Pers.* Syn. i. 455.—Am. austr.
integerrima, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 197.—Bras.
integrifolia, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 321. —Malaya.
intermedia, Vog. in Linnaea, xi. (1837) 309 = rufa.
involucellata, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) 11. t. 73.—Burma.
isopetala, *Grif. Notul.* iv. 451.—Burma.
japonica, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 401.—Japon.
Kappleri, *Sagot, in Ann. Sc. Nat.* Sér. VI. xiii. (1882) 317.—Guiana.
khasiana, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 281.—Ind. or.
Kirkii, *Oliver, Fl. Trop. Afr.* ii. 288.—Afr. trop.
Kockiana, *Korth. Verh. Nat. Gesch. Bot.* (1839–42) 87. —Sumatra.
Krugii, *Urban, in Berl. Deutsch. Bot. Ges.* iii. (1885) 88.—Porto-Rico.
Kunthiana, *Vog. in Linnaea*, xiii. (1839) 312.—Am. austr.
Lamarckiana, *DC. Prod.* ii. 512.—Am. austr.
Langsdorffiana, *Bong. in Mém. Acad. Pétersb.* iv. (1836) 109.—Bras.
latifolia, *Cav. Ic.* v. 4. t. 405.—Mexic.
latisiliqua, *Cav. l. c.* 5.—Ins. Philipp.
Lawii, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 277.—Ind. or.
Leichhardtii, *F. Muell. in Trans. Phil. Inst. Victoria*, iii. (1859) 50.—Austral.
leiopetala, *Benth. in Mart. Fl. Bras.* xv. ii. 209.—Bras.
leptopetala, *Moç. & Sessé, ex DC. Prod.* ii. 513.—Mexic.
Lianea, *DC. l. c.* 516.—Malaya.
longicuspis, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 185.—Bras.
longiflora, D. Dietz. Syn. Pl. ii. 1475 = forficata.
longifolia, *D. Dietr. l. c.* 1476.—Bras.
longipetala, *Walp. Rep.* i. 852.—Guiana.
longistyla, Benth. in Mart. Fl. Bras. xv. ii. 185 = holophylla.
lucida, Wall. Cat. n. 5779 A = emarginata.
lunaria, *Cav. Ic.* v. 4. t. 407.—Mexic.
lunarioides, *A. Gray, in S. Wats. Bibliog. Ind. N. Am. Bot.* i. 206.—N. Mexic.
macrantha, *Oliver, Fl. Trop. Afr.* ii. 289.—Afr. trop.
macranthera, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 338.—Mexic.
macrophylla, *Poir. Encyc. Suppl.* i. 600.—Am. austr.
macrostachya, *Benth. in Hook. Journ. Bot.* ii. (1840) 96.—Guiana.
macrostachya, *Wall. Cat.* n. 5774.—Ind. or.
maculata, *Tenore, Cat. Hort. Neap.* (1845) 79.—Hab.?
madagascariensis, *Desv. Journ. Bot.* i. (1814) 74.—Madag.
malabarica, *Roxb. Hort. Beng.* [81]; *Fl. Ind.* ii. 321. —Ind. or.; Burma.
marginata, *D. Dietr. Syn. Pl.* ii. 1475.—Bras.
Maximiliani, *Benth. in Mart. Fl. Bras.* xv. ii. 202.—Am. trop.
megaladena, *Griseb. Fl. Brit. W. Ind.* 218.—Ind. occ.
membranacea, *Benth. in Mart. Fl. Bras.* xv. ii. 187.—Bras.
mexicana, *Vog. in Linnaea*, xiii. (1839) 299.—Mexic.
microphylla, *Vog. l. c.* 301.—Am. austr.

BAUHINIA :—
mollis, *D. Distr. Syn. Pl.* ii. 1475.—Bras.
mollissima, Wall. Cat. n. 5782 = elongata.
Mombassæ, Vatke, in Oestr. Bot. Zeitschr. xxx. (1880) 279.—Afr. trop.
monandra, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 78.—Burma.
mucora, C. Bolle, in Peters, Reise Mossamb. Bot. i. 22 = tomentosa.
multinervia, *DC. Prod.* ii. 515.—Venezuela.
natalensis, *Oliver, in Hook. Bot. Mag.* t. 6066.—Afr. austr.
nervosa, *Wall. Cat.* n. 5777.—Ind. or.
nitida, *Benth. in Mart. Fl. Bras.* xv. ii. 184.—Bras.
notophila, *Griseb. in Goett. Abh.* xxiv. (1879) 116.—Reg. Argent.
obcuata, *Vog. in Linnæa*, xlii. (1839) 305.—Bras.
odoratissima, *Moric. Pl. Nov. Am.* 76. t. 52.—Bras.
odoratissima, *Newm. ex G. Don, Gen. Syst.* ii. 461.—Bras.
Oliveriana, *Vog. in Linnæa*, xlii. (1839) 305.—Bras.
ornata, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 78.—Burma.
Ouémoulia, Anbl. Fl. Gui. i. 375. t. 144 = guianensis, macrophylla.
ovata, *Vog. in Linnæa*, xlii. (1839) 300.—Bras.
paucinervata, Spreng. Syst. ii. 334 = B. Pauletia.
parviflora, Vahl, Symb. Bot. iii. 55 = racemosa
parvifolia, Hochst. ex Field. & Gardn. Sert. t. 10 = rufescens.
parvifolia, Seem. Bot. Voy. Herald, 113 = B Pauletia.
parvifolia, *Teijsm. & Binn in Tijdschr. Nederl. Ind.* xxix. (1867) 357.—China.
Pauletia, *Pers. Syn.* i. 455.—Panama
pentandra, *Vog. ex D. Distr. Syn. Pl.* ii. 1475.—Bras.
Perrichii, *F. Muell. in Wing, South Sc. Record* (Feb. 1885), 35.—Austral.
Pervilleana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 374.—Madag.
Pes-caprae, *Cav. Ic.* v. 8. t. 404.—Mexic.
Petersiana, *C. Bolle, in Peters, Reise Mossamb. Bot.* i. 24.—Afr. trop.
petiolata, *Triana, ex Hook. Bot. Mag.* t. 6277.—N. Granat.
phoenicea, *Heyne, in Wall. Cat.* n. 5800.—Ind. or.
picta, *DC. Prod.* ii. 515.—N. Granat.
pinnata, *Blanco, Fl. Filip.* 334.—Ins. Philipp.
piperifolia, Roxb. Fl. Ind. ii. 327 = anguina ?
platisiliqua, Stend. Nom. ed. II. i. 195 = reticulata.
platycalyx, *Benth. in Mart. Fl. Bras.* xv. ii. 207.—Bras.
platypetala, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 389.—Mexic.
platyphylla, *Benth. in Mart. Fl. Bras.* xv. ii. 185.—Bras.
platyphylla, Zipp. ex Span. in Linnæa, xv. (1841) 201 = purpurea
platypitiflora, Perr. ex DC. Prod. ii. 515 = reticulata.
Poltesmana, *Vog. in Linnæa*, xlii. (1839) 309.—Guiana.
polyantha, *Vog. l. c.* 310.—Bras.
polyacantha, Stend. Nom. ed. II. i. 192 = polyantha.
polycarpa, *Wall. Cat.* n. 5787.—Burma.
porrecta, *Benth. in Mart. Fl. Bras.* xv. ii. 185.—Bras.
Pottsii, G. Don, Gen. Syst. ii. 463 = ferruginea.
pruinosa, *Vog. in Linnæa*, xlii. (1839) 301.—Bras.
pubescens, DC. Lég. Mém. xiii. 483 = aculeata.
pulchella, *Benth. in Mart. Fl. Bras.* xv. ii. 190.—Bras.
punctata, *Bolle, in Peters, Reise Mossamb. Bot.* i. 23.—Afr. trop.
punctata, *Burch. ex Benth. in Mart. Fl. Bras.* xv. ii. 211.—Bras.
purpurea, DC. ex Walp. Ann. ii. 448 = malabarica.
purpurea, *Linn. Sp. Pl.* 375.—Ind. or. ; Burma ; China.
purpurea, Wall. Cat. n. 5797 = acuminata, variegata.
pyrrhasecura, *Korth. Verh. Nat. Gesch. Bot.* (1839-42) 80.—Sumatra.
pyrrhocarpa, Hochst. in Flora, xxvii. (1844) 99 = reticulata.
racemiflora, *Desf. Journ. Bot.* iii. (1814) 74.—Am. austr.
racemosa, *Lam. Encyc.* i. 390.—Ind. or. ; Malaya ; China.
racemosa, Vahl, Symb. Bot. iii. 56. t. 29 = Vahlii.
Raddiana, *Benz. in Mém. Acad. Pétersb.* Sér. VI. iv. (1836) 111.—Bras.

BAUHINIA :—
radiata, *Vell. Fl. Flum.* 170 ; iv. t. 81.—Bras.
ramosissima, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 389.—Mexic.
reniformis, Royle, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 277 = malabarica.
reticulata, *DC. Lég. Mém.* xiii. 484.—Senegal ; Aegypt.
retusa, Poir. Encyc. Suppl. i. 699 = Lamarckiana, purpurea.
retusa, *Roxb. Hort. Beng.* [31] ; *Fl. Ind.* ii. 322.—Reg. Himal.
rhodacantha, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 499.—Bras.
Richardiana, *DC. Prod.* ii. 517.—Guiana.
Riedeliana, *Bong. in Mém. Acad. Pétersb.* Sér. VI. iv. (1836) 118.—Bras.
Riedelii, *Baker, in Journ. Linn. Soc.* xv. (1877) 98.—Celebes.
riparia, Mart. ex Benth. in Mart. Fl. Bras. xv. ii. 190 = pulchella.
rosea, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 72.—Burma.
rosea, *Mig. in Linnæa*, xviii. (1844) 568.—Surinam.
rotundifolia, Pers. Syn. i. 455 = subrotundifolia.
Rothrightiana, Voigt, Hort. Suburb. Calc. 354 = retusa.
rubescens, Pers. Syn. i. 455 = rufescens.
rubiginosa, *Bong. in Mém. Acad. Pétersb.* Sér. VI. iv. (1836) 112.—Bras.
rufa, *Desv. in Wall. Cat.* n. 5798.—Ind. or.
rufa, *Steud. Nom.* ed. II. i. 193 ; *et D Distr. Syn. Pl.* ii 1476.—Bras.
rufescens, *Lam. Encyc.* i. 391.—Afr. austr. et trop.
rufiearpa, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 480 = phoenicea.
Rutenbergiana, *Vatke, in Bremen Abh.* vii. (1882) 249.—Afr. trop.
rutilans, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 206.—Bras.
scandens, Blanco, Fl. Filip. ed. I. 333 = Cumingiana.
scandens, Burm. f. Fl. Ind. 94 = corymbosa.
scandens, Linn. Sp. Pl. 374 = anguina.
scandens, Roxb. Hort. Beng. [31] ; Fl. Ind. ii. 326 = racemosa, racemosa.
Schlechtendaliana, *Mart. & Gal. in Bull. Brux.* x. (1843) 11. 208.—Mexic.
Sellowiana, Vog. in Linnæa, xlii. (1839) 306 = cujabensis.
semibifida, *Roxb. Hort. Beng.* [31] ; *Fl. Ind.* ii. 330.—Bras ; Malaya.
Serpes, *Choulte & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 20.—Afr. trop.
smilacifolia, *Burch. ex Benth. in Mart. Fl. Bras.* xv. ii. 212.—Bras.
smilacina, *Steud. Nom.* ed. II. i. 192.—Bras.
spathacea, *Seem, ex DC. Prod.* ii. 512.—Mexic.
speciosa, Bonb. in Wall. Cat. n. 5791 = tomentosa.
spicata, Vog. in Linnæa, xlii. (1839) 314 = superba.
spicata, Koen. ex Wall. Cat. n. 5789 = racemosa.
spicata, Kottl. ex Spreng. Fl. Hal. Mant. 39 ; Syst. ii. 333 = racemosa.
spicata, *Vog. in Linnæa*, xlii. (1839) 310.—Hab. ?
spinosa, Poir. Encyc. Suppl. i. 699 = B. Pauletia.
splendens, *H. B. & K. Nov. Gen. et Sp.* vi. 319.—Am. austr.
Sprucei, *Benth. in Mart. Fl. Bras.* xv. ii. 205.—Bras.
stipularis, *Korth. ex Verh. Nat. Gesch. Bot.* (1839-42) 93.—Sumatra.
suaveolens, *H. B. & K. Nov. Gen. et Sp.* vi. 321.—Am. austr.
subcalvata, *Benth. in Mart. Fl. Bras.* xv. ii. 188.—Bras.
subrotundifolia, *Cav. Ic.* v. t. 406.—Mexic.
superba, *Steud. Nom.* ed. II. i. 192.—Guiana.
tamarindacea, Dalla, Voy. *Mérod*, 24 = reticulata.
Teysmanniana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 19.—N. Guinea.
Thonningii, Schum. Beskr. Guin. Pl. 208 = reticulata.
tenella, *Benth. in Mart. Fl. Bras.* xv. ii. 195.—Bras.
tenuis, Span. Ic. 36 ; et in Linnæa, xv. (1841) 202 = racemosa.
tiomarioana, Steud. Nom. ed. II. i. 192 = seq.
timorensis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 466 = racemosa.
tomentosa, *Linn. Sp. Pl.* 375.—Ac. et Afr. trop.
tomentosa, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 72 = acuminata.

BAUHINIA :—
tomentosa, Vell. Fl. Flum. 171 ; iv. t. 84 = Langsdorffiana.
tomentosa, Wall. Cat. n. 5790. F & G = malabarica.
triandra, Roxb. Hort. Beng. [31] ; Fl. Ind. ii. 320 = purpurea.
unguicalyx, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 340.—Mexic.
unguiculata, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 277.—Siam.
Ungula, Jacq. Fragm. 22. t. 15 = aculeata.
Ungula, Willd. ex Steud. Nom. ed. II. i. 192 = Riedemeyeri.
ungulata, *Linn. Sp. Pl.* 374.—' Amer.'
uruguayensis, *Benth. in Mart. Fl. Bras.* xv. ii. 209.—Bras.
Vahlii, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 72 = Cumingiana.
Vahlii, *Wight & Arn. Prod.* i. 297.—Ind. or.
variegata, *Linn. Sp. Pl.* 375.—Ind. or. ; Burma ; China.
velutina, *Wall. Cat.* n. 5781.—Burma.
verrucosa, *Vog. in Linnæa*, xlii. (1839) 311.—Bras.
vesicolor, *Bertol. Hort. Bonon. Pl. Nov.* i. 7.—Mexic.
viridescens, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 499.—Ind. or.
Vogeliana, Steud. Nom. ed. II. i. 192 = spicata.
Welwitschii, *Oliver, Fl. Trop. Afr.* ii. 287.—Afr. trop.
Williamsii, *F. Muell. Descr. Pl. Papuan.* iv. 61.—N. Guinea.
xanthonevra, Korth. ex Miq Fl. Ind. Bat. i. 68 = pyrrhaneura.

BAUKEA, Vatke, in Linnæa, xlii. (1880-82) 104.
LEGUMINOSÆ.
insignis, *Vatke, l. c.*—Madag.
maxima, Baill. in Bull. Soc. Linn. Par. i. (1883) 383 = insignis.

BAUMANNIA, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 582 (Not. Pl. Rar. Jard. Genèv. vi. t. 1, 25) = Damnacanthus, Gaertn. (Rubiac.).
geminiflora, DC. ll. cc. = D. indicus,

BAUMANNIA, Spach, Hist. Vég. Phan. iv. 351 (1835) = Oenothera, Linn. (Onagrar.).
Douglasiana, Spach, l. c. 353 = O. albicaulis.
Nuttalliana, Spach, l. c. = O. albicaulis.
pinnatifida, Spach, l. c. 353 = O. pinnatifida.

BAUMEA, Gandich. in Freyc. Voy. Bot. 416. t. 29 (1826) = Cladium, P. Br. (Cyperac.).
acuminata, Regel, Cat. Pl. Hort. Alsakov. 15.—Hab. ?
arthrophylla, Boeck. in Linnaea, xxxviii. (1874) 243 = C. arthrophyllum.
Brownii, Boeck. l. c. = C. glomeratum.
crassa, Thw. Enum. Pl. Zeyl. 353 = C. riparium.
Deplanchei, Boeck. in Flora, lxi. (1878) 148.—N. Caled.
falcata, Nees, in Hook. Kew Journ. vi. (1854) 29.—Ins. Philipp.
flexuosa, Boeck. in Bremen Abh. vii. (1880) 86.—Madag.
glomerata, Gandich. in Freyc. Voy. Bot. 416. t. 29.—Ins. Moluccc.
viridifolia, Boeck. in Linnaea, xxxviii. (1874) 244.—Madag.
laxa, Boeck. l. c. 245 = C. laxum.
laxiflora, Boeck. l. c. 243 = C. articulatum.
longifolia, Boeck. l. c. 244 = C. Filum.
mariscoides, Gandich. in Freyc. Voy. Bot. 417 = C. mariscoides.
Meyenii, Knuth, Enum. Pl. ii. 314.—Ins. Sandvic.
Preissii, Nees, in Lehm. Pl. Preiss. ii. 75 = C. laxiusculum.
riparia, Boeck. in Linnaea, xxxviii. (1874) 246 = C. riparium.
rubiginosa, Boeck. l. c. 241 = C. glomeratum.
schoenoides, Boeck. l. c. 246 = C. schoenoides.

BAUMGARTENIA, Spreng. Anleit. ii. t. 238 (1817) ; Syst. ii. 91 (1825) = Borya, Labill. (Liliac.).
bacca, Steud. Nom. ed. II. i. 195 = Borya nitida.
nitida, Spreng. Syst. ii. 91 = Borya nitida.
sphaerocephala, Spreng. l. c. = Borya nitida.

BAUMGARTIA, Moench, Meth. 650 (1794) = **Cocculus**, DC. (Menisp.).
scandens, Moench, l. c. = C. carolinus.

BAURSEA, Hort. ex Hoffmgg. Verz. Pfl. 42 (1824) = **Philodendron**, Schott (Aroid.).
glauca, Hoffmgg. l. c.—Hab.?
nervosa, Hort. Angl. ex Hoffmgg. l. c. = P. Karstenianum?

BAURSIA, Schott, Syn. Aroid. i. 73 (1856) = **Philodendron**, Schott (Aroid.).

BAUSCHIA, Scub. ex Warm. in Kjoeb. Vidensk. Meddel. (1872) 123 = **Anellema**, R. Br. (Commel.).
bracteolata, Seub. ex Warm. l. c. 124 = A. ovato-oblongum.

BAUXIA, Neck Elem. iii. 160 (1790) = **Marica**, Ker-Gawl. (Irid.).

BAVERA, Poir. in Dict. Sc. Nat. iv. Suppl. 52 (1816) = **Bauera**, Banks (Saxif.).

BAXTERA, Reichb. Consp. 181 (1828) = **Maradenia**, R. Br. (Asclep.).
lonicerodes, Steud. Nom. ed. II. i. 192 = M. lonicerodes.

BAXTERIA, R. Br. ex Hook. Lond. Journ. Bot. ii. (1843) 494. t. 18–15. *YUNCACEÆ*, Benth. & Hook. f. iii. 866.
australis, *R. Br. l. c.*—Austral.

BAZIASA, Steud. Nom. ed. II. i. 192 (1840) = **Babasia**, Cass. (Compos.).
humilis, Steud. l. c. = S. humilis.
microglossa, Steud. l. c. = S. microglossa.
sarmentosa, Steud. l. c. = S. sarmentosa.
urticæfolia, Steud. l. c. = S. urticæfolia.

BDALLOPHYTON, Eichl. in Bot. Zeit. xxx. (1872) 709. t. 8 = **Cytinus**, Linn.
Andrewsii, Eichl. l. c. 715. t. 8 = C. Andrieuxii.
ceratanthemum, Eichl. l. c. = C. americanus.

BEATONIA, Herb. in Bot. Mag. sub t. 3779 (1839) = **Tigridia**, Ker-Gawl. (Irid.).
atrata, Herb. in Lindl. Bot. Reg. (1843) Misc. 72 = T. atrata.
bracteolata, Klatt, in Linnæa, xxxiv. (1865–66) 733.—Bolivia.
coelestina, Klatt, l. c. xxxi. (1861–62) 567.—Mexic.
curvata, Herb. in Lindl. Bot. Reg. (1843) Misc. 74 = T. curvata.
grandiflora, Klatt, in Abh. Naturf. Ges. Halle, xii. (1873) 269 = T. lutea.
Houttei, Klatt, l. c. = T. Houttei.
lutea, Klatt, in Linnæa, xxxi. (1861–62) 566 = T. lutea.
purpurea, Herb. in Bot. Mag. sub t. 3799 = T. violacea.

BEATSONIA, Roxb. in Beats. St. Hel. App. 800 (1816) = **Frankenia**, Linn.
portulacifolia, Roxb. l. c. = F. portulacæfolia.

BEAUCARNEA, Lem. Illustr. Hortic. viii. (1861) Misc. 59 = **Nolina**, Michx. (Liliac.).
Bigelovii, Baker, in Journ. Bot. x. (1872) 326 = N. Bigelovii.
crumpens, Baker, l. c. = N. crumpens.
gracilis, Lem. Illustr. Hortic. viii. (1861) Misc. 61 = N. Hartwegiana.
Hartwegiana, Baker, in Journ. Bot. x. (1872) 327 = N. Hartwegiana.
Hooberi, Baker, l. c. = Dasylirion Hookeri.
humilis, Baker, in Journ. Linn. Soc. xvii. (1881) 327 = N. humilis.
Lindheimeriana, Baker, in Journ. Bot. x. (1872) 328 = N. Lindheimeriana.
longifolia, Baker, l. c. 324 = N. longifolia.
microcarpa, Baker, in Journ. Linn. Soc. xvii. (1881) 326 = N. microcarpa.
Palmeri, Baker, l. c. 325 = N. Palmeri.
parviflora, Baker, in Journ. Bot. x. (1872) 328 = N. parviflora.
recurvata, Lem. Illustr. Hortic. (1861) Misc. 59 = N. recurvata.
stricta, Lem. l. c. = N. recurvata.

BEAUCARNEA :—
texana, Baker, in Journ. Linn. Soc. xvii. (1881) 326 = Nolina texana.
Watsoni, Baker, l. c. = N. Watsoni.

BEAUFORTIA, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 418 (1812). *MYRTACEÆ*, Benth. & Hook. f. i. 705.
SCHIZOPLEURA, Endl. Gen. 1228 (1840).
anisandra, *Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 13. t. 1.—Austral.
carinata, *A. Cunn. ex Lond. Hort. Brit.* 319.—Austral.
cyrtodonta, *Benth. Fl. Austral.* iii. 167.—Austral.
Dampieri, *A. Cunn. in Hook. Bot. Mag.* t. 3272.—Austral.
decussata, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.
elegans, *Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 16.—Austral.
empetrifolia, Schau. l. c. 17 = micrantha.
heterophylla, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 345.—Austral.
inops, Schau. in Nov. Act. Nat. Cur. xxi. (1844) 17 = Regelia inops.
interstans, *F. Muell. Fragm.* x. 80.—Austral.
macrostemon, *Lindl. Swan Riv. Bot.* 10. n. 31.—Austral.
micrantha, *Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 18.—Austral.
microphylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 24.—Austral.
orbifolia, *F. Muell. Fragm.* iii. 110.—Austral.
pinifolia, Cels, Cat. (1817) 8 = Calothamnus quadrifida.
planifolia, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 345.—Austral.
purpurea, *Lindl. Swan Riv. Bot.* 10. n. 30.—Austral.
Schaueri, *Preiss, ex Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 14.—Austral.
sparsa, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.
spiralaris, Baxter, in Lond. Hort. Brit. Suppl. i. 582 = sparsa.
squarrosa, *Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 11.—Austral.
velutina, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 345.—Austral.

BEAUHARNOISIA, Ruiz & Pav. in Ann. Mus. Par. xi. (1808) 71. t. 9 = **Tovomita**, Aubl. (Guttif.).
fructipendula, Ruiz & Pav. l. c. = T. fructipendula.

BEAUMARIA, Deless. ex Steud. Nom. ed. II. i. 192 (1840) = **Aristotelia**, L'Hérit. (Tiliac.).
Macqui, Deless. ex Steud. l. c. = A. Macqui.

BEAUMONTIA, Wall. Tent. Fl. Nep. 14. t. 7 (1824). *APOCYNACEÆ*, Benth. & Hook. f. ii. 721.
grandiflora, *Wall. l. c.* 15. t. 7.—Ind. or.
Jerdoniana, Wight, *Ic.* iv. t. 1314.—Ind. or.
khasiana, *Hook. f. Fl. Brit. Ind.* iii. 661.—Reg. Himal.
longiflora, Hook. f. l. c. 660, in syn. = grandiflora.
longifolia, Lodd. ex Lond. Hort. Brit. 87; Hook. Bot. Mag. sub t. 3213 = grandiflora.
multiflora, Teijsm. & Binn. in Nederl. Kruidk. Arch. iii. (1853) 392 = khasiana.
Wallichii, Walp. Ann. iii. 89 = Wrightia tomentosa.

BEAUMULIX, Willd. ex Poir. in Dict. Sc. Nat. iv. Suppl. 54 (1816) = **Beaumuria**, Linn. (Tamariscin.).

BEAUPREA, Brongn. & Gris. in Bull. Soc. Bot. Fr. xviii. (1871) 248. *PROTEACEÆ*, Benth. & Hook. f. iii. 375.
Balansæ, *Brongn. & Gris, l. c.* 246.—N. Caled.
diversifolia, *Brongn. & Gris, l. c.* 244.—N. Caled.
gracilis, *Brongn. & Gris, l. c.* 243.—N. Caled.
Pancheri, *Brongn. & Gris, l. c.* 245.—N. Caled.
spathulæfolia, *Brongn. & Gris, l. c.* 244.—N. Caled.

BEAUTEMPSIA, Gaudich. Voy. Bonite, t. 56 (1844 ?) = **Capparis**, Linn.
asiaticæfolia, Gaudich. l. c.—Hab.?

BEBBIA, Greene, in Bull. Calif. Acad. i. (1885) 179. *COMPOSITÆ*.
striplicifora, *Greene, l. c.* 181.—Calif.
juncea, *Greene, l. c.* 179.—Calif.

BECCABUNGA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophul.).
Anagallis, Fourr. l. c. = V. Anagallis.
vulgaris, Fourr. l. c. = V. Beccabunga.

BECHIUM, DC. Prod. v. 70 (1836) = **Wernonia**, Schreb. (Compos.).
foliosum, Klatt, in Linnæa, xxxvii. (1871–73) 508.—Madag.
rubricaule, DC. Prod. v. 70 = V. Candollei.
scopiformis, DC. l. c. 71.—Madag.

BECHONNERIA, Hort. ex Carr. in Rev. Hortic. (1867) 320 = **Beschorneria**, Kunth (Amaryll.).

BECKEA, Pers. Syn. i. 284 (1805) = BAECKEA, Burm. f. = **Brunia**, Linn.

BECKEA, St. Hil. in Dict. Sc. Nat. iv. 205 (1804) = **Baeckea**, Linn. (Myrtac.).

BECIUM, Lindl. Bot. Reg. (1842) Misc. 42 = **Ocimum**, Tourn. (Labiat.).
bicolor, Lindl. l. c. = O. filamentosum.

BECKERIA, Fresen. in Mus. Senckenb. ii. (1837) 182. t. 8. *GRAMINEÆ*, Benth. & Hook. f. iii. 1117.
dieica, Heynh. Nom. ii. 68 = Panicum adoense.
glabrescens, *Steud. Syn. Pl. Gram.* 117.—Abyssin.
gracilis, *Hochst. in Flora*, xxxviii. (1855) 199.—Abyssin.
mutica, *Hochst. l. c.* xxvii. (1844) 514.—Abyssin.
nubica, *Hochst. l. c.* 512.—Abyssin.
petiolaris, *Hochst. l. c.*—Abyssin.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 358.—Abyssin.
polystachia, *Fresen. in Mus. Senckenb.* ii. (1837) 182.—Abyssin.
Schimperi, *Hochst. in Flora*, xxvii. (1844) 512.—Abyssin.
unisaria, Steud. Syn. Pl. Gram. 118 = Penniscum unisetum.

BECKERIA, Bernh. Syst. Verz. Erf. 20 (1800) = **Melica**, Linn. (Gramin.).
ciliata, Bernh. l. c. 40 = M. ciliata.

BECKERIA, Heynh. Nom. ii. 68 (1846) = **Beckera**, Fresen. (Gramin.).

BECKEROPSIS, Fig. & De Not. in Mem. Acc. Torin. Ser. II xiv. (1854) 365 = **Pennisetum**, Pers. (Gramin.)
nubica, Fig. & De Not. l. c. 366.—Nubia.
petiolaris, Fig. & De Not. l. c. 368.—Abyssin.

BECKIA, Rafin. Sylva Tellur. 103 (1838) = **Osbeckia**, Linn. (Melast.).

BECKMANNIA, Host, Gram. Austr. iii. 5. t. 6 (1805). *GRAMINEÆ*, Benth. & Hook. f. iii. 1099.
BRUCKMANNIA, Nutt. Gen. Am. i. 48 (1818).
JOACHIMIA, Tenore, ex Roem. & Schult. Syst. ii. 695 (1817).
erucæformis, *Host, l. c.*—Europ.; As. temp.; Am. bor.
erucoides, Beauv. Agrost. t. 19. f. 6 = erucæformis.

BECLARDIA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 69 = Cryptopus, Lindl. (Orchid.).
brachystachya, A. Rich. l. c. 71 = Aeonia brachystachya.
elata, A. Rich. l. c. 69 = C. elatus.
macrostachya, A. Rich. l. c. 70 = Aeonia macrostachya.

BECQUERELA, Ness, in Linnæa, ix. (1834) 304 = seq.

BECQUERELIA, Brongn. in Duperr. Voy. Coq. Bot. Phaner. 161. t. 37 (1829). *CYPERACEÆ*, Benth. & Hook. f. iii. 1069.
cymosa, *Brongn. l. c.*—Bras.
cymosa, Kunth, Enum. Pl. ii. 361 = Merkeliana.
discolor, *Kunth, l. c.* 362.—Bras.
gigantea, *Hochst. ex Steud. Syn. Pl. Cyp.* 180.—Guiana.
glomerata, Kunth, Enum. Pl. ii. 364 = seq.

BECQUERELIA :—
glomerulata, Brongn. in Duperr. Voy. Coq. Bot. Phan. 168 = Calyptrocarya Palmetto.
Martiolana, *Nees*, in *Mart. Fl. Bras.* ii. 1. 191.—Bras.
muricata, *Nees*, *l. c.* 192.—Bras.
verruculosa, *Nees*, *l. c.* 191.—Bras.

BEDDOMEA, Hook. f. in Benth. & Hook. f. Gen. i. 886 (1862). *MELIACEÆ*, Benth. & Hook. f. l. c.
indica, *Hook. f. l. c.*—Ind. or.
simplicifolia, *Bedd. Fl. Syle.* i. t. 135.—Ind. or.

BEDFORDIA, DC. in Guillem. Arch. Bot. ii. 332 (1838). *COMPOSITÆ*, Benth. & Hook. f. ii. 460.
linearis, *DC. l. c.* 333 ; *et Prod.* vi. 441.—Tasmania.
salicina, *DC. ll. cc.*—Austral.

BEDOUSIA, Desnst. Schluss. Hort. Malab. 31 (1818) = **Casearia**, Jacq. (Samyd.).
aromatica, Rafin. Sylva Tellur. 11 = C. tomentosa.
malabarica, Desnst. Schluss Hort. Malab. 31 = C. tomentosa.

BEDUSIA, Rafin. Sylva Tellur. 11 (1838) = praec.

BEEHSA, Endl. Gen. 1354 (1841) = BEESHA, Kunth = **Bambusa**, Linn. (Gram.).

BEERA, Beauv. in Lestib. Essai Fam. Cyp. 43 (1819) = **Hypolytrum**, Rich. (Cyperac.).
carpitosa, Schrad. ex Kunth, Enum. Pl. ii. 270 = H. strictum.
laxa, Schrad ex Kunth, l. c. = H. laxum.
silvatica, Schrad. ex Nees, in Mart. Fl Bras. ii. t. 67 = H. fuscum.

BEESHA, Kunth, Syn. i. 258 (1822) = **Bambusa**, Linn. (Gram.).
baccifera, Kunth, l. c. = Melocanna bambusoides.
Fax, Schult. f. in Roem. & Schult. Syst. vii. 1336 = Melocanna humilis.
humilis, Kunth, Enum. Pl. i. 434 = Melocanna humilis.
Rheedii, Kunth, Rev. Gram. i 141 = Melocanna bambusoides.
travancorica, Bedd. Forest. Man. Bot. 234 = Melocanna travancorica.

BEESHA, Munro, in Trans. Linn. Soc. xxvi. 144 (1868) = **Oohlandra**, Thw. (Gram.).
capitata, Munro, in Trans. Linn. Soc. xxvi. (1868) 145. —Madag.
elegantissima, Kurz, ex Munro, l. c 146.—Java.
stridula, Munro, l. c. 145 = O. stridula.

BEETHOVENIA, Endl. in Linnæa, xxxiii. (1864–65) 677 = Ceroxylon, Humb. (Palm.).
cerifera, Engl. l. c. = C. andicola.

BEFARIA, Mutis, ex Linn. Mant. ii. 152 (1771). *ERICACEÆ*, Benth. & Hook. f. ii. 599.
ACUNNA, Ruiz & Pav. Prod. 69. t. 12 (1794).
BEJARIA, Zea, in Anal. Clenc. Nat. iii. (1801) 151, in nota ; Vent. Jard. Cels (1800), sub t. 51.
JURGENSENIA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) *t.* 151.
asetuana, *Linn. Mant.* ii. 242.—N. Granat.
caxamarcensis, *H. B. & K. Nov. Gen. et Sp.* iii. 294.—Peruv.
cinnamomea, *Lindl. in Gard. Chron.* (1848) 175.—Peruv.
coarctata, Hook. Bot. Mag. t. 4483 = Lindeniana.
coarctata, *Humb. & Bonpl. Pl. Aequin.* ii. 125. t. 121. —Peruv.
cubensis, *Griseb. Cat. Pl. Cub.* 59.—Cuba.
densa, *Planch. ex Hérincq*, in *Rev. Hortic.* Sér. III. iv. (1850) 185.—Hab.?
dentiniana, *Remy*, in *Ann. Sc. Nat.* Sér. III. viii. (1847) 290.—Bolivia.
discolor, *Benth. Pl. Hartw.* 65.—Mexic.
dryanidia, *Linden*, ex *Hérincq*, in *Rev. Hortic.* Sér. III. iv. (1850) 185.—N. Granat.
floribunda, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 544 = discolor.
glabra, Mart. & Gal. l. c. = laevis.
glauca, *Humb. & Bonpl. Pl. Aequin.* ii. 118. t. 117.—N. Granat.

BEFARIA :—
grandiflora, *Humb. & Bonpl. Pl. Aequin.* ii. 222. t. 119.—Ecuador.
guyanensis, *Klotzsch*, in *Schomb. Fauna et Fl. Guy.* 1088.—Guiana.
hispida, *Koepp. & Endl. Nov. Gen. et Sp.* i. 24. t. 39. —Chili.
laevis *Benth. Pl. Hartw.* 65.—Mexic.
lancolata, *Ruiz & Pav. Fl. Per.* iv. t. 413. f. b.—Peruv.
leolicifolia, *Humb. & Bonpl. Pl. Aequin.* ii. 124. t. 120. —N. Granat
Lindeniana, *Hérincq*, in *Rev. Hortic.* Sér. III. iv. (1850) 181.—Venezuela.
mexicana, *Benth. Pl. Hartw.* 15. n. 92.—Mexic.
Mathewsii, *Field. & Gardn. in Bot. Mag.* t. 4981.—Peruv.
microphylla, Paxt. & Lindl. Flow. Gard. i. 86 = densa.
myrtifolia, *Hérincq*, in *Rev. Hortic.* Sér. III. iv. (1850) 184.—N. Granat.
oblonga, *Ruiz & Pav. Fl. Per.* iv. t 413 ; *Pers. Syn.* ii. 8.—Venezuela.
pallens, *Remy*, in *Ann. Sc. Nat.* Sér. III. viii. (1847) 289.—Peruv.
paniculata, Michx. Fl. Am. Bor. i. 280 = racemosa.
parviflora, *Benth. Pl. Hartw.* 143.—Peruv.
phyllireaefolia, *Benth. l. c.* 225.—N. Granat.
racemosa, *Vent. Jard. Cels*, t. 51.—Am. bor.
resinosa, *Linn. f. Suppl* 246.—N. Granat.
Schomburgkiana, *Klotzsch*, in *Schomb. Fauna et Fl. Guy.* 1088.—Guiana.
Spruceі, *Meissn. in Mart. Fl. Bras.* vii. 171.—Peruv
subsessilis, *Benth. Pl. Hartw.* 143.—Peruv.
tricolor, *Linden*, ex *Hérincq*, in *Rev. Hortic.* Sér. III. iv. (1850) 185.—N. Granat.

BEGONIA, [Tourn.] Linn Gen. ed. II. 516 (1742). *BEGONIACEÆ*, Benth & Hook. f. i. 841
AUGUSTIA, BARYA, CASPARYA, CYATHOCNEMIS, DONALDIA, DORATOMETRA, EWALDIA, GAREIDIA, GIREOUDIA, GURLTIA, HAGGEA, HUSSIA, ISOPTERIX, ISOPTERYX, KNESEBECKIA, LAUCHEA, LEPSIA, MAGNUSIA, MITSCHERLICHIA, MOSCHKOWITZIA, NEPHROMISCHUS, PETERMANNIA, PILDERIA, PLATYCENTRUM, PRITZELIA, PUTZEYSIA, RACHIA, REICHENHEIMIA, REISSIA, ROSSMANNIA, SASSEA, SAUERIA, SCHMIDWEILERIA, STEINERIA, STIRADIOPTERA, TITTELBACHIA, TRACHELANTHUS, TRENDELENBURGIA, WAGENERIA, et WEILBACHIA, Klotzsch, in Abh. Akad. Berl. (1854) 17–134 tt. 1–12.
CLADOMISCHUS, Klotzsch, ex DC. l. c. 378 (1864).
DIPLOCLINIUM, Lindl. Veg. Kingd. 819 (1847).
EUPETALUM, Lindl. Introd. Nat. Syst. ed. II. 440 (1836).
FALKEA, Koen. ex Stend. Nom. ed. II. i. 194, 625 (1840).
IRMA, Bonst. ex A. DC. Prod. xv. i. 406 (1864).
MITCHERLICHIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 124.
MEZIERRA, Gaudich. Voy. Bonite, t. 82 (1839).
PLATYCLINIUM, T. Moore, in Moore & Ayres, Gard. Mag. Bot. ii. (1850) 153.
SPHENDANTHERA, Hassk. in Veral. Kon. Akad. Wetensch. ix. (1855) 139.
STIRADOTHECA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127.
TRACHELOCARPUS, C. Muell. in Walp. Ann. iv. 909 (1858).
TRITOMISA, Rafin. Fl. Tellur. ii. 91 (1836).
WAGNERIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 126.
acerifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 186.—Ecuador.
acetosa, *Vell. Fl. Flum.* x. t. 50.—Bras.
acida, *A. DC. Prod.* xv. i. 364.—Bras.
acida, Vell. Fl. Flum. x. t. 49 = praec. ?
acomitifolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 197.—Bras.
aculeata, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 409 = maculata.
acuminata, *Dryand. in Trans. Linn. Soc.* i. (1791) 166. t. 14.—Jamaica.
acutifolia, Balb. ex A. DC. Prod. xv. i. 295 = portoricensis.
acutifolia, Jacq. Coll. i. 123 = acuminata.
acutifolia, Herb. Vindob. ex A. DC. Prod. xv. i. 378 = dentatiloba.

BEGONIA :—
acutiloba, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 14.—Mexic.
adenopoda, Lem. Jard. Fleur. ii. Misc. 17 = verticillata.
aequata, *A. Gray, Bot. U. St. Expl. Exped.* i. 658.—Ins. Luzon.
alaecida, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 637.—Burma.
albido-setulosa, *Hassk. Cat Hort. Bog. Alt.* 318 = Hab.?
albiflora, *Hort. ex Klotzsch*, in *Monatsb. Berl. Acad* (Maerz 1854) 126, nomen.—Hab.?
albo-coccinea, *Hook. Bot. Mag.* t. 4172.—Ind. or.
alchemilloides, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi (1859) 126.—Bras.
almifolia, *A. DC. l. c.* 182.—N. Granat.
alto-peruviana, *A DC. l. c.* 123.—Peruv.
amabilis, *Linden, Hort. Linden.* (1859) 1.—Assam.
amoena, *Wall. Cat. n.* 3682.—Neg. Himal
ampla, *Hook f. in Oliver, Fl. Trop. Afr.* ii. 574.—Afr. trop.
anamalaiensis, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 48.—Ind. or.
anamalayana, Bedd in Trans. Linn. Soc. xxv. (1866) 217 = prsec.
andamensis, *Parish, ex C. B. Clarke, in Hook. f. Fl. Brit Ind.* ii. 650.—Ind. or.
anemonoides, *Asara, ex Steud. Nom.* ed. II. i. 198.—Hab. ?
angularis, *Rhdsi, in Mem. Mod.* xviii. *Fis.* (1820) 407. —Bras.
angulata, *Vell Fl. Flum.* x. t. 52.—Bras.
angustiloba, *Blume, Enum. Pl. Jav.* 97.—Java.
angustiloba, *A. DC in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126 —Mexic.
anisosepala, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 576. —Afr. trop.
anuobonensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 127.—Afr. trop.
annulata, *C. Koch, in Berl. Gartenz.* i. (1857) 76 — Reg. Himal.
anodaefolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 198.—Mexic.
apterа, *Blume, Enum. Pl. Jav.* 97.—Ins. Celebes.
apterа, *Roxb. Hort. Beng* 104 ; *Fl. Ind.* iii. 650.—Ins. Mascar. ; Timor. ; Ins. Philip.
arborescens, *Radd.*, in *Mem. Mod.* xviii. *Fis.* (1820) 408.—Bras.
areolata, *Miq. Pl. Junghh.* 417.—Java.
argentea, Van Houtte, ex A. DC. Prod. xv. i. 318 = maculata.
argyrocoelis, *Hérincq, in Rev. Hortic.* Sér. III. i. (1847) 111, 165.—Hab.?
argyrostigma, Fisch. ex Link & Otto, Ic. Sel. i. 28. t. 10 = maculata.
Arnottiana, A. DC. Prod. xv. i. 392 = cordifolia.
asarifolia, Liebm. in Kjoeb. Videnak. Meddel. 1852 (1853) 8 = hydrocotilifolia.
ascotiensis, *Y. B. Weber, in Rev. Hortic.* (1874) 17.—Hab. ?
asplenifolia, *Hook. f. ex A. DC. Prod.* xv. i. 292.—Afr. occ.
atrocha, *Miq. ex A. DC. l. c.* 321.—Sumatra.
attenuata, *A. DC. l. c.* 380.—Bras.
attenuata, Klotzsch, ex Regel, Ind. Sem. Hort. Petrop. (1866) 101 = hederacea.
aucubaefolia, Hort. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 129 = incarnata.
aurantiaca, Hort. ex Planch. in Fl. des Serres, Sér. I. v. (1849) 530 = cinnabarina.
auriculata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 577.—Afr. trop.
auriformis, Hort. Berol. ex Klotzsch, in Abh. Akad. Berl. (1855) 38 = incana.
baccata, *Hook. Bot. Mag.* t. 5554.—Afr. trop.
bahiensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 187.—Bras.
balmisiana, *Ruiz, ex Klotzsch, in Abh. Akad. Berl.* (1855) 38.—Mexic.
barbata, *Wall. Cat.* n. 3679, A. & B. partim.—Ind. or. ; Burma.
barbata, Wall. l. c. = rubro-venia.
Barkeri, *Knowl. & Westc. Flor. Cab.* iii. 179. t. 185.—Mexic.
Beddomei, *Hook. f. Bot. Mag.* t. 6767.—Assam.
Bernieri, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Madag.

BEGONIA :—

bederiaefolia, *Schott, in Spreng. Syst.* iv. 403.—Bras.
bedentata, *Raddi, in Mem. Mod.* xviii. *Pis.* (1820) 408.—Bras.
bipetala, *Ledd. ex Otto & Dietr. Allg. Gartenz.* ix. (1841) 86.—Ind. or.
biserrata, *Lindl. in Journ. Hort. Soc.* ii. (1847) 313.—Guatemala.
Boissieri, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 196.—Mexic.
Bolviniana, *A. DC. l. c.* 130.—Madag.
boliviensis, *A. DC. l. c.* 122.—Bolivia.
bomblycina, Blume, Enum. Pl. Jav. i. 97 = inopters.
bonnensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Borneo.
Bonchenna, *A. DC. Prod.* xv. i. 372.—Venezuela.
Bowringiana, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 120 = laciniata.
bractenta, *Yack, in Malay. Misc.* ii. (1822) VII. 18.—Sumatra.
bracteosa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 132.—Peruv.
Brandtiana, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 58; *et in Flora*, liv. (1871) 395.—Burma.
brasilia, A. DC. in Mém. Soc. Phys. Genèv. vii. (1836) 295 = hirtella.
brasiliensis, Schrank, ex Steud. Nom. ed. II. i. 198 = prsec.
brasiliensis, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 155, nomen.—Bras.
brevicaulis, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi. (1859) 134.—Reg. Himal.
Bridgesii, *A. DC. l. c.* 132.—Bolivia.
Brongniartiana, *Lem. Hort. Univ.* v. 256.—Hab. ?
brousonetiæfolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 139.—Guatemala.
buddleiæfolia, *A. DC. l. c.* 141.—Peruv.
bulbifera, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 155 = discolor.
bulbifera, Ledd. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 88 = bulbillifera.
bulbillifera, *Link & Otto, Ic. Pl. Rar.* 89. t. 45.—Mexic.
bulbillifera, Moric. ex A. DC. Prod. xv. i. 310 = gracilis.
caespitosa, *Yack, in Malay. Misc.* ii. (1822) VII. 8.—Sumatra.
cafra, Meisn. in Linnæa, xiv. (1840) 501 = Dregei.
canarana, *Miq. Anal. Bot. Ind.* iii. 18.—Ind. or.
candicans, A. DC. Prod. xv. i. 400, lapsu, *vide* Bisgnonia candicans.
capensis, Blanco, Fl. Filip. ed. I. 724 = rhombicarpa.
capensis, Linn. f. Suppl. 309 = dipters.
cardiocarpa, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 18.—Nicaragua.
carminata ×, *Hort. Fl. Mag* t. 551 (err. typ. 451).
carnosa, *Teijsm. & Binn. Epim. Lugd. Bat.* (1863) 4.—Malaya.
caroliniæfolia, *Regel, Gartenfl.* i. (1852) 250. t. 25.—Mexic.
carpinifolia, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 20.—Costa Rica.
castaneæfolia, *Otto & Dietr. Allg. Gartenz.* iv. (1836) 356.—Bras.
castaneæfolia, Schott, in Spreng. Syst. iv. App. 407 = fruticosa.
Cathcartii, *Hook. f. & Thoms. Illustr. Pl. Himal.* t. 18.—Reg. Himal.
ciliata, Hort. ex Steud. Nom. ed. II. i. 198, 194 = insignis.
ciliata, H. B. & K. Nov. Gen. et Sp. vii. 186 = hirtella.
ciliata, Pav. ex A. DC. Prod. xv. i. 331 = subciliata.
cinnabarina, *Hook. Bot. Mag.* t. 4488.—Bolivia.
circumlobata, *Hance, in Journ. Bot.* xxi. (1883) 203.—China.
clandestina, *Baker, in Journ. Bot.* xx. (1882) 113.—Madag.
Clarkei, *Hook. f. Bot. Mag.* sub t. 5663 et t. 5675.—Bolivia.
clypeifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 576.—Afr. trop.
coccinea, *Hook. Bot. Mag.* t. 3990.—Bras.
coccinea, *Raiz, ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 127; *A. DC. Prod.* xv. i. 373.—Peruv.
colcmanaris, *Benth. Pl. Hartw.* 181.—Peruv.

BEGONIA :—

columnaris, Ruiz, ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127 = cordifolia.
complicata, *A. DC. Prod.* xv. i. 299.—Hab. ?
concanensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 136.—Ind. or.
concholifolia, *A. Dietr. Allg. Gartenz.* xix. (1851) 268.—Costa Rica.
concinna, Schott, in Spreng. Syst. iv. App. 408 = peltata ?
confertiflora, Gardn. in Hook. Lond. Journ. Bot iv. (1845) 134 = arborescens.
convolvulacea, *A. DC. in Mart. Fl. Bras.* iv. i. 367.—Bras.
corallina, *Carr. in Rev. Hortic.* (1875) 89.—Bras.
cordata, *Vell. Fl. Flum.* x. t. 84.—Bras.
cordifolia, *Thw. Enum. Pl. Zeyl.* 129.—Ind. or.
coriacea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 122.—Bolivia.
coriacea, Hassk. Cat. Hort Bogor. Alt. 311 = peltata.
crassicaulis, *Lindl. in Bot. Reg.* (1842) *Misc.* 22.—Guatemala.
crenata, *Dryand. in Trans. Linn. Soc.* i. (1791) 162. t. 14. f. 3.—Ind. or.
crenata, Herb. Finlays. ex A. DC. Prod. xv. i. 401 = prolifera ?
crenatifera, *A. DC. Prod.* xv. i. 306.—Mexic.
crenatifolia, Hemsl. Biol. Centr. Am. Bot. i. 494 = crenatifera.
crenulata, Schott, ex A. DC. in DC. Prod. xv. i. 358 = sagittaria.
crinita, *Oliver, in Hook. Bot. Mag.* t. 5897.—Peruv.
cruenta, *R. Grah. ex Steud. Nom.* ed. II. i. 193.—Bras.
cubensis, Hassk. Cat. Hort. Bogor. Alt. 322 = Wrightiana.
cubincola, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 131.—Cuba.
cucullata, Ruiz, ex A. DC. Prod. xv. i. 374 = columnaris.
cucullata, *Willd. Sp. Pl.* iv. 414.—Bras.
cucullifera, Hassk. Cat. Hort. Bogor. Alt. 311 = semperflorens.
Cumingii, A. Gray, Bot. U. St. Expl. Exped. i. 558 = philippinensis.
Cummingiana, *A. DC. Prod.* xv. i. 320.—Ins. Fiji.
cuneata, Walp. Rep. ii. 214 = spathulata.
cyathophora, *Poepp. & Endl. Gen. et Sp.* i. 7. t. 14.—Peruv.
daedalea, Lem. Illustr. Hortic. vii. (1860) 54. t. 269 = strigillosa.
dasyrana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 127.—Bras.
dasypoda, Meisn. ex A. DC. Prod. xv. i. 299 = hirtella.
Dawsoniana, Godefr. in Rev. Hortic. (1880) 390 = Pellionia Daveesiana.
Davisii, *Veitch, in Hook. f. Bot. Mag.* t. 6252.—Peruv.
dealbata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 5.—Mexic.
decandra, *Pav. ex A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 129.—Mexic.
declinata, *Vell. Fl. Flum.* x. t. 44.—Bras.
delicatula, *Parish, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 652.—Burma.
dentata, Pav. ex A. DC. Prod. xv. i. 307 = angustiloba.
dentatiloba, *A. DC. in Ann. Sc. Nat. Sér.* IV. xi. (1859) 140.—Bras.
denticulata, *H. B. & K. Nov. Gen. et Sp.* vii. 182.—Venezuela.
depauperata, Schott, in Spreng. Syst. iv. App. 408 = Olfersiana.
dervyeiana, Lem. Hort. Univ. (1845-44) Misc. 855, fide A. DC. Prod. xv. i. 343 = nelumbiifolia.
Dielsiana, *Lindau, in Illustr. Hortic.* xxix. (1882) t. 446.—Borneo.
dichotoma, *Yacq. Coll.* iii. 250.—Venezuela.
digitata, *Raddi, in Mem. Mod.* xviii. *Pis.* (1820) 406.—Bras.
Digwelliana, *Carr. in Rev. Hortic.* (1876) 450.—Hab. ?
dimidiata, Vell. Fl. Flum. x. t. 46 = arborescens.
dioica, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 222 = scutata ?
diptenia, R. Grah. in Bot. Mag. t. 3849 = malabarica.
diptera, *Dryand. in Trans. Linn. Soc.* i. (1791) 170.—Bras.

BEGONIA :—

discolor, [R. Br. in] Ait. Hort. Kew. ed. II. v. 164 = Evansiana.
disticha, Hort. Berol. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 132 = suaveolens.
distichoa, Link, Enum. Hort. Berol. ii. 296 = stipulacea.
divaricata, *Hook. f. Bot. Mag.* t. 6252.—Peruv.
diversifolia, R. Grah. in Edinb. N. Phil. Journ. (Jan.-Apr. 1830) 188 = gracilis.
domingensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 194.—Ins. S. Doming.
dominicalis, *A. DC. Prod.* xv. i. 366.—Ins. Dominica.
Donkelaariana, *Lem. Jard. Fleur.* i. (1851) *Misc.* 84.—Mexic.
Dregei, *Otto & Dietr. Allg. Gartenz.* iv. (1836) 357.—Bras.
dubia, *Vell. Fl. Flum.* x. t. 42.—Bras.
Dux, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 637.—Burma.
echinata, Royle, Illustr. Bot. Himal. 318. t. 80. fig. 1 = picta.
echinosepala, *Regel, in Act. Hort. Petrop.* i. (1871) 91.—Bras.
elaeagnifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 579.—Afr. trop.
elata, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 129.—Bras.
elatior, Hort. ex Steud. Nom. ed. II. i. 198 = Dregei.
elatostemmoides, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 576.—Afr. trop.
elegans, *H. B. & K. Nov. Gen. et Sp.* vii. 182.—N. Granat.
elliptica, H. B. & K. l. c. 188 = scandens.
elongata, Wall. Cat. n. 6391 = sinuata.
Epibaterium, *Mart. ex A. DC. in Mart. Fl. Bras.* iv. i. 363.—Bras.
epiphytica, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 580.—Afr. trop.
episcopalis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 644.—Reg. Himal.
erecta, *Vell. Fl. Flum.* x. t. 45.—Bras.
eriocaulis, Vis. Orto Padov. (1842) 135 = tomentosa.
eriocaulon, Neumann, in Rev. Hortic. Sér. III. i. (1847) 166 = tomentosa ?
Ernesti, Klotzsch, in Abh. Akad. Berl. (1855) 36 = ulliginosa.
erminea, *L'Herit. Stirp. Nov.* 97. t. 47.—Madag.
eroso, *Blume, Enum. Pl. Jav.* i. 96.—Java.
eroso, Wall. Cat. n. 3688, partim = picta, amoena.
erythrocarpa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—Bras.
erythrophylla, *Neuman, in Rev. Hortic.* Sér. III. i. (1847) 166.—Hab. ?
Evansiana, *Andr. Bot. Rep* t. 627.—Malaya; China; Japoc.
excelsa, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 571.—Afr. trop.
eximia, *Hort. ex Lem. Illustr. Hortic.* vii. (1860) t. 238.—Hab. ?
fagifolia, *Yack, in Otto & Dietr. Allg. Gartenz.* iv. (1836) 356.—Bras.
fagopyrolden, *Kunth & Bouché, Ind. Pl. Hort. Berol.* (1848).—Venezuela.
falcifolia, *Hook. f. Bot. Mag.* t. 5707.—Afr. trop.
falciloba, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 15.—Mexic.
fallax, *A. DC. Prod.* xv. i. 329.—Ind. or.
fasciculata, *Yack, in Malay. Misc.* ii. (1822) VII. 12.—Sumatra.
ferruginea, *Linn. f. Suppl.* 419.—N. Granat.
fibrosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 652.—Burma.
filipes, *Semtk. Bot. Voy. Sulph.* 101.—Panama.
fimbriata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 18.—Mexic.
fimbristipula, *Hance, in Journ. Bot.* xxi. (1883) 202.—China.
Finlaysoniana, Wall. Cat. n. 3684 = prolifera ?
Fischeri, *Otto & Dietr. Allg. Gartenz.* iv. (1836) 354.—Bras.
Fischeri, Schrank, Pl. Rar. t. 89 = macroptera.
flaccidissima, *Kurz, in Journ. As. Soc. Beng.* xl. (1872) ii. 306.—Burma.
flexuosa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 142.—Panama.
flocciflora, *Bedd. Ic. Pl. Ind. Or.* i. 28.—Ind. or.
floribunda, *Carr. in Rev. Hortic.* (1875) 450.—Hab. ?

BEGONIA :—
foliosa, *H. B. & K. Nov. Gen. et Sp.* vii. 188. t. 642 —N. Granat.
foliosa, Poepp. ex A. DC. Prod. xv. i. 376 = Poeppigiana.
Francoana, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 21.—Mexic.
frigida, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 148.—Hab. ?
Frobelii, *A. DC. in Gard. Chron.* (1874) ii. 552.— Ecuador.
fruticosa, *A. DC. in Mart. Fl. Bras.* iv. i. 377.— Bras.
fusca, *Liebm. ex Klotzsch, in Abh. Akad. Berl.* (1855) 89.—Bras.
fuchsioides, *Hook. Bot. Mag.* t. 4981.—Mexic.
furfuracea, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 571.— Afr. trop.
fusca, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 7.—Mexic.
Galeottii, Hort. Berol. ex Klotzsch, in Abh. Akad. Berl. (1855) 39 = lobata.
Galeottiana, *Lem. Jard. Fleur.* i. (1851) *Misc.* 14.— Bras.
Gardneri, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 188.—Bras.
Gaudichaudii, *Walp. Rep.* v. 769.—Peruv.
gemmipara, *Hook. f. & Thoms. Illustr. Him. Pl.* t. 14. —Reg. Himal.
geniculata, Vell. Fl. Flum. x. t. 51 = convolvulacea.
geniculata, Jack, in Malay. Misc. (1822) vii. 15 = isoptera.
geranifolia, *Hook. Bot. Mag.* t. 3357.—Peruv.
geranioides, *Hook. f. t. c.* t. 5585.—Afr. austr.
gigantea, *Wall. Cat.* n. 3677.—Reg. Himal.
gigantea, Wall. l. c. n. 3677 B = silhetensis.
glabra, Aubl. Pl. Gui. ii. 916 = scandens.
glabra, *Kuntz, ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 128.—Peruv.
glandulifera, *Griseb. Fl. Brit. W. Ind.* 304.—Ind. occ.
glandulosa, *A. DC. Prod.* xv. i. 339.—Am. centr.
glauca, *Ruiz & Pav. ex A. DC. l. c.* 330.—Peruv.
goegoensis, *N. E. Br. in Gard. Chron.* (1882) ii. 71.— Sumatra.
goniotis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 646.—Burma.
Goudotii, *A. DC. in Ann Sc. Nat.* Sér. IV. xi. (1859) 130.—Madag.
gracilis, *H. B. & K. Nov. Gen. et Sp.* vii. 184.— Mexico.
gracillima, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Peruv.
Grahamiana, Wight, Ic. t. 1811 = albo-coccinea.
grandiflora, Jacq. Coll. i. 198 = dipetera.
grandiflora, Knowl. & West. Flor. Cab. i. (1837) 51 = octopetala.
grandis, Dryand. in Trans. Linn. Soc. i. (1791) 163 = Evansiana.
grandis, Otto, ex A. DC. in Mart. Fl. Bras. iv. i. 369 = vitifolia.
Griffithii, *Hook. Bot. Mag.* t. 4984.—Reg. Himal.
grisea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 188.—Bras.
guadnensis, *H. B. & K. Nov. Gen. et Sp.* vii. 178.— N. Granat.
guineensifolia, *Linden & André, Illustr. Hortic.* xxii. (1875) 106. t. 212.—N. Granat.
guttata, *Wall. Cat.* n. 3671 A.—Malaya.
guttata, Wall. l. c. n. 3671 B = sinuata.
gymnomela, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 142.—Guiana.
haematotricha, Brâm. ex A. DC. Prod. xv. i. 298 = humilis.
Hamiltoniana, Lehm. in E. Otto, Hamb. Allgem. Gartenz. vi. (1850) 456 = acuminata.
Hasskarli, *Zoll. & Mor. Syst. Verz. Zoll.* 31.—Java.
Hasskarliana, *Miq. ex A. DC. Prod.* xv. i. 339.— Sumatra.
hastata, Vell. Fl. Flum. x. t. 54 = angularis.
Hatacoa, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 232. —Nepal.
hederacea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 150.—N. Granat.
hederaefolia, *Lindeu, ex A. DC. Prod.* xv. i. 401.— Mexic.
hemicladifolia, *Chem. & Schlecht. in Linnaea*, v. (1830) 603.—Mexic.
herbacea, *Vell. Fl. Flum.* x. t. 53.—Bras.

BEGONIA :—
hernandiaefolia, Hort. ex Klotzsch, in Monatsb. Berl. Akad. (Maerz 1854) 132 = palmatifolia.
hernandiaefolia, Hook. Bot. Mag. t. 4676 = peltata.
hernandiaefolia, Seem. ex A. DC. Prod. xv. i. 339 = glandulosa.
heterophylla, *Hort. Schoenb. ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 122.—Mexic.
heteropoda, *Baker, in Journ. Linn. Soc.* xxi. (1884) 247.—Madag.
Hilariana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 125.—Bras.
hirsuta, *Audl. Pl. Gui.* ii. 918. t. 348.—Guiana.
hirsuta, Hort. ex A. DC. Prod. xv. i. 298 = humilis.
hirsuta, Knir, ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127 = cordifolia
hirsutula, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 575.— Afr. trop.
hirtella, *Link, Enum. Hort. Berol.* ii. 296.—Bras.
hispida, *Schott, ex A. DC. Prod.* xv. i. 368.—Bras.
holosericea, *Teijsm. & Binn. in Epem. Lugd. Bat.* (1868) 5.—Ins. Ternate.
Holtonis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 141.—N. Granat.
homonyma, Steud. Nom. ed. II. i. 194 = Dregei.
Hookeri, Sweet, Hort. Brit. ed. II. 487 = nitida.
Hookeriana, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 135.—Bras.
Horsfieldii, *Miq. ex A. DC. Prod.* xv. i. 397.—Sumatra.
Hugelii, *Hort. Berol. ex A. DC. l. c.* 364.—Bras.
humilis, *Dryand.* in] *Ait. Hort. Kew.* ed. I. iii. 353.— Mexic.
humilis, Bonpl. ex Spreng. Syst. ii. 626 = hirsuta.
humilis, Dryand. in Trans. Linn. Soc. i. (1791) 166 = fisherana.
humilis, Ker-Gawl. in Bot. Reg. t. 284 = dipters.
hydrocotylifolia, *Otto, ex Hook. Bot. Mag.* t. 3968.— Mexic.
hydrophila, Miq. Anal. Bot. Ind. iii. 18 = malabarica.
ignea, *Warsc. in A. DC. Prod.* xv. i. 306.—Guatemala.
imperialis, *Lem. Illustr. Hortic.* vii. (1860) 58.— Mexic.
incana, *Lindl. in Bot. Reg.* (1841) *Misc.* 39.— Mexic.
incarnata, *Link & Otto, Ic. Pl. Sel.* 37. t. 19.— Mexic.
incarnata, Pav. ex A. DC. Prod. xv. i. 373 = cordifolia.
incarnata, Seem. Bot. Voy. Herald, 128 = Seemanniana.
incisa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 129.—Ins. Philipp.
incisa-serrata, *A. DC. Prod.* xv. i. 371.—Bras.
inflata, *C. B. Clarke, in Hook.f. Fl. Brit. Ind.* ii. 636. —Reg. Himal.
insignis, R. Grah. in Bot. Mag. t. 3960 = incarnata.
integerrima, *Spreng. Neue Entdeck.* ii. 174.—Bras.
integrifolia, *Dals. in Hook. Kew Journ.* iii. (1851) 230. —Ind. or.
involucrata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 15.—Costa-Rica.
isoptera, *Dryand. in Sm. Ic.* t. 43 ; *et Trans. Linn. Soc.* i. (1791) 150.—Java.
jamaicensis, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 134 = acutifolia.
Jazmesoniana, *A. DC. l. c.* 139.—Ecuador.
jenmanica, Hort. ex Klotzsch, in Abh. Akad. Berl. (1855) 39 = heracleifolia.
Josephi, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 136.—Reg. Himal.
Junghuhniana, *Miq. Pl. Jungh.* 418.—Java.
Karwinskyana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Mexic.
khasiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 636.—Reg. Himal.
Kunthiana, *Walp. Ann.* ii. 650.—Bras.
lacinista, *Reich. Hort. Bras.* 68 ; *Fl. Ind.* iii. 649.— Reg. Himal. ; Burma ; China.
laciniosa, *A. DC. Prod.* xv. i. 340.—Am. centr.
lasteviron, *Van Houtte, ex A. DC. l. c.* 400.—Hab. ?
lasseoides, *Vell. Fl. Flum.* x. t. 53.—Bras.
Langsdorffii, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 140 = hirsutula.
lanuginosifolia, *A. DC. l. c.* 141.—N. Granat.
langginosa, *A. DC. l. c.* 131.—N. Granat.
laurina, Hort. ex A. DC. Prod. xv. i. 292 = Ottonis.
Lauuii, Linden, in Belg. Hortic. viii. (1858) 166 = xanthina.
lepida, *Blume, Enum. Pl. Jav.* 93.—Java.

BEGONIA :—
lepidota, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 17 = manicata.
leprosa, *Hance, in Journ. Bot.* xxi. (1883) 202.—China.
Liebmannii, *A. DC. Prod.* xv. i. 345.—Mexic.
Lindeniana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Cuba.
Lindleyana, *Walp. Rep.* ii. 209.—Guatemala.
Lindleyana, *Warc. ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 132.—Am. calid.
linosta, *N. E. Br. in Gard. Chron.* (1882) ii. 199.— Java.
lobata, Schott, in Spreng. Syst. iv. App. 408 = multinervia.
Lobbiana, A. DC. Prod. xv. i. 355 = paniculifera.
Lobbii, *A. DC. l. c.* 390.—Java.
lobulata, *A. DC. l. c.* 339.—Mexic.
locellata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 137.—Mexic.
longifolia, *Blume, Cat. Gew. Buitenz.* 103.—Java.
longipes, *Hook. Bot. Mag.* t. 8001.—Mexic.
longipila, Lem. Illustr. Hortic. vii. (1860) 307 = heracleifolia.
longirostris, *Benth. Pl Hartw.* 185.—Ecuador.
longovillosa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Ins. Philip.
lorentzioides, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 580. —Afr. trop.
Lubbersii, *E. Morr. in Belg. Hortic.* xxxiii. (1883) 155. t. 18.—Bras.
lucida, Ekw. Saxifr. Enum 197 = humilis.
lucida, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 16 = Kunthiana.
lucida, Otto & Dietr. Allg. Gartenz. xvi. (1848) 162 = scandens.
lucida, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 208 ; *Contrib.* 57.—Paraguay.
ludica, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 133.—Mexic.
luxurians, *Schnidw. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 131.—Bras.
Lyallii, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 144.—Madag.
Lyncheana, *Hook. f. Bot. Mag.* t. 6758.—N. Granat.
macra, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—N. Granat.
macrophylla, *Dryand. in Trans. Linn. Soc.* i. (1791) 164.—Jamaica.
macrophylla, Lam. Encyc. i. 394 = dipters.
macroptera, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 132.—Bras.
macrota, *Vis. in Act. Ist. Ven. Sc.* Ser. III. iv. (1858–59) 138.—Hab. ?
maculata, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 406. —Bras.
magnifica, *Linden, in Belg. Hortic.* vi. (1856) 5.— N. Granat.
magnifica, Warsc. ex A. DC. Prod. xv. i. 269 = farinulosa.
magnifolia, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
malabarica, *Lam. Encyc.* i. 393.—Ind. or.
malabarica, Roxb. Hort. Beng. 68 ; Fl. Ind. iii. 648 = Roxburghii.
malabarica, Buch.-Ham. in Wall. Cat. n. 3676 D = megaptera.
malvacea, *Klotzsch, in Abh. Akad. Berl.* (1855) 38.— Bras.
manicata, *Cels, ex Vis. Orto Padov.* (1842) 185.— Mexic.
manillensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 129.—Ins. Philipp.
Mannii, *Hook. f. Bot. Mag.* t. 5484.—Afr. trop.
marcayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 209 ; *Contrib.* 58.—Paraguay.
marmorata, Van Houtte, ex Planch. in Fl. des Serres, ix. (1853–54) 243. t. 952–53 = xanthina.
martabanica, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 136.—Burma.
Martiana, *Link & Otto, Ic. Pl. Select. Hort. Berol.* 49. —Mexic.
Martiana, Schlecht. in Linnaea, v. (1830) 604 = incarnata.
martinicensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 128.—Ins. Martinic.
mascarinsis, *Boj. Hort. Maurit.* 271.—Ins. Mascar.
Masraudiee, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 119.—N. Granat.

BEGONIA :—
maxime, *Hort. Berol. ex Klotzsch, in Abh. Akad. Berl.* (1855) 39.—Mexic.
maynensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126.—Peruv.
megaphylla, *A. DC. l. c.* 122.—Bras.
megaptera, *A. DC. l. c.* 184.—Reg. Himal. ; Burma.
Meissneri, *Wall. Cat.* n. 6294.—Ind. or.
membranacea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 139.—Bras.
meridensis, *A. DC. l. c.* 123.—Venezuela.
metallica, L. Sm. in Floral Mag. (1876) t. 197 = incarnata.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 409.—Bras.
Meyeri, *Hook. Bot. Mag.* t. 4100.—Mexic.
Meyeri, Otto & Dietr. Allg. Gartenz. iv. (1836) 349 = tomentosa.
micrantha, Steud. Nom. ed. II. i. 194 = parviflora.
micranthera, *Grisch. in Goett. Abh.* xix. (1874) 148.—Reg. Argent.
microcarpa, *A. DC. Prod.* xv. i. 311.—Ecuador.
microphylla, *A. DC. l. c.* 375.—Reg. Granat.
microphylla, Willd. ex Klotzsch, in Abh. Akad. Berl 1854 (1855) 62 = foliosa.
microptera, *Hook. f. Bot. Mag.* t. 4974.—Borneo.
minuta, Planch. & Linden, in Fl. des Serres, Sér. I viii. (1852–53) 105 = fuchsioides.
minima, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 48 = crenata.
minor, Jacq. Coll. i. 126 = nitida.
modesta, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 20.—Mexic.
modestiflora, *Kunz, in Flora*, liv. (1871) 296.—Reg. Himal. ; Burma.
mollis, *A. DC. Prod.* xv. i. 391.—Java.
monadelpha, *Ruiz & Pav. ex A. DC. l. c.* 286.—Peruv.
monophylla, *Pav. ex A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—Mexic. ?
monophylla, Pourr. ex A. DC. Prod. xv. i. 403 (ex Orchidea).
monoptera, *Link & Otto, Ic. Pl. Rar.* t. 14.—Mexic.
Morisiana, Klotzsch, in Abh. Akad. Berl. (1854) 31 = tuvaensis.
Moritziana, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 16 = scandens.
mouinensiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 649.—Burma.
multangula, *Blume, Enum. Pl. Jav.* i. 96.—Java.
multibulillosa, *Mart. ex Steud. Nom.* ed. II. i. 194. —Bras.
multiflora, *Benth. Pl. Hartw.* 185.—N. Granat.
multinervia, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 18.—Costa Rica.
muricata, Blume, Cat. Gew. Buitenz. 108 = tuberosa.
muricata, *Schrädw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 158.—Brasil.
nana, *L'Hérit. Stirp. Nov.* 99. t. 48.—Madag.
natalensis, *Hook. Bot. Mag.* t. 4841.—Afr. austr.
neglecta, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 139.—Bras.
nelumbiifolia, *Cham. & Schlecht. in Linnaea*, v. (1830) 604.—Mexic.
nemophila, Kunz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 106, 108 = Cathcartii.
neocomensium, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Bras.
nervosa, [Desf.] Hort. Par. ex H. B. & K. Nov. Gen. et Sp. vii. 177 = cucullata.
nigrescens, *Van Houtte.—Cf. Hamb. Gartenz.* viii. (1852) 3.—Hab. ?
nigritarum, Steud. Nom. ed. I. 104 ; ed. II. i. 194 = malabarica ?
nigro-venia, Regel, Gartenfl. (1867) 168 = glandulosa.
nitida, [*Desrand. in*] *Ait. Hort. Kew.* ed. I. iii. 352.—Jamaica.
nossibea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Ins. Nossibé.
Novo-Granatæ, A. DC. Prod. xv. i. 317 = rosacea.
nivea, *Parish, ex Kurz, in Journ. As. Soc. Beng* xlii. (1873) ii. 81.—Tenasserim.
nummulariæfolia, *Putnoys.—Cf. Bot. Zeit.* xi. (1853) 716—Hab. ?
oaxacana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 127.—Mexic.
obliqua, L'Hérit. Stirp. Nov. ii. 46 = nitida.
obliqua, Jacq. Obs. ex Steud. Nom. ed. II. i. 194 = macrophylla.

BEGONIA :—
obliqua, Linn. Sp. Pl. 1056 = acuminata.
obliqua, Ruiz, ex A. DC. Prod. xv. i. 333 = cyathophora.
obliqua, Thunb. Fl. Jap. 231 = Evansiana.
obliqua, *Vell. Fl. Flum* x. t. 48.—Bras.
octopetala, *L'Hérit. Stirp. Nov.* 103.—Peruv.
odorata, Willd. Enum. Hort. Berol. Suppl. 64 = diptera.
olita, *Kerchove, in Rev. Hortic. Belge* (1883), 241—Bras.
oligophylla, *Blume, ex Miq. Fl. Ind. Bat.* i. 692.—Java.
opuliflora, *Puts. in Fl. des Serres*, Sér. I. x (1854–55) 71. t. 995.—N. Granat.
opuliifolia, Loud. Encyc. Pl. 1540 = prænc.?
orbiculata, *Yack, in Malay. Misc.* ii. (1812) vii. 9. —Sumatra.
oregana, *Hort ex Klotzsch, in Monatsb. Berl. Acad.* (Mærz 1854) 126.—Bras.
Ottoniana ×, *A. DC. Prod.* xv. i. 399.
Ottonis, *Walp. Rep.* ii. 212.—N. Granat.
ovatifolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi (1859) 132.—Reg. Himal.
oxylobe, *Weiw. ex Hook. f. in Oliver, Fl. Trop. Afr.* ii. 578.—Afr. trop.
oxyphylla, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 140.—Bras.
oxysperma, *A. DC. l. c.* 122.—As. trop.
palaciosa, *Kurz, in Flora*, liv. (1871) 297.—Burma.
paleata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 137.—Bras.
palmaria, *A. DC. l. c.* 126.—Mexic.
palmata, D. Don, Prod. Fl. Nep. 223 = laciniata ?
palmata, Pav. ex A. DC. Prod. xv. i. 307 = palmaria.
paludosa, *Benth. Pl. Hartw.* 184. n. 1022.— N. Granat.
panduiata, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 209 ; *Contrib.* 59.—Paraguay.
pantherina, *Puts. ex Linden, Cat.* n. 17 (1862) 2.—Mexic.
papilosa, R. Grah. in Hook. Bot. Mag. t. 2846 = incarnata.
papillosa, *Lindl. Bot. Reg.* (1841) *Misc.* 89.—Hab. ?
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 207 ; *Contrib.* 56.—Paraguay.
Parishii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 651.—Burma.
parviflora, Liebm. ex A. DC. Prod. xv. i. 383 = Francisca.
parviflora, *Poepp. & Endl. Nov. Gen. et Sp.* i. 7. t. 12. —Chili.
parviflora, Steud. Nom. ed. II. i. 194 = parvifolia.
parvifiora, Wall. Cat. n. 3674 = modestiflora.
parvifolia, R. Grah. in Bot. Mag. t. 3720 = Dregei.
parvifolia, *Klotzsch, in Abh. Akad. Berl.* (1855) 33.—Bras.
parvifolia, *Schott, in Spreng. Syst. iv. App.* 408.—Bras.
partita, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 139.—Bras.
parvulifiora, *A. DC. l. c.* 136.—Burma.
pastoensis, *A. DC. l. c.* 121.—N. Granat.
patens, Grisch. ex A. DC. Prod. xv. i. 377 = arborescens.
patula, Haw. Pl. Succ. Suppl. 100 = macroptera.
paucicrenata, Lindl. Bot. Reg. t. 471 = macroptera.
paulensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 125.—Bras.
Pavoniana, *A. DC. l. c.* 142.—Peruv.
Pearcei, *Hook. f. Bot. Mag.* t. 5845.—Bolivia.
pedata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 10.—Mexic.
peduncularis, *Wall. Pl. As. Rar.* i. 82. t. 97.—Reg. Himal.
pedunculosa, Wall. Cat. n. 3673 B = khasiana.
peltata, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1844) 138 = Hasskarlii.
peltata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 136.—Bras.
peltata, Otto & Dietr Allg. Gartenz. ix. (1841) 58 = incana.
peltifolia, Schott, in Spreng. Syst. iv. App. 408 = parvipeltata.
pentaphylla, Walp. Rep. ii. 209 = muricata.
peperomioides, *Hook. f. in Oliver, Fl. Trop Afr.* ii. 575.—Afr. trop.

BEGONIA :—
peponifolia, *Vis. Ind. Sem. Hort. Patav.* (1847).—*Cf. Linnæa*, xxiv. (1851) 180.—Jamaica.
perpusilla, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 143.—Ins. Nossibé.
peruviana, *A. DC. l. c.* 138.—Peruv.
petaloides, *Lindl. Bot. Reg.* t. 1757.—Bras.
philippinensis, *A. DC. in DC. Prod.* xv. i. 320.—Ins. Philipp.
phyllomaniaca, *Mart. in Ann. Sc. Nat.* Sér. III. xix. (1853) 306.—Bras.
physalifolia, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 19 = scandens.
picta, Hort. Henders. ex A. DC. Prod. xv. i. 350 = Griffithii.
picta, *Sw. Exot. Bot.* ii. 81. t. 101.—Reg. Himal.
Pilderi, *A. DC. Prod.* xv. i. 380.—Venezuela.
pilifera, *A. DC. l. c.* 387.—Am. centr.
pilosa, *Yack, in Malay. Misc.* ii. (1822) vii. 13.—Sumatra.
pinetorum, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 131.—Mexic.
plagiata, Linden, ex A. DC. Prod. xv. i. 312 = oaxacana.
plantanifolia, *Schott, in Spreng. Syst.* iv. *App.* 407.—Bras.
plebeja, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 8.—Nicaragua.
pleiopetala, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 131.—Bras.
Plumieri, *Kunth, in DC. Prod.* xv. i. 295.—Ins. S. Doming.
pocnifera, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 574.—Afr. trop.
poecila, *C. Koch, in C. Koch & Fint. Wochenschr* (1858) 338.—Hab. ?
Poeppigiana, *A. DC. Prod.* xv. i. 376.—Bras.
Pohliana, Klotzsch, in Abh. Akad. Berl. (1855) 33 = macroptera.
polygonata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 12.—Mexic.
polygonifolia, *A. DC. in Mart. Fl. Bras.* iv. i. 362.—Bras.
polygonoides, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 571. —Afr. trop.
polypetala, *A. DC. — Cf. ' Garden'* (1878) 531.—Peruv.
populifolia, H. B. & K. Nov. Gen. et Sp. vii. 142 = Balmisiana.
populifolia, Klotzsch, in Abh Akad. Berl. (1855) 30 = tuvaensis.
populifolia, Schott, in Spreng. Syst iv. App. 408 = scandens.
populnea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 128.—Bras.
Porteriana, Fisch. Mey. & Avé-Lall Ind. Sem Hort. Petrop. viii. 51 = itumilia.
portoricensis, *A. DC. Prod.* xv. i. 295.—Porto Rico.
Prieurei, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 142.—Guiana.
Princeps, *Hort. Berol. ex Klotzsch, in Abh. Akad. Berl.* (1855) 39.—Bras.
priamatocarpa, *Hook. Bot. Mag.* t. 6307.—Afr. trop.
procrictifolia, *Wall. Cat.* n. 6292.—Burma.
procumbens, *Vell. Fl. Flum.* x. t. 36.—Bras.
prolifera, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 135.—Burma ; Malaya.
prolnata, *A. DC. Prod.* xv. i. 338.—Costa Rica.
pulchella, *Raddi, in Mem. Mod.* xviii. *Fas.* (1820) 407. —Bras.
pulchra, Herb. Schreb. ex A. DC. Prod. xv. i. 298 = nitida.
punctata, Hort. ex Steud. Nom. ed. II. i. 194 = maculata.
punctata, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Berol. i. 16 = heracleifolia.
Purdieana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Jamaica.
purpurea, Sw. Prod. Veg. Ind. Occ. 85 = nitida.
pusilla, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Bolivia.
pusulata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 6.—Mexic.
Putseysiana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Venezuela.
querciifolia, *A. DC. l. c.* 130.—Ins. Philipp.
racemosa, *Yack, in Malay. Misc.* ii. (1822) vii. 14. —Sumatra.

BEGONIA :—
radiata, R. Grah. in Edinb. N. Phil. Journ. (July 1838) 132 = heracleifolia.
radicans, *Vell. Fl. Flum.* x. t. 89.—Bras.
ramentacea, *Pax. Mag. Bot.* x. 78.—Bras.
reniformis, Becd. in Madr. Journ. Sc. xxii. (1861) 72 = zoalmaiaiensis.
reniformis, Dryand. in Trans. Linn. Soc. i. (1791) 161. t. 14. f. 1-2 = longipes
reniformis, Hook. Bot. Mag. t. 3225 = vitifolia.
reniformis, Hort. Berol. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 24 = Dregei.
reniformis, Pav. ex A. DC. Prod. xv. i. 308 = Balmisiana.
reniformis, *Vell. Fl. Flum.* x. t. 40.—Bras.
repanda, *Blume, Enum. Pl. Jav.* i. 97.—Sumatra
repens, Blume, l. c. 96 = stolifa.
repens, Lam. Encyc. i. 394 = rotundifolia.
repens, Hemsl. Biol. Centr. Am. Bot. i. 497 = Liebmanni.
repens, Ruiz & Pav. ex A. DC. Prod. xv. i. 333 = Rosamanniae.
repens, Schott, ex A. DC. l. c. 365 = convolvulacea.
repens, *Vell. Fl. Flum.* x. t. 85.—Bras.
reptans, *Benth. Pl. Hartw.* 62.—Mexic.
reptans, Liebm. in Kjoeb. Vidensk Meddel. 1852 (1853) b = Liebmanni.
reticulata, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 134.—Bras.
Rex, *Puis. in Fl. des Serres*, ii. (1857) 141. t. 1255.—Reg. Himal.
rhizocarpa, *Fisch. ex A. DC. Prod* xv. i. 388.—Bras.
rhizocaulis, *Hort. Berol. ex A. DC. l. c.* 340 — Am. centr.
rhombacarpa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 329.—Ins. Luzon.
Richardii x, [*C. Moore*] *Cat. Bot. Gard. Sydney* (1857), 9.—Hab.!
Richardsiana, *T. Moore, in Gard. Chron.* (1871) 1065. —Afr. austr.
Richardsoniana, *Houll. in Rev. Horlic.* (1872) 333.—Hab.!
ricinifolia x, A. Dietr. Allg. Gartenz. xv. (1847) 252.
Riedelii, *A. DC. in Ann. Sc. Nat* Sér. IV. xi. (1859) 137.—Bras.
rigida, *Regel, ex A. DC. Prod.* xv. i. 378.—Bras.
robusta, *Blume, Enum. Pl. Jav.* i. 96.—Java.
robusta, Zoll. ex A. DC. Prod. xv. i. 275 = multangula.
Rockwelli, *Hort —Cf. Gard. Chron.* (1867) 11. 481 — Hab.!
Roezli, Regel, Gartenfl. (1876) 194; et in Act. Hort. Petrop. iv. (1876) 324 = Lynchiana.
ROSACEA, *Puis. in Belg. Hortic.* vii. (1857) 196; *et in Fl. des Serres*, Sér. II. ii. (1857) 25. t. 1194. —N. Granat.
rosaeflora, *Hook. f. Bot. Mag.* t. 5680.—Peruv.
rosea, *A. DC. Prod.* xv. i. 299.—Am. centr.
rosea, Pav. ex A. DC. l. c. 278 = cordifolia.
Rosamaniae, *A. DC. l. c.* 333.—Peruv.
rostrata, *Welw. ex Hook. f. in Oliver, Fl. Trop. Afr.* ii. 578.—Afr. trop.
rotata, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 11 = carolinaefolia.
rotunda, *Vell. Fl. Flum.* x. t. 38.—Bras.
rotundifolia, *Lam. Encyc* i. 394.—Ind. occ.
Roxburghii, *A. DC. Prod.* xv. i. 398.—Reg. Himal.; Burma.
Royiai, *Linden, in Belg. Hortic.* vii. (1857) 197.—Ind. or.
rubella, *Buch.-Ham. ex D. Don, Prod. Fl. Nep* 223.—Nepal.
rubella, Miq. Anal. Bot. Ind. iii. 18 = canarana.
rubella, Wall. Cat. n. 3687 = scutata.
rubicunda, Hort. Turic. ex A. DC. Prod. xv. i. 384 = Dregei.
rubra, *Blume, Enum. Pl. Jav.* i. 96.—Java.
rubricaulis, *Hook. Bot. Mag.* t. 4181.—Peruv.
rubro-nervia, *Hort. ex Klotzsch, in Abh. Akad. Berl.* (1855) 40.—Ind. or.
rubro-palens, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 195.—Bras.
rubro-setulosa, *A. DC. Prod.* xv. i. 390.—Hab.?
rubro-venia, *Hook. Bot. Mag.* t. 4869.—Reg. Himal.
rufa, *Thunb. in Flora*, iv. (1821) 330.—Bras.
rugosa, *Hort. Schoenb. ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 126.—Hab.!
rupestris, *Bunt. ex Lond. Hort. Brit. Suppl.* iii. 498. —Bras.

BEGONIA :—
rupicola, *Miq. Pl. Yungh.* 418.—Java.
rutilans, *Hort. Van-Houtte, ex A. DC. Prod.* xv. i. 373.—Hab.!
sagittata x, Hort.—Cf. Gard. Chron. (1868) 1. 1317.
salicifolia, *A. DC. in Ann. Sc. Nat* Sér. IV. xi. — (1859) 128.—Bras.
sandalifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 649.—Burma.
Sandereii, *Hort. Kew. ex A. DC. Prod.* xv. i. 400, nonen.—Hab.!
sanguinea, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 409 —Bras.
sanguinea, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 12.—Mexic.
Sartorii, Liebm. l. c. 14 = pruec.
Sattrea, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 638. —Reg. Himal.
saxatilis, *Blume, Enum. Fl. Jav.* 95.—Ins. Philipp
saxicola, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 144.—Bras.
Saxifraga, *A. DC. l. c.* 138.—Bras.
scabrida, *A. DC. Prod.* xv. i. 367.—Venezuela.
scandens, *Vell. Fl. Flum* x. t. 41.—Bras.
scandens, Hort. Schoenb. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 126 = convolvulacea.
scandens, *Sw. Prod. Veg. Ind. Occ.* 80.—Am. austr.
scapigera, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 572.— Afr. trop.
schnizlepis, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 17 = manicata.
Schlumbergeriana, *Lem. Illustr. Horlic.* (1858) *Misc.* 61.—Bras.
Schmidti, Hort. ex Wien. Illustr. Gart. Zeit. (1880) 250 = Schmidtiana.
Schmidtiana, *Regel, Gartenfl.* (1879) 321.—Bras.
Schottiana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 140.—Bras.
scutata, *Wall. Cat.* n. 3686. partim.—Reg. Himal.
scutata, Wall. l. c. A = Josephi.
scutellata, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 9 = conchaefolia.
scutifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 572.— Afr. trop.
Scutiflora, *Hook. f. l. c.* 575.—Afr. trop.
Sedeni x, Hort.—Cf. Gard. Chron. (1870) 38.
Seemanniana, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi (1859) 138.—Am. centr.
Sellol, *Ladd. ex Steud. Nom.* ed. II. i. 194—Bras.
Sellowii, Klotzsch, in Abh. Akad. Berl. (1855) 28 = semperflorens.
semiovata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 22.—Am. centr.
semperflorens, Hook. Bot. Mag. t. 2920 = Hookeri.
semperflorens, *Link & Otto, Ic. Pl. Rar.* t. 9.—Bras.
sericoneura, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 13.—Am. centr.
sericoneura, Seem. Bot Voy. Herald, 128 = pilifera.
serotina, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—Guayaquil
sessilifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 577. —Afr. trop.
setaria, Hort. Angl. ex A. DC. Prod. xv. i. 293 = semperflorens.
setifera, *A. DC. l. c.* 338.—Am. centr.
setosa, *Klotzsch, in Abh. Akad. Berl.* (1855) 31.—Bras.
setulosa, *Bertol. in Nov. Comm. Bon.* iv. (1840) 437.— Guatemala.
sikkimensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Reg. Himal.
silhetensis, *C. B Clarke, in Hook. f. Fl. Brit. Ind.* ii. 636.—Reg. Himal.
sinensis, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 125 = Evansiana.
sinuata, N. Grah. in Edinb. N. Phil. Journ. (Jan. 1838) 192 = Dregei.
sinuata, *E. Mey. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 257.—Afr. austr.
sinuata, *Wall. Cat.* n. 3680.—Burma ; Malaya.
smilacina, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 137.—Bras.
socotrana, *Hook. f. in Gard. Chron.* (1881) 1. 8.—Ins. Socotr.
solanantherа, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126.—Bras.
sparsipila, *Baker, in Saund. Ref. Bot.* t. 340.—Am. centr. !
spathulata, Ludd. Bot. Cab. t. 107 = cucullata.

BEGONIA :—
speciosa, Hort. Van Houtte, ex A. DC. Prod. xv. i. 292 = nitida.
spilotophylla, *F. Muell. Not. Papuan Pl.* iv. 67.—N. Guinea.
splendens, Hort. Turic. ex A. DC. Prod. xv. i. 375 = Putzeyiana.
splendida, Hort. Rollison, ex A. DC. l. c. 275 = robusta.
Spruceana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 142.—Peruv.
squamulosa, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 579. —Afr. trop.
squarrosa, *Liebm. in Kjoeb Vidensk. Meddel.* 1852 (1853) 7.—Mexic.
squarrosa, Secm. Bot. Voy. Herald, 128 = stigmosa.
Stella, *Regel, Gartenfl.* (1875) 88.—Hab.!
stenophylla, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi. (1859) 137.—Bras.
stictopoda, *Miq. ex A. DC. Prod.* xv. i. 391.—Sumatra.
stigmosa, *Lindl. Bot. Reg.* (1840) *Misc.* 32.— Mexic.
stipulacea, Willd. Sp. Pl. iv. 414 = angularis.
strigillosa, *A. Dietr. Allg. Gartenz.* xix. (1851) 380.— Am. centr.
strigulosa, *A. DC. Prod.* xv. i. 399.—Hab.!
suaveolens, Haw. in Lodd. Bot. Cab. t. 69 = diptera.
suaveolens, Klotzsch, in Abh. Akad. Berl. (1855) 26 = nitida.
subcillata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 132.—Peruv.
subhumilis, *A. DC. l. c.* 124.—Guayaquil
sublobata, *Yack, in Malay. Misc.* ii. (1822) n. vii. 10. —Sumatra.
subnuta, Wall Cat. n. 3688 = ovatifolia.
subpeltata, Regel, Ind. Sem. Hort. Petrop. (1866) 102 = incarnata.
subpeltata, *Wight, Ic.* t. 1812.—Ind. or.
subperfoliata, *Parish, ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 81.—Burma.
subrotunda, Wall. Cat. n. 6298 = sinuata.
subvillosa, *Klotzsch, in Abh. Akad. Berl.* (1855) 32.— Bras.
suffruticosa, *Mexin. in Linnaea*, xiv. (1840) 502.— Ait. austr.
suffruticosa, Pav. ex A. DC. Prod. xv. i. 311 = Boissieri.
sulcata, *Schnidc. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 130.—N. Granat.
surculigera, *Kurz, in Flora*, liv. (1871) 296.—Burma.
Sutherlandi, *Hook. f. Bot. Mag.* t. 5689.—Afr. austr.
sylvatica, *A. DC. Prod.* xv. i. 365.—Bras.
sylvestris, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi. (1859) 140.—Bras.
tanacetifolia, Hort. ex Steud. Nom. ed. II. i. 194 = heracleifolia.
Telfariae, Steud. l. c. sphalm. vide Bignonia Telfairiae.
tenella, D. Don, Prod. Fl. Nep. 223 = amoena.
tenera, *Dryand. in Trans. Linn. Soc.* i. (1791) 169. t. 16.—Ceylan.
tenuicaulis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Bolivia.
tenuifolia, *Dryand. in Trans. Linn. Soc.* i. (1791) 162. t. 14.—Java.
tessaricarpa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 636.—Assam.
Teuscheri, *Linden, ex André, in Illustr. Hortic.* (1879) 137. t. 358.—Malaya.
Thomsoni, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 124.—Reg. Himal.
Thunbergii, Hook. Bot. Mag. t. 4092 = teneva.
tomentosa, *Schott, in Spreng. Syst.* iv. *App.* 408.— Bras.
tovarensis, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 122.—Mexic.
trachyptera, *Benth. Pl. Hartw.* 184.—N. Granat.
trichocarpa, *Dels. in Hook. Ken Journ.* iii. (1851) 230. —Ind. or.
trichopoda, *Miq. Fl. Ind. Bat.* i. 1. 1098.—Sumatra.
tricuspidata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 637.—Burma.
triradiata, *C. B. Clarke, l. c.*—Burma.
tropaeolifolia, *A. DC. in Ann. Sc. Nat.* Sér. IV xi. (1859) 120.—N. Granat.
truncata, Vell. Fl. Flum. x. t. 47 = tomentosa vitifolia.
tuberosa, Herb. Wight, ex Wall. Cat. n. 3678 = malabarica.

BEGONIA:—
tuberosa, Heyne, ex A. DC. Prod. xv. 1. 291 = dipetala.
tuberosa, *Lam. Encyc.* i. 298.—Ins. Molucc.
tuberosa, Pav. ex A. DC. Prod. xv. 1. 311 = bulbillifera.
tuberosa, Ruiz, ex A. DC. l. c. 281 = geranifolia.
uliginosa, *Klotzsch*, in *Abh. Akad. Berl.* (1856) 85.—Bras.
ulmnolia, *Willd. Sp. Pl.* iv. 418.—Venezuela.
umbellata, *H. B. & K. Nov. Gen. et Sp.* vii. 148.—N. Granat.
umbilicata, Hort. ex Planch. in Fl. des Serres, Sér. I. viii. (1852–53) sub t. 810 = hernandiæfolia.
umbrata, *Hort. ex A. DC. Prod.* xv. 1. 296.—Hab. ?
uncinata, Hort. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 134 = Dregei.
undulata, *Schott, in Sprengl. Syst. iv. App.* 406.—Bras.
uropbylla, *Hook. Bot. Mag.* t. 4855.—Mexic.
Urticæ, *Linn. f. Suppl.* 420.—N. Granat.
urticæfolia, Sm. Ic. Fasc. ii. 45 = tuberosa.
urticæfolia, Hort. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127 = B. Pildera.
urticifolia, F. G. Dietr. Vollst. Gartenlex. ii. 177 = Urticæ.
vaginata, *Vell. Fl. Flum.* x. t. 87.—Bras.
valdensium, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Bras.
variana, *A. DC. l. c.* 135.—Java.
Veitchii, *Hook. f. in Gard. Chron.* (1867) 734.—Peruv.
velleræ, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 122.—Bras.
Vellosiana, Walp. Rep. ii. 216 = repens.
velutina, Hort. Vindob. ex Klotzsch, in Abh. Akad. Berl. (1856) 41 = lobata.
velutina, Pariah, ex Kurs, in Journ. As. Soc. Beng. xiii. (1873) ii. 81 = parvulifora.
verricosa, Hort. Berol. ex Klotzsch, in Abh. Akad. Berl. (1855) 40 = lobata.
Verschaffeltiana ×, Lem. Illustr. Hortic. ii. (1855) t. 68 = Verschaffeltii.
Verschaffeltii ×, *Regel, Gartenfl.* (1855) 248. t. 189.
verticillata, Vell. Fl. Flum. x. t. 45 = digitata.
verticillata, *Hook. Ic. Pl.* t. 811.—Burma.
villosa, Gardn. in Hook. Lond. Journ. i. (1842) 186 = macroptera.
villosa, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 123 = subvillosa.
villosa, Lindl. Bot. Reg. t. 1952 = hirtella.
violæfolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Mexic.
viridiflora, *A. DC. l. c.* 133.—Peruv.
visifolia, Lindl. Bot. Reg. (1852) Misc. 21 = Lindleyana.
vitifolia, *Schott, in Sprengl. Syst.* iv. *Cur. Post.* 407.—Bras.
Wagneriana, *Hook. Bot. Mag.* t. 4968.—Bras.
Wallichiana, *Steud. Nom.* ed. II. i. 394; *Schim. in Linnæa*, xxiv. (1851) 218 (sp. dubia).—Ind. or.
Walpersii, *Heynh. Nom.* ii. 68.—Venezuela.
Warscewiczii, Neuman, in Rev. Hortic. Sér. III. i. (1847) 186 = conchæfolia.
Weddelliana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 122.—Bolivia.
weltoniensis ×; *Hort.*—*Cf. Y. B. Weber, in Rev. Hortic.* (1876) 106.
Wightiana, Wall. Cat. n. 3673 = albo-coccinea.
Wrightiana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 128.—Cuba.
xanthina, *Hook. Bot. Mag.* t. 4683 et t. 5107.—Reg. Himal.
zebrina, Hort. Angl. ex Lond. Encyc. Pl. Suppl. ii. 1506 = angularis.
Zollingeriana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 135.—Java.

BEGONIELLA, Oliver, in Trans. Linn. Soc. xxviii. (1873) 513. *BEGONIACEÆ.*
angustifolia, *Oliver, in Hook. Ic. Pl.* t. 1487.—N. Granat.
Kaltenyeri, *Oliver, l. c.* t. 1352.—N. Granat.
Whitei, *Oliver, in Trans. Linn. Soc.* xxviii. (1873) 513.—N. Granat.

BEHAMIA, Grimb. Cat. Pl. Cub. 77 (1859). *LEGUMINOSÆ*, Benth. & Hook. f. i. 1002.
cubensis, *Griseb. l. c.* 76.—Cuba.

BEHEN, Hill, Veg. Syst. iv. 41 (1762) = **Vernonia**, Schreb. (Compos.).
Yacca, Hill, Hort. Kew. 65 = Centaurea Jacea.
acmebarensis, Hill, l. c. = V. novebarcensis.
prasalium, Hill, l. c. = V. novebarcensis et alicissima.

BEHEN, Moench, Meth. 709 (1794) = **Silene**, Linn. (Caryoph.).
angustifolius, Schott, ex Steud. Nom. ed. II. i. 194 = S. Cucubalus.
glareosus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 348 = S. Cucubalus ?
puberulus, Fourr. l. c. = S. Cucubalus.
trachiurus, Fourr. l. c. = S. Cucubalus.
vesicarius, Fourr. l. c. = S. Cucubalus.
vulgaris, Moench, Meth. 709 = S. Cucubalus.

BEHENANTHA, Schur, in Verh. Naturf. Ver. Bruenn, xv. ii. (1877) 130 = **Silene**, Linn. (Caryoph.).
Carei, Schur, l. c. 132 = S. Cucubalus.
glaucescens, Schur, l. c. = S. Cucubalus.
inflata, Schur, l. c. 131 = S. Cucubalus.
saponariæfolia, Schur, l. c. 133 = S. Cucubalus.

BEHNIA, F. Didricha. in Kjoeb. Vidensk. Meddel. (1854) 182. *LILIACEÆ*, Benth. & Hook. f. iii. 766.
DICTYOPSIS, Harv. in Bot. Mag. t. 5635 (1867).
HYLONOMA, Baker, in Journ. Linn. Soc. xiv. (1875) 561.
reticulata, *F. Didrichs. l. c.* 183.—Afr. austr.

BEHRINIA, Sieber, ex Steud. Nom. ed. I. 105 (1821) = BERINIA, Brign. = Crepis, Linn. (Compos.).

BEHURIA, Cham. in Linnæa, ix. (1834) 376. *MELASTOMACEÆ*, Benth. & Hook. f. i. 748.
insignis, *Cham. l. c.* 377.—Bras.

BEILIA, Eckl. Verz. 42 (1827) = **Watsonia**, Mill. (Irid.).
spicata, Eckl. l. c. = W. spicata.
tritticea, Eckl. l. c. = W. plantaginea.

BEILSCHMIEDIA, Nees, in Wall. Pl. As. Rar. ii. 69 (1831). *LAURINEÆ*, Benth. & Hook. f. iii. 193.
BEILSCHMEDIA, Panch. & Sebert, Not. Bois Nouv. Caléd. 176 (1874).
HUFELANDIA, Nees, Prod. Laurin. ii. (1833); et Laurin. Expos. (1836) 197, 674.
NESODAPHNE, Hook. f. Fl. N. Zel. i. 217 (1853).
assamica, *Meisen. in DC. Prod.* xv. i. 64.—Assam; Burma.
Bailloni, *Panch. & Sebert, Not. Bois Nouv. Caled.* 176.—N. Caled.
calcarea, *Kurz, f. in Ann. Jard. Buitenz.* i. (1876) 67.—N. Guin.
chinensis, *Hance, in Journ. Bot.* xx. (1882) 79.—China.
fagifolia, *Nees, in Wall. Pl. As. Rar.* ii. 69.—Ind. or. globularia, *Kurz, For. Fl. Brit. Burma*, ii. 294.—Burma.
javanica, *Mig. Fl. Jungh.* i. 176.—Java.
lanceolata, *Panch. & Sebert, Not. Bois Nouv. Caled.* 177.—Caled.
lancifolia, *Mig. Fl. Jungh.* 177.—Java.
macrophylla, *Meisen. in DC. Prod.* xv. i. 63.—Burm
Madang, *Blume, Mus. Bot. Lugd. Bat.* i. 332.—Java.
oppositifolia, *Benth. & Hook. f. Gen.* iii. 152.—Zeylan.
pendula, *Hemsl. Biol. Centr. Am. Bot.* iii. 70.—Mexic.
Roxburghiana, Dalz. & Gibs. Bomb. Fl. 232 = fagifolia.
Roxburghiana, *Nees, in Wall. Pl. As. Rar.* ii. 61.—Assam; Burma.
Thomsea, *Benth. & Hook. f. Gen.* iii. 152.—Ind. oc.
undulata, *Mig. Fl. Ind. Bat.* i. 1. 920.—Java.
Wightii, *Benth. & Hook. f. Gen.* iii. 152.—Ind. or.

BEJARIA, Zea, in Anal. Clenc. Nat. iii. (1801) 151, in nota; Vent. Jard. Cels, sub t. 51 = **Befaria**, Mutis (Ericac.).

BEJUCO, Loefl. Itin. 314 (1758) = **Hippocratea**, Linn.
pendulus, Loefl. l. c. = H. scandens.

BELAIRIA, A. Rich. Ess. Fl. Cub. 511. t. 40 (1845). *LEGUMINOSÆ*, Benth. & Hook. f. i. 558.
macrostata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 178.—Cuba.
spinosa, *A. Rich. Ess. Fl. Cub.* 513.—Cuba.
ternata, *Wright, ex Griseb. Cat. Pl. Cub.* 81.—Cuba.

BELAMCANDA, Adans. Fam. ii. 60 (1763) (Belamcanda). *IRIDIÆ*, Benth. & Hook. f. iii. 697.
PARDANTHUS, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 246.
aristata, Moench, Meth. Suppl. 214 = Sparaxis grandiflora.
bulbifera, Moench, l. c. = Sparaxis bulbifera.
chinensis, [DC. in] Red. Lil. iii. t. 121 = punctata.
fenestrata, Moench, Meth. 529 = Tritonia crocata.
punctata, *Moench, l. c.*—China.
semiflavescens, Moench, Meth. Suppl. 214 = Sparaxis grandiflora.
tripedalis, *Flack. ex Roem. & Schult. Syst.* i. *Mant.* 306 (sp. dub.).
villosa, Moench, Meth. Suppl. 214 = Babiana villosa.

BELANGERA, Cambess. Canon. Syn. Bras. Mer 8 (1829). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 650.
LAMANONIA, Vell. Fl. Flum. v. t. 104 (1827).
POLYSTEMON, D. Don, in Edinb. N. Phil. Journ. ix. 95 (1830).
cuneata, *Cambess. l. c.*—Bras.
denticulata, *Moric. Pl. Nouv. Am.* 158. t. 90.—Bras.
glabra, *Cambess. Canon.-Syn. Bras. Mer.* (1829) 8.—Bras.
intermedia, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 95.—Bras.
Lamanonia, *Steud. Nom.* ed. II. i. 195.—Bras.
Riedeliana, Cæsar. ex Eichl. in Mart. Fl. Bras. xiv. ii. 165 = speciosa.
speciosa, *Cambess. Canon. Syn. Bras. Mer.* (1829) 4.—Bras.
tomentosa, *Cambess. l. c.*—Bras.

BELANTHERIA, Nees, in DC. Prod. xi. 96 (1847) = Brillantaisia, Beauv. (Gramin.).
Beloisiana, Nees, l. c. 97 = Brillantaisia owariensis.

BELENCITA, Karst. in C. Koch, Allg. Gartenz. (1857) 26. *CAPPARIDEÆ.*
Hagenii, *Karst. l. c.* 24.—N. Granat.

BELENIA, Decne. in Jacquem. Voy. Bot. 112. t. 120 (1844) = Physochlaina, D. Don (Solanac.).
grandiflora, Decne. l. c. 114. t. 120 = P. praealta.

BELENIDIUM, Am. ex DC. Prod. viii. 292, in syn. (1844) = Hymenatherum, Cass. (Compos.).
Candollеanum, Am. ex DC. l. c. = H. Belenidium.

BELEROPONE, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 524 (1885), aphalm. = Beloperone, Nees (Acanth.).

BELHARNOSIA, Adans. Fam. ii. 428 (1763) = Sanguinaria, Linn. (Papav.).

BELIA, Steller, ex S. G. Gmel. Fl. Sibir. iv. 88 (1769) = Claytonia, Gron. (Portulac.).

BELILLA, Adans. Fam. ii. 150 (1763) = **Mussenda**, Burm. (Rubiac.).

BELIS, Salisb. in Trans. Linn. Soc. viii. (1807) 315, nomen prius = Cunninghamia, R. Br. (1826) (Coniferæ).
jacuifolia, Salisb. l. c. = C. sinensis.
lanceolata, Sweet, Hort. Brit. ed. II. 475 = C. sinensis.

BELLADONA, Adans. Fam. ii. 219 (1763) = **Atropa**, Linn. (Solanac.).

BELLADONNA, Sweet, Hort. Brit. ed. II. 506 (1830) = Amaryllis, Linn.
blanda, *gallica*, *pudica*, *purpurascens*, Sweet, l. c. = A. Belladonna.

BELLADONNA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 252 (1745) = **Atropa**, Linn. (Solanac.).
baccifera, Lam. Fl. Fr. ii. 256 = A. Belladonna.
peruviana, Hort. Par. ex Adans. Fam. ii. 219 = Nolana prostrata.
trichotoma, Scop. Fl. Carn. ed. II. i. 160 = A. Belladonna.

BELLARDIA, All. Fl. Pedem. i. 61 (1786) = **Bartsia**, Linn. (Scrophul.)
Trixago, All. l. c. = Bartsia Trixago.
viscosa, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 4 = Bartsia viscosa.

BELLARDIA, Colla, in Mem. Acc. Torin. xxxviii. (1835) 40. t. 34 = **Microseris**, D. Don (Compos.).
pusilla, Colla, l. c. 43. t. 34.—Chili.

BELLARDIA, Schreb. Gen. i. 790 (1789) = **Coccocephalum**, R. Br. (Rubiac.).
mollis, Willd. ex DC. Prod. iv. 397 = C. canescens.
repens, Willd. Sp. Pl. i. 926 = C. Tontanea.
Tontanea, Roem. & Schult. Syst. iii. 305 = C. Tontanea.

BELLENDENA, R. Br. in Trans. Linn. Soc. x. (1810) 166. *PROTEACEÆ*, Benth. & Hook. f. iii. 179.
montana, *R. Br. l. c.* 48, 166.—Tasmania.

BELLENDENIA, Rafin. ex Endl. Gen. 169 (1837), in syn. = **Tritonia**, Ker-Gawl. (Irid.).

BELLENDENIA, Endl. Gen. 840 (1888) = **Bellendena**, R. Br. (Proteac.).

BELLEVALIA, Delile, ex Endl. Gen. 281 (1836) = **Altheniа**, Petit (Naïad.).
australis, Delile, ex Parl. Fl. Ital. iii. 648 = A. filiformis.

BELLEVALIA, Lapeyr. in Journ. Phys. lxvii. (1808) 435 = **Hyacinthus**, Tourn. (Liliac.).
abortiva, Gren. in Gren. & Godr. Fl. Fr. iii. 217 = H. trifoliatus.
aleppica, Boiss. Diagn. Ser. II. iv. 111 = H. nervosus.
appendiculata, Lapeyr. Hist. Abr. Pl. Pyr. 186 = H. romanus.
atrovioiacea, Regel, in Act. Hort. Petrop. viii. (1884) 654.—As. centr.
aurea, Boiss. Fl. Orient. v. 807 = H. ciliatus.
Battandieri, Freyn, in Flora, lxviii. (1885) 25.—Afr. bor.
Boissieri, Freyn, l. c. 95.—Graecia.
Bourgæana, Heldr. ex Nym. Consp. 733.—Graecia.
calandriniana, Nym. Consp. 732 = Muscari comosum.
caucasica, Griseb. Spicil. Fl. Rumel. ii. 387 = Muscari caucasicum.
ciliata, T. Ness, Gen. 4. t. 8; Kunth, Enum. Pl. iv. 308 = H. ciliatus.
Clusiana, Griseb. Spicil. Fl. Rumel. ii. 387 = H. dubius.
comosa, Kunth, Enum. Pl. iv. 306 = Muscari comosum.
Cupaniana, Nym. Consp. 733 = Muscari Cupanianum.
curta, Nym. l. c. = Muscari Holzmanni.
cyanescens, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 151 = H. romanus.
densiflora, Boiss. Diagn. Ser. I. vii. 109 = H. densiflorus.
dubia, Reichb. Fl. Germ. Excurs. 105 = H. dubius.
fallax, Pomel, Nouv. Mat. Fl. Atl. 254.—Algeria.
Heldreichi, Boiss. Diagn. Ser. I. xiii. 86 = H. Szovitsianus.
Pontæanæi, Nym. Consp. 733.—Creta.
glauca, Kunth, Enum. Pl. iv. 306 = H. ciliatus.
graeca, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 238 = Muscari graecum.
graminifolia, Nym. Consp. 733.—Ins. Graec.
Gussoniana, Griseb. Spicil. Fl. Rumel. ii. 387.—Sicil.
Hackeli, Freyn, in Oestr. Bot. Zeitschr. xxvii. (1877) 289.—Europ.
Haynei, Boiss. Fl. Orient. v. 806 = H. Haynei.
Heldreichii, Boiss. Diagn. Ser. II. iv. 111 = H. tenuiflorus.
hispida, J. Gay, in Bull. Soc. Bot. Fr. iii. (1856) 240 = H. hispidus.
Holzmanni, Heldr. in Atti Congr. Bot. Firenze, 1874 (1879). 238 = Muscari Holzmanni.
leucophæa, Boiss. Fl. Orient. v. 806—H. leucophaeus.
linearis, Kunth, Enum. Pl. iv. 309 = H. linearis.

BELLEVALIA:—
macrobotrys, Boiss. Diagn. Ser. I. xiii. 86 = Hyacinthus macrobotrys.
maritima, Kunth, Enum. Pl. iv. 307 = Muscari maritimum.
mauritanica, Pomel, Nouv. Mat. Fl. Atl. 255.—Algeria.
micrantha, Boiss. Diagn. Ser. I. v. 68 = H. micranthus.
monophylla, J. Gay, ex Boiss. Fl. Orient. v. 294 = Muscari latifolium.
montana, Boiss. l. c. 804 = Muscari montanum.
muscarioides, Mast. in Journ. Linn. Soc. iii. (1859) 113 = Muscari latifolium.
nervosa, Boiss. Fl. Orient. v. 806 = H. nervosus.
nivalis, Boiss. & Kotschy, Diagn. Ser. II. iv. 110 = H. nivalis.
operculata, Lapeyr. in Journ. Phys. lxvii. (1808) 426, cum. tab. = H. romanus.
pallens, Vis. in Mem. Ist. Venet. xvi. (1871) 67 = H. dalmaticus.
paradoxa, Boiss. Fl. Orient. v. 808 = H. paradoxus.
persica, Boiss. l. c. = H. persicus.
pharmacusana, Nym. Consp. 733 = Muscari Sartorianum.
Pinardi, Boiss. Diagn. Ser. I. v. 62 = Muscari Pinardi.
pseudo-muscari, Boiss. & Buhse, Diagn. Ser. II. iv. 110 = H. paradoxus.
romana, Sweet, Hort. Brit. ed. I. 419; Reichb. Fl. Germ. Excurs. 105 = H. romanus.
Sartoriana, Nym. Consp. 733 = Muscari Sartorianum.
sessiliflora, Kunth, Enum. Pl. iv. 309 = H. sessiliflorus.
spicata, Boiss. Diagn. Ser. I. vii. 110 = H spicatus.
syriaca, Herb. in Lindl. Bot. Reg. (1844) Misc. 88 = H. trifoliatus.
tenuiflora, Nym. Consp. 732 = Muscari caucasicum.
theracea, Nym. l. c. 733 = Muscari theraeum.
trifoliata, Kunth, Enum. Pl. iv. 308 = H. trifoliatus.
turkestanica, Franch. in Ann. Sc. Nat. Sér. VI. xviii. (1884) 256.—Turkest.
trifoliifolia, Freyn, in Flora, lxviii. (1885) 29.—Alger.
Webbiana, Parl. Nouv. Gen. e Spec. 19 = Hyacinthus dubius.
Wetsiei, Freyn, in Oestr. Bot. Zeitschr. xxviii. (1878) 87 = Muscari Weissii.

BELLIDIASTRUM, Mich. ex Hall. Enum. Stirp. Helv. ii. 793 (1742) = **Aster**, Tourn. (Compos.).
Michelii, Fiori. Enxayo Fl. Gall. 259 = Bellis sylvestris.
Micheli, Cass. in Dict. Sc. Nat. iv. 70 = A. Bellidiastrum.
montanum, Hoppe, ex DC. Prod. v. 226 = A. Bellidiastrum.
subalpinum, Schrank, Fl. Mon. iii. 251 = A. Bellidiastrum.

BELLIDIASTER, Dum. Fl. Belg. 66 (1827) = **Aster**, Tourn. (Compos.).
montanus, Dum. l. c. = A. Bellidiastrum.

BELLIDIASTRUM, [Vaill.] Less. Syn. Comp. 388 (1829) = **Osmites**, Linn. (Compos.).
osmitoides, Less. l. c. 384 = O. Bellidiastrum.

BELLIDIASTRUM, Cass. in Bull. Soc. Philom. (1816) 199 = **BELLIDIASTRUM, Mich.** = **Aster**, Tourn. (Compos.).

BELLIDISTRUM, Reichb. Consp. 109 (1828) = prasc.

BELLIDIOPSIS, Spach, Hist. Vég. Phan. x. 25 (1841) = **Osmites**, Linn. (Compos.)

BELLIDIUM, Bertol. Fl. Ital. ix. 816 (1853) = **Bellis**, Tourn. (Compos.).
bellidioides, Bertol. l. c. = Bellis sylvestris.
rotundifolium, Bertol. l. c. = Bellis rotundifolia.

BELLILLA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 83 = **BELLILLA, Rheede** = **Mussaenda**, Burm. (Rubiac.).

BELLINIA, Roem. & Schult. Syst. iv. 56 (1819) = **Saracha**, Ruiz & Pav. (Solanac.).
biflora, Roem. & Schult. l. c. 668 = S. procumbens.
contorta, Roem. & Schult. l. c. = S. contorta.
dentata, Roem. & Schult. l. c. = S. dentata.
micrantha, Roem. & Schult. l. c. 690 = S. micrantha.

BELLINIA:—
peruviana, Steud. Nom. ed. I. 105 = *Saracha peruviana* (nomen delendum).
procumbens, Roem. & Schult. Syst. iv. 687 = S. procumbens.
pubescens, Roem. & Schult. l. c. 689 = S. pubescens.
punctata, Roem. & Schult. l. c. 688 = S. punctata.
umbellata, Roem. & Schult. l. c. = S. umbellata.

BELLIOPSIS, Pomel, Mat. Fl. Atl. 7 (1860) = **Bellium**, Linn. (Compos.).
cordifolia, Pomel, Nouv. Mat. Fl. Atl. 286.—Algeria.
rotundifolia, Pomel, Mat. Fl. Atl. 7 = Bellium rotundifolium.

BELLIS, [Tourn.] Linn. Gen. ed. I. 856 (1737). *COMPOSITÆ*, Benth. & Hook. f. ii. 265.
ASTRANTHIUM, Nutt. in Trans. Am. Phil. Soc. ser II. vii. (1841) 312.
KYBERIA, Neck. Elem. i. 42 (1790).
SEUBERTIA, H. Watt. in Hook. Lond. Journ. Bot iii. (1844) 602.
aculeata, Labill. Nov. Holl. Pl. ii. 55. t. 206 = Brachycome Billardieri.
alpina, Hegetschw. Fl. Schw. 831 = perennis.
americana, Mirb. ex Stead. Nom. ed. II. i. 195 = integrifolia.
annua, *Linn. Sp. Pl.* 887.—Reg. Mediterr.
arabica, *Roxusch. Nom.* ed. III. 246, nomen.—Arabia.
armena, *Boiss. Fl. Orient.* iii. 174.—Armenia.
atlantica, *Boiss. & Reut. Pugill.* 54.—Hispan.
azorica, *Hochst. in Seub. Fl. Azor.* 31. t. 9. fig. 1–3.—Ins. Azores.
Bernardi, *Boiss. & Reut. Pugill.* 56.—Hispan.
caerulescens, *Cass. & Bal. in Bull. Soc. Bot. Fr.* xxii (1875) 60, nomen.—Marocco.
campestris, *Vell. Fl. Flum.* viii. t. 194.—Bras.
ciliaris, Labill. Nov. Holl. Pl. ii. 56. t. 209 = Brachycome ciliaris.
ciliata, *Rafin. New Fl. Am.* ii. 24.—Am. bor.
cordifolia, *Willk. Illustr. Fl. Hisp.* 4. 47.—Hispan.
dentata, DC. Prod. v. 304 = annua.
drosrarifolia, Gouan, Illustr. 69 = Bellium bellidioides.
dubia, *Spreng. Syst.* iii. 582.—Hab. ?
gelida, Banks & Soland. ex Hook. Fl. N. Zel. i. 125 = Lagenophora Forsteri.
glandulosa, Klatt, in Leopoldina, xix. (1883) 32 = Brachycome glandulosa.
graminea, Labill. Nov. Holl. Pl. ii. 54. t. 204 = Brachycome graminea.
hirta, Host, Fl. Austr. ii. 501 = sylvestris!
hortensis, Mill. Gard. Dict. ed. VIII. n. 8 = perennis.
hybrida, Tenore, Fl. Nap. v. 238. t. 194 = perennis.
integrifolia, DC. in Lam. Encyc. v. 7 = perennis.
integrifolia, *Michx. Fl. Bor. Am.* ii. 131.—Am. bor.
javanica, Blume, Bijdr. 917 = Myriactis Wightii.
longifolia, *Boiss. & Heldr. Diagn.* Ser. I. xi. 1.—Ins. Creta.
longifolia, Orph. ex Nym. Consp. 390 = sylvestris.
magellanica, DC. in Lam. Encyc. v. 7 = Lagenophora Commersonii.
maritima, *A. Gray, Pl. Wright.* i. 98.—Mexic.
microcephala, Lange, in Kjoeb. Vidensk. Meddel. (1861) 66 = annua.
muscarioides, Klatt, in Leopoldina, xix. (1883) 32 = Brachycome mossambicensis.
nutans, *Rafin. New Fl. Am.* ii. 23.—Am. bor.
pappulosa, Boiss. ex DC. Prod. vii. 276 = sylvestris.
parviflora, *Rafin. New Fl. Am.* ii. 23.—Am. bor.
pedunculata, *Vell. Fl. Flum.* viii. t. 125.—Bras.
perennis, *Linn. Sp. Pl.* 886.—Europ. occ.
pilosa, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 126 = Lagenophora lanata.
pilosa, Blume, Bijdr. 917 = Myriactis pilosa.
prostrata, *Pomel, Nouv. Mat. Fl. Atl.* 287.—Algeria.
racemosa, Stend. Nom. ed. I. 105, sphalm. = seq.
remosa, Jacq. Enum. Pl. Carib. 28 = Eclipta punctata.
ramosa, Lam. Fl. Fr. ii. 122 = annua.
repens, Lam. l. c. = annua.
rotundifolia, *Boiss. & Reut. Pugill.* 55.—Algeria.
scandens, *Vell. Fl. Flum.* viii. t. 126.—Bras.
scoposa, Gilib. Fl. Lituan. i. 213 = perennis.
stipitata, Labill. Nov. Holl. Pl. ii. 55. t. 105 = Lagenophora Billardieri.
sylvestris, *Cyrill. Pl. Rar.* ii. 29. t. 4.—Reg. Mediterr.
valentina, *Pomel, Nouv. Mat. Fl. Atl.* 85.—Algeria.
xanthocomoides, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 118.—Mexic.

BELLIUM, Linn. Mant. 167 (1771). *COMPOSITAE*, Benth. & Hook. f. ii. 365.
Bellidiopsis, Pomel, Mat. Fl. Atl. 7 (1860).
bellidioides, Desf. Fl. Atlant. ii. 279 = Bellis dentata.
bellidioides, *Linn. Mant.* 285.—Reg. Mediterr.
bellidioides, Urv. in Mém. Soc. Linn. Par. i. (1822) 368 = minutum.
bellioides, G. Don, in Sweet, Flow. Gard. ii. 175 = bellidioides.
cordifolium, Kunze, in Flora, xxix. (1846) 703 = Bellis rotundifolia.
crassifolium, *Moris, Pl. Sard. Elench.* i. 26.—Sardinia.
dentatum, Viviani, Fragm. 8. t. 10 = Bellis annua.
droseraefolium, Labill. ex Steud. Nom. ed. I. 105 = bellidioides.
gigantеum, Cass. in Dict. Sc. Nat. xxxvii. 456 = rotundifolium.
minutum, *Linn. Mant.* 286.—Archip. Graec.
nivale, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 383 = bellidioides.
pappulosum, Kunze, in Flora, xxix. (1846) 703 = Bellis sylvestris.
pumilum, Salisb. Prod. 209 = bellidioides.
rotundifolium, *DC. Prod.* v. 308.—Algeria.

BELLOA, Remy, in C. Gay, Fl. Chil. iii. 336. t. 38 (1849) = **Gnaphalium**, *Linn.* (Compos.).
chilensis, Remy, l. c. 337. t. 38. f. 2.—Chili.
subspicata, Wedd. Chlor. And. i. 159.—Bolivia.

BELLONIA, [Plum.] Linn. Gen ed. I. 50 (1737). *GESNERACEAE*, Benth. & Hook. f. ii. 996.
BELONIA, Adans. Fam. ii. 158 (1763).
aspera, *Linn. Sp. Pl.* 172.—Ind. occ.
spinosa, *Sw. Prod. Veg. Ind. Occ.* 41.— Ins. S. Doming.

BELLOTA, C. Gay, Fl. Chil. v. 297. t. 59 (1849) = **Cryptocarya**, R. Br. (Laurin.).
Miersii, C. Gay, l. c. 298.—Chili.
nitida, Phil. in Linnaea, xxix. (1857–58) 39 = C. nitida.
pauciflora, Phil. in Anal. Univ. Chil. (1865) i. 649.—Chili.

BELLOTA, A. Rich. ex Phil. in Linnaea, xxix. (1857–58) 40 = **Boldu**, Nees (Laurin.).

BELLUCCIA, Adans. Fam. ii. 344 (1763) = **Ptelea**, Linn. (Rutac.).

BELLUCIA, Meisn. Gen. Comm. 46, in syn. = praec.

BELLUCIA, Neck. Elem. ii. 142 (1790). *MELASTOMACEAE*, Benth. & Hook. f. i. 763.
Apatitia, Desv. in Hamb. Prod. Fl. Ind. Occ. 42 (1825).
Axinanthera, Karst. in Linnaea, xxx. (1859–60) 157.
Ischyranthera, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 102.
Werrea, J. F. Gmel. Syst. 820 (1790).
arborescens, Baill. Hist. Pl. vii. 34 = Loreya arborescens.
Aublettii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 102.—Guiana.
Aublettii, Seem. Bot. Voy. Herald, 122 = superba.
Axinanthera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 142.—N. Granat.
brasiliensis, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 104.—Bras.
Cacatin, *Sagot, l. c.* Sér. VI. xv. (1883) 382.—Guiana.
grossularioides, Triana, in Trans. Linn. Soc. xxviii. (1871) 141 = Aublettii.
Hostmannii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 103.—Guiana.
macrophylla, Triana, in Trans. Linn. Soc. xxviii. (1871) 143 = superba.
multiflora, Karst. in Linnaea, xxx. (1859–60) 158 = Aublettii.
nervosa, Rafin. Sylva Tellur. 93 = Blakea trinervia.
pentamera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 105.—Perv.
pluriculis, Spruce, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 142 = Hostmannii.
riparia, Spruce, l. c. = Hostmannii.
superba, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 104.—Mexic.

BELLYNKXIA, Muell. Arg. in Flora, lviii. (1875) 465 = **Appunia**, Hook. f. (Rubiac.).
angulata, Muell. Arg. l. c. = A. angulata.

BELMONTIA, E. Mey. Comm. Pl. Afr. Austr. 188 (1837). *GENTIANEAE*, Benth. & Hook. f. ii. 804.
Exochaenium, Griseb. in DC. Prod. ix. 56 (1845).
Parasia, Rafin. Fl. Tellur. iii. 78 (1836).
cordata, *E. Mey. l. c.*—Afr. austr.
gracilis, *Wolw. in Trans. Linn. Soc.* xxvii. (1869) 47. —Afr. trop.
grandis, *E. Mey. Comm. Pl. Afr. Austr.* 183.—Afr. austr.
Ohlendorffii, *Griseb. in DC. Prod.* ix. 54.—Afr. austr.
spathulata, *E. Mey. Comm. Pl. Afr. Austr.* 183.—Afr. austr.

BELOAKON, Rafin. Fl. Tellur. iii. 87 (1836) = **Phlomis**, Linn. (Labiat.).
luteum, Rafin. l. c. = P. fruticosa.
tomentosum, *Rafin. l. c.* = P. Samia.

BELOANTHERA, Hassk. in Flora, xxv. (1842) Beibl. ii. 55. *'GENUS DUBIUM.'*
oppositifolia, *Hassk. l. c.*—Java.

BELOÈRE, Shuttlew. ex A. Gray, Pl. Wright. i. 21 (1852) = **Abutilon**, Gaertn. (Malvac.).
cistiflora, Shuttlew. ex A. Gray, l. c. = A. indicum.
crispa, Shuttlew. ex A. Gray, l. c. in nota = A. crispum.

BELONIA, Adans. Fam. ii. 158 (1763) = **Bellonia**, Plum. (Gesner.).

BELONITES, E. Mey. Comm. Pl. Afr. Austr. 187 (1837) = **Psohypodium**, Lindl. (Asclep.).
bispinosa, *h.* Mey. l. c. 188 = P. bispinosum.
succulenta, E. Mey. l. c. 187 = P. sacciantum.

BELONOPHORA, Hook. f. Ic. Pl. t. 1127 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 109.
coffeoides, *Hook. f. l. c.*—Afr. trop.

BELOPERONE, Nees, in Wall. Pl. As. Rar. iii. 76, 102 (1832). *ACANTHACEAE*, Benth. & Hook. f. 1110.
Beloperonides, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 162.
Nuestera, Kegel, Gartenfl. vi. (1857) 345.
Simonisia, Nees, in Mart. Fl. Bras. ix. 144. t. 28 (1847).
Amherstiae, Nees, l. c. 102 = Dianthera nodosa.
appendiculata, *Nees, in DC. Prod.* xi. 423.—Peruv.
atropurpurea, *Nees, in Mart. Fl. Bras.* ix. 138.—Bras.
aurantiaca, *Benth. & Hook. f. Gen.* ii. 1111.—Am. mexic.
auriculata, *Nees, in Mart. Fl. Bras.* ix. 136.—Bras.
bullata, *Nees, l. c.* 141.—Bras.
californica, *Benth. Bot. Voy. Sulph.* 38.—Am. bor. occ.
calycina, Nees, in Hook. Lond. Journ. Bot. iv. (1845) 637 = Dianthera calycina.
carnea, *Nees, in Jacq. Ecleg.* ii. 2, in obs.—Venezuela.
carthaginensis, Benth. & Hook. f. Gen. ii. 1111 = violacea.
ciliata, Hook. Bot. Mag. t. 5868 = Dianthera ciliata.
ciliata, *Nees, in DC. Prod.* xi. 422.—Peruv.
comosa, *Nees, l. c.* 418.—Mexic.
cordata, Mart. ex Nees, l. c. 361 (= *Sericographis cordata*).—Bras.
cristata, Mart. ex Nees, l. c. 419 = Dianthera nodosa.
denudata, *Nees, l. c.* 423.—Peruv.
dicliptereoides, *Nees, in Mart. Fl. Bras.* ix. 136.—Bras.
Eustachyana, *Benth. & Hook. f. Gen.* ii. 1111.—Ind. occ.
fimbriata, *Nees, in DC. Prod.* xi. 416.—Mexic.
formosa, *Fenzl, in Jacq. Eclog.* ii. *in Ind. (Justicia, t. 192).*—Venezuela.
fragilis, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 140.—Bras.
fulgida, Hassk. Cat. Hort. Bog. Alt. 180 = Climacanthes Burmanni.
glandulosa, *Nees, in DC. Prod.* xi. 414.—Bras.

BELOPERONE:—
glomerata, *Nees, in Mart. Fl. Bras.* ix. 137.—Bras.
hirsuta, *Nees, l. c.* 143.—Bras.
involucrata, *Nees, l. c.*—Bras.
Lamarckiana, *Nees, in DC. Prod.* xi. 415.—Ind. occ.
lanceolata, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 141. —Bras.
longipetiolata, *Nees, l. c.* 137.—Bras.
macrantha, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 516.—Mexic.
Mathewsiana, *Nees, in DC. Prod.* xi. 781.—Peruv.
Meyeniana, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 381.—Bras.
microcarchya, *Nees, in Mart. Fl. Bras.* ix. 137.—Bras.
mollis, *Nees, l. c.* 142.—Bras.
monticola, *Nees, l. c.*—Bras.
nemorosa, Nees, in DC. Prod. xi. 415 = Adhatoda tetramera.
nodicaulis, *Nees, in Mart. Fl. Bras.* ix. 140.—Bras.
oblongata, *Lindl. Bot. Reg.* t. 1657.—Bras.
oblongata, Nees, in Wall. Pl. As. Rar. iii. 102 = plumbaginifolia.
obtusifolia, Nees, l. c. 339 = ovata.
plumbaginifolia, *Nees, in Wall. Pl. As. Rar.* iii. 102.—Bras.
portoricensis, *Nees, in DC. Prod.* xi. 414.—Porto Rico.
pulchella, Linden, Cat. n. 17 (1862) 2; ex Regel, Gartenfl. xiv. (1865) 289 = violacea.
Riedeliana, *Benth. & Hook. f. Gen.* ii. 1111.—Bras.
Rohrii, *Nees, in DC. Prod.* xi. 415.—Am. austr.
rosea, *Benth. & Hook. f. Gen.* ii. 1111.—Bras.
Schomburgkiana, *Nees, in Hook. Lond. Journ. Bot.* iv. (1845) 636.—Guiana.
scorpioides, *Nees, in DC. Prod.* xi. 422.—Bras.
Selloviana, *Nees, in Mart. Fl. Bras.* ix. 136.—Bras.
spathulata, *Nees, l. c.*—Bras.
sphaerosperma, *Benth. & Hook. f. Gen.* ii 1111.—Ind. occ.
strobilacea, Mart. ex Nees, in DC. Prod. xi. 358 (= *Amphiscopia strobilacea*).—Bras.
surinamensis, *Miq. Stirp. Surin. Sel.* 130.—Guiana.
trifoliata, *Nees, in Mart. Fl. Bras.* ix. 141.—Bras.
villosa, Mart. ex Nees, in DC. Prod. xi. 417 = hirsuta.
violacea, *Planch. & Linden, Hort. Linden.* i. 5. t. 3.—Mexic.

BELOPERONIDES, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 162 = **Beloperone**, Nees (Acanth.).
macrantha, Oerst. l. c. = Beloperone macrantha.

BELOSPIS, Rafin. Fl. Tellur. iii. 90 (1836) = **Salvia**, Tourn. (Labiat.).
lovigata, Rafin. l. c. 91 = S. involucrata.

BELOSTEMMA, Wall. ex Wight, Contrib. Bot. Ind. Or. 53 (1834) = **Tylophora**, R. Br. (Asclepiad.).
hirsutum, Wall. ex Wight, l. c. = T. Belostemma.

BELOSVNAPSIS, Hask. in Flora, liv. (1871) 289 = **Cyanotis**, D. Don (Commelin.).
brownsis, Hassk. l. c. = C. kewensis.

BELOTIA, A. Rich. Ess. Fl. Cub. 207. t. 21 (1845). *TILIACEAE*, Benth. & Hook. f. i. 253.
Adenodiscus, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) ii. 804.
Gallezotti, *Turcz. l. c.*—Mexic.
grevisefolia, *A. Rich. Ess. Fl. Cub.* 209. t. 22.—Cuba.
insignis, *Baill. Adansonia*, x. (1872) 182.—Mexic.

BELOU, Adans. Fam. ii. 408 (1763) = **Aegle**, Correa (Rutac.).

BELOVIA, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1859) 462 = **Suaeda**, Forsk. (Chenopod.).
baccifera, Moq. in DC. Prod. xiii. ii. 168 = S. setigera.
paradoxa, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1859) 462 = S. setigera.

BELTOKON, Rafin. Fl. Tellur. iii. 86 (1836) = **Origanum**, Tourn. (Labiat.).
Tournefortii, Rafin. l. c. = O. Tournefortii.

BELUTTA, Rafin. Fl. Tellur. iii. 39 (1836) = **Allmania**, R. Br. (Amarant.).
pedunculata, Rafin. l. c. = A. nodiflora.
sessilis, Rafin. l. c. = A. albida, pyramidalis.

BELVALA, Adans. Fam. ii. 285 (1763) = **Struthiola**, Linn. (Thymel.).
spicata, Rafin. Fl. Tellur. iv. 105 = S. virgata ?

BELVALIA, Delile, sec. Flore, xiii. (1830) ii. 455 = **Althenia**, Petit (Naiad.).
australis, Delile, l. c. = A. filiformis.

BELVEDERE, Clayt. ex Gronov. Fl. Virgin. 25 (1739) = **Galax**, Linn. (Diapens.).

BELVISIA, Desv. Journ. Bot. iv. (1814) 130 = **Napoleona**, Beauv. (Myrtac.).
caerulea, Desv. l. c. = N. imperialis.

BEMBECODIUM, Lindl. Veg. Kingd. 712 (1847) = BEMBYCODIUM, Kunze = **Athanasia**, Linn. (Compos.).

BEMBICIA, Oliver, in Hook. Ic. Pl. t. 1404. *SAMYDACEÆ.*
axillaris, *Oliver, l. c.*—Malaya.

BEMBICIUM, Mart. ex Baker, in Mart. Fl. Bras. v. ii. 341 (1876) = **Rupestorium**, Tourn. (Compos.).
gracile, Mart. ex Baker, l. c. 349 = E. kleinioides.

BEMBIX, Lour. Fl. Cochinch. 346 (1790). *MALPIGHIACEÆ* !—Cf. Benth. & Hook. f. Gen. i. 251.
tectoria, *Lour. l. c.*—Cochinch.

BEMBYCODIUM, Kunze, in Linnæa, xvi. (1842) 816 = **Athanasia**, Linn. (Compos.).
Athanasiæ, Kunze, l. c.—Afr. austr.

BEMSETIA, Rafin. Sylva Tellur. 12 (1838) = **Ixora**, Linn. (Rubiac.).
paniculata, Rafin. l. c. = I. stricta.

BENAUREA, Rafin. Fl. Tellur. ii. 77 (1836) = **Mussœnda**, Dum. (Campan.).
sempervirens, Rafin. l. c. 78 = M. aurea.

BENCOMIA, Webb & Berth. Phyt. Canar. ii. 10. t. 89 (1846). *ROSACEÆ*, Benth. & Hook. f. i. 624.
caudata, *Webb & Berth. l. c.* 11.—Ins. Canar. Moquiniana, *Webb & Berth. l. c.*—Ins. Canar.

BENEDICTA, Bernh. Syst. Verz. Erf. 108 (1800) = Carbenia, Adans. (Compos.).
officinalis, Bernh. l. c. 159 = C. benedicta.

BENINCASA, Savi, in Bibl. Ital. ix. 158 (1818); (Mem. sopra Piant. Cucurb. i. (1818) 6). *CUCURBITACEÆ*, Benth. & Hook. f. i. 824.
cerifera, *Savi, ll. cc.*—Ins. et Afr. trop. (cult).
cylindrica, Hort. ex Ser. in DC. Prod. iii. 308 = cerifera.
hispida, Cogn. in DC. Monog. Phan. iii. 518 = cerifera.

BENITZIA, Karst. in Linnæa, xxviii. (1856) 490 = Gymnosiphon, Blume (Burman.).
Poeppigiana, Karst. l. c. 491.—Peru.
sanguinea, Karst. l. c. 490.—Venezuela.

BENJAMINA, Vell. Fl. Flum. 98 (1825); ii. t. 139 = Dictyoloma, DC. (Simarub.).
alata, Vell. ii. cc. = D. incanescens.

BENJAMINIA, Mart. ex Benj. in Mart. Fl. Bras. x. 256 (1847) = QUINQUELOBUS, Benj. (= Utricularia ?).
Glabra, Benj. in Linnæa, xx. (1847) 761 = Limnophila Griffithii.
minor, Benj. l. c. (= *Q. minor*).—As. trop.
splendens, Benj. l. c. (= *Q. splendens*).—As. trop.
utriculariiformis, Mart. ex Benj. in Mart. Fl. Bras. x. 256 (= *Q. utriculariiformis*).—Bras.

BENKARA, Adans. Fam. ii. 85 (1763) = **Malastoma**, Burm.
galia, Rafin. Sylva Tellur. 98.—Hab. ?

BENNETTIA, DC. Prod. vi. 581, in syn. (1837) = BENNETTIA, S. F. Gray = **Saussurea**, DC. (Compos.).

BENNETTIA, Rafin. in Ser. Bull. Bot. i. (1830) 220 = Sporobolus, R. Br. (Gramin.).
juncea, Rafin. l. c. = S. junceus.

BENNETTIA, R. Br. in Benn. Pl. Jav. Rar. 249. t. 50 (1838) = **Galearia**, Zoll. (Euphorb.).
filiformis, R. Br. l. c. 251 = G. affinis.
aristifera, Muell. Arg. in DC. Prod. xv. ii. 1089 = G. stipitifera.
filiformis, Muell. Arg. in Linnæa, xxxiv. (1865–66) 204. —Ind. or.
Finlaysoniana, R. Br. in Benn. Pl. Jav. Rar. 251 = G. Finlaysoni.
fulva, Muell. Arg. in Linnæa, xxxiv. (1865–66) 205 = G. fulva.
Jackiana, R. Br. in Benn. Pl. Jav. Rar. 251 = G. Jackiana.
javanica, R. Br. l. c. 249 = G. pedicellata.
Lindleyana, Muell. Arg. in Linnæa, xxxiv. (1865–66) 205.—Ind. or.
pedicillata, R. Br. in Benn. Pl. Jav. Rar. 251 = G. pedicellata.
phlebocarpa, R. Br. l. c. = G. phlebocarpa.
splendens, Muell. Arg. in Linnæa, xxxiv. (1865–66) 205 = G. splendens.
subulata, Muell. Arg. in DC. Prod. xv. ii. 1089.—Ins. Penang.
Wallichii, R. Br. in Benn. Pl. Jav. Rar. 251 = G. Wallichii.

BENNETTIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 440 (1821) = **Saussurea**, DC. (Compos.).
alpina, S. F. Gray, l. c. = S. alpina.

BENNETTIA, Miq. Fl. Ind. Bat. i. ii. 105 (1859). *BIXINEÆ*, Benth. & Hook. f. i. 128.
Hornfeldii, *Miq. l. c.*—Java.

BENTECA, Adans. Fam. ii. 166 (1763) = **Ambelania**, Aubl. (Apocyn.).
odorata, Rafin. Sylva Tellur. 13 = A. acida.

BENTEKA, Adans. Fam. ii. 525 (1763).—Cf. præc. = Ambelania, Aubl. (Apocyn.).
Rheedii, Koem. & Schult. Syst. iv. 706 = 'Malabaria.'

BENTHAMIA, Lindl. Bot. Reg. t. 1579 (1833) = Cornus, Tourn.
fragifera, Lindl. l. c. = C. capitata.
japonica, Sieb. & Zucc. Fl. Jap. i. 88. t. 16 = C. Kousa.

BENTHAMIA, Lindl. Nat. Syst. ed. I. 241 (1830) = Amalnokia, Lehm. (1831) (Borragin.).
lycopsoides, Lindl. ex A. DC. Prod. x. 118 = A. lycopsoides.

BENTHAMIA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 37. t. 7 = **Habenaria**, Willd. (Orchid.).
latifolia, A. Rich. l. c. 38 = H. chloranthа.
spiralis, A. Rich. l. c. 39 = Spiranthes aracana.

BENTHAMANTHA, Alef. in Bonplandia, x. (1862) 264 = Cracca, Benth. (Legum.).
glabrescens, Alef. l. c. = C. glabrescens.
glandulifera, Alef. l. c. = C. glandulifera.
Grayi, Alef. l. c. = C. sericea.
mollis, Alef. l. c. = C. mollis.
ochroleuca, Alef. l. c. = C. ochroleuca.

BENTHAMIDIA, Spach, Hist. Vég. Phan. viii. 106 (1839) = Cornus, Tourn.
florida, Spach, l. c. 107 = C. florida.

BENTHAMIELLA, Speg. in Anal. Soc. Cient. Argent. xv. (1883) 109. *SOLANACEÆ.*
patagonica, *Speg. l. c.* 110.—Reg. Argent.

BENTHECA, Neck. Elem. ii. 84 (1790) = BENTECA, Adans. ! = Ambelania, Aubl. (Apocyn.).

BENTHEKA, Neck. ex A. DC. Prod. viii. 321, in syn. (1844) = **Willughbeia**, Roxb. (Apocyn.).

BENTINCKIA, Berry, ex Roxb. [Hort. Beng. 68 (1814) et] Fl. Ind. iii. 621 (1832). *PALMÆ*, Benth. & Hook. f. iii. 916.
KEPPLERIA, Mart. ex Endl. Gen. 251 (1838).
ceramensis, Miq. in Verh. Kon. Akad. Wetensch. xi (1868) n. v. 8 = Cyrtostachya ceramica.
ceramica, H. Wendl. in Kerch. Palm. 235 = præc. Condapanna, *Berry, ex Roxb.* [*Hort. Beng.* 68 *et*] *Fl. Ind.* iii. 621.—Ind. or.
nicobarica, *Becc. in Ann. Jard. Buitens.* ii. (1885) 165.—Ins. Nicob.
Rendu, Mart. Hist. Nat. Palm. iii. 316 = Cyrtostachys Renda.

BENZOIN, Hayne, Arzneigew. xi. t. 24 (1829) = **Styrax**, Linn.
officinale, Hayne, l. c. = S. Benzoin.

BENZOIN, Nees, in Wall. Pl. As. Rar. ii. 68 (1831) = Lindera, Thunb. (Laurin.).
austrinum, Nees, Syst. Laurin. 495 = L. Benzoin.
citriodorum, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 206, 78 (= *Aperula citriodora*).—Japon.
Diospyros, Sweet, Hort. Brit ed. III. 581 = L. melissæfolia.
geniculatum, G. Don, in Sweet, l. c. = L. geniculata.
glaucescens, Nees, ex Meisn. in DC. Prod. xv. i. 193 = Litsea glaucescens.
glaucum, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 205 = L. glauca.
melissæfolium, Nees, Syst. Laurin. 494 = L. melissæfolia.
Neesianum, Nees, in Wall. Pl. As. Rar. ii. 68 = L. Neesiana.
odoriferum, Nees, l. c. = L. Benzoin.
praecox, Sieb. & Zucc. in Abh. Akad. Muench. iv iii. (1846) 205 = L. praecox.
sericeum, Hook. f. & Thoms. ex Meisn. in DC. Prod. xv. i. 246 = L. heterophylla.
sericeum, Sieb. & Zucc in Abh. Akad. Muench. iv. iii (1846) 204 = L. sericea.
Thunbergii, Sieb. & Zucc. l. c. = L. umbellata.
trilobum, Sieb. & Zucc. l. c. 202 = L. triloba.

BENZOINA, Rafin. Sylva Tellur. 138 (1838) = **Styrax**, Linn.
vera, Rafin. l. c. = S. Benzoin.

BENZONIA, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 118 (1827). *RUBIACEÆ* ! Benth. & Hook. f. ii. 29.
corymbosa, *Schum. l. c.* 114.—Afr. trop.

BERARDIA, Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 360. t. 87 = **Nebelia**, Neck. (Bruniac.).
affinis, Brongn. l. c. 361 = N. affinis.
affinis, Sond. in Harv. & Sond. Fl. Cap. ii. 320.—Afr. austr.
angulata, Sond. l. c.—Afr. austr.
aspera, Sond. l. c. 321.—Afr. austr.
Dregeana, Sond. l. c.—Afr. austr.
fragarioides, Schlecht. in Linnæa, vi. (1831) 190.—Afr. austr
globosa, Sond. in Harv. & Sond. Fl. Cap. ii. 320.—Afr. austr.
laevis, E. Mey. ex Sond. l. c. 319.—Afr. austr.
microphylla, Sond. l. c. 320.—Afr. austr.
palearea, Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 361 = N. palearea.
phylicoides, Brongn. l. c. = N. phylicoides.
sphaerocephala, Sond. in Harv. & Sond. Fl. Cap. ii. 319.—Afr. austr.

BERARDIA, Vill. Prosp. 27 (1779). *COMPOSITÆ*, Benth. & Hook. f. ii. 474.
ARCTIO, Lam. Encyc. i. 285 (1783).
ARCTIUM, Lam. Fl. Fr. ii. 70 (1778).
VILARIA, Gouan. Mém. Mineral. Dauph. i. 108. ii. t. 19 (1779).
subacaulis, *Vill. l. c.* 28.—Gallia.

BERBERIDOPSIS, Hook. f. Bot. Mag. t. 5343 (1863). *BERBERIDEÆ*, Benth. & Hook. f. i. 44.
corallina, *Hook. f. l. c.*—Chili.

BERBERIS, [Tourn.] Linn. Gen. ed. I. 94 (1737).
BERBERIDEÆ, Benth. & Hook. f. i. 43.
MAHONIA, Nutt. Gen. Am. i. 211 (1818).
ODOSTEMON, Rafin. in Am. Monthly Mag. (1817) 192.
abortiva, Renault, Fl. Dép. Orne, 179 = vulgaris.
acanthifolia, Wall. ex Walp. Rep. i. 108 = nepalensis.
acida, Gilib. Exercit. i. 324 = vulgaris.
actinacantha, *Mart. ex Schult. f. Syst.* vii. 19.—Chili.
acutifolia, *Pranti, Ind. Sem. Wirceburg.* (1878) 7.—Hab.?
aetnensis, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 901 = vulgaris.
aetnensis, *Presl, Fl. Sic.* i. 98.—Cors.; Sardin.; Sicil.
affinis, G. Don, Gen Syst. i. 115 = aristata.
africana, *Hohenstr. & Ludw. ex Schult. f. Syst.* vii. 11.—Afr. austr.
agapetensis, *Lechl. Berb. Am. Austr.* 44.—Peruv.
alaica, Pall. Fl. Ross. ii. 41. t. 67 = sibirica.
amurensis, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1857) 260.—Manchuria.
Andrieuxii, *Hook. & Arn. in Bot. Beech. Voy.* 318, adnot.—Mexic.
angulosa, *Wall. Cat.* n. 1475.—Ind. or.
angustifolia, *Benth. Pl. Hartw.* 34, 270.—Mexic.
angustifolia, Roxb. Hort. Beng. [87]; Fl. Ind. ii. 188 = B. Lycium.
apyrena, Hort. ex C. Koch, Dendrol. i. 395 = vulgaris.
Aquifolium, *Pursh, Fl. Am. Sept.* i. 219. t. 4.—Am. bor.
arborescens, Hort. ex C. Koch, Dendrol. i. 393 = vulgaris.
aristata, Sims, Bot. Mag. t. 2549 = umbellata.
aristata, *DC. Syst.* ii. 8.—Ind. or.
articulata, *Lois. in Dict. Sc. Nat.* lxviii. 218 — Gallia.
asiatica, Griff. Ic. Pl. Ind. Or. iv. t. 648 = Wallichiana.
asiatica, *Roxb. ex DC. Syst.* ii. 18.—Reg. Himal.
asperma, Pgit. & Turp. in Duham. ed. Nov. iv. 12 = vulgaris.
atrovirens, Wall. ex G. Don, Gen. Syst. i. 117 = Wallichiana.
atroviridis, Steud. Nom. ed. II. i. 196 = Wallichiana.
aurahuascensis, *Lem. in Fl. des Serres*, Sér. I. iv. (1848) 334.—N. Granat.
aurea, Tausch, in Flora, xvii. (1834) 495 = vulgaris.
Banlei, *Fortune, in Gard. Chron.* (1850) 212.—Japon.
bidentata, *Lechl. Berb. Am. Austr.* 12.—Chili.
Bigelovii, Schrad. in Linnæa, xii. (1838) 362 = vulgaris.
boliviana, *Lechl. Berb. Am. Austr.* 21.—Bolivia.
brachybotrys, Edgew. in Trans. Linn. Soc. xx. (1851) 29 = vulgaris.
brachybotrys, *C. Gay, Fl. Chil.* i. 31.—Chili.
brachypoda, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 308.—China.
brevifolia, Hort. ex C. Koch, Dendrol. i. 398 = canadensis.
Buxensis, C. Koch, ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 253 = aristata.
buxifolia, *Lam. Illustr.* t. 253. f. 3; *Poir. Encyc.* viii. 619.—Reg. Magell.
callicbotrys, Bienert, ex Aitch. in Journ. Linn. Soc. xix. (1882) 151 = vulgaris.
canadensis, Guimp. Abbild. Fremd. Holz. 79. t. 63 = Guimpeli.
canadensis, Hayne, ex Steud. Nom. ed. II. i. 196 = spathulata.
canadensis, Mill. Gard. Dict. ed. VIII. n. 2 = vulgaris.
canadensis, *Pursh, Fl. Am. Sept.* 219.—Am. bor.
caragnaefolia, Banks, ex DC. Syst. ii. 18 = Caragana Chamlagu.
carinata, *Lechl. Berb. Am. Austr.* 41.—Peruv.
caroliniana, Sweet, Hort. Brit. ed. III. 19 = canadensis.
cerasina, *Schrad. in Linnæa*, xii. (1838) 379.—Am. bor.
ceratophylla, G. Don, Gen. Syst. i. 115 = aristata.
chilensis, *Gill. in Hook. Bot. Misc.* iii. (1833) 135.—Chili.
BERBERIS:—
chinensis, Hort. ex C. Koch, Dendrol. i. 408 = Guimpeli.
chinensis, Poir. Encyc. viii. 617 = sinensis.
Chitria, D. Don, Prod. Fl. Nep. 204 = aristata.
Chococa, *Schlecht. in Bot. Zeit.* xii. (1854) 652.—Mexic.
ciliaris, *Lindl. in Journ. Hort. Soc.* v. (1850) 5, in adnot.—Bolivia.
coccinea, Hort. ex C. Koch, Dendrol. i. 406 = aristata.
coletioides, *Lechl. Berb. Am. Austr.* 38.—Chili.
concinna, *Hook. f. Bot. Mag.* t. 4744.—Reg. Himal.
conferta, *H. B. & K. Nov. Gen. et Sp.* v. 69. t. 480.—Ecuador.
congestiflora, *C. Gay, Fl. Chil.* i. 75. t. 8.—Chili.
consanguinea, *Fortune, in Journ. Hort. Soc.* vii. (1852) 226.—China.
coriacea, *St. Hil. Fl. Bras. Mer.* i. 46.—Bras.
coriaria, Royle, ex Lindl. Bot. Reg. xxvii. (1841) t. 46 = aristata.
corymbosa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 135.—Ins. Juan Fernand.
crataegina, DC. Syst. ii. 9 = vulgaris.
crenulata, Schrad. in Linnæa, xii. (1838) 362 = vulgaris.
cretica, Falk, Beitr. ii. 164 = sibirica.
cretica, Linn. Sp. Pl. 331 = vulgaris.
cretica, Thunb. Fl. Jap. 146 = sinensis.
cretica, Viv. Fl. Cors. 5 = actnensis.
crispa, *C. Gay, Fl. Chil.* i. 86.—Chili.
cuneata, *DC. Syst.* ii. 16.—Am. austr.
cuneata, Hort. ex C. Koch, Dendrol. i. 415 = empetrifolia.
Darwini, *Hook. Ic. Pl.* t. 672.—Chili.
dasystachya, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 306.—China.
dasyrica, *Lodd. ex Lond. Hort. Brit.* 189.—Davuria.
dealbata, *Lindl. Bot. Reg.* t. 1750.—Mexic.
declinata, *Schrad. in Linnæa*, xii. (1838) 368.—China?
densa, Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat. Sér. VI. xvii. (1862) 54 = rigidifolia.
densiflora, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 9 = vulgaris.
densiflora, *Rafin. Sylva Tellur.* 68.—Hab.?
dentata, Tausch, in Flora, xxi. (1838) 729 = vulgaris.
diaphana, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 306.—China.
diffusa, *C. Gay, Fl. Chil.* i. 80.—Chili.
discolor, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 287.—Venezuela.
discolor, Willd. ex Steud. Nom. ed. II. i. 197 = rigidifolia.
diversifolia, *Steud. l. c.*—Reg. Argent.
dulcis, Hort. ex C. Koch, Dendrol. i. 395 = vulgaris.
dulcis, Sweet, Brit. Flow. Gard. Ser. I. ii. t. 100 = buxifolia.
dumetorum, Gouan, Fl. Monsp. 312 = vulgaris.
edulis, Hort. ex C. Koch, Dendrol. i. 395 = vulgaris.
Ehrenbergii, *Kunze, in Linnæa*, xx. (1847) 45.—Mexic.
elegans, Hort. ex C. Koch, Dendrol. i. 409 = aristata.
emarginata, Willd. Enum. Hort. Berol. 895 = vulgaris.
empetrifolia, *Lam. Illustr.* t. 253. f. 4.—Reg. Magellan.
fascicularis, Sims, Bot. Mag. t. 2396 = pinnata.
fasciculata, Schult. f. Syst. vii. 17 = pinnata.
Fendleri, *A. Gray, Pl. Fendl.* 5.—Am. bor. occ.
feroz, *C. Gay, Fl. Chil.* i. 80.—Chili.
ferruginea, *Lechl. Berb. Am. Austr.* 9.—Bolivia.
Fischeri, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
flexuosa, *Ruiz & Pav. Fl. Per.* iii. 52. t. 281. f. A.—Peruv.
floribunda, Wall. Cat. n. 1474; et G. Don, Gen. Syst. i. 115 = aristata.
floribunda, Hort.—Cf. Gard. Chron. (1869) 512 = B. Aquifolium.
florida, *Phil. in Linnæa*, xxxiii. (1864–65) 5.—Chili.
Fortunei, *Lindl. in Journ. Hort. Soc.* i. (1846) 231, 300.—China.
frustrifolia, *Hook. Ic. Pl.* t. 329.—Mexic.
Fremontii, *Torr. in Bot. Mex. Bound.* 30.—Am. bor. occ.
glauca, Benth. Pl. Hartw. 195. n. 710 = Jamesoni.
glauca, *DC. Syst.* ii. 19; *H. B. & K. Nov. Gen. et Sp.* v. 71.—N. Granat.
BERBERIS:—
glauca, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
glaucocarpa, *St. Hil. Fl. Bras. Mer.* i. 45.—Bras.
globosa, *Benth. Pl. Hartw.* 158.—N. Granat.
glomerata, *Hook. & Arn. Bot. Beech. Voy.* 5.—Chili.
glumacea, Spreng. Syst. ii. 130 = nervosa.
Gondoti, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 52.—N. Granat.
gracilis, Hort. ex Lindl. in Journ. Hort. Soc. v. (1850) 13 = aristata.
gracilis, *Harbw. ex Benth. Pl. Hartw.* 34. n. 271.—Mexic.
gracillima, C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 253 = aristata.
grandiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii (1854) ii. 287.—Ecuador.
gratissima, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 180.—Himal.
Grevilleana, *Gill. in Hook. Bot. Misc.* iii. (1833) 136.—Chili.
Griesbachii, *Lechl. Berb. Am. Austr.* 34.—Chili.
Gullache, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 54.—N. Granat.
Guimpeli, *C. Koch & Bouché, in App. Ind. Hort. Berol.* (1854) 13.—Am. bor.
Haenkeana, *Presl, ex Schult. f. Syst.* vii. 1616.—Chili.
Hartwegi, *Benth. Pl. Hartw.* 34. n. 272.—Mexic.
heterophylla, Hort. ex C. Koch, Dendrol. i. 395 = vulgaris.
heterophylla, *Juss. in Poir. Encyc.* viii. 622.—Reg. Magellan.
heteropoda, Schrenk, Enum. Pl. Nov. 109 = vulgaris.
hispanica, Boiss. & Reut. Pugill. Pl. Nov. (1852) 8 = vulgaris.
Hookeri, Hort. ex Lem. Illustr. Hortic. vi. (1859) t. 207 = Wallichiana.
horrida, *C. Gay, Fl. Chil.* i. 84.—Chili.
horrida, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 42 = Wallichiana.
hypoleuca, Lindl. in Journ. Hort. Soc. ii. (1847) 246 = aristata.
iberica, Sweet, Hort. Brit. ed. I. 18 = vulgaris.
ilicifolia, Bert. ex Steud. Nom. ed. II. i. 197 = glomerata, stipulacea.
ilicifolia, Booth, ex Steud. l. c. = vulgaris.
Ilicifolia, *Forst. in Comm. Gott.* ix. (1789) 28.—Reg Magellan.
ilicifolia, Hort. ex C. Koch, Dendrol. i. 402 = actnensis.
ilicifolia, Roxb. in As. Res. vi. (1799) 357 = asiatica.
ilicifolia, Scheele, in Linnæa, xxi. (1848) 591 = trifoliolata.
Ilicina, *Hemsl. Biol. Centr. Am. Bot.* i. 23.—Mexic.
inermis, *Perrs. Syn.* i. 387.—Reg. Magell.
innominata, Klein. Dec. Rar. Pl. Theb. 18 = vulgaris.
insignis, *Hook. f. & Thoms. Fl. Ind.* i. 226.—Reg. Himal.
integerrima, Bunge, in Linnæa, xviii. (1844) 149 = vulgaris.
integerrima, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
intermedia, Hort. ex C. Koch, l. c. 404 = Guimpeli.
intermedia, Hort. Angl.—Cf. Gard. Chron. (1869) 612 = nepalensis.
irritabilis, Salisb. Prod. 218 = vulgaris.
Jacquini, Hort. ex C. Koch, Dendrol. i. 394 = vulgaris.
Jamesoni, *Türcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 286.—Ecuador.
Jamesonii, *Lindl. in Journ. Hort. Soc.* v. (1850) 8; *Lem. Illustr. Hortic.* vi. (1859) t. 201.—Ecuador.
japonica, *R. Br. in Tuckey, Congo Exped. App.* 33.—Japon.
japonica, Lindl. ex Lindl. & Paxt. Flow. Gard. i. (1850–51) 11 = Banlei.
kamaonica, *Rupr. Sert. Tianschk.* 38.—As. temp.
Klotzschiana, *Warcz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 393, nomen.—Bolivia.
kumaonensis, Royle, Illustr. Bot. Himal. 64 = vulgaris.
Knightii, Hort. ex C. Koch, Dendrol. i. 413 = Banlei.
Lagenaria, Pohr. Encyc. viii. 619 = ilicifolia.
lanceolata, *Benth. Pl. Hartw.* 34.—Mexic.
latifolia, *Ruiz & Pav. Fl. Per.* iii. 52. t. 283.—Peruv.
lanuina, *Thunb. Fl. Bras.* i. 8.—Bras.

BERBERIS:—
laxa, *Rafin. Sylva Tellur.* 68.—Hab.?
laxiflora, Schrad. in Linnæa, xii. (1838) 367 = vulgaris.
Laschenaultiana, Wall. Cat. n. 1479 = nepalensis.
Ligonii, Hort. ex C. Koch, Dendrol. ii. 411 = concinna.
linearifolia, *Phil. in Linnæa*, xxviii. (1856) 663.—Chili.
littoralis, *Phil. Fl. Atac.* 7.—Chili.
littoralis, *Rupr. Fl Cauc.* 49, in textu.—Reg. Caucas.
loxensis, *Benth. Pl. Hartw.* 126.—Peruv.
lucida, *Schrad. in Linnæa*, xii. (1838) 368.—Iberia.
lutea, *Ruiz & Pav. Fl. Per.* iii. 51. t. 280.—Ecuador.
Lycium, Hort. ex C. Koch, Dendrol. i. 393 = vulgaris.
Lycium, *Royle, Illustr. Bot. Himal.* 64.—Reg. Himal.
macracantha, Hort. ex C. Koch, Dendrol. i. 400 = canadensis.
macracantha, Schrad. in Linnæa, xii. (1838) 366 = vulgaris.
macrocarpa, Schrad. l. c. 370 = canadensis.
macrophylla, Hort. ex Lindl. in Journ. Hort. Soc. v. (1850) 4 = Wallichiana.
macrophylla, Hort. ex C. Koch, Dendrol. i. 408 = asiatica.
mactosepala, *Hook. f. & Thoms. Fl. Ind.* i. 228.—Reg. Himal.
macrotheca, Hort. ex C. Koch, Dendrol. i. 400 = canadensis.
madrensis, *Lowe, in Hook. Kew Journ.* viii. (1856) 289.—Ins. Madera.
magellanica, Hort. ex C. Koch, Dendrol. i. 414 = Bealei.
marginata, *C. Gay, Fl. Chil.* i. 88.—Chili.
marginata, Hort. ex C. Koch, Dendrol. i. 394 = vulgaris.
Maximowiczii, Regel, Gartenfl. (1872) 288 = vulgaris.
melanocarpa, *Schau. Jahrb. Ges. Vat. Cult.* xxviii. 180.—Am. bor.
Micrii, Buch.-Ham. in D. Don, Prod. Fl. Nep. 205 = nepalensis.
microcarpa, *Warcz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 295.—Bolivia.
microphylla, F. G. Dietr. Vollst. Lex. Gaertn. ii. 184 = vulgaris.
microphylla, Forst. in Comm. Gott. ix. (1789) 29 = buxifolia.
microphylla, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
minor, Forst. ex DC. Syst. ii. 15 = buxifolia.
mitis, Schrad. in Linnæa, xii. (1838) 371 = vulgaris.
monosperma, Hort. ex C. Koch, Dendrol. i. 405 = sinensis.
monosperma, *Ruiz & Pav. Fl. Per.* iii. 52.—Peruv.
montana, *C. Gay, Fl. Chil.* i. 90.—Chili.
mornensis, *Schult. f. Syst.* vii. 17.—Mexic.
multiflora, *Benth. Pl. Hartw.* 124.—Peruv.
mutabilis, *Phil. in Anal. Univ. Chil.* (1872) 665.—Chili.
neglecta, *Hort. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 284.—Hab.?
nepalensis, Hort. ex C. Koch, Dendrol. i. 393 = vulgaris.
nepalensis, *Spreng. Syst.* ii. 120.—Reg. Himal.
nervosa, *Pursh, Fl. Am. Sept.* 219. t. 5.—Am. bor. occ.
Neubertii ×, Hort. ex Lem. Illustr. Hortic. i. (1854) Misc. 111. t. c.
nigra, *Rafin. Sylva Tellur.* 67.—Am. bor.
nitens, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
nitens, Schrad. in Linnæa, xii. (1838) 382 = vulgaris.
nitida, Willd. ex Steud. Nom. ed. II. i. 197 = quinquenervia.
Nummularia, Bunge, in Linnæa, xvili. (1844) 149 = vulgaris.
obvata, Schrad. l. c. xii. (1838) 385 = vulgaris?
orthobotrys, *Bienert, ex Attch. in Journ. Linn. Soc.* xix. (1882) 150.—Persia.
pallida, *Hartw. ex Benth. Pl. Hartw.* 84.—Mexic.
paniculata, Ell. B. & K. ex DC. Syst. ii. 10 = glauca.
paniculata, *Ruiz. ex DC. l. c.* 12.—Peruv.
paniculata, *Hemsl. Biol. Centr. Am. Bot.* i. 94.—Costa-Rica.
paniculata, *Phil. in Anal. Univ. Chil.* (1872) 664.—Chili.

BERBERIS:—
parvifiora, *Lindl. in Journ. Hort. Soc.* ii. (1847) 243. —Am. austr.
paucifiora, Salisb. Prod. 218 = vulgaris.
Pearcei, *Phil. in Linnæa*, xxxiii. (1864–65) 4.—Chili.
petiolaris, Wall. ex G. Don, Gen. Syst. i. 116 = aristata.
pichinchensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 286.—Ecuador.
pichinchensis, *Weddl. Chlor. And.* ii. 295 (= præc.?).—Ecuador.
pinnata, Banks, ex DC. Syst. ii. 20 = B. Aquifolium.
pinnata, Buch. ex DC. l. c. 21 = nervosa.
pinnata, H. B. & K. Nov. Gen. et Sp. v. 434 = moranensis.
pinnata, *Lag. Elench. Hort. Madr.* (1803) 6.—Mexic.
pinnata, Roxb. Fl. Ind. ii. 184 = nepalensis.
pinifera, Rafin. Sylva Tellur. 66 = canadensis.
polymorpha, *Phil. in Linnæa*, xxviii. (1856) 664.—Chili.
polyspatala, *Phil. in Anal. Univ. Chil.* (1872) 666.—Chili.
provincialis, Audib. ex Schrad. in Linnæa, xii. (1838) 381 = vulgaris.
psilopoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 285.—N. Granat.
purpurea, *Rafin. Sylva Tellur.* 67.—Hab.?
quindiuensis, *H. B. & K. Nov. Gen. et Sp.* v. 70. t. 432. —Ecuador.
racemosa, Stokes, Bot. Mat. Med. ii. 296 = vulgaris.
rariflora, *Lechl. Berb. Am. Austr.* 33.—Bolivia.
repens, Lindl. Bot. Reg. t. 1176 = B. Aquifolium.
reticulata, *Steud. Nom.* ed. II. i. 197.—Chili.
retinervia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 52.—N. Granat.
revoluta, Sm. ex DC. Syst. ii. 16 = empetrifolia.
rigida, Willd. ex Steud. Nom. ed. II. i. 197 = vulgaris.
rigidifolia, *H. B. & K. Nov. Gen. et Sp.* v. 70. t. 431. —Am. austr.
Roemeriana, Schreib, in Linnæa, xxii. (1849) 352 = trifoliata.
rotundifolia, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 68.—Chili.
rubra, Poit. & Turp. in Duham. ed. Nov. iv. 12 = vulgaris.
rubra, *Lam. Illustr.* t. 259. f. 2.—Reg. Argent.
sanguinea, Hort. ex C. Koch, Dendrol. i. 394 = vulgaris.
sanguinolenta, Hort. ex C. Koch, l. c. = vulgaris.
sanguinolenta, *Schrad. in Linnæa*, xii. (1838) 378.—Hab.?
saxicola, *Lechl. Berb. Am. Austr.* 42.—Peruv.
Schiedeana, *Schlecht. in Bot. Zeit.* xii. (1854) 654.—Mexic.
serotina, *Lange, Ind. Sem. Haun.* (1872) 5, adnot.—Hab.?
serratifolia, Hort. ex C. Koch, Dendrol. i. 406 = aristata.
serratodentata, *Lechl. Berb. Am. Austr.* 16.—Chili.
serrulata, Rafin. Sylva Tellur. 68 = canadensis.
sibirica, Hort. ex Schult. f. Syst. vii. 2 = vulgaris.
sibirica, *Pall. Fl. Ross.* ii. 42. t. 67.—Sibir. altaic.
sicula, Jan, Elench. ex Schult. f. Syst. vii. 3 = actinensis.
Siebo1dii, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 69.—Japon.
sinensis, *Desf. Tabl. ed.* I. (1804) 150.—China.
sinensis, DC. Syst. ii. 6 = Guimpelii.
spathulata, *Schrad. in Linnæa*, xii. (1838) 376.—Am. bor.
sphærocarpa, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 876 = vulgaris.
spinosa, Comm. ex Decne. in Bot. Voy. Astrol. & Zél. 65 = buxifolia.
spinulosa, *Sr. Hil. Fl. Bras. Mer.* i. 45.—Bras.
stenophylla, *Hance, in Journ. Bot.* xx. (1882) 257.—China.
stenophylla ×, Hort.—Cf. Gard. Chron. (1864) 460; (1868) ii. 613; (1882) ii. 466; C. Koch, Dendrol. i. 413 = Darwinii.
stipulacea, *Steud. Nom.* ed. II. i. 197.—Chili.

BERBERIS:—
sulcata, *C. Koch & Bouché, in App. Ind. Sem. Hort. Berol.* (1854) 13.—Hab.?
Swaseyi, *Buckley, in Young, Fl. Texas.* 152.—Texas.
tenuifolia, *Lindl. Bot. Reg.* (1838) *Misc.* 64.—Mexic.
Thunbergii, DC. Syst. ii. 9 = vulgaris.
tinctoria, Leschen. in Mem. Mus. Par. ix. (1822) 306 = aristata.
tolucensis, *Hort.—Cf. Gard. Chron.* (1868) 435.—Am. austr.
tolimensis, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 52.—N. Granat.
tomentosa, *Ruiz & Pav. Fl. Per.* iii. 52. t. 282 b.—Chili.
toriusa, Domb. ex DC. Syst. ii. 13 = flexuosa.
tragacanthoides, *DC. Syst.* ii. 18.—Armenia.
trucupidata, Sm. ex DC. l. c. 17 = heterophylla.
trifolia, *Schrad. f. Syst.* vii. 1616.—Mexic.
trifoliata, Hartw. ex Lindl. Bot. Reg. xxvii. (1841) Misc. 68 = sep.
trifoliolata, *Moric. Pl. Nouv. Am.* 113. t. 69.—Texas.
trifurca, *Lindl. in Paxt. Flow. Gard.* iii. 57 (1852–53).—China.
trigona, *Kunze, in Poepp. & Endl. Nov. Gen et Sp.* ii. 68. t. 167.—Chili.
truxillensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 285.—Venezuela.
Tschonoskyana, *Regel, in Act. Hort. Petrop.* ii. (1873) 421.—Japon.
turcomannica, Kar. ex Ledeb. Fl. Ross. i. 79 = vulgaris.
ulcina, *Hook. f. & Thoms. Fl. Ind.* i. 227.—Tibet. occ.
umbellata, Lindl. Bot. Reg. xxx. (1844) t. 44 = aristata.
umbellata, *Phil. in Anal. Univ. Chil.* (1872) 663.—Chili.
umbellata, *Well. Cat. n.* 1475; *et ex G. Don, Gen. Syst.* i. 116.—Reg. Himal.
undulata, Hort. ex C. Koch, Dendrol. i. 408 = asiatica.
undulata, *Lindl. in Journ. Hort. Soc.* v. (1850) 7.—Peruv.
valdiviana, *Phil. in Linnæa*, xxviii. (1856) 609.—Chili.
verticillata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 286.—N. Granat.
violacea, Poit. & Turp. in Duham. ed. Nov. iv. 12; Hort. ex C. Koch, Dendrol. i. 394 = vulgaris.
virgata, Hort. ex Lindl. in Journ. Hort. Soc. v. (1850) 6 = parviflora.
virgata, Hort. ex C. Koch, Dendrol. i. 418 = actinacantha.
virgata, *Ruiz & Pav. Fl. Per.* iii. 51. t. 281. f. B.—Peruv.
vulgaris, *Linn. Sp. Pl.* 330.—Europ.; As. temp.
vulgaris, Vell. Fl. Flum. iii. t. 149 = glaucescens.
Wallichiana, *DC. Prod.* i. 107.—Reg. Himal.
Weddelli, *Lechl. Berb. Am. Austr.* 21.—Bolivia.
xanthoxylon, Hassk. Cat. Hort. Bog. Alt. 180 = Wallichiana?

BERCHEMIA, Neck. Elem. ii. 122 (1790). *RHAMNEÆ*, Benth. & Hook. f. i. 377.
ONOPLEA, Hedw. f. Gen. i. 151 (1806).
OPNOPLIA, Schult. ex Roem. & Schult. Syst. v. 382 (1819).
affinis, *Hassk. Hort. Bog. Descr.* 117.—Malaya.
Burmanniana, *DC. Prod.* ii. 28.—Zeylan.
calophylla, G. Don, Gen. Syst. ii. 28 = Zizyphus calophylla.
congesta, S. Moore, in Journ. Bot. xlii. (1875) 226 = Microrhamnus frangulioides.
corollata, F. Muell. Census, 60 = ecorollata.
discolor, *Hemsl. in Oliver, Fl. Trop. Afr.* i. 581.—Afr. trop.
Edgeworthii, *Lawson, in Hook. f. Fl. Brit. Ind.* i. 636. —Reg. Himal.
ecorollata, *F. Muell. Fragm.* ix. 141.—Austral.
flavescens, *Well. Cat. n.* 4955; *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 357.—Reg. Himal.
floribunda, *Well. Cat. n.* 4256; *Brongn. l. c.*—Reg. Himal.
Fournieri, *Pauch. & Sabert, Not Bois. Nouv. Caléd.* 296.—N. Caled.

BERCHEMIA:—
hamosa, Wall. Cat. n. 4253 = Sageretia hamosa.
laxa, Wall. l. c. n. 4257 = floribunda.
lineata, Benth. in Hook. Kew Journ. iv. (1852) 42 = racemosa.
lineata, *DC. Prod.* ii. 23.—China.
Loureiriana, *DC. l. c.*—China.
oppositifolia, Wall. Cat. n. 4254 = Sageretia oppositifolia.
parviflora, Wall. l. c. n. 4258 ; Wight, Ic. t. 19 = Sageretia oppositifolia.
Poiretiana, DC. Prod. ii. 23 = lineata.
polyphylla, *Wall. Cat.* n. 4259.—Burma.
pubiflora, *Miq. Fl. Ind. Bat.* i. 1. 645.—Malaya.
racemosa, *Sieb. & Zucc in Abh. Akad. Muench.* iv. 11. (1845) 147.—Japon.
repanda, *Rafin. Sylva Tellur.* 33.—Hab. ?
sessiliflora, *Benth. Fl. Hongk.* 68.—China.
trichantha, *Miq. Fl. Ind Bat. Suppl* i. 831.—Malaya.
undulata, *Rafin. Sylva Tellur.* 33.—Hab. ?
volubilis, *DC. Prod.* ii. 23.—Am. bor.

BERCHTOLDIA, Presl, Rel. Haenk. 223. t. 43 (1830) = **Chaetium**, Nees (Gramin.).
bromoides, Presl, l. c. 224 = C. bromoides.
holciformis, Fourn. ex Benth. in Journ. Linn. Soc. xix. (1881) 46 = Opliemenus holciformis.
junciformis, Hemsl. Biol. Centr. Am. Bot. iii. 501, lapsu = Opliemenus holciformis.
opliemenoides, Fourn. ex Benth. in Journ. Linn. Soc. xix. (1881) 46 = Echinochloae sp. ?

BERCKHEYA, Pers. Syn. ii. 480 (1807) = **Berkheya**, Ehrh. (Compos.).

BEREBERA, Baker, in Oliver, Fl. Trop. Afr. ii. 131 (1871), sphalm = BERREBERA, Hochst. = **Millottia**, Wight & Arn. (Leguminos.).

BERENDTIA, A Gray, in Proc. Am. Acad. vii. (1868) 379. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 942.
Coulteri, *A. Gray, l. c.*—880.—Mexic.
Gheisbrechtii, A. Gray, l. c. = rugosa.
rugosa, *A. Gray, l. c.*—Mexic.

BERENICE, Salisb. Gen. Pl. Fragm. 89 (1866) = **Allium**, Linn. (Liliac.).
Victorialis, Salisb. 490 = A. Victorialis.

BERENICE, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 156. *SAXIFRAGEAE*, Benth. & Hook. f. i. 646.
arguta, *Tul. l. c.*—Ins. Borbon.

BERGENA, Adans. Fam. ii. 245 (1763) = **Leoythis**, Linn. (Myrtac.).

BERGENIA, Moench, Meth. 664 (1794) = **Saxifraga**, Tourn.
bifora, Moench, l. c. = S. crassifolia.
cordyfolia, Sternb. Rev. Sax. Suppl. ii. 2 = S. crassifolia.
purpurascens, Engl. in Verh. Zool. Bot. Ges. Wien, xix. (1869) 549 = S. purpurascens.

BERGENIA, Neck. Elem. ii. 108 (1790) = **Lythrum**, Linn.

BERGENIA, Rafin. Sylva Tellur. 102 (1838) = **Cuphea**, P. Br. (Lythrac.).
alphilitica, Rafin. l. c.—Hab. ?

BERGERA, Koen. ex Linn. Mant. ii. 563 (1771) = **Murraya**, Linn. (Rutac.).
compressa, Blanco, Fl. Filip. ed. I. 360.—Ins. Philipp.
inodora, Blanco, l. c. 361.—Ins. Philipp.
integerrima, Buch.-Ham. ex Roxb. Hort. Beng. 32 ; Fl. Ind. ii. 376 = Micromelum pubescens.
intermedia, Zoll. & Moritzi, Syst. Verz. Zoll. 24.—Java.
Koenigii, Linn. Mant. 565 = Murraya Koenigii.
nitida, Thw. Enum. Pl. Zeyl. 46 = Clausena indica.
ternata, Blanco, Fl. Filip. ed. I. 360 = Glycosmis bilocularis.
villosa, Wall. Cat. n. 6373 = Micromelum pubescens.
Wallichii, A. DC. ex Oliver, in Journ. Linn. Soc. v. Suppl. ii. (1861) 35 = Clausena Wallichii.

BERGERETIA, Desv. Journ. Bot. iii. (1813) 161 = **Clypeola**, Linn. (Crucif.).
echinata, Desv. l. c. = C. lasiocarpa.

BERGERIA, Koen. ex Steud. Nom. ed. II. i. 249, in syn. (1840) = **Koenigia**, Linn. (Polygon.).

BERGERONIA, Michell, in Mém. Soc. Phys. Genèv. xxviii. (1883) n. 7 (Contr. Fl. Parag. 38). *LEGUMINOSAE.*
sericea, *Michell, il. cc.* 39.—Paraguay.

BERGHAUSIA, Endl. Gen. Suppl. iii. 57 (1843) = **Garrotia**, Incogn. (Gramin.).
barbulata, Endl. ex Miq. Anal. Bot. Ind. 32.—Malaya.
courtallensis, Nees, ex Miq. l. c. = G. courtallensis.
elata, Arn. ex Miq. l. c.—Malaya.
Emodi, Endl. ex Miq. l. c.—Malaya.
mutica, Munro, in Proc. Am. Acad. iv. (1864) 362.—Hongkong.
patiens, Arn. ex Miq. Anal. Bot. Ind. 32.
patula, Munro, in Proc. Am. Acad. iv. (1864) 362 = G. patula.
tenella, Arn. ex Miq. Anal. Bot. Ind. 34.—Malaya.

BERGHEMIA, Nees, in Linnaea, xx. (1847) 701. *RUBIACEAE* ‡, Benth. & Hook. f. ii. 29.
coccinea, *Nees, l. c.* 702.—Mexic.

BERGHIAS, Juss. in Mém. Mus. Par. vi. (1820) 399 = BERGKIAS, Sonn. = **Gardenia**, Ellis (Rubiac.).

BERGIA, Linn. Mant. ii. 252. n. 1309 (1771). *ELATINEAE*, Benth. & Hook. f. i. 188.
BERGIERA, Neck. Elem. ii. 327 (1790).
LANCRETIA, Del. Fl. Egypte, 218. t. 25. f. 1 (1813).
MERIMEA, Cambess. in St. Hil. Fl. Bras. Mer. ii. 160. t. 107 (1829).
SPINDLOCOCCA, Willd. ex Roem. & Schult. Syst. vi. p. lxx. 799 (1820).
abyssinica, *A. Rich. Tent. Fl. Abyss.* i. 49.—Abyss.
aestivosa, *Steud. Nom.* ed. II. i. 198.—Ind. bor. occ.
ammannioides, *Roxb. Hort. Beng.* 34 ; *Fl. Ind.* ii. 456. —As. Afr. et Austral. trop.
anagalloides, *Walp. Rep.* ii. 286.—Afr. austr.
aquatica, Roxb. Pl. Corom. ii. 22. t. 142 = verticillata.
arenarioides, *Fenzl, in Denkschr. Baier. Bot. Ges.* iii. (1841) 183.—Bras.
capensis, *Linn. Mant.* 241.—Afr. austr.
decumbens, *Planch. ex Harv. Thes. Cap.* i. 15. t. 24. —Afr. austr.
erecta, *Guill. & Perr. Tent. Fl. Senag.* 44.—Afr. trop.
glandulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 371.—Ins. Luson.
glomerata, *Linn. f. Suppl.* 248.—Afr. austr.
odorata, *Edgew. in Journ. As. Soc. Beng.* vii. (1838) 765.—Afr. trop. ; Persia ; Ind. bor. occ.
oryxetorum, *Fenzl, ex Zoll. Syst. Verz. Ind. Archip.* 62. —Malayen.
pedicellaris, *F. Muell. Fragm.* ii. 145.—Austral.
pentandra, Cambess. ex Guill. & Perr. Tent. Fl. Senag. 43. t. 12 = ammanioides.
pepliodes, *Guill. & Perr. l. c.* 44.—Afr. trop.
perennis, *F. Muell. Fragm.* ii. 146.—Austral.
polyantha, *Sond. in Linnaea*, xxiii. (1850) 16.—Afr. austr.
pusilla, *Benth. Fl. Austral.* i. 180.—Austral.
repens, *Blume, Bijdr.* 62.—Java.
serrata, *Blanco, Fl. Filip.* ed. I. 337.—Ins. Philipp.
sessiliflora, *Griseb. Cat. Pl. Cub.* 40.—Cuba.
suffruticosa, Fenzl, in Denkschr. Baier. Bot. Ges. iii. (1841) 183 = odorata.
texana, *Seub. ex Walp. Rep.* i. 285.—Texas.
trimera, Flech. & Mey. in Linnaea, x. (1836) 76 = ammanioides.
tripetala, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 66.—Austral.
verticillata, *Willd. Sp. Pl.* ii. 770 = capensis.

BERGIERA, Neck. Elem. ii. 327 (1790) = prec.

BERGINIA, Harv. ex Benth. & Hook. f. Gen. ii. 1096 (1876). *ACANTHACEAE*, Benth. & Hook. f. i. c.
virgata, *Harv. ex Benth. & Hook. f. l. c.* 1097.—Calif.

BERGKIAS, Sonner. Voy. 48. t. 17 (1776) = **Gardenia**, Ellis (Rubiac.).

BERGSMIA, Blume, Rumphia, iv. 28. t. 176 C. f. 2 (1846). *BIXINEAE*, Benth. & Hook. f. i. 139.
acuminata, *Miq. Fl. Ind. Bat. Suppl.* i. 289.—Sumatra.
javanica, *Blume, Rumphia*, iv. 28.—Java.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* i. 289.—Sumatra.

BERHARDIA, C. Muell. in Walp. Ann. v. 89 (1846), sphalm. = **Berardia**, Vill. (Compos.).

BERIESA, Steud. Nom. ed. II. i. 199 (1840) = SIEBERA, Presl = **Anredera**, Juss. (Chenop.).
baselioides, Steud. l. c. = A. spicata.

BERINGERIA, Neck. Elem. i. 312 (1790) = **Ballota**, Linn. (Labiat.).
acetabulosa, Neck. l. c. = Ballota acetabulosa.
africana, Neck. l. c. = Ballota africana.
cinerea, G. Don, in Sweet, Hort. Brit. ed. III. 545 = Ballota hirsuta.
crispa, G. Don, l. c. = Ballota africana.
frutescens, Reichb. Fl. Germ. Excurs. 325 = Ballota spinosa.
hirsuta, Neck. Elem. 312 = Ballota hirsuta.
hispanica, Neck. l. c. = Ballota hispanica.
mollissima, Heynh. Nom. ii. 86 = Ballota mollissima.
orientalis, G. Don, in Sweet, Hort. Brit. ed. III. 548 = Ballota orientalis (sp. dubia).
pseudodictamnus, Neck. Elem. i. 312 ; Link, Handb. 478 = Ballota pseudodictamnus.
rugosa, Heynh. Nom. ii. 66 = Ballota saxatilis.

BERINIA, Brignol. Pl. Forojul. 50 (1810) = **Crepis**, Vaill. (Compos.).
abyssinica, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 315 = C. abyssinica.
acuminata, Sch. Bip. l. c. = C. acuminata.
alpestris, Sch. Bip. l. c. = C. alpestris.
andryaloides, Brignol. Pl. Forojul. 50 = C. chondrilloides.
aurea, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 318 = C. aurea.
auriculaefolia, Sch. Bip. l. c. 315 = C. auriculaefolia.
biennis, Sch. Bip. l. c. 318 = C. biennis.
chondrilloides, Sieb. ex Sch. Bip. l. c. 315 = C. chondrilloides.
chrysantha, Sch. Bip. l. c. 318 = C. chrysantha.
ciliata, Sch. Bip. l. c. 317 = C. ciliata.
crocea, Sch. Bip. l. c. = C. crocea.
foetida, Sch. Bip. l. c. 315 = C. foetida.
grandiflora, Sch. Bip. l. c. 317 = C. grandiflora.
heterantha, Sch. Bip. l. c. = Lactuca Dubyaea.
heterotricha, Sch. Bip. l. c. = C. heterotricha.
Jacquini, Sch. Bip. l. c. 315 = C. Froelichii ?
jubata, Sch. Bip. l. c. = C. jubata.
lacera, Sch. Bip. l. c. 315 = C. lacera.
nicaeensis, Sch. Bip. l. c. 318 = C. nicaeensis.
occidentalis, Sch. Bip. l. c. 315 = C. occidentalis.
oporinioides, Sch. Bip. l. c. = C. oporinioides.
pannonica, Sch. Bip. l. c. = C. rigida.
pinnatifida, Sch. Bip. l. c. = C. pinnatifida.
polytricha, Sch. Bip. l. c. 317 = C. polytricha.
runcinata, Sch. Bip. l. c. 315 = C. runcinata.
Sibthorpiana, Sch. Bip. l. c. = C. Sibthorpiana.
stricta, Sch. Bip. l. c. 317 = C. virens ? vel tectorum.
tenuifolia, Sch. Bip. l. c. 316 = C. tenuifolia.

BERKHEYA, Ehrh. Beitr. iii. 187 (1788). *COMPOSITAE*, Benth. & Hook. f. ii. 460.
AGRIPHYLLUM, Juss. Gen. 190 (1789).
APULEIA, Gaertn. Fruct. ii. 439. t. 171 (1791).
ARELINA, Neck. Elem. i. 88 (1790).
ASPIDALIS, Gaertn. Fruct. ii. t. 171 (1791).
BASTERIA, Houtt. Nat. Hist. Pl. iv. 158 (1776).
BASTERA, J. F. Gmel. Syst. 1871 (1791).
BERCKEYA, Schreb. Gen. ii. 877 (1791).
CROCODILOIDES, Adans. Fam. ii. 127 (1763).
CUPIDIA, Gaertn. Fruct. ii. 454. t. 171 (1791).
ENOPIA, Cass. in Bull. Soc. Philom. (Févr. 1818) 25 ; et in Dict. Sc. Nat. xvi. 25 (1820).
GORTERIA, Linn. Sp. Pl. ed. II. 1262 (1762), pro parte ; Linn. Ilustr. t. 702 (1789 ?).
HETEROACHYS, Sch. Bip. in Flora, xxvii. (1844) 775.
ROHRIA, Vahl, in Skr. Naturh.-Selsk. Kjoeb. iii. (1793) 97.
STOBAEA, Thunb. Prod. Pl. Cap. 161 (1800).
ZARABELLIA, Neck. Elem. i. 16 (1790).

BERKHEYA :—
alhida, Lichtst. ex DC. Prod. vii. 86 = Dicoma capensis.
anaxetens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 309.—Afr. austr.
bisulca, Pers. Syn. ii. 486 = Cullumia bisulca.
canescens, *DC. Prod.* vi. 507.—Afr. austr.
carduiformis, *DC. l. c.*—Afr. austr.
cerinoides, *Willd. Sp. Pl.* iii. 2275.—Afr. austr.
carthamoides, *Willd. l. c.* 2274.—Afr. austr.
cernua, R. Br. in Ait. Hort. Kew. ed. II. v. 139 = Didelta cernua.
ciliaris, Willd. Sp. Pl. iii. 2273 = Cullumia ciliaris.
coriacea, *Harv. in Harv. & Sond. Fl. Cap.* iii. 506.—Afr. austr.
corymbosa, *DC. Prod.* vi. 507.—Afr. austr.
crucifera, *Willd. Sp. Pl.* 2276.—Afr. austr.
cuneata, *Willd. l. c.* 2270.—Afr. austr.
cynaroides, *Willd. l. c.* 2275.—Afr. austr.
decurrens, Willd. l. c. 2276 = Stephanocoma berkheyoides.
Dregei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 507.—Afr. austr.
Ecklonis, *Harv. l. c.* 504.—Afr. austr.
eryngiifolia, *Less. Syn. Comp.* 703.—Afr. austr.
fruticosa, *Ehrh. Beitr.* iii. 138.—Afr. austr.
gazanioides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 508.—Afr. austr.
gorterioides, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 429.—Afr. trop.
grandiflora, *Willd. Sp. Pl.* iii. 2271.—Afr. austr.
hirpicioides, *Sond. in Harv. & Sond. Fl. Cap.* iii. 509.—Afr. austr.
hispida, Willd. Sp. Pl. iii. 2274 = Cullumia hispida.
incana, Willd. l. c. 2269 = fruticosa.
Kraussii, Sch. Bip. in Flora, xxvii. (1844) 774 = cynaroides.
lanceolata, *Willd. Sp. Pl.* iii. 2270.—Afr. austr.
obovata, *Willd. l. c.* 2269.—Afr. austr.
palmata, *Willd. l. c.* 2271.—Afr. austr.
patula, Willd. l. c. 2272 = Cullumia patula.
pectinata, Willd. l. c. = Cullumia pectinata.
pinnata, *Less. Syn. Comp.* 78.—Afr. austr.
pungens, *Willd Sp. Pl.* iii. 2275.—Afr. austr.
purpurea, *Benth. & Hook. f. ex Mast. in Gard. Chron.* (1872) 1392.—Afr. austr.
scolymoides, *DC. Prod.* vi. 504.—Afr. austr.
serrulata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 507.—Afr. austr.
setifera, *DC. Prod.* vi. 507.—Afr. austr.
setosa, Willd. Sp. Pl. iii. 2273 = Cullumia setosa.
Spekeana, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 100.—Afr. trop.
spinosissima, *Willd. Sp. Pl.* iii. 2270.—Afr. austr.
squarrosa, Willd. l. c. 2272 = Cullumia bisulca.
stobaeoides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 505.—Afr. austr.
subulata, *Harv. l. c.* 507.—Afr. austr.
sulcata, Willd. Sp. Pl. 2271 = Cullumia sulcata.
umbellata, *DC. Prod.* vi. 507.—Afr. austr.
uniflora, *Willd. Sp. Pl.* iii. 2274.—Afr. austr.
Zeyheri, *Harv. & Sond. Fl. Cap.* iii. 496.—Afr. austr.

BERLANDIERA, DC. Prod. v. 517 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 350.
incisa, Torr. & Gray, Fl. N. Am. ii. 282 = lyrata.
longifolia, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 349 = texana.
lyrata, *Benth. Pl. Hartw.* 18.—Am. bor. occ.
pumila, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 343 = tomentosa.
subacaulis, *Nutt. l. c.* 343.—Florida.
texana, *DC. in Deless. Ic. Sel.* iv. t. 96.—Am. bor.
tomentosa, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 343.—Am. bor.

BERLIERA, Buch.-Ham. ex Wall. Cat. sub n. 6235 (1832) = **Myrtomarvan**, R. Br. (Rubiac.).
mutans, Buch.-Ham. l. c. = M. mutans.

BERLINIA, Soland. ex Hook. f. & Benth. in Hook. Niger Fl. 296 (1849). *LEGUMINOSAE*, Benth. & Hook. f. i. 579.
scandens, *Soland. ex Hook. f. & Benth. l. c.*—Afr. trop.
angolensis, *Welw. ex Benth. in Trans. Linn. Soc.* xxv. (1866) 310.—Afr. trop.
auriculata, *Benth. l. c.* 309.—Afr. trop.
bracteosa, *Benth. in Trans. Linn. Soc.* xxv. (1866) 309.—Afr. trop.
Heudelotiana, *Baill. Adansonia*, vi. (1865) 185.—Afr. trop.
paniculata, *Benth. in Trans. Linn. Soc.* xxv. (1866) 311.—Afr. trop.
stipulacea, *Benth. l. c.* 310.—Afr. trop.

BERMUDIANA, [Tourn.] Linn. Syst. ed. I (1735) = **Sisyrinchium**, Linn. (Irid.).
graminea, Gaertn. Fruct. i. 32 = S. Bermudiana.
graminifolia, Medic. Malv. 77 = S. anceps
guttata, Stokes, Bot. Mat. Med i. 109 = Moraea chinensis.
iridifolia, Medic. Malv. 77 = S. Bermudiana.
striata, Moench, Meth. Suppl 118 = S. striatum.

BERNARDIA, Endl. Gen. 469, in syn. (1838) = **Berardia**, Vill. (Compos.).

BERNARDIA, Houst. ex P. Br. Nat. Hist. Jamaic. 361 (1756). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 308.
ALEVIA, Baill. Etud. Gén. Euph. 508 (1858).
BIVONIA, Spreng. Neue Entdeck. ii. 116 (1821).
PASSARA, Baill. Etud. Gén. Euph. 507. t. 18 (1858).
PHAEDRA, Klotzsch, in Endl. Gen. Suppl iv. III. 88 (1847).
POLYBASA, Klotzsch, l. c. (1847).
TRAGANTHUS, Klotzsch, in Wiegm. Archiv, vii. (1841) 188. t. 9.
TYRIA, Klotzsch, in Endl. Gen. Suppl iv. III. 88 (1847).
axillaris, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 174.—Bras.
brevipes, *Muell. Arg. l. c.* 176.—Bras.
caperonioefolia, *Muell. Arg. in DC. Prod.* xv. II. 920.—Reg. Argent.
capetiifolia, Muell. Arg. in Linnaea, xxxiv. (1865–66) 176 = celastrinea.
carpinifolia, *Griseb. Fl. Brit. W. Ind.* 46.—Ind. occ.
celastrinea, *Muell. Arg. in DC. Prod.* xv. II. 921.—Bras.
confertifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 175.—Bras.
corensis, *Muell. Arg. l. c.* 173.—Ind. occ.
crassifolia, *Muell. Arg. in Mart. Fl. Bras.* xi II. 394.—Bras.
dichotoma, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 172.—Ind. occ.
fasciculata, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 158.—Am. bor. occ.
gambas, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 391.—Bras.
Gardneri, *Muell. Arg. l. c* 394.—Bras.
grandifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 178.—Guiana.
hirsutissima, *Muell. Arg. in DC. Prod.* xv. II. 922.—Bras.
intermedia, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 160 = mexicana.
interrupta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 171.—Mexic.
Jacquiniana, *Muell. Arg. l. c.* 173.—Venezuela.
leprosa, *Muell. Arg. l. c.* 172.—Ins. S. Doming.
Lorentzii, *Muell. Arg. in Journ. Bot. xii.* (1874) 229.—Reg. Argent.
megalophylla, Muell. Arg. in Linnaea, xxxiv. (1865–66) 178 = Adenophaedra megalophylla.
mexicana, *Muell. Arg. l. c.* 173.—Mexic.
microphylla, *Muell. Arg. l. c.*—Cuba.
multicaulis, *Muell. Arg. l. c.* 177.—Bras.
myricoides, *Benth. & Hook. f. Gen.* iii. 308.—Am. bor.
oligandra, Muell. Arg. in Linnaea, xxxiv. (1865–66) 175 = tamandoana.
peduncularis, *Muell. Arg. l. c.* 176.—Bras.
polichella, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 392.—Bras.
scabra, *Muell. Arg. l. c.* 396.—Bras.
Sellowii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 177.—Bras.
sidoides, *Muell. Arg. l. c.*—Guiana.
spartioides, *Muell. Arg. l. c.*—Bras.
tamandoana, *Muell. Arg. in DC. Prod.* xv. II. 920.—Bras.
venosa, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 159.—Ind. occ.

BERNARDINIA, Planch. in Linnaea, xxiii. (1850) 412. *CONNARACEAE*, Benth. & Hook. f. i. 431.
fluminensis, *Planch. l. c.*—Bras.

BERNBUXIA, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 159. *DIAPENSIACEAE*, Benth. & Hook. f. ii. 691.
thibetica, *Decne. l. c.*—Tibet.

BERNIERA, DC. Prod. vii. 18 (1838) = **Gorbera**, Gronov. (Compos.).
nepalensis, DC. l. c. = G. macrophylla.

BERNIERIA, Baill. in Bull. Soc. Linn. Par. i. (1884) 434. *LAURINEAE*.
madagascariensis, *Baill. l. c.*—Madag.

BERNOULLIA, Oliver, in Hook. Ic. Pl. t. 1169 (1873). *STERCULIACEAE*.
flammea, *Oliver, l. c.* t. 1169–70.—Guatemala.

BERREBERA, Hochst. in Flora, xxvii. (1844) 100; xxix. (1846) 597 = **Millettia**, Wight & Arn. (Legum.).
caffra, Hochst. ll. cc. 100 et 598 = M. caffra.
ferruginea, Hochst. ll cc. 100 et 597 = M. ferruginea.

BERTERIA, Rozb. Hort. Beng. 42 (1814); Pl. Coromand. iii. 60. t. 264 (1819). *TILIACEAE*, Benth. & Hook. f. i. 232.
ESPERA, Willd. in Ges. Naturf. Fr. Berl. Neue Schr. iii. (1801) 449, nomen prius.
HEXAGONOTHECA, Turcz. in Bull. Soc. Nat. Mosc. (1846) ii. 505.
Ammonilla, *Rosb Hort. Beng.* 42; *Fl. Ind.* ii. 639.—Zeylan.; Ind or.
mollis, Wall. Cat. n. 1186 = B. Ammonilla.
Vescoana, *Baill. Adansonia*, x. (1872) 240.—Ins. Tahiti.

BERRYA, DC. Prod. i. 517 (1824) = prael.

BERRYA, Klein, ex Willd. Sp. Pl. ii. 840, in syn. (1800) = **Litsea**, Lam (Laurin.)
chinensis, Klein, ex Willd l. c. = L. sebifera.

BERSAMA, Fresen. in Mus. Senckenb. ii. (1837) 280. t. 17. *SAPINDACEAE*, Benth. & Hook. f. i. 412.
NATALIA, Hochst. in Flora, xxiv. (1841) 668.
RHAGANUS, E. Mey. Zwei Pfl. Docum. 216 (1843).
abyssinica, *Fresen. l. c.* 281.—Abyss.
integrifolia, *A. Rich. Tent. Fl. Abyss.* i. 107.—Abyss.
maxima, *Baker, in Oliver, Fl. Trop. Afr.* i. 434.—Afr. trop.
paullinioides, *Baker, l. c.* 435.—Afr. trop.
serrata, *A. Rich. Tent. Fl. Abyss.* i. 107.—Abyss.

BERTERA, Steud. Nom. ed. II. i. 199 (1840) = **Gladiolus**, Tourn. (Irid.).

BERTERoA, DC. in Mém. Mus. Par. vii. (1821) 282; et Syst. ii. 390 (1821) = **Alyssum**, Tourn. (Crucif.).
ascendens, Koch, in Linnaea, xv. (1841) 257 = A. mutabile.
graeca, Boiss. & Heldr. Diagn. Ser. II. i. 35 = A. mutabile.
incana, DC. Syst. ii. 291 = A. incanum.
mutabilis, DC. l. c. 292 = A. mutabile.
obliqua, DC. l. c. = A. mutabile.
orbiculata, DC. l. c. 293 = A. orbiculatum.
peruviana, DC. l. c.—Peru.
Potanini, Maxim. in Bull. Acad. Pétersb. xxvi. (1880) 422.—Mongolia.
procumbens, Portenschl. Enum. Pl. Dalm. 15 = A. mutabile.
spathulata, C. A. Mey. in Ledeb. Fl. Alt. iii. 48 = A. spathulatum.
stricta, Boiss. Diagn. Ser. II. i. 35 = A. orbiculatum.
thessala, Heldr. ex Nym. Consp. 51 = A. orbiculatum.
viridis, Tausch, in Syll. Ratisb. ii. (1828) 245 = A. incanum.

BERTHELOTIA, DC. Prod. v. 375 (1836) = **Fluchea**, Cass. (Compos.).
lanceolata, DC. l. c. 376 = P. lanceolata.

BERTHIERA, Vmt. Tabl. ii. 572 (1799) = **Bertiera**, Aubl. (Rubiac.).

BERTHOLETIA, Reichb. Nom. 91 (1841) = BERTHELOTIA, DC. = **Fluchea**, Cass. (Compos.).

BERTHOLETIA, Bronga. Enum. Gen. 123 (1843) = **Bertholletia**, Humb. & Bonpl. (Myrtac).

BERTHOLLETIA, Humb. & Bonpl. Pl. Aequin. i. 192. t. 56 (1808). *MYRTACEAE*, Benth. & Hook. f. i. 732.
TONCA, Rich. Anal. Fruit. 84 (1811).
excelsa, Berg, in Mart. Fl. Bras. xiv. i. 478. t. 60, 61 = nobilis.
excelsa, *Humb. & Bonpl. Pl. Aequin.* i. 192. t. 56.—Bras.
nobilis, *Miers, in Trans. Linn. Soc.* xxx. (1874) 197.—Guiana.

BERTHRA, Aubl. Pl. Gui. i. 180. t. 69 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 77.
POMATIUM, Gaertn. f. Fruct. iii. 252. t. 225 (1805).
ZALUZANIA, Comm. ex Gaertn. f. Fruct. iii. 74. t. 193 (1805).
aethiopica, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 88.—Afr. trop.
africana, *Rich. in Mém. Soc. Hist. Nat. Par.* v. (1830) 356.—Afr. trop.
angustifolia, *Benth. Bot. Voy. Sulph.* 108.—Ins. Cocos.
blattifolata, Boj. Hort. Maurit. 166 = B. Zaluzania.
borbonica, Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 354 = B. Zaluzania.
bracteolata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 84.—Afr. trop.
brevifiora, *Hiern, l. c.* 85.—Afr. trop.
chrysantha, *Hassk. Retzia*, i. 80.—Java.
coccinea, G. Don, Gen. Syst. iii. 506 = Mussaenda elegans.
ferruginea, Willd. ex Roem. & Schult. Syst. v. 227 = Psychotria rufescens.
gonzaleoides, *Griseb. Cat. Pl. Cub.* 124.—Cuba.
guianensis, *Aubl. Pl. Gui.* i. 180. t. 69.—Guiana.
lateriflora, *Blume, Bijdr.* 987.—Java.
laxa, *Benth. in Hook. Niger Fl.* 394.—Afr. trop.
macrocalyx, *Hassk. in Nederl. Kruidk. Arch.* iv. (1859) 7.—Malaya.
macrocarpa, *Benth. in Hook. Niger Fl.* 394.—Afr. trop.
montana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 83.—Afr. trop.
mucronata, *Comm. ex Gaertn. f. Fruct.* iii. 74. t. 192.—Hab.!
palustris, *Rich. in Mém. Soc. Hist. Nat. Par.* v. (1830) 354.—Guiana.
Pomatium, *Benth. in Hook. Niger Fl.* 394.—Afr. trop.
rufa, Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 354 = B. Zaluzania.
subsessilis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 83.—Afr. trop.
Zaluzania, *Comm. ex Gaertn. f. Fruct.* iii. 74. t. 192.—Ins. Mascar.

BERTIERA, Blume, Bijdr. 987 (1826) = **Adenosacme**, Wall. (Rubiac.).
fasciculata, Blume, l. c. = A. longifolia.
javanica, Blume, l. c. = A. longifolia.
lateriflora, Blume, l. c.—Java.

BERTOLONIA, DC. in Ann. Mus. Par. xix. (1812) t. 5 (14) = **Lavoheria**, Lag. (Compos.).
purpurea, DC. l. c. = L. purpurea.

BERTOLONIA, Raddi, in Mem. Mod. xviii. Fis. 384 (1820). *MELASTOMACEAE*, Benth. & Hook. f. i. 765.
TRIBLEMMA, R. Br. ex DC. Prod. iii. 119 (1828).
acaulis, Cham. in Linnaea, ix. (1834) 388 = Eriocnema Hilariana.
acuminata, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 544.—Bras.
aenea, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 318 = marmorata.
guttata, Hook. f. Bot. Mag. t. 5894 = Gravesia guttata.
hirsuta, Benth. Bot. Voy. Sulph. 94 = Triolaena hirsuta.
Houtteana, *Van Houtte, in Fl. des Serres*, Sér. II. x. (1874) 109.—Hab.!
leucsana, *DC. Prod.* iii. 119.—Bras.
longifolia, *Cham. in Linnaea*, ix. (1834) 384 = Salpinga longifolia.
BERTOLONIA :—
maculata, *DC. Prod.* iii. 114.—Bras.
margaritacea, Hort. Bull. ex Naud. in Fl. des Serres, xvi. (1865–67) 168 = Salpinga margaritacea.
marmorata, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 318.—Bras.
Mirandaei x, *Van Houtte, in Fl. des Serres*, xxi. (1875) 135.
nymphaeaefolia, *Raddi, in Mem. Mod. xviii. Fis.* (1820) 384.—Bras.
ovata, *DC. Prod.* iii. 118.—Bras.
primulaeflora, Hort. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 80 = Monolena primulaeflora.
scabra, Rafin. ex j. C. Schau. in DC. Prod. xi. 584 = Lippia lanceolata.
scorpioides, Baill. Hist. Pl. vii. 11 = Triolena scorpioides.
superbissima, *Hort.—Cf. Gard. Chron.* (1873) 5.—Hab.?

BERTOLONIA, Rafin. in Am. Monthly Mag. (1818) 267 = Lippia, Linn. (Verben.).

BERTOLONIA, Moç. & Sessé, ex DC. Prod. iii. 589 (1828) = Cercocarpus, H. B. & K. (Resac.).
guierioides, Moç. & Sessé, l. c. = C. fothergilioides.

BERTOLONIA, Spinola, Cat. Jard. St. Sebast. (1809) 24 = **Myoporum**, Banks.
glandulosa, Spinola, l. c. = M. viscosum.

BERTOLONIA, Spreng. Neue Entdeck. ii. 110. t. 1. f. 1 (1821) = **Chrysochlamys**, Poepp. (Guttif.).
paniculata, Spreng. l. c.—Bras.

BERTUCHIA, Dennst. Schluess. Hort. Malab. 30 (1818) = **Gardenia**, Ellis (Rubac.).
speciosa, Dennst. l. c.—Ind. or.

BERTYA, Planch. in Hook. Lond. Journ. iv. (1845) 472. t. 16. *EUPHORBIACEAE*, Benth. & Hook. f. ii. 394.
australiana, *Planch. l. c.* 473.—Austral.
dimerostigma, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 98.—Austral.
Findlayi, *F. Muell. Fragm.* viii. 141.—Austral.
gummifera, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 473. t. 16.—Austral.
Mitchelli, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 63.—Austral.
oblongifolia, Muell. Arg. in Flora, xlvii. (1864) 471 = pomaderroides.
oleaefolia, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 478.—Austral.
oppositifolia, *F. Muell. & O'Shanesy, in Wing, South Sc. Record*, ii. (1882) 98.—Austral.
pedellata, *F. Muell. Fragm.* iv. 148.—Austral.
pinifolia, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 478.—Austral.
polymorpha, *Baill. Adansonia*, vi. (1865–66) 298.—N. Zel.
pomaderroides, *F. Muell. Fragm.* iv. 84.—Austral.
spiculata, Muell. Arg. in Flora, xlvii. (1864) 471 = Ricinocarpus pelloculatus.
quadrisepala, *F. Muell. Fragm.* x. 52.—Austral.
rosmarinifolia, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 473.—Austral.
rotundifolia, *F. Muell. Fragm.* iv. 84.—Austral.
tasmanica, Muell. Arg. in Linnaea, xxxiv. (1865–66) 68 = rosmarinifolia.

BERULA, Hoffm. ex Bess. Enum. Pl. Volh. 44 (1822) = Sium, Tourn. (Umbellif.).
angustifolia, Mert. & Koch, i Rochl. Deutschl. Fl. ii. 488 = S. erectum.
lancifolia, Bess. Enum. Pl. Volh. 44 = S. lancifolium.

BERYLLIS, Salisb. Gen. Pl. Fragm. 33 (1866) = Ornithogalum, Tourn. (Liliac.).
pyrenaica, Salisb. l. c. = O. pyrenaicum.
stachyoides, Salisb. l. c. = O. narbonense?

BESLERIA, Bronga. in Ann. Sc. Nat. Sér. I. viii (1826) 370. t. 85. *BRUNIACEAE*, Benth. & Hook. f. i. 671.
HETESODON, Meisn. Gen. 72 (1837).
RAEHNIOSTIA, Reichb. Nom. 159 (1841).
abrotanoides, *Bronga. l. c.* 371.—Afr. austr.
BESLERIA :—
alopecuroides, *Sond. in Harv. & Sond. Fl. Cap.* ii. 310.—Afr. austr.
arachnoides, *Eckl. & Zeyh. Enum.* 138.—Afr. austr.
brevifolia, Eckl. & Zeyh. l. c. = abrotanoides.
callunoides, *Oliver, in Journ. Linn. Soc.* ix. (1867) 338.—Afr. austr.
commutata, *Sond. in Harv. & Sond. Fl. Cap.* ii. 310.—Afr. austr.
comosa, Eckl. & Zeyh. Enum. 137 = commutata.
cordifolia, *Schlecht. in Linnaea*, vi. (1831) 189.—Afr. austr.
ericoides, Eckl. & Zeyh. Enum. 137 = intermedia.
formosa, Eckl. & Zeyh. l. c. 138 = abrotanoides.
globosa, G. Don, Gen. Syst. iii. 46 (= *Berardia globosa*, Sond.).—Afr. austr.
intermedia, *Schlecht. in Linnaea*, vi. (1831) 188.—Afr. austr.
lanuginosa, *Bronga. in Ann. Sc. Nat.* Sér. I. viii. (1826) 372. t. 85.—Afr. austr.
rubra, Schlecht. in Linnaea, vi. (1831) 189 = squarrosa.
squarrosa, *Sond. in Harv. & Sond. Fl. Cap.* ii. 312.—Afr. austr.
superba, *Eckl. & Zeyh. Enum.* 138.—Afr. austr.
Wendlandiana, Eckl. & Zeyh. l. c. 137 = intermedia.

BERZELIA, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 292 = **Hermbstaedtia**, Reichb. (Amarant.).
glauca, Mart. l. c. = H. glauca.

BESCHORNERIA, Kunth, Enum. Pl. v. 844 (1850). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 738.
bracteata, *Yacobi, Index*, 12; *Baker, in Bot. Mag.* t. 6641.—Mexic.
Decosteriana, Baker, in Bot. Mag. t. 6768 = sec.
Dekosteriana, *C. Koch, Wochenschr.* vii. (1864) 187.—Mexic.
dubia, *Carr. in Rev. Hortic.* (1877) 155.—Texas.
floribunda, *Hort. ex C. Koch, Wochenschr.* v. (1862) 199.—Hab.!
Galleottii, *Yacobi, Index*, 12, nomen.—Mexic.?
multiflora, *Hort. ex C. Koch, Wochenschr.* iii. (1860) 63.—Hab.!
Parmentieri, *Yacobi, in Hamb. Gartens.* xx. (1864) 508.—Hab.!
pumila, *Yacobi, Index*, 12, nomen.—Hab.!
Tonelliana, Jacobi, l. c. = sec.
Tonelii, *Yacobi, in Hamb. Gartens.* xx. (1864) 508. nomen.—Mexic.
Schlechtendalii, *Yacobi, Index*, 12, nomen.—Hab.?
tubiflora, *Kunth, Enum. Pl.* v. 844.—Mexic.
Verlindeiana, *Yacobi, Index*, 12, nomen.—Hab.!
yuccoides, Hort. ex Hook. Bot. Mag. t. 5903 = sec.
yuccoides, *C. Koch, Wochenschr.* ii. (1859) 387; iii. (186) 63; vi. (1864) 188.—Mexic.

BESENNA, A. Rich. Tent. Fl. Abyss. i. 253 (1847) = **Albinia**, Durr. (Legumin.).
anthelmintica, A. Rich. l. c. = A. anthelmintica.

BESHA, D. Dietr. Syn. i. 189 (1839) = **Oehlandra**, Thw. (Gramin.).

BESLERIA, Plum. ex Linn. Gen. ed. I. 161 (1737). *GESNERACEAE*, Benth & Hook. f. ii. 1015.
CRANTZIA, Neck. Elem. i. 266 (1790).
CREMOSPERMA, Benth. Pl. Hartw. 284 (1846).
CYRTANTHEMUM, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 130.
DRUPINA, Linn. Pl. Surin. 11 (1775); Amoen. Acad. viii. 259 (1785).
ERIPHIA, P. Br. Hist. Jamaic. 270 (1756).
FLIMNOBOLINA, Rafin. Sylva Tellur. 71 (1838).
GASTERANTHOPSIS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 129.
GASTERANTHUS, Benth. Pl. Hartw. 283 (1846).
HEMATOPHYLA, Rafin. Sylva Tellur. 71 (1838).
PARABESLERIA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 126.
PSEUDOBESLERIA, Oerst. l. c. 128 (1861).
acutifolia, *Benth. Pl. Hartw.* 287.—N. Granad.
aggregata, *Hanst. in Mart. Fl. Bras.* viii. 400.—Bras.
ampla, *Vell. Fl. Flum.* 262; vi. t. 64.—Bras.
arborea, *Vell. fl. cv.* 261; vi. t. 92.—Bras.
ardens, *Decne. ex Lindex, Cat.* n. 5 (1880) 1.—Venezuela.

BESLERIA :—
barbata, *Hanst. in Linnaea*, xxxiv. (1865–66) 327.—Peruv.
barbensis, *Hanst. l. c.* 319.—Costa Rica.
Berteriana, *DC. Prod.* vii. 538.—Jamaica.
bicolor, *H. B. & K. Nov. Gen. et Sp.* ii. 398.—N. Granat.
bicolor, Schott, in Oestr. Med. Jahrb. vi. ii. 65 = Allopilectus bicolor.
bifora, Forst. f. Fl. Ins. Austr. 43. n. 236 = Cyrtandra biflora.
bivalvis, Linn. f. Suppl. 280 (= *Picria surinamensis*). —Guiana.
bonodora, Vell. Fl. Flum. 261; vi. t. 80 = Brunfelsia latifolia.
carrulea, Aubl. Pl. Gui. ii. 631 = violacea.
calcarata, *H. B. & K. Nov. Gen. et Sp.* ii. 399.—N. Granat.
campanulata, *Linden, Cat.* n. 5. (1850) 2.—Venezuela.
capitata, *Poepp. & Endl. Nov. Gen. et Sp* iii 2.— Peruv.
coccinea, Aubl. Pl. Gui. ii. 632 = Allopilectus coccineus.
columneoides, *Hanst. in Linnaea*, xxxiv. (1865–66) 325.—Costa Rica.
costaricensis, *Hanst. l. c.* 330.—Costa Rica.
crassifolia, *Schott, Neu. Bras. Pfl.* i. 7.—Bras.
cristata, Linn. Sp. Pl. 619 = Allopilectus cristatus.
cristata, Moç. & Sesse, ex DC. Prod. vii. 545, 540 = Allopilectus glaber.
cuneata, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 548.—Bras.
cymosa, Forst. f. Fl. Ins. Austr. 44. n. 237 = Cyrtandra cymosa.
Cyrtanthemum, *Hanst. in Linnaea*, xxxiv. (1865–66) 321.—Mexic.
deflexa, *Hanst. l. c.*—Mexic.
dichrus, Spreng. Syst. ii. 840 = Allopilectus dichrous.
divaricata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 2.— Peruv.
Drymonia, Steud. Nom. ed. II. i. 200 = Drymonia calcarata.
elegans, *H. B. & K. Nov. Gen. et Sp.* ii. 397.—N. Granat.
fasciculata, *Wawra, in Oestr. Bot. Zeitschr.* xiii. (1863) 142.—Bras.
flavoviresa, *Mart. in Nov. Act. Nat. Cur.* xi. (1823) 49.—Bras.
geminiflora, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 548.—Bras.
gibbosa, *Hanst. in Mart. Fl. Bras.* viii. 421.—Bras.
glabra, *Hanst. in Linnaea*, xxxiv. (1865–66) 325.— Mexic.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* ii. 401.—N. Granat.
grandifolia, *Schott, Neu. Bras. Pfl.* i. 5. n. 5.—Bras.
guadalupensis, *DC. Prod.* vii. 535.—Guadalup.
hirsuta, *Hanst. in Linnaea*, xxxiv. (1865–66) 326.— Mexic.
hirtella, Schott, in Oestr. Med. Jahrb. vi. ii. 66 = Allopilectus sparsiflorus.
hispida, H. B. & K. Nov. Gen. et Sp. ii. 39 = Allopilectus hispidus.
Imray, *Hook. Bot. Mag.* t. 6841.—Ind. occ.
incarnata, *Aubl. Pl. Gui.* ii. 635.—Guiana.
inodora, Vell. Fl. Flum. 261; vi. t. 81 = Brunfelsia pauciflora.
insignis, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) ii. 87 = Solenophora insignis.
labiosa, *Hanst. in Linnaea*, xxxiv. (1865–66) 324.—N. Granat.
laxiflora, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 361.—Guiana.
leucostoma, *Hanst. in Linnaea*, xxxiv. (1865–66) 326. —N. Granat.
lucida, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 2.—Peruv.
lutea, *Linn. Sp. Pl.* 619.—Ind. occ.
lutea, Spreng. Syst. ii. 841 = Berteriana.
lutea-virens, Mart. Nov. Gen. iii. 45 = flavoviresa.
melittifolia, Linn. Sp. Pl. 619 = Episcia melittifolia.
mellis, *Hook. ex Steud. Nom.* ed. II. i. 200.—Am. austr.
mollissima, Spreng. ex Steud. Nom. ed. I. 107 = preec. ?
BESLERIA :—
pentaphylla, *Vell. Fl. Flum.* 262; vi. t. 88.—Bras.
Princeps, *Hanst. in Linnaea*, xxxiv. (1865–66) 317.— Costa Rica.
pulchella, Donn, Hort. Cantab. 140, ex Sims, Bot. Mag. t. 1146 = Tussacia pulchella.
ruiteusis, *Hanst. in Linnaea*, xxxiv. (1865–66) 324.— Ecuador.
Rhytidophyllum, *Hanst. l. c.* 324.—N. Granat.
Riedeliana, *Hanst. in Mart. Fl. Bras.* viii. 399.— Bras.
sanguinea, Pers. Syn. ii. 165 = Allopilectus sanguineus.
scandens, Moç. & Sesse, ex D. Don, in Trans. Linn. Soc. xv. (1827) 349 = Maurandia erubescens?
Selloana, *Klotzsch & Hanst. in Mart. Fl. Bras.* viii. 398.—Bras.
serrulata, *Vacq. Hort. Schœrnb.* iii. 21.—Ind. occ.
solanoides, *H. B. & K. Nov. Gen. et Sp.* ii. 398.—Am. austr.
spectabilis, H. B. & K. l. c. 400 = Drymonia spectabilis.
splendens, Hort. ex Steud. Nom. ed. II. i. 200 = Drymonia spectabilis.
surinamensis, *Miq. in Linnaea*, xxii. (1849) 471.— Guiana.
Symphytum, *Klotzsch & Hanst. ex Hanst. in Mart. Fl. Bras.* viii 399.—Bras.
tetragonifolia, *Ruiz, ex Hanst. in Linnaea*, xxxiv. (1865–66) 331.—Peruv.
trifora, *Hanst. l. c.* 329.—Costa Rica.
umbellata, Herb. Banks, ex Sims, Bot. Mag. t. 1146 = Episcia pulchella.
umbrosa, *Mart. Nov. Gen.* iii. 44. t. 218.—Bras.
violacea, *Aubl. Pl. Gui.* ii. 630. t. 254.—Guiana.
Wendlandiana, *Hanst. in Linnaea*, xxxiv. (1865–66) 318.—Costa Rica.

BESSERA, Schult. f. in Linnaea, iv. (1829) 121. *LILIACEÆ*, Benth. & Hook. f. iii. 801.
PHARIUM, Herb. in Bot. Reg. t. 1546 (1832).
Bessera, *Schult. f. l. c.*—Mexic.
fistulosa, Pritz. Ic. Ind. 147 = elegans.
Herberti, G Don, in Sweet, Hort. Brit. ed. III. 694 = elegans.
minuta, Lem. in Fl. des Serres, Sér. I. iv. (1848) t. 424 = elegans.
multiflora, Mart. & Gal. in Bull. Acad. Bras. ix. (1842) (suppl. 14) = elegans.

BESSERA, Schult. Obs. Bot. 27 (1809) = **Pulmonaria**, Tourn. (Borag.).
azurea, Schult. l. c. = P. azurea.

BESSERA, Spreng. Pugill. ii. 90 (1815) = **Xyloma**, Forst. (Bixin.).
inermis, Spreng. l. c. = Fleggia microcarpa.
spinosa, Spreng. l. c. 91 = X. nitidum.

BESSERA, Vell. Fl. Flum. 147 (1825); iv. t. 2 = **Pisonia**, Linn. (Nyctagin.).
calycantha, Vell. Il. cc. = P. Olfersiana.

BESSIA, Rafin. Sylva Tellur. 18 (1838) = **Corypha**, Linn. (Palm.).
sanguinolenta, Rafin. l. c. = C. umbraculifera.

BESTRAM, Adans. Fam. ii. 354 (1763) = **Antidesma**, Linn. (Euphork.).

BETA, [Tourn.] Linn. Syst. ed. I (1735). *CHENOPODIACEÆ*, Benth. & Hook. f. i. 52.
alba, DC. Fl. Fr. iii. 388 = vulgaris.
altissima, Hort. ex Steud. Nom. ed. II. i. 200 = vulgaris.
bengalensis, Roxb. Hort. Beng. 21; Fl. Ind. ii. 59 = vulgaris.
Bourgeui, *Coss. Notes Crit.* 44.—Hispan.
campanulata, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 66.—Marocco.
carnosa, *Gren. in Gren. & Godr. Fl. Fr.* iii. 16.— Gallia.
chilensis, *Hort.-Cf. Gard. Chron.* (1870) i. 449.— Chili.
Cicla, Georgi, Beschr. Russ. Reich. iii. iv. 848 = trigyna.
Cicla, Linn. Syst. ed. XII. 195 = vulgaris.
crispa, Trott. Arch. vii. 117 = vulgaris.
Cycla, Pall. Tabl. Taur. 46 = trigyna.
decumbens, Moench, Meth. 382 = vulgaris.

BETA :—
diffusa, *Coss. Notes Crit.* 178.—Hispan.
esculenta, Salisb. Prod. 193 = vulgaris.
foliosa, Ehrenb. ex Steud. Nom. ed. II. i. 200 = vulgaris.
hastata, Desf. Cat. Hort. Par. ed. III. (1829) 389 = procumbens.
hortensis, Mill. Gard. Dict. ed. VIII. n. 2 = vulgaris.
hybrida, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1886) 134, nomen.—Rossia.
incarnata, Hort. ex Steud. Nom. ed. II. i. 200 = vulgaris.
intermedia, *Bunge, ex Boiss. Fl. Orient.* iv. 900.— Anatolia.
lomatogona, *Fisch. & Mey. in Bull. Soc. Nat. Mosc* (1836) 360.—Aa. Min.; Syria.
longispicata, Moq. Chenop. Enum. 18 = lomatogona.
lutea, Hort. ex Steud. Nom. ed. II. i. 200 = vulgaris.
macrocarpa, Guss. Fl. Sicul. Prod. i. 302 = vulgaris.
macrorhiza, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 267.—Reg. Caucas.; Persia.
marina, Crantz, Inst. i. 120 = vulgaris.
maritima, Linn. Sp. Pl. ed. II. 322 = vulgaris.
nana, *Boiss. & Heldr. Diagn.* Ser. I. vii. 82.— Graecia.
Notana, Bunge, ex Boiss. Fl. Orient. iv. 899 = vulgaris.
orientalis, Roth, Nov. Pl. Sp. 181 = vulgaris.
patellaris, *Moq. in DC. Prod.* iii. ii. 97.—Ins. Teneriff.
patula, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 315.—Ins. Madera.
procumbens, Lange, in Willk. & Lange, Prod. Fl. Hisp. i. 274 = diffusa.
procumbens, *Chr. Sm. in Hornem. Hort. Hafn. Suppl.* 81.—Ins. Canar.
pumila, Link, in Buch, Beschr. Canar. Ins. 141 = Webbiana.
purpurea, *Hort. ex Steud. Nom.* ed. II. i. 200 = vulgaris.
Rapa, Dum. Fl. Belg. 21 = vulgaris.
rapacea, Hegetschw. Fl. Schw. 256 = vulgaris.
rosea, Hort. ex Steud. Nom. ed. II. i. 200 = vulgaris.
rubra, *Delile, Fl. Aeg. Illustr.* 57.—Aegypt.
rubra, *Noronha, in Verh. Batav. Gen. v.* (1790) ed. I. Art. iv. 8.—Malaya.
sativa, Benth. Syst. Verc. Ecf. 162 = vulgaris.
stricta, C. Koch, in Linnaea, xxii. (1849) 180 = vulgaris.
sulcata, Gasp. in Guss. Fl. Sicul. Syn. ii. 800 = vulgaris.
sylvestris, Hort. Panorm. ex Webb & Berth. Phyt. Canar. iii. 197 = procumbens.
trifora, Salisb. Prod. 193 = vulgaris.
trigyna, *Waldst. & Kit. Pl. Rar. Hung.* 84. t. 85.— Hungar.; As. Min.
vulgaris, *Linn. Sp. Pl.* 322.—Europ.; Afr. bor.; Orient. Webbiana, *Moq. Chenop. Enum.* 16.—Ins. Canar.

BETCKEA, DC. Mém. Valer. 18; et Prod. iv. 642 (1839) = **Plectritis**, DC. (Rubiac.).
caucasica, Boiss. Fl. Orient. iii. 94.—Reg. Caucas.
Gillesii, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 366.—Reg. Argent.
heterophylla, Phil. in Linnaea, xxviii. (1856) 704.— Chili.
major, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 30 = P. samolifolia.
samolifolia, DC. Prod. iv. 642 = P. samolifolia.
samolifolia, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 366 = Gilliesii.

BETELA, Rafin. Sylva Tellur. 85 (1838) = **Piper**, Linn.
mastica, Rafin. l. c. = P. Betle.

BETENCOURTIA, St. Hil. Voy. i. 376 (1833) = **Galactia**, P. Br. (Legumin.).
rhynchosioides, St. Hil. l. c. = G. angustifolia.

BETHENCOURTIA, Choisy, in Buch, Beschr. Canar. Ins. 148 (1825). = **Senecio**, Tourn. (Compos.).
palmensis, Choisy, l. c. in obs. = S. palmensis.

BETONICA, [Tourn.] Linn. Syst. ed. I (1735) = **Stachys**, Tourn. (Labiat.).
affinis, Wender. in Flora, ix. (1826) 558 = S. Betonica.
alopecurus, Noé, in Bull. Soc. Bot. Fr. ii. (1856) 552 = S. Betonica.
Alopecuros, Linn. Sp. Pl. 578 = S. Alopecuros.

BETONICA :—
alpestris, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 93 = Stachys densiflora.
alpigena, Schur, Enum. Pl. Transs. 540.—Hungaria.
alpina, Mill. Gard. Dict. ed. VIII. n. 2 = S. Alopecuros.
angustifolia, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 97 = S. Betonica.
annua, Linn. Sp. Pl. 573 ; Moench, Meth. 395 = S. annua.
brachystachya, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 98 = S. Betonica.
Brantii, Boiss. Fl. Orient. iv. 750 = S. Brantii.
capensis, Burm. f. Fl. Cap. Prod. 16.—Afr. austr.
Clementei, Parre-Lara, in Anal. Soc. Esp. Hist. Nat. xi. (1882) 400.—Hispan.
danica, Mill. Gard. Dict. ed. VIII. n. 2 = S. densiflora.
decumbens, Moench, Meth. 396 = S. recta.
densiflora, Schur, Enum. Pl. Transs. 890, Index = S. densiflora.
divulsa, Tenore, Syll. Fl. Neapol. App. 614 = S. Betonica.
drymophila, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 100 = S. Betonica.
elongata, Willd. ex Stend. Nom. ed. II. i. 200 = S. Betonica.
flava, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = S. Alopecuros.
foliosa, Presl, *Fl. Sicul.* 87 = S. Betonica.
folosa, Rupr. Sert. Tianseh. 66.—As. centr.
frutescens, Bontekoe, Jav. 146 = Justicia Betonica.
glabrata, C. Koch, in Linnæa, xxi. (1848) 684 = S. Betonica.
grasca, Boiss. & Sprun. Diagn. Ser. I. v. 27 — Graecia.
grandiflora, Thuill. Fl. Par. ed. II. 293 = S. longifolia.
grandiflora, Steph. ex Willd. Sp. Pl. iii. 96 = S. grandiflora.
grandifolia, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 101 = S. Betonica.
hervelen, Linn. Mant. i. 83 = S. integrifolia.
hirsuta, Linn. f. c. ii. 248 = S. densiflora.
hirsuta, C. A. Mey. Verz. Pfl. Cauc. 95 = S. longifolia.
hirta, *Leyss. Fl. Halens.* 109 = S. Betonica.
hirta, Linn. Cent. ii. 20 ; Gouan, Hort. Monsp. 276 = S. recta.
hylebium, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 101 = S. Betonica.
incana, Mill. Gard. Dict. ed. VIII. n. 5 = S. densiflora.
Jacquini, Gren. & Godr. Fl. Fr. ii. 694 = S. Alopecuros.
laevigata, D. Don, Prod. Fl. Nep. 110 = Nepeta spicata.
lasiata, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 96 = S. Betonica.
legitima, Link, Handb. 476 = S. Betonica.
leiocalyx, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 95 = S. Betonica.
lutea, Lam. Fl. Fr. ii. 404 = S. Alopecuros.
macrantha, C. Koch, in Linnæa, xxi. (1848) 688 = S. grandiflora.
macrostachys, Wender. in Flora, ix. (1826) 358 = S. densiflora.
Monnieri, Gouan, Illustr. 86 = S. densiflora.
montana, Lej. Fl. Spa, ii. 24 = S. Betonica.
monticola, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 95 = S. Betonica.
nemorosa, Jord. & Fourr. l. c. 98 = S. Betonica.
nivea, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 266 = S. nivea.
nutans, Kit. ex Schult. Oestr. Fl. ed. II. ii. 146 = S. Betonica.
occitana, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 97 = S. Betonica.
officinalis, Linn. Sp. Pl. 578 = S. Betonica.
orientalis, Linn. l. c. = S. longifolia.
orientalis, Thuill. Fl. Par. 293 = S. grandiflora.
Orphanidea, Heldr. ex Boiss. Fl. Orient. iv. 750 = S. Alopecuros.
parvula, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 94 = S. Betonica.
polyclada, Jord. & Fourr. l. c. 103 = S. Betonica.
pradica, Zanted. ex Nym. Consp. 577 = S. densiflora.
pratensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 96 = S. Betonica.
psilostachys, Jord. & Fourr. l. c. 97 = S. Betonica.
purpurea, Gilib. Fl. Lituan. i. 69 = S. Betonica.
pyrenaica, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 94 = S. Betonica.

BETONICA :—
recurva, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 102 = Stachys Betonica.
recurvidens, Peterm. Fl. Lips. Excurs. 444 = S. Betonica.
rigida, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 103 = S. Betonica.
rosea, Hort. ex Heynh. Nom. ii. 67 = S. grandiflora.
rubicunda, Wender. Ind. Sem. Hort. Marb. ex Benth. in DC. Prod. xii. 460 = S. densiflora.
rusticana, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 96 = S. Betonica.
sabauda, Jord. & Fourr. l. c. 98 = S. Betonica.
scardica, Griseb. Spicil. Fl. Rumel. ii. 126 = S. grasca.
serotina, Host, Fl. Austr. ii. 165 = S. Betonica.
stricta, [Soland. in] Ait. Hort. Kew. ed. I. ii. 299 = S. Betonica.
strictiscalis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 99 = S. Betonica.
subcarnea, Jord. & Fourr. l. c. 98 = S. Betonica.
sylvulinea, Jord. & Fourr. l. c. 100 = S. Betonica.
valdepubens, Jord. & Fourr. l. c. 103 = S. Betonica.
vaisdula, Jord. & Fourr. l. c. = S. Betonica.
virescens, Jord. & Fourr. l. c. 99 = S. Betonica.
virgultorum, Jord. & Fourr. l. c. 100 = S. Betonica.
vulgaris, Rota, Prosp. Plant. Pav. 272 = S. Betonica.

BETULA, [Tourn.] Linn. Syst. ed. *I (1735). CUPULIFERAE*, Benth. & Hook. f. ui. 404.
BETULASTER, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 189.
acuminata, Ehrh. Beitr. vi. 98 = alba.
acuminata, Wall. Pl. As. Rar. ii 7. t. 109 = alnoides.
aetnensis, Raân. Giorn. i. 17 ; J. & C. Presl, Delic. Prag. i. 144 = alba.
affinis, Endl. Gen. Suppl. iv. 20 = alnoides.
alba, *Linn. Sp. Pl.* 982.—Europ. ; As. bor. ; Am. bor.
Alnobetula, Ehrh. Beitr. ii. 73 = Alnus viridis.
alnoides, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 58. —Reg. Himal.
Alnus, Linn. Sp. Pl. 983 = Alnus glutinosa.
Alnus, Tausch. Fl. Jap. 76 = Alnus maritima.
Alnus-crispa, Steud. Nom. ed. II. i. 201 = Alnus crispa.
Alnus-incana, Du Roi, Obs. Bot. 31 = Alnus incana.
Alnus-rugosa, Du Roi, l. c. 32 = Alnus serrulata.
alpestris, *Fries, Summ. Veg. Scand.* i. 212.—Europ. bor. ; Groenland.
alpina, Borckh. ex Reichb. Fl. Germ. Excurs. 174 = Alnus viridis.
ambigua, Hampe, ex Reichb. Ic. Fl. Germ. xii. 2 = alba.
angulata, *Hort. ex Steud. Nom.* ed. II. i. 201.—Am. bor.
antarctica, Forst. in Comm. Gott. ix. (1789) 45 = Fagus betuloides.
arguta. Schlecht. in Linnæa, vii. (1832) 189 = Alnus acuminata.
asplenifolia, Hort. ex Regel, in DC. Prod. xvi. II. 167 = alba.
aurata, Borkh. Forstb. i. 378 = alba.
aurea, Steud. Nom. ed. II. i. 201 = alba.
Bhojpattra, Wall. Pl. As. Rar. ii. 7 = utilis.
borealis, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 196 = pumila.
braccembergensis, Becbst. Forstb. 297 = alba.
carpathica, Waldst. & Kit. ex Willd. Sp. Pl. iv. 464 = alba.
canadensis, Hort. ex C. Koch, Dendrol. ii. 11. 656 = alba.
carpinifolia, Ehrh. Beitr. vi. 99 = lenta.
carpinifolia, *Sieb. & Zucc. in Abh. Acad. Muench.* iv. *Abth.* iii. (1846) 228.—Japon.
chinensis, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 47.—China.
ciliata, Stend. Nom. ed. II. i. 201 = Populus ciliata.
cordata, Loisel. Notul. 139 = Alnus cordifolia.
cordifolia, Regel, in Nouv. Mém. Soc. Nat. Mosc. xiii. (1861) 86 = alba.
corifolia, Tausch, ex Regel, in DC. Prod. xvi. II. 168 = alba.
corylifolia, *Regel & Maxim. in Bull. Soc. Nat. Mosc.* xxxvii. (1865) ii. 417.—Japon.
costata, Trautv. in Maxim. Prim. Fl. Amur. 258 = alnifolia.
crispa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 339 = Alnus viridis.
cryptocarpa, Laestad. in Bot. Notiser, 1836 (1837) 109 = alba.

BETULA :—
cuspidata, Schrad. ex Regel, in DC. Prod. xvi. II. 164 = alba.
cycovicensis, Steud. Nom. ed. I. 107, sphalm. = cycovicnsis.
cylindrostachya, Wall. Fl. As. Rar. ii. 7 = alnoides.
dalecarlica, Linn. f. Suppl. 416 = alba.
davurica, *Pall. Fl. Ross.* i. 60. t. 39. fig. A.—Sibir. ; Am. bor.
dilcica, *Pall. Reise*, iii. 321.—Sibir. transbaical.
divaricata, Ledeb. in Denkschr. Bater. Bot. Ges. iii. (1841) 59 = fruticosa.
emarginata, Ehrh. Beitr. iv. 158 = Alnus glutinosa
Ermani, *Cham. in Linnæa*, vi. (1831) 587. t. 6. fig. D. —As. bor. or.
ethnensis, Raân. ex Regel, in DC. Prod. xvi. II. 168 = alba.
exaltata, *S. Moore, in Journ. Linn. Soc.* xvii. (1880) 386. t. 16.—China.
excelsa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 337 = papyracea !
excelsa, Hook. Fl. Bor. Am. ii. 156 = lenta.
excelsa, Pursh, Fl. Am. Sept. ii. 621 = lutea.
Friesii, Lasm. Fl. Vermi. ed. II. ex Hartm. Scand. Fl. ed. XIII. 376 = alba.
fruticans, *Pall. Reise*, iii. t. kk = fruticosa.
fruticosa, Pries, in Flora, xxxiii. (1850) 182 = alpestris.
fruticosa, *Pall. Fl. Ross.* i. t. 62. t. 40. f. A. B. C.—Europ. ; As. bor. ; Sibir. ; Manchur.
fruticosa, Willd. Sp. Pl. iv. 467 = humilis.
fusca, Pall. Reise, iii. 278 = fruticosa.
glabra, *Dum. Fl. Belg.* 11.—Belgia.
glandulosa, *Michx. Fl. Am. Bor.* ii. 180.—Am. bor.
glauca, Wender. in Bot. Zeit. iv. (1846) 291 = alba.
glutinosa, Lam. Encyc. i. 454 = Alnus glutinosa.
glutinosa, Wallr. Sched. Crit. 497 = alba.
Gmelini, Bunge, Verz. Suppl. Fl. Alt. 118 = fruticosa.
grandis, Schrad. in Ind. Sem. Hort. Gott. (1838) 2 = alba.
Grayii, Regel, in Bull. Soc. Nat. Mosc. xxxvii. (1865) ii. 406 = pumila.
grossa, *Sieb. & Zucc. in Abh. Acad. Muench.* iv. *Abth.* iii. (1846) 228.—Japon.
gummifera, Bertol. Fl. Ital. x. 229 = alba.
Hackelii, Opiz, ex Steud. Nom. ed. II. i. 201 = alba
harcynica, Wender. ex Steud. l. c. = alba.
humilis, *Fries*, ex Regel, in DC. Prod. xvi. II. 172 = alpestris.
humilis, *Schrank, Bayer. Fl.* i. 420.—Europ. bor. ; As bor. ; Am. bor.
hybrida, Blom. in Vet. Acad. Nya Handl. vii. (1786) 186 t. 6. f. 18 = alba.
hybrida, *Regel, in Nouv. Mém. Soc. Nat. Mosc.* xiii. (1861) 94.—Rossia.
incana, Roth, Tent Fl. Germ. iii. 477 = Alnus incana. intermedia, *Thomas, ex Reichb. Ic. Fl. Germ.* xii. t. 624. fig. 1288.—Europ.
Jacquemontii, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 189 = utilis.
japonica, Siebold, in Verh. Batav. Gen. xii. (1830) 25 = alba.
kamtschatica, Buek, in DC. Prod. Ind. iv. 52 = humilis.
laciniata, Blom, in Vet. Acad. Nya Handl. vii. (1786) 186. t. 6. f. 18 = alba.
laciniata, Ehrh. Beitr. iii. 21 = Alnus glutinosa.
laciniata, Thunb. Diss. Bet. 10 = alba.
lagopina, Hartm. ex Regel, in DC. Prod. xvi. II 178 = alpestris.
lanulosa, Michx. Fl. Bor. Am. ii. 181 = nigra.
latifolia, Tausch, in Flora, xxi. (1838) 731 = alba.
lenta, Darol. Harbk. Baume. i. 93 = alba.
lenta, *Linn. Sp. Pl.* 983.—Am. bor.
lenta, Regel, in Nouv. Mém. Soc. Nat. Mosc. xiii. (1861) 126 = carpinifolia.
leptophylla, Regel, in Prod. xvi. II. 181 = Alnus nepalensis.
leptostachya, Wall. Cat. n. 2799 = Alnus nepalensis.
Littelliana, Tuckerm. in Am. journ. Sc. xlv. (1843) 80 = glandulosa.
lobulata, *Kit. in Linnæa*, xxxii. (1863) 951.—Europ.
longifolia, Bové, ex Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 208 = Alnus orientalis.
lutea, *Michx. Hist. Arb.* iii. 188. t. 5.—Am. bor.
Maackii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 366 = davurica.
major, Gilib. Exercit. ii. 400 = alba.

BETULA:—
macrostachya, Schrad. ex Regel, in DC. Prod. xvi. 167; Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 7 = alba.
Maximowiciana, Regel, in DC. Prod. xvi. 180 = Maximowiczii.
Maximowiczii, *Regel, in Bull. Soc. Nat. Mosc.* xxxviii (1866) ii. 418.—Japon.
Mandshurica, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 189 = davurica.
Maximowiczkii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 485 = dahurica.
media, Laestad. in Bot. Notiser, 1856 (1857) 107 = alba.
megalocarpa, Laestad. l. c. 106 = alba.
Michauxii, *Spach, in Ann. Sc. Nat.* Sér. II. xv. (1841) 196.—Am. bor.
microphylla, *Bunge, Verz. Suppl. Fl. Alt.* 119; *in Mém. Acad. Pétersb.* (1835) 506.—Sibir. altaic.
Middendorffii, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 84).—Manchuria.
Murithii, Gaudin, ex Regel, in DC. Prod. xvi. ii. 168 = alba.
myrsinoides, Tausch, in Flora, xi. (1828) 758 = humilis.
nana, Hook. Fl. Bor. Am. ii. 156 = glandulosa.
nana, *Linn. Sp. Pl.* 983.—Reg. bor.
nana, Michx. Fl. Bor. Am. ii. 180 = Michauxii.
nigra, Dubam. Arb. ed. Nov. iii. 208. t. 51 = alba.
nigra, Durol, Harbk. Baumz. i. 98 = lenta.
nigra, *Linn. Sp. Pl.* 982.—Am. bor.
nigra, Murith, Guide Bot. Valais, 55 = intermedia
nigricans, Wender. in Bot. Zeit. iv. (1846) 291 = alba.
nitida, D. Don, Prod. Fl. Nep. 58 = alnoides.
oblongata, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 338 = Alnus glutinosa.
occidentalis, *Hook. Fl. Bor. Am.* ii. 155.—Am. bor. occ.
odorata, Bechst. Forstb. 294 = alba.
ovalifolia, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 376 = humilia.
ovata, C. Koch, in Linnaea, xxii. (1849) 338 = alba.
ovata, Schrank, Baier. Fl. i. 419 = Alnus viridis.
oycoviensis, Bess. Prim Fl. Galic. ii. 289 = alba.
oycoviensis, Laestad. ex Regel, in DC. Prod. xvi. ii. 171 = intermedia.
oycoviensis, Reichb. Ic. Fl. Germ. xii. t. 622. f. 1281 = humilis.
palmaeformis, Laestad. in Bot. Notiser, 1856 (1857) 106 = alba.
palmata, Borkh. Handb. i. 500 = alba.
palustris, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 376 = humilis.
palustris, Salisb. Prod. 396 = Alnus glutinosa.
papyracea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 337.—Am. bor.
papyrifera, Michx. Fl. Bor. Am. ii. 180 = papyracea.
pendula, Roth, Tent. Fl. Germ. i. 405 = alba.
persicifolia, Hort. ex C. Koch, Dendrol. ii. ii. 641 = lutea.
pinnata, Landmark, in Vet. Acad. Nya Handl. Stockh. (1790) 181. f. 5 = Alnus incana.
pirifolia, Hort. ex C. Koch, Dendrol. *ii.* ii. 646 = papyracea.
pontica, *Dcsf. ex Steud. Nom.* ed. II. i. 301.—As. Min.
pontica, Hort. ex Loud. Encyc. Pl. 720 = alba.
populifolia, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 336 = alba.
pubescens, Ehrh. Beitr. iv. 160; et vi. 88 = alba.
pumila, *Linn. Mant.* i. 134.—Am. bor.
pumila, J. Vahl, ex Lange, Comp. Fl. Groenl. i. 114 = alpestris ?
quebecensis, Burg. in Ges. Naturf. Fr. Berl. Schr. v. (1784) 196 = humilis.
reticulata, Laestad. ex Regel, in DC. Prod. xvi. ii. 169 = alba.
reticulata, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 378 = humilis.
rhombifolia, Tausch, in Flora, xxxviii. (1838) 753 = alba.
rotundifolia, Regel & Tiling, Fl. Ajan. 118 = Middendorffii.
rotundifolia, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 194 = glandulosa.
rubra, Host. ex Regel, in DC. Prod. xvi. ii. 167 = alba.

BETULA:—
rubra, Michx. Hist. Arb. ii. 143. t. 8 = nigra.
rugosa, Ehrh. Beitr. iii. 21 = Alnus serrulata.
rustica, Laestad. in Bot. Notiser, 1856 (1857) 104 = alba.
Schmidtii, *Regel, in Bull. Soc. Nat. Mosc.* xxxviii. (1865) ii. 412.—Manchuria.
serrulata, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 338 = Alnus serrulata.
sibirica, Lodd. ex Regel, in DC. Prod. xvi. ii. 173 = humilis.
silvatica, Laestad. in Bot. Notiser, 1856 (1857) 108 = alba.
silvestris, Laestad. l. c. 107 = alba.
Socolowii, Jacq. f. ex Regel, in DC. Prod. xvi ii. 178 = humilis.
Sokolowii, Hort. Petrop. ex Regel, l. c. 168 = alba.
subaequalis, Laestad. in Bot. Notiser, 1856 (1857) 109 = alba.
subalpina, Laestad. l. c. 110 = alba.
tianschanica, *Rupr. Sert. Tianach.* 72.—As. centr.
tiliaefolia, Laestad. in Bot. Notiser, 1856 (1857) 106 = alba.
tomentosa, Retz. & Abel, Beschr. Holsarten, i. 17. t. 16 = alba.
torfacea, Schleich. Cat. Pl. Helv. ed. IV. 9 = alba.
tortuosa, Ledeb. Fl. Ross. iii. 653 = alba.
transsilvanica, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 68.—Hungar.
triphylla, *Thunb. Diss. Betul.* 12.—Hab ?
tristis, *Wormsk. ex Link, Enum. Hort. Berol.* ii. 401. —Kamtschat.
turfosa, Weigel. ex Reichb. Fl. Germ. Excurs. 174 = fruticosa.
ulmifolia, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 228.—Japon.
urticaefolia, Curtois, ex Steud. Nom. ed. II. i. 301 = alba ?
urticifolia, *Regel, in Nouv. Mém. Soc. Nat. Mosc.* xiii. (1861) 115.—Suecis (cult.).
utilis, *D. Don, Prod. Fl. Nep.* 58.—Reg. Himal.; Japon.
verrucosa, Ehrh. Beitr. v. 161; vi. 98 = alba.
virgata, Salisb. Prod. 395 = alba.
virgultosa, Fries, ex Regel, in Nouv. Mém. Soc. Nat. Mosc. xiii. (1861) 112 = urticifolia.
viridis, Chaix, in Vill. Hist. Pl. Dauph. iii. 789 = Alnus viridis.

BETULASTER, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 195 = **Betula**, Tourn. (Cupulif.).
acuminata, Spach, l. c. = Betula alnoides.
affinis, Spach, l. c. 199 = Betula alnoides.
cylindrostachya, Spach, l. c. 198 = Betula alnoides.
nitida, Spach, l. c. 200 = Betula alnoides.

BEURERIA, Ehret, Pl. & Pap. Rar. t. 18 (1755) = **Calycanthus**, Linn.

BEURRERIA, Jacq. Enum. Pl. Carib. 2 (1760) = **Bourreria**, P. Br. (Borag.).

BEVERNA, Adans. Fam. ii. 90 (1763) = **Iris**, Tourn.?

BEYERIA, Miq. in Ann. Sc. Nat. Sér. III. i. (1844) 350. t. 15. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 268.
BEYERIOPSIS, Muell. Arg. in Linnaea, xxxiv. (1865) 66.
CALYPTROSTIGMA, Klotzsch, in Lehm. Pl. Preiss. i. 175 (1844).
Backhousii, Hook. f. Fl. Tasm. i. 239 = opaca.
brevifolia, *Baill. Adansonia*, vi. (1865–66) 309.—Austral.
cinerea, *Baill. l. c.*—Austral.
cyanescens, *Benth. Fl. Austral.* vi. 66.—Austral.
cygnorum, *Baill. Adansonia*, vi. (1865–66) 309.—Austral.
Drummondii, *F. Muell. Census*, 18.—Austral.
lasiocarpa, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 59.—Austral.
latifolia, *Baill. Adansonia*, vi. (1865–66) 304.—Austral.
ledifolia, Sond. in Linnaea, xxvii. (1856) 565 = opaca.
lepidopetala, *F. Muell. Fragm.* i. 230.—Austral.
Leschenaultii, Baill. Adansonia, vi. (1865–66) 307 = opaca.

BEYERIA:—
lorantholides, *Baill. Etud. Gén. Euph.* 408.—Austral.
oblongifolia, Sond. in Linnaea, xxvii. (1856) 564 = viscosa.
opaca, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 16.—Austral.
similis, *Baill. Adansonia*, vi. (1865–66) 309.—Austral
tristigma, *F. Muell. Fragm.* vi. 181.—Austral.
uncinata, *F. Muell. ex Baill. Adansonia.* vi. (1865–66) 306.—Austral.
viscosa, *Miq. in Ann. Sc. Nat.* Sér. III. i. (1844) 350. t. 15.—Austral.

BEVERIOPSIS, Muell Arg. in Linnaea, xxxiv. (1865) 66 = **Beyeria**, Miq. (Euphorb.).
brevifolia, Muell. Arg. l. c. 68 = Beyeria brevifolia.
cinerea, Muell. Arg. l. c. 67 = Beyeria cinerea.
cyanescens, Muell. Arg. in DC. Prod. xv. ii. 200 = Beyeria cyanescens.
cygnorum, Muell. Arg. in Linnaea, xxxiv. (1865) 66 = Beyeria cygnorum.
Drummondi, Muell. Arg. l. c. 58 = Beyeria Drummondi.
lasiocarpa, Muell. Arg. l. c. 59 = Beyeria lasiocarpa.
latifolia, Muell. Arg. l. c. 57 = Beyeria latifolia.
lepidopetala, Muell. Arg. l. c. = Beyeria lepidopetala.
similis, Muell. Arg. l. c. 58 = Beyeria similis.

BEYRICHIA, Cham. & Schlecht. in Linnaea, iii. (1828) 21. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 949
ACHETARIA, Cham. & Schlecht. in Linnaea, ii. (1827) 586, nomen prius
arborescens, Benth. Scroph. Ind. 9, in nota = ocymoides.
floribunda, *Benth. in DC. Prod.* x. 279.—Bras.
ocymoides, *Cham. & Schlecht. in Linnaea*, iii. (1828) 21.—Bras.
scutellarioides, *Benth. Scroph. Ind.* 9, in nota.—Bras.
villosa, *Benth. l. c.*—Bras.

BEYTHEA, Endl. Gen. 1011 (1840) = **Blasicoarpus**, Linn. (Tiliac.).
bifida, Endl. l. c. = E. bifidus.

BEZANILLA, Remy, in C. Gay, Fl. Chil. iv. 109. t. 46 (1849) = **Petiocarphus**, Nutt. (Compos.).
chilensis, Remy, l. c. 110.—Chili.

BHESA, Buch.-Ham. ex Arn. in Edinb. Phil. Journ. xvi. (1834) 315 = **Kurrimia**, Wall. (Celastr.).
Moja, Buch.-Ham. l. c. = K. pulcherrima.
paniculata, Arn. l. c. = K. paniculata.

BIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 189 = **Tragia**, Linn. (Euphorb.).
alienata, F. Didr. in Kjoeb. Videnak. Meddel. (1857) 181 = T. Sellowiana.
Lessertiana, Baill. Etud. Gén. Euph. 502 = T. Lessertiana.
Lhotskyana, Klotzsch, in Wiegm. Archiv, vii. (1841) 190 = T. Sellowiana.
Sellowiana, Klotzsch, l. c. = T. Sellowiana.

BIANCAEA, Todaro, Nuov. Gen. 21 (1858) = **Caesalpinia**, Linn. (Legum.).
mimosoides, Todaro, Hort. Bot. Panorm. 3 = C. mimosoides.
Sappan, Todaro, l. c. = C. Sappan.
scandens, Todaro, Nuov. Gen. 22; et l. c. 4. t. 1 = C. sepiaria.

BIARUM, Schott, Melet. i. 17 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 986.
CYLLENIUM, Schott, Gen. Aroid. t. 9 (1858).
HOMAID, Adans. Fam. ii. 470 (1763), nomen prius.
ISCHARUM, Blume, Rumphia, i. 144. t. 29 (1835).
LAETOPETION, Schott, Gen. Aroid. t. 8 (1858).
STENURUS, Sallab. Gen. Pl. Fragm. t. 5 (1866).
abbreviatum, Schott, Prod. Aroid. 63 = tenuifolium.
alexandrinum, Boiss. Diagn. Ser. I. xiii. 6 = Olivieri.
Anguillarae, Schott, Prod. Aroid. 62 = tenuifolium.
angustatum, *N. E. Br. in Journ. Linn. Soc.* xvii. (1880) 255.—Orient; Syria.
arundanum, Boiss. & Kent. Pugill. 110 = tenuifolium.

BIARUM :—
Bovei, *Blume, Rumphia*, i. 114. t. 29.—Syria; As. Min.
carduchorum, *Engl. in DC. Monog. Phan.* ii. 575.—As. Min.; Syria.
constrictum, C. Koch, Ind. Sem. Hort. Berol. (1858) 19 = tenuifolium.
eximium, *Engl. in DC. Monog. Phan.* ii. 576.—Cilicia.
Frasaianum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1880) 254.—Oriens.
gramineum, Schott, Melet. i. 17 = tenuifolium.
Haenseleri, *Willk. in Bot. Zeit.* v. (1847) 49. t. 2.—Hispan.
Homeld, *Blume, Rumphia*, i. 115.—Oriens.
Lehmanni, Bunge, in Mém. Sav. ltr. Pétersb. vii. (1851) 237 = Helicophyllum Lehmanni.
longifolium, *Pomel, Nouv. Mat. Fl. Atl.* 391.—Algeria.
macroglossum, *Pomel, l. c.* 390.—Algeria.
numidicum, *Parl. Fl. Ital.* ii. 243.—Numidia.
Olivieri, *Blume, Rumphia*, i. 115.—Aegypt.
Pyrami, *Engl. in DC. Monog. Phan.* ii. 576.—Cilicia; Syria.
rhopalospadix, C. Koch, Ind. Sem. Hort. Berol. (1853) 12 = Sprunerl.
rupestre, *Pomel, Nouv. Mat. Fl. Atl.* 391.—Algeria.
Russellianum, *Schott, Prod. Aroid.* 63.—Syria.
Sewerswi, *Regel, in Act. Hort. Petrop.* vi. (1879) 489. —Caucasus.
Sprunerl, *Boiss. Diagn.* Ser. I. xiii. 5.—Graecia.
Sprunerl, Schott, in Gen. Aroid. t. 7 = abbreviatum.
tenuifolium, Heldr. ex Nym. Consp. 756 = abbreviatum.
tenuifolium, *Schott, Melet.* i. 17.—Hispan.
Zeleborii, Schott, in Oestr. Bot. Wochenbl. (1857) 245 = tenuifolium.

BIASLIA, Vand. Fl. Lusit. et Bras. Spec. 4. t. 1. f. 2 (1788) = **Mayaca**, Aubl.

BIASOLETTIA, Koch, in Flora, xix. (1836) i. 163; Bertol. Fl. Ital. iii. 191 (1837) = **Chaerophyllum**, Linn. (Umbellif.).
congesta, Nym. Syll. Suppl. 38.—Graecia.
cretica, Nym. Consp. 303 = C. creticum.
divaricata, Nym. Syll. Suppl. 38.—Graecia.
macrocarpa, Nym. Consp. 303.—Graecia.
nodosa, Bertol. Fl. Ital. iii. 191 = C. nodosum.
parnassica, Nym. Syll. Suppl. 37.—Graecia.
pumila, Nym. Consp. 302.—Graecia.
tuberosa, Koch, in Flora, (1836) i. 163 = Carum cylindricum!

BIASOLETTIA, Pohl, ex Baker, in Mart. Fl. Bras. vi. ii. 311 = **Bupatorium**, Linn. (Compos.).
elliptica, Pohl, ex Baker, l. c. = E. trixoides.

BIASOLETTIA, Presl, Rel. Haenk. ii. 141 (1840 !) = **Hernandia**, Linn. ! (Laurin.).
nymphaeaefolia, Presl, l. c.—Ins. Marian.

BIASSOLETTIA, Endl. Gen. 1004 (1840) = BIASOLETTIA, Presl (= **Hernandia**, Linn. ! Laurin.).

BIATHERIUM, Desv. Opusc. 72 (1821) = **Gymnopogon**, Beauv. (Gramin.).
foliosum, Desv. l. c. = G. foliosum.

BICCHIA, Parl. Fl. Ital. iii. 396 (1858) = **Habenaria**, Willd. (Orchid.).
albida, Parl. l. c. 397 = H. albida.

BICHEA, Stokes, Bot. Mat. Med. ii. 564 (1812) = **Cola**, Schott (Stercul.).
solitaria, Stokes, l. c. 565 = C. acuminata.

BICHENIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 236 = **Trichocline**, Cass. (Compos.).
aurea, D. Don, l. c. 237.—Chili.
auriculata, Wedd. Chlor. And. i. 26.—Bolivia.
Berteriana, Steud. Nom. ed. II. i. 203 = *aurea*.
Cineraria, D. Don, in Phil. Mag. (1828) 391 = T. Cineraria.
crenata, Remy, in C. Gay, Fl. Chil. 302.—Chili.
dealbata, D. Don, ex Hook. Comp. Bot. Mag. i. (1835) 104 = T. dealbata.

BICHENIA :—
plicata, D. Don, ex Wedd. Chlor. And. i. 26 = Trichocline plicata.
reptans, Wedd. l. c. 25.—Bolivia.
sinuata, D. Don, Phil. Mag. (1832) 391 = T. incana.

BICORNELLA, Lindl. Bot. Reg. sub t. 1701 (1735). *ORCHIDEAE*, Benth. & Hook. f. Gen. iii. 634.
gracilis, *Lindl. Gen. et Sp. Orch.* 334.—Madag.
longifolia, *Lindl. l. c.* 335.—Madag.
parviflora, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 300. —Madag.

BICORONA, A. DC. Prod. viii. 330 (1844) = **Melodinus**, Forst. (Apocyn.).
phyllireoides, A. DC. l. c. = M. phillyraeoides.

BICUCULLA, Borckh. in Roem. Arch. i. ii. 46 (1797) = **Adlumia**, Raf. (1808) (Papav.).
fumariodes, Borckh. l. c. = A. cirrhosa.

BICUCULLARIA, Juss. ex Steud. Nom. ed. II. i. 202 (1840) = **Dicentra**, Benth. (Papav.).
canadensis, Juss. ex Steud. l. c. 201 = D. Cucullaria.

BIDARIA, Decne. in DC. Prod. viii. 623 (1844) = **Gymnema**, N. Br. (Asclep.).
elegans, Dalz. & Gibs. Bomb. Fl. 151 = G. montanum.
elegans, Decne. in DC. Prod. viii. 623 = G. elegans.
erecta, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 59 = G. etenophyllum.
inodora, Decne. in DC. Prod. viii. 624 (Sp. dub.).—Ind. or.
leptophylla, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 60 = Marsdenia leptophylla.
pergularioides, Thw. Enum. Pl. Zeyl. 196 = G. pergularioides.
pubiflora, Miq. Fl. Ind. Bat. ii. 497.—Java.
syringaefolia, Decne. in DC. Prod. viii. 623 = Marsdenia syringaefolia.
tingens, Decne. l. c. = G. montanum.
trinervia, Decne. l. c. 624 = G. trinerve.

BIDENS, (Tourn.) Linn. Gen. ed. I. 348 (1737) = *COMPOSITAE*, Benth. & Hook. f. ii. 337.
CERATOCEPHALUS, Vaill. ex Cass. in Dict. Sc. Nat. vii. 432 (1817).
DELISEA, DC. Prod. v. 683 (1836).
EDWARSIA, Neck. Elem. i. 87 (1790).
KERNERIA, Moench, Meth. 595 (1794).
PLUMBIDENS, Neck. Elem. i. 86 (1790).
Abadiae, *DC. Prod.* v. 601.—Peruv.
abortiva, Schum. & Thonn. Beskr. Guin. Pl. 381 = pilosa.
abyssinica, Sch. Bip. ex Walp. Rep. vi. 167 = pilosa, quadriseta.
acaulis, *Baker, in Mart. Fl. Bras.* vi. iii. 247.—Para.
Acmella, Lam. Encyc. i. 415 = Spilanthes Acmella.
acmelioides, Berg. in Vet. Acad. Handl. xxix. ii. (1768) 246 = Spilanthes Acmella.
adhaesicarpa, *Vell. Fl. Flum.* 348; viii. t. 88.—Bras.
alausanaia, *H. B. & K. Nov. Gen. et Sp.* iv. 285.—Recent.
alba, DC. Prod. v. 606 = pilosa.
andicola, *H. B. & K. Nov. Gen. et Sp.* iv. 257.—Ecuador.
angustifolia, Lam. Encyc. i. 415 = Spilanthes urens.
angustifolia, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 389.—Ins. Sandvic.
angustiana, *H. B. & K. Nov. Gen. et Sp.* iv. 236.—Mexic.
angustissima, Sprong. Syst. iii. 453 = andicola.
anthicoides, *DC. Prod.* v. 600.—Mexic.
aplifolia, Linn. Syst. ed. X. 1206; Fl. Jam. Pugill. 22 = Chrysanthellum procumbens.
arborea, Roxb. in Hexb. App. 301, 325 = Petrobium arboreum.
arguta, H. B. & K. Nov. Gen. et Sp. iv. 231 = heterophylla.
artemisiaefolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 49. —Peruv.
atriplicifolia, Linn. Cent. ii. 30; Amoen. Acad. iv. 329 = Isocarpha atriplicifolia.
atrosanguinea, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 269 = Dahlia Zimapani.

BIDENS :—
atrosanguinea, Ortg. ex Regel, Gartenfl. x. (1861) 406 t. 347 = Cosmos diversifolius.
australis, Sprong. Syst. iii. 458 = Coreopsis micrantha.
Bardanna, Wall. Cat. n. 3221 = Chrysanthellum indicum.
Beckii, *Torr. in Sprong. Neue Entdeck.* ii. 135.—Am. bor.
Berteriana, Sprong. Syst. iii. 454 = Cosmos caudatus.
Bigelovii, *A. Gray, in Torr. Bot. Mex. Bound.* 91.—Am. bor. occ.
bimucronata, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 184.—Ins. S. Doming.
bipinnata, Ball. Hist. Pl. viii. 50 = Cosmos bipinnatus.
bipinnata, Hook. & Arn. in Bot. Beech. Voy. i. 30 = chilensis.
bipinnata, *Linn. Sp. Pl.* 832.—Reg. trop.
Bonplandii, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos tenellus.
brachycarpa, *DC. Prod.* v. 600.—Mexic.
Buchingeri, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop 142 = Schimperi.
bullata, *Linn. Sp. Pl.* 833.—Am. bor.; Europ.
californica, DC. Prod. v. 599 = pilosa.
calva, C. B. Clarke, Comp. Ind. 141 = Cosmos sulphureus.
canescens, *Bertol. Fl. Guatimal.* 31.—Guatemala.
cannabina, Lam. Fl. Fr. ii. 44 = tripartita.
caracasana, *DC. Prod.* v. 600.—Venezuela.
carnea, *Herr. Ind. Sem. Hort. Turic.* (1842).—Guiana.
carvifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos carvifolius.
cascalides, *DC. Prod.* v. 604.—Mexic.
caudata, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos caudatus.
cernua, *Linn. Sp. Pl.* 832.—Reg. bor. temp.
Cervantesii, Baill. Hist. Pl. viii. 305 = Dahlia coccinea.
chilensis, *DC. Prod.* v. 602.—Chili.
chinensis, Willd. Sp. Pl. iii. 1719 = pilosa.
chrysantha, *DC. Prod.* v. 605.—Ind. occ.
chrysanthemoides, *Michx. Fl. Bor. Am.* ii. 136.—Am. bor.
cicutaefolia, *Tausch, in Flora*, xix. (1836) 395.—Hab. ?
ciliata, *Hoffmgg. ex Flech. & Mey. Ind. Sem. Hort. Petrop.* vi. 46.—Hab. ?
coccinea, Baill. Hist. Pl. viii. 49 = Dahlia coccinea.
connata, *Muhl. in Willd. Sp. Pl.* iii. 1716.—Am. bor.
consolidaefolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 185.—Peruv.
coordifolia, *Sch. Bip. in Flora*, xxxix. (1856) 361.—Ins. Pacif.
coreopsidis, DC. Prod. v. 599 = rubifolia.
coronata, Fisch. ex Steud. Nom. ed. II. i. 202 = Coreopsis coronata.
Cosmenthus, *Griseb. in Gentt. Abh.* xix. (1874) 185.—Reg. Argent.
costaricensis, *Benth. ex Oerst. in Kjoeb. Videnskf. Meddel.* (1859) 94.—Costa Rica.
crithmifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 234.—Cuba.
crocata, Cav. Ic. i. 66. t. 99 = Verbesina crocata.
cynaefolia, *H. B. & K. Nov. Gen. et Sp.* iv. 235.—Cuba.
daucifolia, *DC. Prod.* v. 601.—Mexic.
decussata, *H. B. & K. Nov. Gen. et Sp.* iv. 233.—Mexic.
decomposita, Wall. Cat. n. 3223 = pilosa.
decussata, Pav. ex Steud. Nom. ed. II. i. 202 = hispida.
delphinifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 236.—Mexic.
denudata, Turcz. in Bull. Soc. Nat. Mosc. xviii. (1851) i. 153 = Glossogyne tenuifolia.
dichotoma, *Desf. Tabl. ed.* I. (1804) 204.—Brus.
Dilleniana, *Hill, Veg. Syst. iii.* 138.—Hab. ?
diversifolia, *Ortg. in Regel, Gartenfl.* (1861) 403.—Mexic.
diversifolia, Hort. ex DC. Prod. v. 602 = grandiflora.
diversifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 306 = Cosmos diversifolius.
diversifolia, Willd. ex DC. Prod. v. 602 = alausanaia.
dondiaefolia, *Less. in Linnaea*, v. (1830) 156.—Mexic.

BIDENS:—
elongata, *Tausch, in Flora*, xix. (1836) 896.—Hab.? exsiliata, *DC. Prod.* v. 600.—Mexic.
fastigiata, Michalet, in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 20 = tripartita.
ferulæfolia, *DC. Prod.* v. 603.—Mexic.
ferulæfolia, Hemsl. Biol. Centr. Am. Bot. ii. 202, partim = procera.
feroida, Lam. Encyc. i. 415 = Spilanthes Acmella.
fœtulcus, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 250.—Bras.
fœa, Hook. f. Fl. Brit. Ind. iii. 307, sphalm. = fusca.
flagellaris, *Baker, in Mart. Fl. Bras.* vi. III. 248.—Bras.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iv. 238.—Ecuador.
foeniculifolia, DC Prod. v. 603 = procera?
foliosa, *Willd. Enum. Hort. Berol. Suppl.* 66, nomen. —Hab.?
frondosa, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 309 = tripartita.
frondosa, *Linn. Sp. Pl.* 832.—Am. bor.
frutescens, Mill. Gard. Dict. ed. VIII. n. 4 = Salmea grandiceps.
fruticosa, *DC. Prod.* v. 596.—Hab.?
fruticosa, Linn. Sp. Pl. 883 = Wulffia stenoglossa.
fruticulosa, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 371.—Peruv.
fusca, Lam. Encyc. i. 416 = Spilanthes Acmella.
Gardneri, *Baker, in Mart. Fl. Bras.* vi. III. 246.—Bras.
glaberrima, *DC. Prod.* v. 601.—Am. austr.
glycinæfolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 247.—Bras.
gracilis, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 368.—Ins. Sandvic.
gracilis, Torr. in Ann. Lyc. N. York, ii. (1828) 215 = Thelesperma gracile.
grandiflora, *Balb. Cat. Hort. Taur.* (1812) 19.—Am. austr.
graveolens, *Mart. in Isis*, (1834) 590.—Bras.
hawaiensis, *A. Gray, in Proc. Am. Acad.* v. (1861) 128. —Ins. Hawai.
helianthoides, H. B. & K. Nov. Gen. et Sp. iv. 230 = chrysanthemoides.
heterophylla, *Ortog. Hort. Matr.* 99. t. 12.—Am. bor. occ.; Mexic.
heterosperma, *A. Gray, Pl. Wright.* ii. 90.—Arizona.
hirsuta, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 369 = pilosa.
hirsuta, Swartz, Prod. Veg. Ind. Occ. 110 = Salmea hirsuta.
hirta, Jord. in Gren. & Godr. Fl. Fr. ii. 168 = bullata.
hirtella, *H. B. & K. Nov. Gen. et Sp.* iv. 232.—Mexic.
hispida, *H. B. & K. l. c.* 237.—Venezuela.
Humboldtii, Sch. Bip. in Seem. Bot. Voy. Herald, 307 = Cosmos parviflorus.
humilis, *H. B. & K. Nov. Gen. et Sp.* iv. 234.—Ecuador.
hybrida, Thuill. Fl. Par. ed. II. 422 = tripartita.
incisa, G. Don, in Sweet, Hort. Brit. ed. III. 360 = rubifolia.
insipida, Lam. Encyc. i. 416 = Spilanthes insipida.
intermedia, Opiz, ex Nym. Consp. 848 = tripartita.
Jardinii, *Sch. Bip. in Flora*, xxxix. (1856) 360.—Ins. Pacif.
Katzkyi, Sch. Bip. ex Walp. Rep. vi. 168 = bipinnata.
laciniata, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 283, nomen.—Afr. trop.
lantanoides, *A. Gray, in Proc. Am. Acad.* v. (1861) 128.—Ins. Pacif.
Leuconoi, *A. Gray, Syn. Fl. N. Am.* i. ii. 297.—Arizona.
leucantha, Poepp. ex DC. Prod. v. 604 = chilensis.
leucantha, Willd. Sp. Pl. iii. 1719 = pilosa.
leucorrhiza, *DC. Prod.* v. 605.—China.
Leybaldii, *Phil. in Anal. Univ. Chil.* ii. (1865) 336.—Chili.
Lindleyi, Sch. Bip. in Seem. Bot. Voy. Herald, 307 = Cosmos tenuifolius.
linearifolia, Sch. Bip. l. c. = Cosmos linearifolius.
linearioba, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 99. t. 90.—Afr. trop.
longifolia, DC. Prod. v. 597 = heterophylla.
luxurians, Willd. Enum. Hort. Berol. 847 = heterophylla.
Macrachæna, *Griseb. in Goett. Abh.* xix. (1874) 186.—Reg. Argent.

BIDENS:—
macrosperma, Fisch. Hort. Gorenk. (1808) 59; ed. II. (1812) 87 = parviflora.
marginata, Perr. ex DC. Prod. v. 581 = Chrysanthellum indicum.
megapotamica, *Spreng. Syst.* iii. 454.—Am. austr.
Mesersehmidii, Turcz. ex Ledeb. Fl. Ross. ii. 518 = parviflora.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 371.—China.
micrantha, Gaudich. Voy. Freyc. Bot. 464, 85 = Coreopsis micrantha.
minima, Huds. Fl. Angl. 310 = cernua.
mollis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 49.—Ecuador.
monticola, Poepp. & Endl. l. c. 49. t. 255 = rubifolia.
multifida, Desf. Cat. Hort. Par. ed. III. 186 = parviflora.
mutica, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 368 = Coreopsis macrocarpa.
myrrhidifolia, *Tausch, in Flora*, xix. (1836) 894.—Hab.?
nivea, Lam. Sp. Pl. 838 = Melanthera hastata.
nivea, Sw. ex Steud. Nom. ed. II. i. 203 = Melanthera deltoidea.
nodiflora, *Linn. Sp. Pl.* 882.—Ind. or.
Novæ-Caledoniæ, *Forst. ex Schlecht. in Linnæa*, x. (1835–36) 480 (Cf. Glossogyne pinnatifida).—N. Hebrid.
nudiflora, Steud. Nom. ed. II. i. 203 sphalm. = nodiflora.
ocymifolia, Lam. Encyc. i. 416, Illustr. t. 668. f. 3 = Spilanthes alba.
odorata, *Cav. Ic.* i. 9. t. 13.—Mexic.
odorata, Dum.-Cours. Bot. Cult. iv. 241; G. Don, in Loud. Hort. Brit. 286 = grandiflora.
oleracea, Cav. ex Steud. Nom. ed. II. i. 203 = Spilanthes Acmella.
ostruthioides, *Sch. Bip. in Seem. Bot. Voy. Herald*, 308.—Mexic
oxyodonta, *DC. Prod.* v. 600.—Ins. S. Doming.
palescens, *Vis. in Nuov. Saggi Acc. Padov.* v. (1840) 266.—Nubia.
paniculata, *Hook. & Arn. Bot. Beech. Voy.* 66.—Ins. Tahiti.
paradoxa, *D. Don, ex Hook. & Arn. in Hook. Journ. Bot.* iii. (1840) 319.—Chili.
parviflora, *Willd. Enum. Hort. Berol.* 840.—Sibiria; China.
patula, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 405 = rubifolia.
pauciflora, Pohr. Encyc. Suppl. i. 630 = parviflora.
pectinata, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 528, nomen. —Bolivia.
pedunculata, *Gaudich. Voy. Freyc. Bot.* 464.—Ins. Sandvic.
perianthomega, Steud. Nom. ed. II. i. 203, sphalm. = Bignonia perianthomega.
petiolata, Nutt. in Journ. Acad. Philad. vii. (1834) 99 = connata.
pilosa, *Linn. Sp. Pl.* 832.—Ind. occ.; Am. austr.
pinnata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
pinnatifida, Forst. ex DC. Prod. v. 598 = Coreopsis micrantha.
pinnatifida, Buch.-Ham. in Wall. Cat. n. 3190 = Glossogyne pinnatifida.
platycephala, Gerst. Ind. Sem. Haun. (1859) 27; et in Linnæa, xxx. (1859–60) 753 = tripartita.
polycephala, *Sch. Bip. in Flora*, xxxix. (1856) 360.—Ins. Pacif.
portoricensis, *Spreng. in DC. Prod.* v. 601.—Porto Rico.
procera, *D. Don, in Bot. Reg.* t. 684.—Mexic.; Arizona.
pseudamica, *H. B. & K. Nov. Gen. et Sp.* iv. 232.—Mexic.
pulcherrima, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 528, nomen.—Bolivia.
pumila, Steud. Nom. ed. I. 108 = tripartita.
purpurea, DC. Prod. v. 604 = Cosmos purpureus.
quadrangularis, DC. l. c. 600 = pilosa.
quadriradiata, DC. l. c. 595 = chrysanthemoides.
quadriseta, *Hochst. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 392.—Afr. trop.
quinquenudata, Zea, ex Steud. Nom. ed. I. 108 = grandiflora.
radiata, Thuill. Fl. Par. ed. II. 422 = tripartita.
Remillei, Wall. Cat. n. 3191 = Heliopsis platyglossa.
reflexa, *Link, Enum. Hort. Berol.* ii. 305.—Mexic.
regens, D. Don, Prod. Fl. Nep. 180 = tripartita.

BIDENS:—
reptans, G. Don, in Sweet, Hort. Brit. ed. III. 360 = tripartita.
reptans, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos diversifolius.
rhombifolia, *Sch. Bip. in Bonplandia*, iv. (1856) 54.—Peruv.
Riedelii, *Baker, in Mart. Fl. Bras.* vi. III. 246.—Bras.
rigida, Hort. Calc. ex C. B. Clarke, Comp. Ind. 141 = Glossogyne pinnatifida.
riparia, *H. B. & K. Nov. Gen. et Sp.* iv. 236.—Venezuela.
rosea, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos pilosus.
rubifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 237. t. 381. —Ecuador.
rugulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 184.—Peruv.
sambucifolia, *Cav. Ic.* iii. 15. t. 219.—Peruv.; Mexic.
sandwicensis, Less. in Linnæa, vi. (1831) 508 = peduncularis.
Sartorii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 307.—Mexic.
scabioscides, *Orig. in Regel, Gartenfl.* (1861) 408.—Mexic.
scabiosoides, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos scabiosoides.
scandens, Linn. Sp. Pl. 838 = Salmea grandiceps.
scandicina, *H. B. & K. Nov. Gen. et Sp.* iv. 235.—Ecuador.
Schimperi, *Sch. Bip. ex Walp. Rep.* vi. 168.—Abyss.
scorzoneræfolia, *Baker, in Mart. Fl. Bras.* vi. III. 247. —Bras.
Seemannii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 307. —Mexic.
serrata, *Pav. ex DC. Prod.* v. 597.—Peruv.
serrulata, Ds.-el. Cat. Hort. Par. ed. III. (1829) 186 = Coreopsis aurea.
serrulata, *Sch. Bip. in Flora*, xxxix. (1856) 361.—Ins. Pacif.
silvaticus, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 246 = rubifolia.
speciosa, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 126.—Bras.
squarrosa, *H. B. & K. Nov. Gen. et Sp.* iv. 238.—Venezuela.
squarrosa, Less. in Linnæa, v. (1830) 156 = rubifolia.
striata, Schott, ex Sweet, Brit. Flow. Gard. t. 287 = pilosa.
subalternans, DC. Prod. v. 600 = pilosa.
sulphurea, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos sulphureus.
sundaica, Blume, Bijdr. 913 = pilosa.
tenella, *Linn. Fil. Par. Afr.* 17; *Amœn. Acad.* vi. 96. —Afr. austr.
tenuifolia, Lehill. Sert. Austr. Caled. 44. t. 45 = Glossogyne tenuifolia.
tenuifolia, *Tausch, in Flora*, xix. (1836) 895.—Hab.?
tenuiseeta, *A. Gray, Pl. Fendl.* 86.—Am. bor. occ.
tereticaulis, DC. Prod. v. 598 = rubifolia.
tetragona, *DC. l. c.* 596.—Mexic.
tinctoria, Banll. Hist. Fl. viii. 305 = Coreopsis tinctoria.
tridentata, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 183 = Glossogyne tridentata.
trifida, Roxb. Fl. Ind. iii. 411 = tripartita.
trifoliata, Gueldenst. ex Ledeb. Fl. Ross. ii. 516 = tripartita.
tripartita, Bigel. Fl. Bost. ed. II. 294 = connata.
tripartita, Boj. Hort. Maurit. 188 = pilosa.
tripartita, *Linn. Sp. Pl.* 831.—Europ.; Sibir.
tripliner via, *H. B. & K. Nov. Gen. et Sp.* iv. 231.—Mexic.
tuberosa, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 528, nomen.—Bolivia.
valladolidensis, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos crithmifolius.
vulparadisæa, Colla, in Mem. Acc. Torin. xxviii. (1825) 12. t. 24 = chilensis.
verlaskia, Baill. Hist. Pl. viii. 305 = Dahlia variabilis.
vernan, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 405 = gravcolens.
verticillata, Baill. Hist. Pl. viii. 305 = Coreopsis verticillata.
verticillata, *Linn. Sp. Pl.* 833.—Mexic.
Wallichii, DC. Prod. v. 598 = bipinnata.
Warszewiczianus, *Regel, in Flora*, xxxii. (1849) 188.—Guatemala.

BIDWILLIA, Herb. ex Lindl. Veg. Kingd. 806 (1847) nomen. (Liliac.). Benth. & Hook. f. Gen. iii. 708.

BIEBERSTEINIA, Steph. ex [Fisch.] Hort. Gorenk. ed. I. 88 (1808); et in Mém. Soc. Nat. Mosc. i. (1811) 89. t. 9. *GERANIACEAE*, Benth. & Hook. f. i. 271. DAYENIA, Michx. ex Jaub. & Spach, in Ann. Sc. Nat. Sér. III. vi. (1846) 187. — *ambigua*, Jaub. & Spach, in Ann. Sc. Nat. Sér. III. vi. (1846) 141 = multifida. — *Auckovi*, Jaub. & Spach, Illustr. ii. 110 = multifida. — *brachypetala*, Jaub. & Spach, l. c. 113. t. 192 = multifida. — Emodii, *Jaub. & Spach, l. c.* 109.—Tibet. occ. leiosepala. *Jaub. & Spach, l. c.* 114. t. 198.—Persia. — multifida, *DC. Prod.* i. 708.—Syria; Persia; Soongaria. — *odora*, Royle, Illustr. Bot. Himal. i. 168. t. 80 = Emodii. — odora, *Steph. ex* [*Fisch.*] *Hort. Gorenk.* ed. I. (1808) 88; *et ex Mém. Soc. Nat. Mosc.* i. (1811) 89.—Persia. — Orphanidis, *Boiss. Diagn.* Ser. II. i. 118.—Graecia; Cilicia. — *Ruckeri*, Walp. Ann. i. 153 = multifida.

BIENERIA, Reichb. f. in Bot. Zeit. xi. (1853) 3. t. 1 = **Chloraea**, Lindl. (Orchid.). — *boliviana*, Reichb. f. l. c.—Bolivia.

BIENNERTIA, Bunge, in Act. Hort. Petrop. vi. (1879) ii. 425; et in Boiss. Fl. Orient. iv. 945 (1879). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 68. cyclopters, *Bunge, ii. cc.*—Persia.

BIFOLIUM, G. Gaertn. Mey. & Scherb. Fl. Wett. i. 178 (1799) = **Maianthemum**, Wiggers (Liliac.). — *cordatum*, G. Gaertn. Mey. & Scherb. l. c. 209 = M. Bifolium.

BIFORA, Hoffm. Umb. Gen. ed. II. 191 (1816). *UMBELLIFERAE*, Benth. & Hook. f. i. 927. ATHAMA, DC. Prod. iv. 250 (1830). BIFORIS, Spreng. Anleit. ii. 685 (1817); et in Roem. & Schult. Syst. vi. 448 (1820). CORION, Link & Hoffngg. Fl. Port. ii. 457 (1820). — americana, *Benth. & Hook. f. ex S. Wats. Bibl. Index*, 415.—Am. bor. — *benghalensis*, Wall. [Cat. n. 888] ex DC. Prod. iv. 140 = Oenanthe benghalensis. — dicocca, *Hoffm. Umb. Gen.* ed. II. 192.—Europ.; Oriens. — *flexuosa*, Bieb. Fl. Taur. Cauc. iii. 324 = dicocca. — *glauca*, Wall. Cat. n. 887 = Oenanthe benghalensis. — Louveiri, *Kostel. Allg. Med.-Pharm. Fl.* iv. 1188.—Cochinch. — radiana, *Bieb. Fl. Taur. Cauc. Suppl.* 283.—Europ. austr.; Oriens. — *testiculata*, DC. Prod. iv. 249 = dicocca. — testiculata, *Roth, Enum. Pl. Phan. Germ.* i. 388.— Reg. Mediterr.

BIFORIS, Spreng. Anleit. ii. 685 (1817); et in Roem. & Schult. Syst. vi. 448 (1820) = **Bifora**, Hoffm. (Umbellif.).

BIFRENARIA, Lindl. Gen. et Sp. Orchid. 152 (1838). *ORCHIDEAE*, Benth. & Hook. f. Gen. iii. 546. ADIPE, Rafin. Fl. Tellur. ii. 101 (1836). STENOCORYNE, Lindl. Bot. Reg. xxix. (1843) Misc. 68. — albiflora, *Rodrig. Orch. Nov.* ii. 212, nomen.—Bras. — atropurpurea, *Lindl. Gen. et Sp. Orch.* 153.—Bras. — aurantiaca, *Lindl. Bot. Reg.* (1836) t. 1875.—Ind. occ. — aurea, *Rodrig. Orch. Nov.* ii. 212.—Bras. — aureofulva, *Lindl. Bot. Reg.* (1843) *Misc.* 69.—Bras. — *bella*, Lem. Jard. Fleur. iii. (1853–53) t. 295 = Coelia bella. — bicornata, *Reichb. f. in Hamb. Gartenz.* xix. (1863) 19.—Bras.

BIFRENARIA;—
— chloroleuca, *Rodrig. Orch. Nov.* ii. 212, nomen.—Bras. — calcarata, *Rodrig. l. c.* 213.—Bras. — clavigera, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 296.—Bras. — *fragrans*, Rodrig. Orch. Nov. ii. 213 = inodora. — *Hadwenii*, Lindl. in Paxt. Fl. Gard. ii. 67 (1851–52) = Scuticaria Hadwenii. — Harrisoniae, *Reichb. f. in Bonplandia*, iii. (1855) 217. —Hab.! — inodora, *Lindl. Bot. Reg.* xxix. (1843) *Misc.* 48.—Bras. — leucopetala, *Rodrig. Orch. Nov.* ii. 212, nomen.—Bras. — leucorrhoda, *Reichb. f. in Otto, Hamb. Gartenz.* xv. (1859) 54.—Hab.! — longicornis, *Lindl. Bot. Reg.* (1838) *Misc.* 93.— Guiana. — melanopoda, *Klotzsch, in Otto & Dieter. Allg. Gartenz.* xxiii. (1855) 105.—Bras. — mellicolor, *Reichb. f. in Gard. Chron.* (1878) i. 622.—Bras.? — parvula, *Reichb. f. in Walp. Ann.* vi. 547.—Bras. — racemosa, *Lindl. Bot. Reg.* (1843) *Misc.* 52.—Bras. — sabolosa, *Rodrig. Orch. Nov.* i. 111.—Bras. — vinosa, *Rodrig. l. c.* ii. 213, nomen.—Bras. — vitellina, *Lindl. Bot. Reg.* (1843) *Misc.* 59.—Bras. — tyrianthina, *Reichb. f. Xenia*, i. 61, 228.—Bras. — Wagneri, *Reichb. f. in Bonplandia*, iii. (1854) 17.— Venezuela.

BIGAMEA, Koen. ex Endl. Gen. 1188 (1840) = **Ancistrocladus**, Wall. (Dipterocarp.).

BIGELONIA, Rafin. in Journ. Phys. lxxxix. (1819) 289. sphaim. = BIGELOWIA, Rafin. = **Axenaria**, Rupp. (Caryoph.). — *elongata*, Rafin. l. c. = A. lanuginosa. — *montana*, Rafin. l. c. = A. montana.

BIGELOVIA, Spreng Syst. i. 866, 404 (1827) = **Spermacoce**, Linn. (Rubiac.). — *alata*, Reichb. ex Steud. Nom. ed. II. i. 208 = S. latifolia, longifolia. — *angustifolia*, Span. in Linnaea, xv. (1841) 320 = S. stricta I. — *connatuata*, Spreng. Syst. i. 404 = Diodia maritima. — *cornifolia*, Heynh. Nom. i. 121 = S. capitata. — *corymbosa*, Bartl. ex DC. Prod. iv. 550 = S. corymbosa. — *discolor*, Bartl. ex DC. l. c. (= *Borreria discolor*). — *distans*, Hook. & Arn. Bot. Beech. Voy. 295 = S. distans. — *glata*, Bartl. ex DC. Prod. iv. 544 = S. latifolia, longifolia. — *eryngioides*, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 360 = S. diffusa. — *gracilis*, Spreng. Syst. i. 404 = S. oxymoides. — *gracilis*, Miq. in Verh. Nederl. Inst. Reeks III. v. (1859) 18 = Malaya. — *Klotzschii*, Hook. f. Fl. Ind. 437 = S. stricta. — *laevicaulis*, Miq. Fl. Ind. Bat. ii. 335 = S. oxymoides. — *laniocarpa*, Wight & Arn. Prod. 437 = S. stricta. — *linearis*, Spreng. Syst. i. 405 = Diodia hyssopifolia. — *linifolia*, Spreng. l. c. 404 = S. laevis. — *mitracarpoides*, Miq. Fl. Ind. Bat. ii. 334 = S. stricta. — *myrianthea*, Miq. l. c. = S. stricta. — *oxymoides*, Miq. l. c. 335 = S. oxymoides. — *parviflora*, Sieber, ex DC. Prod. iv. 543 = S. oxymoides. — *parviflora*, Spreng. Syst. i. 405 = S. oxymoides. — *pavifolioides*, Bartl. ex DC. Prod. iv. 547 = S. tenella. — *pumila*, Span. in Linnaea, xv. (1841) 320 = S. pumila. — *pusilla*, Spreng. Syst. i. 406 = S. pusilla. — *Roxburghiana*, Wight & Arn. Prod. 437 = S. stricta. — *Roxburghiana*, Wight, ' Cat. n. 1350 ' = S. oxymoides. — *setidens*, Miq. Fl. Ind. Bat. ii. 336 = S. stricta. — *socialis*, Span. in Linnaea, xv. (1841) 320.—Ins. Timor. — *striata*, Span. l. c. 321 = S. stricta. — *stricta*, Blume, Bijdr. 946 = S. stricta. — *suaveolens*, Spreng. Syst. i. 404 = S. tenella. — *tenera*, Heynh. Nom. i. 122 = S. tenera. — *tenuis*, Heynh. l. c. = S. tenuis. — *umbellatae*, Spreng. Syst. i. 405.—Bras. — *vaginata*, Spreng. l. c. = S. laevis. — *verticillata*, Spreng. l. c. 404 = S. verticillata. — *Zollingeriana*, Miq. Fl. Ind. Bat. ii. 336.—Malaya.

BIGELOWIA, Spreng. Neue Entdeck. ii. 150 (1821) = **Stamydia**, Link. — *brasiliensis*, Spreng. l. c. = S. Selloi.

BIGELOVIA, Sm. in Rees, Cyclop. xxxix. (1819) = **Forestiera**, Poir. (Oleac.). — *acuminata*, Sm. l. c. = F. acuminata. — *cassinioides*, Sm. l. c. = F. cassinioides. — *ligustrina*, Sm. l. c. = F. ligustrina. — *porulosa*, Sm. l. c. = F. porulosa.

BIGELOWIA, DC. ex Ging. in DC. Prod. i. 390 (1824) = **Holostylis**, H. B. & K. (Violar.).

BIGELOWIA, DC. Prod. iv. 540 (1830) = BIGELOVIA, Spreng. = **Spermacoce**, Linn. (Rubiac.).

BIGELOWIA, DC. Prod. v. 329 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 292. CHRYSOTHAMNUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 393. ISOCOMA, Nutt. l. c. 390 (1841). LINOSYRIS, Torr. & Gray, Fl. N. Am. ii. 393 (1841) acradenia, *Greene, in Bull. Torrey Club*, x. (1883) 196.—Am. bor. occ. — albida, *M. E. Jones, ex A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 209.—Am. bor. occ. — arborescens, *A. Gray, l. c.* viii. (1873) 640.—Calif. — Bigelovii, *A. Gray, l. c.* xvii. (1881–82) 209.—N Mexic.; Colorado. — Bolanderi, *A. Gray, l. c.* viii. (1873) 641.—Calif. — brachylepis, *A. Gray, in Bot. Calif.* i. 514.—Calif. — *carnosa*, Benth. & Hook. f. Gen. ii. 293, 255 = Aster carnosus. — ceramicosa, *A. Gray, in Proc. Am. Acad.* viii. (1873) 648.—Calif. — Cooperi, *A. Gray, l. c.* 640.—Calif. — coronopifolia, *A. Gray, l. c.* 638.—Texas. — depressa, *A. Gray, l. c.* 643.—Am. bor. occ. — diffusa, *A. Gray, l. c.* 640.—Calif. — *Douglasii*, A. Gray, l. c. 644 = viscidiflora. — dracunculoides, *DC. Prod.* v. 329.—Am. bor. — Drummondii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 639.—Texas. — Engelmanni, *A. Gray, l. c.* xi. (1876) 75.—Am. bor. occ. — filaginea, *A. Gray, l. c.* viii. (1873) 638.—Ecuador. — furfuracea, *Greene, in Bull. Calif. Acad.* i. n. 3 (1885) 87.—Calif. — *graveolens*, A. Gray, in Proc. Am. Acad. viii. (1873) 644 = dracunculoides. — Greenei, *A. Gray, l. c.* xi. (1876) 75.—Colorado. — Hartwegi, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 112.—Mexic. — Howardi, *A. Gray, in Proc. Am. Acad.* viii. (1873) 641.—Colorado. — hypoleuca, *A. Gray, l. c.* 638.—Ecuador. — *intricata*, A. Gray, l. c. xvii. (1881–82) 208 = Aster carnosus. — junces, *Greene, in Coult. Bot. Gazette*, vi. (1881) 184. —Arizona. — *lanceolata*, A. Gray, in Proc. Am. Acad. viii. (1873) 639 = viscidiflora. — leiosperma, *A. Gray, Syn. Fl. N. Am.* i. ii. 139.— Am. bor. occ. — Mandonii, *Benth. & Hook. f. Gen.* ii. 293, 255.— Bolivia! — Menziesii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 638.—Am. bor. occ. — *microcrinota*, DC. Prod. v. 329 = dracunculoides. — mohavensis, *Greene, ex A. Gray, Syn. Fl. N. Am.* i. ii. 136.—Am. bor. occ. — nevadensis, *A. Gray, l. c.* 136.—Calif. — nudata, *DC. Coll. Mém.* ix. t. 8.—Am. bor. or. — oppositifolia, *A. Gray, in Proc. Am. Acad.* xv. (1880) 92.—Mexic. — paniculata, *A. Gray, l. c.* viii. (1873) 644.—Am. bor. occ. — Parishii, *Greene, in Bull. Torrey Club*, ix. (1882) 66. —Calif. — Parryi, *A. Gray, in Proc. Am. Acad.* viii. (1873) 643. —Colorado. — pluriflora, *A. Gray, l. c.* 639.—Colorado! — pulchella, *A. Gray, l. c.* 642.—Texas; N. Mexic. — ramulosa, *Benth. & Hook. f. Gen.* ii. 293, 255.—Am. bor. — *rupestris*, Greene, in Coult. Bot. Gazette, vi. (1881) 186 = Aplopappus cuneatus. — spathulata, A. Gray, in Proc. Am. Acad. xi. (1876) 74 = Aplopappus cuneatus. — tenuifolia, *A. Gray, l. c.* viii. (1873) 644.—Calif.; Arizona. — *triderantata*, Greene, in Bull. Torrey Club, x. (1883) 126 = vaseyi.

BIGELOWIA:—
umbigulata, DC. Prod. v. 329 = Solidago neglecta.
Vaseyi, *A. Gray*, *in Proc. Am. Acad.* xii. (1877) 58.—Colorado.
venets, *A. Gray*, *l. c.* viii. (1873) 688.—Calif.; Mexic.
virgata, DC. Prod. v. 329 = nudata.
viscidiflora, *DC. l. c.* vii. 279.—Am. bor.
Wrightii, *A. Gray*, *in Proc. Am. Acad.* viii. (1873) 689.—Am. bor. occ.

HIGGINA, Rafin. Alsog. Am. 14 (1838) = **Salix**, Tourn.
angustata, Rafin. l. c. = S. cordata ?
argentea, Rafin. l. c. = S. argentea.
conifera, Rafin. l. c. = S. humilis.

BIGLANDULARIA, Karst. in Linnaea, xxviii. (1856) 416. *GENTIANEAE.*
azurea, *Karst. l. c.* 417.—N. Granat.

BIGLANDULARIA, Seem. in Gard. Chron. (1868) 738 = **Sinningia**, Nees (Generac.).
conspicua, Seem. l. c.—Bras.

BIGNONIA, [Tourn.] Linn. Syst. ed. I (1735) *BIGNONIACEAE*, Benth. & Hook. f. ii. 1038.
AMISOSTICHUS, Bur. Monog. Bign. 48. t. 6 (1864).
CLEMATITARIA, Bur. l. c. 44 (1864).
CREMASTUS, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 187.
CUPULIBIA, Rafin. Fl. Tellur. ii. 57 (1836).
CYDISTA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 191.
DOXANTHA, Miers, l. c. 188 (1863).
ENDOLOMA, Rafin. Sylva Tellur. 79 (1838).
HIERANTHES, Rafin. l. c. (1838).
ODISCA, Rafin. l. c. 80 (1838).
PANTERPA, Miers, in Proc. Roy Hort. Soc. iii. (1863) 195.
PARAGONIA, Bur. in Bull. Soc. Bot. Fr. xix. (1872) 17.
PENTELESIA, Rafin. Sylva Tellur. 146 (1838).
PETASTOMA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 193.
PHAEDRANTHUS, Miers, l. c. 182 (1863).
PLEONOTOMA, Miers, l. c. 184 (1863).
PONGELIA, Rafin. Sylva Tellur. 78 (1838).
PROTERPIA, Rafin. l. c. 80 (1838).
PYROSTEGIA, Presl, in Abh. Boehm. Ges. V. iii (1845) 528 (Bot. Bemerk 98).
SERERRA, Rafin. Sylva Tellur. 107 (1838).
STIZOPHYLLUM, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 197.
TEMNOCYDIA, Mart. ex DC. Prod. ix. 143 (1845).
acutiflora, Mart. ex DC. Prod. ix. 181 = Lundia nitidula.
acutissima, *Cham. in Linnaea*, vii. (1832) 691.—Bras.
acutistipula, *Schlechl. l. c.* xxvi. (1858–59) 375.—Mexic.
adenophylla, Wall. Cat. n. 6502 = Heterophragma adenophyllum.
adhaerodactylis, *Jacq. Fragm.* 61.—Venezuela.
aequinoctialis, *Linn. Sp. Pl.* 628.—Am. trop.
aequinoctialis, Sieber, ex DC. Prod. ix. 146 = B. Unguis-cati.
aesculifolia, H. B. & K. Nov. Gen. et Sp. iii. 140 = Tabebuia aesculifolia.
Aesculus, Desf. Cat. Hort. Par. ed. III. 297 = Tecoma Aesculus.
affinis, *G. Don, Gen. Syst.* iv. 217.—Bras.
africana, *Lam. Encyc.* i. 424.—Afr. trop.
Agnus-castus, Cham. in Linnaea, vii. (1832) 678 = Arrabidaea Agnus-castus.
alata, Pav. ex DC. Prod. ix. 224 = Tecoma Gaarume.
alba, Aubl. Fl. Gui. ii. 658 = Spathodea bracteosa.
alba, Blume, in Verh. Bat. Gen. ix. (1825) 195 = Aeschynanthus purpurascens.
alba, Stend. Nom. ed. II. i. 304 = Aeschynanthus angustifolia.
albiflora, *Salzm. ex DC. Prod.* ix. 167.—Bras.
allamandaeflora, *Benth. ex Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 186, nomen.—Bras.
allacoa, *Lam. Encyc.* i. 181.—Guiana.
amara, Lond. Hort. Brit. 483 ; ex G. Don, Gen. Syst. iv. 223 = Stereospermum hypostictum.
amarae, Wall. Cat. n. 6572 ; ex Fl. As. Rar. ii. 76. t. 158 = Stereospermum hypostictum.
anastomosans, *DC. Prod.* ix. 144.—Madag.
Andrieuxii, *DC. Prod.* ix. 156.—Mexic.
angrensis, *Vell. Fl. Flum.* 246 ; vi. t. 28.—Bras.
angustifolia, Blume, Cat. Hort. Buitenz. 89 = Aeschynanthus angustifolia.
anisophlebia, Mart. Reise Bras. i. 283 = Cybistax antisyphilitica.
apurensis, *H. B. & K. Nov. Gen. et Sp.* iii. 138 — Venezuela.
aquatilis, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 780 = Tabebuia fluviatilis.
araliacea, Chamn. in Linnaea, vii. (1832) 688 = Tecoma araliacea.
argyrea, *André, in Rev. Hortic.* (1865) 251.—Hab. ?
argyreo-violascens, *Hort. ex Lem. Illustr. Hortic.* (1866) t. 469.—Hab. ?
Arrabidae, *Steud. Nom.* ed. II. i. 204.—Bras.
Arthersoni, Mart. Herb. Fl. Bras. 287 = Distictis Arthersoni.
articulata, Desf. in Poir. Encyc. Suppl. i. 686 = Phyllarthron Noronhianum.
arvensis, Vell. Fl. Flum. 250 ; vi. t. 40 = Anemopaegma mirandum.
atrovirens, DC. Rev. Bign. ex Prod. ix. 171 = Tecoma atrovirens.
atrovirens, Roth, Nov. Sp. 284 = Dolichandrone falcata.
aurea, Manso, Enum. 40 = Tecoma aurea
auriculigera, Mart. ex DC. Prod. ix. 194 = Pithecoctenium stipulaceum.
australis, Ait. Hort. Kew. ed. II. iv. 84 = Tecoma australis.
Ausdarachta, Kon. & Sims, Ann. Bot. i. (1805) 278 = Millingtonia hortensis.
Baclei, DC. Rev. Bign. ex Prod. ix. 171 = Pithecoctenium cynanchoides.
Balbisiana, *DC. Prod.* ix. 153.—Bras.
Banalbana, Callery, ex Bur. in Baill. Adansonia, ii. (1861–62) 195 = Stereospermum Banalbanai.
barbata, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 795 = Tecoma barbata.
Bellus, *Sellow, ex Steud. Nom.* ed. II. i. 204.—Bras.
bibracteata, Bert. ex DC. Prod. ix 240 = Bravaisia floribunda.
Bifwillana, *Regel, Cat. Pl. Hort. Askachev.* 19.—Hab. ?
bijuga, *Vahl, Symb. Bot.* ii. 68.—Madag.
binata, Thunb. Dias. Pl. Bras. iii. 35. n. 27 = Anemopaegma nocnosum.
bipinnata, Salzm. ex DC. Prod. ix. 230 = Jacaranda nitida.
Blanchetii, *DC. l. c.* 164.—Bras.
Bojeri, *DC. l. c.* 166.—Madag.
botryoides, Cham. in Linnaea, vii. (1832) 100 = Pithecoctenium botryoides.
brachycalyx, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1085.—Guiana.
brachypoda, *DC. Prod.* ix. 145.—Bras.
bracteata, Cham. in Linnaea, vii. (1832) 692 = Adenocalymna bracteatum.
bracteolata, *DC. Prod.* ix. 157.—Bras.
bracteosa, *DC. l. c.* 165.—Madag.
brasiliana, Lam. Encyc. i. 426 = Jacaranda brasiliana.
buccinatoria, *Meisel, ex DC. Prod.* ix. 195.—Mexic.
caerulea, Linn. Sp. Pl. 626 = Jacaranda caroliniana.
callistegioides, Cham. in Linnaea, vii. (1832) 712 = Cuspidaria callistegioides.
calycina, *DC. Prod.* ix. 160.—Bras.
campanulata, Cham. in Linnaea, vii. (1832) 711 = Cuspidaria campanulata.
candicans, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Arrabidaea candicans.
capensis, Thunb. Prod. Pl. Cap. 105 = Tecoma capensis.
capreolata, *Linn. Sp. Pl.* 624.—Am. bor.
carichensis, *H. B. & K. Nov. Gen. et Sp.* iii. 137.—Venezuela.
Carobe, Vell. Fl. Flum. 250 ; vi. t. 43 = Jacaranda Carobe.
Caroltonea, *Lindl. Bot. Reg.* (1842) sub t. 45.—Hab. ?
caryophyllea, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 293.—Porto Rico.
cassinoides, Lam. Encyc. i. 418 = Tabebuia cassinoides.
castanaefolia, *DC. Prod.* ix. 145.—Peruv.
Catalpa, Linn. Sp. Pl. 622 = Catalpa bignonioides.
Catharinae, DC. Rev. Bign. ex Prod. ix. 172 = Pithecoctenium Catharinae.
caudata, DC. Prod. ix. 166 = Stereospermum caudatum.
caudata, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 269 = Stereospermum chelonoides.
cauliflora, Sieber, ex DC. Prod. ix. 141 = Colea floribunda.
Chamberlaynii, *Sims, Bot. Mag.* t. 2148.—Bras.
chelonoides, Linn. f. Suppl. 289 = Stereospermum chelonoides.
Cherere, Lindl. Bot. Reg. t. 1301 = bucinatoria.
Chica, *Humb. & Bonpl. Pl. Aequin.* i. 107. t. 31.—Venezuela.
chinensis, Lam. Encyc. i. 428 = Tecoma grandiflora.
chondrogona, *Miq. in Linnaea*, xviii. (1844) 611.—Guiana.
chrysantha, Jacq. Hort. Schoenb. ii. 45 = Tecoma chrysantha.
chrysoleuca, *H. B. & K. Nov. Gen. et Sp.* iii. 134.—Venezuela.
Cicutaria, Koen. ex Mart. in Denkschr. Akad. Muench. vi. (1820) 153 = Millingtonia hortensis.
ciliata, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 287.—Bras.
cinerea, DC. Rev. Bign. ex Prod. ix. 172 = Pithecoctenium cinereum.
cinnamomea, *DC. Prod.* ix. 164.—Bras.
citrifolia, *Vitm. Summa Veg.* iii. 511.—Hab. ?
Clematis, *H. B. & K. Nov. Gen. et Sp.* iii. 141.—Venezuela.
coccinea, Steud. Nom. ed. I. 109 = Tecoma radicans.
coccinea, *Vell. Fl. Flum.* 250 ; vi. t. 42.—Bras.
cognata, *Cham. in Linnaea*, vii. (1832) 708.—Bras.
Colei, Vell. Fl. Flum. 248 ; vi. t. 19 = Spathodea Colei.
Colei, Buch.-Ham. ex Wall. Cat. sub n. 6501 = Stereospermum chelonoides.
Colei, Boj. ex Hook. Bot. Mag. t. 2817 = Colea mauritiana.
colorata, Poir. Encyc. Suppl. i. 684 = Arrabidaea candicans.
comosa, Cham. in Linnaea, vii. (1832) 698 = Adenocalymna comosum.
comosa, *Roxb. Hort. Beng.* [95] ; *Fl. Ind.* iii. 108.—Ins. Molucc.
compressa, Lam. Encyc. i. 424 = Colea decora.
conjugata, Vell. Fl. Flum. 243 ; vi. t. 18 = Arrabidaea conjugata.
conspicua, Rich. ex DC. Prod. ix. 231 = Tecoma conspicua.
convoluta, Vell. Fl. Flum. 248 ; vi. t. 31 = Cuspidaria pubercula.
Copaia, Aubl. Pl. Gui. ii. 650. t. 262 = Jacaranda Copaia.
corallina, *Jacq. Fragm.* 87.—Venezuela.
corchoroides, Cham. in Linnaea, vii. (1832) 706 = Arrabidaea corchoroides.
cordata, Vell. Fl. Flum. 247 ; vi. t. 29 = Lundia cordata.
cordifolia, *DC. Prod.* ix. 162.—Bras.
corsaca, Sellow, ex Steud Nom. ed. II. i. 204, 205 = Selloi.
corymbifera, *Vahl, Ecl. Am.* ii. 45. t. 17.—Am. austr.
craterophora, *DC. Prod.* ix. 147.—Bras.
crenata, *Ladd. ex Loud. Hort. Brit.* 242.—Ind. or.
crispa, Buch.-Ham. ex Roxb. Hort. Beng. 47 ; Fl. Ind. iii. 108 = Dolichandrone crispa.
crucifera, Bert. ex DC. Prod. ix. 218 = Tabebuia rufinervis.
cruenta, Linn. Sp. Pl. 624 = capreolata.
cujabana, *DC. Prod.* ix. 164.—Bras.
cunelifolia, *DC. l. c.* 157.—Bahia.
cupreo, *Cham. in Linnaea*, vii. (1832) 666.—Bras.
cupulata, *Spitzg. in Hoen. & De Vriese, Tijdschr.* ix. (1843) 6.—Bras.
curialis, Vell. Fl. Flum. 258 ; vi. t. 55 = Jacaranda curialis.
cuspidata, Boj. ex DC. Prod. ix. 248 = Perfiblema cuspidata.
Cymbalum, *Cham. in Linnaea*, vii. (1832) 716.—Bras.
cymosa, *Vell. Fl. Flum.* 249 ; vi. t. 36.—Bras.
dasyantha, DC. Rev. Bign. ex A. DC. Prod. ix. 172 (= *Pachyptera dasyantha*).—Bras.
decipiens, *Ladd. ex Loud. Hort. Brit.* 249.—Am. austr.
decomposita, *Miq. in Linnaea*, xviii. (1844) 260.—Guiana.
densiflora, Mart. ex DC. Prod. ix. 181 = Lundia densiflora.
dentata, *DC. l. c.* 163.—Bras.

BIGNONIA :—
dichotoma, *Jacq. Enum. Pl. Carib.* 25.—Am. austr.
dichotoma, *Vell. Fl. Flum.* 248 : vi. t. 32.—Bras.
difflexilis, *Cham. in Linnaea*, vii. (1832) 714.—Bras.
digitalis, Vell. Fl. Flum. 251 ; vi. t. 47 = Zeyheria montana.
digitata, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 783 = Tecoma Meyeriana.
discolor, Boj. ex DC. Prod. ix. 165 = Bojeri.
discolor, *R. Br. in Salt, Voy. Abyss. App* 64, nomen. —Abyss.
discolor, Rich. Sert. Astrol. p. xxix = moluccana.
diversifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 183.—Mexic.
dolichoides, *Cham. in Linnaea*, vii. (1832) 696.—Bras.
Dombeyana, DC. Rev. Bign. ex A. DC. Prod. ix. 172 = Delostoma nervosum ?
Donianа, *DC. Prod.* ix. 145.—Bras.
dubia, *Linn. Pl. Surin.* n. 77 ; *Amoen. Acad.* viii. 258. —Guiana.
echinata, Aubl. Pl. Gui. ii. 648 = Pithecoctenium Aubletii
echinata, Jacq. Enum. Pl. Carib. 25 = Pithecoctenium muricatum.
ehretioides, *Cham. in Linnaea*, vii.(1832) 704.—Bras.
elegans, *Cham. l. c.* 702.—Bras.
elegans, Vell. Fl. Flum. 247 ; vi. t. 37 = exserta.
elliptica, Cham. in Linnaea, vii. (1832) 696 = Tecoma elliptica.
elliptica, *Thumb. Diss. Pl. Bras* iii. 84.—Bras.
elliptica, Vell. l. c. 250 ; vi. t. 44 = Jacaranda elliptica.
elongata, *Vahl. Ecl.* ii. 45. t. 16.—Am. austr.
erubescens, *DC. Prod.* ix. 157.—Bras.
euphorioides, Boj. Hort. Maurit. 219 = Stereospermum euphorioides.
exoleta, *Vell. Fl. Flum.* 248 ; vi. t. 30.—Bras.
exserta, *DC. Prod.* ix. 158.—Bras.
extensa, *Soland. ex Mors, in Proc. Roy. Hort. Soc.* iii. (1863) 196, nomen.—Bras.
fagoides, *Cham. in Linnaea*, vii. (1832) 680.—Bras.
falcata, Koen. ex DC. Prod. ix. 206 = Dolichandrone falcata.
falcata, *Vell. Fl. Flum.* 245 ; vi. t. 17.—Bras.
fallax, *Cham. in Linnaea*, vii. (1832) 717.—Bras.
farinosa, Salisb. ex DC. Prod. ix. 185 = Arrabidaea viraecea.
fasciculata, Vell. Fl. Flum. 247 ; vi. t. 25 = Arrabidaea fasciculata.
Ferdinandi, *Weiw. Apont.* 584.—Afr. trop.
ferruginea, *Tuerdic, ex Mors, in Proc. Roy. Hort. Soc.* iii. (1863) 190, nomen.—Reg. Argent.
ficifolia, Steud. Nom. ed. ii. i. 304, sphal. = ilicifolia.
ilicifolia, Anders. in Trans. Soc. Arts. xxv. (1807) 200 = jacaranda filicifolia.
filiformis, A. Cunn. ex DC. Prod. ix. 209 = Spathodea filiformis.
fimbriata, Wall. Cat. n. 6500 = Stereospermum fimbriatum.
flava, *DC. Prod.* ix. 168.—Bras.
flavescens, Vell. Fl. Flum. 252 ; vi. t. 51 = Tecoma flavescens.
flavida, *DC. Prod.* ix. 168.—Bras.
floribunda, G. Don, Gen. Syst. iv. 216 = Doniana.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iii. 134.— Mexic.
florida, Salisb. Prod. 106 = Tecoma radicans.
fluminensis, *Vell. Fl. Flum.* 246 ; vi. t. 34.—Bras.
fluviatilis, Aubl. Pl. Gui. ii. 655 = Tabebuia fluviatilis.
fluviatilis, H. B. & K. Nov. Gen. et Sp. ii. 189 = Tecoma barbata.
fluviatilis, G. F. W. Mey. Prim. Fl. Esseq. 219 = Tecoma Meyeriana.
Focicana, *Miq. in Linnaea*, xviii. (1844) 609.— Guiana.
fraxinea, *Desf. Cat. Hort. Par.* ed. III. 396.—Hab. ?
fraxinifolia, Spreng. Syst. ii. 833 = Spathodea fraxinifolia.
fraxinoides, *Perr. in Mém. Soc. Linn. Par.* iii. (1824) 103.—Java.
frutescens, Mill. Gard. Dict. ed. VIII. n. 3 = Tecoma stans.
fulva, Cav. Ic. iii. 58 = Tecoma fulva.
Gaudichaudi, DC. Rev. Bign. ex Prod. ix. 172 = Landia pubescens.
Ghorta, Buch.-Ham. ex Wall. Cat. n. 6510 = Stereospermum Ghorta.

BIGNONIA :—
Glesbreghtii, *Heller, ex Peyr. in Linnaea*, xxx. (1859–60) 45.—Mexic.
gigantea, *Norenha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 8.—Malaya.
glabertiana, *Cham. in Linnaea*, vii. (1832) 677.— Bras.
glabrata, *H. B. & K. Nov. Gen. et Sp.* iii. 135.— Am. austr.
glandulosa, *Mart. ex DC. Prod.* ix. 149.—Bras.
glandulosa, Schum. Beskr. Guin. Pl. 274 = Spathodea adenantha.
glauca, Decne. in Jacquem. Voy. Bot. 187. t. 142 = Tecoma undulata.
glauca, *Wall. Cat.* n. 6506.—Burma.
glutinosa, *DC. Prod.* ix. 163.—Bras.
guaphalantha, *A. Rich. Fl. Cub. Fanerog.* ii. 105.— Cuba.
gracilis, Lodd. Bot. Cab. t. 1705 = B. Unguis-cati.
grandiflora, Spreng. Syst. ii. 830 = grandifolia.
grandiflora, Thunb. Fl. Jap. 253 = Tecoma grandiflora.
grandifolia, *Jacq. Hort. Schoenb.* iii. 19. t. 287.—Am. austr.
grandifolia, Vell. Fl. Flum. 247 ; vi. t. 28 = Adenocalymna grandifolium et subsessilifolium.
gratissima, Koen. ex Wall. Cat. sub n. 6499 ; et DC. Prod. ix. 211 = Stereospermum suaveolens.
Guaraunci, Domb. ex A. DC. l. c. 235 = Tecoma Guaraunci.
guayaquilensis, *DC. l. c.* 155.—Ecuador.
harmandiana, Bert. ex Spreng. Syst. ii. 832 = Tabebuia haemantha.
hebantha, Mart. ex DC. Prod. ix. 181 = Landia hebantha.
heptaphylla, Vell. Fl. Flum. 251 ; vi. t. 46 = Tecoma heptaphylla.
heterophylla, Willd. Sp. Pl. iii. 298 = B. Kerere.
heterophylla, DC. Rev. Bign. ex Prod. ix. 173 = Tecoma heteropoda.
heterotricha, DC. l. c. = Tabebuia heterotricha.
hexagona, DC. l. c. = Pithecoctenium hexagonum.
hibiscifolia, *Cham. in Linnaea*, vii. (1832) 705.— Bras.
hirsuta, Lam. Encyc. i. 222 = Tecoma hirsuta.
hirta, *Vell. Fl. Flum.* 249 ; vi. t. 35.—Bras.
hispida, *DC. Prod.* ix. 152.—Bras.
hondensis, *H. B. & K. Nov. Gen. et Sp.* iii. 135.—N. Granat.
Hostmanni, *E. Mey. in Nov. Act. Nat. Cur.* xii. (1825) ii. 779.—Guiana.
Hymenaea, *DC. Prod.* ix. 158.—Bras.
igasa, *Vell. Fl. Flum.* 244 ; iv. t. 15.—Bras.
ilicifolia, Pers. Syn. ii. 170 = Tabebuia ilicifolia.
inaequalis, *DC. Prod.* ix. 170.—Guiana.
Imperatoris-Maximiliana, *Wawra, Bot. Ergeb. Maxim. Bras.* 73.—Bras.
incarnata, *Aubl. Pl. Gui.* ii. 645. t. 261.—Guiana.
incisa, Hort. ex DC. Prod. ix. 224 = Tecoma stans.
indica, Linn. Sp. Pl. 625 = Oroxylum indicum.
indica, Lour. Fl. Cochinch. 460 = Pajanelia multijuga.
infusa, *Griseb. in Goett. Abh.* xxiv. (1879) 256.—Reg. Argent.
jasminifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 141.— Venezuela.
jasminoides, A. Cunn. ex DC. Prod. ix. 225 = Tecoma jasminoides.
jasminoides, *Thunb. Pl. Bras.* iii. 86. n. 28.—Bras.
Kerere, *Aubl. Pl. Gui.* ii. 644.—Guiana.
labiata, *Cham. in Linnaea*, vii. (1832) 701.—Bras.
lactiflora, Vahl, Symb. Bot. iii. 80. t. 66 = Dianthera lactiflora.
lacta, *Wall. Cat.* n. 6505.—Burma.
lanata, Steud. Nom. ed. II. i. 205 = discolor.
lanceolata, *DC. Prod.* ix. 168.—Bras.
lanuginosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 461.— Mexic.
laseripitifolia, *Mart. ex DC. Prod.* ix. 168.—Bras.
latifolia, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Callichlamys riparia.
latiuliana, *Poir. Encyc. Suppl.* i. 682.—Guiana.
laurifolia, *Vahl. Ecl. Am.* ii. 44.—Guiana.
lexiflora, DC. Rev. Bign. 21 ; Prod. ix. 178 = Pitheocctenium lanifforum.
lenta, *Mart. ex DC. Prod.* ix. 159.—Bras.
lepidophylla, *A. Rich. Fl. Cub. Fanerog.* ii. 104.— Isa. Pine. (Ind. occ.).
lepidota, H. B. & K. Nov. Gen. et Sp. iii. 139 = Tecoma lepidota.

BIGNONIA :—
lepidota, *Seem. Bot. Voy. Herald*, 197.—Mexic.
leucantha, Vell. Fl. Flum. 251 ; vi. t. 49 = Sparattosperma lithontripticum.
leucopogona, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = scq.
Leucopogon, *Cham. in Linnaea*, vii. (1832) 707.— Bras.
leucoxylla, Vell. Fl. Flum. 252 ; vi. t. 54 = Tabebuia leucoxyla.
Leucoxylon, DC. Cat. Bign. ex Prod. ix. 173 = Tecoma Salamanni.
Leucoxylon, Linn. Sp. Pl. 624 = Tecoma Leucoxylon.
Lindleyi, *DC. Prod.* ix. 147.—Reg. Argent.
linearis, Cav. Ic. iii. 35. t. 269 = Chilopsis saligna.
littoralis, *H. B. & K. Nov. Gen. et Sp.* iii. 139.— Mexic.
longa, Vell. Fl. Flum. 249 ; vi. t. 37 = Lundia longa.
longiflora, DC. Prod. ix. 206, sphalm. = Pajanelia Rheedii
longiflora, Vell. Fl. Flum. 252 ; vi. t. 52 = Tecoma speciosa.
longiflora, *Cav. Ic.* vi. 58. t. 581.—Gasyaquil.
longifolia, Willd. Sp. Pl. iii. 306 = Pajanelia Rheedii.
longistiliqua, Vell. Fl. Flum. 247 ; vi. t. 26 = Arrabidae.
longistiliqua, *Bert. ex Spreng. Syst.* ii. 830.—Venezuela
longissima, Jacq. Enum. Pl. Carib. 25 = Catalpa longissima.
longissima, Lour. Fl. Cochinch. ii. 580 = Dolichandrone Rheedii.
lucida, *Lodd. ex Loud. Hort. Brit.* 242.—Am austr.
lugubris, Salisb. Prod. 106 = Oroxylum indicum.
Lundii, DC. Rev. Bign. 21, ex Prod. ix. 179 = Pithecoctenium Lundii.
macrophylla, Cham. in Linnaea, vii. (1832) 689 = Adenocalymna macrophyllum.
macrostachya, *Wall. Cat.* n. 6504.—Burma.
magnifica, *Bull, in Gard. Chron.* (1879) ii. 78.—N Granat.
magnoliaefolia, *H. B. & K. Nov. Gen. et Sp.* iii. 136.— Venezuela.
Mansoana, *DC. Prod.* ix. 157.—Bras.
marashamaensis, *DC. l. c.* 165.—Bras.
marginata, Cham. in Linnaea, vii. (1832) 695 = Adenocalymna marginatum.
marmoratum, *Hort. ex André, in Rev. Hortic.* (1865) 268.—Hab. ?
Martini, *DC. Prod.* ix. 152.—Guiana.
Martiusiana, *DC. l. c.* 156.—Bras.
megapotamica, *Spreng. Syst.* iv. 287.—Bras.
meonantha, Link, Enum. ii. 130 = Tecoma australis.
meridionalis, Salisb. Prod. 106 = Tecoma Leucoxylon.
Meyeniana, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 366.—Peruv.
microcalyx, *G. F. W. Mey. Prim. Fl. Esseq.* 211.— Guiana.
microphylla, *Lam. Encyc.* i. 418.—Ins. S. Doming.
Mina, *Robin, Voy. Louis.* iii. 408.—Am. bor.
miranda, Cham. in Linnaea, vii. (1832) 687 = Anemopaegma mirandum.
mollis, *Vahl, Ecl. Am.* ii. 46.—Guiana.
mollissima, *H. B. & K. Nov. Gen. et Sp.* iii. 138.— Venezuela.
molusacea, *DC. Prod.* ix. 144.—Ins. Amboina.
moringaefolia, *DC. l. c.* 170.—Guiana.
multifida, *Lodd. ex Loud. Hort. Brit.* 242.—Ind. or.
multiflora, *Salzm. ex DC. Prod.* ix. 160.—Bras.
multijuga, Wall. Cat. n. 6508 ; Pl. Rar. As. i. 81. t. 95, 96 = Pajanelia Rheedii.
muricata, DC. Rev. Bign. 22, ex Prod. ix. 173 = Pithecoctenium muricatum.
muricata, Herb. Madr. ex Wall. Cat. n. 6511 = Stereospermum xylocarpum.
myriantha, Cham. ex Steud. Nom. ed. II. i. 205 = Tecoma myriantha.
nematocarpa, Boj. Hort. Maurit. 219 = Stereospermum nematocarpum.
nervosa, Domb. ex DC. Prod. ix. 196 = Delostoma nervosum.
nigricans, *Regel, Cat. Pl. Hort. Absolon.* 19.—Hab. ?
nitidissima, *DC. Prod.* ix. 180.—Venezuela.
nitidula, Mart. ex DC. l. c. 181 = Landia nitidula.
noctiflora, Buch.-Ham. ex Wall. Cat. sub n. 6499 = Stereospermum suaveolens.
nodosa, *Mexter, Enum. Cat. Bras.* 40.—Bras.
noterophila, *DC. Prod.* ix. 148.—Bras.

BIGNONIA :—
oblique, *H. B. & K. Nov. Gen. et Sp.* iii. 185.—Venezuela.
obovata, Hook. & Arn. Bot. Beech. Voy. 489 = Stemmadenia pubescens.
obovata, Spreng. Syst. ii. 830 = Spathodea obovata.
obovata, Vell. Fl. Flum. 251; vi. t. 45 = Jacaranda subrhombea.
obtusifolia, Lam. Encyc. i. 418 = Spathodea obtusifolia *seu* Tabebuia ulginosa.
occidentalis, *Salsm. ex DC. Prod.* ix. 160.—Bras.
odorata, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 98.—Porto Rico.
ophthalmica, *Chisholm, ex Steud. Nom.* ed. II. i. 205. —Guiana.
orbiculata, Jacq. Enum. Pl. Carib. 25 = Anemopaegma orbiculatum.
ornata, *Linden, Cat.* n. 21 (1867) 1; *E. Morr. Belg. Hortic.* xvii. (1867) 102.—Brsa.
oxyphylla, *DC. Prod.* ix. 169 (*Cf. Fl. Brit. Ind.* iv. 876).—Ind. or.?
pachyptera, Dc. Rev. Bign. 21; ex Prod. ix. 174 (= *Pachyptera pubernula*).—Bras.
Paiannila, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 515 = Pajanella Rheedii.
pallida, *Lindl. Bot. Reg.* t. 965.—Ind. occ.
pandorana, Andr. Bot. Rep. t. 86 = Tecoma australis.
Pandorea, Vent. Jard. Malm. t. 48 = Tecoma australis.
paniculata, Linn. Sp. Pl. 623 = Amphilophum paniculatum.
pannosa, *DC. Prod.* ix. 148.—Peruv.
Parkeri, *DC. l. c.* 157.—Guiana.
parviflora, Rottl. ex Wall. Cat. sub n. 6501 = Stereospermum chelonoides.
parviflora, Salsm. ex DC. Prod. ix. 183 = Arrabidaea Salzmanni.
patellifera, *Schlecht. in Linnaea*, viii. (1833) 516.—Mexic.
Patrisiana, DC. Prod. ix. 174 = Tecoma Patrisiana.
pedunculata, *Vell. Fl. Flum.* 250; vi. t. 41.—Bras.
pentandra, Lour. Fl. Cochin. ii. 379 = Oroxylum indicum.
pentaphylla, Linn. Sp. Pl. ed. II. 870 = Tabebuia pentaphylla.
pentaphylla, Salzm. ex DC. Prod. ix. 219 = Tecoma Salzmanni.
perforata, *Cham. in Linnaea*, vii. (1832) 667.—Bras.
perianthomega, *Vell. Fl. Flum.* 248; vi. t. 34.—Bras.
peruviana, *Linn. Sp. Pl.* 625.—Peruv.
petiolaris, DC. Cat. Bign. 22; ex Prod. ix. 174 = Tecoma Leucoxylon.
phaseoloides, *Cham. in Linnaea*, vii. (1832) 698.—Bras
physaloides, *Cham. l. c.* 668.—Bras.
picta, *H. B. & K. Nov. Gen. et Sp.* iii. 186.—Guiana.
picta, Lindl. Bot. Reg. (1842) t. 45 = Lindleyi.
pilosa, Dietr. ex Steud. Nom. ed. II. i. 205 = villosa.
pilulifera, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guiana.
platyphylla, Cham. in Linnaea, vii. (1832) 679 = brachypoda.
Poeppigii, DC. Rev. Bign. ex Prod. ix. 174 = Anemopaegma Poeppigii.
popaifolia, *DC. Prod.* ix. 160.—Bras.
Porteriana, Wall. Cat. n. 6509 = Stereospermum glanduliosum.
Prieurei, *DC. Prod.* ix. 154.—Guiana.
prunea, Willd. Sp. Pl. iii. 307 = Jacaranda Copaia.
pseudonugis, *Desf. Cat. Hort. Par.* ed. III. 397.—Ind. occ.
pterocarpa, Cham. in Linnaea, vii. (1832) 673 = Caspidaria trifoliata.
pubescens, *Linn. Sp. Pl.* ed. II. 870.—Mexic.
pubescens, Pav. ex DC. Prod. ix. 148 = pannosa.
pulchella, *Cham. in Linnaea*, vii. (1832) 663.—Bras.
pulchra, *Cham. l. c.* 708.—Bras.
punctata, DC. Rev. Bign. ex Prod. ix. 160 = occidentalis.
punicea, *Mart. Herb. Fl. Bras.* 255.—Bras.
purpurea, *Ledd. ex Sweet, Hort. Brit.* 283, nomen; *Bot. Mag.* t. 3800.—Am. austr.
pyramidata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Tabebuia pyramidica.
quadrangularis, *Lechert, ex Steud. Nom.* ed. II. i. 205. —Ins. Trinitat.
quadrilocularis, Roxb. Fl. Corom. ii. 24. t. 145 = Hexaphragma Roxburghii.
quadripinnata, Blanco, Fl. Filip. ed. I. 499 = Oroxylum indicum.

BIGNONIA :—
quadrivalvis, *Jacq. Fragm.* 97.—Venezuela.
Quercus, Lam. Encyc. i. 417 = Catalpa longissima.
quinquefolia, Vell. Fl. Flum. 253; vi. t. 50 = Cybistax antisyphilitica.
racemosa, *Lam. Encyc.* i. 424.—Madag.
radiata, Linn. Sp. Pl. 624 = Argylia Feuillei.
radicans, Linn. l. c. = Tecoma radicans.
ramiflora, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 381.—Coles mauritians.
Regnelliana, *Sond. in Linnaea*, xxii. (1849) 558.—Bras.
Rego, Vell. Fl. Flum. 249 = Arrabidaea Sego.
reticulata, Hb. Madr. ex Wall. Cat. n. 6511 = Stereospermum xylocarpum?
Revelli, *Hamon, in Bull. Soc. Hortic. Prat. Rhône*, (1851) 149. t. 3.—Hab.?
rhodoantha, *Zipp. ex Blume, Rumphia*, iv. 35.—N Guin.
rigescens, Jacq. Hort. Schoenb. h. 44. t. 210 = Distictis rigescens.
riparia, *H. B. & K. Nov. Gen. et Sp.* iii. 188.—N. Granat.
rivularis, Hort. Kew. (1830) ex DC. Prod ix. 199 = Cybistax antisyphilitica.
rosea, DC. Rev. Bign. ex Prod. ix. 174 = Arrabidaea rosea.
rosea, Pav. ex DC. Prod. ix. 198 = Delostoma denitatum.
rotundata, *DC. l. c.* 168.—Bras.
rufescens, DC. Rev. Bign. ex Prod. ix. 174 = Pithecoctenium rufescens.
rufescens, Hoffingg. ex DC. Prod. ix. 218 = Callichlamys riparia.
rugosa, *Schlecht. in Linnaea*, xxvi. (1853–55) 656.—N. Granat.
rupestris, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 179.—Bras.
Sagraeana, *DC. Prod.* ix. 148.—Cuba.
salicifolia, Buch.-Ham. ex Wall. Cat. n. 6498 = Tecoma undulata.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 183.—Venezuela.
Salzmanni, DC. Rev. Bign. ex Prod. ix. 174 = Adenocalymna Salzmanni.
sanyidana, *Cham. in Linnaea*, vii. (1832) 669.—Bras.
sambucifolia, *Hort. ex Carr. Rev. Hortic.* (1878) 50.—Hab.?
sarmentosa, Bertol. f. Fl. Guatim. 25 = sequinoctialis.
scandens, Aca. Vell. Fl. Flum. 246; vi. t. 22 = Anemopaegma racemosum.
Sceptrum, *Cham. in Linnaea*, vii. (1832) 710.—Bras.
Sego, Vell. Fl. Flum. vi. t. 39 = Arrabidaea Agnuscastus.
Selloi, *Spreng. Syst.* ii. 831.—Bras.
sempervirens, Linn. Sp. Pl. 623 = Gelsemium sempervirens.
serrata, Pav. ex G. Don, Gen. Syst. iv. 328 = castaneaefolia.
serratifolia, Vahl, Ecl. Am. ii. 46 = Tecoma serratifolia.
Serratula, Wall. ex DC. Prod. ix. 206 = Stereospermum serrulatum.
serrulata, Steud. Nom. ed. II. i. 205, sphalm. = prace.
Sieberi, DC. Rev. Bign. ex Prod. ix. 175 = Arrabidaea Sieberi.
simplex, *A. Rich. Fl. Cub. Fanerog.* ii. 104.—Cuba.
simplicifolia, Pav. ex DC. Prod. ix. 198 = Delostoma integrifolium.
Sinclairii, *Benth. Bot. Voy. Sulph.* 129 = laurifolia.
sordidula, Sallsb. Prod. 106 = Tecoma stans.
sordida, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1158.—Guiana.
spathacea, Linn. f. Suppl. 288 = Dolichandrone Rheedii.
spathacea, Roxb. Pl. Corom. ii. 24. t. 144 = Dolichandrone falcata.
speciosa, *R. Grah. in Bot. Mag.* t. 8888.—Bras.
spectabilis, Sallsb. Prod. 106 = Catalpa bignonoides.
spectabilis, Vahl, Symb. Bot. lii. 80 = aequinoctialis.
spicata, *Lechhart, ex Steud. Nom.* ed. II. i. 205.—Ins. Trinitat.
Splendor-sylvae, *Mart. ex DC. Prod.* ix. 154.—Bras.
squalea, Vell. Fl. Flum. 244; vi. t. 18 = Pitheoctenium squalum.
squamellulosa, DC. Rev. Bign. ex Prod. ix. 174 = Tecoma squamellulosa.
stanifera, *Lam. Encyc.* i. 421.—Ins. S. Doming.
stans, Linn. Sp. Pl. ed. II. 871 = Tecoma stans.
stipulata, Roxb. Hort. Beng. 47; Fl. Ind. iii. 108 = Dolichandrone stipulata.

BIGNONIA :—
striata, DC. Rev. Bign. ex Prod. ix. 174 (= *Pachyptera striata*).—Bras.
suaveolens, Roxb. Hort. Beng. 47; Fl. Ind. iii. 104 Stereospermum suaveolens.
suberosa, Roxb. ll. cc. 47, 104 = Millingtonia hortensis
subincana, *Hensl. Bisl. Centr. Am. Bot.* ii. 491.—Panama.
subsericosa, DC. Rev. Bign. ex Prod. ix. 220 = Tecoma subsericosa.
Tabebuya, Vell. Fl. Flum. 251; vi. t. 53 = Tabebuia ulginosa.
Tagoada, Roxb. ex Wall. Cat. n. 6501 = Stereospermum chelonoides.
tecomoides, DC. Prod. ix. 166 = Tecoma mollis.
tenuiflora, *DC. l. c.*—Bolivia.
Telfairiae, *Boy. ex Hook. Bot. Mag.* t. 2976.—Madag
tenuisiliqua, *Vahl, Ecl. Am.* ii. 43.—Am. austr.
ternata, G. Don, Gen. Syst. iv. 221 = maranhamensis.
ternata, *Vell. Fl. Flum.* 246; vi. t. 21.—Bras.
tetragona, Wall. ex DC. Prod. ix. 211 = Stereospermum tetragonum.
tetragonocaulos, *DC. Prod.* ix. 169.—Bras.
tetragonolobe, *Jacq. Fragm.* 98.—Venezuela.
tetraquetra, *Cham. in Linnaea*, vii. (1832) 675.—Bras
thyrsoides, *DC. Prod.* ix. 158.—Bras.
tiliaefolia, *H. B. & K. Nov. Gen. et Sp.* iii. 186.—Am. austr.
tinctoria, *Arruda, in Koster, Trav. Bras.* 497.—Bras.
tomentosa, Pav. ex DC. Prod. ix. 219 = Tecoma heteropoda.
tomentosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = mollis.
tomentosa, Thunb. Fl. Jap. 252 = Paulownia imperialis.
triantha, DC. Rev. Bign. ex Prod. ix. 148 = exoleta.
trichocladu, *DC. l. c.* 158.—Bras.
trifora, Pav. ex DC. l. c. 148 (est Rubiaces).
trifoza, *Vell. Fl. Flum.* 249; vi. t. 38.—Bras.
trifoliata, Vell. l. c. 245; vi. t. 16 = Adenocalymna longerraceanum.
triphylla, Vell. l. c. 244; vi. t. 14 = triternata.
triphylla, Linn. Sp. Pl. ed. II. 870 = Tabebuia triphylla.
triphylla, Willd. ex DC. Prod. ix. 151 = B. Chica.
tripinnata, *Noronha, in Verh. Batow. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
triplinervia, *Mart. ex DC. Prod.* ix. 153.—Bras.
triternata, *Steud. Nom.* ed. II. i. 205.—Bras.
Tropaeolum, Jacquem. ex DC. Prod. ix. 228, in nota = Tecoma undulata.
tuberculata, Roxb. ex DC. Prod. ix. 177 = Oroxylum indicum.
tuberculosa, Vell. Fl Flum. 251; vi. t. 46 = Zeyheria montana.
tuberculosa, Schum. Beskr. Guin. Pl. 278 = Spathodea tulipifera.
Tweedlana, *Lindl. Bot. Reg.* (1840) t. 45.—Bras.
ulginosa, Gomez, Obs. ll. 7 t. 2; Mem. Ac. Lisbon. ill. (1812) Mem. dos Corresp. 57 = Tabebuia ulginosa
umbellulata, *DC. Prod.* ix. 148.—Bras.
umbrosa, *H. B. & K. Nov. Gen. et Sp.* iii. 188.—N. Granat.
uncata, Andr. Bot. Rep. t. 580 = Macfadyena uncinata.
uncata, Sims, Bot. Mag. t. 1511 = Spathodea uncata.
unguiculata, G. F. W. Mey. Prim. Fl. Esseq. 210 = Macfadyena uncinata.
undulata, Sm. Exot. Bot. i. 35 = Tecoma undulata.
unguiculata, *Vell. Fl. Flum.* 248; vi. t. 32.—Bras.
unguiculata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = B. Unguis-cati.
Unguis, Linn. Syst. ed. X. xiii. n. 8 = B. Unguis-cati
Unguis-cati, *Linn. Sp. Pl.* 623.—Am. trop.
Vargasiana, DC. Rev. Bign. ex Prod. ix. 175 = Anemopaegma Vargasianum.
variabilis, *Jacq. Hort. Schoenb.* ii. 45. t. 212.—Venezuela.
variabilis, Mart. Herb. Fl. Bras. 256 = varians.
variabilis, Sieber, ex E. Mey. in Nov. Act. Nat. Cur xii. (1825) 779 = Hostmanni.
variana, *DC. Prod.* ix. 149.—Bras.
velutina, *DC. l. c.* 162.—Bras.
venusta, *Kw-Gaul. in Bot. Reg.* t. 249.—Bras.
verrucifera, *Schlecht. in Linnaea*, xxvi. (1853–55) 655. N. Granat.
verrucosa, *H. B. & K. Nov. Gen. et Sp.* iii. 187.—Venezuela.
villosa, Bert. ex Spreng. Syst. ii. 830 = Bolbisiana.
villosa, *Vahl, Ecl. Am.* ii. 43.—Venezuela.

BIGNONIA :—
viminalis, H. B. & K. Nov. Gen. et Sp. iii. 182 = Tecoma viminalis.
violacea, *DC. Prod.* ix. 186.—Bolivia.
virginalis, Mart. ex DC. l. c. 181 = Lundia virginalis.
viridiflora, Lodd. Bot. Cab. xi. (1825) t. 1026 = Cybistax antisyphilitica.
Vitalba, Cham. in Linnæa, vii. (1832) 699 = Pithecoctenium Vitalbæ.
xylocarpa, Roxb. Hort. Beng. 47 ; Fl. Ind. iii. 108 = Stereospermum xylocarpum.

BIKAI, Adans. Fam. ii. 67 (1763) = **Heliconia**, Linn. (Scitam.).

BIHANIA, Meisn. in DC. Prod. xv. t. 96 (1861) = **Buxideroxylon**, Teijsm. (Laurin.).
borneensis, Meisn. l. c.—Borneo.

BIKERA, Adans. Fam. ii. 180 (1763) = **Tetragonotheca**, Dill. (Compos.).

BIKKIA, Reinw. ex Blume, Bijdr. 1017, in syn. (1826) ; et in Syll. Ratisb. ii. (1828) 8. *RUBIACEÆ*, Benth. & Hook. f. ii. 46.
BIKKIOPSIS, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 404.
CORMIGONUS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 83.
GRISIA, Brongn. in Bull. Soc. Bot. Fr. xii. (1865) 405.
TATEA, Seem. Fl. Vit. 125 (1866).
australis, DC. Prod. iv. 403 = grandiflora.
Forsteriana, *Brongn. in Bull. Soc. Bot. Fr.* xiii. (1866) 42.—Ins. Fiji.
Gaudichaudiana, *Brongn. l. c.* 43.—N. Guinea.
grandiflora, *Reinw. ex Blume, Bijdr.* 1017, *in syn. ; et in Syll. Ratisb.* ii. (1828) 8.—Java ; Ins. Amic.
Guillovieana, *Brongn. in Bull. Soc. Bot. Fr.* xiii. (1866) 43.—N. Guinea.
Hornbroniana, *Brongn. l. c.* 43.—Ins. Pacif.
mariannensis, *Brongn. l. c.* 42.—Ins. Marian.

BIKKIOPSIS, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 404 = **Bikkia**, Reinw. (Rubiac.).
Pancheria, Brongn. l. c.—N. Caled.

BIKUKULLA, Adans. Fam. ii. 23 (1763) = **Diolytra**, Borckh. (Papav.).

BILABIUM, Miq. Fl. Ind. Bat. ii. 730 (1856). *DIDYMOCARPEÆ*, Benth. & Hook. f. Gen. ii. 996.
limaus, *Miq. l. c.*—Malaya.

BILABRELLA, Lindl. Bot. Reg. sub t. 1701 (1835) = **Habenaria**, Willd. (Orchid.).
falciornis, Lindl. l. c. = H. tetrapetala.
falciformis, Bolus, in Journ. Linn. Soc. xix. (1882) 340 = prasc.
salisicornis, Steud. Nom. ed. II. i. 205, sphalm. = prasc.

BILAMISTA, Rafin. New Fl. Am. iv. 92 (1836) = **Gentiana**, Tourn.
grandiflora, Rafin. l. c. 98 = G. grandiflora !

BILDERDYKIA, Dum. Florul. Belg. 18 (1827) = **Polygonum**, Tourn.
Convolvulus, Dum. l. c. = P. Convulvulus.
dumetorum, Dum. l. c. = P. dumetorum.

BILITALIUM, Buch.-Ham. Journ. Mysore, 185 (1807), nomen.
farinosum, Buch.-Ham. l. c. (Quid ?).—Ind. or.

BILLARDIERA, Moench, Meth. 869 (1794) = **Verbena**, Linn.
explanata, Moench, l. c. = V. Aubletia.

BILLARDIERA, Sm. Bot. Nov. Holl. i. t. 1 (1798). *PITTOSPOREÆ*, Benth. & Hook. f. i. 182.
LABILLARDIERA, Roem. & Schult. Syst. v. p. xxvi. 380 (1819).
angustifolia, DC. Prod. i. 345 = scandens.
brachyantha, F. Muell. ex Klatt, in Linnæa, xxviii. (1856) 671 = scandens.
canariensis, Wendl. Beobacht. 48 = scandens.
coriacea, *Benth. Fl. Austral.* i. 124.—Austral.
cymosa, *F. Muell. in Trans. Vict. Inst.* i. (1855) 29. —Austral.

BILLARDIERA :—
cymosa, Klatt, in Linnæa, xxviii. (1856) 571 = cymosa.
daphnoides, *Knowles & Westc. Flor. Cab.* ii. (1838) 60. —Austral.
elegans, *F. Muell. Pl. Vict.* i. 78.—Austral.
floribunda, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 1.—Austral.
fusiformis, Labill. Nov. Holl. Pl. t. 90 = Sollya fusiformis.
grandiflora, Putterl. Stirp. Nov. Decad. 48 = scandens.
Hambrechtiana, Seem. & Schmidt, in Flora, xxvii. (1844) 11. 496 = Sollya heterophylla.
latifolia, Putterl. Stirp. Nov. Decad. 47 = scandens.
Lehmanniana, *F. Muell. Pl. Vict.* i. 78.—Austral.
longiflora, *Labill. Nov. Holl. Pl.* i. 64. t. 89.—Tasmania.
macrantha, Hook. f. Fl. Tasm. i. 87 = longiflora.
melocarpa, *Hort. ex Steud. Nom.* ed. II. i. 206.—Austral.
mutabilis, *Salisb. Parad. Lond.* t. 48.—Austral.
ovalis, Lindl. Bot. Reg. t. 1719 = longiflora.
parviflora, DC. Prod. i. 345 = Marianthus tenuis.
pseudocymosa, Klatt, in Linnæa, xxviii. (1856) 571 = cymosa.
rosmarinifolia, *DC. Prod.* i. 345.—Austral.
scandens, *Sm. Exot. Bot.* i. t. 1.—Austral.
sericophora, F. Muell. in Linnæa, xxv. (1852) 371 = cymosa.
variifolia, *DC. Prod.* i. 346.—Austral.
versicolor, F. Muell. ex Klatt, in Linnæa, xxviii. (1856) 751 = cymosa.

BILLARDIERA, Vahl, Eclog. Am. i. 18. t. 10 (1796) = **Coussaria**, Aubl. (Rubiac.).
paniculata, Vahl, l. c. = C. Fruelichia.

BILLBERGIA, Thunb. Dec. Pl. Bras. 30 (1823). *BROMELIACEÆ*, Benth. & Hook. f. iii. 664.
CREMOSTYRS, Beer, in Flora, xxxvii. (1854) 348.
EUCALIAS, Rafin. Fl. Tellur. iv. 25 (1836).
HELICODEA, Lem. Illustr. Hortic. xi. (1864) t. 421.
JONGHRA, Lem. Jard. Fleur. ii. sub t. 180-181 (1852).
PANETA, Lem. Illustr. Hortic. ii. (1855) t. 48.
aculis, Hort. ex Vis.—Cf. Otto & Dietr. Allgem. Gartenz. xvi. (1848) 29 = Cryptanthus sonatus.
amabilis, Beer, Bromel. 118 = vittata.
amœna, Lindl. Bot. Reg. sub t. 1068 = speciosa.
angustifolia, *C. Koch, Wochenschr.* ix. (1866) 181.—Hab. ?
atro-rosea, *Drapiez, Encyc. et Fl. des Serr. Angl.* (1834) f. 1.—Bras.
Bakeri, *E. Morr. in Belg. Hortic.* xxx. (1880) 166. t. 8. —Bras.
Baraquiniana, Lem. Illustr. Hortic. (1864) 421 = decora.
bicolor, Lodd. Bot. Cab. t. 1819 = pyramidalis.
bicolor, Schult. f. Syst. vii. 1256 = Rhodostachys bicolor.
bifrons, Lindl. in Journ. Hort. Soc. viii. (1853) 54 = Pitcairnea bracteata !
bivittata, Hook. Bot. Mag. t. 5270 = Cryptanthus bivittatus.
Beauveana x, André, in Rev. Hortic. (1884) 117.
Brongniarti, Regel, Gartenfl. (1875) 106 ; et in Act. Petrop. iii. 11. (1875) 233 = Fortes kermesina.
Chantini, *Carr. in Rev. Hortic.* (1878) 112.—Bras.
chlorocyanea, De Vriese, Hort. Acad. Lugd. Bat. Pl. Rar. (1854).—Cf. Linnæa, xxvi. (1853–55) 755 = speciosa.
chloratica, Hort. Saunders ; ex Gard. Chron. (1871) 1426 = Saundersii.
clavata, Lindl. Bot. Reg. sub t. 1068 = Aechmea bromeliæfolia.
coelestis, Hort. ex Morr. in Fl. des Serres, xxi. (1875) 2195 = Aechmea coelestis.
Colleei x, A. Van Geert, Cat. n. 81. p. 91 ; Belg. Hortic. xxxii. (1882) 334.
commixta, C. Koch, Wochenschr. iii. (1860) 146 = zebrina !
contracta, Mart. ex Schult. f. Syst. vii. 11. 1263 = Aechmea contracta.
corymbosa, *Mart. ex Schult. f. l. c.* 1266.—Bras.
Creyiana, De Jongh, ex Lem. Jard. Fleur. iv. (1854) t. 418 = pyramidalis.
cruenta, *Hook. Bot. Mag.* lvi. t. 2892.—Bras.
decora, *Poepp. & Endl. Nov. Gen.* 48.—Peruv.
dicolor, Beer, Bromel. 121 = speciosa.
distichoclastchya, Lem. Jard. Fleur. ii. (1852) Misc. 96 = Aechmea distichoclastcha.

BILLBERGIA :—
elegans, Mart. ex Schult. f. Syst. vii. 11. 1265 = speciosa.
Euphemiæ, *E. Morr. Belg. Hortic.* xxii. (1872) 1. 1.—Bras.
excellens, *Miq. in Journ. Bot. Néerl.* i. (1861) 84.—Guiana.
fasciata, Lindl. Bot. Reg. xiii. t. 1120 = Aechmea fasciata.
fastuosa, Beer, Bromel. 110 = thyrsoidea.
filicaulis, *Griseb. in Goett. Nachr.* (1864) 18.—Venezuela.
Glasioviana, Regel, Gartenfl. (1885) 280 = Quesnelia strobilispica.
Glymiana, De Vriese, Jaarb. der Kon. Ned. Maatsch Tuinb. (1858) 87 = Morelli.
Glymiana, Beer, Bromel. 119 = Morelli.
gracilis, Poepp. ex Beer, l. c. 254 = Aechmea bilianidoides.
granulosa, Brongn. in C. Koch, Wochenschr. iii. (1860) 145 = decora.
horrida, *Regel, Ind. Sem. Hort. Petrop.* (1856) 17 ; *Gartenfl.* (1857) 148.—Bras.
ianthina, Hort. ex E. Morr. Belg. Hortic. (1871) 2 = Leopoldi.
incarnata, *Schult. f. Syst.* vii. 11. 1261.—Peruv.
iridifolia, *Lindl. Bot. Reg.* t. 1068.—Bras.
Ivanwilleı, Van Houtte, ex E. Morr. in Belg. Hortic. (1876) 161 = Rhodostachys pitcairniæfolia.
Jolibeiı x, A. Van Geert, Cat. n. 81. p. 81 ; Belg. Hortic. xxxii. (1882) 334.
lanuginosa, Hort. Hamb. ex C. Koch, Ind. Sem. Hort. Berol. (1856) App. 7 = Aechmea nudicaulis.
latifolia, Hort. Belg. ex Heynh. Nom. ii. 60 = Aechmea sp. !
Leopoldiana, Hort. ex Baker, in Saund. Ref. Bot. t. 285 = Orgiesiæ sp.
Leopoldi, *Linden, in Rev. Hortic.* (1869) 87 ; *E. Morr. in Belg. Hortic.* xxi. (1871) 1. t. 1.—N. Granat. ; Bras
Leopoldi, Hort. Belg. ex C. Koch, Ind. Sem. Hort. Berol. (1856) 5 = vittata.
Liboniana, *De Jongh. Journ. Hort. Prat.* (Mars. 1851) ; *et Lem. Yard. Fleur.* ii. (1852) t. 197.—Bras.
Lietzei, *E. Morr. in Belg. Hortic.* xxxi. (1881) 97. t. 5-7.—Bras.
Loddigesii, Steud. Nom. ed. II. i. 206 = pyramidalis.
longifolia, C. Koch & Bouché, Ind. Sem. Hort. Berol (1856) 5 = thyrsoides.
lutea, Schult. f. Syst. vii. 1258 = Aechmea nudicaulis.
macrocalyx, *Hook. Bot. Mag.* t. 5114.—Bras.
marmorata, *Lem. Illustr. Hortic.* ii. (1855) t. 48.—Bras.
Martinezii, Miq. in Linnæa, xviii. (1844) 376 = Aechmea Martenzii.
Meyendorfii, Regel, Ind. Sem. Hort. Petrop. (1857) 37 = Karatas Carolinæ.
miniato-rosea, Lem. Illustr. Hortic. ii. (1855) Misc. 94 = thyrsoides.
mitis, *Mart. ex Schult. f. Syst.* vii. 1267.—Bras.
Moreliana, Brongn. ex Rev. Hortic. Sér. III. iii. (1849) 89 = Morelli.
Morsliana, Lem. Jard. Fleur. ii. (1852) t. 188 = vittata.
Morelli, *Brongn. in Portef. Hort.* ii. (1848) 97 ; *et E. Morr. Belg. Hortic.* x. (1860) 161.—Bras.
nudicaulis, Lindl. Bot. Reg. sub t. 1068 = Aechmea nudicaulis.
nuptialis, Hort. ex E. Morr. in Belg. Hortic. (1871) 2 = Leopoldi.
nutans, *H. Wendl. in Regel, Gartenfl.* (1869) 162.—Bras.
Oberthuri x, A. Van Geert. Cat. n. 81. p. 92 ; Belg. Hortic. xxxii. (1882) 334.
obtecta, Mart. ex Schult. f. Syst. vii. 1256 (= Aechmea sp.).—Bras.
odora, Miq. in Linnæa, xviii. (1844) 377 = Aechmea odora.
olens, Hook. f. Bot. Mag. t. 5502 = Karatas Carolinæ.
pallescens, Baker, in Hook. f. l. c. t. 6949 = Bakeri.
pallescens, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1856) App. 5 = speciosa.
pallida, Beer, Bromel. 121 = speciosa.
pallidiflora, *Liebm. Ind. Sem. Hort. Hafn.* (1854) 26.—Nicaragua.
paniculata, Mart. ex Schult. f. Syst. vii. 11. 1263 = Aechmea paniculata.
parviflora, Mart. ex Schult. f. l. c. 1270 = Aechmea parviflora.

BILLBERGIA :—
patentissima, Mart. ex Schult. f. Syst. vii. 11. 1270 = Aechmea patentissima.
Pazzensi, Beer, Bromel. 118 = thyrsoidea.
polystachya, Lindl. & Paxt. in Paxt. Flow. Gard. iii. (1853–58) 88. t. 80 = Aechmea distichantha.
Porteana, *Brongn. ex Beer, Bromel.* 115.—Bras.
pulcherrima, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1856) App. 6 = vittata.
punicea, *Beer, Bromel.* 119.—Hab.?
purpurea, *Beer, l. c.* 126 (≠ Pitcairniae sp. ?).—Am. trop.
purpuro-rosea, Hook. Bot. Mag. t. 3304 = Aechmea suavolens.
pyramidalis, Forbes, Hort. Woburn. 64 = Aechmea nudicaulis.
pyramidalis, *Lindl. Bot. Reg.* sub t. 1068.—Bras.
pyramidata, Beer, Bromel. 122 = Aechmea nudicaulis.
Quesneliana, *A. Brongn. in Ann. Sc. Nat.* Sér. II. xv. (1841) 371.—Guiana.
Rancogneri v., André, in Rev. Hortic. (1884) 524.
Reichardtii, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 115.—Bras.
rhodocneis v., A. Van Geert, Cat. n. 81. p. 92; Belg. Hortic. xxxii. (1882) 884.
rhodocyanea, Lem. in Fl. des Serres, Sér. I. iii. (1847) 207 = Aechmea fasciata.
Rohaniana, De Vriese, in Linnaea, xxvi. (1853–55) 756 = vittata.
rosea, Beer, Bromel. 123 = Porteana.
roseo-marginata, Hort. ex C. Koch, Wochenschr. (1867) 141 = Quesnelia rufa.
rubro-marginata, Hort. ex Carr. in Rev. Hortic. (1880) 70 = Quesnelia rufa.
Sanderiana, *S. Morr. in Belg. Hortic.* xxxiv. (1884) 17. —Bras.
Saundersii, *Hort. Bull, ex Floral Mag.* N. S. t. 106.—Bras.
sextilis, *Beer, Bromel.* 127.—Bras.
Shinneri, Hort. ex Carr. in Rev. Hortic. (1880) 70 = Quesnelia rufa.
speciosa, *Thunb. Dec. Pl. Bras.* iii. 30.—Bras.
sphaeolata, Schult. f. Syst. vii. 1969 = Greigia sphaeolata.
splendida, Lem. Jard. Fleur. ii. (1852) t. 180–181 = thyrsoidea.
tetrantha, Beer, Bromel. 197 = Tillandsia tetrantha.
thyrsoidea, *Mart. ex Schult. f. Syst.* vii. 11. 1260.—Bras.
tillandsioides, Mart. ex Schult. f. l. c. 1269 = Aechmea tillandsioides.
tinctoria, Mart. ex Schult. f. l. c. 1256 = Aechmea bromeliaefolia.
variegata, Schult. f. l. c. 1262 = speciosa.
variegata, *Morr. in Belg. Hortic.* xxxi. (1881) 78.—Bras.
violacea, *Beer, Bromel.* 116.—Guiana.
viridiflora, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 164.—Hab.?
vittata, *Brongn. ex Morel, Portef. Hortic.* ii. (1848) 253.—Bras.
Wetherilli, *Hook. Bot. Mag.* t. 4895.—Bras.
zebrina, *Lindl. Bot. Reg.* sub t. 1068.—Bras.
zonata, Linden, Cat. n. 6 (1851) 20 = vittata.

BILLIA, Peyr. in Bot. Zeit. xvi. (1858) 153; xvii. (1859) 221 = Aesculus, Linn. (Sapind.).
columbiana, Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 367.—N. Granat.
Hippocastanum, Peyr. in Bot. Zeit. xvi. (1858) 153 = Aesculus mexicana.

BILLIETIA, G. Don, Gen. Syst. iii. 560 (1834) = Billiottia, DC. (Rubiac.).

BILLIETIA, Reichb. Nom. 176 (1841) = BILLOTTIA, R. Br. = Agonis, Lindl. (Myrtac.).

BILLIOTTIA, Endl. Gen. 1229 (1840) = BILLOTTIA, R. Br. = Agonis, Lindl. (Myrtac.).

BILLIOTTIA, DC. Prod. iv. 618 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 80.
Billiotia, G. Don, Gen. Syst. iii. 560 (1834).
Melanospidium, Colla, Hort. Ripul. 88. t. 85 (1824).

BILLIOTTIA :—
Viviania, Colla, in Mém. Soc. Linn. Par. iv. (1826) 25. t. 9.
polytrichoides, Stend. Nom. ed. II. ii. 114, sphalm. = seq.
psychotrioides, *DC. Prod.* iv. 618.—Bras.

BILLOTIA, G. Don, Gen. Syst. iii. 610, 827 (1832) = BILLOTTIA, R. Br. = Agonis, Lindl. (Myrtac.).

BILLOTIA, Sch. Bip. in Herb. & Winck. Jahrb. Pharmac. iv. (1841) in nota 155 = Crepis, Linn. (Compos.).
alpina, Sch. Bip. l. c. = C. alpina.

BILLOTTIA, R. Br. in Journ. Roy. Geogr. Soc. i. (1832) 19 = Agonis, DC. (Myrtac.).
elliptica, Sweet, Hort. Brit. ed. III. 249 = Leptospermum ellipticum.
flexuosa, R. Br. in Journ. Roy. Geogr. Soc. i. (1832) 19 = A. flexuosa.
hypericifolia, Heynh. Nom. iii. 69 = A. hypericifolia.
linearifolia, R. Br. in Journ. Roy. Geogr. Soc. i. (1832) 19 = A. linearifolia.
marginata, R. Br. l. c. 19 = A. marginata.
ovatifolia, Sweet, Hort. Brit. ed. III. 249.—Austral.
saligna, Sweet, l. c.—Austral.
theaeformis, G. Don, ex Lond. Encyc. Pl. Suppl. ii. 1380 = A. hypericifolia.

BILLOTTIA, Colla, Hort. Ripul. 20. t. 23 (1824) = Calothamnus, Labill. (Myrtac.).
acerosa, Colla, l. c. = C. quadrifida.

BIMA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1 (Quid?).

BINARIA, Rafin. Sylva Tellur. 122 (1838) = Bauhinia, Linn. (Legumin.).
cumanensis, Rafin. l. c. = Bauhinia cumanensis.

BINDERA, Rafin. New Fl. Am. iv. 71 (1836) = Aster, Tourn. (Compos.).
ciliata, Rafin. l. c. = A. ciliatus.

BINECTARIA, Forsk. Fl. Aegypt. Arab. 82 (1775) = Imbricaria, Comm. (Sapot.)

BINNENDIJKIA, Kurz, in Tijdschr. Nederl. Ind. xxviii. (1865) 164 = Leptonychia, Turcz. (Sterculiac.).
trichostylis, Kurz, l. c. = L. glabra.

BIONDEA, Usteri, Ann. Bot. x. (Neue Ann. iv.) 90 (1794) = BLONDEA, Rich. = Sloanea, Linn. (Tiliac.).

BIONIA, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 130 = Camptosema, Hook. & Arn. (Legumin.).
acuminata, Benth. l. c. = C. acuminatum.
bella, Mart. in Walp. Rep. i. 764 = C. bellum.
coccinea, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 360 = C. coccineum.
coriacea, Benth. l. c. 130 = C. coriaceum.
marginata, Mart. ex Benth. l. c. = C. coriaceum.
nitens, Benth. l. c. = C. coccineum.
rigida, Benth. l. c. = C. coriaceum.

BIOPHYTUM, DC. Prod. i. 689 (1824). *GERANIACEAE*, Benth. & Hook. f. i. 276.
abyssinicum, *Steud. ex A Rich. Tent. Fl. Abyss.* i. 122. —Abyss.
adiantoides, *Wight, ex Edgew. & Hook. f. in Fl. Brit. Ind.* i. 487.—Ind. or.
Apodiscias, *Edgew. & Hook. f. l. c.*—Ind. or.
Candolleanum, Wight, Ill. i. 101. t. 62 = sensitivum.
castum, G. Don, Gen. Syst. i. 758 = Oxalis casta.
Cumingianum, Edgew. & Hook. f. in Fl. Brit. Ind. i. 487 = sensitivum.
Cumingii, Klotzsch, in Peters, Reise Mossamb. Bot. 85 = sensitivum.
dendroides, *DC. Prod.* i. 690.—N. Granat.
dormiens, G. Don, Gen. Syst. i. 758 = Oxalis dormiens.
fruticosum, *Blume, Bijdr.* 242.—Java.
intermedium, *Wight, Illustr.* i. 102.—Zeylan.
mimosoides, G. Don, Gen. Syst. i. 758 = Oxalis mimosoides.

BIOPHYTUM :—
nervifolium, Thw. Enum. Pl. Zeyl. 64 = sensitivum.
nudum, *Edgew. & Hook. f. Fl. Brit. Ind.* i. 488.—Zeylan.
Petersianum, Klotzsch, in Peters, Reise Mossamb. Bot. 81. t. 15 = sensitivum.
polyphyllum, *Munro, in Wight, Illustr.* i. 102.—Ind. or.
poteriocides, Edgew. ex Hook. f. Fl. Brit. Ind. i. 487 = sensitivum.
proliferum, *Edgew. & Hook. f. l. c.* 488.—Zeylan.
Reinwardtii, *Klotzsch, in Peters, Reise Mossamb. Bot.* 85.—Ax. trop.
sensitivum, *DC. Prod.* i. 690.—Cosmopol. trop.
sebanioides, Edgew. ex Hook. f. Fl. Brit. Ind. i. 487 = sensitivum.
sessile, Wall. Cat. n. 4344 = B. Apodiscias.
sommieri, G. Don, Gen. Syst. i. 758 = Oxalis sommieri.
Umbraculum, *Welw. Apont.* 590.—Afr. trop.
verticillatum, Wight, Illustr. i. 102 = intermedium.

BIOTA, D. Don, in Lamb. Pinet. ed. II. ii. 129 (1828); Endl. Conif. 46 (1847) = Thuya, Linn. (Conif.).
corsicana, Steb. ex Gord. Pinet. Suppl. 17 = T. orientalis.
dumosa, Hort. ex Carr. Conif. ed. II. 96 = T. orientalis.
cricoides, Hort. ex Carr. l. c. 141 = T. occidentalis.
excelsa, Hort. ex Gord. Pinet. Suppl. 17 = T. orientalis.
falcata, Hort. ex Carr. Conif. ed. II. 96 = T. orientalis.
Fortunei, Hort. ex Carr. l. c. 94 = T. orientalis.
frenelioides, Hort. Belg. ex Gord. Pinet. 85 = T. orientalis.
funiculata, Hort. ex Gord. l. c. ed. II. 52 = T orientalis.
glauca, Hort. ex Carr. Conif. ed. II. 100 = T. orientalis.
gracilifolia, Knight, Syn. Conif. 16 = T. orientalis.
intermedia, Hort. ex Gord. Pinet. 85 = T. orientalis.
japonica, Steb. ex Gord. l. c. 88 = T. orientalis.
macrocarpa, Hort. ex Gord. l. c. ed. II. 422 = T. orientalis.
meldensis, Laws. ex Gord. l. c. 87 = T. meldensis.
nepalensis, Endl. ex Gord. l. c. 85 = T. orientalis.
orientalis, Endl. Conif. 47 = T. orientalis.
pendula, Endl. l. c. 49 = T. orientalis.
prostrata, Hort. ex Gord. Pinet. ed. II. 401 = T. orientalis.
pyramidalis, Carr. Conif. ed. I. 96 = T. orientalis.
semperaurea, Hort.—Cf. Gard. Chron. (1874) 80 = T. orientalis.
stricta, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 205 = T. orientalis.
tatarica, Lindl. & Gord. l. c. = T. orientalis.
variegata, Hort. ex Gord. Pinet. 84 = T. orientalis.
Warreana, Hort. ex Gord. l. c. Suppl. 104 = T. orientalis.
Zuccarini, Siebold, ex Carr. Conif. ed. II. 94 = T. orientalis.

BIOTIA, Cass. in Dict. Sc. Nat. xxxiv. 308 (1825) = Madia, Molina (Compos.).
viscosa, Cass. l. c. 309 = M. viscosa.

BIOTIA, DC. Prod. v. 264 (1856) = Aster, Tourn. (Compos.).
commixta, DC. Prod. v. 265 = A. Herveyi.
corymbosa, DC. l. c. = A. corymbosus.
crenata, Steud. Nom. ed. II. i. 454, in syn. = Mairia crenata.
discolor, Maxim. Prim. Fl. Amur. 146 = A. scaber. Thunb.
glomerata, DC. Prod. v. 265 = A. macrophyllus.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 170.—Japon.
latifolia, DC. Prod. v. 264 = A. macrophyllus.
macrophylla, DC. l. c. 265 = A. macrophyllus.
Schreberi, DC. l. c. 264 = A. macrophyllus.

BIPINNULA, Comm. ex Juss. Gen. 65 (1789). *ORCHIDEAE*, Benth. & Hook. f. iii. 618.
bonariensis, *Spreng. Syst.* iii. 745.—Reg. Argent.
Commersonii, Lindl. in Quart. Journ. Roy. Inst. N. S i. (1827) 52 = bonariensis.

BIPINNULA :—
Gibérti, *Reichb. f. in Linnæa*, xli. (1877) 51.—Reg. Argent.
myriachna, *Lindl. in Hook. Journ. Bot.* i. (1834) 5.—Chili.
pinnata, *Lindl. in Quart. Journ. Roy. Inst.* N. S. i. (1827) 51.—Chili.

BIPONTINIA, Alef. in Pollichia, xxii-xxiv. (1866) 121 = **Psoralea**, Linn. (Leguminos.).
bituminosa, Alef. l. c. 123 = *P. bituminosa*.
corylifolia, Alef. l. c. 124 = *P. corylifolia*.

BIPOREIA, Thou. Gen. Nov. Madag. 14. (1806) = **Samadera**, Gaertn. (Simarub.).

BIRAMIA, Neraud, in Freyc. Voy. Bot. 28 (1826), nomen = **Maclaeanis**, Hook. (Vaccin.).
tuberosa, Auct. ex Walp. Ann. i. 478 = M. insignis.

BIRCHEA, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 66. t. 10 = **Luisia**, Gaudich. (Orchid.).
teretifolia, A. Rich. l. c. 67. t. 10 = L. tenuifolia.

BIRIS, Medic. in Staatsw. Cories. Charpf. Phys.-Oek. Ges. i. (1791) 357; et Phil. Bot. ii. 97 (1791) = **Iris**, Tourn.
sibirica, Medic. ll. cc. = I. sibirica.

BIRNBAUMIA, Kostel. Ind. Pl. Hort. Prag. (1844) 21 = **Anisacanthus**, Nees (Acanth.).
quadrifida, Kostel. l. c. = A. virgularis.

BIROLIA, Bell. in Mém. Acad. Turin, Ser. I. xviii. (1808) 408. t. 11 = **Blatine**, Linn.
palludosa, Bell. l. c. = E. triandra.

BIROLIA, Rafin. Sylva Tellur. 163 (1838) = **Clusia**, Linn. (Guttif.).
alba, Rafin. l. c. = C. alba.

BISCHOFIA, Blume, Bijdr. 1168 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 281.
MICROELUS, Wight & Arn. in Edinb. N. Phil. Journ. xix. (1835) 298.
STYLODISCUS, Benn. Pl. Jav. Rar. 138. t. 29 (1838).
Cumminghiana, Decne. in Jacquem. Voy. Bot. 153 = javanica.
javanica, *Blume, Bijdr.* 1168.—Ind. or.; Malaya; Ins. Pacif.
leptopoda, *Muell. Arg. in DC. Prod.* xv. ii. 479.—Ins. Pacif.
oblongifolia, Decne. in Jacquem. Voy. Bot. 152. t. 154 = javanica.
Roperiana, Decne. l. c. 153 = javanica.
Towi, Decne. l. c. = javanica.
trifoliata, Hook. Ic. Pl. t. 844 = javanica.

BISCUTELLA, Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 91.
DITHYREA, Harv. in Hook. Lond. Journ. Bot. iv. (1845) 77. t. 5.
IONDRABA, Medic. Pflanzeng. 27 (1792).
THLASPIDIUM, Tourn. ex Adans. Fam. ii. 422 (1763).
algeriensis, Jord. Diagn. i. 318 = didyma.
alpestris, Waldst. & Kit. Pl. Rar. Hung. iii. 258 = laevigata.
alpicola, Jord. Diagn. i. 292 = laevigata.
alsatica, Jord. l. c. 300 = laevigata.
ambigua, DC. Mon. Bisc. 9. t. 11. f. 1; Syst. ii. 415 = auriculata.
anchusaefolia, Marz. ex Reichb. Fl. Germ. Excurs. 660 = laevigata.
anchusaefolia, Tenore, ex Stend. Nom. ed. II. i. 206 = sempervirens.
angulata, Steud. l. c. = Megacarpaea angulata.
angustifolia, *Schkuhr, Handb.* iii. 285.—Europ.
apetala, Walt. Fl. Carol. 174 = Sanchiera didyma.
apricorum, Jord. Diagn. i. 307 = laevigata.
apula, Gaertn. Fruct. ii. 279 = leiocarpa.
apula, Lam. Encyc. iii. 617 = ciliata.
apula, Linn. Mant. 354 = didyma.
apula, Webb. ex Willk. & Lange, Prod. Fl. Hisp. iii. 761 = scutulata.

BISCUTELLA :—
arvernensis, Jord. Diagn. i. 298 = laevigata.
auriculata, Herb. Un. ex Steud. Nom. ed. II. i. 207 = cichoriifolia.
auriculata, *Linn. Sp. Pl.* 652.—Reg. Mediterr.
austriaca, *Bonnet, in Bull. Soc. Dauph.* fasc. vi. 222. —Europ.
balearica, L'Hérit. ex DC. Syst. ii. 643 = Succowia balearica.
balearica, Jord. Diagn. i. 295, 328 = auriculata !
bastica, Bolss. & Reut. Diagn. Ser. II. i. 49 = laevigata.
Bourgaei, Jord. Diagn. i. 320 = lyrata.
brevicaulis, Jord. l. c. 303 = laevigata.
Burseri, Jord. l. c. 326 = auriculata.
californica, *Benth. & Hook. f. ex S. Wats. Bot. Calif.* i. 48.—Calif.
Candollii, Jord. Diagn. i. 326 = auriculata.
chilensis, Lag. ex DC. Mon. Bisc. 6 = Cremolobus chilensis.
Chouletti, Jord. Diagn. i. 319 = didyma.
cichoriifolia, *Lousel. Add. Fl. Gall.* 167.—Mont. Fyren.
ciliata, *DC. Mon. Bisc.* 6; *Syst.* ii. 410.—Ital.; Hispan.
collina, Jord. Diagn. i. 299 = laevigata.
Columnae, Tenore, Prod. Fl. Nap. 38 = didyma.
confusa, *Pomel, Nouv. Mat. Fl. Alt.* 331.—Afr. bor.
controversa, Bor. Fl. Centr. Fr. ed. III. 56 = laevigata.
coronaria, *Ehrenb. ex Willd. in Ind. Sem. Horpat. Adn.* (1868).—Hab. ?
coronopifolia, Willd. Sp. Pl. iii. 474 = ciliata.
coronopifolia, Linn. Mant. ii. 356; All. Fl. Pedem. i. 247 = laevigata.
cuneata, Lag. Cat. Hort. Madr. (1816) 20 = Cremolobus chilensis.
depressa, Thom. ex DC. Syst. ii. 415 = laevigata.
depressa, Willd. Enum. Hort. Berol. 678 = didyma.
didyma, *Linn. Sp. Pl.* 653.—Reg. Mediterr.; Persia.
didyma, Scop. Fl. Carn. ed II. ii. 14 = laevigata.
didyma, Willd. Enum Hort. Berol. 678 = ciliata.
dilatata, Vis. Stirp. Dalm. 14 = cichoriifolia.
divisocarpa, Jord. Diagn. i. 305 = laevigata.
dubia, Steud. Nom. ed. II. i. 207 = Cremolobus linearifolius.
erigerifolia, DC. Mon. Bisc. 4; Syst. ii. 408 = auriculata.
ericarpa, DC. Mon. Bisc. 7. t. 9. f. 2; Syst. ii. 411 = didyma.
ernstiana, *Reichb. Ic. Fl. Germ.* vii. 5.—Europ.
flexuosa, Jord. Diagn. i. 300 = laevigata.
frutescens, *Coss. Notes Crit.* ii. 97.—Hispan.
glabra, [Clairv.] Man. Herb. 216 = laevigata.
glacialis, Jord. Diagn. i. 310 = laevigata.
glareosa, Jord. l. c. 304 = laevigata.
granitica, Bor. ex Perar. in Bull. Soc. Bot. Fr. xvi. (1869) 353 = laevigata.
Guilloni, Jord. Diagn. i. 309 = laevigata.
hispida, DC. Mon. Bisc. 4. t. i. f. 1; Syst. ii. 408 = cichoriifolia.
incana, *Tenore, Fl. Neap. Prod. App.* 5te. (1826).—Italia.
intermedia, Gouan, Illustr. 42 = laevigata.
intricata, Jord. Diagn. i. 308 = laevigata.
laciniata, Willd. ex DC. Syst. ii. 417 = Megacarpaea laciniata.
laevigata, *Linn. Mant.* 235.—Reg. Medit.
laevigata, Sibth. & Sm. Fl. Graec. Prod. ii. 16 = didyma.
Lamarckii, Jord. Diagn. i. 323 = auriculata.
Lamottei, Jord. l. c. 302 = laevigata.
laxa, *Boiss. & Reut. Diagn.* Ser. II. i. 48.—Hispan.
laxiflora, Presl, ex Spreng. Syst. iv. Cur. Post. 239 = didyma.
leiocarpa, *DC. Mon. Bisc.* 8; *Syst.* ii. 413.—Orient.
Linae, Reichb. Fl. Germ. Excurs. 661, sub n. 4306 = laevigata.
longa, Dalac, Fl. Hautes-Pyr. 190 = cichoriifolia.
longifolia, Vill. Prosp. 38 = laevigata.
lucida, Balb. ex DC. Mon. Bisc. 9. t. 7; Syst. ii. 414 = laevigata.
lusitanica, Jord. Diagn. i. 315 = laevigata.
lyrata, Linn. Mant. 354 = didyma.
macrocarpa, Hort. ex DC. Syst. ii. 409 = cichoriifolia.
macrocarpa, *Bieb. ex Tenacher, in Mém. Soc. Nat. Mosc.* iv. (1812) 283.—Transsyl.
major, Schkuhr, Handb. ii. 283. n. 1822 = auriculata.
marginata, Tenore, Prod. Fl. Nap. 38 = didyma.
maritima, Tenore, l. c. = didyma.
mauritanica, Jord. Diagn. i. 325 = auriculata.

BISCUTELLA :—
maxima, Pall. ex DC. Syst. ii. 417 = Megacarpaea laciniata.
mediterranea, Jord. Diagn. i. 318 = laevigata !
Megacarpaea, Boiss. & Reut. Diagn. Ser. II. i. 44 = laevigata.
megalocarpa, Fisch. ex DC. Mon. Bisc. 5. t. 3 = Megacarpaea laciniata.
microcarpa, DC. l. c. 7; Syst. ii. 411 = didyma.
minor, Jord. Diagn. i. 297 = laevigata.
mollis, Loisel. Not. 168 = laevigata.
montana, Cav. Ic. ii. 59. t. 177 = laevigata.
neustriaca, Ed. Bonnet, in Act. Soc. Dauph. (1879) 196 = laevigata.
nicaensis, Jord. Diagn. i. 314 = laevigata ?
obcordata, Reichb. lc. Fl. Germ. vii. 8. f. 615 = laevigata.
obovata, *Hort. Par. ex DC. Mon. Bisc.* 8; *Syst.* ii. 413 —Hab. ?
orcelitana, Lag. ex DC. Syst. ii. 408 = auriculata.
oreites, Jord. Diagn. i. 294 = laevigata.
orientalis, [Desf.] ex DC. Mon. Bisc. 8 = leiocarpa.
patulipes, Jord. Diagn. i. 320 = didyma.
perennis, Spach, Hist. Vég. Phan. vi. 571 = ciliata, laevigata, sempervirens.
peruviana, Lam. Encyc. iii. 620 = Cremolobus peruvianus.
petraea, Jord. Diagn. i. 306 = laevigata.
picridifolia, Lapeyr. Hist. Abr. Pl. Pyr. 375 = cichoriifolia.
pinnatifida, Hort. ex Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 302 = laevigata !
pinnatifida, Jord. Diagn. i. 311 = praec. !
polycicada, Jord. l. c. 312 = laevigata.
pratensis, *Timb. in Bull. Soc. Hist. Nat. Toul.* iii. (1869) 114.—Gallia.
pumila, Balb. ex DC. Mon. Bisc. 6; et Syst. ii. 411 = didyma.
pyrenaica, *Huet, in Ann. Sc. Nat. Sér. III.* xix. (1853) 252.—Hispan.
radicata, *Coss. & Dur. in Bull. Soc. Bot. Fr.* xix. (1872) 234.—Afr. bor.; Hispan.
radicosa, Willk. & Lange, Prod. Fl. Hisp. iii. 778 = radicata.
raphanifolia, Poir. Voy. Barb. ii. 198 = didyma.
rosularis, *Boiss. & Reut. Diagn.* Ser. II. i. 45 — Hispan.
rupestris, *Ehrh. Beitr.* v. 177, nomen.—German.
saxatilis, Schleich. ex DC. Prod. i. 188 = laevigata.
saxatilis, Endr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 765 = pyrenaica.
scabra, Salisb. Prod. 368 = didyma.
sclerocarpa, *Revel, in Bull. Soc. Bot. Fr.* xxv. (1878) *Bibl.* 78.—Gallia.
scutulata, *Boiss. & Reut. Diagn.* Ser. II. i. 41.—Hispan
secunda, Jord. Diagn. i. 310 = laevigata.
sempervirens, DC. Syst. ii. 416 = laxa.
sempervirens, *Linn. Mant.* 355.—Hispan.; Lusitan.
sinuata, Jord. Diagn. i. 301 = laevigata.
spathulata, DC. Prod. i. 188, in syn. = laevigata.
stenophylla, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 299 = laevigata.
stricta, Jord. Diagn. i. 312 = laevigata.
subpatulata, Lam. Encyc. iii. 620 = laevigata.
suffruticosa, *Coss. Notes Crit.* 97.—Hispan.
suffruticosa, DC. Mon. Bisc. 5. t. 5 = Cremolobus suffruticosus.
tergestina, Jord. Diagn. i. 295 = laevigata.
tenuicaulis, Jord. l. c. 310 = laevigata.
tomentosa, Lag. ex DC. Syst. ii. 416 = laevigata.
tumidula, Lag. ex DC. l. c. 411 = didyma.
varia, Dum. Fl. Belg. Prod. 114 = laevigata.
variabilis, Loisel. Fl. Gall. ii. 56 = laevigata.
variegata, Boiss. & Reut. Diagn. Ser. II. i. 44 = laevigata.
verna, Math. ex Nym. Consp. 59 = laevigata.
virgata, Jord. Diagn. i. 309 = laevigata.
Wiulzeni, *Benth. & Hook. f. ex S. Wats. Bot. Calif.* i. 483.—Am. bor. occ.

BISERRULA, Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 507.
PELECINUS, Tourn. ex Medic. in Vorles. Churpf. Phys. Oec. ii. (1787) 378.
leiocarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 194 = B. Pelecinus.
pelecina, Linn. Fl. Fr. ii. 684 = B. Pelecinus.
Pelecinus, *Linn. Sp. Pl.* 762.—Reg. Mediterr.

BISMALVA, Medic. Malv. 39 (1787) = **Malva**, Tourn.
Alcea, Medic. l. c. = M. Alcea.
fastigiata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 260 = M. Alcea.
laciniata, Fourr. l. c. = M. moschata.
moschata, Medic. Malv. 39 = M. moschata.
Tournefortiana, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195 = M. Tournefortiana.

BISMARCKIA, Hildebr. & H. Wendl. in Bot. Zeit. xxxix. (1881) 89, 90, 93. *PALMAE*, Benth. & Hook. f. Gen. iii. 882.
nobilis, *Hildebr. & H. Wendl. l. c.* 94.—Madag.

BISPHAERIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1 (Gen. dub.).

BISTANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 64 (1827) (Gen. dub.).

BISTELLA, Adans. Fam. ii. 226 (1763) (an Rubiaceæ ?).

BISTELLA, Delile, in Caill. Voy. Méroé (Cent. Pl. Afr.) 97, t. 63 (1826) = **Vahlia**, Thunb. (Saxifr.).
geminiflora, Caill. & Delile, Voy. Méroe, 97, t. 2. f. 2 = V. viscosa.

BISTORTA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 115 (1737) = **Polygonum**, Tourn.
americana, Rafin. Fl. Tellur. iii. 19 = P. viviparum ?
major, S. F. Gray, Nat. Arr. Brit. Pl. ii. 267 = P. Bistorta.
officinalis, Rafin. Fl. Tellur. iii. 12 = B. Bistorta.
vivipara, S. F. Gray, Nat. Arr. Brit. Pl. ii. 268 = P. viviparum.

BISTROPOGON, Auct. ex Steud. Nom. ed. II. i. 207. (1840) = **Bysteropogon**, L'Hérit. (Labiat.).

BISWARRA, Cogn in Comptes-rend. Soc. Bot. Belg. xxi. (1882) 16. *CUCURBITACEAE*.
Warra, C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 227.
tonglensis, *Cogn. l. c.*—Reg. Himal.

BIVINIA, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 78. *SAMYDACEAE*, Benth. & Hook. f. i. 800.
Jaubertia, *Tul. l. c.*—Madag.

BIVONAEA, DC. Syst. ii. 554 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 88.
lutea, *DC. Syst. ii.* 555.—Sicily.
praecox, *Bertol. Fl. Ital.* x. 520.—Sicil.
Savinna, *Carusel, Prod. Fl. Tosc.* 47.—Ital.

BIVONAEA, Moç. & Sesse, ex DC. Prod. iii. 372 (1828) = **Pentacosaea**, Bartl. (Illecebr.).
multicaulis, Moç. & Sesse, ex DC. l. c. 378 = P. polycnemoides.

BIVONEA, Rafin. Fl. Ludov. 138 (1817) = **Jatropha**, Linn. (Euphorb.).
stimulosa, Rafin. l. c. = J. urens.

BIVONIA, Spreng. Neue Entdeck. ii. 116 (1821) = **Bernardia**, P. Br. (Euphorb.).
axillaris, Spreng. l. c. = B. axillaris.

BIWALDIA, Scop. Introd. 282 (1777) = **Garcinia**, Linn. (Guttif.).

BIXA, Linn. Coroll. Gen. 8 (1737). *BIXINEAE*, Benth. & Hook. f. i. 125.
Urucu, Adans. Fam. ii. 331 (1763).
acuminata, Boj. Hort. Maurit. 20 = Orellana.
americana, Poir. Encyc. vi. 229 = Orellana.
odorata, Rniz & Pav. Fl. Per. v. t. 460, ex G. Don, Gen. Syst. i. 295 = Orellana !
Orellana, *Linn. Sp. Pl.* 512.—Am. austr.
orleana, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 8 = Orellana.
platycarpa, Ruiz & Pav. Fl. Per. v. t. 459, ex G. Don, Gen. Syst. i. 294 = Orellana.
purpurea, Sweet, Hort. Brit. ed. I. 83; G. Don, Gen. Syst. i. 295 = Orellana !
sphaerocarpa, *Triana, in Bull. Soc. Bot. Fr.* v. (1858) 849.—N. Granad.
tinctoria, Salish. Prod. 369 = Orellana.
Urucurana, Willd. Enum. Hort. Berol. 565 = Orellana.

HIXAGREWIA, Kurz, in Journ. Bot. xiii. (1875) 325 = **Trichospermum**, Blume (Tiliac.).
nicobarica, Kurz, l. c.—Ins. Nicobar.

BLABEROPUS, A. DC. Prod. viii. 410 (1844) = **Alstonia**, R. Br. (Apocyn.).
lucidus, A. DC. l. c. 411 = Trachelospermum fragrans.
neriifolius, A. DC. l. c. = A. neriifolia.
Sebusi, Heurck & Muell. Arg. in Heurck, Obs. Bot. ii. 188 = A. neriifolia.
sericeus, A. DC. Prod. viii. 411 = A. sericea.
venenatus, A. DC. l. c. = A. venenata.
villosus, Miq. Fl. Ind. Bat. ii. 440 = A. villosa.

BLACHIA, Baill. Etud. Gén. Euph. 388. t. 19 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 301.
calycina, *Benth. in Journ. Linn. Soc.* xvii. (1880) 296.—Ind. or.
denudata, *Benth. l. c.*—Ind. or.
Pentzii, *Benth. l. c.*—China.
reflexa, *Benth. l. c.*—Ind. or.
umbellata, *Baill. Etud. Gin. Euph.* 387. t. 19. f. 18–20. —Ind. or.

BLACKBOURNEA, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 356 = seq.

BLACKBURNIA, Forst. Char. Gen. t. 6 (1776) = **Zanthoxylum**, Linn. (Rutac.).
monadelpha, Roxb. Hort. Beng. (88); Fl. Ind i. 415 = Dolsera Roxburghii.
oppositifolia, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 191 = Dolsera Roxburghii.
pinnata, Forst. Char. Gen. t. 6 = Z. Blackburnia.

BLACKIA, Schrank, ex DC. Prod. iii. 165 (1828) = **Myriaspora**, DC. (Melastom.).
egensis, Schrank, ex DC. l. c. = M. egensis.

BLACKSTONIA, Huds. Fl. Angl. ed. I. 146 (1762), nomen prius = **Chlora**, Ren. ex Adans. Fam. ii. 508 (1763).
perfoliata, Huds. l. c. = C. perfoliata.

BLACKSTONIA, A. Juss. in Orbign. Dict. vi. 482 (1849) = BLAKSTONIA, Scop. = **Moronobea**, Aubl. (Guttif.).

BLACKWELLIA, Gaertn. Fruct. ii. t. 117 (1791) (Blakwellia, p. 169) (Genus dubium).
antarctica, Gaertn. l. c.—Hab.?

BLACKWELLIA, Comm. ex Juss. Gen. 343 (1789) (Blakwellia) = **Homalium**, Jacq. (Samyd.).
acuminata, Wall. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 596 = H. nepalense.
africana, Hook. f. in Hook. Niger Fl. 361 = H. africanum.
austro-caledonica, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 107 = H. austro-caledonicum !
axillaris, Lam. Encyc. i. 428 = H. villare.
brachybotrya, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 48 = H. brachybotrys.
brachystylis, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 61.—Madag.
caryophyllacea, Zoll. & Mor. Syst. Verz. Zoll. 33 = H. caryophyllaceum.
cerasifolia, Vent. Choix, 56. t. 56.—Madag.
ceylanica, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 452 = H. zeylanicum.
chinensis, Lindl. ex Steud. Nom. ed. II. i. 306 = H. fagifolium.
cochinchinensis, Blume, Mus. Bot. Lugd. Bat. ii. 27.—Cochinchina.
dasycnetha, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 610 = H. Griffithianum.
decurrens, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 106.—N. Caled.
denticulata, Harv. in Harv. & Sond. Fl. Cap. ii. 585.—Afr. austr.
Deplanchei, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 110.—N. Caled.
erianthe, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 62.—Madag.
fagifolia, Lindl. in Trans. Hort. Soc. Lond. vi. (1826) 389 = H. fagifolium.
foetida, Wall. Cat. n. 4899 = H. foetidum.

BLACKWELLIA :—
Jussieua, J. F. Gmel. Syst. 768 = Homalium paniculatum !
glauca, Vent. Choix, 55. t. 55 = H. paniculatum.
gracilis, Blume, Mus. Bot. Lugd. Bat. ii. 26 = H. axillare.
grandiflora, Lindl. ex Loud. Hort. Brit. 193 = H. fagifolium.
Guillaini, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 106.—N. Caled.
integrifolia, Lam. Encyc. i. 428 = H. paniculatum.
intermedia, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 109.—N. Caled.
kanalensis, Vieill. l. c. 112.—N. Caled.
longiflora, Miq. Fl. Ind. Bat. i. t. 715 = H. corallanthus.
Lowrea, Benth. in Hook. Lond. Journ. Bot. i. (1842) 483 = H. fagifolium.
macrostachya, Turcz. in Bull. Soc. Nat. Mosc. (1863) 610 = H. longifolium.
micrantha, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 63 = H. micranthum.
Mathieuana, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 108.—N. Caled.
molaccana, Blume, Mus. Bot. Lugd. Bat. ii. 27.—Ins. Moluccæ.
monostachya, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 610.—Penang.
montana, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 110.—N. Caled.
nepaulensis, DC. Prod. iii. 54 = seq.
nepalensis, Wall. Pl. As. Rar. t. 179 = H. nepalense.
padifera, Lindl. Bot. Reg. t. 1308 = H. fagifolium.
padifolia, Steud. Nom. ed. II. i. 208 = H. fagifolium.
paniculata, Lam. Encyc. i. 428 = H. paniculatum.
planiflora, Boiv ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 64.—Madag.
polystachya, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 107.—N. Caled.
propinqua, Wall. Cat. n. 4896 = H. propinquum.
pubiflora, Walp. Rep. i. 548 = H. fagifolium.
purpurea, Steud. Nom. ed. II. i. 308, sphalm. = H. propinquum.
rivularis, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 111.—N. Caled.
rubiginosa, Vieill. l. c. 108.—N. Caled.
rufescens, Presl, Bot. Bemerk. 42 = H. rufescens.
spiralis, Wall. in As. Res. xiii. (1820) 400 = H. tomentosum.
spiralis, Wall. Cat. n. 4897, partim = H. propinquum.
tetrandra, Wight, Ic. v. t. 1851 = H. zeylanicum.
Thouarsiana, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 60.—Madag.
tomentosa, Vent. Choix, 57. t. 57 = H. tomentosum.

BLADHIA, Thunb. Nov. Gen. 6 (1781), nomen prius = **Ardisia**, Sw. (1788) (Myrsin.).
crispa, Thunb. Fl. Jap. 97 = A. crenata, *Roxb.*
glabra, Thunb. in Trans. Linn. Soc. ii. (1794) 331 = A. japonica.
japonica, Blanco, Fl. Filip. ed. I. 126 = A. pyramidalis.
japonica, Thunb. Nov. Gen. 7 = A. japonica.
villosa, Thunb. Fl. Jap. 96 = A. pusilla.

BLAERIA, Linn. Coroll. Gen. 3 (1737). *ERICACEAE*, Benth. & Hook. f. ii. 592.
Kolbia, Adans. Fam. ii. 164 (1763).
aggregata, Wendl. ex Steud. Nom. ed. II. i. 208; ii. 589 = Simochellus multiflorus.
articulata, Linn. Mant. ii. 198 (= *Syndesmanthus articulatus*).—Afr. austr.
articulata, Willd. ex Steud. Nom. ed. II. i. 208 = Sympiesa capitellata.
barbigera, *G. Don, Gen. Syst.* iii. 805.—Afr. austr.
bicolor, Klotzsch, in Linnaea, viii. (1833) 660 (err. typ. 606) = Simochellus bicolor.
bractosta, Wendl. Collect. ii. 1. t. 37 = Sympiesa capitellata.
brunneifolia, G. Don, Gen. Syst. iii. 805 = Erica brunneifolia.
cadmea, *Thunb. Diss. Blaeria*, 10.—Afr. austr.
campanulata, *Benth. in DC. Prod.* viii. 698.—Afr. austr.
carnea, Klotzsch, in Linnaea, viii. (1833) 661 = Simochellus carneus.
ciliaris, Linn. f. Suppl. 122 = Griesbachia ciliaris.

BLAERIA :—
ciliciflora, G. Don, Gen. Syst. iii. 805 = Grisebachia velleriflora.
coccinea, *Klotzsch, in Linnaea*, viii. (1833) 657.—Afr. austr.
condensata, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 13 = spicata.
depressa, Drège, ex Benth. in DC. Prod. vii. 706 = Sympiesa Kunthii.
depressa, Lichst. ex Roem. & Schult. Syst. iii. 168 = Simochellus depressus.
dumosa, Wendl. Collect. ii. 8. t. 38 = purpurea.
equisetifolia, G. Don, Gen. Syst. iii. 805 = purpurea.
eriantha, Willd. ex Steud. Nom. ed. II. i. 908 (= *Syndesmanthus articulatus*).—Afr. austr.
ericoides, *Linn Sp. Pl.* 162.—Afr austr.
fasciculata, Willd. Sp. Pl. i. 629 = Simochellus glabellus.
fastigiata, *Benth. in DC. Prod.* vii. 697.—Afr. austr.
flexuosa, *Benth. l. c.* 698.—Afr. austr.
floccifera, G. Don, Gen. Syst. iii. 805 (= *Syndesmanthus pauciflorus*).—Afr. austr.
fuscescens, *Klotzsch, in Linnaea*, viii. (1833) 657.—Afr. austr.
glabella, Drège, ex Benth. in DC. Prod. vii. 698 = campanulata.
glabella, Willd. Sp. Pl. i. 631 = Simochellus glabellus.
glabra, *Thunb. Diss. Blaeria*, 10.—Afr. austr.
gracilis, Bartl. in Linnaea, vii. (1832) 650 = Sympiesa Kunthii.
hirsuta, Lichst. in Roem. & Schult. Syst. iii. 170 (= *Syndesmanthus pauciflorus*).—Afr. austr.
hirsuta, *Thunb. Diss. Blaeria*, 8.—Afr. austr.
incana, Bartl. in Linnaea, vii. (1832) 650 = Grisebachia incana.
jucunda, Reichb. ex Steud. Nom. ed. II. i. 908 = purpurea.
Krauseana, *Klotzsch, in Walp. Rep.* ii. 728.—Afr. austr.
leucocephala, *Bory, Voy.* iii. 172.—Ins. Borbon.
multiflora, Klotzsch. in Linnaea, viii. (1833) 661 = Simochellus multiflorus.
muscosa, Ait. Hort. Kew. i. 150 = Scyphogyne inconspicua.
nodiflora, G. Don, Gen. Syst. iii. 805 = Grisebachia plumosa.
nudiflora, *Thunb. Diss. Blaeria*, 6.—Afr. austr.
paniculata, *Thunb. l. c.* 10.—Afr. austr.
parviflora, Klotzsch, in Linnaea, viii. (1833) 665 (= *Codonanthemum parviflorum*).—Afr. austr.
pauciflora, G. Don, Gen. Syst. iii. 804, sphalm. = *paucifolia*.
paucifolia, Wendl. Collect. ii. 17. t. 48 (= *Syndesmanthus pauciflorus*).—Afr. austr.
pilosa, E. Mey. in Drège, ex Benth. in DC. Prod. vii. 701 = Grisebachia hispida.
pubescula, Klotzsch, in Linnaea, viii. (1833) 661 (= *Codonanthemum puberulum*).—Afr. austr.
purpurea, Berg. Descr. Pl. Cap. 24 = Simochellus glabellus.
purpurea, *Linn. f. Suppl.* 122.—Afr. austr.
pusilla, *Klotzsch, in Linnaea*, viii. (1833) 659.—Afr. austr.
pusilla, *Linn. Mant.* 139 ('sp. dubia').—Afr. austr.
revoluta, Bartl. in Linnaea, vii. (1832) 650 = barbigera.
rubra, Hort. ex Stend. Nom. ed. II. i. 908 = ericoides.
scabra, Drège, ex Benth. in DC. Prod. vii. 698 = ericoides.
scabra, Willd. Sp. Pl. i. 629 (= *Syndesmanthus scabra*).—Afr. austr.
serrata, *Thunb. Diss. Blaeria*, 6.—Afr. austr.
sessiliflora, Wendl. ex Steud. Nom. ed. II. i. 908 = Simochellus multiflorus.
spicata, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 13.—Abyss.
Thunbergii, G. Don, Gen. Syst. iii. 805 = Grisebachia ciliaris.
trigyna, Wendl. ex Steud. Nom. ed. II. i. 908 = fuscescens.
turmalis, G. Don, Gen. Syst. iii. 805 = Erica turmalis.
serunthemifolia, G. Don, l. c. = Eremia lanata.

BLAINVILLEA, Cass. in Dict. Sc. Nat. xxix. 493 (1823). *COMPOSITAE*, Benth. & Hook. f. ii. 369.
CALYPTOCARPUS, Less. Syn. Comp. 231 (1832).
EISENMANNIA, Sch. Bip. in Flora, xxv. (1842) ii. 418.

BLAINVILLEA, Galophthalmum, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 7. t. 2.
OLIGOGYNE, DC. Prod. v. 699 (1837).
UCACEA, Cass. in Dict. Sc. Nat. xxvii. 9 (1823).
alba, Edgew. in Trans. Linn. Soc. xx. (1851) 70 = rhomboidea.
amazonica, *Benth. & Hook. f. Gen.* ii. 370.—Reg. Amazon.
bahiensis, *Baker, in Mart. Fl. Bras.* vi. iii. 177.—Bras.
blariasta, *DC. Prod.* v. 493.—Bras.
Gayana, Benth. in Hook. Niger Fl. 484, in syn. = Aspilia Gayana.
Gayana, *Cass. in Dict. Sc. Nat.* xlvii. 90.—Afr. trop.
hispida, Edgew. in Trans. Linn. Soc. xx. (1851) 70 = rhomboidea.
lanceolata, *Baker, in Mart. Fl. Bras.* vi. iii. 176.—Bras.
latifolia, DC. in Wight, Contrib. Bot. Ind. Or. 71 = rhomboidea.
polycephala, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 89 = rhomboidea.
Prieureana, Benth. in Hook. Niger Fl. 483, in syn. = Aspilia Prieureana.
Prieureana, *DC. Prod.* v. 492.—Afr. trop.
racemosa, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 89 = rhomboidea.
rhomboides, *Cass. in Dict. Sc. Nat.* xxix. 493.—Cosmop. trop.; Ind. or.
synrefolioides, *Benth. & Hook f. Gen.* ii. 370.—Hab.? tampicana, *Benth. & Hook. f. c.*—*Cf. Hemsl. Biol. Centr. Am. Bot.* ii. 169.—Mexic.
tenuicaulis, *Benth. & Hook. f. Gen.* ii. 370.—Ins. Galap.

BLAIRIA, Houst. ex Linn. Gen. ed. I. 334, in syn. (1737) = Evea, Adans. (Verben.).
cordifolia, Moench, Meth. 493 = P. hispida.
javanica, Garcin. Fruct. i. t. 56 = Lippia javanica.
lappulacea, Stend. Nom. ed. II. i. 908 = P. echinata.
maritima, Garcin. Fruct. i. 265. t. 56. f. 1 = P. hispida.
nodiflora, Gaertn. l. c. 266. t. 56 = Lippia nodiflora.

BLAIRIA, Gled. in Mém. Acad. Berl. (1751) 125 = **Blaeria**, Linn. (Ericac.).

BLAKBURNIA, J. F. Gmel. Syst. 258 (1791) = BLACKBURNIA, Forst. = Xanthoxylum, Linn. (Rutac.).

BLAKEA, P. Br. Hist. Jamaic. 238 (1756). *MELASTOMACEAE*, Benth. & Hook. f. i. 770.
PYXIDANTHUS, Nand. in Ann. Sc. Nat. Sér. III. xviii. (1852) 150. t. 6.
VALDESIA, Ruiz & Pav. Prod. 67 (1794).
brevipes, Nand. in Ann. Sc. Nat. Sér. III. xviii. (1852) 144 = repens.
Cacatin, *D. Don, in Mem. Wern. Soc.* iv. (1823) 324. —Guiana.
caudata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 148.—N. Granat.
glaberrima, *Benth. Bot. Voy. Sulph.* 94.—Peruv.
gracilis, *Hemsl. Diag. Pl. Nov.* 13.—Costa Rica.
granatensis, *Naud. in Ann. Sc. Nat. Sér.* III. xviii. (1852) 143.—N. Granat.
grandiflora, *Hemsl. Diag. Pl. Nov.* 13.—Costa Rica.
hirsuta, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 148.—Bras.
laevigata, D. Don, in Mem. Wern. Soc. iv. (1823) 327 = Topobea laevigata.
latifolia, *D. Don, l. c.* 323.—Peruv.
laurifolia, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 143 = pulverulenta.
Lindeniana, *Triana in Trans. Linn. Soc.* xxviii. (1871) 149.—Venezuela.
macrophylla, D. Don, in Mem. Wern. Soc. iv. (1823) 326 = Bellucia superba.
mexicana, D. Don, l. c. 325 = repens.
multiflora, D. Don, l. c. 326 = Topobea multiflora.
ovalis, *D. Don, l. c.* 326.—Peruv.
ovalis, Griseb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 148 = repens.
parasitica, D. Don, in Mem. Wern. Soc. iv. (1823) 327 = Topobea parasitica.
podagrica, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 149.—N. Granat.
pulverulenta, DC. Prod. iii. 196 = Conostegia subhirsuta.

BLAKEA, pulverulenta, *Vahl, Symb. Bot.* iii. 61.—Am. austr.
pyxidanthus, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 149.—N. Granat.
quadrangularis, *Triana, l. c.*—N. Granat.
quinquenervia, Aubl. Pl. Gui. i. 325. t. 210 = Bellucia Aublettii.
repens, *D. Don, in Mem. Wern. Soc.* iv. (1823) 325. —Peruv.
repens, Griseb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 149 = ovalis.
rosea, *D. Don, in Mem. Wern. Soc.* iv. (1823) 325. —Peruv.
rostrata, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 148.—Peruv.
rotundifolia, *D. Don, in Mem. Wern. Soc.* iv. (1823) 326.—Peruv.
Schlimii, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 149.—Venezuela.
sessiliflora, *Naud. in Ann. Sc. Nat. Sér.* III. xviii (1852) 144.—Peruv.
sessilifrons, Poepp. ex Triana, in Trans. Linn. Soc xxviii. (1871) 148 = hirsuta.
subconnata, *O. Berg, ex Triana, l. c.*—Ecuador.
trinervia, *Linn. Sp. Pl.* ed. II. 635.—Jamaica.
trinervia, Pav. ex Triana, in Trans. Linn. Soc. xxviii (1871) 149 = Topobea laevigata.
trinervis, Pav. ex D. Don, in Mem. Wern. Soc. iv (1823) 327 = Topobea laevigata.
trijuinervia, Linn. Amoen. Acad. viii. 257; Pl Surinam. n. 59 = Bellucia Aublettii.

BLAKSTONIA, Scop. Introd. 276 (1777) = **Moronobea**, Aubl. (Guttif.).

BLAKWELLIA, Comm. ex Juss. Gen. 348 (1789) (Blackwellia, ex J. F. Gmel. Syst. 768 (1791)) = **Homalium**, Jacq. (Samyd.).

BLAKWELLIA, Gaertn. Fruct. ii. 170 (1791) (Blackwellia, t. 117) (Genus dubium).

BLAKWELLIA, Scop. Introd. 326 (1777) = **Aralia**. Toxm.

BLANCHEA, Boiss. Fl. Orient iii. 212, in syn. (1875) = Xphiona, Cass. (Compos.).
iphimoidea, Boiss. l. c.—Syria.

BLANCHETIA, DC. Prod. v. 75 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 296.
heterosticha, *DC. l. c.*—Bras.

BLANCKIA, Neck. Elem. i. 347 (1790) = **Conobea**, Aubl. (Scrophul.).

BLANCOA, Blume, Rumphia, ii. 128 (1836) = **Didymosperma**, H. Wendl. (Palm.).

BLANCOA, Blume, Rumphia, iii. 181 (1847) = **Harpullia**, Roxb. (Sapind.).
arborea, Blume, l. c. = H. cupaniodes.

BLANCOA, Lindl. Swan Riv. App. 45 (1839). *HAEMODORACEAE*, Benth. & Hook. f. iii. 676.
canescens, *Lindl. l. c.*—Austral.

BLANDFORDIA, Andr. Bot. Rep. t. 348 (1804) = **Gaisa**, Linn. (Diapens.).
caudata, Stend. Nom. ed. II. i. 209, sphalm. = seq.
cordata, Andr. Bot. Rep. t. 348 = G. sphylla.

BLANDFORDIA, Sm. Exot. Bot. i. 5. t. 4 (1804). *LILIACEAE*, Benth. & Hook. f. iii. 774.
aurea, *Hook. f. Bot. Mag.* t. 5809.—Austral.
Backhousii, Lindl. Bot. Reg. (1844) sub t. 18 = grandiflora.
Cunninghamii, Lindl. l. c. = grandiflora.
Cunninghamii, Van Houtte, Fl. des Serres, t. 1339 = flammea.
flammea, *Lindl. in Journ. Hort. Soc.* v. (1850) 22.—Austral.
grandiflora, *R. Br. Prod.* 296.—Tasmania.
grandiflora, Hook. f. Fl. Tasm. ii. 49 = marginata.
grandiflora, Lindl. Bot. Reg. t. 924 = nobilis?
intermedia, *Herb. in Lindl. Bot. Reg.* (1846) *Misc.* 64. —Tasmania.
marginata, *Herb. l. c.* (1842) *Misc.* 64.—Austral.

BLANDFORDIA :—
noblilis, *Sm. Exot. Bot.* i. 5. t. 4.—Austral.
Princeps, W. G. Smith, in Flor. Mag. N. S. (1875) t. 170 = flammea.
punicea, Sweet, Hort. Brit. ed. II. 517 = grandiflora.
speciosa, Hort. Angl. ex Link, Enum. Hort. Berol. i. 330 = grandiflora!
uniflora, Willd. ex Kunth, Enum. Pl. iv. 478 = Milla biflora.

BLANDFORTIA, Poir. Encyc. Suppl. iv. 116 (1814) = **Blandfordia**, Sm. (Liliac.).

BLANDINA, Rafin. Fl. Tellur. iii. 88 (1836) = **Leucas**, Burm. (Labiat.).
biflora, Rafin. l. c. = L. biflora.
chinensis, Rafin. l. c. = L. chinensis.

BLANDOWIA, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 10. t. 4. f. 2 = **Apinagia**, Tul (Podost.).
Preissii, Wedd. in DC. Prod. xvii. 88 = A. Preissii.
striata, Corda, in Sturm, Deutschl. Fl. Abth. ii. Heft v–vi. p. 107. t. 38 = A. Preissii.
striata, Lehm ex Wedd. in DC. Prod. xvii. 71 = Dicraea Wallichii.
strata, Willd. in Ges Naturf. Fr. Berl. Mag. iii. (1809) 100. t. 4 = A. Willdenowii.

BLANISIA, Pritz. Ind. Ic. 164 (1855), sphalm. = POLANISIA, Rafin. = **Cleome**, Linn. (Cappar.).

BLASTANIA, Kotschy & Peyr. Pl. Tinn. 15. t. 7 (1865–66). *CUCURBITACEAE*.
CTENOLEPIS, Hook. f. in Benth. & Hook. f. Gen. i. 832 (1867).
CTENOPSIS, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 12.
fimbristipula, *Kotschy & Peyr. l. c.*—Afr. trop.
Garcini, *Cogn. in DC. Monog. Phan.* iii. 630.—Ind. or.

BLASTEMANTHUS, Planch. in Hook. Lond. Journ. Bot. v. (1846) 644. *OCHNACEAE*, Benth. & Hook. f. i. 320.
gemmiflorus, *Planch. l. c.* 645.—Bras.
grandiflorus, *Spruce, ex Engl. in Mart. Fl. Bras.* xii. ii. 866.—Bras.

BLASTOTROPHE, F. Didrichs. in Kjoeb. Vidensk. Meddel. (1854) 195 = **Alafia**, Thou. (Apocyn.).
scandens, F. Didrichs. l. c. 198.—Afr. trop.

BLASTUS, Lour. Fl. Cochinch. 526 (1790). *MELASTOMACEAE*, Benth. & Hook. f. i. 769.
cochinchinensis, *Lour. l. c.* 527.—Cochinch.
Hindsii, Hance, in Journ. Linn. Soc. xiii. (1873) 108 = Allomorphia pauciflora.
parviflorus, Triana, in Trans. Linn. Soc. xxviii. (1871) 74 = cochinchinensis.

BLATTARIA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 247 (1745) = **Verbascum**, Linn. (Scrophul.).
alba, Medic. in Vorles. Churpf. Phys. Ges. iv. i. (1789) 280 = V. Lychnitis.
Bastardi, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 125 = V. Bastardi.
virgata, Fourr. l. c. = V. virgatum.
vulgaris, Fourr. l. c. = V. Blattaria.

BLATTI, Adans. Fam. ii. 88 (1763) = **Sonneratia**, Linn. f. (Lythrac.).

BLAXIUM, Cass. in Dict. Sc. Nat. xxx. 358 (1824) = **Dimorphothea**, Vaill. (Compos.).
decumbens, Cass. l. c. = D. fruticosa.

BLEASDALEA, F. Muell. Fragm. v. 90 (1865) = **Grevillea**, R. Br. (Proteac.).
cupaneoides, F. Muell. l. c. = G. Bleasdalea.

BLECHUM, P. Br. Hist. Jamaic. 261 (1756). *ACANTHACEAE*, Benth. & Hook. f. ii. 1062.
ALVARESIA, Pav. ex Nees, in DC. Prod. xi. 465 (1847).
angustifolium, *R. Br. Prod.* i. 478.—Ind. occ.
angustum, *Nees, in DC. Prod.* xi. 467.—N. Granat.
axillaripilosum, Juss. in Ann. Mus. Par. ix. (1807) 270 = Phaylopsis longifolia.

BLECHUM :—
brasiliense, Mart. ex Nees, in DC. Prod. xi. 284 = Stenandrium mandiocanum.
brasiliense, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 345.—Bras.
Brownei, *Juss. in Ann. Mus. Par.* ix. (1807) 270.—Ind. occ.
Brownei, H. B. & K. Nov. Gen. et Sp. ii. 238 = Linnaei.
costaricense, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 168.—Costa Rica.
grandiflorum, *Oerst. l. c.*—Am. centr.
Haenkei, *Nees, in DC. Prod.* xi. 467.—Mexic.
hamatum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 220.—Afr. trop.
laxiflorum, *Juss. in Ann. Mus. Par.* ix. (1807) 270.—Jamaica.
Linnaei, *Nees, in DC. Prod.* xi. 465.—Mexic.
luxonium, Nees, l. c. 466 = Brownei.
mexicanum, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 189.—Mexic.
Sternbergii, Nees, in DC. Prod. xi. 468 = Tetramerium nervosum.
trinitense, *Nees, l. c.* 466.—Ins. Trinit.
Tweedii, *Nees, l. c.*—Mexic.

BLEEKERIA, Hask. Retzia, i. 38 (1855) = **Oohrosia**, Juss. (Apocyn.).
calocarpa, C. Muell. in Walp. Ann. v. 492 = seq.
kalocarpa, Hask. Retzia, i. 40 = O. elliptica.
salubris, Hask. l. c. 41 = O. borbonica.

BLEEKERIA, Miq. Fl. Ind. Bat. i. ii. 407 (1859) = **Alohornea**, Sw. (Euphorb.).
Zollingerii, Miq. l. c. = A. villosa.

BLEEKRODEA, Blume, Mus. Bot. Lugd. Bat. ii. 87. t. 28 (1852) [1856]. *URTICACEAE*, Benth. & Hook. f. iii. 358.
insignis, *Blume, l. c.* 88.—Borneo.
madagascariensis, *Blume, l. c.*—Madag.

BLENCOCOËS, Rafin. Fl. Tellur. iii. 75 (1836) = **Macrargia**, Ruiz & Pav. (Scrophulac.).
filicaulis, Rafin. l. c. 76 = N. linearifolia.
longiflora, Rafin. l. c.—Hab.!

BLENNODERMA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 406 = **Oenothera**, Linn. (Onagrar.).
Drummondii, Spach, l. c. 407 = Oe. Spachiana.

BLENNODIA, R. Br. in Sturt, Exped. ii. App. 67 (1849). *CRUCIFERAE*, Benth. & Hook. f. i. 69.
alpestris, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 100.—Austral.
biennodes, F. Muell. l. c.—lasiocarpa.
brevipes, *F. Muell. l. c.*—Austral.
canescens, *R. Br. in Sturt, Exped. ii. App.* 67.—Austral.
cardaminoides, *Benth. Fl. Austral.* i. 75.—Austral.
Cunninghamii, *Benth. l. c.* 76.—Austral.
curvipes, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 100.—Austral.
crenulata, *Benth. Fl. Austral.* i. 74.—Austral.
filifolia, *Benth. l. c.* 73.—Austral.
lasiocarpa, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 100.—Australia.
nasturioides, *Benth. Fl. Austral.* i. 74.—Australia.
triacata, *Benth. l. c.*—Austral.

BLENNOSPERMA, Less. Syn. Comp. 267 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 404.
APALUS, DC. Prod. v. 507 (1836).
CONIOTHELE, DC. Prod. v. 581 (1836).
UNXIA, Colla, in Mem. Acc. Torin. xxxvii. (1835) 87. t. 22.
californicum, *Torr. & Gray, Fl. N. Am.* ii. 272.—Calif. chilense, *Less. Syn. Comp.* 267.—Chili.

BLENNOSPORA, A. Gray, in Hook. Kew Journ. iii. (1851) 172 = **Calocephalus**, R. Br. (Compos.).
Drummondii, A. Gray, l. c. 178 = C. Drummondii.

BLEPETALON, Rafin. Sylva Tellur. 80 (1838) = **Scutia**, Comm. (Rhamn.).
aculeatum, Rafin. l. c. = S. Commersoni.

BLEPHANTHERA, Rafin. Fl. Tellur. ii. 59 (1836) = **Bulbine**, Linn. (Liliac.).
depressa, Rafin. l. c. = Bulbine bulbosa.
Hookeri, Rafin. l. c. (= *Anthericum Hookeri*).—N. Zel.

BLEPHARACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836) = **Blepharis**, Juss. (Acanth.).
capensis, Nees, l. c. = Blepharis capensis.
cardaufolius, Nees, ex Steud. Nom. ed. II. i. 209 = Blepharis cardaufolia.
furcatus, Nees, ex Steud. l. c. = Blepharis furcata.
integrifolius, Nees, ex Steud. l. c. = Blepharis saturejaefolia.
procumbens, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 = Blepharis glomerata.
verticillatus, Steud. Nom. ed. II. i. 209 = Cardanthera verticillata.

BLEPHARADEN, Dulac, Fl. Hautes-Pyr. 449 (1867) = **Swertia**, Linn. (Gentianac.).
perennis, Dulac, l. c. = S. perennis.

BLEPHARANDRA, Griseb. in Linnaea, xxii. (1849) 7. *MALPIGHIACEAE*, Benth. & Hook. f. i. 324.
hypoleuca, *Griseb. l. c.*—Guiana.

BLEPHARANTHEMUM, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. 20 (1841) = **Plagianthus**, Forst. (Malvac.).
sidoides, Klotzsch, l. c. = P. sidoides.

BLEPHARANTHES, Sm. Gramm. Bot. 188 (1821) = **Modecca**, Lam. (Passif.).

BLEPHARIGLOTIS, Rafin. Fl. Tellur. ii. 88 (1836) = **Habenaria**, Willd (Orchid.).
albiflora, Rafin. l. c. = H. albida!
bicolor, Rafin. l. c. 39.—Hab.!
cristata, Rafin. l. c. = H. cristata.
flaviflora, Rafin. l. c.—Hab.!
longicornis, Rafin. l. c. 38 = H. longicornis.

BLEPHARIPAPPUS, Hook. Fl. Bor. Am. i. 316 (1833). *COMPOSITAE*, Benth. & Hook. f. ii. 392.
PTILONELLA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 385.
glandulosus, Hook. l. c. = Layia glandulosa.
scaber, *Hook. l. c.*—Am. bor. occ.

BLEPHARIS, Juss. Gen. 108 (1789). *ACANTHACEAE*, Benth. & Hook. f. ii. 1089.
ACANTHODIUM, Delile, Fl. Egypte, 241. t. 33 (1813).
BLEPHARACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836).
ablephars, *Nees, in DC. Prod.* xi. 267.—Java.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 160 = boerhaviaefolia.
acanthodioides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 212.—Afr. trop.
angusta, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 35.—Afr. austr.
arborea, *Sprong. Syst.* ii. 819.—Arabia.
asperrima, *Nees, in DC. Prod.* xi. 267.—Ind. or.
boerhaaviifolia, Roth, Nov. Sp. 320 = boerhaviaefolia.
boerhaviaefolia, *Pers. Syn.* ii. 180.—Ind. or.
calaminthaefolia, *Pers. l. c.*—Hab.!
capensis, *Pers. l. c.*—Afr. austr.
cardufolia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 35.—Afr. austr.
carnesiana, *Wats. ex S. Moore, in Journ. Bot.* xviii. (1880) 230.—Afr. trop.
edulis, *Pers. Syn.* ii. 180.—Arab.; Persia; Ind. bor. occ.
furcata, *Pers. l. c.*; *T. Anders. in Journ. Linn. Soc.* vii. (1864) 35.—Afr. austr.
furcata, Eckl. ex Nees, in DC. Prod. xi. 276; et Zeyh. ex Nees, l. c. 277 = capensis.
glauca, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 36.—Afr. austr.
glomerata, *Pers. Encyc. Suppl.* i. 89.—Afr. austr.
glumacea, *S. Moore, in Journ. Bot.* xviii. (1880) 232. —Afr. trop.
grossa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 36.—Afr. austr.
Guinzii, *T. Anders. l. c.* 34.—Afr. austr.
hirtinervia, *T. Anders. l. c.* 36.—Afr. austr.

BLEPHARIS :—
integrifolia, E. Mey. in Drège, Zwei Pfl. Docum. 168 = saturejaefolia.
involucrata, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 107.—Afr. trop.
linariaefolia, *Pers. Syn.* ii. 180.—Afr. trop.
maderaspatensis, Heyne, ex Roth, Nov. Sp. 320 = boerhaviaefolia.
molluginifolia, *Pers. Syn.* ii. 180.—Ind. or.
Noli-me-tangere, *S. Moore, in Journ. Bot.* xviii. (1880) 321.—Afr. trop.
pratensis, *S. Moore, l. c.* xv. (1877) 294.—Afr. trop.
procumbens, Heyne, ex Roth, Nov. Sp. 320 = boerhaviaefolia.
procumbens, Pers. Syn. ii. 180 = glomerata.
procurrens, Nees, in DC. Prod. xi. 267 = boerhaviaefolia.
pungens, *Klotzsch, in Peters, Reise Mossamb. Bot.* 311.—Afr. trop.
repens, Roth, Nov. Sp. 321 = molluginifolia.
rubiaefolia, Schum. in Dansk. Vidensak. Selsk. Afh. iv. (1858) 66 = boerhaviaefolia.
saturejaefolia, *Pers. Syn.* ii. 180.—Afr. austr.
scindica, Stocks, ex T. Anders. in Journ. Linn. Soc vii. (1864) 36 = linariaefolia.
serrulata, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 24.—Afr. trop.
setosa, *Nees, in DC. Prod.* xi. 265.—Afr. austr.
zeodica, Stocks, ex T. Anders. in Journ. Linn. Soc. ix. (1867) 500 = linariaefolia.
spathularis, *T. Anders. l. c.* vii. (1864) 35.—Afr. austr.
spicifolia, *Buhf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 85.—Ins. Socotr.
squarrosa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 35.—Afr. austr.
Togodelia, *Solms, ex Schweinf. Beitr. Fl. Aethiop* 108.—Abyss.
verticillata, Heyne, ex Wall. Cat. n. 2409 = Cardanthera verticillata.
Welwitschii, *S. Moore, in Journ. Bot.* xvii. (1880) 281.—Afr. trop.

BLEPHARISPERMUM, Benth. in DC. Prod. vii. 709 (1839) = BLEPHAROPHYLLUM, Klotzsch = **Scyphogyne**, Brongn. (Ericac.).

BLEPHARISPERMUM, Wight, ex DC. in Wight, Contrib. Bot. Ind. 11 (1834). *COMPOSITAE*, Benth. & Hook. f. ii. 393.
LEUCOBLEPHARIS, Arn. in Mag. Zool. & Bot. ii. (1838) 492.
petiolare, *DC. in Wight, l. c.* 12.—Ind. or.
spinulosum, *Oliver & Hiern, in Oliver. Fl. Trop. Afr.* iii. 335.—Afr. trop.
subsessile, *DC. in Wight, Contrib. Bot. Ind.* 12.—Ind. or.
zanguebaricum, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 336.—Afr. trop.

BLEPHARISTEMMA, Benth. in Wall. Cat n. 6230 (1832). *RHIZOPHOREAE*, Benth. & Hook. f. i. 682.
corymbosum, *Benth. l. c.*—Ind. or.

BLEPHAROCALYX, Berg. in Linnaea, xxvii. (1854) 412 = **Myrtus**, Tourn.
acuminatissimus, Berg. l. c. 418; et in Mart. Fl. Bras. xiv. 1. 491.—Reg. Argent.
acuminatus, Berg. ll. cc. 414, 426 (= Myrcia acuminata?).—Bras.
affinis, Berg, in Mart. Fl. Bras. xiv. 1. 505.—Bras.
amarus, Berg. l. c. 422.—Reg. Argent.
angustifolius, Berg. l. c. 421.—Bras.
angustissimus, Berg. l. c.—Bras.
apiculatus, Berg. l. c. 425.—Bras.
brunneus, Berg. l. c. 427.—Bras.
camecrus, Berg. l. c. 426.—Bras.
cisplatensis, Griseb. in Goett. Abh. xxiv. (1879) 126. —Reg. Argent.
cuspidatus, Berg, in Mart. Fl. Bras. xiv. 1. 494.—Bras.
depauperatus, Berg, in Linnaea, xxvii. (1854) 415; et in Mart. l. c. 426.—Bras.
ellipticus, Berg, ll. cc. 418 et 422.—Bras.
lanceolatus, Berg, in Mart. Fl. Bras. xiv. 1. 422.—Bras.
longipes, Berg, l. c. 423.—Bras.
parvifolius, Berg, l. c. 506.—Bras.
piscocarpus, Berg, l. c. 427.—Bras.
pilosus, Berg, l. c. 422.—N. Granat.
ramosissimus, Berg, l. c. 424.—Bras.

BLEPHAROCALYX —
salicifolia, Berg, in Linnaea, xxvii. (1854) 418 = Myrtus salicifolia.
Serra, Berg, in Mart. Fl. Bras. xiv. 1. 506.—Bras.
sessilifolius, Berg, l. c. 423.—Bras.
strictus, Berg, l. c. 423.—Bras.
Tweedei, Berg, in Linnaea, xxvii. (1854) 415; et in Mart. Fl. Bras. xiv. 1. 429.—Bras.
villosus, Berg, in Mart. Fl. Bras. xiv. 1. 425.—Bras.
Widgreni, Berg, l. c. 427.—Bras.

BLEPHAROCARYA, F. Muell. Fragm. ix. 15 (1878). *SAPINDACEAE*.
involucigera, *F. Muell. l. c.* 16.—Austral.

BLEPHAROCHLAMYS, Presl, Epim. Bot. 245 (1849) = **Myrcropetalon**, Harv. (Balanoph.).
capensis, Presl, l. c. 246 = M. Thomii.

BLEPHAROCHLOA, Endl. Gen. 1352 (1841) = **Lasiola**, Sw. (Gramin.).
ciliata, Endl. l. c. = L. ciliata?

BLEPHARODON, Decne. in DC. Prod. viii. 608 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 756.
PYCHOLEPIS, Griseb. in Lechl. Berb. Am. Austr. 58 (1857).
amplifiorus, *Fourn. in Mart. Fl. Bras.* vi. iv. 304.—Bras.
asperuloides, *Decne. in DC. Prod.* viii. 604.—Bras.
asterios, *Fourn. in Mart. Fl. Bras.* vi. iv. 307.—Bras.
bicolor, *Decne. in DC. Prod.* viii. 604.—Bras.
bicuspidatus, *Fourn. in Mart. Fl. Bras.* vi. iv. 306.—Bras.
bracteatus, *Fourn. l. c.* 307.—Bras.
Decaisnei, *Fourn. l. c.* 305.—Bras.
diffusus, *Decne. in DC. Prod.* viii. 605.—Bras.
grandiflorum, *Benth. Pl. Hartw.* 215.—N. Granat.
laurifolium, Decne. in DC. Prod. viii. 608 = Nephradenia laurifolia.
linearis, Decne. l. c. = Nephradenia linearis.
linearifolium, Benth. & Hook. f. Gen. ii. 756, sphalm. = lanceolatum.
longipedicellatum, *Fourn. in Mart. Fl. Bras.* vi. iv. 306.—Bras.
mucronatum, *Decne. in DC. Prod.* viii. 608.—Mexic.
nervifolium, Decne. l. c. = Nephradenia nervifolia
pallidum, *Decne. l. c.*—Bras.
salicinum, *Decne. l. c.* 604.—Bras.?
Spruceanum, *Fourn. in Mart. Fl. Bras.* vi. iv. 307.—Bras.
triplinerve, *Turcz. in Bull. Soc. Nat./Mosc.* xxv. (1852) ii. 318.—Venezuela.

BLEPHAROLEPIS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 443 (1836) = Polypoda, Presl (Filicid.).
Zeyheriana, Nees, l. c. = P. capensis.

BLEPHAROLEPIS, Nees, in Mart. Fl. Bras. ii. 1. 91 (1843) = Scirpus, Linn. (Cyperac.).
leucocephalaria, Nees, l. c.—Bras.

BLEPHAROPHYLLUM, Klotzsch, in Linnaea, xii. (1838) 216 = **Scyphogyne**, Brongn. (Ericac.).
divaricatum, Klotzsch, l. c. 215 = S. divaricata.

BLEPHAROSTEMMA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 398 = **Aspertula**, Linn. (Rubiac.).
taurinum, Fourr. l. c. = A. taurina.

BLEPHILIA, Raf. in Am. Monthly Mag. (1819) 190; et Journ. Phys. lxxxix. (1819) 98. *LABIATAE*. Benth. & Hook. f. ii. 1198.
Beckl, *Raf. in Atl. Journ.* 146.—Am. bor.
brevifolia, Raf. in Am. Fl. Am. iv. 96 = ciliata.
brevipes, *Raf. l. c.*—Am. bor.
ciliata, *Raf. in Am. Monthly Mag.* (1819) 190.—Am. bor.
heterophylla, Raf. in Am. Fl. Am. iv. 96 = ciliata?
hirsuta, *Benth. Lab. Gen. et Sp.* 330.—Am. bor.
lanceolata, Raf. in Am. Fl. Am. iv. 96 = hirsuta?
nepetoides, Raf. l. c. = hirsuta.
prostrata, Raf. l. c. = ciliata.

BLEPHILOMA, Raf. Fl. Tellur. iii. 96 (1836) = Phlomis, Tourn. (Labiat.).
ampifolia, Raf. l. c.—Hab.?

BLEPHISTELMA, Raf. Fl. Tellur. iv. 108 (1836) = **Passiflora**, Linn.
aurantia, Raf. l. c. = P. aurantia.

BLETIA, Ruiz & Pav. Prod. 119. t. 26 (1794). *ORCHIDEAE*, Benth. & Hook. iii. 513.
BLETIANA, Raf. in Am. Monthly Mag. (1818) 268
BLETILLA, Reichb. f. in Fl. des Serres, viii. (1853) 246.
DILOMILIS, Raf. Fl. Tellur. iv. 48 (1836).
GYAS, Salisb. in Trans. Hort. Soc. i. (1812) 299.
HECASE, Raf. Fl. Tellur. iv. 44 (1836).
JIMENSIA, Raf. l. c. 38 (1836).
THIERAUTIA, Colla, Hort. Rupul. 189 (1824).
acaulis, Reichb. f. in Walp. Ann. vi. 485 = Brassavola acaulis.
acuminata, Reichb. f. l. c. 427 = Laelia acuminata.
acutipetala, Hook. Bot. Mag. t. 5217 = verecunda.
adenocarpa, *Reichb. f. in Bonplandia*, iv. (1856) 216 —Mexic.
albida, Reichb. f. in Walp. Ann. vi. 428 = Laelia albida.
amazonica, Reichb. l. c. 434 = Brassavola amazonica
ambonensis, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 180 = Phaius ambonensis.
ancops, Reichb. f. in Walp. Ann. vi. 418 = Laelia ancepa.
angustata, Gaudich. in Freyc. Voy. Bot. 421 = Spathoglottis plicata.
angustata, Reichb. f. in Walp. Ann. vi. 434 = Brassavola angustata.
anomala, *Reich. & Gal. in Ann. Sc. Nat.* Sér. III. i (1845) 28.—Mexic.
aphylla, Nutt. Gen. Am. ii. 194 = Hexalectris aphylla.
attenuata, *Reichb. f. Xenia Orchid.* ii. 64.—Ind. occ.
autumnalis, La Llave & Lex. Nov. Veg. Desc. fasc ii. (Orch. Opusc.) 19 = Laelia autumnalis.
bicallosa, D. Don, Prod. Fl. Nep. 30 = Liparis bituberculata?
Boothiana, Reichb. f. Xenia Orchid. ii. 51 = Laelia Boothiana.
bracteosa, *Grisb. Notul.* iii. 226.—Ind. or.
campanulata, *La Llave & Lex. Nov. Veg. Desc.* fasc ii. (*Orch. Opusc.*) 17.—Mexic.
capitata, [R. Br. in] Ait. Hort. Kew. ed. II. v. 206 = Eulophia capitata.
Casperiana, Reichb. f. in Walp. Ann. vi. 428 = Laelia purpurata.
catenulata, *Ruiz & Pav. Syst.* 229.—Peruv.
caulescens, Reichb. f. in Walp. Ann. vi. 431 = Laelia caulescens.
Celebella, Reichb. f. l. c. 435 = Brassavola Cebolletta.
ciliata, *Pursh, Fl. Hasmh.* 99.—Peruv.
cinnabarina, Reichb. f. in Walp. Ann. vi. 430 = Laelia cinnabarina.
cinnamomea, Reichb. f. l. c. 418 = Laelia cinnamomea.
coccinea, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 16.—Mexic.
cordata, Reichb. f. in Walp. Ann. vi. 436 = Brassavola nodosa.
crispa, Reichb. f. l. c. 428 = Laelia crispa.
crispina, Reichb. f. l. c. 419 = Schomburgkia crispa.
cucullata, Reichb. f. l. c. 433 = Brassavola cucullata.
Dabia, D. Don, Prod. Fl. Nep. 30 = Eulophia campestris?
densiflora, *Grisb. Notul.* iii. 226.—Ind. or.
Digbyana, Reichb. f. in Walp. Ann. vi. 422 = Laelia Digbyana.
domingensis, Reichb. f. l. c. 432 = Broughtonia lilacina.
elegans, Reichb. f. l. c. 427 = Laelia elegans.
emiformis, *Ruiz & Pav. Syst.* 230.—Peruv.
Eulophiae, Reichb. f. in Walp. Ann. vi. 439 = Tetramicra teneru.
exspatha, Reichb. f. l. c. 424 = Laelia exspatha.
flabellata, Reichb. f. l. c. 422 = Laelia xanthina.
flava, Reichb. f. l. c. 421 = Laelia flava.
flava, Wall. ex Lindl. Gen. et Sp. Orch. 227 = Phaius maculatus.
florida, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 206.—Ind. occ.
fulgens, *Reichb. f. in Bonplandia*, iii. (1855) 221.—Mexic.
furfurea, Reichb. f. in Walp. Ann. vi. 438 = Laelia furfuracea.
gebina, Lindl. in Journ. Hort. Soc. ii. (1847) 307 = lyrachellina.
glauca, Reichb. f. in Walp. Ann. vi. 433 = Laelia glauca.

BLETIA :—
gloriosa, Reichb. f. in Walp. Ann. vi. 419 = Schomburgkia crispa.
gracilis, *Ladd. Bot. Cab.* t. 1977.—Mexic.
graminifolia, D. Don, Prod. Fl. Nep. 29 = Arundina bambusifolia.
grandiflora, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 17 = Laelia grandiflora.
grandis, Reichb. f. in Walp. Ann. vi. 424 = Laelia grandis.
guineensis, *G. Don, ex Sweet, Hort. Brit.* ed. I. 492, nomen.—Guinea.
havanensis, *Lindl. Bot. Reg.* (1838) *Misc.* 38.—Cuba.
Humboldtii, Reichb. f. in Walp. Ann. vi. 430 = Schomburgkia Humboldtii.
humilis, Link & Otto, Ic. Pl. Rar. 53. t. 27 = Maxillaria gracilis.
hyacinthina, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 206. —China; Japon.
irrorata, Reichb. f. in Walp. Ann. vi. 426 = Laelia irrorata.
jucunda, *Linden & Reichb. f. in Bonplandia*, iii. (1855) 221.—Mexic.
Landsbergii, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 317.—Am. trop.
latifolia, *Gaudich. Bot. Freyc. Voy.* 421.—Ins. Molucc.
lilacina, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 23.—Mexic.
Lindenii, Reichb. f. in Walp. Ann. vi. 431 = Laelia Lindenii.
Lindleyana, Reichb. f. Xenia Orchid. ii. 65 = Laelia Lindleyana.
linearis, Reichb. f. in Walp. Ann. vi. 426 = Brassavola linearis.
lobata, Reichb. f. l. c. 424 = Laelia Boothiana
longipes, Reichb. f. Xenia Orchid. ii. 59 = Laelia longipes.
Lundii, Reichb. f. & Warm. in Otia Bot. Hamb. 92 = Laelia Lundii.
Lyonii, Reichb. f. in Walp. Ann. vi. 418 = Schomburgkia Lyonsia.
marginata, Reichb. f. l. c. 419 = Schomburgkia marginata.
Martiana, Reichb. f. l. c. 434 = Brassavola Martiana.
Mauwa, D. Don, Prod. Fl. Nep. 30 = Calanthe Masuca.
mentana, Reichb. f. in Walp. Ann. vi. 446 = Octadesmia serratifolia.
nipponica, *Franch. & Sav. Enum. Pl. Jap.* ii. 511.—Japon.
nodosa, Reichb. f. in Walp. Ann. vi. 437 = Brassavola nodosa.
obcordata, *Lindl. Gen. et Sp. Orch.* 123.—Ind. or. Origistana, *Reichb. f. in Otto, Hamb. Gartenz.* xvi. (1860) 420.—Hab.?
pallida, Lodd. Bot. Cab. t. 939 = florida.
Parkinsoni, *Hook. Bot. Mag.* t. 3736.—Mexic.
parviflora, Reichb. f. in Walp. Ann. vi. 439 = Tetramicra parviflora.
parviflora, *Ruiz & Pav. Syst. Veg.* 231.—Peruv.
patula, *Hook. Bot. Mag.* t. 3518.—Ins. S. Doming.
Paulinae, F. Muell. Fragm. vi. 95 = Spathoglottis Paulinae.
pedunculata, Reichb. f. in Walp. Ann. vi. 426 = Laelia pedunculata.
Perrinii, Reichb. f. l. c. 421 = Laelia Perrinii.
platyphylla, Reichb. f. ex Griseb. Cat. Pl. Cub. 264 = sarcophylla.
praetansa, Reichb. f. in Walp. Ann. vi. 425 = Laelia praetansa.
pratensis, Reichb. f. l. c. 439 = Tetramicra rigida.
pulchella, Hort.—Cf. Gard. Chron. (1850) 168 = verecunda.
pumila, Reichb. f. in Walp. Ann. vi. 421 = Laelia pumila.
Pumilio, Reichb. f. l. c. 436 = Brassavola Pumilio.
punctata, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (Orch. Opusc.) 15.—Mexic.
purpurata, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 23 = Archaea rosea.
purpurata, Reichb. f. in Walp. Ann. vi. 423 = Laelia purpurata.
purpurea, DC. Huit. Not. Pl. Rar. Jard. Genèv. 23 = verecunda.
rafflesia, *Lindl. Bot. Reg.* xxi. (1836) t. 1760.—Peruv.
rapanda, *Ruiz & Pav. Syst. Veg.* 230.—Peruv.
rubens, Reichb. f. in Walp. Ann. vi. 435 = Brassavola nesua.

BLETIA :—
rhopalorrhachis, Reichb. f. l. c. 437 = Brassavola rhopalorrhachis.
rigida, Reichb. f. l. c. 438 = Tetramicra rigida.
Roezlii, *Reichb. f. in Linnaea*, xli. (1877) 7.—Mexic.
rosea, Reichb. f. in Walp. Ann. vi. 420 = Schomburgkia rosea.
rubescens, Reichb. f. l. c. 425 = Laelia rubescens.
rupestris, Reichb. f. l. c. 431 = Laelia rupestris.
sanguinea, Poepp. & Endl. Nov. Gen. et Sp. i. 56. t. 95 = catenulata.
sarcophylla, *Reichb. f. in Flora*, xlviii. (1865) 278.—Ind. occ.
Schilleriana, Reichb. f. in Walp. Ann. vi. 424 = Laelia Schilleriana.
schomina, Reichb. f. l. c. 439 = Tetramicra schoenina.
Schomburgki, Reichb. f. l. c. = Tetramicra rigida.
secunda, *Lindl. Bot. Reg.* (1840) *Misc.* 57.—Mexic.
Shepherdii, Hook. Bot. Mag. t. 3319 = verecunda.
Sherrattiana, *Batem. in Hook. l. c.* t. 5646.—N. Granat.
speciosa, H. B. & K. Nov. Gen. et Sp. i. 343 = Laelia grandiflora.
splendens, *Linden, Illustr. Hortic.* xxvii. (1881) 64, nomen.—Hrsa.
striata, *C. Presl, Rel. Haenk.* 98.—Ins. Luzon.
superbiens, Reichb. f. in Walp. Ann. vi. 418 = Laelia superbiens.
sylvatica, Boj. Hort. Maurit. 218 = Calanthe sylvatica.
Tankervilleae, R. Br. in Bot. Mag. t. 1924 = Phaius grandiflorus.
tenera, A. Rich. Fl. Cub. Fanerog. ii. 242 = Tetramicra tenera.
tibicinis, Reichb. f. in Walp. Ann. vi. 429 = Schomburgkia tibicina.
tuberculata, Reichb. f. l. c. 434 = Brassavola tuberculata.
tuberculosa, Spreng. Syst. iii. 744 = Phaius tuberculosus.
undulata, Reichb. f. in Walp. Ann. vi. 420 = Schomburgkia undulata.
uniflora, *Ruiz & Pav. Syst. Veg.* 230.—Peruv.
venosa, Reichb. f. in Walp. Ann. vi. 438 = Brassavola venosa.
verecunda, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 206. —Florida; Ind. occ.
villosa, Spreng. Syst. iii. 743 = Phaius villosus.
violacea, Reichb. f. in Walp. Ann. vi. 426 = Laelia violacea.
Wagenerii, *Reichb. f. in Bonplandia*, ii. (1854) 22.—Venezuela.
Wallisii, *Reichb. f. in Linnaea*, xli. (1877) 115.—N. Granat.
Wendlandi, Reichb. f. in Walp. Ann. vi. 431 = Laelia Wendlandi.
Woodfordii, Hook. Bot. Mag. t. 2719 = Phaius maculatus.
xanthina, Reichb. f. in Walp. Ann. vi. 425 = Laelia xanthina.

BLETIANA, Rafin. in Am. Monthly Mag. (1818) 268 = **Bletia**, Ruiz & Pav. (Orchid.).

BLETILLA, Reichb. f. in Fl. des Serres, Sér. I. vii. (1851–52) 246 = **Bletia**, Ruiz & Pav. (Orchid.).
florida, Reichb. f. l. c. = Bletia florida.
Gebinae, Reichb. f. l. c. = Bletia hyacinthina.

BLETTI, Steud. Nom. ed. II. i. 210 (1840) = BLATTI, Adans. = **Bonnetarias**, Lian. f. (Lythrar.).

BLIGHIA, Kon. in Kon. & Sims, Ann. Bot. ii. (1806). 571. t. 16, 17. *SAPINDACEAE*. [Benth. & Hook. f. i. 399.]
AKEA, Stokes, Bot. Mat. Med. ii. 353 (1812).
AKEESIA, Tussac, Fl. Ant. i. 66 (1808).
sapida, *Kon. in Kon. & Sims, Ann. Bot.* ii. (1806) 571. —Afr.; Ind. occ.
unijugata, Baker, in Oliver, Fl. Trop. Afr. i. 427.—Afr. trop.
xanthocarpa, *Baker, l. c.*—Afr. trop.

BLINKWORTHIA, Choisy, in Mém. Soc. Phys. Genève. vi. (1833) 430. t. 4. *CONVOLVULACEAE*; Benth. & Hook. f. ii. 869.
lycioides, *Choisy, l. c.*—Burma.

BLITANTHUS, Reichb. Cat. Hort. Dresd. (1824); Handb. 238 (1837) = **Acroglochin**, Schrad. (Chenopod.).
sagittatus, Reichb. ii. cc. = A. chenopodioides.

BLITUM, [Tourn.] Linn. Gen. ed. I. 20 (1737) = **Chenopodium**, Linn.
acuminatum, Schur, Enum. Pl. Transs. 571 = C. rubrum.
antarcticum, Hook. f. Fl. Antarct. ii. 549 = C. antarcticum.
atriplicinum, F. Muell. in Trans. Vict. Inst. (1855) 138 = C. striatiplicatum.
Bonus-Henricus, C. A. Mey. Fl. Alt. i. 11, note = C. Bonus-Henricus.
botryoides, Dne. in Fl. Dan. t. 2804 = C. rubrum.
californicum, S. Wats. in Proc. Am. Acad. ix. (1874) 101 = C. californicum.
capitatum, Linn. Sp. Pl. 4 = C. capitatum.
carinatum, C. A. Mey. in Ledeb. Fl. Alt. i. 11 adnot. = C. carinatum.
chenopodioides, Bernh. ex Steud. Nom. ed. II. i. 210 = petiolare.
chenopodioides, Lam. Encyc. ii. 11 = C. capitatum.
chenopodioides, Nutt. Gen. Am. i. 4 = Monolepis chenopodioides.
crassifolium, Reichb. Fl. Germ. Excurs. 582 = C. rubrum.
cristatum, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 78 = C. crustatum.
gangeticum, Moench, Meth. 359 = Amaranthus gangeticus.
graecum, Koch, ex Buek, Ind. DC. Prod. iii. 64 (est sectio generis)
glandulosum, Moq. in DC. Prod. xiii. ii. 82 = C. carinatum
glaucum, Koch, in Sturm, Deutschl. Fl. Heft 75 = C. glaucum.
graecisanz, Moench, Meth. 359 = Amaranthus Blitum.
lividum, Moench, l. c. = Amaranthus lividus.
maritimum, Nutt. Gen. Am. ii. Add. post Ind. [12] = C. rubrum.
mauve, Scop. Fl. Carn. ed. II. ii. 287 = Amaranthus Blitum.
melancholicum, Moench, Meth. 359 = Amaranthus gangeticus.
Nuttallianum, Roem. & Schult. Mant. i. 65 = Monolepis chenopodioides.
oleraceum, Moench, Meth. 359 = Amaranthus oleraceus.
petiolare, Link, Enum. Hort. Berol. i. 8.—Teneria!
polymorphum, C. A. Mey. in Ledeb. Fl. Alt. i. 18 = C. rubrum.
Pumilio, C. A. Mey. l. c. 11, adnot. = C. Pumilio.
rubrum, C. A. Mey. l. c. = C. rubrum.
Sanguisorba, Pritz. Ic. Ind. 155, sphalm. = Poterium Sanguisorba.
tataricum, Mill. Gard. Dict. ed. VIII. n. 3 = C. capitatum.
tenue, Moq. in DC. Prod. xiii. ii. 81 = Roubieva multifida.
terminale, Stokes, Bot. Comment. i. 11 = C. capitatum.
virgatum, Linn. Sp. Pl. 4 = C. capitatum.
viride, Moench, Meth. 359 = C. album.

BLOCHMANNIA, Reichb. Consp. 168 (1828) = **Triplaris**, Linn. (Polygon.).
peruviana, Poepp. ex Meisn. in DC. Prod. xiv. 178 = T. Poeppigiana.
Weigeltiana, Reichb. Consp. 168 = T. surinamensis.

BLONDEA, Rich. in Act Soc. Hist. Nat. Par. i. (1792) 110 = **Sloanea**, Linn. (Tiliac.).
latifolia, Rich. l. c.—Guiana.

BLONDIA, Neck. Elem. ii. 115 (1790) = **Tiarella**, Linn. (Saxifr.).
biternata, Rafin. Fl. Tellur. ii. 75 = Astilbe decandra.
trifoliata, Rafin. l. c. = T. trifoliata.

BLOOMERIA, Kellogg, in Proc. Calif. Acad. ii. 11 (1863). *LILIACEAE*, Benth. & Hook. f. Gen. iii. 801.
aurea, *Kellogg, l. c.*—Calif.
Clevelandi, *S. Wats. in Proc. Am. Acad.* xx. (1885) 376.—Calif.

BLUFFIA, Delile, Sem. Hort. Hamb. (1838) (Cf. Linnaea, x. (1836) 114); Nees, Fl. Afr. Austr. Gram. (1836) = **Panlouim**, Linn. (Gramin.).
Ecklonicana, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447.—Afr. austr.

BLUMEA, G. Don, Gen. Syst. i. 378 (1831) = BLUMIA, Spreng. = **Saurauja**, Willd. (Ternstrocm.).

BLUMEA, Reichb. Consp. 209 (1828) = **Nocca**, Blume (Malvac).
altissima, Reichb. ex Steud. Nom. ed. II. i. 211 = N. altissima.

BLUMEA, DC. in Guill. Arch. Bot. ii. (1833) 514.
COMPOSITAE, Benth. & Hook. f. ii. 289.
DOELLIA, Sch. Bip. in Walp. Rep. ii. 958 (1843).
PLACUS, Lour. Fl. Cochinch. 496 (1790).
abyssica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 393.—Abyss.
acutata, DC. in Nouv. Ann. Mus. Par. iii. (1834) 408 = laciniata.
alata, DC. Prod. v. 448 = Laggera alata.
amethystina, Hance, in Journ. Bot. vi. (1868) 178 = glomerata.
amplectens, *DC. in Wight, Contrib. Bot. Ind.* 18.—Ind. or.
amplectens, Thw. Enum. Fl. Zeyl. 163 = bifoliata.
anagalidifolia, DC. in Wight, Contrib. Bot. Ind. 18 = amplectens.
angustifolia, *Thw. Enum. Pl. Zeyl.* 164.—Zeylan.
appendiculata, *DC. Prod.* v. 447 —Java.
arenaria, DC. in Wight, Contrib. Bot. Ind. 18 = amplectens.
arfakiana, *Martelli, in Nuov. Giorn. Bot.* xv. (1883) 292.—N. Guin.
Arnottiana, Steud. Nom. ed. II. i. 210 = Inula Cappa.
aromatica, *DC. Prod.* v. 446.—Reg. Himal. ; Burma.
aurita, DC. in Wight, Contrib. Bot. Ind. 16 = Laggera aurita.
axillaris, DC. Prod. v. 434 = lacera.
baccharoides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 391 = Pluches Dioscoridis.
balsamifera, *DC. Prod.* v. 447.—Ind. or. ; Burma ; Malaya.
barbata, *DC. in Wight, Contrib. Bot. Ind.* 14.—Ind. or.
Belangeriana, *DC. Prod.* v. 444.—Ind. or.
bifoliata, *DC. in Wight, Contrib. Bot. Ind.* 14.—Ind. or. ; Burma : Malaya.
Bowei, Vatke, in Linnaea, xxxix. (1875) 485 = abyssinica.
bovina, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 178 = oxyodonta.
cernua, DC. Prod. v. 436 = hieraciifolia.
Chamissoniana, DC. l. c. 434 = Wightiana.
chinensis, *DC. l. c.* 444.—Ind. or. ; China ; Java.
chinensis, Hook. & Arn. Bot. Beech. Voy. 195 = Inula Cappa.
chinensis, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 294 = hieraciifolia.
cicharioïdes, DC. ex Decne. in Nouv. Ann. Mus. Par. iii. (1834) 410 = laciniata.
cinerascens, DC. Prod. v. 439 = lacera.
Clarkei, Hook. f. Fl. Brit. Ind. iii. 267 = elongata.
Commersonii, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 88.—Ind. or.
crassifolia, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 394.—Abyss.
crepidifolia, DC. Prod. v. 437 = laciniata.
crinita, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 848. —Zeylan.
crinita, Steets, in Seem. Bot. Voy. Herald, 387 = hieraciifolia.
cuneifolia, DC. Prod. v. 441 = lacera.
Cunninghami, *DC. l. c.* 435.—Austral.
densiflora, *DC. l. c.* 446.—Ind. or. ; Malaya ; Ins. Pacif.
diffusa, *R. Br. ex Benth. Fl. Austral.* iii. 525.—Austral.
diffusa, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 187.—Burma.
Dregeana, *DC. Prod.* v. 436.—Afr. austr.
dregeanoides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 393 = lacera.
elongata, *DC. Prod.* v. 445.—Reg. Himal. ; Burma ; Penang.
eriantha, *DC. in Wight, Contrib. Bot. Ind.* 15.—Ind. or.
excisa, DC. Prod. v. 446 = densiflora.
fasciculata, DC. l. c. 442 = glomerata.
fatuloza, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 187 = glomerata.
flava, DC. Prod. v. 439 = Laggera flava.
flexuosa, *C. B. Clarke, Compos. Ind.* 96.—Ind. or.
fruticulosa, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 178 = lacera.

BLUMEA :—
garispina, *DC. Prod.* v. 448.—Afr. austr.
glandulosa, Benth. Fl. Hongk. 177 = laciniata.
glandulosa, DC. in Wight, Contrib. Bot. Ind. 14 = lacera.
glandulosa, Thw. Enum. Pl. Zeyl. 163 = membranacea.
glomerata, *DC. in Wight, Contrib. Bot. Ind.* 15.—Ind. or. ; Malaya ; China.
glutinosa, *DC. Prod.* v. 434.—Madag.
gracilis, DC. l. c. 443 = lacera.
grandiflora, Zipp. ex Span. in Linnaea, xv. (1841) 105 = Capparis spinosa.
grandis, DC. Prod. v. 447 = densiflora.
guineensis, DC. l. c. 449 = Laggera aurita.
Hamiltoni, DC. l. c. 439 = hieraciifolia.
Heyneana, DC. in Wight, Contrib. Bot. Ind. 15 = lacera.
hieraciifolia, *DC. l. c.* 15.—Ind. or. ; Burma ; Malaya.
hieraciifolia, Hook. f. & Thoms. ex Fl. Brit. Ind. iii. 267 = elongata.
hispidula, DC. ex C. B. Clarke, in Fl. Brit. Ind. iii. 272 = membranacea.
holosericea, Dals. & Gibs. Bomb. Fl. 126 = glomerata.
holosericea, DC. Prod. v. 443 = lacera.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 269.—Reg. Himal.
hymenophylla, DC. Prod. v. 440 = membranacea.
integrifolia, *DC. Prod.* v. 433.—Austral.
Jacquemontii, *Hook. f. Fl. Brit. Ind.* iii. 265.—Ind. or.
javanica, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie*, ii. (1845) 239.—Java.
Kingii, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 265. —Ind. or.
lacera, *DC. in Wight, Contrib. Bot. Ind.* 14.—Ind. or. ; Malaya ; Ins. Mascar.
laciniata, *DC. Prod.* v. 436.—Ind. or. ; Malaya ; China.
lactucaefolia, DC. l. c. 435 = lacera.
lapsanoides, DC. in Wight, Contrib. Bot. Ind. 15 = virens.
leptoclada, DC. Prod. v. 443 = glomerata.
leptoclada, Dals. & Gibs. Bomb. Fl. 125 = lacera.
Leschenaultiana, DC. Prod. v. 435 = Wightiana.
leucanthema, *DC. l. c.* 436 (Dekha).—Nepal.
longifolia, DC. l. c. 446 = myriocephala ?
macrophylla, *DC. l. c.* 446.—Java.
macrostachya, DC. l. c. 442 = hieraciifolia.
malabarica, *Hook. f. Fl. Brit. Ind.* iii. 267.—Ind. or.
Malcolmii, *Hook. f. l. c.* 266.—Ind. or.
manillensis, *DC. Prod.* v. 435.—Ins. Luzon.
membranacea, *DC. l. c.* 440.—Ind. or. ; Malaya.
Metziana, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 266 = Belangeriana.
Milnei, Sezn in Bonplandia, ix. (1861) 257 ; Fl. Vit. 141 = densiflora.
muralis, DC. Prod. v. 440 = membranacea.
Munro, DC. l. c. 435 = lacera.
myriocephala, *DC. l. c.* 445.—Reg. Himal. ; Burma.
myriocephala, Thw. Enum. Pl. Zeyl. 163 = spectabilis.
napifolia, DC. l. c. 440 = lacera.
nataiensis, *Sch. Bip. in Walp. Rep.* ii. 971.—Afr. austr.
nelligherrcala, *Hook. f. Fl. Brit. Ind.* iii. 261.—Ind. or.
nitida, Wall. ex C. B. Clarke, Comp. Ind. 85 = chinensis.
nitida, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie*, ii. (1845) 244.—Java.
nodiflora, *Hook. f. Fl. Brit. Ind.* iii. 262.—Burma.
obovata, C. B. Clarke, Comp. Ind. 77 = Jacquemontii.
obovata, DC. Prod. v. 446.—Nepal.
oligocephala, DC. in Wight, Contrib. Bot. Ind. 13 = bifoliata.
oloptera, *DC. Prod.* v. 448.—Afr. trop.
oxyodonta, *DC. in Wight, Contrib. Bot. Ind.* 15.—Ind. or. ; Burma ; Malaya ; China.
parviflora, DC. Prod. v. 437 = Wightiana.
pauciflora, DC. l. c. 440 = membranacea.
Perrottetiana, DC. l. c. 443 = lacera.
Petitiana, *A. Rich. Tent. Fl. Abyss.* i. 395.—Abyss.
phagnaloïdes, A. Rich. l. c. 398 = Phagnalon sceparium.
phyllostachya, DC. Prod. v. 438 = Wightiana.
procera, *DC. l. c.* 445.—Ind. or. ; Burma.
pteroclonia, DC. in Wight, Contrib. Bot. Ind. 15 = Laggera pteroclonia.
pubiflora, DC. Prod. v. 434 = amplectens.

BLUMEA :—
purpurascens, A. Rich. Tent. Fl. Abyss. i. 395 = Laggera pteroclonia.
purpurea, DC. Prod. v. 442 = glomerata.
racemosa, DC. l. c. 442 = glomerata.
riparia, DC. l. c. 444 = chinensis.
runcinata, DC. l. c. 436 = laciniata.
runcinata, Wall. Cat. n. 3057 = lacera.
salvifolia, *DC. Prod.* v. 448.—Ins. Borbon.
salvifolora, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie*, ii. (1845) 246.—Java.
saxatilis, *Zoll. & Mor. l. c.* 248.—Java.
semisecrisa, DC. Prod. v. 445 = procera.
senecioides, DC. l. c. 439 = membranacea.
senecioides, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 88 = Pulicaria foliolosa.
senegalensis, DC. Prod. v. 449 = Conyza senegalensis.
sericana, *Hook. f. Fl. Brit. Ind.* iii. 262.—Burma.
sessiliflora, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 410.—Ins. Timor.
sessilifolia, DC. Prod. v. 447 = myriocephala.
sikkimensis, *Hook. f. Fl. Brit. Ind.* iii. 268.—Reg. Himal.
soldanelloides, *DC. Prod.* v. 443.—Afr. trop.
sonchifolia, DC. l. c. 438 = laciniata.
spectabilis, *DC. l. c.* 445.—Ind. or.
spinellosa, DC. l. c. 443 = hieraciifolia ?
squarrosa, C. B. Clarke, Comp. Ind. 87 = myriocephala.
subcapitata, *DC. Prod.* v. 439.—Reg. Himal.
subcrispiter, DC. l. c. 441 = membranacea.
sylvatica, *DC. l. c.* 447.—Java.
tenella, DC. in Wight, Contrib. Bot. Ind. 13 = bifoliata ?
tetraptera, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 313 —Ins. Philipp.
thyrsoidea, Sch. Bip. in Bot. Zeit. xxiv. (1866) 166 = lacera.
timorensis, *DC. ex Decne. in Nouv. Ann. Mus. Par* iii. (1834) 411.—Ins. Timor.
tomentosa, *A. Rich. Tent. Fl. Abyss.* i. 394.—Abyss.
trichophora, DC. Prod. v. 436 = Wightiana.
trigona, DC. l. c. 437 = lacera.
triulca, DC. l. c. 440 = Conyza stricta.
turneroïdes, DC. l. c. 447 = Laggera alata.
villosa, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 268 = lacera.
vimines, *DC. Prod.* v. 443.—Ins. Timor.
virens, *DC. in Wight, Contrib. Ind. Bot.* 14.—Ind. or.
viscosula, DC. Prod. v. 441 = membranacea.
Wallichii, *C. B. Clarke, Comp. Ind.* 87.—Assam.
Wightiana, *DC. in Wight, Contrib. Bot. Ind.* 14.—Geront. trop.
Zollingeriana, C. B. Clarke, Comp. Ind. 90 = balsamifera.

BLUMENBACHIA, Koel. Gram. Gall. et Germ. 28 (1802) = **Borghkum**, Moench (Gramin.).
halepensis, Koel. l. c. 29 = S. halepense.

BLUMENBACHIA, Schrad. in Goett. Gel. Anz. (1826) 1705 ; Blumenb. (1827) 6. t. 1. *LOASKAE*, Benth. & Hook. f. i. 806.
CAJOPHORA, Endl. Gen. 981 (1839).
GRIFIDIA, Miers, in Trans. Linn. Soc. xxv. (1866) 227. t. 28.
RAPHIBANTHE, Lilja, in Linnaea, xv. (1841) 262.
cardiifolia, *Ball, in Journ. Linn. Soc.* xxii. (1885) 89. —Peruv.
cornea, *Griseb. in Goett. Abh.* xix. (1874) 152.—Reg. Argent.
chuquitensis, *Hook. f. Bot. Mag.* t. 6148.—Peruv.
conferta, Hook. f. l. c. t. 6184 = grandiflora.
corensis, *Haage & Schmidt, in Ken. Hortic.* (1874) 58.—Chili.
Domeykoana, *Phil. in Linnaea*, xxxiii. (1864–65) 75. —Chili.
Espigneura, *C. Gay, Fl. Chil.* ii. 434.—Chili.
grandiflora, *G. Don, Gen. Syst.* iii. 62.—Peruv.
Hieronymi, *Urb. in Bichl. Fahrb. Berl.* iii. (1884) 249. —Reg. Argent.
insignis, *Schrad. in Goett. Gel. Anz.* (1828) 1706.—Reg. Argent.
latifolia, *Cavanilles, in St. Hil. Fl. Bras. Merr.* ii. 208. —Paraguay.
micrantha, *G. Don, Gen. Syst.* iii. 62.—Peruv.
multifida, Griseb. in Goett. Abh. xxiv. (1879) 189 = Hieronymi.

BLUMENBACHIA:—
multifida, *Hook. Bot. Mag.* t. 3599.—Reg. Argent.
palmata, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 208.—Paraguay.
parviflora, Gill. ex G. Don, Gen. Syst. iii. 69 = insignis.
Prietea, *C. Gay, Fl. Chil.* ii. 485.—Chili.
pterosperma, *G. Don, Gen. Syst.* iii. 69.—Peruv.
punices, *G. Don, l. c.*—Peruv.
scandens, *Meyen, Reise*, i. 810.—Chili.
sepiaria, *G. Don, Gen. Syst.* iii. 69.—Peruv.
syivestris, *Poepp. Fragm. Syn. Fl. Chil.* 25.—Chili.

BLUMEODENDRON, Kurz, in journ. As. Soc. Beng. xlii. (1873) ii. 245 = **Mallotus**, Lour. (Euphorb.).
Muelleri, Kurz, l. c. = Paracroton pendulus.
Tokbrai, Kurz, l. c.—Ins. Andaman.

BLUMIA, Nees, ex Blume, in Verh. Bat. Gen. ix. (1823) 147 = **Talauma**, Juss. (Magnol.).
Candollii, Nees, l. c.; et in Flora, viii. (1825) 152 = T. Candollii.

BLUMIA, Spreng. Syst. iii. 126 (1826) = **Saurauja**, Willd. (Ternstr.).
japonica, Steud. Nom. ed. II. i. 212, sphalm. = seq
javanica, Spreng. Syst. iii. 126 = S. Blumia.

BLUTAPARON, Rafin. New Fl. Am. iv. 45 (1836) **Philoxerus**, R. Br. (Amarant.).
brevifolium, Rafin. l. c. = P. vermicularis.
repens, Rafin. l. c. 46 = P. vermicularis.

BLYSMUS, Panz. ex Roem. & Schult. Syst. ii. Mant. 41 (1824) = **Scirpus**, Tourn. (Cyperac.).
Bonanni, Steud. Nom. ed. II. i. 212 = Schoenus Bonanni.
brevifolius, Decne. in Jacquem. Voy. Bot. 178. t. 178 = S. Caricis.
compressus, Panz. in Link, Hort. Berol. i. 278 = S. Caricis.
rufus, Link, l. c. = S. rufus.

BLYTTIA, Arn. in Mag. Zool. et Bot. ii. 430 (1838) = **Vincetoxicum**, Moench (Asclep.).
arabica, Arn. l. c. = V. fruticulosum.

BLYTTIA, Fries, Novit. Fl. Suec. Mant. ii. 2 (1839) = **Cinna**, Linn. (Gramin.).
suaveolens, Fries, l. c. = C. pendula.

BLYXA, Noronha, ex Thou. Gen. Nov. Madag. 4 (1806). *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 451.
DIPLOSIPHON, Decne. in Jacquem. Voy. Bot. 166. t. 167 (1844).
SAIVALA, Jones, in As. Res. iv. (1799) 275.
Aubertii, *Rich. in Mem. Inst. Fr.* (1811) 4.—Madag.
graminea, *Thou. ex Steud. Nom.* ed. II. i. 212.—Madag.
javanica, *Hassk. in Hove. & De Vriese, Tijdschr.* x. (1843) 121.—Java.
octandra, *Planch. ex Thw. Enum. Pl. Zey.* 282.—Zeylan.
Roxburghii, *Rich. in Mém. Inst. Fr.* (1811) 77. t. 5.—As. trop.; Austral.
Saivala, *Steud. Nom.* ed. II. i. 212.—Ind. or.

BOADSCHIA, All. Fl. Pedem. i. 248 = BOHADSCHIA, Crantz = **Peltaria**, Linn. (Crucif.).

BOARIA, DC. Prod. viii. 299 (1844) = **Maytenus**, Molina (Celastr.).
Molinae, DC. l. c. = M. Boaria.

BOBAEA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1890) 215 = **Bobea**, Gaudich. (Rubiac.).

BOBARTIA, Linn. Dias. Dens. 3 (1747); Fl. Zeyl. 17 (1747); Salish. in Trans. Hort. Soc. i. (1812) 313. *IRIDEAE*, Benth. & Hook. f. iii. 693.
HECASTA, Soland. ex Schum. in Skr. Naturk-Selsk. iii. (1792) 19.
aphylla, *Rev.-Suppl. Gen. Irid.* 90.—Afr. austr.
aurantiaca, Sweet, Hort. Brit. ed. III. 662 = Homeria aurantiaca.
filiformis, *Ker-Gawl. Gen. Irid.* 30.—Afr. austr.

BOBARTIA:—
gladiata, *Ker-Gawl. Gen. Irid.* 30; *Sweet, Hort. Brit.* ed. I. 396.—Afr. austr.
indica, *Linn. Sp. Pl.* 54.—Afr. austr.
juncea, Salisb. in Trans. Hort. Soc. i. (1812) 313 = indica.
Linnaei, Pritz. Ind. Ic. ii. 47 perperam = indica.
spathacea, Ker-Gawl. Gen. Irid. 30 = indica.
umbellata, *Ker-Gawl. l. c.* 31.—Afr. austr.

BOEBA, Gaudich. in Freyc. Voy. Bot. 473. t. 93 (1826). *RUBIACEAE*, Benth. & Hook. f. ii. 101.
BOBAEA, A. Rich. in Mém. Soc. Hist Nat. Par. v. (1830) 215.
BURNEYA, Cham. & Schlecht. in Linnaea, iv. (1829) 190.
brevipes, *A. Gray, in Proc. Am. Acad.* iv. (1860) 86.—Ins. Sandvic.
elatior, Gaudich. in Bot. Freyc. Voy. 473. t. 93 = Timonius Gaudichaudi.
glabra, Korth. in Nederl. Kruidk. Arch. ii. (1851) 11. 211 = Timonius Jambosella.
hirsutiuscula, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 247 = Polyphragmon sericeum.
inaequispepala, *Miq. Fl. Ind. Bat. Suppl.* 545.—Ins. Banca.
mutabilis, Korth. in Nederl. Kruidk. Arch. ii. (1851) 11. 212 = Polyphragmon mutabile.
putaminosa, F. Muell Fragm. iv. 92 = Guettardella putaminosa.
sericantha, *Miq. Fl. Ind. Bat.* ii. 855.—Malaya.
Wallichiana, Korth. in Nederl. Kruidk. Arch. ii. (1851) 11. 211 = Timonius Rumphii.

BOBU, Adans. Fam. ii. 88 (1763) = **Symplocos**, Linn. (Styrac.).

BOBUA, DC. Prod. iii. 23 (1828) = **Symplocos**, Linn. (Styrac.).
arborea, Miers, in Journ. Linn. Soc. xvii. (1879) 305 = S. arborea.
brevoridii, Miers, l. c. 304 = S. spicata.
baptica, Miers, l. c. 305 = S. baptica.
carnulescens, Miers, l. c. = S. caerulescens.
cerasifolia, Miers, l. c. 304 = S. cerasifolia.
glaucescens, Miers, l. c. 305 = S. glaucescens.
gracilis, Miers, l. c. = S. gracilis.
japonica, Miers, l. c. 306 = S. japonica.
lanceifolia, Miers, l. c. = S. lanceifolia.
laurina, DC. Prod. iii. 24 = S. spicata.
Lenormandiana, Miers, in Journ. Linn. Soc. xvii. (1879) 304 = S. Lenormandiana.
leptostachya, Miers, l. c. 306 = S. lanceifolia.
lucida, Miers, l. c. 305 = S. lucida.
montana, Miers, l. c. = S. montana.
myrtacea, Miers, l. c. 306 = S. myrtacea.
nerrifolia, Miers, l. c. = S. nerrifolia.
nitida, Miers, l. c. 305 = S. nitida.
oligostachya, Miers, l. c. 304 = S. spicata.
prunifolia, Miers, l. c. 306 = S. caudata.
rotundifolia, Miers, l. c. 305 = S. rotundifolia.
stravadioides, Miers, l. c. 304 = S. stravadioides.
theophrastaefolia, Miers, l. c. 306 = S. theophrastaefolia.
Vieillardi, Miers, l. c. 305 = S. Vieillardi.

BOCA, Vell. Fl. Flum. 283; v. t. 113 (1825) = **Banara**, Aubl. (Samyd.).
serrata, Vell. Il. cc. = Banara Vellozii.

BOCAGEA, St. Hil. Fl. Bras. Mer. i. 41. t. 9 (1825). *ANONACEAE*, Benth. & Hook. f. i. 29.
OXANDRA, A. Rich. Ess. Fl. Cub. 45. t. 8 (1842).
SAGERASEA, Dalz. in Hook. Kew journ. iii. (1851) 207.
acuminata, Steud. Nom. ed. II. i. 212 = Orophea acuminata.
alba, *St. Hil. Fl. Bras. Mer.* i. 42.—Bras.
aromatica, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 86.—N. Granat.
canescens, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 70.—Am. austr.
celebica, *Blume, Fl. Jav. Anon.* 88.—Ins. Celebes.
coriacea, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 96.—Zeylan.
corymbosa, *Blume, Fl. Jav. Anon.* 85.—Java.

BOCAGEA:—
Dalzellii, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 92.—Ind. or.
elliptica, *Hook. f. & Thoms. l. c.*—Burma; Malaya.
enneandra, *Spreng. Syst. iv. Cur. Post.* 144.—Java.
espintana, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 71.—Peruv.
Gaudichaudiana, *Baill. Adansonia*, viii. (1867–68) 183.—Cochinch.
heterantha, *Baill. l. c.* 178.—Madag.
hexandra, Blume, Fl. Jav. Anon. 83 = javensis.
javensis, *Spreng. Syst. iv. Cur. Post.* 144.—Java.
latifolia, *Blume, Fl. Jav. Anon.* 89.—Java.
leucoderma, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 71.—Bras.
longipedunculata, *Mart. Fl. Bras.* xiii. i. 45.—Bras.
lutea, Baill. Hist. Pl. i. 215 = Alphonsea lutea.
multiflora, *Mart. Fl. Bras.* xiii. i. 45.—Bras.
nervosa, Pierre, Fl. For. Cochin. sub t. 16 = Kingstonia nervosa.
obliqua, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 93.—Zeylan.
philastreana, *Pierre, Fl. For. Cochin.* t. 16.—Cochinch.
pisocarpa, *Blume, Fl. Jav. Anon.* 90. t. 45.—Java.
polyandra, *C. Presl, Rel. Haenk.* ii. 77.—Ins. Luzon.
polycarpa, Steud. Nom. ed. II. i. 212 = Orophea polycarpa.
Thwaitesii, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 92.—Zeylan.
ventricosa, Baill. Hist Pl. i. 215 = Alphonsea ventricosa.
verrucosa, Baill. l. c. sphalm. = praec.
viridis, *St. Hil. Fl. Bras. Mer.* i. 42. t. 9.—Bras.
Weddelliana, *Baill. Adansonia*, viii. (1867–68) 170.—Bras.

BOCCO, Steud. Nom. ed. I. 112 (1821) sphalm. = **Booea**, Aubl.

BOCCONIA, Plum. ex Linn. Gen. ed. I. 29 (1737). *PAPAVERACEAE*, Benth. & Hook. f. i. 53.
MACLEAYA, R. Br. App. Denh. & Clapp. Trav. 218, in adnot. (1826).
cordata, *Willd. Sp. Pl.* ii. 841.—China; Japon.
ferruginea, *Roesl, in Belg. Hortic.* xxiv. (1874) 39.—Peruv.
frutescens, *Linn. Sp. Pl.* 505.—Mexico; Peruv.
glauca, Salisb. Prod. 377 = frutescens.
integrifolia, *Humb. & Bonpl. Pl. Aequin.* i. 119. t. 35.—Mexico; Peruv.
japonica, Hort. ex André, in Rev. Hortic. (1866) 369 = cordata.
jodotnsis, Carr. l. c. 340 = cordata.
quercifolia, Moench, Meth. Suppl. 229 = frutescens.
sinuatifolia, Stokes, Bot. Mat. Med. iii. 8 = frutescens.
subtomentosa, L'Hérit. ex Stahl, Estud. Fl. Puertorico, Foll. ii. 34 = frutescens.

BOCKIA, Scop. Introd. 106 (1777) = **Mourirla**, Juss. (Melast.).

BOCOA, Aubl. Pl. Gui. Suppl. 38. t. 391 (1775) = **Inocarpus**, Forst. (Legum.).
edulis, Baill. Adansonia, ix. (1868–70) 287 = I. edulis.
guianensis, Steud. Nom. ed. I. 113 = seq.
prouacensis, Aubl. Pl. Gui. Suppl. 38.—Guiana.

BOCQUILLONIA, Baill. Adansonia, ii. 225 (1861–62). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 313.
brachypoda, *Baill. Adansonia*, xi. (1873) 127.—N. Caled.
brevipes, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 166.—N. Caled.
grandidens, *Baill. Adansonia*, xi. (1873) 128.—N. Caled.
sessiliflora, *Baill. l. c.* ii. (1861–62) 226.—N. Caled.
spicata, *Baill. l. c.* 227.—N. Caled.

BODWICHIA, Walp. Rep. i. 806 (1842), sphalm. = Bowdichia, H. B. & K. (Legum.).

BOEA, Comm. ex Lam. Encyc. i. 401 (1788). *GESNERACEAE*, Benth. & Hook. f. ii. 1028.
BAEA, Comm. ex Juss. Gen. 131 (1789).
CHLETERUS, Rafin. Prim. Somiol. 28 (1814).

BOEA:—
DOXOCERAS, Bunge, Enum. Pl. Chin. Bor. 54 (1833).
alata, Pers. Syn. i. 15 = Calceolaria petiolaris.
amplexicaulis, Parish, ex C. B. Clarke, Comm. et Cyrt. Beng. t. 84 = Phyllobora amplexicaulis.
bornensis, Scheff. ex H. O. Forbes, in Journ. Linn. Soc. xix. (1882) 298 = Didymocarpus Scheefferi.
Cavaleriei, F. Muell. ex C. B. Clarke, in DC. Monog. Phan. v. 145 = Conmersonii.
cochinchinensis, *C. B. Clarke, l. c.* 143.—Cochinch.
Commersonii, *R. Br. in Benn. Pl. Jav.* 120.—China.
dictyoneura, *Hance, in Journ. Bot.* xxi. (1883) 169.—China.
flosculosa, C. B. Clarke, Comm. et Cyrt. Beng. t. 83 = multiflora.
Hancei, C. B. Clarke, in DC. Monog. Phan. v. 144 = dictyoneura.
herbacea, *C. B. Clarke, l. c.* 142.—Ins. Formosa, hygrometrica, *R. Br. in Benn. Pl. Jav.* 120.—China.
hygroscopica, *F. Muell. Fragm.* iv. 146.—Austral.
magellanica, Lam. Illustr. i. 38. t. 15 = Commersonii.
Minahassae, *Teijsm. & Binn. in Tydschr. Nederl. Ind.* xxv. (1868) 415.—Ins. Celebes.
multiflora, *R. Br. in Benn. Pl. Jav.* 120.—Ind. or. philippensis, *C. B. Clarke, in DC. Monog. Phan.* v. 146.—Ins. Philipp.
plantaginea, Pers. Syn. i. 15 = Calceolaria plantaginea.
prasinuana, J. St. Hil. Expos. i. 279 = magellanica.
primuloides, Miq. in Ann. Mus. Lugd. Bat. iii. 190 = Oreocharis primuloides.
prolixa, *C. B. Clarke, in DC. Monog. Phan.* v. 143.—Burma.
punctata, Pers. Syn. i. 15 = Calceolaria punctata.
rufescens, *Franch. in Bull. Soc. Linn. Par.* i. (1885) 449.—China.
Swinhoii, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 291.—China.
Treubii, *H. O. Forbes, in Journ. Linn. Soc.* xix. (1882) 267.—Sumatra.
triandra, Pers. Syn. i. 15 = Calceolaria triandra.
Urvillei, *C. B. Clarke, in DC. Monog. Phan.* v. 147.—Ins. Walgiou.
violacea, Pers. Syn. i. 15 = Calceolaria violacea.
Wallichii, *R. Br. in Benn. Pl. Jav.* 120.—Burma.

BOEBERA, Willd. Sp. Pl. iii. 2125 (1803) = **Dyssodia**, Cav. (Compos.).
alternifolia, Moç. & Sessé, ex DC. Prod. v. 689 = Adenophyllum Porophyllum.
Cavanillesii, Spreng Syst. iii. 544 = Adenophyllum coccineum.
chrysanthemoides, Willd. Sp. Pl. iii. 2125.—D. chrysanthemoides.
fastigiata, H. B. & K. Nov. Gen. et Sp. iv. 198 = D. chrysanthemoides.
glandulosa, Pers. Syn. ii. 459 = D. chrysanthemoides, glandulosa.
incana, Lindl. Bot. Reg. t. 1602 = D. pubescens.
Porophyllum, H. B. & K. Nov. Gen. et Sp. iv. 198 = Adenophyllum Porophyllum.
pubescens, Spreng. Syst. iii. 544 = D. pubescens.
subintegerrima, Spreng. l. c. = D. subintegerrima.
tagetiflora, Spreng. l. c. = D. chrysanthemoides.

BOECKHIA, Kunth, Enum. Pl. iii. 448 (1841) = **Hypodiscus**, Nees (Restiac.).
laevigata, Kunth, l. c. 450 = H. albo-aristatus.
striata, Kunth, l. c. 449 = H. striatus.

BOEHMERIA, Jacq. Enum. Pl. Carib. 9 (1760).
URTICACEAE, Benth. & Hook. f. iii. 387.
GYMNOVINUS, F. Dietr. Ind. Sem. Hann. (1850).—Cf. Linnaea, xxix. (1857-58) 737.
DURETIA, Gaudich. in Bot. Freyc. Voy. 500 (1826).
SPITURERBERA, Miq. Comm. Phytogr. 153. t. 14 (1840).
acalyphoides, Blume, Mus. Bot. Lugd. Bat. ii. 225 = Pouzolzia scalyphoides.
acuminata, Hook. & Arn. ex Wedd. in DC. Prod. xvi. i. 285 = Pipturus albidus.
acuminata, Pers. Syn. ii. 556 = Elatostema acuminatum.
acuminata, *Wedd. in DC. Prod.* xvi. i. 209.—N. Caled.
albida, Hook. & Arn. Bot. Beech. Voy. 96 = Pipturus albidus.
alienata, Willd. Sp. Pl. iv. 341 = Pouzolzia indica.

BOEHMERIA:—
amplissima, Blume, Mus. Bot. Lugd. Bat. ii. 219 = stipularis.
ampullacea, H. B. & K. Nov. Gen. et Sp. ii. 35 = candata.
angustata, Hassk. Cat. Hort. Bog. Alt. 79 = Debregeasia velutina.
angustifolia, H. B. & K. Nov. Gen. et Sp. ii. 34 = Phenax angustifolius.
arachnoidea, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 423 = Pouzolzia macrura.
arborea, Desf. Cat. Hort. Par. ed. III. 348 = Gesnouinia arborea.
arborescens, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 541 = candata.
argentea, Guill. ex Wedd. in DC. Prod. xvi. i. 285[19] = Pipturus argenteus.
argentea, *Linden, in Belg. Hortic.* viii. (1858) 166.—Mexic.
argutа, *Mart. ex Miq. in Mart. Fl. Bras.* iv. i. 186.—Am. trop.
aspera, Blume, Mus. Bot. Lugd. Bat. ii. 206, in nota = Pouzolzia aspera.
aspera, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855-56) 349. t. 11.—N. Granat.
australis, *Endl. Prod. Fl. Norfolk.* 38.—Ins. Norfolk.
ballotaefolia, H. B. & K. Nov. Gen. et Sp. ii. 35 = Phenax ballotaefolius.
bifida, Blume, Mus. Bot. Lugd. Bat. ii. 222 = bilobа, Wedd.
bilobа, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 100.—Java.
bilobа, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 199.—Java.
Blumei, *Wedd. in DC. Prod.* xvi. i. 204.—Ins. Philipp.
Boiviniana, Blume, Mus. Bot. Lugd. Bat. ii. 227 = Pouzolzia Boiviniana.
boiviniana, Blume, l. c. 206, in nota = Pouzolzia Boiviniana.
brevirostris, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 201.—Peruv.
ballata, *H. B. & K. Nov. Gen. et Sp.* ii. 35.—Ecuador.
caerulea, Blume, Mus. Bot. Lugd. Bat. ii. 209 = Cyphophus coeruleus.
calophlebia, *C. Moore & F. Muell. in F. Muell. Fragm.* viii. 11.—Austral.
candicans, Hassk. Pl. Jav. Rar. 208; Cat. Hort. Bog. Alt. 79 = nivea.
candidissima, Hassk. Cat. Hort. Bog. Alt. 79 = Leucosyke javanica.
Candolleana, Gaudich. in Freyc. Voy. Bot. 499 = Pipturus argentea.
canescens, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 200 = platyphylla.
capensis, *Spreng. Syst.* iii. 844.—Afr. austr.
castaneaefolia, A. Cunn. ex Bureau, in DC. Prod. xvii 250 = Pseudomorus Brunoniana.
caudata, Poir. in Bonpl. Jard. Malm. t. 15 = platyphylla.
candata, *Sw. Prod. Veg. Ind. Occ.* 84.—Am. trop.
caudigera, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 199 = platyphylla.
celebica, *Blume, Mus. Bot. Lugd. Bat.* ii. 217.—Ins. Celebes.
celtidifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 34.—Ecuador.
cephalota, Pers. Syn. ii. 556 = Procris cephalida.
chinensis, Pers. l. c. 557 = Procris cantonensis.
cinerascens, Blume, Mus. Bot. Lugd. Bat. ii. 208 = Pouzolzia cinerascens.
cinerascens, Hassk. Cat. Hort. Bog. Alt. 79 = Pipturus argenteus.
cladomoides, Miq. Pl. Jungh. 34 = platyphylla.
cochinchinensis, Pers. Syn. ii. 557 = Procris cochinchinensis.
cochinchinensis, Spreng. Syst. iii. 844 = Pouzolzia indica.
cuneata, *Wedd. in DC. Prod.* xvi. i. 208 = sidaefolia.
compacta, *Blume, Mus. Bot. Lugd. Bat.* ii. 210.—Java.
conglobata, Blume, l. c. 204 = Pouzolzia conglobata.
cornuta, Blume, l. c. 208 = Cyphophus macrocephalus?
Cummingiana, Blume, l. c. ii. 199 = heterophylla.
cuspidata, *Blume, l. c.* 218 (sp. dubia).—Nepal.
cuspidata, Wedd. in Arch. Mus. Hist. Nat. Par. viii. (1855-56) 345 = ramiflora.
cylindrica, Sw. ex Blume, Mus. Bot. Lugd. Bat. ii. 218 = elongata.

BOEHMERIA:—
cylindrica, *Sw. Prod. Veg. Ind. Occ.* 84.—Am. bor. et austr.
dasyoda, Miq. in Mart. Fl. Bras. iv. i. 188 = cylindrica.
densiflora, *Hook. & Arn. Bot. Beech. Voy.* 271.—China.
depauperata, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 202 = malabarica.
dichotoma, Hassk. Cat. Hort. Bog. Alt. 79 = Debregeasia velutina.
didymogyne, *Wedd. in DC. Prod.* xvi. i. 204.—Burma
diffusa, Wedd. in Arch. Mus. Hist. Nat. Par. viii. (1855-56) 356 = sidaefolia.
discolor, Poepp. ex Blume, Mus. Bot. Lugd. Bat. ii. 206 = Pouzolzia discolor.
diversiflora, Miq. in Zoll. Syst. Verz. Ind. Archip. 101, 104 = Zollingeriana.
diversifolia, Miq. Fl. Jungh. 34 = Maoutia diversifolia.
diversifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 202 = Pavonii.
Drummondiana, Wedd. l. c. 201 = cylindrica.
elliptica, Wedd. l. c. 200 = platyphylla.
elongata, *Blume, Mus. Bot. Lugd. Bat.* ii. 218.—Hab ?
elongata, Hort. ex Steud. Nom. ed. II. i. 212 = platyphylla.
erythropoda, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 101, 104.—Malaya.
excelsa, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 202.—Ins. Juan Fernand.
fagifolia, Pers. Syn. ii. 556 = Elatostema fagifolium.
fallax, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855-56) 346.—Peruv.
Fernandesiana, C. Gay, Fl. Chil. v. 360 = Urtica glomerulacflora.
fagellifformis, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 310 = candata.
florida, *Miq. in Mart. Fl. Bras.* iv. i. 187.—Am. trop.
formicaria, Poepp. ex Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 201 = Pouzolzia formicaria.
frondosa, D. Don, Prod. Fl. Nep. 59 = Pouzolzia vimines.
frutescens, D. Don, l. c. = Maoutia Puya.
frutescens, Thunb. Fl. Jap. 367. n. 98 = Villebrunea frutescens.
glabra, Steud. Nom. ed. II. i. 192 = Neraudia melastomaefolia.
glochidiata, Hassk. Pl. Jav. Rar. 206 = Pouzolzia indica.
glomerulifera, Miq. in Zoll. Syst. Verz. Ind. Archip. 101, 104 = malabarica.
grandifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 199 = platyphylla.
grevicaefolia, *Presl, Bot. Bemerk.* 110.—Hab.?
guatemalana, Blume, Mus. Bot. Lugd. Bat. ii. 206 = Pouzolzia guatemalana.
Hamiltoniana, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 199 = platyphylla.
Harveyi, Seem. Fl. Vit. t. 62 = Cyphophus heterophyllus.
Helferi, *Blume, Mus. Bot. Lugd. Bat.* ii. 201.—Burma.
heterodon, *Blume, l. c.* 216 (sp. dubia).—Ind. or.
heterophylla, Blume, l. c. 204 = Blumei.
heterophylla, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855-56) 351.—Ins. Philipp.
hirta, Roth, ex Blume, Mus. Bot. Lugd. Bat. ii. 231 = Pouzolzia Rothiana.
hirta, Sw. Prod. Veg. Ind. Occ. 84 = Phenax urticifolius.
hirta, *Wedd. in Ann. Sc. Nat. Sér.* IV. i. (1854) 202.—Peruv.
hispidula, *Blume, Mus. Bot. Lugd. Bat.* ii. 228.—Sumatra.
holosericea, *Blume, l. c.* 221.—Japon.
Hugeliana, *Blume, l. c.* 218 (sp. dubia).—Ind. or.
humilis, Miq. Fl. Jungh. 33 = platyphylla.
hypoleuca, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 264 = Debregeasia hypoleuca.
incana, Hassk. Cat. Hort. Bog. Alt. 79 = Pipturus velutinus.
interrupta, Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 182 = platyphylla.
interruptа, Willd. Sp. Pl. iv. 340 = Fleurya interrupta.
Jacquiniana, *Blume, Mus. Bot. Lugd. Bat.* ii. 219.—Hab.?
Jamaicae-ferae, Bichold, in Verh. Bat. Gen. xii. (1830) 70 = Debregeasia edulis.

BOEHMERIA:—
japonica, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 131. —Japon.
javanica, Hassk. Cat. Hort. Bog. Alt. 79 = Fleurya interrupta.
laevis, Blume, Mus. Bot. Lugd. Bat. ii. 206 in nota = Pouzolzia virgata.
lateriflora, Muhl. ex Willd. Sp. Pl. iv. 842 = cylindrica.
latifolia, Blume, Mus. Bot. Lugd. Bat. ii. 207 = Cyphophalus latifoliatus.
Lindeniana, *Blume, l. c.* 226.—N. Granat.
littoralis, Sw. Prod. Veg. Ind. Occ. 84 = cylindrica.
longispica, Steud. in Flora, xxxiii. (1850) 260 = japonica.
lutescens, Blume, Mus. Bot. Lugd. Bat. ii. 209 = Cyphophalus lutescens.
macrophylla, Blume, l. c. 208, in obs. = Cyphophalus macrocephalus.
macrophylla, *Hornem. Hort. Hafn.* ii. 890.—Hab.?
macrophylla, *D. Don, Prod. Fl. Nep.* 60.—Reg. Himal.; Burma.
macrophylla, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 215 = japonica.
macrostachya, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 199 = platyphylla.
macrothyrsa, Miq. in Zoll. Syst. Verz. Ind. Archip. 100, 103 = Laportia decnmana.
maculata, Pers. Syn. ii. 556 = Elatostema maculatum, malabarica, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855–56) 355.—Ind. or.; Malaya.
mauritiana, Blume, Mus. Bot. Lugd. Bat. ii. 216 = platyphylla.
mauritiana, Wedd in Ann. Sc. Nat. Sér. IV. i. (1854) 200 = platyphylla.
melanocarpa, Blume, Mus. Bot. Lugd. Bat. ii. 210 = Cyphophalus melanocarpus.
melastomaefolia, Hook. & Arn. Bot. Beech. Voy. 96 = Neraudia melastomaefolia.
melastomoidea, Griff. Notul. iv. 886 = Pouzolzia pentandra.
microcarpa, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 201 = platyphylla.
microceltis, Miq. in Zoll. Syst. Verz. Ind. Archip. 100, 104 = Pouzolzia vimínea.
mollicoma, *Miq. in Zoll. l. c.*—Java.
mollis, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 208 = Cyphophalus macrocephalus.
moluccana, Blume, Mus. Bot. Lugd. Bat. ii. 208 = Cyphophalus macrocephalus.
moluccana, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 201 = Cyphophalus macrocarpus.
monticola, *Blume, Mus. Bot. Lugd. Bat.* ii. 227.—Sumatra.
nana, D. Don, Prod. Fl. Nep. 60 = Pouzolzia indica.
neglecta, *Blume, Mus. Bot. Lugd. Bat.* ii. 200.—Java.
nepalensis, Wedd. in Arch. Mus. Hist. Nat. Par. viii. (1855–56) 383 = Villebrunea frutescens.
nervosa, Madden, in Journ. As. Soc. Beng. xvii. (1848) 1. 408 = rugulosa.
nivea, *Gaudich. in Bot. Freyc. Voy.* 499.—As. trop. (enlt.).
nudiflora, *Willd. Sp. Pl.* iv. 843.—Venezuela.
oblíqua, Poepp. ex Blume, Mus. Bot. Lugd. Bat. ii. 206, adnot. = Pouzolzia obliqua.
odontophylla, Miq. Fl. Jungh. 35 = Maoutia odontophylla.
ouranthe, Miq. l. c. 33 = platyphylla.
ovalis, Miq. in Zoll. Syst. Verz. Ind. Archip. 100 = Pouzolzia vimínea.
ovata, Blume, Mus. Bot. Lugd. Bat. ii. 206, adnot. = Pouzolzia ovata.
ovata, Steud. Nom. ed. II. i. 212 = Neraudia melastomaefolia.
palmata, *Hassk. Cat. Hort. Bog. Alt.* 79.—Hab.?
parvifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 208 = Droguetia diffusa.
pauciflora, Blume, Mus. Bot. Lugd. Bat. ii. 201 = Droguetia diffusa.
Pavonii, *Wedd. in Ann. Sc. Nat. Sér.* IV. i. (1854) 202.—Peruv.
pendulifloræ, Wedd. l. c. 199 = macrophylla.
peruviana, Blume, Mus. Lugd. Bat. ii. 234 = candata.
petiolaris, H. B. & K. Nov. Gen. et Sp. ii. 36 = candata.
phyllostachya, Miq. in Mart. Fl. Bras. iv. i. 187 = cylindrica.
plicatinoda, Hassk. Cat. Hort. Bog. Alt. 79 = platyphylla.

BOEHMERIA:—
platanifolia, *Franch. & Sav. Enum. Pl. Jap.* i. 440.—Japon.
platyphylla, *D. Don, Prod. Fl. Nep.* 60.—As. et Afr. trop.; Ins. Pacif.
polystachya, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 200.—Reg. Himal.; Burma.
procridioides, Blume, Mus. Bot. Lugd. Bat. ii. 204 = Pouzolzia procridioides.
propinqua, Decne. Herb. Timor. 168 = Pipturus argenteus.
prostrata, Blume, Mus. Bot. Lugd. Bat. ii. 209 = Cyphophalus prostratus.
Puya, Hook. in Hook. Kew Journ. iii. (1851) 316 = Maoutia Puya.
racemiflora, Forst. ex Blume, Rumphia, iv. 37 = Phytocrene hirsuta.
radicans, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 85 = Pilea rhizobola.
ramentora, Bedd. Forest. Man. Bot. t. 27. f. 2 = malabarica.
ramiflora, *Jacq. Enum. Pl. Carib.* 31.—Ind. occ.; Mexic.
repanda, Hassk. Cat. Hort. Bog. Alt. 79 = Pipturus repandus.
repens, Nees & Mart in Nov. Act. Acad. Nat. Cur. xi. (1823) 86 = Pilea hirtella.
repens, *Wedd. in DC. Prod.* xvi. t. 201.—Cuba.
rhynchophylla, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 202.—Am. austr.
rigida, *Benth. in Hook. Niger Fl.* 519.—Afr. trop.
rotundifolia, D. Don, Prod. Fl. Nep. 60 = platyphylla.
rubescens, *Hassk. Cat. Hort. Bog. Alt.* 79 = Villebrunea rubescens.
rubescens, Jacq. Fragm. 8 = Gesnouinia arborea.
rugosa, Pers. Syn. ii. 556 = Phenax balloteafolius.
rugosissima, Miq. Fl. Jungh. i. 32 = platyphylla.
rugosissima, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 200.—Java.
rugulosa, *Wedd. l. c.* 200.—Reg. Himal.
salicifolia, D. Don, Prod. Fl. Nep. 60 = Debregeasia hypoleuca.
sanguinea, Hassk. Cat. Hort. Bog. Alt. 79 = Pouzolzia vimínea.
scabra, *Hassk. l. c.*—As. trop.?
scabrella, Gaudich. Bot. Freyc. Voy. 499 = platyphylla.
scabrella, Hort. ex Blume, Mus. Bot. Lugd. Bat. ii. 215 = candata.
sericea, Blume, Mus. Bot. Lugd. Bat. ii. 208 = Cyphophalus moluccanus.
sidæfolia, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 208.—Reg. Himal.; Burma; Java.
Siebaldiana, Blume, Mus. Bot. Lugd. Bat. ii. 220 = platyphylla.
spicata, Thunb. in Trans. Linn. Soc. ii. (1794) 330 = platyphylla.
spiciflora, *Blume, Mus. Bot. Lugd. Bat.* ii. 217 (sp. dubia).—Ind. or.
spicigera, *Spreng. Syst.* iii. 845.—Venezuela.
squamigera, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 208 = Debregeasia cuspidata.
stipularis, *Wedd. l. c.* 200.—Ins. Mascar.
strigosa, Blume, Mus. Bot. Lugd. Bat. ii. 226 = Pavonia.
strigosula, Blume, l. c. 203 = Pouzolzia vimínea.
subperforata, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855–56) 383 (sp. dubia).—Ind. or.
sumatrana, Miq. Fl. Jungh. 32 = platyphylla.
sylvatica, Hassk. Cat. Hort. Bog. Alt. 79 = Villebrunea sylvatica.
taitensis, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 200 = platyphylla.
tenacissima, Gaudich. Bot. Freyc. Voy. 500 = nivea.
ternifolia, D. Don, Prod. Fl. Nep. 59 = platyphylla.
tomentosa, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 200 = platyphylla.
travancorica, Bedd. Forest. Man. Bot. 225 = malabarica.
trinervata, Pers. Syn. ii. 556 = Gyrotaenia trinervata.
trinervis, Miq. Fl. Jungh. 36 = Pipturus repandus.
uliginosa, Mor. ex Wedd. in DC. Prod. xvi. 1. 236* = Phenax uliginosus.
ulmifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 209 = fallax.
urentissima, Blume, Mus. Bot. Lugd. Bat. ii. 221 = platyphylla.

BOEHMERIA:—
urticæfolia, Pers. Syn. ii. 556 = Phenax urticæfolius.
urticæfolia, Spreng. Syst. iii. 844 = Pilea rupipendia.
volutina, Decne. Herb. Timor. 168 = Pipturus velutinus.
vestita, Blume, Mus. Bot. Lugd. Bat. ii. 209 = Cyphophalus vestitus.
vialis, Cassr. ex Wedd. in DC. Prod. xvi. t. 235* = Phenax Sonnerattii.
viminea, Blume, Mus. Bot. Lugd. Bat. ii. 202 = Pouzolzia vimínea.
virgata, Guillem. in Ann. Sc. Nat. Sér. II. vii. (1837) 189 = platyphylla.
Waitsiana, Blume, Mus. Bot. Lugd. Bat. ii. 203 = Pouzolzia vimínea.
Wightiana, Blume, l. c. 217 = platyphylla.
Zollingeriana, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 201.—Java.

BOEHMERIA, C. B. Clarke, Comm. et Cyrt. Beng. 118. t. 85 ad 87 (1874). *GESNERACEÆ*, Benth. & Hook. f. ii. 1028
BARCA, C. B. Clarke, in DC. Monog. Phan. v. 134 (1883).
filiformis, *C. B. Clarke, Comm. et Cyrt. Beng.* 118. t. 87.—Reg. Himal.
fuiva, *C. B. Clarke, l. c.* t. 85.—Reg. Himal.
Griffithii, *C. B. Clarke, l. c.* t. 86.—Assam.
hirsuta, *C. B. Clarke, in DC. Monog. Phan.* v. 136.—Assam?
porosa, *C. B. Clarke, l. c.*—Burma.

BOELIA, Webb, Otia Hisp. 20. t. 15, 16 (1853) = **Genista**, Linn. (Legumin.).
sphærocarpa, Webb, l. c.—G. sphærocarpa.

BOENNINGHAUSENIA, Reichb. Consp. 197 (1828). *RUTACEÆ*, Benth. & Hook. f. i. 287.
PODOSTAURUS, Jungh. in Nat. en Geneesk. Arch. Nederl. Indie ii. (1845) 45.
albiflora, *Reichb. l. c.*—Reg. Himal.; Japon.
japonica, Sieb. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 21 = albiflora.

BOENNINGHAUSIA, Spreng. Syst. iii. 853, 845 (1826) = **Chastocalyx**, DC. (Legumin.).
vincentina, Spreng. l. c. 245 = C. vincentinus.

BOERHAVIA, Linn. Fl. Zeyl. 4 (1747) = **Boerhaavia**, Vaill. (Nyctag.).

BOERHAAVIA, Vaill. ex Linn. Syst. ed. I (1735); Linn. Gen. ed. I. 8 (1737). *NYCTAGINEÆ*, Benth. & Hook. f. iii. 5.
adscendens, Willd. Sp. Pl. i. 19 = repens.
africana, *Lour. Fl. Cochinch.* i. 16.—Afr. trop.
aggregata, Pav. ex Choisy, in DC. Prod. xiii. ii. 429 = Nyctaginia capitata.
angustifolia, *Linn. Syst.* ed. XII. 52 (sp. dubia).—Hab.?
angustifolia, Span. in Linnaea, xv. (1841) 342 = repens.
anisophylla, *Torr. in Bot. Mex. Bound.* 171.—Texas.
arborea, Roem. & Schult. Syst. i. 66 = Fiscnia hirtella.
arborescens, Lag. & Rodr. in Anal. Cienc. Nat. iv. (1801) 257 = Fiscnia hirtella.
ascendens, Schum. Beskr. Guin. Pl. 17 = repens.
bastica, Buch.-Ham. ex Wall. Cat. sub n. 6769 = repanda.
bicolor, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 168, 169 = repens.
bracteosa, *S. Wats. in Proc. Am. Acad.* xx. (1885) 370. —Texas.
Burchellii, *Choisy, in DC. Prod.* xiii. ii. 455.—Afr. austr.
caribæa, Jacq. Obs. Bot. iv. 5 = repens.
chaerophylloides, Willd. Sp. Pl. i. 22 = Astrephia chaerophylloides.
coccinea, Mill. Gard. Dict. ed. VIII. n. 4 = repens.
Commersonii, *Basill. in Bull. Soc. Linn. Par.* i. (1885) 484.—Madag.
cortensis, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 168 = repens.
crispa, *Heyne, in Wall. Cat.* n. 6771.—Ind. or.
decumbens, Vahl, Enum. i. 204 = paniculata.

BOERHAAVIA :—
depressa, *C. Sm. in Tuckey, Congo*, 249, nomen.—Ins. Cap. Verd.
diandra, Ambl. Pl. Gui. i. 4 = paniculata !
diandra, Burm. f. Fl. Ind. 3. t. 1 = erecta.
diandra, Linn. Sp. Pl. 1194 = hirsuta, paniculata.
dichotoma, *Hochst. ex Walp. Ann.* iii. 398.—Abyss.
dichotoma, Schimp. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 142 = grandiflora.
diffusa, Engelm. & A. Gray, in Boston, Journ. Nat. Hist. v. (1847) 259 = viscosa.
diffusa, Linn. Sp. Pl. 3 = repens.
diffusa, Sw. Obs. 10 = paniculata.
discolor, *H. B. & K. Nov. Gen. et Sp.* ii. 215.—Ecuador.
eglandulosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 169 = repens.
elegans, *Choisy, in DC. Prod.* xiii. 11. 453.—Nubia ; Arabia ; Ind. bor. occ.
elongata, Salisb. Prod. 56 = erecta.
erecta, Burm. f. Fl. Ind. t. 3. f. 2 = glutinosa.
erecta, Kll. Sketch, i. 41 = repens.
erecta, *Linn. Sp. Pl.* 3.—Am. bor. ; Ind. occ.
erionelena, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 292.—Mexic.
excelsa, Willd. Phyt. 1. n. 5 ; Sp. Pl. i. 21 = Oxyphaphus microcarpus !
fruticosa, *Dals in Dals. & Gibs. Fl. Bomb.* 213.—Ind. or.
fruticosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 167 = plumbaginea.
gibbosa, Pav. ex A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 293 = Senkenbergia annulata.
gibbosa, Pav. ex Choisy, in DC. Prod. xiii. 11. 467 = Tinantia gypsophiloides.
glabrata, Blume, Bijdr. 733 = repens.
glandulosa, *Anderss. Om Oarn. Galapagos.* 64.—Ins. Galap.
glomerata, *Klotzsch, in Schomb. Fauna et Fl. Guy.* 1181, nomen.—Guiana.
glutinosa, Miers, ex Bert. in Merc. Chil. (1829) 596 = discolor.
glutinosa, Vahl, Enum. i. 287 = diffusa !
Grahamii, A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 293 = scandens.
grandiflora, A. Rich. Tent. Fl. Abyss. ii. 209 = plumbaginea.
Helenæ, Koem. & Schult. Syst. Mant. i. 78 = verticillata.
hirsuta, *Linn. Mant.* 1170.—Ind. occ. ; Am. bor.
hirsuta, Willd. Phyt. 1 = repens.
insularis, Boj. Hort. Maurit. 264 = repens.
laxa, Pers. Syn. i. 36 = paniculata.
leiocolena, *Torr. Bot. Mex. Bound.* 172.—Texas.
libyca, *Pomel, Nouv. Mat. Fl. Atl.* 336.—Algeria.
linearifolia, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 293.—N. Mexic.
littoralis, *H. B. & K. Nov. Gen. et Sp.* ii. 216.—Peruv.
maroccana, Ball, in Journ. Bot. xiii. (1875) 176 = repens.
microphylla, Boiv, ex Steud. Nom. ed. II. i. 213 (orbo ignot.).—Ins. Maurit.
minutiflora, Span. in Linnæa, xv. (1841) 342 = repens.
nudicaulis, R. Br. Prod. 422 = repens.
nantocana, Schan. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 402 = viscosa.
nudicaulis, *Phil. in Linnæa*, xxix. (1857-58) 37.—Chili.
obtusifolia, Lam. Illustr. i. 10 = patula.
obtusifolia, Steud. Nom. ed. II. i. 213 = paniculata.
Palmeri, *S. Wats. in Proc. Am. Acad.* xvii. (1882-83) 149.—Texas.
paniculata, Lam. Illustr. i. 10 = hirsuta, erecta.
paniculata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 105.—Texas ; Mexic.
patula, *Dondo, ex Vahl, Enum.* i. 287.—Peruv.
pedunculosa, *A. Rich. Tent. Fl. Abyss.* ii. 210.—Abyss.
pentandra, Burch. Trav. i. 489 = Burchellii.
periplocæfolia, *Comm. ex Vahl, Enum.* i. 290.—Madag.
periplocifolia, Sieber, ex Choisy, in DC. Prod. xiii. 11. 453 = repens.
pilosa, *Boj. Sme Rapp. Soc. Hist. Nat. Maurit.* 43 (sp. dub.).—Maurit. !
plumbaginea, *Cav. Ic.* ii. 7. t. 112.—Hispan. ; Afr. trop. et austr. ; Syria.
polymorpha, Kth. in Act. Soc. Hist. Nat. Par. i. (1792) 105 = caribæa, hirsuta, repens.
procumbens, Banks, ex Roxb. Fl. Ind. i. 146 = repens.
pterocarpa, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 376.—Am. bor.
pubescens, R. Br. Prod. i. 422 = repens.
pulchella, *Griseb. in Goett. Abh.* xix. (1874) 88.—Reg. Argent.
pulverulenta, Dupuis, in Vahl, Enum. i. 291 = dichotoma.
purpurascens, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 291.—N. Mexic.
Rebouillana, *Pomel, Nouv. Mat. Fl. Atl.* 335.—Algeria.
repanda, Kotschy, ex Choisy, in DC. Prod. xiii. 11. 454 = verticillata.
repanda, Roxb. Hort. Beng. 2 ; in Beatson, St. Helena Tracts, 302 = verticillata.
repanda, *Willd. Sp. Pl.* i. 22.—As. trop. ; Austral.
repens, *Linn. Sp. Pl.* 3.—Cosmop. trop.
repens, Sieber.—Cf. Schweinf. Beitr. Fl. Aethiop. 168 = plumbaginea.
rhomboidea, Humb. in Spreng. Jahrb. Gew. i. 111. 66 = Cryptocarpus globosus.
Rottleri, Steud. Nom. ed. II. i. 213 = crispa !
rubicunda, Steud. l. c. = elegans.
rupens, Kotth. ex Wall. Cat. sub n. 6771 = crispa.
rupestris, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 169 = repens.
sarmentosa, Willd. Sp. Pl. i. 21 = scandens.
scabrida, Steud. & Hochst. ex Steud. Nom. ed. II. i. 213 = discolor.
scandens, Drège, ex Choisy, in DC. Prod. xiii. 11. 455 = Burchellii.
scandens, Forsk. Fl. Aegypt. Arab. 3 = plumbaginea.
scanderes, Gibs. Cat. Bomb. Fl. 167 = verticillata.
scandens, *Linn. Sp. Pl.* 3.—Am. bor. ; Ind. occ.
Schomburgkiana, *Oliver, in Hook. Ic. Pl.* t. 1325.—Austral.
sessilifora, Steud. Nom. ed. II. i. 213 = repens.
spicata, *Choisy, in DC. Prod.* xiii. 11. 456.—Am. bor. occ. stellata, *Boj. Sme Rapp. Soc. Hist. Nat. Maurit.* 43.—Ins. Maurit. !
stellata, Wight, Ic. t. 876 = verticillata.
suberosa, C. Sm. in Tuckey, Congo, 249 = repens.
sumata, Zipp. ex Span. in Linnæa, xv. (1841) 342 = repens.
surinamensis, Miq. l. c. xviii. (1844) 244 = repens.
tetrandra, *Forst. f. Prod.* 2. n. 5.—Ins. Societ.
tomentosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 168 = repens.
tuberosa, *Lam. Illustr.* i. 10.—Peruv.
umbellata, Wight, ex Hook. f. Fl. Brit. Ind. iv. 709 = repanda.
undulata, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 168 = repens.
verticillata, Boiss. Fl. Orient. iv. 1044 = plumbaginea.
verticillata, *Poir. Encyc.* v. 56.—Afr. trop. ; Syria ; Ind. or.
verticillata, Rottl. in Wall. Cat. n. 6772 = crispa.
virgata, *H. B. & K. Nov. Gen. et Sp.* ii. 218.—N. Mexic.
viscosa, Fresen. in Mus. Senckenb. i. (1834) 75 = repens.
viscosa, *Lag. & Rodr. in Anal. Cienc. Nat.* iv. (1801) 256. n. 13.—Peruv.
viscosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 167 = plumbaginea.
vulcanicifolia, Pelt. Encyc. v. 55 = repens.
Wrightii, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 292.—N. Mexic.

BOGENHARDIA, Reichb. Nom. 200 (1841) = Abutilon, Tourn. (Malvac.).
crispa, Reichb. l. c. in nota = A. crispum.

BOHADSCHIA, Crantz, Stirp. Austr. ed. I. fasc. i. 2 (1762) ; ed. II. 5 (1769) = Peltaria, Linn. (Crucif.).
alliacea, Crantz, l. c. = P. alliacea.

BOHADSCHIA, Presl, Rel. Haenk. ii. 98. t. 68 (1831) = Turnera, Linn.
humifusa, Presl, l. c. = T. diffusa.
microphylla, Griseb. Cat. Fl. Cub. 114 = T. diffusa.

BOHADSCHIA, F. W. Schmidt, Samml. Phys. Aufs. i. 279 (1795) = Hyoseris, Linn. &c. (Compos.).
hirta, F. W. Schmidt, l. c. 280 = Leontodon hirtus.
lucida, F. W. Schmidt, l. c. 279 = H. lucida.

BOHERAVIA, Pavoli, in Anal. Soc. Cient. Argent. v. 210 (1878) ; Contrib. 59 = Boerhaavia, Vaill. (Nyctag.).

BOISDUVALIA, Spach, Hist. Vég. Phan. iv. 388 (1835) = Oenothera, Linn. (Onagrar.).
andina, Phil. in Linnæa, xxxiii. (1864-65) 70 (= O. andina !).—Chili.
elevistegama, Curran, in Bull. Calif. Acad. (1884) 12.—Calif.
concinna, Spach, Hist. Vég. Phan. iv. 384 = O. concinna.
densiflora, S. Wats. Bot. Calif. i. 238 = O. densiflora.
Douglasii, Spach, Hist. Vég. Phan. iv. 385 = O. densiflora.
glabella, Walp. Rep. ii. 89 = O. glabella.
Tocornalii, C. Gay, Fl. Chil. ii. 344.—Chili.
Torreyi, S. Wats. Bot. Calif. 238 = O. Torreyi.
Valchmanni, Phil. in Linnæa, xxx. (1859-60) 187.—Chili.

BOISSIAEA, Lem. in Orb. Dict. ii. 338 (1842) = Bossiaea, Vent. (Legumin.).

BOISSIERA, Domb. ex DC. Syst. i. 512, in syn. (1818) = Lardizabala, Ruiz & Pav. (Berberid.).
trifoliata, Domb. ex DC. l. c. 513 = L. trifoliolata.
triternata, Domb. ex DC. l. c. 512 = L. biternata.

BOISSIERA, Haenseler, ex Willk. & Lange, Prod. Fl. Hisp. i. 218 (1861) = Gagea, Salisb. (Liliac.).
baetica, Haenseler, l. c. = G. polymorpha.

BOISSIERA, Hochst. & Steud. in Flora, xxi. (1838) i. Intell. 25. *GRAMINEÆ*, Benth. & Hook. f. iii. 1176.
SCHNEIZLEINIA, Steud. Nom. ed. II. i. 218, in syn. (1840).
WIBERIA, Boiss. Fl. Orient. v. 659 (1884).
bromoides, *Hochst. & Steud. l. c. ; Syn. Pl. Gram.* 200.—Arab. ; Syria ; Persia ; Afghan.
Danthoniæ, A. Braun, Ind. Sem. Hort. Berol. (1857) 3 = Bromus macrostachys.
Pumilio, *Staff, in Denkschr. Acad. Wien*, i. (1885) 9.—Persia.

BOJERIA, DC. Coll. Mém. ix. 18 ; et Prod. v. 94 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 331.
glabra, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 364.—Afr. trop.
speciosa, *DC. Coll. Mém.* ix. 18. t. 3.—Madag.

BOJERIA, Rafin. Fl. Tellur. iv. 50 (1836) = Amomum, Linn. (Scitam.).
magnifica, Rafin. l. c. = A. magnificum.

BOLANDRA, A. Gray, in Proc. Am. Acad. viii. (1868) 341. *SAXIFRAGEÆ*.
californica, *A. Gray, l. c.*—Calif.
oregana, *S. Wats. l. c.* xiv. (1879) 292.—Am. bor. occ.

BOLANOSA, A. Gray, Pl. Wright. i. 82 (1852). *COMPOSITÆ*, Benth. & Hook. f. ii. 332.
Coulteri, *A. Gray, l. c.*—Mexic.

BOLAX, Comm. ex Juss. Gen. 236 (1789) = Azorella, Lam. (Umbell.).
angulatus, Presl, ex DC. Prod. iv. 80 = Mulinum angustum.
arctioides, Spreng. Umb. Sp. 11 = A. arctioides.
caespitosus, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zel. 63 = A. caespitosa.
caespitosus, Spreng. in Roem. & Schult. Syst. vi. 356 = A. arctioides.
cladorrhizus, Spreng. l. c. 362 = A. cladorrhiza.
conglomerata, Spreng. Umb. Sp. 9, 10 = A. caespitosa.
crenatus, Spreng. Syst. i. 880 = A. crenata.
Echinus, Presl, ex DC. Prod. vi. 79 = Mulinum Echinus.

BOLAX :—
flamentosus, Spreng. in Roem. & Schult. Syst. vi. 359 = A. filamentosa.
floccipes, Sieber, ex Benth. Fl. Austral. iii. 380 = Astrotricha floccosa.
geranifolius, Presl, ex DC. Prod. iv. 81 = Huanaca geranifolia.
Gilliesii, Hook. Bot. Misc. i. (1830) 325 = A. Gilliesii.
Glebaria, Comm. ex Lam. Encyc. iii. 156 = A. caespitosa.
gummifer, Spreng. Syst. i. 879 = A. caespitosa.
integrifolius, Spreng. ex Steud. Nom. ed. II. i. 214 = A. filamentosa.
imbifolius, Sieber, ex DC. Prod. v. 74 = Astrotricha ledifolia.
lycopodioides, Spreng. in Roem. & Schult. Syst. vi. 861 = A. corymbosa.
multifidus, Spreng. Syst. i. 880 = A. multifida.
pedunculatus, Spreng. Umb. Sp. 10 = A. multifida.
pilifera, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 82.—Peruv.
prolifer, Spreng. in Roem. & Schult. Syst. vi. 861 = Mulinum proliferum.
spinosus, Spreng. l. c. 862 = Mulinum spinosum.
tridentatus, Spreng. Umb. Prod. 84 = Hydrocotyle tridentata.
trifurcatus, Spreng. ex Steud. Nom. ed. II. i. 214 = A. caespitosa.
trilobus, Spreng. Umb. Prod. 84 = Hydrocotyle triloba.

BOLBIDIUM, Lindl. Veg. Kingd. 161 (1847), nomen (Orchid.). In Herb. Lindl. includit species Dendrobii (D. subacaule & prasinum).

BOLBOPHYLLARIA, Reichb. f. in Bot. Zeit. x. (1852) 984 = Bulbophyllum, Thou. (Orchid.).
aristata, Reichb. f. Beitr. Orch. Centr. Am. 60 = Bulbophyllum aristatum.
biseta, Reichb. f. in Walp. Ann. vi. 242 = Bulbophyllum bisetum.
bracteolata, Reichb. f. in Bot. Zeit. x. (1852) 984 = Bulbophyllum bracteolatum.
clavata, S. Moore in Baker, Fl. Maurit. 346 = Bulbophyllum clavatum.
Oerstedii, Reichb. f. in Bonplandia, iii. (1855) 228 = Bulbophyllum Oerstedii.
pachyrhachis, Reichb. f. in Walp. Ann. vi. 241 = Bulbophyllum pachyrhachis.
sordida, Reichb. f. l. c. 242 = Bulbophyllum bracteolatum.

BOLBOPHYLLOPSIS, Reichb. f. in Bot. Zeit. x. (1852) 953 = Cirrhopetalum, Lindl. (Orchid.).
maculosa, Reichb. f. in Walp. Ann. vi. 243 = C. maculosum.
morphologorum, Reichb. f. in Bot. Zeit. x. (1852) 953 = C. maculosum.

BOLBOPHYLLUM, Spreng. Syst. iii. 681, 732 (1826) = Bulbophyllum, Thou. (Orchid.).

BOLBORCHIS, Zoll. & Moritzi, Syst. Verz. Zoll. 89 (1845–48) = Coelogyne, Lindl. (Orchid.).
crocyfrons, Zoll. & Moritzi, l. c. = C. javanica.

BOLBOSTYLIS, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 467 = Bulbostylis, Tourn. (Compos.).

BOLDEA, Juss. in Ann. Mus. Par. xiv. (1809) 134 = Peumus, Molina (Monim.).
fragrans, C. Gay, Fl. Chil. v. 358; et Tul. in Arch. Mus. Par. viii. (1855) 412. t. 31 = P. Boldus.

BOLDOA, Endl. Gen. Suppl. i. 1378 (1841) = BOLDEA, Juss. = Peumus, Molina (Monim.).

BOLDOA, Cav. Cat. Hort. Madr. (1803) t. 7; Lag. Gen. et Sp. Nov. 10 (1816). *NYCTAGINEÆ*, Benth. & Hook. f. ill. 5.
SALPIANTHUS, Humb. & Bonpl. Pl. Aequin. i. 154. t. 44 (1808).
lanceolata, *Lag. Gen. et Sp. Nov.* 10.—Mexic.
ovatifolia, Lag. l. c. = Cryptocarpus globosus.
paniculata, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 256 = Cryptocarpus globosus.
purpurascens, Cav. Hort. Madr. t. 7 = Cryptocarpus globosus.
repens, *Spreng. Syst.* i. 179.—Braz.

BOLDU, Feuill. ex Adans. Fam. ii. 446 (1763) = Peumus, Molina (Monim.).

BOLDU, Nees, Progr. 11 (1833); Syst. Laurin. 177 (1836) = Cryptocaarya, R. Br. (Laurin.).
chilanum, Nees, Syst. Laurin. 178 = Peumus Boldus.
chilense, Poepp. ex Meisn. in DC. Prod. xv. i. 67 = pruce.
nitidum, Meisn. l. c. 506 = C. nitida.

BOLDUCIA, Neck. Elem. iii. 32 (1790) = Dipteryx, Schreb. (Legumin.).

BOLDUS, Schult. f. in Roem. & Schult. Syst. vii. p. x. 56 (1829) = Peumus, Molina (Monim.).
chilensis, *Schult. f. l. c.* 56.—P. Boldus.

BOLELIA, Rafin. Atl. Journ. 130 (1832) = Downingia, Torr. (Campan.).

BOLEMUM, Desv. Journ. Bot. iii. (1814) 163. t. 26. *CRUCIFERÆ*, Benth. & Hook. f. i. 98.
asperum, *Desv. l. c.* 163, 175.—Hispan.

BOLINA, Rafin. Sylva Tellur. 101 (1838) (Melastomac.).
conferta, Rafin. l. c. = Brachyotum confertum.
divaricata, Rafin. l. c. = Pterogastra divaricata.
ericæfolia, Rafin. l. c. = Centronia ericæfolia.

BOLIVARIA, Cham. & Schlecht. in Linnæa, i. (1826) 207. t. 4 = Manodora, Humb. & Bonpl. (Oleaceæ).
chloranganthe, Remy, in Ann. Sc. Nat. Sér. III. viii. (1847) 225.—Bolivia.
decandra, Gill. ex Hook. Journ. Bot. i. (1834) 284.—Chili.
Griesebachii, A. Scheele, in Linnæa, xxv. (1852) 254 = M. heterophylla.
integrifolia, Cham. & Schlecht. l. c. i. (1826) 208 = M. integrifolia.
mexicana, A. DC. Prod. viii. 315 = M. mexicana.
pinnatifida, Schlecht. in Linnæa, xxvi. (1853–55) 474 = M. pinnatifida.
robusta, Benth. in Hook. Lond. Journ. Bot. v. (1846) 190. t. 8 = M. robusta.
trifida, Cham. & Schlecht. in Linnæa, i. (1826) 209 = M. trifida.

BOLLÆA, Parl. in Bull. Soc. Bot. Fr. v. (1858) 509 = Pancratium, Linn. (Amaryll.).
canariensis, Parl. l. c. = P. canariense.

BOLLEA, Reichb. f. in Bot. Zeit. x. (1852) 667 = Zygopetalum, Hook. (Orchid.).
coelestis, Reichb. f. in Gard. Chron. (1876) i. 756 = Z. coelestre.
guianensis, Klotzsch, in Schomb. Fauna & Fl. Guy. 1206, nomen.—Guiana.
Lalindri, Reichb. f. in Gard. Chron. (1874) ii. 33 = Z. Lalindei.
Lawrenceana, Reichb. f. l. c. (1878) ii. 266 = Z. Lawrenceana.
pallens, Reichb. f. l. c. (1881) i. 462.—Hab.?
Patini, Reichb. f. l. c. (1874) ii. 34 = Z. Patini.
pulvinaris, Reichb. f. in Linnæa, xli. (1877) 107 = Z. pulvinare.
violacea, Reichb. f. in Bot. Zeit. x. (1852) 668 = Z. violaceum.

BOLOPHYTA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 347 = Parthenium, Linn. (Compos.).
alpina, Nutt. l. c. 348 = P. alpinum.

BOLOSIA, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 211 (1870) = Hispidella, Barnad. (Compos.).
piloselloides, Pourr. ex Willk. l. c. = H. hispanica.

BOLTONIA, L'Hérit. Sert. Angl. 27 (1788). *COMPOSITÆ*, Benth. & Hook. f. ii. 269.
ASTEROMOEA, Blume, Bijdr. 901 (1827).
DICHAETOPHORA, A. Gray, Fl. Fendl. 73 (1849).
HISUTOSA, DC. Prod. vi. 44 (1837).
MADIA, L'Hérit. ex DC. Prod. v. 201 (1836).
asteroides, Sims, Bot. Mag. t. 2554 = diffusa.
asteroides, *L'Hérit. Sert. Angl.* 27.—Am. bor.
campestris, *Hemsl. Biol. Centr. Am. Bot.* ii. 190.—Texas.
castaneiensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 398.—Japon.
diffusa, *Ell. Sketch*, ii. 400.—Am. bor.

BOLTONIA :
glastifolia, L'Hérit. Sert. Angl. 27 = asteroides.
incisa, *Benth. Fl. Hongk.* 178 = Aster incisus.
indica, *Benth. l. c.* 174 = Aster indicus.
integrifolia, Benth. & Hook. f. Gen. ii. 269 = pekinensis.
japonica, *Benth. & Hook. f. l. c.*—Japon.
latisquama, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 238.—Am. bor.
lautureana, Debeaux, in Act. Linn. Soc. Bord. xxxi. (1877) 215 = Aster lautureanus.
pekinensis, *Hance, in Journ. Bot.* v. (1867) 370.—China.
Seemannii, Benth. & Hook. f. Gen. ii. 269 = Achaetogeron Seemannii

BOLUSIA, Benth. in Hook. Ic. Pl. xii. 57. t. 1163 (1878). *LEGUMINOSÆ*.
capensis, *Benth. l. c.*—Afr. austr.

BOMAREA, Mirb. in Hist. Nat. Pl. ix. (1804) 71. *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 736.
COLLANIA, Herb. Amaryll. 103. t. 8 (1837).
DANBYA, Salisb. Gen. Pl. Fragm. 57 (1866).
DODECASPERMA, Rafin. Fl. Tellur. iv. 25 (1836).
SPHAERINE, Herb. Amaryll. 106. t. 19, 18, 16, 18 (1837)
VANDESIA, Salisb. in Trans. Hort. Soc. i. (1812) 282.
WICHURARA, M. Roem. Syn. Ensat. 277 (1847).
acutifolia, *Herb. Amaryll.* 112.—Mexic.
affinis, Kunth, Enum. Pl. v. 794 = edulis.
amoena, *M. Roem. Syn. Ensat* 274.—Peruv.
anceps, Herb. Amaryll. 116 = fimbriata.
andimarcana, *Baker, in Journ. Bot.* xx. (1882) 201.—Peruv.
Andreana, *Baker, l. c.* 205.—N. Granat.
angulata, *Benth. Pl. Hartw.* 156.—Peruv.
angustifolia, *Benth. l. c.*—Peruv.
angustipetala, *Baker, in Journ. Bot.* xx. (1882) 202.—N. Granat.
aurantiaca, *Herb. Amaryll.* 899. t. 46. f. 1.—Peruv.
brachysepala, *Benth. Pl. Hartw.* 157.—Peruv.
bractiata, Herb. Amaryll. 112. t. 8. f. 3 = edulis.
Brauniana, Schenk, in Mart. Fl. Bras. iii. t. 168 = edulis.
Bredemeyerana, Herb. Amaryll. 118 = multiflora.
brevis, *Baker, in Journ. Bot.* xx. (1882) 202.—Peruv.
Caldasiana, *Herb. Amaryll.* 119.—Guatemala; Peruv.
caracasensis, Herb. l. c. 111. t. 16. f. 2 = edulis.
Carderi, *Mast. in Gard. Chron.* (1876) i. 793, 795.—N. Granat.
chontalensis, Seem. in Gard. Chron. (1871) 479 = edulis.
coccinea, *Baker, in Journ. Bot.* xx. (1882) 202.—Peruv.
conferta, Benth. Pl. Hartw. 259 = patacocensis.
cordifolia, *Herb. Amaryll.* 118.—Peruv.
cornigers, *Herb. l. c.* 116. t. 17. f. 2.—Peruv.
cornuta, Herb. l. c. 114. t. 17. f. 4 = edulis.
crinita, *Herb. l. c.* 119. t. 15. f. 4.—Peruv.
croces, *Herb. l. c.* 119.—Peruv.
cumbrensis, Herb. l. c. 115. t. 18. f. 2 = torta.
decimata, Klotzsch, ex Kunth, Enum. Pl. v. 802 = dispar.
densiflora, Herb. Amaryll. 899. t. 46. f. 4 = tomentosa.
diffracta, *Baker, in Journ. Bot.* xx. (1882) 206.—N. Granat.
denticulata, *Herb. Amaryll.* 118.—Peruv.
dispar, *Herb. l. c.* 113. t. 17. f. 1.—Peruv.
disstitifolia, *Baker, in Journ. Bot.* xx. (1882) 203.—Ecuador.
distichophylla, *Baker, l. c.* 203.—Peruv.
edulis, *Herb. Amaryll.* 111.—Am. trop.
Fanningiana, Herb. l. c. 116. t. 16. f. 3 = multiflora.
fimbriata, *Herb. l. c.* 116.—Peruv.
floribunda, Herb. l. c. = fimbriata.
formosissima, *Herb. l. c.* 111. t. 14. f. 4.—Peruv.
frondea, *Mast. in Gard. Chron.* (1883) i. 668.—N. Granat.
furcata, Klotzsch, ex Kunth, Enum. Pl. v. 792 = edulis.
fuscata, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1190, nomen.—Guiana.
glaucescens, *Baker, in Journ. Bot.* xx. (1882) 201.—Ecuador.
glomerata, *Herb. Amaryll.* 115. t. 15. f. 1.—Peruv.
gomjocalon, *Baker, in Journ. Bot.* xx. (1882) 204.—Ecuador.

BOMAREA:—
gloriosa, M. Roem. Syn. Ensat. 269 = edulis.
granatensis, M. Roem. l. c. 264 = B. Salsilla.
grandifolia, Herb. Amaryll. 113 = edulis.
Hallians, *Herb. Amaryll.* 117. t. 10. f. 3.—N. Granat.
Hartwegii, *Baker, in Journ. Bot.* xx. (1882) 203.—Ecuador.
hirsuta, *Herb. Amaryll.* 114.—N. Granat.
hirta, *Schenk, in Mart. Fl. Bras.* iii. t. 169.—Bras.
hirtella, Herb. Amaryll. 113 = edulis.
hispida, *Baker, in Journ. Bot.* xx. (1882) 202.—N. Granat.
Hookeriana, *Herb. Amaryll.* 398. t. 46. f. 8.—Peruv.
involucrosa, *Baker, in Journ. Bot.* xx. (1882) 201.—Peru.
Jacquesiana, *Kunth, Enum. Pl. v.* 800.—Bras.
jamirensis, M. Roem. Syn. Ensat. 267 = edulis.
Kalbreyeri, *Baker, in Journ. Bot.* xx. (1882) 204.—N. Granat.
lancifolia, *Baker, l. c.* 202.—Ecuador.
latifolia, Herb. Amaryll. 113 = edulis.
Lehmanni, *Baker, in Journ. Bot.* xxi. (1883) 373.—Am. austr.
linifolia, *Baker, l. c.* xx. (1882) 201.—N. Granat.
longipes, *Baker, l. c.* 204.—Ecuador.
lutea, *Herb. Amaryll.* 120. t. 5.—Peruv.
lyncina, *Herb. l. c.* 398. t. 46. f. 8.—Peruv.
Mankiana, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xx. (1852) 387.—Venezuela.
Macleanica, Herb. in Lindl. Bot. Reg. (1842) Misc. 66 = aurantiaca.
macrocarpa, Herb. Amaryll. 14 = edulis.
maranensis, *Herb. l. c.* 17.—Bras.
Martiana, Schenk, in Mart. Fl. Bras. iii. t. 170 = edulis.
miniata, Kunth, Enum. Pl. v. 792 = edulis.
minima, *Baker, in Journ. Bot.* xx. (1882) 202.—N. Granat.
Moritziana, Klotzsch, ex Kunth, Enum. Pl. v. 797 = edulis.
multiflora, *Mirb. Hist. Nat. Pl.* ix. (1804) 72.—N. Granat.
multiplex, *Benth. Pl. Hartw.* 156.—Peruv.
nervosa, *Baker, in Journ. Bot.* xx. (1882) 202.—Peruv.
nobilis, Herb. Amaryll. 422 = ovata.
obovata, Herb. l. c. 113. t. 14. f. 2 = edulis.
oculata, M. Roem. Syn. Ensat. 268 = B. Salsilla.
oligantha, *Baker, in Gard. Chron.* (1877) ii. 648.—Peruv.
ovata, *Mirb. Hist. Nat. Pl.* ix. (1804) 72.—Am. austr.
pachyphlebia, *Baker, in Journ. Bot.* xx. (1882) 204.—Ecuador.
pardina, *Herb. Amaryll.* 120. t. 14. f. 1.—N. Granat.
patersumensis, Herb. l. c. 422 fimbriata.
patacocensis, *Herb. l. c.* 120. t. 14. f. 3.—Ecuador; N. Granat.
paucifora, Herb. l. c. 112 = edulia.
platypetala, Benth. Pl. Hartw. 156 = uncifolia.
podopetala, *Baker, in Journ. Bot.* xx. (1882) 202.—Ecuador.
polygonatoides, *Baker, l. c.*—Ecuador.
praecipua, Herb. Amaryll. 421 = B. Salsilla.
pudibunda, *Planch. & Linden, Cat.* n. 10. (1855) 4.—N. Granat.
pumila, *Grisob. in Lechl. Berb. Am. Austr.* 56, nomen.—Peru.
punctata, *Herb. in Lindl. Bot. Reg.* (1842) *Misc.* 66.—Peruv.
purpurea, *Herb. Amaryll.* 118, 399.—Peruv.
rosea, *Herb. l. c.* 118.—Peruv.
Salsilla, *Mirb. in Hist. Nat. Pl.* ix. (1804) 71.—Chili.
Salsilla, Vell. Fl. Flum. iii. t. 80 = edulis.
salsilloides, M. Roem. Syn. Ensat. 264 = edulis.
secundifolia, *Baker, in Journ. Bot.* xx. (1882) 202.—Peruv.
setacea, Herb. Amaryll. 117 = tomentosa.
Shuttleworthii, *Mast. in Gard. Chron.* (1882) i. 76.—N. Granat.
simplex, Benth. Pl. Hartw. 156 = Hartwegii.
simplex, Herb. Amaryll. t. 15; et in Bot. Mag. t. 3868 = rosea.
spectabilis, Schenk, in Mart. Fl. Bras. iii. t. 169 = edulis.
subfalcata, Herb. Amaryll. 422; Pritz. Ic. Ind. 158 = B. Salsilla.
subglobosa, Herb. in Lindl. Bot. Reg. (1842) Misc. 66 = fimbriata.

BOMAREA:—
superba, *Herb. Amaryll.* 117. t. 6. f. 1.—Peruv.
Tatiana, Herb. l. c. 423 = ovata.
tomentosa, *Herb. l. c.* 117.—Peruv.
torta, *Herb. l. c.* 115.—Ecuador.
Turneriana, Herb. in Lindl. Bot. Reg. (1842) Misc. 66 = formosissima.
uncifolia, *Herb. l. c.*—Peruv. vel Bolivia.
variabilis, Herb. l. c. = rosea.
villosa, M. Roem. Syn. Ensat. 268 = ovata.
vitellina, *Mast. in Gard. Chron.* (1882) i. 148.—N. Granat.
Williamsiae, *Mast. l. c.* (1882) ii. 553.—N. Granat.

BOMARIA, Kunth, Enum. Pl. v. 787 (1850) = Bomarea, Mirb. (Amaryll.).

BOMBAX, Linn. Sp. Pl. ed. I. 511 (1753). *MALVACEAE*, Benth. & Hook. f. i. 210.
ERIOTHECA, Schott & Endl. Meletem. 35 (1832).
SALMALIA, Schott & Endl. l. c. (1832).
aculeatum, *Vell. Fl. Flum.* 289; vii. t. 56.—Bras.
asculifolium, H. B. & K. Nov. Gen. et Sp. v. 298 = Eriodendron asculifolium.
asilitare, Moç. & Sessé, ex DC. Prod. i. 479 = Eriodendron asculifolium.
Barrigon, *Decne. in Fl. des Serres*, Sér. II. xiii. (1860) 44.—Panama.
buonoponense, *Beauv. Fl. Owar.* ii. 42. t. 88.—Afr. trop.
carolinodes, Donn, Cat. Hort. Cant. ed. IV. 156 = Pachira minor.
carolinum, *Vell. Fl. Flum.* 289; vii. t. 57.—Bras.
Criba, Burm. f. Fl. Ind. 145 = malabaricum.
Ceiba, *Linn. Sp. Pl.* 511.—Am. austr.
Congo, Burm. f. Fl. Ind. 145 = Cochlospermum Gossypium.
coriaceum, *Mart. Nov. Gen. et Sp.* i. 93, in obs.—Bras.
cumanense, *H. B. & K. Nov. Gen. et Sp.* v. 300.—Venezuela.
decaphyllum, Decne. in Fl. des Serres, Sér. II. xiii. (1860) 44 = endecaphyllum.
discolor, *H. B. & K. Nov. Gen. et Sp.* v. 299.—N. Granat.
ellipticum, *H. B. & K. l. c.*—Mexic.
emarginatum, *Decne. in Fl. des Serres*, Sér. II. xiii. (1860) 44.—Bras.
endecaphyllum, *Vell. Fl. Flum.* 288; vii. t. 50.—Bras.
Erianthus, Cav. Diss. v. 294 = Eriodendron leiantherum.
Fendleri, *Benth. in Journ. Linn. Soc.* vi (1862) 108.—Panama.
festivum, Wall. Cat. n. 1841 = insigne.
glabrescens, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 211.—Bras.
globiferum, Raeusch. Nom. ed. III. 105 = globosum.
globosum, *Aubl. Pl. Gui.* ii. 703. t. 281.—Guiana.
Gossypium, Linn. Syst. ed. XII. 517 = Cochlospermum Gossypium.
grandiflorum, Cav. Diss. v. 295 = insigne.
grandiflorum, Sonner. Voy. Ind. ii. 235. t. 133 = Cochlospermum Gossypium.
guineense, Schum. & Thonn. Beskr. Guin. Pl. 302 = Eriodendron anfractuosum.
heptaphyllum, Cav. Diss. v. 296; Roxb. Fl. Corom. iii. 43. t. 247 = malabaricum.
heptaphyllum, Linn. Sp. Pl. 960 = septenatum.
hexaphyllum, *Vell. Fl. Flum.* 288; vii. t. 52.—Bras.
hibiscifolium, Willd. ex Steud. Nom. ed. II. i. 815 = Cochlospermum hibiscoides.
Hilarianum, *Decne. in Fl. des Serres*, Sér. II. xiii. (1860) 44.—Bras.
humile, *Benth. in Journ. Linn. Soc.* vi. (1862) 108.—Bras.
insigne, *Wall. Pl. As. Rar.* i. 74. t. 79 et 80.—Ind. or.
jasminiflorum, Beill. Hist. Pl. iv. 119 = Eriodendron jasminodorum.
malabaricum, *DC. Prod.* i. 479.—Ind. or.; Malaya; China.
mediterraneum, *Arruda, in Koster, Trav. Bras.* 489.—Bras.
mexicanum, *Hemsl. Diag. Pl. Nov.* 4.—Mexic.
momponense, *H. B. & K. Nov. Gen. et Sp.* v. 300.—Venezuela.
Munguba, *Mart. Nov. Gen. et Sp.* i. 93.—Bras.
occidentale, Spreng. Syst. iii. 124 = momponense; Eriodendron caribaeum.
octophyllum, *Vell. Fl. Flum.* 288; vii. t. 51.—Bras.
orientale, Spreng. Syst. iii. 124 = Eriodendron anfractuosum.

BOMBAX:—
orinoccense, H. B. & K. Nov. Gen. et Sp. v. 300 = Cochlospermum orinoccense.
parviflorum, *Mart. Nov. Gen. et Sp.* i. 91. t. 57.—Bras.
pentandrum, Jacq. Stirp. Am. 191 = Eriodendron caribaeum.
pentandrum, Linn. Sp. Pl. 511 = Eriodendron anfractuosum.
pentaphyllum, *Vell. Fl. Flum.* 289; vii. t. 55.—Bras.
pleiostemon, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 8 = Eriodendron anfractuosum †
pubescens, *Mart. Nov. Gen. et Sp.* i. 91. t. 58.—Bras.
pyramidale, Cav. Diss. v. 294. t. 153 = Ochroma lagopus.
pyramidatum, Steud. Nom. ed. I. 114 = praec.
quinatum, Jacq. Stirp. Am. 192 = B. Ceiba.
religiosum, Linn. Sp. Pl. 513 = Cochlospermum Gossypium.
retusum, *Mart. Nov. Gen. et Sp.* i. 92. t. 59.—Bras.
rubrinervis, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 69.—Bras.
Samauma, Spreng. Syst. iv. Cur. Post. 259 = Eriodendron Samauma.
septenatum, *Jacq. Enum. Pl. Carib.* 26.—N. Granat.
serrata, *Desf. Cat. Hort. Par.* ed. III. 407.—Hab.?
sexdigitatum, *Vell. Fl. Flum.* 289; vii. t. 54.—Bras.
tomentosum, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* i. 268.—Bras.
ventricosum, Arruda, in Koster, Trav. Bras. 489 = Chorisia crispiflora.
villosum, *Mill. Gard. Dict.* ed. VIII. n. 8.—Am. austr.
vitifolium, Willd. Enum. Hort. Berol. 720 = Cochlospermum vitifolium.

BOMBIX, Medic. Malv. 44 (1787) = **Hibiscus**, Linn (Malvac.).
phoenicea, Medic. l. c. = H. phoeniceus.

BOMBYCELLA, Lindl. Veg. Kingd. 870 (1847) = **Hibiscus**, Linn. (Malvac.).
betulina, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 241 (= H. betulinus?).—Porto Rico.
bicolor, Welw. Apont. 589.—Afr. trop.
phoenicea, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 241 = H. phoeniceus.

BOMBYCIDENDRON, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 14 = **Hibiscus**, Linn. (Malvac.).
grewiaefolium, Zoll. & Mor. l. c. = H. grewiaefolius.

BOMBYCOSPERMUM, Presl, Rel. Haenk. ii. 137. t. 71 (1831) = **Ipomoea**, Linn. (Convolv.).
mexicanum, Presl, l. c. = I. bombycina.

BOMBYNIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1; ed. II. 64 (1827) (Quid?).

BOMBYX, Moench, Meth. 616 (1794) = **Hibiscus**, Linn. (Malvac.).
phoenicea, Moench, l. c. = H. phoeniceus.

BONA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 380; et Phil. Bot. i. 205 (1789) = **Vicia**, Linn. (Legumin.).
narbonensis, Medic. l. c. 380 = V. narbonensis.
speciosa, Medic. l. c. 361 = V. narbonensis.

BONAFIDIA, Neck. Elem. iii. 46 (1790) = **Amorpha**, Linn. (Legumin.).

BONAFOUSIA, A. DC. Prod. viii. 359 (1844) = **Tabernaemontana**, Linn. (Apocyn.).
alternata, Miers, Apocyn. S. Am. 51.—Am. austr.
guyanensis, Miers, l. c. = T. guyanensis.
latiflora, Miers, l. c. 50 = T. flavicans.
oblique, Miers, l. c. 48.—Am. austr.
oblongifolia, Miers, l. c. 50 = T. oblongifolia.
olivacea, Miers, l. c. 52 = T. olivacea.
Perrottetii, Miers, l. c. 51 = T. Perrottetii.
polyneura, Miers, l. c. 53.—Am. austr.
rariflora, Miers, l. c.—Am. austr.
rupicola, Miers, l. c. 52 = T. rupicola.
undulata, A. DC. Prod. viii. 359 = T. undulata.

BONAGA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 345 ; Phil. Bot. 206 (1789) = **Ononis**, Linn (Leguminos.).
antiquorum, Medic. l. c. 346 = O. antiquorum.
arvensis, Medic. Phil. Bot. 206 = O. arvensis.
Columnae, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 346 = O. Columnae.
hircina, Medic. l. c = O. hircina.
minutissima, Medic. l. c. = O. minutissima.
mitis, Medic. l. c. = O. arvensis.
spinosa, Medic. l. c. = O. spinosa.

BONAMIA, A. Gray, in Proc. Am. Acad. v. (1862) 337 = **Breweria**, K. Br. (Convolv.).
aquatica, A. Gray, Man. Bot. N. U. St. ed. V. 376 = Breweria aquatica.
humistrata, A. Gray, in Proc. Am. Acad. v. (1862) 337 = Breweria humistrata.
Mensiesii, A. Gray, l. c.—Ins. Sandvic.
Pickeringii, A. Gray, Man. Bot. N. U. St. ed. V. 376 = Breweria Pickeringii.

BONAMIA, Thou. in Dict. Sc. Nat. v. 145 (1804) ; et Hist. Veg. Isl. Afr. 17. t. 8 (1806). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 877.
madagascariensis, *Poir. Encyc. Suppl.* i. 677.—Madag.

BONAMICA, Vell. Fl. Flum. 21 (1825) ; t. t. 50 = **Linociera**, Sw. (Oleac.).
filiformis, Vell. l. c.—Bras.

BONANIA, A. Rich. Fl. Cub. Fanerog. ii. 201. t. 68 (1853). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 335.
adenodon, *Benth. & Hook. f. Gen.* iii. 335.—Cuba.
cubana, *A. Rich. Fl. Cub. Fanerog.* ii. 201.—Cuba.
cubensis, A. Rich. l. c. t. 68 = cubana.
emarginata, *Wright, ex Griseb. in Goett. Nachr.* (1866) 78.—Cuba.
erythrosperma, *Benth. & Hook. f. Gen.* iii. 335.—Cuba.
myricifolia, *Benth. & Hook. f. l. c.*—Cuba.

BONANNIA, Gaza. Fl. Sic. Syn. i. 355 (1842). *UMBELLIFERAE*, Benth. & Hook. f. i. 910.
retziifera, *Guss. l. c.*—Sicilia.
resinosa, Strobl, in Oestr. Bot. Zeitschr. xxxiv. (1884) 174 = resinifera.

BONANNIA, Presl, Fl. Sic. i. 99 (1826) = **Brassica**, Linn. (Crucif.).
dissecta, Presl, l. c. 100 = Brassica dissecta.
officinalis, Presl, l. c. 99 = Brassica alba.

BONANNIA, Rafin. Specchio, iv. 115 (1814) = **Blighia**, Kon. (Sapind.).
nitida, Rafin. l. c. 116 = Blighia sapida.

BONANOX, Rafin. Fl. Tellur. 77 (1836) = **Ipomoea**, Linn. (Convolv.).
indica, Rafin. l. c.—Ind. or.
muricata, Rafin. l. c. = I. Bona-nox.
orbiculata, Rafin, l. c. = I. orbiculata.
riparia, Rafin. l. c. = I. riparia.

BONAPARTEA, Haw. Syn. Pl. Succ. 68 (1812) = **Agave**, Linn. (Amaryll.).
filamentosa, Boucen. in Rev. Hortic. (1868) 65 = A. filifera.
flagelliformis, Henck. in Flora, iii. (1820) 45 = A. geminiflora.
gracilis, Sweet, Hort. Brit. ed. II. 397 = Dasylirion acrotichum.
Hystrix, Hort. ex Pasq. Cat. Orto Bot. Nap. 111 = A. striata.
juncea, Haw. Syn. Pl. Succ. 68 = A. geminiflora.
stricta, Hort. ex Vukot. in Rad. Jugos. Akad. Zagrab, xxvii. (1874) 213 = A. striata.

BONAPARTEA, Ruiz & Pav. Fl. Per. iii. 86 (1802) = **Tillandsia**, Linn. (Bromel.).
juncea, Ruiz & Pav. l. c. = T. setacea.
strobilanthe, Ruiz & Pav. l. c. 89 = T. strobilanthe.
vittata, Mant. ex Schult. f. Syst. vii. 1198 = T. vittata.

BONAROTA, Adans. Fam. ii. 209 (1763) = **Pasderota**, Linn. (Scrophul.).
chamaedrifolia, Scop. Fl. Carn. ed. II. i. 92 = P. Ageria.
rotundifolia, Caruel, in Parl. Fl. Ital. vi. 582 = P. Bonarota.

BONATEA, Willd. Sp. Pl. iv. 48 (1805) = **Habenaria**, Willd. (Orchid.).
Bilabrella, Lindl. Gen. et Sp. Orch. 328 = H. tetrapetala.
Boltoni, Harv. Thes. Cap. i. 55. t. 88 = H. Boltoni.
cassidea, Sond. in Linnaea, xix. (1847) 81 = H. cassidea.
cirrhata, Lindl. Gen. et Sp. Orch. 327 = H. cirrhata.
clavata, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 208 = H. clavata.
Darwinii, Weale, in Journ. Linn. Soc. x. (1869) 470 = H. camidea.
densiflora, Sond. in Linnaea, xix. (1847) 80 = H. densiflora.
flexuosa, Lindl. Gen. et Sp. Orch. 328 = H. flexuosa.
foliosa, Lindl. l. c. 329 = H. foliosa.
gracilis, Lindl. l. c. = H. gracilis.
herbacea, Wall. ex Lindl. l. c. 307 = H. digitata.
incarnata, Lindl. l. c. 327 = Disa incarnata.
macilenta, Lindl. ex Benth. in Hook. Lond. Journ. Bot. ii. (1843) 673 = H. macilenta.
micrantha, Lindl. Gen. et Sp. Orch. 329 = H. arenaria.
panduana, Lindl. in Wall. Cat. n. 7063 = H. digitata.
pauciflora, Lindl. Gen. et Sp. Orch. 329 = H. pauciflora.
pratensis, Lindl. l. c. 328 = H. pratensis.
speciosa, Willd. Sp. Pl. iv. 48 = H. Bonatea.
tetrapetala, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 208 = H. tetrapetala.

BONAVERIA, Scop Introd. 310 (1777) = **Securigera**, DC. (1806) (Legum.).
Securidaca, Desv. Journ. Bot. i. (1813) 120 = S Coronilla.
Securigera, Endl. ex Heynh. Nom. ii. 78 = S. Coronilla.

BONDUC, Adans. Fam. ii. 318 (1763) = **Caesalpinia**, Linn. (Legum.).
canaderae, Medic. Theod. 118 = C. Bonduc.
majus, Medic. l. c. 48 = C. Bonduc.
minus, Medic. l. c. 41 = C. Bonducella.

BONELLIA, Bert. ex Colla, Hort. Ripul. 21 (1826) = **Jacquinia**, Jacq. (Myrsin.).
Cavanillesii, Bert. l. c. = J. macrocarpa.

BONGARDIA, C. A. Mey. Verz. Pfl. Cauc. 174 (1831). *BERBERIDEAE*, Benth. & Hook. f. i. 48.
chrysogonum, Boiss. Fl. Orient. i. 99 = Rauwolfii.
Olivieri, C. A. Mey. Verz. Pfl. Cauc. 175 = Rauwolfii. Rauwolfii, *C. A. Mey. l. c.* 174.—Graecia ; Orient ; Afghan.

BONIFACIA, Manso, ex Steud. Nom. ed. II. i. 216 (1840) = **Augusta**, Pohl (Rubiac.).
riparia, Manso, ex Steud. l. c. = A. lanceolata.

BONJEANEA, Reichb. Fl. Germ. Excurs. 507 (1832) = **Dorycnium**, Linn. (Legum.).
cinerascens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 12 = D. hirsutum.
graeca, Griseb. Spicil. Fl. Rumel. i. 43 = D. latifolium.
hirsuta, Reichb. Ic. Fl. Germ. x. 32 = D. hirsutum.
hirta, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 11 = D. hirsutum.
italica, Jord. & Fourr. l. c. 12 = D. hirsutum.
Kotschyi, Boiss. & Kent. ex Boiss. Fl. Orient. ii. 161 = D. Kotschyi.
microphylla, Reichb. Fl. Germ. Excurs. 507, in nota = Indigofera gracilis.
prostrata, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 11 (emend. fasc. ii. 138) = D. hirsutum.
recta, Reichb. Fl. Germ. Excurs. 507 = D. rectum.
sericea, Jord. & Fourr. Ic. Fl. Eur. i. 24. t. 61 = D. hirsutum.
syriaca, Boiss. Diagn. Ser. I. ix. 31 = D. hirsutum.
ornata, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 12 = D. hirsutum.

BONNAYA, Link & Otto, Ic. Pl. Sel. 25. t. 11 (1820). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 956.
alata, *Sprengl. Syst.* i. 41.—Ins. Moluce.

BONNAYA:—
alterniflora, *Wright, in Sauv. Fl. Cub.* 101.—Cuba.
bifolia, Sprengl. Syst. iv. Cur. Post. 15 = Chirita bifolia.
brachiata, *Link & Otto, Ic. Pl. Sel.* ii. 25. t. 11.—Ind. or. ; Malaya.
brachycarpa, Cham. & Schlecht. in Linnaea, ii. (1827) 568 = Ilysanthes capensis.
bracteata, Griff. Notul. iv. 107 = veronicaefolia.
ciliata, Sprengl. Syst. i. 41 = reptans.
clausa, *F. Muell. ex Benth. Fl. Austral.* iv. 490.—Austral.
cordifolia, Sprengl. Syst. i. 42 = Vandellia pedunculata.
cyanea, Griff. Notul. iv. 107 = veronicaefolia.
grandiflora, Sprengl. Syst. i. 41 = veronicaefolia.
hispidula, G. Don, Gen. Syst. iv. 538 (Gen. dubium).—Ind. or.
humilis, Sprengl. Syst. iv. Cur. Post. 15 = Chirita pumila.
hyssopoides, Benth. in Wall. Cat. n. 3866 = Ilysanthes hyssopioides.
integrifolia, Koxtel. Allg. Med.-Pharm. Fl. iii. 893 = Vandellia erecta.
marginata, Sprengl. Syst. i. 41 = veronicaefolia.
minima, G. Don, Gen. Syst. iv. 538 = oppositifolia.
minima, Wight, Ic. iii. t. 888 = Ilysanthes minima.
oxymoloides, G. Don, Gen. Syst. iv. 538 (= Vandelliae sp.).—Ind. or.
oppositifolia, *Sprengl. Syst.* i. 41.—Ind. or.
originifolia, Spreng. l. c. 42 = Vandellia diffusa.
parviflora, Benth. in Wall. Cat. n. 3867 = Ilysanthes parviflora.
peduncularis, *Benth. in Wall. l. c.* n. 3865 ; *et Scroph. Ind.* 34.—Ind. or.
penicillata, *Nees, in DC. Prod.* xi. 161.—Japon.
personata, Hassk. in Flora, xlvii. (1864) 56 = brachiata.
procumbens, Benth. in Wall. Cat. n. 3859 = veronicaefolia.
pulegiifolia, Sprengl. Syst. i. 41 = veronicaefolia.
pumila, Sprengl. Syst. iv. Cur. Post. 15 = Chirita pumila.
punctata, G. Don, Gen. Syst. iv. 538 = Limnophila conferta.
pusilla, Benth. in Wall. Cat. n. 3861 ; Scroph. Ind. 33 = brachiata.
pusilla, Griff. Notul. iv. 107 = tenuifolia.
pusilla, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 121. —Afr. trop.
repens, G. Don, Gen. Syst. iv. 537 = seq.
reptans, *Sprengl. Syst.* i. 41.—Ind. or. ; Malaya ; Ins. Maurit.
rigida, Benth. in Wall. Cat. n. 3860 = veronicaefolia.
Rothii. A. Dietr. Sp. Pl. i. 556 = veronicaefolia.
rotundifolia, Benth. Scroph. Ind. 34 = Ilysanthes rotundifolia.
Roxburghiana, A. Dietr. Sp. Pl. i. 559 = veronicaefolia.
rueliioides, Spreng. Syst. i. 41 = reptans.
serrata, A. Dietr. Sp. Pl. i. 556 ; Griff. Notul. iv. 106 = brachiata.
tenuifolia, *Sprengl. Syst.* i. 42.—Ind. or. ; China.
trichotoma, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 121.—Afr. trop.
Vahlii, G. Don, Gen. Syst. iv. 538 = Vandellia scabra.
verbenaefolia, Spreng. Syst. i. 42 = veronicaefolia.
veronicaefolia, *Sprengl. l. c.* 41.—Ind. or. ; Malaya ; China.
Wightii, Benth. in Wall. Cat. n. 3864 = veronicaefolia.

BONNETIA, Mart. & Zucc. Nov. Gen. et Sp. i. 114. t. 100 (1824). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 187.
KIESKIRIA, Nees, in Neuwied, Reise nach Brasil. ii. 338 (1821).
anceps, *Mart. & Zucc. Nov. Gen. et Sp.* i. 115.—Bras.
bahiensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 246.—Bras.
coriacea, Sprengl. syst. iv. Cur. Post. 207 = Kielmeyera coriacea.
corymbosa, Spreng. l. c. = Kielmeyera corymbosa.
meridionalis, *Sw. Fl. Ind. Occ.* ii. 997.—Am. trop.
palustris, Vahl, Eclog. Am. ii. 42 = meridionalis.
paniculata, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 83.—Peruv.
parviflora, *Spruce, ex Benth. l. c.*—Peruv.
petiolaris, Spreng. Syst. iv. Cur. Post. 207 = Kielmeyera petiolaris.
rosea, Spreng. l. c. = Kielmeyera rosea.

BONNETIA:—
sessilis, *Benth. in Hook. Lond. Journ. Bot.* ii.(1843) 368.—Guiana.
stricta, *Mart. in Nov. Act. Nat. Cur.* xii. (1825) 87.—Bras.
trifoliata, Auct. ex Walp. Rep. i. 499 = Galipea officinalis.
variabilis, Spreng. Syst. iv. Cur. Post. 207 = Kielmeyera variabilis.
venulosa, *Mart. & Zucc. Nov. Gen. et Sp.* i. 115.—Bras.

BONNETIA, Neck. Elem. i. 868 (1790) = **Buchnera**, Linn. (Scrophular.).

BONNETIA, Schreb. Gen. 368 (1789) = **Mahurea**, Aubl. (Ternstroem.).
palustris, J. F. Gmel. Syst. 814 = M. palustris.
racemosa, Sw. Fl. Ind. Occ. ii. 966 = *M. speciosa*.

BONPLANDIA, Cav. in Anal. Hist. Nat. ii. (1800) 131. t. 20. *POLEMONIACEAE*, Benth. & Hook. f. ii. 924.
CALDASIA, Willd. Hort. Berol. t. 71 (1807).
geminiflora, *Cav. l. c.* 132.—Mexic.

BONPLANDIA, Willd. in Mém. Acad. Berl. (1802) 24 = **Galipea**, Aubl. (Rutac.).
Angustura, Rich. in Mém. Math.-Phys. Inst. Paz. (1811) 63. t. 10 = G. Cusparia.
Candolleana, Spreng. Syst. i. 781 = G. Candolleana.
cuneifolia, Spreng. l. c. = G. cuneifolia.
Erythrochiton, Spreng. l. c. 780 = G. pentandra.
Fontanesiana, Spreng. l. c. = G. Fontanesiana.
heterophylla, Spreng. l. c. = G. heterophylla.
Laxiostemon, Spreng. l. c. = G. Laxiostemon.
pentagyna, Spreng. l. c. = G. pentagyna.
resinosa, Spreng. l. c. = G. resinosa.
trifoliata, Willd. in Mém. Acad. Berl. (1802) 25 = G. Cusparia.

BONTIA, Linn. in Loefl. Iter, 193 (1758) = **Aviosmina**, Linn. (Verbenac.).
germinans, Linn. Sp. Pl. ed. II. 891 = A. officinalis.

BONTIA, Linn. Syst. ed. I (1735). *MYOPORINEAE*, Benth. & Hook. f. ii. 1126.
daphnifolia, Sallsb. Prod. 109 = seq.
daphnoides, *Linn. Sp. Pl.* 885.—Ind. occ.
minor, Gaertn. f. Fruct. iii. 169. t. 213. f. 3 = daphnoides.

BONTUNIA, Schomb. Reise, iii. 1082, nomen; Prog. in Mart. Fl. Bras. vi. i. 267. t. 72 (1868). *LOGANIACEAE*, Benth. & Hook. f. ii. 796.
antoniaefolia, *Prog. in Mart. Fl. Bras.* vi. i. 268.—Bras.
superba, *Schomb. Reise*, iii. 1082.—Guiana.

BOOPHONE, Herb. App. 18 (1821) = **Buphane**, Herb. (Amaryll.).

BOOPIS, Juss. in Ann. Mus. Par. ii. (1803) 350. t. 58. *CALYCERAE*, Benth. & Hook. f. ii. 561.
ACARPHA, Griseb. in Goett. Abh. vi. (1858–59) 125.
CHIONOPHILA, Miers, ex Lindl. Veg. Kingd. 701 (1847).
GAMOCARPHA, DC. Prod. v. 2 (1836).
NASTANTHUS, Miers, ex Lindl. Veg. Kingd. 701 (1847), nomen; et in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 185.
annalis, *Phil. in Anal. Univ. Chil.* (1861) 40.—Chili.
alpina, *Poepp. ex Less. in Linnaea*, vi. (1831) 358.—Chili.
anthemoides, *Juss. in Ann. Mus. Par.* ii. (1803) 350.—Reg. Argent.
araucana, *Phil. in Anal. Univ. Chil.* (1872) 786.—Chili.
australis, *Decne. Voy. Pôle Sud Bot.* ii. 87. t. 18, B.—Reg. Magell.
balsamitaefolia, Juss. in Ann. Mus. Par. ii. (1803) 350 = Calycera balsamitaefolia.
bellidifolia, *Phil. in Anal. Univ. Chil.* (1872) 786.—Chili.
boplearoides, *C. A. Muell. in Mart. Fl. Bras.* vi. iv. 255.—Bras.
caespitosa, *Phil. Fl. Atac.* 26.—Chili.
compacta, *Phil. in Linnaea*, xxviii. (1856) 709.—Chili.
crassifolia, *A. Gray, in Proc. Am. Acad.* v. (1861) 321.—Patagon.
Diezi, *Phil. in Linnaea*, xxxiii. (1864–65) 106.—Chili.
Gayana, *Phil. l. c.* xxviii. (1856) 709.—Chili.
gracilis, Phil. l. c. 707 = leucanthema.
graminea, *Phil. l. c.*—Chili.
laciniata, *Ball, in Journ. Linn. Soc.* xxi. (1884) 220.—Patagon.
leucanthema, *Poepp. ex Less. in Linnaea*, vi. (1831) 359.—Chili.
multicaulis, *Phil. in Linnaea*, xxviii. (1856) 706.—Chili.
poaeoeformis, *Phil. in Anal. Univ. Chil.* (1872) 786.—Chili.
pusilla, *Phil. in Linnaea*, xxxiii. (1864–65) 105.—Chili.
rigidula, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 284.—Patag.
sanjuanina, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 31.—Reg. Argent.
scapigera, *Remy, in C. Gay, Fl. Chil.* iii. 250.—Chili.
spathulata, *Phil. in Linnaea*, xxviii. (1856) 708.—Chili.
squarrosa, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 286 = australis.
tridactyloides, Spreng. Syst. iii. 674 = Acicarpha tribuloides.

BOOTHIA, Dougl. ex Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 405 = **Platystemon**, Benth. (Papaveracˇ.).
californica, Dougl. ex Lood. Encyc. Pl. Suppl. i. 1214 = P. californicus.

BOOTIA, Adans. Fam. ii. 320 (1763) = **Borbonia**, Linn. (Legumin.).

BOOTIA, Bigel. Fl. Boston. ed. II. 206 (1824) = **Potentilla**, Linn. (Rosac.).
sylvestris, Bigel. l. c. = P. arguta.

BOOTIA, Neck. Delic. Gallo-Belg. i. 193 (1768) = **Saponaria**, Linn. (Caryophyll.).
glutinosa, Neck. ex Reichb. Ic. Fl. Germ. v. 41 = S. glutinosa.
ocymoides, Neck. ex Reichb. l. c. = S. ocymoides.
Saponaria, Neck. in Act. Acad. Theod. Pal. ii. (1770) 484 = S. officinalis.
vulgaris, Neck. Delic. Gallo-Belg. i. 193 = S. officinalis.

BOOTROPHIS, Stend. Nom. ed. II. i. 22 (1840) = BOTROPHIS, Rafin. = Climatrugna, Linn. (Rosmac.).

BOOTTIA, Ayres, ex Baker, Fl. Maurit. 49 (1877) = **Pleurostylia**, Wight & Arn. (Celastrac.).
obovata, Ayres, ex Baker, l. c. = P. leucocarpa.

BOOTTIA, Wall. Pl. As. Rar. i. 51. t. 65 (1829). *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 453.
cordata, *Wall. l. c.*—Burma.

BOPUSIA, Presl, Bot. Bemerk. 91 (1844) = **Gradaria**, Benth. (1845) (Scrophul.).
scabra, Presl, l. c. = G. scabra.

BOQUILA, Decne. in Comptes Rend. (1837) 394; et in Arch. Mus. Par. i. (1839) 188. t. 11 *H. BERBERIDEAE*, Benth. & Hook. f. i. 45.
discolor, *Decne. in Ann. Sc. Nat.* Sér. II. xii. (1839) 108.—Chili.
trifoliata, *Decne. l. c.*—Chili.

BORABORA, Stend. Syn. Pl. Cyp. 71 (1855) = **Marisous**, Gaertn. (Cyperac.).
cyperoides, Stend. l. c. = M. macrophyllus.

BORAEVA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 383 = **Boraeva**, Jaub. & Spach (Crucif.).

BORAGO, Linn. Sp. Pl. i. 137 (1753). *BORAGINEAE*, Benth. & Hook. f. ii. 854.
BORRAGO, Tourn. ex Linn. Syst. ed. I (1735).
BUGLOSSITES, Moris, in Enum. Sem. Hort. Taur. (1845).—Cf. Ann. Sc. Nat. Sér. III. v. (1846) 365.
adscens, Gilib. Fl. Lithuan. Suppl. 272 = officinalis.
africana, Linn. Sp. Pl. 138 = Trichodesma africanum.
arabica, Ehrenb. ex Boiss. Fl. Orient. iv. 281 = Trichodesma Ehrenbergii.
aspera, Gilib. Fl. Lithuan. i. 25 = officinalis.
constantinopolitana, *Hill. Veg. Syst.* vii. 45.—Oriens.
cordifolia, Moench, Meth. Suppl. 275 = orientalis.
crassifolia, Vent. Jard. Cels, 100. t. 100 = Caccinia glauca.
cretica, *Willd. Sp. Pl.* i. 778.—Graecia.
gruina, *C. Sm. in Tuckey, Congo*, 250, nomen.—Ins. Cap. Virid.
indica, Blanco, Fl. Filip. ed. I. 81 = Trichodesma zeylanicum.
indica, Linn. Sp. Pl. 137 = Trichodesma indicum.
inflata, Salisb. Prod. 116 = orientalis.
lasiflora, *Willd. Enum. Hort. Berol. Suppl.* 9.—Corsica.
longifolia, *Poir. Voy. Barb.* ii. 119.—Afr. bor.
macrantha, (Soland. in) Russell, Aleppo, ed. II. ii. 246 = Caccinia Russellii.
officinalis, *Linn. Sp. Pl.* 137.—Europ.; Afr. bor.; As. Min.
orientalis, *Linn. l. c.* 138.—As. Min.; Reg. Caucas.
spinulosa, Roxb. Hort. Beng. [13]; Fl. Ind. i. 459 = Trichodesma indicum.
verrucosa, Forsk. Fl. Aegypt. Arab. 41 = Trichodesma africanum.
zeylanica, Burm. f. Fl. Ind. 41 = Trichodesma zeylanicum.

BORASSUS, Linn. Mus. Cliff. 18 (1736). *PALMAE*, Benth. & Hook. f. iii. 939.
LONTARUS, Adans. Fam. ii. 25 (1763).
acthiopicum, Mart. Hist. Nat. Palm. iii. 231 = flabellifer
caudatum, Lour. Fl. Cochinch. 760 = Wallichia caudata duchotomus, *White, in Y. Grah. Cat. Bomb. Pl.* (1839) 226.—Ind. or.
flabellifer, *Linn. Sp. Pl.* 1187.—Ind. or.
flabelliformis, Mart. Syst. ed. XIII. 937 = flabellifer.
Gomutus, Lour. Fl. Cochinch. ii. 618 = Arenga saccharifera.
Ihur, Gaertn. Linn. Prael. 87 = Phoenicoraphis Ihur.
madagascariensis, *Baj. Hort. Maurit.* 308.—Madag.
pinnatifrons, Jacq. Hort. Schoenb. ii. 85 = Chamaedorea gracilis.
secundiflorus, *Hort. ex H. Wendl. in Kerch. Palm.* 285, nomen.—Hab.?
Sonnerati, Gaertn. Linn. Prael. 86 = Lodoicea seschellarum.
tunicatus, Lour. Fl. Cochinch. 618 = Pholidocarpus tunicatus.

BORBONIA, Linn. Coroll. Gen. 12. n. 971 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 475.
BOOTIA, Adans. Fam. ii. 320 (1763).
alata, Willd. Hort. Berol. i. 45 = parviflora.
alpestris, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 461.—Afr. austr.
angustifolia, *Lam. Encyc.* i. 436.—Afr. austr.
axillaris, Lam. l. c. 438 = Amphithalea dense.
barbata, *Lam. l. c.* 436.—Afr. austr.
Candolleana, *Echl. & Zeyh. Enum.* 163.—Afr. austr
capitata, Poir. Encyc. Suppl. i. 680 = Priestleya capitata.
ciliata, DC. Prod. ii. 190 = Candolleana.
ciliata, *Willd. Sp. Pl.* iii. 923.—Afr. austr.
complicata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 462.—Afr. austr.
cordata, Andr. Bot. Rep. t. 81 = Rafnia triflora.
cordata, *Linn. Sp. Pl.* 707.—Afr. austr.
cordifolia, Lam. Encyc. i. 436 = cordata.
crenata, *Linn. Sp. Pl.* 708.—Afr. austr.
cronata, Willd. ex Stend. Nom. ed. II. i. 217 = parviflora.
dasiphylla, E. Mey. Comm. Pl. Afr. Austr. i. 15 = lanceolata.
ericifolia, Linn. Pl. Rar. Afr. 14; Amoen. Acad. vi. 92 = Amphithalea ericaefolia.
graminifolia, Lam. Encyc. i. 437 = Priestleya graminifolia.
hirsuta, *Poir. Encyc. Suppl.* i. 680 = Priestleya hirsuta.
laevigata, Linn. Mant. 100 = Priestleya Thunbergii.

BORBONIA :—
lanceolata. *Linn. Sp. Pl.* 707.—Afr. austr.
latifolia, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 462.—Afr. austr.
monosperma, *DC. Prod.* ii. 130.—Afr. austr.
myrtifolia, Poir. Encyc. Suppl. i. 680 = Priestleya myrtifolia.
obcordata, Spreng. Syst. iv. Cur. Post. 267 = Tephrosia obcordata.
ovata, Andr. Bot. Rep. t. 81 = Rafnia trifora.
parviflora, *Lam. Encyc.* i. 497.—Afr. austr.
perfoliata, Lam. i. c. = ciliata.
perfoliata, Thunb. Prod. Pl. Cap. 122 = Rafnia perfoliata.
perforata, *Eckl. & Zeyh. Enum.* 168.—Afr. austr.
perforata, Thunb. Prod. Pl. Cap. 122 = ciliata.
pungens, Mundt, ex Benth. in Hook. Lond. Journ. Bot. ii. (1843) 461 = trinervia.
ruscifolia, Sims, Bot. Mag. t. 2128 = parviflora.
sericea, Lam. Encyc. i. 498 = Priestleya sericea.
serrulata, Thunb. Diss. Borbonia, 6 = parviflora.
sphaerica, Lam. Encyc. i. 497 = Liparia sphaerica.
sphaerosperma, Spreng. Syst. iv. Cur. Post. 267 = Requienia sphaerosperma.
tecta, Poir. Encyc. Suppl. i. 680 = Priestleya tecta.
teres, Poir. l. c. = Priestleya teres.
tomentosa, Linn. Sp. Pl. 707 = Priestleya villosa.
trifiora, Auct. ex Steud. Nom. ed. II. i. 217 = Rafnia trifiora.
trinervia, E. Mey. Comm. Pl. Afr. Austr. i. 15 = lanceolata.
trinervia, Linn. Sp. Pl. 707 = Cliffortia ruscifolia.
trinervia, *Thunb. Prod. Pl. Cap.* 122.—Afr. austr.
umbellifera, *Poir. Encyc. Suppl.* i. 680.—*Afr. austr.*
undulata, *Thunb. Prod. Pl. Cap.* 122.—Afr. austr.
vestita, Poir. Encyc. Suppl. i. 680 = Priestleya vestita.
villosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 38.—Afr. austr.
villosa, Thunb. Fl. Cap. 560 = Priestleya angustifolia.

BORBONIA, Plum. ex Adans. Fam. ii. 341 (1763) (Laurin.).
carolinensis, J. S. Presl, Rostl. ii. 57 = Persea caroliniensis.
cupularis, Gaertn. f. Fruct. iii. 225. t. 222 (= *Mespilodaphne cupularis*).—Maurit.
exaltata, J. S. Presl, Rostl. ii. 57 = Nectandra exaltata.
fortens, J. S. Presl, l. c. = Ocotea foetens.
globosa, Gaertn. f. Fruct. iii. 224. t. 222 = Nectandra sanguinea.
indica, J. S. Presl, Rostl. ii. 57 = Persea indica.
martinicensis, J. S. Presl, l. c. (= *Oreodaphne Jacquiniana*).—Ind. occ.
membranacea, J. S. Presl, l. c. = Nectandra membranacea.
peruviana, Juss. ex Steud. Nom. ed. II. i. 217 (= *Mespilodaphne pretiosa*).—Peruv.

BORBOYA, Rafin. Fl. Tellur. iii. 52 (1836) = **Hyacinthus**, Tourn. (Liliac.).
spicata, Rafin. l. c. = H. spicatus.

BORCKHAUSENIA, Gaertn. Mey. & Scherb. Fl. Wett. iii. 4 (1801) = **Corydalis**, Vent. (1808) (Papav.).
capnoides, Fuss, Fl. Transs. 43 = C. capnoides.
cava, Gaertn. Mey. & Scherb. Fl. Wett. iii. 17 = C. bulbosa.
lutea, Gaertn. Mey. & Scherb. l. c. 19 = C. lutea.
ochroleuca, Fuss, Fl. Transs. 43 = C. ochroleuca.
solida, Gaertn. Mey. & Scherb. Fl. Wett. iii. 16 = C. Halleri.

BORCKHAUSENIA, Roth, Catalecta, fasc. ii. 56 (1800) = **Teedia**, Rudolphi (Scrophul.).
lucida, Roth, l. c. = T. lucida.

BORDEREA, Mirbel. in Bull. Soc. Bot. Fr. xiii. (1866) 374 = **Dioscorea**, Linn.
pyrenaica, Mirbel. l. c. = D. pyrenaica.

BOREA, Zipp. ex Mackl. in Bijdr. Nat. Wet. v. (1836) 148, nomen (absque descr.).

BOREA, Meisn. Gen. Comm. 321 (1836-42), sphalm. = ROVEA, Decne. = **Lindenbergia**, Lehm. (Scrophul.).

BOREAVA, Jaub. & Spach, Illustr. i. 3. t. 2 (1841). *CRUCIFERAE*, Benth. & Hook. f. i. 95.
MARTINSIA, Godr. in Mém. Acad. Stanislas, Sér. III. [xvii.] 1853 (1854) 378.
aptera, *Boiss. & Heldr. Diagn.* Ser. I. viii. 49.—As. Min.
orientalis, *Jaub. & Spach, Illustr.* i. 3.—As. Min.; Syria.

BORELLIA, Neck. Elem. i. 275, 276 (1790) = **Cordia**, Linn. (Boragin.).
aspera, Rafin. Sylva Tellur. 41 = C. aspera.

BORETTA, Neck. Elem. i. 212 (1790) = **Daboëcia**, D. Don (Ericac.).

BORISSA, Rafin. ex Steud. Nom. ed. II. i. 217 (1840) = **Lysimachia**, Linn. (Primulac.).
Linum-stellatum, Rafin. ex Steud. l. c. = L. Linum-stellatum.

BORITH, Adans. Fam. ii. 262 (1763) = **Anabasis**, Linn. (Chenopod.).

BORKHAUSIA, Nutt. Gen. Am. ii. 125 (1818) = BARKHAUSIA, Moench = **Crepis**, Linn. (Compos.).

BORONELLA, Baill. Adansonia, x. (1872) 302. *RUTACEAE*.
Pancheri, *Baill. l. c.* 303.—N. Caled.

BORONIA, Sm. Tracts, 288 (1798); et in Trans. Linn. Soc. viii. (1807) 285. t. 8-7. *RUTACEAE*, Benth. & Hook. f. i. 291.
CYANOTHAMNUS, Lindl. Swan River Bot. 18 (1839).
affinis, *R. Br. ex Benth. Fl. Austral.* i. 311.—Austral.
alata, *Sm. in Trans. Linn. Soc.* viii. (1807) 283 — Austral.
alhiflora, *R. Br. ex Benth. Fl. Austral.* i. 317.—Austral.
algida, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 100.—Austral.
alulata, *Soland. ex Benth. Fl. Austral.* i. 313.—Austral.
anemonifolia, *A. Cunn. in Field, New South Wales*, 326.—Austral.
anemonifolia, Paxt. Mag. Bot. ix. 128 = Fraseri.
anethifolia, A. Cunn. ex Endl. in Enum. Pl. Hueg. 16 = anemonifolia.
arenicola, F. Muell. Fragm. i. 100 = Zieria Smithii.
artemisiaefolia, *F. Muell. l. c.* 66.—Austral.
artemisioides, F. Muell. in Hook. Kew Journ. ix. (1857) 196 = prace.
baeckeaceae, *F. Muell. Fragm.* iv. 28.—Austral.
Barkeriana, *F. Muell. l. c.* xi. 96.—Austral.
bicolor, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 166 = tetrandra.
bipinnata, Lindl. in Mitch. Journ. Trop. Austr. 225 = anemonifolia.
Bowmani, *F. Muell. Fragm.* iv. 135.—Austral.
brachyphylla, F. Muell. l. c. i. 99 = oxyantha.
Bussellians, *F. Muell. l. c.* ix. 113.—Austral.
caerulea, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 11.—Austral.
calophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 160.—Austral.
Candollei, G. Don, Gen. Syst. i. 793 = alata.
capitata, *Benth. Fl. Austral.* i. 323.—Austral.
chironiifolia, Bartl. in Lehm. Pl. Preiss. i. 167 = denticulata.
citriodora, Gunn, ex Hook. f. Fl. Tasm. i. 68 = pinnata.
clavellifolia, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 12 = inornata.
colorata, Lehm. Pl. Preiss. ii. 226 = viminea.
crassifolia, *Bartl. in Lehm. l. c.* i. 169.—Austral.
crassiuscula, *Bartl. l. c.* 168.—Austral.
crenulata, *Sm. in Trans. Linn. Soc.* viii. (1807) 284.—Austral.
cymosa, *Endl. in Enum. Pl. Hueg.* 16.—Austral.
deflexa, *F. Muell. Fragm.* ix. 113.—Austral.
denticulata, *Sm. in Trans. Linn. Soc.* viii. (1807) 284. —Austral.
dentigera, F. Muell. in Trans. Inst. Vict. i. (1855) 82 = anemonifolia.
dichotoma, Lindl. Bot. Reg. (1841) sub t. 47 = spathulata.

BORONIA :—
Drummondii, Planch. in Fl. des Serres, Sér. I. ix. (1853-54) 69. t. 881 = pulchella.
Edwardsii, *Benth. Fl. Austral.* i. 312.—Austral.
elatior, *Bartl. in Lehm. Pl. Preiss.* i. 170.—Austral.
erianthe, *Lindl. in Mitch. Journ. Trop. Austr.* 226.—Austral.
ericifolia, *Benth. Fl. Austral.* i. 313.—Austral.
ericifolia, *A. Cunn. in Lindl. Bot. Reg.* (1841) sub t. 47.—Austral.
fasciculifolia, F. Muell. Fragm. i. 99 = thymifolia.
fastigiata, *Bartl. in Lehm. Pl. Preiss.* i. 168.—Austral.
filicifolia, *A. Cunn. ex Benth. Fl. Austral.* i. 311.—Austral.
filifolia, *F. Muell. Fragm.* i. 3.—Austral.
flexuosa, Bartl. in Lehm. Pl. Preiss. i. 166 = spathulata.
floribunda, Siebert, ex Spreng. Syst. iv. Cur. Post. 148 = pinnata.
Fraseri, *Hook. Bot. Mag.* t. 4052.—Austral.
gracilipes, *F. Muell. Fragm.* ii. 99.—Austral.
grandisepala, *F. Muell. l. c.* i. 66.—Austral.
granulata, F. Muell. l. c. 101 = Zieria granulata.
Gunnii, Hook. f. Fl. Tasm. i. 68. t. 10 = pinnata.
heterophylla, *F. Muell. Fragm.* ii. 98.—Austral.
hirsuta, F. Muell. l. c. i. 101 = Zieria pilosa.
humilis, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 160 = crassifolia.
hypericifolia, Kegel, Gartenfl. (1857) 152 = fastigiata.
hyssopifolia, Nieber, in Spreng. Syst. iv. Cur. Post. 148 = polygalifolia.
inconspicua, *Benth. Fl. Austral.* i. 318.—Austral.
inornata, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 164.—Austral.
junces, *Bartl. in Lehm. Pl. Preiss.* i. 166.—Austral.
laevigata, F. Muell. Fragm. i. 101 = Zieria laevigata.
lanceolata, *F. Muell. l. c.* 66.—Austral.
lanifera, Bartl. in Lehm. Pl. Preiss. i. 165 = junces.
lanuginosa, *Endl. in Enum. Pl. Hueg.* 16.—Austral.
ledifolia, *Y. Gay, in Mém. Mus. Par.* vii. (1821) 450, in nota.—Austral.
ledophylla, F. Muell. Fragm. i. 67 = ledifolia.
lepidota, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 596.—Austral.
leptophylla, Turcz. l. c. xxv. (1852) II. 164 = inornata.
Machardianus, *F. Muell. Fragm.* ix. 115.—Austral.
macra, Bartl. in Lehm. Pl. Preiss. i. 167 = spathulata.
magnaguma, *Neve, ex Bartl. in Lehm. Pl. Preiss.* ii. 227.—Austral.
microphylla, *Sieber, ex Spreng. Syst. iv. Cur. Post.* 148.—Austral.
minutiflora, F. Muell. Fragm. i. 100 = Zieria obcordata.
mollis, *A. Cunn. in Lindl. Bot. Reg.* (1841) sub t. 47. —Austral.
multicostis, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 161 = crassifolia ?
nana, Hook. ic. Pl. t. 270 = polygalifolia.
nematophylla, *F. Muell. Fragm.* ii. 100.—Austral.
ovata, *Lindl. Bot. Reg.* (1841) sub t. 47.—Austral.
oxyantha, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 165.—Austral.
palefolia, A. Cunn. in Enum. Pl. Hueg. 16 = falcifolia.
paradoxa, DC. Prod. i. 722 = ledifolia.
parviflora, *Sm. Tracts*, 295. t. 6.—Austral.
penicillata, *Benth. Fl. Austral.* i. 323.—Austral.
pilonema, Lahill. Nov. Holl. Pl. i. 98. t. 126 = parviflora.
pilosa, *Labill. l. c.* 97. t. 124.—Austral.
pinnata, *Sm. Tracts*, 290.—Austral.
platyrrhachis, *F. Muell. Fragm.* viii. 37.—Austral.
polygalifolia, *Sm. Tracts*, 297. t. 7.—Austral.
pooraleoides, DC. Prod. i. 721 = elatior.
pteropoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 596.—Austral.
pubescens, Bartl. in Lehm. Pl. Preiss. ii. 227 = lanuginosa ?
pulchella, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 162.—Austral.
ramosa, *Benth. Fl. Austral.* i. 320.—Austral.
rhomboidea, *Cheel. Jc. Pl.* t. 732.—Austral.
Runzii, *Hueg. ex Heynh. Nom.* ii. 78, nomen.—Austral.
rosmarinifolia, A. Cunn. ex Endl. in Enum. Pl. Hueg. 16 = ledifolia.
rubiginosa, A. Cunn. ex Endl. l. c. = ledifolia.
salicifolia, *A. Cunn. ex Heynh. Nom.* ii. 78, nomen.—Austral.
scabra, *Lindl. Swan River App.* 76.—Austral.
semifertilis, F. Muell. Fragm. ii. 98 = elatior.
serrulata, *Sm. Tracts*, 292. t. 5.—Austral.

BORONIA:—
spathulata, *Lindl. Swan River App.* 78.—Austral.
spinescens, *Benth. Fl. Austral.* i. 319.—Austral.
stricta, Bartl. in Lehm. Pl. Preiss. i. 269 = lanuginosa.
subcaerulea, F. Muell. Fragm. ii. 100 = ramosa.
subsessilis, *Benth. Fl. Austral.* i. 322.—Austral.
tenuifolia, Hartl. in Lehm. Pl. Preiss. i. 168 = viminea.
tenuis, *Benth. Fl. Austral.* i. 320.—Austral.
teretifolia, Lindl. Swan River App. 76 = cymosa.
ternata, *Endl. Nov. Stirp. Dec.* 6.—Austral.
tetrandra, Hook. Journ. Bot. ii. (1840) 419 = pinnata.
tetrandra, *Labill. Nov. Holl. Pl.* i. 98. t. 125.—Austral.
tetrandra, Lindl. & Paxt. Flow. Gard. i. (1850-51) 35. t. 8 = pulchella.
tetrathecoides, DC. Prod. i. 723 = polygalifolia.
thymifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 165.—Austral.
triphylla, Sieber, ex Spreng. Syst. iv. Cur. Post. 148 = latifolia.
trutta, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 162 = megastigma.
variabilis, Hook. Journ. Bot. i. (1834) 255 = anemonifolia.
veronicea, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 11 = Zieria veronicea.
viminea, *Lindl. Swan River App.* 74.—Austral.

BORRAGINOIDES, Moench, Meth. 515 (1794) = **Trichodesma**, R. Br. (Borragin.).
aculeata, Moench, l. c. 516 = T. africanum.
sagittata, Moench, l. c. = T. indicum.

BORRAGO, Tourn. ex Linn. Syst. ed. I (1735) = **Borago**, Linn.

BORRERA, Spreng. Gen. i. 93 (1830) = BORRERIA, G. F. W. Mey. = **Spermacoce**, Linn. (Rubiac.).

BORRERIA, G. F. W. Mey. Prim. Fl. Esseq. 79. t. 1 (1818) = **Spermacoce**, *Linn.* (Rubiac.).
adscendens, Cham. & Schlecht. in Linnæa, iii. (1828) 340 = S. adscendens.
adsens, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 46 = Diodia verticillata?
affinis, DC. Prod. iv. 546 = S. diffusa.
alata, DC. l. c. 544 = S. alata.
alternans, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 283.—Porto Rico.
angustifolia, Cham. & Schlecht. in Linnæa, iii. (1828) 330.—Bras.
anthospermoides, DC. Prod. iv. 550.—Am. austr.
aralioides, Cham. & Schlecht. in Linnæa, iii. (1828) 339 = *umbellata*?
arenicoides, Griseb. Cat. Pl. Cub. 286.—Cuba.
argentea, Cham. in Linnæa, ix. (1834) 210.—Bras.
arida, DC. Prod. iv. 549.—Porto-Rico.
asclepiadea, Cham. & Schlecht. in Linnæa, iii. (1828) 330 = S. Poaya.
aspera, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 130 = S. aspera.
asperula, DC. Prod. iv. 548 = Diodia rigida?
asurgens, Griseb. in Goett. Abh. xxiv. (1879) 156 = Diodia dasycephala.
aturensis, Cham. & Schlecht. in Linnæa, iii. (1828) 345 = S. aturensis.
Barthingiana, DC. Prod. iv. 544 = S. latifolia.
basalis, Anders. Om Galap.-Oarnes Vegetat. 191.—Ins. Galapag.
brachystemonoides, Cham. & Schlecht. in Linnæa, iii. (1828) 314.—Bras.
capitata, DC. Prod. iv. 545 = S. capitata.
capitellata, Cham. & Schlecht. in Linnæa, iii. (1828) 319 = S. laevis.
centranthoides, Cham. & Schlecht. l. c. 327.—Bras.
commutata, Spreng. Syst. i. 404 = S. verticillata.
coriacea, Scheele, in Linnæa, xvii. (1843) 389 = S. tenella.
cornifolia, DC. Prod. iv. 545 = S. capitata.
corymbosa, DC. l. c. 550 = S. corymbosa.
crassifolia, C. A. Mey. ex Heynh. Nom. ii. 74, nomen.—Am. austr.
cupularia, DC. Prod. iv. 548 = S. pusilla.
cymosa, Cham. & Schlecht. in Linnæa, iii. (1828) 328 = S. cymosa.
densiflora, DC. Prod. iv. 542 = S. longifolia, *L'Hérit.*
dichotoma, Cham. & Schlecht. in Linnæa, iii. (1828) 340 = Diodia glabra.
diffusa, DC. Prod. iv. 545 = S. diffusa.

BORRERIA:—
discolor, Benth. in Hook. Kew Journ. iv. (1852) 199 = Spermacoce semierecta.
discolor, DC. Prod. iv. 545.—Ins. Philipp.
dispersa, Hook. f. in Trans. Linn. Soc. xx. (1849) 217. —Ins. Galapag.
distans, Cham. & Schlecht. in Linnæa, iii. (1828) 340 = S. distans.
divaricata, Hook. f. in Trans. Linn. Soc. xx. (1859) 219.—Ins. Galapag.
domingensis, Griseb. Cat. Pl. Cub. 141 = S. ocymoides.
dubia, Cham. in Linnæa, ix. (1834) 216 = *valerianoides*.
elongata, DC. Prod. iv. 547 = S. capitata.
equisetoides, Cham. & Schlecht. in Linnæa, iii. (1828) 333.—Bras.
ericæfolia, Hook. f. in Trans. Linn. Soc. xx. (1859) 218.—Ins. Galapag.
ericoides, Cham. & Schlecht. in Linnæa, iii. (1828) 326 = S. corymbosa.
eritrichoides, Griseb. Cat. Pl. Cub. 142.—Cuba.
eryngioides, Cham. & Schlecht. in Linnæa, iii. (1828) 315 = S. diffusa.
eupatorioides, Cham. & Schlecht. l. c. 327.—Bras.
falcifolia, Hook. f. in Trans. Linn. Soc. xx. (1859) 219. —Ins. Galapag.
ferruginea, DC. Prod. iv. 547 = S. capitata.
ferruginea, Mart. & Gal. in Bull. Acad. Brux. Sér. I. xi. i. (1844) 128 = Mitracarpum villosum.
flavoviridis, DC. Prod. iv. 548 = S. Poaya.
Fochsona, Miq. in Linnæa, xviii. (1844) 299 = S. latifolia.
globulariodes, Cham. & Schlecht. in Linnæa, iii. (1828) 312 = S. verticillata.
gracilis, Miq. ex Hook. f. Fl. Brit. Ind. iii. 200 = S. stricta.
gracilis, Scheele, in Linnæa, xvii. (1843) 340 = S. tenella.
gracillima, DC. Prod. iv. 548 = S. gracillima.
graminifolia, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 129 = S. verticillata.
gymnocephala, DC. Prod. iv. 549.—Bras.
Haenkeana, DC. l. c. 547 = S. tenella.
hebecarpa, Hook. ex A. Rich. Tent. Fl. Abyss. i. 347 = S. stricta.
humifusa, Mart. in Flora, xxiv. (1841) Beibl. ii. 68.—Bras.
Kleinii, Walp. Rep. ii. 464 = S. stricta.
kohautiana, Cham. & Schlecht. in Linnæa, iii. (1828) 311 = S. verticillata.
laevigata, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 130 = S. laevigata.
laevis, Griseb. in Goett. Abh. vii. (1857) 231 = S. laevis.
lasiocarpa, Walp. Rep. ii. 464 = S. stricta.
lasia, Cham. & Schlecht. in Linnæa, iii. (1828) 337.—Bras.
linearifolia, Hook. f. in Trans. Linn. Soc. xx. (1859) 217.—Ins. Galapag.
linifolia, DC. Prod. iv. 551 = S. laevis.
linoides, DC. l. c. 548 = S. tenella.
longiseta, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 128 = S. longiseta.
lutescens, DC. Prod. iv. 547 = S. tenella.
micrantha, Torr. & Gray, Fl. N. Am. ii. 98 = S. ocymoides.
microphylla, DC. Prod. iv. 546 = S. ocymoides.
minima, DC. l. c. 542 = S. verticillata.
Neesiana, DC. l. c. 543 = S. Neesiana.
neglecta, A. Rich. Tent. Fl. Abyss. i. 177.—Abyssin.
nervosa, DC. Prod. iv. 548 = S. Poaya.
oaxacana, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 130 = S. verticillata.
ocymoides, DC. Prod. iv. 546 = S. ocymoides.
organensis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 111.—Bras.
ovalifolia, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 129 = S. ocymoides.
ovalis, Anders. Om Galap.-Oarnes Vegetat. 192.—Ins. Galapag.
parviflora, G. F. W. Mey. Prim. Fl. Esseq. 88. t. 1 = S. ocymoides.
parvifolia, Hook. f. in Trans. Linn. Soc. xx. (1859) 218.—Ins. Galapag.
patula, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 128 = S. parviflora.
pentelitata, Miq. in Verh. Nat. Wet. Haarl. vii. (1851) 176 = S. latifolia.

BORRERIA:—
perpusilla, Hook. f. in Trans. Linn. Soc. xx. (1859) 218.—Ins. Galapag.
Perrottetii, DC. Prod. iv. 545.—Spermacoce latifolia.
platyphylla, DC. l. c. 549 = S. Poaya.
Poaya, DC. l. c. = S. Poaya.
podocephala, DC. l. c. 542 = S. verticillata.
Pohliana, DC. l. c. 550 = *eupatorioides*.
prasinata, DC. l. c. 548 = S. Poaya.
procera, Cham. in Linnæa, ix. (1834) 222 = S. tenella.
prostrata, Miq. in Verh. Nat. Wet. Haarl. vii. (1851) 177 = S. ocymoides.
psyllioides, DC. Prod. iv. 548 = S. tenella.
pterophora, Presl, Bot. Bemerk. 86.—Bras.
pumila, DC. Prod. iv. 548 = S. pumila.
pusilla, DC. l. c. = S. stricta.
radiata, DC. l. c. 542 = S. radiata.
radicans, Griseb. Cat. Pl. Cub. 142 = Nodocarpæa radicans.
ramisparsa, DC. Prod. iv. 544 = S. ocymoides.
remotifolia, DC. l. c. = Mitracarpum villosum.
repens, DC. l. c. 543 = S. ocymoides.
Rosburghiana, Walp. Rep. ii. 464 = S. stricta.
rubrostipulata, DC. Prod. iv. 546 = S. tenella.
saponariæfolia, Cham. & Schlecht. in Linnæa, iii. (1828) 335.—Bras.
saturejæfolia, Cham. & Schlecht. l. c. 313 = S. tenella.
scabiosoides, Cham. & Schlecht. l. c. 318.—Bras.
scabrida, DC. Prod. iv. 548 = S. latifolia.
scandens, DC. l. c. 546 = S. scandens.
setosa, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 131 = S. setosa.
Sideritis, Cham. & Schlecht. in Linnæa, iii. (1828) 301 = S. latifolia.
simplex, Griseb. Fl. Brit. W. Ind. 350.—Jamaica.
sphærica, DC. Prod. iv. 547 = S. laevis.
spinosa, Cham. & Schlecht. in Linnæa, iii. (1828) 340 = S. spinosa.
squamosa, Griseb. Cat. Pl. Cub. 142.—Cuba.
stricta, G. F. W. Mey. Prim. Fl. Esseq. 88. t. 1. f. 1—3 = S. verticillata.
strictissima, DC. Prod. iv. 547 = S. tenella.
strumpfioides, Griseb. Cat. Pl. Cub. 142.—Cuba.
suaveolens, G. F. W. Mey. Prim. Fl. Esseq. 81. t. 1 = S. tenella.
suberecta, Hook. f. in Trans. Linn. Soc. xx. (1859) 217. —Ins. Galapag.
subulata, DC. Prod. iv. 543 = Cruces subulata.
tampicana, DC. l. c. 544, pro parte = S. laevis.
tenella, Cham. & Schlecht. in Linnæa, iii. (1828) 317 = S. tenella.
tenera, DC. Prod. iv. 548 = S. tenera.
tenuis, DC. l. c. = S. tenuis.
tenuissima, Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr. iii. 324 = S. tenuissima.
tetracocca, Thw. Enum. Pl. Zeyl. 242 = Fergusonia zeylanica.
tetraphora, Miq. in Linnæa, xviii. (1843) 70 = S. latifolia.
thymifolia, Griseb. Fl. Brit. W. Ind. 350.—Ind. occ.
thymocephala, Griseb. Cat. Pl. Cub. 143 = S. verticillata.
trichantha, Miq. in Verh. Nat. Wet. Haarl. vii. (1851) 178.—Guiana.
umbellata, Spreng. Neue Entdeck. ii. 144.—Bras.
vaginata, Cham. & Schlecht. in Linnæa, iii. (1828) 340 = S. laevis.
valerianoides, Cham. & Schlecht. l. c. 336.—Bras.
verbenoides, Cham. & Schlecht. l. c. 331; iv. (1829) 550.—Bras.
verticillata, G. F. W. Mey. Prim. Fl. Esseq. 88 = S. verticillata.
virgata, Cham. & Schlecht. in Linnæa, iii. (1828) 324 = S. virgata.
Wydleriana, DC. Prod. iv. 545.—Porto Rico.

BORRICHIA, Adans. Fam. ii. 130 (1763). *COMPOSITÆ*, Benth. & Hook. f. ii. 266.
ADELMANNIA, Reichb. Consp. 130 (1828).
DIOMEDEA, Cass. in Bull. Soc. Philom. (1815) 176.
HELICTA, Cass. l. c. (1815) 167.
TRIMETRA, Moq. ex DC. Prod. vii. 263 (1838).
arborescens, *DC. Prod.* v. 462.—Am. trop.
argentea, *DC. l. c.*—Am. trop.
demissa, *DC. l. c.*—N. Granat.
frutescens, *DC. l. c.*—Am. bor.
parvulina, *DC. l. c.*—Peruv.

BORSCZOWIA, Bunge, in Act. Hort. Petrop. v. (1877) 648. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 68.
aralo-caspica, *Bunge, l. c.*—Reg. Casp.

BORSZCZOWIA, Bunge, in Act. Hort. Petrop. vi. (1879) 424 = prsec.

BORYA, Labill. Nov. Holl. Pl. i. 81. t. 107 (1804). *LILIACEAE*, Benth. & Hook. f. iii. 797.
BAUMGARTENIA, Spreng. Syst. ii. 91 (1825).
DAVIESIA, Poir. Encyc. Suppl. v. 480 (1817).
Cataractae, Endl. in Lehm. Pl. Preiss. ii. 43 = nitida.
gracilis, Endl. l. c. = nitida.
lucens, Poir. Encyc. viii. 615 = nitida.
nitens, Steud. Nom. ed. II. i. 192 = seq.
nitida, *Labill. Nov. Holl. Pl.* i. 81. t. 107.—Austral.
scirpoidea, Lindl. Swan River App. 57. t. 9 = nitida.
septentrionalis, *F. Muell. Fragm.* v. 41.—Austral.
sphaerocephala, R. Br. Prod 286 = nitida.
sublaxa, F. Muell. ex Benth. Fl. Austral. vii. 71 = nitida.

BORYA, Willd. Sp. Pl. iv. 711 (1805) = **Forestiera**, Poir. (Oleac.).
acuminata, Willd. l. c. = F. acuminata.
cassinoides, Willd. l. c. = F. acuminata.
distichophylla, Nutt. Gen. Am. ii. 232 = Buckleya distichophylla.
ligustrina, Willd. l. c. = F. ligustrina.
nitida, Willd. Enum. Hort. Berol. Suppl. [66] nomen = F. acuminata.
obovata, Desf. Cat. Hort. Par. ed. III. 88 = F. pinoides.
porulosa, Willd. Sp. Pl. iv. 711 = F. porulosa.
prinoides, Willd. Enum. Hort. Berol. Suppl. [66]; ex Spreng. Syst. iii. 900 = F. prinoides.
retusa, Willd. l. c. [66] = F. retusa.

BOSCA, Vell. Fl. Flum. 150. (1825); iv. t. 11 = **Funifera**, Leandr. ! (Thymel.).
stupacea, Vell. ll. cc.—Bras.

BOSCHERIA, Carr. in Rev. Hortic. (1872) 199 = BOSSCHERIA, De Vriese & Teijsm. = **Ficus**, Tourn. (Urtic.).

BOSCHIA, Korth. Verh. Nat. Gesch. Bot. 257. t. 69 (1849). *MALVACEAE*, Benth. & Hook. f. i. 213
HETEROPYXIS, Griff. Notul. iv. 524. t. 594 (1854).
acutifolia, *Mast. in Journ. Linn. Soc.* xiv. (1875) 508. —Borneo.
excelsa, *Korth. Verh. Nat. Gesch. Bot.* 258. t. 69.—Sumatra.
grandiflora, *Mast. in Journ. Linn. Soc.* xiv. (1875) 508.—Borneo.
Griffithii, *Mast. l. c.* 508.—Malaya.

BOSCHNIAKIA, C. A. Mey. (Bong. Veg. Sitka, 41) in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 157. *OROBANCHACEAE*, Benth. & Hook. f. ii. 985.
STELLARA, Fisch. & Kent. in DC. Prod. xi. 89 (1847).
glabra, *C. A. Mey. ex Bong. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 157.—Sibir.; Japon.; Am. bor.
himalaica, *Hook. f. & Thoms. in Fl. Brit. Ind.* iv. 327. —Reg. Himal.
Hookeri, *Walp. Rep.* iii. 479.—Am. bor. occ.
strobilacea, *A. Gray, in Pacif. Rail. Rep.* iv. 118.—Calif.

BOSCIA, Lam. Illustr. t. 895 (1804?). *CAPPARIDEAE*, Benth. & Hook. f. i. 108.
PODORIA, Pers. Syn. ii. 5 (1807).
angustifolia, Harv. in Harv. & Sond. Fl. Cap. i. Add. 19 = Niebuhria angustifolia.
angustifolia, *A. Rich. in Guill. & Perr. Tent. Fl. Senegj.* i. 26.—Afr. trop.
caffra, *Sond. in Linnaea*, xxiii. (1850) 8.—Afr. austr.
firma, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* xiv. (1884) 69, in nota.—Afr. trop.
integrifolia, Swana. in Flora, xxiii. (1840) ii. Beibl. 32 = angustifolia.
intermedia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 27 = angustifolia.
microphylla, *Oliver, Fl. Trop. Afr.* i. 93.—Afr. austr.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 164.—Afr. trop.
octandra, Fenzl, ex Radlk. in Sitzb. Math.-Phys. Acad. Munich. xiv. (1884) 62 = firma.
octandra, *Hochst. ex Radlk. l. c.*—Afr. trop.
reticulata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 28. —Abyss.
salicifolia, *Oliver, Fl. Trop. Afr.* i. 93.—Afr. trop.
senegalensis, Hochst. ex Walp. Ann. iii. 824 = angustifolia.
senegalensis, *Lam. Illustr.* t. 395.—Afr. trop.
urens, *Welw. in Oliver, Fl. Trop. Afr.* i. 93.—Afr. trop.

BOSCIA, Thunb. Prod. Pl. Cap. 32 (1794) = **Toddalia**, Juss. (Rutac.).
undulata, Thunb. l. c. = T. lanceolata.

BOSEA, Linn. Coroll. 5 (1737); (Bosia, Hort Cliff. 84 (1737).] *AMARANTACEAE*, Benth. & Hook. f. iii. 26.
ROBETIA, Moq. in DC. Prod. xiii. pt. 323 (1849)
Amherstiana, *Hook. f. Fl. Brit. Ind.* iv. 716.—Reg. Himal.
cannabina, *Lour. Fl. Cochinch.* i. 176.—Cochinch.
cypria, *Boiss. ex Hook. f. Fl. Brit. Ind.* iv. 716, in nota. —Cyprus.
humilis, Salisb. Prod. 154 = B. Vervamora.
irmeria, Roxb. Hort. Beng. 86; Fl. Ind. ii. 87 = Celtis Wightii.
Vervamora, *Linn. Sp. Pl.* 225.—Ins. Canar.

BOSISTOA, F. Muell. ex Benth. Fl. Austral. i. 359 (1863). *RUTACEAE*, Benth. & Hook. f. i. 390.
eudoliformis, *F. Muell. Fragm.* ix. 174.—Austral.
pentacocca, Baill. Hist. Pl. iv. 470 = aspindiformis.
aspindiformis, *F. Muell. ex Benth. Fl. Austral.* i. 359. —Austral.

BOSQUIERA, Thou. ex Baill. Adansonia, iii. (1862–63) 338. t. 10. *URTICACEAE*, Benth. & Hook. f. iii. 370.
Boiviniana, *Cord. ex Baill. l. c.* iii. (1862–63) 340. —Madag.
gymnadena, *Baker, Fl. Maurit.* 283.—Ins. Mascar.
Phoberos, *Baill. Adansonia*, viii. (1867) 72.—Afr. trop.
Thouarsiana, *Cord. ex Baill. l. c.* iii. (1862–63) 339.—Madag.

BOSSCHERIA, De Vriese & Teijsm. in Tijdschr. Nederl. Ind. xxiii. (1861) 218 = **Ficus**, Tourn. (Urtic.).
Minahassae, De Vriese & Teijsm. l. c. 214 = F. Minahassae.

BOSSIAEA, Vent. Jard. Cels, i. 7. t. 7 (1800). *LEGUMINOSAE*, Benth. & Hook. f. i. 478.
LALAGE, Lindl. Bot. Reg. t. 1722 (1834).
SCOTTEA, K. Br. in Ait. Hort. Kew. ed. II. iv. 269 (1812).
aculeata, *F. Muell. Fragm.* ii. 120.—Austral.
Aspinifolium, *Benth. Fl. Austral.* ii. 157.—Austral.
Armitii, *F. Muell. Fragm.* ix. 44.—Austral.
biloba, *Benth. in Enum. Pl. Hueg.* 36.—Austral.
bracteosa, *F. Muell. ex Benth. Fl. Austral.* ii. 166.—Austral.
Brownei, *Benth. l. c.* 163.—Austral.
buxifolia, *A. Cunn. in Field, N. S. Wales*, 348.—Austral.
calycina, *Benth. Fl. Austral.* ii. 159.—Austral.
carinalis, *Benth. in Mitch. Journ. Trop. Austr.* 290. —Austral.
cinerea, *K. Br. in Ait. Hort. Kew.* ed. II. iv. 266.—Austral.
coccinea, Bonpl. Jard. Malm. 128. t. 52 = cinerea.
conchinna, *Benth. Fl. Austral.* ii. 161.—Austral.
cordifolia, Sweet, Fl. Austral. t. 20 = cinerea.
cordigera, *Benth. ex Hook f. Fl. Tasm.* i. 95. t. 16.—Austral.
decumbens, F. Muell. Fragm. i. 9 = buxifolia.
dentata, *Benth. Fl. Austral.* ii. 156.—Australia.
distichа, *Lindl. Bot. Reg.* (1841) *Misc.* 88.—Austral.
distichoides, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 89 = foliosa.
divaricata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 285.—Austral.
egena, *F. Muell. in Hook. Kew Journ.* viii. (1856) 48. —Austral.
Endlicheri, *Meisen. in Lehm. Pl. Preiss.* i. 83.—Austral.
enaeta, Hook. f. Fl. Tasman. i. 94 = riparia.
ensata, *Sieber, ex DC. Prod.* ii 117.—Austral.
eriocarpa, *Benth. in Enum. Pl. Hueg.* 36.—Austral.
foliosa, *A. Cunn. in Field, N. S. Wales*, 347.—Australia.
Gilbertii, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 286 = eriocarpa.
Hendersonii, *Hort. ex Regel, Gartenfl.* (1866) 322. —Austral.
heterophylla, *Vent. Jard. Cels*, t. 7.—Austral.
humilis, Meisen. in Lehm. Pl. Preiss i. 85 = prostrata.
kiamensis, *Benth. Fl. Austral.* ii. 158.—Austral.
lalagoides, F. Muell. Fragm. iv. 12 = divaricata.
lanceolata, Sims, Bot. Mag. t. 1144 = heterophylla.
lenticularis, Lodd. Bot. Cab. t. 1238 = rhombifolia.
lenticularis, *Sieber, ex DC. Prod.* ii. 117.—Austral.
linnaeoides, G. Don, Gen. Syst. ii. 129 = prostrata.
linophylla, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 268.—Austral.
microphylla, *Sm. in Trans. Linn. Soc.* ix. (1808) 303. —Austral.
neo-anglica, *F. Muell. Fragm.* v. 106.—Austral.
nervosa, Meisen. in Bot. Zeit. xiii. (1855) 81 = eriocarpa.
Nummularia, Endl. Nov. Stirp. Dec. 27 = prostrata.
ornata, *Benth. Fl. Austral.* ii. 158.—Austral.
ovalifolia, Endl. Nov. Stirp. Dec. 26 = eriocarpa.
ovata, G. Don, Gen. Syst. ii. 128 = heterophylla.
ovata, Sm. in Trans. Linn. Soc. ix. (1808) 303 = prostrata.
oxyclada, Turcz. in Bull Soc. Nat. Mosc. xxvi. (1858) i. 284 = rufa.
paucyfolia, Benth. ex Lindl. Bot Reg. (1841) Misc. 53 = rufa.
pedunculariа, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 287.—Austral.
phylloclada, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 52.—Austral.
Preissi, *Messm. in Lehm. Pl. Preiss.* i. 82.—Austral.
prostrata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 268.—Australia.
pulchella, *Meisen. in Lehm. Pl. Preiss.* i. 84.—Australia.
rhombifolia, *Sieber, ex DC. Prod.* ii. 117.—Austral.
rigida, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1858) i. 285 = Preissii.
riparia, *A. Cunn. ex Benth. Fl. Austral.* ii. 166.—Austral.
Rossii, F. Muell. Fragm. iii. 94, 168 = Templetonia sulcata.
rotundifolia, DC. Prod. ii. 117 = rhombifolia.
rufa, *R. Br. in Ait. Hort Kew.* ed. II. iv. 266.—Austral.
rufa, Maund, Botanist, t. 81 = ensata.
ruptcola, *A. Cunn. ex Benth. Fl. Austral.* ii. 162.—Austral.
scolopendria, *Sm. in Trans. Linn. Soc.* ix. (1808) 308. —Australia.
Scortechinii, *F. Muell. Census*, 140, nomen.—Austral.
spinescens, Meisen. in Lehm. Pl. Preiss. i. 82 = rufa.
stenophylla, *F. Muell. Fragm.* i. 9.—Austral.
strigillosa, *Benth. Fl. Austral* ii. 157.—Austral.
sulcata, *Meisen. in Lehm. Pl. Preiss.* i. 81.—Austral.
tenuicaulis, R. Grah. in Edinb. N. Phil. Journ. xxix. (July 1840) 171 = cinerea.
virgata, Hook. Bot. Mag. t. 3986 = rufa.
Walkeri, *F. Muell. Fragm.* ii. 120.—Austral.
Webbii, *F. Muell. in Melb. Chem. et Drug.* (Dec. 1882).—Austral.

BOSSIENA, Pers. Syn. ii. 286 (1807) = **Bossiaea**, Vent. (Legum.).

BOSTRYCHANTHERA, Benth. in Benth. & Hook. f. Gen. ii. 1216 (1876). *LABIATAE*, Benth. & Hook. f. i. c.
deflexa, *Benth. l. c.*—China.

BOSWELLIA, Roxb. ex Colebr. in As. Res. ix. (1807) 379; Fl. Coromand. iii. 4. t. 207 (1819). *BURSERACEAE*, Benth. & Hook. f. i. 323.
LIBANUS, Colebr. in As. Res. ix. 382, in nota (1807), i. 6. f. 1.
PLORELLIA, Endl. Nov. Stirp. Dec. 39 (1839).

BOSWELLIA —
Ameero, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 505.—Ins. Socotra.
balsamifera, Spreng. Syst. ii. 313 = glabra.
Bhaw-Dajiana, *Birdw. in Trans. Linn. Soc.* xxvii. (1870) 144. t. 31.—Afr. trop.
Carterii, *Birdw. l. c.* 143.—Arabia.
clongata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 505.—Ins. Socotra.
Frereana, *Birdw. in Trans. Linn. Soc.* xxvii. (1870) 146. t. 31.—Afr. trop.
glabra, Roxb. Hort. Beng. 90; Fl. Ind. ii. 384 = serrata.
hirsuta, Sm. in Rees, Cyclc. xxxix. n. 2 = Canarium hirsutum.
integra, Bianco, Fl. Filip. ed. II. 242 = Bursera javanica.
javanica, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 619.—Java
neglecta, *S. Moore, in Journ. Bot.* xv. (1877) 67. t. 185.—Afr. trop.
papyrifera, *Hochst. in Flora*, xxvi. (1843) 81.—Abyss.
sacra, Flueck. Lehrb. Pharmak. (1867) 81 = Carterii.
serrata, *Roxb. ex Colebr. in As. Res.* ix. (1807) 279. t. 5.—Ind. or.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 505.—Ins. Socotra.
thurifera, Roxb. ex Flem. in As. Res. xi. (1810) 168 = serrata.

BOTELUA, Lag. in Varied. Cienc. ii. iv. (1805) 134 = Boutelous, Lag. (Gramin.).

BOTHERBE, Steud. ex Klatt, in Linnaea, xxxi. (1861-62) 569 = Calydorea, Herb. (Irid.).
bulbosa, Steud. ex Klatt, l. c. = C. speciosa.
campestris, Klatt, l. c. 563 = C. campestris.
gracilis, Klatt, l. c. 565 = C. nuda.
longispatha, Klatt, l. c. 564 = Sphenostigma longispatha.
luteola, Klatt, l. c. 563 = C. luteola.

BOTHRIOCHILUS, Lem. Jard. Fleur. iii. (1852-53) t. 295; Illustr. Hortic. iii. (1856) Misc. 80, 81 = Coelia, Lindl. (Orchid.).
bellus, Lem. Il. cc. = C. bella.

BOTHRIOCLINE, Oliver, ex Benth. in Hook. Ic. Pl. xii. 80. t. 1183 (1878). *COMPOSITAE*, Benth. & Hook. f. ii. 226.
Schimperi, *Oliver & Hiern, ex Benth. l. c.*—Abyssin.

BOTHRIOSPERMUM, Bunge, Enum. Pl. Chin. Bor. 47 (1832). *BORAGINEAE*, Benth. & Hook. f. ii. 858.
asperrimoides, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 150 = tenellum.
bicarinatum, Bunge, ex Steud. Nom. ed. II. i. 219 = sep.
bicornuculatum, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. (1835) 28 = chinense.
chinense, *Bunge, Enum. Pl. Chin. Bor.* 47.—China bor.
chinense, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 28 = Kusnetzowii.
Kusnetzowii, *Bunge, ex DC. Prod.* x. 116, cum cit. falsa.—Mongolia.
marifolium, A. DC. Prod. x. 118 = Hellotropium marifolium.
peruvianum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 96 = tenellum.
secundum, *Maxim. Prim. Fl. Amur.* 302.—China.
tenellum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. (1835) 24.—China bor.

BOTHRIOSPORA, Hook. f. in Hook. Ic. Pl. xi. 55. t. 1069 (1870). *RUBIACEAE*, Benth. & Hook. f. ii. 76.
corymbosa, *Hook. f. l. c.*—Am. austr.

BOTOR, Adans. Fam. ii. 326 (1763) = Psophocarpus, Neck. (Legumin.).

BOTRIA, Lour. Fl. Cochinch. 153 (1790) = Vitis, Tourn. (Ampel.).
africana, Lour. l. c. 154 = V. africana.

BOTROPHIS, Rafin. Med. Fl. i. 85 (1828) = Cimicifuga, Linn. (Ranunc.).
actaeoides, Rafin. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 = C. racemosa.
pumila, Rafin. Med. Fl. i. 86 = C. racemosa?
serpentaria, Rafin. l. c. 85 = C. racemosa.

BOTRYA, Juss. in Mém. Mus. Par. iii. (1817) 444 = BOTRIA, Lour. = Vitis, Tourn. (Ampelid.).

BOTRYADENIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 30 (1835) = Myriacotis, Less. (Compos.).
Gmelini, Fisch. & Mey. l. c. = M. Gmelini.

BOTRYANTHE, Klotzsch, in Wiegm. Archiv, ii. (1841) 190. t. 9 = Phiskanetia, Linn. (Euphorb.).
concolor, Klotzsch, l. c. 191.—Bras.
discolor, Klotzsch, l. c.—Bras.

BOTRYANTHUS, Kunth, Enum. Pl. iv. 210 (1843) = Muscari, Mill. (Liliac.).
alboviresc, Tod. Nuovi Gen. 76 = M. commutatum.
alpestris, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
atlanticus, Nym. Consp. 734 = M. racemosum.
Auckeri, Boiss. Diagn. Ser. I. v. 63 = M. Auckeri.
autumnalis, Heynh. Nom. ii. 76 = M. parviflorum.
Boraeanus, Jord. & Fourr. Ic. Pl. Eur. ii. 22 = M. botryoides.
breviscapus, Tod. Hort. Panorm. t. 5.—Hab.?
candidus, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
commutatus, Kunth, Enum. Pl. iv. 211 = M. racemosum.
compactus, Jord. & Fourr. Ic. Pl. Eur. ii. 24 = M. botryoides.
covicus, Jord. & Fourr. l. c. = M. botryoides.
dilutus, Elwes, ex Baker, in Gard. Chron. (1878) i. 708 = M. dilutum.
festivus, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
Gussonei, Tod. Ind. Sem. Panorm. (1872) 38; Nuov. Giorn. Bot. v. (1873) 157 = M. Gussonii.
heterophyllus, Jord. & Fourr. Ic. Pl. Eur. ii. 24 = M. Heldreichii.
Kerneri, Marches.—Cf. Oestr. Bot. Zeitschr. xxxii. (1882) 339 = M. botryoides.
Lelievrei, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
micranthus, Hort. Berol. ex Baker, in Gard. Chron. (1878) i. 799 = M. micranthum.
Mordoanus, Nym. Consp. 734 = M. Mordoanum.
neglectus, Jord. & Fourr. Ic. Pl. Eur. ii. 24 = M. neglectum.
neglectus, Kunth, Enum. Pl. iv. 679 = M. neglectum.
odorus, Kunth, l. c. 211 = M. racemosum.
paleus, Kunth, Enum. Pl. iv. 213 = M. pallens.
paradoxus, Kunth, l. c. = Hyacinthus paradoxus.
parviflorus, Kunth, l. c. = M. parviflorum.
pulchellus, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
racemosus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 160 = M. racemosum.
Sartori, Tod. Hort. Panorm. t. 5 = M. racemosum.
Sauli, Jaub. & Spach, Illustr. iv. 44 = M. moschatum.
speciosus, Marches.—Cf. Oestr. Bot. Zeitschr. xxxii. (1882) 339 = M. neglectum.
stenophyllus, Herbich. Stirp. Rar. Bucov. 30 (1855) 183 = Hyacinthus leucophaeus.
Strangwaysii, Kunth, Enum. Pl. iv. 679 = M. racemosum.
Szovitsianus, Hort. Bonn. ex Baker, in Gard. Chron. (1878) i. 799 = M. Szovitsianum.
vulgaris, Kunth, Enum. Pl. iv. 211 = M. botryoides.

BOTRYCARPUM, A. Rich. Bot. Méd. ii. 467 (1828), in adnot. = Ribes, Linn. (Saxifrag.).
nigrum, A. Rich. l. c. 490 = R. nigrum.

BOTRYCERAS, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 396. *ANACARDIACEAE*, Benth. & Hook. f. i. 496.
DARWINIA, Sprang. Syst. i. 370 (1825).
LAUROPHYLLUS, Roem. & Schult. Syst. iii. 80 (1815).
LAUROPHYLLUS, Thumb. Nov. Gen. vi. 304 (1792), nomen prius.
laurinum, *Willd. l. c.* 397.—Afr. austr.

BOTRYCOMUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 160 = Muscari, Mill. (Liliac.).
vulgaris, Fourr. l. c. = M. comosum.

BOTRYDIUM, Spach, Hist. Vég. Phan. v. 298 (1836) = Chenopodium, Linn.
arenarium, Spach, l. c. 299 = C. Botrys.
Schraderi, Spach, l. c. = C. foetidum.

BOTRYMOBUS, Miq. Fl. Ind. Bat. i. ii. 382 (1858) *ARTOCARPEAE*.
paniculata, *Miq. l. c.* 283.—Malaya.

BOTRYODENDRUM, Endl. Prod. Fl. Norf. 62 (1833) = Meryta, Forst. (Araliac.).
angustifolium, Endl. l. c. = M. angustifolia.
capitatum, Endl. ex Heynh. Nom. ii. 78 = Oreopanax capitatum.
coriaceum, Panch. ex Baill. Adansonia, xii. (1878) 153 = M. coriacea.
latifolium, Endl. Prod. Fl. Norf. 62 = M. latifolia.
macrophyllum, Rich, ex A. Gray, Bot. U. St. Expl. Exped. i. 723. t. 97 = M. macrophylla.
Sinclairii, Hook. f. Fl. N. Zel. 97 = M. Sinclairii.
taitense, Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 349.—Ins. Tahiti.

BOTRYOLOTUS, Jaub. & Spach, Illustr. i. 134. t. 63 = Trigonella, Linn. (Legumin.).
cachemyrianus, Jaub. & Spach, l. c. 135 = T. Emodi.
persicus, Jaub. & Spach, l. c. 134. t. 63 = T. elliptica.

BOTRYOPANAX, Miq. in Ann. Mus. Bot. Lugd. Bat i. 5 (1863-64) = Polyscias, Forst. (Araliac.).
borbonica, Miq. l. c. = Ins. Borbon.

BOTRYOPSIS, Miers, in Ann. Nat. & Mag. Hist. Ser. II. vii. 48 (1851) = Chondodendron, Ruiz & Pav (Menisperm.).
platyphylla, Miers, l. c. = C. platyphyllum.
Spruceana, Eichl. in Flora, xlvii. (1864) 394 = C. cretosum.

BOTRYOROPIS, Presl, Epim. Bot. 220 (1849) = Barringtonia, Forst. (Myrtac.).
insonensis, Presl, l. c. = B. insonensis.

BOTRYOSICYOS, Hochst. in Flora, xxvii. (1844) Bes. Beil. 8 = Dendrosicyos, Linn.
pentaphyllum, Hochst. l. c. = D. pentaphylla.

BOTRYPHILE, Salisb. Gen. Pl. Fragm. 25 (1866) = Muscari, Mill. (Liliac.).
botryoides, Salisb. l. c. = M. botryoides.
parviflorus, Salisb. l. c. = M. parviflorum.
racemosus, Salisb. l. c. = M. racemosum.

BOTRYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 138 = Teucrium, Linn. (Labiat.).
chamaedryoides, Fourr. l. c. = T. Botrys.

BOTTIONEA, Cels, in Mem. Acc. Torin. xxxvii. (1834) 43. t. 1. *LILIACEAE*, Benth. & Hook. f. iii. 789.
ENDOCOMA, Rafin. Fl. Tellur. ii. 57 (1836).
TRICHOPETALUM, Lindl. Bot. Reg. t. 1856 (1833).
thysanthoides, *Colla, l. c.* 45. t. 1.—Chili.

BOUCHEROSIA, Wight & Arn. Contrib. Bot. Ind. 34 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 784.
ASTERANTHES, Miken, in Nov. Act. Nat. Cur. xvii. (1835) 593. t. 41.
DESMIDOCHUS, Reichb. Consp. 120 (1828).
DESMIDORCHIS, Ehrenb. in Linnaea, iv. (1829) 94.
HUTCHINSIA, Wight & Arn. Contrib. Bot. Ind. 34 (1834).
HUTCHINSIA, D. Dietr. Syn. i. 879 (1839).
Aucosia, *Hart, in Trans. Roy. Irish Acad.* xxviii. (1868) 496.—Palaestin.
acutangula, *Decne. in DC. Prod.* viii. 668.—Afr. trop.
adscendens, Wall. Cat. n. 6146, partim = Cosmiana adulis.
Aucheriana, *Decne. in DC. Prod.* viii. 649.—Arabia; Persia; Ind. bor. occ.
campanulata, Wight, Ic. vi. t. 1997 = umbellata.
canescens, *Wight & Arn. Contrib. Bot. Ind.* 34.—Ind. or.

BOUCHEROSIA :—
cylindrica, *Brongn. in Bull. Soc. Bot. Fr.* vii. (1860) 900.—Abyss.
Decaisniana, *Linn. Herb. Gén. Amat. Sér.* II. iv. t. 21 ; *Decne. in DC. Prod.* viii. 648.—Afr. trop.
diffusa, *Wight, Ic.* vi. t. 1987.—Ind. or.
edulis, Edgew. in Journ. Linn. Soc. vi. (1862) 205 = Carallume edulis.
Forskaëli, *Decne. in DC. Prod.* viii. 648.—Arabia.
Gussoniana, *Hook. f. Bot. Mag.* t. 6137 in textu.—Algeria.
Hitchinsii, Hook. f. Fl. Brit. Ind. iv. 78, lapsu = B. Hutchinia.
Hutchinia, *Decne. in DC. Prod.* viii. 649.—Ind. or. incarnata, *N. E. Br. in Journ. Linn. Soc.* xvii. (1878) 166.—Afr. austr.
lasiantha, Wight, Ic. vi. t. 1986 = crenulata.
mammillaris, *N. E. Br. in Journ. Linn. Soc.* xvii. (1878) 166.—Afr. austr.
marocana, *Hook. f. Bot. Mag.* t. 6137.—Marocco.
Munbyana, *Decne. ex Munby, Fl. Algér.* 26.—Algeria.
pauciflora, *Wight, in Madras Journ. Sc.* vi. (1837) t. ad p. 473.—Ind. or.
quadrangula, Decne. in DC. Prod. viii. 664, in syn. = Forskaëli.
Russeliana, *Courb. ex Brongn. in Bull. Soc. Bot. Fr.* vii. (1860) 900.—Orient.
sinuica, *Decne. in DC. Prod.* viii. 649.—Arabia.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 79.—Ins. Socotr.
Stackiana, Boiss. Fl. Orient. iv. 68 = Carallume edulis.
umbellata, *Wight & Arn. Contrib. Bot. Ind.* 84.—Ind. or.

BOUCHARDATIA, Baill. Adansonia, vii. (1867) 350 = **Melicope**, Forst. (Rutac.).
austrotis, Baill. l. c. 351. C. 10 = M. neurococca.
neurococca, Baill. l. c. ix. (1868) 110 = M. neurococca.

BOUCHEA, Cham. in Linnaea, vii. (1832) 222. *VERBENACEAE*, Benth. & Hook. f. ii. 1144.
CHASCANUM, E. Mey. Comm. Pl. Afr. Austr. 275 (1835).
PLEUROSTIGMA, Hochst. ex Schnizl. in Flora, (1842) t. Beib. 144.
PLEXIPUS, Rafin. Fl. Tellur. ii. 104 (1836).
adenostachya, *Schau. in DC. Prod.* xi. 560.—Afr. trop.
agrestis, *Schau. l. c.* 558.—Bras.
cernua, *Schau. l. c.* 559.—Afr. austr.
copaiferoides, C. Gay, Fl. Chil. v. 26. t. 55 = Priva laevis.
cuneifolia, *Schau. in DC. Prod.* xi. 559.—Afr. austr.
Ehrenbergii, *Cham. in Linnaea*, vii. (1832) 253.—Ins. S. Doming.
grazopensis, *Schau. in DC. Prod.* xi. 560.—Afr. austr.
Hanningtonii, *Oliver, in Hook. Ic. Pl.* t. 1446.—Afr. trop.
hederacea, *Sond. in Linnaea*, xxiii. (1850) 36.—Afr. austr.
hydrastoides, *Walp. Rep.* iv. 12.—Ind. or.
incrassata, *Lange, Ind. Sem. Haun.* (1870) 31.—Hab. ?
lactevirens, *Schau. in DC. Prod.* xi. 557.—Bras.
latifolia, *Harv. Thes. Cap.* ii. 57.—Afr. austr.
linifolia, *A. Gray, in Am. Journ. Sc. Ser.* II. xvi. (1853) 98.—N. Mexic.
marrubiifolia, *Schau. in DC. Prod.* xi. 558.—Afr. trop.; Arab.; Scind.
pinnatifida, *Schau. l. c.* 560.—Afr. austr.
pseudogervão, *Cham. in Linnaea*, vii. (1832) 252.—Bras.
pterygocarpa, *Schau. in DC. Prod.* xi. 558.—Nubia.
pubescens, Schau. l. c. 560 = pumila.
pumila, *Schau. l. c.*—Afr. austr.
sessilifolia, *Vatke, in Linnaea*, xliii. (1880-82) 529.—Afr. trop.
spathulata, *Torr. Bot. Mex. Bound.* 126.—Am. bor. occ.

BOUCHERIA, DC. ex Dun. in DC. Prod. xiii. t. 589 (1852). *SOLANACEAE*, Benth. & Hook. f. ii. 906.
LEUCANTHERA, Schlecht. in Linnaea, xxv. (1852) 358.
ovata, *DC. ex Dun. l. c.*—Texas; Mexic.
procumbens, *DC. ex Dun. l. c.*—Mexic.

BOUEA, Meisn. Gen. 75 ; Comm. 55 (1837). *ANACARDIACEAE*, Benth. & Hook. f. i. 420.
CAMBESSEDEA, Wight & Arn. Prod. i. 170 (1834).
angustifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 204.—Borneo.
Brandisiana, Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 39.—Burmanica.
burmanica, *Griff. Pl. Cantor*, 14.—Burma ; Malaya.
diversifolia, Miq. Fl. Ind. Bat. Suppl. i. 522 = microphylla.
Gandaria, Blume, Mus. Bot. Lugd. Bat. i. 204 = macrophylla ?
macrophylla, *Griff. Pl. Cantor*, 15 ; *Notul.* iv. 420.—Malaya.
microphylla, *Griff. Il. cc.* 15, 423.—Malaya.
myrsinoides, Blume, Mus. Bot. Lugd. Bat. i. 204 = microphylla ?
oppositifolia, Meisn. Gen. 75 ; Walp. Rep. i. 556 = burmanica.

BOUGAINVILLAEA, Comm. ex Juss. Gen. 91 (1789) (Bugiovillaea) ; Choisy, in DC. Prod. xiii. II. 437 (1849). *NYCTAGINEAE*, Benth. & Hook. f. iii. 7.
JOSEPHA, Vell. Fl. Flum. 154 (1825) ; iv. t. 16.
bracteata, Pers. Syn. i. 418 = spectabilis.
brasiliensis, Lund, ex Choisy, in DC. Prod. xiii. II. 437 = virescens.
brasiliensis, Raznsch. Nom. ed. III. 112 ; [Nees, in] Princ. Neuw. Reise, i. 44 = spectabilis.
frondosa, *Griseb. in Goett. Abh.* xix. (1874) 89.—Reg. Argent.
glabra, *Choisy, in DC. Prod.* xiii. II. 437.—Bras.
infesta, *Griseb. in Goett. Abh.* xxiv. (1879) 40.—Reg. Argent.
patagonica, *Decne. in Orbign. Voy.* t. 8.—Patagon.
peruviana, *Humb. & Bonpl. Pl. Aequin.* i. 173. t. 49. —Peru.
peruviana, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 89 = spectabilis.
pomacea, *Choisy, in DC. Prod.* xiii. II. 438.—Bras.
praecox, *Griseb. in Goett. Abh.* xxiv. (1879) 40.—Reg. Argent.
racemosa, Blanco, Fl. Filip. ed. I. 307 = Pisonia excelsa.
rubicunda, Schott, ex Rohrb. in Mart. Fl. Bras. xiv. II. 351 = glabra.
speciosa, Schnizl. Ic. Fam. Nat. t. 104 = spectabilis.
spectabilis, *Willd. Sp. Pl.* ii. 348.—Bras.
stipitata, *Griseb. in Goett. Abh.* xix. (1874) 88.—Reg. Argent.
virescens, *Choisy, in DC. Prod.* xiii. II. 437.—Bras.

BOUGAINVILLEA, Spach, Hist. Vég. Phan. x. 516 (1841) = **Bougainvillaea**, Juss. (Nyctag.).

BOUGUERIA, Decne. in Ann. Sc. Nat. Sér. II. v. (1836) 193. *PLANTAGINEAE*, Benth. & Hook. f. ii. 1224.
nubicola, *Decne. l. c.*—Peruv. ; Boliv.

BOULARDIA, F. Schultz, Arch. Fl. Fr. et Allem. 108 (1847) = **Orobanche**, Tourn.
latisquama, F. Schult. l. c. 104.—Algeria.

BOUPHON, Lem. in Orb. Dict. ii. 686 (1842) = seq.

BOUPHONE, Lem. in Orb. Dict. ii. 778 (1842) = **Buphane**, Herb. (Amaryll.).

BOURGAEA, Cos. Notes Crit. 39 (1849) = **Cynara**, Vaill. (Compos.).
humilis, Cos. l. c. = C. humilis.
Tournefortii, Willk. in Linnaea, xxv. (1852) 39 = C. Tournefortii.

BOURGAEA, Sosp. Introd. 158 (1777). *BORAGIN.* ?

BOURJOTIA, Pomel, Nouv. Mat. Fl. Atl. 89 (1874) = Heliotropium, Linn. (Borragin.).
Kralikii, Pomel, l. c. = H. Kralikii.
pterocarpa, Pomel, l. c. 90 = H. pterocarpum.

BOURRERIA, P. Br. Hist. Jamaic. 168 (1756). *BORAGINEAE*, Benth. & Hook. f. ii. 840.
HEURERIA, Jacq. Enum. Pl. Carib. 2 (1760).
CREMATOMIA, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. iii. (1869) 300.

BOURRERIA :—
HYMENEMYTES, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 26. t. 6.
MORELOSIA, La Llav. & Lex. Nov. Veg. Descr. i. 1 (1824).
Andrieuxii, *Hemsl. Biol. Centr. Am. Bot.* ii. 369.—Mexic.
aspera, G. Don, Gen. Syst. iv. 390 = Ehretia laevis.
baccata, *Raffn. Sylva Tellur.* 42.—Hab. ?
calophylla, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 528.—Cuba.
casasinifolia, *Griseb. l. c.*—Cuba.
cestroides, *Spreng. Syst.* iv. *Cur. Post.* 66.—Am. austr.
clariuscula, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 304.—Am. austr.
dichotoma, Rottl. ex G. Don, Gen. Syst. iv. 390 = Ehretia laevis ?
divaricata, *G. Don, l. c.* 387.—Cuba.
domingensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 528.—Ind. occ.
exsucca, *Vacq. Enum. Pl. Carib.* 14.—Am. austr.
formosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 369.—Mexic.
glabra, G. Don, Gen. Syst. iv. 390 = Cordia Myxa.
grandiflora, *Bertol. Pl. Guatim.* 10.—Guatemala.
havanensis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 307.—Cuba.
Hinckia, *Hemsl. Biol. Centr. Am. Bot.* ii. 370.—Mexic.
laevis, G. Don, Gen. Syst. iv. 390 = Ehretia laevis.
laxa, *G. Don, l. c.*—Ins. Madagar.
linearis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 310.—Ind. occ.
microphylla, *Griseb. Cat. Pl. Cub.* 210.—Cuba.
ovata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 308.—Ind. occ.
punctata, G. Don, Gen. Syst. iv. 390 = Ehretia laevis.
Radula, *G. Don, l. c.*—Ins. S. Doming.
recurva, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 308.—Ind. occ.
reticulata, *Griseb. Cat. Pl. Cub.* 210.—Cuba.
retusa, G. Don, Gen. Syst. iv. 390 = Ehretia retusa.
revoluta, *H. B. & K. Nov. Gen. et Sp.* iii. 67.—Mexic.
rigida, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 304.—Ind. occ.
setosa, G. Don, Gen. Syst. iv. 390 = Ehretia setosa.
spathulata, *Hemsl. Biol. Centr. Am. Bot.* ii. 370.—Mexic.
spinifex, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 528.—Cuba.
succulenta, *Vacq. Enum. Pl. Carib.* 14.—Ind. occ.
tomentosa, *G. Don, Gen. Syst.* iv. 390.—Ind. occ.
uniflora, G. Don, l. c. = Ehretia uniflora.
virgata, *G. Don, l. c.* 389.—Ind. occ.

BOUSSINGAULTIA, H. B. & K. Nov. Gen. et Sp. vii. 194. t. 646 (1825). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 77.
TANDONIA, Moq. in DC. Prod. xiii. II. 296 (1849).
baselloides, *H. B. & K. Nov. Gen. et Sp.* vii. 196. t. 646.—Ecuador.
cordifolia, *Tenore, Ind. Sem. Neap.* (1852) ex *Ann. Sc. Nat.* Sér. III. xix. (1853) 355.—Hab. ?
floribunda, *Moq. in DC. Prod.* xiii. II. 229.—N. Granat.
gracilis, *Miers, in Journ. Bot.* ii. (1864) 161. t. 18.—Bras.
Lachaumei, *Carr. in Rev. Hortic.* (1879) 279.—Cuba.
leptostachys, Moq. in DC. Prod. xiii. II. 229 = baselloides.
ramosa, *Hemsl. Biol. Centr. Am. Bot.* iii. 27.—Guatemala.

BOUTELOUA, Lag. in Varied. Cienc. ii. iv. (1806) 184 (Boteloa). *GRAMINEAE*, Benth. & Hook. f. iii. 1166.
ACTINOCHLOA, Willd. ex Roem. & Schult. Syst. ii. 23, 417 (1818).
ARISTIDIUM, Endl. Gen. 94 (1836).
ATHEROPOGON, Muhl. in Willd. Sp. Pl. iv. 907 (1806).
CHONDROSIUM, Desv. Journ. Bot. iii. (1813) 68.
CORETHRUM, Vahl, in Skr. Naturk. Selsk. Kiobenh. vi. (1810) 85.
DIMERA, DC. Cat. Hort. Mons. 104 (1813), pro parte.
EUTRIANA, Trin. Fund. Agrost. 161 (1820).

BOUTELOUA:—
Heterostega, Desv. in Nov. Bull. Soc. Philom. ii. (1810, 188; Desv. Journ. Bot. iii. (1812) 68.
Heterostoma, Kunth, in Mém. Mus. Par. ii. (1815) 73.
Nuttelera, Willd. ex Steud. Nom. ed. II. ii. 192 (1841).
Polydon, H. B. & K. Nov. Gen. et Sp. i. 174. t. 55 (1815).
Triaena, H. B. & K. l. c. 178. t. 61 (1815).
Triathera, Desv. in Nov. Bull. Soc. Philom. ii. (1810) 188.
Triplathera, Endl. Gen. 94 (1836).
aristidoides, *Griseb. Fl. Brit. W. Ind.* 537, in obs.—Mexic.
barbata, *Lag. Varied. Cienc.* ii. iv. (1805) 141; *Gen. et Sp. Nov.* 5.—Mexic.; Ins. Philipp.
brevifolia, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 93 = eriopoda.
bromoides, *Lag. Gen. et Sp. Nov.* 5.—Mexic.
Buckei, *Scribn. ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 179.—Texas.
chondrosioides, Benth. ex S. Wats. l. c. = ovata.
ciliata, *Griseb. in Goett. Abh.* xxiv. (1879) 302.—Reg. Argent.
curtipendula, Torr. in Emory, Notes Mil. Reconnois. 153 = racemosa.
distichn, *Benth. in Journ. Linn. Soc.* xix. (1882) 106.—Am. trop.
elatior, *Griseb. Fl. Brit. W. Ind.* 537.—Ind. occ.
eriopoda, *Torr. in Pacif. Rail. Rep.* iv. *Bot.* 155.—Am. bor.
fonsa, Bigelow, in Whipple Exped. Add. p. ii = hirsuta.
gracilis, Steud. Nom. ed. II. i. 719 = oligostachya.
gracilis, Vasey, in Wheeler, Exped. 287 = aristidoides.
Havardi, *Vasey, ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 179.—Texas.
hirsuta, *Lag. Varied. Cienc.* ii. iv. (1805) 141; *Gen. et Sp. Nov.* 5.—Mexic.
hirta, Hort. Madr. ex Lag. l. c. 141 = hirsuta.
Humboldtiana, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 583 = bromoides.
humilis, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 496.—Reg. Argent.
juncifolia, *Lag. Gen. et Sp. Nov.* 5.—Mexic.; Ind. occ.
juncifolia, Vasey, Cat. Gr. Un. St. 62 = bromoides.
litigiosa, Lag. Gen. et Sp. Nov. 5 = juncifolia.
lophostachya, *Griseb. in Goett. Abh.* xxiv. (1879) 302.—Reg. Argent.
melicaeformis, Hornem. ex Steud. Nom. ed. I. 91, in syn. = racemosa.
meliceides, Steud. l. c. 115 = racemosa.
multiseta, *Griseb. in Goett. Abh.* xxiv. (1879) 303.—Bras.; Reg. Argent.
nana, *Griseb. l. c.*—Reg. Argent.
oligostachya, *Torr. ex A. Gray, Man.* ed. II. 558; *et in Pacif. Rail. Rep.* iv. *Bot.* 155.—Am. bor.
ovata, *Lag. Gen. et Sp. Nov.* 5.—Mexic.
pendula, Hort. Madr. ex Lag. in Varied. Cienc. ii. iv. (1805) 141 = racemosa.
pilosa, *Benth. ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 179.—Ins. Galap.
polystachya, *Torr. in Pacif. Rail. Rep.* v. 366. t. 10.—Am. bor.
porphyrantha, Wright, in Sauv. Fl. Cub. 192 = juncifolia.
prostrata, *Lag. in Varied. Cienc.* ii. iv. (1805) 141; *Gen. et Sp. Nov.* 5.—Mexic.
pumila, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 93 = polystachya.
pusilla, *Vasey, in Bull. Torrey Bot. Club*, xi. (1884) 6.—N. Mexic.
racemosa, *Lag. in Varied. Cienc.* ii. iv. (1805) 141; *Gen. et Sp. Nov.* 5.—Mexic.
Rothrockii, *Vasey, Grasses U. St.* 33, nomen.—Am. bor.
scorpioides, *Lag. Gen. et Sp. Nov.* 5; *S. Wats. in Proc. Am. Acad.* xviii. (1883) 178.—Mexic.
simplex, *Lag. in Varied. Cienc.* ii. iv. (1805) 141.—Peruv.; Ins. Philipp.
tenuis, *Griseb. in Goett. Abh.* xxiv. (1879) 303.—Mexic.; Reg. Argent.
texana, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 180.—Texas.
Triathera, *Benth. in Journ. Linn. Soc.* xix. (1882) 104.—Mexic.
trifida, *Thurb. ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 177.—Mexic.

BOUTONIA, Boj. Hort. Maurit. 282 (1837) = **Mallotus**, Lour. (Euphorb.).
acuminata, Baill. Etud. Gén. Euphorb. 401 = M. Bellicianus.
mascarenia, Boj. Hort. Maurit. 282 = M. integrifolius.

BOUTONIA, DC. Rev. Bign. 18 (1838) = **Feriblasma**, DC. (Bignon.).
cuspidata, DC. l. c. = P. cuspidatum.

BOUTONIA, Hort. Erfurt. ex Steud. Nom. ed. II. i. 219 (1840) = **Geodiantia**, Sav.
pomifera, Hort. Erfurt. ex Steud. l. c. = G. paniculata.

BOUVARDIA, Salisb. Parad. Lond. t. 88 (1805).
RUBIACEAE, Benth. & Hook. f. ii. 86.
Argunbeia, Cav. Ic. vi. 61. t. 572 (1801).
Bouwardia, Polak. in Linnaea, xli. (1877) 565.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 384.—Mexic.
bicolor, *Kunze, in Linnaea*, xx. (1847) 24.—Mexic.
candidissima ×, Hort.—Cf. Gard. Chron. (1878) 714.
Cavanillesii, *DC. Prod.* iv. 366.—Mexic.
chlorantha, *Bert. ex Roem. & Schult. Syst.* iii. *Mant.* 116.—Am. austr.
chrysantha, *Mart. Del. Sem. Hort. Monac.* (1846) 4.—Mexic.
coccinea, Link, Enum. Hort. Berol. i. 189 = triphylla.
coccinea, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 272.—Guiana.
cordifolia, *DC. Prod.* iv. 366.—Mexic.
corymbosa, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 46.—Nicarag.
crocata, *Van Houtte, Cat.* (1846).—Hab.?
Davisoni, *Hort.—Cf. Gard. Chron.* (1872) 7.—Hab.?
discolor, Hook. & Arn. Bot. Beech. Voy. 438 = Rondeletia dubia.
ferruginea, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 273.—Mexic.
flava, *Decne. in Van Houtte, Fl. des Serres*, Sér. I. i. (1845) 90. t. 38.—Mexic.
glaberrima, *Engelm. in Wislis. Tour N. Mexico*, 106.—Mexic.
glabra, *Polak. in Linnaea*, xli. (1877) 565.—Costa Rica.
havanensis, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 272.—Ind. occ.
hirsuta, *A. Rich. l. c.*—Ind. occ.
hirtella, *H. B. & K. Nov. Gen. et Sp.* iii. 384.—Mexic.
Houttenas, *Schlecht. in Fl. des Serres*, Sér. I. x. (1854-55) 149.—Mexic.
Humboldti, *Hort.—Cf. Gard. Chron.* (1872) 714, 716.—Hab.?
hypoleuca, *Benth. Pl. Hartw.* 288.—Mexic.
jacquinii, H. B. & K. Nov. Gen. et Sp. iii. 385 = triphylla.
jasminiflora, *Hort.—Cf. Gard. Chron.* (1872) 215.—Hab.?
laevis, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 286.—Mexic.
leiantha, *Benth. Pl. Hartw.* 85; *Decne. in Rev. Hortic. Sér.* III. v. (1851) 81.—Guatemala; Mexic.
linearis, *H. B. & K. Nov. Gen. et Sp.* iii. 383.—Mexic.
longiflora, H. B. & K. l. c. 386 = Houstonia longiflora.
microphylla, *Schlecht. in Linnaea*, xxvi. (1853-55) 113.—Mexic.
mollis, *Linden, Cat.* (1848).—*Cf. Schlecht. l. c.* 55.—Mexic.
multiflora, Schult. in Roem. & Schult. Syst. iii. 118 = Cavanillesii.
mutabilis, Hort. Berol. ex Walp. Ann. v. 127 = versicolor.
myrtifolia, *Schlecht. in Linnaea*, xxvi. (1853-55) 121.—Mexic.
obovata, Benth. Pl. Hartw. 16 = scabra.
obovata, *H. B. & K. Nov. Gen. et Sp.* iii. 385.—Mexic.
Oriana ×, *Pars. ex Van Houtte, Fl. des Serres*, xii. (1857) 139. t. 1265.
ovata, *A. Gray, Pl. Wright.* ii. 67.—N. Mexic.
subrubra, Decne. in Fl. des Serres, Sér. I. i. (1845) 90, in descr. = flava.
quadriformis, Steud. Nom. ed. II. i. 200, in syn. = seq.
quaternifolia, *DC. Prod.* iv. 365.—Mexic.
quinquefolia, *Dehnh. Revista Napol.* i. iii. 167.—Mexic.

BOUVARDIA:—
racemosa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834).—Peruv.
rosea, *Schlecht. in Linnaea*, xxvi. (1853-55) 116.—Calif.
scabra, *Hook. & Arn. Bot. Beech. Voy.* 427.—Mexic.
scabrida, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 287.—Mexic.
scandens, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 273.—Hab.?
Schiedeana, *Schlecht. in Linnaea*, xxvi. (1853-55) 123.—Mexic.
splendens, R. Grah. in Bot. Mag. t. 3781 = triphylla.
strigillosa, [Bast. in] Loud. Hort. Brit. Suppl. iii. 502 = seq.!
strigosa, Benth. Pl. Hartw. 75 = Rondeletia strigosa.
tenuiflora, *Hort. Berol. ex Schlecht. in Linnaea*, xxvi. (1853-55) 97.—Mexic.
ternifolia, Schlecht. l. c. 98 = triphylla.
tolucana, *Hook. & Arn. Bot. Beech. Voy.* 427.—Mexic.
triflora, H. B. & K. Nov. Gen. et Sp. iii. 386. t. 289 = Houstonia triflora.
triphylla, *Salisb. Parad. Lond.* t. 88.—Mexic.
umbellata, *Hort.—Cf. Gard. Chron.* (1872) 714.—Hab.?
uniflora, *Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 272.—Am. trop.
versicolor, *Ker-Gawl. Bot. Reg.* t. 345.—Am. austr.
viminalis, *Schlecht. in Linnaea*, xxvi. (1853-55) 130.—Mexic.
viperalis, *Schlecht. l. c.* 114.—Mexic.
Vreelandi, Hort. ex Fl. et Pomol. (1872) 174 = Davisoni.
xylosteoides, *Hook. & Arn. Bot. Beech. Voy.* 426.—Mexic.

BOUZETTA, Montrous. in Mém. Acad. Lyon, x. (1860) 192. *RUTACEAE*?, Benth. & Hook. f. i. 989.
maritima, *Montrous. l. c.*—N. Caled.

BOVEA, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 253 = Lindenbergia, Lehm. (Scrophulac.).
sinuata, Decne. l. c. = L. sinuata.

BOWDICHIA, H. B. & K. Nov. Gen. et Sp. vi. 276 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 597.
Sebipira, Mart. Reise, 787 (1823).
brevipes, Benth. in Ann. Wien. Mus. ii. (1838) 89 = virgilioides.
ferruginea, Benth. l. c. = virgilioides.
ferruginea, Walp. Rep. i. 808 = virgilioides.
floribunda, Benth. in Ann. Wien. Mus. ii. (1838) 89 = virgilioides.
major, Mart. Syst. Mat. Med. Bras. 58 = virgilioides.
nitida, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. i. 313.—Bras.
pubescens, Benth. in Ann. Wien. Mus. ii. (1838) 89 = virgilioides.
Sebipira, Steud. Nom. ed. II. i. 220 = virgilioides.
virgilioides, *H. B. & K. Nov. Gen. et Sp.* vi. 276.—Am. austr.

BOWENIA, Hook. Bot. Mag. t. 5828 (1863) et 6008. *CYCADACEAE*, Benth. & Hook. f. iii. 446.
spectabilis, *Hook. l. c.* 5398.—Austral.

BOWIEA, Harv. ex Hook. f. in Bot. Mag. t. 5619 (1867). *LILIACEAE*, Benth. & Hook. f. iii. 786.
volubilis, *Harv. l. c.*—Afr. austr.

BOWIEA, Haw. in Phil. Mag. lxiv. (1824) 399 = **Aloë**, Tourn. (Liliac.).
africana, Haw. l. c. = A. Bowiea.
myriacantha, Haw. l. c.; et N. S. i. (1827) 123 = A. myriacantha.

BOWKERIA, Harv. Thes. Cap. i. 24. t. 37 (1859). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 987.
triphylla, *Harv. l. c.*—Afr. austr.

BOWLESIA, Ruiz & Pav. Prod. Fl. Per. 44. t. 64 (1794). *UMBELLIFERAE*, Benth. & Hook. f. i. 878.
Drusa, DC. in Ann. Mus. Par. x. (1807) 468. t. 33.
Elsneria, Walp. Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 346. t. 3.

BOWLESIA :—
HOMALOCARPUS, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 348.
acutangula, *Benth. Pl. Hartw.* 186.—Peruv.
crenata, Desv. ex DC. Prod. iv. 70 = Dysopis Guyana.
dichotoma, *Poepp. ex DC. l. c.* 76.—Chili.
diversifolia, Meyen, ex Walp. Rep. v. 843 = lobata.
elata, *Clos, in C. Gay, Fl. Chil.* iii. 74.—Chili.
elegans, *Clos*, l. c. 74. t. 90.—Chili.
flexilis, *Meyen, Reise*, i. 348.—Chili.
geniculata, Spreng. Umbellif. 14. t. 5 = Angelica geniculata.
geranifolia, *Schlecht. in Linnaea*, i. (1826) 382.—Chili.
incana, *Ruiz & Pav. Fl. Per.* iii. 28.—Peruv.
integerrima, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i. 170.—Chili.
lobata, *Ruiz & Pav. Prod.* 44. t. 34 ; *Fl. Per.* iii. 28. —Peruv.
multiradiata, *Colla, in Mem. Acc. Torino*, xxxvii. (1835) 31.—Chili.
nigripetala, *Clos, in C. Gay, Fl. Chil.* iii. 71.—Chili.
nodiflora, Presl, ex DC. Prod. iv. 75 = tenera.
oppositifolia, *Buch, in Abh. Acad. Berl.* 1816–17. (1819) 370.—Ins. Canar.
palmata, *Ruiz & Pav. Fl. Per.* iii. 28.—Peruv.
pulchella, *Wedd. Chlor. And.* ii. 138.—Peruv. ; Boliv.
rotundifolia, *Phil. in Linnaea*, xxviii. (1856) 651.—Chili.
tenella, *Meyen, Reise*, i. 466.—Peruv.
tenera, *Spreng. Syst.* i. 880.—Bras.
tripartita, *Clos, in C. Gay, Fl. Chil.* iii. 76.—Chili.
tropæolifolia, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 326.—Chili.
uncinata, *Colla, in Mem. Acc. Torino*, xxxvii. (1835) 81. —Chili.

BOWMANIA, Gardn. in Hook. Ic. Pl. 519 (1843) ; et in Hook. Lond. Journ. Bot. ii. (1843) 9 = Trixis, P. Br. (Compos.).
verbascifolia, Gardn. ll. cc. 519 ; 10 = T. Bowmanii.

BOWRINGIA, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 75. *LEGUMINOSÆ*, Benth. & Hook. *f.* i. 552.
callicarpa, *Champ. ex Benth. l. c.*—Hongkong.

BOYKINIA, Nutt. in Journ. Acad. Philad. vii. (1834) 113. *SAXIFRAGEÆ*, Benth. & Hook. f. i. 626.
THEROPON, Raf. in New Fl. Am. iv. 66 (1836).
aconitifolia, *Nutt. l. c.*—Am. bor.
humilis, Raf. in Ann. Bot. 9 ex S. Wats. Bibl. Ind. N. Am. Bot. i. 380 = Ammannia humilis.
major, *A. Gray, in S. Wats. Bot. Calif.* i. 196.—Calif.
occidentalis, *Torr. & Gray, Fl. N. Am.* i. 577.—Am. bor. occ.
ranunculifolia, A. Gray, in Am. Journ. Sc. Ser. I. xlii. (1842) 21 = Saxifraga ranunculifolia.
Richardsoni, *A. Gray, l. c.*—Am. bor. occ.
rotundifolia, *Parry, ex A. Gray, in Proc. Am. Acad.* xiii. (1878) 371.—Calif.
trispora, Nutt. ex Arn. in Hook. Journ. Bot. iii. (1840) 278 = Cayaponia Boykinii.

BOYMIA, A. Juss. in Mém. Mus. Par. xii. (1825) 507 (Mem. Rut. 124) = Zwodia, Forst. (Rutac.).
glabrifolia, Champ. in Hook. Kew Journ. iii. (1851) 330 = E. meliæfolia.
marginicrenata, G. Don, Gen. Syst. i. 805.—Ins. Martinic.
rutæcarpa, A. Juss. in Mém. Mus. Par. xii. (1825) 507 = E. rutæcarpa.

BRABEJARIA, Burm. f. Fl. Cap. Prod. 30, *err. typ.* 26 (1768) (Gen. dub.).
verticillata, Burm. f. l. c.—Afr. austr.

BRABEJUM, Linn. Gen. ed. I. 31 (1737). *PROTEACEÆ*, Benth. & Hook. f. iii. 177.
BRABYLA, Linn. Mant. 187 (1767).
cuticulatum, Blanco, Fl. Filip. ed. II. 39 = Loranthus Blancoanus.
concostenatum, Blanco, l. c. 40 = Alyxia laurina.
lucidum, Blanco, l. c. = Alyxia stellata.
pinnatum, Blanco, l. c. = Alyxia odorata.
stellatifolium, *Linn. Sp. Pl.* 131.—Afr. austr.
stellare, Knight, Prot. 98 = stellatifolium.
stellatum, Thunb. Prod. Fl. Cap. 81 = stellatifolium.
stellulifolium, Murr. Syst. Veg. ed. XIII. 764 = stellatifolium.

BRABYLA, Linn. Mant. 187 (1767) = Brabejum, Linn. (Proteac.).
capensis, Linn. l. c. = Brabejum stellatifolium.

BRACERA, Engelm. in Buchner, Rep. Pharm. lxviii. (1839) 369 = Brayera, Kunth (Rosac.).

BRACHANTHEMUM, DC. Prod. vi. 44 (1837) = Chrysanthemum, Tourn. (Compos.).
fruticulosum, DC. l. c. 45 = C. fruticulosum.

BRACHATERA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = Triodia, R. Br. (Gramin.).
decumbens, Desv. l. c. = T. decumbens.

BRACHEILEMA, R. Br. in Salt, Abyss. App. 65 (1814) = Vernonia, Schreb. (Compos.).
paniculatum, R. Br. l. c. = V. amygdalina.

BRACHIARIA, Griseb. in Ledeb. Fl. Ross. iv. 469 (1853) = Panicum, Linn. (Gramin.).
cruciformis, Griseb. l. c. = P. cruciforme.
prostrata, Griseb. in Goett. Abh. vii. (1857) 263 = P. prostratum.

BRACHILOBOS et BRACHILOBUS, Desv. Journ. Bot. iii. 270 (1814) = seq.

BRACHILOROS, Ait. Fl. Pedem. i. 278 (1785) = Nasturtium, R. Br. (Crucif.).
amphibius, All. l. c. = N. amphibium.
anceps, Link, Handb. ii. 311 = N. anceps.
armoracioides, Schur, Enum. Pl. Transs. 40 = N. armoracioides.
austriacus, Schur, l. c. 41 = N. austriacum.
barbarœfolius, Desv. Journ. Bot. iii. (1814) 170 = N. palustre.
brevistylus, Schur, Enum. Pl. Transs. 39 = N. brevistylon.
dominganensis, Desv. Journ. Bot. iii. (1814) 188 = N. pyrenaicum.
hispidus, Desv. l. c. = N. palustre.
hybridus, Schur, Enum. Pl. Transs. 40 = N. austriacum.
lippicensis, Link, Handb. ii. 311 = N. lippicense.
palustris, [Clairv.] Man. Herb. 218 ; Schur, Enum. Pl. Transs. 39 = N. palustre.
pyrenaicus, All. Fl. Pedem. i. 278 = N. pyrenaicum.
riparius, Schur, Enum. Pl. Transs. 40 = N. amphibium.
sylvestris, All. Fl. Pedem. i. 278 = N. sylvestre.
tanacetifolius, Link, Handb. ii. 311 = N. tanacetifolium.

BRACHILOBUS, Bernh. Syst. Verz. Erf. 181 (1800) = præc.

BRACHIONIDIUM, Lindl. Fol. Orch. (1859). *ORCHIDEÆ*, Benth. & Hook. f. iii. 491.
concolor, *Lindl. l. c.*—Bolivia.
parvifolium, *Lindl. l. c.*—N. Granat.
tuberculatum, *Lindl. l. c.*—N. Granat.

BRACHISTUS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iii. (1849) 264. *SOLANACEÆ*, Benth. & Hook. f. ii. 893.
FREGIRARDIA, Dun. Ind. Sem. Monsp. (1849). Cf. Ann. Sc. Nat. Sér. III. xii. (1849) 366.
SICHLERA, Sendtn. in Flora, xxix. (1846) 194 (*err. typ.* 178), nomen prius.
ciliatus, *Miers, l. c.* 265.—N. Granat.
dimorphus, *Miers, l. c.* 267.—N. Granat.
diversifolius, *Miers, l. c.* 268.—Mexic.
dumetorum, *Miers, l. c.* 265.—N. Granat.
hebephyllus, *Miers, l. c.* 266.—N. Granat.
Hookerianus, *Miers, l. c.* 266.—Ecuador.
lanceseolius, *Miers, l. c.* 267.—Peruv.
Linnæanus, Miers, l. c. 269 = Physalis arborescens.
macrophyllus, *Miers, l. c.* 265.—N. Granat.
ocellis, *Miers, l. c.* 264.—Peruv.
Neesianus, Miers, l. c. 268 = Physalis arborescens.
oblongifolius, *Miers, l. c.* 266.—N. Granat.
rhombodeus, *Miers, l. c.* 264.—N. Granat.
riparius, *Miers, l. c.* 265.—N. Granat.
solanaceus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 424.—Nicaragua.
straminifolius, *Miers, Illustr. S. Am. Pl.* ii. 7.—Mexic.

BRACHTIA, Reichb. f. in Linnæa, xxii. (1849) 858 *ORCHIDEÆ*, Benth. & Hook. f. iii. 561.
ONCODIA, Lindl. Fol. Orch. (1853).
sadina, *Reichb. f. in Bonplandia*, iv. (1856) 322.—N. Granat.
glumacea, *Reichb. f. in Linnæa*, xxii. (1849) 858 — N. Granat.
sulphurea, *Reichb. f. in Bonplandia*, ii. (1854) 14.—N. Granat.

BRACHYACHYRIS, Spreng. Syst. iii. 574 (1826) = BRACHYRIS, Nutt. = Gutierrezia, Lag. (Compos).
Euthamiæ, Spreng. l. c. = G. Euthamiæ.

BRACHYACTIS, Ledeb. Fl. Ross. ii. 495 (1846). *COMPOSITÆ*, Benth. & Hook. f. ii. 279.
ciliata, Ledeb. l. c. = Aster angustus.
frondosa, A. Gray, in Proc. Am. Acad. viii. (1873) 647 = Aster frondosus.
gymnocephala, *Rupr. Sert. Tianschi.* 51.—As. centr.
Ilienais, *Rupr. l. c.*—As. centr.
indsca, C. B. Clarke, Compos. Ind. 49 = menthodora.
menthodora, *Benth. in Hook. Ic. Pl.* t. 1106.—Reg. Himal.
obovata, *Benth. l. c.* sub t. 1106.—Oriens.
pubescens, *Aitch. & Clarke, in Journ. Linn. Soc.* xviii. (1881) 68.—Ind. or.
robusta, *Benth. in Hook. Ic. Pl.* sub t. 1106.—Reg. Himal.
umbrosa, *Benth. l. c.*—Reg. Himal. ; As. centr.
wongtsuensis, Hook. f. Fl. Brit. Ind. iii. 253 = umbrosa.

BRACHYANDRA, Naud. in Ann. Sc. Nat. Sér. III. ii. 148 (1844) = Pterolepis, Miq. (Melastom.).
perpusilla, Naud. l. c. = P. perpusilla.
pusilla, Naud. l. c. xii. (1844) 360 = P. perpusilla.

BRACHYANDRA, Phil. Fl. Atac. 34. t. 4 (1860) *COMPOSITÆ*, Benth. & Hook. f. ii. 244.
macrogyne, *Phil. l. c.*—Chili.

BRACHYANTHES, Cham. ex Dunal, in DC. Prod. xiii. i. 575 (1852) = Fetunia, Juss. (Solanac.).
nierembergensis, Cham. ex Dunal, l. c. = P. parviflora.

BRACHYBOTRYS, Maxim. ex Oliver, in Hook. Ic Pl. xiii. 48. t. 1254 (1878). *BORAGINEÆ*.
paridiformis, *Maxim. ex Oliver, l. c.*—China.

BRACHYCARPÆA, DC. in Mém. Mus. Par. vii (1821) 347 ; et Syst. Veg. ii. 698 (1821). *CRUCIFERÆ*, Benth. & Hook. f. i. 87.
laxa, *Sond. in Harv. & Sond. Fl. Cap.* i. 33.—Afr. austr.
linifolia, *Eckl. & Zeyh. Enum.* 3.—Afr. austr.
polygaloides, *Eckl. & Zeyh. l. c.* 7.—Afr. austr.
variana, *DC. Syst.* ii. 699.—Afr. austr.

BRACHYCENTRUM, Meisn. Gen. 114 (1837) = Centronia, D. Don (Melastom.).
excelsum, Meisn. Gen. Comm. 81 = C. excelsa.

BRACHYCHÆTA, Torr. & Gray, Fl. N. Am. ii. 194 (1841). *COMPOSITÆ*, Benth. & Hook. f. ii. 256.
cordata, *Torr. & Gray, l. c.* 195.—Am. bor.

BRACHYCHEILA, Harv. ex Eckl. & Zeyh. in Linnæa, xx. (1847) 192 = Bucles, Linn. (Ebenac.).
pubescens, Harv. l. c. = E. polyandra.

BRACHYCHITON, Schott & Endl. Meletem. 34 (1832) = Sterculia, Linn.
acerifolium, F. Muell. Fragm. i. 1 = S. acerifolia.
Bidwilli, Hook. Bot. Mag. t. 5138 = S. Bidwilli.
Delabechei, F. Muell. Pl. Vict. i. 157 = S. rupestris.
discolor, F. Muell. Fragm. i. 1 = S. discolor.
diversifolium, R. Br. in Benn. Pl. Jav. Rar. 224 = S. caudata.
Gregorii, F. Muell. in Hook. Kew Journ. ix. (1857) 199 = S. diversifolia.
incanum, R. Br. in Benn. Pl. Jav. Rar. 224 = S. incana.
luridum, C. Moore, ex F. Muell. Fragm. i. 1 = S. lurida.
paradoxum, Schott, Meletem. 34 = S. ramiflora.

BRACHYCHITON :—
platanoides, R. Br. in Benn. Pl. Jav. Rar. 234 = Sterculia trichosiphon.
populneum, R. Br. l. c. = S. diversifolia.
ramiflorum, R. Br. l. c. = S. ramiflora.

BRACHYCLADOS, D. Don, in Phil. Mag. (1832) 391. *COMPOSITAE*, Benth. & Hook. f. ii. 495.
lycioides, *D. Don, l. c.*—Chili.

BRACHYCODON, Prog. in Mart. Fl. Bras. vi. i. 229. t. 42 (1865) = **Pagaea**, Griseb. (Gentian.).
pumilus, Prog. l. c. 230.—Bras.
ramosissimus, Prog. l. c.—Bras.

BRACHYCOME, Cass. in Dict. Sc. Nat. xxxvii. 491 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 364.
BRACHYSCOME, Cass. in Bull. Soc. Philom. (1816) 199.
BRACHYSTEPHIUM, Less. Syn. Comp. 388 (1832).
CTENOSPERMA, F. Muell. ex Pfeiff. Nom. i. 936 (1874).
PAQUERINA, Cass. in Dict. Sc. Nat. xxxvii. 492 (1825).
SILPHIOSPERMA, Steetz, in Lehm. Pl. Preiss. i. 483 (1844).
STERIGLOSSA, DC Prod. vi. 88 (1838).
aculeata, Less. Syn. Comp. 192 = Sieberi.
angustifolia, *A. Cunn. in DC. Prod.* v. (1854) 306.—Austral.
asasmica, *C. B. Clarke, Compos. Ind.* 40.—Assam.
basalticæ, *F. Muell. Fragm.* i. 50.—Austral.
bellidioides, Steetz, in Lehm. Pl. Preiss. i. 426 = pusilla.
Billardieri, *Benth. Fl. Austral.* iii. 518.—Austral.
calocarpa, *F. Muell. in Linnaea*, xxv. (1852) 399.—Austral.
capillacea, Walp. Rep. ii. 584 = Iberidifolia.
cardiocarpa, *F. Muell. ex Benth. Fl. Australi.* iii. 517.—Austral.
cheilocarpa, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 172.—Austral.
chrysoglossa, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 44 = marginata.
ciliaris, *Less. Syn. Comp.* 192.—Austral.
collina, *Benth. Fl. Austral.* iii. 521.—Austral.
debilis, *Sond. in Linnaea*, xxv. (1852) 477.—Austral.
decipiens, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 114.—Austral.
discolor, *C. Stuart, ex Benth. Fl. Australi.* iii. 520.—Austral.
diversifolia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 31.—Austral.
Drummondii, Walp. Rep. ii. 584 = ciliaris.
exilis, *Sond. in Linnaea*, xxv. (1852) 478.—Austral.
glabra, Benth. in Enum. Pl. Hueg. 59 = multifida.
glandulosa, *Benth. Fl. Australi.* iii. 520.—Austral.
glauca, Walp. in Linnæa, xiv. (1840) 815 = stricta.
goniocarpa, *Sond. & F. Muell. in Linnaea*, xxv. (1852) 476.—Austral.
graminea, *F. Muell. Fragm.* i. 49.—Austral.
heterodonta, *DC. Prod.* v. 306.—Austral.
heterophylla, *Benth. in Enum. Pl. Hueg.* i. 50.—Austral.
iberidifolia, *Benth. l. c.* 59.—Australi.
lanuginosa, Steetz, in Lehm. Pl. Preiss. i. 427 = ciliaris.
latisquamea, *F. Muell. Fragm.* xi. 16.—Australi.
leptocarpa, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 43 = exilis.
leucanthemifolia, Benth. in Enum. Pl. Hueg. 60 = stricta.
linearifolia, *DC. Prod.* v. 306.—Austral.
linearifolia, Hook. f. Fl. Tasm. i. 185 = cardiocarpa.
marginata, *Benth. in Enum. Pl. Hueg.* 60.—Austral.
melanocarpa, *F. Muell. & Sond. in Linnaea*, xxv. (1852) 476.—Austral.
mexicana, Klatt, in Leopoldina, xix. (1883) 92 = Bellis mexicana.
microcarpa, *F. Muell. Fragm.* i. 50.—Australasian.
mossambicensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 306.—Afr. trop.
Muelleri, *Sond. in Linnaea*, xxv. (1852) 475.—Austral.
multicaulis, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 43 = ciliaris.
multifida, *DC. Prod.* v. 306.—Austral.

BRACHYCOME :—
nivalis, *F. Muell. in Trans. Phil. Soc. Viet.* i. (1855) 48.—Austral.
obcongifolia, Benth. in Enum. Pl. Hueg. 60 = stricta.
odorata, *Hook. f. Handb. N. Zeal. Fl.* 188.—N. Zel.
pachyptera, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 175.—Austral.
parvula, *Hook. f. Fl. Tasm.* i. 185.—Australi.
pinnata, *Hook. f. Handb. N. Zeal. Fl.* 1881.—N. Zel.
ptychocarpa, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 48.—Austral.
pumila, Walp. Rep. ii. 584 = Billardieri.
pusilla, *Steetz, in Lehm. Pl. Preiss.* i. 427.—Australi.
radicans, *Steetz, l. c.* 429.—Austral.
radicata, *Hook. f. Fl. N. Zel.* i. 127.—N. Zel.
scaptiformis, *DC. Prod.* v. 306.—Austral.
scapigera, *DC. l. c.* viii. 277.—Austral.
segmentosa, *F. Muell. Fragm.* viii. 65; 144.—Ins. Dom. Howe.
Sieberi, *DC. Prod.* v. 306.—Austral.
simplicifolia, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 338.—N. Zel.
Sinclairii, *Hook. f. Handb. N. Zeal. Fl.* 187.—N. Zel.
squalida, Hook. f. in Hook. Lond. Journ. Bot. v. (1846) 116 = ciliaris.
stricta, *DC. Prod.* v. 305.—Austral.
strongyiospermoides, Walp. in Linnæa, xiv. (1840) 805 = ciliaris.
Stuartii, *Benth. Fl. Australi.* iii. 513.—Austral.
sylvestris, Klatt, in Leopoldina, xix. (1883) 92 = Bellis sylvestris †
tenuifolia, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 176 = collina.
tenera, Benth. in Enum. Pl. Hueg. 59 = multifida.
tenuiscapa, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 114 = scaptiformis.
Thomsonii, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 372.—N. Zel.
trachycarpa, *F. Muell. in Linnaea*, xxv. (1852) 389.—Austral.
xanthocomoides, Less. in Linnæa, ix. (1834) 265 = Bellis xanthocomoides.
xeranthemoides, Less. ex Steud. Nom. ed. II. i. 220 = Xanthocephalum Alamani.

BRACHYCOME, Gaudich. Freyc. Voy. Bot. 467 (1826) = **Vittadinia**, Rich. (Compos.).
dentata, Gaudich. l. c. 468 = V. australis †
spathulata, Gaudich. l. c. = V. australis.
triloba, Gaudich. l. c. 467 = V. australis.

BRACHYCORYS, Schrad. Cat. Hort. Goetting. (1830).—Cf. Linnæa, vi. (1831) Litt. 72 = **Lindenbergia**, Linn. (Scrophular.).
parviflora, Schrad. l. c. = L. urticæfolia.

BRACHYCORYTHIS, Lindl. Gen. et Sp. Orch. 262 (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 633.
Baibisella, *Reichb. f. in Flora*, l. (1867) 116 (sphalm. 161).—Afr. austr.
Gerardi, *Reichb. f. l. c.*—Afr. austr.
Kalbreyeri, *Reichb. f. l. c.* ixl. (1878) 77.—Afr. trop.
Mac-Owaniana, *Reichb. f. Otia Bot. Hamb.* 104.—Afr. austr.
pleistophylla, *Reichb. f. l. c.*—Afr. trop.
ovata, *Lindl. Gen. et Sp. Orch.* 263.—Afr. austr.
pubescens, *Harv. Thes. Cap.* i. 35.—Afr. austr.
Pumilio, *Reichb. f. in Flora*, lxv. (1882) 581.—Afr. austr.
Schweinfurthii, *Reichb. f. Otia Bot. Hamb.* 89.—Afr. trop.
tenuior, *Reichb. f. in Flora*, xlviii. (1865) 188.—Afr. trop.
Tysoni, *Bolus, in Journ. Linn. Soc. xx.* (1884) 485.—Afr. austr.
Welwitschii, *Reichb. f. in Flora*, l. (1867) 99.—Afr. trop.
Zeyheri, Reichb. f. l. c. 117 = Schizochilus Zeyheri.

BRACHYDEREA, Cass. in Dict. Sc. Nat. xlviii. 429 (1827) = **Crepis**, Linn. (Compos.).
alpestris, Sch. Bip. ex Nym. Consp. 456 = C. alpestris.
aurea, Sch. Bip. in Follichia, xxii-iv. (1866) 318 = C. aurea.
biennis, Sch. Bip. l. c. = C. biennis.

BRACHYDEREA :—
chondrilloides, Sch. Bip. ex Nym. Consp. 456 = Crepis chondrilloides.
grandiflora, Sch. Bip. ex Nym. l. c. 455 = C. grandiflora.
heterantha, Sch. Bip. in Follichia, xxii-iv. (1866) 318 = Lactuca Dubyana.
Jacquinii, Sch. Bip. l. c. = C. Jacquinii.
jubata, Sch. Bip. l. c. = C. jubata.
nicæensis, Sch. Bip. l. c. = C. nicæensis.
rigida, Cass. in Dict. Sc. Nat. xlviii. 430 = C. rigida.
Rueppellii, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 283 = C. Rueppellii.
Schultzii, Sch. Bip. l. c. = C. Schultzii.
tenerrima, Sch. Bip. l. c. = C. carbonaria.
tenuifolia, Sch. Bip. in Bonplandia, x. (1862) 351 = C. tenuifolia.
xylorrhiza, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 284.—Afr. trop.

BRACHYELYTRUM, Beauv. Agrost. 89. t. 9. f. 2 (1819). *GRAMINEAE*, Benth. & Hook. f. Gen. iii. 1144.
aristatum, Roem. & Schult. Syst. ii. 613 = erectum.
erectum, *Beauv. Agrost.* 89, 155.—Am. bor.

BRACHYGLOTTIS, Forst. Char. Gen. 91. t. 46 (1776). *COMPOSITAE*, Benth. & Hook. f. ii. 444.
Bidwillii, Hook. f. Fl. N. Zel. ii. 342 = Senecio Bidwillii.
Caledoniae, *DC. Prod.* v. 311.—N. Caled.
elaeagnifolia, Hook. f. Fl. N. Zel. i. 41 = Senecio elaeagnifolius.
Forsteri, *DC. Prod.* v. 311.—N. Hebrid.
Forsteri, Hook. f. Fl. N. Zel. ii. 342 = repanda.
perdicioides, Hook. f. l. c. = Senecio perdicioides.
Rani, A. Cunn. in Ann. Nat. Hist. ii. (1839) 192 = Olearia Cunninghami.
Rangiora, *Y. Buch. in Trans. N. Z. Inst.* xiv. (1882) 327.—N. Zel.
repanda, *Forst. Char. Gen.* 92.—N. Zel.
rotundifolia, Forst. l. c. = Senecio rotundifolius.

BRACHYGYNE, Cass. in Dict. Sc. Nat. l. (1827) 498 = **Eriocephalus**, Linn. (Compos.).

BRACHYLAENA, R. Br. in Trans. Linn. Soc. xii. (1816) 115. *COMPOSITAE*, Benth. & Hook. f. ii. 388.
OLIGOCARPHA, Cass. in Bull. Soc. Philom. (1817) 161.
dentata, *DC. Prod.* v. 430.—Afr. austr.
discolor, *DC. l. c.*—Afr. austr.
elliptica, *Less. Syn. Comp.* 208.—Afr. austr.
grandiflora, *DC. Prod.* v. 430.—Afr. austr.
natalensis, Sch. Bip. ex Walp. Rep. ii. 972 = discolor.
nerifolia, *R. Br. in Trans. Linn. Soc. xii.* (1816) 116 —Afr. austr.
racemosa, *Less. Syn. Comp.* 208.—Afr. austr.
trinervia, *Sond. in Linnaea*, xxiii. (1850) 63.—Afr. austr.
uniflora, *Harv. in Harv. & Sond. Fl. Cap.* iii. 117.—Afr. austr.

BRACHYLEPIS, Wight & Arn. Contrib. Bot. Ind. 68 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 742.
nervosa, *Wight & Arn. l. c.*—Ind. or.

BRACHYLEPIS, C. A. Mey. in Ledeb. Fl. Alt. i. 370 (1829) = **Anabasis**, Linn. (Chenoped.).
glatior, C. A. Mey. in Bull. Sc. Acad. Pétersb. viii. (1840) 341 = A. phyllophora.
oriopoda, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 360 = A. phyllophora.
kirghisina, Bunge, in Act. Hort. Petrop. vi. (1879) 444. —Turkestan.
humilis, Less. in Linnæa, ix. (1834) 128 = salsa.
intermedia, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 422 = A. phyllophora.
jaxartica, Bunge, in Act. Hort. Petrop. vi. (1879) 449. —Reg. Casp.
salsa, C. A. Mey. in Ledeb. Fl. Alt. i. 372.—Reg. Casp.
truncata, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. iii. (1844) 183 = A. truncata.

BRACHYLEPIS, Hook. & Arn. in Hook. Journ. Bot. (1834) 290 = **Malmia**, Desne. (Asclep.).
Candolleana, Hook. & Arn. l. c. 291 = M. Candolleana.

BRACHYLOBOS, Desv. Journ. Bot. iii. (1814) 170 = seq.

BRACHYLOBUS, Link, Handb. vii. 311 (1831) = BRACHIOLOBOS, All. = **Nasturtium**, R. Br. (Crucif.).

BRACHYLOBUS, Dulac, Fl. Hautes-Pyr. 279 (1867) = **Melilotus**, Tourn. (Legumin.).
officinalis, Dulac, l. c. = M. officinalis.

BRACHYLOMA, Hanst. in Linnaea, xxvi. (1853) 203 = **Isoloma**, Decne. (1848) (Gesnerac.).
elongatum, Hanst. l. c. xxix. (1857-58) 532.—Ecuador!
erianthum, Hanst. l. c. 530 = I. erianthum.
hirsutum, Hanst. in Linnaea, xxvi. (1853) 208; xxix. (1857-58) 526 = I. hirsutum.
hondense, Hanst. l. c. xxvi. (1853) 208, 213. t. f. 17 = I. hondense.
incurvum, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 105 = I. incurvum.
Karstenianum, Hanst. in Karst. Fl. Colomb. i. 101. t. 50.—N. Granat.
leucomallon, Hanst. in Linnaea, xxix. (1857-58) 528.—N. Granat.
longiflorum, Hanst. l. c. 530 = I. longipes.
longifolium, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 104 = I. longifolium.
longipes, Hanst. in Linnaea, xxix. (1857-58) 530 = I. longipes.
molle, Hanst. l. c. 528 = I. molle.
Moritzianum, Bouché & Hanst. l. c. 524.—N. Granat.
petiolare, Hanst. l. c. 534 = I. petiolare.
pictum, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 104 = I. pictum.
pilosum, Oerst. l. c. 103 = I. pilosum.
rhodomallon, Hanst. in Linnaea, xxix. (1857-58) 526.—Venezuela.
rhynchocarpum, Oerst. in Vidensk. Selsk. Skr. V. v (1861) 103 = I. rhynchocarpum.
rubricaule, Hanst. in Linnaea, xxix. (1857-58) 526 = I. rubricaule.
strictum, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 108 = I. strictum.
Triane, Hanst. in Linnaea, xxix. (1857-58) 532 = I. Trianaei.
tubiflora, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 106 = I. tubiflorum.
ventricosum, Hanst. in Linnaea, xxix. (1857-58) 528.—Hab.?
vestitum, Hanst. l. c. 530 = I. vestitum.

BRACHYLOMA, Sond. in Lehm. Pl. Preiss. i. 304 (1845). *EPACRIDEAE*, Benth. & Hook. f. ii. 618.
ciliatum, *Benth. Fl. Austral.* iv. 178.—Austral.
concolor, *F. Muell. Fragm.* vi. 89.—Austral.
daphnoides, *Benth. Fl. Austral.* iv. 178.—Austral.
depressum, *Benth. l. c.*—Austral.
ericoides, *Sond. in Linnaea, xxvi.* (1853-55) 247.—Austral.
Preissii, *Sond. in Lehm. Pl. Preiss.* i. 305.—Austral.
Scortechinii, *F. Muell. Fragm.* xi. 121.—Austral.

BRACHYMERIS, DC. Prod. vi. 76 (1837) = **Marasmodes**, DC. (Compos.).
scoparia, DC. l. c.—Afr. austr.

BRACHYPHEMA, Benth. in Trans. Linn. Soc. xxii. (1859) 125. t. 22. *EBENACEAE*, Benth. & Hook. f. ii. 666.
ramiflorum, *Benth. l. c.* 126.—Bras.

BRACHYNEMA, Griff. Notul. iv. 176 (1854) = **Sphenodesma**, Jack (Verben.).
ferruginea, Griff. l. c. = S. microstylis.

BRACHYNEMA, F. Muell. Fragm. iii. 90 (1862) = **Abrophyllum**, Hook. f. (Saxifrag.).
ornans, F. Muell. l. c. = A. ornans.

BRACHYOGLOTIS, Lam. Encyc. i. 459 (1783) = **Brachyglottis**, Forst. (Compos.).

BRACHYOLOBOS, DC. Fl. Fr. v. 661 (1805) = BRACHIOLOBOS, All. = **Nasturtium**, R. Br. (Crucif.).

BRACHYOTUM, Triana, in Benth. & Hook. f. Gen. i. 743 (1867). *MELASTOMACEAE*, Benth. & Hook. f. i. c.
Alifanu, Rafin. Sylva Tellur. 101 (1838).
Benthamianum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 49.—Peruv.
campanulare, *Triana, l. c.* 48.—Peruv.
campylanthum, *Triana, l. c.* 49.—Peruv.
caesecens, *Triana, l. c.* 48.—N. Granat.
cernuum, *Triana, l. c.*—N. Granat.
confertum, *Triana, l. c.* 49.—Peruv.
floribundum, *Triana, l. c.*—Peruv.
gracilescens, *Triana, l. c.*—Peruv.
hermanioides, *Triana, l. c.*—Bolivia.
Jamesonii, *Triana, l. c.*—Ecuador.
ledifolium, *Triana, l. c.* 48.—Peruv.
lutescens, *Triana, l. c.*—Peruv.
lycopodioides, *Triana, l. c.* 49.—Peruv.
Microdon, *Triana, l. c.*—Bolivia.
microphyllum, *Triana, l. c.*—Peruv.
Naudini, *Triana, l. c.* 48.—Ecuador.
Pentlandi, *Triana, l. c.* 49.—Bolivia.
quinqueneve, *Triana, l. c.* 48.—Peruv.
Rachela, *Triana, l. c.*—Peruv.
rosmarinifolium, *Triana, l. c.* 49.—Peruv.
rostratum, *Triana, l. c.* 48.—Peruv.
sanguineolutum, *Triana, l. c.* 49.—Bolivia.
strigosum, *Triana, l. c.*—N. Granat.
trichocalyx, *Triana, l. c.* 48.—Ecuador.

BRACHYPAPPUS, Sch. Bip. in Flora, xxxviii. (1855) 119 = Senecio, Tourn. (Compos.).
candicans, Sch. Bip. l. c. 120 = S. candicans.
Smithii, Sch. Bip. l. c. = S. Smithii.

BRACHYPODIUM, Beauv. Agrost. 100 partim (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1201. DISTICHELA, Ehrh. Beitr. vi. 148 (1789).
HEMIBROMUS, Steud. Syn. Pl. Gram. 317 (1854).
TRACHYNIA, Link, Hort. Berol. i. 42 (1827).
acuminatum, *Dum. Obs. Gram. Belg.* 99.—Belgium.
Allionii, Presl, Fl. Sic. p. xlv, in nota = ramosum.
asperum, Roem. & Schult. Syst. ii. 742 = distachyon.
Barrelieri, Roem. & Schult. l. c. 741 = pinnatum.
biflorum, Kunth, Enum. Pl. i. 643 (Ind.) = Agropyron biflorum.
bivascule, Roem & Schult. Syst. ii. 746 = Festuca Pos.
Boissieri, Nym. Syll. 425 = ramosum.
brevistum, Roem. & Schult. Syst. ii. 743 = Triticum brevistum.
caespitosum, Presl, Fl. Sic. p. xlv, nomen = ramosum.
caespitosum, Roem. & Schult. Syst. ii. 737 = pinnatum.
cenisium, Beauv. Agrost. 155, nomen = Poa cenisia.
chinense, *S. L. Moore, in Journ. Bot.* xlii. (1878) 280.—China.
ciliatum, Beauv. Agrost. 101 = distachyum.
commutatum, Beauv. l. c. = Bromus racemosus.
contractum, Presl, Cyp. et Gram. Sic. 40 = pinnatum.
corniculatum, Lam. Fl. Fr. v. 608 = pinnatum.
dalmaticum, *Vis. Fl. Dalm. Suppl.* 23. t. 1.—Europ. austr.
distachyum, *Beauv. Agrost.* 101.—Reg. Mediterr.; Orient.
distachyon, Roem. & Schult. Syst. ii. 741 = prsec.
divaricatum, Presl, Cyp. et Gram. Sic. 40 = Festuca divaricata.
Festuca, *Beauv. Agrost.* 101.—Hab.?
festucoides, Beauv. l. c. 155, nomen = Festuca Poa.
festucoides, Link, Enum. Hort. Berol. i. 95 = Festuca tenella.
flexum, *Nees, Fl. Afr. Gram.* 456.—Afr. austr.
Fontanesianum, Nees, l. c. 457 = ramosum.
fragile, Beauv. Agrost. 156, nomen = Triticum fragile.
geniculatum, C. Koch, in Linnaea, xxi. (1848) 422 = distachyum.
gracile, Roem. & Schult. Syst. ii. 743 = Triticum gracile.
gracile, Beauv. Agrost. 101 = sylvaticum.
Halleri, Beauv. l. c. = Festuca Poa.
hispanicum, Retchb. Fl. Germ. Excurs. 140 = Festuca Poa.
Hostii, Link, Hort. Berol. i. 40 = ramosum.
intermedium, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 89.—Transsylv.
interruptum, Roem. & Schult. Syst. ii. 747 = Apera interrupta.
involutum, *Buese, in Miq. Pl. Jungh.* 347.—Malaya.
japonicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 256.—Japon.

BRACHYPODIUM :—
Kotschyi, *Boiss. Fl. Orient.* v. 659.—Cilicia.
latifolium, *Fourn. Mex. Pl. Enum. Gram.* 125 : *ex Hemsl. Biol. Centr. Am. Bot.* iii. 584.—Mexic.
littorale, Roem. & Schult. Syst. ii. 740 = ramosum.
loliaceum, Fries, Mant. iii. 15 = Glyceria fluitans.
loliaceum, Link, Hort. Berol. i. 42 = Festuca elatior.
loliaceum, Roem. & Schult. Syst. ii. 740 = Demaseria loliacea.
longearistatum, Boiss. Diagn. Sér. I. vii. 127 = Agropyron longearistatum.
longifolium, Beauv. Agrost. 101 = Bromus longifolius
macropodum, *Hack. in Oestr. Bot. Zeitschr.* xxvii (1877) 48.—Europ. austr.
macrostachyum, Bess. ex Roem. & Schult. Syst. Mant. iii. 651 = distachyum.
maritimum, Roem. & Schult. Syst. ii. 743 = Cutandia maritima.
megastachyum, Bess. ex Roem. & Schult. Syst. Mant. iii. 651 = distachyum.
mexicanum, *Link, Hort. Berol.* i. 41.—Mexic.
monostachyum, Nym. Consp. 848 = distachyum.
montanum, Nym. Syll. 426 = Festuca montana.
mucronatum, *Willd. in Willd. & Lange, Prod. Fl. Hisp.* i. 111.—Hispan.
Nardus, Beauv. Agrost. 155, nomen = Festuca tenuiflora.
Nardus, Link, Enum. Hort. Berol. i. 95 = Festuca Poa.
nepalense, *Nees, ex Steud. Syn. Pl. Gram.* 316.—Ind. or.
nigricans, Roem. & Schult. Syst. ii. 743 = Triticum nigricans.
obtusifolium, Link, Enum. Hort. Berol. i. 95 = ramosum.
patens, Nym. Syll. 425 = Festuca loliacea.
pentastachyum, Nym. Consp. 848 = distachyum.
peruvianum, Kunth, Enum. Pl. 643 (Ind.) = Agropyron peruvianum.
phleoides, Beauv. Agrost. 155, nomen = Koeleria phleoides.
phornicodes, Roem. & Schult. Syst. ii. 740 = ramosum.
pinnatum, *Beauv. Agrost.* 101.—Europ.; Reg. Mediterr.
Plukenetii, Beauv. l. c.; Link, Hort. Berol. i. 40 = ramosum.
Poa, Beauv. l. c.; Roem. & Schult. Syst. ii. 746 = Festuca Poa.
psilanthum, Link, Gram. Europ. Austr. Obs. Bot. 298 = Festuca tenuiflora.
ramosum, S. F. Grey, Nat. Arr. Brit. Pl. ii. 113 = Glyceria maritima!
ramosum, *Roem. & Schult. Syst.* ii. 737.—Reg. Mediterr.
retusum, Beauv. Agrost. 101 = ramosum.
rigidulum, *Opiz & Bercht. Oek. Fl. Boehm.* i. 392.—Bohem.
rigidum, Link, Enum. Hort. Berol. i. 96 = Triticum asperum.
Rottboelia, Beauv. Agrost. 155, nomen = ramosum.
rupestre, Roem. & Schult. Syst. ii. 736 = pinnatum.
sanctum, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 161.—Macedon.
scaberrimum, Wight & Arn. ex Steud. Nom. ed. II. i. 231 = Triticum scaberrimum.
Schrenkianum, *Ledeb. Fl. Ross.* iv. 346.—Sibiria.
seisifolium, *Schur, Enum. Pl. Transs.* 800.—Hungar.
strigosum, Schult. in Roem. & Schult. Syst. Mant. ii. 404 = Triticum strigosum.
subtile, De Not. Prosp. Fl. Ligust. 49 = distachyum
subulatum, *Fourn. Mex. Pl. Enum. Gram.* 125 ; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 584.—Mexic.
sylvaticum, *Beauv. Agrost.* 101.—Europ.; Oriens.
tataricum, Munro, ex Aitch. in Journ. Linn. Soc. (1880) 109 = Agropyron longearistatum.
tenellum, Beauv. Agrost. 101 = Festuca tenuiflora.
tenellum, Roem. & Schult. Syst. ii. 745 = Festuca montana!
Tenorianum, Schult. in Roem. & Schult. Syst. Mant. iii. 605 = Triticum Barrelieri.
tenue, Beauv. Agrost. 155, nomen = Lolium perenne.
tenue, Tineo, Cat. Hort. Panorm. (1827) 46 = Festuca incrassata.
tenuiculum, Roem. & Schult. Syst. ii. 744 = Festuca Poa.
tenuiflorum, Roem. & Schult. l. c. 745 = Festuca tenuiflora.
unilateral, Beauv. Agrost. 155; Roem. & Schult. l. c. 747 = Triticum tenuiflorum.
unioloides, Link, Enum. Hort. Berol. i. 195 = Demaseria sicula.

BRACHYPTERUM, Benth. in Ann. Wien. Mus. ii. (1888) 101 = **Derris**, Lour. (Legumin.).
Benthamii, Thw. Enum. Fl. Zeyl. 98 = D. paniculata.
casuarinoe, Dals. & Gibs. Bomb. Fl. 76 = D. casuarinoia.
elegans, Thw. Enum. Fl. Zeyl. 93 = D. parviflora.
floribundum, Miq. Fl. Ind. Bat. i. 139 = D. floribunda.
microphyllum, Miq. Fl. Ind. Bat. Suppl. 296 = D. microphylla.
polyphyllum, Miq. Fl. Ind. Bat. i. 139 = D. polyphylla.
robustum, Dals. & Gibs. Bomb. Fl. 77 = D. robusta.
scandens, Benth. in Ann. Wien. Mus. ii. (1838) 101 = D. scandens.
timorense, Benth. in Miq. Pl. Jungh. i. 255 = D. timorensis.

BRACHYPTERYS, A. Juss. in Deless. Ic. Sel. iii. 20 (1837). *MALPIGHIACEAE*, Benth. & Hook. f. i. 256.
australis, *A. Juss. l. c.*—Bras.
borealis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 356. t. 2. f. 16.—Am. centr.

BRACHYPUS, Ledeb. Fl. Ross. i. 133 (1842) = **Lunaria**, Tourn. (Crucif.).
asper, Ledeb. l. c. (= *Fibigia suffruticosa*).—Rossia.

BRACHYRAMPHUS, DC. Prod. vii. 176 (1838) = **Lactuca**, Tourn. (Compos.).
cavibasus, DC. l. c. 177 = L. intybacea.
goroatensis, DC. l. c. = L. goroeënsis.
Heyneanus, Wight, Ic. t. 1145 = L. Heyneana.
intybaceus, DC. Prod. vii. 177 = L. intybacea.
lactucoides, T. Anders. in Journ. Linn. Soc. Bot. v. Suppl. i. (1860) 28 = Heterachaena massaviensis.
nudicaulis, Klatt, in Ann. Sc. Nat. Sér. V. xviii. (1873) 327 = Launaea nudicaulis.
obtusus, DC. in Deless. Ic. Sel. iv. t. 96 = Launaea nudicaulis.
ramosissimus, Benth. in Hook. Journ. Bot. i. (1834) 459 = L. denticulata.
sinicus, Miq. in Journ. Bot. Néerl. i. (1861) 105 = L. denticulata.
sonchifolius, DC. Prod. vii. 177 = L. remotiflora.
sonchifolius, Thw. Enum. Pl. Zeyl. 168 = L. Heyneana.
taraxacoides, DC. Prod. vii. 177 = L. Heyneana.

BRACHYRHYNCHOS, Less. Syn. Comp. 392 (1833) = **Senecio**, Tourn. (Compos.).
albicaulis, DC. Prod. vi. 438 = S. diversifolius, *Harv. cymbalarifolius*, DC. l. c. = S. cymbalarifolius.
diversifolius, DC. l. c. 439 = S. diversifolius, *Harv.*
elongatus, Less. Syn. Comp. 393 = S. diversifolius, *Harv.*
eupatorioides, DC. Prod. iv. 439 (non seq.).—Afr. austr.
eupatorioides, Less. Syn. Comp. 393 = S. othonnaeflorus.
junceus, Less. l. c. = S. junceus.
brachycarpus, DC. Prod. vi. 438 = S. incertus.
tuberosus, DC. l. c. = S. incertus.

BRACHYRIS, Nutt. Gen. Am. ii. 168 (1818) = **Gutierrezia**, Lag. (Compos.).
californica, DC. Prod. v. 313 = G. californica.
divaricata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 313 = G. Esthamiae.
dracunculoides, DC. in Mém. Soc. Phys. Genève, vii. (1836) 1. 265. t. 1 = Amphiachyris dracunculoides.
Esthamiae, Nutt. Gen. Am. ii. 168 = G. Esthamiae.
floribunda, Phil. in Linnaea, xxxiii. (1864–65) 137.—Chili.
Gayana, Remy, in C. Gay, Fl. Chil. iv. 84.—Chili.
linoni, Phil. in Anal. Univ. Chil. (1865) ii. 835.—Chili.
Maudonii, Sch. Bip. in Linnaea, xxxiv. (1865–66) 534.—Bolivia.
microcephala, DC. Prod. v. 313 = G. Esthamiae.
microcephala, Hook. Ic. Pl. t. 147 = G. texana.
Nuttallii, DC. Prod. v. 313.—Mexic.
ovatifolia, DC. l. c. = Brachychaeta cordata.
paniculata, DC. l. c. = G. paniculata.
ramosissima, Hook. Ic. Pl. t. 143 = Amphiachyris dracunculoides.
scotifolia, Cas. in Atti Soc. Acc. Sc. Nap. v. (1871) vii. 11.—Chili.
spathulata, Phil. in Anal. Univ. Chil. (1865) ii. 836.—Chili.

BRACHYSCOME, Cass. in Bull. Soc. Philom. (1816) 199.—Cf. Dict. Sc. Nat. v. Suppl. 63 (1817) = **Brachyeome**, Cass. (Compos.).
Billardieri, Cass. in Dict. Sc. Nat. v. Suppl. 63 = Brachycome heterodonta.

BRACHYSCYPHA, Baker, in Journ. Linn. Soc. xi. (1871) 393 = **Lachenalia**, Jacq. (Liliac.).
undulata, Baker, l. c. 394 = L. pusilla.

BRACHYSEMA, R. Br. in Ait. Hort. Kew. ed. II. iii. 10 (1811). *LEGUMINOSAE*, Benth. & Hook. f. i. 467.
Subgenus, *Eu-Brachysema*, Benth. & Hook. f. i. 467.
Subgenus, *Eutaxia*, F. Muell. Fragm. i. 222 (1860?).
Subgenus, *Kalencienkia*, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 252.
Subgenus, *Leptosema*, Benth. in Ann. Wien. Mus. ii. (1838) 84.
acuminatum, *Trusf. in Rev. Hortic.* (1866) 413.—Hab. ?
aphyllum, *Hook. Bot. Mag.* t. 4491.—Austral.
bossiaeoides, *Benth. Fl. Austral.* ii. 12.—Austral.
bracteolosum, *F. Muell. Fragm.* iv. 10.—Austral.
Celsianum, Lem. Hortic. Univ. v. n. 2. 58, ex Walp. Rep. v. 428 = lanceolatum.
Chamberlaii, *F. Muell. ex Benth. Fl. Austral.* ii. 13.—Austral.
daviesoides, *Benth. l. c.*—Austral.
lanceolatum, *Meissn. in Lehm. Pl. Preiss.* i. 24.—Austral.
latifolium, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 10.—Austral.
macrocalyx, *Benth. Fl. Austral.* ii. 13.—Austral.
melanopetalum, F. Muell. Fragm. iv. 11 = undulatum.
oxyloboides, *Benth. Fl. Austral.* ii. 12.—Austral.
praemorsum, *Meissn. in Lehm. Pl. Preiss.* i. 25.—Austral.
subcordatum, *Benth. Fl. Austral.* ii. 11.—Austral.
tomentosum, *Benth. l. c.* 13.—Austral.
undulatum, *Ker-Gawl. in Bot. Reg.* t. 542.—Austral.
uniflorum, *R. Br. ex Benth. Fl. Austral.* ii. 12.—Austral.

BRACHYSIPHON, A. Juss. in Ann. Sc. Nat. Sér. III. vi. (1846) 24. t. 2 = **Saroocolla**, Linn. (Penaeac.).
acutus, A. Juss. l. c. 25 = S. acuta.
ericoeformis, A. Juss. l. c. 24.—Afr. austr.
imbricatus, A. Juss. l. c. 25 = S. imbricata.
Mundii, Sond. in Linnaea, xxiii. (1850) 102.—Afr. austr.
rupestris, Sond. l. c. 101 = S. rupestris.
speciosus, Sond. l. c. 108 = S. speciosa.

BRACHYSPATHA, Schott, Syn. Aroid. 35 (1856) = **Amorphophallus**, Blume (Aroid.).
consimilis, Schott, Prod. Aroid. 127 = Hydrosme consimilis.
Konjac, C. Koch, in C. Koch & Fint. Wochenschr. (1858); Gart. Nachr. n. 4. p. 13 = A. Rivieri.
Muelleri, Schott, Syn. Aroid. 36 = A. Mulleri.
variabilis, Schott, l. c. 35 = Synantherias sylvatica.
variabilis, Schott, l. c. = A. variabilis.
zeylanica, Schott, l. c. 36 = A. zeylanicus.

BRACHYSTACHYS, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 47 = Croton, Linn. (Euphorb.).
hirta, Klotzsch, l. c. = C. glanduloeus.

BRACHYSTEGIA, Benth. in Benth. & Hook. f. Gen. i. 583 (1865). *LEGUMINOSAE*, Benth. & Hook. f. i. 468.
appendiculata, *Benth. in Trans. Linn. Soc.* xxv. (1866) 313. t. 42.—Afr. trop.
floribunda, *Benth. in Hook. Ic. Pl.* sub t. 1359.—Afr. trop.
globiflora, *Benth. l. c.*—Afr. trop.
longifolia, *Benth. l. c.* t. 1369.—Afr. trop.
spiceiformis, *Benth. in Trans. Linn. Soc.* xxv. (1866) 312.—Afr. trop.
tamarindoides, *Welw. ex Benth. l. c.*—Afr. trop.

BRACHYSTELMA, R. Br. in Bot. Mag. t. 2543 (1823). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 781.
Subgenus, *Eu-Brachystelma*, Benth. & Hook. f. ii. 781.
Subgenus, *Etaopetalum*, Wight, Centrib. Bot. Ind. 35 (1834).
Subgenus, *Micrositer*, Harv. Gen. Fl. Afr. Austr. ed. II. 342 (1868).
Arnottii, *Baker, in Saund. Ref. Bot.* t. 9.—Afr. austr.

BRACHYSTELMA:—
attenuatum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
Barberiae, *Harv. ex Hook. f. Bot. Mag.* t. 5607.—Afr. austr.
Beddomei, *Hook. f. Fl. Brit. Ind.* iv. 66.—Ind. or.
caudatum, N. E. Br. in Journ. Linn. Soc. xvii. (1880) 169 = crispum.
circinnatum, *S. Mey. Comm. Pl. Afr. Austr.* 196.—Afr. austr.
Comaru, *E. Mey, l. c.* 195.—Afr. austr.
crispum, *R. Grah. in Edinb. Phil. Journ.* (1830) 170.—Afr. austr.
elliptioum, *A. Rich. Tent. Fl. Abyss.* ii. 49.—Abyssin.
filiforme, *Horv. Thes. Cap.* i. 58. t. 93.—Afr. austr.
Gerrardi, *Harv. l. c.* ii. 61. t. 196.—Afr. austr.
glabrum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
hirsutum, *E. Mey. Comm. Pl. Afr. Austr.* 197.—Afr. austr.
laevigatum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
linearo, *A. Rich. Tent. Fl. Abyss.* ii. 49.—Abyssin.
macrorhizum, *E. Mey. Comm. Pl. Afr. Austr.* 197.—Afr. austr.
maculatum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
micranthum, *E. Mey. Comm. Pl. Afr. Austr.* 196.—Afr. austr.
ovatum, *Oliver, in Saund. Ref. Bot.* t. 226.—Afr. austr.
parviflorum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
plocamoides, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 112.—Afr. trop.
spathulatum, *Linell. Bot. Reg.* t. 1113.—Afr. austr.
tuberosum, *R. Br. in Bot. Mag.* t. 2343.—Afr. austr
volubile, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.

BRACHYSTEMMA, D. Don, Prod. Fl. Nep. 216 (1288) = **Arenaria**, Rupp. (Caryophyll.).
calycinum, D. Don, l. c. = A. nepalensis.

BRACHYSTEMUM, Michx. Fl. Bor. Am. ii. 5. t. 81, 82 (1803) = **Pycnanthemum**, Michx. (Labiat.).
lanceolatum, Willd. Enum. Hort. Berol. 622 = P. lanceolatum.
linifolium, Willd. l. c. = P. linifolium.
muticum, Michx. Fl. Bor. Am. ii. 6. t. 82 = P. muticum.
verticillatum, Michx. l. c. 6. t. 81 = P. lanceolatum.
virginicum, Michx. l. c. 6 = P. lanceolatum.

BRACHYSTEPHANUS, Ness, in DC. Prod. xi. 511 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1105.
Lyallii, *Ness, l. c.*—Madag.

BRACHYSTEPHIUM, Less. Syn. Comp. 388 (1832) = **Brachycome**, Cass. (Compos.).
leucanthemoides, Less. l. c. 389 = B. diversifolia.
scapigerum, DC. Prod. vi. 304 = B. scapigera.

BRACHYSTEPIS, Pritz. Ind. Ic. 102 (1855) = **Aeonia**, Lindl. (Orchid.).

BRACHYSTYLIS, E. Mey. ex DC. Prod. vi. 76 (1837) = **Marasmodes**, DC. (Compos.).
scoparia, E. Mey. ex DC. Prod. vi. 76 (= *Brachymeris scoparia*).—Afr. austr.

BRACHYSTYLUS, Dulac, Fl. Hautes-Pyr. 85 (1867) = **Koeleria**, Pers. (Gramin.).
cristatus, Dulac, l. c. = K. cristata.
setaceus, Dulac, l. c. = K. setacea.

BRACHYTOME, Hook. f. Ic. Pl. xi. 70. t. 1086 (1871). *RUBIACEAE*, Benth. & Hook. f. ii. 87.
Wallichii, *Hook. f. ll. cc.*—Reg. Himal.

BRACHYTROPIS, Reichb. Consp. 120 (1828) = **Polygala**, Tourn.
microphylla, Willk. in Oestr. Bot. Zeitschr. xxix. (1879) 286 = P. microphylla.

BRACKENRIDGEA, A. Gray, Bot. U. St. Expl. Exped. i. 461. t. 42 (1854). *OCHNACEAE*, Benth. & Hook. f. i. 218.
australiana, *F. Muell. Fragm.* v. 20.—Austral.
fascicularis, *Villar, in Blanco, Fl. Phil.* ed. III. *Nov. App.* 40.—Ins. Philipp.

BRACKENRIDGEA :—
Hookerl, *A. Gray, in Proc. Am. Acad.* iii. (1857) 51. —Ins. Fiji.
nitida, *A. Gray, Bot. U. St. Expl. Exped.* i. 362.—Ins. Fiji.
zanguebarica, *Oliver, in Hook. Ic. Pl.* t. 1096.—Afr. trop.

BRACONOTIA, Godr. Fl. Lorr. ed. I. iii. 191 (1844) [Ed. ii. (1861) Braconnotia.] = **Agropyron**, J.Gaertn. (Gramin.).
acuta, Godr. l. c. 192 = A. acutum.
elymoides, Godr. l. c. 193 = A. caninum.
glauca, Godr. l. c. 192 = A. glaucum.
juncea, Godr. l. c. = A. junceum.
officinarum, Godr. l. c. = A. repens.
pungens, Godr. l. c. = A. pungens.
rigida, Godr. l. c. = A. glaucum et rigidum.
stricta, Godr. l. c. = A. strictum.

BRACTEARIA, DC. ex Steud. Nom. ed. II. i. 222 (1840) = **Pithouchina**, Aubl. (Melast.).
Kappleri, Steud. ex Cogn. in Mart. Fl. Bras. xiv. iii. 575 = T. sapera.

BRACTEOLARIA, Hochst. in Flora, xxiv. (1841) ii. 658 = **Baphia**, Afzel. (Legum.).
polygalacea, Hook. f. in Hook. Niger Fl. 321 = Baphia polygalacea.
racemosa, Hochst. in Flora, xxiv. ii. (1841) 658 = Baphia racemosa.

BRADBURIA, Torr. & Gray, Fl. N. Am. ii. 250 (1841). *COMPOSITAE*, Benth. & Hook. f. ii. 251.
hirtella, *Torr. & Gray, l. c.*—Texas.

BRADBURIA, Spreng. Syst. iii. 255 (1826) = seq.

BRADBURYA, Rafin. Fl. Ludov. 104 (1817) = **Wistaria**, Nutt. (Legumin.).
erecta, Rafin. l. c. 105 = W. speciosa.
scandens, Rafin. l. c. 104 = W. speciosa.

BRADDLEYA, Vell. Fl. Flum. 98 (1825); ii. t. 140 = **Amphirrhox**, Spreng. (Violar.).
tegalis, Vell. l. c. 94; ii. t. 140 = A. longifolia.

BRADLAEIA, Neck. Elem. i. 169 (1790) = **Stier**, Scop. (Umbellif.).

BRADLEA, Adans. Fam. ii. 324 (1763) = **Aptos**, Tourn. (Legumin.).

BRADLEIA, Banks, ex Gaertn. Fruct. ii. 127. t. 109 (1791) = **Glochidion**, Forst. (Euphorb.).
acuminata, Wall. Cat. sub n. 7855 = G. lanceolarium.
arborescens, Steud. Nom. ed. II. i. 222 = G. arborescens. *Blumei*, Steud. l. c. = G. molle.
coccinea, Wall. Cat. n. 7865 = G. coccineum.
coriacea, Wall. l. c. 7872 = Chastocarpus castanocarpus.
coronata, Wall. l. c. 7857 = G. lævigatum.
disica, Gaertn. ex Vahl, fide Baill. Adansonia, i. (1860) 60 = Flueggia microcarpa.
Finlaysoniana, Wall. Cat. n. 7860 = G. superbum.
glauca, Lablll. Sert. Austr. Caled. 76. t. 77 = G. glaucum.
glaucophylla, Hassk. Cat. Hort. Bog. Alt. 242 = G. glaucum.
Glochidion, Gaertn. Fruct. ii. 128. t. 109 = G. ramiflorum.
hirsuta, Roxb. Hort. Beng. 104; Fl. Ind. iii. 699 = G. hirsutum.
impubes, Roxb. ll. cc. 104, 698 = G. Lambertianum.
Kingeriana, Steud. Nom. ed. II. i. 222 = G. glaucum.
laevigata, Wall. Cat. n. 7858.—Ind. or.
lanceolaria, Roxb. Hort. Beng. 69; Fl. Ind. iii. 697 = G. lanceolarium.
lanceolaria, Wall. Cat. n. 7855 = G. Hohenackeri et sphaerogynum.
littorea, Steud. Nom. ed. II. i. 222 = G. littorale.
lucida, Steud. l. c. = G. lucidum.
lutescens, Steud. l. c. = G. lutescens.
macrocarpa, Steud. l. c. = G. macrocarpum.
macrophylla, Lablll. Sert. Austr. Caled. 77. t. 78 = G. macrophyllum.
mollis, Steud. Nom. ed. II. i. 222 = G. molle.

BRADLEIA :—
mollis, Wall. Cat. n. 7858 = Glochidion hirsutum.
moluccana, Steud. Nom. ed. II. i. 222 = G. moluccanum.
multiloculoris, Spreng. Syst. iii. 19 = G. multiloculare.
Nephroia, Steud. Nom. ed. I. 116 = Cocculus Nephroia.
nitida, Roxb. Hort. Beng. 69; Fl. Ind. iii. 699 = G. zeylanicum.
nitida, Wall. Cat. n. 7867 = G. multiloculare.
obliqua, Spreng. Syst. iii. 19 = G. zeylanicum.
obliqua, Wight, ex Wall. Cat. n. 7947 = Phyllanthus Wightianus.
obtusa, Wall. l. c. 7869 = G. littorale.
ovata, Wall. l. c. 7052 = G. velutinum.
philippensis, Bech.-Ham. in Wall. l. c. 7871 = G. multiloculare.
philippensis, Willd. Sp. Pl. iv. 599 = G. philippinense.
philippica, Cav. Ic. iv. 48. t. 371 = G. philippinense.
pinnata, Roxb. Hort. Beng. 69; Fl. Ind. iii. 700 = G. obscurum.
pubera, Roxb. ll. cc. 104; 698 = G. multiloculare.
rubra, Steud. Nom. ed. II. i. 222 = G. rubrum.
sinensis, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 128 = G. obovatum.
sinica, Gaertn. Fruct. ii. 127 t. 109. f. 1 = G. sinicum.
sinica, Hort. ex Muell. Arg. in DC. Prod. xv. ii. 414 = Phyllanthus nobilis.
Smeringia, Steud. Nom ed. II. i. 222 = G. zeylanicum.
Wightiana, Wall. Cat. n. 7862 = G. ellipticum.
zeylanica, Gaertn. Fruct. ii. 128. t. 109 = G. zeylanicum.
zeylanica, Lablll. Sert. Austr. Caled. 76. t. 76 = G. philippinense.
zeylanica, Wall. Cat. n. 7855 G. = G. lanceolarium.

BRAGANTIA, Lour. Fl. Cochinch. 528 (1790). *ARISTOLOCHIACEAE*, Benth. & Hook. f. iii. 122.
APAMA, Lam. Encyc. i. 91 (1783).
ASIPHONIA, Griff. in Trans. Linn. Soc. xix. (1845) 338. t. 37.
CERAMIUM, Blume, Bidr. 1154 (1826).
CYCLODISCUS, Klotzsch, in Monatsb. Acad. Berl. (1859) 591.
MUNNICKIA, Reichb. Consp. 85 (1828).
STRAKARA, Presl, Epim. Bot. 221 (1850).
TRIMERIZA, Lindl. Bot. Reg. sub t. 1548 (1832).
VANHALLIA, Schult. f. Syst. vii. 18 (1829).
corymbosa, *Griff. in Trans. Linn. Soc.* xix. (1845) 335. —Malaya.
hispida, Thw. ex Dsch. in DC. Prod. xv. i. 480 = Wallichii.
khasyana, Griff. in Trans. Linn. Soc. xix. (1845) 336 = tomentosa.
latifolia, Lindl. Bot. Reg. sub t. 1548 = Wallichii.
melastomaefolia, Dsch. in DC. Prod. xv. i. 429 = corymbosa.
racemosa, *Lour. Fl. Cochinch.* 528.—China.
tomentosa, *Blume, Enum. Pl. Jav.* 82.—Ind. or.; Malaya.
Wallichii, *R. Br. in Wall. Cat.* n. 7415.—Ind. or.

BRAGANTIA, Vand. Fasc. Fl. Nov. (1771) 6 = **Gaspharnia**, Linn. (Amarant.).
Vandelli, Steud. Nom. ed. II. i. 222 = G. officinalis.

BRAHEA, Mart. Hist. Nat. Palm. iii. 248, 319. t. 187, 193 (1840). *PALMAE*, Benth. & Hook. f. iii. 926.
armata, S. Wats. in Proc. Am. Acad. xi. (1876) 146 = Erythea armata.
calcarata, Liebm. ex Linden, Cat. n. 87 (1871), spinosa, seq.!
calcarea, *Liebm. in Mart. Hist. Nat. Palm.* iii. 319.—Mexico.
conduplicata, *Linden, Illustr. Hortic.* xxviii. (1881) 16. —Mexico.
dulcis, J. Cooper, in Smithson. Rep. (1860) 442 = Washingtonia filifera.
dulcis, *Mart. Hist. Nat. Palm.* iii. 244. t. 187, 192.—Peru.
edulis, H. Wendl. ex S. Wats. in Proc. Am. Acad. xi. (1876) 190, 146 = Erythea edulis.
filamentosa, Hort. ex S. Wats. l. c. 147 = Washingtonia filifera.
mexicana, H. Wendl. in Kerch. Palm. 235 = Sabal Adansoni!
Roezli, *Linden, Illustr. Hortic.* xxviii. (1881) 88.—Arizona.
serrulata, *H. Wendl. in Kerch. Palm.* 235.—Georgia.

BRAMI, Adans. Fam. ii. 308 (1763) = **Herpestis**, Gaertn. f. (1805) (Scrophul.).

BRAMIA, Lam. Encyc. i. 459 (1783), nomen prius = **Herpestis**, Gaertn. f. (1805) (Scrophul.).
floribunda, F. Muell. Fragm. ix. 167 = H. floribunda.
indica, Lam. Encyc. i. 459 = H. Monniera.
lanigera, G. Don, Gen. Syst. iv. 546 = H. lanigera.
semierrata, Mart. Amoen. Monac. t. 8 = H. gratioloides.

BRANCIONA, Saliab. Gen. Pl. Fragm. 36 (1866) = **Albuca**, Linn. (Liliac.).
setosa, Saliab. l. c. = A. setosa.

BRANDESIA, Mart. Nov. Gen. et Sp. ii. 25. t. 125 (1826) = **Pelanthera**, R. Br. (Amarant.).
alba, Mart. ex Moq. in DC. Prod. xiii. ii. 311 = Achyranthes alba.
echinocephala, Hook. f. in Trans. Linn. Soc. xx. (1851) 189 = T. echinocephala.
elongata, Mart. Beitr. Amarant. 105. n. 4 = T. elongata.
lanceolata, Benth. Pl. Hartw. 247 = T. lanceolata.
Martiana, Gill. ex Moq. in DC. Prod. xiii. ii. 363 = T. phlioxeroides.
mexicana, Schlecht. in Linnaea, vii. (1832) 392 = T. mexicana.
porrigens, Mart. Beitr. Amarant. 106 = T. porrigens.
puberula, Mart. Nov. Gen. et Sp. ii. 27 = puberula.
pubiflora, Benth. in Bot. Voy. Sulph. 157 = T. pubiflora.
pycnantha, Benth. l. c. = T. pycnantha.
rufa, Mart. Nov. Gen. ii. 26. t. 125 = T. rufa.
serpyllifolia, Mart. Beitr. Amarant. 106 = T. serpyllifolia.
villosa, Mart. Nov. Gen. et Sp. ii. 28. t. 128 = T. Martii.

BRANDISIA, Hook. f. & Thoms. in Journ. Linn. Soc. viii. (1865) 11. t. 4. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 935.
discolor, Hance, in Journ. Bot. xviii. (1880) 299 = Hancei.
discolor, *Hook. f. & Thoms. in Journ. Linn. Soc.* viii. (1865) 11.—Burma.
Hancei, *Hook. f. Fl. Brit. Ind.* iv. 257.—China.

BRANDONIA, Reichb. Consp. 127 (1828) = **Pinguicula**, Linn. (Lentibul.).

BRANDTIA, Kunth, Rev. Gram. ii. 511. t. 170 (1830) = **Arundinella**, Raddi (Gramin.).
holcoides, Kunth, l. c. = A. holcoides.

BRANICA, Endl. Ench. 389 (1841), sphalm. = BRAMIA, Lam. = **Herpestis**, Gaertn. f. (Scrophul.).

BRANICIA, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 476 = **Senecio**, Tourn. (Compos.).
insignis, Andrz. l. c. = S. racemosus.

BRANDESIA, Baill. Adansonia, ix. (1869) 217. filicifolia, *Baill. l. c.*—Ins. Mascar.

BRASENIA, Schreb. Gen. 372 (1789). *NYMPHAEACEAE*, Benth. & Hook. f. i. 46.
HYDROPELTIS, Michx. Fl. Bor. Am. i. 323. t. 29 (1803).
IXODIA, Soland. ex DC. Syst. ii. 37 (1821).
RONDACHINE, Bosc; Encyc. Agric. vi. (1816) 180 (nomen gallicum, tamen citatur ab DC. Syst. et Steud.).
Hydropeltis, Muhl. Cat. 55; Rafin. Med. Fl. i. 90. f. 17 = Schreberi.
nymphoides, Benth. Hist. Pl. iii. 82 = Schreberi.
peltata, Pursh, Fl. Bor. Am. ii. 389 = Schreberi.
purpurea, Casp. in Jorn. Sc. Acad. Lisb. iv. (1873-74) 213 = Schreberi.
Schreberi, *Y. F. Gmel. Syst.* 853.—Ind. or.; Japon; Austral.; Afr. trop.; Am. bor.

BRASILIASTRUM, Lam. Encyc. i. 462 (1783).
americanum, Lam. l. c. = Comocladia Brasiliastrum et Picramnia Antidesma.

BRASILIUM, J. F. Gmel. Syst. 417 (1791).
glabrum, J. F. Gmel. l. c. = Picramnia Antidesma.
hirsutum, J. F. Gmel. l. c. = Comocladia Brasiliastrum.

BRASSAIA, Endl. Nov. Stirp. Dec. i. 89 (1839). *ARALIACEÆ*, Benth. & Hook. f. i. 943.
actinophora, *Endl. l. c.*—Austral.
actinophylla, *F. Muell. Fragm.* iv. 121.—Austral.
capitata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 732. —Reg. Himal.; Burma.
littorea, *Seem. in Journ. Bot.* ii. (1864) 244 — Amboina.
macrostachya, *Seem. l. c.*—N. Guinea.
palmata, Decne. & Planch. in Rev. Hort.ℓ. Sér. IV. iii. (1854) 106 = Trevesia sundaica.
sessilis, *Seem. in Journ. Bot.* ii. (1864) 244.—Sumatra.

BRASSAIOPSIS, Decne. & Planch in Rev. Hortic. Sér. IV. iii. (1854) 106. *ARALIACEÆ*, Benth. & Hook. f. i. 945.
ARALIOPSIS, Kurz, in Andam. Rep. 39 (1870).
aculeata, *Seem. in Journ. Bot.* ii. (1864) 293.—Reg. Himal.
alpina, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 736. —Reg. Himal.
confluens, Seem. in Journ. Bot. ii. (1864) 292 = B. Hainla et Trevesia palmata.
cyrtostyla, *Seem. l. c.* 293.—Sumatra.
floribunda, Seem. l. c. 292 = speciosa.
glomerulata, Regel, Gartenfl. (1863) 275. t. 411 = speciosa.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 736.—Reg. Himal.
Hainla, *Seem. in Journ. Bot.* ii. (1864) 291.—Reg. Himal.
hispida, *Seem. l. c.* 292.—Reg. Himal.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 737.—Reg. Himal.
mitis, *C. B. Clarke, l. c.* 736.—Reg. Himal.
palmata, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 77.—Reg. Himal.; Burma.
ricinifolia, Seem. in Journ. Bot. ii. (1864) 291 = Acanthopanax ricinifolium.
simplicifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 735.—Reg. Himal.
speciosa, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 106.—Reg. Himal.

BRASSAVOLA, Adans. Fam. ii. 127 (1763) = **Helenium**, Linn. (Compos.).

BRASSAVOLA, R. Br. in Ait. Hort. Kew. ed. II. v. 216 (1813). *ORCHIDEÆ*, Benth. & Hook. f. iii. 533.
LYSIMNIA, Rafin. Fl. Tellur. i. 43 (1836).
TULEXIS, Rafin. l. c. 42 (1836).
acaulis, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 152. f. 216.—Am. centr.
amazonica, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 3.— Bras.
angustata, *Lindl. Bot. Reg.* (1838) *Misc.* 41.— Guiana.
appendiculata, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 23 = cucullata.
Cebolletta, *Reichb. f. in Bonplandia*, iii. (1855) 221.— Hab.?
cordata, *Lindl. Bot. Reg.* t. 1913.—Bras.
cucullata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 216.— Ind. occ.
cuspidata, Hook. Bot. Mag. t. 3729 = cucullata.
Digbyana, Lindl. Bot. Reg. (1846) t. 53 = Laelia Digbyana.
elegans, Hook. Bot. Mag. t. 3098 = Tetramicra rigida.
filifolia, *Linden, Illustr. Hortic.* xxviii. (1881) 64.— N. Granat.
flagellaris, *Rodrig. Orch. Nov.* ii. 161.—Bras.
fragrans, Lem. in Jard. Fleur. iii. (1853) Misc. 78 = tuberculata.
fragans, *Rodrig. Orch. Nov.* i. 65.—Bras.
Gibsoniana, Hort. ex Nichols. Dict. Gard. i. 209 = tuberculata.
glauca, Lindl. Bot. Reg. (1839) Misc. 47 = Laelia glauca.
grandiflora, Lindl. l. c. 16 = nodosa.
linearis, *Hook. Bot. Mag.* t. 4784.—Am. centr.
Martiana, *Lindl. Bot. Reg.* (1836) sub t. 1914.— Bras.
Mathieuana, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 290 = linearis.
nodosa, Hook. Bot. Mag. (1833) t. 3229 = subulifolia.

BRASSAVOLA:—
nodosa, *Lindl. Gen. et Sp. Orch.* 114.—Ind. occ. *odoratissima*, Regel, Gartenfl. (1852) 295. t. 33 = cucullata.
Perrinii, *Lindl. Bot. Reg.* t. 1861.—Bras.
Pumilio, *Reichb. f. in Linnaea*, xviii. (1844) 402 — Mexic.
retusa, *Lindl. Bot. Reg.* (1847) sub t. 20.—Venezuela.
revoluta, *Rodrig. Orch. Nov.* ii. 161.—Bras.
rhopalorrhachis, *Reichb. f. in Bot. Zeit.* x. (1852) 855. —Guatemala.
Sloanei, Grisb. Fl. Brit. W. Ind. 620 = subulifolia.
Sloanei, Lindl. ex Heynh. Nom. ii. 76, nomen (= nodosa?).—Jamaica.
suaveolens, Gal. ex Hemsl. in Gard. Chron. (1879) i. 385, 396 = Epidendri sp. ?
subulifolia, *Lindl. Gen. et Sp. Orch.* 115.—Ins. Nevis.
surinamensis, Focke, in Tijdschr. Nat. Wetens. iv. (1851) 37 = angustata.
tuberculata, *Hook. Bot. Mag.* t. 3878.—Bras.
venosa, *Lindl. Bot. Reg.* (1840) *Misc.* 20. t. 39.—N. Granat.
vomeriformis, *Reichb. f. ex Griseb. Fl. Brit. W. Ind.* 621.—Ind. occ.

BRASSAVOLAEA, Poepp. & Endl. Nov. Gen. et Sp. ii. 3 (1838) = praec.

BRASSAVOLEA, Spreng. Syst. iii. 744 (1826) = praec.

BRASSENIA, Heynh. Nom. i. 130 (1840) = **Brassenia**, Schreb. (Nymphaeac.).

BRASSIA, R. Br. in Ait. Hort. Kew. ed. II. v. 215 (1813). *ORCHIDEÆ*, Benth. & Hook. f. iii. 564.
angusta, Lindl. Bot. Reg. (1844) Misc. 5 = Lawrenceana.
antherotes, *Reichb. f. in Gard. Chron.* (1879) ii. 782. —Ecuador.
arachnoides, *Rodrig. Orch. Nov.* i. 97.—Bras.
arcuigera, *Reichb. f. in Gard. Chron.* (1869) 388.— Peruv.
aristata, Lindl. Bot. Reg (1844) Misc. 7 = verrucosa.
bidens, *Lindl. l. c.* 6.—Hab.?
brachiata, *Lindl. in Benth. Pl. Hartw.* 94.—Guatemala.
brachypus, *Reichb. f. in Gard. Chron.* (1875) i. 136.— Ecuador.
candata, *Lindl. Bot. Reg.* t. 832.—Ind. occ.
chloroleuca, *Rodrig. Orch. Nov.* i. 97.—Bras.
cinnabarina, *Linden, ex Lindl. Fol. Orchid. Brassia*, 6.—N. Granat.
cryptophthalma, *Reichb. f. in Gard. Chron.* (1879) ii. 594.—Peruv.
elegantula, *Reichb. f. l. c.* (1885) ii. 616.—Mexic.
encoles, *Reichb. f. l. c.* (1880) i. 680.—N. Granat.
fasinifera, *Linden & Reichb. f. l. c.* (1870) 953.— Ecuador.
Gireoudiana, *Reichb. f. & Warsc. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 273.—Bras.
glumacea, *Lindl. Orch. Linden.* 17.—Venezuela.
guttata, Lindl. in Benth. Pl. Hartw. 94 = maculata.
Helenae, *Reichb. ex Linden, Illustr. Hortic.* xxviii. (1881) 64.—Ecuador.
hexodonta, *Reichb. f. in Bonplandia*, ii. (1854) 98.— Peruv.
imbricata, *Lindl. Fol. Orchid. Brassiæ*, 5.—N. Granat.
Jonstiana, *Reichb. f. in Regel, Gartenfl.* iii. (1854) 209.—Bras.
Kelliana, *Reichb. f. ex Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 114.—N. Granat.
Lanceana, *Lindl. Bot. Reg.* t. 1754.—Guiana.
Lawrenceana, *Lindl.* l. c. (1841) *Misc.* 2. t. 18.— Bras.
longicuspis, *Reichb. f. Otia Bot. Hamb.* 28.— Ecuador.

BRASSIA:—
longilobe, DC. in Mém. Soc. Phys. Genèv. xi. (1846) ii. 427. t. 1 = verrucosa.
macrostachya, Lindl. Sert. Orch. t. 6 = Lanceana.
maculata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 215.— Jamaica.
neglecta, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 322.—Am. centr.
ocanensis, *Lindl. Fol. Orch. Brassia*, 5.—N. Granat.
odontoglossoides, Klotzsch & Karst. in Otto & Dietr. Allg. Gartenz. xv. (1847) 330 = verrucosa.
peruviana, *Poepp. & Endl. Nov. Gen. et Sp.* ii. t. 117. —Peruv.
pumila, *Lindl. Bot. Reg.* (1845) *Misc.* 60 = Lanceana.
signata, *Reichb. f. in Gard. Chron.* (1881) ii. 6. Hab.?
spathacea, *Linden & Reichb. f. in Illustr. Hortic* xxviii. (1881) 64, nomen.—N. Granat.
thyrsodes, *Reichb. f. in Gard. Chron.* (1866) 849.— Peruv.
verrucosa, *Batem. Orch.* t. 23.—Mexic.
villosa, *Lindl. Fol. Orchid. Brassia*, 4.—Peruv.
Wagneri, *Reichb. f. in Bonplandia*, ii. (1854) 14.—N Granat.
Warscewiczii, *Reichb. f. in Bot. Zeit* x. (1852) 698.— Ecuador.
Wrayae, Skinner, ex Hook. Bot. Mag. t. 4008 = maculata.

BRASSICA, [Tourn.] Linn. Syst. ed. I (1735) *CRUCIFERÆ*, Benth. & Hook. f. i. 84.
AGRONIMUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 329.
BRASSICASTRUM, Link, Handb. ii. 318 (1831).
BRASSICELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 330.
BONANNIA, Presl, Fl. Sic. i. 99 (1826).
CORYNELOBUS, R. Roem. in Linnaea, xiv. (1840) 7.
ERUCASTRUM, Presl, Fl. Sic. i. 92 (1826).
GUENTHERA, Andrz. ex Bess. Enum. Pl. Volh. 83 (1822).
HETEROCRAMBE, Coss. & Dur. Ind. Sem. Hort. Burdig. (1867) 25.
HIRSCHFELDIA, Moench, Meth. 264 (1794).
MELANOSINAPIS, Schimp. & Spenn. in Spenn. Fl. Frib. iii. 944 (1829).
MICROPODIUM, Reichb. Consp. 185 (1828).
MUTARDA, Bernh. Syst. Verz. Erf. 184 (1800).
NAPUS, Schimp. & Spenn. in Spenn. Fl. Frib. iii 939 (1829).
RAMPHOSPERMUM, Andrz. ex Reichb. Moesl. Handb. ed. II. i. p. lvii (1827).
RAPA, Tourn. ex Linn. Syst. ed. I (1735).
RHAMPHOSPERMUM, Andrz. ex Bess. Enum. Pl Volh. 83 (1822).
SINAPI, Dulac, Fl. Hautes-Pyr. 194 (1867).
SINAPIDENDRON, Lowe, in Trans. Camb. Phil. Soc. iv. 86 (1831).
SINAPIS, Linn. Syst. ed. I (1735).
SINAPISTRUM, Spach, Hist. Vég. Phan. vi. 343 (1838).
STERANGALIS, Dulac, Fl. Hautes-Pyr. 195 (1867).
adpressa, *Boiss. Voy. Espagne*, ii. 58; *Vis. Fl. Dalm.* iii. 359.—Europ.; Reg. Mediterr.
agrestis, Stend. Nom. ed. I. 116, 117 = Conringia orientalis.
alba, *Boiss. Voy. Espagne*, ii. 39.—Europ.; Orient; Ind. or. (cult.).
alba, Gilib. Fl. Lituan. ii. 63 = Conringia orientalis.
Allionii, Moretti, Ind. Sem. Hort. Taur. (1839) (fide DC. sched.) = oleracea.
alpina, Linn. Mant. 95 = Conringia alpina.
amplexicaulis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 93.—Abyssin.
amplexicaulis, *Yanbs, in Turnites. Fluestet*, vi. (1883) 177.—Europ.
amplexicaulis, *Pomel, Mat. Fl. Alt.* 15.—Algeria.
anisorrhiza, Molina, Sagg. Chile, ed. II. 278 = magellanica.
anticaria, *Reuy & Yanbs, in Turnites. Fluestet*, vi. (1882) 177.—Europ.
apula, *Yanbs, l. c.* 180.—Europ.
arborea, Stend. Nom. ed. I. 116, 117 = campestris.
arvensis, Jord. Diag. i. 181 = B. Cheiranthus.
armoracioides, *Czern. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 611.—Ucrania.
arvensis, Bald. Taur. 146 = campestris.

BRASSICA :—

arvensis, Linn. Mant. 95 = Moricandia arvensis.

asperifolia, Lam. Encyc. i. 746 = campestris.

atlantica, Coss. & Dur. ex Munby, Cat. Pl. Algér. 2 = Gravinae.

Aucheri, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 88 = Raphanus Aucheri.

austriaca, Jacq. Fl. Austr. iii. 45 = Coringia austriaca.

azurea, *Uspenzky*, *in Bull. Soc. Nat. Mosc.* vii. (1834) 276, nomen.—Kossia.

balearica, Loisel. Fl. Gall. ii. 69 = Robertiana.

balearica, *Pers. Syn.* ii. 206.—Ins. Balear.

barbaraefolia, *Ball*, *in Journ. Linn. Soc.* xvi. (1878) 328 :—Marocco.

Barrelieri, *Yanka*, *in Termész. Füzetek*, vi. (1882) 179. —Europ.

Besseriana, *Andrz. ex Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) i. 184.—Rossia; An. temp.

Blancoana, *Boiss. Diagn. Ser.* II. i. 29.—Hispan.

boetica, *Boiss. Voy. Espagne*, ii. 36. t. 9.—Hispan.

Bolzaneri, *Munby*, *in Bull. Soc. Bot. Fr.* ii. (1855) 288. —Afr. bor.

botrytis, Mill. Gard. Dict. ed. VIII. n. 3 = oleracea.

Botteri, *Vis. Fl. Dalm.* iii. 135.—Dalmat.

brachyloma, *Boiss. & Reut. Diagn.* Ser. II. i. 30.— Marocco.

bracteata, *Yanka*, *in Termész. Füzetek*, vi. (1882) 178. —Europ.

bracteolata. *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 33.—Aegypt.

brevipes, Syme, Engl. Bot. ed. III. i. 140 = Diplotaxis muralis, viminea.

Briggsii, Varenne, in Journ. Bot. xix. (1881) 360 = B. Kaja.

bullata, Pasq. Cat. Ort. Bot. Nap. 17 = oleracea.

Bunias, *DC. Syst.* ii. 606.—Orient.

campestris, *Linn. Sp. Pl.* 666.—Orb. Vet. (Cosmop. cult.).

campestris, Oed. in Fl. Dan. x. t. 550 = Turritis glabra.

capitata, Hort. ex Pers. Syn. ii. 206, in syn. = oleracea.

carinata, *A. Braun*, *in Flora*, xxiv. (1841) 267.— Abyss.

catholica, Brot. ex Willk. & Lange, Prod. iii. 867 = Diplotaxis catholica.

caulorapa, Pasq. Cat. Ort. Bot. Nap. 17 = oleracea.

cespitosa, *Pomel*, *Nouv. Mat. Fl. Atl.* 861.—Afr. bor.

cheiranthifolia, DC. Syst. ii. 603 = B. Cheiranthus.

Cheiranthus, *Vill. Prosp.* 40; *Hist. Pl. Dauph.* iii. 832. t. 36.—Europ. occ. ; Afr. bor.

chinensis, Linn. Cent. 19 ; in Amoen. Acad. iv. 280 = campestris.

corynelobа, *Nym. Consp.* 47.—Hispan.

Cossoniana, *Boiss. & Reut. Diagn.* Ser. II. i. 31.— Hispan.

crassifolia, Forsk. Fl. Aegypt. Arab. 118 = Eruca crassifolia.

cretica, Gaus. Cat. Hort. Boccadif. (1821) 11 = rupestris.

cretica, *Lam. Encyc.* i. 747.—Graecia.

cretica, Moris, Stirp. Sard. Elench. i. 3 = insularis.

crispa, *Raſin. Specch.* ii. 69.—Sicilia.

cyrenaica, *Spreng. Syst.* ii. 911.—Cyrenaic.

defiexa, *Boiss. in Ann. Sc. Nat. Sér.* II. xvii. (1842) 87.—Syria ; Persia.

densiflora, Jord. Diag. i. 186 = B. Cheiranthus.

dimorpha, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 306.—Algeria.

dissecta, *Boiss. Voy. Espagne*, ii. 40.—Hispan.

elata, *Ball*, *in Journ. Bot.* xi. (1873) 295.—Marocco.

elongata, *Ehrh. Beitr.* vii. 159.—Europ. or. ; Reg. Caucas. ; Persia.

enifera, *Dum. Fl. Belg. Prod.* 122.—Belgium.

eruca, *Turca. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 211.—Afr. austr.

Eruca, Boiss. Voy. Bot. Espagne, ii. 41 = Eruca longirostris.

Eruca, Linn. Sp. Pl. 667 = Eruca sativa.

Erucastrum, Prsl. Fl. Palat. ii. 340 = B. Cheiranthus.

Erucastrum, *Linn. Sp. Pl.* 667.—Europ.

erucoides, Boiss. Voy. Bot. Espagne, ii. 38 = Diplotaxis erucoides.

erucoides, Roxb. Hort. Beng. [48] ; Fl. Ind. iii. 117 = Eruca sativa.

erysimoides, Pourr. in Mém. Acad. Toul. iii. (1788) 308 = B. Cheiranthus.

erysimoides, *Sieber*, *ex Spreng. Syst.* ii. 912.—Aegypt.

fimbriata, Hort. ex Steud. Nom. ed. II. i. 223 = oleracea.

floridana, *Raſin. Fl. Journ.* 147.—Am. bor.

fostida, *Yanka*, *in Termész. Füzetek*, vi. (1882) 177. —Europ.

foliosa, Willd. Enum. Hort. Berol. 686 = alba.

Forskahli, *Schult. Obs.* 181. n. 1012.—Orient.

fragilis, Sieber, ex Spreng. Syst. ii. 912 = Diplotaxis acris.

frutescens, *Soland. ex DC. Syst.* ii. 624.—Ins. Madera.

fruticulosa, *Cyril. Pl. Rar.* ii. 7. t. 1.—Reg. Mediterr.

geniculata, Ball, in Journ. Linn. Soc. xvi. (1878) 327 = adpressa.

glaberrima, *Pomel*, *Nouv. Mat. Fl. Atl.* 864.— Algeria.

glareosa, Jord. Diagn. i. 183 = B. Cheiranthus.

gongylodes, Mill. Gard. Dict. ed. VIII. n. 8 = oleracea.

gracilis, *Pomel*, *Nouv. Mat. Fl. Atl.* 325.—Algeria.

grandiflora, Balansa, ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 221 = sabularia.

Gravinae, *Tenore*, *Prod. Fl. Nap.* 39. t. 62.—Ital. ; Afr. bor.

Griffithii, Hook. f. & Thoms. in Journ. Linn. Soc v. (1861) 171 = Diplotaxis Griffithii.

Havardi, *Pomel*, *Nouv. Mat. Fl. Atl.* 863.—Algeria.

helenioides, *Burch. ex Loud. Hort. Brit.* 264.—Ins. S. Helen.

heterophylla, Bolss. ex Nym. Syll. 197 = adpressa.

hirta, *Moench*, *Meth. Suppl.* 84.—Hab.?

hispida, *Boiss. Voy. Espagne*, ii. 40.—Hispan. ; Afr. bor.

hispida, Tenore, Prod. Fl. Nap. 39 = Eruca sativa.

humilis, Boiss. Voy. Espagne, 35 = latuliluca.

humilis, *DC. Syst.* ii. 598.—Gallia.

hybrida, *Tenore*, *Prod. Fl. Nap.* 59.—Italia.

incana, Mabl. Icann. Stirp. Austr. 272 = Botteri.

incana, *Tenore*, *Prod. Fl. Nap.* 59.—Italia.

insularis, *Moris*, *Fl. Sard.* i. 168.— Sardinia.

intricata, *Yanka*, *in Termész. Füzetek*, vi. (1882) 179. —Europ.

japonica, Schiedl. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 74 = auriculata ?

juncea, *Coss. in Bull. Soc. Bot. Fr.* vi. (1859) 609.— An. temp. et trop. (cult.).

lacviata, Steud. Nom. ed. I. 116, 117 = oleracea.

laevigata, *Lag. in Varied. Cienc.* ii. iv. (1805) 40.— Hispan.

lanceolata, *Lange*, *in Fl. Dan.* t. 2656.—Dania.

lanceolata, *Sauv. Fl. Cub.* 2.—Cuba

latuliluca, *Boiss. & Reut. Diagn.* Ser. II. i. 30.— Hispan.

latuvalvis, Boiss. Diagn. Ser. I. vi. 72 = Eruca latuvalvis.

leptocarpa, *Boiss. Fl. Orient.* i. 892.—Persia.

leptopetala, *Sond. in Harv. & Sond. Fl. Cap.* i. 81.— Afr. austr.

leucophaea, Boiss. ex Nym. Consp. 45 = nivea.

loncholoma, *Pomel*, *Nouv. Mat. Fl. Atl.* 860.— Algeria.

longirostra, *Boiss. Voy. Espagne*, ii. 40.—Hispan.

lutea, Gilib. Fl. Lituan. ii. 62 = Coringia austriaca.

lyrata, *Desf. Fl. Atlant.* ii. 96. t. 166.—Afr. bor.

macloviana, Urv. in Mém. Soc. Linn. Par. iv. (1826) 616 = Arabis macloviana.

macrocarpa, *Guss. in Presl, Fl. Sic.* i. 92.—Sicilia.

macrorhiza, S. F. Gray, Nat. Arr. Brit. Pl. ii. 682 = campestris.

magellanica, Gaudich. Freyc. Voy. Bot. i. 186 in nota = Arabis macloviana.

magellanica, *Pers. Syn.* ii. 207.—Reg. Magellan.

maritima, Tardent, Ess. Hist. Nat. Bernar. 79 = oleracea.

maurorum, *Durr. in Duch. Rev. Bot.* ii. (1847) 433. —Algeria.

mesopotamica, Bernh. ex Steud. Nom. ed. II. i. 223 = B. Sinapistrum.

mollis, *Vis. Fl. Dalm.* iii. 369.—Europ. austr

monensis, *Huds. Fl. Angl.* ed. II. 291.—Europ. occ.

montana, DC. Fl. Fr. iv. 681 = B. Cheiranthus.

montana, Raſin. Specch. ii. 69 = rupestris.

Moricandia, Boiss. Voy. Espagne, ii. 341 = Moricandia arvensis.

moricandioides, Boiss. Elench. 10 = Moricandia Ramburei.

muralis, Huds. Fl. Angl. ed. II. 291 = Diplotaxis tenuifolia, muralis.

muscosita, Steud. Nom. ed. I. 117 = oleracea.

BRASSICA :—

Napella, Chaix, in Vill. Hist. Pl. Dauph. iii. 834 = campestris.

napobrassica, Mill. Gard. Dict. ed. VIII. n. 2 = campestris.

Napus, Linn. Sp. Pl. 666 = campestris.

nervosa, *Bali*, *in Journ. Bot.* xi. (1873) 299 (sp. dubia). —Marocco.

nigra, *Koch*, *in Roehl. Deutschl. Fl.* ed. III. iv. 718.— Orb. vet. ; (Cosmop. cult.).

nivalis, *Boiss. & Heldr. in Boiss. Diagn.* Ser. II. i 32. —Graecia.

nivea, *Boiss. & Sprun. l. c.* Ser. I. i. 72.—Graecia.

nudicaulis, *Pomel*, *Nouv. Mat. Fl. Atl.* 861.—Algeria.

obtusangula, *Reichb. in Mossl. Handb.* ed. II. ii. 1188. —Europ.

obtusangulata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = B. Erucastrum.

ochroleuca, Soyer-Willem. in Ann. Sc. Nat. ii. (1824) 116 = B. Cheiranthus.

odorata, *Schrank*, *ex Steud. Nom.* ed. I. 117.—Europ.? *oleifera*, Moench, Meth. 253 = campestris.

oleracea, *Linn. Sp. Pl.* 667.—Europ. occ. (Cosmop. cult.).

oleracea, Lour. Fl. Cochinch. 396 = campestris.

oleraceoides, *A. Savat*, *ex Foucaud*, *in Bull. Soc. Bot. Fr.* xxvi. (1879) 74.—Gallia.

orientalis, Linn Sp. Pl. 666 = Coringia orientalis.

ovalis, *Viv. Fl. Libyc. Spec.* 35 t. 16. f. 3.—Libya.

oxyrrhina, *Coss. Notes Crit.* ii 26.—Hispan.

oxyrrhynchа, Cosa. ex Regel, Ind. Sem. Hort. Petrop. (1856) 34 = oxyrrhina.

palustris, Bivona, Fl. Forojulienisis Syllab. (1855) 18 = elongata.

papillaris, Boiss. Elench. 10 = sabularia.

pendula, *Boiss. Voy. Espagne*, ii. 35.—Hispan.

peregrina, Steud. Nom. ed. I. 117 = oleracea.

perfoliata, Crantz, Stirp. Austr. ed. II. 87 ; Lam. Fl. Fr. iii. 487 = Coringia orientalis.

persica, Boiss. & Hohen. Diagn. Ser. I. viii. 26 = elongata.

petraea, Jord. Diagn. i. 185 = B. Cheiranthus.

pinnata, *Spreng. Syst.* ii 911.—Afr. bor.

pinnatifida, *Desf Fl. Atlant.* ii. 95. t. 165.—Hispan. Afr. bor.

Pollichii, Shuttl. in Mag. Zool. Bot. ii. (1838) 581 = B. Cheiranthus.

polymorpha, Murr. in Nov. Comm. Gott. vii. (1777) 35. t. 6 = juncea.

polymorpha, Syme, Engl. Bot. ed. III. i. 138 = campestris.

praecox, Waldst. & Kit. ex DC. Syst. ii. 598 = campestris.

Presili, *Yanka*, *in Termész. Füzetek*, vi. (1882) 182.— Europ.

Prolongi, *Boiss. Voy. Espagne*, ii. 38.—Hispan.

propera, Jord. Diagn. i. 182 = B. Cheiranthus.

pseudoerucastrum, *Brot. Fl. Lusit.* i. 561.—Lusitan.

psammophila, *Pomel*, *Nouv. Mat. Fl. Atl.* 863.— Algeria.

purpurascens, [*Soland. in*] *Russell*, *Aleppo*, ed. II. ii. 258.—Syria.

purpurea, Mill. Gard. Dict. ed. VIII. n. 6 = Moricandia arvensis.

pyrenaea, Jord. Diagn. i. 187 = B. Cheiranthus.

quadrivalvis, *Hook. f. & Thoms. in Journ. Linn. Soc. v.* (1861) 170.—Ind. or. (cult.).

racemifera, Jord. Diagn. i. 184 = B. Cheiranthus.

Repa, Linn. Sp. Pl. 666 = campestris.

rectangularis, *Vis. in Linnaea*, i. (1826) 502.—Corsica.

recurvata, Jord. Diagn. i. 183 = B. Cheiranthus.

repanda, *DC. Syst.* ii. 598.—Gallia.

resyanensis, *Bali*, *in Journ. Bot.* xi. (1873) 299.— Marocco.

Richerii, Lange, in Fl. Dan. xiv. t. 2656 = lanceolata, *Lange*.

Richerii, *Vill. Prosp.* 40.—Gallia.

Robertiana, *J. Gay*, *in Ann. Sc. Nat.* Sér. I. vii. (1826) 416.—Europ.

rotunda, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 500.—Ins. Socotr.

Rouyana, *Yanka*, *in Termész. Füzetek*, vi. (1882) 181. —Europ.

rubra, Hort. ex Steud. Nom. ed. II. i. 224 = oleracea.

rupestris, *Raſin. Caratteri*, n. 191.—Sicil.

rupicola, *Pomel*, *Nouv. Mat. Fl. Atl.* 862.—Algeria.

sabellica, Hort. ex Pers. Syn. ii. 206, in syn. = oleracea.

sabularia, *Brot. Fl. Lusit.* i. 552.—Lusitan. ; Afr. bor.

BRASSICA :—
subularia, Willk. ex Willk. & Lange, Prod. Fl. Hisp. iii. 854 = oxyrrhina.
Sarsuna, Siebold, in Verh. Bat. Gen. xii. (1830) 52 = campestris !
Schimperi, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 86.—Arabia.
scopose, *Panáa, in Termész. Füzetek*, vi. (1882) 179.—Europ.
sempervirens, *Tenore, ex Schrank, Hort. Monac.* t. 10. —Europ. austr.
setigera, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 857.—Hispan.
sinaica, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 85 = Moricandia sinaica.
Sinapistrum, *Boiss. Voy Espagne*, ii. 39.—Europ.
Sinapis, *Vis. Fl. Dalm.* iii. 136.—Europ. (Cosmop. cult.).
stapeliroides, Fisch. ex DC. Prod. i. 217 = Tournefortii.
sphaerorhiza, S. F. Gray, Nat. Arr. Brit. Pl. ii. 688 = campestris.
spinescens, *Pomel, Nouv. Mat. Fl. Atl.* 364.—Algeria.
Stocksii, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 171 = Tournefortii.
stricta, Neail. ex DC. Syst. ii. 598 = campestris.
strigosa, *DC. l. c.* 608.—Afr. austr.
subhastata, Willd. Sp. Pl. iii. 550 = Sisymbrium orientale.
subspinescens, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1839) ii. 145 = Sisymbrium subspinescens.
subularia, Pers. Syn. ii. 207 (sphalm.) = sabularia.
suffruticosa, Desf. Fl. Atlant. ii. 94 = Moricandia arvensis.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 4. = oleracea.
taraxacifolia, Boiss. Elench. 11 = Sisymbrium crassifolium.
tenuifolia, Balll. Hist. Pl. iii. 227 = Diplotaxis tenuifolia.
teretifolia, Desf. Fl. Atlant. ii. 94 = Moricandia teretifolia.
Tigridia, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 87 = deflexa.
timoriana, *F. Muell. Papuan Pl.* ii. 4.—Ins. Timor.
Thai, *Tod. Ind. Sem. Panorm.* (1861) 23.—Hab. ?
torulosa, Dur. in Duch. Rev. Bot. ii. (1847) 434 = Diplotaxis silicalis.
Tournefortii, *Gouan, Illustr.* 44. t. 20. f. A.—Reg. Mediterr. ; Orient ; Ind. or.
trilocularis, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 169.—Reg. Himal.
trinastria, Boiss. Prod. Fl. Monast. 201 = campestris.
tuberosa, Salisb. Prod. 273 = campestris.
turgida, Pers. Syn. ii. 207 = Eruca sativa.
Turritis, Wulf. Obs. 32 = Conringia orientalis.
Uechtritziana, *Panáa, in Termész. Füzetek*, vi. (1882) 182.—Europ.
valentina, *DC. Syst.* ii. 608.—Hispan.
varia, *Dur. in Duch. Rev. Bot.* ii. (1847) 434.—Afr. bor.
vesicaria, Linn. Sp. Pl. 668 = Eruca vesicaria.
villosa, Bivon. ex Spreng. Syst. iv. Cur. Post. 243 = incana.
viminea, Boiss. Voy. Espagne, ii. 38 = Diplotaxis viminea.
violacea, Burm. f. Fl. Ind. 140 = campestris.
violacea, *Linn. Sp. Pl.* 667 (sp. dub.).—China.
virgata, Boiss. Voy. Espagne, ii. 35 = Diplotaxis virgata.
Washitana, Mubl. Cat. 68 = Streptanthes maculatus.
Willdenowii, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 88 = juncea.

BRASSICARIA, Pomel, Mat. Fl. Atl. 15 (1860) = Brassica, Tourn. (Crucifer.).
humilis, Lamotte, Prod. i. 87 = B. humilis.

BRASSICASTRUM, Link, Handb. ii. 318 (1831) = Brassica, Tourn. (Crucifer.).
elongatum, Link, l. c. = B. elongata.

BRASSICELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 230 = Brassica, Tourn. (Crucifer.).
arvensis, Fourr. l. c. = B. Cheiranthus.
Cheiranthus, Fourr. l. c. = B. Cheiranthus.

BRATHYDIUM, Spach, Hist. Vég. Phan. v. 443 (1836); et in Ann. Sc. Nat. Sér. II. v. (1836) 365 = Hypericum, Tourn.
canadense, Spach, l. c. 446 = H. pyramidatum !
Chamaeriinum, Steud. Nom. ed. II. i. 324, sphalm. = seq.
Chamaeriorium, Spach, Hist. Vég. Phan. v. 445 = H. sphaerocarpum.
grandiflorum, Spach, l. c. 443 = H. dolabriforme.
hyssopifolium, Spach, l. c. 445 = H. ciatifolium.
sphaerocarpum, Spach, l. c. 444 = H. sphaerocarpum.

BRATHYS, Mutis, ex Linn. f. Suppl. 49 (1781) = Hypericum, Tourn.
acrosa, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 366 = H. laricifolium.
acicularis, Spach, l. c. = H. aciculare.
Billardieri, Spach, l. c. 367 = H. gramineum.
caespitosa, Blume, Mus. Bot. Lugd. Bat. ii. 20 = H. japonicum.
canadensis, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. canadense.
caracasana, Spach, l. c. 366 = H. caracasanum.
connata, Spach, l. c. = H. connatum.
debilis, Blume, Mus. Bot. Lugd. Bat. ii. 20 = H. japonicum.
denticulata, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. denticulatum.
Drummondii, Spach, l. c. = H. Drummondii.
Erythraea, Spach, l. c. 367 = H. angulosum.
euphorbioides, Spach, l. c. = H. mutilum ?
Forsteri, Spach, l. c. = H. gramineum.
gentianoides, Spach, l. c. = H. Sarothra.
humifusa, Spach, l. c. = H. japonicum.
indecora, Spach, l. c. = H. thesifolium.
japonica, Blume, Mus. Bot. Lugd. Bat. ii. 19 = H. japonicum.
juniperina, Linn. f. Suppl. 268 = H. Brathys.
lanceolata, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. angulosum.
laricifolia, Spach, l. c. 366 = H. laricifolium.
lasa, Blume, Mus. Bot. Lugd. Bat. ii. 19 = H. japonicum.
linoides, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. angulosum.
moranensis, Spach, l. c. = H. moranense.
Mutisiana, Spach, l. c. 366 = H. mexicanum.
nepalensis, Blume, Mus. Bot. Lugd. Bat. ii. 19 = H. japonicum.
oryzatum, Blume, l. c. 20 = H. japonicum.
paniculata, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. paniculatum.
pauciflora, Spach, l. c. = H. pauciflorum.
prolifica, Payer, Organ. 8. t. ? = H. densiflorum.
quinquenervia, Spach, l. c. = ri. mutilum.
radicans, Blume, Mus. Bot. Lugd. Bat. ii. 20 = H. japonicum.
struthiolaefolia, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 366 = H. struthiaefolium.
tarquinensis, Spach, l. c. 367 = H. thesifolium.
thesifolia, Spach, l. c. = H. thesiifolium.
thuyoides, Spach, l. c. 366 = H. thuyoides.
thymifolia, Spach, l. c. = H. Hamboldtianum.
tomentosa, Spach, l. c. 367 = H. setosum.

BRAUNEA, Willd. Sp. Pl. iv. 797 (1805), *ex parte* = Villaresia, Colebr. (Meniosperm.).
menispermoides, Willd. l. c. = T. racemosa.

BRAUNERIA, Neck. Elem. i. 17 (1790) = Echinacea, Moench (Compos.).

BRAUNLOWIA, A. DC. Prod. xvii. 290 (1873) = Brownlowia, Roxb. (Tiliac.).

BRAVAISIA, DC. in Bibl. Univ. xvii. (1838) 124 (Rev. Hign. 16). *ACANTHACEAE*, Benth. & Hook. f. ii. 1084.
CYPHOMACANTHUS, Nees, in DC. Prod. xi. 217 (1847). floribunda, *DC. Prod.* ix. 240.—Mexic.

BRAVOA, Lex. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 6 (1824). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 727.
COSTOCAPNIA, Link & Otto, Ic. Pl. Rar. 38 (1828).
ZETOCAPNIA, Link & Otto, l. c. i. t. 18 (sphalm.) (1828).

BRAVOA :—
Bullana, *Baker, in Gard. Chron.* (1884) ii. 228.—Am. austr.
Costacapsis, M. Roem. Syn. Ensat. 245 = geminiflora.
geminiflora, *Lex. in La Llave & Lex. Nov. Veg. Desc* fasc. i. 6.—Mexic.
sessiliflora, *Hemsl. Diag. Pl. Nov.* 54.—Mexic.

BRAXIPIS, Rafin. Sylva Tellur. 72 (1838) = Sterculia, Linn.
grandiflora, Rafin. l. c. = S. grandiflora.
nitida, Rafin. l. c. 73 = Cola acuminata.

BRAXIREON, Rafin. Fl. Tellur. iv. 23 (1836) = Tapeinanthus, Herb. (1837) (Amaryll.).
humile, Rafin. l. c. = T. humile.

BRAXYLIS, Rafin. Sylva Tellur. 51 (1838) = Ilex, Tourn.
obcordata, Rafin. l. c. = I. obcordata.

BRAYA, Sternb. & Hoppe, in Denkschr. Regensb. i. (1815) 65. *CRUCIFERAE*, Benth. & Hook. f. i. 82
APHRAGMUS, Andrz. ex DC. Prod. i. 209 (1824).
EUDEMA, Humb. & Bonpl. Pl. Aequin. ii. 133. t. 135 (1809), nomen prius.
ORBAS, Cham. & Schlecht. in Linnaea, i. (1826) 29. t. 1.
OROBIUM, Reichb. Consp. 185 (1828).
PLATYPETALUM, R. Br. in Parry, Voy. App. 266 (1824).
aenea, Bunge, Del. Sem. Hort. Dorp. (1841) 8 = alpina.
alpina, *Sternb. & Hoppe, in Denkschr. Regensb.* i. i (1815) 66.—Alp. Europ. ; Reg. arct. ; Tibet.
arctica, Hook. in Parry, Voy. App. 387 = alpina.
calycina, *Wedd. Chlor. And.* t. 85.—Reg. Andin.
dubia, R. Br. ex Hook. f. in Trans. Linn. Soc. xxiii (1861) 319 = alpina.
Eschscholtziana, *Benth. & Hook. f. ex S. Wats. Bibl Ind. N. Am. Bot.* 51.—Am. arct.
falcata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 17 = Sisymbrium falcatum.
glabella, Richards. in Frankl. 1st Journ. 743 = alpina.
imbricatifolia, A. Gray, Bot. U. St. Expl. Exped. i 58 = Sisymbrium imbricatifolium.
involucrata, *Lodd. Fl. Ross.* i. 194.—Sibir. altaic.
Limosella, Bunge, Del. Sem. Hort. Dorp. (1841) 8 = Sisymbrium Limosella.
limosetioides, Bunge, ex Ledeb. Fl. Ross. i. 194 = Sisymbrium Limosella.
Meyeri, Bunge, ex Walp. Rep. ii. 755 (= Parrya sp.). —Sibir. altaic.
nana, *Bunge, Del. Sem. Hort. Dorp.* (1841) 8.—Sibir. baical.
Novae-Zelandiae, *Hook. f. Handb. New Zeal. Fl.* 13.—N. Zel.
oregonensis, *A. Gray, in Proc. Am. Acad.* xviii. (1881-83) 199.—Am. bor. occ.
oxycarpa, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 169 = alpina.
pilosa, *Hook. Fl. Bor. Am.* i. 65. t. 17.—As. et Am. arct.
pinnatifida, Koch, Syn. Fl. Germ. ed. I. 50 = Sisymbrium pinnatifidum.
purpurascens, Bunge, ex Ledeb. Fl. Ross. i. 195 = alpina.
psedla, *A. Gray, Bot. U. St. Expl. Exped.* i. 57.—Chili.
rosea, *Bunge, Del. Sem. Hort. Dorp.* (1841) 8.—As. arct. ; Reg. altaic. et Himal.
Schanginii, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 241.—Turkestan.
silicuosa, *Bunge, Del. Sem. Hort. Dorp.* (1839) 7.—*Cf. Linnaea*, xiv. 118.—Sibir. altaic.
supina, Koch, Syn. Fl. Germ. ed. I. 50 = Sisymbrium supinum.
Thomsoni, Hook. f. in Journ. Linn. Soc. v. (1861) 168 = alpina.
thetica, *Hook. f. & Thoms.* l. c.—Tibet. occ.
Tilingii, *Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc.* xi. (1859) 61.—Kams. bor.
uniflora, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 168.—Tibet. occ.
vernicolor, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 281.—Sibir. baical.

BRAVERA, Kunth, in Bruyer, Not. Vermif. (1824). *ROSACEAE*, Benth. & Hook. f. i. 622.
BAMBEKIA, Bruce, Travels, v. 23, 33 (1790); Ed. II. vii. 161 (1805).
BRACERA, Engelm. in Buchner, Rep. Pharm. lxviii. (1839) 269.
HAGEA, Poir. in Dict. Sc. Nat. xii. 253 (1818).
HAGENIA, J. F. Gmel. Syst. 618 (1791).
anthelmintica, *Kunth, l. c.*—Abyssin.

BRAZORIA, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (Pl. Lindh.) 255 (1845). *LABIATAE*, Benth. & Hook. f. ii. 1204.
Roemeriana, *Scheele, in Linnaea*, xxii. (1849) 590.—Texas.
scutellarioides, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 257.—Texas.
truncata, *Engelm. & Gray, l. c.* 256.—Texas.

BREHISSONIA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 175 = **Fuchsia**, Linn. (Onagrar.).
microphylla, Spach, l. c. = F. microphylla.
thymifolia, Steud. Nom. ed. II. i. 225 = F. thymifolia.

BREIDENBACHIA, Willd. in Ges. Naturf. Freunde Neue Schr. iii. (1801) 412. t. 6. *POLYGALEAE*, Benth. & Hook. f. i. 128.
CATOCOMA, Benth. in Hook. Journ. Bot. iv. (1842) 101.
HIALAINIA, Phil. in Linnaea, xxxiii. (1864–65) 18.
acuminata, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. 111. 51 = myrtifolia.
acuminata, *Poepp. ex Klotzsch, in Peters, Reise Mossamb. Bot.* 117.—Afr. trop.
altissima, *A. W. Benn. in Mart. Fl. Bras.* xiii. 111. 50. —Am. austr.
bracteata, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 187.—Guiana.
brevifolia, *Klotzsch, ex A. W. Benn. in Mart. Fl. Bras.* xiii. 111. 49.—Bras.
cuneata, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 187.—Guiana.
densiflora, *A. W. Benn. in Mart. Fl. Bras.* xiii. 111. 52.—Bras.
floribunda, *Willd. in Ges. Naturf. Freunde Neue Schr.* iii. (1801) 412.—Am. austr.
Hilariana, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. 111. 53 = laurifolia.
Kunthiana, *Klotzsch, ex A. W. Benn. l. c.* 53.—Bras.
laurifolia, *Klotzsch, ex A. W. Benn. l. c.* 52.—Bras.
Lhotskyana, Klotzsch, in Peters, Reise Mossamb. Bot. 117 = floribunda.
lucida, *A. W. Benn. in Mart. Fl. Bras.* xiii. 111. 51.—Bras.
lucida, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 189.—Guiana.
Mansonii, Klotzsch, in Peters, Reise Mossamb. Bot. 117 = lucida.
Martiana, *A. W. Benn. in Mart. Fl. Bras.* xiii. 111. 49. —Bras.
microphylla, *Hieron. in Roca, Exped. Rio-Negro*, ii. 188.—Patagonia.
Morikiana, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 188.—Guiana.
myrtifolia, *A. W. Benn. in Mart. Fl. Bras.* xiii. 111. 50.—Bras.
parviflora, *A. W. Benn. l. c.* 51.—Am. trop.
revoluta, *A. W. Benn. l. c.* 49.—Bras.
Sellowii, *Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 189.—Bras.
velutina, *A. W. Benn. in Mart. Fl. Bras.* xiii. 111. 53. —Bras.

BRENDIA, Blume, Mus. Bot. Lugd. Bat. i. 24. f. 4 (1849). *MELASTOMACEAE*, Benth. & Hook. f. i. 753.
hirsuta, *Blume, l. c.* 25.—Japon.
Oldhami, *Hook. f. ic. Pl.* t. 1068.—Ins. Formosa.

BREZA, Lam. Syn. Comp. 9 (1698) = **Cnicus**, Linn. (Compos.).
Breza, Lam. t. c. 10 = C. arvensis.

BREHMIA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 88 = **Simpelmannia**, Linn. (Logan.).
spinosa, Harv. l. c. 89 = S. spinosa.

BREHMIA, Schrank, in Syll. Ratisb. i. (1824) 85 = **Pavonia**, Cav. (Malvac.).
arborescens, Schrank, l. c. = P. velutina.

BREHNIA, Baker, in Journ. Linn. Soc. xiv. (1875) 561 sphalm. = **Behnia**, Didrichs. (Liliac.).

BREMONTIERA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 93. *LEGUMINOSAE*, Benth. & Hook. f. i. 464.
Ammoxylon, *DC. l. c.*—Ins. Mascar.

BRENNIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 390. *RUBIACEAE*, Benth. & Hook. f. ii. 22. madagascariensis, *A. Rich. l. c.*—Madag.

BREPHOCTON, Rafin. Fl. Tellur. ii. 49 (1836) = **Ecchonia**, DC. (Compos.).
cuneatum, Rafin. l. c. = R. cuneata.

BRETEUILLIA, Buchoz, ex DC. Prod. vi. 508, in syn. (1837) = **Didelta**, Less. (Compos.).

BREVIGLANDIUM, Dulac, Fl. Hautes-Pyr. 423 (1867) = **Hottonia**, Linn. (Primul.).
palustre, Dulac, l. c. = H. palustris.

BREVOORTIA, Wood, in Proc. Acad. Sc. Philad. (1867) 81. *LILIACEAE*, Benth. & Hook. f. iii. 800.
coccinea, S. Wats. in Proc. Am. Acad. xiv. (1879) 289 = Ida-Maia.
Ida-Maia, *Wood, in Proc. Acad. Sc. Philad.* (1867) 82.—Calif.

BREWERIA, R. Br. Prod. 467 (1810). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 876.
BONAMIA, A. Gray, in Proc. Am. Acad. v. (1862) 337.
CALYCOBOLUS, Willd. ex Roem. & Schult. Syst. v. 2 (1819).
CODONANTHUS, G. Don, Gen. Syst. iv. 166 (1856).
DETHARDINGIA, Nees, in Nov. Act. Nat. Cur. xi. (1823) 80.
DUPOUREA, H. B. & K. Nov. Gen. et Sp. iii. 118. t. 214 (1818).
PREVOSTEA, Choisy, in Ann. Sc. Nat. Sér. I. iv. (1825) 107.
REINWARDTIA, Spreng. Syst. i. 597 (1825).
SIDDERA, Hochst. in Flora, xxvii. (1844) Bes. Bell 7. t. 5.
STYLISMA, Rafin. Neogenyt. 2 (1825).
TRICHANTHA, Karst. & Triana, in Linnaea, xxviii. (1856) 487.
WILHELMSIA, Hook. f. ex Planch. in Hook. Ic. Pl. t. 796 (1848).
abscisa, *Choisy, in DC. Prod.* ix. 438.—Madag.
africana, *Benth. & Hook. f. Gen.* iii. 877.—Afr. trop.
aquatica, *A. Gray, Syn. Fl. N. Am.* ii. 1. 217.—Am. bor.
brevifolia, *Benth. Fl. Austral. iv.* 426.—Austral.
Burchellii, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 498.—Bras.
Cholsyana, *Steud. Nom.* ed. II. i. 294.—Ind. or.
cordata, *Blume, Bijdr.* 725.—Ind. or. ; Malaya.
elegans, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 494.—Burma.
evolvuloides, Choisy, l. c. = Choisyana.
evolvuloides, *R. Br. in Salt, Abyss. App.* 65.—Abyss.
fastigiata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 83.—Ins. Socotr.
ferruginea, *Benth. & Hook. f. Gen.* iii. 877.—Bras.
glomerata, *Balf. f. l. c.*—Ins. Socotr.
grandiflora, *A. Gray, in Proc. Am. Acad.* xv. (1880) 51. —Florida.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 522. —Afr. trop.
hispida, *Franch, in Rétocil, Camail*, 43.—Afr. trop.
humistrata, *A. Gray, Syn. Fl. N. Am.* ii. 1. 217.—Am. bor.
Langsdorffii, *Meisen. in Mart. Fl. Bras.* vii. 325.—Bras.
latifolia, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 234.—Afr. bor. ; Arab. ; Ind. bor. occ.
linearis, *R. Br. Prod.* 488.—Austral.
linifolia, Spreng. Syst. i. 614 = Wahlenbergia linifolia.
madagascariensis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 498.—Madag.

BREWERIA:—
malvacea, Klotzsch, in Peters, Reise Mossamb. Bot. 245 = Convolvulus malvaceus.
media, *R. Br. Prod.* 488.—Austral.
Menziesii, *Benth. & Hook. f. Gen.* iii. 877.—Ins Sandvic.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* ii. 400.—Mexic.
minima, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 238.—Calif.
ovalifolia, *A. Gray, Syn. Fl. N. Am.* ii. 1. 217.—Am. bor. occ.
oxycarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 76 = evolvuloides.
pannosa, *R. Br. Prod.* 488.—Austral.
parviflora, Am. ex Steud. Nom. ed. II. i. 224 = Convolvulus parviflorus.
pedunculata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 83.—Ins. Socotr.
Pickeringii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 217.—Am. bor.
rosea, *F. Muell. Fragm.* i. 232.—Austral.
Roxburghii, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 498 = cordata.
scoparia, *Lindl. Fl. Med.* 400.—Hab. ?
secunda, *Benth. in Hook. Niger Fl.* 470 (err. typ. 70). —Afr. trop.
sericea, *Griseb. in Goett. Abh.* xix. (1874) 229.—Reg. Argent.
somaliensis, *Vatke, in Linnaea*, xliii. (1880–82) 522.—Afr. trop.
spectabilis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 498.—Madag.
valerianoides, *Villar, in Blanco, Fl. Philip.* ed. III. *Nov. App.* 143.—Ins. Philipp.
venulosa, *Meisen in Mart. Fl. Bras.* vii. 326.—Bras.
virgata, *Vatke, in Linnaea*, xliii. (1880–82) 522.—Afr. austr.

BREWERINA, A. Gray, in Proc. Am. Acad. viii. (1872) 620 = **Arenaria**, Rupp. (Caryophyll.).
suffruticosa; A. Gray, l. c. = A. congesta.

BREWSTERA, M. Roem. Syn. i. Hesp. 122, 141 (1846) = **Izoganthes**, Jack (Linnac.).
crenata, M. Roem. l. c. 141 = I. locosandra.

BREYNIA, Noronha, ex Thou. Gen. Nov. Madag. 20 (1806). *SAXIFRAGEAE*, Benth. & Hook. f. i. 645.
VENANA, Lam. Illustr. ii. 99. t. 181 (1800), nomen prius.
acanthifolia, Noronha, ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 159 = madagascariensis.
amplifolia, Noronha, l. c. = madagascariensis.
caulifloru, *Tul. l. c.* 161.—Madag.
chrysophylla, Sweet, Hort. Brit. ed. I. 492 = madagascariensis.
digyna, Noronha, ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 159 = madagascariensis.
heterophylla, Boj. Hort. Maurit. 52 = madagascariensis.
Lucasiana, *Lubbers, Cat. Pl. Rer.* 6.—Hab. ?
madagascariensis, *Thou. ex Ker-Gawl. in Bot. Reg.* t. 790.—Madag.
microcarpa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 160.—Ins. Seychell.
ovatifolia, Noronha, ex Tul. l. c. 159 = madagascariensis.
serrata, Presl, Rep. Bot. Syst. i. 190 = madagascariensis.
spinosa, Lindl. in Bot. Reg. t. 878 = madagascariensis.

BREYNIA, Forst. Char. Gen. 145. t. 73 (1776). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 276.
FORESTERIA, Scop. Introd. 98 (1777).
MELANTHESA, Blume, Bijdr. 590 (1825).
MELANTHESIOPSIS, Mosll. Arg. in Linnaea, xxxii. (1863) 74.
acuminata, *Muell. Arr. in DC. Prod.* xv. ii. 442.—Ins. Philipp.
axillaris, Spreng. Pugill. ii. 92 = disticha.
cernua, *Muell. Arg. in DC. Prod.* xv. ii. 439.—Austral. ; Ins. Timor.
cinerascens, Baill. Adansonia, vi. (1866) 344 = oblongifolia.
discigera, *Muell. Arg. in DC. Prod.* xv. ii. 440.—Malaya.
disticha, *Forst. Char. Gen.* 146. t. 73.—N. Caled.

BREYNIA:—
microphylla, *Muell. Arg. in DC. Prod.* xv. 11. 442.—Java.
Muelleriana, Baill. Adansonia, vi. (1866) 344 = stipitata.
oblongifolia, *Muell. Arg. in DC. Prod.* xv. 11. 440.—Austral.
paniculata, Spreng. Pugill. ii. 93 (gen. dub.).—Ind. or.
racemosa, *Muell. Arg. in DC. Prod.* xv. 11. 441.—Java.
rhamnoides, *Muell. Arg. l. c.* 440.—As. trop.
rhynchocarpa, *Benth. Fl. Austral.* vi. 114.—Austral.
rubra, *Muell. Arg. in DC. Prod.* xv. 11. 438.—Java.
stipitata, *Muell. Arg. l. c.* 442.—Austral.
virgata, *Muell. Arg. l. c.* 441.—Java.

BREYNIA, [Plum.] Linn. Gen. ed. I. 118 (1737) = Capparis, Tourn.
indica, Linn. Sp. Pl. 503 = C. Breyniana.

BREZIA, Moq. in DC. Prod. xiii. ii. 167 (1849) = Bumelia, Forsk. (Chenop.).
heterophylla, Moq. l. c. = S. heterophylla.

BRICKELLIA, Ell. Sketch, ii. 290 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 247.
BULIOSTYLIS, DC. Prod. v. 128 (1836).
CLAVIGERA, DC. l. c. 127 (1836).
COLEOSANTHUS, Cass. in Bull. Soc. Philom. (1817) 67.
ISMARIA, Rafin. Sylva Tellur. 117 (1838).
ROSALSIA, La Llav. & Lex. Nov. Veg. Desc. i. 9 (1824).
atractyloides, *A. Gray, in Proc. Am. Acad.* viii. (1870) 290.—Am. bor. occ.
baccharides, *A. Gray, Pl. Wright.* i. 87.—Texas; Arizona.
betonicaefolia, *A. Gray, l. c.* ii. 79.—N. Mexic.; Arizona.
brachyphylla, *A. Gray, l. c.* i. 84.—Am. bor. occ.
californica, *A. Gray, Pl. Fendl.* 84.—Am. bor. occ.
Cavanillesii, *A. Gray, Pl. Wright.* i. 85.—Mexic.
cedrosensis, *Greene, in Bull. Torrey Club*, x. (1883) 86.—Ins. Cedros.
cordifolia, *Ell. Sketch*, ii. 290.—Am. bor.
corymbosa, *A. Gray, Pl. Wright.* i. 84.—Mexic.
Coulteri, *A. Gray, l. c.* 86.—Am. bor. occ.
Cusmingii, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 325 = Chrysopsis oregana.
cylindracea, *A. Gray, ex Engelm. in Proc. Am. Acad.* i. (1846) 46.—Texas.
dentata, *Sch. Bip. in Seem. Bot. Voy. Herald*, 301.—Texas; Arizona.
diffusa, *A. Gray, Pl. Wright.* i. 86.—Am. trop.
Fendleri, *A. Gray, Pl. Fendl.* 83.—N. Mexic.
floribunda, *A. Gray, Pl. Wright.* ii. 73.—Arizona.
frutescens, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 207.—Calif.
Galeottii, *A. Gray, Pl. Wright.* i. 85.—Mexic.
grandiflora, *Nutt. in Trans. Am. Phil. Soc. Ser.* II. vii. (1841) 387.—Am. bor. occ.
Greenei, *A. Gray, in Proc. Am. Acad.* xii. (1877) 88.—Calif.
Hartwegi, *A. Gray, Pl. Wright.* i. 85.—Mexic.
hastata, *Benth. Bot. Voy. Sulph.* 21.—Calif.
hebecarpa, *A. Gray, Pl. Wright.* i. 85.—Mexic.
hymenoclaena, *A. Gray, in Proc. Am. Acad.* xv. (1880) 99.—Mexic.
insana, *A. Gray, l. c.* vii. (1868) 350.—Calif.
lacviusta, A. Gray, Pl. Wright. i. 87 = dentata.
lanata, *A. Gray. l. c.* 84.—Mexic.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 206.—Calif.
linearifolia, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 325 = Bigelovia Douglasii.
linifolia, *Eaton, in Bot. King. Exp.* 187. t. 15. f. 1–6.—Am. bor. occ.
longifolia, *S. Wats. in Am. Natural.* vii. (1873) 301.—Am. bor. occ.
microphylla, *A. Gray, Pl. Wright.* i. 85.—Am. bor. occ.
mohavensis, *A. Gray, Syn. Fl. N. Am.* i. 11. 104.—Calif.
multiflora, *Kellogg, in Proc. Calif. Acad.* vii. (1877) 46.—Calif.
Nevinii, *A. Gray, in Proc. Am. Acad.* xx. (1885) 297.—Calif.

BRICKELLIA:—
oblongifolia, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 388.—Am. bor. occ.
odontophylla, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 206.—Mexic.
oliganthes, *A. Gray, Pl. Wright.* i. 84.—Arizona; Mexic.
Palmeri, *A. Gray, in Proc. Am. Acad.* xv. (1880) 90.—Mexic.
Parryi, *A. Gray, l. c.* 91.—Mexic.
parvula, *A. Gray. Pl. Wright.* i. 84.—Texas, paucidentata, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 326.—Ins. S. Cruc.
pendula, *A. Gray, Pl. Wright.* i. 85.—N. Mexic.
pinifolia, *A. Gray, l. c.* 84.—Bras.
Pringlei, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 206.—Arizona.
ramiformis, A. Gray, Pl. Wright. i. 86 = Wrightii.
Riddellii, *A. Gray, l. c.* 83.—Texas.
Rosalesia, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am.* Bot. ii. 106.—Mexic.
Rusbyi, *A. Gray, Syn. Fl. N. Am.* i. 11. 106.—N. Mexic.
scoparia, *A. Gray, Pl. Wright.* i. 84.—Mexic.
secundiflora, *A. Gray, l. c.* 85.—Mexic.
Seemanni, *A. Gray, in Proc. Am. Acad.* xv. (1880) 90.—N. Mexic.
simplex, *A. Gray, Pl. Wright.* ii. 73.—Arizona.
spinulosa, *A. Gray, l. c.* i. 84.—Mexic.
squamulosa, *A. Gray, in Proc. Am. Acad.* xv. (1880) 90.—Mexic.
texana, A. Gray, Fl. Wright. ii. 72 = Wrightii.
thyrsiflora, *A. Gray, in Proc. Am. Acad.* xv. (1880) 90.—Mexic.
tomentosa, *A. Gray, Pl. Wright.* i. 85.—Mexic.
veronicaefolia, *A. Gray, l. c.*—Mexic.
Wialisani, *A. Gray, Pl. Fendl.* 84.—Arizona; Mexic.
Wrightii, Durand & Hilg. in Pacif. Rail. Rep. v. 8 = californica.
Wrightii, *A. Gray, Pl. Wright.* ii. 72.—Am. bor. occ.

BRICKELLIA, Rafin. in Med. Repos. New York, v. (1808) 358 = Gilia, Ruiz & Pav. (Polemon.).

BRICOUR, Adans. Fam. ii. 423 (1763) = **Myagrum**, Tourn. (Crucifer.).

BRIDELIA, Willd. Sp. Pl. iv. 978 (1805) (Briedelia). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 267.
CANDELARIA, Hochst. in Flora, xxvii. (1843) 79.
CLUYTIA, Roxb. ex Steud. Nomn. ed. II. i. 225 (1840).
PENTAMERIA, Klotzsch, ex Baill. Etud. Gén. Euph. 584 (1858).
ZARCOA, Lisnoe, in Bot. Zeit. xv. (1857) 423.
acuminata, Wall. Cat. n. 7885 = Phyllanthus bicolor.
alnifolia, *Griff. Notul.* iv. 481.—Burma.
amorna, Wall. ex Baill. Etud. Gén. Euph. 584 = retusa.
angolensis, *Muell. Arg. in Seem. Journ. Bot.* ii. (1864) 327.—Afr. trop.
atroviridis, *Muell. Arg. l. c.*—Afr. trop.
attenuata, *Wall. ex Voigt, Hort. Suburb. Calc.* 156.—Ind.
Bernieriana, *Baill. Adansonia*, ii. (1861–62) 39.—Madag.
Berryana, Wall. Cat. n. 7876, 7960 = Phyllanthus indicus.
buxifolia, *Baill. Adansonia*, ii. (1861–62) 230.—N. Caled.
calsacrifica, *Bertol. f. in Mem. Acc. Sc. Bolog.* v. (1854) 476.—Afr. trop.
chartacea, Wall. Cat. n. 7881 = Cleistanthus chartaceus.
coccolobaefolia, *Kurz, in Journ. Linn. Soc. xxi.* (1885) 441.—Madag.
collina, Hook. & Arn. Bot. Beech. Voy. 211 = Cleistanthus collinus.
crenulata, Roxb. Hort. Beng. 70; Fl. Ind. iii. 784 = retusa.
dasyalyx, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 241.—Burma.
divaricata, *Hook. & Arn. Bot. Beech. Voy.* 211.—Malaya.
elegans, *Muell. Arg. in Seem. Journ. Bot.* ii. (1864) 327.—Afr. occ.

BRIDELIA:—
exaltata, *F. Muell. Fragm.* iii. 32.—Austral.
fagines, *F. Muell. ex Benth. Fl. Austral.* vi. 120.—Austral.
ferruginea, *Benth. in Hook. Niger Fl.* 511.—Afr. trop.
gambecola, *Baill. Adansonia*, i. (1860–61) 79.—Afr. trop.
glauca, *Blume, Bijdr.* 597.—Java.
glauca, Wall. Cat. n. 7875 = prace. ? (Glochidionia sp.).—Penang.
Hamiltoniana, *Wall. l. c.* 7882.—Ind. or.
heterantha, Wall. l. c. 7878 = Glochidion Wallichianum.
horrida, Dillw. Review Hort. Malab. 18 = Scleropyron Wallichianum.
insulana, *Hance, in Journ. Bot.* xv. (1877) 337.—China.
lanceafolia, Roxb. Hort. Beng. 70; Fl. Ind. iii. 787 = tomentosa.
lanceolata, *Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 45.—Malaya.
lancifolia, Buch.-Ham. in Wall. Cat. n. 7884 = tomentosa.
laurina, *Baill. Adansonia*, ii. (1861–62) 229.—N. Caled.
Leichardtii, Baill. Etud. Gén. Euph. 584 = fagines.
Lourreiri, Hook. & Arn. Bot. Beech. Voy. 211 = tomentosa.
melanthesoides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 103.—Afr. trop.
micrantha, *Baill. Adansonia*, iii. (1862–63) 164.—Afr. trop.
montana, Wall. Cat. n. 7879 = retusa.
montana, *Willd. Sp. Pl. iv.* 978.—Ind. or.
Moonii, *Thw. Enum. Pl. Zeyl.* 279.—Zeylan.
multiflora, *Zipp. in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 119.—Amboina.
oblongifolia, Hook. & Arn. Bot. Beech. Voy. 212 = Cleistanthus chartaceus.
ovata, *Decne. in Nouv. Ann. Mus. Par.* iii. (1835) 484.—Austral.; Burma; Malaya; Ins. Timor.
patula, Hook. & Arn. Bot. Beech. Voy. 213 = Cleistanthus patulus.
Perrilleana, *Baill. Adansonia*, ii. (1861–62) 38.—Madag.
polystachya, Baill. Etud. Gén. Euph. 584 = Cleistanthus polystachyus.
pubescens, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 241.—Reg. Himal.; Burma.
retusa, Baill. Etud. Gén. Euph. 584, nomen = Moonii.
retusa, A. Juss. Tent. Euphorb. 27. t. 7 = stipularis.
retusa, *Spreng. Syst.* iii. 48.—Ind. or.; Malaya.
rhamnoides, Griff. Notul. iv. 480 = tomentosa.
rhomboidalis, *Baill. Adansonia*, ii. (1861–62) 37.—Ins. Borbon.
rugosa, *Miq. Fl. Ind. Bat. Suppl.* 445.—Sumatra.
scandens, Willd. Sp. Pl. iv. 979 = stipularis.
sclerocuera, *Muell. Arg. in Flora*, xlvii. (1864) 515.—Afr. trop.
sinica, J. Grah. Cat. Bomb. Pl. 179 = Cleistanthus malabaricus?
speciosa, *Muell. Arg. in Seem. Journ. Bot.* ii. (1864) 327.—Afr. trop.
spinosa, Hort. ex DC. in Mém. Soc. Phys. Genèv. vi. (1833) 563 = Damascanthus indicus.
spinosa, Willd. Sp. Pl. iv. 979 = retusa.
stenocarpa, *Muell. Arg. in Flora*, xlvii. (1864) 515.—Afr. trop.
stipitata, *Baill. Adansonia*, ii. (1861–62) 229.—N. Caled.
stipularis, *Blume, Bijdr.* 597.—Ind. or.; Malaya; Afr. trop.
stipularis, Hook. & Arn. Bot. Beech. Voy. 211 = Cleistanthus stipularis.
tenuifolia, *Muell. Arg. in Journ. Bot.* ii. (1864) 328.—Afr. trop.
tomentosa, *Blume, Bijdr.* 597.—Ind. or.; Malaya; Austral.
Tulasneana, *Baill. Adansonia*, ii. (1861–62) 40.—Madag.
urticoides, Griff. Notul. iv. 481 = tomentosa?
Zollingeri, Miq. Fl. Ind. Bat. i. 11. 564 = stipularis.

BRIDGESIA, Bert. ex Cambess. in Nouv. Ann. Mus. Par. iii. (1834) 324. t. 13. *SAPINDACEAE*, Benth. & Hook. f. i. 398.
TRITTOSOCARPUS, Mchm. Gen. 52 (1837).
insularis, *Bert. ex Cambess. l. c.* 325.—Chili.

BRIDGESIA, Hook. Bot. Misc. ii. (1831) 222. t. 92 = **Polyachyrus**, Lag. (Compos.).
echinopoides, Hook. l. c. = P. sphaerocephalus.

BRIDGESIA, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 168. t. 102 = **Brellia**, A. Juss. (Phytolacc.).
spicata, Hook. & Arn. l. c. 169 = E. volubilis.

BRIEDELIA, Willd. Sp. Pl. iv. 978 (1805) = **Bridelia**, ‘Willd.’ (Euphorb.).

BRIGHAMIA, A. Gray, in Proc. Am. Acad. vii. (1867) 185. *CAMPANULACEAE*, Benth. & Hook. f. ii. 545.
insignis, *A. Gray, l. c.* —Ins. Hawaii.

BRIGNOLIA, Bertol. in Desv. Journ. Bot. ii. (iv.) (1815) 76 = **Kundmannia**, Scop. (Umbellif.).
panacifolia, Heynh. Nom. i. 131, sphalm. = sco.
pastinacifolia, Bertol. in Desv. Journ. ii. (iv.) (1815) 76 = K. sicula.

BRIGNOLIA, DC. Prod. iv. 444 (1830) = **Isertia**, Schreb. (Rubiac.).
acuminata, DC. l. c.—Ins. Trinit.
pubigera, Benth. in Hook. Journ. Bot. iii. (1840) 219. —Guiana.

BRILLANTAISIA, Beauv. Fl. Owar. ii. 67. t. 100 (1805). *ACANTHACEAE*, Benth. & Hook. f. ii. 1078.
ELLANTHERIA, Nees, in DC. Prod. xi. 96 (1847).
LUCORRAPHIS, Nees, l. c. 97 (1847).
alata, *T. Anders. ex Oliver, in Trans. Linn. Soc.* xxix. (1875) 125.—Afr. trop.
hirsuta, *T. Anders. in Journ. Agri-Hort. Soc. Ind.* N. S. i. (1858) 367.—Afr. trop.
Lamium, *Benth. in Hook. Niger Fl.* 477.—Afr. trop.
owariensis, *Beauv. Fl. Owar.* ii. 68. t. 100. f. 2.—Afr. trop.
patula, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 21. —Afr. trop.
pubescens, *T. Anders. ex Oliver, in Trans. Linn. Soc.* xxix. (1875) 125.—Afr. trop.
Rutenbergiana, *Vatke, in Bremen Abh.* ix. (1885) 181 —Afr. trop.
Vogeliana, *Benth. in Hook. Niger Fl.* 477.—Afr. trop.

BRIMEURA, Salisb. Gen. Pl. Fragm. 26 (1866) = **Hyacinthus**, Tourn. (Liliac.).
amethystina, Salisb. l. c. = H. amethystinus.

BRIMYS, Scop. Introd. 223 (1777) err. typ. = **Drimys**, Forst. (Magnol.).

BRINDONIA, Thou. in Dict. Sc. Nat. v. 389 (1804) = **Garcinia**, Linn. (Gutti.).
celsica, Thou. l. c. 544 = G. cornea !
cochinchinensis, Thou. l. c. = G. cochinchinensis.
indica, Thou. l. c. 540 = G. indica.

BRISEGNOA, Remy, in C. Gay, Fl. Chil. v. 291 (1849) = **Oxythece**, Nutt. (Polygon.).
chilensis, Remy, l. c. 292 = O. dendroides.

BRISEIS, Salisb. Gen. Pl. Fragm. 92 (1866) = **Allium**, Tourn. (Liliac.).
triquetra, Salisb. l. c. 93 = A. triquetrum.

BRISSONIA, Neck. Elem. iii. 86 (1790); Desv. [Journ. i. (1814) 78]; in Ann. Sc. Nat. Sér. I. ix. (1826) 411 = **Indigofera**, Linn. et **Tephrosia**, Pers. (Legumin.).
coronillaefolia, Desv. Journ. l. (1814) 78 = T. coronillaefolia.
stipularis, Desv. l. c. = T. stipularis.
trapezicarpa, Desv. l. c. = sco. !
trapezicarpa, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 411 = I. senegalensis.

BRITOA, Berg, in Linnaea, xxvii. (1854) 485 = **Campomanesia**, Ruiz & Pav. (Myrtac.).
acida, Berg, l. c. 486 ; et in Mart. Fl. Bras. xiv. 1. 461. —Bras.
dichotoma, Berg, ll. cc. 486, 462.—Bras.
eriantha, Berg, ll. cc. = C. eriantha.
pedicellata, Berg, ll. cc. 486, 463.—Bras.
rugosa, Berg, in Mart. Fl. Bras. xiv. 1. 612.—Bras.
Sellowiana, Berg, in Linnaea, xxvii. (1854) 487 ; et in Mart. Fl. Bras. xiv. 1. 468.—Bras.
sessiliflora, Berg, ll. cc. 486, 461.—Bras.
trifiora, Berg, ll. cc. 486, 462 = C. trifiora.

BRIZA, Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1194.
CALOTHECA et CHAECOLYTRUM, Desv. in Nov. Bull. Soc. Philom. ii. (1810) 190.
POA, Adans. Fam. ii. 34 (1763).
TREMULARIA, Heist. Syst. 12 (1748).
aequa, Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 44 = media.
aspera, Knapp, Gram. Brit. t. 61 = minor.
australis, Trev. ex Steud. Syn. Pl. Gram. 288 = triloba.
barbata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 366.—Bras.
Berteroniana, *Steud. Syn. Pl. Gram.* 288.—Chili.
bipinnata, Linn. Syst. Nat. x. 875 = Eragrostis cynosuroides.
canadensis, *Michx. Fl. Bor. Am.* i. 71.—Am. bor.
canadensis, Nutt. Gen. Am. i. 69 pro parte = Glyceria nervata.
capensis, Schrank, Hort. Monac. t. 43 = maxima.
capensis, Thunb. Prod. Fl. Cap. 21 = Eragrostis brizoides.
caroliniana, Lam. Encyc. i. 465 ; Illustr. i. 187. t. 45. f. 3 = Uniola paniculata.
caroliniana, Walt. Fl. Carol. 79 = Eragrostis major.
Clusii, Schult. in Roem. & Schult. Mant. ii. 294 = media.
cynosuroides, Scop. Delic. Insub. ii. 21 = Demaseria sicula.
deltoides, *Burm. f. Fl. Cap. Prod.* 3.—Afr. austr.
distichia, Moench, Meth. 195 = Demaseria sicula.
dura, *Desv. Opusc.* 106.—Am. austr.
elatior, Sibth. & Sm. Fl. Graec. t. t. 75 = media.
elegans, *Doell, in Mart. Fl. Bras.* ii. iii. 135.—Bras.
Eragrostis, Desf. Fl. Atlant. t. 78 = Eragrostis minor.
Eragrostis, Linn. Sp. Pl. 70 = Eragrostis major.
erecta, *Lam. Illustr. Gen.* i. 187.—Reg. Argent.
fisra, *Desv. Opusc.* 106.—Am. austr.
geniculata, *Thunb. Prod. Pl. Cap.* 21.—Afr. austr.
grandis, Sallsb. Prod. 21 = maxima.
humilis, Bieb. Fl. Taur. Cauc. i. 66 = spicata.
imbricata, Steud. in Flora, xii. (1829) 489 = Poa Uniolae.
Lamarckiana, Cham. & Schlecht. in Linnaea, vi. (1831) 39 = rotundata.
Lamarckiana, *Nees, Agrost. Bras.* 481.—Am. austr.
lutescens, [Foucault, in] Desv. Journ. iii. (1814) t. 24. f. 2 = media.
macrocalyx, *Steud. Nom.* ed. II. i. 225.—Peruv.
major, Desv. ex Kunth, Enum. Pl. i. 371 = pelchella.
major, Presl, Cyp. et Gram. Sic. 42 = maxima.
maxima, *Linn. Sp. Pl.* 70.—Reg. Mediterr. ; Afr. austr.
media, *Linn. l. c.*—Europ. ; Orient ; Aa. bor. (Amphig. advent.).
megastachya, Hort. ex Steud. Nom. ed. II. i. 225 = Eragrostis major.
microstachys, *Steud. l. c.*—Peruv.
minima, Hort. ex Nichols. Dict. Gard. i. 312 = minor.
minor, *Linn. Sp. Pl.* 70.—Europ. ; Orient ; As. bor. (Amphig. advent.).
monspessulana, Gouan, Hort. Monsp. 45 = maxima.
montana, *Brouss. Cat. Monsp.* (1804) ; *Roem. & Schult. Syst.* ii. 631.—Teneriffa !
mucronata, Lam. Encyc. i. 466 = Lamarckiana.
multiflora, *Forst. ex Benew. Agrost.* 155, nomen.—Orient.
Neesii, *Doell, in Mart. Fl. Bras.* ii. iii. 132.—Bras.
nigra, *Burch. ex Steud. Nom.* ed. II. i. 225.—Afr. austr.
oblonga, Moench, Meth. 195 = Eragrostis major.
pannifolia, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 56.—Europ.
pilosa, *Schur, Enum. Pl. Transs.* 766.—Hungaria.
Poa, Spreng. ex Steud. Syn. Pl. Gram. 283 = Neesii.
pelchella, *Kunth, Rev. Gram.* i. 190.—Ins. Teneriff.
purpurascens, *Muhl. Descr. Uber.* 154.—Carolina.
ramigera, *Steud. Nom.* ed. II. i. 225.—Peruv.
rotundata, *Steud. Syn. Pl. Gram.* 284.—Mexic.
rubella, *Steud. l. c.* 282.—Afr. trop.
rubens, Poir. Suppl. i. 699 = maxima.
rubra, Lam. Illustr. i. 187 ; Pers. Syn. i. 98 = maxima.
rufa, Steud. Nom. ed. II. i. 225.—Peruv.
scabra, Nees, ex Steud. Syn. Pl. Gram. 276. n. 191 = Neesii.
Sellowii, *Nees, ex Steud. l. c.* 284. n. 36.—Bras.
serotina, Dum. Obs. Gram. Belg. 110 = media.
spicata, *Burm. f. Fl. Cap. Prod.* 3.—Afr. austr.

BRIZA, —
spicata, Lam. Encyc. i. 465 = Poa Michauxii.
spicata, *Sibth. & Sm. Fl. Graec.* i. t. 77.—Graecia ; As. Min.
spicigera, *Steud. Nom.* ed. II. i. 225.—Peruv. ; Chili.
stricta, *Steud. l. c.*—Chili.
suburistata, *Lam. Illustr.* i. 187.—Reg. Argent.
toluccensis, *H. B. & K. Nov. Gen. et Sp* i. 145.—Mexic.
tremula, Lam. Fl. Fr. iii. 587 = media.
trichotoma, *Steud. in Flora*, xxix. (1846) 90.—Japon
triloba, *Nees, Agrost. Bras.* 482.—Reg. Argent.
Uniolae, Nees, ex Steud. Syn. Pl. Gram. 283 = Eragrostis Uniolae.
violacea, Steud. l. c. = suburistata.
virens, Linn. Sp. Pl. ed. II. 108 = minor.
virens, Walt. Fl. Carol. 79 = Poa obtusa.
virens, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 362 = media.
viridis, Pall. ex Steud. Nom. ed. II. i. 226 = minor.

BRIZOPHILE, Salisb. Gen. Pl. Fragm. 84 (1866) = **Ornithogalum**, Tourn. (Liliac.).
nutans, Salisb. l. c. = O. nutans.

BRIZOPYRUM, Link, Hort. Berol. i. 159 (1827) = **Demaseria**, Dum. (Gramin.).
sicutum, Link, l. c. = D. sicula.

BRIZOPYRUM, J. Presl, in Rel. Haenk. 280 (1830) = **Distichlis**, Rafin. (Gramin.).
acutiflorum, Nees, Fl. Afr. Austr. 371 = Eragrostis acutiflora.
alternans, Nees, l. c. 369 = Poa alternans.
americanum, Link, Hort. Berol. i. 160 = Distichlis maritima.
borcale, Presl, Rel. Haenk. 28 = Distichlis maritima.
bromoides, Presl, l. c. = Poa bromoides.
calycinum, Presl, l. c. 281 = Poa calycina.
capense, Nees, Fl. Afr. Austr. 371 = Poa Uniolae.
condensatum, Fourn. Mex. Pl. Enum. Gram. 121 ; ex Hemsl. Biol. Centr. Am. Bot. iii. 578 = Distichlis condensata.
cyperoides, Nees, Fl. Afr. Austr. 374 = Eragrostis cyperoides.
Douglasii, Hook. & Arn. Bot. Beech. Voy. 404 = Poa Douglasii.
obtusiflorum, Fourn. Mex. Pl. Enum. Gram. 190 ; ex Hemsl. Biol. Centr. Am. Bot. iii. 579 = Distichlis thalassica.
ovatum, Nees, ex Steud. Syn. Pl. Gram. 282.—Peruv.
pilosum, J. Presl, in Rel. Haenk. i. 280 = Distichlis thalassica.
prostratum, Benth. ex Fourn. Mex. Pl. Enum. Gram. 190 ; ex Hemsl. Biol. Centr. Am. Bot. iii. 578 = Distichlis prostrata.
scirpoidcum, Steud. Syn. Pl. Gram. 282 = Schedonorus scirpoideus.
spicatum, Hook. & Arn. Bot. Beech. Voy. 408 = Distichlis maritima.
subspicatum, J. Presl, in Rel. Haenk. 281 = Poa subspicata.

BRIZULA, Hieron. in Bot. Zeit. xxx. (1872) 206 = Aphelia, R. Br. (Centrolepid.).
Drummondii, Hieron. in Abh. Naturf. Ges. Halle, xii. (1873) 206 = A. Drummondii.
gracilis, Hieron. l. c. = A. gracilis.
Muelleri, Hieron. l. c. 207 = A. Brizula.
Pumilio, Hieron. l. c. = A. Pumilio.

BROCCHIA, Mauri, ex Tenore, Cat. Ort. Napoli, 80 (1845) = **Simmondsia**, Nutt. (Euphorb.).
dichotoma, Mauri, l. c. = S. californica.

BROCCHIA, Vis. Pl. Aeg. et Nub. 85 (1836) = **Cotula**, Linn. (Compos.).
cinerea, Vis. l. c. = C. cinerea.
Kotschyi, Schult. ex Schweinf. & Aschers. Enum. 283 = C. Kotschyi.

BROCCHINIA, Schult. f. Syst. vii. 1250 (1830). *BROMELIACEAE*, Benth. & Hook. f. Gen. iii. 666. cordylinoides, *Baker, in Journ. Bot.* xx. (1882) 330.—Guiana.
paniculata, *Schult. f. Syst.* vii. 1250.—Bras.
Plumierii, *Griseb. Fl. Brit. W. Ind.* 593.—Ind. occ.
reducta, *Baker, in Journ. Bot.* xx. (1882) 331.—Guiana.

BROCHOSIPHON, Nees, in DC. Prod. xi. 492 (1847) = **Dicliptera,** Juss. (Acanthac.).
australis, Nees, l. c. = D. glabra.

BRODIAEA, Sm. in Trans. Linn. Soc. x. (1811) 2. t. 1.
LILIACEAE, Benth. & Hook. f. iii. 800.
CALLIPRORA, Lindl. Bot. Reg. t. 1390 (1832).
DICHELOSTEMMA, Kunth, Enum. Pl. iv 469 (1843). *Gardinia*, Benth. ‘in Merc. Chil. i. (1839)’.—*Cf.* Férussac, Bull. xx. (1830) 111.
HESPEROCORDUM, Lindl. Bot. Reg. sub t. 1293 (1829).
HOOKERA, Salisb. Parad. Lond. t. 98 (1808), nomen prius.
SCADIAKINTOS, Rafin. Fl. Tellur. iv. 23 (1836).
SAURETTIA, Kunth, Enum. Pl. iv. 475 (1843).
THEMIS, Salisb. Gen. Pl. Fragm. 85 (1866).
TRITELEIA, Dougl. ex Lindl. Bot. Reg. sub t. 1293 (1829).
VEATCHIA, Kellogg, in Proc. Calif. Acad. ii. (1859) 11.

alliacea, Mors, ex Baker, in Journ. Linn. Soc. xi. (1871) 375 = Leucocoryne alliacea.
Bridgesii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 287.—Calif.
californica, Lindl. ex Lem. in Fl. des Serres, Sér. I. v. (1849) 428 b = grandiflora.
capitata, *Benth. Pl. Hartw.* 339.—Am. bor. occ.
coccinea, *A. Gray, in Proc. Am. Acad.* vii. (1868) 389. —Calif.
congesta, *Sm. in Trans. Linn. Soc.* x. (1811) 3. t. 1.—Am. bor. occ.
crocea, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 288. —Calif.
Douglasii, *S. Wats. l. c.* 287.—Am. bor. occ.
filifolia, *S. Wats. l. c.* xvii. (1881–82) 381.—Calif.
gracilis, *S. Wats. l. c.* xiv. (1879) 288.—Calif.
grandiflora, Pursh, Fl. Am. Sept. i. 228 = lactea.
grandiflora, *Sm. in Trans. Linn. Soc.* x. (1811) 2.—Am. bor. occ.
grandiflora, Torr. in Stansb. Rep. 397 = Douglasii.
Howellii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 301.—Am. bor.
iridoides, Sims, Bot. Mag. t. 2532 = Leucocoryne iridoides.
iridoides, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 288. —Am. bor. occ.
lactea, *S. Wats. l. c.*—Am. bor. occ.
laxa, *S. Wats. l. c.* 287.—Calif.
Lemmonae, *S. Wats. l. c.* xx. (1885) 376.—Arizona.
minor, *S. Wats. l. c.* xiv. (1879) 286.—Am. bor. occ.
multiflora, *Benth. Pl. Hartw.* 339.—Am. bor. occ.
parviflora, Torr. & Gray, in Pacif. Rail. Rep. ii. (1854) 125 = multiflora.
pedunculata, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 287.—Calif.
stellaris, *S. Wats. l. c.* xvii. (1881–82) 380.—Calif.
terrestris, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 6. —Calif.
Torreyi, Wood, in Proc. Acad. Sc. Philad. (1868) 173 = terrestris.
volubilis, Baker, in Journ. Linn. Soc. xi. (1871) 377 = Strophobilrion californicum.

BROGNIARTIA, Walp. Rep. ii. 994 (1843) = **Brongniartia,** H. B. & K. (Leguminos.).

BROMBYA, F. Muell. Fragm. v. 4 (1866). *RUTACEAE*, Benth. & Hook. f. i. 991.
platynema, *F. Muell. l. c.*—Australia.

BROMELIA, [Plum.] ex Linn. Syst. ed. I (1735). *BROMELIACEAE*, Benth. & Hook. f. iii. 660.
AGALLOSTACHYS, Beer, Bromel. 35 (1857).
ANANAS, Gaertn. Fruct. i. 30. t. 11 (1788).
PINGUIN, [Dill.] Adans. Fam. ii. 67 (1763).
PREDOMELIA, Neck. Elem. iii. 150 (1790).
Acanga, Linn. Syst. ed. XII. 332 = B. Plagula et Karatas Plumieri.
Acanga, Schult. f. Syst. vii. 11. 1231 = Arseococcus micranthus.
Acorna, Thunb. ex Schult. f. l. c. 1232, nomen.—Hab.?
acaulis, Stokes, Bot. Mat. Med. ii. 304 = Karatas Plumieri.
agavaefolia, Brongn. in Rev. Horticle. (1875) 247 = Karatas agavaefolia.
agavoides, *Carr. l. c.* (1881) 31.—Hab.?

BROMELIA:—
albo-bracteata, *Steud. in Lechl. Berb. Am. Austr.* 58, nomen.—Chili.
albo-rosea, Lem. Illustr. Hortic. ii. (1855) Misc. 64 = Aechmea stuevoleana.
Ananas, Linn. Sp. Pl. 285 = Ananas sativa.
anticacantha, Bertol. in Virid. Bonon. Veg. (1824) 4; Colla, Hort. Ripul. App. ii. 343 = fastuosa.
aquilega, Salisb. Parad. Lond. t. 40 = Aechmea aquilega.
arenaria, Vell. Fl. Flum. 180; iii. t. 114 = Quesnelia rufa.
bicolor, Ruiz & Pav. Fl. Peruv. Chil. iii. 83 = Rhodostachys bicolor.
Binoti, Hort. Jacob-Makoy.—*Cf.* Gard. Chron. (1880) ii. 151 = B. Pinguin.
bracteata, Schult. f. Syst. vii. 11. 1280, ex parte = Aechmea laxiflora.
bracteata, Sw. Prod. Veg. Ind. Occ. 56 = Aechmea bractesta.
Caratas, Hill, Veg. Syst. vii. 19 = Karatas Plumieri.
carnea, Beer, Bromel. 31 = Rhodostachys andina.
Carolinae, Beer, l. c. 29; Orig. ex Regel, Gartenfl. vi. (1857) 361 = Karatas Carolinae.
chrysantha, *Jacq. Hort. Schaenb.* i. 28.—Am. austr.
clandestina, Hort. ex Carr. in Rev. Hortic. (1880) 256 = Greigia sphacelata.
comata, *Beer, Bromel.* 34.—Bras.
Commeliniana, De Vriese, Del. Sem. Hort. Amst. 1844; ex Hoev. & De Vriese, Tijdschr. xii. (1845) 1. fastuosa.
comosa, Hill, Veg. Syst. vii. 18 = Ananas sativa.
concentrica, Beer, Bromel. 29 = Karatas Laurentii?
crassa, Steud. in Lechl. Berb. Am. Austr. 58 = Greigia laxiflora.
cruenta, R. Grah. in Edinb. N. Phil. Journ. (Oct.–Dec. 1828) 174 = Billbergia cruenta.
denticulata, *C. Koch, Wochenschr.* ii. (1859) 151.—Bras.
discolor, *Lindl. Bot. Reg.* (1838) *Misc.* 45.—Am. austr.
edulis, Salisb. Prod. 247 = Ananas sativa.
elegans, *Hort. Hamb. ex Schult. f. Syst.* vii. 11. 1283. —Hab.?
exudans, Lodd. Bot. Cab. t. 801 = Aechmea exsudans.
fastuosa, *Lindl. Collect.* i.—N. Granat.
Fernandae, *Morr. in Illustr. Hortic.* xviii. (1871) 114. t. 65.—Bras.
gigantea, *Hort. Vindob. ex Schult. f. Syst.* vii. 11. 1283.—Hab.?
glabra, Schult. f. l. c. 1285 = Ananas sativa.
hemisphaerica, Lam. Encyc. i. 145 = Karatas humilis.
Hookeri, Sweet, Hort. Brit. ed. I. 425 = Aechmea nudicaulis.
humilis, Linn. Syst. ed. XII. 332; Jacq. Enum. Vindob. App. 306 = Karatas humilis.
ignea, Beer, Bromel. 35 = B. Pinguin.
incarnata, Ruiz & Pav. Fl. Per. iii. 82 = Billbergia incarnata.
inermis, Steud. Nom. ed. II. i. 236 = Ananas sativa.
iridifolia, Ness & Mart. in Nov. Act. Nat. Cur. xi. (1823) 16 = Billbergia iridifolia.
Itatiaiae, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 114.—Bras.
Joinvillei, Z. Morr. ex C. H. Morr. in Belg. Hortic. xxvi. (1876) 161. t. 10, 11 = Rhodostachys pitcairniaefolia.
Karatas, Linn. Sp. Pl. 285 = Karatas Plumieri.
lacinicora, *Mart. ex Schult. f. Syst.* vii. 11. 1278.—Bras.
laevis, *Hort. Cortevuh. ex Schult. f. l. c.* 1282.—Hab.?
Landbeckii, Lechl. ex Phil. in Linnaea, xxxiii. (1864–65) 246 = Greigia Landbeckii.
lanuginosa, Beer, Bromel. 32 = Pitcairnia lanuginosa.
laxiantha, Willd. ex Schult. f. *Syst.* vii. 11. 1283 = Puya laxiata.
latifolia, *Willd. ex Schult f. l. c.* 1283.—Isa. Antigua.
Lindleyana, Lem. Jard. Fleur. iii. (1853–53) t. 288, in obs. = Arseococcus micranthus.
lingulata, Linn. Sp. Pl. 285 = Aechmea lingulata.
longifolia, Lindl. in Past. Flow. Gard. ii. t. 65 = Rhodostachys andina.
longifolia, *Rudge, Gui.* i. 31. t. 49.—Guiana.
longifolia, *Klotzsch, in Schomb. Reise*, iii. 992, nomen. —Guiana.
lucida, *Willd. Enum. Hort. Berol.* 343 = Ananas sativa.

BROMELIA:—
lutea, G. F. W. Mey. Prim. Fl. Essequ. 145 = Aechmea nudicaulis.
malacantha, Ker-Gawl. in Bot. Reg. t. 766 = Aechmea bromeliaefolia.
Mertensii, G. F. W. Mey. Prim. Fl. Essequ. 144 = Aechmea Mertensii.
mexicana, *Nois. ex Schult. f. Syst.* vii. 11. 1283, nomen.—Mexic.
Moensi, *Linden, Cat.* (1872) n. t. 6.—Bras.
muricata, Aruda, Diss. Pl. Bras. 21 = Ananas bracteata?
nudicaulis, Linn. Sp. Pl. 286 = Aechmea nudicaulis.
nudicaulis, Reichb. Ic. Exot. 135 = Billbergia pyramidalis.
pallida, Ker-Gawl. in Bot. Reg. t. 344 = Billbergia speciosa.
paniculata, J. F. Gmel. Syst. 529 = Aechmea paniculigera.
paniculigera, Reichb. Ic. Exot. t. 239, 240 = Aechmea exsudans.
paniculigera, Sw. Prod. Veg. Ind. Occ. 56 = Aechmea paniculigera.
pauciflora, *C. Koch, Wochenschr.* ix. (1866) 183.—Hab.?
Pinguin, *Linn. Sp. Pl.* 285.—Am. trop.
pitcairnifolia, C. Koch, Wochenschr. xi. (1868) 825 = Rhodostachys pitcairniaefolia.
pumila, *Hort. ex Schult. f. Syst.* vii. 11. 1283.—Hab.?
pyramidalis, Sims, Bot. Mag. t. 1782 = Billbergia pyramidalis.
pyramidata, Beer, Bromel. 34 = Puya pyramidata.
Rohaniana, Walp. Ann. vi. 71 = Billbergia vittata.
rubra, Hort. ex Schult. f. Syst. vii. 11. 1286 = Ananas sativa.
Sagenaria, Arruda, Diss. Pl. Bras. 14 = Ananas bracteata.
scarlatina, Hort. ex Hérincq, Hortic. Franç. (1869) 246 = Disticacanthus scarlatinus.
scariatina, *E. Morr. ex C. H. Morr. in Belg. Hortic.* xxxi. (1881) 164.—Bras.
Scepitrum, Hueg. Parad. Vind. fasc. viii. (sic) = fastuosa.
semiserrata, Willd. Enum. Hort. Berol. 346 = Ananas sativa.
sepiaria, *Hort. ex Schult. f. Syst.* vii. 11. 1283.—Hab.?
Serra, *Griseb. in Gott. Abh.* xxiv. (1879) 328.—Reg. Argent.
sessiliflora, *Lodd. ex Loud. Hort. Brit.* 118.—Am austr.
silvestris, Vell. Fl. Flum. 129; iii. t. 113 = Ananas sativa.
sphacelata, Ruiz & Pav. Fl. Per. iii. 82. t. 274 = Greigia sphacelata.
spicata, *Lam. Encyc.* i. 146.—Ins. Martinic.
subpinnea, Wendl. ex Schult. f. Syst. vii. 11. 1286 = Ananas sativa.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 578.—Guiana.
sylvestris, *Willd. ex Sims, Bot. Mag.* t. 2392.—Am. trop.
thyrsiflora, Willd. ex Schult. f. Syst. vii. 11. 1282 = Aechmea spicata.
tinctoria, Mart. Reise Bras. ii. 554 = Aechmea bromeliaefolia.
tristis, *Beer, Bromel.* 30.—Bras.
variegata, Arruda, Diss. Pl. Bras. 7 = Billbergia speciosa.
viridans, Hort. ex Schult. f. Syst. vii. 11. 1285 = Ananas sativa.
viridis, Hort. ex Schult. f. l. c. = Ananas sativa.
zebrina, Herb. in Bot. Mag. t. 2656 = Billbergia zebrina.

BROMFELDIA, Neck. Elem. ii. 347 (1790) = Jatropha?, Linn. (Euphorb.).

BROMHEADIA, Lindl. Bot. Reg. (1841) App. 90. *ORCHIDEAE*, Benth. & Hook. f. iii. 540.
sporoides, *Reichb. f. Otia Bot. Hamb.* 64.—Burma.
Finlaysoniana, Reichb. f. in Walp. Ann. vi. 690, 692 = palustris.
palustris, *Lindl. Bot. Reg.* (1841) *Misc.* 99.—Malaya.

BROMIDIUM, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (Pl. Meyen.) 184 (1843) = Dayeuxia, Clar. (Gramin.).

BROMIDIUM :—
anomalum, Doell, in Mart. Fl. Bras. ii. III. 103.—Bras.
brevifolium, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. 155 = D. brevifolia.
caespitosum, Nees & Meyen, l. c. = Agrostis Presili.
hygrometricum, Nees & Meyen, l. c. 154.—Bras.
laxatum, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 415 = D. quadriseta.
quadriseta, Nees, l. c. 416 = D. quadriseta.
rigescens, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. 155 = Agrostis rigescens.
spectabile, Nees & Meyen, l. c. 156.—Peruv.

BROMOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 187 = **Bromus**, Dill. (Gramin.).
aspera, Fourr. l. c. = Bromus asper.
erecta, Fourr. l. c. = Bromus erectus.

BROMUS, Dill. ex Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1200.
ANISANTHA, C. Koch, in Linnaea, xxi. (1848) 394.
BROMOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 187.
CERATOCHLOA, Beauv. Agrost. 75. t. 15. f. 7 (1812).
LASIOPOA, Ehrh. Beitr. iv. 147 (1789).
LIBERTIA, Lej. in Nov. Act. Nat. Cur. xii. (1825) 755. t. 65.
MICHELARIA, Dum. Obs. Gram. Belg. 77. t. 16 (1823).
SCHEDONORUS, Beauv. Agrost. 99. t. 19. f. 2 (1812), partim.
SERRAFALCUS, Parl. Fl. Nov. 75 (1842).
TRIBULUS, Steud. Syn. Pl. Gram. 328 (1854).
ZERNA, Panz. in Denkschr. Acad. Muench. 1813 (1814) 296, partim.
abortiflorus, St. Amans, Fl. Agen. 44 = tectorum.
abocanis, *Hochst. ex Steud. Syn. Pl. Gram.* 326.—Abyss.
aegyptiacus, Tausch, in Flora, xx. (1837) 124 = brachystachys.
affinis, Dum. Obs. Gram. Belg. 118 = mollis ?
agrarius, Dum. l. c. = diffusus.
agrarius, Hornung, ex Steud. Nom. ed. II. i. 226 = racemosus.
agrestis, All. Fl. Pedem. ii. 249 = erectus.
albidus, *Bieb. Fl. Taur. Cauc.* iii. 79.—Reg. Caucas.
alectorus, *Trin. ex Ledeb. Fl. Ross.* iv. 361.—Unalaska.
alopecuroides, Lag. Gen. et Sp. Nov. 4 = Kocleria phleoides.
alopecuroides, Moris, ex Steud. Nom. ed. II. i. 226, 228 = intermedius.
alopecuroides, Poir. Encyc. Suppl. i. 703 = seq.
Alopecurus, *Poir. Voy. Barb.* ii. 100; *Vahl, Symb. Bot.* ii. 32.—Reg. Mediterr.
Alopecurus, Pers. Syn. i. 95 = prasc.
alpestris, *Steud. in Lechl. Berb. Am. Austr.* 52, nomen. —Chili.
altissimus, Gilib. Exercit. ii. 587 = arvensis.
altissimus, *Pursh, Fl. Am. Sept.* ii. 738.—Am. bor.
altissimus, [Weber, in] Wigg. Prim. Fl. Hols. 9 = asper.
amblyneus, Jord. ex Nym. Consp. 891 = rigidus.
ambiguus, Cyrill. Pl. Rar. Isac. i. 10 = Festuca bromoides.
amplus, C. Koch, in Linnaea, xxi. (1848) 418 = sterilis.
anatolicus, Boiss. & Heldr. Diagn. Ser. I. xiii. 63 = patulus.
angustifolius, Schrank, Baier. Fl. i. 366 = erectus.
angustifolius, *Hornem. Hort. Hafn. Suppl.* 13.—Hab. ?
angustissimus, C. Koch, in Linnaea, xxi. (1848) 421 = Bieberstenii.
anomalus, *Rupr. in Bull. Acad. Brux.* ix. II. (1842) 266.—Mexic.
antarcticus, Hook. f. Fl. Antarct. 97 = Danthonia bromoides.
aquaticus, Schrank, Baier. Fl. i. 367 = Festuca gigantea.
arduennensis, *Dum. Obs. Gram. Belg.* 77.—Belgium.
arenarius, Fourr. in Mém. Acad. Toul. iii. (1788) 308 = tectorum.
arenarius, *Labill. Nov. Holl. Pl.* i. 23. t. 28.—Austral.; N. Zel.
arenarius, Thomine Desm. in Mém. Soc. Linn. Calv. (1824) 40 = mollis.
arfakiana, *Steud. Syn. Pl. Gram.* 330.—Persia.
armenus, *Boiss. Fl. Orient.* v. 643.—Armenia.
arrhenatheroides, *Baker, in Journ. Linn. Soc.* xx. (1884) 301.—Madag.
arundinaceus, Roth, Tent. Fl. Germ. ii. 141 = Festuca elatior.
arvensis, Knapp, Gram. Brit. t. 32 = racemosus.
arvensis, Lam. Fl. Fr. iii. 607 = erectus.
arvensis, *Linn. Sp. Pl.* 77.—Europ.; Orient.; As. bor.
arvensis, Ord. in Fl. Dan. t. 305 = secalinus.
asper, *Murr. Prod. Gotting.* 42.—Europ.; Orient.; As. bor.
asper, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. i. 73 = erectus.
asperipes, Jord. ex Nym. Consp. 891 = rigidus.
asperrimus, *Tenore, Fl. Nap.* t. 105.—Italia.
atticus, Steud. ex Nym. Consp. 892 = mollis.
auleticus, *Trin. ex Nees, in Mert. Fl. Bras.* ii. 468.—Reg. Argent.
auriculatus, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 439 = arduennensis.
australis, R. Br. Prod. i. 178 = arenarius.
avenaceus, Lam. Illustr. i. 194 = tectorum.
avenaeformis, Steud. Syn. Pl. Gram. 326 = Triactum flavescens.
avenoides, *Baker, in Journ. Linn. Soc.* xx. (1884) 302.—Madag.
badensis, *C. C. Gmel.* Fl. Bad. iv. 75. t. 5 = secalinus.
barbatus, Savi, Bot. Etrusc. ii. 61 = Festuca Alopecuros.
Baumgartenii, Steud. Syn. Pl. Gram. 330 = rigidus.
Berterianus, *Colla, in Mem. Acc. Torin.* xxxix. (1835) 35.—Chili.
biaspis, Nees, ex Steud. Syn. Pl. Gram. 322.—Chili.
Biebersteinii, Roem. & Schult. Syst. iii. 686 = albidus.
bifidus, Thunb. Fl. Jap. 53 = Triactum cernuum.
Billotii, Sch. Bip. in Flora, xxxii. (1849) 233 = secalinus.
bohemicus, F. W. Schmidt, ex Mert. & Koch, in Roehl. Deutschl. Fl. ed. III. i. 683 = Festuca Myurus.
Borael, Jord. ex Nym. Consp. 891 = rigidus.
brachyanthecus, *Doell, in Mart. Fl. Bras.* ii. III. 110.—Bras.
brachystachys, *Hornung, in Flora*, xvi. (1833) II. 417. —Oriens; Aegypt.
breviaristatus, *Buchl. in Proc. Acad. Sc. Philad.* '1862' (1863) 49.—Am. bor. occ.
brevis, *Steud. Syn. Pl. Gram.* 326.—Mendoza.
brevisetus, *Dum. Fl. Belg.* 155.—Belgium.
brizaeformis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 30.—Reg. Caucas.; Persia.
brizaeformis, Lam. Illustr. i. 193. n. 1060 = Briza Lamarckiana.
brizoides, Willd. ex Steud. Nom. ed. II. i. 227 = Briza rotundata.
Brongniartii, *Kunth, Enum. Pl.* i. 491.—Bras.
Bucheanus, Tenore, ex Steud. Nom. ed. II. i. 227 = Brachypodium distachyum.
caerulescens, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 9.—Hab. ?
caespitosus, Host, Gram. Austr. iv. t. 18 = Brachypodium pinnatum.
caespitosus, Steud. Nom. ed. I. 119 = Brachypodium ramosum.
Caldasii, H. B. & K. Nov. Gen. et Sp. i. 151 = Festuca Caldasii.
canadensis, Michx. Fl. Bor. Am. i. 65 = ciliatus.
canariensis, Zuccagni, Cent. n. 27; in Roem. Coll. 135 = lanceolatus.
cancscens, Viv. Fl. Lyb. Spec. 5 = rubens.
canus, Willd. ex Steud. Nom. ed. II. i. 227 = Festuca mollis.
capensis, Steud. in Flora, xli. (1829) 491 = vestitus.
capillaris, Moench, Meth. 194 = Leptochloa dominguensis.
capillatus, *Vahl, ex Hornem. Hort. Hafn.* i. 101.—Hab. ?
capillifolia, *Dufour, ex Roem. & Schult. Syst.* ii. 736. —Hispan.
cappadocicus, *Boiss. & Bal. in Bull. Soc. Bot. Fr.* iv. (1857) 306.—As. Min.
capreorum, Hackl. in Oestr. Bot. Zeitschr. xxix. (1879) 209 = caprinus.
caprinus, *Kerner, ex Hack. l. c.*—Italia.
carinatus, *Hook. & Arn. Bot. Beech. Voy. Suppl.* 408.—Am. bor. occ.
carthaticus, Vahl, Symb. Bot. ii. 32 = unioloides.
cascasicus, *Fisch. ex Steud. Nom.* ed. II. i. 227.—Reg. Caucas.
caudatus, *Brot. Phyt. Lusit.* t. 90.—Lusitan.
Cavanillesii, *Willd. ex Willd. & Lange, Prod. Fl. Hisp.* i. 101.—Hispan.
Cebadilla, *Steud. Syn. Pl. Gram.* 321.—Chili.
Chiappirianus, De Not. ex Nym. Consp. 824 = patulus.
chilensis, *Trin. in Linnaea*, x. (1836) 304.—Chili.
chloroticus, Bieb. ex Kunth, Enum. Pl. i. 490 = albidus.
Chrysopogon, Viv. Fl. Lyb. Spec. 4. t. 2 = scoparius.
ciliatus, Huds. Fl. Angl. ed. I. 40 = madritensis.
ciliatus, Lam. Fl. Fr. iii. 609 = Brachypodium distachyum.
ciliatus, *Linn. Sp. Pl.* 76.—Am. bor.
ciliatus, Mahl. Desc. Gram. 169 = Kalmii.
cinereus, S. G. Gmel. ex Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 66 = Triactum Gmelinii.
cincinnatus, Tenore, Lez. Bot. iv. t. 52; Fl. Nap. iii. 85 = Avena bromoides.
circinnatus, Kunth, Enum. Pl. i. 423 = prasc.
coarctatus, Hornem. ex Kunth, l. c. 303 = Gaudinia geminiflora.
cognatus, *Steud. Syn. Pl. Gram.* 321.—Abyssin.
collinus, *Phil. in Linnaea, xxix. (1857–58)* 102.—Chili.
coloratus, Baumg. ex Nym. Consp. 824 = squarrosus.
coloratus, *Steud. Syn. Pl. Gram.* 329.—Reg. Magellan.
communatus, Bieb. Fl. Taur. Cauc. iii. 77 = patulus.
communatus, Gnss. ex Steud. Syn. Pl. Gram. 325 = scabratus.
communitus, Schrad. Fl. Germ. i. 358 = racemosus.
compactus, Hort. ex Steud. Nom. ed. II. i. 227 = mollis.
compressus, Lag. Elench. 4 = ciliatus.
condensatus, *Hack. in Oestr. Bot. Zeitschr.* xxix. (1879) 209.—Europ.
confertus, Bieb. Fl. Taur. Cauc. i. 71 = scoparius.
confertus, Bor. Fl. Centr. Fr. ed. II. 566 = mollisformis.
confinis, *Nees, ex Steud. Syn. Pl. Gram.* 320.—Ind. or.
conformis, *Steud. l. c.* 323.—Japon.
contortus, Desf. Fl. Atlant. i. 95 = B. Alopecuros.
cornicularus, Lam. Fl. Fr. iii. 608 = Brachypodium pinnatum.
cruinus, *Boiss. & Hohen. Diagn.* Ser. I. xiii. 64.—Pers.; Afghan.; Turkest.
cristatus, Linn. Sp. Pl. 78 = Agropyron cristatum.
cristatus, Spreng. Nachr. i. Bot. Gart. Halle, 15 = Kocleria phleoides.
cultus, *Steud. Syn. Pl. Gram.* 323.—Europ.
cylindricus, *Brot. Phyt. Lusit.* 98.—Lusitan.
Cyrs, Trin. in C. A. Mey. Vers. Pfl. Cauc. 24 = patulus.
dactyloides, Roth, Nov. Sp. 60 = Kocleria phleoides.
Danthoniae, Trin. in C. A. Mey. Vers. Pfl. Cauc. 24 = macrostachys.
decipiens, Hort. Madr. ex Steud. Nom. ed. II. i. 227 = madritensis.
decolorans, Reichb. Fl. Germ. Excurs. 44 = Festuca elatior.
decumbens, Koel. Gram. 242 = decumbens.
denudatus, *Steud. Syn. Pl. Gram.* 322.—' Terra nova.'
depauperatus, *J. Presl, in Rel. Haenk.* i. 263.—Am. bor. occ.
dertonensis, All. Fl. Pedem. ii. 249 = Festuca bromoides.
diandrus, Curt. Fl. Lond. fasc. vi. t. 5 = madritensis.
diandrus, Roth, Bot. Abh. 44 = pilosus.
diffusus, *Dum. Obs. Gram. Belg.* 118.—Belgium.
digitatus, Poir. ex Beauv. Agrost. 156 = madritensis.
dilatatus, Lam. Encyc. i. 468 = madritensis.
dimidiiflorus, *Baker, in Journ. Linn. Soc.* xx. (1884) 301.—Madag.
distachyos, Linn. Cent. ii. 8; Amoen. Acad. iv. 304 = Brachypodium distachyum.
distichus, Georgi, Bemerk. Reise Russ. Reich. i. 197 = Agropyron cristatum.
distichus, Moench, Meth. 192 = sterilis.
diversiflorus, Rhode, ex Lolael. in Desv. Journ. Bot. ii. (1809) 314 = macrostachys.

BROMUS:—
dortonensis, Kunth, Enum. Pl. i. 397, in syn. = Festuca bromoides.
dumetorum, Lam. Fl. Fr. iii. 605 = asper.
dumosus, Vill. Hist. Pl. Dauph. ii. 119 = Brachypodium sylvaticum.
Ehrhartii, Gaud. in Roem. Collect. 19 = secalinus.
elatior, Koel. Gram. 214 = Festuca elatior.
elatior, Spreng. Syst. i. 359 = Festuca arundinacea.
elatus, Gueldenst. ex Ledeb. Fl. Koss. iv. 358 = Festuca elatior.
elongatus, Gaud. Agrost. i. 305 = racemosus.
erectus, *Huds. Fl. Angl.* ed. 1. 39.—Europ.; As. Min.
erectus, Moris, Stirp. Sard. Elench. i. 50 = scoparius.
erectus, Ledeb. Fl. Koss. iv. 358 = inermis.
exaltatus, *Bernh. in Linnaea*, xv. (1841) *Litt.* 90.—Mexic.
fasciicularis, Tenore, Fl. Nap. iv 17, 161 = fasciculatus.
fasciculatus, *Presl, Cyp. et Gram. Sic.* 39.—Reg. Mediterr. or.
Ferronii, Mabelli, in Act. Soc. Linn. Bord. xxv. (1864) 613.—Cf. Bull. Soc. Bot. Fr. xiv. (1867) Bibl. 188 = mollis.
festucoides, *Link, in Schrad. Journ.* ii. (1799) 315.—Lusitan.
fibrosus, *Hack. in Oestr. Bot. Zeitschr.* xxix. (1879) 300.—Europ. or.
flabellatus, *Hack. ex Boiss. Fl. Orient.* v. 648.—Palaestin.
flavescens, *Tausch, in Flora*, xx. (1837) i. 124.—Sicilia.
Fonki, *Phil. in Linnaea*, xxix. (1857–58) 102.—Chili.
fragilis, Schrank, Baier. Fl. i. 370 = Brachypodium pinnatum.
fragilis, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 90.—Europ.
frigidus, *Ball, in Journ. Linn. Soc.* xxii. (1885) 68.—Peruv.
frigidus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 645.—Persia.
Gaudini, Roem. & Schult. Syst. ii. 682 = racemosus.
geniculatus, Linn. Mant. i. 33 = Festuca geniculata.
geniculatus, *Steud. Syn. Pl. Gram.* 315.—Oriens.
giganteus, Leers, Fl. Herborn. 33 = Festuca elatior.
giganteus, *Linn. Sp. Pl.* 77.—Europ.; Sibir.
gigantinus, Vill. Hist. Pl. Dauph. ii. 118 = asper.
Gilliesii, *Nees, ex Steud. Syn. Pl. Gram.* 328.—Chili.
glaber, Reich. Fl. Franco-Moen. i. 19 = Festuca gigantea.
glaber, Scop. Fl. Carn. ed. II. i. 84 = Agropyron repens.
glaber, Willd. ex Steud. Nom. ed. II. i. 227 = pitensis.
glaberrimus, C. Koch, in Linnaea, xxi. (1848) 420 = erectus.
glaucus, Hort. ex Reichb. Fl. Germ. Excurs. 38 = Festuca ovina.
glaucus, Lapeyr. Hist. Abr. Pl. Pyr. t. 58 = erectus.
glaucus, Spreng. in Link, Hort. Berol. i. 170 = Festuca rubra.
glomeratus, Scop. Fl. Carn. ed. II. i. 76 = Dactylis glomerata.
glomeratus, Tausch, in Flora, xx. (1837) 124 = mollis.
gracilis, Leyss. Fl. Hal. 26. n. 116 = Brachypodium pinnatum.
gracilis, Weigel, Obs. 15 = Brachypodium sylvaticum.
gracillimus, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 537 = erubens.
grandiflorus, *C. Koch, in Linnaea*, xxi. (1848) 419 (sp. dub.).—Pontus.
grandiflorus, Weigel, Obs. 9 = sterilis, tectorum.
grossus, DC. Fl. Fr. iii. 68 = secalinus.
grossus, Desf. in C. C. Gmel. Fl. Bad. iv. t. 3, 3 = velutinus.
Gussoni, Parl. Pl. Nov. 66 = rigidus.
gyganteus, Crantz, Inst. t. 390 = giganteus.
Gynendrus, Roth, in Roem. & Ust. Mag. iv. 10[mo] St. (1790) 20 = maditensis.
Haenkeanus, Kunth, Enum. Pl. i. 416 = unioloides.
Haussknechtii, *Boiss. Fl. Orient.* v. 648.—Mesopotam.
hirsutissimus, Cyrill. ex Roem. & Schult. Syst. ii. 643 = asper.
hirsutus, Curt. Fl. Lond. fasc. ii. t. 8 = asper.
hirsutus, Schrank, in Denkschr. Bot. Ges. Regensb. (1818) 161 = squarrosus.
hirtus, Lichtst. in Roem. & Schult. Syst. ii. 654.—Afr. austr.

BROMUS:—
hispidus, Savi, Bot. Etrusc. ii. 62 = Koeleria hispida.
Hohenackeri, Hochst. ex Ledeb. Fl. Ross. iv. 416 = Ventenata macra.
holciformis, *Steud. & Hochst. ex Steud. Syn. Pl. Gram.* 324.—Chili.
Hookeri, Fourn. Mex. Pl. En. Gram. 197; ex Hemsl. Biol. Centr. Am. Bot. iii. 583 = ciliatus.
Hoserianus, Thurber, in Bot. U. St. Expl. Exped. ii. 493 = virens.
hordeaceus, C. C. Gmel. Fl. Bad. iv. 68 = secalinus.
hordeiformis, Linn. Sp. Pl. 77 = racemosus, mollis.
hordeiformis, Lam. Illustr. i. 105 = Festuca bromoides.
Hughi, Tod. ex Nym. Consp. 833 = intermedius.
humilis, Cav. Ic. vi. 65 = scoparius.
hyalinus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 90 = arvensis.
hybridus, *Schur, Enumer. Pl. Transs.* 801.—Transsylv.
imperialis, Hort. ex Steud. Nom. ed. II. i. 227 = ciliatus.
incrassatus, Lam. Encyc. i. 468 = rigidus, Festuca geniculata.
indicus, Hort. ex Steud. Nom. ed. II. i. 227 = rigidus.
inermis, *Leyss. Fl. Hal.* 16; *Linn. Mant.* ii. 186.—Europ.; Reg. Caucas.; As. Bor.
inermis, Sten. in Mém. Soc. Nat. Mosc. iii. (1812) 252 = albidus.
insignis, *Buese, in Mém. Pl. Yungh.* 247.—Java.
intermedius, *Guss. Fl. Sic. Prod.* i. 114.—Reg. Mediterr. intermedius, *Zumag. Fl. Pedem.* i. 166.—Pedemont.
japonicus, *Thunb. Fl. Jap.* 51. t. 11.—Japon.
jubatus, Tenore, Fl. Nap. 10 = sterilis.
jubatus, *Vill. Hist. Pl. Dauph.* ii. 118.—Gallia.
Kalmii, *A. Gray, Man.* 600.—Am. bor.
Kochii, C. C. Gmel. Fl. Bad. iv. 76. t. 6 = patulus.
Krauei, *Regel, in Act. Hort. Petrop.* vii. (1880) 600.—As. centr.
lanatus, H. B. & K. Nov. Gen. et Sp. i. 150 = Festuca lanata.
lanceolatus, Roth, Catalect. i. 18 = macrostachys.
lanuginosus, Poir. Encyc. Suppl. i. 708 = macrostachys.
lateriparous, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = tectorum.
latifolius, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 865.—Sibir.
laxiflorus, *Spreng. in Zeph. ex Steud. Nom.* ed. II. i. 228.—Afr. austr.
laxus, Hornem. Hort. Hafn. ii. 954 = erectus.
laxus, Reichb. lc. Fl. Germ. i. 32. t. 77. f. 1605 = inermis.
lenis, *Y. Presl, in Rel. Haenk.* i. 262.—Peruv.
leptoclados, *Nees, Fl. Afr. Austr. Gram.* i. 458.—Afr. austr.
leptostachys, Ehrenb. ex Boiss. Fl. Orient. v. 654 = brachystachys.
leptostachys, [Pieri,] in Ionios Anthol. iii. (1834) 680 = squarrosus.
leptostachyus, *Steud. Syn. Pl. Gram.* 324.—Gallia.
liguisticus, All. Fl. Pedem. ii. 249 = rubens, scoparius, Festuca geniculata.
lithobius, *Trin. in Linnaea*, x. (1836) 303.—Chili.
littoralis, *Steud. Syn. Pl. Gram.* 325.—' Austr. littoralis.'
littoreus, Georgi, Beschr. Russ. Reich. Nachtr. 257 = inermis.
littoreus, Retz. Fl. Scand. Prod. ed. I. 19 = Festuca elatior.
lividus, H. B. & K. Nov. Gen. et Sp. i. 150 = Festuca livida.
loliaceus, Tenore, Fl. Nap. iii. 91 = Triticum Barrelieri.
longearistatus, Griseb. ex Ledeb. Fl. Ross. iv. 396 = sterilis.
longiflorus, Spreng. Fugill. ii. 23 = tectorum.
longifolius, *Schousb. Vestr. Marokko*, 44.—Marocco.
longifolius, Kummer & Sendt. in Flora, xxxii. (1849) 757 = tectorum.
lusonensis, J. Presl, in Rel. Haenk. i. 262 = Triticum lusonense.
macranthos, Meyen, Reise, i. 311, nomen = setifolius.
macranthus, E. Desv. in C. Gay, Fl. Chil. vi. 448 = praec.
macrochaetus, Boiss. Diagn. Ser. I. xiii. 64 = arvensis.
macrostachys, *Desf. Fl. Atlant.* i. 96. t. 19.—Reg. Mediterr.
macrostachyus, Cass. Fl. Sic. Prod. 115 = lanceolatus.
madritensis, Cav. ex Kunth, Enum. Pl. i. 413 = maximus.
madritensis, *Linn. Cent. Pl.* i. 5; *Amoen. Acad.* iv. 295.—Europ.; Afr. bor.; Oriens.
malacanthus, Trin. ex Ledeb. Fl. Ross. iv. 361 = ciliatus.

BROMUS:—
Mango, *B. Dew. in C. Gay, Fl. Chil.* vi. 440.—Chili.
marginatus, Steud. Syn. Pl. Gram. 329 = ciliatus.
massiliensis, *Porsk. Fl. Aegypt. Arab.* 208.—Massilia.
matritensis, 'Linn.' ex Roem. & Schult. Syst. ii. 651 = maditensis.
matritensis, Cav. Ic. vi. 67 = rigidus.
Matthewsii, *Steud. Syn. Pl. Gram.* 325.—Peruv.
maximus, Desf. Fl. Atlant. i. 95. t. 26 = rigidus.
maximus, Gilib. Exercit. ii. 535 = secalinus.
maximus, Roth, N. Beitr. i. 114 = pilosus.
megapotanicus, *Spreng. Syst. iv. Cur. Post.* 86.—Bras.
mexicanus, Hort. ex Steud. Nom. ed. II. i. 228 = Triticum mexicanum.
Michaëli, Savi, Bot. Etrusc. i. 78 = festucoides, Avellinia Michell.
modensis, *Steud. Syn. Pl. Gram.* 324.—Gracia.
molliformis, *Lloyd, Fl. Loire-Inf.* 315.—Gallia.
mollis, *Linn. Sp. Pl.* ed. II. 112.—Europ.; Afr. et As bor.
mollis, Thunb. Prod. Pl. Cap. 22 = vestitus.
mollissimus, *Hornem. Hort. Hafn. Suppl.* 13.—Hab. ?
monostachyus, Steud. Nom. ed. II. i. 228 = Brachypodium distachyum.
montanus, Gaertn. Mey. & Scherb. Fl. Wett. ex Kunth, Enum. Pl. i. 413 = erectus.
montanus, Scop. Fl. Carn. ed. II. i. 82 = asper.
montanus, *Retz. Prod. Scand.* 124.—Europ. bor.
muceosus, Willd. ex Steud. Nom. ed. II. i. 228 = unioloides.
multiflorus, Host, Gram. Austr. i. t. 11 = patulus.
multiflorus, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 865 = macrostachys.
multiflorus, Roth, Tent. Fl. Germ. i. 47 = racemosus.
multiflorus, Sm. in Trans. Linn. Soc. iv. (1798) 283 = secalinus.
multiflorus, Weigel, Obs. 2. t. 1. f. 1 = arvensis.
multiflorus, Willd. Sp. Pl. i. 428 = commutatus.
multipaicatus, Roem. & Schult. Syst. ii. 650 = maditensis.
Munroi, *Beiss. Fl. Orient.* v. 643.—Afghanist.
muralis, Huds. Fl. Angl. ed. II. i. 50 = maditensis.
murorum, Sprun. ex Roem. & Schult. Syst. ii. 651 = pilosus, tectorum.
mutabilis, *F. Schultz, in Flora*, xxxii. (1849) 284.—Europ.
nanus, Weigel, Obs. 8 = mollis.
neglectus, Nym. Syll. 419 = racemosus.
nemoralis, Huds. Fl. Angl. ed. II. i. 61 = asper.
nemorosus, Vill. Hist. Pl. Dauph. ii. 117 = asper.
nitidus, Clarke, Trav. ii. 260 = rigidus.
nitidus, Dum. Obs. Gram. Belg. 119 = secalinus.
Notensis, Boiss. Diagn. Ser. II. iv. 141 = squarrosus.
nototropus, *Kuper. Syst. Tianach.* 78.—As. centr.
odoratus, Gouan, Herbor. 11 = erectus.
oreaemis, *Steud. Syn. Pl. Gram.* 319.—Afr. bor.
Orcuttianus, *Vasey, in Coult. Bot. Gaz.* x. (1885) 223.—Am. bor.
ovatus, Cav. Ic. vi. 67 = Trisetum ovatum.
ovatus, J. Gaertn. in Nov. Comm. Acad. Petrop. xiv. (1770) 587. t. 19. f. 1 = secalinus.
ovinus, Scop. Fl. Carn. ed. II. i. 77 = Festuca ovina.
oxyodon, Schrenk, Enum. Pl. Nov. ii. 1 = macrostachys.
oxyphlorus, Payne, Palest. Expl. Rep. (1874) 128 = macrostachys.
pallens, *Cav. Ic.* vi. 68. t. 591. f. 1.—Ins. Philipp.
pannonicus, *Kummer & Sendt. in Flora*, xxxii. (1849) 757.—Hungar.
parodianus, Presl, Fl. Sicul. i. p. xlv, in nota = ponticus.
parviflorus, *Desf. Cat. Hort. Par.* ed. III. 367.—Hab. ?
patagonicus, *Phil. in Anal. Univ. Chil.* (1873) 577.—Chili.
patentissimus, Weigel, Obs. 13 = Festuca sylvatica.
patulus, *Mert. & Koch. in Roehl. Deutschl. Fl.* i. 635.—Europ.; Orient.; Afghanist.
pauciflorus, *Schous. in Fl. Dan.* t. 1888.—Island.
pauciflorus, Willd. ex Steud. Nom. ed. II. i. 228 = Festuca mollis.
pauper, J. F. Gmel. Syst. 191 = Brachypodium distachyum.
pauper, Schrank, Beier. Fl. i. 370 = Brachypodium sylvaticum.
pectinatus, *Thunb. Prod. Pl. Cap.* 1. 22.—Afr. austr.

BROMUS :—
pendulinus, Desf. ex Steud. Nom. ed. I. 119, 120 = commutatus.
pendulinus, Hort. ex Schrad. N. Journ. iv. (1810) 70 = ciliatus.
pendulus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 90 = patulus.
pentastachyus, Tineo, Pl. Rar. Sic. Pugill. i. 4 = Brachypodium distachyum.
perennis, Vill. Hist. Pl. Dauph. ii. 122 = erectus.
persicus, Boiss. Diagn. Ser. I. xiii. 66; Ledeb. Fl. Ross. iv. 357 = tomentosus.
Petitianus, *A. Rich. Tent. Fl. Abyss.* ii. 488.—Abyss.
phleoides, L. F. Gmel. Syst. 190 = Koeleria phleoides.
phleoicoides, Steud. Nom. ed. I. 120 = Brachypodium pinnatum.
Phoenix, *Y. F. Gmel. Syst.* 190.—Hab.?
phrygius, Boiss. Diagn. Ser. II. iv. 140 = patulus.
pictus, *Hook. f. Fl. Antarct.* 387.—Chili.
pilosus, *F. G. Dietr. Vollst. Lex. Gaertn.* i. 558. —Europ. austr.
pilosus, Willd. in Spreng. Syst. i. 358 = trichodes.
pinnatus, Linn. Sp. Fl. 78 = Brachypodium pinnatum.
pinnatus, Ocd. in Fl. Dan. t. 164 = Brachypodium sylvaticum.
pitensis, *H. B. & K. Nov. Gen. et Sp.* i. 152.— Ecuador.
platystachys, Desf. Cat. Hort. Par. ed. III. 387 = arduennensis.
platystachyus, Lam. Illustr. i. 195 = Brachypodium distachyum.
Plachetzkii, All. Fl. Pedem. ii. 250 = Brachypodium ramosum.
poaeformis, *Forsk. Fl. Aegypt. Arab.* 28.—Aegypt.
poaeformis, Spreng. Neehr. i. Bot. Gart. Halle, 15; Fl. Hal. Mant. 84 = Diplachne fascicularis.
poaeformis, *Vahl, ex Hornem. Hort. Hafn.* i. 102.— Hab.?
polymorphus, Hohen. Enum. Talusch. 19 = patulus.
polymorphus, Hort. ex Steud. Nom. ed. II. i. 228 = arvensis.
polymorphus, Scop. Fl. Carn. ed. II. i. 79; Huds. Fl. Angl. ed. II. i. 48 = mollis, racemosus, secalinus.
polystachios, Forsk. Fl. Aegypt. Arab. 28 = Diplachne fusca.
polystachyus, DC. Fl. Fr. vi. 276 = madritensis.
polystachyus, Kunth, Enum. Pl. i. 447 = Brachypodium distachyum, Triticum ciliatum.
polystachyus, Steud. Nom. ed. II. i. 228 = arduennensis.
pratensis, Ehrh. Beitr. vi. 84, nomen = racemosus.
pratensis, Lam. Encyc. i. 468 = erectus.
pratensis, Spreng. Syst. i. 359 = Festuca elatior.
Presli, *Kunth, Enum. Pl.* i. 416.—Peruv.
procerus, H. B. & K. Nov. Gen. et Sp. i. 150 = Festuca eminens.
procerus, Spreng. Syst. i. 357 = quitensis.
propendens, Jord. ex Nym. Consp. 821 = rigidus.
pseudoarvensis, Koel. Gram. 241 = erectus.
pseudoasper, Schur, Enum. Pl. Transs. 804 = asper.
pseudoinermis, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 90.—Transsylv.
pubescens, C. Koch, in Linnaea, xxi. (1848) 490 = variegatus.
pubescens, Muhl. in Willd. Enum. Hort. Berol. 120 = ciliatus.
pubescens, [Pierl,] in Ionios Anthol. iii. (1884) 680 = squarrosus.
palchellus, *Fig. & De Not. Agr. Fragm.* 16 (sp. dub.). —Arabia.
purgans, Linn. Sp. Fl. 78 = ciliatus.
purgans, Torr. Fl. N. York. ii. 468 = Kalmii.
purpurascens, Delile, Fl. Aeg. Illustr. 53 = rubens.
purpurascens, Turcz. ex Ledeb. Fl. Ross. iv. 357 = inermis.
quitensis, *Willd. ex Steud. Nom.* ed. II. i. 228.— Ecuador.
racemiferus, *Steud. Syn. Pl. Gram.* 328.—Japon.
racemosus, *Linn. Sp. Pl.* ed. II. 114.—Europ.
ramosus, Linn. Mant. i. 34 = Brachypodium ramosum.
ramosus, Huds. Fl. Angl. 40 = asper.
ramosus, Schrank. Verz. Marok. 44 = Brachypodium pinnatum.
repens, Boeh. ex Nym. Consp. 823 = vernalis.
Requieni, Lois. Fl. Gall. i. 90 = intermedius.
rethaeus, Pers. Syn. i. 96 = Brachypodium ramosum.
Richardsoni, *Link, Hort. Berol.* ii. 281.—Am. bor.
rigens, Linn. Mant. 28 = rigidus.

BROMUS :—
rigens, Vandelli, in Roem.-Script. Pl. Hisp. 49 = Koeleria phleoides.
rigidus, C. A. Mey. Vers. Pfl. Casc. 28 = rigidus.
rigidus, Reichb. Ic. Fl. Germ. i. 31, n. 1586 = rubens.
rigidus, *Roth, in Roem. & Ust. Mag. Bot.* iv. (1790) 21.—Europ.; Reg. Mediterr. et Caucas.
riparius, *Rehm. Notiz. Vog. Gest. Schwarz. Meer.* 8.— Rossia.
Rochelianus, Opiz, Natural. n. 9 (1825) 119 = madritensis.
rotundatus, H. B. & K. Nov. Gen. et Sp. i. 152 = Brizа rotundata.
rubens, Cav. Ic. i. t. 45 = scoparius.
rubens, Delile, Fl. Aeg. Illustr. 52 = fasciculatus.
rubens, Desv. ex Willk. & Lange, Prod. Fl. Hisp. i. 98 = madritensis.
rubens, Host. Gram. Austriac. i. t. 18 = rigidus.
rubens, *Linn. Cent.* i. 5.—Reg. Mediterr.
rupestris, Host, Gram. Austriac. iv. t. 17 = Brachypodium pinnatum.
sabulosus, *Guss. Fl. Sic. Prod.* i. 120.—Sicilia.
sardous, *Spreng. Syst. iv. Cur. Post.* 36.—Sardin.
scabrerrimus, Tenore, Fl. Nap. iii. 89 = madritensis.
scabratus, *Link, in Linnaea*, xvii. (1843) 402.—Italia.
scabridus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 281.—Afr. trop.
scabriflorus, Opiz, Natural. n. 9 (1825) 119 = tectorum.
Scheuchzeri, C. C. Gmel. Fl. Bad. iv. 79. t. 7 = Festuca gigantea.
Schraderi, Kunth, Enum. Pl. i. 416 = unioloides.
sclerophyllus, *Boiss. Diagn.* Ser. I. xiii 65.—Lydia.
scoparius, Guss. Fl. Sic. Prod. 10 = alopecuroides.
scoparius, *Linn. Cent Pl* i. 6; *Amoen. Acad.* iv. 266. —Reg. Mediterr.; As. Min.; Persia.
secalinus, Bieb. Fl. Taur. Cauc. i. 71 = patulus.
secalinus, *Linn. Sp. Pl.* 76.—Europ.; Reg. Mediterr.; As. bor.
secalinus, Savi, Fl. Pis. i. 120 = arvensis.
secundus, *Y. Presl, in Rel. Haenk.* i. 263.—Sinus Nootk.
segetalis, A. Braun, ex Nym. Consp. 822 = secalinus.
segetum, Chaub. & Schlecht. in Linnaea, vi. (1831) 38 = ciliatus.
segetum, *H. B. & K. Nov. Gen. et Sp.* i. 152.— Am. austr.
sericeus, Tenore, Fl. Nap. Prod. 10 = alopecuroides.
serotinus, Beneken, in Bot. Zeit. iii. (1845) 724 = asper.
setaceus, Buckl. in Proc. Acad. Sc. Philad. '1863' (1863) 98 = sterilis.
setifolius, *Y. Presl, in Rel. Haenk.* i. 261.—Chili; Abyss.
Sewerzowi, *Regel, in Act. Hort. Petrop.* vii. (1880) 601.—As. centr.
silvaticus, J. F. Gmel. Syst. 190 = sylvaticus.
simplex, Caud. Agrost. Helv. i. 296 = racemosus.
simplex, *Pers. Syn.* i. 96.—Hab.?
simplicissimus, Cesati, ex Nym. Consp. 823 = mollis.
sipyleus, *Boiss. Diagn.* Ser. I. xiii. 65.—As. Min.
sitchensis, *Bongard, Veget. Ins. Sitcha*, 178.—Ins. Sitcha.
speciosus, *Nees, Fl. Afr. Austr.* i. 454.—Afr. austr.
spicatus, Nees, in Mart. Fl. Bras. ii. 471 = Diplachne simplex.
spiculiformata, Knapp, Brit. Gram. t. 81 = arvensis.
squarrosus, Brot. Fl. Lusit. i. 112 = macrostachys.
squarrosus, *Linn. Sp. Pl.* 76.—Europ. austr.; Orient; As. bor.
squarrosus-muticus, C. A. Mey. Vers. Pfl. Cauc. 24 = brisiformis.
staphyoides, *Link, Hort. Berol.* ii. 285.—Hab.?
staminea, E. Desv. in C. Gay, Fl. Chil. vi. 440 = unioloides.
stenophyllus, Link, Enum. Hort. Berol. i. 94 = erectus.
stenostachyus, *Boiss. Fl. Orient.* v. 648.—Afghanist.
sterilis, Gaus. Fl. Sic. Prod. i. 116 = madritensis.
sterilis, Lang, ex Kumm. & Sendt. in Flora, xxxiii. (1849) 758 = tectorum.
sterilis, *Linn. Sp. Pl.* 77.—Europ.; Orient; As. bor.
Steudelii, Frank, ex Steud. Nom. ed. II. i. 229 = ciliatus.
stipoides, Linn. Mant. iii. 567 = Festuca geniculata.
striatus, Brongn. in Duperr. Voy. 45 = Brongniartii.
striatus, *Kunth, Rev. Gram.* i. 134.—Ins. Borbon.
strictus, Scop. Fl. Carn. ed. II. i. 79 = Diplachne serotina.
strictus, Hall. f. in Roem. Arch. i. ii. 10 = Brachypodium pinnatum?

BROMUS :—
strigosus, Bieb. Fl. Taur. Cauc. iii. 81 = Triticum strigosum.
strigosus, Lam. Illustr. 191. n. 1068 = Festuca gigantea.
subalpinus, Rupr. in Bull. Acad. Brux. ix. (1842) 2, 286 = exaltatus.
submuticus, *Steud. Syn. Pl. Gram.* 321.—Missouri.
subulatus, Griseb. in Ledeb. Fl. Ross. iv. 358 = Melica acuminata.
Sukedorfi, *Vasey, in Coult. Bot. Gaz.* x. (1885) 228.— Am. bor.
sylvaticus, Pollich, Hist. Pl. Palat. i. 118 = Brachypodium sylvaticum.
sylvaticus, Vogler, Schediasma, 18 = asper.
syriacus, Boiss. & Blanche, Diagn. Ser. II. iv. 139 = erectus.
Tacna, *Steud. in Lechl. Berb. Am. Austr.* 56.— Peruv.
subrectus, Fries, Mant. iii. 131 = erectus.
tectorius, Dulac, Fl. Hautes-Pyr. 96 = tectorum.
tectorum, *Linn. Sp. Pl.* 77.—Europ.; Orient; As. bor.
temulentus, Bernh. Syst. Verz. Erf. 47 = Lolium temulentum.
Tenorianus, Roem. & Schult. Syst. Mant. iii. 860, obs. = sterilis.
tenuiflorus, Viv. Fl. Lyb. Spec. 5. t. 1. f. 1 = fasciculatus.
tenuis, *Y. Presl, in Rel. Haenk.* i. 262.—Peruv.
tenuis, Tineo, Rar. Pl. Sic. Pugill. i. 3 = Festuca incrassata.
Thominii, Ard. ex Nym. Consp. 823 = mollis.
Tigridis, Boiss. & Noë, Diagn. Ser. II. iv. 141 = brachystachys.
tigurinus, Suter, Fl. Helv. i. 65 = Brachypodium pinnatum.
tomentellus, *Boiss. Diagn.* Ser. I. vii. 126.—Graecia; As. Min.; Persia.
tomentosus, Roxb. in Schrad. N. Journ. iii. (1809) 42 = macrostachys?
tomentosus, *Trin. in Mém. Acad. Pétersb.* (1818-14) t. 9.—Persia.
tortilis, *Presl, ex Steud. Nom.* ed. II. i. 229.—Europ. austr.
transsilvanicus, Schur, in Oestr. Bot. Zeitschr. x. (1860) 227 = fibrosus.
tranquebaricus, Steud. Syn. Pl. Gram. 320 = prsec.
triaristatus, Loisel. Fl. Gall. i. 89 = arduennensis.
trichodes, *Kunth, Rev. Gram.* i. 135.—Ins. Mascar.
trichopodes, *A. Rich. Tent. Fl. Abyss.* ii. 457.— Abyss.
trifforus, Ehrh. Beitr. i. 90 = Festuca sylvatica.
trifforus, Linn. Sp. Pl. ed. II. 115 = Festuca gigantea.
trifforus, Pollich, Hist. Pl. Palat. i. 119 = Ventenata avenacea.
Trinianus, *Schult. Mant.* ii. 254.—Persia.
Trinii, *E. Desv. in C. Gay, Fl. Chil.* vi. 441.— Chili.
tristachyus, *Steud. ex Steud. Nom.* ed. II. i. 229, nomen. —Hab.?
triticeus, *Raeusch. Nom.* ed. III. 27.—Helvet.
trivialis, Savi, Fl. Pis. i. 124 = Koeleria phleoides.
truncatus, Scop. Fl. Carn. ed. II. i. 85 = Agropyron junceum.
tunicatus, *Phil. in Linnaea*, xxxiii. (1864-65) 298.— Chili.
turgidus, Pers. Syn. i. 95 = macrostachys.
unilateralis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 90 = patulus.
unioloides, *H. B. & K. Nov. Gen. et Sp.* i. 151.— Am. calid.
valdivianus, *Phil. in Linnaea*, xxix. (1857-58) 101.— Chili.
variegatus, Fl. Danub. ex Boiss. Fl. Orient. v. 645 = fibrosus.
variegatus, *Bieb. Fl. Taur. Cauc.* iii. 79.—Reg. Caucas.; Persia.
vernus, Brot. Fl. Lusit. i. 113 = rigidus.
velutinus, Schrad. Fl. Germ. i. 349. t. 6. f. 3 = secalinus.
Ventolense, Hort. ex Steud. Nom. ed. II. i. 229 = racemosus.
Ventolensa, Schleich. ex Steud. l. c. = arvensis.
vernalis, *Pancl. ex Hach. in Oestr. Bot. Zeitschr.* xxix. (1879) 208.—Europ.
versicolor, Pollich, Hist. Pl. Palat. i. 190 = arvensis.
verticillatus, *Cav. Ic.* vi. 66. t. 590.—Hispan.

BROMUS:—
vestitus, *Schrad. in Goett. Gel. Anz.* (1821) iii. 2074. —Afr. austr.
willdenowii, Steud. Syn. Pl. Gram. 236 = japonicus.
villosulus, *Steud. l. c.* 257.—Graecia.
villosus, Forsk. Fl. Aegypt. Arab. 23 = maditensis, rigidus.
villosus, C. C. Gmel. Fl. Bad. i. 229 = squarrosus.
villosus, Scop. Fl. Carn. ed. II. i. 83 = Agropyron repens.
virens, *Buchl. in Proc. Acad. Sc. Philad.* '1862' (1863) 98.—Am. bor. occ.
virgatus, *J. Presl, in Rel. Haenk.* i. 263.—Peruv.; Mexic.
vitiosus, Weigel, Obs. 4 = secalinus.
Willdenowii, Kunth, Rev. Gram. i. 184 = unioloides.
wulfenii, Willd. Enum. Hort. Berol. Suppl. 9; Jacq. Ecl. Gram. 20. t. 15 = squarrosus.

BROMUS, Scop. Introd. 74 (1777) = **Tritticum**, Linn. (Gramin.).

BRONGNIARTIA, Blume, Bijdr. 485 (1825) = **Kibara**, Endl. (Monim.).
corsacea, Blume, l. c. = K. coriacea.

BRONGNIARTIA, H. B. & K. Nov. Gen. et Sp. vi. 465. t. 587 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 493.
MEGASTEGIA, G. Don, Gen. Syst. ii. 468 (1832).
PERALTEA, H. B. & K. Nov. Gen. et Sp. vi. 469. t. 589 (1823).
Benthamiana, *Hemsl. Biol. Centr. Am. Bot.* i. 252. —Mexic.
foliolosa, *Benth. ex Hemsl. Diag. Pl. Nov.* 7.—Mexic.
galegoides, *Presl, Symb. Bot.* ii. 21. t. 67.—Am. trop.
glabrata, *Hook. & Arn. Bot. Beech. Voy.* 238.—Mexic.
gracilis, *Hemsl. Biol. Centr. Am. Bot.* i. 253.—Mexic.
intermedia, *Moric. in Mém. Soc. Phys. Genèv.* vii. (1836) 253.—Mexic.
mexicana, *Schlecht. in Linnaea*, xii. (1838) 338. —Mexic.
minutifolia, *S. Wats. in Proc. Am. Acad.* xx. (1885) 366.—Texas.
mollis, *H. B. & K. Nov. Gen. et Sp.* vi. 465. t. 587.—Mexic.
obliqua, *Schlecht. in Linnaea*, xii. (1838) 339.—Mexic.
oligosperma, *Baill. Adansonia*, ix. (1868–70) 240.—Mexic.
oxyphylla, *Hemsl. Biol. Centr. Am. Bot.* i. 254.—Mexic.
Parryi, *Hemsl. l. c.*—Mexic.
pedalyrioides, *H. B. & K. Nov. Gen. et Sp.* vi. 465. t. 588.—Mexic.
retusa, *Benth. in Hemsl. Diag. Pl. Nov.* 8.—Mexic.
robinioides, *Kunth, Ind. Sem. Hort. Berol.* (1845) 12.—Chili.
sericea, *Schlecht. in Linnaea*, xii. (1838) 338.—Mexic.
stipitata, *Hemsl. Diag. Pl. Nov.* 8.—Mexic.
thermoides, *Sprang. Syst.* iv. *Cur. Post.* 287.—Mexic.
vicioides, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 49.—Mexic.

BRONNIA, H. B. & K. Nov. Gen. et Sp. vi. 53. t. 528 (1823) = Fouquieria, H. B. & K. (Tamaric.).
spinosa, H. B. & K. l. c. = F. spinosa.

BROOKEA, Benth. in Benth. & Hook. f. Gen. ii. 989 (1876). *SCROPHULARINEAE*, Benth. & Hook. f. l. c.
dasyantha, *Benth. in Hook. Ic. Pl.* xii. 84. t. 1197.—Borneo.
tomentosa, *Benth. l. c.*—Borneo.

BROSIMUM, Sw. Prod. Veg. Ind. Occ. 12 (1788). *URTICACEAE*, Benth. & Hook. f. iii. 360.
GALACTODENDRUM, Humb. Relat. Hist. ii. 108 (1819).
PERAVINERA, Aubl. Pl. Gui. ii. 888. t. 340 (1775), nomen prius.
Alicastrum, *Sw. l. c.*—Am. trop.

BROSIMUM:—
Aublettii, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 54. t. 148. f. a.—Am. trop.
costaricanum, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 234.—Costa Rica.
discolor, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 408.—Bras.
echinocarpum, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 54. t. 148.—Peruv.
Galactodendron, *D. Don, ex Sweet, Hort. Brit.* ed. II. 462.—Guiana.
Gaudichaudii, *Trécul, in Ann. Sc. Nat.* Sér. III. viii. (1847) 139.—Bras.
Humboldtii, Carr. in Rev. Hortic. (1874) 314. f. 49, 48 = B. Galactodendron.
microcarpum, Schott, in Spreng. Syst. iv. Cur. Post. 408 = B. Alicastrum.
refractum, Mart. ex Miq. in Mart. Fl. Bras. iv. i. 108 = Gaudichaudii.
spurium, *Sw. Prod. Veg. Ind. Occ.* 12.—Jamaica.
turbinatum, *Spruce, ex Baill. Adansonia*, xi. (1875) 299.—Bras.

BROSSAEA, [Plum.] Linn. Gen. ed. I. 366 (1737) = Gaultheria, Kalm. (Ericac.).
anastomosans, Griseb. Fl. Brit. W. Ind. 142 = G. anastomosans.
coccinea, Linn. Sp. Pl. 1190 = G. cordifolia.

BROSSARDIA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 183. *CRUCIFERAE*, Benth. & Hook. f. i. 91.
pygmaea, *Boiss. l. c.* 184.—Persia.
retusa, *Boiss. Fl. Orient.* i. 886.—Persia.

BROTERA, Cav. in Anal. Cienc. Nat. i. (1800) 32; Ic. v. 51. t. 483. (1799) = Malbantia, Forsk. (Sterculiac.).
bractosa, Guill. & Perr. Fl. Seneg. Tent. i. 86 = M. Denhami.
Leprieurii, Guill. & Perr. l. c. = Pentapetes Leprieurii.
ovata, Cav. Ic. v. 20 = M. abyssinica.

BROTERA, Spreng. in Schrad. Journ. Bot. iv. (1800) t. 5 = Flavaria, Juss. (Compos.).
chilensis, Spreng. l. c. = F. repanda.
Sprengelii, Cass. in Dict. Sc. Nat. xxxiv. 306 = F. repanda.
trinervata, Pers. Syn. ii. 498 = F. repanda.

BROTERA, Spreng. Nachr. i. Bot. Gart. Halle, (1801) 15; et in Trans. Linn. Soc. vi. (1802) 151. t. 12 = Lygtis, Jacq. (Labiat.).
peraica, Spreng. ll. cc. = H. pectinata.

BROTERA, Vell. Fl. Flum. 323 (1825); vii. t. 168 (1827) = Lychnophora, Mart. (Tulisc.).
martiana, Vell. ll. cc. 323; t. 168 = L. speciosa.
mediterranea, Vell. ll. cc. 323; t. 164 = L. divaricata.

BROTERA, Willd. Sp. Pl. iii. 3299 (1803) = Cardopatium, Juss. (Compos.).
corymbosa, Willd. l. c. = C. corymbosum.

BROTEROA, DC. Prod. v. 635 (1836) = Flaveria, Juss. (Compos.).
trinervata, DC. l. c. 636 = F. repanda.

BROUGHTONIA, R. Br. in Ait. Hort. Kew. ed. II. v. 217 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 531.
alba, Spreng. Syst. iii. 735 = Maxillaria alba.
aurea, Lindl. Bot. Reg. (1840) Misc. 19 = Epidendrum xanthinum.
chinensis, Lindl. in Hook. Lond. Journ. Bot. i. (1842) 492 = Epidendrum nonchinense.
coccinea, Hook. Bot. Mag. t. 3036 = sanguinea.
grandiflora, Spreng. Syst. iii. 735 = Maxillaria grandiflora.
Hiachne, *Hensfr. in Moore & Ayres, Gard. Mag. Bot.* iii. (1851) 201.—ind. occ.?
maculata, Spreng. Syst. iii. 735 = Maxillaria maculata.
sanguinea, *R. Br. in Ait. Hort. Kew.* ed. II. v. 217.—Ind. occ.
tetragona, Spreng. Syst. iii. 734 = Phajus tetragonus.
violacea, Hort. ex Moore & Ayres, Gard. Mag. Bot. iii. (1851) 201 = Hiachne.

BROUGHTONIA, Wall. ex Lindl. Gen. et Sp. Orch. 35 (1830) = Otochilus, Lindl. (Orchid.).
amoena, Wall. l. c. = O. fusca.
nitida, Herb. ex Sweet, Hort. Brit. ed. III. 641, nomen (= Coelogynes sp. ?).—Hab. ?
linearis, Wall. l. c. 42 = Coelogyne ovalis.
pendula, Wall. l. c. 36 = O. alba.
pilosa, Hook. ex Steud. Nom. ed. II. i. 230.—Ind. or.

BROUSSAISIA, Gaudich. Freyc. Voy. Bot. 479. t. 69 (1826). *SAXIFRAGEAE*, Benth. & Hook. f. i. 641
arguta, *Gaudich. l. c.*—Ins. Sandvic.
pellucida, *Gaudich. Bot. Voy. Bonite*, t. 9.—Ins. Sandvic.

BROUSSONETIA, Orteg. Hort. Matr. 61. t. 7 (1798) = Sophora, Linn. (Legumin.).
secundiflora, Orteg. l. c. = S. secundiflora.
speciosa, Orteg. l. c. t. 7 = S. secundiflora.

BROUSSONETIA, L'Hérit. ex Vent. Tabl. Regn Veget. iii. 547 (1799). *URTICACEAE*, Benth. & Hook. f. iii. 361.
PAPYRIUS, Lam. Illustr. t. 769 (1798).
STENOCHASMA, Miq. Fl. Jungh. i. 45 (1850).
Billardii, Carr. in Rev. Hortic. (1878) 375 = papyrifera.
brasiliensis, Mart. in Flora, xxiv. (1841) Beibl. 10 = Maclura brasiliensis.
cordata, *Blume, Mus. Bot. Lugd. Bat.* ii. 87.—China
cordata, Hort. ex C. Koch, Dendrol. ii. ii. 440 = papyrifera.
cucullata, Hort. ex Steud. Nom. ed. II. i. 230 = papyrifera.
disseeta, Hort. ex Bur. in DC. Prod. xvii. 225 = papyrifera.
elegans, Hort. ex C. Koch, Dendrol. ii. ii. 440 = papyrifera.
integrifolia, *Buch.-Ham. in Trans. Linn. Soc.* xvii. (1837) 195.—Ind. or.
Kaempferi, *Siebold, in Verh. Bat. Gen. zizi.* (1830) 28 —Japon.
Kaempferi, Siebold, ex Blume, Mus. Bot. Lugd. Bat. ii. 86 = papyrifera.
Kaempferi, Siebold, *in Verh. Bat. Gen.* xii. (1830) 28.—Japon.
laxonica, *Bur. in DC. Prod.* xvii. 224.—Ins. Philipp.
luzoniensis, Blanco, Fl. Filip. ed. II. 488 = Allaeanthus luzonicus.
maculata, Hort. ex Steud. Nom. ed. II. i. 230 = papyrifera.
mexicana, *Hance, in Journ. Bot.* xx. (1882) 294.—China.
novicularis, Lodd. ex C. Koch, Dendrol. ii. ii. 440 = papyrifera.
nana, Hort. ex Bur. in DC. Prod. xvii. 225 = papyrifera.
novicularis, Lodd. ex Bur. l. c. = papyrifera.
novifolia, Hort. ex Steud. Nom. ed. II. i. 230 = papyrifera.
papyrifera, *Vent. Tabl. Regn. Veget.* iii. 547.—Malaya; Ins. Pacific.
Pinnerii, Spreng. Syst. iii. 901 = Chlorophora tinctoria.
Siebaldii, Blume, Mus. Bot. Lugd. Bat. ii. 86 = B. Kaempferi.
spathulata, Hort. ex Steud. Nom. ed. II. i. 230 = papyrifera.
tinctoria, Blanco, Fl. Filip. ed. II. 488 = Cudrania javanensis.
tinctoria, H. B. & K. Nov. Gen. et Sp. ii. 32 = Chlorophora tinctoria.
tinctoria, Spreng. Syst. iii. 901 = Maclura tinctoria.
tricolor, Hort. ex C. Koch, Dendrol. ii. ii. 440 = papyrifera.
xanthoxylon, Mart. in Flora, xxiv. (1841) Beibl. 10 = Maclura xanthoxylon.

BROUVALEA, Adans. Fam. ii. 311 (1763) = seq.

BROVALLIA, Linn. Orb. Nat. (1764) = seq.

BROWALLIA, Linn. Gen. ed. I. 246 (1737). *SOLANACEAE*, Benth. & Hook. f. ii. 910.
abbreviata, *Benth. Bot. Voy. Sulph.* 144.—Peruv.
alienata, Linn. Syst. ed. X. 1117 = Roellia pentlandii.
americana, Linn. Sp. Pl. 631 = demissa.
cordata, *G. Don, Gen. Syst.* iv. 478.—Peruv.

BROWALLIA:—
demissa, *Linn. Syst.* ed. X. 1116.—Am. austr.
elata, Linn. l. c.=demissa.
elongata, H. B. & K. Nov. Gen. et Sp. ii. 373=demissa.
grandiflora, *R. Grah. in Edin. N. Phil. Journ.* 'Jan.-Apl. 1830' (1829).—Peruv.
grandiflora, Lindl. Bot. Reg. t. 1884=cordata.
humifusa, Forsk. Fl. Aegypt. Arab. 112=Striga humifusa.
Jamesonii, *Benth. in DC. Prod.* x. 197.—Peruv.; N. Granat.
lactea, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 508=demissa.
Linnaeana, Spreng. Syst. ii. 808=demissa.
nervosa, Miers, in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 306=demissa.
peduncularis, *Benth. Bot. Voy. Sulph.* 148.—Peruv.
pulchella, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 176=viscosa.
pulchella, *Lehm. in Hamb. Gartens.* x. (1854) 375.—Hab.?
speciosa, *Hook. Bot. Mag.* t. 4839.—N. Granat.
tenella, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. v. (1850) 208.—Bras.
tenuna, Torr. Bot. Mex. Bound. 156=Nierembergia viscosa.
viscosa, *H. B. & K. Nov. Gen. et Sp.* ii. 373.—Am. austr.

BROWNEA, Jacq. Enum. Pl. Carib. 6 (1760) (Brownaea). *LEGUMINOSAE*, Benth. & Hook. f. i. 577.
HERMESIAS, Loefl. Iter, 278 (1758).
antioquensis, Linden, Cat. (1859) 3=macrophylla.
Arisa, *Benth. Pl. Hartw.* 171.—N. Granat.
Arisa, Lindl. & Paxt. Flow. Gard. ii. 107. t. 59=grandiceps.
Birchellii, *Hook. f. Bot. Mag.* t. 5998.—Venezuela.
capitella, *Jacq. Fragm.* 26. t. 18, 19.—Venezuela.
cauliflora, *Poepp. & Endl. Gen. et Sp. Pl. Chil.* iii. 32.—Chili.
coccinea, *Jacq. Enum. Pl. Carib.* 26.—Venezuela.
coccinea, Loefl. ex Griseb. Fl. Brit. W. Ind. 219=B. Rose-de-monte.
erecta, Hort. Belg. ex Planch. in Fl. des Serres, vi. (1850-51) 154 (=Theophrasta pinnata).
grandiceps, *Jacq. Collect.* iii. 287. t. 22.—Am. austr.
guianensis, *Nietzsch, in Schomb. Faun. & Fl. Guy.* 1209, nomen.—Guiana.
latifolia, *Jacq. Fragm.* 25. t. 17.—Venezuela.
leucantha, *Jacq. l. c.* 26.—Venezuela.
macrophylla, *Linden, Cat.* n. 18 (1868) 11; *Mast. in Gard. Chron.* (1878) 777, f. 149.—N. Granat.
negrensis, *Benth. in Mart. Fl. Bras.* xv. ii. 213.—Bras.
pauciflora, Willd. Sp. Pl. iii. 716=Palowea guianensis.
Princeps, Linden, ex Otto, in Otto & Dietr. Allg. Gartenz. xxiii. (1856) 147=B. Ariza.
racemosa, *Jacq. Fragm.* 25. t. 16.—Venezuela.
Rosa, Pers. Syn. ii. 237=B. Rosa-de-monte.
Rosa-de-monte, *Berg. in Phil. Trans.* (1771) 171. t. 8, 9.—Am. austr.
rosea, Otto, in Otto & Dietr. Allg. Gartenz. xxiii. (1856) 147=B. Rosa-de-monte.
speciosa, Reichb. ex DC. Prod. ii. 477=B. Rosa-de-monte.

BROWNETERA, Rich. in Ann. Mus. Hist. Nat. Par. xvi. (1810) 296=Phyllocladus, Rich. (Conifer.).
asplenifolia, Rich. l. c.=Ph. rhomboidalis.

BROWNLLINIA, Harv. ex Lindl. in Hook. Lond. Journ. Bot. i. (1842) 16. *ORCHIDEAE*, Benth. & Hook. f. iii. 531.
caerulea, *Harv. ex Lindl. l. c.*—Afr. austr.
macrocerus, Sond. in Linnaea, xix. (1847) 106=caerulea.
parviflora, *Harv. ex Lindl. in Hook. Lond. Journ. Bot.* i. (1842) 16.—Afr. austr.
recurvata, *Sond. in Linnaea*, xix. (1847) 107.—Afr. austr.

BROWNLOWIA, Roxb. Fl. Comom. iii. 61. t. 265 (1819). *TILIACEAE*, Benth. & Hook. f. i. 331.
BURMEA, Roxb. Fl. Ind. ii. 640 (1832).
argentea, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 97.—Ind. or.
elata, *Roxb. Fl. Comom.* iii. 61. t. 265.—Ind. or.; Burma.
lanceolata, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 37.—Ind. or.; Burma.
peltata, *Benth. l. c.* 36.—Borneo.

BROWNLOWIA:—
Riedelii, *Hemsl. Bot. Voy. Challenger*, i. 111. 128.—Ins. Moluce.

BRUCEA, J. S. Mill. Fasc. t. 25 (1780). *SIMARUBEAE*, Benth. & Hook. f. i. 311.
GONUS, Lour. Fl. Cochinch. ii. 658 (1790).
NIMIA, Buch.-Ham. ex A. Juss. in Mém. Mus. Par. xii. 516 (1825).
abyssinica, Spreng. Pugill. ii. 90, in obs.=ferruginea.
antidysenterica, Lam. in Mém. Acad. Par. (1784) 342; Encyc. i. 472=ferruginea.
dubia, Steud. Nom. ed. II. i. 330=Picrasma nepalensis.
ferruginea, *L'Hérit. Stirp. Nov.* 19. t. 10.—Abyss.
glabrata, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 447.—Ins. Timor.
gracilis, DC. Prod. ii. 58=sumatrana.
guineensis, G. Don, Gen. Syst. i. 801=paniculata.
macrophylla, *Oliver, Fl. Trop. Afr.* i. 310.—Afr. trop.
mollis, *Wall. Cat.* n. 8483.—Reg. Himal.
paniculata, *Lam. Encyc.* i. 472.—Afr. trop.
quercifolia, *Seem. Fl. Vit.* 38.—Ins. Fiji.
sumatrana, *Roxb. Hort. Beng.* 12; *Fl. Ind.* i. 449.—As. trop.; Austral.
sumatrensis, Spreng. Pugill. ii. 90=sumatrana.
trichotoma, *Spreng. Syst.* i. 441.—Cochinch.
trifoliata, Spreng. l. c. iv. Cur. Post. 48=Evodia simplex.
virginica, *Hornem. ex Steud. Nom.* ed. I. 122, nomen.—Virginia.

BRUCHMANNIA, Nutt. Gen. Am. i. 48 (1818)=Beckmannia, Host (Gramin.).
cruciformis, Nutt. l. c.=Beckmannia cruciaeformis.

BRUCKENTHALIA, Reichb. Fl. Germ. Excurs. 414 (1831). *ERICACEAE*, Benth. & Hook. f. ii. 591.
spiculifolia, *Reichb. l. c.*—Europ. austr. or.; As. Min.

BRUEA, Gandich. in Freyc. Voy. Bot. 211 (1826). *URTICACEAE*? (Gen. dubium) Benth. & Hook. f. iii. 351.
bengalensis, *Gaudich. l. c.* (=Laportea terminalis?).—Ind. or.

BRUECKEA, Klotzsch & Karst. in Karst. Ausw. Gew. Venez. 31. t. 10 (1848)=**Asgiphila**, Jacq. (Verbenac.).
grandifolia, Klotzsch & Karst. l. c. 32=A. verrucosa.

BRUENNICHIA, Willd. Sp. Pl. ii. 751 (1800)=Brunnichia, Banks (Polygon.).

BRUGMANSIA, Blume, in Van Hall, Bijdr. Nat. Wet. ii. (1828) 422. *CYTINACEAE*, Benth. & Hook. f. ii. 119.
MYCETANTHE, Reichb. Nom. 61 (1841).
ZIPPELIA, Reichb. ex Endl. Gen. Suppl. ii. 8 (1842).
Leschenaultii, Decne. in Arti Soc. Ital. Sc. Nat. xi. (1868) 198=Zippelii.
Zippelii, *Blume, in Van Hall, Bijdr. Nat. Wet.* ii. (1828) 422.—Java.

BRUGMANSIA, Pers. Syn. i. 216 (1805)=Datura, Linn. (Solanac.).
arborea, Steud. Nom. ed. II. i. 330=D. arborea.
aurantiaca, Hort. ex Walp. Rep. iii. 934=Juanulloa parasitica.
bicolor, Pers. Syn. i. 216=D. sanguinea.
candida, Pers. l. c.=D. arborea.
floribunda, Paxt. Mag. Bot. ix. 8=Juanulloa aurantiaca.
Knightii, Hort. ex Dun. in DC. Prod. xiii. i. 690=D. sanguinea.
parviflora, Hort. ex Walp. Rep. iii. 934=Juanulloa parasitica.
sanguinea, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. t. 272=D. sanguinea.
suaveolens, Bercht. & Presl, Rostl. i. Solanac. 45=D. suaveolens.
Waymanni, Paxt. Mag. Bot. iv. 241=D. fastuosa.

BRUGUIERA, Lam. Encyc. iv. 626 (1796). *RHIZOPHORACEAE*, Benth. & Hook. f. i. 679.
KANDELIA, Blume, Mus. Bot. Lugd. Bat. i. 140 (1849).
PALETUVIERA, Thou. ex DC. Prod. iii. 22 (1828).
Arnottiana, Wight, ex Arn. in Ann. Nat. Hist. i. (1838) 364=Ceriops Candolleana.
australis, A. Cunn. ex Arn. l. c. 366=Rhizophora Candel, Steud. Nom. ed. II. i. 331=Kandelia Rheedii.

BRUGUIERA:—
caseosa, Blume, Mus. Bot. Lugd. Bat. i. 137=gymnorrhiza.
caryophylleoides, *Blume, Enum. Pl. Jav.* i. 93.—Ind. or.; Malaya.
cylindrica, Blume, l. c. i. 91=caryophylleoides, malabarica.
decanginata, Griff. Notul. iv. 669=Rhizophora Mangle.
eriopetala, *Wight & Arn. in Ann. Nat. Hist.* i. (1838) 368.—As. trop.
gymnorrhiza, *Lam. Illustr.* t. 397; *Blume, Mus. Bot. Lugd. Bat.* i. 136.—Geront. trop.
malabarica, *Arn. in Ann. Nat. Hist.* i. (1838) 369.—Ind. or.
nemorosa, Blanco, Fl. Filip. ed. II. 375=Carallia integerrima.
nudicaula, Pritz. Ic. Ind. ii. 51, sphalm.=Bougueria nubicola.
obtusa, Steud. Nom. ed. II. i. 331 (=*Problastes cuneifolia*).—Java.
oxyphylla, *Miq. Fl. Ind. Bat. Suppl.* i. 824.—Sumatra.
parviflora, *Wight & Arn. Prod.* 311.—Ind. or.; Malaya.
Rheedii, *Blume, Enum. Pl. Jav.* i. 93.—Geront. trop.
Rumphii, Blume, Mus. Bot. Lugd. Bat. i. 138=Rheedii.
sexangula, Poir. Encyc. Suppl. iv. 362=Rhizophora sexangula.
sexangularis, Spreng. Syst. ii. 602=Rhizophora sexangula.
timoriensis, Wight & Arn. Prod. 311=Ceriops Candolleana.
Wightii, Blume, Mus. Bot. Lugd. Bat. i. 138=gymnorrhiza.
Zippelii, Blume, l. c. 139=Rheedii.

BRUGUIERA, Thou. Gen. Nov. Madag. 21 (1806)=Lumnitzera, Willd. (Combret.).
littorea, Steud. Nom. ed. II. i. 331=L. coccinea.
madagascariensis, DC. Prod. iii. 23=L. racemosa.

BRUGUIERA, Rich. ex DC. Prod. iii. 178, in syn. (1828)=Conostegia, D. Don (Melastom.).

BRUINSMANIA, Miq. in Linnaea, xvii. (1843) 72=Isertia, Schreb. (Rubiac.).
tortilobata, Miq. l. c. 78.—Guiana.

BRUNELLA, Tourn. ex Linn. Syst. ed. I (1735)=Prunella, Linn. (Labiat.).

BRUNELLIA, Ruiz & Pav. Prod. Fl. Per. 71. t. 12 (1794). *SIMARUBEAE*, Benth. & Hook. f. i. 313.
acuminata, *Ruiz & Pav. Fl. Per.* iv. 71. t. 413; *Syst.* 127.—Peruv.
acutangula, *Humb. & Bonpl. Pl. Aequin.* i. 216. t. 63.—N. Granat.
comocladifolia, *Humb. & Bonpl. l. c.* i. 211. t. 59.—Peruv.
Funkiana, *Tul. in Ann. Sc. Nat. Sér.* III. vii. (1847) 267.—N. Granat.
Goudotii, *Tul. l. c.* 270.—N. Granat.
inermis, *Ruiz & Pav. Fl. Per.* iv. 71. t. 414; *Syst.* 127.—Peruv.
ovalifolia, *Humb. & Bonpl. Pl. Aequin.* i. 216. t. 61.—Peruv.
propinqua, *H. B. & K. Nov. Gen. et Sp.* vii. 45.—Am. austr.
quadriloccularis, *Hook. & Arn. Bot. Beech. Voy.* 282.—Mexic.
racemifera, *Tul. in Ann. Sc. Nat. Sér.* III. vii. (1847) 269.—N. Granat.
rugosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 274.—Bras.
sandwicensis, Gandich. Freyc. Voy. Bot. 98; ex Hook. & Arn. Bot. Beech. Voy. 60=Pelea sandwicensis.
tomentosa, *Humb. & Bonpl. Pl. Aequin.* i. 214. t. 60.—Am. austr.

BRUNFELSIA, Plum. ex Linn. Hort. Cliff. 5 (1737) (Brunsfelsia); Gen. ed. I. 332. *SOLANACEAE*, Benth. & Hook. f. ii. 911.
FRANCISCEA, Pohl, Pl. Bras. Ic. i. 1. t. 1—7 (1827).
acuminata, Benth. in DC. Prod. x. 199=ramosissima.
americana, *Linn. Sp. Pl.* 191.—Am. trop.

BRUNFELSIA :—
australis, Benth. in DC. Prod. x. 200 = Hopeana.
bahiensis, *A. DC. Prod.* x. 590.—Am. sentr.
calycina, Benth. in DC. Prod. x. 199 = pauciflora.
capitata, *Benth. l. c.* 198.—Bras.
cestroides, *A. Rich. Fl. Cub. Fanerog.* ii. 151.—Cuba.
confertiflora, Benth. in DC. Prod. x. 199 = ramosissima.
cuneifolia, *J. A. Schmidt, in Mart. Fl. Bras.* viii. 259. —Bras.
eximia, Bosse, Handb. Blumeng. 524 = macrophylla.
falcata, *Regel, Cat. Pl. Hort. Absakov.* 21.—Hab.?
fallax, *Duchass. ex Griseb. in Goett. Abh.* vii. (1857) 249.—Ind. occ.
gracilis, *Regel, Cat. Pl. Hort. Absakov.* 21.—Hab.?
grandiflora, *D. Don, in N. Edin. Phil. Journ.* (Apr.–Oct. 1839) 86 —Peruv.
guianensis, *Benth. in DC. Prod.* x. 200.—Guiana.
Hopeana, *Benth. l. c.*—Bras.
hydrangeaeformis, *Benth. l. c.* 198.—Bras.
inodora, *Mart. Del. Sem. Hort. Monac.* (1847).—*Cf. Linnæa*, xxiv. (1851) 180.—Bras.
jamaicensis, *Griseb. Fl. Brit. W. Ind.* 432.—Ind. occ.
latifolia, *Benth. in DC. Prod.* x. 199.—Bras.
latifolia, Hort. ex Steud. Nom. ed. II. i. 231 = americana.
Lockhartii, *Hort. Belg. ex Heynh. Nom.* ii. 79, nomen. —Ins. Trinitat.!
longiflora, *Regel, Cat. Pl. Hort. Absakov.* 21.—Hab.? longituba, *Lem. Jard. Fleur.* iv. (1854) *Misc.* 61.—Cuba.
macrophylla, *Benth. in DC. Prod.* x. 198.—Bras.
maritima, *Benth. l. c.* 200.—Bras.
montana, *Lodd. ex Loud. Hort. Brit.* 253, nomen.—Ind. occ.
multiflora, *Regel, Cat. Pl. Hort. Absakov.* 21.—Hab.? mutabilis, *Poit. in Rev. Hortic. Sér.* II. i. (1842) 261. —Hab.?
nitida, *Benth. in DC. Prod.* x. 201.—Am. trop.
obovata, *Benth. l. c.* 199.—Bras.
parviflora, A. Rich. Fl. Cub. Fanerog. ii. 237 = seq.
parvifolia, A. Rich. l. c. 151 = nitida.
pauciflora, *Benth. in DC. Prod.* x. 199.—Bras.
purpurea, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 228.—Ind. occ.
ramosissima, *Benth. in DC. Prod.* x. 199.—Bras. Schomburgkiana, *Klotzsch, in Schomb. Fauna et Fl. Guy.* 1158, nomen.—Guiana.
Siebert, *Regel, Cat. Pl. Hort. Absakov.* 21.—Hab.? sinuata, *A. Rich. Fl. Cub. Fanerog.* ii. 151.—Am. trop.
spinosa, *Jacq. Enum. Pl. Carib.* 14.—Ind. occ.
Spruceana, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 258 = maritima.
terminalis, Salisb. Prod. 109 = americana.
undulata, *Sw. Prod. Veg. Ind. Occ.* 90.—Jamaica.
uniflora, *D. Don, in Edinb. N. Phil. Journ.* (Apr.–Oct. 1839) 88.—Hab.?
visciflora, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 228.—Ind. occ.
viscosa, *Lodd. Bot. Cab.* t. 772.—Ind. occ.

BRUNIA, Linn. Syst. ed. I (1735). *BRUNIACEÆ*, Benth. & Hook. f. i. 671.
BARCKIA, Burm. f. Prod. Fl. Cap. 12 (1768).
abrotanifolia, F. G. Dietr. Vollst. Lex. Gaertn. ii. 320 = Berzelia abrotanoides.
abrotanoides, Linn. Sp. Pl. 199 = Berzelia abrotanoides.
alopecuroides, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 375.—Afr. austr.
alopecuroides, Thunb. Prod. Pl. Cap. 187 = Berzelia alopecuroides.
arachnoides, Wendl. Coll. ii. 62 = Berzelia squarrosa.
brevifolia, D. Dietr. Syn. Pl. i. 848 = Berzelia abrotanoides.
candicans, *Hort. ex Steud. Nom.* ed. II. i. 231.—Afr. austr.
capitata, Desf. Tabl. Hort. Par. ed. II. 322 = Staavia capitella.
capitella, Thunb. Diss. Brun. 5 = Staavia capitella.
capitellata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 316 = staavioides.
ciliata, *Linn. Sp. Pl.* 199.—Afr. austr.
cuneata, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 17 = Berzelia lanuginosa.
cordata, Walp. Rep. i. 844 = racemosa.
cordifolia, D. Dietr. Syn. Pl. i. 848 = Berzelia cordifolia.
coronata, Linn. Mant. ii. 310, in syn. = Staavia glutinosa.
cupressina, Linn. l. c. ii. 349 = Diosma cupressina.

BRUNIA :—
densa, *Thunb. Prod. Pl. Cap.* 187.—Afr. austr.
densa, Willd. in Denkschr. Acad. Muench. (1806) 197. t. 7. f. 2 = Nebelia phylicoides.
Dregeana, *Presl, Bot. Bemerk.* 39.—Afr. austr.
elegans, *Dum.-Cours. Bot. Cult.* ed. I. iii. 616.—Afr. austr.
ericoides, Wendl. Coll. ii. 57 = Berzelia squarrosa.
flagelliformis, *Hort. ex Steud. Nom.* ed. II. i. 231 (quid ?).—Hab.?
formosa, D. Dietr. Syn. Pl. i. 848 = Berzelia abrotanoides.
formosa, *Dum.-Cours. Bot. Cult.* ed. I. iii. 616.—Afr. austr.
fragarioides, Willd. Sp. Pl. i. 1143 (= *Berardia globosa*). —Afr. austr.
glabrata, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 18.—Afr. austr.
globosa, Eckl. & Zeyh. Enum. 139 = laevis.
globosa, Thunb. Diss. Brun. 4 (= *Berardia globosa*).—Afr. austr.
glutinosa, Berg. Desc. Pl. Cap. 57 = Staavia glutinosa.
hirsuta, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 19.—Afr. austr.
imbricata, Sweet, Hort. Brit. ed. II. 115 = racemosa.
imbricata, *Wendl. f. ex Hoffm. Vers. Pfl. Nachtr.* i. 228, ii. & iii. 26, 71.—Afr. austr.
intermedia, D. Dietr. Syn. Pl. i. 848 = Berzelia intermedia.
laevis, *Thunb. Prod. Fl. Cap.* 187.—Afr. austr.
lanciflolia, *Walp. Rep.* i. 844.—Afr. austr.
lanuginosa, Linn. Sp. Pl. 199 = Berzelia lanuginosa.
lanuginosa, Thunb. Prod. Pl. Cap. 41 = Berzelia squarrosa.
laurifolia, Sond. in Harv. & Sond. Fl. Cap. ii. 315 = lanuginosa.
laxa, Thunb. Prod. Pl. Cap. 187 = Tittmannia laxa.
Levisanus, Linn. Sp. Pl. 199 = Leucadendron Levisanus.
macrocephala, *Willd. in Denkschr. Acad. Muench.* i. (1806) 152.—Afr. austr.
microcephala, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 320 (= *Berardia sphaerocephala*).—Afr. austr.
microphylla, Thunb. Prod. Pl. Cap. 187 (= *Berardia microphylla*).—Afr. austr.
nodiflora, *Linn. Sp. Pl.* 199.—Afr. austr.
palaeoca, Berg. Desc. Pl. Cap. 56 = Nebelia palaeoca.
palaeoca, Wendl. Coll. t. 3 = Berzelia intermedia.
passerinoides, *Schlecht. in Linnaea*, vi. (1831) 190.—Afr. austr.
phylicoides, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 321 (= *Berardia Dregeana*).—Afr. austr.
phylicoides, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 18 = Nebelia phylicoides.
pinifolia, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 374.—Afr. austr.
pinnaea, Linn. Amœn. Acad. Encyc. i. 475 = Berzelia squarrosa.
pungens, Berg. Desc. Pl. Cap. ii. 45, ex. typ. = *Fragarioides. Protea*, Crantz, Inst. i. 357 = Berzelia abrotanoides.
racemosa, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 374.—Afr. austr.
racemosa, Sond. in Harv. & Sond. Fl. Cap. ii. 315 = lanciflolia.
radiata, Berg. Desc. Pl. Cap. 58 = Staavia glutinosa.
radiata, Linn. Mant. ii. 309 = Staavia radiata.
rubra, Willd. in Denkschr. Acad. Muench. i. (1809) 131. t. 7 = Berzelia squarrosa.
sericea, Hort. ex Dum.-Cours. Bot. Cult. ed. II. vii. 329 = Berzelia superba.
speciosa, Hort. ex Dum.-Cours. l. c. vi. 379; vii. 329 = Berzelia superba.
squalida, *Sond. in Harv. & Sond. Fl. Cap.* ii. 315.—Afr. austr.
squarrosa, Sw. ex Harv. & Sond. l. c. 311 = Berzelia squarrosa.
squarrosa, Thunb. Diss. Brun. 5 = Berzelia squarrosa.
staavioides, *Sond. in Harv. & Sond. Fl. Cap.* ii. 316.—Afr. austr.
superba, Donn, Ind. Hort. Cantab. 25 = Berzelia lanuginosa.
superba, Krauss, ex Harv. & Sond. Fl. Cap. ii. 314 = laevis.
tenuifolia, Willd. in Denkschr. Acad. Muench. i. (1809) 129 = Berzelia lanuginosa.
teres, *Oliver, in Journ. Linn. Soc.* ix. (1867) 333.—Afr. austr.
Thunbergiana, D. Dietr. Syn. Pl. i. 849 = Nebelia palaeoca.
thymophora, *Walp. Rep.* i. 844.—Afr. austr.

BRUNIA :—
uniflora, Linn. Sp. Pl. 199 = Diosma cupressina.
verticillata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 322 = Staavia nuda.
verticillata, *Linn. f. Suppl.* 156.—Afr. austr.
verticillata, Thunb. Prod. Pl. Cap. 41 = virgata.
villosa, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 315.—Afr. austr.
virgata, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 376.—Afr. austr.

BRUNIERA, Franch., in Billiotia, (1864) 25 = Wolfia, Horkel. (Lemnac.).
viridpurea, Franch. l. c. = W. Michelli.

BRUNNERA, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 583 = Anchusa, Linn. (Boragin.).
myosotidiflora, Stev. l. c. = A. myosotidiflora.
sibirica, Stev. l. c.—Sibir.

BRUNNICHIA, Banks, ex Gaertn. Fruct. i. 213. t. 45 (1788). *POLYGONACEÆ*, Benth. & Hook. f. iii. 103.
FALLOPIA, Adans. Fam. ii. 274, 277 (1763).
RAJANIA, Walt. Fl. Carol. 247 (1788).
africana, *Wats. in Trans. Linn. Soc.* xxvii. (1872) 61. —Afr. trop.
cirrhosa, *Gaertn. Fruct. i.* 214. t. 45.—Am. bor.

BRUNONIA, Sm. in Trans. Linn. Soc. x. (1811) 365. t. 26, 29. *GOODENOVIEÆ*, Benth. & Hook. f. ii. 541.
australis, *Sm. l. c.* 367. t. 28.—Austral.
sericea, Sm. l. c. 367. t. 29 = australis.
simplex, Lindl. in Mitch. Trop. Austr. 89 = australis.

BRUNSFELSIA, Plum. ex Linn. Hort. Cliff. 5 (1737) = Brunfelsia, Linn. (Solanac.).

BRUNSVIA, Neck. Elem. iii. 337 (1790) = Croton, Linn. (Euphorb.).

BRUNSVIGIA, Heist. Beschr. Brunsv. Braunschweig (1755), 8, cum ic. *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 728.
albiflora, *Eckl. Topogr. Vers.* 7, nomen.—Afr. austr.
Burchelliana, Herb. Amaryll. t. 23. f. 2 = radulosa.
ciliaris, [Ker-Gawl. in] Bot. Reg. sub t. 192, 193 = Buphane disticha.
Cooperi, *Baker, in Saund. Ref. Bot.* t. 380.—Afr. austr.
coranica, *Sweet, Hort. Brit.* ed. I. 404.—Am. austr.
disticha, Sweet, l. c. = Buphane disticha.
falcata, [*Ker-Gawl. in*] *Bot. Mag.* t. 1443.—Afr. austr.
fulva, Steud. Nom. ed. II. i. 331 = Stenosmesson incarnatum.
gigantea, *Heist. Beschr. Brunsv. Braunschweig*, 3, *cum ic. fide Schult. f. Syst.* vii. 844 (nobis latet).—Afr. austr.
glauca, Salisb. Gen. Pl. Fragm. 117 = Josephinae.
grandiflora, *Lindl. Bot. Reg.* t. 1335.—Afr. austr.
Josephinae, [*Ker-Gawl. in*] *Bot. Reg.* t. 192, 193.—Afr. austr.
latiscrua, Sweet, Hort. Brit. ed. I. 404 = Nerine lucida.
lucida, Herb. App. 16 = Nerine lucida.
magnifica, Linden & Rodigas, in Illustr. Hortic. xxiii. (1876) t. 833 = Crinum Forbesianum.
marginata, Ait. Hort. Kew. ed. II. ii. 330 = Nerine marginata.
minor, *Lindl. Bot. Reg.* t. 964.—Afr. austr.
multiflora, Ait. Hort. Kew. ed. II. ii. 330 = gigantea.
orientalis, Ait. ex Eckl. Topog. Vers. 7 = gigantea.
Radula, *Ait. Hort. Kew.* ed. II. ii. 330.—Afr. austr.
radulosa, *Herb. Amaryll.* 181. t. 23. f. 3.—Afr. austr.
rubriscabia, M. Roem. Syn. Enast. 57 = gigantea.
Staceana, *Benth. & Hook. f. Gen.* iii. 727.—Afr. austr.
striata, *Ait. Hort. Kew.* ed. II. ii. 331.—Afr. austr.
tenuiorata, [Ker-Gawl. in] Bot. Reg. t. 367 = Buphane disticha.
vaginata, Schult. f. Syst. vii. 853 = Hessea vaginata.

BRUSCHIA, Bertol. in Mem. Acad. Sc. Bolog. viii. (1857) 286 (Misc. Bot. xviii. 16. t. 2) = Hymenanthe, Lem. (Oleac.).
macrocarpa, Bertol. ll. cc. 286; 17 = H. Arbor-tristis.

BRUXANELIA, *Donnat*, Schinzia, Hort. Malab. 31 (1818). *RUBIACEÆ* (Gen. dub.).
indica, *Donnat. l. c.*—Ind. or.

BRYA, Vell. Fl. Flum. 146 (1825); iv. t. 1 (1827) = **Hirtella**, Linn. (Rosac.).
racemosa, Vell. ll. cc. 147; iv. t. 1.—Bras.

BRYA, P. Br. Hist. Jamaic. 299. t. 31 (1756). *LEGUMINOSAE*, Benth. & Hook. f. i. 514.
Ebenus, *DC. Prod.* ii. 421.—Ind. occ.
leonensis, *Lodd. ex Loud. Hort. Brit.* 385, nomen.—Afr. trop.
nicaraguensis, *Oerst. in Kjoeb. Vidensh. Meddel.* (1855) 18.—Nicaragua.
parviflora, *DC. Prod.* ii. 421.—Bras.

BRYANTEA, Rafin. Sylva Tellur. 165 (1838) = **Litsea**, Lam. (Laurin.).
dealbata, *Rafin. l. c.* = L. dealbata.

BRYANTHUS, S. G. Gmel. Fl. Sibir. iv. 183. t. 57. f. 3 (1769). *ERICACEAE*, Benth. & Hook. f. ii. 596.
aleuticus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 368.—As. bor. or.; Am. bor. occ.
Breweri, *A. Gray, l. c.* 367.—Calif.
empetriformis, *A. Gray, l. c.*—Am. bor. occ.
erectus, Lindl. & Paxt. Flow. Gard. i. (1850–51) 97. glanduliflorus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 368.—Am. bor. occ.
Gmelini, *D. Don, in Edinb. N. Phil. Journ.* xvii. (July 1834) 160.—Sibir.; Kamtsch.
intermedius, A. Gray, in Proc. Am. Acad. vii. (1868) 367 = empetriformis.
Stelleri, D. Don, in Edinb. N. Phil. Journ. xvii. (July 1834) 160 = Cassiope Stelleriana.
taxifolius, *A. Gray, in Proc. Am. Acad.* vii. (1868) 368.—Reg. bor. frigid.

BRYANTIA, Webb, ex Gaudich. Bot. Voy. Bonite, t. 20 (1844 ?); Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 287. t. 15 = **Fandanus**, Linn.
butyrophora, Webb, l. c. = P. butyrophorus.
oblonga, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 288; in Comptes-Rendus, lxxx. (1875) 1196 = P. oblongus.
viscida, Brongn. ll. cc. 287; 1196 = P. viscidus.

BRYLKINIA, F. Schmidt, Reisen Amurl. 199. t. 8 (1868). *GRAMINEAE*, Benth. & Hook. f.' iii. 1192.
caudata, *F. Schmidt, l. c.*—Japon; Ins. Sachal.

BRYOBIUM, Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = **Bria**, Lindl. (Orchid.).
pubescens, Lindl. Bot. Reg. (1838) Misc. 79 = E. retusa.

BRYOCARPUM, Hook. f. & Thoms. in Hook. Kew Journ. ix. (1857) 200. t. 5. *PRIMULACEAE*, Benth. & Hook. f. ii. 684.
himalaicum, *Hook. f. & Thoms. l. c.*—Reg. Himal.

BRYOCLES, Salisb. in Trans. Hort. Soc. i. (1812) 335 = Punkia, Spreng. (Liliac.).
ventricosa, Salisb. l. c. = P. ovata.

BRYODES, Benth. in DC. Prod. x. 433 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 957.
micrantha, *Benth. l. c.*—Ins. Maurit.

BRYOMORPHA, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 172 = **Thylacospermum**, Fenzl (Caryophyll.).
rupifragum, Kar. & Kir. l. c. = T. rupifragum.

BRYOMORPHE, Harv. Thes. Cap. ii. 33. t. 151 (1863). *COMPOSITAE*, Benth. & Hook. f. ii. 334.
Zeyheri, *Harv. l. c.*—Afr. austr.

BRYONIA, Linn. Syst. ed. I (1735). *CUCURBITACEAE*, Benth. & Hook. f. i. 829.
abyssinica, Gouan, in Rev. Hortic. (1868) 61 = Maximowiczia Lindheimeri.
abyssinica, Lam. Encyc. i. 497 = Cocciata abyssinica.
acerifolia, D. Dietr. Syn. Pl. v. 367 = Coccinia indica.
acuta, Desf. Fl. Atlant. ii. 360 = dioica.
acutangula, Thunb. Prod. Fl. Cap. 13 = (Sicoecios sp.).
africa, Knill. Prod. Fl. Norfolk. 126 = Bryonopsis ciliata.

BRYONIA:—
africana, E. Mey. Zwei Pfl. Docum. 169 = Kedrostis grossulariaefolia.
africana, Linn. Sp. Pl. 1012 = Kedrostis africana.
africana, Thunb. Prod. Pl. Cap. 13 = Kedrostis nana.
agrestis, Raeusch. Nom. Bot. ed. II. 288 = Blastania Garcini.
alba, Bull. Herb. t. 55 = dioica.
alba, *Linn. Sp. Pl.* 1012.—Europ.; Reg. Caucas.; Persia.
alcaefolia, Willd. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 235 = Coccinia indica.
althaeoides, Ser. in DC. Prod. iii. 306 = Melothria maderaspatana.
americana, Descourt. Fl. Antill. ii. 281. t. 136 = Melothria pervaga.
americana, Lam. Encyc. i. 498 = Cayapoina americana.
amplexicaulis, Lam. l. c. 496 = Melothria amplexicaulis.
angulata, Thunb. Prod. Fl. Cap. 13 = Melothria punctata.
arguta, *Zipp. ex Span. in Linnaea*, xv. (1841) 206.—Malaya.
aspera, Stev. in Cat. Hort. Gorenk. (1812) 32 = alba.
asperifolia, Presl, Bot. Bemerk. 72 = Cayapoina americana.
attenuata, Hook. & Arn. Bot. Beech. Voy. Suppl. 424 = Cayapoina attenuata.
barbata, Buch.-Ham. ex Cogn. in DC. Monog. Phan. iii. 580 = Coccinia indica.
bifora, Buch.-Ham. in Wall. Cat. n. 6707 = Melothria odorata.
Blumei, Ser. in DC. Prod. iii. 305 = Melothria Rumphiana.
bonariensis, Mill. Gard. Dict. ed. VIII. n. 6 = Cayapoina ficifolia.
Boykinii, Torr. & Gray, Fl. N. Am. i. 540 = Cayapoina ficifolia.
Cabocla, Vell. Fl. Flum. x. t. 88 = Cayapoina globosa.
callosa, Heyne, ex Wall. Cat. sub n. 6708 = Melothria maderaspatana.
callosa, Kottl. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 210; Wall. Cat. n. 6710 = Cucumis trigonus.
capillacea, Schpm. & Thonn. Beskr. Guin. Pl. 430 = Melothria triangularis.
cheirophylla, Wall. Cat. n. 6715 A = Luffa aegyptiaca.
ciliata, Moench, Meth. 653 = Kedrostis africana.
cissoides, Wall. Cat. n. 6698 = Melothria perpusilla.
cochinchinensis, Lour. Fl. Cochinch. 595 = Gymnopetalum cochinchinense.
collosa, Rottl. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 210 = *sphalm. vide* callosa.
convolvulifolia, Schlecht. in Linnaea, xxvi. (1853–55) 646 = Melothria guadalupensis.
convolvuloides, A. Rich. Tent. Fl. Abyss. i. 388 = Coccinia adoensis.
cordata, Thunb. Fl. Cap. 149 = Melothria cordata.
cordatifolia, Godoy Torres, in Patriota, iii. n. 87 (1814) 71 = Cayapoina Tayuya.
cordifolia, Walp. Rep. ii. 109 = prsec.
cordifolia, Linn. Sp. Pl. 1012 = Coccinia indica.
cordifolia, Russ. ex Steud. Nom. ed. II. i. 282 = Melothria maderaspatana.
cretica, *Linn. Sp. Pl.* 1013.—Graecia et Ins. Creta.
cucumeroides, Ser. in DC. Prod. iii. 308 = Trichosanthes cucumeroides.
deltoidea, Am. Pugill. 19; in Nov. Act. Nat. Cur. xvii. (1836) 357 = Melothria scylanica.
deltoidea, Schum. & Thonn. Beskr. Guin. Pl. 429 = Melothria deltoidea.
dentata, E. Mey. Zwei Pfl. Docum. 169 = Melothria punctata.
digitata, Thunb. Prod. Pl. Cap. 13 = Kedrostis digitata.
digyna, Pomel, Nouv. Mat. Fl. Atl. 155 = dioica.
dioica, Bieb. Fl. Taur. Casc. iii. 625 = alba.
dioica, Bot. Hort. Maurit. 146 = Bryonopsis laciniosa.
dioica, *Jacq. Fl. Austr.* ii. 59. t. 199 (err. typ. dioica.).—Europ.; Reg. Mediterr.
dioscoria, Thunb. Prod. Pl. Cap. 13 = Kedrostis africana.
divisa, *Willd. ex Steud. Nom.* ed. II. i. 282, nomen.—Hab.?
epigaea, Blume, Bijdr. 925 = Melothria Rumphiana.
epigaea, Rottl. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 228 = Corallocarpus epigaeus.

BRYONIA:—
ficifolia, Descourt. Fl. Antill. vi. 89. t. 391 = Cayaponia americana.
ficifolia, Lam. Encyc. i. 498 = Cayaponia ficifolia.
filiculifolia, Wall. Cat. n. 6713 = Kedrostis rostrata.
filifolia, Spreng. Syst. iii. 16 = Cayaponia ficifolia.
filiformis, Roxb. Hort. Beng. 70; Fl. Ind. iii. 737 = Melothria heterophylla?
fimbristipula, Fenzl, ex Hook. f. in Oliver, Fl. Trop. Afr. ii. 558 = Blastania fimbristipula.
fluminensis, Vell. Fl. Flum. x. t. 87 = Cayaponia fluminensis.
fortidissima, Schum. & Thonn. Beskr. Guin. Pl. 428 = Kedrostis foetidissima.
Garcini, Willd. Sp. Pl. iv. 628 = Blastania Garcini.
geminata, Blume, Bijdr. 924 = Melothria indica.
glabra, Roxb. Hort. Beng. 104 = Corallocarpus epigeus.
glandulosa, *Gill. in Hook. Bot. Misc.* iii. (1833) 324.—Chili.
glandulosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 56. t. 175 = Cayaponia glandulosa.
gracilis, Wall. Cat. n. 6714 = Melothria maderaspatana.
grandis, Linn. Mant. i. 126 = Coccinia indica.
grandis, Wall. Cat. n. 6700 K = Gymnopetalum cochinchinense.
grandis, Wall. l. c. L = Momordica dioica.
grossulariifolia, E. Mey. Zwei Pfl. Docum. 169 = Kedrostis grossulariaefolia.
guadalupensis, Spreng. Syst. iii. 15 = Melothria pervaga.
guineensis, G. Don, Gen. Syst iii. 82 (sp. dub.).—Afr. trop.
hastata, Lour. Fl. Cochinch. ii. 594 = Melothria heterophylla.
hederaefolia, Jacq. Fragm. 78. n. 280. t. 113 = verrucosa.
heterophylla, Blume, Bijdr. 925 = Melothria Rumphiana.
heterophylla, Raeusch. Nom. ed. III. 282; Steud. Nom. ed. I. 192 = Melothria heterophylla.
heterophylla, Wall. Cat. n. 6704 = Melothria marginata?
hispida, Salisb. Prod. 158 = Melothria maderaspatana.
Hookeriana, Wight & Arn. Prod. 345 = Melothria perpusilla.
japonica, Thunb. Fl. Jap. 325 = Melothria japonica.
jatrophaefolia, A. Rich. Tent. Fl. Abyss. i. 389 = Coccinia jatrophaefolia.
Johnstoni, Cusent, ex Seem. in Journ. Bot. ii. (1864) 47 = Melothria Grayana.
laciniosa, Linn. Sp. Pl. 1013 = Bryonopsis laciniosa.
lawii, Thunb. Prod. Pl. Cap. 13 = Melothria velutina.
lagenaria, E. Mey. Zwei Pfl. Docum. 169 = Coccinia sessilifolia.
latebrosa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 384 = Cayaponia latebrosa.
leiosperma, Wight & Arn. Prod. 345 = Melothria leiosperma.
leucocarpa, Blume, Bijdr. 924 = Melothria odorata.
Lhotskyana, Cogn. in Mart. Fl. Bras. vi. iv. 88 = Cayaponia Lhotskyana.
lutea, Bast. ex Ser. in DC. Prod. iii. 307 = dioica.
macrophylla, *Kotschy, ex Boiss. Fl. Orient.* ii. 761.—Armenia.
macrophylla, *Ser. in DC. Prod.* iii. 308 (sp. dub.).—Hab.?
maderaspatana, Berg. Descr. Pl. Cap. 351 = Melothria punctata.
maderaspatana, Lam. Encyc. i. 496 = Melothria maderaspatana.
marginata, Blume, Bijdr. 924 = Melothria Rumphiana.
mayaroensis, Wall. Cat. n. 6705 = Melothria maderaspatana.
mayaroensis, Wight, Ic. iii. t. 7581 = Melothria perpusilla.
Melothria, Crantz, Inst. i. 173 = Melothria pendula.
micrantha, Boiss. Diagn. Ser. I. x. 7 = syriaca.
micrantha, Hochst. ex Cogn. in DC. Monog. Phan. iii. 632 = Melothria maderaspatana.
microcoda, E. Mey. Zwei Pfl. Docum. 169 = Melothria maderaspatana.
Moimoi, Ser. in DC. Prod. iii. 305 = Coccinea Moghadd.
mucronata, Blume, Bijdr. 923 = Melothria mucronata.
multifida, E. Mey. Zwei Pfl. Docum. 169 = Kedrostis africana.
multiflora, *Boiss. & Heldr. Diagn.* Ser. I. x. 8.—Oriens; Ind. bor. occ.

BRYONIA :—
myosvensis, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 628 = Melothria leiosperma.
myosvensis, Walp. Rep. ii. 198 = Melothria macrostachya.
myosvensis, Wight & Arn. Prod. i. 345 = Melothria perpusilla.
nana, Lam. Encyc. i. 497 = Kedrostis nana.
nepaulensis, Ser. in DC. Prod. iii. 307 = Melothria heterophylla.
nigra, Gilib. Fl. Lituan. i. 60 = alba.
nitida, Link, Enum. Hort. Berol. ii. 404 = dioica.
obtusa, A. Rich. Tent. Fl. Abyss. i. 288 = Mukia scabrella.
obtusiloba, E. Mey. Zwei Pfl. Docum. 169 = Melothria obtusiloba.
odorata, Buch.-Ham. ex Wall. Cat. n. 6706 B, C = Melothria odorata.
odorata, Wall. Cat. n. 6705 D = Melothria Wallichii.
oxyphylla, Wall. Cat. n. 6698 = Melothria perpusilla.
palmata, Linn. Sp. Pl. 1012 = Bryonopsis laciniosa.
palmata, Wall. Cat. n. 6711 A, B, C = Coccinia indica.
palmata, Wall. l. c. D = Corallocarpus epigaeus.
Panckeri, Naud. in Ann. Sc. Nat. Sér. IV. ix. (1858) 298 = Bryonopsis affinis.
pectinata, E. Mey. Zwei Pfl. Docum. 169 = Trochomeria pectinata.
pedata, Hamilt. in Flora, xxv. (1842) Beibl. 49 = Bryonopsis laciniosa.
peduncolosa, Ser. in DC. Prod. iii. 306 = Herpetospermum caudigerum.
perpusilla, Blume, Bijdr. 926 = Melothria perpusilla.
Perrottetiana, Ser. in DC. Prod. iii. 304 = Kedrostis foetidissima.
pilosa, Vell. Fl. Flum. x. t. 86 = Cayaponia diffusa.
pilosa, Roxb. Hort. Beng. 104; Fl. Ind. iii. 726 = Kedrostis rostrata.
pinnatifida, Vell. Fl. Flum. x. t. 90 = Cayaponia diversifolia.
pinnatifida, Burch. Trav. i. 547 = Kedrostis africana.
pinnatiloba, M. Roem. Syn. fasc. ii. Pepon. 40 = Cayaponia diversifolia.
prostrata, D. Dietr. Syn. Pl. v. 363 = Kedrostis rostrata.
polymorpha, *Fenzl. ex Tranto. in Act. Hort. Petrop.* viii. (1883) 353, nomen.—Hab.?
pubescens, *Pohr. Encyc. Suppl.* i. 731 (sp. dub.)—Oriens.
punctata, Buch.-Ham. ex Wall. Cat. sub n. 6706 = Melothria maderaspatana.
punctata, Thunb. Prod. Fl. Cap. 13 = Melothria punctata.
quinquefolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 48 = Bryonopsis laciniosa.
quinquelobata, Thunb. Prod. Fl. Cap. 13 = Coccinia quinquelobata.
racemosa, *Mill. Gard. Dict.* ed. VIII. n. 4.—Ind. occ.
racemosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 56 = Cayaponia Poeppigii.
racemosa, Sw. Prod. Veg. Ind. Occ. = Cayaponia racemosa.
reniformis, Roxb. ex Wight & Arn. Prod. 344 = Blastania Garcini.
repanda, Blume, Bijdr. 928 = Melothria punctata.
Rheedei, Blume, l. c. 925 = Melothria heterophylla.
rostrata, Rottl. in Ges. Naturf. Freunde Neue Schr. iv. (1803) 213 = Kedrostis rostrata.
Rottleri, Spreng. Syst. iii. 15 = Melothria maderaspatana.
ruderalis, Salisb. Prod. 158 = dioica.
ruderalis, *Zipp. ex Span. in Linnaea*, xv. (1841) 206, nomen.—Malaya.
sagittata, Blume, Bijdr. 926 = Melothria heterophylla.
scabra, Rottl. ex Wight & Arn. Prod. i. 345 = Melothria maderaspatana.
scabra, Thunb. Prod. Fl. Cap. 13 = Melothria punctata.
scabrata, Blume, Bijdr. 923 = Melothria perpusilla.
scabrella, Linn. f. Suppl. 434 = Melothria maderaspatana.
scabrella, Sieber, ex Cogn. in DC. Monog. Phan. iii. 736 = Cayaponia americana.
scabrella, Wall. Cat. n. 6706 = Melothria leiosperma.
scariatina, Dum. Fl. Belg. 54 = dioica.
scrobiculata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 367 = Melothria scrobiculata.
septemlobata, *Vell. Fl. Flum.* x. t. 85.—Bras.
sicula, Guss. Fl. Sic. Syn. ii. 631 = dioica.
stenuata, Wall. Cat. n. 6711 D = Corallocarpus epigaeus.
sinuosa, Wall. Cat. n. 6716 = Coccinia indica et Melothria heterophylla.

BRYONIA :—
Solena, Mirb. ex Steud. Nom. ed. II. i. 283 = Melothria heterophylla.
stipulacea, Willd. Sp. Pl. iv. 620 = Blastania Garcini.
subpinnata, Steud. Nom. ed. II. i. 283 = Cayaponia diversifolia.
syriaca, *Boiss. Diagn.* Ser. II. ii. 59.—Syria; Palaestin.
Tayuya, Mart. in Linnaea, xii. (1838) Litt. 86 = Cayaponia Martiana.
Tayuya, Vell. Fl. Flum. x. t. 89 = Cayaponia Tayuya.
Teedonda, Roxb. ex Willd. Sp. Pl. iv. 618 = Melothria heterophylla.
tenella, Roxb. Hort. Beng. 70; Fl. Ind. iii. 725 = Melothria indica.
tenuifolia, Gilib. in Hook. Bot. Misc. iii. (1833) 323 = Abobra tenuifolia.
tenuifolia, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 56.—Chili.
tenuis, Klotzsch, in Peters, Reise Mossamb. Bot. 150 = Bryonopsis laciniosa.
ternata, Vell. Fl. Flum. x. t. 91 = Cayaponia ternata.
Thunbergiana, Dietr. ex Steud. Nom. ed. I. 228 = Melothria cordata.
Tinni, Huet, ex Cogn. in DC. Monog. Phan. iii. 471 = dioica.
tomentosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
tricuspidata, Lesch. ex Steud. Nom. ed. II. i. 275 = Passiflora Leschenaultii.
trifora, Wall. Cat. n. 6707 = Melothria leucocarpa.
triloba, Lour. Fl. Cochinch. ii. 595 = Blastania Garcini.
triloba, Thunb. Prod. Fl. Cap. 12 = Kedrostis nana.
trilobata, Ser. in DC. Prod. iii. 307 = Kedrostis nana.
tuberosa, Wight & Arn. Prod. 347 = Gymnopetalum Wightii.
umbellata, Klein, ex Willd. Sp. Pl. iv. 618 = Melothria heterophylla.
umbellata, Wall. Cat. n. 6705 D = Trichosanthes cucumerina.
umbellata, Wall. l. c. K, L = Melothria amplexicaulis.
variegata, Mill. Gard. Dict. ed. VIII. n. 5 = Bryonopsis laciniosa.
verrucosa, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 385.—Ins. Canar.
vitifolia, *Hort. Belg. ex Heynh. Nom.* ii. 79, nomen.—Hab.?
vittata, *Zipp. ex Span. in Linnaea*, xv. (1841) 206, nomen.—Malaya.
vulgaris, Gaudichaud, ex Ledeb. Fl. Ross. ii. 1401 = alba.
Wallichiana, Ser. ex DC. Prod. iii. 309 = Melothria heterophylla.
Wightiana, Wall. Cat. n. 6708 = Melothria leiosperma.

BRYONOPSIS, Arn. in Hook. Journ. Bot. iii. (1841) 274. *CUCURBITACEÆ*, [Benth. & Hook. f. i. 374].
affinis, *Cogn. in DC. Monog. Phan.* iii. 479.—Ins. Norfolk.
Bonnetiana, Miq. Fl. Ind. Bat. i. i. 657 = Cerasicarpum zeylanicum.
courtallensis, Arn. in Hook. Journ. Bot. iii. (1841) 274 = laciniosa.
erythrocarpa, Naud. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 194 = laciniosa.
laciniosa, *Naud. l. c.* Sér. V. vi. (1866) 30.—As. Afr. et Austral. trop.
leucocarpa, Miq. Fl. Ind. Bat. i. i. 657 = Melothria odorata.
Panckeri, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 30 = affinis.
pinnata, Hassk. Cat. Pl. Hort. Bog. Alt. 189 = laciniosa.

BRYOPHTHALMUM, E. Mey. ex Presl. Pflanseng. 101 (1839) = **Momordica**, Salisb. (Ericac.).
uniflorum, E. Mey. l. c. = M. uniflora.

BRYOPSYLLUM, Salisb. Parad. Lond. t. 8 (1806). *CRASSULACEÆ*, Benth. & Hook. f. i. 656.
PHYLLOCALCIUM, Vent. in Flora, iii. (1820) 409.
CRASSULA, Comm. ex Lam. Encyc. ii. 141 (1786).
calycinum, *Salisb. l. c.*—Mexic.
cochlearium, Lem. Illustr. Hortic. vi. (1859) Misc. 100 = Kalanchoë cochlearium.
crenatum, *Baker, in Journ. Linn. Soc.* xx. (1884) 189. —Madag.
gemmiparum, Blanco, Fl. Filip. ed. II. 220 = calycinum.
jongatum, Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 85 = calycinum.
proliferum, *Bowie, ex Hook. Bot. Mag.* t. 5147.—Afr. austr.

BRYOPHYLLUM :—
serratum, Blanco, Fl. Filip. ed. II. 220 = Kalanchoe laciniata.
triangulare, Blanco, l. c. 221 = Kalanchoë spathulata.
tubiflorum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 380. —Afr. austr.

BUBALINA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 86 = **Burchellia**, R. Br. (Rubiac.).

BUBANIA, Girard, in Mém. Acad. Montp. (Sect. Sc.) i. (1848) 182 = **Limnastrastrum**, Moench (Plumbagin.). *Feei*, Girard, l. c. 185 = L. Gyrostanum.
monopetala, Girard, l. c. 188 = L. articulatum.

BUBON, Linn. Gen. ed. I. 389 (1737) = **Smyrnium**, Linn (Umbellif.).
aphyllus, Cham. & Schlecht. in Linnaea, i. (1826) 389 (= *Daucus tortuosus*).—Aegypt.
buchtormensis, Spreng. Pugill. ii. 55 = S. buchtormense.
capensis, Sond. in Harv. & Sond. Fl. Cap. iii. 561.— Afr. austr.
cuneifolius, Spreng. Syst. i. 900 = S. cuneifolium.
dichotomus, Link, Enum. Hort. Berol. i. (1821) 278 = S. dichotomum.
eriocephalus, Pall. ex Spreng. Syst. i. 900 = S. Lessingianum.
Galbanum, Hort. ex Steud. Nom. ed. II. i. 633 (= *Meloxanthum decipiens*).—Madera.
Galbanum, Linn. Sp. Pl. 364 = Pencedanum Galbanum.
garganicus, Tenore, Fl. Nap. i. 153 = Athamanta sicula.
glaucus, Spreng. Umb. Spec. 186 = S. petraeum.
globosus, Stev. ex Ledeb. Fl. Ross. ii. 274 = S. rigidum.
gummifer, Linn. Sp. Pl. 254 = S. gummiferum.
humilis, Salisb. Prod. Stirp. 165 = Athamanta macedonica.
hypoleucus, Meisn. in Hook. Lond. Journ. Bot. ii. (1840) 536.—Afr. austr.
laevigatus, Ait. Hort. Kew. ed. II. v. i. 259.—Afr. austr
latifolius, Salisb. Prod. 165 = *laevigatus*.
Libanotis, Dum. Fl. Belg. 79 = S. Libanotis.
macedonicus, Linn. Sp. Pl. 253 = Athamanta macedonica.
montanus, Sond. in Harv. & Sond. Fl. Cap. iii. 560.— Afr. austr.
multiflorus, Moench, Meth. 95 = Athamanta macedonica.
perviscus, Pall. ex Ledeb. Fl. Ross. ii. 329 = Zosima orientalis.
pseudamplexifolius, Spreng. Syst. i. 900 = S. rigidum.
pimpinellifolius, Eckl. & Zeyh. Enum. 258.—Afr. austr.
proliferus, Burm. f. Fl. Cap. Prod. 7 = Lichtensteinia pyrethrifolia.
pyrenaicus, Dum. Fl. Belg. 79 = S. vulgare.
rigidior, Linn. Sp. Pl. 254 = Ferula communis.
rigidus, Spreng. Pugill. ii. 53 = S. gummiferum.
rigidus, Spreng. Syst. i. 900 = S. rigidum.
siculus, Spreng. in Roem. & Schult. Syst. vi. 499 = S. Brocconi.
tenuifolius, Salisb. Prod. 165 = S. gummiferum.
tenuifolius, Sond. in Harv. & Sond. Fl. Cap. ii. 560.— Afr. austr.
tortuosus, Desf. Fl. Atlant. i. 257 (= *Daucus tortuosus*). —Afr. bor.

BUBONIUM, Hill, Veg. Syst. ii. 74 (1764); Hort. Kew. 13 (1769) (Compos.).
aquaticum, Hill, ii. cc. = Odontospermum aquaticum.
arborescens, Hill, Hort. Kew. 13 = Borrichia arborescens.
frutescens, Hill, Veg. Syst. ii. 74; Hort. Kew. 13 = Borrichia frutescens.
maritimum, Hill, Hort. Kew. 13 = Odontospermum maritimum.
spinosum, Hill, Veg. Syst. ii. 74 = Pallenis spinosa.

BUBROMA, Schreb. Gen. 538 (1791) = **Guazuma**, Plum. (Sterculiac.).
crinitum, Stend. Nom. ed. II. i. 398 = G. crinita.
grandiflorum, Willd. ex Steud. l. c. = G. grandiflora.
Guazuma, Willd. Sp. Pl. iii. 1469 = G. ulmifolia.
lumbos, Willd. Enum. Hort. Berol. 996 = G. ulmifolia.
polybotrys, Willd. l. c. = G. polybotrya.
tomentosum, Spreng. Syst. iii. 225 = G. ulmifolia.

BUCAFER, Adans. Fam. ii. 439 (1763) = **Tanghinia**, Thou. (Island.).

BUCANION, Sav. in Bull. Soc. Nat. Mosc. (1851) i. 568 = **Heliotropium**, Tourn. (Boragin.).
Saovitsii, Sav. i. c. = *H. Saovitsii*.

BUCCAFERREA, Babani, in Nuovo Giorn. Bot. Ital. v. (1878) 316 = **Potamogeton**, Tourn. (Naiad.).
Hornemanni, Babani, l. c. = P. Hornemanni.
obtusifolia, Babani, l. c. 317 = P. obtusifolius.

BUCCAFERREA, Mich. ex Petagna, Inst. Bot. v. 1826 (1787) = **Ruppia**, Linn. (Naiad.).
cirrhosa, Petagna, l. c. = R. maritima.

BUCCO, Wendl. Coll. 18, tt. 2, 3, 18, 28, 77 (1808) = **Agathosma**, Wendl. (Rutac.).
acuminata, Wendl. Coll. i. t. 28 = A. imbricata.
barbata, Roem. & Schult. Syst. v. 445 = Macrostylis lanceolata.
betulina, Roem. & Schult. l. c. 448 = Barosma betulina.
brevifolia, Roem. & Schult. l. c. 446 = A. erecta.
Cerefolium, Roem. & Schult. l. c. 439 = A. Cerefolium.
crenata, Roem. & Schult. l. c. 444 = Barosma crenata.
cuspidata, Wendl. Coll. iii. 17. t. 31 = A. cuspidata.
erecta, Wendl. l. c. i. 17. t. 3 = A. erecta.
hamata, Wendl. ex Bartl. & Wendl. f. Diosmeae, 114 = Barosma diolca.
hirta, Roem. & Schult. Syst. v. 446 = A. hirta.
hispida, Roem. & Schult. l. c. = A. hispida.
imbricata, Wendl. Coll. i. 28. t. 9 = A. imbricata.
Lamarckiana, Roem. & Schult. Syst. v. 447 = A. virgata.
linifolia, Lichtst. in Roem. & Schult. l. c. 448 = A. linifolia.
obtusa, Wendl. Coll. i. 45 = A. obtusa.
obtusata, Wendl. l. c. iii. 7. t. 76 = A. imbricata.
ovata, Roem. & Schult. Syst. v. 442 = Barosma ovata.
prolifera, Wendl. Coll. iii. t. 77 = A. prolifera.
pulchella, Roem. & Schult. Syst. v. 449 = Barosma pulchella.
rubra, Roem. & Schult. l. c. 441 = Diosma vulgaris.
scabra, Wendl. ex Bartl. & Wendl. f. Diosmeae, 156 = A. ciliata.
serpyllacea, Roem. & Schult. Syst. v. 447 = A. serpyllacea.
spicata, Roem. & Schult. l. c. 448 = A. spicata.
tetragona, Roem. & Schult. l. c. 444 = Acmadenia tetragona.
Ventenatiana, Roem. & Schult. l. c. 449 = A. Ventenatiana.
ovatifolia, Roem. & Schult. l. c. 447 = A. imbricata.
villosa, Wendl. Coll. i. 14. t. 2 = A. villosa.

BUCCULINA, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 209 = **Holothrix**, Rich. (Orchid.).
aspera, Lindl. l. c. = H. aspera.

BUCEPHALANDRA, Schott, Gen. Aroid. t. 56 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 984. Mottleyana, *Schott, l. c.*—Borneo.

BUCEPHALON, [Plum.] Linn. Gen. ed. I. 375 (1737) = **Tropha**, Linn. (Urtic.).
racemosum, Linn. Sp. Pl. 1190 = T. americana.

BUCERAS, Hall. ex All. Fl. Pedem. i. 313 (1785) = **Trigonella**, Linn. (Legumin.).
corniculata, All. Fl. Pedem. i. 313 = T. corniculata.
elliptica, Moench, Meth. 148 = T. monspeliaca.
Foenum-graecum, All. Fl. Pedem. i. 313 = T. Foenum-graecum.
grandiflora, Moench, Meth. 148 = T. corniculata.
monspeliaca, All. Fl. Pedem. i. 313 = T. monspeliaca.
nautica, Moench, Meth. 149 = T. polyceratia.
odoratissima, Medic. in Vorles. Churpf. Phys. Gen. ii. (1787) 338 = T. Foenum-graecum.
platycarpa, Moench, Meth. Suppl. 51 = Medicago platycarpa.
polyceratia, All. Fl. Pedem. i. 313 = M. polyceratia.
spinosa, Moench, Meth. 148 = T. spinosa.
torosa, Moench, Meth. Suppl. 51 = T. striata.
tortuosa, Stend. Nom. ed. II. i. 283 = T. striata.

BUCERAS, P. Br. Hist. Jamaic. t. 28 (1756) = **Turnimalia**, Linn. (Combretac.).
Bucida, Cranz, Inst. i. 128 = T. Catappa.

BUCEROSIA, Endl. Gen. 896 (1836) = **Boucerosia**, Wight & Arn. (Asclep.).

BUCETUM, Parnell, Grasses of Scotl. 104-109 (1843) = **Festuca**, Tourn. (Gramin.).
elatius, Parnell, l. c. 107 = F. elatior.
gigantheum, Parnell, l. c. 108 = F. gigantea.
loliaceum, Parnell, l. c. 104 = F. loliacea.
pratense, Parnell, l. c. 105 = F. elatior.

BUCHANANIA, Sm. Exot. Bot. ii. tab. solum, 115 (1805) = **Colebrookea**, Sm. (Labiat.).
oppositifolia, Sm. l. c. = C. oppositifolia.

BUCHANANIA, Spreng. in Schrad. Journ. (1800) ii. 284; Roxb. Fl. Corom. iii. 79. t. 283 (1819). *ANACARDIACEAE*, Benth. & Hook. f. i. 491.
CAMBESSEDEA, Kunth, in Ann. Sc. Nat. ii. (1824) 366.
CONIOGETON, Blume, Bijdr. 1156 (1826).
LANZANA, Stokes, Bot. Mat. Med. ii. 548 (1812).
LAUNEAN, Buch.-Ham. in As. Res. v. (1799) 138.
LAUNZEA, Endl. Ench. 599 (1841).
LUNDIA, Puer. ex DC. Prod. ii. 64, in syn. (1825).
acuminata, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 472.—Burma; Malaya.
acuminata, Wall. Cat. n. 981 = lanceolata.
amboinensis, *Miq. in Ann. Mus. Lugd. Bat.* iv. 117. —Amboina.
angustifolia, Benth. Fl. Austral. i. 490 = Muelleri.
angustifolia, Roxb. Hort. Beng. 32; Fl. Ind. ii. 386 = B. Lanzan.
arborescens, Blume, Mus. Bot. Lugd. Bat. i. 183 = floridа.
arborescens, F. Muell. Fragm. vii. 23 = Muelleri.
auriculata, Blume, l. c. 185 = Campnosperma auriculata.
bancana, Miq. Fl. Ind. Bat. Suppl. i. 528 = floridа.
ferruginea, *Engl. in DC. Monog. Phan.* iv. 194.—Ind. or.
floridа, A. Gray, in Bot. U. St. Expl. Exped. i. 366 = vitiensis.
floridа, *Schum. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1846) 483.—As. trop.; Malaya.
glabra, *Wall. Cat.* n. 984.—Burma.
Hainaheirae, *Miq. in Ann. Mus. Lugd. Bat.* iv. 117. —Malaya.
insignis, *Blume, Mus. Bot. Lugd. Bat.* i. 184.—Borneo.
intermedia, Wight, Ic. i. t. 81 = floridа.
lanceolata, Roxb. Hort. Beng. 32 = lancifolia.
lanceolata, *Wight, Ic.* i. t. 387.—Ind. or.
lancifolia, *Roxb. Fl. Ind.* ii. 386.—Ind. or.; Burma.
Lanzan, *Spreng. in Schrad. Journ.* (1800) ii. 284.—Ind. or.
latifolia, *Roxb. Hort. Beng.* 32; *Fl. Ind.* ii. 385.—Ind. or.; Burma.
laxiflora, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 304.—Burma.
longifolia, Blume, Mus. Bot. Lugd. Bat. i. 184 = floridа.
longifolia, *Span. in Linnaea*, xv. (1841) 168.—Ins. Timor.
lucida, Blume, Mus. Bot. Lugd. Bat. i. 184 = floridа.
macrophylla, Blume, l. c. 185 = Campnosperma macrophylla.
mangoides, *F. Muell. Fragm.* vii. 23.—Ins. Pacif.
microphylla, *Engl. in DC. Monog. Phan.* iv. 185.—Ins. Philipp.
Muelleri, *Engl. l. c.* 190.—Austral.
nitida, *Engl. l. c.* 192.—Ins. Philipp.
obovata, *Engl. l. c.* 187.—Austral.
oxyphylla, *Miq. Fl. Ind. Bat. Suppl.* i. 522.—Sumatra.
oxyrhachis, Miq. l. c. 524 = Campnosperma oxyrhachis.
polymorpha, Blume, Mus. Bot. Lugd. Bat. i. 186 = floridа.
paniculata, Roxb. ex Steud. Nom. ed. II. i. 233 = Amoora cucullata.
petiolaris, Miq. Fl. Ind. Bat. i. ii. 587 = floridа.
platyneura, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 196.—Ins. Nicobar.
polyhastya, Miq. Fl. Ind. Bat. i. ii. 588 = floridа.
racemiflora, Miq. Fl. Ind. Suppl. i. 523 = Campnosperma macrophylla.
reticulata, *Hance, in Journ. Bot.* xv. (1877) 332.—Cochinch.
semilifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 184.—Malaya.
sinuosa, *Miq. in Ann. Mus. Lugd. Bat.* iv. 118.—Siam.
spicata, Roxb. ex Wall. Cat. n. 4895 = Amoora cucullata.
splendens, *Miq. Fl. Ind. Bat. Suppl.* i. 524.—Sumatra.
subobovata, Griff. Notul. iv. 613 = floridа.

BUCHANANIA:—
vitiensis, *Engl. in DC. Monog. Phan.* iv. 186.—Ins. Fiji.
zeylanica, Blume, Mus. Bot. Lugd. Bat. i. 185 = Mangifera zeylanica.

BUCHAREA, Rafin. Fl. Tellur. iv. 84 (1836) = **Convolvulus**, Tourn.
atlantica, Rafin. l. c.—Afr. bor.
maderensis, Rafin. l. c.—Ins. Madera.

BUCHENAVIA, Eichl. in Flora, xlix. (1866) 164 = **Terminalia**, Linn. (Combret.).
capitata, Eichl. l. c. 165 = T. Hilarians.
macrophylla, Eichl. l. c. 166 = T. macrophylla.
megalophylla, Heurck & Muell. Arg. in Heurck, Obs. Bot. 211.—Am. trop.
octopetala, Eichl. in Flora, xlix. (1866) 165.—Am. trop.
oxycarpa, Eichl. l. c. 166 = T. oxycarpa.
punctata, Eichl. l. c. = T. punctata, *Spruce*.
reticulata, Eichl. l. c.—Am. trop.
suaveolens, Eichl. l. c. = T. suaveolens.
tomentosa, Eichl. l. c. = T. tomentosa, *Mart.*

BUCHENROEDERA, Eckl. & Zeyh. Enum. 194 (1836). *LEGUMINOSAE*, Benth. & Hook. f. i. 478. COLOBOTUS, E. Mey. Comm. Pl. Afr. Austr. 156. (1836).
alpina, *Eckl. & Zeyh. Enum.* 195.—Afr. austr.
carnulecona, *Presl, Rel. Bemerk.* 47.—Afr. austr.
gracilis, *Eckl. & Zeyh. Enum.* 195.—Afr. austr.
holosericea, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 581.—Afr. austr.
lanceolata, *Presl, Rel. Bemerk.* 47.—Afr. austr.
Meyeri, *Presl, l. c.*—Afr. austr.
multiflora, *Eckl. & Zeyh. Enum.* 196.—Afr. austr.
spicata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 93.—Afr. austr.
tenuifolia, *Eckl. & Zeyh. Enum.* 196.—Afr. austr.
tereticaulis, Eckl. & Zeyh. l. c. = Aspalathus armata.
trichodes, *Presl, Rel. Bemerk.* 47.—Afr. austr.
umbellata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 93.—Afr. austr.
vimines, *Presl, Rel. Bemerk.* 47.—Afr. austr.

BUCHERIA, Heynh. Nom. ii. 80 (1846) = **Thrypticomaena**, Endl. (Myrtac.).
saxicola, Heynh. l. c. = T. saxicola.

BUCHIA, D. Dietr. Syn. Pl. iii. 870 (1843) = **Bouchea**, Cham. (Verbenac.).

BUCHIA, H. B. & K. Nov. Gen. et Sp. ii. 269. t. 192 (1817) = **Perama**, Aubl. (Rubiac.).
plantaginea, H. B. & K. l. c. 270. t. 193 = P. dichotoma.

BUCHINGERA, Boiss. & Hohen. Diagn. Ser. I. viii. 29 (1849). *CRUCIFERAE*, Benth. & Hook. f. i. 72. axillaris, *Boiss. & Hohen. l. c.*—Persia.

BUCHINGERA, F. Schultz, in Jahrb. Pharm. (Mar. 1847).—Cf. Bot. Zeit. vi. (1848) 760 = Cuscuta, Linn. (Convolv.).
suaveolens, F. Schultz, l. c. = C. racemosa.

BUCHLOË, Engelm. in Trans. Acad. Sc. St. Louis (1859), 433. t. 19, 14. *GRAMINEAE*, Benth. & Hook. f. iii. 1173.
BULBILIS, Rafin. in Journ. Phys. lxxxix. (1819) 262.
CALANTHERA, Nutt. ex Hook. Kew Journ. viii. (1856) 18.
SESLERIA, Nutt. Gen. Am. i. 64 (1818).
dactyloides, *Engelm. l. c.*—Texas; Mexic.

BUCHNERA, Linn. Hort. Cliff. 301 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 968.
BONNETIA, Neck. Elem. i. 368 (1790).
CRYPTA, Gaertn. f. Fruct. iii. 134. t. 214 (1805).
MICAILIA, Rafin. Fl. Tellur. ii. 104 (1836).
PIRIPEA, Aubl. Pl. Gui. ii. 697. t. 259 (1775).
abyssinica, Linn. Mant. ii. 251 = Chaenostoma asbiopicum.
africana, Linn. Fl. Afr. Rar. 13; in Amoen. Acad. vi. 89 = Trianga Apala.
americana, *Linn. Sp. Pl.* 630.—Am. bor.
anathystina, Cham. & Schlecht. in Linnaea, ii. (1827) 585 = lobelioides.

BUCHNERA :—
angustifolia, D. Don, Prod. Fl. Nep. 91 = Striga euphrasioides.
angustifolia, *Rafin. New Fl. Am.* ii. 32.—Am. bor.
aquatica, Wight, ex Steud. Nom. ed. II. i. 384 = Striga lutea.
arguta, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 374 = ramosissima.
asiatica, Linn. Sp. Pl. 630; Vahl, Symb. Bot. iii. 31 = Striga densiflora.
aspera, Schum. & Thonn. Beskr. Guin. Pl. 380 = Striga aspera.
asperata, R. Br. Prod. 436 = linearis.
aurantica, Burch. Trav. i. 388 = Lyperia multifida.
aurea, Banks, ex Benth. in Hook. Comp. Bot. Mag. i. (1835) 372 = Polycenea aurea.
bifida, Buch.-Ham. in Wall. Cat. n. 3872 = Striga euphrasioides.
bilabiata, Thunb. Prod. Pl. Cap. 100 = Striga Thunbergii.
canadensis, Linn. Mant. 88 = Verbena Aubletia.
capensis, Linn. l. c. = Polycarena capensis.
capillaris, *Desv. in Ham. Prodr. Fl. Ind. Occ.* 46.—Guiana.
capitata, *Benth. in DC. Prod.* x. 495.—Madag.
capitata, *Burm. f. Fl. Cap. Prod.* 17.—Afr. austr.
cernua, Houtt. Handleid. ii. 549 = Bouchea cuscifolia.
cernua, Linn. Mant. ii. 251 = Bouchea cernua.
coccinea, Benth. Scroph. Ind. 40 = Striga lutea.
coccinea, Soop. Delic. Insob. ii. 10 = Achimenes nocciuea.
cordifolia, Linn. f. Suppl. 287 = Streptium asperum.
cruciata, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 91.—Au. trop.; Malaya.
cuneifolia, Thunb. Prod. Pl. Cap. 100 = Bouchea cuscifolia.
curvifora, R. Br. Prod. 433 = Striga curvifora.
densiflora, Benth. Scroph. Ind. 41 = Striga densiflora.
densiflora, Hook. & Arn. Bot. Beech. Voy. 203 = cruciata.
denticulata, Buch.-Ham. ex Wall. Cat. n. 3872 = Striga euphrasioides.
distichа, *H. B. & K. Nov. Gen. et Sp.* ii. 340.—N. Granat.
divaricata, Linn. ex Walp. Rep. iii. 308 = Zaluzianskya pusilla.
dura, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 366.—Afr. austr.
elegans, D. Dietr. Syn. Pl. iii. 525 = Striga elegans.
elongata, [Benth.] ex Griseb. Fl. Brit. W. Ind. 428 = longifolia.
elongata, *Sw. Prod. Veg. Ind. Occ.* 92.—Am. calid.
erinoides, *Yarous, Pl. Nov. Cap.* 19.—Afr. austr.
euphrasioides, Vahl, Symb. Bot. iii. 31 = Striga euphrasioides.
exserta, *Fauc. in H. O. Forbes, Nat. Wand. East. Archip.* 512.—Ins. Timor.
foetida, Andr. Bot. Rep. t. 80 = Chaenostoma hispidum.
foetida, Jacq. Hort. Schoenb. iv. 23. t. 448 = Chaenostoma foetidum.
foetida, *Schrad. Journ.* 1800. ii. (1801) 196.—Hab. ?
Forbesii, D. Dietr. Syn. Pl. iii. 526 = Striga Forbesii.
gunneroides, Willd. Sp. Pl. iii. 338 = Striga orobanchoides.
glabrata, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 366.—Afr. austr.
gracilis, *R. Br. Prod.* 437.—Austral.
grandiflora, Linn. f. Suppl. 287 = Escobedia scabrifolia.
Hermonthica, Delile, Fl. Egypte, 245. t. 34 = Striga Hermonthica.
Heyneana, D. Dietr. Syn. Pl. iii. 525 = Striga densiflora.
hirsuta, Wall. Cat. n. 3869 = Striga euphrasioides.
hispida, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 91.—Ind. or. ; Afr. trop.
humifusa, Vahl, Symb. Bot. iii. 31 = Striga humifusa.
hydrobadensis, Roth, Nov. Sp. Pl. 293 = Striga orobanchoides.
Juncea, *Cham. & Schlecht. in Linnaea*, ii. (1827) 590.—Bras.
lavandulacea, Cham. & Schlecht. l. c. 589 = rosea.
leptostachya, *Benth. in DC. Prod.* x. 497.—Afr. trop.
levicaulis, *Rafin. New Fl. Am.* ii. 29.—Am. bor.
lilacina, Zipp. ex Span. in Linnaea, xv. (1841) 331 = ramosissima.
linearifolia, *Schum. & Thonn. Beskr. Guin. Pl.* 379.—Afr. trop.
linearis, *R. Br. Prod.* 437.—Austral.

BUCHNERA :—
lithospermifolia, H. B. & K. Nov. Gen. et Sp. ii. 341 = elongata.
lobelioides, *Cham. & Schlecht. in Linnaea*, xi. (1837) 589.—Bras.
longiflora, Am. in Nov. Act. Nat. Cur. xviii. (1836) 358 = Rhamphicarpa longiflora.
longifolia, H. B. & K. Nov. Gen. et Sp. Pl. ii. 340 = elongata.
longifolia, *Klotzsch, in Peters, Reise Mossamb. Bot.* 225.—Afr. trop.
macrantha, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 366.—Afr. trop.
macrocarpa, *A. Rich. Tent. Fl. Abyss.* ii. 198.—Abyssin.
macrocarpa, *H. B. & K. Nov. Gen. et Sp.* ii. 342.—N. Granat.
major, *Poiret. in Linnaea*, xli. (1877) 588.—Costa Rica.
Mauritia, Buch.-Ham. ex Benth. Scroph. Ind. 41 = Striga Mauritia.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* ii. 457.—Mexic.
minarica, *Rafin. New Fl. Am.* ii. 32.—Am. bor.
montevidensis, Spreng. ex Walp. Rep. iv. 37 = Verbena venosa.
Mooreria, Steud. Nom. ed. II. i. 384 = Striga Mauritia.
mossambica, *Klotzsch, in Peters, Reise Mossamb. Bot.* 224.—Afr. trop.
multifora, D. Dietr. Syn. Pl. iii. 525; F. Muell. Cens. 98 = Striga multiflora.
nigrescens, Span. Ic. 69; et in Linnaea, xv. (1841) 331 = strigula.
obliqua, *Benth. in DC. Prod.* x. 498.—Mexic.
oppositifolia, Hort. ex Steud. Nom. ed. II. i. 384 = Chaenostoma hispidum.
orobanchoides, R. Br. in Salt, Abyss. App. 64 = Striga orobanchoides.
palustris, *Spreng. Syst.* ii. 805.—Guiana.
parviflora, R. Br. Prod. 438 = Striga parviflora.
pedunculata, Andr. Bot. Rep. t. 54 = Lyperia pedunculata.
Peripea, Steud. Nom. ed. I. 124 = palustris.
phoenicea, Wall. in Benth. Scroph. Ind. 41 = Striga lutea.
pilosa, Benth. Bot. Voy. Sulph. 144 = elongata.
pinnatifida, Linn. f. Suppl. 288 = Bouchea pinnatifida.
Pirrpea, Raesch. Nom. ed. III. 179 = palustris.
pubescens, Benth. in Hook. Comp. Bot. Mag. i. (1835) 365 = ramosissima.
pusilla, *H. B. & K. Nov. Gen. et Sp.* ii. 340.—N. Granat.
ramosissima, *R. Br. Prod.* 438.—Austral.
rosea, *H. B. & K. Nov. Gen. et Sp.* ii. 342.—Venezuela.
rupestris, Sw. Fl. Ind. Occ. ii. 1068 = Rachicallis rupestris.
Schimperiana, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 198 = hispida.
stricta, Benth. in Hook. Comp. Bot. Mag. i. (1835) 367 = cruciata.
sumatrana, Miq. Fl. Ind. Bat. ii. 703 = cruciata.
tenella, *R. Br. Prod.* 437.—Austral.
tenuifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 340.—Am. austr.
tetragona, *R. Br. Prod.* 437.—Austral.
tetrastichа, *Wall. Cat.* n. 3878 ; *et in Benth. Scroph. Ind.* 41.—Assam ; Burma.
Thunbergii, D. Dietr. Syn. Pl. iii. 525 = Striga Thunbergii.
timorensis, *Fauc. in H. O. Forbes, Nat. Wand. East. Archip.* 513.—Ins. Timor.
tinctoria, *Bertol. Fl. Guatim.* 26.—Guatemala.
tomentosa, *Blume, Bijdr.* 741.—Java.
urticifolia, *R. Br. Prod.* 437.—Austral.
usaramdo, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 121.—Afr. trop.
verbenoides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 226.—Afr. trop.
virgata, *H. B. & K. Nov. Gen. et Sp.* ii. 342.—N. Granat.
viscosa, [Roland. in] Alt. Hort. Kew. ed. I. ii. 257 = Sphenandra viscosa.
Wallichii, Benth. in Wall. Cat. n. 3870 = Striga Masuria.

BUCHOLTZIA, Meisn. Gen. 313 ; Comm. 330 (1841) = BUCHOLZIA, Mart. = Telanthera, P. Br. (Amaranth.).

BUCHOLZIA, Mart. Nov. Gen. et Sp. ii. 49. t. 147 (1826) = Telanthera, R. Br. (Amaranth.).
brachiata, Schrad. ex Seub. in Mart. Fl. Bras. v. 178 = T. ficoides.
diffusa, Mart. Nov. Gen. et Sp. ii. tt. 149, 151 = T. polygonoides.
ficoides, Mart. l. c. 52 = T. ficoides.
filifolia, Hook. f. in Trans. Linn. Soc. xx. (1851) 192 = T. filifolia.
frutescens, Mart. Nov. Gen. et Sp. ii. 52, in obs. = T. frutescens.
glauca, Hook. f. in Trans. Linn. Soc. xx. (1851) 191 = T. giauscescens.
lupulina, Mart. Beitr. Amarant. 108 = T. lupulina.
maritima, Mart. Nov. Gen. et Sp. ii. 50. t. 147 = T. maritima.
nudicaulis, Hook. f. in Trans. Linn. Soc. xx. (1851) 191 = T. nudicaulis.
obovata, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 348 = T. obovata.
philcoroides, Mart. Beitr. Amarant. 107. n. 5 = T. philoxeroides.
polygonoides, Mart. Nov. Gen. et Sp. ii. 51 = T. polygonoides.
spinescens, Gill. ex Moq. in DC. Prod. xiii. ii. 359 = Alternanthera Achyranthа.

BUCHOLZIA, Stadtm. ex Willem. in Ust. Ann. Bot xvii. (1796) 54 = Combretum, vel Terminalia, Linn.

BUCHOZIA, L'Hérit. ex Juss. in Dict. Sc. Nat. v. (1804) 409 ; et in Mém. Mus. Par. vi. (1820) 395 = Barleas, Comm. (Kubiac.).
coprosmoides, L'Hérit. ex DC. Prod. iv. 575 = S foetida.

BUCHOZIA, Vell. Fl. Flum. 33 (1825) = Hoteranthera, Ruiz & Pav. (Pontederc.).
aquatica, Vell. l. c. 33 ; i. c. 80 = H. reniformis.

BUCIDA, Linn. Syst. ed. X. 1025 (1759) = Terminalia, Linn. (Combret.).
angustifolia, DC. Prod. iii. 10 = T. Buceras.
angustifolia, Spruce, ex Eichl. in Mart. Fl. Bras. xiv. ii. 96 = T. Hilariana.
Buceras, Linn. Syst. ed. X. 1095 = T. Buceras.
Buceras, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = T. Catappa.
Buceras, Vell. Fl. Flum. 172 ; iv. t. 87 = Laguncularia racemosa.
capitata, Vahl, Eclog. Am. i. 50. t. 8 = T. Hilariana.
Comintsna, Blanco, Fl. Filip. ed. I. 356 = T. Catappa.
megaptamica, Spreng. Syst. iv. Cur. Post. 177.—Bras
nitida, Hassk. in Flora, xxvii. (1844) 605.—Ins Sandvic.

BUCKINGHAMIA, F. Muell. Fragm. vi. 248 (1868).
PROTEACEAE, Benth. & Hook. f. iii. 182.
celsissima, *F. Muell. l. c.*—Austral.

BUCKLANDIA, R. Br. in Wall. Cat. n. 7414 (1833).
HAMAMELIDEAE, Benth. & Hook. f. i. 666.
populifolia, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 86 = populnea.
populnea, *R. Br. in Wall. Cat.* n. 7414.—Reg. Himal.

BUCKLEYA, Torr. in Am. Journ. Sc. xly. (1843) 170. *SANTALACEAE*, Benth. & Hook. f. ii. 236.
DARBYA, A. Gray, in Am. Journ. Sc. Ser. II. i. (1846) 388.
NESTRONIA, Rafin. New Fl. Am. iii. 19 (1836).
QUADRILAS, Sieb. & Zucc. in Abh. Akad. Munch. iv. ii. (1846) 194 ; (Fam. Nat. Fl. Jap. i. 96. t. 9). distichophylla, *Torr. l. c.*—Am. bor.

BUCKNERA, Michx. Fl. Bor. Am. ii. 18 (1804) = Buchnera, Linn. (Scrophal.).

BUCQUETIA, DC. Prod. iii. 110 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 736.
glutinosa, *DC. l. c.*—N. Granat.
nigrisilla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 28.—Ecuador.

BUDA, Adans. Fam. ii. 507 (1763); Dum. Fl. Belg. 110 (1827) = **Spergularia**, C. & J. Presl (1817) (Caryoph.).
marina, Dum. l. c. = S. marina.
media, Dum. l. c. = S. marina.
rubra, Dum. l. c. = S. rubra.

BUDDLEIA, Houst. ex Linn. Gen. ed. I. 26 (1737). *LOGANIACEÆ*, Benth. & Hook. f. ii. 792.
ADENOPLEA, Radlk. in Bremen Abh. viii. (1883) 406.
ROMANA, Vell. Fl. Flum. 54, pro parte (1825).
abbreviata, *H. B. & K. Nov. Gen. et Sp.* ii. 353.—Mexic.
acuminata, R. Br. in Salt, Abyss. App. 62 = polystachya.
acuminata, H. B. & K. l. c. 349 = Humboldtiana.
acuminata, *Poir. Encyc. Suppl.* i. 745.—Ind or.
acuminatissima, Blume, Bijdr. 748 = asiatica.
alpina, *Cerst. in Kjoed. Vidensk. Meddel.* (1858) 25.—Costa Rica.
alternifolia, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 494.—China.
americana, H. B. & K. Nov. Gen. et Sp. ii. 351 = spicata.
americana, *Linn. Sp. Pl.* 119.—Am. austr.
angustata, Benth. in DC. Prod. x. 443 = vetula.
araucana, *Phil. in Anal. Univ. Chil.* (1873) 525.—Chili.
arborea, Meyen, Reise, i. 478 = pichinchensis.
aromatica, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 227.—Bolivia.
asiatica, *Lour. Fl. Cochinch.* 72.—As. trop.; Malaya.
auriculata, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 60.—Afr. austr.
australis, Vell. Fl. Flum. 41; i. t. 104 = brasiliensis.
axillaris, Willd. in Roem. & Schult. Syst. iii. 97 = Adenoplusia Willdenowii.
barbata, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 19.—Mexic.
betonicæfolia, *Lam. Illustr.* i. 291.—Peruv.
bracteata, *Cham. & Schlecht. in Linnæa*, ii. (1827) 599.—Bras.
brasiliensis, *Vacq. f. Eclog.* t. 158.—Bras.
brunfelsia, Willd. ex Roem. & Schult. Syst. iii. Mant. 97 = abbreviata.
bullata, *H. B. & K. Nov. Gen. et Sp.* ii. 348.—Ecuador.
callicarpoides, H. B. & K. l. c. 350 = americana.
calycina, *Benth. in DC. Prod.* x. 440.—Ecuador.
campestris, *Walp. Rep.* iii. 330.—Bras.
cana, Willd. ex Roem. & Schult. Syst. iii. Mant. 94 = verbascifolia.
capitata, Jacq. Ic. Rar. ii. 9; Coll. ii. 332 = globosa.
carnea, Carr. in Rev. Hortic. (1879) 90 = curviflora.
cestriflora, *Cham. in Linnæa*, viii. (1833) 20.—Bras.
Colvilei, *Hook. f. Illustr. Himal. Pl.* t. 18.—Reg. Himal.
comorensis, *Baker, in Journ. Linn. Soc.* xx. (1884) 206.—Ins. Comor.
connata, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 358 = brasiliensis.
connata, *Ruiz & Pav. Fl. Per.* i. 52. t. 18.—Peruv.
cordata, *H. B. & K. Nov. Gen. et Sp.* ii. 348. t. 185.—Mexic.
cordobensis, *Griseb. in Gentt. Abh.* xix. (1874) 213.—Reg. Argent.
coriacea, *Kenny, in Ann. Sc. Nat.* Sér. III. viii. (1847) 226.—Bolivia.
crispa, Benth. in Wall. Cat. n. 6404; et Scroph. ad. 48 = paniculata.
cronsoldes, *A. Gray, in Proc. Am. Acad.* v. (1861) 305.—Calif.
cuneata, *Cham. in Linnæa*, viii. (1833) 17.—Bras.
curviflora, *Hook. & Arn. Bot. Beech. Voy.* 367.—China.
curviflola, *Hort. ex Carr. in Rev. Hortic.* (1866) 339.—Hab.?
decurrens, Cham. & Schlecht. in Linnæa, v. (1830) 106 = americana.
densiflora, *Blume, Bijdr.* 748.—Java.
densata, *H. B. & K. Nov. Gen. et Sp.* ii. 352.—Venezuela.
diffusa, *Ruiz & Pav. Fl. Per.* i. 53. t. 20.—Peruv.
discolor, Roth, Nov. Pl. Sp. 88 = asiatica.
dioscoreifolia, Vahl, Symb. Bot. iii. 15 = Nicodemia diversifolia.
elegans, *Cham. & Schlecht. in Linnæa*, ii. (1827) 594.—Bras.

BUDDLEIA:—
ellipticus, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 24.—Mexic.
ferruginea, Pohl, ex J. A. Schmidt, in Mart. Fl. Bras. 264 = elegans.
flocculosa, *Kunth, in Ind. Sem. Hort. Berol.* (1844). —*Cf. Linnæa*, xviii. (1844) 501.—Mexic.
floribunda, *H. B. & K. Nov. Gen. et Sp.* ii. 352.—Mexic.
foliata, L. B. ex Benth. in DC. Prod. x. 446 = polystachya.
fusca, *Baker, in Journ. Linn. Soc.* xx. (1884) 205.—Madag.
Gayana, *Benth. in DC. Prod.* x. 442.—Chili.
glaberrima, Hort. Amat. Cat. Hort. Taurin. 18 = Freylinia castroides.
glabra, Hort. ex C. Koch, Dendrol. ii. 11. 295 = curviflora.
glabrata, Spreng. Syst. i. 481 = Cephalanthus Sarandi.
globifera, Mirb. in Duham. Arb. ed. Nov. i. t. 25 = globosa.
globosa, *Hope, in Verh. Hariem.* xx. ii. (1782) 417. t. 11.—Chili; Peruv.
glomerata, *Wendl. ex Benth. in DC. Prod.* x. 447, nomen.—Afr. austr.
gracilis, *Kunth, in Ind. Sem. Hort. Berol.* (1844). —*Cf. Linnæa*, xviii. (1844) 500.—Mexic.
grandiflora, *Cham. & Schlecht. in Linnæa*, ii. (1827) 596.—Bras.
guaranica, *Roxb. Hort. Beng.* [10], nomen.—Am. austr.
heterophylla, Lindl. Bot. Reg. t. 1259 = madagascariensis.
Humboldtiana, *Roem. & Schult. Syst.* iii. 98.—Texas; Mexic.
incana, *Ruiz & Pav. Fl. Per.* i. 52. t. 80.—Peruv.
incompta, *Linn. f. Suppl.* 123.—Afr. austr.
indica, Lam. Encyc. i. 513 = Nicodemia diversifolia.
insignis, *A. Van Geert, Cat.* n. 78 (1879–80), 8; *Carr. in Rev. Hortic.* (1878) 330.—Hab.?
integrifolia, Willd. ex Roem. & Schult. Syst. iii. Mant. 96 = abbreviata.
intermedia, *H. B. & K. Nov. Gen. et Sp.* ii. 350.—Mexic.
internodia, *Lorens, in Roca, Exped. Rio-Negro*, ii. 255, nota.—Patagon.
interrupta, Buch.-Ham. ex Wall. Cat. sub n. 6401 = asiatica.
interrupta, *H. B. & K. Nov. Gen. et Sp.* ii. 348.—Ecuador.
Jamesonii, *Benth. in DC. Prod.* x. 441.—Ecuador.
japonica, *Linden*, Cat. n. 19 (1865).—Japon.
lanata, *Benth. in DC. Prod.* x. 441.—Peruv.
lanceolata, *Benth. Pl. Hartw.* 48.—Am. centr.
lanceolata, Heyne, ex Wall. Cat. sub n. 6401 = asiatica.
Lindenii, *Benth. in DC. Prod.* x. 438.—N. Granat.
Lindleyana, *Fortune, in Lindl. Bot. Reg.* (1844) *Misc.* 91.—China.
longifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 349.—Peruv.
Lucæ, *Niederl. in Roca, Exped. Rio-Negro*, ii. 254.—Patagon.
macrantha, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 385 = grandiflora.
macrostachya, *Kunth, in Ind. Sem. Hort. Berol.* (1844).—*Cf. Linnæa*, xviii. (1844) 500.—Hab.?
macrostachya, *Benth. in Wall. Cat.* n. 6407; *et Scroph. Ind.* 43.—Reg. Himal.
madagascariensis, *Lam. Encyc.* i. 513.—Ins. Mascar.; Madag.
marrubiifolia, *Benth. in DC. Prod.* x. 441.—Texas; Mexic.
Martii, J. A. Schmidt, in Journ. Bot. vi. (1868) 228, 345 = macrostachya.
mellodora, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 10.—Mexic.
mendozinensis, *Gill. ex Benth. in DC. Prod.* x. 443.—Chili.
microphylla, *H. B. & K. Nov. Gen. et Sp.* ii. 353.—Mexic.
Missionis, *Benth. in Wall. Cat.* n. 6405; *et Scroph. Ind.* 43.—Ind. or.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 349.—Ecuador.
Nappii, *Lorens, in Roca, Exped. Rio-Negro*, ii. 254.—Patagon.
Neemda, Buch.-Ham. in Wall. Cat. n. 6401 = asiatica.
Neemda, Hort. ex Link, Enum. Hort. Berol. i. 126 = brasiliensis.

BUDDLEIA:—
nepalensis, Colla, in Mem. Acc. Torin. xxxv. (1833) 183 = paniculata.
nitida, *Benth. in DC. Prod.* x. 437.—Mexic.
oblonga, *Benth. l. c.* 442.—Bras.
obtusifolia, Mart. & Gal. in Bull. Acad. Brux. xii. 11. (1845) 26 = microphylla.
occidentalis, *Linn. Sp. Pl.* ed. II. 162.—Jamaic.; Peruv.
occidentalis, Ruiz & Pav. Fl. Per. i. 53. t. 82. f. a = americana.
officinalis, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 496.—China.
otophylla, *Hassk. in Versl. en Med. Kon. Akad. Wetensch.* v. (1857) 97.—Malaya.
ovalifolia, *Kunth, in Ind. Sem. Hort. Berol.* (1844). —*Cf. Linnæa*, xviii. (1844) 500.—Mexic.
paniculata, *Wall. Cat* n. 6403; *et in Roxb. Pl. Ind. ed. Carey*, i. 412.—Reg Himal.; Burma.
parviflora, *H. B. & K. Nov. Gen. et Sp.* ii. 353.—Mexic.
parvifolia, Willd. ex Roem. & Schult. Syst. iii. Mant. 96 = microphylla.
perfoliata, *H. B. & K. Nov. Gen. et Sp.* ii. 346.—Mexic.
pichinchensis, *H. B. & K. l. c.*—Ecuador.
Poiretii, *Spreng. Syst.* i. 480.—Ind. or.
polycephala, *H. B. & K. Nov. Gen. et Sp.* ii. 346.—Peruv.
polystachya, *Fresen. in Flora*, xxi. (1838) 605.—Abyssin.
Pringlei, *A. Gray, in Proc. Am. Acad.* xix. (1883) 86.—Arizona.
propinqua, *Kunth, Ind. Sem. Hort. Berol.* (1844). —*Cf. Linnæa*, xviii. (1844) 501.—Mexic.
pseudoverticillata, Mart. & Gal. in Bull. Acad. Brux. xii. 11. (1845) 24 = sessiliflora.
racemosa, *Torr. Bot. Mex. Bound.* 121.—Texas
rondeletiæflora, *Benth. in DC. Prod.* x. 445.—Ins. Comor.
rufa, *Fresen. in Flora*, xxi. (1838) 606.—Abyss.
rufescens, Willd. ex Roem. & Schult. Syst. iii. Mant. 97 = americana.
rupcola, *H. B. & K. Nov. Gen. et Sp.* ii. 347.—N. Granat.
salicifolia, Jacq. Hort. Schoenb. i. 12 = Chilianthus arborcus.
salicifolia, Vahl, Symb. Bot. iii. 14 = thyrsoides.
salsicina, Lam. Illustr. i. 291 = asiatica.
saligna, Willd. Enum. Hort. Berol. 159 = Chilianthus arborea.
Saltiana, Steud. Nom. ed. II. i. 285 = polystachya.
salvifolia, *Lam. Encyc.* i. 513.—Afr. austr.
scordioides, *H. B. & K. Nov. Gen. et Sp.* ii. 345. t. 183.—Texas; Mexic.
serrulata, Roth, Nov. Pl. Sp. 88 = asiatica.
sessiliflora, *H. B. & K. Nov. Gen. et Sp.* ii. 345. t. 182.—Mexic.
sinuata, *Willd. ex Roem. & Schult. Syst. Mant.* iii. 97.—Madag.
spectabilis, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 11.—Mexic.
sphærantha, Cham. & Schlecht. in Linnæa, v. (1830) 104 = perfoliata.
sphærocephala, *Baker, in Journ. Linn. Soc.* xxi. (1885) 425.—Madag.
spicata, *Ruiz & Pav. Fl. Per.* i. 53. t. 81.—Peruv.
stachyoides, *Cham. & Schlecht. in Linnæa*, ii. (1827) 597.—Bras.
stæveolens, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 11.—Chili.
subserrata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 92 = asiatica.
sundaica, *Blume, Bijdr.* 743.—Java.
tenuifolia, *Griseb. in Gentt. Abh.* xix. (1874) 213.—Am. austr.
ternata, *Lour. Fl. Cochinch.* 72.—Cochinch.
thapsoides, Desf. Cat. Hort. Par. ed. III. 106 = brasiliensis.
thyrsoides, *Lam. Illustr.* i. 291.—Reg. Argent.
tubiflora, *Benth. in DC. Prod.* x. 448.—Bras.
tucumanensis, *Griseb. in Gentt. Abh.* xix. (1874) 213.—Tucuman.
venusta, *Kunth, in Ind. Sem. Hort. Berol.* (1844).—*Cf. Linnæa*, xviii. (1844) 500.—Mexic.
Verbasci, Pers. Syn. i. 135 = americana.
verbascifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 351.—N. Granat.

BUDDLEIA :—
verticillata, *H. B & K. Nov. Gen. et Sp.* 346. t. 184.—Mexic.
vetule, *Cham. in Linnaea*, viii. (1833) 18.—Bras.
virgata, Blanco, Fl. Filip. ed. 1. 57 = salatica.
virgata, Lin. f. Suppl. 133 = Gomphostigma scoparioides.
volubilis, *Lam. Illustr.* i. 391.—Ins. Mascar.

BUECHNERIA, Roth, Nov. Fl. Sp. 292 (1821) = **Buchnera**, Linn. (Scrophul.).

BUECKIA, A. Rich. in Orb. Dict. ii. 759 (1849) = BUKEIA, Nees = **Blynanthus**, Linn. (Cyperac.).

BUKEIA, Gieseke, Prael. Linn. 199, 206 (1792) = **Alpinia**, Linn. (Scitamin.).
malaccensis, Raeusch. Nom. ed. III. 1 = A. malaccensis.

BUKEIA, Nees, in Linnaea, ix. (1834) 300 = **Blynanthus**, Linn. (Cyperac.).
punctoria, Nees, l. c. = E. punctorius.

BUELOWIA, Schum. & Thonn. Beskr. Guin. Pl. 246 (1827) = **Smeathmannia**, Soland. (Passifor.).
illustris, Schum. & Thonn. l. c. 247 = S. pubescens.

BUENA, Cav. in Anal. Hist. Nat. ii. (1800) 276 ; Ic. vi. 49 = GONZALAGUNIA, Ruiz & Pav. = **Gonsales**, Pers. (Rubiac.).
panamensis, Cav. ii. cc. 279 ; 50 = G. panamensis.

BUENA, Pohl, Pl. Bras. Ic. i. 8 (1827) ; Wedd. in Journ. Linn. Soc. xi. (1869) 186 = **Cosmibuena**, Ruiz & Pav. et Cascarilla, Wedd. (Rubiac.).
acuminata, DC. Prod. iv. 356 = Cosmibuena acuminata.
acutifolia, Wedd. in Journ. Linn. Soc. xi. (1869) 186 = Cascarilla acutifolia.
bullata, Wedd. l. c. 187 = Cascarilla bullata.
calycina, Wedd. l. c. = Cascarilla calycina.
Cornea, Wedd. l. c. = Cascarilla Cornea.
citrifolia, Wedd. l. c. = Cascarilla citrifolia.
crassifolia, Wedd. l. c. = Cascarilla calyptrata.
Gaudichaudiana, Wedd. l. c. = Cascarilla Gaudichaudiana.
heterophylla, Wedd. l. c. 186 = Cascarilla heterophylla.
hexandra, Pohl, Pl. Bras. Ic. i. 10. t. 8 = Cascarilla hexandra.
Hookeriana, Wedd. in Journ. Linn. Soc. xi. (1869) 187 = Cascarilla Hookeriana.
Lambertiana, Wedd. l. c. = Cascarilla Lambertiana.
latifolia, Benth. Pl. Hartw. 191 = Cosmibuena latifolia.
macrocarpa, Benth. Bot. Voy. Sulph. 104. t. 38 = Cosmibuena macrocarpa.
macrocarpa, Wedd. in Journ. Linn. Soc. xi. (1869) 187 = Cascarilla macrocarpa.
magnifolia, Wedd. l. c. 186 = Cascarilla magnifolia.
mucronata, Wedd. l. c. 187 = Cascarilla mucronata.
nitida, Wedd. l. c. 186 = Cascarilla nitida.
obtusifolia, DC. Prod. iv. 356 = Cosmibuena obtusifolia.
Pavonii, Wedd. in Journ. Linn. Soc. xi. (1869) 186 = Cascarilla Pavonii.
Riedeliana, Wedd. l. c. = Cascarilla Riedeliana.
Riparensis, Wedd. l. c. = Cascarilla magnifolia.
Roraimae, Wedd. l. c. 187 = Cascarilla Roraimae.
Skinneri, Cerut. in Kjoeb. Vidensk. Meddel. (1852) 48 = Cosmibuena Skinneri.
stenocarpa, Wedd. in Journ. Linn. Soc. xi. (1869) 186 = Cascarilla stenocarpa.
trifora, Benth. in Hook. Journ. Bot. iii. (1841) 216 = Cosmibuena trifora.
undata, Wedd. in Journ. Linn. Soc. xi. (1869) 187 = Cascarilla undata.

BUERGERIA, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 53 (1867) = **Cladrastis**, Raf. (Leguminos.).
floribunda, Miq. l. c. = C. amurensis.

BUERGERIA, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1845) 186. t. 9 (Fl. Jap. Fam. i. 78. t. 2) = Talauma, Juss. (Magnol.).
obovata, Sieb. & Zucc. ll. cc. 187 = Magnolia obovata.
salicifolia, Sieb. & Zucc. l. c. = T. salicifolia.
stellata, Sieb. & Zucc. l. c. 186 = T. stellata.

BUETTNERA, J. F. Gmel. Syst. ii. 404 (1791) = seq.

BUETTNERIA, Loefl. Iter Hisp. 318 (1758) (Byttneria) ; Linn. Syst. ed. X. ii. 939 (1759). *STERCULIACEAE*, Benth. & Hook. f. i. 236.
BYTTNERIA, Loefl. Iter Hisp. 318 (1758).
CHANTARA, Jacq. Enum. Pl. Carib. 2 (1760).
HETEROPHYLLUM, Boj. ex Hook. Bot. Misc. i. (1830) 291.
PENTACEROS, G. F. W. Mey. Prim. Fl. Esseq. 186 (1818).
TELFAIRIA, Newm. ex Hook. Bot. Misc. i. (1830) 291.
abutiloides, St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 31 = hirsuta.
aculeata, Domb. ex Lam. Encyc. i. 528 = cordata.
aculeata, Jacq. Stirp. Am. 76 = carthagenensis.
acuminata, *Bred. ex Roem. & Schult. Syst.* v. 470.—Venezuela.
afinis, Pohl, Pl. Bras. ii. 78 = melastomaefolia.
africana, *Mast. in Oliver, Fl. Trop. Afr.* i. 239.—Afr. trop.
andamanensis, *Kurz, in Journ. As. Soc.* Beng. ii. (1872) 47 ; *et Flora*, liv. (1871) 377.—Burma.
angulata, *Nasth. Cat. Pl. Hort. Bog.* 204.—Java.
aspera, *Colebr. in Roxb. Fl. Ind. ed. Carey*, ii. 388.—Reg. Himal. ; China.
australis, Sieber, ex Steud. Nom. ed. II. i. 241 = pannosa.
austrocalis, St. Hil. Fl. Bras. Mer. i. 145 = tereticaulis.
bilobe, *Baill. Adansonia*, x. (1872) 178.—Madag.
brasiliensis, *Sprang. Syst.* i. 790.—Bras.
brevipes, *Benth. Pl. Hartw.* 114.—Am. trop.
caledonica, *Turcz. in Bull. Soc. Nat. Mosc.* (1868) i. 566.—N. Caled.
carthagenensis, *Jacq. Stirp. Am. Pict.* 41.—N. Granat.
catapifolia, *Vacq. Hort. Schoenbr.* i. 21.—Am. trop.
catalpifolia, Wall. Cat. n. 1152 = crenulata ?
celtoides, *St. Hil. Fl. Bras. Mer.* i. 141.—Bras.
corchorifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 152.—Am. austr.
cordifolia, *Sagot, in Ann. Sc. Nat. Sér.* VI. xi. (1881) 155.—Guiana.
cordata, *Lam. Encyc.* i. 528.—Peruv.
coryllifolia, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 470.—Am. austr.
crenulata, *Wall. Cat.* n. 1150.—Reg. Himal. ; Burma.
dasyphylla, J. Gay, in DC. Prod. i. 486 = Rullagia pannosa.
denisia, *Pohl, Pl. Bras.* ii. 79.—Bras.
discolor, *Benth. in Hook. Kew Journ.* iii. (1851) 165.—Bras.
divaricata, *Benth. in Hook. Journ. Bot.* iv. (1842) 194.—Guiana.
echinata, Wall. Cat. n. 1149 = crenulata ?
elliptica, *Mast. in Hook. f. Fl. Brit. Ind.* i. 377.—Java.
elliptica, Pohl, Pl. Bras. ii. 71 = melastomaefolia.
flaccida, *Spach, Ic.* n. 18 ; *in Linnaea*, xv. (1841) 174.—Ins. Ymor.
fulva, *Presp. & Endl. Nov. Gen. et Sp.* iii. 37.—Bras.
Gayana, *St. Hil. Fl. Bras. Mer.* i. 146.—Bras.
geminiflola, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 153.—Ecuador.
genistella, *Triana & Planch. in Ann. Sc. Nat. Sér.* IV. xvii. (1863) 338.—N. Granat.
glabrescens, *Benth. Bot. Voy. Sulph.* 71.—Am. trop.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 502.—Madag.
grandifolia, DC. Prod. i. 486 = aspera.
herbacea, *Nees, Fl. Corom.* i. 25. t. 20.—Ind. or.
hermanniaefolia, J. Gay, in DC. Prod. i. 486 = Rullagia hermanniaefolia.
heterophylla, *Hook. Bot. Misc.* i. (1830) 291.—Madag.
hirsuta, *Ruiz & Pav. Fl. Per.* iii. 10.—Peruv.
hypoleuca, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 207 = Commersonia platyphylla.
incdora, J. Gay, ex DC. Prod. i. 486 = Rullagia pannosa.
Jacksiana, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 386.—Ins. Penang.
jucuilifolia, *Pohl, Pl. Bras.* ii. 77.—Bras.
javanica, Sprang. ex Miq. Illustr. Fl. Ind. Arch. 91 = nagalata.
laevigata, Schott, in Sprang. Syst. Veg. iv. App. 404 = Gayana.
Lescolaia, *Seem, ex DC. Prod.* i. 487.—Mexic.

BUETTNERIA, l.—
lanciflola, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 158.—Borneo.
lanciflola, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 32.—Bras.
lazarala, *Presl, Rel. Haenk.* ii. 145.—Mexic.
lobata, *Baill. in Bull. Soc. Linn. Par.* i. (1886) 503.—Madag.
longicuria, *Baill. l. c.* 502.—Madag.
longifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 154.—Venezuela.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* v. 215.—N Granat.
Maingayi, *Mast. in Hook. f. Fl. Brit. Ind.* i. 377.—Malacca.
melastomaefolia, *St. Hil. Fl. Bras. Mer.* i. 144.—Bras
melastomoides, G. Don, Gen. Syst. i. 824 = pract.
microphylla, *Vacq. Hort. Vindob.* i. 10. t. 29 ; *Linn. Mant.* 309.—Ind. or.
mollis, *H. B. & K. Nov. Gen. et Sp.* v. 214. t. 481 a et b.—N. Granat.
morifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér IV. xvii. (1863) 332.—N. Granat.
nepalensis, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 207 = aspera.
oblique, *Benth. in Hook. Journ. Bot.* iv. (1842) 194.—Guiana.
oblongata, *Pohl, Pl. Bras.* ii. 78.—Bras.
ovata, *Lam. Encyc.* i. 523.—Peruv.
pannosa, Bauer, ex Sheetz, in Lahm. Pl. Preiss. ii. 350 = Keraudrenia Hookeriana.
pannosa, DC. Prod. i. 486 = Rullagia pannosa.
parvifiora, *Benth. Pl. Hartw.* 114.—Ecuador.
peruviana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 207.—Peruv.
pilosa, *Roxb. Hort. Beng.* 18 ; *Fl. Ind.* i. 618.—Ind or. ; Burma.
Pohliana, *Steud. Nom.* ed. II. i. 249.—Bras.
polycantha, Wall. Cat. n. 1151 = crenulata ?
ramosissima, Pohl, Pl. Bras. ii. 78 = scabra.
Reinwardtii, *Korth. in Nederl. Kruidk. Arch.* (1848) i. 310.—Borneo.
reticulata, Cav. Diss. 292, sphalm. = tereticaulis.
rhamnifolia, *Benth. in Hook. Kew Journ.* iii. (1851) 164.—Bras.
rivularis, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 338.—Bras.
rubricaulis, *Presl, Rel. Haenk.* ii. 145.—Mexic.
sagittaefolia, *St. Hil. Fl. Bras. Mer.* i. 143. t. 27.—Bras
salicifolia, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 470.—Venezuela.
salicifolia, *Presl, Rel. Haenk.* ii. 144.—Mexic.
scabra, *Linn. Syst. ed.* X. 197.—Am. austr.
scabra, Pohl, Pl. Bras. ii. 81 = Pohliana.
scaipellata, Pohl, l. c. 78 = scabra.
scorpiura, *Wright, ex Griseb. Cat. Pl. Cub.* 29.—Cuba.
sidaefolia, St. Hil. Fl. Bras. Mer. i. 146 = catalpaefolia.
sulcata, *Ruiz & Pav. Fl. Per.* iii. 10.—Peruv.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 68.—Ins. Tahiti.
tereticaulis, *Lam. Encyc.* i. 528.—Peruv.
tiliaefolia, *Presl, Rel. Haenk.* ii. 144.—Mexic.
uncinata, *Mast. in Hook. f. Fl. Ind.* i. 377.—Malacca.
velutina, Wall. Cat. nn. 1148, 1925 = pilosa.
virgata, Pohl, Pl. Bras. ii. 80 = scabra.
vitifolia, *Baill. Adansonia*, x. (1872) 177.—Madag.
Vouilly, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 502.—Madag.

BUFFONIA, Koch, Syn. Fl. Germ. ed. I. 108 (1836) = **Buffonia**, Linn. (Caryophyll.).

BUFONIA, Sauv. ex Lam. Dict. Encyc. Bot. Dupl. 3 (1747) ; Annon. Acad. i. 286 (1749). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 151.
alsinoides, Hill, Veg. Syst. xi. 27 = annua ?
angustifolia, Cav. ex Willk. & Lange, Prod. Fl. Hisp. iii. 616 = tenuifolia.
annua, DC. Fl. Fr. iv. 768 = tenuifolia et macrosperma.
brachyphylla, *Boiss. & Heldr. Diagn. Ser. I.* viii. 94.—Graecia et Ins.
calycinata, *Boiss. & Bal. l. c. Ser.* II. v. 58.—Cilicia.
Davallieri, *Batt. & Trab. in Bull. Soc. Bot. Fr.* xxvi. (1879) 56.—Afr. bor.
elata, *Boiss. Diagn. Ser.* I. i. 44 = macrocarpa.
exserta, *Boiss. l. c.*—Persia.

BUFONIA :—
hebecalyx, *Boiss. Diagn.* Ser. I. i. 45.—Persia.
Kotschyana, *Boiss. l. c.* Ser. II. i. 84.—Persia.
macrocarpa, *Ser. in DC. Prod.* i. 388.— Persia; Afghan.
macropetala, *Willd. in Flora*, xxxiv. (1851) 604 ; *et lc. Pl. Hisp.* 119.—Hispan.
macrosperma, *J. Gay, in Mutel, Fl. Dauph.* ed. II. 88. —Europ.; As. Min. ; Syria.
multiceps, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 259.—Arabia.
Olivieriana, *Ser. in DC. Prod.* i. 388.—Mesopot. ; Persia : Afghan.
paniculata, Dabois, in Delarb. Fl. Auvergne, ed. II. 300 = tenuifolia.
parviflora, Griseb. Spicil. Fl. Rumel. i. 197 = tenuifolia.
perennis, *Pourr. in Mém. Acad. Toul.* iii. (1788) 309.— Gallia.
rotundifolia, Buch.-Ham. ex Steud. Nom. ed. II. i. 235 = Drymaria cordata.
spicata, Delarb. Fl. Auvergne, ed. II. 299 = macrosperma.
tenuifolia, *Linn. Sp. Pl.* 123. — Europ. austr. ; Marocco ; Reg. Caucas.
tenuifolia, Moq. & Sessé, ex DC. Prod. i. 390 = Hymenella mochringioides
tenuifolia, Vill. Hist. Pl. Dauph. iii. 650 = macrosperma.
virgata, *Boiss. Fl. Orient.* i. 666.—As. Min. ; Syria.
Willkommiana, Boiss. Diagn. Ser. II. i. 83 = perennis.

BUFORRESTIA, C. B. Clarke, in DC. Monog. Phan. iii. 333. t. 6, 7 (1881). *COMMELINACEAE*, Benth. & Hook. f. Gen. iii. 856.
Candolleana, *C. B. Clarke, l. c.* 333.—Guiana.
imperforata, *C. B. Clarke, l. c.* 334.—Afr. trop.
Mannii, *C. B. Clarke, l. c.* 333.—Afr. trop.
tenuis, *C. B. Clarke, l. c.* 334.—Afr. trop.

BUGENVILLEA, Endl. Gen. 312 (1837) = seq.

BUGINVILLAEA, Comm. ex Juss. Gen. 91 (1789) = **Bougainvillaea**, Juss. (Nyctag.).

BUGLOSSA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 351 (1821) = **Lycopsis**, Linn. (Boragin.).
arvensis, S. F. Gray, l. c. = L. arvensis.
sempervirens, S. F. Gray, l. c. 352 = Anchusa sempervirens.

BUGLOSSITES, Morz, in Enum. Sem. Hort. Taur. (1845).—Cf. Ann. Sc. Nat. Sér. III. v. (1846) 365 = **Borago**, Linn.
laxiflora, Moris, l. c. = Borago laxiflora.

BUGLOSSOIDES, [Riv.]. Moench, Meth. 418 (1794) = **Lithospermum**, Linn. (Borag.).
ramosissima, Moench, l. c. = L. tenuiflorum.

BUGLOSSUM, [Tourn.]. Adans. Fam. ii. 178 (1763) ; Gaertn. Fruct. i. 332. t. 67 (1788) = **Anchusa**, Linn. (Boragin.).
amoenum, Gaertn. l. c. = A. italica.
angustifolium, All. Fl. Pedem. i. 47 = A. undulata.
angustifolium, Moench, Meth. 418 = A. angustifolia.
asperrimum, Gilib. Fl. Lituan. i. 91 = Anchusa officinalis.
Barrelieri, All. Fl. Pedem. i. 48 = A. Barrelieri.
caeruleum, Pers. Syn. i. 199 = A. italica.
echinatum, Tausch, in Flora, vii. (1824) 280 = A. strigosa.
elatum, Moench, Meth. 418 = A. italica.
italicum, Tausch, in Flora, vii. (1824) 280 = A. italica.
officinale, Lam. Fl. Fr. ii. 278 = A. officinalis.
paniculatum, Tausch, in Flora, vii. (1824) 238 = A. italica.
sempervirens, All. Fl. Pedem. i. 48 = A. sempervirens.
syriacum, Tausch, in Flora, vii. (1824) 281 = A. strigosa.
tinctorium, Lam. Fl. Fr. ii. 278 = Alkanna tinctoria.
vulgare, Tausch, in Flora, vii. (1824) 229 = A. italica.

BUGRANOPSIS, Pomel, Nouv. Mat. Fl. Atl. 171 (1874) = **Cneorella**, Lhé. (Leguma.).
alopecuroides, Pomel, l. c. 173 = O. alopecuroides.
cephalanthus, Pomel, l. c. = O. cephalanthus.

BUGRANOPSIS :—
crinita, Pomel, Nouv. Mat. Fl. Atl. 171 = O. crinita.
euphrasiaefolia, Pomel, l. c. 172 = O. euphrasiaefolia.
megalostachys, Pomel, l. c. 171 = O. megalostachys.
mitissima, Pomel, l. c. 172 = O. mitissima.
rosea, Pomel, l. c. 171 = O. roseo.
Salemaniana, Pomel, l. c. 172 = O. Salzmanniana.
stricta, Pomel, l. c. = O. stricta.
Tournefortii, Pomel, l. c. = O. Tournefortii.
variegata, Pomel, l. c. = O. variegata.

BUGULA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Ajuga**, Linn. (Labiat.).
alpina, Alt. Fl. Pedem. i. 45 = A. genevensis.
arvensis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 368 = A. Chamaepitys.
Chamaepitys, Scop. Fl. Carn. ed. II. i. 416 = A. Chamaepitys.
decumbens, Mill. Gard. Dict. ed. VIII. n. 2 = A. reptans.
densiflora, Benth. in DC. Prod. xii. 595, in syn. = A reptans.
genevensis, Crantz, Inst. i. 516 = A. genevensis.
nana, Gilib. Fl. Lituan. i. 94 = A. pyramidalis.
obliqua, Moench, Meth. 381 = A. orientalis.
orientalis, Crantz, Inst. i. 518 = A. orientalis.
pyramidalis, Crantz, l. c. = A. pyramidalis.
reptans, Crantz, l. c. = A. reptans.
tomentosa, Gilib. Fl. Lituan. i. 94 = A. genevensis.
villosa, Steud. Nom. ed. II. i. 235 = A. pyramidalis.

BUHSIA, Bunge, Del. Sem. Hort. Dorpat. (1859) 41.—Cf. Linnaea, xxx. (1859–60) 752 = **Cleome**, Linn. (Capparid.).
coluteoides, Bunge, l. c. = C. coluteoides.

BUINALIS, Rafin. New Fl. Am. iv. 40 (1836) = **Paronychia**, Linn. (Illecebrac.).
floridana, Rafin. l. c. = P. Baldwinii.

BUJACIA, E. Mey. Comm. Pl. Afr. Austr. 127 (1835) = **Glycine**, Linn. (Legumin.).
anonychina, E. Mey. l. c. = G. javanica.
genuina, Benth., E. Mey. l. c. = Teramnus labialis.

BULBEDULIS, Rafin. Fl. Tellur. ii. 26 (1836) = **Camassia**, Lindl. (Liliac.).

BULBILIS, Rafin. in Am. Monthly Mag. (1819) 190, nomen prius ; et in Journ. Phys. lxxxv. (1819) 262 = **Buchloë**, Engelm. (Gramin.).
dactyloides, Rafin. in Am. Monthly Mag. (1819) 190 = Buchloe dactyloides.

BULBILLARIA, Zucc. in Abh. Akad. Muench. iii. (1848) 229. t. 2 = **Gagea**, Salisb. (Liliac.).
gageoides, Zucc. l. c. = G. persica.

BULBINE, Linn. Hort. Cliff. 123 (1737) ; Gen. Pl. ed. I. 95 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 784.
Blephanthera, Rafin. Fl. Tellur. iii. 59 (1836).
Nemopogon, Rafin. l. c. 37 (1836).
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 334 = asphodeloides.
acaulis, Linn. Sp. Pl. ed. II. 446 = alooides.
alooides, *Willd. Enum. Hort. Berol.* 372.—Afr. austr.
annua, *Willd. l. c.*—Afr. austr.
asiatica, Gaertn. Fruct. i. 42. t. 13 = Crinum asiaticum.
asphodeloides, *Spreng. Syst.* ii. 85.—Afr. austr.
australis, Spreng. Syst. ii. 86 = bulbosa.
bisulca, *Haw. in Phil. Mag.* (Feb. 1827) 121.—Afr. austr.
brevifolia, Schult. f. Syst. vii. 450 = Anthericum brevifolium.
bulbosa, *Haw. Rev. Pl. Succ.* 38.—Austral.
canaliculata, Spreng. Syst. ii. 86 = Anthericum canaliculatum.
Caudafolia, Schult. f. Syst. vii. 450 = Bulbinella caudata.
caudata, Spreng. Syst. ii. 86 = Bulbinella caudata.
caulescens, Linn. ex Steud. Nom. ed. I. 125 ; ed. II. i. 286 = frutescens.
ciliata, Link, Enum. Hort. Berol. i. 329 = Anthericum ciliatum.
crispa, Spreng. Syst. ii. 86 = Chlorophytum crispum.
densiflora, *Baker, in Journ. Linn. Soc.* xv. (1876) 347.—Afr. austr.

BULBINE :—
dubia, *Schult. f. Syst.* vii. 450.—Afr. austr.
falcata, Schult. f. l. c. 451 = Anthericum falcatum.
favosa, *Schult. f. l. c.* 444.—Afr. austr.
filifolia, *Baker, in Journ. Linn. Soc.* xv. (1876) 344. —Afr. austr.
floribunda, Schrad. ex Benth. Fl. Austral. vii. 35 = semiberbata.
frutescens, *Willd. Enum. Hort. Berol.* 372.—Afr. austr.
graminea, Haw. Revis. Pl. Succ. 38 = asphodeloides.
hispida, Sweet, Hort. Brit. ed. I. 416 = Anthericum hispidum.
Hookeri, Kunth, Enum. Pl. iv. 566 = bulbosa.
incurva, Spreng. Syst. ii. 86 = rostrata.
latifolia, *Spreng. l. c.*—Afr. austr.
laxiflora, *Baker, in Journ. Linn. Soc. xv.* (1876) 347. —Afr. austr.
longiscapa, *Willd. Enum. Hort. Berol.* 372.—Afr austr.
Mackenii, Hook. f. Bot. Mag t. 5955 = Eriospermum Mackenii.
macrophylla, Salm-Dyck, Hort. Dyck. 333 = latifolia.
mesembryanthemoides, *Haw. in Phil. Mag.* (July 1826) 31.— Afr. austr.
Mettingkii, Tenore, Syll. 563 = longiscapa.
minima, *Baker, in Journ. Linn. Soc. xv.* (1876) 344 —Afr. austr.
narcisifolia, *Salm-Dyck, Hort. Dyck.* 334.—Afr. austr.
nutans, *Spreng. Syst.* ii. 86.—Afr. austr.
nutans, Zeyh. ex Baker, in Journ. Linn. Soc. xv (1876) 347 = Zeyheri.
pallida, *Baker, in Journ. Bot.* xiv. (1876) 184.— Afr. austr.
planifolia, Spreng. Syst. ii. 86 = Simethis bicolor.
praemorsa, *Spreng. l. c.*—Afr. austr.
pugioniformis, *Link, Enum. Hort. Berol.* i. 329.—Afr austr.
rostrata, *Willd. Enum. Hort. Berol.* 372.—Afr. austr.
scabra, Spreng. Syst. ii. 86 = Anthericum scabrum.
semibarbata, *Haw. Revis. Pl. Succ.* 38.—Austral.
semibarbata, Hook. Bot. Mag. t. 3129 = bulbosa.
suavis, Lindl. Bot. Reg. (1838) Misc. 45 = bulbosa.
subbarbata, Steud. Nom. ed. II. i. 74, sphalm. = bulbosa.
triquetra, Spreng. Syst. ii. 86 = Bulbinella triquetra.
uncinata, Moench, Meth. 641 = Crinum americanum.
undulata, *Spreng. Syst.* ii. 86 = Anthericum undulatum.
unifolia, Schult. f. ex Baker, in Journ. Linn. Soc. xv. (1876) 368 = Eriospermum abyssinicum.
urginoides, *Baker, l. c.* 345.—Afr. austr.
Zeyheri, *Baker, l. c.* 347.—Afr. austr.

BULBINELLA, Kunth, Enum. Pl. iv. 569 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 784.
Chrysobactron, Hook. f. Fl. Antarct. i. 72. t. 44 (1845).
brevifolia, Kunth, l. c. 573 = Anthericum brevifolium.
Burkei, *Benth. & Hook. f. Gen.* iii. 784.—Afr. austr.
capillaris, Kunth, Enum. Pl. iv. 573 = triquetra.
caudata, *Kunth, l. c.*—Afr. austr.
ciliolata, *Kunth, l. c.* 570.—Afr. austr.
filiformis, Kunth, l. c. 572 = triquetra.
gracilis, *Kunth, l. c.* 571.—Afr. austr.
latifolia, *Kunth, l. c.* 572.—Afr. austr.
ornithogaloides, *Kunth, l. c.* 699.—Afr. austr.
peronata, *Kunth, l. c.* 570.—Afr. austr.
robusta, *Kunth, l. c.* 571.—Afr. austr.
setifolia, Kunth, l. c. 569 = triquetra.
squamea, Kunth, l. c. 573 = Anthericum hispidum.
triquetra, *Kunth, l. c.*—Afr. austr.

BULBISPERMA, Reinw. ex Blume, Cat. Gew. Buitenz. 59 (1823) = **Peliosanthes**, Andr. (Hæmod.).
ovigera, Reinw. l. c. = P. javanica.

BULBOCAPNOS, Bernh. in Linnaea, viii. (1833) 469 = **Corydalis**, Vent. (Papaverac.).
bracteatus, Bernh. l. c. = C. bracteata.
cavus, Bernh. l. c. = C. cava.
digitatus, Bernh. l. c. = C. Halleri.
fabaceus, Bernh. l. c. = C. fabacea.
Halleri, Spach, Hist. Vég. Phan. vii. 77 = C. Halleri.
longiflorus, Bernh. in Linnaea, viii. (1833) 469 = C. longiflora.

BULBOCAPNOS:—
Marschallianus, Bernh. in Linnaea, viii. (1833) 469 = C. Marschallianus.
nobilis, Bernh. l. c. = C. nobilis.
tuberosus, Spach, Hist. Vég. Phan. vii. 78 = C. cava.

BULBOCASTANUM, Lag. Amen. Acad. Madr. 99 (1821) = **Conopodium**, Koch (Umbellif.).
majus, Lag. l. c. = C. denudatum.

BULBOCASTANUM, Schur, Enum. Pl. Transs. 249 (1866) = **Carum**, Rupp. (Umbellif.).
denudatum, Nym. Comp. 304 = C. denudatus.
ferulaceum, Nym. l. c. 303.—Europ.
incrassatum, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 88 = C. incrassatus.
Linnæi, Schur, Enum. Pl. Transs. 249 = C. Bulbocastanum.
mauritanicum, Willk. in Linnæa, xl. (1876) 84 = C. mauritanicus.

BULBOCODIUM, Linn. Coroll. Gen. 6 (1737). *LILIACEÆ*, Benth. & Hook. f. iii. 821.
ABANDIUM, Adans. Fam. ii. 54 (1763).
alpinum, Mill. Gard. Dict. ed. VIII. n. 1 = Lloydia alpina.
atticum, Nym. Syll. 379 = Merendera attica.
autumnale, Lapeyr. Hist. Abr. Pl. Pyr. 202 = Merendera Bulbocodium.
balearicum, Nym. Syll. 379 = Merendera filifolia.
Broteri, Welw. ex Baker, in Journ. Linn. Soc. xvii. (1880) 441 = Merendera Bulbocodium.
caucasicum, Endl. ex Heynh. Nom. ii. 81 = Merendera caucasica.
colchicoides, Nym. Syll. 379 = Merendera Bulbocodium.
dentatum, *Schur, in Verh. Siebenb. Ver. Naturw.* ii. (1851) 166.—Transylv.
edentatum, Schur, in Oestr. Bot. Zeitschr. x. (1860) 355 = ruthenicum.
Eichleri, Regel, in Act. Hort. Petrop. v. (1877) 638 = Merendera Eichleri.
Fontanesii, *Heynh. Nom.* ii. 81.—Ins. Majore.; Algeria.
hastulatum, Frivald. in Flora, xix. (1836) 634 = Merendera sobolifera.
lusitanicum, *Heynh. Nom.* ii. 81.—Lusitania.
montanum, Fisch. Ind. Pl. Hort. Goreak. (1812) 10 = ruthenicum.
persicum, *Regel, Gartenfl.* xxx. (1881) 294. t. 1058.—Persia.
robustum, *Bunge, ex Regel, in Act. Hort. Petrop.* vi. (1879) 491.—Turkestan.
ruthenicum, *Bunge, in Del. Sem. Hort. Dorpat.* (1837) 11.—Transylv.; Reg. Caucas.
serotinum, Linn. Sp. Pl. 294 = Lloydia alpina.
sobolifèrum, Heynh. Nom. ii. 81 = Merendera sobolifera.
trigynum, Nym. Syll. 379 = Merendera sobolifera.
trigynum, Adam, in Weber & Mohr, Cat. i. 40; ex Schult. f. Syst. vii. 1524 = Merendera caucasica.
trigynum, Griseb. Spicil. Fl. Rumel. ii. 380 = Merendera sobolifera.
vernum, Bieb. Fl. Taur. Cauc. i. 261 = ruthenicum.
vernum, Desf. Fl. Atlant. i. 284 = Merendera filifolia.
vernum, Linn. Sp. Pl. 294 = ruthenicum.
versicolor, *Sprengl. Syst.* ii. 40.—Russia austr.

BULBOPHYLLARIA, S. Moore, in Baker, Fl. Maurit. 344 (1877), sphalm. = BOLBOPHYLLARIA, Reichb. f. = **Bulbophyllum**, Thou. (Orchid.).

BULBOPHYLLUM, Thou. Orch. Ins Afr. Tab. Esp. 8 et lc. t. 93-97 (1822). *ORCHIDEÆ*, Benth. & Hook. f. iii. 501.
ANISOPETALUM, Hook. Exot. Fl. t. 149 (1825).
BOLBOPHYLLARIA, Reichb. f. in Bot. Zeit. x. (1852) 934.
BOLBOPHYLLUM, Spreng. Syst. iii. 681 (1826).
COCHLIA, Blume, Bijdr. 320. t. 59 (1825).
DIDACTYLE, Lindl. Fol. Orch. (1852).
DIPHYES, Blume, Bijdr. 310. t. 56 (1825).
EPICRANTHES, Blume, l. c. 306. t. 9 (1825).
EPICRIANTHES, Blume, Fl. Jav. Pracf. p. vii (1828).
GERSINIA, Neraud, in Gaudich. Bot. Freyc. Voy. 27 (1826).
IONE, Lindl. Fol. Orch. (1853).
LYRAEA, Lindl. Gen. et Sp. Orch. 46 (1830).
MACROLEPIS, A. Rich. Sert. Astrol. 25. t. 19 (1833).
MALACHADENIA, Lindl. Bot. Reg. (1839) Misc. 67.

BULBOPHYLLUM:—
ODONTOSTYLES, Breda, Orch. Kuhl et Hass. t. sine n. (1827).
OXYEPAXA, Wight, Ic. v. 17. t. 1786 (1852).
SARCOSODIUM, Beer, Prakt. Orch. 306 (1854).
SESTOCHILOS, Breda, Orch. Kuhl et Hass. t. sine n. (1827).
TAURORAXIS, Reichb. f. in Bot. Zeit. x. (1852) 933.
TRIBRACHIA, Lindl. Bot. Reg. sub t. 832 (1824).
XIPHIDIUM, Reichb. f. in Bot. Zeit. x. (1852) 919.
abbreviatum, Reichb. f. in Gard. Chron. (1881) ii. 70 = Cirrhopetalum abbreviatum.
acutiflorum, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 18 (Cirrhopetali sp.).—Ind. or.
adenopetalum, *Lindl. Bot. Reg.* (1842) *Misc.* 85.—Malaya.
affine, *Lindl. in Wall. Cat.* n. 1982.—Reg. Himal.
alcicorne, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 151.—Burma.
alopecurum, *Reichb. f. in Gard. Chron.* (1880) ii. 70.—Burma.
amplebracteatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 307.—Malaya.
amplum, Reichb. f. in Walp. Ann. vi. 244 = Dendrobium amplum.
Andersonii, *Kurz, Rep. Andam.* 51, nomen.—Ins. Andam.
andongense, *Reichb. f. in Flora*, xlviii. (1865) 184.—Afr. occ.
angustifolium, *Lindl. Gen. et Sp. Orch.* 57.—Java.
antenniferum, *Reichb. f. in Walp. Ann.* vi. 250.—Ins. Philipp.
anteniniferum, Reichb. f. l. c. 248 = geminiflorum.
apetalum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 127.—Afr. trop.
apiferum, *Batem. ex Steud. Nom.* ed. II. i. 241.—Afr. occ.
appressum, *Reichb. f. in Linnæa*, xli. (1877) 42.—Ins. Norfolc.
aristatum, *Hemsl. Biol. Centr. Am. Bot.* iii. 213.—Am. centr.
atropurpureum, *Rodrig. Orch. Nov.* i. 41.—Bras.
aurantiacum, *F. Muell. Fragm.* iii. 39.—Austral.
aurantiacum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 219.—Afr. occ.
auratum, *Reichb. f. in Walp. Ann.* vi. 261.—Manilla.
auricomum, *Lindl. in Wall. Cat.* n. 1985.—Burma.
Bailey*i*, *F. Muell. Fragm.* ix. 5.—Austral.
balaeniceps, Reichb. f. in Hamb. Gartenz. xix. (1863) 208 = Napelli.
barbatum, *Rodrig. Orch. Nov.* ii. 119.—Bras.
barbigerum, *Lindl. Bot Reg.* xxiii. (1837) t. 1942.—Afr. trop.
Baroni, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 468.—Madag.
Beccarii, *Reichb. f. in Gard. Chron.* (1879) i. 41.—Borneo.
Berenicis, *Reichb. f. l. c.* xiv. (1880) ii. 588.—Hab.?
Betchei, *F. Muell. in Wing. South. Sc. Record*, i. (1881) 178.—Ins. Samoa.
bicolor, *Lindl. Gen. et Sp. Orch.* 49.—China.
bifarium, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 219.—Afr. trop.
bifforum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 397.—Java.
biseum, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 186.—Reg. Himal.
Blepharistes, *Reichb. f. in Flora*, lv. (1872) 278 (Cirrhopetali sp.).—Burma.
Bonkeitae, *F. M. Bailey, in Proc. Roy. Soc. Queensl.* i. (1884) 89.—Austral.
bootanense, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 158 = Cirrhopetalum bootanense.
Bowringianum, *Reichb. f. in Gard. Chron.* (1881) i. 814.—Assam.
brassicanum, *Reichb. f. in Linnæa*, xli. (1877) 117.—Ins. Philipp.
bracteolatum, *Lindl. Bot. Reg.* xxiv. (1838) 57.—Guinea.
Buchi, Reichb. f. in Walp. Ann. vi. 258 = Megaclinium Buchi.
caespiticum, *Thou. Orch. Afr.* t. 108.—Ins. Borbon.
calamarium, *Lindl. Bot. Reg.* (1848) *Misc.* 70.—Afr. trop.
capillipes, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 150.—Burma.
capitatum, *Lindl. Gen. et Sp. Orch.* 56.—Malaya.

BULBOPHYLLUM:—
Careyanum, *Spreng. Syst.* iii. 782.—Reg. Himal.
carinaeflorum, Reichb. f. in Walp. Ann. vi. 258 = biseum.
caudatum, *Lindl. in Wall. Cat.* n. 1983.—Reg. Himal
cernuum, *Lindl. Gen. et Sp. Orch.* 48.—Java.
Cheirl, *Lindl. Bot. Reg.* (1844) *Misc.* 44.—Manilla.
chinense, *Reichb. f. in Walp. Ann.* vi. 260.—China.
chloroglossum, *Reichb. f. in Gard. Chron.* (1871) 1195.—Bras.
chloropterum, *Reichb. f. in Linnæa*, xxii. (1849) 885.—Bras.
ciliatum, *Lindl. Gen. et Sp. Orch.* 48.—Java.
cirrhopetaloides, Griff. Notul. iii. 290 = biseum.
clandestinum, *Lindl. Bot. Reg.* (1841) *Misc.* 71.—Ind. or.; Malaya.
Clausseni, *Reichb. f. in Linnæa*, xix. (1847) 871.—Bras.
clavatum, *Thou. Orch. Afr.* t. 99.—Ins. Mascar.
clavigerum, *F. Muell. Census, Suppl.* i. 3.—Austral.
cochleatum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 125.—Afr. trop.
cocoinum, *Batem. ex Lindl. Bot. Reg.* xxiii. (1837) t. 1964.—Afr. trop.
colubrinum, Reichb. f. in Walp. Ann. vi. 257 = Megaclinium colubrinum.
comatum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 128.—Afr. trop.
Commersonii, *Thou. Orch. Afr.* t. 97.—Ins. Borbon.
compressum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 307.—Sumatra.
conchiferum, *Reichb. f. in Walp. Ann.* vi. 258.—Reg Himal.
conicum, Thou. Orch. Afr. Tab. Esp. 8 = clavatum
cornifera, Thou. l. c. t. 100 = clavatum.
cornutum, Reichb. f. in Walp. Ann. vi. 247 = Cirrhopetalum cornutum.
coriscense, *Reichb. f. Otia Bot. Hamb.* 118.—Afr. trop
crassifolium, *Thw. ex Trimen, in Journ. Bot.* xxii. (1885) 244.—Zeylan.
croceum, *Lindl. Gen. et Sp. Orch.* 57.—Java.
Cumingii, *Reichb. f. in Walp. Ann.* vi. 261.—Ins Philipp.
cupreum, Hook. Bot. Mag. t. 5316 = Careyanum.
cupreum, *Lindl. Bot. Reg.* (1838) *Misc.* 92.—Burma.
cuspidilligua, *Reichb. f. in Walp. Ann.* vi. 264.—Ins. Philipp.
cylindraceum, *Lindl. Gen. et Sp. Orch.* 53.—Reg Himal.
Dayanum, *Reichb. f. in Gard. Chron.* (1866) 484.—Burma.
delitescens, Hance, in Journ. Bot. xiv. (1876) 44 = Cirrhopetalum delitescens.
densum, *Thou. Orch. Afr.* t. 108.—Ins. Mascar.; Madag.
dianaea, *Lindl. in Journ. Linn. Soc.* vi. (1862) 125.—Afr. trop.
Dalliei, *Reichb. f. in Regel, Gartenfl.* (1885) 90.—Afr. trop.
elatidium, *Lindl. in Journ. Linn. Soc.* vi. (1862) 127.—Afr. trop.
elegans, *Gardn. ex Thw. Enum. Pl. Zeyl.* 298.—Zeylan.
Elisae, *F. Muell. Fragm.* vi. 130.—Austral.
Elitae, Reichb. f. in Walp. Ann. vi. 268 = Cirrhopetalum Wightii.
elongatum, *Hassk. Cat. Hort. Bog. Alt.* 39.—Malaya.
epiphytum, *Rodrig. Orch. Nov.* i. 40.—Bras.
erectum, *Thou. Orch. Afr.* t. 96.—Madag.
exiliphantum, *Reichb. f. in Walp. Ann.* vi. 252.—Reg. Himal.
exaltatum, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 186.—Guiana.
exiguum, *F. Muell. Fragm.* ii. 72.—Austral.
falcatum, Reichb. f. in Walp. Ann. vi. 258 = Megaclinium falcatum.
falciperalatum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 128.—Afr. trop.
fimbriatum, Reichb. f. in Walp. Ann. vi. 260 = Cirrhopetalum fimbriatum.
flavescens, *Lindl. Gen. et Sp. Orch.* 54.—Java.
flavidum, *Lindl. Bot. Reg.* (1840) *Misc.* 88.—Afr. trop.
foveiscrii, Par. ex Reichb. f. in Bot. Zeit. xxiii. (1865) 99 = auricomum.
fuscescens, Reichb. f. Walp. Ann. vi. 244 = Dendrobium fuscescens.
fusco-purpureum, *Wight, Ic.* v. t. 1661.—Ind. or.
fuscum, *Lindl. Bot. Reg.* (1839) *Misc.* 9.—Afr. trop.

BULBOPHYLLUM:—
Gaboniæ, *Lindon & Reichb. f. in Flora*, xlviii. (1865) 185.—Aff. oco.
galeatum, *Lindl. Gen. et Sp. Orch.* 57.—Aff. trop.
geneense, *Reichb. f. ex Rodrig. Orch. Nov.* ii. 125.—Bras.
gibbosum, *Lindl. Gen. et Sp. Orch.* 54.—Java.
gladiatum, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 186.—Bras.
gracile, *Lindl. Gen. et Sp. Orch.* 50.—Java.
gracile, Thou. Orch. Afr. t. 101 = Thouarsii.
gracile, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 159.—Burma.
grandiflorum, *Blume, Rumphia*, iv. 42.—N. Guinea.
grandiflorum, Griff. Notul. iii. 293 = reptans.
grmuloso, *Rodrig. Orch. Nov.* i. 41.—Bras.
gravidum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 126.—Afr. trop.
Griffithianum, Par. & Reichb f. in Trans. Linn. Soc. xxx. (1874) 158 = Cirrhopetalum gamosepalum.
Griffithii, *Reichb. f. in Walp. Ann.* vi. 247.—Reg. Himal.
Henshallii, Hort. ex Gard. Chron. (1852) 492 = Lobbii.
hermistoschyus, *Reichb f. in Walp. Ann.* vi. 254.—Afr. trop.
Hildebrandtii, *Reichb. f. Otia Bot. Hamb.* 74; *et in Bot. Zeit.* xxxix. (1881) 449.—Madag.
hirsutum, *Lindl. Gen. et Sp Orch.* 49.—Java.
hirtum, *Lindl. in Wall. Cat.* n. 1980.—Reg. Himal.
imbricatum, Griff. Notul. iii. 289 = cylindraceum.
imbricatum, *Lindl. Bot. Reg.* (1841) *Misc.* 87.—Afr. trop.
inæquale, *Lindl. Gen. et Sp. Orch.* 49.—Java.
incurvum, *Thou. Orch. Afr.* t. 95.—Ins Mascar ; Madag.
inera, *Reichb. f. in Gard. Chron.* (1880) i. 776.—Assam ?
inops, *Reichb. f. l. c.* (1880) ii. 620.—Hab.?
insectiferum, *Rodrig. Orch. Nov.* ii. 120.—Bras.
intertextum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 127.—Afr. trop.
javanicum, *Mig. Fl. Ind. Bat.* iii. 650.—Java.
kaduense, Reichb. f. in Walp. Ann. vi. 262 = Cirrhopetalum nilghernense.
khasyanum, Griff. Notul. iii. 284 = cylindraceum.
khasyanum, Reichb. f. in Trans. Linn Soc. xxx. (1874) 158 = repens.
ladanthum, *Lindl. in Gard. Chron.* (1855) 58.—Sumatra.
lasiochilum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 158.—Burma.
laxiflorum, *Lindl. Gen. et Sp. Orch.* 57.—Java.
lemniscatum, *Par. ex Hook. f. Bot. Mag.* t. 5961.—Burma.
leopardinum, *Lindl. in Wall. Cat.* n. 1981.—Reg. Himal.
lichenastrum, *F. Muell. Fragm.* vii. 60.—Austral.
limbatum, *Lindl. Bot. Reg.* (1840) *Misc.* 74.—Singapora.
limbatum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 159 (non Lindl.).—Burma.
Lindleyanum, *Grif. Notul.* iii. 287.—Burma.
Lobbii, *Lindl. Bot. Reg.* (1847) t. 29.—Burma; Malaya.
longiflorum, Thou. Orch. Afr. t. 98 = Cirrhopetalum Thouarsii.
longipes, *Reichb. f. in Walp. Ann.* vi. 258.—Burma.
Lucknithii, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 71.—Ins. Solomon.
Lundianum, *Reichb. f. & Warm. in Otia Bot. Hamb.* 94.—Bras.
lupulinum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 126.—Afr. trop.
Macraei, Reichb. f. in Walp. Ann. vi. 263 = Cirrhopetalum Wightii.
macranthum, *Lindl. Bot. Reg.* (1844) t. 13.—Singapora.
macrocnema, *Rodrig. Orch. Nov.* ii. 118.—Bras.
maculatum, *Rezadi, ex Nanez, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 236.—Ins. Philipp.
Makoyanum, *Reichb. f. in Gard. Chron.* (1879) t. 284 in note.—Bras.
mandibulare, *Reichb. f. l. c.* (1862) t. 866.—Borneo.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 218.—Afr. trop.
Mannii, Reichb. f. in Flora, lv. (1872) 275 (= Cirrhopetali sp.).—Reg. Himal.
maxillare, *Reichb. f. in Walp. Ann.* vi. 248.—Ins. Philipp.

BULBOPHYLLUM:—
maximum, Reichb. f. in Walp. Ann. vi. 259 = Megaclinium maximum.
Meduse, *Reichb. f. in Walp. Ann.* vi. 262.—Malaya.
megalanthum, *Grif. Notul.* iii. 286.—Malaya.
megalonyx, *Reichb. f. Otia Bot. Hamb.* 74; *et in Bot. Zeit.* xxxix. (1881) 449.—Ins. Comor.
melanorrhachis, Reichb. f. in Gard. Chron. (1875) ii. 162 = Megaclinium melanorrhachis.
Melleri, Reichb. f. in Flora, lxi. (1878) 78 = Megaclinium Mellerl.
membranaceum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 897.—Java.
mentosum, *Rodrig. Orch. Nov.* i. 42.—Bras.
mesgunese, Par. & Reichb f. in Trans. Linn Soc. xxx. (1874) 154 (= Cirrhopetali sp.).—Burma.
meridense, *Reichb. f. in Linnæa*, xxii. (1849) 836 — Venezuela.
micranthum, *Rodrig. Orch Nov.* i 39.—Bras.
microcaulon, *Lindl. in Journ. Linn Soc.* vi. (1861) 127.—Afr. trop.
microcaulon, *Rodrig. Orch. Nov.* ii. 121.—Bras.
microcaulon, *Reichb. f. in Walp. Ann.* vi. 258.—Burma.
minutum, *Thou. Orch. Afr* t 110.—Madag
minutissimum, *F. Muell. Fragm.* xi. 53.—Austral.
monilliforme, *F. Muell. in Gard. Chron.* (1878) ii. 818 —Austral.
monsiforme, R. King, in Gard. Chron. (1879) i. 55, 149 = minutissimum.
mosillifer, *Par. & Reichb. f. in Trans. Linn. Soc* xxx. (1874) 151 —Burma
monosepalum, *Rodrig. Orch. Nov.* ii. 121.—Bras.
monticolum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 219.—Afr. trop.
moulmeinense, Reichb. f. in Walp. Ann. vi. 249 = Trias oblonga.
mucronatum, *Lindl. Gen. et Sp. Orch.* 50.—Java.
mucronifolium, *Reichb. f. & Warm. in Otia Bot Hamb.* 94.—Bras.
multiflorum, *Ridley, in Journ Linn Soc.* xxi. (1885) 463.—Madag.
muscicolum, Reichb. f. in Flora, lv. (1872) 275 = Cirrhopetalum Wallichii.
mutabile, *Lindl. Gen. et Sp. Orch.* 48.—Java.
Napelli, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 185.—Bras.
nasutum, *Reichb. f. in Gard. Chron.* (1871) 148.—Burma.
neilgherrense, *Wight, Ic.* v. 1, t. 1680.—Ind. or.
nematopodum, *F. Muell. Fragm.* viii. 80.—Austral.
nutans, *Thou. Orch. Afr.* t. 107.—Ins. Mascar.; Madag.
nutans, *Reichb. f. in Walp. Ann.* vi. 260.—Ins. Philipp.
nutans, Thou. Orch. Afr. 63. t. 8. n. 2 = variegatum.
oblongum, Reichb. f. in Walp. Ann. vi. 249 = Trias oblonga.
obtusum, *Lindl. Gen. et Sp. Orch.* 56.—Java.
occidentale, Spreng. Syst. iii. 732 = Epidendrum polybulbon.
occlusum, *Ridley, in Journ. Linn. Soc.* xxi. (1885) 464.—Madag.
ocenlium, *Thou. Orch. Iles Afr.* t. 93.—Ins. Mascar.
oculatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 809.—Malaya.
odoratissimum, *Lindl.* in *Wall. Cat.* n. 1987.—Reg. Himal.; Burma.
odoratum, *Lindl. Gen. et Sp. Orch.* 54.—Java.
Oerstedii, *Hemsl. Biol. Centr. Am. Bot.* iii. 218.—Nicaragua.
oligoglossum, *Reichb. f. in Hamb. Gartenz.* xxi. (1885) 297.—Burma.
orecomedes, *Reichb. f. Otia Bot. Hamb.* 118.—Afr. trop.
ovalifolium, *Lindl. Gen. et Sp. Orch.* 49.—Java.
oxyglossum, Reichb. f. in Walp. Ann. vi. 258 = Megaclinium oxyptcrum.
pachyrrhachis, *Griseb. Fl. Brit. W. Ind.* 618.—Ind. occ.
Pahudi, *Reichb. f. in Walp. Ann.* vi. 264.—Malaya.
palmosum, *Benth. & Hook. f. Gen.* iii. 508.—Reg. Himal.
Pangerangi, *Reichb. f. in Bonplandia*, v. (1857) 57.—Malaya.
parviflorum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 152.—Burma.
parvulum, *Lindl. Gen. et Sp. Orch.* 49.—Java.

BULBOPHYLLUM:—
pavimentatum, *Lindl. in Journ Linn. Soc.* vi. (1861) 128.—Afr. trop
pendulum, *Thou. Orch Afr* t. 104.—Ins. Mascar.
Penicillium, *Par & Reichb. f in Trans. Linn. Soc.* xxx. (1874) 151 — Burma
petiolare, *Thw. Enum Pl. Zeyl.* 298.—Zeylan.
pictum, Par. & Reichb f. in Trans. Linn. Soc. xxx (1874) 150 = Drymoda picta.
picturatum, Reichb f. in Walp. Ann. vi. 262 = Cirrhopetalum picturatum.
pilentum, *Lindl. Bot. Reg.* (1844) *Misc.* 73.—Malaya.
Pipio, *Reichb. f. in Linnæa*, xli (1877) 92.—Afr. trop.
polyrrhizum, *Lindl. Gen. et Sp. Orch.* 53.—Reg Himal.
Prentiesi, *F. Muell. in Wing, South. Sc. Record*, i. (1881) 173.—Austral
primaticum, *Thou. Orch. Afr* t. 102.—Ins. Mascar.
psittacoglossum, *Reichb. f. in Bot Zeit.* xxi. (1863) 236.—Burma
Psychoon, *Reichb. f. in Gard. Chron.* (1878) ii. 170.—Assam.
Pumilio, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 153 = Cirrhopetalum Pumilio.
pumilum, *Lindl. Gen. et Sp Orch.* 54.—Afr. trop.
punctatum, *Fitzgerald, in Journ. Bot.* xxi. (1883) 205.—Austral.
punctatum, *Rodrig. Orch. Nov.* i. 40.—Bras.
purpurascens, *F. Muell. Bailey, in Proc. Roy Soc. Queensl.* i. (1884) 88.—Austral.
purpurascens, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 808.—Java.
purpureum, *Thw. Enum Pl. Zeyl* 298.—Zeylan.
pusillum, *Thou. Orch. Afr* t. 102.—Ins. Maurit.
pygmæum, *Lindl. Gen et Sp Orch* 58.—N. Zeland.
quadriseture, *Lindl. in Hook Lond. Journ. Bot.* ii (1843) 670.—Guiana
radiatum, *Lindl. in Wall. Cat.* n. 1986.—Burma; China.
recurvum, *Lindl. Gen. et Sp. Orch.* 53.—Afr trop.
refractum, Reichb. f. in Walp Ann. vi. 259 = Cirrhopetalum refractum.
Regnelli, *Reichb. f. in Linnæa*, xxii. (1849) 835.—Bras.
Reinwardtii, *Reichb. f. in Walp. Ann.* vi. 246.—Malaya.
repens, *Griff. Notul.* iii. 293.—Reg Himal.
repens, *Lindl. in Wall Cat.* n. 1988.—Reg. Himal.
reticulatum, *Batem. in Hook. f. Bot. Mag.* t. 5605.—Borneo.
retususculum, Reichb f. in Gard. Chron. (1869) 1182 = Cirrhopetalum retususculum.
Rhizophorae, *Lindl in Journ. Linn. Soc.* vi. (1861) 125.—Afr. trop.
rigens, *Reichb. f. in Hamb. Gartens.* xxi. (1885) 296.—Burma.
rostriceps, *Reichb. f. Otia Bot. Hamb.* 55.—Ins. Fiji
rotundatum, Reichb. f. in Walp. Ann. vi. 244 = Dendrobium rotundatum.
Roxburghii, Reichb. f. l. c. 263 = Cirrhopetalum Roxburghii.
rufinum, *Reichb. f. Xenia Orch.* iii. 46.—Burma.
rupicola, *Rodrig. Orch. Nov.* i. 39.—Bras.
rupicola, *Reichb. f. in Flora*, xlviii. (1865) 184.—Afr. trop.
salaccense, *Reichb. f. in Bonplandia*, v. (1857) 57.—Java.
saltatorium, *Lindl. Bot. Reg.* xxiii. (1837) t. 1970.—Afr. trop.
Sanderson, Reichb. in Flora, lxi. (1878) 78 = Megaclinium Sandersoni.
sarcoscapum, *Teijsm & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 242.—Java.
scabratum, Reichb. f. in Walp. Ann. vi. 259 = Cirrhopetalum cæspitosum.
Sceptrum, *Reichb. f. l. c.* 254.—Ins. Sunda.
Schilleriauum, Reichb. f. in Hamb. Gartenz. xvi. (1860) 423 = Shepherdi.
Schmidtianum, *Reichb. f. l. c.* xxii. (1866) 357.—Ind. or.
sciadanthum, *F. Muell. in Wing, South. Sc. Record.* ii. (1882) 95.—Ins. Samoa.
serpens, Lindl. Gen. et Sp. Orch. 57 = sutcnse.
setigerum, *Lindl. Bot. Reg.* (1838) *Misc.* 21.—Guiana.
Seychellarum, *Reichb. f. in Linnæa*, xli. (1877) 93.—Ins. Mascar.
Shepherdi, *F. Muell. Fragm.* iii. 40 —Austral.

BULBOPHYLLUM :—
siamense, Reichb. f. in Gard. Chron. (1867) 572 = Lobbii.
sicyobulbon, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 152.—Burma.
Sillemianum, *Reichb. f. in Gard. Chron.* (1884) ii. 166.—Burma.
simillimum, Par. & Reichb. f. Otia Bot. Hamb. 49 = Cirrhopetalum simillimum.
sordidum, Lindl. Bot. Reg. (1840) Misc. 89 = bracteolatum.
stenobulbon, *Par & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 153.—Burma.
striatum, *Reichb. f. in Walp. Ann.* vi. 257.—Reg. Himal.
suave, Griff. Notul. iii. 292 = birtum.
sulcatum, *Lindl. Gen. et Sp. Orch.* 57.—Java.
taeniophyllum, Par & Reichb. f. in Journ. Bot. xii (1874) 198 (= Cirrhopetalli sp.).—Moulmein
tahitense, *Nadeaud, Enum. Pl. Tahiti*, 36.—Ins. Pacif.
Taylori, *F. Muell. Fragm.* viii. 150.—Austral.
tenellum, *Lindl. Gen. et Sp. Orch.* 49.—Java.
tentaculigerum, Reichb. f. in Flora, lxi. (1878) 77 = Megaclinium tentaculigerum.
tenuisale, *Lindl. in Journ. Linn. Soc.* vi. (1861) 126. —Afr. trop.
tenuifolium, *Lindl. Gen. et Sp. Orch.* 50.—Java.
tetragonum, *Lindl. l. c.* 58.—Afr. trop.
Thompsoni, *Ridley, in Journ. Linn. Soc.* xxi. (1885) 464.—Madag.
Thouarii, *Stend. Nom.* ed. II. i. 214.—Ins. Maurit.
Thwaitesii, Par. & Reichb. f. in Journ. Bot. xii (1874) 199 (= Cirrhopetali sp.).—Zeylan.
tigridum, *Hance, in Journ. Bot.* xxi. (1883) 232.—China.
tortuosum, *Lindl. Gen. et Sp. Orch.* 50.—Java.
tremulum, *Wight, Ic.* v. t. 1749.—Ind. or.
triadenium, *Reichb. f. in Walp. Ann.* vi. 257.—Java.
trigonopus, Reichb. f. in Gard. Chron. (1881) ii. 71 = Cirrhopetalum abbreviatum.
tripetalum, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 185.—Bras.
tripedianus, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 154 = Cirrhopetalum refractum.
tristotum, Griff. Notul. iii. 283 = clandestinum.
truncatum, *Reichb. f. in Walp Ann.* vi. 258.—Burma.
tuberculatum, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 336.—N. Zel.
umbellatum, Lindl. in Wall. Cat. n. 1984 = Cirrhopetalum maculosum.
unguiculatum, *Reichb. f. in Linnaea*, xxii. (1849) 886. —Java.
uniflorum, *Griff. Notul.* iii. 293.—Reg. Himal.
uniflorum, *Hassk. Cat. Hort. Bogor. Alt.* 39.—Malaya.
vaginatum, Reichb. f. in Walp. Ann. vi. 261 = Cirrhopetalum vaginatum.
variegatum, *Thou. Orch. Afr.* t. 105, 106.—Ins. Borbon.
velutinum, Reichb. f. in Walp. Ann. vi. 258 = Megaclinium velutinum.
violaceum, *Lindl. Gen. et Sp. Orch.* 50.—Java.
vittatum, *Reichb. f. & Warm. in Otia Bot. Hamb.* 95. —Bras.
vittatum, *Teijsm. & Binn in Tydschr. Nederl. Ind.* xxiv. (1862) 308.—Java.
Wallichii, Reichb. f. in Walp. Ann. vi. 259 = Cirrhopetalum Wallichii.
Weddelii, *Reichb. f. l. c.* 251.—Bras.
Wightii, Reichb. f. l. c. 262 = Cirrhopetalum grandiflorum.
xylophyllum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 151.—Burma.

BULBOSPERMUM, Blume, Enum. Pl. Jav. 15 (1828) = Pellionanthes, Andr. (Hesmodor.).
javanicum, Blume, l. c. = P. javanica.

BULBOSTYLIS, DC. Prod. v. 138 (1836) = Brickellia, Ell. (Compos.).
annua, Nutt. in Journ. Acad. Philad. Ser. II. i. (1847) 179 = Psathyrotes annua.
californica, Torr. & Gray, Fl. N. Am. ii. 78 = Brickellia californica.
Cavanillesii, DC. Prod. v. 138 = Brickellia californica.
diffusa, DC. l. c. vii. 268 = Brickellia diffusa.
glabra, DC. l. c. v. 139 = Eupatorium Vauthierianum.
glabriuscula, C. Koch, in App. Sem. Hort. Berol. (1856) 13.—Am. trop.
hastata, Walp. Rep. Ind. p. 279 = Brickellia hastata.
hebecarpa, DC. Prod. v. 138 = Brickellia hebecarpa.

BULBOSTYLIS :—
lanata, DC. Prod. vii. 268 = Brickellia lanata.
microphylla, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 286 = Brickellia microphylla.
nepetaefolia, DC. Prod. v. 139 = Brickellia californica.
oliganthes, DC. l. c. = Brickellia oliganthes.
pauciflora, DC. l. c. = Eupatorium paciflorum.
pedunculosa, DC. in Deless. Ic. Sel iv. t. 17 = Eupatorium longipes.
pendula, DC. Prod. v. 138 = Brickellia pendula.
reticulata, DC. l. c. vii. 268 = Brickellia reticulata.
rigida, Hook. & Arn. Bot Beech. Voy. 297 = Brickellia lanata.
scorodonicefolia, Kunth, Ind. Sem. Hort. Berol. (1846) 13 = Brickellia californica ?
secundiflora, DC. Prod. v. 138 = Brickellia secundiflora.
spinaciaefolia, DC. l. c. 139 = Eupatorium spinaciaefolium.
subuligera, S. Schau. in Linnaea, xix. (1847) 718 = Barrocita subuligera.
triangularis, DC. Prod. vii. 268 = Carphephorus triangularis.
veronicaefolia, DC. l. c. v. 139 = Brickellia veronicaefolia.

BULBOSTYLIS, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 467 = **Eupatorium**, Tourn. (Compos.).
Cassiniana, Gardn. l. c. 472 = E. Vauthierianum.
deltoidea, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 456 = E. ageratifolium.
elegans, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 467 = E. elegans ?
glandulosa, Gardn. l. c. 469 = E. amygalinum.
micrantha, Gardn. l. c. 469 = E. amygdalinum.
microcephala, Gardn. l. c. 468 = E. amygdalinum.
oblongifolia, Gardn. l. c. 469 = E. trixolides.
pumila, Gardn. l. c. 470 = E. amphidictyum.
ramosissima, Gardn. l. c. = E. ramosissimum.
scandens, Gardn. l. c. = E. Vitalbae.
tomentosa, Gardn. l. c. 468 = E. amygdalinum.

BULBOSTYLIS, Kunth, Enum. Pl. ii. 205 (1837) = **Fimbristylis**, Vahl (Cyperac.).
capillaris, Nees, in Mart. Fl. Bras. ii. t. 84, in obs. = F. capillaris.
ciliata, Vahl, ex Nees, l. c. 84, in obs.—Hab. ?
consanguinea, Nees, l. c.—Bras.
hirtella, Nees, l. c. 85.—Bras.
laxiculm, Nees, l. c. 93.—Bras.
tenuifolia, Nees, l. c. 83.—Bras.
trifida, Nees, l. c. 84, in obs. = F. capillaris.

BULBOSTYLIS, Stev. in Mém. Soc. Nat. Mosc. v. (1814) 355 = **Blasocharis**, R. Br. (Cyperac.).
capitatus, Stev. l. c. = E. capitata.
geniculatus, Stev. l. c. = E. geniculata.
maculosus, Stev. l. c. = E. maculosa.
ovatus, Stev. l. c. = E. ovata.
palustris, Stev. l. c. = E. palustris.
tuberculatus, Stev. l. c. = E. tuberculosa.

BULLIARDA, DC. in Bull. Soc. Philom. iii. (1801) n. 69. Ja. = **Pilisa**, Mich. (Crassul.).
abyssinica, A. Rich. Tent. Fl. Abyss. i. 306 = T. aquatica.
alpina, Harv. in Harv. & Sond. Fl. Cap. ii. 330.—Afr. austr.
aquatica, DC. in Bull. Soc. Philom. (1801) n. 69. p. 2 = T. aquatica.
bonariensis, DC. l. c. = T. pedunculata.
brevifolia, Eckl. & Zeyh. Enum. 290.—Afr. austr.
caponis, DC. in Bull. Soc. Philom. (1801) n. 69. p. 2 = T. caponis.
decumbens, DC. l. c. = T. decumbens.
Dregei, Harv. in Harv. & Sond. Fl. Cap. ii. 330 = Crassula prostrata, *E. Mey.*
elatinoides, Eckl. & Zeyh. Enum. 290.—Afr. austr.
filiformis, Eckl. & Zeyh. l. c. = T. filiformis.
Linnaei, Spreng. Syst. i. 497 = T. aquatica.
magellanica, Comm. ex DC. in Bull. Soc. Philom. (1801) n. 69. p. 2 = T. moschata.
moschata, Urv. in Mém. Soc. Linn. Par. iv. (1826) 618 = T. moschata.
pedunculata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = T. Vaillantii.
perfoliata, DC. in Bull. Soc. Philom. (1801) n. 69. p. 2 = T. caponis.
prostrata, DC. l. c. = T. prostrata.

BULLIARDA :—
prostrata, Weinm. Fl. Petrop. 21 = T. aquatica.
recurva, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 472 [bis] = T. recurva.
Schkuhrii, Spreng. Syst. i. 498 = T. prostrata.
Sieberi, Spreng. Syst. iv. Cur. Post. 48 = T. verticillata
trichotoma, Eckl. & Zeyh. Enum. 290 = T. decumbens.
Vaillantii, DC. in Bull. Soc. Philom. (1801) n. 69. p 1 = T. Vaillantii.
Vaillantii, Schlmp. ex Britten, in Oliver, Fl. Trop. Afr ii. 388 = T. aquatica.

BULLIARDA, Neck. Elem. ii. 321 (1790) = **Unona**, Linn. f. et **Xylopia**, Linn. (Anonac.).

BULNESIA, C. Gay, Fl. Chil. i. 474. t. 15 (1846 *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 268.
GONOPTERA, Turcz. in Bull. Soc. Nat. Mosc. xx (1847) i. 150.
bonariensis, *Griseb. in Goett. Abh.* xix. (1874) 105.—Reg. Argent.
chilensis, *C. Gay, Fl. Chil.* i. 475.—Chili.
foliosa, *Griseb. in Goett. Abh.* xix. (1874) 106.—Reg. Argent.
macrocarpa, *Phil. Sert. Mend. Alt.* 9.—Chili.
Retamo, *Griseb. in Goett. Abh.* xix. (1874) 106.—Reg. Argent.
Sarmienti, *Lorentz, ex Griseb. l. c.* xxiv. (1879) 75.—Reg. Argent.
Schickendantzii, *Hieron. ex Griseb. l. c.*—Reg. Argent

BULOWIA, Hook. Bot. Mag. sub t. 4864 (1848) = BUELOWIA, Schur = **Smeathmannia**, Soland (Passifl.).

BULWERIA, F. Muell. Fragm. iv. 147 (1864) = **Diplanthera**, R. Br. (Pignon.).
nobilissima, F. Muell. l. c. = D. tetraphylla.

BUMALDA, Thunb. Nov. Gen. 62 (1783) = **Staphylea**, Linn. (Sapind.).
trifolia, Thunb. l. c. 63 = S. Bumalda.

BUMELIA, Sw. Prod. Veg. Ind. Occ. 49 (1788) *SAPOTACEAE*, Benth. & Hook. f. ii. 660.
SCLEROCLADUS et SYONDOGONA, Rafin. Sylva Tellur 35 (1838).
amoena, Tenore, ex A DC. in DC. Prod. v. 189 = lycioides.
angustifolia, *Nutt. Sylv. Am.* iii. 88. t. 98.—Am. bor., Mexic.
arachnoides, Rafin. New Fl. Am. iii. 28 = lanuginosa
arborea, Buckl. in Proc. Ac. Sc. Philad. 1861 (1862) 461 = lanuginosa.
argentea, Roem. & Schult. in Ges. Naturf. Fr. Berl Mag. iii. (1809) 67 ; Syst. iv. 499 = Heeria argentea
Axudus, *Roem. & Schult. Syst.* iv. 499.—Ins. b Doming.
borbonica, *Lodd. ex Loud. Hort. Brit.* 89.—Ins Borbon.
buxifolia, Roem. & Schult. Syst. iv. 802 = obtusifolia.
buxifolia, *Willd. ex H. B. & K. Nov. Gen. et Sp.* vi. t. 547.—Am. austr.
celastrina, *H. B. & K. l. c.* 212.—Mexic.
chrysophylloides, Wats. Dendrol. i. 10 = tenax.
crenulata, *Spreng. Syst.* i. 565.—Porto Rico.
Cruegerii, *Griseb. Fl. Brit. W. Ind.* 401.—Ind. occ.
cubensis, *Griseb. Cat. Pl. Cub.* 104.—Cuba.
cuneata, Sw. Fl. Ind. Occ. i. 496 = Sideroxylon obovatum, *Lam.*
cuneifolia, Rudge, Pl. Guian. i. 30. t. 47 = Chrysophyllum cuneifolium.
denticulata, *Rafin. New Fl. Am.* iii. 29.—Am. bor.
dulcifera, Schum. & Thonn. Beskr. Guin. Pl. 130 = Sideroxylon dulciferum.
Dumastii, *A. DC. in DC. Prod.* viii. 191.—Am. austr.
egregia, Poepp. ex A. DC. l. c. 192 = Sideroxylon rugosum.
excelsa, A. DC. l. c. 192 = obtusifolia.
feroz, *Chaun. & Schlecht. in Linnaea*, vi. (1831) 392. —Mexic.
ferruginea, Nutt. Sylv. Am. 88 = lanuginosa.
fosteriadenus, Willd. Sp. Pl. i. 1086 = Sideroxylon fosteriadenum.
glomerata, *Griseb. in Mem. Am. Acad. N. S.* viii. (1863) 518.—Ind. occ.
Hayesii, *Hemsl. Biol. Centr. Am. Bot.* ii. 297.—Guatemala.

BUMELIA :—
horrida, *Griseb. Cat. Pl. Cub.* 165.—Cuba.
laete-virens, *Hemsl. Biol. Centr. Am. Bot.* ii. 298.—Mexic.
lanuginosa, *Pers. Syn.* i. 237.—Am. bor.
lucida, Roem. & Schult. Syst. iv. 499 = Sideroxylon lucidum.
lycioides, *Willd. Enum. Hort. Berol.* 249.—Am. bor.
macrocarpa, Nutt. Sylva Am. iii. 87 = lanuginosa ?
Mangillio, Willd. Sp. Pl. i. 1087 = Myrsine Mangillia.
Mastichodendrum, Roem. & Schult. Syst. iv. 498 = Sideroxylon Mastichodendron.
micrantha, Willd. ex Roem. & Schult. l. c. 802 = Achlsis micrantha.
microphylla, *Griseb Cat. Pl. Cub.* 165.—Cuba.
montana, Sw. Prod. Veg. Ind. Occ. 49 = Dipholis montana.
monticola, *Buchl. in Bull. Torrey Club*, x. (1883) 91.—Texas.
multiflora, Roem. & Schult. Syst. iv. 498 (non Sapotaceæ).—Afr. austr.
myrsinifolia, A. DC. in DC. Prod. viii. 102 = Sideroxylon obovatum, *Lam.*
nervosa, Vahl, Eclog. Am. i. 28 = Lucuma Rivicoa.
nigra, Bert. ex Griseb. Fl. Brit. W. Ind. 400 = Dipholis salicifolia.
nigra, Sw. Prod. Veg. Ind. Occ. 49 = Dipholis nigra.
oblongifolia, Nutt. Gen. Am. i. 135 = lanuginosa.
obovata, *A. DC. in DC. Prod.* viii. 191.—Am. austr.
obtusifolia, *Roem. & Schult Syst.* vi. 802.—Bras.
obtusifolia, Willd. ex Steud. Nom ed. II. i. 837 = rotundifolia.
occidentalis, *Hemsl. Biol. Centr. Am. Bot.* ii. 298 — Mexic.
pallida, Sw. Prod. Veg. Ind. Occ. 49 = Sideroxylon Mastichodendron.
parvifolia, *A. DC. in DC. Prod.* viii. 190.—Ind. occ.
parvifolia, Chapm. Fl. S. U. St. 275 = angustifolia.
pauciflora, Engelm. ex A. Gray, Syn. Fl. ii. 68 = lanuginosa.
pauciflora, Roem. & Schult. Syst. iv. 498 = Sideroxylon pauciflorum.
pentagona, Sw. Prod. Veg. Ind. Occ. 50 = Dipholis salicifolia.
persimilis, *Hemsl. Biol. Centr. Am. Bot.* ii. 298.—Mexic.
pubescens, Tenore, Sem. Hort. Neap. (1837) = lycioides.
punctata, Roem. & Schult. Syst. iv. 498 = Myrsine Rapanea ?
rectinata, Torr. Bot. Mex. Bound. 109 = angustifolia.
reclinata, *Vent. Choix*, 23.—Am. bor.
retusa, Balbis, ex A. DC. Prod. viii. 194 = Ilicis sp.
retusa, *Sw. Prod. Veg. Ind. Occ.* 49.—Ind. occ.
rhamnoides, Cæsar, Stirp. Ibus. Dec. 64 = obtusifolia.
rotundifolia, H. B. & K. Nov. Gen. et Sp. iii. 287 = obtusifolia.
rotundifolia, *Sw. Prod. Veg. Ind. Occ.* 50.—Ind. occ.
rufa, *Raſin. New Fl. Am.* iii. 29.—Am. bor.
salicifolia, Bert. ex A. DC. in Trans. Linn. Soc. xvii. (1837) 108 = Myrsine salicifolia.
salicifolia, *Sw. Prod. Veg. Ind. Occ.* 50 = Dipholis salicifolia.
sartorum, *Mart. Horb. Fl. Bras.* 333.—Bras.
serrata, Pursh, Fl. Am. Sept. i. 155 = Prunus caroliniana.
serrulata, *Raſin. New Fl. Am.* iii. 29.—Am. bor.
spinifiora, *A. DC. Prod.* viii. 191.—Mexic.
spinosa, *A. DC. l. c.*—Mexic.
spinosa, S. Wats. in Proc. Am. Acad. xviii. (1883) 112 = lanuginosa.
strigosa, *Spreng. Syst.* i. 855.—Hab. ?
subcessiflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 299.—Mexic.
tenax, *Willd. Sp. Pl.* i. 1085.—Am. bor.
texana, *Buchl. in Bull. Torrey Club*, x. (1883) 90.—Texas.
tomentosa, A. DC. Prod. viii. 190 = lanuginosa.
tortuosa, *Wright, in Sauv. Fl. Cub.* 87.—Cuba.
undulata, *Raſin. New Fl. Am.* iii. 28.—Am. bor.

BUNBURIA, Harv. Gen. S. Afr. Pl. ed. I. 416 (1868) = **Vinetoxicum**, Moench (Asclep.).
elliptica, Harv. l. c. 417.—Afr. austr.

BUNBURYA, Meisn. ex Hochst. in Flora, xxvii. (1844) 560 = **Trichylus**, A. Rich. (Rubiac.).
capensis, Meisn. ex Hochst. l. c.—Afr. austr.

BUNCHOSIA, Rich. ex Juss. in Ann. Mus. Par. xvii. (1811) 481. *MALPIGHIACEÆ*, Benth. & Hook. f. i. 259.
MALACMARIA, Griseb. in Linnæa, xiii. (1839) 248.
acutifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 336.—Mexic.
angustifolia, *A. Juss. l. c.* 331.—Peruv.
argentea, *DC. Prod.* i. 582.—Venezuela.
armeniaca, *DC. l. c.*—Peruv.
bilocellata, Hemsl. Biol. Centr. Am. Bot. i. 147 = sec.
biocellata, *Schlecht. in Linnæa*, x. (1836) 241.—Mexic.
Bonplandiana, *A. Juss. in Arch. Mus. Par.* iii (1843) 333.—Am. trop.
bracteosa, Juss. in St. Hil Fl. Bras. Mer. iii 79 = Dicella bracteosa.
cancscens, *DC. Prod.* i. 582.—Ind. occ
cornifolia, *H. B. & K. Nov. Gen. et Sp.* v. 154.—N. Granat.
cumanensis, *G. Don, in Loud. Hort. Brit.* 170.—Venezuela.
deflexa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 310.—N. Granat.
discolor, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) 1. 266 = biocellata.
eglandulosa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 330.—Hab ?
elliptica, *Tod. Ind. Sem. Hort. Panorm.* (1877) 38.—Hab. ?
emarginata, *Regel, Ind. Sem. Hort. Petrop.* (1856) 35—Cuba.
fluminensis, *Griseb. in Mart. Fl. Bras.* xii i. 31.—Bras.
Gaudichaudiana, A. Juss. in Arch. Mus. Par. iii. (1843) 334 = fluminensis.
glandulifera, *H. B. & K. Nov. Gen. et Sp.* v. 154.—Venezuela.
glandulosa, *DC. Prod.* i. 581.—Am. trop.
glauca, *H. B. & K. Nov. Gen. et Sp.* v. 155.—N. Granat.
glauca, Seem. ex Hemsl. Biol. Centr Am. Bot. i. 147 = cornifolia.
grandifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 341.—Ins. Martinic.
Hartwegiana, *Benth. Pl. Hartw.* 164.—N. Granat
holosericea, Mart. ex A Juss. in Arch. Mus. Par. iii. (1843) 344 = Dicella holosericea.
Hookeriana, *A. Juss. l. c.* 337.—Peruv.
hypoleuca, *Mog. Stirp. Surin. Sel.* 77.—Guiana.
lanceolata, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) 1. 266 = discolor.
laxiflora, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 168 = Tetrapteris Guilleminiana.
Landeniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 335.—Am. trop.
Martiana, A. Juss. l. c. 334 = fluminensis.
media, *DC. P. l. i.* 581.—Ind. occ.
mollis, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 127.—Guiana.
mollis, Seem. ex Hemsl. Biol. Centr. Am. Bot. i. 146 = pilosa.
montana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 340.—Mexic.
multiflora, *Hook. f. Arn. in Hook. Bot. Misc.* iii. (1833) 157.—Am. austr.
nitida, *Juss. in Ann. Mus. Par.* xviii. (1811) 481.—Mexic.; Peruv.; Ind. occ.
odorata, Turcz. l. c.—N. Granat.
palustris, Poepp. ex A. Juss. in Arch. Mus. Par. iii. (1843) 311 = Burdachia prismatocarpa.
paniculata, DC. Prod. i. 582 = Tetrapteris ovæ'a.
pilosa, *H. B. & K. Nov. Gen. et Sp.* v. 154.—N. Granat.
polystachya, *DC. Prod.* i. 581.—Ind. occ.
retusa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 311.—N. Granat.
rhombifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 583.—Venezuela.
sessilifolia, *DC. Prod.* i. 582.—Mexic.
sphærocarpa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 318.—Bras.
squarrosa, Griseb. in Linnæa, xxii. (1849) 11 = Tetrapteryx squarrosa.
strigosa, *Schlecht. in Linnæa*, x. (1836) 242.—Mexic.
Swartziana, *Griseb. Fl. Brit. W. Ind.* 115.—Jamaic.
tuberculata, *DC. Prod.* i. 581.—Venezuela.

BUNGEA, C. A. Mey. Verz. Pfl. Canc. 106 (1831). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 976.
Sheareri, *S. Moore, in Journ. Bot.* xiii. (1875) 229.—China.
trifida, *C. A. Mey. Verz. Pfl. Canc.* 108.—As. Min.; Persia.
turkestanica, *Maxim. in Mém. Acad. Pétersb. Sér.* VII. xxix. No. 111. 61.—Turkestan.

BUNIAS, [Tourn.] Linn. Syst. ed. I. (1735). *CRUCIFERÆ*, Benth. & Hook f. i. 97.
ERUCAGO, Tourn ex Adans. Fam. ii. 428 (1763).
LAELIA, Adans. Fam. ii 428 (1763); Desv. Journ Bot. iii. (1813) 160.
ægyptiaca, Linn. Syst. Nat. ed. XII. 231 = Ochthodium segyptiacum.
altaica, Schang. ex DC. Syst. ii. 688 = Hutchinsia calycina.
americana, Rafin. Med. Fl. ii. 209 = Cakile maritima.
arvensis, *Jord. Cat. Gr. Yard. Dijon* (1848) 18.—Gallia.
aspera, Retz. Obs. ii. 21 = B. Erucago.
balearica, Linn. Syst. Nat. ed. XII. 446 = Succowia balearica.
Cakile, Linn. Sp. Pl. 670 = Cakile maritima.
cochlearioides, Bieb. Fl. Taur. Cauc. ii. 87 = Calepina Corvini.
cochlearioides, *Murr. in Comm. Gott.* viii. (1777) 42. t. 3.—Subsr. altaic.
cornuta, Linn. Sp. Pl. 669 = Fugionium cornutum.
crustata, Desv. Journ. Bot. iii. (1813) 168 = B. Erucago.
edentula, Bigel. Fl Bost. ed. III. 267 = Cakile maritima.
Erucago, *Linn. Sp. Pl.* 670.—Europ. austr.; As. Min.; Syria.
glomerata, Lapeyr. Hist. Abr. Pl. Pyr. 862 = Senebiera Coronopus.
littoralis, Salisb. Prod. 278 = Cakile maritima.
macropterа, Reichb. Fl. Germ. Excurs. 654 = B. Erucago.
myagroides, Linn. Mant. i. 96 = Erucaria aleppica.
orientalis, *Linn. Sp. Pl.* 670.—Europ. or.; As. Min.
ovalis, Viv. Fl. Lib. Spec. t. 16 = Cakile cyrenaica.
paniculata, L'Hérit ex DC. Syst. ii. 630 = Neslia paniculata.
paradoxa, Banks, ex DC. l. c. 677 = Chamira cornuta.
perennis, Moench, Meth. Suppl 91 = orientalis.
perennis, Sibth. & Sm. Fl. Græc. Prod. ii. 2 = Rapistrum perenne.
prostrata, Desf. Fl. Atlant. ii. 76 = Muricaria prostrata.
pyramidata, *Fisch. Hort. Gorenk.* ed. II. (1812) 49, nomen.—Hab. ?
raphanifolia, Sibth. & Sm. Fl. Græc. t. 612 = Rapistrum orientale.
rostrata, L'Hérit. ex DC. Syst. ii. 632 = Kucklidium syriacum.
silvicola, *Linn. Amœn. Acad.* viii. 8.—Sibir.
spinosa, Linn. Mant. 96 = Zilla myagroides.
syriaca, Gaertn. Fruct. ii. 290 = Euclidium syriacum.
tatarica, Willd. Sp. Pl. iii. 413 = Euclidium tataricum.
tcheliensis, *Debeaux, in Actes Linn. Soc. Bord.* xxxiii. (1879) 85.—China.
tenuifolia, Sibth. & Sm. Fl. Græc. Prod. ii. 8 (= *Didesmus tenuifolius*).—Græcia.
Tournefortii, Sibth. ex DC. Syst. ii. 659 (= *Didesmus tenuifolius*).—Græcia.
turcmica, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 82.—Hispan.
verrucosa, Moench, Meth. 277 = Rapistrum orientale.
virgata, Sibth. & Sm. Fl. Græc. t. 613 = Ochthodium segyptiacum.
vulgaris, Andrz. ex Steud. Nom. ed. II. i. 237, 238 = B. Erucago.
Winterii, Schult. Obs. 121 = orientalis.

BUNION, St. Lag. in Ann. Soc. Lyon, vii. 85 (1880) = seq.

BUNIUM, Koch, Syn. Fl. Germ. ed. II. 315 (1843) = **Pimpinella**, Riv. (Umbellif.).
montanum, Koch, l. c.—Europ.

BUNIUM, Linn. Gen. ed. I. 72 (1737) = **Carum**, Linn. (Umbellif.).
acaule, Bieb. Fl. Taur. Cauc. i. 212 = Chamascidium acaule.

BUNIUM :—
alpinum, Waldst. & Kit. Pl. Rar. Hung. ii. 199. t. 182 = C. alpinum.
ammoides, Cesati, in Linnaea, xi. (1837) 323 = C. ammoides.
ammoides, Link, ex Spreng. Syst. i. 908 = Conopodium denudatum.
aphyllum, Jan, ex DC. Prod. iv. 117.—Sicilia.
arcuatum, Griseb. ex Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 387.—Europ.
aromaticum, Linn. Mant. 218 = C. copticum.
brevifolium, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) Repr. 21.—Ins. Mader.
Bulbocastanum, Bert. ex DC. Prod iv. 668 = Diposis Bulbocastanum.
Bulbocastanum, Huds Fl. Angl. 122 = Conopodium denudatum.
Bulbocastanum, Linn. Sp Pl. 243 = C. Bulbocastanum.
bulbosum, Dulac, Fl. Hautes-Pyr. 351 = C. Bulbocastanum.
bulnioides, Boiss. & Reut. ex Boiss. Voy. Bot. App. 736.—Hispan.
capillifolium, Bertol. Fl. Ital. iii. 219.—Italia.
capillifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 428 = Carum setaceum.
Carvi, Bieb. Fl Taur. Cauc. i. 211 = C. Carvi.
carvifolium, DC. Prod. iv. 116 = C. flexuosum.
cazzium, Boiss. Diagn. Ser. I. x. 24 = C. elegans.
cilicicum, Fenzl, in Tchihat. Asie Min. Bot. i. 411 = C. Pestalozzae.
collinum, Albert, in Bull. Soc. Sc. Draguign. (1884) repr. 23.—Europ.
copticum, Spreng. Prod. Umb. 28 = C. copticum.
corydalinum, DC. Prod. iv 117.—Ins. Corsic.
creticum, Mill. Gard. Dict. ed. VIII. n. 2 = C. ferulaefolium.
creticum, Urv. in Bull. Soc. Linn. Par. i. (1822) 287 (= Conopodii sp.).—Graecia ; As. Min. ; Syria.
cynapioides, Bertol. Fl. Ital. iii. 217 = C. cylindricum ?
denudatum, DC. Fl. Fr. iv. 525 = Conopodium denudatum.
divaricatum, Bertol. Fl. Ital. iii. 222 (= Pimpinellae sp.).—Ital.
divaricatum, Cesati, in Linnaea, xi. (1837) 314 = C. ferulaefolium.
Falcaria, Bieb. Fl. Taur. Cauc. i. 211 = Falcaria Rivini.
ferulaceum, Sibth. & Sm. Fl. Graec. prod. i. 186 = C. ferulaefolium.
ferulaefolium, Desf. in Ann. Mus. Par. xi. (1808) 275 ; Choix, 55. t. 48 = C. ferulaefolium.
flexuosum, Brot. Fl. Lusit. i. 437 = Conopodium capillifolium.
flexuosum, Wrib. Bot. Arr. Brit. Pl. ed. II. 276 = Conopodium denudatum.
glaberrimum, DC. Prod. iv. 118.—Algeria.
glaucescens, DC. l. c. 117 (= *Scaligeria glaucescens*). —Armenia ; Persia.
glaucocarpum, Boiss. Diagn. Ser. I. x. 25 = C. Pestalozzae.
junceum, Margot & Reut. in Mém. Soc. Phys. Genèv. viii. (1839) 300 = C. creticum.
ligusticifolium, Cesati, ex Boiss. Fl. Orient. ii. 882 = C. multiflorum.
luteum, Hoffm. Gen. Umb. 108 = Muretia tanacetensis.
Macuca, Boiss. Elench. 44 ; Voy. Espagne, ii. 289. t. 66 = C. Macuca.
mayus, Bieb. Fl. Taur. Cauc. i. 210 = C. ferulaefolium.
majus, Gouan, Illustr. 10 = Conopodium denudatum.
majus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 514 = C. Bulbocastanum.
minus, Gouan, Illustr. 10 = C. Bulbocastanum.
minus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 514 = Conopodium denudatum.
minutifolium, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 177 = Feuseduanum vittijugum.
napiforme, Willd. ex Roem. & Schult. Syst. vi. 539 = C. creticum.
nepaleuse, Jacquem. ex Walp. Rep. ii. 423 = Chaerophyllum capnoides.
nivale, Boiss. Elench. 45 ; Voy. Espagne, ii. 240. t. 67 = C. nivale.
paucifolium, DC. Prod. iv. 117 = C. elegans.
Pestalozzae, Boiss. Diagn. Ser. I. x. 24 = C. Pestalozzae.

BUNIUM :—
petraeum, Loisel. Fl. Gall. ed. II. i. 195 = *corydalinum*.
petraeum, Tenore, Prod. Fl. Neap. p. lix = Carum alpinum.
peucedanoides, Bieb. Fl. Taur. Cauc i. 211 = Silaus carvifolius.
peucedanoides, DC. Prod. iv. 116.—Reg. Caucas.
pilosum, Willd. ex Roem. & Schult. Syst. vi. 500 = Conopodium denudatum.
pityranthon, Spreng. Syst. i. 901 = Pituranthus denudatus.
pumilum, Sibth. & Sm. Fl. Graec. iii. 67. t. 274 (= *Bunolattia pumila*).—Graecia.
pyrenaeum, Loisel. Fl. Gall. 161. t. 5 = Conopodium denudatum.
rupens, Spreng. in Roem. & Schult. Syst. vi. 500.—Afr. austr.
rigidulum, Caruel, Prod. Fl. Tosc. 272 = C. rigidulum.
saxatile, Mill. Gard. Dict. ed. VIII. n. 3 = C. alpinum.
strictum, Griseb. Spicil. Fl. Rumel. i. 344 = C. strictum.
subcarinum, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 14 = Conopodium subcarinum.
tenuifolium, Salzm. ex DC. Prod. iv. 118 = Conopodium capillifolium.
tenuissimum, Griseb. ex Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 267.—Europ.
ternatum, Pers. Syn. i. 308 = Conopodium denudatum.
thalictrifolium, Boiss. Elench. 45.—Hispan.
verticillatum, Gren. & Godr. Fl. Fr. i. 729 = C. verticillatum.
virescens, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 197 = Silaus carvifolius.

BUNNVA, F. Muell. Fragm. v. 86. t. 39 (1865) = **Cyanotaenia**, Turcz. (Verbenac).
cyanocalyx, F. Muell. l. c. = C. Bunnyana.

BUNOPHILA, Willd. ex Roem. & Schult. Syst. iii. Mant. 128, et Ind. (1827) (Ord. Nat.?).
PUNOPHILA, Willd. l. c. 8 (1827).
lycioides, Willd. l. c. 128.—Am. austr.

BUONAPARTEA, G. Don, in Sweet, Hort. Brit. ed. III. 706 (1839) = BONAPARTEA, Ruiz & Pav. = **Tillandsia**, Linn. (Bromel.).

BUPHANE, Herb. in Bot. Mag. sub t. 2578 (1825) ; et Amaryll. 289. t. 22, 23. f. 7, 8 (1826) ; prius Boophane, ejusd. App. 18 (1821) ; cetus Buphone in Bot. Mag. sub t. 2593 in nota (1825) *AMARYLLIDEAE*, Benth. & Hook. f. ii. 730.
CROSSYNE, Salisb. Gen. Pl. Fragm. 116 (1866).
amarythides, M. Roem. Syn. Ensat. 61 = ciliaris.
angolensis, Baker, in journ. Bot. xvi. (1878) 197 = Crinum angolense.
ciliaris, Herb. (Boophane).—Afr. austr.
distichа, *Herb. in Bot. Mag.* sub t. 2578.—Afr. austr.
guttata, Herb. Amaryll. 240. t. 22. f. 1 = ciliaris.
intermedia, M. Roem. Syn. Ensat. 59 = distichа.
toxicaria, Herb. App. 18 (Boophane) = distichа.

BUPHTHALMUM, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 388.
ATHALMUM, Neck. Elem. i. 20 (1790).
BUSTIA, Adans. Fam. ii. 127 (1763).
MOLPADIA, Cass. Dict. Sc. Nat. xxiii. 565 (1822).
TELEKIA, Baumg. Fl. Transylv. iii. 149 (1816).
angustifolium, Pursh, Fl. Am. Sept. ii. 564 = Marshallia angustifolia.
aquaticum, Linn. Sp. Pl. 908 = Odontospermum aquaticum.
arabicum, Delile, in Laborde, Voy. Arab. Pétr. 84 = Anvillea Garcini.
arborescens, Linn. Syst. ed. X. 1227 = Borrichia arborescens.
arvense, *Vell. Fl. Flum.* viii. t. 128.—Bras.
asperrimum, Spreng. Neue Entdeck. ii. 140 (= *Anomostephium buphthalmoides*).—Guadeloup.
asperum, Hornem. Hort. Hafn. Suppl. 102 = Wedelia hispida.
asteriscoides, Baill. Hist. Pl. viii. 301 = Osmites asteriscoides.
astwoidеum, Viv. Fl. Lib. Spec. 57. t. 25. f. 2 = Pallenis spinosa.
australe, Spreng. Fl. Hal. Mant. 58 = Wedelia aristata.

BUPHTHALMUM :—
aureum, Salzm. ex Steud. Nom. ed. II. i. 288 = Pallenis spinosa.
bipinnatum, Gilib. Fl. Lituan. i. 218 = Anthemis tinctoria.
bonariense, Pers. Syn. ii. 474 = Acicarphos spathulata ?
canum, L'Hérit. ex DC. Prod. v. 489 = Borrichia argentea.
capense, Linn. Syst. ed. X. 1227 = Cedera prolifera.
chrysanthemoides, Poir. Encyc. Suppl. i. 747 = Odontospermum pinifolium.
cordifolium, Waldst. & Kit. Pl. Rar. Hung. ii. 117 = speciosum.
coriaceum, Hort. ex Lond. Hort. Brit. 357 = Jasonia lacvigata.
crassifolium, Rafin. Specch. 191 = Odontospermum maritimum.
dentatum, Baill. Hist. Pl. viii. 301 = Osmites dentata.
difusum, Vahl, ex DC. Prod. v. 491 = Eclipta erecta.
durum, Linn. Sp. Pl. ed. II. 1275 = Osteospermum spinosum.
elegans, Brouss. ex Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 138 = Odontospermum imbricatum.
equisum, Vell. Fl. Flum. viii. t. 131.—Bras.
flexile, *Bertol. Fl. Ital.* ix. 413.—Ital.
flosculosum, Vent. Jard. Cels, 25 = Anvillea Garcini *frutescens*, Linn. Sp. Pl. 903 = Borrichia frutescens.
Garcini, Burm. f. Fl. Ind. t. 60 = Anvillea Garcini.
grandiflorum, Linn. Sp. Pl. 904 = salicifolium.
graveolens, Forsk. Fl. Aegypt. Arab. 151 = Odontospermum graveolens.
graveolens, Schimp. ex Boiss. Fl. Orient. iii. 880 (= *Asteriscus Schimperi*).—Arab.
Gussoni, *Pasquale, Notizie*, (1878) 2.—Sicil. ?
helianthoides, Forst. f. Prod. 57 = Wedelia aristata.
helianthoides, Linn. Sp. Pl. 904 = Heliopsis laevis.
heterophyllum, Willd. ex DC. Prod. v. 484 = Wedelia pedunculosa.
imbricatum, Cav. in Anal. Cienc. Nat. iv. (1801) 94 = Odontospermum imbricatum.
incanum, Mill. Gard. Dict. ed VIII. n. 10 = Borrichia frutescens.
inuloides, *Moris, Sard. App.* (1826).—Sardinia.
laevigatum, Brouss. ex Willd. in Ges. Naturf. Fr Berl. Mag. i. (1807) 138 = Jasonia laevigata.
leucifolium, *Burm. f. Fl. Cap. Prod.* 27.—Afr. austr
linearе, Willd. Sp. Pl. iii. 2280 = Borrichia frutescens.
littorale, *Vell. Fl. Flum.* viii. t. 124.—Bras.
longipes, *Comm. ex Cass. in Bull. Soc. Philom.* (1822) 144.—Madag.
maritimum, Linn. Sp. Pl. 908 = Odontospermum maritimum.
mediterraneum, *Vell. Fl. Flum.* viii. t. 135.—Bras.
melissoides, Poir, Encyc. Suppl. i. 747 = Heliopsis scabra.
melitense, *Forsk. Fl. Aegypt. Arab.* 218, nomen.—Ins. Melita.
mutans, Vitm. Summa Veg. v. 128 = Bidens cernua.
odorum, Schousb. Vexr. Marokko, 199 = Odontospermum odorum.
oleraceum, *Lour. Fl. Cochinch.* ii. 506.—China.
Ortega, *Spreng. Nachr. i. Bot. Gart. Halle*, 16, nomen.—Hab. ?
peruvianum, Lam. Encyc. i. 515 = Borrichia peruviana.
pratense, Vahl, Symb. Bot. i. 75 = Cervana pratensis.
procumbens, Desf. ex Steud. Nom. ed. I. 197 = Wedelia carnosa.
ramosum, *Forsk. Fl. Aegypt. Arab.* 151.—Arab.
Remirlio, Banch.-Ham. ex Wall. Cat. sub n. 3191 = Guizotia oleifera.
repens, Lam. Encyc. i. 515 = Wedelia carnosa.
sagittatum, Pursh, Fl. Am. Sept. ii. 564 = Balsamorhiza sagittata.
salicifolium, *Linn. Sp. Pl.* 904.—Europ. austr.
scabrum, Cav. Ic. ii. 58 = Wedelia hispida.
scandens, *Vell. Fl. Flum.* viii. t. 126.—Bras.
scandens, Schum. & Thonn. Beskr. Guin. Pl. 392 = Melanthera Brownei.
sericeum, Linn. f. Suppl. 379 = Odontospermum sericeum.
sessile, Mill. Gard. Dict. ed. VIII. n. 5 = Odontospermum maritimum.
silphioides, *Hornem. ex Steud. Nom.* ed. II. i. 289.—Hab. ?
speciosissimum, *Ard. Spec.* i. 26 ; *Linn. Mant.* i. 117.—Europ.
speciosum, *Schrad. Diss.* 11. t. 6.—Orient.
spinosum, Linn. Sp. Pl. 904 = Pallenis spinosa.

BUPHTHALMUM :—
stenophyllum, Link, in Buch, Beschr. Canar. Ins. 150 = Odontospermum sericeum.
strigosum, Spreng. Neue Entdeck. ii. 140 = Wedelia pedunculosa.
succisæfolium, Britting. in Flora, xxiv. (1841) i. 188 = salicifolium.
tenuifolium, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
trifolium, Poepp. ex DC. Prod. v. 577 = Oyedaea buphthalmoides.
umbrosum, Forst. f. Prod. 541 = Wedelia Forsteriana.

BUPLEUROIDES, [Boerh.] Moench, Meth. 294 (1794) = **Phyllis**, Linn. (Rubiac.).
Nobla, Moench, l. c. = P. Nobla.

BUPLEURUM. [Tourn.] Linn. Syst. ed. I (1735) *UMBELLIFERÆ*, Benth. & Hook. f. i. 886.
AGROSTANA, Hill, Veg. Syst. v. 32 (1772).
BUPRESTIS, Spreng. Pl. Umb. Prod. 27 (1813).
DIAPHYLLUM, Hoffm. Gen. Umb. 115 (1814).
DIATROPA, Dum. Fl. Belg. 76 (1827).
ISOPHYLLUM, Hoffm Gen. Umb. 115 (1814).
ONONTEA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 391.
ODONTITES, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 256.
PERFOLIATA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 390.
TENORIA, Spreng. Pl. Umb. Prod. 82 (1813).
TRACHYPLEURUM, Reichb. Consp 148 (1828).
acerosum, *E. Mey. in Drège, Zwei Pfl. Docum.* 170, nomen.—Afr. austr.
adiphyllum, *Webb & Berth. Phyt. Canar.* ii. 154. t. 77. —Ins. Canar.
acutifolium, Boiss. Elench. 47 = pancalicum.
aegyptiacum, Willd. ex Steud. Nom. ed. II. i. 239 = protractum.
æneum, *Boiss. & Huet, Diagn.* Ser. II. ii. 85.—Armenia.
affine, *Sadl. Fl. Pesth.* 204.—Europ. austr. ; Taurica.
aleppicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 148.—Syria.
alpigenum, Jord. & Fourr. Brev. Fl. Nov. fasc. i. 85 = falcatum.
alpinum, Nym. Consp. 311 = falcatum.
altaicum, Pall. ex Roem. & Schult. Syst. vi. 867, 868 = triradiatum.
amplexicaule, [Clairv.] Man. Herb. 87 = pyrenaicum.
angulosum, Cham. & Schlecht. in Linnæa, i. (1826) 883 = ranunculoides.
angulosum, *Linn. Sp. Pl.* 236.—Reg. arct.
angustifolium, *Ledeb. Fl. Rossc.* ii. 265.—Sibir. altaic.
angustifolium, *Pall. Reise Statth. Russ. Reich.* i. 345. —Reg. Caucas.
apiculatum, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 147 = semidiaphanum.
apiculatum, *Friwald. in Flora*, xviii. (1835) 335.—Oriens.
arborescens, Jacq. Coll. ii. 343 = gibraltaricum.
arborescens, Thunb. Prod. Pl. Cap. 50 = Heteromorpha arborescens.
arixtatum, *Bartl. in Reichb. Ic. Fl. Germ.* ii. 70. t. 178. —Europ. ; Reg. Caucas.
armenum, Boiss. & Kotschy, ex Boiss. Fl. Orient. ii. 845 = kurdicum.
asperuloides, *Heldr. in Boiss. Diagn.* Ser. II. vi. 76.—Graecia.
Auckeri, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 146 = brevicaule.
aureum, *Fisch. in Hoffm. Gen. Umb.* 115.—Sibir.
australe, Jord. Pugill. 72 = Gerardi.
badium, Rochel & Sag. ex Saut. in Flora, lx. (1836) 177 = junceum.
Balansæ, *Boiss. & Reut. Diagn.* Ser. II. ii. 83.—Afr. bor.
baldense, *Host, Syn.* 141.—Europ. ; Oriens.
baldense, Turr. in Giorn. Ital. i. (1766) 130 = aristatum.
baldense, Waldst. & Kit. Ic. Pl. Hung. Rar. iii. 285. t. 257 = junceum.
Barcellii, *Cosn. ex Willk. in Linnæa*, xl. (1876) 83.—Afr. bor.
Baumgartenianum, Schur, Enum. Pl. Transs. 252 = petræum.
bicaule, Balb. Stirp. Nov. 4. t. 2 = stellatum.
bicaule, Hoffm. in Mém. Soc. Nat. Mosc. ii. (1809) 196. t. 9 = baldense.
Bourgæi, *Boiss. & Reut. Diagn.* Ser. II. ii. 84.—Hispan.
brachiatum, *C. Koch, ex Boiss. Fl. Orient.* ii. 844.—As. Min.
Brasinum, *Timb. in Mém. Acad. Toulouse*, Sér. VIII. iv. (1892) 125.—Gallia.
brevicaule, *Schlecht. in Linnæa*, xvii. (1843) 124.—Syria.
Burseranum, Willd. Enum. Hort. Berol. Suppl. 14 = angulosum.
canalense, Wulf. ex Spreng. in Roem. & Schult. Syst. vi. 371 = ranunculoides.
canariense, Spreng. Syst. i. 882 = Astydamia canariensis.
Candollii, *Wall. Cat.* n. 552.—Reg. Himal.
canescens, *Schousb. Vetr. Marokkko*, 197.—Marocco.
capillare, *Boiss & Heldr. Diagn.* Ser. II. ii. 82.—Graecia.
capistratum, Thunb. Prod. Pl. Cap. 50 = Hermas capitata.
cappadocicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 146.—As. Min. ; Syria.
carieciifolium, Willd. Sp. Pl. i. 1373 = ranunculoides
carsicinum, Reichb Ic. Crit. ix. f. 1109, 1110 = ranunculoides
cernuum, Tenore, App. i. Cat. Hort. Neap. ; DC. Prod. iv. 131 = baldense
chinense, *Franch. in Nouv Arch. Mus. Par.* Sér. II. vi. (1883) 18.—China.
chimense, DC. Prod. iv. 128 = falcatum.
chlorocarpum, *Timb. & Seanb. in Mém. Acad. Toulouse*, Sér. VI. v. (1867) 471, nomen.—Gallia.
Chouletii, *Pomel, Nouv. Mat. Fl. Atl.* 140.—Algeria.
ciliatum, Thunb. Prod. Pl. Cap. 50 = Hermas ciliata.
citrinum, *Hochst. in Lorent, Wanderungen*, 387.—Armenia.
collinum, D. Dietr. Syn. Pl. ii. 951 = Heteromorpha arborescens.
coloratum, Schur, Enum. Pl. Transs. 254 = aureum.
Columnæ, Guss. Fl. Sic. Prod. 70 = tenuissimum.
commutatum, *Boiss. & Bal. Diagn.* Ser. II. vi. 75.—Macedon. ; As. Min.
compositum, *Spreng. in Ges. Naturf. Fr. Berl. Mag.* vi. (1812) 258.—Hab. ?
corbariense, *Timb. in Mém. Acad. Toulouse*, Sér. VIII. v. ii. (1883) 139.—Mont. Pyren.
coriaceum, L'Hérit. Stirp. Nov. 139. t. 67 = gibraltaricum.
corsicum, *Cos. & Kral. Notes Crit.* 56.—Corsica.
crocetum, *Fenzl, Pugill. Pl. Nov. Syr.* 16.—As. Min. ; Syria.
cuspidatum, Bunge, in Mém. Acad. Pétersb. vii. (1854) 599 ; (Lehm. Rel. 128) = linearifolium.
dahuricum, *Fisch. & Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 720, 721.—Dahuria.
dendriticum, *Rupr. Sert. Tianschk.* 47.—As. centr.
denticulatum, Bellard, ex Spreng. in Roem. & Schult. Syst. vi. 396 = petræum.
dianthifolium, *Guss. Suppl. Fl. Sic. Prod.* i. 71.—Sicilia.
dichotomum, *Boiss. Diagn.* Ser. I. x. 28.—Syria.
dichotomum, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 343, in textu = Gerardi.
difforme, *Linn. Sp. Pl.* 238.—Afr. austr.
dilatatum, Schur, Enum. Pl. Transs. 253 = falcatum.
disticbophyllum, *Wight & Arn. Prod.* i. 370.—Ind. or.
divaricatum, Lam. Fl. Fr. iii. 410 = B. Odontites.
divaricatum, Steph. ex Ledeb. Fl. Ross. ii. 262 = gracile.
diversifolium, Rochel, Banat. 68. t. 28 = baldense.
dumosum, *Cosn. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 249 ; xxii. (1875) 59.—Marocco.
elatum, *Guss. Fl. Sic. Prod.* i. 316.—Sicil.
erubescens, *Boiss. Fl. Orient.* ii. 841.—Lycia.
exaltatum, Bieb. Beschr. Casp. 156 = baldense.
exaltatum, Schur, Enum. Pl. Transs. 253 = subfalcatum.
falcatum, Berg. Desc. Pl. Cap. 76, partim = Mundtii.
falcatum, Dals. & Gibs. Bomb. Fl. 108 = mucronatum.
falcatum, *Linn. Sp. Pl.* 237.—Europ. ; Oriens ; As. bor. ; Reg. Himal.
falcatum, Turra. in Bull. Soc. Nat. Mosc. xvii. (1844) 721—dahuricum.
filicaule, *Brot. Fl. Lusit.* i. 452.—Lusitan.
flavicans, *Boiss. & Heldr. Diagn.* Ser. II. vi. 74.—Aetolia.
flavum, *Forsk. Fl. Aegypt. Arab.* 205 (sp. dub.) = glumaceum ?
flexuosum, *Ledeb. Fl. Ross.* ii. 267.—Dahuria.
flexuosum, Moench, Meth. 88 = falcatum.
flexuosum, Wall. Cat. n. 554 = tenue.
foliosum, *Salzm. ex DC. Prod.* iv. 133.—Marocco.
Fontanesii, Guss. Ind. Sem. Hort. Boce. (1825) = B. Odontites.
frutescens, Hill, Veg. Syst. vi. 21 = fruticosum.
fruticans, Stend. Nom. ed. I. 128 = difforme.
fruticescens, Bourg. ex Nym. Consp. 310 = foliosum fruticescens, *Linn. Cent. Pl.* i. 9.—Reg. Mediterr. occ
fruticosum, *Linn. Sp. Pl.* 238.—Reg. Mediterr.
Gerardi, Colm. ex Nym. Consp. 312 = junceum.
Gerardi, Gaertn. Mey. & Scherb. Fl. Wett. i. 408 = tenuissimum.
Gerardi, *Jacq. Fl. Austr.* iii. 31. t. 256.—Europ. austr. ; Oriens.
Gerardi, Pollini, Viag. Lag. Gard. 15, 31 = B. Odontites.
Gerardi, Sibth. & Sm. Fl. Græc. t. 262 = trichopodum.
Gerhardi, Georgi, Beschr. Russ. Reich. iii. iv. 826 = Gerardi.
gibraltaricum, *Lam. Encyc.* i. 690.—Lusitan. ; Marocco.
gigantæum, Thunb. Prod. Pl. Cap. 50 = Hermas gigantea.
glaucum, *Robill & Cast. ex DC. Fl. Fr. Suppl.* 515.—Reg. Mediterr.
glumaceum, *Sibth. & Sm. Fl. Græc. Prod.* i. 177.—Graecia ; As. Min
gracile, DC. Prod. iv. 128 = Marschallianum.
gracile, Urv. in Mém. Soc. Linn. Par. i. (1822) 98 = glumaceum.
gracillimum, Klotzsch, in Bot. Ergeb. Waldem. Reise, 148 = falcatum.
gramineum, Vill. Prosp. 28 = baldense.
graminifolium, Favre, in Bull. Soc. Murith. v. (1875) 85, 250 = stellatum.
graminifolium, Georgi, Beschr. Russ. Reich. iii. iv. 826 = exaltatum.
graminifolium, Lapeyr. Hist. Abr. Pl. Pyr. 189 = ranunculoides.
gramsinifolium, Salisb. Prod. 160 = B. Odontites.
graminifolium, Vahl, Symb. Bot. iii. 48 = petræum.
Hausknechtii, *Boiss. Fl. Orient.* ii. 848.—Kurdistan.
Heldreichii, *Boss. & Bal. Diagn.* Ser. II. vi. 75.—As. Min.
heterophyllum, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 675 = jucundum.
heterophyllum, Link, Enum. Hort. Berol. i. 262 = protractum.
heterophyllum, Rochel, Banat. 28 = baldense.
himalayense, Klotzsch, in Bot. Ergeb. Waldem. Reise, 146 = longicaule.
hispanicum, Pers. Syn. i. 305 = fruticescens.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 147 = falcatum.
humile, Vest, ex Reichb. Ic. Fl. Germ. ii. 71 = aristatum.
imaicolum, Kern. in Ber. Naturw. Ver. Innsbruck, i. (1870) 115 (Nov. Pl. Sp. Dec. ii. 19) = falcatum.
incurvum, Bell. App. Fl. Pedem. 77. t. 11 = petræum.
intermedium, Stend. Nom. ed. I. 128 = protractum.
irregulare, *Boiss & Kotschy, ex Boiss. Fl. Orient.* ii. 850.—Syria.
Jacquinianum, Jord. Pugill. 71 = Gerardi.
jucundum, *Kurz, in Journ. Bot.* v. (1867) 240.—Himal. bor. occ. ; Reg. Amur.
juncetum, Bieb. Fl. Taur. Cauc. i. 304 = affine.
juncetum, *Linn. Sp. Pl.* ed. II. 343.—Europ. austr.
juncetum, Pall. Ind. Taur. ex Bieb. Beschr. Casp. 166 = exaltatum.
juncetum, Pollich, Fl. Palat. i. 264 = tenuissimum.
Kargli, *Vis. Fl. Dalm.* iii. 35.—Dalmat.
Koechelli, *Fenzl, Pugill. Pl. Nov. Syr.* 17.—As. Min.
kokanicum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 29.—As. centr.
Kotschyanum, Boiss. Diagn. Ser. I. x. 29 = linearifolium.
kurdicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 146.—As. Min. ; Syria ; Persia.
lancifolium, *Wall. Cat.* n. 551.—Reg. Himal.
lancifolium, Hornem. Cat. Hort. Hafn. i. 267 = protractum.

BUPLEURUM :—
- laricense, *Gaut. & Timb. in Mém. Acad. Toulouse*, Sér. VIII. iv. (1883) 180.—Gallia.
- lateriflorum, *Cass. in Bull. Soc. Bot. Fr.* xxii. (1875) 59.—Marocco.
- leucocladum, *Boiss. in Ann. Sc. Nat. Sér.* III. i. (1844) 144.—Assyria.
- libanoticum, *Boiss. & Blanche, Diagn.* Ser. II. ii. 82. —Syria.
- linearifolium, *DC. Prod.* iv. 131.— Persia ; Arab. ; Turkest.
- *longepetiolatum*, Weiss, in Verh. Zool.-Bot. Ges. Wien, xvi. (1866) 581 = Kargli.
- longeradiatum, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 719.—Sibir. balcal.
- longicaule, *Wall. Cat. n.* 557.—Reg. Himal.
- *longifolium*, Falk, Beitr. ii. 144 = aureum.
- longifolium, *Linn. Sp. Pl.* 287.—Europ.
- lophocarpum, *Boiss. & Bal. Diagn.* Ser. II. v. 97.— Cilicia.
- Maddeni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 678.—Reg. Himal.
- *marsoticum*, Delile, ex DC. Prod. iv. 129 = nodiflorum.
- *marginatum*, Wall. Cat. n. 566 = falcatum.
- Marschallianum, *C. A. Mey. Verz. Pfl. Cauc.* 123.— Graecia ; As. Min. ; Persia.
- *maximum*, Clarke, Trav. ii. 844 ; Spreng. Neue Entdeck. iii. 163 = nodiflorum.
- montanum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iii. (1856) 706.—Algeria.
- *mucronatum*, Brouss. ex Spreng. in Roem. & Schult. Syst. vi. 877 = fruticescens.
- mucronatum, *Wight & Arn. Prod.* 370.—Ind. or.
- multinerve, *DC. in Mém. Phys. Soc. Genèv.* iv. (1828) 500.—Sibir. altaic.
- Mundtii, *Cham. & Schlecht. in Linnaea*, i. (1826) 384. —Afr. austr.
- *nanum*, Poir. Encyc. Suppl. i. 750 = nodiflorum.
- *nebulosum*, Hausskn. ex Boiss. Fl. Orient. ii. 847 = cappadocicum.
- *neglectum*, Cesati, in Linnaea, xi. (1837) 314 = baldense.
- *nervosum*, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 97 = polyphyllum.
- *nervosum*, Moon, Cat. 22 = mucronatum.
- *nervosum*, Schoush. ex Ball, in Journ. Linn. Soc. xvi. (1878) 466 = paniculatum.
- *nervosum*, Trev. Synsb. Phytol. i. 22 = multinerve.
- *nigrocarpa*, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 678 = falcatum.
- nodiflorum, *Sibth. & Sm. Fl. Graec.* v. 260.—Graecia ; Syria ; Aegypt.
- *Nordmannianum*, Bal. ex Boiss. Fl. Orient. ii. 851 = polyphyllum.
- *Nordmannianum*, Ledeb. Fl. Ross. ii. 265 = baldense.
- nudum, [*Soland. in*] *Ait. Hort. Kew.* ed. 1. i. 381.— Afr. austr.
- *obliquatum*, Schoush. ex Ball, in Journ. Linn. Soc. xvi. (1878) 466 = foliosum.
- *obliquum*, Vahl, Symb. Bot. i. 24 = gibraltaricum.
- oblongifolium, *Ball*, in *Journ. Linn. Soc.* xvi. (1878) 466.—Marocco.
- *obtusatum*, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. ii. 48 = angulosum.
- *obtusifolium*, Tenore, Prod. Fl. Nap. p. xviii = baldense.
- octoradiatum, *Bunge*, in *Mém. Sav. Etr. Pétersb.* ii. (1830) 106.—China bor.
- Odontites, *Linn. Sp. Pl.* 287.—Europ. ; Reg. Mediterr.
- *Odontites*, Sm. Engl. Bot. t. 2468 (1815) ; Lapeyr. Hist. Abr. Pl. Pyr. 141 (1813) = aristatum.
- *odontitum*, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 67, 121 = B. Odontites.
- oligactis, *Boiss. Diagn.* Ser. II. ii. 84.—Hispan. ; Marocco.
- *olympicum*, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 149 = baldense.
- *opacum*, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 73 = aristatum.
- oppositifolium, *Lapeyr. Hist. Abr. Pl. Pyr.* 141.—Mont. Pyren.
- pachnospermum, *Panč. Fl. Princ. Serbiae* (1874) 239. —Banat.
- *panacifolium*, Hornem. ex Steud. Nom. ed. ii. i. 240 = heterophyllum.
- *paniculatum*, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 74 = Bourgaei.

BUPLEURUM :—
- paniculatum, *Brot. Fl. Lusit.* i. 455.—Lusit. ; Marocco.
- *paniculatum*, Willk. ex Willk. & Lange, Prod. Fl. Hisp. iii. 74 = rigidum.
- papillosum, *DC. Prod.* iv. 127.—Mesopotam.
- paucidradiatum, *Fenzl, in Tchihat. Asie Min. Bot.* i. 418.—Cilicia.
- *perfoliatum*, Lam. Fl. Fr. iii. 405 = rotundifolium.
- *persicum*, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 150 = baldense.
- *petiolare*, Lapeyr. Hist. Abr. Pl. Pyr. 141 = falcatum.
- *petraeum*, Genera. ex DC. Prod. iv. 131 = angulosum.
- *petraeum*, Georgi, It. i. 204 = exaltatum.
- petraeum, *Linn. Sp. Pl.* 236.—Europ.
- *petraeum*, Wulf. in Jacq. Coll. ii. 209 = graminifolium.
- *petrogenes*, Jord. ex Nym. Consp. 311 = falcatum.
- *petraeifolioides*, Spreng. Pl. Umb. Prod. 39 = Carum peregrinum.
- plantagineum, *Desf. Fl. Atlant.* i. 283. t. 57.— Marocco.
- plantaginifolium, *Wight, Ic.* t. t. 281.—Ind. or.
- *Pallichii*, C. C. Gmel. Fl. Bad. i. 613 = tenuissimum.
- polyphyllum, *Ledeb. in Mém. Acad. Petersb.* v. (1815) 538.—Reg. Caucas.
- *procumbens*, Desf. Fl. Atlant. i. 280. t. 86 = tenuissimum.
- *proliferum*, Delile, Fl. Egypte, 205. t. 22 = nodiflorum.
- prostratum, *Ledeb. ex Loud. Hort. Brit. Suppl. ii.* 615. —Sibir. altaic.
- protractum, *Hoffmgg. & Link, Fl. Port.* ii. 387.— Reg. Mediterr. or.
- provinciale, *Vent. ex Trinb. in Mém. Acad. Toulouse*, Sér. VIII. iv. (1882) 181.—Gallia.
- *prunosum*, Cesati, ex Boiss. Fl. Orient. ii. 849 = baldense.
- pulchellum, *Boiss. & Heldr. Diagn.* Ser. I. x. 27.— Pamphylia.
- *pyrenaicum*, Willd. Sp. Pl. i. 1371 = angulosum.
- *pyrenaeum*, Gouan, Illustr. 8. t. 4 = angulosum.
- *quinqueden*[*ta*]*tum*, Thunb. Prod. Pl. Cap. 50 = Hermas quinquedentata.
- *quinquenervatum*, Steud. Nom. ed. ii. i. 240, sphalm. = junceo.
- *ramosissimum*, Wight & Arn. Prod. 370 = mucronatum.
- ramosum, *Gaut. ex Timb. in Mém. Acad. Toulouse*, Sér. VIII. iv. (1882) 186.—Gallia.
- *ranunculiformis*, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 67 = sec.
- ranunculoides, *Linn. Sp. Pl.* 287.—Reg. bor. temp.
- *ranunculoides*, Sibth. & Sm. Fl. Graec. Prod. i. 178 = baldense.
- *repens*, Lapeyr. Hist. Abr. Pl. Pyr. 140 = ranunculoides.
- *rigidum*, Desv. ex Reichb. Ic. Fl. Germ. xxi. 21 = baldense.
- *rigidum*, Georgi, Beschr. Russ. Reichb. iii. tw. 898 = Gerardi.
- rigidum, *Linn. Sp. Pl.* 238.—Gall. austr.
- *Rissoni*, Rich. ex DC. Prod. iv. 126 = affine.
- rotundifolium, *Linn. Sp. Pl.* 236.—Europ. ; Reg. Caucas. ; Persia.
- *rotundifolium*, Schoush. Vext. Marokko, 126 = protractum.
- *rupestre*, Edgew. in Trans. Linn. Soc. xx. (1851) 52 = longicaule.
- rupestre, *Kaſin. in Med. Repos. N. York*, v. (1808) 352, nomen.—Hab.?
- *sachalinense*, F. Schmidt, Reisen Amurl. 185 = jucundum.
- salicifolium, *Salomd. ex Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) repr. 21.—Ins. Madera.
- Savignonii, *De Not. in Mem. Acc. Tor.* ix. (1848) 208. —Italia.
- *saxatile*, Pall. Reise, ii. 676 = baldense.
- *scabrum*, Griseb. Spicil. Fl. Rumel. i. 348 = brevicaule.
- *Schefferi*, Hampe, ex Nym. Consp. 312 = filicaule.
- *Schimperi*, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 151 = linearifolium.
- *scorzonerafolium*, Willd. Enum. Hort. Berol. Suppl. 30 = falcatum.
- *semicompositum*, Hochst. & Steud. ex Steud. Nom. ed. ii. i. 240 = aristatum.
- semicompositum, *Linn. Amoen. Acad.* iii. 405.—Europ. austr.
- *semicompositum*, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. i. 208 = junceum.
- *semicompositum*, Sibth. & Sm. Fl. Graec. Prod. i. 178 = glaucum.
- semidiaphanum, *Boiss. Diagn.* Ser. II. vi. 78.— Graecia et Ins.

BUPLEURUM :—
- *semiperfoliatum*, Griseb. Spicil. Fl. Rumel. i. 350 = crocum.
- setaceum, *Fenzl, in Tchihat. Asie Min. Bot.* i. 418.— Bithynia ; Himal. bor. occ.
- *sibiricum*, Vest, ex Roem. & Schult. Syst. vi. 868 = angulosum.
- *Sibthorpianum*, DC. ex Boiss. Fl. Orient. ii. 849 = linearifolium.
- *Sibthorpianum*, Sibth. & Sm. Fl. Graec. Prod. i. 179 = baldense.
- sphnosum, *Gouan, Illustr.* 8.—Hispan. ; Afr. bor. occ
- *Sprunerianum*, Hampe, in Flora, xxv. (1842) 1. 68 = B. Odontites.
- *stellatum*, Lapeyr. Hist. Abr. Pl. Pyr. 139 = angulosum.
- stellatum, *Linn. Sp. Pl.* 236.—Europ. austr.
- *striatum*, Steph. ex Roem. & Schult. Syst. vi. 867 = multinerve.
- subfalcatum, *Schur, Enum. Pl. Transs.* 253. — Transsylv.
- *subovatum*, Link, ex Spreng. Sp. Umb. 19 = protractum
- *subpinnatum*, Ledeb. in Eichw. Pl. Nov. Casp. Cauc 13. t. 11 = Carum segetum.
- *subrubrum*, Dulac, Fl. Hautes-Pyr. 346 = ranunculoides.
- subuniflorum, *Boiss. & Heldr. Diagn.* Ser. I. x. 28. —Pamphylia.
- *sulcatum*, Steud. Nom. ed. II. i. 240 = Mundtii.
- sulphureum, *Boiss. & Bal. Diagn.* Ser. II. vi. 74.— As. Min.
- *tauricum*, Grem. ex Nym. Consp. 311 = ranunculoides
- tenue, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 182.— Reg. Himal.
- *tenue*, Salisb. Prod. 160 = tenuissimum.
- *tenuifolium*, Pourr. ex Nym. Consp. 311 = baldense.
- tenuissimum, *Linn. Sp. Pl.* 238.—Europ. ; Afr. bor.
- *termonale*, Salisb. Prod. 160 = fruticosum.
- Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 675.—Reg. Himal.
- tortuosum, *Schousb. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 466.—Marocco.
- *transsilvanicum*, Schur, Enum. Pl. Transs. 253 = baldense.
- trichopodum, *Boiss. & Sprun. in Ann. Sc. Nat.* Sér. III. i. (1844) 145.—Graecia ; As. Min.
- *trifolium*, Tenore, Prod. Fl. Nap. 18 = junceum.
- *trifoliatum*, Wendl. & Bartl. Beitr. ii. 18 = Heteromorpha trifoliata.
- triradiatum, *Adams, ex Hoffm. Umb.* 115.—Sibir.
- *vaginatum*, Vill. Prosp. 28 = longifolium.
- Vaucheriſ, *Mudl. Arg. in Heurck, Obs. Bot.* 207.— China.
- *veronense*, Turra, Fl. Ital. Prod. 64 = B. Odontites.
- *verticale*, Orteg. Fl. Esp. ex DC. Prod. iv. 133 = gibraltaricum.
- *villosum*, Linn. Sp. Pl. 238 = Hermas villosa.
- *virgatum*, Brot. ex Steud. Nom. ed. II. i. 240 = fruticosa.
- *virgatum*, Cav. Desc. i. 121 = Gerardi.
- *virgatum*, Wall. Cat. n. 555 = falcatum.
- *virgatum*, Wight & Arn. Prod. 370 = mucronatum.
- *Wittmanni*, Siev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 342 = rotundifolium.

BUPRESTIS, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 258 = Bupleurum, Tourn. (Umbell.).
- *arborescens*, Spreng. l. c. 259 = Heteromorpha arborescens.
- *coriacea*, Spreng. l. c. = B. gibraltaricum.
- *difformis*, Spreng. l. c. = B. difforme.
- *exaltata*, Spreng. l. c. = B. baldense.
- *fruticosa*, Spreng. l. c. = B. fruticosum.
- *frutiscea*, Spreng. l. c. = B. fruticosum.
- *Gerardi*, Spreng. l. c. = B. Gerardi.
- *gigantea*, Spreng. l. c. = Hermas gigantea.
- *juncea*, Spreng. l. c. = B. junceum.
- *latifolia*, Spreng. l. c. = Astydamia canariensis.

BURASAIA, Baill. Adansonia, xi. (1873) 88. *EUPHORBIACEAE*, Benth. & Hook. f. ill. 390. caruncula, *Baill. l. c.* 84.—N. Caled. clusiacea, *Baill. l. c.*—N. Caled.

BURASAIA, Thou. Gen. Madag. 18 (1806) ; et in Roem. Collect. 209 (1809) ; DC. Syst. Veg. i. 534 (1818). *MENISPERMACEAE*, Benth. & Hook. f. i. 36, 960.

BURASAIA :—
congesta, *Decne. in Ann. Sc. Nat.* Sér. II. xii. (1839) 108.—Madag.
gracilis, *Decne. l. c.*—Madag.
madagascariensis, *DC. Syst.* i. 514.—Madag.
Vitis, *Mello, in Journ. Linn. Soc.* xiv. (1875) 4.—Bras.

BURBIDGEA, Hook. f. Bot. Mag. t. 6408 (1879).
SCITAMINEAE, Benth. & Hook. f. iii. 647.
nitida, *Hook. f. l. c.*—Borneo.

HURCARDA, J. F. Gmel. Syst. 514 (1791) = seq.

BURCARDIA, Schreb. Gen. i. 330 (1789) = **Piriqueta**, Aubl. (Turnerac.).
aromatica, Rafin. Sylva Tellur. 106.—Hab. ?
grandiflora, Rafin. l. c.—Hab. ?
villosa, J. F. Gmel. Syst. ii. 514 = P. villosa.

BURCHARDIA, [Heist.] Duham. Arb. i. 111, t. 44 (1755) = **Calliscarpa**, Linn. (Verbenac.).
americana, Duham. l. c. = C. americana.
Calliscarpa, Crantz, Inst. ii. 360 = C. americana.

BURCHARDIA, R. Br. Prod. 272 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 822.
congesta, Lindl. Swan River App. 58 = umbellata.
multiflora, Lindl. l. c. = umbellata.
umbellata, *R. Br. Prod.* 273.—Austral.

BURCHELLIA, R. Br. in Bot. Reg. t. 466 (1820). *RUBIACEAE*, Benth. & Hook. f. ii. 86.
RUBALINA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 86.
Bubalina, Sims, Bot. Mag. t. 2339 = capensis.
capensis, *R. Br. in Bot. Reg.* t. 466.—Afr. austr.
Kreussii, Hochst. in Flora, xxv. (1842) 237 = capensis.
major, Hort. ex Heynh. Nom. ii. 88 = capensis.
parviflora, Lindl. Bot. Reg. t. 891 = capensis.
speciosa, Hort. ex Heynh. Nom. ii. 88 = capensis.

BURDACHIA, Mart. ex A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 329. *MALPIGHIACEAE*, Benth. & Hook. f. i. 352.
macrocarpa, Benth. ex Griseb. in Mart. Fl. Bras. xii. 1. 23 = Glandonia macrocarpa.
prismatocarpa, *Mart. ex A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 330.—Bras.
pyramidata, Mart. ex Griseb. in Mart. Fl. Bras. xii. 1. 23 = prismatocarpa.
sphaerocarpa, *A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 330.—Bras.

BUREAVA, Baill. Adansonia, i. 71 (1860–61) = Combretum, Linn.
crotonoides, Baill. l. c. 72 = C. altum.

BURGESIA, F. Muell. Fragm. i. 232 (1859) = **Brachysaccus**, R. Br.
homalocalada, F. Muell. l. c. = Brachysema aphyllum.

BURGHARTIA, Scop. Introd. 229 (1777) = **Piriqueta**, Aubl. (Turnerac.).
villosa, J. F. Gmel. Syst. 514 = P. villosa.

BURGLARIA, Wendl. ex Steud. Nom. ed. I. 239 (1821) = **Ilex**, Tourn.
lucida, Wendl. l. c. = I. salicifolia.

BURGSDORFIA, Moench, Meth. 392 (1794) = **Sideritis**, Tourn. (Labiat.).
montana, Reichb. Fl. Germ. Excurs. 327 = S. montana.
rigida, Moench, Meth. 392 = S. romana.
romana, Hoffmgg. & Link, Fl. Port. i. 97 = S. romana.

BURKEA, Benth. in Hook. Ic. Pl. t. 598 (1848). *LEGUMINOSAE*, Benth. & Hook. f. i. 587.
africana, *Hook. l. c.* tt. 598, 594.—Afr. austr.
Capensegun, *Baill. Hist. Pl.* ii. 107.—Afr. trop.

BURKHARDIA, Benth. & Hook. f. Gen. i. 1007 (1867) = **BURGHARTIA**, Scop. = **Piriqueta**, Aubl. (Turnerac.).

BURLINGTONIA, Lindl. Bot. Reg. t. 1927 (1837) = **Rodriguesia**, Ruiz & Pav. (Orchid.).
Batemani, Linden, Illustr. Hortic. xxviii. (1881) 64 = R. Batemanni.
candida, Lindl. Bot. Reg. xxiii. (1837) sub t. 1927 = R. candida.
carnea, Lindl. ex Linden, Illustr. Hortic. xxviii. (1881) 80 = R. carnea.
decora, Lem. Jard. Fleur. ii. (1852) Misc. 96 = R. decora.
fragrans, Lindl. Bot. Reg. xxiii. (1837) sub t. 1927 = R. fragrans.
granadensis, Lindl. Orchid. Linden. 24 = R. granadensis.
maculata, Lindl. Bot. Reg. xxv. (1839) 44 = R. maculata.
obtusifolia, Lindl. Sert. Orch. (1840) sub t. 36 = R. obtusifolia.
pubescens, Lindl. & Paxt. Flow. Gard. i. (1851) 158 = R. pubescens.
refracta, Lindl. ex Linden, Illustr. Hortic. xxviii. (1881) 80 = R. refracta.
rigida, Lindl. ex Lem. in Fl. des Serres, i. (1845) 11. t. 9 = R. rigida.
rubescens, Lindl. Bot. Reg. xxiii. (1837) sub t. 1927.—Peruv.
sivinitiulana, Lem. Jard. Fleur. iii. (1852–53) Misc. 66. —Bras.
venusta, Lindl. ex Lem. Illustr. Hortic. v. (1858) t. 188 = R. venusta.

BURMANNIA, Linn. Syst. ed. I (1735). *BURMANNIACEAE*, Benth. & Hook. f. iii. 457.
CYANANTHUS, Miers, in Wall. Cat. nn. 9007–8(1849).
CYANOTIS, Miers, l. c. n. 9008 (1849).
CRYPTONEMA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 590.
GONIANTHUS, Blume, Cat. Gew. Buitenz. (1823) 19.
MARIANTHA, Thou. Gen. Nov. Madag. 4 (1806).
NEPHROCOELIUM, Turcz. in Bull. Soc. Nat. Mosc. (1858) t. 287.
TETRAPTERA, Miers, in Lindl. Veg. Kingd. 172 (1847).
TRIPTERANTHUS, Wall. ex Miers, l. c. (1847).
TRIPTERELLA, Michx. Fl. Bor. Am. i. 19. t. 3 (1803).
VOGELIA, J. F. Gmel. Syst. 107 (1791).
alba, *Mart. Nov. Gen. et Sp.* i. 12.—Bras.
asurea, Griff. Notul. iii. 236 = coelestis.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 617.—Ins. Banca.
bicolor, *Mart. Nov. Gen. et Sp.* i. 10. t. 5.—Bras.
biflora, *Linn. Sp. Pl.* 287.—Am. bor.
brachyphylla, *Willd. ex Schult. f. Syst.* vii. ii. p. lxxv. —Venezuela.
brachystachya, *Miq. in Linnaea*, xix. (1847) 141.—Guiana.
capensis, *Mart. Nov. Gen. et Sp.* i. 12.—Afr. austr.
capitata, *Mart. l. c.*—Am. bor.
Championi, *Thw. Enum. Pl. Zeyl.* 325.—Zeylan.
coelestis, *D. Don, Prod. Fl. Nep.* 44.—As. trop.; Austral.
dasyantha, *Mart. Nov. Gen. et Sp.* i. 11.—Bras.
distachya, R. Br. Prod. 305 = disticha.
distichia, *Linn. Sp. Pl.* 287.—As. trop.; Austral.
flava, *Mart. Nov. Gen. et Sp.* i. 11. t. 5. f. 2.—Bras.
flavula, *Wright, in Sauv. Fl. Cub.* 168.—Cuba.
Goetzeiklana, *Becc. Malesia*, i. (1878) 244.—Malaya.
javanica, *Blume, Enum. Pl. Yav.* fasc. i. 28.—Java.
juncea, *Soland. ex R. Br. Prod.* 305.—Austral.
Kalbreyeri, *Oliver, in Hook. Ic. Pl.* t. 1857.—N. Granat.
longifolia, *Becc. Malesia*, i. (1878) 244.—Malaya.
lutescens, *Becc. l. c.* 246.—Malaya.
madagascariensis, *Baker, in Journ. Linn. Soc.* xx. (1884) 268.—Madag.
madagascariensis, *Mart. & Zucc. Nov. Gen. et Sp.* i. 12.—Madag.
pantenista, *Willd. ex Schult. f. Syst.* vii. ii. p. lxxiv. —Madag.
pusilla, *Thw. Enum. Pl. Zeyl.* 325.—Zeylan.
quadrifiora, *Willd. ex Schult. f. Syst.* vii. 79. p. lxxiv.—Guiana.
selebica, *Becc. Malesia*, i. (1878) 243.—Ins. Celebes.
Sellowiana, *Seub. in Mart. Fl. Bras.* iii. 57. t. 7. f. 2. —Bras.
sphagnoides, *Becc. Malesia*, i. (1878) 246.—Malaya.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 617 = distichia.
tenella, *Benth. in Hook. Kew Journ.* vii. (1855) 12. —Bras.
tridentata, *Becc. Malesia*, i. (1878) 246.—Malaya.

BURMANNIA :—
triflora, Roxb. Hort. Beng. [87]; Fl. Ind. ii. 117 = coelestis.
tuberosa, *Becc. Malesia*, i. (1878) 245.—Malaya.
uniflora, Rottl. ex Spreng. Syst. iv. Cur. Post. 23 = coelestis.

BURMEISTERA, Karst. & Triana, in Linnaea, xxviii. (1856) 444. *CAMPANULACEAE*, Benth. & Hook. f. ii. 547.
acuminata, *Karst. l. c.* 445.—N. Granat.
glabrata, *Benth. & Hook. f. Gen.* iii. 547.—N. Granat
lbaguensis, *Karst. in Linnaea*, xxviii. (1856) 444.—N. Granat.
lacerata, *Karst. l. c.* 445.—N. Granat.
Matthiasi, *Benth. & Hook. f. Gen.* ii. 547.—Peruv.
marginata, *Karst. in Linnaea*, xxviii. (1856) 445.—N. Granat.
pomifera, *Karst. l. c.*—N. Granat.
succulenta, *Karst. l. c.*—N. Granat.
tomentonula, *Karst. l. c.*—N. Granat.
virescens, *Benth. & Hook. f. Gen.* ii. 547.—Guatemala.

BURNATIA, Micheli, in DC. Monog. Phan. iii. 81 (1881). *ALISMACEAE*, Benth. & Hook. f. iii. 1007.
enneandra, *Micheli, l. c.*—Nubia.

BURNETTIA, Lindl. Gen. et Sp. Orch. 517 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 612.
cuneata, *Lindl. l. c.* 518.—Tasmania.

BURNEYA, Cham. & Schlecht. in Linnaea, iv. (1829) 189 = **Pimonia**, Rumph. et **Bobea**, Gaudich. (Rubiac.).
Forsteri, Cham. & Schlecht. l. c. = T. Forsteri.
Gaudichaudii, Cham. & Schlecht. l. c. 190 = B. Gaudichaudi.

BURRIELIA, DC. Prod. v. 668 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 398.
anthemoides, *A. Gray, in Torr. Bot. Mex. Bound.* 97. —Am. bor.
chrysostoma, Torr. & Gray, Fl. N. Am. ii. 379 = Baeria chrysostoma.
Fremontii, Benth. & Hook. Fl. Hartw. 317 = Baeria Fremontii.
gracilis, DC. Prod. v. 664 = Baeria gracilis.
hirsuta, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 381 = Baeria gracilis.
lanosa, A. Gray, in Pacif. Rail. Rep. iv. 107 = Eriophyllum lanosum.
leptalea, A. Gray, in Proc. Am. Acad. vi. (1864–65) 546 = Baeria leptalea.
longifolia, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 380 = Baeria gracilis.
maritima, A. Gray, in Proc. Am. Acad. vii. (1868) 358 = Baeria maritima.
microglossa, *DC. Prod.* v. 664.—Calif.
nana, D. Eaton, in Bot. King, Exp. 174 = Eatonella nivea.
parviflora, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 381 = Baeria gracilis.
platycarpha, A. Gray, in Torr. Bot. Mex. Bound. 97 = Baeria platycarpha.
tenerrima, DC. Prod. v. 664 = Baeria gracilis ?

BURSA, [Weber, in] Wigg. Prim. Fl. Hols. 47 (1780) = **Capsella**, Medic. (1792) (Crucif.).
astoris, [Weber,] l. c. = C. Bursa-pastoris.

BURSA-PASTORIS, [Tourn.] Rupp. Fl. Jen. 77. ed. Hall. (1745) 87 = **Capsella**, Medic. (1792) (Crucif.).

BURSARIA, Steud. Nom. ed. II. i. 241 (1840) sphalm. = **Bursaria**, Thou. (Menisp.).

BURSARIA, Cav. Ic. iv. 80. t. 350 (1797). *PITTOSPOREAE*, Benth. & Hook. f. i. 182.
dicarnoides, Putterl. Syn. Pittosp. 20 = Marianthus procumbens.
incana, Lindl. in Mitch. Journ. Trop. Austral. 224 = spinosa.
inermis, *Blanco, Fl. Filip.* ed. II. 134.—Ins. Philipp.
procumbens, Putterl. Syn. Pittosp. 20 = Marianthus procumbens.
spinosa, *Cav. Ic.* iv. 80. t. 350.—Austral.
Stuartiana, F. Muell. ex Klatt, in Linnaea, xxviii. (1856) 586 = Marianthus procumbens.

BURSERA, ' Jacq.' ex Linn. Sp. Pl. ed. II. 471 (1762).
BURSERACEÆ, Benth. & Hook. f. i. 324.
BUSSERIA, Cramer, Disp. Syst. 144 (1808).
DAMMARA, Gaertn. Fruct. fil. 100. t. 103 (1791).
ELAPHRIUM, Jacq. Enum. Pl. Carib. 3 (1760); Stirp. Am. i. 105. t. 71 (1763), nomen prius.
ICICA, Aubl. Pl. Gui. i. 387. t. 150 ad 152 (1775).
MARIGNIA, Comm. ex Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 351.
acuminata, Willd. Sp. Pl. iv. 1120 = Dacryodes hexandra.
acutifolia, Sieber, ex Steud. Nom. ed. II. i. 241 = Sorindeia madagascariensis.
Aloexylon, *Engl. in DC. Monog. Phan.* iv. 52.—Mexic.
altissima, *Baill. Hist. Pl.* v. 297.—Guiana.
angustata, *Griseb. Cat. Pl. Cub.* 65.—Cuba.
bahiensis, Salzm. ex Engl. in DC. Monog. Phan. iv. 277, 557 = Tapiria guianensis.
balsamifera, Pers. Syn. i. 413 = Hedwigia balsamifera.
bicolor, *Engl. in DC. Monog. Phan.* iv. 53.—Mexic.
bipinnata, *Engl. l. c.* 49.—Mexic.
caudata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 614 = Protium Sagotianum.
cinerea, *Engl. in DC. Monog. Phan.* iv. 48.—Mexic.
cuneata, *Engl. l. c.* 56.—Mexic.
decandra, *Baill. Hist. Pl.* v. 302, 297.—Guiana.
Delpechiana, *Poiss. ex Engl. in DC. Monog. Phan.* iv. 53.—Mexic.
excelsa, *Engl. l. c.* 57.—Mexic.
fagaroides, *Engl. l. c.* 48.—Mexic.
Galeottiana, *Engl. l. c.* 47.—Mexic.
glabra, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 304.—N. Granat.
glauca, *Griseb. Cat. Pl. Cub.* 66.—Cuba.
gracilis, *Engl. in DC. Monog. Phan.* iv. 50.—Mexic.
grandifolia, *Engl. l. c.* 45.—Mexic.
graveolens, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 303.—Peruv.
guianensis, *Baill. Hist. Pl.* v. 297.—Guiana.
gummifera, *Linn. Sp. Pl.* ed. II. 471.—Am. trop.
heptaphylla, *Wright, in Sauv. Fl. Cub.* 90.—Cuba.
heterophylla, *Engl. in DC. Monog. Phan.* iv. 46.—Mexic.
Hindsiana, *Engl. l. c.* 58.—N. Granat.
Icicariba, *Baill. Hist. Pl.* v. 297.—Bras.
integerrima, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 303.—N. Granat.
javanica, *Baill. Hist. Pl.* v. 297.—Java.
jorullensis, *Engl. in DC. Monog. Phan.* iv. 57.—Mexic.
Karsteniana, *Engl. l. c.* 41.—Venezuela.
Karwinskii, *Engl. l. c.*—Mexic.
Kerberi, *Engl. l. c.*—Mexic.
lancifolia, *Engl. l. c.* 49.—Mexic.
lanuginosa, *Engl. l. c.* 58.—Mexic.
leptophlœos, *Mart. in Flora*, xi. (1828) ii. 393.—Bras.
macrostachya, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 613.—Guiana.
Martiana, *Engl. in Mart. Fl. Bras.* xii. ii. 253.—Bras.
mexicana, *Engl. in DC. Monog. Phan.* iv. 51.—Mexic.
microphylla, *A. Gray, in Proc. Am. Acad.* v. (1861) 158.—Mexic.
multijuga, *Engl. in DC. Monog. Phan.* iv. 42.—Mexic.
obovata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 614.—Mexic.
obtusifolia, Lam. Encyc. ii. 768 = Protium obtusifolium.
orinocensis, *Engl. in Mart. Fl. Bras.* xii. ii. 254.—Bras.
ovalifolia, *Engl. in DC. Monog. Phan.* iv. 40.—Mexic.
paniculata, Lam. Encyc. ii. 768 = Canarium mauritianum.
pannosa, *Engl. in DC. Monog. Phan.* iv. 54.—Mexic.
penicillata, *Engl. l. c.* 52.—Mexic.
Schiedeana, *Engl. l. c.* 57.—Mexic.
Schlechtendalii, *Engl. l. c.* 41.—Mexic.
Sellowii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 613.—Bras.
serrata, Wall. ex Coleb. in Trans. Linn. Soc. xv. (1827) 361. t. 4 = Protium serratum.

BURSERA:—
sessiliflora, *Engl. in DC. Monog. Phan.* iv. 55.—Mexic.
simplicifolia, *DC. Prod.* ii. 78.—Jamaic.
submonoliformis, *Engl. in DC. Monog. Phan.* iv. 55.—Mexic.
Tacahamaca, Baill. Hist. Pl. v. 297 = Protium heptaphyllum.
Tatamaco, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 303.—N. Granat.
tomentosa, *Triana & Planch. l. c.* 304.—N. Granat.

BURSERIA, Loefl. It. Hisp. 194 (1758) = **Verbena**, Linn.

BURSERIA, Jacq. Stirp. Am. 94. t. 65 (1763) = **Bursera**, Linn.

BURSHIA, Rafin. in Med. Repos. N. York, v. (1808) 357 = **Myriophyllum**, Linn. (Haloragˢ.).
humilis, Rafin. l. c. et vi. (1808) 422 = M. ambiguum.

BURSINOPETALUM, Wight, Ic. t. 956 (1847) = **Mastixia**, Blume (Cornac.).
arborea, Wight, l. c. = M. arborea.
tetrandrum, Wight, ex Thw. Enum. Pl. Zeyl. 42 = M. tetrandra.

BURTONIA, R. Br. in Ait Hort. Kew. ed. II. iii. 12 (1811). *LEGUMINOSÆ*, Benth. & Hook. f. i. 468.
WEINERA, Reichb. Consp. 212ᵇ (1828).
conferta, *DC. Prod.* ii. 106.—Austral.
disomaefolia, Benth. Enum. Pl. Hueg. 82 = Latrobea diosmifolia.
foliolosa, *Benth. Fl. Austral.* ii 50.—Austral.
gompholobioides, *F. Muell. Fragm.* x. 84.—Austral.
Hendersonii, *Benth. Fl. Australˢ* ii. 51.—Austral.
minor, DC. Prod. ii. 106 = Gompholobium minus.
polycarpa, *Benth. Fl. Australˢ* ii. 51.—Austral.
pulchella, Meisn. in Lehm. Pl. Preiss. i. 41 = scabra.
scabra, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 12.—Austral.
sessilifolia, DC. Prod. ii. 106 = scabra.
solanacea, *Hort. ex Steud. Nom.* ed. II. i. 241.—Austral.
subalpina, F. Muell. in Trans. Phil. Soc. Victoria, i. (1855) 89 = Pultenæa rosea.
subulata, *Benth. Fl Australˢ* ii. 50.—Austral.
tenella, Meisn. in Lehm. Pl. Preiss. i. 43 = Latrobea tenella.
villosa, *Meisn. l. c.* 41.—Austral.

BURTONIA, Salisb. Parad. Lond. t. 78 (1807) = **Hibbertia**, Andr. (Dillen.).
grossulariæfolia, Salisb. l. c. = H. grossulariæfolia.

BUSBECKEA, Endl. Prod. Fl. Ins. Norf. 64 (1833) = **Capparis**, Tourn.
arborea, F. Muell. Fragm. i. 163 = C. nobilis.
canescens, F. Muell. l. c. v. 104 = C. canescens.
corymbifloraˢ, F. Muell. l. c. vii. 163 = C. lucida.
humistrata, F. Muell. l. c. v. 156 = C. humistrata.
inconstifolia, F. Muell. l. c. i. 164 = C. lomatiifolia.
Mitchelliiˢ, F. Muell. l. c.; et Vict. Pl. i. 53 = C. Mitchelliii.
nobilis, Endl. Prod. Fl. Ins. Norf. 64 = C. ornans.
ornans, F. Muell. Fragm. v. 104 = C. ornans.
Shanesiana, F. Muell. l. c. x. 94 = C. Shanesiana.
Thozetiana, F. Muell. l. c. v. 104 = C. Thozetiana.
umbonata, F. Muell. l. c. i. 164; v. 104 = C. umbonata.

BUSBECKEA, Mart. Cat. Hort. Monac. (1829) 89 = **Salpichroa**, Miers (Solanac.).
radicosa, Mart. l. c. = S. rhomboides.

BUSBEQUIA, Salisb. Gen. Pl. Fragm. 95 (1866) = **Hyacinthus**, Tourn. (Liliac.).
ciliaris, Salisb. l. c. = H. ciliatus.

BUSEA, Miq. Fl. Ind. Bat. fl. 782 (1856) = **Cyrtandromœa**, Zoll. (Gesnerac.).
acuminata, Miq. l. c. 783 = C. Miqueliana.
angustifolia, Miq. l. c. 784 = C. angustifolia.
asperifolia, Miq. l. c. 783 = Dichrotrichum asperifolium.

BUSEA:—
decurrens, Miq. Fl. Ind. Bat. ii. 783 = C. decurrens.
subsessilis, Miq. l. c. = C. acuminata.

BUSIPHO, Salisb. Gen. Pl. Fragm. 76 (1866) = **Aloë**, Tourn. (Liliac.).
ferox, Salisb. l. c. = A. ferox.

BUSSERIA, Cramer, Disp. Syst. 144 (1808) = **Bursera**, Jacq.

BUSTAMENTA, Alam. ex DC. Prod. v. 166 (1836) = **Bupatorium**, Tourn. (Compos.).
cordata, Alam. ex DC. l. c. = E. petiolare.
ovata, Mairet, ex DC. l. c. 168 = E. Bustamenta.

BUSTELINA, Fourn. in Mart. Fl. Bras. vi. iv. 288 (1885). *ASCLEPIADEÆ*.
Warmingii, *Fourn. l. c.*—Bras.

BUSTIA, Adans. Fam. ii. 127 (1763) = **Buphthalmum**, Linn. (Compos.).

BUSTILLOSIA, Clos, in C. Gay, Fl. Chil. iii. 106. t. 32 (1847) = **Asteriscium**, Cham. (Umbellif.).
chilensis, Clos, l. c. 108 = A. chilense.

BUTEA, Koen. ex Roxb. Fl. Corom. i. 22. t. 21 (1795) *LEGUMINOSÆ*, Benth. & Hook. f. i. 588.
MEGALOTROPIS, Griff. Notul. iv. 441 (1854).
MEIZOTROPIS, Volgt, Hort. Suburb. Calc. 289 (1845).
PLASO, Adans. Fam. ii. 325 (1763).
acuminata, Wall. Cat. n. 5443 = Spatholobus acuminatus.
Brasiliana, *DC. Prod.* ii. 415.—China.
frondosa, *Roxb. Pl. Corom.* i. 21.—Ind. or.; Burma.
frondosa, Wall. Cat. n. 5569, partim = superba.
gyrocarpa, R. Grah. in Wall. l. c. n. 5442 = Spatholobus gyrocarpus.
Loureiri, *Spreng. Syst.* iii. 186.—Cochinch.
minor, *Buch.-Ham. ex Wall. Cat.* n. 5439.—Reg. Himal.
parviflora, Roxb. Pl. Coromˢ. ii. 23 = Spatholobus Roxburghii.
pelsata, Pers. Syn. ii. 279 = Rudolphia pelsata.
scandens, Wall. Cat. n. 5441 = Spatholobus Roxburghii.
superba, *Roxb. Pl. Corom.* i. t. 22.—Ind. or.
volubilis, Vers. Syn. ii. 279 = Rudolphia volubilis.

BUTERAEA, Nees, in Wall. Pl. As. Rar. iii. 75, 196 (1832) = **Strobilanthes**, Blume (Acanthac.).
rhamnifolia, Nees, in DC. Prod. xi. 726 = S. rhamnifolius.
ulmifolia, Nees, in Wall. Pl. As. Rar. iii. 84 = S. rufescens.

BUTINIA, Boiss. Elench. 58 (1838) = **Conopodium**, Koch (Umbellif.).
buniosides, Boiss. l. c. 54 = Conopodium betinioides.
capnoides, Decne. in Jacquem. Voy. Bot. 70 = Chærophyllum capnoides.
cretica, Boiss. & Heldr. Diagn. Ser. I. x. 50 = Carum creticum.
glaucescens, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 61 (= *Biasoletia glaucescens*).—Persia.
libanotica, Boiss. l. c. 62 (= *Biasoletia meifolia*).—Syria.
macrocarpa, Boiss. & Sprun. in Ann. Sc. Nat. Sér. III. ii. (1844) 62 = Carum macrocarpum.
nodosa, Boiss. l. c. 61 (= *Biasoletia nodosa*).—Persia.
parnassica, Heldr. ex Nym. Consp. 302 (= *Biasoletia parnassica*).—Græcia.
stylosa, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 62 (= *Preyera stylosa*).—Græcia.

BUTNERIA, Duham. Arb. i. 128. t. 45 (1755) = **Calycanthus**, Linn.

BUTOMISSA, Salisb. Gen. Pl. Fragm. 91 (1866) = **Allium**, Tourn. (Liliac.).
tatarica, Salisb. l. c. = A. tataricum.

BUTOMOPSIS, Kunth, Enum. Pl. iii. 164 (1841). *ALISMACEÆ*. Benth. & Hook. f. iii. 1063.
TENAGOCHARIS, Hochst. in Flora, xxiv. (1841) 369.
cordofana, Kunth, ex Walp. Ann. i. 708 = Limnocharis

BUTOMOPSIS :—
lanceolata, *Kunth, Enum. Pl.* iii. 165.—Austral.
latifolia, Kunth, l. c. = lanceolata.

BUTOMUS, Linn. Syst. ed. I (1735). *ALISMACEAE*, Benth. & Hook. f. iii. 1006.
Caroliniani, Nesk. Delic. Gallo-Belg. i. 189 = umbellatus.
floridus, Gaertn. Fruct. t. 74. t. 19. f. 8 = umbellatus.
junceus, Turra. in Bull. Soc. Nat. Monc. (1837) n. VII. 187; (1838) 101, nomen; et xxvii. (1854) ii. 60 = umbellatus.
lanceolatus, Roxb. Hort. Beng. 89; Fl. Ind. ii. 315 = Butomopsis lanceolata.
latifolius, D. Don, Prod. Fl. Nep. 22 = Butomopsis lanceolata.
umbellatus, *Linn. Sp. Pl.* 372.—Europ.; As temp.
vulgaris, Gueldenst. Iter, ii. 22 = umbellatus.

BUTONICA, Lam. Encyc. i. 515 (1783); Juss. Gen. 326 (1789) = **Barringtonia**, Forst. (Myrtac).
acutangula, Lam. l. c. 591 = Barringt. acutangula.
alata, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 70 = Barringt. conoidea.
alba, Miers, l. c. 65 = Barringt. racemosa.
apiculata, Miers, l. c. 78.—Madag.
capsa, Miers, l. c. 77 = Barringt. racemosa.
calyptrata, Miers, l. c. 76 = Barringt. speciosa ?
ceylanica, Miers, l. c. 77 = Barringt. ceylanica.
edulis, Miers, l. c. 76 = Barringt. edulis.
incisa, Miers, l. c. 71 = Barringt. racemosa.
intermedia, Miers, l. c. 73 = Barringt. intermedia.
procera, Miers, l. c. 74.—Ins. Pacif.
racemosa, Juss. Gen. 326; Miers, l. c. 65 = Barringt. racemosa.
rosata, Miers, l. c. 72.—Ins. Philipp.
rubra, Miers, l. c. 70 = Barringt. racemosa.
Rumphiana, Miers, l. c. 68 = Barringt. litoniica.
samoensis, Miers, l. c. 75 = Barringt. samoensis.
speciosa, Lam. Encyc. i. 521 = Barringt. speciosa.
terratra, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 69 = Barringt. elongata.

BUTTNERIA, Duham. Arb. ed. II. 217. t. 47 (1801) = **Calycanthus**, Linn.

BUTTNERIA, Schreb. Gen. i. 145 (1789) = **Buettneria**, Loefl.

BUTTONIA, MacKen, ex Benth. in Hook. Ic. Pl. xi. 63. t. 1080 (1871). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 970.
natalensis, *MacKen. ex Benth. in Hook. l. c.*—Afr. austr.
natalis, MacKen, ex Benth. in Hook. l. c. = natalensis.

BUTTROSPERMUM, Kotschy, in Sitzb. Acad. Wien, Math. Nat. i. Abth. i. (1865) 857. *SAPOTACEAE*, Benth. & Hook. f. ii. 661.
MICADANIA, R. Br. in Denh. et Clapp. Voy. App. 239 (1826).
Kirkii, *Baker, in Oliver, Fl. Trop. Afr.* iii. 504.—Afr. trop.
niloticum, Kotschy, in Sitzb. Acad. Wien, Math. Nat. i. Abth. i. (1865) 858. t. 2 = seq.
Parkii, *Kotschy, l. c.* 859. t. 2.—Afr. trop.

BUXUS, Linn. Syst. ed. I (1735). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 366.
CRANTZIA, Sw. Prod. Veg. Ind. Occ. 86 (1788).
TRICERA, Schreb. Gen. 630 (1791); Sw. Fl. Ind. Occ. 331. t. 7 (1797).
acuminata, *Muell. Arg. in DC. Prod.* xvi. i. 15.—Cuba.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 2 = sempervirens.
aquartziana, Rich. ex Baill. Monog. Bux. 69 = citrifolia.
arborescens, Mill. Gard. Dict. ed. VIII. n. 1; Lam. Fl. Fr. ii. 303 = sempervirens.
argentea, Hort. ex Steud. Nom. ed. II. i. 242 = sempervirens.
aurea, Hort. ex Steud. l. c. = sempervirens.
australis, *A. Cunn. ex Steud. l. c.*—Austral.
balearica, *Lam. Encyc.* i. 511.—Ins. Balear.
californica, Hort. ex Baill. Monog. Bux. 66 = Simmondsia californica ?

BUXUS :—
caucasica, Hort. ex C. Koch, Dendrol. ii. ii. 476 = sempervirens.
chinensis, Link, Enum. Hort. Berol. ii. 386 = Simmondsia californica.
citrifolia, *Spreng. Syst.* iii. 847.—Am. austr.
cordifolia, Spreng. Syst. iii. 847 = Tricera cordifolia.
coriacea, Spreng. Syst. iv. Cur. Post. 314 = Sarcococca pruniformis.
crenata, Hort. ex C. Koch, Dendrol. ii. ii. 476 = sempervirens.
cruciata, Rich. ex Baill. Monog. Bux. 67 = Vahlii.
cubana, *Baill. l. c.* 71.—Cuba.
cuculiata, Hort. ex C. Koch, Dendrol. ii. ii. 476 = sempervirens.
dioica, Forsk. Fl Aegypt. Arab. 159 = Myrsine lutescens.
elegantissima, Hort. ex C. Koch, Dendrol. ii. ii. 477 = sempervirens.
Fortunei, *Carr. in Rev. Hortic* (1870–71) 519 — China.
glomerata, *Muell. Arg. in DC. Prod.* xvi. i. 17 — Cuba.
gonaïoides, *Muell Arg. l. c* 16 — Cuba.
haleppica, Hort. ex C. Koch, Dendrol. ii. ii. 477 = longifolia.
Handsworthii, Hort. ex C. Koch. l. c. 476 = sempervirens.
Harlandi, *Hance, in Journ. Linn. Soc.* xiii. (1873) 123.—China.
Hildebrandtii, *Baill. Adansonia*, xi. (1873–76) 268.—Afr. trop.
japonica, *Muell. Arg in DC. Prod.* xvi. i. 20.—Japon.
laevigata, *Spreng. Syst.* iii 847.—Jamaic.
longifolia, *Boiss Diagn.* Ser. I. xii. 107.—Syria.
madagascarica. *Baill. Monog. Bux.* 65.—Madag.
marginata, Hort. ex Steud. Nom ed. II. i. 242 = sempervirens.
microphylla, Sieb. & Zucc. in Abh. Akad. Muench. iv. (1846) 142 = japonica.
mucronata, Hort. ex Baill. Monog. Bux. 62 = sempervirens.
myrtifolia, Lam. Encyc. i. 511 = sempervirens.
obcordata-variegata, Fortune, in Gard. Chron. (1861) 735 = microphylla.
ovalifolia, Siebold, ex C. Koch, Dendrol. ii. ii. 479 = microphylla.
pulchella, *Baill. Monog. Bux.* 68.—Jamaic.
Purdieana, *Baill. l. c.* 70 (1859).—Jamaic.
rosea, *Muell. Arg. in DC. Prod.* xvi. i. 16.—Cuba.
rosmarinifolia, Hort. ex Baill. Monog. Bux. 62 = sempervirens.
rotundifolia, Hort. ex C. Koch, Dendrol. ii. ii. 479 = microphylla.
salicifolia, Hort. ex C. Koch, l. c. 476 = sempervirens.
saligna, D. Don, Prod. Fl. Nep. 63 = Sarcococca pruniformis.
sempervirens, *Linn. Sp. Pl.* 983.—Europ.; Orient; As. temp.
sempervirens, Thunb. Fl. Jap. 77 = japonica.
stenophylla, *Hance, in Journ. Bot.* vi. (1868) 331.—China.
subcolumnaris, *Muell. Arg. in DC. Prod.* xvi. i. 14.—Ind. occ.
suffruticosa, Mill. Gard. Dict. ed. VIII. n. 3 = sempervirens.
tenuifolia, Hort. ex Baill. Monog. Bux. 61 = sempervirens.
Vahlii, *Baill. Monog. Bux.* 67.—Ind. occ.
variegata, Hort. ex Steud. Nom. ed. II. i. 242 = sempervirens.
virens, Thunb. Fl. Jap. 77 = microphylla.
Wallichiana, Baill. Monog. Bux. 63 = sempervirens.
Wrightii, *Muell. Arg. in DC. Prod* xvi. i. 17.—Cuba.

BYBLIS, Salisb. Parad. Lond. t. 95 (1808). *DROSERACEAE*, Benth. & Hook. f. i. 664.
caerulea, R. Br. ex Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 306 = linifolia.
filifolia, Planch. l. c. 305 = linifolia.
gigantea, *Lindl. Swan River App.* 92.—Austral.
linifolia, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 307 = gigantea.
linifora, *Salisb. Parad. Lond.* t. 95.—Austral.

BYGNONIA, Barcena, in La Naturaleza, ii. (1873) 178 = **Bignonia**, Linn.

BYRONIA, Endl. in Ann. Wien. Mus. i. (1836) 184. *ILICINEAE*, Benth & Hook. f. i. 357.
POLYSTIGMA, Meisn. Gen. 252; Comm. 161 (1839).
arnhemensis, *F. Muell. Fragm.* ii. 119.—Australi.
Hookeri, Steud. Nom. ed. II. i. 242 = seq.
sandwicensis, *Endl. in Ann. Wien. Mus.* i. (1836) 184.—Ins. Sandvic.
taitensis, *A. Gray, Bot. U. St Expl. Exped.* i. 297. t. 25. —Ins. Tahiti.

BYRSA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. i. (Gen. ignot.).

BYRSANTHES, Presl, Prod. Monog. Lobel. 41 (1836) = **Siphocampylus**, Pohl (Campanul.).
Hallianus, Presl, l. c. 42 = S. Hallianus.
Humboldtiana, Presl, l. c. = S. niveus.
nivea, A. DC. in DC. Prod. vii. 408 = S. niveus.

BYRSANTHUS, Guillem. in Deless Ic. Sel. iii. 30. t. 62 (1837). *SAMYDACEAE*, Benth. & Hook. f. i. 800.
ANETIA, Endl. Gen. 923 (1839).
Brownii, *Guillem. l. c.*—Afr. trop.
epigynus, *Mast. in Oliver, Fl. Trop. Afr* ii. 499.—Afr. trop.

BYRSOCARPUS, Schum. & Thonn. Beskr. Guin. Pl. 236 (1827) *CONNARACEAE*, Benth. & Hook. f. i. 431.
Boivinianus, Baill. Adansonia, vii. (1866) 231 = Rourea Boivinianea.
coccineus, *Schum. & Thonn. Beskr. Guin. Pl.* 226.—Afr. trop.
maximus, *Baker, in Oliver, Fl. Trop Afr.* i. 453.—Afr. trop.
orientalis, *Baill. Adansonia*, vii. (1866) 230.—Afr. trop.
ovatifolius, *Baker, in Oliver, Fl. Trop. Afr.* i. 452.—Afr. trop.
parviflorus, Baill. Adansonia, vii. (1866) 229 = coccineus
parvifolius, Planch. in Linnaea, xxiii. (1850) 412 = coccineus.
Pervilleanus, Baill. Adansonia, vii. (1866) 232 = Rourea Pervilleana.
puniceus, Schum. & Thonn. Beskr. Guin. Pl. 227 = coccineus.

BYRSONIMA, Rich. ex Juss. in Ann. Mus. Par. xviii. (1811) 481. *MALPIGHIACEAE*, Benth. & Hook. f. i. 251.
adenophylla, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 314.—N. Granat.
Aerugo, *Sagot, l. c.* Sér. VI. xii. (1881) 178.—Guiana.
altissima, *DC. Prod.* i. 578.—Guinea.
amazonica, *Griseb. in Mart. Fl. Bras.* xii. i. 14.—Bras.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* v. 158. t. 448.—Venezuela.
arborea, *Mart. in Mart. Fl. Bras.* xii. i. 14.—Bras.
arthropoda, A. Juss. in Arch. Mus. Par. iii. (1843) 307 = laxiflora.
basiloba, *A. Juss. l. c.* 279.—Bras.
Beyerovana, A. Juss. l. c. 303 = spicata.
bicornicula, *A. Juss. l. c.* 284.—Bras.
biflora, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 167.—Cuba.
Blanchetiana, Miq. in Linnaea, xxii. (1849) 799 = bicornicula.
brachystachya, *DC. Prod.* i. 581.—Bras.
brachystachya, Poepp. ex Griseb. in Mart. Fl. Bras. xii. i. 10 = crispa.
bracteolaris, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 128.—Guiana.
brasiliensis, Griseb. in Linnaea, xiii. (1839) 255 = sericea.
bumelisefolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 79.—Bras.
cerantheres, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 121.—Guiana.
chrysophylla, *H. B. & K. Nov. Gen. et Sp.* v. 150.—Venezuela.
chrysophylla, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 81 = sericea.
cinerea, *DC. Prod.* i. 580.—Am. trop.
Clausseniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 281.—Bras.

BYRSONIMA:—
coccolobaefolia, *H. B. & K. Nov. Gen. et Sp.* v. 148.—Am. austr.
colocostachys, *Griseb. in Linnaea*, xxii. (1849) 3.—Guiana.
concinna, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 122.—Guiana.
contophylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 310.—Bras.
coriacea, DC. Prod. i. 580 = spicata.
corneaefolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 78. —Bras.
cotinifolia, Benth. Pl. Hartw. 277 = oaxacana.
cotinifolia, *H. B. & K. Nov. Gen. et Sp.* v. 152. t. 447. —Mexic.
crassifolia, *H. B. & K. Nov. Gen. et Sp. Pl.* v. 149.—Guiana.
crassifolia, Lunan, ex Griseb. Fl. Brit. W. Ind. 114 = cinerea.
crassifolia, Stend. Nom. ed. II. i. 242 = verbascifolia.
crispa, *A. Juss. ex Arch. Mus. Par.* iii. (1843) 306.—Bras.
cujabensis, *A. Juss. ex Mart. Herb. Fl. Bras.* 344. n. 657, nomen.—Bras.
cubensis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 292. —Cuba.
Cuminginna, A. Juss. l. c. 282 = crassifolia.
cuprea, *Griseb. in Mart. Fl. Bras.* xii. t. 19.—Bras.
cydoniaefolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 77.—Bras.
cydoniaefolia, Mart. in Flora, xxiv. (1841) Beibl. ii. 104 = Clusseniana.
dealbata, *Griseb. in Linnaea*, xiii. (1839) 253.—Bras.
densa, *DC. Prod.* i. 580.—Guiana.
diphylla, *DC. l. c.* 581.—N. Granat.
dispar, *Griseb. in Mart. Fl. Bras.* xii. t. 10.—Bras.
eglandulosa, A. Juss. in Arch. Mus. Par. iii. (1843) 278 = sericea.
elegans, *DC. Prod.* i. 580.—Guiana.
eriopoda, *DC. l. c.* 579.—Am. austr.
fasciculata, Mart. in Flora, xxiv. (1841) ii. Beibl. 61 = laevigata.
Fendleri, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 582.—Panama.
ferruginea, H. B. & K. Nov. Gen. et Sp. v. 151 = crassifolia.
frondosa, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 303.—Guiana.
Gardneriana, *A. Juss. l. c.* 296.—Bras.
guadalupensis, *G. Don, Gen. Syst.* i. 637.—Ind. occ.
Guilleminiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 276.—Bras.
gymnocalycina, *A. Juss. l. c.* 304.—Guiana.
Hostmanni, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 120.—Guiana.
hypargyrea, *Planch. ex Lindrn, Cat.* n. 8 (1853) 2.— N. Granat.
hypoleuca, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 391.—N. Granat.
intermedia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 82. —Bras.
inundata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 122.—Bras.
japurensis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 305.—Bras.
Karwinskiana, *A. Juss. l. c.* 288.—Mexic.
laevigata, *DC. Prod.* i. 580.—Guiana.
lanceolata, DC. l. c. = crassifolia.
lancifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 275. —Bras.
laurifolia, *H. B. & K. Nov. Gen. et Sp.* v. 147.—Venezuela.
laxiflora, *Griseb. in Linnaea*, xiii. (1839) 256.—Bras.
leucophlebia, *Griseb. in Mart. Fl. Bras.* xii. t. 15.— Bras.
ligustrifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 82. —Bras.
ligustrifolia, Mart. Herb. Fl. Bras. n. 1176 = intermedia.
linearifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 279.—Bras.
longibracteata, A. Juss. ex Mart. in Flora, xxiv. (1841) ii. Beibl. 61 = stipulacea.
lucida, *DC. Prod.* i. 580.—Am. bor.; Ind. occ.
Martiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 286. —Bras.
microphylla, *A. Juss. l. c.* 296.—Bras.
mollis, *DC. Prod.* i. 581.—Guiana.
montana, H. B. & K. Nov. Gen. et Sp. v. 149 = crassifolia.
monticola, A. Juss. & Mart. ex Griseb. in Mart. Fl. Bras. xii. t. 17 = coccolobaefolia.
Moritziana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 391.—Venezuela.
Mourella, Loud. Hort. Brit. 183 = crassifolia.
myricifolia, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 125.—Bras.
nervosa, *DC. Prod.* i. 579.—Bras.
nitida, *D. Don, Gen. Syst.* i. 636.—Peruv.
nitidifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 296.—Bras.
nitidissima, *H. B. & K. Nov. Gen. et Sp.* v. 152.— Venezuela.
oaxacana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 288. —Mexic.
oblongifolia, *A. Juss. l. c.* 300.—Bras.
obversa, *Mag. Surin. Surin. Sal.* 76.—Guiana.
Orbignyana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 284.—Bolivia.
oxyphylla, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 84.— Bras.
pachyphylla, Griseb. in Linnaea, xiii. (1839) 250 = verbascifolia.
pachyphylla, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 77.—Bras.
pallida, *DC. Prod.* i. 581.—Guiana.
paraensis, *Bruning, in Vet. Akad. Handl. Stockh.* (1854) 117.—Panama.
Pauliana, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 85.— Bras.
perseaefolia, *Griseb. in Linnaea*, xiii. (1839) 257.— Bras.
peruviana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 289. —Peruv.
phlomoides, *G. Don, Gen. Syst.* i. 636.—Bras.
pieuhensis, A. Juss. in Arch. Mus. Par. iii. (1843) 274 = sericea.
pinetorum, *Griseb. Cat. Pl. Cub.* 42.—Cuba.
Poeppigiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 308.—Bras.
propinqua, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 120.—Guiana.
psilandra, *Griseb. in Linnaea*, xiii. (1839) 251.—Bras.
pulchra, *DC. Prod.* i. 580.—Mexic.
punctulata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 309.—Bras.
reticulata, DC. Prod. i. 581 = Heteropteris platyptera.
reticulata, *Mutisech & Karst. ex Griseb. in Linnaea*, xxii. (1849) 6.—N. Granat.
rhombifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 299.—Bras.
rhopalaefolia, H. B. & K. Nov. Gen. et Sp. v. 148 = crassifolia.
rigida, *A. Juss. in Arch. Mus. Par.* iii. (1843) 298.— Bras.
rotunda, *Griseb. in Mart. Fl. Bras.* xii. t. 20.—Bras.
rufescens, *Bertol. Fl. Guatim.* 18.—Guatemala.
rugosa, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 118.—Guiana.
Salzmanniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 277.—Bras.
Schomburgkiana, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 123.—Guiana.
sericea, *DC. Prod.* i. 580.—Bras.
sessilifolia, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 124.—Guiana.
spanidilis, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 391.—Bras.
spicata, *Rich. ex Juss. in Ann. Mus. Par.* xviii. (1811) 481; *DC. Prod.* i. 580.—Am. trop.
spicata, Poepp. ex Spreng. Syst. ii. 383 = Heteropteris platyptera.
stigmatophora, *Schlecht. in Linnaea*, x. (1836) 241.— Mexic.
stipulacea, *A. Juss. in Arch. Mus. Par.* iii. (1843) 287.—Bras.
trinitensis, *A. Juss. l. c.* 296.—Ind. occ.
trioptericifolia, *A. Juss. l. c.* 286.—Am. austr.
umbellata, *Mart. ex A. Juss. l. c.* 301.—Bras.
vacciniaefolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* i. 84.—Bras.
variabilis, *A. Juss. l. c.* 78.—Bras.
verbascifolia, *Rich. ex Juss. in Ann. Mus. Par.* xviii. (1811) 481; *DC. Prod.* i. 579.—Am. trop.
viminifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 83. —Bras.
volubilis, DC. Prod. i. 581 = Hiraea Stanleana.

BYRSOPHYLLUM, Hook. f. in Benth. & Hook. f. Gen. ii. 83 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
ellipticum, *Hook f. l. c.*—Zeylan.
tetrandrum, *Hook. f. l. c.*; *Bedd. Forest. Man.* 184/2.— Ind. or.

BYSTROPOGON, L'Hérit. Sert. Angl, 19. t. 22, 23 (1788). *LABIATAE*, Benth. & Hook. f. ii. 1184.
ASTRMON, Regel, Ind. Sem. Hort. Petrop. (1860) 30.
MINTOSTACHYS, Spach, Hist. Vég. Phan. ix. 164 (1840).
bipinnatus, Roth, Nov. Pl. Sp. 225 = Lavandula Burmanni.
canariensis, *L'Hérit. Sert. Angl.* 20.—Ins. Canar.
canus, *Benth. Lab. Gen. et Sp.* 326.—Peruv.
coarctatus, *Thunb. & Schum. in Vidensk. Selsk. Skr.* iv. (1828) 84 = Hyptis pectinata.
confertus, Willd. ex Steud. Nom. ed. II. i. 248 = Mentha mollis.
dentatus, *Poir. Encyc. Suppl.* i. 755.—Peruv.
erianthus, Benth. Lab. Gen. et Sp. 737 = Cunnia erianthe.
glabrescens, *Benth. Pl. Hartw.* 145.—Peruv.
graveolens, Blume, Bijdr. 824 = Hyptis suaveolens.
maderensis, *Webb & Berth. Phyt. Canar.* iii. 66.—Ins. Madera.
marifolius, Jusa. ex Steud. Nom. ed. II. i. 248, 521 = Micromeria marifolia.
Meridiani, *C. Bolle, in Bonplandia*, viii. (1860) 281.— Ins. Canar.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 317.—N. Granat.
mollis, Salisb. Prod. 80 = canariensis.
odoratissimus, *C. Bolle, in Bonplandia*, viii. (1860) 281.—Ins. Canar.
origanifolius, *L'Hérit. Sert. Angl.* 20.—Ins. Canar.
pectinatus, L'Hérit. l. c. 19 = Hyptis pectinata.
piperina, *Lour., in Hook. Kew Journ.* viii. (1856) 299. —Ins. Madera.
plumosus, *L'Hérit. Sert. Angl.* 20. t. 22.—Ins. Canar.
punctatus, *L'Hérit. l. c.* 20. t. 23.—Ins. Madera.
reticulatus, Willd. ex Steud. Nom. ed. II. i. 248 = Mentha mollis.
serrulatus, *Webb & Berth. Phyt. Canar.* iii. 65. t. 151. —Ins. Canar.
sidaefolius, L'Hérit. Sert. Angl. 19 = Hyptis polyantha.
Smithii, Webb & Berth. Phyt. Canar. iii. 281 = punctata.
spicata, *Benth. Lab. Gen. et Sp.* 326.—Peruv.
suaveolens, L'Hérit. Sert. Angl. 19 = Hyptis suaveolens.
tomentosus, *Benth. Lab. Gen. et Sp.* 326.—Peruv.

BYTTHOPHYTON, Hook. f. Fl. Brit. Ind. iv. 286 (1884). *SCROPHULARINEAE* indicum, *Hook. f. l. c.*—Reg. Himal.

BYTNERIA, Jacq. Hort. Vindob. i. 10 (1770) = seq.

BYTTNERIA, Loefl. Iter Hisp. 313 (1758) = Buettneria, Loefl. (Sterculiac.).

BYTTNERIA, Stend. Nom. ed. II. i. 248 (1840) = Calyoanthus, Linn.

C

CAAPEBA, Plum. ex Adans. Fam. ii. 357 (1763) = Cissampelos, Linn. (Menisp.).

CABALLERIA, Ruiz & Pav. Prod. Fl. Per. et Chil. 141. t. 30 (1794) = Myrrhine, Linn.
coriacea, G. F. W. Mey. Prim. Fl. Esseq. 118 = M. floribunda.
dentata, Ruiz & Pav. Syst. Veg. 281 = M. dentata.
dependens, Ruiz & Pav. l. c. = M. dependens.
ferruginea, Ruiz & Pav. l. c. 280 = M. ferruginea.
latifolia, Ruiz & Pav. l. c. 279 = M. latifolia.
myrtifolia, Ruiz & Pav. ex DC. Prod. viii. 102 = M. dependens.
oblonga, Ruiz & Pav. Syst. Veg. 280 = M. Mangilla.
pulhuide, Ruiz & Pav. l. c. = M. pulhuide.
venosissima, Ruiz & Pav. l. c. 279 = M. venosissima.

CABANISIA, Klotzsch, ex Schlecht. in Abh. Naturf. Ges. Halle, vi. (1862) 176 = **Richhornia**, Kunth (Ponted.).
caracasana, Schlecht. l. c. = E. paradoxa.
Martiana, *Schlecht. l. c.* = E. paniculata.

CABOMBA, Aubl. Pl. Gui. i. 321. t. 124 (1775). *NYMPHÆACEÆ*, Benth. & Hook. f. i. 46.
NECTRIS, Schreb. Gen. i. 287 (1789).
VILLARSIA, Neck. Elem. ii. 110 (1790).
aquatica, *Aubl. l. c.*—Mexic.
aquatica, Juss. Gen. Pl. ii. 86 = caroliniana.
Aublettii, Michx. Fl. Bor. Am. i. 206 = caroliniana.
australis, *Speg. in Anal. Soc. Cient. Argent.* x. (1880) 219.—Reg. Argent.
caroliniana, *A. Gray, in Ann. Lyc. N. York*, iv. (1848) 46.—Am. bor.
furcata, *Schult. f. Syst.* vii. 1379.—Bras.
peltata, F. Muell. Native Pl. Vict. 15 = Brasenia peltata.
planhiensis, *Gardn. in Hook. Ic. Pl.* vii. t. 641.—Bras.
pinnata, Schult. f. Syst. vii 1379 = caroliniana.
Warmingii, *Casp. in Mart. Fl. Bras.* iv. ii. 142.—Bras.

CABRALBIA, A. Juss. Mém. Mus. Par. xix. (1830) 229. t. 16. *MELIACEÆ*, Benth. & Hook. f. i. 3 3.
affinis, A. Juss. l. c. 230 = polytricha.
Burchellii, *C. DC. in Mart. Fl. Bras.* xi. t. 171.—Bras.
Cangerana, *Saldanha, in Ann. Sc. Nat.* Sér. V. xix. (1874) 210.—Bras.
Clausseniana, *C. DC. in Mart. Fl. Bras.* xi. t. 173.—Bras.
corcovadensis, *C. DC. l. c.* 177.—Bras.
coriacea, *C. DC. l. c.* 181.—Bras.
Eichleriana, *C. DC. l. c.* 174.—Bras.
estrellensis, *C. DC. l. c.* 171.—Bras.
Gaudichaudii, *C. DC. l. c.* 172.—Bras.
glaberrima, *A. Juss. in Mém. Mus. Par.* xix. (1830) 270.—Bras.
Glaziovii, *C. DC. in Mart. Fl. Bras.* xi. t. 172.—Bras.
inaequilatera, *Casar. Decad. Nov. Stirp. Bras.* ii. 22.—Bras.
Jussiæana, *C. DC. in Mart. Fl. Bras.* xi. t. 171.—Bras.
laevis, *C. DC. l. c.* 176. t. 53.—Bras.
lagoënsis, *C. DC. l. c.* 178.—Bras.
Lundii, *C. DC. l. c.* 180.—Bras.
macrophylla, *C. DC. l. c.* 174.—Bras.
montana, *C. DC. l. c.*—Bras.
multijuga, *C. DC. l. c.* 176.—Bras.
oblongifolora, *C. DC. l. c.* 180.—Bras.
oligotricha, A. Juss. in Mém. Mus. Par. xix. (1830) 230 = polytricha.
pallescens, *C. DC. in Mart. Fl. Bras.* xi. t. 175. t. 52.—Bras.
pedunculata, *C. DC. l. c.* 179.—Bras.
piloza, *C. DC. l. c.* 177.—Bras.
Poeppigii, *C. DC. in DC. Monog. Phan.* i. 471.—Am. trop.
polytricha, *A. Juss. in Mém. Mus. Par.* xix. (1830) 230.—Bras.
Richardiana, *C. DC. in Mart. Fl. Bras.* xi. t. 176.—Bras. (vel Cuba).
Riedelii, *C. DC. l. c.* 180.—Bras.
rubiginosa, *C. DC. l. c.* 179.—Bras.
Sellowii, *C. DC. l. c.* 176.—Bras.
silvatica, *C. DC. l. c.* 178.—Bras.
sulcata, *C. DC. l. c.* 171.—Bras.
tomentosa, Casar. Decad. Nov. Stirp. Bras. ii. 22 = polytricha.
Warmingiana, *C. DC. in Mart. Fl. Bras.* xi. t. 178.—Bras.

CABRERA, Lag. Gen. et Sp. Nov. 5 (1816) = **Pasopalum**, Linn. (Gramin.).
chrysoblepharis, Lag. l. c. = P. exasperatum.

CACABUS, Berah. in Linnæa, xiii. (1839) 860. *SOLANACEÆ*, Benth. & Hook. f. ii. 896.
DICTYOCALYX, Hook. f. in Trans. Linn. Soc. xx. (1847) 302.
STYLETOTHECA, Regel, Gartenfl. (1858) 292. t. 68.
TRIPOGITON, Benth. Bot. Voy. Sulph. 148 (1844).

CACABUS:—
inflatus, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 255.—Peruv.
maximus, *Benth. & Hook. f. Gen.* ii. 897.—N. Granat.
mexicanus, *S. Wats. in Proc. Am. Acad.* xviii. (1883–83) 127.—Mexic.
nolanoides, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 255.—Peruv.
prostratus, *Bernh. in Linnæa*, xiii. (1839) 860.—Peruv.

CACALIA, Linn. Sp. Pl. 834 (1753) = **Senecio**, Tourn.
abyssinica, R. Br. in Salt, Abyss. App. 65 = S. abyssinicus.
abyssinica, Walp. Ann. ii. 908 = S. tuberosus.
acaulis, Linn. f. Suppl. 364 = S. acaulis, *Sch. Bip.*
acerifolia, Siebold, ex Maxim. in Bull. Acad. Pétersb. xix. (1874) 487, nomen.—Japon.
acetosæfolia, Jacq. ex Steud. Nom. ed. II. ii. 243 = S. articulatus.
aconitifolia, Bunge, Enum. Pl. Chin. Bor. 37 = S. aconitifolioides.
aconitifolia, Miq. ex Maxim. in Bull. Acad. Pétersb. xix. (1874) 487 = S. adenostyloides.
acuminata, Pisch. ex Herd. in Bull. Soc. Nat. Mosc. xlii. (1870) 1. 215 (= *Molgedium hastatum*).
adenostyloides, Franch. & Sav. ex Maxim. in Bull. Acad. Pétersb. xix. (1874) 486 = S. adenostyloides.
albida, Cass. ex Schur, Enum. Pl. Transs. 301 = Adenostyles alba.
albifrons, Linn. f. Suppl. 358 = Adenostyles alba.
Allioriæ, Gouan, Illustr. 65 = Adenostyles alba.
alliariæfolia, Lam. Encyc. i. 592 = Adenostyles viridis.
alliariæfolia, Poir. Voy. Barb. ii. 235 = Petasites fragrans.
alpina, Linn. Sp. Pl. 886 = Adenostyles viridis.
alpina, Mill. Gard. Dict. ed. VIII. n. 1 = Adenostyles alba.
amplifolia, Pres. Prod. vi. 328 = S. amplifolius.
angulata, Vahl, Symb. Bot. iii. 92 = Emilia angulata.
angulata, Vell. Fl. Flum. 386; viii. t. 55 = Mikania cordifolia.
anguina, Wall. Cat. n. 3152 = Gynura angulosa.
angustifolia, H. B. & K. Nov. Gen. et Sp. iv. 159 = S. Humboldtianus.
angustifolia, Wall. Cat. n. 3163 = Emilia prenanthoidea.
Antenphorbium, Linn. Sp. Pl. 834 = S. Anteuphorbium.
appendiculata, Linn. f. Suppl. 362 = S. populifolius.
arborea, H. B. & K. Nov. Gen. et Sp. iv. 163 (= *Psacalium arborescens*).—Am. austr.
arbuscula, Thunb. Prod. Pl. Cap. 142 = Othonna arbuscula.
arenaria, H. B. & K. Nov. Gen. et Sp. iv. 165 = S. sabulosus.
articularis, Salisb. Prod. 187 = seq.
articulata, Linn. f. Suppl. 364 = S. articulatus.
asarifolia, Steud. Nom. ed. II. i. 248 = Faujasia asarifolia.
asclepiadea, Linn. f. Suppl. 352 = S. asclepiadeus.
atriplicifolia, Lam. Sp. Pl. 835 = S. atriplicifolius.
aurantiaca, Blume, Bijdr. 908 = Gynura aurantiaca.
aurantiaca, Wall. Cat. n. 3146 = Gynura nepalensis.
auriculata, DC. Prod. vi. 299 = S. daburicus.
baccharidifolia, Poepp. ex Phil. f. Cat. Pl. Vasc. Chil. 156 = S. baccharidifolius.
Beriandieri, DC. Prod. vi. 328 = S. desertorum.
bicolor, Roxb. Hort. Beng. 61; Fl. Ind. iii. 412 = Gynura bicolor.
bipinnata, Linn. f. Suppl. 353 = S. bipinnatus.
bulbosa, Lour. Fl. Cochinch. 485 = Gynura ovalis.
cancellata, Rudolphi, in Ledeb. Diss. Fl. Doming. (1806) = S. cancellatus.
candicans, Poepp. ex DC. Prod. vi. 325 = Culcitium Poeppigii.
candicans, Vahl, Symb. Bot. iii. 91 = S. candicans.
canescens, Poepp. ex DC. Prod. vi. 414 = S. Neaei.
canescens, Willd. Enum. Hort. Berol. Suppl. 427 = S. Haworthii.
Cattiarica, Gouan, Illustr. 65 = Adenostyles alpina.
carnosa, Ait. Hort. Kew. iii. 156 = S. elegans.
caucasica, Hoffm. Hort. Mosq. (1808) n. 598 = S. platyphyllus.
cervariæfolia, DC. Prod. vi. 328 = S. cervariæfolius.

CACALIA:—
chiapensis, A. Gray, in Proc. Am. Acad. xix. (1883) 58 = Senecio chiapensis.
cinerarioides, H. B. & K. Nov. Gen. et Sp. iv. 166 = S. Bonplandianus.
cursifolia, Hook. & Arn. Bot. Beech. Voy. 486 = S. cirsiifolius.
cissampelina, DC. Prod. vi. 331 = S. cissampelinus.
citrina, Boj. ex DC. l. c. 302 = Emilia citrina.
Clematites, Poir. Encyc. Suppl. ii. 4 = Faujasia flexuosa.
coccinea, Sims, Bot. Mag. t. 564 = Emilia flammea.
cordata, Vell. Fl. Flum. 386; viii. t. 53 = Mikania cordifolia.
Cor-Jesu, Vell. l. c. 340; viii. t. 71 = Mikania officinalis.
cordifolia, H. B. & K. Nov. Gen. et Sp. iv. 168. t. 360 = S. cardiophyllus.
cordifolia, Linn. f. Suppl. 351 = Mikania cordifolia.
cordiformis, Lam. Fl. Fr. ii. 76 = Adenostyles alpina.
crassinervia, Wall. Cat. n. 3155 = Notonia crassinervia.
crenata, Vell. Fl. Flum. 385; viii. t. 51.—Bras.
cuneifolia, Linn. Mant. i. 110 = S. cuneifolius.
Cunninghamii, D. Don, Prod. Fl. Nep. 179 = Gynura angulosa.
cuspidata, Bertol. Fl. Guatimal. 32 = S. guatimalensis.
cuspidata, Klatt, in Ann. Sc. Nat. Sér. V. xviii. (1873) 874.—Ins. Borbon.
cylindrica, Lam. Encyc. i. 529 = Othonna cylindrica.
cylindrifiora, Wall. Cat. n. 3150 = Gynura sarmentosa.
decomposita, A. Gray, Pl. Wright. ii. 99 = S. Grayanus.
decurrens, Vell. Fl. Flum. 340; viii. t. 72 = Baccharis genistelloides.
delphinifolia, Sieb. & Zucc. in Abh. Acad. Muench. iv. III. (1846) 190 = S. Zuccarinii.
denticulata, Hook. & Arn. Bot. Beech. Voy. 29 = S. denticulatus.
dichroa, Bong. Nov. Comp. Bras. 40. t. 7 = S. dichrous.
discolor, Griseb. Cat. Pl. Cub. 157 = S. discolor.
distichæ, Heyne, ex DC. Prod. v. 384 = Conyza Wallichii.
diversifolia, Torr. & Gray, Fl. N. Am. ii. 436 = S. floridanus.
dulcis, Salisb. Prod. 187 = S. ficoides.
echinata, Linn. f. Suppl. 353 = S. echinatus.
elata, Spreng. ex DC. Prod. vi. 314 = S. Ligularia.
emarginata, Vell. Fl. Flum. 336; viii. t. 53.—Bras.
farfaraefolia, Sieb. & Zucc. in Abh. Acad. Muench. iv. III. (1846) 190 = S. farfaraefolius.
ficoides, Linn. Sp. Pl. 834 = S. ficoides.
fimbrillifera, Cass. in Dict. Sc. Nat. xlviii. 460 = S. deltoideus.
Finlaysoniana, Wall. Cat. n. 3162 = Gynura sarmentosa.
flexuosa, Wall. Cat. n. 3160 = Faujasia flexuosa.
flexuosa, Willd. ex DC. Prod. vii. 69 = Trixis divaricata.
floridana, A. Gray, in Proc. Am. Acad. xix. (1883) 52 = S. floridanus.
foetens, Wall. Cat. n. 3156 = Gynura nepalensis.
fruticosa, Vell. Fl. Flum. 338; viii. t. 64.—Bras.
gigantea, Nees, Del. Sem. Hort. Vrat. (1841) 8 = S. stipitiflorus.
glabra, Ledeb. Ind. Sem. Hort. Dorpat. (1830) 16 = S. sagittatus.
glabra, Heyne, in Wall. Cat. n. 3145 = Emilia sonchifolia.
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = Adenostyles alpina.
glabra, Vill. Prosp. 30 = Adenostyles viridis.
glabrata, H. B. & K. Nov. Gen. et Sp. iv. 161 (= *Psacalium glabratum*).—Am. trop.
glandulosa, Salisb. Prod. 187 = Porophyllum ruderale.
glauca, Salisb. l. c. = S. succulentus, *Sch. Bip.*
gracilifiora, Wall. Cat. n. 3149 = S. gracilifiorus.
gracilis, Boj. ex DC. Prod. vi. 302 = Emilia gracilis.
grandiflora, Wall. Cat. n. 3147 = Notonia grandiflora.
hastata, Linn. Sp. Pl. 835 = S. sagittatus.
Haworthii, Sweet, in Loud. Hort. Brit. 336 = S. Haworthii.
heterogama, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1857) 107 = S. heterogamus.
heterophylla, Poepp. ex DC. Prod. vi. 414 = S. pilquensis.

CACALIA :—
hieracioides, Willd. Sp. Pl. iii. 1731 = Gynura auriculata.
hirsuta, Vill. Prosp. 31 = Adenostyles alba.
humifusa, Boj. ex DC. Prod. vi. 302 = Emilia humifusa.
humilis, H. B. & K. Nov. Gen. et Sp. iv. 166 = Senecio humilis.
hybrida, Steud. Nom. ed. II. i. 244 = Adenostyles alpina, etc.
iatrophoides, H. B. & K. Nov. Gen. et Sp. iv. 169 = S. jatrophoides.
ilicifolia, H. B. & K. l. c. 168 = S. scrubicarina.
incana, Bosc, ex DC. Prod. vi. 330 = S. Boscianus.
incana, Linn Sp. Pl. ed. II. 1169 = Gynura nitida.
indica, Lam. Encyc. t. 830 —Ind. or.
involucrata, H. B. & K. Nov. Gen. et Sp. iv. 166 = S. involucratus.
Kleinia, J. Grah. Cat. Bomb. *Pl.* 98 = Notonia balsamica.
Kleinia, Herb. Madr. ex C. B. Clarke, Comp. Ind. 176 = Notonia grandiflora.
Kleinia, Linn. Sp. Pl. 834 = S. Kleinia.
Kleinsana, Hill, Veg. Syst. iii. 144 = præc.
hirsinodes, H. B. & K. Nov. Gen. et Sp. iv. 164 (= *Pucalium hirsinoides*).—Am. trop.
laciniata, Jacq. Ic. Rar. 1. t. 68 = S. articulatus.
lacviniata, Wall. Cat. n. 3158 = Gynura lycopersicifolia.
laeta, Vell. Fl. Flum. 341; viii. t. 76.— Bras.
lanata, H. B. & K. Nov. Gen. et Sp. 162 = S. lanatus.
lanceolata, Nutt. Gen. Am. ii. 138 = S. Boscianus.
laurifolia, Linn. f. Suppl. 351 = Mikania laurifolia.
ledifolia, H. B. & K. Nov. Gen. et Sp. iv. 162 = S. ledifolius.
leucopappa, DC. Prod. vi. 330 = S. leucopappus.
leucophylla, Willd. Sp. Pl. iii. 1736 = Adenostyles candidissima.
Linaria, Cav. Ic. iii. 29. t. 257 = Porophyllum Linaria.
linearis, Labill. Nov. Holl. Pl. ii. 86. t. 178 = Bedfordia linearis.
linifolia, Heyne, ex DC. Prod. v. 25 = Vernonia cinerea.
linifolia, Linn. Mant. 109 = Porophyllum linifolium.
lobata, Moris, in Flora, xv. (1832) Lit. ii. 79.—Chili.
longifolia, Haw. Rev. Pl. Succ. 193 = S. pugioniformis.
lutea, Mill. Gard. Dict. ed. VIII. n. 10.—Ins. S. Helena.
Lycopersicum, Heyne, ex DC. Prod. vi. 301 = Gynura lycopersicifolia.
macrophylla, Bieb. Fl. Taur. Cauc. ii. 286 = S. platyphyllus.
macrophylla, Rafin, in Med. Repos. N. York, v. (1808) 354, nomen.—Hab. ?
maculata, Buch.-Ham. ex Steud. Nom. ed. II. i. 244 = Gynura sinuata.
Mentrasto, Vell. Fl. Flum. 339; viii. t. 69 = Ageratum conyzoides.
mikaniæfolia, DC. Prod. vi. 328 = S. mikaniæfolia.
mucronata, Heyne, in Wall. Cat. n. 3161 = Emilia flammea.
multiflora, Anct. ex Walp. Rep. vi. 266 = S. multiflorus.
napeæfolia, DC. Prod. vi. 328 = S. napeæfolius.
napellifolia, S. Schan. in Linnæa, xix. (1847) 732 = S. napellifolius.
Nardosmia, A. Gray, in Proc. Am. Acad. vii. (1868) 361 = Cacaliopsis Nardosmia.
nepalensis, Spreng. ex DC. Prod. vi. 300 = Gynura nepalensis.
nipponica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 181.—Japon.
nitida, H. B. & K. Nov. Gen. et Sp. iv. 162 = S. nitidus.
nitida, Vell. Fl. Flum. 337; viii. t. 59.—Bras.
nivea, Kunze, ex DC. Prod. vi. 325 = Calcitium Poeppigii.
obatata, Vell. Fl. Flum. 337; viii. t. 57 = Mikania biformis.
odora, Forsk. Fl. Aegypt. Arab. 146 = S. odorus.
odorata, Desf. Cat. Hort. Par. ed. III. 165, 400 = S. odoratus.
oligocephala, Scheele, in Linnæa, xviii. (1844) 480.—Hab. ?
oppositifolia, Poir. Encyc. Suppl. ii. 7 = Porophyllum oppositifolium.
ovalifolia, Salisb. Prod. 186 = S. Antenphorbium.

CACALIA :—
ovalis, Ker-Gawl. in Bot. Reg. t. 101 = Gynura pseudochina.
ovata, Wall. Fl. Carol. 196 = Senecio Walteri ?
packyphylla, Sch. Bip. ex Seem. Bot. Voy. Herald, 310 = S. sclerophyllus.
paniculata, Rafin. Ann. Nat. (1820) 14 = S. Nuttallii.
papillaris, Linn. Sp. Pl. 834 = S. papillaris.
parasitica, Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot. ii. 344 = S. parasiticus.
patens, H. B. & K. Nov. Gen. et Sp. iv. 164 = S. patens.
pedunculata, Vell. Fl. Flum. 338; viii. t. 66.—Bras.
peltata, H. B. & K. Nov. Gen. et Sp. iv. 170. t. 361 = S. peltiferus.
pendula, Forsk. Fl. Aegypt. Arab. 145 = S. pendulus.
penduliflora, A. Gray, in Proc. Am. Acad. xix. (1883) 58 = S. subpeltatus, *Sch. Bip.*
penicillata, Cass. in Dict. Sc. Nat. xlviii. 460 = S. penicillatus.
Petasites, Lam. Encyc. i. 531 = Adenostyles alba.
peucedanifolia, Jacq. Coll. Suppl. 149 = S. paniculatus.
pilosa, Vell. Fl. Flum. 338; viii. t. 61 = Mikania cordifolia.
pinnata, Willd. Enum. Hort. Berol. 850 = S. Othonnæ.
pinnatifida, Berg. Desc. Pl. Cap. 230 = S. pinnatifidus.
pinnatifida, Linn. Mant. ii. 281 = S. bipinnatus.
pinnatifida, Lour. Fl. Cochinch. ii. 486 = Gynura pinnatifida.
pinnatifida, Wight, ex DC. Prod. vi. 301 = Gynura lycopersicifolia.
Porophyllum, Linn. Sp. Pl. 834 = Porophyllum rudderalis.
praestans, Baker, in Mart. Fl. Bras. vi. iii. 384, sphalm. = Trixis mollissima.
prenoides, H. B. & K. Nov. Gen. et Sp. iv. 167 = S. albiflorus.
prenanthoides, Sieber, ex DC. Prod. vi. 302 = Emilia sonchifolia.
procumbens, Lour. Fl. Cochinch. ii. 485 = Gynura sarmentosa.
Pteranthus, Rafin. Ann. Nat. (1820) 14 = S. Nuttalli.
pubescens, Rafin. Fl. Ludov. 81 = Eupatorium hodivicianum.
pugioniformis, Salzm. ex Link, Enum. Hort. Berol ii. 310 = S. pugioniformis.
pulchella, H. B. & K. Nov. Gen. et Sp. iv. 160 = S. pulchellus.
pumila, Boj ex DC. Prod. vi. 302 = Emilia pumila.
punctulata, Vell. Fl. Flum. 339; viii. t. 66.—Bras.
pungens, H. B. & K. Nov. Gen. et Sp. iv. 160 = S. pungens.
purpurascens, Wall. Cat. n. 3157 = Gynura purpurascens.
quenqueloba, Thunb. Prod. Pl. Cap. 142 = S. quinqueloba.
racemosa, DC. Prod. vi. 330 = S. exsertus.
racemosa, Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot. ii. 341 = S. heleches.
radicans, Linn. f. Suppl. 354 = S. radicans.
radulæfolia, H. B. & K. Nov. Gen. et Sp. iv. 169 = S. tæniæfolius.
ramosa, Spreng. ex DC. Prod. vi. 326 = S. triligulatus.
reclinata, Roxb. Hort. Beng. 101; Fl. Ind. iii. 412.—Ins. Ind. or.
reclinata, Wall. Cat. n. 3151 = Gynura sarmentosa.
regia, Baker, in Mart. Fl. Bras. vi. iii. 384, sphalm. = Trixis divaricata.
remiformis, Lam. Fl. Fr. ii. 75 = Adenostyles viridis.
reniformis, Muhl. in Willd. Sp. Pl. iii. 1735 = S. Muehlenbergii.
repens, Linn. Mant. 110 = S. succulentus.
reticulata, Vahl, Symb. Bot. iii. 91 = Fajasia reticulata.
reticulata, Vell. Fl. Flum. 337; viii. t. 60.—Bras.
rhombifolia, Willd. Enum. Hort. Berol. 850 = S. platyphyllus.
rigida, Thunb. Prod. Pl. Cap. 149 = S. spinescens.
Rio-marinus, Vell. Fl. Flum. 339; viii. t. 67.—Bras.
rotundifolia, Willd. Phyt. 11. t. 6. f. 1 = Vernonia cinerea.
ruderalis, Sw. Prod. Veg. Ind. Occ. 110 = Porophyllum ruderale.

CACALIA :—
runcinata, H. B. & K. Nov. Gen. et Sp. iv. 168 = Senecio roseus.
runcinata, Lam. Encyc. i. 529 = S. articulatus.
sagittaria, Heyne, in Wall. Cat. n. 3159 = Gynura pseudochina.
sagittata, Vahl, Symb. Bot. iii. 91 = Emilia flammea.
sagittifolia, Mertens, in Linnæa, v. (1830) 68, nomen.—Ins. Koragin.
salicina, Labill. Nov. Holl. Pl. ii. 87. t. 179 = Bedfordia salicina.
sarmentosa, Blume, Bijdr. 907 = Gynura sarmentosa.
sarmentosa, Lasch. ex Hook. f. Fl. Brit. Ind. iii. 352 = S. candicans.
Sarracenia, Bianco, Fl. Filip. ed. I. 618 = Gynura angulosa.
sarracenica, Linn. Sp. Pl. ed. II. 1169 = S. Cacalaster.
scabra, Vahl, Symb. Bot. iii. 92 = Trixis brasiliensis.
scandens, Alt. Hort. Kew. iii. 167.—Afr. austr.
scandens, Thunb. Prod. Pl. Cap. 142 = S. deltoideus.
scandens, Zeyh. ex DC. Prod. vi. 304 = S. volubilis.
Schaffneri, A. Gray, in Proc. Am. Acad. xix. (1883) 53.—Mexic.
semperviræa, Spreng. ex DC. Prod. vi. 442 = Notonia grandiflora.
semperviræa, Vahl, Symb. Bot. iii. 92 = S. semperviræ.
semperviræa, Forsk. Fl. Aegypt. Arab. 146 = S. semperviræ.
senecioides, H. B. & K. Nov. Gen. et Sp. iv. 168 = S. senuayensis.
septemnata, Vell. Fl. Flum. 338; viii. t. 63.—Bras.
sessilifolia, Hook. & Arn. Bot. Beech. Voy. 436 = S. sessilifolius.
sessilis, Vell. Fl. Flum. 341; viii. t. 78 = Baccharis articulata.
sinuata, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 25 = S. albo-luteocens.
sonchifolia, Forsk. Fl. Aegypt. Arab. p. cxix = Emilia angulata.
sonchifolia, Hort. ex Steud. Nom. ed. II. i. 244 = Emilia sagittata.
sonchifolia, Linn. Sp. Pl. 835 = Emilia sonchifolia.
sonchifolia, Wall. Cat. n. 3144 D = Lactuca remotiflora.
spathulata, Moç. & Sesse, ex D. Don, in Trans. Linn. Soc. xvi. (1830) 250 = Chaptalia spathulata.
suaveolens, Linn. Sp. Pl. 835 = S. suaveolens.
suffruticosa, Linn. Mant. i. 109 = Porophyllum linifolium.
tabularis, A. Gray, in Proc. Am. Acad. xix. (1883) 52 = S. tabularis.
tenuissima, Boj. in Rapp. Soc. Maur. vi. 27 = Emilia gracilis.
teres, Wall. Cat. n. 3164 = Emilia prenanthoides.
tertifolia, H. B. & K. Nov. Gen. et Sp. iv. 159 = S. teretifolius.
terminalis, Salisb. Prod. 187 = S. Kleinia.
ternata, Vell. Fl. Flum. 336; viii. t. 56 = Mikania spiffola.
thyrsoidea, Spreng. ex DC. Prod. vi. 368 = S. alatus.
tuberosa, DC. Prod. vi. 328 = S. calliosus.
tomentosa, Eckl. ex DC. l. c. 403 = S. senposus.
tomentosa, Haw. Misc. 189 = S. Haworthii.
tomentosa, Jacq. Fl. Austr. iii. 30 = Adenostyles alba.
tomentosa, Linn. f. Suppl. 353 = S. oligantha.
tomentosa, Vell. Fl. Flum. 341; viii. t. 75.—Bras.
tomentosa, Vill. Prosp. 31 = Adenostyles candidissima.
triangularis, Vell. Fl. Flum. 338; viii. t. 62 = Mikania cordifolia.
trichobata, Vell. l. c. 336; viii. t. 54 = Mikania glomerata.
trinervia, Vell. l. c. 338; viii t. 65.—Bras.
tripkylla, Vell. l. c. 341; viii. t. 74.—Bras.
tuberosa, Bosc, ex DC. Prod. vi. 30 = Krigia Dandelion.
tuberosa, Dellie, in Ferr. & Gall. Voy. Abyss. iii. 142 = S. tuberosus.
tuberosa, Nutt. Gen. Am. ii. 138 = S. Nuttallii ?
tussilaginoides, H. B. & K. Nov. Gen. et Sp. iv. 168 = S. farfarus.
umbifera, Schum. & Thonn. Beskr. Guin. Pl. 382 = Gynura cernua.
vaccinioides, H. B. & K. Nov. Gen. et Sp. iv. 163 (= *Pucalium vaccinioides*).—Am. trop.

CACALIA :—
verbascifolia, Sibth. & Sm. Fl. Graec. ix. 35 = Senecio thapsoides.
villosa, Jacq. Coll. ii. 347 = S. purpureus.
villosa, Vell. Fl. Flum. 337 ; vili. t. 56.—Bras.
volubilis, Blume, Bijdr. 903 = S. araneosus.
volubilis, Buch.-Ham. ex DC. Prod. vi. 370 = S. Boinalis.
Wightiana, Wall. ex DC. Prod. vi. 370 = S. scandens.

CACALIOPSIS, A. Gray, in Proc. Am. Acad. xix. (1883) 50. *COMPOSITAE*.
Nardosmia, *A. Gray, l. c.*—Am. bor. occ.

CACAO, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Theobroma**, Linn. (Sterculiac.).
bicolor, Poit. Encyc. Suppl. ii. 7 = T. bicolor.
guianensis, Aubl. Pl. Gui. ii. 683 = T. Cacao.
minor, Gaertn. Fruct. ii. 190 = T. Cacao.
sativa, Aubl. Pl. Gui. ii. 689 = T. Cacao.
sylvestris, Aubl. l. c. 687 = Genipa Merianae.
Theobroma, Tussac, Fl. Med. Ant. t. 13 = T. Cacao.

CACARA, Rumph. ex Thou. in Dict. Sc. Nat. vi. 85 (1805) = **Pachyrhizus**, Rich. (Legumin.).

CACATALI, Adans. Fam. ii. 218 (1763) = **Pedalium**, Royen (Pedalin.).

CACOINIA, Savi, Cose Bot. I. t. 1 (1832). *BORAGINEAE*, Benth. & Hook. f. ii. 846.
ANISANTHERA, Rafin. Fl. Tellur. iii 80 (1838).
Actinobole, *Bunge, Heliac.* 8.—Persia.
Celsii, Boiss. Diagn. Ser. I. vi. 133 = glauca.
debia, *Bunge, Lehm. Rel.* (1847) 242.—As. centr.
glauca, *Savi, Cose Bot.* I. t. 1. f. 1–6 (1832).—Persia ; Beluchist.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 31.—Persia.
Rauwolfii, A. DC. in DC. Prod. x. 167 = Russellii.
Rauwolfii, *C. Koch, in Linnaea*, xvii. (1843) 308.—As. Min. ; Persia.
Russellii, *Boiss. Diagn.* Ser. I. xi. 134.—Syria.
strigosa, *Boiss. l. c.*—Persia.

CACHRIS, D. Dietr. Syn. Pl. i. 586 (1839) = **Cachrys**, Tourn. (Umbellif.).

CACHRYDIUM, Link, Handb. i. 339 (1829) = **Hippomarathrum**, Link (Umbellif.).
Libanotis, Link, Handb. i. 339 = H. Libanotis.

CACHRYS Linn. Syst. ed. I (1735) ; Gen. ed. I. 75 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 904.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 338 = Diplolophium africanum.
acaulis, DC. Prod. iv. 238 = Prangos Szovitzii.
alata, Hoffm. Umb. i. 176 ; Bieb. Fl. Taur. Cauc. i. 217 = Prangos ferulacea.
alpina, *Bieb. l. c.*—Tauria.
ambigua, Salzm. ex DC. Prod. iv. 241 = Magydaris panacina.
amplexifolia, Pritz. Ind. Ic. 179 = seq.
amplifolia, Ledeb. in Eichw. Casp. 6. t. 9 = Hippomarathrum crispum.
athamanthoides, Bieb. Fl. Taur. Cauc. iii. 217 = Rumia athamanthoides.
calliosperma, Pall. Reise, iii. 668 = odontalgica.
cheilanthifolia, *Boiss. Fl. Orient.* ii. 988.—Persia.
cretica, Lam. Encyc. i. 359 = Leoskia cretica.
crispa, Pers. Syn. i. 311 = Hippomarathrum crispum.
crispata, *Pomel, Nouv. Mat. Fl. Atl.* 151.—Algeria.
cristata, DC. Prod. iv. 238 = Hippomarathrum cristatum.
cylindracea, Guss. ex DC. l. c. 239 = Prangos cylindracea.
decussata, Hornem. Hort. Hafn. ii. 960. Add. = Siler caucasicum.
dichotoma, Spreng. Pl. Umb. Prod. 21 = Capnophyllum dichotomum.
didyma, *Regel, in Act. Hort. Petrop.* v. (1877) 601.—Turkestan.
echinophora, Guss. Fl. Sic. Syn. i. 347 = Hippomarathrum cristatum.
eriantha, *DC. Prod.* iv. 238.—Persia.
fasciculata, *Boiss. Diagn.* Ser. I. x. 58.—Syria.
ferulacea, *Regel, in Act. Hort. Petrop.* v. (1877) 601.—Turkest.
hirta, *Bertol. Disquisit. de Pl. Nov.* 6.—Hab. ?

CACHRYS:—
humilis, *Schousb. Vextr. Marokko*, 183.—Marocco.
hungarica, Mill. Gard. Dict. ed. VIII. n. 5 = Magydaris panacina.
involucrata, Pall. ex Willd. fide Roem. & Schult. Syst. vi. 447 = Polylophium orientale.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 602.—Turkestan.
laevigata, Brouss. ex Ball, in Journ. Linn. Soc. xvi. (1878) 468 = Hippomarathrum cristatum.
laevigata, *Lam. Encyc.* i. 356.—Lusitan.
lanata, Hort. ex Steud. Nom. ed. II. i. 245 = Magydaris panacina, tomentosa.
latifolia, Bieb. Fl. Taur. Cauc. i. 219 = Siler caucasicum.
Libanotis, Georgi, Beschr. Russ. Reich. iii. iv. 845 = odontalgica.
Libanotis, Gouan, Illustr. 12 = laevigata.
Libanotis, Guss. Fl. Sic. Prod. i 358 = Prangos ferulacea.
Libanotis, Lina. Sp. Pl. 346 = Hippomarathrum Libanotis.
Libanotis, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 468 = Hippomarathrum cristatum.
Linearia, Mill. Gard. Dict. ed. VIII. n. 4 = Hippomarathrum Libanotis.
littoralis, Spreng. Umbell. Prod. iv. 20 = Daucus bessarabicus.
longiloba, DC. Prod. iv. 237 = Hippomarathrum crispum.
macrocarpa, *Ledeb Fl. Alt* i. 364.—Sibir. altaic.
maritima, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 259 = Crithmum maritimum.
microcarpa, Bieb. Fl. Taur. Cauc. i. 218 = Hippomarathrum crispum.
Morisoni, All. Auctuar. Fl. Pedem. 28 = laevigata.
nudicaulis, Gouan, ex DC. Prod. iv. 237 = Hippomarathrum crispum.
odontalgica, *Pall. Reise*, iii. 720.—Reg. Caucas. ; Sibir.
panacifolia, Vahl, Symb. Bot. i. 25 = Magydaris panacina.
panax-siculum, Ucria, in Roem. Arch. i. 68 = Magydaris tomentosa
papillaris, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 78.—Mesopotam.
pastinacea, Lam. Encyc. i. 360 = Magydaris panacina.
peregrina, Spreng. in Ges. Naturf. Fr Berl. Mag. vi. (1812) 259 = Capnophyllum dichotomum.
peucedanoides, *Desf. Fl. Atlant.* i. 250.—Afr. bor.
prangoides, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 76.—Persia.
pterocarpa, DC. Prod. iv. 237 = Hippomarathrum pterocblaenum.
pungens, Jan, ex Guss. Fl. Sic. Prod. i. App. 7 = Hippomarathrum cristatum.
sesseloides, Bieb. Fl. Taur. Cauc. iii. 217 = Rumia sesseloides.
sibirica, Steph. ex Fisch. in Cat. Sem. Hort. Petrop. (1828) = Phlodicarpos dahurica.
sibirica, Turcz. ex Ledeb. Fl. Ross. ii. 331 = Phlodicarpos villosa.
sicula, Gmelinest. ex Ledeb. Fl. Ross. ii. 357 = Hippomarathrum crispum.
sicula, Guss. Fl. Sic. Prod. i. 359 = Hippomarathrum cristatum.
sicula, Linn. Sp. Pl. ed. II. 355 = Hippomarathrum pterochlaenum.
sicula, Urv. in Mém. Soc. Linn. Par. i. (1822) 287 = Hippomarathrum Boccconi.
sphaerocarpa, Tenore, Prod. Fl. Nap. xix. ; Syn. 120 = Hippomarathrum Libanotis.
taurica, Willd. Sp. Pl. i. 1410 = Rumia taurica.
tomentosa, Desf. Fl. Atlant. i. 249 = Magydaris tomentosa.
tortuosa, Webb & Berth. Phyt. Canar. iii. 231 = Ferula tomentosa.
trifida, Mill. Gard. Dict. ed. VIII. n. 1 = laevigata.
triquetra, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1813) 259 = Prangos triquetra.
vaginata, Ledeb. Fl. Alt. i. 366 = Hippomarathrum vaginatum.

CACHVRIS, Zumag. Fl. Pedem. i. 39, 435 (1849) = prec.

CACONAPEA, Cham. in Linnaea, viii. (1833) 28 = **Herpestis**, Gaertn. (Scrophularia.).
gratioloides, Cham. l. c. 29 = H. gratioloides.

CACONOBEA, Walp. Rep. iii. 281 (1845) = prec.

CACOSMANTHUS, Miq. Fl. Ind. Bat. ii. 1040 (1856) = KAKOSMANTHUS, Hasak. = **Payena**, Linn. (Sapotac.).

CACOSMIA, H. B. & K. Nov. Gen. et Sp. iv. 289. t. 404 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 396.
CLAIRVILLEA, DC. Prod. v. 686 (1836).
XANTHOLEPIS, Willd. ex Less. in Linnaea, iv. (1829) 338.
quinquenervia, Steud. Nom. ed. II. i. 377, in syn. = seq. rugosa, *H. B. & K. Nov. Gen. et Sp.* iv. 289.—Peruv.

CACOTANIS, Rafin. New Fl. Am. ii. 43 (1836) = **Eclipta**, Linn. ? (Compos.).
ciliata, Rafin. l. c. 44.—Am. bor.

CACOUCIA, Aubl. Pl. Gui. i. 350. t. 129 (1775). *COMBRETACEAE*, Benth. & Hook. f. i. 688.
HAMBERGERA, Scop. Introd. 106 (1777).
SCHOUSBOEA, Willd. Sp. Pl. ii. 578 (1799).
bracteata, *Laws. in Oliver, Fl. Trop. Afr.* ii. 434.—Afr. trop.
chinensis, *A. Juss. ex DC. Prod.* iii. 22.—China.
coccinea, *Aubl. Pl. Gui.* i. 350. t. 129.—Guiana.
cordifolia, *Walp. Rep.* ii. 68.—Afr. trop.
exanulata, *O. Hoffm. in Linnaea*, xliii. (1880–82) 132.—Afr. trop.
lucida, Hasak. in Flora, xxvi. (1844) ii. 607 = Combretum trifoliatum.
panniculata, *Laws. in Oliver, Fl. Trop. Afr.* ii. 434.—Afr. trop.
platypters, *Welw. in Trans. Linn. Soc.* xxvii. (1868) 24.—Afr. trop.
trifoliata, DC. Prod iii. 22 = Combretum trifoliatum.
volutina, *S. Moore, in Journ. Bot.* xvii. (1880) 2.—Afr. trop.
villosa, *Laws. in Oliver, Fl. Trop. Afr.* ii. 433.—Afr. trop.

CACTUS, Linn. Syst. ed. I (1735) = *CACTEAE*, Benth. & Hook. f. Gen. i. 1008.
abnormis, Willd. Enum. Hort. Berol. Suppl. 31 = Cereus peruvianus.
Ackermannii, Haw. in Bot. Reg. t. 1331 = Phyllocactus Ackermanni.
acuelatissimus, Zeyh. ex Steud. Nom. ed. I. 181 (quid ?).
alatus, Hook. Bot. Mag. t. 2320 = Rhipsalis pachyptera.
alatus, Sw. Prod. Veg. Ind. Occ. 77 = Rhipsalis Swartziana.
alatus, Willd. Enum. Hort. Berol. Suppl. 35 = Phyllocactus phyllanthoides.
albisetosus, Haw. ex Spreng Syst. ii. 496 = Cereus albisetosus.
albisetus, Steud. Nom. ed. II. i. 341.—Peruv.
ambiguus, Bonpl. Jard. Malm. t. 36 = Cereus amboguus.
ambiguus, Gill. ex Lond. Hort. Brit. 194 = Mammillaria ambigua.
americanus, Vitm Summa Pl. iii. 209 (quid ?).
anisogonus, Hort. Angl. ex Foerst. Handb. Cact. ed. II. i. 764 = Cereus triangularis.
arboreus, Vell. Fl. Flum. 207 ; v. t. 26 = Opuntia arbores.
aurantiacus, Gill. ex Lond. Hort. Brit. Suppl. ii. 652 = Opuntia aurantiaca.
aurantiformis, Thicry, ex Steud. Nom. ed. I. 181 ; ed. II i. 245 (quid ?).
aureus, Meyen, Reise, i. 447 = Echinocactus aureus.
bicolor, Berland. ex Engelm. Cact. Mex. Bound. 21 = Echinocactus uncinatus.
Bleo, H. B. & K. Nov. Gen. et Sp. vi. 69 = Pereskia Bleo.
Bleo, Torr. in Ann. Lyc. N. York, ii. (1828) 202 = Opuntia arborescens.
boliviensis, Lem. Cact. 38.—Bolivia.
Bonplandii, H. B. & K. Nov. Gen. et Sp. vi. 69 = Opuntia Tuna.
brachypus, Lehm. Ind. Sem. Hort. Hamb. (1826) 17 = Cereus semilis.
brasiliensis, Willd. Enum. Hort. Berol. Suppl. 33 = Opuntia brasiliensis.
bulbispinus, Lem. Cact. 38.—Hab. ?
californicus, Torr. & Gray, *Fl. N. Am.* i. 555 = Opuntia serpentina ?

CACTUS :—
campechianus, Thiery, ex Steud. Nom. ed. I. 181 ; ed. II. i. 343 (quid ?).
caracasanus, Moç. & Sesse, ex DC. Prod. iii. 459 = Mammillaria lanifera.
caripensis, H. B. & K. Nov. Gen. et Sp. vi. 66 = Cereus caripensis.
cassythoides, Moç. & Sesse, ex DC. Prod. iii. 476 = Rhipsalis Cassytha.
caudatus, Gill. ex Loud. Hort. Brit. 194 = Mammillaria caudata.
chiloënsis, Colla, ex Steud. Nom. ed. II. i. 345 = Cereus chiloënsis.
chinensis, Roxb. Hort. Beng. [37] ; Fl. Ind. ii. 476 = Opuntia Dillenii.
chlorocarpus, H. B. & K. Nov. Gen. et Sp. vi. 67 = Cereus chlorocarpus.
clavatus, Lem. Cact. 88 = Opuntia clavata, *Engelm.*
coccinellifer, Steud. Nom. ed. I. 181, vide cochenillifer.
coccineus, Gill. ex Loud. Hort. Brit. 194 = Mammillaria coccinea.
cochenillifer, Linn. Sp. Pl. 468 = Nopalea coccinellifera.
cochenillifer, Moç. & Sesse, ex DC. Prod. iii. 474 = Opuntia Hernandesii.
columnaris, Moç. & Sesse, ex DC. Prod. iii. 459 = Mammillaria acanthoplegma.
communis, Steud. Nom. ed. II. i. 345 = Melocactus communis.
compressus, Salisb. Prod. 348 = Opuntia vulgaris.
coquimbanus, Molina, Saggio Chile, 170 = Cereus eburneus.
cornigerus, Moç. & Sesse, ex DC. Prod. iii. 461 = Echinocactus cornigerus.
coronarius, Cav. Descr. 490 = Cereus cylindricus.
coronatus, Lam. Encyc. i. 537 = Melocactus communis.
coronatus, Willd. Enum. Hort. Berol. Suppl. 30 = Mammillaria coronaria.
corrugatus, Gill. ex Loud. Hort. Brit. 194 = Echinocactus corrugatus.
crispatus, Moç. & Sesse, ex DC. Prod. iii. 462 = Echinocactus strigens.
cruciformis, Vell. Fl. Flum. 207 ; v. t. 29 = Cereus cruciformis.
curassavicus, Linn. Sp. Pl. 469 = Opuntia curassavica.
curvispinus, Bert. in 'Merc. Chil. 1829.'—Cf. Bull. Férussac, xx. (1830) 108.—Chili.
cylindricus, James, Cat. 182 = Opuntia arborescens.
cylindricus, Lam. Encyc. i. 539 = Opuntia cylindrica.
cylindricus, Orteg. Hort. Matr. 128. t. 16 = Mammillaria coronaria.
cylindrus, Vell. Fl. Flum. 207 ; v. t. 31 = Rhipsalis cylindrica.
decumanus, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia decumana.
depressus, DC. Cat. Hort. Monsp. 84 = Mammillaria discolor.
depressus, Haw. Syn. Pl. Succ. 178 = Echinocactus depressus.
Dillenii, Ker-Gawl. in Bot. Reg. t. 255 = Opuntia Dillenii.
divaricatus, Lam. Encyc. i. 540 = Cereus divaricatus.
eburneus, Link, Enum. Hort. Berol. ii. 22 = Cereus eburneus.
eburneus, Salm. ex Spreng. Syst. ii. 494 = Cereus eburneus.
echinocarpus, Lem. Cact. 88.—Hab. ?
elatior, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia elatior.
elegans, Link, Enum. Hort. Berol. ii. 25 = Phyllocactus phyllanthoides.
elongatus, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia decumana ?
Emoryi, Lem. Cact. 88 = Cereus Emoryi.
enniformis, *Biden, in Gard. Chron.* xx. (1883) 53.—Hab. ?
Epidendrum, Linn. Amoen. Acad. viii. 257 = Rhipsalis undulata ?
erinaceus, Haw. Suppl. 74 = Cereus erinaceus.
euphorbioides, Spreng. Syst. ii. 496 = Cereus euphorbioides.
fasciculatus, Meyen, Reise, i. 447 = Echinocactus fasciculatus.
fasciculatus, Willd. Enum. Hort. Berol. Suppl. 33 = Rhipsalis fasciculata.
ferox, Nutt. Gen. Am. i. 296 = Opuntia missouriensis.
ferox, Willd. Enum. Hort. Berol. Suppl. 33 = Opuntia ferox.

CACTUS :—
Ficus, Thunb. Fl. Jap. 198 = seq.
Ficus-indica, Linn. Sp. Pl. 468 = Opuntia Ficus-indica, decumana.
fimbriatus, Desc. Fl. Antill. vi. 160 = Cereus Swartzii.
fimbriatus, Lam. Encyc. i. 539 = Cereus fimbriatus.
fimbriatus, Moç. & Sesse, ex DC. Prod. iii. 475 = Pereskia lychnidiflora.
flagelliformis, Linn. Sp. Pl. 467 = Cereus flagelliformis.
flavescens, DC. Cat. Hort. Monsp. 83 = Mammillaria flavescens.
flavispinus, Colla, Hort. Rip. 24 = Cereus flavispinus.
foliosus, Gill. ex Loud. Hort. Brit. 194 = Echinocactus foliosus.
foliosus, Willd. Enum. Hort. Berol. Suppl. 35 = Opuntia foliosa.
fragilis, Nutt. Gen. Am. i. 296 = Opuntia fragilis.
fruticescens, Moç. & Sesse, ex Pfeiff. Enum. Diag. Cact. 178 = Pereskia rotundifolia.
fulvispinosus, Spreng. Syst. ii. 497 = Cereus Royeni.
funalis, Spreng. l. c. 479 = Rhipsalis funalis.
gariepinus, Kunth, Syn. Pl. Aeq. iii. 270 = Cereus caripensis.
gibbosus, Haw. in Bot. Reg. t. 187 = Echinocactus gibbosus.
glomeratus, Lam. Encyc. i. 257 = Mammillaria glomerata.
grandiflorus, Linn. Sp. Pl. 467 = Cereus grandiflorus.
grandifolius, Link, Enum. Hort. Berol. ii. 25 = Pereskia grandifolia.
Haworthii, Spreng. Syst. ii. 495 = Cereus Haworthii.
Helicteres, Moç. & Sesse, ex DC. Prod. iii. 460 = Mammillaria Helicteres.
heptagonus, Linn. Sp. Pl. 466 = Cereus heptagonus.
heptagonus, Vell. Fl. Flum. 205 ; v. t. 19.—Bras.
heteropterus, DC. Prod. iii. 470, sphalm. = praec.
hexagonus, Lam. Encyc. i. 538 = Melocactus monoclonos.
hexagonus, Linn. Sp. Pl. 466 = Cereus hexagonus.
hexagonus, Vell. Fl. Flum. 205 ; v. t. 18.—Bras.
Histrix, Salm-Dyck, Obs. Bot. (1822) = Cereus Hystrix.
horridus, H. B. & K. Nov. Gen. et Sp. vi. 70 = Pereskia horrida.
horridus, Salisb. Prod. 348 = Opuntia Tuna.
Humboldtii, H. B. & K. Nov. Gen. et Sp. vi. 66 = Cereus Humboldtii.
humifusus, Raf. Ann. Nat. (1820) 15 = Opuntia vulgaris.
humilis, Desf. ex Steud. Nom. ed. II. i. 346 = Opuntia vulgaris.
humilis, Haw. Misc. 187 = Opuntia Tuna.
humilis, Lag. Gen. et Sp. Nov. 16 = Opuntia foliosa.
hybridus, Hort. Angl. ex F. C. Van Geel, Sert. Bot. iv. (1832) t. 118.—Hab. ?
hystricacanthus, Lem. ex Steud. Nom. ed. II. i. 346 = Echinocactus hystricacanthus.
icosagonus, H. B. & K. Nov. Gen. et Sp. vi. 67 = Cereus icosagonus.
imbricatus, Lem. Cact. 88.—Hab. ?
indicus, Roxb. Hort. Beng. [37] ; Fl. Ind. ii. [475] = Opuntia Dillenii.
intortus, Mill. Gard. Dict. ed. VIII. n. 2 = Echinocactus intortus.
italicus, Tenore, ex Steud. Nom. ed. II. i. 346 (quid ?).
Jamacaru, Kostel. Allg. Med.-Pharm. Fl. iv. 1893 = Cereus Jamacaru.
Kegeneichii, Gmel. ex Salm-Dyck, fide DC. Prod. iii. 463 = Cereus multangularis.
lactus, H. B. & K. Nov. Gen. et Sp. vi. 68 = Cereus lactus.
Lemairii, Colla, Hort. Ripol. App. iii. 15. t. 7 = Melocactus communis.
lanatus, H. B. & K. Nov. Gen. et Sp. vi. 68 = Cereus lanatus.
lanceolatus, Haw. Misc. 188 = Opuntia lanceolata.
Langsdorffii, Lehm. Ind. Sem. Hort. Hamb. (1826) = Echinocactus Langsdorffi.
lanuginosus, Linn. Sp. Pl. 467 = Cereus repandus.
latispinus, Haw. in Phil. Mag. lxiii. (1824) 41 = Echinocactus latispinus.
Lecchii, Colla, Hort. Ripul. App. iv. 6. t. 2 = Cereus Lecchii.
Lemairii, Monvill. ex 'Lem. Cact. Gen. 105' = Echinocactus Lemairii.
Limbi, Lehm. Ind. Sem. Hort. Hamb. (1827) 16 = Cereus Limbi.
lucidus, Salisb. Prod. 349 = Pereskia aculeata.
lutzes, Thiery, ex Steud. Nom. ed. I. 183 ; ed. II. i. 346 (quid ?).

CACTUS :—
lyratus, Vell. Fl. Flum. 205 ; v. t. 21 = Rhipsalis salicornioides.
macracanthus, Salm-Dyck, Obs. (1820) 1 = Melocactus macracanthus.
magnimamma, Salm-Dyck, ex Steud. Nom. ed. II. i. 346 = Mammillaria magnimamma.
Mallisoni, Hort. ex Loud. Encyc. Pl. Suppl. i. 1302 = Cereus Smithii.
mammillaris, Linn. Sp. Pl. 466 = Mammillaria simplex.
mammillaris, Nutt. Gen. Am. i. 295 = Mammillaria missourensis.
melocactoides, Hoffmgg. Verz. Pfl. Nachtr. iii. 24 = Melocactus melocactoides.
Melocactus, Linn. Sp. Pl. 466 = Melocactus communis.
mensarum, Thiery, ex Steud. Nom. ed. I. 182 ; ed. II. i. 346 (quid ?).
meonacanthus, Lem. ex Steud. Nom. ed. II. i. 346 = Echinocactus meonacanthus.
micranthus, H. B. & K. Nov. Gen. et Sp. vi. 65 = Rhipsalis micrantha.
microdasys, Lehm. Ind. Sem. Hort. Hamb. (1827) 16 = Opuntia microdasys.
microhele, Spreng. Syst. ii. 494 = Mammillaria parvimamma.
monacanthos, Willd. Enum. Hort. Berol. Suppl. 33 = Opuntia monacantha.
moniliformis, Lam. Encyc. i. 541 = Cereus moniliformis.
moniliformis, Linn. Sp. Pl. 468 = Opuntia moniliformis.
multangularis, Moç. & Sesse, ex DC. Prod. iii. 462 = Echinocactus melocactiformis.
multangularis, Volgt, ex Steud. Nom. ed. I. 182 = Echinocactus recurvus.
multangularis, Willd. Enum. Hort. Berol. Suppl. 33 = Cereus multangularis.
nanus, H. B. & K. Nov. Gen. et Sp. vi. 68 = Cereus nanus.
nanus, Hort. ex Steud. Nom. ed. II. i. 346 = Opuntia vulgaris.
Napoleonis, Hort. ex Loud. Encyc. Pl. Suppl. i. 1302 = Cereus Napoleonis.
neglectus, Dehnhardt, Revist. Napol. i. 111. 166 (= Pereskiae sp.).
niger, Spreng. Syst. ii. 495 = Cereus niger.
nigricans, Haw. Misc. 187 = Opuntia nigricans.
nobilis, Haw. Syn. Pl. Succ. 174 = Cereus reductus.
nobilis, Lam. Encyc. i. 537 = Echinocactus intortus.
nobilis, Linn. Mant. 243 = Echinocactus recurvus.
nudus, Moç. & Sesse, ex DC. Prod. iii. 460 = Mammillaria nuda.
oboallatus, Moç. & Sesse, ex DC. l. c. 463 = Echinocactus oboallatus.
octogonus, Page, ex Steud. Nom. ed. II. i. 346 = Cereus hexagonus.
Opuntia, Blanco, Fl. Filip. ed. I. 414 = Nopalea coccinellifera.
Opuntia, Gaza. Fl. Sicul. Prod. 559 = Opuntia Ficus-indica.
Opuntia, Linn. Sp. Pl. 468 = Opuntia vulgaris.
opuntiaeflorus, Moç. & Sesse, ex Pfeiff. Enum. Diag. Cact. 178 = Pereskia opuntiaeflora.
Ottonis, Lehm. Ind. Sem. Hort. Hamb. (1827) 16 = Echinocactus Ottonis.
ovatus, Gill. ex Loud. Hort. Brit. 195 = Cereus ovatus.
ovoides, Lem. Cact. 88.—Hab. ?
oxypetalus, Moç. & Sesse, ex DC. Prod. iii. 470 = Phyllocactus latifrons.
paniculatus, Lam. Encyc. i. 540 = Cereus paniculatus.
parodanus, Hornem. Hort. Hafn. ii. 449 = Opuntia brasiliensis.
parasiticus, Lam. Encyc. i. 541 = Rhipsalis fasciculata.
parasiticus, Linn. Syst. ed. X. 1054 = Rhipsalis baccifera.
Parryi, Lem. Cact. 88.—Hab. ?
parvifolius, Ehrh. ex F. G. Dietr. Vollst. Lex. Gaertn. ii. 416.—Hab. ?
parvipinus, Haw. Suppl. 73 = Echinocactus parvispinus.
pendulinus, Sieber, ex DC. Prod. iii. 476 = Rhipsalis Cassyta.
pendulus, Sw. Prod. Veg. Ind. Occ. 77 = Rhipsalis Cassutha.
pentagonus, Linn. Sp. Pl. 467 = Cereus pentagonus.
pentagonus, Vell. Fl. Flum. 205 ; v. t. 35.—Bras.
Pentlandii, Lem. Cact. 88.—Hab. ?

CACTUS :—
Pereskia, Linn. Sp. Pl. 469 = Pereskia aculeata.
peruvianus, DC. Pl. Grasses, t. 68 = Cereus hexagonus.
peruvianus, Linn. Sp. Pl. 467 = Cereus peruvianus.
peruvianus, Willd. Enum. Hort. Berol. Suppl. 32 = Cereus chrysacanthus.
phyllanthoides, DC. Hort. Monsp. 84 = Phyllocactus phyllanthoides.
Phyllanthus, Linn. Sp. Pl. 469 = Phyllocactus Phyllanthus.
Phyllanthus, Vell. Fl. Flum. 208 ; v. t. 38.—Bras.
Pitajaya, Blanco, Fl. Filip. ed. II. 289 = Cereus triangularis.
Pitajaya, Jacq. Enum. Pl. Carib. 28 = Cereus variabilis.
placentiformis, Lehm. Ind. Sem. Hort. Hamb. (1826) 17 = Melocactus placentiformis.
polyanthus, Sims, Bot. Mag. t. 2691 = Opuntia polyanthus.
polygonus, Lam. Encyc. t. 539 = Cereus polygonus.
polymorphus, Gill. ex Loud. Hort. Brit. 195 = Cereus polymorphus.
portulacaefolius, Linn. Sp. Pl. 469 = Pereskia portulacaefolia.
prismaticus, Desf. Tabl. Hort. Par. 165 = Cereus triqueter.
prismaticus, Willd. Enum Hort. Berol. Suppl. 32 = Cereus pentagonus, prismaticus.
prolifer, Willd. ex Pfeiff. Enum. Diagn. Cact. 9 = Mammillaria parvimamma.
proliferus, Mill. Gard. Dict. ed. VIII. n. 6 = Mammillaria prolifera.
proteiformis, Desf. Tabl. Hort. Par. ed. II. 192 = Cereus proteiformis.
pruinosus, Monvill. ex Steud. Nom. ed. II. i. 246 = Echinocactus pruinosus.
pseudococcinelliger, Bertol. ' Exc. 11 virid. (1824) 4 '; ex DC. Prod. iii. 473 = Opuntia nigricans.
pseudomammillaria, Desf. Tabl. Hort. Par. ed. II. 191 = Mammillaria discolor.
Pseudotuna, Hort. ex Steud. Nom. ed. II. i. 246 = Opuntia Pseudotuna.
pubescens, Lem. Cact. 87.—Hab. ?
pusillus, DC. Cat. Hort. Monsp. 184 = Mammillaria pusilla.
pusilus, Haw. Misc. 188 = Opuntia fragilis.
pyramidalis, Salm-Dyck, Obs. (1820) 4 = Melocactus pyramidalis.
quadrangularia, Haw. Syn. Pl. Succ. 181 = Cereus caripensis.
quadrifus, Gill. ex Loud. Hort. Brit. 194 = Mammillaria quadrata.
recurvus, Mill. Gard. Dict. ed. VIII. n. 8 = Echinocactus recurvus.
reductus, Link, Enum. Hort. Berol. ii. 21 = Echinocactus gibbosus.
regalis, Spreng. Syst. ii. 496 = Cereus regalis.
repandus, Linn. Sp. Pl. 467 = Cereus gracilis.
reptans, Salm-Dyck, ex DC. Prod. iii. 468 = Cereus pentagonus.
reptans, Willd. Enum. Hort. Berol. Suppl. 33 = Cereus radicans.
reticulatus, Desc. Fl. Antill. t. t. 68.—Ind. occ.
Rosa, Vell. Fl. Flum. 206 ; v. t. 37 = Pereskia grandifolia.
rotundifolius, Mog. & Sesse, ex DC. Ill. 475 = Pereskia rotundifolia.
Royeni, Linn. Sp. Pl. 467 = Cereus Royeni.
Royeni, Willd. Enum. Hort. Berol. Suppl. 32 = Cereus crenulatus.
salicornioides, Steud. Nom. ed. II. i. 246 = Rhipsalis salicornioides.
Salmianus, Lem. Cact. 87.—Hab. ?
Scopa, Link, Enum. Hort. Berol. ii. 21 = Cereus Scopa.
senilis, Haw. in Phil. Mag. lxiii. (1824) 41 = Cereus senilis.
septum, H. B. & K. Nov. Gen. et Sp. vi. 66 = Cereus septum.
setiger, Gill. ex Loud. Hort. Brit. 196 = Opuntia setosa.
serpens, H. B. & K. Nov. Gen. et Sp. vi. 68 = Cereus serpens.
serpentinus, Lag. & Rodr. in Anal Cienc. Nat. iv. (1801) 261 = Cereus serpentinus.
serpentinus, Pers. Syn. ii. 99 = Cereus flagelliformis.
solitarius, Gill. ex Loud. Hort. Brit. 194 = Mammillaria solitaria.
speciosissimus, Desf. in Mém. Mus. Par. iii. (1817) 198, t. 9 = Cereus speciosissimus.

CACTUS :—
speciosus, Cav. in Anal. Cienc. Nat. vi. (1803) 339 = Cereus speciosissimus.
speciosus, Desf. Tabl. Hort. Par. 191 = Phyllocactus phyllanthoides.
speciosus, Gill. ex Sweet, Hort. Brit. ed. II. 235 = Mammillaria speciosa.
speciosus, Hook. Bot. Mag. t. 2820 = Rhipsalis Swartziana.
Spinii, Colla, Antol. vi. 501 = Mammillaria discolor.
spinosissimus, Martyn, Cat. Hort. Cant. 88 = Opuntia spinosissima.
spinosus, Gill. ex Loud. Hort. Brit. 194 = Mammillaria spinosa.
stellaris, Salm-Dyck, Hort. Dyck. ed. II. 7 = Mammillaria pusilla.
stellatus, Willd. Enum. Hort. Berol. Suppl. 30 = Mammillaria pusilla.
stramineus, Spreng. Syst. ii. 494 = Mammillaria flavescens.
strictus, Haw. Misc. 188 = Opuntia stricta.
strictus, Willd. Enum. Hort. Berol. Suppl. 32 = Cereus strictus.
subinermis, Link, ex Steud. Nom. ed. II. i. 246 = Opuntia coccinellifera.
subquadriflorus, Moç. & Sesse, ex DC. Prod. iii. 471 = Opuntia rosea.
subrepandus, Spreng. Syst. ii. 495 = Cereus subrepandus.
subsulcatus, Gill. ex Loud. Hort. Brit. 196 = Opuntia subsulcata.
sylvestris, Thiery, ex Steud. Nom. ed. I. 189 ; ed. II. i. 246 (quid ?).
tenuis, Schott, ex DC. Prod. iii. 469 = Cereus tenuis.
tetra, Vell. Fl. Flum. 207 ; v. t. 80 = Rhipsalis fasciculata.
tetragonus, Linn. Sp. Pl. 466 = Cereus tetragonus.
tomentosus, Link, Enum. Hort. Berol. ii. 24 = Opuntia tomentosa.
torquatus, Hort. Lugd. ex Walp. Rep. ii. 342 = Rhipsalis rhombea.
triacanthos, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia triacantha.
triangularis, Linn. Sp. Pl. 468 = Cereus compressus, trigonus.
triangularis, Vell. Fl. Flum. 206 ; v. t. 24 = Cereus triangularis.
trichotomus, Tenore, ex Steud. Nom. ed. II. i. 246 (quid ?).
triquetor, Haw. Misc. 189 = Cereus triqueter.
triquetor, Vell. Fl. Flum. 206 ; v. t. 25 = Cereus Arrabidae.
triqueter, Willd. Enum. Hort. Berol. Suppl. 33 = Cereus trigonus.
truncatus, Link, Enum. Hort. Berol. ii. 24 = Epiphyllum Altensteinii ?
tuberculatus, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia tuberculata.
Tuna, Linn. Sp. Pl. 468 = Opuntia Tuna.
Tuna-elatior, Roxb. Hort. Beng. [87] nomen.—Am. austr.
Tuna-major, Roxb. l. c.—Am. austr.
tunicatus, Lehm. Ind Sem. Hort. Hamb. (1827) 17 = Opuntia tunicata.
undulosus, Kostel. Allg. Med.-Pharm. Flora, iv. 1393 = Cereus variabilis.
Urumbeba, Vell. Fl. Flum. 207 ; v. t. 32 = Opuntia triacantha.
Urumbella, Steud. Nom. ed. II. i. 246 = prsec.
verticillatus, Brot. ex Heynh. Nom. fl. 103, nomen (quid ?).
viviparus, Nutt. Gen. Am. i. 295 = Mammillaria vivipara.
zinniaeflorus, Moç. & Sesse, ex DC. Prod. iii. 475 = Pereskia zinniaeflora.

CACUCIA, J. F. Gmel. Syst. 698 (1791) = Cacoucia, Aubl. (Combret.).

CACUVALLUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 399 = Muouna, Adans. (Legum.).
altissimum, Medic. l. c. 400 = M. urens.

CADABA, Forsk. Fl. Aegypt. Arab. 67 (1775). *CAPPARIDEAE*, Benth. & Hook. f. i. 108.
DENOCARPUS, Wall. Cat. n. 6373 (1832).
MACRONEMIUM, Burch. Trav. i. 388 (1825) ; ii. 615.
SCHEFFERIA, Neck. Elem. iii. 67 (1790).
STROMERIA, Roxb. Fl. Ind. ii. 78 (1832).
STROMESIA, Vahl, Symb. Bot. i. 19 (1790).

CADABA :—
capparoides, *DC. Prod.* i. 244.—Ins. Molucc.
dubia, DC. l. c. = farinosa.
farinosa, *Forsk. Fl. Aegypt. Arab.* 68.—Ind. bor. occ. ; Arab. ; Afr. trop.
glandulosa, *Forsk. l. c.*—Arabia.
glandulosa, Miq. in Verh. Nederl. Inst. III. v. (1852) 2 = Miqueliana.
heterotricha, *Stocks, in Hook. Ic. Pl.* t. 889.—Sindh.
indica, *Lam. Encyc.* i. 544.—Ind. or.
jonces, *Harv. Gen. S. Afr. Pl.* ed. II. 12.—Afr. austr.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 90.—Afr. trop.
longifolia, *DC. Prod.* i. 244.—Abyss.
madagascariensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 464.—Madag.
Miquelliana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 321.—Malaya.
monopetala, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1212 = glandulosa.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 8.—Natal.
rotundifolia, *Forsk. Fl. Aegypt. Arab.* 68.—Arab. ; Abyss.
somalensis, *Franch. in Revoil, Çomali*, 19.—Afr. trop.
trifoliata, *Wight & Arn. Prod.* 24.—Ind. or.
trinervia, DC. Prod. i. 244 = Cleome coluteoides.
triphylla, Wight, in Hook. Bot. Misc. iii. (1833) 296 = trifoliata.
virgata, *Boj. in Ann. Sc. Nat. Sér.* II. xx. (1843) 58.—Madag.

CADALVENA, Fenzl, in Sitz. Akad. Wien, Math. Nat. ii. Abth. ii. (1865) 139 = **Kaempferia**, Linn. (Scitamin.).
spectabilis, Fenzl, l. c. 140.—Afr. trop.

CADAMBA, Sonner. Voy. ii. 228. t. 128 (1782) = **Guettarda**, Linn. (Rubiac.).
jasminiflora, Sonner. l. c. ii. t. 128 ; iii. 253 = G. speciosa.

CADELARI, Adans. Fam. ii. 268 (1763) = **Pupalia**, Juss. (Amarantac.).
lappacea, Medic. Malv. 92 = P. atropurpurea.

CADELARIA, Rafin. Fl. Tellur. iii. 39 (1836) = **Achyranthes**, Linn. (Amarantac.).
indica, Rafin. l. c. = A. aspera.
punctata, Rafin. l. c. = A. aspera ?
sicula, Rafin. l. c. = A. aspera.

CADELIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 353 ; Phil. Bot. i. 206 (1789) = **Phaseolus**, Tourn. (Legumin.).
nigrum, Medic. l. c. 353 = P. Max.
viride, Medic. l. c. = P. Max.

CADELLIA, F. Muell. Fragm. ii. 25 (1860). *SIMARUBEAE*, Benth. & Hook. f. i. 313.
GUILFOYLIA, F. Muell. Fragm. viii. 33 (1873).
monostylis, *Benth. Fl. Austral.* i. 376.—Austral. peninsula, *F. Muell. Fragm.* ii. 25. t. 13.—Austral.

CADETIA, Gaudich. in Freyc. Voy. Bot. 422. t. 83 (1826) = **Dendrobium**, Sw. (Orchid.).
angustifolia, Blume, Rumphia, iv. 39 = D. angustifolium ?
biloba, Blume, Mus. Bot. Lugd. Bat. i. (1849) 30 = D. bilobum.
recurvata, Blume, l. c.—N. Guinea.
similis, Blume, Rumphia, iv. 39, in obs.—N. Guinea.
umbellata, Gaudich. in Freyc. Voy. Bot. 422. t. 83 = D. umbellatum.

CADIA, Forsk. Fl. Aegypt. Arab. 90 (1775). *LEGUMINOSAE*, Benth. & Hook. f. i. 560.
PANCIATICA, Pieete. Hort. Panciat. 9 (1788).
SPAENDONCEA, Desf. Dec. Phil. vii. 259 ; et in Usteri, N. Ann. xiii. (1796) 15.
anonala, *Vathe, in Linnaea*, xliii. (1880–82) 327.—Madag.
arabica, Raeusch. Nom. ed. III. 117 = purpurea.
Commersoniana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 370.—Madag.
Ellisiana, *Baker, in Journ. Linn. Soc.* xx. (1884) 185.—Madag.
pedicellata, *Baker, l. c.* 186.—Madag.
pubescens, *Boj. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 256.—Madag.
purpurea, *Forsk. Fl. Aegypt. Arab.* 90.—Arab. felix.
varia, L'Hérit. Dis. in Mag. Enc. v. 29 = purpurea.

CADISCUS, E. Mey. ex DC. Prod. vii. 254 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 408.
aquaticus, *E. Mey. l. c.* 255.—Afr. austr.

CADSURA, Spreng. Syst. ii. 642 (1827) = **Kadsura**, Juss. (Magnoliac.).

CAELESTINA, Cass. in Dict. Sc. Nat. vi. Suppl. 8. t. 93 (1817) = **Ageratum**, Linn. (Compos.).
ageratoides, H. B. & K. Nov. Gen. et Sp. iv. 151 = A. corymbosum.
albida, DC. Prod. v. 107 = A. albidum.
caerulea, Cass. in Dict. Sc. Nat. vi. Suppl. 8 = A. corymbosum.
cordata, Pohl, ex Baker, in Mart. Fl. Bras. vi. ii. 345 = Eupatorium macrophyllum.
corymbosa, Benth. Bot. Voy. Sulph. 111 = A. salicifolium.
corymbosa, DC. Prod. v. 108 = A. corymbosum.
Hartwegi, Walp. Rep. ii. 545 = Eupatorium Hartwegi.
hastata, Pohl, ex Baker, in Mart. Fl. Bras. vi. ii. 363 = Eupatorium betonicaeforme.
isocarphoides, DC. Prod. v. 107 = A. laocarphoides.
latifolia, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 71 = A. latifolium.
Lasiognaa, Klotzsch, ex Walp. Rep. ii. 545 = A. corymbosum.
linearifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 191 = Alomia polyphylla.
longifolia, Pohl, ex Baker, l. c. 361 = Eupatorium organense.
maritima, Torr. & Gray, Fl. N. Am. ii. 64 = A. littorale.
micrantha, Spreng. Syst. iii. 446 = A. coelestinum.
microcarpa, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 72 = A. microcarpum.
palaceus, J. Gay, ex DC. Prod. v. 107 = A. palaceum.
parvifolia, DC. l. c. 108.—Am. austr.
petiolata, Hook. & Arn. Bot. Beech. Voy. 138 = A. petiolatum.
Pohlii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 190 = Alomia Pohlii.
repanda, Pohl, ex Baker, l. c. 309 = Eupatorium viridiflorum.
repens, Sch. Bip. in Schomb. Fauna et Flora Guy. 1138.—Guiana.
scabriuscula, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 72 = A. scabriusculum.
suffruticosa, Sweet. Hort. Brit. ed. I. 229 = A. corymbosum.
tomentosa, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 71 = A. tomentosum.
urticaefolia, Pohl, ex Baker, in Mart. Fl. Bras. vi. ii. 369 = Eupatorium betonicaeforme.
viscosa, Pohl, ex Baker, l. c. = Eupatorium spiculatum.

CAELIA, G. Don, in Sweet, Hort. Brit. ed. III. 637 (1839) = **Coelia**, Lindl.

CAELOCLINE, Auct. ex Steud. Nom. ed. II. i. 247 (1840) = COELOCLINE, A. DC. = **Xylopia**, Linn. (Anonac.).

CAELODEPAS, Benth. & Hook.f. Gen. iii. 318 (1880) = **Coelodepas**, Hassk. (Euphorb.).

CAELOGLOSSUM, Steud. Nom. ed. II. i. 247 (1840) = COELOGLOSSUM, Lindl. = **Habenaria**, Willd. (Orchid.).

CAELOGYNE, Wall. ex Steud. Nom. ed. II. i. 247 (1840) = **Coelogyne**, Lindl. (Orchid.).

CAELOSPERMUM, Blume, Bijdr. 994 (1826), vide **Coelospermum**, Blume.

CAENOTUS, Rafin. Fl. Tellur. ii. 50 (1836) = **Erigeron**, Linn. (Compos.).
canadensis, Rafin. l. c. = E. canadensis.
longifolius, Rafin. l. c. = E. longifolius ?
pusillus, Rafin. l. c. = E. canadensis.

CAESALPINA, Plum. ex Linn. Gen. ed. I. 363 (1753) = seq.

CAESALPINIA, Linn. Sp. Pl. 380 (1753). *LEGUMINOSAE*, Benth. & Hook. f. i. 585.
BALSAMOCARPON, Cloa, in C. Gay, Fl. Chil. ii. 226. t. 20 (1846).
BIANCAEA, Todaro, Nuov. Gen. 21 (1858).
BONDUC, Adans. Fam. ii. 318 (1763).
CAPRETIA, Adans. l. c. (1763).
CINCLIDOCARPUS, Zoll. in Nat. en Genees. Arch. Neêrl. Indie, iii. (1846) 81.
CLADOTRICHIUM, Vog. in Linnaea, xi. (1837) 401.
COULTERIA, H. B. & K. Nov. Gen. et Sp. vi. 328. t. 568 (1823).
ERYTHROSTEMON, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. i. 67. t. 39 (1841).
GUILANDIA, P. Br. Hist. Jamaic. 228 (1756).
GUILANDINA, Linn. Gen. ed. I. 124 (1737).
LIBIDIBIA, Schlecht. in Linnaea, v. (1830) 192.
POMARIA, Cav. Ic. v. i. t. 409 (1799).
TARA, Molina, Saggio Chile, ed. II. 153 (1810).
TICANTO, Adans. Fam. ii. 319 (1763).
acanthobotrys, *Miq. Fl. Ind. Bat. Suppl.* i. 292.—Sumatra.
acinaciformis, *Mart. ex Steud. Nom.* ed. II. i. 247.—Bras.
affinis, *Hemsl. Diag. Pl. Nov.* 8.—Guatemala.
angulicaulis, *Clos, in C. Gay, Fl. Chil.* ii. 223.—Chili.
angustifolia, Salisb. Prod. 326 = C. Sappan.
arborea, Zoll. ex Miq. Fl. Ind. Bat. i. 112 = Peltophorum ferrugineum.
Aristæ, Mill. Gard. Dict. ed. VIII. n. 2, sphalm. = Crista.
armata, R. Grah. in Wall. Cat. n. 5340 = mimosoides.
axillaris, DC. Prod. ii. 481 = C. sepiaria ? et Mezoneurum cucullatum.
bahamensis, Lam. Encyc. i. 461 = Crista.
Barbon, *Bert. in 'Merc. Chil.* (1829)'; *et in Bull. Férussac*, xx. (1830) 109.—Chili.
Berteri, Spreng. ex DC. Prod. ii. 482 = bijuga.
bijuga, Sch. Obs. 166.—Ind. occ.
bijuga, Wall. Cat. n. 5338 = C. Nuga.
Bonduc, *Roxb. Hort. Beng.* 32; *Fl. Ind.* ii. 362.—Cosmop. trop.
Bonduc, Wight & Arn. Prod. 280 = C. Bonducella.
Bonducella, *Fleming, in As. Res.* xi. (1810) 159.—Cosmop. trop.
bracteosa, *Tul. in Arch. Mus. Par.* iv. (1844) 141.—Bras.
brasiliensis, *Linn. Sp. Pl.* 380.—Am. trop.
Brea, Gill. ex Steud. Nom. ed. II. i. 247 = praecox.
brevifolia, *Baill. Hist. Pl.* ii. 168.—Chili.
Cacalaco, *Humb. & Bonpl. Pl. Aeq.* ii. 178. t. 137.—Mexic.
calycina, *Benth. in Mart. Fl. Bras.* xv. ii. 71.—Am. trop.
cassioides, *Willd. Enum. Hort. Berol.* 444.—Am. austr.
Cavanillesii, Spreng. Syst. v. Ind. 128 = bijuga.
chinensis, Roxb. Hort. Beng. 32; Fl. Ind. ii. 361 = C. Nuga.
Cinclidocarpus, *Miq. Fl. Ind. Bat.* i. 110.—Ind. or.; Malaya.
coluteifolia, *Griseb. in Gott. Abh.* xxiv. (1879) 111.—Reg. Argent.
compressa, *D. Dietr. Syn. Pl.* ii. 1494.—Mexic.; Bras.
coriacea, Poit. in Rev. Hortic. Sér. II. ii. (1848–44) 269 = coriaria.
coriaria, *Willd. Sp. Pl.* ii. 532.—Am. austr.
corymbosa, *Benth. Pl. Hartw.* 117.—Guayaquil.
coulteriioides, *Griseb. in Gott. Abh.* xxiv. (1879) 113.—Reg. Argent.
Courbonilana, *Baill. Adansonia*, x. (1871) 108.—Mare rubrum.
Crista, *Linn. Sp. Pl.* 380.—Ind. occ.
Crista, Thunb. Fl. Jap. 179 = sepiaria.
cucullata, Roxb. Hort. Beng. 32; Fl. Ind. ii. 358 = Mezoneurum cucullatum.
dasyrachis, Miq. Fl. Ind. Bat. Suppl. i. 292 = Peltophorum dasyrachis.
desertorum, Mart. ex G. Don, Gen. Syst. ii. 481 = bracteosa.
digyna, *Rottl. in Ges. Naturf. Freunde Neue Schr.* iv. (1803) 200.—Ind. or.; Malaya.
dubia, Spreng. Syst. ii. 348 = Peltophorum Vogelianum.
Ebano, *Kunt. Pl. Columb.* ii. 87. t. 129.—N. Granat.
echinata, *Lam. Encyc.* i. 461.—Bras.
elata, Sw. Obs. 166 = Poinciana elata.
CAESALPINIA :—
enneaphylla, Roxb. Hort. Beng. 32; Fl. Ind. ii. 363 = Mezoneurum enneaphyllum.
eriostachys, *Benth. Bot. Voy. Sulph.* 88.—Am. centr.
exilifolia, *Griseb. in Gott. Abh.* xix. (1874) 128.—Reg. Argent.
excelsissima, *Moç. & Sesse, ex DC. Prod.* ii. 488.—Nicaragua.
feros, Hassk. Ind. Sem. Hort. Amst. (1841); Pl. Jav Rar. 400 = sepiaria.
feros, Hohen. ex Baker, in Hook. f. Fl. Brit. Ind ii. 261 = Wagatea spicata.
ferrea, *Mart. Reise Bras.* ii. 611.—Bras.
ferruginea, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 462 = Peltophorum ferrugineum.
fimbriata, *Tul. in Arch. Mus. Par.* iv. (1844) 145.—Bolivia.
Finlaysoniana, R. Grah. in Wall. Cat. n. 5971 = Peltophorum dasyrachis.
Fischeri, *Kegel & Koern. in Gartenfl.* (1858) 251.—Ind. or. ?
flavicans, R. Grah. in Wall. Cat. n. 5825 = digyna.
floribunda, *Tul. in Arch. Mus. Par.* iv. (1844) 140.—Bras.
furfuracea, Wall. Cat. n. 5835 = Mezoneurum glabrum Gardneriana, *Benth. in Mart. Fl. Bras.* xv. ii. 68.—Bras.
Gilliesii, *Wall. ex Hook. Bot. Misc.* i. (1830) 129.—Am. austr.
glabrata, *H. B. & K. Nov. Gen. et Sp.* vi. 326.—Peruv.
glandulosa, Sesse, ex DC. Prod. ii. 482 = bijuga.
Glenieii, *Thw. Enum. Pl. Zeyl.* 414 = Peltophorum ferrugineum.
gracilis, *Benth ex Hemsl. Diag. Pl. Nov.* 9.—Mexic.
gracilis, Miq. Fl. Ind. Bat. i. 110 = digyna.
grandis, *Heyne, in Wall. Cat.* n. 5830.—Ind. or.
Heyneana, Steud. Nom. ed. II. i. 247 = Wagatea spicata.
Hildebrandtii, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 871.—Afr. trop.
hispida, *D. Dietr. Syn. Pl.* ii. 1495.—Mexic.
horrida, Herb. Madr. ex Wall. Cat. n. 5836 = mimosoides.
horrida, A. Rich. Ess. Fl. Cub. 411 = sepiaria.
horrida, *Spreng. Syst. iv. Cur. Post.* 169.—N. Granat.
hymenocarpa, Wall. Cat. n. 5832 = Mezoneurum pubescens.
ignota, Blanco, Fl. Filip. ed. I. 336 = Mezoneurum pubescens.
inermis, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 367 = Peltophorum ferrugineum.
insignis, *Steud. Nom.* ed. I. 138; *D. Dietr. Syn. Pl.* ii. 1494.—Reg. Amazon.
japonica, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1845 ?) 117 = sepiaria.
kavalensis, *H. Mann, in Proc. Am. Acad.* vii. (1867) 164.—Ins. Hawaii.
Kilarae, Span. in Linnaea, xv. (1841) 200 = Peltophorum ferrugineum.
lacerans, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 367 = Pterolobium indicum.
languida, *Hofmgg. Verz. Pfl. Nachtr.* ii. 72.—Ind. or.
latifolia, Sallsb. Prod. 326 = vesicaria.
laxa, *Benth. Pl. Hartw.* 60.—Mexic.
laxiflora, *Tul. in Arch. Mus. Par.* iv. (1844) 143.—Bras.
lebbekhioides, DC. Prod. ii. 483 = Albizzia Millettii.
ligulata, Heyne, ex DC. Prod. ii. 484 = Pterolobium lacerans.
Machiottii, Zipp. ex Miq. Fl. Ind. Bat. i. 112 = Peltophorum ferrugineum.
macracantha, *Delile, Ind. Sem. Hort. Monsp.* (1838) 8.—Chili.
marginata, *Tul. in Arch. Mus. Par.* iv. (1844) 147.—Bras.
melanocarpa, *Griseb. in Gott. Abh.* xix. (1874) 128.—Reg. Argent.
Melanosticta, Spreng. Syst. iv. Cur. Post. 169 = Melanosticta Burchellii.
mexicana, *A. Gray, in Proc. Am. Acad.* v. (1861) 157.—Mexic.
microphylla, Buch.-Ham. in Wall. Cat. n. 5826 = C. Cinclidocarpa.
microphylla, *Mart. Reise Bras.* ii. 611.—Bras.
Millettii, *Hook. & Arn. Bot. Beech. Voy.* 163.—China.
mimosifolia, *Griseb. in Gott. Abh.* xix. (1874) 128.—Reg. Argent.

CAESALPINIA :—
mimosoides, Heyne, in Wall. Cat. n. 5837 = Wagatea spicata.
mimosoides, *Lam. Encyc.* i. 452.—Ind. or. ; Burma.
Minax, *Hance, in Journ. Bot.* xxii. (1884) 365.—China.
mollis, *Spreng. Syst.* iv. *Cur. Post.* 169.—Am. trop.
monosperma, Buch.-Ham. ex Wall. Cat. sub n. 5828 = Mezoneurum cucullatum.
monosperma, *Tul. in Arch. Mus. Par.* iv. (1844) 148.—Ind. occ.
mucronata, *Willd. Enum. Hort. Berol.* 444.—Bras.
nitida, *Hassk. Cat. Hort. Bog. Alt.* 235.—Malaya.
Nuga, *Alt. Hort. Kew.* ed. II. iii. 32.—As. trop. ; Austral. ; Polynes.
obliqua, Vog. in Linnæa, xi. (1837) 407 = echinata.
oleosperma, Roxb. Hort Beng. 32 ; Fl. Ind. ii. 357 = digyna.
Paipae, *Ruiz & Pav. Fl. Per.* iv. t. 375.—Peruv.
paniculata, Desf. Tabl. Hort. Par. ed. II. 210 = C. Nuga.
Parakyhda, Allem. in Trab. Soc. Vell. 56 = Schizolobium excelsum.
parvifolia, Steud. Nom. ed. II. i. 247 = C. Cinclidocarpa.
pauciflora, *Benth. & Hook. f. ex Chapm. Fl. S. U. St. Suppl.* 18.—Am. bor. ; Ind. occ.
pectinata, Cav. Desc. 467 = tinctoria.
pellucida, *Vog. in Linnæa*, x. (1836) 601.—Ins. S. Doming.
peltophoroides, *Benth. in Mart. Fl. Bras.* xv. ii. 72.—Bras.
pilosa, *Benth. l. c.* 73.—Bras.
pluviosa, *DC. Prod.* ii. 483.—Bras.
polyphylla, *Roxb. Hort. Beng.* 32, nomen.—Pegu.
porcina, *Mart. ex Steud. Nom.* ed. II. i. 247.—Bras.
praecox, *Ruiz & Pav. Fl. Per.* iv. t. 376.—Chili.
procera, *Poepp. ex Spreng. Syst.* ii. 343.—Cuba.
pubescens, Buch.-Ham. ex Wall. Cat. sub n. 5834 = sepiaria.
pulcherrima, *Sw. Obs.* 166.—Cosmop. trop.
Pumilio, *Griseb. in Goett. Abh.* xxiv. (1879) 113.—Reg. Argent.
punctata, *Willd. Enum. Hort. Berol.* 445.—Bras.
pyramidalis, *Tul. in Arch. Mus. Par.* iv. (1844) 139.—Bras.
regia, D. Dietr. Syn. Pl. ii. 1495 = Poinciana regia.
resupinata, Roxb. Hort. Beng. 32 ; Fl. Ind. ii. 262 = mimosoides.
Roxburghii, D. Dietr. Syn. Pl. ii. 1494 = Peltophorum ferrugineum.
rubicunda, *Benth. in Mart. Fl. Bras.* xv. ii. 73.—Bras.
Sappan, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art iv. 12 = C. Sappan.
Sappan, *Linn. Sp. Pl.* 381.—As. trop.
scandens, *Heyne, ex Roth, Nov. Pl. Sp.* 209.—Ind. or.
scandens, Koen. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 266 = Nuga.
Sellot, G. Don, Gen. Syst. ii. 481 = Peltophorum Vogelianum.
sepiaria, *Roxb. Hort. Beng.* 32 ; *Fl. Ind.* ii. 260.—As. trop.
sepiaria, Wall. Cat. n. 5834 H. = Mezoneurum ennesphyllum.
Sirmora, Buch.-Ham. ex Roxb. Hort. Beng. 32 ; Fl. Ind. ii. 259 = mimosoides.
smilacea, Hort. ex Steud. Nom. ed. II. i. 246 = Coccinlus carolinus.
spicata, Dals. in Hook. Kew Journ. iii. (1851) 89 = Wagatea spicata.
stipularis, *Benth. in Mart. Fl. Bras.* xv. ii. 72.—Bras.
suberosa, *Lodd. ex Loud. Hort. Brit.* 167.—Ind. or.
sumatrana, Roxb. Hort. Beng. 32 ; Fl. Ind. ii. 266 = Mezoneurum sumatranum.
sumatrana, Wall Cat. n. 5831 B, C = Nuga.
Tara, *Ruiz & Pav. Fl. Per.* iv. t. 374.—Peruv.
thomasa, Spreng. Syst. ii. 343 = coriaria.
tinctoria, *Domb. ex DC. Prod.* ii. 481.—N. Granat.
torquata, Blanco, Fl. Filip. ed. I. 336 = Mezoneurum glabrum.
tortuosa, *Roxb. Hort. Beng.* 32 ; *Fl. Ind.* ii. 265.—Burma ; Malaya ; Sumatra.
tortuosa, Wall. Cat. n. 5832 D = C. Cinclidocarpa.
trichocarpa, *Griseb. in Goett. Abh.* xxiv. (1879) 113.—Reg. Argent.

CAESALPINIA :—
ulosperma, *Hort. ex Steud. Nom.* ed. II. i. 248.—Hab. ?
ventricosa, Griseb. Fl. Brit. W. Ind. 205 = bijuga.
vernalis, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 77.—China.
vesicaria, Lam. Encyc. i. 461 = bijuga.
vesicaria, *Linn. Sp. Pl.* 381.—Bras.
vesicaria, Vell. Fl. Flum. 172 ; iv. t. 89 = echinata.
wyasadensis, Heyne, ex Wall. Cat. sub n. 5830 = grandis.

CAESAREA, Cambess. in Mém. Mus. Par. xviii. (1829) 378. t. 18 = **Viviania**, Cav. (Geraniac.).
albiflora, Cambess. l. c. 374.—Bras.
grandiflora, St. Hil. ex Steud. Nom. ed. II. i. 248.—Bras.
hypoleuca, Mart. ex Steud. l. c.—Bras.
lanceolata, Klotzsch, in Linnæa, x. (1836) 487.—Bras.
montevidensis, Klotzsch, l. c. 486.—Reg. Argent.
ovata, Klotzsch, l. c.—Bras.
petiolata, Klotzsch, l. c. 487.—Bras.
rubriflora, Cambess. in Mém. Mus. Par. xviii. (1829) 374.—Bras.

CAESIA, R. Br. Prod. 277 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 792.
acanthoclada, F. Muell. Fragm. i. 215 = Corynotheca acanthoclada.
africana, *Baker, in Trans. Linn. Soc.* xxix. (1875) 160.—Afr. trop.
alpina, Hook. f. Fl. Tasm. ii. 373 = Chlorophytum alpinum.
asparagoides, *Eckl. ex Steud. Nom.* ed. II. i. 248.—Afr. austr.
chlorantha, F. Muell. Fragm. i. 63 = vittata.
coarctata, Spreng. Syst. ii. 38 = Eccremis coarctata.
comosa, Spreng. l. c. = Chlorophytum comosum.
corymbosa, R. Br. Prod. 277 = Chamaescilla corymbosa.
cyanea, A. Cunn. ex Baker, in Journ. Linn. Soc. xv. (1876) 358 = vittata.
dichotoma, F. Muell. Fragm. i. 215 = Corynotheca dichotoma.
Dregeana, Kunth, Enum. Pl. iv. 611 = Anthericum Dregeanum.
Eckloniana, Schult. f. Syst. vii. 1691 = Anthericum scilliflorum.
Helleri, *Hueg. ex Heynh. Nom.* ii. 103, nomen.—Austral.
hirsuta, Lindl. Swan Riv. App. 57. n. 278 = Agrostocrinum stypandroides.
lateriflora, R. Br. Prod. 277 = Corynotheca lateriflora.
micrantha, Lindl. Swan Riv. App. 57 = parviflora.
occidentalis, R. Br. Prod. 277 = parviflora.
paradoxa, Endl. in Lehm. Pl. Preiss. ii. 34 = Chamaescilla spiralis ?
parviflora, *R. Br. Prod.* 277.—Austral.
physodes, Spreng. Syst. ii. 88 = Urginea physodes.
pusilla, Spreng. l. c. = Urginea pusilla.
rigidifolia, *F. Muell. Fragm.* x. 48.—Australia.
scabra, Baker, in Journ. Linn. Soc. xv. (1876) 359 = Agrostocrinum stypandroides.
settiera, *Baker, l. c.*—Austral.
spiralis, Endl. in Lehm. Pl. Preiss. ii. 34 = Chamaescilla spiralis.
Thunbergii, Schult. f. Syst. vii. 1692 = Anthericum contortum.
vernicolor, Lindl. Swan Riv. App. 57 = Chamaescilla corymbosa.
vittata, *R. Br. Prod.* 277.—Austral.

CAESIA, Vell. Fl. Flum. 107 (1825) ; ill. t. 23 (1827) = **Cormonema**, Reissek (Rhamnac.).
spinosa, Vell. l. c. = C. spinosum.

CAESULIA, Roxb. Pl. Corom. i. 64. t. 93 (1795). *COMPOSITAE*, Benth. & Hook. f. ii. 318.
axillaris, *Roxb. l. c.*—Ind. or.
lancifolia, *Sweet, Hort. Brit.* ed. II. 288 (quid ?).—Ind. or.
radicans, Willd. Sp. Pl. iii. 1797 = Cryphiospermum repens.
scandens, Poepp. ex DC. Prod. v. 564 = Wulffia scandens.

CAETA, Stend. Nom. ed. II. i. 248 (1840) = **Torenia**, Linn. (Scrophular.).

CAETOCAPNIA, Endl. Gen. 180 (1837) = COETOCAPNIA, Link & Otto = **Bravoa**, Lex.

CAFFEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 12, an **Coffea**, Linn.

CAHOTA, Karst. in Linnæa, xxviii. (1856) 448 = **Clusia**, Linn. (Guttifer.).
carachensis, Karst. l. c.—Venezuela.

CAIDBEJA, Forsk. Fl. Aegypt. Arab. 82 (1775) = **Forskohlea**, Linn. (Urticac.).
adhaerens, Forsk. l. c. = F. tenacissima.

CAILLIEA, Guillem. & Perr. Tent. Fl. Seneg. 239 (1833) = **Dichrostachys**, DC. (Legumin.).
dichrostachys, Guillem. & Perr. l. c. 240 = D. nutans.
macrostachya, Steud. in Flora, xxvi. (1843) 759 = Pentaclethra filamentosa.

CAINA, Panch. ex Baill. in Bull. Soc. Linn. Par. i. (1880) 264 = **Couthovia**, A. Gray (Loganiac.).
erratica, Panch ex Baill. l. c. = C. corynocarpa.

CAINITO, Plum. ex Adans. Fam. ii. 166 (1763) ; Tuss. Fl. Antill. iii. 41. t. 9 (1824) = **Chrysophyllum**, Linn. (Sapotac.).
pomiformum, Tuss. Fl. Antill. iii. 41. t. 9 = Chrysophyllum Cainito.

CAIOPHORA, Presl, Rel. Haenk. ii. 41. t. 56 (1831) = **Blumenbachia**, Schrad. (Loas.).
absinthifolia, Presl, l. c. 42 = B. coronata.
cardui folia, Presl, l. c. = B cardui folia.
cirsui folia, Presl, l. c. t. 56.—Peruv.
contorta, Presl, l. c. 42 = Loasa compressa.
coronata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 327 = B. coronata
Herberti x, G. Don, ex Lour. Encyc. Pl. Suppl. ii. 1438.—Hab. ?
lateritia, Benth. in Maund. & Hensl. Botanist, 111. t. 119.—Chili.
Pentlandii, G. Don, ex Loud. Encyc. Pl. Suppl. ii 1438.—Peruv.
scandens, Mey. & Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 389 = B. scandens.

CAJAN, Adans. Fam. ii. 326 (1763) = **Cajanus**, DC (Legumin.).
inodorum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 368 = Cajanus indicus.
scarabaeoides, Thou. in Dict. Sc. Nat. vi. 167 = Atylosia scarabaeoides.

CAJANUM, Rafin. Sylva Tellur. 25 (1838) = seq.
Thora, Rafin. l. c. = Cajanus indicus.

CAJANUS, DC. Cat. Hort. Monsp. 85 (1813). *LEGUMINOSAE*, Benth. & Hook. f. i. 541.
albicans, R. Grah. in Wall. Cat. n. 5582 = Atylosia albicans.
argenteus, Spreng. Syst. iii. 248 = Argyrolobium Linnaeanum.
bicolor, DC. Cat. Hort. Monsp. 85 = indicus.
candicans, Wall. Cat. n. 5576 = Rhynchosia avensis.
cinereus, *F. Muell. Census*, 41.—Austral.
confertiflorus, *F. Muell. Pl. Pitcairn*, 9.—Austral.
flavus, DC. Cat. Hort. Monsp. 85 = indicus.
glandulosus, Dals. & Gibs. Bomb. Fl. 78 = Atylosia mollis.
goensis, Dals. in Hook. Kew Journ. ii. (1850) 264 = Atylosia barbata.
grandiflolius, *F. Muell. Pl. Pitcairn*, 9.—Austral.
heterus, Spreng. Syst. iii. 248 = Glycine helvola.
indicus, *Spreng. l. c.*—Ind. or.
kulnensis, Dals. in Hook. Kew Journ. ii. (1850) 264 = Atylosia kulnensis.
linearius, R. Grah. in Wall. Cat. n. 5578 = Atylosia linearis.
lutens, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 260 = indicus ?
marmoratus, *F. Muell. Census*, 41.—Austral.
megalanthus, Spreng. ex DC. Prod. ii. 218 = Astragalus megalanthus.
niveus, R. Grah. in Wall. Cat. n. 5581 = Atylosia nivea.
quinquefolius, Blanco, Fl. Filip. ed. II. 417 = Desmodium Cephalotes.

CAJANUS:—
reticulatus, *F. Muell. Census*, 41.—Austral.
scarabaeoides, Thon. ex R. Grah. in Wall. Cat. n. 5580 = Atylosia scarabaeoides.
striatus, Boj. Hort. Maurit. 109 = Indicus.
suaveolens, R. Grah. in Wall. Cat. n. 5579 = Rhynchosia suaveolens.
sibilis, Blanco, Fl. Filip. ed. II. 417 = Atylosia mollis.
Wightianus, R. Grah. in Wall. Cat. n. 5583 = Atylosia albicans.
wolgaricus, Spreng. ex Steud. Nom. ed. II. i. 348 = Calophaca wolgarica.

CAJU-PUTI, Adans. Fam. ii. 84 (1763) = **Melaleuca**, Linn. (Myrtac).

CAKILE, Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 99.
aegyptia, Spreng. Syst ii. 852 = Didesmus aegyptiacus.
aegyptiaca, Willd. Sp. Pl. iii. 617 = maritima.
aequalis, L'Hérit. ex DC. Syst. ii. 630 = maritima.
americana, Nutt. Gen. Am. ii. 62 = maritima.
bipinnata, Spreng. Syst. ii. 852 = Didesmus bipunnatus.
clavata, Spreng. l. c. = Rapistrum clavatum.
crenata, Jord. Diagn. i. 346 = maritima.
cubrani, H. B. & K. Nov. Gen. et Sp. v. 75 = maritima. cyrenaica, *Spreng. Syst.* ii. 852.—Cyrenia.
domingensis, Tuss. Fl. Ant. i. 119 = maritima.
edentula, Hook. Fl. Bor. Am. i. 59 = maritima.
hispanica, L'Hérit. ex DC. Syst. ii. 619 = Brassica incana.
hispanica, Jord. Diagn. i. 345 = maritima.
latifolia, Poir. Encyc. Suppl. ii. 58 = maritima.
littoralis, Jord. Diagn. i. 345 = maritima.
maritima, *Scop. Fl. Carn.* ed. II. ii. 85.—Europ.; Afr. bor.; Am. bor.
monosperma, *Lange, Pl. Nov. Hisp.* fasc. i. & t. 7.—Hispan.
myagroides, Poir. Encyc. Suppl. ii. 58 = Erucaria aleppica.
perennis, L'Hérit. ex DC. Syst. ii. 631 = Rapistrum perenne.
perfoliata, L'Herit. ex DC. l. c. 575 = Coringia orientalis.
pinnatifida, Stokes, Bot. Mat. Med. iii. 484 = maritima.
rugosa, L'Hérit. ex DC. Syst. ii. 632 = Rapistrum rugosum.
Serapionis, Gaertn. Fruct. ii. 267 = maritima.
sinuatifolia, Stokes, Bot. Mat. Med. iii. 485 = maritima.
striata, L'Hérit. ex DC. Syst. ii. 659 = Didesmus tenuifolius.

CALACANTHA, T. Anders. ex Benth. & Hook. f. Gen. ii. 1088 (1876). *ACANTHACEAE*, Benth. & Hook. f. i. c.
Dalzelliana, *T. Anders. ex Benth. & Hook. f. l. c.*—Ind. or
grandiflorus, Radlk. in Sitzb. Math.-Phys. Acad. Muench. xiii. ' 1884 ' (1883) 279 = prac.

CALACINUM, Rafin. Fl. Tellur. iii. 83 (1838) = **Muehlenbeckia**, Meisn. (Polygon.).
adpressum, Rafin. l. c. 84 = M. adpresse.
australis, Rafin. l. c. = M. australis.

CALADENIA, R. Br. Prod. 322 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 512.
LEPTOCERAS, Lindl. Gen. et Sp. Orch. 415 (1840).
alata, R. Br. l. c. 324 = carnea.
alba, R. Br. l. c. 324 = carnea.
angustata, Hook. f. Fl. Tasm. ii. 30. t. 125 = carnea.
angustata, Lindl. Gen. et Sp. Orch. 430 = testacea.
antarctica, Reichb. f. Beitr. Syst. Pfl. 67 = Lyperanthus antarcticus.
aphylla, *Benth. Fl. Austral.* vii. 537.—Austral.
armeria, *Fitzg. Austral. Orch.* i. vii. t. 1.—Austral.
Barbarossa, *Reichb. f. Beitr. Syst. Pfl.* 64.—Austral.
barbata, Lindl. Gen. et Sp. Orch. 418 = deformis.
Behrii, Schlecht. in Linnaea, xx. (1847) 589 = Patersonii.
bifolia, *Hook. f. Fl. Nov. Zel.* i. 247.—N. Zel.
Brunonis, Reichb. Beitr. Syst. Pfl. 67 = Glossodia Brunonis.
caerulea, *R. Br. Prod.* 324.—Austral.
Cairnsiana, *F. Muell. Fragm.* vii. 31.—Austral.
carnea, *R. Br. Prod.* 324.—Austral.

CALADENIA:—
clavigera, *A. Cunn. ex Lindl. Gen. et Sp. Orch.* 432. —Austral.
concolor, *Fitzg. Austral. Orch.* i. vii. t. 1.—Austral.
congesta, *R. Br. Prod.* 324.—Austral.
cornuta, Reichb. f. Beitr. Syst. Pfl. 67 = Chiloglottis cornuta.
cucullata, *Fitzg. Austral. Orch.* i. ii. t. 4.—Austral.
cuneata, Reichb. f. Beitr. Syst. Pfl. 67 = Burnettia cuneata.
deformis, *R. Br. Prod.* 324.—Austral.
denticulata, Lindl. Swan Riv. App. 52 = filamentosa.
dilatata, R. Br. Prod. 325 = Patersonii.
dimorpha, *Fitzg. Austral. Orch.* i. t. 3.—Austral.
diphylla, Reichb. f. Beitr. Syst. Pfl. 67 = Chiloglottis diphylla.
discoidea, *Lindl. Swan Riv. App.* 52.—Austral.
Drummondii, *Benth. Fl. Australe.* vi. 383.—Austral.
elegans, Reichb. f. Beitr. Syst. Pfl. 67 = Lyperanthus elliptica.
elongata, Lindl. Swan Riv. App. 52 = latifolia.
emarginata, Reichb. f. Beitr. Syst. Pfl. 67 = Glossodia emarginata.
filamentosa, Lindl. Gen. et Sp. Orch. 421 = Patersoni. filamentosa, *R. Br. Prod.* 324.—Austral.
filifera, Lindl. Swan Riv. App. 52 = filamentosa.
fimbriata, *Reichb. f. Beitr. Syst. Pfl.* 65.—Austral.
flava, *R. Br. Prod.* 324.—Austral.
gemmata, *Lindl. Swan Riv. App.* 52.—Austral.
Gigas, *Reichb. f. in Linnaea*, xli. (1877) 56.—N. Caled.
gracilis, R. Br. Prod. 324 = testacea.
Gunnii, Reichb. f. Beitr. Syst. Pfl. 67 = Chiloglottis Gunnii.
hirta, *Lindl. Swan Riv. App.* 52.—Austral.
ixioides, *Lindl. l. c.*—Austral.
javanica, *Benn. ex Ridl. in H. O. Forbes, Nat. Wand. East. Archip.* 518.—Timor.
latifolia, *R. Br. Prod.* 324.—Austral.
leptoclada, *Fitzg. in Gard. Chron.* (1882) i. 462.— Austral.
lobata, *Fitzg. l. c.* 461.—Austral.
longicauda, Lindl. Swan Riv. App. 52. t. 8 = Patersoni.
Lyallii, *Hook. f. Fl. Nov. Zel.* i. 247.—N. Zel.
macrophylla, R. Br. Prod. 325 = Menziesii.
macrostylis, *Fitzg. in Gard. Chron.* (1882) i. 462.— Austral.
major, Reichb. f. Beitr. Syst. Pfl. 67 = Glossodia major.
marginata, Lindl. Swan Riv. App. 51 = latifolia.
Menziesii, *R. Br. Prod.* 325.—Austral.
minor, *Hook. f. Fl. Nov. Zel.* i. 247.—N. Zel.
minor, Reichb. f. Beitr. Syst. Pfl. 67 = Glossodia minor.
mollis, Endl. in Lehm. Pl. Preiss. ii. 8 = hirta.
mollis, Lindl. Swan Riv. App. 51 = latifolia.
multisetis, *Reichb. f. Beitr. Syst. Pfl.* 64.—Austral.
nana, Endl. in Lehm. Pl. Preiss ii. 7 = reptans.
nigricans, Reichb. f. Beitr. Syst. Pfl. 67 = Lyperanthus nigricans.
Nortoni, *F. Muell. Census*, 118.—Austral.
oblonga, Reichb. f. Beitr. Syst. Pfl. 67 = Cyrtostylis oblonga.
ochreata, Lindl. Swan Riv. App. 51 = latifolia.
pallida, Lindl. Gen. et Sp. Orch. 421 = Patersoni.
paniculata, *Fitzg. in Gard. Chron.* (1882) i. 461.— Austral.
Patersonii, *R. Br. Prod.* 324.—Austral.
pellita, Endl. in Lehm. Pl. Preiss. ii. 8 = gemmata.
plicata, *Fitzg. in Gard. Chron.* (1882) i. 461.— Austral.
Preissii, Endl. in Lehm. Pl. Preiss. ii. 7 = reptans.
pulchernima, F. Muell. Fragm. v. 93 = Patersonii.
remiformis, Reichb. f. Beitr. Syst. Pfl. 67 = Cyrtostylis remiformis.
reptans, *Lindl. Swan Riv. App.* 52.—Austral.
reticulata, *Fitzg. in Gard. Chron.* (1882) i. 462.— Austral.
Rost, *Benth. Fl. Australe.* vi. 383.—Austral.
rotundifolia, Reichb. f. Beitr. Syst. Pfl. 67 = Cyrtostylis rotundifolia.
saccharata, *Reichb. f. l. c.* 63.—Austral.
serieea, *Lindl. Swan Riv. App.* 52.—Austral.
serrata, *Reichb. f. Beitr. Syst. Pfl.* 67.—Austral.
suaveolens, *Reichb. f. l. c.*—Austral.
sulphurea, A. Cunn. in Field, N. S. Wales, 361 = suaveolens.

CALADENIA:—
tentaculata, Schlecht. in Linnaea, xx. (1847) 571 = Patersonii.
tenuis, *Fitzg. in Gard. Chron.* (1882) i. 462.— Austral.
tessellata, *Fitzg. Austral. Orch.* i. ii. t. 3.—Austral.
testacea, *R. Br. Prod.* 324.—Austral.
unguiculata, Lindl. Swan Riv. App. 51 = deformis.
unita, *Fitzg. in Gard. Chron.* (1882) i. 461.— Austral.
variegata, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 248.—N. Zel.

CALADIUM, Vent. in Roem. Arch. ii. (1800) 847. *AROIDEAE*, Benth. & Hook. f. iii. 976.
AROSMA, Rafin. Fl. Tellur. iii. 66 (1836).
CYRTOSPADIX, C. Koch, Ind. Sem. Hort. Berol. (1858).
acre, R. Br. Prod. 336 = Colocasia antiquorum.
aculeatum, F. G. Dietr. Vollst. Lexik. Gaertn. i. 616 = Xanthosoma belophyllum.
alho-punctatissimum, J. Makoy, ex Karst. in C. Koch & Fint. Wochenschr. (1858) 331 = bicolor ?
amabile, *Versch. Cat. n.* 70 (1862) 2.—Bras.
amoenum, Hort. ex Engl. in DC. Monog. Phan. ii 461 = bicolor.
appendiculatum, Hort. ex Engl. l. c. 476 = Xanthosoma atrovirens.
Appianianum, Hort. ex Engl. l. c. 466 = picturatum.
arborescens, *Vent. Yard. Cels*, sub t. 30; *Willd. Sp Pl.* iv. 490.—Ind. or.
arboreum, *H. B. & K. Nov. Gen. et Sp.* i. 80.— N. Granat.
argyrites, Lem. in Illustr. Hortic. (1858) t. 185. f. 3 Misc. 57 = Humboldtii.
argyrospilum, Lem. l. c. 59 = bicolor.
argyrostigma, *Schott, Melet.* i. 16, nomen.—Am trop.
asperulum, Schott, in Oestr. Bot. Wochenbl. (1858) 313 = smaragdinum.
auritum, Vent. Jard. Cels, sub t. 30; Willd. Sp. Pl. iv. 491 = Syngonium auritum.
Baraquini, Hériucq, in Fl. des Serres, Sér. II. iii. (1858) 165 = bicolor.
Barilletii, *Hort.—Cf. Gard. Chron.* (1867) 461.— Hab. ?
Bauerii, *Reichb. Hort. Bot.* t. 160.—Bras.
Belleyenei, Hort. Chantin, ex Lem. in Illustr. Hortic. (1860) t. 324 = picturatum.
belophyllum, Willd. Enum. Hort. Berol. 987 = Xanthosoma belophyllum.
bicolor, *Vent. Yard. Cels*, t. 30.—Am. austr.
Bongniartii, C. Koch & Fint. Wochenschr. (1858) 287, sphalm. = bicolor.
Bowei, Steud. Nom. ed. II. i. 249 = Blurium Bowei.
brasilianum, Desf. Cat. Hort. Par. ed. III. 7, 386 = Xanthosoma hastifolium.
Brongniartii, Lem. in Illustr. Hortic. (1858) Misc. 58 = bicolor.
caracasanum, Engl. in DC. Monog. Phan. ii. 666, Index = Xanthosoma belophyllum.
Chantinii, Lem. in Illustr. Hortic. (1858) Misc. 58 = bicolor.
Colocasia, Schott, ex Wight, Ic. t. 787 = Alocasia cucullata.
colocasioides, Bronga. in Nouv. Ann. Mus. Par. iii. 156 = Colocasia antiquorum.
concolor, *C. Koch, in Beri. Allg. Gartenz.* (1857) 135. —Venezuela.
Consentartii, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
cordatum, Hort. Versch. ex Engl. l. c. 456 = Schomburgkii.
crassipes, Hort. ex Engl. l. c. 369 = Philodendron cannaefolium.
cucullatum, Pers. Syn. ii. 575 = Alocasia cucullata.
cupreum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1854) 6 = Alocasia cuprea.
Curmudalii, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
Devestianum, Lem. in Illustr. Hortic. (1862) t. 222. f. 1 = bicolor.
digitatum, *Blanco, Fl. Filip.* ed. II. 490.—Ins. Philipp.
discolor, Hort. ex Engl. in DC. Monog. Phan. ii. 462 = bicolor.
Duchartrei, Hort. ex Engl. l. c. = bicolor.

CALADIUM:—
Eckhartii, Lem. ex Engl. in DC. Monog. Phan. ii. 453 = bicolor.
edule, G. F. W. Mey. Prim. Fl. Essequ. 272 = Xanthosoma edule.
Engelii, *Karst. in C. Koch & Fint. Wochenschr.* (1858) 330.—Venezuela.
Enbeanum, C. Koch, Wochenschr. (1861) 267 = bicolor.
erythropus, Mart. ex Engl. in Mart. Fl. Bras. iii. 11. 172 = Adelonema erythropus.
esculentum, Vent. Jard. Cels, sub t. 80; et in Roem. Arch. ii. 111. 851 = Colocasia antiquorum.
firmulum, Schott, in Oestr. Bot. Wochenbl. (1854) 417 = bicolor.
fragrantissimum, Hook. Bot. Mag. t. 3314 = Philodendron fragrantissimum.
Gaerdtii, C. Koch & Fint. Wochenschr. (1858) 288 = bicolor.
giganteum, Blume, Cat. Gew. Buitenz. 108 = Colocasia indica.
glycorrhizum, Fraser, in Hook. Bot. Misc. i. (1830) 329 = Alocasia macrorrhiza.
grandifolium, Sims, Bot. Mag. t. 2648 = Philodendron Simsii.
grandifolium, Willd. Sp. Pl. iv. 490 = Philodendron grandifolium.
grisea argentea, Hort. ex Engl. in DC. Monog. Phan. ii. 468 = bicolor.
guttiferum, Poepp. ex Kunth, Enum. Pl. iii. 52 = Philodendron guttiferum.
Haageanum, C. Koch, Wochenschr. (1862) 136 = bicolor.
haematostigma, Kunth, Enum. Pl. iii. 42 = bicolor.
hastatum, Lem. in Illustr. Hortic. (1858) Misc. 61 = picturatum.
hastatum, Willd. ex Engl. in DC. Monog. Phan. ii. 279 = Urospatha Poeppigiana.
hastifolium, Steud. Nom. ed. II. i. 249 = Xanthosoma hastifolium.
helleborifolium, Vent. Jard. Cels, sub t. 80; et in Roem. Arch. ii. 111. 851 = Xanthosoma helleborifolium.
Hendersoni, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
heterophyllum, *Presl, Reliq. Haenk.* ii. 148.—Ins. Luzon.
Houlyanum, Hort. ex Engl. in DC. Monog. Phan. ii. 464 = bicolor.
Houlletii, Lem. in Illustr. Hortic. (1858) 59 = bicolor.
Humboldtii, *Schott, in Oestr. Bot. Wochenbl.* (1854) 417.—Bras.
Imbe, Hort. ex Kunth, Enum. Pl. iii. 50 = Philodendron Imbe.
indica, Hort. ex C. Koch, Berl. Allg. Gartenz. (1857) 136 = Alocasia indica.
Ketteleri, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
Kochii, Hort. Versch. ex C. Koch, Wochenschr. v. (1862) 135 = bicolor.
Kramerianum, Hort. ex Engl. in DC. Monog. Phan. ii. 459 = bicolor.
lacerum, Willd. Sp. Pl. iv. 491 = Philodendron lacerum.
Lauchaeanum, C. Koch, Wochenschr. (1861) 266 = bicolor.
Lemairenanum, Barraq. in Illustr. Hortic. (1862) t. 311 = picturatum.
Leopoldii, Hort. Versch. ex Engl. in DC. Monog. Phan. ii. 460 = bicolor.
Lindeni, Hort. ex Engl. l. c. 462 = bicolor.
lividum, *Neee, in Nouvied, Reise nach Brasil.* ii. 238.—Bras.
lividum, Lodd. Bot. Cab. t. 1590 = Staurostigma concinnum.
Louvi, Lem. in Illustr. Hortic. (1863) t. 360 = Alocasia Lowii.
luridum, Kunth, Enum. Pl. iii. 50 = Staurostigma concinnum.
macrophyllum, Lem. in Illustr. Hortic. (1863) t. 816, 822 = bicolor.
macropus, Hort. Petrop. ex Kunth, Enum. Pl. iii. 47 = Philodendron consanguineum.
macrorrhizon, R. Br. Prod. 336 = Alocasia macrorrhiza.
macrotiza, *Schott, in Oestr. Bot. Wochenbl.* (1858) 20.—Bras.
maculatum, Lodd. Bot. Cab. t. 608 = Dieffenbachia Seguina.

CALADIUM:—
Mafaffa, Hort. ex Engl. in DC. Monog. Phan. ii. 471 = Xanthosoma Mafaffa.
marantaefolium, Zipp. ex Kunth, Enum. Pl. iii. 55 = Aglaonema marantifolium.
marginatum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1854) 6 = bicolor.
marmoratum, *L. Math. ex C. Koch, l. c.*—Ecuador.
Martenstigianum, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
metallicum, Hamburg. Gart. Blumenzeit. (1858) 317 = Alocasia cuprea.
metallicum, Hort. ex Engl l. c. 502 = Alocasia indica.
mirabile, Lem. Illustr. Hortic. (1862) t. 854 = bicolor.
Mooreanum, Hort. ex Engl. in DC. Monog. Phan. ii. 462 = bicolor.
myrmecophyllum, C. Koch, Wochenschr. v. (1862) 135 = Humboldtii.
Newmanni, Lem. in Illustr. Hortic. (1858) 58 = bicolor.
nymphaefolium, Vent. Jard. Cels, sub t. 80; Willd. Sp. Pl. iv. 488 = Colocasia antiquorum.
odoratissimum, Hort. ex C. Koch, Berl. Allg. Gartenz. (1857) 30 = Alocasia odora.
odoratum, Lodd. Bot. Cab. t. 416 = Alocasia macrorrhiza.
odorum, Lindl. Bot Reg. t 641 = Alocasia macrorrhiza.
Ottonis, Hort. ex Engl. in DC. Monog. Phan. ii. 464 = bicolor.
ovatum, Vent. Jard. Cels, sub t. 80; et in Roem. Arch. ii. 111. 851 = Lagenantha toxicaria.
pallidinervium, Hort. ex Engl. in DC. Monog. Phan. ii. 468 = bicolor.
pallidum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1856) 6 = smaragdinum.
pedatum, Hook. Exot. Fl. t. 206 = Philodendron lacinistum.
pellucens, DC. in Pl. Jard. Genèv. (1826) 88 = bicolor.
pendulinum, *Manso, Enum.* 21.—Bras.
pendulum, *Blume, Cat. Gew. Buitenz.* 108.—Malaya.
Perrotteti, Lem. in Illustr. Hortic. (1858) 59 = bicolor.
petiolatum, Hook. Bot. Mag. t. 3428 = Anchomanes difformis.
pictum, DC. 4ème Note, Pl. Jard. Genèv. 88 = bicolor.
pictum, Lodd. Bot. Cab. t. 908 = Dieffenbachia picta.
picturatum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1854) 6.—Am. austr.
pinnatifidum, Willd. Sp. Pl. iv. 487 = Philodendron pinnatifidum.
platynervium, Hort. ex Kunth, Enum. Pl. iii. 47 = Philodendron crassinervium.
platypeevron, *Salm-Dyck, Hort. Dyck.* (1834) 334.—Bras.
plumbeum, C. Koch, Berl. Allg. Gartenz. (1857) 136 = Alocasia cuprea.
poecile, Schott, Melet. i. 18 = bicolor.
porphyroneuron, C. Koch, Wochenschr. v. (1862) 135 = picturatum.
Princeps, Hort. ex Kunth, Enum. Pl. iii. 55 = Aglaonema simplex.
pumilum, D. Don, Prod. Fl. Nep. 21 = Gonatanthus pumilus.
punctatissimum, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
punctatum, *Desf. Tabl. Hort. Par.* ed. II. 6, nomen.—Hab.?
Purdieanum, Schott, in Oestr. Bot. Zeitschr. (1859) 38 = bicolor.
pusillum, C. Koch, in Fl. des Serres, Sér. II. iii. (1858) 110 = bicolor.
regale, Lem. in Illustr. Hortic. (1863) t. 816 = bicolor.
Reichenbachianum, Stange, ex Engl. in Mart. Fl. Bras. iii. 33 = bicolor.
reticulatum, *Desf. Tabl. Hort. Par.* ed. II. 6, nomen.—Hab.?
Rougieri, [*Veresch.*].—*Cf. Gard. Chron.* (1864) 414.—Bras.
rubellum, C. Koch & Fint. Wochenschr. (1858) 288 = bicolor.
rubricaule, Hort. ex Lem. in Fl. des Serres, Sér. II. iii. (1858) 109 = bicolor.
rubroveninm, Hort. ex Engl. in DC. Monog. Phan. ii. 460 = bicolor.
rugosum, Desf. Cat. Hort. Par. ed. III. 386 = Alocasia cucullata.
sagittaefolium, Sieber, ex Engl. in DC. Monog. Phan. ii. 464 = bicolor.

CALADIUM:—
sagittaefolium, Vent. Jard. Cels, sub t. 80; et in Roem. Arch. ii. 111. 851 = Xanthosoma sagittifolium.
sarmentosum, Hort. ex Kunth, Enum. Pl. iii. 36 = Gonatanthus sarmentosus.
scandens, Willd. Sp. Pl. iv. 489 = Culcasia scandens.
Schmidtii, Lem. Illustr. Hortic. (1861) t. 297. f. 3 = Schomburgkii.
Schoolleri, Lem. l. c. f. 2 = Schomburgkii.
Schomburgkii, *Schott, in Oestr. Bot. Wochenschr* (1858) 122.—Guiana.
seguinum, Vent. Jard. Cels, sub t. 80; Willd. Sp. Pl. iv. 490 = Dieffenbachia Seguine.
Seiboldii, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
simplex, Blume, Cat. Gew. Buitenz. 108 = Aglaonema simplex.
Simsii, Hook. Bot. Mag. t. 3845 = Philodendron Simsii.
smaragdinum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1853) 13.—Am. trop.
sororium, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 88.—Bras.
splendens, C. Koch & Fint. Wochenschr. (1858) 288 = bicolor.
splendidum, *Versch. Cat.* n. 70 (1862) 4.—Bras.
Spruceanum, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 38 = bicolor.
Stangeanum, C. Koch, Wochenschr. (1861) 136 = bicolor.
steudnerifolium, *Engl. Bot. Jahrb.* vi. (1885) 284.—N. Granat.
strictipes, *Schott, Syn. Aroid.* 51.—Bras.
subrotundum, Lem. in Illustr. Hortic. (1858) Misc. 60 = Schomburgkii.
surinamense, Miq. Hort. Amstelod. (1858) = bicolor. Thelemanni, *Versch. Cat.* n. 72 (1868) 2.—Hab.?
thripedeum, Lem. in Illustr. Hortic. v. (1858) Misc. 59 = marmoratum.
trifoliatum, Desf. Cat. Hort. Par. ed. III. 386 = acq.
tripartitum, Willd. Sp. Pl. iv. 491 = Philodendron tripartitum.
Troubetskoyi, Hérincq, in Fl. des Serres, Sér. II. iii. (1858) 164 = picturatum.
undulatum, Steud. Nom. ed. II. ii. 219 = Peltandra undulata.
utile, Hort. ex Engl. in DC. Monog. Phan. ii. 469 = Xanthosoma sagittifolium.
variegatum, Desf. Cat. Hort. Par. ed. III. 7, 385 = Xanthosoma helleborifolium.
Veitchii, Henders. ex Hook. f. Bot. Mag. t. 5876 = Alocasia Lowii.
Veitchii, Lindl. in Gard. Chron. (1859) 740 = Alocasia cuprea.
Vellosianum, Schott, in Bonplandia, vii. (1859) 163 = bicolor.
Verschaffeltii, Lem. in Illustr. Hortic. (1858) 59. t. 185 = bicolor.
violaceum, Desf. Cat. Hort. Par. ed. III. 7 = acq. †
violaceum, Hort. ex Engl. in DC. Monog. Phan. ii. 492 = Colocasia antiquorum.
virginicum, Hook. Exot. Fl. t. 182 = Peltandra undulata.
viviparum, Nees, in Schott, Melet. i. 18; Lodd. Bot. Cab. t. 981 = Remusatia vivipara.
Wagneri, Hort. ex Engl. in DC. Monog. Phan. ii. 464 = bicolor.
Wightii, Lem. Illustr. Hortic. (1861) 294 = bicolor.
xanthorrhizon, Boos, Cat. 846 = Xanthosoma jacquini.
xanthorrhizum, Willd. Sp. Pl. iv. 490 = Xanthosoma sagittifolium.
zamiaefolium, Lodd. Bot. Cab. t. 1408 = Zamioculcas Loddigesii.

CALAENA, Schlecht. in Linnaea, ii. (1827) 528, sphalm. = Calaena, R. Br. (Orchid.).

CALAIS, DC. Prod. vii. 85 (1839) = **Microcseris**, D. Don (Compos.).
aphanocarpha, A. Gray, in Proc. Am. Acad. vi. (1864–65) 559 = M. aphanocarpha.
Bigelovii, A. Gray, Pacif. Rail. Rep. iv. 113 = M. Bigelovii.
Bolanderi, A. Gray, in Proc. Am. Acad. vii. (1868) 355 = M. Bolanderi.
cyclocarpha, A. Gray, in Pacif. Rail. Rep. iv. 113 = M. Douglasii.

CALAIS :—
Douglasii, DC. Prod. vii. 85 = Microseris Douglasii.
eriocarpha, A. Gray, in Proc. Am. Acad. vi. (1864–65) 559 = M. Douglasii.
glauca, A. Gray, l. c. vii. (1868) 364 = M. procera.
gracilisloba, Kellogg, in Proc. Calif. Acad. v. (1873) 48 = M. major.
laciniata, A. Gray, in Pacif. Rail. Rep. iv. 113 = M. laciniata.
Lindleyi, DC. Prod. vii. 85 = M. Lindleyi.
linearifolia, DC. l. c. = M. linearifolia.
macrochaeta, A. Gray, Pl. Fendl. 112 = M. macrochaeta.
major, A. Gray, in Pacif. Rail. Rep. iv. 113 = M. major.
nutans, A. Gray, l c. = M. nutans.
Parryi, A. Gray, l. c. 112 = M. Parryi.
platycarpha, A. Gray, l. c. 113 = M. platycarpha.
sylvatica, A. Gray, l. c. 114 = M. sylvatica.
tenella, A. Gray, l. c. = M. aphantocarpha.

CALAMAGROSTIS, Adans. Fam. ii. 31 (1763).
GRAMINEAE, Benth. & Hook. f. iii. 1150.
AMAGRIS, Rafin. Princ Somiol. 27 (1814); et in Am. Monthly Mag. (1819) 100.
ATHERNOTUS, Dulac, Fl. Hautes-Pyr. 74 (1867).
LECHLERA, Steud. Syn. Pl. Gram. 101, in nota (1855).
PTERODIPHUM, Willd. ex Steud. Nom. ed. II. ii. 414 (1841).
abortiva, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 155 = Deyeuxia acutiflora.
scrathera, *Peterm. in Flora*, xxvii. (1844) 225.—Saxonia.
acutiflora, DC Fl. Fr. v. 255 = Deyeuxia acutiflora.
aemula, Steud. Nom. ed. II. i. 249; Syn. Pl. Gram. 192 = Deyeuxia Forsteri.
agapates, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
agrostoides, *Boiss. Fl. Orient.* v. 522.—Reg. Caucas.
agrostoides, Spreng. Syst. i. 252 = Deyeuxia canadensis.
airoides, Steud. Nom. ed. II. i. 249 = Arundo airoides.
alba, *Steud. l. c.*—Peruv.
albicans, Buckl. in Proc. Acad. Sc. Philad. '1862' (1863) 92 = Deyeuxia aleutica.
aleutica, Bong. Veg. Ins. Sitcha, 171 = Deyeuxia aleutica.
alpina, Hoet, Gram. iv. t. 51 = tenella.
altissima, Host, Fl. Austr. i. 101 = Arundo mauritanica.
ambigua, *Hartm. Skand. Fl.* ed. V.—Scandin.
andina, *Nutt. in Journ. Acad. Sc. Philad.* N. S. i. (1847) 187.—Calif.
anomala, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
anthoxanthoides, *Regel*, in *Act. Hort. Petrop.* vii. (1880) 640.—Turkestan.
Antoniana, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
arenaria, Roth, Tent. Fl. Germ. i. 34 = Ammophila arundinacea.
argentea, DC. Fl. Fr. iii. 25 = Stipa Calamagrostis.
arundinacea, Roth, Tent. Fl. Germ. i. 33 = Deyeuxia sylvatica.
arundinacea, Wibel, Prim. Fl. Werth. 122 = Deyeuxia acutiflora.
Arundo, J. F. Gmel. Syst. 171 = Stipa Calamagrostis.
Arundo, Roth, Tent. Fl. Germ. i. 38 = Aira caespitosa.
australis, *Buese, in Mig. Pl. Yungh.* 843.—Malaya.
aurea, Steud. Syn. Pl. Gram. 189 = Agrostis pallida.
Balansae, *Boiss. Fl. Orient.* v. 522.—Fontus.
baltica, Trin. Gram. Unifi. 227; Hartm. Skand. Fl. ed. II. 50 = Ammophila baltica.
Beyrichiana, *Ness, ex Doell, in Mart. Fl. Bras.* ii. III. 53.—Bras.
bicolor, J. F. Gmel. Syst. i. 172 = Ampelodesma bicolor.
Billardieri, Steud. Nom. ed. II. i. 249 = Deyeuxia Billardieri.
Bolanderi, Thurb. in S. Wats. Bot. Calif. ii. 280 = Deyeuxia Bolanderi.
borealis, Laest. Bidr. Vaextl. Torn. Lappm. Upa. 44 = Deyeuxia neglecta.
brachyrrhiza, *Steud. Syn. Pl. Gram.* 189.—Japon.
brevifolia, Steud. Nom. ed. II. i. 249 = Deyeuxia brevifolia.
brevifolia, Beck, Bot. N. & Mid. St. 401 = Ammophila brevipilis.

CALAMAGROSTIS :—
Breweri, Thurb. ex Boland. in Trans. Calif. Agr. Soc. (1864–65) 186 = Deyeuxia Breweri.
bromoides, Nutt. Gen. Am. i. 44 = Stipa Aristella.
caespitosa, Steud. Syn. Pl. Gram. 189 = Agrostis pallida.
canadensis, Beauv. Agrost. 157, nomen = Deyeuxia canadensis.
canadensis, Nutt. Gen. Am. i. 46 = Deyeuxia Nuttaliana.
cenenses, Beauv. ex Steud. Nom. ed. II. i. 249, 250 = Epigeios.
capillaris, *Nees, ex Steud. Nom.* ed. II. i. 219; *Syn. Pl. Gram.* 188.—Bras.
caucasica, Trin. in C. A. Mey. Verz. Pfl. Cauc. 15 = Deyeuxia acutiflora.
chalybaea, *Fries, Summa Pl.* 241.—Norveg.
Chamissonis, Steud. Nom. ed. II. i. 250 = Deyeuxia Chamissonis.
chilensis, *Phil. in Linnaea*, xxix. (1857–58) 89.—Chili.
chrysantha, Steud. Nom. ed. II. i. 250 = Deyeuxia chrysantha.
ciliata, *Steud. Syn. Pl. Gram.* 193.—Nepal.
cinnoides, Spreng. Syst. i. 252 = Deyeuxia Nuttalliana.
Clarionis, Loisel. Fl. Gall. 719 = Deyeuxia Halleriana.
coarctata, Steud. Nom. ed. II. i. 250 = Deyeuxia coarctata.
coarctata, Torr. Fl. N. York, ii. 444 = Deyeuxia Nuttalliana.
colorata, Nutt. Gen. Am. i. 46 = Phalaris intermedia.
colorata, Sibth. Fl. Oxon. 27 = Phalaris arundinacea.
columbiensis, Nutt. ex A. Gray, in Proc. Acad. Sc. Philad. '1862' (1863) 334 = Deyeuxia Langsdorfii.
compressa, *Doell, in Mart. Fl. Bras.* ii. III. 58.—Bras.
confinis, *Nutt. Gen. Am.* i. 47.—Am. bor.
conspicua, Bergeret, Fl. Pyr. i. 60 = Stipa Calamagrostis.
conspicua, J. F. Gmel. Syst. 172 = Arundo conspicua.
crassiglumis, Thurb. ex S. Wats. in Bot. Calif. ii. 281 = Deyeuxia crassiglumis.
centrorum, Dum. Obs. Gram. Belg. 126 = stricta ?
cuspidata, *Spreng. Syst.* i. 253.—Mexic.
densiflora, C. Muell. in Walp. Ann. vi. 986 = phragmitoides.
densiflora, Steud. Nom. ed. II. i. 250 = Deyeuxia densiflora.
deschampsioides, Trin. Ic. Gram. ibi. t. 354 = Deyeuxia deschampsioides.
dioica, *Steud. Syn. Pl. Gram.* 193.—Cochinch.
donacyformis, Loisel. Fl. Gall. i. 58 = Arundo Phinlana.
dubia, *Bunge, Lehm. Rel.* (1847) 548.—As. centr.
effusa, Steud. Nom. ed. II. i. 250 = Deyeuxia effusa.
elata, Blytt, Norges Fl. i. 92 = phragmitoides.
eminens, *Steud. Nom.* ed. II. i. 250 = Deyeuxia eminens.
emodensis, *Griseb. in Gottt. Nachr.* (1868) 80.—Reg. Himal.
epigeios, Huds. Fl. Angl. 48; ed. II. 54 = lanceolata.
epigeios, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1842) 329 = laxa.
epigeios, *Roth, Tent. Fl. Germ.* i. 34.—Europ.; As. temp.
epigeios, Steud. Nom. ed. II. i. 250, sphalm. = Saccharum cylindricum.
erianthe, Steud. Syn. Pl. Gram. 190 = Deyeuxia erianthe.
Falklandiae, *Steud. l. c.* 192.—Ins. Falkland.
farcta, *Y. F. Gmel. Syst.* 173.—Guiana.
filiformis, *Steud. in Gottt. Nachr.* (1868) 79.—Ind. or.
flexuosa, Rupr. Beitr. Pfl. Russl. iv. 34 = phragmitoides.
Forsteri, Steud. Nom. ed. II. i. 250 = Deyeuxia Forsteri.
Funckii, *Steud. Syn. Pl. Gram.* 192.—Venezuela.
fusca, Steud. Nom. ed. II. i. 250 = Deyeuxia fusca.
Gaudiniana, Reichb. Fl. Germ. Exc. 27 = lanceolata.
georgica, C. Koch, in Linnaea, xxi. (1848) 587 = epigeios.
glauca, Reichb. Exsm. Talassch. 14 = laxa.
glauca, Reichb. Fl. Germ. Excurs. 27 = epigeios.
glauca, Steud. Nom. ed. II. i. 250 = Deyeuxia Nuttalliana.
glauca, Steud. Syn. Pl. Gram. 187 = littorea.
glaucescens, C. Muell. in Walp. Ann. vi. 986 = phragmitoides.
glomerata, Boiss. & Buhse, in Mém. Soc. Nat. Mosc. xii. (1860) 229 = epigeios.

CALAMAGROSTIS :—
graniescens, *Blytt, Norges Fl.* i. 88.—Norveg.
gracilis, Seemus, Reise Isl. 78 = Stipa Aristella.
gracilis, Schumach. ex Nym. Consp. 800 = lanceolata groenlandica, *Kunth, Rev. Gram.* i. 75.—Groenl; Labrad.
hakonensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 599.—Japon.
Halleriana, DC. Fl. Fr. Suppl. 256 = Deyeuxia Halleriana.
Halleriana, Fries, Summa Veg. 240 = gracilescens.
Hartmaniana, Fries, l. c. 241 = ambigua.
hirtiguma, *Steud. Syn. Pl. Gram.* 188.—Am. bor.
holciformis, *Jaub. & Spach, Illustr.* iv. 61.—Tibet.
Hookeriana, *Steud. Syn. Pl. Gram.* 192.—Zeylan.
Howellii, Vasey, in Coult. Bot. Gaz. vi. (1881) 271 = Deyeuxia Howellii.
Huebneriana, Reichb. Fl. Germ. Excurs. 27 = epigeios.
Humboldtiana, *Steud. Nom.* ed. II. i. 250.—N. Granat.
hyperbores, *Lange, in Fl. Dan.* t. 2942.—Groenl.
inexpansa, A. Gray, ex Torr. Fl. N. York, ii. 445. t 152 = Deyeuxia confinis.
intermedia, Lindeb. ex Nym. Consp. 799 = Hartmaniana.
intermedia, Steud. Nom. ed. II. i. 250 = Deyeuxia intermedia.
Jamesoni, *Steud. Syn. Pl. Gram.* 191.—Ecu dor.
javana, *Steud. l. c.* 193.—Java.
junciformis, Steud. Nom. ed. II. i. 250 = Deyeuxia junciformis.
Karka, J. F. Gmel. Syst. 172 = Arundo Karka.
laguroides, *Regel, in Act. Hort. Petrop.* vii. (1880) 641. —Turkestan.
Lagurus, Koel. Desc. Gram. 112 = Imperata arundinacea.
lanceolata, *Roth, Tent. Fl. Germ.* i. 34.—Europ.; As. temp.
lanceolata, Stokes, in With. Arr. Brit. Pl. ed. III. ii 122 = epigeios.
lanceolata, Trin. ex Steud. Nom. ed. II. i. 250 = purpurea.
Langsdorffii, Auct. ex Nym. Consp. 799 = phragmitoides.
Langsdorffii, Trin. Gram. Unifi. 225. t. 4. f. 10 = Deyeuxia Langsdorffii.
lapponica, Hartm. Scand. Fl. ed. I. 46; Trin. Gram. Unifi. 226 = Deyeuxia lapponica.
lapponica, Hook. Brit. Fl. ed. IV. 34 = Deyeuxia neglecta.
latifolia, Blytt, Norges Fl. i. 90 = phragmitoides.
laxa, *Hegetschw. Fl. Schw.* 65.—Helvet.
laxa, Host, Gram. iv. 25 = littorea.
laxior, C. Muell. in Walp. Ann. vi. 1115 = Deyeuxia Hartmanniana.
Lersii, Koel. Desc. Gram. 107 = Deschampsia juncea
Lersii, Trin. ex Steud. Nom. ed. II. i. 316, nomen. = Stipa Calamagrostis.
lenkoranensis, *Steud. Syn. Pl. Gram.* 188.—Reg. Caucas.
leucantha, *Delile, ex Steud. Nom.* ed. II. i. 250 = Stipa Calamagrostis.
lithuanica, Bess. in Roem. & Schult. Add. ad Mant. ii. 602 = phragmitoides.
littorea, *DC. Fl. Fr.* vi. 255.—Europ.; As. temp.
longearistata, *Fisch. ex Ledeb. Fl. Ross.* iv. 488, nomen. —Hab. ?
longifolia, Hook. Fl. Bor. Am. ii. 241 = Ammophila longifolia.
macloviana, *Steud. Syn. Pl. Gram.* 192.—Ins. Falkl.
maritima, *Kunth, Rev. Gram.* i. 75.—Scania.
mauritanica, Spreng. Syst. i. 252 = Arundo mauritanica.
megapotamica, Steud. Syn. Pl. Gram. 192 = Andropogon laguroides.
mexicana, Kunth, Syn. iv. 478 = Muehlenbergia calamagrostis.
mexicana, Nutt. Gen. Am. i. 46 (excl. syn.) = Deyeuxia canadensis.
Michauxii, Trin. ex Steud. Nom. ed. II. i. 250 = Deyeuxia canadensis.
montana, *DC. Fl. Fr.* vi. 254.—Germ.; Austr.; Gall.
montana, Host, Gram. i. 146 = Deyeuxia sylvatica.
montevidensis, *Nees, in Mart. Fl. Bras.* ii. 401.—Reg. Argent.
Munroana, *Boiss. Fl. Orient.* v. 523.—Afghan.
Munroi, *Baker, l. c.* 525.—Afghan.
mutica, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.

CALAMAGROSTIS:—
Nessii, Steud. Nom. ed. II. i. 250; *Syn. Pl. Gram.* 193. —Ind. or.
neglecta, Gaertn. Mey. & Scherb. Fl. Wett. i. 94 = Deyeuxia neglecta.
nepalensis, Nees, ex Steud. Syn. Pl. Gram. 193.—Nepal.
nigricans, Mérat, Fl. Paris, ed. II. ii. 33 = Phragmites communis.
nipponica, Franch. & Sav. Enum. Pl. Jap. ii. 599 = Japon.
nutans, Sart. ex Reichb. Fl. Germ. Exc. 26 = Deyeuxia Halleriana.
nuttabnata, Steud. Syn. Pl. Gram. 190 = Deyeuxia nutkaensis.
Nuttalliana, Steud. Nom. ed. II. i. 251 = Deyeuxia Nuttalliana.
obscura, Downar, in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 605.—Podol.
obtusata, Trin. Gram. Unifl. 225. t. 4. f. 11.—Sibir.
obtusata, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104; xxix. (1856) i. 26 = Deyeuxia deschampsioides.
odessana, Bess. ex Andrz. Rys Botan. ii. (1830) 26.—Tauria.
olympica, Boiss. Diagn. Ser. I. v. 70.—Reg. Caucas.; Persia.
Onaei, Franch. & Sav. Enum. Pl. Jap. ii. 598.—Japon.
oregonensis, Buckl. in Proc. Acad. Sc. Philad. '1862' (1863) 92 = Deyeuxia Langsdorffii.
Orizabae, Steud. Syn. Pl. Gram. 193 = Deyeuxia Orizabae.
ovata, Steud. Nom. ed. II. i. 251 = Deyeuxia ovata.
pallens, Steud. l. c. = Deyeuxia pallens.
pallida, C. Muell. in Walp. Ann. vi. 986 = phragmitoides.
pallidiflora, C. Muell. l. c. = phragmitoides.
parviflora, Rupr. in Beitr. Pfl. Russ. Reich. iv. 36 = Deyeuxia sylvatica
patula, Steud. Syn. Pl. Gram. 422.—Chili.
persica, Boiss. Diagn. Ser. I. vii. 120 = littorea.
phalaroides, Regel, in Act. Hort. Petrop. vii. (1880) 641, in textu.—Turkestan.
phleoides, Steud. Nom. ed. II. i. 251.—Am. austr.?
phragmitoides, Hartm. Scand. Fl. ed. IV. 25.—Reg. bor.
Pickeringii, A. Gray, Man. Bot. U. St. ed. I. 547 = Deyeuxia Pickeringii.
piscatoria, Steud. Syn. Pl. Gram. 193.—Cochinch.
planifolia, Trin. ex Steud. Nom. ed. II. i. 251 = Deyeuxia planifolia.
Plinii, J. F. Gmel. Syst. 172 = Arundo mauritanica.
plumosa, Spreng. Syst. i. 258 = Stipa papposa.
poaeoides, Steud. Syn. Pl. Gram. 423.—Reg. Magell.
poaeoides, Trin. ex Steud. Nom. ed. II. i. 251 = Poa subuniflora.
Porteri, A. Gray, in Proc. Am. Acad. vi. (1862) 79 = Deyeuxia Porteri.
Pseudophragmites, Aschers. Fl. Brandenb. 821 = littorea.
Pseudophragmites, Blytt, Norges Fl. i. 89 = phragmitoides.
Pseudophragmites, Link, ex Reichb. Fl. Germ. Exc. 26 = Deyeuxia Halleriana.
Pseudophragmitis, Koel. Desc. Gram. 106 = littorea.
pulchella, Blytt, Norges Fl. i. 93 = phragmitoides.
pulchella, *Griseb. in Goett. Nachr.* (1868) 78.—Ind. or.
pulchella, Sart. ex Reichb. Fl. Germ. Excurs. 26 = Deyeuxia Halleriana.
puznila, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 176. —Ins. Galap.
purpurascens, R. Br. in Richards. App. Frankl. Journ. 8 = Deyeuxia sylvatica.
purpurea, *Trin. Gram. Unifl.* 219.—Sibir. baical.
pyramidalis, Host, Gram. iv. t. 49 = Deyeuxia sylvatica.
quadriseta, Spreng. Syst. i. 258 = Deyeuxia quadriseta.
quitensis, H. B. & K. Nov. Gen. et Sp. i. 133 = Muhlenbergia calamagrostides.
ramosa, Host, Gram. iv. t. 44 = lanceolata.
rariflora, C. Muell. in Walp. Ann. vi. 986 = Deyeuxia Halleriana.
rariflora, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 156 = Deyeuxia sylvatica.
recta, Trin. ex Steud. Nom. ed. II. i. 251 = Deyeuxia recta.
retroflexa, Link, ex Steud. l. c. = Deyeuxia Forsteri.
rigens, Trin, Summa Veg. 240 = epigeios.

CALAMAGROSTIS:—
rigida, Trin. ex Steud. Nom. ed. II. i. 251 = Deyeuxia rigida.
robusta, C. Muell. in Walp. Ann. vi. 985 = Deyeuxia Halleriana.
robusta, *Franch. & Sav. Enum. Pl. Jap.* ii. 600.—Japon.
Roylcana, *Steud. Nom.* ed. II. i. 251.—Ind. or.
Royleі, Steud. Syn. Pl Gram. 193 = prasec.
rubella, Boiss. Fl. Orient. v. 625 = littorea.
rubescens, Buckl. in Proc. Acad. Sc. Philad. '1862' (1863) 92 = Deyeuxia rubescens.
rubicunda, Blytt, Norges Fl. i. 91 = phragmitoides.
rubra, Bosc, ex Kunth, Enum. Pl. 478 = Eriauthus saccharoides.
rudis, Steud. Syn. Pl. Gram. 193 = Deyeuxia scabra.
rupestris, *Trin. ex Steud Nom.* ed. II. i. 251, *nomen*; *Doell. in Mart. Fl. Brasil.* ii. 111. 53.—Bras.
sachalinensis, *F. Schmidt, Reisen Amuri.* 202.—Terr. Amur.
scabra, *Presl, Rel. Haenk.* i. 284.—Sinus Nootka.
scabrescens, *Griseb. in Goett. Nachr.* (1868) 79.—Ind. or.
Schiedeana, Steud. Syn. Pl. Gram. 193 = Deyeuxia Schiedeana.
Schimperiana, *Hochst. in Flora*, xxxviii. (1855) 202. —Abyss.
Schlechteri, Bess. Prim. Fl. Gallic. i. 69 = lanceolata.
Schwabi, Spreng. Syst. i. 253 = Gastridium australe.
sciuroides, *Franch. & Sav. Enum. Pl. Jap.* ii. 600.—Japon.
sedenensis, Loisel. Fl. Gall. ed. II. i. 54 = varia.
sesquiriflora, Steud. Syn. Pl. Gram. 190 = Deyeuxia toluccana.
simplex, *Boiss. Fl. Orient.* v. 623.—Reg. Caucas.
sitchensis, *Trin. ex Ledeb. Fl. Ross.* iv. 433.—Ins. Sitcha.
speciosa, Host, Gram. iv. t. 45 = Stipa Calamagrostis.
spicigera, Steud. Nom. ed. II. i. 251 = Deyeuxia spicigera.
splendens, *Steud. l. c.*—Ins. Catharin.
stricta, Koel. Desc. Gram. 105 = Deyeuxia neglecta?
stricta, *Trin. Gram. Unifl.* 226.—Europ.; Am. bor.
strigosa, Hartm. Skand. Fl. ed. I. 46 = Deyeuxia strigosa.
subalpina, Schur, Enum. Pl. Transs. 740 = Deyeuxia sylvatica.
subspicata, Desv. Opusc. 60 = Trisetum subspicatum?
subulata, Dum. Agr. Belg. t. 9. f. 36 = montana.
subulata, Steud. Nom. ed. II. i. 251 = varia.
sylvatica, *Bess. Prim. Fl. Gallic.* i. 70.—Europ.
sylvatica, DC. Fl. Fr. v. 253 = Deyeuxia arundinacea.
sylvatica, Host, Gram. iv. t. 48 = varia.
sylvatico-stricta, Knehlew. ex Ledeb. Fl. Ross. iv. 429 = Deyeuxia neglecta.
tenella, *Host, Gram.* iv. t. 50.—Eur. med.; As. bor.
tenuiflora, H. B. & K. Nov. Gen. et Sp. i. 134 = Muehlenbergia tenuiflora.
Thrysoides, C. Koch, in Linnaea, xxi. (1848) 388 = epigeios.
tianschanica, *Rupr. Sert. Tianach.* 34.—Turkestan.
toluccana, Trin. ex Steud. Nom. ed. II. i. 250 = Deyeuxia toluccana.
transylvanica, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 154 = laxa.
Trinii, Rupr. in Beitr. Pfl. Russ. Reich. iv. 36 = Deyeuxia acutiflora.
varia, *Beauv. Agrost.* 157.—Europ.; As. bor.
varia, Boland. ex Bot. Calif. ii. 280 = Deyeuxia Bolanderi.
varia, Gren. & Godr. Fl. Fr. iii. 477 = montana.
varia, Host, Gram. iv. t. 47 = Deyeuxia Halleriana.
variegata, With. Bot. Arr. Brit. Pl. ed. III. ii. 124 = Phalaris arundinacea.
variflora, C. Muell. in Walp. Ann. vi. 1115 = Deyeuxia Halleriana.
velutina, *Steud. Nom.* ed. II. i. 251.—Chili.
villosa, J. F. Gmel. Syst. 172 = Deyeuxia sylvatica.
villosa, Mutel, Fl. Fr. iv. 41 = Deyeuxia Halleriana.
vilmorini, Bess. ex Roem. & Schult. Add. ad Mant. ii. 602 = phragmitoides.
vinealis, *Baumg. ex Steud. Syn. Pl. Gram.* 186.—Transylv.
viridescens, Steud. l. c. 192 = seq.
viridiflavescens, Steud. Nom. ed. II. i. 251 = Deyeuxia viridiflavescens.
Villdenowii, Steud. Syn. Pl. Gram. i. 193 = Deyeuxia Forsteri.

CALAMINA, Beauv. Agrost. 128. t. 23. f. 1. (1812) = **Apludа**, Linn. (Gramin.).
gigantea, Beauv. l. c. 129 = Aplueda mutica.
gigantea, Roem. & Schult. Syst. ii. 810 = Anthistiria gigantea.
humilis, Presl, Rel. Haenk. i. 344 = Aplueda mutica.
imberbis, Beauv. Agrost. 128 = Anthistiria imberbis.
mutica, Beauv. l. c. = Aplueda mutica.
Schima, Beauv. l. c. 157 = Ischaemum laxum.
Themeda, Beauv. l. c. = Anthistiria imberbis.

CALAMINTHA, Adans. Fam. ii. 192 (1763) = **Nepeta**, Linn. (Labiat.).

CALAMINTHA, [Tourn.] Lam. Fl. Fr. ii. 393 (1778). *LABIATAE*, Benth. & Hook. f. ii. 1190.
ACINOS, Moench, Meth. 407 (1794).
CLINOPODIUM, Tourn. ex Linn. Gen. ed. I. 170 (1737), nomen prius.
FAUCHARDIA, Dulac, Fl. Hautes-Pyr. 402 (1867).
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 191.—Abyssin.
acinoides, Nym. Consp. 589 = suaveolens.
Acinos, [*Claire.*] *Man. Herb.* 197 (*err. typ.* 917).—Europ.; Reg. Caucas.
acuminata, Frivald. in Flora, xviii. (1835) 332 = suaveolens.
adscendens, Willk. & Lange, Prod. Fl. Hispan. ii. 413 = officinalis.
aegyptiaca, Lam. ex Schur, Enum. Pl. Transs. 531 = C. Clinopodium.
aetnensis, Strobl, in Oestr. Bot. Zeitschr. xxiv. (1874) 29 = alpina.
alba, Lam. ex Steud. Nom. ed. II. i. 251 = Melissa alpina.
alba, Reichb. Fl. Germ. Exc. 828 = Micromeria rupestris
alpina, *Lam. Fl. Fr.* ii. 394.—Europ. austr.; Reg Mediterr.
annua, *Schrenck, in Bull. Sc. Acad. Pétersb.* x. (1842 358.—Rossia.
arundana, *Benth. in DC. Prod.* xii. 233.—Hispan.
arvensis, Lam. Fl. Fr. ii. 394 = C. Acinos.
asperata, Jord. Obs. Pl. Crit. iv. (1846) 8 = officinalis
athonica, Reichb. Fl. Germ. Exc. 829 = C. Nepeta.
atlantica, Ball, in journ. Bot. xiii. (1875) 175 = C. Clinopodium.
atlantica, Coss. in Bull. Soc. Bot. Fr. ix. (1862) 176 = grandiflora.
baetica, Boiss. & Reut. Pugill. Pl. Nov. (1852) 92 = officinalis.
betulifolia, *Boiss. & Bal. Diagn.* Ser. II. iv. 14.—Cilicia.
Bieberstеinii, C. Koch, ex Ledeb. Fl. Ross. iii. 355 = umbrosa.
bysantina, C. Koch, in Linnaea, xxi. (1848) 672 = C. Nepeta.
caerulescens, *Wedd. Chlor. And.* ii. 149. t. 62.—N. Granat.
cana, Auct. ex Heynh. Nom. ii. 104 = graveolens.
candidissima, *Munby, ex Benth. in DC. Prod.* xii. 226. —Algeria.
canescens, Presl, Fl. Sicul. 37 = C. Nepeta.
canescens, Torr. & Gray, ex Benth. in DC. Prod. xii. 229 = Conradina canescens.
caroliniana, *Sweet, Hort. Brit.* 409.—Am. bor.
chinensis, *Benth. in DC. Prod.* xii. 233.—China.
Clinopodium, *Benth. l. c.*—Reg. bor. temp.
coccinea, *Benth. l. c.* 229.—Am. bor.
confinis, *Hance, in Journ. Bot.* vi. (1868) 381.—China.
corsica, *Benth. in DC. Prod.* xii. 231.—Ins. Corsica.
cretica, *Benth. l. c.* 227.—Ins. Creta.
cretica, Lam. Fl. Fr. ii. 395 = Micromeria marifolia.
crootica, Host, Fl. Austr. ii. 132 = Micromeria Piperella.
cryptantha, *Vatke, in Linnaea*, xxxvii. (1871-73) 328. —Abyss.
debilis, *Benth. in DC. Prod.* xii. 232.—Sibiria.
dentata, *Chapm. Fl. S. U. St.* 318.—Florida.
depauperata, *Bunge, Lab. Pers.* 35.—Persia.
dilatata, Schrad. ind. Sem. Hort. Goetting. (1835) = C. Nepeta.
Elaeoleuca, *F. Schultz, Arch. Fl. Fr. et Allem.* 178.—Europ.
exigua, Nym. Consp. 589 = graveolens.
Fenalii, Vis. Revis. Pl. Min. Cogn. (1855) 7 = subnuda.
florida, *Boiss. Diagn.* Ser. I. xii. 90.—Cilicia.
foliosa, *Opiz, Natural.* iii. (1829) [20].—Europ.

CALAMINTHA:—
fruticosa, Stead. Nom. ed. II. i. 251 = Micromeria marifolia.
glabella, *Benth. in DC. Prod.* xii. 280.—Am. bor.
glabella, A. Gray, Man. ed. II. 307 = Nuttallii.
glabrescens, Boiss. Fl. Orient. iv. 580 = staminea.
glandulosa, *Benth. Lab. Gen. et Sp.* 387.—Corsica.
glomerulosa, Boiss. & Heldr. ex Benth. in DC. Prod. xii. 284 = origanifolia.
gracilis, *Benth. l. c.* 282.—Java.
granatensis, Boiss. & Reut. Pugill. Pl. Nov. 94 = alpina.
grandiflora, *Moench, Meth.* 408.—Europ.
grandiflora, Pursh, Fl. Am. Sept. ii. 414 = caroliniana.
graveolens, *Benth. in DC. Prod.* xii. 281.—Reg. Mediterr. ; Ab. Min.
graveolens, Coss. ex Willk. & Lange, Prod. Fl. Hispan. ii. 415 = umbrosa.
Gussonei, Tod. ex Nym. Consp. 588 = C. Nepeta.
Hausknechtii, *Bunge, Lab. Pers.* 36.—Persia.
hederacea, Soop. Fl. Carn. ed. II. i. 423 = Nepeta Glechoma.
Herba-barona, Heynh. Nom. i. 144 = Thymus Herba-barona.
heterophylla, Heynh. i. c. = C. Acinos.
heterotricha, *Boiss. & Reut. Pugill. Pl. Nov.* 93.—Hispan.
hispidula, *Boiss. & Reut. l. c.*—Hispan.
ilicifolia, A. Gray, in Proc. Am. Acad. viii. (1873) 366 = Acanthomentha ilicifolia.
incana, *Boiss. ex Benth. in DC. Prod.* xii. 286.—Graecia et Ins. ; Syria.
Jahniana, *Simk. in Termtsz. Füzetek*, ix. (1885) 25.—Europ.
Langei, Nym. Consp. 589 = suaveolens.
longicaulis, *Benth. in DC. Prod.* xii. 284.—Reg. Himal.
macrostema, *Benth. l. c.* 280.—Mexic.
marifolia, Stead. Nom. ed. II. i. 251 = Micromeria marifolia.
mentha, *Benth. in DC. Prod.* xii. 281.—Thracia.
menthaefolia, Host, Fl. Austr. ii. 120 = officinalis.
meridionalis, Nym. Consp. 589 = alpina.
mimuloides, *Benth. Pl. Hartw.* 331.—Calif.
mollis, Jord. ex Lamotte, Prod. ii. 539 = C. Nepeta.
moluccana, *Miq. Fl. Ind. Bat.* ii. 968.—Ins. Moluce. ; Timor.
montana, Lam. Fl. fr. ii. 396 = officinalis.
multicaulis, *Maxim. in Bull. Acad. Pétersb.* xx. (1875) 466.—Japon.
nebrodensis, Kern. & Strobl, in Oestr. Bot. Zeitschr. xxiv. (1874) 171 = alpina.
neglecta, *Opiz, Natural.* iii. (1828) [30].—Europ.
nepetoides, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 58 = umbrosa.
Nepeta, *Savi, Fl. Pis.* (1798) 197.—Europ. ; Afr. bor. ; Oriens.
Nepeta, Willk. in Flora, xxxv. (1852) 277 = officinalis.
nepetoides, Jord. Obs. Pl. Crit. iv. (1846) 16 = C. Nepeta.
nervosa, *Pomel, Nouv. Mat. Fl. Atl.* 307.—Afr. bor.
nivea, *Boiss. in Ann. Sc. Nat. Sér.* IV. ii. (1854) 253. —As. Min.
Nuttallii, *Benth. in DC. Prod.* xii. 280.—Am. bor.
oblique, Host, Fl. Austr. ii. 131 = C. Nepeta.
officinalis, *Moench, Meth.* 409.—Europ. ; Oriens.
origanifolia, *Boiss. Diagn.* Ser. I. xii. 50.—As. Min. ; Syria.
origanifolia, Host, Fl. Austr. ii. 130 = Micromeria Pulegium.
origanifolia, Vis. Fl. Dalm. ii. 199 = Micromeria origanifolia.
Palmeri, *A. Gray, in Proc. Am. Acad.* xi. (1876) 100. —Calif.
pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. xii. 92.—Pamphylia.
paradoxa, *Vahl, in Linnaea*, xxxvii. (1871–72) 327.—Abyssin.
parviflora, Lam. Fl. fr. ii. 396 = C. Nepeta.
patavina, Heldr. ex Nym. Consp. 588 = suaveolens.
patavina, *Host, Fl. Austr.* ii. 132.—Oriens ; Europ. austr. or.
Pinpinella, Reichb. Fl. Germ. Excurs. 328 = Thymus Pimpinella.
Pinpinella, Schkuhr, & Vahot. Syll. Fl. Croat. 80 = Micromeria croatica.
pipenelloides, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 96.—Lycia.
Pulegium, Reichb. f. Ic. Fl. Germ. xviii. 45. f. 1978 = Micromeria Pulegium.
purpurascens, *Benth. in DC. Prod.* xii. 281.—Hispan.

CALAMINTHA:—
repens, Benth. l. c. 288 = umbrosa.
rotundifolia, Benth. l. c. 282 = umbrosa.
rotundifolia, Ces. Pass. & Gib. Comp. Fl. Ital. 303 = alpina.
rotundifolia, Host, Fl. Austr. ii. 131 = C. Nepeta.
rotundifolia, Loec. & Pard. Ser. Inconf. 82 = Ziziphora sclinoides.
rotundifolia, Reichb. Ic. Fl. Germ. xviii. 43 = patavina.
rupestris, Host, Fl. Austr. ii. 131 = Micromeria rupestris.
simensis, *Benth. in DC. Prod.* xii. 280.—Abyssin.
spicigera, C. Koch, in Linnaea, xxi. (1848) 671 = Micromeria serpyllifolia.
Sprunerî, Boiss. Diagn. Ser. I. xii. 53 = C. Nepeta.
staminea, *Boiss. l. c.* 51.—Kurdistan.
stenostoma, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 96.—Lycia.
stricta, Reichb. Ic. Fl. Germ. xviii. 44 = officinalis.
suaveolens, *Boiss. Diagn.* Ser. I. xii 54.—Graecia.
subnuda, Host, Fl. Austr. ii. 130 = Micromeria Pulegium.
subnuda, Reichb. Fl. Germ. Excurs. 329 = thymifolia.
sylvatica, Bromf. in Engl. Bot. Suppl. t. 2897 = officinalis.
thymifolia, *Host, Fl. Austr.* ii. 132.—Carniola.
thymifolia, Reichb. Fl. Germ. Excurs. 328 = Micromeria rupestris.
trichotoma, Moench, Meth. 409 = C. Nepeta.
umbrosa, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 6 ; ix. *Suppl.* 8.—Oriens ; Ind. or.
umbrosa, Reichb. Fl. Germ. Excurs. 329 = officinalis.
usuriensis, *Regel & Maack, in Regel, Tent. Fl. Ussur.* 116.—Reg. Amur.
vulgaris, [Clairv.] Man. Herb. 197 (err. typ. 917) = C. Nepeta.

CALAMOCHLOA, Fourn. in Bull. Soc. Bot. Fr. xxix. (1882) 178. *GRAMINEAE*, Benth. & Hook. f. iii. 1178.
filifolia, *Fourn. l. c.*—Mexic.

CALAMOCHLOË, Reichb. Consp. 52 (1828) = Arundinella, Raddi (Gramin.).

CALAMOSAGUS, Griff. in Calc. Journ. Nat. Hist. v. (1845) 22 = Korthalsia, Blume (Palm.).
harinaefolius, Griff. Palms Brit. E. Ind. 29. t. 184 = K. wallichiaefolia.
lacinulosus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 38 = K. laciniosa.
ochriger, Griff. Palms Brit. E. Ind. 31. t. 216 = K. pectinata.
polystachys, H. Wendl. in Kerch. Palm. 235, sphalm. = K. polystachys.
scaphiger, Griff. Palms Brit. E. Ind. 30. t. 148 a = K. wallichiaefolius.
wallichiaefolius, Griff. in Calc. Journ. Nat. Hist. v. (1845) 24 = K. wallichiaefolia.

CALAMPELIS, D. Don, in Edinb. N. Phil. Journ. vii. (1829) 89 = Eccremocarpus, Ruiz & Pav. (Bignoniac.).
scaber, D. Don, l. c. = E. scaber.

CALAMUS, Auct. ex Linn. Gen. ed. I. 104 (1737) = Acorus, Linn. (Aroid.).
aromaticus, Gueldenst. ex Ledeb. Fl. Ross. iv. 18 = Acorus Calamus.

CALAMUS, Linn. Sp. Pl. 325 (1753). *PALMAE*, Benth. & Hook. f. iii. 931.
ROTANG, Adans. Fam. ii. 24 (1763).
ROTANGA, Boerh. ex Crantz, Inst. i. 197 (1766).
acanthophis, Griff. Palms Brit. E. Ind. t. 216 = Dasmonorops malaccensis.
acanthospathus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 20.—Reg. Himal.
accedens, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 38 = Daemonorops Draco.
adspersus, Blume, Rumphia, iii. 46 = Daemonorops adspersa.
albus, *Pers. Syn.* i. 382.—Amboina.
amarus, Lour. Fl. Cochinch. 210 = tenuis ?
ambinervis, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 30 = Daemonorops calapparius.
americanus, Hort. ex Stend. Nom. ed. II. i. 262 = Desmoncus americanus.

CALAMUS:—
amplectens, *Becc. Malesia*, ii. (1884) 78.—Borneo.
ancepe, Blume, Rumphia, iii. 65. t. 149 = melanoloma.
andamanicus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 311.—Ins. Andaman.
angustifolius, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 39 = Daemonorops angustifolius.
arborescens, *Griff. l. c.* 33.—Burma.
asperrimus, *Blume, in Schult. f. Syst.* vii. 1837.—Java.
assamicus, *Linden, Illustr. Hortic.* xxvIII. (1881) 16, nomen.—Assam.
aurvus, Reinw. ex Mart. Hist. Nat. Palm. iii. 341 = ornatus.
australis, *Mart. l. c.* 342.—Austral.
barbatus, Mackint. in Bijdr. Nat. Wetensch. v. (1880) 178, nomen ; Blume, Rumphia, iii. 42 = Daemonorops barbatus.
Blancoi, *Kunth, Enum. Pl.* iii. 505.—Ins. Philipp.
bornaensis, Miq. Anal. Bot. Ind. 4 = javensis.
brevifrons, Mart. Hist. Nat. Palm. iii. 338 = Blancoi.
bureensis, *Mart. l. c.* 336.—Ins. Buro.
caesius, *Blume, Rumphia*, iii. 57.—Malaya.
calapparius, Mart. Hist. Nat. Palm. ed. prior, 209 (iii 331) = Daemonorops calapparius.
calicarpus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 23 = Daemonorops calicarpus.
calolepis, Miq. Pl. Jungh. 159 = melanoloma.
caryotoides, *A. Cunn. ex Mart. Hist. Nat. Palm.* iii. 338.—Austral.
castaneus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 38.—Malaya.
Cawa, *Blume, Rumphia*, iii. 31.—Amboina.
ciliaris, *Blume, in Schult. f. Syst.* vii. 1830.—Malaya.
cochlearus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 29 = Daemonorops cochlearus.
collinus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 31.—Ind. or.
concinnus, *Mart. Hist. Nat. Palm.* iii. 332. t. 116.—Burma.
crinitus, Miq. Anal. Bot. Ind. 6 = Daemonorops crinitus.
cuspidatus, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 436. t. 41 = Eremospatha cuspidata.
deultuca, Hort. ex H. Wendl. in Kerch. Palm. 236 = Acanthophoenix rubra.
decrustus, *Mann & H. Wendl. in Trans. Linn. Soc.* xxiv. (1864) 439.—Afr. trop.
delicatulus, *Thw. Enum. Pl. Zeyl.* 330.—Zeylan.
depressiusculus, *Teijsm. & Binn. Cat. Hort. Bog.* 74. —Sumatra.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 594.—Malaya.
dioicus, *Lour. Fl. Cochinch.* 211.—Cochinch.
discolor, Mart. Hist. Nat. Palm. iii. 341 = siphonospathus.
Draco, Willd. Sp. Pl. ii. 203 = Daemonorops Draco.
elegans, *Hort. ex H. Wendl. in Kerch. Palm.* 236.—Hab. ?
elongatus, Miq. Anal. Bot. Ind. 6 = Daemonorops elongatus.
ephippiatus, *Mart. Hist. Nat. Palm.* iii. 336.—Java.
equestris, Blume, in Schult. f. Syst. vii. 1830 = javensis.
equestris, *Willd. Sp. Pl.* ii. 204.—Ins. Moluce.
erectus, *Roxb. Hort. Beng.* 73 ; *Fl. Ind.* iii. 774.—Reg. Himal.
exilis, *Griff. ex Mart. Hist. Nat. Palm.* iii. 333.—Malaya.
extensus, Mart. Hist. Nat. Palm. iii. 300, partim, t. 116. f. 4.=viminalis.
extensus, Roxb. Hort. Beng. 73 ; Fl. Ind. iii. 777 = palustris !
farinosus, *Linden, in Illustr. Hortic.* xix. (1872) t. 199. —Sumatra.
fasciculatus, Roxb. Hort. Beng. 73 ; Fl. Ind. iii. 779 = viminalis.
Fernandesii, H. Wendl. in Kerch. Palm. 236 = Oxleyanus.
flavus, Miq. Anal. Bot. Ind. 6 = Daemonorops flavus.
Flagellum, Griff. ex Mast. Hist. Nat. Palm. iii. 338 = polygamus.
floribundus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 36.—Reg. Himal.
Gaudichaudii, H. Wendl. in Kerch. Palm. 236 = Daemonorops Gaudichaudii.
grandiflorus, Griff. ex Mart. Hist. Nat. Palm. iii. 336 = Plectocomia grandiflora.
guruinsitus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 27 = Daemonorops guruinsitus.

CALAMUS:—
glaucescens, Blume, Rumphia, iii. 55 = cassia.
glaucescens, D. Dietr. Syn. Pl. ii. 1064 = Cerasolobus glaucescens.
gracilipes, Miq. in Verh. Kon. Akad. Wetensch. xi. (1858) n. v. 18 = Daemonorops longipes, *Miq.*
gracilis, Blanco, Fl. Filip. ed. 1. 267 = Blancoi.
gracilis, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 781.—Reg. Himal.
gracilis, Thwaites, Enum. Pl. Zeyl. 330 = pachystemonus.
graminoxus, *Blume, Rumphia*, iii. 31.—Amboina.
grandiflorus, Steud. Nom. ed. II. i. 552, sphalm. = Ancistrophyllum secundiflorum.
grandis, Griff. in Calc. Journ. Nat. Hist. v. (1845) 84 = Daemonorops grandis.
Griffithianus, *Mart. Hist. Nat. Palm.* iii. 382.—Malaya.
Guruba, *Buch.-Ham. ex Kunth, Enum. Pl.* iii. 210.—Reg. Himal.; Burma.
Hasskarianus, Mart. Hist. Nat. Palm. iii. 387 = mollis.
Helferianus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 218.—Burma.
Helostropium, Bech.-Ham. ex Kunth, Enum. Pl. iii. 210 = tenuis.
heteracanthus, *Maclot, in Bijdr. Nat. Wetensch.* v. (1830) 178.—N. Guinea.
heteroideus, Blume, Rumphia, iii. 46 = Reinwardtii.
hirsutus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1858) n. v. 28 = Daemonorops Hystrix.
Hookeri, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 434 = Eremospatha Hookeri.
horsfa, *Blume, Rumphia*, iii. 43.—Java.
hystrix, Wall. ex Voigt, Hort. Suburb. Calc. 639; Miq. Fl. Ind. Bat. iii. 113, nomen = arborescens.
Hasgelianus, *Mart. Hist. Nat. Palm.* iii. 388.—Malabar.
humilis, Roxb. Hort. Beng. 73; Fl. Ind. iii. 778 = latifolius?
hygrophilus, Griff. Palms Brit. Ind. t. 318 C = Daemonorops hygrophilus.
hypolencus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 208.—Burma.
Hystrix, Griff. in Calc. Journ. Nat. Hist. v. (1845) 70 = Daemonorops Hystrix.
inermis, T. Anders. in Journ. Linn. Soc. xi. (1871) 11 = latifolius.
insignis, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 59.—Malaya.
intermedius, Griff. l. c. 86 = Daemonorops intermedius.
javensis, *Blume, Rumphia*, iii. 62.—Malaya.
Jenkinsianus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 81 = Daemonorops Jenkinsianus.
Jenkinsianus, Griff. Palms Brit. Ind. 40. f. 186 A. f. 3 (non p. 89) = polygamus.
Korthalsii, Miq. Anal. Bot. Ind. i. 6 = Daemonorops Korthalsii.
laevigatus, *Mart. Hist. Nat. Palm.* iii. 389.—Malaya.
laevis, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 430. t. 38 (Ancistrophyllit sp.).—Afr. trop.
latifolius, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 20. t. 31. A. = palustris.
latifolius, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 775.—Burma.
latisplum, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1858) n. v. 29.—Sumatra.
leptopus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 73 = Daemonorops leptopus.
leptospadix, *Griff. l. c.* 49.—Reg. Himal.
Lewisianus, Griff. l. c. 87 = Daemonorops Lewisianus.
Lindeni, *Rodigas, in Illustr. Hortic.* xxx. (1883) t. 499.—Malaya.
Horsfieldii, Blume, Rumphia, iii. 43 = viminalis.
longipes, Griff. in Calc. Journ. Nat. Hist. v. (1845) 68 = Daemonorops longipes, *Mart.*
longisetus, *Griff. l. c.* 88.—Burma.
macrocanthus, T. Anders. in Journ. Linn. Soc. xi. (1871) 10 = latifolius.
macrocarpus, Griff. ex Mart. Hist. Nat. Palm. iii. 383 = arenius.
macrocarpus, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 426 = Eremospatha macrocarpa.
macrophyllus, Miq. in Verh. Kon. Acad. Wetensch. xi. (1858) n. v. 26.—Sm. Celebes.
Manna, *Miq. Fl. Ind. Bat. Suppl.* 596.—Sumatra.

CALAMUS:—
manicatus, Teijsm. & Binn. ex Miq. Fl. Ind. Bat. iii. 135 = Daemonorops crinitus.
manillensis, *H. Wendl. in Kerch. Palm.* 287.—Ins. Philipp.
Mannii, H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 436 = Acoccalamus Mannii.
Margaritae, *Hance, in Journ. Bot.* xii. (1874) 266.—China.
marginalis, *Mart. Hist. Nat. Palm.* iii. 342.—Borneo.
maricinus, *Blume, Rumphia*, iii. 31, in adnot.—Java.
Materniensus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 76 = C. Guruba.
maximus, Blanco, Fl. Filip. ed. I. 266 = manillensis?
maximus, Reinw. ex Schult. f. Syst. vii. 1333 = Plectocomia elongata.
melanacanthus, *Mart. Hist. Nat. Palm.* iii. 383. t. 116. f. 13.—Burma.
melanochaetes, Miq. in Verh. Kon. Akad. Wetensch. xi. (1858) n. v. 28 = Daemonorops melanochaetes.
melanolepis, H. Wendl. in Kerch. Palm. 287 = Hugelianus.
melanoloma, *Mart. Hist. Nat. Palm.* iii. 399, 389. t. 116. f. 2.—Java.
Metzianus, *Schlecht. in Linnaea*, xxvi. (1853-55) 727. —Ind. or.
Meyenianus, *Schau. in Nov. Act. Nat. Cur.* xix. Suppl. i. (1843) 425.—Ins. Philipp.
micracanthus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 62.—Malacca.
microstylus, *Blume, Rumphia*, iii. 53.—Malaya.
mirabilis, Mart. ex Kunth, Enum. Pl. iii. 218 = Daemonorops mirabilis.
mishmeensis, Griff. in Calc. Journ. Nat. Hist. v. (1845) 84.—Mishmudus.
mollis, *Blanco, Fl. Filip.* ed. I. 264.—Ins. Philipp.
monoecus, Roxb. Hort. Beng. 73; Fl. Ind. iii. 783 = C. Rotang.
montanus, T. Anders. in Journ. Linn. Soc. xi. (1871) 9 = acanthosphathus.
monticolus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 90 = Daemonops monticolus.
Muelleri, *H. Wendl. & Drude, in Linnaea*, xxxix. (1875) 198.—Austral.
nigar, Willd. Sp. Pl. ii. 208 = Daemonorops niger.
nitidus, *Mart. Hist. Nat. Palm.* iii. 384.—Burma; Ind. or.
nivalis, Thw. ex Trimen, in Journ. Bot. xxiii. (1885) 268, sphalm. = rivalis.
nutantiflorus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 79 = Daemonorops Jenkinsianus.
oblongus, *Reinw. in Schult. f. Syst.* vii. 1329.—Java.
obtrurus, F. Muell. Fragm. v. 48 = australis.
ochreatus, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1858) n. v. 29.—Sumatra.
opacus, *Blume, Rumphia*, iii. 50.—Sumatra.
opacus, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 431. t. 41 = Lacosperma opaca.
ornatus, *Blume, ex Schult. f. Syst.* vii. 1326.—Malaya.
ovatus, Reinw. ex Kunth, Enum. Pl. iii. 205 = ceramicus.
ovoideus, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 269.—Zeylan.
Oxleyanus, *Teijsm. & Binn. Cat. Hort. Bog.* 75, nomen = Malaya.
pachystemonus, *Thw. Enum. Pl. Zeyl.* 331.—Zeylan.
palembanicus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1858) n. v. 29 = Daemonorops palembanicus.
pallens, Blume, Rumphia, iii. 51 = Reinwardtii.
palustris, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 60.—Burma.
paniculatus, *Roxb. Hort. Beng.* [106], nomen.—Penang.
parvicornus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 213. t. 29, 30.—Burma.
penicillatus, Roxb. Fl. Ind. iii. 781 = javensis?
periacanthus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1858) n. v. 28 = Daemonorops periacanthus.
petiolaris, Griff. in Calc. Journ. Nat. Hist. v. (1845) 93 = Daemonorops petiolaris.
petraeus, *Lour. Fl. Cochinch.* 209.—Cochinch.
philippensis, *Lindley, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Ins. Philipp.

CALAMUS:—
pisicarpus, *Blume, Rumphia*, iii. 39.—Ins. Moluco.
platyacanthos, Mart. ex Kunth, Enum. Pl. iii. 205 = oblongus.
platyspathus, *Mart. ex Kunth, l. c.* 209.—Burma.
plicatus, *Blume, Rumphia*, iii. 57.—Ins. Celebes.
polygamus, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 780.—Reg. Himal.
Pseudo-Rotang, Mart. ex Kunth, Enum. Pl. iii. 207 = viminalis.
pulcher, *Miq. Anal. Bot. Ind.* i. 3.—Borneo.
quinquenervius, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 777.—Reg. Himal.
radiatus, *Thw. Enum. Pl. Zeyl.* 331.—Zeylan.
radicalis, *H. Wendl. & Drude, in Linnaea*, xxxix. (1875) 195.—Austral.
ramosissimus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 78.—Malaya.
Reinwardtii, *Mart. Hist. Nat. Palm.* iii. 398. t. 119.—Malaya.
Rheedei, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 78, nota.—Malabar.
rhomboideus, *Blume, in Schult. f. Syst.* vii. 1333.—Malaya.
Riedelianus, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1858) n. v. 18.—Ins. Celebes.
rivalis, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 268, 305.—Zeylan.
Rotang, *Linn. Sp. Pl.* 325.—Ind. or.
Roxburghii, Griff. in Calc. Journ. Nat. Hist. v. (1845) 48 = C. Rotang.
Royleanus, Griff. l. c. 40 = tenuis.
rubr, Reinw. ex Kunth, Enum. Pl. iii. 208 = Daemonorops Draco.
rudentum, *Lour. Fl. Cochinch.* 209.—Cochinch.
rudentum, Mart. Hist. Nat. Palm. iii. 340 = rivalis.
rudentum, Roxb. Hort. Beng. 72; Fl. Ind. iii. 776 = albus.
Rumphii, Blume, Rumphia, iii. 38 = Daemonorops Rumphii.
Salakka, Willd. ex Steud. Nom. ed. II. i. 252 = Zalacca edulis.
schistacanthos, *Blume, Rumphia*, iii. 49.—Sumatra.
schizopethus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 83 = arcuatus.
Scipionum, Lam. Encyc. vi. 304 = C. Rotang.
Scipionum, *Lour. Fl. Cochinch.* 210.—Malaya.
secundiflorus, Benav. Fl. Owar. i. 15. t. 9, 10 = Ancistrophyllum secundiflorum.
siphonospathus, *Mart. Hist. Nat. Palm.* iii. 342.—Ins. Philipp.
spectabilis, *Blume, Rumphia*, iii. 55.—Sumatra.
stoloniferus, Teijsm. & Binn. Cat. Hort. Bog. 75 = horsfa.
strictus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1858) n. v. 28 = Daemonorops longipes.
subangulatus, *Miq. Fl. Ind. Bat. Suppl.* 594.—Sumatra.
symphysipus, *Mart. Hist. Nat. Palm.* iii. 386.—Ins. Celebes.
tenuis, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 780.—Reg. Himal.; Burma; Cochinch.
tetradactylus, *Hance, in Journ. Bot.* xiii. (1875) 289. —China.
tetrastichus, Blume, Rumphia, iii. 58 = javensis.
thyrsoclepis, *Hance, in Journ. Bot.* xii. (1874) 266.—China.
tigrinus, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 211. t. 25 = longisetus.
trichrous, Miq. in Verh. Kon. Akad. Wetensch. xi. (1858) n. v. 28 = Daemonorops trichrous.
unifarius, *H. Wendl. in Bot. Zeit.* xvii. (1859) 158.—Ins. Nicobar.; Java.
usitatus, Blanco, Fl. Filip. ed. I. 265 = Daemonorops Gaudichaudii.
usitatus, Mart. Hist. Nat. Palm. iii. 340 = Blancoi.
Verschaffelti, Hort. ex H. Wendl. in Kerch. Palm. 288 = Acanthophoenix crinita.
verticillaris, Griff. in Calc. Journ. Nat. Hist. v. (1845) 68 = Daemonorops verticillaris.
verus, Lour. Fl. Cochinch. 210 = pisicarpus.
viminalis, *Reinw. ex Mart. Hist. Nat. Palm.* iii. 385. t. U. 133.—Celebes; Java?
viminalis, *Willd. Sp. Pl.* ii. 208.—Ind. or.; Malaya.
Walkeri, *Hance, in Journ. Bot.* xii. (1874) 266.—China.
Wightii, Griff. Palms Brit. Ind. 102. t. 216 G = Hugelianus.

CALAMUS :—
Zalacca, Gaertn. Fruct. ii. 207. t. 189 = Zalacca edulis.
Zalacca, Roxb. Hort. Beng. 72; Fl. Ind. iii. 773 = Zalacca Wallichiana.

CALANCHOË, Pers. Syn. i. 445 (1805) = **Kalanchoë**, Adans. (Crassulac.).

CALANDARIUM, Juss. ex Steud. Nom. ed. II. i. 252 (1840) = sec.

CALANDRINIA, H. B. & K. Nov. Gen. et Sp. vi. 77. t. 526 (1823). *PORTULACEAE*, Benth. & Hook. f. i. 155.
BAITARIA, Ruiz & Pav. Prod. 63. t. 86 (1794).
CISTANTHE, Spach, Hist. Vég. Phan. v. 229 (1838).
CISMIA, Domb. ex Juss. Gen. 313 (1789).
DIAZIA, Phil. Fl. Atac. 23. t. 2 E (1880).
GEUNSIA, Moç. & Sesse, ex DC. Prod. iii. 358 (1828).
PHACOSPERMA, Haw. in Phil. Mag. i. (1827) 124.
RHODOPSIS, Lilja, Fl. Sverig. Suppl. i. 42 (1840).
TEGNERIA, Lilja, Fl. Sverig. edl. Vext. 76 (1839).
acaulis, *H. B. & K. Nov. Gen. et Sp.* vi. 78.—Ecuador.
acuta, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 340 = bifora.
adscendens, Hort. ex Phil. f. Pl. Vasc. Chil. 85 = pilosiuscula.
affinis, *Gill. ex Arn. in Chenk, Edinb. Journ.* iii. (1831) 358.—Chili.
alba, *DC. Prod.* iii. 359.—Peruv.
amoena, *Vis. in Nuov. Sagg. Padov.* vi. (1847) 94.—Hab. ?
andicola, *Gill. in Hook. Bot. Misc.* iii. (1833) 332.—Chili.
Andrewsii, *Sweet, Hort. Brit.* ed. II. 219.—Ind. occ.
Andrewsii, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 865 = Talinum patens.
angulata, Hort. ex Flach. & Mey. Ind. Sem. Hort. Petrop. xi. 53 = procumbens.
arenaria, *Cham. in Linnaea*, vi. (1831) 568.—Chili.
aurea, *Barn. in C. Gay, Fl. Chil.* ii. 510.—Chili.
axilliflora, *Barn. l. c.* 486.—Chili.
balonensis, *Lindl. in Mitch. Journ. Trop. Austral.* 148.—Austral.
Berteroana, *Phil. in Linnaea*, xxviii. (1856) 648.—Chili.
bifora, *Meyen, Reise*, i. 349.—Chili.
brevipedata, *F. Muell. Fragm.* x. 69.—Austral.
Breweri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 124.—Am. bor. occ.
Burdigi, *Hort. ex Regel, Gartenfl.* (1855) 4.—Am. austr.
cachinalensis, *Phil. Fl. Atac.* 20.—Chili.
caespitosa, *Gill. ex Arn. in Chenk, Edinb. Journ.* iii. (1831) 356.—Chili.
calycina, *Phil. in Linnaea*, xxxiii. (1864–65) 75.—Chili.
calyptoticha, *Phil. l. c.* xxx. (1859–60) 188.—Chili.
calyptrata, *Hook. f in Ic. Pl.* t. 296.—Austral.
capitata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 334.—Chili.
caulescens, *H. B. & K. Nov. Gen. et Sp.* vi. 78. t. 526.—Ecuador; Peruv.
Chamissoi, *Barn. in C. Gay, Fl. Chil.* ii. 497.—Chili.
chromantha, *Griseb. in Goett. Abh.* xxiv. (1879) 90.—Reg. Argent.
ciliata, *DC. Prod.* iii. 359.—Peruv.
cistiflora, *Gill. ex Arn. in Chenk, Edinb. Journ.* iii. (1831) 355.—Chili.
colchaguensis, *Barn. in C. Gay, Fl. Chil.* ii. 488.—Chili.
compacta, *Barn. l. c.* 481.—Chili.
composita, *Benth. Fl. Austral.* i. 176.—Austral.
compressa, *Schrad. ex DC. Prod.* iii. 359.—Chili.
conferta, *Gill. ex Arn. in Chenk, Edinb. Journ.* iii. (1831) 356.—Chili.
coquimbensis, *Barn. in C. Gay, Fl. Chil.* ii. 488.—Chili.
corrigioloides, *F. Muell. ex Benth. Fl. Austral.* i. 175.—Austral.
corymbosa, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 341.—Chili.
Cotyledon, *S. Wats. in Proc. Am. Acad.* xx. (1885) 355.—Calif.
Cuningii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 334.—Chili.

CALANDRINIA :—
curvifolia, Hort. ex Heynh. Nom. ii. 105 = procumbens.
densiflora, *Phil. in Linnaea*, xxviii. (1856) 644.—Chili.
densiflora, *Barn. in C. Gay, Fl. Chil.* ii. 508.—Chili.
densiflora, *Phil. in Linnaea*, xxviii. (1856) 642.—Chili.
denticulata, *Gill. in Hook. Bot. Misc.* iii. (1833) 333.—Chili.
dianthoides, *Phil. in Anal. Univ. Chil.* (1872) 717.—Chili.
diffusa, *Gill. ex Arn. in Chenk, Edinb. Journ.* iii. (1831) 355.—Chili.
discolor, Lindl. Bot. Reg. (1839) t. 4 = Lindleyana.
discolor, *Schrad. in Linnaea*, viii. (1833) Lit. 22.—Chili.
elegans, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 149 = discolor.
elegans, Spach, Hist. Vég. Phan. v. 232 = Menziesii.
erythrocoma, *Phil. in Linnaea*, xxxiii. (1864–65) 76.—Chili.
fasciculata, *Phil. l. c.* xxvii. (1856) 645.—Chili.
Fensiii, *Barn. in C. Gay, Fl. Chil.* ii. 493.—Chili.
ferruginea, *Barn. l. c.* 513.—Chili.
floribunda, *Phil. in Linnaea*, xxxiii. (1864–65) 78.—Chili.
frigida, *Barn. in C. Gay, Fl. Chil.* ii. 495.—Chili.
fruticulosa, Kunze, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 340 = trifida.
Gandichaudii, *Barn. in C. Gay, Fl. Chil.* ii. 490.—Chili.
Gayana, Barn. l. c. 503 = bifora.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 333.—Chili.
glandulosa, Hort. ex Steud. Nom. ed. II. i. 253 = procumbens.
glauca, Schrad. ex DC. Prod. iii. 359 = grandiflora.
glomerata, *Phil. in Linnaea*, xxviii. (1856) 644.—Chili.
gracilis, *Benth. Fl. Austral.* i. 173.—Austral.
graminifolia, *Phil. in Linnaea*, xxviii. (1856) 690.—Chili.
grandiflora, *Lindl. Bot. Reg.* t. 1194.—Chili.
granuliferа, *Benth. Fl. Austral.* i. 176.—Austral.
hirtella, *Phil. in Linnaea*, xxviii. (1856) 691.—Chili.
Landbeckii, *Phil. l. c.* xxxiii. (1864–65) 77.—Chili.
Leana, *Porter, in Coult. Bot. Bull.* i. (1876) 49.—Calif.
Lehmanni, *Endl. in Lehm. Pl. Preiss.* ii. 285.—Austral.
leucocephala, *Phil. Fl. Atac.* 21.—Chili.
leucotricha, *Phil. in Anal. Univ. Chil.* (1861) 91.—Chili.
Lindleyana, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 162 = discolor.
Lindleyana, *Walp. Rep.* ii. 335.—Hab. ?
linearifolia, *DC. Prod.* iii. 359.—Mexic. ?
lingulata, *DC. l. c.* 358.—Peruv.
lingulata, Hort. ex Steud. Nom. ed. II. i. 253 = procumbens.
liniflora, *Fensi, in Enum. Pl. Huag.* 52.—Austral.
litoralis, *Phil. Fl. Atac.* 20.—Chili.
Lloydii, Lond. Hort. Brit. Suppl. i. 616, sphalm. = Cuphea Llavea.
Lockhartі, *Sweet, Hort. Brit.* ed. II. 219.—Ins. Trinit.
longiscapa, *Barn. in C. Gay, Fl. Chil.* ii. 492.—Chili.
maritima, *Barn. l. c.* 483.—Chili.
maritima, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 197.—Calif.
megalyza, *Hemsl. Diag. Pl. Nov.* 28.—Guatemala.
Menziesii, *Torr. & Gray, Fl. N. Am.* i. 197.—Am. bor. occ.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 341.—Chili.
micrantha, *Schlecht. in Linnaea*, xiii. (1839) Lit. 97.—Mexic.
minima, *Bert. ex Steud. Nom.* ed. II. i. 253.—Chili.
minutissima, *Barn. in C. Gay, Fl. Chil.* ii. 504.—Chili.
modesta, *Phil. Fl. Atac.* 20.—Chili.
monandra, DC. Prod. iii. 359 = Monocosmia corrigioloides.
monticulata, *Meyen, Reise*, i. 346.—Chili.
multicaulis, *Phil. in Linnaea*, xxviii. (1856) 641.—Chili.
nevadensis, *A. Gray, in Proc. Am. Acad.* viii. (1878) 232.—Am. bor. occ.
nitida, *DC. Prod.* iii. 359.—Chili.

CALANDRINIA :—
oblongifolia, *Barn. in C. Gay, Fl. Chil.* ii. 482.—Chili.
oculta, *Phil. Fl. Atac.* 20.—Chili.
oppositifolia, *S. Wats. in Proc. Am. Acad.* xx. (1885) 355.—Calif.
paniculata, *DC. Prod.* iii. 358.—Peruv.
parviflora, Hort. ex Steud. Nom. ed. II. i. 253 = procumbens.
Phacosperma, *DC. Prod.* iii. 359.—Peruv.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 144.—Chili.
picta, Gill. ex Arn. in Chenk, Edinb. Journ. iii. (1831) 356 = arenaria.
pilosiuscula, *DC. Prod.* iii. 359.—Chili.
pleiopetala, *F. Muell. Fragm.* x. 70.—Austral.
Poeppigiana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 340.—Chili.
pogonophora, *F. Muell. Fragm.* x. 69.—Austral.
polyandra, *Benth. Fl. Austral.* i. 172.—Austral.
polycarpoides, *Phil. in Linnaea*, xxviii. (1856) 644.—Chili.
polyclados, *Phil. l. c.* xxxiii. (1864–65) 77.—Chili.
polypetala, *Fensi, in Enum. Pl. Huag.* 51.—Austral.
potentilloides, *Barn. in C. Gay, Fl. Chil.* ii. 512.—Chili.
procumbens, *Moris, in Mem. Acc. Tor.* xxxvii. (1884) 100.—Chili.
prostrata, *Phil. in Linnaea*, xxviii. (1864–65) 76.—Chili.
ptychosperma, *F. Muell. Fragm.* iv. 137.—Austral.
pulchella, Lilja, in Linnaea, xvii. (1843) 109.—Menziesii.
pumila, *F. Muell. Fragm.* x. 68.—Austral.
pusilla, *Barn. in C. Gay, Fl. Chil.* ii. 485.—Chili.
pusilla, *Lindl. in Mitch. Trop. Austral.* 360.—Austral.
pygmaea, *A. Gray, in Proc. Am. Acad.* viii. (1873) 623.—Am. bor. occ.
pygmaea, *F. Muell. Fragm.* i. 175.—Austral.
quadrivalvis, *F. Muell. l. c.* 176.—Austral.
racemosa, Steud. Nom. ed. II. i. 253 = arenaria.
ramosissima, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 334.—Chili.
rupestris, *Barn. in C. Gay, Fl. Chil.* ii. 485.—Chili.
salsoloides, *Barn. l. c.* 502.—Chili.
Saxifraga, *Barn. l. c.* 500.—Chili.
scrices, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 334.—Chili.
serrata, Poepp. ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 340 = Poeppigiana.
setosa, *Phil. in Linnaea*, xxviii. (1856) 690.—Chili.
speciosa, Lehm. in Linnaea, vi. (1831) Lit. 74 = discolor
speciosa, Lindl. Bot. Reg. t. 1360 = Menziesii.
spectabilis, *Otto & Distr. Allg. Gartenz.* i. (1888) 161.—Chili.
spergularina, *F. Muell. Fragm.* i. 175.—Austral.
spicata, *Phil. Fl. Atac.* 21.—Chili.
splendens, *Barn. in C. Gay, Fl. Chil.* ii. 499.—Chili.
strophiolata, *F. Muell. Fragm.* xi. 82.—Austral.
tenuifolia, Hook. Bot. Beech. Voy. 24 = pilosiuscula.
tenuifolia, *Phil. in Linnaea*, xxviii. (1856) 691.—Chili.
tricolor, *Phil. l. c.* 649.—Chili.
trifida, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 333.—Chili.
tuberosa, *Benth. Pl. Hartw.* 9.—Mexic.
umbellata, *DC. Prod.* iii. 358.—Chili.
umbellulata, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 334 = Gilliesii.
uniflora, *F. Muell. in Trans. Phil. Inst. Vict.* vi. (1862) 41.—Austral.
venulosa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 336 = arenaria.
venusta, Steud. Nom. ed. II. i. 253 = arenaria.
viridulans, Walp. Rep. iii. 326 = arenaria.
virgata, *Phil. in Linnaea*, xxviii. (1856) 649.—Chili.
volubilis, *Benth. Fl. Austral.* i. 174.—Austral.

CALANTHE, R. Br. in Bot. Reg. sub t. 379 (1821). *ORCHIDEAE*, Benth. & Hook. f. iii. 530.
AMBLYGLOTTIS, Blume, Bijdr. 368. t. 84 (1825).
CRYPTOPUS, Theo. Orch. Ins. Afr. t. 25, 26 (1822).
GHIESBREGHTIA, A. Rich. in Ann. Sc. Nat. Sér. III. iii. (1845) 28.
PREPTANTHUS, Reichb. f. in Fl. des Serres, Sér. I. viii. (1853) 245.
STYLOGLOSSUM, Breda, Orch. Kruid. et Hass. (1827).
ZEUXINE, Bech.-Hain. ex Meisn. Gen. Comm. 280 (1842).

CALANTHE :—
abbrevista, *Lindl. Gen. et Sp. Orch.* 251.—Java.
alismaefolia, *Lindl. Fol. Orchid. Calanth.* 8.—Reg. Himal.
alpina, *Hook. f. ex Lindl. l. c.* 4.—Reg. Himal.
alta, *Reichb. f. Otia Bot. Hamb.* 58.—Ins. Fiji.
anchorifera, *Reichb. f. in Gard. Chron.* (1888) ii. 166. —Polynesia.
angustifolia, *Reichb. f. in Linnaea*, xli. (1877) 76.—N. Caled.
angusta, *Lindl. Fol. Orphid. Calanth.* 7.—Reg. Himal.
angustifolia, *Lindl. Gen. et Sp. Orch.* 250.—Malaya.
aristulifera, *Reichb. f. in Bot. Zeit.* xxxvi. (1878) 74.—Japon.
australis, Ait. ex Lond. Hort. Brit. Suppl. ii. 615 = Spiranthes australis.
Barbersana x ; Reichb. f. in Gard. Chron. (1881) i. 136.
bella x ; Reichb. f. l. c. 234.
bicalar, Lindl. Sert. Orch. sub t. 9 = striata.
biloba, *Lindl. Fol. Orchid. Calanth.* 3.—Reg. Himal.
bracteosa, *Reichb. f. in Flora*, lxv. (1882) 533.—Ins. Samoa.
brevicornu, *Lindl. in Wall. Cat.* n. 7388.—Reg. Himal.
cattilligera, *Reichb. f. in Bonplandia*, v. (1857) 37.—Java.
Ceciliae, *Hort. Low.—Cf. Gard. Chron.* (1888) i. 492.—Malaya.
chloroleuca, *Lindl. Fol. Orchid. Calanth.* 10.—Reg. Himal.
citrina, *Scheidw. in L'Hortic. Belge*, v. (1838) 185. t. 104.—Hab.?
clavata, *Lindl. in Wall. Cat.* n. 7843.—Reg. Himal. ; Burma.
colorans, *Reichb. f. in Gard. Chron.* (1885) ii. 860.—Hab.?
comosa, Reichb. f. in Linnaea, xix. (1847) 874 = veratrifolia.
conspicua, *Lindl. Fol. Orchid. Calanth.* 4.—Malaya.
corymbosa, *Lindl. in Journ. Linn. Soc.* vi. (1862) 120. —Afr. trop.
curculigoides, *Lindl. in Wall. Cat.* n. 7340.—Malaya.
Curtisii, *Reichb. f. in Gard. Chron.* (1884) ii. 252. —Ins. Sondaic.
densiflora, *Lindl. in Wall. Cat.* n. 7344.—Reg. Himal.
dipteryx, *Reichb. f. in Gard. Chron.* (1884) ii. 394. —Ins. Sondaic.
discolor *Lindl. Sert. Orch.* sub t. 9.—Japon.
Dominii x ; Lindl. in Gard. Chron. (1858) 4.
emarginata, *Lindl. Gen. et Sp. Orch.* 249.—Java.
flava, *Morr. in L'Hortic. Belge*, ii. (1884) 238. t. 46.—Java.
Foerstermanni, *Reichb. f. in Gard. Chron.* (1888) i. 514.—Burma.
fulgens, Lindl. Fol. Orchid. Calanth. 10 = C. Masuca.
furcata, *Batem. ex Lindl. Bot. Reg.* (1838) *Misc.* 28. —Ins. Luzon.
galeata, Lindl. Fol. Orchid. Calanth. 5 = chloroleuca.
Gebina, Lodd. Cat. ex Lindl. l. c. 11 = Bletia hyacinthina.
gracilis, *Lindl. in Wall. Cat.* n. 7341.—Reg. Himal. ; China.
gracillima, *Lindl. Fol. Orchid. Calanth.* 8.—Ins. Carolin.
granatensis, *Reichb. f. in Bonplandia*, iv. (1856) 322. N. Granat.
grandiflora, *Nadeaud, Enum. Pl. Tahiti*, 37.—Ins. Pacif.
Griffithii, *Lindl. in Past. Fl. Gard.* iii. (1852–53) 87. —Reg. Himal.
herbacea, *Lindl. Fol. Orchid. Calanth.* 10.—Reg. Himal.
hololeuca, *Reichb. f. in Seem. Fl. Vit.* 298.—Ins. Fiji.
japonica, *Blume, in Mtq. Ann. Mus. Bot. Lugd. Bat.* ii. 205.—Japon.
Jumelli, *Bossil, ex Naves, in Blanco, Fl. Filip.* ed. III. *Nov. App.* 247.—Ins. Philipp.
labrosa, *Reichb. f. in Gard. Chron.* (1883) i. 44.—Burma.
Langei, *F. Muell. in Wing, South Sc. Record*, N. S. i. (1885).—N. Caled.
lentiginosa x, Reichb. f. in Gard. Chron. (1883) i. 44.
lycoglossa, *Reichb. f. Otia Bot. Hamb.* 58.—Afr. trop.
Mannii, *Lindl. in Wall. Cat.* n. 7337.—Ind. or. ; Malaya.
maculosa, *Reichb. f. in Linnaea*, xvii. (1844) 406.—Manil.
CALANTHE :—
natalensis, *Reichb. f. in Bonplandia*, iv. (1856) 322.—Afr. austr.
nivalis, *Boxall, ex Naves, in Blanco, Fl. Filip.* ed. III. *Nov. App.* 247.—Ins. Philipp.
occidentalis, Lindl. Fol. Orchid. Calanth. 3 = tricarinata.
odora, *Grif. Notul.* iii. 365.—Reg. Himal.
parviflora, *Lindl. in Past. Fl. Gard.* iii. (1852–53) 87, sub t. 61.—Java.
Perrottetii, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 68 = veratrifolia.
Petri, *Reichb. f. in Gard. Chron.* (1880) ii. 326.—Polynesia.
phajoides, Reichb. f. in Bonplandia, v. (1857) 37 = angustifolia.
plicata, *Lindl. Fol. Orchid. Calanth.* 11.—Sumatra.
plantaginea, Griff. Notul. iii. 365 = Griffithii.
plantaginea, *Lindl. in Wall. Cat.* n. 7346.—Reg. Himal.
pleiochroma, *Reichb. f. in Gard. Chron.* (1871) 938. —Japon.
porphyrea x , Reichb. f. l. c. (1884) i. 76.
proboscidea, *Reichb. f. l. c.* 476.—Ins. Sondaic.
puberula, *Lindl. in Wall. Cat.* n. 7342.—Reg. Himal.
pulchra, *Lindl. Gen. et Sp. Orch.* 250.—Java.
purpurea, *Lindl. l. c.* 249.—Zeylan.
reflexa, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 68.—Japon.
Regnieri, *Reichb. f. in Gard. Chron.* (1888) i. 274.—Cochinch.
rosea, *Benth. in Journ. Linn. Soc. Lond.* (1880) 309.—Burma.
Sandhamiastiana x ; Gosse, ex Reichb. f. in Gard. Chron. (1881) i. 395.
Sedeni x , Reichb. f. l. c. (1878) i. 168.
Sieboldii, *Decne. ex Regel, Ind. Sem. Hort. Petrop.* (1868) 80.—Japon.
speciosa, *Lindl. Gen. et Sp. Orch.* 250.—Java.
speciosa, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 44.—Caled.
Stevensiana, *Régnier, ex Reichb. f. in Gard. Chron.* (1888) i. 274, in nota.—Java ?
striata, *R. Br. in Bot. Reg.* 578, in nota.—China ; Japon.
sulphurea, *Hort. Hueg. ex Heynh. Nom.* ii. 105, nomen. —Ind. or.
tamaranae, *Blanco, ex Boerl. in Hasselt & Boerl. Bijdr. Midd. Sum.* 33 —Sumatra.
sylvatica, *Lindl. Gen. et Sp. Orch.* 250.—Madag.
sylvestris, Lindl. ex Steud. Nom. ed. II. i. 253 = sylvatica.
tahitiensis, *Nadeaud, Enum. Pl. Tahiti*, 37.—Ins. Tahiti.
Textori, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 204.—Japon.
triantheriformis, *Nadeaud, Enum. Pl. Tahiti*, 38.—Ins. Tahiti.
tricarinata, *Lindl. in Wall. Cat.* n. 7339.—Reg. Himal.
Turneri, *Reichb. f. in Gard. Chron.* (1883) i. 274. —Java ?
uncata, *Lindl. Fol. Orchid. Calanth.* 6.—Reg. Himal.
vaginata, *Lindl. l. c.* 7.—Assam.
variegata, *Scheidw. in L'Hortic. Belge*, v. (1838) 185. t. 103.—Hab.?
Veitchii x , Hort.—Cf. Gard. Chron. (1859) 1016.
Ventilabrum, *Reichb. f. in Seem. Fl. Vit.* 298.—Ins. Fiji.
veratrifolia, *R. Br. in Bot. Reg.* sub t. 573.—Ind. or. ; Malaya ; Austral.
veratrifolia, Hook. Bot. Beech. Voy. 71 = gracillima.
verecunda, Lindl. Sert. Orch. 42 = C. Masuca.
vestita, *Lindl. in Wall. Cat.* n. 7345.—Burma ; Malaya.
viridi-fusca, Hook. Bot. Mag. t. 4569 = Tainia viridifusca.
Wykehii, Reichb. f. in Walp. Ann. vi. 932 = C. Masuca.
Zollingeri, *Miq. Fl. Ind. Bat.* iii. 711.—Java.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 38.—Java.

CALANTHERA, Knuth (?), ex Hook. Kew Journ. vii. (1856) 18 = Stuhlbloß, Engelm. (Gramin.).
dactyloides, Knuth, l. c. = Buchloë dactyloides.

CALANTHUS, Gerst. in Videnak. Selsk. Ser. V. v. (1861) 191 ; (Gen. Centr. Am. 47. t. 4) = Alloploctus, Mart. (Gesneracˆ.).
multiflorus, Gerst. l. c. = A. multiflorus.

CALANTICA, Jaub. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 74. *SAMYDACEAE*, Benth. & Hook. f. i. 799.
cerasifolia, *Tul. l. c.* 76.—Madag.
grandiflora, *Jaub. ex Tul. l. c.* 75.—Madag.
Zalberti, Balli. Hist. Pl. iv. 377 = Bivinia Jaubertia.

CALASIAS, Rafin. Fl. Tellur. iv. 64 (1836) (nom. prius) = **Anisotes**, Nees (Acanthac.).
bracteata, Rafin. l. c. = A. trisulcus.

CALATHEA, G. F. W. Mey. Prim. Fl. Esseq. 6 (1818). *SCITAMINEAE*, Benth. & Hook. f. iii. 653.
ENDOCODON, Rafin. Fl. Tellur. iv. 49 (1836).
GOEPPERTIA, Nees, in Linnaea, vi. (1831) 837.
MONOSTICHE, Koern. in Regel, Gartenfl. (1858) 88.
PSYDARANTA ?, Neck. Elem. iii. 145 (1790).
ZELMIRA, Rafin. Fl. Tellur. iv. 50 (1836).
Ackermanni, *Koern. in Bull. Soc. Nat. Mosc.* xxxv (1862) i. 133.—Bras.
aemula, *Koern. l. c.* 131.—Bras.
albicans, *A. Brongn. ex Gris, in Ann. Sc. Nat.* Sér. IV. xii. (1859) 198.—Hab.?
affinis, *Fenzl, ex Regel, Gartenfl.* (1879) 294.—Bras.?
Allouya, Steud. Nom. ed. II. i. 253 = C. Alloula.
Allonia, *Lindl. Bot. Reg.* sub t. 1210.—Guiana.
altissima, *Horan. Prod. Scitam.* 13 ; *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 141.—Am. trop.
angustifolia, *Koern. in Regel, Gartenfl.* (1858) 87. —Am. trop.
applicata, *Y. Makoy, ex E. Morr. in Belg. Hortic.* xxiv. (1874) 228.—Bras.
argyraea, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 142.—Bras.
Arnaldia, *Koern. l. c.* 144.—Bras.
arrecta, *Lindl. & André, in Illustr. Hortic.* (1871) 174. t. 77.—Ecuador.
Backeniana, *E. Morr. in Belg. Hortic.* xxv. (1875) 271. f. 4.—Bras.
bambusacea, Poepp. & Endl. Nov. Gen. et Sp. ii. 93 = Ischnosiphon bambusaceus.
belle, *Regel, Gartenfl.* (1879) 297.—Hab.?
bellula, *Linden, Cat.* n. 89 (1872) 2.—Reg. Amazon.
bicolor, Steud. Nom. ed. II. i. 253 = Maranta bicolor.
blanda, *Stroud. l. c.*—Bras.
brasiliensis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 118.—Bras.
brevipes, *Koern. l. c.* 140.—Bras.
Cachibou, Lindl. ex Horan. Prod. Scitam. 12 = lutea.
capitata, *Lindl. Bot. Reg.* t. 1210.—Peruv.
Casupito, *G. F. W. Mey. Prod. Fl. Esseq.* 10.—Venezuela.
chrysantha, *Horan. Prod. Scitam.* 13.—Bras.
chimborazensis, *Linden, Illustr. Hortic.* (1870) t. 6.—Ecuador.
chlorosticta, Hort. Vind. ex Regel, Gartenfl. (1879) 294 = angustifolia.
chrysoleuca, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 142.—Peruv.
cinerea, *Regel, Gartenfl.* (1876) 2.—Reg. Amazon.
colorata, *Benth. & Hook. f. Gen.* iii. 654.—Bras.
comosa, *Lindl. Bot. Reg.* sub t. 1210.—Ind. occ.
composita, Klotzsch, in Schomb. Brit. Guian. iii. 918 = Ischnosiphon Parkeri.
confecta, *Benth. in Benth. & Hook f. Gen.* iii. 653.—Afr. trop.
crocata, *E. Morr. & Yorius. in Belg. Hortic.* xxv. (1875) 141. t. 3.—Bras.
densa, *Regel, Ind. Sem. Hort. Petrop.* (1866) 83.—Bras.
dictyphala, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 127.—Peruv.
discolor, *G. F. W. Mey. Prim. Fl. Esseq.* 7.—Guiana.
dubia, Lindl. Bot. Reg. sub t. 1210 = Ischnosiphon spicatus.
eximia, *Koern. ex Regel, in Gartenfl.* (1858) 87.—Am. trop.
exscapa, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 123.—Peruv.
fasciata, *Regel & Koern. in Gartenfl.* (1858) 848.—Bras.
fasciculata, Presl, Reliq. Haenk. i. 108. t. 16 = Phrynium fasciculatum.
flavescens, *Lindl. Bot. Reg.* t. 983.—Bras.

CALATHEA:—
Glaziovi, *Benth. in Benth. & Hook. f. Gen.* iii. 654.—Bras.
grandifolia, *Lindl. Bot. Reg.* t. 1210.—Bras.
hieroglyphica, *Linden & André, Illustr. Hortic.* (1878) 63. t. 128.—N. Granat.
Jagoriana, *Regel, Gartenfl.* (1879) 297.—Bras.
Lancea, Spreng. Syst. i. 8 = Ischnosiphon Arouma.
Kappleriana, *Koern. ex Horan. Prod. Scitam.* 12.—Bras.
Koernickeana, Horan. l. c. = brasiliensis.
Koernickiana, *Regel, Gartenfl.* (1874) 88.—Bras.
Kummeriana, *E. Morr. in Belg. Hortic.* xxv. (1875) 270. t. 15–17. t. 1.—Bras.
lanceolata, Korn. in Bull. Soc. Nat. Mosc. xxxv. (1862) 1. 184 = longifolia.
lateralis, *Lindl. Bot. Reg.* sub t. 1910.—Peruv.
latifolia, *Klotzsch, in Schomb. Faun. et Fl. Guian.* 918.—Guiana.
laxa, Poepp. & Endl. Nov. Gen. et Sp. ii. 22. t. 130 = Ischnosiphon plurispicatus.
Legrelliana, *Regel, Gartenfl.* (1879) 301.—Ecuador.
leopardina, *Regel, l. c.* (1877) 35. t. 898.—Bras.
leucocephala, D. Dietr. Syn. Pl. i. 7 = Ischnosiphon leucophaeus.
leuconeura, Hort. ex Nichola. Dict. Gard. i. 287 = Maranta leuconeura.
leucophaea, Poepp. & Endl. Nov. Gen. et Sp. ii. 21. t. 130 = Ischnosiphon leucophaeus.
leucostachys, *Hook. f. Bot. Mag.* t. 6905.—Costa Rica.
Lietzei, *E. Morr. in Belg. Hortic.* xxv. (1875) 273. f. 8.—Bras.
Lindeni, Wallis & André, Illustr. Hortic. (1871) 211. t. 82 = seq.
Lindeniana, *Wallis, in Belg. Hortic.* xvi. (1866) 200.—Peruv.
littoralis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 144.—Bras.
longibracteata, *Lindl. Bot. Reg.* t. 1020.—Bras.
longifolia, *Klotzsch, ex Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 134.—Bras.
lutea, *G. F. W. Mey. Prod. Fl. Esseq.* 10.—Venezuela.
maculosa, *Lindl. Bot. Reg.* sub t. 1210.—Bras.
macrostachya, *Griseb. in Goett. Abh.* xxiv. (1879) 385.—Reg. Argent.
macrostachya, Klotzsch, in Schomb. Faun. et Fl. Brit. Guian. 918 = Ischnosiphon obliquus.
Makoyana, *E. Morr. in Belg. Hortic.* xxii. (1872) 251. t. 94 & 95.—Bras.
Mandioccae, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 138.—Bras.
Mannii, *Benth. in Benth. & Hook. f. Gen.* iii. 653.—Afr. trop.
Mansonis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 119.—Bras.
marantina, C. Koch, Allg. Gartenz. (1857) 168 = C. Campto.
medio-picta, *Makoy, ex E. Morr. in Belg. Hortic.* xxiv. (1874) 238.—Bras.
metallica, *Koern. ex Regel, Gartenfl.* (1858) 88.—Hab. ?
micans, *Koern. l. c.* 87.—Peruv.
microcephala, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 135.—Peruv.
microstachya, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 135.—Peruv.
mirabilis, *Makoy, ex E. Morr. in Belg. Hortic.* xxiv. (1874) 238.—Bras.
modesta, *A. Brongn. ex Gris, in Ann. Sc. Nat.* Sér. IV. xii. (1859) 198.—Hab. ?
monophylla, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 144.—Bras.
Myrosma, *Koern. in Regel, Gartenfl.* (1858) 87.—Guiana.
nigro-costata, *Linden & André, Illustr. Hortic.* xx. (1873) t. 144.—N. Granat.
nobilis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 144.—Hab. ?
oblonga, *Koern. l. c.* 138.—Bras.
oliveria, Hort. ex Nichols. Dict. Gard. i. 288 = Makoyana.
Oppenheimiana, *E. Morr. in Belg. Hortic.* xxv. (1875) 271. f. 3.—Bras.
orbiculata, *Ledd. Bot. Cab.* t. 1879.—Ind. occ.
ornata, *Koern. in Regel, Gartenfl.* (1858) 87.—Am. trop.
ovata, *Lindl. Bot. Reg.* sub t. 1910.—Bras.
pachystachya, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 142.—Peruv.
pallida, *Linden & André, Illustr. Hortic.* xix. (1872) t. 101.—Peruv.

CALATHEA:—
pardina, Planch. & Linden, in Fl. des Serres, Sér. II. i. (1856) 53 = villosa.
Parkeri, Poepp. & Endl. Nov. Gen. et Sp. ii. 22 = Ischnosiphon Parkeri.
Pavonii, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 116.—Peruv.
pavonina, *C. Koch & Linden, Belg. Hortic.* xv. (1865) 99.—Bras.
peruviana, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 128.—Peruv.
picturata, *C. Koch & Linden, ex C. Koch, Wochenschr.* vi. (1868) 346.—Bras.
polyphylla, Poepp. & Endl. Nov. Gen. et Sp. ii. 23. t. 131 = Ischnosiphon polyphyllus.
Princeps, *Regel, Gartenfl.* (1879) 302.—Reg. Amazon.
propinqua, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 122.—Peruv.
pulchella, *Koern. in Regel, Gartenfl.* (1858) 87.—Bras.
pumila, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 134.—Bras.
Rindeliana, Paull, ex Regel, in Act. Hort. Petrop. iii. (1875) 11. 284 = Koernickiana.
rosea-picta, *Regel, Ind. Sem. Hort. Petrop.* (1869) 12.—Am. trop.
Rossii, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 144.—Bras.
rotundifolia, *Koern. l. c.* 125.—Hab. ?
rotundifolia, Poepp. & Endl. Nov. Gen. et Sp. ii. 23 = Ischnosiphon rotundifolius.
ruiberba, *Fenzl, in Regel, Gartenfl.* (1879) 294.—Bras. ?
Sellowii, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 120.—Bras.
smaragdina, *Linden & André, Illustr. Hortic.* (1870) 78.—Ecuador !
spicata, *Steud. Nom.* ed. II. i. 258.—Guiana; Bras.
stenolithica, *Miq. in Linnaea*, xxii. (1849) 79.—Surinam.
taenioza, *G. Joriss. in Belg. Hortic.* xxvi. (1876) 88. t. 5.—Bras.
tridactyla, *Koern. in Regel, Gartenfl.* (1858) 86.—Guiana.
tuberosa, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 104.—Bras.
tubispatha, *Hook. Bot. Mag.* t. 5542.—Bolivia.
umbrosa, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 137.—Bras.
undulata, *Regel, Illustr. Hortic.* xix. (1872) t. 98.—Ecuador.
Vandenheckei, *Regel, Ind. Sem. Hort. Petrop.* (1869) 14.—Hab. ?
variana, *C. Koch & Math. in App. Cat. Sem. Hort. Berol.* (1855) 12.—Am. trop.
variegata, *Koern. in Regel, Gartenfl.* (1858) 86.—Am. trop.
Veitchiana, *Veitch, ex Hook. f. Bot. Mag.* t. 5535.—Bolivia.
velutina, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 127.—Peruv.
vestita, *Baker, in Saund. Ref. Bot.* v. t. 311.—Bras.
villosa, *Lindl. Bot. Reg.* (1834) *Misc.* 61.—Guiana.
violacea, *Lindl. l. c.* sub t. 982; t. 962.—Bras.
virginalis, *Linden, ex Regel, Gartenfl.* (1879) 299.—Bras.
viridis, *Koern. l. c.* (1858) 88.—Hab. ?
Wallisi, *Regel, Ind. Sem. Hort. Petrop.* (1869) 14.—Reg. Amazon.
Warscewiczii, *Kobrn. in Regel, Gartenfl.* (1858) 87.—Am. trop.
Widgreni, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 117.—Bras.
Wioletae, *Makoy, ex E. Morr. in Belg. Hortic.* xxiv. (1874) 238.—Bras.
zebrina, *Lindl. Bot. Reg.* sub t. 1210.—Bras.
zingiberina, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 133.—Bras.

CALATHIANA, Delarb. Fl. Auv. ed. II. i. 28 (1800) = Gentiana, Tourn.
stenalis, Delarb. l. c. = G. nivalis.

CALATHINUS, Raf. Fl. Tellur. iv. 21 (1836) = Narcissus, Tourn. (Amaryllid.).
cernuus, Raf. l. c. = N. calathinus.
multiflorus, Raf. l. c. = N. Tenetta !
serotinus, Raf. l. c. = N. serotinus.

CALATHODES, Hook. f. & Thoms. Fl. Ind. i. 40 (1855). *RANUNCULACEAE*, Benth. & Hook. f. i. 7.
palmata, *Hook. f. & Thoms. l. c.* 41.—Reg. Himal.

CALATHOSTELMA, Franch. in Mem. Fl. Bras. vi. iv. 319 (1885). *ASCLEPIADEAE.*
ditassoides, *Fourn. l. c.*—Bras.

CALBOA, Cav. Ic. v. 51. t. 476 (1799) = **Ipomoea**, Linn. (Convolvul.).
globosa, Lindl. in Journ. Hort. Soc. v. (1850) 82 = I. Llaveana.
wrightia, Cav. Ic. v. 51. t. 476 = I. peduncularis.

CALCALIA, Krock. Fl. Siles. ii. 11. 881 = CACALIA, Linn. = **Adenostyles**, Cass. (Compos.).
alpina, Krock. l. c. = A. glabra.

CALCARUNIA, Rafin. Med. Fl. ii. 106 (1830) = **Pontedaria**, Linn.
hastata, Rafin. l. c. = P. cordata.

CALCATRIPPA, Heist. Syst. 8 (1748) = **Delphinium**, Linn. (Ranunculac.).

CALCEARIA, Blume, Bijdr. 417. t. 83 (1825) = **Coryanthes**, R. Br. (Orchid.).
fornicata, Blume, l. c. = Cor. fornicata.
picta, Blume, l. c. = Cor. picta.

CALCEOLARIA, Heist. Syst. 5 (1748) = **Cypripedium**, Linn. (Orchid.).

CALCEOLARIA, Linn. Mant. ii. 148 (1771). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 992. FAGELIA, Schwencke, in Verh. Bat. Gen. Roterd. i. (1774) 474. t. 13.
JOVELLANA, Ruiz & Pav. Fl. Per. i. 12 (1798).
adscendens, *Lindl. Bot. Reg.* t. 1915.—Chili.
alba, *Ruiz & Pav. Fl. Per.* i. 19. t. 27.—Chili.
alternifolia, *Cav. Ic.* v. 28. t. 445. f. 3.—Peruv.
ambigua, *Phil. in Linnaea*, xxix. (1857–58) 92.—Chili.
amplexicaulis, H. B. & K. Nov. Gen. et Sp. ii. 884.—Ecuador.
andina, *Benth. in DC. Prod.* x. 219.—Chili.
angustiflora, Hook. Bot. Mag. t. 3094 = verticillata.
angustifolia, *Ruiz & Pav. Fl. Per.* i. 17, 18. t. 28.—Ecuador.
angustifolia, Sweet, Hort. Brit. ed. II. 396 = integrifolia.
anomala, Pers. Syn. i. 16 = crenatiflora.
aquatica, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1855) 2.—Hab. ?
arachnoidea, *R. Grah. in Edinb. N. Phil. Journ.* (Sept. 1858) 371; *et Bot. Mag.* t. 5274.—Chili.
arucana, *Phil. in Linnaea*, xxix. (1857–58) 33.—Chili.
argentea, *H. B. & K. Nov. Gen. et Sp.* ii. 887. t. 170.—Peruv.
Athinsiana, Sweet, Brit. Flow. Gard. Ser. II. i. 188.
bertalafolia, *Wedd. Chlor. And.* ii. 134.—Bolivia.
bellidifolia, *Gill. ex Benth. in DC. Prod.* x. 207.—Chili.
bellidifolia, Salisb. Prod. 89 = Fothergillii.
Benthami, *Phil. in Anal. Univ. Chil.* (1873) 583.—Chili.
Bergii, *Hieron. in Bol. Acad. Nac. Cordoba*, iii. (1879) 383.—Reg. Argent.
Berterii, *Colla, in Mem. Acc. Tor.* xxxix. (1836) 138.—Chili.
bicolor, *Ruiz & Pav. Fl. Per.* i. 16. t. 25.—Peruv.
biennata, *Ruiz & Pav. l. c.* 15. t. 24.—Peruv.
bifida, Willd. ex Link, in Spreng. Jahrb. i. iii. (1830) 58 = graminifolia.
bifora, Lam. Ill. 59; Encyc. i. 556 = plantaginea.
bifora, Ruiz & Pav. Fl. Per. i. 16. t. 20 = ovata.
biplanatifida, *Phil. Pl. Atac.* 46.—Chili.
brachiata, Willd. ex Link, in Spreng. Jahrb. i. iii. (1830) 62 = floccosa.
brevistyla, *Clos, in C. Gay, Fl. Chil.* v. 173.—Chili.
bridgesii, *Kunze, in Linnaea*, xx. (1847) 35.—Chili.
calycina, *Benth. in DC. Prod.* x. 211.—Peruv.
cana, *Cav. Ic.* v. 27. t. 443. f. 2.—Chili.
candicans, Willd. ex Link, in Spreng. Jahrb. i. iii. (1830) 62 = argentea.
canescens, Humb. ex Link, l. c. 54 = hypophylla.

CALCEOLARIA:—
canescens, Willd. ex Roem. & Schult. Syst. Mant. i. 155.—Am. exact.
Cavanillesii, Pers. ex Steud. Nom. ed. I. 185 = Fothergilli.
Cavanillesii, *Phil. in Anal. Univ. Chil.* (1878) 582.—Chili.
cerasifolia, Benth. in DC. Prod. x. 218.—Peruv.
chelidonioides, H. B. & K. Nov. Gen. et Sp. ii. 378.—Ecuador.
chilensis, Lindl. Bot. Reg. t. 1476 = dentata.
chimborazensis, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 59 = mazsilla.
collina, Phil. in Linnæa, xxix. (1857–58) 80.—Chili.
concinna, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 51 = cheliodonioides.
connata, Hook. ex R. Grah. in Edinb. N. Phil. Journ. (Sept. 1838) 372 = petiolaris.
corymbosa, *Ruiz & Pav. Fl. Per.* i. 14. t. 20.—Chili.
crassifolia, *Phil. in Anal. Univ. Chil.* (1878) 582.—Chili.
crenata, Bot. Reg. t. 790 = adscendens.
crenata, *Lam. Encyc.* i. 556.—Ecuador.
crenatiflora, *Cav. Ic.* v. 26. t. 446.—Ins. Chiloe.
cuneiformis, *Ruiz & Pav. Fl. Per.* i. 19. t. 27.—Ecuador.
Cunninghamii, *Vathe, in Linnæa*, xl. (1876) 222.—Chili.
Darwinii, *Benth. in DC. Prod.* x. 207.—Reg. Magellan.
deflexa, *Ruiz & Pav. Fl. Per.* i. 18. t. 30.—Ecuador.
deflexa, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 52 = padifolia.
dentifolia, *Phil. in Linnæa*, xxxiii. (1864–65) 216.—Chili.
dentata, *Ruiz & Pav. Fl. Per.* i. 18. t. 29.—Chili.
denticulata, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 53 = stricta.
dichotoma, Lam. Illustr. i. 82 = ovata.
diffusa, Lindl. Bot. Reg. t. 1374 = bicolor.
dilatata, *Benth. in DC. Prod.* x. 211.—Ecuador.
divaricata, *H. B. & K. Nov. Gen. et Sp.* ii. 381.—Peruv.
elatior, *Griseb. in Gott. Abh.* xxiv. (1879) 237.—Reg. Argent.
elliptica, *Wedd. Chlor. And.* ii. 187.—Peruv.; Boliv.
ericoides, *Juss. ex Vahl, Enum.* i. 190.—Ecuador.
erioclada, *Benth. in DC. Prod.* x. 215.—Ecuador.
excelsa, *Hort. ex Court. in Mag. Hortic.* i. (1833) 64.—Hab.?
exoniensis x, Hort. Angl. ex Steud. Nom. ed. II. i. 254.
extensa, *Benth. in DC. Prod.* x. 214.—Peruv.
fasciculata, *Willd. ex Link, in Spreng. Jahrb.* i. III. (1820) 58.—Peruv.
ferruginea, *Cav. Ic.* v. 27. t. 445. f. 1.—Chili.
ferruginea, Colla, in Mém. Acc. Tor. xxxvii. (1885) 137. t. 46 = integrifolia.
ferruginea, Kunze, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 397 = integrifolia.
filicaulis, *Clos, in C. Gay, Fl. Chil.* v. 181.—Chili.
flexuosa, *Ruiz & Pav. Fl. Per.* i. 17, 26.—Peruv.
floribunda, *H. B. & K. Nov. Gen. et Sp.* ii. 383.—Ecuador.
floribunda, Hook. Bot. Mag. t. 4154 = crenata.
floribunda, Lindl. Bot. Reg. t. 1214 = petiolaris.
foliosa, *Griseb. in Gott. Abh.* xix. (1874) 210.—Reg. Argent.
foliosa, Meyen, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 396 = cuneiformis.
foliosa, *Phil. in Anal. Univ. Chil.* (1878) 581.—Chili.
Fothergilli, [*Saland. in*] *Ait. Hort. Kew.* ed. I. i. 80. t. 1.—Ins. Falkl.
fuchsiæfolia, *Hort.—Cf. Gard. Chron.* (1881) i. 269.—Hab.?
Gellendiana, Lindl. ex Steud. Nom. ed. II. i. 254 = sec.
Gellendiana, Penny, in Loud. Hort. Brit. Suppl. i. 568 = purpurea.
gunnellifolia, *Cav. Ic.* v. 30. t. 450. f. 1.—Peruv.
glabesia, *Phil. in Linnæa*, xxix. (1857–58) 81.—Chili.
glacialis, *Wedd. Chlor. And.* ii. 148.—Bolivia.
glandulosa, Benth. in DC. Prod. x. 210 = pseudoglandulosa.
glandulosa, *Poepp. ex Benth. l. c.*—Chili.
glauca, *Ruiz & Pav. Fl. Per.* i. 17. t. 27.—Peruv.
glutinosa, *Herr & Regel, Ind. Sem. Hort. Turic.* (1848) 4.—*Cf. Linnæa*, xxiv. (1851) 196.—Guatemala.

CALCEOLARIA:—
gnidiifolia, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 54 = lavandulæfolia.
gonnyana, *Benth. in DC. Prod.* x. 238.—N. Granat.
gracilis, *H. B. & K. Nov. Gen. et Sp.* ii. 379.—Ecuador.
graveolens, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 53 = ovata.
graminifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 386.—Ecuador.
guttata, *Willmore, ex Knowles & Westc. Flor. Cab.* i. (1837) 21.—Hab.?
Hartwegii, *Benth. Pl. Hartw.* 147.—Peruv.
helianthemoides, *H. B. & K. Nov. Gen. et Sp.* ii. 387 (err. typ. hellanthoides).—Peruv.
Henrici, *Hook. f. Bot. Mag.* t. 5772.—Ecuador.
Herbertiana, Lindl. Bot. Reg. t 1815 = racemosa.
heterophylla, *Ruiz & Pav. Fl. Per.* i. 16. t. 21.—Peruv.
heterophylla, Willd. Enum. Hort. Berol. 29 = scabiosæfolia.
hians, Penny, ex G. Don, Gen. Syst. iv. 603 = corymbosa ?
hispida, *Benth. in DC. Prod.* x. 212.—Peruv.
Hopeana, Morrison, ex G. Don, Gen. Syst. iv. 606 = corymbosa ?
Hoppeana, Morrison, ex Loud. Hort. Brit. Suppl. i. 583 = plantaginea.
hypericina, *Poepp. ex Benth. in DC. Prod.* x. 222.—Chili.
hypoleuca, *Benth. l. c.*—Ecuador.
hypoleuca, *Meyen, Reise*, i. 324; *Walp. & Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 396.—Chili.
hyssopifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 386.—Ecuador.
inflexa, Ruiz & Pav. Fl. Per. i. 16. t. 25 = serrata.
integrifolia, Linn. f. Suppl. 82 = ovata.
integrifolia, *Murr. Syst. ed.* XIII. 61.—Chili.
involuta, *Ruiz & Pav. Fl. Per.* i. 15. t. 23.—Ecuador.
Kellyana, *Hort.—Cf. Gard. Chron.* (1883) ii. 12.—Hab.?
knyperlensis, Hort. ex Steud. Nom. ed. II. i. 254 = crenatiflora.
lamiifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 383.—Ecuador.
lanata, *H. B. & K. l. c.*—Ecuador.
lanceolata, *Cav. Ic.* v. 27. t. 444. f. 2.—Patagon.
lanceolata, Link, in Spreng. Jahrb. i. III. (1820) 5 = lamiifolia.
Landbeckii, *Phil. in Linnæa*, xxxiii. (1864–65) 215.—Chili.
lanigera, *Phil. in Anal. Univ. Chil.* (1884) reimpr. 9.—Chili.
latifolia, *Benth. in DC. Prod.* x. 212.—Chili.
lavandulæfolia, *H. B. & K. Nov. Gen. et Sp.* ii. 386.—Ecuador.
laxa, *Benth. in DC. Prod.* x. 238.—Peruv.
linearis, *Ruiz & Pav. Fl. Per.* i. 19.—Peruv.
linifolia, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 54 = hyssopifolia.
lobata, *Cav. Ic.* v. 26. t. 448. f. 1.—Peruv.
Lorentzii, *Griseb. in Gott. Abh.* xix. (1874) 210.
Martinesiana x, Low, ex Sweet, Brit. Fl. Gard. Ser. II. t. 169.
Matthewsii, *Benth. in DC. Prod.* x. 207.—Peruv.
meliamœfolia, *Benth. l. c.* 214.—Peruv.
mendocina, *Phil. in Anal. Univ. Chil.* (1892) ii. 404.—Chili.
mexicana, *Benth. Pl. Hartw.* 47.—Mexic.
Meyeniana, *Phil. in Linnæa*, xxix. (1857–58) 82.—Chili.
mimuloides, *Clos, in C. Gay, Fl. Chil.* v. 174.—Chili.
mirabilis, Knowl. & Westc. Flor. Cab. i. (1837) 25 = crenatiflora.
mollissima, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 396 = integrifolia.
monanthos, Pohr. Encyc. Suppl. ii. 24 = sinuosa.
montana, *Cav. Ic.* v. 27. t. 444. f. 1.—Chili.
Morisii, *Walp. Rep.* iii. 167.—Chili.
Morrisoni, G. Don, in Sweet, Hort. Brit. ed. III. 521 = plantaginea.
mulliflora, *Cav. Ic.* v. 29. t. 449.—Peruv.
Nahuelbutæ, *Phil. in Anal. Univ. Chil.* (1878) 581.—Chili.
nana, *Sm. Ic. Ined.* i. t. 1.—Reg. Magellan.
Neuana, Spreng. Syst. i. 44 = Fothergilli.

CALCEOLARIA:—
nitida, Colla, in Mem. Acc. Tor. xxxix. (1886) 138 = adscendens.
nivalis, *H. B. & K. Nov. Gen. et Sp.* ii. 381.—Chili.
nivalis, Steud. & Hochst. ex Benth. in DC. Prod. x. 219 = adscendens.
nudicaulis, *Benth. l. c.* 208.—Chili.
nudicaulis, Meyen, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 397 = plantaginea.
obovata, Stokes, Bot. Comment. i. 119 = Fothergilli.
oblonga, *Ruiz & Pav. Fl. Per.* i. 15. t. 24.—Peruv.
obtusifolia, Kunze, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 397 = plantaginea.
oligantha, *Phil. in Linnæa*, xxxiii. (1864–65) 216.—Chili.
ovata, *Sm. Ic. Ined.* i. 3. t. 3.—Peruv.
padifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 380.—Peruv.
Paipæ, Steud. & Hochst. ex Benth. in DC. Prod. x. 219 = thyrsiflora.
panicea, Roem. & Schult. Syst. i. 181, sphalm. = punicea.
paniculata, Herb. ex G. Don, Gen. Syst. iv. 606 = petiolaris.
paniculata, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 58 = floribunda.
pannosa, *Phil. in Linnæa*, xxxiii. (1864–65) 219.—Chili.
paposana, *Phil. Fl. Atac.* 46.—Chili.
parælensis, Spreng. Syst. i. 45 = sec.
paralia, *Cav. Ic.* v. 28. t. 447.—Chili.
paralia, Hook. Exot. Fl. t. 75 = corymbosa.
pardanthera x, W. Baxt. ex Loud. Hort. Brit. Suppl. ii. 616.
parviflora, *Gill. ex Benth. in DC. Prod.* x. 207.—Chili.
parvifolia, *Wedd. Chlor. And.* ii. 184.—Boliv.
Pavonii, *Benth. in DC. Prod.* x. 211.—Peruv.
pendula, Sweet, Brit. Flow. Gard. Ser. II. 155 = crenatiflora.
perfoliata, *Linn. f. Suppl.* 86.—N. Granat.
perfoliata, Ruiz & Pav. Fl. Per. i. 15. t. 21. f. a = Pavonii.
petiolaris, *Cav. Ic.* v. 30, 451.—Ecuador.
petiolaris, Vahl, Enum. i. 186 = prænc.
picta, *Phil. in Linnæa*, xxix. (1857–58) 84.—Chili.
pinifolia, *Cav. Ic.* v. 26. t. 442. f. 2.—Chili.
pinnata, *Linn. Mant.* 71.—Peruv.
pinnata, Ruiz & Pav. Fl. Per. i. 14. t. 19 = scabiosæfolia.
placosmensis, *Meyen, Reise*, i. 469.—Peruv.
plantaginea, *Sm. Ic. Ined.* i. 2. t. 2.—Reg. Magellan.
plectranthifolia, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 394.—Peruv.
polifolia, *Hook. Bot. Mag.* t. 3897.—Chili.
polyantha, Penny, ex Loud. Hort. Brit. Suppl. i. 568 = petiolaris.
polyrrhiza, *Cav. Ic.* v. 25. t. 441.—Patagon.
pratensis, *Phil. in Linnæa*, xxix. (1857–58) 83.—Chili.
pseudoglandulosa, *Clos, in C. Gay, Fl. Chil.* v. 176.—Chili.
pulchella, Salisb. Prod. 89 = pinnata.
pulverulenta, *Ruiz & Pav. Fl. Per.* i. 19.—Peruv.
punctata, *Vahl, Enum.* i. 177.—Peruv.
punctiniculata, *Phil. in Linnæa*, xxix. (1857–58) 86.—Chili.
punicea, *Ruiz & Pav. Fl. Per.* i. 15. t. 24.—Peruv.
purpurea, *R. Grah. in Bot. Mag.* t. 3775.—Peruv.
quadrindiata, *Phil. in Linnæa*, xxix. (1857–58) 81.—Chili.
racemosa, *Cav. Ic.* v. 29. t. 448.—Patagon.
refugens, Sweet, Brit. Flow. Gard. Ser. II. t. 227 = sanchoides.
repens, *Hook. f. Fl. Nov. Zel.* i. 187.—N. Zel.
robusta, A. Dietr. in Otto & Dietr. Allg. Gartenz. i. (1833) 78 = integrifolia.
romaniifolia, *Lam. Encyc.* i. 556.—Ecuador.
rotundifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 382.—Peruv.
rugosa, Hook. Exot. Fl. t. 99 = adscendens.
rugosa, Ruiz & Pav. Fl. Per. i. 19. t. 28; Lodd. Bot. Cab. t. 942 = integrifolia.
salicifolia, Colla, in Mem. Acc. Tor. xxxix. (1886) 139 = semilis ?
salicifolia, *Ruiz & Pav. Fl. Per.* i. 18. t. 19.—Peruv.
salsoloefolia, Pers. Syn. i. 17 = integrifolia.
saxatilis, *H. B. & K. Nov. Gen. et Sp.* ii. 383.—Ecuador.
scabiosæfolia, Nees, ex G. Don, Gen. Syst. iv. 606 = rugosa.
scabiosæfolia, Roem. & Schult. Syst. i. 187 = cheliodonioides.

CALCEOLARIA :—
scabiosæfolia, *Sims, Bot. Mag.* t. 2405.—Peruv.
scabra, *Ruiz & Pav. Fl. Chil.* i. 19. t. 29.—Peruv.
scapiflora, *Benth. in DC. Prod.* x. 206.—Peruv.
Segethi, *Phil. in Linnæa*, xxix. (1857–58) 29.—Chili.
serrata, *Lam. Encyc.* i. 555.—Icosand.
sessilis, *Ruiz & Pav. Fl. Per.* i. 18.—Chili.
sibthorpioides, *H. B. & K. Nov. Gen. et Sp.* ii. 382.—Peruv.
Sinclairii, *Hook. Ic. Pl.* t. 561.—N. Zel.
sparsiflora, *Kunze, in Linnæa*, xxiv. (1851) 298.—Boliv.
spectabilis, Kunze, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 396 = dentata.
Standishi x , Past. Mag. ix. t. 75.
stricta, *H. B. & K. Nov. Gen. et Sp.* ii. 380.—N. Granat.
tenella, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 76. t. 287.—Chili.
tenera, *Phil. in Linnæa*, xxix. (1857–58) 34.—Chili.
tenuis, *Benth. in DC. Prod.* x. 205.—Chili.
ternifora, *Cav. Ic.* v. 30. t. 450. f. 2.—Peruv.
tetragona, *Benth. in DC. Prod.* x. 218.—Peruv.
tetraphylla, *Phil. in Linnæa*, xxix. (1857–58) 35.—Chili.
teucrioides, *Griseb. in Gott. Abh.* xix. (1874) 212.—Am. austr.
thyrsiflora, *R. Grah. in Edinb. N. Phil. Journ.* (Sept 1828) 372 (err. typ. 272).—Chili.
tinctoria, Gill. ex G. Don, Gen. Syst. iv. 606 = arachnoidea.
tomentosa, *Ruiz & Pav. Fl. Per.* i. 15. t. 22.—Peruv.
trachelifolia, Mart. & Gal. in Bull. Acad. Brux. xii. (1845) 1pt. 2 = mexicana.
triandra, *Vahl, Enum.* i. 181.—Peruv.
trifida, *Ruiz & Pav. Fl. Per.* i. 17. t. 26.—Peruv.
trilobata, *Hemsl. Biol. Centr. Am. Bot.* ii. 489.—Guatemala.
tripartita, *Ruiz & Pav. Fl. Per.* i. 14. t. 20.—Peruv.
umbellata, *Wedd. Chlor. And.* ii. 142.—Boliv.
undulata, *Benth. in DC. Prod.* x. 212.—Chili.
uniflora, Lam. Tabl. Meth. t. 52 = nana.
unifiora, *Ruiz & Pav. Fl. Per.* i. 16. t. 20.—Peruv.
utricularioides, *Hook. ex Benth. in DC. Prod.* x. 218.—Peruv.
valdiviana, *Phil. in Linnæa*, xxix. (1857–58) 35.—Chili.
verbascifolia, *Phil. l. c.* 30.—Chili.
verbascifolia, *Bert. in 'Merc. Chil. 1829' in Bull. Férussac*, xx. (1830) 109, nomen.—Chili.
vernicosa, *Phil. in Linnæa*, xxix. (1857–58) 29.—Chili.
verticillata, *Ruiz & Pav. Fl. Per.* i. 17.—Ecuador.
violacea, *Cav. Ic.* v. 31. t. 452.—Chili.
virgata, *Ruiz & Pav. Fl. Per.* i. 20. t. 31.—Peruv.
viscosa, *Ruiz & Pav. l. c.* 15. t. 20.—Peruv.
viscosissima, *Lindl. Bot. Reg.* t. 1611.—Chili.
Volckmanni, *Phil. in Linnæa*, xxxiii. (1864–65) 218.—Chili.
Wheeleri, Sweet, Brit. Flow. Gard. Ser. II. t. 130 = corymbosa.
Youngii x , Lindl. Bot. Reg. t. 1448.

CALCEOLUS, [Tourn.] Adans. Fam. ii. 70 (1763) = **Cypripedium**, Linn. (Orchid.).
alternifolius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 62, 124 = C. Calceolus.
Marianus, Crantz, Stirp. Austr. ed. II. fasc. vi. 454 = C. Calceolus.

CALCITRAPA, Hall. Enum. Stirp. Helv. ii. 689 (1742) = **Centaurea**, Linn. (Compos.).
Adami, Schur, Enum. Pl. Transs. 409 = Cent. solstitialis.
aegyptiaca, Sweet, Hort. Brit. ed. I. 218, nomen = Cent. aegyptiaca.
altissima, Lam. Fl. Fr. ii. 81 = Microlonchus salmanticus.
aspera, Cass. in Dict. Sc. Nat. viii. 252 = Cent. aspera.
Balsamita, Sweet, Hort. Brit. ed. I. 218 = Cent. Balsamita.
benedicta, Sweet, l. c. = Carbenia benedicta.
brevispina, Moench, Meth. 568 = Microlonchus salmanticus.
Calcitrapa, Hill, Hort. Kew. 62 = Cent. Calcitrapa.
calcitrapoides, Hill. l. c. = Cent. calcitrapoides.
centauroides, Sweet, Hort. Brit. ed. I. 218 = Cent. centauricoides.
cicheracea, Lam. Fl. Fr. ii. 82 = Cent. Seridis.

CALCITRAPA :—
collina, Lam. Fl. Fr. ii. 88 = Centaurea collina.
conferta, Moench, Meth. 564 = Cent. melitensis.
cucullata, Moench, l. c. 563 = Cent. Crocodylium.
elongata, Moench, l. c. = Volutarella muricata.
eriophora, Moench, l. c. 564 = Cent. eriophora.
ferox, Moench, Meth. Suppl. 229 = Cent. ferox.
fusca, Moench, Meth. 564 = Cent. sicula.
Galactites, Lam. Fl. Fr. ii. 80 = Galactites tomentosa.
heterophylla, Moench, Meth. Suppl. 230 = Cent. aspera.
Hippophaestrum, Gaertn. Mey. & Scherb. Fl. Wett. iii. 178 = Cent. Calcitrapa.
hybrida, Sweet, Hort. Brit. ed. I. 218 = Cent. Calcitrapa.
Hypophaestum, Gaertn. Fruct. ii. t. 163 = Cent. Calcitrapa.
iberica, Trev. ex Spreng. Syst. iii. 406 = Cent. iberica.
lanceolata, Lam. Fl. Fr. ii. 84 = Cent. calcitrapoides.
lanuginosa, Steud. Nom. ed. II. i. 845 = Kentrophyllum lanatum.
lanuginosa, Lam. Fl. Fr. ii. 85 = Carbenia benedicta.
longifolia, Willk. & Lange, Prod. Fl. Hisp. ii. 141, sphalm. = Cent. scabiosifolia.
lutea, Delarb. Fl. Auv. ed. II. 190 = Cent. solstitialis.
myacantha, Cass. in Dict. Sc. Nat. viii. 251 = Cent. Calcitrapa.
napifolia, Moench, Meth. 564 = Cent. napifolia.
nudicaulis, Lam. Fl. Fr. ii. 80 = Serratula nudicaulis.
parviflora, Lam. l. c. 82 = Cent. aspera.
patibilarensis, H. B. & K. Nov. Gen. et Sp. iv. 28 = Cent. americana.
phœnicea, Sweet, Hort. Brit. ed. I. 218 = Cent. calcitrapoides.
pubescens, Sweet, l. c. = Cent. incana.
reflexa, Sweet, l. c. = Cent. reflexa.
rigida, Sweet, l. c. = Cent. rigida.
rupestris, Sweet, l. c. = Cent. rupestris.
sessiliflora, Lam. Fl. Fr. ii. 85 = Cent. melitensis.
solstitialis, Lam. l. c. 84 = Cent. solstitialis.
sonchifolia, Lam. l. c. 82; Moench, Meth. 564 = Cent. sonchifolia.
sordida, Sweet, Hort. Brit. ed. I. 218 = Cent. sordida.
stellaris, Hill, Herb. Brit. t. 76 = Cent. Calcitrapa.
stellata, Lam. Fl. Fr. ii. 84 = Cent. Calcitrapa.
subspinosa, Moench, Meth. Suppl. 228 = Cent. dilutа.
Verutum, Moench, Meth. 563 = Cent. Verutum.
vulgaris, Bernh. Syst. Verz. Erf. 158 = Cent. Calcitrapa.

CALCOA, Salisb. Gen. Pl. Fragm. 67 (1866) = **Geitonoplesium**, A. Cunn. (Liliac.).

CALDASIA, Lag. Amen. Nat. ii. 98 (1821) = **Oreomyrrhis**, Endl. (Umbellif.).
andicola, Lag. ex DC. Coll. Mém. v. t. 2 = O. andicola, *andicola*, Hook. f. in Hook. Ic. Pl. iii. t. 300 = O. andicola.
brachycarpa, Hook. f. l. c. nota = O. andicola.
charophylloides, DC. Prod. iv. 229 = O. andicola.
charophylloides, Lag. Amen. Nat. ii. 99 = O. andicola.
dasycoides, Hook. f. Fl. Antarct. ii. t. 101 = O. andicola.
eriopoda, DC. Coll. Mem. v. 60 t. 2 ; et Prod. iv. 229 = O. andicola.
eriopus, DC. Coll. Mém. v. 75 = O. andicola.
lasiopetala, Lag. ex DC. Prod. iv. 229 = O. andicola.

CALDASIA, Mutis, ex Caldas, in Seman. Nuov. Gran. II. 28 (1610) = **Helosis**, Rich. (Balanoph.).
cayennensis, Mutis, ex Steud. Nom. ed. II. i. 255 = H. guyanensis.
funiformis, Mutis, ex Eichl. in DC. Prod. xvii. 136 = H. guyanensis.

CALDASIA, Willd. Hort. Berol. t. 71 (1807) = **Bonplandia**, Cav. (Acanthac.).
heterophylla, Willd. l. c. = B. geminiflora.

CALDCLUVIA, D. Don, in Edinb. N. Phil. Journ. ix. (1830) 92. *SAXIFRAGEÆ*, Benth. & Hook. f. i. 653.
DIDYSERICA, Ser. in DC. Prod. iv. 8 (1830), paniculata, *D. Don, l. c.*—Chili.

CALDENBACHIA, Pohl, ex Nees, in DC. Prod. xi. 281 (1847) = **Stenandrium**, Nees (Acanthac.).
elegans, Pohl, ex Nees, l. c. 282 = S. Pohlii.
macrogranum, Nees, l. c. 282 = S. dulce.

CALDESIA, Parl. Fl. Ital. iii. 398 (1858) = **Alisma**, Linn.
parnassifolia, Parl. l. c. 399 = A. parnassifolium.

CALEA, Linn. Sp. Pl. ed. II. 1179 (1763). *COMPOSITÆ*, Benth. & Hook. f. ii. 390.
ALLOCARPUS, H. B. & K. Nov. Gen. et Sp. iv. 291, t. 405 (1820).
ALLOISPERMUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 130.
AMPHICALEA, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 411.
CALEACTE, R. Br. in Trans. Linn. Soc. xii. (1818) 109.
CALEBRACHYS, Cass. in Dict. Sc. Nat. lv. 277 (1828)
CALYDERMOS, Lag. Gen. et Sp. Nov. 24 (1816).
LEMMATIUM, DC. Prod. v. 669 (1836).
LEONTOPHTHALMUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 140.
MEYERIA, DC. Prod. v. 670 (1836).
MOCINNA, Lag. Gen. et Sp. Nov. 31 (1816).
OTEIZA, La Llave, Reg. Trim. Mex. (1832) 41.
STENOPHYLLUM, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 391 (1873).
TETRACHYRON, Schlecht. in Linnæa, xix. (1847) 744.
acaulis, *Baker, in Mart. Fl. Bras.* vi. III. 266.—Bras
aculeata, Labill. Nov. Holl. Pl. ii. 41. t. 185 = Cassinia aculeata.
albida, *A. Gray, in Proc. Am. Acad.* xv. (1880) 88 = Mexic.
Amellus, Linn. Sp. Pl. ed. II. 1179 = Salmea scandens.
angustifolia, Gardn. in Hook. Lond. Journ. Bot. vii (1848) 417 = multiplinervia.
angustifolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi III. 256.—Bras.
aspera, Jacq. Collect. ii. 290 = Melananthera deltoidea.
axillaris, *DC. Prod.* v. 673.—Mexic.
Berteriana, *DC. l. c.* 672.—N. Granat.
brachiata, *DC. l. c.* 673 —Panama.
cacomoides, *Less. in Linnæa*, v. (1830) 157 = Mexic.
Candolleana, *Baker, in Mart. Fl. Bras.* vi. III. 256 — Bras.
carolinians, *Walt. Fl. Carol.* 200 —Am. bor.
Clausseniana, *Baker, in Mart. Fl. Bras.* vi. III. 265.—Bras.
clematides, *Baker, l. c.* 262.—Bras.
collina, *Klatt, in Leopoldina*, xx. (1884) 95.—N. Granat.
cordata, *Lour. Fl. Cochinch.* ii. 488.—China.
cordifolia, Sw. Fl. Ind. Occ. iii. 1326 = jamaicensis.
coriacea, *DC. Prod.* v. 675.—Peruv.
crenata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 256 = Candolleana.
cuneifolia, *DC. Prod.* v. 674.—Bras.
cymosa, *Less. in Linnæa*, v. (1830) 158.—Bras.
denudata, *Klatt, in Leopoldina*, xx. (1884) 96.—Peruv.
discolor, *A. Gray, in Proc. Am. Acad.* xv. (1880) 88.—Mexic.
divaricata, *Benth. in Hook. Journ. Bot.* ii. (1840) 44.—Guiana.
divergens, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 262.—Bras.
elegans, *DC. Prod.* v. 674.—Mexic.
elongata, *Baker, in Mart. Fl. Bras.* vi. III. 255.—Bras.
eupatorioides, Gardn. in Hook. Lond. Journ. Bot. vii (1848) 417 = asperata.
ferruginea, *Baker, in Mart. Fl. Bras.* vi. III. 261.—Bras.
fruticosa, Benth. & Hook. f. Gen. ii. 390 = rotundifolia.
Gardneriana, *Baker, in Mart. Fl. Bras.* vi. III. 255.—Bras.
gemmicoides, DC. Prod. v. 672 = Gelasopappus gemmicoides.
glabra, *DC. l. c.* 674.—Bras.
Grayii, *Klatt, in Leopoldina*, xx. (1884) 96.—Peruv.
hispida, *Baker, in Mart. Fl. Bras.* vi. III. 261.—Bras.
hymenolepis, *Baker, l. c.* 265.—Bras.
hypericifolia, *Baker, l. c.*—Bras.
integrifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 206.—Mexic.
jamaicensis, *Linn. Sp. Pl. ed. II.* 1179.—Jamaica.
lantanoides, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 416.—Bras.
lanuginosissima, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 262.—Bras.

CALEA:—
Leontophthalmum, *Less. Syn. Comp.* 241.—Peruv.
leptophylla, Forst. f. Prod. 56 = Casinia leptophylla.
lobata, Sw. Prod. Veg. Ind. Occ. 113 = Neurolaena lobata.
longifolia, *Baker*, *in Mart. Fl. Bras.* vi. 111. 260.—Bras.
longifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 418 = multiplinervia.
manicata, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 306.—Mexic.
Martiana, *Baker*, *in Mart. Fl. Bras.* vi. 111. 256.—Bras.
martinicensis, *Hort. ex Steud. Nom.* ed. II. i. 255.—Ins. Martinic.
melissæfolia, *Baker*, *in Mart. Fl. Bras.* vi. 111. 257.—Bras.
microphylla, *Baker*, *l. c.* 259.—Bras.
montana, *Klatt*, *in Leopoldina*, xx. (1884) 96.—Peruv.
multiplinervia, *Less. in Linnæa*, v. (1830) 159.—Bras.
myrtifolia, *Benth. & Hook. f. Gen.* ii. 393; *Baker*, *in Mart. Fl. Bras.* vi. 111. 259.—Bras.
nitida, *Less. in Linnæa*, v. (1830) 158.—Bras.
oligocephala, *DC. Prod.* v. 678.—Bras
oppositifolia, Linn. Sp. Pl. ed. II. 1179 = Isocarpha oppositifolia.
oxylepis, *Baker*, *in Mart. Fl. Bras.* vi. 111. 254.—Bras.
parvifolia, *Baker*, *l. c.* 259.—Bras.
paucidentata, Sch. Bip. ex Benth. & Hook. f Gen. ii. 391 = myrtifolia.
peduncularis, *H. B. & K. Nov. Gen. et Sp.* iv. 205. t. 408.—Mexic.
pedunculosa, DC. Prod. v. 673 = uniflora.
perfoxa, *Klatt*, *in Leopoldina*, xx. (1884) 95.—N. Granat.
phyllolepis, *Baker*, *in Mart. Fl. Bras.* vi. 111. 260.—Bras.
pilosa, *Baker*, *l. c.* 257.—Bras.
pinifolia, Forst. f. Prod. 55. n. 288 = Ozothamnus pinifolius.
pinnatifida, *Banks*, *ex Steud. Nom.* ed. I. 136.—Bras.
platylepis, *Sch. Bip. ex Baker*, *in Mart. Fl. Bras.* vi. 111. 267.—Bras.
Poblana, *Sch. Bip. ex Baker*, *l. c.* 266.—Bras.
pranifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 294. t. 406.—N. Granat.
ramosissima, *Baker*, *in Mart. Fl. Bras.* vi. 111. 257.—Bras.
reticulata, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 416.—Bras.
rotundifolia, *Benth. & Hook. f. Gen.* ii. 390; *Baker*, *in Mart. Fl. Bras.* vi. 111. 253.—Bras.
rugosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 306.—Mexic.
sabaloides, *Hemsl. l. c.*—Mexic.
salmifolia, *Hemsl. l. c.*—Mexic.
scabertima, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 111. 255 = elongata.
scabriolia, *Benth. & Hook. f. Gen.* ii. 391.—Mexic.
scandens, *Griseb. in Goett. Abh.* xxiv. (1879) 199.—Reg. Argent.
scoparia, Linn. Syst. Nat. ed. XII. 234 = Baccharis scoparia.
senecioides, *Baker*, *in Mart. Fl. Bras.* vi. 111. 258.—Bras.
Senecionis, Benth. & Hook. f Gen. ii. 391 = prænc.
serrata, F. G. Dietr. Vollst. Lex. Gaertn. i. 642 = cordifolia.
serrata, *Less. in Linnæa*, v. (1830) 158.—Bras.
sessilifora, *Less. l. c.*—Mexic.
sessiliflora, Stokes, Bot. Mat. Med. iv. 172 = Pacourina edulis.
solidaginea, *H. B. & K. Nov. Gen. et Sp.* iv. 295.—Venezuela; Ind. occ.
spectabilis, Labill. Nov. Holl. Pl. i. 49. t. 166 = Apalochlamis spectabilis.
stenophylla, *Baker*, *in Mart. Fl. Bras.* vi. 111. 255.—Bras.
subrotunda, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 415 = lantanoides.
Suriani, Cass. in Dict. Sc. Nat. vi. Suppl. 32 = Neurolæna integrifolia.
terrifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 294.—N. Granat.
teucrifolia, *Baker*, *in Mart. Fl. Bras.* vi. 111. 259.—Bras.
tomentosa, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 418.—Bras.

CALEA:—
tomentosa, *A. Gray*, *in Proc. Am. Acad.* xv. (1880) 88.—Mexic.
trifida, Moench, Meth. 581 = jamaicensis.
uniflora, *Less. in Linnæa*, v. (1830) 159.—Bras.
urticæfolia, *DC. Prod.* v. 674.—Mexic.
verbenæfolia, *DC. l. c.* 678.—Am. trop.
villosa, *Sch. Bip. ex Baker*, *in Mart. Fl. Bras.* vi. 111. 262.—Bras.
Zacatechichi, *Schlecht. in Linnæa*, ix. (1884) 589.—Mexic.

CALEA, Sw. Prod. Veg. Ind. Occ. 113 (1788) = Neurolaena, R. Br. (Compos.).
lobata, Sw. l. c. = N. lobata.

CALEACTE, R. Br. in Trans. Linn. Soc. xii. (1818) 109 = Calea, Linn. (Compos.).
pinnatifida, R. Br. l. c. = Calea pinnatifida.
rotundifolia, Less. in Linnæa, v. (1830) 158 = Calea rotundifolia.
urticifolia, R. Br. in Trans. Linn. Soc. xii. (1818) 109 = Calea urticæfolia.

CALEANA, R. Br. Prod. 329 (1810). *ORCHIDEÆ*, Benth. & Hook. f. iii. 610.
CALEYA, R. Br. in Ait. Hort. Kew. ed. II. v. 204 (1813).
major, *R. Br. Prod* 329.—Austral.
minor, *R. Br. l. c.*—Austral.
nigrita, *Lindl. Swan Riv. App.* 54.—Austral.

CALEBRACHYS, Cass. in Dict. Sc. Nat. lv. 277 (1828) = Calea, Linn. (Compos.).
pedunculariis, Cass. l. c. = Calea peduncularis.

CALECTASIA, R. Br. Prod. i. 268 (1810); et in Flind. Voy. ii. 609. t. 9 (1814). *JUNCACEÆ*, Benth & Hook. f. iii. 866.
HYPIA, Presl, ex Hook. Bot. Mag. t. 3834 (1840).
cyanea, *R. Br Prod.* 264.—Austral.
grandiflora, Presl, in Lehm. Pl. Preiss. ii. 53 = cyanea.
intermedia, Sond. in Linnæa, xxviii. (1856) 222 = cyanea.

CALENDULA, Linn. Syst. ed. I (1735). *COMPOSITÆ*, Benth. & Hook. f. ii. 454.
CALTHA, Tourn. ex Adans. Fam. ii. 126 (1763).
aegyptiaca, *Desf. Tabl. Hort. Par.* ed. I. 100.—Reg. Mediterr. or. ; Mesopot.
algeriensis, Boiss. Diagn. Ser. II. vi. 106 = suffruticosa.
algeriensis, *Boiss. & Reut. in Boiss. l. c.* 109.—Algeria.
amœna, *Willd. Gard. Dict.* ed. VIII. n. 10.—Mexic.
amplexicaulis, Thunb. Prod. Pl. Cap. 164 = Tripteris amplexicaulis.
amplexifolia, Reichb in Flora, xiii. (1830) 852 = arvensis.
anemonefiora, *Brrit. ex DC. Prod.* vi. 454, nomen.—Hab. ?
arborescens, Jacq. Collect. iii. 167 = Tripteris arborescens.
arragonica, *Fisch. ex Steud. Nom.* ed. II. i. 256.—Hispan.
arvensis, Bieb. Fl. Taur. Cauc. iii. 596 = persica.
arvensis, Boiss. Voy. Bot. Espagne, ii. 338 = aegyptiaca.
arvensis, *Linn. Sp. Pl.* ed. II. 1308.—Europ. ; Orient.
aspera, Thunb. Prod. Pl. Cap. 164 = Osteospermum asperum.
Asterias, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 31 = stellata.
aurantiaca, Kotschy, ex Boiss. Diagn. Ser. I. vi. 96 = sinuata.
aurea, *Hort. ex Steud. Nom.* ed. II. i. 256.—Afr. austr.
Balansæ, *Boiss. & Reut. Diagn.* Ser. II. vi. 107.—Hispan.
bicolor, Rafin. Caratt. 82. n. 199 = arvensis.
brachypiana, Rupr. in Ind. Sem. Hort. Petrop. (1852) ; ex Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 230 = arvensis.
byzantina, DC. Prod. vi. 453 = arvensis.
carinata, *Boiss. Diagn.* Ser. I. x. 82.—As. Min.
ceratosperma, Viv. Fl. Lib. Spec. 59. t. 20 = palestina.
chrysanthemifolia, Vent Jard. Malm. 56 = Dimorphotheca chrysanthemifolia.
cornuta, Polr. Encyc. vii. 277 = stellata.

CALENDULA:—
Cristagalli, Viv. Fl. Lib. Spec 59. t. 26. f. 2 = sicula, stellata.
cuneata, Thunb. Prod. Pl. Cap. 164 = Dimorphotheca cuneata.
Dalgetiarum, Hort. ex DC. Prod. vi. 452 = marginata.
decumbens, *Mill. Gard. Dict.* ed. VIII. n. 9.—Mexic.
decurrens, Thunb. Prod. Pl. Cap. 163 = Dimorphotheca pluvialis.
dentata, Hort. ex Steud. Nom. ed. I. 136 = denticulata.
denticulata, *Schousb. in Willd. Enum. Hort. Berol.* 926.—Mauri.
diffusa, Salisb. Prod. 209 = Dimorphotheca fruticosa.
echinata, DC. Prod. vi. 453 = arvensis.
eriocarpa, *DC. l. c.*—Hab. ?
flaccida, Vent. Jard. Malm. 20 = Dimorphotheca aurantiaca.
Forskhlœana, Hort. ex Steud. Nom. ed. II. i. 256 = stellata.
fruticosa, Hort. ex DC. Prod. vi. 452 = suffruticosa.
fruticosa, Linn. Fl. Afr. Rar. 25 (Amœn. Acad. v. 25) = Dimorphotheca fruticosa.
fulgida, Rafin. Caratt. 88. n. 200 = suffruticosa.
glabrata, Thunb. Prod. Pl. Cap. 163 = Osteospermum glabratum.
glauca, *Salisb. Prod.* 209.—Hab. ?
gracilis, DC. Prod. vi. 453 = persica.
graminifolia, Linn. Sp. Pl. 922 = Dimorphotheca graminifolia.
grandiflora, *Hort. ex Steud. Nom.* ed. II. i. 256.—Afr. austr.
hispanica, Hort. ex DC. Prod. vi. 452 = suffruticosa.
hispida, Thunb. Prod. Pl. Cap. 164 = Aster hispidus.
hybrida, Linn. Sp. Pl. 921 = Dimorphotheca hybrida.
hybrida, Sweet, Brit. Flow. Gard. i. t. 89 = Dimorphotheca pluvialis.
incana, Willd. Sp. Pl. iii. 2341 = suffruticosa.
incanescens, *Schrad. ex Steud. Nom.* ed. II. i. 256.—Hab. ?
linifolia, *Wendl. ex DC. Prod.* vi. 454, nomen.—Hab. ?
lusitanica, Boiss. Diagn. Ser. I. x. 83 = suffruticosa.
maderensis, *DC. Prod.* vi. 454.—Ins. Madera.
magellanica, Willd. Sp. Pl. iii. 2344 = Lagenophora Commersonii.
malacitana, Boiss. & Reut. Pugill. (1852) 61 = aegyptiaca.
malvacarpa, *Pomel, Nouv. Mat. Fl. Atl.* 33.—Algeria.
marginata, Willd. Enum. Hort. Berol. 935 = suffruticosa.
maritima, *Guss. Ind. Sem. Hort. Boce.* (1825).—Sicilia.
maritima, Lowe, ex DC. Prod. vi. 454 = maderensis.
maroccana, *Ball, in Journ. Bot.* xi. (1873) 365 — Marocco.
micrantha, Boiss. & Noé, Diagn. Ser. II. vi. 111 = aegyptiaca.
micrantha, Tineo & Guss. Fl. Sic. Syn. ii. 874 = arvensis.
microcephala, Kral. ex Reichb. Ic. Fl. Germ. xv. 99. t. 891 = aegyptiaca.
microcephala, Lange, ex Ficalho, in Bol. Soc. Broter. i. (1883) 51 = suffruticosa ?
microphyla, *Lange, ex Ficalho, l. c.*—Hispan.
Monardi, *Boiss. & Reut. Diagn.* Ser. II. vi. 108.—Hispan.
monstrosa, Burm. f. Fl. Cap. Prod. 28 = Tripteris scariosa.
muricata, Thunb. Prod. Pl. Cap. 164 = Tripteris arborescens.
Notana, Boiss. Fl. Orient. iii. 416 = suffruticosa.
nudicaulis, Linn. Sp. Pl. 922 = Dimorphotheca nudicaulis.
officinalis, Bieb. in Bull. Soc. Nat. Mosc. (1838) 256 = persica.
officinalis, *Linn. Sp. Pl.* 921.—Europ. austr.
oppositifolia, Ait. Hort. Kew. iii. 272 = Dimorphotheca oppositifolia.
palestina, *Boiss. Diagn.* Ser. I. x. 83.—Palestin. ; Mesopotam.
parviflora, DC. partim, ex Boiss. Fl. Orient. iii. 419 = aegyptiaca.
parviflora, Rafin. Caratt. 88. n. 201 = stellata.
parviflora, Thunb. Prod. Pl. Cap. 163 = Oligocarpus calendalaceus.

CALENDULA:—
persica, *C. A. Mey. Verz. Pfl. Cauc.* 72.—Reg. Cancas.; Persia; Afghan.
hispanica, Thunb. Prod. Pl. Cap. 164 = Chrysanthemum bellieckioides.
platycarpa, Cos. in Bull. Soc. Bot. Fr. iii. (1856) 655 = stellata.
pluvialis, Linn. Sp. Pl. 921 = Dimorphotheca pluvialis.
pluvialis, Thunb. Fl. Cap. 704 = Dimorphotheca clandestina.
prolifera, Hort. ex Stend. Nom. ed. II. i. 256 = officinalis.
pumila, Forst. f. Prod. 57 = Lagenophora Commersonii.
pusilla, Thou. Fl. Trist. Acun. 40 = Lagenophora Commersonii.
repanda, Boiss. & Noé, Diagn. Ser. II. vi. 110 = palestina.
rigida, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 371 = Tripteris arborescens.
rosmarinifolia, Houtt. Handl. xi. 84 = Osteospermum polygaloides.
sancta, Linn. Sp. Pl. ed. II. 1304 = aegyptiaca ?
scabra, Berg. Desc. Pl. Cap. 310 = Dimorphotheca pluvialis.
sicula, DC. Prod. vi. 452 (pl. aegypt.) = aegyptiaca.
sicula, Cyrill. ex Balb. Hort. Taur. (1800) = stellata.
sicula, Poir. Encyc. vii. 377 = praec. ?
sinuata, *Boiss. & Gaill. Diagn.* Ser. II. vi. 109.—Syria; Persia.
spatulata, Hoffmgg. Verz. Pfl. Nachtr. iii. 24 = stellata.
stellata, *Cav. Ic.* i. 8. t. 5.—Reg. Mediterr.
stellata, Lowe, ex DC. Prod. vi. 454 = maderensis.
stipitata, Spreng. Syst. iii. 624 = Lagenophora Billardieri.
sublacerata, *Pomel, Nouv. Mat. Fl. Atl.* 84.—Algeria.
sublanata, Relchb. f. Ic. Fl. Germ. xv. 99. t. 891 = arvensis.
suffruticosa, *Vahl, Symb. Bot.* ii. 94.—Reg. Mediterr. occ.
thapsiascarpa, *Pomel, Nouv. Mat. Fl. Atl.* 84.—Algeria.
tomentosa, Desf. Fl. Atlant. ii. 305. t. 245 = suffruticosa.
tomentosa, Thunb. Prod. Pl. Cap. 163 = Tripteris tomentosa.
Tragus, Alt. Hort. Kew. iii. 371; Jacq. Hort. Schoenb. iii. 14. t. 258 = Dimorphotheca Tragus.
Tragus, Curt. Bot. Mag. t. 408 = Dimorphotheca aurantiaca.
tripterocarpa, *Rupr. in Ind. Sem. Hort. Petrop.* (1858); *ex Bull. Phys.-Math. Acad. Pétersb.* xiv. (1856) 231.—Mesopot.
undulata, J. Gay, ex Gaud. Fl. Helv. v. 417 = arvensis.
undulata, Raf. Caratt. 81. n. 198 = stellata.
versicolor, Salisb. Prod. 309 = Dimorphotheca hybrida.
viscosa, Andr. Bot. Rep. t. 412 = Dimorphotheca cuneata.

CALENFINA, Adans. Fam. ii. 428 (1763); DC. Syst. Veg. ii. 646. *CRUCIFERÆ*, Benth. & Hook. f. i. 96.
cochlearioides, Dum. Fl. Belg. 121 = Banias cochlearioides.
Corvini, *Desv. Journ. Bot.* iii. (1814) 158.—Europ. med.; Reg. Caucas.; Persia.
prostrata, Spreng. Syst. ii. 850 = Muricaria prostrata.

CALESIA, Raf. Princ. Somiol. 29 (1814) = seq.

CALESIAM, Adans. Fam. ii. 446 (1763) = **Odina**, Roxb. (Anacard.).

CALETIA, Baill. Etud. Gén. Euph. 558. t. 26 (1858) = **Microcanthium**, Desf. (Euphorbiac.).
discreticistima, Muell. Arg. in Linnaea, xxxii. (1863) 79 = Ferdanthus discreticistimus.
hexandra, Muell. Arg. in DC. Prod. xv. ii. 194 = M. hexandrum.
linearis, Muell. Arg. in Linnaea, xxxii. (1863) 79 = Ferdanthus orientalis.
microcanthoides, Baill. Etud. Gén. Euph. 554. t. 26. f. 1 = M. hexandrum.
orientalis, Baill. Adansonia, vi. (1866–66) 337 = Ferdanthus orientalis.

CALETIA:—
ovalifolia, Muell. Arg. in Linnaea, xxxiv. (1865–66) 55 = Pseudanthus ovalifolius.
Wilhelmi, F. Muell. ex Muell. Arg. in DC. Prod. xv. ii. 194 = Pseudanthus ovalifolius.

CALEYA, R. Br. in Ait. Hort. Kew. ed. II. v. 204 (1813) = **Calæana**, R. Br. (Orchid.).
major, R. Br. l. c. = Calæana major.
minor, Sweet, Hort. Brit. ed. I. 385 = Calæana minor.
Sullivani, F. Muell. in Meib. Chemiz (1883) 44.—Australia.

CALIBRACHOA, Cav. in Le Llave & Lex. Nov. Veg. Desc. fasc. ii. (1825) 8 = **Petunia**, Juss. (Solanac.).
mexicana, Lex. l. c. = P. parviflora.
procumbens, Cav. l. c. = P. viscidula ?

CALICERA, Cav. Ic. iv. 54. t. 358 (1797) = **Calycera**, Cav.

CALICOREMA, Hook. f. in Benth. & Hook. f. Gen. iii. 84 (1880). *AMARANTACEÆ*, Benth. & Hook. f. i. c.
capitata, *Hook. f. l. c.* 85.—Afr. austr.

CALICOTOME, Link, in Schrad. Neues Journ. ii. (1808) ii. 50 = **Calycotome**, Link (Legumin.).

CALIGULA, Klotzsch, in Linnaea, xxiv. (1851) 28 = **Agapetes**, G. Don (Vaccinac.).
odontocera, Klotzsch, l. c. = A. variegata.
pulcherrima, Klotzsch, l. c. = A. variegata.

CALIMERIS, Nees, Gen. et Sp. Ast. 225 (1833) = **Aster**, Tourn. (Compos.).
Alberti, Regel, Gartenfl. (1884) 130.—Turkestan.
altaica, Nees, ex Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop. viii. 52 = A. altaicus.
alyssoides, DC. Prod. v. 259 = A. alyssoides.
amplexifolia, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 182 = A. trinervius.
biennis, Ledeb. Fl. Ross. ii. 483 = Heteropappus hispidus.
biennis, Nees, Gen. et Sp. Ast. 231 = A. altaicus.
canescens, Nees, l. c. 225 = A. altaicus.
cassiarabica, G. Don, in Sweet, Hort. Brit. ed. III. 243 = Aster cassiarabicus
chinensis, Sch. Bip. in Flora, xxxv. (1852) 48 = A. altaicus.
ciliata, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 394 = A. altaicus.
ciliata, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 61 = A. altaicus.
cordata, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126, nomen.—Malaya.
coronata, Sch. Bip. in Flora, xxxv. (1852) 48 = A. altaicus.
distorta, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) ii. 153 = A. altaicus.
exilis, Lindl. ex DC. Prod. v. 259 = A. altaicus.
fastigiata, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126 = A. fastigiatus.
flexuosa, Royle, ex Lindl. in DC. Prod. v. 258 = A. Thomsoni.
hispida, Lindl. l. c. 259 = A. altaicus.
hispida, Nees, Gen. et Sp. Ast. 237 = Heteropappus hispidus.
incisa, DC. Prod. v. 258 = A. incisus.
indica, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 125 = Boltonia indica.
integrifolia, Turcz. in DC. Prod. v. 259.—Reg. Amur.
japonica, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126 = A. fastigiatus.
microcephala, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 169.
platycephala, Nees, Gen. et Sp. Ast. 236 = A. incisus.
rosea, DC. Prod. v. 260 = A. roseus.
tatarica, Lindl. ex DC. Prod. v. 259 = Heteropappus hispidus.

CALINEA, Aubl. Pl. Gui. i. 336. t. 131 (1775) = **Dolichocarpus**, Roland. (Dilleniac.).
scandens, Aubl. l. c. = D. Calinea.

CALINUX, Raf. in Med. Repos. New York, v. (1808) 352 = **Pyrularia**, Michx. (Santalac.).

CALIPEURURIA, Herb. in Bot. Reg. (1844) Misc. 87. *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 721. Hartwegiana, *Herb. l. c.*—N. Granat.
subdentata, Baker, in Bot. Mag. t. 6989 = Eucharis subedentata.

CALIPOGON, Raf. Atl. Journ. 148 (1832) = Calopogon, R. Br. (Orchid.).

CALISAVA, Hort. ex Pav. Nuev. Quinol. sub t. 5. p. 6 (1862); Howard, Quinol. E. Ind. 86 (1869) = Cinchona, Linn. (Rubiac.).
anglica ×, Hort. ex Howard, l. c. 87.
javanica, Hort. ex Howard, l. c. 86 = Cinch. Calisaya.

CALISPERMUM, Lour. Fl. Cochinch. 157 (1790) = **Bimbella**, Juss. (Myrsin.).
scandens, Lam. l. c. = E. Ribes.

CALISTA, Rafg. in Marb. Ges. Schrift. ii. 128 (1831) = CALLISTA, Lour. = **Dendrobium**, Sw. (Orchid.).

CALISTACHYA, Rafin. in Med. Repos. New York, v (1808) 352 = **Veronica**, Tourn. (Scrophular.).
alba, Rafin. l. c. = V. virginica.

CALIUS, Blanco, Fl. Filip. ed. I. 698 (1837) = **Streblus**, Lour. (Urticac.).
lactescens, Blanco, l. c. = S. asper.

CALIXNOS, Rafin. Fl. Tellur. iv. 62 (1836) (quid ?).
trinervius, Rafin. l. c. = Convolvulus trinervis; ? non Convolvulacea, Sm. Cheiv.

CALLA, Linn. Syst. ed. I (1735). *AROIDEÆ*, Benth. & Hook. f. iii. 989.
CALLARIA, Rafin. in Am. Monthly Mag. (1818) 267
HOUTTINIA, Neck. Elem. iii. 291 (1790).
PROVENSALIA, Adans. Fam. ii. 468 (1763).
æthiopica, Gaertn. Fruct. ii. 20. t. 84 = palustris.
æthiopica, Linn. Sp. Pl. 968 = Zantedeschia æthiopica.
ambigua, Salisb. Prod. 262 = Zantedeschia æthiopica
angustifolia, Jack, in Malay. Misc. i. (1820) n. 1. p. 24 = Chamaecladon angustifolium.
aromatica, Roxb. Hort. Beng. 65; Fl. Ind. iii. 513 = Homalomena aromatica.
Badian, Blanco, Fl. Filip. ed. I. 658 = Alocasia indica.
calyptrata, Roxb. Hort. Beng. 65; Fl. Ind. iii. 514 = Schismatoglottis calyptrata.
cordifolia, Stoken, Bot. Mat. Med. iv. 326 = palustris.
Dracontium, G. F. W. Mey. Prim. Fl. Esseq. 197 = Monstera pertusa.
Gaby, *Blanco, Fl. Filip.* ed. I. 659.—Ins. Philipp.
humilis, Jack, in Malay. Misc. i. (1820) n. 1. p. 22 = Chamaecladon humile.
maxima, *Blanco, Fl. Filip.* ed. I. 658.—Ins. Philipp
monstera, Blume, Cat. Gew. Buitenz. 62 = Anadendrum montanum *sec* Rhaphidophora montana.
moschata, Moench, Meth. 246 = Richardia elliptica.
nitida, Jack, in Malay. Misc. i. (1820) n. 1. p. 24 = Aglaonema nitidum.
oblongifolia, Roxb. Hort. Beng. 65; Fl. Ind. iii. 516 = Aglaonema marantifolium.
occulta, Lodd. Bot. Cab. t. xii. = Homalomena aromatica.
occulta, Lour. Fl. Cochinch. ii. 593 = Zantedeschia occulta.
oculata, Lindl. in Gard. Chron. (1859) 788 = Richardia melanoleuca.
orientalis, Linn. Sp. Pl. ed. II. 1378 = Biarum Bovei.
ovatifolia, Gilib. Exercit. ii. 462 = palustris.
palmæfolia, *Linn. Sp. Pl.* 968.—Europ.; As. bor.; Am. bor.
pertusa, Kunth, Syn. i. 199 = Monstera pertusa.
picta, Roxb. Hort. Beng. 65; Fl. Ind. iii. 516 = Aglaonema pictum.
polyphylla, *Blanco, Fl. Filip.* ed. I. 659.—Ins. Philipp.
rudicrana, Raf. ex Engl. in DC. Monog. Phan. ii. 377 = Philodendron deltoideum.
rubescens, Roxb. Hort. Beng. 65; Fl. Ind. iii. 515 = Homalomena rubescens.
sagittifolia, Michx. Fl. Bor. Am. ii. 187 = Peltandra alba.
spicuatris, Blume, Cat. Gew. Buitenz. 62 = Rhaphidophora angustifolia.
virginica, Michx. Fl. Bor. Am. ii. 187 = Peltandra virginica.
virosa, Roxb. Hort. Beng. 65; Fl. Ind. iii. 517 = Colocasia virosa.

CALLAEOCARPUS, Miq. Pl. Jungh. 13 (1851–55) = Castanopsis, Spach (Cupulif.).
rhamnifolius, Miq. Fl. Ind. Bat. Suppl. 858 = Cast. rhamnifolia.
sumatrana, Miq. Pl. Jungh. 14 = Cast. sumatrana.

CALLAELEPIUM, Karst. Fl. Colomb. ii. 123. t. 165 (1869 !) = Pimbristemma, Turcz. (Asclepiad.).
Warscewiczii, Karst. l. c. = P. Warscewiczii.

CALLANTHUS, Reichb. Consp. 44 (1828) = Watsonia, Mill. (Irid.).

CALLARIA, Rafin. in Am. Monthly Mag. (1818) 267 = Calla, Linn. (Aroid.).

CALLERVA, Endl. Gen. Suppl. iii. 104 (1843) = Millettia, Wight & Arn. (Legumin.).
tomentosa, Endl. l. c. = M. nitida.

CALLIACHYRIS, Torr. & Gray, in Journ. Bost. Soc. Nat. Hist. v. (1844) 110 = Layia, Hook. (Compos.).
Fremontii, Torr. & Gray, l. c. = L. Fremontii.

CALLIANASSA, Webb & Berth. Phyt. Canar. iii. 143 (1836–50) = Isoplexis, Lindl. (Scrophularin.).
canariensis, Webb & Berth. l. c. 144 = I. canariensis.
isabelliana, Webb & Berth. l. c. 145.—Ins. Canar.

CALLIANDRA, Benth. in Hook. Journ. Bot. ii. (1840) 138. *LEGUMINOSAE*, Benth. & Hook. f. i. 596.
ANNESLIA, Salisb. Parad. Lond. t. 64 (1807), nomen prius.
CLELIA, Cassar. Nov. Stirp. Dec. 88 (1842).
CODONANDRA, Karst. Fl. Colomb. ii. 48. t. 122 (1869 !).
abbreviata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 108.—Bras.
aculeata, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 541.—Ecuador.
acschymenoides, *Benth. in Mart. Fl. Bras.* xv. ii. 415.—Am. trop.
alba, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = portoricensis.
alternans, *Benth. in Trans. Linn. Soc.* xxx. (1875) 548. —Madag.
amazonica, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 94.—Bras.
angelica, *Benth. l. c.* 100.—Mexic.
angustifolia, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 539.—Am. trop.
aspleniodes, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = mertensioides.
axillaris, *Benth. in Mart. Fl. Bras.* xv. ii. 415.—Bras.
bella, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 110.—Am. trop.
bicolor, *Benth. in Hook. Journ. Bot.* ii. (1840) 130.— Bras.
Blanchetii, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 109.—Bras.
bombycina, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 538.—Am. trop.
bracteosa, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 104.—Bras.
brevicaulis, *Michell, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. vii. 61.—Paraguay.
brevipes, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.— Bras.
californica, *Benth. Bot. Voy. Sulph.* 14. t. 11.— Calif.
Callistemon, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 527.—Mexic.
calothyrsus, *Meisn. in Linnaea*, xxi. (1848) 251.— Guiana.
calycina, *Benth. in Mart. Fl. Bras.* xv. ii. 419.—Am. trop.
canescens, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 96.—Mexic.
capillata, *Benth. l. c.* 98.—Mexic.
carassana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 543.—Am. trop.
carbonaria, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 95.—N. Granat.
Chamaedrys, Engelm. ex A. Gray, Pl. Fendl. 39 = eriophylla.
chilensis, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 108.—Chili.

CALLIANDRA :—
clavellina, Karst. Fl. Colomb. i. 159. t. 79 = Pardiaei.
Codonandra, *Benth. in Trans. Linn. Soc.* xxx. (1875) 547.—Am. trop.
collettioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 180.—Cuba.
comosa, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 104.—Am. trop.
conferta, *Benth. in A. Gray, Pl. Wright.* i. 68.—Am. bor. occ.
coriacea, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 98 = emarginata.
corotensis, Karst. Fl. Colomb. ii. 41 = purpurea.
Coulteri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 353.—Am. bor. occ.
crassipes, *Benth. in Trans. Linn. Soc.* xxx. (1875) 555. —Am. trop.
Crugerii, *Griseb. Fl. Brit. W. Ind.* 224.—Ind. occ.
Cumingii, *Benth. in Hook. Journ. Bot.* ii. (1840) 140. —N. Granat.
cylindrocarpa, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 98 = Harrisii.
cynometroides, Bedd. Fl. Sylv. t. 317 = Inga cynometroides.
dasyantha, Walp. Rep. i. 926, sphalm. = dysantha.
decaptera, *Benth. in Trans. Linn. Soc.* xxx. (1875) 546.—Am. trop.
diademata, Lem. Jard. Fleur. iii. (1852–53) t. 305 = bicolor.
dysantha, *Benth. in Hook. Journ. Bot.* ii. (1840) 138. —Bras.
emarginata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 98.—Mexic.
eriophylla, *Benth. l. c.* 105.—N. Mexic.
expansa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 549. —Am. trop.
falcata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 97.—Am. cent.
fasciculata, *Benth. in Hook. Journ. Bot.* ii. (1840) 140. —Bras.
filipes, *Benth. l. c.* 139.—Am. trop.
foliolosa, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 110.—Am. trop.
foliosa, *Benth. l. c.*—Bras.
formosa, *Benth. l. c.* 98.—Am. trop.
Gardneri, *Benth. l. c.* 108.—Bras.
geminata, Benth. in Trans. Linn. Soc. xxx. (1875) 548 Pithecolobium geminatum.
glomerulata, *Karst. Fl. Colomb.* ii. 5. t. 108.—N. Granat.
glyphoxylon, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 539.—Ecuador.
gracilis, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 181.—Jamaica.
gracilis, *Klotzsch, ex Baker, in Saund. Refug. Bot.* t. 294.—Am. trop.
grandiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 139. —Am. trop.
Griffithii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 537. —Reg. Himal.
Guildingii, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 96.—Bras.
haematocephala, *Hassk. Retzia*, 216.—Hab. !
haematoma, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 108.—Am. trop.
Harrisii, *Benth. l. c.* 98.—Mexic.
herbacea, Engelm. ex A. Gray, Pl. Fendl. 39 = humilis.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 110. —Ins. Mascar.
hirsuta, *Benth. in Trans. Linn. Soc.* xxx. (1875) 554. —Mexic.
hirtiflora, *Benth. l. c.* 550.—Am. trop.
Hookeriana, *Schomb. in Linnaea*, xx. (1847) 754 = rigida.
Houstoni, *Benth. in Hook. Journ. Bot.* ii. (1840) 139. —Mexic.
humilis, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 103.—N. Mexic.
hymenaeoides, *Benth. in Trans. Linn. Soc.* xxx. (1875) 537.—Am. trop.
Hystrix, Griseb. Cat. Pl. Cub. 88 = Pithecolobium Hystrix.
Kunthii, Benth. in Hook. Journ. Bot. ii. (1840) 139 = grandiflora.
Lambertiana, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 100.—Mexic.

CALLIANDRA :—
lanata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 553.— Am. trop.
lasiopus, Griseb. Fl. Brit. W. Ind. 211 = Pithecolobium caulifloruin.
latifolium, Griseb. l. c. 225 = Pithecolobium latifolium.
laxa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 551.— Am. trop.
leptopoda, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 101.—Bras.
linearis, *Benth. l. c.* v. (1846) 103.—Bras.
longipes, *Benth. in Trans. Linn. Soc.* xxx. (1875) 538. —Bras.
longipinna, *Benth. l. c.* 546.—Am. trop.
macrocephala, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—Bras.
Magdalenae, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 102.—Am. trop.
malacophylla, *Benth. l. c.* iii. (1844) 100.—Mexic.
marginata, Benth. in Trans. Linn. Soc. xxx. (1875) 536 = Pithecolobium marginatum.
marginata, Griseb. Fl. Brit. W. Ind. 210 = formosa.
mertensioides, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 106.—Am. trop.
metrosideriflora, *Benth. l. c.* i. (1842) 527.—Mexic.
microphylla, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Bras.
mollissima, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 97.—Bras.
myriophylla, *Benth. l. c.* 111.—Bras.
obovalis, Benth. l. c. 101 = Pithecolobium lusorium.
obtusifolia, Karst. Fl. Colomb. ii. 41. t. 121 = purpurea.
Pacara, *Griseb. in Gott. Abh.* xxiv. (1879) 123.—Reg Argent.
pallens, Benth. in Hook. Lond. Journ. Bot. v. (1846) 102 = Pithecolobium albicans.
parviflora, *Benth. l. c.* iii. (1844) 112.—Bras.
Patrisii, *Sagot, in Ann Sc. Nat.* Sér. VI. xiii. (1882) 324.—Guiana.
pauciflora, *Griseb. Cat. Pl. Cub.* 284.—Cuba.
Peckoltii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 555. —Bras.
pedicellata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 102.—Ins. S. Doming.
portoricensis, *Benth. l. c.* 99.— Am. bor.; Ind. occ.
prehensilis, Wright, in Sauv. Fl. Cub. 85 = Pithecolobium prehensile.
prostrata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 554.—Peruv.
Purdiaei, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 104.—N. Granat.
purpurea, *Benth. l. c.* iii. (1844) 104.—Am. trop.
quadrangularis, Walp. Rep. i. 927 = tetragona.
reticulata, *A. Gray, Pl. Wright.* ii. 58.—N. Mexic.
revoluta, Griseb. Cat. Pl. Cub. 88 = Pithecolobium obovale.
rigida, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 103.—Guiana.
Rutenbergiana, *Vatke, in Bremen Abh.* vii. (1882) 250.—Madag.
Samaan, Griseb. Fl. Brit. W. Ind. 225 = Pithecolobium Samaan.
Sancti-Pauli, Hassk. Retzia, i. 214 = Tweedii.
Schottii, *Torr. ex S. Wats. in Proc. Am. Acad.* xx. (1885) 364.—Arizona.
scutellifera, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Am. trop.
Seemanni, *Benth. in Seem. Bot. Voy. Herald*, 116. t. 23.—Venezuela.
sessilis, *Benth. in Hook. Journ. Bot.* ii. (1840) 141.— Bras.
Spruceana, Benth. in Trans. Linn. Soc. xxx. (1875) 598 = Pithecolobium parvifolium.
squarrosa, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 104.—Am. trop.
stenophylla, *Wawra, in Oestr. Bot. Zeitschr.* xxix. (1879) 215.—Bras.
stipulacea, *Benth. in Hook. Journ. Bot.* ii. (1840) 137. —Guiana.
submervosa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 540.—Am. trop.
subsp̄icata, *Benth. l. c.* 556.—Am. trop.
surinamensis, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 105.—Guiana.
taxifolia, *Benth. l. c.* 104.—N. Granat.

CALLIANDRA:—
tenuiflora, *Benth. in Trans. Linn. Soc.* xxx. (1875) 547.—Am. trop.
tergemina, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 96.—Am. trop.
tetragona, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Am. trop.
tetraphylla, *Benth. in Trans. Linn. Soc.* xxx. (1875) 544.—Mexic.
Thouarsiana, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 865.—Madag.
trinervia, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 94.—Bras.
tubulosa, Benth. l. c. 101 = Pithecolobium Saman.
turbinata, *Benth. l. c.* 109.—Bras.
Tweedii, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—Bras.
umbellifera, *Benth. l. c.* 141.—Bras.
umbrosa, *Benth. in Benth. & Hook. f. Gen.* i. 597.—Ind. or.; Penang.
ungulata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = portoricensis.
vespertina, Benth. l. c. = portoricensis.
virgata, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—Bras.
viridiflora, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 98.—Bras.
viscidula, *Benth. l. c.* 109.—Bras.
Wendlandi, *Benth. in Trans. Linn. Soc.* xxx. (1875) 556.—Guatemala.
xalapensis, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 105 = laxa.

CALLIANIRA, Miq. Syst. Pip. 244 (1843) = **Piper**, Linn.
melastomoides, Miq. l. c. 345 = P. melastomoides.

CALLIANTHEMUM, C. A. Mey. in Ledeb. Fl. Alt. ii. 336 (1830). *RANUNCULACEÆ*, Benth. & Hook. f. i. 5.
anemonoides, Endl. ex Heynh. Nom. ii. 106 = rutæfolium.
bipinnatum, Dulac, Fl. Hautes-Pyr. 216 = rutæfolium.
cachemirianum, *Cambess. in Jacquem. Voy. Bot.* 5.—Reg. Himal.
coriandrifolium, Reichb. Fl. Germ. Excurs. 737 = rutæfolium.
Endlicheri, Walp. Rep. i. 33 = Oxygraphis polypetala.
pimpinelloides, Royle, Illustr. Bot. Himal. 45 (Hook. f. & Thoms. Fl. Ind. 36) = cachemirianum.
rutæfolium, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 336.—Sibir.
rutæfolium, Reichb. Fl. Germ. Excurs. 737 = Ranunculus anemonoides.
semlverticillatum, *Phil. in Linnæa*, xxxiii. (1864–65) 8.—Chili.

CALLIAS, Cass. in Dict. Sc. Nat. xxiv. 327 (1823) = KALLIAS, Cass. = **Heliopsis**, Pers. (Compos.).

CALLICARPA, Linn. in Act. Soc. Ups. (1741) 80. *VERBENACEÆ*, Benth. & Hook. f. ii. 1150.
AGANON, Rafin. Sylva Tellur. 161 (1838).
AMICTONIS, Rafin. l. c. (1838).
BUCCHARDIA, [Heist.] Duham. Arb. i. 111. t. 44 (1755).
ILLA, Adans. Fam. ii. 446 (1763).
JOHNSONIA, T. Dale, ex Mill. Gard. Dict. ed. VI. App. 73 (1763).
PORPHYRA, *Lour. Fl. Cochinch.* 89 (1790).
SPONDYLOCOCCA, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. viii. (1748) 218.
TOMEA, Linn. Diss. Dans. 5 (1747) (Amoen. Acad. i. 389).
aculeolata, *Schau. in DC. Prod.* xi. 642.—Ins. S. Deming.
acuminata, *H. B. & K. Nov. Gen. et Sp.* ii. 252.—N. Granat.
acuminata, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 394.—Ind. or.
acuminatissima, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 409 = Geunsia farinosa.
acutidens, *Schau. in DC. Prod.* xi. 645.—Cochinch.
adenanthera, K. Br. Prod. 513 = cana.
albida, *Blume, Bijdr.* 818.—Java.
americana, Blanco, Fl. Filip. ed. I. 517 = Blanconi.
americana, *Linn. Sp. Pl.* 111.—Am. bor.
americana, Lour. Fl. Cochinch. 70 = cana.

CALLICARPA:—
ampla, *Schau. in DC. Prod.* xi. 642.—Jamaica.
angusta, *Schau. l. c.*—Ins. Philipp.
arborea, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 567 = lanata.
arborea, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 390.—As. trop.
arborea, Wall. Cat. n. 1826, partim = vestita.
attenuata, Wall. l. c. n. 1825 = longifolia.
bicolor, Juss. in Ann. Mus. Par. vii. (1806) 77 = cana.
bicolor, F. Vill. in Blanco, Fl. Philipp. ed. III. Nov. App. 158 = Blanconi.
Blancoi, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 315.—Ins. Philipp.
Blumei, *Zoll. & Moritzi, Syst. Verz. Zoll.* 53.—Malaya.
Bonplandiana, Schult. Mant. iii. 59 = acuminata.
brevicepa, Hance, in Journ. Linn. Soc. xiii. (1873) 116 = brevipes.
brevipes, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 338.—China.
cana, Dalz. & Gibs. Bomb. Fl. 209 = lanata.
cana, Gamble, Darjeeling List, 60; Ind. Timbers, 258 = macrophylla.
cana, *Linn. Mant.* ii. 198.—As. trop.; Austral.
cana, Wall. Cat. n. 1824 = pedunculata et longifolia.
cinerea, *A. Rich. Fl. Cub. Fanerog.* ii. 145.—Cuba.
cordifolia, Ruiz & Pav. Fl. Per. i. 50. t. 57. f. a = Aegiphila Mutisii.
Cumingiana, Schau. in DC. Prod. xi. 644 = Geunsia Cumingiana.
cuspidata, Roxb. Hort. Beng. 83; Fl. Ind. i. 394 = pedunculata.
dentata, Roth, Nov. Pl. Sp. 81; Wall. Cat. n. 6819 = pedunculata.
dentata, Roxb. ex Wall. l. c. p. 87. n. 1824 = cana.
dentata, Wall. l. c. n. 6819, pro parte = longifolia.
dichotoma, Raeusch. Nom. ed. III. 87 = purpurea.
discolor, Willd. ex Steud. Nom. ed. II. i. 257 = Aegiphila arborescens.
erosicona, Schau. in DC. Prod. xi. 643 = cana.
farinosa, Roxb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 567 = lanata.
ferruginea, *Sw. Prod. Veg. Ind. Occ.* 81.—Jamaica.
formosana, *Rolfe, in Journ. Bot. xx.* (1885) 358.—Ins. Formosa.
fulva, *A. Rich. Fl. Cub. Fanerog.* ii. 146.—Ins. Cuba.
globiflora, *Ruiz & Pav. Fl. Per.* i. 49. t. 57. f. b.—Am. austr.
gracilis, *Steb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 154.—Japon.
hexandra, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 410 = Geunsia farinosa ?
Heynii, Roth, Nov. Fl. Sp. 93 = cana.
Horsfieldii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 217.—Java.
incana, Roxb. Hort. Beng. [10]; Fl. Ind. i. 393 = macrophylla.
integerrima, *Champ. ex Benth. in Hook. Kew Journ.* v. (1853) 135.—China.
integrifolia, Jacq. Enum. Pl. Carib. 13 = Aegiphila arborescens.
japonica, *Thunb. Fl. Jap.* 60.—Japon.
lanata, Gamble, Darjeeling List, 60 = vestita.
lanata, Lam. ex Steud. Nom. ed. I. 137 = Premna lanata (sp. ignot.).
lanata, *Linn. Mant.* ii. 291.—Ind. or.
lanata, *Schau. in DC. Prod.* xi. 644 = pedunculata.
lanceolaria, Roxb. Hort. Beng. [10]; Fl. Ind. i. 395 = longifolia.
latifolia, *Kopp. ex Span. in Linnæa*, xv. (1841) 880, nomen.—Ins. Timor.
lobata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 566.—Reg. Himal.
longifolia, *Lam. Encyc.* i. 563.—Malaya; Austral.
macrocarpa, Raeusch. Nom. ed. III. 87 = cana ?
macrophylla, *Vahl, Symb. Bot.* iii. 13. t. 58.—Ind. or.
magna, *Schau. in DC. Prod.* xi. 643.—Ins. Philipp.
micrantha, *Vidal, Phan. Cuming. Philipp.* 187.—Ins. Philipp.
Mimurensis, Hassk. Cat. Hort. Bog. Alt. 136 = japonica.
mollis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 156.—Japon.
mollis, Willd. ex Schau. in DC. Prod. xi. 644 = acuminata, *H. B. & K.*

CALLICARPA:—
nudiflora, Hook. & Arn. Bot. Beech. Voy. 206 = Reevesii.
oblongifolia, Hassk. Cat. Hort. Bog. Alt. 136 = longifolia.
paniculata, Lam. Encyc. i. 563 = Chilianthus arboreus.
parvifolia, *Hook. & Arn. Bot. Beech. Voy.* 205.—Mexic.
pedunculata, *R. Br. Prod.* 513.—Malaya; Austral.
pentandra, Roxb. Hort. Beng. 83; Fl. Ind. i. 395 = Geunsia farinosa.
pallocalyx, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 569.—Reg. Himal.
purpurea, *Juss. in Ann. Mus. Par.* vii. (1806) 67.—China.
Reevesii, *Wall. Cat.* n. 1830.—China.
reticulata, Poepp. ex Walp. Rep. iv. 72 = Petitia Poeppigii.
reticulata, *Sw. Prod. Veg. Ind. Occ.* 81.—Jamaica.
Rheedii, Kostel. Allg. Med.-Pharm. Fl. iii. 939 = cana.
rhynchophylla, *Miq. Fl. Ind. Bat.* ii. 888.—Malaya.
Roxburghiana, Schult. Mant. iii. 54 = longifolia.
Roxburghii, Wall. Cat. n. 1833 = macrophylla.
rubella, *Lindl. Bot. Reg.* t. 888.—Reg. Himal.; China.
Schlimii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 217.—N. Granat.
serrata, Moench, Meth. 468 = americana.
sessiliflora, Wall. Cat. n. 1827 = rubella.
sinensis, Hort. ex Steud. Nom. ed. II. i. 257 = cana.
subintegerrima, H. B. & K. Nov. Gen. et Sp. ii. 252 = acuminata, *H. B. & K.*
subpubescens, *Hook. & Arn. Bot. Beech. Voy.* 205.—Mexic.
sumatrana, Miq. Fl. Ind. Bat. ii. 888 = cana.
tectonæfolia, Wall. Cat. n. 1827 = arborea.
tenuiflora, Champ. ex Benth. in Hook. Kew Journ. v. (1853) 135 = rubella.
tiliæfolia, Teijsm. & Binn. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 569 = pedunculata.
tomentosa, Koen. ex Vahl, Symb. Bot. iii. 13 = macrophylla.
tomentosa, Lam. Encyc. i. 563 = cana.
tomentosa, Murr. Syst. Veg. ed. XIII. 158 = lanata
tomentosa, *Willd. Enum. Hort. Berol.* 158; *Hook. & Arn. Bot. Beech. Voy.* 205.—China.
Towes, Poir. Encyc. Suppl. ii. 32 = lanata.
triflora, *Lour. Fl. Cochinch.* 70.—Cochinch.
umbellata, *Lour. l. c.*—Cochinch.
velutina, *Presl, Bot. Bemerk.* 98.—Bras.
vestita, *Wall. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 567.—Reg. Himal.
viburnifolia, Salisb. Prod. 83 = americana.
villosa, Roxb. Hort. Beng. [10] = arborea.
villosa, *Vahl, Symb. Bot.* iii. 14.—Ind. or. ?
Wallichiana, Walp. Rep. iv. 125 = lanata.
Zollingeriana, Schau. in DC. Prod. xi. 640 = mollis.

CALLICEPHALUS, C. A. Mey. Verz. Pfl. Cauc. 66 (1831) = **Centaurea**, Linn. (Compos.).
nitens, C. A. Mey. l. c. = Cent. nitens.

CALLICHLAMYS, Miq. in Linnæa, xviii. (1844) 254. *BIGNONIACEÆ*, Benth. & Hook. f. ii. 1039.
riparia, *Miq. l. c.*—Panama.
rubiginosa, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 201 = riparia.
ruginervis, Miers, l. c. 201 = riparia.
sphendida, Miers, l. c. = riparia.

CALLICHLOË, Pfeiff. Nom. i. 586 (1871) = seq.

CALLICHLOEA, Spreng. ex Stend. Nom. ed. II. i. 257 (1840) = **Hillontus**, Humb. & Bonpl. (Gramin.).
elegans, Spreng. l. c. = E. elegans.

CALLICHROA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 31 (1835) = **Lagrea**, Kunth. & Am. (Compos.).
angustifolia, Walp. Rep. ii. 628 = L. platyglossa.
angustifolia, Walp. l. c. iii. 386 = Castilleja angustifolia.
Bradburii, Walp. l. c. = Castilleja Bradburii.
Douglasii, Torr. & Gray, Fl. N. Am. ii. 396 = L. Douglasii.
hirsuta, Walp. Rep. ii. 628 = L. platyglossa.
platyglossa, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 31 = L. platyglossa.

CALLICOCCA, Schreb. Gen. i. 126 (1789) = **Cephaëlis**, Sw. (Rubiac.).
alba, J. F. Gmel. Syst. 371 = Ceph. alba.
alata, J. F. Gmel. l. c. 372 = Ceph. alata.
glabra, J. F. Gmel. l. c. = Ceph. glabra.
guianensis, J. F. Gmel. l. c. = Ceph. involucrata.
Ipecacuanha, Brot. in Roem. Arch. Ili. iii. 455 = Psychotria Ipecacuanha.
mucosa, Steud. Nom. ed. II. i. 287 = Ceph. mucosa.
purpurea, J. F. Gmel. Syst. 371 = Ceph. purpurea.
tomentosa, J. F. Gmel. l. c. = Psychotria tomentosa.

CALLICOMA, Andr. Bot. Rep. t. 566 (1809). *SAXIFRAGEAE*, Benth. & Hook. f. i. 649.
CALYCOMIS, R. Br. in Flind. Voy. App. 549 (1814).
STUTZERIA, F. Muell. Fragm. v. 81 (1865).
Billardieri, *D. Don, in Edinb. N. Phil. Journ.* (Apr.–June 1830) 94.—Austral.
ferruginea, D. Don, l. c. = serratifolia.
serratifolia, *Andr. Bot. Rep.* t. 566.—Austral.
Stutzeri, *F. Muell. Fragm.* v. 84.—Austral.
ternata, *Montr. in Mém. Acad. Lyon*, x. (1860) 211.—N. Caled.

CALLICORE, Link, Handb. i. 198 (1829) = **Amaryllis**, Linn.
crocata, Link, l. c. = Hippeastrum rutilum.
reticulata, Link, l. c. = Hippeastrum reticulatum.
rosea, Link, l. c. = A. Belladonna.
rutila, Link, l. c. 194 = Hippeastrum rutilum.

CALLICORNIA, Bu.m. f. Prod. Fl. Cap. 24 (1768) = **Leysseria**, Linn. (Compos.)
gnaphaloides, Burm. f. l. c. = L. gnaphaloides.
tenuifolia, Burm. f. l. c.—Afr. austr.
trifora, Burm. f. l. c.—Afr. austr.

CALLICYSTHUS, Endl. Prod. Fl. Norf. 90 (1838) = **Vigna**, Savi (Legumin.).
catubilis, Endl. l. c.—Ins. Norfolk.

CALLIDRYNOS, Neraud, in Freyc. Voy. Bot. 29 (1829), nomen (Sapind.).

CALLIGLOSSA, Hook. & Arn. Bot. Beech. Voy. 356 (1840) = **Layia**, Hook. & Arn. (Compos.).
Douglasii, Hook. & Arn. l. c. = L. Calliglossa.

CALLIGONUM, Linn. Gen. ed. I. 345 (1737). *POLYGONACEAE*, Benth. & Hook. f. iii. 95.
CALLIPHYSA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 24 (1835).
GIBSONIA, Stocks, in Proc. Linn. Soc. i. (1848) 367.
PALLASIA, Linn. f. Suppl. 87 (1781).
PTEROCOCCUS, Pall. Reise, i. App. 788. t. 8 (1776); Reise, ii. App. 48 (1777).
TRAXILIRA, Rafin. Sylva Tellur. 161 (1838).
xanthopterum, *Borszcz. Aralo-Casp. Callig.* 84.—Reg. Aral. Casp.
anfractuosum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 487 (*Lehm. Rel.* 311).—As. centr.
arolense, *Borszcz. Aralo-Casp. Callig.* 81.—Reg. Aral. Casp.
asperum, Lour. Fl. Cochinch. 343 = Tetracera sarmentosa.
buconiadum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 485 (*Lehm. Rel.* 309).—As. centr.
Bungei, *Boiss. Fl. Orient.* iv. 999.—Persia.
Calliphysa, *Bunge, Del. Sem. Dorpat.* (1839) 8.—Reg. Casp.; Turkest.
caucasicum, Pursh, Fl. Am. Sept. ii. 370 = Atriplex canescens.
Capot-Medusæ, *Schrenk, Enum. Pl. Nov.* i. 9.—Soongaria.
colabrinum, *Borszcz. Aralo-Casp. Callig.* 86.—Reg. Aral. Casp.
comosum, *L'Hérit. in Trans. Linn. Soc.* i. (1791) 180. —Aegypt.; Arab.; Persia.
crinitum, *Boiss. Diagn. Ser.* II. iv. 77.—Baluchist.
crispum, *Bunge, Del. Sem. Dorp.* (1839) 8.—Hab.?
cristatum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 485 (*Lehm. Rel.* 309).—As. centr.
desertum, *Borszcz. Aralo-Casp. Callig.* 86.—Reg. Aral. Casp.
denticulatum, *Bunge, ex Boiss. Fl. Orient.* iv. 999.—Persia.
crinaceum, *Borszcz. Aralo-Casp. Callig.* 87.—Reg. Aral. Casp.

CALLIGONUM:—
eriopodum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 485, 493 (*Lehm. Rel.* 309, 310).—Turkest.
flavidum, *Bunge, Del. Sem. Dorp.* (1839) 8.—Hab.?
gobicum, Bunge, ex Meisn. in DC. Prod. xiv. 29 = mongolicum.
horridum, *Borszcz. Aralo-Casp. Callig.* 82.—Reg. Aral. Casp.
leucocladum, *Bunge, in Mém. Sav. Etr. Pétersb.* (1851) 486 (*Lehm. Rel.* 309).—As. centr.
macrocarpum, *Borszcz. Aralo-Casp. Callig.* 85.—Reg. Aral. Casp.
medusocephalum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69.—C. Caput-Medusæ.
microcarpum, *Borszcz. l. c.* 41.—Reg. Aral. Casp.
mongolicum, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 204.—Mongolia.
Murex, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 485, 486 (*Lehm. Rel.* 309, 310).—As. centr.
Pallasii, *L'Hérit. in Trans. Linn. Soc.* i. (1791) 180. —Reg. Casp.; Turkest.
Panderi, *Fisch. ex Sweet, Hort. Brit.* ed. II. 440, nomen. —Reg. Casp.
paniculatum, *Borszcz. Aralo-Casp. Callig.* 82.—Reg. Aral. Casp.
persicum, *Boiss. Fl. Orient.* iv. 999.—Persia.
platyacanthum, *Borszcz. Aralo-Casp. Callig.* 87.—Reg. Aral. Casp.
polygonoides, *Linn. Sp. Pl.* 530.—Armenia; Persia; Ind. bor. occ.
polygonoides, Pall. Reise, iii. 580 = C. Pallasia.
Rotula, *Bosscs. Aralo-Casp. Callig.* 85.—Reg. Aral. Casp.
rubiaceum, *Bunge, Del. Sem. Dorp.* (1839) 8.—Hab.?
songaricum, *Endl. Gen. Suppl.* iv. 11. 50.—Soongaria.
stenopterum, *Bunge, ex Boiss. Fl. Orient.* iv. 999.—Persia; Afghan.
tetrapterum, *Jaub. & Spach, Illustr.* v. 80. t. 471.—Mesopotam.
tripterum, Bunge, ex Boiss. Fl. Orient. iv. 999 = Bungei.

CALLIGONUM, Lour. Fl. Cochinch. 342 (1790) = **Tetracera**, Linn. (Dilleniac.).
asperum, Lour. l. c. = T. sarmentosa.

CALLILEPIS, DC. Prod. v. 671 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 339.
glabra, *DC. l. c.*—Afr. austr.
hispida, *DC. l. c.*—Afr. austr.
Lawreola, *DC. l. c.*—Afr. austr.
leptophylla, *Harv. in Harv. & Sond. Fl. Cap.* iii. 186. —Afr. austr.
salicifolia, *Oliver, in Hook. Ic. Pl.* t. 1482.—Afr. austr.

CALLIOPEA, D. Don, in Edinb. N. Phil. Journ. (1828–29) 309 = **Crepis**, Linn. (Compos.).
aurea, D. Don, l. c. = Crepis aurea.

CALLIOPSIS, Reichb. Ic. et Descr. Pl. t. 70 (1824) = **Coreopsis**, Linn. (Compos.).
Athinsoniana, Hook. Fl. Bor. Am. i. 310 = Cor. Atkinsoniana.
atropurpurea, Hort. ex Steud. Nom. ed. II. i. 258 = Cor. tinctoria.
basalis, A. Dietz. in Otto & Dietr. Allgem. Gartenz. iii. (1835) 330.—Am. bor.
bicolor, Reichb. Ic. et Descr. Pl. t. 70 = Cor. tinctoria.
cardaminefolia, DC. Prod. v. 568 = Cor. cardaminefolia.
Drummondi, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 315 = Cor. Drummondii.
fimbriata, Lange, Sem. Hort. Haun. Adn. (1866) 5.—Hab.?
nudata, Spreng. Syst. iii. 611 = Cor. nudata.
palmata, Spreng. l. c. = Cor. palmata.
rosea, Spreng. l. c. = Cor. rosea.
tinctoria, DC. Prod. v. 568 = Cor. tinctoria.

CALLIPELTIS, Stev. in Mém. Soc. Nat. Mosc. vii. (1829) 276 (Obs. Pl. Ross. 69). *RUBIACEAE*, Benth. & Hook. f. ii. 148.
aperta, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 110.—Aegypt.; Persia.
Cucullaria, *Stev. in Mém. Soc. Nat. Mosc.* vii. (1829) 276.—Orient; Reg. Mediterr.
microstegia, *Boiss. Fl. Orient.* iii. 84.—Persia.

CALLIPHYSA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 24 (1835) = **Calligonum**, Linn (Polygonac.).
junceum, Fisch. & Mey. l. c. = Calligonum Calliphysa.
polygonea, C. A. Mey. ex Borszcz. Aralo-Casp. Callig. 48 = Calligonum Calliphysa.

CALLIPRENA, Salisb. Gen. Pl. Fragm. 89 (1865) = **Allium**, Tourn. (Liliac.).
cernua, Salisb. l. c. = A. cernuum.

CALLIPRORA, Lindl. Bot. Reg. t. 1590 (1833) = **Brodiæa**, Sm. (Liliac.).
aurantica, Kellogg, in Proc. Calif. Acad. ii. (1863) 20 = B. ixioides.
flava, Hort. ex Steud. Nom. ed. II. i. 258 = B ixioides.
lutea, Lindl. Bot. Reg. t. 1590 = B. ixioides.

CALLIPSYCHE, Herb in Bot. Reg. (1842) Misc 49. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 731.
aurantiaca, *Baker, in Saund. Ref. Bot.* t. 107.—Ecuador.
auratum, Hort. ex Gard. Chron. (1871) 6 = praec.
cucrosioides, *Herb. in Bot. Reg.* (1842) *Misc.* 50.—Ecuador.
mexicana, M. Roem. Syn. Ensat. 155 = cucrosioides.
mirabilis, *Baker, in Saund. Ref. Bot.* t. 168.—Peruv.

CALLIRHOË, Nutt. in Journ. Acad. Philad. ii. (1821) 181. *MALVACEAE*, Benth. & Hook. f. i. 201.
MONOLIX, Rafin. Cat. 14 (1824).
NUTTALLIA, Bart. Fl. N. Am. ii. 74. t. 62 (1822).
alceoides, *A. Gray, Pl. Fendl.* 18.—Am. bor.
digitata, *Nutt. in Journ. Acad. Philad.* ii. (1821) 181. —Am. bor. occ.
involucrata, *A. Gray, Pl. Fendl.* 16.—Am. bor. occ.
linearifolia, *A. Gray, in Proc. Am. Acad.* xix. (1883) 74.—Am. bor.
macrorrhiza, *A. Gray, Pl. Fendl.* 18 = alceoides.
palmata, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 449 = involucrata.
Papaver, *A. Gray, Pl. Fendl.* 17.—Am. bor. occ.
pedata, *A. Gray, l. c.*—Am. bor. occ.
spicata, Kegel, Gartenfl. (1872) 291. t. 737 = Sidalcea malvaeflora.
triangulata, *A. Gray, Pl. Fendl.* 16.—Am. bor.
verticillata, Groenl. in Rev. Hortic. (1862) 171 = involucrata.

CALLIRRHOË, A. Gray, Pl. Fendl. 16 (1849) = praec.

CALLISACE, Fisch. in Hoffm. Gen. Umb. ed. II. 170 = **Angelica**, Linn. (Umbellif.).
canadensis, Hoffm. ex Steud. Nom. ed. I. 762, 138 = Peucedanum palustre
dahurica, Fisch. in Hoffm. Gen. Umb. ed. II. 170 = A. dahurica.
ruthenica, Fisch. l. c. = Peucedanum ruthenicum?
Schieferecki, Hoffm. l. c. = Peucedanum palustre.

CALLISEMA, Steud. Nom. ed. II. i. 258 (1840) = seq.

CALLISEMAEA, Benth. in Ann. Wien. Mus. ii. (1838) 105 = **Platypodium**, Vog. (Legumin.).
elegans, Walp. Rep. i. 803 = P. elegans.
grandiflora, Benth. in Ann. Wien. Mus. ii. (1838) 105 = P. grandiflorum.
pubescens, Benth. l. c. = P. elegans.
sericea, Benth. l. c. = P. elegans.
viridis, Walp. Rep. i. 803 = P. viride.

CALLISIA, Linn. in Loefl. It. Hisp. 305 (1758). *COMMELINACEAE*, Benth. & Hook. f. iii. 654.
HAPALANTHUS, Jacq. Stirp. Am. ii. t. 11 (1763).
ciliata, H. B. & K. Nov. Gen. et Sp. i. 261 = Tradescantia elongata.
ciliata, Pers. Syn. i. 56 = repens.
delicatula, Kunth, Enum. Pl. iv. 63 = umbellulata.
diffusa, Pohl, ex Seub. in Mart. Fl. Bras. iii. i. 251 = Tradescantia effusa.
glandulosa, Pohl, ex Seub. l. c. 255 = Dithyrocarpos glandulosus.
insignis, *C. B. Clarke, in Hemsl. Diagn. Pl. Mex.* 56. —Mexic.

CALLISIA :—
Martensiana, *C. B. Clarke, in DC. Monog. Phan.* iii. 813.—*Mexico.*
melandra, Wright, in Sauv. Fl. Cub. 159 = Tradescantia cordifolia.
monandra, Schult. f Syst. vii. 1179 = umbellulata.
orientalis, Bech.-Ham. ex Wall. Cat. n. 6522 = Ancilema Hamiltonianum.
repens, *Linn. Sp. Pl. ed.* II. 62.—Am. austr.
rosea, Pohl, ex Seub. in Mart. Fl. Bras. iii. t. 352 = Tradescantia Gaudichaudiana.
umbellulata, *Lam. Illustr.* t. 130. t. 85. f. 2.—Ins. Martinic.

CALLISTA, Lour. Fl. Cochinch. 519 (1790) = **Dendrobium**, Sw. (Orchid.).
amabilis, Lour. l. c.—Cochinch.

CALLISTA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 104 = **Brica**, Tourn.
acuminata, G. Don, Gen. Syst. iii. 813 = E. Massoni.
armata, G. Don, l. c. = E. Sparmanni.
Baudonia, G. Don, l. c. = E. Irbyana.
Bedfordiana, G. Don, l. c. 814 = E. Massoni.
blanda, G. Don, l. c. 815 = E. blanda.
Broadleyana, G. Don, l. c. 813 = E. Broadleyana.
buccinæflora, G. Don, l. c. = E. Massoni.
calostema, G. Don, l. c. 815 = E. Massoni.
Celsiana, G. Don, l. c. 813 = E. Thunbergii.
Clifftoniana, G. Don, l. c. 815 = E. tenuiflora.
comata, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. comosa.
Comptoniana, G. Don, Gen. Syst. iii. 815 = E. curviflora.
Cooreyvana, G. Don, l. c. 815 = E. pavettæflora.
daphnifora, G. Don, l. c. = E. daphnifora.
daphnoides, G. Don, l. c. = E. præstans.
denticulata, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. denticulata.
fastigiata, G. Don, Gen. Syst. iii. 814 = E. fastigiata.
ferruginea, G. Don, l. c. 813 = E. squarrosa.
fragrans, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. fragrans.
Humeana, G. Don, Gen. Syst. iii. 813 = Erica fastigiata.
hyacinthoides, G. Don, l. c. 814 = E. præstans.
infiata, G. Don, l. c. = E. infiata.
infundibuliformis, G. Don, l. c. 813 = E. pavettæflora.
Juliana, G. Don, l. c. = E. Walkeriana.
Lawsoniana, G. Don, l. c. = E. Lawsoniana.
magnifica, G. Don, l. c. 813 = E. Massoni.
matulæflora, G. Don, l. c. = E. Massoni.
moschata, G. Don, l. c. 815 = E. Muscari.
mundula, G. Don, l. c. = E. fastigiata.
Muscari, G. Don, l. c. 814 = E. Muscari.
nidiflora, G. Don, l. c. 815 = E. Muscari.
Parmentieri, G. Don, l. c. = E. præstans.
pavettæflora, G. Don, l. c. = E. pavettæflora.
pellucida, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. Walkeriana.
pellucida, G. Don, Gen. Syst. iii. 814 = E. venusta.
prægnans, G. Don, l. c. 813 = E. ventricosa.
præstans, G. Don, l. c. 815 = E. præstans.
primuloides, G. Don, l. c. 816 = E. fastigiata.
quadrangularis, G. Don, l. c. = E. Massoni.
rigida, G. Don, l. c. 815 = E. Massoni.
Russeliana, G. Don, l. c. = E. Russeliana.
Saundersiana, G. Don, l. c. 813 = E. infiata.
stellifera, G. Don, l. c. 815 = E. Massoni.
struthiolæflora, G. Don, l. c. = E. tereticaula.
Swainsoniana, G. Don, l. c. 816 = E. Massoni.
Templana, G. Don, l. c. 814 = E. Massoni.
tenuiflora, G. Don, l. c. 812 = E. tenuiflora.
tetragona, G. Don, l. c. = E. tetragona.
thymifolia, Steud. Nom. ed. II. i. 788 = E. thymifolia.
truncata, G. Don, Gen. Syst. iii. 815 = E. Muscari.
undulata, G. Don, l. c. 815 = E. Massoni.
varia, G. Don, l. c. = E. varia.
ventricosa, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. ventricosa.
venusta, G. Don, Gen. Syst. iii. 815 = E. venusta.
verecunda, G. Don, l. c. = E. Meurori.
Walkeria, G. Don, l. c. 814 = E. Walkeriana.

CALLISTACHYA, Rafin. in Med. Repos. New York v. (1808) 60 = **Veronica**, Tourn. (Scrophular.).
sibirica, Rafin. l. c. = V. sibirica.
virginica, Rafin. l. c. = V. virginica.

CALLISTACHYA, Sm. in Trans. Linn. Soc. ix. (1808) 267 = CALLISTACHYS, Vent.

CALLISTACHYS, Henffel, in Flora, xxvii. (1844) 528 = **Carex**, Linn. (Cyperac.).
pyrenaica, Henffel, l. c. = C. pyrenaica.

CALLISTACHYS, Vent. Jard. Malm. 115. t. 115 (1803) = **Oxylobium**, Andr. (Legumin.).
carinata, Meissn. in Lehm. Pl. Preiss. i. 97 = O. carinatum.
cuneata, Heynh. Nom. i. 148 = Isotropis stricta.
cuneifolia, Sm. in Trans. Linn. Soc. ix. (1808) 266 = Isotropis striata.
elliptica, Vent. Jard. Malm. 115 = O. ellipticum.
lanceolata, Vent. l. c. = O. Callistachys.
linarifolia, Hort. Mackay, ex Sweet, Hort. Brit. ed. I. 474, nomen = O. lineare.
linariæfolia, G. Don, Gen. Syst. ii. 117 = O. lineare.
linearis, Benth. in Enum. Pl. Hueg. 27 = O. lineare.
longifolia, Pxzl. Mag. viii. 81 = O. Callistachys.
ovata, Sims, Bot. Mag. t. 1925 = O. Callistachys.
oxylobioides, Meissn. in Lehm. Pl. Preiss. i. 27 = O. capitatum.
parviflora, Benth. in Enum. Pl. Hueg. 27 = O. lineare.
purpurea, Van Houtte, ex Heynh. Nom. ii. 107 = O. ellipticum.
retusa, Lodd. Bot. Cab. t. 1983 = O. Callistachys.
sparsa, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 69 = O. Peltenea.
spathulata, Meissn. in Lehm. Pl. Preiss. ii. 208 = O. spathulatum.
tetragona, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) t. 349 = O. retusum.

CALLISTEMMA, Boiss. Fl. Orient. iii. 146 (1875) = **Boaldoa**, Tourn. (Dipsac.).
brachiata, Boiss. l. c. = S. brachiata.
Sibthorpianum, Nym. Consp. 341 = S. brachiata.

CALLISTEMMA, Cass. in Dict. Sc. Nat. vi. Suppl. 45 (1817), nomen prius = **Callistephus**, Cass. (Compos.).
Capistrum, Reichb. Hort. Bot. i. t. 84, sphalm. = *Callistemma capitatum*.
hortense, Cass. in Dict. Sc. Nat. vi. Suppl. 45; et Bull. Soc. Philom. (1817) 32 = Callistephus chinensis.
indicum, G. Don, in Lond. Hort. Brit. 348 = Boltonia indica.

CALLISTEMON, R. Br. in App. Flind. Voy. ii. 547 (1814). *MYRTACEÆ*, Benth. & Hook. f. i. 704.
acerosus, Miq. in Nederl. Kruidk. Arch. iv. (1859) 141 = brachyandrus.
acerosus, Tausch, in Flora, xix. (1836) 411 = pinifolius.
amœnus, *Lem. in Illustr. Hortic.* vii. t. 247.—Hab.?
arborescens, F. Muell. in Linnæa, xxv. (1852) 388 = brachyandrus.
brachyandrus, *Lindl. in Journ. Hort. Soc.* iv. (1849) 113.—Austral.
capitatus, Reichb. Ic. Exot. i. 59. t. 84 = Kunzea capitata.
coccineus, *F. Muell. Fragm.* i. 18.—Austral.
Cunninghami, *Hort. Berol. ex C. Koch, Wochenschr.* vi. (1863) 294.—Austral.
flavescens, *Regel, Gartenfl.* (1861) 53.—Austral.
glaucus, Sweet, Fl. Austral. sub t. 29 = speciosus.
Hainesii, F. Muell. l. c. iii. 158 = Kunzea Baxteri.
hybridus, *DC. Prod.* iii. 294.—Austral.
intermedius, *Hort. Berol. ex Steud. Nom.* ed. II. i. 389.—Austral.
juniperinus, *Heynh. Nom.* i. 148.—Austral.
lanceolatus, *Sweet, Fl. Austral.* sub t. 29; *DC. Prod.* iii. 293.—Austral.
lophanthus, Sweet, Fl. Austral. t. 29 = salignus.
macrostachyus, Lindl. Bot. Reg. (1838) t. 7 = Kunzea Baxteri.
marginatus, Sweet, Hort. Brit. ed. I. 155; DC. Prod. iii. 294 = lanceolatus.
macrostachyus, Lindl. Bot. Reg. (1838) t. 7 = Kunzea Baxteri.

CALLISTEMON :—
nervosus, Lindl. in Mitch. Journ. Trop. Austral. 235 = Melaleuca Leucadendron.
pallidus, DC. Prod. iii. 293 = salignus.
paludosus, F. Muell. Fragm. i. 14 = salignus.
Panchæi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 188; *in Ann. Sc. Nat.* Sér. V. ii. (1864) 188.—N. Caled.
pendulus, *Regel, Gartenfl.* (1858) 288.—Austral.
phœniceus, *Lindl. Swan River App.* 10.—Austral
pinifolius, *Sweet, Hort. Brit.* ed. I. 155; *DC. Prod.* iii. 293.—Austral.
Pithyoides, Miq. in Nederl. Kruidk. Arch. iv. (1859) 143 (sp. dub.).—Austral.
pungens, *G. Don, ex Loud. Hort. Brit.* 197.—Austral
rigidus, *R. Br. in Bot. Reg.* t. 893.—Austral.
rugulosus, DC. Prod. iii. 293 = rigidus.
rugulosus, Miq. in Nederl. Kruidk. Arch. iv. (1859) 141 = coccineus.
salignus, *Sweet, Fl. Austral.* sub t. 29; *DC. Prod.* iii. 293.—Austral.
scaber, Lodd. Bot. Cab. t. 1988 = lanceolatus.
sempervirens, *Lodd. l. c.* t. 523.—Austral.
Sieberi, DC. Prod. iii. 293 = salignus.
speciosus, *DC. l. c.* 294.—Austral.
suberosus, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 188.—N. Caled.
tereticaulus, *F. Muell. in Linnæa*, xxv. (1852) 387.—Austral.
viminalis, G. Don, ex Loud. Hort. Brit. 197 = rigidus.
viridiflorus, Sweet, Hort. Brit. ed. I. 155 = salignus.

CALLISTEPHANA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 367 = Coronilla, Linn. (Legumin.).
varia, Fourr. l. c. = C. varia.

CALLISTEPHUS, Cass. in Dict. Sc. Nat. xxxvi. 491(1825). *COMPOSITÆ*, Benth. & Hook. f. ii. 270
CALLISTEMMA, Cass. l. c. vi. Suppl. 45 (1817), nomen prius.
hirsutus, Lindl. ex DC. Prod. v. 275 = Heteropappus hispidus.
chinensis, Nees, Gen. et Sp. Ast. 222 = hortensis
concolor, Dalz. in Hook. Kew Journ. ii. (1850) 344 = Pulicaria angustifolia.
Heyneanus, DC. Prod. v. 275 = Pulicaria Wightiana.
hortensis, *Cass. in Dict. Sc. Nat.* xxxvii. 491.—Sibiria; China.
indicus, DC. Prod. v. 275 = Boltonia indica.
Wightianus, DC. l. c. = Pulicaria Wightiana.

CALLISTHENE, Mart. Nov. Gen. et Sp. i. 123. t. 75, 76 (1824). *VOCHYSIACEÆ*, Benth. & Hook. f. i. 976.
Blanchetii, *Warm. in Mart. Fl. Bras.* xiii. ii. 26.—Am trop.
erythrocalda, *Warm. l. c.* 28.—Am. trop.
fasciculata, *Mart. Nov. Gen. et Sp.* i. 126.—Bras.
major, *Mart. l. c.* i. 124. t. 75.—Bras.
microphylla, *Warm. in Mart. Fl. Bras.* xiii. ii. 28.—Am. trop.
minor, *Mart. Nov. Gen. et Sp.* ii. 126. t. 76.—Bras.
mollissima, *Warm. in Mart. Fl. Bras.* xiii. ii. 26.—Am. trop.

CALLISTHENIA, Spreng. Gen. i. 8 (1830) = prec.

CALLISTROMA, Fenzl, in Flora, xxvi. (1843) 467; Endl. Gen. Suppl. iii. 83 (1843) = **Olivaria**, Vent. (Umbell.).
crubescens, Fenzl, l. c. = O. orientalis.

CALLITHAMNA, Herb. Amaryll. 225 (1837) = **Stenomesson**, Herb. (Amaryllid.).
angustifolium, Herb. in Bot. Mag. t. 3696. f. b = S. viridiflorum.
Hartwegianum, Van Houtte, ex Heynh. Nom. ii. 108. nomen = S. aurantiacum!
spathulatum, Herb. Amaryll. 225.—! Trusillo.!
viridiflorum, Herb. l. c. = S. viridiflorum.

CALLITRICHE, Linn. Syst. ed. VI. 32 (1748). *HALORAGEÆ*, Benth. & Hook. f. i. 676.
STELLARIA, Hall. Enum. Stirp. Helv. i. 366 (1742).
STELLINA, Babani, in Nuov. Giorn. Bot. Ital. v. (1873) 218.
aestivalis, Thuil. Fl. Par. 2 = stagnalis.

CALLITRICHE:—
alpina, Schur, in Oestr. Bot. Zeitschr. (1858) 408 = verna.
amblycarpa, Scheidw. ex Hegelm. Monog. Callit. 61 = truncata.
androgyna, Linn. Cent. Pl. Rar. i. 31 = verna.
angustifolia, Gilib. Exercit. ii. 421 = autumnalis.
angustifolia, Hoppe, Taschenb. (1792) 160 = verna.
antarctica, *Engelm. ex Hegelm. in Verh. Bot. Ver. Brandenb.* ix. (1867) 20.—Reg. Antarct.
aquatica, Huds. Fl. Angl. ed. II. 489 = verna ?
Asagraei, *Hegelm. Monog. Callit.* 54. t. 8. f. 9.—Am. bor.
attenuata, *Delastre, Pl. Dép. Vienne*, 144.—Gallia.
autumnalis, A. Gray, ex Hegelm. Monog. Callit. 55 = Asagraei.
autumnalis, Kuetz. in Linnaea, vii. (1832) 186 = hamulata.
autumnalis, Bertol. ex Steud. Nom. ed. II. i. 359 = Bertoniana.
autumnalis, Brot. ex Stend. l. c. = latifolia.
autumnalis, *Linn. Fl. Suec.* ed. II. n. 4.—Europ. ; Am. bor.
autumnalis, Michx. Fl. Bor. Am. i. 3 = Peplis diandra.
autumnalis, Pollich, Fl. Palat. 3 = verna.
Bertoniana, *Steud. Nom.* ed. II. i. 359.—Chili.
Bolanderi, *Hegelm. in Verh. Bot. Ver. Brandenb.* x. (1868) 116.—Am. bor. occ.
brachycarpa, *Hegelm. l. c.* 115.—Austral.
brevifolia, *Pursh, Fl. Am. Sept.* i. 3.—Am. bor.
brutia, Petagna, Inst. Bot. ii. 10 = hamulata.
capillaris, *Purl. Fl. Ital. iv.* 423.—Italia.
carpitana, Schult, Prod. Fl. Stargard. 3 = verna.
confervoides, Thuil. ex Link, in Spreng. Jahrb. Gew. iii. 80 = autumnalis.
cornuta, Rafin. in Med. Rep. New York, Ser. III. ii. (1811) 408 = hamulata.
cophocarpa, Sendtn. ex Hegelm. Monog. Callit. 55 = verna.
cruciata, Lebel, ex Hegelm. in Verh. Bot. Ver. Brandenb. ix. (1867) 38 = truncata.
cruciata, *Rafin. in Med. Rep. New York*, Ser. III. ii. (1811) 408.—Am. bor.
cuneifolia, A. Br. ex Hegelm. in Verh. Bot. Ver. Brandenb. ix. (1867) 21 = verna.
cyclocarpa, *Hegelm. l. c.* x. (1868) 116.—Austral.
decussata, Link, in Spreng. Jahrb. Gew. iii. 80 = autumnalis.
decussata, Schur, Enum. Pl. Transs. 217 = hamulata.
deflexa, *A. Br. in Hegelm. Monog. Callit.* 58. t. 3.—Am. bor. et austr.
dolca, *Petagn. Inst. Bot.* ii. 10.—Italia.
dioica, Schur, in Verh. Siebenb. Ver. Naturw. (1859) 98 = stagnalis.
Drummondi, *Hegelm. Monog. Callit.* 60.—Austral.
dubia, *Hoffm. Obs. Bot.* n. 1 ; *ex Roth, Tent. Fl. Germ.* i. 353 = verna.
eunutumnalis, Syme, Engl. Bot. ed. III. viii. 122 = autumnalis.
eurverna, Syme, l. c. 119 = verna.
foliosa, Rafin. in Med. Repos. New York, V. (1808) 354 = autumnalis.
fontana, Scop. Fl. Carn. ed. II. ii. 251 = verna.
genuina, Ducommun, Taschenb. 348 = hamulata.
glomerata, Thuill. ex Link, in Spreng. Jahrb. Gew. iii. 80 = autumnalis.
graminea, Link, l. c. = truncata.
hamulata, *Kuetz. ex Koch, Syn. Fl. Germ.* ed. II. 871.—Europ.
hermaphrodita, Linn. Cent. Pl. Rar. i. 31 = verna.
heterophylla, Pursh, Fl. Am. Sept. i. 3 = verna.
heteropoda, *Engelm. ex Hegelm. in Verh. Bot. Ver. Brandenb.* ix. (1867) 40.—Am. austr.
intermedia, Hoffm. Fl. Germ. (1791) 2 = hamulata.
intermedia, *Hegge, Taschenb.* (1792) 159 = verna.
japonica, *Engelm. ex Hegelm. in Verh. Bot. Ver. Brandenb.* x. (1868) 118.—Japon.
Kuetzingi, Rayr. Fl. Hgric. 872 = stagnalis.
latifolia, Gilib. Exercit. ii. 421 = hamulata vel verna.
latifolia, *Link, ex Steud. Nom.* ed. II. i. 359.—Lusitan.
linearis, Pursh, Fl. Am. Sept. i. 3 = autumnalis.
macropteryx, *Hegelm. Monog. Callit.* 59.—Austral.
marginata, *Torr. in Pacif. Rail. Rep.* iv. 135.—Calif. ; Chili.
Mandonis, *Hieronch & Muell. Arg. in Heurch, Obs. Bot.* 128.—Ecuador.
microscypa, *Engelm. ex Hegelm. in Verh. Bot. Ver. Brandenb.* x. (1868) 113.—Cuba.
minima, Hoppe, Taschenb. (1792) 157 = verna.

CALLITRICHE:—
minor, Hellynck, Fl. Namur (1855), 232 = stagnalis ?
Muelleri, *Sond. in Linnaea*, xxviii. (1856) 229.—Austral.
Nuttalli, *Torr. in Pacif. Rail. Rep.* iv. 135.—Am. bor.
obtusangula, *Le Gall, ex Hegelm. Monog. Callit.* 54.—Europ.
occidentalis, *Hegelm. l. c.* 57. t. iv. 5.—Cuba.
ovata, *Rafin. in Med. Rep. New York*, Ser. III. ii. (1811) 408.—Am. bor.
pallens, Bleb. Fl. Taur. Casuc. iii. 6 = hamulata.
pallens, Goldb. in Mém. Soc. Nat. Mosc. v. (1817) 118 = verna.
palustris, Linn. Sp. Pl. 969 = autumnalis.
pedunculata, DC. Fl. Fr. iv. 415 = hamulata.
pedunculata, A. Gray, Man. ed. Hegelm. Monog. Callit. 58 = deflexa.
pendunculata, F. Muell. ex Hegelm. l. c. 61 = truncata.
pedunculosa, *Nutt. in Trans. Am. Phil. Soc.* N. S. v. (1837) 140.—Am. bor.
peploides, *Nutt. l. c.* 141.—Am. bor
platycarpa, Kuetz. in Linnaea, vii (1832) 174 = stagnalis.
plena, *Rafin. in Med. Rep. New York*, Ser. III. ii. (1811) 408.—Am. bor.
polymorpha, *Loennroth, Obs. Crit. Pl. Suec.* 19 (1854).—Scania.
reflexa, Lange, in Cutanda, Fl. Matrit. 292 = hamulata
sepulta, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 298.—Am. bor. occ.
sessilis, DC. Fl. Fr. iv. 414 = autumnalis, verna.
Sonderi, *Hegelm. in Verh. Bot. Verh. Brandenb.* ix. (1867) 31.—Austral.
spuria, *Rafin. in Med. Repos. New York*, v. (1808) 354.—Am. bor.
stagnalis, *Scop. Fl. Carn.* ed. II. ii. 251.—Europ ; Indi.
Stellaria, Rafin. in Med. Rep. New York, Ser. III. ii. (1811) 408 = autumnalis.
stenocarpa, *Hegelm. in Verh. Bot. Ver. Brandenb.* x. (1868) 114.—Calif.
stellata, Hoppe, Taschenb. (1792) 158 = verna.
tenuifolia, Fries, Novit. Fl. Suec. 280 = hamulata.
tenuifolia (Gennar), ex Parl. Fl. Ital. iv. 434 = capillaris.
tenuifolia, Thuil. ex Pers. Syn. i. 6 = verna.
terrestris, DC. Prod. iii. 71 = deflexa.
terrestris, *Kuhl. ex Rafin. l. c.* Ser. III. ii. (1811) 408.—Am. bor.
terrestris, Rafin. in Med. Repos. New York, v. (1808) 358 = brevifolia.
transitanica, Schur, Enum. Pl. Transs. 216 = stagnalis.
truncata, *Guss. Pl. Rar.* 4.—Italis.
turfosa, *Bert. in Am. Journ. Sc.* xix. (1831) 808, nomen ; *Hegelm. Monog. Callit.* 59.—Chili.
turfosa, *Schur, Enum. Pl. Transs.* 216.—Transylv.
umbonata, *Hegelm. in Verh. Bot. Ver. Brandenb.* ix. (1867) 19.—Tasmania.
verna, Hook. f. Fl. Antarct. i. 11 = antarctica.
verna, Linn. f. Suppl. Pl. ed. II. n. 8.—Europ ; Am. bor.
verna, Linn. l. c. partim = polymorpha.
verna, Huds. Fl. Pls. i. 4 = stagnalis.
vernalis, Kuetz. in Linnaea, vii. (1832) 175 = verna.
virens, Goldb. in Mém. Soc. Nat. Mosc. v. (1817) 119 = autumnalis.
Wightiana, Wall. Cat. n. 7006 = stagnalis.

CALLITRIS, Vent. Decad. (1808) 10. *CONIFERAE*, Bentth. & Hook. f. iii. 424.
FRENELA, Mirb. in Mém. Mus. Par. xiii. (1826) 30.
FRESNELIA, Steud. Nom. ed. II. i. 648 (1840).
LEICHHARDTIA, Sheph. Cat. Pl. Colt. Sydn. (1851) 15.
PACHYLEPIS, Brongn. in Ann. Sc. Nat. Sér. 1. xxx. (1833) 189.
OCTOCLINIS, R. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 31.
PAROLINIA, Endl. Gen. Suppl. i. 1372 (1841).
WIDDRINGTONIA, Endl. l. c. ii. 25 (1843).
Actinostrobus, F. Muell. Rep. Burdek. Exp. 19 = Actinostrobus pyramidalis.
acuminata, *F. Muell. Census*, 109.—Austral.
arborea, *Schrad. ex E. Mey. Zwei Pfl. Docum.* 73, nomen.—Afr. austr.
arenosa, Sweet, Hort. Brit. ed. II. 478, nomen = rhomboides.
articulata, Hort. ex Gord. Pinet. ed. II. 117 = rhomboides.
australis, Sweet, Hort. Brit. ed. II. 474 = rhomboides.
calcarata, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 74.—Austral.
capensis, Schrad. ex Gord. Pinet. 338 = cupressoides.

CALLITRIS:—
columellaris, F. Muell. Fragm. v. 198 = robusta.
conglobata, *Sieber, ex Heynh. Nom.* i. 148.—Austral.
cupressiformis, Vent. Decad. 10 = rhomboides.
cupressoides, *Schrad. ex E Mey. Zwei Pfl. Docum.* 126.—Afr. austr.
Drummondii, *Benth. & Hook. f. ex F. Muell. Census.* 109.—Austral.
elegans, *Sieber, ex Heynh. Nom.* i. 149.—Austral.
Fothergilli, Loud. Encycl. Trees, 1072 = australis.
fruticosa, R. Br. ex Rich. Conif 49 = calcarata.
glauca, R. Br. ex Mirb. in Mém. Mus. Par. xiii. (1825) 74 = robusta.
Gunnii, Hook. f. in Hook. Lond. Journ. iv. (1845) 147 = australis.
Hugelii, *Knight, Syn. Conif.* 14.—Austral.
Maciesyana, *F. Muell. in Rep. Burdek. Exped.* 17, *Extra-Trop. Pl. Am.* ed. (1854) 85.—Austral.
macrostachya, Hort. ex Steud. Nom. ed. II. i. 260 = quadrivalvis.
montana, *Sieber, ex Heynh. Nom.* i. 149.—Austral.
Muelleri, *Benth. & Hook. f. ex F. Muell. Census*, 109 —Austral.
oblonga, *Rich. Conif.* 49. t. 18. f. 2.—Austral.
Parlatorei, *F. Muell. Fragm.* v. 186.—Austral.
Preissii, Miq. in Lehm. Pl. Preiss. i. 643 = robusta.
propinqua, R. Br. ex Endl. Syn. Conif. 88 = robusta.
pyramidalis, Sweet, Hort. Brit. ed. II. 478 = calcarata.
quadrivalvis, *Vent. Decad.* 10.—Afr. bor.
rhomboides, *R. Br. in Rich. Conif.* 47. t. 18.—Austral
robusta, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 74.—Austral.
Roei, *Benth. & Hook. f. ex F. Muell. Census*, 109 —Austral.
stricta, Schrad. ex Carr. Conif. ed. I. 66 = cupressoides.
tripartita, Hort. ex Loud. Encyc. Trees, 1072 = quadrivalvis !
tuberculata, R. Br. ex Endl. Syn. Conif. 88 = *Ventenatii*, R. Br. ex Mirb. in Mém. Mus. Par. xiii. (1825) 74 = rhomboides.
verrucosa, R. Br. ex Mirb. l. c. = robusta.

CALLIXENE, Comm. ex Juss. Gen. 41 (1789) = Luxuriaga, Ruiz & Pav. (Liliac.).
magellanica, Raensch. Nom. ed. III. 98 = L. marginata.
marginata, Lam Illustr. t. 248 = L. marginata.
melantha, Colenso, in Trans. N. Z. Inst. xvii. (1885) 250.—N. Zel.
parviflora, Hook. f. in Hook. Ic. Pl. t. 682 = L. parviflora.
polyphylla, Hook. l. c. t. 674 = L. erecta.

CALLOPISMA, Mart. Nov. Gen et Sp. ii. 107. t. 188, 184 (1827) = Deianira, Cham. & Schlecht. (Gentian.).
amplexifolium, Mart. l. c. 109 = D. nervosa.
cordifolium, Lhotsky, ex Griseb. in DC. Prod. ix. 48 = D. erubescens.
perfoliatum. Mart. Nov. Gen. et Sp. ii. 108 = D. erubescens.

CALLOSMIA, Presl, Bot. Bemerk. 108 (1844) = Anneslea, Wall. (Ternstroem.).
fragrans, Presl, l. c. = A. fragrans.

CALLOSTYLIS, Blume, Bijdr. 340. t. 74 (1825) = Eriis, Lindl. (Orchid.).
rigida, Blume, l. c. = E. pulchella.

CALLOTROPIS, G. Don, Gen. Syst. ii. 228 (1832) = Galega, Linn. (Legumin.).
tricolor, G. Don, l. c. = G. officinalis.

CALLUNA, Salisb. in Trans. Linn. Soc. vi. (1802) 317. *ERICACEAE*, Bentth. & Hook. f. ii. 589.
atlantica, Seem. in Journ. Bot. iv. (1866) 306 = vulgaris.
ciliaris, Schur, Enum. Pl. Transs. 447 = vulgaris.
Erica, DC. Fl. Fr. iii. 680 = vulgaris.
genuina, Ducommun, Taschenb. 491 = vulgaris.
sagittifolia, S. F. Gray, Nat. Arr. ii. 399 = vulgaris.
vulgaris, *Salisb. in Trans. Linn. Soc.* vi. (1802) 317.—Europ. ; Am. bor.

CALOBOTA, Eckl. & Zeyh. Enum. 192 (1836) = **Lebeckia**, Thunb. (Legumin.).
cytisoides, Eckl. & Zeyh. l. c. 191 = L. cytisoides.
decipiens, Benth. in Ann. Wien. Mus. ii. (1838) 142 = L. decipiens.
decutiens, Benth. l. c. = L. decutiens.
flexuosa, Benth. l. c. = L. flexuosa.
multiflora, Benth. l. c. = L. multiflora.
pulchella, Eckl. & Zeyh. Enum. 191 = L. cytisoides.

CALOBOTRYA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 21 = **Ribes**, Linn. (Saxifr.).
sanguinea, Spach, l. c. = R. sanguineum.

CALOCEDRUS, Kurz, in Journ. Bot. xi. (1873) 196. t. 183 = **Libocedrus**, Endl. (Conifer.).
macrolepis, Kurz, l. c. = L. macrolepis.

CALOCEPHALUS, R. Br. in Trans. Linn. Soc. xii. (1817) 106. *COMPOSITAE*, Benth. & Hook. f. ii. 520.
ACHRYSUM, A. Gray, in Hook. Kew Journ. iv. (1852) 228.
RENNOSPORA, A. Gray, l. c. iii. (1851) 172.
LEUCOPHYTA, R. Br. in Trans. Linn. Soc. xii. (1817) 106.
PACHYSURUS, Steetz, in Lehm. Pl. Preiss. i. 441 (1845).
serusoides, *Benth. Fl. Austral.* iii. 576.—Austral.
angianthoides, *Benth. l. c.* 575.—Austral.
Brownii, *F. Muell. Rep. Babb. Exp.* 13.—Austral.
chrysanthus, Schlecht. in Linnaea, xx. (1847) 592 = Craspedia chrysantha.
citreus, *Less. Syn. Comp.* 371.—Austral.
Drummondii, *Benth. Fl. Austral.* iii. 574.—Austral.
Franciali, *Benth. l. c.* 576.—Austral.
gnaphalioides, Hook. in Mitch. Journ. Trop. Austral. 378 = Helipterum moschatum.
lacteus, *Less. Syn. Comp.* 371.—Austral.
multiflorus, *Benth. Fl. Austral.* iii. 576.—Austral.
platycephalus, *Benth. l. c.*—Austral.
Sonderi, *F. Muell. 2nd Gen. Rep.* (1854) ; *et Rep. Babb. Exp.* 13.—Austral.

CALOCHILUS, R. Br. Prod. 320 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 614.
campestris, *R. Br. l. c.*—Austral.
herbaceus, Lindl. Gen. et Sp. Orch. 459 = campestris.
paludosus, *R. Br. Prod.* 320.—Austral.
Robertsoni, *Benth. Fl. Austral.* vi. 315.—Austral.

CALOCHLAMYS, Presl, Bot. Bemerk. 148 (1844) = **Congea**, Roxb. (Verbenac.).
capitata, Presl, l. c. 149 = Congea tomentosa.

CALOCHORTUS, Pursh, Fl. Am. Sept. i. 240 (1814). *LILIACEAE*, Benth. & Hook. f. iii. 820.
CYCLOBOTHRA, G. Don, in Sweet, Brit. Flow. Gard. t. 278 (1829).
albus, *Dougl. ex Benth. in Maund & Hensl. Botanist*, t. 98.—Calif.
apiculatus, *Baker, in Journ. Linn. Soc.* xvii. (1875) 306.—Am. bor. occ.
aureus, *S. Wats. in Am. Natural.* vii. (1873) 303.—Calif.
Barnardi, *Dougl. ex Steud. Nom.* ed. II. i. 260.—Am. bor. occ.
Benthami, *Baker, in Journ. Linn. Soc.* xiv. (1875) 304.—Calif.
Bonplandianus, *Schult. f. Syst.* vii. 1532.—Mexic.
caeruleus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 263.—Calif.
Catalinae, *S. Wats. l. c.* 263.—Calif.
citrinus, *Baker, in Bot. Mag.* t. 6900.—Calif.
clavatus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 265. —Calif.
Douglasianus, *Schult. f. in Van Hall, Bijdr.* iv. 127. —Am. bor. occ.
elegans, Baker, in Journ. Linn. Soc. xiv. (1875) 305 = Illacinus, nudus et Tolmiei.
elegans, Hook. f. Bot. Mag. t. 5976 = Maweanus.
elegans, *Pursh, Fl. Am. Sept. i.* 240.—Am. bor. occ.
eurycarpus, S. Wats. in Bot. King, Exp. 348 = nitidus.
flavus, *Schult. f. Syst.* vii. 1533.—Mexic.
flexuosus, *S. Wats. in Am. Natural* vii. (1873) 303.— Am. bor. occ.
fuscus, *Schult. f. Syst.* vii. 1534.—Mexic.

CALOCHORTUS:—
Ghiesbreghtianus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 268.—Mexic.
gracilus, Nuttall, Gartenfl. (1875) 360 ; et in Act. Hort. Petrop. iii. 11. (1875) 285 = caeruleus.
Greenei, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 264. —Am. bor. occ.
Gunnisoni, *S. Wats. in Bot. King, Exp.* 348.—Am. bor. occ.
Hartwegii, *Benth. Pl. Hartw.* 26.—Mexic.
Kennedyi, *Porter in Coult. Bot. Gaz.* ii. (1877) 79.—Calif.
Leichtlinii, Hook. f. Bot. Mag. t. 5862 = Nuttallii.
lilacinus, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 5.—Calif.
longebarbatus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 381.—Am. bor. occ.
luteus, *Dougl. ex Kunth, Enum. Pl.* iv. 288.—Calif.
luteus, Nutt. in Journ. Acad. Phil. vii. (1834) 58 = Nuttallii.
Lyallii, Baker, in Journ. Linn. Soc. xiv. (1875) 305 = elegans.
macrocarpus, *Dougl. in Trans. Hort. Soc.* vii. (1830) 276. t. 5.—Am. bor. occ.
Maweanus, *Leichtl. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 305.—Am. bor. occ.
monophyllus, *Lem. in Fl. des Serres*, Sér. I. v. (1849) 480 b.—Calif.
nitidus, *Dougl. in Trans. Hort. Soc.* vii. (1830) 277. t. 8.—Calif.
nitidus, Torr. Pacif. Rail. Rep. iv. 146 = Benthami.
nudus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 263.— Calif.
Nuttallii, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 124. —Am. bor. occ.
Palmeri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 266. —Calif.
pallidus, Schult. f. Syst. vii. 1538 = flavus.
parviflorus, Nutt. ex Baker, in Journ. Linn. Soc. xiv. (1875) 306 = Nuttallii.
pulchellus, *Dougl. ex Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 412.—Calif.
purpureus, Baker, in Journ. Linn. Soc. xiv. (1875) 308, pro parte = Bonplandianus.
pusillus, *Dougl. ex Steud. Nom.* ed. II. i. 260.—Am. bor. occ.
spatulatus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 267.—Mexic.
splendens, *Dougl. ex Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 411. t. 15.—Calif.
Tolmiei, *Hook. & Arn. Bot. Beech. Voy.* 398.—Calif.
umbellatus, Wood, in Proc. Nat. Sc. Philad. (1868) 168 = lilacinus.
uniflorus, *Hook. & Arn. Bot. Beech. Voy. Suppl.* 398. —Calif.
uniflorus, Hook. f. Bot. Mag. t. 5804 = lilacinus.
venustus, *Dougl. ex Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 412 t. 15. f. 2.—Calif.
vestitus, *Benth. ex Steud. Nom.* ed. II. i. 260.— Calif.
Wendii, Wood, in Proc. Nat. Sc. Philad. (1868) 169 = luteus.

CALOCOCCUS, Kurz, ex Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind. xxvi. (1864) 48 = Phyllanthus, Linn. (Euphorbiac.).
sundaicus, Kurz, l. c. = P. sundaicus.

CALODENDRUM, Thunb. Nov. Gen. ii. 41 (1782). *RUTACEAE*, Benth. & Hook. f. i. 288.
PALLASIA, Houtt. Handlid. ii. 383 (1775).
capense, *Thunb. l. c.* 43 ; *et Fred. Pl. Cap.* 44.— Afr. austr.

CALODIUM, Lour. Fl. Cochinch. 247 (1790) = Cassytha, Linn. (Laurin.).
cochinchinensis, Lour. l. c. = Cas. filiformis.

CALODONTA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vi. (1841) 448 = Tolpis, Adans. (Compos.).
asperica, Nutt. l. c. = T. virgata.

CALODRACON, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 125, 127 = Cordyline, Comm. (Liliac.).
helicoinstylis, Planch. l. c. = Cord. helicoinstylis.
Jacquinii, Planch. l. c. = Cord. Jacquinii.
nubilis, Planch. l. c. vii. (1851–52) 106. t. 622, 623.— Japon.
Siebori, Planch. l. c. vi. (1850–51) 128 = Cord. Siebori.

CALODRACON :—
Siebolditi, Planch. in Fl. des Serres, Sér. I. vi. 127 = Dracaena Sieboldii.
terminalis, Planch. l. c. = Cord. terminalis.
venusta, Planch. l. c. 132 = Dracaena venusta.

CALODRYUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 401. t. 51 = **Quivisia**, Comm. (Mellac.).
tubiflorum, Desv. l. c. = Q. lanceolata.

CALOGYNE, R. Br. Prod. 579 (1810). *GOODENOVIEAE*, Benth. & Hook. f. ii. 589.
DIETVIA, Gaudich. in Freyc. Voy. Bot. 460. t. 80 (1826).
Berardiana, *F. Muell. Fragm.* vi. 7.—Austral.
chinensis, *Benth. in Journ. Linn. Soc.* v. (1861) 78.— China.
dietyfia, F. Muell. Fragm. vi. 6 = Berardiana.
heteroptera, *F. Muell. Fragm.* x. 43.—Austral.
pilosa, *R. Br. Prod.* 579.—Austral.
purpurea, *F. Muell. Fragm.* viii. 57.—Austral.

CALOMECON, Spach, Hist. Vég. Phan. vii. 7 (1839 = **Papaver**, Tourn.
bracteatum, Spach, l. c. 9 = P. bracteatum.
orientale, Spach, l. c. = P. orientale.

CALOMERIA, Vent. Jard. Malm. 78. t. 78 (1804) = **Rumfea**, Sw. (Compos.).
amaranthoides, Vent. l. c. = H. elegans.

CALOMORPHE, Kunze, ex Walp. Rep. i. 680 (1840) = **Lesmes**, Klotzsch (Legumin.).
Ehrenbergiana, Kunze, ex Walp. l. c. = L. robinioides.

CALOMYRTUS, Blume, Mus. Bot. Lugd. Bat. i. 76, in obs. (1849) = **Myrtus**, Tourn.
excelsa, Blume, l. c. = M. excelsa.
ovalifolia, Blume, l. c. = M. ovalifolia.

CALONNEA, Bachox, Icon. (1796) t. 136 = **Gaillardia**, Fouger. (Compos.).
pulcherrima, Bachox, l. c. = G. pulchela.

CALONYCTION, Choisy, in Mém. Soc. Phys. Genèv vi. 441 (1833) (Conv. Or. 59) = **Ipomoea**, Linn (Convolvul.).
acanthocarpum, Choisy, in DC. Prod. ix. 346 = I. acanthocarpa.
asperum, Choisy, Conv. Or. 60 = I. grandiflora.
Bona-nox, Bojer, Hort. Maurit. 228 = I. Bona-nox.
capillatum, Miq. Fl. Ind. Bat. ii. 598 = I. capillata.
chenopodiifolium, Mart. & Gal. in Bull. Acad. Brux xii. ii. (1845) 369 = I. chenopodiifolia.
cissoides, G. Don, Gen. Syst. iv. 264.—Ecuador.
comoremis, Boj. Hort. Maurit. 228 = I. grandiflora.
comosperma, Boj. l. c. = I. grandiflora.
diversifolium, Hask. Pl. jav. Rar. 528 = I. tricho sperma.
dubium, Mart. & Gal. in Bull. Acad. Brux. xii. ii (1845) 368 = I. dubia.
Galeottii, Martens, ex Mart. & Gal. l. c.—Mexic.
grandiflorum, Choisy, Conv. Or. 60 = I. grandiflora.
Jacquinii, G. Don, Gen. Syst. iv. 264 = I. grandiflora.
longiflorum, Hask. Pl. Jav. Rar. 528 = I. grandiflora.
macranthoides, Colla, N. Sp. Calon. 15 = I. Bona-nox.
macranthum, Van Houtte, ex Pois. in Rev. Hortic. Sér. II. ii. (1846–48) 498.—Hab. ?
magnolocarpum, A. Rich. Fl. Cub. Fanerog. ii. 129 = I. Bona-nox.
malitiosum, Zoll. Syst. Verz. Ind. Archip. 128, 181. —Java.
muricatum, G. Don, Gen. Syst. iv. 264 = I. Bona-nox.
muticum, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 380 = I. grandiflora.
noctoniscum, G. Don, in Sweet, Hort. Brit. ed. III. 483 = I. latifolia.
praetermissum, Mart. & Gal. in Bull. Acad. Brux. xii. ii. (1845) 368 = I. proximum.
pseudomuricatum, G. Don, Gen. Syst. iv. 264 = I. grandiflora.
purpurea, G. Don, l. c.—Ecuador.
pterodes, Choisy, in DC. Prod. ix. 349 = praec.
reticulorrimosum, Pavoll. Contrib. 13.—Paraguay.
Rheedii, Colla, H. Sp. Calon. 15 = I. Bona-nox.
Roxburghii, G. Don, Gen. Syst. iv. 263 = I. Bona-nox.
sanguineum, Hort.—Cf. Gard. Chron. (1868) 934 = Loasanula Kersti.
sanguineolentum, Gaillon, in Rev. Hortic. (1860) 20. —Hab. ?

CALONYCTION :—
speciosum, Choisy, Conv. Or. 59 = I. Bona-nox.
trichospermum, Choisy, l. c. 60 = I. trichosperma.
Tuba, Colla, Nov. Sp. Calon. 15 = I. grandiflora †
venustum, Mart. & Gal. in Bull. Acad. Brux. xii. ii. (1845) 269 = I. venustum.

CALOPAPPUS, Meyen, Reise, i. 315 (1834) = Nassauvia, Juss. (Compos.).
acanthifolius, Remy, in C. Gay, Fl. Chil. iii. 398.— Chili.
acerosus, Meyen, Reise, i. 315 = N. acerosa.

CALOPETALON, J. Drum. ex Harv. in Hook. Kew Journ. vii. (1855) 59 = Marianthus, Hueg. (Pittospor.).
ringens, J. Drum. l. c. 58 = M. ringens.

CALOPHACA, Fisch. Hort. Gorenk. ed. II. 67 (1812); et in DC. Prod. ii. 270 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 606.
CHESNEYA, Lindl. in Endl. Gen. 1275 (1840).
Hovenii, *Schrenk, Enum. Pl. Nov.* 74.—Soongaria.
juncea, Stend. Nom. ed. II. i. 360, sphalm. = Caligonum Calliphysa.
soongorica, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 401 = Hovenii.
volgarica, *Fisch. Hort. Gorenk.* ed. II. (1812) 68.— Ross. austr.

CALOPHANES, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 181 (1833). *ACANTHACEAE*, Benth. & Hook. f. ii. 1077.
CHARTACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836).
DYSCHORISTE, Nees, in Wall. Pl. As. Rar. iii. 75 (1832), nomen prius.
LINOTYLES, Fenzl, in Linnaea, xviii. (1850) 94.
adscendens, *Hochst. ex Nees, in DC. Prod.* xi. 111.— Abyss.
amoenus, *Nees, Mert. Fl. Bras.* ix 27.—Bras.
angustus, *A. Gray, Syn. Fl. N. Am.* ii. i. 456.—Florida.
angustifolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 502.— Mexic.
bifrons, Shuttlew. ex Nees, in DC. Prod. xi. 107 = oblongifolius.
bilabiatus, *Seem. Bot. Voy. Herald*, 324. t. 65.— Mexic.
buchenavii, *Vatke, in Bremen Abh.* ix. (1885) 181.— Madag.
Barkei, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 24. —Afr. austr.
capitatus, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 121.—Mexic.
Clarkei, *Vatke, in Bremen Abh.* ix. (1886) 182.—Ins. Mascar.
ciliatus, *Nees, in DC. Prod.* xi. 110.—Peruv.
costatus, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. austr.
crinitus, *Nees, in Mart. Fl. Bras.* ix. 26.—Bras.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 160.—Cuba.
Dalzellii, *T. Anders. ex Bedd. Ic. Ind. Or.* i. 60.— Ind. or.
decumbens, *A. Gray, Syn. Fl. N. Am.* ii. i. 325.— Arizon. ; Mexic.
depressus, T. Anders. in Thw. Enum. Pl. Zeyl. 225 = Nagchana.
gracilis, *Nees, in DC. Prod.* xi. 111.—Madag.
Hendelotianus, *Nees, l. c.* 113.—Afr. trop.
Hildebrandtii, *S. Moore, in Journ. Bot.* xvii. (1880) 8, 197.—Afr. trop.
hirsutissimus, *Nees, in DC. Prod.* xi. 109.—Mexic.
hirsutus, *Oerst. ex Hiorm, in Kjoeb. Vidensk. Meddel.* (1877–78) 71.—Mexic.
humistratus, *Shuttlew. ex Nees, in DC. Prod.* xi 108. —Ann. bor.
hygrophiloides, *Nees, in Mart. Fl. Bras.* ix. 26.— Bras.
hysopifolius, *Nees, in DC. Prod.* xi. 111.—Abyss.
Jamilum-mexicanum, *Nees, l. c.* 110.—Mexic.
lavandulaceus, *Nees, in Mart. Fl. Bras.* ix. 27.—Bras.
linearis, *A. Gray, Syn. Fl. N. Am.* ii. i. 324.—Texas ; Mexic.
littoralis, *T. Anders. in Thw. Enum. Pl. Zeyl.* 225.— Ind. or.
madagascariensis, *Nees, in DC. Prod.* xi. 112.— Madag.
Marschallia, *Nees, in Mart. Fl. Bras.* ix. 25.—Bras.

CALOPHANES :—
mauritianus, T. Anders. in Journ. Linn. Soc. vii. (1864) 23, nomen = Hypoëstes serpens.
microphyllus, *Nees, in DC. Prod.* xi. 113.—Mexic.
multicaulis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Abyss.
Nagchana, *Nees, in DC. Prod.* xi. 109.—Ind. or.
natalensis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. austr.
oblongifolius, *D. Don, in Sweet, Flow. Gard.* t. 181. —Ann. bor.
oblongifolius, Torr. Bot. Mex. Bound. 122 = decumbens.
ovatus, Benth. Pl. Hartw. 89 = linearis.
Perrottetii, *Nees, in DC. Prod.* xi. 111.—Afr. trop.
Persoonii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. austr.
Pulegium, *Nees, in Mart. Fl. Bras.* ix. 25.—Bras.
quadrangularis, *Oerst. in Kjoeb. Videnak. Meddel.* (1854) 120.—Mexic.
quitensis, *Nees, in DC. Prod.* xi. 110.—Ecuador.
radicans, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. trop.
repens, *Nees, in DC. Prod.* xi. 109.—Peruv.
rigidus, Dalz. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 411 = Dalzellii.
Schiedeanus, *Nees, in DC. Prod.* xi. 111.—Mexic.
Serpyllum, *Nees, in Mart. Fl. Bras* ix. 26.—Bras.
setosus, *Nees, in DC. Prod.* xi. 112.—Afr. austr.
siphonanthus, *Nees, l. c.*—Madag.
thunbergiaeflora, *S. Moore, in Journ. Bot.* xviii. (1880) 9, 7.—Afr. trop.
trichocalyx, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 126.—Afr. trop.
Tweedieanus, *Nees, in DC. Prod.* xi. 106.—Reg. Argent vagans, *Wight, Ic.* xi. t. 1528.—Ind. or.
verticillaris, *T. Anders. ex Oliver, in Trans. Linn. Soc.* xxix. (1874) 126, in nota.—Afr. trop.

CALOPHTHALMUM, Reichb. Nom. 82 (1841) : GALOPHTHALMUM, Nees & Mart.

CALOPHYLICA, Presl, Bot. Bemerk. 89 (1844) = Phyllica, Linn. (Rhamn.).
gnidioides, Presl, l. c. = P. gnidioides.
juniperifolia, Presl, l. c. = P. gnidioides.

CALOPHYLLOIDES, Smeathm. ex DC. Prod. iii. 272 (1868) = Rugenia, Mich. (Myrtac.).
lucida, Smeathm. l. c. = E. Calophylloides.

CALOPHYLLUM, Linn. Gen. ed. I. 154 (1737). *GUTTIFERAE*, Benth. & Hook. f. i. 175.
APOTERIUM, Blume, Bijdr. 218 (1825).
AUGIA, Lour. Fl. Cochinch. 387 (1790).
acuminatum, Lam. Encyc. i. 653 = spectabile.
acuminatum, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 80 = Rheedii acuminata.
Akara, Burm. f. Fl. Ind. 121 = Tetracera laevis.
amoenum, Wall. Cat. 4849 = retusum.
amplexicaule, *Choisy, ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 281.—Ins. Philipp.
angustifolium, *Roxb. Hort. Beng.* 41 ; *Fl. Ind.* ii. 608. —Penang.
apetalum, *Blanco, Fl. Filip.* ed. II. 429.—Ins. Philipp.
apetalum, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 79 = C. inophyllum †
Augia, *Stend. Nom.* ed. II. i. 260.—China.
bancanum, Miq. Fl. Ind. Bat. Suppl. i. 499 = pulcherrimum.
Bingotan, Roxb. Hort. Beng. 41 ; Fl. Ind. ii. 607 = C. Inophyllum.
Blancoi, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 272.—Ins. Philipp.
Blumei, *Wight, Illustr.* i. 128.—Java.
brachystum, *Thw. Enum. Pl. Zeyl.* 51.—Zeylan.
brasiliense, *Camb. in St. Hil. Fl. Bras. Mer.* i. 321. t. 67.—Bras.
Burmanni, Wight, Illustr. i. 129 = C. Calaba, *Linn.*
Calaba, *Jacq. Am.* 259. t. 165.—Ins. Carib.
Calaba, *Linn. Sp. Pl.* 514.—Zeylan.
calaboides, *G. Don, Gen. Syst.* i. 622.—Ind. or.
caledonicum, *Vieill. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 291.—N. Caled.
canum, *Hook. f. Fl. Brit. Ind.* i. 271.—Malacca.
chinense, T. Anders. in Hook. f. Fl. Brit. Ind. i. 270 = Ochrocarpus longifolius.
cordato-oblongum, *Thw. Enum. Pl. Zeyl.* 407.— Zeylan.

CALOPHYLLUM :—
Caminghi, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 289.—Ins. Philipp.
canellifolium, *Thw. Enum. Pl. Zeyl.* 51, 407.—Zeylan.
Cupi, H. B. & K. Nov. Gen. et Sp. v. 208 = Rheedia Capi.
Cuspa, G. Don, Gen. Syst. i. 628 = prsec.
cymosum, Miq. Fl. Ind. Bat. Suppl. i. 497 = spectabile.
dasypodium, *Miq. Fl. Ind. Bat.* i. ii. 511.—Java.
decipiens, Thw. Enum. Pl. Zeyl. 51 = Thwaitesii.
decipiens, Wight, Illustr. 128 ; Ic. t. 108 = Wightianum.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* i. 497.— Sumatra.
dongnaiense, *Pierre, Fl. For. Cochinch.* t. 108.— Cochinch.
dryobalanoides, *Pierre, l. c.* t. 106.—Cochinch.
dubium, Stend. Nom. ed. II. i. 261 = Ochrocarpus longifolius.
edule, Seem. Bot. Voy. Herald, 89 = Rheedia edulis.
elatum, Bedd. Fl Sylv. ic. t. 250 = tomentosum.
exrelsum, Zoll. & Morr. in Nat. en Geneesk. Arch. Neerl. Indie, ii. (1845) 582 = Ochrocarpus ovalifolius.
floribundum, *Hook. f. Fl. Brit. Ind.* i. 272.—Malacca.
gracile, Miq. Fl. Ind. Bat. Suppl. i. 498 = pulcherrimum.
Griffithii, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 273. —Malacca.
Hasskarlii, *Teijsm. & Binn. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 273.— Java.
hexapetalum, *Hook. f. in Trans. Linn. Soc.* xxiii (1862) 168.—Borneo.
hirtellum, *Miq. Pl. Jungh.* 291.—Java.
Inophyllum, Lam. Encyc. i. 652 = C. Tacamahaca.
Inophyllum, *Linn. Sp. Pl.* 513.—Geront. trop.
Inophyllum, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = C. Calaba, *Jacq.*
javanicum, *Miq. Pl. Jungh.* 293.—Java.
lanceolatum, Roxb. Hort. Beng. 41 ; Fl. Ind. ii. 608 = C. Tacamahaca.
lanceolatum, Blume, Bijdr. 217 = C. Tacamahaca.
lanceolatum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 395.—Java.
lanigerum, *Miq. Fl. Ind. Bat. Suppl.* i. 498.—Ins. Banca.
longifolium, Wall. Cat. n. 4851 = Ochrocarpus longifolius.
longifolium, *Willd. Ges. Naturf. Fr. Berl. Mag.* v. (1811) 80.—N. Granat.
Lowei, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 271.—Borneo.
lucidum, *Bentb. in Hook. Lond. Journ. Bot.* ii. (1843) 270.—Guiana.
macrocarpum, *Hook. f. Fl. Brit. Ind.* i. 273.—Malacca.
macrophyllum, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 406.—Malaya.
Madruka, H. B. & K. Nov. Gen. et Sp. v. 202 = Rheedia Madruno.
marginatum, *Wall. Cat.* n. 4845 (sp. ?).—Ins. Penang.
Matlae, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 301.—Am. austr.
membranaceum, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 309.—China.
metusefolium, Wall. Cat. n. 4850 = pulcherrimum.
microphyllum, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 272.—Malaya.
microphyllum, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 282 = parvifolium.
microphyllum, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 406.—Ins. Gebeh.
montanum, *Vieill. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 292.—N. Caled.
Moonii, Wight, Illustr. i. 129 = spectabile.
Nagassarium, Burm. f. Fl. Ind. 121 = Mesua ferrea.
oborale, *Miq. Fl. Ind. Bat. Suppl.* i. 498.—Sumatra.
odoratissimum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 12 = spectabile.
ovatifolium, Noronha, l. c. 18 = C. Inophyllum.
pachyphyllum, *Planch. & Triana, in Ann. Sc. Nat* Sér. IV. xv. (1861) 285.—Bras.
parviflorum, *Boj. ex Baker, Fl. Maurit.* 16.—Ins. Maurit.
parvifolium, *Choisy, in Mém. Soc. Hist. Nat. Par.* i. (1823) 329.—Ins. Molucc.
pisiferum, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 294 = retusum.
plicipes, Miq. Fl. Ind. Bat. Suppl. i. 499 = pulcherrimum.

CALOPHYLLUM:—
polyanthum, *Wall. Cat.* n. 4844; *Planch. & Triana, in Ann. Sc. Nat. Sér.* IV. xv. (1861) 278.—Ind. or.
pseudotacamacha, *Planch. & Triana, l. c.* 270.—Ind. or.
pulcherrimum, *Wall. Cat.* n. 4848; *Planch. & Triana, l. c.* 274.—Malaya.
retusum, *Wall. Cat.* n. 4846; *Planch. & Triana, l. c.* 265.—Malaya.
rigidum, *Miq. Fl. Ind. Bat. Suppl.* i. 497.—Sumatra.
salgoense, *Pierre, Fl. For. Cochinch.* t. 105.—Cochinch.
Sonaiatti, Burm. f. Fl. Ind. 121 = spectabile.
spectabile, Hook. & Arn. Bot. Beech. Voy. 174 = membranaceum.
spectabile, *Willd. in Ges. Naturf. Fr. Berl. Mag.* v. (1811) 80.—Malaya; Polynes.
spurium, Boj. Hort. Maurit. 89 = parviflorum.
spurium, Choisy, in Mém. Soc. Hist. Nat. Par. i. (1823) 229 = C. Inophyllum.
Suriga, *Buch.-Ham. ex Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 608.—Ind. or.
Suringa, Buch.-Ham. ex Steud. Nom. ed. II. i. 361 = pracc.
Tacamahaca, *Willd. in Ges. Naturf. Fr. Berl. Mag.* v. (1811) 79.—Borbon.; Madag.
tetrapetalum, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 608 = spectabile.
tetrapetalum, Wall. Cat. n. 4848 = Wallichianum.
tetrapterum, *Miq. Pl. Jungh.* 291.—Sumatra.
Teysmanni, *Miq. Fl. Ind. Bat. Suppl.* i. 499.—Sumatra.
Teysmannii, Zoll. ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 275 = lanceolatum.
thuriferum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 16.—Peruv.
Thorelii, *Pierre, Fl. For Cochinch. t.* 108.—Cochinch.
Thwaitesii, *Planch. & Triana, in Ann. Sc. Nat. Sér.* IV. xv. (1861) 260.—Zeylan.
tomentosum, *Wight, Illustr.* i. 128.—Ind. or.
trapezifolium, *Thw. Enum. Pl. Zeyl.* 51.—Zeylan.
venulosum, *Zoll. Syst. Verz. Ind. Archip.* 149, 150.—Java.
Walkeri, *Wight, Illustr.* i. 128.—Ind. or.
Wallichianum, *Planch. & Triana, in Ann. Sc. Nat. Sér.* IV. xv. (1861) 277.—Malaya.
Wightianum, *Wall. Cat.* n. 4847; *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 256.—Ind. or.

CALOPHYSA, DC. Prod. iii. 166 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 766.
ciliata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 140.—Am. trop.
dentata, *Triana, l. c.*—Peruv.
dispar, *Triana, l. c.* 141.—Peruv.
ferruginea, *Triana, l. c.*—Venezuela.
heterophylla, *Triana, l. c.* 140.—Peruv.
pilosa, *Triana, l. c.*—Am. trop.
setosa, *Triana, in Journ. Bot.* v. (1867) 209.—Nicaragua.
testiculata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 140.—N. Granat.
tococoides, *DC. Prod.* iii. 166.—Bras.
vesiculosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 140.—Mexic.

CALOPLECTUS, Cerst. in Videnak. Selsk. Skr. V. v. (1861) 119 = **Alloplectus**, Mart. (Melastom.).
macrophyllus, Cerst. l. c. = *A. macrophyllus.*

CALOPOGON, R. Br. in Ait. Hort. Kew. ed. II. v. 204 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 515.
CATHEA, Salisb. in Trans. Hort. Soc. i. (1812) 300, nomen prius.
Lesueurii, Bronga. Bot. Voy. Coquille, 188. t. 37 = Pogonia tetraphylla.
multiflorus, *Lindl. Gen. et Sp. Orch.* 425.—Am. bor.
pallidus, *Chapmann, Fl. S. U. St.* 457.—Am. bor.
parviflorus, *Lindl. Gen. et Sp. Orch.* 424.—Am. bor.
parviflorus, *Rafin. Atl. Journ.* 148 (= pracc. ?).—Am. bor.
pulchellus, *R. Br. in Ait. Hort. Kew.* ed. II. v. 204.—Am. bor.

CALOPOGONIUM, Desv. in Ann. Sc. Nat. Sér. 1. ix. (1826) 422. *LEGUMINOSAE*, Benth. & Hook. f. i. 534.
CYANOTREMMA, Benth. ex Hook. & Arn. Bot. Beech. Voy. 415 (1841).

CALOPOGONIUM:—
STENOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 136.
brachycarpum, Benth. ex Hemsl. in Biol. Centr. Am. Bot. i. 300 = mucunoides.
caeruleum, *Hemsl. l. c.* 301.—Mexic.
galactoides, *Benth. ex Hemsl. l. c.*—Mexic.
mucunoides, *Desv. in Ann. Sc. Nat.* Sér. 1. ix. (1826) 423.—Guiana.

CALOPSIS, Beauv. ex Juss. in Dict. Sc. Nat. xlv. 272 (1827); et ex Desv. in Ann. Sc. Nat. Sér. 1. xiii. (1828) 44. t. 2 = **Leptocarpus**, R. Br. (Restiac.).
chilensis, Steud. Syn. Pl. Cyp. 256 = L. chilensis.
festucacea, Kunth, Enum. Pl. iii. 425 = L. incurvatus.
hirtella, Kunth, l. c. 426 = L. peronatus.
incurvata, Kunth, l. c. 427 = L. incurvatus.
neglecta, Hochst. in Flora, xxviii. (1845) 388 = L. neglectus.
oxylepis, Kunth, Enum. Pl. iii. 427 = L. oxylepis.
paniculata, Desv. in Ann Sc. Nat. Sér. 1. xiii. (1828) 44. t. 2. f. 2 = L. paniculatus.
peronata, Kunth, Enum. Pl. iii. 426 = Restio ocreatus et L. peronatus.
ramiflora, Kunth, l. c. 428 = L. paniculatus.
triticea, Kunth, l. c. 424 = Restio triticeus.

CALOPTILLUM, Lag. Amen. Nat. i. 34 (1811) = **Nassauvia**, Juss. (Compos.).
Lagasca, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 96 = N. Lagascae.

CALOPYXIS, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 86 = **Combretum**, Linn.
alata, Tul. l. c. 89.—Madag.
eriantha, Tul. l. c. 90.—Madag.
oxygonia, Tul. l. c. 88.—Madag.
sphaerodes, Tul. l. c. 86.—Madag.
velutina, Tul. l. c. 87.—Madag.

CALORCHIS, Rodrig. Orch. Nov. i. 195 (1877). *ORCHIDEAE.*
phaenoleuca, *Rodrig. l. c.* 196.—Bras.

CALORHABDOS, Benth. Scroph. Ind. 44 (1835). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 963.
Brannoniana, *Benth. l. c.*—Reg. Himal.
caulopteris, *Hance, in Journ. Bot.* xv. (1877) 298.—China.

CALOROPHUS, Labill. Nov. Holl. Pl. ii. 78. t. 228 (1806) = **Hypolaena**, R. Br. (Restiac.).
axpor, Nees, in Lehm. Pl. Preiss. ii. 67 = Restio Loxocarya.
crispatus, Nees, l. c. = Restio crispatus.
dense, Nees, l. c. = H. dense.
dentatus, Nees, l. c.—Austral.
elongatus, Labill. Nov. Holl. Pl. ii. 78. t. 228 = H. elongata.
elongatus, F. Muell. Fragm. viii. 86 = H. elongata.
elongatus, Nees, in Lehm. Pl. Preiss. ii. 68 = H. gracillima.
flexuosus, Nees, in Lehm. Pl. Preiss. ii. 68 = H. flexuosa.
minor, Hook. f. Fl. N. Zel. i. 267 = H. elongata.
Sieberianus, Steud. Syn. Pl. Cyp. 265 = Restio fastigiatus.

CALOSACME, Wall. Cat. nn. 800–806 (1829) = **Chirita**, Buch.-Ham. (Gesnerac.).
acuminata, Wall. l. c. n. 802 = Ch. acuminata et Didymocarpus acuminata.
amplexicaulis, Wall. l. c. n. 806 = Ch. bifolia.
dimidiata, Wall. l. c. n. 803 = Ch. dimidiata.
flava, Wall. l. c. n. 801 = Ch. pumila.
grandiflora, Wall. l. c. n. 800 = Ch. urticaefolia.
macrophylla, Wall. l. c. n. 805 = Ch. macrophylla.
polycarpa, Wall. l. c. n. 804 = Lysionotus serrata.

CALOSANTHES, Blume, Bijdr. 760 (1826) = **Croxyluna**, Vest. (Bignoniac.).
indica, Blume, l. c. = O. indicum.

CALOSANTHOS, Reichb. in Mossel. Handb. ed. II. i. 80 (1837) = **KALOSANTHES**, Haw. = **Rochea**, DC. (Crassul.).

CALOSCIADIUM, Endl. Gen. Suppl. iv. 111. 7 (1856) = **Aciphylla**, Forst. (Umbellif.).
antipodum, Endl. ex Walp. Ann. iii. 702.—N. Zel.
latifolium, Endl. ex Walp. l. c.—N. Zel.

CALOSCILLA, Jord. & Fourr. Ic. Fl. Eur. ii. 14. t. 246 (1869) = **Scilla**, Linn. (Liliac.).
Clusii, Jord. & Fourr. l. c. = S. peruviana.
elegans, Jord. & Fourr. l. c. 16. t. 250 = S. peruviana.
flexuosa, Jord. & Fourr. l. c. 17. t. 254 = S. peruviana.
grandiflora, Jord. & Fourr. l. c. 16. t. 252 = S. peruviana.
hispanensis, Jord. & Fourr. l. c. 17. t. 255 = S. peruviana.
Hughii, Jord. & Fourr. l. c. 15. t. 246 = S. peruviana.
lirioides, Jord. & Fourr. l. c. = S. peruviana.
pallidiflora, Jord. & Fourr. l. c. 17. t. 248 = S. peruviana.
subcilioides, Jord. & Fourr. l. c. 16. t. 253 = S. peruviana.
subcurnosa, Jord. & Fourr. l. c. t. 250 = S. peruviana.
venusta, Jord. & Fourr. l. c. 15. t. 247 = S. peruviana.

CALOSCORDUM, Herb. in Lindl. Bot. Reg. (1844) Misc. 66 = **Nothoscordum**, Kunth (Liliac.).
neriniflorum, Herb. l. c. 67 = N. neriniflorum.

CALOSERIS, Benth. Pl. Hartw. 88 (1841) = **Onoseris**, DC. (Compos.).
rupestris, Benth. l. c. = O. isotypus.

CALOSMON, J. S. Presl, Rostl. li. 71 (1823) = **Lindera**, Thunb. (Laurin.).
aestivale, Kostel. Allg. Med.-Pharm. Fl. ii. 478 = L. Benzoin.
Benzoin, J. S. Presl, Rostl. ii. 71 = L. Benzoin.
Nessiana, Kostel. Allg. Med.-Pharm. Fl. ii. 478 = L. Nessiana.

CALOSPHACE, Rafin. Fl. Tellur. iii. 91 = **Salvia**, Tourn. (Labiat.).

CALOSTELMA, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. sub t. 184 (1833) = **Lidaria**, Schreb. (Compos.).
elegans, D. Don, l. c. = L. elegans.

CALOSTEMMA, R. Br. Prod. 297 (1810) *AMARYLLIDEAE*, Benth. & Hook. f. iii. 735.
album, *R. Br. l. c.* 298.—Austral.
aureum, Link, Enum. Hort. Berol. i. 319 = luteum.
candidum, Lindl. in Mitch. Three Exped. i. 54 = luteum.
carneum, Lindl. l. c. ii. 39 = purpureum.
Cunninghami, *Ait. ex Loud. Hort. Brit. Suppl.* ii. 616. —Austral.
luteum, *Sims, Bot. Mag.* t. 2101.—Austral.
purpureum, *R. Br. Prod.* 298.—Austral.

CALOSTEPHANE, Benth. in Hook. Ic. Pl. t. 1111 (1872). *COMPOSITAE*, Benth. & Hook. f. ii. 335.
divaricata, *Benth. l. c.*—Afr. trop.

CALOSTIGMA, Deene. in Ann. Sc. Nat. Sér. II. ix. (1838) 348. t. 12. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 792.
Burchellii, *Fourn. in Mart. Fl. Bras.* vi. iv. 286.—Bras.
glabrum, *Decne. in DC. Prod.* viii. 561.—Bras.
Glaziovii, *Fourn. in Mart. Fl. Bras.* vi. iv. 286.—Bras.
Guillemininum, *Decne. in DC. Prod.* viii. 561.—Bras.
insigne, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 348.—Bras.

CALOSTIGMA, Schott, Meiet. Bot. 19 (1832) = **Philodendron**, Schott (Aroid.).
Imbe, Schott, l. c. = P. Imbe.

CALOSTIMA, Rafin. Fl. Tellur. iii. 47 (1836) = **Urera**, Gaudich. (Urtic.).
aculeata, Rafin. l. c. 46 = U. baccifera.

CALOSTROPHUS, F. Muell. Fragm. viii. 86 (1875) = **Hypolaena**, R. Br. (Restiac.).
elongatus, F. Muell. l. c. = H. longissima.
elongatus, Mast. in DC. Monog. Phan. i. 872 = H. laterifloru.

CALOSTROPHUS:—
annuicus, F. Muell. Census, 194 = H. exsulca.
fastigiatus, F. Muell. in Papers Roy. Soc. Tasm. (1878) 117 = H. fastigiata.
gracillimus, F. Muell. Fragm. viii. 88 = H. gracillima.
laterifloros, F. Muell. l. c. 87 = H. laterifiora.
Siberianus, Mast. in DC. Monog. Phan. i. 378 = H. lateriflora.

CALOTHAMNUS, Labill. Nov. Holl. Pl. ii. 25. t. 164 (1806). *MYRTACEAE*, Benth. & Hook. f. i. 706.
BAUDINIA, Leschen. ex DC. Prod. iii. 211 (1828).
BILLOTTIA, Colla, Hort. Ripul. 20. t. 28 (1824).
affinis, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 346 = microcarpus ?
asper, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 35.—Austral.
Baxteri, Hort. Angl. ex Klotzsch, in Otto & Dietr. Allg. Gartenz. iv. (1836) 115 = Kunzea Baxteri.
blephaxrantherus, *F. Muell. Fragm.* iii. 112.—Austral.
blephaxspermus, *F. Muell. l. c.* 111.—Austral.
chrysantherens, *F. Muell. l. c.* 112.—Austral.
clavatus, Mackay, ex Lodd. Bot. Cab. t. 1447 = quadrifidus.
ericacpus, Lindl. Swan River App. 9 = sanguineus.
glibbosus, *Benth. Fl. Austral.* iii. 176.—Austral.
Gilesii, *F. Muell. Fragm.* x. 81.—Austral.
gracilis, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.
homalophyllus, *F. Muell. Fragm.* iii. 111.—Austral.
Huegelii, Schau. in Nov. Act. Nat. Cur. xxi. (1845) 80 = lateralis.
Knightii, Hort. ex Schau. l. c. 28 = robusta.
lanuginosus, Schau. l. c. 26 = quadrifidus.
lateralis, *Lindl. Swan River App.* 9.—Austral.
laxus, Kunze, in Linnæa, xx. (1847) 56 = Preissii.
Lehmanni, *Schau. in Nov. Act. Nat. Cur.* xxi. (1845) 97.—Austral.
longifolius, Lehm. Del. Sem. Hort. Hamb. (1842) 7 = lateralis.
longissimus, *F. Muell. Fragm.* iii. 113.—Austral.
microcarpus, *F. Muell. l. c.* 113.—Austral.
nodosus, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 168 = lateralis.
Oldfieldii, *F. Muell. Fragm.* iii. 113.—Austral.
pachystachyus, *Benth. Fl. Austral.* iii. 178.—Austral.
pinifolius, *F. Muell. Fragm.* iii. 158.—Austral.
planifolius, *Lehm. Del. Sem. Hort. Hamb.* (1842) 7.—Austral.
plumosus, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 168 = Lehmanni.
Preissii, *Schau. in Nov. Act. Nat. Cur.* xxi. (1845) 27.—Austral.
purpureus, Endl. in Enum. Pl. Hueg. 48 = quadrifidus.
quadrifidus, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.
robustus, Schau. in Nov. Act. Nat. Cur. xxi. (1845) 22 = villosus.
rupestris, *Schau. l. c.*—Austral.
sanguineus, *Labill. Nov. Holl. Pl.* ii. 25. t. 164.—Austral.
Schaueri, *Lehm. Del. Sem. Hort. Hamb.* (1842) 7.—Austral.
schanostylus, Schau. in Nov. Act. Nat. Cur. xxi. (1845) 29 = Schaueri.
spathulatus, Hort. ex Steud. Nom. ed. II. i. 361 = Kunzea Baxteri.
suberosus, Schau. in Lehm. Pl. Preiss. i. 156 = Melaleuca suberosa.
torulosus, *Schau. in Nov. Act. Nat. Cur.* xxi. (1845) 21.—Austral.
villosus, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.

CALOTHECA, Desv. in Nov. Bull. Soc. Philom. ii. (1810) 190 = **Briza**, Linn. (Gramin.).
barbata, Nees, Agrost. Bras. 477 = B. barbata.
bricoides, Beauv. Agrost. 86. t. 17. f. 6 = B. erecta.
brizoides, Desv. Journ. Bot. i. (1813) 71 = B. Lamarckiana.
bromoides, Lejeune, Messager des Sciences (1828); et in Nov. Act. Nat. Cur. xii. (1825) 757 = Bromus arduennensis.
dilatata, Link, Hort. Berol. i. 157 = B. erecta.
elegans, Beauv. Agrost. 157. t. 17 = B. elegans.
macrostachya, Presl, Rel. Haenk. i. 268 = B. macrostachya.
microstachya, Presl, l. c. = B. microstachya.
montevidensis, Spreng. ex Steud. Nom. ed. II. i. 261, 225 = B. subaristata.
reniformis, Presl, Rel. Haenk. i. 268 = B. reniformis.
rotundata, Roem. & Schult. Syst. ii. 682 = B. rotundata.
sabulosa, Steud. in Flora, xli. (1829) ii. 488 = Sporobolus virginicus.
stricta, Hook. & Arn. Bot. Beech. Voy. 50 = B. stricta.
triloba, Kunth, Enum. Pl. i. 374 = B. triloba.

CALOTHYRSUS, Spach, in Ann. Sc. Nat. Sér. II. ii. (1844) 69 = **Acaulus**, Linn. (Sapindac.).
californica, Spach, l. c. = A. californica.

CALOTIS, R. Br. in Bot. Reg. t. 504 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 267.
CHRYSOLOMA, F. Muell. in Linnæa, xxv. (1852) 401.
GONIOPOGON, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 178.
HUENEFELDIA, Walp. in Linnæa, xiv. (1840) 807.
anthemoides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 44.—Austral.
brevicula, *Benth. in Enum. Pl. Hueg.* 60.—Austral.
cuneifolia, *R. Br. in Bot. Reg.* t. 504.—Austral.
cymbacantha, *F. Muell. in Linnæa*, xxv. (1852) 400.—Austral.
dentex, *R. Br. in Bot. Reg.* t. 504.—Austral.
dilatata, A. Cunn. in DC. Prod. v. 302 = cuneifolia.
erinacea, *Steetz, in Lehm. Pl. Preiss.* i. 424.—Austral.
glandulosa, *F. Muell. in Trans. Vict. Inst.* (1855) 129.—Austral.
hispidula, *F. Muell. l. c.* 130.—Austral.
Kempei, *F. Muell. in Trans. Roy. Soc. S. Austral.* iv. (1881) 112.—Austral.
lappulacea, *Benth. in Enum. Pl. Hueg.* 60.—Austral.
microcephala, *Benth. Fl. Austral.* iii. 504.—Austral.
microphylla, *Benth. in Enum. Pl. Hueg.* 60.—Austral.
Muelleri, Sond. in Linnæa, xxv. (1852) 470 = scabiosifolia.
palmata, *Gray, in Proc. Am. Acad.* v. (1861) 191.—Austral.
plumulifera, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 57.—Austral.
polycnemoides, Sond. in Linnæa, xxv. (1852) 470 = lappulacea.
porphyroglossa, *F. Muell. ex Benth. Fl. Austral.* iii. 505.—Austral.
pterosperma, *R. Br. ex Benth. l. c.*—Austral.
scabiosifolia, *F. Muell. in Linnæa*, xxv. (1852) 471.—Austral.
scapigera, *Hook. in Mitch. Journ. Trop. Austral.* 75.—Austral.
tropica, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 58 = brevisceta.

CALOTROPIS, R. Br. in Mem. Wern. Soc. i. (1809) 39. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 754.
Acia, *Buch.-Ham. in Trans. Linn. Soc.* xiv. (1825) 247.—Ind. or.
gigantea, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 78.—Ind. or.
Hamiltonii, Wight, Contrib. Ind. Bot. 58 = procera.
herbacea, Wight, l. c. 54 = C. Acia.
heterophylla, Wall. Cat. n. 2317 = procera.
procera, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 78.—Peruv.; Afr. trop.; Ind. or.
Susuuka, G. Don, Gen. Syst. iv. 147 = Hoya imperialis ?
Wallichii, Wight, Contrib. Ind. Bot. 58 = procera.

CALPANDRIA, Blume, Bijdr. 178 (1825) = **Camellia**, Linn. (Ternstroem.).
lanceolata, Blume, l. c. = Cam. lanceolata.
Quiescensure, Korth. in Verh. Nat. Gesch. Bot. 149 = Cam. Quinosaura.

CALPICARPUM, G. Don, Gen. Syst. iv. 100 (1837) = **Kopsia**, Blume (Apocyn.).
albiflorum, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 402.—Malaya.
arcuatum, G. Don, Gen. Syst iv. 100 = K. fruticosa.
Roxburghii, G. Don, l. c. = K. fruticosa.

CALPIDIA, Thou. Hist. Vég. Iles Afr. Austr. (1805) 37. t. 10 = **Plectia**, Linn. (Nyctagin.).
costata, Boj. Hort. Maurit. 265 = P. costata.
lanceolata, Thou. Hist. Vég. Iles Afr. Austr. 37. t. 10 = P. Calpidia.
macrophylla, G. Don, Gen. Syst. iv. 100 = P. macrophylla.
ovalifolia, Boj. l. c. = P. Calpidia.

CALPIGYNE, Blume, Mus. Bot. Lugd. Bat. ii. 198 (1856). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 323.
fruticosa, *Blume, l. c.*—Borneo.

CALPURNIA, E. Mey. Comm. Pl. Afr. Austr. 2 (1836). *LEGUMINOSAE*, Benth. & Hook. f. i. 554.
aurea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 282.—Afr. trop.
aurea, Benth. in Ann. Wien. Mus. ii. (1840) 90 = laiogyne.
capensis, Steud. Nom. ed. II. i. 392 = Virgilia capensis.
floribunda, *Harv. in Harv. & Sond. Fl. Cap.* ii. 267.—Afr. austr.
intrusa, *E. Mey. Comm. Pl. Afr. Austr.* 2.—Afr. austr.
laiogyne, *E. Mey. l. c.* 3.—Afr. austr.
robinioides, *E. Mey. l. c.*—Afr. austr.
sericea, *Harv. in Harv. & Sond. Fl. Cap.* ii. 267.—Afr. austr.
sylvatica, *E. Mey. Comm. Pl. Afr. Austr.* 2.—Afr. austr.
villosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 268.—Afr. austr.

CALSIAMA, Rafin. Sylva Tellur. 12 (1838) = **Odina**, Roxb. (Anacard.).
malabarica, Rafin. l. c. = O. Wodier.

CALTHA, Tourn. ex Adans. Fam. ii. 296 (1763) = **Calendula**, Linn. (Compos.).
arvensis, Moench, Meth. 586 = Calend. arvensis.
dentata, Moench, Meth. Suppl. 238 = Calend. stellata.
graveolens, Gilib. Fl. Lituan. i. 212 = Calend. arvensis.
officinalis, Moench, Meth. 586 = Calend. officinalis.
officinalis, Scop. Fl. Carn. ed. II. ii. 167 = Calend. arvensis.
stellata, Moench, Meth. Suppl. 238 = Calend. stellata.

CALTHA, Tourn. ex Adans. Fam. ii. 296 (1763) = **Calendula**, Linn. (Compos.).
arvensis, Moench, Meth. 586 = Calend. arvensis.
dentata, Moench, Meth. Suppl. 238 = Calend. stellata.
graveolens, Gilib. Fl. Lituan. i. 212 = Calend. arvensis.
officinalis, Moench, Meth. 586 = Calend. officinalis.
officinalis, Scop. Fl. Carn. ed. II. ii. 167 = Calend. arvensis.
stellata, Moench, Meth. Suppl. 238 = Calend. stellata.

CALTHA, [Rupp.] Linn. Syst. ed. I (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 6.
PSYCHROPHILA, Bercht. & Presl, Rostl. i. Ranunc. 80 (1832).
PSYCROPHILA, Rafin. Atl. Journ. 144 (1832).
THACLA, Spach, Hist. Vég. Phan. vii. 295 (1839).
alba, Jacquem. Voy. Bot. 6 = palustris.
alpestris, Schott, Nym. & Kotschy, Analect. Bot. 33 = palustris.
alpina, Schur, Enum. Pl. Transs. 25 = palustris.
andicola, *C. Gay, Fl. Chil.* i. 49. t. 2.—Chili.
appendiculata, Lechl. ex C. Muell. in Walp. Ann. vii. 26 = limbata.
appendiculata, *Porr. Syn.* ii. 107.—Reg. Magell.
arctica, R. Br. in Parry, 1st Voy. App. 265 = palustris.
asarifolia, DC. Syst. i. 309 = palustris.
baicalensis, Demid. ex Steud. Nom. ed. II. i. 262 = natans.
biflora, DC. Syst. i. 310 = palustris.
Bisma, Buch.-Ham. in Brewst. Edinb. Journ. Sc. i. (1824) 351 = Aconitum palustrum.
camtschatica, Spreng. Syst. ii. 660 = Oxygraphis glacialis.
Codua, Buch.-Ham. in Brewst. Edinb. Journ. Sc. i. (1824) 351 = Aconitum ferox.

CALTHA:—
cornuta, Schott, Nym. & Kotschy, Analect. Bot. 31 = palustria.
coronata, *Sckur, Enum. Pl. Transs.* 35.—Transsylvan.
dentata, Mühl. Cat. ed. II. 56 = palustria.
De-Ranco, *Stend. in Flora*, xxxix. (1856) 407.—Chili.
dionaeafolia, *Hook. f. in Hook. Lond. Journ. Bot.* ii. (1843) 806.—Reg. Magell.
emodorum, Spreng. Syst. iv. Cur. Post. 220 = palustris.
ficarioides, Pursh, Fl. Am. Sept. ii. 389 = palustris.
flabellifolia, Pursh, l. c. 390. t. 17 = palustris.
glabra, Gilib. Fl. Lituan. ii. 379 = palustris.
glacialis, Spreng. Syst. ii. 660 = Oxygraphis glacialis.
Gouaniana, Wall. Cat. n. 4710 = palustris.
grosse-serrata, Pantot. in Verh. Ver. Naturw. Heilk. Presb. ii. (1871–72) 86 = palustris.
Guerangeri, Bor. in Billot, Annot. 11 = palustris.
himalensis, D. Don, Prod. Fl. Nep. 195 = palustris.
holophylla, *Leyb. in Anal. Univ. Chil.* (1859) 678.—Chili.
integerrima, Pursh, Fl. Am. Sept. ii. 390 = palustris.
intermedia, Schott, Nym. & Kotschy, Analect. Bot. 32 = palustris.
introloba, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 98.—Austral.
kamtschatica, Spreng. Syst. ii. *666 = asarifolia.*
laeta, Schott, Nym. & Kotschy, Analect. Bot. 32 = palustris.
laxopetala, Steud. Nom. ed. II. i. 189, sphalm. = leptosepala.
latifolia, Schott, Nym. & Kotschy, Analect. Bot. 32 = palustris.
leptosepala, *DC. Syst.* i. 310.—Am. bor. occ.
leptostachya, G. Don, Gen. Syst. i. 44 = prsec.
limbata, *Schlecht. in Linnaea*, xxvii. (1854) 556.—Chili.
major, Mill. Gard. Dict. ed. VIII. n. 1 = palustris.
minor, Mill. l. c. n. 2 = palustris.
multicapsularis, Sond. ex DC. Prod. i. 44 = sagittata.
natans, *Pall. Reise*, iii. 246.—Reg. bor.
Nirbisia, Buch.-Ham. in Brewst. Edinb. Journ. Sc. i. (1824) 251 = Aconitum ferox.
Novae-Zelandiae, *Hook. f. Fl. Nov. Zeal.* i. 12. t. 6.—N. Zel.
orthorhyncha, *Rupr. Fl. Caucas.* 28.—Reg. Caucas.
pallidiflora, Maxim. in Bull. Soc. Bot. Fr. (1862) 180 = palustris.
palustris, *Linn. Sp. Pl.* 558.—Reg. bor. temp. et arct.
paniculata, Wall. Cat. n. 4711 = palustris.
paradoxa, T. F. Forst. in Trans. Linn. Soc. viii. (1807) 324 = appendiculata.
parnassifolia, Raf. in Med. Rep. New York v. (1808) 361 = palustris.
polypetala, *Hochst. in Lorent, Wanderungen*, 289.—Ab. Min. ; Persia.
Populago, Golan, Hort. Monsp. 268 = palustris.
pumila, Schur, Enum. Pl. Transs. 26 = palustris.
radicans, *T. F. Forst. in Trans. Linn.* Soc. viii. (1807) 324. t. 17.—Scotia.
ranunculoides, Schur, Enum. Pl. Transs. 26 = palustris.
riparia, G. Don, Gen. Syst. i. 44 = palustris.
sagittata, *Cav. Ic.* v. 8, 456. t. 414.—Reg. Magell.
scaposa, *Hook. f. & Thoms. Fl. Ind.* i. 40.—Reg. Himal.
vulgaris, Pall. ex Stend. Nom. ed. II. i. 269 = palustris.
vulgaris, Schott, Nym. & Kotschy, Analect. Bot. 33 = palustris.

CALTHOIDES, B. Juss. ex DC. Prod. vi. 473 (1837) = Othonna, Linn. (Compos.).

CALUCECHINUS, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 19 (1853) = **Fagus**, Tourn. (Cupulif.).
antarctica, Hombr. & Jacquinot, l. c. = F. antarctica.
Menziesii, Hombr. & Jacquinot, l. c. = F. antarctica.

CALUSIA, Bert. ex Steud. Nom. ed. II. i. 292 (1840) = Myrospermum, Jacq. (Legumin.).
emarginata, Bert. ex Steud. l. c. ii. 175 = M. frutescens.

CALUSPARASSUS, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 20 (1853) = **Fagus**, Tourn. (Cupulif.).

CALUSPARASSUS :—
betuloides, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 20 = F. betuloides.
Forsteri, Hombr. & Jacquinot, l. c. = F. betuloides.
Pumilio, Hombr. & Jacquinot, l. c. 21 = F. betuloides.

CALVARIA, Comm. ex Gaertn. f. Fruct. Suppl. 116. t. 200, 201 (1808) (Gen. dub.).
cerebellina, Comm. ex Gaertn. f. l. c.—Hab. ?
globosa, Gaertn. f. l. c. 117.—Hab. ?
hexangularis, Gaertn. f. l. c.—Hab. ?
major, Gaertn. f. l. c. 116 = *cerebellina* ?

CALVELIA, Moq. in DC. Prod. xiii. ii. 167 (1849) = Suaeda, Forsk. (Chenopod.).
pterantha, Moq. l. c. 168.—Soongar.

CALVOA, Hook. f. in Benth. & Hook. f. Gen i. 755 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
crassinodis, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 76.—Afr. trop.
hirsuta, *Hook. f. ex Triana, l. c.*—Afr. trop.
sinuata, *Hook. f. ex Triana, l. c.*—Afr. trop.

CALYCADENIA, DC. Prod. v. 695 (1836) = Hemisonia, DC. (Compos.).
Cephalea, DC. l. c. = H. multiglandulosa.
Fremontii, A. Gray, in Torr. Bot. Mex. Bound. 101, in note = H. multiglandulosa.
Hartwegii, Lem. Illustr. Hortic. (1855) Misc. 9 = Odontadenia speciosa.
mollis, A. Gray, in Proc. Am. Acad. vii. (1868) 360 = H. mollis.
multiglandulosa, DC. Prod. v. 695 = H. multiglandulosa.
pauciflora, A. Gray, in Torr. Bot. Mex. Bound. 101, in note = H. paucifloras.
pinnosa, Kellogg, in Proc. Calif. Acad. v. (1873) 49 = H. pinnosa.
tenella, Torr. & Gray, Fl. N. Am. ii. 409 = H. tenella.
truncata, DC. Prod. v. 695 = H. truncata.
villosa, DC. l. c. = H. Douglasii.

CALYCAMPE, Berg. in Linnaea, xxvii. (1854) 129 = Myrcia, DC. (Myrtac.).
angustifolia, Berg, l. c. 131 = M. bebepetala.
latifolia, Berg, l. c. 130.—Guiana.

CALYCANDRA, Lepr. ex A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 50 et 322. t. 9 (1832) = Cordyla, Lour. (Legumin.).
pinnata, A. Rich. l. c. 51. t. 9 = Cord. africana.

CALYCANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 243. t. 40 (1861) = Ipomoea, Linn. (Convolvul.).
leucanthemum, Klotzsch, l. c. 244.—Afr. trop.

CALYCANTHUS, Linn. Syst. ed. X. 1066 (1759).
CALYCANTHACEAE, Benth. & Hook. f. l. 16.
BASTERIA, Mill. ex Adans. Fam. ii. 394 (1763).
BEURRERIA, Ehret, Pl. et Pap. Rar. t. 13 (1765).
BUTNERIA, Duham. Arb. i. 113. t. 42 (1765).
POMPADOURA, Buchoz, ex DC. Prod. iii. 2 (1828).
acuminatus, Hort. ex C. Koch, Hort. Dendrol. 185 = floridus.
asplenifolius, Hort. ex C. Koch, Dendrol. i. 365 = floridus.
ballatus, Hort. ex C. Koch, l. c. = floridus.
fenzlianus, Fl. Bor. Am. i. 205 = laevigatus.
fertilis, Andr. Rep. viii. t. 539 = laevigatus.
fertilis, Walt. Fl. Carol. 151 = glaucus.
floridus, *Linn. Syst. ed. X.* 1066.—Am. bor.
glaucus, *Willd. Enum. Hort. Berol.* 559.—Am. bor.
inodorus, Ell. Sketch. i. 576 = laevigatus.
laevigatus, *Willd. Hort. Berol.* t. 80; *Enum. Hort. Berol.* 559.—Am. bor.
macrophyllus, Hort. ex C. Koch, Dendrol. i. 364 = occidentalis.
mollis, *Rafin. Alsog. Am.* 7.—Am. bor.
nanus, Loisel. in Duham. ed. Nov. i. 219 = laevigatus.
oblongifolius, *Loud. Hort. Brit.* 214.—Am. bor.
occidentalis, *Hook. & Arn. Bot. Beech. Voy.* 340. t. 84.—Calif.
pensylvanicus, *Ladd. in Loud. Hort. Brit.* 214.—Am. bor.

CALYCANTHUS :—
philadelphicus, *Nois. ex Steud. Nom.* ed. II. i. 282.—Am. bor.
praecox, Linn. Sp. Pl. ed. II. 718 = Chimonanthus fragrans.
pumilionis, *Sweet, Hort. Brit.* ed. II. 195, nomen.—Carolina.
reticulatus, *Rafin. Alsog. Am.* 7.—Am. bor.
sterilis, Walt. Fl. Carol. 151 = floridus.
suaveolens, Salisb. Prod. 379 = Chimonanthus fragrans.
tomentosus, *Rafin. Alsog. Am.* 7.—Am. bor.
verrucosus, *Rafin. l. c.* 8.—Am. bor.
versicolor, *Regel, Cat. Pl. Hort. Absakov.* 25.—Am. bor.

CALYCERA, Cav. Ic. iv. 84. t. 358 (1797) (Calicera) *CALYCEREAE*, Benth. & Hook. f. ii. 462.
ANOMOCARPUS, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 351 ; Contrib. Bot. ii. 38.
DISCOPHYTUM, Miers, ex Lindl. Veg. Kingd. 701 (1847).
LYCNOCAULUS, Phil. in Linnaea, xxviii. (1856) 705.
LEUCOCERA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 382.
balsamitaefolia, *Rich. in Mem. Mus. Par.* vi. (1820) 77.—Chili.
Calcitrapa, *Griseb. in Gentt. Abh.* xix. (1874) 163.—Reg. Argent.
Cavanillesii, Rich. in Mem. Mus. Par. vi. (1820) 77 = herbacea.
eryngioides, *Remy, in C. Gay, Fl. Chil.* iii. 254.—Chili.
herbacea, Cav. ic. iv. 84. t. 358.—Chili.
intermedia, *Phil. in Anal. Univ. Chil.* (1870) i. 174.—Chili.
involucrata, *Phil. Sert. Mend. Alt.* 16.—Chili.
multicaulis, Phil. ex Miers, in Ann. & Mag. Nat. Hist Ser. III. vi. (1860) 399.—viridiflora.
palvinata, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 362.—Bolivia.
sessiliflora, *Phil. in Linnaea*, xxviii. (1856) 706.—Chili.
sinuata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III vi. (1860) 398.—Chili.
spinulosa, *Gill. ex Miers, l. c.* 399.—Chili.
squarrosa, *Miers, l. c.* 396.—Chili.
tenuifolia, *Phil. ex Miers, l. c.* 356.—Chili.
ventosa, *Meyen, Reise*, i. 356.—Chili.
viridiflora, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 399.—Chili.

CALYCIUM, Ell. Sketch. ii. 339 (1824) = Heterotheca, Cass. (Compos.).
divaricatum, Ell. l. c. 339, 338 = H. Lamarckii.
scabrum, Ell. l. c. 339 = H. Lamarckii.

CALYCOBOLUS, Willd. ex Roem. & Schult. Syst. v. 2 (1819) = Breweria, R. Br. (Convolvul.).
emarginatus, Willd. l. c. 4.—N. Granat.
pulchellus, Willd. l. c.—N. Granat.

CALYCOCARPUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 48 (1838). *MENISPERMACEAE*, Benth. & Hook. f. i. 85.
Lyonii, *Nutt. ex A. Gray, Gen. Am. Bor.* i. 76.—Am. bor.

CALYCOCORSUS, F. W. Schmidt, Samml. Phys. Aufs. i. 271 (1795) = Chondrilla, Linn. (Compos.).
Stenacoleium, F. W. Schmidt, l. c.—Ch. Fabiformis.

CALYCOCRADOPHNE, Boj. Hort. Maurit. 278 (1837) = Cootoea, Adrib. (Laurin.).
cupularis, Boj. l. c. = C. floribunda.
dysoxylon, Boj. l. c. = C. floribunda.
floribunda, Boj. l. c. = C. floribunda.

CALYCODON, Nutt. in Journ. Acad. Phil. Ser. II. i. 186 (1847) = Muehlenbergia, Schreb. (Gramin.).
montanum, Nutt. l. c. = M. gracilis.

CALYCOGONIUM, DC. Prod. iii. 186 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 761.
CALYCOSTEMA, Rich. ex DC. Prod. iii. 186 (1828).
angulatum, *Griseb. Cat. Pl. Cub.* 93.—Cuba.
dichaenicoides, *Griseb. l. c.* 94.—Cuba.
ciliosum, *Griseb. l. c.* 93 = dichaenicoides.

CALYCOGONIUM:—
cubense, A. Rich. Ess. Fl. Cub. 555 = Pachyanthus Poiretii.
echinatum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 184.—Cuba.
elliptícum, *Sauvalle, Fl. Cub.* 80.—Cuba.
glabratum, *DC. Prod.* iii. 168.—Jamaic.
Griesbachii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 97.—Cuba.
heterophyllum, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 85.—Cuba.
heterotrichum, *Triana, in Trans. Linn Soc.* xxviii. (1871) 96.—Cuba.
hypoglaucum, *Wright, ex Griseb. Cat. Pl. Cub.* 94.—Cuba.
insolúcratum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 184 = Cesaea involucrata.
lanceolatum, Griseb. Cat. Fl. Cub. 98 = glabratum.
Lima, Griseb. l. c. 95 = Oxymeris asperifolia.
Lindenianum, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 84.—Cuba.
microphyllum, Sauvalle, Fl. Cub. 80 = glabratum.
muricatum, Griseb. Cat. Fl. Cub. 95 = Octopleura muricata.
pauciflorum, Triana, in Trana. Linn. Soc. xxviii. (1871) 96 = echinatum.
plicatum, *Griseb. Cat. Pl. Cub.* 94.—Cuba.
productum, Sauvalle, Fl. Cub. 49 = heterophyllum.
rhamnoídeum, Griseb. Cat. Pl. Cub. 98 = glabratum.
rhamnoídeum, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 85.—Jamaic.
rosmarinifolium, *Griseb. Cat. Pl. Cub.* 98.—Cuba.
stellatum, *DC. Prod.* iii. 168.—Ins. S. Doming.
stellatum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 183 = Griesebachii.
verrucosum, *Griseb. Cat. Pl. Cub.* 96.—Cuba.
Wrightianum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 184 = glabratum.

CALYCOLPUS, Berg, in Linnaea, xxvii. (1854) 878. *MYRTACEAE*, Benth. & Hook. f. i. 713.
calophyllus, *Berg, in Mart. Fl. Bras.* xiv. i. 412.—Bras.
glaber, *Berg, l. c.* 413.—Peruv.
Goetheanus, *Berg, l. c.* 412.—Bras.
Kegelianus, *Berg, in Linnaea*, xxvii. (1854) 861.—Guiana.
ovalifolius, *Berg, l. c.* 879.—Ind. occ.
parviflorus, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 181.—Guiana.
revolutus, *Berg; in Linnaea*, xxvii. (1854) 883.—Guiana.
Schomburgkianus, *Berg, l. c.* 880.—Guiana.
Warscewiczianus, *Berg, l. c.* 882.—N. Granat.

CALYCOMELIA, Kostel. Alg. Med.-Pharm. Fl. iii. 1003 (1884) = **Fraxinus**, Linn. (Oleac.).
acuminata, Kostel. l. c. 1004 = F. americana.
alba, Kostel. Ind. Hort. Prag. 26 = F. alba.
americana, Kostel. Alg. Med.-Pharm. Fl. iii. 1004 = F. americana.
caroliniana, Kostel. Ind. Hort. Prag. 26 = F. platycarpa.
elíptica, Kostel. l. c. = F. elliptica.
epíptera, Kostel. l. c. = F. epiptera.
expansa, Kostel. l. c. = F. pubescens.
juglandifolia, Kostel. Alg. Med.-Pharm. Fl. iii. 1004 = F. americana.
lancea, Kostel. Ind. Hort. Prag. 26 = F. lancea.
nigra, Kostel. l. c. = F. nigra.
ovata, Kostel. l. c. = F. ovata.
pallida, Kostel. l. c. = F. pallida.
platycarpa, Kostel. l. c. = F. platycarpa.
pubescens, Kostel. l. c. = F. pubescens.
quadrangulata, Kostel. Alg. Med.-Pharm. Fl. iii. 1004 = F. quadrangulata.
Richardi, Kostel. Ind. Hort. Prag. 26 = F. Richardi.
timentosa, Kostel. Alg. Med.-Pharm. Fl. iii. 1004 = F. pubescens.
viridis, Kostel. Ind. Hort. Prag. 26 = F. viridis.

CALYCOMIS, R. Br. in Flind. Voy. App. 549 (1814) = **Callicoma**, Andr. (Saxifrag.).

CALYCOMIS, D. Don, in Edinb. N. Phil. Journ. ix. (1830) 93 = **Astrophyllum**, Benth. (Saxifr.).
verticillatum, D. Don. l. c. = A. venosum.

CALYCOMORPHUM, Presl, Symb. Bot. i. 50 (1830) = **Trifolium**, Tourn. (Legumin.).
globosum, Presl, l. c. = T. globosum.
subterraneum, Presl, l. c. = T. subterraneum.

CALYCOPEPLUS, Planch. in Bull. Soc. Bot. Fr. viii. (1861) 80. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 361.
ephedroides, *Planch. l. c.* 81.—Austral.
marginatus, *Benth. Fl. Australica*. vi. 53.—Austral.
paucifolius, Baill. Adansonia, vi. (1865–66) 319 = ephedroides.

CALYCOPHYLLUM, DC. Prod. iv. 367 (excl. sp. 2 & 3) (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 26.
EUKYLISTA, Benth. in Hook. Kew Journ. v. (1853) 230.
candidissimum, *DC. Prod.* iv. 367.—Am. austr.
coccineum, DC. l. c. = Warscewiczia coccinea.
grandiflorum, Meyen, Reise, ii. 324 = Mussaenda grandiflora.
multiflorum, *Griseb. in Goett. Abh.* xxiv. (1879) 165.—Reg. Argent.
Stanleyanum, *Schomb. in Hook. Lond. Journ. Bot.* iii. (1844) 692.—Guiana.
tubulosum, DC. Prod. iv. 367 (Pogonopt sp.).—Bras.
tubulosum, Seem. Bot. Voy. Herald, 135 = Pogonopus exsertus.

CALYCOPHYSUM, Karst. & Triana, in Linnaea, xxviii. (1856) 427. *CUCURBITACEAE*, Benth. & Hook. f. i. 828.
gracile, *Cogn. in Bull. Acad. Belg.* Sér. II. xlix. (1880) 191.—N. Granat.
pedunculatum, *Karst. & Triana, in Linnaea*, xxviii. (1856) 436.—N. Granat.

CALYCOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 130 = **Alloplectus**, Mart. (Gesner.).
concolor, Oerst. l. c. = A. dichrous.
dichrous, Oerst. l. c. = A. dichrous.
Pinelianus, Oerst. l. c. = A. Pinelianus.

CALYCOPTERIS, Lam. Illustr. t. 857 (1794†). *COMBRETACEAE*, Benth. & Hook. f. i. 686.
GETONIA, Roxb. Fl. Corom. i. 61. t. 87 (1795).
floribunda, *Lam. Tabl. Encyc.* ii. 468.—Ind. or.; Malaya.
nutans, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 59 = floribunda.

CALYCOPTERIS, Rich. ex DC. Prod. iii. 168 (1828) = **Calycoogonium**, DC. (Melastom.).

CALYCORECTES, Berg, in Linnaea, xxvii. (1854) 817. *MYRTACEAE*, Benth. & Hook. f. i. 720.
SCHIZOCALYX, Berg, l. c. 818 (1854).
cubensis, *Griseb. Cat. Pl. Cub.* 90.—Cuba.
grandifolius, *Berg, in Linnaea*, xxvii. (1854) 817.—Guiana.
Langsdorffi, *Berg, in Mart. Fl. Bras.* xiv. i. 597.—Bras.
latifolius, *Berg, in Linnaea*, xxx. (1859–60) 701.—Guiana.
Martianus, *Berg, in Mart. Fl. Bras.* xiv. i. 596.—Bras.
mexicanus, *Berg, in Linnaea*, xxvii. (1854) 818.—Mexic.
protensus, *Griseb. Cat. Pl. Cub.* 284.—Cuba.
Riedelianus, *Berg, in Mart. Fl. Bras.* xiv. i. 596.—Bras.
Schottianus, *Berg, l. c.* 857.—Bras.
Sellowianus, *Berg, l. c.* 856.—Bras.

CALYCOSERIS, A. Gray, Pl. Wright. ii. 104. t. 14 (1853). *COMPOSITAE*, Benth. & Hook. f. ii. 523.
Parryi, *A. Gray, in Torr. Bot. Mex. Bound.* 106.—Californ.
Wrightii, *A. Gray, Pl. Wright.* ii. 104. t. 14.—N. Mexic.

CALYCOSORUS, Endl. Gen. 499, in syn. (1838) = CALYCOCORSUS, F. W. Schmidt = Chondrilla, Linn. (Compos.).

CALYCOSIA, A. Gray, in Proc. Am. Acad. iv. (1860) 47. *RUBIACEAE*, Benth. & Hook. f. ii. 192.
Hunteri, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1884) 204.—Ins. Fiji.

CALYCOSIA:—
Milnei, *A. Gray, in Proc. Am. Acad.* iv. (1860) 207.—Ins. Fiji.
petiolata, *A. Gray, l. c.* 48.—Ins. Fiji.
pubiflora, *A. Gray, l. c.* 206.—Ins. Fiji.
sessilis, *A. Gray, l. c.* 48.—Ins. Samoa.

CALYCOSTEGIA, Lem. in Orb. Dict. iii. 86 (1849) = **Calystegia**, R. Br. (Convolv.).

CALYCOSTEMMA, Hanst. in Linnaea, xxix. (1857–58) 506 = **Isoloma**, Decne. (Gesnetrac.).
Lindenii, Hanst. l. c. 516.—N. Granat.

CALYCOTHRIX, Meisan. Gen. 107 (1837) = **Calythrix**, Labill. (Myrtac.).

CALYCOTOME, E. Mey. Comm. Pl. Afr. austr. 113 (1835) = **Diahilua**, DC. (Legumin.).
pusilla, E. Mey. l. c. 114 = D. gracilis.

CALYCOTOME, Link, in Schrad. Neues Journ. ii. (1808) 11. 50 (Calicotome). *LEGUMINOSAE*, Benth. & Hook. f. i. 481.
cretica, Presl, ex Nym. Consp. 168 = villosa.
infesta, *Guss. Fl. Sic. Syn.* ii. 247.—Dalmatia.
intermedia, Boiss. ex Reichb. f. Ic. Fl. Germ. xxii. 38. t. 2094 = villosa.
spinosa, *Link, Enum. Hort. Berol.* ii. 225.—Lusitan.
villosa, *Link, in Schrad. Neues Journ.* ii. (1808) ii. 51. —As. Min.

CALYCOTOMON, Hoffmg. Verz. Pfl. 166 (1824) = **Calicotome**, Link (Legumin.).

CALYCOTOMUS, Rich. ex DC. Prod. iii. 178 (1828) = **Conostegia**, D. Don (Melastom.).

CALYCOTROPIS, Turcz. in Bull. Soc. Nat. Mosc. xxxv. (1862) ii. 327. *PARONYCHIEAE* vel *ELATINEAE* ? Benth. & Hook. f. Gen. i. 978.
minuartioides, *Turcz. l. c.* 328.—Mexic.

CALYDERMOS, Lag. Gen. et Sp. Nov. 24 (1816) = **Calea**, Linn. (Compos.).
atriplicifolius, Spreng. Syst. iii. 457 = Isocarpha atriplicifolia.
jamaicensis, Spreng. Syst. iii. 457 = Equatorium Dalea.
longifolius, Lag. Gen. et Sp. Nov. 24 = Calea pedunculata.
pedunculatus, DC. Prod. v. 569 = Calea pedunculata.
repens, Spreng. Neue Entdeck. i. 278 = Otospermum involucratum.
rugosus, DC. Prod. v. 570 = Calea rugosa.
salmeafolius, DC. l. c. = Calea salmiaefolia.
scaber, Lag. Gen. et Sp. Nov. 24 = Calea peduncularis.
spilanthoides, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 514.—Cuba.

CALYDERMOS, Ruiz & Pav. Fl. Per. ii. 43 (1799) = **Nioandra**, Adans. (Solanac.).
erosus, Ruiz & Pav. l. c. = N. physaloides.

CALYDOREA, Herb. in Lindl. Bot. Reg. (1843) Misc. 85. *IRIDEAE*, Benth. & Hook. f. iii. 695.
BOTHREE, Steud. ex Klatt, in Linnaea, xxxi. (1860 61) 582.
ROTREEE, Klatt, in Mart. Fl. Bras. iii. 1. 543. t. 71 (1871).
azurea, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 287.—Hab. ?
campestris, *Baker, in Journ. Bot.* xiv. (1876) 188.—Am. trop.
cipuroides, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 287.—Hab. ?
furcata, *Baker, in Journ. Bot.* xix. (1876) 189.—Reg. Argent.
Gardneri, *Baker, l. c.*—Bras.
longispatha, Baker, l. c. = Sphenostigma longispatha.
lutcola, *Baker, l. c.*—Bras.
nuda, *Baker, l. c.*—Reg. Argent.
pallens, *Griseb. in Goett. Abh.* xxiv. (1879) 234.—Reg. Argent.
punctata, Baker, in Journ. Bot. xiv. (1876) 189 = Nemastylis punctata.
speciosa, *Herb. in Bot. Reg.* (1843) *Misc.* 85.—Chili.
texana, *Baker, in Journ. Bot.* xiv. (1876) 189.—Texas.

CALYGOGONIUM, G. Don, Gen. Syst. ii. 781 (1838) = **Calycogonium**, DC. (Melastom.).

CALYGONIUM, D. Dietr. Syn. ii. 1376 (1840) = prsec.

CALYLOPHIS, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 272 = **Oenothera**, Linn. (Onagrar.).
Berlandieri, Spach, l. c. 278 = Oe. serrulata ?
Drummondiana, Spach, l. c. = Oe. serrulata.
Drummondii, Spach, ex Steud. Nom. ed. II. i. 262 = prsec.
Nuttallii, Spach, Hist. Vég. Phan. iv. 350 = Oe. serrulata.

CALYLOPHUS, Spach, Hist. Vég. Phan. iv. 351 = prsec.

CALYMENIA, Pers. Syn. i. 86 (1805) = **Oxybaphus**, Vahl (Nyctagin.).
aggregata, Pers. Syn. i. 87 = O. aggregatus.
albida, Nutt. Gen. Am. i. 96 = O. aggregatus.
angustifolia, Nutt. in Fras. Cat. (1813); Gen. Am. i. 96 = O. angustifolius.
Cervantesii, Heynh. Nom. ii. 110 = O. Cervantesii.
chilensis, Heynh. l. c. = O. chilensis.
corymbosa, Pers. Syn. i. 87 = O. glabrifolius.
decumbens, Nutt. Gen. Am. i. 96 = O. decumbens.
expansa, Pers. Syn. i. 86 = O. expansus.
hirsuta, Nutt. Gen. Am. i. 96 = O. hirsutus.
linearis, Nutt. l. c. = O. angustifolius.
nyctaginea, Nutt. l. c. = O. nyctagineus.
ovata, Pers. Syn. i. 86 = O. ovatus.
paniculata, Heynh. Nom. ii. 110 = O. glabrifolius.
pilosa, Nutt. Gen. Am. i. 96 = O. ovatus.
prostrata, Pers. Syn. i. 86 = O. prostratus.
viscosa, Pers. l. c. = O. viscosus.

CALYMMANDRA, Torr. & Gray, Fl. N. Am. ii. 262 (1841) = **Evax**, Gaertn. (Compos.).
candida, Torr. & Gray, l. c. = E. candida.

CALYMNANDRA, Lindl. Veg. Kingd. 710 (1847) = prsec.

CALYNUX, Rafin. in Journ. Phys. lxxxix. 257 (1819) = **Fyrularia**, Rich. (Santalac.).

CALYPLECTUS, Ruiz & Pav. Prod. 73. t. 13 (1794) = **Lafoensia**, Vand. (Lythrar.).
acuminatus, Ruiz & Pav. Fl. Per. iv. t. 423 = L. acuminata.
adenophyllus, Schott, ex Steud. Nom. ed. II. i. 263 = L. replicata.
dependens, Ruiz, ex Koehne, in Engl. Bot. Jahrb. iii. (1882) 155 = L. acuminata.
punicaefolius, Bert. ex Koehne, l. c. 154 = L. punicaefolia.
speciosus, H. B. & K. Nov. Gen. et Sp. vi. 183. t. 548 = L. speciosa.

CALYPSO, Salisb. Parad. Lond. t. 89 (1806).
ORCHIDEAE, Benth. & Hook. f. iii. 496.
CYTHEREA, Salisb. in Trans. Hort. Soc. i. (1812) 301.
NORNA, Wahlenb. Fl. Suec. 561 (1826).
ORCHIDIUM, Sw. in Svensk Bot. t. 518 (1819).
americana, R. Br. in Ait. Hort. Kew. ed. II. v. 208 = borealis.
borealis, *Salisb. Parad. Lond.* t. 89.—Reg. bor. temp. *bulbosa*, Reichb. f. lc. Fl. Germ. xiii. 158. t. 187 = borealis ?

CALYPSO, Thou. Hist. Vég. Afr. Austr. i. 33. t. 6 (1806) = **Salacia**, Linn. (Celastr.).
africana, G. Don, Gen. *Syst.* i. 629 = S. africana.
campestris, Cambess. in St. Hil. Fl. Bras. Mer. ii. 111. t. 104 = S. campestris.
debilis, G. Don, Gen. Syst. i. 629 = S. debilis.
erecta, G. Don, l. c. = S. erecta.
madagascariensis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 111. t. 104 = S. madagascariensis.
pyriformis, G. Don, Gen. Syst. i. 629 = S. pyriformis.
salaciaides, Thou. ex Cambess. in St. Hil. Fl. Bras. Mer. ii. 111. t. 104 = S. Calypso.
senegalensis, Cambess. l. c. = S. senegalensis.
sylvestris, Mart. in Flora, xx. (1837) ii. Beibl. 96 = S. sylvestris.

CALYPSODIUM, Link, Handb. i. 262 (1829) = **Calypso**, Salisb. (Orchid.).
boreale, Link, l. c. = Calypso borealis.

CALYPSOGYNE, Neraud, in Freyc. Voy. Bot. 35, 96 (1826), nomen.

CALYPTERIOPETALON, Hassk. in Flora, xl. (1857) 581 = Croton, Linn. (Euphorbiac.).
brastianum, Hassk. l. c. = Croton urticaefolius.

CALYPTERIS, Zipp. ex Mackl. in Alg. Konst. en Letter-boek, i. (1828) 398 (Ord. Nat. ?).
minata, *Zipp. ex Mackl. l. c.* nomen.—Hab. ?

CALYPTHRANTES, Raeusch. Nom. ed. III. 144 (1797) = **Calyptranthes**, Sw. (Myrtac.).

CALYPTOCARPUS, Less. Syn. Comp. 221 (1832) = **Blainvillea**, Cass. (Compos.).
bahiensis, Sch. Bip. in Bot. Zeit. xxiv. (1866) 165 = B. bahiensis.
Burchellii, Sch. Bip. l. c. = B. rhomboides.
megapotamica, Sch. Bip. l. c. = B. biaristata.
vialis, Less. Syn. Comp. 221 = B. tampicana.
Wendlandii, Sch. Bip. in Bot. Zeit. xxiv. (1866) 165. —Costa Rica.

CALYPTRANTHES, Sw. Prod. Veg. Ind. Occ. 79 (1788). *MYRTACEAE*, Benth. & Hook. f. i. 717.
CRYPTERIA, Adans. Fam. ii. 80 (1763).
MITRANTHES, Berg, in Linnaea, xxvii. (1854) 316.
SUZYGIUM, P. Br. ex Adans. Fam. ii. 244 (1763).
affinis, *Berg, in Mart. Fl. Bras.* xiv. i. 43.—Bras.
ancepa, *Berg, l. c.* 40.—Bras.
aromatica, Blume, Bijdr. 1092 (= *Caryophyllus antisepticus*).
aromatica, *St. Hil. Pl. Us. Bras.* fasc. iii. t. 14.—Bras.
axillaris, *Berg, in Mart. Fl. Bras.* xiv. i. 53.—Bras.
binarginata, *Berg, l. c.* 48.—Bras.
bipennis, *Berg, in Linnaea*, xxxi. (1861–62) 248.—Peruv.
Blanchetiana, *Berg, in Mart. Fl. Bras.* xiv. i. 40.—Bras.
brasiliensis, *Spreng. Syst.* ii. 499.—Bras.
bullata, *DC. Prod.* iii. 258.—Honduras.
calyptrata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 181.—Cuba.
capitellata, Buch.-Ham. ex Wall. Cat. n. 3560 B = Eugenia Jambolana.
caryophyllata, Pers. Syn. 92 = Eugenia caryophyllaea.
caryophyllifolia, Willd. in Usteri, Ann. xvii. (1796) 22 = Eugenia Jambolana.
candata, *Garden. in Hook. Lond. Journ. Bot.* iv. (1845) 102.—Bras.; N. Granat.
cephalantha, *Berg, in Mart. Fl. Bras.* xiv. i. 46.—Bras.
Chytracuila, *Sw. Prod. Veg. Ind. Occ.* 79.—Jamaica.
cinnaleafolia, *Berg, in Mart. Fl. Bras.* xiv. i. 39.—Bras.
concinna, *DC. Prod.* iii. 258.—Bras.
cordata, *Berg, in Mart. Fl. Bras.* xiv. i. 48.—Bras.
cordifolia, Moon, Cat. Fl. Ceyl. 39 = Eugenia Neesiana sed cordifolia.
corymbosa, Blume, Bijdr. 1091 (= *Syzygium corymbosum*).
costaricensis, *Berg, in Linnaea*, xxvii. (1854) 90.—Costa Rica.
costata, Buch.-Ham. in Wall. Cat. n. 3556 = Eugenia operculata.
cubensis, *Berg, in Linnaea*, xxix. (1857–58) 215.—Ind. occ.
Cumini, Pers. Syn. ii. 92 = Eugenia Jambolana.
cuminoides, Stokes, Bot. Mat. Med. iii. 67 = Eugenia caryophyllifolia.
cuneata, Buch.-Ham. ex Wall. Cat. n. 3557 = Eugenia operculata.
cuprea, *Berg, in Linnaea*, xxx. (1859–60) 653.—Cuba.
cupulata, *Mart. ex DC. Prod.* iii. 256.—Bras.
Dancea, Buch.-Ham. ex Wall. Cat. n. 3560 = Eugenia Heyneana.
decandra, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 181.—Cuba.
densa, *DC. Prod.* iii. 257.—Bras.
densiflora, *Poepp. ex Berg, in Linnaea*, xxvii. (1854) 90.—Peruv.

CALYPTRANTHES:—
dichotoma, *Casar. Decad. Nov. Stirp. Bras.* v. 47.—Bras.
ennesatha, *Wright, in Sauv. Fl. Cub.* 40.—Cuba.
eriopoda, *DC. Prod.* iii. 257.—Bras.
eugenioides, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 370.—Bras.
eugenioides, *Seem. Fl. Vit.* 81.—Ins. Fiji.
fasciculata, *Berg, in Linnaea*, xxvii. (1854) 81.—Guiana.
fastigiata, *Berg, in Mart. Fl. Bras.* xiv. i. 50.—Bras.
fastigiata, Blume, Bijdr. 1090 (= *Caryophyllus fastigiatus*).
floribunda, Blume, l. c. 1091 = Eugenia floribunda.
Forsteri, *Berg, in Linnaea*, xxvii. (1854) 38.—Ins. S. Lucia.
fragrans, Ruiz, ex Berg, l. c. 20 = paniculata.
glomerata, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 371 —Bras.
gracilipes, *Wright, in Sauv. Fl. Cub.* 39.—Cuba.
grandiflora, *Berg, in Mart. Fl. Bras.* xiv. i. 542.—Bras.
grandifolia, *Berg, l. c.* 48.—Bras.
grandis, Buch.-Ham. ex Wall. Cat. n. 3554 = Eugenia operculata.
guineensis, *Willd. Sp. Pl.* ii. 974.—Afr. trop.
Jambolana, Moon, Cat. Fl. Ceyl. 39 = Eugenia sylvestria.
Jambolana, Willd. in Usteri, Ann. xvii. (1796) 38 = Eugenia Jambolana.
jambolifera, Stokes, Bot. Mat. Med. iii. 68 = Eugenia Jambolana.
Karwinskyana, *Berg, in Linnaea*, xxix. (1857–58) 214 —Mexic.
lanceolata, *Berg, in Mart. Fl. Bras* xiv. i. 51.—Bras.
Langsdorffi, *Berg, l. c.* 539.—Bras.
laterifiora, DC. Prod. iii. 258 (= *Rubachia laterifiora*) —Bras.
latifolia, *Poir. Encyc. Suppl.* ii. 42.—Ins. Maurit.
laxiflora, *Blume, Bijdr.* 1090.—Java.
Lindeniana, *Berg, in Linnaea*, xxix. (1857–58) 213.—Mexic.
longifolia, *Berg, in Mart. Fl. Bras.* xiv. i. 46.—Bras.
lorentziflora, *DC. Prod.* iii. 258.—Bras.
lucida, *Mart. ex DC. l. c.*—Bras.
macrophylla, *Berg, in Mart. Fl. Bras.* xiv. i. 45.—Bras.
Makui, Raeusch. Nom. ed. III. 144 (= *Syzygium Makui*).
Makui, Blanco, Fl. Filip. ed. I. 419 = Eugenia operculata.
marginifera, Hance, ex Walp. Ann. ii. 629 = Eugenia operculata.
Martiniana, *DC. Prod.* iii. 257.—Bras.
maschalantha, *Berg, in Mart. Fl. Bras.* xiv. i. 53.—Bras.
melanocoida, *Berg, l. c.* 50.—Bras.
micrantha, *Wright, ex Griseb. Cat. Pl. Cub.* 85.—Cuba.
multiflora, *Poepp. ex Berg, in Mart. Fl. Bras.* xiv. i. 42.—Bras.
muscifora, *Berg, l. c.* 54.—Bras.
mutabilis, *Berg, l. c.* 540.—Bras.
myrtoides, *Seem. Fl. Vit.* 81.—Ins. Fiji.
nigricans, *DC. Prod.* iii. 258.—Bras.
nitida, Raeusch. Nom. ed. III. 144 = *Syzygium lucidum*.
Nummularia, *Berg, in Linnaea*, xxx. (1859–60) 653. —Ins. S. Domingo.
obscura, *DC. Prod.* iii. 257.—Bras.
obscura, Mart. in Flora, xx. (1837) Beibl. ii. 87 = tuberculata.
obtusa, *Benth. in Hook. Journ. Bot.* ii. (1840) 319.—Guiana.
obtusa, Miq. in Linnaea, xxii. (1849) 173 = Marlieria suffruticosa.
obtusifolia, Buch.-Ham. ex Wall. Cat. n. 3560 B = Eugenia inophylla.
obovata, *Berg, in Mart. Fl. Bras.* xiv. i. 51.—Bras.
obtusa, Wight, Illustr. ii. 15 = Eugenia myrtifolia.
Ottonis, *Wright, in Sauv. Fl. Cub.* 40.—Cuba.
ovalifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 371. —Bras.
ovata, *Berg, in Mart. Fl. Bras.* xiv. i. 56.—Bras.
pallens, *Griseb. in Gentt. Am.* vii. (1857) 215.—Ind. occ.

CALYPTRANTHES:—
paniculata, Raeusch. Nom. ed. III. 144 = Eugenia paniculatum.
paniculata, *Ruiz & Pav. Fl. Per.* iv. t. 424; *Syst.* 181.—Peruv.
paniculata, Spreng. Neue Entdeck. ii. 170 = brasiliensis.
pauciflora, *Berg*, in *Mart. Fl. Bras.* xiv. i. 543.—Bras.
pendula, Berg, in Linnaea, xxvii. (1854) 21 = C. Chytraculia.
platyphylla, *Berg*, in *Mart. Fl. Bras.* xiv. i. 539.—Bras.
Poeppigiana, *Berg*, l. c. 45.—Bras.
Pohliana, *Berg*, l. c. 42.—Bras.
pollicina, Willem. in Usteri, Ann. xvii. (1796) 39 = Eugenia obovata.
polyantha, *Berg*, in *Mart. Fl. Bras.* xiv. i. 541.—Bras.
pteropoda, *Berg*, l. c. 47.—Bras.
pulchella, *DC. Prod.* iii. 257.—Bras.
punctata, *Griseb.* in *Mem. Am. Acad.* N. S. viii. (1861) 181.—Cuba.
pyrifolia, Blume, Bijdr. 1090 = Eugenia pyrifolia.
racemosa, Blume, l. c. 1089 (= *Syzygium racemosum*).
ramiflora, Blanco, Fl. Filip. ed. I. 420 = Eugenia bracteata.
Regeliana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 542.—Bras.
rigida, Mnch. ex Griseb. Fl. Brit. W. Ind. 238 = C. Syzygium.
rigida, *Sw. Prod. Veg. Ind. Occ.* 80.—Jamaic.; Cuba.
rigida, Tussac, Fl. Antill. iii. t. 26 = Tussaccana.
rostrata, Blume, Bijdr. 1092 (= *Syzygium rostratum*).
rostrata, *Griseb.* in *Mem. Am. Acad.* N. S. viii. (1861) 181.—Cuba.
rotundata, *Griseb. l. c.*—Cuba.
rufa, *Berg*, in *Mart. Fl. Bras.* xiv. i. 541.—Bras.
Ruiziana, *Berg*, in *Linnaea*, xxvii. (1854) 23.—Peruv.
Schiediana, *Berg*, l. c. 28.—Mexic.
Schlechtendaliana, *Berg*, l. c. 39.—Mexic.
scoparis, *Berg*, l. c. xxxi. (1861–62) 247.—Bras.
Seemanni, *Seem. Fl. Vit.* 81.—Ins. Fiji.
sericea, *Griseb. Fl. Brit. W. Ind.* 238.—Ind. occ.
speciosa, *Sagot*, in *Ann. Sc. Nat.* Sér. VI. xx. (1885) 187.—Guiana.
Spruceana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 45.—Bras.
strigipes, *Berg*, l. c. 540.—Bras.
Syzygium, *Sw. Fl. Ind. Occ.* ii. 919.—Ind. occ.
Taina, Buch -Ham. ex Wall. Cat. n. 5555 = Eugenia operculata.
tenuis, Buch.-Ham. ex Wall. l. c. sub n. 3570 = Eugenia tenuis.
tetrapteris, *Berg*, in *Mart. Fl. Bras.* xiv. i. 58.—Bras.
Thomasiana, *Berg*, in *Linnaea*, xxvii. (1854) 26.—Ind. occ.
tuberculata, *Berg*, in *Mart. Fl. Bras.* xiv. i. 52.—Bras.
Tussaccana, *Berg*, in *Linnaea*, xxvii. (1854) 25.—Ind. occ.
uniflora, *Spreng. in Flora*, xx. (1837) *Beibl.* ii. 87.—Bras.
variabilis, *Berg*, in *Mart. Fl. Bras.* xiv. i. 49.—Bras.
Widgreniana, *Berg*, l. c. 39.—Bras.
Wilsonii, *Griseb. Fl. Brit. W. Ind.* 238.—Ind. occ.
Zuygium, Blanco, Fl. Filip. ed. II. 393 = Eugenia operculata.
Zuygium, Sw. Prod. Veg. Ind. Occ. 79 = C. Syzygium.

CALYPTRANTHUS, Thou. Obs. Pl. Isles Afr. 26 (1811) = **Capparis**, Tourn.

CALYPTRANTHUS, Blume, Bijdr. 1089 (1826) = **Eugenia**, Mich. (Myrtac.).

CALYPTRANTHUS, Juss. in Dict. Sc. Nat. vi. 274 (1817) = **Calyptranthes**, Sw. (Myrtac.).

CALYPTRARIA, Nand. in Ann. Sc. Nat. Sér. III. xvii. (1852) 182 = **Centronia**, D. Don (Melastom.).
africana, DC. Prod. ix. 469, in syn. = Varronia abyssinica.
brachycera, Nand. in Ann. Sc. Nat. Sér. III. xvii. (1852) 184 = Cent. brachycera.
eximia, Nand. l. c. 183 = Cent. eximia.
hexmeantha, Planch. & Linden, in Fl. des Serres, Sér. I. ix. (1853–54) t. 924 = Cent. hexmeantha.
insignis, Nand. in Ann. Sc. Nat. Sér. III. xvii. (1852) 183 = Cent. insignis.
Mutisii, Nand. l. c. 182 = Cent. Mutisii.

CALYPTRELLA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 115. *MELASTOMACEAE*, Benth. & Hook. f. i. 750.
cucullata, *Triana*, in *Trans. Linn. Soc.* xxviii. (1871) 72.—Peruv.
Galeotti, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 115.—Mexic.
gracilis, *Triana*, in *Trans. Linn. Soc.* xxviii. (1871) 72.—Peruv.
tristis, *Triana*, l. c.—Peruv.

CALYPTRIDIUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 198 (1838). *PORTULACEAE*, Benth. & Hook. f. i. 159.
monandrum, *Nutt. l. c.*—Calif.
quadripetalum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 356.—Calif.
roseum, *S. Wats. in Bot. King, Exp.* 44. t. 6.—Calif.

CALYPTRION, Ging. in DC. Prod. i. 288 (1824) = **Corynocarpus**, Mart. (Violac.).
Aubletii, Ging. l. c. 289 = Cor. Hybanthus.
Berterii, Ging. l. c. = Cor. Berterii.
citrifolium, Ging. l. c. = Cor. Loeflingii.
diandrum, Ging. l. c. = Cor. diandra.
frangulaefolium, Ging. l. c.—Peruv.
nitidum, Benth. in Hook. Journ. Bot. iv. (1842) 106 = Cor. Hybanthus.
orinocense, H. B. & K. Nov. Gen. et Sp. v. 334.—Venezuela.
pyrifolium, Sweet, ex Steud. Nom. ed. II. i. 263 = Anchietea pyrifolia.

CALYPTRIOPETALUM, Hassk. ex Muell. Arg. in DC. Prod. xv. ii. 518 (1866) = **Croton**, Linn. (Euphorb.).

CALYPTROCALYX, Blume, Rumphia, ii. 103. t. 102 D, 113, 161 (1836). *PALMAE*, Benth. & Hook. f. Gen. iii. 902.
LACCOSPADIX, H. Wendl. & Drude, in Linnaea, xxxvii. (1875) 203.
australiacus, *Scheff. in Ann. Jard. Bot. Buitenz.* i. (1876) 132, 140.—Austral.
spicatus, *Blume, Rumphia*, ii. 108.—Amboina.

CALYPTROCARPUS, Reichb. Nom. 82 (1841) = CALYPTOCARPUS, Less. = **Blainvillea**, Cass. (Compos.).

CALYPTROCARYA, Nees, in Linnaea, ix. (1834) 304. *CYPERACEAE*, Benth. & Hook. f. Gen. iii. 1069.
angustifolia, *Nees, l. c.*—Guiana.
bicolor, Nees, l. c. = Poeppigiana.
brevicaulis, *Nees, l. c.* 305.—Guiana.
fragilens, *Nees, l. c.* 304.—Am. trop.
longifolia, Boeck. in Linnaea, xxxviii. (1874) 431 = C. Palmetto.
longifolia, *Kunth, Enum. Pl.* ii. 365.—Guiana.
Martii, *Nees, in Mart. Fl. Bras.* ii. i. 195.—Bras.
Palmetto, *Nees, l. c.*—Am. trop.
Poeppigiana, *Kunth, Enum. Pl.* ii. 364.—N. Granat.
Schottmelleriana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 509.—Bras.

CALYPTROCORYNE, Schott, in Oestr. Bot. Wochenschr. (1857) 262 = **Theriophonum**, Blume (Aroid.).
cochlichinensis, Schott, Prod. Aroid. 105 = T. Wightii.
minuta, Schott, l. c. = T. Wightii.
Wightii, Schott, in Oestr. Bot. Wochenschr. (1857) 262 = T. Wightii.

CALYPTROGYNE, H. Wendl. in Bot. Zeit. xvii. (1859) 72. *PALMAE*, Benth. & Hook. f. iii. 914. CALYPTRONOMA, Griseb. Fl. Brit. W. Ind. 518 (1864).
Ghiesbrechtiana, *H. Wendl. l. c.*—Mexic.
Ghiesbreghtii, H. Wendl. in Kerch. Palm. 238 = Ghiesbrechtiana.
glanca, *H. Wendl. l. c.*—Nicaragua.
sarapiquensis, *H. Wendl. l. c.*—Costa Rica.
spicigera, *H. Wendl. in Bot. Zeit.* xvii. (1859) 72.—Guatemala.

CALYPTROLEPIS, Steud. Syn. Pl. Cyp. 151 (1855) = **Rynchospora**, Vahl (Cyperac.).
junciformis, Steud. l. c. = R. oligantha.

CALYPTROMYRCIA, Berg, in Linnaea, xxvii. (1854) 34 = **Myrcia**, DC. (Myrtac.).
cordata, Berg, in Mart. Fl. Bras. xiv. i. 56 = M. cordata.
costata, Berg, l. c. = M. costata.
cymosa, Berg, l. c. 58.—Bras.
elegans, Berg, l. c. = M. elegans.
eugenioides, Berg, l. c. 56 = M. eugenioides.
paniculata, Berg, l. c. 57.—Bras.
puberula, Berg, l. c. = M. puberula.
Spiriana, Berg, l. c. 59 = M. Spixiana.
venosa, Berg, l. c. 57.—Bras.

CALYPTRONOMA, Griseb. Fl. Brit. W. Ind. 518 (1864) = **Calyptrogyne**, H. Wendl. (Palm.).
dulcis, H. Wendl. in Kerch. Palm. 238 (= *Geonoma dulcis*).
intermedia, H. Wendl. l. c. nomen.—Hab.?
robusta, Trail, in Journ Bot. xiv. (1876) 330.—Bras.
Schwartzii, H. Wendl. in Kerch. Palm. 238 = seq.
Swartzii, Griseb. Fl. Brit. W. Ind. 518 (= *Geonoma Swartzii*).

CALYPTROON, Miq. Fl. Ind. Bat. Suppl. 471 (1860) = **Baccaurea**, Lour. (Euphorbiac.).
sumatranum, Miq. l. c. 472 = B. sumatrana.

CALYPTROPSIDIUM, Berg, in Linnaea, xxvii. (1854) 849 = **Psidium**, Linn. (Myrtac.).
Friedrichsthalianum, Berg, l. c. 850.—Guatemala.

CALYPTROSPATHA, Klotzsch, ex Baill. Etud. Gén. Euph. 440 (1858); et in Peters, Reise Mossamb. Bot. t. 18 (1861) = **Acalypha**, Linn. (Euphorb.).
pubiflora, Klotzsch, l. c. 97. t. 18 = A. pubiflora.

CALYPTROSPERMUM, A. Dietr. Sp. Pl. i. 926 (1831) = **Monodora**, Humb. & Bonpl. (Oleac.).
integrifolium, A. Dietr. l. c. 227 = M. integrifolia.
trifidum, A. Dietr. l. c. = M. trifida.

CALYPTROSTEGIA, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 72 pro parte = **Pimelea**, Banks (Thymel.).
angustifolia, C. A. Mey. l. c. 74 = P. angustifolia.
argentea, C. A. Mey. l. c. = P. argentea.
brevifolia, C. A. Mey. l. c. = P. brevifolia.
cernua, C. A. Mey. l. c. = P. spathulata.
clavigera, Walp. Ann. iii. 324 = P. scrpyllifolia.
Cornucopiae, Endl. Gen. Suppl. iv. ii. 60 = P. Cornucopiae.
curvifloca, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. curviflora.
Drummondii, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 178 = P. suaveolens.
flava, Endl. Gen. Suppl. iv. ii. 60 = P. flava.
glauca, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. glauca.
gracilifloca, Endl. Gen. Suppl. iv. ii. 61 = P. sylvestris.
gracilis, Endl. l. c. = P. curviflora.
hypericina, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. ligustrina.
intermedia, C. A. Mey. l. c. = P. glauca.
lanata, Endl. Gen. Suppl. iv. ii. 61 = P. hispida.
latifolia, Endl. l. c. = P. latifolia.
Lehmanniana, Endl. l. c. = P. Lehmanniana.
ligustrina, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. ligustrina.
linifolia, C. A. Mey. l. c. = P. linifolia.
linoides, Endl. Gen. Suppl. iv. ii. 61 = P. linifolia.
longiflora, Endl. l. c. = P. longiflora.
macrocephala, Walp. Ann. iii. 326 = P. suaveolens.
Menkenna, Endl. Gen. Suppl. iv. ii. 61 = P. suaveolens.
microcephala, Endl. l. c. = P. microcephala.
myrianthe, Endl. l. c. = P. argentea.
nana, Endl. l. c. = P. imbricata.
nervosa, Walp. Ann. iii. 324 = P. nervosa.
phylicoides, Walp. l. c. = P. phylicoides.
punicea, Endl. Gen. Suppl. iv. ii. 60 = P. punicea.
Schlechtendaliana, Walp. Ann. iii. 324 = P. glauca.
spathulata, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. spathulata.
spicata, Endl. Gen. Suppl. iv. ii. 61 = P. spicata.

CALYPTROSTEGIA :—
suaveolens, Endl. Gen. Suppl. iv. II. 61 = Pimelea suaveolens.
sulphurea, Walp. Ann. iii. 325 = P. sulphurea.
sylvestris, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. sylvestris.
tinctoria, Endl. Gen. Suppl. iv. II. 61 = P. suaveolens.
trichostachya, Walp. Ann. iii. 325 = P. trichostachya.
villifera, Walp. l. c. 324 = P. villifera.
villosa, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 178 = P. longiflora.

CALYPTROSTIGMA, Klotzsch, in Lehm. Pl. Preiss. i. 175 (1844) = **Bayeria**, Miq. (Euphorbiac.).
ledifolium, Klotzsch, l. c. 176 = B. Drummondii.
oblongifolium, Klotzsch, l. c. = B. viscosa.
viscosum, Klotzsch, l. c. = B. viscosa.

CALYPTROSTIGMA, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 46) (1847) = **Diervilla**, Tourn. (Caprifol.).
Middendorffianum, Trautv. & Mey. l. c. = D. Middendorffiana.

CALYPTROSTYLIS, Nees, in Linnæa, ix. (1834) 295 = **Rynchospora**, Vahl (Cyperac.).
articulata, Nees, in Hook. Journ. Bot. ii. (1840) 394, in obs. = R. aurea.
asperula, Nees, in Mart. Fl. Bras. ii. i. 139 = R. aurea.
capitata, Nees, l. c. 140 = R. minarum.
divergens, Nees, l. c. 139 = R. aurea.
fasciculoris, Nees, l. c. 139 = R. aurea.
florida, Nees, l. c. 138 = R. aurea.
Gaudichaudii, Nees, in Linnæa, ix. (1834) 295 = R. aurea.
gigantea, Nees, in Mart. Fl. Bras. ii. i. 137 = R. gigantea.
longirostra, Nees, in Hook. Journ. Bot. ii. (1840) 394 = R. amazonica.
paniculata, Liebm. in Videnak. Selsk. Skr. V. ii. (1851) 252 = R. Schiedeana.
Rudgei, Nees, in Linnæa, ix. (1834) 295 = R. gigantea.
Schiedeana, Liebm. in Videnak. Selsk. Skr. V. ii. (1851) 252 = R. Schiedeana.
terminalis, Nees, ex Steud. Syn. Pl. Cyp. 143 = R. minarum.
asylanica, Nees, in Hook. Journ. Bot. ii. (1840) 394, in obs. = R. trifida.

CALYSACCION, Wight, Illustr. i. 130 (1840) = **Oohroacarpus**, Thou. (Guttifer.).
chinense, Walp. in Bot. Zeit. ix. (1851) 868 = O. longifolius.
Horsfii, Teijsm. & Binn. Cat. Hort. Bog. 205.—Borneo.
longifolium, Wight, Illustr. i. 130 = O. longifolius.
obovale, Miq. Fl. Ind. Bat. Suppl. i. 500 = O. ovalifolius.
ovalifolium, Choisy, Guttif. Ind. 45 = O. ovalifolia.
siamense, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 7, 209 = O. siamensis.
tinctorium, Seem. Fl. Vit. 13.—Ins. Fiji.

CALYSERICOS, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 578 (1856–57) = **Cryptadenia**, Meisn. (Thymelæac.).
argentea, Eckl. & Zeyl. l. c. 582 = Gnidia Cephalotes.
canaliculata, Eckl. & Zeyh. l. c. 578 = Cr. grandiflora.
grandiflora, Eckl. & Zeyh. l. c. = Cr. grandiflora.
linarifolia, Eckl. & Zeyh. l. c. 583 = Gnidia linoides.
parviflora, Eckl. & Zeyh. l. c. 574 = Cr. brevifiora.
subulata, Eckl. & Zeyh. l. c. = Cr. brevifiora.
tubularis, Eckl. & Zeyh. l. c. = Cr. brevifiora.
typica, Eckl. & Zeyl. l. c. 581 = Gnidia pubescens.
uniflora, Eckl. & Zeyh. l. c. 578 = Gnidia uniflora.

CALYSPHYRUM, Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) 107 ; Enum. Pl. Chin. Bor. 33 = **Diervilla**, Tourn. (Caprifol.).
floridum, Bunge, l. c. = D. florida.
pauciflorum, Bunge, in Walp. Rep. ii. 450 = D. florida.
roseum, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. xiii. (1855) 220 = D. rosea.

CALYSTEGIA, R. Br. Prod. 483 (1810). *CONVOLVULACEÆ*, Benth. & Hook. f. ii. 874.
Melmania, Raſn. Fl. Tellur. iv. 71 (1838).
Volvulus, Medic. in Theatrw. Vorles. Churpf. Phys. Oek. Ges. i. (1791) 202.

CALYSTEGIA :—
acetosifolia, Turcz. Fl. Baic. Dahur. ii. II. 289 = hederacea.
affinis, *Endl. Prod. Fl. Norf.* 51.—Ins. Norfolk.
asserifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 844 = C. Soldanella.
barbara, *Pomel, Nouv. Mat. Fl. Atl.* 88.—Afr. bor.
Basiniana, Turcz. ex Ledeb. Fl. Ross. iii. 93 = pellita.
Catesbeiana, *Pursh, Fl. Am. Sept. Supplt.* ii. 729.—Am. bor.
chinensis, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 284.—China.
dahurica, Choisy, in DC. Prod. ix. 433 = pellita.
dahurica, *Pusch. ex Planch. in Fl. des Serres*, x. (1854–55) 345, t. 1075.—Sibir.
grandiflora, *G. Don, Gen. Syst.* iv. 297.—Ind. or.
Hantelmanni, *Phil. in Linnæa*, xxx. (1859–60) 196.—Chili.
hederacea, *Wall. Cat.* n. 1838.—Ind. or. ; China ; Afghan.
inflata, G. Don, in Sweet, Hort. Brit. ed. III. 486 = sepium.
japonica, *Choisy, in Zoll. Syst. Vers. Ind. Archip.* 130, 182.—Japon.
Kerianæ, Sweet, Hort. Brit. ed. I. 287 = Hewittia bicolor.
Krauseana, *Phil. in Linnæa*, xxxiii. (1864–65) 154.—Chili.
lucens, G. Don, Gen. Syst. iv. 296 = sylvestris.
marginata, *R. Br. Prod.* 484.—Australas.
Maximiliana, *Nees, in Nesswied, Reise Nord. Am.* ii. 443.—Am. austr.!
mucronata, Spreng. ex Choisy, in DC. Prod. ix. 430 = Ipomoea martinicensis.
multiflora, Raſn. Fl. Ludov. 46 = sepium !
obtusa, *Pomel, Nouv. Mat. Fl. Atl.* 295.—Afr. bor.
ochroleuca, *Boj. Hort. Maurit.* 231.—Afr. trop.
palmato-partita, Benth. & Hook. f. Gen. ii. 874 = sep.
palmatipartita, Meisn. in Mart. Fl. Bras. vii. 317 = Ipomoea palmato-pinnata.
paradoxa, Pursh, Fl. Am. Sept. Suppl. ii. 729 = Convolvulus hirsutus.
pellita, *G. Don, Gen. Syst.* iv. 296.—Sibir. baical.
physoides, *Pomel, Nouv. Mat. Fl. Atl.* 294.—Afr. bor.
pubescens, Lindl. in Journ. Hort. Soc. i. (1846) 70 = hederacea.
reniformis, R. Br. Prod. 483 = C. Soldanella.
riparia, *Raſn. New Fl. Am.* ii. 29.—Am. bor.
rosea, *Wall. in Linnæa*, xxix. (1857–58) 15.—Chili.
rugosa, *Ledeb. Ind. Hort. Dorp.* (1821) 4.—Hab. !
sagittata, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 256.—Am. bor.
Scammonia, Pritz. Ind. Ic. ii. 58 = Convolvulus Scammonia.
sepium, *R. Br. Prod.* 483.—Cosmop. temp.
sethifera, Meisn. in Mart. Fl. Bras. vii. 318 = Ipomoea setifera.
silvatica, *Choisy, in DC. Prod.* ix. 433.—Europ. austr. or ; Reg. Caucas. ; Afr. bor.
Soldanella, *R. Br. Prod.* 483.—Cosmop. temp.
spithamæa, *Pursh, Fl. Am. Sept.* i. 143.—Am. bor.
stans, Steud. Nom. ed. II. i. 364 = spithamæa ?
subacualis, *Hook. & Arn. Bot. Beech. Voy.* 368.—Calif.
subvolubilis, G. Don, Gen. Syst. iv. 296 = pellita.
sylvestris, *Roem. & Schult. Syst.* iv. 188.—Oriens.
tomentosa, Pursh, Fl. Am. Sept. i. 143 = spithamæa.
tuguriorum, *R. Br. ex Hook. Fl. Nov. Zel.* i. 328. t. 87.—N. Zeland. ; Chili.
valdiviana, *Miq. in Lechl. Berl. Ahm. Austr.* 54.—Chili.
villosa, *Kellogg, in Proc. Calif. Acad.* v. (1873) 17.—Calif.
villosa, *Raſn. Fl. Ludov.* 45.—Am. bor.

CALYTHRIX, Labill. Nov. Holl. Pl. ii. 8. t. 146 (1806) (Calytrix). *MYRTACEÆ*, Benth. & Hook. f. i. 699.
Calycothrix, Meisn. Gen. 107 (1837).
Trichocalyx, Schau. Myrt. Xeroc. 86, adnot. (1840).
achaeta, *F. Muell. in Trans. Phil. Viet. Inst.* iii. (1859) 48.—Australas.
angulata, *Lindl. Swan River App.* 5.—Australas.
arborescens, *F. Muell. in Trans. Phil. Viet. Inst.* iii. (1859) 48.—Austral.
asperula, *Schau. in Lehm. Pl. Preiss.* i. 104.—Australas.
aurea, *Lindl. Swan River App.* 5. t. 2.—Australas.
Banksii, Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 361 = tetragona.

CALYTHRIX :—
Behriana, Schlecht. in Linnæa, xx. (1847) 650 = tetragona.
Billardieri, Schau. in Nov. Act. Nat. Cur. xix. Suppl ii. (1841) 345 = tetragona.
Birdii, *F. Muell. Fragm.* x. 26.—Australi.
brachychaeta, *F. Muell. in Trans. Phil. Inst. Vict.* iii (1859) 48.—Austral.
brachyphylla, *A. Cunn. ex Steud. Nom. ed.* II. i. 264 —Australi.
brachyphylla, *Turcz. in Bull. Soc.Nat. Mosc.* xx. (1847) i. 161.—Austral.
brevifolia, *Meissn. in Journ. Linn. Soc.* i. (1857) 46 —Austral.
breviseta, *Lindl. Swan River App.* 5.—Austral.
Brownii, Schau. in Nov. Act. Nat. Cur. xix. Suppl ii. (1841) 360 = tetragona.
brunioides, A. Cunn. in Bot. Mag. sub t. 3823 = tetragona.
Candollenana, Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 360 = C. decandra.
ciliata, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847 i. 161 = empetroides.
conanthera, F. Muell. Fragm. i. 146 = decandra.
conferta, *A. Cunn. in Bot. Mag.* sub t. 3823.—Austral.
Creswelli, *F. Muell. Fragm.* x. 27.—Australi.
cupressoides, A. Rich. Sert. Astrol. 41. t. 16 = microphylla.
cuprestoides, A. Rich. l. c. 43 = microphylla.
curtophylla, A. Cunn. in Bot. Mag. sub t. 3823. flavescens.
curtophylla, Schau. in Nov. Act. Nat. Cur. xix. Suppl ii. (1841) 242 = Leschenaultii.
cuspidata, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i 162 = breviseta.
decandra, *DC. Prod.* iii. 208.—Australi.
depressa, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i 162.—Australi.
diversifolia, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 238.—Austral.
Drummondii, Meisn. in Journ. Linn. Soc. i. (1857) 47 = flavescens.
empetroides, *Schau. in Nov. Act. Nat. Cur.* xix. Suppl. ii. (1841) 354.—Australi.
ericoides, A. Cunn. in Field, N. S. Wales, 350 = tetragona.
exstipulata, DC. Prod. iii. 208 = microphylla.
flavescens, *A. Cunn. in Bot. Mag.* sub t. 3823.—Austral.
Fraseri, *A. Cunn. l. c.*—Austral.
glabra, R. Br. in Bot. Reg. t. 409 = tetragona.
glutinosa, *Lindl. Swan River App.* 5.—Austral.
gracilis, *Benth. Fl. Australi.* iii. 45.—Australi.
granulosa, *Benth. l. c.* 49.—Australi.
laticina, *R. Br. ex Benth. l. c.* 52.—Australi.
lastantha, Meisn. in journ. Linn. Soc. i. (1857) 46 = strigosa.
lasiocalyx, F. Muell. Fragm. i. 224 = sapphirina.
leptophylla, *Benth. Fl. Australi.* iii. 50.—Australi.
Leschenaultii, *Schau. in Lehm. Pl. Preiss.* i. 104.—Austral.
linearifolia, Miq. in Nederl. Kruidk. Arch. iv. (1859) 117 = tetragona.
longiflora, *F. Muell. Fragm.* i. 12.—Austral.
lutea, Schau. in Lehm. Pl. Preiss. i. 106 = flavescens.
megaphylla, *F. Muell. Fragm.* i. 146.—Australi.
microphylla, *A. Cunn. in Bot. Mag.* sub t. 3823.—Austral.
monticola, Miq. in Nederl. Kruidk. Arch. iv. (1859) 118 = tetragona.
Muelleri, Miq. l. c. 119 = tetragona.
muricata, *F. Muell. Fragm.* i. 224.—Australi.
Oldfieldii, *Benth. Fl. Australi.* iii. 46.—Australi.
pinnulosa, *F. Muell. Fragm.* x. 27.—Australi.
propinqua, A. Cunn. ex Walp. Rep. ii. 138 = tetragona.
pulchella, *Meisn. in journ. Linn. Soc.* i. (1857) 47.—Austral.
pulchella, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 238.—Australi.
rosea, Meisn. in journ. Linn. Soc. i. (1857) 46 = tenuifolia.
rosea, Miq. in Nederl. Kruidk. Arch. iv. (1859) 117 = tetragona.
sapphirina, *Lindl. Swan River App.* 5.—Australi.
scabra, DC. Prod. iii. 208 = tetragona.

CALYTHRIX :—
Schlechtendahlii, Miq. in Nederl. Kruidk. Arch. iv. (1859) 116 = tetragona.
simplex, *Lindl. Swan River App.* 5.—Austral.
sparsa, A. Cunn. ex Walp. Rep. ii. 160 = strigosa.
squarrosa, Miq. in Nederl. Kruidk. Arch. iv. (1859) 118 = tetragona.
strigosa, *A. Cunn. in Bot. Mag.* sub t. 3828.—Austral. Sullivani, *F. Muell. Fragm.* ix. 1.—Austral.
tenuifolia, Meisn. in Journ. Linn. Soc. i. (1857) 47 = flavescens.
tenuifolia, *Meisn. l. c.* 46.—Austral.
tenuiramea, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 20.—Austral.
tetragona, *Labill. Nov. Holl. Pl.* ii. 8. t. 146.—Austral.
tetragonophylla, Meisn. in Journ. Linn. Soc. i. (1857) 47 = flavescens.
tetraptera, DC. Prod. iii. 308 = tetragona.
variabilis, *Lindl. Swan River App.* 5.—Austral.
virgata, A. Cunn. in Bot. Mag. sub t. 3828 = tetragona.

CALYTRIPLEX, Ruiz & Pav. Prod. 96 (1794) = **Herpestis**, Gaertn. (Scrophular.).
obovata, Ruiz & Pav. Syst. Veg. 166 = H. Monniera.

CALYXHYMENIA, Orteg. Hort. Matr. Dec. 5. t. 1 (1798) = **Corybanthus**, Vahl (Nyctagin.).
aggregata, Orteg. l. c. 31. t. 11 = O. aggregatus.
Cervantesii, Desf. Cat. Hort. Par. ed. III. 390 = O. Cervantesii.
chilensis, Desf. ex Steud. Nom. ed. II. i. 364 = O. chilensis.
expansa, Ruiz & Pav. Fl. Per. i. 45 = O. expansus.
glabrifolia, Orteg. Hort. Matr. 5. t. 1 = O. glabrifolius.
ovata, Ruiz & Pav. Fl. Per. i. 45 = O. ovatus.
paniculata, Desf. Cat. Hort. Par. ed. III. 390 = O. ovatus.
pilosa, Choisy, in DC. Prod. xiii. II. 438 = O. angustifolius.
prostrata, Ruiz & Pav. Fl. Per. i. 46 = O. prostratus.
viscosa, Orteg. Hort. Matr. Dec. i. 6, in obs. = O. viscosus.

CAMACUM, 'Adans.' ex Steud. Nom. ed. II. ii. 174, in syn. (1841) = **Myristica**, Linn.

CAMAION, Rafin. Sylva Tellur. 75 (1838) = **Helicteres**, Linn. (Sterculiac.).
hirsuta, Rafin. l. c. = H. hirsuta.
undulata, Rafin. l. c. = Sterculia lanceolata.

CAMARA, Plum. ex Linn. Syst. ed. I (1735); Adans. Fam. ii. 199 (1763) = **Lantana**, Linn. (Verbenac.).
tiliaefolia, Benth. in Hook. Journ. Bot. ii. (1840) 58 = L. tiliaefolia.
vulgaris, Benth. Bot. Voy. Sulph. 154 = L. Camara.

CAMARRA, St. Hil. in Bull. Soc. Philom. (1822) 188. *MALPIGHIACEAE*, Benth. & Hook. *f.* i. 361.
affinis, *St. Hil. Pl. Rem.* 57.—Bras.
axillaris, *St. Hil. in Bull. Soc. Philom.* (1822) 188.—Bras.
discolor, *Griseb. in Mart. Fl. Bras.* xii. 1. 105.—Bras.
ericoides, *St. Hil. in Bull. Soc. Philom.* (1822) 188.—Bras.
hirsuta, *St. Hil. l. c.*—Bras.
juncea, *Griseb. in Mart. Fl. Bras.* xii. 1. 106.—Bras.
linearifolia, *St. Hil. in Bull. Soc. Philom.* (1822) 188.—Bras.
phylicoides, Mart. ex Walp. Rep. v. 351 = axillaris.
pulchella, *Griseb. in Mart. Fl. Bras.* xii. 1. 106.—Bras.
sericea, *St. Hil. in Bull. Soc. Philom.* (1822) 188.—Bras.
tripbylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 601.—Bras.

CAMARTIDIUM, Lindl. Bot. Reg. sub t. 844 (1824). *ORCHIDEAE*, Benth. & Hook. *f.* Gen. iii. 554.
AMARIDIUM, Hort. ex Lubbers, Cat. Pl. Rar. San Donato, 15 (1860).
elatum, *Lindl. in Benth. Pl. Hartw.* 158.—Ecuador.
Arbuscula, *Lindl. l. c.*—Ecuador.
brevifolium, *Lindl. l. c.* 154.—N. Granad.

CAMARIDIUM :—
graminifolium, *Reichb. f. in Linnaea*, xxii. (1849) 857.—Peruv.
inflexum, *Lindl. in Ann. Nat. Hist.* v. (1840) 116.—Ins. Dominica.
lanciflolium, *Reichb. f. in Linnaea*, xxii. (1849) 857.—Peruv.
longibracteatum, *Lindl. in Benth. Pl. Hartw.* 154.—Peruv.
luteo-rubrum, *Lindl. Orch. Linden.* 22.—Venezuela.
lutescens, *Reichb. f. in Walp. Ann.* vi. 541.—Bras.
ochroleucum, Lindl. Bot. Reg. x. (1824) 844 = Maxillaria Camaridii.
pendulum, *Rodrig. Orch. Nov.* i. 128.—Bras.
purpuratum, *Lindl. Orch. Linden.* 22.—Venezuela.
purpureum, Spreng. Syst. iii. 735 = Cymbidium ensifolium.
robustum, *Rodrig. Orch. Nov.* ii. 210.—Bras.
vestitum, Lindl. in Ann. & Mag. Nat. Hist. Ser. III. i. (1858) 353 = Ornithidium vestitum.

CAMARILLA, Salisb. Gen. Pl. Fragm. 91 (1866) = **Allium**, Tourn. (Liliac.).
obliqua, Salisb. l. c. = A. obliquum.

CAMAROTIS, Lindl. Gen. et Sp. Orch. 219 (1833) = **Sarcochilus**, R. Br. (Orchid.).
apiculata, Reichb. f. in Bonplandia, v. (1857) 89.—Malaya.
cochinchinensis, Reichb. f. in Gard. Chron. (1876) ii. 740.—Cochinch.
obtusa, Lindl. Bot. Reg. xxx. (1844) Misc. 71 = S. obtusus.
pallida, Lindl. in Journ. Linn. Soc. iii. (1859) 87 = S. pallidus.
philippinensis, Lindl. l. c. = S. philippinensis.
purpurea, Lindl. in Wall. Cat. n. 7329 = S. purpureus.
rostrata, Reichb. f. in Walp. Ann. vi. 881 = S. purpureus.

CAMASSIA, Lindl. Bot. Reg. t. 1486 (1832). *LILIACEAE*, Benth. & Hook. *f.* iii. 815.
BULBEDILIS, Rafin. Fl. Tellur. ii. 26 (1836).
CYANOTRIS, Rafin. in Am. Monthly Mag. (1811) 356; (1819) 192, nomen prius.
LEMOTRIS, Rafin. Fl. Tellur. ii. 26 (1836).
QUAMASIA, Rafin. Cat. 15 (1824); QUAMASSIA, Rafin. Fl. Tellur. ii. 25 (1836).
SITOCODIUM, Salisb. Gen. Pl. Fragm. 27 (1866).
alba, Hort.—Cf. Gard. Chron. (1875) i. 662 = Leichtlinii.
esculenta, *Lindl. Bot. Reg.* t. 1486.—Am. bor. occ.
Fraseri, *Torr. in Rep. Facif. Rail.* iv. 91.—Am. bor. occ.
Leichtlinii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 376.—Calif.

CAMAX, Schreb. Gen. i. 144 (1789) = **Ropourea**, Aubl. (Verbenac.).
frazinea, Willd. Sp. Pl. i. 1117 = R. guianensis.
guianensis, J. F. Gmel. Syst. 408 = R. guianensis.

CAMBANIA, Comm. ex M. Roem. Syn. Hesper. 88, 102 (1846) = **Dyscolyum**, Blume (Meliac.).
Fraxerriana, M. Roem. l. c. 102 = D. Fraxeranum.

CAMBDERIA, Steud. Nom. ed. II. i. 364 (1840) = CAMPDERIA, Rich. = **Vellozia**, Vand. (Amaryll.).

CAMBEA, Endl. Gen. 1284, in Syn. (1840) sphalm. = CUMBIA, Buch.-Ham. = **Careya**, Roxb. (Myrtac.).

CAMBESSEDEA, Wight & Arn. Prod. i. 170 (1834) = **Bouea**, Meisn. (Anacard.).
oppositifolia, Wight & Arn. l. c. = B. burmanica.

CAMBESSEDEA, Kunth, in Ann. Sc. Nat. Sér. I. ii. 336 (1824) = **Buchanania**, Roxb. (Anacard.).
axillaris, Kunth, l. c. = B. angustifolia.

CAMBESSEDESIA, DC. Prod. iii. 110 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 787.
ACIPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 377, pro parte.
adamantium, *DC. Prod.* iii. 111.—Bras.
adamantium, Miq. in Linnaea, xxii. (1849) 587 = Ilicifolia.
arenaria, *Cogn. in Mart. Fl. Bras.* xiv. III. 15.—Bras.
balsamifera, DC. Prod. iii. 110 = Microlicia balsamifera.
bidentata, DC. l. c. 111 = Hilariana.
corymbosa, *DC. l. c.* 110.—Bras.
corymbosa, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 61 = rugosa.
crenulata, DC. Prod. iii. 111 = Microlicia crenulata.
Eichleri, *Cogn. in Mart. Fl. Bras.* xiv. III. 11.—Bras.
Espora, *DC. Prod.* iii. 111.—Bras.
Espora, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 62 = Ilicifolia.
Hilariana, *DC. Prod.* iii. 111.—Bras.
ilicifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 24.—Bras.
intermedia, Mart. Herb. Fl. Bras. n. 940 = rugosa.
latevenosa, *DC. Prod.* iii. 110.—Bras.
membranacea, *Gardn. in Field. Sert. Pl.* t. 64.—Bras.
paraguayensis, *Hook. f. Bot. Mag.* t. 6604.—Paraguay.
purpurata, *DC. Prod.* iii. 110.—Bras.
Regnelliana, *Cogn. in Mart. Fl. Bras.* xiv. III. 19.—Bras.
rugosa, *Cogn. l. c.* 14.—Bras.
setacea, *Cogn. l. c.* 23.—Bras.
sincorensis, DC. Prod. iii. 111 = Microlicia sincorensis.
terminalis, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 61 = Hilariana.
Weddelii, *Naud. l. c.*—Bras.

CAMBOGIA, Linn. Fl. Mart. Burser. (1745); Dios. Dess. 18 (1747) (Amoen. Acad. i. 171, 402) = **Garcinia**, Linn. (Guttifer.).
Buncca, Blanco, Fl. Filip. ed. I. 484 = G. Cambogia.
crassifolia, Blanco, l. c. ed. II. 304 = G. Cowa.
Gutta, Linn. Gen. ed. V. Add. Fin. = G. Cambogia.
solitaria, Stokes, Bot. Mat. Med. iii. 175 = G. Cambogia.
venulosa, Blanco, Fl. Filip. ed. I. 435 = G. venulosa.

CAMDENIA, Scop. Introd. 190 (1777) = **Evolvulus**, Linn. (Convolvul.).

CAMDERIA, Dum. Anal. Fam. 80 (1829) = **Lachnanthes**, Ell. (Haemod.).

CAMELINA, Crantz, Strip. Austr. ed. I. fasc. I. 18 (1762); ed. II. 18 (1769). *CRUCIFERAE*, Benth. & Hook. f. i. 88.
CRAMARLINUM, Host, Fl. Austr. ii. 224 (1831).
LINOSTROPHUM, Schrank, Fl. Prim. Salisb. 168 (1792).
SINISTROPHORUM, Schrank, ex Endl. Gen. 882 (1840).
albiflora, Kotschy, ex Boiss. Fl. Orient. i. 312 = sylvestris.
ambigua, Bess. ex Steud. Nom. ed. II. i. 264 = sativa.
anomala, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 313. —Syria.
apetala, Brot. Fl. Lusit. i. 560 = Nasturtium amphibiam.
armeniaca, Desv. Journ. Bot. iii. (1814) 169 = sylvestris.
austriaca, Pers. Syn. ii. 191 = Nasturtium austriacum.
barbarefolia, DC. Syst. ii. 517 (= *Tetrapoma barbarae-folium*.).
Caisir, Wall. Cat. n. 4803 = Cochlearia flava.
Camisir, Buch.-Ham. ex Steud. Nom. ed. II. i. 264, sphalm. = prsec.
campestris, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 959 = sylvestris.
dentata, Pers. Syn. ii. 191 = sativa.
edentula, Desv. Journ. Bot. iii. (1814) 170 = Alyssum edentulum.
foetida, *Fries, Nov. Mant.* iii. 70.—Europ.
grandiflora, *Boiss. Diagn. Ser.* I. v. 38.—As. Min.
hirsuta, Bernh. Syst. Verz. Erf. 190 = sativa!
hispida, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 176.—Armenia; Persia.
Isaiocarpa, *Boiss. & Bal. ex Boiss. Fl. Orient.* i. 312.—Syria.
laxa, *C. A. Mey. Verz. Pfl. Cauc.* 198.—As. Min.; Persia.
Linicola, *Schimp. & Spenn. in Spenn. Fl. Friburg.* iii. 958.—Europ.

CAMELINA :—
macrocarpa, Wierzb. ex Reichb. Ic. Fl. Germ. ii. 10 = sativa.
microcarpa, Andrz. ex DC. Syst. ii. 617 = sylvestris.
myagrodes, Moretti, in Giorn. Fis. Pav. (1820) 279 = Cochlearia saxatilis.
pinnatifida, Hornem. Hort. Hafn. ii. 598 = foetida.
sagittata, Moench, Meth. 265 = sativa.
sativa, Boiss. Voy. Bot. Espagne, 49 = sylvestris.
sativa, *Crantz, Stirp. Austr.* ed. I. fasc. i. 14; ed. II. 18.—Europ. : As. temp.
saxatilis, Pers. Syn. ii. 191 = Cochlearia saxatilis.
sylvestris, *Wallr. Sched. Crit.* 347.—Europ. ; As. Min.

CAMELLIA, Linn. Syst. ed. I (1735). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 167.
CALPANDRIA, Blume, Bijdr. 178 (1825).
DESMITUS, Rafin. Sylva Tellur. 189 (1838).
DRUPIFERA, Rafin. l. c. 140 (1836)
KEMELIA, Rafin. l. c. 138, 139 (1836).
SALCEDA, Blanco, Fl. Filip. ed. II. 374 (1854).
SASANQUA, Nees, in Flora, iv. (1824) Lit. 144.
THEA, Linn. Gen. ed. I. 154 (1737).
THEAPHYLLA, Rafin. in Loud. Gard. Mag. vii. (1832) 246.
TSIA, Adans. Fam. ii. 450 (1763).
TSUBAKI, Adans. l. c. 399 (1763).
assimilis, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 209.—Hongkong.
axillaris, Roxb. ex Ker-Gawl. in Bot. Reg. t. 349; et Sims, Bot. Mag. t. 2047 = Gordonia anomala.
axillaris, Wall. Cat. n. 158 = Adinandra acuminata.
Bohea, Lindl. in Loud. Encycl. Pl. 892; Griff. Notul. iv. 558 = C. Thea.
caudata, Griff. Notul. iv. 559 = drupifera.
caudata, *Wall. Cat.* n. 978.—Reg. Himal.
Chamgota, Buch.-Ham. ex Wall. in As. Res. xiii. (1820) 429 = C. Sasanqua.
drupifera, *Lour. Fl. Cochinch.* ii. 611.—Reg. Himal. ; Burma ; China.
Edithae, *Hance, in Ann. Sc. Nat. Sér.* IV. xv. (1861) 221.—China.
euryoides, *Lindl. Bot. Reg.* t. 988.—China.
florida, Salisb. Prod. 370 = japonica.
fraterna, *Hance, in Ann. Sc. Nat. Sér.* IV. xviii. (1862) 218.—China.
grandiflora, *Forsk. Fl. Aegypt. Arab.* 196.—Arab. Grijsli, *Hance, in Journ. Bot.* xvii. (1879) 9.—China.
hongkongensis, *Seem. in Trans. Linn. Soc.* xxii. (1859) 342. t. 40.—China.
integrifolia, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 150 = Actinodaphne chinensis.
japonica, Champ. in Hook. Kew Journ. iii. (1851) 209 = hongkongensis.
japonica, *Linn. Sp. Pl.* 698.—Japon.
japonica, Wall. Cat. n. 3667 = Adinandra integerrima.
Kaempferia, *Retzius, in Act. Soc. Reun. Sc. Ital.* (1841) 494.—Hab. ?
Kwina, Buch.-Ham. in D. Don, Prod. Fl. Nep. 224 = drupifera.
Kissi, Wall. in As. Res. xiii. (1820) 429 = drupifera.
lanceolata, *Seem. in Trans. Linn. Soc.* xxii. (1859) 345. —Malaya.
latescens, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 293.—Reg. Himal.
maliflora, Lindl. Bot. Reg. t. 1078 = roseaflora.
Masterzia, Griff. Notul. iv. 559 = drupifera.
mutabilis, Paxt. Mag. Bot. ii. 122 = japonica.
oleifera, Abel, Journ. in China, 174 = C. Sasanqua.
oleifera, Wall. Cat. n. 976 = drupifera.
oleosa, Lour. Fl. Cochinch. 339 = C. Sasanqua.
planipetala, Lam. Illustr. Hortic. (1866) t. 496 = japonica.
Quinoaana, *Seem. in Trans. Linn. Soc.* xxii. (1859) 345.—Java.
reticulata, *Lindl. Bot. Reg.* t. 1078.—China.
roseaflora, *Hook. Bot. Mag.* t. 5044.—China.
salicifolia, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 209.—Hongkong.
Sasanqua, Blanco, Fl. Filip. ed. I. 530 = drupifera.
Sasanqua, *Thunb. Fl. Jap.* 273. t. 30.—Japon. ; China.
Scottiana, Choisy, Mém. Terntr. 90 = Adinandra dumosa.
Scottiana, Wall. Cat. n. 3668 = C. Thea.
simplicifolia, Griff. Notul. iv. 560. t. 604. f. 2 = drupifera.

CAMELLIA :—
spectabilis, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 210 = reticulata.
symplocifolia, Griff. Itin. Notes, 40. n. 652 = drupifera.
Thea, *Link, Enum. Hort. Berol.* ii. 78.—China.
theifera, Griff. Notul. iv. 558. t. 601. f. 1, 3 = C. Thea.
theifolia, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 221 = euryoides.
Tsubakki, Crantz, Inst. ii. 172 = japonica.
viridis, Link, Enum. Hort. Berol. ii. 78 = C. Thea.

CAMERARIA, Dill. ex Moench, Meth. 630 (1794) = **Montia**, Linn. (Portulac.).
fontana, Moench, l. c. = M. fontana.

CAMBRARIA, [Plum.] Linn. Syst. ed. I (1735). *APOCYNACEAE*, Benth. & Hook. f. ii. 701.
angustifolia, *Linn. Sp. Pl.* 210.—Am. sept.
dubia, Sims, Bot. Mag. t. 1646 = Wrightia dubia.
guianensis, Aubl. Pl. Gui. i. 262 = Malouetia Tamaquarina.
latifolia, *Linn. Sp. Pl.* 210.—Ind. occ.
lutea, Lam. Encyc. i. 578 = Malouetia Tamaquarina.
obesa, Spreng. Syst. i. 641 = Adenium obesum.
retusa, *Griseb. Fl. Brit. W. Ind.* 410.—Ind. occ.
Tamaquarina, Aubl. Pl. Gui. i. 256 = Malouetia Tamaquarina.
myrtifolia, Retz. Obs. iv. 24 = Hunteria corymbosa.
myrtifolia, Wight, ex Wall. Cat. n. 4459 = Strophanthus Wightianus.

CAMERIDIUM, Reichb. f. in Linnaea, xxii. (1849) 857, sphalm. = **Camaridium**, Lindl. (Orchid.).

CAMFOROSMA, Spreng. Syst. i. 453 (1825) = **Camphorosma**, Linn. (Chenop.).

CAMIRIUM, Gaertn. Fruct. ii. 194. t. 125 (1791) = **Alouirites**, Forst. (Euphorbiac.).
cordifolium, Gaertn. l. c. 195 = A. moluccana.
oleosum, Reinw. ex Blume, Cat. Gew. Buitenz. 104 = A. moluccana.

CAMISSONIA, Link, in Jahrb. Gew. i. (1818) 1. 186 = **Oenothera**, Linn. (Onagrar.).
flava, Link, l. c. = O. dentata.

CAMMARUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 287 = **Aconitum**, Tourn. (Ranuncul.).
paniculatum, Fourr. l. c. = A. paniculatum.

CAMOCLADIA, Linn. Syst. ed. X. 861 (1756) = **Comocladia**, P. Br. (Anacard.).

CAMOENSIA, Welw. ex Benth. & Hook. f. Gen. i. 557 (1865). *LEGUMINOSAE*, Benth. & Hook. f. i. c.
brevicalyx, *Benth. in Trans. Linn. Soc.* xxv. (1866) 309.—Afr. trop.
maxima, *Welw. ex Benth. l. c.* 309. t. 36.—Afr. trop.

CAMOMILLA, Gilib. Exercit. i. 178 (1792) = **Matricaria**, Tourn. (Compos.).
defexa, Gilib. l. c. 179 = M. suaveolens.
inodora, Gilib. l. c. = M. inodora.
patens, Gilib. l. c. 178 = M. Chamomilla.

CAMONEA, Rafin. Fl. Tellur. iv. 81 (1836) = **Ipomoea**, Linn. (Convolvul.).
bifida, Rafin. l. c. = I. cymosa.

CAMPANEA, Decne. in Rev. Hortic. (1849) 241. t. 13. *GESNERACEAE*, Benth. & Hook. f. ii. 1003.
grandiflora, *Decne. l. c.*—N. Granat.
Himalayana, *Hanst. in Linnaea*, xxvi. (1855) 214. t. 1. f. 24.—Costa Rica.
Ourstedii, *Oerst. in Videnskab. Selsk. Skr.* V. v. (1861) 108.—Costa Rica.

CAMPANULA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 46 (1737). *CAMPANULACEAE*, Benth. & Hook. f. ii. 561.
APENULA, Neck. Elem. i. 284 (1790).
BALLELA, Rafin. Fl. Tellur. iv. 70 (1836).
CERVENIA, Opiz, Seznam, 57 (1852).
DECAPHYSSA, Rafin. l. c. ii. 78 (1836).
DEPIERREA, Anon. ex Schlecht. in Linnaea, xvi. (1842) 874.

CAMPANULA :—
DRYMOCODON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 111.
ERINIA, Noulet, Fl. Sous-Pyr. 407.
HECALE, Rafin. Fl. Tellur. ii. 79 (1836).
HETEROCODON, Nutt. in Trans. Am. Phil. Soc Ser. II. viii. (1842) 265.
LACARA, Rafin. Fl. Tellur. iv. 70 (1836).
LOBELIA, Rafin. l. c. iii. 82 (1836).
MARIANTHEMUM, Schrank, in Denkschr. Regensb. Ges. ii. (1822) 84.
MEDIUM, [Tourn.] Fisch. ex A. DC. Monog. Camp. 218 (1830).
NENNINGIA, Opiz, Seznam, 68 (1852).
PENTROPIS, Rafin. Fl. Tellur. ii. 80 (1836).
QUINQUELOCULARIA, C. Koch, in Linnaea, xxiii (1850) 680.
RAPUNCULUS, Fourr. in Ann. Soc. Linn. Lyon, N. S xvii. (1869) 111.
ROUCELIA, Willk. in Willk. & Lange, Prod. Fl Hisp. ii. 389 (1870).
ROUCELIA, Dum. Comm. Bot. 14 (1822).
SYKORAEA, Opiz, Seznam, 94 (1852).
SYNCODON, Fourr. in Ann. Soc. Linn. Lyon, N S xvii. (1869) 111.
TALANELIS, Rafin. Fl. Tellur. iv. 70 (1836).
TRACHELIOPSIS, Opiz, Seznam, 98 (1852).
WEITENWEBERA, Opiz, l. c. 104 (1852).
abietina, *Griseb. & Schenk, in Wiegm. Archiv*, xv (1852) 1. 333.—Europ.
acuminata, Michx. Fl. Bor. Am. i. 108 = americana.
acutangula, *Leresche & Levier, in Journ. Bot.* xvii (1879) 196.—Hispan.
acutiloba, *Vahl, in Linnaea*, xxxviii. (1874) 709.—Kurdistan.
Adami, Bieb. Fl. Taur. Cauc. i. 153 = bellidifolia.
Adami, Willd. ex Steud. Nom. ed. II. i. 285 = lasiocarpa.
adharensi, Ehrenb. ex Boiss. Fl. Orient. iii. 948 = retrorsa.
adpressa, Linn. f. Suppl. 140 = Wahlenbergia adpressa adscendens, *Vest. ex Roem. & Schult. Syst.* v. 90.—Sibir.
adscendens, *Leresche & Levier, in Journ. Bot.* xvii. (1879) 199.—Hispan.
affinis, Fisch. ex Kupr. in Bull. Acad. Petersb. xi (1867) 305 = Aucheri.
affinis, Reichb. ex Nym. Consp. 480 = Portenschlagiana affinis, *Roem. & Schult. Syst.* v. 140.—Hispan.
algasica, *Pomel, Nouv. Mat. Fl. Atl.* 257.—Afr. bor.
alpa, Cav. in Anal. Clenc. Nat. iii. (1801) 21 = dichotoma.
aggregata, Willd. Enum. Hort. Berol. Suppl. 10 = glomerata.
agrestis, Wall. Cat. n. 1292 = Wahlenbergia gracilis Aixoon, *Boiss. & Sprun. Diagn.* Ser. I. iv. 84.—Graecia
ajugaefolia, *Sestini, ex Spreng. Syst.* i. 732 (sp. dub.). —As. Min.
alata, *Desf. Fl. Atlant.* i. 178. t. 50.—Afr. bor.
Alberti, *Trautv. in Act. Hort. Petrop.* vi. (1879) 83.—Turkestan.
albiflora, C. Koch, in Linnaea, xxiii. (1850) 684 = sarmatica.
alburnica, Brigant. Pempt. i. 3. t. 2 = Phyteuma amplexicaule.
algida, Fisch. ex A. DC. Monog. Camp. 338 = lasiocarpa.
alliariaefolia, Reichb. Pl. Crit. Cent. iii. 345 = Symphyandra pendula.
alliariaefolia, *Willd. Sp. Pl.* i. 910.—Reg. Caucas. ; As. Min.
Allionii, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
Allionii, *Vill. Fl. Delph.* 18; *Prosp.* 22.—Europ.
allophylla, Rafin. ex A. DC. Monog. Camp. 332 = rotundifolia.
alopecuroides, Willd. ex Steud. Nom. ed. II. i. 285 = Phyteuma Sibthorpianum.
alpestris, All. Fl. Pedem. 113. t. 6. f. 3 = Allionii.
Alphonsii, *Wall. Cat.* n. 1296.—Ind. or.
alpigena, C. Koch, in Linnaea, xxii. (1849) 638 = Aucheri.
alpina, *Jacq. Enum. Vind.* 26; 210.—Europ.
Alpini, Linn. Sp. Pl. ed. II. 1662 = Adenophora lilifolia.
Alpini, Ptchl. ex Vis. Fl. Dalm. Suppl. 74 = Phyteuma amplexicaule.
alsinoides, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 94.—Reg. Himal.
altaica, A. DC. Monog. Camp. 220. t. 20 = pilosa.

CAMPANULA:—
altaica, Ledeb. Ind. Sem. Hort. Dorpat. (1824) 2 = Steven.
altiflora, Pers. Syn. i. 192 = Prismatocarpus altiflorus.
americana, Hort. ex Steud. Nom. ed. II. i. 266 = Phyteuma canescens.
americana, *Linn. Sp. Pl.* 164.—Am. bor.
americana, Mill. Gard. Dict. ed. VIII. n. 18 = planiflora.
amplexicaulis, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma amplexicaule.
amplexicaulis, Michx. Fl. Am. Bor. i. 108 = Specularia perfoliata.
amygdalifolia, Salisb. Prod. 226 = persicaefolia.
anagalloides, Royle, Illustr. Bot. Himal. 253 = Cephalostigma hirsutum.
anchusiflora, *Sibth. & Sm. Fl. Graec. Prod.* i. 141. —Archip. Graec.
andina, Rupr. in Bull. Acad. Pétersb. xi. (1867) 215 = gumbetica.
Andrewsii, A. DC. Monog. Camp. 220 = tomentosa.
androsacea. D. Dietr. Syn. Pl. i. 753 = Wahlenbergia androsacea.
angulata, Kaſin. Fl. Ludov. 55 = Specularia perfoliata.
angustifolia, Lam. Tabl. Encyc. ii. 58 = linifolia.
anisocrata, Kunze, ex A. DC. in DC. Prod. vii. 491 = Specularia perfoliata.
antirrhina, Schleich. Cat. Pl. Helv. ed. III. 10 = caespitosa, rotundifolia.
aparinoides, *Pursh, Fl. Am. Sept.* i. 159.—Am. bor.
appendiculata, A. DC. Prod. vii. 462 = tomentosa.
arborescens, P. Br. (err. R. Br.) ex Steud. Nom. ed. II. i. 266 = Hamelia ventricosa.
arcuata, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 138 = rotundifolia.
ardonensis, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 212.—Reg. Caucas.
argaea, *Boiss. & Bal. Diagn. Ser.* II. vi. 119.—Cappadoc.
argenica, *Lam. Encyc.* i. 584.—Armenia.
argunensis, Rupr. in Bull. Acad. Pétersb. xi. (1867) 209 = Aucheri.
argyrotricha, *Wall. Cat.* n. 7138.—Reg. Himal.
arida, H. B. & K. Nov. Gen. et Sp. iii. 301 = Wahlenbergia linarioides.
arista, *Wall. Cat.* n. 1291; *et in Roxb. Fl. Ind. ed. Carey*, ii. 98.—Reg. Himal.
armena, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 286 = Symphyandra armena.
arvatica, Lag. in Varied. Cienc. ii. iv. (1805) 40 = Wahlenbergia hederacea.
arvensis, Pers. Syn. i. 198 = Specularia Speculum.
aspera, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma asperum.
aspera, Donn, ex A. DC. Monog. Camp. 290 = sparinoides.
aspera, *Moench, Meth. Suppl.* 188.—Hab.?
asperrima, Zuccagni, in Koem. Collect. 238 = sjuganefolia.
asteroides, Lam. Tabl. Encyc. ii. 55 = americana.
athoa, Boiss. & Heldr. Diagn. Ser. II. iii. 110 = C. Trachelium.
attenuata, Ledeb. ex Spreng. Syst. i. 739 = persicifolia.
attica, Boiss. & Heldr. Diagn. Ser. I. xi. 67 = drabaefolia.
Aucheri, *A. DC. in DC. Prod.* vii. 460.—Reg. Caucas.; As. Min.
aurea, Linn. f. Suppl. 141 = Musschia aurea.
axillaris, *Boiss. & Bal. Diagn.* Ser. II. iii. 109.—Cilicia.
azurea, Soland. & Banks, ex Sims, Bot. Mag. t. 651 = rhomboides.
baicalensis, Pall. ex A. DC. Prod. vii. 479 = silenifolia.
Balansae, *Boiss. & Reut. Fl. Orient.* iii. 981.—Cappadoc.
baldensis, Balb. Cat. Hort. Taur. (1813) 20 = ramosissima.
Banksiana, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia Banksiana.
barbata, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
barbata, *Linn. Syst.* ed. X. 937.—Europ.
barbata, Spreng. Fl. Halens. 77 = glomerata.
Barrelieri, *Marnock, Floric. Mag.* i. (1886) 161. t. 12. —Italia.
Barrelieri, Prusl, Symb. Bot. 30. t. 19 = fragilis.
Baumgarteni, Becker, Fl. Francof. i. 364 = rotundifolia.

CAMPANULA:—
Bayeriana, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 214.—Reg. Caucas.
bellardi, All. Fl. Pedem. i. 108 = caespitosa.
bellidifolia, *Adam, in Web. & Mohr, Beitr.* i. (1805) 47.—Reg. Caucas.
bellidifolia, Frivald. ex Nym. Consp. 476 = Orphanidea.
bellidifolia, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 36 = patula.
Benthamii, Wall. Cat. n. 1290 = canescens.
Bergiana, D. Dietr. Syn. Pl. i. 757 = Prismatocarpus subulatus.
betonicaefolia, Biehl. Pl. Nov. Hort. Spreng. 14 = sarmatica.
betonicaefolia, Gilib. Fl. Lituan. i. 53 = glomerata.
betonicaefolia, *Sibth. & Sm. Fl. Graec. Prod.* i. 141.—As. Min.
betulaefolia, *C. Koch, in Linnaea*, xxiii. (1850) 685.—Armenia.
bicaulis, Lapeyr. Hist. Abr. Pl. Pyr. 107. t. 7 = speciosa.
bicolor, Boiss. Fl. Orient. iii. 981 = camptoclada.
Biebersteiniana, C. A. Mey. Verz. Pfl. Cauc. 88, pro parte = Aucheri.
Biebersteinsana, Roem. & Schult. Syst. v. 147 = tridentata.
Bielzii, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 97 = rotundifolia.
biflora, Ruiz & Pav. Fl. Per. ii. 55 = Specularia biflora.
Billardieri, A. DC. Monog. Camp. 303. t. 13 = C. Cymbalaria.
biserrata, C. Koch, in Linnaea, xix. (1847) 29 = lactiflora.
bithynica, A. DC. in DC. Prod. vii. 460 = tridentata.
Bocconi, J. F. Gmel. Syst. 348 = rhomboides.
Bocconi, Vill. Hist. Pl. Dauph. ii. 502 = pusilla.
Bosseri, Sprun. ex Boiss. Fl. Orient. iii. 915 = radicosa.
Boissieri, Vatke, in Linnaea, xxxviii. (1874) 708 = Phyteuma asperum.
Boissieri, *Lojac. in Anal. Soc. Esp. Hist. Nat.* viii. (1879) 451.—Hispan.
bononiensis, *Linn. Sp. Pl.* 165.—Europ. austr.; Reg. Caucas.; Sibir.
brachiata, Schm. ex A. DC. Prod. vii. 462 = dichotoma.
brachiata, Seidl, ex Opiz, Boehems Gew. 168; Naturul. ii. (1823) (8) = patula.
brachyscpala, Vukot. in Linnaea, xxvi. (1853–55) 825 = cenisia.
bracteata, Thunb. in Hoffm. Phytogr. Blaett. i. (1803) 20 = Roella bracteata.
brevibracteata, D. Dietr. Syn. Pl. i. 757 = Microcodon glomeratum.
Brounsiana, Roem. & Schult. Syst. v. 104 = Loeflingii.
Brunonis, Wall. Cat. n. 1285 = latifolia.
Buchii, D. Dietr. Syn. Pl. i. 758 = Wahlenbergia pilosa.
caespitosa, *Scop. Fl. Carn.* ed. II. i. 143. t. 4.—Europ.
calamenthifolia, Aucher, ex A. DC. Prod. vii. 462 = tomentosa.
calamenthifolia, *Lam. Encyc.* i. 585.—Ins. Naxos.
calaminthifolia, Gmelin. Spicil. Fl. Rumel. i. 286 = drabaefolia.
calycantha, Turcz. ex Herd. in Act. Hort. Petrop. i. (1873) 288, 296 = pilosa, silenifolia.
calycina, Boeber, ex Roem. & Schult. Syst. v. 104 = C. Rapunculus.
camptoclada, *Boiss. Diagn.* Ser. I. xi. 68.—Syria.
camtschatica, *Pall. ex Roem. & Schult. Syst.* v. 152.—Kamtschat.
cana, *Wall. Cat.* n. 1284; *et in Roxb. Fl. Ind. ed. Carey*, ii. 101.—Reg. Himal.
canaria, *Schrank, ex Steud. Nom.* ed. II. i. 266.—Ins. Canar.
canariensis, Linn. Sp. Pl. 168 = Canarina campanulata.
candida, *A. DC. Monog. Camp.* 234.—Kurdistan.
Candolleana, D. Dietr. Syn. Pl. i. 756 = Prismatocarpus Candolleanus.
canescens, Roth, Enum. Pl. Phan. Germ. i. 216 = Phyteuma canescens.
canescens, *Wall. Cat.* n. 1289.—Ind. or.; Afghan.
capensis, Linn. Sp. Pl. 169 = Wahlenbergia capensis.
caperonioides, Klotzsch, in Bot. Zeqeh. Waldem. Reise, 78 = silvatica.

CAMPANULA:—
capillacea, Thunb. Prod. Pl. Cap. 38 = Wahlenbergia capillacea.
capillaris, Lodd. Bot. Cab. t. 1406 = Wahlenbergia gracilis.
capitata, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = C. Cervicaria.
capitata, Sims, Bot. Mag. t. 811 = lingulata.
cappadocica, Boiss. Diagn. Ser. I. xi. 78 = Phyteuma cappadocica?
carienxis, A. DC. in DC. Prod. vii. 459 = betonicaefolia.
carnica, *Schiede, ex Mert. & Koch, Deutschl. Fl.* ii. 158 —Carniola.
carnosa, Wall. Cat. n. 1282; et in Roxb. Fl. Ind. ed. Carey, ii. 102 = Pentaraxpa carnosa.
carpatica, *Jacq Hort. Vind.* i. 22. t. 57.—Hungar.
cashmiriana, *Royle, Illustr. Bot. Himal.* i. 254. t. 62.—Reg. Himal.
caucasea, Bieb. Casp. 140; App. n. 14 = sibirica?
caudata, *Vis. Fl. Dalm.* ii. 126. t. 33.—Dalmat.
Cavolini, Tenore, Fl. Nap. t. 213 = fragilis.
celsiica, D. Dietr. Syn. Pl. i. 758 = Campanumoea celsiica.
Celsii, A. DC. Monog. Camp. 217 = tomentosa.
cenisia, Linn. Sp. Pl. ed. II. iii. 111 = lactiflora.
cenisia, *Linn. Sp. Pl.* ed. II. Suppl. 1669.—Europ.
cephalantha, Fisch. ex A. DC. Monog. Camp. 254 = glomerata.
Cephalaria, *Vukot. in Linnaea*, xxvi. (1853–55) 333.—Hungar.
Cephalotes, Fisch. ex A. DC. Monog. Camp. 254 = glomerata.
cernua, Ilort. ex Hornem. Hort. Hafn. 201 = sibirica.
cernua, Thunb. Prod. Pl. Cap. 39 = Wahlenbergia cernua.
cerviana, Pall. Reise, iii. 687 = C. Cervicaria.
Cervicaria, *Linn. Sp. Pl.* 167.—Europ.
cervicarioides, Roem. & Schult. Syst. v. 128 = glomerata.
Cervicina, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia Cervicina.
chilensis, Bert. ex Steud. Nom. ed. II. i. 266 = Wahlenbergia hirsuta, reflexa.
chilensis, Molina, Sagg. Chile, ed. II. 231 = Wahlenbergia linarioides.
chondrophylla, D. Dietr. Syn. Pl. i. 758 = Wahlenbergia chondrophylla.
cichoracea, Sibth. & Sm. Fl. Graec t. 209 = lingulata.
cichoriformis, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma cichoriforme.
ciliaris, Salisb. Prod. 127 = Roella ciliata.
ciliata, Patr. ex A. DC. Prod. vii. 479 = silenifolia.
ciliata, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 256. —Reg. Caucas.
ciliata, Thunb. Fl. Cap. ii. 24 = Lightfootia tenella.
cilicia, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma amplexicaule.
cinerea, Linn. f. Suppl. 139 = Lightfootia grisea.
circeoides, *F. Schmidt, Reisen Amurl.* 154.—Japon.
cissophylla, *Boiss. & Hausskn. ex Boiss. Pl. Or. Nov. Dec.* i. 2.—Kurdistan.
citiophylla, *Moretti, ex Reichb. Ic. Fl. Germ.* xix. t. 1611.—Hab.?
civsova, Banks, ex A. DC. Monog. Camp. 161 = Wahlenbergia civsova.
clivsova, Herb. Roxb. ex A. DC. Prod. vii. 486 = Wahlenbergia linifolia.
coa, D. Dietr. Syn. Pl. i. 758 = Specularia pentagonia.
coarctata, Gilib. Fl. Lituan. i. 48 = C. Rapunculus.
coccinea, *Sieber, ex A. DC. Prod.* vii. 486 (quid?).— Ins. Maurit.
cochlearifolia, Lam. Encyc. i. 578 = caespitosa.
cochlearifolia, Vahl, Symb. Bot. 18 = fragilis.
collina, *Bieb. Fl. Taur. Cauc.* i. 153.—Reg. Caucas.
coloradoënsis, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 460 = Specularia Lindheimeri.
colorata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 101.—Ind. or.; Afghan.
commutata, Roem. & Schult. Syst. v. 143 = sarmatica.
compacta, *Boiss. & Heldr. Diagn.* Ser. I. xi. 71.—Pisidia.
conierta, *A. DC. Prod.* vii. 468.—Cappadoc.
confertifolia, Kaut. ex Nym. Consp. 479 = rotundifolia.
congesta, Kocm. & Schult. Syst. v. 125 = glomerata.
conglomerata, Gaudenst. ex Ledeb. Fl. Ross. ii. 880 = glomerata.

CAMPANULA :—
consanguinea, *Schott, Nym. & Kotschy, Analect. Bot.* 8.—Europ.
contracta, Muhl, ex Nym. Consp. 478 = rapunculoides.
controversa, Boiss. Diagn. Ser. II. iii. 115 = Phyteuma rigidum.
cordata, *Peterm. Fl. Lips. Enecurs.* 187.—Saxonia.
cordata, Tausch, in Flora, xxv. (1842) 1. 288 = Adenophora cordata.
cordata, Vis. Stirp. Dalm. 5 = Specularia Speculum.
cordifolia, Coch, in Linnæa, xix. (1847) 29 = rapunculoides.
cordifolia, *Vukot. in Linnæa*, xxvi. (1853) 826.—Europ.
coriacea, Boiss. & Kotschy, ex Fl. Orient. iii. 910 = C. Radula.
coronata, Ker-Gawl. in Bot. Reg. t. 149 = Adenophora marsupiflora.
coronopifolia, Fisch. ex Roem. & Schult. Syst. v. 157 = Adenophora coronopifolia.
corymbifera, Poir. Encyc. Suppl. ii. 64 = sec.
corymbosa, Desf. in Ann. Mus. Par. xi. (1808) 189 = peiviformis.
corymbosa, Tenore, Prod. Fl. Nap. 15 = versicolor.
Costæ, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 294 = patula.
crassifolia, Nees, in Syll. Ratisb. i. (1824) 6 = fragilis.
crassipes, *Heuff. in Oestr. Bot. Zeitschr.* viii. (1858) 27. —Europ.
crenata, Link, Enum. Hort. Berol. i. 214 = rapunculoides.
cretica, *D. Distr. Syn. Fl.* i. 758.—Ins. Creta.
crispa, Banks, ex A. DC. Prod. vii. 444 = Prismatocarpus crispus.
crispa, *Lam. Encyc.* i. 581.—Armenia.
Cymbalaria, *Sibth. & Sm. Fl. Græc. Prod.* i. 139. —As. Min.; Syria.
dalmatica, D. Dietr. Syn. i. 752 = Wahlenbergia dalmatica.
dalmatica, Tausch, in Flora, x. (1827) 246 = C. Cervicaria.
damascena, *Labill. Ic. Pl. Syr. Dec.* v. 7. t. 5.—Syria.
dasyantha, Bieb. Fl. Taur. Cauc. iii. 147 = pilosa.
dasycarpa, Kit. ex Schult. Oestr. Fl. ed. II. 404 = persicifolia.
dasycarpa, Schur, Enum. Pl. Transs. 440 = carpatica.
davurica, *Sieb. in Pall. N. Nord. Beitr.* vii. 218.— Sibir.
debilis, D. Dietr. Syn. Pl. i. 752 = Wahlenbergia debilis.
decipiens, Roem. & Schult. Syst. v. 142 = dichotoma.
declinata, Moench, Meth. Suppl. 187 = americana.
Declextiana, Ortm. in Westenw. Beitr. iii. (1838) 220 = rotundifolia.
decumbens, A. DC. Monog. Camp. 334. t. 11 = Loeflingii.
decurrens, Linn. Sp. Pl. 164 = patula, persicæfolia.
decurrens, Thore, Ess. Chlor. 64 = C. Rapunculus.
decurrens, Zuccagni, in Roem. Collect. 128 = peregrina.
dehiscens, Roxb. Hort. Beng. 85; Fl. Ind. i. 505 = Wahlenbergia gracilis.
delicatula, *Boiss. Diagn.* Ser. I. xi. 67.—As. Min.
delicatula, Jord. ex Nym. Consp. 479 = cæspitosa.
denticulata, Boiss. & Huet, Diagn. Ser. II. iii. 107 = betulæfolia.
denticulata, Burch. Trav. i. 538 = Lightfootia capillaris.
denticulata, Spreng. Syst. i. 785 = Adenophora tricuspidata.
denudata, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia denudata.
desertorum, *Weinm. in Bull. Soc. Nat. Mosc.* (1837) n. VII. 58.—Ruthenia.
dichotoma, *Linn. Cent. Pl.* ii. 10; (*Amœn. Acad.* iv. 306).—Reg. Mediterr. occ.
diffusa, Banks, ex A. DC. Prod. vii. 436 = Wahlenbergia nudicaulis.
diffusa, D. Dietr. Syn. i. 752, 756 = Microcodon lineare.
diffusa, Vahl, Symb. Bot. i. 48 = fragilis.
dilecta, Schott, Nym. & Kotschy, Analect. Bot. 8 = rotundifolia.
dimorphantha, *Schweinf. Beitr. Fl. Aethiop.* 140.— Ægypt.
dissoluta, Schott & Kotschy, ex Boiss. Fl. Orient. iii. 943 = retrorsa.

CAMPANULA :—
divaricata, *Michx. Fl. Bor. Am.* i. 109.—Am. bor
divergens, Willd. Enum. Hort. Berol. i. 212 = sibirica.
diversifolia, D. Dietr. Syn. Pl. i. 762 = Wahlenbergia diversifolia.
diversifolia, Dum. Fl. Belg. 58 = rotundifolia.
Draba, *Bunn. f. Fl. Cap. Prod.* 5.—Afr. austr.
drabæfolia, Frivald. ex Boiss. Fl. Orient. iii. 922 = scutellata.
drabæfolia, *Sibth. & Sm. Fl. Græc.* iii. 11. t. 215.— Græcia.
drabæfolia, Stend. Nom. ed. II. i. 367 = fastigiata.
dracunculifolia, Boiss. & Kotschy, Fl. Orient. iii. 923 = Phyteuma rigidum.
duhsa, A. DC. Monog. Camp. 386 = rotundifolia.
dulcis, *Decne. in Ann. Sc. Nat.* Sér. II. ii. (1884) 258. —Arab.
dumetorum, Boiss. Fl. Orient. iii. 922 = rapunculoides.
Dunantii, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia Dunantii.
Duriæi, Boiss. Voy. Bot. Espagne, 401 = Loeflingii.
echitifolia, Gilib. Fl. Lituan. i. 56 = C. Cervicaria.
Ecklonis, D. Dietr. Syn. Pl. i. 758 = Wahlenbergia Ecklonii.
edulis, *Forsk. Fl. Aegypt. Arab.* 106.—Arab.
elatina, Hill, Veg. Syst. viii. 13 = C. Elatines.
Elatines, Bont. ex Willk. & Lange, Prod. Fl. Hisp. ii. 295 = specularioides.
Elatines, *Linn. Syst.* ed. X. 937.—Europ. austr.
Elatines, Petag. Inst. Bot. ii. 867 = garganica.
Elatines, Pollini, Fl. Veron. i. 378 = elatinoides.
elatinoides, *Moretti, in Brugnat. Giorn. Fis.* II. v. (1822) 110.—Italia.
elatior, Link & Hoffegg. Fl. Port. ii. 11. t. 80 = C. Rapunculus.
elegans, *Roem. & Schult. Syst.* v. 105.—Sibir.
ellipticæ, Kit. in Schult. Oestr. Fl. ed. II. 410 = glomerata.
elongata, Banks, ex A. DC. Prod. vii. 436 = Wahlenbergia nudicaulis.
elongata, Fortensch. ex A. DC. Monog. Camp. 377 = rapunculoides.
elongata, Willd. Enum. Hort. Berol. Suppl. 10 = Wahlenbergia capensis.
emarginata, Lam. Encyc. i. 582 = Heterochasmia ensifolia.
ephesia, *Boiss. Fl. Orient.* iii. 898.—Lydia.
ernantha, Hampe, in Flora, xxv. (1842) i. 76 = tomentosa.
ericoides, Lam. Tabl. Encyc. ii. 63 = Prismatocarpus paniculatus.
erinoides, Cav. in Anal. Cienc. Nat. iii. (1800) 20 = Loeflingii.
erinoides, *Linn. Mant.* i. 44.—Hisp.; Afr. bor.
erinoides, Mühl. Cat. n. 25 = spartioides.
Erinus, *Linn. Sp. Pl.* 169.—Reg. Mediterr.
eriocarpa, Bieb. Fl. Taur. Cauc. i. 149 = latifolia.
erysimoides, Vest, ex Roem. & Schult. Syst. v. 102 = Adenophora Gmelini.
esculenta, *A. Rich. Tent. Fl. Abyss.* ii. 4.—Abyssin.
esculenta, Sallsb. Prod. 126 = C. Rapunculus.
euclasta, *Boiss. Diagn.* Ser. I. xi. 70.—Syria.
evolvulacea, Royle, Illustr. Bot. Himal. i. 254 = cachemiriana.
exalta, *Schleich. ex Murith, Guide Val.* 33, 35.— Helvet.
exilis, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia exilis.
expansa, *Frivald. in Flora*, xix. (1836) ii. 434.— Thessalia; Macedon.
expansa, Rudolph, in Mém. Acad. Pétersb. iv. (1809) 340. t. 2. f. 1 = Wahlenbergia homallanthina.
Exul, *Schott, Nym. & Kotschy, Analect. Bot.* 9.— Austria.
falcata, Roem. & Schult. Syst. v. 154 = Specularia falcata.
fallax, Rupr. in Bull. Acad. Pétersb. xi. (1867) 305, in textu = Aucheri.
farinosa, Andrz. ex Bess. Enum. Pl. Volh. 10 = glomerata.
fasciculata, Linn. f. Suppl. 189 = Lightfootia fasciculata.
fastigiata, *Dufour, ex A. DC. Monog. Camp.* 340. t. 12. —Hispan.; Afr. bor.; Reg. Caucas.
fastigiata, S. G. Gmel. Itin, i. 158. t. 38 = C. Rapunculus.
Feldschekiana, *Transc. in Act Hort. Petrop.* vi. (1879) 77.—As. temp.

CAMPANULA :—
fernandeziana, D. Dietr. Syn. Pl. i. 744 = Wahlenbergia fernandeziana.
fenestrellata, Tineh. in Mém. Acad. Toulouse, Sér. V. vi (1862) 58 = linifolia.
filicaulis, *Dur. Espl. Algér.* t. 62.—Algeria.
filifolia, Kellogg, in Proc. Calif. Acad. ii. (1863) 5 = pensantholdes.
filiformis, Gilib. Fl. Lituan. i. 53 = rotundifolia.
filiformis, Moretti, in Brugnat. Giorn. Fis. II. ix. (1826) 185 = Morettiana.
filiformis, Ruiz & Pav. Fl. Per. ii. 55 = Wahlenbergia linarioides.
firma, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma Aucheri
Firmiana, Vand. Fasc. Pl. 9 = barbata.
Fischeri, Roem. & Schult. Syst. v. 116 = Adenophora communis.
Fischeriana, Spreng. Syst. iv. Cur. Post. 77 = Adenophora coronopifolia.
flaccida, D. Dietr. Syn. Pl. i. 752 = Wahlenbergia flaccida.
flaccida, *Vatke, in Linnæa*, xxxvIII. (1874) 712.— Mesopot.
flagellaris, H. B. & K. Nov. Gen. et Sp. iii. t. 265 = Specularia perfoliata.
flexuosa, Michx. Fl. Bor. Am. i. 109 = divaricata.
flexuosa, Waldst. & Kit. Pl. Rar. Hung. ii. 145 = Waldsteiniana.
floribunda, Viv. Fl. Lib. Spec. 67 = isophylla.
florida, Salisb. Prod. 127 = C. Medium.
floridana, *S. Wats. ex A. Gray, Syn. Fl. N. Am.* ii. 18.—Floridæ.
foliosa, *Tenore, Fl. Nap.* i. 71. t. 18.—Italia.
Fontanesiana, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma lanceolatum.
Foudrasi, Jord. Pugill. Pl. Nov. (1852) 124 = cæspitosa.
fragilis, *Cyrill. Pl. Fasc.* i. 32. t. 11. f. 2.—Italia.
frigida, Pall. ex A. DC. Prod. vii. 461 = pilosa.
Frivaldskyi, Stend. Nom. ed. II. i. 367 = expansa.
fruticosa, Banks, ex A. DC. Prod. vii. 444 = Prismatocarpus fruticosus.
fruticosa, Hill, Syst. Veg. viii. t. 8 = Prismatocarpus paniculatus.
fruticosa, Linn. Sp. Pl. 168 = Lightfootia subulata.
fulgens, *Wall. Cat.* n. 1288; *et in Roxb. Fl. Ind. ed Carey*, ii. 99.—Ind. or.
garganica, *Tenore, Cat. Sem. Hort. Nap.* (1827) 3; *et Fl. Nap.* iii. 208.—Italia; Græcia.
Gautieri, Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat Toul. iii. (1875–76) 621 = Scheuchzeri.
genuinoides, Lam. Encyc. i. 581 = Platycodon grandiflorum.
ghilanensis, Pall. ex Roem. & Schult. Syst. v. 154 = Specularia ghilanensis.
ghilanensis, *Schlecht. ex Boiss. Fl. Orient.* iii. 948.— Persia.
Gieseckiana, Vest, ex Roem. & Schult. Syst. v. 89 = uniflora.
gilanica, Rupr. ex Boiss. Fl. Orient. iii. 905 = Ruprechtii Gisleri x, *Bruegg. in Yahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 104.—Europ.
glabrata, Banks, ex A. DC. Prod. vii. 435 = Wahlenbergia undulata.
glabricarpa, Schleich. Cat. Pl. Helv. ed. IV. 10 = rapunculoides.
glacialis, *Shuttl. in Mag. Zool. & Bot.* ii. (1838) 168.— Helvetia.
glandulosa, Banks, ex A. DC. Monog. Camp. 326 = C. Rapunculus.
glareosa, Huet, ex Nym. Consp. 480 = cæspitosa.
glauca, *Thunb. Fl. Jap.* 88.—Japon.
glaucophylla, Schloss. & Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 111 = glomerata.
glomerata, *Linn. Sp. Pl.* 166.—Europ.; As. temp.
Gmelini, Koch, in Linnæa, viii. (1833) 688 = Adenophora marsupiflora.
Gmelini, Spreng. Fl. Hal. Mant. 36 = Adenophora Gmelini.
gracilis, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma lobelioides.
gracilis, Bert. ex Colla, in Mem. Acc. Tor. xxxvIII. (1835) 119 = Wahlenbergia Bertoroi.
gracilis, Forst. f. Prod. 15 = Wahlenbergia gracilis.
gracilis, Jord. Pugill. Pl. Nov. (1852) 123 = cæspitosa.
graminifolia, Host, Fl. Austr. i. 388 = Wahlenbergia tenuifolia.

CAMPANULA:—
graminifolia, Linn. Sp. Pl. 166 = Wahlenbergia graminifolia.
graminifolia, Waldst. & Kit. Pl. Rar. Hung. ii. 166 = Wahlenbergia Kitaibelii.
graminifolia, Willerm. ex A. DC. Monog. Camp. 254 = glomerata.
grammocarpala, Vukot. in Linnaea, xxvi. (1853–55) 328 = spp. plurimi.
grandiflora, Jacq. Hort. Vindob. iii. 4 = Platycodon grandiflorus.
grandiflora, Lam. Fl. Fr. iii. 334 = C. Medium.
grandiflora, Pourr. ex Nym. Consp. 478 = speciosa.
grandis, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 84 = latiloba.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 23.—Afghan.
Grosseckii, *Herdf. in Flora*, xvi. (1833) i. 358.—Hungar.
gumbetica, *Boiss. Fl. Orient.* iii. 914.—Reg. Caucas.
gummifera, Willd. ex Roem. & Schult. Syst. v. 144 = sarmatica.
Gundelia, C. Koch, in Linnaea, xxiii. (1850) 634 = sibirica.
hagielia, *Boiss. Fl. Orient.* iii. 899.—Ins. Rhodos.
hastaefolia, Salisb. Prod. 127 = Canarina campanulata.
Haussknechtii, Schott, Nym. & Kotschy, Analect. Bot. 10 = rotundifolia.
Haussmanni ×, Reichb. f. in Flora, lx. (1877) 31.
havlodgensis, *Hort.—Cf. Gard. Chron.* (1888) ii. 147.—Hab. ?
hederacea, Linn. Sp. Pl. 169 = Wahlenbergia hederacea.
hederaefolia, Salisb. Prod. 127 = Wahlenbergia hederacea.
hemschinica, *C. Koch, in Linnaea*, xxiii. (1850) 644.—As. Min.
Hendersoni, *Hort.—Cf. Gard. Chron.* (1882) ii. 502.—Hab. ?
Herminii, Bory & Chaub. ex Boiss. Fl. Orient. iii. 937 = Sprunerl.
Herminii, *Hoffmgg. & Link, Fl. Port.* ii. 9. t. 79.—Lusitan.
hesperidifolia, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma lobelioides.
heterodoxa, Bong. Veg. Sitcha, 144 = rotundifolia.
heterodoxa, Vest, in Roem. & Schult. Syst. v. 98 = linifolia.
heterophylla, Baumg. Fl. Transs. Suppl. iii. 342 = Wanneri.
heterophylla, S. F. Gray, Nat. Arr. Brit. Pl. ii. 408 = rotundifolia.
heterophylla, Linn. Sp. Pl. ed. I. errata, et 169 (sine nom. triv.).—Archip. Graec.
hieracioylvaitana, *Boiss. Diagn.* Ser. I. xi. 62.—Palaestin.
Hilsenbergi, Boj. ex A. DC. Prod. vii. 429 = Wahlenbergia Hilsenbergii.
himalayensis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 78 = colorata.
hirsuta, Gmel. Syn. Fl. Sic. i. 250, in syn. = Specularia Speculum.
hirsuta, *Pantoč. in Verh. Ver. Natur- u. Heilk. Presb.* ii. (1871–72) 54.—Europ.
hirsutissima, Guss. ex A. DC. Monog. Camp. 316 = peregrina.
hirta, Roem. & Schult. Syst. v. 153 = Specularia Speculum.
hispanica, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 291.—Hispan.
hispida, Eckl. ex A. DC. Prod. vii. 421 = Microcodon glomeratum.
hispida, Fisch. ex Hornem. Hort. Hafn. 199 = lactiflora.
hispida, Lejeune, Fl. Spa, Suppl. ii. 298 = persicifolia.
hispida, Portensch. ex A. DC. Monog. Camp. 304 = Morettiana.
hispida, Spreng. ex Steud. Nom. ed. II. i. 368 = ciliata.
hispidissima, Hochst. in Lorent, Wanderungen, 331 = propinqua.
hispidula, Eckl. ex A. DC. Monog. Camp. 137 = Microcodon glomeratum.
hispidula, Link, Enum. Hort. Berol. i. 215 = Microcodon sparsiflorum.
hispidula, Linn. f. Suppl. 143 = Microcodon depressum.
Hochstetteri, Schott, Nym. & Kotschy, Analect. Bot. 18 = cespitosa.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 74 = colorata.

CAMPANULA:—
Hohenackeri, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 9 = sibirica.
homallanthina, Ledeb. in Mém. Acad. Pétersb. iv. (1811) 340 = Wahlenbergia homallanthina.
Hornungiana, *Schur, Enum. Pl. Transs.* 442.—Transsylv.
hortensis, Meerb. Pl. Sel. 10 = rapunculoides.
Hostii, Baumg. Enum. Stirp. Transs. iii. 342 = rotundifolia.
humilis, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia humilis.
humilis, Schur, Enum. Pl. Transs. 438 = persicifolia.
humillima, *A. DC. Prod.* vii. 474.—Persia.
hungarica, Pers. Syn. i. 191 = lingulata.
hybrida, Bont. ex Willk. & Lange, Prod. Fl. Hisp. ii. 295 = Loeflingii.
hybrida, Linn. Sp. Pl. 168 = Specularia hybrida.
hybrida, Rodigas.—Cf. Illustr. Hortic. xxx. (1883) 118. t. 499 = Van-Houttei.
hygrophila, Rupr. in Bull. Acad. Pétersb. xi. (1867) 209 = Aucheri.
hypopolia, *Trautv. in Act. Hort. Petrop.* iv. (1876) 389.—As. temp.
hyrcana, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, i. (1888) 69.—Persia.
illinoisana, Trelease, Fresen. Ind. Sem. Hort. Franc. (1836) = americana.
imbricata, Hayne, ex A. DC. Monog. Camp. 284 = carpatica.
imeretina, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 217.—Reg. Caucas.
incanescens, *Boiss. Diagn.* Ser. I. vii. 16.—Persia.
incanescens, Schott, Nym. & Kotschy, Analect. Bot. 10 = Scheuchzeri.
inconspicua, Banks, ex A. DC. Monog. Camp. 157 = Wahlenbergia lobelioides.
incurva, Aucher, ex A. DC. Prod. vii. 464 = Leutweinii.
indica, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia indica.
infundibuliformis, Sims, Bot. Mag. t. 2682 = rapunculoides.
Infundibulum, Reichb. Ic. Pl. Crit. i. t. 75. f. 158 = allenifolia.
Infundibulum, Vest, ex Roem. & Schult. Syst. v. 106 = Steveni.
integerrima, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 153 = sylvatica.
intermedia, Engelm. ex Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 256 = bifora.
intermedia, Roem. & Schult. Syst. v. 110 = Adenophora communis.
interrupta, Pers. Syn. i. 192 = Prismatocarpus interruptus.
Intybus, Schott & Kotschy, ex Tchihat, Asie Min. Bot. ii. 402 = trachyphylla.
involucrata, *Aucher, ex A. DC. Prod.* vii. 467.—As. Min.; Persia.
isophylla, *Moretti, in Brugnat. Giorn. Fis.* ii. vii. (1824) 44, 98 (*App. Schemb. Fl. Ital.* 22).—Italia.
Jacobaea, *C. Sm. ex Hook. Niger Fl.* 148. t. 12.—Ins. Cap. Virid.
Jacquinii, A. DC. Monog. Camp. 251 = Trachelium Jacquinii.
japonica, Vatke, in Linnaea, xxxviii. (1874) 705 = Phyteuma japonicum.
jasionefolia, A. DC. in DC. Prod. viii. 468 = stricta.
Jaubertiana, *Timb. in Bull. Soc. Bot. Fr.* xv. (1868) p. xlviii.—Gallia.
javanica, D. Dietr. Syn. Pl. i. 759 = Campanumoea javanica.
juncea, D. Dietr. l. c. 757 = Lightfootia juncea.
juncea, *Hill, Veg. Syst.* viii. 14.—German.
juncea, Hort. ex A. DC. Monog. Camp. 157 = Wahlenbergia lobelioides.
juncea, *Wettstein, in Stapf, in Denkschr. Acad. Wien*, i. (1885) 117.—Lycia.
juncifolia, Hort. ex A. DC. Monog. Camp. 157 = Wahlenbergia lobelioides.
jurjurensis, *Pomel, Nouv. Mat. Fl. Atl.* 257.—Afr. bor.
kamtschatica, Pall. ex Ledeb. Fl. Ross. ii. 889 = camtschatica.
kashmiriana, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 24 = cashmiriana.
khasiana, *Hook. f. & Thoms. l. c.* 25.—Reg. Himal.
Kitaibelliana, *Roem. & Schult. Syst.* v. 90.—Banatus.
kolenatiana, *C. A. Mey. ex Rupr. in Bull. Acad. Pétersb.* xi. (1867) 216.—Reg. Caucas.
Kotschyana, *A. DC. Prod.* vii. 481.—Cilicia.

CAMPANULA:—
Krokeri, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia Zeyheri.
Kremeri, Boiss. & Reut. Pugill Pl Nov. 75 = dichotoma
kryophila, Rupr. in Bull. Acad. Pétersb. xi. (1867) 211 = arvicensis.
laciniata, Andr. Bot. Rep. t. 385 = tomentosa.
lacinitata, *Linn. Sp. Pl.* 165.—Archip. Graec.
lactiflora, *Bieb. Fl. Taur. Cauc.* i. 153.—Reg. Caucas.
Lamarckii, D. Dietr. Syn. Pl. i. 755 = Adenophora Lamarkii.
Lamborliana, A. DC. Monog. Camp 327 = C. Rapunculus.
lamiifolia, Bieb. Fl. Taur. Cauc. i 154 = alliariaefolia.
lanata, *Fryvald. in Flora*, xix. (1836) 484.—Thracia.
lanceolata, Boiss. & Kotschy, ex Willk. & Lange, Prod. Fl. Hisp. ii. 291 = hispanica.
lanceolata, *Lapeyr. Hist. Abr. Pl. Pyr.* 105.—Mont. Pyren.
lanceolata, Pall. ex A. DC. Monog. Camp. 321 = Steveni.
lanceolata, J. & C. Presl, Fl. Cech. 49 = persicifolia.
lanceolata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = rotundifolia.
lancifolia, Koxb. Fl. Ind. ed. Carey, ii. 96 = Campanumoea celebica.
lancifolia, Schur, Enum. Pl. Transs. 445 = rotundifolia.
Langsdorffiana, Fisch. ex A. DC. Monog. Camp. 279 = linifolia.
Langsdorffiana, A. Gray, in Am. Journ. Sc. Ser. II. xxxiv. (1862) 254 = Parryi.
laniflora, *Hornem. ex Steud. Nom.* ed. II. i. 368.—Hab. ?
lanuginosa, Hort. ex Steud. l. c. = divergens.
lanuginosa, Lam. Encyc. i. 584 = peregrina.
lanuginosa, Willd. Enum. Hort. Berol. 213 = rupestris.
Larreinii, Bert. ex Colla, in Mem. Acc. Tor. xxxviii. (1835) 118. t. 35 = Wahlenbergia fernandezlana.
lasiocarpa, *Cham. in Linnaea*, iv. (1829) 89.—Am. bor. occ.
latifolia, *Linn. Sp. Pl.* 165.—Europ.; Oriens; As. temp.
latiloba, *A. DC. Prod.* vii. 478.—Bithynia.
lanandulaefolia, Reinw. ex Blume, Bijdr. 720 = Wahlenbergia gracilis.
Ledebouriana, *Trautv. in Act. Hort. Petrop.* ii. (1873) 477.—Armenia.
Ledebourii, Boiss. Fl. Orient. iii. 905 = pracc.
Lehmanniana, *Bunge, in Lehm. Reliq. Hot.* 211.—Turkestan.
leptocarpa, Engelm. ex A. Gray, in Proc. Am. Acad. xi. (1876) 82 = Specularia leptocarpa.
leptophylla, Ehrenb. ex Boiss. Fl. Orient. iii. 952 = Phyteuma rigidum.
leucanthemifolia, Pourr. in Mém. Acad. Toul. iii. (1788) 309 = caespitosa.
leucocladia, *Boiss. Diagn.* Ser. II. iii. 109.—Afghan.
leucosiphon, *Boiss. & Heldr. Diagn.* Ser. I. xi. 94.—As. Min.
Leutweinii, *Heldr. in Ann. Sc. Nat. Sér.* IV. xiii. (1860) 270.—Cilicia.
libanotica, *A. DC. Prod.* vii. 463.—Syria.
ligularia, *Lam. Encyc.* i. 585.—Europ.
ligulata, Pers. Syn. i. 191 = lingulata.
ligulata, Steud. Nom. ed. II. i. 365 = C. Cervicaria.
lilifiora, Roth, Enum. Pl. Phan. Germ. i. 715 = Adenophora communis
lilifolia, Lam. Encyc. i. 581 = Adenophora Lamarkii.
lilifolia, Linn. Sp. Pl. 165 = Adenophora communis.
limonia, Murr. Syst. Veg. ed. XIII. 176 = seq.
limonifolia, Linn. Sp. Pl. ed. II. 239 = Phyteuma limonifolium.
linarioides, Lam. Encyc. i. 580 = Wahlenbergia linarioides.
linarifolia, D. Dietr. Syn. Pl. i. 756 = Prismatocarpus campanuloides.
linearis, Linn. f. Suppl. 140 = Microcodon lineare.
lingulata, Reichb. Ic. Pl. Crit. vi. 13 = C. Cervicaria.
lingulata, *Waldst. & Kit. Pl. Rar. Hung.* i. 65. t. 64.—Europ. austr. or.
linifolia, *Scop. Ann.* ii. *Hist. Nat.* 47.—Europ.
linifolia, Schrank, Prim. Fl. Sal. 70 = rotundifolia.
linifolia, Pall. ex A. DC. Monog. Camp. 303 = Adenophora marsupiifolia.
linnaeifolia, *A. Gray, in Proc. Am. Acad.* vii. (1868) 366.—Calif.
linoides, Presl, ex A. DC. Monog. Camp. 158 = Wahlenbergia linarioides.

CAMPANULA :—
littoralis, Labill. Nov. Holl. Pl. i. 49. t. 70 = Wahlenbergia gracilis.
lobata, *Sieblas. & Vukot., in Rad Jugos. Akad. Zagreb.* xxxi. (1875) 105.—Europ.
lobelioides, Linn. f. Suppl. 140 = Wahlenbergia lobelioides.
lobelioides, Spreng. Syst. i. 736 = Wahlenbergia undabunda.
lobelioides, Vatke, in Linnæa, xxxvlii. (1874) 704 = Phyteuma lobelioides.
Loeflingii, *Brot. Fl. Lusit.* i. 287; *Phyt. Lusit.* i. 41. t. 18.—Lusitan. ; Marocco.
longebracteata, D. Dietr. Syn. Pl. i. 757 = Trichelia longebracteata.
longifolia, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
longipes, Coss. ex Nym. Consp. 485 = Loeflingii.
longirostris, Banks, ex A. DC. Monog. Camp. 168 = Prismatocarpus crispus.
Loreyi, A. Blanco, ex Nym. Consp. 269 = Loeflingii.
Loreyi, *Pollini, Elem. Bot.* ii. 149 (Lorel).—Europ. austr.
Lostritti, *Tenore, Syll.* 96.—Italia.
lourica, *Bouss. Diagn.* Ser. I. xi. 70.—Persia.
ludmiciana, Torr. ex A. Gray, Syn. Fl. N. Am. ii. i. 11 = Specularia biflora.
lunarnifolia, Reichb. Ic. Fl. Germ. vi. 750 = rapunculoides.
lunariæfolia, Willd. ex Roem. & Schult. Syst. v. 92 = Phyteuma campanuloides.
lusitanica, Linn. in Loefl. It. Hisp. 111, 126 = Loeflingii.
Lychnitis, Hornem. Hort. Hafn. i. 199 = bononiensis.
lycia, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma lycium.
lyræfolia, Saliab. Prod. 127 = Michauxia campanuloides.
lyrata, Bourg. ex Nym. Consp. 476 = hagielia.
lyrata, *Lam. Encyc.* i. 588.—Aa. Min. ; Reg. Caucas.
macedonica, Boiss. Fl. Orient. iii. 923 = scutellata.
macedonica, Boiss. & Orph. ex Boiss. l. c. 925 = multiflora.
macrantha, Fisch. ex Hornem. Hort. Hafn. Suppl. 23 = latifolia.
macrocalyx, Schur, Enum. Pl. Transs. 441 = pulla.
macrocephala, Willd. ex Steud. Nom. ed. II. i. 268 = Wahlenbergia tenuifolia.
macrochlamys, *Boiss. & Huet, Diagn.* Ser. II. iii. 111. —Armenia.
macrophylla, Sims, Bot. Mag. t. 912 = alliariæfolia.
macrorhiza, Vukot. in Linnæa, xxvi. (1853–55) 329 = spp. plurim.
macrorrhiza, *Y. Gay, ex A. DC. Monog. Camp.* 302.—Gallia.
macrostachya, Panz. ex Roem. & Schult. Syst. v. 121 = rapunculoides.
macrostachya, Willd. Enum. Hort. Berol. i. 213 = multiflora.
macrostyla, *Boiss. & Heldr. Diagn.* Ser. I. xi. 65.— Phrygia.
madagascariensis, D. Dietr. Syn. Pl. i. 752 = Wahlenbergia oppositifolia.
Malyi, Schott, Nym. & Kotschy, Analect. Bot. 11 = rotundifolia.
marginata, Thunb. Fl. Jap. 89 = Wahlenbergia gracilis.
maroccana, *Ball, in Journ. Linn. Soc.* xvi. (1878) 584.—Morocco.
Marschalliana, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma campanuloides.
marsupiflora, Fisch. ex Roem. & Schult. Syst. v. 116 = Adenophora marsupiflora.
Massonii, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia Massonii.
Mathonnti, Jord. ex Gren. & Godr. Fl. Fr. ii. 418 = cæspitosa.
matritensis, A. DC. Monog. Camp. 332 = Loeflingii.
mauritanica, *Pomel, Nouv. Mat. Pl. Alt.* 8.—Afr. bor.
Medium, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa. Medium, *Linn. Sp. Pl.* 167.—Europ. austr.
Meyeriana, Rupr. in Bull. Acad. Pétersb. xi. (1867) 502 = Auctori.
michauxioides, *Boiss. Diagn.* Ser. I. iv. 86.—Caria.
micrantha, Schur, Enum. Pl. Transs. 438 = bononiensis.
Microcodon, D. Dietr. Syn. Pl. i. 757 = Microcodon glomeratum.
microphylla, Cav. in Anal. Clenc. iii. (1800) 19 = mollis.
microphylla, Kit. ex Schult. Oestr. Fl. ed. II. 400 = Kitaibeliana.
Minæ, *F. G. Strobl, in Flora*, lxvi. (1883) 549.—Sicil.
CAMPANULA :—
minor, [Honck.] Vers. aller Gew. Teutschl. 617 = cæspitosa.
minor, Lam. Fl. Fr. iii. 389 = rotundifolia.
minuta, Agardh, ex Roem. & Schult. Syst. v. 87 = pusilla.
minuta, Savi, Cent. ii. Fl. Etrusc. 54 = rotundifolia.
minutissima, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 47.—Transs.
mixta, *Hegetschw. Fl. Schw.* 231.—Helvet.
modesta, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 34.—Reg. Himal.
modesta, Schott, Nym. & Kotschy, Analect. Bot. 13 = cæspitosa.
mollis, *Linn. Sp. Pl.* ed. II. 237.—Reg. Mediterr. occ.
monadelpha, Pall. ex A. DC. Monog. Camp. 868 = Adenophora coronopifolia.
monantha, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 47 = Scheuchzeri.
monanthos, Pantot. in Verh. Ver. Natur- u. Heilk. Presb. ii. (1871–72) 54 = præc. ?
monocephala, *Trautv. in Act. Hort. Petrop.* vi. (1879) 54.—Turkestan.
montana, *Dulard. Fl. Auvergne*, ed. II. 40.—Gallia.
montevidensis, Spreng. Syst. i. 738 = Specularia biflora ?
Moorcroftiana, Wall. Cat. n. 1288 = colorata.
Moretti ana, *Reichb. Ic. Fl. Germ.* iv. 18.—Dalmat.
morifolia, Salisb. Prod. 126 = rapunculoides.
multicaulis, Boiss. Diagn. Ser. I. vi. 19 (= *Podanthum persicum*).—Persia.
multiflora, Hort. ex A. DC. Monog. Camp. 271 = bononiensis.
multiflora, *Waldst. & Kit. Pl. Rar. Hung.* iii. 292.— Hungar.
multiflora, Willd. ex Roem. & Schult. Syst. v. 130 = polyantha.
muralis, Portenschl. ex A. DC. Monog. Camp. 308 = Portenschlagiana.
musarum, Heldr. ex Nym. Consp. 477 = rupicola.
myosotidifolia, Boiss. in Tchihat. Asie Min. Bot. ii. 392 = trojanica.
muscosa, [*Pieri*,] *in Ionios Anthol.* v. 204.—Ins. Corcyra.
myrtifolia, Boiss. & Heldr. Diagn. Ser. I. xi. 69 = Trachelium myrtifolium.
nana, Lam. Encyc. i. 585 = Allionii.
napuligera, Schur, Enum. Pl. Transs. 444 = rotundifolia.
neglecta, Bess. Cat. Hort. Cremen. (1816) 28 = rapunculoides.
neglecta, Bowmar, in Bull. Soc. Nat. Mosc. xxxiv. (1861) 1, 182 = præc. ?
neglecta, Roem. & Schult. Syst. v. 104 = patula.
nemorosa, A. DC. Monog. Camp. 271 = rapunculoides.
nervosa, Royle, Illustr. Bot. Himal. 253 = colorata.
nicaeensis, Risso, ex A. DC. Monog. Camp. 302 = macrorrhiza.
nicaeensis, Roem. & Schult. Syst. v. 126 = glomerata.
nitida, [Soland. in] Ait. Hort. Kew. i. 246 = planiflora.
nobili-macrantha, *Regel, Ind. Sem. Hort. Petrop.* (1869) .... Fish. ?
nobilis, Lindl. in Journ. Hort. Soc. i. (1846) 222 = punctata.
notata, Schott, Nym. & Kotschy, Analect. Bot. 13 = cæspitosa.
nudicaulis, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia nudicaulis.
numidica, *Jour. Exptl. Algér.* t. 62.—Afr. bor.
nutabunda, Guss. in Tenore, Append. v. ad Fl. Nap. Prod. 8 = Wahlenbergia nutabunda.
nutans, Lam. Fl. Fr. ii. 236 = rapunculoides.
nutans, Sieber, ex A. DC. Monog. Camp. 366 = Symphyandra cretica.
nutans, Vahl, ex Hornem. Hort. Hafn. 201 = sibirica.
Nuttalli, *Penny, ex Loud. Hort. Brit. Suppl.* i. 558.— Am. bor.
obliqua, Jacq. Hort. Schoenb. iii. 47 = americana.
obliquifolia, Tenore, Fl. Nap. Prod. 18 = bononiensis.
Olivieriana, *Boiss. Diagn.* Ser. I. xi. 66.—Persia.
omnipontana, Moretti, ex A. DC. Monog. Camp. 269 = rapunculoides.
oligocantha, Schur, Enum. Pl. Transs. 436 = C. Trachelium.
Olivieri, A. DC. Monog. Camp. 283. t. 9 = calamenthifolia.
olympica, *Boiss. Diagn.* Ser. I. iv. 84.—Aa. Min.
orbelica, *Panč. Elem.* 48.—Europ.
CAMPANULA :—
oreadum, *Boiss. & Heldr. Diagn.* Ser. II. iii. 107.— Thessalia.
orophila, Schur, Enum. Pl. Transs. 441 = carpatica.
orientalis, Steud. Nom. ed. II. i. 269 = peregrina ?
Orphanidea, *Boiss. Fl. Orient.* iii. 897.—Macedon.
ossetica, Bieb. Fl. Taur. Cauc. iii. 145 = Symphyandra ossetica.
Ottoniana, Roem. & Schult. Syst. v. 116 = Lightfootia oxyococcoides.
ovata, Spreng. Syst. iv. Cur. Post. 78 = Wahlenbergia ovata.
oxyphylla, Vukot. in Linnæa, xxvi. (1853–55) 331 = spp. plurim.
pacnina, *Reut. ex Tissière*, Bot. Gr. St. Bernard, 83.— Helvet.
Pallasiana, Roem. & Schult. Syst. v. 138 = pilosa.
pallida, Wall. in Aa. Research. xiii. (1820) 375 = colorata.
paniculata, Linn. f. Suppl. 139 = Wahlenbergia paniculata.
paniculata, Pohl, ex A. DC. Monog. Camp. 244 = sibirica.
paniculata, Sm. ex A. DC. l. c. 154 = Wahlenbergia Banksiana.
paniculata, *Turra, ex Sacc. in Atti Ist. Venet.* Ser. III ix. (1863–64) 486.—Italia.
paniculata, Wall. Cat. n. 1295 A, B = Cephalostigma paniculatum.
pannonica, Bernh. ex Schrank, in Denkschr. Regensb. Ges. ii. (1822) 34 = sibirica.
parnassica, *Boiss. & Spruu. Diagn.* Ser. I. vii. 17.— Græcia.
Parryi, *A. Gray, in Syn. Fl. N. Am.* ii. i. 895.—Am. bor. occ.
parviflora, Lam. Encyc. i. 588 = sibirica.
parviflora, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = C. Erinus.
parviflora, Saliab. Prod. 126 = Wahlenbergia lobelioides.
parvula, Jord. Pugill. Pl. Nov. (1852) 128 = cæspitosa.
patens, Gueldenst. ex Ledeb. Fl. Ross. ii. 887 = patula.
patula, *Linn. Sp. Pl.* 168.—Europ.
patula, Sibth. & Sm. Fl. Græc. Prod. i. 127 = Sprunerlana.
pauciflora, Desf. Choix Pl. 36. t. 26 = Sprunerlana.
paucifloru, Lam. ex Steud. Nom. ed. II. i. 269 = *planiflora*, Lam.
pedunculuris. Wall. Cat. n. 7139 = Wahlenbergia pedunculuria.
pedunculata, J. F. Gmel. Syst. 852 = Wahlenbergia paniculata.
petrescifolia, Spreng. Syst. i. 735 = Adenophora latifolia.
peliviformis, *Lam. Encyc.* i. 586.—Ins. Creta.
pendula, Bieb. Fl. Taur. Cauc. i. 154 = Symphyandra pendula.
pennica, Reut. ex Payot, in Bull. Soc. Bot. Fr. xxiii. (1883).—Cf. Bibl. 43 = seq.
pennina, Reut. in Comptes-Rend. Soc. Hallér. (1854–56) 115 = rotundifolia.
Pentagonia, Linn. Sp. Pl. 169 = Specularia Pentagonia.
pentagonophylla, Vukot. in Linnæa, xxvi. (1853–55) 325 = Wahlenbergia hederacea.
peregrina, Hoffmagg. & Link, Fl. Port. ii. 15. t. 83 = primulæfolia.
peregrina, *Linn. Mant.* ii. 204.—Aa. Min. ; Syria.
Pereskia, Fisch. ex A. DC. Monog. Camp. 357 = Adenophora latifolia.
pereskiæfolia, Fisch. ex Roem. & Schult. Syst. v. 116 = Adenophora latifolia.
perfoliata, Linn. Sp. Pl. 169 = Specularia perfoliata.
perfoliatifolia, Linn. Amœn. i. 280 = Adenophora perfoliatifolia.
perosphecta, *Schott, Nym. & Kotschy, Analect. Bot.* 11.—Austria.
Perpenline, Moretti, ex A. DC. Monog. Camp. 305 = Kaiserli.
perpusilla, *A. DC. Prod.* vii. 474.—Persia.
persepolitana, *Kotschy, ex Boiss. Diagn.* Ser. I. vii. 19. —Persia.
persica, A. DC. Prod. vii. 463 (= *Podanthum persicum*). —Persia.
persicifolia, *Linn. Sp. Pl.* 164.—Europ. ; Aa. bor.
Pestalozzæ, Boiss. Diagn. Ser. I. xi. 69 = propinqua.
petiolata, A. DC. Monog. Camp. 278 = rotundifolia.
petræa, All. Fl. Pedem. i. 113 = glomerata.

CAMPANULA:—
petraea, Habl. Taur. 157 = bononiensis.
petraea, Kotschy, ex A. DC. Prod. vii. 467 = involucrata.
petræa, *Linn. Syst.* ed. X. 937.—Italia.
petræa, Zantedeschi, ex Moretti, in Syll. Ratisb. i. (1824) 154 = stationidea.
petrophila, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 212.—Reg. Caucas.
phrygia, *Yaub. & Spach, Illustr. Pl. Or.* iii. 42.—Macedon.; As. Min.
phyctidocalyx, *Boiss. & Noë, Diagn.* Ser. II. iii. 114.—Armenia.
phyteumoides, Sibth. ex Zuccagni, in Roem. Collect. 128 = Phyteuma limonifolium.
Pichleri, Vis. in Mem. Ist. Venet. xvi. (1871) 106 = Phyteuma amplexicaule.
pilosa, Less. ex Ledeb. Fl. Ross. ii. 894 = Adenophora tricuspidata.
pilosa, *Pall. ex Roem. & Schult. Syst.* v. 148.—Sibir.; Am. bor. occ.
pinifolia, Vukot. in Rad Jugos. Akad. Zagreb. lvii. (1881) 92 = patula.
planiflora, Engelm. in Coult. Bot. Gaz. vii. (1882) 5 = Parryi.
planiflora, Lam. Encyc. i. 580 = pyramidalis?
planiflora, Willd. Enum. Hort. Berol. 210 = versicolor.
plicata, Pers. Syn. i. 192 = Prismatocarpus crispus.
plicatula, Dum. Fl. Belg. 58 = C. Trachelium.
podanthoides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 938.—Catsonia.
podocarpa, *Boiss. Diagn.* Ser. I. xi. 68.—Pamphylia.
polyantha, *Roem. & Schult. Syst.* v. 130.—Sibiria.
polymorpha, Banks & Soland. ex A. DC. Monog. Camp. 142 = Wahlenbergia gracilis.
pomponiifolia, Fisch. ex A. DC. l. c. 363 = Adenophora Gmelini.
porua, Thunb. Prod. Pl. Cap. 39 = Salmolus Valerandi.
Portenschlagiana, *Roem. & Schult. Syst.* v. 93.—Dalmatia.
portensis, Linn. in Loefl. It. Hisp. 111, 127 = C. Erinus.
Pourretii, *Yeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* vi. (1888–89) 158.—Gallia.
pratensis, Lapyl. ex A. DC. Monog. Camp. 287 = rotundifolia.
pratensis, Reut. ex Nym. Consp. 478 = C. Trachelium.
precatoria, Timb. in Mém. Acad. Sc. Toul. Sér. VII. v. (1878) 271 = lanceolata.
Preissii, De Vriese, in Lehm. Pl. Preiss. ii. 241 = Wahlenbergia gracilis.
prenanthoides, *Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 88.—Calif.
primulæfolia, *Brot. Fl. Lusit.* i. 288.—Lusitan.
Prismatocarpus, L'Hérit. Sertum, t. 3 = Prismatocarpus sibthn.
procumbens, Linn. f. Suppl. 141 = Wahlenbergia procumbens.
propinqua, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1836) 32.—As. Min.; Persia.
prolnosa, [*Pierr*] *in Jonios Anthol.* v. 204.—Ins. Corcyra.
prostrata, Dulac, Fl. Hautes-Pyr. 457 = Scheuchzeri.
pseudo-carpatica, Schur, Enum. Pl. Transs. 441 = reniformis.
pseudolanceolata, *Pantoč. in Magyar Növent. Lapok*, vi. (1882) 103.—Hungar.
pseudo-pulla, Schur, Enum. Pl. Transs. 441 = pulla.
pseudo-saldensis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = rotundifolia.
pulloatachya, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 118.—Cilicia.
pumicæfolia, *Lam. Encyc.* i. 570.—As. Min.
pumonea, Reichb. ex Nym. Consp. 479 = rotundifolia.
pubescens, Schmidt, Fl. Boëm. Cent. ii. 67 = caespitosa.
pubiflora, Rupr. in Bull. Acad. Pétersb. xi. 207 = Ancheri.
pulchella, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma pulchellum.
pulchella, Jord. Fugill. Nov. Pl. (1852) 196 = caespitosa.
pulchella, Sellsb. Prod. 127 = Specularia Speculum.
pulcherrima, *Schrenk & Zeyh. ex Steud. Nom.* ed. II. i. 369.—Stab.?
pulla, Baumg. ex Steud. Nom. ed. II. i. 268, 269 = linifolia.
pulla, *Guldenst. It.* ii. 24; *ex Ledeb. Fl. Ross.* ii. 891.—Reg. Caucas.
pulla, *Linn. Sp. Pl.* 168.—Europ.
pulla, Parol. ex Follini, Fl. Ver. i. 268 = Moretiana.
pumila, Curt. Bot. Mag. t. 512 = caespitosa.
pumila, Schmidt, Fl. Boem. ii. 71 = persicifolia.
Pumilio, Portenschl. ex Roem. & Schult. Syst. v. 136 = Wahlenbergia Pumilio.
punctata, *Lam. Encyc.* i. 586.—Sibir.; Japon.
pundwana, D. Dietr. Syn. Pl. i. 757 = Campanumoea punctata.
purpurea, Spreng. Syst. iv. Cur. Post. 78 = Codonopsis purpurea.
pusilla, Haenke, in Jacq. Collect. ii. 79 = caespitosa.
pygmæa, D. Dietr. Sp. Pl. i. 757 = Microcodon glomeratum.
pygmæa, Lam. ex DC. Fl. Fr. iii. 705 = Borago laxiflora.
pyramidalis, Gilib. Fl. Lituan. i. 52 = bononiensis.
pyramidalis, *Linn. Sp. Pl.* 164.—Europ.
pyramidata, Gilib. Anat. Pl. Eur. ii. 209 = pracc.
pyramidiflora, Reichb. Cat. Sem. Hort. Dresd. (1822) = rapunculoides.
pyrenaica, *A. DC. Monog. Camp.* 324.—Mont. Pyren.
pyrenaica, 'Hechtl. Wildf.' ex Steud. Nom. ed. II. i. 269, 270 = rhomboides.
quadrifida, R. Br. Prod. 561 = Wahlenbergia gracilis.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* ii. 6.—Abyss.
Rahelaiana, Roem. & Schult. Syst. v. 158 = Adenophora Gmelini.
racemosa, G. Emel. It. i. 9 = sibirica.
racemosa, Opiz, ex Steud. Nom. ed. II. i. 269 = rapunculoides.
racemosa, Vukot. in Linnaea, xxvi. (1851) 332 = spp. plurim.
Raddeana, *Trautv. in Bull. Acad. Pétersb.* x. (1866) 895.—Reg. Caucas.
radicosa, *Bory & Chaub. Fl. Péiop.* 14. t. 7.—Graecia.
Radula, *Fisch. ex Tchihat. Asie Min. Bot.* ii. 895.—Kurdistan.
Rainerii, *Perpenf. in Bibl. Ital.* v. (1817) 184.—Italia.
ramosissima, Griseb. Spicil. Fl. Rumel. ii. 282 = phrygia.
ramosissima, Host, Fl. Austr. i. 264 = Loreyi.
ramosissima, *Sibth. & Sm. Fl. Graec. Prod.* i. 137.—Graecia.
ramossissima, Sprun. ex Nym. Consp. 481 = Sprunneriana.
ramosissima, Willd. ex Steud. Nom. ed. II. i. 269 = Loeflingii.
ramulosa, Wall. Cat. n. 1286; in Roxb. Fl. Ind. ed. Carey. ii. 100 = colorata.
ranunculoides, Uspensky, in Bull. Soc. Nat. Mosc. vii. (1834) 369 = rapunculoides.
rapunculoides, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = C. Rapunculus.
rapunculoides, *Linn. Sp. Pl.* 165.—Europ.; As. temp.
rapunculoides, Pall. ex Steud. Nom. ed. II. i. 270 = bononiensis.
Rapunculus, *Linn. Sp. Pl.* 164.—Europ.; Orient.; Orient.; Afr. bor.; As. bor.
Rapunculus, O. F. Muell. Fl. Dan. t. 855 = rotundifolia.
Ravaji, *Boiss. Diagn.* Ser. I. iv. 82.—Caria.
Rebeudiana, Gren. & Godr. Fl. Fr. ii. 415 = rotundifolia.
Rebeudiana, *Pomel, Nouv. Mat. Fl. Atl.* 2.—Algeria.
recta, Dulac, Fl. Hautes-Pyr. 458 = linifolia.
Redowskiana, Cham. & Schlecht. in Linnaea, iv. (1829) 41 = Wahlenbergia hemiathantha.
Redowskyi, Fisch. ex Herd. in Act. Hort. Petrop. i. (1873) 288 = pilosa.
redux, Schott, Nym. & Kotschy, Analect. Bot. 9 = Schanschieri.
reflexa, Hausm. ex Nym. Consp. 480 = carnica.
reflexa, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = caespitosa.
remotiflora, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 180.—Japon.
reniformis, *Schur, Enum. Pl. Transs.* 441.—Transylv.
repanda, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma limonifolium.
repens, Host. ex Steud. Nom. ed. II. i. 270 = Wahlenbergia procumbens.
repens, *Lour. Fl. Cochinch.* 139.—Cochinch.
retrorsa, *Labill. Ic. Pl. Syr. Dec.* v. 5.—Syria; Cilicia.
Reverchoni, *A. Gray, Syn. Fl. N. Am.* ii. 1. 396.—Texas.
Reuteriana, *Boiss. & Bal. Diagn.* Ser. II. iii. 108.—As. Min.; Persia.
Reyneri, D. Dietr. Syn. Pl. i. 740 = Raineri.
rhodensis, A. DC. Monog. Camp. 297 = drabæfolia.
Rhodii, Loisel. Fl. Gall. ed. II. i. 140 = linifolia.
rhomboidalis, Auct. ex Steud. Nom. ed. II. i. 270 = lanceolata.
rhomboidalis, Gorter, Fl. Ingr. 34 = rapunculoides.
rhomboidalis, *Linn. Sp. Pl.* 165.—Europ.
rhomboidalis, Schur, Enum. Pl. Transs. 448 = arenata.
rhomboides, Endr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 291 = lanceolata.
rhomboides, *Murr. Syst.* ed. XIII. 173.—Europ.
rigescens, Pall. ex Roem. & Schult. Syst. v. 102 = sileniifolia.
rigida, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma rigidum.
rigida, Gilib. Fl. Lituan. i. 51 = rapunculoides.
rigidipila, *Steud. & Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 8.—Abyss.
rimarum, *Boiss. Fl. Orient.* iii. 931.—Lycia.
riparia, Boj. ex A. DC. Prod. vii. 432 = Wahlenbergia emirnensis.
riparia, Leprieur & Perr. ex A. DC. l. c. 435 = Wahlenbergia ripens.
Rochelii, Schur, in Oestr. Bot. Zeitschr. (1861) 45 = Stevenii.
rotiloides, Juss. ex A. DC. Monog. Camp. 165 = Prismatocarpus paniculatus.
Rœsii, Kegel, Gartenfl. (1872) 289; et in Act. Hort. Petrop. ii. (1872) 310 = prenanthoides.
Rohelii, Loisel. Fl. Gall. ed. II. i. 140. t. 24 = linifolia.
Rosani, Tenore, Fl. Nap. iii. 205. t. 120 = versicolor.
rosea, *Noronha, in Verh. Batav. Gen. v.* (1790) ed. I. Art. iv. 32.—Malaya.
rosulata, Vukot. in Linnaea, xxvi. (1853-55) 326 = cenisia.
rotunda, Gilib. Fl. Lituan. i. 46 = rotundifolia.
rotundifolia, All. ex Steud. Nom. ed. II. i. 270 = rhomboides.
rotundifolia, Boiss. Voy. Espagne, 309 = macrorrhiza.
rotundifolia, Ledeb. Fl. Alt. i. 288 = linifolia.
rotundifolia, *Linn. Sp. Pl.* 168.—Reg. bor. temp.
rotundifolia, Pall. ex A. DC. in DC. Prod. vii. 484 = adscendens.
rubioides, Banks, ex A. DC. Monog. Camp. 187 = Microcodon linearc.
rubra, Hort. journ. & Fl. des Jard. (1832) 120. t. 120 = rhomboidalis.
ruderalis, *Asteh. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 174.—Afghan.
rumeliana, *Vatke, in Linnaea*, xxxviii. (1874) 705.—Rumel.
rupestris, Adams, ex Ledeb. Fl. Ross. ii. 876 = bellidifolia.
rupestris, Bieb. Fl. Taur. Cauc. i. 154 = Bieberteiniana.
rupestris, Host, Fl. Austr. i. 263 = Waldsteiniana.
rupestris, Kaso, ex A. DC. Monog. Camp. 302 = macrorrhiza.
rupestris, Sibth. & Sm. Fl. Graec. Prod. i. 142 = tomentosa.
rupestris, Sprun. ex Nym. Consp. 477 = rupicola.
rupicola, *Boiss. & Sprun. Diagn.* Ser. I. vii. 17.—Graecia.
Ruprechtii, *Boiss. & Bal. Diagn.* iii. 905.—Persia.
ruscinonensis, Timb. in Mém. Acad. Sc. Toul. Sér. VII. v. (1878) 276 = linifolia.
Russelliana, Roem. & Schult. Syst. v. 142 = strigosa.
ruthenica, Bieb. Fl. Taur. Cauc. i. 151 = bononiensis.
sabatia, *De Not. Prosp. Fl. Ligust.* 52.—Italia.
sajanensis, [Fisch.] Hort. Gorenk. ed. I. (1808) 52 = Adenophora marsupiflora.
sajanensis, Spreng. ex Steud. Nom. ed. II. i. 270 = lasiocarpa.
salicifolia, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma canescens.
salicifolia, Boiss. & Kotschy, ex Boiss. Fl. Orient. iii. 949 = Phyteuma amplexicaule.
salicifolia, Juss. ex Ledeb. Fl. Ross. ii. 894 = Adenophora coronopifolia.
sanensis, Fisch. Cat. Hort. Gorenk. (1808), ex A. DC. in Prod. vii. 491 = pilosa.
sarmatica, *Ker-Gawl. in Bot. Reg.* t. 237.—Reg. Caucas.

CAMPANULA:—
sarmentosa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 4. —Abyssin.
sarmentosa, Fenzl & Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 105 = lobata.
Sartorii, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iii. 919. —Archip. Graec.
saxatilis, Georgi, Beschr. Russ. Reich. iii. iv. 776 (ap. dub.).—Russia.
saxatilis, *Linn. Sp. Pl.* 167.—Ins. Cretæ.
saxicola, R. Br. Prod. 561 = Wahlenbergia saxicola.
Saxifraga, *Biebst. Fl. Taur. Cauc.* i. 165.—Reg. Caucas.
scabrella, *Engelm. in Coult. Bot. Gazette*, vi. (1881) 237.—Am. bor. occ.
scabrida, *Hochst. in Lorent, Wanderungen*, 331.—Mesopotam.
scandens, *Pall. ex Roem. & Schult. Syst.* v. 130.—Sibr.
Scheuchzeri, A. Gray, Syn. Fl. N. Am. ii. i. 12 = Parryi.
Scheuchzeri, Lodd. Bot. Cab. t. 485 = rotundifolia.
Scheuchzeri, *Vill. Prosp.* 22.—Mont. Pyren.
Schimperi, Vatke, in Linnæa, xxxviii. (1874) 712 = rigidipila.
Schlicherei, Suter, Fl. Helv. i. 124 = linifolia.
Schotti, *Hort.—Cf. Gard. Chron.* (1886) ii. 147.—Hab.?
sclerotricha, *Boiss. Diagn.* Ser. I. xi. 66.—Persia.
Sconleri, *Hook. ex A. DC. Monog. Camp.* 319.—Am. bor. occ.
scutellata, *Griseb. Spicil. Fl. Rumel.* ii. 282.—Macedon.
secunda, Schmidt, Fl. Boem. ii. 74 = rapunculoides.
secundiflora, Vis. & Panč. in Mem. Ist. Venet. x. (1861) 442 = rapunculoides.
seminuda, Vest, in Roem. & Schult. Syst. v. 91 = Steveni.
serpyllifolia, *Vis. Fl. Dalm.* ii. 136. t. 15.—Dalmatia.
serratifolia, Vukot. in Linnæa, xxvi. (1853–55) 335 = C. Trachelium, latifolium.
sessiliflora, *Boiss. Fl. Orient.* iii. 947.—Armenia.
sessiliflora, D. Dietr. Syn. Pl. i. 756 = Prismatocarpus campanuloides.
sessiliflora, C. Koch, in Linnæa, xix. (1847) 80 = latifolia.
sessilifolia, Linn. f. Suppl. 139 = Lightfootia longifolia, subulata.
sessilifora, Vukot. in Linnæa, xxvi. (1853–55) 337 = Raineri.
sessilis, D. Dietr. Syn. Pl. i. 756 = Prismatocarpus sessilis.
setacea, D. Dietr. l. c. 754 = Wahlenbergia capillacea.
setosa, Fisch. ex DC. Monog. Camp. 369 = rapunculoides.
setosa, *Wendl. Beobacht.* 42.—Hab.?
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 188.—As. temp.
sibirica, *Linn. Sp. Pl.* 167.—Europ.; As. temp.
Sibthorpiana, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma Sibthorpianum.
sidonicensis, *Boiss. & Blanche, Diagn.* Ser. II. iii. 114.—Syria.
Sieberi, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia gracilis.
silenifolia, *Fisch. ex A. DC. Monog. Camp.* 320.—As. bor.
silenifolia, Host, Fl. Austr. i. 368 = Wahlenbergia Pumilio.
simplex, Lam. ex DC. Fl. Fr. iii. 730 = bononiensis.
simplex, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 255 = Steveni.
Sinai, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma lanceolatum.
sinensis, D. Dietr. Syn. Pl. i. 755 = Adenophora sinensis.
singarensis, *Boiss. & Hausskn. ex Boiss. Pl. Or. Nov. Dec.* i. 2.—Mesopotam.
sparsiflora, D. Dietr. Syn. Pl. i. 757 = Microcodon sparsiflorum.
spathulæfolia, Sprun. ex Nym. Consp. 488 = ramosissima.
spathulata, Ehrenb. ex Boiss. Fl. Orient. iii. 918 = trichopoda.
spathulata, Sibth. & Sm. Fl. Græc. Prod. i. 137 = Spruneriana.
spathulata, Waldst. & Kit. Pl. Rar. Hung. ii. 206. t. 186 = stellaris.
speciosa, Gilib. Fl. Lituan. i. 50 = persicifolia.

CAMPANULA:—
speciosa, Hornem. Hort. Hafn. ii. 957, Add. = glomerata.
speciosa, *Pourr. in Mém. Acad. Sc. Toul.* iii. (1788) 309.—Mont. Pyren.
speciosa, Willd. ex Spreng. Syst. i. 729 = elegans.
spectaticoides, *Cosa. Notes Crit.* 41.—Hispan.
Speculum, Linn. Sp. Pl. 166 = Specularia Speculum.
spherothrix, Griseb. Spicil. Fl. Rumel. ii. 280 = expansa.
spicata, *Linn. Sp. Pl.* 166.—Europ.
spicata, Geners. Elench. Scepus. n. 192; ex Wahlenb. Fl. Carp. 62 = C. Cervicaria.
spinosa, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia spinulosa.
spreta, Roem. & Schult. Syst. v. 128 = Adenophora communis.
Sprunerl, Boiss. Fl. Orient. iii. 927 = seq.
Spruneriana, *Hampe, in Flora*, xxv. (1842) i. 76.—Græcia et Ins.
spuria, Pall. ex Roem. & Schult. Syst. v. 154 = Specularia hybrida.
St.-Helenæ, D. Dietr. Syn. Pl. i. 755 = Wahlenbergia linifolia.
stellarioides, D. Dietr. l. c. 758 = Wahlenbergia stellarioides.
stellaris, *Boiss. Diagn.* Ser. I. xi. 68.—Syria; Palestin.
stellata, *Lam. & Lamb. in Redoute. Phyt. Biastr.* i. (1808) 20.—Afr. austr.
Stelleri, Steph. ex Herd. in Act. Hort. Petrop. i. (1871) 299 = lasiocarpa.
stenanthina, Ledeb. in Mém. Acad. Pétersb. v. (1814) 525 = Adenophora marsupiflora.
stenocarpa, *Traute. & Mey. in Middend. Reise (Fl. Ochot.* 51).—As. temp.
stenocodon, Boiss. & Reut. Diagn. Ser. II. iii. 112 = rotundifolia.
stenophylla, Boiss. & Heldr. l. c. Ser. I. xi. 77 = Phyteuma linifolium.
stenosiphon, Boiss. & Heldr. l. c. vii. 18 = glomerata.
stenosiphon, Reichb. f. ex Nym. Consp. 479 = rotundifolia.
Steveni, *Bieb. Fl. Taur. Cauc.* iii. 138.—Reg. Caucas.; Persia.
Steverana, Roem. & Schult. Syst. v. 91 = prænc.
stoloniifera, *Migu. in Bull. Soc. Bot. Fr.* xii. (1865) 242.—Hispan.
stricta, Labill. In. Pl. Syr. Dec. v. 6 = libanotica.
stricta, *Linn. Sp. Pl.* ed. II. 238.—As. Min.
stricta, Sibth. & Sm. Fl. Græc. Prod. i. 140 = lyrata.
strictiflora, Wall. in As. Research. xiii. (1820) 374 = sylvatica.
strigillosa, *Boiss. in Ann. Sc. Nat.* Ser. IV. ii. (1854) 251.—Cappadocia.
strigosa, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 246.—Cilicia; Syria.
strigosa, *Vahl, Symb. Bot.* iii. 34.—As. Min.
stylaticoides, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma lobelioides.
stylosa, Bess. Cat. Hort. Crem. (1816) = Adenophora communis.
stylosa, Lam. Encyc. i. 580 = Adenophora stylosa.
styriaca, Schott, Nym. & Kotschy, Analect. Bot. 9 = Scheuchzeri.
suaveolens, *Bieb. Fl. Taur. Cauc.* iii. (1807) 215.—Reg. Caucas.
suaveolens, Gilib. Fl. Lituan. i. 50 = Adenophora communis.
saavecolens, *Wender. ex Steud. Nom.* ed. II. i. 370.—Hab.?
subalpina, *Delponte & Gras, Ind. Sem. Hort. Taurin.* (1872) 25.—Hab.?
subpyrenaica, *Timb. in Mém. Acad. Toul. Sér.* IV. v. (1855) 387.—Hispan.
subramulosa, Jord. in Gren. & Godr. Fl. Fr. ii. 418 = conglobata.
subulata, Bess. ex A. DC. Prod. vii. 479 = americana.
subulata, Sprong. Syst. i. 809 = Lightfootia subulata.
subulata, Thunb. Prod. Pl. Cap. 38 = Prismatocarpus subulatus.
subuniflora, Lam. Tabl. Encyc. ii. 38 = linifolia.
sulphurea, *Boiss. Diagn.* Ser. I. xi. 64.—Syria.
swellendamensis, D. Dietr. Syn. Pl. i. 758 = Wahlenbergia swellendamensis.
sylvatica, *Wall. Cat.* n. 1293; *et in Roxb. Fl. Ind.* ii. 97.—Reg. Himal.

CAMPANULA:—
syngenesiflora, Pall. ex A. DC. Monog. Camp. 363 = Adenophora marsupiflora.
syriaca, Ehrenb. ex Boiss. Fl. Orient. iii. 928 = stellaris.
syriaca, Willd. ex Roem. & Schult. Syst. v. 133 = Specularia falcata.
syspirensis, *C. Koch, in Linnæa*, xxiii. (1850) 639.—Armenia.
tauricola, Boiss. & Bal. Diagn. Ser. II. iii. 116 = Phyteuma rigidum.
telephioides, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. Dec.* i. 1.—Catasonia.
tenella, Jord. Pugill. Pl. Nov. (1852) 125 = cæspitosa.
tenella, Lange, ex Nym. Consp. 479 = hispanica.
tenella, Linn. f. Suppl. 141 = Lightfootia tenella, Thunbergiana.
tenerrima, D. Dietr. Syn. Pl. i. 757 = Prismatocarpus tenerrimus.
Tenorii, Moretti, Prosp. Geogr. Bot. Ital. Schouw, 19 = vernicolor.
tenuiflora, Schur, Enum. Pl. Transs. 438 = bononiensis.
tenuiflora, Tenore, Fl. Nap. iii. 207. t. 210 = linguata.
tenuifolia, Hoffm. Fl. Germ. i. 100 = rotundifolia.
tenuifolia, Waldst. & Kit. Pl. Rar. Hung. ii. 168 = Wahlenbergia tenuifolia.
tetraphylla, Thunb. Pl. Jap. 87 = Adenophora verticillata.
tauricoides, *Boiss. Diagn.* Ser. I. iv. 88.—Lydia.
Teymonsii, *Hort.—Cf. Gard. Chron.* (1885) ii. 147, 182.—Hab.?
Thaliana, Wallr. Sched. Crit. 86 = bononiensis.
thalictrifiora, Sprengel, Syst. iv. Cur. Post. 77 = Codonopsis thalictrifolia.
Thomasii, Tenore, ex A. DC. Monog. Camp. 309 = vernicolor.
Thomsonii, Hook. f. in Journ. Linn. Soc. ii. (1858) 25 = Phyteuma Thomsoni.
Thunbergii, Roem. & Schult. Syst. v. 135 = Lightfootia ciliata.
thyrsoides, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
thyrsoides, *Linn. Sp. Pl.* 167.—Europ.
tomentosa, *Lam. Encyc.* i. 584.—As. Min.
tomentosa, *Vent. Hort. Cels*, 18.—Græcia et Ins.
Tommasiniana, Koch, ex F. Schultz, Arch. Fl. Fr. et Allem. 239; Reut. in Cat. Genev. (1865) 4 = Waldsteiniana.
trachelioides, Bieb. Fl. Taur. Cauc. i. 150 = rapunculoides.
trachelioides, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 285.—Afr. bor.
Trachelium, Host. Fl. Austr. i. 327 = primulæfolia.
Trachelium, Bell. Herb. Fr. t. 319 = rapunculoides.
Trachelium, *Linn. Sp. Pl.* 166.—Europ.; As. temp.; Afr. bor.
Trachelium, Thunb. Fl. Jap. 88 = punctata.
trachyphylla, *Schott & Kotschy, ex Boiss. Fl. Orient* iii. 926.—Cilicia.
transalvanica, *Schur, ex Andrae, in Bot. Zeit.* xiii. (1855) 938.—Transylv.
trichocalycina, Tenore, Fl. Nap. i. 67. t. 16 = Physparnea amplexicaule.
trichopoda, *Boiss. Diagn.* Ser. I. xi. 68.—Syria.
tricuspidata, Ehrenb. ex Roem. & Schult. Syst. v. 158 = Adenophora denticulata.
tridens, Rapr. in Bull. Acad. Pétersb. xi. (1867) 204 = tridentata.
tridentata, *Schrenk. Is.* 3. t. 2; *Linn. Mant.* i. 44.—As. Min.
trigona, Ehrenb. ex Boiss. Fl. Orient. iii. 959 = Specularia Speculum.
trilocularis, Turra, Prod. Fl. Ital. 64 = Allionii.
triphylla, Thunb. Fl. Jap. 87 = Adenophora triphylla.
truncata, D. Dietr. Syn. Pl. i. 757 = Campanumœa celebica.
tubiflora, Tausch, ex Ledeb. Fl. Ross. ii. 880 = glomerata.
tubulosa, *Linn. Encyc.* i. 586.—Ins. Cretæ.
turbinata, Schott, Nym. & Kotschy, Analect. Bot. 14 = carpatica.
turczaninowii, Schott, Nym. & Kotschy, l. c. 13 = cæspitosa.
turczaninowii, Bess. Cat. Hort. Cremen. (1816) 26 = rapunculoides.
umbelliflora, Vukot. in Linnæa, xxvi. (1853–55) 333 = pyramidalis.

CAMPANULA:—
umbrosa, F. G. Dietr. Vollst. Lex. Gaertn. i. 675 = Adenophora communis.
undulata, Linn. f. Suppl. 142 = Wahlenbergia undulata.
undulata, Moench, Meth. Suppl. 189 = albirica.
unidentata, Linn. f. Suppl. 139 = Lightfootia unidentata.
uniflora, Georgi, Beschr. Russ. Reich. iii. iv. 772 = silenifolia.
uniflora, Gorter, Fl. Ingr. 83 = rotundifolia.
uniflora, Honck. Syn. Pl. Germ. ii. 251 = cenisia.
unifloras, *Linn. Sp. Pl.* 168.—Reg. Arctic.
uniflora, Schult. Oestr. Fl. ed. II. 400 = caespitosa.
uniflora, Vill. Prosp. 32 = linifolia.
urticaefolia, All. Fl. Pedem. i. 110 = latifolia.
urticaefolia, Gilib. Fl. Lithuan. i. 52 = bononiensis.
urticaefolia, Turra, Prod. Fl. Ital. 64 = rapunculoides.
urticifolia, Schmidt, Fl. Boem. ii. 78 = C. Trachelium.
waldensis, All. Fl. Pedem. i. 109 = linifolia.
Van-Houttei, *Carr. in Rev. Hortic.* (1878) 430.—Hab.! Vandesi, *G. Den, ex Loud. Hort. Brit.* 77.—Hab.!
varrifolia, Salisb. Prod. 128 = rotundifolia.
Vayredae, *Lessecke & Lesser, in Journ. Bot.* xvii. (1879) 139.—Hispan.
velata, *Pomel, Nouv. Mat. Fl. Atl.* 2.—Algeria.
velebica, *Borb.—Cf. in Oestr. Bot. Zeitschr.* xxxiii. (1883) 201.—Austria.
velutina, Desf. Fl. Atlant. i. 180. t. 51 = mollis.
venosa, Willd. Enum. Hort. Berol. Suppl. 10 = rhomboidalis.
venusta, Schur, Enum. Pl. Transs. 442 = caespitosa.
verbenaefolia, *Sieber, in Ann. Fl. Pom.* (1838) 375 — Hab.!
veronicifolia, *Hance, in Journ. Bot. etc.* (1871) 138.—China.
verrucifolia, Hoffmgg. & Link, Fl. Port. ii. 12. t. 81 = C. Rapunculus.
versicolor, *Sibth. & Sm. Fl. Graec. Prod.* i. 138.—Italia : Graecia.
verticillata, Gmel. Fl. Rar. 92 = versicolor.
verticillata, *Hill, Veg. Syst.* viii. 10.—Am. bor.
verticillata, Pall. Reise, iii. 719 = Adenophora verticillata.
vesula, All. Fl. Pedem. i. 108 = persicifolia.
Vidalii, *H. C. Watt. in Hook. Ic. Pl.* t. 684.—Ins. Azores.
vincaeflora, Vent. Jard. Malm. 12 = Wahlenbergia gracilis.
Violae, Pers. Syn. I. 192 = seq.
violaefolia, *Lam. Encyc.* i. 587.—Sibiria.
virgata, A. DC. Prod. vii. 483 = C. Rapunculus.
virgata, Labill. Ic. Pl. Syr. Dec. ii. 11. t. 6 = Phyteuma limonifolium.
virgata, Rafin. ex Roem. & Schult. Syst. v. 100 = divaricata, linifolia.
viridis, Spreng. Syst. iv. Cur. Post. 78 = Codonopsis viridis.
Vitrophoana, Roem. & Schult. Syst. v. 102 = Steveni, persicifolia.
Vlachovae, Orphan. ex Boiss. Fl. Orient. iii. 927 = glomerata.
volubilis, Willd. ex Roem. & Schult. Syst. v. 98 = lactiflora.
vulgaris, *Gueldenst. It.* ii. 10, *ex Ledeb. Fl. Ross.* ii. 891.—Reg. Caucas.
Waldsteiniana, *Roem. & Schult. Syst.* v. 99.—Croatia.
Wanneri, Rochel, Pl. Banat. Rar. 41. t. 5. f. 12 = Symphyandra Wanneri.
Welandii, Heuff. in Oestr. Bot. Wochenschr. (1857) 118 = expansa.
Willdenowiana, Roem. & Schult. Syst. v. 107 = versicolor.
Willdenowii, Boiss. Diagn. Ser. II. iii. 117 = Phyteuma lobelioides.
Zeyheri, D. Distr. Syn. Pl. i. 753 = Wahlenbergia Zeyheri.
Zoysii, *Wulf. in Jacq. Collect.* ii. 122.—Europ. austr.

CAMPANULOIDES, Hort. Kew. ex A. DC. Monog. Camp. 107 (1886) = Lightfootia, L'Hérit. (Campan.),
subulata, Hort. Kew. ex A. DC. l. c. = L. subulata.

CAMPANULOPSIS, Zoll. & Mor. in Nat. en Genees. Arch. Nederl. Indië, i. (1844) 454, nomen. *CAMPANULACEAE*
cyanea, *Zoll. & Mor. l. c.*—Malaya.

CAMPANUMOEA, Blume, Bijdr. 726 (1825).
CAMPANULACEAE, Benth. & Hook. f. ii. 557.
CYCLOCODON, Griff. Notul. iv. 277 (1854).
celebica, *Blume, l. c.* 727.—Ind. or.; Malaya; China.
cordata, Maxim. in Bull. Acad. Pétersb. xii. (1868) 68 = javanica.
Infra, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 436. —Reg. Himal.
japonica, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 67.—Japon
japonica, Siebold, ex Morr. Belg. Hortic. (1862) 337 = lanceolata ?
javanica, *Blume, Bijdr.* 726.—Reg. Himal. ; Java.
lanceolata, Sieb. & Zucc. Fl. Japon. i. 174. t. 91. = Codonopsis lanceolata.
leucocarpa, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 436 = celebica.
parviflora, *Benth. & Hook. f. Gen.* ii. 558.—Reg. Himal.
pilosula, *Franch. Pl. Davidian.* i. (1884) 192.—Mantchur.
truncata, Endl. Gen. Plant. i. 515 ; C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 436 = celebica.

CAMPBELLIA, Wight, Ic. iv. 5. t. 1424, 1425 (1850) = **Christisonia**, Gardn. (Orobanch.).
alboides, Benth. & Hook. f. Gen. ii. 967 = Christisonia albicida.
aurantiaca ?, Wight, Ic. iv. t. 1424 = Chr. neilgherrica.
cytinoides, Wight, l. c. 1425 = Chr. neilgherrica.
Lawii, Benth. & Hook. f. Gen. ii. 967 = Chr. Lawii.
subacaulis, Benth. & Hook. f. l. c. = Chr. subacaulis.

CAMPDERIA, Benth. Bot. Voy. Sulph. 159. t. 52. (1844). *POLYGONACEAE*, Benth. & Hook. f. iii. 102. floribunda, *Benth. l. c.* 160.—Guatemala.
gracilis, *Meisn. in Mart. Fl. Bras.* v. i. 23.—Bras. Lindeniana, *Benth. & Hook. f. Gen.* iii. 102.—Mexic.
mexicana, Meisn. in DC. Prod. xiv. 171 = floribunda.
nematostachya, *Griseb. in Bonplandia*, vi. (1858) 4.—Panama.

CAMPDERIA, Lag. Amen. Nat. iii. 99 (1821) = **Kundmannia**, Scop. (Umbellif.).
acuta, Lag. l. c. = K. sicula.

CAMPDERIA, A. Rich. in Bull. Soc. Philom. (1822) 79 = **Barbaeonia**, Vand. (Amaryllid.).

CAMPE, Dulac, Fl. Hautes-Pyr. 199 (1867) = **Barbarea**, R. Br. (Crucifer.).
praecox, Dulac, l. c. = B. praecox.
vulgaris, Dulac, l. c. = B. vulgaris.

CAMPECARPUS, H. Wendl. ex Benth. & Hook. f. Gen. iii. 898 (1883) = **Cyphophoenix**, H. Wendl. (Palm.).
fulcitus, H. Wendl. l. c. = Cyph. fulcita.

CAMPECIA, Adans. Fam. ii. 567 (1763) = **Cassalpinia**, Linn. (Legumin.).

CAMPELEPIS, Falc. in Trans. Linn. Soc. xix. (1845) 101 = **Periploca**, Linn. (Asclepiad.).
viminea, Falc. l. c. = P. aphylla.

CAMPELIA, Kunth, Enum. Pl. i. 286, 287 (1833) = CAMPELLA, Link.

CAMPELLA, Rich. Anal. Fruit. 46 (1808). *COMMELINACEAE*, Benth. & Hook. f. iii. 855.
GONATANDRA, Schlecht. in Linnaea, xxiv. (1851) 659.
SARSIA, Rafin. Fl. Tellur. ii. 16 (1836).
ZANONIA, [Flum.] Cram. Diss. Syst. 75 (1808).
bibracteata, Wied-Neuwied, Beitr. Bras. 15. t. 2. f. 4 = Zanonia.
Boucheana, Schult. f. Syst. vii. 1150 = C. Zanonia.
fastigiata, Schlecht. in Linnaea, xxv. (1852) 188 = C. Zanonia.
Fendleri, Hasak. in Flora, xlviii. (1865) 102 = C. Zanonia.
glabrata, Hasak. in Miq. Pl. Jungh. 154 = Forrestia glabrata.
glabrata, Kunth, Enum. Pl. iv. 109 = C. Zanonia.
Hofmanni, Hasak. in Flora, xlviii. (1865) 105 = C. Zanonia.
marginata, Blume, Enum. Pl. Jav. i. 7 = Forrestia marginata.
marginata, Wall. Cat. n. 8977 = Forrestia Hookeri.

CAMPELIA:—
mexicana, Mart. ex Kunth, Enum. Pl. iv. 109 = C. Zanonia.
mollissima, Blume, Enum. Pl. Jav. i. 7 = Forrestia mollis.
Pseudo-Zanonia, Kunth, Enum. Pl. iv. 108 = C. Zanonia.
scandens, *Hassk. in Flora*, xlviii. (1865) 103.—Peruv.
Zanonia, *H. B. & K. Nov. Gen. et Sp.* i. 264.—Am. austr.

CAMPELLA, Link, Hort. Berol. i. 122 (1827) = **Daschampsia**, Beauv. (Gramin.).
caespitosa, Link, l. c. 122 = D. caespitosa.
media, Link, l. c. 123 = D. juncea.

CAMPHORA, [Bauh.] Linn. Syst. ed. I (1735) ; Nees, Syst. Laurin. 37 = **Cinnamomum**, Tourn. (Laurin.).
angustifolia, Rafin. Sylva Tellur. 136.—Hab.!
chinensis, Nees, in Otto, Gartenz. (1833) 91 ; Syst. Laurin. 92 = Cinn. Parthenoxylon.
glandulifera, Nees, in Wall. Pl. As. Rar. ii. 72 = Cinn. glanduliferum.
glauca, G. Don, in Sweet, Hort. Brit. ed. III. 582 = Litsea glauca.
inodora, Blume, ex Miq. Fl. Ind. Bat. i. 904.—Arch. Sunda.
inuncta, Nees, in Wall. Pl. As. Rar. iii. 82 = Cinn. inunctum.
japonica, Rafin. Sylva Tellur. 136 = Cinn. pedunculatum ?
officinalis, Steud. Nom. ed. II. i. 271 = Cinn. Camphora.
officinarum, Nees, ex Wall. Pl. As. Rar. ii. 72 = Cinn. Camphora.
Parthenoxylon, Nees, l. c. = Cinn. Parthenoxylon.
porrecta, Voigt, Hort. Suburb. Calc. 808 = Cinn. Parthenoxylon.
Pseudo-Sassafras, Miq. in Zoll. Syst. Verz. Ind. Archip. 113, 114 = Cinn. Parthenoxylon.
sumatrana, Miq. Fl. Jungh. 174.—Sumatra.
vera, Rafin. Sylva Tellur. 136 = Cinn. Camphora.

CAMPHORATA, Tourn. ex Crantz, Inst. i. 118 (1766) = **Camphoromaea**, Linn. (Chenopod.).
acuta, Crantz, l. c. 114 = Halimocnemis juniperina.
glabra, Crantz, l. c. = Camphoromaea glabrum.
hirsuta, Moench, Meth. 700 = Camphoromaea monspeliacum.
monspeliensium, Crantz, Inst. i. 114 = Camphoromaea monspeliacum.

CAMPHORINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1 = **Cinnamomum**, Tourn. ? (Laurin.).

CAMPHOROMOEA, Nees, Laurin. Expos. 16 (1833) = **Cocotes**, Aubl. (Laurin.).
divaricata, Nees, Syst. Laurin. 467.— Bras.
laxa, Nees, l. c. 468.—Bras.
litseaefolia, Meisn. in DC. Prod, xv. i. 145.—Bras.
ovalifolia, Meisn. l. c. 144.—Bras.
rhamnoides, Meisn. l. c. 145.—Bras.
subtriplinervia, Nees, Syst. Laurin. 466.—Bras.
swinamensis, Meisn. in DC. Prod. xv. i. 145.—Guinea.
tenuiflora, Meisn. l. c. 144.—Bras.
ovmulosa, Nees, Syst. Laurin. 469.—Bras.
staphyloides, Meisn. in DC. Prod. xv. i. 145.—Bras.

CAMPHOROMYRTUS, Schau. in Linnaea, xvii. (1843) 240 = **Basaekia**, Linn. (Myrtac.).
Behrii, Schlecht. l. c. xx. (1847) 651 = B. Behrii.
Brownii, Schau. l. c. xvii. (1843) 240 = B. camphorata.
crenulata, F. Muell. in Trans. Vict. Inst. i. (1855) 138 = B. virgata.
pluriflora, F. Muell. l. c. = B. virgata.

CAMPHOROSMA, Linn. Diss. Dass. 8 (1747) (Amoen. Acad. i. 392). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 56.
CAMPHORATA, Tourn. ex Crantz, Inst. i. 118 (1766).
acutum, Falk, Beitr. ii. 199 = Nanophyton caspicum.
acutum, Linn. Sp. Pl. 322 = Halimocnemis juniperina.
acutum, Pollich, Fl. Palat. iii. 817 = Kochia arenaria.
annua, *Pall. Illustr. Pl.* 67. t. 58.—Rossia.
glabrum, Auct. ex DC. Prod. iii. 368 = Siphonychia americana.
glabrum, *Linn. Sp. Pl.* 323.—Helvetia.
hirsutum, Stokes, Bot. Mat. Med. i. 232 = monspeliacum.
monandrum, *Bunge, ex Boiss. Fl. Orient.* iv. 922.—Afghan.

CAMPHOROSMA :—
monspeliacum, *Linn. Sp. Pl.* 122.—Reg. Mediterr.
monspeliacum, Pall. Reise, ii. 471 = ruthenicum.
monspeliacum, Pollich, Pl. Palat. i. 165 = Kochia arenaria.
monspeliense, Murr. Syst. Veg. ed. XIII. 139 = monspeliacum.
ovatum, Hasel. ex Moq. in DC. Prod. xiii. ii. 126 = monspeliacum.
ovatum, *Waldst. & Kit. Pl. Rar. Hung.* i. 64. t. 63.—Europ. austr. or.
ovatum, Bung. & Mey. Suppl. ii. Fl. Alt. 68. p. 274 = annuum.
palmarum, Linn. f. Suppl. 128 = Monetia paleacea.
perennis, *Pall. Illustr. Pl.* iii. 66. t. 57.—Reg. Caucas.; Persia.
polygamum, *Bunge, ex Boiss. Fl. Orient.* iv. 921.—Afghan.
Pteranthus, Linn. Mant. i. 41 = Pteranthus echinatus.
ruthenicum, *Bieb. Fl. Taur. Cauc.* iii. *Suppl.* 112.—Europ. austr. or.; As. bor.
sabulosa, *Kit. ex Moq. in DC. Prod.* xiii. ii. 126.—Hungar
songoricum, *Bunge, in Act. Hort. Petrop.* vi. (1879) 415.—Sibir.
vingintimillieis, *Tineo, Cat. Hort. Panorm.* 277.—Sicil.

CAMPHUSIA, De Vilese, in Nederl. Kruidk. Arch. ii. (1850) 148 = **Soaevola**, Linn. (Goodeniac.).
glabra, De Vriese, l. c. 149.—Malaya.

CAMPHYLEIA, Spreng. Gen. ii. 486 (1831) = **Striga**, Lour. (Scrophulac.).

CAMPIA, 'Domb.' ex Endl. Ench. 91, in syn. (1841) = CAPIA, Juss. = **Lagenaria**, Ruiz & Pav. (Liliac.).

CAMPILOSTACHYS, A. Juss. in Orb. Dict. xii. 88 (1849) = **Campylostachys**, Kunth (Verbenac.).

CAMPNOSPERMA, Thw. in Hook. Kew Journ. vi. (1854) 65. t. 1. *ANACARDIACEAE*, Benth. & Hook f. i. 426.
auriculata, *Hook. f. Fl. Brit. Ind.* ii. 41.—Ind. or.
Griffithii, *March. Rev. Anac.* 174.—Penang.
gummifera, *March. l. c.* 178.—Bras.
macrophylla, *Hook. f. Fl. Brit. Ind.* ii. 41.—Malacca. Micrantela, *March. Rev. Anac.* 179.—Madag.
oxyrhachis, *Engl. in A. DC. Monog. Phan.* iv. 319.—Sumatra.
seychellarum, *March. Rev. Anac.* 173.—Ins. Mascar.
zeylanica, *Thw. in Hook. Kew Journ.* vi. (1854) 65.—Zeylan.

CAMPOMANESIA, Ruiz & Pav. Prod. 72. t. 13 (1797). *MYRTACEAE*, Benth. & Hook. f. i. 712.
ABREVILLEA, ACRANDRA, BRITOA, LACERDAEA, Berg, in Linnaea, xxvii. (1854) 425–437.
adamantium, *Blume, Mus. Bot. Lugd. Bat.* i. 73.—Bras.
affinis, *Berg, in Mart. Fl. Bras.* xiv. i. 446.—Bras.
aprica, *Berg, l. c.* 459.—Bras.
arenaria, *Berg, l. c.* 448.—Reg. Argent.
aromatica, *Griseb. Fl. Brit. W. Ind.* 242.—Ind. occ.
aurea, *Berg, in Mart. Fl. Bras.* xiv. i. 454.—Bras.
australis, *Berg, l. c.* 445.—Bras.
caerulea, *Berg, l. c.* 455.—Bras.
caerulescens, *Berg, l. c.* 612.—Bras.
Cambessedeana, *Berg, l. c.* 457.—Bras.
campestris, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
cerasoides, *A. Gray, Bot. U. St. Expl. Exped.* i. 549.—Bras.
ciliata, *Berg, in Mart. Fl. Bras.* xiv. i. 458.—Bras.
cosetanea, *Berg, l. c.* 444.—Am. trop.
cornifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 150. t. 547.—N. Granat.
corymbosa, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
crassifolia, *Benth. Bot. Voy. Sulph.* 97. t. 37.—N. Granat.
crenata, *Berg, in Mart. Fl. Bras.* xiv. i. 456.—Bras.
cuneata, *Berg, l. c.* 446.—Bras.
cyanea, *Berg, l. c.* 454.—Bras.
dentata, *Berg, l. c.* 457.—Bras.
desertorum, *Berg, l. c.* 450.—Bras.
dimorpha, *Berg, l. c.* 609.—Bras.
discolor, *Berg, l. c.* 459.—Bras.

CAMPOMANESIA :—
eriantha, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
eugenioides, *Blume, l. c.*—Bras.
fruticosa, *Berg, in Mart. Fl. Bras.* xiv. i. 458.—Bras.
fusca, *Berg, l. c.* 447.—Bras.
Gardneriana, *Berg, l. c.* 455.—Bras.
glabra, Benth. in Hook. Journ. Bot. ii. (1843) 319 = Calycolpus glaber.
glabra, *Berg, in Mart. Fl. Bras.* xiv. i. 450.—Bras.
Goethsiia, Hemsl. Biol. Centr. Am. Bot. i. 407 = Calycolpus glaber.
grandiflora, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 182.—Guiana.
Guaviroba, *Benth. & Hook. f. Gen.* i. 713.—Bras.
guazumaefolia, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
heterophylla, *Berg, in Mart. Fl. Bras.* xiv. i. 440.—Bras.
hirsuta, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 353.—Bras.
Houlletii, *Berg, in Mart. Fl. Bras.* xiv. i. 449.—Bras.
lanceolata, *Berg, l. c.* 612.—Bras.
Langsdorffii, *Berg, l. c.* 610.—Bras.
laurifolia, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 353.—Bras.
linearifolia, *Ruiz & Pav. Fl. Per.* iv. t. 422; *Syst.* 138.—Peruv.
mallifolia, *Berg, in Mart. Fl. Bras.* xiv. i. 452.—Bras.
Martiana, *Berg, l. c.* 610.—Bras.
mediterranea, *Berg, l. c.* 458.—Bras.
microcarpa, *Berg, l. c.* 455.—Bras.
multiflora, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
obscura, *Berg, in Mart. Fl. Bras.* xiv. i. 451.—Bras.
obversa, *Berg, l. c.* 445.—Bras.
ovalifolia, *Berg, l. c.* 452.—Bras.
Pohliana, *Berg, l. c.* 441.—Bras.
Polteaui, *Berg, in Linnaea*, xxvii. (1854) 432.—Bras.
pubescens, *Berg, in Mart. Fl. Bras.* xiv. i. 448.—Bras.
racemosa, *Berg, l. c.* 458.—Bras.
repanda, *Berg, l. c.* 456.—Bras.
reticulata, *Berg, l. c.* 439.—Bras.
rhombea, *Berg, l. c.* 451.—Bras.
rhytidophylla, *Berg, l. c.* 608.—Bras.
Riedeliana, *Berg, l. c.* 610.—Bras.
rigens, *Berg, l. c.* 430.—Bras.
salvaefolia, *Berg, l. c.* 442.—Bras.
suaveolens, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
suffruticosa, *Berg, in Mart. Fl. Bras.* xiv. i. 448.—Bras.
synchrona, *Berg, l. c.* 444.—Bras.
tenuifolia, *Berg, l. c.* 462.—Bras.
tomentosa, *H. B. & K. Nov. Gen. et Sp.* vi. 151. t. 547.—N. Granat.
transalpina, *Berg, in Mart. Fl. Bras.* xiv. i. 459.—Bras.
trifiora, *Baill. Hist. Pl.* vi. 342.—Bras.
vaccinioides, *Berg, in Mart. Fl. Bras.* xiv. i. 450.—Bras.
velutina, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
virescens, *Berg, in Mart. Fl. Bras.* xiv. i. 611.—Bras.
Widgreniana, *Berg, l. c.* 447.—Bras.
xanthocarpa, *Berg, l. c.* 451.—Bras.

CAMPSANTHUS, Steud. Nom. ed. II. i. 272 (1840), aphalm. = COMPSANTHUS, Spreng. = **Trioyrtis**, Wall. (Liliac.).

CAMPSIANDRA, Benth. in Hook. Journ. Bot. ii. (1840) 93. *LEGUMINOSAE*, Benth. & Hook. f. i. 564.
angustifolia, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 56.—Am. trop.
comosa, *Benth. in Hook. Journ. Bot.* ii. (1840) 93.—Guiana.
laurifolia, *Benth. l. c.* 94.—Bras.
rosea, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 62.—Bras.

CAMPSIDIUM, Seem. in Bonplandia, x. (1862) 147; in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 32 = **Tecoma**, Juss. (Bignon.).
chilense, Reiss. & Seem. in Bonplandia, x. (1862) 147.—Chili.
flicifolium, A. Van Geert, Cat. n. 71 (1874); *Fl. et Pom.* (1874) 280.—Ins. Fiji.

CAMPSIS, Lour. Fl. Cochinch. 377 (1790) = **Tecoma** Juss. (Bignon.).
adrepens, Lour. l. c. = Incarvillea sinensis.
amboinensis, Seem. in Journ. Bot. v. (1867) 374 = T. amboinensis.
Curtisii, Seem. l. c. 371 = T. radicans ?
dendrophila, Seem. l. c. 373 = T. dendrophila.
Fortunei, Seem. l. c.—China.
Hausmannii, F. Muell. Fragm. iv. 148 = Haussmannia jucunda.
radicans, Seem. in Journ. Bot. v. (1867) 372 = T. radicans.

CAMPTEDERIA, Steud. Nom. ed. II. i. 272 (1840), aphalm. = CAMPDERIA, A. Rich.

CAMPTOCARPUS, Decne. in DC. Prod. viii. (1844) 498. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 744
Bojerianus, *Decne. l. c.* 494.—Madag.
crassifolius, *Decne. l. c.*—Madag.
linearis, *Decne. l. c.*—Madag.
mauritianus, *Decne. l. c.* 498.—Ins. Borbon.

CAMPTOCARPUS, C. Koch, in Linnaea, xvii. (1842) 804 = **Alkanna**, Tausch (Borragin.).
orientalis, C. Koch, l. c. = A. orientalis.

CAMPTOLOMA, Benth. in DC. Prod. x. 480 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 960.
rotundifolia, *Benth. l. c.* 481.—Afr. austr.
villosa, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 84.—Ins. Socotra.

CAMPTOPUS, Hook. f. Bot. Mag. t. 5755 (1869) = **Cephaëlis**, Sw. (Rubiac.).
Mannii, Hook. f. l. c. = Cephaëlis Mannii.

CAMPTOSEMA, Hook. & Arn. in Hook. Bot. Misc iii. (1833) 200. *LEGUMINOSAE*, Benth. & Hook. f. i. 536.
BIONIA, Mart. ex Benth. in Ann. Wien. Mus. ii (1838) 180.
acuminatum, *Benth. in Mart. Fl. Bras.* xv. i. 155.—Am. austr.
bellum, *Benth. l. c.* 156.—Am. austr.
coccineum, *Benth. l. c.* 154.—Am. austr.
cortaceum, *Benth. l. c.* 155.—Am. austr.
erythrinoides, *Benth. l. c.* 156.—Am. austr.
grandiflorum, *Benth. in Linnaea*, xxii. (1849) 516.—Bras.
pedicellatum, *Benth. in Hook. Journ. Bot.* ii. (1840) 60.—Guiana.
pinnatum, *Benth. in Mart. Fl. Bras.* xv. i. 525.—Bras.
rubicundum, *Hook. & Arn. in Hook. Bot. Misc.* iii (1833) 201.—Reg. Argent.
tomentosum, *Benth. in Mart. Fl. Bras.* xv. i. 155.—Am. austr.

CAMPTOSTEMON, Mast. in Hook. Ic. Pl. xii. 18. t. 1119 (1879). *MALVACEAE*.
Schultzii, *Mast. l. c.*—Austral.

CAMPTOTHECA, Decne. in Bull. Soc. Bot. Fr. xx (1873) 157. *CORNACEAE*.
acuminata, *Decne. l. c.*—Tibet.

CAMPULEIA, Thou. Gen. Nov. Madag. 7 (1806) = **Striga**, Lour. (Scrophularin.).
coccinea, Hook. Exot. Fl. t. 203 = S. lutea.
coerulea, Boj. ex Benth. l. c. = S. orobanchioides.
rosea, Boj. ex Benth. l. c. = S. orobanchioides.

CAMPULOA, Desv. Journ. Bot. i. (1813) 69 = **Cteniium**, Pers. (Gramin.).
brachystachya, Nees, ex Doell, in Mart. Fl. Bras. ii. 1. 73 = Ct. brachystachyum.
chapadensis, Trin. ex Doell, l. c. 74 = Ct. chapadense.
elegans, J. Gay, ex Kunth, Rev. Gram. i. 99 = Ct. elegans.
gracilis, Desv. Journ. Bot. i. (1813) 69 = Ct. corolinianum.
hirsuta, Desv. l. c. (= *Campuloaus yclustus*).—Afr. austr.
monostachya, Rosm. & Schult. Syst. ii. 818 = Ct. corolinianum.

CAMPULOCLINIUM, DC. Prod. v. 136 (1836) = **Eupatorium**, Tourn. (Compos.).
alternifolium, Gaud. in Hook. Lond. Journ. Bot. vi. (1847) 488 = Ageratum alternifolium.
arenarium, Gardn. l. c. v. (1846) 467.—Bras.
ascendens, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 368 = E. betonicæfolium.
corymbosum, Sch. Bip. ex Baker, l. c. 356 = E. Burchellii.
crenatum, Schlecht. in Mart. Herb. Fl. Bras. n. 668 = E. sordescens.
decumbens, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 344 = E. decumbens.
frutescens, Bartl. Ind. Sem. Hort. Goett. (1843).—Hab.?
hirsutum, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 488 = E. hirsutum.
klotzschioides, DC. Prod. v. 137 = E. klotzschioides.
macrocephalum, DC. l. c. = E. macrocephalum.
palustre, DC. in Deless. Ic. Sel. iv. t. 16 = E. palustre.
Pohlianum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 358 = E. macrocephalum.
polyphyllum, Sch. Bip. ex Baker, l. c. 354 = E. megacephalum.
regnebeckioides, Sch. Bip. ex Baker, l. c. 359 = E. hirsutum.
strigosum, Sch. Bip. ex Baker, l. c. 358 = E. macrocephalum.
subparniculatum, DC. Prod. v. 137 = E. sordescens.
surinamense, Miq. in Linnæa, xvii. (1843) 59 = E. Vitalbæ.
turbinatum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 357 = E. chlorolepis.
urticæfolium, DC. Prod. v. 137 = E. urticæfolium.
viridiflorum, Bartl. ex Baker, in Mart. Fl. Bras. vi. ii. 309 = E. viridiflorum.

CAMPULOSUS, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = **Ctenium**, Panz. (1814) (Gramin.).
acicularius, Trin. Sp. Gram. iii. t. 301 = Ct. elegans.
aromaticus, Trin. ex Steud. Nom. ed. II. i. 273 = Ct. carolinianum.
brachystachyus, Nees, Agrost. Bras. 417 = Ct. brachystachyum.
chapadensis, Trin. Sp. Gram. iii. 26. t. 308 = Ct. chapadense.
cirrhosus, Nees, Agrost. Bras. 416 = Ct. cirrhosum.
concinnus, Steud. Nom. ed. II. i. 272 = Ct. concinnum.
dolichostachys, Trin. ex Steud. l. c.—Hab.?
falcatus, Beauv. Agrost. 64.—Afr. austr.
gracilior, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = Ct. carolinianum.
gracilis, Bertol. in Mem. Acad. Sc. Bolog. ii. (1850) 602.—Am. bor.
hirsutus, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = *falcatus*.
macrostachys, Steud. Nom. ed. II. i. 29 = Ct. carolinianum.
monostachyus, Beauv. Agrost. 64. t. 13. f. 1 = Ct. carolinianum.
planifolius, Presl, Reliq. Haenk. i. 287 = Ct. planifolium.
tricorys, Trin. ex Steud. Nom. ed. II. i. 279.—Hab.?

CAMPYDORUM, Salisb. Gen. Pl. Fragm. 64 (1866) = **Polygonatum**, Adans. (Liliac.).
verticillatum, Sallab. l. c. = P. verticillatum.

CAMPYLANDRA, Baker, in Journ. Linn. Soc. xiv. (1875) 582. t. 20 = **Tupistra**, Ker-Gawl. (Liliac.).
aurantiaca, Baker, l. c. 582 = T. aurantiaca.
dasystachys, Wall. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 256 = Dasystachys campanulata.

CAMPYLANTHERA, Hook. Ic. Pl. t. 82 (1837) = **Pronaya**, Haeg. (Pittospor.).
ericoides, Lindl. in Mitch. Three Exped. ii. 277 = Marianthus procumbens.
Fraseri, Hook. Ic. Pl. t. 83 = P. elegans.

CAMPYLANTHERA, Schott, Melet. 35 (1832) = **Briodendron**, DC. (Meliac.).
Sennensis, Schott, l. c. = K. Senaarum.

CAMPYLANTHUS, Roth, Nov. Pl. Sp. 4 (1821). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 961.
Benthamii, *Webb*, *in Hook. Niger Fl.* 168.—Ins. Cap. Verd.
CAMPYLANTHUS:—
glabrs, *Benth. in DC. Prod.* x. 508.—Ins. Cap. Verd.
junceus, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) 1217.—Ind. or.
ramosissimus, *Wight*, *Ic.* iv. t. 1416.—Ind. or.; Beluchist.
salsoloides, *Roth*, *Nov Pl. Sp.* 4.—Ins. Teneriff.
salsoloides, Stocks, ex Boiss. Fl. Orient. iv. 433 = ramosissimus.
salsoloides, Webb, Phyt. Canar. iii. 126 = Benthamii.
spinosus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 84.—Ins. Socotra.

CAMPYLEIA, Spreng. Syst iv. Cur. Post. 222 (1831) = CAMPULEIA, Thou. = **Strigs**, Lour. (Scrophular.).

CAMPYLIA, Lindl. ex Sweet, Geran. sub t. 48 (1820) = **Pelargonium**, L'Hérit. (Geraniac.).
blatteria, Sweet, Geran. t. 88 = P. ovale.
cana, Colv. ex Sweet, l. c. t. 114 = P. ovale.
capillaris, Eckl. & Zeyh. Enum. 74 = P. capillare.
carinata, Lindl. ex Sweet, Geran. sub t. 48 = P. carinatum.
cartilaginea, Eckl. & Zeyh. Enum. 74 = P. cartilagineum.
coronopifolia, Sweet, Hort. Brit. ed. I. 76 = P. coronopifolium.
elata, Eckl. & Zeyh. Enum. 74 = P. elatum.
elegans, Colv. ex Sweet, Geran. t. 229 = P. elegans.
eriostemon, Eckl. & Zeyh. Enum. 73 = P. ovale.
holosericea, Sweet, Geran. t. 75 = P. ovale.
laciniata, Sweet, l. c. t. [401] = P. laciniatum.
Ornithrae, Sweet, Hort. Brit. ed. I. 75 = P. Oenotherae.
staticephylla, Eckl. & Zeyh. Enum. 74 = P. staticæfolium.
tricolor, Eckl. & Zeyh. l. c. = P. violaceum.
variegata, Sweet, Geran. t. 286 = P. marginatum.
verbascifiora, Sweet, l. c. ii. t. 157 = P. ovale.
veronicæfolia, Eckl. & Zeyh. Enum. 73 = P. veronicæfolium.

CAMPYLOBOTRYS, Lem. in Fl. des Serres, Sér. I. iii. (1847) sub t. 260 = **Hoffmannia**, Sw. (Rubiac.).
argyronnura, Linden, in Belg. Hortic. viii. (1858) 166.—Mexic.
discolor, Hook. Bot. Mag. t. 4530 = seq.?
discolor, Lem. in Fl. des Serres, Sér. I. iii. (1847) sub t. 260 = H. discolor.
Ghiesbreghtii, Lem. Illustr. Hortic. viii. (1861) t. 279 = H. Ghiesbregtii.
pyrophylla, Linden, Cat. n. 16 (1861) 2.—Mexic.
regalis, Linden, l. c. n. 15 (1860) = H. regalis.

CAMPYLOCARYUM, DC. ex A. DC. in Prod. x. 97, in syn. (1846) = **Alkanna**, Tausch (Borragin.).
smyrnaeum, Boiss. ex A. DC. Prod. x. 98 = A. Pinardi.
syriacum, Boiss. Fl. Orient. iv. 226 = A. syriaca.
syriacum, Boiss. & Hohen. ex A. DC. Prod. x. 101 = A. strigosa.
verrucosum, Boiss. ex A. DC. l. c. 102 = A. incana.

CAMPYLOCENTRON, Benth. in Journ. Linn. Soc. xviii. (1881) 337. *ORCHIDEÆ*, Benth. & Hook. f. iii. 595.
Todaroa, A. Rich. in Ann. Sc. Nat. Sér. III. iii. (1845) 28.
Schiedei, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* iii. 393.—Mexic.

CAMPYLOCERA, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1842) 267 = **Specularia**, Heist. (Campanul.).
leptocarpa, Nutt. l. c. = S. leptocarpa.

CAMPYLOCHINIUM, Endl. Gen. 869 (1838) = CAMPULOCHINIUM, DC. = **Eupatorium**, Tourn. (Compos.).

CAMPYLONEMA, Poir. Illustr. 598 (1823); Schult. f. Syst. vii. 96, 1507 (1830) = **Campynema**, Labill. (Amaryllid.).
linearе, Schult. f. l. c. = Campynema lineare.

CAMPYLOPTERA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1841) 194 = **Asthionema**, R. Br. (Crucif.).
heterocarpa, Baill. Hist. Pl. iii. 291 = Ae. campylopterum.
syriaca, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1841) 194 = Ae. campylopterum.

CAMPYLOPUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 360 = **Hypericum**, Tourn.
cerastioides, Spach, Hist. Vég. Phan. v. 412 = H. rhodopeum.

CAMPYLOSIPHON, Benth. in Hook. Ic. Pl. t. 1384 (1880). *BURMANNIACEÆ*, Benth. & Hook. f. iii. 456.
purpurascens, *Benth. l. c.*—Am. austr.

CAMPYLOSIPHON, St. Lag. in Ann. Soc. Bot. Lyon, *vii.* (1880) 135 = **Siphocampylus**, Pohl (Campanul.).
lycioideus, St. Lag. l. c. = S. lycioides.

CAMPYLOSPORUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 363 = **Hypericum**, Tourn.
angustifolius, Spach, Hist. Vég. Phan. v. 425 = H. lanceolatum.
madagascariensis, Spach, l. c. 426 = H. madagascariense.
reticulatus, Spach, l. c. 424 = H. lanceolatum.

CAMPYLOSTACHYS, Kunth, in Abh. Akad. Berl. (1831) 206. *VERBENACEÆ*, Benth. & Hook f. ii. 1127.
Campilostachys, A. Juss. in Orb. Dict. xii. 38 (1849).
abbreviata, *E. Mey. Comm. Pl. Afr. Austr.* i. 279.—Afr. austr.
cernua, *Kunth*, *in Abh. Akad. Berl.* (1831) 201.—Afr. austr.
cernua, Zeyh. ex DC. Prod. xii. 607 = Stilbe pinastra.
phylicoides, Sond. in Linnæa, xix. (1847) 209 = Stilbe phylicoides.

CAMPYLOSTACHYS, E. Mey. in Drège, Zwei Pfl. Docum. 83 (1849) = **Fimbristylis**, Vahl (Cyperac.).
abbreviata, E. Mey. l. c. = F. ferruginea.

CAMPYLOSTEMON, E. Mey. in Drège, Zwei Pfl. Docum. 170 (1848), nomen = **Justicia**, Linn. (Acanthac.).
campanulatus, E. Mey. l. c. (= *Leptostachya Campylostemon*).—Afr. austr.
paniculatus, Nees, in DC. Prod. xi. 378 = prsec.

CAMPYLOSTEMON, Welw. in Benth. & Hook. f. Gen. i. 898 (1862). *CELASTRINEÆ*, Benth. & Hook. f. i. c.
sagolense, *Welw. ex Oliver, in Journ. Linn. Soc.* x. (1869) 44.—Afr. trop.

CAMPYLOTHECA, Cass. in Dict. Sc. Nat. ii. 476 (1827) = **Corvispota**, Linn. (Compos.).
angustifolia, Walp. Rep. ii. 618 = Bidens angustifolia.
australis, Less. in Linnæa, vi. (1831) 509 = Cor. micrantha.
gracilis, Walp. Rep. ii. 618 = Bidens gracilis.
grandiflora, DC. Prod. v. 598.—Ins. Sandvic.
micrantha, Cass. in Dict. Sc. Nat. lix. 321 = Cor. micrantha.
mutica, Walp. Rep. ii. 618 = Bidens mutica.

CAMPYLOTROPIS, Bunge, Pl. Mongholico-Chin. 6 (1835) = **Lespedeza**, Michx. (Legumin.).
chinensis, Bunge, l. c. 7 = L. macrocarpa.
cytisoides, Benth. in Pl. Jungh. i. 230.—Java.
macrostyla, Lindl. ex Miq. Fl. Ind. Bat. i. 280 = L. eriocarpa.
virgata, Miq. l. c. 280 = L. virgata.

CAMPYLUS, Lour. Fl. Cochinch. 113 (1790) (Genus dub.).
sinensis, *Lour. l. c.*—China.

CAMPYNEMA, Labill. Nov. Holl. Pl. i. 98. t. 131 (1804). *AMARYLLIDEÆ*, Benth. & Hook. f. Gen. iii. 716.
Campylonema, Poir. Illustr. 598 (1823); Schult. f. Syst. vii. 96, 1507 (1830).
lineare, *Labill. l. c.*—Ins. Tasman.
pygmaeum, *F. Muell. ex Benth. Fl. Austral.* vi. 415.—Ins. Tasman.; Austral.

CAMUNIUM, Adans. Fam. ii. 166 (1763) = **Aglaia**, Lour. (Meliac.).
beaguirae, Buch.-Ham. ex Wall. Cat. n. 1279 = A. edulis.
chinense, Roxb. Hort. Beng. 18; Fl. Ind. i. 626 = A. odorata.

CAMUTIA, Bonat. ex Stend. Nom. ed. II. i. 278 (1840) = **Malampodium**, Linn. (Compos.).
perfoliata, Bonat. ex Stend. l. c. = M. perfoliatum.

CANABIS, Roth, Tent. Fl. Germ. i. 422 (1788), sphalm. = **Cannabis**, Tourn. (Urtic.).

CANAHIA, Steud. Nom. ed. I. 146 (1821) = **Kanahia**, R. Br. (Asclepiad.).
laniflora, Steud. l. c. = K. laniflora.

CANALA, Pohl, Pl. Bras. Ic. ii. 62. t. 142 (1831) = **Spigelia**, Linn. (Loganiac.).
heliotropioides, Pohl, l. c. 63. t. 143 = S. Olfersiana.
macrophylla, Pohl, l. c. 64 = S. macrophylla.
rubiaefolia, Pohl, l. c. 65 = S. Schlechtendaliana.

CANALIA, F. W. Schmidt, Neue u. Selt. Pfl. 30 (1793) = **Gnidia**, Linn. (Thymel.).
daphnoides, F. W. Schmidt, l. c. = G. pinifolia.

CANANGA, Aubl. Pl. Gui. i. 607. t. 244 (1775) = **Guatteria**, Ruiz & Pav. (1794) (Anonac.).
Candolleana, Warm. in Kjoeb. Vidensk. Meddel. (1879) 145 = G. Candolleana.
Klotzschiana, Warm. l. c. = G. Klotzschiana.
lanceа, Poit. ex DC. Syst. i. 506 = G. virgata.
Ouregon, Aubl. Pl. Gui. i. 608 = G. Ouregon.
pseudolanceа, Poit. ex DC. Syst. i. 507 = G. laurifolia.
Schomburgkiana, Baill. Hist. Pl. i. 204 = G. Schomburgkiana.
Sellowiana, Warm. in Kjoeb. Vidensk. Meddel. (1878) 144 = G. Sellowiana.
villosissima. Warm. l. c. = G. villosissima.

CANANGA, Rumph. ex Hook. f. & Thoms. Fl. Ind. i. 129 (1855). *ANONACEAE*, Benth. & Hook. f. i. 24.
MONOCARPERIA, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 57.—Malacca.
odorata, *Hook. f. & Thoms. Fl. Ind.* i. 130.—Burma; Malaya.
virgata, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 57.—Malaya.

CANARIA, Linn. Mant. ii. 148 (1767) = **Canarina**, Linn. (Campanul.).
campanulata, Linn. l. c. = Canarina campanulata.

CANARINA, Linn. Mant. 588 (1767). *CAMPANULACEAE*, Benth. & Hook. f. ii. 558.
CANARIA, Linn. Mant. ii. 148 (1767).
PERSETVA, Scop. Introd. 180 (1777).
Campanula, Lam. Encyc. i. 598 = sect. campanulata, *Linn. Mant.* ii. 225.—Ins. Canar.
laevigata, G. Don, ex Loud. Hort. Brit. 189 = campanulata.
moluccana, *Rozb. Hort. Beng.* 67, nomen.— Ins. Molucc.
Zangueбar, *Lour. Fl. Cochinch.* 195.—Afr. trop.

CANARION, St. Lag. in Ann. Soc. Bot. Lyon, vii. 121 (1880) = **Canarium**, Rumph. (Burserac.).
pimelum, St. Lag. l. c. = Canarium Pimela.

CANARIOPSIS, Miq. Fl. Ind. Bat. i. ii. 651 (1859) = **Canarium**, Rumph. (Burserac.).
altissima, Miq. l. c. = Canarium altissimum.
angustifolia, Miq. l. c. 653 = Canarium angustifolium.
aspera, Miq. l. c. 653 = Canarium asperum.
decumana, Miq. l. c. = Canarium decumanum.
denticulata, Miq. l. c. 654 = Canarium denticulatum.
gimbra, Miq. l. c. 652 = Canarium balsamiferum.
hirsuta, Miq. l. c. 653 = Canarium hirsutum.
hispida, Miq. l. c. = Canarium hispidum.
paucijuga, Miq. l. c.—Ins. Molucc.
villosa, Miq. l. c. 652 = Canarium villosum.

CANARIUM, [Rumph.] Linn. Herb. Amboin. 9 (1754) (Amoen. Acad. iv. 121); Mant. 127 (1767). *BURSERACEAE*, Benth. & Hook. f. i. 394.
CANARIOPSIS, Miq. Fl. Ind. Bat. i. ii. 651 (1859).
CENARIUM, Linn. Amoen. Acad. iv. 121, in nota (1759).
COLOPHONIA, Comm. ex Kunth, in Ann.ˢ Sc. Nat. Sér. I. ii. (1824) 862.
MEHENBETHENE, Besl. ex Gaertn. Fruct. ii. 98 (1788); Walp. Ann. ii. 291 (1851).

CANARIUM:—
PACHYLOBUS, G. Don, Gen. Syst. ii. 69 (1832).
PIMELA, Lour. Fl. Cochinch. 407 (1790).
SCUTINANTHE, Thwaites, in Hook. Kew Journ. viii. (1856) 267. t. 8.
acutum, *Engl. in DC. Monog. Phan.* iv. 113.—Borneo.
album, Blanco, Fl. Filip. ed. I. 798 = luzonicum.
album, *Raeusch. Nom. ed.* III. 287; *DC. Prod.* ii. 80. —Cochinch.
altissimum, *Blume, Bijdr.* 1168.—Java.
altissimum, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 211 = Santiria laevigata.
angustifolium, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 117.—N. Guin.
asperum, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 215.—N. Guin.
australianum, *F. Muell. Fragm.* iii. 15.—Austral.
Balanae, *Engl. in DC. Monog. Phan.* iv. 117.—N. Caled.
balsamiferum, Moon, Cat. 68 = zeylanicum.
balsamiferum, *Willd. Sp. Pl.* iv. 760.—Ins. Molucc.
Beccarii, *Engl. in DC. Monog. Phan.* iv. 107.— Borneo.
bengalense, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 136.— Ind. or.
Bennettii, *Engl. in DC. Monog. Phan.* iv. 119.— Burma. Malacca.
Boivini, *Engl.* i. c. 110.—Madag.
brunneum, *Bedd. Fl. Sylv.* t. 127.—Zeylan.
coccineocrinitum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) 11. 296.—Ins. Andaman.
Colophania, Baker, Fl. Maurit. 44 = mauritianum.
commune, Blanco, Fl. Filip. ed. I. 791 = Cumingii.
commune, *Linn. Mant.* i. 127.—Ins. Molucc.
commune, Wight & Arn. Prod. 175 = zeylanicum.
Cumingii, *Engl. in DC. Monog. Phan.* iv. 152.—Ins. Philipp.
decumanum, *Gaertn. Fruct.* ii. 99. t. 102.—Ins. Molucc.
denticulatum, *Blume, Bijdr.* 1162.—Java.
dichotomum, *Miq. Fl. Ind. Bat.* i. ii. 648.—Sumatra.
divergens, *Engl. in DC. Monog. Phan.* iv. 148.— Borneo.
edule, *Hook. f. in Hook. Niger Fl.* 285.—Afr. trop.
euphyllum *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 295.—Ins. Andaman.
expteron, *Miq. Fl. Ind. Bat.* i. ii. 648.—Sumatra.
fastetiplum, *Miq. i. c. Suppl.* 525.—Sumatra.
fragile, *Engl. in DC. Monog. Phan.* iv. 138.—Borneo.
fuscum, *Engl. l. c.* 116.—Java.
giganteum, *Engl. l. c.* 106.—Borneo.
grisescens, Miq. Fl. Ind. Bat. i. ii. 650 = Santiria rostrata.
glaucum, *Blume, Mus. Bot. Lugd. Bat.* i. 219.— Sumatra.
gracile, *Engl. in DC. Monog. Phan.* iv. 140.—Ins. Philipp.
grandiflorum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 533.—Malacca.
Harami, *Boj. Hort. Maurit.* 33.—Madag.
Harveyi, *Seem. Fl. Vit.* 35.—Ins. Tonga.
hirsutum, *Willd. Sp. Pl.* iv. 760.—Ins. Molucc.
hirtellum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 534.—Penang.
hispidum, *Blume, Bijdr.* 1168.—Java.
Ierana, Engl. in DC. Monog. Phan. iv. 150 = Harami.
incurvatum, *Engl. l. c.* 138.—Borneo.
Kadondon, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 535.—Malacca.
Kipella, *Miq. Fl. Ind. Bat.* i. ii. 646.—Java.
Kitenga, *Afz.* t. 647.—Java.
Kerthalsii, Miq. l. c. 645 = Santiria tomentosa.
laevigatum, Miq. l. c. 648 = Santiria laevigata.
laxiflorum, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 477 = microcarpum.
laxiflorum, *Zipp. ex Blume, Mus. Bot. Lugd. Bat.* i. 221.—Ins. Molucc.
lazum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 535. —Malacca.
legitimum, *Miq. Fl. Ind. Bat.* i. ii. 647.—Ins. Amboina.
littorale, *Blume, Bijdr.* 1164.—Java.
longiflorum, *Zipp. ex Miq. Fl. Ind. Bat.* i. ii. 647.— Ins. Molucc.
luzonicum, *Engl. in DC. Monog. Phan.* iv. 146.—Ins. Philipp.
luzonicum, *Miq. Fl. Ind. Bat.* i. ii. 651.—Ins. Philipp.

CANARIUM:—
macrophyllum, *Oliver, Fl. Trop. Afr.* i. 327.—Afr. trop.
madagascariense, *Engl. in DC. Monog. Phan.* iv. 111. —Madag.
Mahassan, *Miq. Fl. Ind. Bat. Suppl.* 526.—Sumatra.
mauritianum, *Blume, Mus. Bot. Lugd. Bat.* i. 217.— Ins. Maurit.
Mehenbethene, *Gaertn. Fruct.* ii. 98. t. 102.—Ins. Molucc.
microcarpum, *Willd. Sp. Pl.* iv. 760.—Ins. Molucc.
minutiflorum, *Engl. in DC. Monog. Phan.* iv. 123.— Ins. Philipp.
molle, *Engl.* i. c. 109.—Ins. Borbon.
moluccanum, Blume, Mus. Bot. Lugd. Bat. i. 216 = C. Mehenbethene.
montanum, Korth. ex Blume, l. c. 212 = Santiria montana.
Motleyanum, *Engl. in DC. Monog. Phan.* iv. 133.— Borneo.
muticiflorum, *Engl.* l. c. 128.—Madag.
multipinnatum, *Llanos, Fragm.* 107.—Ins. Philipp.
nigrum, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 138.—Ins. Amboina.
nitidum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 533 —Malaya.
oblongifolium, Miq. Fl. Ind. Bat. i. ii. 645 = Santiria oblongifolia.
odontophyllum, *Miq. l. c. Suppl.* 525.—Sumatra.
oleiferum, *Baill. Adansonia*, x. (1871–72) 341.—N. Caled.
ovatum, *Engl. in DC. Monog. Phan.* iv. 110.—Ins. Philipp.
paniculatum, Benth. ex Engl. l. c. 124 = mauritianum.
parvifolium, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 536.—Malacca.
patentinervium, *Miq. Fl. Ind. Bat. Suppl.* 526.— Sumatra.
pilosum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 533 —Malacca.
Pimela, Blanco, Fl. Filip. ed. II. 545 = villosum.
Pimela, Blume, Bijdr. 1163 = C. Kipella.
Pimela, *Kon. in Kon. & Sims, Ann. Bot.* i. (1805) 361.—China.
Pimela, Zoll. ex Engl. in DC. Monog. Phan. iv. 127 = Zollingeri.
pruinosum, *Engl.* l. c. 106.—borneo.
purpurascens, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 535.—Malacca.
Riedelianum, *Engl. in DC. Monog. Phan.* iv. 130.— Ins. Celebes.
rigidum, *Zipp. ex Miq. Fl. Ind. Bat.* i. ii. 648.—N. Guinea.
rostratum, *Zipp. ex Blume, Mus. Bot. Lugd. Bat.* i. 221.—Ins. Molucc.
rostriferum, *Miq. Fl. Ind. Bat. Suppl.* 526.— Sumatra.
rubiginosum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 535.—Malacca.
rubiginosum, Miq. Fl. Ind. Bat. i. ii. 651 = Santiria rubiginosa.
rufescens, Miq. l. c. 650 = Santiria rufescens.
rufum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 533. —Malacca.
rugosum, Miq. Fl. Ind. Bat. i. ii. 649 = Santiria rugosa.
Safiga, Boch.-Ham. ex Wight & Arn. Prod. 175 = Turpinia pomifera.
samoense, *Engl. in DC. Monog. Phan.* iv. 134.—Ins. Samoa.
Schweinfurthii, *Engl.* l. c. 145.—Afr. trop.
secundum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 532.—Malaya.
serrulatum, *Miq. l. c.* 646.—Java.
simplicifolium, *Engl. in DC. Monog. Phan.* iv. 146.— Ins. Amboina.
spectabile, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 116.— Sumatra.
strictum, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 139.—Ind. or.
subrepandum, *Miq. Fl. Ind. Bat. Suppl.* 525.— Sumatra.
subtrinervatum, *Engl. in DC. Monog. Phan.* iv. 109.— Java.
sylvestre, *Gaertn. Fruct.* ii. 99. t. 102.—Ins. Amboina.
tomentosum, *Blume, Mus. Bot. Lugd. Bat.* i. 219.— Malaya.
triandrum, *Engl. in DC. Monog. Phan.* iv. 145.— Malaya.

CANARIUM:—
trifoliolatum, *Engl. in DC. Monog. Phan.* iv. 124.—N. Caled.
villosum, *Benth. & Hook. f. ex Naves, in Blanco, Fl. Philipp.* ed. III. *App.* 46.—Ins. Philipp.
virgatum, Miq. Fl. Ind. Bat. i. II. 650 = Santiria virgata.
vitiense, *A. Gray, Bot. U. St. Expl. Exped.* i. 378.—Ins. Fiji.
Vriesianum, *Engl. in DC. Monog. Phan.* iv. 142.—Ins. Celebes.
zephyrinum, March. in Baill. Adansonia, viii. (1867–68) 55 = commune!
zeylanicum, *Blume, Mus. Bot. Lugd. Bat.* i. 218.—Zeylan.
Zollingeri, *Engl. in DC. Monog. Phan.* iv. 127.—Java.

CANAVALI, Adans. Fam. ii. 325 (1763); Thou. in Desv. Journ. Bot. i. (1813) 77 = **Canavalia**, DC. (Legumin.).

CANAVALIA, DC. Mém. Lég. ix. 375; Prod. ii. 403 (1825). *LEGUMINOSÆ*, Benth. & Hook. f. i. 537.
CANAVALI, Adans. Fam. ii. 325 (1763); Thou. in Desv. Journ. Bot. i. (1813) 77.
CLEMENTEA, Cav. in Anal. Cienc. Nat. vii. (1804) 63. t. 47.
MALOCCHIA, Savi, in Nuov. Giorn. Pisa, vii. (1824) 113.
WENDEROTHIA, Schlecht. in Linnæa, xii. (1838) 330.
alcicalma, *Macfad. Fl. Jamaic.* i. 292.—Jamaica.
Bauerriana, Endl. Prod. Fl. Norf. 91 = obtusifolia.
bonariensis, *Lindl. Bot. Reg.* t. 1199.—Bras.
Bonquete, *Montr. in Mém. Acad. Lyon*, x. (1860) 197. —N. Caled.
brasiliensis, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 135 = ensiformis.
cathartica, *Thou. in Desv. Journ. Bot.* i. (1813) 81.—Madag.
cryptodon, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 96 = bonariensis.
cubensis, *Griseb. in Mem Am. Acad.* N. S. viii. (1861) 178.—Cuba.
dolichoides, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 185 = Vigna dolichoides.
emarginata, G. Don, Gen. Syst. ii. 362 = obtusifolia.
ensiformis, *DC. Prod.* ii. 404.—Cosmop. trop.
Finlaysoniana, *R. Grah. in Wall. Cat.* n. 5585.—Siam.
galeata, *Gaudich. Freyc. Voy. Bot.* 486.—Ins. Sandvic.
Gaudichaudii, Endl. in Ann. Wien. Mus. i. (1836) 186 = galeata.
gladiata, DC. Prod. ii. 404 = ensiformis.
grandiflora, *Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Bras.
grandis, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 185 = Phaseolus velutinus.
incurva, DC. Prod. ii. 404 = ensiformis.
incurva, Thou. in Desv. Journ. Bot. i. (1813) 80 = ensiformis.
lenta, *Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Bras.
linearis, DC. Prod. ii. 404 = obtusifolia.
Lowreiri, G. Don, Gen. Syst. ii. 363 = ensiformis.
lucens, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 185 = Vigna lucens.
mackenroides, Steud. Nom. ed. II. i. 273 = cathartica.
maritima, Thou. in Desv. Journ. Bot. i. (1813) 80 = obtusifolia.
maxima, *Thou. l. c.*—Ind. or.
miniata, DC. Prod. ii. 404 = obtusifolia.
mollis, Wall. Cat. n. 5533 = ensiformis.
monets, *Wehw. Apont.* 666, n. 92.—Afr. trop.
monodon, E. Mey. Comm. Pl. Afr. Austr. 169 = bonariensis.
multiflora, Hook. & Arn. Bot. Beech. Voy. 416 = villosa.
obcordata, Volgt, Hort. Suburb. Calc. 288 = obtusifolia.
obtusifolia, *DC. Prod.* ii. 404.—Cosmop. trop.
paranensis, Benth. in Ann. Wien. Mus. ii. (1838) 135 = parviflora.
paranensis, Mook. & Arn. in Hook. Bot. Misc. iii. (1833) 200 = bonariensis.
parviflora, *Benth. in Mart. Fl. Bras.* xv. 1. 177.—Am. trop.

CANAVALIA:—
picta, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Bras.
polystachya, Schweinf. Rel. Kotsch. 25. t. 20 = ensiformis.
pubescens, Hook. & Arn. Bot. Beech. Voy. 81 = galeata.
rosea, DC. Prod. ii. 404 = obtusifolia.
rostrata, Benth. in Ann. Wien. Mus. ii. (1838) 135 = villosa.
rutilans, DC. Prod. ii. 404 = obtusifolia.
sericea, *A. Gray, Bot. U. St. Expl. Exped.* i. 440.—Ins. Fiji.
Stocksii, Dals. & Gibs. Bomb. Fl. 69 = ensiformis.
turgida, R. Grah. in Wall. Cat. n. 5534 = ensiformis.
villosa, *Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Mexic.
virosa, *Naves, ex Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 64.—Ins. Philipp.
virosa, Wight & Arn. Prod. 253 = ensiformis.

CANBYA, Parry, ex A. Gray, in Proc. Am. Acad. xii. (1877) 51. *PAPAVERACEÆ*.
candida, *Parry, ex A. Gray, l. c.*—Calif.

CANCELLARIA. Sch. Bip. ex Oliver, Fl. Trop. Afr. iii. 321 (1877) = **Adelostigma**, Steetz (Compos.).
senegalensis, Sch. Bip. ex Oliver, l. c. = A. senegalensis.

CANCRINIA, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 194. *COMPOSITÆ*, Benth. & Hook. f. ii. 424.
chrysocephala, *Kar. & Kir. l. c.* 125.—As. centr.

CANDARUM, Schott, Melet. i. 17 (1832) = **Amorphophallus**, Blume (Aroid.).
Hookeri, Schott, l. c. = A. campanulatus.
Roxburghii, Schott, l. c. = A. campanulatus.
Rumphii, Schott, l. c. = A. campanulatus.

CANDELABRIA, Hochst. in Flora, xxvi. (1843) 79 = **Bridelia**, Willd. (Euphorbiac.).
micrantha, Hochst. l. c. = B. micrantha.
polystachya, *Fisch. in Mem Am. Acad.* N. S. viii. (1861) 178.—Cuba.
dolichoides, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 185 = Vigna dolichoides.
emarginata, G. Don, Gen. Syst. ii. 362 = obtusifolia.
ensiformis, *DC. Prod.* ii. 404.—Cosmop. trop.
Finlaysoniana, *R. Grah. in Wall. Cat.* n. 5585.—Siam.
galeata, *Gaudich. Freyc. Voy. Bot.* 486.—Ins. Sandvic.
Gaudichaudii, Endl. in Ann. Wien. Mus. i. (1836) 186 = galeata.
gladiata, DC. Prod. ii. 404 = ensiformis.
grandiflora, *Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Bras.
grandis, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 185 = Phaseolus velutinus.
incurva, DC. Prod. ii. 404 = ensiformis.
incurva, Thou. in Desv. Journ. Bot. i. (1813) 80 = ensiformis.
lenta, *Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Bras.
linearis, DC. Prod. ii. 404 = obtusifolia.
Lowreiri, G. Don, Gen. Syst. ii. 363 = ensiformis.
lucens, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 185 = Vigna lucens.
mackenroides, Steud. Nom. ed. II. i. 273 = cathartica.
maritima, Thou. in Desv. Journ. Bot. i. (1813) 80 = obtusifolia.
maxima, *Thou. l. c.*—Ind. or.
miniata, DC. Prod. ii. 404 = obtusifolia.
mollis, Wall. Cat. n. 5533 = ensiformis.
monets, *Wehw. Apont.* 666, n. 92.—Afr. trop.
monodon, E. Mey. Comm. Pl. Afr. Austr. 169 = bonariensis.
multiflora, Hook. & Arn. Bot. Beech. Voy. 416 = villosa.
obcordata, Volgt, Hort. Suburb. Calc. 288 = obtusifolia.
obtusifolia, *DC. Prod.* ii. 404.—Cosmop. trop.
paranensis, Benth. in Ann. Wien. Mus. ii. (1838) 135 = parviflora.
paranensis, Mook. & Arn. in Hook. Bot. Misc. iii. (1833) 200 = bonariensis.
parviflora, *Benth. in Mart. Fl. Bras.* xv. 1. 177.—Am. trop.

CANDOLLEA :—
diuroides, F. Muell. Census, 86 = Stylidium diuroides
diversifolia, F. Muell. l. c. = S. diversifolia.
eglandulosa, F. Muell. l. c. = S. eglandulosa.
elongata, F. Muell. l. c. 85 = S. elongata.
emarginata, F. Muell. l. c. 86 = S. emarginata.
eriorrhiza, F. Muell. l. c. = S. eriorrhiza.
falcata, F. Muell. l. c. = S. falcata.
fasciculata, F. Muell. l. c. = S. fasciculata.
filifera, F. Muell. l. c. = S. filifera.
fusiloba, F. Muell. l. c. = S. fusiloba.
Floodii, F. Muell. l. c. = S. Floodii.
floribunda, F. Muell. l. c. = S. floribunda.
glandulosa, F. Muell. l. c. = S. glandulosa.
glauca, Labill. in Ann. Mus. Par. vi. (1805) 454 = S. glaucum.
graminifolia, F. Muell. Census, 85 = S. graminifolia.
guttata, F. Muell. l. c. = S. guttata.
hirsuta, F. Muell. l. c. = S. hirsuta.
imbricata, F. Muell. l. c. 86 = S. imbricata.
juncea, F. Muell. l. c. 85 = S. juncea.
laricifolia, F. Muell. l. c. 86 = S. laricifolia.
lepida, F. Muell. l. c. = S. lepida.
leptophylla, F. Muell. l. c. = S. leptophylla.
leptorrhiza, F. Muell. l. c. = S. leptorrhiza.
limbata, F. Muell. l. c. = S. limbata.
linearis, F. Muell. l. c. 86 = S. linearis.
linearis, F. Muell. l. c. 86 = S. linearis.
lobuliflora, F. Muell. l. c. = S. lobuliflora.
longituba, F. Muell. l. c. = S. longituba.
lutea, F. Muell. l. c. = S. lutea.
muscicola, F. Muell. l. c. = S. muscicola.
pachyrrhiza, F. Muell. l. c. = S. pachyrrhiza.
pedunculata, F. Muell. l. c. = S. pedunculata.
perpusilla, F. Muell. l. c. = S. perpusilla.
petiolaris, F. Muell. l. c. = S. petiolaris.
pilosa, Labill. in Ann. Mus. Par. vi. (1805) 453 = S. pilosum.
Preissii, F. Muell. Census, 86 = S. Preissii.
pubigera, F. Muell. l. c. = S. pubigera.
pulchella, F. Muell. l. c. = S. pulchella.
pychnostachya, F. Muell. l. c. = S. pychnostachya.
pygmæa, F. Muell. l. c. = S. pygmæa.
redupliçata, F. Muell. l. c. 85 = S. redupliçata.
repens, F. Muell. l. c. = S. repens.
rhynchocarpa, F. Muell. l. c. 86 = S. rhynchocarpa.
rotundifolia, F. Muell. l. c. = S. rotundifolia.
rupestris, F. Muell. l. c. = S. rupestris.
scabrida, F. Muell. 85 = S. scabrida.
scandens, F. Muell. l. c. 86 = S. scandens.
schizantha, F. Muell. l. c. = S. schizantha.
serrulata, Labill. in Ann. Mus. Par. vi. (1805) 454 = S. graminifolium.
setacea, Labill. l. c. 455 = S. spinulosum!
sobolifera, F. Muell. Census, 86 = S. sobolifera.
spathulata, F. Muell. l. c. = S. spathulata.
spinulosa, F. Muell. l. c. 85 = S. spinulosa.
squamellosa, F. Muell. l. c. 86 = S. squamellosa.
streptocarpa, F. Muell. l. c. = S. streptocarpa.
striata, F. Muell. l. c. = S. striata.
tenerrima, F. Muell. l. c. = S. tenerrima.
trichopoda, F. Muell. l. c. = S. trichopoda.
uliginosa, F. Muell. l. c. = S. uliginosa.
umbellata, Labill. in Ann. Mus. Par. vi. (1805) 456 = S. graminifolium.
uniflora, F. Muell. Census, 86 = S. uniflora.
utricularioides, F. Muell. l. c. = S. utriculariodes.
verticillata, F. Muell. l. c. = S. pulcillata.
violacea, F. Muell. l. c. = S. violacea.

CANDOLLEA, Labill. Nov. Holl. Pl. ii. 38 t. 176 (1806). *DILLENIACEÆ*, Benth. & Hook. f. i. 14.
assimilis, Steud. in Lehm. Pl. Preiss. i. 278 = pedunculata.
calycina, Steud. l. c. 274 = tetrandra.
cuneiformis, *Labill. Nov. Holl. Pl.* ii. 34. t. 176.—Austral.
Cunninghamii, Benth. in Mausd. Bot. ii. t. 85 = Hibbertia Cunninghamii.
cygnorum, *Steud. in Lehm. Pl. Preiss.* i. 275.—Austral.
dennophylla, *Benth. Fl. Austral.* i. 43.—Austral.
exasperata, *Steud. in Lehm. Pl. Preiss.* i. 276.—Austral.
fasciculata, *R. Br. ex DC. Syst.* i. 434.—Austral.
glaberrima, *Steud. in Lehm. Pl. Preiss.* i. 274.—Austral.

CANDOLLEA:—
glomerosa, *Benth. Fl. Austral.* i. 48.—Austral.
helianthemoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 8.—Austral.
Hueglii, *Benth. in Enum. Pl. Hueg.* 4.—Austral.
hochstedes, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 7 = fasciculata.
latifolia, Steud. in Lehm. Pl. Preiss. i. 273 = tetrandra.
pachyrrhiza, *Benth. Fl. Austral.* i. 44.—Austral.
parviflora, Steud. in Lehm. Pl. Preiss. i. 276 = pedunculata.
pedunculata, *R. Br. ex DC. Syst.* i. 424.—Austral.
Preissiana, *Steud. in Lehm. Pl. Preiss.* i. 274.—Austral.
ravenaea, Endl. in Enum. Pl. Hueg. 5 = pedunculata.
rupestris, Steud. in Lehm. Pl. Preiss. i. 275 = glaberrima.
striata, Steud. in Lehm. Pl. Preiss. i. 274 = Hueglii.
subnigricata, Steud. l. c. 276 = glaberrima.
teretifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 6.—Austral.
tetrandra, *Lindl. Bot. Reg.* (1842) *Misc.* 40.—Austral.
tridentata, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 140 = pedunculata.
uncinata, *Benth. Fl. Austral.* i. 46.—Austral.
vaginata, *Benth. l. c.* 45.—Austral.

CANDOLLEA, Steud. Nom. ed. II. i. 278 (1840) = DECANDOLIA, Bast. = **Agrostis**, Linn. (Gramin.).

CANELLA, P. Br. Hist. Jamaic. 275 t. 27. f. 3 (1756); Sw. in Trans. Linn. Soc. i. (1791) 96. t. 8. *CANELLACEÆ*, Benth. & Hook. f. i. 191.
WINTERANIA, Linn. Hort. Cliff. 488 (1737); et Syst. ed. X. (1759) 1045 (Wintieania).
alba, *Murr. Syst.* ed. XIV. 448.—Ind. occ.
alba, Nees, Pl. Officin. t. 418 = Cinnamodendron corticosum.
axillaris, Nees & Mart. in Neuwied, Reise nach Brasil. ii. 326 = Cinnamodendron axillare.
laurifolia, Lodd. Cat. ex Sweet, Hort. Brit. 65 = alba.
obtusifolia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 348.—Venezuela.
Winterana, Gaertn. Fruct. i. t. 77 = alba.

CANEPHORA, Juss. Gen. 908 (1789). *RUBIACEÆ*, Benth. & Hook. f. ii. 74.
axillaris, Willd. Sp. Pl. i. 976 = madagascariensis.
capitata, Willd. l. c. = Barchellia capensis.
madagascariensis, *Y. F. Gmel. Syst.* 872.—Madag.

CANICIDIA, Vell. Fl. Flum. 184 (1825); iv. t. 139 (1827) = **Connarus**, Linn.
rostrata, Vell. Il. cc. = Connarus cymosa.

CANIDIA, Salisb. Gen. Pl. Fragm. 92 (1866) = **Allium**, Tourn. (Liliac.).
magica, Salisb. l. c. = A. magicum.

CANIRAM, Thou. ex Steud. Nom. ed. II. i. 274 (1840) = **Strychnos**, Linn. (Loganiac.).

CANISTRUM, E. Morr. Belg. Hortic. (1878) 257. t. 15 = **Aechmea**, Ruiz & Pav. (Bromeliac.).
aurantiacum, E. Morr. l. c. = A. aurantiaca.
eburneum, E. Morr. l. c. xxix. (1879) 168. t. 13, 14.—Bras.
fuscum, E. Morr. l. c. xxxiii. (1883) 195.—Bras.
purpureum, E. Morr. l. c.—Bras.
roseum, E. Morr. l. c. xxix. (1879) 301.—Bras.
viride, E. Morr. ex Ch. Morr. l. c. xxiv. (1874) 376. t. 16 = A. viridis.

CANKRIENIA, De Vriese, in Jaarb. Nederl. Maatsch. Tuinb. (1850) = **Primula**, Tourn.
chrysantha, De Vriese, l. c. 80 = P. imperialis.
farinosa, Zoll. Syst. Verz. Ind. Archip. 89.—Java.

CANNA, Linn. Syst. ed. I (1735); Gen. ed. I. 1 (1737). *SCITAMINEÆ*, Benth. & Hook. f. iii. 654.
ACHIRIDA, Horan. Prod. Scitam. 18. t. 2 (1862).
CANNACORUS, Tourn. ex Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 272.
DISTEMON, Bonché, in Linnæa, xviii. (1844) 494.
EURYSTYLUS, Bouché, l. c. (1844) 485.

CANNA:—
Katurala, Adans. Fam. ii. 67 (1763).
Xyphostylis, Rafin. Fl. Tellur. iv. 52 (1836).
Achiras, *Gill. in Bot. Reg.* t. 1255.—Chili.
Altenstenii, *Bouché, in Linnæa*, xi. (1837) 236.—Am. austr.
angustifolia, *Linn. Sp. Pl.* 1.—Bras.
angustifolia, Wall. Fl. Carol. 59 = flaccida.
aurantiaca, Rosc. Scit. t. 21 = lutea.
aurantiaca ×, *Tineo, ex Tod. in Ann. Sc. Nat. Sér.* IV. xi. (1859) 277.—Hab.?
aurea, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Hab.?
aureo-vittata, Lodd. Bot. Cab. t. 449 = limbata.
barbadica, Hort. ex Bouché, in Linnæa, viii. (1833) 145 = occidentalis.
bidentata, *Bertol. in Mem. Acc. Sc. Bolog.* x. (1859) 93.—Afr. trop.
bifida, Herb. ex Roem. & Schult. Syst. i. Add. i. 69 = speciosa.
brasiliensis, Hort. Vind. ex Bouché, in Linnæa, xii. (1838) 148 = Jacquini.
brasiliensis, *Rosc. ex Spreng. Syst.* iv. *Cur. Post.* 5.—Bras.
Buchii, Weinm. in Syll. Ratisb. i. (1824) 119; Flora, vii. (1824) 10 = pedunculata.
caripensis, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Hab.?
carnea, *Rosc. Scit.* t. 15.—Bras.
chinensis, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 170 = orientalis.
cinnabrina, *Bouché, in Linnæa*, xviii. (1844) 490.—Mexic.
coccinea, Link, Enum. Hort. Berol. i. 1 = patens.
coccinea, *Lodd. Bot. Cab. Ind.* VIII. n. 3.—Ind. or.
commutata, Bouché, in Linnæa, viii. (1833) 147 = lutea.
compacta, Hort. ex Bouché, l. c. 145 = occidentalis.
compacta, *Rosc. Scit.* t. 22.—Hab.?
concinna, *Bouché, in Linnæa*, xviii. (1844) 491.—Am. trop.
coreo, *Lag. ex Reichb. Hort. Bot.* t. 148.—Hab.?
crocea, Roem. & Schult. Syst. i. 11 = patens.
curviflora, *Hort. ex Horan. Prod. Scitam.* 18.—Am. centr.
denudata, *Bouché, in Linnæa*, xviii. (1844) 489.—China.
denudata, *Rosc. Scit.* t. 28.—Bras.
discolor, *Lindl. Bot. Reg.* t. 1251.—Ins. Trinitat.
edulis, *Ker-Gawl. in Bot. Reg.* ix. t. 775.—Am. trop.
Ehrenbergii, *Bouché, in Linnæa*, viii. (1833) 150.—Ins. S. Thomas.
elegans, Rafin. Fl. Ludov. 148 = flaccida?
elliptica, *Italia*, Stokes, Bot. Mat. Med. i. 2 = indica.
excelsa, Lodd. Cat. ex Lond. Hort. Brit. 1 = edulis.
excelsa, *Lodd. Bot. Cab.* n. 743*.—Bras.
exigua, *Bouché, in Linnæa*, xviii. (1844) 490.—Nepal.
eximia, *Hort. ex Horan. Prod. Scitam.* 18.—Bras.
Fintelmanni, *Bouché, in Linnæa*, xviii. (1844) 487.—Hab.?
flaccida, *Rosc. in Trans. Linn. Soc.* viii. (1807) 339.—Am. bor.
flava, Michx. ex Lam. in Journ. Nat. Hist. Par. i. 418 = flaccida.
flavescens, *Link, Handb.* i. 296.—Ind. or.
floribunda, *Bouché, in Linnæa*, xviii. (1844) 489.—Hab.?
formosa, *Bouché, l. c.* 491.—Bras.
fulgida, *Bouché, l. c.* 490.—Ins. S. Barthol.
gramilia, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 23 = gigantea.
gigantea, *Desf. Tabl. Hort. Par.* ed. I. 32.—Bras.
glauca, *Linn. Sp. Pl.* 1.—Ind. occ.; Mexic.
glauca, Wall. Pl. Carol. 59 = flaccida.
hallcomiæfolia, *Hort. Berol. ex Bouché, in Linnæa*, viii. (1833) 134.—Mexic.
humilis, *Bouché, l. c.* 152.—China.
indica, Curt. Bot. Mag. t. 454 = patens.
indica, [DC. in] Red. Liliac. t. 501 = speciosa.
indica, *Linn. Sp. Pl.* 1.—Cosmop. trop.
indica, Rosc. in Trans. Linn. Soc. viii. (1807) 338 = orientalis.
indica, Ruiz & Pav. l. c.—Peruv.
iridiflora, Willd. Enum. Hort. Berol. Suppl. 1 = gigantea.

CANNA:—
Jacquini, *Bouché, in Linnæa*, xii. (1838) 148.—Bras.
janea, *Rets. Obs.* fasc. i. 9.—China.
laeta, *Bouché, in Linnæa*, xviii. (1844) 489.—Hab.?
lagunensis, Lindl. Bot. Reg. t. 1311 = lutea.
Lamberti, *Lindl. l. c.* t. 476.—Ind. occ.; Guiana.
lanceolata, *Lodd. ex Lond. Hort. Brit.* 1.—Bras.
lanuginosa, *Rosc. Scit.* t. 16.—Am. trop.
latifolia, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. bor.
latifolia, Rosc. Scit. t. 4 = gigantea.
leptochila, *Bouché, in Linnæa*, viii. (1833) 152.—Hab.?
lescoarpa, *Bouché, l. c.* xviii. (1844) 492.—Am austr.
lilliiflora, *Warscew. ex Planch. in Fl. des Serres*, Sér. I. x. (1854–55) 211.—Am. centr.
limbata, *Rosc. Scit.* t. 9.—Bras.
Linkii, *Bouché, in Linnæa*, viii. (1833) 166.—Bras.
liturata, Link, ex A. Dietr. Sp. Pl. i. 12 = glauca.
longifolia, *Bouché, in Linnæa*, xviii. (1844) 486.—Mexic.
lutea, *Mill. Gard. Dict.* ed. VIII. n. 4.—Am. trop.
macrocarpa, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Am. trop.
macrophylla, *Hort. ex Horan. l. c.*—Am. trop.
maculata, *Link, Handb.* i. 297.—Am. trop.
maxima, Lodd. ex Rosc. Scit. t. 5 = Lamberti.
mexicana, A. Dietr. Sp. Pl. i. 11 = glauca.
minista, *Bouché, in Linnæa*, viii. (1833) 165.—Bras.
montana, *Blume, Enum. Pl. Jav.* 85.—Java.
Moritziana, *Bouché, in Linnæa*, xviii. (1844) 492.—Venezuela.
nepalensis, Wall. Cat. n. 6622 = indica.
neglecta, Hort. ex Steud. Nom. ed. II. i. 274 = lanuginosa.
neglecta, Weinm. in Flora, iii. (1820) 607 = gigantea.
nepalensis, Hort. Par. ex D. Dietr. Syn. Pl. i. 4 = indica
nudata, *Lodd. Cat. ex Steud. Nom.* ed. II. i. 274.—Hab.?
occidentalis, *Rosc. in Bot. Reg.* t. 772.—Ind. occ.
orientalis, Bouché, in Linnæa, viii. (1833) 152 = pulchra.
orientalis, *Rosc. Scit.* t. 12.—Ind. or.
Ottonia, *Bouché, ex Heynh. Nom.* ii. 116, nomen.—Venezuela.
ovata, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Hab.?
pallida, Rosc. Scit. t. 19 = lutea.
paniculata, *Ruiz & Pav. Fl. Per.* i. 1.—Peruv.
patens, Hook. Exot. Fl. t. 228 = limbata.
patens, *Rosc. in Trans. Linn. Soc.* viii. (1807) 338.—Hab.?
pedicellata, *Presl, Reliq. Haenk.* 106.—Chili.
pedunculata, *Sims, Bot. Mag.* t. 2323.—Am. trop.
pentaphylla, D. Dietr. Syn. Pl. i. 5 = platyphylla.
petiolata, *Wall. ex Horan. Prod. Scitam.* 17, nomen.—Hab.?
platyphylla, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 22.—Bras.
Poeppigii, *Bouché, in Linnæa*, xii. (1838) 143.—Bras.
polyclada, *Wawra, in Oestr. Bot. Zeitschr.* xiii. (1863) 7.—Bras.
polymorpha, *Lodd. ex Loud. Hort. Brit.* 1; *Bouché, in Linnæa*, viii. (1833) 151.—Am. austr.
portoricensis, *Bouché, l. c.* xii. (1838) 147.—Porto Rico.
prunosa, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 79.—Bras.
pulchra, *Bouché, ex Horan. Prod. Scitam.* 18.—Ind. or.
pulchra, *Hassk. Cat. Bogor. Alt.* 52; *et Pl. Jav. Rar.* i. 140.—Java.
recurvata, *Bouché, in Linnæa*, xviii. (1844) 488.—Hab.?
reflexa, Nees, ex D. Dietr. Syn. Pl. i. 4 = pedunculata.
Reevesii, *Lindl. Bot. Reg.* t. 2004.—China.
Rozconea, *Hort. Berol. ex Bouché, in Linnæa*, viii. (1833) 146.—Am. trop.
rotundifolia ×, E. André, in Rev. Hortic. (1892) 371.
rubra, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 169 = coccinea.
rubricaulis, Link, Enum. Hort. Berol. i. 1 = edulis.
sanguinea, Hort. Angl. ex Bouché, in Linnæa, viii. (1833) 154 = speciosa.
sanguinea, *Warsz. ex* Otto & Dietr. Allg. Gartenz. xix. (1851) 390 = *Warszewiczii*.
sanguinea-rubra, *Bouché, ex C. Koch, Allg. Gartenz.* (1858) 396.—Hab.?

CANNA:—
Schlechtendallana, *Bouché, in Linnæa*, xviii. (1844) 487.—Hab. ?
Schubertii, *Hort. ex Horan. Prod. Scitam.* 18.—Venezuela.
Sellowi, *Hort. Berol. ex Bouché, in Linnæa*, viii. (1833) 163.—Bras.
speciosa, Herb. ex Sims, Bot. Mag. t. 2317 = coccinea.
speciosa, Rosc. Scit. t. 17 = indica.
spectabilis, *Bouché, in Linnæa*, xviii. (1844) 487.—Hab. ?
stolonifera, [*Hort. Berol. ex*] *A. Dietr. Syn. Pl.* i. 11.—Reg. Argent.
stricta, *Bouché, in Linnæa*, xii. (1838) 144.—Hab. ?
sulphurea, Hort. ex Bouché, l. c. vill. (1838) 150 = lutea.
surinamensis, *Bouché, in Linnæa*, xviii. (1844) 491.—Guiana.
sylvestris, *Rosc. Scit.* t. 10.—Am. austr.
tenuiflora, *Bouché, in Linnæa*, viii. (1838) 150.—Ind. occ.
thyrsiflora, Hegetschw. Comm. Bot. t. 1. f. 8 = patens.
Tinei, *Ten. Cat. Hort. Panorm.* (1858) 26.—Hab. ?
tubiflora, *Regel, Cat. Pl. Hort. Absahor* 27.—Hab. ?
variabilis, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 169 = carnea.
variegata, *Bouché, in Linnæa*, xviii. (1844) 487.—Hab. ?
ventricosa, *Bouché, l. c.* 488.—Bras.
violacea, *Bouché, l. c.* xii. (1838) 146.—Hab. ?
vitellina, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Hab. ?
Warszewiczii, *Hort. ex Otto & Dietr. Allg. Gartenz.* xix. (1851) 289.—Am. centr.
zalebensis, Hort. Berol. ex Bouché, in Linnæa, viii. (1833) 163 = patens.

CANNABINA, Tourn. ex Medic. Phil. Bot. i. 53 (1789) = Dattisco, Caertn.
lævis, Moench, Meth. 363 = D. Cannabina.

CANNABIS, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 304 (1737). *URTICACEÆ*, Benth. & Hook. f. iii. 357.
chinensis, Delile, Ind. Sem. Hort. Monsp. (1849) = sativa.
erratica, Siev. in Pall. N. Nord. Beitr. vii. 174 = sativa.
fœtens, Gilib. Exercit. ii. 450 = sativa.
indica, Lam. Encyc. i. 695 = sativa.
Lupulus, Scop. Fl. Carn. ed. II. ii. 265 = sativa.
macrosperma, Stokes, Bot. Mat. Med. iv. 589 = sativa.
sativa, *Linn. Sp. Pl.* 1027.—As. centr.; Himal. bor. occ.

CANNACORUS, Tourn. ex Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 378 = **Canna**, Linn. (Scitamin.).
indicus, Medic. l. c. 379 = Canna indica.
ovatus, Moench, Meth. 526 = Canna indica.

CANNELLA, Schott, ex Meisn. in DC. Prod. xv. i. 168, in syn. (1864) = **Cocotes**, Aubl. (Laurin.).
Sassafras, Schott, l. c. = O. indecora.

CANNOMOIS, Beauv. ex Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 48. t. 8. f. 1. *RESTIACEÆ*, Benth. & Hook. f. iii. 1086.
CICULIFERA et MESANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 451 (1836).
Cephalotes, *Beauv. l. c.*—Afr. austr.
robustus, Hochst. in Flora, xxviii. (1845) 840 = C. Cephalotes.
scirpoides, *Hochst. l. c.*; *Mast. in Journ. Linn. Soc.* x. (1869) 236.—Afr. austr.
simplex, *Kunth, Enum. Pl.* iii. 448.—Afr. austr.
striatus, Hochst. in Flora, xxviii. (1845) 840 = simplex ?
virgatus, Hochst. l. c. = C. Cephalotes.

CANONANTHUS, G. Don, Gen. Syst. iii. 718 (1834) = Siphocampylus, Pohl (Campanulac.).
campanulatus, G. Don, l. c. 719 = S. campanulatus.

CANOPHOLIS, G. Don, Gen. Syst. iv. 324 (1837), sphalm. = Conopholis, Wallr. (Orobanch.).

CANOPUS, Presl, Epim. Bot. 248 (1849) = **Exocarpus**, Labill. (Santalac.).
laxorensis, Presl, l. c. = E. latifolius.

CANSCHI, Adans. Fam. ii. 448 (1763) = **Trewia**, Linn. (Euphorb.).

CANOTHUS, Rafin. in Med. Repos. New York, v. (1808) 360 = Ceanothus, Linn. (Rhamn.).

CANOTIA, *Torr.* in Pacif. Rail. Rep. iv. 68 (1856). *RUTACEÆ*, Benth. & Hook. f. i. 364, 816.
holacantha, *Torr. l. c.*—Calif.

CANSCORA, Lam. Encyc. i. 601 (1783). *GENTIANEÆ*, Benth. & Hook. f. ii. 811.
COMBAMBA, Blanco, Fl. Filip. 510 (1837).
FLEMINGIA, Roxb. ex Steud. Nom. ed. II. i. 643 (1840).
HETEROCANSCORA, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 431.
ORTHOTHEMON, R. Br. Prod. 451 (1810).
PHYLLOCYCLIS, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 285.
PLADERA, Soland. ex Roxb. Fl. Ind. ed. Carey, i. 400 (1832).
POOTIA, Dennst Schluess. Hort. Malab. 27 (1818).
alata, Wall. Cat. n. 4368 = perfoliata.
andrographoides, *Griff. ex C. B. Clarke, in Journ. Linn. Soc.* xiv (1875) 431.—Reg. Himal.; Burma.
concanensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 104.—Ind. or.
decurrens, *Dals. in Hook. Kew Journ.* ii. (1850) 136.—Ind. or.
decussata, *Schult. Mant.* iii. 229.—Ind. or.; Afr. trop.
diffusa, *R. Br. Prod.* 451.—Geront. trop.
diffusa, Wall. Cat. n. 4361 K = andrographioides.
divaricata, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 103 = decurrens.
foliosa, D. Don, in G. Don, Gen. Syst. iv. 199 = diffusa ?
grandiflora, Wight, Ic. iv. t. 1326 = perfoliata.
Helferiana, *Wall. ex C. B. Clarke, in Journ Linn. Soc.* xiv. (1875) 432.—Burma.
justicioides, *Griff. ex Voigt, Hort. Suburb. Calc.* 520.—Ind. or.
lancifolia, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 103 = pauciflora.
Lawii, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 430 = pauciflora.
Lawii, Wight, Ic. iv. t. 1327 = diffusa.
macrocalyx, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 104 = perfoliata.
macrocalyx, *Miq. Fl. Ind. Bat.* ii. 558.—Java.
Parishii, *Hook. f. Bot. Mag.* t. 5429.—Burma.
pauciflora, *Dals. in Hook. Kew Journ.* ii. (1850) 136.—Ind. or.
perfoliata, *Lam. Encyc.* i. 601.—Ind. or.
perfoliata, Wall. Cat. n. 4365 = Wallichii.
pusilla, Roem. & Schult. Syst. iii. 301 = Hoppea dichotoma.
Roxburghii, *Arn. ex Miq. Anal. Bot. Ind.* iii. 11.—Ind. or.
Schultesii, *Wall. Cat.* n. 4366.—Burma.
sessiliflora, *Schult. Mant.* iii. 230.—Ind. or.
tenella, Wall. Cat. n. 4362; Wight, Ic. t. 1327 = diffusa.
ventricosa, J. F. Gmel. Syst. 262 = perfoliata.
Wallichii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 103.—Ind. or.

CANSENIA, Rafin. Sylva Tellur. 122 (1838) = Bauhinia, Linn. (Legumin.).
angulosa, Rafin. l. c. = B. angulosa.
tomentosa, Rafin. l. c. = B. tomentosa.

CANSJERA, Juss. Gen. 448 (1789). *OLACINEÆ*, Benth. & Hook. f. i. 349.
OCTOPLIS, Rafin. Fl. Tellur. iv. 106 (1836).
grossibracteata, *Blanco, Fl. Filip.* ed. I. 78.—Ins. Philipp.
lanceolata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 491 = Rheedii.
leptostachya, *Benth. l. c.* ii. (1843) 231.—Austral.
madagascariensis, Spreng. Syst. i. 453 = Potamia Thouarsii.
malabarica, Lam. Encyc. iii. 462 = Rheedii.
manillana, *Blume, Mus. Bot. Lugd. Bat.* i. 246.—Ins. Philipp.

CANSJERA:—
martabanica, Wall. Cat. n. 7266 = Rheedii.
monostachya, M. Roem. Syn. Hesper. 16 = Rheedii.
parvifolia, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 298.—Burma.
pentandra, Blanco, Fl. Filip. ed. I. 78 = Antidesma Camingii.
pentandra, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 190 = Antidesma Ghesembilla.
polystachya, M. Roem. Syn. Hesper. 144 = Rheedii.
Rheedii, Blanco, Fl. Filip. ed. I. 78 = Antidesma Ghesembilla.
Rheedii, *Y. F. Gmel. Syst.* i. 280.—As. et Austral. trop.
Rheedii, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 190 = Antidesma Camingiana.
scandens, Roxb. Pl. Corom. ii. [2]. t. 103 = Rheedii.
timorensis, *Decne. Bot. Venus*, 12. t. 4.—Ins. Timor.
zeaphifolia, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 336 = Rheedii.

CANTALEA, Rafin. Sylva Tellur. 54 (1838) = Lycium, Linn. (Solanac.).

CANTHAROSPERMUM, Wight & Arn. Prod. 255 (1834) = Atylosia, Wight & Arn. (Legumin.).
albicans, Wight & Arn. l. c. 256 = A. albicans.
calycinum, Muq. Anal. Bot. Ind. i. 8.—Borneo.
distans, Royle, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 216 = A. platycarpa.
nervosum, Royle, ex Baker, l. c. = A. scarabæoides.
pauciflorum, Wight & Arn. Prod. 255 = A. scarabæoides.
scarabæoidum, Baill. in Bull. Soc. Linn. Par. i. (1883) 884 = A. scarabæoides.

CANTHIOPSIS, Seem. Fl. Vit. 166. t. 46 (1865–73) = **Bandia**, Linn. (Rubiac.).
odorata, Seem. l. c.—Ins. Fiji.

CANTHIUM, Lam. Encyc. i. 602 (1783) = **Plectronia**, Linn (Rubiac.).
acuminatum, D. Dietr. Syn. Pl. i. 770 = P. acuminata.
acutiflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 136.—Afr. trop.
Afzelianum, Hiern, l. c. 142.—Afr. trop.
angulosum, Wall. Cat. n. 8285 = Randia longiflora.
angustifolium, Roxb. Hort. Beng. 85; Fl. Ind. i. 533 = P. angustifolia.
anomocarpum, DC. Prod. iv. 476.—Guiana.
anonæfolium, Webb, in Hook. Kew Journ. ii. (1850) 370.—Ins. Cap. Verd.
approximatum, Korth. in Nederl. Kruidk. Arch. ii. 11 (1851) 284.—Borneo.
arborum, Vidal, Phan. Cuming. Philipp. 181.—Ins. Philipp.
attenuatum, R. Br. ex Benth Fl. Austral. iii. 421.—Austral.
Balansæ, Baill. Adansonia, xii. (1878) 184.—N. Caled.
barbatum, Seem. Fl. Vit. 132.—Ins. Societ.
Barteri, Hiern, in Oliver, Fl. Trop. Afr. iii. 143.—Afr. trop.
Beccheyi, Steud. Nom. ed. II. i. 275.—Ins. Societ.
Benthamianum, Baill. Adansonia, xii. (1878) 190 = Vangueria canthioides.
bibracteatum, Hiern, in Oliver, Fl. Trop. Afr. iii. 145 = P. bibracteatum.
Billardieri, D. Dietr. Syn. Pl. i. 770 = Coprosma Billardieri.
Broeni, Baill. Adansonia, xii. (1878) 221.—Madag.
buxifolium, Benth. Fl. Austral iii. 423 = P. buxifolia.
campanulatum, Thw. Enum. Pl. Zeyl. 152 = P. campanulatum.
cardiocarpum, Baill. Adansonia, xii. (1878) 185.—N. Caled.
caudatiflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 137.—Afr. trop.
celastroides, Baill. Adansonia, xii. (1878) 190 = P. celastroides.
chinense, Pers. Syn. i. 200 = Randia dumetorum.
confertum, Korth. in Nederl. Kruidk. Arch. ii. ii. (1851) 235.—Borneo.
congense, Hiern, in Oliver, Fl. Trop. Afr. iii. 141.—Afr. trop.
coprosmoides, F. Muell. Fragm. ii. 134 = *vaccinifolium*.

CANTHIUM :—
Cornelia, Cham. & Schlecht. in Linnaea, iv. (1829) 14.—Afr. trop.
corneliauum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = prec.
coromandelicum, Lam. Encyc. i. 602 = Randia dumetorum.
corymbosum, D. Dietr. Syn. Pl. i. 779 = P. ventosa, Celastrus Plectronia.
corymbosum, Pers. Syn. i. 200 = Webera corymbosa.
crassum, Hiern, in Oliver, Fl. Trop. Afr. iii. 145.—Afr. trop.
cymosum, Pers. Syn. i. 200 = P. didyma.
Deplanchei, Baill. Adansonia, xii. (1878) 183, 186 = Cyclophyllum Deplanchei.
didymum, Gaertn. f. Fruct. iii. 94. t. 196 = P. didyma.
discolor, Benth. in Hook. Niger Fl. 409.—Afr. trop.
dubium, Lindl. Bot. Reg. t. 1026 = Diplospora viridiflora.
edule, Baill. Adansonia, xii. (1878) 191 = Vangueria edulis.
ellipticum, Hochst. ex Delile, in Ferr. & Galin. Voy. Abyss. iii. 136 = Feretia spodanthera.
erythroxyloides, Baill. Adansonia, xii. (1878) 220.—Madag.
euonymoides, Baill. l. c. 189 = Vangueria euonymoides.
fasciculatum, Blume, Bijdr. 967.—Java.
floriforme, Hook. f. Fl. Brit. Ind. iii. 133.—Ind. or.
flavidum, Seem. Fl. Vit. 132.—Ins. Fiji.
floribundum, Wall. ex Voigt, Hort. Suburb. Calc. 389, nomen.—Hab.?
foetidum, Hiern, in Oliver, Fl. Trop. Afr. iii. 142.—Afr. trop.
fraternum, Miq. Fl. Ind. Bat. iii. 253.—Malaya.
furcatum, Miq. l. c. 252.—Malaya.
Gaertneri, Steud. Nom. ed. II. i. 675 = Damnacanthus indicus.
glabriflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 140.—Afr. trop.
glabrum, Blume, Bijdr. 967 = P. glabra.
glaucum, Hiern, in Oliver, Fl. Trop. Afr. iii. 134.—Afr. trop.
glomeratum, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 505 (aphalm. *glomerulatum*) = seq.
glomerulatum, Miq. Fl. Ind. Bat. Suppl. 545 = *confertum*.
gracile, Hiern, in Oliver, Fl. Trop. Afr. iii. 139.—Afr. trop.
gracilipes, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 214 = P. gracilipes.
Guersii, Sond. in Linnaea, xxiii. (1850) 54.—Natal.
Gynochthodes, Baill. Adansonia, xii. (1878) 199.—Ins. Philipp.
hebecladum, DC. Prod. iv. 474.—Ins. Luzon.
Henriettae, Baill. Adansonia, xii. (1878) 186.—N. Caled.
Heudelotii, Hiern, in Oliver, Fl. Trop. Afr. iii. 139.—Afr. trop.
hispidum, Benth. in Hook. Niger Fl. i. 409.—Afr. trop.
horizontale, Benth. l. c. 408 = *Afzelianum*.
horizontale, Hiern, in Oliver, Fl. Trop. Afr. iii. 137.—Afr. trop.
horridum, Benth. Fl. Hongk. 159 = P. parvifolia.
horridum, Blume, Bijdr. 966 = P. horrida.
indicum, D. Dietr. Syn. Pl. i. 779 = Damnacanthus indicus.
infaustum, Baill. Adansonia, xii. (1878) 191 = Vangueria infausta.
Korthalsianum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 255.—Sumatra.
laeve, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 32.—Ins. Banca.
lagenae, Baill. Adansonia, xii. (1878) 190 = Vangueria euonymoides ?
lamprophyllum, F. Muell. Fragm. ii. 133 = *lucidum*.
lanceolatum, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 842 = P. didyma.
lanciflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 146.—Afr. trop.
lasianthoides, Miq. Fl. Ind. Bat. ii. 253.—Java.
latifolium, F. Muell. ex Benth. Fl. Austral. iii. 421.—Austral.
Leschenaultii, Wight & Arn. Prod. 426 = P. angustifolia.
leucophlorum, Baill. Adansonia, xii. (1878) 191 = Fadogia leucophloea.

CANTHIUM :—
lividum, Hiern, in Oliver, Fl. Trop. Afr. iii. 144.—Afr. trop.
lucidulum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 254 = Plectronia didyma.
lucidum, R. Br. in Salt, Abyss. App. lxii.—Abyssin.
lucidum, Hook. & Arn. Bot. Beech. Voy. 65 = Beecheyi.
lucidum, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 132 = P. umbellata ?
lycioides, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 188.—Ins. Philipp.
macrocalyx, Baill. Adansonia, xii. (1878) 191 = Vangueria macrocalyx.
macrocarpum, Bolv. ex Baill. l. c. 226 = *pallens*.
macrocarpum, Thw. Enum. Pl. Zeyl. 152 = P. macrocarpa.
Mannii, Hiern, in Oliver, Fl. Trop. Afr. iii. 143.—Afr. trop.
microphyllum, F. Muell. Fragm. ii. 134 = *vaccinifolium*.
mite, Bartl. in DC. Prod. iv. 474.—Ins. Luzon.
moluccanum, Roxb. Hort. Beng. 85 ; Fl. Ind. i. 536.—Ins. Moluce.
mombazense, Baill. Adansonia, xii. (1878) 188.—Afr. trop.
monoflorum, Blanco, Fl. Filip. ed. I. 166 = *pedunculare*.
montanum, Thw. Enum Pl. Zeyl. 152.—Zeylan.
multiflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 144.—Afr. trop.
Mundtianum, Cham. & Schlecht. in Linnaea, iv. (1829) 131 = P. Mundtiana.
neglectum, Hiern, in Oliver, Fl. Trop. Afr. iii. 135.—Afr. trop.
neilgherrense, Wight, Ic. t. 1064 *bis* = P. neilgherrensis.
nervosum, Hiern, in Oliver, Fl. Trop. Afr. iii. 143.—Afr. trop.
nitens, DC. Prod. iv. 474.—Ind. or.
nitens, Hiern, in Oliver, Fl. Trop. Afr. iii. 135.—Afr. trop.
obovatum, Klotzsch, in Eckl. & Zeyh. Enum. 361.—Afr. austr.
odoratum, Seem. Fl. Vit. 132.—Ins. Fiji.
oleifolium, Hook. in Mitch. Journ. Trop. Austr. 397.—Austral.
oligocarpum, Hiern, in Oliver, Fl. Trop. Afr. iii. 138.—Afr. trop.
pallens, Baill. Adansonia, xii. (1878) 226.—Ins. Noudbe.
Pancheri, Baill. l. c. 186.—N. Caled.
parviflorum, Bartl. ex DC. Prod. iv. 474 = *hebecladum*.
parviflorum, Lam. Encyc. i. 602 = P. parviflora.
parviflorum, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 131 = Randia malabarica.
parvifolium, Baill. Adansonia, xii. (1878) 191 = Vangueria parvifolia.
parvifolium, Roxb. Hort. Beng. 15 ; Fl. Ind. i. 534 = P. parviflora.
pauciflorum, Baill. Adansonia, xii. (1878) 189 = P. pauciflora.
paucipilosum, Blanco, Fl. Filip. ed. I. 165 = P. horrida.
pedunculare, Cav. Ic. v. 21. t. 486.—Ins. Philipp.
Perrillanum, Baill. Adansonia, xii. (1878) 197.—Madag.
phyllanthoideum, Baill. l. c. 220.—Afr. trop.
polyanthum, Miq. Fl. Ind. Bat. Suppl. 545.—Sumatra.
polycarpum, Schweinf. ex Oliver, Fl. Trop. Afr. iii. 139.—Afr. trop.
puberulum, Thw. ex Hook. f. Fl. Brit. Ind. iii. 134.—Zeylan.
pyrifolium, Klotzsch, ex Eckl. & Zeyh. Enum. 361.—Afr. austr.
quadrifidum, Labill. Nov. Holl. Pl. i. 89 = Coprosma Billardieri.
racemosum, Wall. Cat. n. 6284 = Randia malabarica.
Rheedii, DC. Prod. iv. 474 = P. Rheedii.
rubens, Hiern, in Oliver, Fl. Trop. Afr. iii. 142.—Afr. trop.
ruminatum, Baill. Adansonia, xii. (1878) 200.—Afr. trop.
sagittatum, Baill. l. c. 184.—N. Caled.
scandens, Blume, Bijdr. 966.—Java.
Schimperianum, A. Rich. Tent. Fl. Abyss. i. 350 = P. Schimperi.
senegalense, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 188 = P. hirsuta.
sessiliflorum, A. Gray, in Proc. Am. Acad. iv. (1860) 39.—Ins. Fiji.

CANTHIUM :—
setiflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 134.—Afr. trop.
setosum, Hiern, l. c. 141.—Afr. trop.
spiratylum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 251.—Sumatra.
subcordatum, DC. Prod. iv. 473.—Afr. trop.
sumatranum, Miq. Fl. Ind. Bat. ii. 254.—Sumatra.
tetraphyllum, Baill. Adansonia, xii. (1878) 192 = Vangueria tetraphylla.
Thonningii, Benth. in Hook. Niger Fl. 410.—Afr. trop.
Thunbergianum, Cham. & Schlecht. in Linnaea, iv. (1829) 130 = Wendlandia thyrsoides.
thyrsoideum, Roem. & Schult. Syst. vi. 207 = Wendlandia Notoniana.
tomentosum, D. Dietr. Syn. Pl. i. 779 (= *Gynopachis tomentosa*).—Malaya.
travancoricum, Bedd. Ic. Pl. Ind. Or. t. 329 = P. travancorica.
triacanthum, Webb, in Hook. Kew Journ. ii. (1850) 371.—Ins. Cap. Verd.
umbellatum, Wight, Ic. t. 1084 ; Spicil. Neilgh. i. 81. t. 97 = P. umbellata.
umbelligerum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 252.—Borneo.
umbellulatum, Korth. in Nederl. Kruidk. Arch. ii. ii. (1851) 235.—Borneo.
undulatum, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 199 = P. didyma.
vaccinifolium, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 47.—Austral.
vanguerioides, Hiern, in Oliver, Fl. Trop. Afr. iii. 146.—Afr. trop.
Vatkeanum, Hiern, l. c. 136.—Afr. trop.
venosum, Hiern, l. c. 144.—Afr. trop.
vensulum, Bolv. ex Baill. Adansonia, xii. (1878) 221.—Ins. Comor.
Viridiardi, Baill. l. c. 185.—N. Caled.
Villarii, Vidal, Phan. Cuming. Philipp. 182.—Ins. Philipp.
violaceum, Zoll. & Moritzi, Syst. Verz. Zoll. 64.—Java.
viridiflorum, D. Dietr. Syn. Pl. i. 780 = Diplospora viridiflora.
zanzibaricum, Klotzsch, in Peters, Reise Mossamb. Bot. 291 = P. zanzibarica.
zizyphinum, Wall. Cat. n. 6288 A, C = P. horrida.

CANTHOPSIS, Miq. Fl. Ind. Bat. ii. 256 (1856). *RUBIACEAE*, Benth. & Hook. f. ii. 113. pubiflora, *Miq. l. c.* 257.—Ins. Timor.

CANTUA, Juss. Gen. 136 (1789). *POLEMONIACEAE*, Benth. & Hook. f. ii. 824.
PERIPHRAGMOS, Ruiz & Pav. Prod. 28. t. 24 (1794).
aggregata, Pursh, Fl. Am. Sept. i. 147 = Gilia aggregata.
bicolor, *Lem. in Fl. des Serres*, Sér. I. iii. (1847) 242, verso.—Bolivia.
brevifolia, Juss. in Ann. Mus. Par. iii. (1804) 119 = Gilia laciniata.
brevifolia, Steud. Nom. ed. II. i. 275 = prec.
buxifolia, *Lam. Encyc.* i. 603.—Peruv.
caerulea, Poir. Encyc. Suppl. ii. 80 = Loeselia caerulea.
coccinea, Poir. l. c. = Loeselia coccinea.
cordata, *Juss. in Ann. Mus. Par.* iii. (1804) 119.—Peruv.
coronopifolia, Willd. Sp. Pl. i. 879 = Gilia coronopifolia.
cunelifolia, *Juss. ex Roem. & Schult. Syst.* iv. 269.—Hab.?
dependens, Pers. Syn. i. 187 = buxifolia.
elegans, Poir. Encyc. Suppl. ii. 80 = Gilia coronopifolia.
fasciculata, Humb. ex Roem. & Schult. Syst. iv. 269 = Fouquiera speciosa.
flexuosa, Pers. Syn. i. 187 = pyrifolia.
floridana, Nutt. ex A. Gray, Syn. Fl. N. Am. ii. 1. 145 = Gilia coronopifolia.
foetida, Pers. Syn. i. 187 = Ventia lycioides.
glandulosa, Poir. Encyc. Suppl. ii. 80 = Loeselia glandulosa.
glomerifiora, Juss. in Ann. Mus. Par. iii. (1804) 119 = Cellonia Cavanillesiana.
glutinosa, *Presl, Epim. Bot.* 218.—Chili.
Hottaia, Willd. Sp. Pl. i. 878 = Loeselia coccinea.
incisifolia, Poir. Encyc. Suppl. ii. 83 = Gilia laciniata.
ligustrifolia, Juss. in Ann. Mus. Par. iii. (1804) 116 = Ventia lycioides.

CANTUA:—
longiflora, Torr. in Ann. Lyc. N. York, iii. (1828) 221 = Collomia longiflora.
laxensis, Willd. ex Roem. & Schult. Syst. iv. 389 = pyrifolia.
megapotamica, *Spreng. Syst.* iv. *Cur. Post.* 64 (quid!). —Bras.
ovata, *Cav. Ic.* iv. 48.—Peruv.
parviflora, Pursh, Fl. Am. Sept. ii. 730 = Gilia parviflora.
peruviana, j. F. Gmel. Syst. 347 = pyrifolia.
pinnatifida, Lam. Illustr. i. 473 = Gilia coronopifolia.
pungens, Torr. in Ann. Lyc. N. York. ii. (1826) 221 = Gilia pungens.
pyrifolia, *Juss. in Ann. Mus. Par.* iii. (1804) 117.—Peruv.
quercifolia, *Juss. l. c.* 118.—Peruv.
thesefolia, *D. Don, in Edinb. Phil. Journ.* (Apl.-Oct. 1832) 289.—Peruv.
thyrsoidea, Juss. in Ann. Mus. Par. iii. (1804) 119 = Gilia coronopifolia.
tomentosa, Cav. Ic. iv. 48 = buxifolia.
tuberosa, Roem. & Schult. Syst. iv. 793 = Ipomoea muricata.
umbiflora, Pers. Syn. i. 187 = ovata.

CANTUFFA, j. F. Gmel. Syst. 677 (1791) = **Pterolobium**, R. Br. (Legumin.).
exosa, j. F. Gmel. l. c. = P. lacerans.

CAOUTCHOUA, j. F. Gmel. Syst. 1007 (1791) = **Hevea**, Aubl. (Euphorbiac.).
elastica, j. F. Gmel. l. c. = H. guianensis.

CAPANEA, Decne. in Van Houtte, Fl. des Serres, t. 499 (1850) = **Campanea**, Decne. (Generac.).

CAPANEMIA, Rodrig. Orch. Nov. i. 137 (1877). *ORCHIDEAE.*
carinata, *Rodrig. l. c.* ii. 243.—Bras.
micromera, *Rodrig. l. c.* i. 138.—Bras.
Theresiae, *Rodrig. l. c.* ii. 244.—Bras.
uliginosa, *Rodrig. l. c.* i. 137.—Bras.

CAPASSA, Klotzsch, in Peters Reise Mossamb. Bot. 27. t. 5 (1861) = **Lonchocarpus**, Humb. & Bonpl. (Legumin.).
violacea, Klotzsch, l. c. 26 = L. violaceus.

CAPELLENIA, Hassk. Cat. Hort. Bog. Alt. 178 (1844) = CAPELLIA, Blume = **Wormia**, Rottb. (Dilleniac.).

CAPELLENIA, Teijsm. & Binn. in Tijdschr., Nederl. Ind. xxix. (1866) 239 = **Endosperm**um, Benth. (Euphorb.).
malaccana, Teijsm. & Binn. l. c.—Ins. Molucc.

CAPELLIA, Blume, Bijdr. 5 (1825) = **Wormia**, Rottb. (Dilleniac.).
biflora, A. Gray, Bot. U. St. Expl. Exped. i. 15. t. 1 = W. biflora.
membranifolia, A. Gray, l. c. 17 = W. membranifolia.
multiflora, Blume, Bijdr. 5.—Java.
paucifiora, Zoll. & Moritzi, Syst. Verz. Zoll. 35.—Java.

CAPERONIA, St. Hil. Pl. Rem. Bras. 244 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 304.
ANDROPHORANTHUS, Karst. in C. Koch & Fint. Wochenschr. (1859) 5.
CAVANILLA, Vell. Fl. Flum. 298 (1825).
LEPIDOCOCCA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 588.
MITERANA, Rafin. Sylva Tellur. 65 (1838).
acalyphoïdia, *Griseb. in Goett. Abh.* xviv. (1879) 58. —Reg. Argent.
aculeolata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 152.—Bras.
angustissima, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 96.—Guiana.
baklousii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 235.—Bras.
buettneriana, *Muell. Arg, l. c.* 230.—Bras.
castaneaefolia, Miq. in Linnaea, xxi. (1848) 478 = corchorifolia.
castaneaefolia, *St. Hil. Pl. Rem. Bras.* 245.—Java.
corchorifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 153.—Guiana.
cordata, *St. Hil. Pl. Rem. Bras.* 245.—Bras.

CAPERONIA:—
Gardneri, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 221. —Bras.
heteropetala, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1857) 148.—Bras.
heteropetaloides, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 152.—Bras.
Langsdorffii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 219.—Bras.
linearifolia, *St. Hil. Pl. Rem. Bras.* 246.—Bras.
malticostata, *Muell. Arg. in Mart. Fl. Bras.* xi. ii 228.—Bras.
nervosa, A. Rich. Fl. Cub. Fanerog. ii. 213 = castaneaefolia.
paludosa, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 51.—Mexic.
palustris, *St. Hil. Pl. Rem. Bras.* 245.—Ind. occ.; Mexic.
palustris, H. B. & K. Nov. Gen. et Sp. ii. 71 = castaneaefolia.
paraensis, Klotzsch, in Seem. Bot. Voy. Herald, 108 = paludosa.
Regnellii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 221.—Bras.
Rutenbergii, *Muell. Arg. in Bremen Abh.* vii. (1880) 25.—Madag.
senegalensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 153.—Afr. trop.
serrata, Baill. in Adansonia, i. (1860–61) 66 = senegalensis.
serrata, *Presl, Epim. Bot.* 213.—Afr. trop.
stenophylla, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 226. —Bras.
Velloziana, *Muell. Arg. l. c.* 223.—Bras.

CAPIA, Domb. ex Juss. in Dict. Sc. Nat. vi. 500 (1805) = **Lapageria**, Ruiz & Pav. (Liliac.).

CAPIRONA, Spruce, in Journ. Linn. Soc. iii. (1859) 200. *RUBIACEAE*, Benth. & Hook. f. iii. 39.
Boiviniana, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 270.—Madag.
decorticans, *Spruce, in Journ. Linn. Soc.* iii. (1859) 200.—Peruv.

CAPITELLARIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 109 = **Sagraea**, DC. (Melastom.).
Benthami, Naud. l. c. = S. capitata.

CAPNITES, Dum. Fl. Belg. 117 (1827) (err. typ. 177) = **Corydalis**, Vent. (Papaverac.).
alpestris, Rupr. Fl. Caucas. 59 = Cor. pauciflora.
angustifolia, Rupr. l. c. 57 = Cor. angustifolia.
araratica, Rupr. l. c. 61.—Armenia.
Bayeriana, Rupr. l. c. 59. t. 4 = Cor. pallida.
caucasica, Rupr. l. c. 56 = Cor. caucasica.
cava, Dum. Fl. Belg. 117 (err. typ. 177) = Cor. tuberosa, bulbosa.
conorhiza, 'Ledeb.' ex Rupr. Fl. Caucas. 61 = Cor. conorhiza.
digitata, Dum. Fl. Belg. 117 (err. typ 177) = Cor. solida.
Emanueli, Rupr. Fl. Caucas. 58 = Cor. pauciflora.
fabacea, Dum. Fl. Belg. 117 (err. typ. 177) = Cor. fabacea.
macrosepala, Rupr. Fl. Caucas. 61.—Reg. Caucas.
Marschalliana, Rupr. l. c. 55 = Cor. Marschalliana.
ochroleuca, Rupr. l. c. 61 = Cor. lutea.
pallidiflora, Rupr. l. c. 58 = Cor. pallida.
roseo-purpurea, Rupr. l. c. 288.—Reg. Caucas.
solida, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 299 = Cor. solida.

CAPNITIS, E. Mey. Comm. Pl. Afr. Austr. 81 (1835) = **Lotononis**, Eckl. & Zeyh. (Legumin.).
clandestina, E. Mey. l. c. = L. clandestina.
porrecta, E. Mey. l. c. = L. porrecta.

CAPNOCYSTIS, Juss. in Ann. Mus. Par. xviii. (1811) 478 = CYSTICAPNOS, Boerh. = **Corydalis**, Vent. (Papaverac.).

CAPNOGONIUM, Bernh. Ind. Sem. Hort. Erfurt. (1844) ex Endl. Gen. Suppl. ii. 107 = **Corydalis**, Vent. (Papaverac.).
nobile, Bernh. l. c. = Cor. nobilis.

CAPNOIDES, Tourn. ex Adans. Fam. ii. 431 (1763) = **Corydalis**, Vent. (Papaverac.).
albida, Bernh. ex Steud. Nom. ed. II. i. 276 = Cor. lutea.
cava, Moench, Meth. 52 = Cor. tuberosa.
glauca, Moench, l. c. = Cor. glauca.
lutea, Gaertn. Fruct. ii. 153 = Cor. lutea.
minor, Schum. ex DC. Syst. ii. 119 = Cor. fabacea.
nobilis, Moench, Meth. 52 = Cor. nobilis.
rigidula, Bernh. ex Steud. Nom. ed. II. i. 276 = Cor. ochroleuca.
scandens, Moench, Meth. Suppl. 215 = Adlumia cirrhosa.
*semperviren*s, Borckh. in Roem. Arch. i. ii. 44 = Cor. glauca.
solida, Moench, Meth. 52 = Cor. tuberosa.

CAPNOPHYLLUM, Gaertn. Fruct. ii. 32. t. 85 (1791). *UMBELLIFERAE*, Benth. & Hook. f. i. 908.
ACTINOCLADUS, E. Mey. Ind. Sem. Hort. Regiom. (1846).
KRUBERA, Hoffm. Gen. Umb. 108 (1814).
SCLEROCIADIUM, Koch, in DC. Coll. Mem. v. 48 (1829); Prod. iv. 140 (1830).
ULOSPERMUM, Link, Enum. Hort. Berol. i. 267 (1822).
africanum, *Gaertn. Fruct.* ii. 32. t. 85.—Afr. austr.
dichotomum, *Lag. Gen. et Sp. Nov.* 13.—Reg. Mediterr.
Jacquini, *DC. Prod.* iv. 187.—Afr. austr.
peregrinum, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 33 = dichotomum.
simplex, Lag. Gen. et Sp. Nov. 13 = africanum.

CAPNORCHIS, Borck. in Roem. Arch. i. ii. 46 (1797) = **Dicentra**, Bernh. (Papaverac.).
chrysantha, Planch. in Fl. des Serres, Sér. I. viii. (1852–53) 193. t. 820 = D. chrysantha.
Cucullaria, Planch. l. c. 193 = D. Cucullaria.
eximia, Planch. l. c. = D. eximia.
formosa, Planch. l. c. = D. formosa.
spectabilis, Borck. in Roem. Arch. i. ii. 46 = D. spectabilis.

CAPNOREA, Rafin. Fl. Tellur. iii. 74 (1836) = **Nicotiana**, Linn. (Solanac.).
nana, Rafin. l. c. 75 = N. nana.

CAPPARIS, [Tourn.] Linn. Syst. ed. I (1735). *CAPPARIDEAE*, Benth. & Hook. f. i. 108.
BEAUTEMPSIA, Gaudich. Bot. Voy. Bonite, t. 56 (1844).
BUSBECKIA, Endl. Prod. Fl. Ins. Norf. 64 (1838).
CALYPTRANTHUS, Thou. Obs. Pl. Isles Afr. 26 (1811).
COLICOSPERMUM, Mart. Herb. Fl. Bras. 201 (1837).
DESTRUGUEZIA, Gaudich. Bot. Voy. Bonite, t. 57 (1844).
HOMBAX, Adans. Fam. ii. 408 (1763).
INTUITS, MARSESINA, OCTANEMA, OLIGOLORON, OLOFUTON, PLEUTRON, TRICLANTHERA, Rafin. Sylva Tellur. 108–112 (1838).
LINDACKERA, Sieber, ex Endl. Gen. 893, in syn. (1841).
PETERSIA, Klotzsch, in Peters, Reise Mossamb. Bot. 188. t. 80 (1861).
QUADRELLA, Meisn. Gen. 15 (1867).
SODADA, Forsk. Fl. Aegypt. Arab. 81 (1775).
UTERVERIA, Bertol. Pl. Nov. Hort. Bonon. ii. 8 (1839).
aculeata, Steud. Nom. ed. I. 147 = spinosa.
acuminata, Lindl. Bot. Reg. t. 1230 = acutifolia.
acuminata, Roxb. Hort. Beng. 41; Fl. Ind. ii. 566 = zeylanica.
acuminata, *Willd. Sp. Pl.* ii. 1131.—Ind. or.
acuminata, Willd. ex Wall. Cat. n. 6990 C = horrida.
acutifolia, *Sweet, Hort. Brit. ed.* II. 525.—China.
aegyptia, Lam. Encyc. i. 605 = spinosa.
aegyptia, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 374 = Pontanesii.
Afzelii, DC. Prod. i. 246 = erythrocarpos.
Agaba, Banks, ex DC. Prod i. 246 = Roxburghii.
albitronea, *Burch. Trav.* i. 248.—Afr. austr.
ambigua, *Kurz, For. Fl. Brit. Burma*, i. 65 — Ins. Andam.
amplissima, *Lam. Encyc.* i. 607.—Ind. occ.; Mexic.
amygdalifolia, *Vacq. Enum. Pl. Cuvit.* 24.—Ind. occ.
amygdalina, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 155 = Griesebachii.

CAPPARIS:—
amygdalina, Lam. Encyc. i. 606 = Capparis Breynia.
angulata, *Ruiz & Pav. Fl. Per.* t. 481 *ex DC. Prod.* i. 253.—Ecuador.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* v. 96. t. 438. —Mexic.
anisacanthera, *Bull. Soc. Linn. Par.* i. (1885) 462.—Madag.
apetala, Roth, Sp. Pl. Nov. 238 = Niebuhria linearis.
aphylla, *Roth, l. c.*—Afr. bor.; Arab.; Ind. or.
arborescens, Mill. Gard. Dict. ed. VIII. n. 8 = pulcherrima.
Arrabidæa, *Steud. Nom.* ed. II. i. 276.—Bras.
artensis, *Mentr. in Mém. Acad. Lyon*, x. (1860) 177.—N. Caled.
asperifolia, *Presl, Rel. Haenk.* ii. 86.—Mexic.
assamica, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 177. —Assam.
aurantioides, *Presl, Rel. Haenk.* ii. 86.—Ins. Luzon.
avicennifolia, *H. B. & K. Nov. Gen. et Sp.* v. 94.—Am. austr.
Baducca, Blanco, Fl. Filip. ed. I. 438 = spinosa.
Baducca, Linn. Sp. Pl. 504 = zeylanica ?
barcellensis, H. B. & K. Nov. Gen. et Sp. v. 92 = Breynia.
Billardieri, *DC. Prod.* i. 247.—Ins. Molucc.
bispernna, Roxb. Hort. Beng. [98]; Fl. Ind. ii. 568 = grandis.
bispermo, Wight, ex Hook. f. Fl. Brit. Ind. i. 176 = Roxburghii.
brachycarpa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 323.—Java.
brasiliana, *DC. Prod.* i. 249.—Bras.
Brasiâ, *DC. l. c.* 248.—Am. austr.
brevipes, Benth. Bot. Voy. Sulph. 65 = verrucosa.
brevis, Spreng. Fl Hal. Mant. 43 = eustachiana.
brevistiliqua, *Moç. & Sesse, ex DC. Prod.* i. 251.—Mexic.
brevispina, DC. Prod. i. 246 = zeylanica.
brevispina, Wight & Arn. Prod. 24 = divaricata.
Breynia, H. B. & K. Nov. Gen. et Sp. v. 97 = odoratissima.
Breynia, *Linn. Syst.* ed. X. 1071; *Yacq. Am. Select.* 161. t. 103.—Am. austr.
caerulea, Heyne, ex Wall. Cat. n. 6986 = diversifolia.
callophylla, *Blume, Bijdr.* 58.—Java.
callosa, Blume, l. c. = micracantha.
canescens, *Banks, ex DC. Prod.* i. 246.—Austral.
canescens, G. Don, Gen. Syst. i. 281 = Dodsæa.
cantoniensis, *Lour. Fl. Cochinch.* 830.—China.
capensis, Thunb. Prod. Fl. Cap. 92 = citrifolia.
Carandas, *Burm. f. Fl. Ind.* 118.—Ind. or.
cardiophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 325.—Venezuela.
cartilaginea, Decne. in Ann. Sc. Nat. Sér. II. iii. (1885) 373 = galeata.
celebica, *Miq. Illustr. Fl. Arch. Ind.* 26.—Ins. Celebes.
cerasifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 71.—Archip. Sulu.
chinensis, G. Don, Gen. Syst. i. 278 = acutifolia.
chrysomela, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 59.—Madag.
citrifolia, *Lam. Encyc.* i. 606.—Afr. austr.
clutiæfolia, *Burch. ex DC. Prod.* i. 248.—Afr. austr.
coconolobifolia, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. t. 284.—Am. trop.
coloratoides, Boiss. Diagn. Ser. I. xiii. 88 = Cleome coluteoides.
commutata, Spreng. Neue Entdeck. iii. 97 = frondosa.
comosa, *Yacq. Enum. Pl. Carib.* 23.—Ins. S. Doming.
conferta, Mill. Gard. Dict. ed. VIII. n. 8 = laurifolia.
conspicua, Wall. Cat. n. 6991 = micracantha.
cordata, Ruiz & Pav. Fl. Per. t. 432; *ex DC. Prod.* i. 251 = crotonoides.
cordifolia, Lam. Encyc. i. 609 = spinosa.
corlacea, *Burch. ex DC. Prod.* i. 248.—Afr. austr.
corymbifera, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 62. —Afr. austr.
corymbosa, *Lam. Encyc.* i. 605.—Afr. trop.
corymbosa, Roxb. Hort. Beng. [98]; Fl. Ind. ii. 569 = Roxburghii.
crassifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 71.—Archip. Sulu.
crassifolia, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 237.—Burma.
crotonoides, *H. B. & K. Nov. Gen. et Sp.* v. 95. t. 437. —N. Granat.
cumeata, DC. Prod. i. 249 = frondosa.
Cynophallophora, Linn. Sp. Pl. ed. I. 504 = jamaicensis.

CAPPARIS:—
Cynophallophora, *Linn. Sp. Pl.* ed. II. 723; *Jacq. Am. Select.* t. 98.—Am. austr.
Dahl, *Forsk. Fl. Aegypt. Arab.* 212 (Zygophylli sp. ?). —Arab.
daceyzetas, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 322.—Java.
dealbata, *DC. Prod.* i. 246.—Ins. Timor.
decidua, Edgew. in Journ. Linn. Soc. vi. (1862) 184 = aphylla.
declinata, Vell. Fl. Flum. 230; v. t. 111 = C. Cynophallophora.
detonsa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 80.—N. Granat.
didymobotrys, *Ruiz & Pav. Fl. Per.* t. 423; *ex DC. Prod.* i. 253.—Ecuador.
dispersa, Walp. Rep. i. 199 = grandis.
disticha, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 70.—Burma.
divaricata, Lam. Encyc. i. 606. ; Klein, ex Wall. Cat. n. 6988 *A* = diversifolia.
diversifolia, *Wight & Arn. Prod.* 27.—Ind. or.
domingensis, *Spreng. ex DC. Prod.* i. 253.—Ins. S. Doming.
Donians, *D. Dietr. Syn. Pl.* iii. 227.—Afr. trop.
Eichleriana, *Urban, in Linnæa*, xliii. (1880–82) 265. —Bras.
elæagnifolia, *Jacq. Enum. Pl. Carib.* 23.—Ind. occ.
elegans, *Mart. in Flora*, xxii (1839) i. *Beibl.* 24.—Bras
elliptica, Span. in Linnæa, xv. (1841) 166 = Roxburghii ?
emarginata, *Presl, Reliq. Haenk.* ii. 85.—Ins. Luzon.
emarginata, A. Rich. Ess. Fl. Cub. 78 = jamaicensis.
emarginata, Zipp. ex Span. in Linnæa, xv. (1841) 166 = sepiaria.
erythrocarpos, *Isert, in Ges. Naturf. Fr. Berl. Schrift.* ix. (1789) 334. t. 9.—Guiana.
erythrocarpos, Moç. Fl. Jungh. 257 = horrida.
eustachiana, *Jacq. Enum. Pl. Carib.* 23.—Ins. S. Eustach.
exstipulata, *J. F. Gmel. Syst.* 807 = inermis ?
falcata, Lour. Fl. Cochinch. 330 = Crataeva falcata.
fascicularis, DC. Prod. i. 248 = corymbosa.
ferruginea, *Linn. Syst* ed X. 1071.—Ind. occ.
ferruginea, Sieber, ex Eichl. in Mart. Fl. Bras. xiii. t. 271 = C. Breynia.
ferruginea, Willd. ex Eichl. l. c. = odoratissima.
Finlaysoniana, *Wall. Cat.* n. 6992 B.—Malaya.
Finlaysoniana, Wall. l. c. *A* = micracantha.
flaviana, *Wall. l. c.* n. 7003.—Burma.
flexicaulis, *Hance, in Journ. Bot.* vi. (1878) 235.—China.
flexuosa, *Blume, ex Hassk. Pl. Jav.-Rar.* 179.—Java.
flexuosa, Linn. Sp. Pl. ed. II. 722 = C. Cynophallophora.
flexuosa, Vell. Fl. Flum. 230; v. t. 108 = Arrabidæa.
floribunda, Lepr. ex Walp. Rep. i. 197 = tomentosa.
floribunda, *Wight, Illustr. Ind. Bot.* i. 35. t. 14.—Ind. or.; Ins. Philipp.
foetida, *Blume, Bijdr.* 52.—Java.
Fontanesii, *DC. Prod.* i. 245.—Ins. Maurit.
formosa, Dalz. in Hook. Kew Journ. ii. (1850) 40 = Heyneana.
formosa, Wall. Cat. n. 6982 *A* = horrida.
Forsteniana, *Miq. Illustr. Fl. Arch. Ind.* 32.—Malaya.
frondosa, *Yacq. Enum. Pl. Carib.* 24; *Linn. Sp. Pl.* ed. II. 720. 1674.—Am. austr.
fruticosa, *Mill. Gard. Dict.* ed. VIII. n. 7.—Am. austr.
furfuracea, *Ruiz & Pav. ex DC. Prod.* i. 252.—Mexic.
galeata, *Presen. in Mus. Senchenb.* ii. (1837) 111.—Aegypt.
Gaudichaudiana, *Eichl. in Mart. Fl. Bras.* xiii. t. 273. —Peruv.
glandulosa, *Desf. Cat. Hort. Par.* ed. III. 405.—Hab. ?
glauca, *Wall. Cat.* n. 7008.—Burma.
globifera, *Delile, in Rochet, Sec. Voy. Choa*, 240.—Afr. trop.
Gomesiana, Steud. Nom. ed. II. i. 277 = pumila.
Granddidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 463.—Madag.
grandiflora, *Wall. Cat.* n. 6984.—Ind. or.
grandis, *Linn. f. Suppl.* 268.—Ind. or.; Burma.
grandis, Rusa, ex Wall. Cat. n. 7001 = Russeliana.
Griesbachii, *Eichl. in Mart. Fl. Bras.* xiii. t. 275.—Cuba.
guayaquilensis, H. B. & K. Nov. Gen. et Sp. v. 89 = G. Cynophallophora.
Gscneeli, *Sond. in Harv. & Sond. Fl. Cap.* i. 62.—Afr. austr.

CAPPARIS:—
Hasseliana, *Miq. Illustr. Fl. Arch. Ind.* 24. t. 13.—Java.
hastata, Jacq. Enum. Pl. Carib. 23 = C. Cynophallophora.
hastigera, Hance, in Journ. Bot. vi. (1868) 296 = Swinhoii.
herbacea, Willd. Enum. Hort. Berol. 560 = spinosa.
hetercarantha, *Du. Prod.* i. 246.—Mesopotam.
heterolitis, Roxb. Hort. Beng. 41; Fl. Ind. ii. 570 = Maerua arenaria.
heterogenea, Roxb. ex Steud. Nom. ed. II. i. 277 = Maerua oblongifolia.
heterophylla, *Ruiz & Pav. ex DC. Prod.* i. 250.—Ecuador.
Heyneana, *Wall. Cat.* n. 6985.—Ind. or.
horrida, Banks, ex Wight & Arn. Prod. 25 = stylosa.
horrida, *Linn. f. Suppl.* 264.—As. trop.; Malaya. Humblotii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 463. —Madag.
humistrata, *F. Muell. Fragm.* v. 156.—Austral.
hypericoides, *Hochst. in Flora*, xxvii. (1844) i. 290.—Afr. occ.
hypoleuca, *Presl, Rel. Haenk.* ii. 87.—Peruv.
incana, *H. B. & K. Nov. Gen. et Sp.* v. 94.—Am austr.
incanescens, DC. Prod. i. 247 = sepiaria.
inermis, *Forsk. Fl. Aegypt. Arab.* 100.—Arab.
intermedia, H. B. & K. Nov. Gen. et Sp. v. 96 = odoratissima.
ishmensis, *Eichl. in Mart. Fl. Bras.* xiii. t. 269.—Am trop.
Jacobinæ, *Moric. ex Eichl. l. c.* 277.—Bras.
jamaicensis, *Yacq. Enum. Pl. Carib.* 23.—Jamaica.
Karwinskiana, *Schlecht. in Linnæa*, x. (1836) 237.—Mexic.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 98.—Afr. trop.
Korthalsiana, *Miq. Illustr. Fl. Arch. Ind.* 31.—Borneo.
laeta, H. B. & K. Nov. Gen. et Sp. v. 88 = verrucosa.
lasteviroi, Mart. in Flora (1839) i. Beibl. 23 = C. Cynophallophora.
laevigata, Mart. l. c. 21 = C. Cynophallophora.
lanceolaris, *Lesch. ex Steud. Nom.* ed. II. i. 277.—Java.
lanceolata, Ruiz & Pav. Fl. Per. t. 433; ex DC. Prod. i. 249 = C. Cynophallophora.
lasiantha, *R. Br. ex DC. Prod.* i. 247.—Austral.
lasiopoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 322.—Ins. Luzon.
laurifolia, *Mill. Gard. Dict.* ed. VIII. errat.—N. Granat
laurina, *H. B. & K. Nov. Gen. et Sp.* v. 93.—Am. austr.
leucophylla, DC. Prod. i. 246 = spinosa.
linearifolia, Hook. f. in Hook. Niger Fl. 317 = Thonningii.
linearifolia, *Linden, ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 324.—Venezuela.
linearis, Blanco, Fl. Filip. ed. I. 438 = viminea.
linearis, *Yacq. Enum. Pl. Carib.* 24.—Am. austr.
linearis, *Pers. Syn.* ii. 90.—Bras.
linifolia, Roxb. Hort. Beng. 41 = Niebuhria linearis.
Lobbiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 323.—Singapore.
Loddigesii, *Steud. Nom.* ed. II. i. 277.—Hab. ?
longifolia, *Sw. Prod. Veg. Ind. Occ.* 91.—Ind. occ.
longispina, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 176 —Burma.
lorsanthifolia, *Lindl. in Mitch. Trop. Austral.* 220.—Austral.
lucida, *Desv. ex DC. Prod.* i. 254.—Austral.
lusonensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 324.—Ins. Luzon.
macrocarpa, Ruiz & Pav. ex DC. Prod. i. 250 = petiolaris.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* v. 91.—N. Granat.
macrosperma, *Delile, in Rochet, Sec. Voy. Choa*, 244. —Afr. trop.
magna, Lour. Fl. Cochinch. 330 = Crataeva magna.
maidaeria, Wight, ex Steud. Nom. ed. II. i. 277 = Heyneana.
malabarica, Wall. Cat. n. 7000 = floribunda.
mariana, Jacq. Hort. Schoenbr. i. 57. t. 109 = spinosa.
maxima, Roth, Sp. Pl. Nov. 237 = grandis.
membrenacea, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 241.—China.
monbrasifolia, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 70.—Burma.
micracantha, *DC. Prod.* i. 247.—Burma; Malaya.

CAPPARIS:—
micracantha, Teijsm. ex Miq. Illustr. Fl. Arch. Ind. 88 = Forsteniana.
micrantha, *A. Rich. Tent. Fl. Abyss.* i. 81.—Abyss.
Mitchellii, *Lindl. in Mitch. Three Exped.* i. 316.—Austral.
mitchidatica, *Forsk. Fl. Aegypt. Arab.* 99.—Arabia.
mollis, *H. B. & K. Nov. Gen. et Sp.* v. 88.—Peruv.
mollisacula, Wall Cat. n. 7006 (= Cansjerae sp.).—Ind. or.
Moonii, *Wight, Illustr. Ind. Bot.* i. 85.—Zeylan.
Morisonia, Sw. Obs. 273 = Morisonia americana.
mucronifolia, Boiss. Diagn. Ser. I. i. 8 = spinosa.
multiflora, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 178. —Reg. Himal.; Burma.
Murrayana, J. Grah. in Cat. Pl. Bomb. 9 = spinosa.
Murrayi, Stew. ex Dalz. in Journ. Linn. Soc. xiii. (1873) 74 = spinosa ?
nectaria, *Veil. Fl. Flum.* 330; v. t. 107.—Bras.
negrorosa, *Blanco, Fl. Filip.* ed. I. 438.—Ins. Philipp.
nemorosa, *Jacq. Enum. Pl. Carib.* 24.—N. Granat.
nepaulensis, DC. Prod. i. 246; Wall. Cat. n. 6979 = spinosa.
nerilloila, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* xiv. (1884) 180.—Hab. ?
nigricans, *Span. in Linnaea*, xv. (1841) 165.—Ins. Timor.
nitida, *Ruiz & Pav. ex DC. Prod.* i. 252.—Peruv.
nobilis, *F. Muell. ex Benth. Fl. Austral.* i. 95.—Austral.
noctiflora, Spreng. ex Eichl. in Mart. Fl. Bras. xiii. 1. 285 = nitida.
nummularia, *DC. Prod.* i. 246.—Austral.
oblongifolia, *Forsk. Fl. Aegypt. Arab.* 99.—Arabia.
obovata, Buch.-Ham. ex DC. Prod. i. 248 = grandis.
obovata, Royle, Illustr. Bot. Himal. 73 = spinosa.
obovatifolia, H. B. & K. Nov. Gen. et Sp. v. 92 = tenuisiliqua.
obtusa, Rafin. Sylva Tellur. 109 = C. Cynophallophora.
octandra, Jacq. Am. Select. 180. t. 100 = ferruginea.
odontia, *Blanco, Fl. Filip.* ed. I. 439.—Ins. Philipp.
odoratissima, *Jacq. Hort. Schoenb.* t. 110.—Venezuela.
oleifolia, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 178. —Reg. Himal.
oleoides, *Burch. ex DC. Prod.* i. 248.—Afr. austr.
oligandra, Griff. Notul. iv. 577 = floribunda.
orbiculata, *Wall. ex Hook. f. & Thoms. in Fl. Brit. Ind.* i. 176.—Burma.
ornans, *F. Muell. ex Benth. Fl. Austral.* i. 95.—Austral.
ovalifolia, Ruiz & Pav. Fl. Per. t. 422, ined. ex DC. Prod. i. 258 = sviceminifolia.
ovalis, Risso, Fl. Nice, 44 = spinosa.
ovata, Bieb. Fl. Taur. Cauc. ii. 1 = spinosa.
ovata, Desf. Fl. Atlant. i. 404 = Fontanesii.
oxyphylla, *Miq. Pl. Jungh.* 397.—Java.
oxyphylla, Wall. Cat. n. 6997 = tenera.
oxypepala, *Wright, ex Radlk. in Sitzb. Math.-Phys. Acad. Muench.* xiv. (1884) 173.—Nicarag.
Pachaca, *H. B. & K. Nov. Gen. et Sp.* v. 93.—Venezuela.
pandurifformis, Lam. Encyc. i. 609 = Thylachium pandurifforme.
pandurifformis, Thou. Obs. Afr. Austr. 26 = Thylachium heterophyllum.
paradoxa, Jacq. Hort. Schoenb. i. 58 = Steriphoma paradoxum.
parvifiora, Boiss. Diagn. Ser. I. i. 4 = spinosa.
parvifiora, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 176. —Ind. or.
pauciflora, H. B. & K. Nov. Gen. et Sp. v. 89 = C. Cynophallophora.
pauciflora, Presl, Rel. Haenk. ii. 86 = inconspicua.
pedunculata, *C. & Y. Presl, Delic. Prag.* i. 90.—Sicilia.
pedunculosa, *Wall. Cat.* n. 6999.—Ind. or.
pelitata, G. Don, Gen. Syst. i. 835.—Ins. Trinitat.
pendula, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 76.—N. Granat.
persicifolia, A. Rich. Tent. Fl. Abyss. i. 81 = tomentosa
petiolaris, Benth. ex DC. Prod. i. 249 = frondosa.
petiolata, *H. B. & K. Nov. Gen. et Sp.* v. 91.—Guayaquil.
platyacantha, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 393.—Java.
Plumbella, Blanco, Fl. Filip. ed. 2. 230 = C. Cynophallophora.

CAPPARIS:—
polyantha, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 76.—N. Granat.
polymorpha, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 227.—Burma.
polymorpha, Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 34 = tomentosa.
populifolia, Willd. ex Eichl. in Mart. Fl. Bras. xiii. 1. 274 = crotonoides.
pruinosa, *Griseb. in Gortt. Abh.* xxiv. (1879) 18.—Reg. Argent.
pubernla, DC. Prod. i. 248 = tomentosa.
pubiflora, *DC. l. c.* 246.—Malaya.
pulcherrima, *Jacq. Enum. Pl. Carib.* 24.—N. Granat.
punila, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 260.—China; Ind. or.
punuisia, *Burch. Trav.* i. 492.—Afr. austr.
punctata, Wall. Cat. n. 7002 = pumila.
pyracantha, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 59.—Madag.
pyrifolia, *Lam. Encyc.* i. 606.—Ind. or.
pyrifolia, Wight & Arn. Prod. 25 = grandiflora.
quadrifaria, DC. Prod. i. 247 = horrida.
quinifolia, *DC. l. c.*—Austral.
quinquefiora, Steud. Nom. ed. II. i. = quinifiora.
racemifiora, DC. Prod. i. 248 = grandis.
racemosa, *Mill. Gard. Dict. ed.* VIII. n. 5.—Mexic.
racemulosa, *DC. Prod.* i. 248.—Afr. austr.
radiatiflora, Ruiz & Pav. Fl. Per. v. t. 433 = Crataeva radiatiflora.
reflexa, *Thonn. & Schum. in Dan. Vid. Sel. Afh.* iv. (1828) 11.—Guinea.
reticulata, Klein, ex Wall. Cat. n. 6986 = diversifolia.
retusa, *Griseb. in Goett. Abh.* xxiv. (1879) 18.—Am. austr.
retusifolia, Thw. Enum. Pl. Zeyl. 16, 400 = sepiaria.
Rheedii, DC. Prod. i. 246 = zeylanica.
Richardi, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 463. —Madag.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 69.—Ins. Fiji.
rosea, *Oliver, Fl. Trop. Afr.* i. 99.—Afr. trop.
Rothii, *Oliver, l. c.* 97.—Afr. trop.
rotundifolia, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 185 = zeylanica.
Roxburghii, *DC. Prod.* i. 247.—Ind. or.
roydiaefolia, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 69.—Burma.
rufescens, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 321.—Ins. Samoa.
rugulosa, *Turcz. l. c.* 326.—Venezuela.
rupestris, Sibth. & Sm. Fl. Graec. t. 487 = spinosa.
Russeliana, *Wall. Cat.* n. 7001.—Ind. or.
sabiaefolia, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 179. —Reg. Himal.
salaccensis, *Blume, Bijdr.* 54.—Java.
salicifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 17.—Reg. Argent.
saligna, Vahl, Symb. Bot. iii. 66 = C. Cynophallophora.
sandwicensis, Walp. Rep. v. 104 = seq.
sandwichiana, DC. Prod. i. 245 = spinosa.
sarmentosa, *A. Cunn. ex Benth. Fl. Austral.* i. 95.—Austral.
sativa, Pers. Syn. ii. 59 = spinosa.
saxatilis, *H. B. & K. ex DC. Prod.* i. 250.—Venezuela.
scabrida, *H. B. & K. Nov. Gen. et Sp.* v. 95.—Am. austr.
scandens, Vell. Fl. Flum. 330; v. t. 109 = linearis.
Schimperi, *Steud. Nom.* ed. II. i. 278.—Arab.
sclerophiila, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 206 = pumila.
securidocses, *Triana & Planch. l. c.* Sér. IV. xvii. (1863) 77.—N. Granat.
sepiaria, *Linn. Syst.* ed. X. 1071.—Ind. or.; Malaya.
sepiaria, Wall. Cat. n. 6999 G = pedunculosa.
sessilis, *Benth. ex DC. Prod.* i. 249.—N. Granat.
Shanzeriana, *F. Muell. Fragm.* x. 94.—Austral.
siamensis, *Kurz, For. Fl. Brit. Burma*, i. 53.—Burma.
sicula, Duham. Arbr. i. 159 = spinosa.
sideroxylla, Ruiz & Pav. ex DC. Prod. i. 252 = crotonoides.
sikkimensis, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 121.—Reg. Himal.
silfiuensis, Linn. Syst. ed. X. 1071 = jamaicensis.
Sinclairii, *Benth. Bot. Voy. Sulph.* 68. t. 27.—Am. austr.

CAPPARIS:—
singalensis, Korth. ex Miq. Illustr. Fl. Arch. Ind. 88 = pubiflora.
Sodada, R. Br. in Denh. Clapp. & Oudn. Trav. 225 = aphylla.
solanoides, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 59 = chrysomela.
speciosa, *Griseb. in Goett. Abh.* xxiv. (1879) 18.—Reg. Argent.
speciosa, Moric. ex Eichl. in Mart. Fl. Bras. xiii. 1. 272 = C. Yco.
spinosa, *Linn. Sp. Pl.* 503.—Reg. Mediterr. ; Ind. or. ; Oriens.
Spruceі, *Eichl. in Mart. Fl. Bras.* xiii. 1. 281.—Peruv.
stylosa, DC. Prod. i. 246 = diversicosta.
subacuta, *Miq. Fl. Ind. Bat.* i. 1. 101.—Borneo.
subbiloba, H. B. & K. Nov. Gen. et Sp. v. 90 = C. Cynophallophora.
subcordata, *Span. in Linnaea*, xv. (1841) 166.—Ins. Timor.
subspinosa, *Roxb. Hort. Beng.* [98]; *Fl. Ind.* ii. 568.—Ins. Moluce.
Swinhoii, *Hance, in Journ. Bot.* vi. (1868) 296.—China.
tarapotensis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 284.—Peruv.
tenera, *Dals. in Hook. Kew Journ.* ii. (1850) 41.—Ind. or.
tenuiflora, Fl. Brit. Ind. i. 178, sphalm. = ternifiora.
tenuifolia, J. F. Gmel. Syst. 1519, sphalm. = seq.
tenuisiliqua, *Jacq. Enum. Pl. Carib.* 24.—N. Granat.
ternifiora, DC. Prod. i. 247 = horrida.
tetrasperma, Thw. Enum. Pl. Zeyl. 15 = tenera.
Thonningii, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 286.—Afr. trop.
Thorelиana, *F. Muell. Fragm.* v. 104.—Austral.
tomentosa, *Lam. Encyc.* i. 606.—Afr. trop.
torulosa, Griseb. ex Hemsl. Biol. Centr. Am. Bot. i. 44 = odoratissima.
torulosa, Sw. Prod. Veg. Ind. Occ. 81 = jamaicensis.
trapeziflora, *Span. in Linnaea*, xv. (1841) 165.—Ins. Timor.
trifiora, Mill. Gard. Dict. ed. VIII. n. 10 = frondosa.
trifoliata, Roxb. Hort. Beng. 41; Fl. Ind. ii. 571 = Crataeva religiosa.
trifoliata, Spreng. ex DC. Prod. i. 248 = Crataeva tapioides.
trinervis, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 176. —Burma.
triphylla, *Thunb. Prod. Pl. Cap.* 92.—Afr. austr.
Tweediana, *Eichl. in Mart. Fl. Bras.* xiii. 1. 273.—Reg. Argent.
tylophylla, *Spreng. Syst.* iv. *Cur. Post.* 204.—Java.
uberiflora, *F. Muell. Fragm.* ix. 172.—Austral.
umbellata, *R. Br. in DC. Prod.* i. 247.—Austral.
umbellata, Willd ex Eichl. in Mart. Fl. Bras. xiii. 1. 273 = angustifolia.
umbonata, *Lindl. in Mitch. Journ. Trop. Austral.* 257.—Austral.
uncinata, Edgew. in Journ. As. Soc. Beng. xvi. (1847) ii. 1218 = spinosa.
uncinata, Lodd. ex Eichl. in Mart. Fl. Bras. xiii. 1. 270 = jamaicensis.
uncinata, Wall. Cat. n. 6998 = tenera.
undulata, Lodd. ex Loud. Hort. Brit. 215 = Loddigesii.
undulata, *Zeyh. ex Eckl. & Zeyh. Enum.* 14.—Afr. austr.
variabilis, Wall. Cat. n. 7004 (non Capparid.).—Ind. or.
Velloziana, Mart. in Flora, xxii. (1839) 1. Beibl. 23 = C. Cynophallophora.
verrucosa, *Jacq. Enum. Pl. Carib.* 23.—Ind. occ.; N. Granat.
verrucosa, Wight, ex Wall. Cat. n. 6980 = styiosa.
vesicolor, *Griff. Notul.* iv. 577.—Burma.
verticillaris, *Turcz. in Bull. Soc. Nat. Mosc.* (1863) i. 553.—Bras.
viminea, *Hook. f. & Thoms in Fl. Brit. Ind.* i. 179.—Reg. Himal.; Burma.
viridiflora, H. B. & K. Nov. Gen. et Sp. v. 92 = tenuisiliqua.
Volkmannia, *DC. Prod.* i. 247.—Afr. austr.
Wallichiana, Wight & Arn. Prod. 25 = zeylanica.
Wightiana, Wall. Cat. n. 6987 = zeylanica.
Yco, *Mart. Reise*, ii. 599; *Mart. & Eichl. in Mart. Fl. Bras.* xiii. 1. 272.—Bras.

CAPPARIS:—
Zeyheri, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 224.—Afr. austr.
zeylanica, *Linn. Sp. Pl. ed.* II. 720.—Ind. or.
zeylanica, Roxb. Hort. Beng. 47; Fl. Ind. ii. 567 = horrida.
zeylanica, Wight & Arn. Prod. 25 = acuminata.
Zippeliana, *Miq. Illustr. Fl. Arch. Ind.* 35. t. 14.—N. Guinea.

CAPRAEA, Opiz, Seznam, 28 (1852) = **Salix**, Tourn.
acuminata, Opiz, l. c. = S. acuminata.
aurita, Opiz, l. c. = S. aurita.
cinerea, Opiz, l. c. = S. cinerea.
grandifolia, Opiz, l. c. = S. grandifolia.
vulgaris, Opiz, l. c. = S. Caprea.

CAPPARIA, [Tourn.] Linn. Gen. ed. I. 98, 75 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. 2. 992.
POGOSTOMA, Schrad. Ind. Sem. Hort. Goett. (1831).
XUARESIA, Ruiz & Pav. Prod. 24. t. 4 (1794).
aegyptiaca, Steud. & Hochst. ex Endl. Nov. Doc. 28 = Anticharis arabica.
arabica, Steud. & Hochst. ex Steud. Nom. ed. II. i. 278 = Anticharis arabica.
bifora, *Linn. Sp. Pl. Pl.* 628.—Am. trop.
calycina, A. Gray, in Proc. Am. Acad. vi. (1862) 49 = Myoporum debile.
crustacea, Linn. Mant. 87 = Vandellia crustacea.
cunaata, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 45 = saxifragaefolia.
diffusa, Roxb. Hort. Beng. 47; Fl. Ind. iii. 93 = Ebermaiera thysoidea.
diosoeta, Delile, Fl. Egypte, 289. t. 82. f. 2 = Sutera glandulosa.
duranteaefolia, Linn. Syst. ed. X. 1116 = Stemodia duranteaefolia.
gratioloides, Linn. l. c. 1117 = Ilysanthes gratioloides.
gratissima, Roxb. Hort. Beng. 47; Fl. Ind. iii. 92 = Limnophila Roxburghii.
hirsuta, H. B. & K. Nov. Gen. et Sp. ii. 355 = saxifragaefolia.
humifusa, Buch.-Ham. ex Wall. Cat. n. 3883 = Centranthera hamiflora.
humilis, [Soland. in] Ait. Hort. Kew. ed. I. ii. 354 = Stemodia parviflora.
integerrima, *Miq. in Linnaea*, xxii. (1849) 476.—Guiana.
integrifolia, *Mart. & Gal. in Bull. Acad. Brux.* xii. ii. (1846) 90.—Mexic.
lanceolata, Hort. ex Steud. Nom. ed. II. i. 278 = salicifolia.
lanceolata, Linn. f. Suppl. 284 = Freylinia lanceolata.
lanceolata, *Vahl, Ecl. Am.* ii. 47.—N. Granat.
lucida, [Soland. in] Ait. Hort. Kew. ed. I. ii. 353 = Teodia lucida.
maritima, Moric. ex Benth. in DC. Prod. x. 429 = bifora.
Mouriera, Roxb. Hort. Beng. 47 = Herpestis Monniera.
multifida, Michx. Fl. Bor. Am. ii. 22. t. 35 = Conobea multifida.
multiflora, Steud. Nom. ed I. 148 = Conobea multifida.
peruviana, Benth. in DC. Prod. x. 480 = bifora.
pusilla, Torr. in Ann. Lyc. N. York, i. (1824) 86 = Mimulus floribundus.
riparia, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 301 = Centranthera hispida.
rigida, *Thunb. Prod. Pl. Cap.* 108.—Afr. austr.
salicifolia, Link & Otto, Ic. Fl. Rar. Hort. Berol. 11. t. 4 = Freylinia costricidea.
salicifolia, Salisb. Prod. 94 = Freylinia lanceolata.
saxifragaefolia, *Cham. & Schlecht. in Linnaea*, v. (1830) 106.—Mexic.
semiserrata, *Vahl, Ecl. Am.* ii. 47.—Am. mer.
undulata, *Linn. f. Suppl.* 284.—Afr. austr.
uniflora, *Burm. f. Fl. Ind.* 138.—Ind. or.

CAPRIFICUS, Gasp. Nov. Gen. Fic. 6 (1844) = **Ficus**, Tourn. (Urticac.).
gigantea, Gasp. Ric. Caprif. 30 = F. fulva.
insectifera, Gasp. Nov. Gen. Fic. 6 = F. Carica.
leucocarpa, Gasp. Ric. Caprif. 30. t. 3, 3 = F. albescens.
oblongata, Gasp. l. c. = F. neapolitana.
pulmeulata, Gasp. l. c. = F. pulmeulata, *Miq.*
rugosa, Gasp. l. c. = F. rugosa, *Miq.*
sphaerocarpa, Gasp. l. c. = F. globosa, *Miq.*

CAPRIFOLIUM, Tourn. ex Linn. Syst. ed. I (1735) = Lonicera, Linn. (Caprifol.).
alpigenaeus, Gaertn. Fruct. i. 126 = L. alpigena.
alpinum, Lam. Fl. Fr. iii. 266 = L. alpigena.
arboreum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9.—Malaya.
atropurpureum, Hort. ex C. Koch, Dendrol. ii. ii. 8 = L. Caprifolium.
balearicum, Dum.-Cours. Bot. Cult. ed. II. iv. 358 = L. implexa.
brachaceum, Michx. Fl. Bor. Am. i. 105 = L. glauca.
caerulenm, Lam. Fl. Fr. iii. 266 = L. coerulea.
chinense, Watt. ex Lond. Hort. Brit. 79 = L. japonica.
ciliosum, Pursh, Fl. Am. Sept. i. 160 = L. ciliosa.
cristatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9.—Malaya.
dentatum, Raf. Atl. Journ. 151.—Am. bor.
dioicum, Roem. & Schult. Syst. v. 260 = L. glauca.
distinctum, Moench, Meth. 501 = L. Periclymenum.
Douglasii, Lindl. in Trans. Hort. Soc. Lond. vii. (1830) 244 = L. hirsuta.
dumetorum, Lam. Fl. Fr. iii. 267 = L. Xylosteum.
etruscum, Roem. & Schult. Syst. v. 261 = L. etrusca.
flavum, Ell. Sketch, i. 271 = L. flava.
flexuosum, Hort. ex Steud. Nom. ed. II. i. 278 = L. japonica.
Fraseri, Pursh, Fl. Am. Sept. i. 160 = L. flava.
germanicum, Delarb. Fl. Auv. ed. II. 128 = L. Caprifolium.
germanicum, Quer, Fl. Esp. t. 79 = L. etrusca.
germanicum, Roehl. Deutschl. Fl. i. 175 = L. Periclymenum.
glaucum, Moench, Meth. 502 = L. glauca.
grataum, Pursh, Fl. Am. Sept. i. 161 = L. grata.
hirsutum, Desnoe. ex Loud. Hort. Brit. Suppl. iii. 509 = L. hirsuta.
hispidum, Dougl. ex Lindl. Bot. Reg. t. 1761 = L. hispidula.
hortense, Lam. Fl. Fr. iii. 265 = L. Caprifolium.
implexum, Roem. & Schult. Syst. v. 261 = L. implexa.
italicum, Medic. Ph. Bot. 97 = L. Caprifolium.
japonicum, D. Don, Prod. Fl. Nep. 140 = L. macrantha.
japonicum, Roem. & Schult. ex Steud. Nom. ed. II. i. 278 = L. longiflora.
javanicum, Blume, Bijdr. 653 = L. javanica.
longiflorum, Sabine, in Bot. Reg. t. 1232 = L. longiflora.
Lonicera, Blume, Bijdr. 653 = L. confusa.
macranthum, D. Don, Prod. Fl. Nep. 140 = L. macrantha.
nepalense, G. Don, ex Loud. Hort. Brit. 79 = L. macrantha.
occidentale, Lindl. Bot. Reg. t. 1457 = L. ciliosa.
ovatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9 = Clerodendron serratum.
pallidum, Schnr, Enum. Pl. Transs. 243 = L. Caprifolium.
paniculatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9.—Malaya.
parviflorum, Pursh, Fl. Am. Sept. i. 161 = L. glauca.
periclymenum, Roehl. Deutschl. Fl. i. 175 = L. Caprifolium.
Periclymenum, Delarb. Fl. Auv. ed. II. 129 = L. Periclymenum.
pilosum, H. B. & K. Nov. Gen. et Sp. iii. 427. t. 298 = L. pilosa.
pubescens, Goldie, in Edinb. Phil. Journ. (Apr. 1822) 298 = L. hirsuta.
pyrenaicum, Lam. Fl. Fr. iii. 266 = L. pyrenaica.
quercifolium, Malgen, Syst. Veg. 55 = L. Periclymenum.
reclinatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9.—Malaya.
rosseum, Lam. Fl. Fr. iii. 268 = L. japonica.
rotundifolium, Moench, Meth. 501 = L. Caprifolium.
rubrum, Raf. Fl. Ludov. 78.—Am. bor.
rupestre, Raf. in Nov. Fl. Am. iii. 17.—Am. bor.
scandicinum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9.—Malaya.
semperrirens, Moench, Meth. Suppl. 194 = L. sempervirens.
serotinum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9.—Malaya.
sylvaticum, Lam. Fl. Fr. iii. 265 = L. Caprifolium.
viticiflorum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9.—Malaya.
vulgare, Medic. Ph. Anat. 267 = L. Caprifolium.
Xylosteum, Gaertn. Fruct. i. 125 = L. Xylosteum.

CAPRIOLA, Adans. Fam. ii. 81 (1763) = **Cynodon**, Pers. (Gramin.).

CAPROXYLON, Tussac, Fl. Antill. iv. t. 20 (1827) = **Hedwigia**, Sw. (Burserac.).
Hedwigii, Tussac, l. c. = H. balsamifera.

CAPSELLA, Medic. Pflanzeng. i. 35 (1792). *CRUCIFERAE*, Benth. & Hook. f. i. 86.
BURSA, [Web. in] Wigg. Prim. Fl. Holsat. 47 (1780).
BURSA-PASTORIS, [Tourn.] Rupp. Fl. Jen. ed. Hall. 97 (1745), nomen prius.
HYSTEROLOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 117 (1838).
MARSYPOCARPUS, Neck. Elem. iii. 91 (1790).
MICROLEPIDIUM, F. Muell. in Linnaea, xxv. (1852) 371.
acutifolia, Rafin. New Fl. Am. ii. 27 = C. Bursa-pastoris!
agrestis, Jord. Diagn. i. 839 = C. Bursa-pastoris.
agrestis-rubella, Paill. in Mém. Soc. Emul. Doubs, Sér. IV. vi. (1872) 80 = C. Bursa-pastoris.
amblodes, Rafin. New Fl. Am. ii. 28 = C. Bursa-pastoris.
Andreasea, *F. Muell. in Wing, South. Sc. Record* (March, 1885).—Austral.
antipoda, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 94.—Austral.
antipoda, F. Muell. Pl. Vict. i. 44 = australis.
apetala, Opiz, in Flora, v. (1822) 369 = C. Bursa-pastoris.
australis, *Hook. f. ex Benth. Fl. Austral.* i. 81.—Austral.
Berengeriana, Hort. Petrop. ex Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 284 = procumbens.
bifida, *Rafin. New Fl. Am.* ii. 28 = C. Bursa-pastoris!
blennodina, F. Muell. Pl. Vict. i. 42 = Blennodia alpestris.
Bursa, Rafin. New Fl. Am. ii. 28 = C. Bursa-pastoris. Bursa-pastoris, *Medic. Pflanzeng.* 35.—Reg. temp.
ceratocarpa, Moench, Meth. 271 = Thlaspi ceratocarpon.
cochlearina, *F. Muell. Pl. Vict.* i. 51.—Austral.
cornigera, Medic. Pflanzeng. 56 = Thlaspi ceratocarpon.
dentata, Rafin. New Fl. Am. ii. 28 = C. Bursa-pastoris!
divaricata, *Walp. Rep.* i. 175.—Am. bor. occ.
Drummondii, *F. Muell. Fragm.* xi. 26.—Austral.
elliptica, A. Mey. in Ledeb. Fl. Alt. iii. 199 = procumbens.
erecta, Walp. Rep. i. 175 = divaricata.
furcata, Rafin. New Fl. Am. ii. 28 = C. Bursa-pastoris!
gracilis, Gren. in Mém. Soc. Emul. Doubs, Sér. III. ii. (1858) 408 = C. Bursa-pastoris.
grandiflora, *Boiss. Diagn.* Ser. I. i. 76.—Archip. Graec.
humistrata, *F. Muell. Fragm.* xi. 25.—Austral.
hybrida, *Albert, in Bull. Soc. Jc. Draguignan* (1884) [reimpr. 9].—Gallia.
integrifolia, Rafin. New Fl. Am. ii. 27 = C. Bursa-pastoris.
mexicana, *Hemsl. Diagn. Pl. Nov.* 19.—Mexic.
pastoris, Rupr. Fl. Caucas. 128 = C. Bursa-pastoris.
pauciflora, Koch, in Sturm, Deutschl. Fl. Heft x. t. 66 = Hutchinsia pauciflora.
pastoralis, Dulac, Fl. Hautes-Pyr. 189 = C. Bursa-pastoris.
pilosula, *F. Muell. Virt.-Pl.* i. 44.—Austral.
polymorpha, Cav. Descr. 411 = C. Bursa-pastoris.
poimenanthelantica, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 71 = C. Bursa-pastoris.
procera, Jord. Diagn. i. 845 = C. Bursa-pastoris.
procumbens, *Brias, Nouvl. Fl. Sous. Mont.* i. 14.—Reg. temp.
pubens, *Bastd. & Chaub. f. ex S. Wats. Bibliog. Ind. N. Am. Bl.*—Am. bor. occ.
puberula, *Rouy. & Foucaud, Fl. Centre*, 190.—Reg. Caucas.
rubella, *Reut. in Compter-Rend. Soc. Malléy.* (1858-54) 18.—Europ.
rubella-agrestis, Paill. in Mém. Soc. Emul. Doubs, Sér. IV. vi. (1872) 80 = C. Bursa-pastoris.
rubescens, Parsonsi, in Bull. Soc. Bot. Fr. xi. (1864) 511 = rubella.
rudcralis, Jord. Diagn. i. 840 = C. Bursa-pastoris.
setosissima, Jord. i. c. 841 = C. Bursa-pastoris.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 368.—Mexic.
spinosa, Moench, Meth. Suppl. 69 = Lepidium Draba.

CAPSELLA :—
sessocarpa. *Timb. in Bull. Soc. Hist. Nat. Toulouse*, iii. (1849) 115.—Gallia.
tasmanica, *P. Muell. Fragm.* xi. 26.—Ins. Tasman. Thomsoni. *Hook. f. in Journ. Linn. Soc.* v. (1861) 172.—Tibet. occ.
triangularis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69 = C. Bursa-pastoris.
virgata, Jord. Diagn. i. 239 = C. Bursa-pastoris.

CAPRICUM, [Tourn.] Linn. Syst. ed. I (1735). *SOLANACEÆ*, Benth. & Hook. f. ii. 892.
abyssinicum, *A. Rich. Tent. Fl. Abyss.* ii. 96.—Abyssin.
agrestinum, *Willd. ex Roem. & Schult. Syst.* iv. 809. —Am. bor.
anguinum, Mill. Gard. Dict. ed. VIII. n. 4 = annuum ?
angustifolium, *Dun. in DC. Prod.* xiii. i. 420.—Ind. or.
annuum, *Linn. Sp. Pl.* 188.—Reg. trop.
anomalum, *Franch. & Sav. Enum. Pl. Jap.* ii. 452.—Japon.
Azi, Vell. Fl. Flum. 61 ; ii. t. 6 = annuum.
baccatum, Hort. Genev. ex Dun. in DC. Prod. xiii. i. 420 = angustifolium.
baccatum, Buch.-Ham. ex Wall. Cat. n. 3644 = minimum.
baccatum, H. B. & K. Nov. Gen. et Sp. iii. 39 = cumanense.
baccatum, *Linn. Mant.* i. 48.—Reg. trop.
baccatum, Roxb. Obs. 38 = globiferum.
baccatum, Vell. Fl. Flum. 60 ; ii. t. 3 = frutescens.
bicolor, *Dun. in DC. Prod.* xiii. i. 428.—Hab. ?
bicolor, Jacq. Fragm. 66. t. 99. f. 1 = nigrum.
caerulescens, *Sess. Cat. Hort. Crem.* 27.—Hab. ?
campylopodium, *Sendt. in Mart. Fl. Bras.* x. 144.—Bras.
cerasiforme, *Link*, *Enum. Hort. Berol.* i. 190.—Hab. ?
cerasiforme, Hort. ex Dun. in DC. Prod. xiii. i. 420 = baccatum.
cerasiforme, Mill. Gard. Dict. ed. VIII. n. 5 = annuum ?
cerasiforme, Willd. Enum. Hort. Berol. i. 242 = grossum.
ceratocarpum, *Fingerh. Monog.* 22.—Hab. ?
cerealum, Bertol. Hort. Bonon. Pl. Nov. i. 6. t. 2 = frutescens.
chamaecrassa, Nees, in Trans. Linn. Soc. xvii. (1837) 65 = grossum.
chinense, *Jacq. Hort. Vindob.* iii. 38. t. 67.—Hab. ?
chlorocladum, *Dun. in DC. Prod.* xiii. i. 415.—Mexic.
ciliare, Willd. Enum. Hort. Berol. 248 = microcarpum.
Comarim, Vell. Fl. Flum. 60 ; ii. t. 2 = frutescens.
conicum, Lam. Illustr. n. 2890 = conoideum.
conicum, *G. F. W. Mey. Prim. Fl. Essq.* 112.—Guiana.
conicum, Vell. Fl. Flum. ii. t. 9 = baccatum.
conoides, Roem. & Schult. Syst. iv. 562 = frutescens.
conoideum, Mill. Gard. Dict. ed. VIII. n. 1 = annuum ?
cordiforme, Mill. l. c. n. 3 = annuum ?
crispum, *Dun. in DC. Prod.* xiii. i. 415.—Ins. Maurit.
cumanense, *Fingerh. Monog.* 17.—Venezuela.
curvipes, *Dun. in DC. Prod.* xiii. i. 428.—Guiana.
cydoniforme, Hort. ex Roem. & Schult. Syst. iv. 561 = tetragonum.
dichotomum, *Vell. Fl. Flum.* 61 ; ii. t. 9.—Bras.
dulce, *Hort. ex Dun. in DC. Prod.* xiii. i. 428.—Reg. trop.
fastigiatum, Blume, Bijdr. 705 = minimum.
fasciculatum, *Sendt. in Mart. Fl. Bras.* x. 143.—Bras.
frutescens, *Linn. Sp. Pl.* 189.—Reg. trop.
frutescens, Rudolph. Obs. 36 = conicum.
glandulosum, *Dun. in DC. Prod.* xiii. i. 417.—Bras.
globiferum, *G. F. W. Mey. Prim. Fl. Essq.* 113.—Guiana.
globosum, Bess. Cat. Hort. Crem. 27 = grossum.
grandiflorum, *Dun. in DC. Prod.* xiii. 416.—Bras.
grossum, *Linn. Mant.* i. 47 ; *Roxb. Hort. Beng.* 17 ; *Fl. Ind.* i. 574.—Reg. trop.
grossum, Willd. Enum. Hort. Berol. 241 = annuum.
Hamiltonii, G. Don, Gen. Syst. iv. 647 = nigrum.
humosum, *H. B. & K. Nov. Gen. et Sp.* iii. 40.—Am. austr.
hispidum, *Dun. in DC. Prod.* xiii. i. 419.—Texas.
Humboldtianum, *Dun.* l. c. 420.—Hab. ?

CAPSICUM :—
inaequale, *Vell. Fl. Flum.* 61 ; ii. t. 5.—Bras.
laurifolium, *Dun. in DC. Prod.* xiii. i. 418.—Ind. occ. ; Bras.
leucocarpon, *Mill. ex Fingerh. Monog.* 32.—Hab. ?
longum, Bouton, ex Dun. in DC. Prod. xiii. i. 414 = conoides.
longum, DC. Cat. Hort. Monsp. (1813) 86 = annuum.
luteum, *Lam. Illustr.* n. 2892.—Ind. or.
Maximowiczii, *Regel & Rach, Ind. Sem. Hort. Petrop.* (1858) 40.—Hab. ?
microanthum, *Link*, *Enum. Hort. Berol.* i. 190.—Bras.
microcarpum, *Cav. Dec.* 371 ; *DC. Cat. Hort. Monsp.* —Ind. occ.
microphyllum, Dun. in DC. Prod. xiii. i. 421 = baccatum.
Millori, Roem. & Schult. Syst. iv. 563 = annuum ?
minimum, *Blanco, Fl. Filip.* ed. I. 185.—Ins. Philipp.
minimum, Mill. Gard. Dict. ed. VIII. n. 10 = baccatum.
minimum, *Roxb. Hort. Beng.* 17 ; *Fl. Ind.* i. 574.—Reg. trop.
mirabile, *Mart. Fl. Bras.* x. 144.—Bras.
Narunca, Hort. Matr. ex Dun. in DC. Prod. xiii. i. 414 = nigrum.
nigrum, *Willd. Enum. Hort. Berol.* 242 —Hab. ?
odoratum, Stend. Nom. ed. II. i. 279 = frutescens.
odoriferum, Vell. Fl. Flum. 61 ; ii. t. 8 = frutescens.
oleaeforme, Mill. Gard. Dict. ed. VIII. n. 6 = annuum ?
ovatum, *DC. Cat. Hort. Monsp.* (1813) 86.—Hab. ?
oxycarpum, *Dun. in DC. Prod.* xiii. i. 426.—Bras.
parvifolium, *Sendt. in Mart. Fl. Bras.* x. 145.—Bras.
pendulum, *Willd. Enum. Hort. Berol.* i. 242.—Cuba.
pomiferum, Mart. ex Steud. Nom. ed. II. i. 279 = grossum.
pubescens, Dun. in DC. Prod. xiii. i. 415 = chlorocladum.
pubescens, *Ruiz & Pav. Fl. Per.* ii. 80.—Peruv.
pulchellum, Salisb. Prod. 134 = baccatum.
purpureum, Vahl, ex Homem. Hort. Hafn. i. 224 = grossum.
pyramidale, Mill. Gard. Dict. ed. VIII. n. 7.—Aegypt.
quitense, Willd. ex Roem. & Schult. Syst. iv. 809 = violaceum.
Rabenii, *Sendt. in Mart. Fl. Bras.* x. 145.—Bras.
salicifolium, *Dun. in DC. Prod.* xiii. i. 418.—Bras.
Schottianum, *Sendt. in Mart. Fl. Bras.* x. 148.—Bras.
silvestre, Vell. Fl. Flum. 60 ; ii. t. 1 = annuum.
sinense, Murr. Syst. ed. XIV. 226 = chinense.
sphaerium, Willd. Enum. Hort. Berol. i. 241 = annuum ?
striatum, *Fingerh. Monog.* 21.—Am. austr.
testiculatum, *Vis. ex Dun. in DC. Prod.* xiii. i. 424.—Hab. ?
tetragonum, Mill. Gard. Dict. ed. VIII. n. 8 = annuum ?
tomaeforme, Fingerh. ex Steud. Nom. ed. II. i. 279 = dulce.
torulosum, Hort. Matr. ex Hornem. Hort. Hafn. Suppl. 57 = pyramidale.
torulosum, *Vell. Fl. Flum.* 60 ; ii. t. 4.—Bras.
Tournefortii, Bess. Cat. Hort. Crem. (1811) 27 = annuum.
tortaeforme, Poepp. ex Fingerh. Monog. 32 = frutescens.
umbilicatum, Vell. Fl. Flum. 61 ; ii. t. 7 = annuum.
utriculosum, *Pant. Mag. Bot.* v. 197.—Chili.
villosum, *Sendt. in Mart. Fl. Bras.* x. 144.—Bras.
violaceum, Desf. Tabl. Hort. Par. ed. 1. 79 ; Desv. in Ham. Prod. 25 = nigrum.
violaceum, *H. B. & K. Nov. Gen. et Sp.* iii. 40.—Ecuador.

CAPURA, Blanco, Fl. Filip. ed. I. 264 (err. typ. 644). *SAPINDACEÆ*, Benth. & Hook. f. i. 405.
OVOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 572.
OTOPHORA, Blume, Rumphia, iii. 149 (1850).
nigrescens, Vidal, Sinopsis, i. 24. f. K = planuta.
planuta, *Blanco, Fl. Filip.* ed. I. 264.—Ins. Philipp.

CAPURA, Linn. Mant. ii. 149 (1771) = Wikstroemia, Endl. (Taymnhea.).
purpurascens, Linn. l. c. 235 = W. indica.

CAQUEPIRIA, J. F. Gmel. Syst. 651 (1791) = Gardenia, Ellis (Rubiac.).
Berghia, J. F. Gmel. l. c. = G. Thunbergia.

CARACALLA, Tod. Ind. Sem. Hort. Panorm. (1861) 32 ; et in Lem. Illustr. Hortic. ix. (1862) Misc. 18 = Phaseolus, Tourn. (Legumin.).
pulcherrima, Tod. ll. cc. = P. Caracalla.

CARACHERA, Juss. in Dict. Sc. Nat. vii. 11 (1817) = Charachera, Forsk. (Inc. sed.).

CARADESIA, Rafin. New Fl. Am. iv. 80 (1836) = Rupatorium, Tourn. (Compos.).
pauciflora, Rafin. l. c. = E. pauciflorum, *Rafin.*

CARAGANA, Lam. Encyc. i. 615 (1783). *LEGUMINOSÆ*, Benth. & Hook. f. i. 506.
acaulis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 44. —Afghanist.
Altagana, Poir. Encyc. Suppl. ii. 89 = microphylla.
ambigua, *Stocks, in Hook. Kew Journ.* iv. (1852) 145. —Beluchist.
arborescens, *Lam. Encyc.* i. 615.—Sibir.
arenaria, *Down., ex Sims, Bot. Mag.* t. 1886.—Sibir.
argentea, Lam. Encyc. i. 616 = Halimodendron argenteum.
brevispina, *Benth. in Royle, Illustr. Bot. Himal.* 198. —Reg. Himal. ; Afghan.
Bungei, *Ledeb. Fl. Alt.* iii. 264.—Sibir. alt.
Chamlagu, *Lam. Encyc.* i. 616.—China.
chinensis, Turcz. ex Maxim. Prim. Fl. Amur. 470 = C. Chamlagu.
confertа, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 118.—Tibet. occ.
crassicaulis, *Benth. ex Baker, l. c.* 117.—Reg. Himal.
cuneata, *Baker, l. c.*—Reg. Himal.
cuneata, Moench, Meth. 138 = frutescens.
digitata, Lam. Encyc. i. 616 = frutescens.
ferox, Lam. l. c. 816 = spinosa.
flava, Poir. Encyc. Suppl. ii. 90 = Robinia flava.
frutescens, *Medic. in Vorles. Churpf. Phys. Oec.* ii. (1787) 265.—Reg. Caucas. ; Soongaria.
frutex, C. Koch, Dendrol. i. 48 = seq. ?
fruticosa, Bess. Cat. Hort. Crem. 118 = arborescens.
Gerardiana, *Benth. in Royle, Illustr. Bot. Himal.* 198.—Reg. Himal.
glomerata, Hort. ex C. Koch, Dendrol. i. 48 = microphylla.
gracilis, *Hort. ex C. Koch, l. c.* 52.—Hab. ?
grandiflora, *DC. Prod.* ii. 268.—As. Min. ; Reg. Casp.
Halodendrum, Hoffmgg. Verz. Pfl. 46 = Halimodendron argenteum.
inermis, Moench, Meth. 138 = arborescens.
jubata, *Poir. Encyc. Suppl.* ii. 89.—Sibir.
microphylla, *Lam. Encyc.* i. 615.—Sibir. alt.
mollis, Bess. Enum. Pl. Volh. 29 = frutescens.
mongolica, *Fisch. ex Steud. Nom.* ed. II. i. 279.—Mongolia.
mongolica, Hort. ex C. Koch, Dendrol. i. 47 = arborescens.
Moorcroftiana, Benth. in Royle, Illustr. Bot. Himal. 198 = Sophora Moorcroftiana.
muhgrna, Bunge, Monog. Astrag. 240 = crassicaulis.
parvifolia, *Hoffmgg. Verz. Pfl.* viii. *Anh.* 89 ; *ex Walp. Rep.* i. 681.—Hab. ?
polyacantha, *Benth. in Royle, Illustr. Bot. Himal.* 198.—Reg. Himal.
pygmaea, *DC. Prod.* ii. 268.—Reg. Himal. ; As. bor.
quadrifolia, Medic. Pfl.-Anat. 68 = frutescens.
Redowskii, *DC. Mém. Lég.* 94. t. 11. f. 46.—Sibir.
rosea, Turcz. ex Maxim. in Prim. Fl. Amur. 470 = frutescens.
sibirica, *Medic. in Vorles. Churpf. Phys. Oec.* ii. (1787) 265.—Sibir.
sophorafolia, Hort. ex C. Koch, Dendrol. i. 47 = arborescens.
sophorafolia, *Tausch, in Flora*, xxi. (1838) t. *Beibl.* 77.—Hab. ?
spinosa, *DC. Prod.* ii. 269.—Sibir.
spinosa, Kit. ex DC. l. c. 814 = Pictetia squamata.
spinescissima, Benth. in Royle, Illustr. Bot. Himal. 198 = Gerardiana.
tenuis, *Hoffmgg. Pretiv. Pfl.* viii. *Anh.* 40 ; *ex Walp. Rep.* i. 681.—Hab. ?
tortuosa, *Boiss. ex Tchihat. Asie Min. Bot.* i. 49, nomen.—As. Min.

CARAGANA :—
tragacanthoides, *Poir. Encyc. Suppl.* ii. 90.—Sibir.
trifoliata, Lindl. Bot. Reg. (1845) Misc. 41 = hveriaginea ?
ulicina, *Stocks, in Hook. Kew Journ.* iv. (1852) 145.—Beloochist.
versicolor, Benth. in Royle, Illustr. Bot. Himal. 198. t. 34 = pygmaea.

CARAGUATA, [Flum.] Lindl. Bot. Reg. sub t. 1068 (1827). *BROMELIACEAE*, Benth. & Hook. f. iii. 668.
DEVILLEA, Bert. ex Schult. f. Syst. vii. p. lxvii. 1229 (1830).
MARANGIA, E. Morr. in Belg. Hortic. (1877) 59, 199. t. 8.
Andreana, *E. Morr. in Rev. Hortic.* (1884) 247.—N. Granat.
angustifolia, *Baker, in Gard. Chron.* (1884) ii. 616.—N. Granat.
Augustae, *Benth. & Hook. f. Gen.* iii. 668.—Guiana.
Berteroniana, *Schult. f. Syst.* vii. ii. 1229.—Porto Rico.
cardinalis, E. André, in Illustr. Hortic. xxvii. (1880) t. 874 ; Rev. Hortic. (1883) 12 = lingulata.
coriostachya, *Griseb. in Goett. Nachr.* (1864) 21.—Venezuela.
fulgens, *Griegt. l. c.*—Venezuela.
Fuerstenbergiana, *Kirch. & Wittm. in Berl. Gartenzeit.* (1883) 299.—Ecuador.
latifolia, Beer, Bromel. 78 = lingulata.
lingulata, *Lindl. Bot. Reg.* sub t. 1068.—Am. trop.
macrostachya, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 122.—Porto Rico.
musaica, *E. André, in Illustr. Hortic.* xxiv. (1877) t. 268.—N. Granat.
nitida, *Griseb. in Goett. Nachr.* (1864) 21.—Venezuela.
Osyana, *E. Morr. in Belg. Hortic. Annex.* (1885) 254.—Ecuador.
Pesacckii, *E. Morr. l. c.* 82.—Reg. Andin.?
sanguinea, *E. André, in Rev. Hortic.* (1883) 468.—N. Granat.
serrata, Hort. Petrop. ex Antoine, Bromel. 46 = Karatas Schwenziewii.
serrata, Schult. f. Syst. vii. ii. 1231 = Tillandsia serrata.
sphaerocephala, *Gaudich. Bot. Voy. Bonite*, t. 61.—Hab.?
splendens, Hort. ex Planch. in Fl. des Serres, (1856) 31. t. 1001 = lingulata.
Van-Volxemi, *E. André, in Illustr. Hortic.* xxv. (1878) t. 226.—N. Granat.
Zahnii, *Hook. f. Bot. Mag.* t. 6059.—Am. centr.

CARAIPA, Aubl. Pl. Gui. i. 561 (1775), ex parte. *TERNSTROMIACEAE*, Benth. & Hook. f. i. 188.
africana, *Oliver, Fl. Trop. Afr.* i. 172.—Afr. trop.
angustifolia, *Aubl. Pl. Gui.* i. 561.—Guiana.
calophylla, *Benth. in Journ. Linn. Soc.* v. (1860) 61.—Bras.
calophylla, Spreng. ex Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 166 = glabrata.
costata, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1860) 61.—Am. austr.
densiflora, *Mart. Nov. Gen. et Sp.* i. 105.—Bras.
lasciculata, *Cambess. in Mém. Mus. Par.* xvi. (1828) 416.—Guiana.
glabrata, *Mart. Nov. Gen. et Sp.* i. 105.—Bras.
grandiflora, *Mart. l. c.*—Bras.
latifolia, *Aubl. Pl. Gui.* i. 561.—Guiana.
laurifolia, *Spruce, ex Choisy, in Mém. Soc. Phys. Genève.* xiv. (1855) 166.—Bras.
laxiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 264.—Guiana.
leiantha, *Benth. l. c.*—Guiana.
longifolia, *Aubl. Pl. Gui.* i. 561.—Guiana.
myrciaephylla, Spruce, ex Benth. in Journ. Linn. Soc. v. (1860) 61 = fascicularis.
paniculata, *Mart. Nov. Gen. et Sp.* i. 104.—Bras.
parvifolia, *Aubl. Pl. Gui.* i. 561.—Guiana.
racemosa, *Cambess. in Mém. Mus. Par.* xvi. (1828) 415. t. 8.—Bras.
Richardiana, *Cambess. l. c.* 414.—Guiana.
Sellowii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 578.—Bras.
sessilana, *Planch. ex Benth. in Journ. Linn. Soc.* v. (1860) 61.—Guiana.
surinamensis, *Miq. Stirp. Surin. Sel.* 75.—Guiana.

CARAIPA :—
tereticaulis, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 241.—Guiana.
variabilis, *Cambess. in Mém. Mus. Par.* xvi. (1828) 416.—Guiana.

CARAJAEA, Wedd. in DC. Prod. xvii. 84 (1873) = Castelnavia, Tul. (Podostemac.).
orthocarpa, Wedd. l. c. 85 = Castelnavia orthocarpa.

CARALLIA, Roxb. ex R. Br. in Flind. Voy. ii. 549 (1814). *RHIZOPHOREAE*, Benth. & Hook. f. i. 680.
BARRALDEIA, Thos. Gen. Nov. Madag. 24 (1809).
BARAULTIA, Spreng. Syst. ii. 313 (1825).
CATALIUM, Buch.-Ham. ex Wall. Cat. sub n. 4580 (1831?).
DEMIDOFIA, Dennst. Schluss. Hort. Malab. 31 (1818).
DIATOMA, Lour. Fl. Cochinch. 296 (1790), nomen prius.
KARE-KANDEL, Adans. Fam. ii. 88 (1763).
PETALOTOMA, DC. Prod. iii. 294 (1828).
SYMMETRIA, Blume, Bijdr. 1180 (1826).
Baraldiela, *Arn. in Ann. Nat. Hist.* i. (1838) 371.—Madag.
calycina, *Benth. in Journ. Linn. Soc.* iii. (1859) 75.—Zeylan.
celebica, *Blume, Mus. Bot. Lugd. Bat.* i. 131.—Ins. Celebes.
cerisopifolia, Miq. Anal. Bot. Ind. iii. 8 = integerrima.
ceylanica, Arn. in Ann. Nat. Hist. i. (1838) 371 = integerrima.
confinis, Blume, Mus. Bot. Lugd. Bat. i. 129 = lanceaefolia.
corymbosa, Arn. in Ann. Nat. Hist. i. (1838) 371 = integerrima.
cuspidata, *Blume, Mus. Bot. Lugd. Bat.* i. 129.—Borneo.
densiflora, *Grisf. Notul.* iv. 668.—Burma.
integerrima, *DC. Prod.* iii. 33.—As. et Austral. trop.
lanceaefolia, *Roxb. Hort. Beng.* [87] ; *Fl. Ind.* ii. 481. = Burma : Malaya.
lanceolaria, Wall. Cat. n. 4881 = lanceaefolia.
lucida, Roxb. ex Kurz, For. Fl. Brit. Burm. i. 451 = integerrima.
lucida, Roxb. Hort. Beng. [92] ; Fl. Ind. ii. 481 = lanceaefolia.
madagascariensis, *Tul. in Ann. Sc. Nat.* Sér. IV. iv. (1856) 117. t. 6.—Madag.
multiflora, *Blume, Mus. Bot. Lugd. Bat.* i. 131.—Borneo.
multiflora, Miq. Fl. Ind. Bat. i. i. 594 = floribunda.
obcordata, Wight, ex Walp. Rep. ii. 71 = integerrima.
octopetala, F. Muell. Pl. Austr. Trop. Occid. ex Benth. in Journ. Linn. Soc. iii. (1859) 74 = integerrima.
sinensis, Arn. in Ann. Nat. Hist. i. (1838) 371 = integerrima.
symmetria, Blume, Mus. Bot. Lugd. Bat. i. 130 = integerrima.
timorensis, Blume, l. c. 128 = integerrima.
Tulasnei, *Baill. Adansonia*, iii. (1862-63) 20.—Ins. Marian.
zeylanica, Arn. in Wight, Illustr. Ind. Bot. t. 90 = integerrima.

CARALLUMA, R. Br. in Mem. Wern. Soc. i. (1809) 25. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 792.
adscendens, *R. Br. i. c.*—Ind. or.
attenuata, *Wight, ic.* iv. 15. t. 1965.—Ind. or.
crenulata, Wall. Pl. As. Rar. i. 6. t. 7 = Boucerosia crenulata.
edulis, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 76.—Ind. or.
fimbriata, *Wall. Pl. As. Rar.* i. 7. t. 8.—Ind. or. ; Burma.
subulata, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 267.—Arabia.
umbellata, Haw. Syn. Pl. Succ. 47 = Boucerosia umbellata.

CARAMANICA, Tineo, Pl. Rar. Sic. 3 (1847 ?) = Tarenaya, Lkn. (Compos.).
tarenacoides, Tineo, l. c. 4 = T. minimum.

CARAMBOLA, Adans. Fam. ii. 306 (1763) = Averrhoa, Lkn. (Oxalid.).

CARANDA, Gaertn. Fruct. ii. 17. t. 88. f. 5 (1791) (Gen. dub. fructa deformato).
pedunculata, Gaertn. l. c.—Zeylan.

CARANDAS, Rumph. ex Adans. Fam. ii. 171 (1763) = Carissa, Lkn. (Apocyn.).

CARANGA, Vahl, Enum. i. 100 (1805) = Curanga, Juss. (Scrophular.).
amara, Vahl, l. c. = Curanga amara.

CARAPA, Aubl. Pl. Gui. ii. Suppl. 32. t. 387 (1775). *MELIACEAE*, Benth. & Hook. f. i. 388.
PERSOONIA, Willd. Sp. Pl. ii. 881 (1800).
RACAPA, M. Roem. Syn. Hesper. 129 (1846).
TOULOUCOUNA, M. Roem. l. c. (1846).
XYLOCARPUS, Koen. ex A. Juss. in Mém. Mus. Par. xix. (1830) 343.
ZELSA, Hort. ex Tenore, in Atti 8ª Reun. Sc. Ital. (1841) 504.
ZUELOA, Tenore, l. c. 503 (1841).
caniflora, Poepp. & Endl. Nov. Gen. et Sp. iii. 38 = Gaurea Poeppigii.
guianensis, *Aubl. Pl. Gui. Suppl.* 32. t. 387.—Guiana ; Afr. trop.
guineensis, Sweet, Hort. Brit. ed. I. 72 ; Juss. in Mém. Mus. Par. xix. (1830) 343 = procera.
guyanensis, Oliver, Fl. Trop. Afr. i. 336 = procera.
indica, A. Juss. in Dict. Sc. Nat. vii. 31 = moluccensis.
latifolia, Willd. ex C. DC. in DC. Monog. Phan. i. 718 = guianensis.
moluccensis, *Lam. Encyc.* i. 621.—Geront. trop.
nicaraguensis, *C. DC. in DC. Monog. Phan.* i. 717.—Am. centr.
obovata, Blume, Bijdr. i. 179 = moluccensis.
procera, *DC. Prod.* i. 626.—As. et Afr. trop.
Rumphii, Kostel. Allg. Med.-Pharm. Fl. v. 1988 = moluccensis !
sericea, Poepp. & Endl. Nov. Gen. et Sp. iii. 39. t. 245 = Guarea truncifiora.
surinamensis, *Miq. Stirp. Surin. Sel.* 75. t. 19.—Guiana.
Toulouconna, Guill. & Perr. Fl. Seneg. Tent. i. 128 = procera.
trifuga, Willd. ex C. DC. in DC. Monog. Phan. i. 564 = Guarea humilis.

CARAPICHEA, Aubl. Pl. Gui. 167. t. 64 (1775) = Cephaëlis, Sw. (Rubiac.).
Aublett, DC. Prod. iv. 536 = Ceph. involucrata.
guianensis, Aubl. Pl. Gui. i. 166 = Ceph. involucrata.
Kappleri, Miq. Stirp. Surin. Sel. 181.—Guiana.
Patrisii, DC. Prod. iv. 536 = Ceph. Patrisii.

CARARA, Medic. Pflanzeng. 84 (1792) = Senebiera, DC. (1799) (Cruciffer.).
Coronopus, Medic. l. c. 85 = S. Coronopus.

CARAXERON, Vaill. ex Rafin. Fl. Tellur. iii. 88 (1836) = Phillonerus, R. Br. (Amaranthac.).
brasiliense, Rafin. l. c. = Telanthera dentata.
conicus, Rafin. l. c. = P. conicus.
diffusus, Rafin. l. c. = Compsoma diffusa.
vermicularis, Rafin. l. c. = P. vermicularis.

CARBENIA, Adans. Fam. ii. 116 (1763) (Carbeni). *COMPOSITAE*, Benth. & Hook. f. ii. 468.
BENEDICTA, Bernh. Syst. Verz. Erf. 106 (1800).
benedicta, *Adans. l. c.* 116.—Reg. Mediterr. et Cancas.

CARCIA, Raeusch. Nom. ed. III. 275 (1797) = Garcia, Rohr. (Euphorb.).

CARDA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid ?).

CARDAMINDUM, Tourn. ex Adans. Fam. ii. 368 (1763) = Trapeolum, Lkn. (Geraniac.).
majus, Mouch, Meth. Suppl. 20 = T. majus.
minus, Mouch, l. c. 21 = T. minus.

CARDAMINE, [Tourn.] Lkn. Syst. ed. I (1735) ; Gen. ed. I. 276 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 68.
DENTARIA, [Tourn.] Lkn. Syst. ed. I (1735).
HETEROCARPUS, Pall. in Nov. Spec. xiv. (1856) 241.
KARDAMOULTHOS, Schlecht. in Linnaea, xxviii. (1856) 472.
PTEROSIPHONIA, DC. Prod. i. 220 (1824).

CARDAMINE:—
acaulis, *Berg, in Bot. Zeit.* xiv. (1856) 874.—German.
acris, *Griseb. Spicil. Fl. Rumel.* i. 253.—Macedon.; Armen.
affinis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 137.—Chili.
africana, *Linn. Sp. Pl.* 655.—Ind. or.; Afr.; Japon.
alleria, Comm. ex DC. Syst. ii. 252 = africana.
alpina, *Willd. Sp. Pl.* iii. 481.—Europ.
amara, Bieb. Fl. Taur. Cauc. ii. 109 = uliginosa.
amara, Lam. Encyc. ii. 186 = pratensis.
amara, *Linn. Sp. Pl.* 656.—Europ.; Aa. bor.
amplexicaulis, *Hornem. in Bot. Zeit. iv.* (1846) 812.—Hispan.
angulata, *Hook. Bot. Misc.* i. (1830) 348. t. 69.—Am. bor. occ.
angulata, Regel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) ii. 173 = hirsuta.
angulata, Torr. in Pacif. Rail. Rep. iv. 66, 77 = paucisecta.
angustifolia, *Rafin. Fl. Ludov.* 84.—Louisiana.
antemiquana, Burch. ex DC. Syst. ii. 259 = africana.
anticorbutica, Banks, ex Griseb. in Goett. Abh. vi. (1854) 112 = plicatula.
apetala, Gilib. Fl. Lituan. ii. 67 = impatiens.
apetala, Moench, Meth. 259 = hirsuta.
appendiculata, *Franch. & Sav. Enum. Pl. Jap.* ii. 281.—Japon.
arenosa, Roth, Man. Bot. 926 = Arabis arenosa.
armoracioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 298.—Venezuela.
articulata, Pursh, Fl. Am. Sept. ii. 430 = Parrya nudicaulis.
asarifolia, *Linn. Sp. Pl.* 654.—Europ.
auriculata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 319.—Calif.
axillaris, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 290.—Bolivia.
bellidifolia, Hill, Herb. Brit. ii. 275 = bellidifolia.
belgaumensis, Dalz. ex Hook. in Hook. Kew Journ. iv. (1852) 294 = subumbellata.
bellidifolia, All. Fl. Pedem. i. 260. t. 18. f. 3 = alpina.
bellidifolia, *Linn. Sp. Pl.* 654.—Reg. bor. et arct.
Bergeriana, Andrz. ex DC. Syst. ii. 266 = glanca Berro, *Stend. Nom.* ed. II. i. 360.—Ins. Juan. Fern.
bicolor, Opiz, ex J. & C. Presl, Fl. Cech. 136 = amara.
Bielzii, Schan. in Verh. Siebenb. Ver. Naturw. iv. (1853) 94 = amara.
*Boccon*i, *Vis.* Prod. Fl. Cors. App. 4 = thalictroides.
bonariensis, Juss. ex Pers. Syn. ii. 195 = flaccida.
borbonica, Boj. Hort. Maurit. 11 = hirsuta.
borbonica, Pers. Syn. ii. 195 = africana.
borealis, Andrz. ex DC. Syst. ii. 256 = prorepens.
borealis, Laest. ex Nym. Consp. 37 = amara.
Boryi, Boiss. Elench. 9 = Sisymbrium pinnatifidum.
brachycarpa, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 58.—Japon.
brachycarpa, Opiz, Natural. xi. (1836) 411 = impatiens.
brachiata, *S. Moore, in Journ. Bot.* xvi. (1878) 130.—Japon.
brasiliensis, *D. Dietr. Syn. Pl.* iii. 698.—Bras.
Brewer, *S. Wats. in Proc. Am. Acad.* x. (1875) 389.—Am. bor. occ.
buchtormensis, Willd. ex DC. Syst. ii. 258 = pratensis.
bulbifera, *[R. Br. in] Ait. Hort. Kew.* ed. II. iv. 101.—Europ.
Burchellii, Spreng. Syst. ii. 886 = africana.
calabrica, *Archang. in Nuov. Giorn. Bot.* x. (1878) 168.—Italia.
calbucana, *Phil. in Anal. Univ. Chil.* (1873) 668.—Chili.
caldcirarum, *Guthn. in Seub. Fl. Asor.* 48.—Ins. Azores.
carnea, *Waldst. & Kit. Pl. Rar. Hung.* ii. 187. t. 169.—Transalp.
casparioides, Willd. ex Ledeb. Fl. Ross. i. 126 = uliginosa.
chaldaeoides, S. G. Gmel. It. iv. 106 = impatiens.
chaldaeoides, Lam. Encyc. ii. 186 = latifolia.
chaldaeoides, *Linn. Sp. Pl.* 656.—Italia; Hungar.
chaldaeoides, Pall. Reise, iii. 94 = macrophylla.
chaldaeoides, *S. Moore, in Journ. Bot.* xvi. (1878) 130.—Japon.
chenopodifolia, Pers. Syn. ii. 196.—Am. austr.
chilensis, DC. Syst. ii. 254.—Chili.
circaeoides, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 146.—Sikk. Himal.
Clematitis, *Skuttl. ex S. Wats. Bibliog. Ind. N. Am. Bot.* i. 85.—Am. bor.
cognata, *Stend. in Flora*, xxxix. (1856) 409.—Chili.
colchagnensis, *Barn. in C. Gay, Fl. Chil.* i. 115.—Chili.
cordata, *Barn. l. c.* 109.—Chili.
cordifolia, *A. Gray, Pl. Fendl.* 5.—N. Mexic.
corsica, Jord. ex Nym. Consp. 38 = graeca.
corsica, *Sieber, ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 293.—Ins. Corsica.
corydaloides, Gmel. ex Spreng. Syst. ii. 888 = glauca.
corymbosa, Hook. f. in Ic. Pl. t. 686 = hirsuta.
crassifolia, Opiz, Natural. xi. (1836) 412 = amara.
cretica, Nym. Consp. 38 = graeca.
croatica, Schott, Nym. & Kotschy, Analect. Bot. 46 = carnosa.
cunesata, *Greene, in Bull. Calif. Acad.* n. 3 (1885) 74.—Calif.
Cupani, Jord. ex Nym. Consp. 38 = graeca.
curvisiliqua, *Skuttl. ex Chapm. Fl. S. U. St. Suppl.* 504.—Florida.
dalmatica, Nym. Consp. 38 = maritima.
dasycarpa, Bieb. Fl. Taur. Cauc. iii. 437 = C. impatiens.
dasyclada, Bernh. ex Steud. Nom. ed. II. i. 280 = macrophylla.
debilis, Banks, ex DC. Syst. ii. 265 = hirsuta.
debilis, D. Don, Prod. Fl. Nep. 201 = hirsuta.
decumbens, *Barn. in C. Gay, Fl. Chil.* i. 109.—Chili.
decurrens, *Zoll. & Moritzi, Syst. Verz. Zoll.* 85. - Java.
dentata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 60.—N. Granat.
dentariaefolia, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 145 = macrophylla.
dentata, Schult. Obs. 136 = pratensis.
depressa, *Hook. f. Fl. Antarct.* 6.—N. Zel.
dictyosperma, *Hook. Journ. Bot.* i. (1834) 246.—Austral.
digitata, Richards. in Frankl. Narr. 1st Journ. 748 = pratensis?
diphylla, *Wood, Bot. & Fl.* 87.—Am. bor.
divaricata, Hook. f. Fl. N. Zel. i. 13 = stylosa.
diversifolia, Sternb. & Hoppe, in Denkschr. Bot. Ges. Regensb. i. (1816) 156. t. 2 = Arabis Halleri.
dryoneja, Schur, Enum. Pl. Transs. 47 = hirsuta.
duriaeiana, *Kenal, ex Der Moul. in Actes Linn. Soc. Bord.* xx. (1855) 466.—Gallia.
elegantula, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 146.—Reg. Himal.
Engelmanniana, Ind. Sem. Hort. Berol. (1840); ex Heynh. Nom. ii. 118 = Arabis ludoviciana.
enneaphylla, *[R. Br. in] Ait. Hort. Kew.* ed. II. iv. 101.—Europ.
Eschscholtziana, Andrz. ex Ledeb. Fl. Ross. i. 128 = purpurea.
exstylis, *F. Muell. in Trans. Vict. Inst.* i. (1854) 114.—Austral.
fabrotensis, Hornem. in Fl. Dan. t. 1292 = Arabis petraea.
fagetana, Schur, Enum. Pl. Transs. 47 = umbrosa.
fastigiata, *Hook. f. Handb. New Zeal. Fl.* 18.—N. Zel.
flaccida, Bert. ex C. Muell. in Walp. Ann. vii. 100 = macrorrhiza.
flaccida, *Cham. & Schlecht. in Linnaea*, i. (1826) 21.—Chili.
flavescens, *Phil. in Anal. Univ. Chil.* (1872) 667.—Chili.
flexuosa, With. Bot. Arr. Brit. Pl. ed. III. iii. 578 = sylvatica.
fluminensis, *Richt. in Kjoob. Vidensk. Meddel.* (1870) 165. t. 2.—Bras.
foliosa, Wall. Cat. n. 4779 = macrophylla.
fontana, Lam. Fl. Fr. ii. 499 = Nasturtium officinale.
fontinalis, Schur, Enum. Pl. Transs. 48 = pratensis.
foveicola, Godet, Suppl. Fl. Jura, 13 = pratensis.
Gambelii, *S. Wats. in Proc. Am. Acad.* xi. (1876) 147.—Calif.
gelida, *Schott, in Oestr. Bot. Wochenbl.* (1855) 145.—Tyrol.
geranifolia, *DC. Syst. ii.* 268.—Reg. Magellan.
gilanensis, Willd. ex DC. l. c. 262 = impatiens.
glacialis, *DC. l. c.* 264.—Reg. Magellan.
glandulosa, *Blanco, Fl. Filip.* ed. I. 821.—Ins. Philipp.
glauca, *Spreng. ex DC. Syst. ii.* 268.—Calabria.
glaucescens, Reichb. Ic. Fl. Crit. v. Index, 4 (Quid f.).—Europ.
Gmelini, Tausch, in Flora, xix. (1836) 402 = macrophylla.
gongylodes, *Phil. in Linnaea*, xxviii. (1856) 664.—Chili.
graeca, Jenks, ex Nym. Consp. 38 = longirostris.
graeca, *Linn. Sp. Pl.* 655.—Reg. Mediterr. or.
grandiflora, Hallier, in Bot. Zeit. xxiv. (1866) 209 = pratensis.
grandiflora, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 338 = tenuifolia, *Turcz.*
grandis, Schur, Enum. Pl. Transs. 49 = amara.
granulata, *Phil. in Linnaea*, xxxiii. (1864–65) 8.—Chili.
granulosa, All. Auct. Fl. Pedem. 16 = pratensis.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 146.—Sikk. Himal.
Hamiltonii, G. Don, Gen. Syst. i. 167 = hirsuta.
hamulosa, Bertol. Mant. Fl. Alp. Apuan. 48 = resedifolia.
hastata, Willd. ex Ledeb. Fl. Ross. i. 125 = uliginosa.
hastulata, Sm. Engl. Bot. t. 469 = Arabis petraea.
hastulata, Stend. Nom. ed. II. i. 280 = resedifolia.
hederacea, DC. Syst. ii. 264 = thalictroides.
herbinga, Jord. Diagn. i. 129 = pratensis.
heterophylla, Bury, in Ann. Sc. Gén. Phys. iii. (1820) 6 = Sisymbrium pinnatifidum.
heterophylla, Hook. Comp. Bot. Mag. i. (1835) 278 = hirsuta?
heterophylla, Host, Syn. Fl. Austr. 366 = resedifolia.
heterophylla, *Lapeyr. Hist. Abr. Pl. Pyr.* 377.—Mont. Pyren.
heterophylla, *Wood, Bot. & Flor.* 38.—Am. bor.
Heyneana, Welw. ex Reichb. Fl. Germ. Excurs. 676 = pratensis.
Hilariana, *Walp. Rep.* i. 137.—Bras.
hirsuta, *Linn. Sp. Pl.* 655.—Reg. temp. et subtrop.
hirsuta, Oed. in Fl. Dan. t. 148 = amara.
hirsuta, Pall. ex DC. Syst. ii. 256 = prorepens.
Huetii, Boiss. Diagn. Ser. II. v. 18 = Sisymbrium Huetii.
humilis, *Kit. in Linnaea*, xxxii. (1863) 496.—Hungar.
Ibaguense, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 60.—N. Granat.
impatiens, *Linn. Sp. Pl.* 655.—Europ.; As. bor.; Reg. Himal.
impatiens, O. F. Muell. in Fl. Dan. t. 735 = hirsuta.
indica, *Burm. f. Fl. Ind.* 140.—Java.
integrifolia, Gilib. Fl. Lituan. ii. 68 = pratensis.
intermedia, Hook. Ic. Pl. t. 358 = tenuifolia?
intermedia, *Stend. in Flora*, xxxix. (1856) 410.—Chili.
Jamesoni, *Hook. Lond. Journ. Bot.* vi. (1847) 293.—Ecuador.
javanica, *Miq. Illustr. Fl. Archip.* t. 10.—Java.
Kochii v. A. Kern. in Zeitschr. Ferdinand. Tirol. III. xv. (1870) 280.—Europ.
kopaonikensis, Pand. ex Nym. Consp. 37 = glauca.
laciniata, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 34.—Austral.
lacinulata, Stend. Nom. ed. II. i. 281 = Thelypodium laciniatum.
lasiolata, *Wood, Bot. & Flor.* 38.—Am. bor.
Lamoriili, *Hance, in Journ. Bot.* xiv. (1876) 363.—China.
latesiliqua, *Cheverm. in Trans. N. Z. Inst.* xv. (1883) 298.—N. Zel.
latifolia, Lej. Fl. Spa, ii. 348 = pratensis.
latifolia, *Vahl, Symb. Bot.* ii. 77.—Europ. austr.
laza, *Bruch. Pl. Hartw.* 158.—N. Granat.
Lechleriana, *Steud. in Flora*, xxxix. (1856) 409.—Chili.
leptonensis, *Rouy, Ind. Sem. Hort. Genev.* (1861) 4.—Hispan.
lereensis, Andrz. in Ledeb. Fl. Alt. iii. 38 = bellidifolia.
Libertiana, Lej. Rev. Fl. Spa, 135 = amara.
lilacina, Hook. Comp. Bot. Mag. i. (1835) 278 = tenuifolia.
litoralis, *Phil. in Anal. Univ. Chil.* (1865) ii. 214.—Chili.
lobata, Moench, Meth. 260 = P. graeca.
longirostris, *Jenka*, ' *Adat. Magyar.* 164.' —*Cf. Yust, Bot. Jahresb.* (1876) 1061.—Banat.
lucorum, Boiss. & Heut, ex Boiss. Fl. Orient. i. 161 = pectinata.
ludoviciana, Hook. Journ. Bot. i. (1834) 191 = Arabis ludoviciana.

CARDAMINE:—
Lunaria, Linn. Sp. Pl. 656 = Ricotia Lunaria.
lyrata, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 573.—China.
macrophylla, Schur, Enum. Pl. Transs. 49 = amara.
macrophylla, *Willd. Sp. Pl.* iii. 484.—Reg. Himal. ; Sibir. ; Japon.
macrorrhiza, *Bert. ex Steud. Nom.* ed. II. i. 281.—Chili.
magellanica, *Phil. in Anal. Univ. Chil.* (1873) 666.—Chili.
marginata, *Phil. l. c.* (1865) ii. 324.—Chili.
maritima, *Portenschl. Enum. Pl. Dalm.* 15.—Dalmat.
Matthioli. Moretti, ex Comol. Fl. Com. v. 157 = pratensis.
maxima, *Wood, Bot. & Flor.* 38.—Am. bor.
melananthera, Steud. Nom. ed. II. i. 280, 281 = seq.
melanthera, Stokes, Bot. Mat. Med. iii. 445 = amara.
Menziesii, DC. Syst. ii. 267 = Sisymbrium canescens.
micrantha, Spenn. Fl. Frib. iii. 922 = hirsuta.
microphylla, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 111.—Sibir. or.
microphylla, J. & C. Presl, Delic. Prag. 15 = thalictroides.
minima, *Steud. in Flora*, xxxix. (1856) 410.—Pers.
minuta, Willd. ex DC. Prod. i. 150 = microphylla.
monticola, *Timb. in Bull. Soc. Hist. Nat. Toulouse*, iii. (1869) 119.—Gallia.
multicaulis, Hoppe, ex Schur, Enum. Pl. Transs. 47 = hirsuta.
multifida, Pursh, Fl. Am. Sept. ii. 440 = Sisymbrium canescens.
maltifida, *Wood, Bot. & Flor.* 38.—Am. bor.
muscosa, Vahl, ex DC. Syst. ii. 260 = hirsuta.
nana, *Barn. in C. Gay, Fl. Chil.* i. 108.—Chili.
nana, *C. Muell. in Walp. Ann.* vi. 106.—Perav.
nasturtiana, Thuill. Fl. Par. ed. II. i. 330 = amara.
Nasturtii, Sprang. Syst. iv. Cur. Post. 241 = nasturtioides, *D. Don.*
nasturtiifolia, Steud. Nom. ed. II. i. 281 = nasturtioides, *Bert.* ; C. Berro.
nasturtioides, *Bert. in 'Merc. Chil.* (1829)' ; *Bull. Férussac*, xx. (1830) 109 ; *Barn. in C. Gay, Fl. Chil.* i. 118.—Chili.
nasturtioides, Camb. in St. Hil. Fl. Bras. Mer. ii. 89 = Hilariana.
nasturtioides, *D. Don, Prod. Fl. Nep.* 201.—Nepal.
nasturtioides, Schur, Enum. Pl. Transs. 48 = pratensis.
Nasturtium, Wallr. Beitr. Fl. Hercyn. 235 = amara ?
nemophila, *Phil. in Linnæa*, xxx. (1859–60) 136.—Chili.
nemorosa, *Lej. Fl. Spa*, ii. 62.—Belgium.
nervatensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 295.—Hispan.
nilagirica, Schlecht. ex Hook. f. & T. Anders. in Fl. Brit. Ind. i. 158 = subumbellata.
nipponica, *Franch. & Sav. Enum. Pl. Jap.* i. 281.—Japon.
nivalis, *Gill. ex Hook. Bot. Misc.* iii. (1833) 136.—Chili.
nivalis Pall. Reise, ii. 740 = Macropodium nivale.
nivalis, Schur, Enum. Pl. Transs. 46 = sessilifolia.
niwæ, Hook. Comp. Bot. Mag. i. (1835) 378 = dictyosperma.
nudicaulis, Linn. Sp. Pl. 654 = Parrya macrocarpa.
nudicaulis, Pall. ex DC. Syst. ii. 654 = Macropodium nivale.
nudicaulis, S. Wats. Bibl. Ind. N. Am. Bot. 54 = Parrya nudicaulis.
obliqua, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 19.—Abyssin.
occulta, *Harvem. Hort. Hafn. Suppl.* 71.—China.
oligosperma, A. Gray, in Proc. Am. Acad. viii. (1873) 376 = Breweri.
oligosperma, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 85.—Am. bor. occ.
olympica, Boiss. Diagn. Ser. I. viii. 19 = uliginosa.
Opizii, C. & J. Presl, Fl. Cech. 126 = amara.
orophila, *Timb. in Bull. Soc. Hist. Nat. Toulouse*, iii. (1869) 119.—Gallia.
ovata, *Benth. Pl. Hartw.* 158.—N. Granat.
oxycarpa, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 56 = pectinata.
oxyphylla, Andrz. ex Ledeb. Fl. Ross. i. 128 = pratensis ?
palustris, Kaaf, in Flora, xxix. (1846) 293 = pratensis.
palustris, Peterm. in Rabenh. Bot. Centralbl. i. (1846) 47 = pratensis.

CARDAMINE:—
paradoxa, *Hance, in Journ. Bot.* vi. (1868) 111.—China.
parviflora, Linn. Syst. ed. X. 1181 = hirsuta.
parviflora, Suter, Fl. Helv. ii. 59 = tetrandra.
pascigugo, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 296 = hirsuta.
pancisecta, *Benth. Pl. Hartw.* 297.—Calif.
pectinata, Pall. ex DC. Syst. ii. 264 = impatiens.
pedata, *Bert.* & *Tilings in Nouv. Mém. Soc. Nat. Mosc.* xi. (1858) 47.—Sibir.
pennsylvanica, Muhl. Cat. 60, ex Willd. Sp. Pl. iii. 486 = hirsuta.
pentaphylla, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 101 = pratensis.
pentaphylla, *Phil. in Linnæa*, xxxiii. (1864–65) 6.—Chili.
petiolaris, DC. Syst. ii. 264 = graeca.
petræa, Linn. Sp. Pl. 654 = Arabis petræa.
petræa, Townson, Trav. Hung. 490, 348 = Arabis arenosa.
picta, Hook. Lond. Journ. Bot. vi. (1847) 292. t. 12 = Porphyrocodon pictum.
pilosella, Willd. ex DC. Syst. ii. 256 = prorepens.
pinnata, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 101.—Europ.
Plumieri, Vill. Prosp. 38 = thalictroides.
polyphylla, D. Don, Prod. Fl. Nep. 201 = macrophylla.
præcox, Pall. ex Ledeb. Fl. Ross. i. 127 = hirsuta.
pratensis, Hook. f. Fl. Tasm. i. 19 = tenuifolia.
pratensis, *Linn. Sp. Pl.* 656.—Reg. temp.
præticola, Jord. Diagn. i. 128 = pratensis.
propinqua, Carm. in Trans. Linn. Soc. xii. (1819) 507 = hirsuta.
prorepens, *Fisch. ex DC. Syst.* ii. 256.—Sibir.
pseudopratensis, Schur, ex Nichols. in Journ. Bot. xvii. (1879) 200 = pratensis.
pubescens, Stev. ex DC. Syst. ii. 256 = prorepens.
punicea, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 296.—Venezuela.
purpurea, *Cham. & Schlecht. in Linnæa*, i. (1826) 20.—Am. bor. occ.
purpurea, Torr. & Gray, Fl. N. Am. i. 86 = pauciflora.
pusilla, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 18.—Abyssin.
pusilla, *Phil. in Linnæa*, xxviii. (1856) 665.—Chili.
pusilla, Schur, Enum. Pl. Transs. 47 = hirsuta.
radicata, *Hook. f. in Ic. Pl.* t. 982.—Tasman.
ramosissima, *Steud. in Flora*, xxxix. (1856) 400.—Chile.
raphanifolia, Pourr. in Mém. Acad. Toul. iii. (1788) 310 = latifolia.
reflexa, *Raffn. Fl. Ludov.* 84.—Am. bor.
Regnellana, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 78 = hirsuta.
reniformis, *Phil. in Anal. Univ. Chil.* (1865) ii. 313.—Chili.
repanda, Sm. ex DC. Syst. ii. 251 ; Prod. i. 150 = flaccida.
resedifolia, *Linn. Sp. Pl.* 654.—Europ.
rhomboidea, *DC. Syst.* ii. 246.—Am. bor.
rhomboidea, Durand, Fl. Utah. 160 = cordifolia.
rigida, *Bioh. Pl. Taur. Cauc.* iii. 499.—Reg. Caucas.
riparia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 61 = pratensis.
rostrata, *Griseb. in Goett. Abh.* vi. (1854) 115.—Chili.
rotundifolia, Bigel. Fl. Bost. ed. II. 352 = bellidifolia.
rotundifolia, Torr. & Gray, Fl. N. Am. i. 88, pro parte = rhomboidea.
rubifolia, Sm. ex DC. Syst. ii. 252 ; Prod. i. 150 = flaccida.
runcinata, Pourr. in Mém. Acad. Toul. iii. (1788) 310 = Cardamine, Sisymbrium hesperidiformis.
sarmentosa, *Forst. f. Prod. App.* 92, nomen.—Ins. Pacif.
saxatilis, Salisb. Prod. 369 = impatiens.
scaposa, *Franch. Pl. Davidian.* i. (1884) 33.—Mongol.
scaturiginosa, Wahlenb. ex Steud. Nom. ed. II. i. 281 = pratensis.
Schafneri, Hook. f. in Hemsl. Diagn. Fl. Nov. 2 = Gambelii.
scutata, *Thunb. in Trans. Linn. Soc.* ii. (1794) 332.—Japon.
sessilifolia, *Franch. & Sav. Enum. Pl. Jap.* ii. 250.—Japon.

CARDAMINE:—
serbica, Panč. ex Achers. & Kan. Cat. Corn. Serb. 76 = carnosa.
setigera, *Tausch, in Flora*, xix. (1836) 404.—Hab. ?
silvatica, Auct. vide *sylvatica*.
simensis, *Hochst. ex Oliver, Fl. Trop. Afr.* i. 61.—Abyssin.
Solisi, *Phil. in Anal. Univ. Chil.* (1856) 410.—Chili.
spathulata, *Michx. Fl. Bor. Am.* ii. 30.—Carolina.
stellata, Hook. f. Fl. Antarct. 7. t. 4 A = depressa.
stolonifera, Scop. Fl. Carn. ed. II. ii. 32 = Arabis stolonifera.
stolonifera, Tausch, Cat. 5 = pratensis.
striatula, *Steud. in Flora*, xxxix. (1856) 410.—Reg. Magell.
stylosa, *DC. Syst.* ii. 248.—Austral. ; N. Zel.
subipsata, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 78 = Nasturtium subhyratum.
subumbellata, *Hook. ex Hook. f. & T. Anders. in Fl. Brit. Ind.* i. 158.—Ind. or.
sylvatica, Bess. Primit. Fl. Galic. ii. 76 = pratensis.
sylvatica, Link, in Hoffm. Phyt. Blaett. i. (1808) 50 = hirsuta.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* i. 26.—Japon.
tenella, Clark, Trav. ii. 117 = hirsuta.
tenera, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 56 = pectinata.
tenera, *S. G. Gmel. ex C. A. Mey. Vers. Pfl. Casc.* 179.—Reg. Caucas. ; Persia.
tenuifolia, *Hook. Journ. Bot.* i. (1834) 347.—Austral.
tenuifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 238.—Sibir.
tenuirostris, *Hook. f. & Ars. Bot. Beech. Voy.* 6.—Chili.
teres, Michx. Fl. Bor. Am. ii. 29 = Nasturtium tanacetifolium.
ternata, Bory, ex DC. Syst. ii. 252 = africana.
tetrandra, *Hegetschw. Fl. Schw.* 641.—Helvet.
thalictroides, *All. Fl. Pedem.* i. 261.—Europ. austr.
tolimensis, *Planch. & Linden, in Ann. Sc. Nat. Sér.* IV. xvii. (1863) 39.—N. Granat.
trichocarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 18 = graeca.
trifolia, *Linn. Sp. Pl.* 654.—Europ. austr.
trifolia, Pall. Reise, iii. 316 = amara.
trifolia, Thunb. Fl. Jap. 260 = scutata.
trifoliata, Wahlenb. Fl. Lapp. 179 = amara.
trifoliolata, *Hook. f. & Thoms. in Journ. Linn. Soc. v.* (1861) 145.—Reg. Himal.
triphylla, Pall. Reise, ii. 35 = amara.
tuberosa, Bert. ex Steud. in Flora, xxxix. (1856) 409 = cognata.
tuberosa, *DC. Syst.* ii. 264.—Chili.
uliginosa, Jord. Diagn. i. 130 = pratensis.
uliginosa, *Bieb. Fl. Taur. Cauc.* iii. 488.—Ross. austr. ; Orient.
umbrosa, *Andrz. in DC. Syst.* ii. 260.—Volhynia.
umbrosa, Lej. Fl. Spa, ii. 62 = amara.
unalaschensis, Andrz. ex Ledeb. Fl. Ross. i. 127 = hirsuta.
undulata, *Laromb. & Timb. in Bull. Soc. Bot. Fr.* xiv. (1867) 69.—Gallia.
uniflora, Michx. Fl. Bor. Am. ii. 29 = Leavenworthia Michauxii.
valdiviana, *Phil. in Anal. Univ. Chil.* (1865) ii. 314.—Chili.
variabilis, *Phil. in Linnæa*, xxxiii. (1864–65) 5.—Chili.
violacea, *Wall. Cat.* n. 4782.—Reg. Himal.
virginica, Linn. Sp. Pl. 656 = Arabis indovicicana.
virginica, Michx. Fl. Bor. Am. ii. 29 = hirsuta.
viscosa, S. G. Gmel. ex DC. Syst. ii. 216 = Arabis albida.
Volckmanni, *Phil. in Linnæa*, xxxiii. (1864–65) 7.—Chili.
vulgaris, Phil. l. c. xxviii. (1856) 665 = pratensis.
Wichmanniana, *Boiss. Fl. Orient.* i. 166.—Anatolia.
Wightiana, Wall. Cat. n. 6780 = africana.
Willdenowii, Tausch, in Flora, xix. (1836) 403 = macrophylla.
yezoensis, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 277.—Japon.
Zellingeri, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 294.—Java ?

CARDAMINUM, *Moench*, Meth. 262 (1794) = Sisymbrium, *L. Ex. (Candia).*
Nasturtium, *Moench*, l. c. = *N. officinale*.

CARDAMOMUM, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 ; Salisb. in Trans. Hort. Soc. i. (1812) 393 = **Alsteria**, Matoe (Scitamin.).
maldaricum, Pritz. Ic. Ind. 207 = E. Cardamomum.
medium, Schult. Mant. i. 24 = Amomum costatum.
officinale, Salisb. in Trans. Hort. Soc. i. (1812) 282 = E. Cardamomum.

CARDAMON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 388 = **Lepidium**, Linn. (Crucif.).
sativum, Fourr. l. c. = L. sativum.

CARDANTHERA, Buch.-Ham. ex Nees, in DC. Prod. xi. 67 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1074.
ADENORMA, Nees, in Wall. Pl. As.˙Rar. iii. 75 (1839).
SYNNEMA, Benth. in DC. Prod. x. 538 (1846), nomen prius.
africana, *Benth. & Hook. f. Gen.* ii. 1075.—Afr. trop.
avana, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv.* 405.—Burma.
balsamica, *Benth. l. c.* 404.—Ind. or.
glutinosa, Buch.-Ham. ex Nees, in DC. Prod. xi. 68 = triflora.
Griffithii, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 404.—Assam ; Burma.
justicioides, *S. Moore, in Journ. Bot.* xviii. (1880) 8.—Afr. trop.
longifolia, Buch.-Ham. ex Nees, in DC. Prod. xi. 90 = Hygrophila phlomoides.
pinnatifida, *Benth. : C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 405.—Ind. or.
Thwaitesii, *Benth. & Hook. f. Gen.* ii. 1075.—Zeylan.
triflora, *Buch.-Ham. ex Nees, in DC. Prod.* xi. 68.—Ind. or. ; Malaya.
uliginosa, *Buch.-Ham. l. c.* 69.—Ind. or.
verticillata, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 405.—Ind. or.

CARDARIA, Desv. Journ. Bot. iii. (1813) 168 = **Lepidium**, Linn. (Crucif.).
amplexicaulis, Spach, Hist. Vég. Phan. vi. 544 = L. amplexicaule.
brachypetala, Opiz, Seznam, 26 = L. Draba.
cochlearia, Spach, Hist. Vég. Phan. vi. 542 = L. Draba.
crassifolia, Spach, l. c. 545 = L. crassifolium.
Draba, Desv. Journ. Bot. iii. (1813) 168 = L. Draba.
latifolia, Spach, Hist. Vég. Phan. vi. 546 = L. latifolium.

CARDERINA, Cass. in Dict. Sc. Nat. xxxv. 272 (1825) = **Senecio**, Tourn. (Compos.).
reclinata, Cass. l. c. xlviii. 447 = S. reclinatus.

CARDIA, Dulac, Fl. Hautes-Pyr. 387 (1867) = **Veronica**, Tourn. (Scrophularin.).
agrestis, Dulac, l. c. 388 = V. agrestis.
alpina, Dulac, l. c. 391 = V. alpina.
amplexicaulis, Dulac, l. c. 390 = V. Anagallis.
aphylla, Dulac, l. c. = V. aphylla.
arvensis, Dulac, l. c. 388 = V. arvensis.
Beccabunga, Dulac, l. c. 390 = V. Beccabunga.
cilista, Dulac, l. c. = V. Chamaedrys.
didyma, Dulac, l. c. 388 = V. didyma.
elliptica, Dulac, l. c. 390 = V. anagalloides.
filiformis, Dulac, l. c. 389 = V. Tournefortii.
fruticulosa, Dulac, l. c. 391 = V. fruticulosa.
latifolia, Dulac, l. c. 390 = V. latifolia.
montana, Dulac, l. c. 389 = V. montana.
multiflora, Dulac, l. c. 391 = V. serpyllifolia.
Nummularia, Dulac, l. c. 392 = V. Nummularia.
obtusifolia, Dulac, l. c. 391 = V. bellidioides.
officinalis, Dulac, l. c. 390 = V. officinalis.
orbiculata, Dulac, l. c. 388 = V. acinifolia.
plana, Dulac, l. c. 391 = V. Teucrium.
praecox, Dulac, l. c. 389 = V. praecox.
prostrata, Dulac, l. c. 391 = V. prostrata.
quadrifolia, Dulac, l. c. 389 = V. hederaefolia.
scutellata, Dulac, l. c. = V. scutellata.
serrata, Dulac, l. c. 391 = V. Persoonii.
spicata, Dulac, l. c. 392 = V. spicata.
verna, Dulac, l. c. 388 = V. verna.

CARDIACA, [Tourn.] Linn. Syst. ed. I (1735) = **Leonurus**, Linn. (Labiat.).
arvensis, Linn. Fl. Fr. ii. 388 = Stachys arvensis.

CARDIACA :—
crispa, Moench, Meth. 401 = Leonurus Cardiaca.
glabra, Gilib. Fl. Lituan. i. 85 = L. tataricus.
Marrubiastrum, Medic. Beobacht. (1783) 127 = L. Marrubiastrum.
quinquelobata, Gilib. Fl. Lituan. i. 84 = L. Marrubiastrum.
Stachys, Medic. Beobacht. (1783) 126 = L. Cardiaca.
sylvatica, Lam. Fl. Fr. ii. 384 = Lamium Galeobdolon.
tricuspis, Lam. l. c. 383 = L. Cardiaca.
vulgaris, Moench, Meth. 401 = L. Cardiaca.

CARDIACANTHUS, Schau. in Linnaea, xx. (1847) 714 ; et in DC. Prod. xi. 681 (1847) = **Jacobinia**, Moric. (Acanthac.).
Nessianus, Schau. l. c. 715 = J. Nessiana.

CARDIANDRA, Sieb. & Zucc. Fl. Jap. 119. t. 65, 66 (1835). *SAXIFRAGEAE*, Benth. & Hook. f. i. 648.
alternifolia, *Sieb. & Zucc. l. c.* 121.—Japon.

CARDIANTHERA, Hance, in Journ. Linn. Soc. xiii. (1873) 134, sphalm. = **Cardanthera**, Buch.-Ham.

CARDINALIS, Rupp. Fl. Jen. ed. Hall. 248 (1745) = **Lobelia**, Linn.

CARDIOCARPUS, Reinw. in Syll. Ratisb. ii. (1828) 14 = **Soulamea**, Lam. (Simarub.).
amara, Reinw. l. c. = S. amara.

CARDIOCHLAMYS, Oliver, in Hook. Ic. Pl. t. 1408 (1888). *CONVOLVULACEAE*.
madagascariensis, *Oliver, l. c.*—Madag.

CARDIOCRINUM, Lindl. Veg. Kingd. 205 (1847) = **Lilium**, Tourn.

CARDIOGYNE, Bar. in DC. Prod. xvii. 282 (1873) = **Pleocospermum**, Trèc. (Urticac.).
africana, Bur. l. c. 288.—Afr. trop.

CARDIOLEPIS, Rafin. Neogen. 1 (1825) = **Rhamnus**, Tourn.
nigra, Rafin. l. c. 1 = R. lanceolata.
obtusa, Rafin. Sylva Tellur. 25 = R. Purshiana.
rubra, Rafin. Neogen. 1 = R. lanceolata.
spinosa, Rafin. l. c.—Am. bor.

CARDIOLEPIS, Wallr. Sched. Crit. 349 (1822) = **Lepidium**, Linn. (Crucif.).
denitata, Wallr. l. c. = L. Draba.

CARDIOLOCHIA, Rafin. ex Reichb. Consp. 85 (1828) = **Aristolochia**, Tourn.

CARDIOLOPHUS, Griff. in Madr. Journ. Sc. iv. (1836) 276 = **Herpestis**, Gaertn. (Scrophular.).
decussata, Griff. l. c.= H. Hamiltoniana.

CARDIONEMA, DC. Prod. iii. 372 (1828) = **Pentaosena**, Bartl. (Illecebrac.).
muticuale, DC. l. c. = P. polycnemoides.

CARDIOPETALUM, Schlecht. in Linnaea, ix. (1834) 228 = **Duguetia**, St. Hil. (Anonac.)
calophyllum, Schlecht. l. c.—Bras.

CARDIOPHORA, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 216 = **Soulamea**, Lam. (Simarub.).
Hindsii, Benth. l. c. = S. amara.

CARDIOPTERIS, Wall. Cat. n. 8083 (1886 ?). *OLACINEAE*, Benth. & Hook. f. i. 353.
PERIPTERYGIUM, Hamk. in Hoev. & De Vriese, Tijdschr. x. (1843) 142.
Stoja, Buch.-Ham. ex Lindl. Introd. Nat. Syst. Bot. ed. II. 92 (1836).
hamulosa, Griff. Notul. iv. 542 ; Ic. Pl. Asiat. t. 598 = lobata.
javanica, Blume, Rumphia, iii. 206 = lobata.
lobata, *Wall. Cat.* n. 8083.—Ind. or.
moluccana, Blume, Rumphia, iii. 207 = lobata.
quinquelobata, Hamk. Retzia, i. 84 = lobata.
Rumphii, Baill. Adansonia, x. (1871–72) 280 = lobata.
subhamata, Wall. ex Baill. in DC. Prod. xvii. 26 = lobata.

CARDIOSPERMUM, Linn. Syst. ed. I (1735). *SAPINDACEAE*, Benth. & Hook. f. i. 398.
acuminatum, Miq. in Linnaea, xvii. (1843) 579 = microcarpum.
altissimum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 38.—Bras.
anomalum, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 351 —Bras.
barbicaule, *Baker, in Oliver, Fl. Trop. Afr.* i. 418.—Afr. trop.
canescens, *Wall. Pl. As. Rar.* i. 14.—As. et Afr. trop.
clematideum, *A. Rich. Tent. Fl. Abyss.* i. 100.—Abyss.
coluteoides, *H. B. & K. Nov. Gen. et Sp.* v. 100.—Am. austr.
Corindum, Linn. Sp. Pl. 526 = C. Halicacabum.
corycedes, *Kunze, in Linnaea*, xviii. (1844) 359.—Cuba.
Dusrtenum, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 349. —Bras.
elegans, *H. B. & K. Nov. Gen. et Sp.* v. 99.—Peruv.
ferrugineum, *A. Rich. Enc. Fl. Cub.* 276.—Cuba.
glabrum, Schum. & Thonn. Beskr. Guin. Pl. 197 = C. Halicacabum.
grandiflorum, *Sw. Prod. Veg. Ind. Occ.* 64.—Jamaica.
Halicacabum, *Linn. Sp. Pl.* 366.—Reg. trop.
hirsutum, *Willd. Sp. Pl.* iii. 467.—Afr. occ.
hispidum, *H. B. & K. Nov. Gen. et Sp.* v. 100.—Am. austr.
inflatum, Salisb. Prod. 279 = C. Halicacabum.
inflatum, *Vell. Fl. Flum.* 158 ; iv. t. 32.—Bras.
integerrimum, *Radlk. in Sitzb. Math.-Phys. Acad. Munich.* viii. (1878) 260.—Bras.
Lusarthianum, G. Don, Gen. Syst. i. 657, sphalm. = Duarteanum.
loxense, *H. B. & K. Nov. Gen. et Sp.* v. 104.—Peruv.
macrolophum, *Radlk. in Sitzb. Math.-Phys. Acad. Munich.* viii. (1878) 261.—Venezuela.
macrophyllum, *H. B. & K. Nov. Gen. et Sp.* v. 100.—Venezuela.
microcarpum, H. B. & K l. c. 104 = C. Halicacabum.
microspermum, E. Mey. in Drege, Zwei Pfl. Docum. 170 = C. Halicacabum.
mollis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 100 = clematideum.
molle, *H. B. & K. Nov. Gen. et Sp.* v. 104.—Mexic.
moniliferum, *Sw. ex Strud. Nova.* ed. II. i. 150.—Ind. occ.
oblongum; A. Rich. Tent. Fl. Abyss. i. 101 = canescens.
ovatum, Wall. Cat. n. 8081 = canescens.
pauciflorum, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 351. —Bras.
pilosum, *Vell. Fl. Flum.* 159 ; iv. t. 26.—Bras.
pilosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 895.—Bras.
procumbens, *Radlk. in Sitzb. Math.-Phys. Acad. Munich.* viii. (1878) 262.—Bras.
pubescens, Griff. Journ. 108 = canescens.
pubescens, Lag. Gen. et Sp. Nov. 14 = molle.
Schmiedelia, Dalz. & Gibs. Bomb. Fl. 34 = Schmiedelia villosa.
striatum, *Radlk. in Sitzb. Math.-Phys. Acad. Munich.* viii. (1878) 262.—Bras.
tortuosum, *Benth. Bot. Voy. Sulph.* 9.—Calif.
triphyllum, *Vell. Fl. Flum.* 159 ; iv. t. 25.—Bras.
truncatum, A. Rich. Tent. Fl. Abyss. i. 101 = C. Halicacabum.
velutinum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 158.—Bras.
verticillatum, Humb. Rel. Hist. i. 89 = coluteoides.
villosum, Dalz. & Gibs. Bomb. Fl. 34 = Schmiedelia villosa.
villosum, *Macfad. Fl. Jamaic.* i. 154.—Ind. occ.
villosum, Mill. ex DC. Prod. i. 602 = C. Halicacabum.

CARDIOSTEGIA, Presl, Epim. Bot. 249 (1849) = **Malhania**, Forsk. (Sterculiac.).
Ketschyi, Presl, l. c. 249 = M. Deshasni.

CARDIOSTIGMA, Baker, in Journ. Linn. Soc. xvi. (1878) 102 = Sphenocostigma, Baker (Irid.).
longipetalum, Baker, l. c. = S. longipetalum.
nuda, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 288 = Calydorea nuda.

CARDIOTHECA, Ehrenb. ex Steud. Nom. ed. II. i. 382 (1840) = **Anarrhinum**, Desf. (Scrophularin.).
pubescens, Ehrenb. ex Steud. l. c. = A. pubescens.
virgatum, Ehrenb. ex Boiss. Fl. Orient. iv. 862 = A. orientale.

CARDOPATIUM, Juss. in Ann. Mus. Par. vi. (1805) 324 in adn. *COMPOSITAE*, Benth. & Hook. f. ii. 468.
BROTERA, Willd. Sp. Pl. iii. 2899 (1803).
amethystinum, *Spach, in Ann. Sc. Nat.* Sér. III. v. (1846) 343.—Afr. bor.
apulum, Spach, l. c. 340 = corymbosum.
Billardieri, Spach, l. c. 347 = corymbosum.
Boryi, *Spach, l. c.* 345.—As. Min.
corymbosum, *Perr. Sym.* ii. 500.—Reg. Mediterr. or.; As. Min.
Fontanesii, *Spach, in Ann. Sc. Nat.* Sér. III. v. (1846) 344.—Reg. Tunetan.
orientale, Juss. & Spach, Illustr. t. 496 = corymbosum.
orientale, *Spach, in Ann. Sc. Nat.* Sér. III. v. (1846) 337.—Graecia.
Vrienii, Heldr. in Boiss. Fl. Orient. iii. 443 = corymbosum.

CARDUNCELLUS, Adans. Fam. ii. 116 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 483.
LAMOTTEA, Pomel, Mat. Fl. Atl. 3 (1860 f).
ONOBROMA, Gaertn. Fruct. ii. 380. t. 160 (1791).
acaulis, Presl, Fl. Sic. i. p. xxx = pinnatus.
ARASCOSA, *Boiss. & Reut. Pl. Nov. Hisp. Diagn.* 18.—Hispan.
arborescens, Sweet, Hort. Brit. ed. I. 216 = Carthamus arborescens.
atlanticus, *Coss & Dur. ex Coss. in Ann. Sc. Nat.* Sér. IV. i. (1854) 240.—Algeria.
atractyloides, *Coss. & Dur. ex Pomel, Nouv. Mat. Fl. Atl.* 278.—Algeria.
caerulens, *Less. Syn. Comp.* 6.—Reg. Mediterr.
calvus, *Boiss. & Reut. Pugill.* 64.—Afr. bor.
dianius, Webb, Iter Hisp. 33 = caerulens.
eriocephalus, *Boiss. Diagn.* Ser. I. x. 100.—Afr. bor.
nispanicus, Boiss. ex DC. Prod. vii. 304 = caerulens.
ilicifolius, *Pomel, Nouv. Mat. Fl. Atl.* 277.—Afr. bor.
lucens, *Ball, in Journ. Bot.* xi. (1873) 370.—Marocco.
mitissimus, *DC. Fl. Fr.* iv. 78.—Gallia.
monspeliensis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = seq.
monspeliensium, *All. Fl. Pedem.* i. 154.—Gall. austr.
pectinatus, *DC. Prod.* vi. 615.—Afr. bor.
pinnatus, *DC. l. c.* 614.—Oriens.
plumosus, *Pomel, Nouv. Mat. Fl. Atl.* 277.—Afr. bor.
rhaponticoides, *Coss. & Dur. ex Pomel, Nouv. Mat. Fl. Atl.* 278.—Algeria.
tingitanus, DC. ex Duby, Bot. Gall. i. 381 = caerulens.
vulgaris, Loud. Hort. Brit. 382 = monspeliensium.

CARDUUS, (Tourn.) Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 467.
ALFREDIA, Cass. in Bull. Soc. Philom. (1815) 175.
CLAVENA, DC. Prod. vi. 623 (1837).
ONOPYXUM, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 319.
POLYACANTHA, Hill, Herb. Brit. i. 69 (17 9).
PTERNIX, Hill, Veg. Syst. iv. 18 (1762).
abyssinicus, *Sch. Bip. in Linnaea*, xix. (1847) 382.—Abyssin.
acanthifolius, Lam. Encyc. i. 708 = Cnicus oleraceus.
Acanthium, Beill. Hist. Pl. vii. 389 = Onopordon Acanthium.
acanthocephalus, *C. A. Mey. Verz. Pfl. Casc.* 74.—Reg. Caucas.
acanthoides, Gren. & Godr. Fl. Fr. ii. 231, sec. Nym. Consp. 411 = nutans.
acanthoides, Hornem. in Fl. Dan. t. 1341 = Cnicus palustris.
acanthoides, Huds. Fl. Angl. ed. II. ii. 351 = pycnocephalus.
acanthoides, *Linn. Sp. Pl.* 821.—Europ.; Reg. Caucas.
acanthoides, Pall. ex Bieb. Fl. Taur. Cauc. ii. 268 = hamulosus.
acanthoides, Urv. in Mém. Soc. Linn. Par. i. (1822) 361 = nutans.
acanthoides, *Benth. in Benth. & Hook. f. Gen.* ii. 468. —As. centr.
Acarna, Linn. Sp. Pl. 820 = Cnicus Acarna.
acaulis, Linn. l. c. 1199 = Cnicus acaulis.
acaulis, Thunb. Fl. Jap. 306 = Cnicus japonicus.
acicularis, *Bertol. in Ann. Sc. Nat.* Sér. I. i. (1829) 374.—Italia.
acuminatus, Gaud. ex Reichb. Ic. Fl. Germ. xv. 90. t. 873 = acanthoides.
adpressus, *C. A. Mey. Verz. Pfl. Casc.* 71.—Reg. Caucas.

CARDUUS:—
afer, Jacq. Hort. Schoenb. ii. 10 = Cnicus afer.
affinis, *Guss. Pl. Rar.* 884. t. 57.—Neapol.
aggregatus, Sehlsch. Cat. Pl. Helv. ed. IV. 11 = nutans?
agrestis, Presl, Fl. Sicul. p. xxx, in nota = corymbosus.
Aitonii, Seaton, Nom. ed. III. 239 (Quid?).
alatus, D. Don, Prod. Fl. Nep. 167 = Jurinea alata.
albidus, Bieb. Fl. Taur. Cauc. ii. 269 = pycnocephalus.
albidus, Lam. Fl. Fr. ii. 15 = pratc.?
algeriensis, Munby, in Bull. Soc. Bot. Fr. ii. (1855) 285. —Algeria.
alpestris, Waldst. & Kit. Pl. Rar. Hung. iii. 296. t. 267 = arctioides.
altaicus, Patrin, ex S. G. Gmel. Fl. Sib. t. 39. f. 2 = Serratula glauca.
altissimus, Gilib. Fl. Lituan. iii. 186 = Cnicus palustris.
altissimus, Linn. Sp. Pl. 824 = Cnicus altissimus.
amblyrepus, Pers. Syn. ii. 389 = Cnicus heterophyllus.
ammophilus, *Hoffegg. & Link, Fl. Port.* ii. 187.—Lusitan.
amplexicaulis, *Nevoska, in Verh. Batow. Gen.* v. (1790) ed. I. Art. iv. 11.—Malaya.
Amstenii x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.-xxiv. (1880) 106.—Helvet.
anglicus, Lam. Encyc. i. 705 = Cnicus pratensis.
angustifolius, *Lam. l. c.* 708.—Hab.?
antarcticus, Auct. ex Steud. Nom. ed. I. 153 = autarcticus.
apherusius, Jan, ex DC. Prod. vi. 622 = nutans.
arabicus, Jacq. Coll. i. 56 = pycnocephalus.
arachnoideus, Bieb. ex Steud. Nom. ed. II. i. 288 = Cnicus arachnoideus.
aranuosus, Stend. Nom. ed. I. 151 = Cnicus aranosus.
arctioides, Vill. Hist. Pl. Dauph. ii. 93 = Personata.
arctioides, *Willd. Sp. Pl.* iii. 1656.—Europ.
arenarius, DC. Fl. Fr. vi. 457 = Candollei.
arenarius, *Desf. Fl. Atlant.* ii. 99.—Afr. bor.
Argemone, Fourr. in Lam. Encyc. i. 700 = carlinaefolius.
argentatus, *Linn. Mant.* ii. 280.—Reg. Mediterr. or.; Mesopot.
argyreolatus, Willd. ex Steud. Nom. ed. II. i. 288 = pycnocephalus.
argyrus, Nym. Consp. 413 = argyros.
argutus, Sweet, Hort. Brit. ed. I. 213 = Serratula tinctoria.
argyracanthus, Wall. Cat. n. 2908 = Cnicus Wallichii.
argyros, *Bivona, Stirp. Rar. Sicil.* i. [7].—Sicilia.
argyrus, Kunze, in Flora, xxix. (1846) 760 = malacitanus.
armatus, Boiss. & Heldr. Diagn. Ser. I. vi. 104 = cronius.
armatus, Steud. Nom. ed. I. 151 = Cnicus armatus.
armenus, Boiss. Fl. Orient. iii. 516 = nutans.
armenus, Rohb. Brit. Fl. 163 = Cnicus arvensis.
Aschersonianus x, *Rohb. in Bichl. Yahrb. Berl.* i. (1881) 229.—Germaniae.
astracanicus, *Spreng. Fl. Hal. Mant.* 49.—Astracan.
atlanticus, *Pomel, Nouv. Mat. Fl. Atl.* 22.—Afr. bor.
atriplicifolius, Trevir. Hort. Wratisl. 1820 = Serratula atriplicifolia.
auriculatus, Wall. Cat. n. 2899 = Saussurea hypoleuca.
auriculus, Chaix, ex Vill. Fl. Delph. 90 sphalm. = seq.
auroticus, *Vill. Hist. Pl. Dauph.* iii. 7.—Gallia.
australis, Jord. Gr. Rec. Bot. Gren. (1849) 14 = tivrariensis.
australis, Linn. f. Suppl. 346 = pycnocephalus.
autarcticus, Vill. Hist. Pl. Dauph. iii. 13 = Cnicus heterophyllus.
auillarii, Gaud. Fl. Helv. ii. 169 = acanthoides.
baccophilus, *Webb & Berth. Phyt. Canar.* ii. 278.—Ins. Canar.
baeticus, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 63.—Hispan.
Balansae, *Boiss. Diagn.* Ser. II. iii. 44.—Algeria.
Ballii, *Hook. f. in Journ. Bot.* xi. (1873) 363.—Marocco.
Barrelieri, *Bertol. ex Nym. Consp.* 411.—Liguria.
benedictus, Auct. ex Steud. Nom. ed. I. 151 = Carbenia benedicta.
bicolor, Visiani, Fl. Dalm. ii. 43. t. 49. fig. 2 = argyros.
biharicus x, Simk. in Termész. Füzetek, v. (1881) 51. —Transylv.

CARDUUS:—
bojnicti, Pill. & Mitterp. Iter Poseq. i. 143. t. 13 = Cnicus arachnoideus.
Bourgaei, Savi, Fl. Pis. ii. 248 = Cnicus tenuis.
Bourgaeanus, Boiss. & Reut. Pugill. Pl. Nov. (1852) 62 = Reuterianus.
Bourgeanus, *Sch. Bip. ex Boiss. Diagn.* Ser. II. iii. 44. —Ins. Canar.
brachycephalus, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 58 = collinus.
bulbosus, Lam. Encyc. i. 705 = Cnicus tuberosus.
caeruleus, *Brot. Fl. Lusit.* i. 342.—Lusitan.
casuius, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 498, nomen.—Rossia.
Cattichii x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.-xxiv. (1880) 106.—Helvet.
callosus, *Steud. Nom.* ed. II. i. 283.—Hab.?
camporum, *Boiss. Diagn.* Ser. II. iii. 45.—Rumel.
canaliculatus, Dulac, Fl. Hautes-Pyr. 322 = carlinaefolius.
candicans, Waldst. & Kit. Pl. Rar. Hung. i. t. 88 = collinus.
Candollei, *Moretti, Pl. Ital. Dec.* ii. 10.—Gallia; Ital.
canus, Linn. Mant. 105 = Cnicus canus.
Candnelii, Gren. in Billotia, i. (1864) 14 = arctioides.
carlinaefolius, *Lam. Encyc.* i. 700.—Europ.
carlinaefolius, Nym. Consp. 413 = defloratus.
carlinaefolius, Tenore, Prod. Fl. Neap. 48 = chrysacanthus.
carlinoides, Dur. ex DC. Prod. vi. 620 = conferta.
carlinoides, *Gouan, Illustr.* 69. t. 28.—Mont. Pyren.
carlinoides, Steud. Nom. ed. II. i. 283 = Cnicus carlinoides.
carlinoides, Tenore, Prod. Fl. Neap. 48 = chrysacanthus.
carnuolicus, Pers. Syn. ii. 388 = Cnicus rufescens.
carolinianus, Wall. Fl. Carol. 195 = Cnicus muticus.
Carolorum x, Hoppe, in Jenner, in Trans. Bot. Soc. Edinb. ix. (1868) 260.—Scotia.
carpetanus, Boiss. & Reut. Diagn. Pl. Hisp. 19. n. 35 = Gayanus.
carthamoides, Steud. Nom. ed. I. 151 = Leuzea carthamoides.
Casabonae, Linn. Sp. Pl. 822 = Cnicus Casabonae.
cascacicus, Adam, in Weber & Mohr, Beitr. i. 64 = Cnicus horridus.
caulescens, Pers. ex Steud. Nom. ed. I. 151 = Cnicus acaulis.
centaurioides, Buch.-Ham. ex DC. Prod. vi. 643 = Volutarella divaricata.
centauroides, D. Don, Prod. Fl. Nep. 167 = Serratula centauroides.
centauroides, Hoppe, ex DC. Prod. vi. 628 = arctioides.
centauroides, Linn. Sp. Pl. ed. II. 1148 (= *Cnicus cynaroides*).
cephalanthus, *Viv. Fl. Cors. Sp. Nov.* 14.—Sardin.
cerinthifolius, Vill. Prosp. 20 = Serratula nudicaulis.
cerinthoides, Willd. Sp. Pl. iii. 1660 = Serratula nudicaulis.
cernuus, *Bertol. Fl. Guatimal.* 32.—Guatemal.
cernuus, *Steud. Nom.* ed. I. 151.—Sibir.
Chaillei, Godr. Fl. Lorr. ii. 49 = Cnicus palustris.
chamaecephalus, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 434.—Abyssin.
chionaeis, DC. Prod. vi. 629 = Cnicus chionaeis.
chlus, *Spag. Hort. Vindel.* iii. 7.—Ins. Chios.
chrysacanthus, *Tenore, Sem. Hort. Neap.* (1826) 13.—Hispan.; Italia.
cichoraceus, Cav. Fl. Rar. Neap. Fasc. ii. 17. t. 7 = Serratula cichoracea.
ciliatus, Murr. in Comm. Gott. (1784) 35. t. 5 = Cnicus ciliatus.
ciliatus, Poir. ex Willd. & Lange, Prod. Fl. Hisp. ii. 198 = Gayanus.
ciliatus, Vill. Prin. Voy. 45 = Cnicus spathulatus.
cinnamomeus, *Hort. Huber, in Rev. Horticole.* (1872) 60 —Am. bor.!
cinereus, Bieb. Fl. Taur. Cauc. ii. 270 = pycnocephalus.
circilensis, *Vahnl. in Rad Jugosc. Akad. Zagreb.* xxxix. (1877) 206.—Europ.
circioides, Vill. Prosp. 20 = defloratus.
Clementa, Mill. Gard. Dict. ed. VIII. n. 5 = Cnicus diffusus.
clavulatus, *Link, in Buch, Beschr. Canar. Inss.* 247.—Ins. Canar.
collinus, *Waldst. & Kit. Pl. Rar. Hungr.* iii. 287. t. 265.—Europ. austr. et.

CARDUUS :—
comosus, Lam. Encyc. i. 703 = Cnicus spinosissimus.
confertus, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 194 = hastiens.
confertus, Moris, Elench. Pl. Sard. ii. 5 = myriacanthus.
congestus, Gmel. in Tenore, Syll. 412 = cephalanthus.
corlarmiais, *Timb. & Theven. in Mém. Acad. Toulouse*, Sér. VII. vi. (1874) 647.—Europ.
coronatus, D. Don, Prod. Fl. Nep. 167 = Serratula coronata.
corsicus, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276 = clavulatus.
corymbosus, *Tenore, Prod. Fl. Neap.* 48.—Italia.
Cosmeli, Adam, ex DC. Prod. vi. 637, 688 = Cnicus lappaceus.
crassifolius, Willd. Enum. Hort. Berol. 886 = defloratus.
creticus, Lam. Encyc. i. 701 = Cnicus pangens.
crispus, Gouan, Fl. Monsp. 327 = tenuiflorus.
crispus, Guir. ex Nym. Consp. 418 = Resterianus.
crispus, Hnds. Fl. Angl. ed. II. ii. 350 = acanthoides.
crispus, *Linn. Sp. Pl.* 821.—Europ. ; Aa. bor.
cronius, *Boiss. & Heldr. Diagn.* Ser. I. vi. 105.—Graecia ; As. Min.
cyanoides, Linn. Sp. Pl. 822 = Jurinea cyanoides.
cyanoides, Pall. Reise, iii. 594 = Jurinea polyclonos.
Cynara, Lam. ex Gmel. Stend. Nom. ed. II. i. 358 = Cnicus cynaroides.
cynaroides, Lam. Encyc. i. 702 = Cnicus cynaroides.
Decandollei, Auct. ex Steud. Nom. ed. II. i. 388 = Candollei.
deflоratus, *Linn. Syst. ed.* X. 1200.—Europ.
demidoviarrensx, Franch. Fl. Cher-et-Loire, 317.—Gallia.
dentatus, Pers. Syn. ii. 389 = Kentrophyllum dentatum.
desertorum, Fisch. ex Steud. Nom. ed. II. i. 388 (cf. *Cirsium desertorum*).—Sibir.
discanthe, Labill. Ic. Pl. Syr. dec. i. 7 = Cnicus discanthus.
discolor, Hook. Fl. Bor. Am. i. 302 = Cnicus Hookerianus et undulatus.
discolor, Nutt. Gen. Am. ii. 130 = Cnicus altissimus.
dissectus, Linn. Sp. Pl. 822 = Cnicus dissectus.
dissectus, Vill. Hist. Pl. Dauph. iii. 15 = Cnicus pratensis.
divaricatus, Gillh. *Fl. Lituan.* i. 187 = Cnicus lanceolatus.
diversifolius, *Wall. Beitr. Fl. Hercyn.* 367.—Europ.
dubius x, Balb. Cat. Hort. Taur. (1813) = crispus.
Durieui, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 64.—Algeria.
echinatus, Desf. Fl. Atlant. ii. 247 = Cnicus echinatus.
echinocephalus, Pers. ex Stend. Nom. ed. I. 353 = Cnicus echinocephalus.
elegans, *Pall. ex Georgi, Beschr. Russ. Reich.* iii. IV. 1224.—Rossia.
elegans, Salisb. Prod. 188 = Cnicus Cambonae.
elegans, Stephan, ex Ledeb. Fl. Ross. ii. 678 = Cousinia wolgensis.
elegans, Willd. ex Stend. Nom. ed. II. i. 388 = Cnicus iguarica.
elongatus, Wall. Cat. n. 2900 = Tricholepis elongata.
eriophorus, Linn. Sp. Pl. 822 = Cnicus eriophorus.
eriophorus, Pall. ex Bieb. Fl. Taur. Cauc. ii. 276 = Cnicus fimbriatus.
eriophorus, Thunb. Fl. Jap. 305 = Cnicus japonicus.
Erithales, Jacq. Enum. Vindob. 146, 279 = Cnicus ochroleucus.
Erithales, Vill. Hist. Pl. Dauph. ii. 26 = Cnicus rivularis.
eruciginosus, Lam. Encyc. i. 764 = Cnicus ochroleucus ?
estroalcuicus, Boiss. Diagn. Ser. I. x. 90 = argematicus.
fasciculiflorus, *Viv. App. Alt. Fl. Cors.* 8. t. 1.—Corsica ; Sardinia.
fastigiatus, Moris, Elench. Fl. Sard. ii. 5 = argyrus.
ferox, Vill. Prosp. 30 ; Lam. Encyc. i. 702 = Cnicus ferox.
fimbriatus, Steud. Nom. ed. II. i. 384 = Cnicus fimbriatus.
firmus, Steud. l. c. ed. I. 352 = Cnicus firmus.
flavescens, *C. Koch, in Linnaea*, xxiv. (1851) 994.—Ponina.
flavescens, Linn. Sp. Pl. 826 = Serratula flavescens.
foliosus, Hook. Fl. Bor. Am. i. 306 = Cnicus foliosus.
Forsteri x, Bab. Man. Brit. Bot. ed. I. 178.—Anglia.
fruticosus, Desf. Tabl. Mort. Fav. ed. I. 91 = Cnicus fruticosus.

CARDUUS :—
fruticosus, Poir. Encyc. Suppl. ii. 199 = Carlina xeranthemoides.
Galactites, Bory & Chaub. Fl. Pélop. 56 = Galactites tomentosa.
Galactites, Lam. Encyc. i. 701 = pycnocephalus.
Gayanus, *Dur. ex DC. Prod.* vi. 636.—Hispan.
gelsilus, *Pomel, Nouv. Mat. Fl. Atl.* 275.—Afr. bor.
giganteus, Desf. Fl. Atlant. ii. 25. t. 221 = Cnicus giganteus.
Gigas, Uria, in Roem. Arch. i. 70 = Cnicus giganteus.
gleber, Buch.-Ham. ex Wall. Cat. n. 2907 = Tricholepis radicans.
glaber, Nutt. Gen. Am. ii. 129 = Cnicus muticus ?
glaber, Steud. Nom. ed. I. 352 = Cn. spinosissimus ?
glaucophyllus, Desf. ex DC. Prod. vi. 638 = deflоratus.
glaucus, Baumg. Transs. iii. 58 = deflоratus.
glaucus, Cav. Ic. iii. 13. t. 236 = Serratula radicalis.
glaucus, *Ledeb. Hort. Dorp. Suppl.* i. 2.—Rossia.
glomeratus, Lam. Fl. Fr. ii. 20 = Cnicus pratensis.
glutinosus, Beck, Bot. N. & Mid. St. 178 = Cnicus muticus.
glycacanthus, Sibth. & Sm. Fl. Graec. Prod. ii. 150 = Jurinea glycacantha.
Gmelini, Steud. Nom. ed. I. 352 = Cnicus acaulis.
gnaphaloides, Cyr. Pl. Rar. Neap. Fasc. i. 27 = Cnicus gnaphaloides.
Gouani, Steud. Nom. ed. I. 352 = Cnicus tuberosus.
granatensis, Willk. in Linnaea, xxx. (1859–60) 118 = platypus.
Grenieri, Sch. Bip. ex Nym. Consp. 412 = C. Personata.
haenseleri-undalis, Auct. ex DC. Prod. vi. 643 = Cnicus arvensis.
hamulosus, *Ehrh. Beitr* vii. 166.—Europ. austr. or. ; Reg. Cauc.
hamulosus, Tenore, Prod. Fl. Neap. 48 = nutans.
hastatus, Lam. Encyc. i. 704 = Cnicus heterophyllus.
helenifolius, Salisb. Prod. 185 = Cnicus heterophyllus.
helenioides, Linn. Sp. Pl. 825 = Cnicus heterophyllus.
heteromallus, D. Don, Prod. Fl. Nep. 166 = Saussurea candicans.
heterophyllus, Linn. Sp. Pl. 824 = Cnicus heterophyllus.
Hippolyti, Bory & Chaub. Exp. Mor. 241. t. 29 = Cnicus pungens.
hispanicus, Bory, ex DC. Prod. vii. 804 = carlinoides.
hispanicus, Lam. Encyc. i. 701 = Cnicus hispanicus.
horridulus, Pers. Syn. ii. 390 = Cnicus horridulus.
horridus, Adam, in Weber & Mohr, Beitr. i. 66 = Cnicus muticus.
horridus, Steud. Nom. ed. I. 352 = Cnicus ochroleucus.
hymenolepis, *Transc. in Act. Hort. Petrop.* iii. ii. (1875) 273.—Daghestan.
hyssopifolius, Schur, Enum. Pl. Transs. 895, Index = Serratula heterophylla.
Hystrix, *C. A. Mey. Verz. Pfl. Cauc.* 71.—Persia.
ignitarius, Pall. ex DC. Prod. vi. 657 = Cnicus ignitarius.
illyricus, Bail. Hist. Pl. viii. 289 = Onopordon illyricum.
inclinans, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. iii. 870 = crispus.
inclinatus, Balac, Fl. Hautes-Pyr. 532 = medius.
indicus, Klein, ex Wall. in DC. Prod. vi. 564 = Tricholepis angustifolia.
inermis, Pers. Syn. ii. 388 = Cnicus centauroides.
intermedius, Heller, ex Steud. Nom. ed. I. 352 = Cnicus intermedius.
introcutus, Reichb. f. Ic. Fl. Germ. xv. 91. t. 866 = Candollei.
intricatus, *Timb. in Bull. Soc. Bot. Fr.* xv. (1868) p. lxxxvi.—Mont. Pyren.
involucratus, Wall. Cat. n. 2902 = Cnicus involucratus.
italicus, Savi, Bot. Etrur. iii. 140 = Cnicus italicus.
Jacquini, Steud. Nom. ed. I. 352 = Cnicus pyrenicus.
japonicus, Franch. Pl. Davidian. i. 178 = Cnicus japonicus.
Kalbinos, *Pomel, Nouv. Mat. Fl. Atl.* 275.—Afr. bor.
Killiasi x, *Brueggr. in Yahresb. Naturf. Ges. Graub.* II. xxiii.–xxiv. (1880) 106.—Europ.
Kosmidii, Adam, in Weber & Mohr, Beitr. i. 66 = Cnicus lappaceus.
laciniatus, Lam. Encyc. i. 706 = Cnicus palustris.
lacinulatus, Lodd. Fl. Brit. ii. 732 = adpressus.
lanceolatus, Stokes, Bot. Mat. Med. iv. 141 = Silybum Marianum.
lanatus, *Brot. Fl. Lusit.* 342.—Lusitan.

CARDUUS :—
lanatus, ? Roxb. ex Poir. Encyc. vi. 565 = Cnicus arvensis.
lanceatus, Hill, Veg. Syst. iv. 14 = Cnicus lanceolatus.
lanceolatus, Linn. Sp. Pl. 821 = Cnicus lanceolatus.
lanceolatus, Lour. Fl. Cochinch. ii. 538 = chinensis.
lanifloras, Steud. Nom. ed. I. 352 = Cnicus lanifloras.
lanuginosus, *Willd. Sp. Pl.* iii. 1648.—As. Min.
lappaceus, Bieb. Casp. 193 = Cnicus lappaceus.
lappoides, Less. in Linnaea, v. (1830) 129 = Cnicus lappoides.
latifolius, Banks, ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 28 (= *Cirsium latifolium*).—Ins. Madera.
latiflorus, *Tausch, in Flora*, xii. (1829) Erg. 41.—Austria.
leptacanthus, *Fresen. in Mus. Senckenb.* iii. (1845) 70.—Abyssin.
leptolenlus, *Dur. in Duch. Rev. Bot.* i. (1845–46) 302.—Algeria.
leptophyllus, Gaud. Fl. Helv. v. 171 = deflоratus.
leucanthemos, Schur, Enum. Pl. Transs. 417 = C. Personata.
leucanthus, Cav. Ic. ii. 52. t. 165 = Serratula flavescens.
leucocephalus, Pers. Syn. ii. 387 = Cnicus leucocephalus.
leucographus, Linn. Sp. Pl. 820 = Tyrimnus leucographus.
leucophyllus, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 194.—Mongolia.
linearis, Thunb. Fl. Jap. 305 = Cnicus linearis.
litigiosus, Moris, Elench. Pl. Sard. i. 36 = pycnocephalus.
litigiosus, Nocca & Balb. Fl. Ticin. ii. 98. t. 12 = Candollei.
littoralis x, *Borb. in Magyar Növeny. Lapok*, ix. (1885) 59.—Europ.
Lomonosowii, *Transle. in Act. Hort. Petrop.* i. (1871–72) 168.—Mongolia.
lucidus, Moench, Meth. 554 = pycnocephalus.
lucidus, Wall. Cat. n. 2895 = nutans.
lycopifolius, Vill. Prosp. 30 = Serratula heterophylla.
macranthus, Habl. ex Boiss. Fl. Orient. iii. 496 = Cousinia Hablitzii.
macrocephalus, Cess. in Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 198 = chrysacanthus.
macrocephalus, Desf. Fl. Atlant. ii. 245 = nutans.
macrocephalus, Wall. Cat. n. 2904 = Jurinea macrocephala.
macrolepis, Peterm. in Flora, xxvii. (1844) 492 = nutans ?
macropterus, Fisch. ex Spreng. Pugill. i. 54 = Jurinea polyclonos.
maleitanus, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 92.—Hispan.
Marisc, Crantz, Inst. i. 248 = Silybum Marianum.
Marianus, Linn. Sp. Pl. 823 = Silybum Marianum.
marmoratus, *Boiss. & Heldr. Diagn.* Ser. II. iii. 48.—Peloponnes.
Martrinii, Timb. in Martr. Fl. Tarn, 377 = nigrescens.
medius, *Gouan, Illustr.* 62. t. 24.—Mont. Pyren.
meditus, Steud. Nom. ed. I. 352 = Cnicus tuberosus.
meonanthus, *Hoffmgg. & Link, Fl. Port.* ii. 136.—Hispan.
mexicanus, Moric. ex DC. Prod. vi. 637 = Cnicus mexicanus.
microcephalus, Gaud. Fl. Helv. v. 168 = tenuiflorus.
microcephalus, *Tenore, Syll. App.* iv. *ex Fl. Nap.* v. 206.—Italia.
mitis, Steud. Nom. ed. I. 353 = Cnicus mitis.
mollis, Bieb. Fl. Taur. Cauc. ii. 272 = Jurinea arachnoides.
mollis, Gouan, Illustr. 68 = Jurinea humilis.
mollis, Linn. Cent. Pl. Rar. ii. 30 = Jurinea mollis.
mollis, Pollich, Fl. Palat. ii. 421 = Jurinea cyanoides.
mollis, Vill. Hist. Pl. Dauph. ii. 17 = Cnicus acaulis.
monocephalus, *Schur, in Verh. Siebenb. Ver. Naturw.* iii. (1852) 87.—Europ.
monspessulanus, Linn. Sp. Pl. 822 = Cnicus monspessulanus.
montanus, Pers. Syn. ii. 388 = Cnicus montanus.
montanus, Steff. ex DC. Prod. vi. 636 = Candollei.
montanus, Pollini, Viag. Lag. Gard. 115 = deflоratus.
Morisii, Balb. in Moris, Elench. Pl. Sard. 36 = fasciculiflorus.
Mortisii x, Brueggr. in Jahresb. Naturf. Ges. Graub. II. xxiii.–xxiv. (1880) 106.—Europ.

CARDUUS:—
meschatus, Gmel. Ind. Sem. Hort. Boccadif. (1825) = Jurinea mollis.
multiflorus, Gand. Fl. Helv. v. 166 = acanthoides.
multiflorus, Pand. ex Nym. Consp. 414 = Pascicli.
multiflorus, C. Koch, in Linnæa, xxiv. (1851) 392 = adpressus.
mutieus, Pers. Syn. ii. 386 = Cnicus muticus.
myriacanthus, Boiss. Voy. Bot. Espagne, ii. 361 = Reuterianus.
myriacanthus, *Salsm. in DC. Prod.* vi. 634.—Reg. Mediterr. occ.
Nagueli x, Bragg. in Jahresb. Nat. Ges. Graub. II. xxiii.–xxiv. (1880) 107.—Europ.
neglectus, Steud. Nom. ed. I. 158 = Cnicus uliginosus.
neglectus, Tenore, Ind. Sem. Hort. Neap. (1830) 14 = acicularis.
nepalensis, Spreng. ex DC. Prod. vi. 642 = Jurinea macrocephala.
nervosus, *C. Koch, in Linnæa*, xvii. (1843) 42.—Armenia.
nidulans, *Rupr. Sert. Tianach.* 55.—As. centr.
nigrescens, Mutel, ex Nym. Consp. 412 = vivariensis.
nigrescens, *Vill. Prosp.* 30.—Europ.
nitidus, Waldst. & Kit. Pl. Rar. Hung. i. 52 = Serratula heterophylla.
nitidus, Wall. Cat. n. 2908 = Trichelopis angustifolia.
nivus, *Benth. in Benth. & Hook. f. Gen.* ii. 468.—As. centr.
nudicaulis, Pers. Syn. ii. 386 = Serratula nudicaulis.
numidicus, *Coss. & Dur. Explor. Pl. Algér.* t. 49.—Algeria.
nutans, Boiss. ex Nym. Consp. 411 = platypus.
nutans, *Linn. Sp. Pl.* 821.—Europ. ; As. temp.
obvallatus, *Bieb. ex Willd. Enum. Hort. Berol.* 849.—Reg. Caucas.
obvallatus, Steud. Nom. ed. I. 158 = Cnicus obvallatus.
obvallatus, Wall. Cat. n. 2905 = Saussurea obvallata.
occidentalis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 418 = Cnicus occidentalis.
ochroleucus, Pers. Syn. ii. 389 = Cnicus ochroleucus.
odoratus, Muhl. Cat. Dart. Fl. Cestr. 85 = Cnicus pumilus.
oleraceus, Vill. Hist. Pl. Dauph. iii. 21 = Cnicus oleraceus.
olympicus, *Boiss. Diagn.* Ser. II. iii. 48.—As. Min. onopordoides, *Fisch. ex Bieb. Fl. Taur. Cauc.* iii. 552. —Reg. Caucas.
Onopyrox, *Hill, Veg. Syst.* iv. 13.—German.
orgyalis, Steud. Nom. ed. I. 158 = Cnicus orgyalis.
orientalis, Adam, in Weber & Mohr, Beitr. i. 68 = Cousinia cacuminosa.
orientalis, Pers. Syn. ii. 387 = Galactites tomentosa.
orthocephalus x, Wallr. Beitr. Fl. Hercyn. 268 ; et in Linnæa, xiv. (1840) 68 = acanthoides.
osseticus, Adam, in Weber & Mohr, Beitr. i. 65 = Cnicus fimbriatus.
palustris x, Sch. Bip. ex Nym. Consp. 413 = acanthoides.
palustris, Linn. Sp. Pl. 822 = Cnicus palustris.
palustris, Savi, Bot. Etrusc. iii. 141 = Cnicus polyanthemus.
Pancicii, *Sch. Bip. ex Nym. Consp.* 414.—Serbia.
pancicalatus, *Ait. Hort. Kew.* ed. I. iii. 143.—Europ.
paniculatus, Dalac, Fl. Hautes-Pyr. 522 = carlinoides.
paniculatus, Vahl, Symb. Bot. i. 56 = Cnicus flavispinus.
pannonicus, Linn. f. Suppl. 346 = Cnicus pannonicus.
pannonicus, Schleich. ex DC. Prod. vi. 629 = tenuifolius.
pannosus, *Trautv. in Act. Hort. Petrop.* iii. ii. (1875) 271.—Armenia.
parviflorus, Buch, Beschr. Ins. Cæsar. 147 = pycnocephalus.
parviflorus, Linn. Mant. 279 = Cnicus parviflorus.
pauciflorus, Pers. Syn. ii. 386 = Cnicus pauciflorus.
pectinatus, Linn. Mant. 279 = deflexatus.
pedemontanus, Pers. Syn. ii. 389 = Cnicus tuberosus.
peregrinus, Retz. Obs. i. 27 = pycnocephalus.
Perezanus, *Pacq. Fl. Austr. iv.* 25. t. 548.—Europ.
perezanus, Guerin. Pract. ii. 278 = præc.
petrophilus, *Timb. in Bull. Soc. Bot. Fr.* xv. (1868) p. lxxxvii.—Mont. Pyren.
pinnatifidus, Cav. Ic. i. 55. t. 83 = Serratula pinnatifida.
Pitcheri, Steud. Nom. ed. II. i. 284 = Cnicus Pitcheri.
platylepis, Sauter, in Flora, xiii. (1830) 410 = nutans.
platypus, *Lange, Ind. Sem. Hort. Havn.* (1857) 36.—Hispan.
podacanthus, DC. Fl. Fr. iii. 80 = sarcodus.

CARDUUS:—
poliochrus, *Trautv. in Act. Hort. Petrop.* iii. ii. (1875) 272.—Reg. Caucas.
polyacanthus, Cart. Fl. Lond. fasc. vi. t. 54 = acanthoides.
polyacanthus, Lam. Fl. Fr. ii. 90 = Cnicus Casabonæ.
polyanthemos, *Hegel. ex Herr, Fl. Schw.* 801.—Helvet.
polyanthemus, Schrad. Enum. Pl. Transs. 417 = collinus.
polyanthemus, Boeber, in Pall. N. Nord. Beitr. vi. 261 ; Bieb. Casp. App. 210 (= *Cirsium elodes*).—Reg. Caucas.
polyanthemus, Linn. Mant. i. 109 = Cnicus polyanthemus.
polyanthemus x, Schleich. Spicil. 15 = nutans.
polyanthos, Boiss. Fl. Orient. iii. 648, 988 = Cnicus pungens.
polyanthos, Schreb. Spicil. 15 = acanthoides.
polycinos, Willd. Sp. Pl. iii. 1655 = Jurinea polycinos.
polymorphus, Lapeyr. in Mem. Acad. Toulouse, i. (1782) 217. t. 10, 20 = Cnicus heterophyllus.
Pontederæ, Steud. Nom. ed. I. 158 = Cnicus rivularis.
Poselii x, Bragg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 107.—Europ.
pratensis, Huds. Fl. Angl. ed. II. 353 ; Lam. Encyc. i. 700 = Cnicus pratensis.
pseudosyriacus, *Lojac. in Naturol. Sicil.* iv. (1885) 109.—Sicil.
ptarmicæfolius, Mill. Gard. Dict. ed. VIII. n. 1 = Cnicus stellatus.
ptarmicanthus, Dur. in Duch. Rev. Bot. i. (1845–46) 363 = Reuterianus.
ptarmicoides, Steph. ex DC. Prod. vi. 653 = Cousinia arachnoidea.
pulcher, *Clark, Trav.* i. 789, nomen.—Tauris.
pumilus, *D. Don, Prod. Fl. Nep.* 167.—Nepal.
pumilus, Hook. f. Bot. Am. i. 302 = Cnicus Drummondii.
pumilus, Nutt. Gen. Am. ii. 130 = Cnicus pumilus.
pumilus, Vill. Hist. Pl. Dauph. ii. 17 = Cnicus heterophyllus.
punctatus, Herb. Madr. ex Wall. Cat. n. 2907 = Trichelopis radiosa.
pungens, Pers. ex DC. Prod. vi. 643 = Cnicus pungens.
purpureus, *Vill. Fl. Delphin.* 90.—Gallia.
pycnocephalus, *Linn. Sp. Pl.* ed. II. 1151 ; *Jacq. Hort. Vind.* i. t. 17. t. 44.—Europ. ; Reg. Mediterr. ; Orient.
pycnocephalus, Spreng. Syst. iii. 385 = cephalanthus.
pycnocephalus-alatus, Stev. ex DC. Prod. vi. 653 = Cousinia arachnoidea.
pygmæus, Jacq. Enum. Vindob. 147, 282 = Saussurea pygmæa.
pyrenaicus, Gouan, Illustr. 68 = Cnicus pyrenaicus.
pyrenaicus, *Yacq. Obs. Bot.* iv. 11. t. 94.—Mont. Pyren.
pyrochros, Less. in Linnæa, v. (1830) 180 = Cnicus conspicuus.
quinquefolius, D. Don, Prod. Fl. Nep. 167 = Serratula quinquefolia.
radiatus, Waldst. & Kit. Pl. Rar. Hung. i. 9 = Serratula radiata.
radicans, Roxb. Hort. Beng. 60 ; Fl. Ind. iii. 408 = Trichelopis radicans.
ramosus, Roxb. ll. cc. 101, 407 = Volutarella divaricata.
ramosissimus, *Pant. Bianch. Cruc-Gors*, 51.—Monteneg.
recurvatus, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 869, in obs. = nigrescens.
recurvatus, Wallr. in Linnæa, xxx. (1859–60) 113 = granatensis
remotifolius, Hook. Fl. Bor. Am. i. 303 = Cnicus remotifolius.
repandus, Pers Syn. ii. 386 = Cnicus repandus.
Reuterianus, *Boiss. Diagn.* Ser. II. iii. 44.—Hispan.
rhaticus, *Dalla Torre, Alt. Alpenpfl.* 136.—Europ.
rhegninus, Steud. Nom. ed. I. 158 = Cnicus strictus.
rigens, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 144 = Cnicus rigens.
rigens x, *Godr. Fl. Lorr.* ii. 40.—Gallia.
rivularis, Jacq. Fl. Austr. i. 57 = Cnicus rivularis.
Rœmii, Vill. Fl. Delphin. 91 = Cnicus acaulis.
rudecalis, Tausch, ex Nym. Consp. 413 = acanthoides.
rufescens, Pers. Syn. ii. 389 = Cnicus rufescens.
salinus, Bieb. ex Steud. Nom. ed. II. i. 253 = Centaurea serratuloides.
salisburgensis, Pers. Syn. ii. 386 = Cnicus rivularis.
samniticus, Tenore, Sem. Hort. Neap. (1826) 12 = Cnicus italicus.
Sancto-Balsano, *Lisiel. Nouv. Not.* 84.—Candollei.
sardous, DC. Prod. vi. 636 = pycnocephalus.
scaber, Polr. Voy. Barb. ii. 241 = Cnicus giganteus.

CARDUUS:—
Schimperi, *Sch. Bip. in Linnæa*, xix. (1847) 884.—Abyssin.
Schmidianus x, Ruhm. in Eichl. Jahrb. i. (1881) 240.—Europ.
Scolymus, Balbi, Hist. Pl. vii. 7, 289 = Cynara Cardunculus.
Scordium, Stokes, Bot. Mat. Med. iv. 142 = Serratula Scordium.
segetum, Frauch. Pl. Davidian. i. 178 = Cnicus segetum
semiandus, *Bieb. Fl. Taur. Cauc.* ii. 271.—Reg. Caucas. et Casp.
semipectinatus, Lam. Encyc. i. 706 = Cnicus semipectinatus.
semipinnatus, Desf. ex Steud. Nom. ed. I. 158 = Cnicus pratensis.
serratuloides, Jacq. Fl. Austr. 197 = Cnicus pannonicus.
serratuloides, Linn. Sp. Pl. 825 = Cnicus serratuloides.
serratuloides, Neck. Delic. Gallo-Belg. 338 = Cnicus arvensis.
serrulatus, Steud. Nom. ed. I. 154 ; ed. II. i. 285 = Cnicus serrulatus.
setosus, Bab. in Gard. Chron. (1844) 718 = Cnicus arvensis !
simplicifolius, *Sang. in Atti Acc. Pont. Lincei*, Ser. I xviii. (1865) t. 2.—Italia.
sinuatus, Gilib. Fl. Lituan. i. 187 = acanthoides.
Spachianus, *Dur. in Duch. Rev. Bot.* i. (1845–46) 361 —Afr. bor.
spinigerus, Jord. Obs. Pl. Crit. iii. (1846) 215 = hamulosus.
spinosissimus, Gueldenst. ex Ledeb. Fl. Ross. ii. 743 = Cnicus acaulis.
spinosissimus, C. A. Mey. in Beitr. Pfl. Russ. Reich. iv. 48 = Cnicus esculentus.
spinosissimus, Steud. Nom. ed. II. i. 385 = Cnicus echinocephalus.
spinosissimus, Vill. Hist. Pl. Dauph. iii. 11 = Cnicus spinosissimus.
spinosissimus, Walt. Fl. Carol. 194 = Cnicus horridulus.
spinulosus, Bertol. Amoen. Ital. 41 = deflexatus.
spinosus, Gueldenst. ex Ledeb. Fl. Ross. ii. 743 = Cnicus acaulis.
spurius, Linn. Hort. Upa. 240 = Cnicus eriophorus.
squarrosus, *DC. ex Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimpr. 16.—Madera.
Stangii, Buek, ex Nym. Consp. 411 = nutans.
stellatus, Linn. Sp. Pl. 823 = Cnicus stellatus.
stenolepis, *Benth. in Benth. & Hook. f. Gen.* ii. 468.—As. centr.
strictus, Steud. Nom. ed. I. 154 = Cnicus strictus.
strigosus, Bieb. Tabl. Casp. 194 = Cnicus strigosus.
suaveolens, *Benth. in Benth. & Hook. f. Gen.* ii. 468. —As. centr.
subcoriaceus, Less. in Linnæa, v. (1830) 180 = Cnicus subcoriaceus.
subdecurrens, *Bertol. Fl. Ital.* viii. 238.—Liguria.
sumanus, Pollini, in Brugnat. Giorn Fis. ix. (1816) 96 = deflexatus.
syncephalus, Reichb. f. Ic. Fl. Germ. xv. 87. t. 864 = pycnocephalus.
syriacus, Linn. Sp. Pl. 823 = Notobasis syriaca.
tataricus, Lam. Encyc. i. 708 = Cnicus rigens.
tataricus, Linn. Sp. Pl. 825 = Cnicus oleraceus !
tauricus, Boeber, ex Georgi, Beschr. Russ. Reich. iii. iv. 1294 = Cnicus echinocephalus.
taygeteus, Boiss. & Heldr. Diagn. Ser. II. iii. 42 = nutans.
tectus, Wall. Cat. n. 2906 = Saussurea obvallata.
tenuifolius, Curt. Fl. Lond. fasc. vi. t. 55 = pycnocephalus.
tenuifolius, Gand. Fl. Helv. v. 174 = deflexatus.
thessalus, Boiss. & Heldr. Diagn. Ser. II. iii. 46 = acanthoides.
Thorenori, Weinm. in Bull. Soc. Nat. Mosc. (1837) 2. VII. 69 = nutans.
Thomasii, *Hook. f. Fl. Brit. Ind.* iii. 561.—Tibet. occ.
Timbali, Martr. Fl. Tarn, 377 = nigrescens.
tinctorius, Falk, Beitr. i. 387 = Carthamus tinctorius.
tinctorius, Savi, Fl. Pis. ed. II. ii. 332 = Serratula tinctoria.
tmoleus, *Boiss. Diagn.* Ser. I. iv. 21.—As. Min.
tomentosus, Gilib. Fl. Lituan. i. 185 = Cnicus eriophorus.
transalpinus, Setter, Fl. Helv. 373 = deflexatus.
trichophalodes, Lam. Encyc. i. 708 = Cnicus stellatus.
trichocephalus, Wall. Cat. n. 2904 = Trichelopis floccosa.
trichodus, Buch.-Ham. ex [Don] Prod. vii. 269 = Cnicus involucratus.

CARDUUS:—
tuberosus, Georgi, It. i. 238 (= *Cirsium setigerum*).
tuberosus, Jacq. Fl. Austr. 330 = Cnicus canus.
tuberosus, Linn. Sp. Pl. 824 = Cnicus pratensis.
tuberosus, Vill. Hist. Pl. Dauph. iii. 15 = Cnicus tuberosus.
uliginosus, Host ex Stend. Nom. ed. I. 154 = Cnicus Willdenowii.
uliginosus, Stend. l. c. = Cnicus uliginosus.
uncinatus, *Bieb. Fl. Taur. Cauc.* iii. 258.—Reg. Caucas. ; Taurin.
undulatus, Nutt. Gen. Am. ii. 130 = Cnicus undulatus.
uniflorus, Turcz. ex Ledeb. Fl. Ross. ii. 756 = Serratula glauca.
valentinus, *Boiss. & Reut. Diagn.* Ser. II. iii. 45.—Hispan.
Vardii, *Carr. in Rev. Hortic.* (1870–71) 520.—Hispan.
vericolor, Saltsb. Prod. 185 = Silybum Marianum.
virginianus, Bosc, ex DC. Prod. vi. 651 = Cnicus Nuttallii.
virginiensis, Linn. Sp. Pl. 824 = Cnicus virginianus.
virginianus, Walt. Fl. Carol. 195 = Cnicus repandus.
viridis, *A. Kern. Sched. Fl. Exsicc.* i. 74.—Europ.
virsiensis, *Verd. Obs.* iii. 213.—Gallia.
vulgaris, Savi, Fl. Pis. ii. 241 = Cnicus italicus.
Winklerianus ×, Dalla Torre, in Lotos, xxvii. 1877 (1878) 45.—Europ.
wolgensis, Bieb. ex Willd. Enum. Hort. Berol. 889 = Cousinia wolgensis.
xanthacanthus, *Freyn, in Flora*, lxiv. (1881) 211.—Lusitan.
xeranthemoides, Sweet, Hort. Brit. ed. I. 218 = Serratula xeranthemoides.

CARDWELLIA, F. Muell. Fragm. v. 28 (1865). *PROTEACEAE*, Benth. & Hook. f. iii. 189.
sublimis, *F. Muell. l. c.* 24, 88, 78.—Austral.

CARELIA, Adans. Fam. ii. 128 (1763) = **Ageratum**, Linn. (Compos.).

CARELLIA, Less. Syn. Comp. 156 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 241.
cistifolia, *Less. l. c.*—Bras.

CAREX, [Dill.] Linn. Syst. ed. I (1735). *CYPERACEAE*, Benth. & Hook. f. iii. 1073.
CALLISTACHYS et CRYPTOGLOCHIN, Heuff. in Flora, xxvii. (1844) 538.
CARICELLA, Ehrh. Beitr. iv. 146 (1789).
DISTINAX, Rafin. ex Steud. Nom. ed. II. i. 521 (1840).
DORNERA, Heuff. ex Schur, Enum. Pl. Transs. 697 (1866).
DRYMEIA, Ehrh. Beitr. iv. 148 (1789).
GEWERBICHIA, Heuff. in Flora, xxvii. (1844) 538.
HELEONASTES, Ehrh. Beitr. iv. 147 (1789).
LEPTOSTACHYS, Ehrh. l. c. (1789).
LEUCOGLOCHIN, Heuff. in Flora, xxvii. (1844) 538.
LIMONASTES, Ehrh. Beitr. iv. 148 (1789).
MAUKSCHIA, Heuff. in Flora, xxvii. (1844) 537.
MONDO, Adans. Fam. ii. 496 (1763).
NELLREICHIA, Kotula, in Sprav. Kom. Fis. Krakow, xvii. (1883) 186.
PHYLLOSTACHYS, Torr. ex Steud. Nom. ed. II. i. 997, in syn. (1840) ; ii. 328.
PRYNGLOCHIN, Neck. Elem. iii. 345 (1790).
POLYGLOCHIN, Ehrh. Beitr. iv. 146 (1789).
PSEUDOCAREX, Miq. in Ann. Mus. Lugd. Bat. ii. 146 (1866).
PSYLLOPHORA, Ehrh. Beitr. iv. 146 (1789).
PYACOSTA, Ehrh. l. c. 147 (1789).
SCHIZAMMERIA, Moench, Meth. Suppl. 119 (1802).
SCURIA, Rafin. in Journ. Phys. lxxxix. (1819) 106.
TRAUSI, S. F. Gray, Nat. Arr. Brit. Pl. ii. 53 (1821).
TRIODUS, Rafin. in Journ. Phys. lxxxix. (1819) 106.
TRIPLISMA, Rafin. in Ann. Monthly Mag. (1819) 195.
ULVA, Rafin. Enum. Stirp. Helv. i. 249 (1742).
UVA, Stend. Nom. ed. II. i. 856 (1840).
VIGNAMTHEA, Schur, Enum. Pl. Transs. 706 (1866).
VIGNEA, Bouvy. in Lestib. Rami Fam. Cyp. 22 (1812).
alcoviensis, Pauc. ex Boott, in Ann. Nat. Hist. xvii. (1846) 87 = Torreyi.
alpicola, *Steud. Syn. Pl. Cyp.* 260.—Missouri.
amalla, *Trin. in Mém. Sav. Etran. Pan.* iv. (1836) 299. —Ins. Falkl.
CAREX:—
aciculata, *Boott, in Hook. Fl. N. Zel.* 280. t. 63.—Austral. ; N. Zel.
acicularis, *Turcz. ex Steud. Nom.* ed. II. i. 385.—Sibir. ; Reg. Caucas.
acrodira, *Schur, Enum. Pl. Transs.* 699.—Transylv.
acrolepis, Lecbl. in Denkschr. Bair. Bot. Ges. iii. (1841) 56 = Gmelini.
acrolepis, *Liebm. in Vidensh. Selsk. Skr.* V. ii. (1851) 271.—Costa-Rica.
acuminata, Reichb. Ic. Fl. Germ. viii. 28. t. 367 = nutans.
acuminata, Willd. Sp. Pl. iv. 300 = glauca.
acuta, All. Fl. Pedem. ii. 247 = riparia.
acuta, Curt. Fl. Lond. i. 281 = paludosa.
acuta, Darling. Florul. Cestrie. 98 = virginiana.
acuta, Dewey, ex Hook. Fl. Bor. Am. ii. 218 = angustata.
acuta, *Linn. Sp. Pl.* 976.—Reg. bor. temp. ; Austral.
acuta, Mnhl. Cat. 84 ; ed. (1813) 55, nomen = commutata.
acuta, Suter, Fl. Helv. ii. 261 = glauca.
acutaeformis, Brot. Fl. Lusit. i. 66 = echinata.
acutata, *Boott, in Proc. Linn. Soc.* i. (1846) 267.—Am. austr.
acutiformis, Ehrh. Beitr. iv. 48 = paludosa.
acutiuscula, Déségl. in Loisel. Fl. Gall. ed. I. 628 = pyrenaica.
Æderi, Desv. Fl. Anj. 74, sphalm. = Oederi.
adusta, *Boott, in Hook. Fl. Bor. Am.* ii. 215.—Am. bor.
adusta, Carey, ex Boott, Illustr. Carex, iii. 118 = foenea.
aemotrynchа, Desv. in C. Gay, Fl. Chil. vi. 224 = filiformis.
aemulans, *Liebm. & Drej. in Flora*, xxii. (1839) 142 = stricta.
aequabilis, *Boott, Illustr. Carex*, i. 66. t. 178.—Ins. S. Helenæ.
aequata, Nees, ex Boott, l. c. ii. 83 = myosurus.
aestivalis, *M. A. Curt. ex A. Gray, in Am. Journ. Sc.* xlii. (1842) 28.—Am. bor.
aethiopica, *Schkuhr, Riedgr.* i. 107. t. 2. f. 83.—Afr. bor.
aethostachya, Schkuhr, t. c. ii. 53 = strata.
affinis, Ehrh. Br. in Richards. in Frankl. Narr. 1st Journ. 750 = obtusata.
agastachys, Linn. f. Suppl. 414 = pendula.
Aitchisonii, Boeck. in Flora, lxiii. (1880) 456 = fissirostris.
alascana, *Boeck. in Engl. Bot. Jahrb.* vii. (1885) 277. —Am. bor. occ.
alata, Torr. in Ann. Lyc. N. York, iii. (1836) 396 = straminea.
alba, Bast. Ess. Fl. Maine-et-Loire, 388 = Bastardiana.
alba, Dewey, in Am. Journ. Sc. vii. (1824) 266 = eburnea.
alba, *Scop. Fl. Carn.* ed. II. ii. 218.—Europ. ; Am. bor.
albata, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 193, nomen.—Japon.
albescens, F. Nyland. Spicil. Pl. Fenn. ii. n. 90 = gracilis.
albicans, *Willd. ex Spreng. Syst.* iii. 818.—Am. bor.
albo-atra, Willd. ex Kunth, Enum. Pl. ii. 492, in obs. = fusca.
albo-brunescens, Schwein. in Ann. Lyc. N. York, i. (1824) 60 = straminea.
algida, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 104 ; xxvii. (1855) 1. 341.—Sibir. balcal.
alligata, *Boott, Illustr. Carex*, iv. 139.—Ins. Sandvic.
allomacros, *Steud. Syn. Pl. Cyp.* 159.—Chili.
alopecoides, *Tuckerm. Enum. Caric.* 18.—Am. bor.
alopecuroides, *D. Don, in Trans. Linn. Soc.* xiv. (1823) 393 ; *Prod. Fl. Nep.* 48.—Reg. Himal.
Alopecurus, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 141 = Eriophorum latifolium.
alpestris, All. Fl. Pedem. ii. 270 = Halleriana.
alpestris, Dewey, in Am. Journ. Sc. vii. (1824) 268 = varia.
alpestris, Gaud. Etrennes Fl. 179 = sempervirens.
alpestris, Lam. Encyc. iii. 289 = cæspitosa.
alpigena, Kern. ex Dalla Torre, Atl. Alpenfl. 234 = ferruginea.
alpina, Hoppe, Bot. Zeit. i. 374 = vulgaris.
alpina, Hoppe, ex Kunth, Enum. Pl. ii. 466 = ferruginea.
alpina, Schrank, Bair. Fl. i. 296 = sempervirens.
alpina, Sw. ex Wahlenb. Vet. Akad. Nya Handl. Stockh. (1803) 160 = Vahlii.
CAREX:—
alsophila, *F. Muell. Fragm.* viii. 257.—Austral.
alta, *Boott, in Proc. Linn. Soc.* i. (1849) 254 ; *et in Trans. Linn. Soc. xx.* (1846) 180.—Java.
alvesia, *Boott, Illustr. Carex*, i. 21. t. 56.—Texas.
ambligna, *Link, in Schrad. Journ.* ii. (1799) 306.—Lusitan. ; Maroc.
ambigua, Moench, Meth. 324 = acuta.
amblocarpa, Willd. Sp. Pl. iv. 307 = glauca.
amblyolepis, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 99).—Sibir.
ambusta, *Boott, Illustr. Carex*, i. 64. t. 172.—Alaska.
amgunensis, *F. Schmidt, Reisen Amurl.* 69.—Reg. Amur.
amicta, *Boott, Illustr. Carex*, iv. 131.—N. Granat.
ammophila, Willd. Sp. Pl. iv. 236 = divisa.
amoena, *Boott, Illustr. Carex*, iii. 106.—Ind. or.
amphibola, *Steud. Syn. Pl. Cyp.* 284.—Am. bor.
amphigena, C. Koch, in Linnæa, xxi. (1848) 615 = incurva.
amphora, *Franch. & Sav. Enum. Pl. Jap.* ii. 568.—Japon.
amplifolia, *Boott, in Hook. Fl. Bor. Am.* ii. 228.—Am. bor.
ampullacea, *Good. in Trans. Linn. Soc.* ii. (1794) 207. Reg. bor. temp.
ampullacea, Wulf. ex Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1808) 155 = Michellii.
anceps, *Muhl. ex Schkuhr, Riedgr.* ii. 66. t. F. ff. fg. 122.—Am. bor.
anceps, Willd. ex Kunth, Enum. Pl. ii. 457 = grisea.
Andersonii, Boott, in Hook. f. Fl. Antarct. 364 = decidua.
Andoiniana, *De Not. Sem. Hort. Genov.* (1847) ; *ex Steud. Syn. Pl. Cyp.* 238.—Europ.
androgyna, Balb. Elenchv, 97 = bicolor.
Angstræ, *Steud. Syn. Pl. Cyp.* 190.—Sibir.
angustata, Drej. in Fl. Dan. t. 2846 ; Revis. Caric. 86 = aquatilis.
angustata, Boott, in Hook. Fl. Bor. Am. ii. 218 = stricta.
angustifolia, Linn. ex Steud. Nom. ed. I. 156 = filiformis.
angustifolia, Seguier, ex Boott, Illustr. Carex, iv. 190 = Schreberi.
angustifolia, Sm. Engl. Fl. iv. 137 = vulgaris.
anisostachya, *Liebm. in Vidensh. Selsk. Skr.* V. ii. (1851) 266.—Mexic.
anomala, Boott, ex Perry, Jap. Exp. ii. 327 = gibba.
anomala, *Yanka, in Linnæa*, xxx. (1859–60) 609.—Europ. austr.
anomala, Pall. ex Kunth, Enum. Pl. ii. 479 = Halleriana.
anomala, *Steud. Syn. Pl. Cyp.* 230.—Abyssin.
anthericoides, *Presl, Rel. Haenk.* iii. 204.—Am. bor. occ.
anthoxantha, *Presl, l. c.* 208.—Alaska.
antucensis, Kunze, ex Kunth, Enum. Pl. ii. 412 = vulgaris.
aperta, Boott, in Hook. Fl Bor. Am. ii. 218 = stricta.
aphanandra, *Franch. & Sav. Enum. Pl. Jap.* ii. 564. —Japon.
aphanolepis, *Franch. & Sav. l. c.* 580.—Japon.
aphylla, *Kunth, Enum. Pl.* ii. 421.—Chili.
aporandra, Kunze, ex Boott, Illustr. Carex, ii. 92 = Shortiana.
apporus, R. Br. Prod. 242 = paniculata.
approximata, All. Fl. Pedem. ii. 267 = eriostorum.
approximata, Hoppe, ex DC. Fl. Fr. Suppl. 290 = tagopina.
approximata, Willd. ex Kunth, Enum. Pl. ii. 411 = cæspitosa.
aprica, *Turcz. ex Bess. in Flora*, xvii. (1834) 1. *Beibl.* 27.—Sibir. balcal.
aquatilis, Tenore, Fl. Nap. iv. 188 = microcarpa.
aquatilis, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 165.—Reg. bor.
Arcberi, Boott, in Hook. f. Fl. Tasm. ii. 98. t. 150 = acicularis.
arctic, Boott, Illustr. Carex, iv. 156 = canescens.
arctata, *Boott, in Hook. Fl. Bor. Am.* ii. 227.—Am. bor.
arctica, Delnb. in Fl. Dan. t. 101 ; Fries, Summ. Scand. i. 222 = stenophylla.
arctica, Dewey, in Am. Journ. Sc. xxvii. (1835) 239 = Perryana.

CAREX :—
arctophila, Nyland. ex Fries, Summ. Scand. 561 = aquatilis.
Ardoiniana, De Not. in Ann. Sc. Nat. Sér. III. ix. (1848) 294 = olbiensis.
arenaria, Dubois, ex Steud. Nom. ed. II. i. 286 = praecox.
arenaria, Lapeyr. ex Kunth, Enum. Pl. ii. 389 = paniculata.
arenaria, Leers, Fl. Herborn. 195. t. 14. f. 2 = disticha.
arenaria, *Linn. Sp. Pl.* 978.—Reg. bor. temp.
arenaria, Mohr, ex Boott, Illustr. Carex, iv. 210 = incurva.
arenicola, *F. Schmidt, Reisen Amuri.* 195.—Ins. Sachalin.
argentea, J. F. Gmel. Syst. 143 = alba.
argentea, Vill. Fl. Hist. Pl. Dauph. ii. 266 = humilis.
argyraeensis, *Turcz. ex Ledeb. Fl. Ross.* iv. 267.—Daburia.
argyrantha, Tuckerm. ex Boott, Illustr. Carex, iii. 119 = adusta.
argyroglochin, Hornem. in Fl. Dan. x. 7. t. 1710 = leporina.
argyrolepis, Maxim. in Franch. & Sav. Enum. Pl. Jap. i. 136 = albata.
arida, Schleich. ex Kunth, Enum. Pl. ii. 464 = semper-virens.
arida, Schwein. & Torr. in Ann. Lyc. N. York, i. (1824) 312. t. 24 = muskingumensis.
aristata, R. Br. in Richards. in Frankl. Narr. 1st Journ. 751 = trichocarpa.
aristata, Clairv. ex Steud. Nom. ed. II. i. 286 = Uncinia microglochin.
aristata, Dewey, in Am. Journ. Sc. vii. (1824) 277. t. A. f. 1 = Daviesi.
aristata, Honck. Syn. Pl. Germ. 861 = praecox.
aristata, Siegert, ex Wimm. Fl. Schles. ed. III. (1857) 72 = hirta.
aristata, Urv. in Mém. Soc. Linn. Par. iv. (1826) 599 = trifida.
armena, Boiss. Fl. Orient. v. 426 = fulva.
Arnottiana, *Boott, in Ann. & Mag. Nat. Hist.* xvii. (1846) 56.—Ind. or.; Malaya.
aspera, Willd. Prod. Fl. Berl. 32 = glauca.
asperifolia, *Steud. Syn. Pl. Cyp.* 195.—Ins. Borbon.
asperula, *Nees, in Wight, Contrib.* 134.—Reg. Himal.
asperula, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104; xxvii. (1855) 1. 847 = Meyeriana.
assimilis, *Steud. Syn. Pl. Cyp.* 195.—Ins. Maurit.
assiniboinensis, *Boott, in Coult. Bot. Gas.* ix. (1884) 91.—Am. bor.
astrocarpica, Willd. ex Kunth, Enum. Pl. ii. 385 = muricata.
asturica, *Boiss. Pugill. Pl. Nov.* 117.—Hispan.
atavina, *Hoppe, in Denkschr. Bot. Ges. Regensb.* (1815) 1. B. f. o, p, q.—Alp. Europ.
atavrina, Koch, Syn. Fl. Germ. ed. II. 874 = atrata.
atherodes, Frank, ex Kunth, Enum. Pl. ii. 495 = stenolepis.
atherodes, Sprang. Syst. iii. 828 = aristata.
athrostachya, *Olney, in Proc. Am. Acad.* viii. (1868) 293.—Am. bor. occ.
atrata, Banks, ex Boott, in Proc. Linn. Soc. i. (1849) 257 = Bankii.
atrata, *Linn. Sp. Pl.* 976.—Reg. bor. et arct.
atrofusca, Schkuhr, Riedgr. 106 = tenuiseta.
atrofusca, Sieber, ex Kunth, Enum. Pl. ii. 435 = fuliginosa.
atrofusca, Stev. in Mém. Soc. Nat. Mosc. iv. (1812) 67 = atrata.
atrofusca, Willd. ex Steud. Nom. ed. II. i. 286 = pichinchensis.
atrosicta, *Steud. Syn. Pl. Cyp.* 204.—Reg. Magell.
atropurpurea, *Boeck. in Linnaea*, xxxix. (1875) 150.—Am. austr.
atropurpurea, Fisch. ex Trevir. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 536 = atrata.
atrorufa, Willd. ex Kunth, Enum. Pl. ii. 393 = pichinchensis.
attenuata, R. Br. in Richards. in Frankl. Narr. 1st Journ. 750 = rupestris.
aurea, *Nutt. Gen. Am.* ii. 205.—Am. bor.
aureolensis, *Steud. Syn. Pl. Cyp.* 223.—Am. bor.
australasica, Schkuhr, Riedgr. ii. 10. t. 2 qq. f. 167 = divisa.

CAREX :—
axillaris, *Good. in Trans. Linn. Soc.* ii. (1794) 151. t. 19. f. 1.—Europ.
axillaris, Link, Enum. Hort. Berol. ii. 329 = diluta.
axillaris, Linn. Sp. Pl. ed. II. 1382 = remota.
axillaris, Schrank, ex Steud. Nom. ed. I. 155 = loliacea.
azorica, *Y. Gay, in Ann. Sc. Nat.* Sér. II. xi. (1839) 155.—Ins. Azoric.
baenzana, *Steud. Syn. Pl. Cyp.* 236.—Texas.
baccans, *Nees, in Wight, Contrib.* 132.—Ind. or.
Backiana, Dewey, in Am. Journ. Sc. xxix. (1836) 250 = obtusata.
Backii, *Boott, in Hook. Fl. Bor. Am.* ii. 210.—Am. bor.
badia, Pers. Syn. ii. 541 = canescens.
baetica, Asterw. ex Willk. in Bot. Zeit. vi. (1848) 414 = distans.
Balbisii, Schkuhr, ex Spreng. Pugill. ii. 86 = extensa, paucinata.
baldensis, *Linn. Syst. Pl.* ii. 32.—Europ.
baldensis, Vill. Hist. Pl. Dauph. ii. 196 = foetida.
Baldwiniana, Dewey, in Am. Journ. Sc. xxvi. (1834) 107. t. 1. f. 61 = Elliottii.
Baldwiniana, Steud. Nom. ed. II. i. 286 = praec.
Baltzellii, *Chapm. List of Florida Pl.* (1845).—Am. bor.
banata, Sin. in Rees, Cyclop. xxxix. = inouva.
banatica, *Hersff. in Verh. Zool.-Bot. Ges. Wien*, viii. (1858) 233.—Banat.
Banksii, *Boott, in Trans. Linn. Soc.* xx. (1851) 119; *ex Hook. f. Fl. Antarct.* 485. t. 142.—Reg. Magell.
Barbarae, Dewey, ex Torr. in Bot. Mex. Bound. 221 = Pratensis.
barbata, Boott, Illustr. Carex, i. 68 = Gunniana.
Baroni, *Baker, in Journ. Linn. Soc.* xxi. (1885) 451.—Madag.
Barrattii, Torr. ex Schwein. in Ann. Lyc. N. York, i. (1824) 361 = littoralis.
basiantha, *Steud. Syn. Pl. Cyp.* 232.—Am. bor.
basilaris, *Nord. Obs. Pl. Crit.* iii. (1846) 246.—Europ. austr. occ.; Marocco.
Bastardiana, DC. Fl. Fr. vi. 298 = pilulifera.
Beckeri, *A. Mey. ex Claus, in Beitr. Pfl. Russ. Reich*, vii. 58.—Rossia.
Beckyana, Boott, Illustr. Carex, i. 67 = Hookeri.
Bellardi, All. Fl. Pedem. ii. 264 = Elyna spicata.
Bella-Villa, *Dewey, in Am. Journ. Sc.* Ser. II. xli. (1866) 239.—Am. bor.
bengalensis, *Roxb. Hort. Beng.* 103; *Fl. Ind.* iii. 572.—Ind. or.
Benthamiana, Boott, in Royle, Illustr. Bot. Himal. (Pref.) ex Boott, Illustr. Carex, i. 8 = notha.
beringiana, *Cham. ex Steud. Syn. Pl. Cyp.* 229.—Alaska.
bermudiana, *Hemsl. in Journ. Bot.* xxi. (1883) 260.—Ins. Bermuda.
Bertevana, Desv. in C. Gay, Fl. Chil. vi. 196 = aphylla?
Bertoroana, Hohen. ex Boott, Illustr. Carex, iv. 149 = scitiofia.
Bertoniana, *Steud. in Flora*, xxv. (1842) 604.—Ins. Juan. Fernand.
Bertolonii, Schkuhr, Riedgr. ii. 8. t. D. f. 18 = divisa.
Beyrichiana, *Boeck. in Linnaea*, xli. (1877) 289.—Ecuador.
Bichenoviana, *Boott, in Hook. f. Fl. Tasm.* ii. 101.—Tasmania.
bicolor, *All. Fl. Pedem.* ii. 267.—Europ. ; Am. bor.
bicolor, Nylandr. ex Nym. Consp. 776 = salina.
bicostata, *Olney, Car. Bor. Am.* 9.—Am. bor.
bifida, *Boott, in Proc. Am. Acad.* vii. (1868) 294.—Am. bor. occ.
bifida, *Roth, ex Steud. Nom.* ed. I. 158.—Hab.?
bifurca, Moench, Meth. 326 = riparia.
bifurca, Schrank, Beitr. Fl. I. 304 = ampullacea.
Bigelowii, Torr. & Schwein. ex Boott, Illustr. Carex, iv. 167 = vulgaris.
bilgolarta, DC. Fl. Fr. v. 296 = laevigata.
binis, Schkuhr, Riedgr. i. 36 = distachya.
binata, *Poir. Encyc. Suppl.* iii. 257.—Afr. bor.
binervis, Dewey, in Am. Journ. Sc. xxx. (1836) 61 = laevigata.
binervis, *Sm. in Trans. Linn. Soc.* v. (1800) 268.—Europ.
binervis, Wahlenb. ex Kunth, Enum. Pl. ii. 461 = fulva.
binervis, Willd. ex Kunth, l. c. 447 = extensa.

CAREX :—
bipartita, All. Fl. Pedem. ii. 265. t. 89. f. 3 = Kobresia caricina.
bipartita, F. G. Dietr. Vollst. Lex. Gaertn. Nachtr. ii. 16 = bifida, *Roth*.
bispicata, Hook. & Arn. in Bot. Beech. Voy. 118. t. 28 = longerostata.
blanda, Dewey, in Am. Journ. Sc. x. (1826) 45 = laxiflora.
blepharophora, A. Gray, in Ann. Lyc. N. York, iii. (1835) 235 = flexilis.
Blyttii, F. Nyland. Spicil. Fl. Fenn. ii. 85 = tenella.
Boenninghauseniana, Kunth, Enum. Pl. ii. 404 = axillaris.
Boenninghausiana, Weihe, in Flora, ix. (1826) 759 = axillaris.
bohemica, Schreb. Beschr. Graes. ii. 53 = cyperoides.
Bolanderi, *Steud. Syn. Pl. Cyp.* 215.—Calif.
Bolanderi, *Olney, in Proc. Am. Acad.* vii. (1868) 293.—Calif.
bolina, Lang, in Linnaea, xxiv. (1851) 581 = vulgaris.
bollviensis, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 32.—Bolivia.
Bolliana, *Boeck. in Flora*, lxi. (1878) 40.—Am. bor. bonariensis, *Desf. in Poir. Encyc. Suppl.* iii. 250.—Reg. Argent.
bonariensis, Schlecht. in Linnaea, x. (1836) 116 = involucrata.
Bongardi, *Boott, in Proc. Linn. Soc.* i. (1846) 260 = Ins. Loo-choo.
Bongardiana, *C. A. Mey. in Middend. Reise (Fl. Ochot* 101).—Sibir. or.
Bonplandii, *Kunth, Enum. Pl.* ii. 380.—Am. bor. occ. et austr.
Boottiana, Benth. ex Boott, in Bost. Journ. Nat. Hist. v. (1845) 112 = picta.
Boottiana, Hook. & Arn. Bot. Beech. Voy. 373 = Bongardi.
borbonica, *Lam. Encyc.* iii. 387.—Ins. Mascar.
borealis, Lang, in Flora, xxvi. (1843) 142 = aquatilis.
borealicola, *Regel, in Act. Hort. Petrop.* vii. (1880) 566.—As. centr.
Boryana, *Schkuhr, Riedgr.* ii. 43.—Ins. Mascar.
Boscii, *Willd. ex Spreng. Syst.* iii. 812.—Am. bor
brachyclinna, *Griseb. in Goett. Abh.* xxiv. (1879) 315. —Reg. Argent.
brachycephala, Poepp. ex Boott, Illustr. Carex, i. 67 = Indecora.
brachylepis, Turcz. ex Steud. Nom. ed. II. i. 286; et ex Boott, l. c. iii. 118 = alpina.
brachyphylla, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104; xxvii. (1855) 1. 846 = ericetorum.
brachyrhyncha, *Gaudin, in Oestr. Bot. Zeitschr.* xx (1870) 199, 293.—Europ.
brachystachya, Schrank, Beitr. Fl. I. 294 = tenuis.
bracteata, Giesecke, ex Boott, Illustr. Carex, iv. 160 = salina.
bracteata, Seter, Fl. Helv. ii. 250 = mucronata.
bracteosa, *Kunze, in Schkuhr, Riedgr. Suppl.* 12.—Chili.
bracteosa, Schwein. in Ann. Lyc. N. York, i. (1824) 55 = xylophloides.
brasiliensis, St. Hil. Voy. Bras. i. 369 = procera.
brasiana, Boeck. in Flora, xxxix. (1856) 321 = bananensis.
Brenkii, Schulzi. ex Lange, Consp. Fl. Groenl. i. 136 = canescens.
brevicaulis, *Thom. ex Steud. Nom.* ed. II. i. 287.—Ins. Trist. d'Acunha.
brevicollis, *DC. Fl. Fr.* vi. 295.—Europ. austr.; As. Min.
brevicostata, *R. Br. Prod.* 242.—Reg. Himal.; Japon.; Australas.
brevicuminis, Thw. Enum. Pl. Zeyl. 356 = Thwaitesii.
brevifolia, Host, Gram. iv. 50. t. 89 = ferruginea.
brevigumis, *Le Gall, Fl. Morb.* 458.—Gallia.
brevipes, *S. Wats. in Bot. Calif.* ii. 246.—Calif.
brevirostrata, Pois. Encyc. Suppl. iii. 278 = obtusata.
brevirostris, *Blytt, in Bot. Notiser*, (1857) 133.—Europ.
brevirostris, Cedercr. in Vet. Acad. Oefv. Stockh. xiv. (1857) 199 = leucoglochin.
Breweri, *Boott, Illustr. Carex*, iv. 149. t. 455.—Am. bor. occ.
brizoides, Gaeaen. Klinsch. v. 267 = hyperbo.
brizoides, Huds. Fl. Angl. 349; ed. II. 406 = canescens.

CAREX :—
brizoides, Lam. Encyc. iii. 333 = praecox.
brizoides, *Linn. Cent. Pl.* i. 31.—Europ.
brizoides, Pall. ex Boott, Illustr. Carex, iv. 199 = Schreberi.
Brizopyrum, Kunze, in Schkuhr, Riedgr. Suppl. i. 169. t. 46 = alita.
bromoides, Dubois, ex Boott, Illustr. Carex, iv. 199 = Schreberi.
bromoides, *Schkuhr, Riedgr.* ii. 8.—Am. bor.
Bromgniartii, *Kunth, Enum. Pl.* ii. 380.—Chili.
Brownei, Steud. Syn. Pl. Cyp. 209 = longifolia.
Brownii, Tuckerm. Enum. Car. 21 = laciostoma.
brunnea, *Thunb. Fl. Jap.* 35.—Reg. bor. et austr.
brunnescens, Poir. Encyc. Suppl. iii. 286 = Gebhardi.
Buchanani, *Berggr. in Journ. Bot.* xviii. (1880) 104.—N. Zel.
Buchtyi, Wimm. Fl. Am. Journ. Sc. xlvii. (1845) 143 = canescens.
Buekii, Wimm. Fl. Schles. ed. III. (1857) 81 = stricta.
bulbosa, *Gueldenst. ex Ledeb. Fl. Ross.* iv. 320.—Ross. austr.
bulbosa, *Booch. in Flora*, xxxviii. (1855) 597.—Am. bor.
bullata, Ell. Sketch, ii. 556 = striata.
bullata, Dewey, in Am. Journ. Sc. ix. (1825) 71 = Tuckermani.
bullata, *Rudolph, ex Georgi, Beach. Russ. Reich.* iii. v. 1266.—Tauria.
bullata, *Schkuhr, Riedgr.* ii. 85.—Am. bor.
Bungeana, *Debeaux, in Act. Linn. Soc. Bord.* xxxiii. (1879) 68.—China.
Burchelliana, *Boeck. in Linnaea*, xli. (1877) 284.—Afr. austr.
Buxbaumii, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1808) 168 = fusca.
caduca, *Boott, Illustr. Carex*, iv. 157.—Am. austr.
caeria, Gmelin. Spicil. Fl. Rumel. ii. 412 = tomentosa.
caespitila, *Nees, in Wight, Contrib.* 127.—Reg. Himal.
caespitosa, Good. in Trans. Linn. Soc. ii. (1794) 195. t. 21 = vulgaris.
caespitosa, Hook. Fl. Scot. 286 = saxatilis.
caespitosa, Huds. Fl. Angl. ed. II. 412 = stricta.
caespitosa, Ledeb. Fl. Ross. iv. 310 = cicuinoides.
caespitosa, *Linn. Sp. Pl.* 978.—Reg. bor. et austr.
caespitosa, Soland. ex Boott, Illustr. Carex, i. 63 = decidua.
caespitosa, Tenore, Viaggio in Calab. 129 = frigida.
caespitosa, Thunb. Fl. Jap. 39 = Thunbergii ?
Caflischii x, Bruegg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxix. (1880) 119.—Europ.
campestris, Hort. ex Boott, Illustr. Carex, i. 161 = obesa.
campestris, Host, Gram. iv. t. 88 = supina.
Campoli, *Boiss. & Reut. Pugill. Pl. Nov.* 117.—Hispan.
Campyloxys, *Steud. Syn. Pl. Cyp.* 213.—Reg. Magell.
camtschatcensis, Kunth, Enum. Pl. ii. 477 = longerostrata.
canadensis, *Dewey, in Am. Journ. Sc.* Ser. II. xli. (1866) 229.—Am. bor.
canescens, Host, Gram. i. t. 57 = paradoxa.
canescens, Huds. Fl. Angl. ed. II. 406 = divulsa.
canescens, Leers, Fl. Herborn. 712. t. 14. f. 8 = muricata.
canescens, *Linn. Sp. Pl.* 974.—Reg. temp. bor. et austr.
canescens, Poll. Palat. Hist. Pl. ii. 571 = elongata.
capensis, *Schkuhr, Riedgr.* ii. 89.—Afr. austr.
capensis, Thunb. Prod. Pl. Cap. 14 = Schoenoxyphium Thunbergii.
capillacea, *Boott, Illustr. Carex*, i. 44. t. 110.—Reg. Himal. ; Austral.
capillaris, Leers, Fl. Herborn. t. 15. f. 2 = sylvatica.
capillaris, *Linn. Sp. Pl.* 977.—Reg. bor. temp. et arct.
capillaris, Willd. ex Kunth, Enum. Pl. ii. 464 = firma.
capillata, Schkuhr, ex Kunth, l. c. 490 = hirsuta.
capillipes, *Dref. in Fl. Dan.* t. 2844 ; *Rovis. Caric.* 50.—Ins. Island.
capitata, All. Fl. Pedem. ii. 264 = Davalliana.
capitata, *Linn. Syst. ed. X.* 1263.—Europ. ; Am. bor.
capitata, F. Muell. Fragm. viii. 251 = Cephalotes.
capitata, Ord. in Fl. Dan. t. 572 = Enockiorhiza sp.
capitata, Soland. ex Boott, Illustr. Carex, iv. 148 = diclina.
capitellata, *Boiss. & Bal. ex Boiss. Fl. Orient.* v. 299.—Ins. Min.

CAREX :—
cardiolepis, *Nees, in Wight, Contrib.* 127.—Reg. Himal.
Careyana, *Torr. ex Dewey, in Am. Journ. Sc.* Ser. I. xxx. (1836) 60.—Am. bor.
Carlantonensis, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 293 = enervis.
carlottensis, *Dewey, in Am. Journ. Sc.* Ser. I. xxviii. (1835) 338.—Am. bor.
carolinian, *Buckley, l. c.* xlv. (1843) 173.—Am. bor.
caryophyllaea, J. F. Gmel. Syst. 1413 = seq.
caryophyllata, Kunth, Enum. Pl. ii. 487 = emarginata.
caryophyllea, Latour. Chlor. 57 = praecox.
Castanea, Ell. Sketch, ii. 546 = Elliottii.
Castanea, Micliich. in Flora, xxxii. (1849) 665 = capillaris.
Castriana, Heer, ex Boott, Illustr. Carex, iv. 143 = dioica.
cataractae, R. Br. Prod. 242 = flava.
caucasica, *Stev. in Mém. Soc. Nat. Mosc.* iv. (1813) 68.—Reg. Caucas.
celsa, *Boott, Illustr. Carex*, iii. 108.—Reg. Himal.
cenisia, Balb. in Roem. Arch. iii. 187 = bicolor.
cephaloides, *Dewey, ex Boott, Illustr. Carex*, iii. 123.—Am. bor.
cephalophora, *Muhl. ex Schkuhr, Riedgr.* ii. 4.—Am. bor.
Cephalotes, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 110.—Austral.
cerina, Dum. Fl. Belg. 147 ; Boott, Illustr. Carex, iv. 170 = vulgaris.
cernua, *Host, l. c.* 171.—Reg. Himal. ; Japon.
cernua, *Phil. in Linnaea*, xxix. (1857–8) 58.—Chili.
cernua, *Raousch. Nom.* ed. III. 270.—' Helvet.'
ceylanica, *Boeck. in Linnaea*, xl. (1876) 241.—Ind. or.
Chaberti, F. Schult. in Flora, liv. (1871) 21 = divulsa.
chaetophylla, Steud. Syn. Pl. Cyp. 187 = divisa.
chaetorhiza, *Franch. & Sav. Enum. Pl. Jap.* ii. 552.—Japon.
chalaros, *Steud. Syn. Pl. Cyp.* 231.—Am. bor.
Chamissoi, *Boeck. in Linnaea*, xli. (1877) 145.—Heb. !
Chapmanni, Torr. ex Boott, Illustr. Carex, i. 16 = Baltzellii.
Chapmanni, Sartwell, ex Dewey, in Am. Journ. Sc. Ser. I. xix. (1855) 254 = tenax.
Chapmanni, *Steud. Syn. Pl. Cyp.* 222.—Florida.
Cheesemanii, *Petrie, in Trans. N. Z. Inst.* xv. (1883) 358.—N. Zel.
cherokeensis, *Schwein. & Torr. in Ann. Lyc. N. York*, i. (1824) 369. t. 25. f. 1.—Am. bor.
chilensis, Bronge. in Duperr. Voy. Bot. 156 = riparia.
chilluensis, *Phil. in Anal. Univ. Chil.* ii. (1862) 384.—Chili.
chinensis, *Retz. Obs.* iii. 42.—China.
chlorantha, *R. Br. Prod.* 242.—Austral.
chlorocarpa, Wimm. in Flora, xxxiii. (1849) 620 = vulgaris.
chlorocarpus, Liebm. in Videnak. Selsk. Skr. V. ii. (1851) 265 = Liebmanni.
chlorocystis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 520. Hongkong.
chlorolepis, *Steud. Syn. Pl. Cyp.* 204.—Chili.
chlorostachys, D. Don, in Trill. Mag. (Dec. 1823) 455 = japonica.
chlorostachys, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 68 = tempervirens.
chordalis, *Liebm. in Videnask. Selsk. Skr.* V. ii. (1851) 269.—Mexic.
chordorrhiza, *Linn. f. Suppl.* 414.—Europ. ; Am. bor.
Christyana, Boott, in Bot. Journ. Nat. Hist. v. (1847) 114 = cherokeensis.
chrysolepis, *Franch. & Sav. Enum. Pl. Jap.* ii. 571.—Japon.
ciliata, Wild. in Mém. Acad. Berol. 1794 (1797) 47. t. 2. f. 2 = eriostoma.
ciliolea, *Boiss. Diagn.* Ser. II. iv. 134.—Cilicia.
cinerea, Poll. Hist. Pl. Palat. ii. 573 = canescens.
cinnamomea, Boott, in Proc. Linn. Soc. i. (1846) 257 ; Trans. Linn. Soc. xx. (1851) 136 = nivalis.
cinnamomea, *Chamm. in Trans. N. Z. Inst.* xiv. (1882) 301.—N. Zel.
cinnamomea, *Olney, in Proc. Am. Acad.* vii. (1868) 206.—Calif.
circhnata, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 209. t. 6.—Unalaschka.
circhnata, *Berggr. in Minnesohr. Fisiog. Sallsk. Lund* (1877) n. viii. 20.—N. Zel.

CAREX :—
cirrhulosa, *Nees, in Hook. Kew Journ.* vi. (1854) 20.—Ins. Philipp.
cladostachya, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1808) 149 = polystachya.
clandestina, Good. in Trans. Linn. Soc. ii. (1794) 167 = humilis.
clavaeformis, *Hoppe, in Sturm, Deutschl. Fl. Heft* 61 (1833).—Carinthia.
clavata, *Thunb. Prod. Pl. Cap.* 14.—Afr. austr.
coacta, Boott, in Proc. Linn. Soc. i. (1846) 285 ; et Trans. Linn. Soc. xx. (1846) 133 = curaica.
coarctata, *Kit. in Linnaea*, xxxii. (1863) 319.—Hungar.
cognata, *Kunth, Enum. Pl.* ii. 502.—Afr. austr.
colchica, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 303 = arenaria.
Colensoi, *Boott, in Hook. f. Fl. N. Zeal.* 281. t. 83.—N. Zel.
collata, Boott, in Hook. Lond. Journ. Bot. iii. (1844) 417 = virgata.
collecta, *Dewey, in Am. Journ. Sc.* xi. (1826) 314. tab. N. fig. 44.—Am. bor.
collina, Willd. Sp. Pl. iv. 260 = montana.
Colinsii, Nutt. Gen. Am. ii. 205 = subulata.
columbiana, Dewey ex Torr. in Ann. Lyc. N. York, iii. (1836) 406 = Mertensii.
comana, *Bergg. in Minnesohr. Fisiog. Sallsk. Lund* (1877) n. viii. p. 28.—N. Zel.
Commersoniana, Sieber, ex Kunth, Enum. Pl. ii. 391 = brunnea.
commista, *Steud. Syn. Pl. Cyp.* 207.—Java.
commutata, *J. Gay, in Ann. Sc. Nat.* Sér. II. xi. (1839) 198.—Am. bor.
commutata, *Torr. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 1. 205.—Soongaria.
comosa, Boott, in Trans. Linn. Soc. xx. (1846) 117 = C. Pseudo-cyperus.
compacta, *R. Br. in App. Ross, Voy.* 163.—Am. arct.
compacta, Krock. ex Hoppe, Caric. 37 = vulgaris.
compacta, Lam. Fl. Fr. ii. 172 = vulpina.
compacta, Pur. Encyc. Suppl. iii. 282 = Uncinia compacta.
compacta, *Wimm. in Flora*, xxxiii. (1850) 618.—Europ.
complanata, Torr. in Ann. Lyc. N. York, iii. (1836) 408 = triceps.
composita, *Boott, Illustr. Carex*, i. 3.—Reg. Himal.
compressa, Gand. Itinenes Fl. 111 = stricta.
compressa, *Host, in Usteri, Ann. Bot.* xxi. (1797) 33.—Europ.
compressa, Kit. ex Willd. Sp. Pl. iv. 276 = brachystachys.
compressa, Vahl, ex Kunth, Enum. Pl. ii. 378 = bonariensis.
condinsa, *R. Br. in Richards. Frankl. Narr.* 1st *Journ.* 751.—Am. arct.
concolor, *R. Br. in Parry, Voy. App.* 283.—Am. arct.
concolor, Nees, in Wight, Contrib. 125 = speciosa.
condensata, *Nees, in Wight, Contrib.* 133.—Reg. Himal.
confecta, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 512.—Abyss.
confertiflora, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 418.—Japon.
congesta, C. A. Mey. ex Boott, Illustr. Carex, i. 17 = Hoodii.
conglobata, All. Fl. Pedem. ii. 268 = montana.
conglobata, Kit. in Willd. Sp. Pl. iv. 231 = obesa.
conica, *Boott, ex Parry, Proc. Exped.* ii. 295.—Japon.
conjugata, Willd. ex Kunth, Enum. Pl. ii. 391 = Commersoniana.
conjuncta, *Boott, Illustr. Carex*, iii. 122.—Am. bor.
connidea, Mühl. Descr. Gram. Uber. 248 = laxiflora.
conoidea, *Schkuhr, Riedgr.* ii. 67.—Am. bor.
consanguinea, Kunth, Enum. Pl. ii. 374 = divisa.
consocialis, *Steud. Syn. Pl. Cyp.* 232.—Japon.
constricta, Hoppe, in Sturm, Deutschl. Fl. Heft 61 (1833) = muricata.
contracta, *F. Muell. Fragm.* viii. 258.—Austral.
convexa, *Kit. in Linnaea*, xxxii. (1863) 317.—Hungaria.
Cooleyi, Dewey, in Ann. Journ. Sc. Ser. I. xlviii. (1845) 144 = hyperlenica.
coriophora, Fisch. & Mey. ex Kunth, Enum. Pl. ii. 463 = ustulata.
corrahensis, *Phil. in Linnaea*, xxxiii. (1864–65) 273.—Chili.

CAREX :—
corsica, Degl. ex Loisel. Fl. Gall. ii. 307 = microcarpa.
corvicensa, Link, Enum. Hort. Berol. i. 368 = punctata.
Cortesii, *Liebm. in Videnskh. Selsh. Shr.* V. ii. (1851) 285.—Mexic.
corynophora, Peterm. in Flora, xxvii. (1844) 385 = acuta.
costata, *Wahl. & C. Presl, Fl. Čech.* 101.—Bohemia.
costata, Schwein. in Ann. Lyc. N. York, i. (1824) 67 = virescens.
costata, Turcz. ex Ledeb. Fl. Ross. iv. 506 = obesa.
Coulteri, *Boott, ex Hemsl. Biol. Centr. Am. Bot.* iii. 473.—Mexic.
courtallensis, *Nees, ex Boott, Illustr. Carex*, i. 52.—Ind. or.
crassa, Ehrh. Beitr. iv. 43 = riparia.
crassipes, *Borch. in Linnæa*, xi. (1876) 339.—Ind. or.
Crawei, *Dewey, ex Torr. Fl. N. York*, ii. 408.—Am. bor.
crinalis, *Boott, in Proc. Linn. Soc.* i. (1846) 256.—N. Granat.
crinigera, *Boott, Illustr. Carex*, ii. 102.—Ins. Borbon.
crinita, *Lam. Encyc.* iii. 293.—Am. bor.
cristata, [Clairv.] Man. Herb. 291 = ferruginea.
cristata, Schwein. in Ann. Lyc. N. York, i. (1824) 66 = tribuloides.
cruciata, *Nees, ex Wight, Contrib.* 123.—Ind. or.
cruciata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 149 = ramosa.
cruenta, *Nees, in Wight, Contrib.* 128.—Reg. Himal.
Cruse-corvi, *Skuttl. ex Kunze, in Schkuhr, Riedgr. Suppl.* 128. t. 20.—Am. bor.
cryptandra, Franch. & Sav. Enum. Pl. Jap. ii. 148 = macroglossa.
cryptocarpa, *Chesterm. in Trans. N. Z. Inst.* xvi. (1884) 412.—N. Zel.
cryptocarpa, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 326 = halophila.
cryptostachys, *Brongn. in Duperr. Voy. Bot.* 153.—N. Guinea.
Curtisii, Nym. Consp. 778 = Cæstzii.
Cumingiana, *Steud. Syn. Pl. Cyp.* 206.—Ins. Philipp.
Cunninghamii, Boott, Illustr. Carex, iv. 171. t. 379 = acuta.
curvicia, Franch. & Sav. Enum. Pl. Jap. ii. 252 = chætorhiza.
curvicia, Kunth, Enum. Pl. ii. 375 = vulpinaris.
curta, Good. in Trans. Linn. Soc. iii. (1794) 145 = canescna.
curvata, *Boott, Illustr. Carex*, i. 2.—Reg. Himal.
curvata, Knaf, in Flora, xxx. (1847) 184 = Schreberi.
curviculis, *Franch. & Sav. Enum. Pl. Jap.* ii. 279.—Japon.
curvirostra, Hartm. Skand. Fl. ed. II. 321 = vaginata.
curvirostris, Kunze, in Schkuhr, Riedgr. Suppl. 79. t. 20 = buccana.
curvula, *All. Fl. Pedem.* ii. 264. t. 92. f. 3.—Europ.
curvula, Lam. Encyc. iii. 280 = Schreberi.
curvula, Sav. in Mém. Soc. Nat. Mosc. vi. (1838) 67 = Elyna schœnoides.
curvula, Willd. ex Kunth, Enum. Pl. ii. 269 = Davalliana.
cuspidata, Bertol. in Mem. Soc. Med. Ensal. Gen. ii. (1809) 145 = divisa.
cuspidata, Degl. in Loisel. Fl Gall. 294 = echinata.
cuspidata, Host, Gram. i. 71. t. 97 = glauca.
cuspidata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 164 = salina.
Custoriana, Heer, ex Hegetschw. Fl. Schw. 906 = dioica.
cylindrica, Michlch. ex Kunth, Enum. Pl. ii. 484 = strata.
cylindrica, Schwein. in Ann. Lyc. N. York, i. (1824) 71 = bullata.
cylindrica, Tuckerm. ex Boott, in Hook. Lond. Journ. Bot. v. (1846) 74 = Tuckermani.
cyperoides, Dewey, in Am. Journ. Sc. Ser. II. iii. (1847) 171 = cychocephala.
cyperoides, *Murr. in Linn. Syst.* ed. XIII. 708.—Europ. ; As. temp.
Castzii, *Semler, in Oestr. Bot. Zeitschr.* viii. (1858) 98.—Europ.
dacica, Heuff. in Flora, xviii. (1835) 247 = cæspitosa.
Daltonii, *Boott, Illustr. Carex*, i. 5.—Ind. or.

CAREX :—
Darlingtonii, Schwein. ex M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 410 = scativalis.
Darwinii, *Boott, in Proc. Linn. Soc.* i. (1845) 261.—Reg. Magell. ;
dasycarpa, *Muhl. Descr. Gram. Uber.* 286.—Am. bor.
Davalliana, Dewey, ex Steud. Nom. ed. II. i. 388 = pyrenaica.
Davalliana, *Sm. in Trans. Linn. Soc.* v. (1800) 266.—Reg. bor. temp.
Davalliana, Willd. ex Kunth, Enum. Pl. ii. 368 = dioica.
Davidi, *Franch. Pl. Davidian.* i. (1884) 319.—China.
Dawsii, Dewey, in Am. Journ. Sc. x. (1826) 279 = varia.
Davisi, *Schwein. & Torr. in Ann. Lyc. N. York*, i. (1824) 236.—Am. bor.
debilis, Boiss. & Bal. ex Boiss. Fl. Orient. v. 430 = transylvanica.
debilis, *Forst. f. Prod.* 92 = Forsteri.
debilis, *Michx. Fl. Bor. Am.* ii. 172.—Am. bor.
debilis, Soland. ex Boott, Illustr. Carex, i. 65 = Solandri.
decidua, *Boott, in Proc. Linn. Soc.* i. (1845) 255.—Calif. ; Reg. Magell.
decipiens, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 308 = macrostyla.
decipiens, Turcz. in Bull. Soc. Nat. Mosc. (1838) 109 ; xxviii. (1855) i. 338 = obtusata.
declinata, *Boott, Illustr. Carex*, iv. 171. t. 380.—Austral.
decolorans, Wimm. in Flora, xxxiii. (1850) 622 = vulgaris.
decumbens, *Muhl. Descr. Gram. Uber.* 284.—Am. bor.
decora, *Boott, Illustr. Carex*, i. 5. t. 15.—Reg. Himal.
decurvata, Ehrh. Beitr. vi. 83 = pilulifera.
defexa, *Hornem. in Fl. Dan.* t. 3051.—Am. bor. ; Groenl.
Dimbollinna, J. Gay, in Ann. Sc. Nat. Sér. II. xi. (1839) 188 = stenophylla.
Dematrsnes, *Lagger, in Flora*, xxxviii. (1855) 207.—Europ.
dentata, *Hornem. in Fl. Den.* t. 1845.—Europ.
densiflora, *Presl, Rel. Haenk.* iii. 204.—Ins. Marian.
denudata, Lapeyr. ex Steud. Nom. ed. I. 160 = pyrenaica.
depauperata, *Good. in Trans. Linn. Soc.* ii. (1794) 181.—Europ. ; As. bor.
depauperata, Hornem. Plantelaere, ii. 829 = vaginata.
deprexa, Link, in Schrad. Journ. ii. (1799) 309 = Halleriana.
deprexa, Willd. ex Kunth, Enum. Pl. ii. 479 = alpestris.
deprexa, *Boott, Illustr. Carex*, ii. 83.—Reg. Himal.
Despreauxii, *Steud. Syn. Pl. Cyp.* 227.— "Terra nova."
derbi, *Chesterm. in Trans. N. Z. Inst.* xv. (1883) 301.—N. Zel.
Deweyana, *Schwein. in Ann. Lyc. N. York*, i. (1824) 65, 316.—Am. bor.
Dianæ, *Steud. Syn. Pl. Cyp.* 288.—Ins. S. Helena.
diandra, Roth, Tent. Fl. Germ. ii. 117 = paniculata.
dianthera, Schrank, in Act. Acad. Mogunt. (1788) 49 (err. typ. 57) = tæniuscula.
diaphana, *Boott, Illustr. Carex*, iv. 156.—Austral.
Dicksonii, *Franch. & Sav. Enum. Pl. Jap.* ii. 581.—Japon.
dicliua, *Phil. in Linnæa*, xxxiii. (1864-65) 271.—Chili.
Dietrichiæ, Bocck. in Flora, lvii. (1875) 122 = famlis ?
digitalis, Schwein. & Torr. in Ann. Lyc. N. York, i. (1824) 224. t. 27 = gracillima.
digitalis, *Willd. Sp. Pl.* iv. 298.—Am. bor.
digitata, *Linn. Sp. Pl.* 975.—Europ. ; As. temp.
dilatata, Hohen. ex Boiss. Fl. Orient. v. 493, sphalm. = sq.
dilata, *Sieb. Fl. Tour. Cauc.* ii. 586.—Europ. sustr. ; As. temp.
dimorpha, Brot. Fl. Lusit. i. 84 = Linkii.
dimorphalepis, Steud. Syn. Pl. Cyp. 214 = pruinosa.
dimorphostoca, *Steckligl. in Bull. Soc. Nat. Mosc.* (1854) 206.—Sibir. alpina.
diœca, Leyss, ex Boott, Illustr. Carex, iv. 140 = chordorhiza.

CAREX :—
diœica, Host, ex Steud. Nom. ed. II. i. 388 = Davalliana.
diœica, Lam. ex Steud. l. c. = Elyna spicata.
diœica, *Linn. Sp. Pl.* 973.—Reg. bor.
diœicostigma, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = Davalliana.
diœicostrongyla, St. Lag. l. c. = diœica.
dipsacea, *Bergg. in Minnesbr. Fisiogr. Sallsk. Lund* (1877) n. viii. p. 26.—N. Zel.
discoidea, Boott, in Mem. Am. Acad. N. S. vi. (1858-59) 419 = brevicuinis.
discolor, *Bosch. in Engl. Bot. Jahrb.* v. (1884) 519.—Japon.
discolor, F. Nyland. Spicil. Pl. Fenn. iii. n. 19 = salina.
discolor, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 143, partim = virgata.
dispalnaha, *Boott, in Perry, Fap. Exp.* ii. 295.—Japon.
dispersa, Dewey, in Am. Journ. Sc. viii. (1820) 266. t. A. f. 8 = tenella.
dispersa, Herb. Un. It. ex Kunth, Enum. Pl. ii. 480 = Steudeli.
dissitiflis, *Y. F. Gmel. Syst.* 144.—Helvet.
dissita, *Soland. ex Hook. f. Fl. N. Zel.* 284.—N. Zel.
distachya, Desf. Fl. Atlant. ii. 336 = Linkii.
distachya, Willd. Sp. Pl. iv. 230 = strata.
distans, Bert. ex Steud. Nom. ed. II. i. 388 = ludecora.
distans, Host, Gram. i. 67. t. 77 = fulva.
distans, Lightf. Fl. Scot. ii. 561 = binervis.
distans, *Linn. Syst.* ed. X. 1268.—Europ. ; As. occ. ; Am. bor.
distans, Willd. ex Kunth, Enum. Pl. ii. 488 = sutans.
distenta, Kunze, ex Kunth, l. c. 449 = indecora.
disticha, A. Gray, Man. N. U. St. ed. V. 674 = Sartwelli.
distichna, *Huds. Fl. Angl.* ed. I. 347.—Europ.
divergens, Thuill. Fl. Par. ii. ed. 481 = elongata.
diversicolor, *Crantz, Inst.* i. 405.—Italia ?
diversiflora, Host, Gram. i. 53. t. 70 = Hallerlana.
divisa, *Huds. Fl. Angl.* ed. I. 346 ; ed. II. 405.—Reg. bor. temp.
divisa, Ood. in Fl. Dan. t. 371 = flava.
divulsa, Good. Fl. Helv. vi. 47 = muricata.
divulsa, *Good. in Trans. Linn. Soc.* ii. (1794) 160.—Reg. bor. temp.
Doenitzii, *Bosch. in Flora*, lxv. (1882) 61.—Japon.
Doniana, Spreng. Syst. iii. 825 = japonica.
Douglasii, *Boott, in Hook. Fl. Bor. Am.* ii. 218.—Am. bor. occ.
Dregeana, *Kunth, Enum. Pl.* ii. 511.—Afr. austr.
Dreyeri, Lange, in Flora, xxv. (1842) 548 = cæspitosa.
Drejeriana, *Lange, in Fl. Dan.* t. 2975.—Groenl.
Drummondiana, Dewey, in Am. Journ. Sc. xxix. (1836) 351. t. V. f. 68 = rupestris.
Drymeia, Linn. f. Suppl. 414 = sylvatica.
drymophila, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 104.—Sibir.
dubia, Gand. Eremus Fl. 178 = strata.
dubia, *Hose, in Usteri, Ann. Bot.* xxi. (1797) 84.—Europ.
dubia, Sibth. in Sibth. & Sm. Fl. Græc. Prod. i. 58 = Scslerla alba.
dubitata, Dewey, ex Boott, Illustr. Carex, iv. 167 = vulgaris.
Duftneri, Lapeyr. Hist. Abr. Fl. Pyr. Suppl. 140 = rupestris.
dura, Boott, in Proc. Linn. Soc. i. (1845) 255 = plebeiocheata.
Duriesi, *Steud. Nom.* ed. II. i. 389.—Hispan.
duriusculis, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 214. t. 8 = stenophylla.
Duvaliana, *Franch. & Sav. Enum. Pl. Jap.* ii. 588.—Japon.
ebenacuta, *Phil. in Linnæa*, xxxiii. (1864-65) 271.—Chili.
ebenacuta, *Trauto. in Act. Hort. Petrop.* v. (1877) 155.—Sibir.
eburnea, *Boott, in Hook. Fl. Bor. Am.* ii. 236.—Am. bor.
echinata, Lam. Fl. Fr. ii. 177 = fusca.
echinata, *Murr. Prod. Stirp. Gotting.* 76.—Reg. bor. temp. ; Austral.
echinata, Roth, Tent. Fl. Germ. i. 386 = stellulata.
echinochlon, *Kunze, in Schkuhr, Riedgr. Suppl.* 67. t. 18.—Abyssin.
Eckloni, *Nees, in Linnæa*, x. (1836) 203.—Afr. austr.

CAREX :—

effata, Lasstad. ex Ledeb. Fl. Ross. iv. 391 = paniceæ.
egyptiaca, Steud. Syn. Pl. Cyp. 230 = brevicularis.
Ehrenbergiana, *Boeck. in Linnæa*, xli. (1877) 175.—Mexic.
Ehrhartiana. Hoppe, ex Boott, Illustr. Carex, iv. 145 = tereticaulis.
elata. All. Fl. Pedem. ii. 272 = stricta.
elata, *Lowe, in Trans. Camb. Phil Soc.* iv. (1831) 11. —Ins. Madera.
elata, Turner, ex Boott, Illustr. Carex, i. 66 = præcluta.
elatior, *Boeck. in Bremen Abh.* vii. (1880) 41.—Madag.
elegans, Willd. Prod. Fl. Berl. 84 = limosa.
elongatomodes, Steud. Syn. Pl. Cyp. 213 = Darwinii.
eleusinoides, *Turcz. ex Kunth, Enum. Pl.* ii. 407.—Sibir. baical.
Elliottii, *Torr. in Ann. Lyc. N. York*, i. (1824) 357.—Am. bor.
elongata, Leers, Fl. Herborn. 197. t. 14. f. 7 = canescens.
elongata, *Linn. Sp. Pl.* 974.—Europ. ; Sibir.
elytroides, Fries, in Bot. Notiser, (1846) 106 = saxatilis.
emarcida, Suter, Fl. Helvet. ii. 368 = sylvatica.
emarginata, Willd. Sp. Pl. iv. 262 = montana.
eminens, *Nees, in Wight, Contrib.* 122.—Reg. Himal.
emirnensis, *Baber, in Journ. Bot.* xxi. (1883) 129.—Madag.
Emmonsii, Dewey, in Ann. Lyc. N. York, iii. (1836) 411 = varia.
enodarum, Spreng. Syst. iii. 818 = alopecuroides.
Emoryi, *Dewey, ex Torr. Bot. Mex. Bound.* 230.—Am. bor. occ.
enervis, Dulac, Fl. Hautes-Pyr. 53 = brizoides.
enervis, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 209.—Sibir. altaic.
enervisformis, Turcz. ex Ledeb. Fl. Ross. iv. 302 = saxatilis.
epigejos, Fries, Summa, 228 = salina.
erecta, DC. Fl. Fr. iii. 120 = sempervirens.
erecta, Gilib. Exercit. ii. 349 = canescens.
ericetorum, *Poll. Hist. Pl. Palat.* ii. 380.—Europ. ; As. temp.
erinacea, Cav. Ic. v. 40 = Uncinia erinacea.
erinacea, Muhl. ex Steud. Nom. ed. II. i. 388 = hystericina.
eriophora, *Fisch. ex Steud. l. c.* 289.—Dahuria.
erythrea, *Schrad. in Linnæa*, xi. (1837) 87.—Reg. Himal.
erythrorhiza, *Boeck. in Linnæa*, xxxix. (1875) 103.—Abyss.
erythrorhiza, Steud. Syn. Pl. Cyp. 194 = Koestlini.
erythrostachys, Hoppe, in Sturm, Deutschl. Fl. Heft 69 (1885) = glauca.
erythrostachys, Kral. ex Nym. Consp. 774 = serrulata.
Esenbeckiana, *Boeck. in Linnæa*, xl. (1876) 872.—Afr. austr.
Esenbeckii, *Kunth, Enum. Pl.* ii. 522.—Reg. Himal.
Essenbeckii, Boott, Illustr. Carex, i. 52 = linearis.
evoluta, *Hartm. in Vet. Akad. Handl. Stockh.* (1818) 196.—Europ.
exaltata, *Kit. in Linnæa*, xxxii. (1863) 322.—Hungaria.
exaltata, Peterm. in Flora, xxvii. (1844) 340 = riparia.
excelsa, Poepp. ex Kunth, Enum. Pl. ii. 502 = C. Pseudocyperus.
exilis, *Boott, ex Perry, Jap. Exp.* ii. 326.—Japon.
exserverus, Cham. ex Steud. Syn. Pl. Cyp. 228 = podocarpa.
exserva, *Boott, Illustr. Carex*, i. 57.—Reg. Himal.
exigua, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 514.—Zeylan.
exilis, *Dewey, in Am. Journ. Sc.* xiv. (1828) 351.—Am. bor.
extensa, *Good. in Trans. Linn. Soc.* ii. (1794) 178. t. 21. f. 7.—Reg. temp. bor. ; Am. austr.
extensa, C. A. Mey. Vers. Pfl. Casc. 21 = dilata.
Fabri, *Hance, in Journ. Linn. Soc.* xiii. (1873) 90.—China.
falcata, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104 ; xxviii. (1855) t. 8 = saxatilis.
fallax, *Steud. Syn. Pl. Cyp.* 198.—Java.
familiaris, *Steud. l. c.* 230.—Texas.
fasciculariis, Sloyan, ex Trevir. in Bull. Soc. Nat. Mosc. suppl. (1868) t. 5 = cæspitosa.

CAREX :—

fascicularis, Soland. ex Hook. f. Fl. N. Zel. i. 288 = C. Pseudocyperus.
fasciculata, Link, in Schkuhr, Riedgr. i. 119 = echinata.
fasciculata, *Willd. Sp. Pl.* iv. 248.—Lusitan.
Faurise, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 89.—Japon.
fecunda, *Steud. Syn. Pl. Cyp.* 194.—Am. trop.
Fedia, Nees, ex Wight, Contrib. 129 = Wallichiana.
feminea, Steud. ex Boott, Illustr. Carex, iv. 138 = acuta.
Fendleriana, *Boeck. in Linnæa*, xxxix. (1875) 185.—Am. bor.
ferax, Retzsch. Nom. ed. III. 270 = erecta.
ferruginea, Schkuhr, Riedgr. i. 92 = sempervirens.
ferruginea, *Scop. Fl. Carn.* ed. II. ii. 225.—Europ.
ferruginea, Stev. in Mém. Soc. Mat. Mosc. iv. (1818) 68 = sempervirens.
festiva, *Dewey, in Am. Journ. Sc.* xxix. (1836) 246.—Reg. bor. ; ins. Sandvic.
festucacea, Schkuhr, Riedgr. ii. 28 = straminea.
fibrillosa, *Franch. & Sav. Enum. Pl. Jap.* ii. 564.—Japon.
filicina, *Nees, in Wight, Contrib.* 128.—Ind. or. ; Malaya.
filiculmis, *Franch. & Sav. Enum. Pl. Jap.* ii. 563.—Japon.
filifolia, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 125 = Durieui.
filifolia, Nutt. Gen. Am. ii. 204 = Uncinia breviseta.
filiformis, Leers, Fl. Herborn. t. 16. f. 8 = præcox.
filiformis, *Linn. Sp. Pl.* 976.—Europ. ; Am. bor. et austr. ; As. bor.
filiformis, Thuill. Fl. Par. 440 = tomentosa.
filiformis, Vahl, in Fl. Dan. t. 1048 = pilulifera.
filipendula, Drej. l. c. t. 2870 = halophila.
filipes, *Franch. & Sav. Enum. Pl. Jap.* ii. 576.—Japon.
fimbriata, *Schkuhr, Riedgr.* ii. 61.—Europ.
finitima, *Boott, Illustr. Carex*, i. 44.—Reg. Himal.
firma, *Host*, Syn. 509.—Europ.
Fischeriana, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 324 = Schmidtii.
fistilis, *Boott, Illustr. Carex*, ii. 86. t. 245.—Austral. ; N. Caled.
fistulosa, *Ball, in Journ. Bot.* xiii. (1875) 206.—Marocco ; Afghanist.
flacca, Carey, ex Boott, Illustr. Carex, i. 69 = Barrattii.
flacca, Schreb. Spicil. 689 = glauca.
flaccida, Sw. ex Kunth, Enum. Pl. ii. 150 = cladostachya.
flaccidula, *Steud. Syn. Pl. Cyp.* 199.—Am. bor.
flacciformis, Hoffmgg. ex Kunth, Enum. Pl. ii. 498 = laevigata.
flaccosperma, *Dewey, in Am. Journ. Sc.* Ser. II. ii. (1846) 346.—Am. bor.
flagellifera, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 342.—N. Zel.
Flagellum, J. F. Gmel. Syst. 138 = Scleria Flagellum.
flava, Boiss. Voy. Espagne, ii. 581 = Oederi.
flava, *Linn. Sp. Pl.* 976.—Reg. bor. temp. ; Tasman.
flavescens, Host, Gram. iv. 58. t. 96 = fulva.
flavicans, J. Nyland. Spicil. N. Fenn. iii. 15 = salina.
flavocuspis, *Franch. & Sav. Enum. Pl. Jap.* ii. 574.—Japon.
flava, Berzl. in Bot. Notiser (1853) 87 = flava.
flectens, *Boott, Illustr. Carex*, iv. 171.—Japon.
flexilis, D. Don, in Trtl. Mag. (Dec. 1823) 455 = gracilis.
flexilis, Rudge, in Trans. Linn. Soc. vii. (1804) 98. t. 10 = castanea.
flexuosa, Muhl. ex Schkuhr, Riedgr. ii. 74 = debilis.
floerkeiana, Hochst. ex Sanb. Fl. Azor. 22. t. 3. f. 1 = Vulcani.
floribunda, *Boeck. in Linnæa*, xl. (1876) 885.—Reg. Himal.
floridana, *Schwein. in Ann. Lyc. N York*, i. (1824) 66. —Florida.
floridula, *Boott, Illustr. Carex*, iv. 172.—Burma.
fluminea, Steud. Syn. Pl. Cyp. 209 = rostata.
foenea, *Mühl. ex Torr. in Ann. Lyc. N. York*, iii. (1836) 313.—Am. bor.
foenea, Willd. Enum. Hort. Berol. ii. 957 = straminea.
foetida, *All. Fl. Pedem.* ii. 265.—Europ.
foliosa, All. l. c. 270 = flava, cæspitosa.
foliosa, *D. Don, Prod. Fl. Nep.* 42.—Reg. Himal.
follicostima, *F. Schmidt, Reisen Amurl.* 196.—Reg. Amur.

CAREX :—

follicolacea, *Crantz, Inst.* i. 405.—Hab. ?
folliculata, Ell. Sketch, ii. 545 = turgescens.
folliculata, *Linn. Sp. Pl.* 978.—Am. bor.
folliculata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1808) 159 = intumescens.
Fonki, *Phil. in Linnæa*, xxix. (1857-58) 83.—Chili.
Fontanesiana, DC. Fl. Fr. iii. 103 = pyrenaica.
Fontanesii, Petr. Encyc. Suppl. iii. 257 = divisa.
Forbesii, *Franch. & Sav. Enum. Pl. Jap.* ii. 557.—Japon.
formosa, *Dewey, in Am. Journ. Sc.* viii. (1824) 98. t. B. fig. 6.—Am. bor.
Forsteri, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 154 = C. Pseudo-cyperus.
fragilis, *Boott, Illustr. Carex*, i. 7.—Reg. Himal.
Frankii, Kunth, Enum. Pl. ii. 498 = stenolepis.
Frankii, *Steud. Nom.* ed. II. i. 390.—Am. bor.
Franklinii, *Boott, in Hook. Fl. Bor. Am.* ii. 217.—Am. bor.
Fraseri, *Andr. Bot. Rep.* t. 689.—Am. bor.
Fraseriana, Ker-Gawl. in Bot. Mag. t. 1891 = Fraseri.
fraterna, *Rupr. Symb. Pl. Ross.* 208.—Rossia.
Fredrichsthaliana, Steud. Syn. Pl. Cyp. 211 = cæspitosa.
frigida, *All. Fl. Pedem.* ii. 270.—Reg. bor.
frigida, Vill. Hist. Pl. Dauph. ii. 313 = ferruginea.
frigida, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 154 = fuliginosa.
frigida, Willd. ex Kunth, Enum. Pl. ii. 464. 465 = firma, fimbriata.
frisica, H. Koch, in Flora, xxix. (1846) 273 = trinervia.
fulvicoides, *Gaudich. Freyc. Voy. Bot.* 412.—Ins. Marian.
fuliginosa, Host, Gram. iv. 52. t. 92 = fimbriata.
fuliginosa, *Schkuhr, Riedgr.* i. 91.—Lappon.
fulva, Boiss. Voy. Bot. Espagne, ii. 581 = Hornschuchiana.
fulva, Gaerta. Mey. & Scherb. Fl. Wett. iii. 330 = distans.
fulva, *Good. in Trans. Linn. Soc.* ii. (1794) 177. t. 20. f. 5.—Europ. ; Am. bor.
fulva, Hornem. in Fl. Dan. t. 1768 = binervis ?
fulva, Host, Gram. iv. 58. t. 95 = varia.
fulva, Muhl. Desc. Gram. Uber. 246 = Elliotti.
fulva, Thuill. Fl. Par. 483 = tereticaula, paradoxa.
fulvescens, Dewey, in Am. Journ. Sc. xxix. (1836) 249 – chordorrhiza.
funiformis, [Clairv.] Man. Herb. 267 = chordorrhiza.
furcata, Ell. Sketch. ii. 552 = conoea.
furcata, *Lapeyr. Hist. Abr. Pl. Pyr.* 568.—Mont. Eyren.
furculata, Peterm. in Flora, xxvii. (1844) 329 = muricata.
furva, Webb. It. Hisp. 5 = lagopina.
fusca, All. Fl. Pedem. ii. 269 = cæspitosa.
fusca, Schkuhr, Riedgr. i. 64 = pulla.
fusca, Soland. ex Boott, Illustr. Carex, iv. 217 = pumila.
fuscescens, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 517. —Japon.
fuscescens, Willd. ex Kunth, Enum. Pl. ii. 378 = divisa.
fuscicalla, *Boeck. in Linnæa*, xl. (1876) 876.—Peruv.
fuscula, Urv. in Mém. Soc. Linn. Par. iv. (1826) 509 = indecora.
fusiformis, Chapm. ex Boott, Illustr. Carex, i. 37 = laxiflora.
fusiformis, *Nees, in Wight, Contrib.* 128.—Reg. Himal.
gaimardioides, *Desv. in C. Gay, Fl. Chil.* vi. 202.—Chili.
Galeottiana, C. A. Mey. in Bull. Acad. Brux. ix. (1842) ii. 248 = Jamesoni.
Gasparrinii, *Parl. Fl. Ital.* ii. 149.—Italia.
Gaudichaudiana, Kunth, Enum. Pl. ii. 417 = vulgaris.
Gaudichaudii, *Gudnich, in Flora*, xv. (1832) 241.—Helvet.
Gayana, *Desv. in C. Gay, Fl. Chil.* vi. 206.—Chili.
Gayliana, Steud. Nom. ed. II. i. 390 = saturata.
Gebhardi, Willd. Sp. Pl. iv. 240 ; Schkuhr, Riedgr. ii. 22 = elongata.
Gebhardii, Hoppe, Caric. t. 30 = canescens.
Gebleri, Prescott, ex Boott, in Proc. Linn. Soc. i. (1846) 265 = nongarica.

CAREX:—
gemella, *Hochst. ex Steud. Syn. Pl. Cyp.* 214.—Ind. or.
geminata, Schkuhr, Riedgr. i. 65 = temaria.
geniculata, Host, Gram. iv. 52. t. 94 = frigida.
geniculata, Schleich. ex Kunth, Enum. Pl. ii. 466 = ferruginea.
genuensis, DC. Cat. Hort. Monsp. 87 = glauca.
georgiana, *Dewey, ex Steud. Syn. Pl. Cyp.* 226.—Am. bor.
germana, *Boott, in Hook. f. Fl. Antarct.* 466.—Chili.
Gerneri, Suter, Fl. Helv. i. 369 = humilis.
Geyeri, *Boott, in Trans. Linn. Soc.* xx. (1846) 118.—Am. bor. occ.
gibba, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1808) 148.—Japon.
Gibsoni, Bab. in Jard. Ann. Nat. Hist. xi. (1843) 166. t. 8 = vulgaris.
gigantea, *Rudge, in Trans. Linn. Soc.* vii. (1804) 99. t. 10. f. 3.—Am. bor.
Gilliesii, *Phil. in Anal. Univ. Chil.* (1873) 556.—Chili.
Girardiana, *Steud. Syn. Pl. Cyp.* 209.—Ins. Borbon.
glabra, Boott, Illustr. Carex, ii. 93 = vesuta.
glandulosa, Ell. ex Boott, i. c. 89 = glaucescens.
glareosa, *Schkuhr, ex Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 146.—Scandinav.
glauca, Bosc, ex Boott, Illustr. Carex, ii. 89 = glaucescens.
glauca, *Scop. Fl. Carn.* ed. II. ii. 223.—Europ.; Afr. bor.; Am. bor.
glaucescens, *Ell. Sketch*, ii. 553.—Am. bor.
glaucina, *Boeck. in Linnaea*, xl. (1876) 353.—Ind. or.
glaucodes, *Tuckerm. ex Olney, in Proc. Am. Acad.* vii. (1868) 395.—Am. bor.
Glazioviana, *Boeck. in Flora*, lxv. (1882) 59.—Bras.
Glehni, *F. Schmidt, Reisen Amurl.* 194.—Reg. Amur.
globosa, *Boott, in Proc. Linn. Soc.* i. (1846) 259.—Calif.
globularis, Leyss. ex Kunth, Enum. Pl. ii. 439 = prascox.
globularis, *Linn. Sp. Pl.* 976.—Scandinav.
globularis, Oed. in Fl. Dan. t. 448 = panices.
globularis, Roth, Tent. Fl. Germ. i. 397 = obesa.
globularis, Suter, Fl. Helv. ii. 349 = ericetorum.
globularis, Vahl, in Skr. Naturh. Selsk. iii. t. (1792) 45 = rotundata.
globularis, Vill. Hist. Pl. Dauph. ii. 211 = canescens.
globularis, Willd. in Ges. Naturf. Fr. Neue Schr. i. (1794) 45. t. 2. f. 1 = montana.
glomerata, Gilib. Exercit. ii. 548 = vulpina.
glomerata, Host, Gram. i. t. 44 = stenophylla.
glomerata, Schkuhr, Riedgr. i. 79. t. 1. f. 41 = obesa.
glomerata, *Thunb. Prod. Pl. Cap.* 14.—Afr. austr.; Am. bor. et austr.
Gmelini, *Hook. & Arn. in Bot. Beech. Voy.* 118. t. 27.—Kamtschatka.
Godefroni, Willem. Phyt. iii. 1114 = laevigata.
Goodenowii, J. Gay, in Ann. Sc. Nat. Sér. II. xi. (1839) 191 = vulgaris.
Goyeni, *Patrie, in Trans. N. Z. Inst.* xiv. (1882) 363.—N. Zel.
gracilescens, Hartm. ex Anderss. Cyp. Scand. 16 = riparia.
gracilescens, *Steud. Syn. Pl. Cyp.* 226.—Am. bor.
gracilipes, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 151.—Japon.
gracilis, *R. Br. Prod.* 242.—Austral.
gracilis, Curt. Fl. Lond. fasc. iv. t. 60 = acuta.
gracilis, Ehrh. Beitr. vi. 83, nomen; Schkuhr, Riedgr. i. 48 = loliacea.
gracilis, A. Gray, in Am. Journ. Sc. Ser. II. iv. (1847) 19 = tenella.
gracilis, Honck. Syn. Pl. Germ. i. 366 = macrostata.
gracilis, Mönch, Meth. 326 = montana.
gracilis, Wimm. in Jahresb. Schles. Ges. (1846) 79; et in Flora, xxxiii. (1850) 630 = stricta.
gracilima, Hoppe, ex Nym. Consp. 775 = panicea.
gracilima, *Schwein. in Ann. Lyc. N. York*, i. (1824) 66.—Am. bor.
gracilima, Steud. & Hochst. ex Steud. Nom. ed. II. i. 390 = ferruginea.
Graeffeana, *Beech. in Flora*, lvii. (1875) 123.—Ins. Fiji.
Grahami, Boott, in Proc. Linn. Soc. i. (1846) 180; et Trans. Linn. Soc. xix. (1846) 215 = saxatilis.
granularioides, Schwein. in Ann. Lyc. N. York, i. (1824) 69 = comosa.

CAREX:—
granularis, *Muhl. ex Schkuhr, Riedgr.* ii. 67.—Am. bor.
Grannoniana, Rabenh. Fl. Lusat.; ex Steud. Syn. Fl. Cyp. 218 = tomentosa.
Grayana, Dewey, in Am. Journ. Sc. xxv. (1834) 141 = livida.
Greyi, *Carey, in A. Gray, Man. Bot. U. St.* ed. I. 568.—Am. bor.
Grevilleana, Dewey, in Am. Journ. Sc. xxx. (1836) 61 = falva.
Grenieri, *Boeck. in Flora*, xli. (1858) 649.—Am. bor.
Griffithii, *Boott, in Proc. Linn. Soc.* i. (1846) 286.—Afghan.
Grisebachi, Roem. in Schkuhr, Riedgr. ii. 76 = grisea.
grisea, *Viv. Ann. Bot.* i. 186.—Reg. Caucas.; Persia.
grisea, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 154.—Am. bor.
Grusechii, Heuff. ex Boott, Illustr. Carex, iv. 148 = pyrenaica.
Gryps, Schkuhr, Riedgr. ii. 18 = schinata.
guayquilica, Boeck. ex Koch, Syn. Fl. Germ. ed. II. 867 = muricata.
Guannana, *Boott, in Proc. Linn. Soc.* i. (1846) 258.—Austral.
Guthnickiana, J. Gay, in Ann. Sc. Nat. Ser. II. x. (1838) 289 = decipiens.
gynandra, *Schwein. in Ann. Lyc. N. York*, i. (1824) 70.—Am. bor.
Gynandra, Sieber, ex Kunth, Enum. Pl. ii. 478 = Linkii.
gynobasis, Vill. Hist. Pl. Dauph. ii. 206 = Halleriana.
gynocrates, Wormsk. in Drej. Rev. Caric. 16 = chordorrhiza.
gynodynama, *Olney, in Proc. Am. Acad.* vii. (1868) 394.—Calif.
Gynomane, Bertol. Pl. Rar. Ital. Dec. ii. 48 = Linkii.
Hanstiana, *Boeck. in Flora*, lxi. (1878) 166.—N. Zel.
haematolepis, Drej. in Fl. Dan. t. 2370 = salina.
haematostoma, *Nees, in Wight, Contrib.* 125.—Nepal.; Reg. Himal.
Haenkeana, Presl, Rel. Haenk. iii. 203 = C. Pseudocyperus.
Hailstoni, Gibs. in Phytol. i. (1841) 870 = axillaris.
hakonensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 550.—Japon.
Halenas, *Olney, Car. Bor. Am.* 6.—Am. bor.
Haleri, Dewey, in Am. Journ. Sc. Ser. II. ii. (1846) 240.—Am. bor. Cruc-corvi.
Halei, *Carey, in Chapm. Fl. S. U. St.* 543.—Am. bor.
Halleri, Vest, ex Steud. Nom. ed. II. i. 390.—Cf. Suteri.
Halleriana, *Asso, ex Henck. Syn.* i. 872.—Europ.; Reg. Casp.; Marocco.
Halliana, *L. Bailey, in Coult. Bot. Gaz.* ix. (1884) 117.—Am. bor. occ.
Hallii, *Olney, in Hayden, Rep.* (1871) 496.—Am. bor. occ.
halmaturina, Boeck. in Linnaea, xxxix. (1875) 100 = pauciflora, tertiscalis.
halophila, Heuff. in Linnaea, xxxi. (1861-62) 695 = falva.
halophila, *F. Nyland. Spicil. Fl. Fenn.* ii. n. 50.—Lappon.
Halsnyana, Dewey, in Am. Journ. Sc. xl. (1836) 315 = polymorpha.
hamata, Forst. ex Steud. Nom. ed. II. i. 390 = Forsteri.
hamata, Sw. Prod. Veg. Ind. Occ. 18 = Uncinia jamaicensis.
Hanseodiana, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) 1. 66.—China.
Harlandi, *Boott, Illustr. Carex*, ii. 87. t. 255.—Hongkong.
Hartii, *Dewey, in Am. Journ. Sc.* Ser. II. xii. (1866) 296.—Am. bor.
Hartwegii, *Boott, in Benth. Pl. Hartw.* 96.—N. Granad.
Haydeniana, *Olney, in Bot. King, Exp.* 366.—Am. bor. occ.
Haydenii, Dewey, in Am. Journ. Sc. Ser. II. xviii. (1854) 103 = aperta.
hebecarpa, Hook. & Arn. Bot. Beech. Voy. 50 = Hookeri.
hebecarpa, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 228. t. 12.—Reg. Himal.
hebetata, *Boott, Illustr. Carex*, iv. 172.—Am. austr.
Heleonastes, *Ehrh. ex Linn. f. Suppl.* 414.—Europ.
Heleonastes, Liljebl. in Vet. Akad. Nya Handl. Stockh. (1798) 73 = glareosa.

CAREX:—
Helferi, *Boeck. in Linnaea*, xl. (1876) 365.—Burma.
helodes, Link, in Schrad. Journ. ii. (1799) 309 = laevigata.
helvola, *Blytt, in Fries, Bot. Notiser* (1849), 58.—Norveg.
helvetica, Honck. Syn. Pl. Germ. i. 378 = frigida.
helvetica, Schleich. ex Kunth, Enum. Pl. ii. 448 = punctata.
Hepburnii, Boott, in Hook. Fl. Bor. Am. ii. 209. t. 207 = sardina.
heptastachys, *Boeck. in Linnaea*, xxxix. (1875) 114.—Am. trop.
hermaphrodita, J. F. Gmel. Syst. 138 = Elyna spicata.
hermaphroditа, Jacq. Coll. iv. 174 = Mariscus Jacquini.
heterolepis, *Bunge, Enum. Pl. Chin. Bor.* 69.—China.
heteronaura, *S. Wats. in Bot. Calif.* ii. 340.—Calif.
heterophylla, *Krech. ex Steud. Nom.* ed. II. i. 390.—Silesia.
heterosperma, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1808) 151 = ancepa.
heterostachya, Bunge, Enum. Pl. Chin. Bor. 69 = nutans.
heterostachya, Torr. in Am. Journ. Sc. Ser. II. ii. (1846) 248 = Crawei.
hians, *Steud. Syn. Pl. Cyp.* 236.—Hab.?
Hilairei, *Boott, Illustr. Carex*, iv. 146.—Bras.
Hildebrandtiana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 516.—Madag.
Hilgendorfiana, *Boeck. l. c.* 518.—Japon.
hirsuta, Suter, Fl. Helv. ed. II. ii. 324 = Halleri.
hirsuta, Willd. Sp. Pl. iv. 293 = tricepa.
hirta, *Linn. Sp. Pl.* 975.—Europ.; As. temp.
hirta, Oed. in Fl. Dan. t. 879 = filiformis.
hirtaeformis, Pers. Syn. ii. 547 = hirta.
hirtella, *Drej. Syn.* 21.—Afghan.; Reg. Himal.
hirtella, J. F. Gmel. Syst. 138 = Scleria hirtella.
hirtisema, *S. Wats. in Bot. Calif.* ii. 347.—Calif.
hispida, *Gaud. Agrost. Helv.* ii. 130.—Helvet.
hispida, Link, Hort. Berol. ii. 331 = laxiculmens.
hispida, Willd. ex Schkuhr, Riedgr. i. 58 = echinata.
hispidula, Gaud. Agrost. Helv. 126 = fimbriata.
Hitchcockiana, *Dewey, in Am. Journ. Sc.* x. (1826) 274.—Am. bor.
Hochstetteri, Frank, ex Steud. Nom. ed. II. i. 391 = polytrichoides.
Hochstetteriana, *Y. Gay, ex Seub. Fl. Asor.* 22. t. 2. f. 2.—Ins. Azoric.
holostoma, *Drej. in Fl. Dan.* t. 2428.—Groenl.
homalocarpa, Peterm. in Flora, xxvii. (1844) 333 = vulgaris.
homodolepis, *Franch. & Sav. Enum. Pl. Jap.* ii. 567.—Japon.
Hoodii, *Boott, in Hook. Fl. Bor. Am.* ii. 211.—Am. arct.
Hookeriana, *Dewey, in Am. Journ. Sc.* xxix. (1836) 246.—Am. arct.
Hookeri, *Kunth, Enum. Pl.* ii. 490.—Chili.
Hoppneri, Boott, in Hook. Fl. Bor. Am. ii. 219 = salina.
hordeiformis, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 152 = hordeistichos.
hordeistichos, *Vill. Prosp.* 18.—Europ.; As. temp.
Hornschuchiana, *Hoppe, in Flora*, vi. (1824) 599.—Europ.
Horsfieldii, *Boott, in Proc. Linn. Soc.* i. (1846) 257.—Java.
Hostеana, DC. Cat. Hort. Monsp. 88 = fulva.
Hostii, Hoppe, ex Kunth, Enum. Pl. ii. 372 = divisa.
Hostii, Schkuhr, Riedgr. Nachtr. 7 = stenophylla.
Houghtonii, *Torr. in Ann. Lyc. N. York*, iii. (1836) 218.—Am. bor. occ.
Hsetiana, *Boiss. Diagn.* Ser. II. iv. 123.—Reg. Caucas.
huttonsis, Steud. Syn. Pl. Cyp. 229 = indocora.
Humboldtiana, *Steud. l. c.* 208.—N. Granat.
humilis, *Leyss. Fl. Hal.* 178.—Europ.; Reg. Caucas.
humilis, Wild. ex Kunth, Enum. Pl. ii. 441 = capitata.
hyalina, Boott, in Bot. Journ. Nat. Hist. v. (1847) 112 = straminea.
hyalinolepis, *Steud. Syn. Pl. Cyp.* 230.—Am. bor.
hybrida, Balev. ex Lam. Encyc. iii. 269 = divisa.
hybrida, Schkuhr, ex Kunth, Enum. Pl. ii. 383 = Kobomia cericea.
hydrophila, Dum. Fl. Belg. 146 = subloculata.
hymalayensis, *Steud. Syn. Pl. Cyp.* 204.—Reg. Himal.
hymenina, *Nees, ex Boott, Illustr. Carex*, i. 67.—Chili.

CAREX:—
hymenocarpa, Drej. in Fl. Dan. t. 2849 = ampullacea.
hymenolepis, Nees, in Wight, Contrib. 126 = setigera.
hypandra, *F. Muell. Fragm.* viii. 259.—Austral.
hyperborea, Drej. Kevis. Carlc. 43 = saxatilis.
hyposanthes, Stend. Syn. Pl. Cyp. 193 = Brongniartii.
hypophylla, *Miq. Fl. Ind. Bat.* iii. 354.—Malaya.
hystericina, *Muhl. ex Schkuhr, Riedgr.* ii. 69.—Am. bor.
Hystrix, A. Gray, ex Boott, Illustr. Carex, i. 26 = Crus-corvi.
Idzuroei, *Franch. & Sav. Enum. Pl. Jap.* ii. 588.—Japon.
ignota, Dewey, in Am. Journ. Sc. Ser II. viii. (1849) 348 = laxiflora.
illegitima, *Cesati, in Frieder. Reise*, 271.—Dalmat. ; Graecia.
illinoensis, *Dewey, in Am. Journ. Sc.* Ser. II. iv. (1847) 245.—Am. bor.
illustranda, Stend. Syn. Pl. Cyp. 190 = bracteosa.
ilsunæ x, Ruhm. in Richl. Jahrb. i. (1881) 258.—Europ.
impuncticta, *Boott, Illustr. Carex*, iii. 107.—Java.
inaequalis, Raeusch. Nom. ed. III. 270 = cacspitosa ?
inanis, *Kunth, Enum. Pl.* ii. 522.—Reg. Himal.
incerta, Stend. Syn. Pl. Cyp. 210 = saxatilis.
incisa, *Boott, in Perry, Yap. Exp.* ii. 327.—Japon.
incisodentata, Leuhl. ex Boott, Illustr. Carex, i. 26 = festiva.
incisodentata, *Steud. Syn. Pl. Cyp.* 189.—Reg. Magellan.
incluna, Turcz. ex Boott, Illustr. Carex, iv. 177 = alba.
inconspicua, Stend. Syn. Pl. Cyp. 221 = indecora.
incrassata, Schlecht. in Linnaea, x. (1836) 119 = riparia.
incrassata, Soland. ex Boott, Illustr. Carex, iv. 138 = trifida.
incurva, *Lightf. Fl. Scot.* ii. 544. t. 24.—Reg. bor. temp. et arct.
incurva, Sibth. & Sm. Fl. Graec. Prod. i. 227 = divisa.
indecora, *Kunth, Enum. Pl.* ii. 448.—Ins. Falk.
indica, Koen. ex Willd. Sp. Pl. iv. 315 = Scleria tuscillata.
indica, *Linn. Mant.* ii. 574.—Ind. or. ; Malaya.
indica, F. Muell. ex Benth. Fl. Austral. vii. 441 = fusilis.
indica, Schkuhr, Riedgr. i. 37 = Uncinia spartea.
indica, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 149 = Wahlenbergiana.
infata, Huds. Fl. Angl. ed. I. 354 = vesicaria.
infecta, Suter, Fl. Helvet. ii. 365 = amplexicaulis.
infuscata, Nees, in Wight, Contrib. 125 = alpina.
innocua, *Wallr. Beitr. Fl. Hercyn.* 299.—Europ.
inops, Kunze, in Schkuhr, Riedgr. Suppl. 131 = tenella.
inornata, Turcz. ex Ledeb. Fl. Ross. iv. 309, nomen = erioterum.
inquina, *Boott, Illustr. Carex*, i. 5.—Reg. Himal.
insulata, *Cormich. in Trans. Linn. Soc.* xii. (1818) 508.—Ins. Trist. d'Acunha.
intermedia, Good. in Trans. Linn. Soc. ii. (1794) 154 = distichea.
intermedia, Mégev. in Bull. Soc. Bot. Fr. x. (1863) 83 = vulgaris.
intermedia, Retz. Fl. Scand. 178 = muricata.
intermedia, Suter, Fl. Helvet. ii. 261, 262 = paludosa.
intermedia, Urv. in Mém. Soc. Linn. Par. i. (1822) 378 = arenaria.
interrupta, *Bosch. in Linnaea*, xl. (1876) 432.—Am. bor.
intricata, Tineo, in Guss. Fl. Sic. Syn. ii. 574 = vulgaris.
intumescens, *Rudge, in Trans. Linn. Soc.* vii. (1804) 97. t. 8.—Am. bor.
inundata, Willd. ex Kunth, Enum. Pl. ii. 492 = glaucescens.
inversa, *R. Br. Prod.* 249.—Austral. ; N. Zeland.
involucrata, *Bosch. in Flora*, xxxvii. (1855) 594.—Texas.
Involucrata, *Boott, Illustr. Carex*, ii. 76.—Am. austr.
iridifolia, Kunth, Enum. Pl. ii. 492 = aethiopica.
irregularis, Schwein. in Ann. Lyc. N. York, i. (1824) 86 = flava.
irigua, *Sm. in Hoppe, Caric.* t. 92.—Reg. bor. temp. et arct.
ischnantha, Steud. Syn. Pl. Cyp. 204 = brunnea.
ischnostachya, *Steud. l. c.* 222.—Japon.
Jackiana, *Boott, in Proc. Linn. Soc.* i. (1846) 260.—Java.
Jamaicensis, Poir. Encyc. iii. 246 = Uncinia jamaicensis.
Jamesii, Schwein. in Ann. Lyc. N. York, i. (1824) 87 = Steudelii.

CAREX:—
Jamesii, *Torr. in Ann. Lyc. N. York*, iii. (1836) 398. —Am. bor.
Jamesoni, *Boott, in Proc. Linn. Soc.* i. (1845) 258.—Ecuador.
japonica, Fisch. ex Boott, l. c. 259 = breviculmis.
japonica, *Thunb. Fl. Jap.* 34.—Japon.
juncea, Scop. Fl. Carn. ed. II. ii. 227 = falva.
juncea, *Willd. Enum. Hort. Berol. Suppl.* 68.—Am. bor.
juncella, T. M. Fries, in Bot. Notiser, (1857) 207 = vulgaris.
juncifolia, All. Fl. Pedem. ii. 264 = incurva.
juncifolia, J. F. Gmel. Syst. 142 = mucronata.
juncifolia, Host, Syn. 804 = stenophylla.
juncoides, *S. & C. Presl, Fl. Cech.* 190.—Bohemia.
jungermannioides, Steud. Syn. Pl. Cyp. 210 = saxatilis.
kaloides, *Petrie, in Trans. N. Z. Inst.* xiii. (1881) 382.—N. Zel.
Kaultii, *Boriaeo, in Magyar Növeny. Lapok*, ix. (1885) 181.—Europ.
kattegatensis, Fries, Ind. Sem. Hort. Upsal. (1857).—Cf. Hartm. Skand. Fl. ed. XIII. 466 = saltna.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 148.—Japon.
Kelloggii, *S. Wats Bot. Calif.* ii. 240.—Calif.
Kelringtoniana, *Steud. Syn. Pl. Cyp.* 215.—Am. bor.
Kerneri, *Kohts, in Oestr. Bot. Zeitschr.* xx. (1870) 164. —Europ.
klotzschi, *Franch. & Sav. Enum. Pl. Jap.* ii. 556.—Japon.
Kirilowii, *Turcz. in Bull. Soc. Nat. Mosc.* xxviii. (1855) i. 340.—As. centr.
Kneiskernii x, Dewey, in Am. Journ. Sc. xxxii. (1846) 247.—Am. bor.
Kochsana, DC. Cat. Hort. Monsp. 8 = paludosa.
Kochiana, Schneb. Fl. Wuerttb. 501 = evoluta.
Koenlini, *Hochst. ex Steud. Syn. Pl. Cyp.* 193.—Abyssin.
Kotschyana, Boiss. & Hohen. Diagn. Ser. I. xiii. 86 = vulgaris.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* i. 551.—Japon.
Krulli, *Boeck. in Flora*, lxv. (1882) 59.—N. Zel.
Kunzei, *Olney, ex A. Gray, in Proc. Am Acad.* viii. (1872) 406.—Am. bor.
Lachenalii, Schkuhr, Riedgr i 51 = lagopina.
lachnosperma, Wall. Cat. n. 3879 = gracilis.
lacinlata, *Boott, Illustr. Carex*, iv. 175.—Calif.
lacistoma, *E. Br. Prod.* 248.—Austral.
lacustris, Balb. ex Kunth, Enum. Pl. ii. 533 = Elyna caricina.
lacustris, *Schkuhr, Riedgr.* ii. 84. t. 6/0. f. 152.—Europ.
lacustris, Willd. Sp. Pl. iv. 305 = riparia.
laeta, *Boott, Illustr. Carex*, i. 69.—Reg. Himal.
laevicaulis, Hochst. in Seub. Fl. Azor. 21 = punctata.
laevi-conica, *Dewey, in Am. Journ. Sc.* xxiv. (1857) 47. —Am. bor.
laevigata, Boiss. Voy. Bot. Espagne, ii. 631 = Camposii.
laevigata, *Sm. in Trans. Linn. Soc.* v. (1799) 272.—Europ.
laevigata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 158 = aethiopica.
laevirostris, Andersa. Cyp. Scand. 17 = rhynchophysa.
laevis, J. F. Gmel. Syst. 138 = Scleria laevis.
laevis, Hoppe, Taschenb. (1800) 243 = dioica.
laevis, *Kit. ex Willd. Sp. Pl. iv.* 292.—Europ. austr. or.
Laggeri, Wimm. in Flora, xxxvii. (1854) 161 = repens.
lagopina, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 145.—Reg. bor.
lagopodioides, Schkuhr, Riedgr. ii. 20 = trifuloides.
Lagopus, Muhl. Desc. Gram. Uber. 265 = Fraseri.
Lamarckii, [W. Wood] in Rees, Cycl. vi. n. 108 = Shea.
Lambertiana, *Boott, ex Hook. f. Fl. N. Zel.* 284.—N. Zel.
lanceata, Dewey, in Am. Journ. Sc. xxix. (1836) 249 = salina.
lanceolata, *Boott, in Perry, Yap. Exp.* ii. 326.—Japon.
lanceolata, *Pelim. in Bull. Soc. Nat. Mosc.* viii. (1835) 238.—Lappon.
Langii, *Steud. Syn. Pl. Cyp.* 227.—Norveg.
Langsdorffii, Boott, in Proc. Linn. Soc. i. (1845) 259 = leucochlora.
lanosa, *Gueldenst. It.* i. 107 ; *ex Ledeb. Fl. Ross.* iv. 320.—Ross. austr.
lanuginosa, *Michx. Fl. Bor. Am.* ii. 175.—Am. bor.

CAREX:—
lapponica, Lang, in Linnaea, xxiv. (1851) 539 = canescens.
lasiocarpa, Ehrh. in Hann. Mag. 9ten Stück (1784), 182 = filiformis.
lasiochlaena, *Kunth, Enum. Pl.* ii. 485.—Sardinia.
latifolia, *Bonss. & Bal. in Boiss. Fl. Orient.* v. 421.—Pontus.
latifolia, J. F. Gmel. Syst. 138 = Scleria latifolia.
latifolia, Moench, Meth. 338 = paniculata.
latifolia, Soland. ex Boott, Illustr. Carex, i. 52 = C. Pseudo-cyperus.
laxa, Dewey, in Am. Journ. Sc. xxvi. (1834) 376 = limosa.
laxa, Sw. ex Kunth, Enum. Pl. ii. 513 = scabrella.
laxa, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 156.—Lappon.
laxiculmis, *Schwein. in Ann. Lyc. N. York*, i. (1824) 70.—Am. bor.
laxiflora, Ell. Sketch, ii. 549 = grisea.
laxiflora, *Lam. Encyc.* iii. 276.—Am. bor.
laxula, Tineo, ex Boott, Illustr. Carex, iv. 202 = sylvatica.
laxiuscula, Boiss. & Bal. in Boiss. Fl. Orient. v. 424 = sempervirens.
Leavenworthii, Dewey, in Am. Journ. Sc. Ser. II. ii. (1846) 246 = cephalophora.
Lechleri, *Phil. in Linnaea*, xxix. (1857–58) 88.—Reg. Magellan.
Lechleri, *Steud. Syn. Pl. Cyp.* 234.—Chili.
Ledebouriana, C. A. Mey. ex Trevir. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 540 = capillaris.
Ledebourii, Boiss. & Balsa, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 222 = nutans.
Leersii, F Schult. in Flora, liii. (1870) 455, & liv. (1871) 25 = muricata.
Leersii, Willd. Prod. Fl. Berl. 28 = stellulata.
Lehmanniana, Boott, ex Walp. Ann. i. 910 = Lemanniana.
Lehmannii, *Drej. Symb.* 18. t. 2.—Ind. or.
leiocarpa, J. Gay, in Ann. Sc. Nat. Ser. II. vi. (1836) 351 = saturica.
leiocarpa, *C. A. Mey. in Mém. Sav. Etr. Petersb.* i. (1831) 208. t. 5.—Unalask.
leiorhyncha, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 217. t. 9.—Dahuria.
Lemaniana, *Boott, in Proc. Linn. Soc.* i. (1846) 256 —Am. austr.
Lemmoni, *W. Boott, in Coult. Bot. Gaz.* ix. (1884) 93. —Am. bor. occ.
lenta, D. Don, ex Spreng. Syst. iii. 811, aphalm = gracilis.
lenticularis, Dewey, in Am. Journ. Sc. vii. (1824) 263 ; ix. (1825) 263 · magellanica.
lenticularis, D. Don, in Trans. Linn. Soc. xvi. (1825) 331 = phaeota.
lenticularis, *Michx. Fl. Bor. Am.* ii. 172.—Am. bor.
leonura, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 161 = crinita, hystericina.
lepida, *Boott, Illustr. Carex*, iv. 211.—Peruv.
lepidocarpa, Tausch, in Flora, xvii. (1834) 179 = flava.
leporina, *Linn. Sp. Pl.* 975.—Reg. bor. temp.
leporina, Michx. Fl. Bor. Am. ii. 170 = scoparia.
leporina, Oed. in Fl. Dan. t. 294 = lagopina.
leporina, Pall. ex Kunth, Enum. Pl. ii. 393 = divisa.
leporina, Schkuhr, ex Ledeb. Fl. Ross. iv. 279 = C. Heleonastes
leptalea, Wahlenb. in Vet. Akad. Nya Handl. Stockh (1803) 139 = polytrichoides.
leptophylla, *Heuff. in Linnaea*, xxxi. (1861–62) 728. t. 6.—Europ.
leptostachya, Boiss. Fl. Orient. v. 418 = seq.
leptostachys, Ehrh. ex Linn. f. Suppl. 414 = strigosa.
Lerchenfeldiana, Schur, Enum. Pl. Transs. 711 = hordeistichos.
Lesoniana, *Steud. Nom.* ed. II. i. 292.—N. Zel.
leucanthe, *Arn. ex Boott, in Proc. Linn. Soc.* i. (1845) 257.—Ind. or.
leucantha, *Schur, Enum. Pl. Transs.* 714.— Europ.
leucocarpa, *Boeck. in Flora*, xxxix. (1856) 227.—Am. bor.
leucocarpa, *Phil. in Linnaea*, xxix. (1857–58) 82.—Chili.
leucochlora, Bunge, in Mém. Sav. Etr. Petersb. ii. (1835) 142 = brevicalmis.
leucoglochin, Wahlenb. in Vet Akad. Nya Handl. Stockh. (1803) 130, 149 = obtusata.
leucoglochin, Linn. f. Suppl. 418 = pauciflora.

CAREX:—

leucolepis, Turcz. ex Steud. Nom. ed. II. i. 292 = tenuiflora.

leucorhiza, Dulac, Fl. Hautes-Pyr. 452 = Davalliana.

Lhotskiana, Nees, ex Steud. Nom. ed. II. i. 293 = Fintelmannia resticides.

Liddoni, *Boott, in Hook. Fl. Bor. Am.* ii. 214.—Am. bor.

Liddoni, Carey, in A. Gray, Man. Bot. U. St. ed. I. 545 = pallida.

Liebmanni, *Wulp. Ann.* iii. 705.—Mexic.

ligata, *Boott, ex Benth. Fl. Hongk.* 402.—China.

ligerica, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 360 = arenaria.

ligerina, Gnep. ex Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 171 = praec.

lignlata, *Nees, in Wight, Contrib.* 127.—Ind. or.

limosa, Engelm. ex Steud. Syn. Pl. Cyp. 209 = abiciena.

limosa, *Linn. Sp. Pl.* 977.—Europ. ; Am. bor.

limosa, Scop. Fl. Carn. ed. II. ii. 222 = utulata.

limula, Fries, Summa Veg. 229 = vulgaris.

Lindleyana, *Nees, in Wight, Contrib.* 121.—Ind. or.

linearis, *Boott, Illustr. Carex*, i. 81.—Reg. Himal.

linearis, Clairv. ex Boott, i. c. iv. 204 = tenuis.

linecla, *Fries, in Bot. Notiser* (1844). 58.—Lappon.

Linkii, *Willd. ex Schkuhr, Riedgr.* ii. 39.—Reg. Mediterr.

Linnaeana, Host, Gram. iii. 77 = dioica.

Linnari, Dégl. in Loisel. Fl. Gall. 627 = dioica.

liparocarpus, Gaud. Etreanes Fl. 153 = obesa.

lipidocarpa, Peterm. Fl. Lips. Excurs. 58 = flava.

lithophila, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 104 ; xxviii. (1855) 1. 328.—Sibir.

lithosperma, Forst. f. Prod. 85 = Scleria margaritifera.

lithosperma, Linn. Syst. ed. XII. 706 = Scleria Flagellum.

lithosperma, Walt. Fl. Carol. 227 = Scleria reticularis.

littoralis, Krock. ex Steud. Nom. ed. II. i. 292 = stricta.

littoralis, *Petrie, in Trans. N. Z. Inst.* xv. (1883) 358. —N. Zel.

littoralis, *Schwein. in Ann. Lyc. N. York*, i. (1824) 70. —Am. bor.

littorea, Labill. Nov. Holl. Pl. ii 69. t. 219 = pumila.

livida, *Willd. Sp. Pl.* iv. 285.—Reg. bor. temp.

lobata, Bell. App. Fl. Pedem. 42 = cervula.

lobata, Lam. Encyc. iii. 279 ; Schkuhr, Riedgr. i. 28 = divisa.

lobata, Schleich. ex Steud. Nom. ed. II. i. 293 = microstyla.

lobata, Vill. Hist. Pl. Dauph. ii. 197 = foetida.

lobata, Wahlenb. ex Steud. Nom. ed. II. i. 292 = curvula.

lobata, Willd. ex Kunth, Enum. Pl. ii. 538 = Elyna caricina.

lobolepis, *F. Muell. Fragm.* viii. 258.—Austral.

lobulirostris, *Drej. Symb.* 27. t. 14.—Zeylan.

loliacea, *Linn. Sp. Pl.* 974.—Reg. bor. et arct.

loliacea, Schkuhr, Riedgr. i. 22 = tenella.

loliacea, Schreb. ex Steud. Nom. ed. II. i. 292 = divulsa.

loliacea, Steph. ex Boott, Illustr. Carex, iv. 187 = elongata.

loliacea, Thuill. ex Boott, Illustr. Carex, iv. 192 = muricata.

lonchicarpa, Willd. ex Kunth, Enum. Pl. ii. 500 = folliculata.

longearistata, Bivona, in Sicil. Sponte Prov. Desc. iv. 8. t. 2 = echinata.

longebrachiata, *Boeck. in Linnaea*, xli. (1877) 282.—Austral.

longebracteata, *Steud. Syn. Pl. Cyp.* 205.—Java.

longerostrata, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1841) 220.—Kamtschatka.

longibracteata, Dulac, Fl. Hautes-Pyr. 57 = C. Pseudocyperus.

longibracteata, Schleich. Cat. Pl. Helv. ed. IV. 11 = intermedia.

longicaulis, *Boeck. in Flora*, lxv. (1882) 62.—Mexic.

longicruris, *Nees, in Wight, Contrib.* 124.—Ind. or.

longicuinis, *Petrie, in Trans. N. Z. Inst.* xiv. (1882) 368.—N. Zel.

longifolia, Host, Gram iv. t. 85 = praecox.

longifolia, Thuill. Fl. Par. i. 490 = ampullacea.

longipes, *D. Don, in Trans. Linn. Soc.* xiv. (1825) 529. —Ind. or.

longirostris, Krock. ex Kunth, Enum. Pl. ii. 498 = laevigata.

longirostris, *Torr. ex Schwein. in Ann. Lyc. N. York*, i. (1824) 71.—Am. bor.

longiseta, *Brot. Fl. Lusit.* i 68.—Lusitan.

Lorentziana, *Griseb. in Goett. Abh.* xix. (1874) 271.—Reg. Antarct.

Lososoli, *Lange, in Kjoeb. Vidensk. Meddel.* (1877-78) 223.—Hispan.

lucida, *Boott, in Journ. Linn. Soc.* xx. (1884) 377.—N. Zel.

lucida, Dicks. ex Steud. Nom. ed. II. i. 292 = Elyna spicata.

lucorum, Sartw. ex Boott, Illustr. Carex, ii. 99 = nigromarginata.

lucorum, *Willd. Enum. Hort. Berol. Suppl.* 68.—Am. bor.

ludibunda, *Y. Gay, in Ann. Sc. Nat.* Sér. II. x. (1838) 357.—Gallia.

lupuliformis, Sartw. ex Dewey, in Am. Journ. Sc. Ser. II. ix. (1850) 29 = lupulina.

lupulina, *Muhl. ex Schkuhr, Riedgr.* ii. 54.—Am. bor.

lurida, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 153 = lupulina.

luxtanica, Schkuhr, Riedgr. ii. 28 = paniculata.

lutensis, *Kunth, Enum. Pl.* ii. 487.—Afr. austr.

lutesia, Nees, ex Boott, Illustr. Carex, iv. 181 = bracteosa.

lurulaefolia, *S. Wats. in Bot. Calif.* ii. 260.—Calif.

lusitina, *Olney, in Proc. Am. Acad.* vii. (1868) 395.—Calif.

Lyallii, *Boott, Illustr. Carex*, iv. 150.—Am. bor. occ.

Lyngbyei, Hornem. in Fl. Dan. t. 1888 = maritima.

Lyoni, *Boott, in Hook. Fl. Bor. Am.* ii. 209.—Am. bor.

Maackii, *Maxim. Prim. Fl. Amur.* 308.—Reg. Amur.

macilenta, *Kunth, Enum. Pl.* ii. 482.—Reg. Argent.

macilenta, Fries, Summa Veg. 224 = gracilis.

macloviana, Urv. in Mém. Soc. Linn. Par. v. (1826) 539 = leporina.

Macounii, *Dewey, in Am. Journ. Sc.* Ser. II. xli. (1866) 238.—Am. bor.

macra, Steud. Syn. Pl. Cyp. 213 = vulpina.

macrocarpa, *Phil. in Linnaea*, xxix. (1857-58) 86.—Chili.

macrocephala, *Willd. ex Kunth, Enum. Pl.* ii. 498.—Sibir. ; Japon.

macrochaeta, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 224. t. 18 = podocarpa.

macroglossa, *Franch. & Sav. Enum. Pl. Jap.* ii. 576. —Japon.

macrogyna, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104 = ferruginea.

macrolepis, *Steud. Syn. Pl. Cyp.* 223.—Am. bor.

Macounii, *Dewey, in Am. Journ. Sc.* Ser. II. xli. (1866) 238.—Am. bor.

macra, Steud. Syn. Pl. Cyp. 213 = vulpina.

macrocarpa, *Phil. in Linnaea*, xxix. (1857-58) 86.—Chili.

macrocephala, *Willd. ex Kunth, Enum. Pl.* ii. 498.—Sibir. ; Japon.

macrochaeta, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 224. t. 18 = podocarpa.

macroglossa, *Franch. & Sav. Enum. Pl. Jap.* ii. 576. —Japon.

macrogyna, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104 = ferruginea.

macrolepis, *Steud. Syn. Pl. Cyp.* 223.—Am. bor.

Macrostachys, *Link, in Schrad. Journ.* ii. (1799) 311. —Lusitan.

macrostachya, Bertol. Amoen. Ital. 421 = ferruginea.

macrostyla, Lapeyr. Hist. Abr. Pl. Pyr. 562 = utulata.

macrothyrsa, *Miq. Fl. Ind. Bat.* iii. 351.—Malaya.

maculata, *Boott, in Trans. Linn. Soc.* xx. (1851) 138. —Austral ; As. or.

maculata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 268 = spilocarpa.

maculata, Turcz. ex Ledeb. Fl. Ross. iv. 295 = utulata.

maculata, Turcz. ex Steud. Nom. ed. II. i. 292 = coriophora.

madagascariensis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 517.—Madag.

magellanica, Banks, ex Boott, in Proc. Linn. Soc. i. (1856) 287 = Banksii.

magellanica, *Lam. Encyc.* iii. 286.—Reg. bor. et austr.

CAREX:—

Mairii, *Coss. & Germ. Obs. Pl. Crit.* 18.—Mont. Pyren.

malacena, Steud. Syn. Pl. Cyp. 211 = vulgaris.

manca, *Boott, ex Benth. Fl. Hongk.* 402.—Hongkong.

manostachys, Spreng. Syst. iii. 824, sphalm. = macrostachys.

Marchandiana, Lapeyr. ex Spreng. l. c 806 = pyrenaica.

marcida, *Boott, in Hook. Fl. Bor. Am.* ii. 212.—Am. bor. occ.

marcida, *Raeusch. Nom.* ed. III. 270 (quid?).—Helvet.

margaritifera, Steud. Nom. ed. II. i. 292 = Scleria Flagellum.

marginata, Gort. Fl. Belg. 247 = divisa, intermedia.

marginata, Muhl. ex Schkuhr, Riedgr. ii. 49 = pensylvanica.

marina, Dewey, in Am. Journ. Sc. xxix. (1836) 347 = C. Heleonastes.

maritima, *O. F. Muell. in Fl. Dan.* t. 703.—Reg. bor. et arct.

Marssoni, Aschersw. in Bot. Zeit. x. (1852) 409 = flava.

Mastertiana, *Boott, Illustr. Carex*, i. 45.—Cochinch.

mauritanica, *Boiss. & Reut. Pugill. Pl. Nov.* 116.—Afr. bor.

mauritanica, *Steud. Syn. Pl. Cyp.* 204.—Ins. Mauric.

maxima, Scop. Fl. Carn. ed. II. ii. 229 = pendula.

Maximowiczii, *Boeck. in Linnaea*, xli. (1877) 237.—Reg. Amur.

Maximowiczii, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 180 = picta.

Maximowiczii, *F. Schmidt, Reisen Amurl.* 71.—Asa temp.

Meadii, Dewey, in Am. Journ. Sc. xliii. (1842) 90 = tetanica.

media, R. Br. in Richards. in Frankl. Narr. 1st Journ. 750 = alpina.

meiogyna, Nees, in Wight, Contrib. 128 = filicina.

melaena, Wimm. in Flora, xxxiii. (1850) 619 = vulgaris.

melananthа, *C. A. Mey. in Ledeb. Fl. Ross. iv.* 216.—As. centr.

melanocarpa, *Cham. in Middend. Sibir. Reise*, i. 11. 21 —Sibiria.

melanocephala, Turcz. ex Kunth, Enum. Pl. ii. 487 = alpina.

melanolepis, Drej. Fl. Lusit. i. 488 = stricta.

melanostylis, Desv. in C. Gay, Fl. Chil. vi. 208 = incurva.

melanosperma, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 270.—Mexic.

melanostachya, Willd. Sp. Pl. iv. 299 = nutans.

melanostylis, *Desv. in. C. Gay, Fl. Chil.* vi. t. 73.—Chili.

melastoma, Fisch. ex Boott, Illustr. Carex, ii. 100 = Tolmiei.

membranacea, Hook. in Parry, 2 Voy. App. 406 = vesicaria.

membranacea, Hoppe, Ic. t. 71 = ericetorum.

mendocinensis, *Olney, in S. Wats. Bot. Calif.* ii. 249. —Calif.

Menziesiana, Sm. ex Boott, Illustr. Carex, i. 27 = macrocephala.

mercarensis, *Hochst. ex Steud. Syn. Pl. Cyp.* 194.—Ind. or.

Mertensii, *Presc. ex Bong. Veg. Sitcha*, 51.—Alaska.

mertensis, Wethe, ex Kunth, Enum. Pl. ii. 503 = vulpina.

Metteniana, Lehm. ex Schur, Enum. Pl. Transs. 697 = dioica.

mexicana, Presl, Rel. Haenk. i. 304 = polystachya.

Meyeni, Nees, in Nov. Act. Nat. Cur. xix. (1843) Suppl. i. 123 = brunnea.

Meyeriana, *Kunth, Enum. Pl.* ii. 488.—Sibir. alt.

micans, *Boott, in Mem. Am. Acad.* N. S. vi. (1858-59) 419 = maculata.

Michauxiana, *Boeck. in Linnaea*, xli. (1877) 306.—Am. bor.

Michauxii, Dewey in Am. Journ. Sc. x. (1826) 278 = subulata.

Michauxii, Schwein. in Ann. Lyc. N. York, i. (1824) 64 = stipitoidea.

Micheliana, Sm. in Trans. Linn. Soc. v. (1800) 270 = glauca.

Micheliі, *Host, Syn. Pl. Austr.* 367.—Europ. austr. ; As. temp.

CAREX :—
microcarpa, *Bertol. ex Moris, Stirp. Sard. Elench.* 48. —Sardinia.
microcephala, C. A. Mey. in Ledeb. Fl. Alt. iv. 205 = obtusata.
microdonton, *Torr. in Ann. Lyc. N. York*, iii. (1886) 228.—Texas.
microglochin, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1800) 146.—Reg. bor. et arct.
microlepis, *Boeck. in Linnæa*, xli. (1877) 269.—Hab.?
micropoda, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1841) 210.—Unalaschk.
microsperma, *Steud. Syn. Pl. Cyp.* 281.—Texas.
microspermæ, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 144 = vulpinoides.
microstachya, *Ehrh. in Hannov. Mag.* 9ten Stück (1784), 132.—Europ. bor.
microstachya, Michx. Fl. Bor. Am. ii. 169 = polytrichoides.
microstyla, *Y. Gay, in Gaud. Fl. Helv.* vi. 87.—Alp. Helv.
Middendorffii, *F. Schmidt, Reisen Amuri.* 70.—Reg. Amur.
Mielichhoferi, Schkuhr, Riedgr. ii. 66 = ferruginea.
Mielichhoferi, Sm. Engl. Bot. t. 2293 = vaginata
Mielichhoferi, Willd. ex Kunth, Enum. Pl. ii. 465 = brachystachys, fimbriata.
milacea, Muhl. ex Schkuhr, Riedgr. ii. 77 = prasina.
milacea, Schrank, Baier. Fl. i. 300 = distans.
miliaris, *Michx. Fl. Bor. Am.* ii. 174.—Am. bor.
minima, *Boulu, in Ann. Soc. Bot. Lyon*, v. (1878) 88. —Gallia.
mirabilis, Dewey, in Am. Journ. Sc. xxx. (1836) 63 = straminea.
mirabilis, Host, Gram. iv. t. 78 = Elyna caricina.
mirabilis, Schleich. Cat. (1821) ex Gaud. Fl. Helv. vi. 65 = curvula.
mirata, *Dewey, in Am. Journ. Sc.* Ser. II. xxxix. (1865) 71.—Am. bor.
misandra, *R. Br. in Parry, Voy. App.* 283.—Am. arct.
miser, Buckl. in Am. Journ. Sc. xlv. (1843) 173 = jancea.
misera, *Phil. Fl. Atac.* 53.—Chili.
Mitchelliana, *M. A. Curt. in Am. Journ. Sc.* xliv. (1843) 84.—Am. bor.
mitis, J. F. Gmel. Syst. 1388 = Scleria mitis.
mixta, J. F. Gmel. l. c. 146 = stricta, paludosa.
mixta, *Midgev. in Bull. Soc. Bot. Fr.* xii. (1865) 348. —Gallia.
modesta, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 304 = disticha.
Monchiana, Wender. in Flora, vi. (1823) 718 = acuta.
monsta, *Kunth, Enum. Pl.* ii. 452.—Reg. Argent.
mollicula, *Boott, Illustr. Carex*, iv. 192.—Ins. Loochoo.
mollis, Gilib. Exercit. ii. 346 = leporina.
mollis, Host, Gram. iv. 46. t. 82 = præcox.
monadelpha, Boott, in Perry, Jap. Exped. ii. 323 = tristachya.
Montesi, *Tineb. in Bull. Soc. Bot. Fr.* iv. (1857) 164.—Gallia.
monile, *Tuckerm. ex Boott, in Hook. Lond. Journ. Bot.* v. (1846) 60.—Am. bor.
monilifera, Thuill. Fl. Par. i. 490 = depauperata.
Montviana, *Lowe, ex Boott, Illustr. Carex*, iv. 174.—Ins. Mader.
monostachya, *A. Rich. Tent. Fl. Abyss.* ii. 512.—Abyssin.
montana, Lecrs, Fl. Herborn. 716. t. 16. f. 6 = ericetorum.
montana, Linn. Sp. Pl. 975 = pilulifera.
montana, *Linn. Fl. Suec. ed.* II. 328.—Europ. ; Reg. Caucas. ; Japon.
montana, Poll. ex Lightf. Fl. Scot. ii. 551 = præcox.
montana, Scop. Fl. Carn. ed. II. ii. 219 = tomentosa.
monticola, Dewey, ex Torr. Bot. Mex. Bound. 229 = triquetra.
Moorcroftii, *Falc. in Proc. Linn. Soc.* i. (1846) 288.—Reg. Himal.
Morrowii, *Boott, in Perry, Jap. Exp.* ii. 326.—Japon.
Motoskei, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 148. —Japon.
mucronata, *All. Fl. Pedem.* ii. 268.—Europ. med. ; Reg. Caucas.
Muehlenbergii, Bronga. Voy. Duperr. Bot. 151 = Brongniartii.
Muehlenbergii, *Schkuhr, Riedgr.* ii. 12.—Am. bor.

CAREX :—
Muelleri, Steud. Syn. Pl. Cyp. 204 = Linkii.
Muhlenbergii, Kunth, ex Boott, Illustr. Carex, iii 125 = vulpinoides.
multicaulis, *L. Bailey, in Coult. Bot. Gaz.* ix. (1884) 118.—Calif.
multiceps, Gaud. Etrennes Fl. 78 = elongata.
multiculmis, Ehrh. Beitr vi. 83 = elongata
multiflora, Muhl. ex Schkuhr, Riedgr. ii. 14 = vulpinoidea.
multiformis, Thuill Fl. Par. i. 479 = disticha
multinervia, *Boeck. in Flora*, xxxviii (1855) 596.—Texas.
multinervis, Krock. ex Steud. Nom. ed II. i. 293 = binervis.
multispicata, *Kunze, in Schkuhr, Riedgr Suppl.* 42 t. 10.—Chili.
munda, *Boott, Illustr. Carex*, i. 7.— Reg. Himal.
muricata, Desf. Fl. Atlant. ii. 335 = divulsa
muricata, Huds. Fl Angl 408 = echinata,
muricata, Jungh in Linnæa, vi (1831) 29 = glomerata.
muricata, Lcers, Fl. Herborn. t 13. f. 8 = stellulata.
muricata, *Linn. Sp. Pl.* 974.—Reg. bor. temp.
muricata, Schleich. in Linnæa, vi. (1831) 29 = Bronguiartii.
musci, *Steud. Syn. Pl. Cyp.* 205.—Ins Borbon.
muskingumensis, *Schwein. in Ann. Lyc. N. York*, i. (1824) 66.—Am. bor.
mutabilis, Willd. Prod. Fl. Berl. 37 = pendula.
mutans, Boott, ex C. B. Clarke, in Journ. Linn. Soc xx. (1883) 388 = Hemicarex Hookeri.
mutica, R. Br. in Richards. in Frankl. Narr. 1st Journ. 731 = aurea.
myosuroides, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1833) 10.—Ins. Azoric.
myosuroides, Vill. Prosp 17 = Elyna spicata.
myosuroides, *Nees, in Wight, Contrib.* 122.—Ind. or.
nana, *Boott, in Mem. Am. Acad.* N. S. vi. (1858-59) 418.—Japon.
nana, *Chem. ex Steud. Syn. Pl Cyp* 228.—Alaska.
nana, Lam. Encyc. iii. 389 = ferruginea.
nardifolia, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 165 = aquatilis.
nardina, *Presc. Nov. Mant* iii. 55.—Europ. et Am. bor et arct.
Naufragii, Hochst. & Steud. Nom. ed II. i. 293 ; Syn. Pl. Cyp. 206 = Illegitima.
Naumanniana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 518.—Japon.
nebrascensis, Dewey, in Am. Journ. Sc. Ser. II. xviii. (1854) 102 = Jamesii.
Neesiana, Arn. ex Boott, in Proc. Linn. Soc. i. (1845) 260 = lobulirostris.
Necdhamii, *Engl. Prod. Fl. Norf.* 34.—Ins. Norfolk.
neglecta, Degl. in Loisel. Fl. Gall. ii. 298 = distans.
neglecta, Peterm. in Flora, xxvii. (1844) 331 = vulgaris.
neglecta, Tuckerm. Enum. 19 ; et in Am. Journ. Sc. xiv. (1843) 40 = rosea.
nematosperma, Willd. ex Kunth, Enum. Pl. ii. 456 = acuta.
nemorensis, J. F. Gmel. Syst. 1413 = pilosa.
nemorosa, J. F. Gmel. l. c. 1520 = pilosa.
nemorosa, Hort. Berol. ex Kunth, Enum. Pl. ii. 388 = laciorhyncha.
nemorosa, Lamarck. ex Honck. Syn. i. 356 = muricata.
nemorosa, Rebent. ex Boott, Illustr. Carex, iii. 132 = vulpina.
nemorosa, Schrank, Baier. Fl. i. 291 = alba.
nemostachys, *Steud. in Flora*, xxix. (1846) 23.—Japon.
nepalensis, Spreng. Syst. iii. 811 = longipes.
nervata, *Franch. & Sav. Enum. Pl. Jap.* ii. 566.—Japon.
nervina, *L. Bailey, in Coult. Bot. Gaz.* x. (1885) 203. —Calif.
nervosa, Desf. Fl. Atlant. ii. 337 = extensa.
nervosa, Willd. ex Kunth, Enum. Pl. ii. 449 = distans
neurocarpa, *Maxim. Prim. Fl. Amur.* 306.—Reg. Amur.
neurochlamys, F. Muell. Fragm. viii. 258 = maculata.
nevadensis, Boiss. & Kent. Pugill. Pl. Nov. 118 = Oederi.
nexa, *Boott, ex Benth. Fl. Hongk.* 402.—China.
nigella, *Boott, ex Hook. Fl. Bor. Am.* ii. 225.—Am. bor.
nigra, All. Fl. Pedem. ii. 267 = atrata.
nigra, Reich. Fl. Mono-Franc. ii. 96 = acuta.
nigra, Schkuhr, Riedgr. ii. 38 = sterrima.

CAREX :—
nigra, Schwein. ex Torr. in Ann. Lyc. N. York, i. (1824) 336 = vulgaris.
nigra, *Suvo. in Pall. N. Nord. Beitr.* vii. 198.—Sibir. baical.
nigricans, Dewey, in Am. Journ. Sc. xxix (1836) 249 = dioica.
nigricans, Hombr. ex Boott, Illustr. Carex, iv. 156 = Darwinii.
nigricans, *Kit in Linnæa*, xxxii (1863) 320.—Europ nigricans, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i (1831) 210. t. 7.—Unalaschk.
nigritella, Drej. ex Fl. Dan. t. 2369 = stylosa.
nigrolutea, Gaud. Agrost. Helv. ii 196 = glauca
nigro-marginata, *Schwein. in Ann Lyc. N. York*, i (1824) 66.—Am. bor.
nikoensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 558.— Japon
nilagirica, *Hochst. ex Steud. Syn. Pl. Cyp.* 207 — Ind or.
nipponica, *Franch. in Bull. Soc. Bot. Fr.* xxvi (1879) 89.—Japon.
nitens, *Phil. in Anal. Univ. Chil* (1873) 557 —Chili.
nitida, Hoppe, Taschenb. (1800) 108 = mucronata.
nitida, *Host, Gram* i 71.—Europ.
nitida, Willd. ex Kunth, Enum. Pl. ii. 411 = caespitosa.
nivalis, *Boott, in Proc Linn. Soc.* i. (1845) 256.— Reg. Himal.
nivalis, Cham. ex Steud. Syn. Pl. Cyp. 184 = micropoda.
nivicola, G Don, ex Boott, Illustr. Carex, iv. 140 = vaginata.
nobilis, *Boott, l. c.* i. 4.—Reg. Himal.
nodiflora, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 516.— Ins. Philipp.
novægica, *Willd. ex Schkuhr, Riedgr.* i. 50.—Europ. et Am. bor
notha, *Kunth, Enum. Pl.* ii 421.—Reg. Himal.
notolena, Nees, in Wight, Contrib. 121 = foliosa.
Novæ-Angliæ, *Schwein. in Ann. Lyc. N. York*, i. (1824) 67 —Am bor.
Novæ-Selandiæ, *Boeck. in Flora*, lxi. (1878) 169.— N Zel.
nubigena, *D. Don, in Trans. Linn. Soc.* xiv. (1825) 226.—Ind. or.
nuda, Lam. Fl. Fr. ii. 172 = leporina.
nudata, *S. Wats. Bot. Calif.* ii. 241.— Calif.
nuptialis, *Boott, Illustr. Carex*, iv 175. —Ins Sandvic
nutans, *Host, Gram.* i. 61. t. 83 — Europ. ; Japon.
nutans, Raeuach. Nom. ed III. 270 = erecta.
Nuttallii, *Dewey, in Am Journ. Sc.* xliii. (1842) 92.— Am bor. occ.
Oakesiana, Torr. in Ann. Lyc. N. York, iii. (1836) 421 = oligosperma.
obesa, *All. Fl. Pedem.* ii. 270 —Europ. ; Tauria.
obesa, Schleich. ex Kunth, Enum. Pl ii. 411 = caespitosa.
obliqua, Turcz. in Bull Soc. Nat. Mosc. (1838) 104 = pediformis.
oblita, *Steud. Syn. Pl. Cyp.* 221.—Am. bor.
oblonga, *F. F. Gmel Syst.* 146.—German.
obovata, Goldb. ex Ledeb. Fl. Ross iv. 290 = pediformis.
obscura, *Nees, in Wight, Contrib.* 126.—Reg. Himal.
obtusa, *Kit. in Linnæa*, xxxii. (1863) 319.—Hungar.
obtusangula, Ehrh Beitr. vi. 82 = ampullacea.
obtusangula, Hort. Berol ex Kunth, Enum. Pl. ii. 459 = glauca.
obtusangula, Salzm. ex Boott, Illustr. Carex, iv. 206 = echinata.
obtusangula, Salzm. ex Nym. Consp. 774 = hispida.
obtusata, *Liljeb. in Vet. Akad. Nya Handl. Stockh.* (1798) 69. t. 4.—Reg. bor. et arct.
ochroleuca, *Cham. ex Steud. Syn. Pl. Cyp.* 216.— Alaska.
odontolepis, *Phil. in Linnæa*, xxix. (1857-58) 82.— Chili.
Oederi, *Retz. ex Ehrh. Beitr.* vi. (1791) 83.—Reg. bor temp
Oederi, Willk. in Flora, xxxv. (1852) 521 = nevadensis.
oedipostyla, *Duval-Jouve, in Mem. Acad. Montp. (Sect. Sc.)* vii. (1869) 442.—Gallia.
ordocarpa, Anders. Cyp. Scand. 26 = Oederi.
Oerstedii, *Liebm. in Videnskk. Selsk. Skr.* V. ii. (1851) 272.—Costa Rica.
obiotica, *Hochst. & Steud. ex Steud. Nom.* ed. II. i. 293.—Am. bor.

CAREX:—
Ohmuelleriana, Lang, in Flora, xxvi. (1843) 240 = arenaria.
Oherieana, Dewey, in Am. Journ. Sc. xiv. (1828) 351 = oligosperma ?
olbiensis, *Jord. Obs. Pl. Crit.* iii. (1846) 241.—Europ.
Olgae, *Regel, Pl. Nov. Fedsch.* 33.—As. centr.
oligandra, F. Muell. ex Boott, Illustr. Carex, iv. 136 = Buxbaumii.
oligantha, *Boott, l. c.* 174. t. 589.—Am. austr.
oligantha, *Steud. Syn. Pl. Cyp.* 208.—Reg. Caucas.
oligocarpa, Hornem. ex Boott, Illustr Carex, iv. 161 = obesa.
oligocarpa, Muhl. Descr. Gram. Uber. 242; ex Boott, l. c. i. 40 = digitalis.
oligocarpa, *Schkuhr, Riedgr.* ii. 58.—Am. bor.
oligogyna, Less. ex Kunth, Enum. Pl. ii. 442 = globularis.
oligosperma, *Michx. Fl. Bor. Am.* ii. 174.—Am. bor.
oligostachya, *Nees, in Hook. New Journ.* vi. (1854) 29.—Ins. Philipp.
olivacea, *Boott, in Proc. Linn. Soc.* i. (1846) 286.—Reg. Himal.
olivacea, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 267.—Mexic.
Oliveri, *Boeck. in Flora*, lxiii. (1880) 455.—Afghanist.
Oliveriana, J. Gay, ex Boiss. Fl. Orient. v. 408 = illegitima.
Olneyi ×, Boott, Illustr. Carex, i. 15.—Am. bor.
olyssiponensis, Steud. Syn. Pl. Cyp. 204 = Linkii
omiana, *Franch. & Sav. Enum. Pl. Jap.* ii. 554.—Japon.
Onoei, *Franch. & Sav. l. c.* i. 551.—Japon.
ontakensis, *Franch. & Sav. l. c.* 550.—Japon.
orbicularis, Boott, in Proc. Linn. Soc. i. (1846) 254 = vulgaris.
oreades, C. A. Mey. in Bull. Acad. Brux. ix. 11. (1842) 248 = festiva.
oregonensis, *Olney, ex A. Gray, in Proc. Am. Acad.* viii. (1872) 407.—Am. bor. occ.
oreophila, *C. A. Mey. Verz. Pfl. Cauc.* 29.—Reg. Caucas.; As. Min.
Orinabar, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 268 = festiva.
orthostachya, *Willd. Sp. Pl.* iv. 255.—Europ.; As. bor.
ornithopodioides, Hausm. in Flora, xxxvi (1853) 225 = ornithopoda.
ornithorhyncha, Boott, Illustr. Carex, i. 26 = C. Cruscorvi.
Orsiniana, Tenore, Fl. Nap. v. 244 = muricata.
orthostachys, C. A. Mey. in Ledeb. Fl. Alt. iv. 231 = striata.
orthostachys, Trev ex Nym. Consp. 769 = hirta.
ovalis, Good. in Trans. Linn. Soc. ii. (1794) 148 = leporina.
ovalis, Willd. ex Kunth, Enum. Pl. ii. 397 = scoparia.
ovata, *Burm. f. Fl. Ind.* 194 (err. typ. 294).—Java.
ovata, Honck. Syn. i. 841 = Halleri.
ovata, C. A. Mey. in Ledeb. Fl. Alt. iv. 207 = curaica.
ovata, Rudge, in Trans. Linn. Soc. vii. (1804) 96. t. 9. f. 1 = strata.
oxylepis, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 409.—Texas.
pachygyna, *Franch. & Sav. Enum. Pl. Jap.* ii. 560.—Japon.
pachystachya, *Cham. ex Steud. Syn. Pl. Cyp.* 197.—Alaska.
pachystylis, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 301 = stenophylla.
pacifica, Drej. Revis. Car. 292 = caespitosa.
Patrowi, F. Schultz, in Flora, ii. (1868) 308 = divulsa.
paleacea, Wahlenb. in Vet. Acad. Nya Handl. Stockh. (1808) 164 = maritima.
paleata, *Boott, in Trans. Linn. Soc.* xx. (1846) 127.—Ins. Juan Fernandez.
pallescens, *Linn. Sp. Pl.* 977.—Reg. bor. temp.
pallida, C. A. Mey. in Mém. Sav. Istr. Pétersb. i. (1831) 212 = stenota.
pallida, Sellah. Prod. 29 = pallescens.
pallidior, Degl. in Loisel. Fl. Gall. ii. 299 = punctata.
paludosa, *Good. in Trans. Linn. Soc.* ii. (1794) 202.—Reg. bor. temp.
palustris, Suter, Fl. Helvet. ii. 261 = paludosa.
panicea, Carey, in A. Gray, Man. Bot. U. St. ed. I. 521 = Meadii.
panicea, Bunge, ex Boott, in Proc. Linn. Soc. i. (1846) 288 = tenuiculmis ?

CAREX:—
paniceа, *Linn. Sp. Pl.* 977.—Reg. bor. temp.
paniculata, *Linn. Cent. Pl.* i. 31 (*Amoen. Acad.* iv. 294).—Reg. temp. bor. et austr.
paniculata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1808) 143 = paradoxa.
paniculata, Willd. ex Kunth, Enum. Pl. ii. 390 = teretiuscula.
panormitana, Guss. Fl. Sic. Syn. ii. 575 = acuta.
papillosa, *Nees, in Hook. Journ. Bot.* ii (1840) 398.—Guiana.
papulosa, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 418.—Japon.
paradoxa, Bentb. Cat. Pl. Pyr. 67 = divisa.
paradoxa, J. F. Gmel. Syst. 145 = secq. ?
paradoxa, *Willd. in Schrift. Naturf. Fr. Berl.* (1794) 30. t. 1.—Europ.; As. bor.
parallela, Laest. in Vet. Akad. Nya Handl. Stockh. (1822) 338 = dioica.
parciflora, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 418.—Japon.
Parkeri, *Petrie, in Trans. N. Z. Inst.* xiii. (1881) 382. —N. Zel.
Parryana, *Dewey, in Am. Journ. Sc.* xxvii. (1835) 239.—Am. bor
parva, *Nees, in Wight, Contrib.* 120.—Reg. Himal.
parvibracteata, *Nees, l. c.* 125.—Reg. Himal.
parviflora, Gaud. Eremnes Fl. 84 = lagopina.
parviflora, Host, Gram. i. 9, 221 = strata.
parviflora, C. A. Mey. Verz. Pfl. Cauc. 30 = alpina.
patula, Host, Gram. i. 9, 221. t. 64 = flava.
patula, Huds. Fl. Angl. ed. II. ii. 402 = pauciflora.
patula, Schkuhr, Riedgr. i. 115. t. B. bb. f. 16 = laevigata.
patula, Scop. Fl. Carn. ed. II. ii. 226 = sylvatica.
pauciflora, Balb. ex Boott, Illustr. Carex, iv. 174 = microglochin.
pauciflora, *Lightf. Fl. Scot.* ii. 548. t. 6. f. 2.—Reg. bor. temp.
paucijlora, Usteri, ex Schkuhr, Riedgr. i. 83 = firma.
paupercula, Michx. Fl. Am. Bor. ii. 172 = magellanica.
pedata, All. Fl. Pedem. ii. 168 = ornithopoda.
pedata, *Linn. Sp. Pl.* ed. II. 1384.—Europ. arct.
pedifcrens, *C. A. Mey, in Mém. Sav. Etr. Pétersb.* i. (1831) 219. t. 10.—As. bor.
peduncularis, Wall. ex Boott, Illustr. Carex, i. 53 = speciosa.
pedunculata, *Muhl. ex Schkuhr, Riedgr.* ii. 89.—Am. bor.
pedunculata, Willd. ex Roxb. in Beata. St. Helena Trans. 1823 = praealta.
pelia, Lang, in Linnaea, xxiv. (1851) 575 = panicea.
pellita, Muhl. ex Schkuhr, Riedgr. ii. 84 = lanuginosa.
pellucida, Turcz. ex Boott, Illustr. Carex, iv. 196 = pediformis.
pendula, Geners. Elench. n. 871 = capillaris.
pendula, *Huds. Fl. Angl.* ed. I. 353; ed. II. 411.—Europ.; As. bor.
pendula, Moench, Meth. 324 = vesicaria.
pennsylvanica, Torr. in Am. Lye. N. York, iii. (1836) 411 = varia.
pensylvanica, *Lam. Encyc.* iii. 388.—Am. bor.
pentacarpa, *Boeck. in Flora*, lviii. (1875) 265.—Java.
pentastachya, *Fisch. ex Steud. Nom.* ed. II. i. 394.—Kamtschatka.
peregrina, *Link, Hort. Berol.* i. 334.—Afr. trop.
perennis, Krock. ex Steud. Nom. ed. II. i. 394 = marginata.
Perraudieriana, *Y. Gay, in Bull. Soc. Bot. Fr.* iii. (1856) 685, nomen.—Ins. Teneriff.
Persooni, Schkuhr, ex Kunth, Enum. Pl. ii. 509 = Wahlenbergiana.
Persoonii, Lang, in Flora, xxv. (1842) 748, 767 = canescens.
Persoonii, Sieber, ex Koch, Syn. Fl. Germ. ed. II. 871 = elongata, Gebhardi.
peruviana, Domb. ex Boott, Illustr. Carex, iv. 188 = acutata.
peruviana, Presl, Rel. Haenk. i. 205 = extensa.
petasata, Dewey, in Am. Journ. Sc. xxix. (1836) 246 = leporina.
Petersii, C. A. Mey. ex F. Schmidt, Reisen Amurl. 194 = falcata.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 513.—Abyssin.
petraea, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1809) 139 = rupestris.

CAREX:—
petricosa, *Dewey, in Am. Journ. Sc.* xxix. (1836) 246. —Am. bor.
Petriei, *Cheeseman. in Trans. N. Z. Inst.* xvi. (1884) 418. —N. Zel.
phaeota, *Spreng. Syst.* iii. 826.—Ind. or.; Malaya.
phaeostachya, Sm. Engl. Bot. Suppl. t. 2791 = vaginata.
phalaroides, *Kunth, Enum. Pl.* ii. 482.—Bras.
pharensis, Vis Fl. Dalm. iii. 146 = illegitima.
phleoides, Cav. Ic. v. 40 = Uncinia phleoides.
phyllomanica, *S. Wats. Bot. Calif.* ii. 238.—Calif.
phyllophora, Kunth, Enum. Pl. ii. 424 = pauciflora.
phyllostachys, *C. A. Mey. Verz. Pfl. Cauc.* 30.—Reg. Cauc.; Persia.
physena, *Dewey, in Am. Journ. Sc.* Ser. II. xxix. (1860) 247.—Am. bor.
physocarpa, Nees, ex Boott, in Proc. Linn. Soc. i. (1846) 287 = acutata.
physocarpa, *Presl, Rel. Haenk.* iii. 202.—Am. arct.
physodes, *Bieb. in Mém. Soc. Nat. Mosc.* ii (1809) 104. t. 7.—Reg. Caucas.; As. bor.
physorhyncha, *Steud. Syn. Pl. Cyp.* 719.—Mexic.
pichinchensis, *H. B. & K. Nov. Gen. et Sp.* i. 238.—Ecuador.
picta, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 418.—Japon.
picta, *Steud. Syn. Pl. Cyp.* 184.—Am. bor.
Pierotti, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 148.—Japon.
pilosa, *Scop. Fl. Carn.* ed. II. ii. 226.—Italia.
pilosiuscula, *Boeck. in Flora*, lxv. (1882) 61.—Am bor.
pilosiuscula x, *Gobi, in Just, Jahresb.* iv. (1876) 1078.—Rossia.
pilulifera, Geners. ex Boott, Illustr. Carex, iv. 216 = praecox.
pilulifera, *Linn. Sp. Pl.* 976.—Europ.; Japon.
pilulifera, *Willd. ex Kunth, Enum. Pl.* ii. 439.—Reg bor. et arct. ?
pinetorum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 268 = festiva.
pinetorum, Willd. ex Kunth, Enum. Pl. ii. 381 = Muehlenbergii.
pinnata, Boec. ex Boott, Illustr. Carex, iii. 124 = Muehlenbergii.
pinnata, Moench, ex Reichb. Fl. Germ. Excurs. 59 = elongata.
pistaliejs, Desv. in C. Gay, Fl. Chil. vi. 201 = setifolia.
pisiformis, *Boott, in Perry, Jap. Exped.* ii. 324.—Japon.
Pissii, *Phil. in Linnaea*, xxxiii. (1864–65) 272.—Chili.
planata, *Franch. & Sav. Enum. Pl. Jap.* ii. 555.—Japon.
plantiolin, *Kunth, in Oestr. Bot. Zeitschr.* xx. (1870) 141.—Europ.
planostachys, Kunze, in Schkuhr, Riedgr. Suppl. 188. t. 35 = Halleriana.
plantaginea, *Lam. Encyc.* iii. 392.—Am. bor.
plantaginea, Muhl. Descr. Gram. Uber. 236 = laxiflora.
platycarpa, Hochst. ex Steud. Syn. Pl. Cyp. 214 = phaeota.
platyphylla, *Carey, in Am. Journ. Sc.* II. iv. (1847) 28.—Am. bor.
platyrhyncha, *Franch. & Sav. Enum. Pl. Jap.* ii. 552. —Japon.
platystachys, Tenore, Fl. Nap. iv. 138 = macrolepis.
plumbea, Willd. Sp. Pl. iv. 308 = riparia.
poaeformis, J. F. Gmel. Syst. 138 = Scleria poaeformis.
pocilliformis, *Boott, Illustr. Carex*, iv. 178.—Japon.
podocarpa, *R. Br. in Richards. in Frankl. Narr. 1st Journ.* 751.—Am. arct.
podogyna, *Franch. & Sav. Enum. Pl. Jap.* ii. 557.—Japon.
podostachys, *Steud. Syn. Pl. Cyp.* 222.—Am. bor.
Poiretl, J. F. Gmel. Syst. 140 = Fulrena pubescens.
polyandra, Schkuhr, Riedgr. i. 59 = cruspitosa.
polyantha, *Franch. & Sav. Enum. Pl. Jap.* ii. 556.—Japon.
polyantha, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 110 = acuta.
polycephala, *Boott, Illustr. Carex*, i. 4.—Reg. Himal.
polygama, Schkuhr, Riedgr. i. 84 = Buxbaumii.

CAREX,—
polymorpha, *Muhl. ex Boott, Illustr. Carex*, i. 21. t. 57.—Am. bor.
polymorpha, Schwein. ex Boott, l. c. iii. 125 = vulpinoides.
polymorpha, Torr. in Ann. Lyc. N. York, iii. (1886) 413 = striata.
polyphylla, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 859.—Aa. centr.
polyrrhiza, Wallr. Sched. Crit. 492 = praecox.
polystachya, Lesson & A. Rich. Voy. Astrol. 118 = Lessoniana.
polystachya, A. Rich. Voy. Astrol. 111. t. 21 = ternaria.
polystachya, *Sw. ex Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1808) 149.—Am. trop.
polysticha, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 157.—Bras.
polytricholdes, *Muhl. ex Schkuhr, Riedgr.* ii. 84.—Am. bor.
Porteri, *Olney, Car. Bor. Am.* 12.—Am. bor.
potamostuarios, *Steud. in Lechl. Berb. Am. Austr.* 52. —Chili.
potosina, *Hemsl. Biol. Centr. Am. Bot.* iii. 474.—Mexic.
praealta, *Boott, Illustr. Carex*, i. 66.—Ins. S. Helen.
praecox, Borck. ex Reichb. Fl. Germ. Excurs. 64 = praecox.
praecox, *Jacq. Fl. Austr.* v. 23.—Reg. bor. temp.
praecox, Jundz. Fl. Lithuan. n. 889 = brizoides.
praecox, Poll. ex Steud. Nom. ed. II. i. 294 = emarginata.
praecox, Schreb. Spicil. 68 = Schreberi.
praegracilis, *W. Boott, in Coult. Bot. Gaz.* ix. (1884) 87.—Am. bor.
Praetutiana, Parl. Fl. Ital. ii. 182 = glauca.
praireа, Dewey, ex Woods, Bot. 760 = teretiuscula.
prairea, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1808) 161 = miliacea.
pratensis, Drej. Revis. Caric. 24 = adusta.
pratensis, *Hose, in Ustери, Ann. Bot.* xxi. (1797) 35.—Europ.
pratensis, *Phil. in Linnaea*, xxix. (1857–58) 81.—Chili.
Prelaii, *Nees, in Lehm. Pl. Preiss.* ii. 94.—Austral.
Prescottiana, *Boott, in Proc. Linn. Soc.* i. (1845) 286. —Calif.; Ins. Hawaii.
Prelii, *Steud. Syn. Pl. Cyp.* 242.—Sinus Nutka.
Princeps, Kunth, Enum. Pl. ii. 489 = praecox.
procera, *Kunth, l. c.* 491.—Bras.
prolifera, *Kit. in Linnaea*, xxxii. (1863) 320.—Hungar.
prolixa, Fries, Nov. Mant. iii. 228 = acuta.
prolixa, Schur, Enum. Pl. Transs. 707 = stricta.
prolixiuscula, Nees & Meyen [Reise, i. 116], ex Kunth, Enum Pl. ii. 396 = festiva.
prostrata, All. Fl. Pedem. ii. 267 = humilis.
prostrata, J. F. Gmel. Syst. 149 = pilosa.
protracta, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
provincialis, Degl. in Loisel. Fl. Gall. ii. 307 = echinata.
pruinosa, *Boott, in Proc. Linn. Soc.* i. (1845) 255.—Java.
psammogaea, Steud. Syn. Pl. Cyp. 187 = incurva.
psammophila, Schur, Enum. Pl. Transs. 700 = brizoides.
pseudo-arenaria, Pers. Syn. ii. 536 = disticha.
pseudo-arenaria, Reichb. Ic. Fl. Germ. viii. 8. t. 908 = armaria.
pseudo-brizoides, *Chavanel, in Bull. Soc. Pharm. Bord.* xiii. (1873) 156.—Gallia.
pseudo-Buxbaumii, *Winkler, in Oestr. Bot. Zeitschr.* xvii. (1868) 72.—Europ.
pseudo-conica, *Franch. & Sav. Enum. Pl. Jap.* ii. 570. —Japon.
pseudo-curaica, *F. Schmidt, Reisen Amurl.* 67.—Reg. Amur.
Pseudo-cyperus, R. Br. Prod. 248 = fascicularis.
Pseudo-cyperus, *Linn. Sp. Pl.* 978.—Reg. temp. bor. et austr.
pseudo-flava, *Schur, Enum. Pl. Transs.* 712.—Transsylv.
pseudo-loliacea, *F. Schmidt, Reisen Amurl.* 193.—Ins. Sachalin.
pseudo-nutans, *Bor. ex Martin, Cat. Remor.* 809.—Gallia.
pseudo-paradoxa, S. Gibson, in Phytol. i. (1844) 778 = teretiuscula.
pseudo-praecox, Schur, Enum. Pl. Transs. 716 = praecox.

CAREX,—
psilostachya, Kit. ex Willd. Sp. Pl. iv. 289 = sylvatica.
psilostachys, *Steud. Syn. Pl. Cyp.* 215.—Chili.
psychrophila, *Nees, in Wight, Contrib.* 127.—Reg. Himal.
psyllophora, Hartm. ex Reichb. Fl. Germ. Excurs. 61 = pauciflora.
psyllophora, Linn. f. Suppl. 413 = pulicaris.
pterolema, Kunze, ex Steud. Syn. Pl. Cyp. 242 = gibba.
pterygocarpa, *Steud. l. c.* 222.—Chili.
ptychocarpa, Link, in Schrad. Journ. ii. (1799) 309 = alpestris.
ptychocarpa, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
pubernula, Boott, in Perry, Jap. Exp. ii. 324 = brevicuinis.
pubescens, Gilib. Exercit. ii. 547 = montana.
pubescens, *Muhl. ex Schkuhr, Riedgr.* ii. 60. t. Exc. fig. 126.—Am. bor.
pubescens, Poir. Voy. Barb. ii. 254 = Fuirena pubescens.
pulchella, *Bergg. in Minnesbr. Fiuog. Sallsk. Lund* (1877) n. VIII. 29.—N. Zel.
pulchella, Nyland. ex Trevir, in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 537 = gracilis.
pulchra, *Boott, Illustr. Carex*, i. 4.—Reg. Himal.
pulicariosdes, Ramond, ex DC. Fl. Fr. iii. 101 = pyrenaica.
pulicaris, All. ex Boott, Illustr. Carex, iv. 174 = microglochin.
pulicaris, *Linn. Sp. Pl.* 972.—Europ.; Aa. bor.
pulicaris, Linn. f. ex Boott, Illustr. Carex, iv. 196 = pauciflora.
pulla, Gaud. in Trans. Linn. Soc. iii. (1797) 78. t. 14 = saxatilis.
pulla, Soland. ex Boott, Illustr. Carex, i. 64 = lucida.
pullulans, Dulac, Fl. Hautes-Pyr. 57 = Oederi.
pumila, Houck. Syn. Fl. Germ i. 273 = montana ?
pumila, *Thunb. Fl. Jap* 39.—China; Japon; Reg. austr. temp.
punctata, *Gaud. Agrost. Helv.* ii. 152.—Europ.; Afr. bor.
punctata, Nees, ex Wight, Contrib. 127 = notha.
punctulata, Leers, Fl. Herborn. 713. t. 14. f. 4 = paniculata.
puncticulata, Lesson, ex A. Rich. Voy. Astrol. Bot. 119. t 21 = C. Pseudo-cyperus.
pungens, *Boeck. in Linnaea*, xxxix. (1876) 46.—Chili.
Purdiei, Boott, Illustr. Carex, i. 96 = Bonplandii.
purpurea, *Boott, l. c.* iv. 157. t. 514.—Hab. ?
purpureovaginata, *Boeck. in Kjoeb. Vidensk. Meddel.* (1879–80) 30.—Bras.
pusilla, Pers. Syn. ii. 536 = bicolor.
pycnostachya, *Desv. in C. Gay, Fl. Chil.* vi. 204.—Chili.
pycnostachya, Kar. & Kir. in Bull. Soc Nat. Mosc. xv. (1842) 592 = C. Curaica.
pygmaea, *Boeck. in Linnaea*, xl. (1876) 379.—N. Granat.
pyrenaica, F. Muell. Fragm. viii. 251 = acicularis.
pyrenaica, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 139.—Europ.; Reg. Caucas.; Am. bor.; N. Zel.
pyriformis, F. Schultz, in Pollichia, xv. (1855) 122 = aurea.
pyriformis, Tausch, in Schultz, Arch. Fl. ii. 285 = flava.
quadrangulata, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 254.—N. Zel.
quaternaria, Sprang. Syst. iii. 809 = loliacea.
quinata, Raeusch. Nom. ed. III. 270 = caespitosa ?
radicalis, *Boott, Illustr. Carex*, i. 56.—Reg. Himal.
Raeana, *Boott, in Richardson, Arctic Exp.* ii. 344.—Am. bor.
Raffeliana, *Boott, in Proc. Linn. Soc.* i. (1845) 285.—Java.
Ramondiana, DC. Fl. Fr. iii. 101 = pyrenaica.
ramosa, Eckl. ex Kunth, Enum. Pl. ii. 581 = Schoenoxiphium Sieberianum.
ramosa, *Schkuhr, Riedgr.* ii. 40.—Ind. or.; China.
Rasulli, *Boott, in Hook. f. Fl. N. Zeal.* 288.—N. Zel.
raphidocarpa, *Nees, ex Wight, Contrib.* 122.—Ind. or.
rara, *Boott, in Trans. Linn. Soc.* xx. (1846) 189.—Zeylan.
rariflora, *Sm. Engl. Fl.* iv. 100.—Reg. bor. et arct.
Raynoldsii, *Dewey, in Am. Journ. Sc. Ser. II.* xxxii. (1861) 39.—Am. bor.
reclinata Facch. in Ambr. Fltr. Mcr. i. (1854) 333, 338 = ornithopoda.
recta, Boott, in Hook. Fl. Bor. Am. ii. 220 = salina.

CAREX,—
recurva, Huds. Fl. Angl. ed. II. 413 = glauca.
recurva, Muhl. Desc. Gram. Uber. n. 55 = cherokeusis.
recurva, Schkuhr, Riedgr. i. 120 = C. Pseudo-cyperus.
recurva, Willd. ex Kunth, Enum. Pl. ii. 418 = acuta.
recurvirostra, Hall. f. ex Steud. Nom. ed. I. 156 = Davalliana.
recurvirostra, *Kunze, in Schkuhr, Riedgr. Suppl.* 79 t. 20.—Java.
Redowskiana, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1881) 207. t. 4 = chordorrhiza.
reducta, Drej. Revis. Caric. 26 = salina.
reflexa, Dietz. ex Kunth, Enum. Pl. ii. 384 = glomerata.
reflexa, Gaud. Etrennes Fl. 53 = Davalliana.
reflexa, Hoppe, Caric. 65 = praecox.
refracta, Koth, Tent. Fl. Germ. ii. ii. 451 = praecox.
refracta, *Schkuhr, Riedgr.* ii. 62.—Mons Cenis.
Reinii, *Franch. & Sav. Enum. Pl. Jap.* ii. 569.—Japon.
remota, *Linn. Cent. Pl.* i. 31.—Reg. bor. temp. et arct.
remota, Richards. in Frankl. Narr. 1st Journ. 760 = Deweyana.
remota, Thunb. Fl. Jap. 37 = gibba.
remotiuscula, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 147 = cristata.
Renschiana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 515. —Mantag.
Rempi, Boeck. in Flora, lviii. (1875) 264 = brunnea.
repens, *Bell. App. Fl. Pedem.* 42.—Italia.
repens, Schleich. ex Kunth, Enum. Pl. ii. 376 = disticha.
reticulosa, *Peterm. in Flora*, xxvii. (1844) 333.—Saxonia.
retrocurva, Dewey, in Am. Journ. Sc. Ser. II. xlii. (1866) 243 = laxiculmis.
retroflexa, Muhl. ex Schkuhr, Riedgr. ii. 11 = rosea.
retrorsa, Nees, in Linnaea, x. (1835–36) 204 = cognata.
retrorsa, *Schwein. in Ann. Lyc. N. York*, i. (1824) 71. —Am. bor.
retusa, Degl. in Loisel. Fl. Gall. 807. t. 20 = echinata.
Retzii, Nees, in Wight, Contrib. 128 = chinensis
Reuteriana, Boiss. & Reut. in Fl. Orient. v. 411 = Huetiana.
Reuteriana, *Boiss. Pugill. Pl. Nov.* 116.—Hispan.
reversa, Gilib. Exercit. ii. 549 = C. Pseudo-cyperus.
reversa, Spreng. Syst. iii. 827 = retrorsa.
rhaphidocarpa, Kunth, Enum. Pl. ii. 512 = raphidocarpa.
rhisantha, J. G. Gmel. Syst. 144 = alpestris.
rhizodes, Hlytt, ex Boott, Illustr. Carex, iv. 196 = pediformis.
rhizomatosa, Steud. Syn. Pl. Cyp. 206 = Cumingiana.
rhynchocarpa, Heuff. in Flora, xvi. (1833) 365 = brevicollis.
rhynchophysa, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl.* 9.—Europ. et Aa. bor.
rhynchophysa, Liebm. in Videnak. Selsk. Skr. V. ii. (1851) 264 = physorhyncha.
rhynchosporoides, *Steud. Syn. Pl. Cyp.* 198.—Ins. Borbon.
Richardi, Mérat. Fl. Par. ii. 170 = tribuloides.
Richardi, Thuill. Fl. Par. i. 482 = canescens.
Richardsonii, *R. Br. in Richards. in Frankl. Narr. 1st Journ.* 761, 35.—Am. arct.
rigens, *L. Bailey, in Coult. Bot. Gaz.* ix. (1884) 117.—Am. bor.
rigens, Boott, in Mem. Am. Acad. N. S. vi. (1858–59) 419 = Brownii.
rigens, Thuill. Fl. Par. i. 488 = paludosa, stricta.
rigida, Clairv. ex Steud. Nom. ed. II. i. 295 = caespitosa.
rigida, Good. in Trans. Linn. Soc. ii. (1794) 198. t. 22 = vulgaris.
rigida, Schrank, Baier. Fl. i. 290 = firma.
rigida, Tuckerm. Enum. Caric. 11 = stylosa.
rigidifolia, Hochst. ex Seub. Fl. Aeor. 21 = punctata.
Ringoldiana, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 419.—Japon.
riparia, *Curt. Fl. Lond. Fasc.* iv. t. 60.—Reg. bor. et austr.
riparia, Gouan. Elench. n. 874 = paludosa.
riparia, Moench, ex Steud Nom. ed. II. i. 295 = acuta.
riparia, Muhl. Desc. Gram. Uber. 259 = laxistris.
riparia, Poir. Encyc. Suppl. iii. 382 = Uncinia riparia.

CAREX :—
rivalis, Willk. & Lange, Prod. Fl. Hisp. i. 180 = divisa.
rivularis, Koch, ex Kunth, Enum. Pl. ii. 487 = paludosa.
rivularis, Schkuhr, Riedgr. i. 30 = divisa.
robusta, F. Nyland, Spicil. Fenn. sub n. 33 = rhynchophysa.
Rochebruni, *Franch. & Sav. Enum. Pl Jap.* ii. 555. —Japon.
Roemeriana, *Scheele, in Linnaea*, xxii. (1849) 346.—Texas.
Romanzowiana, *Cham ex Steud. Syn. Pl. Cyp.* 216 — Alaska.
rosea, *Schkuhr, Riedgr.* ii. 15.—Am. bor.
Rossii, *Boott, in Hook. Fl. Bor. Am.* ii. 222.—Am. bor. occ.
rostellata, Gay, ex Boott, Illustr Carex, i. 8 = notha.
rostrata, Huppe, ex Schkuhr, Riedgr. i. 99 = Michelii.
rostrata, Michx. Fl Bor. Am. ii 173 = Michauxiana.
rostrata, Muhl. ex Schkuhr, l. c. ii. 54 = tentaculata.
rostrata, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. ii. 1050 = ampullacea.
Rotae, De Not. Ind. Sem. Hort. Gen. (1848) 24 = paniculata.
rotundata, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 153 —Lappon.
Royleana, Nees, in Wight, Contrib. 127 = breviculmis.
rubella, *Boott, Illustr. Carex*, iv. 176.—Zeylan.
rubescens, *Borch. in Flora*, lxv (1882) 60.—Japon.
rudis, *Wimm. in Flora*, xxxiii. (1850) 619.—Silesia.
rufa, Lam. Fl. Fr. ii. 184 = paludosa.
rufa, Schrank, Baier. Fl. i. 306 = paludosa.
rufidula, *Steud. Syn. Pl. Cyp.* 220.—Am. bor.
rufina, *Drej. Revis. Crit. Car.* 28.—Norveg.
Rugelisna, Kunze, in Linnaea, xx. (1847) 8 = juncea.
Ruiziana, *Boeck. in Linnaea*, xxxix. (1875) 30.—Peruv.
rupestris, *All. Fl. Pedem.* ii. 264. t. 92. f. 1.—Europ.
rupestris, Turcz. ex Ledeb. Fl. Ross. iv. 267 = argunensis.
Rutenbergiana, *Boeck. in Bremen Abh.* vii. (1880) 40. —Abyssin.
sabuletorum, Bieb. ex Boott, Illustr. Carex, iv. 178 = arenaria.
sabulosa, *Turcz. ex Kunth, Enum. Pl.* ii. 482.—Sibir. baical.
sabynensis, Less. ex Kunth, l. c. 440 = praecox.
sachalinensis, *F. Schmidt, Reisen Amurl.* 194.—Ins. Sachalin.
Sachspata, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
sagittifera, Lowe, in Camb. Phil. Soc. Trans. vi. (1858) 581 = macrostyla.
salina, Boott, in Hook. Fl. Bor. Am. 219 = cryptocarpa.
salina, Bongard, ex Ledeb. Fl. Ross. iv. 281 = canescens.
salina, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1808) 165.—Norveg.
salinoides, Beurl. in Bot. Notiser, (1853) 35 = salina.
Salsinana x, Bruegg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 130.—Europ.
Sampsoni, *Hance, in Journ. Bot.* vi. (1868) 89.—China.
sandwicensis, *Boeck. in Flora*, lvii. (1875) 265.—Ins. Pacif.
sanguinea, *Boott, in Trans. Linn. Soc.* xx. (1846) 187. —Afghanist.
sarda, Link, Hort. Berol. ii. 328 = Linkii.
Sartwelliana, Dewey, in Am. Journ. Sc. xliii. (1842) 90 = disticha.
Sartwelliana, *Olney, in Proc. Am. Acad.* vii. (1868) 396.—Calif.
Sartwellii, Dewey, ex J. Carey, in A. Gray, Man. Bot. U. St. ed. I. 539 = Sartwelliana.
saskatschewana, *Boeck. in Linnaea*, xli. (1877) 159.—Am. bor.
satsumensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 558. —Japon.
saxatilis, Huds. Fl. Angl. 408 = praecox.
saxatilis, *Linn. Sp. Pl.* 976.—Europ.
saxatilis, Matusch. ex Kunth, Enum. Pl. ii. 450 = fulva.
saxatilis, Ced. in Fl. Dan. t. 159 = vulgaris.
saxatilis, Scop. Fl. Carn. ed. II. ii. 221 = atrata.
saxatilis, Suter, ex Boot' Illustr. Carex, iv. 218 = sempervirens.

CAREX :—
scabberrima, *Scheele, in Linnaea*, xxii. (1849) 345.—Texas.
scabra, Hoppe, Taschenb. (1800) 242 = Davalliana.
scabrata, *Schwein. in Ann. Lyc. N. York*, i. (1824) 69. —Am. bor.
scabrella, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1808) 149.—Ind. occ.
scabrifolia, *Steud. Syn. Pl. Cyp.* 287.—Java.
scabrata, Sartwell, ex Boott, Illustr. Carex, iii. 125 = vulpinoidea.
scariosa, Lam. Encyc. iii. 388 = humilis.
Schaffneri, *Boeck. in Flora*, lxi. (1878) 89.—Mexic.
Schaffneri, W. Boott, ex S. Wats. in Proc. Am. Acad. xviii. (1883–83) 172 = potosina.
schedonnatos, *Steud. Syn. Pl. Cyp.* 189 —Reg. Magell.
Schenchseri, Gand. ex Boott, Illustr. Carex, iv. 204 = tenuis.
Schenckerri, J. F. Gmel. Syst. 144 = paludosa!
Schiedeana, *Kunze, in Schkuhr, Riedgr. Suppl.* 119. t. 30.—Mexic.
Schimperiana, *Boeck. in Linnaea*, xl. (1876) 373.—Abyssin.
Schkuhrii, Willd. Sp. Pl. iv. 264 = obesa.
Schmidtii, *Boeck. in Flora*, lxv. (1882) 63.—Ins. Sachalin.
Schmidtii, *Meinshaus. Nachr. Wilmi-Geb.* 224.—Manchuria.
schoenoides, DC. Fl. Fr. iii. 104 = intermedia.
schoenoides, Desf. Fl. Atlant. ii. 336 = divisa.
schoenoides, *Schrank, Baier. Fl.* i. 274.—Europ.
schoenoides, *Thuill. Fl. Par.* 480.—Gallia.
Schotti, Bolss. Diagn. Ser. II. iv. 123 = sempervirens.
Schottii, *Dewey, ex Torr. Bot. Mex. Bound.* 231.—Calif.
Schraderi, Schkuhr, Riedgr. ii. 81 = laevigata.
Schreberi, Desv. Fl. Auv. 78 (Scherberi) = arenaria.
Schreberi, *Schrank, Baier. Fl.* i. 278.—Europ.; As. bor.
Schreberi, Tenore, ex Boott, Illustr. Carex, iv. 186 = divisa.
Schrebers, Willd. in Mém. Acad. Berl. 1794 (1797) 38 = obesa.
Schrenkiana, C. A. Mey. ex Ledeb. Fl. Ross. iv. 317 = vesicaria.
Schummelii, *Siegert, in Verh. Bot. Brandenb.* (1860) 113.—Europ.
Schweinitzii, *Dewey, in Am. Journ. Sc.* ix. (1825) 68. —Am. bor.
scirpina, Nackerm. ex Boott, Illustr. Carex, iv. 150 = scirpoides.
scirpoides, *Michx. Fl. Bor. Am.* ii. 171.—Am. bor.
scirpoides, Schkuhr, Riedgr. ii. 19 = echinata.
scitula, *Boott, Illustr. Carex*, iv. 177.—Reg. Himal.
scoparia, *Schkuhr, Riedgr.* ii. 20.—Am. bor.
scoparia, Torr. in Ann. Lyc. N. York, iii. (1836) 394 = virgata.
Scopoliana, Willd. Sp. Pl. iv. 299 = ferruginea.
Scopolii, Gaud. Agrost. Helv. ii. 168 = ferruginea.
scotica, Spreng. Syst. iii. 819 = vaginata.
scouleri, Torr. in Ann. Lyc. N. York, iii. (1836) 398 = cryptocarpa.
secale, Ledeb. Fl. Ross. iv. 300 = hordeistichos.
secalina, Lapeyr. Hist. Abr. Pl. Pyr. ii. 576 = riparia.
secalina, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1808) 151 = hordeistichos.
secta, *Boott, in Hook. f. Fl. N. Zel.* i. 281.—N. Zel.
sedition, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
Sellowiana, *Schlecht. in Linnaea*, x. (1836) 117.—Bras.
semicylindrica, *Kit. l. c.* xxxii. (1863) 819.—Hungar.
sempervirens, Ell. ex Kunth, Enum. Pl. ii. 460 = communis!
sempervirens, Schwein. in Ann. Lyc. N. York, i. (1824) 70 = glaucescens.
sempervirens, *Vill. Hist. Pl. Dauph.* ii. 214.—Europ.
Semdtneriana x, Bruegg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 119.—Europ.
senta, *Boott, Illustr. Carex*, iv. 174, in nota.—Calif.
serotina, Tenore, Fl. Nap. iv. 133 = muricata.
serratodens, *S. Wats. Bot. Calif.* ii. 246.—Calif.
serrulata, Bivona, ex Spreng. Syst. iii. 827 = glauca.
serrulata, Mutel, Fl. Fr. 396 = echinata.
setacea, Dewey, in Am. Journ. Sc. lx. (1825) 61. t. B. f. 5 = vulpinoidea.
setacea, Frank, ex Kunth, Enum. Pl. ii. 383 = stipata.

CAREX :—
setacea, Froel. ex Kunth, Enum. Pl. ii. 409 = mucronata.
setariaeformis, Turcz. ex Boott, Illustr. Carex, iv. 189 = leiorhyncha.
seticulmis, *Boeck. in Kjoeb. Vidensh. Meddel.* (1869) 156.—Bras.
setifolia, Godr. in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 14 = divisa.
setifolia, *Kunze, Syn. Pl. Am. Austr.* 36.—Chili.
setigera, *D. Don, Prod. Fl. Nep.* 43.—Reg. Himal.
setosa, *Boott, Illustr. Carex*, iii. 108.—Reg. Himal.
setosa, Franch. & Sav. Enum. Pl. Jap. 142 = homoiolepis.
sexspicata, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 345.—N. Zel.
Shortiana, *Dewey & Torr. in Am. Journ. Sc* xxx (1836) 60.—Calif.
Shortii, Steud. Syn. Pl. Cyp. 202 = stenolepis.
Shortii, Torr. in Ann. Lyc. N. York, iii. (1836) 407 = Shortiana.
sibirica, Willd. ex Kunth, Enum. Pl. ii. 406 = loliacea.
sicaeformis, *Boott, in Bost. Journ. Nat. Hist.* v. (1845) 118.—Am. bor.
siccata, *Dewey, in Am. Journ. Sc.* x. (1826) 27.—Am bor.; As. bor. or.
sicula, Presl, ex Boott, Illustr. Carex, iv. 185 = distans.
sicula, *Tineo, Pl. Rar. Sicil.* 30.—Sicilia.
sicyocarpa, Lebel, Obs. Pl. Manche (1848) 18 = praecox.
siderostichs, *Hance, in Journ. Linn. Soc.* xiii. (1873) 89.—China.
Sieberi, Nees, ex Kunth, Enum. Pl. ii. 118 = Maraccs flavus.
Siebersana, Opiz, Natural. xi. (1826) 413 = Davalliana.
Sieboldi, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 150 — Japon.
Siegertiana, Uechtritz, ex Garcke, Fl. Deutschl. ed VIII. 485 = hirta.
sikokiana, *Franch. & Sav. Enum. Pl. Jap.* ii. 573. — Japon.
silaca, *Olney, in Proc. Am. Acad.* vii. (1878) 393.—Am. bor.
simensis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 514 = aethiopica.
similis, Urv. in Mém. Soc. Linn. Par. iv. (1826) 599 = curta.
simplicissima, F. Muell. Fragm. ix. 191 = capillacea simplicissima, *Wahlenb. in Vet. Akad. Nya Handl Stockh.* (1808) 141 (Quid ?).—Anglia.
Sinai, *Boott, in Trans. Linn. Soc.* xx. (1851) 146.—Arab.
sinaica, Nees, ex Steud. Syn. Pl. Cyp. 228 = distans.
sinensis, J. F. Gmel. Syst. 144 = chinensis.
sitchensis, *Presc. ex Bong. Veg. Sitcha*, 51.—Alaska; Am. bor. occ.
Smithii, Tausch, in Flora, iv. (1821) 559 = vaginata.
socia, Boott, in Proc. Linn. Soc. i. (1845) 254 = Arnottiana.
socista, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 490.—China.
Solandri, *Boott, in Hook. f. Fl. N. Zel.* 284.—N. Zel.
Solerioli, DC. & Duby, Bot. Gall. ii. 498 = echinata.
songorica, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 535.—Afghanist.; Soongar.
sordida, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 33.—Kamtschatka.
sororia, *Kunth, Enum. Pl.* ii. 379.—Reg. Argent.
sorsria, Meinsh. Fl. Ingr. 404 = agastra.
Spachiana, *Boott, Illustr. Carex*, i. 43.—China.
spadicea, Gilib. Exercit. ii. 546 = arenaria.
spadicea, J. F. Gmel. Syst. 144 = firma.
spadicea, Host, Gram. iv. 51. t. 91.; DC. Fl. Fr. iii. 130 = ferruginea.
spadicea, Roth, Tent. Fl. Germ. ii. xi. 461 = paludosa.
spadicea, Schkuhr, Riedgr. i. 90 = frigida.
spaniocarpa, Steud. Syn. Pl. Cyp. 228 = obesa.
sparganioides, Le Conte, ex Kunth, Enum. Pl. ii. 382 = stipata.
spargaioides, *Muhl. ex Schkuhr, Riedgr.* ii. 14.—Am. bor.
sparsiflora, Steud. Nom. ed. II. i. 396 = vaginata.
spartea, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 149 = Uncinia spartea.
spatiosa, *Boott, Illustr. Carex*, ii. 26.—Cochinch.

CAREX:—
speciosa, *Kunth, Enum. Pl.* ii. 504.—Ind. or.
spectabilis, Dewey, in Am. Journ. Sc. xxix. (1836) 248. t. X. f. 76 = podocarpa.
spicrostachya, Sw. ex Sm. Engl. Fl. iv. 98 = fulva.
sphaerica, *Lapeyr. Hist. Abr. Pl. Pyr.* 570.—Mont. Pyren.
sphaerocarpa, Ehrh. Beitr. vi. 88 = tomentosa.
sphaerocarpa, Willd. Sp. Pl. iv. 265 = obesa.
sphaerogyne, *Baker, in Journ. Bot.* xxi. (1883) 129.—Madag.
sphaerostachya, *Dewey, in Am. Journ. Sc.* xlix. (1845) 44. t. Es. f. 110.—Am. bor.
spicata, Brot. ex Kunth, Enum. Pl. ii. 373 = divisa.
spicata, Huds. Fl. Angl. ed. I. 358 = muricata.
spicata, Lam. Encyc. iii. 381; Pollich, Palat. ii. 562 = disticha.
spicata, Schkuhr, Riedgr. i. 11 = obtusata.
spicata, Thuill. Fl. Par. 480 = vulpina.
spicigera, *Nees, in Wight, Contrib.* 121.—Zeylan.
spicinata, *Boott, in Trans. Linn. Soc.* xx. (1851) 139.—Reg. Himal.
spiculosa, F. Nyland. Spicil. Pl. Fenn. ii. n. 49 = salina.
spilocarpa, *Steud. Syn. Pl. Cyp.* 195.—Mexic.
spinirostris, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 335.—N. Zel.
splendens, Thuill. ex Pers. Syn. iii. 536 = divisa.
splendida, Willd. Prod. Fl. Berl. 33 = filiformis.
Sprengelii, *Boeck. in Linnaea*, xl. (1876) 871.—Afr. austr.
Sprengelii, Dewey, ex Spreng. Syst. iii. 837 = longirostris.
sporta, Steud. Syn. Pl. Cyp. 212 = caespitosa.
squarrosa, *Linn. Sp. Pl.* 978.—Am. bor.
stans, Drej. Revis. Caric. 40 = aquatilis.
Steenstrupii, *Liebm. in Forhand. Scand. Naturf.* (1840) 231.—Islandia.
stellulata, Bieb. Fl. Taur. Cauc. ii. 388 = muricata.
stellulata, Good. in Trans. Linn. Soc. ii. (1794) 144 = echinata.
stenantha, *Franch. & Sav. Enum. Pl. Jap.* ii. 573.—Japon.
stenocarpa, Turcz ex Steud. Nom. ed. II. i. 296 = frigida.
stenolepis, *Lyss. ex Steud. l. c.*—Norveg.
stenolepis, *Torr. in Am. Journ. Sc.* xxx. (1836) 59.—Am. bor.
stenophylla, Thien. ex Boott, Illustr. Carex, iv. 210 = incurva.
stenophylla, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 142.—Reg. bor. temp.
stenostachys, *Franch. & Sav. Enum. Pl. Jap.* ii. 569.—Japon.
sterilis, *Schkuhr, Riedgr.* ii. 8.—Am. bor.
Steudelii, Kunth, Enum. Pl. ii. 480 = Jamesii.
Steudneri, *Boeck. in Linnaea*, xl. (1876) 864.—Abyssin.
stictocarpa, Sm. Engl. Bot. t. 2772 = glauca.
stipata, *Muhl. ex Schkuhr, Riedgr* ii. 12.—Am. bor.
stipitata, Boeck. in Linnaea, xxxix. (1875) 85 = stipata.
stolonifera, Ehrh. Beitr. vi. 88 = muricata.
stolonifera, Hoppe, Ic. n. 41 = vulgaris.
straminea, *Willd. ex Schkuhr, Riedgr.* i. 49.—Am. bor.
striata, B. Br. Prod. 243 = laciniosa.
striata, Gilib. Exercit. ii. 550 = riparia.
striata, *Michx. Fl. Bor. Am.* ii. 174.—Am. bor.
strata, Torr. ex Boott, Illustr. Carex, i. 22 = polymorpha.
striatula, Michx. Fl. Bor. Am. ii. 178 = laxiflora.
stricta, *Good. in Trans. Linn. Soc.* ii. (1794) 196. t. 21. f. 9.—Europ.; Am. bor.
stricta, Lam. Encyc. iii. 387 = angustata.
strictifolia, Opiz, ex Nym. Consp. 777 = acuta.
strictior, Dewey, ex Wood, Cl. Book. 755 = stricta.
strigosa, All. Fl. Pedem. ii. 270 = tenuis.
strigosa, *Huds. Fl. Angl.* 411.—Europ.; Reg. Caucas.
strigosa, Soter, ex Boott, Illustr. Carex, iv. 208 = firma.
strigosa, Willd. ex Kunth, Enum. Pl. ii. 470 = sylvatica.
strygia, Fries, Mant. iii. 141 = rariflora.
stylosiflora, Buckl. in Am. Journ. Sc. xiv. (1848) 174 = laxiflora.
stylosa, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 332. t. 14.—Unalaschka.
subanceps, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 320.—Japon.

CAREX:—
subbiflora, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
subcompressa, *Steud. l. c.* 231.—Am. bor.
subdola, *Boott, in Proc. Linn. Soc.* i. (1845) 255.—N. Zel.
subdura, Boott, Illustr. Carex, iv. 165 = Pierotii.
subfusca, *S. Wats. in Bot. Calif.* ii. 234.—Calif.
subglobosa, Mielich. in Flora, xxii. (1839) 256 = Oederi.
subhirtella, *Norrl. in Notis. Sællsk. Fauna et Fl. Fenn.* ii. *Haeft* x. 51.—Europ. bor.
subramosa, Willd. ex Kunth, Enum. Pl. ii. 385 = echinata.
subrotunda, *Serres, in Bull. Soc. Bot. Fr.* iv. (1857) 440.—Galilia.
subsalsa, Nym. Syll. Suppl. 69 = fulva.
subsphaltaica, Wormsk. in Fl. Dan. ix. 4. t. 1530 = salina.
subulata, J. F. Gmel. Syst. 138 = Scleria subulata.
subulata, *Michx. Fl. Bor. Am.* ii. 178.—Am. bor.
subulata, Schum. ex Steud. Nom. ed. II. i. 296 = Buxbaumii.
subulata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 152 = Boryana.
subuniflora, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
subvillosa, Bieb. Fl. Taur. Cauc ii. 386 = tomentosa.
sudetica, Opiz, Natural. xii. (1826) 413 = strata.
sudetica, Presl, ex Kunth, Enum. Pl. ii. 451 = vaginata.
suilla, J. Fellm. in Bull. Soc. Nat. Mosc. viii. (1835) 285 = pallida!
sulcata, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 80.—Transsylv.
Sullivantii × Boott, in Am. Journ. Sc. xlii. (1842) 29.—Am. bor.
supina, *Wahlenb. in Akad. Vet. Nya Handl. Stockh.* (1803) 158.—Reg. bor. temp.
Sutera, *Kunth, Enum. Pl.* ii. 523.—Europ.
sychnocephala, *Carey, in Am. Journ. Sc.* Ser II. x (1847) 24.—Am. bor.
sylvatica, Dewey, l. c. Ser. I. x. (1826) 40 = arctata.
sylvatica, *Huds. Fl. Angl.* ed. I. 353.—Europ.; As. temp.
sylvatica, Maxim. ex Boeck. in Linnaea, xli. (1877) 238 = Maximowiczii.
tabina, *Steud. in Lechl. Berb. Am. Austr.* 56.—Am. austr.
tabularis, M. A. Curt. ex Boott, Illustr. Carex, i. 54 = acutivalvis.
tartarea, *Ridl. in Journ. Bot.* xxi. (1883) 262.—Sumatra.
taurorum, Bolss. Fl. Orient. v. 415 = strata.
teinogyna, *Boott, Illustr. Carex*, i. 60.—Reg. Himal.
tenebrosa, *Boott, l. c.* 88. t. 256.—China.
tenax, *Bergg. in Minnesota, Pisog. Salics. Lund* (1877), n. VIII. p. 27.—N. Zel.
tenax, *Chapm. ex Dewey, in Am. Journ. Sc.* Ser. II. xix. (1855) 254.—Am. bor.
tenax, *Kunz. in Comptes-Rend. Soc. Hallér.* (1854–56) 130.—Europ.
tenella, Ehrh. Beitr. vi. 88 = canescens.
tenella, Froel. ex Kunth, Enum. Pl. ii. 445 = supina.
tenella, Pohr. Encyc. Suppl. iii. 292 = Uncinia tenella.
tenella, *Schkuhr, Riedgr.* i. 28.—Am. bor.
tenella, Thuill. Fl. Par. i. 429 = praecox.
teneriformis, Dewey, in Am. Journ. Sc. viii. (1824) 97 = straminea.
Tenorii, Kunth, Enum. Pl. ii. 437 = macrolepis.
tentaculata, *Muhl. ex Willd. Sp. Pl.* iv. 266.—Am. bor.
tenuiflora, Hartm. ex Kunth, Enum. Pl. ii. 405 = loliacea.
tenuiflora, *Wahlenb. in Vet. Acad. Nya Handl. Stockh.* (1803) 147.—Europ.; Am. bor.
tenuifolia, Poir. Voy. Barb. ii. 254 = Scirpus pubescens.
tenuirostris, *Olney, in Am. Natural.* viii. (1874) 214.—Am. bor.
tenuis, Gmel. ex Steud. Nom. ed. II. i. 296 = Scleria tenuis.
tenuis, *Host, Gram. Austr.* iv. 51. t. 92.—Europ. austr.
tenuis, Nees, in Wight, Contrib. 136 = hirtella.
tenuis, Rudge, in Trans. Linn. Soc. vii. (1804) 97. t. 9. f. 3 = debilis.
tenuis, Schleich. ex Steud. Nom. ed. II. i. 296 = leporina.
tenuistylos, Boeck. in Flora, xxxix. (1856) 225 = tristicha, *Boeck.*

CAREX:—
tenuispica, *Steud. Syn. Pl. Cyp.* 219.—Ins. Borbon.
tenuissima, *Boott, in Proc. Linn. Soc.* i. (1845) 288.—Japon.
teres, *Boott, Illustr. Carex*, i. 62.—Reg. Himal.
tereticulis, *F. Muell. Fragm.* viii. 256.—Austral.
teretiuscula, *Good. in Trans. Linn. Soc.* ii. (1794) 163.—Europ.; Am. bor.
teretiuscula, Willd. ex Kunth, Enum. Pl. ii. 389 = paniculata.
teretiusculs, Willk. in Bot. Zeit. vi. (1848) 414 = chordorrhiza.
tergestina, Hoppe, ex Boott, Illustr Carex, iv. 193 = muricata.
ternaria, *Forst. f. Prod.* 92.—N. Zel.
testacea, *Soland. ex Boott, in Hook. f. Fl. N. Zel.* i. 282.—N. Zel.
tetanica, Reichb. Fl. Germ. Excurs. 68 = vaginata.
tetanica, *Schkuhr, Riedgr.* ii. 68.—Groenl.; Am. bor.
tetanica, Schwein. & Torr. in Ann. Lyc. N. York, i. (1824) 347 = conoidea.
tetrastachya, Scheele, in Linnaea, xxii. (1849) 847 = straminea.
tetrastachya, *Traunst. in Flora*, xxxiii. (1850) 866.—Europ.
Textori, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 151 = Inuita.
thecata, Boott, in Trans. Linn. Soc. xx. (1851) 143 = Preissii.
thermarum, *Phil. in Anal. Univ. Chil.* xxi. (1862) 11. 885.—Chili.
Thomsonii, *Boott, Illustr. Carex*, i. 1.—Reg. Himal.
Thouarii, *Carmich. in Trans. Linn. Soc.* xii. (1818) 508.—Ins. Tristan. D'Acunha.
Thunbergii, *Steud. in Flora*, xxix. (1846) 23.—Japon.
Thurberi, *Dewey, ex Torr. Bot. Mex. Bound.* 282.—Am. bor. occ.
thuringiaca, Willd. Sp. Pl. iv. 250 = glauca.
Thwaitesii, *Hance, in Journ. Bot.* v. (1867) 225.—Zeylan.
thyrsiflora, *Boott, Illustr. Carex*, i. 12. t. 34.—Ind. or.
tokiensis, *Boeck. in Flora*, lxv. (1882) 68.—Japon.
Tolmiei, *Boott, in Hook. Fl. Bor. Am.* ii. 224.—Am. bor. occ.
tomentosa, Hornem. in Fl. Dan. t. 1404 = globularis.
tomentosa, Lightf. Fl. Scot. 553 = filiformis.
tomentosa, *Linn. Mant.* 123.—Europ.; Reg. Caucas.
tomentosa, C. A. Mey. Verz. Pfl. Cauc. 192 = grinca.
torfacea, J. F. Gmel. Syst. 145 = rampallacea.
Torreyana, Dewey, in Am. Journ. Sc. x. (1826) 47 = Davisii.
Torreyi, *Tuckerm. Enum. Caric.* 21.—Am. bor.
torta, *Boott, ex Tuckerm. l. c.* 11.—Am. bor.
Tourarginiana, Bor. Fl. Cent. Fr. ed. II. 582 = acuta.
trachyantha, Dorner, ex Heuff. in Verh. Zool. Bot. Ges. Wien, viii. (1858) 219 = praecox.
trachycarpa, Link, in Schrad. Journ. ii. (1799) 810 = glauca.
trachycystis, *Griseb. in Goett. Abh.* xxiv. (1879) 314.—Reg. Argent.
tralstscana, *F. Schmidt, Reisen Amurl.* 192.—Ins. Sachalin.
transversa, *Boott, in Hook. Fl. Bor. Am.* ii. 215. t. 216.—Japon.
transversa, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 148 = rigens.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 80 = Hallerlana!
triandra, *Gronov., ex Trauty. in Act. Hort. Petrop.* ix. (1884) 811, nomen.—Russia.
triangularis, *Boeck. in Flora*, xxxix. (1856) 226.—Texas.
tribuloides, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 145 = lagopodioides.
tricephala, *Boeck. in Flora*, lviii. (1875) 263.—Malaya.
triceps, *Michx. Fl. Bor. Am.* ii. 170.—Am. bor.
triceps, *Schrank, Baier. Fl.* i. 287.—Europ.
trichocarpa, *Muhl. ex Schkuhr, Riedgr.* ii. 47, 81.—Am. bor.
trichocarpa, Schur, Enum. Pl. Transs. 709 = asperrivens.
trichodes, *Steud. in Linnaea*, xxxix. (1875) 29.—Peruv.

CAREX:—
trichostachya, Fisch. ex Boott, Illustr. Carex, iv. 198 = poiocarpa.
trichostyles, *Franch. & Sav. Enum. Pl. Jap.* ii. 581. —Japon.
tricostata, Fries, Novit. Mant. iii. 152 = vulgaris.
trifida, *Cav. Ic.* v. 41. t. 465.—Reg. Magell.
trifera, Schkuhr, Riedgr. i. 94, ii. 66 = depauperata.
trifera, Vahl, ex Kunth, Enum. Pl. ii. 460 = vaginata.
trigona, All. Fl. Pedem. ii. 269 = fulva.
trinervis, *Degl. in Loisel. Fl. Gall.* ii. 781.—Gallia.
trinervis, Nees, in Wight, Contrib. 120 = Esenbeckii.
tripartita, All. Fl. Pedem. ii. 265 = curvula.
tripartita, Lam. Encyc. iii. 339 = divisa.
tripartita, Suter, Fl. Helv. ii. 241 = lagopina.
triquetra, *Boott, in Trans. Linn. Soc.* xx. (1851) 126. —Calif.
trisperma, *Dewey, in Am. Journ. Sc.* ix. (1825) 63.—Am. bor
trisperma, A. Gray, ex Boott, Illustr. Carex, ii. 81 = rosea.
tristachya, *Thunb. Fl. Jap.* 38.—Japon.
tristicha, *Boeck. in Flora*, xli. (1858) 651.—Am. bor.
tristicha, *Spruce, ex Boott, Illustr. Carex*, iv. 153.—Ecuador.
tristis, Bieb. Fl Taur. Cauc. iii. 615 = sempervirens.
tristicha, Wahlenb. ex Steud. Nom. ed. II. i. 299 = clavata.
truncata, *Boeck. in Flora*, xli. (1858) 649.—Am. bor.
truncata, *Fenzl, in Friederichsthal, Reise Griechenl.* 278.—Graecia.
truncatula, *Kit. in Linnaea*, xxxii. (1863) 818.—Hungar.
tuberculata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 270.—Mexic.
tuberculata, Nees, ex Boott, Illustr. Carex, ii. 76 = bonariensis.
tuberosa, Blanco, Fl. Filip. ed. I. 35 = Elecocharis tuberosa.
tuberosa, Degl. in Loisel. Fl. Gall. ii. 629 = Linkii.
tubulata, Schum. ex Boott, Illustr. Carex, iv. 136 = Buxbaumii.
Tuckermanni, *Boott, in Hook. Lond. Journ. Bot.* v. (1846) 78.—Am. bor.
tumida, Bellschmid, in Hornsch. Archiv Skand. ii. 214 = vulgaris.
tumida, *Boott, Illustr. Carex*, i. 66.—Reg. Himal.
tumidicarpa, *Anderss. in Bot. Notiser* (1849); *et Flora*, xxxiii. (1850) 297.—Scandinav.
turbinata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 265.—Mexic.
Turczaninowii, Steud. ex Boott, Illustr. Carex, iv. 161 = obesa.
turfacea, Auct. vide *torfacea*.
turfosa, Fries, Bot. Notiser (1846), 104, 107 = caespitosa.
turfosa, Fries, Summa, 228 = vulgaris.
turgescens, Dewey, in Am. Journ. Sc. Ser. II. iii. (1847) 386 = Halei.
turgescens, *Torr. in Ann. Lyc. N. York*, iii. (1836) 419. —Am. bor.
turkestanica, *Regel, in Act. Hort. Petrop.* vii. (1880) 670.—Turkestan.
Tweediana, *Nees, in Hook. Journ. Bot.* ii. (1840) 398. —Reg. Argent.
typhina, Michx. Fl. Bor. Am. ii. 169 = squarrosa.
typhinoides, Schwein. in Ann. Lyc. N. York, i. (1824) 66 = squarrosa.
typhoides, Bory, Voy. iii. 24 = Boryana.
uda, *Maxim. Prim. Fl. Amur.* 308.—Reg. Amur.
uetlisca, Suter, Fl. Helv. i. 251 = flava.
uliginosa, Linn. Sp. Pl. 978 = Schreus Caricis.
umbellata, *Schkuhr, Riedgr.* ii. 75.—Am. bor.
umbrosa, Host, Gram. i. 52. t. 69 = praecox.
uncifolia, *Cheesem. in Trans. N. Z. Inst.* xvi. (1884) 412.—N. Zel.
uncinata, Linn. f. Suppl. 418 = Uncinia australis.
uncinata, Schkuhr, ex Steud. Nom. ed. II. i. 297 = Uncinia jamaicensis.
uncinioides, *Boott, Illustr. Carex*, i. 6.—Reg. Himal.
undulata, Kunze, in Schkuhr, Riedgr. Suppl. 28. n. 8. t. 4. f. 2 = pallescens.
undulata, Schur, Enum. Pl. Transs. 714 = leucantha.
Urbani, *Boeck. in Engl. Bot. Jahrb.* vii. (1886) 280.—Alaska.
ursina, *Dewey, in Am. Journ. Sc.* xxvii. (1835) 240.—Am. arct.

CAREX:—
Urvillei, Bronga. in Duperr. Voy. Bot. 157 = pumila.
utriculata, *Boott, in Hook. Fl. Bor. Am.* ii. 221.—Am. bor.
vacillans, *Soland. ex Boott, in Hook. f. Fl. N. Zel.* i. 285.—N. Zel.
vacillans, Steud. Syn. Pl. Cyp. 235 = riparia.
vaginata, *Tausch, in Flora*, iv. (1821) 557.—Europ.
Vahlii, *Schkuhr, Riedgr.* i. 87.—Reg. bor. et arct.
valdiviana, *Phil. in Linnaea*, xxix. (1857-58) 84.—Chili.
valesiaca, Suter, Fl. Helv. 259 = tenuis.
valesiaca, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 156 = ambigua.
valida, *Nees, in Wight, Contrib.* 123.—China; Zeylan.
vallicola, *Dewey, in Am. Journ. Sc.* Ser. II. xxxii. (1861) 40.—Calif.
Vanheurckii, *Muell. Arg. in Heurck, Obs. Bot.* 30.—Kamtschatka.
varia, Host, Gram. i. 59. t. 80 = sempervirens.
varia, *Muhl. ex Schkuhr, Riedgr.* ii. 48.—Am. bor.
variegata, Lam. Encyc. iii. 339 = sempervirens.
variegata, *Scheele, in Linnaea*, xxiii. (1850) 565.—Am. bor.
Vasevi, Dewey, in Am. Journ. Sc. Ser. II. xxix. (1860) 347 = monile.
ventricosa, Curt. Fl. Lond. fasc. vi. t. 68 = depauperata.
venusta, *Dewey, in Am. Journ. Sc.* xxvi. (1834) 107.—Am. bor.
verna, Chaix & Vill. Hist. Pl. Dauph. i. 312, ii. 304 = praecox.
verna, Lam. Fl. Fr. ii. 185, partim = glauca.
verna, Schkuhr, Riedgr. i. 89 = obesa.
verna, Vill. ex Kunth, Enum. Pl. ii. 439 = praecox.
verrucosa, Muhl. Desc. Gram. Uber. 261 = glaucescens.
verrucosa, Schwein. ex Torr. in Ann. Lyc. N. York, iii. (1836) 400 = commutata.
versicolor, *Dum. Fl. Belg.* 148.—Belgium.
verticillata, *Boott, Illustr. Carex*, i. 67.—Am. bor.
verticillata, *Zoll. & Moritzi, Syst. Verz. Zoll.* 98.—Java.
vesicaria, Bunge, ex Boott, in Proc. Linn. Soc. i. (1845) 291.—Songorica.
vesicaria, Dewey, ex Steud. Nom. ed. II. i. 297 = bullata.
vesicaria, Laest, Fl. Herborn. t. 16. f. 2. l = riparia.
vesicaria, Lightf. Fl. Scot. ii. 566 = ampullacea.
vesicaria, *Linn. Sp. Pl.* 979.—Reg. bor. temp.
vesicaria, Thunb. Prod. Pl. Cap. 14 = clavata.
vesicifornis, *Peterm. in Flora*, xxvii. (1844) 339.—Saxonia.
vesiculosa, *Boott, Illustr. Caren*, iii. 107.—Ind. or.
vestita, *Willd. Sp. Pl.* iv. 243.—Am. bor.
vestita, Wormsk. ex Boott, Illustr. Carex, iv. 191 = Michelii.
vicinalis, *Boott, l. c.* 188.—Ind. or.
victoriensis, Nees, in Wight, Contrib. 126 = notha.
Villarsii, *Schkuhr, Riedgr.* ii. 5.—Gallia.
villosa, *Boott, in Perry, Jap. Exped.* ii. 327.—Japon.
villosa, Franch. & Sav. Enum. Pl. Jap. ii. 149 = Davidiana.
villosa, Stokes, Bot. Mat. Med. iv. 346 = hirta.
virens, Lam. Encyc. iii. 354 = muricata.
virens, Muhl. Fl. Par. i. 469 = acuta.
virescens, *Muhl. ex Schkuhr, Riedgr.* ii. 45.—Am. bor.
virescens, Willd. ex Kunth, Enum. Pl. ii. 430 = tricepa.
virgata, *Miq. Fl. Ind. Bat.* iii. 561.—Java.
virgata, Soland. ex Hook. f. Fl. N. Zel. i. 282 = paniculata.
virginiana, Woods, in Rees, Cycl. vi. n. 100 = acuta.
virginica, Steud. Syn. Pl. Cyp. 217 = prace.
viridis, Honck. Syn. Pl. Germ. i. 272 = flava.
viridis, *Petrie, in Trans. N. Z. Inst.* xiii. (1881) 332. —N. Zel.
viridis, *Schlecht. & Cham. in Linnaea*, vi. (1831) 80. —Mexico.
viridis, *Spenn. Fl. Friburg.* i. 48.—Europ.
viridula, *Franch. & Sav. Enum. Pl. Jap.* ii. 579.—Japon.
viridula, *Michx. Fl. Bor. Am.* ii. 170.—Am. bor.
viridula, Schwein. & Torr. in Ann. Lyc. N. York, i. (1824) 230 = tricepa.

CAREX:—
vitilis, Fries, Novit. Mant. iii. 137 = canescens.
Vlechii, Schwein. in Ann. Lyc. N. York, i. (1824) 69 = oligocarpa.
Vulcani, *Heckst. ex Soub. Fl. Asor.* 22. t. 8. f. 2.—Ins. Asoric.
vulgaris, *Fries, Nov. Mant.* iii. 138.—Reg. temp. bor. et austr.
vulpina, Carey, ex Dewey, in Am. Journ. Sc. Ser. II. viii. (1849) 348 = conjuncta.
vulpina, Hohen. in Bull. Soc. Nat. Mosc. (1838) 259 = muricata.
vulpina, *Linn. Sp. Pl.* 978.—Europ.; As. temp.
vulpina, Nees, in Hook. Journ. Bot. ii. (1840) 398 = bonariensis.
vulpinaeformis, Tuckerm. Enum. 8 = vulpinoidea.
vulpinaria, Nees, in Wight, Contrib. 121.—Reg Himal.; As. bor.
vulpinoidea, *Michx. Fl. Bor. Am.* ii. 169.—Reg Casc.; Am. bor.
vulpinoidea, Torr. in Ann. Lyc. N. York, i. (1824) 230 = stipata.
Wahlenbergiana, *Boott, Illustr. Carex*, ii. 101. t. 201. —Ins. Maurit.
wahuensis, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 218. t. 10.—Ins. Oahu; Japon.
Wakatipu, *Petrie, in Trans. N. Z. Inst.* xiv. (1882) 363.—N. Zel.
Walkeri, *Arn. ex Boott, in Proc. Linn. Soc.* i. (1845) 267.—Zeylan.
Wallichiana, *Presc. in Wall. Cat.* n. 3380.—Reg. Himal
Wallichiana, Spreng. Syst. iii. 812 = foliosa.
Washingtoniana, Dewey, in Am. Journ. Sc. x. (1826) 278 = vulgaris.
Watsoni, Boott, Illustr. Carex, iv. 208 = Vulcani.
Watsoni, *Olney, in Bot. King, Exp.* 870.—Am. bor. occ
Watsoniana, *Steud. Syn. Pl. Cyp.* 215.—Am. bor.
Waxmanni, *Schrank, Baier. Fl.* i. 297.—Europ.
Weiheana, Boeng. ex Kunth, Enum. Pl. ii. 439 = praecox?
Weiwitschi, *Boiss. ex Steud. Syn. Pl. Cyp.* 230.—Lusitan.
Whitneyi, *Olney, in Proc. Am. Acad.* vii. (1868) 394. —Calif.
Wichurai, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 519. —China.
Wightiana, *Nees, in Wight, Contrib.* 122.—Ind. or.
Wilkesii, *Olney, in Bot. U. St. Expl. Exped.* ii. 477. t. 17.—Hab.?
Willdenowii, *Schkuhr, Riedgr.* ii. 83.—Am. bor.
willisia, *Meinsh. ex Maach, Vilyuisk. Obruge Yakutsk. Obl.* 11. 808.—Sibir.
Wimmeri, *Steud. Syn. Pl. Cyp.* 215.—Silesia.
Winkleri, *Lange, in Kjoeb. Vidensh. Meddel.* (1877-78) 234.—Hispan.
Witheringii, S. F. Gray, Nat. Arr. Brit. Pl. ii. 48 = arenaria.
Wormskioldiana, Hornem. in Fl. Dan. t. 1536 = scirpoides.
Wormskioldii, Drej. Revis. Caric. 18 = scirpoides.
Woodii, Dewey, in Am. Journ. Sc. Ser. II. ii. (1846) 249 = tetanica.
Wrightii, *Dewey, ex Torr. Bot. Mex. Bound.* 285.—Am. bor. occ.
xalapensis, Kunth, Enum. Pl. ii. 380 = Brongniartii.
xanthophysa, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 153 = folliculata.
xerocarpa, *S. H. Wright, in Am. Journ. Sc.* Ser. II. xiii. (1866) 384, in nota.—Am. bor.
yodoënsis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 515. —Japon.
Zollingeri, Kunze, ex Steud. Syn. Pl. Cyp. 221 = nemostachya.
zonata, F. Nyland. Spicil. Pl. Fenn. ii. 19 = vulgaris?

CAREYA, Roxb. Hort. Beng. 52 (1814); Fl. Ind. ii. 636 (1832). *MYRTACEÆ*, Benth. & Hook. f. i. 752.
CAREYA, Wight & Arn. Prod. i. 324 (1834).
CUMBIA, Buch.-Ham. Mysore, iii. 187 (1807).
arborea, *Roxb. l. cc.*—Ind. or.
australis, F. Muell. Fragm. v. 188 = Barringtonia Careya.
herbacea, *Roxb. Hort. Beng.* 52; *Fl. Ind.* ii. 638.—Ind. or.
macrostachya, Jack, in Malay Misc. i. n. v. 67 = Barringtonia macrostachya.

CAREYA :—
orbiculata, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 98 = arbores.
pendula, Griff. Notul. iv. 661 = Barringtonia macrostachya.
sphærica, *Roxb. Hort. Beng.* 52; *Fl. Ind.* ii. 636.—Ind. or.
valida, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 72.—Ins. Andam.

CARGILLA, Adans. Fam. ii. 180 (1763) = Chrysogonum, Linn. (Compos.).

CARGILLIA, R. Br. Prod. 526 (1810) = Diospyros, Lam. (Ebenac.).
arborea, A. Cunn. ex Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 289 = D. pentamera.
australis, R. Br. Prod. 526 = D. Cargillia.
lanx, R. Br. l. c. = D. maritima.
mabacea, F. Muell. Fragm. v. 162 = D. mabacea.
maritima, Hassk. Cat. Hort. Bog. Alt. 159 = D maritima.
megalocarpa, F. Muell. Fragm. v. 163 = D. maritima.
pentamera, F. Muell. l. c. iv. 82 = D. pentamera.

CARICA, Linn. Gen. ed. I. 309 (1737). *PASSIFLOREAE*, Benth. & Hook. f. i. 815.
MOCINNA, Cerv. ex La Llave, in La Naturaleza, vii. (1885) Apénd. 76.
PAPAYA, Tourn. ex Linn. Syst. ed. I (1735).
VASCONCELLEA, A. St. Hil. Mém. Reséd. ii. 13 (1837).
Boissieri, *Hemsl. Biol. Centr.-Am. Bot.* i. 481.—Mexic.
candamarcensis, *Hook. f. Bot. Mag.* t. 6198.—Ecuador.
candicans, *A. Gray, Bot. U. St. Expl. Exped.* i. 640. —Peruv.
caulıflora, *Yacq. Hort. Schœnb.* iii. 33.—Venezuela.
citriformis, *Yacq. f. ex Spreng. Syst.* iii. 905.—Afr. trop.
cundinamarcensis, J. Linden, Cat. n. 87 (1871) = candamarcensis.
digitata, Spreng. Syst. iii. 906 = spinosa.
dodecaphylla, *Vell. Fl. Flum.* x. t. 132.—Bras.
erythrocarpa, *Linden & André, Illustr. Hortic.* xviii. (1871) t. 51.—Ecuador.
glandulosa, *Pav. ex A. DC. in DC. Prod.* xv. i. 417.— Peruv.
gossypiifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 187. —Reg. Argent.
heptaphylla, Vell. Fl. Flum. x. t. 133 = spinosa.
hermaphrodita, Blanco, Fl. Filip. ed. I. 205 = C. Papaya.
heterophylla, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 60. —N. Granat.
Mamaya, *Vell. Fl. Flum.* x. t. 131.—Bras.
microcarpa, *Yacq. Schœnb.* iii. 32.—Am. austr.
monoica, *Desf. in Ann. Mus. Par.* i. (1802) 278. t. 18.—Peruv.
nana, *Benth. Pl. Hartw.* 288.—Mexic.
Papaya, *Linn. Sp. Pl.* 1036.—Am. trop.
pelıata, *Hook. & Arn. Bot. Beech. Voy.* 425. t. 98.— Nicaragua.
Posopora, *Linn. Sp. Pl.* 1036.—Peruv.
pubescens, *Lemsl & C. Koch. in Ind. Sem. Hort. Bot. Berol.* (1854) 12.—Guatemala.
pyriformis, Willd. Sp. Pl. iv. 815 = C. Posopora.
scandens, *A. Gray, Bot. U. St. Expl. Exped.* i. 640.— Peruv.
spinosa, *Aubl. Pl. Gui.* ii. 908.—Guian.; Bras.

CARICELLA, Ehrh. Beitr. iv. 146 (1789) = Carex, Dill.

CARICTERIA, Soop. Introd. 104 (1777) = Corchorus, Tourn. (Tiliac.).

CARIGOLA, Rafin. Fl. Tellur. ii. 10 (1836) = Monochoria, Presl (Pontederac.).
hastata, Rafin. l. c. = M. hastæfolia.

CARIMA, Rafin. Fl. Tellur. iv. 62 (1836) = Adhatoda, Nees (Acanthac.).
sulcata, Rafin. l. c. = A. sulcata.

CARINIANA, Casar. Nov. Stirp. Bras. 35 (1842) = Couratari, Aubl. (Myrtac.).
brasiliensis, Casar. l. c. = Couratari legalis.

CARINIANA :—
domestica, Miers, in Trans. Linn. Soc. xxx. (1874) 286 = Couratari domestica.
excelsa, Casar. Nov. Stirp. Bras. 46 = Couratari estrellensis.
exigua, Miers, in Trans. Linn. Soc. xxx. (1874) 289.— Ind. occ.
pyriformis, Miers, l. c. 290.—Bolivia.
rubra, Miers, l. c. 288 = Couratari rubra.
sahupensis, Miers, l. c. = Couratari naupensis.

CARIONIA, Nand. in Ann. Sc. Nat. Sér. III. xv. (1851) 311. t. 15. *MELASTOMACEAE*, Benth. & Hook. f. i. 769.
elegans, *Nand. l. c.* 312.—Ins. Philipp.
triplinervia, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 310.—Ins. Philipp.

CARISSA, Linn. Mant. i. 7 (1767). *APOCYNACEAE*, Benth. & Hook. f. ii. 695.
ARTUA, Forsk. Fl. Aegypt. Arab. 63 (1775).
ARDUINA, Mill. Ic. t. 200 (1759); Linn. Mant 7 (1767).
abyssinica, *R. Br. in Salt, Abyss. App.* p. lxiv, nomen. —Abyss.
acuminata, *A. DC. Prod.* viii. 335.—Afr. austr.
africana, *A. DC. l. c.* 332.—Afr. trop.
Arduina, *Linn. Encyc.* i. 555.—Afr. austr.
axillaris, *Roxb. Hort. Beng.* 34; *Fl. Ind.* i. 691.—Ins. Molucc.
bispinosa, Desf. Tabl. ed. I. 78 = C. Arduina.
Brownii, F. Muell. Fragm. iv. 45 = lanceolata, ovata, et scabra.
Candollensis, *Jaub. & Spach, Illustr.* v. 107. t. 497.— Abyssin.
Carandas, *Linn. Mant.* 52.—Ind. or.; Malaya.
Carandas, Lodd. Bot. Cab. t. 668 = paucinervia.
Carandas, Lour. Fl. Cochinch. 155 = africana.
congesta, Wight, Ic. iv. t. 1289 = C. Carandas.
coriacea, *Wall. Cat.* n. 1680.—Ind. or.
cornifolia, *Jaub. & Spach, Illustr.* v. 108. i. 498.— Abyssin.
cryptophylla, *Baker, in Journ. Linn. Soc.* xx. (1884) 204.—Madag.
Dalzellii, Bedd. Forest. Fl. Anal. Gen. 156; Man. Bot. 157 = macrophylla.
densiflora, *Baker, in Journ. Linn. Soc.* xx. (1884) 204. —Madag.
diffusa, Roxb. Hort. Beng. 19; Fl. Ind. i. 689 = spinarum.
diffusa, Wall. Cat. n. 1678, partim = paucinervia.
dulcis, Schum. & Thonn. Beskr. Guin. Pl. 146 = edulis.
edulis, *Vahl, Symb. Bot.* i. 23.—Aegypt.
erythrocarpa, *A. DC. Prod.* viii. 335.—Afr. austr.
ferox, *A. DC. l. c.*—Afr. austr.
grandiflora, *A. DC. l. c.*—Afr. austr.
grandis, *Sert. ex A. DC. l. c.* 336.—Ins. Tahiti.
haematocarpa, *A. DC. l. c.*—Afr. austr.
hirsuta, Roth, Nov. Pl. Sp. 128 = spinarum.
inermis, Vahl, Symb. Bot. iii. 43 = macrophylla.
lanceolata, *R. Br. Prod.* 468.—Austral.
lanceolata, Dals. in Dalz. & Gibs. Bomb. Fl. 143 = macrophylla.
latifolia, *Benth. Fl. Austral.* iv. 305.—Austral.
macrocarpa, *A. DC. Prod.* viii. 336.—Afr. austr.
macrophylla, *Wall. Cat* n. 1679.—Ind. or.
madagascariensis, *Thou. Obs. Pl. Iles Afr. Austr.* 80. —Madag.
mitis, Heynh. ex A. DC. Prod. viii. 332 = macrophylla.
mitis, Vahl, Symb. Bot. iii. 44 = Holarrhena mitis.
myrtoides, *Desf. Cat. Hort. Par.* ed. III. 398.—Hab. ?
oblongifolia, *Hochst. in Flora*, xxvii. (1844) ii. 827.— Afr. austr.
ovata, *R. Br. Prod.* 468.—Austral.
paucinervia, *A. DC. Prod.* viii. 338.—Ind. or.
pubescens, *A. DC. l. c.* 334.—Afr. trop.
Richardiana, Jaub. & Spach, Illustr. v. 106 t. 496 = edulis.
salicina, Lam. Encyc. i. 554 = spinarum ?
scabra, *R. Br. Prod.* 468.—Austral.
Schimperi, *A. DC. Prod.* viii. 675.—Abyssin.
sechellensis, *Baker, Fl. Maurit.* 222.—Ins. Seychell.
spinarum, *Linn. Mant.* t. 356.—Ins. Molucc.
spinarum, Lodd. ex A. DC. Prod. viii. 336 = Damnacanthus indicus.
suavissima, *Bedd. ex Hook. f. Fl. Brit. Ind.* iii. 683.— Ind. or.

CARISSA :—
tomentosa, *A. Rich. Tent. Fl. Abyss.* ii. 30.—Abyssin.
villosa, Roxb. Hort. Beng. 19; Fl. Ind. i. 690 = spinarum.
Xylopicron, *Thou. Obs. Pl. Iles Afr. Austr.* 80. t. 2. — Ins. Madag.

CARLEA, Presl, Epim. Bot. 216 (1849) = Symplocos, Linn. (Styrac.).
oblongifolia, Presl, l. c. 217 = S. oblongifolia.

CARLEMANNIA, Benth. in Hook. Kew Journ. v. (1853) 308. *RUBIACEAE*, Benth. & Hook. f. ii. 63.
congesta, *Hook. f. Fl. Brit. Ind.* iii. 85.—Reg. Himal.
Griffithii, *Benth. in Hook. Kew Journ.* v. (1853) 308 —Reg. Himal.
tetragona, *Hook. f. Fl. Brit. Ind.* iii. 85.—Reg. Himal.

CARLINA, Linn. Syst. ed. I (1735). *COMPOSITAE*. Benth. & Hook. f. ii. 465.
ATHAMUS, Neck. Elem. i. 85 (1790).
CARLOWIZIA, Moench, Meth. Suppl. 225 (1802).
CHROMATOLEPIS, Dulac, Fl. Hautes-Pyr. 596 (1867).
MITINA, Adans. Fam. ii. 116 (1763).
acanthifolia, *All. Fl. Pedem.* i. 156. t. 81.—Reg. Mediterr.
Acarna, Bieb. Fl. Taur. Cauc. ii. 282 = Cnicus Acarna.
acaulis, Lam. Encyc. i. 623 = acanthifolia
acaulis, *Linn. Sp. Pl.* 828.—Europ.
acaulis, Stokes, Bot. Mat. Med. iv. 156 = Cnicus acaulis.
aculeata, Burm. f. Fl. Cap. Prod. 28 (= *Stobæa rigida*) —Afr. austr.
aggregata, Willd. Sp. Pl. iii. 1694 = acaulis.
alpina, Jacq. Enum. Stirp. Vindob. 274 = acaulis.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 19.—Algeria.
atractyloides, Linn. Sp. Pl. 1161 (= *Stobæa atractyloides*).—Afr. austr.
Bieberstenii, Benth. ex Hornem Hort. Hafn. Suppl. 94 = vulgaris.
bracteata, J. & C. Presl, Delic. Prag. 103 = sicula.
caulescens, Gilib. Fl. Lituan. iii. 135 = vulgaris.
caulescens, Lam. Fl. Fr. ii. 7 = acaulis.
Chamæleon, Vill. Hist. Fl. Dauph. iii. 80 = acaulis.
Charlousse, Vill. Prosp. 27 = acanthifolia.
complanata, Schur, Enum. Pl. Transs. 418 = vulgaris.
corymbosa, *Linn. Sp. Pl.* 828.—Reg. Mediterr.
corymbosa, Scop. ex Steud. Nom. ed. II. i. 299 = vulgaris.
Curetum, Heldr. in Boiss. Fl. Orient. iii. 449 = corymbosa.
Cynara, Pourr. ex DC. Fl. Fr. iv. 128 = acanthifolia.
discolor, Soleir. ex Nym. Consp. 400 = macrocephala.
Echinus, Bieb. Fl. Taur. Cauc. iii. 288.—Reg. Cauc (= *Cirsium scleranthum*).
elatior, Wallr. in Linnæa, xiv. (1840) 636; et Beitr. Fl. Hercyn. 266 = acaulis.
foliosa, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 450 = oligocephala.
Fontanesii, DC. Prod. vi. 548 = Atractylis macrocephala.
frigida, *Boiss. & Heldr Diagn.* Ser. I. vi. 109.— Græcia.
gorterioides, Lam. Encyc. i. 625 (= Berkheuya sp.).— Afr. austr.
graeca, Heldr. & Sart. ex Boiss. Fl. Orient. iii. 449 = corymbosa.
grandiflora, Moench, Meth. 558 = acaulis.
gummifera, Less. Syn. Comp. 12 = Atractylis gummifera.
hispanica, Lam. Encyc. i. 624 = corymbosa.
intermedia, Schur, Enum. Pl. Transs. 418 = vulgaris.
involucrata, *Poir. Voy. Barb.* ii. 234.—Reg. Mediterr.
lanata, Habl. Taur. 160 = vulgaris.
lanata, *Linn. Sp. Pl.* 828.—Reg. Mediterr.
lanata, Pall. ex DC. Prod. vi. 546 = vulgaris.
libanotica, Boiss. Diagn. Ser. I. x. 95 = corymbosa.
Lobeliana, Tausch, in Flora, xii. (1829) Erg. 41 = corymbosa.
longifolia, Reichb. Ic. Fl. Germ. viii. 25 = vulgaris.
longifolia, Vis. Fl. Lib. Spec. 88 = corymbosa.
lyrata, *Thunb. Prod. Pl. Cap.* 141.—Afr. austr.
macrocephala, Less. Syn. Comp. 12 = Atractylis macrocephala.

CARLINA :—
macrocephala, *Moris, Elench. Sard.* ii. 5.—Reg. Mediterr.
macrophylla, DC. Prod. vi. 648 = Atractylis macrophylla.
nebrodensis, *Guss. in DC. Prod.* vi. 646.—Sicilia.
nebrodensis, Koch, ex Nym. Consp. 401 = vulgaris.
oligocephala, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 51. —As. Min.; Syria.
onopordifolia, Bess. ex DC. Prod. vi. 648 = acanthifolia.
orophila, Lamotte, Prod. ii. 448 = vulgaris.
patula, Burm. f. Fl. Cap. Prod. 23 (= *Stobaeae sp.*).—Afr. austr.
Pola, Hacq. Pl. Alp. Carn. 29. t. 4. f. 3 = lanata.
pyrenaica, Linn. Sp. Pl. 829 = Carduus carlinoides.
racemosa, Gilib. Chlor. Grodn. 34 = vulgaris.
racemosa, Gouan, Illustr. 64 = corymbosa.
racemosa, *Linn. Sp. Pl.* 829.—Reg. Mediterr. occ.
radiata, Viv. Fl. Lib. Spec. 68 = corymbosa.
Reboudiana, *Pomel, Nouv. Mat. Fl. Atl.* 278.—Algeria.
Rothii, Heldr. & Sart. ex Boiss. Fl. Orient. iii. 449 = corymbosa.
salicifolia, *Cav. in Anal. Cienc. Nat.* iv. (1801) 81; *Lasz. in Linnaea*, vi. (1831) 88.—Ins. Madera.
scabra, Burm. f. Fl. Cap. Prod. 23 = Tripteris arborescens.
sicula, *Tenore, Cat. Hort. Neap. App.* (1819).—Sicilia.
simplex, Waldst. & Kit. Pl. Rar. Hung. ii. 164. t. 152 = acaulis.
subacaulis, DC. Fl. Fr. iv. 122 = acaulis.
sulphurea, Desf. Fl. Atlant. ii. 251 = racemosa.
tragacanthifolia, *Klatt, in Leopoldina*, xx. (1884) 94.—Ins. Rhodus.
transsilvanica, Schur, in Verh. Siebenb. Ver. Nat. iv. (1853) 34 = acanthifolia.
tristis, Salisb. Prod. 185 = vulgaris.
Utzka, Hacq. Pl. Alp. Carn. 7. t. 1 = acanthifolia.
vulgaris, *Linn. Sp. Pl.* 828.—Europ.; As. bor.
xeranthemoides, *Linn. f. Suppl.* 349.—Ins. Teneriff.

CARLOTEA, Arruda, ex Koster, Trav. 493 (1816) (Gen. ignot.).
formosissima, Arruda, l. c.—Pernambuc.
speciosa, Arruda, l. c.—Pernambuc.

CARLOWIZIA, Moench, Meth. Suppl. 225 (1802) = Carlina, Linn. (Compos.).
corymbosa, Cass. in Dict. Sc. Nat. xxv. 53 = Carlina salicifolia.
salicifolia, Moench, Meth. Suppl. 226 = Carlina salicifolia.

CARLOWRIGHTIA, A. Gray, in Proc. Am. Acad. xiii. (1878) 364. *ACANTHACEAE.*
arizonica, *A. Gray, l. c.*—Arizona.
linearifolia, *A. Gray, l. c.*—Texas.

CARLUDOVICA, Ruiz & Pav Prod. 146. t. 31 (1794). *CYCLANTHACEAE*, Benth. & Hook. f. iii. 958.
EVODIANTHUS, Oerst. in Kjoeb. Videnak. Meddel. (1875) 194.
LUDOVIA, Pers. Syn. ii. 576 (1807).
SALMIA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 399.
SARCINANTHUS, Oerst. in Kjoeb. Videnak. Meddel. (1875) 196.
acuminata, Poepp. & Endl. Nov. Gen. et Sp. ii. 37 = divergens.
acuminata, *Ruiz & Pav. Syst.* 293.—Am. austr.
angustifolia, Ruiz & Pav. l. c. 292 = Cyclanthus Plumerii.
strovirens, *H. Wendl. Ind. Palm.* (1854) 67.—N. Granat.
brachypus, *Drude, in Mart. Fl. Bras.* iii. 11. 287.—Bras.
chelidonura, *Drude, l. c.* 288.—Bras.
disticha, *Post. in Rev. Hortic. Ser.* III. i. (1847) 86.—Guiana.
divergens, *Drude, in Mart. Fl. Bras.* iii. 11. 241.—Bras.
Drudei, *Mast. in Gard. Chron.* (1877) ii. 714.—Bras.
ensiformis, *Hook. f. Bot. Mag.* t. 6418.—Costa Rica.
funifera, *Kunth, Enum. Pl.* iii. 106.—Guiana.
Gardneri, *Hook. Journ. Bot.* ii. (1840) 29. t. 8, 4.—Bras.
heterophylla, *Mart. ex Drude, in Mart. Fl. Bras.* iii. 11. 241.—Bras.

CARLUDOVICA :—
humilis, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 37.—Peruv.
imperialis, *Linden & André, Illustr. Hortic.* xxi. (1874) t. 165.—Ecuador.
incisa, *H. Wendl. Ind. Palm.* (1854) 67.—Am. centr.
insignis, *Duchass. ex Griseb. Fl. Brit. W. Ind.* 513.—Ind. occ.
Kegeliana, *Lem. Illustr. Hortic.* ii. (1855) *Misc.* 23.—Guiana.
latifolia, *Ruiz & Pav. Syst.* 292.—Ind. occ.
latifrons, *Drude, in Mart. Fl. Bras.* iii. 11. 287.—Bras.
longicrura, *Drude, l. c.* 240.—Bras.
macropoda, *Klotzsch, in Linnaea*, xx. (1847) 469.—N. Granat.
microphylla, *Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 197.—Costa Rica.
Moritziana, *Klotzsch, in Linnaea*, xx. (1847) 468.—N. Granat.
Oerstedii, *Hemsl. Biol. Centr. Am. Bot.* iii. 416.—Costa Rica.
palmata, Griseb. in Goett. Abh. vii. (1857) 259 = insignis.
palmata, *Ruiz & Pav. Syst.* 291.—Am. austr.
pliesta, *Klotzsch, in Linnaea*, xx. (1847) 468.—N. Granat.
plicata. Linden, ex H. Wendl. Ind. Palm. (1854) 67 = sarmentosa.
Plumerii, *Kunth, Enum. Pl.* iii. 106.—Ins. Martinic.
Plumieri, Griseb. Fl. Brit. W. Ind. 513 = gracilis.
purpurascens, *J. Linden*, Illustr. Hortic. xxviii. (1881) 64.—N. Granat.
rupestris, *Klotzsch, in Linnaea*, xx. (1847) 469.—N. Granat.
sarmentosa, *Sagot, ex Drude, in Mart. Fl. Bras.* iii. 11. 239.—Guiana.
Sartoni, *Teijsm. & Binn. Cat. Hort. Bog.* 381, nomen. —Hab. ?
schizophylla, *Drude, in Mart. Fl. Bras.* iii. 11. 235.—Bras.
speciosa, Linden.—Cf. Gard. Chron. (1877) ii. 753 Drudei.
subacaulis, *Kunth, Enum. Pl.* iii. 107.—Guiana.
tetragona, *H. B. & K. Nov. Gen. et Sp.* i. 79.—N. Granat.
tetragonopus, *Mart. ex Drude, in Mart. Fl. Bras.* iii. 11. 238.—Bras.
Trailiana, *Drude, l. c.* 241.—Bras.
trigona, *Ruiz & Pav. Syst.* 293.—Am. austr.
utilis, *Benth. & Hook. f. ex Hemsl. Biol. Am. Centr. Bot.* iii. 416.—Costa Rica.
Wallisi, *Regel, Gartenfl.* (1879) 325.—N. Granat.

CARMELITA, C. Gay, ex DC. Prod. vii. 14 (1838) = Substantiva, Ruiz & Pav. (Compos.).
formosa, C. Gay, l. c. 15 = Chaet. villosa.

CARMENTA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

CARMICHAELIA, Reichb. Nom. 147 (1841) = seq.

CARMICHAELIA, R. Br. in Bot. Reg. xi. t. 912 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 502.
australis, *R. Br. l. c.*—N. Zel.
compacta, *Petrie, in Trans. N. Z. Inst.* xvii. (1885) 272.—N. Zel.
corrugata, *Colenso, l. c. xv.* (1882) 230.—N. Zel.
crassicaulis, *Hook. f. Handb. New Zeal. Fl.* 48.—N. Zel.
Cunninghamii, *Raoul, Choix*, 39.—N. Zel.
Enysii, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 279. —N. Zel.
exsul, *F. Muell. Fragm.* vii. 126.—Ins. Dom. Howe.
flagelliformis, *Colenso, in Hook. f. Fl. N. Zel.* i. 51.—N. Zel.
gracilis, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 336. —N. Zel.
grandiflora, *Hook. f. Handb. New Zeal. Fl.* 49.—N. Zel.
juncea, *Colenso, ex Hook. f. Fl. N. Zel.* i. 51.—N. Zel.
Kirkii, *Hook. f. Ic. Pl.* t. 1322.—N. Zel.
Munroi, *Hook. f. Handb. New Zeal. Fl.* 49.—N. Zel.
nana, *Colenso, ex Hook. f. l. c.*—N. Zel.
odorata, *Colenso, ex Hook. f. Fl. N. Zel.* i. 50.—N. Zel.
pilosa, *Colenso, ex Hook. f. l. c.*—N. Zel.
stricta, *Lehm. in Ind. Hort. Bot. Hamb.* (1852) 8. —*Cf. Hamb. Gartenz.* viii. (1852) 378.—Hab. ?

CARMICHAELIA :—
uniflora, *F. Buch. in Trans. N. Z. Inst.* xvi. (1884) 294.—N. Zel.
uniflora, *T. Kirk, in Trans. N. Z. Inst.* xi. (1884) 1. 512. *Trans. N. Z. Inst.* xvi. (1884) 279.—N. Zel.
Williamsi, *T. Kirk, in Trans. N. Z. Inst.* xii. (1880) 294.—N. Zel.

CARMINATIA, Mog. ex DC. Prod. vii. 267 (1838) *COMPOSITAE*, Benth. & Hook. f. ii. 343.
tenuiflora, *DC. l. c.*; *et in Delessz. Ic. Sel.* iv. t. 98.—Mexic.
tenuifolia, Steud. Nom. ed. II. i. 299, sphalm. = prsec.

CARMONA, Cav. Ic. v. 22. t. 488 (1799) = Ehretia, Linn. (Borag.).
heterophylla, Cav. l. c. 23 = E. buxifolia.
lycioides, G. Don, Gen. Syst. iv. 891 = E. lycioides.
microphylla, G. Don, l. c. = E. buxifolia.
viminea, G. Don, l. c. = Rhadia lycioides.

CARMONEA, Pers. Syn. i. 290 (1805) = prsec.

CARMOREA, Steud. Nom. ed. II. i. 299 (1840) sphalm. = prsec.

CARNARVONIA, F. Muell. Fragm. vi. 80. t. 55, 56 (1867). *PROTEACEAE*, Benth. & Hook. f. iii. 182.
araliaefolia, *F. Muell. l. c.* 81.—Austral.

CAROLI-GMELINA, Gaertn. Mey. & Scherb. Fl Wett ii. 419 (1800) = Nasturtium, R. Br. (Crucif.).
lanceifolia, Gaertn. Mey. & Scherb. l. c. 468 = N. amphibium.
palustris, Gaertn. Mey. & Scherb. l. c. 470 = N. amphibium.
sylvestris, Gaertn. Mey. & Scherb. l. c. 470 = N. sylvestre.

CAROLINEA, Linn. f. Suppl. 51 (1781) = Pachira, Aubl. (Malvac.).
affinis, Mart. Nov. Gen. et Sp. i. 85 = P. affinis.
alba, Lodd. Bot. Cab. t. 732 = P. alba.
armata, G. Don, Gen. Syst. i. 510 = P. armata.
campestris, Mart. Nov. Gen. et Sp. i. 86 = P. campestris.
emarginata, D. Dietr. Syn. Pl. iv. 861 = P. emarginata.
fastuosa, Sesse, ex DC. Prod. i. 478 = P. fastuosa.
grandiflora, Spach, Hist. Veg. Phan. iii. 498; xiv. 206 = P. grandiflora.
insignis, Sw. Prod. Veg. Ind. Occ. 101 = P. insignis.
longiflora, Mart. Nov. Gen. et Sp. i. 86 = P. longiflora.
macrantha, G. Don, Gen. Syst. i. 510 = P. macrantha
macrocarpa, Cham. & Schlecht. in Linnaea, vi. (1831) 628 = P. macrocarpa.
marginata, G. Don, Gen. Syst. i. 511 = P. marginata
minor, Sims, Bot. Mag. t. 1412 = P. minor.
pompifera, Mog. & Sesse, ex DC. Prod. i. 478 = P. minor
Princeps, Linn. f. Suppl. 314 = P. aquatica.
robusta, Lodd. ex Steud. Nom. ed. II. i. 299.—Am. austr.
tomentosa, Mart. Nov. Gen. et Sp. i. 84 = P. tomentosa.
villosa, Mart. ex Steud. Nom. ed. II. i. 300 = P. tomentosa.

CAROMBA, Steud. Nom. ed. II. i. 300 (1840) sphalm. = Cabomba, Aubl. (Nymph.).

CAROPYXIS, Benth. & Hook. f. Gen. i. 668 (1865) sphalm. = CALOPYXIS, Tul. = Combretum, Linn.

CAROSELINUM, Griseb. Spicil. Fl. Rumel. i. 374 (1843) = Petroedanum, Linn. (Umbellif.).
Chabraei, Griseb. l. c. = P. carvifolium.
dioicum, Griseb. l. c. = Johrenia graeca.

CAROTA, Reppr. Sert. Tiansch. 49; in Mém. Acad. Pétersb. Sér. VII. xiv. (1869) = Daucus, Tourn. (Umbellif.).
asclica, Reppr. Fl. Ingric. 468 = D. Carota.
sylvestris, Reppr. Sert. Tiansch. 49 = D. Carota.

CAROXYLON, Thunb. Nov. Gen. & Sp. 27 (1782) = Salsola, Linn. (Chenopod.).
acutiflorum, Mog. in DC. Prod. xiii. 172 = S. Arnicola.

CAROXYLON :—
articulatum, Moq. in DC. Prod. xiii. 11. 175 = Haloxylon articulatum.
Auricula, Moq. l. c. 172 = S. Auricula.
Bottæ, Moq. in DC. l. c. 178.—Afr. austr.
brevifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = S. aphylla.
divaricatum, Moq. in DC. Prod. xiii. 11. 177 = S. longifolia.
foetidum, Moq. l. c. 178 = S. foetida.
glaucum, Moq. l. c. 178 = S. glauca.
Griffithii, Moq. l. c. 175 = Haloxylon Griffithii.
hispidulum, Bunge, Lehm. Reliq. Bot. 293 = S. hispidula.
imbricatum, Moq. in DC. Prod. xiii. 11. 177 = S. gemmascens.
indicum, Wight, Ic. t. 1794 = Haloxylon recurvum.
lanciifolium, Brisa. Diagn. Ser. I. xii. 98 = S. lanciifolia.
nodulosum, Moq. in DC. Prod. xiii. 11. 177 = S. gemmascens.
recurvum, Moq. l. c. 175 = Haloxylon recurvum.
salicornicum, Moq. l. c. 174 = Haloxylon salicornicum.
Salsola, Thunb. Dias. Med. Afric. 8 = S. aphylla.
subaphyllum, Moq. in DC. Prod. xiii. 11. 178 = S. subaphylla.
tamariscifolium, Moq. l. c. 174 = S. genistoides.
tamariscinum, Moq. l. c. = S. tamariscina.
tetragonum, Moq. l. c. 176 = S. tetragona.
tuberculatum, Moq. l. c. 178 = S. tuberculata.
verruconum, Moq. l. c. = S. tuberculata.
Zeyheri, Moq. l. c. 176.—Afr. austr.

CARPACOCE, Sond. in Harv. & Sond. Fl. Cap. iii. 92 (1864–65). *RUBIACEÆ*, Benth. & Hook. f. ii. 141.
LAGOTIS, E. Mey. in Drège, Zwei Pfl. Docum. 197 (1843).
scabra, *Sond. l. c.* 33.—Afr. austr.
Spermacoce, *Sond. l. c.*—Afr. austr.

CARPENTARIA, Beec. in Ann. Jard. Buitenz. ii. (1856) 138. *PALMÆ*.
acuminata, *Becc. l. c.*—Austral.

CARPENTERIA, Torr. in Smithson. Contrib. vi. 12. t. 7 (1854). *SAXIFRAGEAE*, Benth. & Hook. f. i. 643.
californica, *Torr. l. c.*—Calif.

CARPENTIERA, Steud. Nom. ed. II. i. 344 in syn. (1840) sphalm. = **Charpentiera**, Gaudich. (Annarant.).

CARPESIUM, Linn. in Act. Soc. Upsal. (1741) 80. *COMPOSITÆ*, Benth. & Hook. f. ii. 336.
CONYZOIDES, Tourn. ex DC. Prod. vi. 581 (1837).
abrotanoides. *Linn. Sp. Pl.* 860.—Europ.; As. temp. cernuum, *Linn. l. c.* 859.—Europ.; As. temp.
ciliatum, Wall. Cat. n. 3214 = cernuum.
divaricatum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 187.—Japon.
glossophyllum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 479.—Japon.
macrocephalum, *Franch. & Sav. Enum. Pl. Jap.* ii. 406.—Japon.
nepalense, Less. in Linnæa, vi. (1831) 234 = cernuum
pedunculosum, Wall. Cat. n. 3200 = cernuum.
pubescens, Wall. l. c. n. 3199 = cernuum.
racemosum, Wall. l. c. n. 3201 = abrotanoides.
rosulatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 179.—Japon.
scandens, Schult. f. ex Miq. l. c. 173 = Rhynchospermum verticillatum.
Thunbergianum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 187.—Japon.
trachelifolium, *Less. in Linnæa*, vi. (1831) 233.—Reg. Himal.
triste, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 479.—Japon.
Wulfenianum, Schrad. ex DC. Prod. vi. 282 = abrotanoides.

CARPEZIUM, Gronn, Fl. Monsp. 465 (1765) = prec.

CARPHA, Banks & Soland. ex R. Br. Prod. 280 (1810), pro parte. *CYPERACEÆ*, Benth. & Hook. f. iii. 1061.
alpina, *R. Br. Prod.* 280.—Austral. ; N. Zel. ; Patag.
andina, Phil. in Anal. Univ. Chil. (1873) 555 = paniculata.
arundinacea, Brongn. in Duperr. Voy. Bot. 169. t. 30 = Schoenus arundinaceus.
Auberti, Nees, in Linnæa, ix. (1834) 300 = Cyclocampe Auberti.
avenacea, R. Br. Prod. 280 = Cyathochaete avenacea.
capensis, Martens, ex Steud. Nom. ed. II. i. 300 = Ecklonia capensis.
capitellata, *Boeck. in Linnæa*, xxxviii. (1874) 366.—Afr. austr.
clandestina, R. Br. Prod. 281 = Cyathochaete clandestina.
deusta, R. Br. l. c. 280 = Mesomolaena deusta.
diandra, R. Br. l. c. 281 = Cyathochaete diandra.
elongata, Boeck. l. c. xxxvii. (1874) 273 = Asterochaeta elongata.
glomerata, Nees, l. c. vii. (1832) 529 = Asterochaete glomerata.
hexandra, Nees, l. c. ix. (1834) 300 ; x. (1835–36) 193 = Macrochaetium Dregei.
junciformis, Boeck. in Linnæa, xxxvii (1874) 267 = Cladium undulatum.
nivicola, F. Muell. in Trans Phil. Soc. Vict. i. (1855) 111 = alpina.
paniculata, *Phil. in Linnæa*, xxix. (1857–58) 80 — Chili.
schoenoides, Banks & Soland. ex Hook. f. Fl. Antarct. 262. t. 148 = alpina.
Schoenorthuna, *Borck. in Flora*, lxii. (1879) 569.—Afr. trop.
Urvilleana, Gaudich. ex Nees, in Linnæa, ix. (1834) 300 = Asterochaete arundinacea.

CARPHALEA, Juss. Gen. 198 (1789). *RUBIACEÆ*, Benth. & Hook. f. ii. 52.
angulata, *Baill. in Bull. Soc. Linn. Par.* i. (1878) 188. —Madag.
corymbosa, *Willd. Sp. Pl.* i. 583.—Madag.
Kirondron, *Baill. in Bull. Soc. Linn. Par.* i. (1878) 188.—Madag.
madagascariensis, Lam. Illustr. i. 258. t. 59 = corymbosa.
Pervilleana, *Baill. in Bull. Soc. Linn. Par.* i. (1878) 188.—Madag.
pubiflora, Moq. & Sesse, ex DC. Prod. iv. 365 = Bouvardia quaternifolia.

CARPHEPHORUS, Cass. in Bull. Soc. Philom. (1816) 198. *COMPOSITÆ*, Benth. & Hook. f. ii. 249.
atriplicifolius, *A. Gray, in Proc. Am. Acad.* v. (1861) 139.—Calif.
baicalensis, DC. Prod. v. 132 = Saussurea pycnocephala.
bellidifolius, *Torr. & Gray, Fl. N. Am.* ii. 66.—Am. bor.
coridifolius, DC. Prod. vii. 267 = Brickellia pinifolia.
corymbosus, *Torr. & Gray, Fl. N. Am.* ii. 67.—Am. bor.
junceus, *Benth. Bot. Voy. Sulph.* 21.—Calif.
juncous, Durand, in Pacif. Rail. Rep. v. 8 = Linosyris spinulosa.
Pseudo-liatris, *Cass. in Bull. Soc. Philom.* (1816) 149. —Am. bor.
revolutifolius, *DC. Prod.* v. 133.—Mexic.
tenuifolius, *Torr. & Gray, Fl. N. Am.* ii. 66.—Am. bor.
triangularis, *A. Gray, Pl. Wright.* i. 86.—Mexic.

CARPHOBOLUS, Schott, in Spreng. Syst. iv. Cur. Post. 409 (1827) = **Piptocarpha**, R. Br. (Compos.).
asteroidrichius, Sch. Bip. in Pollichia, xx.-xxi. (1863) 426 = P. Vauthieriana.
axillaris, Sch. Bip. l. c. 417 = P. axillaris.
Blanchetianus, Sch. Bip. l. c. 423 = P. Lundianus.
cinereus, Sch. Bip. l. c. 419 = P. cinereus.
cuneifolius, Sch. Bip. l. c. 413 = P. oblongus.
latifolius, Sch. Bip. l. c. 426 = P. opaca.
Lechleri, Sch. Bip. l. c. 426 = P. Lechleri.
lepidotus, Sch. Bip. l. c. 414 = P. oblongus.
leprosus, Sch. Bip. l. c. 419 = P. leprosa.
lucidus, Sch. Bip. l. c. 418 = P. lucida.

CARPHOBOLUS :—
Lundianus, Sch. Bip. in Pollichia, xx.–xxi. (1863) 421 = Piptocarpha Lundiana.
macropodus, Sch. Bip. l. c. 418 = P. macropoda.
neurophyllus, Sch. Bip. l. c. 414 = P. pyrifolia.
notatus, Sch. Bip. l. c. 420 = P. notata.
oblongus, Sch. Bip. l. c. 416 = P. oblonga.
opacus, Sch. Bip. l. c. 427 = P. polycephala.
oxyphyllus, Sch. Bip. l. c. 415 = P. oxyphylla.
pellucidus, Sch. Bip. l. c. 420 = P. pellucida.
Poeppigianus, Sch. Bip. l. c. 422 = P. Poeppigiana.
pyrifolius, Sch. Bip. l. c. 415 = P. pyrifolia.
ramuflorus, Sch. Bip. l. c. 429 = P. ramiflora.
Regnellii, Sch. Bip. l. c. 417 = P. axillaris.
Riedelii, Sch. Bip. l. c. 425 = P. Riedelii.
rotundifolius, Sch. Bip. l. c. 424 = P. rotundifolia.
Sellowii, Sch. Bip. l. c. 421 = P. Sellowii.
semiserrulatus, Sch. Bip. l. c. 416 = P. axillaris.
sessiliflorus, Schott, in Spreng. Syst. iv. Cur. Post. 409 = P. lucida.
teretiscalis, Sch. Bip. in Pollichia, xx.-xxi. (1863) 422 = P. tereticaulis.
umbellatulus, Schott, in Spreng. Syst. iv. Cur. Post. 409 = P. umbellulata.

CARPHOCHAETE, A. Gray, Pl. Fendl. 65 ; in Mem. Am. Acad. N. S. iv. (1849). *COMPOSITÆ*, Benth. & Hook. f. ii. 247.
Bigelovii, *A. Gray, Pl. Wright.* i. 89.—N. Mexic.
Grahami, *A. Gray, l. c.*—Mexic.
Wislizeni, *A. Gray, Pl. Fendl.* 65.—Mexic.

CARPHOLOMA, D. Don, in Mem. Wern. Soc. v. (1826) 565 = **Leochrysum**, Willd. (Compos.).
rigidum, D. Don, l. c. = L. ericaefolium.

CARPHOPAPPUS, Sch. Bip. in Walp. Rep. ii. 954, 973 (1848) = **Iphiona**, Cass. (Compos.).
baccharidifolius, Sch. Bip. l. c. 974 = I. baccharidifolia.

CARPHOSTEPHIUM, Cass. in Dict. Sc Nat. xlv. 62 (1826) = **Tridax**, Linn. (Compos.).
trifidum, Cass. l. c. = T. trifida.

CARPIDOPTERIX, Karst. Fl. Columb. ii. 45. t. 128 (1862?) = **Thounia**, Poit. (Sapindac.).
macroptera, Karst. l. c. = T. macroptera.

CARPINUS, Linn. Gen. ed. I. 292 (1737). *CUPULIFERÆ*, Benth. & Hook. f. iii. 406.
CARPINUM, Rafin. in Am. Monthly Mag. (1818) 268.
DISTEOCARPUS, Sieb. & Zucc. Fam. Nat. Fl. Jap. ii. 102. t. 8 (1846).
americana, *Micha. Fl. Bor. Am.* ii. 201.—Am. bor.
Betulus, *Linn. Sp. Pl.* 998.—Europ. ; Reg. Cauc. ; Persia.
caroliniana, Walt. Fl. Carol. 236 = americana.
Carpinisaa, Kit. in Host, Fl. Austr. ii. 626 = C. Betulus.
compressa, Gilib. Exercit. ii. 399 = C. Betulus.
cordata, *Blume, Mus. Bot. Lugd. Bat.* i. 309.—Japon.
duinensis, Scop. Fl. Carn. ed. II. ii. 243. t. 60 = orientalis.
edentulus, *Kit. in Rochel, Pl. Banat. Rar.* nomen.—Banat.
eroza, *Blume, Mus. Bot. Lugd. Bat.* i. 308.—Japon.
faginea, *Lindl. in Wall. Pl. As. Rar.* ii. 5.—Reg. Himal.
intermedia, Wierzb. ex Reichb. Ic. Fl. Germ. xii. 4. t. 633 = C. Betulus.
italica, Scop. ex Steud. Nom. ed. II. i. 300 = Ostrya carpinifolia.
japonica, *Blume, Mus. Bot. Lugd. Bat.* i. 308.—Japon.
laxiflora, *Blume, l. c.* 309.—Japon.
minor, *Pall. Reise*, ii. 95.—Tauria.
nervata, Dulac, Fl. Hautes-Pyr. 141 = C. Betulus.
nigra, Moench, Vers. Ausl. Baeume, 19 = orientalis.
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 3.—Europ. austr.
Ostrya, Linn. Sp. Pl. 998 = Ostrya virginica.
ostryoides, Rafin. in Med. Repos. N. York, Ser. III. ii. (1811) 353 = americana !
quercifolia, Desf. ex Steud. Nom. ed. II. i. 300 = C. Betulus.

CARPINUS:—
sepium, Lam. Fl. Fr. ii. 219 = C. Betulus.
trifora, Moench, Meth. 694 = Ostrya virginica.
Tschonoskii. *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 534.—Japon.
Turczaninowii, *Hance, in Journ. Linn. Soc.* x. (1869) 303.—China.
ulmifolia, Salisb. Prod. 392 = C. Betulus.
ulmoides, S. F. Gray, Nat. Arr. Brit. Pl. ii. 245 = C. Betulus.
viminea, *Wall. ex Lindl. in Wall. Pl. As. Rar.* ii. 4. t. 106.—Kog. Himal.
virginiana, Michx. Hist. Arb. For. iii. 57. t 8 = americana.
virginiana, Mill. Gard. Dict. ed. VIII. n. 4; Lam. Encyc. i. 708 = Ostrya virginiana.
vulgaris, Mill. l. c. n. 1 = C. Betulus.
yedoënsis, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 535.—Japon.

CARPIPHEA, Rafin. Sylva Tellur. 39 (1838) = **Cordia**, Tourn. (Boragin.).
dentata, Rafin. l. c. = Cordia dentata.

CARPOCALYMNA, Zipp. in Alg. Konst. en Letterbode, i. (1829) 297 = **Epithema**, Blume (Gesneriac.).
monophylla, Zipp. ex Span. in Linnæa, xv. (1841) 331 = E. diffornia.

CARPOCERAS, Link, Handb. ii. 289 (1831) = **Thlaspi**, Tourn. (Crucif.).
cappadocicum, Boiss. Diagn. Ser. II. vi. 19.—Cappadoc.
cilicicum, Boiss. & Kurz, ex Boiss. Fl. Orient. i. 332. —Cilicia.
Griffithianum, Boiss. Diagn. Ser. II. i. 40 = T. alpestre.
hastulatum, Boiss. l. c. Ser. I. viii. 35 = T. hastulatum
oxyceras, Boiss. l. c. 37 = T. oxyceras.
sibiricum, Link, Handb. ii. 289 = T. ceratocarpon.
stenocarpum, Boiss. Diagn. Ser. I. viii. 38 = T. stenocarpum.

CARPOCERAS, A. Rich. Elem. Bot. (1846) 706 = **Martynia**, Linn. (Pedalin.).
longifora, A. Rich. l. c. = M. diandra.

CARPODETES, Herb. App. 41 (1821) = **Stenomesson**, Herb. (Amaryllid.).
recurvata, Herb. l. c. = S. recurvatum.

CARPODETUS, Forst. Char. Gen. 33. t. 17 (1776). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 646.
dentatus, Poir. Encyc. Suppl. ii. 120 = serratus
Forsteri, Roem. & Schult. Syst. v. 604 = serratus.
serratus, *Forst. Char. Gen.* 34. t. 17 A.—N. Zel.

CARPODINUS, R. Br. ex Sabine, in Trans. Hort. Soc. v. (1823) 455. *APOCYNACEÆ*, Benth. & Hook. f. ii. 693.
acida, *Sabine, l. c.* 456.—Afr. trop.
dulcis, *Sabine, l. c.* 455.—Afr. trop.

CARPODIPTERA, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 163. *TILIACEÆ*, Benth. & Hook. f. i. 229.
africana, *Mast. in Oliver, Fl. Trop. Afr.* i. 241.—Afr. trop.
Boivini, *Baill. Adansonia*, x. (1872) 180.—Ins. Comor.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 164.—Cuba.
Schomburgkii, *Baill. Adansonia*, x. (1872) 181.—Guiana.

CARPODON, Spreng. Syst. ii. 634 (1825) = seq.

CARPODONTOS, Labill. Voy. ii. 16. t. 18 (1798) = **Eucryphia**, Cav. (Rosac.).
lucida, Labill. l. c. = E. Billardieri.

CARPOLOBIA, G. Don, Gen. Syst. i. 870 (1831), partim. *POLYGALEÆ*, Benth. & Hook. f. i. 339, 344.
Afzeliana, *Oliver, Fl. Trop. Afr.* i. 136.—Afr. trop.
alba, *G. Don, Gen. Syst.* i. 870.—Afr. trop.
dubia, G. Don, l. c.=Baphia polygalacea.
lutea, *G. Don, l. c.*—Afr. trop.
parvicolor, G. Don, l. c. = Baphia nitida.

CARPOLIZA, Steud. Nom. ed. II. i. 300 (1840) = seq.

CARPOLYZA, Salisb. Parad. Lond. t. 68 (1807). *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 721.
Hessea, Berg, ex Schlecht. in Linnæa, i. (1826) 253, non Herb.
spiralis, *Salisb. Parad. Lond.* t. 68.—Afr. austr.
stellaris, Heynh. Nom. ii. 122 = Hessea stellaris, *Herb.*

CARPONEMA, Eckl. & Zeyh. Enum. 8 (1835). *CRUCIFERÆ*, Benth. & Hook. f. i. 102.
aggregata, Eckl. & Zeyh. l. c. = Heliophila gracilla. filiformis, *Eckl. & Zeyh. l. c.*—Afr. austr.

CARPOPHILLUS, Neck. Elem. ii. 84 (1790) = **Pereskia**, Plum.† (Cactear.).

CARPOPHORA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 139. t. 82 (1862) = **Silene**, Linn. (Caryophyll.). *Hoffmeisteri*, Klotzsch, l. c. = S. Griffithii.

CARPOPHYLLUM, Miq. Fl. Ind. Bat. Suppl. i. 401 (1860) = **Starculia**, Linn.
macropodum, Miq. l. c.—Sumatra.

CARPOPHYLLUM, Neck. Elem. iii. 402, 411 (1790) = CARPOPHILLUS, Neck.

CARPOPODIUM, Eckl. & Zeyh. Enum. i. 13 (1835) = **Heliophila**, Linn. (Crucif.).
carnosum, Eckl. & Zeyh. l. c. = H. carnosa.
cleomoides, Eckl. & Zeyh. l. c. = H. glauca.
Thunbergii, Eckl. & Zeyh. l. c. = H. Thunbergii.

CARPOPOGON, Roxb. Hort. Beng. 54 (1814); Fl. Ind. iii. 183 (1832) = **Mucuna**, Adans. (Legumin.).
anguineum, Roxb. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 135 = M. monosperma.
atropurpureum, Roxb. Hort. Beng. 54; Fl. Ind. iii. 284 = M. atropurpurea.
bracteatum, Roxb. Hort. Beng. 54 = M. bracteata.
capitatum, Roxb. l. c. [98]; Fl. Ind. iii. 284 = M. capitata.
giganteum, Roxb. ll. cc. 54, 287 = M. gigantea.
imbricatum, Roxb. Hort. Beng. 54 = M. imbricata.
monospermum, Roxb. l. c.; Fl. Ind. iii. 283 = M. monosperma.
niveum, Roxb. ll. cc. 54, 285 = M. nivea.
pruriens, Roxb. ll. cc. 54, 283 = M. pruriens.
volubilis, Roxb. ex Steud. Nom. ed. II. i. 300 = M. volubilis.

CARPOTHALIS, E. Mey. in Drège, Zwei Pfl. Docum. 170 (1843) = **Krauseola**, Harv. (Rubiac.).
lanceolata, E. Mey. l. c. = K. lanceolata.

CARPOTRICHE, Reichb. Nom. 189 (1841) = seq.

CARPOTROCHE, Endl. Gen. 918 (1839). *BIXINEÆ*, Benth. & Hook. f. i. 125.
brasiliensis, *Endl. l. c.*—Bras.
denticulata, *Benth. in Hook. Journ. Bot.* iv. (1842) 112.—Guiana.
grandiflora, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiii. i. 437.—Bras.
laxiflora, *Benth. in Hook. Journ. Bot.* iv. (1842) 114. —Guiana.
longifolia. *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 92.—Bras.; Peruv.
paludosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 114. —Guiana.

CARPOXIS, Rafin. New Fl. Am. iii. 89 (1836) = **Forestiera**, Poir. (Oleac.).
inermis, Rafin. l. c. 92.—Am. bor.
nitida, Rafin. l. c. 91 = F. acuminata.
prinoides, Rafin. l. c. 92 = F. prinoides.
retusa, Rafin. l. c. = F. retusa.
spinosa, Rafin. l. c.—Am. bor.

CARPOXYLON, H. Wendl. & Drude, in Kerch. Palm. 388 (1878), nomen. *PALMÆ*.
macrospermum, *H. Wendl. & Drude, l. c.*—Hab.?

CARPUNYA, Presl, Epim. Bot. 228 (1849) = **Piper**, Linn.
Lauxtiana, Presl, l. c. 229 = P. Carpunya.
oblongata, Presl, l. c. = P. oblongatum.
peruviana, Presl, l. c. = P. Carpunya.

CARPUPICA, Rafin. Sylva Tellur. 35 (1838) = **Piper**, Linn.
odorata, Rafin. l. c. = P. Carpunya.

CARRADORIA, A. DC. Prod. xii. 610 (1848) = **Globularia**, Linn. (Selagin.).
incanescens, A. DC. l. c. = G. incanescens.

CARREGNOA, Besler, Hort. Eyst. Vern. ii. 606 (1842) = **Tapeinanthus**, Herb. (Amaryllid.).
dubia, Perez-Lara, in Anal. Soc. Esp. Hist. Nat. xi. (1882) 399 = Narcissus serotinus!
humilis, J. Gay, in Ann. Sc. Nat. Sér. IV. x. (1858) 98 = T. humilis.
lutea, Boiss. Voy. Bot. Espagne, ii. 606 = T. humilis.

CARRIA, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 6 = **Gordonia**, Ellis (Ternstroem.).
speciosa, Gardn. l. c. 7 = G. speciosa.

CARRICHTERA, Adans. Fam. ii. 421 (1763); DC. Syst. Veg. ii. 641. *CRUCIFERÆ*, Benth. & Hook. f. i. 86.
Vella, *DC. l. c.* 642.—Hispan.

CARROA, Presl, Symb. Bot. ii. 25 (1858) = **Dalea**, Linn. (Legumin.).
diffusa, Presl, l. c. = D. diffusa.
grandulosa, Presl, l. c. = D. diffusa.

CARRONIA, F. Muell. Fragm. ix. 171 (1875). *MENISPERMACEÆ*.
multisepalea, *F. Muell. l. c.*—Austral.

CARRUTHERSIA, Seem. Fl. Vit. 155. t. 30 (1865–73). *APOCYNACEÆ*, Benth. & Hook. f. ii. 718.
pilosa, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 129.—Ins. Philipp.
scandens, *Seem. Fl. Vit.* 156. t. 30.—Ins. Fiji.

CARTALINIA, Saov. ex Kunth, Enum. Pl. v. 119 (1850) = **Paris**, Rupp. (Liliac.).

CARTERETIA, A. Rich. Sert. Astrol. 10. t. 4 (1834) = **Doritis**, Lindl. (Orchid.).
paniculata, A. Rich. l. c. = D. paniculata.

CARTESIA, Cass. in Bull. Soc. Philom. (1816) 198 = **Stokesia**, L'Hérit. (Compos.).
centauroides, Cass. l. c. = S. cyanea.

CARTHAMUS, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITÆ*, Benth. & Hook. f. ii. 488.
Durandoa, Pomel, Mat. Fl. Atl. 9 (1860).
Heracantha, [Hoffmgg. &] Link, Fl. Port. ii. 205 (1821).
Heteracantha, Steud. Nom. ed. II. i. 755 (1840).
Honeywartha, Vest, in Flora, iii. (1820) 1.
Kentrophyllum, Neck. Elem. i. 86 (1790).
Onobroma, DC. Prod. vi. 618 (1837).
Phonus, Hill, Veg. Syst. iv. 8 (1762).
acaulis, Presl, ex Steud. Nom. ed. II. i. 300 = Carduncellus pectinatus.
africanus, Lam. Encyc. i. 640 = Berckheya grandiflora.
albus, *Desf. Tabl.* ed. I. 92.—Oriens.
aranosus, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 268 = Carduncellus aranosus.
arborescens, *Linn. Sp. Pl.* 821.—Hispan.
arvalis, Willd. Enum. Hort. Berol. 845 = flavescens.
basticus, Nym. Consp. 419 = lanatus.
brasiliensis, *Vell. Fl. Flum.* 243; vill. t. 77.—Bras.
caerulescens, Linn. Sp. Pl. 820 = Carduncellus caerulescens.
cancellatus, Lam. Fl. Fr. ii. 10 = Atractylis cancellata.
canescens, Lam. l. c. = Cnicus Acarna.
canescens, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 262. —Aleppo.
Carduncellus, Linn. Sp. Pl. 821 = Carduncellus mospelicensium.
carolinianus, Walt. Fl. Carol. 194 = Elephantopus carolinianus!

CARTHAMUS:—
corymbosus, Linn. Sp. Pl. ed. II. 1164 = Cardopatium corymbosum.
creticus, Linn. l. c. 1163 = lanatus.
creticus, Nym. Consp. 419 = glaucus.
cyanens, Banks, ex Steud. Nom. ed. II. i. 300 = Stokesia cyanea.
cynaroides, Bieb. Tabl. Casp. 195 = Cousinia cynaroides.
dentatus, *Vahl, Symb. Bot.* i. 69.—As. Min.
dianius, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 363 = Carduncellus caeruleus.
elatus, *Nym. Consp.* 419.—Italia.
ferox, Lam. Fl. Fr. ii. 31 = Cnicus ferox.
flavescens, *Willd. Sp. Pl.* iii. 1706.—As. Min.; Syria.
fluminensis, *Vell. Fl. Flum.* 342; viii. t. 78.—Bras.
glaber, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
glaucus, *Bieb. Fl. Taur. Cauc.* ii. 284.—Aegypt.; As. Min.; Persia.
gummiferus, Linn. Spec. i. 889 = Atractylis gummifera.
helenioides, *Desf. Fl. Atlant.* ii. 258. t. 230.—Algeria.
hircinus, Lag: ex Spreng. Syst. iii. 391 = arborescens.
hispanicus, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 363 = Carduncellus caeruleus.
humilis, Lam. Encyc. i. 688 = Carduncellus pinnatus.
integrifolius, Hort. ex Webb & Berth. Phyt. Canar. ii. 344 = Carlina salicifolia.
involucratus, Lam. Fl. Fr. ii. 12 = Cnicus spinosissimus.
laevis, Hill, Hort. Kew. 57 = Stokesia cyanea.
lanatus, *Linn. Sp. Pl.* 830.—Europ.; Oriens.
leuccaulos, *Sibth. & Sm. Fl. Graec.* t. 842.—Archip. Graec.
linearifolius, *Molina, Sagg. Chile*, ed. II. 294.—Chili.
maculatus, Lam. Encyc. i. 688 = Silybum Marianum.
magellanicus, *Lam. l. c.* 640.—Reg. Magellan.
marcoticus, *Delile, Fl. Egypte*, 276.—Aegypt.
mitissimus, Linn. Sp. Pl. 831 = Carduncellus mitissimus.
multifidus, Desf. Fl. Atlant. ii. 256 = Carduncellus caeruleus
nitidus, *Bess. Fl. Orient.* iii. 708.—Syria.
Oxyacantha, *Bieb. Fl. Taur. Cauc.* ii. 283.—Reg. Caucas.; As. occ.
pectinatus, Desf. Fl. Atlant. ii. 257 = Carduncellus pectinatus.
persicus, Desf. ex Willd. Sp. Pl. iii. 1707 = flavescens.
persicus, Willd. ex Boiss Fl. Orient. iii. 708 = leucocaulos.
pinnatus, Cstand. Fl. Madrit. 412 = Carduncellus monspelliensium.
pinnatus, Desf. Fl. Atlant. ii. 258 = Carduncellus pinnatus.
polyacanthus, *Bieb. Tabl. Casp.* 58.—Transcauc.
pterocaulos, C. A. Mey. Verz. Pfl. Cauc. 67 = Cousinia Hystrix.
rigidus, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
rigidus, Willd. Enum. Hort. Berol. 845 = arborescens.
salicifolius, Linn. f. Suppl. 350 = Carlina salicifolia.
tauricus, Bieb. Fl. Taur. Cauc. ii. 285 = lanatus.
Teneriffae, *Y. E. Gmel. ex Steud. Nom.* ed. I. 164.—Ins. Teneriffa.
tinctorius, *Linn. Sp. Pl.* 830.—Geront. cult.
tingitanus, Linn. l. c. ed. II. 1163 = Carduncellus caeruleus.
tomentosus, J. F. Gmel. Syst. 1252. Index (quid?).
turbinatus, *Nym. Consp.* 419.—Sicilia.

CARTODIUM, Soland. ex R. Br. in Trans. Linn. Soc. xii. (1817) 106 = **Craspedia**, Forst. f. (Compos.).

CARTONEMA, R. Br. Prod. 571 (1810). *COMMELINACEAE*, Benth. & Hook. f. iii. 859.
brachyantherum, *Benth. Fl. Austral.* vii. 92.—Austral.
parviflorum, *Hassk. in Flora*, iii. (1869) 865.—Austral.
Philydroides, F. Muell. Fragm. i. 63 = spicatum.
spicatum, *R. Br. Prod.* 271.—Austral.
spicatum, F. Muell. Fragm. i. 63 = parviflorum.
tenue, *Cornet, in Nuov. Giorn. Bot. ni.* (1879) 216.—Austral.
trigonospermum, *C. B. Clarke, in DC. Monog. Phan.* iii. 864.—Austral.

CARTREMA, Rafin. Sylva Tellur. 184 (1838) = **Osmanthus**, Lour. (Oleac.).
americana, Rafin. l. c. = O. americanus.

CARUELIA, Parl. Neov. Gen. e Spec. 21 (1854) = Cynthiogalum, Tenore. (Liliac.).
arabica, Parl. l. c. 22 = O. arabicum.

CARUM, Rup. ex Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 890.
ANISACTIS, Dulac, Fl. Hautes-Pyr. 347 (1867).
ATAENIA, Hook. & Arn. Bot. Beech. Voy. 349 (1840).
BULBOCASTANUM, Schur, Enum. Pl. Transs. 249 (1866).
BUNIUM, Linn. Gen. ed. I. 72 (1737).
CHAMAREA, Eckl. & Zeyh. Enum. iii. 346 (1837).
EDOSMIA, Nutt. in Torr. & Gray, Fl. N. Am. i. 612 (1840).
ELWENDIA, Bolss. in Ann. Sc. Nat. Sér. III. i. (1844) 140.
FRORIEPIA, C. Koch, in Linnaea, xvi (1842) 862.
GEOGLYPUM, Cosa. & Dur. Pl. Nouv. et Crit. Exp. 112 (1851).
HLADNIKIA, Reichb Ic. Pl. Crit. ix. 9 (1831); et Fl. Germ. Excurs. 276 (1832).
HUETIA, Bolss. Diagn. Ser. II. ii. 108 (1856).
PETROSELINUM, Hoffm. Umb. i. 78 (1814).
PHYMATIS, E. Mey. in Drège, Zwei Pfl. Docum. 211 (1848).
PTYCHOTIS, Koch, in Nov. Act. Nat. Cur. xii. t. (1824) 194.
RIDOLFIA, Moris, Ind. Sem. Hort. Taur. (1841); Fl. Sard. ii. 215 (1849).
SELINOPSIS, Coss. & Dur. ex Munby, Cat. Pl. Alg. ed. II. 18 (1859), nomen.
SYMPOLIUM, C. Koch, in Linnaea, xvi. (1842) 856.
TRACHYSPERMUM, Link, Enum Hort. Berol. i. 267 (1821).
TRACHYSTIADIUM, Eckl & Zeyh. Enum. 341 (1837).
WYTHLERIA, DC. Coll. Mém. v. 26 (1820).
Ziziа, Koch, in Nov. Act. Nat. Cur. xii. t. (1824) 128.
Ajowan, Benth. & Hook. f. Gen. i. 891 = copticum.
alliaceum, *Franch, in Ann. Sc. Nat. Sér.* VI. xvi. (1883) 294.—Turkest.
alpinum, *Benth. & Hook. f. Gen.* i. 891.—Europ.; Persia.
anmoides, *Benth. & Hook. f. l. c.* 892.—Reg. Mediterr.
anethifolium, *Benth. & Hook. f. l. c.* 891.—Reg. Himal.
Anisum, Baill. Hist. Pl. vii. 119, 178 = Pimpinella Anisum.
apiculatum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 358.—As. centr.
armenum, *Boiss. in Ann. Sc. Nat*, Sér. III. i (1844) 147.—Reg. Caucas.
aromaticum, Salish. Prod. 168 = C. Carvi.
atrosanguineum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 359.—As. centr.
avenaceum, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 888.—Persia.
aureum, Benth. & Hook. f. Gen. i. 891 = Thaspium aureum.
Bourgaei, *Boiss. Fl. Orient.* iii. 885.—As. Min.
brachycarpum, Boiss. Diagn. Sér. I. x. 23 = ferulaefolium.
Bulbocastanum, *Koch, in Nov. Act. Nat. Cur.* xii. t. (1824) 121.—Europ.; As. bor.
Bunius, *Linn. Syst. Nat.* ed. XII. 783.—Europ.
bupleuroides, *Schrenk, ex Fisch. & Mey. in Bull. Phys.-Math. Acad. Petersb.* iii (1845) 305.—Soongar.
buriaeticum, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 713.—Sibir. baical.
calcicolum, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1880) 514.—Ins. Socotra.
capense, *Sond. in Harv. & Sond. Fl. Cap.* ii. 538.—Afr. austr.
Capsul, *Franch. in Ann. Sc. Nat. Sér.* VI. xvi. (1883) 298.—Turkestan.
Carvi, *Linn. Sp. Pl.* 263.—Europ.; Orient; As bor.
carvifolium, *Benth. & Hook. f. Gen.* i. 892.—Afr. austr.
caucasicum, *Boiss. Fl. Orient.* ii. 880.—Reg. Caucas.
chaerophylloides, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 585.—Turkest.
coriandrum, *Benth. & Hook. f. Gen.* i. 891.—Europ.; Afr. bor.; As. bor.
cordatum, Benth. & Hook. f. l. c. 891, 913 = Thaspium trifoliatum.

CARUM:—
cornigerum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii 887.—Kurdistan.
creticum, *Benth. & Hook. f. Gen.* i. 891. —Creta.
cylindricum, *Boiss. & Haussk. Diagn.* Ser. I. x. 23.—Persia; Baluchist.
dancoides, *Boiss. in Ann. Sc. Nat.* Sér. III i. (1844) 139.—Macedon.; Lydia.
decussatum, Gillib. Fl. Lituan. ii. 37 = C. Carvi.
dichotomum, *Benth. & Hook. f. Gen.* i. 891.—Reg. Mediterr. occ.
didymum, *Benth. & Hook. f. l. c.*—Afr. austr.
dissectum, Baill. Hist. Pl. vii. 179 = Pimpinella magna.
divaricatum, Koch, Syn. Fl. Germ. 215 = ferulaefolium.
divergens, *Boiss. & Huet, Diagn.* Ser. II. ii. 80.—Armenia.
diversifolium, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 683.—Reg. Himal.
elegans, *Fenzl, Pugill. Pl. Nov. Syr*, 16.—Oriens.
Elvendia, *Boiss. Fl. Orient.* ii. 888.—Persia.
Falcaria, *Lange, in Willd. & Lange, Prod. Fl. Hisp.* iii. 92.—Hispan.
falcarioides, Bolss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 96 = elegans.
Falconeri, *C. B. Clarke, in Hook. Fl. Brit. Ind.* ii. 688.—Reg. Himal.
ferulaefolium, *Boiss. Diagn.* Ser. I. x. 23.—Reg. Mediterr. or.
flexuosum, Fries, Summa Veg. Scand. 24, 180 = Conopodium denudatum.
flexuosum, *Nym. Consp.* 307.—Italia.
Gairdneri, *A. Gray, in Proc. Am. Acad.* vii (1868) 344.—Am. bor. occ.
gracile, *Lindl. in Royle, Illustr. Bot. Himal.* 232.—Ind. or.
graecum, *Boiss. & Heldr. Diagn.* Ser. I. vi. 58.—Graecia; Serbia.
Hallii, *S. Wats. Bibliog. Ind. N. Am. Bot.* 416.—Calif.
Heldreichii, *Boiss. Diagn.* Ser. II ii 78.—Graecia
heterophyllum, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 586.—As. temp.
hispanicum, *Mill. Gard. Dict.* ed. VIII. n. 9.—Hispan.
hispidum, *Benth. & Hook. f. Gen.* i. 891.—Afr. austr.
Huetii, Boiss. Diagn. Ser. II. ii. 81 = Bourgaei
humile, Boiss. & Bal. ex Bolss. Fl. Orient. ii. 881 = caucasicum.
incrassatum, *Boiss. Voy. Bot. Espagne*, ii. 239, 781.—Hispan.
indicum, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 72.—Soongaria.
inodorum, *Sieb. in Pall. N. Nord. Beitr.* vii. 229.—Rossia.
intermedium, Benth. & Hook. f. Gen. i. 892 = inundatum.
inundatum, *Lespin. in Act. Soc. Bord.* xiv. (1845) 270.—Europ.
involucratum, Baill. Hist. Pl. vii. 179 = Roxburghianum.
Kelloggii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 344.—Calif.
khasianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 682.—Reg. Himal.
latifolium, *Benth. & Hook. f. Gen.* i. 892.—Europ.
leucocoli on, *Boiss. & Huet, Diagn.* Ser. II. ii. 80.—Armenia.
lomatocarum, *Boiss. Fl. Orient.* iii. 879.—Reg. Caucas.
lutescens, Turcz. ex Walp. Rep. v. 866 (= *Cnidium multicaule*).—Sibir.
macrocarpum, *Benth. & Hook. f. Gen.* i. 891.—Graecia.
Maesus, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 81.—Hispan.
magnum, Baill. Hist. Pl. vii. 178 = Pimpinella magna.
mauritanicum, *Boiss. & Reut. Pugill.* (1852) 49.—Afr. bor. occ.
meifolium, Boiss. Fl. Orient. ii. 880 = armenum.
Meum, Stokes, Bot. Mat. Med. ii. 138 = Meum athamanticum.
microcarpum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 137.—Lydia.
multifforum, *Boiss. Fl. Orient.* ii. 882,—Graecia; As. Min.
neurophyllum, *Franch. & Sav. Enum. Pl. Jap.* i. 180.—Japon.
nigrum, Baill. Hist. Pl. vii. 178 = Pimpinella Saxifraga.

CARUM :—
nivale, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 84. —Hispan.
Noëanum, Boiss. Diagn. Ser. II. ii. 77 = elegans.
nothum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 681.—Zeylan.!
officinale, S. F. Gray, Nat. Arr. Brit. Pl ii. 515 = C. Carvi.
oreganum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 365.—Am. bor. occ.
Panatjan, Baill. Hist. Pl. vii. 178 = Pimpinella Panatjan.
Panali, *Griseb. in Goett. Abh* xxiv. (1879) 146.—Reg. Argent.
paucifolium, DC. Prod. iv. 117 = elegans.
peregrinum, *Linn. Disc. Dem. Pl. in Amoen. Acad.* iii. 406 —Peruv.!
persicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 188.—Persia.
Pestalozzæ, *Boiss. Fl. Orient.* ii. 886.—Persia; Beluchist.
Petroselinum, *Benth. & Hook. f. Gen.* i. 891.—Geront. cult.
pimpinelloides, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 514.—Ins. Socotra.
Podagraria, Roth, Enum. Pl. Phan. Germ. i. 246 = Aegopodium Podagraria.
polyphyllum, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 881.—Syria.
purpurascens, Boiss. in Tchihat. Asie Min. Bot. i. 411 = elegans.
rectangulum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 884.—Persia.
Ridolfia, *Benth. & Hook. f. Gen.* i. 891.—Reg. Mediterr.; Orient.
rigidulum, *Koch, in DC. Prod.* iv. 115.—Italia.
rigidum, Steud. Nom. ed. II. i. 301 = rigidulum.
Roxburghianum, *Benth. & Hook. f. Gen.* i. 891.—Ind. or.; Malaya.
rupestre, *Boiss. & Heldr. Diagn.* Ser. II. ii. 79.—Graecia.
Saxifraga, Baill. Hist. Pl. vii. 178 = Pimpinella Saxifraga.
Saxifraga, Benth. & Hook. f. Gen. i. 892 = C. Bunias.
segetum, *Benth. & Hook. f. l. c.*—Europ.
setaceum, *Schrenk, Enum. Pl. Nov.* 61.—Soongaria.
Seweerzowi, *Regel, in Act. Hort. Petrop.* v. (1877) 587, 588.—Turkestan.
simplex, Steph. ex Willd. Sp. Pl. i. 1470 = Sesell annuum.
Sisarum, Baill. Hist. Pl. vii. 179 = Sium Sisarum.
stictocarpum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 681.—Ind. or.
strictum, *Boiss. Fl. Orient.* ii. 887.—Macedon.
sulcatum, Steud. Nom. ed. I. 164, 49 = Selinum Carvifolia.
sylvestre, *Baill. Hist. Pl.* vii. 179.—Ind. or.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* ii. 371.—Japon.
tenuifolium, *Benth. & Hook. f. Gen.* i. 891.—Hispan.; Marocco.
Thorei, Benth. & Hook. f. l. c. 888, in obs. = inundatum.
trichophyllum, *Schrenk, Enum. Pl. Nov.* 61.—Soongaria.
verticillatum, *Koch, in Nov. Act. Nat. Cur.* xii. i. (1824) 122.—Europ.

CARUMBIUM, Kurz, For. Fl. Brit. Burma, ii. 411 (1877) = **Saptium**, P. Br. (Euphorb.).
baccatum, Kurz, l. c. 412 = S. baccatum.
inerme, Kurz, l. c.—Burma.
sebiferum, Kurz, l. c. = S. sebiferum.

CARUMBIUM, Reinw. Elench. Sem. Hort. Leid. ex Isis (1828) 819; et in Syll. Ratisb. ii. (1828) 6 = **Homalanthus**, A. Juss. (Euphorb.).
acuminatum, Muell. Arg. in DC. Prod. xv. ii. 1144. —Ins. Pacif.
amboinicum, Miq. Fl. Ind. Bat. i. ii. 413.—Malaya.
fastuosum, Muell. Arg. in DC. Prod. xv. ii. 1144 = H. fastuosus.
giganteum, Muell. Arg. ex Miq. Fl. Ind. Bat. i. ii. 587 = H. giganteus.
Morrenhoutianum, Muell. Arg. in DC. Prod. xv. ii. 1148.—Ins. Pacif.
nutans, Muell. Arg. i. c. = H. nutans.
pallidum, Muell. Arg. in Linnaea, xxxii. (1863) 85 = H. Leschenaultianus.
pedicellatum, Miq. Fl. Ind. Bat. i. ii. 414 = H. pedicellatus.

CARUMBIUM :—
platyneuron, Muell. Arg. in DC. Prod. xv. ii. 1145 = Homalanthus Leschenaultianus.
polyandrum, Hook. f. Handb. New Zeal. Fl. 248.—N. Zel.
populifolium, Reinw. in Blume, Cat. Hort. Buitenz. 105 = H. Leschenaultianus.
populneum, Muell. Arg. in DC. Prod. xv. ii. 1144 = præc.
Sieberi, Muell. Arg. in Linnaea, xxxii. (1863) 85 = H. Leschenaultianus.
stillingiaefolium, Baill. Adansonia, vi. (1865–66) 225 = H. stillingiaefolius.

CARUNCULARIA, Haw. Syn. Pl. Succ. 333 (1812) = **Stapelia**, Linn. (Asclepiad.).
aperta, Sweet, Hort. Brit. ed. II. 859 = S. aperta.
Jacquini, Sweet, l. c. = S. pedunculata.
Massoni, Sweet, l. c. = S. pedunculata.
pedunculata, Haw. Syn. Pl. Succ. 334 = S. pedunculata.
pendulifera, Sweet, Hort. Brit. ed. II. 339 = S. pendulifloras.
serrata, Decne. in DC. Prod. viii. 658, sphaim. = sec.
serrulata, G. Don, Gen. Syst. iv. 122 = Piaranthus serrulatus.
Simsii, Sweet, Hort. Brit. ed. II. 358 = S. pedunculata.

CARVI, [Tourn.] Benth. Syst. Verz. Erf. 114 (1800) = **Selinum**, Tourn. (Umbellif.).
sulcatum, Benth. l. c. 169 = S. Carvifolia.

CARVIFOLIA, C. Bauh. ex Vill. Hist. Pl. Dauph. ii. 629 (1787) = **Selinum**, Tourn. (Umbellif.).

CARYA, Nutt, Gen. Am. ii. 220 (1818). *JUGLANDEAE*, Benth. & Hook. f. iii. 398.
HICORIUS, Rafin. Fl. Ludov. 109 (1817).
SCORIA, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen prius.
alba, *Nutt. l. c.* 221.—Am. bor.
amara, *Nutt. l. c.* 222.—Am. bor.
angustifolia, Sweet, Hort. Brit. ed. I. 97 = olivaeformis.
aquatica, *Nutt. Gen. Am.* ii. 222.—Am. bor.
compressa, G. Don, in Loud. Hort. Brit. 384 = alba.
cordiformis, C. Koch, Dendrol. i. 597 = sulcata.
glabra, Sweet, Hort. Brit. ed. I. 97 = porcina.
illinoënsis, C. Koch, Dendrol. i. 593 = olivaeformis.
integrifolia, *Spreng. Syst.* iii. 849.—Louisian.
laciniosa, Loud. Hort. Brit. 384, nomen = sulcata.
mexicana, *Ingersoll. ex Hemsl. Biol. Centr. Am. Bot.* iii. 162.—Mexic.
microcarpa, *Nutt. Gen. Am.* ii. 221.—Am. bor.
myristiciformis, *Nutt. l. c.* 222.—Am. bor.
obcordata, Sweet, Hort. Brit. ed. I. 97 = porcina.
olivaeformis, *Nutt. Gen. Am.* ii. 221.—Am. bor.
ovata, C. Koch, Dendrol. i. 598 = alba.
porcina, *Nutt. Gen. Am.* ii. 222.—Am. bor.
pubescens, Sweet, Hort. Brit. ed. I. 97 = sulcata.
sulcata, *Nutt. Gen. Am.* ii. 221.—Am. bor.
tetraptera, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 80 = olivaeformis.
texana, *C. DC. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 34.—Texas.
tomentosa, *Nutt. Gen. Am.* ii. 221.—Am. bor.

CARYOCAR, Linn. Mant. ii. 247. n. 1314 (1771). *TERNSTROMIACEAE*, Benth. & Hook. f. i. 180.
ACARYA, Arruda, ex Koster, Trav. Braz. 491 (1816).
BAROLLARA, Neck. Elem. ii. 822 (1790).
PEKEA, Aubl. Pl. Gui. 594 (1775).
RHIZOBOLUS, Gaertn. ex Schreb. Gen. 360 (1789); et Fruct. ii. 93. t. 98 (1791).
SAOUARI, Aubl. Pl. Gui. 599 (1775).
amygdaliferum, *Cav. Ic.* iv. 57. t. 361, 362.—N. Granat.
amygdaliforme, *Ruiz & Pav. Fl. Per.* v. t. 570.—Peruv.
barbinerve, *Miq. in Linnaea*, xxii. (1849) 802.—Bras. brasiliense, *St.-Hil. Fl. Bras. Mer.* i. 322.—Bras.
butyrosum, *Willd. Sp. Pl.* ii. 1243.—Guiana.
edule, *Casar. Decad. Nov. Stirp. Bras.* viii. 87.—Bras.
glabrum, *Pers. Syn.* ii. 84.—Guiana.
nuciferum, *Linn. Mant.* ii. 247.—Guiana.
tomentosum, *Willd. Sp. Pl.* ii. 1244.—Guiana.
tuberculorum, Baill. Hist. Pl. iv. 251 = tomentosum.
villosum, *Pers. Syn.* ii. 84.—Guiana.

CARYOCHLOA, Spreng. Syst. iv. Cur. Post. 22 (1827) = **Oryzopsis**, Michx. (Gramin.).
chilensis, Spreng. ex Trin. in Mém. Acad. Pétersb. Sér VI. iii. ii. (1835) 128 (= *Stipa Caryochloa*).
monteindesia, Spreng. Syst. iv. Cur. Post. 80 (= *Stipa setifolia*).—Reg. Argent.
refracta, Spreng. ex Trin. in Mém. Acad. Pétersb. Sér VI. iii. ii. (1835) 124 (= *Stipa Caryochloa*).

CARYOCHLOA, Trin. Gram. Panic. Diss. ii. 54 (1826) = **Luziola**, Juss. (Gramin.).
bahiensis, Steud. Syn. Pl. Gram. 5.—Bras.
brasiliensis, Trin. ex Nees, Agrost. Bras. 399.—Bras.
caudata, Trin. ex Steud. Nom. ed. II. i. 301.—Bras.

CARYOCOCCA, Willd. ex Roem. & Schult. Syst. iii. Mant. 181 (1827) = **Gonzalea**, Pers. (Rubiac.).
dependens, Willd. ex Roem. & Schult. l. c. = G. pendula.
multiflora, Willd. ex Roem. & Schult. l. c. 182 = G. pendula.

CARYODAPHNE, Blume, ex Nees, Syst. Laurin. 225 (1888) = **Cryptocarya**, R. Br. (Laurin.).
australis, A. Br. Ind. Sem. Hort. Berol. (1851) App 18 = Crypt. australis.
Brownsiana, Nees, Syst. Laurin. 230 = Crypt. triplinervis.
costata, Miq. in Zoll. Syst. Verz. Ind. Archip. 113, 115 = Crypt. costata.
densiflora, Blume, ex Nees, Syst. Laurin. 228 = Crypt densiflora.
laevigata, Blume, l. c. 227 = Crypt. laevigata.

CARYODENDRON, Karst. Fl. Colomb. i. 91. t. 46 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 314.
CENTRODISCUS, Muell. Arg. in Mart. Fl. Bras. xi. ii. 325. t. 102 (1874).
orinocense, *Karst. l. c.*—N. Granat.

CARYOLOBIS, Gaertn. Fruct. i. 215. t. 45 (1788) = **Doona**, Thw. (Dipteroc.).
indica, Gaertn. l. c. = D. cordifolia!

CARYOLOBIUM, Stev. in Bull. Soc. Nat. Mosc iv (1832) 266 = **Astragalus**, Tourn. (Legumin.).
macrocarpos, Stev. l. c. = A. macrocarpus.

CARYOLOPHA, Fisch. & Trautv. Ind. Sem. Hort Petrop. iii. 31 (1837) = **Anohusa**, Lindn. (Boragin.).
semperviren, Fisch. & Trautv. l. c. = A. semperviren

CARYOPHYLLATA, Tourn. ex Scop. Fl. Carn. ed. II. i. 364 (1772) = **Geum**, Tourn. (Rosac.).
alba, Moench, Meth. 660 = G. album.
aquatica, Lam. Fl. Fr. iii. 123 = G. rivale.
hamtchatica, Lam. Encyc. i. 400 = G. anemonoides.
montana, Scop. Fl. Carn. ed. II. i. 365 = G. montanum
nutans, Lam. Encyc. i. 399 = G. nutans.
nutans, Moench, Meth. 661 = G. rivale.
obliqua, Moench, l. c. Suppl. 280 = G. obliquum.
officinalis, Moench, Meth. 660 = G. urbanum.
potentillaides, Lam. Encyc. i. 400 = Colaria geoides.
prolifera, Gilib. Fl. Lituan. 416.—Europ.
reptans, Lam. Fl. Fr. iii. 122 = G. reptans.
rivalis, Scop. Fl. Carn. ed. II. i. 365 = G. rivale.
urbana, Scop. l. c. 364 = G. urbanum.
virginiana, Lam. Encyc. i. 399 = G. virginianum.
vulgaris, Lam. Fl. Fr. iii. 122 = G. urbanum.

CARYOPHYLLEA, Opiz, Seznam, 27 (1852) = **Aira**, Linn. (Gramin.).
airoides, Opiz, l. c. = A. caryophyllea.
praecox, Opiz, l. c. = A. praecox.

CARYOPHYLLUS, Linn. Syst. ed. I (1735) = **Eugenia**, Linn. (Myrtac.).
acutangulus, Stokes, Bot. Mat. Med. iii. 75 = Barringtonia acutangula.
aeneolepis, Miq. Fl. Ind. Bat. i. 466.—Java.
airoides, Blume, ex DC. Prod. iii. 262.—Java.
aromaticus, Linn. Sp. Pl. 515 = E. caryophyllata.
balsamea, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 186.—N. Caled.
corticosus, Stokes, Bot. Mat. Med. iii. 76 = E. Jambolana.

CARYOPHYLLUS :—
cotinifolius, Mill. Gard. Dict. ed. VIII. n. 4 = Myrcia cotinifolia.
elegans, Bronge. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 184.—N. Caled.
ellipticus, Labill. Sert. Anast. Caled. 64. t. 68.— N. Caled.
fastigiatus, Blume, ex DC. Prod. iii. 269.—Java.
floribundus, Blume, ex DC. l. c.—Java.
fruticosus, Mill. Gard. Dict. ed. VIII. n. 3 = E. biflora.
Gomboi, Stokes, Bot. Mat. Med. iii. 73 = E. Jambos.
hortensis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 11 = E. caryophyllata !
malaccensis, Stokes, Bot. Mat. Med. iii. 72 = E. malaccensis.
Pimenta, Mill. Gard. Dict. ed. VIII. n. 2 = Pimenta officinalis.
pterocarpus, Vieill. ex Panch. & Sebert, Not. Bois. Nouv. Caléd. 256.—N. Caled.
racemosus, Mill. Gard. Dict. ed. VIII. n. 5 = E. acris.
racemosus, Stokes, Bot. Mat. Med. iii. 76 = Barringtonia racemosa.

CARYOPHYLLUS, Tourn. ex Moench, Meth. 58 (1794) = **Dianthus**, Linn. (Caryophyll.)
aridus, Moench, Meth. 59 = D. prolifer
armerius, Moench, l. c. = D. Armeria.
atrorubens, Moench, Meth. Suppl. 28 = D. atrorubens.
barbatus, Moench, Meth. 59 = D. barbatus.
Carthusianorum, Moench, l. c. = D. Carthusianorum.
deltoides, Moench. l. c. = D. deltoides.
glaucus, Moench, l. c. = D. deltoides.
plumarius, Moench, l. c. = D. plumarius.
superbus, Moench, l. c. = D. superbus.

CARYOPTERIS, Bunge, Pl. Mongholico-Chin. 27 (1835). *VERBENACEÆ*, Benth. & Hook. f. ii. 1157.
HARBULA, Lour. Fl. Cochinch. 366 (1790), nomen prius.
MASTACANTHUS, Endl. Gen. 638 (1839).
divaricata, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—Japon.
grata, *Benth. & Hook. f. Gen.* ii. 1158.—Reg. Himal.
incana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 97.—Japon.
nepetæfolia, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—China.
ovata, *Miq. in Journ. Bot. Néerl.* i. (1861) 114.—China.
paniculata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 597.—Reg. Himal.
tangutica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 525.—China.
ternifiora, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 40.—China.
Wallichiana, *Schau. in DC. Prod.* xi. 625.— Reg. Himal.

CARYOSPERMUM, Blume, Mus. Bot. Lugd. Bat. i. 178 (1850). *CELASTRINEÆ*, Benth. & Hook. f. i. 367.
arborescens, *F. Muell. Fragm.* viii. 279.—Austral.
moluccanum, *Blume, Mus. Bot. Lugd. Bat.* i. 176.— Ins. Molluc.
serrulatum, *Miq. Fl. Ind. Bat.* i. ii. 592.—Java.

CARYOTA, Linn. Mus. Cliff. 12 ; et Gen. ed. I. 355 (1737). *PALMÆ*, Benth. & Hook. f. iii. 918.
Alberti, F. Muell. ex H. Wendl. in Linnæa, xxxix. (1876) 321 = Rumphiana.
Cavingi, *Ludd. ex Mart. Hist. Nat. Palm.* iii. 215.— Ins. Philipp.
elegans, *Hort. ex H. Wendl. in Kerch. Palm.* 288, nomen.—Hab. ?
excelsa, *Hort. ex H. Wendl. l. c.* nomen.—Hab. ?
furfuracea, Blume, in Mart. Hist. Nat. Palm. iii. 195 = mitis.
Griffithii, Becc. in Nuov. Giorn. Bot. iii. (1871) 15 = mitis.
horrida, Jacq. Fragm. 20 = Bactris caryotæfolia.

CARYOTA :—
humilis, Reinw. ex Kunth, Enum. Pl. iii. 193 = Didymosperma porphyrocarpa.
javanica, Zipp. ex Miq. Fl. Ind. Bat. iii. 41 = mitis.
majestica, *Linden, Illustr. Hortic.* xxviii. (1881) 16.— Ins. Philipp.
maxima, *Blume, in Mart. Hist. Nat. Palm.* iii. 195.— Java.
mitis, *Lour. Cochinch.* 697.—Malaya ; Cochinch.
nana, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Ins. Philipp.
nana, Wall. Cat. n. 8595 = mitis.
Nó, *Becc. in Nuov. Giorn. Bot.* iii. (1871) 12.— Borneo.
obtusa, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 480.—Reg. Himal.
obtusidentata, Griff. Palms Brit. Ind. t. 236 A, B = pinnc.
ochlandra, Hance, in Journ. Bot. xvii. (1879) 174 = obtusa !
ovata, Blanco, Fl. Filip. ed. I. 741 = Arenga saccharifera.
Palindan, Blanco, l. c. ed. I. 714 ; ed. II. 513 = Orania regalis.
Princeps, *Vieyt. in Syll. Ratisb.* ii. (1828) 51.— Hab. ?
propinqua, Blume, in Mart. Hist. Nat. Palm. iii. 195 = mitis.
Rumphiana, *Mart. l. c.*—Malaya ; Austral.
sobolifera, Wall. in Mart. l. c. 194. t. 107. f. 2 = mitis.
speciosa, *Linden, Illustr. Hortic.* xxviii. (1881) 16.— Ins. Philipp.
tremula, Blanco, Fl. Filip. ed. I. 744 = Didymosperma tremula.
urens, Blanco, l. c. 740 = Cumingii.
urens, Jacq. Fragm. 20. t. 12. f. 1 = mitis.
urens, *Linn. Sp. Pl.* 1189.—Ex. trop.

CARYOTAXUS, Zucc. ex Endl. Syn. Conif. 240 (1847) = **Torreya**, Arn. (Conif.).
grandis, Henk. & Hochst. Nadelhoelz. 367 = T. grandis.
Myristica, Henk. & Hochst. l. c. 368 = T. californica.
nucifera, Zucc. ex Endl. Syn. Conif. 241 = T. nucifera.
taxifolia, Henk. & Hochst. Nadelhoelz. 367 = T. taxifolia.

CASALEA, St. Hil. Fl. Bras. Mer. i. 8. t. 1 (1825) = **Ranunculus**, Tourn.
ascendens, St. Hil. l. c. 11 = R. cordifolius.
biternata, St. Hil. l. c. 9, in obs. = R. biternatus.
breviscapa, St. Hil. l. c. = R. breviscapus.
chilensis, St. Hil. l. c. = R. chilensis.
ficariifolia, St. Hil. l. c. 10 = R. bonariensis.
flagelliformis, St. Hil. l. c. 11 = R. flagelliformis.
hederacea, Mart. ex Eichl. in Mart. Fl. Bras. xiii. i. 154 = R. sessiliflorus.
phyteumæfolia, St. Hil. Bras. Mer. i. 9 = R. bonariensis.
sessiliflora, St. Hil. l. c. = R. platensis.

CASARETTOA, Walp. Rep. iv. 91 (1844) = **Vitex**, Linn. (Verbenac.).
divaricata, Walp. l. c. 92 = V. trifolia.
mollissima, Walp. l. c. 91 = V. polygama.

CASASIA, A. Rich. Fl. Cub. Fanerog. ii. 9. t. 49 (1858). *RUBIACEÆ*, Benth. & Hook. f. ii. 84. calophylla, *A. Rich. l. c.*—Cuba.

CASCABELA, Raf. Sylva Tellur. 162 (1838) = **Cerbera**, Linn. (Apocyn.).
peruviana, Raf. l. c. = Cerb. peruviana.

CASCARILLA, Adans. Fam. ii. 355 (1763) = **Croton**, Linn. (Euphorbiac.).
aromatica, Raf. Sylva Tellur. 62 = Crot. aromaticus.
discolor, Raf. l. c. = Crot. discolor.
linearis, Raf. l. c. = Crot. linearis.
officinalis, Raf. l. c. = Crot. Cascarilla.

CASCARILLA, Ruiz, ex Steud. Nom. ed. I. 165 (1821) = Cinchona, Linn. (Rubiac.).
pallida, Steud. l. c. = Cinch. cordifolia.

CASCARILLA, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 10, 77. *RUBIACEÆ*, Benth. & Hook. f. ii. 83.

CASCARILLA :—
BUENA, Pohl, ex Wedd. in Journ. Linn. Soc. xi. (1871) 185.
LADENBERGIA, Klotzsch, in Hayne, Arzn. Gew. xiv. in not. ad t. 15 (1846).
acutifolia, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 11.—Peruv.
bullata, *Wedd. l. c.* 11.— Peruv.
caducifiora, *Wedd. l. c.* 10.—Peruv.
calycina, *Wedd. l. c.* 12.—N. Granat.
calyptrata, *Wedd. l. c.* 13.—Peruv.
Carua, *Wedd. l. c.* 12.—Peruv.
citrifolia, *Wedd. l. c.* 11.—Venezuela.
Gaudichaudiana, *Wedd. l. c.* 12.—Bras.
gavanensis, *Schlecht. in Linnæa*, xxvi. (1853) 730 — Peruv.
grandiflora, *Linden, in Belg. Hortic.* vii. (1857) 197. —N. Granat.
heterophylla, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 10.—N. Granat.
hexandra, *Wedd. l. c.* 13.—Bras.
Hookeriana, *Wedd. Hist. Nat. Quinq.* 91.—N. Granat. Lambertiana, *Wedd. l. c.* 85.— Bras.
macrocarpa, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 18.—N. Granat. ; Peruv.
magnifolia, *Wedd. l. c.* 10.—Bolivia.
muzonensis, *Wedd. l. c.* 11.—N. Granat.
nitida, *Wedd. l. c.* 11.—Peruv.
oblongifolia, Wedd. l. c. 12 = magnifolia.
officinalis, *Ruiz, Quinol.* 56.—Peruv.
Pavonii, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 11. —Peruv.
Riedeliana, *Wedd. l. c.* 12.—Bras.
Riveroana, *Wedd. l. c.* 11.—Peruv.
Rorainæ, *Wedd. l. c.* 12.—Guiana.
rostrata, *Wedd. l. c.* 10.—Boliv. ; Peruv.
stenocarpa, *Wedd. Hist. Nat. Quinq.* 81.—Peruv.
undata, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 12.—Venezuela.

CASCARONIA, Griseb. in Goett. Abh. xxiv. (1879) 100. *LEGUMINOSÆ*.
astragalina, *Griseb. l. c.*—Reg. Argent.

CASCOELYTRUM, Beauv. Agrost. 85 (1812) = CHASCOLYTRUM, Desv. = **Briza**, Linn. (Gramin.).

CASEARIA, Jacq. Enum. Pl. Carib. 4 (1760). *SAMYDACEÆ*, Benth. & Hook. f. i. 796.
ANAVINGA, Lam. Encyc. i. 147 (1783).
ANTIGONA, Vell. Fl. flum. iv. t. 145 (1827).
ATHENÆA, Schreb. Schreb. Gen. i. 239 (1789).
BEDUSIA, Rafin. Sylva Tellur. 11 (1838).
CHAETOCRATER, Ruiz & Pav. Prod. Fl. Per. 61. t. 26 (1794).
CHETOCRATER, Rafin. Sylva Tellur. 149 (1838).
CLASTA, Comm. ex Vent. Choix, sub t. 47 (1808).
CORIZOSPERMUM, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 255 (1850).
CRATERIA, Pers. Syn. i. 485 (1805).
GUIDONIA, Griseb. Fl. Brit. W. Ind. 24 (1864).
IROUCANA, Aubl. Pl. Gui. i. 328. t. 127 (1775).
LANGLEIA, Scop. Introd. 231 (1777).
LINDLEYA, Kunth, Malv. 10 (1822).
MELISTAURUM, Forst. Char. Gen. 143. t. 72 (1776).
MOELLERIA, Scop. Introd. 235 (1777).
PIPAREA, Aubl. Pl. Gui. ii. App. 30. t. 366 (1775).
PITUMBA, Aubl. l. c. 29. t. 385 (1775).
THIGIA, Griseb. Fl. Brit. W. Ind. 22 (1864).
VALENTINIA, Sw. Prod. Veg. Ind. Occ. 63 (1788).
VAREA, Gaertn. Fruct. i. 290. t. 60 (1788).
ZORLANIA, A. Rich. Ess. Fl. Cub. 86. t. 13 (1845).
aculeata, Jacq. Enum. Pl. Carib. 21 = spinosa.
acuminata, *DC. Prod* ii. 50.—Guiana.
acuminata, Wall. Cat. n. 7198 (non Casearia).— Ind. or.
acuminatissima, *A. Gray, Bot. U. St. Expl. Exped.* i. 80.—Ins. Fiji.
adamantium, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 230.—Bras.
adstringens, *Mart. Mat. Med. Veg. Bras.* 51.—Bras.
affinis, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 229 = sylvestris.
alba, *A. Rich. Ess. Fl. Cub.* 86.—Cuba.
albicans, *Wall. Cat.* n. 7197.—Malaya.
amplissima, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 237.—Madag.
Anavinga, Dalz. & Gibs. Bomb. Fl. 11 = tomentosa.

CASEARIA :—
Anovinga, Pers. Syn. i. 485 = ovata.
angustata, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1856) 400.—Java.
aquifolia, *Sauv. Fl. Cub.* 6.—Cuba.
arguta, H. B. & K. Nov. Gen. et Sp. v. 363 = parvifolia.
astyla, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 608.—Ind. or.
axillaris, Miq. ex Eichl. in Mart. Fl. Bras. xiii. 1. 463 = spinosa.
Barteri, *Mast. in Oliver, Fl. Trop. Afr.* ii. 494.—Afr. trop.
Benthamiana, *Miq. in Ann. Sc. Nat.* Sér. III. i. (1844) 38.—Bras.
Berteriana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 461.—Porto Rico.
Blanchettiana, *Miq. in Linnæa*, xxii. (1849) 801.—Bras.
brasiliensis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 477.—Bras.
brevipes, *Benth. in Hook. Journ. Bot.* iv. (1842) 110.—Guiana; Ind. occ.
Cambessedesii, *Eichl. in Mart. Fl. Bras.* xiii. 1. 475.—Bras.
Canaiala, Buch.-Ham. in Wall. Cat. n. 7192 = tomentosa.
capitata, Spreng. Syst iii. 855 = sylvestris.
capitellata, *Blume, Mus. Bot. Lugd. Bat.* i. 254.—Borneo.
carpinifolia, Benth. in Hook. Journ. Bot. iv. (1842) 112 = parvifolia.
celastroides, *Klotzsch, in Schomb. Faun. et Fl. Guy.* 1167, nomen.—Guiana.
celtidifolia, *H. B. & K. Nov. Gen. et Sp.* v. 362.—Venezuela.
Championi, Thw. Enum. Pl. Zeyl. 19 = esculenta.
cinerea, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 462 = grewiæfolia.
Clausseniana, *Miq. in Linnæa*, xix. (1847) 441.—Bras.
clutiæfolia, *Blume, Mus. Bot. Lugd. Bat.* i. 255.—N. Guinea.
combeyanensis, *Tul in Ann. Sc. Nat.* Sér. III. vii. (1847) 362.—N. Granat.
Commersoniana, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 235.—Bras.
comocladifolia, *Vent. Choix*, t. 44.—Ins. S. Doming.
congestiflora, *Turcz. in Bull Soc. Nat. Mosc.* xxxvi. (1863) i. 608.—Bras.
conterminus, *Miq. Fl. Ind. Bat.* i. i. 707.—Java.
coriacea, *Thw. Enum. Pl. Zeyl.* 20.—Zeylan.
coriacea, *Vent. Choix*, t. 45.—Java.
coriacea, Wall. Cat. n. 7196 = Chætocarpus castanocarpus.
corymbosa, *H. B. & K. Nov. Gen. et Sp.* v. 366.—N. Granat.
crapidata, *Blume, Mus. Bot. Lugd. Bat.* i. 255.—Sumatra.
Dallachii, F. Muell. Fragm. v. 107 = tomentosa.
decandra, Jacq. Enum. Pl. Carib. 31 = parvifolia.
denaifiora, *Benth. in Hook. Journ. Bot.* iv. (1842) 113.—Guiana.
dentata, *Miq. & Sesse, in DC. Prod.* ii. 51.—Mexic.
Hatschek, A. Gray, Bot. U. St. Expl. Exped. i. 81 = C. Melistaurum.
dubia, *Miq. & Sesse, in DC. Prod.* ii. 51.—Mexic.
elliptica, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1167, nomen.—Guiana.
elliptica, Willd. Sp. Pl. ii. 938 = tomentosa.
emarginata, *Griseb. Cat. Pl. Cub.* 10.—Cuba.
eriobotrya, *Wright, ex Griseb. l. c.* 11.—Cuba.
esculenta, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 422.—As. trop.; Austral.
fallax, *Miq. in Linnæa*, xix. (1847) 128.—Guiana.
fascivulata, Boj. Hort. Maurit. 71 = frangila.
flavoviresa, *Blume, Mus. Bot. Lugd. Bat.* i. 254.—Java.
Fockeana, *Miq. in Ann. Sc. Nat.* Sér. III. i. (1844) 39.—Guiana.
fragilis, *Vent. Choix*, 47, verso.—Ins. Mascar.
fuliginosa, *Blanco, Fl. Filip.* ed. II. 262.—Ins. Philipp.
glabra, Kurz, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 601 = glomerata.
glabra, Roxb. Hort. Beng. 88; Fl. Ind. ii. 421 = tomentosa.
glabrata, *Blume, Mus. Bot. Lugd. Bat.* i. 253.—Java.

CASEARIA :—
gladiiformis, *Mast. in Oliver, Fl. Trop. Afr.* ii. 493.—Afr. trop.
glomerata, *Roxb. Hort. Beng.* 88; *Fl. Ind.* ii. 419.—Ind. or.
gonocarpa, *Miq. Fl. Ind. Bat.* i. i. 1093.—Sumatra.
grandiflora, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 232.—Bras.
grandifolia, *Miq. Fl. Ind. Bat.* i. i. 712.—Sumatra.
grewiæfolia, *Dals. in Hook. Kew Journ.* iv. (1852) 107.—Ind. or.
grewiæfolia, *Vent. Choix*, 48.—Malaya.
Guidonia, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 89 = brevipes.
guineensis, *G. Don, Gen. Syst.* ii. 53.—Afr. trop.
Hamiltonii, Wall. Cat. n. 7195 = graveolens.
hexagona, *Decne. in Ann. Nouv. Mus. Par.* iii. (1834) 429.—Ins. Timor.
hirsuta, *Sw. Fl. Ind. Occ.* ii. 755.—Ind. occ.
hirta, *Sw. l. c.* 756.—Ind. occ.
Hostmanniana, Steud. ex Griseb. Fl. Brit. W. Ind. 23 = stipularis.
hypoleuca, *Mart. in Flora*, xx. (1837) *Beibl.* ii. 128.—Bras.
icosandra, *Planch. & Triana, ex Hemsl. Biol. Centr. Am. Bot.* i. 469.—Panama.
illicifolia, *Vent. Choix*, t. 44.—Ins. S. Doming.
impunctata, *Hook. & Arn. Bot. Beech. Voy.* 61.—Ins. Societat.
inaequilatera, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 237.—Bras.
integrifolia, Vahl, ex DC. Prod. ii. 49 = sylvestris.
javitensis, *H. B. & K. Nov. Gen. et Sp.* v. 366.—Am. austr.
Kunzii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 594.—Burma.
lanceolata, Dalz. in Hook Kew Journ. iv. (1852) 108 = esculenta.
lanceolata, Miq. in Linnæa, xviii. (1844) 758 = stipularis.
laniugula, *Eichl. in Mart. Fl. Bras.* xiii. 1. 468.—Bras.
laxiosperma, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii (1862) 113.—N. Granat.
laurifolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 113.—Guiana.
laurina, *Blume, Mus. Bot. Lugd. Bat.* i. 253.—Malaya.
leucoclepus, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 468.—Malaya.
Lingua, Cambess. in St. Hil. Fl. Bras. Mer. ii. 236 = sylvestris.
Lobbiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 463.—Burma; Malaya.
lucida, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 325.—Madag.
lucida, Wall. Cat. n. 7199 (Euphorbiac.).—Ind. or.
macrocarpa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 598.—Penang.
macrogyna, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 463 = graveolens.
macrophylla, *Vahl, Eclog. Am.* ii. 32.—Guiana.
Mannii, *Mast. in Oliver, Fl. Trop. Afr.* ii. 494.—Afr. trop.
mariquitensis, *H. B. & K. Nov. Gen. et Sp.* v. 365.—N. Granat.
Mathewsii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 608.—Peruv
Maximiliana, *Eichl. in Mart. Fl. Bras.* xiii. 1. 467.—Bras.
meliodora, *Eichl. l. c.* 469.—Bras.
Melistaurum, *Spreng. Syst.* ii. 854.—Ins. Pacif.
membranacea, *Hance, in Journ. Bot.* vi. (1868) 113.—China.
micrantha, G. Don, Gen. Syst. ii. 52 = parvifora.
microdon, *Miq. Fl. Ind. Bat. Suppl.* 338.—Sumatra.
microphylla, *Eichl. in Mart. Fl. Bras.* xiii. 1. 474.—Bras
mollis, H. B. & K. Nov. Gen. et Sp. v. 363. i. 480 = hirsuta.
molucensis, *Blume, Mus. Bot. Lugd. Bat.* i. 255.—Amboina.
montana, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 385 = lanigulifera.
multiflora, *Spreng. Pugill.* ii. 61.—Ind. or.
myrianthus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 609.—Panama.
negrensis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 466.—Bras.
nigrescens, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 326.—Madag.

CASEARIA :—
nitida, *Jacq. Enum. Pl. Carib.* 21.—N. Granat.
nitida, Sieber, ex Griseb. Fl. Brit. W. Ind. 23 = parvifolia.
obliqua, Spreng. Syst. ii. 855 = inaequilatera.
oblongifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 234.—Bras.
obovalis, Poepp. ex Griseb. in Goett. Abh. ix. (1860) 27 = obovata.
obovata, *Poepp. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 472.—Peruv.
obovata, *Schlecht. in Linnæa*, xiii. (1839) 434.—Mexic.
odorata, Macfad. Fl. Jamaic. i. 216 = ramiflora.
odorata, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 408.—Java.
oligantha, *Eichl. in Mart. Fl. Bras.* xiii. 1. 476.—Bras.
ovata, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 420 = tomentosa.
ovata, Wall. Cat. n. 7192 E = glomerata.
ovata, *Willd. Sp. Pl.* ii. 629.—Ind. or.
paniculata, Gardn. ex Thw. Enum. Pl. Zeyl 20 = Osmelia zeylanica.
parvifora, J. F. Gmel. Syst. 700 = parvifolia.
parvifora, Willd. Sp. Pl. ii. 627 = sylvestris.
parvifolia, *Willd. l. c.* 628.—Ind. occ.
pauciflora, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 235.—Bras.
petiolaris, *Poepp. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 471.—Peruv.
petraea, *Benth. in Hook. Journ. Bot.* iv. (1842) 111.—Guiana.
Pittumba, DC. Prod. ii. 50, in syn. = macrophylla.
Poeppigii, *Eichl. in Mart. Fl. Bras.* xiii. 1. 475.—Bras.
Pohliana, *Eichl. l. c.* 474.—Bras.
praecox, *Griseb. Cat. Pl. Cub.* 10.—Cuba.
prismatocarpa, *Mast. in Oliver, Fl. Trop. Afr.* ii. 494.—Afr. trop.
propinqua, *Blume, Mus. Bot. Lugd. Bat.* i. 253.—Java.
prunifolia, *H. B. & K. Nov. Gen. et Sp.* v. 362.—Am. austr.
prunifolia, Tul. in Ann. Sc. Nat. Ser. III. vii. (1847) 363 = spinosa.
pubiflora, Benth. Bot Voy. Sulph. 66 = parvifolia.
punctata, Spreng. Neue Entdeck. ii. 154 = parvifolia.
punctata, *Spreng. Syst.* ii. 855.—Am. trop.
quindiuensis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 360.—Ecuador.
ramiflora, Seem. Bot. Voy. Herald, 98 = hirsuta.
ramiflora, *Vahl, Symb. Bot.* ii. 50.—Am. trop.
ramosissima, *Griseb. Cat. Pl. Cub.* 10.—Cuba.
reclinifera, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiii. 1. 466.—Bras.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 82.—Ins. Fiji.
rubescens, *Dals. in Hook. Kew Journ.* iv. (1852) 108.—Ind. or.
rufescens, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 231.—Bras.
ruiginula, *Triana & Planch. in Ann. Sc. Nat.* Ser. IV xvii. (1862) 107.—N. Granat.
rufinervis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 462.—Cuba.
rugulosa, *Blume, Mus. Bot. Lugd. Bat.* i. 255.—Malaya.
rupestris, *Eichl. in Mart. Fl. Bras.* xiii. 1. 468.—Bras.
salacioides, *Blume, Mus. Bot. Lugd. Bat.* i. 252.—N. Guinea.
salicifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 460.—Mexic.
Samyda, *DC. Prod.* ii. 51.—Porto Rico.
Selloana, *Eichl. in Mart. Fl. Bras.* xiii. 1. 468.—Bras.
serrata, Macfad. Fl. Jamaic. i. 215 = serrulata.
serrulata, Sieber, ex Griseb. Fl. Brit. W. Ind. 23 = parvifolia.
serrulata, *Sw. Fl. Ind. Occ.* ii. 754.—Ind. occ.
sessiliflora, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 231.—Bras.
singularis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 473.—Guiana.
spinescens, *Griseb. Cat. Pl. Cub.* 10.—Cuba.
spinosa, Willd. Sp. Pl. ii. 626 = ramiflora.
Spruceana, *Benth. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 466.—Bras.

CASEARIA:—
squarrosa, *Klotzsch, in Schomb. Faun. et Fl. Guy.* 1167, nomen.—Guiana.
stipulata, *Mast. in Oliver, Fl. Trop. Afr.* ii. 498.—Afr. trop.
stipularis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 169 = brasiliensis.
stipularis, *Vent. Choix*, 46. t. 46.—Am. austr.
subcillaris, *Klotzsch, in Schomb. Faun. et Fl. Guy.* 984, nomen.—Guiana.
subcuneata, Miq. Fl. Ind. Bat. i. 706 = grewiaefolia.
subopaca, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xvii. (1862) 110.—N. Granat.
suberbombes, *Hance, in Journ. Bot.* xxiii. (1885) 328. —China.
sylvestris, *Sw. Fl. Ind. Occ.* ii. 752.—Ind. occ.
tinifolia, *Vent. Choix*, 47. t. 47.—Hab.?
tomentosa, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 421.—As. trop.; Austral.
truncata, *Blume, Mus. Bot. Lugd. Bat.* i. 252.— Malaya.
tuberculata, *Blume, l. c.* 254.—Java.
tuberculata, *Blume, l. c.* 255.—Sumatra.
wimifolia, Cambess. in St. Hil. Fl. Bras. Mer. ii. 169 = Cambessedesii.
wimifolia, Vahl, ex Vent. Choix, 46, verso = serrulata.
uniflora, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 428.—Ins. Timor.
Vareca, *Roxb. Hort. Beng.* 33; *Fl. Ind.* ii. 418.—Reg. Himal.; Burma.
variabilis, Blume, Mus. Bot. Lugd. Bat. i. 252 = grewiaefolia.
varians, Thw. Enum. Pl. Zeyl. 19 = esculenta.
velutina, *Blume, Mus. Bot. Lugd. Bat.* i. 253.— Sumatra.
viridiflora, *DC. Prod.* ii. 51.—Ind. or.
viridiflora, Lam. Encyc. vi. 499 = lencolepis.
wynaadensis, *Bedd. Ic. Pl. Ind. Or.* i. 34.—Ind. or.
zeylanica, Thw. Enum. Pl. Zeyl. 19 = esculenta.
zizyphoides, *H. B. & K. Nov. Gen. et Sp.* v. 862.— Am. austr.

CASEOLA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

CASIMIRA, Scop. Introd. 288 (1777) = **Melioocca**, Linn. (Sapind.).

CASIMIROA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 2 (1825). *RUTACEAE*, Benth. & Hook. f. i. 302.
edulis, *La Llave, l. c.*—Mexic.
Sapota, *Oerst. in Kjoerh. Vidensk. Meddel.* (1857) 187. —Nicaragua.
tomentosa, Lomb. ex Miers, in Journ. Linn. Soc. xvii. (1880) 78 = Cervantesia tomentosa.

CASINGA, Griseb. in Goett. Abh. ix. (1861) 27, 29 = **Laetia**, Linn. (Cochl.).
procera, Griseb. Fl. Brit. W. Ind. 710 = L. procera.
suaveolens, Griseb. in Goett. Abh. ix. (1861) 27 = L. suaveolens.

CASIOSTEGA, Rupr. ex Galeotti, in Bull. Acad. Brux. ix. 11. (1842) 325, nomen = **Opizia**, Presl (Gramin.).
anomala, Rupr. l. c. = O. stoloniféra.
dactyloides, Fourn. in Bull. Soc. Bot. Belg. xv. (1876) 470 = Buchloë dactyloides.
Hookeri, Rupr. ex Fourn. l. c. 471 = Buchloë dactyloides.
humilis, Rupr. [ex Munro] in Benth. Pl. Hartw. 347 = Buchloë dactyloides.

CASPAREA, H. B. & K. Nov. Gen. et Sp. vi. 317 (1823) = **Bauhinia**, Linn. (Leguminos.).
acuminata, Heynh. Nom. ii. 132 = B. acuminata.
aurita, Griseb. Fl. Brit. W. Ind. 213 = B. aurita.
castrata, Nimsk. Cat. Hort. Bogor. 288 = B. castrata.
divaricata, H. B. & K. Nov. Gen. et Sp. vi. 317 = B. divaricata.
furcata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 429 = B. furcata.
latifolia, H. B. & K. Nov. Gen. et Sp. vi. 317 = B. latifolia.
lunaria, H. B. & K. l. c. = Bauhinia lunaria.
Pea-capraea, H. B. & K. l. c. 318 = B. Pea-capraea.
porrecta, H. B. & K. l. c. 317 = B. porrecta.
rubrotundifolia, H. B. & K. l. c. = B. rubrotundifolia.

CASPARVA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 137 = **Begonia**, Linn.
anthiopaeuala, A. DC. in Ann. Sc. Nat. Ser. IV. xi. (1859) 116.—N. Granat.
brevipetala, A. DC. l. c. 118.—Venezuela.
coccinea, Klotzsch, in Monatsb. Berl. Acad. (1854) 127 = B. columnaris.
columnaris, Klotzsch, l. c. = B. columnaris.
cordifolia, A. DC. in DC. Prod. xv. 1. 278 = *hirta*.
crassicaulis, A. DC. in Ann. Sc. Nat. Ser. IV. xi. (1859) 119.—Java.
elegans, Klotzsch, in Monatsb. Berl. Acad. (1854) 127 = B. elegans.
erosa, A. DC. in DC. Prod. xv. 1. 276 = B. erosa.
ferruginea, A. DC. l. c. 269 = B. ferruginea.
fuchsioyflora, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 116.—Ecuador.
grewiaefolia, A. DC. l. c. 117.—N. Granat.
hirta, Klotzsch, in Monatsb. Acad. Berl. (1854) 127.— Peru.
longirostris, A. DC. in DC. Prod. xv. 1. 272 = B. longirostris.
montana, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 118.—Venezuela.
multiangula, A. DC. in DC. Prod. xv. 1. 275 = B. multiangula.
oligocarpa, A. DC. in Ann. Nat. Ser. IV. xi. (1859) 118 = B. Roxburghii.
polycarpa, A. DC. l. c. = B. Roxburghii.
robusta, A. DC. in DC. Prod. xv. 1. 275 = B. robusta.
silhetensis, A. DC. l. c. 277 = B. silhetensis.
Teysmanniana, Miq. ex A. DC. l. c. 276.—Sumatra.
trachyptera, A. DC. l. c. 274 = B. trachyptera.
Triana, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 117.—N. Granat.
tripathulata, A. DC. l. c.—N. Granat.
ternicalata, A. DC. l. c. 119.—Java.
umbellata, A. DC. in DC. Prod. xv. 1. 270 = B. umbellata.
Urticae, A. DC. l. c. 274 = B. Urticae.

CASPIA, Pison. ex Scop. Introd. 276 (1777) = **Vismia**, Vell. (Hypericin.).

CASSANDRA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 158. *ERICACEAE*, Benth. & Hook. f. ii. 584.
DIPLARIA, Rafin. ex DC. Prod. vii. 610 (1839).
angustifolia, D. Don, l. c. = calyculata.
calyculata, *D. Don, l. c.* 158.—Europ.; As. et Am. bor.
racemosa, Spach, Hist. Vég. Phan. ix 478 = Lecothoe racemosa.

CASSEBEERIA, Dennst. Schluss. Hort. Malab. 35 (1818) = **Sonerila**, Roxb. (Melast.).
maculata, Dennst. l. c. = S. Wallichii.

CASSELIA, Dum. Comm. Bot. 21 (1822) = **Mertensia**, Roth (Boragin.).
bracteata, Dum. l. c. 25 = M. sibirica.
dawurica, Dum. l. c. 22 = M. dahurica.
denticulata, Dum. l. c. 23 = M. sibirica.
diversifolia, Regel, Ind. Sem. Hort. Petrop. (1857) 46. —Hab.?
gracilis, Dum. Comm. Bot. 23 = M. dahurica.
lanceolata, Dum. l. c. 24 = M. lanceolata.
maritima, Dum. l. c. 25 = M. maritima.
paniculata, Dum. l. c. 23 = M. paniculata.
parviflora, Dum. l. c. 25 = M. maritima.
sibirica, Dum. l. c. 23 = M. sibirica.
simplicissima, Dum. l. c. 24 = M. simplicissima.
villosula, Dum. l. c. 23 = M. villosula.
virginica, Dum. l. c. 24 = M. pulmonarioides.

CASSELIA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 74 t. 6. *VERBENACEAE*, Benth. & Hook. f. ii. 1148.
chamaedryfolia, *Cham. in Linnaea*, vii. (1832) 565.— Bras.
integrifolia, *Nees & Mart. in Nov. Act. Acad. Nat. Cur.* xi. (1823) 76.—Bras.
Mansoi, *Schau. in Mart. Fl. Bras.* ix. 175.—Bras.
serrata, *Nees & Mart. in Nov. Act. Acad. Nat. Cur.* xi. (1823) 75.—Bras.
veronicaefolia, *Cham. in Linnaea*, vii. (1832) 564.— Bras.

CASSIA, Tourn. ex Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 571.
ADIPERA, DIALANTHERA, DIALLOBUS, DISTEREPTA, DIPLOTAX, DITREMEXA, EMELISTA, HEPTHIRECA, HERPETICA, ISANDRINA, NICTITELLA, OCTELISIA, OPHIOCALYON, PANISIA, PSEBANISIA, SCOLODIA, TAGESA, XAMACRISTA, Rafin. Sylva Tellur. 127–129 (1838).
BACTYRILOBIUM, Willd. Enum. Hort. Berol. 439 (1809).
CASSIANA, Rafin. in Am. Monthly Mag. (1818) 266.
CATHARTOCARPUS, Pers. Syn. i 459 (1805).
CHAMAECRISTA, Moench, Meth 272 (1794)
CHAMAEFISTULA, G. Don, Gen. Syst. ii. 451 (1832).
GRIMALDIA, Schrank, in Denkschr. Akad. Muench. (1808) 108. t. 3.
MACLAVEA, Montrouz. in Mém. Acad. Lyon, x. (1860) 198.
SENNA, Tourn. ex Mill. Gard. Dict. ed. VIII. (1768).
abbreviata, *Oliver, Fl. Trop. Afr.* ii. 271.—Afr. trop.
Absus, Aubl. Pl. Gui. i. 383 = lispidula.
Absus, *Linn. Sp. Pl.* 376.—Cosmop. trop.
acanthocladata, *Griseb. in Goett. Abh.* xxiv. (1879) 116. —Reg. Argent.
acapulcensis, H. B. & K. Nov. Gen. et. Sp. vi. 353 = biflora.
acclinus, F. Muell. Fragm. iv. 13 = glauca.
aciphylla, Benth. ex A. Gray, Bot. U. St. Expl. Exped. i. 465 = australis, *Sims*.
acisperma, *Schrank, in Colla, Hort. Ripul. App.* ii. 243.—Hab.?
acosmifolia, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 133.—Bras.
aculeata, *Pohl, ex Benth. l. c.* 128.—Bras.
acuminata, Moench, Meth. 273 = maryiandica.
acuminata, Willd. Sp. Pl. ii. 517 = C. Apoucouita.
acutensis, *Benth. in Mart. Fl. Bras.* xv. ii. 122.— Bras.
acuta, Meyen, Reise, i. 376 = Cumingii.
acutifolia, *Delile, Fl. Egypte*, 219. t. 27.—Aegypt.
acutisepala, *Benth. in Mart. Fl. Bras.* xv. ii. 97.— Am. trop.
adenantha, Zoll. & Moritz. Syst. Verz. Zoll. 2 = divaricata.
adenensis, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 553.—Arab.
adenopoda, Miq. in Linnaea, xix. (1847) 486 = C. Apoucouita.
adiantifolia, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 130.—Bras.
adnata, Willd. ex Vog. Syn. Cass. 18 = bicapsularis.
aegyptiaca, Willd. Enum. Hort. Berol. 442 = C. Sophera.
Aeschynomene, DC. in Collad. Hist. Cass. 127. t. 17 = mimosoides.
aethiopica, *Guibourt, Hist. Drogues*, ed. III. iii. 219.— Aegypt.
affinis, *Benth. in Mart. Fl. Bras.* xv. ii. 98.—Am. trop.
Afzeliana, Vog. in Linnaea, xv. (1841) 70 = podocarpa.
alata, *Linn. Sp. Pl.* 378.—Cosmop. trop.
albida, Orteg. Hort. Matr. Dec 92 = tomentosa.
Alcoparilla, H. B. & K. Nov. Gen. et Sp. vi. 355 = bicapsularis.
Alcaparra, *Phil. in Linnaea*, xxxiii. (1864–65) 61.— Chili.
amoena, Buch.-Ham. in Wall. Cat. n. 5821 = mimosoides.
amplifiora, Steud. in Flora, xxvi. (1843) ii. 760 = multijuga.
anceps, *Benth. in Mart. Fl. Bras.* xv. ii. 166.—Am. trop.
andina, Phil. in Linnaea, xxviii. (1856) 685 = Arnottiana.
Andreuxii, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 548.—Mexic.
andromedea, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 186.—Am. trop.
angulata, *Vog. Syn. Cass.* 16.—Bras.
angustifolia, *Vahl, Symb. Bot.* i. 29.—Afr. trop.
angustisiliqua, Lam. Encyc. i. 649 = biflora.
angustisiliqua, Torr. & Gray, Fl. N. Am. i. 396 = bahamensis.

CASSIA :—
angustissima, Lam. Encyc. i. 650 = mimosoides.
annulata, *Willd. ex Klotzsch, in Schomb. Faun. & Fl. Gui.* 1207, nomen.—Guiana.
aphylla, *Cav. Ic.* vi. 41. t. 561.—Reg. Argent.
apiculata, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 304 = crotalarioides.
Apoucouita, *Aubl. Pl. Gui.* i. 379. t. 146.—Guiana.
appendiculata, *Vog. Syn. Cass* 24.—Bras.
applanata, Anderss. Galap.-Carnes Vegetat. 254 = plicta.
aprica, Vell. Fl. Flum. 169; iv. t. 76 = cathartica.
arachoides, Burch. Trav. i. 341 = obovata
arayatensis, Llanos, Fragm. 71 = timorlensis.
arayatensis, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 71 = alata.
arborea, Macfad. Fl. Jamaic. i. 343 = siamea.
arborescens, Mill. Gard. Dict. ed. VIII. n. 15 = emarginata.
arborescens, Vahl, Symb Bot. iii. 56 = glauca.
arenaria, H. B. & K. Nov Gen. et Sp. vi. 370 = flexuosa.
arequipensis, *Meyen, in Vog. Syn. Cass.* 43.—Peruv.
Arerch, *Delile, Cent. Pl. Afr.* 29.—Aegypt.
argentea, *H. B. & K. Nov. Gen. et Sp.* 406.—Mexic.
aristata, *Benth. in Mart. Fl. Bras.* xv. ii. 170.—Bras.
aristata, Willd. ex Walp. Rep. i. 836 = patellaria.
armata, *S. Wats. in Proc. Am. Acad.* xi. (1876) 136.—Calif.
Armottiana, *Gill. in Hook Bot. Misc.* iii. (1888) 211.—Chili.
Arnottiana, Wight, ex Benth. in Trans. Linn. Soc. xxvii. (1871) 577 = nigricans.
arowana, *Schomb. Faun. & Fl. Gui.* 1206.—Guian.
artemisioides, *Gaudich. ex DC. Prod.* ii. 495.—Austral.
Aschrek, *Forsk. Fl. Aegypt. Arab.* 86.—Arab.
aspera, Muhl. Cat. ex Ell. Sketch, 474 = nictitans.
astroites, Willd. ex Cham. & Schlecht. in Linnaea, ii. (1827) 207 = villosa.
atomaria, *Linn. Mant.* 68.—Am. austr.
atropurpurea, Benth. in Trans. Linn. Soc. xxvii. (1871) 533, sphalm. = seq.
atrovirgata, Span. in Linnaea, xv. (1841) 201 = C. Sophera.
aurantia, *Ruiz & Pav. ex G. Don, Syst.* ii. 442.—Peruv.
aurata, Roxb. Hort. Beng. 32, nomen = laevigata.
auricoma, R. Grah. in Wall. Cat. n. 5822 = mimosoides.
auriculata, *Linn. Sp. Pl.* 379.—Ind. or.
aurita, *Collad. Hist. Cass.* 131.—Bras.
aurivilla, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 152.—Am. trop.
australis, *Reinw. ex Steud. Nom.* ed. II. i. 308, nomen. —Java.
australis, *Sims, Bot. Mag.* t. 2676.—Austral.
australis, Vell. Fl. Flum. 166; iv. t. 65 = appendiculata.
averrifora, Hook. Bot. Mag. t. 2138 = biflora.
bacillaris, *Linn. f. Suppl.* 231.—Am. trop.
bacillaris, Willd. ex Steud. Nom. ed. II. i. 303 = sclerocarpa.
Bacilius, Gaertn. Fruct. ii. 313 = javanica.
bahamensis, *Mill. Gard. Dict.* ed. VIII. n. 9.—Am. trop.
barbata, *Mart. & Nees, in Nov. Act. Nat. Cur.* xii. (1824) 32.—Bras.
Barclayana, Sweet, Fl. Austral. t. 32 = C. Sophera.
Barrenfeldii, Colla, Hort. Ripul. App. ii. 848 = australis.
basifolia, *Vog. Syn. Cass.* 56.—Bras.
batramensis, Steud. Nom. ed. II. i. 308, sphalm. = bahamensis.
bauhiniaefolia, Kunth, Mimos. 122. t. 37 = rotundifolia.
bauhinioides, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 180.—Texas; N. Mexic.
Berlandieri, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 520.—Mexic.
Berryana, Voigt, Hort. Suburb. Calc. 249 = montana.
Berteriana, Balb. ex DC. Prod. ii. 496 = biflora.
Berterii, Colla, Hort. Ripul. 30. t. 24 = bicapsularis.
bicapsularis, *Linn. Sp. Pl.* 376.—Am. trop.
bifda, Zoll. & Mor. in Nat. en Geneak. Arch. Neerl. Ind. iii. (1846) 63, 30 = divaricata.
biflora, Griseb. Cat. Pl. Cub. 79 = robiniaefolia.
biflora, *Linn. Sp. Pl.* 378.—Am. trop.

CASSIA :—
biflora, Mill. Gard. Dict. ed. VIII. n. 14 (= Aeschynomenes sp.).—Ind. occ. ?
bifoliolata, DC. in Collad. Hist. Cass. 130. t. 9 = rotundifolia.
biglandulosa, Bertol. in Bologn. Opusc. Sc. iii. (1819) 409 = patellaria.
bijuga, Vog. Syn. Cass. 17 = speciosa.
bimarginata, *Grisb. in Bonplandia*, vi. (1858) 5.—Panama.
birostris, *Domb. ex Vog. Syn. Cass.* 43.—Chili.
Blanchetti, *Benth. in Hook. Journ. Bot.* ii. (1840) 78. —Bolivia.
bonariensis, Hort. ex Colla, Hort. Ripul. 88 = corymbosa.
Bonplandiana, DC. Prod. ii. 490 = C. Flatula.
borboniodes, Vog Syn. Cass. 53 = Langsdorffii.
bonnemana, *Miq. Anal. Bot. Ind.* i. 9.—Borneo.
Botteriana, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 541.—Mexic.
brachycalyx, *Benth. in Mart. Fl. Bras.* xv. ii. 172.—Am. trop.
brachystachya, *Benth. in Hook. Journ. Bot.* ii. (1840) 78.—Bras.
brachystachya, Moç. & Sesse, ex DC. Prod. ii. 500 = paucifiora.
bracteata, Linn. f. Suppl. 232 = alata.
bracteolata, *Vog. Syn Cass* 52.—Bras.
bracteosa, Welw. ex Oliver, Fl. Trop. Afr. ii. 277 = didymobotrya.
brasiliana, Lam. Encyc. i. 649 = grandis.
brasiliana, Saldanha, Descr. Rio Madeiras Jan. 43 = ferruginea.
brevicalyx, *Benth. in Mart. Fl. Bras.* xv. ii. 134.—Bras.
brevifolia, *Lam. Encyc.* i. 651.—Madag.
brevipes, *DC. in Collad. Hist. Cass.* 119. t. 9.—Panama.
Brewsterii, *E. Mucll. 4th Ann. Rep.* 17.—Austral.
Brongniartu, *Gaudich. Bot Voy. Bonite*, t. 10.—Hab. ?
Browniana, Kunth, Mimos. 135 t. 41 = polyantha.
Buchanani, Kostel. Alg. Med.-Pharm. Fl. iv. 1833 = obtusifolia.
Burchellii, *Benth. in Mart. Fl. Bras.* xv. ii. 165.—Bras.
Burmanni, DC. Prod. ii. 502 = capensis ?
Burmanni, Wight, in Madras Journ. Sc. vi. (1837) t. 5 = obovata.
caespitosa, *Benth. in Mart. Fl. Bras.* xv. ii. 136.—Bras.
caliantha, G. F. W. Mey. Prim. Fl. Esscq. 169 = multijuga.
calycioides, *DC. in Collad. Hist. Cass.* 125. t. 20 B.—Am. trop.
campervan, Benth. in Hook. Journ. Bot. ii. (1840) 79 = paucifiora.
cana, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 34.—Bras.
cana, Schrank, ex Steud. Nom. ed. II. i. 304 = tomentosa.
cana, Wender. in Linnaea, xii. (1838) 22 = holosericea.
canaliculata, K. Br. in Sturt, Centr. Austral. ii. 61 = eremophila.
Canca, Cav. Desc. 132 = C. Sophera.
candelabrica, Benth. *in Mart. Fl. Bras.* xv. ii. 126.—Bras.
candenatensis, Dennst. Schluss. Hort. Malab. 32 = C. Tora.
Candolleana, Vog. Syn. Cass. 42 = bicapsularis.
canescens, H. B. & K. Nov. Gen. et Sp. vi. 357 = emarginata.
capensis, *Thunb. Prod. Pl. Cap.* 79.—Afr. austr.
caracasana, Jacq. Hort. Schoenb. iii. 11 = hirsuta.
cardiosperma, *F. Muell. Progm.* x. 60.—Austral.
carolinlana, Walt. Fl. Carol. 126 = occidentalis.
carthaginensis, Willd. ex Steud. Nom. ed. II. 304 = bacillaris.
Castigliolata, Collad. Hist. Cass. 121 = C. Chamaecrista.
cathartica, *Mart. Reise, Bras.* i. 548.—Bras.
centipeda, Wawra, in Flora, xlvii. (1864) 248 = verrucosa.
cernua, Balb. Cat. Hort. Taur. (1813) 22 = sulcata.
Chamaecrista, Boj. Hort. Masur. 122 = glandulosa.
Chamaecrista, *Linn. Sp. Pl.* 378.—Am. bor. et austr.
Chamaecrista, Mill. Gard. Dict. ed. VIII. n. 17 = procumbens.
Chamaecristoides, Collad. Hist. Cass. 184 = procumbens.
Linn. Herb.
Chateiniana, *Gaudich. in Freyc. Voy. Bot.* 486. t. 111.—Austral.

CASSIA :—
chinensis, Jacq. Coll. i. 64 = C. Sophera.
chinensis, Lam. Encyc. i. 644 = bicapsularis.
chiorlophylla, *Vog. Syn. Cass.* 56.—Bras.
chrysocarpa, *Desv. Journ. Bot.* iii. (1814) 72.—Bras.
chrysogyne, Miq. in Linnaea, xviii. (1844) 579 = chrysocarpa.
chrysoloma, De Not. Ind. Sem. Hort. Gen. (1840) = bicapsularia.
chrysophylla, A. Rich. Ess. Fl. Cub. 500 = emarginata.
chrysotricha, Collad. Hist. Cass. 99. t. 18 = chrysocarpa
chrysotricha, Mart. Herb. Fl. Bras. 108 = angulata.
ciliaris, Collad. Hist. Cass. 98 = C. Tagera.
ciliata, Hoffmgg. Verz. Pfl. Nachtr. i. 308 = sericea.
ciliata, Rafin. Fl. Ludov. 100 = occidentalis.
ciliolata, *Benth. in Mart. Fl. Bras.* xv. ii. 146.—Bras.
cinerascens, *Vog. Syn. Cass.* 60.—Bras.
cinerea, *Cham. & Schlecht. in Linnaea*, v. (1830) 599. —Mexic.
circinnata, *Benth. in Mitch. Journ. Trop. Austral.* 384.—Austral.
Clausseni, *Benth. in Hook. Journ. Bot.* ii. (1840) 79. —Bras.
coccinea, Wall. Cat. n. 5315 = C. Absus.
Colla, G. Don, Gen. Syst. ii. 442 = bicapsularis.
colutevoides, Collad. Hist. Cass. 102. t. 19 = bicapsularis.
comosa, *Vog. Syn. Cass.* 65.—Afr. austr.
conclnna, *Benth. Fl. Austral.* ii. 261.—Austral.
conferta, *Benth. in Mart. Fl. Bras.* xv. ii. 142.—Bras.
conjugata, *Ruiz & Pav. ex Benth. in Trans. Linn Soc.* xxvii. (1871) 540.—Am. trop.
conspicua, Vog. Syn. Cass. 47 = Sieberiana.
contorta, Vog. l. c. 17 = C. Tora.
coquimbensis, *Vog. l. c.* 37.—Chili.
cordistipula, *Mart. ex Benth. in Mart. Fl. Bras.* xv ii. 164.—Bras.
coriacea, *Benth. ex Benth. in Hook. Journ. Bot.* ii (1840) 78.—Bras
corifolia, *Benth. in Mart. Fl. Bras.* xv. ii. 120.—Am trop.
corneliana, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 79.—Afr. trop.
coromandeliana, Jacq Fragm. 67. t. 100 = C. Sophera.
coromiloides, A. Cunn. ex Benth. in Mitch. Journ. Trop. Austral. 384 = australis.
corymbosa, *Lam. Encyc.* i. 644.—Am. calid
corymbosa, Orteg. Hort. Matr. Dec. 124 = laevigata.
corinifolia. *G. Don, Gen. Syst.* ii. 445.—Bras.
Covesii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 399. —Calif.
crassifolia, Orteg. Hort. Matr. Dec. 122 = corymbosa.
crassiramea, *Benth. in Hook. Ic. Pl.* t. 1068.—Bolivia
crassispala, Benth. in Linnaea, xxii. (1849) 537 = bicapsularia.
crenulata, *Benth. in Hook. Journ. Bot.* ii. (1840) 80. —Bras.
Crista, Jaeq. Coll. i. 82 = biflora.
cretacea, Kunth, *Mimos.* 132. t. 40.—Am. austr.
Cruikshankii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 210.—Chili
cubensis, Hoffmgg. Verz. Pfl. Nachtr. i. 209 = corymbosa.
cultrifolia, *H. B. & K Nov. Gen. et Sp.* vi. 368.—Venezuela.
Cumingii, *Hook. tr Arn. in Hook. Bot. Misc.* iii. (1833) 211.—Chili.
cuneata, *DC. in Collad. Hist. Cass.* 121.—Am. austr.
cuneata, Griseb. Cat. Fl. Cub. 80 = linoata.
cunefolia, Vog. Syn. Cass. 49 = viscosa.
curvifolia, *Vog. l. c.* 55.—Bras.
cuspidata, Willd. Enum. Hort. Berol. 448 = pistaciaefolia.
cytisoides, *DC. in Collad. Hist. Cass.* 116. t. 14.—Bras.
dalbergiaefolia, *Benth. in Mart. Fl. Bras.* xv. ii. 149. —Am. trop.
debilis, *Vog. Syn. Cass.* 58.—Bras.
decipiens, Desv. Journ. Bot. iii. (1814) 72 = augustifolia.
decipiens, Griseb. Cat. Pl. Cub. 79 = stenophylla.
decrescens, *Benth. in Hook. Journ. Bot.* ii. (1840) 80. —Bras.
decumbens, *Benth. l. c.* 79.—Am. trop.
delagoënsis, *Harv. in Fl. Cap.* ii. 272.—Afr. austr.
densiflora, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 304.—Mexic.

CASSIA :—
densifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 80.—Bras.
denitata, *Vog. Syn. Cass.* 49.—Bras.
Deplanchei, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 554.—N. Caled.
desertorum, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 133.—Bras
desolata, *F. Muell. in Linnaea*, xxv. (1852) 389.—Austral.
Desvauxii, *Collad. Hist. Cass.* 181.—Am. trop.
didymobotrya, *Fresen. in Flora*, xxii. (1839) 58.—Abyssin.
diffusa, DC. in Mém. Soc. Phys. Genèv. ii. (1824) 130; Prod. ii. 504 = nictitans.
dimidiata, Klein, ex Wall. Cat. n. 5236 = Kleinii.
dimidiata, Roxb. Hort. Beng. 69, nomen = mimosoides.
diphylla, *Linn. Sp. Pl.* 376.—Am. trop.
disadena, Steud. in Flora, xxvi. (1843) ii. 760 = glandulosa.
discolor, Desv. Journ. Bot. iii. (1814) 78 = glauca.
dispar, *Willd. Enum. Hort. Berol.* 441.—Am. austr.
disperma, Vell. Fl. Flum. 167; iv. t. 69 = Peltophorum Vogelianum.
distichoclada, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 168.—Am. trop.
divaricata, *Nees. in Syll. Ratisb.* i. (1824) 94.—Ax. trop.; Malaya.
Dombyana, Vog. Syn. Cass. 28 = aurantia.
domingensis, *Spreng. Neue Entdeck.* iii. 55.—Ind. occ.
dormiens, Vell. Fl. Flum. 167; iv. t. 67 = bicapsularis.
drepanophylla, *Benth. in Mart. Fl. Bras.* xv. ii. 170.—Bras.
dumetorum, Bert. ex DC. Prod. ii. 499 = reticulata.
dysophylla, *Benth. in Mart. Fl. Bras.* xv. ii. 117.—Am. trop.
echinocarpa, *Benth. l. c.* 136.—Am. trop.
edulis, *Posada, in Baill. Adansonia*, x. (1872) 137.—N. Granat.
eglandulosa, *Dum.-Cours. Bot. Cult.* ii. vi. 34.—Hab.?
Ehrenbergii, Bisch. in Bot. Zeit. ii. (1844) 51 = angustifolia
elegans, H. B. & K. Nov. Gen. et Sp. vi. 342 = laevigata.
elegans, *Vogt, in Syll. Ratisb.* ii. (1828) 55.—Hab.?
elliptica, H. B. & K. Nov. Gen. et Sp. vi. 356 = emarginata.
elongata, Lemaire-Lis. in Journ. Pharm. vii. (1821) 345 = angustifolia.
emarginata, Bert. ex Steud. Nom. ed. II. i. 304, 305 = atomaria.
emarginata, Clos, in C. Gay, Fl. Chil. ii. 237 = bicapsularis.
emarginata, *Linn. Sp. Pl.* 376.—Am. trop.
emarginata, Mill. Gard. Dict. ed. VIII. n. 18 = pilosa.
enneaphylla, Koen. ex Wight & Arn. Prod. 289 = glauca.
enxiformis, Vell. Fl. Flum. 170; iv. t. 79 = C. Apoucouita.
erecta, Willd. ex Steud. Nom. ed. II. i. 303 = glandulosa.
eremophila, *A. Cunn. ex Vog. Syn. Cass.* 47.—Austral.
ericifolia, *Benth. in Mart. Fl. Bras.* xv. ii. 145.—Am. trop.
esculenta, Roxb. Hort. Beng. [31], nomen = C. Sophera.
exaltata, *Reinw. in Blume, Cat. Gew. Buitenz.* 68; *Span. in Linnaea*, xv. (1841) 301.—Ins. Timor.
excelsa, H. B. & K. Nov. Gen. et Sp. vi. 339 = C. Fistula.
excelsa, *Schrad. in Gott. Gel. Anz.* i. (1821) 717.—Bras.
exigua, Roxb. Hort. Beng. [31], nomen = C. Absus.
exinclusa, *Benth. in Hook. Journ. Bot.* ii. (1840) 80.—Bras.
fabaginifolia, H. B. & K. Nov. Gen. et Sp. vi. 358 = rotundifolia.
farfugia, Bertol. Fl. Genitim. 14 = oxyphylla.
fagonioides, Vog. Syn. Cass. 50 = hispidula.
falcata, Dum.-Cours. Bot. Cult. ed. II. vi. 35 = corymbosa.

falcata, Linn. Sp. Pl. 377 = occidentalis.
falciculata, *Oliver, Fl. Trop. Afr.* ii. 281.—Afr. trop.
fasciculata, Michx. Fl. Bor. Am. i. 269 = C. Chamaecrista.

CASSIA :—
fastigiata, Nees, in Flora, iv. (1821) 308 = excelsa.
fastigiata, Vahl, Symb. Bot. iii. 57 = glauca.
fastuosa, *Willd. ex Vog. in Linnaea*, xi. (1837) 654.—Am. trop.
felispenna, H. B. & K. Nov. Gen. et Sp. vi. 368 = C. Chamaecrista.
ferruginea, *Schrad. ex DC. Prod* ii. 489.—Am. trop.
Fieldii, Colla, Hort. Ripul. App. iv. 23. t. 11 = fillicifolia.
fillicifolia, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 148.—Bras.
filipendula, Boj. Hort. Maurit. 122 = mimosoides.
filiper, Benth. in Hook. Journ. Bot. ii. (1840) 81 = rotundifolia.
fimbriata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 9.—Malaya.
Fistula, Herbb. ex Oliver, Fl. Trop. Afr. 271 = C. Arereh.
Fistula, *Linn. Sp. Pl.* 377.—Ax. trop.
fistuloides, Collad. Hist. Cass. 87. t. 1 = C. Fistula.
flaccida, Clos, in C. Gay, Fl. Chil. ii. 238 = coquimbensis.
flavicoma, Benth. Pl. Hartw. 171 = trisulcata.
flavicoma, *H. B. & K. Nov. Gen. et Sp.* vi. 366.—Venezuela.
floccosa, A. DC. Not. 7me Pl. Rar. Jard. Genèv. 34. t. 4 = bicapsularis.
flexuosa, Burm. f. Fl. Cap. Prod. 12 = capensis.
flexuosa, *Linn. Sp. Pl.* 379.—Bras.
flexuosa, Mill. Gard. Dict. ed. VIII. n. 16 = serpens.
floribunda, Cav. Desc. 132 = laevigata.
floribunda, Collad. Hist. Cass. 88. t. 11 = corymbosa.
floribunda, Grisch. ex Benth. in Trans. Linn. Soc. xxvii. (1871) 549 = racemosa.
florida, Vahl, Symb. Bot. iii. 57 = siamea.
fluminensis, Vell. Fl. Flum. 168; iv. t. 72 = Dimorphandra exaltata.
Fockeana, Miq. in Linnaea, xviii. (1844) 579 = bacillaris.
foetida, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 443 = stipulacea.
foetida, Sallsb. Prod. 325 = C. Tora.
foetida, Willem. in Usteri, Ann. Bot. xviii. (1796) 32 = occidentalis.
foetidissima, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 441 = mexicana.
foliolosa, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 544.—Guatemala.
foliolosa, Steud. Nom. ed. II. i. 305 = seq.
foliosa, G. Don, Gen. Syst. ii. 745 = tenella?
foliosa, *Phil. in Anal. Univ. Chil.* (1862) ii. 879.—Chili.
Fraseri, A. Cunn. ex Vog. Syn. Cass. 48 = australis.
frassinifolia, H. B. & K. Nov. Gen. et Sp. vi. 349 = pistaciaefolia.
frondosa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 210 = bicapsularis.
frondosa, [Soland. in] Ait. Hort. Kew. ed. I. ii. 58 = biflora.
frutescens, Mill. Gard. Dict. ed. VIII. n. 2 = C. Sophera.
fruticescens, Bert. ex Spreng. Syst. ii. 335 = vimines.
fruticosa, Mill. Gard. Dict. ed. VIII. n. 10 = bacillaris.
fulgens, Macfad. Fl. Jamaic. i. 342 = biflora.
fulgens, Wall. Cat. n. 5310 = multijuga.
fusconeura, *Benth. in Mart. Fl. Bras.* xv. ii. 158.—Am. trop.
galegaefolia, Linn. Syst. Nat. ed. X. 1017 = biflora.
Galeottiana, *Martens, in Bull. Acad. Brux.* x. ii. (1843) 305.—Mexic.
gallinaria, Collad. Hist. Cass. 96 = C. Tora.
Gardneri, *Benth. in Mart. Fl. Bras.* xv. ii. 120.—Bras.
Gaudichaudii, *Hook. & Arn. Bot. Beech. Voy.* 81.—Ins. Sandvic.
gemina, Vell. Fl. Flum. 169; iv. t. 74 = C. Chamaecrista.
geminata, *Benth. in Mart. Fl. Bras.* xv. ii. 135.—Bras.
geminata, Vahl, ex DC. Prod. ii. 505 = mimosoides.
geminiflora, *Collad. Hist. Cass.* t. 8.—Mexic.
geminiflora, Moç. & Sesse, in Collad. Hist. Cass. 103. t. 8 = biflora.
geminiflora, Schrank, Hort. Monac. t. 26 = occidentalis.
geniculata, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 440 = villosa.

CASSIA :—
gigantea, Bert. ex DC Prod. ii. 492 = siamea.
glabra, Collad. Hist. Cass. 131 = cytisoides.
glandulifera, Reinw. ex Steud. Nom. ed. II. i. 305 = seq.
glandulifera, Reinw in Blume, Cat. Gew. Buitenz. 68 = bicapsularis.
glandulosa, Hook. Bot. Mag. t. 3485 = virgata.
glandulosa, *Linn. Syst. Nat.* ed. X. 1017.—Am trop.
glauca, *Lam. Encyc.* i. 647.—As. trop.; Austral.; Polynes.
glaucescens, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 539.—Chili.
glaucescens, *Hoffmgg. Vers. Pfl. Nachtr.* i. 209.—Cuba
glutinosa, *DC. Prod.* ii. 496.—Austral.
goensus, *Dals. in Hook. Kew Journ.* iv. (1852) 112.—Ind or.
gonioides, *A. Cunn. ex Benth. in Hook. Ic. Pl.* xi. 48. t. 1061.—Austral.
gonoclada, *Benth. in Mart. Fl. Bras.* xv. ii. 109.—Bras.
gonoclada, Benth in Trans. Linn. Soc. xxvii. (1871) 572 = choriophylla.
goratensis, *Fresen. in Flora*, xxii. (1839) 53.—Abyssin.
gracilis, *Kunth, Mimos.* 120. t. 36.—Am. trop.
gracillima, *Benth. in Mart. Fl. Bras.* xv. ii. 143.—Bras.
gracillima, Welw. Apont. 590 = mimosoides.
grammica, *Spreng. Neue Entdeck.* iii. 55.—Am. trop.
grandiflora, Desf. Cat. Hort. Par. ed. III. 302 = laevigata.
grandiflora, Pers. Syn. i. 457 = bicapsularis.
grandis, *Linn f Suppl.* 230.—Panama.
grandistipula, Vog Syn. Cass. 58 = rotundata.
Grantii, *Oliver, Fl. Trop. Afr.* ii. 279.—Afr. trop.
graveolens, Colla, Hort. Ripul. App. ii. 343 = atomaria.
Greggii, *A. Gray, Pl. Wright.* i. 59.—Mexic.
grisea, A. Rich. fas. Fl Cub. 408 = emarginata.
gunneusis, G. Don, Gen. Syst ii 449 = mimosoides.
Hartmegi, Benth. Pl. Hartw. 117 = oxyphylla.
hecatophylla, DC. in Collad. Hist. Cass. 124. t. 18 = mimosoides.
hedysaroides, *Vog. Syn. Cass.* 52.—Bras.
heptanthera, *F. Muell. Progn.* t. 8.—Austral.
Herbertiana, Lindl. Bot. Reg. t. 1422 = laevigata.
herpetica, Jacq. Obs. Bot. ii. 24. t. 45. f. 2 = alata.
heterofolia, Lindl. in Mitch. Three Exped. Austral. ed. II. ii. (1838) 122 = eremophila.
hetrophylla, Walp. Rep. i. 838, sphalm. = mimosoides.
Hibbertiana, Steud. Nom ed. II. i. 305, sphalm. = Herbertiana.
Hilariana, *Benth. in Mart. Fl. Bras.* xv. ii. 108.—Am. trop.
Hildebrandtii, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 80.—Afr. trop.
hirsuta, *Linn. Sp. Pl.* 378.—Am. trop.
hirsuta, Vell. Fl. Flum. 170; iv. t. 80 = cathartica.
hirta, *Willd. Enum. Hort. Berol. Suppl.* 28.—Hab.?
hispida, Collad. Hist. Cass. 118 = hispidula.
hispidula, *Vahl, Eclog. Am.* iii. 10.—Mexic.
Hoffmanseggii, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 104.—Bras.
holosericea, *Fresen. in Flora*, xxii. (1839) i. 54.—Abyssin.
homophylla, *Hoffmgg. Vers. Pfl. Nachtr.* i. 209.—Hab.?
Hookeriana, *Gill. in Hook. Bot. Misc.* iii. (1833) 210.—Chili.
Hornemanni, DC. Prod. ii. 507 = C. Chamaecrista.
Horsfieldii, Miq. Fl. Ind. Bat. i. 99 = glauca.
Houttuyana, Collad. Hist. Cass. 122 = *bifora*, Mill.
Humboldtiana, DC. Prod. ii. 489 = spectabilis.
humilis, Collad. Hist. Cass. 96 = C. Tora.
humilis, Willd. ex Steud. Nom. ed. II. i. 305 = C. Tagera.
hybrida, Tenore, Cat. Hort. Neap. 80 = laevigata.
hymenaefolia, *Benth. in Mart. Fl. Bras.* xv. ii. 130.—Am. trop.
hypnotica, Vell. Fl. Flum. 169; iv. t. 75 = C. Chamaecrista.
hypoleuca, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 117.—Bras.
inaequilatera, *Balb. ex DC. Prod.* ii. 490.—N. Granat.
incana, *Vog. Syn. Cass.* 52.—Bras.

CASSIA :—
incarnata, *Pav. ex Benth. in Trans. Linn. Soc.* xxvii. (1871) 545.—Peruv.
incurvata, *Benth. in Mart. Fl. Bras.* xv. ii. 187.—Am. trop.
indecora, H. B. & K. Nov. Gen. et Sp. vi. 818 = bicapsularis.
indica, Poir. Encyc. Suppl. ii. 196 = C. Sophera.
inflata, Spreng. Syst. ii. 536 = bicapsularis.
laldora, *Benth. in Mart. Fl. Bras.* xv. ii. 154.—Bras.
italica, Lam. ex Steud. Nom. ed. I. 166, 167 = C. Senna.
itambana, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 151.—Am. trop
Jacobinea, *Benth. in Mart. l. c.* 132.—Am. trop.
javanica, Arabl. Pl. Gui. i. 382 = grandis.
javanica, *Linn. Sp. Pl.* 379.—Malaya.
javanica, Sieber, ex Benth in Trans. Linn. Soc. xxvii. (1871) 517 = Siebersana.
javanica, Vell. Fl. Flum. 168; iv. t. 78 = ferruginea.
Johannae, *Vatke, in Oestr. Bot Zeitschr.* xxx. (1880) 77.—Afr. trop.
Ketchita, *Hasselq. It. Palaest.* 469.—Palaest.
Kirkii, *Oliver, Fl. Trop. Afr.* ii. 281.—Afr. trop.
kitulensis, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 81.—Afr. trop.
Kleinii, *Wight & Arn. Prod.* 293.—Ind. or. ; Malaya.
Kotschyana, *Oliver, Fl. Trop. Afr.* ii. 271.—Afr. trop.
Kunthiana, Cham. & Schlecht. in Linnaea, v. (1830) 598 = C. Tagera.
lactea, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 78.—Madag.
lacta, *H. B. & K. Nov. Gen. et Sp.* vi. 840.—N. Granat.
laevigata, *Willd. Enum. Hort. Berol.* 441.—Cosmop. trop.
lamprosperma, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 146.—Am. trop.
lanceolata, Colled. Hist. Cass. 93 t. 15 = acutifolia.
lanceolata, Forsk. Fl. Aegypt. Arab. 85 = C. Sophera.
lanceolata, Pers. Syn. i. 456 = uniflora.
lanceolata, Wight & Arn. Prod. 288 = angustifolia.
Langsdorffii, *Kunth, ex Vog. Syn. Cass.* 55.—Bras.
latifolia, *G. F. W. Mey. Prim. Fl. Esseq.* 166.—Guiana.
latistipula, *Benth. in Mart. Fl. Bras.* xv. ii. 156.—Bras.
latopetiolata, *Domb. ex Vog. Syn. Cass.* 70.—Peruv.
lavradioides, *Benth. in Mart. Fl. Bras.* xv. ii. 137.—Am. trop.
laxiflora, *Benth. Fl. Austral.* ii. 283.—Austral.
leiandra, *Benth. in Mart. Fl. Bras.* xv. ii. 94.—Bras.
leiantha, Benth. in Hook. Journ. Bot. ii. (1840) 78 = hispidula.
leiophylla, *Vog. Syn. Cass.* 26.—Bras.
lenitiva, Bisch. in Bot. Zeit. vii. (1850) 885 = acutifolia.
lenticifolia, *Benth. in Mart. Fl. Bras.* xv. ii. 144.—Am. trop.
leptocarpa, *Benth. in Linnaea*, xxii. (1849) 528.—N. Mexic.
leptoclada, *Benth. Fl. Austral.* ii. 290.—Austral.
leptophylla, *Vog. Syn. Cass.* 18.—Bras.
Leschenaultiana, DC. in Mém. Soc. Phys. Genèv. ii. (1824) 182 = mimosoides.
Leschenaultii, Wall. Cat. n. 5825 = mimosoides.
leucoxylon, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1207, nomen.—Guiana.
Liebmanni, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 549.—Mexic.
ligustrina, Forsk. ex Benth. l. c. 538 = C. Sophera.
ligustrina, *Linn. Sp. Pl.* 378.—Am. trop.
ligustrina, Mill. Gard. Dict. ed. VIII. n. 12 = obovata.
ligustrinoides, Schrank, in Denkschr. Akad. Muench. vi. (1816–17) 179 = angustifolia.
limensis, Lam. Encyc. i. 643 = bicapsularis.
Lindheimeriana, *Schlecht. in Linnaea*, xxi. (1848) 457.—Texas.
Lindleyana, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 241 = multijuga.
linearifolia, *G. Don, Gen. Syst.* ii. 446.—Bras.
linearis, Michx. Fl. Bor. Am. i. 261 = occidentalis.
linearis, *Sw. Prod. Veg. Ind. Occ.* 66.—Am. trop.

CASSIA :—
lomatopoda, *Benth. in Mart. Fl. Bras.* xv. ii. 146.—Bras.
longegландulosa, *Benth. l. c.* 119.—Am. trop.
longicuspis, *Benth. l. c.* 150.—Am. trop.
longifolia, Steud. Nom. ed. II. i. 304, in syn. = angustifolia.
longuracemosa, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 80.—Afr. trop.
longistiliqua, Blanco, Fl. Filip. ed. I. 338 = birostra.
longistiliqua, Linn. f. Suppl. 280 = occidentalis.
Lorentzii, *Niedérl. in Roca, Exped. Rio Negro*, ii. 210.—Patagon.
lotoides, H. B. & K. Nov. Gen. et Sp. vi. 861 = hapidula.
lucens, Vog. Syn. Cass. 46 = racemosa.
lucida, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 18.—Hab.?
Lundii, *Benth. in Mart. Fl. Bras.* xv. ii. 143.—Am. trop.
lutescens, G. Don. Gen. Syst. ii. 442 = tomentosa.
machaerifolia, *Benth. in Mart. Fl. Bras.* xv. ii. 151.—Am. trop.
macranthera, *Collad. Hist. Cass.* 182.—Bras.
macranthera, *DC. in Collad. l. c.* 99. t. 8.—Bras.
macrocarpa, *Micheli, in Mém. Soc. Phys. Genève.* xxviii. (1883) n. VII. 43.—Paraguay.
macrophylla, *Kunth, Mimos.* 136. t. 38.—N. Granat.
macrophylla, Kulz & Pav. ex Benth. in Trans. Linn. Soc. xxvii. (1871) 530 = Ruiziana.
madagascariensis, *Bojer. Hort. Mauric.* 190.—Madag.
madagascariensis, Four. ex Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 424 = Swartzia madagascariensis.
magnifica, Mart. Herb. h.t. Bras. 106 = multijuga.
magnifolia, *F. Muell. Fragm.* i. 166.—Austral.
malacophylla, *Vog. Syn. Cass.* 55.—Bras.
Mandoni, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 540.—Peruv.
Mannii, *Oliver, Fl. Trop. Afr.* ii. 272.—Afr. trop.
marginata, *Roxb. Hort. Beng.* [81]; *Fl. Ind.* ii. 338. —Ind. or.
margotta, Willd. Enum. Hort. Berol. 443 = polyphylla.
Marmari, Aubl. Pl. Gui. 382 = biflora.
maritima, Willd. ex Vog. Syn. Cass. 38 = pilifera.
Martiana, *Benth. in Mart. Fl. Bras.* xv. ii. 127.—Am. trop.
maryllodica, *Linn. Sp. Pl.* 378.—Am. bor.
medellinensis, *Posada, in Baill. Adansonia*, x. (1872) 167.—N. Granat.
medica, Forsk. Fl. Aegypt. Arab. p. cxi = angustifolia.
medicinalis, Bisch. in Bot. Zeit. vii. (1850) 887 = angustifolia.
megalantha, *Decne. Herb. Tim. Descr.* 186.—Ins. Timor.
melanocarpa, Bert. ex DC. Prod. ii. 491 = viminea.
mexicana, *Vacq. Hort. Schoenbr.* ii. t. 203.—Am. trop.
micans, Nees, in Flora, iv. (1821) 808 = speciosa.
micrantha, Guill. & Perr. Fl. Seneg. Tent. i. 262 = nigricans.
microphylla, Willd. Sp. Pl. ii. 529 = mimosoides.
Millerii, Colled. Hist. Cass. 192 = pilosa.
Mimosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 9.—Malaya.
mimosoides, DC. Prod. ii. 502 = Kleinii.
mimosoides, *Linn. Sp. Pl.* 379.—Cosmop. trop.
misera, *Phil. Fl. Atac.* 17.—Chili.
mollis, Vahl, Symb. Bot. iii. 57 = grandis.
molliuscula, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 440 = stomaria.
monaden, Vell. Fl. Flum. 166; iv. t. 68 = splendida.
monantha, DC. Prod. ii. 506 = sericea.
monophylla, Vell. Fl. Flum. 166; iv. t. 61 = rotundifolia.
montana, *Heyne, in Roth, Nov. Sp. Pl.* 214.—Ind. or.
montana, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 71 = timoriensis.
montana, *Vell. Fl. Flum.* 169; iv. t. 77.—Bras.
monticola, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 134.—Am. trop.
moschata, Benth. in Hook. Journ. Bot. ii. (1840) 76 = leiandra.
moschata, *H. B. & K. Nov. Gen. et Sp.* vi. 858.—Am. austr.

CASSIA :—
mucronata, *Spreng. Syst.* ii. 341.—Bras.
mucronifera, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 116.—Bras.
multifoliata, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 307.—Am. trop.
multifoliata, Scheele, in Linnaea, xvii. (1843) 337 = excelsa.
multifoliata, Vog. Syn. Cass. 89 = macranthera.
multiglandulosa, Jacq. Ic. Rar. i. t. 72 = tomentosa.
multijuga, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 108.—Guiana.
multinervia, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 166.—Am. trop.
multiseta, *Benth. l. c.* 141.—Am. trop.
multistiqua, J. F. Gmel. Syst. 1521 (quid?).
Mutiana, *Kunth, Mimos.* 142. t. 43.—Am. trop.
myriophylla, Wall. Cat. n. 5836 = mimosoides.
nana, *Benth. in Mart. Fl. Bras.* xv. ii. 110.—Am. trop.
Necsiana, *Mart. ex Benth. l. c.* 148.—Bras.
neglecta, *Vog. in Linnaea*, x. (1836) 894.—Bras.
nemophila, Walp. Rep. i. 827, sphalm. = eremophila.
nemorosa, H. B. & K. Nov. Gen. et Sp. vi. 858 = biflora.
nervosa, Vog. Syn. Cass. 89 = speciosa.
nicaraguensis, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 552.—Am. trop.
nictitans, *Linn. Sp. Pl.* 380.—Am. bor. et austr.
nictitans, Wydl. ex Steud. Nom. ed. II. i. 303, 306 = mimosoides.
nigricans, *Vahl, Symb. Bot.* i. 30.—Geront. trop.
nitida, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = viminea.
nodosa, *Buch.-Ham. ex Roxb. Hort. Beng* [81]; *Fl. Ind.* ii. 336.—As. trop.
notabilis, *F. Muell. Fragm.* iii. 98.—Austral.
nummularia, Vahl, ex DC. Prod. ii. 501 = rotundifolia.
nummulariaefolia, *Benth. in Mart. Fl. Bras.* xv. ii. 143. —Bras.
nutans, Collad. Hist. Cass. 113. t. 4 = stomaria.
obcordata, Sw. ex Wickstr. in Vet. Acad. Handl. (1825) 429 = linearis.
obliqua, *Ruiz & Pav. ex G. Don, Gen. Syst.* ii. 451. —Peruv.
obliquifolia, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 40 = occidentalis.
oblongifolia, *Vog. Syn. Cass.* 23.—Bras.
obovata, *Collad. Hist. Cass.* 92. t. 15.—Geront. trop.
obtecta, *Benth. in Mart. Fl. Bras.* xv. ii. 136.—Am. trop.
obtusa, Clos, in C. Gay, Fl. Chil. ii. 225 = bicapsularis.
obtusa, Roxb. Hort. Beng. [81]; Wight, Ic. t. 757 = obovata.
obtusata, Hayne, Arzneig. ix. t. 43 = obovata.
obtusata, Hochst. & Steud. Nom. ed. II. i. 306 = holosericea.
obtusifolia, Linn. Sp. Pl. 377 = C. Tora.
occidentalis, Hort. ex Steud. Nom. ed. II. i. 306 = hamophylla.
occidentalis, *Linn. Sp. Pl.* 377.—Cosmop. trop.
occidentalis, Naves, in Blanco, Fl. Philipp. ed. III. t. 78 = C. Sophera.
ochnacea, *Vog. Syn. Cass.* 51.—Bras.
odorata, *Morris, Fl. Conspic.* t. 57.—Austral.
oleiphylla, *Vog. Syn. Cass.* 68.—Bras.
oligoclada, *F. Muell. Fragm.* iii. 49.—Austral.
oligophylla, *F. Muell. l. c.*—Austral.
oligosperma, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 147.—Bras.
opaca, R. Grah. ex Vog. Syn. Cass. 42 = stipulacea.
orbiculata, *Benth. in Hook. Journ. Bot.* ii. (1840) 79 —Bras.
orinocensis, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 150.—Bras.
orientalis, Pers. Syn. i. 457 = acutifolia.
ornata, Benth. ex Benth. in Trans. Linn. Soc. xxvii. (1871) 522 = latifolia.
ornata, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1104, nomen.—Guiana.
ornithocladioides, Steud. Nom. ed. I. 266 = neq.
ornithopoides, Lam. Encyc. i. 644 = sericea.
Otterbeynii, G. F. W. Mey. Prim. Fl. Esseq. 169 = glandulosa.
ovalifolia, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 305 = bicapsularis.

CASSIA :—
ovata, *Mirat & Lenz, ex Geiger, Pharm. Bot.* ed. II. ii. 1126.—Arab.
oxyadena, Turcz. Pl. ind. 495 = biflora.
oxyphylla, *Kunth, Mimos.* 139. t. 39.—N. Granat.
pachycalyx, Vog. Syn. Cass. 52 = cathartica.
pachyphylla, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 166.—Bras.
pallida, Vahl, Eclog. Am. iii. 12 = biflora.
palmata, Wall. Cat. n. 5306 = timoriensis.
paniculata, *Benth. in Mart. Fl. Bras.* xv. ii. 153.—Am. trop.
paposana, *Phil. Fl. Atac.* 17.—Chili.
papulosa, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 81.—Bras.
paradyction, *Vog. Syn. Cass.* 45.—Bras.
paraguensis, *Michell, in Mem. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 44.—Paraguay.
Parahybæ, Vell. Fl. Flum. 168 ; iv. t. 71 = Schizolobium excelsum.
paramariboënsis, *Miq. in Bot. Zeit.* i. (1843) 461.—Guian.
paratyensis, *Vell. Fl. Flum.* 168 ; iv. t 70.—Bras.
Parkeriana, DC. Prod. ii. 504 = riparia.
parvifolia, Wender. in Syll. Ratisb. i. (1824) 65 = glandulosa.
parvistipula, *Benth. in Mart. Fl. Bras.* xv. ii. 170.—Am. trop.
pascuorum, *Mart. ex Benth. l. c.* 165.—Am. trop.
patellaria, *DC. in Collad. Hist. Cass.* 125. t. 16.—Am. trop.
patula, [Soland. in] Ait. Hort. Kew. ed. I. ii. 51 = C. Sophera.
pauciflora, *H. B. & K. Nov. Gen. et Sp.* vi. 360.—Am. trop.
Pavoniana, G Don, Gen Syst. ii. 447 = glandulosa.
Pearcii, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 552. —Bolivia.
pedicellaris, *DC. Prod.* ii. 504.—Ins. S. Doming.
pendula, Willd. Enum. Hort. Berol. 440 = bicapsularis.
pentagonia, *Mill. Gard. Dict.* ed. VIII. n. 18.—Am. trop.
pentandra, Raddi, in Mem. Mod. xviii. Fis. (1820) 399 = rotundifolia.
peralteana, *H. B. & K. Nov. Gen. et Sp.* vi. 356.—Mexic.
Persoonii, Collad. Hist. Cass. 119 = uniflora.
peruviana, Vog. Syn. Cass. 39 = obliqua.
Petersiana, Bolle, in Peters, Reise Mossamb. Bot. 18 = delagoensis.
phyllodinea, *R. Br. App Sturt, Centr. Austral.* 15.—Austral.
phyllostachya, *Benth. in Mart. Fl. Bras.* xv. ii. 150. —Am. trop.
picta, *G. Don, Gen. Syst.* ii. 444.—Ecuador ; Ins. Galap.
pisidens, Steud. Nom. ed. II. i. 307 = nictitans.
pilifera, *Vog. Syn. Cass.* 28.—Bras.
pilosa, *Linn. Syst. Nat.* ed. X. 1017.—Am. trop.
pinnata, Hort. ex Volgt, in Syll. Ratisb. ii. (1828) 55 = elegans.
platycalesthia, *H. & K. Nov. Gen. et Sp.* vi. 349.—N. Granat.
planisiliqua, Burm. f. Fl. Ind. 96 = Albizzia Lebbek.
planisiliqua, Linn. Sp. Pl. 377 = occidentalis.
planisiliqua, *Lam. Encyc.* i. 645.—Ins. Guadelup.
platypoda, K. Br. in Sturt, Centr. Austr. ii. 78 = eremophila.
pleurocarpa, *F. Muell. Fragm.* i. 228.—Austral.
Plumieri, *DC. Prod.* ii. 506.—Ins. Guadelup.
plumosa, Vog. Syn. Cass. 54 = mimosoides.
podocarpa, *Guill. & Perr. Fl. Senegʼ. Tent.* 259.—Afr. trop.
Pohliana, *Benth. in Hook. Journ. Bot.* ii. (1840) 80.—Bras.
polyadena, *DC. Pl. Rar. Hort. Genèv.* 2ª *Rapp.* 12.—Ind. occ.
polyanthe, *Moç. & Sesse, ex Collad. Hist. Cass.* 112. t. 2.—Mexic.
polyphylla, *Spacg. Coll.* iv. 104 ; *Ic. Rar.* t. 460.—Ind. occ.
polystachya, *Benth. in Hook. Journ. Bot.* ii. (1840) 77.—Guiana.
portoricensis, Bancroft, ex Macf. Fl. Jamaic. i. 340 = chrysota.
Potentilla, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 168.—Am. trop.
praetexta, *Vog. Syn. Cass.* 67.—Am. trop.
procumbens, Forsk. Fl. Aegypt. Arab. p. cxl = nigricans.
procumbens, Linn. Sp. Pl. ed. I. 380 = nictitans.

CASSIA :—
procumbens, *Linn. Sp. Pl.* ed. II. 548, pro parte, et Herb.—Am. bor.
procumbens, Mill. Gard. Dict. ed. VIII. n. 20 = hispidula.
procumbens, Thunb. Fl. Jap. 179 = mimosoides.
prominens, D. Dietr. Syn. Pl. ii. 1479 = macranthera.
propinqua, Cham. & Schlecht. in Linnæa, v. (1830) 599 = tristicula.
prostrata, H. B. & K. Nov. Gen. et Sp. vi. 369 = glandulosa.
prostrata, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 441 = serpens.
prostrata, Koen. ex Roxb. Hort. Beng. 32 = pumila.
prunosa, *F. Muell. Fragm.* iii. 48.—Austral.
pseudo-nictitans, Hance, in Journ. Linn. Soc. xi. (1871) 78 = nictitans.
psammophila, Benth. in Trans. Linn. Soc. xxvii. (1871) 571 = cordistipula.
pulcherr., Welw. Apont. 567. n. 40 = Sieberiana.
puberula, H. B. & K. Nov. Gen. et Sp. vi. 341 = bacillaris.
pubescens, R. Br. in Salt, Abyss. App 64 = holosericea.
pubescens, Ehrenb. & Hempr. ex Benth. in Proc. Linn. Soc. ii. (1855) 328 = holosericea.
pubescens, *Spacg. Fragm.* 46. t. 57.—Bras.
pubescens, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 442 = tomentosa.
pubigera, Lag. Gen. et Sp. Nov. 14 = hirsuta.
pudibunda, *Mart. ex Benth. in Mart. Fl. Bras*, xv. ii. 102.—Bras.
pulchella, *Boj. Hort. Maurit.* 122.—Ins. Maurit.
pulchella, Salisb. Prod. 326 = C. Chamaecrista.
pulcherrima, *Dehn. Rivista Nap.* i. 111. 174.—Guiana.
pulchra, H. B. & K. Nov. Gen. et Sp. vi. 362 = Desvauxii.
pumila, *Lam. Encyc.* i. 651.—Cosmop. trop.
pumila, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 307 = leiophylla.
pumila, F. Muell. Fragm. iii. 47 = concinna.
Pumilio, *J. Gray, Pl. Lindheim.* 180.—Am. bor. occ.
punctata, Scheele, in Linnæa, xvii. (1843) 336 = sylvestris.
punctata, *Vog. Syn. Cass.* 51.—Bras.
punctulata, Hook. & Arn. Bot. Beech. Voy. 420 = pauciflora.
purgans, *Steud. Nom.* ed. II. i. 307.—Chili.
purpurea, Roxb. Hort. Beng. [31] = C. Sophera.
pygmæa, DC. Prod. ii. 503 = punctata, *Linn. Herb. quadrangularis*, Zoll. & Mor. Syst. Verz. Zoll. 2 = laevigata.
quadriflora, Vog. Syn. Cass. 47 = glutinosa.
Quebracho, Steud. ‘1838’ ex Nom. ed. II. i. 307, 304 = bicapsularis.
quinquangulata, *Rich. in Act. Soc. Hist. Nat. Par.* (1792) 108.—Am. trop.
racemosa, *Mill. Gard. Dict.* ed. VIII. n. 19.—Bras.
ramiflora, Vog. Syn. Cass. 48 = C. Apoucouita.
ramosa, Vog. l. c. 51 = uniflora.
ramosissima, H. B. & K. Nov. Gen. et Sp. vi. 366 = glandulosa.
reclinata, Sallah. Prod. 326 = marylandica.
Reinwardtii, *Hassk. in Hoev. & De Vriese, Tijdschr.* v. (1838-39) 269.—Java.
reniformis, *G. Don, Gen. Syst.* ii. 440.—Bras.
renigera, *Wall. Cat.* n. 5307.—Burma.
repens, *Vog. Syn. Cass.* 60.—Bras.
reticulata, *Willd. Enum. Hort. Berol.* 443.—Am. sunt.
retusa, *Soland. ex Vog. in Linnæa*, xv. (1841) 72.—Austral.
revoluta, F. Muell. in Trans. Vict. Inst. (1855) 120 = australis.
rhombifolia, Roxb. Hort. Beng. [31] ; Fl. Ind. ii. 334 = C. Fistula.
Richardiana, Kunth, Mimos. 139. t. 142 = multijuga.
Riedelii, *Benth. in Mart. Fl. Bras.* xv. ii. 122.—Am. trop.
rigidifolia, *Benth. l. c.* 142.—Bras.
riparia, *H. B. & K. Nov. Gen. et Sp.* vi. 359.—Am. trop.
robiniæfolia, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 548.—Am. trop.
robinioides, Willd. Enum. Hort. Berol. 448 = C. Sophera.
Roemeriana, *Scheele, in Linnæa*, xvi. (1842) 457.—Texas.

CASSIA :—
Roraimæ, *Benth in Trans. Linn. Soc* xxvii. (1871) 574.—Guiana.
rostrata, Voigt, in Syll. Ratisb. ii. (1828) 55 = biflora.
rotundata, *Vog. Syn. Cass* 58.—Bras.
rotundifolia, *Pers. Syn.* i. 466.—Mexic.
Roxburghiana, K. Grah. in Wall. Cat. n. 5328 = mimosoides.
Roxburghii, DC. Prod. ii. 489 = marginata.
Roxburghii, Steud. Nom. ed. II. i. 307, sphalm. = *Roxburghiana*.
rufa, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 306.—Mexic.
rugosa, *G. Don, Gen. Syst.* ii 440.—Bras.
Ruiziana, *Vog. Syn. Cass.* 40.—Am. trop.
rupicola, Benth. in Trans. Linn. Soc. xxvii. (1871) 574 = oleosphylla.
rusa, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 306 = tristicula.
ruscifolia, Jacq. Ic. Rar. t. t. 73 = C. Sophera.
Sabæa, *Delile, Cent. Pl. Afr.* 28.—Aegypt.
Sagera, Lam. Encyc. i. 643 = C. Tora.
Sagotiana, *Benth. in Mart. Fl. Bras.* xv. ii. 93.—Guiana.
Saltiana, Steud. Nom. ed. II. i. 307 = holosericea.
sapindifolia, *Vog. Syn. Cass.* 34.—Bras.
sarmentosa, Miq. in Bot. Zeit. i. (1843) 461 = uniflora.
scabra, *Pohl, ex Benth. in Mart. Fl. Bras.* xv. ii. 135. —Am. trop.
scabrella, *Hoffmgg. Verz Pfl. Nachtr.* ii. 81.—Bras.
scandens, *Ruiz & Pav. ex G. Don, Gen. Syst.* ii. 440. —Ecuador.
Schimperi, Steud. Nom. ed. II. i. 307 = holosericea.
schinifolia, A. DC. 7ª Not. Pl. Rar. Hort. Genèv. 35 = C. Sophera.
Schlimii, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 571.—Am. trop.
Schomburgkii, *Klotzsch, ex Schomb. Faun. & Fl. Gui.* 1207, nomen.—Guiana.
Schulzeii, Colla, Hort. Ripul. App. ii. 348 = australis.
sclerocarpa, *Vog. Syn Cass.* 15.—Bras.
secunda, *Benth. in Mart. Fl. Bras.* xv. ii. 148.—Bras.
Sellii, G. Don, Gen Syst. ii. 442 = multijuga.
Sellii, Spreng ex Vog. Syn. Cass. 39 = speciosa.
semifalcata, Vell. Fl. Flum. 167 ; iv. t. 68 = multijuga.
semperflorens, DC. Cat. Hort. Monsp. 90 = biflora.
Senna, Linn. Sp. Pl. 377 = obovata, acutifolia.
sennoides, Jacq. Coll. i. 74 = bicapsularis.
sensitiva, Jacq. l. c. ii. 307 = sericea.
sensitiva, Roxb. Hort. Beng. 32 = mimosoides.
septentrionalis, *Vie. Elench. Pl. Hort. Dynegro*, 14.—Hab. ?
septentronalis, Zuccagni, in Roem. Collect. 141 = laevigata
sericea, *Sw. Fl. Ind. Occ.* 724.—Am. trop.
serpens, Gmelb. Cat. Pl. Cub. 80 = strigillosa.
serpens, *Linn. Syst. Nat.* ed. X. 1018.—Am. trop.
serzilla, Benth. in Mart. Fl. Bras. xv. ii. 135, in syn. = andromedea.
setigera, DC. Prod. ii. 499 = montana.
setosa, *Vog. Syn. Cass.* 51.—Bras.
sinuosa, *Lam. Encyc.* i. 648.—Ind. or. ; Malaya.
Sieberi, Presl, in Eken, Isis, xxi. (1828) 272 = glauca.
Sieberiana, *DC. Prod.* ii. 489.—Afr. trop.
silvestris, *Vell. Fl. Flum.* 169 ; iv. t. 78.—Bras.
simplicifolia, Desv. Journ. Bot. iii. (1814) 73 (= *Delaria ovalifolia*).—Bras.
sinensis, J. F. Gmel. Syst. 559 = chinensis.
singueana, Delile, in Caill. Voy. Meroë, Bot. iv. 27 (Cent. Pl Afr. 28) = goratensis.
Skinneri, *Benth. in Trans. Linn. Soc. xxvii.* (1871) 542.—Guatemal.
smaragdina, Macfad Fl. Jamaic. i. 347 = nictitans.
Sophera, *Linn. Sp. Pl.* 379.—Geront. trop.
Sophera, Wall. Cat. n. 5317 = occidentalis.
sophoroides, Collad. Hist. Cass. 138 = C. Sophera.
Sophora, Aucт. = C. Sophera.
sophoroides, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 153.—Am. trop.
speciosa, H. B. & K. Nov. Gen. et Sp. vi. 358 = speciabilis.
speciosa, Roxb. Hort. Beng. [31], nomen = glauca.
speciosa, *Schrad. in Goetl. Gel. Anz.* i. (1831) 718.—Bras.
spectabilis, *DC. Cat. Hort. Monsp.* 90.—Am. trop.

CASSIA :—
spinescens, *Hoffmgg. ex Vog. Syn. Cass.* 27.—Bras.
splendida, *Vog. l. c.* 17.—Bras.
Spruceana, *Benth. in Mart. Fl. Bras.* xv. ii. 92.—Bras.
stamínea, Vog. Syn. Cass. 14 = ferruginea.
stenocarpa, *Vog. l. c.* 68.—Bras.
stenophylla, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 535.—Cuba.
stipulacea, [*Soland. in*] *Ait. Hort. Kew.* ed. II. ii. 52. —Chili.
stipulata, G. Don, Gen. Syst. ii. 448 = flavicoma.
striata, Vog. Syn. Cass. 89 = splendida.
stricta, Schrank. Hort. Monac. t. 34 = C. Chamaecrista.
stricta, *Steud. Nom.* ed. II. i. 308.—Afr. austr.
strictifolia, *Benth. in Mart. Fl. Bras.* xv. ii. 145.— Bras.
strigillosa, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 581.—Ind. occ.
strobilacea, H. B. & K. Nov. Gen. et Sp. vi. 347 = reticulata.
Sturtii, *R. Br. in Sturt, Centr. Austral. App.* 14.— Austral.
subtrifolora, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 176.—Bras.
subulata, *Griseb. in Goett. Abh.* xix. (1874) 130.—Am. austr.
succedanea, Bell. ex DC. Prod. ii. 498 = marylandica.
suffruticosa, Koen. ex Roth, Nov. Sp. Pl. 218 = glauca.
sulcata, Bianco, Fl. Filip. ed. II. 236 = hirsuta.
sulcata, *DC. Cat. Hort. Monsp.* 90.—Am. austr.
sulphurea, DC. ex Collad. Hist. Cass. 84 = glauca.
sumatrana, Roxb. Hort. Beng. [31], nomen = siamea.
Sunsuk, Forsk. Fl. Aegypt. Arab. 86 = C. Tora.
supplex, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 163.—Bras.
surattensis, Burm. f. Fl. Ind. 97 = glauca.
surinamensis, *Dahnh. Cat. Pl. Hort. Camald.* ed. II. 18.—Guiana.
Swainsoni, *Benth. in Mart. Fl. Bras.* xv. ii. 167.—Bras.
Swartzii, Wikst. in Vet. Akad. Handl. (1825) 430 = virgata.
sylvestris, Auct. = silvestris.
Tagera, Lam. Encyc. i. 643 = C. Tora.
Tagera, *Linn. Syst. Pl.* 378.—Am. trop.
Tala, Desv. Journ. Bot. iii. (1814) 22 t. 73 = C. Tora.
Tarantan, H. B. & K. Nov, Gen. et Sp. vi. 348 = reticulata.
tarda, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 9.—Malaya.
tecta, *Vog. Syn. Cass.* 66.—Bras.
Telfairiana, Wall. Cat. n. 5324 = mimosoides.
tenella, *H. B. & K. Nov. Gen. et Sp.* vi. 365.— Venezuela.
tenella, Roxb. Hort. Beng. 82, nomen = mimosoides.
tenuifolia, *Vog. Syn. Cass.* 16.—Bras.
tenuisepala, *Benth. in Mart. Fl. Bras.* xv. ii. 164.— Am. trop.
tenuissima, Linn. Sp. Pl. 378 = biflora.
tephrosiaefolia, *Benth. in Mart. Fl. Bras.* xv. ii. 158. —Am. trop.
teretifolia, *A. Cunn. ex Lindl. in Ann. Sc. Nat. Sér.* II. xv. (1841) 60.—Austral.
teretifolia, Lindl. in Mitch. Three Exped. i. 289 = artemisioides.
teretiscandu, F. Muell. in Linnaea, xxv. (1852) 889 = artemisioides.
tetrafoliata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 438 = Langsdorffii.
tetraphylla, Collad. Hist. Cass. 130 = hispidula.
tetraphylla, Desv. journ. Bot. iii. (1814) 72 = Desvauxii.
tettensis, *Bolle, in Peters, Reise Mossamb. Bot.* 14. —Afr. trop.
texana, *Buckl. in Proc. Acad. Sc. Philad.* '1861' (1862) 452 = procumbens, *Linn. Herb.*
Thonningii, Sch. Prod. ii. 500 = C. Absus.
Thunbergiana, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 54 = mimosoides.
timorensis, *DC. Prod.* ii. 499.—Aa. trop. ; Austral.
tomentosa, Ehrenb. ex Hempr. ex Benth. in Proc. Linn. Soc. ii. (1855) 283 = holosericea.
tomentosa, *Linn. f. Suppl.* 231.—Am. trop.
tomentosa, Wall. Cat. n. 5804 = hirsuta.
Tora, *Linn. Sp. Pl.* 376.—Cosmop. trop.
toroides, Roxb. Hort. Beng. [31], nomen = C. Tora.
torosa, Cav. Descr. 191 = C. Sophera.
torulosa, Pois. Encyc. Suppl. ii. 757 = C. Sophera.
trachycarpa, *Vog. Syn. Cass.* 53.—Bras.
CASSIA :—
trachypus, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 122.—Bras.
tragacanthoides, *Mart. ex Benth. l. c.* 166.—Am. trop.
trichopoda, *Benth. l. c.* 163.—Am. trop.
triforo, Jacq. Hort. Schoenb. iv. 40 = C. Chamaecrista.
trifora, *Vahl, Eclog. Am.* iii. 11.—Ins. S. Cruc.
trinitatis, Benth. in Hook. Journ. Bot. ii. (1840) 77 = racemosa.
triniteiis, Reichb. ex DC. Prod. ii. 489 = spectabilis.
triasticula, *H. B. & K. Nov. Gen. et Sp.* vi. 367.—Am. austr.
tropica, Vell. Fl. Flum. 166 ; iv. t. 64 = laevigata.
tuberculata, *Collad. Hist. Cass.* 138.—Bras.
Tula, Desv. ex Steud. Nom. ed. II. i. 308 = C. Tora.
umbellata, Reichb. ic. Exot. t. 206 = australis.
undulata, *Benth. in Hook. Journ. Bot.* ii. (1840) 76. —Am. trop.
unifora. Mill. Gard. Dict. ed. VIII. n. 6 = sericea.
uniflora, *Spreng. Neue Entdeck.* i. 291.—Am. trop.
Urmenetea, *Phil. in Anal. Univ. Chil.* (1872) 708.— Chili.
ursina, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 151. —Am. trop.
Vauthieri, *Benth. l. c.* 138.—Am. trop.
velutina, *Vog. Syn. Cass.* 24.—Bras.
venenata, Rodsch. in G. F. W. Mey. Prim. Fl. Esseq. 167 = hirsuta.
venosa, Castigl. ex Zuccagni, in Roem. Collect. 140 = C. Chamaecrista.
venosa, Desv. Journ. Bot. iii. (1814) 72 = cytisoides.
venulosa, *Benth. in Mart. Fl. Bras.* xv. ii. 170.—Bras.
venusta, *F. Muell. Fragm.* i. 165.—Australia.
venustula, H. B. & K. Nov. Gen.et Sp. vi. 358 = biflora.
vernicosa, Clos, in C. Gay, Fl. Chil. ii. 244 = laevigata.
verrucosa, *Vog. Syn. Cass.* 38.—Bras.
versicolor, Benth. Pl. Hartw. 128 = incarnata.
versicolor, *Meyen, in Vog. Syn. Cass.* 29 ; *et Pl. Meyen*, 39.—Peruv.
vestita, *Vog. Syn. Cass.* 60.—Bras.
vicinifolia, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 544.—N. Granat.
villosa, *Mill. Gard. Dict.* ed. VIII. n. 4.—Mexic.
vimnea, *Linn. Syst. Nat.* ed. X. 1016.—Am. trop.
virgata, Mich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = chrysocarpa.
virgata, Slober, ex Steud. Nom. ed. II. i. 308 = glandulosa.
virgata, *Sw. Prod. Veg. Ind. Occ.* 66.—Ind. occ.
viscida, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indië*, iii. (1846) 80.—Malaya.
viscosa, *H. B. & K. Nov. Gen. et Sp.* vi. 360.—N. Granat.
viscosa, Schum. & Thonn. Beskr. Guin. Pl. 205 = C. Absus.
viscosipilosa, Steud. in Flora, xxvi. (1843) ii. 760 = viscosa.
Vogelsiana, *Schlecht. in Linnaea*, xii. (1838) 342.— Mexic.
vulgaris, *Roxb. ex Steud. Nom.* ed. II. i. 308.—Ind. or.
Wallichiana, DC. in Mém. Soc. Phys. Genèv. ii. (1824) 130 = mimosoides.
Wightiana, R. Grah. in Wall. Cat. n. 5329 = tomentosa.
Wislizeni, *A. Gray, Pl. Wright.* i. 60.—N. Mexic.
Wrightii, *A. Gray, l. c.* ii. 50.—Mexic.
xanthocoma, *Miq. Anal. Bot. Ind.* i. 10.—Borneo.
xiphoidea, Bertol. Fl. Guatim. 14 = biflora.
zambesica, *Oliver, Fl. Trop. Afr.* ii. 280.—Afr. trop.
zanzibarensis, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 77.—Afr. trop.
zygophylla, Benth. in Mitch. Journ. Trop. Austral. 288 = svenophila.

CASSIANA, Rafin. in Am. Monthly Mag. (1818) 266 = **Cassia**, Linn. (Legumin.).

CASSIDA, Tourn. ex Adans. Fam. ii. 188 (1763) = **Scutellaria**, Herm. (Labiat.).
alpina, Moench, Meth. Suppl. 144 = S. alpina.
altissima, Moench, l. c. 145 = S. altissima.
galericulata, Scop. Fl. Carn. ed. II. i. 430 = S. galericulata.
hastifolia, Scop. l. c. = S. hastifolia.
lateriflora, Moench, Meth. Suppl. 145 = S. lateriflora.
major, Gilib. Fl. Lituan. i. 90 = S. galericulata.
minor, Gilib. l. c. 91 = S. minor.
racemosa, Moench, Meth. 414 = S. peregrina.

CASSIDOCARPUS, Presl, ex DC. Prod. iv. 82 (1830) = **Asteriscium**, Cham. & Schlecht. (Umbellif.).
chilensis, Presl, l. c. = A. chilense.

CASSIERA, Raesach. Nom. ed. III. 42 (1797) sphalm. = **Canjera**, Juss. (Olacin.).

CASSINE, Linn. Gen. ed. I. 338 (1737). *CELASTRINEÆ*, Benth. & Hook. f. i. 393.
MAUROCENIA, Linn. Gen. ed. I. 85 (1737).
æthiopica, Eckl. & Zeyh. Enum. 128 = capensis.
æthiopica, *Thunb. Fl. Cap.* ii. 237.—Afr. austr.
affinis, *Sond. in Harv. & Sond. Fl. Cap.* i. 466.—Afr. austr.
albanensis, *Sond. l. c.* 467.—Afr. austr.
articulata, E. Mey. ex Sond. l. c. 467 = Celastrus procumbens.
Barba, Murr. Syst. Veg. ed. XIII. 248, sphalm. = sequ
barbara, *Linn. Mant.* ii. 220.—Afr. austr.
capensis, *Linn. l. c.*—Afr. austr.
caroliniana, Lam. Encyc. i. 652 = Ilex Cassine.
carolinians, Walt. Fl. Carol. 242 = Ilex ambigua.
Colpoon, Eckl. & Zeyh. Enum. 128 = affinis.
Colpoon, Thunb. Prod. Pl. Cap. 62 = capensis.
concava, Lam. Encyc. i. 652 = Celastrus lucidus.
Gongonha, St. Hil. in Guill. Arch. Bot. i. (1832) 31 = Villaresia mucronata.
corymbosa, Mill. Gard. Dict. ed. VIII. n. 1 = Ilex Cassine.
crocea, Presl, Bot. Bemerk. 87 = Elaeodendron capense
discolor, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 378 = Microtropis discolor.
dongonsia, *Spreng. Syst.* i. 939.—Ins. S. Doming.
excelsa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 376 = Ilex excelsa.
Gongonha, Mart. Reise Bras. i. 385 = Ilex Gongonha.
humilis, Hort. ex Steud. Nom. ed. II. i. 308 = Celastrus lucidus.
ilicifolia, Hochst. ex Sond. in Harv. & Sond. Fl. Cap i. 474 = Cassinopsis capensis.
integrifolia, Salisb. Prod. 173 = C. Maurocenia.
Kraussiana, Bernh. ex Harv. & Sond. Fl. Cap. i. 466 = capensis.
Kraussiana, *Hochst. in Flora*, xxvi. (1844) i. 305.— Afr. austr.
laevigata, Lam. Encyc. i. 652 = Pterocelastrus rostratus.
latifolia, *Eckl. & Zeyh. Enum.* 129.—Afr. austr.
magellanica, Lam. Illustr. ii. 92. n. 2690 = Maytenus magellanicus.
Maurocenia, *Linn. Sp. Pl.* ed. II. 385.—Afr. austr.
mucronata, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 455 = Cassinopsis capensis.
oleifolia, Pers. Syn. i. 327 = Celastrus oleoides.
oleoides, Lam. Encyc. i. 652 = Celastrus oleoides.
oppositifolia, *Mill. Gard. Dict.* ed. VIII. n. 3.—Hab.? *parvifolia*, E. Mey. ex Harv. & Sond. Fl. Cap. i. 466 = Elaeodendron croceum.
parvifolia, *Sond. l. c.* 466.—Afr. austr.
Peragua, Linn. Mant. ii. 220 = Ilex Cassine.
ramulosa, Rafin. Fl. Ludov. 110 = Ilex Cassine.
scandens, *Eckl. & Zeyh. Enum.* 128.—Afr. austr.
serratifolia, Salisb. Prod. 173 = capensis.
sphaerocarpa, Cels, ex Steud. Nom. ed. II. i. 308 = capensis.
xylocarpa, Vent. Choix, 28 = Elaeodendron xylocarpum.

CASSINIA, R. Br. in Trans. Linn. Soc. xii. (1817) 196. *COMPOSITÆ*, Benth. & Hook. f. ii. 318.
ACHROMOLAENA, Cass. in Dict. Sc. Nat. lvi. 223 (1828).
APALOCHLAMYS, Cass. l. c. 228 (1828).
CHROMOCHITON, Cass. l. c. 220 (1828).
RHYNEA, DC. in Deless. Ic. Sel. iv. t. 53 (1837).
aculeata, *R. Br. in Trans. Linn. Soc.* xii. (1817) 197. —Austral.
aculeata, A. Cunn. ex DC. Prod. v. 156 = Cunninghamii.
adenota, F. Muell. in Linnaea, xxv. (1852) 496 = aculeata.
affinis, R. Br. in Trans. Linn. Soc. xii. (1817) 197 = aculeata.
arcuata, *R. Br. l. c.*—Austral.
argophylla, DC. Prod. vi. 156 = Helichrysum ferrugineum.
aurea, *R. Br. in Trans. Linn. Soc.* xii. (1818) 197.— Austral.

CASSINIA :—
aurea, R. Br. in Ait. Hort. Kew. ed. II. v. (1813) 185 = Anaphalis tomentosa.
berberifolia, A. Cunn. ex DC. Prod. vi. 155 = denticulata.
cinerea, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 132 = leptophylla.
compacta, *F. Muell. Fragm.* i. 18.—Austral.
cuneifolia, A. Cunn. ex DC. Prod. vi. 155 = Helichrysum Cunninghamii.
Cunninghamii, DC. Prod. vi. 156 = Helichrysum Cunninghamii.
cupres, F. Muell. Fragm. iii. 139 = Helipterum polycephalum.
denticulata, *R. Br. in Trans. Linn. Soc.* xii. (1817) 127.—Austral.
fulvida, *Hook. f. Handb. New Zeal. Fl.* 145.—N. Zel.
glauophylla, *Cass. in Dict. Sc. Nat.* lvi. 210.—Austral.
hygrophila, A. Cunn. in DC. Prod. vi 156 = quinqueseria.
laevis, *R. Br. in Trans. Linn. Soc.* xii. (1818) 127.—Austral.
laevis, Endl. ex DC. l. c. = longifolia.
latifolia, A. Cunn. ex DC. Prod. vi. 155 = Helichrysum latifolium.
leptocephala, *F. Muell. Fragm.* iii. 138.—Austral.
leptophylla, *R. Br. in Trans. Linn. Soc.* xii. (1817) 126.—N. Zel.
longifolia, *R. Br. l. c.* 127.—Austral.
obovata, A. Cunn. in DC. Prod. vi. 155 = Helichrysum obcordatum.
paniculata, Behr & F. Muell. in Linnæa, xxv. (1852) 496 = arcuata.
patens, A. Cunn. ex DC. Prod. vi. 156 = Helichrysum Cunninghamii.
pholidota, F. Muell. Fragm. iii. 131 = Helichrysum pholidotum.
quinquefaria, *R. Br. in Trans. Linn. Soc.* xii. (1817) 127.—Austral.
quinquefaria, Sond. in Linnæa, xxv. (1852) 496 = Homea punctulata.
retorta, *A. Cunn. ex DC. Prod.* vi. 154.—N. Zel.
rosmarinifolia, A. Cunn. ex DC. l. c. 156 = laevis.
spathulata, Endl. ex Steud. Nom. ed. II. i. 309 = spectabilis.
spectabilis, *R. Br. in Trans. Linn. Soc.* xii. (1817) 128.—Austral.
subtropica, *F. Muell. Fragm.* i. 17.—Austral.
tenuifolia, *Benth. Fl. Austral.* iii 585.—Austral.
Theodori, *F. Muell. Fragm.* v. 148.—Austral.
uncata, A. Cunn. in DC. Prod. vi. 156 = aculeata.
Vauvilliersii, *Hook. f. Fl. N. Zel.* 133.—N. Zel.

CASSINIOLA, F. Muell. Fragm. iii. 139 (1863) = **Helipterum**, DC. (Compos.).
cupres, F. Muell. l. c. = H. polycephalum.

CASSINOPSIS, Sond. in Harv. & Sond. Fl. Cap. i. 473 (1860). *OLACINEÆ*, Benth. & Hook. f. i. 354.
capensis, *Harv. & Sond. l. c.* 474.—Afr. austr.
ciliata, *Baker, in Journ. Linn. Soc.* xx. (1884) 118.—Madag.
madagascariensis, *Baill. Adansonia*, xi. (1878–79) 180.—Madag.
tinifolia, *Harv. Thes. Cap.* ii. 44.—Afr. austr.

CASSIOPE, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 157. *ERICACEÆ*, Benth. & Hook. f. ii. 584.
ericoides, *D. Don, l. c.* 158.—Sibiria.
fastigiata, *D. Don, l. c.* 156.—Reg. Himal.
hypnoides, *D. Don, l. c.* 158.—Reg. arct. et subarct.
hypnoidioides, *D. Don, l. c.* 158.—Sibiria; Am. subarct. occ.
Mertensiana, *G. Don, Gen. Syst.* iii. 829.—Am. bor. occ.
oxycoccoides, *A. Gray, in Proc. Am. Acad.* xx. (1885) 300.—Kamtschat.
Redowskii, *G. Don, Gen. Syst.* iii. 829.—Sibiria. or.; Ross.
selaginoides, *Hook. f. & Thoms. in Hook. Kew Journ.* vi. (1854) 196. t. 4.—Reg. Himal.
Stelleriana, *DC. Prod.* vii. 611.—Alaska.
tetragona, *D. Don, in Edinb. N. Phil. Journ.* xvii. (1834) 158.—Reg. arct. et subarct.

CASSIPOUREA, Aubl. Pl. Gui. i. 328. t. 121 (1775). *RHIZOPHOREÆ*, Benth. & Hook. f. i. 682.
LEGNOTIS, Sw. Prod. Veg. Ind. Occ. 84 (1788).
TETA, Scop. Introd. 219 (1777).
africana, Benth. in Hook. Niger Fl. 341 = Weihea africana.
alba, *Griseb. in Goett. Abh.* vii. (1857) 223.—Ind. occ.
congensis, *R. Br. in Tuckey, Voy. Congo*, 18.—Afr. trop.
elliptica, *Poir. Encyc. Suppl.* ii. 131.—Jamaica.
guianensis, *Aubl. Pl. Gui.* i. 329. t. 121.—Am. austr.
gummiflua, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 123 = Dactylopetalum sessiliflorum.
lanceolata, *Tul. l. c.* 121.—Ins. Mascar.
leptoclada, *Tul. l. c.* 125.—Ins Mascar.
macrophylla, *DC. Prod.* iii. 34.—Bras.
madagascariensis, *DC. l. c.*—Madag.
microcarpa, *Tul. in Ann. Sc. Nat* Sér. IV vii. (1857) 162.—Ins. Mascar.
microphylla, *Tul l. c.* vi. (1856) 124.—Madag.
myrtocarya, *Tul. l. c.* viii. (1857) 162.—Madag.
ovata, *Tul. l. c.* vi. (1856) 121.—Ins. Comor.
phæotricha, *Tul l. c.* 122.—Madag.
quadriloculata, Spruce, ex Benth. in Mart. Fl. Bras. xii. 2. 429 = guianensis.
serrata, *Benth in Hook. Journ. Bot.* ii. (1840) 223.—Guiana.
Spruceana, *Benth. in Mart. Fl. Bras.* xii. 2. 429.—Am. trop.

CASSITHA, Hill, Veg. Syst. ix. 17 (1765) = **Cassytha**, Linn. (Laurin.).

CASSUMBIUM, Benth. & Hook. f. Gen. i. 404 (1862) = CUSSAMBIUM, Rumph. = **Schlechtera**, Willd. (Sapind.).

CASSUMUNAR, Colla, Nov. Sclt. Gen. 9 (1830) = **Zingiber**, Adans. (Scitamin.).
Kozburghii, Colla, l. c. 10 = Z. Cassumunar.

CASSUPA, Humb. & Bonpl. Pl. Aequin. i. 43 t. 12 (1808). *RUBIACEÆ*, Benth. & Hook. f. ii. 65.
laevis, *Triana, in Ann. Sc. Nat.* Sér. IV. ix. (1858) 44.—N. Granat.
verrucosa, *Humb. & Bonpl. Pl. Aequin.* i. 43. t. 12.—Mexic.

CASSUTHA, [J. Bauh.] Des Moul. Etud. Org. Cusc. 66 (1853) = **Cuscuta**, Linn. (Convolvul.).
americana, Des Moul. l. c. = Cusc. americana.
arabica, Des Moul. l. c. 72 = Cusc. arabica.
chrysocoma, Des Moul. l. c. 71 = Cusc. australis.
suaveolens, Des Moul. l. c. 66 = Cusc. racemosa.

CASSUVIUM, Lam. Encyc. i. 22 (1783) = **Anacardium**, Roth.
pomiferum, Lam. l. c. = A. occidentale.
reniforme, Blanco, Fl. Filip. ed. I. 322 = A. occidentale.
solitarium, Stokes, Bot. Mat. Med. ii. 475 = A. occidentale.

CASSYTA, Linn. Gen. ed. VI. 291 (1764) = **Cassytha**, Linn. (Laurin.).

CASSYTHA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 846 (1821) = **Cuscuta**, Tourn. (Convolvul.).
major, S. F. Gray, l. c. 846 = Cusc. europæa.

CASSYTHA, Linn. Sp. Pl. ed. I. 35 (1753). *LAURINEÆ*, Benth. & Hook. f. iii. 164.
CALODIUM, Lour. Fl. Cochinch. 247 (1790).
OZANTHUS, Rafin. Fl. Tellur. iv. 92 (1836).
ROMBUT, Rumph. ex Adans. Fam. ii. 384 (1768).
RUMPUTRIS, Rafin. Fl. Tellur. iv. 92 (1836).
VOLUTELLA, Forsk. Fl. Aegypt. Arab. 84 (1775).
americana, Nees, Syst. Laurin. 644 = filiformis.
aphylla, Raeusch. Nom. ed. III. 116 = filiformis.
brasiliensis, Mart. ex Nees, Syst. Laurin. 648 = filiformis.
capensis, *Meisn. in DC. Prod.* xv. 1. 254.—Afr. austr.
capillaris, *Meisn. l. c.* 252.—Zeylan.; Borneo.
Cassurinae, Nees, in Lehm. Pl. Preiss. i. 619 = glabella.
ceratopoda, Meisn. in DC. Prod. xv. 1. 257 = pomiformis.
ciliolata, *Nees, Syst. Laurin.* 646.—Afr. austr.
cornicula, *Burm. f. Fl. Ind.* 98.—Ind. or.
cornuta, *Nees, in Lehm. Pl. Preiss.* i. 620.—Austral.
digitata, Nees, l. c. = racemosa.

CASSYTHA :—
dispar, Schlecht. in Linnæa, xx. (1847) 578 = glabella.
dissitiflora, *Meisn. in Kjoeb. Vidensk. Meddel.* (1870) 146.—Bras.
filiformis, *Linn. Sp. Pl.* 35.—Cosmop. trop.
filiformis, Thunb. Prod. Pl. Cap. 78 = glabella.
flava, *Nees, in Lehm. Pl. Preiss.* i. 620.—Austral.
glabella, *R. Br. Prod.* 404.—Austral.
glabella, Sieber, ex Meisn. in DC. Prod. xv. 1. 256 = paniculata.
guineensis, Schum. & Thonn. Beskr. Guin. Pl 190 = filiformis.
melantha, *R. Br. Prod.* 404.—Austral.
micrantha, *Meisn. in DC. Prod.* xv. 1. 256.—Austral.
microcephala, Meisn. l. c. 258 = glabella.
Muelleri, Meisn. l. c. 257 = racemosa.
multiflora, Nees, in Lehm. Pl. Preiss. i. 621 = pomiformis.
nodiflora, *Meisn. in DC. Prod.* xv. 1. 252.—Austral.
paniculata, *R. Br. Prod.* 404.—Austral.
pharolasia, *F. Muell. Fragm.* v. 167.—Austral.
pilgera, Schlecht in Linnæa, xxi. (1848) 446 = pubescens.
pomiformis, *Nees*, in *Lehm. Pl. Preiss.* i. 620.—Austral.
pubescens, *R. Br. Prod.* 404.—Austral.
racemosa, *Nees, in Lehm. Pl. Preiss.* i. 621.—Austral.
remotiflora, F. Muell. ex Meisn. in DC. Prod. xv. 1. 256 = paniculata.
robusta, Meisn. l. c. 255 = melantha.
rubiginosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 171, nomen.—Afr. austr.
ruguloza, Meisn. in DC. Prod. xv. 1. 255 = pubescens.
subcapitata, Meisn. l. c. 253 = pomiformis, racemosa.
tasmanica, Meisn. l. c. 252 = pubescens.
triflora, E. Mey. in Drège, Zwei Pfl. Docum. 171 = capensis.
umbellata, Meisn. in DC. Prod xv. 1. 257 = racemosa ?
zeylanica, Gaertn. Fruct. i. 134 = filiformis ?

CASSYTHA, Mill. Gard. Dict. ed. VIII (1768) = **Rhipsalis**, Gaertn (Cactese).
baccifera, Mill ex DC. Prod iii. 476 = seq.
filiformis, Mill. Gard. Dict. ed. VIII = R. Cassytha.
polysperma, Ait. ex Gaertn. Fruct. i. 137 = R. Cassytha.

CASTALIA, Salisb. Parad. Lond. i. t. 34 (1805); et in Kon. & Sims, Ann Bot. ii. (1806) 71 = **Nymphæa**, Linn.
ampla, Salisb. ll. cc. t. 14, 73 = N. ampla.
caerulea, Tratt. in Flora, v. (1822) 604 = N. stellata.
edulis, Salisb. Parad. Lond. i. sub t. 14; et in Kon. & Sims, Ann. Bot. ii. (1806) 73 = N. Lotus.
Lotus, Tratt. in Flora, v. (1822) 598 = N. Lotus.
magnifica, Salisb. Parad Lond. i. t. 14 = N. Lotus.
mystica, Salisb. l. c.; et in Kon. & Sims, Ann. Bot. ii. (1806) 73 = N. Lotus.
pubescens, Hume, Bjdr. 48 = N. Lotus.
pudica, Salisb. in Kon. & Sims, Ann. Bot. ii. (1806) 72 = N. odorata.
pygmæa, Salisb. Parad. Lond. (1806) i. 2. t. 68 = N. tetragona.
rubra, Tratt. in Flora, v. (1822) 595 = N. Lotus.
Rudgeana, Tratt. l. c. 598 = N. tetragona.
sacra, Salisb. Parad. Lond. i. sub t. 14 = N. Lotus.
scutifolia, Salisb. l. c.; et in Kon. & Sims, Ann. Bot. ii. (1806) 72 = N. stellata.
speciosa, Salisb. in Kon. & Sims, Ann. Bot. ii. (1806) 72 = N. alba.
stellaris, Salisb. Parad. Lond. i. sub t. 14 = N. stellata.
stellata, Blume, Bjdr. 49 = N. stellata.
versicolor, Tratt. in Flora, v. (1822) 600 = N. stellata.

CASTALIS, Cass. in Dict. Sc. Nat. xxx. 382 (1824) = **Dimorphotheca**, Moench (Compos.).
Ventenati, Cass. l. c. = D. aurantiaca.

CASTANEA, Tourn. ex Linn. Syst. ed. I (1735). *CUPULIFERÆ*, Benth. & Hook. f. iii. 409.
CASTANOPHORUM, Neck. Elem. iii. 287 (1790) (Castanopborum).
acuminatissima, Blume, Mus. Bot. Lugd. Bat. i. 283 = Quercus Junghuhnii.
alnifolia, Nutt. Gen. Am. ii. 217 = pumila.
americana, G. Don, in Sweet, Hort. Brit. ed. III. 614 = sativa.
americana, Rafin. Fl. Ludov. 134; Now Sylva, 82 = sativa.
argentea, Blume, Bijdr. 535 = Castanopsis argentea.

CASTANEA:—
argyrophylla, Buek, in DC. Prod. Index, iv. 67 = Castanopsis argentea.
brevicuspis, Miq. Fl. Ind. Bat. i. 1. 866 = Castanopsis javanica.
Bungeana, Blume, Mus. Bot. Lugd. Bat. i. 284 = sativa.
buruana, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 377 = Castanopsis buruana.
castanicarpa, Oerst. l. c. = Castanopsis castanicarpa.
Chinquapin, Hort. ex C. Koch, Dendrol. ii. ii. 24 = pumila.
chinensis, Hassk. Cat. Hort. Bog. Alt. 73 = sativa.
chinensis, Spreng. Syst. iii. 856 = Castanopsis tribuloides.
chrysophylla, Dougl. ex Hook. Fl. Bor. Am. ii. 159 = Castanopsis chrysophylla.
concinna, Champ. ex Benth. in Hook. Kew Journ. vi. (1854) 115 = Castanopsis concinna.
cooperta, *Oerst. in Vidensk. Selsk. Skr.* V. ix. (1873) 374.—Ins. Philipp.
costata, Blume, Mus. Bot. Lugd. Bat. i. 284 = Castanopsis javanica.
crenata, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 324 = sativa.
divaricata, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Castanopsis argentea.
diversifolia, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 198.—Ind. or.
echidnocarpa, Hook. f. et Thoms. ex A. DC. in Prod. xvi. ii. 112 = Castanopsis tribuloides.
Fagus, Scop. Fl. Carn. ed. II. ii. 242 = Fagus sylvatica.
Falconeri, Hance, in Journ. Bot. xiii. (1875) 367 = Castanopsis armata.
fusfurella, Miq. Fl. Ind. Bat. Suppl. 852 = Castanopsis Hystrix.
glomerata, Blume, Mus. Bot. Lugd. Bat. i. 283 = Castanopsis sumatrana.
Hystrix, Hook. f. & Thoms. ex Miq. in Ann. Mus. Bot. Lugd. Bat. i. 119 = Castanopsis Hystrix.
indica, Roxb. Hort. Beng. 68; Fl. Ind. iii. 643 = Castanopsis indica.
inermis, Lindl. in Wall. Pl. As. Rar. ii. 6 = Castanopsis sumatrana.
japonica, Blume, Mus. Bot. Lugd. Bat. i. 284 = sativa.
javanica, Blume, Bijdr. 525 = Castanopsis javanica.
lanceaefolia, Kurz, For. Fl. Brit. Burma, ii. 482 = Quercus lanceaefolia.
latifolia, *Blume, Bijdr.* 526.—Java.
martabanica, Wall. Cat. n. 2764; Pl. As. Rar. ii. t. 107 = Castanopsis argentea.
microcarpa, Lindl. ex Wall. Cat. n. 8735 = Castanopsis tribuloides.
mollissima, *Blume, Mus. Bot. Lugd. Bat.* i. 286.—China.
montana, Blume, Bijdr. 526 = Castanopsis javanica.
nana, Mubl. Cat. 86 = pumila.
pumila, Hamb. Cat. Hort. Bog. Alt. 78 = sativa.
pumila, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. bor.
pumila, Siebold, ex Blume, Mus. Bot. Lugd. Bat. i. 286 = Castanopsis tribuloides.
purpurella, Miq. Fl. Ind. Bat. Suppl. 852 = Castanopsis Hystrix.
regia, Hance, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 280 = Quercus finas.
rhamnifolia, Kurz, For. Fl. Brit. Burma, ii. 481 = Castanopsis rhamnifolia.
rhamnifolia, *Oerst. in Vidensk. Selsk. Skr.* V. ix. (1873) 378.—Am. centr.
Roxburghii, Lindl. in Wall. Pl. As. Rar. ii. 6 = Castanopsis castanicarpa.
sativa, *Mill. Gard. Dict.* ed. VIII. n. 1.—Europ.; Japon.; Am. bor.
sempervirens, Kellogg, in Proc. Calif. Acad. i. 71, ed. II. (1873) 75 = Castanopsis chrysophylla.
wenf[olia], Blume, Mus. Bot. Lugd. Bat. i. 284 = Quercus Junghuhnii.
Sloanea, Mill. Gard. Dict. ed. VIII. n. 3 = Sloanea dentata.
spectabilis, Miq. Fl. Ind. Bat. i. 1. 866 = Castanopsis sumatrana.
sphaerocarpa, Lindl. in Wall. Pl. As. Rar. ii. 5 = Castanopsis armata.
stricta, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 225 = sativa.
sumatrana, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 376 = Castanopsis sumatrana.

CASTANEA:—
tribuloides, Lindl. in Wall. Pl. As. Rar. ii. 6 = Castanopsis tribuloides.
tribuloides, Wall. cat. n. 2765 B. = Quercus lanceaefolia.
Tungurrut, Blume, Bijdr. 526 = Castanopsis Tungurrut.
vesca, Gaertn. Fruct. i. 181. t. 37 = sativa.
vulgaris, Lam. Encyc. i. 708 = sativa.

CASTANELLA, Spruce, ex Benth. & Hook. f. Gen. i. 894 (1862). *SAPINDACEAE*, Benth. & Hook. f. i. c.
grandistamis, Planch. & Linden, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 365 = Paullinia granatensis.
paullinioides, *Spruce, ex Radlk. Monog. Serjan.* 76.—Bras.
riparia, *Spruce, ex Radlk. l. c.*—Peruv.

CASTANOCARPUS, Sweet, Hort. Brit. ed. II. 389 (1830) = **Castanospermum**, A. Cunn. (Leguminos.). *australis*, Sweet, l. c. = Castanosp. australe.

CASTANOLA, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 503 = **Connarus**, Linn.
trimervis, Llanos, l. c. = Conn. monocarpus.

CASTANOPHORUM, Neck. Elem. iii. 257 (1790) = **Castanea**, Tourn. (Cupulif.).

CASTANOPSIS, Spach, Hist. Vég. Phan. xi. 185 (1842). *CUPULIFERAE*, Benth. & Hook. f. iii. 409. CALLAICOCARPUS, Miq. Fl. Jungh. i. 18 (1851).
acuminatissima, A. DC. ex Hance, in Journ. Bot. i. (1863) 182 = Quercus lanceaefolia.
argentea, *A. DC. in Journ. Bot.* i. (1863) 182.—Burma; Malaya.
armata, *Spach, Hist. Vég. Phan.* xi. 185.—Reg. Himal.; Burma.
brevicuspis, A. DC. ex Hance, in Journ. Bot. i. (1863) 183 = javanica.
buruana, *Miq. Ann. Mus. Lugd. Bat.* i. 120.—Ins. Molucc.
castanicarpa, *Spach, Hist. Vég. Phan.* xi. 185.—Reg. Himal.
caudata, *Franch. Pl. Davidian.* i. 277.—Mongol.
chinensis, Hance, in Journ. Linn. Soc. x. (1869) 199 = tribuloides.
chrysophylla, *A. DC. in Journ. Bot.* i. (1863) 182.—Am. bor. occ.
concinna, *A. DC. l. c.*—Hongkong.
costata, A. DC. l. c.—javanica.
discocarpa, Hance, l. c. xvi. (1878) 201 = Quercus discocarpa.
echidnocarpa, A. DC. l. c. i. (1863) 162 = tribuloides.
echinocarpa, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 119, sphalm. = pmec.
Fabri, *Hance, in Journ. Bot.* xxii. (1884) 230.—China.
Falconeri, Hance, l. c. xiii. (1875) 368 = armata.
ferox, Spach, Hist. Vég. Phan. xi. 185 = tribuloides.
Fordii, *Hance, in Journ. Bot.* xxii. (1884) 230.—China.
Hystrix, *A. DC. l. c. i.* (1863) 182.—Reg. Himal.; Malaya.
indica, *A. DC. l. c.*—Reg. Himal.
inermis, Benth. & Hook. f. Gen. iii. 409 = sumatrana.
javanica, *A. DC. in Journ. Bot.* i. (1863) 182.—Malaya.
jucunda, *Hance, l. c.* xxii. (1884) 230.—China.
Lamontii, *Hance, l. c.* xiii. (1875) 369.—China.
lanceaefolia, Kurz, For. Fl. Brit. Burma, ii. 482 = Quercus lanceaefolia.
mitiflora, Hance, in Journ. Bot. xvi. (1878) 200 = sumatrana.
Pierrei, *Hance, l. c.* xiii. (1875) 369.—Cambodia.
rhamnifolia, *A. DC. Prod.* xvi. ii. 113.—Burma; Sumatra.
Scheferiana, *Hance, in Journ. Bot.* xvi. (1878) 200.—Sumatra.
sessilifolia, A. DC. l. c. i. (1863) 182 = Quercus Junghuhnii.
spectabilis, A. DC. l. c. = javanica.
sumatrana, *A. DC. l. c.*—Burma; Malaya.
tibetana, *Hance, l. c.* xiii. (1875) 367.—Tibet; China.
tribuloides, *A. DC. l. c. i.* (1863) 183.—Reg. Himal.; Burma.
triqueroma, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 502; in Flora, liii. (1870) 252 = javanica.
Tungurrut, *A. DC. in Journ. Bot.* i. (1863) 182.—Malaya.

CASTANOSPERMUM, A. Cunn. in Hook. Bot. Misc. i. (1830) 241. t. 51. *LEGUMINOSAE*, Benth. & Hook. f. i. 565.
CASTANOCARPUS, Sweet, Hort. Brit. ed. II. 389 (1830).
VIELLARDIA, Montr. in Mém. Acad. Lyon, x. (1860) x. 196.
australe, *A. Cunn. & Fraser, in Hook. i. c.*—Austral.

CASTANOSPORA, F. Muell. Fragm. ix. 92 (1875). *SAPINDACEAE*.
Alphandi, *F. Muell. l. c.*—Austral.

CASTELA, Turp. in Ann. Mus. Par. vii. (1806) 78. t. 5. *SIMARUBEAE*, Benth. & Hook. f. i. 310.
alaternifolia, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 569.—Chili.
coccinea, *Griseb. in Goett. Abh.* xix. (1874) 107.—Reg. Argent.
depressa, *Turp. in Ann. Mus. Par.* vii. (1806) 79. t. 5 A.—Ins. S. Doming.
erecta, *Turp. l. c.* 80. t. 5 B.—Ind. occ.
Galapagei, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 229.—Ins. Galap.
laevigata, *Zipp. ex Span. in Linnaea*, xv. (1841) 186, nomen.—Hab.?
lychnophoroides, *Liebm. in Kjoeb. Vidensk. Meddel.* (1853) 111.—Mexic.
Nicholsoni, *Hook. Bot. Misc.* i. (1830) 271. t. 56.—Ind. occ.
retusa, *Liebm. in Kjoeb. Vidensk. Meddel.* (1853) 110.—Mexic.
tortuosa, *Liebm. l. c.* 108.—Mexic.
Tweediei, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 569.—Reg. Argent.

CASTELIA, Cav. in Anal. Cienc. Nat. iii. (1801) 134 = **Priva**, Adans. (Verbenac.).
cuneata-ovata, Cav. l. c. 135 = laevis.

CASTELIA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 108 = **Castela**, Turp. (Simarub.).

CASTELLIA, Tin. Pl. Rar. Sicul. 17 (1817) = **Festuca**, Linn. (Gramin.).
tuberculata, Tin. l. c. 18 = F. tuberculata.

CASTELNAVIA, Tul. & Wedd. in Ann. Sc. Nat. Sér. III. xi. (1849) 108. *PODOSTEMACEAE*. Benth. & Hook. f. iii. 114.
CARAJASIA, Wedd. in DC. Prod. xvii. 84 (1873).
fimbriata, *Tul. & Wedd. l. c.*—Bras.
fluitans, *Tul. & Wedd. l. c.* 109.—Bras.
fruticulosa, *Tul. & Wedd. l. c.* 110.—Bras.
monandra, *Tul. & Wedd. l. c.* 109.—Bras.
multipartita, *Tul. & Wedd. l. c.*—Bras.
orthocarpa, *Tul. & Wedd. l. c.* 110.—Bras.
Princeps, *Tul. & Wedd. l. c.* 108.—Bras.
pusillina, *Tul. & Wedd. l. c.* 109.—Bras.
serpens, *Tul. & Wedd. l. c.*—Bras.

CASTIGLIONIA, Ruiz & Pav. Prod. 139. t. 37 (1794) = **Jatropha**, Linn. (Euphorbiac.).
lobata, Ruiz & Pav. l. c.; Syst. Veg. 277 = J. Curcas.

CASTILLA, Cerv. in Supl. Gaz. Lit. Mexico (Jul. 7, 1794), 7 = **Castillon**, [1] Cerv.[1] (Urticac.).

CASTILLEJA, Mutis, ex Linn. f. Suppl. 47 (1781). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 973. EUCHROMA, Nutt. Gen. Am. ii. 54 (1818).
acuminata, Spreng. Syst. iii. 778 = pallida.
acuminata, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 231 = pallida.
affinis, Benth. Pl. Hartw. 329 = stenantha.
affinis, *Hook. & Arn. Bot. Beech. Voy.* 154.—Calif.
ambigua, Hook. & Arn. l. c. = Orthocarpus zastrulis.
angustifolia, G. Don, Gen. Syst. iv. 616 = parviflora.
angustifolia, A. Gray, in Torr. Bot. Mex. Bound. 118 = integra.
angustifolia, *Mart. & Gal. in Bull. Acad. Brux.* xii. ii. (1845) 10.—Mexic.
anthemoidifolia, Benth. in DC. Prod. x. 528 = tenuiflora.
arvensis, *Cham. & Schlecht. in Linnaea*, v. (1830) 108.—Mexic.
Breadburii, G. Don, Gen. Syst. iv. 616 = parviflora.
breviflora, *Benth. in DC. Prod.* x. 534.—Venezuela.
brevifolia, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiv. (1862) 250.—Am. bor. occ.

CASTILLEJA:—
canderi, Durand & Hilg. in Pacif. Rail. Rep. v. 12 = linariæfolia.
canescens, *Benth. in DC. Prod.* x. 583.—Mexic.
cinerea, *A. Gray, in Proc. Am. Acad.* xix. (1883) 93.—Calif.
coccinea, Dougl. in Lindl. Bot. Reg. t. 1136 = parviflora.
coccinea, *Spreng. Syst.* ii. 775.—Am. bor.
communis, *Benth. in DC. Prod.* x. 529.—Am. trop.
coronopifolia, *Vent. Choix*, 59.—N. Granat.
deserlorum, Geyer, in Hook. Kew Journ. v. (1853) 258 = parviflora.
divaricata, *Benth. in DC. Prod.* x. 584.—Venezuela.
Douglasii, Benth. l. c. 580 = parviflora.
fasciculata, *Linn. f. Suppl.* 293.—N. Granat.
flava, *S. Wats. in Bot. King, Exped.* 280.—Am. bor. occ.
foliolosa, *Hook. & Arn. Bot. Beech. Voy.* 164.—Calif
fulgens, Nutt. ex A. Gray, in Torr. Bot. Mex. Bound. 119 = linariæfolia ?
gracilis, *Benth. in DC. Prod.* x. 528.—Mexic.
grandiflora, Spreng. Syst. ii. 755 = sessiliflora.
hirsuta, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 29.—Mexic.
hispida, Benth. in Hook. Fl. Bor. Am. ii. 105 = parviflora.
hyssopifolia, Willd. ex G. Don, Gen. Syst. iv. 615 = lithospermoides.
indivisa, *Engelm. in Engelm. & A. Gray, Pl. Lindh.* i. 47.—Texas.
integra, *A. Gray, in Torr. Bot. Mex. Bound.* 119.—N. Mexic.
integrifolia, H. B. & K. Nov. Gen. et Sp. ii. 330 = canescens.
integrifolia, *Linn. f. Suppl.* 293.—Am. trop.
insaniens, *Orrst. in Kjoeb. Vidensk. Meddel.* (1858) 27. —Costa Rica.
laciniata, *Hook. & Arn. Bot. Beech. Voy.* 60.—Chili.
lanata, *A. Gray, in Torr. Bot. Mex. Bound.* 118.—N. Mexic.
latifolia, *Hook. & Arn. Bot. Beech. Voy.* 164.—Calif.
laxa, *A. Gray, in Torr. Bot. Mex. Bound.* 119.—N. Mexic.
Lemmoni, *A. Gray, Syn. Fl. N. Am.* ii. 1. 297.—Calif.
linariæfolia, *Benth. in DC. Prod.* x. 583.—Mexic.
Lindheimeri, *A. Gray, Syn. Fl. N. Am.* ii. 1. 298.—Texas.
lineolata, *A. Gray, l. c.* 299.—Am. bor. occ.
lithospermoides, *H. B. & K. Nov. Gen. et Sp.* ii. 331. t. 164.—Mexic.
longibracteata, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 26.—Mexic.
longiflora, *Kunze, in Linnæa*, xvi. (1842) 812.—Mexic.
macrocarpa, Benth. Scroph. Ind. 18 = latifolia.
minlata, *Dougl. in Hook. Fl. Bor. Am.* ii. 106.—Am. bor. occ.
minor, *A. Gray, Bot. Calif.* i. 573.—Am. bor. occ.
monarensis, *H. B. & K. Nov. Gen. et Sp.* 329.—Mexic.
nubigena, *H. B. & K. l. c.* 330.—Ecuador.
oblongifolia, *A. Gray, Syn. Fl. N. Am.* ii. 1. 296.—Calif.
obovata, Benth. in DC. Prod. x. 528 = hirsuta.
occidentalis, Torr. in Ann. Lyc. N. York, ii. (1828) 230 = pallida.
Orinoban, Benth. in DC. Prod. x. 583 = pectinata.
pallida, *Kunth, Syn. Pl. Aequin.* ii. 100.—Sibir.; Am. bor.
pallida, Spreng. Syst. ii. 774 = septentrionalis, sibirica.
parviflora, *Bong. in Mém. Acad. Pétersb.* Sér. VI. v. 11. (1840) 157.—Am. bor. occ.
pectinata, *Mart. & Gall. in Bull. Acad. Brux.* xii. 11. (1845) 27.—Mexic.
plagiotoma, *A. Gray, in Proc. Am. Acad.* xix. (1883) 93.—Calif.
pumila, Wedd. Chlor. And. t. 61 = fastigiata.
purpurea, *G. Don, Gen. Syst.* iv. 615.—Am. bor. occ.
Schaffneri, *Hemsl. Biol. Centr. Am. Bot.* ii. 402.—Mexic.
scorzoneræfolia, *H. B. & K. Nov. Gen. et Sp.* ii. 331. —Mexic.
septentrionalis, *Lindl. Bot. Reg.* t. 926.—Am. bor.
serrata, *Hort. ex Lond. Hort. Brit. Suppl.* iii. 511.—Hab. ?
sessiliflora, *Pursh, Fl. Am. Sept.* ii. 735.—Am. bor. occ.
sibirica, Lindl. Bot. Reg. t. 926 = pallida.

CASTILLEJA:—
speciosa, Mart. & Gal. in Bull. Acad. Brux. xii. 11. (1845) 30 = scorzoneræfolia.
stenantha, *A. Gray, Syn. Fl. N. Am.* ii. 1. 295.—Am. bor. occ.
stricta, *Benth. in DC. Prod.* x. 584.—Ecuador.
tenuiflora, *Benth. Pl. Hartw.* 52.—Mexic.
tenuifolia, *Mart. & Gal. Bull. Acad. Brux.* xii. 11. (1845) 30.—Mexic.
toluccensis, Cham. & Schlecht. in Linnæa, ii. (1827) 579 = parviflora ?
toluccensis, *H. B. & K. Nov. Gen. et Sp.* ii. 329.—Mexic.
tomentosa, A. Gray, in Torr. Bot. Mex. Bound. 118 = integra.
trifida, Willd. ex Walp. Rep. iii. 394 = rubigena.
viscidula, *A. Gray, Syn. Fl. N. Am.* ii. 1. 297.—Am. bor. occ.

CASTILLOA, Cervant. in Supl. Gaz. Lit. Mexic. (Jul. 2, 1794) 7 (Castilla). *URTICACEAE*, Benth. & Hook. f. iii. 372.
costaricana, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 319 = Costa Rica.
elastica, *Cerv. in Supl. Gaz. Lit. Mexico* (Jul. 2, 1794), 7.—Mexic.
Markhamiana, *Collins, Rep. Caoutch. Comm.* 12. t. 3. —Am. trop.

CASTRA, Vell. Fl. Flum. 242 (1825); viii. 79–81 = **Trixis**, P. Br. (Compos.).
illustris, Vell. Fl. Flum. 343; viii. t. 81 = T. brasiliensis.
nobilis, Vell. l. c. 344; viii. t. 81 = T. glaberrima.
pruriens, Vell. l. c. 343; viii. t. 80 = T. mollissima.
regia, Vell. l. c. 343; viii. t. 79 = T. divaricata

CASTRAETALLA, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 189. *MELASTOMACEAE*, Benth. & Hook. f. i. 736.
piloselloides, *Naud. l. c.*—N. Granat.

CASTREA, A. St. Hil. Morphol. 451 (1840) = **Phoradendron**, Nutt. (Loranthac.).
falcata, St. Hil. l. c.—Bras. !

CASTRONIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 = **Helicia**, Lour. (Protenc.).

CASUARINA, Linn. Amœn. Acad iv. 143, nomen (1759); Adans. Fam. ii. 481 (1768); Forst. Char. Gen. 108. t. 52 (1776). *CASUARINEAE*, Benth. & Hook. f. iii. 402.
acuaria, *F. Muell. in Journ. Bot.* v. (1867) 212.—Austral.
acutivalvis, *F. Muell. Fragm.* x. 61.—Austral.
africana, Lour. Fl. Cochinch. ii. 549 = equisetifolia.
angulata, *Poiss. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 49.—N. Caled.
Baxteriana, Miq. Rev. Crit. Casuar. 37. t. 3 C = distyla.
bicudatata, *Benth. Fl. Austral.* vi. 202.—Austral.
Brunoniana, *Miq. Rev. Crit. Casuar.* (1848) 21. t. 1. C. —Java.
Chamæcyparis, *Poiss. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 52.—N. Caled.
collina, *Poiss. ex Pancb. & Sebert, Not. Bois Nouv. Caled.* 174.—N. Caled.
cornicuiata, *F. Muell. Fragm.* x. 62.—Austral.
cristata, Miq. Rev. Crit. Casuar. 70. t. 10. A = stricta.
Cunninghamiana, *Miq. l. c.* 56. t. 6 A.—Austral.
Decaisneana, *F. Muell. Fragm.* i. 61.—Austral.
decussata, *Benth. Fl. Austral.* vi. 200.—Austral.
Deplancheana, *Miq. in DC. Prod.* xvi. ii. 342.—N. Caled.
distyla, *Vent. Jard. Cels*, 63. t. 62.—Austral.
Drummondiana, *Miq. Rev. Crit. Casuar.* 26. t. 1 D.—Australia.
dumosa, A. Cunn. ex Miq. in DC. Prod. xvi. ii. 338 = distyla.
echinata, R. Br. ex Benth. Fl. Austral. vi. 202 = thuyoides.
equisetifolia, Blanco, Fl. Filip. ed. I. 601 = sumatrana.
equisetifolia, *Linn. Amœn. Acad.* iv. 143; *Forst. Char. Gen.* 108. f. 53.—Malaya; Ins. Pacif.

CASUARINA:—
excelsa, Dehnhardt, ex Miq. Rev. Crit. Casuar. 28. t. 1 F = equisetifolia.
excelsa, Salisb. Prod. 2 = stricta.
Fraseriana, *Miq. Rev. Crit. Casuar.* 59. t. 6. D.—Austral.
glauca, *Sieber, in Spreng. Syst. iii.* 803.—Austral.
Gunnii, Hook. f. ex Miq. in Neder. Kruidk. Arch. iv. (1859) 100 = stricta.
hexagona, *Dehnhardt, in Revista Napolitana*, i. 111. 176.—Austral.
humilis, Otto & Dietr. Allg. Gartenz. ix. (1841) 163 = Fraseriana.
incana, A. Cunn. ex Benth. Fl. Austral. vi. 197 = equisetifolia.
indica, Pers. Syn. ii. 581 = equisetifolia.
inophloia, *F. Muell. & Bailey, in Melb. Chemist* (Apr. 1869).—Austral.
Junghuhniana, *Miq. Pl. Jungh.* 7.—Malaya.
laterifiora, Poir. Encyc. Suppl. ii. 658 = equisetifolia.
Lehmanniana, Miq. in Lehm. Pl. Preiss. i. 639 = humilis.
lepidophloia, *F. Muell. Fragm.* x. 115.—Austral.
leptoclada, Miq. Rev. Crit. Casuar. 41. t. 4. C = suberosa.
leucodon, *Poiss in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 52.—N. Caled.
littoralis, Salisb. Prod. 2 = equisetifolia.
lucida, *Dehnhardt, in Revista Napolitana*, i. 111. 175. —Austral.
lugubris, Salisb. Prod. 2 = tornlosa
macrocarpa, A. Cunn. ex Miq. in DC. Prod. xvi. ii. 338 = stricta.
Mertensiana, Rupr. ex Miq. l. c. 339 = equisetifolia.
microstachya, *Miq. in Lehm. Pl. Preiss.* i. 642.—Austral.
Miquelii, Hook. f. ex Miq. in DC. Prod. xvi. ii. 337 = suberosa.
moesta, *F. Muell.* ex Miq. in Neder. Kruidk. Arch. iv. (1859) 98 = suberosa.
montana, *Leschen. ex Miq. in Zoll. Syst. Verz. Ind. Archip.* 86.—Malaya.
Muelleriana, Miq. in Neder. Kruidk. Arch. iv. (1859) 99 = distyla.
muricata, Herb. Paris. ex Miq. in DC. Prod. xvi. ii 335 = montana.
muricata, Roxb. Hort. Beng. 66; Fl. Ind. iii. 519 = equisetifolia.
nana, A. Cunn. ex Benth. Fl. Austral. vi. 200 = humilis.
nana, Schlecht. in Linnæa, xx. (1847) 574 = Fraseriana.
nana, *Sieber, ex Spreng. Syst.* iii. 804.—Austral.
nodiflora, *Forst. f. Prod. Fl. Ins. Austr.* 64.—Ins. N. Hebrid.
obesa, Miq. in Lehm. Pl. Preiss. i. 643 = Huegeliana.
obtusa, Hort. ex Miq. in DC. Prod. xvi. ii. 334 = glauca.
oxyclada, Miq. l. c. 344 = acuaria.
paludosa, Sieber, ex Spreng. Syst. iii. 803 = distyla.
Pauper, *F. Muell. ex Miq. in Neder. Kruidk. Arch. iv.* (1859) 100.—Austral.
Preissiana, Miq. in Lehm. Pl. Preiss. i. 640 = humilis.
pumila, Otto & Dietr. in Allg. Gartenz. ix. (1841) 162 = distyla.
quadridentata, Desf. Tabl. Cat. Hort. Par. ed. II. 346 = quadrivalvis
quadrivalvis, Labill. Nov. Holl. Pl. ii. 67. t. 218 = stricta.
ramulifiora, Otto & Dietr. Allg. Gartenz. ix. (1841) 163 = humilis.
repens, Hort. ex Hoffmgg. Verz. Pfl. Nachtr. 58 = equisetifolia.
rigida, Miq. in DC. Prod. xvi. ii. 337 = distyla.
rigida, Miq. Rev. Crit. Casuar. 61. t. 7. D = suberosa.
rugosa, *Roxb. Hort. Beng.* 66, nomen.—Ind. or. ?
Rumphiana, *Miq. in Flora*, xlviii. (1865) 23, 38.—Amboina.
selaginoides, Miq. in Lehm. Pl. Preiss. i. 643 = humilis.
sparsa, Tausch, in Flora, xxii. (1839) 480 = equisetifolia ?
stricta, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 320.—Austral.
stricta, Miq. in DC. Prod. xvi. ii. 336 = distyla.

CASUARINA:—
suberosa, *Otto & Dietr. Allg. Gartenz.* (1841) 155.—Austral.
sumatrana, *Jungh. in Horv. & De Vriese, Tijdschr.* xi. (1844) 115.—Sumatra.
tenuissima, Sieber, ex Spreng. Syst. iii. 804 = torulosa.
taphrosperma, Lehm. Hort. Hamb. ex Miq. Rev. Crit. Casuar. 31 = humilis.
thuyoides, *Miq. in Lehm. Pl. Preiss.* i. 641.—Austral.
tortuosa, Henry, in Nov. Act. Nat. Cur. xx. 1. (1847) 217. t. 19 = stricta ?
torulosa, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 320 —Austral.
torulosa, Miq. Rev. Crit. Casuar. 75. t. 11 B = glauca.
torulosa, Miq. in Lehm. Pl. Preiss. i. 639 = Fraseriana.
trichodon, *Miq. l. c.* 641.—Austral.
truncata, Willd. Enum. Hort. Berol. Suppl. 68 = equisetifolia ?
verticillata, Lam. Encyc. ii. 502 = nodiflora.
Webbiana, *Miq in Flora*, xlviii. (1865) 33.—Borneo; N. Caled. ?

CATABROSA, Beauv. Agrost. 97. t. 19. f. 8 (1812). *GRAMINEÆ*, Benth. & Hook. f. iii. 1186.
agrostidea, Presl, Fl. Sicul. p. xliv = Airopsis agrostidea.
algida, Fries, Nov. Fl. Suec. Mant. iii. 174 = Phippalia algida.
altaica, Boiss. Fl. Orient. v. 579 = Colpodium altaicum.
antarctica, *Hook. f. Fl. Antarct.* 102.—N. Zel.
aquatica, *Beauv. Agrost.* 97. t. 19. f. 8.—Reg. bor. temp. Balansæ, *Boiss. in Bull. Soc. Bot. Fr.* xxi. (1874) 15. —As. Min.
Capnii, *Franch. in Ann. Sc. Nat. Sér.* VI. xviii. (1884) 272.—Turkest.
concinna, T. Fries, in Öfver. Vet. Akad. xxvi. (1869) 140 = Phippalia algida ?
distans, Link, ex Heynh. Nom. ii. 126 = Glyceria distans.
elodes, Auct. = helodes.
Engelmanni, Link, ex Heynh. l. c. = Glyceriae sp. ?
—Hab. ?
fibrosa, *Boiss. Fl. Orient.* v. 577.—Armenia.
frigida, *Phil. Fl. Atac. Bot.*—Chili.
glaucescens, Phil. in Anal. Univ. Chil. (1873) 569 = = Glyceria glaucescens.
helodes, Roem & Schult. Syst. ii. 686 = Poa nemoralis.
humilis, *Trin. Fund. Agrost.* 136.—Persia; Reg. Casp.
latifolia, Fries, Summ. Veg. Scand. i. 245 = Arctagrostis latifolia.
magellanica, Hook. f. Fl. Antarct. 387 = Glyceria magellanica.
micrantha, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 496 = Eragrostis interrupta.
minuta, Trin. Fund. Agrost. 136 = Aira minuta.
ochroleuca, Dum. Obs. Gram. Belg. 108 = aquatica.
parviflora, *Boiss. Fl. Orient.* v. 578.—Persia.
pontica, *Balansa, in Bull. Soc. Bot. Fr.* xxi. (1874) 15. —As. Min.
pumila, Roem. & Schult. Syst. ii. 698 = Poa annua.
scabra, *Kunth, Rev. Gram.* i. 119.—Hab. ?
stricta, *Link, ex Stend. Nom.* ed. II. i. 310.—Hab. ?
tenuifolia, *Ş. Presl, in Rel. Haenk.* i. 256.—Chili.
vaillanti, T. Fries, in Öfver. Vet. Akad. Stockh. xxvi. (1869) 142 = Phippsia algida.
variegata, *Boiss. Diagn.* Ser. i. v. 71.—As. Min.
vernicolor, Boiss. Fl. Orient. v. 579 = Colpodium Stevenii.
verticillata, *Beauv. Agrost.* 157, nomen.—Hab. ?
wilfoides, Anders. ex Malm. in Öfver. Vet. Akad. Stockh. xix. (1862) 254 = Glyceria wilfoides.
violacea, Boiss. Diagn. Sér. 1. vii. 126 = Colpodium violaceum.
viridula, Roem. & Schult. Syst. ii. 697 = Aira viridula.

CATABROSIA, Roem. & Schult. Syst. ii. 696 (1817) = **Catabrosa**, Beauv. (Gramin.).

CATACHAENIA, Griseb. in Goett. Abh. ix. (1861) 54 = **Miconia**, Ruiz & Pav. (Melastom.).
alternifolia, Griseb. l. c. 55 = M. Grisebachii.
parviflora, Griseb. Cat. Pl. Cub. 102 = M. Grisebachii.
rufa, Griseb. l. c. = M. rufa.

CATACHAETUM, Hoffmgg. Verz. Orch. 22 (1842); ex Lindau, xvi. (1842) Litt. 321 = **Catasetum**, Rich. (Orchid.).

CATACLINE, Edgew. in Journ. As. Soc. Bengal, xvi. (1847) 1214 = **Tephrosia**, Pers. (Legumin.).
sericea, Edgew. l. c. = T. sericea.

CATACOMA, Walp. Rep. i. 243 (1842) = CATOCOMA, Benth. = **Bredemeyera**, Willd. (Polygal.).

CATAGYNA, Beauv. in Lestib. Essai Fam. Cyp. 26 (1819) = **Eriospora**, Hochst. (Cyperac.).

CATAKIDOZAMIA, T. Hill, in Gard. Chron. (1865) 1107 = **Macrozamia**, Miq. (Cycad.).
Hopei, T. Hill, l. c. = M. Perowskiana.
Macleayi, Hill, ex Miq. in Verh. Akad. Wetensch. II. iii. (1869) 56 = M. Perowskiana.

CATALIUM, Buch.-Ham. ex Wall. Cat. sub n. 4880 (1831) = **Carallia**, Roxb. (Rhizoph.).
nitidum, Buch.-Ham. l. c. = Carallia lanceafolia.

CATALPA, Scop. Introd. 170 (1777). *BIGNONIACEÆ*, Benth. & Hook. f. ii. 1041
CATALPIUM, Rafin. Princ. Somiol. 27 (1814); Fl. Ludov. 139 (1817).
CUMBALU, Adans. Fam. ii. 199 (1763).
bignonioides, *Walt. Fl. Carol.* 64.—Am. bor.
Bungei, *C. A. Mey. in Bull. Sc. Acad. Pétersb.* ii. (1837) 51.—China.
cassinoides, *Spreng. Syst.* i. 70.—Bras.
communis, Dum.-Cours. Bot. Cult. ed. II. iii. 242 = bignonioides.
cordifolia, Jaume, in Duham. Arb. ed. Nov. ii. 13. t. 5 = bignonioides.
hirsuta, *Spreng. Syst.* i 70.—Bras.
Kaempferi, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 142.—Japon.
longisiliqua, Cham. in Linnæa, vii. (1832) 720 = longissima.
longissima, *Sims, Bot. Mag.* sub t. 1094.—Ind. occ.
microphylla, *Spreng. Syst.* i. 70.—Ins. S. Doming.
ovata, G. Don, Gen. Syst. iv. 280 = Kaempferi.
Pottsii, *Seem. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 321.—Mexic.
punctata, *Griseb. Cat. Pl. Cub.* 102.—Cuba.
purpurea, *Griseb. l. c.*—Cuba.
speciosa, *Warder, ex Engelm. in Coult. Bot. Gaz.* v. (1880) 1.—Am. bor.
syringæfolia, Bunge, Enum. Pl. Chin. Bor. 45 = Bungei.
syringifolia, Sims, Bot. Mag. t. 1094 = bignonioides.
ternifolia, Cav. Desc. 26 = bignonioides.

CATALPIUM, Rafin. Princ. Somiol. 27 (1814); Fl. Ludov. 139 (1817) = **Catalpa**, Scop. (Bignoniac.).
amena, Rafin. Fl. Ludov. 139 = Catalpa bignonioides.

CATAMIXIS, Thoms. in Journ. Linn. Soc. ix. (1865) 342. t. 4. *COMPOSITÆ*, Benth. & Hook. f. ii. 494.
baccharoides, *Thoms. l. c.* 343. t. 4.—Himal. bor. occ.

CATANANCHE, Linn. Syst. ed. I (1735). *COMPOSITÆ*, Benth. & Hook. f. ii. 505.
PITTOCEPHALUM, Sch. Bip. in Bonplandia (1860) 369.
arenaria, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 253.—Algeria.
caerulea, Georgi, Beschr. Russ. Reich. iii. iv. 1217 = Centaurea pullata.
caerulea, *Linn. Sp. Pl.* 812.—Reg. Mediterr. occ. cæspitosa, *Desf. Fl. Atlant.* ii. 288.—Afr. bor.
carphalepis, Nym. Consp. 473 = caerulea.
graeca, Boiss & Chaub. Nouv. Fl. Pélop. 55 = Hymenonema laconicum.
graeca, Linn. Sp. Pl. 813 = Hymenonema Tournefortii.
lutea, *Linn. l. c.* 812.—Reg. Mediterr.
montana, *Coss. & Dur. in Bull Soc. Bot. Fr.* iii. (1856) *Bibl.* 743.—Afr. bor.
propinqua, *Pomel, Nouv. Mat. Fl. Atl.* 19.—Algeria.
Webbii, Sch. Bip. in Bonplandia, viii. (1860) 369 = montana.

CATANGA, Stend. Nom. ed. II. i. 311 (1840), sphalm. = **Cananga**, Aubl.

CATAPODIUM, Link, Hort. Berol. i. 86, 44 (1827) = **Festuca**, Linn. (Gramin.).
aquaticum, Willk. in Willk. & Lange, Prod. Fl. Hisp i. 77 = Calabrosa aquatica.
durum, Link, in Linnæa, xvii. (1843) 899 = Sclerochloa dura.
fasciforme, Nees, ex Benth. in Journ. Linn. Soc. xix (1881) 128 = Tripogon bromoides.
Halleri, Reichb. Fl. Germ. Excurs. 140 = F. Poa.
loliaceum, Link, Hort. Berol. i. 45 = Desmazeria loliacea.
nepalensis, Link, ex Steud. Nom. ed. II. i. 311 (quod ?) *pungens*, Boiss. Fl. Orient. v. 635.—Afghan.
Salzmanni, Boiss. l. c. 634.—Reg. Mediterr.
siculum, Link, Hort. Berol. ii. 194 = Desmazeria sicula
tuberculosum, Moris, in Atti 2da Riun. Sc. Ital. (1841) 481 = F. tuberculosa.

CATAPPA, Gaertn. Fruct. ii. 206. t. 127 (1791) = **Terminalia**, Linn. (Combretac.).
Benzoon, Gaertn. l. c. 206 = T. angustifolia.
brasiliensis, Raddi, in Mem. Mod. xviii. Fis. (1820) 405 = T. januarensis.
guyanensis, Gaertn. f. Fruct iii. 207 = T. mauritiana
guyanensis, Gaertn. f. l. c. t. 217 = T. Tanibouca.
mauritiana, Gaertn. f. l. c. 207 = T. mauritiana.

CATAPUNTIA, Muell. Arg. in DC. Prod. xv. ii. 1020 (1866), sphalm. = seq.

CATAPUTIA, Ludw. Gen. 81 (1760) = **Ricinus**, Tourn. (Euphorbiac.).
major, Ludw. Ectyp. t. 7 = R. communis.
minor, Ludw. Gen. 81 = R. communis.

CATARIA, Adans. Fam. ii. 192 (1763) = **Nepeta**, Linn (Labiat.).
canescens, Moench, Meth. 388 = N. orientalis.
Nepetella, Moench, l. c. = N. Nepetella.
nuda, Moench, l. c. = N. nuda.
paniculata, Moench, Meth. Suppl. 134 = N. nuda.
tomentosa, Gilib. Fl. Lituan. i. 78 = N. Cataria.
violacea, Moench, Meth. 388 = N. nuda.
vulgaris, Moench, l. c. 387 = N. Cataria.

CATAS, Domb. ex Lam. Encyc. ii. 354 (1786) = **Embothrium**, Forst. (Proteac.).
grandiflora, Juss. ex Lam. Encyc. ii. 354 = E. grandiflorum.
parviflora, Juss. l. c. 355 = E. coccineum.

CATASETUM, Rich. ex Kunth, Syn. Pl. Aequin. i. 330 (1822). *ORCHIDEÆ*, Benth. & Hook f. iii. 551.
CATACHAETUM, Hoffmgg. Verz. Orch. 22 (1842); Cf. Linnæa, xvi. (1842) Litt. 321.
CLOWESIA, Lindl. Bot. Reg. xxix. (1843) Misc. 25. t. 39.
CUCULINA, Rafin. Fl. Tellur. iv. 49 (1836).
MONACHANTHUS, Lindl. Bot. Reg. sub t. 1538 (1832).
MYANTHUS, Lindl. Bot. Reg. sub t. 1538 (1832).
abruptum, *Hook. Bot. Mag.* t. 3929.—Bras.
adnatum, Steud. Nom. ed. II. i. 311, sphalm = stratum.
albovirens, *Rudrig. Orch. Nov.* i. 129.—Bras.
stratum, *Lindl. Bot. Reg.* (1838) *Misc.* 61.—Bras. Barsquinianum, *Lem. Illustr. Hortic.* ix. (1862) *Misc.* 81.—Bras.
barbatum, *Lindl. Bot. Reg.* (1844) *Misc.* 88.—Bras.
bicolor, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 337.—N. Granat.
calceolatum, *Lem. Jard. Fleur.* i. (1851) *Misc.* 45.—Guatemala.
callosum, *Lindl. Bot. Reg.* (1840) *Misc.* 77.—Venezuela.
carunculatum, *Reichb. f. & Warsc. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 274.—Peruv.
cassideum, *Linden & Reichb. f. Xen. Orch.* ii. 171.—Bras.
cernuum, *Reichb. f. in Walp. Ann.* vi. 870.—Bras. Christyanum, *Reichb. f. in Gard. Chron.* (1882) i. 588. —Hab. ?
ciliatum, *Rodrig. Orch. Nov.* i. 130.—Bras. *Cleveringi*, Lindl. ex Van Geel, Sert. Bot. ii. (1839) t. 3 = macrocarpum.
cornutum, *Lindl. Bot. Reg.* (1840) *Misc.* 77.—Guiana.

CATASETUM:—
craniomorphum, *Hoffmgg. ex Heynh. Nom.* ii. 126, nomen.—Hab.?
cristatum, *Lindl. Bot. Reg.* t. 966.—Bras.
deltoideum, *Lindl. l. c.* xxvii. (1840) *Misc.* 157.—Guiana.
dilectum, *Reichb. f. Beitr. Orch. Centr. Am.* 78.—Costa Rica.
discolor, *Lindl. Bot. Reg.* (1844) *Misc.* 84.—Bras.
expansum, *Reichb. f. Otia Bot. Hamb.* 9.—Ecuador.
fimbriatum, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 124.—Hab.?
fimbriatum, Reichb. f. in Saund. Refug. Bot. t. 88 = discolor.
floribundum, Hook. Exot. Fl. ii. 151 = macrocarpum.
fuligineum, *Lindl. Bot. Reg.* (1841) *Misc.* 78.—Hab.?
glaucoglossum, *Reichb f. in Gard. Chron.* (1885) ii. 552.—Mexic.
globiflorum, *Hook. Bot. Mag.* t. 3942.—Bras.
Gnomus, *Linden & Reichb. f. Xen. Orch.* ii. 171.—Bras
heteranthum, Rodrig. Orch. Nov. i. 127 = C. Gnomus.
Hookeri, *Lindl. Bot. Reg.* sub t. 840.—Bras.
inapertum, Steud. Nom. ed. II. i. 811, sphalm. = purum.
incurvum, *Klotzsch, in Otto & Dietr. Allg Gartens.* xxii. (1854) 178.—Peruv.
integerrimum, Hook. Bot. Mag. t. 3823 = maculatum.
labiatum, *Rodrig. Orch. Nov.* ii. 218.—Bras.
laminatum, *Lindl. in Ann. Nat. Hist.* iv. (1840) 384.—Mexic.
lanciferum, *Lindl. Bot. Reg.* xxvii. (1841) 5. f. 5.—Bras.
Landsbergii, *Lindl. & Paxt. Flow. Gard* i. (1850–51) 156.—Venezuela.
luridum, *Hoffmgg. Verz. Orch.* 22; *ex Linnaea*, xvi. (1842) *Litt.* 231.—Bras.
longifolium, *Lindl. Bot. Reg.* (1839) *Misc.* 94.—Guiana.
luridum, *Lindl. Gen. et Sp. Orch.* 156.—Bras.
macrocarpum, *Rich. ex Kunth, Syn. Pl. Aequin.* i. 350.—Guiana.
macroglossum, *Reichb. f. in Gard. Chron.* (1877) ii 552.—Ecuador.
maculatum, *Kunth, Syn. Pl. Aequin.* i. 331.—N. Granat.
medium, *Reichb. f. in Gard. Chron.* (1885) ii. 6.—Hab.?
mentosum, *Lem. Gard. Fleur.* iii. (1852–53) *Misc.* 65.—Bras.
micranthum, *Rodrig. Orch. Nov.* ii. 219.—Bras.
Milleri, *Lodd. ex Lindl. Bot. Reg.* (1838) *Misc.* 80.—Bras.
Naso, *Lindl. Bot. Reg.* (1843) *Misc.* 71.—Venezuela.
ochraceum, *Lindl. l. c.* (1844) *Misc.* 44.—N. Granat.
Oerstedii, *Reichb. f. in Bonplandia*, iii. (1855) 218.—Nicaragua.
ollare, *Linden, Illustr. Hortic.* xxviii. (1881) *Misc.* 80.—Bras.
pallidum, *Klotzsch, in Otto & Dietr. Allg. Gartens.* xviii (1855) 217.—Bras.
Phasma, *Reichb. f. in Gard. Chron.* (1877) ii. 488.—Bras.
pileatum, *Reichb. f. l. c.* (1882) i. 492.—Venezuela.
planiceps, *Lindl. Bot. Reg.* (1843) t. 9.—Guiana.
portiferum, *Lindl. l. c.* (1838) *Misc.* 89.—Guiana.
proboscideum, *Lindl. l. c.* (1839) *Misc.* 86.—Bras.
purpurascens, *Hoffmgg. ex Heynh. Nom.* ii. 126, nomen.—Bras.
purum, *Nees & Sinning, Pl. Hort. Bonn. Nutrit.* i. 1.—Bras.
recurvatum, *Link, Klotzsch & Otto, Ic. Pl.* ii. 105, t. 49.—Guiana.
Regnellii, *Rodrig. Orch. Nov.* ii. 219.—Bras.
rivalatum, *Rodrig. l. c.* i. 130.—Bras.
roseo-album, Lindl. Bot. Reg. (1840) Misc. 65 = discolor.
roseum, *Reichb. f. in Gard. Chron.* (1879) 1003.—Bras.
roseum, *Rodrig. Orch. Nov.* i. 128.—Bras.
Russellianum, *Hook. Bot. Mag.* t. 3777.—Guatemala.
saccatum, *Lindl. Bot. Reg.* (1840) *Misc.* 76.—Peruv.
sanguineum, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 188.—Am. centr.
Scurra, *Reichb. f. in Gard. Chron.* (1878) 1008.—Guiana.
semiglyphum, Hook. Exot. Fl. iii. 213 = purum.
serratum, Lindl. Bot. Reg. (1847) t. 24 = viridi-flavum.
spinosum, Lindl. l. c. (1840) Misc. 65 = barbatum?

CATASETUM:—
squalidum, *Hoffmgg. Verz. Orch.* 23; *ex Linnaea*, xvi. (1842) *Litt.* 231.—Bras.
tabulare, *Lindl. Bot. Reg.* xxx. (1840) *Misc.* 40.—N. Granat.
tigrinum, *Reichb. f. in Gard. Chron.* (1881) i. 40.—Hab.?
thylaciochilum, *Lem. Illustr. Hortic.* iii. (1856) *Misc.* 90.—Mexic.
tricolor, *Reichb. f. in Hamb. Gartenz.* xiii. (1857) 313.—Guatemal.
tridentatum, Hook. Exot. Fl. ii. 90, 91 = macrocarpum.
trifidum, *Hook. Bot. Mag.* t. 2262.—Ind. occ.
trimerochilum, Lem. Illustr. Hortic. ix. (1862) t. 374 = Mormodes lineatum.
triodon, *Reichb. f. in Otto, Hamb. Gartenz* (1857) 318.—Guatemal.
triste, Reichb. f. in Bonplandia, iii. (1855) 218 = Hookeri.
Trulla, *Lindl. Bot. Reg.* (1840) *Misc.* 75.—Am. trop.
turbinatum, *Hoffmgg. ex Heynh. Nom.* ii. 126, nomen. —Bras.
umbrosum, *Rodrig. Orch. Nov.* i. 129.—Bras.
variabile, *Rodrig. l. c.* ii. 217.—Bras.
violascens, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 97.—N. Granat.
viride, *Lindl. Bot. Reg.* (1841) *Misc.* 79.—Bras.
viridi-flavum, *Hook. Bot. Mag.* t. 4017.—Am. centr.
Wailesii, Hook. l. c. t. 3987 = maculatum.
Warszewitzii, *Lindl & Paxt. Flow. Gard.* i. (1850–51) 46. f. 29.—Panama.

CATATEROPHORA, Steud. Nom. ed. II. i. 311 (1840), sphalm. = seq.

CATATHEROPHORA, Steud. in Flora, xii. (1829) 405 = **Pleuniastrum**, Pers. (Gramin.).
hordeiformis, Steud. l. c. = P. macrourum.

CATENARIA, Benth. in Miq. Fl. Jungh. 217, 220 (1852) = **Desmodium**, Desv. (Legumin.).
laburnifolia, Benth. l. c. = D. laburnifolium.

CAEPHA, Leschen. ex Reichb. Handb. 221 (1837) = **Platysace**, Bunge (Umbell.).

CATESBAEA, Linn. Gen. ed. I. 336 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 78.
campanulata, *Sagra, ex DC. Prod.* iv. 401.—Cuba.
elliptica, Spreng. ex. DC. l. c. = Exostemma Vavassorii.
erecta, *Moç. & Sesse, ex DC. l. c.*—Mexic
Grisebachii, *Griseb. in Mem. Acad.* N. S. viii. (1863) 503.—Cuba.
holacantha, *Griseb. Cat. Pl. Cub.* 122.—Cuba.
inermis, Spreng. Syst. i. 418 = Rondeletia buxifolia.
latifolia, *Lindl. Bot. Reg.* t. 858.—Ind. occ.
Lindeniana, *Baxt. ex Loud. Hort. Brit. Suppl.* iii. 511, nomen.—Hab.?
longiflora, Sw. Prod. Veg. Ind. Occ. 80 = spinosa.
longispina, *A. Rich. Fl. Cub. Fanerog.* ii. 12.—Cuba.
macrantha, A. Rich. l. c. = spinosa?
macrantha, *Wright, in Sauv. Fl. Cub.* 59.—Cuba.
parahybensis, *Vell. Fl. Flum.* 64; i. t. 145.—Bras.
parviflora, Griseb. Cat. Pl. Cub. 122 = Scolosanthus parviflorus.
parviflora, Lam. Tabl. i. 288. n. 1481 = Scolosanthus versicolor.
parviflora, Spreng. ex Steud. Nom. ed. II. i. 311 = Rondeletia buxifolia.
parviflora, *Sw. Prod. Veg. Ind. Occ.* 80.—Jamaic.
parvifolia, *DC. Prod.* iv. 401.—Ins. S. Doming.
phlebocalyx, Griseb. Cat. Pl. Cub. 122 = Phyllacantha Grisebachiana.
spinosa, *Linn. Sp. Pl.* 109.—Ind. occ.
strigosa, Willd. ex Steud. Nom. ed. II. i. 311 = Randia tetrandra.
triacantha, Spreng. Neue Entdeck. iii. 47 = Scolosanthus triacanthus.
Vavassorii, Spreng. Syst. i. 416 = Exostemma Vavassorii.

CATEVALA, Medic. Theodora, 67 (1786) = **Aloë**, Linn. (Liliac.).
arachnoidea, Medic. l. c. 68 = Haworthia arachnoides.
arborescens, Medic. l. c. 67 = A. arborescens.
atroviridis, Medic. l. c. 69 = Haworthia atroviresa.
humilis, Medic. l. c. = A. humilis.
retusa, Medic. l. c. 68 = Haworthia retusa.

CATHA, Forsk. Fl. Aegypt. Arab. 63 (1775). *CELASTRINEAE*, Benth. & Hook. f. i. 361
METHYSCOPHYLLUM, Eckl. & Zeyh. Enum. 152 (1836).
TRIGONOTHECA, Hochst. in Flora, xxiv. (1841) 662
edulis, *Forsk. l. c.*—Afr. trop.
Forskalei, A. Rich. Tent. Fl. Abyss. i. 134 = edulis.
inermis, J. F. Gmel. Syst. 411 = edulis.
spinosa, Forsk. Fl. Aegypt. Arab. 64 = Celastrus parviflorus.

CATHA, G. Don, Gen. Syst. ii. 11 (1832) = **Celastrus**, Linn.
acuminata, Presl, Bot. Bemerk. 33 = Cel. acuminatus.
alaternifolia, Tul. in Ann. Sc. Nat. Ser. IV. viii. (1857) 98.—Ins. Comor.
articulata, G. Don, Gen. Syst. ii. 10 = Cel. articulatus.
Benthamii, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 310 = Cel. Championi.
buxifolia, G. Don, Gen. Syst. ii. 10 = Cel. buxifolius.
campestris, Presl, Bot. Bemerk. 34 = Cel. pyracanthus.
cassinoides, Webb & Berth. Phyt. Canar. ii. 142 = Cel. cassinoides.
crenata, A. Gray, Bot. U. St. Expl. Exped. i. 288 in texta = Cel. crenatus.
Cunninghamii, Hook. in Mitch. Journ. Trop. Austral. 387 = Cel. Cunninghamii.
cymea, G. Don, Gen. Syst. ii. 10 = Cel. buxifolius.
decolor, Webb, Fragm. Fl. Aethiop. 60 = Gymnosporia montana.
Dreyeri, Lowe, Man. Fl. Madeir. i. 107.—Ins. Madera.
emarginata, G. Don, Gen. Syst. ii. 9 = Gymnosporia emarginata.
europaea, Boiss. Voy. Espagne, ii. 725 = Gymnosporia montana.
excisa, G. Don, Gen. Syst. ii. 9 = Cel. excisa.
fasciculata, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 98.—Malaya.
flexuosa, G. Don, Gen. Syst. ii. 10 = Cel. flexuosus.
fruticosa, Thw. Enum. Pl. Zeyl. 72 = Gymnosporia emarginata.
Grossularae, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 98.—Madag.
heterophylla, Presl, Bot. Bemerk. 33 = Cel. heterophyllus.
Heyneana, Walp. Rep. i. 532 = Gymnosporia Heyneana.
integra, G. Don, Gen. Syst. ii. 9 = Cel. pyracanthus.
integrifolia, G. Don, l. c. = Cel. integrifolius.
lanceolata, Presl, Bot. Bemerk. 33 = Cel. linearis?
leptopus, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 100.—Madag.
linearis, G. Don, Gen. Syst. ii. 9 = Cel. linearis.
monosperma, Benth. in Hook. Lond. Journ. Bot. i. 483 = Cel. monospermus.
montana, G. Don, Gen. Syst. ii. 10 = Gymnosporia montana.
multiflora, G. Don, l. c. = Cel. multiflorus.
ovata, Walp. Rep. i. 532 = Gymnosporia ovata.
panciulata, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 275 = Cel. panciulatus.
parviflora, G. Don, Gen. Syst. ii. 10 = Cel. parviflorus.
parvifolius, Presl, Bot. Bemerk. 34 = Cel. parvifolius.
patens, Presl, l. c. 33 = Cel. patens.
Pyracantha, G. Don, Gen. Syst. ii. 10 = Cel. pyracanthus.
rigida, G. Don, l. c. 9 = Cel. rigidus.
Rothiana, Walp. Rep. i. 532 = Gymnosporia Rothiana.
rotundifolia, G. Don, Gen. Syst. ii. 10 = Dovyalis celastroides.
rufa, G. Don, l. c. = Gymnosporia rufa.
rupestris, Presl, Bot. Bemerk. 33 = Cel. acuminatus.
senegalensis, G. Don, Gen. Syst. ii. 10 = Gymnosporia montana.
senegalensis, Webb, Fragm. Fl. Aethiop. 61 = Cel. senegalensis.
spathulata, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) 4. 278.—Ind. or.
spathyphylla, Presl, Bot. Bemerk. 33 = Cel. buxifolius.
trigyna, Presl, Bot. Bemerk. 33 = Gymnosporia trigyna.
venenata, Presl, l. c. 34 = Cel. buxifolius.
vitiensis, A. Gray, Bot. U. St. Expl. Exped. i. 287. t. 28.—Ins. Fiji.
Wallichii, G. Don, Gen. Syst. ii. 10 = Gymnosporia Wallichiana.
zeylanica, G. Don, l. c. = Scutia Commersonii.

CATHANTHES, Rich. in Mém. Mus. Par. i. (1815) 365 = **Tetronotum**, Willd. (Naiad.).

CATHARANTHUS, G. Don, Gen. Syst. iv. 95 (1838) = **Vinca**, Linn. (Apocynac.).
pusillus, G. Don, l. c. = V. parviflora.
roseus, G. Don, l. c. = V. roseo.

CATHARTOCARPUS, Pers. Syn. i. 459 (1805) = **Cassia**, Linn. (Legumin.).
Attelah, G. Don, Gen. Syst. iii. 453 = Cassia Arereh.
bacillaris, Pers. Syn. i. 459 = Cassia bacillaris.
Bacillus, Lindl. Bot. Reg. t. 881 = Cassia bacillaris.
brasiliana, Jacq. Fragm. 59. t. 85. f. 8 = Cassia grandis.
Brewsteri, F. Muell. Fragm. i. 110 = Cassia Brewsteri.
conspicua, G. Don, in Loud. Hort. Brit. 167 = Cassia Sieberiana.
excelsa, G. Don. Gen. Syst. iii. 453 = Cassia Fistula.
ferruginea, G. Don, l. c. = Cassia ferruginea.
Fistula, Pers. Syn. i. 459 = Cassia Fistula.
fistuloides, G. Don, in Loud. Hort. Brit. 167 = Cassia Fistula.
grandis, Pers. Syn. i. 459 = Cassia grandis.
Humboldtianus, Loud. Hort. Brit. 167 = Cassia spectabilis.
javanicus, Pers. Syn. i. 459 = Cassia javanica.
lactea, G. Don, Gen. Syst. iii. 454 = Cassia lacta.
marginatus, G. Don, l. c. 458 = Cassia marginata.
moschatus, G. Don, l. c. = Cassia moschata.
nodosus, Scud. Nom. ed. II. i. 311 = Cassia nodosa.
rhomboidea, G. Don, Gen. Syst. iii. 453 = Cassia Fistula.
Roxburghii, Lond. Hort. Brit. 167 = Cassia marginata.
Sieberianus, G. Don, Gen. Syst. iii. 453 = Cassia Sieberiana.
speciosus, G. Don, l. c. = Cassia spectabilis.
Trinitatis, G. Don, in Loud. Hort. Brit. 167 = Cassia spectabilis.

CATHARTOLINUM, Reichb. Handb. 306 (1837) = **Linum**, Linn.
agrestis, Reichb. l. c. 307 = L. angustifolium.
alternum, Reichb. l. c. = L. strictum.
bicolor, Reichb. l. c. = L. bicolor.
corymbuliosum, Reichb. l. c. = L. strictum.
gallicum, Reichb. l. c. = L. gallicum.
prostrat, Reichb. l. c. = L. catharticum.
salsoloides, Reichb. l. c. = L. salsoloides.
sardoum, Reichb. l. c. = L. Muelleri.
spicatum, Reichb. Ic. Fl. Germ. xvi. 62 = L. strictum.
strictum, Reichb. l. c. xvi. t. 257 = L. strictum.
suffruticosum, Reichb. Handb. 307 = L. suffruticosum.
tenuifolium, Reichb. l. c. = L. tenuifolium.
virginianum, Reichb. l. c. = L. virginianum.

CATHAUSTRUM, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 448. *CELASTRINEAE*, Benth. & Hook. f. i. 392.
capense, *Turcz. l. c.*—Afr. austr.

CATHCARTIA, Hook. f. in Bot. Mag. t. 4696 (1851). *PAPAVERACEAE*, Benth. & Hook. f. i. 59.
integrifolia, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 510.—China.
villosa, *Hook. f. in Bot. Mag.* t. 4696.—Reg. Himal.

CATHEA, Salisb. in Trans. Hort. Soc. i. (1812) 300 = Calopogon, R. Br. (Orchid.).
pulchella, Salisb. l. c. = Calop. pulchellus.

CATHEDRA, Miers, in Ann. Nat. Hist. Ser. II. vii. (1851) 452, 457. *OLACINEAE*, Benth. & Hook. f. i. 346.
DIPLOCRATER, Benth. in Hook. Kew Journ. iii. (1851) 267.
acuminata, *Miers, l. c.* Ser. III. iv. (1850) 361.—Bras.
crassifolia, *Miers, l. c.*—Bras.
Gardneriana, *Miers, l. c.* Ser. II. vii. (1851) 459.—Bras.
rubricaulis, *Miers, l. c.* 458.—Bras.

CATHEDRICUM, J. Presl, in Rel. Haenk. i. 294. t. 49 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1192.
erectum, *Vasey & Hack. in Bull. Torrey Bot. Club.* xi. (1884) 87. t. 48.—Mexic.
prostratum, *J. Presl, in Rel. Haenk.* i. 295. t. 49.—Mexic.

CATHETOSTEMMA, Blume, Rumphia, iv. 80 (1848) = **Hoya**, R. Br. (Asclepiad.).
laurifolium, Blume, l. c. = H. laurifolia.

CATHETUS, Lour. Fl. Cochinch. 607 (1790) = **Phyllanthus**, Linn. (Euphorbiac.).
fasciculata, Lour. l. c. 608 = P. cochinchensis.

CATHISSA, Salisb. Gen. Pl. Fragm. 34 (1866) = **Ornithogalum**, Tourn. (Liliac.).
concinna, Salisb. l. c. = O. unifolium.
unifolia, Salisb. l. c. = O. unifolium.

CATHORMION, Hassk. Retzia, i. 281 (1855) = **Pithecolobium**, Mart. (Legumin.).
Jungkuhnianum, Hassk. l. c. 282 = P. Junghuhnianum.
moniliferum, Hassk. l. c. 281 = P. moniliferum.

CATIMBIUM, Juss. Gen. 62 (1789) = **Alpinia**, Linn. (Scitamin.).
erectum, Juss. ex Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 342 = A. calcarata.
exaltatum, Mirb. ex Steud. Nom. ed. II. i. 311 = Renealmia exaltata.
nutans, Juss. ex Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 342 = A. nutans.

CATINGA, Aubl. Pl. Gui. i. 511. t. 208 (1775) = **Eugenia**, Linn. (Myrtac.).
aromatica, Aubl. l. c. f. 2 = E. aromatica.
moschata, Aubl. l. c. f. 1 = E. Catinga.

CATOBLASTUS, H. Wendl. in Bonplandia, viii. (1860) 104, 106. *PALMAE*, Benth. & Hook. f. iii. 901.
maynensis, *Drude, in Mart. Fl. Bras.* iii. ii. 544.—Peruv.
praemorsus, *H. Wendl. in Bonplandia*, viii. (1860) 104.—N. Granat.
pubescens, *H. Wendl. l. c.*—Bras.

CATOCOMA, Benth. in Hook. Journ. Bot. iv. (1842) 101 = **Brodemeyera**, Willd. (Polygal.).
altissima, Poepp. & Endl. Nov. Gen. et Sp. iii. 65. t. 273 = B. altissima.
brevifolia, Benth. in Hook. Journ. Bot. iv. (1842) 102 = B. brevifolia.
emarginata, Spruce, ex A. W. Benn. in Mart. Fl. Bras. xiii. iii. 50 = B. altissima.
floribunda, Benth. in Hook. Journ. Bot. iv. (1842) 102 = B. floribunda.
Kunthiana, Benth. l. c. 103 = B. Kunthiana.
laurifolia, Benth. l. c. = B. laurifolia.
lucida, Benth. l. c. 101 = B. lucida.
Manzonni, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. iii. 51 = B. lucida.
mollis, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) v. 133 = B. altissima.
myrtifolia, Spruce, ex A. W. Benn. in Mart. Fl. Bras. xiii. iii. 51 = B. myrtifolia.
parvifora, Spruce, ex A. W. Benn. l. c. 51 = B. parviflora.

CATOCOEYNE, Hook. f. in Benth. & Hook. f. Gen. i. 768 (1867). *MELASTOMACEAE*, Benth. & Hook. f. i. c.
linnoides, *Hook. f. l. c.*—Peruv.

CATODIACRUM, Dulac, Fl. Hautes-Pyr. 369 (1867) = **Orobanche**, Tourn.
cruentum, Dulac, l. c. 371 = O. cruenta.
fortidum, Dulac, l. c. = O. foetida.
incurvum, Dulac, l. c. 370 = O. caryophyllacea.
laterum, Dulac, l. c. = O. Hederae.
pallidiflorum, Dulac, l. c. = O. pallidiflora.
rubidum, Dulac, l. c. = O. Teucrii.
spectabilorum, Dulac, l. c. = O. Epithymum.
variegatum, Dulac, l. c. = O. variegata.

CATONIA, P. Br. Hist. Jamaic. 143 (1756); Juss. Gen. 441 (1789). *INC. SED.*
Browne), *Roem. & Schult. Syst.* iii. 450.—Jamaica.
marginata, Jamne St. Hill. Expos. ii. 547 = Browne).

CATONIA, Moench, Meth. 535 (1794) = **Crepis**, Linn. (Compos.).
blattarioides, Cass. in Dict. Sc. Nat. v. 374; xxvi. 9 = Crepis blattarioides.
cordifolia, Moench, Meth. 536 = Hieracium amplexicaule.
crocea, Reichb. ex Steud. Nom. ed. II. i. 311 = Crepis Columnae.
sagittata, Moench, Meth. 536 = Crepis blattarioides.

CATONIA, Rafin. Fl. Tellur. ii. 86 (1836) (Quid?).
lantanoides, Rafin. l. c.—Hab.?

CATONIA, Vahl, in Skrivt. Nat. Hist. Selsk. Klobenh vi. (1810) 98 = **Erycibe**, Roxb. (Convolvulac.).
elliptica, Vahl, l. c. 101 = E. paniculata.
glauca, Vahl, l. c. 98 = E. paniculata.

CATONIA, Veil. Fl. Flum. 155 (1825); t. iv. t. 13 (‘Oct. Mon.’ Quid?).
rustica, Vell. l. c.—Bras.

CATOPHERIA, Benth. in Benth. & Hook. f. Gen i. 1173 (1876). *LABIATAE*, Benth. & Hook. f. i. c.
capitata, *Benth. l. c.*—Mexic.
chiapensis, *A. Gray, ex Benth. in Hook. Ic. Pl.* sub t. 1215.—Mexic.
spicata, *Benth. l. c.* t. 1215.—N. Granat.

CATOPHRACTES, D. Don, in Proc. Linn. Soc. i. 6 (1839). Trans. Linn. Soc. xvii. (1841) 306. t. 22. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1048.
Alexandri, *D. Don, ll. cc.* 4; 307.—Afr. austr.
Kolbeana, *Harv. Gen. S. Afr. Pl.* ed. II. 276.—Afr austr.
Welwitschii, *Seem. in Journ. Bot.* iii. (1865) 331.—Afr. trop.

CATOPHYLLUM, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 222 (1876) = **Mikania**, Willd. (Compos.).
deltoideum, Pohl, ex Baker, l. c. = M. officinalis.
tropaeolifolium, Pohl, ex Baker, l. c. = M. officinalis.

CATOPODIUM, Link, Hort. Berol. i. 44 (1827), sphalm = **Catapodium**, Link (Gramin.).

CATOPSIS, Griseb. in Goett. Nachr. 1864 (1865) 21 *BROMELIACEAE*, Benth. & Hook. f. iii. 670.
PODOSPERMUM, Brongn. in Ann. Sc. Nat. Sér. V. i. (1864) 227.
TUSSACIA, Klotzsch, ex Beer, Bromel. 99 (1856).
fulgens, *Griseb. l. c.* 21.—Am. trop.
nitida, *Griseb. Fl. Brit. W. Ind.* 599.—Ind. occ.
nutans, *Griseb. l. c.*—Ind. occ.

CATOSPERMA, Benth. in Hook. Ic. Pl. t. 1028 (1868). *GOODENOVIEAE*, Benth. & Hook. f. ii. 539.
Carvyi, *F. Muell. in Melb. Chemist* (Apr. 1884).—Austral.
Muelleri, *Benth. in Hook. Ic. Pl.* t. 1028.—Austral.

CATOSTEMMA, Benth. in Hook. Lond. Journ. Bot ii. (1843) 365. *MALVACEAE*, Benth. & Hook. f. i. 794.
fragrans, *Benth. l. c.*—Guiana.

CATTLEYA, Lindl. Collect. Bot. tt. 32, 37 (1824). *ORCHIDEAE*, Benth. & Hook. f. iii. 531.
MALKENIA, Dum. in Mém. Acad. Brux. ix. (1836) 18. t. 10.
Achlandiae, Planch. in Fl. des Serres, Sér. I. vii. (1851-52) 88 = seq.
Aclandiae, *Lindl. Bot. Reg.* xxvi. (1840) t. 48.—Bras.
acuminata, Beer, Prakt. Orch. 208 = Laelia acuminata.
albida, Beer, l. c. = Laelia albida.
alutacea, *Rodrig. Orch. Nov.* ii. 157.—Bras.
amethystina, Morr. in Ann. Gand. iv. (1848) 217. t. 201 = intermedia.
amethystoglossa, Linden & Reichb. f. ex Warner, Sel. Orch. Ser. I. t. 5 = guttata.
anceps, Beer, Prakt. Orch. 208 = Laelia anceps.
Arembergii, Scheidw. in Otto & Dietr. Allg. Gartenz. xi. (1843) 109 = Loddigesii.
Aschkenatica, Hoynh. Nom. ii. 136, sphalm. = Aclandiae.

CATTLEYA:—

Auclandi, Beer, Prakt. Orch. 208, sphalm. = Aclandiae.

aurantiaca, P. N. Don, in Florist's Journ. (1840) 185 = Epidendrum aurantiacum.

aurea, *Linden*, *Illustr. Hortic.* xxx. (1883) t. 493.—N. Granat.

autumnalis, Beer, Prakt. Orch. 208 = Laelia autumnalis.

bicolor, *Lindl. Bot. Reg.* (1836) sub t. 1919.—Am. austr.

Bluntei, *Hort. ex Gard. Chron.* (1876) i. 735.—Hab.

bogotensis, Linden, in Belg. Hortic. xv. (1865) 102 = labiata.

Bowringiana, *Veitch, ex Gard. Chron.* (1885) ii. 688. —Honduras.

Brabantiae x, Veitch.—Cf. Gard. Chron. (1864) 413; et Floral Mag. vi. (1867) t. 860.

Bryceyrana x, Reichb. f. in Gard. Chron.(1888) ii. 492.

Brysana, Lem. in Jard. Fleur. iii. (1852–53) t. 275 = Laelia purpurata.

bulbosa, Lindl. Bot. Reg. (1847) t. 42 = Walkeriana.

calammata x, André, in Rev. Hortic. (1888) 564.

candida, Williams, in Gard. Chron. (1851) 453 = Loddigesii.

Carrierii, Houllet, in Rev. Hortic. (1876) 250 = labiata.

cernua, Beer, Prakt. Orch. 209 = Sophronitis cernua.

Chamberlainiana x, Reichb. f. in Gard. Chron. (1881) ii. 427.

chlorantha, Beer, Prakt. Orch. 209 = Sobralia macrophylla.

chocoiana, Linden, in Illustr. Hortic. xvii. (1870) 87 = labiata.

cinnabarina, Beer, Prakt. Orch. 209 = Laelia cinnabarina.

citrina, *Lindl. Coll. Bot.* sub t. 37.—Mexic.

coccinea, Lindl. Bot. Reg. sub t. 1919 = Sophronitis grandiflora.

crispa, Beer, Prakt. Orch. 209 = Schomburgkia crispa.

crispa, Lindl. Bot. Reg. t. 1172 = Laelia crispa.

Dawsoni, Warner, Sel. Orch. Pl. i. t. 16 = labiata.

Deckeri, Klotzsch, in Otto & Dietr. Allg. Gartens. xxiii. (1856) 81 = Skinneri.

decora, Beer, Prakt. Orch. 209 = Sobralia decora.

dessoiniensis x, Hort. ex Floral Mag. N. S. (1881) sub t. 437 = Laelia elegans ?

dolosa, *Reichb. f. Xenia Orchid.* ii. 224, *in synon.*—Bras.

domingensis, Lindl. Gen. et Sp. Orch. 118 = Broughtonia lilacina.

Dominiana x, Lindl. in Gard. Chron. (1859) 948.

Dormaniana x, Reichb. f. in Gard. Chron. (1882) i. 816.

Dowiana, *Batem. l. c.* (1866) 922.—Costa Rica.

elatior, Lindl. Gen. et Sp. Orch. 117 = guttata.

Eldorado, Linden, in Fl. des Serres, Sér. II. vii. (1869–70) 13. t. 18.6 = labiata.

elegans, Morr. in Ann. Gand. iv. (1848) 93. t. 185 = Laelia elegans.

elegantissima, *Linden, Illustr. Hortic.* xxviii. (1881) 80.—Venezuela.

elongata, *Rodrig. Orch. Nov.* i. 72.—Bras.

eximia, *Rodrig. l. c.* 70.—Bras.

exoniensis x, Reichb. f. in Gard. Chron. (1867) 1144.

fausta x, Reichb. f. l. c. (1873) 289.

felix x, Reichb. f. l. c. (1876) ii. 68.

flava, Beer, Prakt. Orch. 210 = Laelia flava.

flavida, Klotzsch, in Otto & Dietr. Allg. Gartens. xxiv. (1856) 73 = luteola.

Forbesii, *Lindl. Coll. Bot.* sub t. 37.—Bras.

fragrans, *Rodrig. Orch. Nov.* i. 72.—Bras.

furfuracea, Beer, Prakt. Orch. 210 = Laelia furfuracea.

fulva, *Beer, l. c.* 210.—Hab.?

Galeottiana, Beer, l. c. = Epidendrum Galeottianum.

Gardneriana, Reichb. f. in Gard. Chron. (1870) 1373, in nota = Walkeriana.

Gaskelliana, Hort. Sander, l. c. (1888) t. 948, 950 = labiata.

gigantea, *Hort. ex P. N. Don, in Florist's Journ.* (1840) 187.—Hab.?

Gigas, Linden & André, Illustr. Hortic. xx. (1873) 70 = labiata.

gloriosa, Carr. in Rev. Hortic. (1885) 338 = labiata.

Grahami, Lindl. Gen. et Sp. Orch. 116 = Laelia majalis.

grandiflora, Beer, Prakt. Orch. 209 = Sophronitis grandiflora.

granulosa, *Lindl. Bot. Reg.* (1842) t. 1.—Am. austr.

guatemalensis, *T. Moore, in Flor. Mag.* (1861) t. 61.—Guatemala.

CATTLEYA:—

guttata, *Lindl. Bot. Reg.* t. 1406.—Bras.

Hardyana, Hort. Sander.—Cf. Gard. Chron. (1888) i. 948 = labiata.

Harrisoniae, Paxt. Mag. Bot. iv. (1838) 247 = Loddigesii.

Harrisoniana, Batem. ex Lindl. in Bot. Reg. sub t. 1919 = Loddigesii.

Harrisonii, Hort. ex Duchartre, in Journ. Soc. Hort. Fr. iii. (1867) 725. t. 7 = praec.

hybrida x, Veitch, in Gard. Chron. (1868) 602.

ianthina, Hort. ex Gard. Chron. (1858) 247, nomen = intermedia ?

imperialis, J. O'Brien, in Gard. Chron. (1888) ii. 404 = labiata.

intermedia, *R. Grah. in Bot. Mag.* t. 2851.—Bras.

intricata, *Reichb. f. in Gard. Chron.* (1884) ii. 7.—Hab.?

iricolor, *Reichb. f. l. c.* (1874) ii. 162.—Hab.?

Isabella, *Reichb. f. in C. Koch & Fint. Wochenschr.* (1859) 286.—Hab.?

isopetala, *Beer, Prakt. Orch.* 211.—Hab.?

Karwinskii, Mart. Auuw. Merkw. Pfl. 14. t. 10 = citrina.

labiata, *Lindl. Coll. Bot.* t. 38.—Am. trop.

laelioides, Lem. Jard. Fleur. iii. (1852–53) Misc. 42, 50 = Skinneri.

Lawrenceana, *Reichb. f. in Gard. Chron.* (1885) i. 338.—Guiana.

Lawrenceana, Warsc. ex Reichb. f. in Gard. Chron (1874) i. 248 = Dowiana.

Leeana, Hort. Sander.—Cf. Gard. Chron. (1883) i. 43 = labiata.

Lemoniana, Lindl. Bot. Reg. (1846) t. 35 = labiata.

Leopoldii, Hort. Versch. ex Illustr. Hortic. i. (1854) Misc. 68; ii. (1855) t. 69 = guttata.

Liliastrum, Beer, Prakt. Orch. 212 = Sobralia Liliastrum.

Lindleyana, *Reichb. f. in C. Koch, Allg. Gartens* (1857) 118.—Bras.

lobata, Lindl. Gard. Chron. (1848) 403 = Laelia Boothiana.

Loddigesii, *Lindl. Coll. Bot.* t. 37.—Bras.

Lucasiana x, Reichb. f. in Gard. Chron. (1885) ii. 456.

Luddemanniana, Reichb. f. Xenia Orch. i. 29 = labiata.

Lueddemanni, *Lindl. in Gard. Chron.* (1853) 774.—Bras.

macrantha, Beer, Prakt. Orch. 212 = Sobralia macrantha

majalis, Beer, l. c. = Laelia majalis.

Mangleni x, Reichb. f. in Gard. Chron. (1880) ii. 556.

Marshalliae x, Seden, ex Reichb. f. l. c. (1878) i. 234.

marginata, Beer, Prakt. Orch. 212 = Schomburgkia marginata.

marginata, Paxt. Mag. Bot. x. (1843) 205 = Laelia pumila.

maritima, Lindl. Bot. Reg. (1836) sub t. 1919 = intermedia.

Mastersoniae x, Seden, ex Reichb. f. in Gard. Chron. (1878) ii. 556.

Mazangeana, Reichb. f. in Gard. Chron. (1888) i. 343 = labiata.

maxima, *Lindl. Gen et Sp. Orch.* 116.—Peruv.

Mendelii, *Hort.—Cf. Gard. Chron.* (1888) ii. 404.—Hab.?

Meyeri, Regel, in Ind. Sem. Hort. Petrop. (1855) 17. Cf. Linnaea, xxviii. (1856) 369 = luteola.

Mitchelli x, Reichb. f. in Gard. Chron. (1876) ii. 386

Morganiae, Warner, Orch. Album, i. (1862) t. 6 = labiata.

Mossiae, Parker, in Hook. Bot. Mag. t. 3669 = labiata.

Naberiana, Reichb. f. in Gard. Chron. (1885) i. 756 = labiata.

nobilior, Reichb. f. in Illustr. Hortic. xxx. (1883) t. 488 = Walkeriana.

odoratissima, P. N. Don, in Florist's Journ. (1840) 185 = Sobralia violacea ?

ovata, Lindl. Bot. Reg. sub t. 1919 = intermedia.

pallida, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 51 = labiata.

paludosa, Beer, Prakt. Orch. 215 = Sobralia paludosa.

papyracinaea, Morr. in Ann. Gand. i. (1846) 57. t. 5 = Loddigesii.

pedunculariis, Beer, Prakt. Orch. 213 = Laelia pedunculariis.

Pentersii, André, in Rev. Hortic. (1886) 271 = labiata.

Percivaliana, Hort.—Cf. Gard. Chron. (1883) ii. 404 = labiata.

CATTLEYA:—

Perrinii, Endl. in Harting. Parad. Vindob. t. 20 = labiata.

Perrinii, Lindl. Bot. Reg. (1838) t. 2 = Laelia Perrinii.

picturata x, Reichb. f. in Gard. Chron. (1877) ii. 584.

Pinelli, Hort. ex Lindl. in Fl. des Serres, Sér. I. vii (1851–52) 43 = Laelia pumila.

piperita, *Hort.—Cf. Gard. Chron.* (1850) 168, nomen —Hab.?

porphyroglossum, *Linden & Reichb. f. in Otto & Dietr. Allg. Gartens.* xxiv. (1856) 98.—Bras.

porphyrophlebia x, Reichb. f. in Gard. Chron. (1885) ii. 552.

Princeps, *Rodrig. Orch. Nov.* i. 68.—Bras.

pterocarpa, Beer, Prakt. Orch. 209 = Sophronitis pterocarpa.

pumila, Hook. Bot. Mag. t. 3656 = Laelia pumila.

purpurata, Beer, Prakt. Orch. 213 = Laelia purpurata.

purpurina, *Rodrig Orch. Nov.* ii. 158.—Bras.

quadricolor, Lindl. in Paxt. Flow. Gard. i. 6 = labiata.

Quindoia, *Hort.—Cf. Gard. Chron.* (1858) 247, nomen. —Hab.?

quinquecolor x, Dombr. in Floral Mag. t. 511.

reflexa, Parment. ex Walp. Ann. vi. 428 = Laelia crispa.

regalis, Hort.—Cf. Gard. Chron. (1883) ii. 404 = labiata ?

Regnellii, Warner, Sel. Orch. Ser. II. t. 22 = Schilleriana.

Reinecheana, Reichb. f. in Bonplandia, iv. (1856) 237 = labiata.

resplendens x, *Reichb. f. in Gard. Chron.* (1885) i. 692.

Roezlii, Reichb. f.—Cf. Gard. Chron. (1882) ii. 457 = labiata.

Rollissonii, T. Moore, in Floral Mag. i. (1861) t. 8 = labiata.

rubescens, Beer, Prakt. Orch. 214 = Laelia rubescens.

Ruckeri, *Linden, in Belg. Hortic. xv.* (1865) 102.—N. Granat.

Ruckerii, Beer, Prakt. Orch. 214 = Sobralia Ruckeri.

Sanderiana, Hort. Low.—Cf. Gard. Chron. (1882) ii. 151 = labiata.

Schilleriana, *Reichb. f. in C. Koch, Allg. Gartens.* (1857) 385.—Bras.

Schofieldiana, *Reichb. f. in Gard. Chron.* (1882) ii. 808.—Hab.?

Schroederiana, *Reichb. f. l. c.* (1883) ii. 102.—Hab.?

Scita x, Reichb. f. l. c. (1885) ii. 489.

sessilis, Beer, Prakt. Orch. 214 = Sobralia sessilis.

Skinneri, *Batem. Orch. Mex. & Guat.* t. 13.—Guatemala.

speciosissima, Hort.—Cf. Gard. Chron. (1868) 404, in nota = labiata.

spectabilis, Hort.—Cf. Florist (1850) 91 = marginata.

sphenophora, Morr. Ann. Gand. iv. (1848) 17. t. 175 = guttata.

stulfurina, Lem. Illustr. Hortic. i. (1854) Misc. 2 = citrina.

superba, Schomb. ex Lindl. Sert. Orch. t. 22 = violacea.

superbiens, Beer, Prakt. Orch. 214 = Laelia superbiens.

tibicinis, Beer, l. c. 215 = Schomburgkia tibicinis.

tichotoma, Beer, l. c. = Sobralia tichotoma.

tigrina, *A. Rich. ex Beer, l. c.* 215.—Hab.?

Trianaei, Linden & Reichb. f. in Bot. Zeit. xviii. (1860) 74 = labiata.

Trianaei, *Linden & Reichb. f. in C. Koch, Wochenschr.* iii. (1860) 67.—N. Granat.

trichopilochila, *Rodrig. Orch. Nov.* i. 70.—Bras.

trilabiata, *Rodrig. l. c.* 69.—Bras.

trichokalma x, Reichb. f. in Gard. Chron. (1888) ii. 596.

tuberculata, *Hort. ex Reichb. f. in Walp. Ann.* xi. 320. —Hab.?

undulata, Beer, Prakt. Orch. 215 = Schomburgkia undulata.

Veitchi, Hort. ex Pfitz, Verg. Morph. Orch. 124 = sec. ?

Veitchiana x, Cf. Gard. Chron. (1874) i. 385, 566 = Laelia Veitchiana.

velutina, *Reichb. f. in Gard. Chron.* (1870) 140, 1373. —Hab.?

venusta, Hoffmgg. Verz.—Cf. Bot. Zeit. i. (1843) 831 = Forbesii.

violacea, Beer, Prakt. Orch. 209 = Sophronitis violacea.

violacea, Beer, l. c. 215 = Sobralia violacea.

virginalis, Linden & André, Illustr. Hortic. xxiii. (1876) t. 257 = labiata.

CATTLEYA:—
Wagneri, Reichb. f. in Bonplandia (1854) 21 = labiata.
Walkeriana, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 662.—Bras.!
Waltini, Linden, in Gard. Chron. (1882) i. 557 = labiata.
Warneri, T. Moore, ex Warner, Orch. Sel. Ser. I. t. 8 = labiata.
Warscewiczii, Reichb. f. in Bonplandia (1854) 119 = labiata.
Whitei, *Hort. Low.*—*Cf. Gard. Chron.* ii. (1882) 151, 566.—Hab.?
Wilsoniana, *Reichb. f. in Gard. Chron.* (1877) ii. 72.—Am. austr.

CATTLEYOPSIS, Lem. Jard. Fleur. iv. (1854) Misc. 59. *ORCHIDEAE.*
delicatula, *Lem. l. c.*—Cuba.

CATU-ADAMBOE, Adans. Fam. ii. 85 (1763) = **Lagerstroemia**, Linn. (Lythrar.) (L. hirsuta).

CATU-NAREGAM, Adans. Fam. ii. 85 (1763) = **Randia**, Houst. (Rubiac.) (R. viross).

CATURUS, Linn. Mant. i. 19 (1767) = **Acalypha**, Linn. (Euphorbiac.).
ramiflorus, Linn. l. c. 127 = Boehmeria ramiflora.
sessilis, Thou. ex Baill. Adansonia, i. (1860–61) 267 = A. integrifolia.
spicifloras, Jacq. f. ex Wedd. in DC. Prod. xvi. 1. 214 = Boehmeria spicifloras.
spicifloras, Linn. Mant. i. 127 = A. hispida.
pedunculatus, Thou. ex Baill. in Adansonia, i. (1860–61) 267 = A. arborea.

CATURUS, Lour. Fl. Cochinch. 612 (1790) = **Malaisia**, Blanco (Urtic.).
scandens, Lour. l. c. = M. tortuosa.
acuminatus, *Cunninghamii*, *Deplanchei*, *fagifolius*, *oblongatus*, *pelagicus*, *torulosus*, *viresceus*, Seem. Fl. Vit. 254 = M. tortuosa.

CATYONA, Lindl. Veg. Kingd. 716 (1847), sphalm. = GATYONA, Cass. = **Crepis**, Linn. (Compos.).

CAUCALIS, Linn. Syst. ed. I (1735); Gen. ed. I. 71 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 928.
AGROCHARIS, Hochst. in Flora, xxvii. (1844) 19.
DAUCALIS, Pomel, Nouv. Mat. Fl. Atl. 148 (1874).
GLOCHIDOTHECA, Fenzl, in Russegg. Reise, i. ii. 970, obs. (1848).
LAPPULARIA, Pomel, Nouv. Mat. Fl. Atl. 149 (1874).
LIBRA, Boiss. in Ann. Sc. Nat. Sér. III. ii. 54 (1844).
TORILIS, Adans. Fam. ii. 99 (1763).
TURGENIA, Hoffm. Gen. Umb. 59 (1814).
TURGENIOPSIS, Boiss. in Ann. Sc. Nat. Sér. III. ii. 53 (1844).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 231 = Daucus abyssinicus.
aequicolorum, All. Fl. Pedem. ii. 33 = Anthriscus vulgaris.
africana, Crantz, Cl. Umb. Emend. 109 = Capnophyllum africanum.
africana, Thunb. Prod. Pl. Cap. 49 = capensis.
angustifolia, *Forsk. Fl. Aegypt. Arab. Suppl.* 208.—Arabia.
Anthriscus, *Huds. Fl. Angl.* 99.—Europ.
arvensis, *Huds. l. c.* 98.—Europ.; As. temp.
aspera, Lam. Fl. Fr. iii. 424 = C. Anthriscus.
asperas, Savi, Fl. Pis. i. 290 = arvensis.
bifrons, *Coss. & Dur. ex Baill. in Bourn. Linn. Soc.* xvi. (1878) 478.—Algeria.
caerulescens, *Boiss. Elench.* 93. p. 53.—Hispan.; Marocco.
canadensis, Crantz, Cl. Umb. Emend. 109 = Sanicula canadensis.
capensis, *Lam. Encyc.* i. 651.—Afr. austr.
capitata, Stokes, Bot. Mat. Med. ii. 64 = Sanicula europaea.
carnosa, Roth, Tent. Fl. Germ. i. 119 = Daucus Carota.
Carota, Huds. Fl. Angl. ed. II. 114 = Daucus Carota.
cenisifolia, Wall. ex DC. Prod. iv. 230 = C. Anthriscus.
cerefolium, Salisb. Prod. Hort. Chap. Allerton, 242 = Anthriscus Cerefolium.
cerefolium, Lam. Fl. Fr. iii. 424 = Anthriscus vulgaris.
cerefolium, Hoffm. Umb. 39 = Chaerophyllum Cerefolium.
cerefolium, Reichb. Fl. Germ. Excurs. 344 = Anthriscus Cerefolium.
chaerophyllum, Linn. Sp. Pl. 241 = Anthriscus Cerefolium.
chaerophyllum, Lam. Fl. Fr. iii. 424 = Anthriscus vulgaris.
chaerophyllum, Roth, Tent. Fl. Germ. i. 119 = Anthriscus Cerefolium.
chaerophyllum, Vill. Hist. Pl. Dauph. ii. 618 = Chaerophyllum temulum.
chaerophyllum, Jacq. Fl. Austr. v. 18 = Chaerophyllum aromaticum.
chaerophyllum, Jacq. Fl. Austr. v. 19 = Chaerophyllum hirsutum.
chaerophyllum, Jacq. Fl. Austr. v. 20 = Chaerophyllum elegans.
chaerophyllum, Jacq. Fl. Austr. v. 21 = Chaerophyllum Villarsii.
chaerophyllum, Jacq. Fl. Austr. v. 22 = Chaerophyllum Cicutaria.
chaerophyllum, Jacq. Fl. Austr. v. 23 = Chaerophyllum Cicutaria.
chaerophyllum, Jacq. Fl. Austr. v. 24 = Chaerophyllum Cicutaria.
chaerophyllum, Jacq. Fl. Austr. v. 25 = Chaerophyllum Cicutaria.
chaerophyllum, Jacq. Fl. Austr. v.

CAULOPHYLLUM, Michx. Fl. Bor. Am. i. 204. t. 21 (1803). *BERBERIDEAE*, Benth. & Hook. f. i. 43.
Pantheirotheca, Maxim. ex Regel, in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 223.
gracile, Dougl. ex Hook. Fl. Bor. Am. i. 30 = Vancouveria hexandra.
robustum, *Maxim. Prim. Fl. Amur* 88.—Manchuria.
thalictroides, *Michx. Fl. Bor. Am.* i. 205.—Am. bor.
thalictroides, Regel, in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 223 = robustum.

CAULOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = **Arabis**, L. n. (Crucif.).
alpina, Fourr. l. c. = A. brassicaeformis.

CAULOTRETUS, Rich. ex Spreng. Syst. iv. Cur. Post. 406 (1827) = **Bauhinia**, Linn. (Legumin.).
cordifolius, Rich. ex DC. Prod. ii. 517 = B. Richardiana.
eriophorus, Tenore, in Rendic. Soc. Borb. Neap. i. (1842) 410 = B. eriophora.
smilacinus, Schott. ex Spreng. Syst. iv. Cur. Post. 406 = B. smilacina.

CAULOTULIS, Rafin. Fl. Tellur. iv. 76 (1836) = **Ipomoea**, Linn. (Convolv.).

CAUSONIA, Rafin. Sylva Tellur. 87 (1838) = seq.

CAUSONIS, Rafin. Med. Fl. ii. 122 (1830) = **Vitis**, Linn. (Ampelid.).
japonica, Rafin. l. c. = V. japonica.
trifolia, Rafin l. c. = V. trifolia.

CAUSTIS, R. Br. Prod.[1] 259 (1810). *CYPERACEAE* Benth. & Hook. f. iii. 1067.
Eurostorhiza, Steud. Syn. Pl. Cyp. 265 (1855).
dioica, *R. Br. l. c.*—Austral.
flexuosa, *R. Br. l. c.*—Austral.
hexandra, Nees, in Lehm. Pl. Preiss. ii. 88 = dioica.
pentandra, *R. Br. Prod.* 239.—Austral.
recurvata, *Spreng. Syst. iv. Cur. Post.* 26.—Austral.
restiacea, *F. Muell. ex Benth. Fl. Austral.* vii. 421.—Austral.
Sieberi, Kunth, Enum. Pl. ii. 307 = Gahnia Sieberi.

CAUTLEA, Royle, Illustr. Bot. Himal. 361 (1839) = **Boscoea**, Sm. (Scitam.).
lutea, Royle, l. c. = K. lutea.

CAVALAM, Adans. Fam. ii. 357 (1763) = **Sterculia**, Linn.

CAVALLIUM, Schott, Melet. 33 (1832) = **Sterculia**, Linn.
comosum, Schott & Endl. l. c. = S. comosa.
urens, Schott & Endl. l. c. = S. urens.

CAVANALIA, Griseb. Cat. Pl. Cub. 76 (1864), sphalm. = **Canavalia**.

CAVANILLA, J. F. Gmel. Syst. 1087 (1791) = **Dombeya**, Cav. (Sterculiac.).
acerifolia, J. F. Gmel. l. c. = Pterospermum acerifolium.
acutangula, J. F. Gmel. l. c. = D. acutangula.
angulata, J. F. Gmel. l. c. = D. acutangula.
decanthera, J. F. Gmel. l. c. = Trochetia decanthera.
ferruginea, J. F. Gmel. l. c. 1088 = D. ferruginea.
ovata, J. F. Gmel. l. c. = D. ovata.
palmata, J. F. Gmel. l. c. 1087 = D. acutangula.
phoenicea, J. F. Gmel. l. c. 1088 = Pentapetes phoenicea.
punctata, J. F. Gmel. l. c. 1087 = D. punctata.
tiliaefolia, J. F. Gmel. l. c. = D. acutangula.
tomentosa, J. F. Gmel. l. c. = D. tomentosa.
umbellata, J. F. Gmel. l. c. = D. umbellata.

CAVANILLA, Salisb. Prod. 385 (1796) = **Stuartia**, Linn. (Ternstroem.).
floride, Salisb. l. c. = S. Malachodendron.

CAVANILLA, Thunb. Nov. Gen. 105 (1795) = **Pyrenacantha**, Hook. (Olacin.).
scandens, Thunb. l. c. = P. scandens.

CAVANILLA, Vell. Fl. Flum. 236 (1825); v. t. 103 (1827) = **Casearia**, St. Hil. (Euphorbiac.).
spinosa, Vell. ll. cc. = Casear. Vellesiana.

CAVANILLEA, Desr. in Lam. Encyc. iii. 663 (1789) = **Diospyros**, Linn. (Ebenac.).
philippinensis, Desr. l. c. = D. Embryopteris.
Mabola, Hiern, Ebenac. 260 = D. discolor.

CAVANILLEA, Medic. Malv. 19 (1787) = **Anoda**, Cav. (Malvac.).
hastata, Medic. l. c. = A. hastata.

CAVANILLESIA, Ruiz & Pav. Prod. 97. t. 20 (1794). *MALVACEAE*, Benth. & Hook. f. i. 211.
Pourretia, Willd. Sp. Pl. iii. 844 (1801).
cordata, *Spreng. Syst.* iii. 125.—Peruv.
platanifolia, *H. B. & K. Nov. Gen. et Sp.* v. 306.—Panama.
tuberculata, Kostel. Allg. Med. Pharm. Fl. v. 1872 = platanifolia ?
umbellata, *Ruiz & Pav. Syst. Veg.* 166.—Peruv.

CAVARIA, Steud. Nom. ed. I. 169 (1821), sphalm. = **Tovaria**, Ruiz & Pav. (Cappand.).

CAVENDISHIA, Lindl. Bot. Reg. sub t. 1791 (1836). *VACCINIACEAE*, Benth. & Hook. f. ii. 570.
Polybaea, Klotzsch, in Linnaea, xxiv. (1851) 80.
Froelesia, Klotzsch, l. c. 82.
Socratesia, Klotzsch, l. c. 22.
acuminata, *Benth ex Hemsl. Biol. Centr. Am. Bot.* ii. 272.—Mexic.
complectens, *Hemsl. l. c.*—Costa Rica.
crassifolia, *Hemsl. l. c* 273.—Mexic.
Endresii, *Hemsl. l. c.*—Costa Rica.
latifolia, *Donnel l. c.*—Mexic.
laurifolia, *Benth. & Hook. f. Gen.* ii. 570.—Guatemala.
melastomoides, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 272.—Costa Rica.
nobilis, *Lindl. Bot Reg.* sub t. 1791.—Am. austr.
pubescens, *Hemsl. Biol. Centr. Am. Bot.* ii. 273.—Panama.
Quercus, *Benth. & Hook. f. Gen.* ii. 570.—N. Granat.
tarapotana, *Benth. & Hook. f. l. c.*—Bras.
verruginosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 273.—Panama.
Warscewiczii, *Hemsl. l. c.* 274.—Guatemala.

CAVINIUM, Thou. Gen. Nov. Madag. 11 (1806) = **Vaccinium**, Linn.
madagascariense, Thou. ex Poir. in Dict. Sc. Nat. vii. 310 = *Agapetes madagascariensis*.

CAVOLINIA, Rafin. in Am. Monthly Mag. (1818) 175 = CAULINIA, Willd. = **Najas**, Linn.

CAYAPONIA, Silva Manso, Enum. Subst. Brasil. 31 (1836). *CUCURBITACEAE*.
Allagosperma, M. Roem. Syn. Pepon. 15 (1846).
Antagonia, Griseb. Pl. Lorentz. 144 (1874).
Alternasemina, Silva Manso, Cathart. Bras. (1836).
Cionandra, Griseb. Fl. Brit. W. Ind. 386 (1860).
Dermophylla, Silva Manso, Enum. Subst. Brasil. 30 (1836).
Druparia, Silva Manso, l. c. 35 (1836).
Periantiopodus, Silva Manso, l. c. 28 (1836)
Trianosperma, Mart. Syst. Mat. Med. Brasil. 79 (1843).
alata, *Cogn. in DC. Monog. Phan.* iii. 746.—Yucatan.
amazonica, *Cogn. l. c.* 797.—Bras.
americana, *Cogn. l. c.* 785.—Am. cent.
Andreana, *Cogn. in Bull. Acad. Belg. Sér. II.* xlix. (1880) 195.—Ecuador.
angustiloba, *Cogn. in DC. Monog. Phan.* iii. 771.—Bras.
attenuata, *Cogn. l. c.* 769.—Mexic.
Bonplandi, *Cogn. l. c.* 796.—Bras.
Boylinii, *Cogn. l. c.* 786.—Am. bor.
Burseavi, *Cogn. l. c.* 790.—N. Granat.
Cabocla, Mart. Syst. Mat. Med. Veg. Bras. 81 = globosa.
calycina, *Cogn. in Mart. Fl. Bras.* vi. iv. 78.—Bras.
citrullifolia, *Cogn. ex Griseb. in Goett. Abh.* xxiv. (1879) 136.—Reg. Argent.
coccinea, André, ex Cogn. in Bull. Acad. Belg. Sér. II. xlix. (1880) 196 = micrantha.
cordifolia, *Cogn. in Mart. Fl. Bras.* vi. iv. 77. t. 21.—Bras.
corlacea, *Cogn. l. c.* 79. t. 23.—Bras.

CAYAPONIA—
Crugeri, *Cogn. in DC. Monog. Phan.* iii. 778.—Ins. Trinit.
diffusa, *Silva Manso, Enum. Subst. Brasil.* 32.—Bras.
diversifolia, *Cogn. in DC. Monog. Phan.* iii 787.—Bras.
elliptica, *Silva Manso, Enum. Subst. Brasil.* 32.—Bras.
Espelinea, *Cogn. in DC. Monog. Phan.* iii. 792.—Bras.
excisa, *Cogn. l. c.* 747.—Cuba.
filifolia, *Cogn. l. c.* 789.—Bras.
floribunda, *Cogn. l. c.* 779.—Bras.
fluminensis, *Cogn. in Mart. Fl. Bras.* vi. iv. 76.—Bras.
glandulosa, *Cogn. in DC. Monog. Phan.* iii. 755.—Peruv.
Glaziovii, *Cogn. in Mart. Fl. Bras.* vi. iv. 74.—Bras.
globosa, *Silva Manso, Enum. Subst. Brasil.* 32.—Bras.
gracillima, *Cogn. in DC. Monog. Phan.* iii. 784.—Bras.
granatensis, *Cogn. l. c.* 794.—N. Granat.
grandiflora, *Cogn. l. c.* 779.—Yucatan.
heterophylla, *Cogn. l. c.* 758.—N. Granat.
hirsuta, *Cogn. in Mart. Fl. Bras.* vi. iv. 76.—Bras.
latebrosa, *Cogn. in DC. Monog. Phan.* iii. 776.—Ins. Canar.
latifolia, *Cogn. l. c.* 752.—Paraguay.
Lhotzkyana, *Cogn. l. c.* 765.—Bras.
longifolia, *Cogn. l. c.* 796.—Bras.
Martiana, *Cogn. l. c* 777.—Bras.
Maximowczii, *Cogn. l. c.* 745.—Mexic.
micrantha, *Cogn. in Bull. Acad. Belg.* Sér. II. xlix. (1880) 196.—N. Granat.
Ottoniana, *Cogn. in DC. Monog. Phan.* iii. 755.—Venezuela.
ovata, *Cogn. l. c* 754.—N. Granat.
palmata, *Cogn. in Bull. Soc. Bot. Belg.* xvii. (1878) 295.—Bras.
pedata, *Cogn. in Mart. Fl. Bras.* vi. iv. 80. t. 23.—Bras.
pentaphylla, *Cogn. in DC. Monog. Phan.* iii. 760.—Bras.
peruviana, *Cogn. l. c.* 795.—Peruv.
petiolulata, *Cogn. in Bull. Soc. Bot. Belg.* xvii. (1878) 295.—Bras.
pisubiensis, *Cogn. in DC. Monog. Phan.* iii. 770.—Bras.
pilosa, Cogn. in Mart. Fl. Bras. vi. iv. 75 = diffusa.
podantha, *Cogn. in DC. Monog. Phan.* iii. 763.—Paraguay.
Poeppigii, *Cogn. l. c.* 756.—Peruv.
racemosa, *Cogn. l. c.* 768.—Porto Rico.
rigida, *Cogn. l. c.* 790.—Guiana.
Ruizii, *Cogn. l. c.* 794.—Peruv.
Sandia, *Cogn. ex Griseb. in Goett. Abh.* xxiv. (1879) 135.—Reg. Argent.
setulosa, *Cogn. in DC. Monog. Phan.* iii. 767.—Bras.
simplicifolia, *Cogn. l. c.* 774.—N. Granat.
Tayuya, *Cogn. l. c.* 772.—Bras.
ternata, *Cogn. in Mart. Fl. Bras.* vi. iv. 79.—Bras.
Tibiricae, *Cogn. in DC. Monog. Phan.* iii. 784.—Bras.
tomentosa, *Cogn. l. c.* 747.—Peruv.
triangularis, *Cogn. l. c.* 774.—Guiana.
trifoliolata, *Cogn. l. c.* 789.—Bras.
trilobata, *Cogn. l. c.* 780.—Bras.
tubulosa, *Cogn. in Mart. Fl. Bras.* vi. iv. 78.—Bras.
villosissima, *Cogn. l. c.* 81. t. 24.—Bras.
Weddellii, *Cogn. in DC. Monog. Phan.* iii. 793.—Bras.

CAYLUSEA, A. St. Hil. 2de Mém. Reséd. 29 (1837). *RESEDACEAE*, Benth. & Hook. f. i. 111.
Stylexia, Rafin. Fl. Tellur. iv. 121 (1836).
Syntrophe, G. Ehrenb. ex Mucil. Arg. in Neue Denks. Schw. Ges. Zuerich, xvi. (1858) 227.
abyssinica, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vii. (1840) 43.—Abyss.
canescens, *A. St. Hil. 2de Mém. Reséd.* 29.—Afr. bor.; Arabia.
Moquiniana, *Webb, Fragm. Fl. Aethiop.* 28.—Afr. bor.

CAYRATIA, Juss. ex Guill. in Dict. Class. Hist. Nat. iv. 846 (1832) = **Vitis**, Linn. (Ampelid.) (V. pedata).

CEANOTHUS, Linn. in Act. Soc. Sc. Upsal. 77 (1741). *RHAMNEAE*, Benth. & Hook. f. i. 378.
FORRESTIA, Raña. in Med. Repos. N. York, II. iii. (1806) 422.
africanus, Linn. Sp. Pl. 196 = Noltea africana.
Alamani, DC. Prod. ii. 31 = Colubrina Alamani.
americanus, *Linn. Sp. Pl.* 195.—Am. bor.
americanus, Mill. Ic. t. 86 = Milleri.
arborescens, Mill. Gard. Dict. ed. VIII. n. 3 = Colubrina ferruginosa!
asiaticus, Linn. Sp. Pl. 196 = Colubrina asiatica.
atropurpureus, Rafin. New Fl. Am. iii. 56 = Nolteae sp.!
azillaris, Carr. in Rev. Horticn. (1876) 87 = azureus!
azureus, *Desf. Tabl.* ed. II. 282.—Mexic.
azureus, Kellogg, in Proc. Calif. Acad. i. (1855) 55 = hirsutus.
Baumannianus, Spach, Hist. Vég. Phan. ii. 460 = ovatus.
Bertini, Carr. in Rev. Horticn. (1872) 440 = azureus!
bicolor, Willd. in Schult. f. Syst. vii. 66 = hirsutus.
Burmannianus, Steud. Nom. ed. II. i. 313, splanin. = Baumannianus.
buxifolius, *Willd. ex Schult. f. Syst.* vii. 62.—Mexic.
caeruleus, *Lag. Gen. et Sp. Nov.* 11.—Mexic.
californicus, Kellogg, in Proc. Calif. Acad. i. (1855) 55 = integerrimus.
capensis, DC. Prod. ii. 30 = Scutia Commersonii.
capsularis, Forst. f. Prod. 18 = Colubrina asiatica.
celtidifolius, Cham. & Schlecht. in Linnaea, v. (1830) 602 = Colubrina celtidifolia.
Chloroxylon, *Vent. Syst. Laurin.* 660.—Jamaica.
circumcisus, Gaertn. Fruct. ii. 110, t. 106 = Scutia Commersonii.
collinus, Dougl. ex Knight & West. Fl. Cab. i. t. 18 = soredians!
colubrinus, Lam. Tabl. Encyc. ii. 90 = Colubrina ferruginosa.
cordulatus, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 124. f. 39.—Calif.
crassifolius, *Torr. in Pacif. Rail. Rep.* iv. 75.—Calif.
cubensis, *Lam. Tabl. Encyc.* ii. 90.—Ind. occ.
cuneatus, A. Gray, in Ives, Rep. Color. Riv. 9 = Greggii.
cuneatus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 267.—Calif.
decumbens, Hort. ex Steud. Nom. ed. II. i. 313 = americanus.
decumbens, *S. Wats. in Proc. Am. Acad.* x. (1875) 336.—Calif.
Delilianus, *Spach, Hist. Vég. Phan.* ii. 459.—Hab.!
dentatus, *Torr. & Gray, Fl. N. Am.* i. 268.—Calif.
depressus, *Benth. Pl. Hartw.* 8.—Mexic.
Dilleniianus, Hort. ex C. Koch, Dendrol. i. 620 = americanus.
discolor, Vent. Jard. Malm. 58 = Pomaderris elliptica.
divaricatus, Boland. Cat. Pl. Calif. 8 = cordulatus.
divaricatus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 266.—Calif.
divergifolius, Kellogg, in Proc. Calif. Acad. i. (1855) 57 = decumbens.
elegans, Lem. Illustr. Hortic. vii. (1860) t. 268 = thyrsiflorus.
elliptícus, *Rafin. New Fl. Am.* iii. 56.—Am. bor.
elongatus, Salisb. Prod. 140 = Noltea africana.
Fendleri, *A. Gray, Pl. Fendl.* 29.—Am. bor. occ.
ferreus, DC. Prod. ii. 30 = Scutia ferrea.
ferrugineus, Wendl. ex Steud. Nom. ed. II. i. 313 = Pomaderris lanigera.
floribundus, *Hook. Bot. Mag.* t. 4806.—Calif.
Fontanesianus, Spach, Hist. Vég. Phan. ii. 460 = ovatus.
glaber, *Spach, l. c.* 459.—Am. bor.
glandulosus, Rafin. New Fl. Am. iii. 57 = ovatus.
glandulosus, *Schlecht. in Linnaea*, xv. (1841) 474.—Mexic.
globulosus, Labill. Nov. Holl. Pl. i. 61. t. 85 = Spyridium globulosum.
glomeratus, Rafin. New Fl. Am. iii. 55 = americanus.
grandis, Dougl. ex Hook. Fl. Bor. Am. i. 125 = velutinus.
granulosus, *Ruiz & Pav. Fl. Per.* iii. t. t. 228.—Peruv.
Greggii, *A. Gray, Pl. Wright.* ii. 23.—Arizona.
Guadalupae, Steud. Nom. ed. II. i. 313 = laevigatus.
guineensis, DC. Prod. ii. 30 = Chailletia toxicaria!
Hartwegii, *Hook. ex Heynh. Nom.* ii. 126, nomen.—Mexic.
herbaceus, Rafin. in Med. Repos. N. York, v. (1808) 360 = americanus.

CEANOTHUS:—
hirsutus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 266.—Calif.
hybridus, Hort. ex C. Koch, Dendrol. i. 620 = americanus.
hypericifolius, L'Hérit. ex DC. Prod. ii. 32 = microphyllus.
incanus, Torr. & Gray, Fl. N. Am. i. 265.—Calif.
infestus, H. B. & K. Nov. Gen. et Sp. vii. 61 = Adolphia infesta.
integerrimus, *Hook. & Arn. Bot. Beech. Voy.* 329.—Am. bor. occ.
intermedius, Hook. Fl. Bor. Am. i. 124 = ovatus.
intermedius, Pursh, Fl. Am. Sept. i. 167 = americanus.
laevigatus, *DC. Prod.* ii. 30.—Ind. occ.
laevigatus, Hook. Fl. Bor. Am. i. 125 = velutinus.
lanciifolius, Moench, Meth. 651 = Noltea africana.
lanigerr, Andr. Bot. Rep. t. 569 = Pomaderris lanigera.
latifolius, *Rafin. New Fl. Am.* iii. 55.—Am. bor.
Leschenaultii, *DC. Prod.* ii. 31.—Zeylan
lewisianus, Rafin. New Fl. Am. iii. 55 = americanus.
Lobbianus, Hook. Bot. Mag. t. 4810 = dentatus.
macrocarpus, *Cav. Ic.* iii. 38, t. 270.—Mexic.
macrocarpus, Hort. ex Steud. Nom. ed. II. i. 313 = americanus.
macrocarpus, Mcq. & Sesse, ex DC. Prod ii. 32, 33 = Mocinianus.
macrocarpus, Nutt. in Torr. & Gray, Fl. N. Am. i. 267 = cuneatus.
macrophyllus, Desf. Tabl. ed. II. 232 = americanus.
magacarpus, Nutt. Syll. Am. ii. 46 = cuneatus.
microphyllus, *Mschx. Fl. Bor. Am.* i. 154.—Am. bor.
Milleri, *Tausch, in Flora*, xxi. (1838) t. *Beibl.* 79.—Am. bor.
Mocinianus, *DC. Prod.* ii. 32.—Mexic.
mystacinus, DC. l. c. 31 = Helmus scandens.
napalensis, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 375 = Rhamnus nipalensis.
Neumanni, *Tausch, in Flora*, xxi. (1838) 788.—Am. bor.
nevadensis, Kellogg, in Proc. Calif. Acad. ii. (1863) 125 = integerrimus.
nitidus, Torr. in Pacif. Rail. Rep. iv. 75 = soredians.
officinalis, Rafin. Med Fl. ii. 205 = americanus.
oligranthus, Nutt in Torr. & Gray, Fl. N. Am. i. 266 = hirsutus.
oreganus, Nutt. l. c. 265 = sanguineus.
ovalifolius, *Wendr. in Schr. Naturf. Ges. Marb.* ii. (1880) 247.—Am. bor.
ovalis, Bigel. Fl. Bost. ed. II. 99 = ovatus.
ovatus, *Desf. Arb.* ii. 881.—Am. bor.
ovatus, Hort. ex Lindl. Bot. Reg. (1840) t. 20 = seq.
pallidus, *Lindl. l. c.*—Hab.?
paniculatus, Roth, Nov. Pl. Sp. 154 = Celastrus paniculata.
papillosus, *Torr. & Gray, Fl. N. Am.* i. 268.—Calif.
pauciflorus, *Mcq. & Sesse, ex DC. Prod.* ii. 33.—Mexic.
perennis, Pursh, Fl. Am. Sept. i. 167 = americanus.
Pitcheri, Pickering, ex Torr. & Gray, Fl. N. Am. i. 264 = americanus.
prunifolius, Hort. ex C. Koch, Dendrol. i. 619 = americanus.
prostratus, *Benth. Pl. Hartw.* 302.—Calif.
pubescens, *Ruiz & Pav. Fl. Per.* iii. t. t. 228.—Peruv.
pubiflorus, DC. Prod. ii. 30 = Zizyphus pubiflora.
pulchellus, Delile, ex Spach, Hist. Vég. Phan. ii. 459 = Delilianus.
reclinatus, Bosc, ex Steud. Nom. ed. II. i. 313 = americanus.
reclinatus, *L'Hérit. Sert. Angl.* 6.—Ind. occ.
rigidus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 268.—Calif.
sanguineus, *Pursh, Fl. Am. Sept.* i. 167.—Am. bor. occ.
sanguineus, Nutt. Gen. Am. i. 153 = americanus.
Sarcomphalus, DC. Prod. ii. 30 = Sarcomphalus laurinus.
scandens, D. Dietr. Syn. Pl. i. 812 = Noltea africana.
serpyllifolius, *Nutt. Gen. Am.* i. 154.—Florida.
soredíatus, *Hook. & Arn. Bot. Beech. Voy.* 328.—Calif.
spathulatus, Labill. Nov. Holl. Pl. i. 60. t. 84 = Trymalium Billardieri.
sphaerocarpus, DC. Prod. ii. 30 = Rhamnus sphaerosperma.
spinosus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 267.—Calif.

CEANOTHUS:—
tardiflorus, Hornem. Hort. Hafn. i. 230 = americanus.
thyrsiflorus, *Eschw. in Mém. Acad. Petersb.* Sér. VI x. (1826) 235.—Calif.
trifloras, Steud. Nom. ed. II. i. 313 = Colubrina triflora.
trinervus, Moench, Meth. 651 = americanus.
triqueter, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 376 = Rhamnus triqueter.
Veitchianus, *Hook. Bot. Mag.* t. 5127.—Calif.
velutinus, *Dougl. in Hook. Fl. Bor. Am.* i. 125. t. 45. —Am. bor. occ.
verrucosus, Nutt. in Torr. & Gray, Fl. N. Am. i. 267 = cuneatus.
virgatus, Rafin. New Fl. Am. iii. 56 = americanus.
Wendlandianus, Roem. & Schult. Syst. v. 299 = Pomaderris ferruginea.
Wightianus, Wall. Cat. n. 4264 = Rhamnus Wightii.
zeylanicus, Heyne, in Roth, Nov. Pl. Sp. 158 = Scutia Commersonii.

CEARIA, Dum. Comm. Bot. 65 in adnot. (1822) = Euryoles, Salisb. (Amaryllid.).
amboinensis, Dum. l. c. = E. sylvestris.

CEBATHA, Forsk. Fl. Aegypt. Arab. 171 (1775) = Cocculus, DC. (Menisperm.) (C. Cebatha).

CECICODAPHNE, Nees, in Wall. Pl. As. Rar. ii. 70 (1831) = Cinnamomum, Blume (Laurin.).
glaucescens, Nees, l. c. = Cinnam. Cecicodaphne.

CECROPIA, Linn. in Loefl. It. 272 (1758). *URTICACEAE*, Benth. & Hook. f. iii. 378.
AMBAIBA, Adans. Fam. ii. 377 (1763).
aculifolia, *Trèc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 81.—Peruv.
adenopus, *Mart. ex Miq. in Mart. Fl. Bras.* iv. i 147. —Am. trop.
albicans, *Trèc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 82.—Peruv.
angustifolia, *Trèc. l. c.* 83.—Peruv.
argentea, *Viviani, Ort. Bot. Padov.* 135.—Hab.?
bicolor, *Klotzsch, in Linnaea*, xx. (1847) 581.—Peruv.
carbonaria, *Mart. ex Miq. in Mart. Fl. Bras.* iv. i 144.—Am. trop.
cinerea, *Miq. l. c.* 142.—Am. trop.
commutata, *Schott, ex Miq. l. c.* 148.—Am. trop.
concolor, Hort. Schoenbr.—ex Miq. l. c.—Schlechtena.
concolor, *Willd. Sp. Pl.* iv. 662.—Bras.
cyrtostachya, *Miq. in Mart. Fl. Bras.* iv. i. 145.—Am. trop.
dentata, *Klotzsch, in Linnaea*, xx. (1847) 588.—Peruv.
digitata, *Tenore, ex Miq. in Mart. Fl. Bras.* iv. i. 149. —Bras.
digitata, Klotzsch, in Linnaea, xx. (1847) 584 = angustifolia.
fragilifyra, Trèc. in Ann. Sc. Nat. Sér. III. viii. (1847) 81 = bicolor.
frigida, *Dufresne, in Illustr. Hortic.* xx. (1873) 70.—Hab.?
Goudotiana, *Trèc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 83.—N. Granat.
hololeuca, *Miq. in Mart. Fl. Bras.* iv. i. 148.—Am. trop.
Humboldtiana, *Klotzsch, in Linnaea*, xx. (1847) 580. —Am. trop.
insignis, *Liebm. in Videnskab. Selsk. Skr.* V. ii. (1851) 218.—Nicaragua.
Klotzschiana, *Miq. in Mart. Fl. Bras.* iv. i. 151.—Am. trop.
latiloba, *Miq. l. c.* 147.—Am. trop.
leucoconia, *Miq. l. c.* 142.—Am. trop.
leucophaea, *Poepp. ex Miq. l. c.* 151.—Am. trop.
lyratiloba, *Miq. l. c.* 144.—Am. trop.
membranacea, *Trèc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 83.—Ind. occ.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* iii. 151.—Mexic.
nivea, *Poepp. ex Klotzsch, in Linnaea*, xx. (1847) 588. —Peruv.
obtusa, *Trèc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 79.—Bras.
obtusifolia, *Bertol. Fl. Guatimal.* 99.—Guatemala.
pachystachya, *Trèc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 80.—Bras.
palmata, *Willd. Sp. Pl.* iv. 662.—Bras.

CECROPIA :—
panamensis, *Hemsl. Biol. Centr. Am. Bot.* iii. 151.—Panama.
pelata, *Linn. Pugill. Pl. Jamaic.* 28 (*Amoen. Acad.* v. 410).—Am. trop.
peltata, G. F. W. Mey. Prim. Fl. Esseq. 280 = surinamensis.
peltata, Ruiz, ex Klotzsch, in Linnæa, xx. (1847) 582 = Ruizeana.
peltata, Schreb. ex Miq. in Mart. Fl. Bras. iv. i. 150 = Schreberiana.
peltata, Vell. Fl. Flum. x. t. 101 = adenopoda.
peltata, Willd. ex Klotzsch, in Linnæa, xx. (1847) 580 = Humboldtiana.
pinnatiloba, *Klotzsch, in Linnæa*, xx. (1847) 588.—Peruv.
propinqua, *Miq. in Mart. Fl. Bras.* iv. i. 149.—Am. trop.
Ruizeana, *Klotzsch, in Linnæa*, xx. (1847) 582.—Peruv.
scabra, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 95.—Bras.
scabra, Ruiz, ex Klotzsch, in Linnæa, xx. (1847) 581 = Klotzschiana.
Schiedeana, *Klotzsch, l. c.*—Mexic.
Schreberiana, *Miq. in Mart. Fl. Bras.* iv. i. 150.—Am. trop.
sciadophylla, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 93.—Bras.
strigosa, *Trèc. in Ann. Sc. Nat.* Ser. III. viii. (1847) 32.—Peruv.
surinamensis, *Miq. in Mart. Fl. Bras.* iv. i. 143.—Am. trop.
tubulosa, *Ruiz, ex Klotzsch, in Linnæa*, xx. (1847) 584.—Peruv.

CEDRELA, P. Br. Hist. Jamaic. 158 (1756). *MELIACEÆ*, Benth. & Hook f. i. 389.
CEDRUS, Mill. Gard. Dict. ed. VII (1759).
CUVERACA, Jones, in As. Res. iv. (1795) 281.
JOHNSONIA, Adans. Fam. ii. 348 (1763).
PTEROSIPHON, Turcz. in Bull. Soc Nat. Mosc. (1858) i. 589.
TOONA, M. Roem. Syn. Hesper. 131 (1846).
adenophylla, Mart. ex C. DC. in DC. Monog. Phan. i. 728 = paraguaiensis.
alternifolia, *Steud. Nom.* ed. I. 170.—Honduras.
angustifolia, *Moç. & Sessé, ex DC. Prod.* i. 624.—Mexic.
australis, F. Muell. Fragm. i. 4 = C. Toona.
bogotensis, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 377.—N. Granat.
brasiliensis, Juss. in St. Hil. Fl. Bras. Mer. ii. 86 = fissilis.
Cedro, Loefl. ii. 183 = odorata.
Degesii, *S. Wats. in Proc. Am. Acad.* xvii. (1882–83) 190.—Mexic.
febrifuga, Forsten, Diss. Cedrel. 16 = C. Toona.
fissilis, *Vell. Fl. Flum.* 75; t. 68.—Bras.
glabra, C. DC. in DC. Monog. Phan. i. 742 = C. Toona.
Glaziovii, *C. DC. in Mart. Fl. Bras.* xi. i. 224. t. 65. f. 2.—Am. trop.
guianensis, *A. Juss. in Mém. Mus. Par.* xix. (1830) 255.—Guiana.
hexandra, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 425 = C. Toona.
inodora, *Hassk. Hort. Bogor.* i. 181.—Java.
Kotschyi, *Schweinf. Reliq. Kotsch.* 36. t. 35.—Afr. trop.
longifolia, Wall. Cat. n. 1278 = C. Toona.
Mahagoni, Linn. Syst. ed. X. 940 = Swietenia Mahagoni.
mexicana, M. Roem. Syn. Hesper. 137 = Swietenia humilis.
microcarpa, C. DC. in DC. Monog. Phan. i. 745 = C. Toona.
montana, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) 415.—Venezuela.
multijuga, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 297 = C. Toona.
odorata, Blanco, Fl. Filip. ed. I. 284 = C. Toona.
odorata, Cham. & Schlecht. in Linnæa, vi. (1831) 422 = Swietenia humilis.
odorata, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 169 = Glaziovii.
odorata, *Linn. Syst.* ed. X. 940.—Am. austr.

CEDRELA :—
odorata, Ruiz & Pav. Fl. Per. iii. 9 = angustifolia.
odorata, Vell. Fl. Flum. 74; ii. t. 67 = Velloziana.
paraguaiensis, *Mart. in Flora*, xx. (1837) *Beibl.* 93.—Paraguay.
Rosmarinus, Lour. Fl. Cochinch. 160 = Bacckia cochinchinensis.
serrata, Royle, Illustr. Bot. Himal. 144. t. 25 = C. Toona.
serrulata, *Miq. Fl. Ind. Bat. Suppl.* i. 508.—Sumatra.
sinensis, *Juss. in Mém. Mus. Par* xix. (1830) 255.—China.
Teysmanni, *Hassk. Hort. Bogor.* i. 181.—Java.
Toona, *Roxb ex Rottl. in Ges. Naturf. Fr. Neue Schr.* ii. (1808) 198.—Malaya; Austral.
Velloziana, *M. Roem. Syn. Hesper.* 137.—Am. trop.
velutina, DC. Prod. i. 625 = Chickrassia tabularis.
villosa, Roxb. Hort. Beng. 18 = Chickrassia tabularis.

CEDRELLA, Scop. Introd. 324 (1777) = **Cedrela**, R. Br.

CEDRONELLA, Riv. ex Rupp, Fl. Jen. ed. Hall. 225 (1745); Moench, Meth. 411 (1794). *LABIATÆ*, Benth. & Hook f. ii. 1200.
brevifiora, *A. Gray, in Proc. Am. Acad.* xx. (1885) 309.—Texas.
cana, *Hook. Bot. Mag.* t. 4618.—Mexic.
canariensis, *Webb & Berth. Phyt. Canar.* iii. 87.—Ins. Canar.
cordata, *Benth. Lab. Gen. et Sp.* 502.—Am. bor.
hastifolia, Regel, Ind. Sem. Hort. Petrop. (1868) 35 = cana.
japonica, *Hassk. in Verh. Natuurk. Ver. Ned. Ind.* i. (1856) 36 — Japon.
mexicana, *Benth. Lab. Gen. et Sp.* 502.—Mexic.
mollis, *A. Gray, in Proc. Am. Acad.* viii. (1870) 369.—Am. bor. occ.
pallida, *Lindl. Bot. Reg.* (1846) t. 29.—Am bor. occ.
triphylla, *Moench, Meth.* 411.—Ins. Canar.

CEDROTA, Schreb. Gen. i. 259 (1789) = **Ocotea**, Aubl. (Laurin.).
guianensis, Blanco, Fl. Filip. ed. II. 218 = Pisonia umbellifera.
guianensis, Raeusch. Nom. ed. III. 111 = O. communata.
longifolia, Willd. Sp. Pl. ii. 338 = O. commutata.

CEDRUS, Mill. Gard. Dict. ed. VII (1759) = **Cedrela**, Linn. (Meliac.).
alternifolia, Mill. l. c. ed. VIII. n. 3 = Cedrela alternifolia.
Mahagoni, Mill. l. c. n. 2 = Swietenia Mahagoni.
odorata, Mill. l. c. n. 1 = Cedrela odorata.

CEDRUS, [Tourn.] Mill. Gard. Dict. ed. III (1737); Loud. Arb. Brit. iv. 2402 (1838). *CONIFERÆ*, Benth. & Hook. f. iii. 439.
africana, Gord. ex Knight, Syn. Conif. 42 = Libani.
argentea, Hort. ex Carr. Conif. ed. I. 286 = Libani.
atlantica, Manetti, Cat. Hort. Madoet. Suppl. 8 = Libani.
Deodara, Loud. Arbor. Brit. iv. 2428 = Libani.
elegans, Knight, Syn. Conif. 42 = Libani.
indica, Chambr. Trait. Prat. Arbr. Réan. Conif. 341 = Libani.
Libani, *Barrel. Ic.* 499.—Aa. Min.; Syria; Afghan.; Reg. Himal.; Alger.
patula, C. Koch, Dendrol. ii. ii. 268 = Libani.

CEIBA, Medic. Malv. 15 (1787); Gaertn. Fruct. ii. 244. t. 138 (1791) = **Briodendron**, DC. (Malvac.).
Caseoria, Medic. l. c. 16 = E. orientale.
pentandra, Gaertn. Fruct. ii. 244. t. 138 = E. anfractuosum.

CELAENA, Wedd. Chlor. And. i. 230 (1855) = **Oligandra**, Less. (Compos.).
pachymorpha, Wedd. l. c. = O. pachymorpha.

CELASINE, Pritz. Ic. Ind. 284, sphalm. (1855) = **Gelasine**, Herb. (Irid.).

CELASTRUS, Linn. Gen. ed. I. 59 (1737). *CELASTRINEÆ*, Benth & Hook. f. i. 364 (365).
CATHA, G. Don, Gen. Syst. ii. 11 (1832).
GUEVINIA, Hort. Par. ex Decne. in Rev. Hortic. Ser. II. iv. (1845–46) 425.
ORIXA, Thunb. Nov. Gen. 66 (1783); Fl. Jap. 3 (1784).
SEMARILLA, Rafin. Sylva Tellur. 146 (1838).
acuminatus, *Linn. f. Suppl.* 154.—Afr. austr.
acuminatus, *Rafin Sylva. Fl. Am.* iii. 14.—Am. bor.
acuminatus, Wall. Cat. n. 4842 = Chailletia gelonioides.
adenophylla, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 85 = Ilex crenata.
alatus, Thunb. Fl. Jap. 98 = Evonymus Thunbergianus.
alnifolius, D Don, Prod Fl. Nep. 190 = paniculatus.
alpestris, *Blume, Bydr.* 1145.—Java.
andongensis, *Oliver, Fl. Trop. Afr.* i. 361.—Afr. trop.
angularis, *Sond. in Harv. & Sond. Fl. Cap.* i. 460.—Afr. austr.
angulata, *Maxim. in Bull. Acad. Petersb.* xxvii. (1881) 455.—China.
aphyllus, *Schlecht. in Linnæa*, xv. (1841) 458.—Mexic.
aquifolius, *Regel, Ind. Sem. Hort. Petrop.* (1856) 36.—Hab.?
arbutifolius, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 133.—Abyss.
articulatus, *Thunb. Fl. Jap.* 97.—Japon.
Atkaio, *A. Rich. Tent. Fl. Abyss.* i. 132.—Abyssin.
attenuatus, Wall. Cat. n 4319 = Gymnosporia neglecta.
auriculatus, Vitm Summa, ii. 31, in syn. = articulatus.
australis, *Harv & F. Muell. in Trans. Phil. Soc. Vict.* i (1855) 41.—Austral.
barbarus, Linn. Mant. ii. 211, in obs. = Cassine barbara.
bilocularis, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 31.—Austral.
brasiens, Jack, in Malay Misc i. (1820) n. v. 19 = Microtropis bivalvis.
Boaria, Dall. Hist. Pl. vi. 26 = Maytenus chilensis.
bullatus, Hort. ex Steud. Nom. ed. II. i. 314 = pyracanthus.
bullatus, Linn. Sp. Pl. 196 - scandens.
buxifolia, Dreg, ex Walp. Ann. i. 189 = Catha patens.
buxifolius, *Linn. Sp. Pl.* 197.—Afr. austr.
buxifolius, Roxb. Hort. Beng. 18, nomen = Gymnosporia Wallichiana.
campestris, Eckl. & Zeyh. Enum. 119 = pyracanthus.
cantoniensis, *Hance, in Journ. Bot.* xxiii. (1885) 325.—China.
capensis, Thunb. ex Steud. Nom. ed. II. i. 314 = excisus.
capitatus, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 458.—Afr. austr.
casmnifolius, Salisb. Prod. 141 = lucidus.
cassinoides, L'Hérit. Sert. Angl. 6. t. 10 = Gymnosporia cassinoides.
cernuus, *Thunb. Prod. Pl Cap.* 42.—Afr. austr.
Championi, *Benth. in Hook. Kew Journ.* iii. (1851) 334 — Ins. Hongkong.
cilioides, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 85 = flagellaris.
collinus, Eckl. & Zeyh. Enum. 119 = undatus.
Coipoon, Hort. ex Steud. Nom. ed. II. i. 315 = pyracanthus
concavus, Lam. Tabl. Encyc. ii. 93 = lucidus.
confertus, *Kurz & Pav. Fl. Per.* iii. 7.—Peruv.
cordatus, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* 454.—Afr. austr.
coriaceus, Gmll. & Perr. Fl. Seneg. Tent. i. 142 = Gymnosporia montana, senegalensis.
cornuculatus, *L'Herit. Sert. Angl.* 6.—Afr. austr.
crassifolius, Salisb. Prod. 141 = cassinoides.
crenatus, *Forst. f. Prod.* 19.—Ins. Marchion.
crenatus, Roth, Nov. Pl. Sp. 186 = Gymnosporia montana.
crenulatus, *Wall. Cat.* n. 4323.—Siam.
crispulus, *Regel, Gartenfl.* (1860) 407.—China.
crispus, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 28 = Euclea ovata.
Cunninghamii, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 80.—Austral.
cymatodes, Spreng. Syst. i. 775 = undatus.

CELASTRUS :—

cymosus, Eckl. & Zeyh. Enum. 121 = heterophyllus.
cymosus, Noland. in Bot. Mag. t. 2070 = buxifolius.
decolor, Delile, Voy. Méroe, 100. t. 64. f. 6 = Gymnosporia montana, senegalensis.
dependens, Ait. Cat. 4302; et in Roxb. Fl. Ind. ed. Carey, i. 389 = paniculatus.
diffusus, G. Don, Gen. Syst. h. 6 = Ventilago leiocarpa.
dilatatus, *Thunb. in Trans. Linn. Soc.* ii. (1794) 332. —Japon.
dispermus, *F. Muell in Trans. Phil. Inst. Vict.* iii. (1859) 31.—Austral.
dubius, *Spreng. Syst.* i. 774.—Ind. occ.
dumetorum, Eckl. & Zeyh. Enum. 119 = undatus.
edulis, Hochst ex A. Rich. Tent. Fl Abyss. i. 131 = serratus.
edulis, Vahl, Symb. Bot. ii. 52 = Catha edulis.
elliptica, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1808) 22.—Afr. austr.
emarginatus, Ruiz & Pav. Fl. Per. iii. 6 = retusus.
emarginatus, Willd. Sp. Pl. i. 1128 = Gymnosporia emarginata.
empleurifolius, Eckl. & Zeyh. Enum. 121 = buxifolius.
enonymifolius, *Burm. f. Fl. Cap. Prod.* 6.—Afr. austr.
enonymoides, *Welw. ex Oliver, Fl. Trop. Afr.* i. 362.—Afr. trop
europæus, Boiss. Elench. 29 = Gymnosporia montana.
excisus, *Thunb. Fl. Cap.* ii. 119.—Afr. austr.
filiformis, Lam. f. Suppl. 153 (= *Mystroxylon filiforme*).—Afr. austr.
Finlaysonianus, *Wall. Cat.* n. 4824.—Siam; China.
flagellaris, *Rupr. in Bull. Phys-Math. Acad. Pétersb.* xv. (1857) 357.—Reg. Amur.
flexuosus, *Thunb. Fl. Cap.* ii. 124.—Afr. austr.
floribundus, *Span. in Linnæa*, xv. (1841) 186.—Hab.?
Fournieri, *Panch. & Sebert, Not. Bois Nouv. Calléd.* 284.—N. Caled.
glaucus, R. Br. in Salt, Voy. Abyss. App. 64 = Gymnosporia montana.
glaucus, Wahl, Symb. Bot. ii. 42 = Elæodendron glaucum.
glomeratus, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 458 = heterophyllus.
glomeratus, E. Mey. ex Walp. Rep. v. 401 = patens.
gomiacualis, Eckl. & Zeyh. Enum. 120 = buxifolius.
gracillipes, *Welw. ex Oliver, Fl. Trop. Afr.* i. 361.—Afr. trop.
Haenkei, Spreng. Syst. iv. Cur. Post. 88 = Schœpfia flexuosa.
Hamelii, Spreng. Syst. i. 774 = Rhamnus racemosus?
heterophyllus, *Eckl. & Zeyh. Enum.* 120.—Afr. austr.
heterophyllus, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 467 = Cassine latifolia.
heterophyllus, *Savi, in Mem. Torino*, xxxviii. (1835) 173.—Hab.?
Heyneana, Roth, in Roem. & Schult. Syst. v. 421 = Gymnosporia Heyneana.
Hindsii, *Benth. in Hook. Kew Journ.* iii. (1851) 334 = monospermus.
hispanicus, Hort. ex DC. Prod. ii. 8 = multiflorus.
huillensis, *Welw. ex Oliver, Fl. Trop. Afr.* i. 364.—Afr. trop.
humilis, Eckl. & Zeyh. Enum. 120 = buxifolius.
ilicifolius, Schrad. in Goett. Gel. Anz. i. (1821) 716 = Maytenus truncatus.
illicinus, *Burch. ex Loud. Hort. Brit.* 83, nomen.—Afr. austr.
illicinus, Kraus. ex Sond. in Harv. & Sond. Fl. Cap. i. 457 = procumbens.
inæquus, Zeyh. ex Sond. l. c. 455 = polyacanthus.
integer, Thunb. Fl. Cap. ii. 123 = pyracanthus.
integrifolius, *Linn. f. Suppl.* 153.—Afr. austr.
Jacksianus, Steud. Nom. ed. II. i. 314 = lucidus.
japonicus, *C. Koch, Dendrol.* i. 695.—Japon.
Jodina, Steud. Nom. ed. II. i. 314 = Iodina rhombifolia.
kirsiana, *Franch. & Sav. Enum. Pl. Jap.* ii. 314.—Japon.
lanceolatus, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 456.—Afr. austr.
lanceolatus, *Spreng. Syst.* i. 775.—Ins. Mascar.
lanciifolius, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 132.—Afr. trop.
laurifolius, *A. Rich. Tent. Fl. Abyss.* i. 130.—Abyssin.

CELASTRUS :—

laurinus, *Thunb. Prod. Pl. Cap.* 42.—Afr. austr.
leptopus, Bernh. ex Walp. Rep. v. 401 = buxifolius.
linearis, Eckl. & Zeyh. Enum. 121 = polyacanthus.
linearis, *Linn. f. Suppl.* 153.—Afr. austr.
lineatus, D. Dietr. Syn. i. 815 = linearis.
lucidus, *Linn. Mant.* i. 49.—Afr. austr.
lucidus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 400.—Ind. or.
luteolus, *Delile, in Ann. Sc. Nat.* Sér. II. xx. (1843) 90.—Afr. trop.
lycioides, *Brouss. ex Willd in Roem. & Schult. Syst.* v. 497.—Ins. Canar.
macrocarpus, *Ruiz & Pav. Fl. Per.* iii. 8. t. 280. f. 16.—Peruv.
macrophyllus, Welw. ex Oliver, Fl. Trop. Afr. i. 364 = buxifolius.
magellanicus, DC Prod. ii. 8 = Maytenus magellanicus.
mauritianus, *Stadtm. ex Willem. in Ust. Ann. Bot* xvii. (1796) 21.—Ins. Maurit.
Maytenus, Willd. Sp. Pl. i. 1127 = Maytenus Boaria.
Metsuanus, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 448 = paniculatus.
mexicanus, *Moç. & Sesse, ex DC. Prod.* ii. 8.—Mexic.
micrantha, *Roxb. Hort. Beng.* 86; *Fl Ind.* i. 625.—Ins. Moluco.
microcarpus, *D. Don, Prod. Fl. Nep.* 191.—Reg. Himal.
microphyllus, Eckl. & Zeyh. Enum. 119 = acuminatus.
microphyllus, *Linn. f. Suppl.* 154.—Afr. austr.
mollis, *Decne. in Rev. Hortic.* Sér. II. iv. (1845-46) 425.—Hab.?
monospermus, *Roxb. Hort. Beng.* 18; *Fl. Ind.* i. 626.—Ind. or.; China.
montanus, Roth, Nov. Pl. Sp. 154 = Gymnosporia montana.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* i. 112.—Afr. trop.
mucronatus, Eckl. & Zeyh. Enum. 119 = acuminatus.
Muelleri, *Benth. Fl. Austral.* i. 399.—Austral.
multiflorus, Eckl. & Zeyh. Enum. 121 = buxifolius.
multiflorus, *Lam. Encyc.* i. 661.—Afr. austr.
multiflorus, Roxb. Hort. Beng. 18; Fl. Ind. 622 = paniculatus.
myrtifolius, *Linn. Sp. Pl.* 196.—Jamaica.
neglectus, Wall. Cat. n. 4341 = Gymnosporia neglecta.
nemorosus, *Eckl. & Zeyh. Enum.* 120.—Afr. austr.
nepalensis, Steud. Nom. ed. II. i. 315 = Pittosporum floribundum.
nossibeus, *O. Hoffm. Sert. Madag.* 12 (= Alsodeia).—Madag.
nutans, Roxb. Hort. Beng. 18; Fl. Ind. i. 623 = paniculatus.
obovatus, Hochst. ex Oliver, Fl. Trop. Afr. i. 362 = Gymnosporia montana.
obscurus, A. Rich. Tent. Fl. Abyss. i. 131 = serratus.
obtusatus, Presl, Bot. Bemerk. 84 = Simmondsia californica.
obtusifolius, Roxb. Hort. Beng. 86; Fl. Ind. i. 625 = Gymnosporia trigyna.
obtusus, Thunb. in Hoffm. Phytogr. Blaetter, i. (1808) 29 = pyracanthus.
octogonus, L'Hérit. Sert. Angl. 6 = Maytenus octogonus.
octogonus, Hort. ex Steud. Nom. ed. II. i. 315 = pyracanthus.
oleifolius, Pers. Syn. i. 297 = oleoides.
oleoides, *Lam. Tabl. Encyc.* ii. 93.—Afr. austr.
opposita, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 398 = Pleurostylia Wightii.
orbicularis, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 493 = Maytenus nitghiosus.
orbiculatus, Lam. Tabl. Encyc. ii. 94 = articulatus.
Orixa, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1846) 150.—Japon.
ovalifolius, Steud. Nom. ed. II. i. 315 = Gymnosporia ovata.
ovatus, *Hill, Veg. Syst.* xiii. 62; *Hort. Kew.* 437. t. 16. —Ins. Guinea.
ovatus, Wall. Cat. n. 4308 = Gymnosporia ovata.
oxyphyllus, Wall. Cat. n. 4312 = Gymnosporia acuminata.
pallidus, Wall. Cat. n. 4307 = Gymnosporia montana.
paniculatus, *Willd. Sp. Pl.* i. 1125.—Asa. trop.; Malaya.
parvifolius, *A. Rich. Ess. Fl. Cub.* 249.—Cuba.
parvifolius, Eckl. & Zeyh. Enum. 121 = heterophyllus.
parvifloras, *Vahl, Symb. Bot.* i. 21.—Arabia.
patens, *Eckl. & Zeyh. l. c.* 120.—Afr. austr.

CELASTRUS :—

pauciflorus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 400.—Ind. or.
pedunculatus, *Sond. in Harv. & Sond. Fl. Cap.* i. 454.—Afr. austr.
pentagynus, *Zipp. ex Span. in Linnæa*, xv. (1841) 186.—Malaya!
peruvianus, F. G. Dietr. Vollst. Lexik. Gaertn. ii. 111 = retusus.
phyllacanthus, L'Hérit. Sert. Angl. 6 = Gymnosporia montana.
Plectronia, DC. Prod. ii. 9 = acuminatus.
polyacanthus, *Sond. in Harv. & Sond. Fl. Cap.* i. 455.—Afr. austr.
polyanthemos, *Eckl. & Zeyh. Enum.* 121.—Afr. austr.
polybotrys, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 449.—Ins Philipp.
populifolius, *Lam. Tabl. Encyc* ii. 94 = acuminatus procumbens, *Linn. f. Suppl.* 153.—Afr. austr.
pterocarpus, DC. Prod. ii. 5 = Pterocelastrus tricuspidatus.
pubescens, *Wall. Cat.* n. 4303.—Ind. or.
punctatus, Thunb. Fl. Jap. 97 = articulatus.
punctulatus, *Sieb. & Zucc. in Abh. Akad. Muench.* iv ii. (1846) 150.—Japon.
pyracanthifolius, Salisb. Prod. 141 = pyracantha.
pyracanthus, *Linn. Sp. Pl.* 197.—Afr. austr.
pyrifolia, Herb. Framp. ex Oliver, Fl. Trop. Afr. i. 361 = serratus.
pyrius, Willem. in Ust. Ann. Bot. xviii. (1796) 21 = Gymnosporia trigyna.
quadrangulatus, *Schrad. in Goett. Gel. Ans.* i. (1821) 716.—Hvas.
racemosus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 599.—Japon.
racemosus, Wall. Cat. n. 4320 = Parastemon urophyllum
racemulosus, *Hassk. Hort. Bogor.* i. 155.—Java.
refractus, E. Mey. ex Sond. & Harv, Fl. Cap. i. 467 = Cassine scandens.
repandus, *Blume, Bijdr.* 1145.—Java.
retusus, *Pair. Encyc. Suppl.* ii. 146.—Peruv.
rhamnoides, *Poir* l. c. 145.—Afr. austr.
rhombifolius, *Eckl. & Zeyh. Enum* 120.—Afr. austr.
rhombifolius, Hook & Arn. in Hook. Bot. Misc. iii (1833) 170 = Iodina rhombifolia.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 289.—Ins. Fiji.
rigidus, Eckl. & Zeyh. Enum. 121 = capitatus.
rigidus, *Thunb. Fl. Cap.* ii. 124.—Afr. austr.
rigidus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 396 = Gymnosporia Wallichiana.
robustus, Roxb. Hort. Beng. 18; Fl. Ind. i. 626. Kurrimia pulcherrima.
rostratus, Thunb. Prod. Pl. Cap. 42 = Pterocelastrus rostratus
Rothsanus, Roem. & Schult. Syst. v. 423 = paniculatus.
Rothianus, Wight & Arn. Prod. i. 159 = Gymnosporia Rothiana.
rotundifolius, Thunb. Prod. Pl. Cap. 42 = Duvyalis celastroides.
Royleanus, Wall. Cat. n. 4317 = Gymnosporia Royleana.
ruber, Gerrard, ex Oliver, Fl. Trop. Afr. i. 363 = mossambicensis.
ruber, *Harv. in Harv. & Sond. Fl. Cap.* ii. 592.—Afr. austr.
rufus, Wall. Cat. n. 4309 = Gymnosporia rufa.
rupestris, Eckl. & Zeyh. Enum. 119 = acuminatus.
saxatilis, *Burch. Trav.* ii. 264.—Afr. austr.
scandens, *Linn. Sp. Pl.* 196.—Am. bor.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 132 = serratus.
semiarillatus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 599.—Java.
senegalensis, Lam. Encyc. i. 661 = Gymnosporia montana.
seplarius, *Dennst. Schlussz. Hort. Malab.* 31.—Ind. or.
serratus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 131.—Abyssin.
serrulatus, R. Br. in Salt, Abyss. App. 64 = serratus.
serrulatus, Roth, Nov. Pl. Sp. 155 = Gymnosporia Heyneana.
Sieberi, Bernh. ex Harv. & Sond. Fl. Cap. i. 473 = Ilex capensis.
sinuato-dentatus, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 130 = luteolus.
spathophyllus, Eckl. & Zeyh. Enum. 121 = buxifolius.

CELASTRUS:—
spicatus, Vell. Fl. Flum. 92; ii. t. 188 = Gouania coryliifolia.
spinosus, Royle, Illust. Bot. Himal. 167 = Gymnosporia Royleana.
stenophyllus, Eckl. & Zeyh. Enum. 122 = linearis.
striatus, Thunb. Fl. Jap. 98 = Enonymus Thunbergianus.
styloses, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 401.—Sapindaceæ.
subpicatus, *Hook. Ic. Pl.* t. 482.—Hab.?
Tatarinowii, Regel, Descr. Pl. Pan.-Mandsh. Acad. Pétersb. xv. (1857) 357 = articulatus.
tenuispinus, *Sond. in Harv. & Sond. Fl. Cap.* i. 456.—Afr. austr.
tetragonus, Thunb. Prod. Pl. Cap. 42 = Cassine acuminata.
trucispidatus, Lam. Tabl. Encyc. ii. 93 = Pterocelastrus tricuspidatus.
trigynus, Lam. l. c. 94 = Gymnosporia trigyna.
Tsaad, *Ferr. & Gal. ex Walp. Ann.* iii. 841.—Abyssin.
umbellatus, Vell. Fl. Flum. 92; ii. t. 187 = Reissekia cordifolia.
uncinatus, Ruiz & Pav. Fl. Per. iii. 7 = Maytenus chilensis.
undatus, *Thunb. Prod. Pl. Cap.* 42.—Afr. austr.
undulatus, Lam. Encyc. i. 662 = Pittosporum Senacia.
venenatus, Eckl. & Zeyh. Enum. 121 = buxifolius.
venulosus, *Wall. Cat.* n. 4821.—Reg. Himal.
verrucosus, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 458.—Afr. austr.
verticillatus, Roxb. Hort. Beng. 18; Fl. Ind. i. 624 = Pittosporum floribundum.
verticillatus, Ruiz & Pav. Fl. Per. iii. 6 = Maytenus verticillatus.
Wallichianus, Spreng. Syst. v. Index, 150 = Gymnosporia Wallichiana.
Wallichianus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 400 = Gymnosporia Heyneana.
Wightianus, Wall. Cat. n. 4822 = Pleurostylia Wightii.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 456.—Afr. austr.
zeylanicus, Roth, ex Roem. & Schult. Syst. v. 427 = Scutia Commersonii.

CELEBNIA, Noronha, in Verh Batav. Gen. v. (1790) ed I. Art. iv. 3 (Quid?).

CELMISIA, Cass. in Dict. Sc. Nat. vii. 356 (1817) = **Alotope**, DC. (Compos.).
rotundifolia, Cass. l. c. 357 = A. tabularia.

CELMISIA, Cass. in Dict. Sc. Nat. xxxvii. 259 (1825). *COMPOSITÆ*, Benth. & Hook. f. ii. 278.
asteliæfolia, Hook. f. Fl. Antarct. 35 = longifolia.
bellidioides, *Hook. f. Handb. New Zeal. Fl.* 135.—N. Zel.
cordatifolia, *T. Buch. in Trans. N. Z. Inst.* xi. (1879) 437. t. 18.—N. Zel.
coriacea, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 119.—N. Zel.
Dallii, *T. Buch. in Trans. N. Z. Inst.* xiv. (1882) 355. t. 16.—N. Zel.
densiflora, *Hook. f. Handb. New Zeal. Fl.* 130.—N. Zel.
discolor, *Hook. f. Fl. N. Zel.* i. 102.—N. Zel.
glandulosa, *Hook. f. l. c.* 104.—N. Zel.
gracilenta, *Hook. f. Fl. Antarct.* 35.—N. Zel.
graminifolia, *Hook. f. l. c.*—N. Zel.
Hastii, *Hook. f. Handb. New Zeal. Fl.* 131.—N. Zel. Hectori, *Hook. f. l. c.* 135.—N. Zel.
hieracifolia, *Hook. f. Fl. N. Zel.* 104. t. 34.—N. Zel.
holosericea, *Hook. f. Fl. Antarct.* 36.—N. Zel.
incana, *Hook. f. Fl. N. Zel.* 103. t. 34.—N. Zel.
lanceolata, *Hook. f. l. c.* 101.—N. Zel.
lateralis, *T. Buch. in Trans. N. Z. Inst.* iv. (1872) 226.—N. Zel.
Lindsayi, *Hook. f. Handb. New Zeal. Fl.* 131.—N. Zel.
linearis, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 387.—N. Zel.
longifolia, *Cass. in Dict. Sc. Nat.* xxxvii. 259.—Austral.
Lyallii, *Hook. f. Handb. N. Zeal. Fl.* 132.—N. Zel.
Mackaui, *Raoul, Choix Pl. Nouv. Zél.* 19. t. 14.—N. Zel.
Munroi, *Hook. f. Handb. New Zeal. Fl.* 133.—N. Zel.
petiolata, *Hook. f. l. c.* 135.—N. Zel.
ramulosa, *Hook. f. l. c.* 733.—N. Zel.
rupestris, *Cheesem. in Trans. N. Z. Inst.* xvi. (1884) 409.—N. Zel.
Lechleri, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
sessiliflora, *Hook. f. Handb. New Zeal. Fl.* 135.—N. Zel.
Sinclairi, *Hook. f. l. c.* 132.—N. Zel.
spathulata, A. Cunn. in DC. Prod. v. 209 = Trichocline scapigera.
spectabilis, *Hook. f. Fl. Antarct* 35.—N. Zel.
Traversii, *Hook. f. Handb. New Zeal. Fl.* 134.—N. Zel.
verbascifolia, *Hook. f. Fl. N. Zel.* 101.—N. Zel.
vernicosa, *Hook. f. Fl. Antarct.* 34. t. 26, 27.—Ins Auckland.
viscosa, *Hook. f. Handb. New Zeal Fl.* 133.—N. Zel.
Walkeri, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 549.—N. Zel.

CELOSIA, Linn. Gen. ed. I. 34 (1737). *AMARANTACEÆ*, Benth. & Hook. f. iii. 24.
AMARANTHUS, Adans. Fam. ii. 269 (1763).
GONUFAS, Rafin. Sylva Tellur. 124 (1838)
KOKERA, Adans. Fam. ii. 269 (1763).
LAGREZIA, Moq. in DC. Prod. xiii. ii. 252 (1849).
LESTIBUDESIA, Thou. Hist. Vég. Iles Afr. Sd. i. 16 (1806).
LOPHOKERA, Rafin. Fl. Tellur. iii. 42 (1836).
SUKANA, Adans Fam. ii. 269 (1763).
acroporoides, *Hochst. ex Benth. & Hook. f. Gen.* iii. 25.—Abyss.
acroprosoides, Hochst. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 140 = trigyna.
adonsis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 211 = trigyna.
albida, Willd. Sp. Pl. i. 1197 = Allmania albida.
allmanoides, *Moq. in DC. Prod.* xiii. ii. 243.—Burma.
altissima, Salzm. ex Moq. l. c. 381 = Telanthera ramosissima.
amarantoides, Medic. Malv. 92 = trigyna.
amphibia, Salzm. ex Moq. in DC. Prod. xiii. ii. 363 = Telanthera philoxeroides.
anthelminthica, *Aschers. in Schweinf. Beitr Fl. Aethiop.* 176.—Abyss.
arborescens, Spreng. Syst. i. 815 = Deeringia altissima.
arenaria, Salzm. ex Seub. in Mart. Fl. Bras. v. i. 170 = Telanthera maritima.
argentea, *Linn. Sp. Pl.* 205.—Cosmop. trop.
aspera, Roth, Nov. Pl. Sp. 178 = Allmania nodiflora.
aurea, T. Moore, Flor. Mag. i. (1861) t. 47 = cristata.
axillaris, Mart. Hort. Erlang. 76 = Alternanthera spinosa.
baccata, Retz. Obs. fasc. v. 23 = Deeringia celosioides.
bicolor, Blanco, Fl. Filip. ed. I. 191 = philippica.
Bolvinii, *Hook. f. in Benth. & Hook. f. Gen.* iii. 25.—Madag.
brasiliensis, *Moq. in DC. Prod* xiii. ii. 239.—Bras.
castrensis, Linn. Sp. Pl. ed. II. 297 = cristata.
caudata, *Vahl, Symb. Bot.* i. 21.—Arabia.
cernua, *Andr. Bot. Rep.* x. t. 685.—Ind. or.
cernua, Mairet, ex Moq. in DC. Prod. xiii. ii. 361 = Alternanthera canescens.
cernua, Roxb. Hort. Beng. 19; Fl. Ind. i. 580 = cristata.
coccinea, Linn. Sp. Pl. ed. II. 297 = cristata.
comosa, Retz. Obs. fasc. vi. 26 = cristata.
corymbifera, *Didr. Ind. Sem. Hort. Havn.* (1849) 13.—Hab.?
corymbosa, Retz. Obs. fasc. i. 13 = Polycarpæa corymbosa.
cristata, *Linn. Sp. Pl.* 206.—Cosmop. trop.
cymosa, *Seub. in Mart. Fl. Bras.* v. i. 246.—Bras.
decumbens, Hort. ex. Moq. in DC. Prod. xiii. ii. 410 = Gomphrena decumbens.
dichotoma, Heyne, in Roth, Nov. Pl. Sp. 172 = Allmania nodiflora.
diffusa, Hoffmgg. ex Seub. in Mart. Fl. Bras. v. i. 179 = Telanthera ramosissima.
echinata, Humb. & Bompl. ex Roem. & Schult. Syst. v. 581 = Alternanthera Achyrantha.
elongata, Spreng. in Schrad. Journ. (1800) ii. 196 = Telanthera porrigens.
eriantha, Vahl, ex Moq. in DC. Prod. xiii. ii. 386 = Gomphrena eriantha.
exstipulata, *Hornem. ex Schrank, in Syll. Ratisb.* i. (1824) 202.—Nepal.
floribunda, *A Gray, in Proc. Am Acad.* v. (1861) 167.—Calif.
glauca, Kottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 198 = patula.
glauca, Wendl. Beobacht. 43; Hort. Herrenh. 4. t. 2 = Hermbstaedtia glauca.
gnaphaloides, Linn. f. Suppl. 161 = Gomphrena gnaphaloides.
grandifolia, *Moq. in DC. Prod.* xiii. ii. 238.—Bras.
humifusa, Willd. ex Roem. & Schult. Syst. v. 581 = Telanthera caracasana.
Huttoni, *Mast. in Gard. Chron.* (1872) 315.—Java.
intermedia, Hochst. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 140 = trigyna.
interrupta, Seub. in Mart. Fl. Bras. v. i. 165 = Froelichia interrupta.
japonica, *Mart. Hort. Belang.* 70.—Japon.?
lactea, Hort. ex Moq. in DC. Prod. xiii. ii. 300 = Aerva sanguinolenta.
lanata, Blanco, Fl. Filip. ed. II. 184 = Aerva lanata.
lanata, Linn Sp. Pl. 205 = Aerva tomentosa.
lanata, Poir. ex Steud. Nom. ed. II. i. 315 = Gomphrena gnaphaloides.
lappacea, Medic Beobacht. (1783) 147 = Pupalia atropurpurea.
latifolia, *Steud. Nom* ed. II. i. 315.—Java.
laxa, *Schur & Thonn. Beskr. Guin. Pf.* 141.—Afr. trop.
leptostachya, *Benth. in Hook. Niger Fl.* 493.—Afr. trop.
linearis, Sweet, ex Hook f. Fl. Brit. Ind. iv. 714 = argentea.
longifolia, *Mart. Nov. Gen et Sp.* ii. 63. t. 157, 158. f. 2 = linea.
madagascariensis, *Poir. Encyc.* v. 39.—Madag.
major, *Grisb. in Goett. Abh.* xix. (1874) 79.—Reg. Argent.
margaritacea, Linn. Sp. Pl. ed. II. 297 = argentea.
maylandica, R. Br. Obs. fasc iii 27 = argentea
maritima, Salzm. ex Moq. in DC. Prod. xiii. ii. 341 = Philoxerus vermicularis.
melanocarpos, Poir Encyc Suppl. iv. 318 = trigyna.
missionis, Wall. Cat. n. 6915 = Banalia thyrsiflora.
Monsonia, Retz. Obs fasc. ii. 18 = Aerva Monsonia.
Moquini, *Guill. ex DC Prod.* xiii. ii. 239.—Mexic.
mucronata, Herb. ex Moq. in DC. l. c. 248 = Allmania albida.
muricata, Spreng. Syst. iv. Cur. Post. 102 = Cladostachys muricata.
nana, Blanco, Fl. Filip. ed. I. 192 = Ammania baccifera.
nitida, Vahl, Symb. Bot. iii. 44 = paniculata.
nodiflora, Linn. Sp. Pl. 205 = Allmania nodiflora.
nodiflora, Salzm. ex Moq. in DC. Prod. xiii. ii. 364 = Telanthera polygonoides.
nodiflora, Wall. Cat. n. 6890 = Allmania albida.
nudiflora, Pav. ex Moq. in DC. Prod. xiii. ii. 341 = Philoxerus vermicularis.
odorata, *Burch. ex Moq. l. c.* 244.—Afr. austr.
pallida, Hort. Vindob. ex Moq. l. c. 248 = Allmania albida.
pallida, Salisb. Prod. 145 = argentea.
Palmeri, *S Wats. in Proc. Am. Acad.* xviii. (1882-83) 148.—Calif.
paniculata, *Linn. Sp. Pl.* 206.—Florida; Ind. occ.
parviflora, Vahl, ex Moq. in DC. Prod. xiii. ii. 346 = Iresine gracilis.
patula, *Willd. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 198.—Ind. or.
peruviana, Van Spand. ex Moq. in DC. Prod. xiii. ii. 365 = Telanthera frutescens.
peruviana, Zuccagni, in Roem. Collect. 133 = Telanthera porrigens.
philippica, *Steud. Nom.* ed. II. i. 316.—Ins. Philipp.
phytolaccæfolia, Juss. ex Moq. in DC. Prod. xiii. ii. 238 = madagascariensis.
piloselloides, Poir. ex Moq. l. c. 337 = Guillemina lanuginosa.
polygonoides, *Retz. Obs. fasc.* ii. 12.—Ind. or.
polygonoides, Wall. Cat. n. 6913 = pulchella.
polysperma, Roxb. Hort. Beng. 84; Fl. Ind. i. 581 = Deeringia polysperma.
populifolia, *Moq. in DC. Prod.* xiii. ii. 239.—Abyss.
procumbens, Hort. ex Moq. l. c. 410 = Gomphrena decumbens.

CELOSIA :—
procumbens, Jacq. Misc. ii. 344 = Froelichia interrupta.
prostrata, Hort. ex Moq. in DC. Prod. xiii. ii. 410 = Gomphrena decumbens.
pulchella, *Moq. l. c.* 238.—Ind. or.
purpurea, St. Hil. ex Steud. Nom. ed. II. i. 315, 316 = cristata.
pyramidalis, Burm. f. Fl. Ind. 62 = cristata, Allmania albida.
recurva, Burch. Trav. ii. 226 = Hermbstaedtia elegans.
sanguinea, *Hoffmgg. Verz. Pfl.* ii. 85.—Ind. or.
sinuata, *Mart. ex Steud. Nom.* ed. I. 171.—Hab.?
sparsa, Forsyth. ex Moq. in DC. Prod. xiii. ii. 250 = Chamissoa altissima.
spathulata, *Boj. Hort. Maurit.* 267.—Madag.
spathulata, Saint-Elme, ex Moq. in DC. Prod xiii. ii. 253 = madagascariensis.
spicata, Andié, ex Moq. l. c. 143 = Anthochlamys polygaloides.
spicata, *Spreng. Syst.* i. 815.—Madag.
spinescens, Kuns. ex Wall. Cat. sub n. 6982 = Achyranthes aquatica.
splendens, *Schum. & Thonn. Beskr. Guin. Pl.* 140.—Afr. trop.
stolonifera, Leprieur, ex Moq. in DC. Prod. xiii. ii. 365 = Telanthera maritima.
stricta, Fisch. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 65 = brasiliensis.
stricta, *Hornem. Hort. Hafn. Suppl.* 28.—Bras.
tetragyna, Roxb. ex Moq. in DC. Prod. xiii. ii. 287 = Decringia tetragyna.
texana, *Scheele, in Linnaea*, xxii. (1849) 148.—Texas.
thyrsiflora, Wall. Cat. n. 6914 = Banalia thyrsiflora.
tomentosa, Willd. in Roem. & Schult. Syst. v. 581 = Chamissoa macrocarpa.
trigyna, *Linn. Mant.* 212.—Afr. trop.
trigyna, Wight, ex Steud. Nom. ed. II. i. 316 = Banalia thyrsiflora.
trigyna, Willd. ex Wall. Cat. n. 6915 = polygonoides.
trilobata, *E. Mey. ex Meisn. in Hook. Lond. Journ. Bot.* ii. (1843) 448.—Afr. austr.
tumida, *Hort. Hall. ex Moq. in DC. Prod.* xiii. ii. 245. —Hab.?
virgata, *Jacq. Coll.* ii. 279.—Am. trop.

CELSIA, Vell. Fl. Flum. 178 (1825); iv. t. 93. *ZYGOPHYLLEAE.*
frutescens, *Vell. ll. cc.* 174.—Bras.

CELSIA, Heist. Syst. 5, 20 (1748) = **Ornithogalum**, Tourn. (Liliac.).

CELSIA, Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 926.
DITOXIA, Rafin. ex Ell. Journ. Bot. iv. (1814) 270.
JANTHE, Griseb. Spicil. Fl. Rumel. ii. 40 (1844).
NEPTEA, Spach, Hist Vég. Phan. ix. 268 (1846).
THAPSANDRA, Griseb. Spicil. Fl. Rumel. ii. 40 (1844).
acaulis, *Bory & Chaub. Expéd. Mor. Bot.* 177. t. 18.—Graecia.
affinis, *A. Rich. Tent. Fl. Abyss.* ii. 110.—Abyssin.
agrimonifolia, C. Koch, in Linnaea, xxii. (1849) 732 = heterophylla.
alpina, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 150.—Syria.
arbuscula, *A. Rich. Tent. Fl. Abyss.* ii. 110.—Abyssin.
arctioria, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 192 = C. Arcturus.
Arcturus, *Jacq. Hort. Vindob.* ii. 58. t. 117.—Creta.
armeniaca, Boiss. Fl. Orient. iv. 861 = aurea.
Aucheri, *Boiss. Diagn.* Ser. I. xii. 29.—Persia.
aurea, *C. Koch, in Linnaea*, xxii. (1849) 731.—Armenia.
australis, Hort. Bonn. ex Flora, xvi. (1833) 125 = cretica.
Bernadesii, *G. Don, Gen. Syst.* iv. 499.—Hispan.; Marocco.
betonicaefolia, *Desf. Fl. Atlant.* ii. 58.—Afr. bor.
Boissieri, *Heldr. & Sart. ex Boiss. Fl. Orient.* iv. 858. —Graecia.
brachypepala, *Fisch. & Trautv. in Ind. Sem. Hort. Petrop.* iv. 33.—As. Min.
bugulifolia, *Yanb. & Spach, Illustr.* v. 7. t. 407.—Rumelia; As. Min.
caduca, Moench, Meth. 447 = orientalis.
CELSIA :—
Cavanillesii, Kunze, in Flora, xxix. (1846) 698 = cretica.
chrysaea, Fenzl, ex Boiss. Fl. Orient. iv. 859 = heterophylla.
cilicica, *Boiss. & Heldr. Diagn.* Ser. I. xii. 30.—Cilicia.
collina, Benth. in DC. Prod. x. 247 = Sowerowiana.
coromandelina, *Vahl, Symb. Bot.* iii. 79.—Ind. or.; Afghanist.
coronopifolia, *Boiss. & Bal. Diagn.* Ser. II. vi. 126.—Phrygia
cretica, *Linn. f. Suppl.* 281.—Reg. Mediterr.
cyllenea, *Boiss. & Heldr. Diagn.* Ser. II. iii. 151.—Graecia.
Duenzeri, *Fauché & Chaub. in Expld. Mor. Bot.* 342.—Graecia.
densiflora, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 208.—Afr. trop.
dentata, Moench, Meth. Suppl. 156 = coromandelina.
Doniana, *Walp. Rep.* iii. 150.—Hab.?
elata, *Fisch. ex Steud. Nom.* ed. I. 172.—Hab.?
erecta, *Fisch. Hort. Gorenk.' ex Steud. Nom.* ed. II. i. 316.—Hab.?
floccosa, *Benth. in DC. Prod.* x. 245.—Abyssin.
glabrata, *Roth, ex Steud. Nom.* ed. II. i. 316.—Hab.?
glandulosa, *Bouché, in Verh. Ges. Naturf. Fr. Berl.* i. (1829) 395.—As. Min.
grandiflora, *Hort. ex Steud. Nom.* ed. II. i. 316.—Asia.
heterophylla, *Desf. ex Pers. Syn.* ii. 161.—Persia; As. Min.; Ind. or.
horizontalis, Moench, Meth. 448 = C. Arcturus.
incana, Benth. in DC. Prod. x. 247 = lanceolata.
incarnata x, Reichb. ex Heynh. Nom. ii. 139, nonen.
interrupta, *Fresen. in Flora*, xvi. (1833) 805.—Abyssin.
intricata, *Benth. in DC. Prod.* x. 247.—Persia.
lacinata, Cass. ex Bail, in Journ Linn. Soc. xvi. (1878) 585 = cretica.
laciniata, Poir. Encyc. Suppl. ii. 147 = Bernadesii.
lanata, Spreng. Syst. ii. 810, err. typ. = sublanata.
lanceolata, *Vent. Jard. Cels*, t. 27.—Mesopot.; Persia.
leptarrat, Schott & Kotschy, in Oestr. Bot. Wochenbl. (1854) 186 = brachypepala.
linearis, Jacq. Coll. ii. 270 = Alonsoa linearis.
Luciliae, *Boiss. Diagn.* Ser. I. iv. 65.—As. Min.
lycia, *Boiss. Fl. Orient.* iv. 857.—Lycia.
lyrata, G. Don, Gen. Syst. iv. 499 = cretica.
maroccana, *Bail, in Journ. Bot.* xiii. (1875) 172.—orientalis, *Linn. Sp. Pl.* 621.—Graecia; As. Min.; Persia.
parviflora, *Decne. in Ann. Sc. Nat.* Sér. II. v. (1836) 354.—Arabia.
parviflora, G. Don, Gen. Syst. iv. 499 = Doniana.
pedunculosa, *Hochst. & Steud. ex Benth. in DC. Prod.* x. 245.—Abyssin.
persica, *C. A. Mey. Verz. Pfl. Casc.* 111.—Reg. Casc.; Persia.
pinetorum, *Boiss. Diagn.* Ser. I. xii. 30.—Cilicia.
pinnatifida, Bolss. & Reut. Diagn. Pl. Nov. Hisp. 22 = Bernadesii.
pontica, *Boiss. in Tchihat. Asie Min. Bot.* ii. 11.—As. Min.
purpurea, *Yanka, in Termész. Füzetek*, iv. (1880) 296.—Hungar.
pyrolliformis, *Boiss. & Heldr. Diagn.* Ser. I. xii. 28.—Lycaonia.
ramosissima, *Benth. in DC. Prod.* x. 244.—Marocco.
scrophulariaefolia, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 113.—Abyssin.
sinuata, Cav. in Anal. Cienc. Nat. iii. (1801) 68 = cretica.
sinuata, *Colla, Hort. Ripul. App.* ii. 344; iii. 30.—Graecia.
speciosa, *Fenzl, in Friedrich. Reise*, 280.—Graecia.
striatiflora, *Hort. ex Steud. Nom.* ed. II. i. 316.—Hab.?
sublanata, Jacq. Fragm. 79. t. 196 = C. Arcturus.
Suwarowiana, *C. Koch, in Linnaea*, xvii. (1844) 384. —Armenia; Persia.
tomentosa, *Hochst. ex Benth. in DC. Prod.* x. 245.—Abyssin.
tomentosa, Zucc. Pl. Nov. Fasc. ii. 22 = Verbascum graecum.
trapaefolia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 88.—Lycia.
CELSIA :—
urticaefolia, Sims, Bot. Mag. t. 417 = Alonsoa incisifolia.
valerianaefolia, *A. Rich. Tent. Fl. Abyss.* ii. 112.—Abyssin.
viscosa, Roth, Catalect. ii. 69 = coromandelina.
viscosa, Wight, ex Benth. in DC. Prod. x. 229 = Verbascum virgatum.

CELTIS, Tourn. ex Linn. Gen. ed. I. 337 (1737) *URTICACEAE*, Benth. & Hook. f. iii. 354.
MERTENSIA, H. B. & K. Nov. Gen. et Sp. ii. 30 (1817).
MOMMIA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 122 (1819).
SAUROBROMA, Rafin. Sylva Tellur. 32 (1838).
SOLENOSTIGMA, Endl. Prod. Fl. Norf. 41 (1839).
Acosta, Buch.-Ham. in Trans. Linn. Soc. xvii. (1837) 211 = australis.
aculeata, *Sw. Prod. Veg. Ind. Occ.* 58.—Ind. occ.; Am. trop.
africana, *Burm. f. Fl. Cap. Prod.* 31.—Afr. austr.
alba, *Rafin. Fl. Ludov.* 25.—Am. bor.
albicans, Willd. ex Steud. Nom. ed. II. i. 316 = Trema micrantha.
alnifolia, *Mig. in Mart. Fl. Bras.* iv. t. 176.—Am. trop
amblyophylla, *F. Muell. Fragm.* ix. 76.—Austral.
amboinensis, Buch.-Ham. ex Wall. Cat. sub n. 8698 = Trema politoria.
amboinensis, Willd. Sp. Pl. iv. 997 = Trema amboinensis.
americana, Hort. ex Planch. in Prod. xvii. 176 = mississipiensis.
americana, *Mill. Gard. Dict.* ed. VIII. n. 4.—Ind. occ.
amplifolia, *Voronkin, in Verh. Batav. Gen.* v. (1790) ed. I. *Art.* iv. 11.—Malaya.
anfractuosa, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 238.—Mexic.
angustifolia, Lindl. in Wall. Cat. n. 8691 = Trema angustifolia.
angustifolia, *Planch. in DC. Prod.* xvii. 186.—Sumatra
appendiculata, E. Mey. ex Planch. in Ann. Sc. Nat Sér. III. x. (1848) 341 = Chaetachme aristata.
arcata, Buch.-Ham. ex Wall. Cat. sub n. 8695 = australis.
aristata, E. Mey. ex Planch. in Ann. Sc. Nat Sér. III x. (1848) 341 = Chaetachme aristata.
aspera, Brongn. in Duperr. Bot. Voy. Coq. 213. t. 48 = Trema aspera.
aspera, Lodd. ex G. Don, in Loud. Hort. Brit. 418 = australis.
asperula, *Mig. in Mart. Fl. Bras* iv. t. 176.—Am. trop
Audibertiana, Spach, in Ann. Sc. Nat. Sér. II. xvi. (1841) 41 = occidentalis.
australis, *Linn. Sp. Pl.* 1043.—Europ.; As. temp.; Ind. or.
australis, A. Rich. Tent. Fl. Abyss. 257 = Kraussiana.
azcurrensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 193.—Reg. Argent.
Balansae, *Planch. in DC. Prod.* xvii. 188.—N. Caled.
Berlandieri, *Klotzsch, in Linnaea*, xx. (1847) 641.—Mexic.
bifera, Ruiz, ex Miq. in Mart. Fl. Bras. iv. t. 182 = dichotoma.
boliviensis, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 311.—Bolivia.
Bonplandiana, *Planch. in DC. Prod.* xvii 190.—Am. austr.
brasiliensis, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 310.—Bras.
brevifolia, *Mig. in Mart. Fl. Bras.* iv. t. 190.—Am. trop.
brevinervis, Blume, Mus. Bot. Lugd. Bat. ii. 67 = paniculata.
brevipes, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 297. —Arizona.
Bungeana, *Blume, Mus. Bot. Lugd. Bat.* ii. 71.—China.
Burmanniі, Planch. in Ann. Sc. Nat. Ser. III. x. (1848) 296 = Kraussiana.
causcorea, H. B. & K. Nov. Gen. et Sp. ii. 28 = Trema micrantha.
canina, *Rafin. in Am. Monthly Mag.* ii. (1817) 48.—Am. bor.
caucasica, Hohen. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 285 = australis.
caucasica, Willd. Sp. Pl. iv. 994 = australis.
caudata, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 241 = Prunus pogonostyla.

CELTIS :—
caudata, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 204.—Mexic
caudata. Wall. ex Planch. in DC. Prod. xvii. 199 = Trema amboinensis.
Chichiltan, Ruiz & Pav. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 335 = Trema orientalis.
Chicopé, Miq. in Mart. Fl. Bras. iv. i. 181 = C. Tala.
chinensis, Hunge, Enum. Pl. Chin. Bor. 61 = Bungeana.
cinerea, Torr. Bot. Mex. Bound. t. 50 = C. Tala.
cinnamomea, *Lindl. in Wall. Cat. n.* 3696.—Ind. or.; Malaya.
Claussentana, *Miq. in Mart. Fl. Bras.* iv. i. 178.—Am. trop.
Commersonii, A. Brongn. in Duperr. Voy. Coq. Bot. 215 = Trema orientalis.
confcrta, *Planch. in DC. Prod.* xvii. 188.—Madag.
cordata, Pers. Syn. i. 292 = occidentalis.
cordifolia, L'Hérit. ex Duham. Arb. ed. Nov. ii. 37 = occidentalis.
crassifolia, Lam. Encyc. iv. 138 = occidentalis.
crenata, Miq. in Mart. Fl. Bras. iv. i. 181 = C. Tala.
crenata, *Planch. in DC. Prod.* xvii. 191.—Bolivia.
Curtondiubo, Gomez, ex Planch. l. c. 204 = Trema micrantha.
Davidiana, *Carr. in Rev. Hortic.* (1868) 800.—China.
dichotoma, *Ruiz, ex Miq in Mart. Fl. Bras.* iv. i. 182.—Peruv.
diffusa, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 314.—Bras
discolor, A. Brongn. in Duperr. Voy. Coq Bot. i. 215. t. 47. f. B = Trema discolor.
disticha, *Parodi, in Anal. Soc Cient. Argent.* v. (1878) 96 (*Contrib.* 44).—Paraguay.
Djungiel, *Planch. in DC. Prod.* xvii 185.— Java.
Douglasii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 293 = reticulata.
dumosa, Casar. ex Planch. in DC. Prod. xvii. 188 = ferruginea.
dysodoxylon, Thw. Enum. Pl. Zeyl. 267 = cinnamomea.
Ehrenbergiana, Liebm. in Dansk. Videnak. Selsk. Skrift. V. ii. (1851) 339 = aculeata.
elongata, Wall. Cat. n. 3692 C = Villebrunea integrifolia.
epiphylladena, Orteg. Hort. Matr. 79 = aculeata.
crianthа, *E. Mey. ex Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 296.—Afr. austr.
eriocarpa, Decne. in Jacquem. Voy. Bot. 150 = australis.
excelsa, Salisb. Prod. 175 = australis.
ferruginea, *Planch. in DC. Prod.* xvii. 188.—Bras.
ferruginea, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 434 = Ziziphus asperata.
flagellaris, Casar. ex Planch. in DC. Prod. xvii. 189 = brasiliensis.
flexuosa, *Miq. in Mart. Fl. Bras.* iv. i. (1852–62) 181. —Am. trop.
flexuosa, Planch. in DC. Prod. xvii. 192 = bolviensis?
floridana, Rafin. New Fl. Am. iii. 37 = occidentalis?
fuscata, Rafin. l. c. 38 = mississipiensis?
Gardneri, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 311.—Bras.
glabra, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. 1. *Art.* iv. 12.—Malaya.
glabra Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 298 = tetrandra.
glabrata, Spreng. Syst. v. 150 = aculeata.
glabrata, Stev. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 285 = australis.
glaucrata, Hochst. in Flora, xxviii. (1845) 87 (= *Sponia guineensis*).—Afr. trop.
glycycarpa, *Mart. ex Miq. in Mart. Fl. Bras.* iv. i. 174.—Am. trop.
Goudotii, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 312.—N. Granat.
grandidentata, Tenore, Ind. Sem. Hort. Neap. (1838) 15 = mississipiensis?
guineensis, *Schum. & Thonn. Beskr. Guin. Pl.* 160.—Afr. trop.
hamata, *Blume, Mus. Bot. Lugd. Bat.* ii. 72.—Malaya.
Hamiltonii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 301 = tetrandra.
Harperi, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 271.—Ins. Fiji.
Hasseltii, *Planch. in DC. Prod.* xvii. 185.—Java.
hebrophylla, Rafin. New Fl. Am. iii. 37 = occidentalis?
Hillebrana, *Planch. in DC. Prod.* xvii. 189.—Bras.
hypoleuca, *Planch. l. c.* 188.—N. Caled.
ingens, F. Muell. Fragm. iv. 66 = paniculata.

CELTIS :—
integrifolia, Lam. Encyc. iv. 140 = mississipiensis.
jamaicensis, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 290.—Jamaica
japonica, *Planch in DC. Prod.* xvii. 172.—Japon.
Kotschyana, Strv. in Bull. Soc. Nat. Mosc. xxx. (1857) i. 392 = australis.
Kraussiana, *Bernh. in Flora*, xxviii. (1845) 87.—Afr. trop. et austr.
laevigata, Salisb. Prod. 175 = orientalis.
laevigata, Spreng. Syst. i 932 = aculeata.
laevigata, Willd. Enum. Hort. Berol. Suppl. 67 = mississipiensis.
Lamarchiana, Roem. & Schult. Syst. vi. 311 = Trema Lamarckiana.
lanceolata, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 150 (*Contrib.* 45).—Paraguay.
lancifolia, *Miq in Mart. Fl. Bras.* iv. i. 175.—Bras.
lanceifolia, *Planch. in DC. Prod.* xvii. 192.—Bras.
latifolia, *Planch. l. c.* 186.—N. Guinea.
laurifolia, *Planch. l. c.* 185.—Java.
Lima, Blanco, Fl. Filip. ed. I. 197 = Trema Blanconi.
Lima, Lam. Encyc. iv. 40 = Trema Lamarckiana.
Limia, Ort. Prod. Veg. Ind. Occ. 53 = Trema Limia.
Lindheimeri, *Engelm. ex C. Koch, Dendrol.* ii. ii. 434. —Am. bor.
littoralis, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 337 = caudata.
longifolia, Rahn. Atl. Journ. 177 = mississipiensis?
lutea, Pers. Syn. i. 292 = australis.
macrophylla, H. B & K. Nov. Gen. et Sp. ii. 24 = Trema macrophylla.
madagascariensis, Boj. Hort. Mauril. 296 = Trema orientalis.
maritima, *Rafin. in Am. Monthly Mag* ii. (1817) 44. —Am. bor.
mauritiana, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 307.—Ins. Mauric.
membranacea, *Miq. in Mart. Fl. Bras.* iv. i. 175.—Bras.
micrantha, Sw. Prod. Veg. Ind. Occ. 58 = Trema micrantha.
microcarpa, Salzm. ex Planch in Ann Sc. Nat. Sér III. x. (1848) 333 = Trema micrantha.
microphylla, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 59 = Trema timorensis.
mississippiensis, *Bosc, Encyc. Agric.* vii. 577.—Am. bor.
mollis, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 996 = Trema micrantha.
mollis, Wall Cat. n. 7208 = tetrandra.
monoica, *Hensl. Bial. Centr. Am. Bot.* iii. 189.— Mexic.
montana, Jungh. ex Planch. in DC. Prod. xvii 198 = Pancheria parviflora.
morifolia, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 311.—Bras.
morifolia, Rafin. New Fl. Am. iii. 36 = occidentalis.
Müll, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 223.—Japon.
napaiensis, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 298 = tetrandra.
obliqua, Moench, Meth. 344 = occidentalis.
occidentalis, *Linn. Sp. Pl.* 1044.—Am. bor.
opaca, C. Moore, ex Benth. Fl. Australiens. vi. 157 = paniculata.
Opegrapha, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 294 = Kraussiana.
orientalis, Hort. ex Steud. Nom. ed. II. i. 316 = occidentalis.
orientalis, Lam. Encyc. iv. 138 = Trema Commersonii.
orientalis, Linn. Sp. Pl. 1044 = Trema orientalis.
orientalis, Mill. Gard. Dict. ed. VIII. n. 3 = australis.
orientalis, Thunb. Fl. Jap. 114 = japonica.
orthacantha, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 309 = aculeata.
pacifica, *Planch. l. c.* 308.—Ins. Marquesas.
pallida, Torr. in Bot. Mex. Bound. 208 = C. Tala.
paniculata, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 305.—Austral.: Ins. Pacific.
parvifolia, Rafin. New Fl. Am. iii. 36 = occidentalis.
parvifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 218.—Cuba.
patula, Rafin. New Fl. Am. iii. 37 = reticulata?
Pavonii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 313 = dichotoma.
philippinensis, *Blanco, Fl. Filip.* ed. I. 197.—Ins. Philipp.; Austral.

CELTIS :—
politoria, Wall. Cat. n. 3693 = Trema politoria.
procera, Salisb. Prod. 175 = occidentalis.
pubescens, *H. B. & K. Nov. Gen. et Sp.* ii 32.—N. Granat.
pumila, Pursh, Fl. Am. Sept. i. 200 = occidentalis.
reticulata, Hook. f. & Thoms. ex Planch. in DC. Prod. xvii. 209 = Gironniera reticulata.
reticulata, *Torr. in Ann. Lyc. N. York*, i. (1824) 247.—Am. bor. occ.
reticulosa, Miq. Fl. Jungh. 69 = cinnamomea.
rhamnoides, DC. ex Planch. in DC. Prod. xvii. 189 = brasiliensis.
rhamnoides, Willd. Sp. Pl. iv. 998 = aculeata.
rigescens, *Planch. in DC. Prod.* xvii. 182.—Sumatra.
rigida, Blume, Bijdr. 486 = Trema rigida.
rigida, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 150 (*Contrib.* 45).—Paraguay.
riparia, H. B. & K. Nov. Gen. et Sp. ii. 28 = Trema riparia.
riparia, Liebm. in Videnak. Selsk. Skr. V. ii. (1851) 337 = caudata.
rostrata, Zipp. ex Span. in Linnaea, xv. (1841) 343 (= *Sponia rostrata*).—Hab.?
Roxburghii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 302 = tetrandra.
rufescens, Banks, ex Planch. l. c. 334 = Trema micrantha.
rugosa, Willd. Sp. Pl iv. 996 = Trema rugosa.
salicifolia, Rafin. New Fl. Am. iii. 34 = mississipiensis?
scabra, Casar. ex Planch. in DC. Prod. xvii. 204 = Trema carinata
Schiedeana, Schlecht. in Linnaea, vii. (1832) 140 = Trema micrantha
Selloviana, *Miq. in Mart. Fl. Bras* iv. i. 170.—Bras.
serotina, Planch. in Ann. Sc Nat Sér. III. x. (1848) 301 = tetrandra.
Shortii, *Planch. l. c.* 292.—Am. bor.
sinensis, Planch. l. c 286 = Bungeana.
sinensis, *Pers* Syn i 292.—China.
sinensis, Willd. Enum. Hort. Berol. Suppl. 67 = japonica.
spinosa, Ruiz, ex Miq. in Mart. Fl. Bras. iv. i. 182 = dichotoma.
spinosa, *Spreng. Syst.* i. 932 (sp. dub.).—Bras.
spinosissima, *Miq. in Mart. Fl. Bras* iv. i. 176.—Bras
stenophloeodes, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 316.—Bras.
subdentata, E. Mey. ex Planch. l. c. 341 = Chaetechme arnstata.
sumatrana, *Planch. in DC. Prod.* xvii. 181.—Sumatra.
Swartzii, *Planch in Ann. Sc. Nat.* Sér. III. x. (1848) 304.—Ins. S. Doming.
Tala, *Gill. ex Planch. l. c.* 310.—Am. austr.
tenuifolia, Nutt. Gen. Am. i. 202 = pumila?
tenuifolia, Rafin. New Fl. Am. iii. 36 = occidentalis?
tetrandra, *Roxb. Hort. Beng.* 21; *Fl. Ind.* ii. 68.—Ind. or.; Malaya.
tetrandra, Wall. Cat. n. 3695 F = Villebrunea integrifolia.
texana, Scheele, in Linnaea, xxii. (1849) 146 = Berlandieri.
timorensis, *Span. l. c.* xv. (1841) 343.—Ins. Timor.
tomentosa, Roxb. Hort. Beng. 86; Fl. Ind. ii. 66 = Trema amboinensis.
Tournefortii, Lam. Encyc. iv. 138 = australis.
triflora, *Ruiz, ex Miq. in Mart. Fl. Bras.* iv. i. 181.—Peruv.
trinervia, Lam. Encyc. iv. 140 (= *Sponia trinervia*).—Ins. S Doming.
trinervia, Roxb. Hort. Beng. 86; Fl. Ind. ii. 65 = tetrandra.
urticifolia, Rafin. New Fl. Am. iii. 33 = occidentalis.
velutina, *Planch. in Ann. Sc. Nat. Sér.* III. x. (1848) 313 —Peruv.
vesiculosa, Hochst. ex Planch. l. c. 295 = Kraussiana.
virgata, Roxb. in Wall. Cat. n. 3694 = Trema timorensis.
Wenzeli, Blume, Mus. Bot. Lugd. Bat. ii. 71 = cinnamomea.
Wallichii, Steud. Nom. ed. II. i. 316 = tetrandra.
Wightii, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 307.—Ind. or.; Malaya; Austral.
Willdenowiana, Roem. & Schult. Syst. vi. 306 = japonica.
Zippelii, *Planch. in DC. Prod.* xvii. 185.—N. Guinea.
zizyphoides, *Spreng. Syst.* i. 932.—N. Granat.

CEMBRA, Opiz, Seznam, 27 (1852) = **Pinus**, Tourn. (Conif.).
montana, Opiz, l. c. = P. Cembra.

CENARIUM, Linn. Amoen. Acad. iv. 121, in nota (1759) = **Canarium**, Rumph. (Burser.).

CENARRHENES, Labill. Nov. Holl. Pl. i. 36. t. 50 (1804). *PROTEACEAE*, Benth. & Hook. f. iii. 176.
nitida, *Labill. l. c.*—Austral.
paniculata, *Brongn. & Gris*, *in Bull. Soc. Bot. Fr.* xii. (1865) 41.—N. Caled.
spathulaefolia, *Brongn. & Gris*, *l. c.*—N. Caled.
spathulata, Panch. & Sebert, Not. Bois Nouv. Caléd. 180 = spathulaefolia.

CENCHRUS, Linn. Coroll. Gen. 20 (1787). *GRAMINEAE*, Benth. & Hook. f. iii. 1105.
ECHISACHYS, Neck. Elem. iii. 228 (1790).
RARAM, Adans. Fam. ii. 35 (1763).
aegyptius, Beauv. Agrost. 157 = Pleurane aegyptiaca.
agrimonoides, *Trin Diss. Gram Panic.* 72.—Ins. Societ.
alopecuroides, *Y. & C. Presl*, *Rel. Haenk.* i. 317.—Am. austr.
alopecuroides, Thunb. Prod. Pl. Cap. 24 = Pennisetum Linnaei.
anceps, *Anderss. in Peters*, *Reise Mossamb. Bot* 558.—Afr. trop.
anomoplexis, Labill. Sert. Austr. Caled. 14. t. 19 = calycinatus.
asperifolius, Desf. Fl. Atlant. ii. 388 = Pennisetum asperifolium.
australis, *R. Br. Prod.* i. 196.—Austral.
barbatus, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 48.—Afr. trop.
biflorus, *Roxb. Hort. Beng.* [81]; *Fl. Ind.* i. 288.—Ind. or.
brevisetus, *Fourn. Mex. Pl. Enum. Gram.* 50; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 506.—Mexic.
Brownei, Roem. & Schult. Syst. ii. 278 = Isiflexus
bulbifer, Hochst. ex Boiss. Fl. Orient. v. 448 = montanus.
bulbosus, Presen. ex Steud Nom. ed. II i. 317 = pennisetiformis.
calyculatus, *Cav. Ic.* v. 39. t. 463.—Polynes.
capitatus, Linn. Sp. Pl. 1049 = Echinaria capitata
cordistanicus, Wall. Pl. Cavol. 79 = tribuloides.
catharticus, *Delile*, *Cat. Hort. Monsp.* (1826).—*Cf. Linnaea*, xiii. (1839) *Litt.* 108.—Ax. et Afr. trop.
Cavanillesii, *Tausch*, *in Flora*, xx. (1837) 97.—Bras.
ciliaris, *Fig. & De Not. in Mem. Acc. Torin.* (1854) 383.—Nubia!
ciliaris, Linn Mant. 302 = Pennisetum cenchroides.
dactyloides, Steud. Syn. Pl. Gram. 100 = viridis.
diffusus, Ehrenb. ex Boiss. Fl. Orient. v. 449 = Pennisetum cenchroides.
distichophyllus, *Griseb. Cat. Pl. Cub.* 234.—Cuba.
echinatus, Cav. Ic. v. 30. t. 462 = Cavanillesii.
echinatus, *Linn. Sp. Pl.* 1050.—Cosmop. trop.
echinatus, A. Rich. Tent Fl. Abyss. ii. 389 = catharticus.
echinatus, Steud. ex Doell, in Mart. Fl. Bras. ii. 11. 309 = viridis.
echinatus, Torr. Fl. N. Am i. 68 = tribuloides.
echinoides, Wight, ex Steud. Nom. ed. II. i. 317 = Pennisetum cenchroides.
Elliottii, *Kunth*, *Rev. Gram.* i. 51.—Am. bor.
elymoides, *F. Muell. Fragm.* viii. 107.—Austral.
frutescens, *Linn. Sp. Pl.* 1050.—[1] Am.
frutescens, *Sieber*, *ex Steud. Nom.* ed. II. i. 317, nomen. —Armenia.
fusiformis, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 170 = agrimonoides.
gemiculatus, Thunb. Prod. Pl. Cap. 24 = Pennisetum Thunbergii.
gracilis, *Beauv. Agrost.* 157, nomen.—Hab.!
granularis, *Anderss. Enum. Pl. Ins. Galap.* 140.—Ins. Galapag.
granularis, Linn. Mant. ii. App. 575 = Manisuris granularis.
heterochaetus, *Steud. Syn. Pl. Gram.* 110.—Hab.!
hexaflorus, Blanco, Fl. Filip. ed. I. 86 = Pennisetum nigricans.
hirsutus, Spreng. Neue Entdeck. iii. 16 = Scleria hirtella.
horderiformis, Rottl. ex Steud. Syn. Pl. Gram. 102 = Pennisetum Alopecuros.

CENCHRUS:—
horderiformis, Thunb. Prod. Pl. Cap. 24 = Pennisetum compressum.
Hystrix, *Fig. & De Not. in Mem. Acc. Torin.* (1854) 382.—Nubia.
incertus, *M. A. Curt. in Bost. Journ. Nat. Hist.* i. (1837) 135.—Am. bor.
inflexus, Poir. Encyc. vi. 50 = Echinolaena hirta.
inflexus, *R. Br. Prod* i. 195.—Austral.
laevigatus, Trin. Fund. Agrost. 172 = Anthephora elegans.
laniflorus, *Steud. Syn. Pl. Gram.* 110.—Ins. Tahiti.
lappaceus, Linn. Sp. Pl. ed. II. 1488 = Centotheca lappacea.
lappaceus, *Tausch*, *in Flora*, xx. (1837) 97.—Hab.!
Lechleri, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
limensis, *Meyen*, *Reise*, ii. 71.—Peruv.
linearis, Lam. Fl. Fr. iii. 631 = Tragus racemosus.
longifolius, *Hochst. ex Steud. Syn. Pl. Gram.* 109 — Arabia.
macrocarpus, Ledeb. ex Steud. Nom. ed. II. i. 317 = echinatus
macrostachyus, *Hochst. ex Steud. Syn. Pl. Gram.* 109. —Abyssin.
marginalis, Rudge, Pl. Guian. 19. t. 25 = Echinolaena hirta.
mitis, *Anderss. in Peters*, *Reise Mossamb. Bot.* 553.—Afr. trop.
montanus, *Nees*, *in Royle*, *Illustr. Bot. Himal.* 416.—Ax. et Afr. trop.
mucronatus, Pers. ex Steud. Nom. ed. I. 172 = murricatus.
multiflorus, *Y. & C. Presl*, *Rel. Haenk.* i. 318.—Mexic.
muricatus, Linn. Mant. 302 = Trachys mucronata.
muricatus, *Phil. Sert. Mend. Ali.* 44.—Chili.
myosuroides, *H. B. & K. Nov. Gen. et Sp.* i. 115. t. 35. —Am. trop.
niloticus, Fig. & De Not. in Mem. Acc. Torin. (1854) 380 = catharticus.
orientalis, Willd. ex Kunth, Enum. Pl i. 162 = Pennisetum orientale.
ovatus, *Lam. ex Poir. Encyc.* vi. 51.—Afr. austr.
pallidus, *Fourn. Mex. Pl. Enum. Gram.* 50; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 506.—Mexic.
parviflorus, Poir. Encyc. vi. 52 = Panicum Ventenatii.
pauciflorus, Benth. Bot. Voy Sulph. 56 = tribuloides.
pennisetiformis, *Hochst. & Steud. ex Steud. Nom.* ed. II. i. 317.—Arabia.
pilosus, *H. B. & K. Nov. Gen. et Sp.* i. 116. t. 36.—Am. austr.
platyacanthus, *Anderss. Enum. Pl. Ins. Galap.* 139.—Ins. Galapag.
pubescens, *Steud. Nom.* ed. II. i. 317.—Afr. austr.
pungens, *H. B. & K. Nov. Gen. et Sp.* i. 115.—Mexic.
purpurascens, Thunb. in Trans. Linn. Soc. ii. (1794) 329 = Cymbrix japonica.
pycnostachyus, *Steud. Syn. Pl. Gram.* 109.—Afr. trop.
racemosus, Linn. Sp. Pl. 1049 = Tragus racemosus.
ramosissimus, Poir. Encyc. vi. 51 = Pennisetum dichotomum.
rigidifolius, *Fig. & De Not. in Mem. Acc. Torin.* (1854) 386.—Nubia.
rigidus, *Willd. ex Doell*, *in Mart. Fl. Bras.* ii. 11. 310, in obs.—Bras.
roseus, *Fourn. Mex. Pl. Enum. Gram.* 50; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 507.—Mexic.
rufescens, Desf. Fl. Atlant. ii. 388 = Pennisetum cenchroides.
Schimperi, Hochst. & Steud. ex Steud. Nom. ed. II. i. 317 = montanus.
setigerus, Steud. l. c. = Pennisetum Vahlii.
setigerus, Vahl, Enum. ii. 395 = agrimonoides.
setosus, Sw. Prod. Veg. Ind. Occ. 26 = Pennisetum setosum.
spicatus, Cav. Desc. 304, 614 = Pennisetum typhoideum.
spinifex, Cav. Ic. v. 38. t. 461 = tribuloides.
strictus, Chapm. in Coult. Bot. Gaz. iii. (1878) 20 = incertus.
taitensis, *Steud. Syn. Pl. Gram.* 419.—Ins. Tahiti.
tomentosus, *Poir. Encyc.* vi. 51.—Afr. austr.
tribuloides, *Linn. Sp. Pl.* 1050.—Am. bor.
tribuloides, Herb. Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 55 = Trachys mucronata.
tripsacoides, R. Br. in Salt, Abyss. App. 62 = montanus.

CENCHRUS:—
uniflorus, Ehrenb. ex Boiss. Fl. Orient. v. 448 = montanus.
vaginatus, *Steud. Syn. Pl. Gram.* 110.—Hab.!
villosus, Spreng. Syst. i. 301 = Anthephora elegans.
viridis, *Spreng. l. c.*—Am. trop.

ČENĚKIA, Opiz, Seznam, 36 (1852) = **Campanula**, Tourn.
rapunculoides, Opiz, l. c. = Camp. rapunculoides.
simplex, Opiz, l. c. = Camp. bononiensis.

CENIA, Comm. ex Juss. Gen. 183 (1789). *COMPOSITAE*, Benth. & Hook. f. ii. 429.
LANCISIA, Gaertn Fruct. ii. 422 (1791).
anthemoides, DC. Prod. vi. 88 = turbinata.
barbata, *Benth. & Hook f. Gen.* ii. 430.—Afr. austr.
debilis, DC. Prod. vi. 83 = turbinata.
discoidea, Less. Syn. Comp. 261 = Cotula nudicaulis.
discolor, DC. Prod. vi. 82 = sericea.
flacculosa, DC. l. c. 83 = microglossa.
geminata, *Kunze*, *Del. Sem. Hort. Lips.* (1846) 1.—Afr austr.
hispida, *Benth. & Hook f. Gen.* ii. 430.—Afr. austr.
microcephala, *E. Mey. ex DC. Prod.* vi. 79.—Afr. austr.
microglossa, *DC. l. c.* 83.—Afr. austr.
nudicaulis, E. Mey. ex Heynh. Nom. ii. 130 = Cotula nudicaulis.
pectinata, *DC. Prod.* vi. 83.—Afr. austr.
prunosa, DC. l. c. 82 = turbinata.
sericea, *DC. l. c.*—Afr. austr.
subheterocarpa, Less. Syn. Comp. 261 = turbinata.
Thunbergii, *Benth. & Hook. f. Gen* ii. 430.—Afr austr.
turbinata, *Pers. Syn.* ii. 465.—Afr. austr.
villosa, Heynh. Nom. ii. 130 = Cotula villosa.

CENOCLINE, C. Koch, in Bot Zeit. i. (1843) 41 = **Cotula**, Tourn. (Compos.).
acutiloba, C. Koch, l. c. = Matricaria acutiloba.
alsuda, C. Koch, l. c. = Matricaria alsuda.
aurea, C. Koch, l. c. = Matricaria aurea.
chamaemelifolia, C. Koch, l. c. = Cotula chamaemelifolia.
cinerea, C. Koch, l. c. = Cotula cinerea
globifera, C. Koch, l. c. = Matricaria globifera.
grandiflora, C. Koch, l. c. = Matricaria grandiflora
paucifora, C. Koch, l. c. = Matricaria discoidea.

CENOLOPHIUM, Koch, Gen. Umbell. 108 (1824) = **Selinum**, Linn. (Umbellif.).
divaricatum, Bess. in Flora, xv. (1832) 11. Beibl. 27 = *Fischeri*.
Fischeri, Koch, Gen. Umbell. 103.—Rossia; Reg. Caucas.; As. bor.
lappanicum, Nyl. ex Nyl. & Sael. Herb. Mus. Fenn. 83 = *Fischeri*.

CENOLOPHON, Blume, Enum. Pl. Jav. 60 (1830) = **Alpinia**, Linn. (Scitamin.).
rubrum, Blume, l. c. = A. rubra.
vitellinum, Horan. Prod. Scitam. 36 = Amomum vitellinum.

CENOSTIGMA, Tul. in Ann. Sc. Nat. Sér. II. xx. 140 t. 3 (1843). *LEGUMINOSAE*, Benth. & Hook. f. i. 564.
angustifolium, Tul. l. c. 141 = Gardnerianum.
Gardnerianum, *Tul. l. c.*—Bras.
macrophyllum, *Tul. l. c.*—Bras.

CENTAUREA, Linn. Gen. ed. I. 263 (1737) (Centauria). *COMPOSITAE*, Benth. & Hook. f. ii. 477.
ACROCENTRON, Cass. in Dict. Sc. Nat. xliv. 87 (1826).
ACROLOPHUS, Cass. l. c. i. 258 (1827).
ACROPTILON, Cass. l. c. 464 (1827).
AEGIALOPHILA, Boiss. & Heldr. Diagn. Ser. I. x. 105 (1849).
AETHEOPAPPUS, Cass. in Dict. Sc. Nat. i. 250 (1827).
ALOPHTUM, Cass. l. c. liv. 493 (1828).
AMBLYOGGON, Fisch. & Mey. ex DC. Prod. vi. 561 (1837).
ANTAUREA, Neckl Elem. i. 70 (1790).
BIBLIZIA, Scher, Esam. Fl. Transs. 409 (1866).

CENTAUREA :—
Calcitrapa, Hall. Enum. Stirp. Helv. ii. 689 (1749).
Calicephalus, C. A. Mey. Verz. Pfl. Cauc. 66 (1831).
Centaurium, Tourn. ex Linn. Syst. ed. I (1735).
Cestrinus, Cass. Bull. Soc. Philom. 83 (1817); et in Dict. Sc. Nat. viii. 24 (1817).
Chartolepis, Cass. Dict. Sc. Nat. xliv. 86 (1826).
Cheirolepis, Boiss. Diagn. Ser. I. x. 106 (1849)
Cheirolophus, Cass. in Dict. Sc. Nat. l. 250 (1827).
Chrysxis, Cass. l. c. lx. 154 (1817).
Colymbada, Hill, Veg. Syst. iv. 31 (1762).
Crepula, Hill. l. c. 9 (1762).
Crocodilium, [Vaill.] Juss. Gen. 173 (1789).
Cyanus, [Tourn.] Linn. Syst. ed. I (1735); Juss. Gen. 174 (1789).
Eriopha, Hill, Hort. Kew. 69 (1769).
Fornicum, Cass. in Bull. Soc. Philom. 93 (1819); et in Dict Sc. Nat. xvii. 249 (1820).
Halocharis, Bieb. ex DC. Prod. vi. 665 (1837).
Heraclea, Hill, Veg. Syst. iv. 37 (1762).
Heterolophus, Cass. in Dict. Sc. Nat. l. 250 (1827).
Hippophaestum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 448 (1821).
Hookia, Neck. Elem. i. 70 (1790).
Hyalea, Jaub. & Spach, Illustr. ni. 19. t. 214, 217, 292 (1847).
Hymenocentron, Cass. in Dict. Sc. Nat. xliv. 37 (1826).
Hymenocephalus, Jaub. & Spach, Illustr in 12. t. 209 (1847).
Jacea, [Tourn.] Linn. Syst. ed. I (1735); Juss. Gen. 173 (1789).
Lepteranthus, Neck. Elem. i. 78 (1790).
Leucantha, S. F. Gray, Nat. Arr. Brit. Pl. ii. 443 (1821).
Lopholoma, Cass. in Dict. Sc. Nat. xliv. 37 (1826).
Malacocephalus, Tausch, in Flora, xi. (1828) ii. 481
Mantisalca, Cass. in Bull Soc. Philom. (1818) 142
Melanoloma, Cass. in Dict. Sc. Nat. xxix. 472 (1823).
Menomphalus, Pomel, Nouv. Mat. Fl. Atl. 32 (1874).
Mesocentron, Cass. in Dict. Sc. Nat. xliv. 38 (1826).
Microlonchus, Cass. l. c. (1826); i. 247 (1827).
Microlophus, Cass. l. c. xlv. 37 (1826).
Odontolophus, Cass. l. c. l. 252 (1827).
Pachycentron, Pomel, Nouv. Mat. Fl. Atl. 32 (1874).
Pectinastrum, Cass. in Dict. Sc. Nat. xliv. 38 (1826).
Phaeopappus, Boiss. Diagn. Ser. I. vi. 122 (1846).
Phalolepis, Cass. in Dict. Sc. Nat. l. 248 (1827).
Philostixus, Cass. l. c. xxxix. 498 (1826).
Phrygia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 441 (1821).
Piptocras, Cass. in Dict. Sc. Nat. l. 469 (1827).
Platylophus, Cass. l. c. xliv. 36 (1826).
Plectocephalus, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. t. 51 (1838).
Phora, Hill, Veg. Syst. iv. 30 (1762).
Polyacantha, S. F. Gray, Nat. Arr. Brit. Pl. ii 448 (1821).
Pterolophus. Cass. in Dict. Sc. Nat. xliv. 34 (1826).
Ptohimopappus, Boiss. Voy. Bot. Espagne, 739 (1839–45).
Pterhellus, Cass. in Dict. Sc. Nat. xliii. 488 (1826).
Podia, Neck. Elem. i. 73 (1790).
Pycnocomus, Hill, Syst. Veg. iv. 38 (1762).
Rhapontica, Hill. l. c. 47 (1763).
Rhapontium, [Vaill.] Adans. Fam. ii. 117 (1763); Juss. Gen. 174 (1789).
Seridia, Juss. Gen. 173 (1789).
Setachna. Dulac, Fl. Hautes-Pyr. 518 (1867).
Sagmen, Hill, Veg. Syst. iv. 35 (1762).
Spilacron, Cass. in Dict. Sc. Nat. l. 236 (1827).
Solstitiaria, Hill, Veg. Syst. iv. 21 (1762).
Sphaerocephala, Hill, l. c. 46 (1763).
Stenolophus, Cass. in Dict. Sc. Nat. l. 499 (1827).

CENTAUREA :—
Stizolophus, Cass. in Dict. Sc. Nat. xliv. 35 (1826), nomen; li. 49 (1827).
Staehe, Hill, Veg. Syst. iv. 33 (1762).
Tetramorphaea, DC. in Guill. Arch. Bot. ii. (1838) 381.
Tomantha, DC. Prod. vi. 564 (1837).
Triplocentron, Cass. in Dict. Sc. Nat. xliv. 38 (1826).
Verutina, Adans. Fam. ii. 116 (1763).
Verutina, Cass. in Dict. Sc. Nat. xliv. 38 (1826).
Xanthopsis, C. Koch, in Linnaea, xxiv. (1851) 422.

abrotanifolia, Lam. Encyc. i. 670 = paniculata.
abyssinica, *Sch. Bip. ex Boiss. Diagn.* Ser. II. ii. 62 —Abyssin.
acanthodes, *Spreng. Syst* iii. 403.—Hab ?
acaulis, Cav. ex Willk. & Lange, Prod. Fl Hisp ii. 150 = Cavanillesiana.
acaulis, Forsk. Fl. Aegypt. Arab. 52 = glomerata.
acaulis, Hacusel. ex Willk. & Lange, Prod. Fl. Hisp. ii. 151 = Hacuseleri.
acaulis, Lag. l. c. = Graelsii.
acaulis. *Linn. Sp. Pl.* 914 —Afr. bor.
achaia, *Boiss. & Heldr. Diagn.* Ser. II. iii. 79.—Graecia.
acicularis, *Sibth & Sm. Fl. Graec* t. 911.—Archip. Graec.
acmophylla, Boiss. Diagn Ser. II. *iii. 68 = axillaris*
acrolepis, *Hoffmgg. ex Heynh. Nom* ii. 181, nomen.—Hab ?
acromacra, *Hoffmgg. ex Heynh. l c.* nomen —Hab.?
acuta, Lam. Fl. Fr. ii. 49 = Crupina vulgaris.
acuta, Vahl, ex Bieb. Casp. 82 = C. Behen.
acutangula, *Boss & Reut. Pugill. Pl. Nov.* (1852) 68. —Hispan.
acutifolia, Jord. Pugill 105 = pectinata.
acutifolia, Schur, Enum. Pl. Transs. 899, Index = phrygia.
acutiloba, DC. Prod. vi. 581 = crustata.
Adami, Willd. Sp. Pl. iii. 2310 = solstitialis.
adenolepis, Reichb in Moesal. Handb. ed. II. ii. 1545 = rupestris.
adpressa, Ledeb. ex Steud. Nom ed. II. i. 316 = stereophylla.
adulterina, Moretti, ex DC. Prod. vi. 596 = C. Calcitrapa.
aegialophila, Boiss. & Heldr. ex Boiss. Fl. Orient. iii. 704 = cretica.
aegyptiaca, *Linn. Mant.* i. 118.—Aegypt.; Arab.
aegyptiaca, Sibth. & Sm. Fl. Graec. t. 907 = iberica
aeolica, Guss. ex DC. Prod. vi. 584 = aplolepus.
affinis, Frivald. in Flora, xix. (1836) 485 = dissecta.
africana, *Lam. Encyc.* i. 664.—Afr. bor.
agregata, *Fisch. & Mey. in DC. Prod.* vi. 585.—As Min. Persia.
Aichangeriana, Welw. ex Reichb. Fl. Germ. Excurs. 846 = variegata.
anatensis, Boiss. Diagn. Ser. I. x. 117 = eryngioides.
alata, Lam Encyc. i. 665 = C. Behen.
alba, Bieb. Fl. Taur. Cauc. ii 853 = sterilis.
alba, Herb. Calc. ex C. B. Clarke, Comp. Ind. 289 = Tricholepis elongata.
alba, *Linn. Sp. Pl.* 914.—Eur. austr. ; Reg. Caucas
alba, Suter, Fl. Helv. ii. 204 = amara.
albida, C. Koch, in Linnaea, xxiv. (1851) 483 = aggregata.
albiflora, C. Koch, l. c. 480 = carduiformis.
alexandrina, *Delile, Fl. Egypte*, 280. t. 49. f. 3.— Aegypt.
algeriensis, *Dur. & Coss. Notes Pl. Crit.* 186.—Algeria.
Alophium, *DC. Prod.* vi. 601.—Hispan.
alpestris, Hegetschw. Fl. Schw. 854 = C. Scabiosa.
alpina, *Linn. Sp. Pl.* 910.—Europ.
altaica, Fisch. ex Spreng. Pugill. i. 59 = Leuzea salina.
amana, Boiss. & Bal. Diagn. Ser. II. vi 112 = amoena.
amara. Habl. Taur. 155 = trinervia.
amara, *Linn. Sp. Pl.* ed. II. 1293.—Europ. austr.; As. Min.
Amberboi, Mill. Gard. Dict. ed. VIII. n. 9 = moschata.
ambigua, *Guss. Ind. Sem. Hort. Boccad.* (1826) 8; *ex DC. Prod.* vi. 592.—Italia.
ambigua, Thom. ex DC. l. c. 573 = nervosa.
ambiensis, *Grenlle, in Mem. Acad. Cienc. Madr.* ii. (1859) 463.—Hispan.
ambylolepis, *Ledeb. Fl. Ross.* ii. 708.—Reg. Caucas.

CENTAUREA :—
ambracea, Sehkuhr, Handb. t. 201 = moschata
americana, *Nutt. in Journ. Acad. Philad.* (1821) 117. —Am. bor.
americana, Spreng. Syst. iii. 407 = melitensis.
ammocyanus, *Boiss. Diagn.* Ser. I. x. 109.—Arab., Palaest.
ammophila, Bess. ex DC. Prod. vi. 586 = diffusa.
amoena, *Boiss. & Bal. Diagn.* Ser. II. vi. 112.— Cappadoc.
Amoi, Campo, ex Campo & Amo, in Revista, v (1855) 56 = Boissieri.
amourensis, *Pomel, Nouv. Mat. Fl. Atl.* 30.— Algeria.
amplexicaulis, S. G. Gmel. ft. i. 186. t. 24 = Jurinea polyclonos.
amplifolia, *Boiss. & Heldr. Diagn.* Ser II. iii. 68.— Graecia.
anatolica, Griseb. Spicil. Fl. Rumel. ii 234 = depressa.
angulosa. *Pomel, Nouv. Mat. Fl. Atl.* 26.—Algeria.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 7 = amara.
angustifolia, Schrank, Baier. Fl. ii 376 = amara.
antennata, *Dufour, in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 158.—Hispan.
antiochia, *Boss. Diagn.* Ser. I x. 115.—Syria.
aphrodisea, *Boiss. l. c. iv* 17.—Caria
apiculata, Ledeb Ind Hort. Dorp. Suppl. (1824) 8 = C. Scabiosa.
apiolepis, *Moretti, in Brugnat. Giorn. Fis.* II. ix. (1826) 154.—Italia.
appendiculata, C. Koch, in Linnaea, xxiv. (1851) 425 = Balansae.
Aquila, Lam. Encyc. i. 674 = melitensis.
arachnoidea, Viv. Ann. Bot. i. ii. 185 = rupestris.
arancosa, *Boiss. Diagn* Ser. I. x. 121.—Syria.
arbores, *Webb & Berth. Phyt. Canar.* ii. 256.—Ins. Canar.
arenaria, *Bieb. Fl. Taur. Cauc.* ii. 347.—Europ. or , Reg. Caucas.
arenaria, Reichb Pl Crit. v. 26. t. 443. f. 634 = Reichenbachii.
arenaria, Urv. in Mém. Soc. Linn. Par. i. (1822) 371 = polyclada.
argentea, Frivald ex Nym. Consp. 429 = rupestris.
argentea, *Linn. Sp. Pl.* 912.—Ins Creta et Istria.
argophylla, Steud. Nom ed. I 173 = argyrophylla.
arguta, *Nees, Hor. Phys Berol* 116, t 25.—Ins. Canar.
argyrophylla, *Willd. Enum. Hort. Berol. Suppl.* 61.— Hab.?
arifolia, *Boiss Diagn* Ser I. x 112.—Syria.
aristata, *Renth & Hook. f. Gen.* ii 480.—Cappadoc.
aristata, Hoffmgg. & Link, Fl. Port. ii. 226 = caerulescens.
Armena, *Boiss. Fl. Orient.* iii. 676.—As. Min.
armoracifolia, *Sibth & Sm. Fl. Graec. Prod.* ii. 205.— Graecia.
arrectispina, Bertol. Fl. Ital. ix. 487 = C. Calcitrapa.
aspera, *Linn. Sp. Pl.* 914.—Europ. austr.
astrachanica, Spreng. Syst. iii. 397 = stereophylla.
athoa, DC. Prod. vi. 586 = rupestris.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 383.—Algeria.
atrata, Willd. Sp. Pl. iii. 2300 = axillaris.
atropurpurea, Griseb. Spicil. Fl. Rumel. ii. 237 = C. Scabiosa.
atropurpurea, Oliver, Voy. i. 218 = Olivieriana.
atropurpurea, *Waldst. & Kit. Pl. Rar. Hung.* ii. 121. —Hungar.
atropurpurea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 65, 122 = C. Centaurium.
attica, Nym. Syll. 38 = graeca.
Auctoriana, *DC. Prod.* vi. 576.—Ausyr.
augustana, Reichb. f. Ic. Fl. Germ. xv. 37. t. 785 = maculosa.
aures, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 265.— Europ. austr.
aurea, Hort. ex Steud. Nom. ed. II. i. 316 = nicaeensis.
auriculata, Balb. ex Pers. Syn. ii. 486 = C. Seridis.
auriculata, Spreng. Pugill. i. 59 = stereophylla.
australis, *Benth & Hook. f. Gen.* ii. 479.—Austral.
australis, Pasa. Elem. 44 = sublanata.
australis, Poepp. ex DC. Prod. vi. 575 = chilensis.
austriaca, C. A. Mey. Verz. Pfl. Cauc. 64 = trichocephala.
austriaca, Reichb. Ic. Fl. Germ. xv. 37. t. 785, 3–4 = phrygia.

CENTAUREA:—
austriaca, Rochel, Pl. Banat. Rar. 76. t. 36. *f.* 5 = nervosa.
austriaca, *Willd. Sp. Pl.* iii 2283.—Europ.; Reg. Caucas.: As bor.
axillaris, *Willd. l. c.* 2290.—Europ.; Orient.
babylonica, Bieb Casp 32 = C. Behen.
babylonica, *Linn. Mant.* ii. 460.—As. Min.; Syria.
badensis, Trat. Arch. Gew. i. 23 = C. Scabiosa.
Balansae, *Boiss. & Reut. Diagn.* Ser. II. iii. 82.—Afr. bor.
baldensis, Pers. Syn. ii. 481 = alpina.
balearica, *Rodrig. in Bull. Soc. Bot. Fr.* xvi. (1869) 237. —Ins. Minorca.
Balsamita, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. t. 255 = Phaeopappus spectabilis.
Balsamita, *Lam. Encyc.* i. 667.—As. Min.; Persia.
banatica, Kern. ex Nym. Consp. 437 = arenaria.
barbara, *Fennel, Nouv. Mat. Fl. Atl.* 27.—Algeria.
Barrelieri, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 301 = Serratula pinnatifida.
Barrelieri, *Spreng. Syst.* iii. 403.—Hungar.
Behen, DC. Prod vi. 567 = Stoebisiana.
Behen, Lam. Encyc. i. 665 = cerinthifolia.
Behen, *Linn Sp Pl.* 914.—As. Min.; Syria; Persia.
Belangeriana, Steph, in Denkschr. Acad. Wien, i. (1885) 66 = phyllocephala.
bella, *Trautv. in Bull. Acad. St. Pétersb.* x. (1866) 394.—Reg. Caucas.
benedicta, Linn Sp. Pl. ed. II. 1296 = Carbenia benedicta.
Bernei, Sieber, ex Steud. Nom. ed. II. i. 319 = nervosa.
Besseriana, *DC. Prod* vi. 585.—Europ. or.
Besseriana, Janka, ex Nym. Consp. 427 = tenuiflora.
bicolor, *C. Koch, in Linnaea*, xxiv. (1851) 428.—Reg. Caucas.
Bieberstinii, DC. Prod. vi. 588 = maculosa.
Bieberstenii, Schur, Verh. Siebenb. Ver. Naturf. (1859) 75 = Reichenbachii.
biflora, *Kalew. in Bull. Soc. Nat. Mosc.* xviii. (1845) i. 337.—Astrakhan.
bimorpha, Vlv. Fl. Lib. Spec. 58. t. 24 = sonchifolia
Boissieri, *DC. Prod.* vi. 608.—Hispan.
Bonarota, Walp. Rep. vi. 266 = graeca.
bombycina, *Boiss. in DC. Prod.* vii. 302.—Hispan.
borealis, Sallsb. Prod. 208 = C. Scabiosa.
borysthenica, *Gruner, in Bull. Soc. Nat. Mosc.* xii. (1868) ii. 426.—Taurica.
Bourgaei, *Boiss. Fl. Orient.* iii. 687.—Lycia
Bourgaei, Schult. ex Nym Consp. 426 = Langei.
Brachiii, Reichb. f. ic. Fl. Germ. xv. 35. t. 786 = maculosa.
brachyptera, DC. Prod. vi. 567 = C. Behen.
bractata, Balb. ex Steud. Nom. ed. II. i. 319 = nigrescens.
bracteata, *Benth. & Hook. f. Gen.* ii. 481.—Syria.
bractoata, Scop. Delic. Insub. ii. 17 = amara.
brevicaulis, Bolss. Diagn. Ser. I. x. 120 = pallescens.
brevipappa, Boiss. & Reut. i. c. Ser. II. iii. 71 = nigrescens.
Brousonna x, *Bonnet, in Bull. Soc. Bot. Fr.* xxvii. (1880) p. xl.—Gallia.
Bubanii, Timb. in Mém. Acad. Sc. Toul. Sec. VIII. t. ii. (1879) 137 = carnicolensis.
budaicacea, *Tardent, Ess. Hist. Nat. Bessar.* 61, nomen.—Bessarab.
bulbosa, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 111.—Chili
busambarensis, *Guss. Fl Sic. Syn.* ii. 878.—Sicilia.
cachinalensis, *Phil. Fl. Atac.* 94.—Chili.
cadmea, *Boiss. Diagn.* Ser. I. iv. 18.—Caria.
caerulescens, *Willd. Sp. Pl.* iii. 2319.—Europ. austr.
caespitosa, Cyrill. Pl. Neap. Nov. Fasc. i. 24 = sphaerocephala.
calalepis, *Boiss. in Tchihat. Asie Min. Bot.* t. 31.—As. Min.
calcaria, Jord. Obs. Pl. Crit. vii. (1849) 32 = C. Scabiosa.
Calcitrapa, Bieb. Fl. Taur Cauc. ii. 594 = iberica.
Calcitrapa, *Linn. Sp. Pl.* 917.—Europ.; Afr. bor.; As. temp.
calcitrapoides, DC. Prod. vi. 597 = iberica.
calcitrapoides, Gouan, Fl. Monsp. 461 = Pourzii.
calcitrapoides, *Linn. Cent.* i. 29; *Amoen. Acad.* iv. 291.—Reg. Caucas.; As. Min.
calcitrapoides, Thuill. Fl. Par. 446 = C. Calcitrapa.
calendulacea, Lam. Encyc. i. 665 = Zoegea leptaurea.

CENTAUREA:—
calliacantha, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 82 = patula.
calocephala, Willd. Enum. Hort. Berol. 928 = atropurpurea.
calolepis, Boiss. Diagn. Ser. I. vi. 129 = diffusa.
calophylla, Steud. Nom. ed. I. 173 = atropurpurea.
calva, *Ledeb. ex DC. Prod.* vi. 583.—Mont. Altai.
calva, Dest. ex Nym. Consp. 430 = alba.
cana, C. A. Mey. ex Steud. Nom. ed. II. i. 320 = Ledebourii.
cana, Sibth. & Sm. Fl. Graec. Prod. ii. 198 = axillaris.
canadensis, *Hort. Huber, ex Regel, Ind. Sem. Hort. Petrop.* (1866) 90.—Am. bor.
canariensis, *Willd. Enum. Hort. Berol.* 928.—Ins. Teneriff.
cancellata, Sieber, ex Spreng. Syst. iv. 406 = aegyptiaca.
candidissima, Lam. Encyc. i. 669 = C. Cineraria.
Candolleana, *Boss. Diagn.* Ser. I. vi. 127.—Persia.
Candollei, *C. Koch, in Linnaea*, xxiv. (1851) 424.—Reg. Caucas.
capillata, *Linn. Sp. Pl.* 910 (sp. dub.).—Sibir.; Hispan. cappadocica, *DC. Prod.* vi. 595.—As. Min.
caprina, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 394 = orbiculata.
cardunfolia, Sallsb. Prod. 208 = C. Calcitrapa.
carduiformis, DC. Prod. vi. 590.—As. Min.; Persia.
carduncellis, DC. l. c. 591 = raphanina.
Cardunculus, *Boss. Diagn.* Ser. I. vi. 135.—Assyria.
Carduncullus, Pall. Reise, i. 500 = Serratula glauca.
Cardwus, Forsk. Fl. Aegypt. Arab. 152 = Atractylis flava.
Carelli, *Uoria, in Roem. Arch.* i. 70.—Sicilia.
carienala, *Boiss. Diagn.* Ser. I. iv. 19.—As. Min.
carnicola, *Boss. l. c.* Ser. II. iii. 88.—Illicia.
carniolica, Host, Fl. Austr. ii. 517 = montana.
carpitmana, Walt. Fl Carol. 214 = americana!
carpathica, (genera. Elem. n. 824 = montana.
carpatica, *Floriano, in Magyar Növönt. Lapok*, ix. (1885) 128.—Hungar.
carpetana, Boiss. & Reut. Pugill. Pl. Nov. (1852) 65 = oligra.
:arratracensis, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 94.—Hispan.
carthaginales, *Benth. & Hook. f. Gen.* ii. 479.—Sibir.
cassia, *Boiss. Diagn.* Ser. I. x. 108.—Syria.
Castellana, *Boss. l. c.* vi. 129.—Hispan.
catanica, *Strobl & Haussk. ex Boiss. Fl. Orient.* iii. 669.—Catania.
caucasica, Bieb. ex DC. Prod. vi. 579 = montana.
Cavanillesiana, *Graells, in Mem. Acad. Cienc. Madr.* ii. (1859) 464.—Hispan.
Ceccarinana, *Boiss. & Heldr. Diagn.* Ser. II. iii. 81.—Macedon.
Centaurium, Host, ex Nym. Consp. 417 = Serratula coronata.
Centaurium, *Linn. Sp. Pl.* 910.—Italia.
centaurioides, Bieb. Fl. Taur. Cauc. ii. 359 = salonitana.
centauroides, *Linn. Sp. Pl.* 918.—Europ.
cephalariaefolia, *Willd. in Flora*, xxxiv. (1851) 762.—Hispan.
ceratophylla, Moris, Elench. i. 55 = filiformis.
ceratophylla, Tenore, Cat. Hort. Neap. (1819) 72 = rupestris.
cerinthifolia, Sibth. & Sm. Fl. Graec. Prod. ii. 197.—Graecia.
chamaerhapsoticum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 526.—Marocco.
Chaubardi, Reichb. f. ic. Fl. Germ. xv. 34. t. 781 = maculosa.
cheiracantha, *Fenzl, ex Boiss. Fl. Orient.* iii. 693.—As. Min.; Syria.
cheiranthifolia, Willd. Phytog. 13 = axillaris.
chilensis, *Bert. ex Bull. Férussac* (1830), 109, nomen; *et ex Hook. & Arn. in Bot. Beech. Voy.* i. 88.—Chili.
chlorantha, Adams, ex Ledeb. Fl. Ross. ii. 699 = montana.
Chouletiana, *Pomel, Nouv. Mat. Fl. Atl.* 31.—Algeria.
chrysanthemifolia, *Tausch, in Flora*, xii. (1829) i. *Erg.* 43.—Hab.!
chrysolecus, *Boiss. Diagn.* Ser. I. iv. 21, in texte; *Fl. Orient.* iii. 678.—As. Min.
chrysolepis, Vis. in Mem. Ist. Venet. ix. (1860) 173 = orientalis.

CENTAUREA:—
chrysolopha, Boiss. & Kotschy, Diagn. Ser. II. iii. 78 = aggregata.
chthonocephala, Boiss. Fl. Orient. iii. 677 = ustulata.
cichoracea, Linn. Syst. ed. V. 1321 = Serratula cichoracea.
cichoracea, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 66 = Serratula cancasica.
cinctuplicata, Hornem. Hort. Hafn. ii. 849 = collina.
ciliata, Frivald. in Flora, xviii. (1835) 334 = pallida.
cilicica, *Boiss. & Bal. Diagn.* Ser. II. v. 113.—Cilicia.
Cineraria, Jacq. ex Nym. Consp. 425 = cinerea.
Cineraria, *Linn. Sp. Pl.* 912.—Europ.; Afr. bor.
Cineraria, Pall. Tabl. Taur. 58 = declinata.
Cineraria, Urv. in Mém. Soc. Linn. i. (1822) 371 = ratifolia.
cinerascens, *Bubani, in Nuov. Giorn. Bot. Ital.* v. (1873) 319.—Hispan.
cinerea, Griseb. Spicil. Fl. Rumel. ii. 336 = diascia.
cinerea, *Lam. Encyc.* i. 669.—Italia.
cirrata, *Reichb. Fl. Germ. Excurs.* 214.—Europ. austr.
Clementei, *Boiss. in DC. Prod.* vii. 303.—Hispan.
coarctata, Spreng. Syst. iii. 407 = glomerata.
cochinchinensis, *DC. Prod* vi. 601.—Cochinch.
coelolepa, Fisch. & *Mey. ex Heynh. Nom.* ii. 130, nomen.—Kossia.
collina, Asso, Syn. 127 = ornata.
collina, *Linn. Sp. Pl.* 918.—Reg. Mediterr.
collina, Schur, Enum. Fl. Transs. 408 = pseudo-collina.
collina, Scop. ex Nym. Consp. 429 = rupestris.
collina, Sibth. & Sm. Fl. Graec. t. 914 = salonitana.
comata, Jord. Pugill. 106 = pectinata.
commutata, Timb. ex Nym. Consp. 421 = nigra.
Compernana, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 393 = diffusa.
concinna, *Trautv. in Act. Hort. Petrop.* ii. (1873) 558. —Armenia.
concinna, Willd. ex Steud. Nom. ed. II. i. 319 = pulcherrima.
congesta, Willd. ex DC. Prod. vi. 593 = mellitensis.
conglomerata, *C. A. Mey. in Beitr. Pfl. Russl.* v. 44. —Rossia.
comifera, *Linn. Sp. Pl.* 915.—Reg. Mediterr.
conocephala, *C. Bolle, in Bonplandsa*, viii. (1860) 184. —Ins. Canar.
consanguinea, *DC. Prod.* vi. 585.—As. Min.
consimilis, Bor. Fl. Cent. Fr. ed. III. ii. 351 = C. Jacea.
contracta, Vis. Fl. Lib. Spec. 58. t. 24. *f.* 1, 2 = Delilei.
Cordeti, Bolss. & Orph. ex Boiss. Fl. Orient. iii. 663 = Ceccarinana.
coriacea, Waldst. & Kit. Pl. Rar. Hung. ii. 214 = C. Scabiosa.
corinthiaca, Boiss. & Heldr. Diagn. Ser. II. vi. 113 = achaia.
Cornuti, Reichb. Hort. t. 146 = alpina.
coronopifolia, *Lam. Encyc.* i. 667.—Iberia.
corymbosa, Pourr. in Mém. Acad. Sc. Toul. iii. (1788) 310 = maculosa.
Cossoniana, *Ball, in Journ. Bot.* xi. (1873) 369.—Marocco.
Costae, *Willk. in Linnaea*, xxx. (1859–60) 115.—Hispan.
crassifolia, *Bertol. in Ann. Stor. Nat. It.* (1839) 359. —Ins. Melita.
cretica, *Nym. Syll.* 84.—Ins. Creta.
cretica, Spreng. Syst. iii. 408 = Carthamus creticus.
crinita, *Willd. ex Spreng. i. c.* 400.—Hab.!
cristata, *Bertl. in Wendl. Beitr.* ii. 119.—Austria.
crithmifolia, *Vis. Fl. Dalmat.* ii. 40.—Dalmatia.
crocodylioides, Boiss. Diagn. Ser. I. x. 118 = C. Crocodylium.
Crocodylium, *Linn. Sp. Pl.* 919.—Syria.
Crupina, Bory & Chaub. Nouv. Fl. Pélop. 59 = Crupina pseudocrupina.
Crupina, Desf. Fl. Atlant. ii. 292 = Crupina intermedia.
Crupina, Linn. Sp. Pl. 909 = Crupina vulgaris.
Crupina, Sibth. & Sm. Fl. Graec. t. 900 = Crupina Crupinastrum.
Crupinastrum, Moris, Ind. Sem. Hort. Taur. (1852) 13; et in Bot. Zeit. i. (1843) 166 = Crupina Crupinastrum.
crupinoides, Desf. Fl. Atlant. ii. 293 = Volutarella bicolor.
Castel, *Borbas, in Értek. Kovl Magyar Tudom. Akad.* ix. (1879) 28.—Hungar.
cucullata, *Spreng. ex Steud. Nom.* ed. I. 174.—Hab.!

CENTAUREA :—
cuneifolia, Orphan. ex Boiss. Fl. Orient. iii. 645 = subinermis.
cuneifolia, *Sibth. & Sm. Fl. Graec.* t. 901.—Macedon.
cuspidata, Aucher, ex DC. Prod. vi. 577 = Aucheriana.
cuspidata, *Vis. in Flora*, xii. (1829) 1. *Erg.* 22.—Dalmatia.
cyanea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 192 = C. Cyanus.
cyanifolia, *C. Koch, in Linnaea*, xxiv. (1851) 436.—Reg. Caucas.
cyanifolia, Folr. Encycl. Suppl. ii. 158 = solstitialis.
cyanoides, *Bergg. & Wahlenb. in Isis*, V. xxi. (1828) 971.—Syria.
cyanoides, DC. Prod. vi. 578 = axillaris.
cyanoides, *C. Sm. in Buch, Beschr. Canar. Ins.* 147.—Ins. Canar.
Cyanus, *Linn. Sp. Pl.* 911.—Europ.; Reg. Caucas.
cylindracarpa, Reichb. f. Ic Fl. Germ. xv 34. t. 781 = cuneifolia et sublanata.
cynaroides, C. Sm. ex Buch, in Berl. Acad. Abh. 1816–17 (1819) 369 = Serratula canariensis.
cynaroides, *Link, in Buch, Beschr. Canar. Ins.* 147.—Ins. Canar.
cyrtolepta, *Ledeb. Fl. Ross.* ii. 709.—As. Min.
damascena, *Boiss. Diagn.* Ser. I. x. 112.—Syria.
dealbata, Moris, ex Nym. Consp. 425 = C. Cineraria.
dealbata, *Willd. Sp. Pl.* iii 2295.—As. Min.; Reg Caucas.; Persia.
Debeauxii, Godr. & Gren. Fl. Fr. ii. 248 = nigra.
decipiens, Thuill. Fl. Par. ed. II. 445 = C. Jacea.
declinata, Aucher, ex DC. Prod. vi 579 = lanigera.
declinata, Bieb. Fl. Taur. Cauc. iii. 590 = leucophylla.
decumbens, Boiss. ex DC. Prod. vii. 308 = Boissieri.
decumbens, DC. Prod. vi. 576 = leucophylla.
decumbens, Dubois, Fl. Orl. ex Pers. Syn. 485 = C. Jacea.
decurrens, *DC. Prod.* vi. 596.—Hab.?
deflexa, *Tatarinow, in Beitr. Kenntn. Russ. Reich.* xx. 295.—Soongaria.
deinacantha, *Boiss. & Haussk. in Fl. Orient.* iii. 665.—Kurdistan.
Delilii, *Godr. in Mem. Acad. Montp. (Sect. Médic.)* i. (1858) 435.—Gallia?
denticuiata, Lejeune, Rev. Fl. Spa, 188 = nigrescens.
depressa, *Bieb. Fl. Taur. Cauc.* ii. 346.—Reg. Caucas.; Iber.
dertosensis, Costa, Fl. Catal. (1864) 141 = podospermifolia.
derventana, Janka, in Oestr. Bot. Zeitschr. xxi. (1871) 287 = Kerneriana.
derventana, *Vis. & Panč. in Mem. Ist. Venet.* xii. (1864) 472.—Oriens.
dermicephala, Fenzl, Ind. Hort. Vind. (1855) = aggregata.
deusta, Tenore, Prod. Fl. Nap. p. ii = alba.
Dewaseri, Bor. ex Nym. Consp. 431 = C. Calcitrapa.
dianthoides, *Remy, in C. Gay, Fl. Chil.* iv. 312.—Chili.
dichroa, *Boiss. & Heldr. Diagn.* Ser. I. x. 111.—Syria.
dichroantha, Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 104 = sordida.
diffusa, *Lam. Encyc.* i. 675.—Europ. austr. or.; As. Min.
dilata, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 361.—Afr. bor.
dimorpha, *Vis. Fl. Lib. Spec.* 58. t. 24. f. 3.—Afr. bor.
dioneaea, *Gasp. Descr. Trem.* 19, 28.—Ital.; Ins. Trem.
dioscidea, Balb. ex DC. Prod. vi. 578 = fosconlosa.
diosoca, *Hill, Veg. Syst. iv.* 33.—Germ.
diosoca, *Tenore, Prod. Fl. Nap.* p. ii—Italia; Graecia.
divaricata, Wall. Cat. n. 3294 = Volutarella divaricata.
divaricata, Lange, in Kjoeb. Vidensk. Meddel. (1861) 87 = Langei.
divergens, *Vis. Fl. Dalmat.* ii. 87. t. 12 b.—Europ. austr.
diversifolia, Amo, ex Campo & Amo, in Revista, v. (1855) 56 = Amoi.
diversifolia, Lag. ex Boiss. Voy. Bot. Espagne, ii. 354 = segetum.
Dorneclina, Amo, Descr. Fl. Nuev. ex Willk. & Lange, Prod. Fl. Hisp. ii. 173 = Serratula audiosula.

CENTAUREA :—
doryphora, Fenzl, ex Boiss. Fl. Orient. iii. 665 = Urvillei.
drabaefolia, Anct.=seq.
drabifolia, *Sibth. & Sm Prod.* ii. 202; *Fl. Graec.* t. 910.—Bithynia.
dracunculifolia, *Dufour, in Ann. Sc. Nat.* xxiii. Sér. I. (1881) 157.—Hispan.
druentica x, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 349, 1848.—Gallia.
dubia, S. G. Gmel. ex Steud. Nom. ed. I. 174 = Xeranthemum annuum.
dubia, Krock. Fl. Sil. Suppl. ii. 231 = humilis?
dubia, Suter, Fl. Helv. ii. 303 = nigrescens.
Dubuisii, Bor. Fl. Centr. Fr. ed. III. ii. 350 = C. Jacea.
Dumortieri, *Thys, in Bull. Soc. Bot. Belg.* viii. (1869) 464.—Belg.
dumulosa, *Boiss. Diagn.* Ser. I. x. 111.—Syria.
dutoserasis, Costa, ex Nym. Consp. 429 = podospermifolia.
ebenoides, *Heldr. ex S. Moore, in Journ. Bot.* xvi. (1878) 183.—Graecia.
elata, *Poir. Encyc. Suppl.* ii. 150.—Mauritan.
elbrusensis, *Boss. & Bukse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 131.—Persia.
elegans, All. Fl. Pedem i. 163 = Galeottes tomentosa.
elegans, Saliab. Prod. 207 = C. Cineraria.
elongata, Desf. ex DC. Prod. vi. 604 (Quid?).
elongata, Schousb. Vextr. Marokko, 199 = diluta.
elongata, *Webb & Berth. Phyt. Canar.* ii. 860.—Ins. Canar.
emigrantis, *Bubani, in Nuov. Giorn. Bot. Ital* v. (1873) 13.—Hispan.
Endressii, Hochst. & Stend. ex Steud. Nom. ed. II. i. 320 = nigra.
eriocephalus, Boiss. & Reut. Diagn. Ser. II. iii. 66 = dimorpha.
eriophora, Forsk. Fl. Aegypt. Arab. 74. n. 465 = aegyptiaca.
eriophora, *Linn. Sp. Pl* 916.—Gall.; Lusit.
eriophylla, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 678 = chrysoleuca.
eriophylla, *Spreng. Syst.* iii. 404.—Hab.?
erucoides, Linn. Sp. Pl. 909 = Volutarella crucifolia.
eryngioides, *Lam. Encyc.* i. 675.—Arab.; Syria.
eryngioides, Sibth. & Sm. Fl. Graec. t. 913 = nchala.
Escolzaria, Cosa. ex Nym. Consp. 426 = castellana.
euphratica, Boiss. Diagn. Ser. I. vi. 126 = polypodifolia.
exaltata, Adam, in Weber & Mohr, Beitr. i. 74 = coronopifolia.
exarata, *Boiss. ex Coss. Notes Pl. Crit* 116.—Afr. bor.
exscapa, Urv. in Mém. Soc. Linn. Par. i. (1822) 373 = axillaris.
Fabrei x, *Bonnet, ex Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 349.—Gallia.
Fenzli, *Reichardt, in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 1089.—As. temp.
Ferdinandi, Gren. Cat. Sem. Hort. Bot. Grat. (1847) Collect. 30 = nervosa.
ferox, *Desf. Fl. Atlant.* ii. 297. t. 242.—Numidia.
filiformis, Bieb. ex DC. Prod. vi. 566 = pulchella.
filiformis, *Vis. Fl. Cors. App.* 6.—Corsic.; Sardin.
Fischeri, Willd. Enum. Hort. Berol. Suppl. 61 = axillaris.
fistulosa, *Wender. ex Steud. Nom.* ed. II. i. 320.—Hab.?
flaviceps, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 149 = C. Jacea.
floccosa, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 210.—Chili.
floccosa, Schloss. ex Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 149 = montana.
flocculosa, Schloss. & Vukot. Syll. Croat. 46 = axillaris.
flosculosa, Balb. ex Nym. Consp. 422 = nervosa.
flosculosa, *Willd. Sp. Pl*, iii. 2285.—Europ. austr.
follosa, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iii. 672.—Syria.
Fontanesii, *Spach, ex Dur. in Duch. Rev. Bot.* ii. (1847) 439.—Algeria.
Forardi, *Hornem. Hort. Hafn. Suppl. Add.* 159.—Italia.
Frassii, Noé & Bal. ex Boiss. Fl. Orient. iii. 681 = monacantha.
Frassii, Sch. Bip. ex Nym. Consp. 429 = seq.
Frantzii, Sch. Bip. ex Tchihat. Asie Min. Bot. ii. 323 = Sprunerl.

CENTAUREA :—
fragilis, *Dur. in Duch. Rev. Bot.* ii. (1847) 439.—Afr. sept.
fraylensis, *Sch. Bip. ex Nym. Consp.* 420.—Hispan.
Friderici, *Vis. Fl. Dalmat.* ii. 40. t. 12 C.—Dalmat.
fruticosa, Linn. Syst. ed. X. 1229 = Stachelina fruticosa.
fuliginosa, Dollin. ex Nym. Consp. 424 = C. Scabiosa.
fulva, Huet, in Ann. Sc. Nat. Sér. III. xix. (1853) 254 = nigra.
Funkii, Boiss. & Reut. Diagn. Ser. II. iii. 83 = macrorrhiza.
Funkii, Sch. Bip. ex Willk. in Linnaea, xxv. (1852) 87 = Bourgeri.
furfuracea, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 368.—Algeria.
fuscata, Desf. Fl. Atlant. ii. 302. t. 244 = nicaeensis.
fuscata, Jord. Pugill. 105 = pectinata.
fuscescens, *Pomel, Nouv. Mat. Fl. Atl.* 28.—Algeria.
Galactites, Linn. Sp. Pl. 919 = Galactites tomentosa.
gargunica, Sch. Bip. ex Nym. Consp. 425 = subcilla.
Gaudini, Boiss. & Reut. Diagn. Ser. II. iii. 70 = amara.
Gayana, *Remy, in C. Gay, Fl. Chil.* iv. 310.—Chili.
geluensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 674.—Persia.
geropholaxis, Boiss. Fl. Orient. iii. 677 = ntulata.
Gerberi, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) ii. 391, in texte.—Ross. austr.
Germanician, Hausak. ex Boiss. Fl. Orient. iii. 682 = C. Behen.
gigantea, *Sch. Bip. ex Boiss. l. c.* 672.—Persia.
glaberrima, *Tausch, in Flora*, x. (1827) ii. 249.—Dalmatia.
glandulifera, *Vukot.—Cf. Oestr. Bot. Zeitschr.* xxx. (1880) 163.—Europ.
glastifolia, *Linn. Sp. Pl.* 915.—As. Min.; Reg. Cauc.; Sibir.
glauca, Desf. ex DC. Prod. vi. 587 = calocephala.
glauca, Puschk. ex Willd. Sp. Pl. iii. 2278 = moschata.
glaucescens, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. i. (1854) 31.—Galatia.
Glehni, *Trautv. in Act. Hort. Petrop.* iv. (1876) 382.—Rossia.
globosa, Hort. ex DC. Prod. vi. 569 = alba.
glomerata, *Vahl, Symb. Bot.* ii. 94.—Aegypt.
glomerata, Webb & Berth. Phyt. Canar. ii. 860 = melitensis.
Gmelini, *Steud. Nom.* ed. II. i. 320.—Sibir.
goniocaula, *Boiss. Fl. Orient.* iii. 647.—Lydia.
gracilior, Reut. ex Nym. Consp. 421 = amara.
gracilis, Bor. ex Nym. l. c. = C. Jacea.
graeca, *Boiss. & Sprun. Diagn.* Ser. I. vi. 128.—Graecia.
graeca, Griseb. Spicil. Fl. Rumel. ii. 342 = Ceccatiniana.
Gradaii, *Nym. Consp.* 429.—Hispan.
Graßana, DC. Prod vi. 587 = sordida.
graminifolia, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 169 = montana.
granatensis, *Boiss. ex DC. Prod.* vii. 308.—Hispan.
grandiflora, Moç. & Sesse, ex DC. Prod. vi. 575 = americana.
grandiflora, Pall. Reise, iii. 287 = monanthos.
grinensis, Reut. Ind. Sem. Hort. Genev. (1857) 4.—Cf. Linnaea, xxix. (1857–58) 723 = C. Scabiosa.
Grisebachii, Nym. Consp. 427 = graeca.
gudrumensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 673.—Kurdist.
Gusciardii, *Boiss. l. c.* 661.—Baeotia.
gymnocarpa, *Moris & De Not. Fl. Insul. Caprar.* 76. t. 2.—Ins. Caprar.
gymnocoala, *Benth. & Hook. f. Gen.* ii. 482.—Mexopot.
Haenseleri, *Boiss. Voy. Bot. Espagne*, ii. 349.—Hispan.
Hamrii, Jord. Obs. Pl. Crit. v. (1847) 70 = coerulescens, limbata.
Haussknechtii, *Boiss. Fl. Orient.* iii. 670.—Syria.
Haynaldi, Borb. ex Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 149 = C. Jacea.
helenioides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 683.—Pontus?
hellenica, *Boiss. & Sprun. Diagn.* Ser. I. vi. 181.—Graecia.
hemiptera x, *Borb. in Oestr. Bot. Zeitschr.* xxviii. (1878) 392.—Europ. austr.

CENTAUREA :—
Herbichii, *Janka, in Flora*, xli. (1858) 441.—Transsylv.
Hermonis, Boiss. Diagn. Ser. I. x. 122 = iberica.
heterocarpa, *Boiss. & Gaill. ex Boiss. Fl. Orient.* iii. 680.—Syria.
heterophylla, Sprun. ex Boiss. l. c. 668 = Sprunerі.
heterophylla, Willd. Sp. Pl. iii. 2314 = aspera.
Heuffelii, Kelchb. f. lc. Fl. Germ. xv. 27. t. 801 = Kotschyana.
hierapolitana, *Boiss. Diagn.* Ser. I. iv. 15.—Caria.
hispida, DC. Prod. vi. 589 = reflexa.
Hladnikiana, *Fisch. ex Heynh. Nom.* ii. 180, nomen. —Europ. austr.
Hochstetteri, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 488.—Afr. trop.
Hoefftiana, *C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* vii. (1855) 7.—Reg. Casp.
Hohenackeri, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) 11. 395 = orina.
Hohenackeriana, Trautv. in Act. Hort. Petrop. viii. (1883) 493 = prasc.
hololenca, *Boiss. Fl. Orient.* iii. 694.—Syria.
horrida, *Badaro, in Brugnat. Giorn. Fis.* (1824) 363.—Liguria; Sardin.
horrida, Tenore, Ind. Sem. Hort. Neap. (1820) 15; Syll. 449 = C. Calcitrapa.
Huetii, *Boiss. Diagn.* Ser. II. iii. 69.—Armenia.
humifusa, Gueldenst. ex Ledeb. Fl. Ross. ii. 696 = decumbens.
humilis, Schrank, Baier. Fl. ii. 376 = pullata †
hyalina, L'Hérit. ex DC. Prod. vi. 566 = pulchella.
hyalolepis, Boiss. Diagn. Ser. I. vi. 133 = pallescens.
hybrida, All. Fl. Pedem. i. 161 = corymbosa.
hybrida, Chaix, in Vill. Hist. Pl. Dauph. iii. 54 = C. Calcitrapa.
hymenolepis, *Trautv. in Act. Hort. Petrop.* iii. (1874) 273.—Reg. Caucas.
hybericifolia, Brouss. ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 255 = canariensis.
hypoleuca, *DC. Prod.* vi. 575.—As. Min.
hyssopifolia, Georgi, Beschr. Russ. Reich. iii. iv. 1263 = trichocephala.
hyssopifolia, *Vahl, Symb. Bot.* i. 75.—Hispan.
Hystrix, Boiss. Voy. Bot. Espagne, ii. 851 = polyacantha.
iberica, *Trevir. ex Spreng. Syst.* iii. 406.—Iber.; Armen.; Caucas.
idaea, *Boiss. & Heldr. Diagn.* Ser. I. x. 119.—Creta.
incana, Burm. f. Fl. Cap. Prod. 28 = Arctotis argentea.
incana, *Desf. Fl. Atlant.* ii. 281.—Marocco.
incana, Lag. Gen. et Sp. Nov. 32 = Lagascae.
incana, Losc. & Pardo, Ser. Inconf. 59 = tenuifolia.
incana, *Tenore, Prod. Fl. Nap.* p. li.—Italia.
incompta, *Vis. Fl. Dalmat.* 86. t. 13b. fig. 1.—Dalmat.
indica, Lour. in Linnaea, vi. (1831) 84 = Goniocaulon glabrum.
indurata, Janka, in Flora, xli. (1858) 444 = nigra.
infestans, *Dur. in Duch. Rev. Bot.* ii. (1847) 480.—Afr. bor.
intacta, *Ledeb. Fl. Ross.* ii. 705.—Reg. Caucas.
integrifolia, *C. A. Mey. Verz. Pfl. Cauc.* 64.—Reg. Caucas.
integrifolia, Tausch, in Flora, xi. (1828) ii. 485 = nigra.
integrifolia, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 153 = C. Scabiosa.
integriuscula, Vukot. l. c. 152 = C. Scabiosa.
intermedia, Cariot, Etud. Fl. ed. V. 317 = axillaris.
intermedia, *Mutel, in Duch. Rev. Bot.* i. (1845-46) 400.—Algeria.
interrupta, *Hoffmgg. & Link, Fl. Port.* ii. 237.—Lusitan.
intricata, *Boiss. Diagn.* Ser. I. vi. 130.—Persia.
intybacea, *Lam. Encyc.* i. 671.—Gallia; Hispan.
involucrata, Fisch. ex Nym. Consp. 418 = trinervia.
inuata, *Hoffmgg. ex Heynh. Nom.* ii. 181, nomen.—Hab.?
involucrata, *Desf. Fl. Atlant.* ii. 295.—Alger.
iserviana, Graelis, ex Willk. & Lange, Prod. Fl. Hisp. ii. 162 = Janerii.
Isernii, *Willk. in Willk. & Lange, l. c.* 155.—Hispan.
Ismaeli, Linn. Sp. Pl. 916 = aspera.
ispahanica, *Boiss. Diagn.* Ser. I. vi. 133.—Persia.
italica, Steud. Nom. ed. II. i. 230 = seq.
italica, Pers. Syn. ii. 434 = C. Scabiosa.

CENTAUREA :—
ixodes, *Pomel, Nouv. Mat. Fl. Atl.* 384.—Algeria.
Jacea, *Linn. Sp. Pl.* 914.—Europ.; As. med.
Jacobaeaefolia, *Lam. Encyc.* i. 676.—Hab.?
jacobaeaefolia, Spreng. Syst. iii. 404 = argyrophylla.
Jacobi, Dufour, in Bull. Soc. Bot. Fr. vii. (1860) 848 = sonchifolia.
Janerii, *Graells, in Mem. Acad. Cienc. Madr.* ii. (1859) 466 (*Ramill.* 8).—Hispan.
Jankeana, *Simk. in Termész. Füzetek*, i. (1877) 168.—Europ. austr.
Jordaniana, *Grev. & Godr. Fl. Fr.* ii. 245.—Gallia.
jurinaeaefolia, Boiss. Diagn. Ser. II. iii. 78 = rutifolia.
Juvenalis x, *Delile, ex Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 483.—Gallia.
karduchorum, *Boiss. Fl. Orient.* iii. 633.—Armenia.
kardobaicensis, Frivald. ex Hampe, in Flora, xx. (1837) i. 228 = napulifera.
Karschiana, Auersw. ex Nym. Consp. 426 = cristata.
Karstiana, Griseb. Spicil. Fl. Rumel. ii. 240 = pallida.
Kartschiana, *Scop. Carn.* ed. II. ii. 140. t. 58.—Carniol.
Kemeriana, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 178.—Thracia.
kestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 225.—Turkest.
kilaea, *Boiss. Fl. Orient.* iii. 643.—Byzant.
Kochii, F. Schultz, ex Nym. Consp. 421 = nigrescens.
Kolczygini, *Kalen. in Bull. Soc. Nat. Mosc.* xviii. (1845) 1. 286.—Reg. Caucas.
Kotschyana, Heuff. in Koch, Syn. Fl. Germ. 487 = C. Scabiosa.
Kralikii, Boiss. Diagn. Ser. II. iii. 84 = dimorpha.
kurdica, *Bornm. & Richardt, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 1042.—Kurdist.
lacuiata, Guss. in DC. Prod. vi. 582 = ambigua.
lacunica, *Boiss. Fl. Orient.* iii. 660.—Laconia.
lactucaefolia, *Boiss. l. c.* 622.—Ins. Rhodus.
Lagascae, *Nym. Syll.* 83.—Hispan.
Lagascana, Graells, in Mem. Acad. Cienc. Madr. ii. (1859) 466 (Ind. 7) = Graelsii.
lanata, DC. Prod. vi. 610 = Carthamus lanatus.
lanata, Roxb. Hort. Beng. 62; Fl. Ind. iii. 444 = C. Cyanus.
lancifolia, *Sieber, ex Spreng. Syst.* iii. 406.—Creta.
Langeana, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 157 = Langei.
Langei, *Nym. Suppl. Syll.* 6.—Hispan.
lanigera, *DC. Prod.* vi. 579.—Cappadoc.
lappacea, Tenore, Ind. Sem. Hort. Neap. (1829) 15 = solstitialis.
lasiorhiza, *DC. Prod.* vi. 600.—Persia.
latifolia, Pers. Syn. ii. 437 = centaureoides.
latifrons, DC. Prod. vi. 589 = salsoloides.
laza, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 640.—Mesopot.
Ledebourii, Steud. Nom. ed. II. i. 230 = cana.
leptolomella, *Boiss. Diagn.* Ser. I. x. 110.—Syria.
leptolomoa, Pané. ex Nym. Consp. 426 = maculosa.
leptophylla, *C. Koch, in Linnaea*, xxiv. (1851) 419. —Lancasian.
leucantha, Pourr. in Mém. Acad. Toul. iii. (1788) 310 = intybacea.
leucanthe, Web. & Mohr, Beitr. I. 74. n. 49 = montana.
leucacephala, Schur, Enum. Pl. Transs. 408 = phrygia.
leucolepis, DC. Prod. vi. 568 = alba.
leucolepis, Ledeb. ex Nym. Consp. 420 = sterilis.
leucolepis, Tenore, ex Nym. l. c. = splendens.
leucophaea, *Jord. Obs. Pl. Crit.* v. (1847) 64.—Gallia.
leucophylla, *Bieb. Fl. Taur. Cauc.* iii. 591.—Reg. Caucas.
leucoides, *Walp. Ann.* i. 447.—Medit.
ligerina x, *Franch. in Bull. Soc. Bot. Fr.* xxvii. (1880) p. —Gallia.
limbata, *Hoffmgg. & Link, Fl. Port.* ii. 241.—Lusitan.
linariifolia, Steud. Nom. ed. II. i. 231 = linifolia.
linearifolia, Lam. Encyc. i. 667 = linifolia.
lingulata, Lag. Gen. et Sp. Nov. 32 = variegata.
linifolia, Lam. Encyc. i. 667 = hymenolepis.
linifolia, *Linn. Mant.* i. 117; *Vahl, Symb. Bot.* i. 75. —Hispan.; Italia.
Linnaeana, Wale. Ann. i. 450 = Fontanesii.
Lippii, Lam. Sp. Pl. 910 = Volutarella Lippii.
livoniana, Walmn. ex Ledeb. Hort. Dorp. 38 = C. Jacea.
longepedunculata, *Sch. Bip. ex Boiss. Fl. Orient.* iii. 668.—Persia.

CENTAUREA :—
lugdunensis, Jord. Obs. Pl. Crit. v. (1847) 49 = montana.
Lauchaniana, *Heimerl. ex Stapf, in Denkschr. Acad. Wien*, i. (1885) 118.—Lycia.
lusitanica, Boiss. & Reut. Diagn. Ser. II. iii. 85 = sphaerocephala.
lutea, *Gueldenst. It.* ii. 48, *ex Ledeb. Fl. Ross.* ii. 712. —Ross. austr.
lycaonica, *Boiss. & Heldr. Diagn.* Ser. II. v. 111.—As. Min.
lycia, *Boiss. Diagn.* Ser. I. x. 109.—Lycia.
lycopifolia, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iii. 693.—Syria.
lydia, *Boiss. Diagn.* Ser. I. iv. 90.—Bithynia.
lyraefolia, Salisb. Prod. 208 = salmantica.
lyrata, Bell. App. Fl. Pedem. 40 = rhaponticoides.
lyrata, *Pers. Syn.* ii. 436.—Uruguay.
lyrata, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 270.—Syria.
lyrophylla, *Griseb. Spicil. Fl. Rumel.* ii. 238.—Europ. austr.
macedonica, *Boiss. Diagn.* Ser. I. vi. 130.—Graecia.
macracantha, Heldr. ex Boiss. Fl. Orient. iii. 690 = iberica.
macracantha, Guss. Fl. Sic. Syn. ii. 518 = calcitrapoides.
macrocarpa, *Boiss. Fl. Orient.* iii. 677.—Persia.
macrocephala, *Puschk. ex Willd. Sp. Pl.* iii. 2299.—Armen.
macrolopha, *Fenzl, ex Boiss. Fl. Orient.* iii. 695.—Orient.
macrorrhiza, *Willk. in Linnaea*, xxv. (1852) 38.—Hispan.
maculata, Andrz. Rye Botan. i. (1828) 12 = maculosa.
maculosa, Eichw. Skizze, 146 = Biebersteinii.
maculosa, Guss. Pl. Rar. 859 = cinerea.
maculosa, *Lam. Encyc.* i. 669.—Gallia; Hispan.; Italia.
maculosa, Noë, ex Nym. Consp. 425 = cuneifolia.
malacitana, Boiss. Diagn. Ser. II. iii. 85 = sonchifolia.
margaritacea, Tenore, Syll. Add. iii. A. 628 = splendens.
marginata, Tenore, Prod. Fl. Nap. p. lii = ticaceensis.
mariana, Nym. Suppl. Syll. 6 = macrorrhiza.
mariolensis, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 110.—Gallia.
maritima, Dufour, in Ann. Sc. Nat. Sér. I. xxiii. (1831) 165 = sonchifolia.
maroccana, *Batt. in Journ. Bot.* xi. (1873) 370.—Marocco.
Marschalliana, *Spreng. Syst.* iii. 393.—Podol.; Caucas.
Mauciana, *Lour, in Hook. Kew Journ.* viii. (1856) 297.—Ins. Maderа.
mathioliifolia, Boiss. Diagn. Ser. I. iv. 19 = axillaris.
maxima, Forsk. Fl. Aegypt. Arab. 152 = verbascifolia.
melitensis, *Linn. Sp. Pl.* 917.—Europ.; Am. austr.
membranacea, Jacquem. Voy. Bot. 97 = Sanssurea denticulata.
membranacea, Lam. Encyc. i. 666 = monanthos.
mentoyerica, Vill. Hist. Pl. Dauph. i. 365; iii. 48 = C. Scabiosa.
Meryenia, DC. Prod. vi. 598 = iberica.
mexicana, DC. l. c. 575 = americana.
micracantha, *Dufour, in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 164.—Hispan.
micranthа, Hoffmgg. & Link, Fl. Port. ii. 220 = castellana.
micrantioa, S. G. Gmel. It. i. t. 28. f. 1 = maculosa.
microcephala, *Benth. & Hook. f. Gen.* iii. 480.—Cappadoc.
microlepis, *Boiss. Fl. Orient.* iii. 651.—Lycia.
microlonchoides, *Boiss. Diagn.* Ser. I. vi. 137.—Persia.
microptilon, Godr. & Gren. Fl. Fr. ii. 242 = nigra.
Mirgiha, Jord. in Cat. Gratn. Grenob. (1850) 15; et Pugill. 110 = paniculata.
Mixae, Heldr. ex Boiss. Fl. Orient. iii. 662 = Gulcharidii.
minor, Willd. ex Spreng. Syst. iii. 408 = polymorpha.
mixta, DC. Prod. vi. 564 = bellenica.
mollis, Bass. [Fl. Gallic. ii. 307 ?] = axillaris.
mollis, Schleich. Cat. Pl. Helv. ed. IV. 13 = C. Jacea.
mollis, Waldst. & Kit. Pl. Rar. Hung. iii. 243 t. 219 = montana.
monanthos, *Boiss. Fl. Orient.* iii. 681.—Thracia.

CENTAUREA :—
monacantha, *Clark, Trav.* ii. 354.—Cyprus.
monantha, *Georgi, Bemerk. Reise Russ. Reich.* i. 321. —Russia.
monocephala, *Remy, in C. Gay, Fl. Chil.* iv. 311.—Chili.
montana, Auct. ex Willk. & Lange, Prod. Fl. Hisp. ii. 159 = variegata, axillaris.
montana, Berm. f. Prod. Fl. Cap. 26 = sempervirens.
montana, Costa, Fl. Catal. 189 = semidecurrens, lagduensis.
montana, *Linn. Sp. Pl.* 911.—Europ.
monticola, *Boiss. ex DC. Prod.* vii. 302.—Hispan.
Moritziana, Hegetschw. Fl. Schw. 851 = phrygia.
Moschata, *Linn. Sp. Pl.* 909.—Orient.
Moschus, Habl. in S. G. Gmel. It. iv. 176 = Jurinea depressa.
Mouelli, *Blanchet, in Bull. Soc. Bot. Fr.* xxviii. (1880) 21.—Gallia.
mucronata, Forsk. Fl. Aegypt. Arab. 151 = pumila.
mucronifera, *DC. Prod.* vi. 599 :—Cappadoc.
multifida, *Pers. Syn.* ii. 487.—Afr. bor.
multiflora, Pers. l. c. 486 = sicula.
Murati, Jord. Pugill. 108 = coerulescens.
muricata, Linn. Sp. Pl. 918 = Volutarella muricata.
muricata, Spreng. Syst. iii. 402 = Volutarella Lippii.
musarum, *Boiss. & Orph. Diagn.* Ser. II. v. 112.—Graecia.
mutabilis, *St. Amans, in Mém. Mus. Par.* i. (1815) 477.—Mont. Pyren.
mycantha, DC. Fl. Fr. iv. 101 = C. Calcitrapa.
mycenia, Boiss. & Sart. Diagn. Ser. II. vi. 118 = Urvillei.
myriocephala, Clark, Trav. i. 23 = diffusa.
myriocephala, *Sch. Bip. ex Boiss. Fl. Orient.* iii. 682. —Syria.
myriostigma, *Lem. in Illustr. Hortic.* (1856) *Misc.* 88. —Hab.?
myriostoma, *Vis. & Panč. in Mem. Ist. Venet.* xii. (1864) 470.—Orient.
nana, Chem. ex Steud. Nom. ed. II. i. 320, 816 = amara.
nana, *Desf. Fl. Atlant.* ii. 296. t. 241.—Mont. Atl.
nana, Sieber, ex Boiss. Fl. Orient. iii. 675 = raphanina, napifolia, *Linn. Sp. Pl.* 916.—Italia ; Graecia ; Afr. bor.
napulifera, *Rochel, in 'Act. Hungar. Echony.'* (1834) t. 8.—Europ.
neapolitana, Boiss. Diagn. Ser. II. iii. 72 = nigrescens.
neglecta, Bess. Enum. Pl. Volh. 85 = centauroides.
neglecta, Martr. ex Nym. Consp. 481 = sonchifolia.
neglecta, Schott, ex Spreng. Syst. iii. 400 = orientalis.
nemophila, Jord. ex Nym. Consp. 421 = C. Jacea.
nemoralis, Jord. Pugill. Pl. Nov. 104 = nigra.
nervosa, Griseb. Spicil. Fl. Rumel. ii. 388 = Kerneriana.
nervosa, Reichb. ex Steud. Nom. ed. II. i. 321 = phrygia.
nervosa, *Willd. Enum. Hort. Berol.* 925.—Europ. austr.
nevadensis, *Boiss. & Reut. Diagn.* Ser. II. iii. 71.—Hispan.
nicaeensis, *All. Fl. Pedem.* i. 162. t. 74. f. 1.—Gall. ; Ital.
nidulans, *Pomel, Nouv. Mat. Fl. Atl.* 280.—Algeria.
Niederi, *Heldr. in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 380.—Graecia.
nigra, Frivald. ex Boiss. Fl. Orient. iii. 680 = Kerneriana.
nigra, Lam. Encyc. i. 666 = pratensis.
nigra, *Linn. Sp. Pl.* 911.—Europ.
nigrescens, DC. Prod. vi. 571 = nigra.
nigrescens, Gren. & Godr. Fl. Fr. ii. 241 = C. Jacea.
nigrescens, *Willd. Sp. Pl.* iii. 2288.—Europ. med. austr.
nigricans, DC. ex Steud. Nom. ed. II. i. 321 = Endressii.
nigricans, Link, ex Spreng. Syst. iii. 401 = nigra.
Nimrodis, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* iii. 684.—Mesopot.
nissana, *Bieb. ex Willd. Sp. Pl.* iii. 2305.—Tauria ; Reg. Cauc.
nitida, Lam. Fl. Fr. ii. 49 = alba.
nitida, Naldi, ex DC. Prod. vi. 601 = crassifolia.
Nobilis, Boiss. Diagn. Ser. II. iii. 84 = iberica.
Novelli ɔc, *Franch. in Bull. Soc. Bot. Fr.* xxvii. (1880) p. xxi.—Gallia.
nudicaulis, Linn. Syst. ed. X. 1329 = Serratula radiosaulis.

CENTAUREA :—
Nuttallii, Spreng. Syst. iv. Cur. Post. 298 = americana.
obscura, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 290 = nigra.
ochroleps, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 149 = axillaris.
ochroleuca, Pushk. ex Willd. Sp. Pl. iii. 2289 = axillaris.
ochrolopha, Costa, Fl. Catal. 140 = leucophaea.
odorata, *Burm. f. Fl. Cap. Prod.* 26.—Afr. austr.
odorata, *C. Koch, in Linnaea*, xxiv. (1851) 417.—Reg. Caucas.
Olivieri, *Pomel, Nouv. Mat. Fl. Atl.* 282.—Algeria.
Olivieriana, *DC. Prod.* vi. 596.—Ins. Naxos.
olympica, *Boiss. Diagn.* Ser. II. iii. 77.—Bithynia.
omphalodes, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 397, nomen.—Algeria.
onopordifolia, *Boiss. Diagn.* Ser. I. x. 114.—Syria.
orientalis, Baumg. ex Schur, Enum. Pl. Transs. 408 = pseudo-collina.
orientalis, *Linn. Sp. Pl.* 918.—Taur. ; Sibir.
ornata, *Willd. Sp Pl* iii. 2320.—Hispan.
Orphanidea, *Heldr. & Sart. ex Boiss. Diagn.* Ser. II. iii. 78.—Graecia.
orthocaantha, *Tausch, in Flora*, xii. (1829) 1. *Erg.* 44. —Hab. ?
ossica, *C. Koch, in Linnaea*, xvi. (1843) 40.—Reg. Cauca.
ovina, Hort. ex Steud. Nom. ed. II. i. 321 = pterolopha.
ovina, Pall. ex Steud. l. c. = diffusa.
ovina, *Pall. ex Willd. Sp. Pl.* iii. 2292.—Taur. ; Sibir.
pallescens, Bové, ex DC. Prod. vi. 597 = alexandrina.
pallescens, *Delile, Fl. Egypte*, 278 t. 49.—Aegypt.
pallida, *Arnold. in Flora*, xviii. (1835) 883.—Europ.
pallida, Wall. Cat. n. 2988 = Serratula pallida.
pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. x. 120.—Pamphylia
paniculata, Auct. Hisp. ex Willk. & Lange, Prod. Fl. Hisp. ii. 157 = castellana.
paniculata, Bieb. Fl. Taur. Cauc. ii. 346 = maculosa.
paniculata, Lag. ex DC. Prod. vi. 588 = coerulescens.
paniculata, *Linn. Sp. Pl.* 912.—Europ. austr
paniculata, Pett. ex Nym. Consp. 426, 427 = cristata, diffusa.
paniculata, Sibth. & Sm. Fl. Graec. Prod. ii. 199 = graeca.
paniculata, Vill. Hist. Pl. Dauph. iii. 53 = leucophaea.
panicea, *DC. Prod.* vi. 582.—Mons Athos.
Parlatorii, Heldr. in Flora, xxvii. (1844) 1. 68 = seq. ?
Parlatorii, Heldr. in Ann. Acc. Asp. Nat. i. (1843) 201 = dissecta.
Parolinii, *DC. Prod.* vi. 592.—Mons Ida ; As. Min.
parviflora, Bieb. Fl. Taur. Cauc. fii. 594 = diffusa.
parviflora, *Desf. Fl. Atlant.* ii. 301.—Pers. ; Afr. bor.
parviflora, Mar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 450 = squarrosa.
parviflora, Lam. Fl. Fr. ii. 82 = aspera.
parviflora, Sm. Fl. Graec. x. 9. t. 912 = diffusa.
patsbilcensis, DC. Prod. vi. 598 = melitensis.
patula, *DC. l. c.* 583.—Persia ; Cappadoc.
pauciflora, C. Koch, in Linnaea, xxiv. (1851) 433 = squarrulosa.
pauciflora, *Trautv. in Act. Hort. Petrop.* ii. (1873) 474.—Daghestania.
pectinata, Auct. Koss. ex Boiss. Fl. Orient. iii. 632 = trichocephala.
pectinata, *Linn. Sp. Pl.* ed. II. 1287.—Gall. austr.
pectinata, Slev. in Pall. N. Nord. Beitr. vii. 328 = calva.
Pelia, *DC. Prod.* vi. 586.—Graecia.
penicillata, Delile, Fl. Aeg. Illustr. 74 = trichacantha.
peregrina, *Linn. Sp. Pl.* 915.—Europ. austr.
pergamacea, *DC. Prod.* vi. 569.—Cappadoc.
Perrottetii, *DC. l. c.* 598.—Afr. trop.
persica, *Boiss. Diagn.* Ser. I. vi. 132.—Persia.
Pestalotii, De Not. ex Cesati, in Bibl. Ital. xci. (1888) 348 = alba.
Pestalozzae, *Boiss. Diagn.* Ser. I. x. 118.—Lycia.
petrocoma, *Trautv. in Act. Hort. Petrop.* ii. (1873) 552.—Armenia.
Petteri, Reichb. f. Ic. Fl. Germ. xv. 36. t. 783 = divergens.
peucedanifolia, *Boiss. & Orph. in Boiss. Fl. Orient.* iii. 647.—Macedon.
phaeolepis, *Coss. Notes Pl. Crit.* 138.—Mauritania.
phaeopbylla, *Wallr. Beitr. Fl. Hercyn.* 270.—Europ.
Pharsonis, *Pomel, Nouv. Mat. Fl. Atl.* 281.—Algeria.

CENTAUREA :—
phlomoides, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* iii. 678.—Persia.
phrygia, Jacq. Hort. Vindob. 167 = austriaca.
phrygia, Lapeyr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 164 = nigra.
phrygia, *Linn. Sp. Pl.* 910.—Europ.
phrygia, Lour. Fl. Cochinch. ii. 681 = cochinchinensis.
phyllocephala, *Boiss. Diagn.* Ser. I. vi. 134.—Afghan.
Pichleri, *Boiss. Fl. Orient.* iii. 688.—Bithynia.
Pieris, *Pall. Tabl. Taur.* 58 ; *Willd. Sp. Pl.* iii. 2302 —Reg. Casp.
Pinardi, *Boiss. Diagn.* Ser. I. iv. 17.—Caria.
pindicola, *Orph. Reise Rumel.* ii. 164.—Maced.
pinnatifida, Schur, Sert. Transs. 85 = axillaris.
pinnatispina, *Turra, Farset. Nov. Gen.* 18.—Sicilia.
pityophala, Brot. Fl. Lusit. i. 369 = Lecuea conifera.
platyptera, Peterm. in Flora, xxvii. (1844) 475 = C. Jacea.
plumosa, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 245 = gymnocarpa.
plumosa, *Kern. in Oestr. Bot. Zeitschr.* xxii. (1872) 44.—Europ. austr.
podospermifolia, *Losc. & Pard. Ser. Inconf.* 58.—Hispan.
polyacantha, Spreng. Syst. iii. 406 = sonchifolia.
polyacantha, *Willd. Sp. Pl.* iii. 2312.—Hispan.
polycephala, Jord. Obs. Fl. Crit. v. (1847) 67 = paniculata.
polyclada, *DC. Prod.* vi. 584.—Graecia.
polymorpha, *Lag. Gen. et Sp. Nov.* 32.—Hispan.
polyphylla, Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb. ii. (1837) 311 = ossica.
polyphylla, *Pomel, Nouv. Mat. Fl. Atl.* 281.—Algeria.
polypodiifolia, *Boiss. Diagn.* Ser. I. v. 126.—Persia.
Postii, *Boiss. Fl. Orient.* iii. 688.—Syria.
Pottii, [Spreng.] Bot. Gart. Halle, 23 = sicula, sulphurea.
Pourretiana, *Timb. & Theven. in Mém. Acad. Toul.* Sér. VII vi. (1874) 645.—Gallia.
Pouzini, DC. Cat. Hort. Monsp. 91 = C. Calcitrapa.
praealta, *Boiss. & Bal. Diagn.* Ser. II. v. 112.—Cilicia.
praecox, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 438.—Afr. trop.
praetermissa, Martr. in Soc. Sc. Tarn et Gar. (1852) ; Pl. Crit. Tarn, 382 = aspera.
praetorta, Reichb. f. Ic. Fl. Germ. xv. 34. t. 886 = Atractylis Preauxiana.
Preauxiana, Webb & Berth. Phyt. Canar. ii. 250 = Atractylis Preauxiana.
princeps, Boiss. & Heldr. Diagn. Ser. II. vi. 111 = alba.
procera, Heyah. Nom. ii. 191 lapsu = Cephalaria procera.
procumbens, *Balb. Misc. Alt.* 31. t. 1.—Pedemont.
procumbens, Habl. in S. G. Gmel. It. iv. 167 = debilata.
procumbens, Jord. Obs. v. 57 = Jordaniana.
procurrens, DC. Prod. vi. 595 = sinaica.
procurrens, *Sieber, ex Spreng. Syst.* iii. 407.—Palaestin ; Arab.
prolifera, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1850) 486 = Delilei.
prolifera, Vent. Jard. Cels, 16 = glomerata.
Prolongi, *Boiss. ex DC. Prod.* vii. 303.—Hispan.
prostrata, *Coss. Notes Pl. Crit.* 118.—Hispan.
prostrata, Huet, ex Nym. Consp. 425 = cinerea.
provincialis, Boiss. ex Lindem. in Bull. Soc. Nat. Mosc. iv. (1860) 11. 181 = jurineaefolia ?
pseudo-behen, *Boiss. Diagn.* Ser. I. vi. 126.—Cappadoc
pseudo-collina, *Schur, Enum. Pl. Transs.* 408.—Transylv.
pseudo-crupina, *Mutel, in Duch. Rev. Bot.* i. (1845-46) 400.—Morea.
pseudo-philostissa, Godr. in Mém. Acad. Stanislas, Sér. III. (xvii) 1853 (1854) 406 = dimorpha.
pseudo-phrygia, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. vi. (1848) 184 = phrygia.
pseudo-scabiosa, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 131 = C. Scabiosa.
pseudo-uniflora, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 85 = nervosa.
pallacantha, *Boiss. & Heldr. Diagn.* Ser. II. iii. 82.—Graecia.

CENTAUREA :—
pterocaula, *Trautv. in Act. Hort. Petrop.* ii. (1873) 552.—Armenia.
pterocaulos, *Pomel, Nouv. Mat. Fl. Atl.* 29.—Algeria.
pterocephala, Hort. ex DC. Prod. vi. 597 = trichascantha.
pterodonta, *Pomel, Nouv. Mat. Fl. Atl.* 28.—Algeria.
pterolopha, *DC. Prod.* vi. 572.—Hab. ?
pubescens, Willd. Sp. Pl. iii. 2292 = incana.
pubigera, Pers. Syn. ii. 488 = Volutarella muricata.
pulchella, *Ledeb. Ic. Pl. Alt.* t. t. 98.—Persia; Caucas.
pulcherrima, Wight, ex DC. Prod. vi. 579 = pulchra.
pulcherrima, *Willd. Sp. Pl.* iii. 2298.—Armenia; Caucas.
pulchra, *DC. Prod.* vi. 578.—Ind. or.
pullata, *Linn. Sp. Pl.* 911.—Europ. austr.
pumila, *Linn. Cent.* i. 30; *Amoen. Acad.* iv. 292.—Aegypt.; Graecia.
pumila, Sprun. ex Nym. Consp. 430 = hellenica.
pumila, Urv. in Mem. Soc. Linn. Par. i. (1822) 374 = raphanina.
Pumilio, Hill, Veg. Syst. iv. 36 = pumila ?
punctata, Dufour, in Ann. Sc. Nat. Sér. I. xxiii. (1831) 159 = hyssopifolia.
punctata, Vis. in Flora, xix. (1839) 28 = glaberrima.
pungens, *Pomel, Nouv. Mat. Fl. Atl.* 28.—Algeria.
pusilla, *Thunll. ex Pers. Syn.* ii. 482.—Aegypt.
pygmaea, *Benth. & Hook. f. Gen.* ii. 482.—Syria.
pygmaea, Chesn. ex DC. Prod. vi. 571 = amara.
pygmaea, Hoffm. Hort. Monac. (1808) n. 788 = depressa.
pyracantha, *Boiss. Fl. Orient.* iii. 685.—Tauria.
pyrenaica, Spreng. Fl. Hal. Mant. 53 = variegata.
pyrrholepha, *Boiss. Diagn.* Ser. II. iii. 66.—Armenia.
Rabenhorstiana, Sch. Bip. ex Nym. Consp. 426 = cristata.
Raddeana, *Trautv. in Act. Hort. Petrop.* ii. (1873) 475.—Armenia.
radiata, Linn. Syst. ed. XII. 656 = Serratula xeranthemoides.
ragusina, Clem. ex Boiss. Voy. Bot. Espagne, ii. 846 = Clementei.
ragusina, *Linn. Sp. Pl.* 912.—Dalmat. ; Creta.
ragusina, Sm. Fl. Graec. x. t. 908 = argentea.
ramosa, *Vukot.—Cf. Oestr. Bot. Zeitschr.* xxx. (1880) 168.—Europ.
ramotisinuma, Tausch, in Flora, xii. (1829) Erg. 45 = sitens.
raphanifolia, Salzm. ex DC Prod. vi. 591 = diluta.
raphanina, *Sibth. & Sm. Fl. Graec.* t. 917.—Creta.
raphanina, Sieber, ex Spreng. Syst. ed. II. i. 822 = nana.
raphanina, Sprun. ex Nym. Consp. 430 = hellenica.
raphanina, Urv. in Mem. Soc. Linn. Par. i. (1822) 373 = Urvillei.
recta, *Kroch. Fl. Sil. Suppl.* ii. 388.—Europ.
recurvata, C. Koch, in Linnaea, xxiv. (1851) 428 = cardaliformis.
reflexa, *Lam. Encyc.* i. 676.—Armenia ; Caucas.
regia, *Boiss. Diagn.* Ser. I. vi. 136.—Assyria.
Reichenbachii, *DC. Prod.* vi. 583.—Europ. ?
reichenbachioides, Schur, Enum. Pl. Transs. 409 = Reichenbachii.
repens, *Linn. Sp. Pl.* ed. II. 1328.—Orient.
resupinata, *Cass. Notes Pl. Crit.* 114.—Hispan.
reticulata, Bieb. ex Willd. Sp. Pl. iii. 2287 = coronopifolia.
Reuteri, Reichb. f. Ic. Fl. Germ. xv. 33. t. 760 = paniculata.
Reuteriana, *Boiss. Diagn.* Ser. I. iv. 16.—Graecia.
rhaetica, Moritz. in Denkschr. Allg. Schw. Ges. Natur. iii. (*Fl.* Graub. 31. t. 3) = phrygia.
rhapontica, Wulf. in Jacq. Coll. I. 238 = rhaponticum.
rhaponticoides, *Benth. & Hook. f. Gen.* ii. 479.—Hispan.
Rhaponticum, *Linn. Sp. Pl.* 915.—Europ.
rhenana, Bor. ex Nym. Consp. 426 = maculosa.
rhizantha, C. A. Mey. Verz. Pfl. Cauc. 84 = sessilis.
rhizantha, Tchihat. Asie Min. Bot. ii. 290 = armena.
rhizocathartica, *C. Koch, in Linnaea*, xxiv. (1851) 427. —Armenia.
rhizocephala, *Olivier & Hiern, in Oliver, Fl. Trop. Afr.* iii. 433.—Afr. trop.
rhizocephala, *Trautv. in Act. Hort. Petrop.* ii. (1873) 476.—Armenia.
rhulenica, Baumg. Enum. Stirp. Transs. iii. 70 = Serratula coronata.
rigens, *Lag. Gen. et Sp. Nov.* 32.—Hispan.

CENTAUREA :—
rigescens, *Hornem. Hort. Hafn.* ii. 859.—Hab. ?
rigida, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 268.—Syria.
rigida, Willd. Enum. Hort, Berol. Suppl. 61 = rigescens.
rigidifolia, Bess. ex DC. Prod. vi. 587 = orientalis.
rigidula, Jord. Obs. Pl. Crit. v. (1847) 69 = paniculata.
riparia, *A. Cunn. ex DC. Prod.* vi. 602.—N. Cambria.
rivularis, Brot. Fl. Lusit. i. 367 = nigra.
romana, *Linn. Sp. Pl.* 916.—Campania.
Ropalon, *Pomel, Nouv. Mat. Fl. Atl.* 25.—Algeria.
rosea, Stend. Nom. ed. i. 175 = trinervia.
rubescens, *Bess. ex DC. Prod.* vi. 567.—Tauria.
rufescens, Jord. Pugill. 107 = pectinata.
rufescens, Noul. ex Nym. Consp. 425, partim = nigra.
rumelica, *Boiss. Diagn.* Ser. II. iii. 78.—Rumelia.
rupestris, Kit. ex Steud. Nom. ed. II. i. 322 = dissecta.
rupestris, *Linn. Sp. Pl.* ed. II. 1298.—Austr. ; Ital.
rupicola, Bad. ex Nym. Consp. 426 = aplolepha.
rupicola, *Pomel, Nouv. Mat. Fl. Atl.* 31.—Algeria.
ruscinonensis, Boiss. Diagn. Ser. II. iii. 70 = C. Jacea ?
ruthenica, Baumg. ex Nym. Consp. 417 = *rhutenica*.
ruthenica, *Lam. Encyc.* i. 668.—Transylv. ; Caucas. ; Sibir.
rutifolia, Boiss. Fl. Orient. iii. 642 = pannosa.
rutifolia, *Sibth. & Sm. Prod. Fl. Graec.* ii. 205. t. 916. —Byzant.
sabulosa, Ledeb. ex Spreng. Syst. iii. 405 = diffusa.
Sadleriana, *Janka, in Termtsz. Füzetek*, ii. (1878) 142. —Europ.
Saharae, *Pomel, Nouv. Mat. Fl. Atl.* 30.—Afr. bor.
salonitana, Bess. ex Nym. Consp. 422 = austriaca.
salicifolia, Bieb. ex Willd. Sp. Pl. iii. 2288 = nigra.
salina, *Benth. & Hook. f. Gen.* ii. 479.—Sibir.
Salisana ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. — Univ. (1860) 110.
salmantica, *Linn. Sp. Pl.* 918.—Europ. ; Afr. bor.
salonitana, *Vis. in Flora*, xii. (1829) 1. *Erg.* 23.— Europ. or.
sanguinea, *Spreng. Pugill.* i. 60.—Hab. ?
saxatilis, Bertol. ex Nym. Consp. 429 = rupestris.
saxatilis, *C. Koch, in Linnaea*, xxiv. (1851) 419.— Creta.
saxicola, *Lag. Gen. et Sp. Nov.* 32.—Hispan.
Scabiosa, Asso, Syn. 126 = cephalariaefolia.
Scabiosa, *Linn. Sp. Pl.* 918.—Europ.
Scabiosa, Sadl. Fl. Pesth. ed. II. 409 = Sadleriana.
scabiosaeformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Scabiosa.
scabiosifolia, St. Lag. l. c. viii. (1881) 208 = praec.
scabra, Preul, Fl. Sicul. p. xxx = sicula.
schemnitzensis, Hort. ex Steud. Nom. ed. II. i. 322 = nigrescens.
Schimperi, *DC. Prod.* vi. 596.—Arab.
schizolepis, *Trautv. in Act. Hort. Petrop.* ii. (1873) 553.—Armenia.
Schmidtiana, *Tausch, in Flora*. xii. (1829) 1. *Erg.* 44. —Hab. ?
Schoenbeol, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 85 (*Pugill.* 186).—Hispan.
Schouwii, DC. Prod. vi. 598 = sicula.
Schwarzenbergiana, Schur, in Verh. Siebenb. Ver. Naturw. vi. (1855) x.—Rumelia.
sciophila, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 180 = nigrescens.
sclerolepis, *Boiss. Diagn.* Ser. I. vi. 136.—Mesopot.
scoparia, *Sieber, ex DC. Prod.* vi. 596.—Palaest. ; Aegypt.
Scopolii, Vest, in Flora, xii. (1829) 1. Erg. 81 = crithmifolia.
scopulorum, *Boiss. & Heldr. Diagn.* Ser. I. x. 115.— Graecia.
scorzonerifolia, Dufour, in Ann. Sc. Nat. xxiii. (1831) 162 = aspera.
segetalis, Salisb. Prod. 207 = C. Cyanus.
semidecurrens, Jord. Obs. Pl. Crit. v. (1847) 52 = axillaris.
sempervirens, *Linn. Sp. Pl.* 913.—Lusit. ; Ital.
senegalensis, *DC. Prod.* vi. 598.—Afr. trop.
Sertilis, *Linn. Sp. Pl.* 915.—Europ. austr.
Seridis, Lolsel. Fl. Gall. ed. II. ii. 212 = aspera.
serotina, Bor. Fl. Cent. Fr. ed. III. ii. 350 = nigrescens.
serrata, Kitt. Taschenb. ed. II. 668 = montana.
serratuloides, *Benth. & Hook. f. Gen.* ii. 479.—As. Min.

CENTAUREA :—
serratuloides, Georgi, Bemerk. Reise Russ. Reich. i. 231 = salina.
serratuloides, Krock. Fl. Sil. ii. 11. 484 (deleuda).—Silesia.
sessiliflora, Lam. Fl. Fr. ii. 35 = melitensis.
sessiliflora, Linn. Sp. Pl. 913 = nigra.
sessilifolia, Hill, Veg. Syst. iv. 32 = praec. ?
sessilis, *Willd. Sp. Pl.* iii. 2300.—Armenia.
setifera, Lag. Gen. et Sp. Nov. 32 = sitena.
setosa, *Cav. in Anal. Ciencias Nat.* iv. (1801) 96.— Marocco.
sexsana, Benth. Cat. Pl. Pyr. 68 = montana.
sexsana, Chaix, in Vill. Hist. Pl. Dauph. iii. 52 = variegata, axillaris.
sibirica, Bieb. Fl. Taur. Cauc. ii. 348 = Marschalliana.
sibirica, *Linn. Sp. Pl.* 918.—Sibiria.
sibirica, Willd. ex Ledeb. Fl. Ross. ii. 695 = declinata.
sicula, Lam. ex Willk. & Lange, Prod. Fl. Hisp. ii. 145 = alaceensis.
sicula, Leyss. ex DC. Prod. vi. 594 = solstitialis.
sicula, *Linn. Sp. Pl.* 918.—Sicilia.
simplex, *Cav. in Anal. Ciencias Nat.* iv. (1801) 95.— Afr. bor.
simplex, Murr. ex Steud. Nom. ed. II. i. 322 = C. Jacea.
simplicicaulis, *Boiss. & Huet, Diagn.* Ser. II. iii. 67.— Armenia.
sinaica, *DC. Prod.* vi. 592.—Arab.
singarensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 654.—Assyria.
Smyrnaea, Boiss. Diagn. Ser. I. iv. 20, in textu = Urvillei.
solstitialis, Asso, Syn. 127 = melitensis.
solstitialis, *Linn. Sp. Pl.* 917.—Europ.
somalensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 438.—Afr. trop.
sonchifolia, Boiss. Voy. Bot. Espagne, ii. 352 = malacitana.
sonchifolia, *Linn. Sp. Pl.* 915.—Europ. austr.
sonchifolia, Welw. ex Nym. Consp. 422 = sphaerocephala.
sordida, Graf, ex DC. Prod. vi. 587 = Grafiana.
sordida, Salisb. Prod. 207 = sempervirens.
sordida, *Willd. Sp. Pl.* iii. 2283.—Austria.
Spachii, *Sch. Bip. ex Willk. & Lange, Prod. Fl. Hisp.* ii. 154.—Hispan.
Sparmanni, *DC. Prod.* vi. 598.—Africa
spathulata, Tenore, Prod. Fl. Nap. p. ii = nigra ?
spathulata, Zersph. Fl. Melit. i. 11 = crassifolia.
speciosa, *Boiss. Diagn.* Ser. I. x. 116.—Palaestina.
speciosa, Mor. ex Nym. Consp. 428 = filiformis.
sphaerocephala, J. Gay, ex Dur. in Duch. Rev. Bot. ii. (1846–47) 429 = Fontanesii.
sphaerocephala, *Linn. Sp. Pl.* 916.—Europ. med.
sphaerocephala, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 581 = fragilis.
spicata, *Boiss. Diagn.* Ser. I. x. 118.—Syria.
Spina-badia, Bubani, ex Timb. in Mém. Acad. Toulouse, Sér. VIII. i. ii. (1879) 187 = coerulescens.
spinosa, Griseb. Spicil. Fl. Rumel ii. 341 = diffusa.
spinosa, *Linn. Sp. Pl.* 912.—Creta ; Oriens.
spinulosa, Rochel, Pl. Ban. Rar. 76. t. 36 = C. Scabiosa.
splendens, *Linn. Sp. Pl.* 914.—Hispan. ; Ital.
splendens, Tenore, Syll. Add. iii. 298 = alba.
Spruneri, *Boiss. & Heldr. Diagn.* Ser. I. vi. 122.— Graecia.
Sprunneri, Boiss. & Orph. ex Boiss. Fl. Orient. iii. 662 = Guiccciardii.
spuria ×, *Kern. in Oestr. Bot. Zeitschr.* xxii. (1872) 51.—Austria.
squarrosa, Roth, Catalecta, fasc. ii. 118 = virgata.
stenolepis, Kern. in Oestr. Bot. Zeitschr. xxii. (1872) 49 = austriaca.
stenophylla, Dufour, in Ann. Sc. Nat. Sér. I. xxiii. (1831) 169 = aspera.
stenophylla, *Walp. Ann.* i. 446.—Persia.
stenophylla, *Bess. Enum. Pl. Volh.* 88.—Podol.
stenophylla, Griseb. Spicil. Fl. Rumel. ii. 337 = C. Scabiosa.
sterilis, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) ii. 390.—Tauria.
Stevani, *Bieb. Fl. Taur. Cauc.* ii. 356.—Reg. Caucas.
Stevenii, Kotschy, ex Boiss. Fl. Orient. iii. 693 = cheiracantha.
Stoebe, Ledeb. Fl. Alt. iv. 40 = calva.
Stoebe, *Linn. Sp. Pl.* 913.—Austria ; Gallia.

CENTAUREA:—
Stoebe, Tausch, in Flora, xii. (1829) 1, Exg. 42 = diffusa.
Stoebe, Tenore, Fl. Nap. ii. 264 = subtilis.
stolenisii, Pand. ex Nym. Consp. 428 = orientalis.
straminea, Willd. Sp. Pl. iii. 2312 = glomerata.
strepens, Hoffmgg. & Link, Fl. Port. ii. 326 = splendens †
stricta, Waldst. & Kit. Pl. Rar. Hung. ii. 194 = axillaris.
strictissima, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 180 = Serratula coriacea.
strobilacea, Scop. Delic. Insub. i. 88 = alba.
suaveolens, Willd. Sp. Pl. iii. 2279 = moschata.
subacaulis, Ledeb. Fl. Ross. ii. 709 = Urvillei.
subalbida, Jord. Pugill. 109 = maculosa.
subalpina, Schur, Enum. Pl. Transs. 407 = atropurpurea.
subcillaris, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iii. 627.—Cephalon.
sublanata, *Boiss. i.* c. 645.—Graecia.
sublanuginosa, Schur, Enum. Pl. Transs. 899, Index = nervosa.
subspinosa, Steud. Nom. ed. II. i. 322 = diluta.
subtilis, *Bertol. Fl. Ital.* ix. 451.—Ital.
sulphurea, Lag. Gen. et Sp. Nov. 32 = dissecta, orientalis.
sulphurea, *Willd. Enum. Hort. Berol.* 930.—Sicilia.
sumensis, *Kalrm. in Bull. Acad. Soc. Nat. Mosc.* xviii. (1845) i. 238.—Reg. Caucas.
supina, Jord. Pugill. 108 = pectinata.
sylvatica, Pourr. in Mém. Acad. Toul. iii. (1788) 310 = C. Scabiosa.
Szovitsiana, *Boiss. Fl. Orient.* iii. 688.—Armenia; Anatol.
tagana, *Iirot. Fl. Lusit.* i. 369.—Lusit.; Afr. bor.
tarascaefolia, Boiss. in Tchihat, Asie Min. Bot. ii. 321 = rhizocalathium.
tarascaefolia, D. Don, Prod. Fl. Nep. 167 = Saussurea eriostemon.
tatarica, Linn. f. Suppl. 388 = orientalis.
tatarica, Pall. Ind. Taur. 58 = centauroides.
taurometana, *Guss. Fl. Sicil. Syn.* ii. 512.—Sicil.
Tauscheri, Kern. in Oestr. Bot. Zeitschr. xxii. (1872) 119 = arenaria.
Tchihatcheffi, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. (1854) i. 31.—Lycaonia.
Teidis, Steud. Nom. ed. II. i. 323 = arguta.
tenella, Hort. ex Spreng. Syst. iii. 401 = salmantica.
Tenoreana, Willd. ex Nym. Consp. 420 = incana.
tenuiflora, *DC. Prod.* vi. 584.—Tauria.
tenuiflora, Fisch. Hort. Gorenk. ed. II. (1812) 36 = salmantica.
tenuifolia, *Dufour, in Ann. Sc. Gén. Phys.* ii. (1820) 303.—Hispan.
tenuifolia, Hort. ex Steud. Nom. ed. II. i. 323 = salmantica.
tenuifolia, Schleich. ex Steud. l. c. = C. Scabiosa.
tenuifolia, Sallsb. Prod. 307 = Crupina vulgaris.
tenuiramea, *Delile, ex Herd. in Ind. Sem. Hort. Petrop.* (1860) 37.—Hab. ?
tenuisecta, Jord. Pugill. 110 = maculosa.
Teydis, C. Sm. ex Buch, in Berl. Acad. Abh. 1816–17 (1819) 369 = arguta.
thiantha, [Spreng.] Bot. Gart. Halle, 24 = orientalis.
Thirkei, *Sch. Bip. in Linnaea*, xix. (1847) 814.—Phrygia.
Thomasiana, *Dalla Torre, Atl. Alpenfl.* 141.—Europ.
thracicaefolia, *DC. Prod.* vi. 590.—Cappadoc.
Timbali, Martr. Fl. Crit. Tarn, 383 = amara.
tinctoria, [Clairv.] Man. Herb. 258 = Serratula tinctoris.
tingitana, Linn. Sp. Pl. ed. II. 1300 = Carduncellus caeruleus.
toletana, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 18.—Hispan.
tomentosa, Sallsb. Prod. 307 = rupestris.
Torreana, Tenore, Ind. Sem. Hort. Neap. (1839) 15 = C. Calcitrapa.
tougourensis, *Boiss. & Reut. Diagn.* Ser. II. iii. 76.—Algeria.
Tournefortii, *Walp. Ann.* i. 448.—Armenia.
transalpina, Schleich. Cat. Pl. Helv. ed. III. 11 = nigrescens.
trapezuntina, Boiss. Fl. Orient. 680 = nigrescens.
trichacantha, Willd. ex Spreng. Syst. iii. 404 = C. Calcitrapa.
trichocephala, *Bieb. ex Willd. Sp. Pl.* iii. 2286.—Tauria.

CENTAUREA:—
trifurcata, *Pomel, Nouv. Mat. Fl. Atl.* 27.—Afr. bor.
trinervia, *Stev. ex Willd. Sp. Pl.* iii. 2301.—Tauria; Canc.
trinitaefolia, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 97 = maculosa.
Trinyfetii, All. Fl. Pedem. i. 168 = variegata.
tuberosa, Vis. Fl. Dalmat. ii. 33. t. 12. f. 2 = napulifera.
tubulosa, Chab. ex Nym. Consp. 421 = nigrescens.
turkestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 325.—Turkestan.
Tweedieii, *Hook. & Arn. in Hook Comp. Bot. Mag.* i. (1836) 110.—Bonaria.
ucranica, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 497, nomen.—Ross. austr.
uliginosa, *Iirot. Fl. Lusit.* i. 368.—Lusitan.
umbrosa, *Hurt, ex Rend. Ind. Sem. Hort. Genev.* (1856) 4.—*Cf. Linnaea*, xxix. (1857–58) 719.—Sicilia.
uniflora, *Linn. Mant* i. 118.—Europ. austr.
uniflora, Pall. Reise, iii. 253 = Serratula glauca.
Urvillei, *DC. Prod.* vi. 592.—Ins. Samos.
ustulata, Bess. Enum. Pl. Volh. 108 = Biebersteinii.
vatulata, *DC. Prod.* vi. 576.—Persia.
vallesiaca, Jord. Pugill. 111 = paniculata.
vallesiaca, *Reichb. f. Ic. Fl. Germ.* xv. 33. t. 783.—Europ.
variabilis, Bartl. ex Steud. Nom. ed. II. i. 323 = rupestris.
variana, *A Rich. Tent. Fl. Abyss.* i. 453.—Abyss.
variegata, *Lam. Encyc.* i. 666.—Europ. austr.
varifolia, Loisel. Not. 130 = C. Scabiosa.
venetum, J. F. Gmel. Syst. 1267, sphalm. = C. Verutum.
verbascifolia, *Vahl, Symb. Bot.* i. 75.—Arab.
Veruum, *Linn. Cent.* i. 30; *Amoen. Acad.* iv. 292.—As. Min.
vesceritensa, *Boss. Diagn.* Ser. II. iii. 70.—Algeria.
Villarsii, *Mutel, Fl. Dauph.* ii. 254.—Gallia.
viminea, Less. in Linnaea, vi. (1831) 83 = pulchella.
viretorum, Jord. ex Nym. Consp. 421 = C. Jacea.
virgata, Bess. Enum. Pl. Volh. 85 = Besseriana.
virgata, Lav. Ic. ii. 16 = incerticeps.
virgata, *Lam. Encyc.* i. 670.—Armenia.
virgata, Portenschl. ex Nym. Consp. 426 = glaberrima.
vochinensis, Bernh. ex Reichb. Ic. Fl. Germ. xv. 15 = nigrescens.
vulgaris, Pers. Syn. ii. 488 = Crupina vulgaris.
vulnerariaefolia, *Pomel, Nouv. Mat. Fl. Atl.* 26.—Afr. bor.
Wallichiana, *Less. in Linnaea* vi. (1831) 84.—Ind. or.
Webbiana, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 258.—Ins. Canar.
Weldeniana, Reichb. Ic. Fl. Germ. xv. 12. t. 753 = amara.
Wiedemanniana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1836) 32.—Anatolia.
Willkommii, *Sch. Bip. ex Willk in Linnaea*, xxv. (1852) 86.—Hispan.
wolgensis, Dc. Prod. vi. 581 = arenaria.
xanthina, Boiss. & Heldr. in Boiss. Fl. Orient. iii. 650 = Pella.
xanthina, Spreng. Syst. iii. 407 = fuscata.
xanthocephala, *Benth. & Hook. f. Gen.* ii. 480.—Armen.
Zanonii, Sebast. & Maur. Fl. Rom. 398 = romana.
Zuccarinana, *DC. Prod.* vi. 574.—Graecia.

CENTAURELLA, Delarb. Fl. Auv. ed. II. 28 (1800) = **Erythraea**, Borckh. (Gentian.).
dichotoma, Delarb. l. c. 99 = E. Centaurium.

CENTAURELLA, Michx. Fl. Bor. Am. i. 97. t. 19 (1803) = **Bartonia**, Muhl. (Gentian.).
aestivalis, Pursh, Fl. Am. Sept. i. 100 = B. verna.
autumnalis, Pursh, l. c. = B. tenella.
Mooriana, Steud. & Hochst. in Grisseb. Gen. et Sp. Gentian. 308 = B. tenella.
paniculata, *Michx. Fl. Bor. Am.* i. 98. t. 12. f. 1 = B. tenella.
verna, Michx. l. c. 98 = B. verna.
vernalis, Pursh, Fl. Am. Sept. i. 99 = B. verna.

CENTAURIUM, Rupp. Fl. Jen. ed. Hall. 22 (1745) = **Erythraea**, Borckh. (Gentian.).

CENTAURIA, Linn. Gen. ed. I. 268 (1737) = **Centaurea**, Linn. (Compos.).

CENTAURIDIUM, Torr. & Gray, Fl. N. Am. ii. 246 (1841) = **Xanthisma**, DC. (Compos.).
Drummondii, Torr. & Gray. l. c. = X. texanum.

CENTAURIUM, Borckh. in Roem. Arch. i. (1796) 29 = **Cnicoseris**, Less. (Gentian.).
malabaricum, Borckh. l. c. = C. sessiliflora.

CENTAURIUM, Gilib. Fl. Lituan. i. 35 (1781); Moench, Meth. 449 (1794) = **Erythraea**, Borkh. (Gentian.).
Erythraea, Rafn, Danm. og Holst. Fl. ii. 75 = E. Centaurium.
inapertum, Rafn, l. c. 77.—Europ.
minus, Moench, Meth. 449 = E. Centaurium.
umbellatum, Gilib. Fl. Lituan. i. 35 = E. Centaurium.
vulgare, Rafn, Danm. og Holst. Fl. ii. 78 = E. Centaurium.

CENTAURIUM, Pers. Syn. i. 137 (1805) = CENTAURELLA, Michx. = **Bartonia**, Muhl. (Gentian.).
autumnale, Pers. l. c. = B. tenella.
vernum, Pers. l. c. = B. verna.

CENTAURIUM, Tourn. ex Linn. Syst. ed. I (1735) = **Centaurea**, Linn. (Compos.).
africanum, Cass. in Dict. Sc. Nat. lviii. 10 = Centaurea africana.
alpina, Pomel, Nouv. Mat. Fl Atl. 32.—Algeria.
moschatum, Cass. in Dict. Sc. Nat. vii. 378 = Centaurea moschata.
officinale, Cass. l. c. = Centaurea Centaurium.
ruthenicum, Cass. l. c. lviii. 10 = Centaurea rhutenica.
suaveolens, Cass. l. c. vii. 379 = Centaurea moschata.
taganum, Pomel, Nouv. Mat. Fl. Atl. 32.—Algeria.
tatarica, Pomel, l. c.—Algeria.

CENTAUROPSIS, Boj. ex DC. Prod. v 93 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 231.
fruticosa, *Boj. l. c.*—Madag.
lanuginosa, *Boj. l. c.*—Madag.
Rutenbergiana, *Vatke, in Bremen Abh.* ix. (1885) 119.—Madag.

CENTELLA, Linn. Pl. Rar. Afr. 28 (1760); Amoen. Acad. vi. 112 (1764) = **Hydrocotyle**, Linn. (Umbellif.).
asiatica, Urban, in Mart. Fl. Bras. xi. i. 287 = H. asiatica.
cunnifolia, F. Muell. in Hook. Kew Journ. vii. (1855) 278. t. 13 = Ascrilla cunnifolia.
glabra, DC. Prod. iv. 69, sphalm. = seq.
glabrata, Linn. Pl. Rar. Afr. 28; Amoen. Acad. vi. 112 = H. Centella.
rensifolia, Urban, in Mart. Fl. Bras. xi. i. 286, in obs. = Micropleura rensfolia.
villosa, Linn. Pl. Rar. Afr. 28; Amoen. Acad. vi. 112 = H. villosa.

CENTEMA, Hook. f. in Benth. & Hook. f. Gen. iii. 31 (1880). *AMARANTACEAE*, Benth. & Hook. f. l. c.
angolensis, *Hook. f. l. c.*—Afr. trop.
Kirkii, *Hook. f. l. c.*—Afr. trop.

CENTHRISCUS, Spreng. ex Steud. Nom. ed. I. 176, 55 (1821) = **Anthriscus**, Bernh. (Umbellif.).
fumarioides, Spreng. l. c. = A. fumarioides.
nemorosa, Spreng. l. c. = A. nemorosa.

CENTINODIA, Reichb. Handb. 236 (1837) = **Polygonella**, Tourn.
agrestina, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 146 = P. aviculare.
arenaria, Fourr. l. c. = P. arenarium.
aviculare, Fourr. l. c. = P. aviculare.
Bellardi, Fourr. l. c. = P. Bellardi.
humifusa, Fourr. l. c. = P. aviculare.
maritima, Fourr. l. c. = P. maritimum.
microperma, Fourr. l. c. = P. aviculare.
ruriveaga, Fourr. l. c. = P. aviculare.

CENTIPEDA, Lour. Fl. Cochinch. 492 (1790). *COMPOSITAE*, Benth. & Hook. f. ii. 480.
MYRIOGYNE, Less. in Linnaea, vi. (1831) 219.
cepensis, Lam. Syn. Comp. 201 = Dichrocephala latifolia.
Cunninghami, *A. Br. & Aschers. Ind. Sem. Hort. Berol.* (1867) *App.* 6 ; *P. Muell. Fragm.* viii. 148.—Austral.
minima, A. Br. & Aschers. l. c. = orbicularis.

CENTIPEDA :—
minuta, C. B. Clarke, Compos. Ind. 151 = orbicularis.
orbicularis, C. B. Clarke, l. c. = Sphaerormphaea Russeliana.
orbicularis, *Lour. Fl. Cochinch.* ii. 493.—As. et Austral. trop. ; Ins. Pacif.
racemosa, *F. Muell. Census*, 84.—Austral.
thespidioides, *F. Muell. Fragm.* viii. 149.—Austral.

CENTOPODIUM, Burch. Trav. i. 340 (1824) = **Emex**, Neck. (Polygon.).
spinosum, Burch. l. c. = E. spinosa.

CENTOTHECA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 (Centostoca). *GRAMINEAE*, Benth. & Hook. f. iii. 1190.
lappacea, *Desv. l. c.*—Geront. trop.
latifolia, Trin. Fund. Agrost 141 = lappacea.
parviflora, *Anderss. in Peters, Reise Mossamb. Bot.* 569.—Afr. trop.

CENTRACHAENA, Less. Syn. Comp. 255 (1832) = seq.

CENTRACHENA, Schott, Cat. Ind. Sem. Hort. Vindob. (1828) ; ex Reichb. ic. Pl. Crit. iv. 47 (1827) = **Chrysanthemum**, Tourn. (Compos.).
viscida, Schott, l. c. ; ex Reichb. l. c. = Chrys. viscosum.

CENTRADENIA, G. Don, Gen. Syst. ii. 765 (1832). *MELASTOMACEAE*, Benth. & Hook. f. i. 789.
DONCKLAERIA, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1350 (1855).
PLAGIOPHYLLUM, Schlecht. in Linnaea, xiii. (1839) 429.
divaricata, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 354.—Am. centr.
floribunda, *Planch. in Fl. des Serres*, Sér. I. v. (1849) t. 458.—Mexic.
grandifolia, *Bndl. ex Walp. Rep.* ii. 119.—Mexic.
inaequilateralis, *G. Don, Gen. Syst.* ii. 765.—Mexic.
Oerstediana, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 354.—Am. centr.
ovata, *Klotzsch, l. c.*—Am. centr.
rosea, Lindl. Bot. Reg. (1848) t. 20 = inaequilaterals.

CENTRANDRA, Karst. in Linnaea, xxviii. (1856) 440 ; = **Julocroton**, Mart. (Euphorbiac.).
hondensis, Karst. l. c. = J. hondensis.

CENTRANTHERA, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 293 = **Pleurothallis**, R. Br. (Orchid.).
punctata, Scheidw. l. c. = P. Centrantherа.

CENTRANTHERA, R. Br. Prod. 338 (1810). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 969.
GUMTROLIS, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 87 (1825).
PURSMIA, Dennst. Schluess. Hort. Malab. 35 (1818).
RAZUMOVIA, Spreng. Syst. ii. 682 (1825).
Brunoniana, Thw. Enum. Pl. Zeyl. 220 = procumbens.
Brunoniana, *Wall. Cat.* n. 8882 ; *Benth. Scroph. Ind.* 50.—Reg. Himal. ; Burma.
grandiflora, *Benth. in Wall. Cat.* n. 8880 ; *Scroph. Ind.* 50.—Reg. Himal.
hispida, Benth. Fl. Hongk. 254 = Brunoniana.
hispida, *R. Br. Prod.* 438.—Austral. ; Afghan.
humifusa, *Wall. Cat.* n. 8883.—Ind. or.
nepalensis, D. Don, Prod. Fl. Nep. 88 = hispida.
procumbens, *Benth. in DC. Prod.* x. 525.—Ind. or.
prostrata, Boj. Hort. Maurit. 246 = Raclamaea prostrata.

CENTRANTHUS, DC. Fl. Fr. iv. 288 (1805) *VALERIANEAE*, Benth. & Hook. f. ii. 153.
HYBIDIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 899.
KENTRANTHUS, Neck. Elem. i. 122 (1790), nomen prius.
angustifolius, Bieb. Fl. Taur. Cauc. i. 67 = longiflorus.
angustifolius, Boiss. Voy. Bot. Espagne, 291 = nevadensis.
angustifolius, *DC. Fl. Fr.* iv. 289.—Europ. austr.
Calcitrapa, *Dufr. Hist. Valér.* 39 ; *DC. l. c.* v. 492.—Europ.
Clusonia, *Pomel, Nouv. Mat. Fl. Atl.* 78.—Afr. bor.
dasycarpus, Kunze, Del. Sem. Hort. Lips. (1847) ; et Linnaea, xxiv. (1851) 176 = macrosiphon.
elatus, Boiss. & Heldr. Diagn. Ser. I. x. 73 = longiflorus.
juncens, *Boiss. & Heldr. l. c.*—Graecia.

CENTRANTHUS:—
latifolius, Duf. Hist. Valér. 38 = ruber.
Lecoqii, Jord. Pugill. Pl. Nov. 76 = angustifolius.
longiflorus, *Stev. Obs. Pl. Ross.* 76.—As. Min. ; Syria ; Reg. Cauca.
macrosiphon, *Boiss. Diagn.* Ser. I. iii. 57.—Hispan.
marinus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 479 = ruber.
maritimus, DC. Prod. iv. 682 = ruber.
nervosus, *Moris, Elench. Sard.* ii. 4.—Corsic. ; Sardin.
nevadensis, *Boiss. Diagn.* Ser. II. ii. 130.—Hispan.
orbiculatus, Duf. Hist. Valér. 39 = C. Calcitrapa.
orbiculatus, *Rouy & Schuit. Syst.* i. 85.—Cyprus.
pinnatifidus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 123 = C. Calcitrapa.
ruber, *DC. Fl. Fr.* iv. 289.—Europ. ; Syria.
Stibthorpii, *Heldr. & Sart. Diagn.* Ser. II. ii. 119.—Graecia.

CENTRAPALUS, Cass. in Dict. Sc. Nat. vii. 382 (1817) = **Vernonia**, Schreb. (Compos.).
galamensis, Cass. l. c. 383 = V. pauciflora.

CENTRATHERUM, Cass. in Bull. Soc. Philom. (1817) 31. *COMPOSITAE*, Benth. & Hook. f. ii. 225.
ANPHERESPHIS, H. B. & K. Nov. Gen. et Sp. iv. 91. t. 314, 315 (1820).
AMPHISBCIS, Schrank, in Syll. Ratisb. i. (1824) 66.
CRANTZIA, Vell. Fl. Flum. Ic. viii. t. 158 (1827).
DECANEURUM, DC. in Guill. Arch. Bot. ii. (1838) 516.
PHYLLOCEPHALUM, Blume, Bijdr. 888 (1826).
ROLFINKIA, Zetsk. Pl. Ind. 13. t. 14 (1837).
SPIXIA, Schrank, Pl. Rar. Hort. Monac. t. 80 (1819).
WIGHTIA, Spreng. ex DC. Prod. v. 67 (1836)
aristata, Cass. in Dict. Sc. Nat. lvii. 342 = punctatum.
brachylepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 12.—Bras.
brevispinum, Cass. in Dict. Sc. Nat. lvii. 346 = intermedium.
chinense, Less. in Linnaea, iv. (1829) 320 = Vernonia chinensis.
coarctatum, *Benth. & Hook. f. Gen.* ii. 225.—Ind. or.
courtallense, C. B. Clarke, Comp. Ind. 8 = Kitchiei.
Hookeri, *C. B. Clarke, l. c.*—Ind. or.
intermedium, *Less. in Linnaea*, iv. (1829) 320.—Bras.
longipinnum, Cass. in Dict. Sc. Nat. lvii. 346 = punctatum.
Metzianum, *Hook. f. Fl. Brit. Ind.* iii. 298.—Ind. or.
molle, *Benth. & Hook. f. Gen.* ii. 225.—Ind. or.
muticum, *Less. in Linnaea*, iv. (1829) 320.—Mexic.
phyllolaenum, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 298.—Ind. or.
pulchellum, *Steud. Nom.* ed. II. i. 324.—N. Zel.
punctatum, *Cass. in Dict. Sc. Nat.* vii. 384.—Venezuela.
reticulatum, *Benth. & Hook. f. Gen.* ii. 225.—Ind. or.
Ritchiei, *Hook. f. Fl. Brit. Ind.* iii. 298.—Ind. or.
scariosum, C. B. Clarke, Comp. Ind. 4 = Vernonia scariosa.
tenue, *C. B. Clarke, l. c.*—Ind. or.

CENTROCARPHA, D. Don, in Sweet, Hort. Brit. ed. II. 310 (1840) ; et in Brit. Fl. Gard. Ser. II. iv. t. 87 (1831) = **Rudbeckia**, Linn. (Compos.).
acutifolia, Sweet, Hort. Brit. ed. II. 310, nomen = R. Newmanni.
aristata, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. sub t. 87 = R. triloba.
chrysantha, Steud. Nom. ed. II. i. 351 = R. Newmanni.
chrysauchen, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. sub t. 87 = R. fulgida.
discolor, D. Don, l. c. = R. fulgida ?
fulgida, D. Don, l. c. = R. fulgida.
gracilis, D. Don, in Sweet, Hort. Brit. ed. II. 310 ; et l. c. = R. hirta.
grandiflora, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. t. 87 = R. grandiflora.
hirta, D. Don, in Sweet, Hort. Brit. ed. II. 310 ; et l. c. = R. hirta.
moschata, J. Cree, ex Steud. Nom. ed. II. i. 351 = R. moschata.
Radula, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. sub t. 87 = Helianthus Radula.
spathulata, D. Don, l. c. = R. spathulata.
subtomentosa, Sweet, Hort. Brit. ed. I. 310, nomen = R. subtomentosa.
triloba, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. sub t. 87 = R. subtomentosa.
triloba, Lond. Hort. Brit. Suppl. i. 564 = R. triloba.

CENTROCHILUS, J. C. Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (Pl. Meyen.) 436. t. 118 (1843) = **Habenaria**, Willd. (Orchid.).
gracilis, Schau. l. c. = H. tipuloides.

CENTROCLINIUM, D. Don, in Trans. Linn. Soc. xvi. (1830) 254 = **Onoseris**, DC. (Compos.).
adpressum, Hook. Bot. Mag. t. 3115 = O. adpressa.
albicans, D. Don, in Trans. Linn. Soc. xvi. (1830) 254 = O. integrifolia.
altissimum, Poepp. & Endl. Nov. Gen. et Sp. iii. 52. t. 259.—Peruv.
hyssopifolium, D. Dietr. Syn. Pl. iv. 1565 = O. hyssopifolia.
reflexum, Hook. Bot. Mag. t. 3115 = O. reflexa.
salicifolium, D. Dietr. Syn. Pl. iv. 1565 = O. salicifolia.

CENTRODISCUS, Muell. Arg. in Mart. Fl. Bras. xi. 21. 295. t. 102 (1874) = **Caryodendron**, Karst. (Euphorb.).
grandifolius, Muell. Arg. l. c. 327.—Bras.

CENTROGLOSSA, Rodrig. Orch. Nov. ii. 234 (1881). *ORCHIDEAE*.
macrocera, *Rodrig. l. c.* 235.—Bras.
tripollinica, *Rodrig. l. c.*—Bras.

CENTROLEPIS, Labill. Nov. Holl. Pl. i. 7. t. 1 (1804). *CENTROLEPIDEAE*, Benth. & Hook. f. iii. 1026.
ALEPYRUM, R. Br. Prod. 258 (1810).
CENTROSEPIS, R. Hedw. Gen. fil. (1806).
DEVAUXIA, R. Br. Prod. 253 (1810) ed. Nees, 108 (1827).
aemula, Rudge, in Trans. Linn. Soc. x. (1811) 264 = strigosa.
aleppyroides, *Walp. Ann.* i. 897.—Austral.
aristata, *Rouy. & Schuit. Syst.* i. 44.—Australi.
Banksii, *Rouy & Schuit. l. c.*—Australi.
brevifolia, Walp. Ann. i. 896 = Drummondii.
cambodiana, *Hance, in Journ. Bot.* xiv. (1876) 14.—Cambodia.
cuspidigera, Rudge, in Trans. Linn. Soc. x. (1811) 258 = fasciculata.
Drummondii, *Walp. Ann.* i. 896.—Austral.
exserta, *Rouy. & Schuit. Syst.* i. 44.—Australi.
fasciculata, *Labill. Nov. Holl. Pl.* i. 7. t. 1.—Austral.
glabra, *Hieron. in Abh. Naturf. Ges. Halle*, xii. (1873) 209.—Australi.
humillima, *F. Muell. ex Benth. Fl. Austral.* vii. 203.—Austral.
longifolia, *Steud. Nom.* ed. II. i. 324 ; *et Kunth, Enum. Pl.* iii. 489.—Austral.
monogyna, *Benth. Fl. Austral.* vii. 205.—Austral.
muscoides, *Hieron. in Abh. Naturf. Ges. Halle*, xii. (1873) 209.—Tasmania.
muticа, *Hieron. l. c.* 211.—Austral.
Petersoni, Roem. & Schult. Syst. i. 43 = strigosa.
pilosa, *Hieron. in Abh. Naturf. Ges. Halle*, xii. (1873) 210.—Austral.
polygyna, Hieron. Centrolep. 96 = polygyna.
polygyna, *Hieron. in Abh. Naturf. Ges. Halle*, xii. (1873) 210.—Australi.
pulchra, Hieron. l. c. 213 = Drummondii.
pulvinata, *Rouy. & Schuit. Syst.* i. 43.—Tasmania.
pusilla, *Rouy. & Schuit. l. c.* 44.—Austral.
strigosa, *Rouy. & Schuit. l. c.* 43.—Austral.
tenuior, Roem. & Schult. l. c. = strigosa.
Urvillei, Hieron. in Abh. Naturf. Ges. Halle, xii. (1873) 214 = Drummondii.

CENTROLOBIUM, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 96. *LEGUMINOSAE*, Benth. & Hook. f. i. 546.
minus, Pesl, Bot. Bemerk. 68 = robustum.
paraënse, *Tul. in Arch. Mus. Par.* iv. (1844) 87.—Am. austr.
robustum, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 96.—Bras.
tomentosum, *Guill. ex Benth. in Hook. Journ. Bot.* ii. (1840) 66.—Bras.

CENTRONIA, Blume, Bijdr. 774 (1826) = **Anginetia**, Linn. (Orobanchac.).
cirrhifolia, Blume, l. c. = A. Centronia.

CENTRONIA, D. Don, in Mem. Wern. Soc. iv. (1823) 314. *MELASTOMACEAE*, Benth. & Hook. f. i. 750.
BRACHYCENTRUM, Meisn. Gen. 114 (1837).
CALYPTRARIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 182.
STEPHANOGASTRA, Karst. & Triana, in Linnaea, xxviii. (1856) 426.
brachycera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 72.—N. Granat.
crasiramea, *Triana, l. c.*—Guiana.
excelsa, *Triana, l. c.*—Peru. ; Ecuador.
eximia, *Triana, l. c.*—Ecuador.
haemantha, *Triana, l. c.*—N. Granat.
insignis, *Triana, l. c.*—Ecuador.
laurifolia, *D. Don, in Mem. Wern. Soc.* iv. (1823) 315. —Peru.
Mutisii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 73.—N. Granat.
phlomoides, *Triana, l. c.*—Costa Rica.
reticulata, *Triana, l. c.* 71.—Peru.

CENTRONOTA, DC. in Meisn. Gen. 303 [213] (1840) ; et Prod. ix. 286 (1843) = **Aeginetia**, Linn. (Orobanchac.).
mirabilis, DC. Prod. ix. 286 = An Centronia.

CENTROPAPPUS, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 124 = **Senecio**, Tourn. (Compos.).
Brunonis, Hook. f. i. c. = S. Centropappus.

CENTROPETALUM, Lindl. Sert. Orch. sub t. 21 (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 570.
FERNANDEZIA, Ruiz & Pav. Prod. Fl. Per. 123. t. 27 (1794).
NASONIA, Lindl. in Benth. Pl. Hartw. 150 (1844).
distichum, *Lindl. l. c.*—Peru.
Myrtillus, *Pfitzer, Verg. Morph. Orch.* 16.—Peru.
Warszewiczii, *Reichb. f. in Bot. Zeit.* x. (1852) 706.—Peru.

CENTROPHORUM, Trin. Fund. Agrost. 106. t. 5 (1820) = **Chrysopogon**, Trin. (Gramin.).
chinense, Trin. l. c. = Chrys. acicularius.

CENTROPHYLLUM, Dum. Anal. Fam 32 (1829) = KENTROPHYLLUM, Neck. = **Carthamus**, Linn. (Compos.).

CENTROPHYTA, Reichb. Nom. 148 (1841) = KENTROPHYTA, Natt. = **Astragalus**, Tourn. (Legum.).

CENTROPODIUM, Lindl. Introd. Nat. Syst. ed. II. 442 (1836), sphalm. = CENTOPODIUM, Burch. = **Ilmex**, Neck. (Polygon.).

CENTROPOGON, Presl. Prod. Monog. Lobel. 48 (1836). *CAMPANULACEAE*, Benth. & Hook. f. ii. 547.
affine, *Mart. & Gal. in Bull. Acad. Brux.* ix. II. (1842) 40.—Mexic.
asmifolius, *Vatke, in Linnaea*, xxxviii. (1874) 716.—Peruv.
Andropogon, *DC. Prod.* vii. 345.—Ecuador.
argentinus, *Griseb. in Goott. Abh.* xxiv. (1879) 219.—Reg. Argent.
Berterianus, *DC. Prod.* vii. 345.—Ins. Guadalup.
bisleriotdes, *DC. l. c.* 346.—N. Granat.
Bonplandianus, *Presl. Prod. Monog. Lobel.* 48.—Am. austr.
calycinus, *Benth. Pl. Hartw.* 219.—N. Granat.
Chamissianus, *Kunth, in Mert. Fl. Bras.* vi. IV. 183. —Bras.
coccineus, *Regel, Gartenfl.* (1858) 374, in textu.—Bras.
cordatus, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 40 = cordifolius.
cordifolius, *Benth. Pl. Hartw.* 77.—Mexico.
costaricanus, *Planch. & Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 156.—Costa Rica.
cuspidatus, *DC. Prod.* vii. 346.—Peruv.
densiflorus, *Benth. Pl. Hartw.* 138.—Peruv.
discolor, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 13.—Venezuela.
edulis, Presl, Prod. Monog. Lobel. 48 = surinamensis.
exasperatus, *Presl, l. c.*—Peruv.
fasciculatus, *Regel, Cat. Pl. Hort. Absnhov.* 29.—Hab. ?

CENTROPOGON:—
fastuosus, *Scheidw. in Otto & Dietr. Allg. Gartens.* ix. (1841) 396.—Mexico.
flocculosus, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 16.—N. Granat.
glandulosus, *Decne. in Rev. Hortic.* Sér. III. ii. (1846) 491.—N. Granat.
grandis, *Presl, Prod. Monog. Lobel.* 48.—N. Granat.
granulosus, *Presl, l. c.* 49.—Peruv.
hirtus, *Presl, l. c.* 48.—Ecuador.
laevigatus, *DC. Prod.* vii. 344.—Guiana.
longipes, *Regel, Gartenfl.* iii. (1854) 3. t. 75.—Am. centr.
nutans, *Planch. & Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 156.—Costa Rica.
oblongus, *Benth. Pl. Hartw.* 214.—N. Granat.
prostratus, *Benth. l. c.* 212.—Peruv.
reflexus, *Presl, Prod. Monog. Lobel.* 49.—Peruv.
scandens, *Planch. & Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 157.—Costa Rica.
solanifolius, *Benth. Pl. Hartw.* 139.—Peruv.
speciosus, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 16.—Venezuela.
surinamensis, *Presl, Prod. Monog. Lobel.* 48.—Am. austr.
tovarensis, *Planch. & Linden, in Fl. des Serres*, Sér. I. vii. (1852–53) 146.—Venezuela.
virescens, Planch. & Oerst. in Kjoeb. Vidensk. Meddel. (1857) 157 = Burmeistera virescens.
Warszewiczii, *Van Houtte, ex Regel, Gartenfl.* (1858) 374.—Hab. ?
Warszewiczii, *Vatke, in Linnaea*, xxxviii. (1874) 716. —Costa Rica.

CENTROPSIS, Endl. Gen. Suppl. ii. 83 (1842) = KENTROPSIS, Moq. = **Sclerolaena**, R. Br. (Chenop.).

CENTROSEMA, Benth. in Ann. Wien. Mus. ii. (1838) 117. *LEGUMINOSAE*, Benth. & Hook. f. i. 527.
CRUMINIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 423.
PILANTHUS, Poit. ex Endl. Gen. 1289 (1841).
PLATYSEMA, Hoffmgg. ex Benth. in Ann. Wien. Mus. ii. (1838) 117.
STEGANOTROPIS, Lehm. Ind. Sem. Tort. Hamb. (1826).
VEXILLARIA, Hoffmgg. Verz. Pfl. 119 (1824) ; Benth. in Ann. Wien. Mus. ii. (1838) 117.
acutifolium, Benth. l. c. 118 = brasilianum.
angustifolium, *Benth. l. c.*—Venezuela.
asperrimum, *Benth. l. c.* 119.—Bras.
bifidum, *Benth. l. c.* 118.—Bras.
biflorum, Mart. ex Benth. l. c. 120 = virginianum.
brachypodum, Benth. in Ann. Nat. Hist. iii. (1839) 435 = brasilianum.
bracteosum, *Benth. in Ann. Wien. Mus.* ii. 119.—Bras.
brasilianum, *Benth. l. c.* 118.—Bras.
conjugatum, *Steud. Nom.* ed. II. i. 325.—Am. austr.
coriaceum, *Benth. in Ann. Wien. Mus.* ii. (1838) 118. —Bras.
cuspidatum, *Hassk. in Flora*, ii. (1866) 27.—Ind. or.
dasyanthum, *Benth. in Ann. Nat. Hist.* iii. (1839) 436.—Bras.
decumbens, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 120 = virginianum.
dubium, Hemsl. Biol. Centr. Am. Bot. i. 294 = hastatum.
fasciculatum, *Benth. in Ann. Wien. Mus.* ii. (1838) 120.—Bras.
ferrugineum, *A. Rich. Ess. Fl. Cub.* 410.—Cuba.
glabrum, Benth. in Ann. Wien. Mus. ii. (1838) 119 = cortaceum.
grandiflorum, *Benth. l. c.*—Bras.
grandiflorum, Walp. Rep. v. 529 = Clitoria grandiflora.
hastatum, *Benth. in Ann. Wien. Mus.* ii. (1838) 120. —Bras.
heptaphyllum, *Moric. Pl. Nouv. Rar. Am.* 96. t. 61.—Cuba.
lanceolatum, Steud. Nom. ed. II. i. 325 = brasilianum.
intermedium, A. Rich. Ess. Fl. Cub. 410 = pubescens.
longifolium, Benth. in Ann. Wien. Mus. ii. (1838) 118 = angustifolium.
macrocarpum, *Benth. in Ann. Nat. Hist.* iii. (1839) 436.—Guiana.
micranthum, Steud. Nom. ed. II. i. 326 in syn. sphalm. = virginianum.

CENTROSEMA:—
molle, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 119 = pubescens.
oblongum, Benth. l. c. 118 = cortaceum.
pascuorum, *Mart. ex Benth. l. c.* 120.—Bras.
perspicuum, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1203, nomen.—Guiana.
platycarpum, *Benth. in Mart. Fl. Bras.* xv. i. 126.—Peruv.
Plumieri, *Benth. in Ann. Wien. Mus.* ii. (1838) 118.—Am. austr.
pubescens, *Benth. l. c.* 119.—Bras.
rigidulum, Benth. in Hook. Journ. Bot. ii. (1846) 58 = sreurarium.
rotundifolium, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 119.—Bras.
Salzmanni, Benth. in Ann. Nat. Hist. iii. (1839) 436 = pubescens.
scabriusculum, Hassk. Pl. Jav. Rar. 377 = virginianum.
sinuatum, *Benth. in Ann. Wien. Mus.* ii. (1838) 120. —Bras.
sordidum, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1203, nomen.—Guiana.
venosum, *Mart. ex Benth. in Mart. Fl. Bras.* xv. i. 133.—Bras.
verticillatum, *Benth. in Hook. Journ. Bot.* ii. (1848) 58.—Guiana.
vestitum, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1203, nomen.—Guiana.
vetulum, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 119.—Bras.
vexillatum, *Benth. in Ann. Nat. Hist.* iii. (1839) 435. —Guiana.
virginianum, *Benth. in Ann. Wien. Mus.* ii. (1838) 120. —Am. trop.
virginianum, Griseb. in Goett. Abh. vii. (1857) 204 = pubescens.

CENTROSEPIS, R. Hedw. Gen. 51 (1806), sphalm. = **Centrolepis**, Labill.

CENTROSIA, A. Rich. in *Mém.* Soc. Hist. Nat. Par. iv. (1828) 89 = **Calanthe**, R. Br. (Orchid.).
Auberti, A. Rich. l. c. 40 = Cal. sylvatica.

CENTROSIS, Thou. Orch. Iles Afr. t. 35, 36 (1822) = **Calanthe**, R. Br. (Orchid.).
corymbosa, Thou. l. c. 1re Tabl. = Cal. sylvatica ?
plantaginea, Thou. l. c. 1re Tabl. = Cal. sylvatica ?
sylvatica, Thou. l. c. tt. 35, 36 = Cal. sylvatica.

CENTROSOLENIA, Benth. in Hook. Lond. Journ. Bot. v. (1846) 262 = **Epicoia**, Mart. (Gesnerac.).
aenea, Linden & André, in Illustr. Hortic. xxii. (1875) t. 222.—N. Granat.
bracteata, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 222.—N. Granat.
bractescens, Hook. Bot. Mag. t. 4675 = E. bractescens.
bullata, Lem. Illustr. Hortic. xvi. (1869) t. 607 = E. tessellata.
glabra, Benth. in Hook. Bot. Mag. t. 4553 = E. glabra.
glabra, Hook. Ic. Pl. t. 873 = E. Hookeri.
hirsuta, Benth. in Hook. Lond. Journ. Bot. v. (1846) 262 = E. hirsuta.
hirtiflora, Sprce, ex Hanst. in Linnaea, xxxiv. (1865–66) 450 = E. hirtiflora.
picta, Hook. Bot. Mag. t. 4611 = E. picta.

CENTROSPERMUM, H. B. & K. Nov. Gen. et Sp. iv. 270. t. 397 (1820) = **Acanthospermum**, Schrank (Compos.).
humile, Less. Syn. Comp. 217 = A. humile.
xanthioides, H. B. & K. Nov. Gen. et Sp. iv. 270 = A. brasilium.

CENTROSPERMUM, Spreng. Nov. Prov. 9 (1818) ; et Syst. iii. 588 (1826) = **Chrysanthemum**, Linn. (Compos.).
Chrysanthemum, Spreng. l. c. = C. viscosum.

CENTROSTACHYS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 497 (1824) = **Achyranthes**, Linn. (Amaranat.).
aquatica, Wall. Cat. n. 6933 = A. aquatica.
diandra, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 504 = A. diandra.
flabelligera, Fenzl, ex Moq. in DC. Prod. xiii. ii. 321 = A. aquatica.

CENTROSTEGIA, A. Gray, ex Benth. in DC. Prod. xiv. 27 (1856); et in Pacif. Rail. Rep. vii. 19 (1857). *POLYGONACEÆ*, Benth. & Hook. f. iii. 88.
leptocarpa, *A. Gray, in Proc. Am. Acad.* viii. (1870) 198.—Calif.
Thurberi, *A. Gray, ex Benth. in DC. Prod.* xiv. 27; *et in Pacif. Rail. Rep.* vii. 90.—Calif.

CENTROSTEMMA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 271. t. 12 = **Hoya**, R. Br. (Asclepiad.).
coriaceum, Meisn. Gen. Comm. 177 = H. coriacea.
Cyrtoceras, Meisn. l. c. = H. multiflora.
elegans, Blume, Mus. Bot. Lugd. Bat. i. 46 (= *Cyrtoceras elegans*).—Borneo.
laurifolium, Blume, l. c. (= *Cyrtoceras laurifolium*).—Java.
Lindleyanum, Decne. in DC. Prod. viii. 634 = H. multiflora.
micranthum, Blume, Mus. Bot. Lugd. Bat. i. 46 (= *Cyrtoceras micranthum*).—Ins. Celebes.
multiflorum, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 272 = H. multiflora.

CENTROSTYLIS, Baill. Etud. Gén. Euph. 469 (1858) = **Adenochlæna**, Baill. (Euphorb.).
scylanica, Baill. l. c. 470. t. 2. f. 26–29 = A. scylanica.

CENTUNCULUS, Adans. Fam. ii. 256 (1763) = **Cerastium**, Linn. (Caryophyll.).
alpinus, Scop. Fl. Carn. ed. II. i. 321 = Cerast. trigynum.
angustifolius, Scop. l. c. 322 = Cerast. strictum.
arvensis, Scop. l. c. 321 = Cerast. arvense.
rigidus, Scop. l. c. 322 = Cerast. strictum.
semidecandrus, Scop. l. c. 321 = Cerast. semidecandrum.

CENTUNCULUS, Dill. ex Linn. Syst. ed. I (1735). *PRIMULACEÆ*, Benth. & Hook. f. ii. 637.
ANAGALLIDASTRUM, Mich. ex Adans. Fam. ii. 230 (1763).
MICROPYXIS, Daby, in DC. Prod. viii. 71 (1844).
erectus, *Phil. Fl. Atac.* 35.—Chili.
lanceolatus, Michx. Fl. Bor. Am. i. 94 = minimus.
minimus, *Linn. Sp. Pl.* 116.—Europ.; Am. bor. et austr.
pentandrus, *R. Br. Prod.* 427.—Am. et Austral. trop.
simplex, Hornem. in Fl. Dan. ix. t. 1567 = minimus.
tenuifus, Daby, in DC. Prod. viii. 72 = pentandrus.

CEODES, Forst. Char. Gen. 141. t. 71 (1776) = **Pisonia**, Linn. (Nyctagin.).
umbellifera, Forst. l. c. 142 = P. umbellifera.

CEPA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 108 (1767) = **Allium**, Tourn. (Liliac.).
alba, Renault, Fl. Dép. Orne, 38 = A. Cepa ?
angulosa, Bernh. Syst. Verz. Erf. 302 = A. angulosum.
ascalonica, Neronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. 11 = A. ascalonicum.
carinata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 185 = A. carinatum.
cornua, Moench, Meth. Suppl. 80 = A. carneum.
cruciata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 188 = A. Cepa.
fistulosa, S. F. Gray, l. c. 184 = A. fistulosum.
flava, Moench, Meth. 248 = A. flavum.
flexuosa, Moench, Meth. Suppl. 80 = A. roseum.
Moly, Moench, Meth. 244 = A. Moly.
obliqua, Moench, l. c. 248 = A. obliquum.
oleracea, Bernh. Syst. Verz. Erf. 302 = A. oleraceum.
pallens, Renault, Fl. Dép. Orne, 38 = A. Cepa ?
paniculata, Moench, Meth. 248 = A. paniculatum.
prolifera, Moench, l. c. 244 = A. Cepa.
rubra, Renault, Fl. Dép. Orne, 38 = A. Cepa ?
Schoenoprasum, Moench, Meth. 244 = A. Schoenoprasum.
senescens, Moench, l. c. = A. senescens.
tenuifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 184 = A. Schoenoprasum.
uvarina, Bernh. Syst. Verz. Erf. 301 = A. ursinum.
vaginicum, Moench, Meth. 244 = A. fistulosum.
Victorialis, Moench, l. c. 248 = A. Victorialis.
vulgaris, Renault, Fl. Dép. Orne, 38 = A. Cepa.

CEPAEA, Casaubon; ex Fourr. in Ann. Sc. Linn. Lyon, N. S. xvi. (1868) 384 = **Sedum**, Tourn. (Crassulac.).
Cæsalpini, Fourr. l. c. = S. Cepæa.

CEPALARIA, Rafin. Fl. Tellur. iv. 96 (1836) = **Cephalaria**, Schrad. (Dipsac.).

CEPHAELIS Sw. Prod. Veg. Ind. Occ. 45 (1788). *RUBIACEÆ*, Benth. & Hook. f. ii. 127.
CALLICOCCA, Schreb. Gen. t. 196 (1789).
CARAPICHEA, Aubl. Pl. Gui. 167. t. 64 (1775).
CHENEA, Scop. Introd. 119 (1777).
EUSMOTIA, Neck. Elem. i. 207 (1790).
EVEA, Aubl. Pl. Gui. 103. t. 39 (1775).
IPECACUANHA, Arruda, Diss. 44 (1810), ex St. Hil. Pl. Bras. t. 6 (1824).
NETTLERA, Rafin. Sylva Tellur. 147 (1838).
TAPOGOMEA, Aubl. Pl. Gui. i. 357. t. 60 ad 63 (1775).
alba, *Willd. Sp. Pl.* i. 978.—Guiana.
ambigua, DC. Prod. iv. 534 = Psychotria Saimannïana.
attenuata, *Miq. in Linnæa*, xix. (1847) 444.—Bras.
axillaris, *Sw. Prod. Veg. Ind. Occ.* 45.—Ind. occ.
Beeri, *Teijsm. & Binn. Cat. Hort. Bog.* 384, nomen.—Hab. ?
Brakeana, Schlecht. in Linnæa, xxviii. (1856) 533 = Psychotria Beckeana.
bidentata, Benth. in Hook. Niger Fl. 421 = peduncularis.
bidentata, Thunb. ex Roem. & Schult. Syst. v. 214 = Psychotria bidentata.
bracteocardia, DC. Prod. iv. 534 = Psychotria bracteocardia.
brasiliensis, *Cham. in Linnæa*, ix. (1834) 287.—Bras.
bubalina, Pers. Syn. i. 303 = Burchellia capensis.
calycina, Lindl. Collect. i. 21 = Psychotria Lindleyana.
Carapichea, Raeusch. ex Steud. Nom. ed. II. i. 326 = Isertia.
Chamissonis, Steud. l. c. = Salzmannia nitida.
clausa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 225.—Afr. trop.
colorata, *Hoffmgg. ex Roem. & Schult. Syst.* v. 213.—Bras.
congesta, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 226.—Afr. trop.
coriacea, D. Dietr. Syn. Pl. i. 774 = Psychotria xanthophylla.
coriacea, G. Don, Gen. Syst. iii. 606 = peduncularis.
cornuta, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 234.—Afr. trop.
costaricensis, *Schlecht. in Linnæa*, xxviii. (1856) 546.—Costa Rica.
croces, *G. F. W. Mey. Prim. Fl. Esseq.* 107.—Guiana.
cuneata, *Korth. in Nederl. Kruidk. Arch.* ii. (1851) 564.—Malaya.
cyanocarpa, Moc. & Sesse, ex DC. Prod. iv. 533 = tomentosa.
cymosa, Spreng. Syst. i. 749 = Psychotria mayparensis.
dichotoma, *Rudge, Pl. Gui.* i. 29. t. 44.—Guiana.
dichotoma, Willd. ex Roem. & Schult. Syst. v. 214 = Psychotria Hoffmannseggiana.
discolor, *Vatke in Linnæa*, xli. (1877) 572.—Costa Rica.
diversifolia, Blume, Bijdr. 1004 = Geophila reniformis.
elata, *Sw. Prod. Veg. Ind. Occ.* 45.—Jamaica.
emetica, Pers. Syn. i. 303 = Psychotria Ipecacuanha.
Evea, *DC. Prod.* iv. 535.—Guiana.
ferruginea, G. Don, Gen. Syst. iii. 605 = Sabicea ferruginea.
fragrans, *Hook. & Arn. Bot. Beech. Voy.* 64. t. 13.—Ins. Societ.
furcata, Bartl. ex DC. Prod. iv. 518 = Psychotria furcata.
fuscescens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 224.—Afr. trop.
gesneriflora, *Willd. ex Roem. & Schult. Syst.* v. 213.—Hab. ?
glabra, *Willd. Sp. Pl.* i. 979.—Guiana.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 178.—Malacca.
herbacea, Kunz, in Journ. As. Soc. Beng. (1877) ii. 140 = Geophila reniformis.
hirsuta, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 185.—Mexic.
hirta, *Miq. in Linnæa*, xvii. (1843) 71.—Guiana.
Hoffmannseggiana, Roem. & Schult. Syst. v. 214 = Psychotria Hoffmannseggiana.
Humboldtiana, Cham. & Schlecht. in Linnæa, iv. (1829) 136 = Psychotria Humboldtiana.
involucrata, *Willd. Sp. Pl.* i. 979.—Guiana.
Ipecacuanha, Rich. in Bull. Fac. Med. iv. (1816) 92 = Psychotria Ipecacuanha.
isticiscifolia, *Rudge, Pl. Gui.* i. 28. t. 43.—Guiana.
latifolia, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 226.—Afr. trop.
ligularia, *Rich. ex DC. Prod.* iv. 533.—Guiana.

CEPHAËLIS:—
Mannii, *Benth. & Hook. f. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 225.—Afr. trop.
maranhatana, *G. Don, Gen. Syst.* iii. 605.—Bras.
microcephala, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 214 = Psychotria microcephala.
microcephala, Miq. in Linnæa, xviii. (1844) 748 = Psychotria Hoffmannseggiana.
muscosa, *Sw. Prod. Veg. Ind. Occ.* 46.—Ins. Martinic.
nuda, *Cham. & Schlecht. in Linnæa*, iv. (1829) 135.—Bras.
oblonga, *DC. Prod.* iv. 535.—Guiana.
ornata, *Spruce, ex Benth. & Hook. f. Gen.* ii. 128, nomen.—Bras.
Patabea, Steud. Nom. ed. I. 177, 596 (= *Patabea coccinea*).—Bras.
Patrisii, *D. Dietr. Syn. Pl.* i. 773.—Guiana.
peduncularis, *Salisb. Parad. Lond.* t. 99.—Afr. trop.
pedunculata, G. Don, Gen. Syst. iii. 605 = pedunicularis.
peruviana, Spreng. Syst. i. 749 = Psychotria capitata.
polycephala, *Schlecht. in Linnæa*, xxviii. (1856) 532.—Peruv.
prunifolia, H. B. & K. Nov. Gen. et Sp. iii. 377 = Psychotria microcephala.
psychotriæfolia, *Seem. Bot. Voy. Herald.* 138.—Panama.
pubescens, Hoffmgg. ex Roem. & Schult. Syst. v. 213 = Psychotria bracteocardia.
punicea, Vahl, Eclog. Am. i. 19 = elata.
purpurea, *Willd. Sp. Pl.* i. 978.—Guiana.
reniformis, H. B. & K. Nov. Gen. et Sp. iii. 377 = Geophila reniformis.
rosea, Benth. in Hook. Journ. Bot. iii. (1841) 224 = Psychotria rosea.
rubescens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 225.—Afr. trop.
rubra, Hoffmgg. ex Roem. & Schult. Syst. v. 214 = Psychotria rubra.
ruelliæfolia, Cham. & Schlecht. in Linnæa, iv. (1829) 134 = Psychotria ruelliæfolia.
salicifolia, *Ib. B. & K. Nov. Gen. et Sp.* iii. 376.—N. Granat.
sessilifora, Willd. Sp. Pl. i. 979 (= *Patabea coccinea*).—Cost. Rica.
spathacea, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 225.—Afr. trop.
speciosa, Spreng. Syst. i. 749 = Psychotria speciosa.
stipulacea, *Blume, Bijdr.* 1005.—Java.
suaveolens, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 224.—Afr. trop.
subcaudata, *Sagot, ex Benth. & Hook. f. Gen.* ii. 128.—Guiana.
subtomentosa, Spreng. Syst. i. 749 = Psychotria subtomentosa.
Swartzii, *DC. Prod.* iv. 534.—Ind. occ.
tenuiflora, D. Dietr. Syn. Pl. i. 774 (= *Patabea tenuiflora*).—Guiana.
tenuis, D. Dietr. l. c. = Psychotria tenerior.
tetranda, Willd. Sp. Pl. i. 979 = C. Evea.
tomentosa, Vahl, Eclog. Am. i. 19 = Psychotria tomentosa.
triplocephala, *Belle, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 282.—Porto Rico.
turbinata, *Benth. & Hook. f. Gen.* ii. 128, nomen.—Bras.
violacea, Sw. Prod. Veg. Ind. Occ. 45 = Swartzii.
violacea, *Willd. Sp. Pl.* i. 977.—Ind. occ.
violæfolia, H. B. & K. Nov. Gen. et Sp. iii. 379 = Geophila violæfolia.

CEPHALANDRA, Schrad. ex Eckl. & Zeyh. Enum. 280 (1836) = Cocotinia, Wight & Arn. (Cucurb.).
decipiens, Hook. f. in Oliver, Fl. Trop. Afr. ii. 502 = Cocc. decipiens.
diversifolia, Nsnd. in Ann. Sc. Nat. Sér. V. v. (1866) 18. t. 7 = Cocc. Hartmannïana.
grandis, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 108 = Cocc. indica.
indica, Nsnd. in Ann. Sc. Nat. Sér. V. v. (1866) 16 = Cocc. indica.
M'Kennii, Nsnd. l. c. 17 = Cocc. Mac-Kennii.
palmata, Sond. in Harv. & Sond. Fl. Cap. ii. 498 = Cocc. palmata.
pubescens, Sond. l. c. = Cocc. adoënsis.
quinquelobata, Schrad. ex Eckl. & Zeyh. Enum. 280 = Cocc. quinquelobata.
Schimperi, Nsnd. in Ann. Sc. Nat. Sér. V. v. (1866) 16 = Cocc. indica.

CEPHALANDRA :—
sessilis, Kleinsch. in Peters, Reise Mossamb. Bot. 151 = Coccinia sessilis.
sessilifolia, Sond. in Harv. & Sond. Fl. Cap. ii. 498 = Cocc. sessilifolia.
Wightiana, Hook. f. Fl. Brit. Ind. ii. Index 757, sphalm. = Cocc. indica.

CEPHALANOPHLOS, Neck. Elem. i. 68 (1790) = Cnicus, Tourn. (Compos.).

CEPHALANTHERA, Rich. in Mém. Mus. Par. iv. (1818) 51. *ORCHIDEÆ*, Benth. & Hook. f. iii. 619. DORYCHILE, Reichb. Nom. 56 (1841).
acuminata, Lindl. Fl. Ross. iv. 78 = pallens.
acuminata, Lindl. in Wall. Cat. n. 7405 = ensifolia.
chloidophylla, *Reichb. f. in Linnæa*, xxv. (1852) 238. —Japon.
comosa, Tineo, in Guss. Fl. Sic. Syn. ii. 877 = rubra.
cucullata, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 13.—Graecia ; As. Min. ; Persia.
ensifolia, *Rich. in Mém. Mus. Par.* iv. (1818) 60.—Europ. ; Afr. bor. ; As. temp.
epipactoides, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 30 = cucullata.
erecta, *Blume, Orch. Archip. Ind.* 188.—Japon.
falcata, *Blume, l. c.* 187.—Japon.
grandiflora, S. F. Gray, Nat. Arr. Brit. Pl. ii. 210 = pallens.
japonica, A. Gray, in Perry, Exp. Jap. ii. 319 = falcata.
lancifolia, *Dum. Fl. Belg.* 134 ; *Tod. Orch. Sic.* 123.—Sicilia.
longibracteata, Reichb. f. Ic. Fl. Germ. xiii. et xiv. 138 = pallens.
longibracteata, *Blume, Orch. Archip. Ind.* 188.—Japon.
Maximowiczii, Tineo, in Guss. Fl. Sic. Syn. ii. 877 = ensifolia.
ochroleuca, Reichb. Fl. Germ. Excurs. 1430 = pallens.
oregana, *Reichb. f. in Linnæa*, xli. (1877) 58.—Am. bor. occ.
pallens, Hohen. in Bull. Soc. Nat. Mosc. (1838) 257 = ensifolia.
pallens, *Rich. in Mém. Mus. Par.* iv. (1818) 60.—Europ. ; Reg. Caucas.
palustris, Rich. ex Prits. Ic. Ind. ii. 71, sphalm. = Epipactis palustris.
platychelia, Reichb. f. in Bot. Zeit. iii. (1845) 335 = falcata.
Royleana, Regel, in Act. Hort. Petrop. vi. (1879) 490 = Epipactis Royleana.
rubra, *Rich. in Mém. Mus. Par.* iv. (1818) 60.—Europ. ; As. Min. ; Persia.
Thomsoni, *Reichb. f. in Linnæa*, xli. (1877) 54.—Reg. Himal.
xiphophylla, Boiss. *Diagn.* Ser. I. vi. Fl. Germ. xiii. et xiv. 135 = ensifolia.

CEPHALANTHUS, Linn. Gen. ed. I. 61 (1737). *RUBIACEÆ*, Benth. & Hook. f. ii. 80. ACRODRYON, AXOLUS, ERESIMUS, GILIPUS, SILAMNUS, Rafin. Sylva Tellur. 60–61 (1838).
acuminatus, *Rafin. New Fl. Am.* iii. 25.—Am. bor.
africanus, Reichb. ex DC. Prod. iv. 545 = Mitragyna africana.
angustifolius, *Lour. Fl. Cochinch.* i. 67.—China.
aralioides, Zoll. & Moritzi, Syst. Verz. Zoll. 61 = nancleoides.
chinensis, Lam. Encyc. i. 678 = Nauclea purpurea.
montanus, *Lour. Fl. Cochinch.* i. 67.—China.
natalensis, *Oliver, in Hook. Ic. Pl.* t. 1331.—Natal.
nauclcoides, *DC. Prod.* iv. 539.—Reg. Himal. ; Malaya ; China.
obtusifolia, *Rafin. New Fl. Am.* iii. 25.—Am. bor.
occidentalis, *Linn. Sp. Pl.* 95.—Am. bor. ; Mexic. ; Ind. occ.
occidentalis, Lour. Fl. Cochinch. i. 67 = Naucica orientalis.
oppositifolius, Moench, Meth. 487 = occidentalis.
orientalis, Blume, Cat. Gew. Buitenz. 36 = Nauclea purpurea.
orientalis, Linn. Sp. Pl. 95 = Anthocephalus moriudaefolius.
orientalis, *Roem. & Schult. Syst.* iii. 105.—China.
pilulifer, Lam. Encyc. i. 679 = Mitragyna parvifolia.
pilulifcrus, *Willd. ex Roem. & Schult. Syst.* iii. 525. —Ind. ?
procumbens, *Lour. Fl. Cochinch.* i. 7.—Cochinch.
pubescens, *Rafin. New Fl. Am.* iii. 5.—Am. bor.

CEPHALANTHUS :—
salicifolius, Humb. & Bonpl. Pl. Aequin. ii. 63. t. 98 = occidentalis.
Sarandi, *Cham. & Schlecht. in Linnæa*, ii. (1827) 610. —Bras.
spathelliferus, *Baker, in Journ. Bot.* xx. (1882) 187.—Madag.
stellatus, *Lour. Fl. Cochinch.* i. 68.—China.

CEPHALARIA, Schrad. Cat. Sem. Hort. Gotting. (1814) ex Roem. & Schult. Syst. iii. 1 (1818). *DIPSACEÆ*, Benth. & Hook. f. ii. 159. CERIONANTHUS, Schott, ex Roem. & Schult. l. c. 43 (1818).
LEPICEPHALUS, Lag. Gen. et Sp. Nov. 7 (1816).
LEUCOPHORA, Rafin. Fl. Tellur. iv. 95 (1836).
PICNOCOMUM, DC. Prod. iv. 647 (1830) in syn.
PTEROCEPHALUS, Adans. Fam. ii. 152 (1763).
PYCNOCOMON, Wallr. Sched. Crit. 46 (1822) in syn.
SICOSIRA, Wallr. l. c. (1822).
acaulis, *Stend. ex A. Rich. Tent. Fl. Abyss.* i. 368.—Abyssin.
albocrem, Roem. & Schult. Syst. iii. 46 = leucantha.
alpina, *Schrad. Cat. Sem. Hort. Gotting.* (1814) ; *ex Roem. & Schult. Syst.* iii. 48.—Europ.
ambrosioides, *Roem. & Schult. l. c.* 45.—Graecia.
amplexicaulis, *Stend. Nom.* ed. II. i. 837.—Graecia ; Oriens ?
appendiculata, Schrad. Cat. Sem. Hort. Gotting. (1814) ; ex Roem. & Schult. Syst. iii. 42 = Dipsacus pilosus.
aristata, *C. Koch, in Linnæa*, xxiv. (1851) 445.—As. Min.
attenuata, *Roem. & Schult. Syst.* iii. 44.—Afr. austr.
balearica, *Coss. ex Willk. in Linnæa*, xl. (1876) 87.—Hispan.
bidens, Roem. & Schult. Syst. iii. 45 = Scabiosa bidens.
bortica, Boiss. Voy. Bot. Espagne, 788 = leucantha.
Boissieri, Reut. in Boiss. Diagn. Ser. II. ii. 122 = syriaca.
cachemirica, Decne. in Jacquem. Voy. Bot. 86. t. 94 = Dipsacus inermis.
calva, Boiss. & Bal. Diagn. Ser. II. vi. 95 = aristata.
centaureifolia, Roem. & Schult. Syst. iii. 49 = cornicalata.
centaurioides, Schur, Enum. Pl. Trans. 294 = laevigata.
cilicica, *Boiss. & Kotschy, Diagn.* Ser. II. v. 106.—Cilicia.
coriacea, Stend. Nom. ed. II. i. 837 = corniculata.
corniculata, *Roem. & Schult. Syst.* iii. 49.—Bunatus.
crotacea, Roem. & Schult. l. c. 51 = corniculata.
decurrens, Roem. & Schult. l. c. 50 = attenuata.
diandra, Roem. & Schult. l. c. 52 = Scabiosa papposa.
dichaetophora, *Boiss. Diagn.* Ser. I. vi. 71.—Persia.
dipsacoides, *Boiss. & Bal. Diagn.* Ser. II. v. 107.—Cilicia ; Syria.
dipsacoides, Kar. & Klr. in Bull. Soc. Nat. Mosc. (1841) 484 = Dipsacus asureus.
elata, Schrad. in Cat. Sem. Hort. Gotting. (1814) ; ex Roem. & Schult. Syst. iii. 51 = tatarica.
Fusziana, Heuff. ex Schur, Enum. Pl. Transs. 293 = radiata.
graeca, *Roem. & Schult. Syst.* iii. 48.—Graecia.
hirsuta, *Stapf, in Denkschr. Acad. Wien*, i. (1885) 55. —Persia.
humilis, Roem. & Schult. Syst. iii. 50 = attenuata.
involucrata, *Roem. & Schult. l. c.* 53.—Creta ; Cyprus.
joppensis, *Coult. ex DC. Prod.* iv. 648.—Italia ; Cilicia ; Syria.
junces, *Boiss. Diagn.* Ser. I. vi. 72.—Persia.
Kotschyi, Boiss. & Hohen. l. c. x. 76 = procera.
laevigata, Pant. ex Nym. Consp. 346 = alpina.
laevigata, Schrad. Cat. Sem. Hort. Gotting. (1814) = leucantha.
lavandulacea, *Stend. in Harv. & Sond. Fl. Cap.* iii. 41. —Afr. austr.
leucantha, Bunge, ex Nym. Consp. 346 = radiata.
leucantha, Ledeb. Fl. Ross. ii. 449 = corniculata.
leucantha, *Schrad. in Cat. Sem. Hort. Gotting.* (1814) ; *ex Roem. & Schult. Syst.* iii. 47.—Hispan. ; Gall.
leucanthema, Roem. & Schult. l. c. 46 = leucantha.
linearifolia, Lange, in Kjoeb. Vidensk. Meddel. (1877–78) 226 = leucantha.
longifolia, E. Mey. Zwei Pfl. Docum. 171 = attenuata.
macrophylla, Griseb. Spicil. Fl. Rumel. ii. 174 = ambrosioides.

CEPHALARIA :—
mauritanica, *Pomel, Nouv. Mat. Fl. Atl.* 66.—Afr. bor.
microcephala, *Boiss. Diagn.* Ser. II. ii. 123.—Persia.
neglecta, *Vvriot, Ind. Sem. Hort. Gratianop.* (1852) 10 ; *et Linnæa*, xxvi. (1853–55) 764.—Hab. ?
papposa, Roem. & Schult. Syst. iii. 52 = syriaca.
pilosa, *Boiss. & Huet, Diagn.* Ser. II. ii. 122.—Armenia.
pilosa, Gren. & Godr. Fl. Fr. ii. 69 = Dipsacus pilosus.
plumosa, Roem. & Schult. Syst. iii. 53 = Scabiosa lustanica ?
procera, *Fisch. & Avé-Lall. Ind. Sem. Hort. Petrop.* vii. 46.—As. Min.
procera, Pant. ex Nym. Consp. 346 = alpina.
radiata, *Griseb. & Schenk, in Wiegm. Archiv*, xviii (1852) 1. 351.—Transsylv.
rigida, Eckl. & Zeyh. Enum. 371 = rigida.
rigida, *Roem. & Schult. Syst.* iii. 44.—Afr. austr.
rupestris, Griseb. ex Nym. Consp. 346 = alpina.
scabra, Eckl. & Zeyh. Enum. 371 = rigida.
scabra, *Roem. & Schult. Syst.* iii. 44.—Afr. austr.
setosa, *Boiss. & Hohen. Diagn.* Ser. I. ii. 107.—Syria.
setulifera, Boiss. & Heldr. in Boiss. Fl. Orient. iii. 194 = graeca.
speciosa, *Boiss. & Kotschy, in Boiss. l. c.* 121.—Armenia.
stellipilla, *Boiss. Diagn.* Ser. I. x. 76.—Syria.
syriaca, *Schrad. Cat. Sem. Hort. Gotting.* (1814) ; *ex Roem. & Schult. Syst.* iii. 45.—Reg. Mediterr. ; Oriens.
tatarica, *Schrad. l. c.* ; *ex Roem. & Schult. l. c.* 51. —Rossia ; As. bor. ; Oriens.
Tchihatchewi, *Boiss. in Tchihat. Asie Min. Bot.* ii. 220. t. 26.—Armenia.
transylvanica, *Schrad. Cat. Sem. Hort. Gotting.* (1814) ; *ex Roem. & Schult. Syst.* iii. 45.—Europ. austr. ; As. Min. ; Reg. Caucas.
uralensis, Roem. & Schult. Syst. iii. 50 = corniculata.
utriulata, E. Mey. Zwei Pfl. Docum. 171 = attenuata.
utulata, *Roem. & Schult. Syst* iii. 43.—Afr. austr.
Vaillantii, Schott, ex Roem. & Schult. l. c. 46 = syriaca.
virginea, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 178 = graeca.
Willkiisi, Roem. & Schult. Syst. iii. 53 = Scabiosa pilumosa.

CEPHALEIS, Vahl, Eclog i. 19 (1796), sphalm. = Cephaëlis, Sw. (Rubiac.).

CEPHALIDIUM, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) = Anthocephalus, A. Rich. (Rubiac.).

CEPHALINA, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 106 (1827) = Sabicoephalus, Afzel. (Rubiac.).
esculenta, Schum. & Thonn. l. c. = S. esculentus.
scandens, DC. Prod. iv. 368, in syn. = S. esculentus.

CEPHALIPTERUM, A. Gray, in Hook. Kew Journ. iv. (1852) 271. *COMPOSITÆ*, Benth. & Hook. f. ii. 321.
Drummondii, *A. Gray, l. c.* 272.—Austral.

CEPHALOCARPUS, Nees, in Mart. Fl. Bras. ii. 1 162. t. 18 (1842). *CYPERACEÆ*, Benth. & Hook. f. iii. 1068.
Dracaenula, *Nees, l. c.*—Bras.

CEPHALOCEREUS, Pfeiff. in Otto & Dietr. Allg. Gartenz. vi. (1838) 142 = Cereus, Haw. (Cactese).
senilis, Pfeiff. l. c. = C. senilis.

CEPHALOCHLOA, Coss. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 229 = Ammochloa, Boiss. (Gramin.).
echinata, Coss. & Dur. l. c. = A. pungens.

CEPHALOCROTON, Hochst. in Flora, xxiv. (1841) 870. *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 207.
albicans, *Muell. Arg. in DC. Prod.* xv. ii. 760.—Malaya.
cordifolius, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 155.—Afr. trop.
discolor, *Muell. Arg. in DC. Prod.* xv. ii. 761.—Malaya.

CEPHALOCROTON :—
indicus, Bedd. Fl. Sylv. ii. t. 261 = Adenochlaena indica.
leucocephalus, *Baill. Adansonia*, v. (1864–65) 148.—Madag.
mollis, *Klotzsch, in Peters, Reise Mosamb. Bot.* 99. t. 17.—Afr. trop.
orientalis, *Mtg. Ann. Mus. Bot. Lugd. Bat.* iv. 120.—Ins. Timor.
socotranus, *Bal.f. in Proc. Roy. Soc. Edinb.* xii. (1884) 95.—Ins. Socotra.
seylanicus, Baill. Adansonia, v. (1864–65) 148 = Adenochlaena seylanica.

CEPHALODES, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159 = **Cephalaria**, Schrad. (Dipsac.).

CEPHALOMA, Neck. Elem. i. 314 (1790) = **Dracocephalum**, Linn. (Labiat.).

CEPHALOMAPPA, Baill. Adansonia, xi. 180 (1874). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 323.
Beccariana, *Baill. l. c.* 181.—Borneo.

CEPHALONOPLOS, Fourr. in Ann. Soc Linn. Lyon, N. S. xvi. (1868) 95 = **Cnicus**, Tourn. (Compos.).
arvens..., Fourr. l. c. = Cnicus arvensis.

CEPHALOPAPPUS, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 5. t. 1. *COMPOSITÆ*, Benth. & Hook. f. ii. 504.
sonchifolius, *Nees & Mart. l. c.* 6.—Bras.

CEPHALOPHORA, Cav. Ic. vi. 79. t. 599 (1801). *COMPOSITÆ*, Benth. & Hook. f. ii. 418.
ACTINEA, A. Juss. in Ann. Mus. Par. ii. (1803) 425. t. 61.
ACTINELLA, Pers. Syn. ii. 469 (1807).
GRAHMIA, Hook. Exot. Fl. t. 189 (1827).
GRAHAMIA, Spreng. Syst. iv. Cur. Post. 297, 299 (1827).
PTILEPIDA, Rafin. in Am. Monthly Mag. (1818) 268.
acaulis, DC. Prod. v. 663 = Actinella acaulis.
anthemoides, Less. in Linnæa, vi. (1831) 518 = Actinella odorata.
aromatica, *Schrad. Ind. Sem. Hort. Gotting.* (1830); *Less. Syn. Comp.* 240.—Chili.
Berteroana, *Phil. in Linnæa*, xxix. (1857–58) 9.—Chili.
collina, *Phil. l. c.* 8.—Chili.
decurrens, Less. l. c. vi. (1831) 517 = Helenium puberulum.
Doniana, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 336.—Reg. Argent.
elongata, D. Don, ex DC. Prod. vii. 298, pro parte = Gaillardia megapotamica.
elongata, D. Don, ex Hook. Journ. Bot. iii. (1841) 324, pro parte = Doniana.
fasciculata, Pohl, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 38 = Hyptis caprariæfolia.
foliosa, *Phil. in Linnæa*, xxix. (1857–58) 7.—Chili.
glauca, *Cav. Ic.* vi. 80. t. 599.—Chili.
grandiflora, *Walp. Ann.* i. 416.—Am. bor. occ.
heterophylla, *Less. Syn. Comp.* 240.—Reg. Argent.
insularis, *Phil. in Anal. Univ. Chil.* (1878) 506.—Chili.
integrifolia, Steud. Nom. ed. II. i. 328 = Helenium integrifolium.
litoralis, *Phil. Fl. Atac.* 84.—Chili.
plantaginea, *DC. Prod.* v. 662.—Chili.
punctata, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 228.—Chili.
radiata, Less. in Linnæa, vi. (1831) 516 = heterophylla.
rigida, *Phil. in Linnæa*, xxxiii. (1864–65) 170.—Chili.
scabiosoides, D. Don, ex DC. Prod. vii. 298 = Gaillardia megapotamica.
scaposa, DC. Prod. v. 663 = Actinella scaposa.
setigera, *Phil. in Anal. Univ. Chil.* (1878) 505.—Chili.
suffruticosa, D. Don, ex Hook. Journ. Bot. iii. (1841) 324 = Doniana.
tenera, Cass. Opusc. iii. 93 = aromatica.
Urmeneta, *Phil. in Anal. Univ. Chil.* (1878) 506.—Chili.

CEPHALOPHORUS, Lem. Cact. Aliq. Nov. Desc. p. xii (1838) = **Cereus**, Haw. (Cactese).
Columna-Trajani, Lem. l. c. = Cereus Columna-Trajani.
senilis, Lem. l. c. = Cereus senilis.

CEPHALOPHYTON, Hook. f. ex Baker, in Journ. Linn. Soc. xx. (1883) 250, nomen = **Thonningia**, Vahl, (Balanoph.).
Parkeri, Hook. f. ex Baker, l. c.—Madag.

CEPHALORKHYNCHUS, Boiss. Diagn. Ser. I. iv. 28 (1844) = **Lactuca**, Tourn. (Compos.).
Aucheri, Boiss. l. c. vii. 11 = L. microcephala.
Candolleanus, Boiss. Fl. Orient. iii. 820 = L. hispida.
glandulosus, Boiss. Diagn. Ser. I. iv. 28.—Aa. Min.
hispidus, Boiss. Fl. Orient. iii. 821 = L. microcephala.
Kotschyi, Boiss. & Reut. ex Boiss. l. c. 816 = L. seticuspis.

CEPHALOSCHOENUS, Nees, in Linnæa, ix. (1834) 295 = **Rynchospora**, Vahl (Cyperac.).
Armeria, Nees, l. c. 296 = R. globosa.
articulatus, Nees, in Edinb. N. Phil. Journ. xvii. (1834) 266 = R. surea.
divergens, Nees, in Linnæa, ix. (1834) 296 = R. aurea.
globosus, Nees, l. c. = R. globosa.
longisetis, Nees, l. c. = R. longiseta.
marginatus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 250 = R. globosa.
oligocephalus, Hochst. in Flora, xxviii. (1845) 759 = R. cyperoides.
parvus, Nees, in Linnæa, ix. (1834) 296 = R. Wallichiana.
Poblicanus, Nees, in Mart. Fl. Bras. ii. t. 131 = R. globosa.
polycephalus, Nees, in Linnæa, ix. (1834) 296 = R. cyperoides.
pseudocurucus, Nees, l. c. = R. gigantea.
tenuirostris, Nees, l. c. = R. cyperoides, tenuirostris.
seylanicus, Nees, in Edinb. N. Phil. Journ. xvii. (1834) 265 = R. trifora.

CEPHALOSCIRPUS, Kurz, in Journ. As. Soc. Beng. xxxvii. (1869) 11. 88 = **Mapania**, Aubl. (Cyperac.).
macrocephalus, Kurz, l. c. 84.—Ins. Moluce.

CEPHALOSERIS, Poepp. ex Reichb. Consp. 212 (1828) = **Polyachyrus**, Lag. (Compos.).
Poeppigii, Kunze, ex Steud. Nom. ed. II. i. 328 = P. Poeppigii.

CEPHALOSORUS, A. Gray, in Hook. Kew Journ. iii. (1851) 152 = **Angianthus**, Wendl. (Compos.).
brevipapposus, F. Muell. Fragm. iii. 159 = A. phyllocephalus.
gnephosoides, F. Muell. l. c. ii. 158 = Gnephosis oyathopappa.
gymnocephalus, A. Gray, in Hook. Kew Journ. iii. (1851) 153 = Gnephosis macrocephala.
leptocladus, F. Muell. Fragm. iii. 158 = Gnephosis leptoclada.
microcephalus, F. Muell. l. c. = A. microcephalus.
phyllocephalus, A. Gray, in Hook. Kew Journ. iii. (1851) 153 = A. phyllocephalus.

CEPHALOSTACHYUM, Munro, in Trans. Linn. Soc. xxvi. (1868) 138. *GRAMINEÆ*, Benth. & Hook. f. iii. 1213.
capitatum, *Munro, l. c.* 139.—Reg. Himal.
Chapelieri, *Munro, l. c.* 140.—Madag.
flavescens, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 252.—Burma.
Griffithii, Kurz, For. Fl. Brit. Burma, ii. 566 = Teinostachyum Griffithii.
latifolium, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 140.—Reg. Himal.
pallidum, *Munro, l. c.* 139.—Reg. Himal.
pergracile, *Munro, l. c.* 141.—Burma.
schizochyloides, *Kurz, For. Fl. Brit. Burma*, ii. 565. —Ins. Andaman.
virgatum, *Kurz, l. c.*—Burma.

CEPHALOSTEMON, Rob. Schomb. Rap. & Sazo-Frid. 9 (1845). *RAPATEACEÆ*, Benth. & Hook. f. iii. 856.
affinis, *Koern. in Linnæa*, xxxvii. (1871–72) 447.—Venezuela.

CEPHALOSTEMON :—
gracilis, *Rob. Schomb. Rap. & Sazo-Frid.* 9.—Bras. Riedelianus, *Koern. in Linnæa*, xxxvii. (1871–72) 445. —Bras.
squarrosus, *Koern. l. c.* 450.—Venezuela.

CEPHALOSTIGMA, A. DC. Monog. Camp. 117 (1830). *CAMPANULACEÆ*, Benth. & Hook. f. ii. 555.
anagalloides, Royle, Illustr. Bot. Himal. 253 = hirsutum.
bahiense, A. DC. Prod. vii. 421 = Perrottetii.
erectum, Vatke, in Linnæa, xxxvlii. (1874) 699 = Schimperi.
flexuosum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 9.—Ind. or.
hirsutum, *Edgew. in Trans. Linn. Soc.* xx. (1851) 81.—Reg. Himal.
hirsutum, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 9 = Schimperi.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 439.—Reg. Himal.
paniculatum, *A. DC. Monog. Camp.* 117.—Burma.
paniculatum, Wall. Cat. n. 1995 C = hirsutum.
Perrottetii, *A. DC. Monog. Camp.* 118.—Afr. trop.
Perrottetii, Hook. f. in Journ. Linn. Soc. vii. (1864) 204 = ramosissimum.
Prieurei, *A. DC. Monog. Camp.* 118.—Afr. trop.
ramosissimum, *Hemsl. in Oliver, Fl. Trop. Afr.* iii. 472. —Afr. trop.
Schimperi, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 2. —Ind. or.; Afr. trop.
spathulatum, Thw. Enum. Pl. Zeyl. 422 = Campanula canescens.

CEPHALOTAXUS, Sieb. & Zucc. ex Endl. Gen. Suppl. ii. 27 (1842). *CONIFERÆ*, Benth. & Hook. f. iii. 430.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 169.—Japon.
drupacea, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 108.—Japon.
Fortuni, Hook. Bot. Mag. t. 4499 = drupacea.
Harringtonia, C. Koch, Dendrol. ii. ii. 102 = pedunculata.
pedunculata, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 108. —Japon.
sumatrana, *Miq. Fl. Ind. Bat.* ii. 1076.—Sumatra.
tardiva, Siebold, ex Endl. Syn. Conif. 239 = Taxus brevifolia.
umbraculifera, Siebold, ex Endl. l. c. = Taxus brevifolia.

CEPHALOTOMANDRA, Karst. & Triana, in Linnæa, xxviii. (1856) 439. *NYCTAGINEÆ*, Benth. & Hook. f. iii. 10.
fragrans, *Karst. & Triana, l. c.* 480.—N. Granat.

CEPHALOTROPHIS, Blume, Mus. Bot. Lugd. Bat. ii. 75. t. 97. 1852 [1856] = **Malaisia**, Blanco (Urticac.).
javanica, Blume, l. c. 76 = M. tortuosa.
puberula, Miq. Fl. Ind. Bat. Suppl. 416 = M. puberula.

CEPHALOTUS, Labill. Nov. Holl. Pl. ii. 6. t. 145 (1806). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 655.
follicularis, *Labill. l. c.* 7.—Austral.

CEPHALOXIS, Desv. Journ. Bot. i. (1806) t. 11 = seq.

CEPHALOXYS, Desv. Journ. Bot. i. (1808) 321 = **Juncus**, Tourn.
flabellata, Desv. l. c. 324 = J. repens.
graminifolia, Nees, ex Kunth, Enum. Pl. iii. 343 = J. cyperoides.

CERADIA, Lindl. Bot. Reg. (1845) Misc. 12 = **Othonna**, Linn. (Compos.).
furcata, Lindl. l. c. (= *Doria Ceradia*).—Afr. austr.

CERAIA, Lour. Fl. Cochinch. ii. 518 (1790) = **Dendrobium**, Sw. (Orchid.).
simplicissima, Lour. l. c. = D. Ceraia.

CERAMANTHE, Dum. Not. Seroph. 7 (1824) = **Scrophularia**, Linn.
vernalis, Dum. l. c. = S. vernalis.

CERAMANTHUS, Hssk. Cat. Hort. Bogor. Alt. 240 (1844) = **Phyllanthus**, Linn. (Euphorbiac.).
gracilis, Hssk. l. c. = P. gracilis.

CERAMIA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 158 = **Erica**, Linn.
aurscularis, G. Don, Gen. Syst. iii. 806 = E. urceolaris.
blenna, G. Don, l. c. 807 = E. Vernix.
calathiflora, G. Don, l. c. = E. bicolor.
cordata, G. Don, l. c. = E. cordata.
helianthemifolia, G. Don, l. c. 806 = E. marifolia.
humifusa, G. Don, l. c. 807 = E. humifusa.
latifolia, G. Don, l. c. = E. latifolia.
marsfolia, G. Don, l. c. 806 = E. marifolia.
obliqua, G. Don, l. c. 807 = E. obliqua.
oxycoccifolia, G. Don, l. c. = E. oxycoccifolia.
planifolia, G. Don, l. c. 806 = E. planifolia.
serpyllifolia, G. Don, l. c. 807 = E. hispidula.
teucrifolia, G. Don, l. c. = E. teucrifolia.
thymifolia, G. Don, l. c. = E. thymifolia.
urceolaris, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 158 = E. urceolaria.

CERAMICALYX, Blume, Mus. Bot. Lugd. Bat. i. 50 (1849) = **Osbeckia**, Linn. (Melastomac.).
calcea, Blume, l. c. = O. rostrata !
pulchellus, Blume, l. c. = O. rostrata.
stellatus, Blume, l. c. = O. crinita.
ternifolius, Blume, l. c. = O. rostrata.

CERAMIOCEPHALUM, Sch. Bip. in Bull. Soc. Bot. Fr. ix. (1862) 284 = **Crepis**, Linn. (Compos.).
patulus, Sch. Bip. l. c. 285 = Crep. patula.

CERAMIUM, Blume, Bijdr. 1134 (1826) = **Bragantia**, Lour. (Aristolochiac.).
tomentosum, Blume, l. c. 1135 = B. tomentosa

CERAMOCARPUM, Nees, ex Meisn. in DC. Prod. xv. 1. 111 (1864) = **Cootes**, Anbl. (Laurin.).
cubense, Nees, l. c. 21 = Phoebe cubensis.
lanatum, Nees, l. c. 102 (= *Mesplodaphne lanata*).—Bras.
marginatum, Nees, l. c. 106 (= *Mespilodaphne marginata*)—Madag.
Maximilianum, Nees, l. c. 108 (= *Mespilodaphne prolifera*).—Bras.
obtusatum, Nees, l. c. 105 (= *Mespilodaphne heteromorpha*).—Ins. Maurit.
polyanthes, Nees, l. c. 133 = O. polyantha.
psychotrioides, Nees, l. c. 124 = O. psychotrioides.
semicompletum, Nees, l. c. 138 (= *Oreodaphne semicompleta*).—Bras.
umbrosum, Nees, l. c. 129 = O. umbrosa, *H. B. & K.*

CERAMOPHORA, Nees, ex Meisn. in DC. Prod. xv. 1. 105, 111 (1864) = **Cootes**, Anbl. (Laurin.).
cymosa, Nees, l. c. 105 (= *Mespilodaphne cymosa*).—Madag.

CERANTHE, Opiz, Seznam, 97 (1852) = **Cerinthe**, Tourn. (Borag.).

CERANTHERA, Beauv. Fl. Ow. et Ben. ii. 10 (1805) = **Alsodeia**, Thou. (Violar.).
dentata, Beauv. l. c. 11. t. 65 = A. gulanensis.
ilicifolia, Welw. in Trans. Linn. Soc. xxvii. (1869) 94. t. 2 = A. ilicifolia.
subintegrifolia, Beauv. Fl. Ow. et Ben. ii. 11. t. 66 = A. owariensis.

CERANTHERA, Ell. Sketch, ii. 93 (1824). *LABIATAE*, Benth. & Hook. f. ii. 1191.
DICRANDRA, Benth. in Bot. Reg. sub t. 1800 (1830).
densiflora, *A. Gray, Syn. Fl.* ii. ii. 365.—Florida.
linearifolia, *Ell. Sketch*, ii. 94.—Am. bor.

CERANTHERA, Endl. Gen. Suppl. ii. 29, in syn. (1842) = CERATANTHERA, Hornem. = **Colebrookia**, Sm. (Labiat.).

CERANTHERA, Rafin. in Am. Monthly Mag. (1818) 176 = **Solanum**, Tourn.
heterandra Rafin. l. c. = S. rostratum.

CERANTHUS, Schreb. Gen. i. 14 (1789) = **Linociera**, Sw. (Oleac.).
cornicularius, Schreb. ex Eichl. in Mart. Fl. Bras. vi. 1. 307 = L. compacta.
Schreberi, J. F. Gmel. Syst. 26 = L. compacta.

CERASEIDOS, Sieb. & Zucc. Abh. Akad. Muench. iii. (1843) 743. t. 5 = **Prunus**, Linn. (Rosac.).
apetala, Sieb. & Zucc. l. c. 744.—Japon.

CERASIOCARPUM, Hook. f. in Benth. & Hook. f. Gen. i. 882 (1867). *CUCURBITACEAE*, Benth. & Hook. f. i. c.
Bennettii, Cogn. in DC. Monog. Phan. iii. 729 = seylanicum.
Maingayi, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 629 = Melothria Rumphiana.
penangerense, C. B. Clarke, l. c. = Melothria Rumphiana.
seylanicum, *Hook. f. in Benth. & Hook. f. Gen.* i. 882. —Zeylan. ; Java.

CERASITES, Stead. Nom. ed. II. i. 528 (1840) = CERASTITES, S. F. Gray = **Papaver**, Tourn.

CERASOPHORA, Neck. Elem. ii. 71 (1790) = **Prunus**, Linn. (Rosac.).

CERASTITES, S. F. Gray, Nat. Arr. Brit. Pl. ii. 703 (1821) = **Papaver**, Tourn.
cambrica, S. F. Gray, l. c. 704 = Meconopsis cambrica.
dubia, S. F. Gray, l. c. = P. dubium.
hybrida, S. F. Gray, l. c. = P. hybridum.
lacuniata, S. F. Gray, l. c. = P. dubium.
macrocephala, S. F. Gray, l. c. 705 = P. Argemone.

CERASTIUM, [Dill.] Linn. Syst. ed. I (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 148.
CENTUNCULUS, Adans. Fam ii. 256 (1763).
DICHODON, Bartl. ex Reichb. Nom 205 (1841).
DUFOUREA, Gren. in Act. Soc. Linn. Bord. ix. (1837) 95.
ESMARCHIA, Reichb. Fl. Germ. Excurs 798 (1832).
LEUCODONIUM, Opiz, Seznam, 59 (1852).
MORNCHIA, Ehrh. Beitr. ii. 177 (1788).
MYOSOTIS, Tourn. ex Moench, Meth. 224 (1794).
PENTAPLE, Reichb. Ic. Fl. Germ. v. 37. t. 227 (1841).
PERVOTIA, Adans. Fam. ii. 256 (1763).
QUATERNELLA, Ehrh. Beitr. iv. 149 (1789).
acutifolium, Schur, Enum. Pl. Transs. 122 = ovatum.
adscendens, *Wender. ex Stend. Nom. ed.* II. i. 328.—Hab. ?
africanum, *Oliver, Fl. Trop. Afr.* i. 141.—Afr. trop.
aggregatum, Dur. ex Brignoli, in Flora, xxiii. (1840) 129 = siculum.
album, Presl, Fl. Sic. 167 = tomentosum.
alpestre, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 151 = triviale.
alpigenum, Schur, l. c. 149 = triviale.
alpinum, All. Fl. Pedem. ii. 116 = arvense.
alpinum, Bieb. Fl. Taur. Cauc. i. 361 = ovatum.
alpinum, Bunge, in Ledeb. Fl. Alt. ii. 180 = glomeratum.
alpinum, *Linn. Sp. Pl.* 438.—Reg. bor. frigid.
alsinifolium, Tausch, in Syll. Ratisb. ii. (1828) 243 = ovatum, arvense.
alsinoides, Pers. Syn. i. 521 = glutinosum, *Fries*.
ambigruum, Fisch. ex Ser. in DC. Prod. i. 419 = arvense.
amplexicaule, Sims, Bot. Mag. t. 1789 = dahuricum.
andinum, *Benth. Pl. Hartw.* 162.—N. Granat.
andinum, *Phil. in Anal. Univ. Chil.* (1862) ii. 816.—Chili.
androsaceum, Ser. in DC. Prod. i. 416 = illyricum.
angustifolium, Vitm. Summa Pl. iii. 187 = arvense.
annuinum, *Waldst. & Kit. ex Willd. Sp. Pl.* ii. 912; *et Pl. Rar. Hung.* i. 21.—Europ. ; Orient.
apetalum, Dum. Comm. Bot. 47 = glomeratum.
apricum, Schleeht. in Linnaea, xii. (1838) 208 = nutans.
apuanum, *Parl. in Nuov. Giorn. Bot. Ital.* vii. (1875) 69.—Italia.
aquaticum, Linn. Sp. Pl. 439 = Stellaria aquatica.
arabicum, *Fisch. & Mey. ex Heynh. Nom.* ii. 133, nomen.—Arab.
arabidis, *E. Mey. ex Fenzl, in Ann. Wien. Mus.* i. (1836) 240.—Afr. austr.
arcticum, Lange, *in Fl. Dan.* t. 2968.—Greenland.
arenarioides, Cranta, Inst. ii. 402 = Arenaria spathulata.
arenarium, Tenore, Fl. Nap. iv. 229 = semidecandrum.
arenosum, Kit. in Linnaea, xxxii. (1863) 518 = glomeratum !
argaeum, *Boiss. & Bal. Diagn.* Ser. II. vi. 38.—Cappadoc.
argenteum, Bieb. Fl. Taur. Cauc. i. 361 = grandiflorum.
armeniacum, *Gren. Monog. Cerast.* 19. t. 1.—Armenia.
arvense, Cham. & Schlecht. in Linnaea, v. (1830) 238 = orithales.
arvense, *Linn. Sp. Pl.* 438.—Reg. temp. bor. et austr.
arvensiforme, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 296.—Bolivia.
asiaticum, *Dur. in Duch. Rev. Bot.* ii. (1846-47) 437. —Hispan.
atratum, Lapeyr. Hist. Abr. Pl. Pyr. 265 = alpinum.
atrovirens, Rab. in Mag. Zool. & Bot. ii. (1838) 317. t. 9 = tetrandrum.
atticum, Boiss. & Heldr. Diagn. Ser. II. i. 93 = brachypetalum.
azoricum, *Hochst. ex Seub. Fl. Azor.* 45.—Ins. Azoric.
banaticum, Kit. in Rochel, Banat. Rar., in ind. = grandiflorum.
barbulatum, Wahlenb. Fl. Carp. 137 = brachypetalum.
berringianum, Cham. & Schlecht. in Linnaea, i. (1826) 69 = alpinum.
Bieberstenii, *DC. in Mem. Soc. Phys. Geneve.* i. (1823) 436.—Tauria.
biflorum, Crantz, Inst. ii. 402 = Stellaria biflora.
biflorum, *Kit. in Linnaea*, xxxii. (1863) 525.—Hungar.
blepharostemon, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 403 = longifolium.
Boissieri, *Gren. in Mem. Soc. Emul. Doubs*, i. (1861) 67.—Hispan.
bombycinum, Schur, Enum. Pl. Transs. 123 = alpinum.
brachycarpum, E. Mey. Zwei Pfl. Docum. 171 = Dregeanum.
brachycarpum, Schur, Enum. Pl. Transs. 124 = arvense.
brachypetalum, *Desp. in Pers. Syn.* i. 520.—Europ. ; Reg Mediterr.
brachypetalum, Stend. Nom. ed. II. i. 328, sphalm. = praec. ?
bracteatum, Rafin. Préc. Découv. 36 = oblongifolium !
brevifolium, Gillib. Fl. Lituan. ii. 158 = semidecandrum.
bulgaricum, *Uechtritz, in Oestr. Bot. Zeitschr.* xxvi. (1876) 221.—Bulgaria.
caeruleum, *Spach, Distrib. Diagn.* Ser. I. i. 53.—As Min.
caespitosum, Gillib. Fl. Lituan. ii. 159 = glomeratum.
caespitosum, Kit. ex Reichb. Fl. Germ. Excurs. 799 = arvense.
caespitosum, Malmgr. in Vet. Akad. Oefvers. xix. (1862) 242 = alpinum.
caespitosum, *Friesa & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 152.—N. Granat.
calycinum, Fenzl, Verbreit. Alsin. tab. ad p. 56 = Arenaria calycina.
campanulatum, *Viviani, in Ann. Bot.* i. ii. 171. t. 1. —Italia ; Hung. ; Graecia.
camphorataefolium, Stend. Nom. ed. I. 178, 179 = arvense.
candicans, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 295.—Ecuador.
canescens, Hornem. ex Ser. in DC. Prod. i. 416 = brachypetalum.
capense, *Sond. in Harv. & Sond. Fl. Cap.* i. 131.—Afr. austr.
cardiopetalum, C. Gay, Fl. Chil. i. 274 = Commersonianum.
carinthiacum, Vest, in Hoppe, Neues Taschenb. (1808) 229 = ovatum.
carnosulum, Turcz. ex Ledeb. Fl. Ross. i. 413 = arvense.
caucasicum, Fisch. ex Ser. in DC. Prod. i. 414 = tomentosum.
chilense, *Bartl. in Presl, Rel. Haenk.* ii. 17.—Chili.

CERASTIUM :—
chlorsefolium, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 34.—As. Min.
ciliatum, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 616 = glomeratum.
ciliatum, Waldst. & Kit. Pl. Rar. Hung. iii. 250 = arvense.
collinum, Hort. Dorpat. ex Flora, xvi. (1833) 122 = purpurascens.
collinum, Saliab. Prod. 300 = arvense.
Colmeanni, *Lahm. ex Spreng. Syst.* ii. 418.—Reg. Magell.
Columnæ, Tenore, Prod. Fl. Nap. p. xxvii = tomentosum.
Commersonianum, *Ser. in DC. Prod.* i. 417.—Reg. Argent.
Commerzonii, Stend. Nom. ed. II. i. 329 = præc.
connatum, Beck, N. et M. St. Bot. 55 = viscosum.
connatum, S. G. Gmel. It. ii. 196 ; ex Ledeb. Fl. Ross. i. 401 = davuricum.
consanguineum, *Wadd. in Ann. Sc. Nat.* Sér. V. i. (1864) 296.—Peruv.
constantinopolitanum, Nym. Consp. 106 = glomeratum.
cordatum, Crantz, Inst. ii. 400 = Drymaria cordata.
cordifolium, Roxb. Hort. Beng. 35 ; Fl. Ind. ii. 458 = Drymaria cordata.
cornicularium, Crantz, Inst. ii. 403 = dichotomum.
coronense, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 154 = ovatum.
corsicum, Soleir. ex Nym. Consp. 108 = arvense.
cossaipes, *Bartl. in Presl, Rel. Haenk.* ii. 18.—Peruv.
cuspidatum, *Hemsl. Diag. Pl. Nov.* 21.—Mexic.
davuricum, *Fisch. ex Spreng. Pugill.* ii. 65.—As. occ. et bor.
dasyphyllum, *Kit. in Linnæa*, xxxii. (1863) 521.—Hungar.
decalvans, Schloss, ex Nym. Consp. 107 = tomentosum.
deflexum, Ser. in DC. Prod. i. 417 = Stellaria aquatica.
densiflorum, *Guss. Suppl. Fl. Sic. Prod.* 126.—Sicilia.
Diezi, *Phil. in Anal. Univ. Chil.* (1862) ii. 391 ; *et in Linnæa*, xxxiii. (1864–65) 21.—Chili.
dichotomum, Crantz, Inst. ii. 401 = Stellaria dichotoma.
dichotomum, *Linn. Sp. Pl.* 438.—Reg. Mediterr. ; As. Min.
dichotomum, Schangin, in Pall. N. Nord. Beitr. vi. 98 = davuricum.
dicrotrichum, *Fenzl, ex Rohrb. in Mart. Fl. Bras.* xiv. ii. 281.—Bras.
diffusum, Pers. Syn. i. 520 = tetrandrum.
dioicum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 120.—Hispan.
divaricatum, *Horbich, in Flora*, vii. (1824) 184.—Neapol.
divaricatum, Kit. in Rochel, Pl. Banat. Rar. in ind. = Kitaibelii.
Dregeanum, *Fenzl, in Ann. Wien. Mus.* i. (1836) 841.—Afr. austr.
echinulatum, *Coss. & Dur. ex Debeaux, in Act. Linn. Soc. Bord.* xxiii. (1860) 180, nomen.—China.
elatum, Huet, ex Nym. Consp. 107 = grandiflorum.
elatum, Tenore, Fl. Nap. iv. 236 = tomentosum.
elbrusense, *Boiss. Fl. Orient.* i. 729.—Persia.
elegans, Fisch. ex Ser. in DC. Prod. i. 400 = trigynum.
elongatum, Bieb. Fl. Taur. Cauc. iii. 316 = nemorale.
elongatum, Pursh, Fl. Am. Sept. i. 221 = arvense.
Endresianum, Prol. ex Willk. & Lange, Prod. Fl. Hisp. iii. 637 = pyrenaicum.
eriophorum, Kit. ex Rochel, Pl. Banat. Rar. in ind. = alpinum.
erectum, Coss. & Germ. Fl. Env. Paris (1845) 89 = quaternellum.
falcatum, *Bunge, Vers. Supp. Fl. Alt.* 87.—Sibir. altaic.
fallax, Guss. Suppl. Fl. Sic. Prod. 129 = semidecandrum.
fasciculatum, *Bartl. in Presl, Rel. Haenk.* ii. 16.—Mexic.
filifolium, *Vest, in Flora*, iii. (1820) 358.—Styria.
filiforme, Schleich. ex Gaud. Fl. Helv. iii. 292, in syn. = latifolium.
filiforme, Ser. in DC. Prod. i. 420 = filifolium.
fimbriatum, Ledeb in Mém. Acad. Pétersb. v. (1815) 540 = Stellaria radians.
Fischeri, Cren. in Mém. Soc. Emul. Doubs, i. (1841) 3 = chlorsefolium.

CERASTIUM :—
Fischerianum, Ser. in DC. Prod. i. 419 = alpinum.
flaccidum, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 155, nomen.—Rossia.
flexuosum, Hegetschw. in Suter, Fl. Helvet. ed. II. App. 454 = latifolium.
floccosum, *Benth. Pl. Hartw.* 169.—N. Granat.
fontanum, Baumg. Enum. Stirp. Transs. i. 425 = triviale.
fragillimum, *Boiss. Diagn.* Ser. I. i. 54.—As. Min. ; Syria.
frigidum, Bieb. Fl. Taur. Cauc. i. 362 = purpurascens.
fulvum, Rafin. Préc. Découv. 36 = glomeratum.
furcatum, Cham. & Schlecht. in Linnæa, i. (1826) 61 = alpinum.
Gayanum, *Boiss. Diagn.* Ser. II. i. 92.—Hispan.
gibraltaricum, Boiss. Elench. 24 = Boissieri.
glabellum, Turcz. ex Ledeb. Fl. Ross. i. 413 = arvense.
glaberrimum, Lapeyr. Hist. Abr. Pl. Pyr. 265 = alpinum.
glabratum, Hartm. Handb. Skand. Fl. ed. I. 180 = alpinum.
glaciale, Gaud. ex Ser. in DC. Prod. i. 419 = latifolium.
glandulosum, *Schur, in Oestr. Bot. Zeitschr.* xix. (1869) 206.—Europ.
glaucum, Gren. in Mém. Soc. Emul. Doubs, i. (1841) 47 = manticum, quaternellum.
glomeratum, *Thuill. Fl. Par.* ed. II. 226.—Reg. temp. et subtrop.
glutinosum, *Fries, Nouv.* ed. II. 132.—Europ. ; As. bor.
glutinosum, Hegetschw. Fl. Schweiz, 488 = latifolium.
glutinosum, *H. B. & K. Nov. Gen. et Sp.* vi. 29.—N. Granat.
glutinosum, Nutt. Gen. Am. i. 291 = nutans.
gnaphaloides, Fenzl, in Russegg. Reis. i. 929 = tomentosum.
gracile, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 304 = Riseï.
gracile, Wallr. Beitr. Fl. Hercyn. 201 = brachypetalum.
gramineum, Crantz, Inst. ii. 401 = Stellaria gramínea.
grandiflorum, D. Don, Prod. Fl. Nep. 216 = glomeratum.
grandiflorum, Gilib. Fl. Lituan. ii. 159 = arvense.
grandiflorum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 168. —Europ. or. ; Reg. Caucas.
grandiflorum, Fourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 636 = Boissieri.
Grenieri, Fl. Schults, in Arch. Fl. Fr. et Allem. 24 = philinosum.
Hausknechtii, *Boiss & Haussk. in Oestr. Bot. Zeitschr.* xxvi. (1876) 411.—Mesopot.
heterophyllum, Schur, Enum. Pl. Transs. 121 = triviale.
heterophyllum, Viviani, Fl. Lib. Spec. 67 ; et Fl. Cors. Sp. Nov. 17 = illyricum.
hirsutum, Crantz, Inst. ii. 400 = Holosteum hirsutum.
hirsutum, Fisch. ex Ser. in DC. Prod. i. 419 = alpinum.
hirsutum, Muhl. Cat. 48 = glomeratum.
hirsutum, Tenore, Prod. Fl. Nap. p. xxvii = arvense.
hirtellum, *Pomel, Nouv. Mat. Fl. Atl.* 206.—Afr. bor.
Hochstetterianum, *Fenzl, Verbreit. Alsin.* tab. ad p. 56, nomen.—Hispan. ; Gall.
Holosteum, Garsault, Hort. Hafn. 484 = nemorale.
holosteiforme, Schur, Enum. Pl. Transs. 119 = semidecandrum.
holosteoides, Fries, Novit. ed. II. 126 = glomeratum.
holosteum, Crantz, Inst. ii. 401 = Stellaria Holostea.
holosteum, Fisch. ex Ser. in DC. Prod. i. 415 = dahuricum.
humifusum, *Camb. in St. Hil. Fl. Bras. Mer.* ii. 166.—Bras.
hybridum, Schur, Enum. Pl. Transs. 125 = tomentosum.
illyricum, *Ard. Animad.* ii. 26.—Græcia ; As. Min.
imbricatum, *H. B. & K. Nov. Gen. et Sp.* vi. 28.—Am. austr.
incanum, Hoffm. Hort. Mosq. 1808. n. 805 ; ex Bieb. Fl. Taur. Cauc. iii. 320 = grandiflorum.
incanum, Ledeb. in Mém. Acad. Pétersb. v. (1815) 540 = arvense.
indicum, *Wight & Arn. Prod.* 42.—Ind. or.
inflatum, *Link, ex Sweet, Hort. Brit.* ed. II. 67 ; Boiss. Fl. Orient. i. 781.—As. Min. ; Persia.
insubricum, Moretti, ex Reichb. Fl. Germ. Excurs. 799 = arvense.

CERASTIUM :—
juvenale, *Godr. in Mém. Acad. Stanislas*, Sér. III. [xvii.] 1853 (1854) 883.—Gallia !
Kablikianum, Wolfner, in Lotos, iv. (1884) 153 = arvense.
Kasbek, *Parrot, Reise Cauc.* ii. t. 8. f. 3.—Reg. Caucas.
Kitaibelii, *Steud. Nom.* ed. II. i. 329.—Bannatus.
Kotschyi, *Boiss. Fl. Orient.* i. 715.—Syria.
Lamottii, Le Grand, ex Nym. Consp. 110 = Riseï.
Lamyi, F. Schults, in Arch. Fl. Fr. et Allem. 24 = littigiosum.
lanatum, Lam. Encyc. i. 680 = alpinum.
lanifrons, Schur, Enum. Pl. Transs. 128 = alpinum.
lanigerum, Clem. in Atti Terza Rian. Sciens. Ital. (1841) 520 = tomentosum.
lanuginosum, Willd. ex Reichb. Fl. Germ. Excurs. 797 = alpinum.
lapponicum, Crantz, Inst. ii. 402 = trigynum.
laricifolium, Vill. Hist. Pl. Dauph. iii. 644 = arvense.
latifolium, Edmonst. in Phytolog. i. (1848) 498 = alpinum.
latifolium, Hartm. Skand. Fl. ed. XI. 239 = arcticum.
latifolium, Lapeyr. Hist. Abr. Pl. Pyr. 265 = pyrenaicum.
latifolium, Linn. Sp. Pl. 439.—Europ. ; Reg. Caucas.
laxiferum, *Schur, Enum. Pl. Transs.* 125.—Transsylvan.
laxum, *Boiss. & Heldr. Diagn.* Ser. I. vi. 25.—Græcia.
Ledebourianum, Ser. in DC. Prod. i. 420 = pilosum.
Lenaii, F. Schults, in Arch. Fl. Fr. et Allem. 24 = littigiosum.
Lerchenfeldianum, Schur, Enum. Pl. Transs. 122 = ovatum.
leguciense, Viv. Cat. Hort. Dinegro, 15 = campanulatum.
linearе, All. Fl. Pedem. ii. 265 = tomentosum.
lithospermifolium, Bunge, in Ledeb. Fl. Alt. ii. 179 = falcatum.
lithospermifolium, *Fisch. in Mém. Soc. Nat. Mosc.* iii (1812) 81.—Sibir.
Loeflingii, *De Lens, in Loisel. Fl. Gall.* i. 228.—Gallia.
Loeflingii, Crantz, Inst. ii. 408 = pentandrum.
longepedunculatum, Muhl. Cat. 47 = nutans.
longifolium, S. G. Gmel. ex Ledeb. Fl. Ross. i. 414 = grandiflorum.
longifolium, Juss. ex Poir. Encyc. Suppl. ii. 164 = Commersonianum.
longifolium, Tenore, Prod. Fl. Nap. p. xxvii = tomentosum.
longifolium, *Willd. Sp. Pl.* ii. 814.—Armenia.
longirostre, Wichura, in Jahresb. Schles. Ges. (1854) 75 = triviale.
lucorum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 150 = triviale.
luridum, Guss. Fl. Sic. Syn. i. 510 = brachypetalum.
macilentum, Aspegr. Blek. Fl. 84 = semidecandrum.
macranthum, *Boiss. Diagn.* Ser. I. v. 85.—As. Min.
macranthum, *Link, ex Steud. Nom.* ed. II. i. 329, nomen.—Hab. ?
macranthum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 14 = latifolium.
macrocarpum, *Boiss. & Haussk. in Boiss. Fl. Orient.* i. 716.—Mesopotam.
macrocarpum, Schur, Enum. Pl. Transs. 120 = triviale.
macrocarpum, Stev. ex Ledeb. Fl. Ross. i. 407 = purpurascens.
magellanicum, *Phil. in Anal. Univ. Chil.* (1872) 678. —Chili.
malachiforme, Schur, in Verh. Naturf. Ver. Brunn. xv. ii. (1877) 149 = triviale.
manticum, *Linn. Cent. Pl.* ii. 18.—Europ. austr.
matronse, Kit. ex Spreng. Pugill. i. 33 = arvense.
maurilanicum, *Pomel, Nouv. Mat. Fl. Atl.* 206.—Afr. bor.
mauritianum, Boston, ex Baker, Fl. Maurit. 14 = glomeratum.
maximum, Gilib. Fl. Lituan. ii. 158 = Stellaria aquatica.
maximum, *Linn. Sp. Pl.* 439.—Sibir. ; Am. sept.
medium, Crantz, Inst. ii. 401 = Spergularia media.
melanandrum, *Maxim. in Bull. Acad. Petersb.* xxvi. (1880) 490.—China.
membranaceum, Jacquem. ex Hook. f. Fl. Brit. Ind. i. 238 = glomeratum ?
Mersidi, *Faillon, in Mém. Soc. Acad. Maine-et-Loire*, xii. (1862) 87.—Gallia.

CERASTIUM:—
Meyerianum, *Rupr. Fl. Cauc.* 221, in textu.—Reg. Caucas.
micranthum, *C. A. Mey. ex Heynh. Nom.* ii. 138, nomen. —Rossia.
microcarpum, *Kit. ex Reichb. Fl Germ. Excurs.* 799. —Croatia.
microphyllum, Schur, Enum. Pl. Transs. 128 = alpinum.
microspermum, *C. A. Mey. Verz. Pfl. Cauc.* 222.— Reg. Caucas.
minus, Schur, in Oestr. Bot. Zeitschr. xxi. (1871) 46 = triviale.
minutulum, Desmoul. ex Steud. Nom. ed. II. i. 329 = glomeratum.
moesiacum, Frivald. in Flora, xix. (1836) 425 = tomentosum.
mollie, *Bartl. in Presl, Rel. Haenk.* ii. 17.—Mexic.
molle, Vill. Hist. Pl. Dauph. iii. 644 = arvense, tomentosum.
mollissimum, *Poir. Encyc. Suppl.* ii. 164.—Peruv.
montanum, *C. Gay, Fl. Chil.* i. 275.—Chili.
montiolides, *C. Gay, l. c.* 279 —Chili.
mosanum, *Cloet, ex Steud. Nom.* ed. II. i. 329.— Belgium.
mucronatum, *Wedd. in Ann Sc. Nat.* Sér. V. i. (1864) 294.—Peruv.; Boliv.
mx. ∂orum, *C. A. Mey. Verz. Pfl. Cauc.* 222.—Reg. Caucas.
murale, Desp. in DC. Fl. Fr. vi. 609 = triviale.
murale, Schur, Enum. Pl. Transs. 119 = glutinosum.
murale, Salisb. Prod. 300 = Holosteum umbellatum.
mutabile, Gren. in Mém. Soc. Emul. Doubs, i. (1841) 68 = arvense.
mutabile, Spenn. Fl. Friburg. iii. 847 = triviale, glomeratum, brachypetalum, semidecandrum.
nemorale, *Bieb. Fl. Taur. Cauc.* iii. 317.—Reg. Caucas.
nemorum, Crantz, Inst. ii. 401 = Stellaria nemorum.
neglectum, Wall. Cat. n. 6238 = glomeratum ?
nervosum, *C. Gay, Fl. Chil.* i. 277.—Chili.
nigrescens, Edmondst. ex Syme, Engl. Bot. ed. III. ii. 87 = alpinum.
nitens, Step. ex Ser. in DC. Prod. i. 415 = nemorale.
nivale, D. Don, ex Nym. Consp. 110 = trigynum.
nutans, *Raf. Prec. Decour.* 96.—Am. bor.
oblongifolium, Anders Cat. Pl. Nevad. 118 = nutans.
oblongifolium, Torr. Pacif. Rail. Rep. iv. 70 = pilosum.
oblongifolium, *Torr. Fl. U. St.* 460.—Am. bor.
obscurum, Chaub. in St. Amans, Fl. Agen. 180 = glutinosum, *Fries*.
obtusatum, *Kit. in Linnaea*, xxxii. (1863) 523.— Hungar.
obtusifolium, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 393.—Sibir. altaic.
obtusifolium, Lam. Fl. Fr. iii. 58 = glomeratum ?
octandrum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 45 = glomeratum.
Opisii, Seidl, in Opiz, Seznam, 28 = glutinosum.
orithales, *Schlecht. in Linnaea*, xii. (1838) 209.— Mexic.
orophyllum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 295.—Peruv.; Bolivia.
ovale, Bess. Prim. Fl Galic. i. 294 = glutinosum.
ovale, Pers. Syn. i. 521 = glomeratum ?
ovatum, *Hoppe, ex Willd. Enum. Hort. Berol.* 493.— Europ.; Reg. Caucas.
Pallasii, Vest, in Flora, iii. (1820) 356 = arvense.
pallens, F. Schultz, Arch. Fl. 124 = littigiosum.
pallidum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 146 = semidecandrum.
palustre, Moris, in Mem. Accad. Torino, xxxviii. (1835) p. xxvii = campanulatum.
pauciflorum, Bunge, in Ledeb. Fl. Alt. ii. 176 = lithospermifolium.
pauciflorum, Stev. ex Ser. in DC. Prod. i. 414 = pilosum.
pauciflorum, Turcz. ex Steud. Nom. ed. II. i. 330 = alpinum.
pedunculare, *Bory & Chaub. Nouv. Fl. Pélop.* 29. t. 13.—Graecia; As. Min.
pedunculatum, Gaud. ex Ser. in DC. Prod. i. 419 = latifolium.
pedunculatum, Willd. ex Steud. Nom. ed. II. i. 330 = natans.
pellucidum, Chaub. in St. Amans, Fl. Agen. 181 = semidecandrum.
pensylvanicum, Hook. Fl. Bor. Am. i. 104 = oblongifolium.
pensylvanicum, Hornem. Hort. Hafn. i. 335 = arvense.

CERASTIUM:—
pentandrum, Bieb. Fl. Taur. Cauc. i. 359 = glutinosum.
pentandrum, Krock. Fl. Siles. ii. i. 81 = semidecandrum.
pentandrum, *Linn. Sp. Pl.* 438.—Europ.
pentandrum, Mora, Fl. Sard. i. 265 = pumilum.
pentandrum, Tenore, Syll. 220 = densiflorum.
pepidisfolium, Kotschy, ex Boiss. Fl. Orient. i. 725 = fragilimanum.
perfoliatum, Falk, Beitr. ii. 182 = davuricum.
perfoliatum, *Linn. Sp. Pl.* 437.—Reg. Mediterr.; As. Min.
persicum, *Boiss. Diagn.* Ser. I. i. 54.—Persia.
petiolare, Hance, in Hook. Kew Journ. i. (1849) 143 = Stellaria aquatica.
petraeum, F. Schultz, Arch. Fl. 124 = glutinosum.
petricola, *Panč. Elem. Fl. Bulg.* 20.—Bulgaria.
petraeum, Schur, Enum. Pl. Transs. 122 = ovatum.
pilosellaefolium, *Kit. in Linnaea*, xxxi. (1862) 519.— Hungar.
pilosum, Fisch. ex Ledeb. Fl. Ross. i. 409 = glomeratum.
pilosum, Hornem. Hort. Hafn. ii. 965 = grandiflorum ?
pilosum, *Ledeb. in Mém. Acad. Petersb.* v. (1815) 539. —Sibir.; Am. bor. occ.
pilosum, Sibth. & Sm. Fl. Graec. Prod. i. 316 = illyricum.
pilosum, Tenore, Pr. d. Fl. Nap. p. xxvii = brachypetalum.
Pohlianum, Fenzl, ex Rohrb. in Mart. Fl. Bras. xiv. ii. 263 = dicrotrichum.
polymorphum, *Rupr. Fl. Cauc.* 237.—Reg. Caucas.
polyphyllum, Step. ex Ledeb. Fl. Ross. i. 418 = arvense.
praecox, Tenore, Prod. Fl. Nap. p xxvii = campanulatum.
procumbens, *Forsk. Fl. Aegypt. Arab.* 211.—Afr. bor
prostratum, Jan, Elench. 7, ex Bertol. Fl. Ital. iv. 756 = campanulatum.
pseudo-sylvaticum, Schur, in Oestr. Bot. Zeitschr. xxi. (1871) 46 = triviale.
pseudo-viscosum, Schur, l. c. 99 = glomeratum.
pubescens, Goldie, in Edinb. Phil. Journ. vi. (1822) 327 = oblongifolium.
pumilum, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 124 = Cavanillesii.
pumilum, *Curt. Fl. Lond. fasc.* vi. t. 30.—Europ.
pumilum, Griseb. Spicil. Fl. Rumel. i. 208 = glutinosum.
pumilum, Rafin. in Am. Monthly Mag. (1818) 266 = glomeratum.
purpurascens, *Adams, in Weber & Mohr, Beitr.* i. 60.— As. Min.; Reg. Caucas.
purpurascens, Fenzl, Verbreit. Alain. tab. ad p. 56 = Arenaria purpurascens.
pusillum, Ser. in DC. Prod. i. 418 = glomeratum.
pyrenaicum, *Y. Gay, in Ann. Sc. Nat.* Sér. I. xxvi. (1832) 321.—Mont Pyren.
quaternellum, *Fenzl, Verbreit. Alsin.* tab. ad p. 56.— Europ.; Am. bor.
racemosum, *Bartl. in Presl, Rel. Haenk.* ii. 18.—Chili.
radians, Crantz, Inst. ii. 401 = Stellaria radians.
ramigerum, *Bartl. in Presl, Rel. Haenk.* ii. 16.— Mexic.
ramosissimum, Boiss. Elench. 23 = Risei.
rectum, *Friwald. in Flora*, xix. (1836) 485.—Rumelia.
refractum, All. Fl. Pedem. ii. 117 = trigynum.
repens, Bertol. Fl. Ital. iv. 767 = sylvaticum.
repens, Bieb. Fl. Taur. Cauc. i. 360 = Biebersteinii.
repens, Boiss. Voy. Bot. Espagne, 105 = Boissieri.
repens, Sp. Pl. 439 = tomentosum.
revolutum, *Meinsh. in Baer & Helm. Beitr. Kenntn. Russ. Reich.* xxvi. 146.—Ins. Koraginsk.
Risei, *Des Moul. ex Y. Gay, in Ann. Sc. Nat.* Sér. II. vi. (1836) 848.—Europ.; As. Min.
rigidum, Ledeb. in Mém. Acad. Pétersb. v. (1815) 538 = alpinum.
rigidum, Vim. Summa Veg. iii. 137 = arvense.
Ripartianum, *F. Schultz, in Flora*, xiv. (1862) 458.— Mexic.
rivulare, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 166.— Bras.
Rossri, Boiss. & Heldr. Diagn. Ser. II. i. 93 = brachypetalum.
rotundatum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 146 = semidecandrum.
rotundifolium, Fisch. Cat. Hort. Gorenk. (1812) 58 = glomeratum.
rotundifolium, Schur, Enum. Pl. Transs. 123 = alpinum.
ruderale, *Bieb. Fl. Taur. Cauc.* i. 357.—Ross.; Reg. Caucas.

CERASTIUM:—
rupestre, Fisch. ex Ser. in DC. Prod. i. 417 = trigynum.
salatavicum, *Rupr. Fl. Cauc.* 220.—Reg. Caucas.
samnianum, Ser. in DC. Prod. i. 421 = grandiflorum.
sanatile, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 616 = alpinum.
sanctjohannis, Schur, in Oestr. Bot. Zeitschr. xxi. (1871) 100 = glutinosum.
scandens, *Lejeune, Fl. Spa*, i. 211.—Belgium.
scaposum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 104.— Ins. Creta.
Scarani, Tenore, Prod. Fl. Nap. p. xxvii = arvense.
Sellei, *Schlecht. ex Rohrb. in Mart. Fl. Bras.* xiv. ii. 288.—Bras.
semidecandrum, Cham. & Schlecht. in Linnaea, v. (1830) 238 = vulcanicum.
semidecandrum, Chaub. Expéd. Sc. Morée, 129 = brachypetalum.
semidecandrum, *Linn. Sp. Pl.* 438.—Europ.; As. Min.
semidecandrum, Sibth. & Sm. Fl. Graec. Prod. i. 315 = glutinosum.
sericeum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 354. —Arizona.
sericeum, *Pourr. in Mém. Acad. Toul.* iii. (1788) 311. —Europ.
serpyllifolium, Hieb. ex Ser. in DC. Prod. i. 417 = glomeratum.
serpyllifolium, Willd. Enum. Hort. Berol. Suppl. 26 = arvense.
sibiricum, Step. ex. Ledeb. Fl. Ross. i. 407 = alpinum.
sibiricum, Turcz. ex Ledeb. l. c. 410 = glomeratum.
siculum, *Guss. Suppl. Fl. Sic. Prod.* 127.—Sicilia.
simense, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 45 = glomeratum.
Soleirolii, Ser. ex Duby, Bot. Gall. i. 87 = arvense.
sorattense, *Rohrb. in Linnaea*, xxxvii. (1871–73) 291. —Bolivia.
spathulatum, *Pers. Syn.* i. 520.—Jamaica.
speciosum, Sprun. ex Boiss. Fl. Orient. i. 727 = grandiflorum.
Sprengelii, *Ser. in DC. Prod.* i. 421.—Hab.?
squalidum, Ram. in Mém. Acad. Paris, vi. (1826) 158 = alpinum.
Stellaria, *Tenore, ex Steud. Nom.* ed. II. i. 330.— Neapol.
stellarioides, Hartm. Handb. Skand. Fl. ed. I. 181 = trigynum.
stellarioides, Moç. ex Ser. in DC. Prod. i. 415 = pilosum.
stenopetalum, *Fenzl, ex Gren. & Godr. Fl.* i. 272.—Ins. Corsica.
strictum, Linn. Sp. Pl. 439 = arvense.
strigosum, Fries, Fl. Halland. 78 = brachypetalum.
subacaule, Hegetschw. in Soter, Fl. Helvet. ed. II. App. 455 = latifolium.
subspicatum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 295.—Peruv.
cuculentum, Crantz, Inst. ii. 400 = Arenaria peploides.
suffruticosum, *Lam. Encyc.* i. 681.—Gallop.; As Min.
suffruticosum, Linn. Sp Pl. 439 = arvense.
sylvaticum, Opiz, Seznam, 28 = triviale.
sylvaticum, Stev. ex Ledeb. Fl. Ross. i. 410 = glomeratum.
sylvaticum, *Waldst. & Kit. Pl. Rar. Hung.* i. 100.— Austria.
szalabaense, *Kit. in Linnaea*, xxxii. (1863) 524.— Hungar.
Szowitsii, *Boiss. Fl. Orient.* i. 717.—Armenia.
tauricum, Spreng. ex Ser. in DC. Prod. i. 415 = brachypetalum.
Tauschianum, *Wolfner, in Lotos*, iv. (1854) 57.— Europ.
tenellum, Gaud. ex Ser. in DC. Prod. i. 416 = glomeratum.
Tenoreanum, Ser. l. c. 421 = brachypetalum.
tenue, Viviani, Fl. Cors. Sp. Nov. 7 = quaternellum.
tenuifolium, Pursh, Fl. Am. Sept. i. 221 = arvense.
tenuifolium, Spreng. in Horsem. Hort. Hafn. Suppl. 138 = Sprengelii.
tenuifolium, Viviani, ex Reichb. Fl. Germ. Excurs. 799 = grandiflorum.
tetragonum, Bunge. Enum. Stirp. Transs. i. 424 = Stellaria crassifolia.
tetrandrum, *Curt. Fl. Lond. fasc.* vi. t. 31.—Europ.
Thomasii, *Tenore, Fl. Neap. Prod. App.* iv. 21.— Europ.
Thomsonii, *Hook. f. Fl. Brit. Ind.* i. 238.—Reg. Himal.
tmoleum, Boiss. Diagn. Ser. I. v. 86 = fragilimum.

CERASTIUM :—
tomentosum, Boj. Hort. Maurit. 24 = glomeratum.
tomentosum, Georgi, Beschr. Russ. Reich. iv. 988 = Biebersteinii.
tomentosum, Hasensel. ex Willk. & Lange, Prod. Fl. Hisp. iii. 686 = Boissieri.
tomentosum, Huds. Fl. Angl. ed. I. 176 = alpinum.
tomentosum, *Linn. Sp. Pl.* 440.—Europ.; As. Min.
Tournefortii, *Gren. in Mém. Soc. Emul. Doubs*, i. (1841) 13. t. 7 —Armenia.
transilvanicum, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1858) 14.—Transylv.
trigynum, *Vill. Prosp.* 45 ; *Hist. Pl. Dauph*, iii. 645.—Reg. hor. temp. et arct.
triviale, *Link, Enum. Hort Berol.* i. 488.—Europ.; As. ; Afr. bor.
uliginosum, Hegetschw. Fl. Schweiz, 437 = triviale ?
umbellatum, Crantz, Inst. ii. 401 = Holosteum umbellatum.
umbellatum, *C. Koch, in Linnaea*, xv. (1841) 708.—Armenia.
umbrosum, Kit. in Linnaea, xxxii. (1863) 515 = triviale ?
uniflorum, Thom. ex Reichb. Fl. Germ. Excurs. 797 = latifolium.
vagans, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimp. 26.—Ins. Madera.
valachicum, *Kit. in Linnaea*, xxxii. (1863) 519.—Hungar.
varians, Coss. & Germ. Obs. Pl. Crit. 26 = glutinosum *Pers.*, semidecandrum.
velutinum, Rafin. in Med. Repos. N. York, v. (1808) 359 = vulgatum ?
Vidali, *Phil. in Anal. Univ. Chil.* (1872) 677.—Chili.
villosum, Baumg. Enum. Stirp. Transs. i. 424 = alpinum.
villosum, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 329 = glomeratum.
viscarium, Schur, Enum. Pl. Transs. 118, in syn. = glutinosum.
viscosum, Link, Enum. Hort. Berol. i. 488 = semidecandrum.
viscosissimum, Schur, Enum. Pl. Transs. 123 = alpinum.
viscosum, Linn. Sp. Pl. 437 = glomeratum.
viscosum, Pollich, Hist. Pl. Palat. i. 448 = brachypetalum.
vulcanicum, *Schlecht. in Linnaea*, xii. (1838) 208.—Mexic.
vulgare, Hartm. Handb. Skand. Fl. ed. I. 182 = triviale.
vulgatum, Linn. Fl. Suec. ii. 158 = glomeratum.
Wahlenbergii, Rupr. Fl. Caucas. 287 = latifolium.
Willdenowii, *H. B. & K. Nov. Gen. et Sp.* vi. 29.—Ecuador.

CERASUS, (Tourn.) Linn. Syst. ed. I (1735) ; Gen. ed. I. 141 (1737) = Prunus, Linn (Rosac.).
acida, Borck. in Roem. Arch. i. ii. 88 = P. Cerasus.
acuminata, Wall. Pl. As. Rar. ii. 78 = P. acuminata.
admophila, M. Roem. Syn. Rosifl. 81 = P. undulata.
affinis, Sweet, Hort. Brit. ed. III. 308.—Hab. ?
americana, Hook. & Arn. in Comp. Bot. Mag. i. (1835) 24 = P. americana.
aspera, Loisel. in Duham. Arb. ed. nov. v. 33 = P. aspera.
asplenifolia, Hort. ex C. Koch, Dendrol. i. 106 = P. Avium.
austera, Borck. in Roem. Arch. i. ii. 88 = P. Cerasus.
avicularis, Dulac, Fl. Hautes-Pyr. 801 = P. Avium.
Avium, Moench, Meth. 672 = P. Avium.
baroniensis, Nym. Consp. 218 = P. Cerasus.
bigarella, Dum. Fl. Belg. 91 = P. Cerasus.
borealis, Michx. Fl. Bor. Am. i. 286 = P. pensylvanica.
brachypetala, Bolss. Diagn. Ser. I. vi. 93 = *incisa*.
brasiliensis, Cham. & Schlecht. in Linnaea, ii. (1827) 549 = P. brasiliensis.
Bungei, Walp. Rep. ii. 9 = P. Cerasus.
campestris, Gueldenst. ex Ledeb. Fl. Ross. ii. 7 = P. Chamaecerasus.
canadensis, Loisel. in Duham. Arb. ed. nov. v. 8 = P. americana.
canadensis, Mill. Gard. Dict. ed. VIII. n. 5 = P. Chamaecerasus.
Capollin, Ser. in DC. Prod. ii. 539 = P. Capollin.
capricida, Wall. Cat. n. 718 = P. undulata.
Caproniana, DC. Fl. Fr. iv. 493 = P. Cerasus.

CERASUS :—
Capuli, Ser. in DC. Prod. ii. 541 = Prunus Capollin.
carolinicna, Michx. Fl. Bor. Am. i. 285 = P. caroliniana.
cerasifera, Eaton & Wright, N. Am. Bot. 189 = P. maritima.
Chamaecerasus, Loisel. in Duham. Arb. ed. nov. v. 29 = P. Chamaecerasus.
Chicasa, Ser. in DC. Prod. ii. 538 = P. Chicasa.
chinensis, G. Don, Gen. Syst. ii. 514 = P. japonica.
collina, Lej. & Court. Comp. Fl. Belg. ii. 130 = P. Cerasus.
cornuta, Wall. Cat. n. 716 = P. Padus.
corymbosa, St. Leg. in Ann. Soc. Bot. Lyon, vii. (1880) 109 = P. Mahaleb.
cuculiata, Hort. ex C. Koch, Dendrol. i. 111 = P. Cerasus.
decumana, Mordant, ex DC. Prod. ii. 536 = P. Avium.
demissa, Nutt. in Torr. & Gray, Fl. N. Am. i. 411 = P. demissa.
densiflora, Spach, Hist. Vég. Phan. i. 415 = P. virginiana.
depressa, Ser. in DC. Prod. ii. 538 = P. pumila.
diffusa, Bolss. & Haussk. ex Bolss. Fl. Orient. ii. 647.—Persia.
domestica, Cat. Perck, ex Wesmael, in Bull. Congr. Bot. Brux. (1864) 255 = P. Cerasus.
Duerrissii, Mart. Bull. Acad. Belg. viii. i. (1841) 68 = P. virginiana.
dulcis, Borck. ex Steud. Nom. ed. II. i. 331 = P. Avium.
dulcis, Gaertn. Mey. & Scherb. Fl. Wett. ii. 181 = P. Cerasus.
duracina, DC. Fl. Fr. iv. 488 = P. Avium.
ebroiensis, Lamotte, Prod. i. 228 = P. Cerasus.
effusa, Lam. Fl. Austriac. ii. 6 = P. Cerasus.
elliptica, Loisel. in Duham. Arb. ed. nov. v. 4 = P. elliptica.
emarginata, Dougl. ex Hook. Fl. Bor. Am. i. 169 = P. emarginata.
erecta, Presl, Epim. Bot. 194 = P. emarginata.
ferruginea, DC. Prod. ii. 540 = P. ferruginea.
fimbriata, Spach, Hist. Vég. Phan. i. 416 = P. virginiana.
floribunda, Hort. ex C. Koch, Dendrol. i. 119 = P. pumila.
Fontanesiana, Spach, Hist. Vég. Phan. i. 410 = P. gracca.
fruticosa, Pall. Fl. Ross. i. 8. = P. Chamaecerasus.
glandulosa, Kellogg, in Proc. Calif. Acad. i. 1855 (1873) 59 = P. emarginata.
glandulosa, Loisel. in Duham. Arb. ed. nov. v. 33 = P. japonica.
glauca, Moench, Meth. 672 = P. pumila.
glaucifolia, Wall. Cat. n. 717 = P. nepaulensis.
graeca, Hort. ex C. Koch, Dendrol. i. 105, Index = P. graeca.
Griffithii, Bolss. Fl. Orient. ii. 648.—Afghanist.
Hesamiana, M. Roem. Syn. Rosifl. 69 = P. Cerasus.
heterophylla, Hort. ex C. Koch, Dendrol. i. 106 = P. Avium.
hirsuta, Spach, Hist. Vég. Phan. i. 417 = P. virginiana.
Hostii, C. Sm. ex Spach, Hist. Nat. Veg. i. 419 = P. Hixa.
hortensis, Mill. Gard. Dict. ed. VIII. n. 3 = P. Cerasus.
humilis, Host, Fl. Austr. i. 7 = P. Chamaecerasus.
humilis, Moris, Elench. Sard. 17 = P. prostrata.
hyemalis, Ser. in DC. Prod. ii. 538 = P. americana.
ilicifolius, Nutt. ex Hook. & Arn. Bot. Beech. Voy. 340. 88 = P. ilicifolia.
incana, Bolss. Fl. Orient. ii. 647 = P. incana.
incana, Spach, in Ann. Sc. Nat. Ser. II. xix. (1843) 126 = P. prostrata.
incisa, Bolss. Fl. Orient. ii. 648.—Persia.
incisa, Loisel. in Duham. Arb. ed. nov. v. 33 = P. incisa.
integerrima, Wall. Cat. n. 723 = P. acuminata.
integrifolia, Presl, Epim. Bot. 194.—Peruv.
intermedia, Host, Fl. Austriac. ii. 7 = P. Avium.
intermedia, Loisel. in Duham. Arb. ed. nov. v. 30 = P. Chamaecerasus.
japonica, Ker-Gawl. ex Steud. Nom. ed. II. i. 331 = P. Kerii.
japonica, Loisel. in Duham. Arb. ed. nov. v. 33 = P. japonica.
javanica, Teijsm. & Binn. in Nederl. Kruidk. Arch. iii. (1855) 412 = P. javanica.
Jenkinsii, Hook. f. & Thoms. ex Fl. Brit. Ind. ii. 317 = P. Jenkinsii.
Juliana, DC. Fl. Fr. iv. 493 = P. Avium.

CERASUS :—
Juliana, Hort. ex Steud. Nom. ed. II. i. 331 = Prunus Cerasus.
Laurocerasus, Carr. in Rev. Hortic. (1872) 198.—Japon.
Laurocerasus, Loisel. in Duham. Arb. ed. nov. v. 6 = P. Laurocerasus.
Lindleyana, Wall. Cat. n. 1008 = P. nepaulensis.
littoralis, Eaton & Wright, N. Am. Bot. 189 = P. maritima.
lusitanica, Loisel. in Duham. Arb. ed. nov. v. 5 = P. lusitanica.
macedonica, M. Roem. Syn. Rosifl. 64 = P. Chamaecerasus.
macrophylla, Sweet, Hort. Brit. ed. I. 485 = P. Avium.
mahaleb, Hort. ex Lavall. Arbor. Segrez. 119 = P. paniculata.
Mahaleb, Mill. Gard. Dict. ed. VIII. n. 4 = P. Mahaleb.
Marasca, Host, Fl. Austriac. ii. 6 = P. Cerasus.
martabanica, Wall. Cat. n. 4902 = P. martabanica.
micrantha, Spach, Hist. Vég. Phan. i. 414 = P. virginiana.
microcarpa, Bolss. Fl. Orient. ii. 646 = P. microcarpa.
minutiflora, A. Gray, Pl. Wright. i. 68 = P. minutiflora.
mollis, Doogl. in Hook. Fl. Bor. Am. i. 169 = P. emarginata.
multicarpa, Hort. ex Rev. Hortic. (1875) 409 = P. Cerasus.
napaulensis, Ser. in DC. Prod. ii. 540 = P. nepaulensis.
nicotianaefolia, Hort. ex DC. Prod. ii. 536 = P. Cerasus.
nigra, Loisel. in Duham. Arb. ed. nov. v. 32 = P. americana.
nigra, Mill. Gard. Dict. ed. VIII. n. 2 = P. Avium.
nigricans, Borck. in Roem. Arch. i. ii. 88 = P. Avium.
obovata, Beck, N. et M. St. Bot. 37 = P. virginiana.
occidentalis, Loisel. in Duham. Arb. ed. nov. v. 4 = P. occidentalis.
opaca, Benth. Pl. Hartw. 172 = P. opaca.
orientalis, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 128 = P. microcarpa.
oxyporma, Spach, Hist. Vég. Phan. i. 418 = P. Padus.
Padus, Delarb. Fl. Auv. ed. II. 323 ; DC. Fl. Fr. iv. 480 = P. Padus.
pallida, M. Roem. Syn. Rosifl. 69 = P. Avium.
paniculata, Loisel. in Duham. Arb. ed. nov. v. 7 = P. paniculata.
Pattoniana, Carr. in Rev. Hortic. (1872) 135 = P. mollis.
pectinata, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 127 = P. pectinata.
pendula, Sweet, Hort. Brit. ed. I. 485 = P. fruticosa.
pensylvanica, Loisel. in Duham. Arb. ed. nov. v. 9 = P. pensylvanica.
persicifolia, Loisel. i. c. = P. pensylvanica.
Pheshia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 239 = P. Puddum.
polygyna, DC. ex Ser. in DC. Prod. ii. 537 = P. Cerasus.
prostrata, Hort. ex C. Koch, Dendrol. i. 119 = P. pumila.
prostrata, Ser. in DC. Prod. ii. 538 = P. prostrata.
Pseudocerasus, G. Don, in Loud. Hort. Brit. 200 = P. Puddum.
pubescens, Ser. in DC. Prod. ii. 536 = P. maritima.
Puddum, Wall. Pl. As. Rar. ii. 87. t. 148 = P. Puddum.
pumila, Borck. in Roem. Arch. i. ii. (1797) 88 = P. Chamaecerasus.
pumila, Michx. Fl. Bor. Am. ii. 286 = P. pumila.
punctata, Hook. f. & Thoms. ex Fl. Brit. Ind. ii. 317 = P. punctata.
pygmaea, Loisel. in Duham. Arb. ed. nov. v. 32 = P. maritima.
racemosa, Gilib. Exercit. i. 349 = P. Padus.
reflexa, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 345 = P. reflexa.
Rhexii, Hort. Gall. ex Van Houtte, Fl. des Serres, Sér. II. vii. (1857–58) 159 = P. Cerasus.
rosea, Hort. ex Steud. Nom. ed. II. i. 331 = P. Cerasus.
rubicunda, Bechst. Forstb. 190, 256 = P. Avium.
rubra, Gilib. Fl. Lituan. ii. 230 = P. Cerasus.
rufa, Wall. Cat. n. 721 = P. rufa.
salicifolia, Hort. ex C. Koch, Dendrol. i. 106 = P. Avium.
salicifolia, Ser. in DC. Prod. ii. 540 = P. salicifolia.

CERASUS :—
salicina, [G. Don, in] Loud. Hort. Brit. 480 = Prunus salicina.
semperfoides, Walp. Rep. iv. Index, 311 = P. semperfoides.
semperflorens, Borckh. in Roem. Arch. i. ii. 38 = P. Cerasus.
serotina, Hook. Fl. Bor. Am. i. 169 = P. virginiana.
serotina, Loisel. in Duham. Arb. ed. nov. v. 3 = P. serotina.
serrulatus, [G. Don, in] Loud. Hort. Brit. 480 = P. paniculata.
sibirica, Hort. ex C. Koch, Dendrol. i. 114 = P. fruticosa.
Sieboldii, Carr. in Rev. Hortic. (1866) 279. ; 370 = P. paniculata.
sinensis, Spach, Hist. Vég. Phan. i. 425 = P. sinensis.
sphærocarpa, Loisel. in Duham. Arb. ed. nov. v. 4 = P. sphærocarpa.
Susquehannæ, Sweet, Hort. Brit. ed. I. 485 = P. pumila.
sylvestris, S. G. Gmel. It. i. 102 = P. Chamæcerasus.
tomentosa, Wall. Cat. n. 715 = P. tomentosa.
tortuosa, Boiss. & Heusak. ex Boiss. Fl. Orient. ii. 647. —As. Min. ; Persia.
tridentina, M. Roem. Syn. Rosifl. 76 = P. Cerasus.
trifora, Wall. ex Steud. Nom. ed. II. i. 331 = P. trifora.
umbellata, Torr. & Gray, Fl. N. Am. i. 409 = Prunus umbellata.
undulata, Ser. in DC. Prod. ii. 540 = P. undulata.
varia, Borckh. in Roem. Arch. i. ii. 38 = P. Avium.
virginiana, Michx. Fl. Bor. Am. i. 285 = P. serotina.
vurginica, Michx. ex Steud. Nom. ed II. i. 331 = P. virginiana.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 1 = P. Cerasus.
Wallichii, M. Roem. Syn. Rosifl. 81 = P. acuminata.
Watsonii, Hort. ex Lavall. Arbor. Segrez. 119 = P. Puddum.

CERATANDRA, Eckl. ex Bauer, Illustr. Gen. Orch. t. 16 (1837). *ORCHIDEÆ*, Benth. & Hook. f. iii. 634.
affinis, *Sond. in Linnæa*, xix. (1847) 108.—Afr. austr.
auriculata, Lindl. Gen. et Sp. Orch. 364 = chloroleuca.
bicolor, *Sond. ex Dreège, in Linnæa*, xx. (1847) 220, nomen.—Afr. austr.
chloroleuca, *Eckl. ex Bauer, Illustr. Gen. Orch.* t. 16. —Afr. austr.
globosa, *Lindl. Gen. et Sp. Orch.* 364.—Afr. austr.
grandiflora, *Lindl. l. c.*—Afr. austr.
Harveyana, *Lindl. l. c.* 366.—Afr. austr.
Harveyana, Sond. in Linnæa, xix. (1847) 108 = bicolor.
ochroleuca, G. Don, in Sweet, Hort. Brit. ed. III. 651 = chloroleuca.
parviflora, *Lindl. Gen. et Sp. Orch.* 364.—Afr. austr.

CERATANTHERA, Hornem. Hort. Hafn. i. 2, 5 (1813) ; Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 355, 361 = **Globba**, Linn. (Scitamin.).
amomoides, Hornem. l. c. i. 5 = G. marantina.
pendula, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 355 = G. pendula.

CERATELLA, Hook. f. Fl. Antarct. i. 25. t. 18 (1845) = **Abrotanella**, Cass. (Compos.).
rosulata, Hook. f. l. c. = A. rosulata.

CERATIA, Adans. Fam. ii. 319 (1763) = **Ceratonia**, Linn. (Leguminos.).

CERATIOLA, Michx. Fl. Bor. Am. ii. 221 (1808). *EMPETRACEÆ*, Benth. & Hook. f. iii. 415.
ericoides, *Michx. l. c.* 222.—Am. bor.

CERATIOSICYOS, Ness, in Eckl. & Zeyh. Enum. 261 (1836). *PASSIFLOREÆ*, Benth. & Hook. f. i. 814.
Ecklonii, *Ness, l. c.*—Afr. austr.

CERATOSYCIOS, Walp. Rep. ii. 222 (1843) = prasc.

CERATIUM, Blume, Bijdr. 341. t. 46 (1825) = **Iria**, Lindl. (Orchid.).
compressum, Blume, l. c. 342 = E. compressa.

CERATOCALYX, Cass. in Ann. Sc. Nat. Sér. III. ix. (1848) 145. t. 10 = **Orobanche**, Linn.
fimbriata, Lange, in Kjoeb. Vidensk. Meddel. (1863) 52.—Hispan.
macrolepis, Cass. in Ann. Sc. Nat. Sér. III. ix. (1848) 146. t. 10 = O. macrolepis.
Reuteri, Lange, in Kjoeb. Vidensk. Meddel. (1863) 52, in obs. = O. gamosepala.

CERATOCAPNOS, Dur. in Parlat. Giorn. Bot. Ital. i. i. (1844) 336 = **Corydalis**, Vent. (Papaverac.).
heterocarpa, Dur. l. c. = C. heterocarpa.
palæstina, Boiss. Diagn. Ser. I. viii. 12.—Syria.
umbrosa, Walp. Ann. ii. 30 = Cor. heterocarpa.

CERATOCARPUS, Burb. ex Linn. Diss. Dass. 28 (1747) ; Amoen. Acad. i. 412 (1749). *CHENOPODIACEÆ*, Benth. & Hook. f. ii. 55.
arenarius, *Linn. Sp. Pl.* 969.—Sibir.
maritimus, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. ii. 357 = Atriplex pedunculata.
salinus, Pall. Reise Statth. Russ. Reich. i. 268, 370 = Atriplex pedunculata.

CERATOCARPUS, Dur. (lapsu) in Parlat. Giorn. Bot. Ital. i. i. (1844) 336 = CERATOCAPNOS, Dur. = **Corydalis**, Vent.

CERATOCARYUM, Ness, in Lindl Nat. Syst. ed. II. 451 (1836). *RESTIACEÆ*, Benth. & Hook. f iii. 1087.
argenteum, *Ness, l. c.*; *Kunth, Enum. Pl.* iii. 483 — Afr. austr.
fatuloxum, *Mart. in Journ. Linn. Soc.* x. (1869) 274 —Afr. austr.
speciosum, Ness, ex Mast. l. c. 273 = argenteum

CERATOCAULOS, Reichb. Handb. 201 (1837) = **Datura**, Linn. (Solanac.).
daturoides, Spach, Hist. Veg. Phan. ix. 69 = D. Ceratocaula.

CERATOCEPHALUS, Moench, Meth. 218 (1794) (Ceratocephala) = **Ranunculus**, Linn.
falcatus, Pers. Syn. i. 341 = R. falcatus
furfuraceus, Pomel, Nouv. Mat. Fl. Atl. 248.—Afr. bor.
incurvus, Stev. in Bull. Soc. Nat. Mosc. xxi. ii. (1848) 269. t. 7 = R. falcatus.
leucorpus, Stev. l. c. = R. falcatus.
orthocerus, DC. Syst. i. 231 = R. falcatus.
platyceras, Stev. in Bull. Soc. Nat. Mosc. xxi. ii. (1848) 269 = R. falcatus
reflexus, Stev. l. c. 268 = R. falcatus.
spicatus, Moench, Meth. 218 = R. falcatus.
syriacus, Stev. in Bull. Soc. Nat. Mosc. xxi. ii. (1848) 268 = R. falcatus.
testiculatus, Roth, Enum. Pl. Phan. Germ. i. i. 1014 = R. falcatus.

CERATOCEPHALUS, Vaill. ex Cass. in Dict. Sc. Nat. vii. 432 (1817) = **Bidens**, Linn. (Compos.).
pilosus, Rich. ex Cass. l. c. = B. pilosa.

CERATOCHILUS, Blume, Bijdr. 358. t. 25 (1825) = **Saccolabium**, Blume (Orchid.).
biglandulosum, Blume, l. c. 359 = S. distichum ?
micranthum, Lindl. Gen. et Sp. Orch. 232 = S. micranthum.
orchideus, Lindl. l. c.—Ind. or.

CERATOCHILUS, Lindl. in Lodd. Bot. Cab. t. 1414 (1828) = **Stanhopea**, Frost (Orchid.).
grandiflorus, Lodd. l. c. = S. eburnea.
insignis, Lindl. ex Steud. Nom. ed. II. i. 331 = S. insignis.
oculatus, Lodd. Bot. Cab. t. 1764 = S. oculata.

CERATOCHLOA, Beauv. Agrost. 75. t. 15. f. 7 (1812) = **Bromus**, Dill. (Gramin.).
australis, Spreng. ex Steud. Nom. ed. II. i. 332 ; 229 = B. unioloides.
Boryana, Roem. & Schult. Syst. ii. 597 = B. strictus.
breviaristata, Hook. Fl. Bor. Am. ii. 253 = B. breviaristatus.
brevis, Ness, ex Steud. Syn. Fl. Germ. 336 = B. brevis.
festucoides, Beauv. Agrost. 75 = B. unioloides.

CERATOCHLOA :—
grandiflora, Hook. Fl. Bor. Am. ii. 253 = Bromus virens.
Haenkeana, J. Presl, in Rel. Haenk. i. 285 = B. unioloides.
marginata, Nees, ex Steud. Syn. Pl. Gram. 322 = B. ciliatus.
pendula, Schrad. Ind. Sem. Hort. Gottng. (1830) ; Linnæa, vi. (1831) Litt. 72 = B. unioloides.
secunda, J. Presl, in Rel. Haenk. i. 285 = B. Preslii.
unioloides, DC. Cat. Hort. Monsp. 92 = B. unioloides.

CERATOCNEMUM, Coss. & Balansa, in Bull. Soc. Bot. Fr. xx. (1873) 289. *CRUCIFERÆ*.
rapistroides, *Coss. & Balansa, l. c.*—Marocco.

CERATOCOCCA, Willd. ex Roem. & Schult. Syst. vi. p. lxx (1820) = **Microtea**, Sw. (Phytolaccac.).
mayurensis, Humb. & Bonpl. ex Roem. & Schult. l. c. 800 = M. mayurensis.

CERATOCOCCUS, Meissn. Gen. Comm. 369 (1843) = **Plukenetia**, Linn. (Euphorbiac.).

CERATOGONON, Meissn. in Wall. Pl. As. Rar. iii. 68 (1832) = **Oxygonum**, Burch. (Polygonac.).
atriplicifolium, Meisn. l. c.—Afr. trop.
cordofanum, Meisn. in DC. Prod. xiv. 39 = *atriplicifolium*.
sinuatum, Hochst & Steud. ex A. Rich. Tent. Fl. Abyss. ii. 231 = *atriplicifolium*.

CERATOGYNE, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 68. *COMPOSITÆ*, Benth. & Hook. f. ii. 481.
DIOTOSPERMA, A. Gray, in Hook Kew Journ. iv. (1852) 275.
obionoides, *Turcz. l. c.* 69.—Austral.

CERATOGYNUM, Wight, Ic. v. ii. 26. t. 1900 (1852) = **Sauropus**, Blume (Euphorbiac.).
concolor, Pritz. Ic. Ind. 245, sphalm. = Ceratolobus concolor.
rhamnoides, Wight, Ic. v. ii. 26. t. 1900 = S. quadrangularis.

CERATOLACIS, Wedd. in DC. Prod. xvii. 68 (1873) *PODOSTEMACEÆ*, Benth. & Hook. f. iii. 118.
erythrolichen, *Wedd. l. c.*—Bras.

CERATOLEPIS, Cass. in Bull. Soc. Philom. (1819) 111 = **Pamphalea**, DC. (Compos.).

CERATOLOBUS, Blume, in Schult. f. Syst. vii. p. lxxx (1830). *PALMÆ*, Benth. & Hook. f. iii. 988.
concolor, *Blume, ex Rumphia*, ii. 165. t. 130. f. 1.—Sumatra.
glaucescens, *Blume, in Schult. f. Syst.* vii. 1384.—Java.
glaucescens, Blume, Rumphia, ii. t. 137 = Korthalsiæ sp. *plicatus*, Zipp. ex Blume, l. c. 171 = Korthalsia Zippelii.
Zippelii, Blume, l. c. = Korthalsia Zippelii.

CERATONIA, Linn. Gen. ed. I. 35 (1754). *LEGUMINOSÆ*, Benth. & Hook. f. i. 574.
CERATIA, Adans. Fam. ii. 319 (1763).
chilensis, Molina, Sagg. Chil. 172 = Prosopis julliflora.
coriacea, Salisb. Prod. 338 = C. Siliqua.
inermis, Stokes, Bot. Mat. Med. i. 487 = C. Siliqua.
Siliqua, *Linn. Sp. Pl.* 1026.—Europ. austr. ; Orient.

CERATONYCHIA, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1215 = **Cometes**, Linn. (Amaranthac.).
Nidus, Edgew. l. c. = Com. surattensis.

CERATOPETALUM, Sm. Bot. N. Holl. 9. t. 3 (1793). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 651.
apetalum, *D. Don, in Edinb. N. Phil. Journ.* ix. (1830) 94.—Austral.
gummiferum, *Sm. Bot. N. Holl.* 9. t. 3.—Austral.
monopetalum, Caley, ex D. Don, in Edinb. N. Phil. Journ. ix. (1830) 94 = apetalum.
montanum, D. Don, l. c. = apetalum.
ovatum, Caley, ex D. Don, l. c. 95 = Schizomeria ovata.

CERATOPHORUS, Hassk. Retzia, i. 100 (1856) (= Keratophorus) = **Payena**, Linn. (Sapotac.).
Leerii, Hassk. l. c. 101 = P. Leerii.
longepetiolatus, Teijsm. & Binn. Cat. Hort. Bog. 161 = P. longepetiolata.
Wightii, Hassk. Retzia, i. 101 = P. lucida †
Wightii, Maing. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 547 = P. malaccensis.

CERATOPHORUS, Sond. in Linnaea, xxiii. (1850) 120 = **Gelonium**, Roxb. (Euphorbiac.).
africanus, Sond. l. c. 121 = G. africanum.

CERATOPHYLLUM, Linn. Syst. ed. I (1735). *CERATOPHYLLEAE*, Benth. & Hook. f. iii. 415.
DICHOTOPHYLLUM, Dill. ex Moench, Meth. 345 (1794).
apiculatum, Cham. in Linnaea, iv. (1829) 504 = demersum.
aquaticum, H. C. Wats. Topogr. Bot. i. 165 = demersum †
asperum, Lam. Fl. Fr. ii. 196 = demersum.
australe, *Griseb. in Goett. Abh.* xxiv. (1879) 14.—Reg. Argent.
chilense, *Leyb. in Anal. Univ. Chil.* (1859) 682.—Chili.
cornutum, Rich. in Ann. Mus. Par. xvi. (1810) 299.—demersum.
cristatum, *Gmill. & Perr. Fl. Senag. Tent.* 295.—Afr. trop.
demersum, Klein, ex Cham. in Linnaea, iv. (1829) 505 = tuberculatum.
demersum, *Linn. Sp. Pl.* 992.—Reg. temp. et trop.
demersum, Sieber, ex Cham. in Linnaea, iv. (1829) 504 = muricatum.
echinatum, *A. Gray, Man. Bot. U. St.* 401.—Am. bor.
gibbum, Laforet, ex Nym. Consp. 251 = demersum.
granulosum, Schur, Enum. Pl. Transs. 218 = submersum.
Hayuadicinum, *Borb. in Magy. Növényt. Lapok*, viii. (1884) 20.—Hungar.
indicum, Willd. ex Cham. in Linnaea, iv. (1829) 504 = tuberculatum.
inerme, Dum. Fl. Belg. 165 = submersum.
laxum, Lam. Fl. Fr. ii. 197 = submersum.
mississ, *Wall. Cat.* sub n. 7007.—Ind. or.
muricatum, *Cham. in Linnaea*, iv. (1829) 504.—As. et Afr. trop.
nudicum, Cham. l. c. 505 = submersum.
oxyacanthum, Cham. l. c. 504 = tuberculatum.
oxyacanthum, Schur, Enum. Pl. Transs. 218 = muricatum.
pentacanthum, *Hayuald, in Magy. Növényt. Lapok*, v. (1881) 109.—Hungar.
platyacanthum, Cham. in Linnaea, iv. (1829) 504 = muricatum.
polyacanthum, Schur, Enum. Pl. Transs. 218 = muricatum.
submersum, *Linn. Sp. Pl.* ed. II. 1409.—Europ.; As. trop.
triacanthum, Schur, Enum. Pl. Transs. 218 = muricatum.
tricorne, Dum. Fl. Belg. 165 = demersum.
tricuspidatum, Dum. l. c. = demersum.
tuberculatum, *Cham. in Linnaea*, iv. (1829) 504.—Ind. or.
unicorne, Dum. Fl. Belg. 165 = demersum.
verrucosum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 554 = submersum.
verruculosum, Rich. in Ann. Mus. Par. xvi. (1810) 299 = submersum.
verticillatum, Roxb Hort. Beng. 68; Fl. Ind. iii. 624 = demersum.
vulgare, Schleid. in Linnaea, xi. (1837) 540 = muricatum, tuberculatum, submersum.

CERATOPSIS, Lindl. Gen. et Sp. Orch. 388 (1840) = **Epipogum**, S. G. Gmel. (Orchid.).
roseo, Lindl. l. c. 384 = E. roseum.

CERATOPTYXIS, Hook. f. ic. Pl. t. 1125 (1873). *RUBIACEAE*, Benth. & Hook. f. iii. 105.
verbenacea, *Hook. f. l. c.*—Cuba.

CERATOSANTHES, Burm. ex Adans. Fam. ii. 139 (1763); Juss. Gen. 396 (1789). *CUCURBITACEAE*, Benth. & Hook. f. i. 833.

CERATOSANTHUS :—
corniculatus, Cogn. in DC. Monog. Phan. iii. 724 = tuberosus.
gracilis, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878); *Diagn. Cucurb.* fasc. ii. 88.—N. Granat.
Hilariana, *Cogn. l. c.* 81.—Bras.
latiloba, *Cogn. l. c.* 58.—Venezuela.
multiloba, *Cogn. l. c.* 29.—Bras.
tomentosa, *Cogn. l. c.*—Bras.
trifoliata, *Cogn. l. c.* 84.—Bras.
tuberosa, *F. F. Gmel. Syst.* 109.—Ind. occ.
Warmingii, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878); *Diagn. Cucurb.* fasc. ii. 80.—Bras.

CERATOSTANTHUS, Schur, in Verh. Sieb. Ver. Naturf. (1858) 46; et Enum. Pl. Transs. 80 (1866) = **Delphinium**, Tourn. (Ranunculac.).
Ajacis, Schur, Enum. Pl. Transs. 80 = D. Ajacis.
Consolida, Schur, in Verh. Sieb. Ver. Naturf. (1858) 46 = D. Consolida.
divaricatus, Schur, l. c. = D. divaricatum.
paniculatus, Schur, l. c. = D. paniculatum.

CERATOSCHOENUS, Nees, in Linnaea, ix. (1834) 296 = **Rynchospora**, Vahl (Cyperac.).
capitatus, Chapman, Fl. S. U. St. 599 = R. cyperoides.
corniculatus, Nees, in Linnaea, ix. (1834) 296 = R. laxa.
longivaginatus, Torr. in Ann. Lyc. N. York, iii. (1836) 269 = R. laxa.
macrostachyus, Tuckerm. in Am. Journ. Sc. Ser. II. vi. (1848) 382 = R. laxa.
thacrostachys, Torr. in Ann. Lyc. N. York, iii. (1836) 269 = R. macrostachya.

CERATOSEPALUM, Oerst. Rech. Fl. Am. Centr. 18. t. 17 (1863) = **Fassifova**, Linn.
micranthum, Oerst. l. c. t. 17 = P. ceratosepala.
parviflorum, Mast. in Mart. Fl. Bras. xiii. t. 555 = P. ceratosepala.

CERATOSPERMUM, Pers. Syn. ii. 551 (1807) = **Eurotia**, Adans. (Chenop.).
papposum, Pers. l. c. 552 = E. ceratoides.

CERATOSTACHYS, Blume, Bijdr. 644 (1825) = **Nyssa**, Linn. (Cornac.).
arborea, Blume, l. c. = N. sessiliflora.

CERATOSTEMA, Juss. Gen. Pl. 168 (1789). *VACCINIACEAE*, Benth. & Hook. f. ii. 570.
SIPHONOSTEMA, Klotzsch, in Linnaea, xxiv. (1851) 24.
SIPHONANDRA, Griseb. in Leohl. Berb. Am. Austr. 88 (1857).
angulatum, Griff. Notul. iv. 802 = Agapetes angulata.
biflorum, Poepp. & Endl. Nov. Gen. et Sp. i. 6. t. 10 = Eurygania biflora.
buxifolium, *Gardn. & Field, Sert. Pl.* t. 7.—Bras.
calycinum, *Benth. & Hook. f. Gen.* ii. 570.—Ecuador.
cordifolium, *Dun. in DC. Prod.* vii. 553.—Peruv.
costatum, *Benth. & Hook. f. Gen.* ii. 570.—Peruv.
ellipticum, *Benth. & Hook. f. l. c.*—Am. austr.
emarginatum, *Ruiz & Pav. Fl. Per. iv.* t. 384.—Peruv.
grandiflorum, *Ruiz & Pav. l. c.* 383.—Peruv.
hirsutum, *Ruiz & Pav. l. c.*—Peruv.
lanceolatum, *Benth. Pl. Hartw.* 142.—Peruv.
longiflorum, *Benth. & Hook. f. Gen.* ii. 570.—Ecuador.
longiflorum, Lindl. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 346 = grandiflorum.
loranthiflorum, *Benth. Pl. Hartw.* 142.—Peruv.
minutum, Griff. Notul. iv. 802 = Agapetes minuta.
myrtifolium, *Griseb. ex Benth. & Hook. f. Gen.* ii. 570, nomen.—Am. austr.
nanum, Griff. Notul. iv. 802 = Agapetes nana.
oblongifolium, *Dun. in DC. Prod.* vii. 553.—Peruv.
odoratissimum, Willd. ex Klotzsch, in Linnaea, xxiv. (1851) 21 = Cavendishia Quercena.
parviflorum, Benth. Fl. Hartw. 220 = Eurygania parviflora.
peruvianum, *Pers. Syn.* i. 480.—Peruv.
pubiflorum, *Wedd. Chlor. And.* ii. 181.—N. Granat.
rigidum, *Benth. Pl. Hartw.* 220.—N. Granat.
Sclaxa, *Benth. l. c.* 141.—Peruv.
speciosum, *Andrz. Illustr. Hortic.* xvii. (1870) t. 9.—Ecuador.

CERATOSTEMA :—
vaccinacrum, Roxb. Hort. Beng. 32; Fl. Ind. ii. 413 = Vaccinium serratum.
variegatum, D. Don, ex Steud. Nom. ed. II. i. 382 = Agapetes variegata.
variegatum, Griff. Notul. iv. 802 = Agapetes grandiflora.
variegatum, Roxb. Hort. Beng. 88; Fl. Ind. ii. 413 = Agapetes macrantha.

CERATOSTIGMA, Bunge, in Mém. Sav. Etr. Acad Pétersb. ii. (1835) 129 (Enum. Pl. Chin. Bor. 55) *PLUMBAGINEAE*, Benth. & Hook. f. ii. 628.
VALGRADIA, Hochst. in Flora, xxv. (1842) 289.
abyssinicum, *Achers. in Schweinf. Beitr. Fl. Aethiop.* 288.—Afr. trop.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 481.—Reg. Himal.
plumbaginoides, *Bunge, Enum. Pl. Chin. Bor.* 55.—China.

CERATOSTYLIS, Blume, Bijdr. 304. t. 56 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 517.
ACORIDIUM, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 181.
ancepa, *Blume, Bijdr.* 305.—Java.
braccata, *Reichb. in Bonplandia*, v. (1857) 58.—Java.
capitata, *Zoll. & Moritzi, in Nat. en Geneesk. Arch Ned. Indie*, i. (1844) 404.—Java.
Cepula, *Reichb. f. in Bonplandia*, v. (1857) 58.—Java.
Gigas, *Reichb. f. l. c.*—Java.
gracilis, *Blume, Bijdr.* 305.—Java.
graminea, *Blume, l. c.* 305.—Java.
latifolia, *Blume, l. c.*—Java.
retiquensis, *Reichb. f. in Bonplandia*, v. (1857) 58.—Ins. Philipp.
senilis, *Reichb. in Otia Bot. Hamb.* 54.—Ins. Philipp.
simplex, *Blume, Bijdr.* 305.—Java.
subulata, *Blume, l. c.* 306.—Java.
teres, *Reichb. f. in Bonplandia*, iii. (1854) 39.—Reg. Himal.

CERATOTHECA, Endl. in Linnaea, vii. (1832) 5. t. 1, 2. *PEDALINEAE*, Benth. & Hook. f. ii. 1059.
SPOELBERGA, Bernh. in Linnaea, xvi. (1842) 61.
Kraussiana, *Bernh. in Linnaea*, xvi. (1842) 62.—Afr. austr.
melanosperma, *Hochst. ex Bernh. l. c.*—Afr. trop.
sesamoides, *Endl. l. c.* vii. (1832) 5. t. 1, 2.—Afr. trop.
triloba, *E. Mey. ex Bernh. l. c.* xvi. (1842) 29.—Afr. austr.

CERATOZAMIA, Brongn. in Ann. Sc. Nat. Sér. III v. (1846) 7. t. 1, partim. *CYCADACEAE*, Benth. & Hook. f. iii. 446.
DIPHACOZAMIA, Lehm. in Lindl. Veg. Kingd. 295 (1847).
angustifolia, *Linden, in Illustr. Hortic.* xxviii. (1881) 93, nomen.—Mexic.
bolviana, *Brongn. in Ann. Sc. Nat.* Sér. III. v. (1846) 9.—Bolivia.
brevifrons, Miq. in Tijdschr. Wis- en Nat. Wetens. i. (1848) 41.—mexicana.
fusco-viridis, D. Moore, in Proc. Roy. Dubl. Soc. (1876) 118.—mexicana.
Ghiesbreghtii, Hort. ex Regel, in Act. Hort. Petrop. iv. (1876) 299.—Miquellana.
intermedia, Miq. in Tijdschr. Wis- en Nat. Wetens. i. (1848) 40.—mexicana.
Kersteniana, Hort. Kew. ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 192.—latifolia.
Kerstenianа, *Regel, in Bull. Soc. Nat. Mosc.* xxx. (1857) t. 187.—Mexic.
latifolia, *Hort. Belg. au Miq. in Tijdschr. Wis- en Nat. Wetens.* i. (1848) 206.—Mexic.
longifolia, Miq. l. c. 40 = mexicana.
mexicana, *Brongn. in Ann. Sc. Nat. Sér.* III. v. (1846) 8.—Mexic.
Miquelliana, *H. Wendl. Ind. Palm.* 69.—Mexic.
muricata, *Miq. in Linden, Illustr. Hortic.* xxviii. (1881) 93, nomen.—Mexic.
robusta, Miq. in Tijdschr. Wis- en Nat. Wetens. i. (1848) 42 = mexicana.

CERAUNIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (1827) (Quid ?).

CERBERA, Linn. Gen. ed. I. 63 (1737). *APOCYNACEAE*, Benth. & Hook. f. ii. 698.
CASCABELA, Rafin. Sylva Tellur. 162 (1838).
ELCANA, Blanco, Fl. Filip. ed. II. 584 (1845).
MANGHAS, Burm. Thes. Zeyl. 160 (1737).
NEIOSPERMA, Rafin. Sylva Tellur. 162 (1838).
ODOLLAM, Adans. Fam. ii. 171 (1763).
ODOLLAMIA, Rafin. Sylva Tellur. 162 (1838).
TANGHINIA, Thou. Gen. Nov. Madag. 10 (1806).
Abouai, Linn. Sp. Pl. 208 = Thevetia Ahouai.
alliodora, Willd. ex Roem. & Schult. Syst. iv. 798 = Thevetia cuneifolia.
batjanica, *Teijsm. & Binn. Cat. Hort. Bog.* 35*5*, nomen. —Malaya.
borbonica, Spreng. Syst. i. 642 = Ochrosia borbonica.
chinensis, Spreng. l. c. = Dissolaena verticillata.
cuneifolia, Jacq. ex Blume, Mus. Bot. Lugd. Bat. i. 158 = Pseudochrosia glomerata.
lactaria, Buch.-Ham. ex D. Dietr. Syn. Pl. i. 628 = C. Odollam.
laurifolia, Lodd. Bot. Cab. t. 989 = C. Odollam.
linearifolia, Stokes, Bot. Mat. Med. i. 490 = Thevetia nereifolia.
Linnaei, *Montr. in Mém. Acad. Lyon*, x. (1860) 233. —N. Caled.
maculata, Willd. Sp. Pl. i. 1223 = Ochrosia borbonica.
Manghas, Linn. Sp. Pl. 208 = C. Odollam, et Tabernaemontana dichotoma.
mauritiformis, Lam. Encyc. i. 62 = Bankeia mauritiiformis.
nereifolia, Zipp. in Bijdr. Nat. Wetensch. v. (1830) 178 = Podocarpus Rumphii.
nitida, H. B. & K. Nov. Gen. et Sp. iii. 225 = Thevetia nitida.
obovata, *Roem. & Schult. Syst.* iv. 798.—Hab. ?
obtusifolia, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 208.—N. Caled.
Odollam, *Gaertn. Fruct.* ii. 193.—Ind. or.; Malaya.
oppositifolia, *Lam. Encyc.* i. 62.—Ins. Moluce.
ovata, Cav. Ic. iii. 58. t. 270 = Thevetia ovata.
parviflora, Forst. f. Prod. 19 = Ochrosia elliptica.
parviflora, Hook. & Arn. Bot. Beech. Voy. 90 = Ochrosia sandwicensis.
parviflora, Wall. Cat. n. 1584 = Ochrosia borbonica.
parvoviana, Pers. Syn. i. 267 = Thevetia nereifolia.
platyspermos, Gaertn. Fruct. ii. 193. t. 134. f. 3 = Ochrosia borbonica.
quaternifolia, *Roxb. Hort. Beng.* 18, nomen.—Sumatra.
sinuosa, Blume, Bijdr. 1082 = oppositifolia.
Tanghin, *Hook. Bot. Mag.* t. 2966.—Madag.
Tanguin, Stend. Nom. ed. II. i. 633 = pruec.
Thevetia, Linn. Sp. Pl. 209 = Thevetia nereifolia.
thevetioides, H. B. & K. Nov. Gen. et Sp. iii. 223 = Thevetia Yecotli.
trifolia, Rudge, Pl. Gui. 31. t. 43 = Couma guianensis.
tubiflora, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 156 = Neuburgia tubiflora.
undulata, Andr. Bot. Rep. t. 130 = Ochrosia borbonica.
venenifera, Steud. Nom. ed. II. i. 633 = C. Tanghin.

CERBERA, Lour. Fl. Cochin. 136 (1790) = **Sonerila**, Linn. (Goodeniac.).
ovata, Sieber, ex DC. Prod. vii. 507 = S. Lobelia.
salutaris, Lour. Fl. Cochinch. 136 = S. Koenigii !

CERBERIOPSIS, Vieill. ex Pasch. & Sebert, Not. Bois Nouv. Calléd. 187 (1874). *APOCYNACEAE.*
candelabra, *Vieill. l. c.*—N. Caled.

CERCESTIS, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 414. *AROIDEAE*, Benth. & Hook. f. iii. 980.
Afzelii, *Schott, l. c.*—Afr. trop.

CERCIDIPHYLLUM, Sieb. & Zucc. in Abh. Math. Acad. Muench. iv. iii. (1846) 228. *MAGNOLIACEAE?*
japonicum, *Sieb. & Zucc. l. c.*—Japan.

CERCIDIUM, Tul. in Arch. Mus. Par. iv. (1844) 132. *LEGUMINOSAE*, Benth. & Hook. f. i. 576.
HOOYESIA, Buckl. in Proc. Acad. Sc. Philad. ' 1861 ' (1862) 458, partim.
RHETINOPHLORUM, Karst. Fl. Columb. ii. 25. t. 118 (1892).
andicolum, *Griseb. in Goett. Abh.* xxiv. (1879) 114.—Reg. Argent.
floridum, *Benth. ex A. Gray, Pl. Wright.* i. 58.—Mexico.
floridum, Torr. in Pacif. Rail. Rep. v. 360. t. 3 = Parkinsonia Torreyana.
spinosum, *Tul. in Arch. Mus. Par.* iv. (1844) 134.—Bras.
texanum, *A. Gray, Pl. Wright.* i. 58.—Mexic.

CERCIS, Linn. Sp. Pl. 374, ed. I (1753). *LEGUMINOSAE*, Benth. & Hook. f. i. 576.
SILIQUASTRUM, Tourn. ex Adans. Fam. ii. 317 (1763).
californica, Torr. ex Benth. Pl. Hartw. 861, nomen = occidentalis.
canadensis, *Linn. Sp. Pl.* 374.—Am. bor.
chinensis, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 95.—China.
floridia, Salisb. Prod. 827 = C. Siliquastrum.
Griffithii, *Boiss. Fl. Orient.* ii. 638.—Afghan.
japonica, Siebold, ex Planch. in Fl. des Serres, Ser. I. vii. (1852–53) 269 = chinensis.
occidentalis, *Torr. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 177.—Texas.
reniformis, Engelm. ex A. Gray, l. c. = occidentalis.
siliquastrum, Sp. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 193 = C. Siliquastrum.
siliquastrum, *Linn. Sp. Pl.* 374.—Europ. ; Oriens.

CERCOCARPUS, H. B. & K. Nov. Gen. et Sp. vi. 232. t. 559 (1823). *ROSACEAE*, Benth. & Hook. f. i. 618.
betuloides, Nutt. ex Hook. Ic. Pl. t. 322 = parvifolius.
betuloides, Nutt. in Torr. & Gray, Fl. N. Am. i. 427 = parvifolius.
brevifolius, *A. Gray, Pl. Wright.* ii. 54.—N. Mexic.
fothergilioides, *H. B. & K. Nov. Gen. et Sp.* vi. 233.—Mexic. ; Am. bor.
fothergilifolius, Torr. in Ann. Lyc. N. York, ii. (1828) 198 = parvifolius.
intricatus, *S. Wats. in Proc. Am. Acad.* x. (1875) 346. —Am. bor. occ.
ledifolius, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 427. —Am. bor. occ.
montanus, *Rafin. Atl. Journ.* 146.—Am. bor.
parvifolius, *Nutt. ex Hook. & Arn. Bot. Beech. Voy.* 337.—Calif.

CERCOCOMA, Wall. Cat. n. 1622 (1832) = Strophanthus, DC. (Apocynac.).
macrantha, Teijsm. & Binn. Cat. Hort. Bog. 126.—Hab. ?
singaporiana, Wall. Cat. n. 1622 = S. brevicaudatus.

CERCOCOMA, Miq. Fl. Ind. Bat. ii. 445 (1856) = Rhynchodia, Benth. (Apocynac.).
Wallichii, Miq. l. c. = R. Wallichii.

CERCODEA, Soland. ex Lam. Encyc. i. 682 (1783) = seq.

CERCODIA, Murr. in Comm. Gotting. iii. (1780) 3. t. 1 = **Haloragis**, Forst.
alternifolia, A. Cunn. in Ann. Nat. Hist. iii. (1839) 29 = H. alata.
erecta, Murr. in Comm. Gotting. iii. (1780) 3. t. 1 = H. alata.
tenuis, A. Cunn. in Ann. Nat. Hist. iii. (1839) 30 = H. tetragyna.
racemosa, DC. Prod. iii. 67 = H. racemosa.

CERCOPHORA, Miers, in Trans. Linn. Soc. xxx. (1874) 801. *MYRTACEAE.*
serrata, *Miers, l. c.* 302.—Mag. Amazon.

CERCOSTYLOS, Less. Syn. Comp. 229 (1832) = Gaillardia, Fouger. (Compos.).
brasiliensis, Less. l. c. 240 = G. megapotamica.
scabiosoides, Arn. ex DC. Prod. vii. 293 = G. megapotamica.
sessiliflora, Steud. Nom. ed. II. i. 633, sphalm. = pruec.

CERDANA, Ruiz & Pav. Prod. 37. t. 6 (1794) = Cordia Linn. (Boragin.).
alliodora, Ruiz & Pav. Fl. Per. ii. 47 = Cordia Cerdana.
cujabensis, Manso, ex DC. Prod. ix. 473 = Cordia cujabensis.

CERDIA, Mog. & Sesse, ex DC. Prod. iii. 377 (1828). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 153.
congestiflora, *Hemsl. Diagn. Pl. Nov.* 23.—Mexic.
glauca, *Hemsl. l. c.* 22.—Mexic.
purpurascens, *Moç. & Sesse, ex DC. Prod.* iii. 377.—Mexic.
virescens, *Moç. & Sesse, l. c.*—Mexic.

CEREFOLIUM, Riv. ex Rupp. Fl. jen. ed. Hall. 288 (1745) = **Anthriscus**, Hoffm. (Umbellif.).
nitidum, Celak. in Arch. Naturw. Landesd. Boehm. iii. (1875) 886 = A. nitida.
sativum, Bess. Prim. Fl. Galic. i. 219 = A. Cerefolium.
sylvestre, Bess. l. c. 218 = A. Cerefolium.
trichospermum, Bess. Enum. Pl. Volh. 44 = A. Cerefolium.

CEREOPSIS, Blanco, Fl. Filip. ed. II. Suppl. 591 (1845), sphalm. = **Coreopsis**, Linn. (Compos.).

CERESIA, Pers. Syn. i. 85 (1805) = **Paspalum**, Linn. (Gramin.).
aristata, Willd. ex Steud. Nom. ed. II. i. 333 = Panicum Thrasya.
elegans, Pers. Syn. i. 85 = Pasp. membranaceum.
fluitans, Ell. Sketch. i. 109 = Pasp. repens.
gracilis, Schlecht. in Linnaea, xxvi. (1853–55) 124 = Pasp. gracile.
membranacea, Beauv. Agrost. 9. t. 5. f. 4 = Pasp. membranaceum.
Wagnerziana, Schlecht. in Linnaea, xxvi. (1853–55) 183 = Pasp. Wagnerianum.

CERBUS, Mill. Gard. Dict. ed. VIII (1768) ; Haw. Syn. Pl. Succ. 176 (1812). *CACTEAE*, Benth. & Hook. f. i. 840.
CEPHALOCEREUS, Pfeiff. in Otto & Dietr. Allg. Gartenz. vi. (1838) 142.
CEPHALOPHORUS, Lem. Cact. Aliq. Nov. Desc. p. xii. (1838).
CLEISTOCACTUS, Lem. Illust. Hortic. viii. (1861) Misc. 85.
ECHINOCEREUS, Engelm. in Wisliz. Tour North Mexico, 91 (1848).
ECHINONYCTANTHUS, Lem. Cact. Gen. Nov. et Sp. 10 (1839).
ECHINOPSIS, Zucc. in Abh. Akad. Muench, ii. (1837) 676.
PILOCEREUS, Lem. Cact. Gen. Nov. et Sp. 6 (1839).
abnormis, Sweet, Hort. Brit. ed. I. 171 = peruvianus.
acifer, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 189.—Hab. ?
acinaciformis, Steud. Nom. ed. II. i. 338 = cinerascens.
acinaciformis, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 101 = cinerascens.
Achermanni, Hort. Berol. ex Pfeiff. l. c. 128 = Phyllocactus Achermanni.
acromelas, Hort. Berol. Index Cact. 1838, ex Pfeiff. Enum. Diagn. Cact. 84 = cremulatus.
aculeatus, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 488, nomen.—Hab. ?
acutangulus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 107.—Mexic.
adustus, *Engelm. in A. Gray, Pl. Fendl.* 50.—N. Mexic.
Aethiops, Haw. in Phil. Mag. (1830) 109 = caerulescens.
affinis, Hort. ex Pfeiff. Enum. Diagn. Cact. 90, 106 = variabilis, virens.
alacriportanus, *Hort. Monac. ex Pfeiff. Enum. Diagn. Cact.* 87.—Bras.
alatus, DC. Prod. iii. 470 = Rhipsalis pachyptera.
albisetosus, *Haw. Suppl. Pl. Succ.* 77.—Ins. S. Doming.
albispinus, *Salm-Dyck, Obs. Bot.* (1822) 5.—Am. austr.
ambiguus, *DC. Prod.* iii. 467.—Hab. !
amblygonus, *G. Don, in Sweet, Hort. Brit.* ed. III. 264, nomen.—Am. austr.
amoenus, *Hemsl. Biol. Centr. Am. Bot.* i. 540.—Mexic.

CEREUS:—
aniscanthus, *DC. in Mém. Mus. Par.* xvii. (1828) 116. —Mexic.
anisogonus, Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 52 = triangularis.
Antonii, Hort. Vindob. ex Pfeiff. Enum. Diagn. Cact. 114 = nycticalus.
aquiculensis, Hort. ex Pfeiff. l. c. 90 = geometrizans.
arequipensis, Meyen, Reise, ii. 41 = peruvianus.
armatus, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 81.—Ins. S. Thom.
Armbidae, *Steud. Nom.* ed. II. i. 388.—Bras.
articulatus, *Pfeiff. Enum. Diagn. Cact.* 103.—Chili.
assurgens, *Wright, ex Griseb. Cat. Pl. Cub.* 116.—Cuba.
atacamensis, *Phil. Fl. Atac.* 23.—Chili.
auratus, *Labour. in Rev Hortic. Sér.* IV. iv. (1855) 27. —Mexic.
aureus, *Salm-Dyck, ex DC. Prod.* iii. 465.—Hab.?
azureus, *Parm. ex Pfeiff. Enum. Diagn. Cact.* 86.—Bras.
bacciferus, Hemsl. Biol. Centr. Am. Bot. i. 548 = Rhipsalis Cassutha.
barbatus, Wendl. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 29 = floccosus.
Baumannii, *Lem. in Hortic. Unio.* (1844) 126, 315; *ex Lem. Jard. Fleur.* i. (1851) t. 46.—Peruv.?
baxaniensis, *Karw. ex Pfeiff. Enum. Diagn. Cact.* 109. —Mexic.
Beneckei, *Ehrenb. in Bot. Zeit.* ii. (1844) 835.—Mexic.
Berlandieri, *Engelm. Cact. Mex. Bound.* 38. t. 58.—Texas.
Bertini, *S. Cels, ex Hérincq, L'Horticult. Franç.* (1868) 251.—Chili.
biformis, Hort. ex Pfeiff. Enum. Diagn. Cact. 114 = radicans.
biformis, Lindl. Bot. Reg. (1843) Misc. 51 = Phyllocactus biformis.
bifrons, Haw. Suppl. Pl. Succ. 76 = coccineus.
Bigelovii, Engelm. in Pacif. Rail. Rep. iv. t. 4. f. 8 = mojavensis.
Blanckii, *Poselg. in Otto & Dietr. Allg. Gartens.* xxi. (1858) 184.—Mexic.
Boeckmanii, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 217.—Mexic.
bonariensis, *Sweet, Hort. Brit.* ed. III. 388.—Reg. Argent.
bonariensis, Hort. Berol. ex Foerst. Handb. Cact. ed. I. 388 = alacriportanus.
Bonplandii, *Parm. ex Pfeiff. Enum. Diagn. Cact.* 108. —Bras.
brachiatus, *Gal. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 195.—Mexic.
bradypus, Steud. Nom. ed. II. i. 388 = senilis.
Brandii, Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 49 = variabilis.
brevispinulus, Salm-Dyck, Hort. Dyck. 339 = nycticalus.
Bridgesii, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 208.—Bolivia.
caerulescens, *Salm-Dyck, Hort. Dyck.* 385.—Bras.
caesius, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 89.—Hab.?
caespitosus, *Engelm. & A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 247.—Texas.
californicus, Torr. & Gray, Fl. N. Am. i. 555 = Opuntia serpentina?
calilloche, Gal. ex Scheidw. in Bull. Acad. Brux. vi. 1. (1839) 88 = Echinocactus myriostigma.
calosceles, DC. in Mém. Mus. Par. xvii. (1828) 116 = peruvianus.
candelabrum, Hort. ex Pfeiff. Enum. Diagn. Cact. 109 = tenellus.
candelaris, Meyen, Reise, i. 447 = peruvianus.
candicans, *Gill. ex Salm-Dyck, Hort. Dyck.* 386.—Chili.
caripensis, *DC. Prod.* iii. 467.—Mexic.
candatus, Gill. ex Sweet, Hort. Brit. ed. III. 385 = Mammillaria caudata.
Cavendishii, *Monv. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 210.—Hab.?
chalibaeus, *Hort. Berol. ex Salm-Dyck, l. c.* 201.—Hab.?
chalybaeus, Otto, ex Walp. Rep. ii. 840 = polychaetus.
chiloensis, *DC. Prod.* iii. 465.—Ins. Chiloe.
chloranthus, *Engelm. Cact. Mex. Bound.* 29.—Texas.
CEREUS:—
chlorocarpus, *DC. Prod.* iii. 466.—Ecuador.
chrysomallus, *Hemsl. Biol. Centr. Am. Bot.* i. 541.—Mexic.
cinerascens, *DC. in Mém. Mus. Par.* xvii. (1828) 11. —Mexic.
cirrhiferus, *Labour. Cact.* 311.—Mexic.
clavarioides, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 178 = Opuntia clavarioides.
clavatus, *Otto & Dietr. Allg. Gartens.* vi. (1838) 28.—Venezuela.
claviformis, *Regel & Klein, Ind. Sem. Hort. Petrop.* (1860) 46.—Mexic.
coccineus, DC. Prod. iii. 469 = setaceus.
coccinnus, Engelm. in A. Gray, Pl. Fendl. 51 = phoeniceus.
coccineus, *Salm-Dyck, ex Pfeiff. & Otto, Cact.* i. t. 15.—Bras.
cognatus, Hort. ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
colicoënse, Booth, Cat. ex Foerst. Handb. Cact. ed. I. 386 = Echinocactus myriostigma.
colubrinus, *Otto, ex Foerst. i. c.* 409; *ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 50, 215.—Hab.?
Columna, Walp. Rep. v. 815 = sep.
Columna-Trajani, *Karw. ex Pfeiff. Enum. Diagn. Cact.* 76.—Mexic.
columnaris, *Lodd. ex Voigt, Hort. Suburb. Calc.* 61, nomen.—Hab.?
Colvillii x, *Sweet, Hort. Brit.* ed. II. 236, nomen, et p. 590.
cometes, *Scheidw. in Otto & Dietr. Allg. Gartens.* viii. (1840) 389.—Mexic.
compressus, *Mill. Gard. Dict.* ed. VIII. n. 10.—Mexic.
concolor, Bigel. in Whipple, Exped. Add. p. ii = caespitosus.
conformis, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 203.—Mexic.
conicus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 97.—Mexic.
conoideus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 36.—Am. bor. occ.
coquimbanus, Hort. ex DC. Prod. iii. 465 = chiloensis.
Coryne, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 206.—Hab.?
cosayrensis, *Tin. ex Todaro, in Ind. Sem. Hort. Panorm.* (1857) 39.—Hab.?
crenatus, Lindl. Bot. Reg. (1844) t. 31 = Phyllocactus crenatus.
crenoides, *Engelm. Cact. Mex. Bound.* 31. t. 42.—Mexic.
crenulatus, *Salm-Dyck, Obs. Bot.* (1822) 6.—Am. austr.
Crimsonii, Pritz. Ic. Ind. 246, sphalm. = Smithii.
crispatus, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 180 = Rhipsalis crispata.
cruciformis, *Steud. Nom.* ed. II. i. 388.—Bras.
cubensis, Zuccar. ex Pfeiff. Enum. Diagn. Cact. 94 = eriophorus.
cupulatus, Hort. ex Pfeiff. l. c. 97 = marginatus.
Curtisii, *Otto, in Otto & Dietr. Allg. Gartens.* i. (1833) 365.—N. Granat.
curvispinus, Pfeiff. Enum. Diagn. Cact. 89 = peruvianus.
cylindricus, Haw. Syn. Pl. Succ. 183 = Opuntia cylindrica.
dasyacanthus, *Engelm. in A. Gray, Pl. Fendl.* 50.—Texas.
decagonus, Hort. ex Pfeiff. Enum. Diagn. Cact. 85 = silvestrianus.
Decandollii, *Pfeiff. ex Foerst. Handb. Cact.* ed. I. 389. —Hab.?
decorus, *Lodd. ex Voigt, Hort. Suburb. Calc.* 62, nomen.—Hab.?
deficiens, *Otto & Dietr. Allg. Gartens.* vi. (1838) 28. —Venezuela.
deflexispinus, Monv. Cat. (1846), ex Labour. Cact. 321 = dasyacanthus.
denudatus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 78.—Bras.
Deppei, Hort. Berol. ex Pfeiff. l. c. 101 = cinerascens.
dichroacanthus, *Mart. ex Pfeiff. l. c.* 76.—Mexic.
divaricatus, *DC. Prod.* iii. 466.—Ins. S. Doming.
divergens, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 95.—Ins. S. Doming.
Donkelaarii, *Salm-Dyck, in Otto & Dietr. Allg. Gartens.* xiii. (1845) 258.—Bras.
dubius, *Engelm. Cact. Mex. Bound.* 36.—Texas.
CEREUS:—
Duledevanti, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 10. —Hab.?
Dumortieri, *Scheidw. in Bull. Acad. Brux.* vi. 1. (1839) 88.—Reg. Argent.
Dyckii, *Mart. ex Pfeiff. Enum. Diagn. Cact.* 87.—Mexic.
eburneus, *Salm-Dyck, Obs. Bot.* (1822) 6.—Ins. Curaçao; Chili.
Ehrenbergii, *Pfeiff. in Otto & Dietr. Allg. Gartens.* viii. (1840) 392.—Mexic.
elegans, Hort. ex Pfeiff. Enum. Diagn. Cact. 138 = Rhipsalis Mittleri.
elegans, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 10.—Peruv.
Emoryi, *Engelm. in Am. Journ. Sc.* Ser. II. xiv. (1852) 388.—Calif.
Engelmanni, *Parry, ex Engelm. l. c.*—Calif.
enneacanthus, *Engelm. in A. Gray, Pl. Fendl.* 50.—Texas.
erectus, *Karw. ex Pfeiff. Enum. Diagn. Cact.* 95.—Mexic.
ericomus, Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 47 = virens.
ericomus, *Reichb. f. in Terschech. Suppl. Cact. Vers.* 3.—Bras.
erinaceus, *Haw. ex Steud. Nom.* ed. II. i. 384.—Ind. occ.
erythrocephalus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 94.—Cuba.
euphorbioides, *Haw. Suppl. Pl. Succ.* 75.—Am. austr.
exsertus, Link, ex Pfeiff. Enum. Diagn. Cact. 99 = virens.
extensus, *Salm-Dyck, ex DC. Prod.* iii. 469.—Am. austr.
Eyriesii, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 72.—Reg. Argent.
farinosus, Haage, ex Salm-Dyck, in Otto & Dietr. Allg. Gartenz. xiii. (1845) 255 = Beneckei.
Fendleri, *Engelm. in A. Gray, Pl. Fendl.* 51.—N. Mexic.
Ferchecki, *Parm. ex Pfeiff. in Hort. Belg.* v. (1838) 66.—Reg. Argent.
fernambucensis, Lem. sphalm. vide pernambucensis.
ferox, *Haw. in Phil. Mag.* (1830) 109.—Bras.
fimbriatus, DC. Prod. iii. 464 = grandispinus.
fimbriatus, Hort. ex Pfeiff. Enum. Diagn. Cact. 95 = divaricatus.
flagelliformis, *Mill. Gard. Dict.* ed. VIII. n. 12.—Am. austr.
flagriformis, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 111. —Mexic.
flavescens, *Otto, ex Pfeiff. l. c.* 79.—Hab.?
flaviconus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 202. —Hab.?
flavispinus, *Haw. ex Steud. Nom.* ed. II. i. 384.—Am. austr.
flavispinus, *Salm-Dyck, Obs. Bot.* (1822) 5.—Am. calid.
floccosus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 81.—Ins. S. Thomas.
fluminensis, *Mig. in Bull. Sc. Phys. Nat. Néerl.* (1838) 48.—Bras.
Forbesii, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 393; *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 206.—Hab.?
formosus, *Hort. ex Foerst. Handb. Cact.* ed. I. 404.—Reg. Argent.
formosus, Monv. Cat. (1854) = Schraderi.
foveolatus, *Hadge, ex Pfeiff. Enum. Diagn. Cact.* 77. —Hab.?
fulgidus, *Hook. f. Bot. Mag.* t. 5856.—Hab.?
fulvibarbis, Otto & Dietr. Allg. Gartenz. vii. (1838) 28 = chiloensis.
fulvispinosus, Haw. Syn. Pl. Succ. 182 = Royeni.
fulvispinus, Salm-Dyck, Cact. Hort. Dyck. ed. II. 46, sphalm. = pmac.
Funkii x, *Hort. Monac. ex Foerst. Handb. Cact.* ed. I. 411.
Garambolli, Haage, ex Foerst. l. c. 438 = geometrizans.
geminiatus, *Reichb. in Terschech. Suppl. Cact. Vers.* 3.—Hab.?
gemmatus, Otto, ex Pfeiff. in Otto & Dietr. Allg. Gartenz. vii. (1838) 214 = turbinatus.
gemmatus, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 96. —Mexic.
geometrizans, *Mart. ex Pfeiff. l. c.* 90.—Mexic.

CEREUS:—
gibbosus, *Sweet, Hort. Brit.* ed. I. 171.—Jamaica.
gigantens, *Engelm. in Emory, Notes Mil. Reconnois.* 158.—Texas.
gilvus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 355.—Hab.?
Gladiator, Otto & Dietr. i. c. vi. (1836) 84 = geometrizans.
gladsatus, Lem. Cact. Alig. Nov. Desc. 28 = candicans.
Gladiger, Cels, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 47 = resupinatus.
glaucescens, *Tweedie, ex Sweet, Hort. Brit.* ed. III. 284, nomen.—Reg. Argent.
glaucus, Hort. ex Pfeiff. Enum. Diagn. Cact. 89 = caesius, variabilis.
glaucus, *Salm-Dyck, Hort. Dyck.* 835.—Bras.
gloriosus, Hort. ex Pfeiff. Enum. Diagn. Cact. 80 = Royeni.
gonacanthus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 33. t. 5. f. 2. 3.—N. Mexic.
gracilis, *How. in Phil. Mag.* (1827) 125.—Hab.?
gracilis, *Mill. Gard. Dict.* ed. VIII. n. 8.—Ind. occ.
gracilis, Salm-Dyck, ex DC. Prod. iii. 468 = humilis.
grandiflorus, *Mill. Gard. Dict.* ed. VIII. n. 11.—Ind. occ.
grandis, *Haw. Suppl. Pl. Succ.* 76.—Bras.
grandispinus, *Haw. in Phil. Mag.* (1830) 109.—Ind. occ.
Greggii, *Engelm. in Wisliz. Rep. N. Mexico*, 102 — Texas.
grisens, Haw. Syn. Pl. Succ. 182 = eburneus.
Haageanus, *Salm-Dyck, ex Foerst. Handb. Cact.* ed. I. 881, nomen.—Hab.?
hamatus, [Scheidw.] Hort. Belg. ex Pfeiff. in Otto & Dietr. Allg. Gartenz. v. (1837) 871 = rostratus.
Hansii x, *Baumann, in Rev. Hortic.* Sér. II. i. (1842) 267.
Haworthii, *DC. Prod.* iii. 465.—Ind. occ.
heptagonus, *Mill. Gard. Dict.* ed. VIII. n. 6; *Haw. Syn. Pl. Succ.* 178.—Am. austr.
hermentianus, *Morev. ex Lem. in Illustr. Hortic.* vi. (1859) *Misc.* 90.—Hab.?
heteracanthus, *Tweedie, ex Sweet, Hort. Brit.* ed. III. 284.—Reg. Argent.
hexaedrus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 34. t. 5. f. 1.—N. Mexic.
hexagonus, *Mill. Gard. Dict.* ed. VIII. n. 1; *Haw. Syn. Pl. Succ.* 179.—Am. austr.
hexangularis, Hort, ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
Histrix, Salm-Dyck, ex DC. Prod. iii. 464.—Cf. Hystrix.
Hoffmannseggii, Hort. ex Foerst. Handb. Cact. ed I. 292 = Echinocactus exsculptus.
Hookeri, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 125.—Bras.
horizontalis, *Gill. ex Sweet, Hort. Brit.* ed. III. 285.—Chili.
horribarbis, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 205.—Hab.?
horridus, *Otto, ex Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 870.—Venezuela.
Humboldtii, *DC. Prod.* iii. 467.—Ecuador.
humilis, *DC. l. c.* 468.—Am. austr.
hybridus, Otto, in Otto & Dietr. Allg. Gartenz. i. (1888) 72 = speciosissimus.
hypogaeus, *Weber, ex Regel, Gartenfl.* (1882) 165. t. 1088.—Hab.?
Hystrix, *Sweet, Hort. Brit.* ed. I. 171.—Ind. occ.
icosigonus, *DC. Prod.* iii. 467.—Ecuador.
imbricatus, Haw. Rev. Pl. Succ. 70 = Opuntia imbricata.
imbricatus, Hort. ex Pfeiff. Enum. Diagn. Cact. 98 = subrepandus.
incrustans, Stend. Nom. ed. II. i. 324 = seq.
incrustatus, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 97 = marginatus.
incurvispinus, Hort. Darnst. ex Otto & Dietr. Allg. Gartenz. iii. (1888) 244 = leucanthus.
Inermis, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 116.—Venezuela.
inermis, Scheidw. in Bull. Acad. Brux. vi. (1839) 88 = Echinocactus myrtostigma.
insularis, *Nemol. Bot. Voy. Challenger*, i. 11. 16.—Moss 3. Mich.

CEREUS:—
intricatus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 194.—Hab.?
inversus, *Hort. Berol. ex Salm-Dyck, l. c.* 219, nomen. —Hab.?
irradians, *Lem. Illust. Hortic.* xi. (1864) *Misc.* 74.—Cuba.
Jamacaru, Hort. Vindob. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 81 = variabilis.
Jamacaru, *DC. Prod.* iii. 467.—Bras.
janthothele, Monv. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 41 = Rhipsalis cereiformis.
jasminosus, Hort. Darmst. ex Pfeiff. Enum. Diagn. Cact. 72 = turbinatus.
Jenkinsonn x, Sweet, Hort. Brit. ed. II. 237, nomen.
jubatus, Salm-Dyck, ex Foerst. Handb. Cact. ed I. 357 = cometes.
Kagenrchii, Gmel. ex Pfeiff. Enum. Diagn. Cact. 77 = multangularis.
Karstenii, Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 207 = lepidotus.
Karstenii, *Salm-Dyck, Cact. Hort. Dyck.* ed. II 218.—N. Granat.
Knightii, Parment. ex Pfeiff. Enum. Diagn. Cact. 139 = Rhipsalis Knightii.
Kunthianus, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 217.—Hab.?
laetevirens, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 99 = lividus.
laetevirens, Salm-Dyck, Hort. Dyck. 836 = variabilis.
laetus, *DC. Prod.* iii. 466.—Ecuador.
laetus, Salm-Dyck, Hort. Dyck. 836 = formosus.
laevigatus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 204.—Hab.?
lagenaeformis, *C. F. Foerst. in Hamb. Gartens.* xvii. (1851) 164.—Peruv.
lamprochlorus, *Lem. Cact. Alig. Nov. Desc.* 30.—Hab.?
lanatus, *DC. Prod.* iii. 464.—Ecuador.
Lanceanus, *Hort. ex G. Don, in Sweet, Hort. Brit.* ed. III. 285, nomen.—Hab.?
Landbecki, *Phil. ex Regel, Gartenfl.* (1875) 162. t. 893.—Reg. Argent.
lanuginosus, Haw. Syn. Pl. Succ. 182 = repandus.
laniginosus, *Mill. Gard. Dict.* ed. VIII. n. 3.—Am. trop.
laterharbatus, Hort. ex Lem. in Rev. Hortic. (1862) 427 — C. Columna-Trajani.
latifrons, Zucc. in Denkschr. Akad. Muench. ii. (1837) 785 = Phyllocactus latifrons.
Lecchi, *Pfeiff. Enum. Diagn. Cact.* 78.—Hab.?
Leeanus, *Hook. Bot. Mag.* t. 4417.—Mexic.
Lehmanni, Hort. ex Foerst. Handb. Cact. ed. I. 245 = Mammillaria Lehmanni.
leiocarpus, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 276.—Porto Rico.
Lemaireii, *Hook. Bot. Mag.* t. 4814.—Hab.?
lepidotus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 207.—Venezuela.
leptacanthus, DC. ex Pfeiff. Enum. Diagn. Cact. 101 = pentalophus.
leptophis, *DC. in Mém. Mus. Par.* xvii. (1828) 117.—Mexic.
leucanthus, *Pfeiff. Enum. Diagn. Cact.* 71.—Chili.
limensis, Lodd. ex Foerst. Handb. Cact. ed. I. 375 = flavescens.
limensis, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 353.—Bolivia.
Linkii, Pfeiff. Enum. Diagn. Cact. 48 = Echinocactus Linkii.
Linneei, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 165.—Mexic.
lividus, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* (1885) 880.—Am. austr.
longifolius, *Karw. ex Sweet, Hort. Brit.* ed. III. 286, nomen.—Mexic.
longipetalus, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 438, nomen.—Hab.?
longiscus, *Engelm. Cact. Mex. Bound.* 32. t. 45.—Texas.
longispinus, *Salm-Dyck, ex Otto & Dietr. Allg. Gartenz.* xiii. (1846) 854.—Bolivia.
lumbricoides, *Lem. Cact. Gen. Nov. et Sp.* 60.—Bras.
lutescens, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 84.—Hab.?
MacDonaldiae, *Hook. Bot. Mag.* t. 4707.—Honduras.

CEREUS:—
macrogonus, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 208.—Hab.?
Maelenii, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 879.—Mexic.
magnus, *Haw. in Phil. Mag.* (1830) 109.—Ins. S. Doming.
Mallisoni, Hort. ex Pfeiff. Enum. Diagn. Cact. 111 = Smithii.
marginatus, *DC. in Mém. Mus. Par.* xvii. (1828) 116. —Mexic.
marginatus, Salm-Dyck, Hort. Dyck. 840 = Hookeri
Mariculi, Hort. ex Pfeiff. Enum. Diagn. Cact. 115 = humilis.
Martianus, *Zucc. ex Pfeiff. l. c.* 110.—Mexic.
Mendory, Hort. ex Pfeiff. l. c. 85 = caerulescens.
mecnacanthus, Link & Otto, ex Steud. Nom. ed. II. i. 334 = Echinocactus mecnacanthus.
mexicanus, *Lem. ex Foerst. Handb. Cact.* ed. I. 430.—Mexic.
micracanthus, *DC. in Mém. Mus. Par.* xvii. (1828) 115.—Mexic.
militaris, *Audot, in Rev. Hortic.* Sér. II. iv. (1845–46) 307.—Mexic.
Mirbelii, Hort. Belg. ex Pfeiff. Enum. Diagn. Cact. 97 = marginatus.
mokoyensis, S. Wats. Bibliog. Ind. Fl. Am. Bor. 398 = seq.
mojavensis, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 33.—Am. bor. occ.
mollis, Hort. ex Pfeiff. Enum. Diagn. Cact. 83 = strictus.
moniliformis, *DC. in Mém. Mus. Par.* xvii. (1828) 60. —Ins S. Doming.
monoclonus, DC. Prod. iii. 464 = peruvianus.
monstrosus, Steud. Nom. ed. II. i. 334 = peruvianus.
montevidensis, Hort. ex Pfeiff. Enum. Diagn. Cact. 65 = Echinocactus exsculptus.
Montezumae, Hort. ex Pfeiff. l. c. 91 = candicans.
Moritzianus, *Otto, ex Pfeiff. l. c.* 84.—Bras.
multangularis, *How. Suppl. Pl. Succ.* 75.—Am. austr.
multiplex, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 70.—Bras.
Mysorus, Salm-Dyck, in DC. Prod. iii. 469 = Rhipsalis Mysorus.
myrsocaulon, Mart. ex Pfeiff. Enum. Diagn. Cact. 115 = humilis.
myriophyllus, Gill. ex Otto, in Otto & Dietr. Allg. Gartenz. i. (1888) 365 = strigosus.
nanus, *DC. Prod.* iii. 470.—Ecuador.
Napoleonis, *R. Grah. in Bot. Mag.* t. 3458.—Ind. occ.
niger, *Salm-Dyck, Obs. Bot.* (1822) 4.—Am. austr.
nigricans, Hort. ex Pfeiff. Enum. Diagn. Cact. 83 = strictus.
nigricans, *Lem. Cact. Gen. Nov. et Sp.* 57.—Hab.?
nigrispilis, *Phil. Fl. Atac.* 23.—Chili.
nitens, *Salm-Dyck, ex Otto & Dietr. Allg. Gartenz.* xiii. (1845) 854.—Bolivia.
nitidus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 211.—Venezuela.
nobilis, Haw. Syn. Pl. Succ. 179 = Haworthii.
nycticalus, *Link, in Verh. Preuss. Ver. Gartenb.* x. (1834) 373. t. 4.—Mexic.
nycticaulon, Sweet, Hort. Brit. ed. III. 285, sphalm. = *myriocaulon*.
obtusus, Haw. Rev. Pl. Succ. 70 = variabilis.
obtusus, Hort. ex Pfeiff. Enum. Diagn. Cact. 114 = nycticalus.
Ocamponis, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 220. —Mexic.; N. Granat.
ochracanthus, Hort. ex Pfeiff. Enum. Diagn. Cact. 78 = multangularis.
octogonus, Hort. ex Otto, in Otto & Dietr. Allg. Gartenz. i. (1888) 365 = allotropus, Curtisii.
Oliveirii, *Salm-Dyck, Hort. Dyck.* 835.—Bras.
olivaceus, *Lem. in Rev. Hortic.* Sér. IV. viii. (1859) 643.—Ind. occ.
Ottonis, Pfeiff. Enum. Diagn. Cact. 47, sphalm. = Echinocactus Ottonis.
ovatus, *G. Don, in Loud. Hort. Brit.* 195; *Pfeiff. Enum. Diagn. Cact.* 102.—Chili.
oxygonus, *Otto, in Otto & Dietr. Allg. Gartenz.* i. (1888) 365.—Bras.
oxygonus, Salm-Dyck, ex Foerst. Handb. Cact. ed. I. 393 = euphorbioides.
oxypetalus, DC. Prod. iii. 470 = Phyllocactus latifrons.

CEREUS :—
oxypetalus, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 97, nomen = Phyllocactus marginatus.
paniculatus, *DC. Prod.* iii. 466.—Ins. S. Doming.
pauciareatus, Cels, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 44 = pycnacanthus.
paradoxus, Hort. ex Steud. Nom. ed. II. i. 335 = Opuntia brasiliensis.
parasiticus, Haw. ex Steud. l. c. = Rhipsalis fasciculata.
parvistus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 79.—Bras.
pasciapinus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 34, 36; *et in Proc. Am. Acad.* iii. (1857) 285.—Texas.
Paxtonianus, *Monv. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 211.—Hab ?
pectinatus, Engelm. in A. Gray, Pl. Fendl. 50 = caespitosus.
pectiniferus, Labour. Monog. Cact. 320 = caespitosus.
pellucidus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 108.—Cuba.
pentagonus, *Haw. Syn. Pl. Succ.* 180.—Am. austr.
pentagonus, Hort. ex Foerst. Handb. Cact. ed. I. 389 = peruvianus.
pentalophus, *DC. in Mém. Mus. Par.* xvii. (1828) 117. —Mexic.
pentapterus, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed II. 220.—N. Granat.
Pepinianus, *Lem. ex Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 354.—Bolivia.
pernambucensis, Lem. Nov. Cact. Gen. et Sp. 58 (err. typ. *fernambucensis*) = variabilis.
Perrottetii, Hort ex Pfeiff. Enum. Diagn. Cact. 98 = lividus.
peruvianus, Hort. ex Foerst. Handb. Cact. ed. I. 392 = eburneus.
peruvianus, *Mill. Gard. Dict.* ed. VIII. n. 4; *Haw. Syn. Pl. Succ.* 179.—Am. austr.
Pfeiffleri. *Parm. ex Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 270.—Reg. Argent.
Phillippi, *Regel & Schmidt, Gartenfl.* (1882) 98.—Chili.
phoenicens, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 34. t. 4.—Am. bor. occ.
phyllanthoides, DC. Prod. iii. 469 = Phyllocactus phyllanthoides.
Phillyanthus, DC. l. c. = Phyllocactus Phyllanthus.
Pitajaya, DC. l. c. 466 = variabilis.
platycarpus, Zucc. ex Foerst. Handb. Cact. ed. I. 452 = Rhipsalis platycarpa.
platygonus, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 199.—Hab. ?
polyacanthus, *Engelm. in A. Gray, Pl. Fendl.* 51.—Texas.
polyacheatus, *Reichb. in Terach. Suppl. Cact. Vers.* 4. —Hab. ?
polygonatus, Hort. ex Pfeiff. Enum. Diagn. Cact. 91 = eburneus.
polygonus, *DC. Prod.* iii. 466.—Ins. S. Doming.
polylophus, *DC. in Mém. Mus. Par.* xvii. (1828) 115. —Mexic.
polymorphus, *G. Don, in Loud. Hort. Brit.* 195.—Chili.
polymorphus, Hort. Monac. ex Foerst. Handb. Cact. ed. I. 472 = articularius.
polyptychus, *Lem. Cact. Gen. Nov. et Sp.* 56.—Hab. ?
Pottsii, Salm-Dyck, Cact. Hort. Dyck. ed. II. 208 = Greggii.
Princeps, *Hort. Wuerteb. ex Pfeiff. Enum. Diagn. Cact.* 108.—Am. bor. occ.
Pringlei, *S. Wats. in Proc. Am. Acad.* xx. (1885) 368. —Am. bor. occ.
prismaticus, Haw. ex Steud. Nom. ed. II. i. 335 = pentagonus.
prismaticus, *Salm-Dyck, in DC. Prod.* iii. 469.—Am. austr.
prismatiformis, Hort. ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
procumbens, *Engelm. in A. Gray, Pl. Fendl.* 50.—Mexic.
procumbens, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 208.—Texas.
propinquus, Salm-Dyck, ex Otto, in Otto & Dietr. Allg. Gartenz. i. (1833) 366 = pentalophus.
protoformis, *Steud. Nom.* ed. II. i. 335.—Hab. ?
pruinosus, Salm-Dyck, Cact. Hort. Dyck. ed. II. 47 = Echinocactus prulnosus.

CEREUS :—
pterandrus, Hemsl. Biol. Centr. Am. Bot. i. 544, sphalm. = seq.
pteranthus, Link, ex Dietr. in Otto & Dietr. Allg. Gartenz. ii. (1834) 309 = nycticalus.
pterocaulis, Hort. ex Pfeiff. Enum. Diagn. Cact. 140 = Rhipsalis paradoxa.
pterogonus, *Lem. Cact. Gen. Nov. et Sp.* 59.—N. Granat.
pugioniferus, Lem. Cact. Aliq. Nov. Desc. 30 = geometrizans, Gladiator.
pulchellus, *Pfeiff. Enum. Diagn. Cact.* 74.—Mexic.
pycnacanthus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 355.—Bolivia.
quadrangularis, Haw. Syn. Pl. Succ. 181 = carpenais.
quadrangularis, Hort. ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
quadrangulipunus, *Lem. ex Ehrenb. in Linnaea*, xix. (1847) 363.—Mexic.
quadricostatus, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 376.—Porto Rico.
Quintero, Hort. Gottlng. ex Pfeiff. Enum. Diagn. Cact. 86 = chilensis.
Quisco, *C. Gay, Fl. Chil.* iii. 19.—Chili.
radicans, *DC. Prod.* iii. 468.—Am. austr.
ramosus, Karw. in Pfeiff. Enum. Diagn. Cact. 108 = baxaniensis.
ramulosus, Salm-Dyck, Hort. Dyck. 840 = Rhipsalis ramulosa.
recurvus, Link & Otto, ex Steud. Nom. ed. II. i. 335 = Echinocactus recurvus.
reductus, DC. Prod. iii. 463 = Echinocactus gibbosus.
reflexus, Hort. ex Steud. Nom. ed. II. i. 335 = virens.
regalis, *Haw. ex Sprengel. Syst.* ii. 496.—Am. austr.
Reichenbachianus, Labour. Monog. Cact. 318 = caespitosus.
repandus, Haw. Syn. Pl. Succ. 183 = gracilis.
repandus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Ind. occ.
reptans, Haw. ex Steud. Nom. ed. II. i. 335 = pentagonus.
reptans, Salm-Dyck, ex DC. Prod. iii. 468 = radicans.
resupinatus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 10.—Hab. ?
retrofractus, Hort. ex Pfeiff. in Otto & Dietr. l. c. iii. (1835) 380 = virens.
rhombeus, Salm-Dyck, Hort. Dyck. 841 = Rhipsalis rhombea.
rigidispinus, *Lem. Hort. Unie.* (1840) 223. t. 1.—Mexic.
rigidispinus, *Muehlenp. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 13.—Mexic.
rigidus, Lem. ex Salm-Dyck, Cat./Hort. Dyck. ed. I. 84; ed. II. 52 = hamilis.
Roemeri, *Engelm. in A. Gray, Pl. Fendl.* 50.—Texas.
Roemeri, *Muehlenp. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 19.—Hab. ?
Roetteri, *Engelm. Cact. Mex. Bound.* 33. t. 41. f. 8–5. —Am. bor. occ.
roridus, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 54. = Blossecatus prulnosus.
rosaceus, Hort. ex Pfeiff. l. c. 114 = nycticalus.
rostratus, *Lem. Cact. Aliq. Nov. Desc.* 29.—Ins. Antigua.
Royeni, Haw. Syn. Pl. Succ. 182 = lanuginosus.
Royeni, Hook. Bot. Mag. t. 3125 = Curtisi.
Royeni, *Mill. Gard. Dict.* ed. VIII. n. 7.—Am. austr.
rupicolus, *Engelm. in A. Gray, Pl. Fendl.* 50.—N. Mexic.
Russellianus, *Forb. Hort. Tour Gerw.* 154, nomen.—Guiana.
Russellianus, Gardn. ex Hook. Bot. Mag. t. 3717 = Ephiphyllum Russellianum.
Russellianus, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 201.—Laguayra.
Salm-Dyckianus, *Hensel. Biol. Centr. Am. Bot.* i. 545. —Mexic.
Salm-Dyckianus, Hort. Vindob. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 81; ed. II. 49 = variabilis.
scandens, *Salm-Dyck, l. c.* ed. II. 219.—Guiana.
Schaerli, *Salm-Dyck, l. c.* 180.—Mexic.
Schelhasii, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iii. (1835) 214.—Hab. ?
Schombergkii, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 219.—Guiana ?

CEREUS :—
Schottii, *Engelm. in Proc. Am. Acad.* iii. (1857) 286 —Am. bor. occ.
Schrankii, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 122.—Mexic.
Scopa, Salm-Dyck, ex DC. Prod. iii. 464 = Echinocactus Scopa.
Siedelii, *Lehm. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 200.—Hab. ?
senilis, *Salm-Dyck, in DC. Prod.* iii. 464.—Am. austr.
septum, *DC. l. c.* 467.—Ecuador.
sericeus, Hort. ex Pfeiff. Enum. Diagn. Cact. 173 = Opuntia clavarioides.
serpens, *DC. Prod.* iii. 470.—Ecuador.
serpentinus, *DC. l. c.* 467.—Mexic.
serrulifolrus, *Haw. in Phil. Mag.* (1830) 109.—Ind. occ.
setaceus, *Salm-Dyck, in DC. Prod.* iii. 469.—Bras.
setiger, *Haw. in Phil. Mag.* (1830) 109.—Bras.
similis, *Meinsh. in C. Koch & First. Wochenschr.* i. (1856) 39.—Mexic.
Smithianus ×, Sweet, Hort. Brit. ed. II. 237, nomen = seq. ?
Smithii ×, *Hort. Angl. ex Pfeiff. Enum. Diagn. Cact.* 111.
Spachianus, *Lem. Hort. Unie.* i. 225.—Mexic.
speciosissimus, *DC. Prod.* iii. 468.—Mexic.
spectatus, Sweet, Hort. Brit. ed. I. 172 = variabilis ?
spinibarbis, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 86.—Chili.
spinosissimus, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 165.—Peruv.
spinulosus, *DC. in Mém. Mus. Par.* xvii. (1828) 117. —Mexic.
splendens, *Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 214.—Hab. ?
splendidus, *Past. ex Steud. Nom.* ed. II. i. 336 (= hplphyll sp. ?).—Mexic.
squamulosus, Salm-Dyck, in DC. Prod. iii. 469 = Rhipsalis Mittleri.
stellatus, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 358.—Mexic.
stramineus, *Engelm. Cact. Mex. Bound.* 35. t. 46.—Texas.
striatus, *DC. Prod.* iii. 465.—Am. austr.
strigosus, *Salm-Dyck, Hort. Dyck.* 384.—Chili.
subinermis, *Hemsl. Biol. Centr. Am. Bot.* i. 546.—Mexic.
sublanatus, *Salm-Dyck, Hort. Dyck.* 837.—Hab. ?
subrepandus, *Haw. Suppl. Pl. Succ.* 78.—Ind. occ.
subrepandus, Hort. ex Pfeiff. Enum. Diagn. Cact. 86, 94 = chilensis, eriophorus.
subsquamatus, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 380 = extensus.
subterranous, Hort. ex Foerst. Handb. Cact. ed. I. 409 = colubrinus.
subuliferus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 354.—Mexic.
superbus, *Ehrenb. in Bot. Zeit.* iv. (1846) 394.—Mexic.
Swartzii, *Griseb. Fl. Brit. W. Ind.* 801.—Ind. occ.
syringacanthus, *Pfeiff. Enum. Diagn. Cact.* 108.—Chili.
tenellus, *Salm-Dyck, ex Pfeiff. l. c.* 109.—Bras.
tenuis, DC. Prod. iii. 469 = Rhipsalis Myosurus.
tenuis, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 407.—Am. austr.
tenuispinus, Haw. in Phil. Mag. (1827) 125 = Rhipsalis Myosurus.
tenuissimus, *G. Don, in Sweet, Hort. Brit.* ed. III. 265, nomen.—Hab. ?
tephracanthus, *Link & Otto, ex Steud. Nom.* ed. II. i. 336.—Am. austr.
Terschreckii, *Parm. ex Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 270.—Reg. Argent.
tetracanthus, *Lahoer, in Rev. Hortic. Sér.* IV. iv. (1855) 25.—Bolivia.
tetragonus, *Mill. Gard. Dict.* ed. VIII. n. 2; *Haw. Syn. Pl. Succ.* 180.—Am. austr.
thalassinus, *Otto & Dietr. Allg. Gartenz.* vi. (1838) 34. —Venezuela.
Thouinii, *Engelm. in Am. Journ. Sc. Ser.* II. xvii. (1854) 234.—N. Mexic.
Ulophorus, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 380 = virens.
Thuri, *Fodera, Ind. Sem. Hort. Panorm.* (1867) 69. —*Cf. in Linnaea*, xxix. (1857–58) 703.—Bras. ?

CEREUS:—
Tonellanus, *Lem. Illustr. Hortic.* ii. (1855) *Misc.* 68.—Mexic.
tortuosus, *Forbes, ex Otto & Dietr. Allg. Gartens.* vi. (1838) 95.—Reg. Argent.
triangularis, Haw. Syo. Fl. Succ. 180 = compressus.
triangularis, *Mill. Gard. Dict. ed.* VIII. n. 9.—Hab. †
trichacanthus, Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 46 = lutescens.
triglochidiatus, *Engelm. in A. Gray, Pl. Fendl.* 51.—N. Mexic.
trigonus, *Haw. Syn. Pl. Succ.* 181.—Ind. occ.
trigonus, Hort. ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
trinitatensis, *Jan & Herm. in Rev. Hortic.* Sér. IV. viii. (1859) 642.—Ind. occ.
tripteris, *Salm-Dyck, in DC. Prod.* iii. 468.—Am. austr.
triqueter, *Haw. Syn. Pl. Succ.* 181.—Am. austr.
truncatus, Sweet, Hort. Brit. ed. I. 172 = Epiphyllum Altensteinii.
tuberosus, *Pfeiff. Enum. Diagn. Cact.* 102.—Mexic.
tuberosus, *Poselg. in Otto & Dietr. Allg. Gartens.* xxi. (1853) 186.—Texas.
tubiflorus, Endl. in Harting. Pazad. Vindob. t. 71 (= *Echinopsis Zuccarini*).—Hab.?
tubiflorus, Pfeiff. Enum. Diagn. Cact. 71 = Echinocactus tubiflorus.
tunicatus, Pfeiff. l. c. 178 in syn. = Opuntia tunicata.
turbinatus, *Pfeiff., in Otto & Dietr. Allg. Gartens.* iii. (1835) 314.—Hab.?
Tweediei, *Hook. Bot. Mag.* t. 4498.—Reg. Argent.
undatus, Haw. in Phil. Mag. (1830) 110 = Napoleonis.
undatus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 94.—Hab.?
undiflorus, *Wright, in Sauv. Fl. Cub.* 59.—Cuba.
undulatus, Hort. Berol. ex D. Dietr. Syn. Pl. iii. 104 = undatus.
undulatus, Hort. Dresd. ex Pfeiff. Enum. Diagn. Cact. 107 = scutangulus.
undulosus, DC. Prod. iii. 467 = variabilis.
urescanthus, *C. F. Foerst. in Hamb. Gartens.* xvii. (1861) 166.—Peruv.
validus, *Haw. in Phil. Mag.* x. (1831) 414.—Am. austr.
variabilis, Engelm. Cact. Mex. Bound. 40 = Princeps.
variabilis, *Pfeiff. Enum. Diagn. Cact.* 108.—Am. austr.
Vaseyi, Young, Fl. Texas, 376 = Princeps.
violaceus, *Lem. Cact. Gen. Nov. et Sp.* 87.—Hab.?
virens, *DC. in Mem. Mus. Par.* xvii. (1828) 116.—Mexico.
viridiflorus, *Engelm. in A. Gray, Pl. Fendl.* 50.—Texas.
xanthochaetus, *Reichb. in Tersch. Suppl. Cact. Vers* 4.—Hab.?

CERINTHE, [Tourn.] Linn. Syst. ed. I. (1735). *BORAGINEÆ*, Benth. & Hook. f. ii. 864.
CERANTHE, Opiz, Seznam, 37 (1852).
acuta, Moench, Meth. 530 = major.
alpina, *Kit. ex Schult. Oestr. Fl.* ed. II. i. 353.—Europ.; As. Min.
alpina, Vis. Fl. Dalmat. Suppl. (1872) 94; fide Nym. Consp. 513 = longiflora.
aperta, [Clairv.] Man. Herb. 53 = major.
aspera, Roth, Catalect. i. 38 = major.
atropurpurea, Vis. ex A. DC. in DC. Prod. viii. 4 = retorta.
auriculata, Tenore, Ind. Sem. Hort., Neap. (1830) 10 = minor.
cilicica, Kotschy, ex Boiss. Fl. Or. Nov. dec. ii. 7 = minor.
cleistostoma, Boiss. & Sprun. Diagn. Ser. i. iv. 44 = minor.
connivens, [Clairv.] Man. Herb. 54 = minor.
echioides, Link, Sp. Pl. 187 = Onosma echioides.
glabra, Mill. Gard. Dict. ed. VIII. n. 2; DC. Fl. Fr. iii. 619 = alpina.
glabra, Scop. ex Nym. Consp. 513 = minor.
glauca, Moench, Meth. 530 = major.
gymnandra, Gasp. in Rendic. Acc. Sc. Napl. i. (1863) 73 = major.
hirsuta, *Wettstein, ex Steitz, in Denkschr. Acad. Wien*, i. (1850) 85.—Lycia.
longiflora, *Vis. App. Alt. ad Fl. Cors.* (1830) 3.—Italia.

CERINTHE:—
lurida, Salisb. Prod. 116 = minor.
macrophylla, Boiss. ex Boiss. Fl. Or. Nov. dec. ii. 7. = minor.
maculata, All. Fl. Pedem. i. 178 = alpina.
maculata, Linn. Sp. Pl. 137 = minor.
major, *Linn. i. c.* 136.—Europ.; As. Min.
minor, *Linn. l. c.* 137.—Europ.; As. Min.; Persia.
orientalis, Linn. Cent. i. 7 = Onosma orientalis.
parade aspera, Schur, Enum. Pl. Transs. 470 = minor.
purpurea, Vis. in Flora, xii. (1829) Erg. 8 = retorta.
quinquemaculata, Wahlenb Fl. Carpat. 51 = alpina.
retorta, *Sibth & Sm. Fl. Graec.* ii. 60. t. 171.—Graecia.
Smithiae, Kern. ex Nym. Consp. 513 = longiflora.
strigosa, Reichb. Fl. Germ. Excurs. 340 = major.
suevica, Mutzen, Reise Vened. i. 44; ex Reichb. Ic. Cnt. v. 40. t. 558 = alpina.
tenuiflora, *Bertol. Fl. Ital.* ii. 325 = longiflora.
vernicolor, Hall. ex Stend. Nom. ed. II. i. i. 336 = major.

CERINTHOPSIS, Kotschy, ex Benth. & Hook. f. Gen. ii. 947 (1876) = **Solenanthus**, Ledeb. (Boragin.).
hurdica, Kotschy, ex Boiss. Fl. Orient. iv. 271 = S. cerinthioides.

CERIONANTHUS, Schott, ex Roem. & Schult. Syst. iii. 48 (1818) = **Cephalaria**, Schrad. (Dipsac.).
alpinus, Schott, ex Roem. & Schult. l. c. = Ceph. alpina.
cornicinus, Schott, ex Roem. & Schult. l. c. 49 = Ceph. corniculata.
cretaceus, Schott, ex Roem. & Schult. l. c. 51 = Ceph. corniculata.
laevigatus, Schott, ex Roem. & Schult. l. c. 49 = Ceph. corniculata.
leucanthus, Schott, ex Roem. & Schult. l. c. 47 = Ceph. leucantha.
syriacus, Schott, ex Roem. & Schult. l. c. 46 = Ceph. syriaca.
tataricus, Schott, *ex Roem. & Schult. l. c. 51* = Ceph. tatarica.
Vaillantii, Schott, ex Roem. & Schult. l. c. 46 = Ceph. syriaca.

CERIOPS, Arn. in Ann. Nat. Hist. i. (1838) 368. *RHIZOPHOREÆ*, Benth. & Hook. f. i. 679.
Bovinianus, *Jul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 115.—Madag.
Candolleana, *Arn. in Ann. Nat. Hist.* i. (1838) 364.—Reg. trop.
Candolleana, Naves, in Blanco, Fl. Filip. ed. III. t. 415 = Roxburghiana.
Forsteniana, *Blume, Mus. Bot. Lugd.-Bat.* i. 143.—Ins. Moluc.
globulifera, Bor. ex Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 112 = Boviniana.
lucida, *Miq. Fl. Ind. Bat. Suppl.* i. 835.—Malaya.
mossambicensis, Klotzsch, in Peters, Reise Mossamb. Bot. i. 71 = Candolleana.
pandifera, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 218.—Polynes.
Roxburghiana, *Arn. in Ann. Nat. Hist.* i. (1838) 364.—Reg. trop.
Zippeliana, *Blume, Mus. Bot. Lugd. Bat.* i. 143.—N. Guinea.

CERISCUS, Gaertn. Fruct. i. t. 28 (1788) = **Randia**, Linn. (Rubiac.).
maleboricus, Gaertn. l. c. = R. dumetorum.

CERISCUS, Nees, in Flora, vii. (1825) 116 = **Taramna**, Gaertn. (Rubiac.).
fragrans, Nees, l. c. (= *Webera fragrans*).—Ind. or.; Malaya.

CERIUM, Lour. Fl. Cochinch. 125. *INC. SED.*
spinatum, *Lour. l. c.* 126.—China.

CEROCARPUS, Hassk. in Flora, xxv. (1842) Beibl. ii. 26 = **Eugenia**, Linn. (Myrtac.).
aqueus, Hassk. l. c. = E. aquea.
biforis, Colebr. in Wall. Cat. n. 3606 = E. Wallichii.

CEROCHILUS, Lindl. in Gard. Chron. (1854) 87 = **Hetaeria**, Blume (Orchid.).
rubens, Lindl. l. c. = H. rubens.

CEROPEGIA, Linn. Gen. ed. I. 65 (1737). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 779.
APEGIA, Neck. Elem. i. 251 (1790).
NIOTA, Adans. Fam. ii. 172 (1763).
SYSTEPHA, Burch. Trav. i. 546 (1822).
abyssinica, *Decne. in DC. Prod.* viii. 644.—Abyss.
acuminata, Dalz. & Gibs. Bomb. Fl. 117 = tuberosa.
acuminata, *Roxb. Pl. Corom.* i. 12. t. 8.—Ind. or.
affinis, *Wathe, in Linnaea*, xl. (1876) 318.—Abyssa.
africana, *R. Br. in Bot. Reg.* t. 626.—Afr. austr.
albiflora, *Hook. f. Fl. Brit. Ind.* iv. 75.—Ind. or.
ampliata, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
angustifolia, Dals. in Hook. Kew Journ. ii. (1850) 259 = attenuata.
angustifolia, Vahl, ex Decne. in DC. Prod. viii. 593 = Secamone saligna.
angustifolia, *Wight, Contrib.* 31.—Reg. Himal.
aphylla, Haw. Syn. Pl. Succ. 13 = dichotoma.
arcta, *Nimmo, in J. Grah. Cat. Bomb. Pl.* 116, nomen. —Ind. or.
arbtolochioides, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 268.—Afr. trop.
Arnottiana, *Wight, Contrib.* 32.—Ind. or.; Burma.
attenuata, *Hook. Ic. Pl.* t. 867.—Ind. or.; Burma.
australis, *Lodd. Cat. ex Loud. Hort. Brit.* 95, nomen. —Austral.
Barklyi, *Hook. f. Bot. Mag.* t. 6315.—Afr. austr.
Beddomei, *Hook. f. Fl. Brit. Ind.* iv. 75.—Ind. or.
biflora, *Linn. Sp. Pl.* 211.—Zeylan.
Bowkeri, *Harv. in Proc. Dubl. Zool. & Bot. Assoc.* i. (1859) 254.—Afr. austr.
brevicollis, *Hook. f. Fl. Brit. Ind.* iv. 74.—Ind. or.
brevituberata, Bedd. Ic. Pl. Ind. Or. t. 88 = Brachystelma Beddomei.
bulbosa, *Roxb. Pl. Corom.* i. 11. t. 7.—Ind. or.
campanulata, *G. Don, Gen. Syst.* iv. 112.—Afr. trop.
candelabriformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Candelabrum.
Candelabrum, Heyne, ex Decne. in DC. Prod. viii. 643 = bulbosa.
Candelabrum, *Linn. Sp. Pl.* 211.—Ind. or.
Candelabrum, Lour. Fl. Cochinch. 114 = Loureiri.
Candelabrum, Roxb. Hort. Beng. 21; Fl. Ind. ii. 27 = tuberosa.
Candelabrum, Thw. Enum. Pl. Zeyl. 199 = intermedia.
carnosa, *E. Mey. Comm. Pl. Afr. Austr.* 193.—Afr. austr.
ciliata, *Wight, Ic.* t. 1262.—Ind. or.
convolvuloides, *A. Rich. Tent. Fl. Abyss.* ii. 47.—Abyssin.
cordata, *Lour. Fl. Cochinch.* 114.—Cochinch.
Cumingiana, *Decne. in DC. Prod.* viii. 643.—Ins. Philipp.
curviflora, *Hassk. in Flora*, xl. (1857) 102.—Java.
Decaisneana, *Wight, Ic.* t. 1259.—Ind. or.
Decaisneana, Miq. ex Hook. f. Fl. Brit. Ind. iv. 42 = Tylophora tenuis.
dichotoma, *Haw. Syn. Pl. Succ.* 13.—Ind. or.
edulis, Hort. ex Decne. in DC. Prod. viii. 643 = bulbosa.
elegans, *Wall. in Bot. Mag.* t. 3015.—Ind. or.
Elliottii, *Hook. f. Fl. Brit. Ind.* iv. 70.—Ind. or.
enatifolia, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 52.—Ind. or.
esculenta, Edgew. in Journ. Linn. Soc. vi. (1862) 204 = bulbosa.
fimbriata, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
fimbriifera, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 53.—Ind. or.
fusca, *C. Bolle, in Bonplandia*, ix. (1861) 51.—Ins. Canar.
Gardneri, *Hook. Bot. Mag.* t. 5806; *Thw. Enum. Pl. Zeyl.* 199.—Zeylan.
gracilis, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 53.—Ind. or.
hirsuta, Hochst. ex Decne. in DC. Prod. viii. 644 = abyssinica.
hirsuta, *Wight & Arn. Contrib.* 30.—Ind. or.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 73. —Reg. Himal.
Horsfieldiana, *Miq. Fl. Ind. Bat.* ii. 528.—Java.
hunudhuliformis, *E. Mey. Comm. Pl. Afr. Austr.* 194. —Afr. austr.
intermedia, *Wight, Ic.* t. 1260.—Ind. or.

CEROPEGIA :—
Jacquemontiana, Decne. in DC. Prod. viii. 641 = hirsuta ?
juncea, *Roxb. Pl. Corom.* i. 12.—Ind. or.
lanceolata, Wight & Arn. Contrib. 31 = longifolia.
Lawii, *Hook. f. Fl. Brit. Ind.* iv. 67.—Ind. or.
linearis, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
longiflora, Poir. Encyc. ii. 177 = bulbosa.
longifolia, Hook. f. & Thoms. in Fl. Brit. Ind. iv. 72 = angustifolia.
longifolia, *Wall. Pl. As. Rar.* i. 56. t. 78.—Reg. Himal.
Loureiri, *G. Don, Gen. Syst.* iv. 112.—Cochinch.
luddu, *Wall. Pl. As. Rar.* ii. 33. t. 139.— Reg. Himal. ; Burma.
Lushii, R. Grah. in Bot. Mag. t. 3800 = bulbosa.
macrantha, *Wight, Contrib.* 31.—Reg. Himal.
maculata, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 52.—Ind. or.
madagascariensis, *Decne. in DC. Prod.* viii. 642.—Madag.
Metziana, *Miq. Anal. Bot. Ind.* iii. 11.—Ind. or.
Meyeri, *Decne. in DC. Prod.* viii. 645.—Afr. austr.
micans, *Nimmo, in Y. Grah. Cat. Bomb. Pl.* 118, nomen.—Ind. or.
macronata, Roth, Nov. Pl. Sp. 179 = tuberosa.
multiflora, *Baker, in Saund. Refug. Bot.* t. 10.—Afr. austr.
Munroi, Wight, Ic. t. 1254 = fimbriflora ?
mysorensis, *Wight, l. c.* t. 846.—Ind. or.
nilotica, *Kotschy, in Sitzb. Acad. Wien. Math. Nat.* ii. (1866) 356.—Afr. trop.
nitida, *Feur. Encyc. Suppl.* ii. 178.—Ins. S. Doming.
obtusa, *Lour. Cochinch.* 114.—Cochinch.
obtusiloba, *Fawc. in Forbes, Nat. Wand. in East. Archip.* 511.—Ins. Timor.
oculata, *Hook. Bot. Mag.* t. 4093.—Ind. or.
odorata, *Nimmo, in Y. Grah. Cat. Bomb. Pl.* 118.—Ind. or.
ophiocephala, Dals. in Hook. Kew Journ. ii. (1850) 259 = hirsuta.
palustris, Pursh, Fl. Am. Sept. i. 184 = Vincetoxicum palustre.
peltata, *Regel, Cat. Pl. Hort. Absahor.* 31.—Hab. ?
paradisica, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 13.—Malaya.
pubescens, *E. Mey. Comm. Pl. Afr. Austr.* 198.—Afr. austr.
pubescens, *Wall. Pl. As. Rar.* ii. 81. t. 187.—Reg. Himal.
pusilla, *Wight & Arn. in Wight, Contrib.* 31.—Ind. or.
ringens, *A. Rich. Tent. Fl. Abyss.* ii. 48.—Abyssin.
sagittata, Linn. Mant. 215 = Microloma sagittatum.
Sandersoni, *Decne. ex Hook. Bot. Mag.* t. 5792.—Natal.
sinuata, *Decne. ex A. Rich. Tent. Fl. Abyss.* ii. 47.—Abyssin.
sinuata, Poir. Encyc. Suppl. ii. 179 = Microloma linearе.
Sororia, *Harv. in Hook. Bot. Mag.* t. 5578.—Afr. austr.
sphæranantha, Wight & Arn. in Wight, Contrib. 31 = elegans.
sphæranthera, Decne. in DC. Prod. viii. 643 = præc.
spiralis, Hook. f. & Thoms. in Fl. Brit. Ind. iv. 67 = attenuata.
spiralis, *Wight, Ic.* t. 1267.—Ind. or.
squamulata, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 368. t. 9.—Arabia.
stapeliæformis, *Haw. in Phil. Mag.* (1827) 121.—Afr. austr.
stellata, *Regel, Cat. Pl. Hort. Absahor.* 31.—Hab. ?
Stephanotis, Poir. ex Decne. in DC. Prod. viii. 631 = Stephanotis lasiura.
Stephanotis, *Room. & Schult. Syst.* vi. 4.—Madag.
Steudneri, *Vatke, in Linnæa*, xl. (1876) 217.—Abyssin.
Stocksii, *Hook. f. Fl. Brit. Ind.* iv. 74.—Ind. or.
tenuiflora, Willd. Sp. Pl. i. 1276 = Microloma linearе.
tenuifolia, Linn. Mant. ii. 245 = Microloma linearе.
Thwaitesii, *Hook. Bot. Mag.* t. 6758.—Zeylan.
torulosa, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
trichanthe, *Hemsl. in Journ. Bot.* xxiii. (1885) 286.—China.
tuberosa, *Roxb. Pl. Corom.* i. 12. t. 9.—Ind. or.
tuberosa, Wall. Cat. n. 6145 A = intermedia.
variegata, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 369.—Arabia.

CEROPEGIA :—
Vignaldiana, *A. Rich. Tent. Fl. Abyss.* ii. 48.—Abyss.
vincæfolia, Hook. Bot. Mag. t. 3740 = hirsuta.
Walkeriæ, *Wight, Ic.* t. 1266.—Zeylan.
Wallichii, *Wight, Contrib.* 32.—Reg. Himal.
Wightii, *R. Grah. in Bot. Mag.* t. 3267.—Ind. or.

CEROPHORA, Rafin. Alsog. Am. 11 (1838) = **Myrios**, Tourn.
angustifolia, Rafin. l. c. = M. cerifera.
inodora, Rafin. l. c. = M. cerifera.
lanceolata, Rafin. l. c. = M. cerifera.
spicans, Rafin. l. c. 12 = M. cerifera.

CEROPHYLLUM, Spach, Hist. Vég. Phan. vi. 152 (1838) = **Ribes**, Linn. (Saxifrag.).
Douglasi, Spach, l. c. 153 = R. cereum.
tenuiriana, Spach, l. c. 154 = R. cereum.

CEROXYLON, Humb. & Bonpl. Pl. Aequin. I. t. 1, 1 b (1808). *PALMÆ*, Benth. & Hook. f. iii. 905.
BERTHOVENIA, Engelm. in Linnæa, xxxiii. (1864–65) 677.
KLOTZSCHIA, Karst. in Linnæa, xxviii. (1856) 251.
alpinum, Steud. Nom. ed. II. i. 337 = seq.
andicolum, *Humb. & Bonpl. Pl. Aequin.* I. t. 1.—N. Granat.
australe, Mart. Hist. Nat. Palm. iii. 314.—Juanis australis.
coarctatum, *H. Wendl. in Kerch. Palm.* 238.—N. Granat.
ferrugineum, *Regel, Gartenfl.* (1879) 163. t. 977.—N. Granat.
interruptum, *H. Wendl. in Bonplandia*, viii. (1860) 70. —N. Granat.
Klopstockia, *Mart. Hist. Nat. Palm.* iii. 314.—Venezuela.
niveum, Hort. Belg. ex H. Wendl. in Kerch. Palm. 239 = Diplothemium caudescens.
parvifrons, *H. Wendl. l. c.*—N. Granat.
pityrhyllum, Mart. ex H. Wendl. l. c. = Cocos pityrophylla.
quindiuense, *H. Wendl. in Bonplandia*, viii. (1860) 70.—N. Granat.
utile, *H. Wendl. l. c.*—N. Granat.
Vogelsianum, *H. Wendl. in Kerch. Palm.* 239.—N. Granat.

CERQUEIRIA, Berg, in Linnæa, xxvii. (1854) 5 = **Myrcia**, DC. (Myrtac.).
Sellowiana, Berg, l. c. 6.—hras.

CERQUEIRIA, Benth & Hook. f. Gen. i. 716 (1865) spnalm. = præc.

CERRARIA, Tausch, in Flora, xvii. (1834) i. 347, spnalm. = CERVARIA, Minuart = **Peucedanum**, Linn. (Umbellif.).

CERRIS, Rafin. Alsog. Am. 29 (1838) = **Quercus**, Linn.
australis, Rafin. l. c. = Q. Cerris.
austriaca, Rafin. l. c. = Q. Cerris.
crinita, Rafin. l. c. = Q. Cerris.
dulcis, Rafin. l. c.—Am. bor.
macrocarpa, Rafin. l. c. = Q. macrocarpa.
microphylla, Rafin. l. c. = Q. microphylla.
oliveformis, Rafin. l. c. = Q. macrocarpa.
palispinea, Rafin. l. c.—Am. bor.
salicifolia, Rafin. l. c. = Q. salicifolia.

CERUANA, Forsk. Fl. Aegypt. Arab. 168 (1775). *COMPOSITÆ*, Benth. & Hook. f. ii. 361.
fruticosa, Less. Syn. Comp. 209 = pratensis.
montana, Aucher, ex DC. Prod. vii. 287 = Chrysophthalmum sternutatorium.
pratensis, *Forsk. Fl. Aegypt. Arab.* p. lxxiv.—Aegypt.
rotundifolia, Cass. in Dict. Sc. Nat. xli. 128 = pratensis.
Schimperi, Boiss. Diagn. Ser. I. xi. 4 = Asteriscus Schimperi.
senegalensis, *DC. Prod.* v. 488.—Afr. trop.

CERUCHIS, Gaertn. ex Schreb. Gen. 548 (1791) = Spilanthes, Jacq. (Compos.).

CERVANTESIA, Ruiz & Pav. Prod. 39. t. 7 (1794). *SANTALACEÆ*, Benth. & Hook. f. iii. 222.
bicolor, *Cav. Ic.* iii. 49. t. 275.—Peruv.
Kunthiana, *Baill. Adansonia*, ii. (1861–62) 378. t. 11.—Peruv.
tomentosa, A. DC. Prod. xiv. 692 = bicolor.
tomentosa, H. B. & K. Nov. Gen. et Sp. vii. 139 = Kunthiana.
tomentosa, *Ruiz & Pav. Prod. Fl. Per.* iii. 20. t. 7.—Peruv.

CERVARIA, Minuart, ex Linn. Diss. Dass. 31 (1747) (Amœn. Acad. i. 142) ; Gaertn. Fruct. i. 90. t. 21 = **Peucedanum**, Linn. (Umbellif.).
alsatica, Gaud. Fl. Helv. ii. 327 = P. alsaticum.
angustifolia, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 295 = P. angustifolium ?
glauca, Gaud. Fl. Helv. ii. 326 = P. Cervaria.
lancea, Gaud. l. c. 327, in obs. = P. Cervaria ?
latifolia, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 295 = P. latifolium.
nigra, Bernh. Syst. Verz. Erf. 171 = P. Cervaria.
Orosselinum, Gaud. Fl. Helv. ii. 324 = P. Oreoselinum.
rigida, Moench, Meth. 98 = P. Cervaria.
Rivini, Gaertn. Fruct. 91. t. 21 = P. Cervaria.

CERVIA, Rodrig. ex Lag. Gen et Sp. Nov. 7 (1816). *CONVOLVULACEÆ* (*Cf.* Benth. & Hook. f. ii. 868).
saturejæfolia, *Rodrig. ex Lag. l. c.*—Hispan.

CERVICINA, Delile, Fl. Egypte, 150 (1813) = **Wahlenbergia**, Schrad. (Campanulac.).
campanuloides, Delile, l. c. t. 8. f. 2 = W. Cervicina.
tomentosa, Perr. ex Steud. Nom. ed. II. i. 337 ; ii. 740. = Vahlia tomentosa.

CERVISPINA, [Dill.] Moench, Meth. 686 (1794) = **Rhamnus**, Tourn.
cathartica, Moench, J. c. = R. cathartica.
infectoria, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 356 = R. infectoria.
saxatilis, Fourr. l. c. = R. saxatilis.
Villarsii, Fourr. l. c. = R. infectoria.

CESATIA, Endl. in Ann. Wien. Mus. ii. (1838) 200 = **Trachymene**, Rudge (Umbellif.).
ornata, Endl. l. c. = T. eriocarpa.
ovata, Steud. Nom. ed. II. i. 337, spnalm. = præc.

CESPEDESIA, Goudot, in Ann. Sc. Nat. Sér. III. ii. (1844) 368. *OCHNACEÆ*, Benth. & Hook. f. i. 320.
Bonplandi, *Goudot, l. c.* 370.—N. Granat.
macrophylla, *Sess. Bot. Voy. Herald*, 97.—Panama.
spathulata, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 647.—Peruv.

CESTICHIS, Thou. Orch. Afr. t. 90 (1822) = **Liparis**, Rich. (Orchid.).

CESTRINUS, Cass. in Dict. Sc. Nat. viii. 24 (1817) = **Centaurea**, Linn. (Compos.).
carthamoides, Cass. l. c. = Cent. Chamaerhaponticum.

CESTRON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = seq.

CESTRUM, Linn. Hort. Cliff. 491 (1737) ; Coroll. Gen. 9 (1737). *SOLANACEÆ*, Benth. & Hook. f. ii. 904.
HABROTHAMNUS, Endl. Gen. 667 (1836).
LOMBRIA, Rafin. Sylva Tellur. 56 (1838).
MEYENIA, Schlecht. in Linnæa, viii. (1833) 251.
PARQUI, Adans. Fam. ii. 219 (1763).
WADEA, Rafin. Sylva Tellur. 56 (1838).
acuminatissimum, *Dun. in DC. Prod.* xiii. 1. 637.—Guiana.
acuminatum, *Sweet, Hort. Brit.* ed. II. 368.—Mexic.
affine, *H. B. & K. Nov. Gen. et Sp.* iii. 60.—Ecuador.
affine, Pers. Syn. i. 230 = auriculatum.
alaternoides, *Desf. Tabl.* ed. I. 70.—Ind. occ.
albo-punctatum, *Dun. in DC. Prod.* xiii. 1. 635.—Peruv.
album, *Ferrero, ex Dun. l. c.* 605.—Cuba.
Amelanchier, *Dun. l. c.* 602.—Hab. ?
amictum, *Schlecht. in Linnæa*, viii. (1833) 64.—Bras.
amygdalifolium, *Dun. in DC. Prod.* xiii. 1. 635.—Mexic.

CESTRUM:—
anagyridium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = aeq.
Anagyris, *Dun. in DC. Prod.* xiii. 1. 608.—Mexic.
angustifolium, *Lodd. Bot. Cab.* t. 818.—Ind. occ.
arcuinervium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Amelanchier.
arvense, *Vell. Fl. Flum.* 101; iii. t. 7.—Bras.
atrovirens, Dun. in DC. Prod. xiii. 1. 648 = Senna atrovirens.
aurantiacum, *Lindl. Bot. Reg.* (1844) *Misc.* 71.—Guatimal.
aurantiacum, Wy. ex Steud. Nom. ed. II. i. 337 = Freylinia cestroides.
auriculatum, *L'Hérit. Stirp. Nov.* 71. t. 35.—Am. austr.
auriculatum, Ruiz & Pav. Fl. Per. ii. 28. t. 155 f. a = hediundinum.
axillare, Vell. Fl. Flum. 101; iii. t. 5 = laevigatum.
Bella-Sombra, *Dun. in DC. Prod.* xiii. 1. 632.—Ins. Canar.
Benthami, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 151.—Mexic.
Billbergianum, *Beurling, in Vet. Acad. Handl.* (1854) 140.—Panama.
bogotense, Willd. ex Roem. & Schult. Syst. iv. 807 = marquitense.
bonariense, *Dun. in DC. Prod.* xiii. 1. 666.—Reg. Argent.
bracteatum, *Link & Otto, Ic. Pl. Rar.* i. 11. t. 6.—Bras.
bupleurifolium, *Dun. in DC. Prod.* xiii. 1. 628.—Bras.
bustilloni, *H. B. & K. Nov. Gen. et Sp.* iii. 57.—N. Granat.
calycinum, *H. B. & K. l. c.* 58.—Peruv.
calycinum, Willd. ex Schlecht. in Linnaea, vii. (1832) 64 = viridiflorum.
campanulatum, Lam. Encyc. i. 688 = Acnistus aggregatus.
campestre, *Griseb. in Goett. Abh.* xxiv. (1879) 244.—Reg. Argent.
cancellatum, *Dun. in DC. Prod.* xiii. 1. 657.—Peruv.
candidum, *Sess. & Schmidt, in Flora*, xxvii. (1844) 496.—Mexic.
caulisforum, Jacq. Hort. Schoenb. iii. 41. t. 326 = Acnistus arborescens.
caulisforum, Sieber, ex Bercht. & Presl, Rostl. i. Solanac. 52 = Acnistus Plumieri.
chlamidatum, *Dun. in DC. Prod.* xiii. 1. 645.—Bras.
chloranthum, *Dun. l. c.* 636.—Ins. Trinitat.
citrifolium, *Retz. in Hofm. Phyt. Blaetter*, i. (1803) 36.—Hab. ?
Clausseni, *Dun. in DC. Prod.* xiii. 1 637.—Bras.
collinum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 160.—Bras.
confertiflorum, *Schlecht. in Linnaea*, xix. (1847) 263.—Mexic.
confertum, Mill. Gard. Dict. ed. VIII. n. 5 = vespertinum.
confertum, *Ruiz & Pav. Fl. Per.* ii. 29. t. 158.—Peruv.
conglomeratum, *Ruiz & Pav. l. c.* 29. t. 156.—Peruv.
corcovadense, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 160.—Bras.
cordatum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 224.—Bras.
coriaceum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 161.—Bras.
corymbosum, *Schlecht. in Linnaea*, vii. (1832) 57.—Bras.
cuspidatum, *Sendt. in Mart. Fl. Bras.* x. 223.—Bras.
cyanense, Schlecht. ex Dun. in DC. Prod. xiii. 1. 673 = Iochroma Schlechtendallianum.
daphnoides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 524.—Ind. occ.
depauperatum, *Dun. in DC. Prod.* xiii. 1. 660.—Ins. Guadalup.
discolor, *Dun. l. c.* 609.—Peruv.
disrumum, Domb. ex Dun. in DC. l. c. 667 = leptanthum.
disrumum, H. B. & K. Nov. Gen. et Sp. iii. 61 = Forsythi.
disrumum, *Linn. Sp. Pl.* 191.—Ind. occ
dubium, *Steud. Nom.* ed. II. i. 337.—Hab. ?
dumetorum, *Schlecht. in Linnaea*, vii. (1832) 61.—Bras.

CESTRUM:—
Ehrenbergi, *Schlecht. ex Dun. in DC. Prod.* xiii. 1. 613.—Mexic.
elegans, *Schlecht. in Linnaea*, xix. (1847) 261.—Mexic.
elongatum, Hort. ex Steud. Nom. ed. II i. 337 = candidum.
Endlicheri, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 151.—Peruv.
eriochiton, *Sendt. in Mart. Fl. Bras.* x. 217.—Bras.
euanthes, *Schlecht. in Linnaea*, vii. (1832) 60.—Bras.
exsiccatum, *Ledeb. in Schrad. N. Journ. Bot.* iv. (1810) 65.—Am. austr.
fasciculare, Otto, ex Dun. in DC. Prod. xiii. 1. 673 = Tetucephalus cestroides.
fasciculatum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 151.—Mexic.
fastigiatum, *Jacq. Hort. Schoenb.* iii. 44. t. 330.—Ind. occ.
fastigiatum, Jan, Cat. Hort. Parm. 189, 10 = diurnum.
Faucheri, *Dun. in DC. Prod.* xiii. 1. 640.—Hab. ?
ferrugineum, *A. Rich. Fl. Cub. Funerog.* ii 126 — Cuba.
floribundum, Willd. ex Roem. & Schult. Syst. iv. 807 = tenuiflorum.
foetidissimum, Jacq. Hort. Schoenb. iii. 43. t. 329 = nocturnum.
foetidum, *Medic. in Act. Acad. Theod Palat.* iv. *Phys.* (1780) 192.—Hab. ?
foetidum, Salisb. Prod. 111 = auriculatum.
Forsythi, *Dun. in DC. Prod.* xiii. 1. 604.—Ind. occ.
fruticosum, *Asubl. Pl. Gui.* i. 238.—Guiana.
Gardneri, *Sendt. in Mart Fl Bras.* x. 208.—Bras.
glaucescens, *Sendt. l. c.* 222.—Bras.
glomeratum, *Schott, ex Sendt. l. c.* 212.—Bras.
graciliflorum, *Dun. in DC. Prod.* xiv. i. 669.—Mexic.
granadense, Willd. ex Roem. & Schult. Syst. iv. 807 = tomentosum.
grandiflorum, Hort. ex Steud. Nom. ed. II. i. 337 = laurifolium.
grandistipulum, *Sendt. in Mart. Fl. Bras.* x. 214.—Bras
Hartwegi, *Dun. in DC. Prod.* xiii. 1 599.—Mexic.
Hedsunda, Lam. Encyc. i. 687 = auriculatum.
Hedsunda, Matr. ex Dun. in DC. Prod xiii. 1. 647 = hediundinum.
hediundinum, *Dun. l. c.*—Am austr.
heteroflexum, Buek, in DC. Prod. Index, iii. 98, sphalm. = retroflexum.
hirsutum, *Jacq. Hort. Schoenb.* iii. 41. t. 324.—Hab. ?
hirtellum, *Schlecht. in Linnaea*, vii. (1832) 62.—Mexic.
hirtum, Sieber, ex Sendt. in Mart. Fl. Bras. x. 210 = Poeppigii.
hirtum, *Sw. Prod. Veg. Ind. Occ.* 49.—Jamaic.
internedium, *Sendt. in Mart. Fl. Bras.* x. 221.—Bras.
involucratum, *Dun. in DC. Prod.* xiii. 1. 610.—Mexic.
jamaicense, Lam. Encyc. i. 687 = C. Parqui, vespertinum.
jasminiflorum, *Moric. ex Dun. in DC. Prod.* xiii. 1. 660.—Hab. ?
Kohautz, Bercht. & Presl, Rostl. i. Solanac. 52 = canliflorum.
laevigatum, *Schlecht. in Linnaea*, vii. (1832) 59.—Bras.
Lamberti, *Dun. in DC. Prod.* xiii. 1. 634.—Hab. ?
lanatum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) (reimpr. 15).—Mexic.
lanceolatum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 159.—Bras.
lanceolatum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 214.—Bras.
lanciflolium, *Schlecht. in Linnaea*, xix. (1847) 267.—Mexic.
lanuginosum, *Ruiz & Pav. Fl. Per.* ii. 30. t. 157.—Peruv.
laniatum, *Dun. in DC. Prod.* xiii. 1. 649.—Peruv.
latifolium, *Lam. Illustr.* ii. 5.—Ins. Trinitat.
latifolium, Sieber, ex Dun. in DC. Prod. xiii. 1. 636 = chlorathum.
laurifolium, *L'Hérit. Stirp. Nov.* 69. t. 34.—Am. austr.
laxiflorum, *Dun. in DC. Prod.* xiii. 1. 655.—Venezuela.
laxum, *Benth. Pl. Hartw.* 33. n. 207.—Mexic.
leptanthum, *Dun. in DC. Prod.* xiii. 1. 666.—Peruv.
leucocarpum, *Dun. l. c.* 650.—Am. austr.

CESTRUM:—
Lindeni, *Dun. in DC. Prod.* xiii. 1. 611.—Venezuela.
longiflorum, *Ruiz & Pav. Fl. Per.* ii. 28. t. 154.—Peruv.
longifolium, Poepp. ex Dun. in DC. Prod. xiii. 1. 657 = umbrosetanum.
Lorentzianum, *Griseb. in Goett. Abh.* xix. (1874) 217 — Am. austr.
lucidum, Pav. ex Dun. in DC. Prod. xiii. 1. 665 = servatum.
Landianum, *Dun. l. c.* 658.—Bras.
lundum, *Dun. l. c.* 659.—Bras.
lycioides, Lichst. ex Roem. & Schult. Syst. iv. 558 = Lycium cinereum.
lycoides, *Sendt. in Mart. Fl. Bras.* x. 208.—Bras.
macrophyllum, Hort. Par. ex Dun. in DC. Prod. xiii 1. 636 = laurifolium.
macrophyllum, Salzm. ex Dun. l. c. = Salzmanni.
macrophyllum, *Vent. Choix* 18.—Ind. occ.
marquitense, *H. B. & K Nov. Gen. et Sp.* iii. 57.—N. Granat.
Martii, *Sendt. in Mart. Fl. Bras.* x. 221.—Bras.
Mathewsii, *Dun. in DC. Prod.* xiii. 1. 637.—Peruv.
megalophyllum, Dun. l. c. 688 = macrophyllum.
melanochloranthum, *Dun. l. c.* 622.—N. Granat.
microphyllum, *Linden, ex Dun. l. c.* 623.—N. Granat.
Miersianum, *Wedd. Chlor. And.* ii. 97.—N. Granat.
Miersi, *Dun. in DC. Prod.* xiii 1. 671.—Bras.
montanum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 161.—Am. austr.
Moquinianum, *Dun. in DC. Prod.* xiii. 1. 646.—Cuba.
Moritzi, *Dun. l. c.* 619.—N. Granat.
mucronatum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 160.—Bras.
multiflorum, *Roem. & Schult. Syst.* iv. 553.—Am. austr.
multiflorum, Schlecht. in Linnaea, vii. (1832) 59 = Schlechtendalii.
multiflorum, Schott, ex Sendt. in Mart. Fl. Bras. x. 216 = laevigatum.
multinervium, Dun. in DC. Prod. xiii. 1. 611 = C. Parqui?
Mutisi, *Willd. ex Roem. & Schult. Syst.* iv. 807.—Am. austr.
nemanthum, *Dun. in DC. Prod.* xii. 1. 687.—Bras.
nervosum, Mill. Gard. Dict. ed. VIII. n. 8 = Tabernaemontana amygdalaefolia.
nitidum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) (reimpr. 19).—Mexic
nocturnum, Lam. Encyc. i. 687 = leucocarpum.
nocturnum, *Linn. Sp. Pl.* 191.—Am. austr.
oblongifolium, *Schlecht. in Linnaea*, xix. (1847) 266.—Mexic.
obovatum, *Sendt. in Mart. Fl. Bras.* x. 209.—Bras.
odontospermum, *Yacq. Hort. Schoenb.* iii. 44. t. 331.—Am. austr.
oliganthum, *Dun. in DC. Prod.* xiii. 1. 634.—Guiana.
oppositifolium, Lam. Illustr. ii t. 112. f. 2. n. 2279 = Acokanthera Lamarkii.
organense, Miers, in Hook. Lond. Journ. Bot. v. (1846) 158 = amictum.
ovatum, *Willd. ex Roem. & Schult. Syst.* iv. 807 = tenuiflorum.
pallidum, *Lam. Encyc.* i. 688.—Jamaic.
paniculatum, *H. B. & K. Nov. Gen. et Sp.* iii. 62.—Venezuela.
paniculatum, Schlecht. in Linnaea, xix. (1847) 262 = sylvaticum.
Parqui, Benth. Pl. Hartw. 24 = terminale.
Parqui, *L'Herit. Stirp. Nov.* 78.—Am. trop.
parviflorum, *Dun. in DC. Prod.* xiii. 1. 624.—Peruv.
parvifolium, *Willd. ex Roem. & Schult. Syst.* iv. 808.—Hab. ?
pauciflorum, *Nees & Mart. in Nov. Act. Acad. Caes. Leop.* xi. (1823) 79.—Bras.
paucyflorum, Willd. ex Roem. & Schult. Syst. iv. 808 = Willdenowii.
pedicellatum, *Sendt. in Mart. Fl. Bras.* x. 224.—Bras.
pedunculare, *Pav. ex Dun. in DC. Prod.* xiii. 1. 618.—Mexic.
pendulinum, Hort. Monsp. ex Dun. in DC. Prod. xiii. 1. 630 = undulatum.
pendulinum, *Yacq. Hort. Schoenb.* iii. 42. t. 327.—N. Granat.
periscaefolium, *Dun. in DC. Prod.* xiii. 1. 608.—Mexic.
peruvianum, Hort. Roth. ex Dun. l. c. 662 = C. Amelanchier.

CESTRUM:—
peruvianum, Willd. ex Roem. & Schult. Syst. iv. 807 = affine.
petiolare, H. B. & K. Nov. Gen. et Sp. iii. 58 = Sessea petiolaris.
Plukenetii, *Tausch, in Flora*, xix. (1836) 387.—Hab.?
Poeppigii, *Schlecht. in Linnaea*, xxvii. (1854) 510.—Ins. Martiniic.
Poeppigii, *Sendt. in Mart. Fl. Bras.* x. 210.—Bras.
polyanthum, *Sendt. in Mart. l. c.* vi. 216.—Bras.
polyanthum, Steud. Nom. ed. II. i. 387 = Schlechtendalii.
porphyreum, *Dun. in DC. Prod.* xii. t. 602.—Hab.?
porphyroneuron, *Schlecht. in Linnaea*, xxv. (1852) 199.—Hab.?
potaliaefolium, *Dun. in DC. Prod.* xiii. t. 638.—N. Granat.
Prieurei, *Dun. l. c.* 685.—Guiana.
propinquum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 147.—Mexic.
Pseudo-Quina, *Mart. in Flora*, xxi. (1838) *Beibl.* 21.—Bras.
psychotrifolium, *Schlecht. in Linnaea*, xix. (1847) 265.—Mexic.
pubescens, *Grisseb. in Goett. Abh.* xix. (1874) 217.—Reg. Argent.
pubescens, Lichst. ex Roem. & Schult. Syst. iv. 558 = Acokanthera pubescens.
racemosum, *Ruiz & Pav. Fl. Per.* ii. 29. t. 154.—Peruv.
racemosum, Tausch, in Flora, xix. (1836) 387 = dubium.
ramosissimum, *Nocca, ex Dun. in DC. Prod.* xiii. t. 678, nomen.—Hab.?
reflexum, *Sendt. in Mart. Fl. Bras.* x. 218.—Bras.
Regelii, *Planch. in Fl. des Serres*, Sér. I. ix. (1853-54) 229. t. 946.—Guatimal.
Requieni, *Dun. in DC. Prod.* xiii. 1. 661.—Hab.?
reticulatum, Willd. ex Roem. & Schult. Syst. iv. 808 = conglomeratum.
retrofractum, *Dun. in DC. Prod.* xiii. t. 655.—Bras.
ruscum, Hort. ex Dun. l. c. 603 = porphyreum.
roseum, *H. B. & K. Nov. Gen. et Sp.* iii. 59. t. 197.—Mexic.
rugosum, Schott, ex Sendt. in Mart. Fl. Bras. x. 215 = lanceolatum.
salicifolium, Hort. Monsp. ex Dun. in DC. Prod. xiii. 1. 618 = foetidissimum.
salicifolium, *Jacq. Hort. Schoenb.* iii. 421.—Venezuela.
salicifolium, Kunth, ex Spreng. Syst. i. 673 = C. Parqui.
Salzmanni, *Dun. in DC. Prod.* xiii. t. 626.—Bras.
scandens, Thib. ex Dun. l. c. 631 = nocturnum.
scandens, *Vahl, Eclog. Am.* i. 24.—Am. austr.
Schlechtendalii, *G. Don, Gen. Syst.* iv. 489.—Bras.
Schottii, Pohl, ex Sendt. in Mart. Fl. Bras. x. 218 = amictum.
Schottii, *Sendt. l. c.* 218.—Bras.
Sellowianum, *Sendt. l. c.* 219.—Reg. Argent.
semivestitum, *Dun. in DC. Prod.* xiii. t. 651.—Mexic.
Sendtnerianum, *Mart. ex Sendt. in Mart. Fl. Bras.* x. 215.—Bras.
servatum, *Dun. in DC. Prod.* xiii. t. 665.—Peruv.
sessiliflorum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 213.—Bras.
spermacoefolium, Willd. ex Roem. & Schult. Syst. iv. 808 = Bouvardia trifolia.
spicatum, *Mill. Gard. Dict.* ed. VIII. n. 4.—N. Granat.
spurium, Hort. Genev. ex Dun. in DC. Prod. xiii. t. 617 = C. Parqui.
stipulatum, Vell. Fl. Flum. 101; iii. t. 5 = bracteatum.
strictum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 206.—Bras.
strigillatum, *Ruiz & Pav. Fl. Per.* ii. 29. t. 156.—Peruv.
suberosum, *Jacq. Hort. Schoenb.* iv. 26. t. 452.—Hab.?
subpyrrherulentum, *Mart. in Flora*, xxi. (1838) 11. *Beibl.* 65.—Bras.
subsessile, *Vell. Fl. Flum.* 102; iii. t. 8.—Bras.
subtriflorum, *Dun. in DC. Prod.* xiii. t. 638.—Ins. Trinitat.
subuniflorum, *Dun. l. c.* 654.—Bras.
sylvaticum, *Dun. l. c.* 603.—Mexic.
tenuiflorum, *H. B. & K. Nov. Gen. et Sp.* iii. 61.—Venezuela.
terminale, *Dun. in DC. Prod.* xiii. t. 606.—Mexic.

CESTRUM:—
thryoideum, *H. B. & K. Nov. Gen. et Sp.* iii. 59.—Mexic.
tinctorium, *Jacq. Hort. Schoenb.* iii. 45. t. 332.—Am. austr.
Tinus, *Dun. in DC. Prod.* xiii. t. 614.—Bras.
tomentosum, *Linn. f. Suppl.* 150.—Am. austr.
tomentosum, Moç. & Sessé, ex Dun. in DC. Prod. xiii. 1. 601 = Benthami.
toxicarium, J. F. Gmel. Syst. 389, 1524 = Antiaris toxicaria.
tubulosum, *Sendt. in Mart. Fl. Bras.* x. 207.—Bras.
umbellatum, E. Mey. ex Meisn. in DC. Prod. xiv. 538 = Peddiea Dregei.
umbellatum, *Pauq. Diss. Bellad.* 16.—Hab.?
undulatum, *Ruiz & Pav. Fl. Per.* ii. 28. t. 155.—Peruv.
unibracteatum, *Dun. in DC. Prod.* xiii. t. 656.—Peruv.
velutinum, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877-78) 68.—Am. trop.
venatorium, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 79 = Acokanthera venenata.
venenatum, Burm. f. Fl. Cap. Prod. 5 = Acokanthera venenata.
venenatum, Lvm. Encyc. i. 688 = laurifolium.
venenatum, Mill. Gard. Dict. ed. VIII. n. 6 = pallidum.
venezum, Willd. ex Roem. & Schult. Syst. iv. 807 = Sessea petiolaris.
vespertinum, Bertero, ex Dun. in DC. Prod. xiii. t. 660 = parqui.
vespertinum, Hort. Val. ex Roem. & Schult. Syst. iv. 806 = Vestia lycioides.
vespertinum, *Linn. Mant.* ii. 206.—Ind. occ.
vespertinum, Poepp. ex Sendt. in Mart. Fl. Bras. x. 218 = Poeppigii.
vestitum, *Schlecht. in Linnaea*, vii. (1832) 65.—Bras.
vestitum, Hook. Ic. Pl. t. 331 = Sessea vestita.
viminale, *Sendt. in Mart. Fl. Bras.* x. 220.—Bras.
virgatum, Ruiz & Pav. Fl. Per. ii. 27 = C. Parqui.
viride, *Moric. ex Dun. in DC. Prod.* xiii. t. 606.—Mexic.
viridiflorum, *Hook. Bot. Mag.* t. 4022.—Bras.
Warscewiczii, *Klotzsch, in Otto & Dietr. Allg. Gartens.* xix. (1851) 362.—Am. austr.
Willdenowii, *Steud. Nom.* ed. II. i. 338.—Hab.?

CETRA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

CEVALLIA, Lag. in Varied. Cienc. fi. iv. (1805) 35. *LOASEAE*, Benth. & Hook. f. i. 808.
PETALANTHERA, Nutt. in Journ. Acad. Philad. vii. (1834) 107.
sinuata, *Lag. l. c.* 36.—N. Mexic.

CEYTOSIS, Munro, in Journ. Linn. Soc. vi. (1862) 54, err. typ. = Cryptsis, Ait. (Gramin.).

CHAEOISSARA, Fourn. Gram. Mex. 112; ex Benth. & Hook. f. Gen. iii. 1096 (1883). Gen. dub. *GRAMIN.*
hirsuta, *Fourn. l. c.*; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 588.—Mexic.

CHABRAEA, Adans. Fam. ii. 287 (1763) = Peplis, Linn. (Lythrar.).

CHABRAEA, DC. in Ann. Mus. Par. xix. (1812) 65. t. 5 = Leucosia, Lag. (Compos.).
abbreviata, Bert. 'Merc. Chil. (1829) 601.'—Cf. Bull. Férussac, xx. (1830) 109.—L. abbreviata.
Barrasiana, Remy, in C. Gay, Fl. Chil. iii. 399.—Chili.
Bertroviana, Steud. Nom. ed. II. i. 338.—Chili.
candidissima, DC. Prod. vii. 59 = L. candidissima.
canescens, Phil. in Linnaea, xxviii. (1856) 716.—Chili.
cinerea, DC. Prod. vii. 60 = L. cinerea.
concinna, Phil. in Linnaea, xxxiii. (1864-65) 138.—Chili.
coquimbana, Phil. in Anal. Univ. Chil. (1861) 50.—Chili.
daucifolia, Wedd. Chlor. And. i. 95.—Peruv.
elongata, Bert. 'Merc. Chil. (1829) 601.'—Cf. Bull. Férussac, xx. (1830) 109 = L. canescides.
fragrans, Phil. in Linnaea, xxix. (1857-58) 6.—Chili.

CHABRAEA:—
Gayana, Remy, in C. Gay, Fl. Chil. iii. 400.—Chili.
glabra, DC. in Deless. Ic. Select. iv. t. 89.—Chili.
glabriuscula, Phil. in Linnaea, xxviii. (1856) 715.—Chili.
glacialis, DC. in Deless. Ic. Select. iv. t. 91 = L. pulchella.
glandulosa, DC. Prod. vii. 59 = L. glandulosa, *D. Don. integrifolia*, Phil. in Linnaea, xxviii. (1856) 716.—Chili.
laciniata, Wedd. Chlor. And. i. 84. t. 10 = L. laciniata.
Landbeckii, Phil. in Linnaea, xxxiii. (1864-65) 192.—Chili.
lithospermifolia, DC. Prod. vii. 60.—Chili.
lyrata, Spreng. Syst. iii. 506 = Gerbera Anandria.
modesta, Phil. Fl. Atac. 98.—Chili.
multifida, DC. Prod. vii. 60.—Patagonia.
nutans, Iiland. C. Gay, Fl. Chil. vii. 397.—Chili.
nutans, Remy, in Ann. Sc. Nat. Sér. III. xii. (1849) 182.—Chili.
oligocephala, Phil. in Anal. Univ. Chil. (1872) 745 = L. oligocephala.
Poeppigii, Phil. in Linnaea, xxix. (1857-58) 3.—Chili.
polycladas, Remy, in C. Gay, Fl. Chil. iii. 399.—Chili.
prenanthoides, Bert. in 'Merc. Chil. 1829.'—*Cf.* Bull. Férussac, xx. (1830) 109.—Chili.
pulchella, Phil. in Anal. Univ. Chil. (1872) 744 = L. pulchella.
purpurea, DC. in Ann. Mus. Par. xix. (1812) 65, 71 = L. purpurea.
rosea, DC. in Deless. Ic. Select. iv. t. 90 = L. runcinata.
runcinata, Hook. Bot. Mag. t. 4116 = L. runcinata.
salina, Remy, in C. Gay, Fl. Chil. iii. 396. t. 41.—Chili.
scrobiculata, DC. Prod. vii. 59 = L. scrobiculata.
suaveolens, DC. l. c. = L. gossypina.
temerifolia, Phil. in Linnaea, xxviii. (1856) 715.—Chili.
tenuior, Bert. in 'Merc. Chil. 1829.'—Cf. Bull. Férussac, xx. (1830) 109 = *Berteroniana*.
tenuisecta, Sch. Bip. in Bonplandia, iv. (1856) 55, nomen.—Chili.
thermarum, Phil. in Anal. Univ. Chil. (1862) 380.—Chili.
tomentosa, DC. Prod. vii. 60 = L. Bridgesii.
viscida, Bert. in 'Merc. Chil. (1829) 601.'—Cf. Bull. Férussac, xx. (1830) 109.—Chili.
viscosa, Bert. ex DC. Prod. vii. 59 = L. runcinata.

CHADARA, Forsk. Fl. Aegypt. Arab. 105 (1775) = Grewia, Linn. (Tiliac.).
arborea, Forsk. l. c. = G. excelsa.
tenax, Forsk. l. c. = G. populifolia.
velutina, Forsk. l. c. = G. velutina.

CHADRA, T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 11, err. typ. = prsec.

CHADSIA, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 104. *LEGUMINOSAE*, Benth. & Hook. f. i. 497.
andravinensis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 891.—Madag.
colutefolia, *Baill. l. c.* 892.—Madag.
flammea, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 106.—Madag.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 891.—Madag.
grantica, *Baill. l. c.* 892.—Madag.
Lantziana, *Baill. l. c.* 891.—Madag.
salicina, *Baill. l. c.* 892.—Madag.
vernicolor, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 106.—Madag.

CHAELANTHUS, Poir. in Dict. Sc. Nat. viii. 50 (1817) spbalm. = Chaetanthus, R. Br. (Restiac.).

CHAELOTHILUS, Neck. Elem. ii. 14 (1790) = Gentiana, Tourn.

CHAENACTIS, DC. Prod. v. 359 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 401.
ACARPHAEA, Harv. & A. Gray, Pl. Fendl. 96 (1849).
ACICARPHAEA, Walp. Ann. ii. 877 (1852).
MACROCARPHUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vi. (1841) 376.
achilleaphila, Hook. & Arn. Bot. Beech. Voy. 354 = Douglasii.

CHAENACTIS:—
acteniaefolia, *A. Gray, in Proc. Am. Acad.* x. (1875) 74.—Calif.
attenuata, *A. Gray, l. c.* 78.—Arizona.
brachypappa, *A. Gray, l. c.* viii. (1872) 390.—Am. bor. occ.
carphoclinia, *A. Gray, in Bot. Mex. Bound.* 94.—Am. bor. occ.
Castelii, *A. Gray, Syn. Fl. N. Am.* ii. ii. 452.—Am. bor. occ.
douglasii, Nutt. in Journ. Acad. Sc. Philad. Ser. II. i. (1847) 177 = glabriuscula.
Douglasii, *Hook. & Arn. Bot. Beech. Voy.* 354.—Am. bor. occ.
filifolia, Harv. in A. Gray, Pl. Fendl. 98 = tenuifolia.
Fremonti, *A. Gray, in Proc. Am. Acad.* xix. (1883) 30.—Calif.
glabriuscula, *DC. Prod.* v. 659.—Calif.
heterocarpha, *A. Gray, Pl. Fendl.* 98.—Calif.
lanosa, *DC. Prod.* v. 659.—Calif.
macrantha, *Eaton, in Bot. King, Exp.* 171. t. 18.—Am. bor. occ.
nevadensis, *A. Gray, in Bot. Calif.* i. 391.—Calif.
Nevii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 30.—Am. bor. occ.
Parishii, *A. Gray, l. c.* xx. (1885) 299.—Calif.
santolinoides, *Greene, in Bull. Torrey Club*, ix. (1882) 17.—Calif.
stevioides, *Hook. & Arn. Bot. Beech. Voy.* 353.—Am. bor. occ.
suffrutescens, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 100.—Calif.
tanacetifolia, A. Gray, l. c. vi. (1864–65) 545 = heterocarpha.
tenuifolia, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 375.—Calif.
thysanocarpha, *A. Gray, in Proc. Am. Acad.* xix. (1883) 30.—Calif.
xantiana, *A. Gray, l. c.* vi. (1865) 545.—Calif.

CHAENANTHE, Lindl. Bot. Reg. (1838) Misc. 88 = Diadenatum, Poepp. & Endl. (Orchid.).
Barkeri, Lindl. l. c. = D. Barkeri.

CHAENANTHERA, Rich. ex DC. Prod. iii. 196 (1828) = Charianthus, D. Don (Melastomac.).
mucronata, Rich. l. c. 197 = Charianthus glaberrimus.

CHAENARRHINUM, Reichb. Consp. 128 (1828) = Linaria, Tourn. (Scrophul.).
creantiifolium, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 579 = L. origanifolia.
exile, Lange, in Kjoeb. Vidensk. Meddel. (1877–78) 228 = L. exilis.
flexuosum, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 580 = L. Langei.
marrocanum, Pomel, Nouv. Mat. Fl. Atl. 98 = L. marrocanum.
macropodum, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 579 = L. macropoda.
minus, Lange, l. c. 577 = L. minor.
origanifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 137 = L. origanifolia.
prostratum, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 578 = L. minor.
rubrifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 127 = L. rubrifolia.
rubrifolium, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 578 = L. rubrifolia.
serpyllifolium, Lange, l. c. = L. serpyllifolium.
tenellum, Lange, l. c. 581 = Anarrhinum tenellum.
thymifolium, Losc. ex Lange, in Kjoeb. Vidensk. Meddel. (1877–78) 228 = L. exilis.
villosum, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 580 = L. villosa.

CHAENESTHES, Mirb. in Hook. Lond. Journ. Bot. iv. (1845) 386 = Lochroma, Benth. (Solanac.).
corollifolia, Mirb. l. c. 388 = L. corollifolia.
fuchsioides, Mirb. l. c. 387 = L. fuchsioides.
gesneroides, Mirb. l. c. 386 = L. gesneroides.
lehmannii, Mirb. l. c. = L. lanceolata.
longipes, Don, in DC. Prod. viii. 469 = L. longipes.
lucinda, Mirb. in Hook. Lond. Journ. Bot. iv. (1845) 387 = L. lucinda.
ovalifolia, Mirb. l. c. 387 = L. ovalifolia.

CHAENOCARPUS, Juss. in Dict. Sc. Nat. viii. 50 (1817) = CHENOCARPUS, Neck. = Spermacoce, Linn. (Rubiac.).

CHAENOCEPHALUS, Griseb. Fl. Brit. W. Ind. 374 (1861). *COMPOSITÆ*, Benth. & Hook. f. ii. 382.
heterophyllus, *Griseb. in Goett. Abh.* xxiv. (1879) 196.—Reg. Argent.
macrophyllus, *Griseb. l. c.*—Reg. Argent.
petrobioides, *Griseb. Fl. Brit. W. Ind.* 374.—Ind. occ.
Sancho, *Griseb. in Goett. Abh.* xxiv. (1879) 196.—Reg. Argent.

CHAENOLOBIUM, Miq. Fl. Ind. Bat. Suppl. i. 302 (1860) = Ormocia, Jack (Legumin.).
decemjugum, Miq. l. c. = O. microsperma.
septemjugum, Miq. l. c. = O. microsperma.

CHAENOMELES, Lindl. Veg. Kingd. 560 (1847) = CHOENOMELES, Lindl. = Pyrus, Linn. (Rosac.).

CHAENOPHORA, Rich. ex Crueger, in Linnæa, xx. (1847) 119, err. typ. = seq.

CHAENOPLEURA, Rich. ex DC. Prod. iii. 197 (1828) = Miconia, Ruiz & Pav. (Melastomac.).
densiflora, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 549 = M. pusilliflora.
ferruginea, H. Cruger, in Linnæa, xx. (1847) 112 = M. hypoleuca.
fulva, Triana, in Goett. Abh. ix. (1861) 55 = M. fulva.
hypoleuca, Benth. in Hook. Journ. Bot. ii. (1840) 315 = M. hypoleuca.
lanceolata, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 549 = M. pusilliflora.
longifolia, Griseb. in Goett. Abh. ix. (1861) 55 = M. longifolia.
parviflora, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 548 = M. pusilliflora.
quadrangularis, Macf. Fl. Jamaic. ii. 96 = M. quadrangularis.
stelligera, Macf. l. c. 97.—Jamaic.
stenobotrys, DC. Prod. iii. 197 = M. quadrangularis.

CHAENORRHINUM, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 577 (1870), err. typ. = CHAENARRHINUM, Reichb. = Linaria, Tourn. (Scrophul.).

CHAENOSTOMA, Benth. in Hook. Comp. Bot. Mag. i. (1835) 374. *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 945.
FALMSTRUCKIA, Retz. f. Obs. Bot. Pugill. 15 (1810).
SUTTERA, Roth, Bot. Bemerk. 172 (1787).
aethiopicum, *Benth. l. c.* 375.—Aff. austr.
affine, *Benth. in Flora*, xxvii. (1844) 834.—Afr. austr.
calycinum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 374.—Aff. austr.
campanulatum, *Benth. l. c.*—Afr. austr.
cordatum, *Benth. l. c.* 377.—Afr. austr.
crenatum, *Benth. l. c.* 375.—Afr. austr.
dendatum, *Benth. l. c.* 375.—Afr. austr.
fastigiatum, *Benth. l. c.* 376.—Afr. austr.
floribundum, *Benth. l. c.*—Afr. austr.
foetidum, *Benth. l. c.* 377.—Afr. austr.
glabratum, *Benth. l. c.* 375.—Afr. austr.
hallmiifolium, *Benth. l. c.*—Afr. austr.
hispidum, *Benth. l. c.* 376.—Afr. austr.
integrifolium, *Benth. l. c.*—Afr. austr.
Krausianum, Bernh. in Flora, xxvii. (1844) 835 = Lyperia Krausiana.
laxiflorum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 374.—Aff. austr.
linifolium, *Benth. l. c.* 375.—Afr. austr.
marifolium, *Benth. l. c.* 376.—Afr. austr.
natalense, *Bernh. in Flora*, xxvii. (1844) 835.—Afr. austr.
parviflorum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 374.—Afr. austr.
pedunculosum, *Benth. l. c.* 377.—Afr. austr.
polyanthum, *Benth. l. c.* 375.—Afr. austr.
procumbens, *Benth. l. c.* 374.—Afr. austr.
pumilum, *Benth. l. c.* 375.—Afr. austr.
racemosum, *Benth. l. c.* 377.—Afr. austr.
revolutum, *Benth. l. c.* 376.—Afr. austr.
rotundifolium, *Benth. l. c.* 374.—Afr. austr.
subspicatum, *Benth. l. c.* 376.—Afr. austr.
villosum, T.C. Don, in Sweet, Hort. Brit. ed. III. 590 = Polycarena capensis.

CHAERADOPLECTRON, Benth. & Hook. f. Gen. iii. 996 (1883), err. typ. = CHOERADOPLECTRON, Schau. = Habenaria, Willd. (Orchid.).

CHAEROPHYLLUM, Linn. Syst. ed. I (1735). *UMBELLIFERÆ*, Benth. & Hook. f. i. 898.
BALANSAEA, Boiss. & Reut. Pugill. Pl. Nov. 49 (1853).
BRACHYTÆNIA, Bertol. Fl. Ital. iii. 191 (1837).
FIBERIA, Opiz, Seznam, 44 (1852).
FREYERA, Reichb. Handb. 291 (1837).
GRAMMOSCIADIUM, DC. Prod. iv. 232 (1830).
PHYSOCAULIS, Tausch, in Flora, xvii. (1834) 343.
RHYNCHOSTYLIS, Tausch, l. c. 343 (1834).
acuminatum, *Lindl. in Royle, Illustr. Bot. Himal.* 232.—Reg. Himal.
affine, Steud. in A. Rich. Tent. Fl. Abyss. i. 332 = Anthriscus sylvestris.
alpestre, Jord. Pugill. 75 = Villarsii.
alpinum, Vill. Prosp. 36 = Anthriscus sylvestris.
angelicaefolium, C. A. Mey. Verz. Pfl. Cauc. 180 = Meyeri.
angelicaefolium, *Bieb. Fl. Taur. Cauc.* iii. 240.—Oriens; Iberia.
angelicaefolium, DC. Prod. iv. 237 = byzantinum.
angulatum, Kit. ex Spreng. in Comm. Gott. Rec. ii. (1812) Umbell. Animad. 7. t. 1. f. 5 = Anthriscus sylvestris.
Anthriscus, Lam. Encyc. i. 335 = Anthriscus vulgaris.
arborescens, Linn. Sp. Pl. 259 = Astella spinosa?
aristatum, Thunb. Fl. Jap. 119 = Osmorhiza Claytoni.
aromaticum, *Linn. Sp. Pl.* 259.—Europ.
articulatum, Boec. ex DC. Prod. iv. 238 = procumbens.
Astrantiae, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 906.—As. Min.
aureum, *Linn. Sp. Pl.* ed. II. 370.—Europ.; As. Min.
aureum, Linn. Mant. ii. 256 = temulum.
australe, Crants, Cl. Umb. Emend. 76 = Scandix australis.
Bieberstenii, *Lag. ex Sweet, Hort. Brit.* ed. I. 190.—Reg. Caucas.
bifidum, Willd. ex DC. Prod. iv. 235 = procumbens.
Boscii, Steud. Nom. ed. II. i. 189 = procumbens.
brachycarpum, Bieb. ex Hoffm. Gen. Umb. i. 212 = millefolium.
bulbosum, *Linn. Sp. Pl.* 258.—Europ.; As. Min.
byzantinum, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 65.—Byzant.; As. Min.
cachemiricum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 891.—Reg. Himal.
cadomense, Schult. ex Steud. Nom. ed. II. i. 339 = Anthriscus sylvestris.
calabricum, *Guss. ex DC. Prod.* iv. 237.—Calabria.
californicum, Torr. in Pacif. Rail. Rep. iv. 98 = Podosciadium californicum.
canadense, Pers. Syn. i. 320 = Cryptotaenia canadensis.
capense, Thunb. Prod. Pl. Cap. 51 = Annesorhiza capensis.
capnoides, *Benth. & Hook. f. Gen.* i. 898.—Reg. Himal.
Cerefolium, Crants, Stirp. Austr. ed. I. fasc. III. 70 = Anthriscus Cerefolium.
chilense, Foir. Encyc. v. 145 = Scandix chilensis.
Cicutaria, Vill. Prosp. 36 = hirsutum.
ciliatum, *Kit. ex Kanitz, Relig. Kitaib.* 103.—Croatia.
Claytoni, Pers. Syn. i. 320 = Osmorrhiza brevistylis.
coloratum, *Linn. Mant.* 57.—Dalmat.; Banat.
cretieum, *Boiss. & Heldr. Diagn.* Ser. I. x. 51.—Ins. Cretæ.
crinitum, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 63.—Mesopot.; Persia.
Cyminum, Fisch. ex DC. Prod. iv. 230 = Conopodium Cyminum.
dasycarpum, Nutt. ex Torr & Gray, Fl. N. Am. i. 639 = Tainturieri.
daucifolium, Desf. Cat. Hort. Par. ed. III. 300, 405 = Anthriscus alpestris.
daucophyllum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 638 = Tainturieri.
divaricatum, Pair. Encyc. Suppl. iv. 84 = Anthriscus fumariioides.
dulce, Fisch. ex Stend. Nom. ed. II. i. 339 = Osmorrhiza Claytoni.
elegans, *Gaud. Fl. Helv.* ii. 364.—Helvet.
Friedrichsthalii, Cesati, in Bibl. Ital. lxxv. 338 = Anthriscus temerrima.
geniculatum, Gilib. Fl. Lituan. ii. 29 = temulum.

CHAEROPHYLLUM :—
glaberrimum, Pers. Syn. i. 320 (= *Bunium glaberrimum*).—Algeria.
gracile, Bess. Hort. Cremeo. (1822) = Conopodium Cynimum.
gracilimum, Klotzsch, in Bot. Ergeb. Waldem. Reise, 149 = Vicatia conifolia.
grandiflorum, Crantz, Cl. Umb. Emend. 77; Lam. Encyc. i. 686 = Scandix grandiflora.
Heldreichii, *Orph. in Boiss. Diagn.* Ser. II. ii. 104.—Graecia.
hirsutum, *Linn. Sp. Pl.* 258.—Europ.; As. Min.
hirsutum, Pall. Tabl. Taur. 49 = temulum.
hirsutum, Vill. ex Koch, Syn. Fl. Germ. ed. I. 317 = Villarsii.
hispidum, Thunb. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 64 = Cascalis Anthriscus.
Hladnikianum, Reichb. ex Heyph. Nom. ii. 185 = Anthriscus fumarioides.
humile, *Bieb. Fl. Taur. Cauc.* iii. 340.—Reg. Caucas.
hybridum, Tenore, Fl. Neap. Prod. p. lxvi = aureum.
infestum, Salisb. Prod. 167 = Anthriscus sylvestris.
Kotschyi, Fenzl, ex Boiss. Fl. Orient. ii. 918 = Anthriscus Kotschyi.
lactescens, Rochel, ex Steud. Nom. ed. II. i. 339 = Anthriscus sylvestris.
laevigatum, Vis. Fl. Dalm. iii. 65 = bulbosum.
lasiolenum, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 903. —As Min.
leucolenum, *Boiss. l. c.* 908.—Armenia.
libanotinum, *Boiss. & Kotschy, Diagn.* Ser. II. vi. 89. —Syria.
lucidum, Desf. Cat. Hort. Par. ed. III. 200 = Anthriscus nemorosa.
macropodum, *Boiss. in Ann. Sc. Nat.* Ser. III. ii. (1844) 64.—Persia.
macrospermum, *Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc.* (1838) 227.—Reg. Caucas.
maculatum, Willd. Enum. Hort. Berol. Suppl. 15 = aureum.
magellense, *Tenore, Prod. Fl. Neap. App.* iv. 15.—Neapol.
Meyeri, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 150.—Persia.
millefolium, *DC. Prod.* iv. 236.—Reg. Caucas.
millefolium, Klotzsch, in Bot. Ergeb. Waldem. Reise, 149 = Vicatia millefolia.
mintinum, *Vend. Fasc. Pl.* 11.—Italia.
monogynum, Kit. ex Link, Enum. Hort. Berol. i. 281 = aureum.
neglectum, *Zinger, in Bull. Soc. Nat. Mosc.* lvi. (1881) i. 213.—Rossia.
nemorosum, Hoffm. Gen. Umb. 45 = Anthriscus nemorosa.
nemorosum, Lag. ex DC. Prod. iv. 234 = Anthriscus Cerefolium.
nitens, Turcz. ex Bernh. in Linnaea, xii. (1838) Litt. 76 = Anthriscus nitens.
nitidum, Wahlenb. Fl. Carp. 88 = Anthriscus alpestris.
nodosum, *Cranz, Cl. Umb. Emend.* 76; *Lam. Encyc.* i. 685.—Europ.; As. Min.; Afr. bor.
odoratum, Cranz, l. c. 75; Lam. l. c. 688; Fl. Fr. iii. 439 = Myrrhis odorata.
orientale, Willd. ex Boiss. Fl. Orient. ii. 906 = byzantinum.
orthostylum, *Transc. in Act. Hort. Petrop.* iii. 11. (1875) 278.—Reg. Caucas.
pallescens, Presl, ex DC. Prod. iv. 237 = hirsutum.
palustre, Lam. Encyc. i. 688 = hirsutum.
Pecten-Veneris, Crantz, Stirp. Austr. ed. I. fasc. ii. 66; ed. II. 139 = Scandix Pecten-Veneris.
pinnatifidum, Poir. Encyc. v. 144 = Scandix pinnatifida.
polonicum, *Zapałowbrodski, ex Rostaf. Fl. Polon. Prod.* 113.—Europ.
Prescottii, DC. Prod. iv. 236 = bulbosum.
procumbens, *Crantz, Cl. Umbell. Emend.* 77.—Am. bor.
reflexum, Aitch. in Journ. Linn. Soc. xiii. (1882) 164 = villosum.
reflexum, *Lindl. in Royle, Illustr. Bot. Himal.* 233.—Reg. Himal.
rigidum, Host, ex Nym. Consp. 300 = Villarsii.
roseum, *Bieb. Fl. Taur. Cauc.* i. 234.—Reg. Caucas.
rostratum, Lam. Encyc. i. 685 = Scandix Pecten-Veneris.
satioum, Bieb. Fl. Taur. Cauc. i. 232 = Cascalis trichosperma.
satioum, Lam. Fl. Fr. iii. 436 = Anthriscus Cerefolium.

CHAEROPHYLLUM :—
scabrum, Thunb. Fl. Jap. 119 = Cascalis Anthriscus.
sicutum, Gmel. Fl. Sic. Prod. i. 362 = Anthriscus sicula.
sphallerocarpus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 367.—Soongaria.
sylvestre, Bieb. Fl. Taur. Cauc. i. 231 = maculatum.
sylvestre, Linn. Sp. Pl. 258 = Anthriscus sylvestris.
syriacum, *Hempr. & Ehrenb. ex Boiss. Fl. Orient.* ii. 904.—Syria.
Teinturieri, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 47.—Texas.
temulentum, Linn. Fl. Suec. 94 = temulum.
temuloides, *Boiss. in Ann. Sc. Nat.* Ser. III. ii. (1844) 64.—Persia.
temulum, *Linn. Sp. Pl.* 258.—Europ.; Afr. bor.; Reg. Caucas.
temulifolium, *Petr. Encyc. Suppl.* iv. 842.—Marocco.
tenuifolium, Stev. in Hoffm. Gen. Umbell. ed. II. 215 = millefolium.
tomentosum, *Petr. ex Steud. Nom.* ed. II. i. 339.—Hab.?
torquatum, DC. Fl. Suppl. 505 = Anthriscus alpestris.
trapezuntinum, Boiss. Fl. Orient. ii. 903 = temuloides.
trichospermum, Lam. Encyc. i. 685 = Cascalis trichosperma.
tuberculosum, Poir. Encyc. Suppl. iv. 842 = Anthriscus fumarioides.
tuberosum, C. A. Mey. ex Regel, in Act. Hort. Petrop. v. (1877) 587 = Carum cylindricum.
tuberosum, *Royle, Illustr. Bot. Himal.* 231.—Reg. Himal.
tumidum, Gilib. Fl. Lituan. ii. 98 = Anthriscus sylvestris.
umbrosum, Jord. Obs. Pl. Crit. vii. (1849) 80 = hirsutum.
verticillatum, Pers. Syn. i. 320 = bulbosum, Anthriscus vulgaris.
Villarsii, *Koch, Syn. Fl. Germ.* ed. I. 317.—Europ.
villosum, *Wall. Cat.* n. 858.—Reg. Himal.
villosum, Wall. l. c. partim = reflexum.

CHAETACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1886) = **Calophanes**, D. Don (Acanthac.).
Burchellii, Nees, in DC. Prod. xi. 462 = Cal. Buricot.
costatus, Nees, l. c. = Calophanes costatus.
glandulosus, Nees, l. c. = Calophanes Persoonii.
Persoonii, Nees, in Linnaea, xv. (1841) 357 = Calophanes Persoonii.
repandus, A. Gray, in Proc. Am. Acad. v. (1861–62) 349.—Ins. Fiji.
setigera, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 = Calophanes Persoonii.

CHAETACHLAENA, D. Don, in Trans. Linn. Soc. xvi. (1830) 156 = Crocosmia, DC. (Compos.).
acerifolia, D. Dietr. Syn. Pl. iv. 1565 = O. acerifolia.
annua, D. Dietr. l. c. = O. annua.
Cumingii, D. Dietr. l. c. = O. Cumingii.
odorata, D. Don, in Trans. Linn. Soc. xvi. (1830) 256 = O. odorata.

CHAETACHME, Planch. in Ann. Sc. Nat. Ser. III. x. (1848) 340. *URTICACEAE*, Benth. & Hook. f. iii. 356.
arietina, *Planch. l. c.* 341.—Afr. austr.
madagascariensis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 443.—Madag.
Meyeri, Harv. Thes. Cap. t. 16 = aristata.
nitida, *Planch. & Harv. in Harv. Thes. Cap.* 16.—Afr. austr.

CHAETADELPHA, A. Gray, ex S. Wats. in Am. Naturd. vii. (1873) 301; et in Proc. Am. Acad. ix. (1874) 218. *COMPOSITAE*, Benth. & Hook. f. ii. 1224.
Wheeled, *A. Gray, ll. cc.*—Arizona.

CHAETAEA, Jacq. Enum. Pl. Carib. 2 (1760) = **Buettneria**, Loefl. (Sterculi.).
aculeata, Jacq. l. c. 17 = B. carthaginensis.

CHAETAGASTRA, Cronq. in Linnaea, xx. (1846) 161, sphalm. = CHAETOGASTRA, DC. = Tithononiina, Aubl. (Meliost.).

CHAETANTERA, Less. Syn. Comp. 111 (1828) = Chaetanthera, Ruiz & Pav. (Compos.).

CHAETANTHERA, Nutt. in Journ. Acad. Philad. vii. (1834) 111 = **Chaetopappa**, DC. (Compos.).
asteroidea, Nutt. l. c. = Chaetopappa asteroides.

CHAETANTHERA, Ruiz & Pav. Prod. Fl. Per. 106. t. 28 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 496.
ALENDATA, Remy, in C. Gay, Fl. Chil. iii. 330. t. 88 (1849).
CARMELITA, C. Gay, in DC. Prod. vii. 14 (1838).
CHERINA, Cass. in Dict. Sc. Nat. viii. 487 (1817).
CHONDROCHILUS, Phil. in Linnaea, xxviii. (1856) 711.
EGANIA, Remy, in C. Gay, Fl. Chil. iii. 324 (1849).
ELACHIA, DC. Prod. vii. 256 (1838).
EUTHERIRIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 257.
MINUTHODES, Phil. ex Benth & Hook. f. Gen. ii. 496 (1879).
ORIASTRUM, Poepp. & Endl. Nov. Gen. et Sp. iii. 50. t. 257 (1847).
PROSELLA, D. Don, in Trans. Linn. Soc. xvi. (1830) 254.
TYLLOMA, D. Don, l. c. 238 (1830).
argentea, D. Don, in Phil. Mag. (1833) 392 = chilensis.
argentea, *Phil. in Anal. Univ. Chil.* (1872) 780.—Chili.
Berteriana, *Less. Syn. Comp.* 111.—Chili.
Berteroana, Hook. & Arn. in Comp. Bot. Mag. i. (1835) 104 = prasea.
Caput-Tringae, Less. ex DC. Prod. vii. 32 = limbata.
chilensis, *DC. in Ann. Mus. Par.* xix. (1812) 70.—Chili.
chilensis, Hook. & Arn. in Bot. Beech. Voy. 29 = lanosa.
ciliaris, Bert. ex Steud. Nom. ed. II. i. 340 = spathulata.
ciliata, *Ruiz & Pav. Syst. Veg. Per. et Chil.* 190.—Chili.
crenata, *Remy, in C. Gay, Fl. Chil.* iii. 302.—Chili.
crispa, *Phil. in Anal. Univ. Chil.* (1872) 780.—Chili.
dealbata, *Hook. & Arn. in Comp. Bot. Mag.* i. (1835) 104.—Chili.
debilis, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 287.—Chili.
elegans, *Phil. in Linnaea*, xxviii. (1856) 718.—Chili.
erysigioides, D. Don, in Phil. Mag. (1833) 391 = tenuifolia.
fabellata, D. Don, l. c. = multicaulis.
glandulosa, *Remy, in C. Gay, Fl. Chil.* iii. 311. t. 85. f. 1.—Chili.
grandiflora, *Steud. Nom.* ed. II. i. 340.—Chili.
humilis, *Phil. in Anal. Univ. Chil.* (1872) 740.—Chili.
incana, *Poepp. ex Less. in Linnaea*, v. (1830) 284.—Chili.
Kunthiana, Less. Syn. Comp. 114 = tenella.
limbata, *Less. l. c.* 116.—Chili.
linearifolia, Poepp. ex Steud. Nom. ed. II. i. 340 = linearis.
linearis, *Poepp ex Less. Syn. Comp.* 112.—Chili.
linifolia, *Bert. ex Less. l. c.*—Chili.
magellanica, Spreng. Syst. iii. 503 = Perezia lactucoides.
microphylla, Hook. & Arn. in Comp. Bot. Mag. i. (1835) 104 = linifolia.
moschiodora, *Less. Syn. Comp.* 113.—Chili.
multicaulis, *DC. Prod.* vii. 31.—Chili.
multiflora, Humb. & Bonpl. Pl. Aequin. ii. 168 = Perezia multiflora.
peruviana, *A. Grey, in Proc. Am. Acad.* v. (1861) 144. —Peruv.
pinnatifida, Humb. & Bonpl. Pl. Aequin. ii. 170 = Perezia pinnatifida.
pratensis, D. Don, in Phil. Mag. (1833) 391 = multicaulis.
pungens, Humb. & Bonpl. Pl. Aequin. 146. t. 127 = Perezia pungens.
pusilla, *Hook. & Arn. in Comp. Bot. Mag.* i. (1835) 106.—Chili.
ramosissima, Hook. & Arn. l. c. 104 = moschiodora, tenella.
rarorensis, Spreng. Syst. iii. 503 = Perezia recurvata.
recurvata, D. Don, in Phil. Mag. (1833) 391 = lanosa.
serrata, Lag. ex DC. Prod. vii. 30 = chilensis.
sonchifolia, Ruiz & Pav. Syst. Veg. Per. et Chil. 191 = chilensis.
spathulata, *Poepp. ex Less. in Linnaea*, v. (1830) 285. —Chili.
spinulosa, Cass. Opusc. ii. 188 = tenuifolia.
tenella, *Less. Syn. Comp.* 114; *DC. in Dietsch. An. Sci.* iv. t. 58.—Chili.
tenuifolia, *D. Don, in Phil. Mag.* (1833) 391.—Chili.
validiflora, *Phil. in Linnaea*, xxviii. (1856) 718.—Chili.
villosa, *D. Don, in Phil. Mag.* (1833) 391.—Chili.
Vollemanni, *Phil. in Anal. Univ. Chil.* (1872) 740.—Chili.

CHAETANTHUS, R. Br. Prod. 251 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1094.
PSEUDOMAPALIUS, Steud. Syn. Pl. Cyp. 266 (1855). leptocarpoides, *R. Br. l. c.*—Austral.

CHAETARIA, Beauv. Agrost. 80. t. 6 (1812) = **Aristida**, Linn. (Gramin.).
Adcensionis, Roem. & Schult. Syst. ii. 390 = A. Adcensionis.
affinis, Roem. & Schult. Syst. ii. Mant. 210 = A. purpurascens.
Antillarum, Roem. & Schult. Syst. ii. 395 = A. Antillarum.
Ascensionis, Beauv. Agrost. 80 = A. Adcensionis.
bipartita, Nees, Fl. Afr. Austr. i. 187 = A. bipartita.
bromoides, Roem. & Schult. Syst. ii. 396 = A. setifolia.
caerulescens, Beauv. Agrost. 80 = A. coerulescens.
calicina, Beauv. l. c. = A. calycina.
canariensis, Beauv. l. c. = A. coerulescens.
capensis, Beauv. l. c. = A. capensis.
capillacea, Beauv. l. c. = A. capillacea.
capillaris, Nees, Agrost. Bras. 358 = A. capillacea.
cearensis, Roem. & Schult. Syst. ii. 396 = A. setifolia.
congesta, Nees, Fl. Afr. Austr. 189 = A. congesta.
curvata, Nees, l. c. 186 = A. coerulescens.
cyanantha, Nees, ex Steud. Syn. Pl. Gram. 141 = A. cyananthe.
deprauta, Beauv. Agrost. 80 = A. Adcensionis.
divaricata, Beauv. l. c. = A. divaricata.
divaricata, Nees, Agrost. Bras. 357 = A. longifolia.
elatior, Beauv. l. c. = A. coerulescens.
elliptica, Nees, Agrost. Bras. 359 = A. elliptica.
fasciculata, Roem. & Schult. Syst. ii. Mant. 378 = A. fasciculata.
festucoides, Beauv. Agrost. 80 = A. Adcensionis.
Forsbolii, Nees, Fl. Afr. Austr. 188 = A. paniculata.
furcata, Beauv. Agrost. 80 (= *Eutriana juncifolia*).—Ind. occ.
gibbosa, Nees, Agrost. Bras. 353 = A. gibbosa.
gigantea, Beauv. Agrost. 80 = A. Adcensionis.
glauca, Nees, in Linnaea, xix. (1847) 668 = A. longiramea.
gossypina, Beauv. Agrost. 80; Roem. & Schult. Syst. ii. 391 = A. gossypina.
Hystrix, Beauv. l. c. = A. Hystrix.
humilis, Roem. & Schult. Syst. ii. 396 = A. humilis.
Hystrix, Roem. & Schult. l. c. 390 = A. Hystrix.
interrupta, Beauv. Agrost. 80 = A. interrupta.
laevis, Nees, Agrost. Bras. ii. 354 = A. laevis.
Lamarckii, Roem. & Schult. Syst. ii. 393 = A. stipiformis.
lanceolata, Beauv. Agrost. 80 = A. lanceolata.
mauritiana, Nees, Fl. Afr. Austr. 186 = A. Adcensionis.
megapotamica, Roem. & Schult. Syst. ii. Mant. 378 = A. megapotamica.
murina, Nees, ex Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1848) 160 = A. Royleana.
nana, Nees, ex Steud. Nom. ed. II. i. 240 = A. nana.
oligantha, Beauv. Agrost. 80 = A. oligantha.
pallens, Beauv. l. c. = A. oligantha.
purpurascens, Beauv. l. c. = A. coerulescens.
racemosa, Beauv. l. c. = A. racemosa.
ramosa, Roem. & Schult. Syst. ii. 397 = A. ramosa.
recurvata, Roem. & Schult. l. c. = A. recurvata.
Royleana, Nees, ex Steud. Syn. Pl. Gram. 148 = A. Royleana.
setacea, Beauv. Agrost. 80 = A. setacea.
setifolia, Roem. & Schult. Syst. ii. 396 = A. setifolia.
Siberiana, Roem. & Schult. Syst. ii. Mant. 378 = A. Hystrix.
spadicea, Roem. & Schult. Syst. ii. 397 = A. spadicea.
spadicea, Nees, Agrost. Bras. 365 = A. tincta.
squarrosa, Roem. & Schult. Syst. ii. Mant. 377 = A. squarrosa.
stipaeformis, Beauv. Agrost. 80 = Avena stipaeformis.
stricta, Beauv. l. c. = A. stricta.
tenuifolia, Beauv. l. c. = Panicum Tenerifae.
tincta, Nees, Agrost. Bras. 366 = A. tincta.
trichoides, Nees, in Hook. Kew Journ. ii. (1850) 101 = A. Cumingiana.
tuberculata, Roem. & Schult. Syst. ii. Mant. 211 = A. Adcensionis.
vaginata, Beauv. Agrost. 80 = A. vaginata.

CHAETHYMENIA, Hook. & Arn. Bot. Beech. Voy. 296. t. 69 (1841) = **Jaumea**, Pers. (Compos.).
pedunculariu, Hook. & Arn. l. c. = J. pedunculariu.

CHANTRIUM, Nees, Agrost. Bras. 369 (1829). *GRAMINEAE*, Benth. & Hook. f. iii. 1104.
BERCHTOLDIA, Presl, Rel. Haenk. 223. t. 45 (1830). bromoides, *Benth. in Journ. Linn. Soc.* xix. (1881) 46.—Mexic.
festucoides, *Nees, Agrost. Bras.* 370.—Bras.

CHAETOBROMUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836) = **Danthonia**, DC. (Gramin.).
Dregeanus, Nees, Fl. Afr. Austr. 248 = D. Dregeana.
fasciculariu, Nees, l. c. 241 = D. fasciculariu.
interceptus, Nees, l. c. 242 = D. interceptu.
involucratus, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 = D. involucrata.

CHAETOCALYX, DC. Mém. Lég. 362 (1826). *LEGUMINOSAE*, Benth. & Hook. f. i. 513.
BONNINGHAUSIA, Spreng. Syst. iii. 245 (1826).
PLANARIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 416.
RHADINOCARPUS, Vog. in Linnaea, xii. (1838) 108.
scutiflora, *Benth. in Mart. Fl. Bras.* xv. i. 75.—Am. austr.
brasiliensis, *Benth. l. c.*—Am. austr.
hebecarpa, *Benth. l. c.* 76.—Am. austr.
latifolia, *Benth. l. c.* 75.—Am. austr.
latifolia, *Benth. Bot. Voy. Sulph.* 81. t. 30.—Panama.
longiflora, *Benth. ex A. Gray, Bot. U. St. Expl. Exped.* i. 168.—Bras.
parviflora, *Benth. in Mart. Fl. Bras.* xv. i. 74.—Am. austr.
polyphylla, *Benth. l. c.* 76.—Am. austr.
pubescens, *DC. Prod.* ii. 248.—Ins. S. Doming. Schottii, *Torr. in Bot. Mex. Bound.* 56.—N. Mexic.
vincentina, *DC. Prod.* ii. 248.—Ins. S. Vincent.
Wrightii, *A. Gray, Pl. Wright.* i. 51.—Am. bor. occ.

CHAETOCAPNIA, Sweet, Hort. Brit. ed. III. 685 (1839) = CORTOCAPNIA, Link & Otto = **Bravoa**, Lex. (Amaryllid.).
geminiflora, Sweet, l. c. = B. geminiflora.

CHAETOCARPUS, Schreb. Gen. 75 (1789) = **Pouteria**, Aubl. (Sapotac.).
Ponteria, J. F. Gmel. Syst. 248 = P. guianensis.

CHAETOCARPUS, Thw. in Hook. Kew Journ. vi. (1854) 200. t. 10. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 232.
RIGNALDIA, Baill. Adansonia, i. (1860–61) 187.
Blanchetii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 507.—Bras.
castanocarpus, *Thw. Enum. Pl. Zeyl.* 275.—Ind. or.; Malaya.
coriaceus, *Thw. l. c.*—Zeylan.
myrsinites, *Baill. Adansonia*, xi. (1873) 95.—Bras.
Pohlii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 508.—Bras.
pungens, Thw. in Hook. Kew Journ. vi. (1854) 201 = castanocarpus, coriaceus.

CHAETOCEPHALA, Barb. Rodr. Orch. Nov. ii. 37 (1881) = **Pleurothallis**, R. Br. (Orchid.).
lonchophylla, Barb. Rodr. l. c. 39.—Bras.
pumila, Barb. Rodr. l. c. 38.—Bras.

CHAETOCHILUS, Vahl Enum. i. 101 (1805) = **Schwenkia**, Linn. (Solanac.).
laterifloru, Vahl, l. c. = S. brasiliensis.

CHAETOCLADUS, Semlin [Nelson]. Pinaceae, 161 (1868) = **Ephedra**, Linn. (Gnetac.).
altissima, Semlin [Nelson], l. c. 162 = E. altissima.
distachya, Semlin [Nelson], l. c. 161 = E. vulgaris.
fragilis, Carr. Conif. ed. II. 360, Index = E. fragilis.
monostachya, Semlin [Nelson]. Pinac. 162 = E. vulgaris.

CHAETOCRATER, Rafn & Paw. Prod. Fl. Por. 61. t. 36 (1799) = **Casearia**, Jacq. (Samydac.).
capitatum, Rafn & Paw. Fl. Por. iv. t. 396; Syst. 10a = Casearia sylvestris.
fasciculatum, Rafn & Paw. ll. cc. 396, 107 = Casearia punctata.

CHAETOCYPERUS, Nees, in Linnaea, ix. (1834) 289 = **Bisocharis**, R. Br. (Cyperac.).
acicularis, Nees, in Mart. Fl. Bras. ii. i. 94 = E. acicularis.
albibracteatus, Nees & Meyen, in Act. Nov. Nat. Cur. xix. Suppl. i. (1843) 85 = E. albibracteata.
arenicola, Steud. Syn. Pl. Cyp. 73 = E. Chaetaria.
Baldwinii, Torr. in Ann. Lyc. N. York, iii. (1836) 295 = E. Baldwinii.
bonariensis, Nees, in Mart. Fl. Bras. ii. i. 96 = E. bonariensis.
capillaceus, Nees, l. c. 93 = E. capillacea.
costulatus, Nees & Meyen, in Act. Nov. Nat. Cur. xix. Suppl. i. (1843) 86 = E. acicularis.
depauperatus, Steud. Syn. Pl. Cyp. 73 = E. Chaetaria.
emarginatus, Nees, in Mart. Fl. Bras. ii. i. 96.—Bras.
Yamesoni, Steud. Syn. Pl. Cyp. 74.—Ecuador.
Linnacharis, Nees, in Wight, Contrib. 96 = E. Chaetaria.
membranaceus, Buckl. in Proc. Acad. Sc. Philad. '1862' (1863) 10 = Scirpus nanus.
nivens, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 243 = E. Chaetaria.
obtusatus, Nees, in Mart. Fl. Bras. ii. i. 94 = E. Chaetaria.
polymorphus, Nees, in Linnaea, ix. (1834) 289 = E. acicularis, ochreata, Scirpus cernuus, nanus.
punctatus, Nees, in Mart. Fl. Bras. ii. i. 93.—Bras.
pusillus, Steud. Syn. Pl. Cyp. 74 = Scirpus nanus.
Ryzmaeus, Torr. ex A. Gray, Man. Bot. N. U. St. 524 = Scirpus nanus.
radicans, Steud. Syn. Pl. Cyp. 74 = E. acicularis.
reptans, Boeck. in Flora, xl. (1857) 84 = E. minima?
rugulosus, Nees, in Bonpiandia, iii. (1855) 86 = E. Chaetaria.
setaceus, Nees, in Mart. Fl. Bras. ii. i. 94 = E. Chaetaria.
stoloniferus, Nees, in Linnaea, xix. (1847) 695 = E. inostana.
subarticulatus, Nees, in Mart. Fl. Bras. ii. i. 96.—Bras.
tenuiculus, Necs, l. c. = E. Rothiana.
trichoides, Steud. Syn. Pl. Cyp. 73 = E. acicularis.
urceolatus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 243 = E. minima.
viviparus, Nees, in Mart. Fl. Bras. ii. i. 93 = E. tenuissima.

CHAETODISCUS, Steud. Syn. Pl. Cyp. 261 (1855) = **Eriosculon**, Linn.
Gilberti, Steud. l. c.—Austral.

CHAETOGASTRA, DC. Prod. iii. 181 (1828); Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 127 = Tibouchina, Aubl. (Melast.).
alpestris, Mart. Nov. Gen. et Sp. iii. 101 = Pterolepis alpestris.
asperior, Cham. in Linnaea, ix. (1834) 485 = T. asperior.
bicolor, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 128 (= *Pleroma bicolor*).—Boliv.
Bonplandiana, Naud. l. c. 127 = Brachyotum lediifolium.
Callichaeta, Benth. in Hook. Journ. Bot. ii. (1840) 291 = Pterolepis glomerata.
campanularis, Benth. Fl. Hartw. 181 = Brachyotum Benthamianum.
campanularis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 130 = Brachyotum campanulare.
canescens, DC. Prod. iii. 134 = Brachyotum canescens.
cardinalis, DC. l. c. = T. cardinalis.
cataphracta, Cham. in Linnaea, ix. (1834) 451 = Pterolepis cataphracta.
cervina, DC. Prod. iii. 136 = Brachyotum cervinum.
Chamaecistus, Griseb. Fl. Brit. W. Ind. 267 (= *Pleroma Chamaecistus*).—Ind. occ.
chrysolaena, Griseb. l. c. (= *Pleroma strigosum*, *chrysolaenum*).—Ind. occ.
ciliaris, DC. Prod. iii. 133 (= *Pleroma ciliare*).—N. Granat.
cistoides, Griseb. Fl. Brit. W. Ind. 267 (= *Pleroma cistoides*).—Ins. S. Vincent.
clinopodifolia, DC. Prod. iii. 138 = T. clinopodifolia.
confertu, DC. l. c. 135 = Brachyotum confertum.
eubensis, Griseb. Cat. Pl. Cub. 109 = Chaetolepis microcladon.

CHAETOGASTRA :—
debilis, Cham. in Linnæa, ix. (1834) 499 = Tibouchina debilis.
debilis, Miq. in Linnæa, xxii. (1849) 540 = T. ursina.
despeserata, Nand. in Ann. Sc. Nat. Sér. III. xiv. (1850) 139 = T. longifolia.
divaricata, DC. Prod. iii. 132 = Pterogastra divaricata.
echinata, DC. Prod. iii. 135 (= *Pterona echinatum*). —Pervv.
ferruginea, Hook. & Arn. Bot. Beech. Voy. 423 = T. longifolia.
floribunda, Griseb. in Lechl. Berb. Am. Austr. 59 = Brachyotum floribundum.
fraterna, DC. Prod. iii. 133 = T. gracilis.
Gardneriana, Schlecht. in Linnæa, xxvii. (1854) 528.— N. Granat.
glomerata, Benth. in Hook. Journ. Bot. ii. (1840) 290 = Pterolepis glomerata.
Gaudichii, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 131 = Brachyotum strigosum.
gracilis, DC. Prod. iii. 133 = T. gracilis.
hawaiensis, DC. l. c. 129 = T. longifolia.
hermanioides, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 153 = Brachyotum hermanioides.
hieracioides, DC. Prod. iii. 132 = T. hieracioides.
hirsuta, DC. l. c. = T. gracilis.
hispida, DC. l. c. 132 = T. longifolia.
hypericoides, DC. l. c. = Desmocelis villosa.
ladanoides, Mart. Herb. Fl. Bras. n. 500 = Pterolepis trichotoma.
lanceolata, DC. Prod. iii. 131 = T. longifolia.
laniphylla, Benth. in Hook. Journ. Bot. ii. (1840) 291 = T. lasiophylla.
lepidota, DC. Prod. iii. 134 = T. lepidota.
Lhotskyana, Steud. in Flora, xxvii. (1844) 720 = T. Lhotskyana.
Lindeniana, Planch. in Fl. des Serres, Sér. I. (1854–55) 118. t. 1011–13 = T. grossa.
longifolia, DC. Prod. iii. 132 = T. longifolia.
lutescens, Griseb. ex Triana, in Trans. Linn. Soc. xxvii. (1871) 48 = Brachyotum lutescens.
lutescens, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 134 = Brachyotum Naudini.
lychnitoides, DC. Prod. iii. 132 = Desmocelis villosa.
Martiana, Benth. in Hook. Journ. Bot. ii. (1840) 290 = T. herbacea.
microdon, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 134 = Brachyotum microdon.
microphylla, Naud. l. c. 136 = Brachyotum microphyllum.
mollis, DC. Prod. iii. 134 (= *Pterona exappendiculatum*).—Pervv.
Montiana, Klotzsch, ex Triana, in Trans. Linn. Soc. xxvii. (1871) 46 (= *Pterona Naudinianum*).—N. Granat.
multiflora, Benth. Pl. Hartw. 181 = Monochaetum Hartwegianum.
muricata, DC. Prod. iii. 134 = T. grossa.
Naudiniana, Decne. in Rev. Hortic. Sér. III. i. (1847) 86 (= *Pterona Naudinianum*).—Mexic.
nigrilla, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 137 = Bucquetia nigrilla.
origanoides, Griseb. Cat. Pl. Cub. 108 = Chaetolepis cubensis.
paniculariis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 139 (= *Pterona paniculare*).—Bolivia.
Pentlandii, Naud. l. c. 133 = Brachyotum Pentlandi.
quinquenervis, Naud. l. c. 130 = Brachyotum quinquenerve.
repanda, Mart. Nov. Gen. et Sp. iii. 100. t. 246 = Pterolepis repanda.
reticulata, DC. Prod. iii. 131 = T. reticulata.
rhynchanthera, Benth. Pl. Hartw. 180 = Monochaetum Bonplandii.
rosea, Regel, Cat. Pl. Hort. Alexhov. 31.—Hab. ?
rosmarinifolia, Benth. Pl. Hartw. 131 = Miconia ledifolia.
rosmarinifolia, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 131 = Brachyotum rosmarinifolium.
rostrata, Naud. l. c. 135 = Brachyotum rostratum.
rufifolia, Schlecht. in Linnæa, xiii. (1839) 490 (= *Pterona rufifolia*).—Mexic.
sanguinolenta, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 131 = Brachyotum sanguinolentum.
saturnioides, DC. Prod. iii. 134 (= *Pterona lasium*).— Pervv.
saturnioides, Griseb. Cat. Pl. Cub. 108 = Chaetolepis saturnioides.

CHAETOGASTRA :—
scabriuscula, Schlecht. in Linnæa, xiii. (1839) 481 (= *Pterona scabriusculum*).—Mexic.
Schiedeana, Cham. & Schlecht. in Linnæa, v. (1830) 565 (= *Pterona ruffifolia*).—Mexic.
sherardioides, DC. Prod. iii. 132 = Pterolepis glomerata.
speciosa, DC. l. c. 131 = Mariania speciosa.
stachyoides, DC. l. c. 132 = Desmocelis villosa.
stricta, Benth. Pl. Hartw. 180 = Brachyotum cernuum.
stricta, DC. Prod. iii. 134 = Brachyotum strigosum.
strigillosa, DC. l. c. 133 = T. gracilis.
strigosa, DC. l. c. iii. 134 (= *Pterona strigosum*).— Am. sept.
strigosa, Mart. Nov. Gen. et Sp. iii. 101. t. 246 = Pterolepis striphnocalyx.
sulphurea, Naud. in Ann. Sc. Nat. Ser. III. xiv. (1850) 135 = Brachyotum latifolium.
tortuosa, DC. Prod. iii. 133 (= *Pterona mexicanum*). —Mexic.
Weddelliana, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 139 (= *Pterona paniculare*).—Bolivia.

CHAETOLEPIS, Miq. Comm. Phyt. ii. 73 (1838). *MELASTOMACEÆ*, Benth. & Hook. f. i. 744.
HAPLODESMIUM, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 154. t. 5.
TRIMERANTHUS, Karst. in Linnæa, xxx. (1859–60) 159.
alpestris, *Triana*, in *Trans. Linn. Soc.* xxvIII. (1871) 51.—Venezuela.
alpina, *Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 140.—Bras.
anisandra, *Naud. l. c.*—Bras.
cubensis, *Triana*, in *Trans. Linn. Soc.* xxvIII. (1871) 51.—Cuba.
ericoides, Triana, l. c. 51. t. 3. f. 35 *g* = Marcetia ericoides.
heterophylla, *Triana*, l. c. 51.—Uruguay.
Lindeniana, *Triana*, l. c.—N. Granat.
loricarella, *Triana*, l. c.—N. Granat.
microphylla, *Miq. Comm. Phyt.* 73.—N. Granat.
saturesioides, *Triana*, in *Trans. Linn. Soc.* xxvIII. (1871) 51.—Cuba.
thymifolia, *Triana*, l. c. 50.—N. Granat.

CHAETONYCHIA, Sweet, Hort. Brit. ed. III. 263 (1839) = Paronychia, Juss.
cymosa, Sweet, l. c. = P. cymosa.

CHAETOPAPPA, DC. Prod. v. 501 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 265.
CHAETANTHERA, Nutt. in Journ. Acad. Sc. Philad. vii. (1834) 111.
CHAETOPHORA, Nutt. ex DC. Prod. v. 501, in syn. (1836).
DIPLOSTELMA, Rafin. ex Torr. & Gray, Fl. N. Am. ii. 187 (1840).
DIPTERIS, DC. Prod. v. 279 (1836).
asteroides, *DC. l. c.* 501.—Am. bor.
modesta, *A. Gray*, in *Proc. Am. Acad.* xvi. (1881) 89. —Mexic.
Parryi, *A. Gray*, l. c.—Texas.

CHAETOPHORA, Nutt. ex DC. Prod. v. 501, in syn. (1836) = Chaetopappa, DC. (Compos.).
asteroides, Nutt. ex DC. l. c. = Chaetopappa asteroides.

CHAETOPTELEA, Liebm. in Kjoeh. Vidensk. Meddel. (1850) 76 = Ulmus, Linn. (Urticac.).
mexicana, Liebm. l. c. 77 = U. mexicana.

CHAETOSCIADIUM, Boiss. Fl. Orient. ii. 1078 (1872) = TORILIS, DC. = Cancalis, Linn. (Umbellif.).
trichospermum, Boiss. l. c. = Canc. trichosperma.

CHAETOSPORA, H. B. & K. Nov. Gen. et Sp. i. 229 (1815) = Rynchospora, Vahl (Cyperac.).
aurea, Cham. & Schlecht. in Linnæa, vi. (1831) 26 = R. aristata.
aurea, H. B. & K. Nov. Gen. et Sp. i. 231 = R. aurea.
capitata, H. B. & K. l. c. 229 = R. capitata.
ferruginea, Cham. & Schlecht. in Linnæa, vi. (1831) 25, partim = R. velutella.
ferruginea, H. B. & K. Nov. Gen. et Sp. i. 25 = R. glauca.
pterocarpa, H. B. & K. Nov. Gen. et Sp. i. 229 = R. pterocarpa.

CHAETOSPORA :—
tricuspa, Cham. & Schlecht. in Linnæa, vi. (1831) 29 = Rynchospora tricuspa.
wiesenaia, Schrad. ex Roem. & Schult. Syst. ii. Mant. 46 = R. glauca.

CHAETOSPORA, R. Br. Prod. 232 (1810) = Schoenus, Linn. (Cyperac.).
albescens, Franch. & Sav. Enum. Pl. Jap. ii. 548 = S. apogon.
alpina, F. Muell. Fragm. ix. 39 = Carpha alpina.
ancepa, R. Br. Prod. 233 = Mesomelaena ancepa.
antarctica, Hook. f. Fl. Antarct. ii. 361. t. 147.— Chili.
arundinacea, A. Dietr. Sp. Pl. ii. 81 = Asterochaete arundinacea.
aurata, Nees, in Ann. Nat. Hist. Ser. I. vi. (1841) 49 = S. curvifolius.
avenacea, F. Muell. Fragm. ix. 40 = Cyathochaeta avenacea.
axillaris, R. Br. Prod. 233 = S. axillaris.
brevisetis, R. Br. l. c. 232 = S. brevisetis.
brevisetis, F. Muell. Fragm. ix. 37 = S. capitatus.
Burmanni, Schrad. Anal. Cap. Cyp. 32. t. 3. f. 4.—Afr. austr.
calostachys, R. Br. Prod. 233 = S. calostachyus.
capillacea, Hook. f. Fl. Tasm. ii. 81. t. 141 A = Elynanthus capillaceus.
capillacea, Nees, in Linnæa, ix. (1834) 299 ; x. (1836) 192 = S. capillaceus.
capillaris, F. Muell. Fragm. ix. 34 = Elynanthus capillaceus.
Capitellum, A. Dietr. Sp. Pl. ii. 27 = Ficinia setiformis.
ciliata, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 128 = Rynchospora elatior.
circinalis, Schrad. in Goett. Gel. Anz. (1821) 111. 1070 = S. circinalis.
clandestina, F. Muell. Fragm. ix. 40 = Cyathochaeta clandestina.
compressa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 71 ; Nees, in Lehm. Pl. Preiss. ii. 85 = Scirpus Caricis.
concava, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 47 = Lepidosperma concavum.
concinna, Hook. f. Fl. N. Zeal. i. 274. t. 62 B = S. concinnus.
cruenta, Nees, in Lehm. Pl. Preiss. ii. 85 = S. cruentus.
curvifolia, R. Br. Prod. 232 = S. curvifolius.
cuspidata, Nees, in Linnæa, vii. (1832) 529 = Elynanthus cuspidatus.
cygnea, Nees, in Ann. Nat. Hist. Ser. I. vi. (1841) 49 = S. brevisetis.
dactyloides, A. Dietr. Sp. Pl. ii. 82 = Asterochaete glomerata.
deformis, R. Br. Prod. 232 = S. deformis.
densata, F. Muell. Fragm. ix. 39 = Mesomelaena densata.
diandra, F. Muell. l. c. = Cyathochaeta diandra.
diodon, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 148 = Rynchospora innata.
dispar, A. Dietr. Sp. Pl. ii. 82 = Ficinia secunda.
distachya, Nees, in Linnæa, x. (1836) 292.—Afr. austr.
distans, F. Muell. Fragm. ix. 35 = S. calostachyus.
effusa, Schrad. ex Kunth, Enum. Pl. ii. 326 = Pleurostachys spaniflora.
elongata, Nees, in Lehm. Pl. Preiss. ii. 275 = Mesomelaena nesea.
ferruginea, Retchb. Fl. Germ. Excurs. 74 = S. ferrugineus.
fimbriolata, Nees, in Linnæa, x. (1836) 192.—Afr. austr.
fimbrisyloides, F. Muell. Fragm. ix. 34 = Cladium undulatum.
flexuosa, Schrad. Anal. Cap. Cyp. 33. t. 3. f. 3 = S. flexuosus.
globosa, H. B. & K. Nov. Gen. et Sp. i. 235 = Rynchospora globosa.
hexandra, Boeck. in Flora, lxi. (1878) 37.—Afr. austr.
imbricata, R. Br. Prod. 233 = S. apogon.
japonica, Franch. & Sav. Enum. Pl. Jap. ii. 548 = S. apogon.
laxa, Kunze, ex Steud. Nom. ed. II. i. 841 = Pimbristylis aspera.
lanata, R. Br. Prod. 233 = S. lanatus.
lana, Hook. f. Fl. Antarct. ii. 361. t. 146 = Carpha paniculata.
laxeva, Poir. in Dict. Sc. Nat. viii. 69 = S. glaucus.
madagascariensis, Steud. Syn. Pl. Cyp. 126.— Madag.

CHAETOSPORA :—
Mauritii, Steud. Syn. Pl. Cyp. 161 = Rynchospora glauca.
microstachya, Nees, in Lehm. Pl. Preiss. ii. 84 = Schoenus Drummondii.
microstachya, Schrad. Anal. Fl. Cap. i. 38, in obs. = S. circinalis.
nana, Nees, in Lehm. Pl. Preiss. ii. 85 = S. nanus.
natans, F. Muell. Fragm. ix. 86 = S. natans.
Neesii, Boeck. in Linnæa, xxxviii. (1874) 297 = Tricostularia Neesii.
nigricans, Kunth, Enum. Pl. ii. 328 = S. nigricans.
nitens, R. Br. Prod. 238 = S. nitens.
oligostachya, Boeck. in Flora, xlii. (1859) 101 = S. nigricans.
oligostachya, F. Muell. Fragm. ix. 88 = S. bifidus.
paludosa, R. Br. Prod. 238 = Tricostularia paludosa.
paniculata, Steud. Syn. Pl. Cyp. 161.—Ins. Borbon.
pedicellata, R. Br. Prod. 232 = S. pedicellatus.
perpusilla, H. B. & K. Nov. Gen. et Sp. i. 230 = Haplostylis barbata.
punctoria, A. Dietr. Sp. Pl. ii. 28 = Bucekia punctoria.
rhynchospermoides, Phil. f. Pl. Vasc. Chil. 306 = seq.
rhynchosporoides, Steud. Syn. Pl. Cyp. 162.—Chili.
robusta, Kunth, Enum. Pl. ii. 325.—Afr. austr.
rufa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 72 = Scirpus rufus.
sphaerocephala, R. Br. Prod. 238 = Mesomelaena sphaerocephala.
spicata, Boeck. in Linnæa, xxxviii. (1874) 297 = Tricostularia compressa.
striata, A. Dietr. Sp. Pl. ii. 27 = Ficinia setiformis.
stygia, R. Br. Prod. 238 = Mesomelaena stygia.
tenax, Hook. f. Fl. N. Zel. i. 273 = S. brevifolius.
Tendo, Hook. f. l. c. = S. Tendo.
tenella, Rupr. Sert. Tianach. 34.—Turkestan.
tenera, A. Dietr. Sp. Pl. ii. 32 = S. tener.
tenuissima, Hook. f. Fl. Tasm. ii. 81. t. 14 = Lepidospora tenuissima.
tenuissima, Steud. Syn. Pl. Cyp. 319, 162 = S. apogon.
tetragona, R. Br. Prod. 238 = Mesomelaena tetragona.
turbinata, R. Br. l. c. 232 = S. turbinatus.
umbellulifera, Boeck. in Flora, lxv. (1882) 28 = S. apogon.
villosa, Nees, ex Steud. Nom. ed. II. i. 341 = S. villosus.

CHAETOSTACHYS, Benth. in Wall. Pl. As. Rar. ii. 19 (1831) = **Lavandula**, Tourn. (Labiat.).
mulithda, Benth. l. c. = L. Burnsanni.

CHAETOSTEMMA, Reichb. Consp. 176 (1828) = seq.

CHAETOSTOMA, DC. Prod. iii. 112 (1828). *MELASTOMACEÆ*, Benth. & Hook. f. i. 787.
acuminatum, *Cogn. in Mart. Fl. Bras.* xiv. iii. 85.—Bras.
ornatum, Cogn. l. c. 81 = diosmoides.
castratum, *Cogn. l. c.* 86.—Bras.
diosmoides, *Mart. Nov. Gen. et Sp.* iii. 180. t. 264.—Bras.
ericoides, DC. Prod. iii. 112 = Marcetia ericoides.
fastigiatum, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 191.—Bras.
Gardneri, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 25. t. 1. f. 8 c.—Bras.
Glaziovii, *Cogn. in Mart. Fl. Bras.* xiv. iii. 80.—Bras.
inerme, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 191.—Bras.
longiflorum, *Cogn. in Mart. Fl. Bras.* xiv. iii. 92.—Bras.
microfoliolatum, Cham. in Linnæa, ix. (1834) 388 = Microlicia Chamissois.
oxyanthemum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 25, pro parte.—Bras.
oxyanthemum, Triana, l. c. pro parte = acuminatum, castratum.
pungens, *DC. Prod.* iii. 113.—Bras.
Riedelianum, *Cogn. in Mart. Fl. Bras.* xiv. iii. 83.—Bras.
tetrastichum, DC. Prod. iii. 113 = Marcetia ericoides, tetrasticha.

CHAETOSUS, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 266. *APOCYNACEÆ*, Benth. & Hook. f. ii. 666.
CHAETOSUS, Benth. in Journ. Linn. Soc. i. (1857) 87.
volubilis, *Benth. l. c.*—N. Guinea.

CHAETOTHYLAX, Nees, in Mart. Fl. Bras. ix. 158. t. 96 (1847). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1107.
CHAETOTHYLOPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 168.
HEINSELIA, Nees, in Mart. Fl. Bras. ix. 159 (1847).
lythroides, *Benth. & Hook. f. Gen.* ii. 1106.—Am. austr.
microanthus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 515.—Nicaragua.
phyllostachyus, *Nees, in DC. Prod.* xi. 813.—Mexic.
tocantinus, *Nees, in Mart. Fl. Bras.* ix. 158.—Bras.
umbrosus, *Nees, in DC. Prod.* xi. 813.—N. Granat.

CHAETOTHYLOPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 168 = **Chaetothylax**, Nees (Acanthac.).
microantha, Oerst. l. c. = Chaetothylax miranthus.

CHAETOTROPIS, Kunth, Rev. Gram. i. 72, 271. t. 47 (1830). *GRAMINEÆ*, Benth. & Hook. f. iii. 1151.
andina, *Ball, in Journ. Linn. Soc.* xxii. (1885) 58.—Peruv.
chilensis, *Kunth, Rev. Gram.* i. 72, 271. t. 47.—Chili.
latifolia, Phil. in Linnæa, xxx. (1859–60) 205 = chilensis.

CHAETOTROPSIS, D. Dietr. Syn. Pl. i. 184 (1839) = præc.

CHAETURUS, Link, in Schrad. Journ. ii (1799) 313. *GRAMINEÆ*, Benth. & Hook. f. iii. 1140.
divaricatus, DC. Cat. Hort. Monsp. 93 = fasciculatus.
fasciculatus, *Link, in Schrad. Journ.* ii. (1799) 313.—Hispan.
prostratus, *Hack. & Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 223.—Lusitan.
spicatus, Schrad. Ind. Sem. Hort. Gotting. ex Steud. Nom. ed. II. i. 342 = fasciculatus.

CHAILLETIA, DC. in Ann. Mus. Par. xvii. (1811) 158. *CHAILLETIACEÆ*, Benth. & Hook. f. i. 341.
DICHAPETALUM, Thou. Gen. Madag. 28 (1806).
LECOSTA, Thou. l. c. (1806).
MESTOTES, Soland. ex Steud. Nom. ed. II. i. 342 (1842).
MORCIERA, Roxb. Hort. Beng. 21 (1814); Fl. Ind. ii. 69 (1832).
PATRISIA, Rohr, ex Steud. Nom. ed. II. i. 342 (1840).
PLAPPERTIA, Reichb. Consp. 146 (1828).
QUILESIA, Blanco, Fl. Filip. ed. I. 176 (1837).
SYMPHYLLANTHUS, Vahl, in Skriv. Nat. Hist. Selsk. vi. 66 (1810).
WAHLENBERGIA, R. Br. in Wall. Cat. p. 950, sub n. 4849 (1831).
affinis, Planch. ex Benth. in Hook. Niger Fl. 276 = toxicaria.
Bangii, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1854) 195.—Afr. trop.
Benthamiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 610.—Ins. Philipp.
Benthamii, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1854) 196.—Afr. trop.
Bojeri, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 86.—Malacca.
brunoniana, Wall. Cat. n. 6088 = Helferiana.
chlorina, *Tul. in Ann. Sc. Nat.* Ser. IV. viii. (1857) 88.—Madag.
cubensis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 41. t. 246. f. 3.—Cuba.
cymosa, *Hook. Ic. Pl.* t. 591.—Afr. austr.
cymulosa, *Oliver, Fl. Trop. Afr.* i. 340.—Afr. trop.
deflexa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 109.—Afr. trop.
deflexifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 611.—Malaya.
Dichapetalum, *DC. Prod.* ii. 57.—Madag.
discolor, *Baker, in Journ. Linn. Soc.* xx. (1883) 119.—Madag.
edulis, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 206 = Sarcosigna edula.
erecta, G. Don, in Edinb. Phil. Journ. xi. (1824) 349 = toxicaria.
fasciculata, *Sprang. Syst.* i. 981.—Madag.
flavicans, *Klotzsch, in Schweinf. Faunæ & Fl. Gui.* 1184, nomen.—Guinea.

CHAILLETIA :—
flexuosa, *Oliver, Fl. Trop. Afr.* i. 340.—Afr. trop.
floribunda, *Planch. in Hook. Ic. Pl.* t. 792.—Afr. trop.
gelonioides, *Hook. f. Fl. Brit. Ind.* i. 670.—Ind. or.; Malaya.
Griffithii, *Hook. f. l. c.* 671.—Malacca.
hainesensis, *Hance, in Journ. Bot.* xxiii. (1885) 323.—China.
Helferiana, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 297.—Burma.
Heudelotii, *Planch. ex Oliver, Fl. Trop. Afr.* i. 344.—Afr. trop.
hirtella, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 88.—Ins. Mascar.
hispida, *Oliver, Fl. Trop. Afr.* i. 340.—Afr. trop.
lanuginosa, Miq. ex Hook. f. Fl. Brit. Ind. i. 672 = Griffithii.
Lauro-cerasus, *Planch. ex Hook. f. l. c.*—Ins. Penang.
Leucosia, *Sprang. Syst.* i. 981.—Madag.
longipetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 611.—Burma.
macrophylla, *Oliver, Fl. Trop. Afr.* i. 343.—Afr. trop.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb Bot.* 108.—Afr. trop.
nigrescens, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 86.—Madag.
oblonga, *Hook. f. ex Benth. in Hook. Niger Fl.* 277.—Afr. trop.
Pachypus, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 87.—Madag.
pallida, *Oliver, Fl. Trop. Afr.* i. 343.—Afr. trop.
paniculata, *Benth. in Hook. Niger Fl.* 279.—Afr. trop.
papuana, *Becc. Malesia*, i. (1877) 176.—N. Guinea.
Parkerii, *Planch. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 379.—Venezuela.
pedunculata, *DC. in Ann. Mus. Par.* xvii. (1811) 154.—Guiana.
reflexa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 109. t. 20.—Afr. trop.
rufa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 86.—Madag.
rufipilla, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 611.—Afr. trop.
sessilifora, DC. in Ann. Mus. Par. xvii. (1811) 154 = Tapura guianensis.
sebaurioides, *Oliver, Fl. Trop. Afr.* i. 344.—Afr. trop.
subcordata, *Hook. f. ex Benth. in Hook. Niger Fl.* 277.—Afr. trop.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 328 = gelonioides.
Thomsoni, *Oliver, Fl. Trop. Afr.* i. 342.—Afr. trop.
timoriensis, *DC. Prod.* ii. 57.—Ins. Timor.
toxicaria, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 348.—Afr. trop.
vestita, *Benth. in Hook. Kew Journ.* iii. (1851) 372.—Bras.
Virchowii, *O. Hoffm. & Hildeb. ex O. Hoffm. Sert. Madag.* 11.—Madag.
vitiensis, *Seem. Viti*, 484, nomen; *Fl. Vit.* 88.—Ins. Fiji.

CHAITAEA, Soland. ex Seem. Fl. Vit. 102 (1865) = **Tacca**, Forst.
Tacca, Soland. ex Seem. l. c. = T. pinnatifida.

CHAITURUS, Willd. Enum. Pl. Berl. 900 (1787) = **Leonurus**, Linn. (Labiat.).
leonuroides, Willd. l. c. 901 = L. Marrubiastrum.
Marrubiastrum, Spenn. in T. Nees, Gen. Fl. Germ. fasc. 18. t. 353 = L. Marrubiastrum.

CHAIXIA, Lapeyr. Hist. Pl. Pyr. Suppl. 38 (1813) = **Ramondia**, Rich. (Gesneriac.).
Myconi, Lapeyr. l. c. 39 = R. pyrenaica.

CHAKIATELLA, DC. Prod. v. 569 (1836) = CHATIAKELLA, Cass. = **Wulffia**, Neck. (Compos.).

CHALARIUM, Pelt. ex DC. Prod. v. 546, in syn. (1836) = OGIERA, Cass. = **Blumtheranthera**, Pelt. (Compos.).

CHALCANTHUS, Boiss. Fl. Orient. i. 811 (1867) = **Hesperis**, Linn. (Crucifer.).
remifolius, Boiss. l. c. 212 = H. remifolius.

CHALCAS, Linn. Mant. 66 (1767) = **Murraya**, Koen. (Rutac.).
Commuxang, Rumph. f. Fl. Ind. 104 = M. exotica.
crenulata, F. Muell. Phyt. N. Hebr. 7 = M. crenulata.
intermedia, Roem. Syn. Hesper. i. 48 = M. exotica.
japonensis, Lour. Fl. Cochinch. 371 = M. exotica.
Koenigii, Kurz, in Journ. As. Soc. Beng. xliv. (1875) ii. 182 = M. Koenigii.
paniculata, Linn. Mant. i. 68 = M. exotica.
sumatrana, Roem. Syn. Hesper. i. 49 = M. exotica.

CHALEPOA, Hook. f. Ic. Pl. t. 1082 (1871) = **Triboles**, Phil. (Pittospor.).
magellanica, Hook. f. l. c. = T. australis †

CHALINOPHYLLUM, Hook. f. Ic. Pl. t. 1148 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 30.
guyanense, *Hook. f. l. c.*—Guiana.

CHALYBEA, Nasd. in Ann. Sc. Nat. Sér. III. xvi. (1851) 99 = **Pachyanthina**, A. Rich. (Melastomac.).
corymbifera, Naud. l. c. 100 = P. corymbosus.
corymbosa, Triana, in Trans. Linn. Soc. xxviii. (1871) 97 = P. corymbosus.

CHAMAEBAINIA, Wight, Ic. vi. 11. t. 1981 (1853). *URTICACEAE*, Benth. & Hook. f. iii. 387.
cuspidata, *Wight, l. c.*—Ind. or.
squamigera, Wall. Cat. n. 4592; Wedd. in DC. Prod. xvi. f. 218 = cuspidata.

CHAMAEBATIA, Benth. Pl. Hartw. 308 (1850). *ROSACEAE*, Benth. & Hook. f. i. 617.
foliolosa, Benth. l. c.—Calif.
foliolosa, Newberry, in Pacif. Rail. Rep. vi. 78 = Spiraea Millefolium.

CHAMAEBATIARIA, Maxim. in Act. Hort. Petrop. vi. (1879) 226 = **Spiraea**, Linn. (Rosac.).
Millefolium, Maxim. l. c. = S. Millefolium.

CHAMAEBETULA, Opiz, in Lotos, v. (1855) 258 = **Betula**, Tourn. (Cupulif.).
acutifolia, Opiz, l. c. 259.—Europ.
glandulosa, Opiz, l. c. = B. glandulosa.
Gmelinii, Opiz, l. c. = B. fruticosa.
Hookeri, Opiz, l. c. = B. glandulosa.
humilis, Opiz, l. c. = B. humilis.
nana, Opiz, l. c. = B. nana.
pumila, Opiz, l. c. = B. pumila.
rotundifolia, Opiz, l. c. = B. glandulosa.

CHAMAEBUXUS, [Tourn.] Spach, Hist. Vég. Phan. vii. 125 (1839) = **Polygala**, Linn.
alpestris, Spach, l. c. 127 = P. Chamaebuxus.
arillata, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 158 = P. arillata.
coriacea, Opiz, Seznam, 28 = P. Chamaebuxus.
harmsiana, Kurz, For. Fl. Brit. Burma, i. 79 = P. karensium.
paniculata, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 154 = P. arillata.
polygaloides, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 11 = P. Chamaebuxus.
pulchra, Hassk. Fl. Jav. Rar. 294, in textu = P. pulchra.
Vayredae, Willk. Illustr. Fl. Hisp. 33. t. 23 = P. Vayredae.
venenosa, Hassk. Fl. Jav. Rar. 294.—Java.
vulgaris, Schur, Enum. Pl. Transs. 90 = P. Chamaebuxus.

CHAMAECALAMUS, Meyen, Reise, i. 456 (1834) = **Dregearia**, Clar. (Graniin.).
spectabilis, Meyen, l. c.; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 156 (= *Bromidium spectabile*).—Peruv.

CHAMAECASSIA, [Breyn] Link, Handb. ii. 139 (1831) = **Cassia**, Tourn. (Leguniin.).
grandiflora, Link, l. c. = Cassia laevigata.
laevigata, Link, l. c. = Cassia laevigata.

CHAMAECERASUS, [Tourn.] Medic. Phil. Bot. i. 126 (1789) = **Lonicera**, Linn. (Caprifol.).
alpigena, Medic. l. c. = L. alpigena.
coerulea, Delarb. Fl. Auv. ed. II. 131 = L. coerulea.
dumetorum, Delarb. l. c. = L. Xylosteum.
fruticosa, Ser. in DC. Prod. ii. 587, sphalm. = Prunus Chamaecerasus.

CHAMAECERASUS:—
nigra, Medic. Phil. Bot. i. 126; Pl. Asiat. 290 = L. nigra.
Stenoblasti, Lescuyer, in L'Horticult. Franç. (1862) 138. t. 10.—Hab. ?
tatarica, Medic. Phil. Bot. i. 126; Pl. Asiat. 290 = Lonicera tatarica.
Xylosteum, Medic. Phil. Bot. i. 126 = L. Xylosteum.

CHAMAECISTUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 401 (1821) = **Lotus**, Leuria. Desv. (Ericac.).
scrpyllifolius, S. F. Gray, l. c. = L. procumbens.

CHAMAECISTUS, Regel, Gartenfl. (1874) 80 = **Rhododendron**, Linn. (Ericac.).
austriacus, Regel, l. c. = R. Chamaecistus.
kamtschaticus, Regel, l. c. = R. kamtschaticum.

CHAMAECLADON, Miq. in Bot. Zeit. xiv. (1856) 564. *AROIDEAE*, Benth. & Hook. f. iii. 988.
angustifolium, *Schott, in Bonplandia*, vi. (1858) 369. —Ins. Penang.
consobrinum, *Schott, in Ann. Mus. Bot. Lugd. Bat.* i. 196.—Sumatra.
Griffithii, *Schott. in Bonplandia*, vi. (1858) 369.—Malaya.
hamile, *Miq. Fl. Ind. Bat.* iii. 318.—Ins. Penang.
lanceolatum, *Miq. in Bot. Zeit.* xiv. (1856) 564.—Java.
metallicum, *N. E. Brown, in Illustr. Hortic.* xxxi. (1884) t. 539.—Borneo.
nigrescens, *Schott, in Ann. Mus. Bot. Lugd. Bat.* i. 196.—Sumatra.
obliquatum, *Schott, in Bonplandia*, vi. (1858) 369.—Ins. Penang.
ovalifolium, *Schott, in Ann. Mus. Bot. Lugd. Bat.* i. 981.—Borneo.
ovatum, *Schott, in Bonplandia*, vii. (1859) 30.—Malaya.
purpurascens, Schott, l. c. vi. (1858) 369 = lanceolatum.
pygmaeum, Engl. in DC. Monog. Phan. ii. 545 = lanceolatum.
rubescens, *Schott, Gen. Aroid.* sub t. 60, in obs.—Hab. ?
sanguinolentum, *Schott, Prod. Aroid.* 816.—Malaya.
saxorum, *Schott, in Ann. Mus. Bot. Lugd. Bat.* i. 980. —Borneo.
subcordatum, *Schott, l. c.* 981.—Borneo.
truncatum, *Schott, in Bonplandia*, vi. (1858) 369.—Burma.

CHAMAECLEMA, Moench, Meth. 393 (1794) = **Nepeta**, Tourn. (Labiat.).
hederacea, Moench, l. c. = N. Glechoma.

CHAMAECNIDE, Nees & Mart. ex Miq. in Mart. Fl. Bras. iv. t. 908 (1858) = **Pilea**, Lindl. (Urticac.).
microphylla, Nees, ex Miq. l. c. = P. muscosa.

CHAMAECRISTA, Moench, Meth. 273 (1794) = **Cassia**, Tourn. (Legumin.).
Burmanni, Eckl. & Zeyh. Enum. 260 = Cassia capensis.
capensis, E. Mey. in Linnaea, vii. (1832) 172 = Cassia capensis.
comosa, E. Mey. Comm. Pl. Afr. Austr. 160 = Cassia comosa.
nictitans, Moench, Meth. 273 = Cassia nictitans.
plumosa, E. Mey. Comm. Pl. Afr. Austr. 159 = Cassia mimosoides.
sericea, Mey. l. c. = Cassia mimosoides.

CHAMAECYPARIS, Spach, Hist. Vég. Phan. xi. (1849) 299 = **Thuya**, Tourn. (Conif.).
andelyensis, Hort. ex Gord. Pinet. ed. II. 365 = *lopiocladia*.
Beaurieri, Carr. Conif. ed. II. 198 = *Lawsoniana*.
Beaurieri, Decne. in Bull. Soc. Bot. Fr. i. (1854) 70 = Juniperus occidentalis.
breviramea, Maxim. in Bull. Acad. Pétersb. x. (1866) 489 = T. obtusa.
decussata, Hort. ex Carr. Conif. ed. II. 140 = T. orientalis.
ericoides, Carr. l. c. = T. orientalis, coniditensis.
exoniea, Finch. ex Carr. l. c. 187 = T. excelsa.
glauca, Hort. ex Carr. l. c. ed. I. 141 = T. orientalis.
Ketteleri, Scand. ex Parl. in DC. Prod. xvi. ii. 466 = T. obtusa.
leucosia, Hort. ex Carr. Conif. ed. I. 189 = T. sphaeroidalis.

CHAMAECYPARIS:—
Lawsoniana, Parl. in Ann. Mus. Stor. Nat. Fir. i. (1864) 181.—Am. bor. occ.
lopicladia, Hook. & Hochst. Nadelhouls. 257.—Japon.
nana, Hort. ex Parl. in DC. Prod. xvi. ii. 464 = Thuya sphaeroidalis.
nootkatensis, Carr. Conif. ed. II. 197 = seq.
nutkaensis, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 207 = T. excelsa.
nutkaensis, Spach, Hist. Vég. Phan. xi. 385 = praec.
obtusa, Sieb. & Zucc. in Endl. Conif. 68 = T. obtusa.
pendula, Maxim. in Bull. Acad. Pétersb. x. (1866) 489 = T. obtusa.
plisfera, Sieb. & Zucc. in Endl. Conif. 64 = T. plisfera.
pseudo-squarrosa, Parl. in DC. Prod. xvi. ii. 467 = T. sphaeroidalis.
pumila, Hort. ex Carr. Conif. ed. II. 194 = T. sphaeroidalis.
sphaeroidea, Spach, Hist. Vég. Phan. xi. 381 = T. sphaeroidalis.
squarrosa, Hort. ex Carr. Conif. ed. I. 65 = T. sphaeroidalis.
squarrosa, Sieb. & Zucc. in Endl. Conif. 65 = T. plisfera.
thujaeformis, R. Smith, ex Gord. Pinet. ed. II. 429.—Hab. ?
thuifera, Endl. Synop. Conif. 62 = Capressus thurifera.
variegata, Hort. ex Carr. Conif. ed. I. 133 = T. sphaeroidalis.

CHAMAECYTISUS, Link, Handb. ii. 154 (1831) **Cytisus**, Linn. (Legumin.).
austriacus, Link, l. c. 155 = Cytisus austriacus.
biflorus, Link, l. c. 154 = Cytisus biflorus.
capitatus, Link, l. c. 155 = Cytisus capitatus.
elongatus, Link, l. c. = Cytisus biflorus.
hirsutus, Link, l. c. = Cytisus hirsutus.
leucanthus, Link, l. c. 154 = Cytisus austriacus.
prolifernus, Link, l. c. = Cytisus prolifernus.
purpureus, Link, l. c. = Cytisus purpureus.
rupinus, Link, l. c. 155 = Cytisus biflorus.

CHAMAECYTISUS, Vis. Fl. Dalm. iii. 272 (1852) = **Argyrolobium**, Eckl. & Zeyh. (Legumin.).
dalmaticus, Vis. l. c. = A. Linnaeanum.

CHAMAEDACTYLIS, T. Nees, Gen. Fl. Germ. Monocot. i. n. 66 (1840?) = **Asturopsus**, Trin. (Gramin.).
maritima, Nees, l. c. = A. littoralis.

CHAMAEDAPHNE, Mitchell, in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) 222, App. = **Mitchellia**, Linn. (Rubiac.).

CHAMAEDAPHNE, Moench, Meth. 457 (1794) = **Cassandra**, D. Don (Ericac.).
calyculata, Moench, l. c. = Cass. calyculata.
crispa, Spach, Hist. Vég. Phan. ix. 477 = Cass. calyculata.

CHAMAEDOREA, Willd. Sp. Pl. iv. 688 (1805); et in Mém. Acad. Berl. 1804 (1807) 40. *PALMAE*, Benth. & Hook. f. iii. 910.
DAEVTSACHYS, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 25.
KUNTHIA, Humb. & Bonpl. Pl. Aequin. ii. 126. t. 122 (1809).
MORENIA, Ruiz & Pav. Prod. 150 (1794).
NUNNEMARIA, Ruiz & Pav. l. c. 147 (1794).
NUNNEZIA, Willd. Sp. Pl. iv. 1154 (1805).
SPATHOSCAPHE, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 29.
STACHYOPHORUS, Liebm. in Oversz. Dansk. Vidensk. Selsk. (1846) 5.
STEPHANOSTACHYS, Klotzsch, ex Oerst. in Kjoeb. Vidensk. Meddel. (1858) 26.
affinis, *Liebm. in Mart. Hist. Nat. Palm.* iii. 208.—Mexic.
alternans, *H. Wendl. in Regel, Gartenfl.* (1880) 104. —Mexic.
amazonica, *Y. Linden, in Illustr. Hortic.* xxviii. (1881) 16, nomen.—Peruv.
Andreana, *Y. Linden, l. c.*, nomen.—N. Granat.
Arenbergiana, *H. Wendl. Pal. Palm.* (1854) 98.—Guatemala!

CHAMAEDOREA:—
atrovirens, Hort. ex H, Wendl. in Kerch. Palm. 289 = Martiana.
atrovirens, *Mart. in Flora*, xxxv. (1859) 721.—Mexic.
aurantiaca, *A. Brongn. in Rev. Hortic. Sér.* III. i. (1847) 86.—Mexic.
Bartlingiana, *H. Wendl. Ind. Palm.* (1854) 60.—N. Granat.
bifida, H. Wendl. l. c. 11 = Bartlingiana.
bifurcata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 13.—Costa Rica.
brachycalyx, *H. Wendl. in Regel, Gartenfl.* (1880) 101. —Panama.
bractesta, *H. Wendl. in Bot. Zeit.* xvii. (1859) 29.—An. centr.
brevifrons, *H. Wendl. Ind. Palm.* (1854) 61.—N. Granat.
Casperiana, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xx. (1852) 368.—Guatemala.
cataractarum, *Mart. Hist. Nat. Palm.* iii. 309.—Mexic.
concolor, *Mart. l. c.* 160.—Mexic.
concorpa, *Mart. Palmet. Orb.* 6.—Bolivia.
costaricana, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 19.—Costa Rica.
Deckeriana, *Hensl. Biol. Centr. Am. Bot.* iii. 404.—Guatemala.
Deppeana, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 369 = elegans.
desmonocoides, *H. Wendl. in Otto & Dietr. l. c.* xxi. (1853) 177.—Mexic.
elatior, *Mart. in Linnaea*, v. (1830) 205.—Mexic.
elegans, *Mart. l. c.* 204.—Mexic.
elegantissima, *Hort. ex H. Wendl. in Kerch. Palm.* 289, no.sen.—Hab.!
Ernesti-Augusti, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 78.—Mexic.
frenestrata, Hort. ex H. Wendl. Ind. Palm. (1854) 28 = Malortiea gracilis.
fenestrata, *Y. Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Mexic.
fibrosa, H. Wendl. Ind. Palm. (1854) 57 = Synechanthus fibrosus.
flavo-virens, *H. Wendl. l. c.* 60.—Hab.!
flexuosa, Hort. ex H Wendl. l. c. 14 = Martiana.
formosa, *Hort.—Cf Gard. Chron.* (1876) i. 794.—Hab.!
fragrans, *Mart. Hist. Nat. Palm.* ii. 4. t. 3. f. 1, 2.—Peruv.
geonomaeformis, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 1.—Guatemala.
geonomoides, *Drude, in Mart. Fl. Bras.* iii. ii. 531.—Peruv.
Ghiesbreghtii, *Hort. ex H. Wendl. in Kerch. Palm.* 289, nomen.—Hab.!
glauca, *Y. Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Venezuela.
glaucifolia, *H. Wendl. Ind. Palm.* (1854) 64.—Mexic.
gracilis, *Wildd. Sp. Pl.* iv. 800.—Am. austr.
graminifolia, *H. Wendl. Ind. Palm.* (1854) 62.—Guatemala.
Helleriana, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 368 = elegans.
humilis, *Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
Karwinskyana, H. Wendl. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 179 = elatior.
Klotzschiana, *H. Wendl. Ind. Palm.* (1854) 58.—Mexic.
lanceolata, *Kunth, Enum. Pl.* iii. 172.—Peruv.
latifrons, H. Wendl. Ind. Palm. (1854) 11 = Arenbergiana.
lepidota, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 183.—Mexic.
Liebmanni, *Mart. Hist. Nat. Palm.* iii. 308 = Mexic.
Lindeniana, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 180.—N. Granat.
linearis, *Mart. Hist. Nat. Palm.* ii. 5.—Peruv.
lunata, Liebm. in Mart. l. c. iii. 307 = oblongata.
macrospadix, *Oerst. in Kjoeb. Vidensk Meddel.* (1858) 20.—Costa Rica.
Martiana, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 187.—Mexic.
membranacea, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 20.—Nicaragua.

CHAMAEDOREA:—
mexicana, Hort. Belg. ex Heynh. Nom. ii. 185 = Sartorii.
microphylla, *H. Wendl. in Bot. Zeit.* xvii. (1859) 102. —Panama.
montana, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308. —Mexic.
oblongata, *Mart. l. c.* 160.—Bras.
onovoides, *H. Wendl. in Kerch. Palm.* 289, nomen.—Hab.!
oreophila, *Mart. Hist. Nat. Palm.* iii 309.—Mexic.
Pazaya, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 12. —Costa Rica.
paradoxa, *H. Wendl. in Bot. Zeit.* xvii. (1859) 29.—Mexic.
pauciflora, *Mart. Hist. Nat. Palm.* ii. 5. t. 3. f. 3.—Bras.
pinnatifrons, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 14 = gracilis.
pochutlensis, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
pulchella, *Linden, Cat.* n. 117, p. 4.—Hab.!
pygmaea, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 217, 249.—N. Granat.
radicalis, *Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
regia, H. Wendl. Ind. Palm. (1854) 11 = desmoncoides.
repens, H. Wendl. l. c = desmoncoides.
resinifera, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 179.—Mexic.
robusta, *Hort. ex H. Wendl. in Kerch. Palm.* 289, nomen —Hab.!
Sartorii, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
scandens, Liebm. in Mart. l. c. = elatior.
Schiedeana, *Mart. in Linnaea*, v. (1830) 204.—Mexic.
simplicifrons, Hort. Belg. ex Heynh. Nom. ii. 185 = Ernesti-Augusti.
speciosa, H. Wendl. Ind. Palm. (1854) 15 = Schiedeana.
spectabilis, *Hort. ex H. Wendl. in Kerch. Palm.* 240, nomen.—Hab.!
tenella, *H. Wendl. in Regel, Gartenfl.* (1880) 102.—Mexic.
Tepejilote, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
velutina, Hort. ex H. Wendl. Ind. Palm. (1854) 13 = lepidota.
Verschaffeltii, *Hort. ex H. Wendl. in Kerch. Palm.* 240, nomen.—Hab.!
Wallisi, *Y. Linden, in Illustr. Hortic.* xxviii. (1881) 16, nomen.—Ecuador.
Warscewiczii, *H. Wendl. in Bonplandia*, x. (1862) 87. —Guatemala.
Wendlandi, *H. Wendl. in Kerch. Palm.* 240.—Mexic.
Wendlandiana, Hensl. Biol. Centr. Am. Bot. iii. 407 = pinea.
Wobstiana, *Linden, Cat.* n. 117, p. 4.—Hab.!

CHAMAEDRYS, Moench, Meth. 388 (1794) = Teucrium, Tourn. (Labiat.).
Botrys, Moench, l. c. = T. Chamaedrys.
capitata, Rafin. Fl. Tellur. iii. 85 = T. Polium.
cinerascens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 47 = T. flavum.
cretica, Rafin. Fl. Tellur. iii. 85 = T. creticum.
flava, Moench, Meth. 389 = T. flavum.
glauca, Jord. & Fourr. Brev. Fl. Nov. fasc. i. 48 = T. flavum.
lacinulata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 369 = T. Botrys.
latifolia, Rafin. Fl. Tellur. iii. 85 = T. fruticans.
lucida, Moench, Meth. 389 = T. lucidum.
Marum, Moench, l. c. 384 = T. Marum.
montana, Rafin. Fl. Tellur. iii. 85 = T. montanum.
Nissoliana, Rafin. l. c. = T. Pseudochamaepitys.
officinalis, Moench, Meth. 388 = T. Chamaedrys.
ovalifolia, Jord. & Fourr. Brev. Fl. Nov. fasc. i. 48 = T. flavum.
palustris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 369 = T. Scordium.
Polium, Rafin. Fl. Tellur. iii. 85 = T. Polium.
pulverulenta, Jord. & Fourr. Brev. Fl. Nov. fasc. i. 47 = T. flavum.
pumila, Rafin. Fl. Tellur. iii. 85 = T. pumilum.
quadrula, Rafin. l. c. = T. quadrastrum?
Scordium, Moench, Meth. 384 = T. Scordium.

CHAMAEDRYS:—
tetonicifolia, Rafin. Fl. Tellur. iii. 85 = Teucrium betonicum.
virescens, Jord. & Fourr. Brev. Fl. Nov. fasc. i. 47 = T. flavum.

CHAMAEFISTULA, G. Don, Gen. Syst. ii. 451 (1832) = **Cassia**, Tourn. (Legumin.).
astroites, G. Don, l. c. = Cass. villosa.
bacillaris, G. Don, l. c. = Cass. Sophera.
chinensis, G. Don, l. c. 462 = Cass. Sophera.
contorta, G. Don, l. c. = Cass. Tora.
coromandelina, G. Don, l. c. = Cass. Sophera.
corymbosa, G. Don, l. c. 451 = Cass. corymbosa.
crotalarioides, G. Don, l. c. 452 = Cass. crotalarioides.
elegans, G. Don, l. c. 451 = Cass. obliqua.
excelsa, G. Don, l. c. 452 = Cass. excelsa.
floribunda, G. Don, l. c. = Cass. laevigata.
gigantea, G. Don, l. c. = Cass. siamea.
Herbertiana, G. Don, l. c. = Cass. laevigata.
inaequilatera, G. Don, l. c. 451 = Cass. inaequilatera.
indecora, G. Don, l. c. 452 = Cass. bicapsularis.
inflata, G. Don, l. c. 451 = Cass. bicapsularis.
laevigata, G. Don, l. c. 452 = Cass. laevigata.
macrophylla, G. Don, l. c. 451 = Cass. macrophylla.
melanocarpa, G. Don, l. c. = Cass. viminea.
obliqua, G. Don, l. c. = Cass. obliqua.
oxyphylla, G. Don, l. c. = Cass. oxyphylla.
pendula, G. Don, l. c. 452 = Cass. bicapsularis.
prominens, G. Don, l. c. 451 = Cass. macranthera.
pubescula, G. Don, l. c. = Cass. bacillaris.
Rusiana, G. Don, l. c. = Cass. Rusiana.
Sophera, G. Don, l. c. 452 = Cass. Sophera.
speciosa, G. Don, l. c. 451 = Cass. speciosa.
torosa, G. Don, l. c. 452 = Cass. Sophera.

CHAMAEGERON, Schrenk in Bull. Phys.-Math. Acad. Pétersb. iii. (1846) 107. *COMPOSITAE*. oligocephalum, *Schrenk. l. c.*—Scongaria.

CHAMAEIRIS, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 417 = **Iris**, Tourn.
angustifolia, Medic. l. c. 418 = I. pumila.
desertorum, Medic. l. c. = I. Gueldenstaedtiana.
foetida, Medic. l. c. = I. foetida.
graminea, Medic. l. c. = I. graminea.
spuria, Medic. l. c. = I. spuria.

CHAMAELAUCIUM, Desf. in Mém. Mus. Par. v. (1819) 39. t. 3. *MYRTACEAE*, Benth. & Hook. f. i. 698.
DECALOPHUM, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 153.
afine, Meisn. in Journ. Linn. Soc. i. (1857) 45 = uncinatum.
axillare, *F. Muell. ex Benth. Fl. Austral.* iii. 88.—Austral.
brevifolium, *Benth. l. c.* 87.—Austral.
Brownii, Desf. in Mém. Mus. Par. v. (1819) 271. t. 19 = Verticordia Brownii.
ciliatum, *Desf. l. c.* 40.—Austral.
Drummondii, *Meisen. in Journ. Linn. Soc.* i. (1857) 44.—Austral.
gracile, *F. Muell. Fragm.* iv. 62.—Austral.
heteranthum, *Benth. Fl. Austral.* iii. 86.—Austral.
megalopetalum, *F. Muell. ex Benth. l. c.* 88.—Austral.
pauciflorum, *Benth. l. c.*—Austral.
plumosum, Desf. in Mém. Mus. Par. v. (1819) 42. t. 4 = Verticordia Brownii.
Thomasii, F. Muell. Fragm. iv. 137. t. 30 = Darwinia Thomasii.
uncinatum, *Schau. in Lehm. Pl. Preiss.* i. 97.—Austral.
verticordinum, F. Muell. Fragm. iv. 57 = Darwinia verticordina.
virgatum, *Endl. in Ann. Wien. Mus.* ii. (1840) 193.—Austral.

CHAMAELE, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 59 (1867). *UMBELLIFERAE*.
Tanakae, Franch. & Sav. Enum. Pl. Jap. i. 186 = Carum Tanakae.
tenera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 60.—Japon.

CHAMAELEA, [Tourn.] Adans. Fam. ii. 345 (1763) = **Cneorum**, Linn. (Simarub.).
tricoccos, Lam. Fl. Fr. ii. 682 = Cneorum tricoccum.

CHAMAELEDON, Link, Enum. Hort. Berol. i. 210 (1821) = **Loiseleuria**, Desv. (Ericac.).
procumbens, Link, l. c. 211 = L. procumbens.

CHAMAELEON, Cass. in Dict. Sc. Nat. xlvii. 509 (1827) = **Atractylis**, Linn. (Compos.).
cauloscens, Cass. l. c. l. 60 = A. macrophylla.
gummifer, Cass. l. c. xlvii. 509 = A. gummifera.
megacephalus, Cass. l. c. l. 59 = A. macrocephala.

CHAMAELINUM, Host, Fl. Austr. ii. 224 (1831) = **Camelina**, Crantz (Crucif.).
austriacum, Host, l. c. 225 = Nasturtium austriacum.
paniculatum, Host, l. c. 224 = Nesilia paniculata.
sativum, Host, l. c. 225 = Cam. sativa.

CHAMAELIRIUM, Willd. in Ges. Nat. Fr. Berl. Mag. ii. (1808) 18. *LILIACEÆ*, Benth. & Hook. f. iii. 826.
DASURUS, Salisb. Gen. Pl. Fragm. 51 (1866).
DICLINOTRYS, Rafin. Neogen. 3 (1825).
OPHIOSTACHYS, [Delile] in Red. Lil. t. 464 (1816).
SIRAITOS, Rafin. Fl. Tellur. iv. 26 (1836).
carolinianum, *Willd. l. c.* 19.—Am. bor.
lutrum, A. Gray, Man. Bot. N. U. States, ed. V. 527 = carolinianum.
luteum, Miq. Ann. Mus. Bot. Lugd. Bat. 144 = Chionographis japonica.

CHAMAELUM, Baker, in Journ. Linn. Soc. xvi. (1877) 190, sphalm. = **Chamelum**, Phil. (Irid.).

CHAMAEMELUM, Lindl. in Trans. Linn. Soc. xiii. (1822) 104. t. 11. *ROSACEAE*, Benth. & Hook. f. i. 628.
corsicum, *Lindl. l. c.*—Ins. Madera.
mexicana, Batll. Adansonia, ix. (1869) 148 = Photinia mexicana.

CHAMAEMELUM, Tourn. ex Adans. Fam. ii. 128 (1763) ; Cass. in Dict. Sc. Nat. xxix. 179, 185 (1823) = **Anthemis**, Linn. (Compos.).
alpinum, All. Fl. Pedem. i. 186 = Achillea oxyloba.
arvense, Schreb. Spicil. 18 = A. arvensis.
austriacum, J. & C. Presl, Fl. Cech. 178 = A. austriaca.
Cota, All. Fl. Pedem. i. 184 = A. Cota.
Cota, C. A. Mey. Verz. Pfl. Cauc. 75 = A. altissima.
Cotula, All. Fl. Pedem. i. 186 = A. Cotula.
discoidewm, All. l. c. 188 = A. tinctoria.
fuscidum, Bunge, Enum. Strp. Transc. iii. 144 = A. Cotula.
incrassatum, Hoffmgg. Fl. Port. ii. 348, ex DC. Prod. vi. 616 = Anacyclus clavatus.
maritimum, All. Fl. Pedem. i. 184 = A. maritima.
Marschallianum, C. A. Mey. Verz. Pfl. Cauc. 75 = A. Biebersteiniana.
mixtum, All. Fl. Pedem. i. 185 = A. mixta.
montanum, All. l. c. 187 = A. montana.
nobile, All. l. c. 185 = A. nobilis.
orientale, Schult. ex Steud. Nom. ed. II. i. 348 = A. montana.
pinnatifidum, Moench, Meth. 605 = A. mixta.
proliferum, Moench, l. c. Suppl. 357 = Cladanthus prolifer.
tinctorium, Schreb. Spicil. 145 = A. tinctoria.
tomentosum, All. Fl. Pedem. i. 184 = Anacyclus clavatus.
Triumfetti, All. l. c. 187 = A. austriaca.
valentinum, All. l. c. = Anacyclus valentinus.

CHAMAEMELUM, Vis. Osserv. Matric. 19 (1845) ; Fl. Dalm. ii. 84 (1847) = **Matricaria**, Linn. (Compos.).
ambiguum, Boiss. Diagn. Ser. I. xi. 20 (= *Pyrethrum ambiguum*).—Orient.
arvense, Reichb. f. ex Willk. in Bot. Zeit. xxii. (1864) 251 = M. glabra.
auriculatum, Boiss. Diagn. Ser. I. xi. 23.—Arabia.
callosum, Boiss. & Heldr. l. c. 22.—As. Min.
caucasicum, Boiss. l. c. 20 = M. grandiflora.
conolinium, Boiss. & Bal. l. c. Ser. II. vi. 98 = M. conolinia.
daghestanicum, Boiss. Fl. Orient. iii. 334.—Daghestan.
decipiens, Boiss. Diagn. Ser. I. xi. 23 = M. decipiens.
discoforme, Vis. Fl. Dalm. ii. 85 = M. corymbifera.
elongatum, Boiss. Diagn. Ser. I. xi. 20 (= *Pyrethrum elongatum*).—Orient.

CHAMAEMELUM :—
grandiflorum, Boiss. & Hausskn. in Boiss. Fl. Orient. iii. 331.—Mesopotam.
heterocarpum, Boiss. l. c. 329 = Matricaria heterocarpa.
inodorum, Vis. Osserv. Matric. 18 ; Fl. Dalm. ii. 85 = M. inodora.
Kochii, Boiss. Fl. Orient. iii. 329 = M. microcephala.
Kotschyi, Boiss. Diagn. Ser. II. iii. 97 = M. Oreades.
lamellatum, Boiss. Fl. Orient. iii. 326 = M. lamellata.
limonum, Maxim. Prim. Fl. Amur. 156 = M. inodora.
maritimum, Boiss. Diagn. Ser. I. xi. 21 = M. maritima.
melanolepis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 118 = M. Oreades.
microcephalum, Boiss. Diagn. Ser. I. xi. 23 = M. microcephala.
monticolum, Boiss. & Huet, l. c. Ser. II. iii. 27.—Armenia.
nanum, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 281.—Kurdistan.
olympicum, Boiss. ex Tchihat. Asie Min. Bot. ii. 258, nomen.—As. Min.
Oreades, Boiss. Diagn. Ser. I. xi. 21 = M. Oreades.
Pichleri, Boiss. Fl. Orient. iii. 332.—Bithynia.
praecox, Vis. Osserv. Matric. 18 ; Fl. Dalm. ii. 86 = M. praecox.
Reuterianum, Boiss. ex Tchihat. Asie Min. Bot. ii. 258, nomen.—As. Min.
roseilum, Boiss. & Orph. in Boiss. Fl. Orient. iii. 334 = M. rosella.
Szovitsii, Boiss. l. c. 332.—Reg. Caucas.
Tchihatchewi, Boiss. in Tchihat. Asie Min. Bot. ii. 255.—As. Min.
tetramastopernum, F. Schmidt, Reisen Amur. 148.—Reg. Amur.
trichophyllum, Boiss. Diagn. Ser. I. xi. 21 = M. trichophylla.
uniglandulosum, Vis. Osserv. Matric. 18 ; Fl. Dalm. ii. 85 = M. trichophylla.

CHAMAEMESPILUS, Medic. Phil. Bot. i. 138, 155 (1780) = **Pyrus**, Linn. (Rosac.).
ambigua, Gren. ex Nym. Consp. 242 = P. Chamaemespilus.
Aria, Gilib. Exerc. Phyt. Roseae, 12 = P. Aria.
humilis, M. Roem. l. c. 131 = P. Chamaemespilus.
Pseudaria, M. Roem. l. c. 132 = P. Aria.
sudetica, M. Roem. l. c. 131 = P. sudetica.

CHAMAEMYRRHIS, Endl. ex Heynh. Nom. ii. 136 (1846) = **Caldasia**, Lag. (Umbellif.).
eriopoda, Endl. l. c. = Cald. eriopoda.

CHAMAENERION, [Tourn.] Adans. Fam. ii. 85 (1763) = **Epilobium**, Linn. (Onagrar.).
americanum, Sweet, Hort. Brit. ed. II. 198 = E. palustre.
angustifolium, Schur, Enum. Pl. Transs. 213 = E. angustifolium.
angustifolium, Scop. Fl. Carn. ed. II. i. 271 = E. Dodonaei.
denticulatum, Schur, Enum. Pl. Transs. 214 = E. angustifolium.
Dodonaei, Schur, ex Fuss. Fl. Transs. 214 = E. Dodonaei.
grandiflorum, Moench, Meth. 677 = E. hirsutum.
halimifolium, Salisb. Parad. Lond. t. 68 = E. latifolium.
Halleri, Sweet, Hort. Brit. ed. II. 198 = E. Dodonaei.
hirsutum, Scop. Fl. Carn. ed. II. i. 270 = E. hirsutum.
latifolium, Sweet, Hort. Brit. ed. II. 198 = E. latifolium.
montanum, Scop. Fl. Carn. ed. II. i. 270 = E. montanum.
obscurum, Schreb. Spicil. 147 = E. tetragonum.
palustre, Schreb. l. c. 23 = E. palustre.
parviflorum, Schreb. l. c. 146 = E. parviflorum.
pubescens, Moench, Meth. 677 = E. parviflorum.
ramosissimum, Moench, l. c. 678 = E. tetragonum.
roseum, Schreb. Spicil. 147 = E. roseum.
roseum×hirsutum, Moench, Meth. Suppl. 288 = E. Dodonaei.
speciosum, Lodd. ex Steud. Nom. ed. II. i. 548 = E. angustifolium.
spicatum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 569 = E. angustifolium.
tetragonum, Scop. Fl. Carn. ed. II. i. 271 = E. tetragonum.

CHAMAENERIUM, Spach, Hist. Vég. Phan. iv. 396 (1835) = prec.

CHAMAEORCHIS, Koch, Syn. Fl. Germ. 692 (1837) = CHAMORCHIS, Rich. = **Herminium**, Willd. (Orchid.).

CHAMAEPEUCE, DC. Prod. vi. 557 (1837) = Cnicus, Tourn. (Compos.).
afra, DC. l. c. 558 = Cn. afer.
Alpini, Jaub. & Spach, Illustr. v. 26. t. 425 = Cn. Chamaepence.
atropurpurea, Boiss. & Heldr. Diagn. Ser. II. iii. 47 = Cn. strictus.
Casabonae, DC. Prod. vi. 558 = Cn. Casabonae.
cynaroides, DC. l. c. 559 = Cn. cynaroides.
diacantha, DC. l. c. = Cn. diacanthus.
echinocephala, DC. l. c. 560 = Cn. echinocephalus.
firma, DC. l. c. 559 = Cn. firmus.
fruticosa, DC. l. c. 558 = Cn. fruticosus.
graphalodes, DC. l. c. = Cn. graphalodes.
hispanica, DC. l. c. = Cn. hispanicus.
horrida, DC. l. c. 560 = Cn. horridus.
imbricata, Rafin. Fl. Tellur. iv. 118 = Cn. Chamaepeuce.
macrantha, Schrank, in Ball. Sc. Acad. Pétersb. x. (1842) 354 = Cousinia Hahlii.
macrostachya, Trautv. in Bull. Phys.-Math. Acad. Pétersb. xvi. (1858) 326.—Reg. Caucas.
mutica, DC. Prod. vi. 557 = Cn. Chamaepence.
nivea, DC. Prod. vi. 559 = Cn. nivea.
polyacantha, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = Cn. Casabonae.
polycephala, DC. Prod. vi. 558 = Cn. Chamaepence.
sinuata, Trautv. in Act. Hort. Petrop. ii. (1873) 477 = Cn. cynaroides.
stellata, DC. Prod. vi. 558 = Cn. stellatus.
stricta, DC. l. c. 559 = Cn. strictus.

CHAMAEPEUCE, Zocc. in Endl. Enchir. 189 (1841) = **Thuya**, Tourn. (Conif.).
obtusa, Zucc. ex Gord. Pinet. Suppl. 98 = T. obtusa.

CHAMAEPHYTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. vii. (1868) 37 = **Potentilla**, Linn. (Rosac.).
supinum, Fourr. l. c. = P. supina.

CHAMAEPITYS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 230 (1745) = **Ajuga**, Tourn. (Labiat.).
arvensis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 60 = A. Chamaepitys.
Ivа, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 138 = A. Iva.
pseudoiva, Fourr. l. c. = A. Iva.
trifida, Dum. Fl. Belg. 42 = A. Chamaepitys.
vulgaris, Link, Handb. i. 458 = A. Chamaepitys.

CHAMAEPLIUM, Wallr. Sched. Crit. i. 376 (1822) = Stegnotheca, Tourn. (Crucifer.).
officinale, Wallr. l. c. 377 = S. officinale.
polyceratium, Wallr. l. c. = S. polyceratium.
supinum, Wallr, l. c. = S. supinum.

CHAMAERANTHEMUM, Nees, in Lindl. Nat. Syst. ed. II. 445 (1836). *ACANTHACEAE*, Benth. & Hook. f. ii. 1095.
Beyrichii, *Nees, l. c.* 445.—Bras.
Gaudichaudii, *Nees, in Mart. Fl. Bras.* ix. 155.—Bras.
igneum, *Regel, Gartenfl.* (1868) 353.—Peruv.
pictum, *Mast. in Gard. Chron.* (1878) i. 527.—Bras.

CHAMAERAPHIS, R. Br. Prod. 192 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1107.
abortiva, Poir. Encyc. Suppl. ii. 159 = spinescens.
aspera, *Nees, in Wall. Cat.* n. 8679 = spinescens.
compressa, Poir. ex Steud. Nom. ed. I. 183 = Pennisetum compressum.
depauperata, Nees, ex Steud. Syn. Pl. Gram. 49 = Panicum asperum, *Wight*.
gracilis, *Hook. in Engl. Bot. Yahrb.* vi. (1885) 236.—Malaya.
hordeacea, *R. Br. Prod.* i. 194.—Austral.
pensilosa, *Poir. Encyc. Suppl.* ii. 159.—Austral.
perpilosa, *Munro, ex Wright, in Sessn. Fl. Cub.* 200, nomen.—Cuba.

CHAMAERAPHIS :—
spinescens, *Poir. Encyc. Suppl.* ii. 189.—As. trop.; Austral.
spinans, Beauv. ex Roem. & Schult. Mant. ii. 258 = spinescens.

CHAMAEREPES, Spreng. Syst. iii. 702 (1826) = **Herminium**, Wlld. (Orchid.).
alpina, Spreng. l. c. = H. alpinum.

CHAMAERHODODENDROS, Tourn. ex Hall. in Rupp. Fl. Jen. ed. Hall. 249 (1745) = **Rhododendron**, Linn. (Ericac.).

CHAMAERHODOS, Bunge, in Ledeb. Fl. Alt. i. 429 (1829). *ROSACEAE*, Benth. & Hook. f. i. 631.
altaica, *Bunge, l. c.*—Sibir. alt. occ.
erecta, *Bunge, l. c.* 430.—Sibir.; Am. bor. occ.
grandiflora, *Ledeb. l. c.* 431, in nota.—Sibir.
mongholica, *Bunge, in Ann. Sc. Nat.* Ser. II. xix. (1843) 117.—Mongolia.
polygyna, G. Don, Gen. Syst. ii. 568 = erecta.
sabulosa, *Bunge, in Ledeb. Fl. Alt.* i. 431.—Sibir.
trifida, *Ledeb. Fl. Ross.* ii. 84.—Sibir.

CHAMAERIPHES, Ponted. ex Gaertn. Fruct. i. 25 (1789) = seq.

CHAMAEROPS, Linn. Mus. Cliff. 10; Gen. ed. I. 354 (1737). *PALMAE*, Benth. & Hook. f. iii. 924. CHAMAERIPHES, Ponted. ex Gaertn. Fruct. i. 25 (1789).
acaulis, Michx. Fl. Bor. Am. ii. 207 = Sabal Adansoni.
Antillarum, Desc. Antill. i. t. 28 (Thrinacis sp. !).—Ind. occ.
arborescens, Steud. Nom. ed. I. 183 = humilis.
arundinacea, Sm. in Rees, Cycl. vii. n. 3 = Rhapidophyllum Hystrix.
Biroo, Siebold, in Mart. Hist. Nat. Palm. 252 = Livistona rotundifolia.
cochinchinensis, Lour. Fl. Cochinch. ii. 557 = Rhapis cochinchinensis.
condulpicata, Kickx, in Bull. Acad. Brux. v. (1838) 61 = humilis.
excelsa, Thunb. Fl. Jap. 139 = Trachycarpus excelsus.
excelsior, *Baj. Hort. Maurist.* 307.—Ins. Maurit.
Fortunei, Hook. Bot. Mag. t. 5221 = Trachycarpus Fortunei.
Ghiesbreghti, Hort. ex H. Wendl. in Kerch. Palm. 240 (= Sabal sp.).—Hab. !
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = Sabal Adansoni.
gracilis, *Lodd. Cat.* (1838) *ex Schult. f. Syst.* vii. 1489, nomen.—Am. austr.
Griffithiana, Wall. ex Voigt, Hort. Suburb. Calc. 641, nomen = seq.
Griffithii, Lodd. Cat. (1841) ex Verlot, in Rev. Hortic. (1870) t. 376 = Trachycarpus khasyanus.
guianensis, *Lodd. Cat. ex Loud. Hort. Brit.* 413, nomen.—Guiana.
humilis, *Linn. Sp. Pl.* 1187.—Reg. Mediterr. occ.
Hystrix, Desf. ex Steud. Nom. ed. I. 183 = Corypha Hystrix.
Hystrix, Pras. ex Parsh, Fl. Am. Sept. i. 240 = Rhapidophyllum Hystrix.
khasyana, Griff. in Calc. Journ. Nat. Hist. v. (1845) 341 = Trachycarpus khasyanus.
Kwamwortzii, Siebold, ex H. Wendl. Ind. Palm. (1854) 24 = Rhapis flabelliformis.
louisiana, Rafin. Fl. Ludov. 159 = Rhapidophyllum Hystrix!
macrocarpa, *Linden, Cat.* n. 87 (1871).—Hab. !
macrocarpa, Tin. in Guss. Fl. Sic. Syn. ii. 683 = humilis.
Martiana, Wall. Pl. As. Rar. iii. 5. t. 211 = Trachycarpus Martianus.
nitida, *Y. Moy. in Mém. Acad. Bert.* (1769) 29.—Hab. !
Mocini, H. B. & K. Nov. Gen. et Sp. i. 300 = Acanthorhiza Mocinni.
napelanaia, Lodd. Cat. (1832) ex Schult. f. Syst. vii. 1489 = Trachycarpus Martianus.
Palmetto, Michx. Fl. Bor. Am. i. 206 = Sabal Palmetto.
radiata, *Yuss. ex Schult. f. Syst.* vii. 1489, nomen.—Hab. !
Ritchieana, Griff. in Calc. Journ. Nat. Hist. v. (1845) 343 = Nannorrhops Ritchieana.
servulata, Michx. Fl. Bor. Am. i. 206 = Brahea serrulata.

CHAMAEROPS :—
Sirotzii, Hort. ex H. Wendl. Ind. Palm. (1854) 17 = Rhapis humilis.
stauracantha, Hort. Belg. ex Heynh. Nom. ii. 136 = Acanthorhiza aculeata.
tomentosa, *Fulchiron, ex Schult. f. Syst.* vii. 1489, nomen.—Hab. !
tomentosa, *Ch. Morr. in Ann. Gand.*, i. (1845) 468.—Hab. !

CHAMAESARACHA, A. Gray, ex Franch. & Sav. Enum. Pl. Jap. ii. 454 (1876). *SOLANACEAE*, Benth. & Hook. f. ii. 891.
conioides, *Benth. & Hook. f. Gen.* ii. 891.—Mexic.
Coronopus, *A. Gray, in Bot. Calif.* i. 540.—Calif.
japonica, *Franch. & Sav. Enum. Pl. Jap.* ii. 454.—Japon.
nana, *A. Gray, in Bot. Calif.* i. 540.—Calif.
physaloides, Greene, in Bull. Torrey Club, ix. (1882) 102 = Physalis lobata.
sordida, *A. Gray, Synop. Fl. N. Am.* ii. i. 282.—Texas; Mexic.

CHAMAESARACHIA, Franch. & Sav. Enum. Pl. Jap. ii. 454 (1876) = praec.

CHAMAESCIADIUM, C. A. Mey. Verz. Pfl. Cauc. 122 (1831). *UMBELLIFERAE*, Benth. & Hook. f. i. 896.
acaule, *C. A. Mey. l. c.*—As. Min.; Reg. Caucas.
albiflorum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 260.—Sibir.
flavescens, C. A. Mey. Verz. Pfl. Cauc. 122 = acaule.
gracile, Boiss. & Heldr. in Boiss. Diagn. Ser. II. ii. 79 = Carum repostre.
Heldreichii, Boiss. l. c. 78 = Carum Heldreichii.

CHAMAESCILLA, F. Muell. Fragm. vii. 68 (1870). *LILIACEAE*, Benth. & Hook. f. iii. 791.
corymbosa, *F. Muell. l. c.*—Austral.
spiralis, *F. Muell. l. c.*—Austral.

CHAMAESPARTIUM, Adans. Fam. ii. 321 (1763) = **Cytisus**, Linn. et **Genista**, Linn. (Legumin.).

CHAMAESPARTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 357 = **Genista**, Linn. (Legumin.).
pilosum, Fourr. l. c. = G. pilosa.

CHAMAESPHACOS, Schrenk, Enum. Pl. Nov. i. 27 (1841). *LABIATAE*, Benth. & Hook. f. ii. 1207.
lilicifolius, *Schrenk, l. c.*—Sibir.

CHAMAESPHAERION, A. Gray, in Hook. Kew Journ. iii. (1851) 176 = **Chthonocephalus**, Steetz (Compos.).
pygmaeum, A. Gray, l. c. 177 = Chthonocephalus pygmaeus.

CHAMAESPILUS, Fourr. in Ann. Soc. Linn. Lyon, N.S. xvi. (1868) 378 = **Pyrus**, Linn. (Rosac.).
sudetica, Fourr. l. c. = P. sudetica.

CHAMAESTEPHANUM, Willd. in Ges. Nat. Fr. Berl. Mag. i. (1807) 140 = **Schkuhria**, Roth (Compos.).

CHAMAESVCE, S. F. Gray, Nat. Arr. Brit. Pl. ii. 260 (1891) = **Euphorbia**, Linn.
maritima, S. F. Gray, l. c. = E. Peplis.

CHAMAEZELUM, Link, Handb. i. 719 (1829) = **Anthemaria**, R. Br. (Compos.).
alpinum, Link, l. c. = A. alpina.
corpaticum, Link, l. c. = A. carpathica.
dioicum, Link, l. c. = A. dioica.
margaritaceum, Link. l. c. = Anaphalis margaritacea.

CHAMAEXEROS, Benth. Fl. Austral. vii. 110 (1878). *JUNCACEAE*, Benth. & Hook. f. iii. 464.
fimbriata, *Benth. l. c.* 111.—Austral.
Serra, *Benth. l. c.* 110.—Austral.

CHAMAEXIPHIUM, Hochst. in Flora, xviii. (1844) 102 = **Pleinia**, Schrad. (Cyperac.).
clandestinum, Hochst. l. c. = P. clandestina.
Dregeanum, Steud. Syn. Pl. Cyp. 89 = P. laevis.

CHAMAGROSTIS, Borkh. ex Wibel, Prim. Fl. Werth. 126 (1799) = **Milium**, Adans. (Gramin.).
minima, Borkh. l. c. = M. vernum.

CHAMALIUM, Juss. in Ann. Mus. Par. vi. (1805) 324 = **Cardopatium**, Juss. (Compos.).

CHAMALIUM, Cass. in Dict. Sc. Nat. xlvii. (1827) 498, 509 = CHAMAELEON, Cass. = **Atractylis**, Linn. (Compos.).

CHAMAREA, Eckl. & Zeyh. Enum. 346 (1837) = **Carum**, Linn. (Umbellif.).
caffra, Eckl. & Zeyh. l. c.—Afr. austr.
capensis, Eckl. & Zeyh. l. c. = Carum capense.

CHAMBEYRONIA, Viell. in Bull. Soc. Linn Normand. Sér. II. vi. 1872 (1873) 229. *PALMAE*. Brésiensii, *Viell. l. c.* 230, nomen.—N. Caled.
macrocarpa, *Viell. l. c.* nomen.—N. Caled.
Moriori, *Viell. l. c.* nomen.—N. Caled.

CHAMEDRYS, Rafin. Fl. Tellur. iii. 85 (1836) = CHAMAEDRYS, Tourn. = **Teucrium**, Tourn. (Scrophul.).

CHAMELUM, Phil. in Linnaea, xxxiii. (1864–65) 250. *IRIDEAE*, Benth. & Hook. f. iii. 700. CHAMALIUM, Baker, in Journ. Linn. Soc. xvi. (1877) 120, sphalm.
andinum, *Benth. & Hook. f. Gen.* iii. 700.—Chili.
luteum, *Phil. in Linnaea*, xxxiii. (1864–65) 250.—Chili.

CHAMIRA, Thunb. Nov. Gen. ii. 48 (1782). *CRUCIFERAE*, Benth. & Hook. f. i. 81.
cornuta, *Thunb. l. c.* 49.—Afr. austr.

CHAMISME, Rafin. ex Steud. Nom. ed. II. i. 776 (in syn.) (1840) = **Houstonia**, Gron. (Rubiac.).

CHAMISSOA, H. B. & K. Nov. Gen. et Sp. ii. 158. t. 125 (1817). *AMARANTACEAE*, Benth. & Hook. f. iii. 27.
KOREA, Adans. Fam. ii. 269 (1763).
acuminata, *Mart. in Nov. Act. Nat. Cur.* xiii. (1826) 286.—Bras.
altissima, Mart. l. c. 287 = Allmania altissima, *H. B. & K. Nov. Gen. et Sp.* 197. t. 125.—Am. austr.
altissima, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 92 = Allmania nodiflora.
angustifolia, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 717 = Allmania nodiflora.
arabica, Spreng. Syst. i. 815 = Digera arvensis.
aspera, Wight, Ic. v. t. 1772 = Allmania nodiflora.
Blanchetii, *Moq. in DC. Prod.* xiii. ii. 251.—Bras.
Brownii, Steud. Nom. ed. II. i. 344 = Allmania nodiflora.
celosioides, *Griseb. in Goett. Abh.* xix. (1874) 79.—Reg. Argent.
ciliata, Spreng. Syst. i. 815 = Digera arvensis.
communis, Spreng. l. c. = Digera arvensis.
dichotoma, Moq. in DC. Prod. xiii. ii. 249 = Allmania nodiflora.
esculenta, Moq. l. c. = Allmania nodiflora.
javanica, Hassk. Fl. Jav. Rar. 484 = Allmania nodiflora.
macrocarpa, *H. B. & K. Nov. Gen. et Sp.* ii. 197.—Am. austr.
Martii, Moq. in DC. Prod. xiii. ii. 253 = altissima.
Maximiliani, *Mart. ex Moq. l. c.* 251.—Bras.
muricata, Spreng. Syst. i. 815 = Digera arvensis.
nodiflora, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 287 = Allmania nodiflora.
ovata, Endl. ex Moq. in DC. Prod. xiii. ii. 252 = Charpentiera ovata.
paniculata, Moq. ex Moq. l. c. 249 = Celosia Moquini.
populifolia, Hochst. ex Moq. l. c. = Celosia populifolia.
pyramidalis, Moq. l. c. 248 = Allmania albida.

CHAMISSONIA, Endl. Gen. 1190, in syn. (1840) = CAMISSONIA, Link = **Oenothera**, Linn. (Onagrar.).

CHAMITEA, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 975 = **Baltia**, Tourn.
reticulata, Kern. l. c. 977 = S. reticulata.

CHAMITIS, Banks, ex Gaertn. Fruct. i. 94. t. 22 (1788) = **Ancrella**, Lam. (Umbell).
complicata, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 286 = A. caespitosa.
integrifolia, Gaertn. Fruct. i. 94. t. 22. f. 4 = A. filamentosa.
tricuspidata, Gaertn. l. c. t. 22 = A. trifurcata.
trifurcata, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 286 = A. trifurcata.
trifurcata, Gaertn. Fruct. i. 95 = A. caespitosa.

CHAMOLETTA, Adans. Fam. ii. 60 (1763) = Iris, Linn.

CHAMOMILLA, [Hall.] S. F. Gray, Nat. Arr. Brit. Pl. ii. 454 (1821); C. Koch, in Linnaea, xvii. (1843) 45 = **Matricaria**, Linn. (Compos.).
brachyglossa, J. Gay, ex Benth. & Hook. f. Gen. ii. 498 = M. brachyglossa.
Courrantiana, C. Koch, in Linnaea, xxiv. (1851) 338, in obs. = M. Chamomilla.
discoidea, J. Gay, ex A. Br. in Bot. Zeit. x. (1852) 650 = M. discoidea.
glabrata, Sch.-Bip. Ueber Tanacet. 21 = M. glabrata.
inodora, C. Koch, in Linnaea, xvii. (1843) 45 = M. inodora.
meridionalis, C. Koch, l. c. = M. Chamomilla.
officinalis, C. Koch, l. c. = M. Chamomilla.
patens, Gilib. Fl. Lituan. iii. 220 = M. Chamomilla.
praecox, C. Koch, in Linnaea, xvii. (1843) 46 = M. inodora.
umbellaralis, C. Koch, l. c. xxiv. (1851) 338.—Reg. Cancas.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 454 = M. Chamomilla.

CHAMOMILLA, Godr. Fl. Lorr. ii. 19 (1843) = **Anthemis**, Linn. (Compos.).
axrea, J. Gay, in Bull. Soc. Bot. Fr. iv. (1857) 278 = Cotula axrea.
fuscata, Gren. & Godr. Fl. Fr. ii. 161 = A. fuscata.
mixta, Gren. & Godr. l. c. = A. mixta.
nobilis, Godr. Fl. Lorr. ii. 19 = A. nobilis.

CHAMORCHIS, Rich. in Mém. Mus. Par. iv. (1818) 49 = **Herminium**, Linn. (Orchid.).
albida, Dum. Fl. Belg. 133 = Habenaria albida.
alpina, Rich. in Mém. Mus. Par. iv. (1818) 57 = Herminium alpinum.
viridis, Dum. Fl. Belg. 133 = Habenaria viridis.

CHAMPACA, Adans. Fam. ii. 365 (1763) = **Michelia**, Linn. (Magnoliac.).
fuscicaula, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 13.—Malaya.
Michelia, Noronha, l. c. 11 = M. Champaca.
turbinata, Noronha, l. c. 12.—Malaya.

CHAMPEREIA, Griff. in Calc. Journ. Nat. Hist. iv. (1843) 227. *SANTALACEAE*, Benth. & Hook. f. iii. 221.
MALOUCHAN, Blanco, Fl. Filip. ed. I. 188 (1837).
gustocarpa, *Kurz, in Journ. Bot.* xiii. (1875) 325.—Isa. Nicobar.
Griffithiana, *Planch. ex Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 154.—Burma; Malaya.
Porretiotiana, Baill. Adansonia, iii. (1862–63) 195 = Scleropyrum Wallichianum.

CHAMPIONIA, C. B. Clarke, Comm. et Cyrt. Beng. 98. t. 88 (1874) = Leptobostesa, Benth. (Geanerac.).
multiflora, C. B. Clarke, l. c. = L. multiflora.

CHAMPIONIA, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 488. *GESNERACEAE*, Benth. & Hook. f. ii. 1094.
reticulata, *Gardn. l. c.*—Zeylan.

CHAMULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

CHAPELIERA, Meisn. Gen. 163; Comm. 116 (1838) = seq.

CHAPELIERIA, A. Rich. in Mem. Soc. Hist. Nat. Par. v. (1830) 253. *RUBIACEAE*, Benth. & Hook. f. ii. 96.
madagascariensis, *A. Rich. l. c.* 253.—Madag.

CHAPELLIERA, Nees, in Linnaea, ix. (1834) 298 = Cladium, P. Br. (Cyperac.).
arthrophylla, Nees, in Lehm. Pl. Preiss. ii. 77 = Clad. arthrophyllum.
articulata, Nees, l. c. 76, in adnot. = Clad. articulatum.
glomerata, Nees, l. c. = Clad. glomeratum.
trifolia, Nees, in Linnaea, ix. (1834) 298 = Clad. flexuosum.
juncea, Nees, in Lehm. Pl. Preiss. ii. 76, in adnot. = Clad. junceum.
laxa, Nees, l. c. 76 = Clad. laxum.
pauciflora, Nees, l. c. 76, in adnot. = Tricostularia pauciflora.
riparia, Nees, l. c. 76 = Clad. riparium.
tereticlia, Nees, l. c. 76, in adnot. = Clad. teretifolium.

CHAPMANNIA, Torr. & Gray, Fl. N. Am. i. 355 (1839). *LEGUMINOSAE*, Benth. & Hook. f. i. 517.
floridana, *Torr. & Gray, l. c.*—Florida.

CHAPTALIA, Royle, Illustr. Bot. Himal. t. 59. f. 2 (1839) = **Gerbera**, Gronov. (Compos.).
gossypina, Royle, l. c. 251 = G. lanuginosa.
maxima, D. Don, Prod. Fl. Nep. 166 = G. macrophylla.

CHAPTALIA, Vent. Jard. Cels, t. 61 (1800). *COMPOSITAE*, Benth. & Hook. f. ii. 498.
LERIA, DC. in Ann. Mus. Par. xix. (1812) 65.
LEBETINA, Cass. in Dict. Sc. Nat. xxvi. 286 (1823).
LOXODON, Cass. l. c. xxvii. 258 (1823).
OXYDON, Less. in Linnaea, v. (1830) 257.
TETRANTHEMA, Neck. Elem. i. 6 (1790).
albicans, *Vent. ex Steud. Nom.* ed. II. i. 844.—Ind. occ.
Anandria, Spreng. Syst. iii. 504 = Gerbera Anandria.
arenosa, *Cass. in Atti 9ª Riun. Sc. Ital.* iii. (1841) 514.—Bras.
dentata, *Cass. in Dict. Sc. Nat.* xxvi. 104.—Ind. occ.
denticulata, Spreng. ex Baker, in Mart. Fl. Bras. vi. iii. 310 = Senecio pulcher.
Ehrenbergii, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
exscapa, *Poepp. ex DC. Prod.* vii. 44, in syn.—Chili.
heterophylla, D. Don, in Trans. Linn. Soc. xvi. (1830) 244 = Trichocline heterophylla.
integrifolia, *Baker, in Mart. Fl. Bras.* vi. iii. 377.—Am. trop.
integrifolia, Vent. ex Spreng. Syst. iii. 504 = tomentosa.
lyrata, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 248.—Mexic.
lyrata, Spreng. Syst. iii. 504 = Gerbera Anandria.
nutans, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
obovata, *D. Don, in Trans. Linn. Soc. xvi.* (1830) 240.—Peruv.
obovata, *Wright, in Sauv. Fl. Cub.* 88.—Ins. Caba.
ovalis, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 241.—Peruv.
piloselloides, *Baker, in Mart. Fl. Bras.* vi. iii. 378.—Bras.
piloselloides, *Vent. ex Steud. Nom.* ed. II. i. 844; *Baker, in Mart. Fl. Bras.* vi. iii. 378.—Reg. Argent.
rotundifolia, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 245.—Peruv.
runcinata, *H. B. & K. Nov. Gen. et Sp.* iv. 6.—N. Granat.
sarmentosa, Vent. ex Steud. Nom. ed. II. i. 844 = Gerbera sarmentosa.
Seemannii, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
sinuata, *Baker, in Mart. Fl. Bras.* vi. iii. 378.—Reg. Uruguay; Argent.
sinuata, *Vent. ex Steud. Nom.* ed. II. i. 844.—Ins. Jamaic.
spathulata, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
tomentosa, *Vent. Jard. Cels*, t. 61.—Am. bor.

CHAQUEPIRIA, Endl. Gen. 862, in syn. (1836) sphalm. = CAQUEPIRIA, J. F. Gmel. = **Gardenia**, Linn. (Rubiac.).

CHARACHERA, Nevsk. Fl. Aegypt. Arab. 116 (1779). *ACANTHACEAE* i, Benth. & Hook. f. ii. 1071. XERALIS, Raffa. Sylva Telnr. 89 (1836).

CHARACHERA:—
spicata, Lam. Encyc. i. 695 = viburnoides.
viburnoides, *Forsk. Fl. Aegypt. Arab.* 116.—Arabia.

CHARACIAS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 259 (1821) = **Euphorbia**, Linn.
amygdaloides, S. F. Gray, l. c. 260 = E. amygdaloides.
purpurea, S. F. Gray, l. c. 259 = E. Characias.

CHARADRA, Scop. Introd. 270 (1777) = CHADARA, Forsk. = **Grewia**, Linn.

CHARDINIA, Desf. in Mem. Mus. Par. iii. (1817) 456. t. 21. *COMPOSITAE*, Benth. & Hook. f. ii. 464.
cylindrica, *Desv. Fl. Anj.* 216 = Xeranthemum cylindraceum.
macrocarpa, C. Koch, in Linnaea, xxiv. (1851) 332 = xeranthemoides.
xeranthemoides, *Desf. in Mém. Mus. Par.* iii. (1817) 456.—Ins. Min.; Persia.

CHARLANTHUS, D. Don, in Mem. Wern. Soc. iv. (1823) 327. *MELASTOMACEAE*, Benth. & Hook. f. i. 762.
CHARNANTHERA, Rich. ex DC. Prod. iii. 126 (1828).
TETRAGYDOX, Rich. ex DC. l. c. 127 (1828).
Berteroanus, Ser. ex DC. l. c. 127 = glaberrimus.
brasiliensis, Cham. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 118 = Miconia brasiliensis.
caerulescens, D. Dietr. Syn. Fl. ii. 1277 = Memecylon Spathandra.
ciliatus, *DC. Prod.* iii. 127.—Ins. Trinitat.
coccineus, *D. Don, in Mem. Wern. Soc. iv.* (1823) 328. —Am. austr.
crinitus, *Naud. in Ann. Sc. Nat.* Ser. III. xviii. (1852) 113.—Ind. occ.
Fadyeni, *Griseb. in Goett. Abh.* ix. (1861) 56.—Jamaic.
glaberrimus, *DC. Prod.* iii. 126.—Ind. occ.
nodosus, Triana, in Trans. Linn. Soc. xxviii. (1871) 99 = ciliatus.
obliquus, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 136 = Miconia obtusa.
obtusus, Triana, in Trans. Linn. Soc. xxviii. (1871) 108 = Miconia obtusa.
purpureus, *D. Don, in Mem. Wern. Soc. iv.* (1823) 329. —Ins. Mauricorat.
tinifolius, *D. Don, l. c.*—Ind. occ.

CHARIDIA, Baill. Etud. Gen. Euph. 572. t. 22 (1858) = **Savia**, Willd. (Euphorb.).
mimosoides, Baill. l. c. 573 = S. mimosoides.
pulchella, Baill. l. c. = S. pulchella.

CHARITEIS, Cass. in Bull. Soc. Philom. (1817) 68; et in Dict. Sc. Nat. viii. 191; xxiv. 369 (1817). *COMPOSITAE*. Benth. & Hook. f. ii. 266.
KAULFUSSIA, Nees, Hor. Phys. Berol. 58. t. 11 (1820).
caerulea, Cass. in Dict. Sc. Nat. xxiv. 373 = heterophylla.
heterophylla, *Cass. in Bull. Soc. Philom.* (1817) 68.—Afr. austr.
Novaë, Cass. l. c. (1821) 15, ex cod. in Dict. Sc. Nat. xxiv. 370, 373 = heterophylla.
pilosella, Baill. Hist. Fl. viii. 139 = heterophylla.

CHARIESSA, Miq. Fl. Ind. Bat. i. 1. 794 (1856) = **Villaresia**, Ruiz & Pav. (Olacin.).
Smythii, Becc. Malesia, i. (1877) 118.—Malaya.
suaveolens, Miq. Fl. Ind. Bat. i. 1. 794.—Malaya.

CHARICOMA, Miers, Apocyn. S. Am. 110 (1878) = **Echites**, Linn. (Apocyn.).
dominguensis, Miers, l. c. 119 = Echites dominguensis.
fama, Miers, l. c. 118 = Dipladenia fama.
macrocalyx, Miers, l. c. 115.—Ins. S. Thom.
nobilis, Miers, l. c. 118 = Dipladenia nobilis.
scandens, Miers, l. c. 114.—Ind. occ.
turrecta, Miers, l. c. 111 = Echites Andrewsii.
verticillata, Miers, l. c. 118.—Jamaic.

CHARLWOODIA, Sweet, Fl. Austral. t. 18 (1827) = Cordyline, Comm. (Liliac.).
angustifolia, Goepp. Beitr. Kenn. Dracaen. 17 = Cord. Pumilio.
australis, G. Don, in Loud. Hort. Brit. 180 = Cord. australis.
congesta, Sweet, Fl. Austral. t. 18 = Cord. Pumilio.
cuneata, Goepp. Beitr. Kenn. Dracaen. 17 = Dianella cuneata?

CHARLWOODIA :—
fragrantissima, Lem. Jard. Flear. iv. (1854) t. 399. Misc. 97 = Cordyline dracaenoides.
indivisa, G. Don, in Loud. Hort. Brit. 180 = Cord. indivisa.
longifolia, Goepp. Beitr. Kenn. Dracaen. 17.—Hab.?
rigidifolia, Koch & Bouche, ind. Sem. Hort. Berol. (1855) 11 = Cord. Pumilio.
rubra, Planch. in Fl. des Serres, vi. (1850–51) 138 = Cord. rubra.
Sellowiana, Planch. l. c. 137 = Cord. Sellowiana.
spectabilis, Planch. l. c. 136 = Cord. dracaenoides.
stricta, Sweet, Fl. Austral. sub t. 16 = Cord. Pumilio.

CHARPENTIERA, Gaudich. in Freyc. Voy. Bot. 444. t. 47. 48 (1826). *AMARANTACEAE*, Benth. & Hook. f. iii. 26.
obovata, *Gaudich. l. c.* t. 48.—Ins. Sandvic.
ovata, *Gaudich. l. c.* t. 47.—Ins. Sandvic.

CHARPENTIERA, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 346 = **Pavetta**, Linn. (Rubiac.).
bracteata, Vieill. l. c. 347.—N. Caled.

CHARTACALYX, Maing. ex Mast. in Hook. f. Fl. Brit. Ind. i. 382 (1874). *TILIACEAE*.
accrescens, *Mast. l. c.*—Malacca.

CHARTOCALYX, Regel, in Act. Hort. Petrop. vi. (1879) 867. *LABIATAE*.
Olgae, *Regel, l. c.* 868.—Turkestan.

CHARTOLEPIS, Cass. in Dict. Sc. Nat. xliv. 36 (1826) Centaurea, Linn. (Compos.).
Boberstrirti, Jaub. & Spach, Illustr. iii. 11 = Cent. glastifolia (ex parte).
cappadocica, Fenzl, in Tchihat. Asie Min. Bot. ii. 328 = Cent. cappadocica.
cappadocica, Jaub. & Spach, Illustr. iii. 137. t. 291 = Cent. macrolopha.
Cauniana, Jaub. & Spach, l. c. 9 = Cent. glastifolia.
cheirolopha, Fenzl, in Tchihat. Asie Min. Bot. ii. 324 = Cent. cheirocatha.
drabaefolia, Jaub. & Spach, Illustr. iii. 136 = Cent. drabifolia.
glastifolia, Cass. in Dict. Sc. Nat. xliv. 36 = Cent. glastifolia.
intermedia, Boiss. Diagn. Ser. II. iii. 64.—Ross. mer.; Sibir.; Soongar.
Kotschyi, Fenzl, in Tchihat. Asie Min. Bot. ii. 323 = Cent. macrolopha.
lancifolia, Fenzl, l. c. 327 = Cent. lancifolia.
libanotica, Jaub. & Spach, Illustr. iii. 136.—Syria.
lyrata, Boiss. & Hausk. ex Boiss. Fl. Orient. iii. 696. —Persia.
saligna, C. Koch, in Linnaea, xxiv. (1851) 421 (= *Phaeopappus saligneus*).—Armenia.
schistolepis, Boiss. ex Boiss. Fl. Orient. iii. 696 (= *Phaeopappus saligneus*).—Kurdist.
Tournefortii, Jaub. & Spach, Illustr. iii. 9 = Cent. glastifolia.

CHARTOLOMA, Bunge, in Bot. Zeit. ii. (1844) 249 = Inula, Tourn. (Crucif.).
platycarpum, Bunge, l. c. 250 = I. platycarpa.

CHASALIA, Comm. ex DC. Prod. iv. 531 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 126.
POLYOSMA, Blume, Bijdr. 947 (1826).
amicorum, *A. Gray, in Proc. Am. Acad.* iv. (1860) 43. —Ins. Fiji.
Borbonica, Comm. ex DC. Prod. iv. 532 = divaricata.
Boryana, *DC. l. c.*—Ins. Mascar.
capitata, *DC. l. c.* 531.—Ins. Maurit.
chondrophylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 208.—Sumatra.
chrysolepis, *DC. Prod.* iv. 532.—Ins. Maurit.
coffeoides, DC. l. c. 531 = divaricata!
Commersonii, Juss. ex Steud. Nom. ed. II. i. 345 = capitata.
curviflora, *Thw. Enum. Pl. Zeyl.* 150.—Ind. or.; Malaya.
divaricata, *DC. Prod.* iv. 532.—Ins. Maurit.
exserens, Miq. Fl. Ind. Bat. ii. 260 = Psychotria exserens.
Fournieri, *DC. Prod.* iv. 531.—Ins. Mascar.
lanciflora, *DC. l. c.* 532.—Ins. Maurit.

CHASALIA:—
Hasseltiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 204. —Java.
lanciflora, Benth. in Hook. Niger Fl. 416 = Psychotria Ansellii.
leucocarpa, Miq. Fl. Ind. Bat. ii. 281 = Psychotria leucocarpa.
lurida, Miq. l. c. 282 = curviflora.
montana, Miq. l. c. 281 = Psychotria montana.
Moonia, *Baker, Fl. Maurit.* 156.—Ins. Mascar.
oxylepis, *Miq. Fl. Ind. Bat.* ii. 283.—Java.
parviflora, *Benth. in Hook. Niger Fl.* 417.—Afr. trop.
perforata, Miq. Fl. Ind. Bat. Suppl. 546 = Psychotria angulata.
psychotrioides, DC. Prod. iv. 531 = Gaertnera psychotrioides.
pyriformis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 44. —Ins. Samoa.
robusta, Miq. Fl. Ind. Bat. ii. 280 = Psychotria robusta.
rostrata, *Miq. l. c.* 281.—Malaya.
sangiana, Miq. Fl. Ind. Bat. Suppl. 546 = curviflora.
stipulacea, DC. Prod. iv. 532 = divaricata.
tetrandra, Miq. Fl. Ind. Bat. ii. 283 = curviflora.
umbraticola, *Vathe, in Oestr. Bot. Zeitschr.* xxv. (1875) 230.—Afr. trop.

CHASALLIA, Comm. ex Poir. in Dict. Sc. Nat. viii. 198 (1817); et Juss. in Mém. Mus. Par. vi. (1820) 379 = prec.

CHASCANUM, E. Mey. Comm. Pl. Afr. Austr. 275 (1837) = Bouchea, Cham. (Verbenac.).
cernuum, E. Mey. l. c. 276 = B. cernua.
cuneifolium, E. Mey. l. c. = B. cuneifolia.
garipense, E. Mey. l. c. 277 = B. garepensis.
laetum, Fenzl, ex Schau. in DC. Prod. xi. 559 = B. hyssopifolia.
marrubiifolium, Fenzl, ex Schau. l. c. 558 = B. marrubiifolia.
pinnatifidum, E. Mey. Comm. Pl. Afr. Austr. 277 = B. pinnatifida.
pumilum, E. Mey. l. c. = B. pumila.

CHASCOLYTRUM, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190 = **Brizа**, Linn. (Gramin.).
barbatum, Kunth, Enum. Pl. i. 373 = B. barbata.
coarctatum, Phil. in Linnaea, xxxiii. (1864–65) 293.—Chili.
elegans, Beauv. Agrost. 86 = B. elegans.
erectum, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190 = B. erecta.
nutans, Lindl. ex Heynh. Nom. ii 137, nomen.—Chili.
rotundatum, Kunth, Rev. Gram. i. 121 = B. rotundata.
rufum, J. & C. Presl, Rel. Haenk. i. 326 = B. rufa.
spicigerum, J. & C. Presl, l. c. 322 = B. spicigera.
strictum, E. Desv. in C. Gay, Fl. Chil. vi. 384 = B. stricta.
subarticulatum, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190 = B. rotundata.
trilobum, E. Desv. in C. Gay, Fl. Chil. vi. 384 = B. trilobа.

CHASMANTHERA, Hochst. in Flora, xxvii. (1844) 21. *MENISPERMACEAE*, Benth. & Hook. f. i. 34.
RHIGIOCARYA, Miers, in Ann. Nat. Hist. Ser. III. xiv. (1864) 100.
dependens, *Hochst. l. c.*—Abyssin.
nervosa, *Miers, Contrib. Bot.* iii. 89.—Afr. trop.
uviformis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 489.—Madag.

CHASMANTHIUM, Link, Hort. Berol. i. 159 (1827) = Uniola, Linn. (Gramin.).
gracile, Link, l. c. = U. gracilis.
ornithorynchа, Nees, ex Steud. Syn. Pl. Gram. 380 = U. ornithorynchа.

CHASME, Salisb. Parad. Lond. sub t. 67 (1807); et in Knight, Prot. 16 (1809) = **Lonchostrobus**, R. Br. (Proteac.).
comosa, Salisb. l. c.; Knight, l. c. 17 = L. comosus.
pinifolia, Knight, l. c.—Afr. austr.
ramentacea, Knight, l. c. = L. acaulus.
spicata, Knight, l. c. 18.—Afr. austr.
ternifolia, Knight, l. c. = L. ulicinus.

CHASMIA, Schott, ex Spreng. Syst. iv. Cur. Post. 409 (1827) = **Arrabidaea**, DC. (Bignoniac.).
ochroleuca, Schott, l. c. = A. Agnus-castus, vel A. Salzmanni.

CHASMONE, E. Mey. Comm. Pl. Afr. Austr. 71 (1835) = **Argyrolobium**, Eckl. & Zeyh. (Legum.).
Andrewsiana, E. Mey. l. c. 74 = A. Andrewsianum.
angustissima, E. Mey. l. c. 75 = A. filiforme.
apiculata, E. Mey. l. c. 73 = A. collinum.
argentea, E. Mey. l. c. 74 = A. Linnaeanum.
ascendens, E. Mey. l. c. 73 = A. ascendens.
baptisioides, E. Mey. l. c. 71 = A. baptisiolides.
barbata, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 77 = A. barbatum.
calycina, E. Mey. Comm. Pl. Afr. Austr. 74, adnot. = A. calycinum.
crassifolia, E. Mey. l. c. 73 = A. crassifolium.
crinita, E. Mey. l. c. 71 = A. crinitum.
cuneifolia, E. Mey. l. c. = A. polyphyllum.
diversifolia, E. Mey. l. c. = A. speciosum.
goodioides, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 75 = A. goodioides.
holosericea, E. Mey. Comm. Pl. Afr. Austr. 73 = A. sericeum.
lanceolata, E. Mey. l. c. 75 = A. lanceolatum.
longifolia, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 74 = A. longifolium.
obcordata, E. Mey. Comm. Pl. Afr. Austr. 73 = A. obcordatum.
petiolaris, E. Mey. l. c. 75 = A. petiolare.
pumila, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 77 = A. pumilum.
rupestris, E. Mey. Comm. Pl. Afr. Austr. 74 = A. rupestre.
sessiliflorum, E. Mey. l. c. 72 = A. candicans.
splendens, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 73 = A. splendens.
stricta, E. Mey. Comm. Pl. Afr. Austr. 75 = A. strictum.
tenuis, *I*. Mey. l. c. 75, adnot. = A. angustifolium.
tuberosa, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 73 = A. tuberosum.
venosa, E. Mey. Comm. Pl. Afr. Austr. 73 = A. molle.
verticillata, E. Mey. l. c. 72 = A. stipulaceum.

CHASMONIA, Presl, Fl. Sic. Praef. p. xxxvii (1826) = **Malucoclia**, Linn. (Labiat.).
incisa, Presl, l. c. = M. spicosa.

CHASSALIA, Comm. ex Poir. in Dict. Sc. Nat. viii. 198 (1817) = **Chasalia**, Comm. (Rubiac.).

CHASSELOUPIA, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 101 = **Syzygioides**, Linn. (Myrac.).
arborea, Vieill. l. c. 102 = S. arborea.
coerulescens, Vieill. l. c. 103 = S. coerulescens, Vieillardi.
gracilis, Vieill. ex Miers, in Journ. Linn. Soc. xvii. (1879) 305 = S. gracilis.
lucida, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 102 = S. glaucescens, lucida.
microphylla, Vieill. l. c. 104.—N. Caled.
montana, Vieill. l. c. = S. montana.
neo-caledonica, Vieill. l. c. 103 = S. stravadioides, Lanceraiodes.
nitida, Vieill. ex Miers, in Journ. Linn. Soc. xvii. (1879) 305 = S. rotundifolia.
tinctoria, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 104 = S. saptica.

CHASTENAEA, DC. Prod. iii. 102 (1828) = **Axinaea**, Ruiz & Pav. (Melastomac.).
affinis, Nand. in Ann. Sc. Nat. Ser. III. xvii. (1852) 130 = A. macrophylla.
coriacea, Nand. l. c. iv. (1845) 55 = A. lepidota.
floribunda, Nand. l. c. xvii. (1852) 129 = A. floribunda.
glandulosa, Nand. l. c. 129 = A. glandulosa.
grandifolia, Nand. l. c. 128 = A. grandifolia.
lanceolata, Nand. l. c. 128 = A. lanceolata.
lepidota, Benth. Pl. Hartw. 129 = A. lepidota.
longifolia, Nand. in Ann. Sc. Nat. Sér. III. iv. (1845) 55 = Marianа Karsteni.
macrophylla, Nand. l. c. = A. macrophylla.
Mertensiana, DC. Prod. iii. 103 = A. Mertensiana.
rigida, Nand. in Ann. Sc. Nat. Sér. III. xvii. (1852) 128 = Marianа rigida.

CHASTOLOMA, Lindl. Veg. Kingd. 855 (1847), sphalm. = CHARTOLOMA, Bunge = Lentis, Linn.

CHATAEA, Soland. ex Sesm. Pl. Vit. 102 (1865) = CHAITAEA, Soland. = Tacca, Forst.
Taeca, Soland. l. c. = T. pinnatifida.

CHATELANIA, Neck. Elem. i. 53 (1790) = Tolpis, Adans. (Compos.).

CHATIAKELLA, Cass. in Dict. Sc. Nat. xxix. 491 (1823) = Wulffia, Neck. (Compos.).
platyglossa, Cass. l. c. xlvi. 402 = W. stenoglossa.
stenoglossa, Cass. l. c. 408 = W. stenoglossa.

CHAUBARDIA, Reichb. f. in Bot. Zeit. x. (1852) 671 = Eygopetalum, Hook. (Orchid.).
surinamensis, Reichb. f. l. c. 672.—Guiana.

CHAULMOOGRA, Roxb. Hort. Beng. 105 (1814); Fl. Ind. iii. 855 (1832) = Gynocardia, R. Br. (Bixin.).
odorata, Roxb. ll. cc. 105, 836 = G. odorata.

CHAUNOCHITON, Benth. in Benth. & Hook. f. Gen. i. 996 (1867). *OLACINEAE*, Benth. & Hook. f. l. c.
loranthoides, *Benth. in Hook. Ic. Pl.* t. 1005.—Bras.

CHAUVINIA, Steud. Syn. Pl. Gram. 362 (1855) = Spartina, Schreb. (Gramin.).
chilensis, Steud. l. c. = S. densiflora.

CHAVANNESIA, A. DC. in DC. Prod. viii. 444 (1844) = Urocola, Roxb. (Apocyn.).
esculenta, A. DC. l. c. = U. esculenta.
javanica, Miq. Fl. Ind. Bat. ii. 458.—Java.
lucida, A. DC. in DC. Prod. viii. 444 = U. lucida.

CHAVICA, Miq. Syst. Pip. 222 (1843) = Piper, Linn.
amboinensis, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 134 = P. amboinensis.
Arnottiana, Miq. Syst. Pip. 268 = P. Hapelum.
auriculata, Miq. l. c. 269 = P. Betle.
Benthamiana, Miq. l. c. 233 = P. Barclayanum.
Betle, Miq. l. c. 238 = P. Betle.
Blumei, Miq. l. c. 246 = P. Malamiri.
bochmeriaefolia, Miq. l. c. 265 = P. bochmeriaefolium.
canaliculata, Presl, Epim. Bot. 224 = P. canaliculatum.
celebica, Miq. Syst. Pip. 276 = P. celebicum.
celtidiformis, Miq. l. c. = P. parvifolium, celtidiforme.
ceramica, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 135 = P. ceramicum.
Chaba, Miq. Syst. Pip. 251 = P. Chaba, *Blume*.
chaerostachya, Pritz. Ic. Ind. ii. 75, sphalm. = *spharostachya*.
Chawya, C. DC. in DC. Prod. xvi. 1. 351 = seq.
Chawya, Miq. Syst. Pip. 267 = P. Betle.
consanguinea, Miq. l. c. 282 = P. Labillardieri.
corylistachya, Miq. l. c. 251 = P. corylistachyon.
densa, Miq. l. c. 253 = P. Betle.
diffusa, Miq. in Zoll. Syst. Verz. Ind. Archip. 84 = P. attenuatum.
exasperata, Miq. Syst. Pip. 275 = P. exasperatum.
frustrata, Miq. Fl. Jungh. 16.—Java.
glandulosa, Presl, Epim. Bot. 223 = P. miniatum.
grandis, Presl, l. c. 224 = P. grande.
Guiguai, Miq. Syst. Pip. 250 = P. brachystachyum, *Wall.*
Haenkeana, Presl, Epim. Bot. 224 = P. Haenkeanum.
Hasseliana, Miq. in Linnaea, xxvi. (1855) 218 = P. Novae-Hollandiae vel P. Hasselianum.
imperialis, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 134 = P. Forsteni.
Labillardieri, Miq. Syst. Pip. 263 = P. Chaba, *Hunter.*
lanceolata, Miq. l. c. 264 = P. miniatum.
leptostachya, Hance, in Journ. Bot. vi. (1868) 301.—China.
Leuconeura, Miq. Syst. Pip. 270 = P. Pseudo-chavica.
macrostachya, Miq. l. c. 236 = P. miniatum.
maculata, Miq. l. c. 256 = P. maculatum.
madagascariensis, Miq. l. c. 277 = P. madagascariense.
majuscula, Miq. l. c. 271 = P. majusculum.
maritima, Miq. l. c. 262 = P. Chaba, *Hunter.*
Melamiris, Miq. l. c. 245 = P. Sirium.
microstigma, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 138 = P. microstigma.
miniata, Miq. Syst. Pip. 234 = P. miniatum.
CHAVICA:—
Mullesua, Miq. Syst. Pip. 280 = Piper brachystachyum, *Wall.*
multinervia, Miq. Fl. Ind. Bat. Suppl. 478 = P. prismamense.
Neesiana, Miq. Syst. Pip. 249 = P. peepuloides.
officinarum, Miq. l. c. 236 = P. Chaba, *Hunter.*
parvifolia, Hassk. in Flora, xlvii. (1864) 59 = P. parvifolium.
peepuloides, Miq. Syst. Pip. 237 = P. peepuloides.
peepuloides, Wight, Ic. t. 1937 = P. Chaba, *Hunter.*
penangensis, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 483 = P. Lonchites.
penangensis, Miq. Syst. Pip. 270 = P. penangense.
petiolata, C. DC. in DC. Prod. xvi. 1. 389.—Ind. or.
populifolia, Miq. Syst. Pip. 248.—Ins. Philipp.
pothoidea, Miq. l. c. 266 = P. pothoideo.
pseudomelamiris, Miq. l. c. 244 = P. Sirium.
puberula, Benth. Fl. Hongk. 335.—China.
pyrifolia, Miq. Syst. Pip. 275 = P. pyrifolium.
radicans, Presl, Epim. Bot. 224 = P. radicans.
Rafflesii, Miq. Syst. Pip. 273 = P. Rafflesii.
Reinwardtiana, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 136 = P. Reinwardtianum.
retrofracta, Miq. Syst. Pip. 275 = P. retrofractum.
Roxburghii, Miq. l. c. 239 = P. longum.
rufinervis, Presl, Epim. Bot. 224 = P. rufinerve.
Rumphii, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 141 = P. Rumphii.
salicina, Presl, Epim. Bot. 224 = P. salicinum.
sarmentosa, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 483 = P. longum.
sarmentosa, Miq. Syst. Pip. 242 = P. sarmentosum.
sinensis, Champ. ex Benth. in Hook. Kew Journ. vi. (1854) 118 = P. sinense.
Siriboa, Miq. Syst. Pip. 224 = P. Betle.
spharostachya, Miq. l. c. 278 = P. brachystachyum, *Wall.*
subcordata, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 138 = P. subcordatum.
subseptemnervia, Miq. Fl. Ind. Bat. Suppl. 473 = P. subseptemnervium.
Suipigua, Miq. Syst. Pip. 275 = P. Suipigua.
sundaica, Miq. l. c. 274 = P. sundaicum.
sylvatica, Miq. l. c. 248 = P. sylvaticum.
syringaefolia, Miq. l. c. 275 = P. syringaefolium.
Teysmanni, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 141 = P. Teysmanni.
thermalis, Miq. Syst. Pip. 276 = P. thermale.
Thomsonii, C. DC. in DC. Prod. xvi. 1. 389.—Ind. or.
viminalis, Miq. in Linnaea, xx. (1847) 181 = P. viminale.
Wallichii, Miq. Syst. Pip. 254 = P. aurantiacum.

CHAVOTA, Jacq. Am. Select. ed. pict. t. 245 (1780) = Sechium, P. Br. (Cucurbitac.).
edulis, Jacq. l. c. = S. edule.

CHEILANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 (genus index. *LABIATARUM*).

CHEILOCLINIUM, Miers, in Trans. Linn. Soc. xxviii. (1872) 430. *HIPPOCRATEACEAE.*
anomalum, *Miers, l. c.*—Bras.

CHEILOCOCCA, Salisb. ex Sm. Spec. Bot. N. Holl. 17 (1793), cum cit. falsa = Platylobium, Sm. (Legumin.).
apocynifolia, Salisb. ex Sm. l. c. = P. formosum.

CHEILODISCUS, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 36 = Proetia, Linn. (Compos.).
littoralis, Triana, l. c.—N. Granat.

CHEILOPSIS, Moq. in Ann. Sc. Nat. Sér. I. xxvii. (1832) 230 = Acanthus, Tourn.
arborea, Nees, in DC. Prod. xi. 273 = A. arborea.
montana, Nees, l. c. = A. montanus.
palystachya, Moq. in Ann. Sc. Nat. Sér. I. xxvii. (1832) 230 = A. arborea.

CHEILOSA, Blume, Bijdr. 613 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 322.
montana, *Blume*, l. c. 614.—Java.

CHEILOSANDRA, Griff. ex Lindl. Veg. Kingd. 672 (1847) = Rhynchotechum, Blume (Gesneriac.).
leptopus, Griff. ex Benth. & Hook. f. Gen. iii. 1025 = Leptobaea glabra.

CHEILOTHECA, Hook. f. in Benth. & Hook. f. Gen. iii. 607 (1876). *MONOTROPEAE*, Benth. & Hook. f. l. c.
khasiana, *Hook. l. c.* 608.—Reg. Himal.

CHEILYCTIS, Benth. Lab. Gen. et Sp. 736 (1836) = Monarda, Linn. (Labiat.).

CHEIRADENIA, Lindl. Fol Orch. Cheirad. (1858) 1. *ORCHIDEAE*, Benth. & Hook. f. iii. 544.
cuspidata, *Lindl. l. c.*—Guiana.

CHEIRANTHERA, Endl. Enchir. 518 (1841) = Cheiro stemon, Humb. & Bonpl. (Sterculiac.).

CHEIRANTHERA, Bronga Voy. Coq. Bot. t. 77 (1826); et A. Cunn. in Bot. Reg. sub t. 1719 (1834). *PITTOSPOREAE*, Benth. & Hook. f. i. 133.
brevifolia, F. Muell. Fragm. i. 97 = filifolia.
cyanea, Bronga. Voy. Coq. Bot. t. 77 = linearis.
filifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 364.—Austral.
linearis, *A. Cunn. in Bot. Reg.* sub t. 1719.—Austral.
parviflora, *Benth. Fl. Austral.* i. 136.—Austral.
Preissiana, *Putterl. in Lehm. Pl. Preiss.* i. 201 (sp. dub.).—Austral.
tortilis, F. Muell. Fragm. ii. 79 = filifolia.
volubilis, *Benth. Fl. Austral.* i. 136.—Austral.

CHEIRANTHODENDRON, Benth. & Hook. f. Gen. i. 212 (1862) = CHIRANTHODENDRON, Cerv. = Cheirostemon, Humb. & Bonpl. (Sterculiac.).

CHEIRANTHODENDRUM, Steud. Nom. ed. I. 186 (1821) = prec.

CHEIRANTHUS, Linn. Gen. ed. I. 195 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 68.
CHEIRI, Adans. Fam. ii. 418 (1763); [Clairv.] Man. Herb. 221 (1811).
DICHROANTHUS, Webb & Berth. Phytog Canar. i. 65. t. 5, 6 (1836).
PHORNICAULIS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 89 (1838).
SCHELHAMERIA, Holst. Syst. 9 (1746).
acaulis, Balb. ex Spreng. Nov. Prov. 10 = Matthiola acaulis.
africanus, Linn. Pl. Afr. 18 (Amoen. Acad. vi. 90) = Hellophila pilosa.
alidifolius, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 133.—Tibet. occ.
alpus, Mill. Gard. Dict. ed. VIII. n. 8 = Matthiola incana.
alpinus, All. Fl. Pedem. iii. t. 20 = Erysimum lanceolatum.
alpinus, Falk, Beitr. ii. 216 = Erysimum altaicum.
alpinus, Jacq. Fl. Austr. i. 48. t. 75 = Erysimum canescens.
alpinus, Lam. Encyc. ii. 716 = Erysimum ochroleucum.
alpinus, *Linn. Mant.* 93 (excl. syn.).—Scandinav.
angulatus, Schult. Obs. 129 = Syrenia sessiliflora.
angustifolius, Gilib. Fl. Lituan. ii. 61 = Erysimum hieraciifolium.
angustifolius, Mill. Gard. Dict. ed. VIII. n. 4 = Erysimum lanceolatum.
annuus, Georgi, Reise, i. 235 = Hesperis aprica.
annuus, *Linn. Sp. Pl.* 663.—Davuria.
apricus, Steph. ex Willd. Sp. Pl. iii. 518 = Hesperis aprica.
arenicolus, Lejeune, Fl. Spa, ii. 68 = Erysimum cheiranthoides.
arborescens, Sieber, ex Link, Enum. Hort. Berol. ii. 166, nomen = Matthiola arborescens.
arborescens, *Sweet, ex Steud. Nom.* ed. II. i. 245.—Aegypt.
arbuscula, *Lowe, in Hook. New Journ.* viii. (1856) 289.—Ins. Canar.
arenarius, Suffren, ex DC. Syst. ii. 171 = Matthiola varia.
armeniacus, Sims, Bot. Mag. t. 835 = Erysimum Ibericum.
arvensis, Bernh. Syst. Verz. Erf. 196 = Erysimum hieraciifolium.
asper, *Chem. & Schlecht. in Linnaea*, i. (1826) 14.—Am. bor. occ.
asper, Nutt. Gen. Am. ii. 436 = Erysimum asperum.
asperrimus, *Bunge, Enum. Pl. Chin. Bor.* 8.—China.
aureus, Willd. ex DC. Syst. ii. 561 = Stenopus elvchrysifolium.

CHEIRANTHUS :—

auriculatus, Lapeyr. Hist. Abr. Pl. Pyr. 362 = Cochlearia auriculata.
axillaris, Bromf. ex Webb & Berth. Phyt. Canar. i. 67 = scoparius.
bicolor, Hornem. Hort. Hafn. ii. 613 = Erysimum bicolor.
bicornis, Sibth. & Sm. Fl. Graec. Prod. ii. 26 = Matthiola bicornis.
bicuspidatus, Willd. Sp. Pl. iii. 519 = Hesperis bicuspidata.
bithynicus, Pers. Syn. ii. 200 = Erysimum cuspidatum, *Bocconi*, All. Fl. Pedem. i 272 = Erysimum canescens.
callosus, Linn. f. Suppl. 296 = Heliophila callosa.
canus, Piller & Mitterp. Iter Posseg. 146. t. 15 = Erysimum angustifolium.
capitatus, Dougl. in Hook. Fl. Bor. Am. i. 38 = asper.
carnosus, Thunb. Prod. Pl. Cap. 108 = Heliophila platyuliqua.
caspicus, Lam. in Pall. Voy. 8° App. 348 = Sterigma tomentosum.
Cheiri, *Linn. Sp. Pl.* 661.—Europ.
chius, Linn. l. c. = Malcolmia chia.
chius, Pall Reise, i. 497 = Chorispora tenella.
cinereus, *Webb & Berth. Phyt. Canar.* i. t. 5.—Ins. Canar.
coccineus, Mill. Gard. Dict. ed. VIII. n. 7 = Matthiola incana.
collinus, Bieb. Fl. Taur. Cauc. ii. 119 = Erysimum collinum.
contortuplicatus, Steph. in Willd. Sp. Pl. iii. 521 = Sisymbrium contortuplicatum.
corinthius, *Boiss. Diagn.* Ser. II. i. 18.—Graecia.
cornutus, Lam. Encyc. ii. 717 = Syrenia sessiliflora.
coronopifolius, Sibth. & Sm. Fl. Graec. Prod. ii. 25 = Matthiola tristis.
crassicaulis, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 45 = Erysimum crassicaule.
Cumbrae, Link, in Buch, Beschr. Canar. Ins. 152 = scoparius.
cuspidatus, Bieb. Fl. Taur. Cauc. ii. 120 = Erysimum cuspidatum.
decumbens, Schleich. ex Willd. Enum. Hort. Berol. 680 = ochroleucus.
Delilianus, *A. Gouanlt, in Rev. Hortic.* Sér. IV. i. (1852) 401.—Hab.?
denticulatus, Willd. ex Steud. Nom. ed. II. i. 346 = Pallasii.
dubius, Hornem. Hort. Hafn. Suppl. 73 = Erysimum dubium.
dubius, Suter, Fl Helv. ii. 65 = ochroleucus.
elongatus, Thunb. Prod. Pl. Cap. 108 = Heliophila linearifolia
erucaefolius, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 798 = Malcolmia lacera.
erysimoides, Gueldenst. ex Ledeb. Fl. Ross. i. 188 = Erysimum leptostylum.
erysimoides, Huds. Fl. Augl. ed. II. 287 = Erysimum cheiranthoides.
erysimoides, Linn. Sp. Pl. 661 = Erysimum lanceolatum.
erysimoides, Pall. Reise, i. 192 = Erysimum Andrzeiskianum.
erysimoides, Vill. ex Willk. & Lange, Prod. Fl. Hisp. iii. 806 = Erysimum ochroleucum.
Farsetia, Bech.-Mann. in Wall. Cat. n. 4801 = Farsetia Hamiltenii.
Farsetia, Linn. Mant. i. 94 = Farsetia aegyptiaca.
femstralis, Linn. Sp. Pl. 1198 = Matthiola fenestralis.
firmus, Schleich. ex Willd. Enum. Hort. Berol. 45 = Erysimum lanceolatum ?
firmus, Willd. Enum. Hort. Berol. Suppl. 45 = Erysimum virgatum.
flexuosus, *Sibth. & Sm. Fl. Graec. Prod.* ii. 24; *Fl. Graec.* t. 634.—Ins. Cyprus.
fragrans, Fisch. Hort. Gorensk. (1812) 51 = Matthiola fragrans.
frutescens, Vent. ex Pers. Syn. ii. 201 = semperflorens.
frutescens, L'Hérit. ex DC. Prod. i. 187 = C. Cheiri.
fruticulosus, Georgi, Beschr. Russ. Reich. iii. 77. 1124 = Sterigma tomentosum.
fruticulosus, L'Hérit. ex DC. *Syst.* ii. 498 = Erysimum suffruticosum.
fruticulosus, Linn. Mant. 94 = C. Cheiri.
fruticulosus, Linn. *Sp. Pl.* 663 = Matthiola tristis.
fruticulosus, Sibth. & Sm. Fl. Graec. Prod. ii. 26 = Erysimum Boryanum.

CHEIRANTHUS :—

glaber, Mill. Gard. Dict. ed. VIII. n. 9 = Matthiola glabra.
glaberrimus, Colla, Antol. Bot. v. 861 = Matthiola glabra.
graecus, [Juss.] Pers. Syn. ii. 201 = Matthiola incana.
gramineus, Thunb. Prod. Pl. Cap. 108 = Heliophila gramines.
Griffithii, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 137 = Erysimum Hookeri.
helveticus, Jacq. Hort. Vindob. iii. 9 = Erysimum murale.
heseperidoides, *Torr. & Gray, Fl. N. Am.* i. 72.—Am. bor.
hieraciifolius, Lam. Encyc. ii. 717 = Erysimum hieraciifolium.
hilaris, Willd. ex Steud. Nom. ed. II. i. 346 = Matthiola tristis.
himalaicus, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 137 = himalayensis.
himalayensis, *Cambess. in Jacquem. Voy. Bot.* 14.—Reg. Himal
hirtus, Schleicht. fide Stev. ex DC. Prod. i. 189 = Hesperis aprica.
hispidus, Adams, in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 250 = Hesperis aprica.
hortensis, Lam. Fl. Fr. ii. 506 = Matthiola incana.
humilis, Clarke, Trav. iii. 42 = Matthiola humilis.
ibericus, Adams, in Weber & Mohr, Beitr. i. 61 = Erysimum ibericum.
ibericus, Willd. Enum. Hort. Berol. 681 = Barbarea arcuata.
incanus, Linn. Sp. Pl. 662 = Matthiola incana.
incrassatus, Bory, ex Steud. Nom. ed. II. i. 346 = Malcolmia flexuosa.
integrifolius, Mill. Gard. Dict. ed. VIII. n. 2 = Erysimum canescens.
integerrimus, Stend. Nom. ed. II. i. 346 = Erysimum canescens.
intermedius, Schleich. Cat. Pl. Helv. ed. IV. 12 = C. Cheiri?
Jacquinii, Clem. Sert. Olymp. 9. t. 4. f. 2 = Erysimum Kotschyanum.
junceus, Kit. Pl. Rar. Hung. iii. 259 = Sisymbrium junceum.
Kevii, Nest. Delic. Gallo-Belg. i. 288 = C. Cheiri.
Knowleanus, D. Distr. Syn. Pl. iii. 688 = ochroleucus.
Kotschyi, Walp. Rep. i. 195 = ochroleucus.
lacerifolius, Pall. Reise Statth. Russ. Reich. i. 124 = Matthiola tatarica.
lacerus, Gouan, Illustr. 44 = Malcolmia parviflora.
incerus, Linn. Sp. Pl. 662 = Malcolmia lacera.
laciniatus, Poir. Encyc. Suppl. ii. 780 = Hesperis laciniata.
laevigatus, Willd. ex DC. Prod. i. 141 = Barbarea arcuata.
lanatus, Gmel. ex DC. Syst. ii. 581 = Sterigma elychrysifolium.
lanceolatus, Willd. Sp. Pl. iii. 515 = Hesperis tristis.
leptophyllus, Bieb. Fl. Taur. Cauc. ii. 119 = Erysimum leptophyllum.
leptophyllus, Willd. ex DC. Syst. ii. 464 = Sisymbrium junceum.
leucanthemus, Steph. ex Willd. Sp. Pl. iii. 521 = Erysimum versicolor.
linariifolius, *Vis. in Nuov. Sagg. Acc. Pad.* v. (1840) 267.—Afr. bor.
linearis, Forsk. Fl. Aegypt. Arab. 119 = Farsetia aegyptiaca.
linearis, Thunb. Prod. Fl. Cap. 108 = Heliophila linearis.
linearis, Vent. Jard. Malm. t. 83 = semperflorens.
linifolius, Pers. Syn. ii. 201 = Erysimum linifolium.
litterosus, All. Fl. Pedem. i. 273 = Malcolmia maritima.
littoreus, Linn. Sp. Pl. ed. II. 925 = Malcolmia littorea.
littoreus, Pall. Reise, ii. 741 = Sterigma tomentosum.
lividus, Delile, Fl. Aeg. Illustr. 67 = Matthiola livida.
loddus, Sieber, ex Steud. Nom. ed. II. i. 346 = Malcolmia aegyptiaca.
longifolius, Vent. Jard. Malm. t. 83 = mutabilis.
longifolius, Vent. Jard. Cels, t. 93 = Matthiola oxyceras.
longisiliquus, Steud. Nom. ed. II. i. 346 = Erysimum longisiliquosum.
lotosus, Dahe, Fl. Hantes-Pyr. 199 = C. Cheiri.

CHEIRANTHUS :—

lyratus, Sibth. & Sm. Fl. Graec. Prod. ii. 24 = Malcolmia lyrata.
maritimus, Linn. Cent. Pl. i. 19 (Amoen. Acad. iv. 280) = Malcolmia maritima.
Marschallii, Stark, ex Moore, Gard. Mag. Bot. i. (1850) 289 = ochroleucus.
Menziesii, *Benth. & Hook. f. Gen.* i. 68.—Am. bor. occ.
mollis, Hornem. Hort. Hafn. ii. 615 = Arabis albida.
montanus, Pall. Reise, i. 496 = Syrenia sifiliculosa.
muralis, Salisb. Prod. 371 = C. Cheiri.
muricatus, Lam. Fl. Fr. ii. 507 = Matthiola sinuata.
muricatus, Welnm. Cat. Hort. Dorpat. (1810) 41 = Dontostemon integrifolius.
mutabilis, Brouss. ex Spreng. Syst. ii. 896 = semperflorens.
mutabilis, *L'Hérit. Stirp. Nov.* 92.—Ins. Mader. et Canar.
nanus, Merk. ex DC. Syst. ii. 486 = Matthiola nana.
nitrarius, *Pall. Reise Statth. Russ. Reich.* i. 108, 124, 141.—Rossia.
nodosus, Lam. Fl. Fr. ii. 508 = Matthiola littorea.
ochroleucus, DC. Syst. ii. 181 = Erysimum ochroleucum.
ochroleucus, *Hall. f. ex Schleich. Cat.* 16.—Europ.
odoratissimus, Bieb. Casp. 116 = Matthiola odoratissima.
odoratus, *Pall. ex Spreng. Syst.* ii. 896.—Persia.
olympicus, *Sibth. ex DC. Syst.* ii. 494.—Graecia.
Pallasii, Pursh, Fl. Am. sept. ii. 486 = pygmaeus.
pallens, Hall. f. ex DC. Syst. ii. 501 = Erysimum ochroleucum.
pallidus, Pall. Reise, iii. 280 = Erysimum canescens.
paniculatus, Cav. Desc. 430 = Erysimum Kunzeanum.
paniculatus, Lam. Encyc. ii. 717 = Erysimum repandum.
paniculatus, Stev. ex DC. Syst. ii. 210 = Arabis incarnata.
Parnassii, Bolss. Diagn. Ser. II. i. 18 = Erysimum Boryanum.
parryoides, *Kurz, ex Hook. f. Fl. Brit. Ind.* i. 182.—Tibet.
parviflorus, Schousb. Vextr. Marokko, 195 = Matthiola parviflora.
parviflorus, Thib. ex DC. Prod. i. 184 = Matthiola coronopifolia.
persicus, Pall. ex DC. l. c. 182 = Anchonium Tournefortii.
persicus, Willd. ex Ledeb. Fl. Ross. i. 109 = Matthiola tatarica.
pinnatifidus, Willd. Sp. Pl. iii. 528 = Dontostemon pectinatus.
planifolius, Waldst. & Kit. ex Bieb. Fl. Taur. Cauc. iii. 448 = Erysimum cuspidatum.
pulchellus, Willd. Sp. Pl. iii. 533 = Erysimum rupestre.
Pumilio, Sibth. & Sm. Fl. Graec. Prod. ii. 26 = Matthiola bicornis.
pumilus, Hornem. Hort. Hafn. ii. 613 = Erysimum pumilum.
pygmaeus, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 114.—Sibir.; Am. bor. occ.
quadrangularis, Stend. Nom. ed. II. i. 346 = seq.
quadrangulus, L'Hérit. Stirp. Nov. 91. t. 44 = Syrenia sessiliflora.
ramosissimus, Lam. Encyc. ii. 717 = Erysimum repandum.
ramosissimus, *Bory & Chaub. Nouv. Fl. Pélop.* 43.—Graecia.
rhaeticus, Schleich. Cat. Pl. Helv. ed. III. 12 = Erysimum ochroleucum.
ruderalis, Pall. ex DC. Syst. ii. 580 = Sterigma tomentosum.
rupestris, Sibth. & Sm. Fl. Graec. Prod. ii. 28; Fl. Graec. t. 633 = Erysimum rupestre.
salinus, *Linn. Mant.* 93.—Sibir.; Tatr.
salinus, Pall. ex DC. Syst. ii. 580 = Sterigma tomentosum.
salinus, Willd. ex DC. l. c. 210 = Arabis incarnata.
scaber, Mnench, Meth. 264 = Malcolmia africana.
scaber, *Webb & Berth. Phyt. Canar.* i. 68.—Ins. Canar.
scoparius, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 113 = Parrya macrocarpa.
scopiurus, Willd. Prod. Fl. Berol. 214. t. 5. f. 10 = Erysimum cheiranthoides.
scoparius, *Brouss. ex Willd. Enum. Hort. Berol.* 681. —Ins. Canar.

CHEIRANTHUS :—
semperflorens, Cons. & Der. in Bull. Soc. Bot. Fr. iv. (1857) 806 = Erysimum longifolium.
semperflorens, *Schmalh. Vestn. Moroškin*, 195. — Marocco.
Sancouri, *Heldr. & Sart. Cat. Hort. Athen.* (1862) *ex Rend. Cat. Gram. Genève* (1866), 4.—Archip. Graec.
sibiricus, Jusa. ex DC. Syst. ii. 492 = Erysimum angustifolium.
silenifolius, Willd. Enum. Hort. Berol. Suppl. 45 = Erysimum murale.
silicularoides, Bieb. Fl. Taur. Cauc. ii. 121 = Syrenia silicifolia.
sinuatus, Linn. Sp. Pl. ed. II. 926 = Matthiola sinuata.
sinuatus, Pall. Reise, ii. 229 = Sterigma tomentosum.
Stewartii, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 132. —Reg. Himal.
Stochsianus, Boiss. Diagn. Ser. II. i. 19 = Erysimum Stocksianum.
strictus, Linn. f. Suppl. 296 = Heliophila callosa.
strictus, Poir. Encyc. Suppl. ii. 781 = Heliophila scoparia.
strigosus, Ledeb. in Mém. Acad. Pétersb. v. (1812) 589 = Erysimum strigosum.
subulatus, Moench, Meth. 264 = Malcolmia chia.
sulphureus, [Soland. in] Russ. Aleppo, ed. II. ii. 257 = Sterigma sulphureum.
sylvestris, Bernh. ex Stend. Nom. ed. II. i. 346 = Erysimum hieracifolium.
sylvestris, Crantz, Stirp. Austr. ed. I. i. 48 ; Lam. Encyc. ii. 716 = Erysimum lanceolatum.
syriacus, DC. Syst. ii. 185 = Coringia orientalis.
syriacus, Willd. ex Stend. Nom. ed. II. i. 346 = Matthiola oxyoceras.
taraxacifolius, Balb. Cat. Hort. Taur. App. (1814) 10 = Malcolmia taraxacifolia.
taraxacifolius, Biasol. ex Reichb. Fl. Germ. Excurs. 687 = Erysimum strictum.
taraxacifolius, Schultes, in Denkschr. Bot. Ges. Regensb. ii. (1816) 164 = Chorispora tenella.
taraxacifolius, *Steph. ex Willd. Sp. Pl.* iii. 525.— Sibir.
tataricus, Pall. ex DC. Syst. ii. 170 = Matthiola tatarica.
tauricus, Lepech. in Nov. Act. Petrop. xiii. (1802) 386. t. 9 = Matthiola odoratissima.
tonsilifolius, *L'Hérit. Stirp. Nov.* 92.—Ins. Madera.
tomentosus, Willd. Sp. Pl. iii. 523 = Sterigma tomentosum.
torulosus, Bieb. Fl. Taur. Cauc. ii. 121 = Sterigma torulosum.
torulosus, Thunb. Prod. Pl. Cap. 108 = Matthiola torulosa.
tricuspidatus, Hads. Fl. Aagl. ed. I. 250 = Matthiola sinuata.
tricuspidatus, Linn. Sp. Pl. 663 = Matthiola tricuspidata.
trilobus, Linn. l. c. 663 = Malcolmia littorea.
tristis, Curt. Bot. Mag. t. 739 = Matthiola varia.
tristis, Forsk. Fl. Aegypt. Arab. 119 = Matthiola livida.
tristis, Linn. Syst. Nat. ed. X. 1134 = Matthiola tristis.
turritoides, Lam. Encyc. ii. 716 = Erysimum cheiranthoides.
uncinatus, Willd. ex Stend. Nom. ed. II. i. 347 = Malcolmia littorea.
valleianus, J. *Gay*, ex DC. Prod. i. 124 = Matthiola varia.
varius, Sibth. & Sm. Fl. Graec. Prod. ii. 25 ; Fl. Graec. t. 696 = Matthiola varia.
versicolor, Bieb. Fl. Taur. Cauc. ii. 119 = Erysimum versicolor.
versicolor, Pall. ex Stend. Nom. ed. II. i. 347 = Hesperis hyrcanica.
villosus, Forsk. Fl. Aegypt. Arab. 120 = Matthiola tricuspidata.
villosus, Spreng. ex DC. Prod. i. 186 = Hesperis laciniata.
virgatus, Poir. ex DC. Syst. ii. 462 = Sisymbrium junceum.
virgatus, Poir. Encyc. Suppl. ii. 781 = Erysimum angustifolium.
viridiflorus, Stend. Nom. ed. II. i. 347 = Matthiola glabra.
viridis, Ehrh. Beitr. vii. 158 = Matthiola incana.
CHEIRI, Adans. Fam. ii. 418 (1763) = **Cheiranthus**, Linn. (Crucif.).
hieracifolium, [Clairv.] Man. Herb. 221 = Erysimum hieracifolium.
montanum, [Clairv.] l. c. = Cheiranthus Cheiri.
vulgare, [Clairv.] l. c. = Cheiranthus Cheiri.
CHEIRINIA, Link, Enum. Hort. Berol. ii. 170 (1822) = **Erysimum**, Tourn. (Crucif.).
altissima, Link, l. c. 171 = E. hieracifolium.
Andreæsowiana, Hort. Berol. ex Stend. Nom. ed. II. i. 347 = E. canescens.
angustifolia, Link, Enum. Hort. Berol. ii. 170 = E. angustifolium.
canescens, Link, l. c. 171 = E. canescens.
cheiranthoides, Link, l. c. 170 = E. cheiranthoides.
crepidifolia, A. Dietr. Fl. Borus. vii. 446 = E. crepidifolium.
cuspidata, Link, Enum. Hort. Berol. ii. 170 = E. cuspidatum.
erysimoides, Dum. Fl. Belg. 123 = E. lanceolatum.
firma, Link, Enum. Hort. Berol. ii. 171 = E. virgatum.
helvetica, Link, l. c. = E. ochroleucum.
hieracifolia, Link, l. c. 170 = E. hieracifolium.
juncea, Link, l. c. 172 = Sisymbrium junceum.
lanceolata, Link, l. c. 171 = E. lanceolatum.
leptostylis, Link, Handb. ii. 215 = E. leptostylum.
longifolia, Link, Enum. Hort. Berol. ii. 171 = E. longifolium.
longisiliqua, Link, Handb. ii. 215 = E. virgatum.
longisiliquosa, Link, Enum. Hort. Berol. ii. 170 = E. longisiliquosum.
odorata, Link, Handb. ii. 215 = E. odoratum.
repanda, Link, Enum. Hort. Berol. ii. 171 = E. repandum.
sessiliflora, Link, l. c. 170 = Syrenia sessiliflora.
stricta, Link, l. c. = E. hieracifolium.
strictissima, Link, l. c. 171 = Sisymbrium strictissimum.
suffruticosa, Link, l. c. 170 = E. murale.
virgata, Link, l. c. = E. virgatum.
CHEIRISANTHERA, Hort. ex Regel, Gartenfl. ii. (1853) 353 = **Hoppiella**, Regel (Gesneriac.).
atrosanguinea, Hort. ex Regel, l. c. = H. atrosanguinea.
coccinea, Hort. ex Regel, l. c. = H. atrosanguinea.
CHEIRODENDRON, Nutt. ex Seem. in Journ. Bot. v. (1867) 286 = **Panax**, Linn. (Araliac.).
Gaudichaudii, Seem. l. c. = P. Gaudichaudii.
laxiceratum, Seem. l. c.—Chili.
platyphyllum, Seem. l. c. = P. platyphyllum.
samoense, Seem. l. c. 287.—Samoa.
valdivianum, Seem. l. c. 286.—Chili.
CHEIROLAENA, Benth. in Benth. & Hook. f. Gen. i. 232 (1862). *STERCULIACEAE*, Benth. & Hook. f. l. c.
Ilondii, *Benth. c.*—Madag.
CHEIROLEPIS, Boiss. Diagn. Ser. I. x. 106 (1849) = Centaurea, Linn. (Compos.).
cappadocica, Boiss. l. c. 107 = Cent. cappadocica.
drabifolia, Boiss. l. c. = Cent. drabifolia.
Kotschyi, Boiss. & Heldr. l. c. = Cent. macrolopha.
libanotica, Boiss. l. c. (= *Phanopappus libanoticus*).— Syria.
persica, Boiss. l. c. 108 = Cent. macrolopha.
CHEIROLOMA, F. Muell. in Linnæa, xxv. (1852) 401 = **Calotis**, R. Br. (Compos.).
hispidulum, F. Muell. l. c. = Cal. hispidula.
CHEIROLOPHUS, Cass. in Dict. Sc. Nat. l. 290 (1827) = Centaurea, Linn. (Compos.).
lanceolatus, Cass. l. c. ii. 96 = Cent. sempervirens.
pinnatifidus, Cass. l. c. = Cent. intybacea.
sempervirens, Pomel, Nouv. Mat. Fl. Atl. 29 = Cent. sempervirens.
CHEIROPETALUM, E. Fries, Ind. Sem. Hort. Upsal. (1857) ex Urban, Addit. Ind. Sem. Hort. Berol. (1861) 11 = Silene, Linn. (Caryoph.).
radians, E. Fries, l. c. = S. odontopetala.
CHEIROPSIS, Bercht. & Presl, Rostl. i. Rannuculus. 11 (1832) = **Clematis**, Tourn. (Ranunculac.).
balearica, Bercht. & Presl, l. c. = Clem. balearica.
cirrhosa, Bercht. & Presl, l. c. = Clem. cirrhosa.
CHEIROPSIS :—
elegans, Spach, Hist. Vég. Phan. vii. 261 = Clematis cirrhosa.
semitriloba, Bercht. & Presl, Rostl. i. Ranunculac. 11 = Clem. semitriloba.
CHEIROPTEROCEPHALUS, Rodrig. Orch. Nov. i. 28 (1877). *ORCHIDEAE*.
sertuliferus, *Rodrig. l. c.* 29.—Bras.
CHEIROSTEMON, Humb. & Bonpl. Pl. Aequin. i. 81. t. 24 (1808). *STERCULIACEAE*, Benth. & Hook. f. i. 212, 232.
CHEIRANTHODENDRON, Cerv. ex Cav. in Anal. Clenc. Nat. vi. (1803) 308, adnot.
aspetalum, Cerv. in Roem. Collect. Bot. 97 = platanoides.
platanoides, *Humb. & Bonpl. Pl. Aequin.* 82. t. 24.— Mexic.
CHEIROSTYLIS, Blume, Bijdr. 413. t. 16 (1826). *ORCHIDEAE*, Benth. & Hook. f. iii. 600.
cochinchinensis, *Blume, Orch. Archip. Ind.* t. 39. f. 2.—Cochinch.
fabellata, *Wight, Ic.* v. 16. (t. 1797, Monochilus).— Ind. or.
grandiflora, *Blume, Orch. Archip. Ind.* 55.—N. Guinea.
Griffithii, *Lindl. in Journ. Linn. Soc.* i. (1857) 188.— Reg. Himal. ; Burma.
gymnochiloides, *Reichb. f. in Flora*, lxviii. (1885) 537. —Madag.
heterosepala, *Reichb. f. Otia Bot. Hamb.* 110.—Afr. trop.
Humblotii, *Reichb. f. in Flora*, lxviii. (1885) 537.— Ins. Comoro.
malleifera, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 141.—Burma.
marmorata, Lindl. ex Lem. in Fl. des Serres, Sér. 1. iv. (1848) t. 370 = Dossinia marmorata.
montana, *Blume, Bijdr.* 413.—Java.
parvifolia, *Lindl. Bot. Reg.* (1839) *Misc.* 19.— Zeylan.
pubescens, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 141.—Burma.
pusilla, *Lindl. Gen. et Sp. Orch.* 489.—Reg. Himal.
CHELIDONIUM, Tourn. ex Linn. Syst. ed. I (1735). *PAPAVERACEAE*, Benth. & Hook. f. i. 58.
asiaticum, Sadeb. Prod. 377 = Glaucium corniculatum.
corniculatum, Donn, Hort. Cant. 100 = Glaucium flavum.
corniculatum, Linn. Sp. Pl. 506 = Glaucium corniculatum.
dahuricum, Hort. ex DC. Prod. i. 128 = majus.
diphyllum, Michx. Fl. Bor. Am. i. 309 = Stylophorum diphyllum.
dodecandrum, Forsk. Fl. Aegypt. Arab. 100 = Roemeria orientalis.
fulvum, Poir. Encyc. Suppl. v. 606 = Glaucium flavum.
glaberum, Mill. Gard. Dict. ed. VIII. n. 5 = Glaucium corniculatum.
Glaucium, Linn. Sp. Pl. 506 = Glaucium flavum.
glaucum, Hill, Veg. Syst. xi. 16 = Glaucium flavum.
grandiflorum, DC. Prod. i. 128 = majus.
hamatodes, Moench, Meth. 249 = majus.
hybridum, Linn. Sp. Pl. 506 = Roemeria hybrida.
japonicum, Thunb. Fl. Jap. 231 = Stylophorum japonicum.
laciniatum, Mill. Gard. Dict. ed. VIII. t. 9 = majus.
littorale, Salisb. Prod. 377 = Glaucium flavum.
luteum, Gilib. Fl. Lituan. ii. 211 = majus.
majus, *Linn. Sp. Pl.* 505.—Europ. ; As. Min.
majus, Lour. Fl. Cochinch. i. 290 = sinense.
multifidum, Moç. & Sesse, ex DC. Prod. iii. 344 = Eschscholtzia californica.
murale, Sen. Fl. Oxon. 180 = majus.
pedatifidum, Spreng. ex Steud. Nom. ed. II. i. 347 = Stylophorum diphyllum.
phœniceum, Lam. Fl. Fr. iii. 189 = Glaucium phœniceum.
quercifolium, Willemet, Phyt. Encyc. ii. 613 = majus.
rubrum, Poir. Encyc. Suppl. v. 606 = Glaucium corniculatum.
rudersale, Salisb. Prod. 377 = majus.
sinense, *DC. Syst.* ii. 106 (sp. dub.).—China.
umbelliferum, Steven, Bot. Mag. Mosc. iii. 186 = majus.

CHELIDONIUM:—
uniflorum, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 169 = Stylophorum japonicum.
violaceum, Lam. Fl. Fr. iii. 690 = Roemeria hybrida.
vitellinum, C. Muell. in Walp. Ann vii. 87 = Glaucium vitellinum.

CHELIDOSPERMUM, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 162, in syn. (1850) = **Pittosporum**, Banks.
verticillatum, Zipp. ex Blume, l. c. = P. Cheilospermum.

CHELIDURUS, Willd. in Cothen. Disp. Meth. 4 (1790) (Gen. ignot.).
ignotus, *J. F. Gmel*, Syst. 105.—Hab. ?

CHELIUSIA, Sch. Bip. in Flora, xxiv. (1841) 1. Intell. 26, nomen; A. Rich. Tent. Fl. Abyss. i. 380 = **Vernonia**, Schreb. (Compos.).
abyssinica, Sch. Bip. l. c. = V. amygdalina.

CHELONANTHERA, Blume, Bijdr. 884. t. 51 (1825) = **Coelogyne**, Lindl. (Orchid.).
gibbosa, Blume, l. c. 383 = Pholidota gibbosa.
incrassata, Blume, l. c. 384 = Coel. incrassata.
longifolia, Blume, l. c. 385 = Coel. longifolia.
minuta, Blume, l. c. = Coel. minuta.
speciosa, Blume, l. c. 384. = Coel. speciosa.
sulphurea, Blume, l. c. 383 = Coel. sulphurea.
ventricosa, Blume, l. c. = Pholidota ventricosa.

CHELONE, Linn. Syst. ed. 1 (1735). *SCROPHULARINEAE*. Benth. & Hook. f. ii. 939.
CHLONANTHUS, Rafin. Princ. Somiol. 26 (1814).
alba, Moench, Meth. 442 = glabra.
albida, Spreng. Syst. ii. 813 = Penstemon albidus.
alpina, Spreng. l. c. iv. Cur. Post. 235 = Penstemon glaber.
angustifolia, H. B. & K. Nov. Gen. et Sp. ii. 365. t. 178 = Pentstemon campanulatus.
angustifolia, Stend. Nom. ed. I. 186 = Pentstemon caeruleus.
atropurpurea, Sweet, Brit. Flow. Gard. Ser. I. t. 235 = Pentstemon campanulatus.
barbata, Cav. Ic. iii. 22. t. 242 = Penstemon barbatus.
Bradburii, Stend. Nom. ed. I. 186 = Penstemon grandiflorus.
caerulea, Spreng. Syst. iii. 813 = Penstemon caeruleus.
campanulata, Cav. Ic. i. 18. t. 29 = Pentstemon campanulatus.
campanuloides, Andr. Bot. Rep. t. 40 = Penstemon campanulatus.
capitata, Rafin. Med. Fl. ii. 118 = glabra.
centranthifolia, Benth. in Trans. Hort. Soc. Lond. N. S. i. (1835) 481 = Penstemon centranthifolius.
cheilanthifolia, Pext. Mag. Bot. ii. t. 50 = prec.
cristata, Spreng. Syst. ii. 813 = Penstemon cristatus.
Digitalis, Sweet, Brit. Flow. Gard. Ser. I. t. 130 = Penstemon laevigatus.
elatior, Rafin. Med. Fl. ii. 118 = glabra.
elegans, H. B. & K. Nov. Gen. et Sp. ii. 364 = Penstemon campanulatus.
erianthera, Stend. Nom. ed. I. 186 = Penstemon cristatus.
filiformis, Buch.-Ham. ex C. B. Clarke, in DC. Monog Phan. v. 134 = Boeica filiformis.
formosa, Wendl. Beobacht. 51 = Pentstemon barbatus.
frutescens, Stend. Nom. ed. I. 186 = Penstemon fruticosus.
gentianoides, H. B. & K. Nov. Gen. et Sp. ii. 364. t. 172 = Pentstemon gentianoides.
glabra, *Linn. Sp. Pl.* 611.—Am., bor.
gracilis, Spreng. Syst. ii. 813 = Penstemon gracilis.
grandiflora, Spreng. l. c. = Penstemon grandiflorus.
hirsuta, Linn. Sp. Pl. 611 = Penstemon pubescens.
imberbis, H. B. & K. Nov. Gen. et Sp. ii. 363 = Pentstemon imberbis.
inermis, Pott. in Dict. Sc. Nat. xxxviii. (1825) 386 = Penstemon imberbis.
laevigata, Pers. Syn. ii. 169 = Penstemon laevigatus.
lanceolata, Rafin. Med. Fl. ii. 117 = glabra.
latifolia, Buch. Mem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 574 = Rhynchotechum latifolium.
latifolia, Mühl. Cat. ex Ell. Sketch, ii. 127 = oblique.
Lyoni, *Pursh, Fl. Am. Sept.* ii. 727.—Am. bor.
maculata, Rafin. Med. Fl. ii. 117 = glabra.
major, Sims, Bot. Mag. t. 1064 = Lyoni.

CHELONE:—
nemorosa, *Dougl. in Lindl. Bot. Reg.* t. 1211.—Am. bor. occ.
oblique, *Linn. Syst.* ed. XII. 468.—Am. bor. occ.
Penstemon, Linn. Mant. 415 = Pentstemon pubescens.
Penstemon, Linn. Sp. Pl. 612, pro parte = Pentstemon laevigatus.
pubescens, Sweet, Hort. Brit. ed. I. 305 = Pentstemon pubescens.
purpurea, Mill. Gard. Dict. ed. VIII. n. 2 = obliqua.
purpurea, Rafin. Med. Fl. ii. 118 = glabra.
rosea, Cerv. ex Sweet, Brit. Flow. Gard. Ser. I. t. 280 = Pentstemon campanulatus.
rubicunda, Buch.-Ham. ex C. B. Clarke, in DC. Monog. Phan. v. 135 = Boeica Griffithii.
rueliodes, Andr. Bot. Rep. t. 34 = Pentstemon barbatus.
rueliodes, Linn. f. Suppl. 271 = Ourisia magellanica.

CHELONOPSIS, Miq. in Ann. Mus. Lugd. Bat. ii. 111 (1865). *LABIATAE*, Benth. & Hook. f. ii. 1204.
moschata, *Miq. l. c.*—Japon.

CHELROSTYLIS, Prits. Ic. Ind. 265 (1855), sphalm. = **Chelrostylis**, Blume.

CHEMNICIA, Scop. Introd. 189 (1777) = **Strychnos**, Linn. (Loganiac.).

CHEMNIZIA, Steud. Nom. ed. II. i. 347 (1840) = **Strychnos**, Linn. (Loganiac.).

CHENOCARPUS, Neck. Elem. i. 202 (1790) = **Spermacoce**, Linn. (Rubiac.).

CHENOLEA, Thunb. Nov. Gen. 10 (1781). *CHENOPODIACEAE*. Benth. & Hook. f. iii. 89.
BASSIA, All. Misc. Taur. iii. (1766) 177. t. 4.
ECHINOPSILON, Moq. in Ann. Sc. Nat. Ser. II. ii. (1834) 257.
ERIOCHITON, F. Muell. 2nd Gen. Report, 15 (1854), nomen.
LONDESIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 40 (1835).
WILLEMETIA, Maerkl. in Schrad. Journ. Bot. iii. (1800) i. 329.
arabica, *Boiss. Diagn.* Ser. I. xii 97.—Syria; Arabia
asphalica, Boiss. l. c. xii. 84 = Suaeda asphalica.
astrocarpa, *F. Muell. Fragm.* xii. 12.—Austral.
bifora, *F. Muell. l. c.* x. 91.—Austral.
canescens, *Moq. Chenop. Enum.* 96.—Morocco; Ins. Canar.
carnea, *Benth. Fl. Australiae*. v. 190.—Austral.
Dallachyana, *Benth. l. c.* 191.—Australia.
discantha, *F. Muell. Fragm.* x. 91.—Austral.
diffusa, *Thunb. Nov. Gen.* 10.—Afr. austr.
echinopsila, *F. Muell. Fragm.* x. 92.—Austral.
eschylmoides, *F. Muell. l. c.*—Austral.
eurotioides, *F. Muell. ex Benth. Fl. Austral.* v. 191.—Austral.
Forrestiana, *F. Muell. Fragm.* xii. 19.—Austral.
hirsuta, *Archang. Comp. Fl. Ital.* 595.—Italia.
lanata, *Jacq. Fragm.* 76.—Afr. austr.
lanata, Jacq. ex Moq. in DC. Prod. xiii. ii. 188 = Kochia prostrata.
lanata, Moq. l. c. 129 = canariensis.
Muelleri, *Benth. Fl. Austral.* v. 191.—Austral.
paradoxa, *F. Muell. Fragm.* x. 91.—Austral.
quinquecuspis, *F. Muell. l. c.*—Austral.
sclerolaenoides, *F. Muell. ex Benth. Fl. Austral.* v. 192.—Austral.
salsuginosa, *F. Muell. Fragm.* x. 92.—Austral.
triconia, *Benth. Fl. Australiae*. v. 191.—Austral.
triscupis, *F. Muell. Fragm.* x. 92.—Austral.
tridens, *F. Muell. l. c.* xii. 19.—Austral.

CHENOPODINA, Moq. in DC. Prod. xiii. ii. 159 (1849) = **Suaeda**, Forsk. (Chenopod.).
acuminata, Moq. l. c. 181 = S. acuminata.
aestuaria, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 229 = S. maritima.
altissima, Moq. in DC. Prod. xiii. ii. 162 = S. altissima.
anonyma, Moq. l. c. 164.—Aegypt.
asphaltica, Boiss. Diagn. Ser. I. xii. 86 = S. asphaltica.
australis, Moq. in DC. Prod. xiii. ii. 163 = S. australis.

CHENOPODINA:—
baccifera, Moq. in DC. Prod. xiii. ii. 160 = Suaeda baccifera.
baccifoemis, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 228 = S. maritima.
Biebersteiniana, Moq. in DC. Prod. xiii. ii. 163 = S. microphylla.
dendroides, Moq. l. c. 164 = S. dendroides.
depressa, Moq. l. c. = S. plattensis.
filiformis, Moq. l. c. = S. maritima.
fruticulosa, Moq. l. c. 165 = S. fruticulosa.
glauca, Moq. l. c. 162 = S. glauca.
Hostii, Moq. l. c. 165.—Dalmat.
indica, Wight, Ic. v. t. 1793 = S. maritima.
leiosperma, Moq. in DC. Prod. xiii. ii. 162 = S. altissima.
linearis, Moq. l. c. 164 = S. linearis.
linearis, Torr. in Stansbury, Rep. 394 = S. Torreyana.
maritima, Coss. ex Willk. & Lange, Prod. Fl. Hisp. i. 361 = S. setigera.
maritima, Moq. in DC. Prod. xiii. ii. 161 = S. maritima.
microphylla, Moq. l. c. = S. microphylla.
microsperma, Moq. l. c. 165 = S. microsperma.
minima, Moq. l. c. 162 = S. minima.
Moquini, Torr. in Pacif. Rail. Rep. vii. 18 = S. Torreyana.
parviflora, Moq. in DC. Prod. xiii. ii. 165 = S. microsperma?
physophora, Moq. l. c. 164 = S. physophora.
prostrata, Moq. l. c. 162 = S. microsperma.
pycnantha, C. Koch, in Linnaea, xxiv. (1851) 90 = Seidlitzia florida.
pygmaea, Moq. in DC. Prod. xiii. ii. 161 = S. microsperma?
salsa, Moq. l. c. 160 = S. salsa.
sativa, Moq. l. c. 165 = S. maritima.
setigera, Moq. l. c. 160 = S. setigera.
spicata, Moq. l. c. 165 = S. maritima.
spiendens, Gren. in Gren. & Godr. Fl. Fr. iii. 30 = S. setigera.
tortuosa, Moq. in DC. Prod. xiii. ii. 163 = S. maritima?
vera, Moq. l. c. 163 = S. vera.

CHENOPODIUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 67 (1737). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 61.
AGATHOPHYTUM, Moq. in Ann. Sc. Nat. Ser. II. i. (1834) 201. t. 10 c.
AMBRINA, Spach, Hist. Vég. Phan. v. 295 (1836).
ANSERINA, Dum. Fl. Belg. 21 (1827).
BLITUM, [Tourn.] Linn. Gen. ed. I. 80 (1737).
BOTRYDIUM, Spach, Hist. Veg. Phan. v. 295 (1836).
LIPANDRA, Moq. Chenop. Enum. 19 (1840).
MOROCARPUS, Adans. Fam. ii. 261 (1763).
OLIGANDRA, Less. in Linnaea, ix. (1834) 199.
OLIGANTHERA, Endl. Gen. Suppl. i. 1377 (1841).
ORTHOSPERMUM, Opiz, Seznam, 70 (1852); Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 63.
ORTHOSPORUM, T. Nees, Gen. Fl. Germ. Monochl. n. 58 (1835).
OXYBASIS, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 788.
TELOXYS, Moq. in Ann. Sc. Nat. Ser. II. i. (1834) 289. t. 10.
acerifolium, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 194, nomen.—Rossia.
acuminatum, Schur, Enum. Pl. Transs. 902 = rubrum ?
acuminatum, *Willd. in Ges. Nat. Fr. Neue Schr.* ii. (1798) 194. t. 5. f. 2.—Senegaria.
acutifolium, Hook. & Arn. Bot. Beech Voy. 307 = acuminatum.
acutifolium, Kit. in Schult. Oestr. Fl. ed. II. i. 458; Sm. Comp. Fl. Brit. 42 = polyspermum.
aegyptiacum, *Hasselq. It. Palaest.* 460 = Suaeda baccata.
album, Boec. ex Moq. in DC. Prod. xiii. ii. 81 = Bocsiacum.
album, *Linn. Sp. Pl.* 319.—Reg. temp. et trop.
alexandrinum, Desf. ex Moq. in DC. Prod. xiii. ii. 155 = Suaeda vermiculata.
altissimum, Linn. Sp. Pl. 321 = Suaeda altissima.
ambiguum, R. Br. Prod. i. 407 = glaucum.
ambroinoides, Bert. ex Steud. Nom. ed. II. i. 346 = citrifolium.
ambroinoides, Hanow, in Journ. Linn. Soc. xiii. (1873) 86 = acuminatum.
ambrosioides, *Linn. Sp. Pl.* 219.—Reg. temp. et trop.
americanum, Spreng. Syst. i. 922 = Suaeda plattensis.

CHENOPODIUM :—
angulatum, Cart. ex Stend. Nom. ed. I. 187 = hybridum.
angulosum, Lam. Fl. Fr. iii. 249 = hybridum.
angustifolium, Gilib. Exercit. ii. 437 = polyspermum.
angustifolium, Pav. ex Moq. in DC. Prod. xiii. ii. 114 = Atriplex linifolia.
antarcticum, *Benth. & Hook. f. Gen.* iii. 52.—Reg. Magellan.
anthelminticum, Bert. ex Stend. Nom. ed. II. i. 348 = chilense.
anthelminticum, Linn. Sp. Pl. 220 = ambrosioides.
arcticum, Hort. ex Moq. in DC. Prod. xiii. ii. 86 = Monolepis chenopodioides.
arenarium, Gaertn. Meyer & Scherb. Fl. Wett. i. 356 = Kochia arenaria.
aristatum, *Linn. Sp. Pl.* 221.—Sibir.
arvicum, Desmaz. Cat. Pl. Belg. 80 = polyspermum.
asphalticum, Boiss. Diagn. Ser. I. xii. 98 = Suaeda asphaltica.
astracanicum, Ledeb. Cat. Hort. Dorpat. (1819) = rubrum.
atriplicinum, *F. Muell. Fragm.* viii. 11.—Austral.
Atriplicis, Lian. f. Suppl. 171 = purpurascens.
augustanum, All. Fl. Pedem. ii. 198. t. 88 = Kochia prostrata.
augustanum, Hort. Par. ex Moq. in DC.-Prod. xiii. ii. 115 = Kochia hyssopifolia.
auricomum, *Lindl. in Mitch. Journ. Trop. Austral.* 94. —Austral.
australasicum, Moq. Chenop. Enum. 20 = Rhagodia linifolia.
australe, R. Br. Prod. 407 = Suaeda australis.
baccatum, Labill. Nov. Holl. Pl. i. 71. t. 96 = Rhagodia Billardieri.
baryosmum, Roem. & Schult. Syst. vi. 269 = Salsola foetida.
benghalense, Spielm. ex Stend. Nom. ed. II. i. 348 = Atriplex horizensis ?
Beriandieri, *Moq. Chenop. Enum.* 23.—Am. centr.
betaceum, *Andrz. ex Trautv. in Act. Hort Petrop.* ix. (1884) 135, nomen.—Ionnis.
bicolor, Boj. ex Moq. in DC. Prod. xiii. ii. 71 = album.
Bieberstejnianum, Roem. & Schult. Syst. vi. 267 = Suaeda microphylla.
biforme, Nees, in Lehm. Pl. Preiss. i. 586 = album.
bipinnatifidum, *Moric. ex Moq. in DC. Prod.* xiii. ii. 76. —Mexic.
Bisseriade, Menyh. Kalocsa Vidék. Növény. 157 = polyspermum.
blitoides, Lejeune, Fl. Spa, 136 = rubrum.
Blitum, F. Muell. Sel. Pl. Indust. Cult. 49 = capitatum.
bonariense, Hort. Par. ex Moq. in DC. Prod. xiii. ii. 67 = purpurascens.
bonariense, Tenore, Ind. Sem. Hort. Neap. (1838) 18 = hirsutum.
Bonus-Henricus, *Linn. Sp. Pl.* 218.—Europ.
Boscianum, *Moq. Chenop. Enum.* 21.—Texas.
botryoides, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Botrys.
botryoides, Rafin. ex Moq. in DC. Prod. xiii. ii. 75 = C. Botrys.
botryodes, Sm. Engl. Bot. t. 2347 = rubrum.
Botrys, *Linn. Sp. Pl.* 219.—Reg. bor.
Brownenum, Roem. & Schult. Syst. vi. 275 = album.
bryoniaefolium, *Bunge, Del. Sem. Hort. Petrop.* (1876) 10 : *ex Trautv. in Act. Hort.-Petrop.* ix. (1884) 398. —Sibiria.
calcosiliforme, Hook. Fl. Bor. Am. ii. 126 = Suaeda calcosiliformis.
californicum, *S. Wats. in Bot. Calif.* ii. 48.—Calif.
camphoratofolium, Pourr. in Mém. Acad. Toul. iii. (1788) 311 = Kochia prostrata.
camphoratoroides, Hort. ex Moq. in DC. Prod. xiii. ii. 128 = prasc.
candicans, Lam. Fl. Fr. iii. 248 = album.
cantabricum, *Lag. Hort. Matrit.* (1815) 15, nomen.—Hispan.
capitatum, *Aschers. Fl. Brandenb.* 572.—Reg. bor. et austr.
carinatum, *R. Br. Prod.* 407.—Austral. ; N. Zel.
carnosulum, *Moq. in DC. Prod.* xiii. ii. 64.—Calif.
Caroxylon, D'Urban, in Journ. Linn. Soc. viii. (1865) 272 = Salsola Caroxylon.
carthagenense, Zuccagni, in Roem. Collect. 183 = murale.
catenulatum, Schleich. ex Stend. Nom. ed. II. i. 348 = album.

CHENOPODIUM :—
caudatum, Jacq. Coll. ii. 295 ; Ic. Pl. Rar. ii. 12. t. 344 = Amaranthus viridis.
Chamissus, Buch.-Ham. in Wall. Cat. n. 6958 = murale.
Chandana, *Buch.-Ham. in Wall. Cat. n.* 6957.—Ind. or.
chilense, Pers. Syn. i. 294 = C. Quinoa.
chilense, *Schrad. Ind. Sem. Hort. Gott.* (1833) 2.—Chili.
chyonolanospermum, Zuccagni, in Roem. Collect. 134 = urbicum.
cinereum, Moq. Chenop. Enum. 27 = Atriplex peruviana.
citriodorum, *Steud. Nom.* ed. II. i. 348.—Ins. Juan Fernand.
Clementeæ, Spreng. Syst. i. 932 = Suaeda setigera.
coccinellum, Hort. ex Willd. Enum. Hort. Berol. 291 = lanceolatum.
concatenatum, Tbuill. Fl. Paris, ed. II. 125 = album.
congestum, Hook. f. in Lond. Journ. Bot. vi. (1847) 280 = Rhagodia Billardieri.
coriospermifolium, *Hort. Amstel.* (1814) 14, nomen.—Hab. ?
cornutum, *Benth. & Hook. f. ex S. Wats. in Bot. Calif.* ii. 492.—Mexic.
coronopifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = seq.
Coronopus, *Moq. in DC. Prod.* xiii. ii. 76.—Ins. Tenerif.
crassifolium, Bieb. Fl. Taur. Casc. iii. 176 = Suaeda microphylla.
crassifolium, Desf. Tabl. ed. II. 51 = Suaeda maritima.
crassifolium, Hornem. Hort. Hafn. 254 = rubrum.
cristatum, *F. Muell. Fragm.* vii. 11.—Austral.
cuneifolium, *Vahl, in West, Fl. S. Cruc.* 278.—Santa Cruz.
cuneifolium, Vent. ex Moq. in DC. Prod. xiii. ii. 73 = spathulatum.
deltoideum, Lam. Fl. Fr. iii. 249 = urbicum.
detestans, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 550.—N. Zel.
divaricatum, *Black. Cat. Hort. Gorenk.* (1808) 24, nomen.—Rossia ?
effusum, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 346 = foetidum.
elatum, Shuttl. ex Moq. in DC. Prod. xiii. ii. 71 = album.
erosum, Bast. in Desv. Journ. Bot. iii. (1814) 20 = opulifolium.
erosum, R. Br. Prod. 407 = murale.
esculentum, Salisb. Prod. 151 = C. Bonus-Henricus.
exocarpum, *Griseb. in Goett. Abh.* xxiv. (1879) 97.—Reg. Argent.
ficifolium, *Sm. Fl. Brit.* i. 276.—Europ.
filifolium, Krock. Fl. Sil. Suppl. i. 309 = prasc.
filifolium, Dum. Fl. Belg. 23 = Suaeda maritima.
filiforme, Moench, Meth. 330 = Suaeda altissima.
flavescens, Roem. & Schult. Syst. vi. 269 = Salsola vermiculata.
flavum, Forsk. Fl. Aegypt. Arab. Suppl. 205 = murale.
fœtidum, Lam. Fl. Fr. iii. 244 = C. Vulvaria.
foetidum, *Schrad. in Gen. Naturf. Fr. Berl. Mag.* ii. (1808) 70.—Reg. trop.
foliosum, Aschers. Fl. Brandenb. 572 = capitatum.
Fremontii, *S. Wats. in Bot. King, Exp.* 387.—Am. bor. occ.
frigidum, *Phil. Fl. Atac.* 47.—Chili.
frutescens, *C. A. Mey. in Ledeb. Fl. Alt.* i. 403.—Sibir. alsatic.
fruticosum, Bieb. Fl. Taur. Casc. i. 181 = Suaeda dendroides.
fruticosum, Linn. Sp. Pl. 221 = Suaeda fruticosa.
furfuraceum, Moq. in DC. Prod. xiii. ii. 64 = auricomum.
Gandhium, Buch.-Ham. in Wall. Cat. n. 6958 = murale.
gigantteum, D. Don, Prod. Fl. Nep. 75 = album.
glandulosum, F. Muell. Fragm. vii. 11 = carinatum.
glaucum, *Linn. Sp. Pl.* 220.—Reg. bor. et austr.
glochidatum, Gilli. ex Moq. in DC. Prod. xiii. ii. 61 = papulosum.
glomeruloseum, Reichb. Fl. Germ. Exsurs. 579 = album.
graveolens, Lag. & Rodr. in Anal. Clenc. Nat. v. (1809) 70 = foetidum.

CHENOPODIUM :—
graveolens, *Willd. Enum. Hort. Berol.* i. 290.—Mexic.
guineense, Jacq. Coll. ii. 346 ; Ic. Pl. Rar. ii. 12. t. 345 = murale.
Guinoa, Krock. Fl. Sil. Suppl. i. 301 = C. Quinoa.
Halimus, Thunb. Prod. Pl. Cap. 48 = Atriplex Halimus.
halophilum, *Phil. in Anal. Univ. Chil.* (1861) i. 67.—Chili.
hastatum, Dum. in Bull. Soc. Bot. Belg. iv. (1865) 339 = Atriplex hastata.
hastatum, *Phil. Fl. Atac.* 47.—Chili.
hircinum, *Schrad. Ind. Sem. Hort. Gott.* (1833) 2.—Bras.
hirsutum, Linn. Sp. Pl. 221 = Kochia hirsuta.
Hookerianum, Moq. in DC. Prod. xiii. ii. 68 = murale.
hortense, Raddi, ex Moq. l. c. 161 = Suaeda maritima.
hortense, Roem. & Schult. Syst. vi. 268 = Suaeda hortensis.
Hostii, Ledeb. Fl. Ross. iii. 788 = Suaeda maritima.
humifusum, Kitt. Taschenb. ed. II. 264 = crassifolium.
humifusum, *Zuccagni, in Roem. Collect.* 134.—Europ.
humile, Hook. Fl. Bor. Am. ii. 127 = rubrum.
hybridum, *Linn. Sp. Pl.* 219.—Reg. temp. hor.
hybridum, Lour. Fl. Cochinch. 174 = Loureirei.
ilicifolium, Griff. Notul. iv. 387 = C. Botrys.
incisum, *Poir. Encyc. Suppl.* i. 392.—Am. centr.
intermedium, Mert. & Koch, in Rochl. Deutschl. Fl. ed. III. ii. 297 = urbicum.
Jacquini, Tenore, Fl. Nap. iii. 258 = Suaeda maritima.
lacerum, *Krock. Fl. Sil. Suppl.* i. 304.—Silesia.
laciniatum, Roxb. Hort. Beng. 21 ; Fl. Ind. ii. 59 = album.
laciniatum, Thunb. Prod. Pl. Cap. 48 = Atriplex dactniata.
lanceolatum, R. Br. Prod. 407 = album.
lanceolatum, Muhl. ex Willd. Enum. Hort. Berol. 291 = album.
lanuginosum, Moench, Meth. 330 = Kochia hyssopifolia.
latifolium, *Pall. Reise Statth. Russ. Reich.* i. 280.—Rossia.
leiospermum, DC. Fl. Fr. iii. 390 = album.
leptophyllum, *Nutt. ex Moq. in DC. Prod.* xiii. ii. 71. —Am. bor.
leucospermum, Schrad. Ind. Sem. Hort. Gott. (1834) 9 = purpurascens.
linearc, Hort. ex Stend. Nom. ed. II. i. 348 = Kochia prostrata.
linifolium, Roem. & Schult. Syst. vi. 271 = Suaeda linifolia.
littorale, *Moq. Chenop. Enum.* 24.—Austral.
littorale, Thunb. in Act. Soc. Sc. Upsal. vii. (1815) 142 = Atriplex littoralis.
Lourcirei, *Stend. Nom.* ed. II. i. 348.—China.
lucidum, Gilli. Exercit. ii. 440 = murale.
macrocarpum, Desv. Journ. Bot. i. (1813) 48 = Suaeda maritima.
macrocarpum, *Hook. f. Fl. Antarct.* ii. 341.—Ins. Falkland.
macrospermum, Hort. Ant. ex Moq. in DC. Prod. xiii. ii. 84 = rubrum.
marginatum, Spreng. ex Hornem. Hort. Hafn. i. 256 = polyspermum.
maritimum, Frivald. ex Ledeb. Fl. Ross. iii. 788 = Suaeda altissima.
maritimum, Linn. Sp. Pl. 221 = Suaeda maritima.
maritimum, Torr. in Ann. Lyc. N. York, ii. (1828) 220 = Suaeda diffusa.
Matthioli, Bertol. ex Moq. in DC. Prod. xiii. ii. 64 = rubrum.
melanospermum, Wallr. Sched. Crit. 113 = urbicum.
mexicanum, *Moq. in DC. Prod.* xiii. ii. 70.—Mexic.
microsathum, *Trautv. in Bull. Soc. Nat. Mosc.* xii. (1839) i. 464.—Rossia.
microphyllum, Bieb. Fl. Taur. Casc. iii. 176 = Suaeda dendroides.
microphyllum, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 74.—Austral.
microphyllum, *Thunb. Prod. Fl. Cap.* 48.—Afr. austr.
microspermum, Wallr. Sched. Crit. 113 = urbicum.

CHENOPODIUM:—
mucronatum, *Thumb. Prod. Pl. Cap.* 48.—Afr. austr.
multifidum, Linn. Sp. Pl. 220 = Roubieva multifida.
multiflorum, Moq. in DC. Prod. xiii. ii. 75 = C. Botrys.
murale, *Linn. Sp. Pl.* 219.—Reg. bor. et austr.
neglectum, *Dum. in Bull. Soc. Bot. Belg.* iv. (1865) 339.—Belgium.
nepalense, Hort. Monsp. ex Moq. in DC. Prod. xiii. ii. 74 = C. Botrys.
nigrum, *Rafin. Atl. Journ.* 146.—Am. bor.
nitidum, Spreng. Syst. i. 921 = Salsola nitida.
nitrariaceum, *F. Muell. ex Benth. Fl. Austral.* v. 158.—Austral.
obovatum, *Moq. in DC. Prod.* xiii. ii. 73.—Am. austr.
oleraceum, *Schur, Enum. Pl. Transs.* 573.—Transylv.
olidum, Curt. Fl. Lond. fasc. v. t. 20 = C. Vulvaria.
olidum, *S. Wats. in Proc. Am. Acad.* ix. (1874) 95.—Am. bor.
oppositifolium, Linn. f. Suppl. 173 = Halogeton oppositiflorus.
oppositiflorum, Willd. Sp. Pl. i. 1307 = Girgensohnia oppositiflora.
opulaceum, Neck. Delic. Gallo-Belg. i. 130 = album.
opulifolium, *Schrad. in DC. Fl. Fr.* v. 372.—Reg. bor.
paganum, Reichb. Fl. Germ. Excurs. 579 = album.
Palasianum, Roem. & Schult. Syst. vi. 271 = Kochia hirsuta.
pallidum, Moq. Chenop. Enum. 30 = glaucum ?
paniculatum, *Hook. Bot. Misc.* ii. (1831) 237.—Peruv.
paniculatum, Salzm. ex Moq. in DC. Prod. xiii. ii. 18 = Microtea paniculata.
papulosum, *Moq. l. c.* 61.—Bras.
parvifolium, Roem. & Schult. Syst. vi. 266 = Suaeda dendroides.
patulum, Mérat, Nouv. Fl. Par. 96 = rubrum.
patulum, *Roth, Nov. Pl. Sp.* 180.—Ind. or.
pauciflorum, Herb. Vind. ex Moq. in DC. Prod. xiii. ii. 89 = Exomis axyrioides.
pauciflorum, *Mart. Hort. Monac.* (1829) 56, nomen.—Hab. ?
Payco, Molina, Sagg. Chile, ed. II. 118 = Roubieva multifida.
pedunculare, Bertol. Fl. Ital. iii. 82 = album.
petiolare, *H. B. & K. Nov. Gen. et Sp.* ii. 191.—Ecuador.
petiolare, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 345 = Cryptocarpus globosus.
physophora, Moq. ex Boiss. Fl. Orient. iv. 939 = Suaeda physophora.
pinnatum, Hort. ex Moq. in DC. Prod. xiii. ii. 93 = Atriplex laciniata.
polyspermum, Neck. Delic. Gallo-Belg. i. 128 = polyspermum.
polyspermum, *Linn. Sp. Pl.* 220.—Europ.; As. bor.
populifolium, Hort. ex Moq. in DC. Prod. xiii. ii. 66 = bicfolium.
portulacoides, Thunb. Fl. Cap. ii. 178 = Atriplex portulacoides.
praestorium, *Dum. in Bull. Soc. Bot. Belg.* iv. (1865) 340.—German.
procerum, *Hochst. ex Moq. in DC. Prod.* xiii. ii. 76.—Abyssin.
prostratum, Hort. ex Moq. l. c. 132 = Kochia prostrata.
prostratum, Jacquem. ex Moq. l. c. 72 = glaucum ?
prostratum, Roem. & Schult. Syst. vi. 272 = Suaeda prostrata.
pseudo-botrys, *Hort. ex Heynh. Nom.* ii. 188, nomen.—Hab. ?
Pumilio, *R. Br. Prod.* i. 407.—Austral.
punctulatum, Scop. Delic. Insub. i. 26. t. 11 = purpurascens.
purpurascens, Bach. Ham. in Wall. Cat. n. 6956 = album.
purpurascens, Jacq. Hort. Vindob. iii. 43. t. 80 = album ?
pusillum, *Hook. f. Handb. New Zeal. Fl.* 281.—N. Zel.
pusillum, *Bertol. in Nov. Comm. Bonon.* iv. (1842) 418.—Fl. Raphrat.
pygmaeum, Mezyh. Kalocsa Vidék. Növeny. 154 = rubrum.
quercifolium, Hort. Monsp. ex Moq. in DC. Prod. xiii. ii. 74 = graveolens.
Quinoa, *Willd. Sp. Pl.* i. 1301.—Am. austr.
radiatum, Schrad. Neues Journ. iii. iii. (1809) 85 = Cyclolomа platyphyllum.

CHENOPODIUM:—
retusum, *Juss. ex Moq. in DC. Prod.* xiii. ii. 78.—Bras.
rhadinostachyum, *F. Muell. in Wing. South Sc. Record*, ii. (1882) 98.—Austral.
rhombifolium, Muhl. ex Willd. Enum. Hort. Berol. 288 = urbicum.
riparium, Boeningh. ex Moq. in DC. Prod. xiii. ii. 71 = album.
rubricaule, Schrad. ex Moq. l. c. 67 = purpurascens.
rubrum, Forsk. ex Steud. Nom. ed. I. 188 = murale.
rubrum, *Linn. Sp. Pl.* 218.—Europ.; As. bor.
rudereale, *Kit. ex Moq. in DC. Prod.* xiii. ii. 77.—Hungaria.
ruderale, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 89 = C. Bonus-Henricus.
sagittatum, Lam. Fl. Fr. ii. 244 = C. Bonus-Henricus.
salsum, Guss. Fl. Sic. Prod. i. 399 = Suaeda maritima.
salsum, Hohen. ex Ledeb. Fl. Ross. iii. 782 = Suaeda altissima.
salsum, Linn. Sp. Pl. 221 = Suaeda salsa.
Sancta-Marsa, Vell. Fl. Flum. 126; iii. t. 104 = ambrosioides.
sandwicheum, *Moq. Chenop. Enum.* 28.—Ins. Sandvic.
Schraderianum, Roem. & Schult. Syst. vi. 260 = foetidum.
scoparia, Linn. Sp. Pl. 221 = Kochia scoparia.
secundiflorum, Viv. Fl. Lib. Spec. 67 = aristatum.
sepium, *Mey. in Abh. Boehm. Ges. Wiss.* (1787) 320.—Bohemia.
sericeum, Spreng. Syst. i. 921 = Kochia sericea.
sericeum, Vitm. Summa Veg. ii. 98 = Kochia prostrata.
serotinum, *Forsk. Fl. Aegypt. Arab.* 205.—Arab.
serotinum, Ledeb. Fl. Alt. i. 406 = album.
serotinum, Linn. Cent. Pl. ii. 12 = ficifolium.
serotinum, Suter, Fl. Helv. i. 148 = hybridum.
setigerum, DC. Cat. Hort. Monsp. 94 = Suaeda setigera.
simplex, *Rafin. Atl. Journ.* 146.—Am. bor.
sinense, Hort. ex Moq. in DC. Prod. xiii. ii. 60 = aristatum.
sinense, Willem. Ess. Med. iii. 283 = Kochia sericea.
sinuatum, Thunb. Fl. Cap. ii. 178 = Atriplex sinuata.
spathulatum, *Sieber, ex Moq. in DC. Prod.* xiii. ii. 73.—Bras.
spicatum, *Lavy, Veg. Orig.* 10.—Hab. ?
spicatum, Roem. & Schult. Syst. vi. 271 = Suaeda maritima.
spinacifolium, Stokes, Bot. Mat. Med. ii. 14 = C. Bonus-Henricus.
spinosum, Hook. Fl. Bor. Am. ii. 137 = Grayia polygaloides.
stellatum, *S. Wats. in Proc. Am. Acad.* xvii. (1882–83) 146.—Mexic.
stramonifolium, Cheval. Fl. Par. ii. 288 = hybridum.
strictum, *Roth, Nov. Pl. Sp.* 180.—Ind. or.
subspicatum, Nutt. Gen. Am. i. 199 = Atriplex hastata.
suffruticosum, Willd. Enum. Hort. Berol. 290 = ambrosioides.
teneriimum, *Krock, Fl. Sil. Suppl.* i. 305.—German.
tenue, Colla, in Mem. Acc. Torin. xxxix. (1836) 9. t. 50 = Roubieva multifida.
tomentum, *Thou. Fl. Tristan.* 38.—Ins. Tristan d'Acugn.
Tournefortianum, Hort. ex Moq. in DC. Prod. xiii. ii. 158 = Suaeda altissima.
Tournefortii, Hort. ex Moq. l. c. = praec.
triandrum, *Forst. f. Prod.* 21.—N. Zel.
triangulare, *R. Br. Prod.* 407.—Austral.
triangulare, Dulac, Fl. Hautes-Pyr. 171 = C. Bonus-Henricus.
triangulare, Forsk. Fl. Aegypt. Arab. Suppl. 205 = murale.
triangulariifolia, Gilib. Exercit. ii. 437 = C. Bonus-Henricus.
trifidum, Trev. Ind. Sem. Vratisl. (1839) = Monolepis chenopodioides.
trigonon, Roem. & Schult. Syst. vi. 275 = triangulare.
trigynum, Roem. & Schult. l. c. 268 = Suaeda altissima.
trilobatum, Hort. ex Moq. in DC. Prod. xiii. ii. 76 = foetidum.
trilobum, Hort. ex Moq. l. c. = foetidum.
trilobum, Schult. ex Moq. l. c. 66 = ficifolium.
tripteris, Dum. Fl. Belg. 22 = Kochia tripteris.
tumidum, *Kit. Cat. Kvasm.* (1814) *Suppl.* 3, nomen.—Hab. ?

CHENOPODIUM:—
Tweedii, *Moq. in DC. Prod.* xiii. ii. 68.—Bras.
urbicum, *Linn. Sp. Pl.* 218.—Europ.; As. bor.
Vachellii, Hook. & Arn. Bot. Beech. Voy. 269 = acuminatum.
variegatum, Gouan, in Cat. Hort. Taur. (1810) = ambrosioides.
vestitum, Thunb. Prod. Pl. Cap. 48 = Atriplex glauca.
villosum, Hort. ex Moq. in DC. Prod. xiii. ii. 132 = Kochia prostrata.
villosum, Lam Encyc. i. 196 = Kochia hyssopifolia.
villosum, Spreng. Syst. i. 924 = Suaeda microphylla.
virens, *Wender. Orn. Mark.* (1836), nomen.—Hab. ?
virgatum, *Thunb. in Nov. Act. Soc. Sc. Ups.* vii. (1815) 148.—Japon.
virgiricum, Linn. Sp. Pl. 222 = aristata.
viride, Curt. Fl. Lond. fasc. ii. t. 16 = ficifolium.
viride, Linn Sp. Pl. 219 = album.
viride, Loisel. Fl. Gall. i. 145 = opulifolium.
vulgare, Gueldenst. ex Ledeb. Fl. Ross. iii. 700 = album.
vulpinum, Buch.-Ham. in Wall. Cat. n. 6954 = album, ambrosioides.
Vulvaria, *Linn. Sp. Pl.* 220.—Europ.; As. bor.
Wolffii, *Symb. in Termész. Fuzetek*, iii. (1879) 164.—Hungar.
Ziegleri, Perrot. ex Moq. in DC. Prod. xiii. ii. 24 = Limeum linifolium.
zostaerifolium, Hook. Fl. Bor. Am. ii. 127 = Atriplex Gmelini.

CHEOBULA, Vell. Fl. Flum. iv. t. 146 (1827) = **Cleobula**, Vell. (Inc. sed.).

CHERINA, Cass. in Dict. Sc. Nat. viii. 487 (1817) = **Chaetanthera**, Ruiz & Pav. (Compos.).
microphylla, Cass. l. c. 488 = Chaet. linifolia.

CHERLERIA, Hall. It. Helv. ii. n. 79. t. 1 (1740); ex Linn. Gen. ed. II. 196 (1742) = **Arenaria**, Rupp. (Caryophyll.).
biparitis, Fisch. ex Steud. Nom. ed. II. i. 349, nomen.—Sibir. baical.
bisulca, Bartl. in Presl, Rel. Haenk. ii. 12.—Peruv.
caespitosa, Lam. Fl. Fr. iii. 46 = A. Cherleria.
dicranoides, Cham. in Linnaea, i. (1826) 62 = Stellaria dicranoides.
grandiflora, D. Don, Prod. Fl. Nep. 214 = A. globiflora.
imbricata, Ser. in DC. Prod. i. 421 = A. aretioides.
juniperina, D. Don, Prod. Fl. Nep. 214 = A. densissima.
laevis, Bartl. in Presl, Rel. Haenk. ii. 12.—Peruv.
nitida, Bartl. l. c.—Peruv.
octandra, Sieber, ex Spreng. Syst. ii. 416 = A. aretioides.
peduncularis, Bunge, ex Steud. Nom. ed. II. i. 349 = *biparitis*.
sedoides, Forsk. Fl. Aegypt. Arab. p. lxvi = A. geniculata.
sedoides, Linn. Sp. Pl. 425 = A. Cherleria.
sedoides, Sibth. & Sm. Fl. Graec. Prod. i. 306 = A. trichocalycina.
sedoides, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 608 = Stellaria petraea.
sibirica, Regel & Tilling, in Nouv. Mém. Soc. Nat. Mosc. xi. (1858) 72.—Sibir. or.
stellata, Clarke, Trav. iv. 211 (= *Alsine parnassica*).—Graecia.

CHEROPHILUM, Nocca, in Usteri, Ann. Bot. v. (1798) 51 = Chaerophyllum, Linn. (Umbellif.).

CHERSYDRIUM, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 73 = **Dracontium**, Linn. (Aroid.).
Yararaca, Schott, l. c. 78 = D. asperum.

CHESNEA, Scop. Introd. 119 (1777) = **Cephalis**, Sw. (Rubiac.).

CHESNEYA, Bertol. in Nov. Comm. Bonon. v. (1842) 437 (Misc. Bot. i. 17. t. 2) = **Pimpinella**, Linn. (Umbellif.).
daucoides, Bertol. l. c. = P. eriocarpa.

CHESNEYA, Lindl. in Endl. Gen. 1275 (1840) = **Calophaca**, Fisch. (Legumin.).
astragalina, Jaub. & Spach, Illustr. i. 94.—Persia.
euphratensis, [Lindl. in] Chesn. Narrat. Euph. (1868) t. ad p. 489 = *rytidosperma*.

CHESNEYA:—
Ketschyii, Boiss. Diagn. Ser. I. vi. 34.—Persia.
microphylla, Jaub. & Spach, Illustr. i. 94 = *rytidosperma*.
mongolica, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 462.—Mongolia.
Olivieri, Jaub. & Spach, Illustr. i. 94.—Mesopotam.
parviflora, Jaub. & Spach, l. c. 95.—Persia.
rytidosperma, Jaub. & Spach, l. c. 93.—Armenia.
turkestanica, Franch. in Ann. Sc. Nat. Sér VI. xv. (1882) 253.—Turkestan.
vaginalis, Jaub. & Spach, Illustr. i. 96. t. 48.—Mesopotam.
velutina, Jaub. & Spach, l. c. 94 = *rytidosperma*.

CHESNYA, Reichb. Nom. 146 (1841) = prsec.

CHETARIA, Steud. Nom. ed. II. 511 (1841) = seq

CHETASTRUM, Neck. Elem. i. 108 (1790) = **Scabiosa**, Tourn. (Dipsac.).

CHETOCRATER, Rafin. Sylva Tellur. 149 (1838) = **Casearia**, Jacq. (Samydac.).
hirta, Rafin. l. c. = Cas. hirta.
javitensis, Rafin. l. c. = Cas. javitensis.
tinifolia, Rafin. l. c. = Cas. tinifolia.

CHETROPIS, Rafin. Fl. Tellur. iii. 80 (1836) = **Arenaria**, Rupp. (Caryophyll.).
setacea, Rafin. l. c. = A. rostrata.

CHEVALIERIA, Gaudich. Voy. Bonite Bot. t. 61, 62 (1852?) = **Asclepias**, Ruiz & Pav. (Bromel.).
crocophylla, E. Morr. in Belg. Hortic. (1885) 81.—Bras.
Germinyana, Carr. in Rev. Hortic. (1861) 230.—N. Granat.
grandiceps, Griseb. in Goett. Abh. xxiv. (1879) 320 = Ac. brasiliensis.
lingulata, Griseb. Fl. Brit. W. Ind. 591 = Ae. lingulata.
ornata, Gaudich. Voy. Bonite Bot. t. 62 = Ae. ornata.
sphærocephala, Gaudich. l. c. t. 61 = Ae. sphærocephala.
Vetchii, E. Morr. in Belg. Hortic. xxviii (1878) 177. t. 9 = Ae. Vetchii.

CHEVALLIERA, Carr. in Rev. Hortic. (1881) 280 = prsec.

CHEVREULIA, Cass. in Bull. Soc Philom. (1817) 69; et Dict. Sc. Nat. viii. 516 (1817). *COMPOSITÆ*, Benth. & Hook. f. ii. 303.
LEUCOPHYLUM, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 124.
acuminata, *Less. in Linnæa*, v. (1830) 261.—Am. austr.
filiformis, Hook. & Arn. in Hook. Comp. Bot Mag. i. (1835) 102 = acuminata.
gnaphalioides, *D. Don, ex Hook. & Arn. in Hook. l. c.*—Chili.
lanceolata, *Remy, in C. Gay, Fl. Chil.* iii. 333.—Chili.
longipes, Wedd. Chlor. And. i. 157 = acuminata.
lycopodioides, *DC. Prod* vii. 46.—Ins. Falkland.
nivea, *Phil. in Anal. Univ. Chil.* ii. (1862) 880.—Chili.
pusilla, DC. Prod. vii. 45 = stolonifera.
stolonifera, *Cass. in Bull. Soc. Philom.* (1817) 69.—Am. austr.
Thouarsii, Remy, in C. Gay, Fl Chil. iii. 332. t. 37. f. 2 = stolonifera.
xeranthemoides, *D. Don, ex Hook. Comp. Bot. Mag.* i. (1835) 102 —Chili.

CHEVNIA, J. Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 56 = **Balaustion**, Hook. (Myrtac.).
pulchella, J. Drumm. l. c. = B. pulcherrimum.

CHIAZOSPERMUM, Bernh. in Linnæa, viii. (1833) 465 = **Hypecoum**, Linn. (Fumariac.).
erectum, Bernh. l. c. xii. (1838) 602 = H. erectum.
lactiflorum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 143.—Soongaria.
pendulum, Bernh. ex Edgew. in Journ. Linn. Soc. vi. (1862) 183 = H. procumbens.

CHIBACA, Bertol. f. in Mem. Acc. Bolog. iv. (1853) 645, nomen (= Gen. dub. LAURIN.) Benth. & Hook. f. iii. 190.
salutaris, *Bertol. f. l. c.; et in Rosenth. Syn. Pl. Diaphor.* i. 238.—Afr. austr.

CHICHAEA, Presl, Rel. Haenk. ii. 140 (1836) = **Sterculia**, Linn.
acerifolia, Presl, l. c. 141 = S. acerifolia.

CHICHARRONIA, A. Rich Ess. Fl. Cub. 329. t. 43 (1845) = **Terminalia**, Linn. (Combretac.).
intermedia, A. Rich. l. c.—Ins. Cuba.

CHICKEBASSIA, A. Juss. in Mém. Mus. Par. xix. (1830) 251. t. 29 (Chokrasia). *MELIACEÆ*, Benth. & Hook. f. i. 339.
PLAGIOTAXIS, Wall. Cat. (nomina) nn. 7269, 7270.
Nimmonii, K. Grah. in Wight, Illustr. 148 = tabularis, tabularis, *A. Juss. in Mém. Mus Par* xix. (1830) 251. —Ind. or.; Malaya.
trilocularis, M. Roem. Syn. Hesper. i. 125 = tabularis.
velutina, Wight & Arn. Prod. 128 = tabularis.

CHICOINAEA, Comm. ex DC. Prod. iv 462 (1830) = **Pasthura**, Comm. (Rubiac.)

CHILECHIUM, Pfeiff. Nom. i. 1. 713 (1873) = CHILOCHIUM, Rafin. = **Echiochilon**, Desf. (Boragin.).

CHILEDRANTHEMUM, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 166. *ACANTHACEÆ*, Benth. & Hook. f. ii. 1119.
trifidum, *Oerst. l. c.* 167. t. 8. f. 30-33.—Mexic.

CHILIADENUS, Cass. in Dict. Sc. Nat. xxxiv. 34 (1825) = **Jasonia**, Cass. (Compos.).
campestratus, Cass. l. c. 35 = J. glutinosa.
glutinosus, Fourr. in Ann. Soc. Linn. Lyon, N.S. xvii. (1869) 93 = J. glutinosa.

CHILIANDRA, Griff. Notul. iv. 150 (1854) = **Rhynchotechum**, Blume (Gesnerac.).
obovata, Griff. l. c. = R. latifolium.

CHILIANTHUS, Burch. Trav. i. 94 (1822). *LOGANIACEÆ*, Benth & Hook. f. ii. 793.
SEMNOS, Rafin. Sylva Tellur. 161 (1838).
arborius, Benth. in DC. Prod. x. 485 = olacacea.
corrugatus, *Benth. l. c.* 486.—Afr. austr.
dysophyllus, *Benth. l. c.*—Afr. austr.
lobulatus, *Benth. l. c.*—Afr. austr.
oleaceus, *Burch. Trav.* i. 94.—Afr. austr.
trophyllus, E. Mey. Zwei Pfl. Docum. 172 = Nuxia floribunda.

CHILIOCEPHALUM, Benth. in Hook. Ic. Pl t. 1137 (1873). *COMPOSITÆ*, Benth. & Hook f. ii. 300.
KRALIKIA, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 151 (1867).
myriocephalum, Baill. Hist. Pl. viii. 173 = Schimperi.
Schimperi, *Benth. in Hook. Ic. Pl. t.* 1137.—Afr. trop.
Schultzii, Buching. ex Baill. Hist. Pl. viii. 173 = densifolium.

CHILIOPHYLLUM, DC. Prod. v. 554 (1836) = **Zalusania**, Pers. (Compos.).
globosum, DC. l. c. = Z. globosa.

CHILIOPHYLLUM, Phil. in Linnæa, xxxiii. (1864-65) 132. *COMPOSITÆ*, Benth. & Hook. f. ii. 258.
densifolium, *Phil. l. c.* 133.—Chili.

CHILIOTRICHUM, Cass. in Dict. Sc. Nat. viii. 576 (1817). *COMPOSITÆ*, Benth. & Hook. f. ii. 276
TROFIDOLEPIS, Tausch, in Flora, xli. (1829) 68.
amelloides, DC. Prod. v. 218 = seq.
amelloideum, *Cass. in Dict. Sc. Nat.* viii. 577.—Reg. Magellan.
Darwinii, Hook. f. Fl. Antarct. 305 = Nardophyllum Darwinii.
Feliciæ, Hombr & Jacquin. ex Decne. Bot. Voy. Astrol. et Zél. 52 = amelloideum.
humile, Hook. f. Fl. Antarct. 304 = Nardophyllum humile.
Kingii, *Hook. f. l. c.* 305 = Nardophyllum Kingii.
ovatifolium, *Hombr. & Jacquin. ex Decne. Bot. Voy. Astrol. et Zél. Dicot. Phan.* t. 28.—Hab. ? rosmarinifolium, *Less. in Linnæa*, vi. (1831) 109.—Chili, virgatum, *Phil. in Anal. Univ. Chil.* (1873) 455.—Chili.

CHILMORIA, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 500 = **Gymnoarvita**, R. Br. (Bixin.).
dodecandra, Buch.-Ham. l. c.—G. odorata.
pentandra, Buch.-Ham. l. c. 501 = Hydnocarpus laurifolia.

CHILOCALYX, Klotzsch, in Peters, Reise Mossamb. Bot. 154 (1861) = **Cleome**, Linn. (Capparid.).
macrophyllus, Klotzsch, l. c. 155 = Cleome Chilocalyx.
tenuifolius, Klotzsch, l. c. = Cleome Chilocalyx.

CHILOCALYX, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 583 = **Atalantia**, Correa (Rutac.).
elliptica, Turcz. l. c. = A. missionis.

CHILOCARPUS, Blume, Cat. Gew. Buitenz. 32 (1823); et Bijdr. 1025 (1826). *APOCYNACEÆ*, Benth. & Hook. f. ii. 691.
atroviridis, *Blume, Mus. Bot. Lugd. Bat.* i. 153.—Burma; Malaya.
australis, *F. Muell. Fragm.* ii. 90.—Austral.
ceylanicus, *Wight, Ic.* t. 1288 = Willughbeia ceylanica
compositus, *Blume, Mus. Bot. Lugd. Bat.* i. 152.—Java.
costatus, *Miq. Fl. Ind. Bat.* ii. 393.—Sumatra.
cuspidatus, Benth. & Hook. f. Gen. ii. 698. in nota sub Hunteria = Melodinus orientalis
decipiens, *Hook. f. Fl. Brit. Ind.* iii. 627.—Malacca.
denaiflorus, *Blume, Mus. Bot. Lugd. Bat.* i. 152.—Sumatra.
denudatus, Blume, Bijdr. 1025 = atroviridis.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 552 — Sumatra.
enervis, *Hook. f. Fl. Brit. Ind.* iii. 626.—Malacca.
flavescens, Kew Gard. Report (1880) 47, ex Fl. Brit Ind. iii. 625 = Willughbeia flavescens.
globuliferus, *Blume, Mus. Bot. Lugd. Bat.* i. 152 — Borneo.
Maingayi, *Dyer, ex Hook. f. Fl. Brit. Ind.* iii. 627.—Malacca.
malabaricus, Bedd. Ic. Pl. Ind. Or. i. 38 = atroviridis.
suaveolens, *Blume, Cat. Gew. Buitenz.* 23; *et Bijdr.* 1025.—Java.
vermicosus, *Blume, Mus. Bot. Lugd. Bat.* i. 152.—Borneo.

CHILOCHIUM, Rafin. in Ann. Gén. Sc. Phys. viii. (1821) 269 = **Echiochilon**, Desf. (Boragin.).

CHILOCHLOA, Beauv. Agrost. 37. t. 7. f. 2 (1812) = **Phleum**, Linn. (Gramin.).
aurasa, Roem. & Schult. Syst. ii. 410 = P. asperum
arenaria, Beauv. Agrost. 37 = P. arenarium.
aspera, Beauv. l. c. = P. asperum.
Boemeri, Beauv. l. c. = P. Boehmeri.
collina, Schur, Enum. Pl. Transs 729 = P. phalaroides.
cuspidata, Beauv. Agrost. 37 = P. Michelii.
cuspidata, Schur, Enum. Pl. Transs. 729 = P. phalaroides.
dentata, Trin. Gram. Unifl. 168 = Phalaris dentata.
explicata, Roem. & Schult. Syst. ii. 410 = Heleochloa alopecuroides.
Gerardi, Bess. ex Schult Mant. ii. 206 = Alopecurus Gerardi.
hirsuta, S. F. Gray, Nat. Arr. Brit. Pl. ii. 189 = P. Michelii.
hispida, Beauv. Agrost. 158, nomen = Arthraxon ciliaris.
Michelii, Trin. Gram. Unifl. 167 = P. Michelii.
paniculata, Beauv. Agrost. 37 = P. asperum.
ventricosa, Beauv. ex Steud. Nom. ed. II. i. 350 = Gastridium australe?

CHILODIA, R. Br. Prod. 507 (1810) = **Prostanthera**, Labill. (Labiat.).
australis, Lond. Hort. Brit. Suppl. i. 585.—Austral.
scutellarioides, R. Br. Prod. 507 = P. empetrifolia.

CHILOGLOSSA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 160 = **Dianthera**, Linn. (Acanthac.).
glabra, Oerst. l. c. 161 = D. glabra.
secunda, Trin. Ic. Ind. ii. 77 = prsec.

CHILOGLOTTIS, R. Br. Prod. 323 (1810). *ORCHIDEÆ*, Benth. & Hook. f. iii. 514.
comuta, *Hook. f. Fl. Antarct.* 69.—N. Zel.
diphylla, *R. Br. Prod.* 323.—Austral.
formicifera, *Fitzgerald, Austral. Orchids*, i. 111. (1877), —Austral.
Gunnii, *Lindl. Gen. et Sp. Orch.* 387.—Austral.
trapeziformis, *Fitzgerald, Austral. Orchids*, i. 111. 1877. —Austral.
trilabra, *Fitzgerald, in Journ. Bot.* xxi. (1883) 204.—Austral.
Trimenii, F. Muell. Veg. Chath.-Isl. 61 = Caladenia bifolia.

CHILOPORUS, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 57 = **Miconia**, Ruiz & Pav. (Melastomac.). *andinus*, Naud. l. c. = M. andina.

CHILOPSIS, D. Don, in Edinb. Phil. Journ. ix. (1823) 261. *BIGNONIACEÆ*, Benth. & Hook. f. ii 1041 glutinosa, *Engelm. in Bot. Wislizen. Mexic. Exped.* 94.—Mexic. bor.

linearis, Sweet, Hort. Brit. ed. I. 383 = saligna.

saligna, *D. Don, in Edinb. Phil. Journ.* ix. (1823) 261. —Mexic.

CHILOSCHISTA, Lindl. Bot. Reg. sub t. 1522 (1832) = **Sarcochilus**, R. Br. (Orchid.).

unicolor, Lindl. l. c. = S. unicolor.

CHILOSTIGMA, Hochst. in Flora, xxiv. (1841) 372 = **Aptosimum**, Burch. (Scrophularin.).

pumilum, Hochst. l. c. = A. pumilum.

CHIMANTHUS, Rafin. Fl. Ludov. 26 (1817) = **Prunus**, Tourn. (Rosac.).

amygdalina, Rafin. l. c. = P. caroliniana.

CHIMAPHILA, Pursh, Fl. Am. Sept. i. 279 (1814). *ERICACEÆ*, Benth. & Hook. f. ii. 603.

CHIMAZA, R. Br. ex DC. Prod. vii. 776 (1839). PYRVA, Rafin. in Journ. Phys. Sc. lxxxix. (1819) 261, in obs.

astyla, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 484. —Japon.

corymbosa, Pursh, Fl. Am. Sept. i. 300 = umbellata.

cymosa, J. & C. Presl, Fl. Cech. 89 = umbellata.

Durandi, *Rafin. Atl. Journ.* 118.—Am. bor.

japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 165.— Japon.

maculata, *Pursh, Fl. Am. Sept.* i. 300.—Am. bor

Menziesii, *Spreng. Syst.* ii. 817.—Am. bor. occ.

umbellata, *Nutt. Gen. Am.* i. 274.—Europ.; As. bor.; Am. bor.

CHIMARHIS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 84 *et seq.*

CHIMARRHIS, Jacq. Stirp. Am. 61 (1763). *RUBIACEÆ*, Benth. & Hook. f. ii. 45.

cymosa, *Jacq. l.*—Ins. Martin.

Gondotii, *Baill. Adansonia*, xii. (1879) 307.—N. Granat.

parsenata, *Baill. l. c.* 308.—Bras.

turbinata, *DC. Prod.* iv. 404.—Guiana.

CHIMAZA, R. Br. ex DC. Prod. vii. 776 (1839) = **Chimaphila**, Pursh (Ericac.).

CHIMOCARPUS, Baill. Dict. i. 784 (1879) = CHYMOCARPUS, D. Don = **Tropæolum**, Linn. (Geraniac.).

CHIMONANTHUS, Lindl. Bot. Reg. sub t. 404 (1819); et t. 451 (1830). *CALYCANTHACEÆ*, Benth. & Hook. f. i. 16.

MERATIA, Loisel. Herb. Gén. Amat. iii. t. 178 (1819).

fragrans, *Lindl. l. c.* t. 451.—Japon.

grandiflorus, Stend. Nom. ed. II. i. 850 = fragrans.

hiemalis, Hort. ex Bielawski, in Ann. Soc. Linn. Maine-et-Loire, ix. (1887) 96 = fragrans.

parviflorus, Rafin. Alsog. Am. 6 = fragrans.

praecox, Lindl. Bot. Reg. sub t. 404 = fragrans.

zeilanicifolius, Steud. Nom. ed. II. i. 850, sphalm. = Myrospyrum zeilanicifolium.

verus, Hort. ex Bielawski, in Ann. Soc. Linn. Maine-et-Loire, ix. (1887) 96 = fragrans.

CHIMOPHILA, Radius, Dis. Fyrol. 7, 53 (1821) = **Chimaphila**, Pursh (Ericac.).

CHINCHARRONIA, A. Rich. Fl. Cub. Fanerog. ii. 226 (1853) = CHICHARRONIA, A. Rich. = **Verbenalia**, Linn. (Combretac.).

CHINCHONA, Howard, in Rep. Bot. Cosgr. London, (1866) 221 = **Cinchona**, Linn. (Rubiac.).

CHIOCOCCA, P. Br. ex Linn. Syst. ed. X. 917 (1759). *RUBIACEÆ*, Benth. & Hook. f. ii. 106.

SIPHONANDRA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 581.

alternifolia, Linn. Syst. ed. XII. 165 = Cestrum vespertinum.

CHIOCOCCA:—

anguifuga, Mart. Reise Bras. i. 806, 554; Mat. Med. Bras. 17. t. 6 = brachiata.

axillaris, Moç. ex DC. Prod. iv. 488 (= *Margaris nudiflora*).—Mexic.

barbata, *Ferst. f. Prod.* 16.—Ins. Pacif.

barbigera, D. Dietr. Syn. Pl. i. 777 (= Symphorocarpos microphyllus.

brachiata, *Ruiz & Pav. Fl. Per.* ii. 67.—Am. trop.

cortacea, *Mart. & Gal. in Bull. Acad. Brux* xi. i. (1844) 231.—Mexic.

densifolia, Mart. Reise Bras. i. 544; Spec. Mat Med. Bras. 17. t. 6 = brachiata.

floridana, Rafin. Alsog. Am. 75 = racemosa.

javanica, *Blume, Bijdr.* 968.—Java.

latifolia, Rafin. Alsog. Am. 75 = racemosa.

macrocarpa, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 230.—Mexic.

malabarica, *Dennst. Schlüssv. Hort. Malab.* 37.—Ind. or.

nitida, *Benth. in Hook. Journ. Bot.* iii. (1841) 256.— Guiana.

nocturna, Jacq. Enum. Pl. Carib. 16 = Cestrum nocturnum.

nudiflora, D. Dietr. Syn. Pl. i. 777 (= *Margaris nudiflora*).—Mexic.

odorata, *Hook. & Arn. Bot. Beech. Voy.* 65.—Ins. Societ.

opulina, Spreng. Syst. i. 756 = Pavetta opulina.

paniculata, Hoffm. ex Roem & Schult. Syst v. 202 = brachiata.

paniculata, Linn. f. Suppl. 145 = Psychotria paniculata.

parviflora, Humb. & Bonpl. ex Roem & Schult. Syst. v. 202 = brachiata.

parvifolia, Wallich. ex Griseb. Fl. Brit. W. Ind. 337 = racemosa.

phaenostemon, *Schlecht. in Linnaea*, ix. (1834) 594.— Mexic.

pubescens, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 202 = brachiata.

racemosa, H. B. & K. Nov. Gen. et Sp. iii. 352 = brachiata.

racemosa, *Linn. Syst.* ed. X. 917; *Jacq. Select. Am.* 68.—Ind. occ.

racemosa, Schlecht. & Cham. in Linnaea, v. (1830) 166 = phaenostemon.

sambucina, Spreng. Syst. i. 756 = Pavetta sambucina.

spicata, Blume, Bijdr. 965 = Xanthophytum spicatum.

staminea, *Mart. & Gal. Bull. Acad. Brux.* xi. i. (1844) 231.—Mexic.

trifora, Spreng. Syst. i. 756 = Pavetta trifora.

trisperma, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 219.—Ins. Galapag.

CHIOGENES, Salisb. in Trans. Hort. Soc. ii. (1817) 94. *VACCINIACEÆ*, Benth. & Hook. f. ii. 577.

GLYCIPHYLLA, Rafin. in Am. Monthly Mag. (1819) 192.

LASIERPA, Torr. in Geol. Rep. N. York (1839), 152. PHALEROCARPUS, G. Don, Gen. Syst. iii. 641 (1834).

hispidula, Torr. & Gray, ex Torr. Fl. N. York, i. 480. t. 68 = serpyllifolia.

japonica, *A. Gray, Syn. Fl. N. Am.* ii. 1. 26.—Japon.

serpyllifolia, *Salisb. in Trans. Hort. Soc.* ii. (1817) 94. —Am. bor.; Japon.

CHIONACENE, R. Br. in Benn. Pl. Jav. Rar. 15 (1838). *GRAMINEÆ*, Benth. & Hook. f. iii. 1113.

barbata, *R. Br. l. c.* 18.—Ind. or.

cynthopoda, *F. Muell. ex Benth. Fl. Austral.* vii. 516. —Austral.

Kornigii, Thw. Enum. Pl. Zeyl. 357 = barbata.

Wightii, *Munro, ex Benth. & Hook. f. Gen.* iii. 1113, nomen.—Ind. or.

CHIONANTHUS, Gaertn. Fruct. i. 189. t. 39 (1788) = **Linociera**, Sw. (Oleac.).

acuminata, Wall. ex DC. Prod. viii. 298 = L. acuminata.

acuminigera, F. Muell. Fragm. viii. 42.—Austral.

albidiflora, Thw. Enum. Pl. Zeyl. 189 = L. albidiflora.

antillaris, R. Br. Prod. 532 = L. albidiflora!

axillaris, Roxb. ex Wall. Cat. sub n. 2894 = L. macrophylla.

callophylla, Blume, Mus. Bot. Lugd. Bat. i. 319.— Malaya.

caribaea, Jacq. Coll. ii. 110 = L. compacta.

compacta, Sw. Prod. Veg. Ind. Occ. 18 = L. compacta.

cotinifolia, Willd. Sp. Pl. i. 47 = L. cotinifolia.

CHIONANTHUS:—

courtallensis, Bedd. For. Man. Bot. 155 = Linociera leprocarpa.

cuspidata, Blume, Mus. Bot. Lugd. Bat. i. 319.—Ins. Celebes.

densiflora, Zoll. & Morr. in Nat. en Geneesk. Arch. Neêrl. Indie, ii (1846) 4 = L. densiflora.

dichotoma, Roxb. Hort. Beng. 3; Fl. Ind. i. 108 = L. purpurea.

dispersifolia, Miq. Fl. Ind. Bat. Suppl. 559.—Sumatra.

domingensis, Lam. Tabl. Encyc. i. 30 = L. latifolia.

effusiflora, F. Muell. Fragm. iv. 33 = L. macrophylla.

elliptica, Blume, Mus. Bot. Lugd. Bat. i. 319.—Malaya.

Ghaeri, Gaertn. Fruct. i. 190 = L. cotinifolia.

glomerata, Blume, Mus. Bot. Lugd. Bat. i. 318.— Sumatra.

guianensis, Pers. Syn. i. 9.—Guiana.

incrassata, Sw. Prod. Veg. Ind. Occ. 18; Willd. Sp. Pl. i. 47 = L. tetrandra.

insignis, Miq. Fl. Ind. Bat. Suppl 559 = L. insignis.

intermedia, Bedd. Fl. Sylv. t. 239 = L. intermedia.

laxiflora, Blume, Mus. Bot. Lugd. Bat. i. 319.—Borneo

leprocarpa, Thw. Enum. Pl. Zeyl. 189 = L. leprocarpa.

ligustrina, Pers. Syn. i. 9 = L. ligustrina.

littorea, Miq. Fl. Ind. Bat. Suppl. 559.—Sumatra.

lunonica, Blume, Mus. Bot. Lugd. Bat. i. 319 = L. lunonica.

macrocarpa, Blume, l. c. = L. insignis.

macrophylla, Blume, l. c. 317; Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 129 = L. macrophylla.

malabarica, Heyne, ex Wall. Cat. sub n. 2898 = L. malabarica.

Mayepea, Vahl, Symb. Bot. ii. 1 = L. tetrandra.

monstiflora, Kurz, For. Fl. Brit. Burma, ii. 159 = L. minutiflora.

montana, Blume, Bijdr. 681 = L. montana.

montana, Kurz, For. Fl. Brit. Burma, ii. 159 = L. insignis.

monticola, Blume, Mus. Bot. Lugd. Bat. i. 317 = L. montana.

palembanica, Miq. Fl. Ind. Bat. Suppl. 558 = L. pauciflora.

picrophloia, F. Muell. Fragm. iii. 189. t. 24 = L. intermedia.

pubescens, H. B & K. Nov. Gen. et Sp. iii. 285.— N. Granat.

purpurea, Lam. Tabl. Encyc. i. 30 = L. purpurea.

quadristaminea, F. Muell. Fragm. viii. 41.—Austral.

ramiflora, Roxb. Hort. Beng. 3; Fl. Ind. i. 107 = L. macrophylla.

rostrata, Miq. Fl. Ind. Bat. ii. 553 = L. rostrata.

rostrata, Thw. Enum Pl. Zeyl. 189 = L. albidiflora.

smilaefolia, Wall in Roxb. Fl. Ind. ed. Carey, i. 108 = Myrospyrum zeilanicifolium.

spicata, Blume, Mus. Bot. Lugd. Bat. i. 318.—Malaya.

sumatrana, Blume, l. c.—Sumatra.

tenuiflora, Wall. ex DC. Prod. viii. 298 = L. macrophylla.

ternifiora, Griff. Notul. iv. 740 = L. ternifiora.

tetrandra, Vahl, Enum. i. 48 = L. tetrandra.

timorensis, Blume, Mus. Bot. Lugd. Bat. i. 318.— Timor.

zeylanica, Heyne, ex Wall. Cat. sub n. 2898 = L. malabarica.

zeylanica, Lam. Tabl. Encyc. i. 30 = L. cotinifolia.

zeylanica, Linn. Sp Pl. 8 = L. purpurea.

CHIONANTHUS, Royen, ex Linn. Gen. ed. I. 886 (1737). *OLEACEÆ*, Benth. & Hook. f. ii. 677.

chinensis, Maxim. in Bull. Acad. Pétersb. xx. (1875) 480 = retusa.

angustifolia, Rafin. New Fl. Am. iii. 88 = virginica.

fraxinifolia, Kellogg, in Proc. Calif. Acad. v. (1873) 16 = Fraxinus dipetala.

fragrans, Edwards, ex Steud. Nom. ed. II. i. 851 = virginica.

heterophyla, Rafin. New Fl. Am. iii. 87 = virginica.

latifolia, Ait. ex Steud. Nom. ed. II. i. 850 = virginica.

longifolia, Rafin. New Fl. Am. iii. 88 = virginica.

maritima, Sweet, Hort. Brit. ed. II. 351 = virginica.

montana, Rafin. New Fl. Am. iii. 87 = virginica.

obovata, Rafin. l. c. = virginica.

retusa, *Paxt. Flow. Gard.* iii. (1852-53) 88.—China.

trifida, Moench, Meth. 478 = virginica.

trifora, Stokes, Bot. Mat. Med. i. 19 = virginica.

vernalis, Salisb. Prod. 14 = virginica.

verna, Baill. Hist. Pl. i. 295 = virginica.

virginica, *Linn. Sp. Pl.* 8.—Am. bor.

CHIONE, DC. Prod. iv. 461 (1830). *RUBIACEÆ*, Benth. & Hook. f. ii. 107.
CRUSEA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 304. t. 19. f. 1.
SACCONIA, Endl. Gen. Pl. 541 (1838).
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 22.—Ins. Cuba.
elliptica, *Griseb. Pl. Wright. Cub.* 507.—Ind. occ.
glabra, *DC. Prod.* iv. 461.—Ind. occ.
lucida, *Griseb. Pl. Wright. Cub.* 507.—Ind. occ.
myrtifolia, *Griseb. l. c.*—Ind. occ.
vaccinifolia, Griseb. Cat. Pl. Cub. 183 = Erithalis vaccinifolia.

CHIONE, Salisb. Gen. Pl. Fragm. 100 (1866) = **Narcissus**, Tourn. (Amaryllid.).
italica, Salisb. l. c. = N. italicus.
papyracea, Salisb. l. c. = N. papyraceus.

CHIONICE, Bunge, ex Ledeb. Fl. Ross. ii. 33 (1844) = **Potentilla**, Tourn. (Rosac.).
dieica, Bunge, l. c. = P. tetrandra.

CHIONODOXA, Boiss. Diagn. Ser. I. v. 61 (1844). *LILIACEÆ*, Benth. & Hook. f. iii. 813.
cretica, *Boiss. & Heldr. l. c.* xiii. 24.—Ins. Creta.
cretica, Jaub. & Spach, Illustr. v. 46 t. 443 = nana.
Forbesii, Baker, in Journ. Linn. Soc. xi. (1871) 436 = Luciliæ.
Luciliæ, *Boiss. Diagn.* Ser. I. v. 61.—As. Min.
nana, *Boiss. & Heldr. l. c.* xiii. 24.—Ins. Creta.

CHIONOGRAPHIS, Maxim. in Bull. Acad. Pétersb. xi. (1867) 485. *LILIACEÆ*, Benth. & Hook. f. iii. 826.
japonica, *Maxim. l. c.* 486.—Japon.

CHIONOLAENA, DC. Prod. v. 297 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 302.
arbuscula, *DC. l. c.*—Bras.
corymbosa, *Hrmsl. Diagn. Pl. Nov.* 32.—Mexic.
Glaziovii, *Baker, in Mart. Fl. Bras.* vi. iii. 130.—Bras.
glomerata, *Baker, l. c.*—Bras.
Isabellae, *Baker, l. c.*—Bras.
latifolia, Baker, l. c. 132 = Leucopholis latifolia.
lavandulaceum, Hemsl. Biol. Centr. Am. Bot. ii. 134 = seq.
lavandulifolium, *Benth. & Hook. f. Gen.* ii. 302.—Mexic.
longifolia, *Baker, in Mart. Fl. Bras.* vi. iii. 131.—Bras.
lychnophoroides, *Sch. Bip. in Pollichia*, xx-xxi. (1863) 391.—Bras.
phyllicoides, Baker, in Mart. Fl. Bras. vi. iii. 131 = Leucopholis phyllicoides.
Wittigiana, *Baker, l. c.* 129.—Bras.

CHIONOPAPPUS, Benth. in Benth. & Hook. f. Gen. ii. 485 (1873). *COMPOSITÆ*, Benth. & Hook. f. l. c.

CHIONOPHILA, Benth. in DC. Prod. x. 331 (1846). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 942.
Jamesii, *Benth. l. c.*—Am. bor. occ.

CHIONOPHILA, Miers, ex Lindl. Veg. Kingd. 701 (1847) = **Boopis**, Juss. (Calycer.).

CHIONOPTERA, DC. Prod. vii. 14 (1838) *COMPOSITÆ*, Benth. & Hook. f. ii. 403.
Gayophyta, *DC. l. c.*—Chili.

CHIONOTHRIX, Hook. f. in Benth. & Hook. f. Gen. iii. 33 (1880). *AMARANTACEÆ*, Benth. & Hook. f. l. c.
somaliensis, *Hook. f. l. c.*—Afr. trop.

CHIONOTRIA, Jack. in Malay. Misc. ii. (1822) vii. 53 = **Glyocosmis**, Correa (Rutac.).
monogyna, Walp. Rep. i. 893 = seq.
rigida, Jack, in Malay Misc. ii. (1822) vii. 54 = G. pentaphylla.

CHIOPHILA, Rafin. Fl. Tellur. iii. 28 (1836) = **Gentiana**, Linn.
nivalis, Rafin. l. c. = G. nivalis.

CHIRANTHODENDRON, Cerv. ex Cav. in Anal. Cienc. Nat. vi. (1803) 303, adnot. = **Chiranthodendron**, Humb. & Bonpl. (Sterculiac.).
californicum, Baill. Hist. Pl. iv. 70 = Fremontia californica.
pentadactylon, Larréat, Desc. Chiranth. ed. Lesc. 17 = Chiranthodendron platanoides.

CHIRATA, G. Don, Gen. Syst. iv. 659 (1837) = **Chirita**, Buch.-Ham.

CHIRATIA, Montr. in Mém. Acad. Lyon. x. (1860) 202 = **Fonsecaea**, Linn. f. (Lythrar.).
leucantha, Montr. l. c. 208.—N. Caled.

CHIRITA, Buch.-Ham. in D. Don, Prod. Fl. Nep. 89 (1825). *GESNERACEÆ*, Benth. & Hook. f. ii. 1092.
BARACTES, DC. Prod. ix. 260 (1845).
CALOSACME, Wall. Cat. nn. 800 ad 806 (1829).
CHIRATA, G. Don, Gen. Syst. iv. 659 (1837).
GNATOSTEMON, Regel, Gartenfl. xv. (1866) 353.
HYPOPTERON, Hassk. Cat. Hort. Bog. Alt. 154 (1844).
LIEBIGIA, Endl. Gen. 1407 (1841).
MORSTDORFFIA, Steud. Nom. ed. II. ii. 161 (1841).
TROSTDORFFIA, Blume, Bijdr. 762 (1826).
acaminata, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Reg. Himal.
acuminata, A. DC. Prod. ix. 269 = Didymocarpus acuminata.
amplectens, Steud. Nom. ed. II. i. 851 = bifolia.
anachoretica, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 231.—China.
bifolia, *D. Don, Prod. Fl. Nep.* 90.—Reg. Himal.
Bilabium, *C. B. Clarke, in A. DC. Monog. Phan.* v. 127.—Sumatra.
Blumei, *C. B. Clarke, l. c.* 122.—Java.
bracteosa, Miq. Fl. Ind. Bat. ii 728 = Horsfieldii.
brevipes, *C. B. Clarke, in A. DC. Monog. Phan.* v. 130.—Ind. or.
caerulea, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Java.
caliginosa, *C. B. Clarke, in A. DC. Monog. Phan.* v. 124.—Malacca.
calva, *C. B. Clarke, l. c.* 126.—Reg. Himal.
Clarkei, *Hook. f. Fl. Brit. Ind.* iv. 350.— Reg. Himal.
communis, Gardn. in Calc. Journ. Nat. Hist. iv. (1846) 481 = zeylanica.
cortusifolia, *Hance, in Journ. Bot.* xxi. (1883) 324.—China.
diaphana, Royle, Illustr. Bot. Himal. 204 = pumila.
dielsiana, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Reg. Himal.
eburnea, *Hance, in Journ. Bot.* xxi. (1883) 168.—China.
Edgeworthii, A. DC. in Prod. ix. 269 = pumila.
Fauriei, *Franch. in Bull. Soc. Linn. Par.* i. (1885) 459.—China.
fimo, K. Br. in Benn. Pl. Jav. Rar. 117 = pumila.
fusca, *C. B. Clarke, in A. DC. Monog. Phan.* v. 129.—Borneo.
glabra, C. B. Clarke, Comm. et Cyrt. Beng. t. 73 = Java.
glabra, *Miq. Fl. Ind. Bat.* ii. 729.—Malaya.
grandiflora, Wall. Pl. As. Rar. i. 48 = urticaefolia.
Griffithii, Hook. f. & Thoms. ex C. B. Clarke, in A. DC. Monog. Phan. v. 60 = Loxostigma Griffithii.
hamosa, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Ind.
Hookeri, *C. B. Clarke, Comm. & Cyrt. Beng.* t. 70.—Reg. Himal.
Horsfieldii, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Java.
Horsfieldii, De Vriese, Pl. Ind. Bat. Reinw. 11 = Blumei.
humilis, *Miq. Fl. Ind. Bat.* ii. 729.—Sumatra.
insignis, *C. B. Clarke, in A. DC. Monog. Phan.* v. 121. —Assam.
Julise, *Hance, in Journ. Bot.* xxi. (1883) 168.—China.
Kurzii, *C. B. Clarke, in Journ. Linn. Soc.* xv. (1876) 145.—Reg. Himal.
lachnocaula, *C. B. Clarke, in A. DC. Monog. Phan.* v. 118.—Reg. Himal.
lilacina, *Lem. Illustr. Hortic.* (1869) t. 668.—Ind. or.
macrosiphon, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 231.—China.
mollis, *Miq. Fl. Ind. Bat.* ii. 728.—Sumatra.
monophylla, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 360.—Reg. Himal.
Moonii, *Gardn. in Calc. Journ. Nat. Hist.* vi. (1846) 479.—Zeylan.
obtusa, *C. B. Clarke, in A. DC. Monog. Phan.* v. 114. —China.
polycarpa, *Steud. Nom.* ed. II. i. 351.—Ind. or.
polymorpha, Miq. Fl. Ind. Bat. ii. 728 = Horsfieldii.
primuloides, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 82. —Reg. Himal.
pumila, *D. Don, Prod. Fl. Nep.* 90.—Reg. Himal.
scaberrima, *R. Br. in Benn. Pl. Jav. Rar.* 117.— Java.
scabra, Royle, ex C. B. Clarke, in A. DC. Monog. Phan. v. 113 = bifolia.
sinensis, *Lindl. Bot. Reg.* (1844) t. 59.—Hongkong.
speciosa, *Kurz, in Journ. Bot.* xi. (1873) 195.— China.
spectabilis, *Miq. Fl. Ind. Bat. Suppl.* 564.—Sumatra.
ternifolia, Bech.-Ham. ex D. Don, Prod. Fl. Nep. 121 = Lysionotus serrata.
urticaefolia, *Buch.-Ham. ex D. Don, l. c.* 90.—Reg. Himal.
vulgaris, Hort. ex Belg. Hortic. iii. (1853) t. [36].— Cf. p. 288 = zeylanica.
Walkeri, *Gardn. in Calc. Journ. Nat. Hist.* vi. (1846) 480.—Zeylan.
Walkeriae, Hook. Bot. Mag. t. 4827 = Walkeri.
zeylanica, *Hook. l. c.* t. 4182.—Zeylan.
Zollingeri, *C. B. Clarke, in A. DC. Monog. Phan.* v. 125.—Java.

CHIROCALYX, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 97 = **Brythrina**, Lino. (Legumin.).
abyssinicus, Hochst. in Flora, xxix. (1846) 600 = E. abyssinica.
Candolleanus, Walp. in Flora, xxxvi. (1853) 148 = E. spathacea.
divaricatus, Walp. l. c. = E. divaricata.
indicus, Walp. l. c. = E. indica.
latifolius, Walp. l. c. = E. senegalensis.
mollissimus, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 98 = E. tomentosa.
pictus, Walp. in Flora, xxxvi. (1853) 148 = E. indica.
pubescens, Walp. in Linnaea, xxiii. (1850) 741 = E. fissa.
rubrinervius, Walp. in Flora, xxxvi. (1853) 148 = E. rubrinervis.
tomentosus, Hochst. in Flora, xxix. (1846) 600 = E. tomentosa.
umbrosus, Walp. in Flora, xxxvi. (1853) 148 = E. umbrosa.
volutinus, Walp. l. c. = E. velutina.

CHIRONIA, Linn. Coroll. Gen. 5 (1737). *GENTIANEÆ*, Benth. & Hook. f. ii. 805.
EUPODIA, Rafin. Fl. Tellur. iii. 29 (1836).
EVALTHA, Rafin. l. c. 77 (1836).
ONEFERA, Rafin. l. c. 80 (1836).
PLOCANDRA, E. Mey. Comm. Pl. Afr. Austr. 181 (1837).
ROELLINIA, Moench, Meth. Suppl. 211 (1802).
ROSLINIA, G. Don, Gen. Syst. iv. 308 (1837).
amoena, Rafin. in Med. Repos. N. York, v. (1808) 350 = Sabbatia gracilis ?
amoena, Salisb. Prod. 127 = linoides.
angularis, Linn. Sp. Pl. 190 = Sabbatia angularis.
angustifolia, Sims, Bot. Mag. t. 818 = frutescens.
arenaria, *E. Mey. Comm. Pl. Afr. Austr.* ii. 180.—Afr. austr.
baccata, Hoffmgg. Vers. Pfl. Nachtr. i. 211 = seq. ?
baccifera, *Linn. Sp. Pl.* 190.—Afr. austr.
baccifera, Zeyh. ex Griseb. in DC. Prod. ix. 41 = linoides.
Barrelyana, Hort. Berol. ex Griseb. Gen. et Sp. Gent. 100 = pedunculata.
brachiata, Willd. ex Griseb. l. c. 145 = Erythraea Roxburghii.
calycosa, Michx. Fl. Bor. Am. i. 147 = Sabbatia calycosa.
campanulata, Linn. Sp. Pl. 190 = Sabbatia gracilis ?

CHIRONIA :—
capensis, Blance, Fl. Filip. ed. I. 102 = Ipomoea peltata.
caryophylloides, Linn. Cent. ii. 12; Amoen. Acad. iv 308 = frutescens.
Centaurium, Schmidt, Fl. Bohem. ii. 81 = Erythraea Centaurium.
centauroides, Roxb. Hort. Beng. 16; Fl. Ind. i. 484 = Erythraea Roxburghii.
chilensis, Willd. Sp. Pl. i. 1067 = Erythraea chilensis
chloroides, Michx. Fl. Bor. Am. i. 147 = Sabbatia chloroides.
cymosa, *Burm. f. Fl. Cap. Prod.* 5.—Afr. austr.
cymosa, Lam. Encyc. i. 479 = Sabbatia lanceolata.
decandra, Walt. Fl. Carol. 95 = Sabbatia chloroides.
decussata, Vent. Jard. Cels, 81 = frutescens.
dichotoma, Walt. Fl. Carol. 95 = Sabbatia calycosa.
dodecandra, Linn. Sp. Pl. 190 = Sabbatia chloroides.
dubia, Willd. ex Griseb. Gen. et Sp. Gent. 146 = Erythraea Muehlenbergii.
emarginata, Jacq. Pl. Nov. Cap. 11 = lychnoides.
Erythraea, Schousb. Vexir. Marok. 95 = Erythraea ramosissima.
exigua, *Oliver, in Hook. Ic. Pl.* t. 1229.—Afr. austr.
filiformis, Desf. Tabl. ed. 1. 77, nomen = Microcala filiformis.
Fischeri, Hort. ex Pant. Mag. Bot. xi. (1844) 287 = floribunda.
floribunda, *Pant. l. c.* 228.—Afr. austr.
frutescens, *Linn. Sp. Pl.* 190; *Berg. Desc. Pl. Cap.* 45. —Afr. austr.
Gerardi, Schmidt, Fl. Bohem. ii. 83 = Erythraea pulchella.
glutinosa, *Pant. Mag. Bot.* xv. (1849) 245.—Austral. ?
gracilis, Michx. Fl. Bor. Am. i. 146 = Sabbatia gracilis.
grandiflora, Salisb. Prod. 137 = frutescens.
heloides, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = Erythraea elodes.
imaperta, Willd. Sp. Pl. i. 1069 = Erythraea ramosissima.
intermedia, Mérat, Fl. Paris, 91 = Erythraea ramosissima.
jasminoides, [Ker-Gawl. in] Bot. Reg. t. 197 = tetragona.
jasminoides, Lam. Illustr. i. 479. t. 108. *f.* 2 = nudicaulis.
jasminoides, *Linn. Pl. Rar. Afr.* 9 (*Amoen. Acad.* vi. 84).—Afr. austr.
Krebsii, *Griseb. Gen. et Sp. Gent.* 98.—Afr. austr.
lanceolata, Walt. Fl. Carol. 95 = Sabbatia lanceolata.
lanceanthera, Blanco, Fl. Filip. ed. II. 71 = Ipomoea peltata.
latifolia, *E. Mey. Comm. Pl. Afr. Austr.* ii. 178.—Afr. austr.
linarifolia, Loisel. Notes, 155 = Erythraea linarifolia.
linarifolia, Steud. Nom. ed. II. i. 852, in syn. sphalm. = pruec.
linifolia, Steud. l. c. 851 = Erythraea linarifolia.
linoides, *Linn. Sp. Pl.* 189.—Afr. austr.
littonia, Turn. in Turn. & Dillw. Bot. Guide, ii. 469 = Erythraea linarifolia.
lutea, Bertol. Rar. It. Pl. dec. ii. 32 = Erythraea maritima.
lychnoides, *Berg. Desc. Pl. Cap.* 45.—Afr. austr.
madagascariensis, *Baker, in Journ. Linn. Soc.* xvii. (1881) 278.—Madag.
maritima, *Eckl. in S. Afr. Quart. Journ.* ii. (1830); *ex Griseb. Gen. et Sp. Gent.* 100.—Afr. austr.
maritima, Georgi, Beschr. Russ. Reich. iii. iv. 791 = Erythraea ramosissima.
maritima, Willd. Sp. Pl. i. 1069 = Erythraea maritima.
melampyrifolia, *Lam. Illustr.* ii. 479.—Afr. austr.
minima, Thuill. Fl. Par. ii. 110 = Cicendia pusilla.
minor, Desf. Cat. Hort. Par. ed. III. 126 = Erythraea ramosissima.
nana, Bast. ex Griseb. in DC. Prod. ix. 57 = Erythraea ramosissima.
nudicaulis, *Linn. f. Suppl.* 151.—Afr. austr.
nummularifolia, Willd. ex Griseb. Gen. et Sp. Gent. 144 = Erythraea diffusa.
occidentalis, DC. Fl. Fr. Suppl. 2782 = Erythraea maritima.
orthostyla, Reichb. Exot. t. 245 = frutescens.
ovata, Spreng. ex Griseb. in DC. Prod. ix. 61 = serpyllifolia.
palustris, *Bunch. Trans. Ill.* (1821) 226.—Afr. austr.
panciciana, Michx. Fl. Bor. Am. i. 146 = Sabbatia paniculata.
parviflora, Salisb. Prod. 136 = lucciflora.

CHIRONIA :—
parvifolia, *E. Mey. Comm. Pl. Afr. Austr.* 180.—Afr. austr.
pedunculata, *Lindl. Bot. Reg.* t. 1803.—Afr. austr.
perfoliata, *Eckl. in S. Afr. Quart. Journ.* ii. (1830), *ex Griseb. Gen. et Sp. Gent.* 104.—Afr. austr.
perfoliata, Salisb. Prod. 137 = Chlora perfoliata.
pubescens, *Baker, in Journ. Bot.* xx (1882) 172.—Madag.
pulchella, Willd. Sp. Pl. i. 1067 = Erythraea ramosissima
pusilla, Steud. Nom. ed. II. i. 851 = Microcala filiformis.
ramosissima, Ehrh. ex Griseb. Gen. et Sp. Gent. 187 = Erythraea ramosissima.
scabrida, *Griseb. l. c.* 103.—Afr. austr.
scabriuscula, Willd. ex Griseb. l. c. 143 = Erythraea linarifolia.
serpyllifolia, *Lehm. Ind. Sem. Hort. Hamb.* (1828).—Afr. austr.
speciosa, *E. Mey. Comm. Pl. Afr. Austr.* 178.—Afr austr.
spicata, Willd. Sp. Pl. i. 1069 = Erythraea spicata
stellata, Muhl. Cat. ex A. Gray, Syn. Fl. N. Am. ii 115 = Sabbatia stellaris.
suffruticosa, Salzm ex Griseb. Gen. et Sp. Gent. 141 = Erythraea Centaurium
tabularis, *Page, ex Steud. Nom.* ed. II. i. 852.—Afr. austr.
tenuiflora, *Link, ex Steud. l. c.*—Hab. ?
tetragona, *Linn. f. Suppl.* 151.—Afr. austr
trinervia, Lam. Sp. Pl. 189 = Exacum zeylanicum.
trinervia, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1306 = pedunculata.
uliginosa, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 39 = Erythraea elodes.
uliginosa, Waldst. & Kit Pl. Rar. Hung. iii 287. t. 259 = Erythraea linarifolia.
uniflora, Eckl. ex Griseb. in DC. Prod. ix. 61 = linifolia.
uniflora, *Lam. Encyc.* i. 787.—Afr. austr.
Vaillantii, F. W. Schmidt, Fl Bohem. ii. 84 = Erythraea ramosissima.
virescens, Willd. ex Steud. Nom. ed. II. i. 852 = Erythraea maritima.
viscosa, Zeyher, ex Griseb. Gen. et Sp. Gent. 105 = tetragona.
vulgaris, Cham. in Linnaea, vi. (1831) 348 = linoides, lychnoides, melampyrifolia.

CHIROPETALUM, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 21 = **Argithamnia**, Sw. (Euphorbiac.).
Berterianum, Schlecht. in Linnaea, xxvi. (1853–55) 637 = Berteroana.
canescens, Phil. Fl. Atac. 49 = A. canescens.
griseum, Griseb. in Goett. Abh. xxiv. (1879) 57.—Reg. Argent.
lanceolatum, C. Gay, Fl. Chil. v. 344 = A. tricuspidata.
lanceolatum, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 21 = A. Berteroana.
lineatum, Klotzsch, in Wiegm. Archiv, vii. (1841) 199 = A. lineata.
mollis, Klotzsch, l. c. = A. mollis.
ovatum, Phil. in Linnaea, xxix. (1857–58) 42 = A. tricuspidata.
peruvianum, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 22 = A. quinquecuspidata.
triandrum, Griseb. in Goett. Abh. xxiv. (1879) 56.—Reg. Argent.
tricuspidatum, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 23 = A. tricuspidata.

CHIROSTEMUM, Cerv. in Anal. Clenc. Nat. vi. (1803) 308 = **Gharcostemon**, Humb. & Bonpl. (Stercullac.).

CHISOCHETON, Blume, Bijdr. 168 (1825). *MELIACEÆ*; Benth. & Hook. f. i. 388.
SCHISOCHITON, Spreng. Syst. iv. Cur. Post. 251 (1827).
amabilis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 27.—As. or.
barbatus, *C. DC. in A. DC. Monog. Phan.* i. 586.—As. or.
canalensis, Baill. Adansonia, xi. (1873–76) 260 = Dysoxylum canalense.
cerasinus, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 29. —Ins. Ceram.

CHISOCHETON :—
costatus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 552.—Ind. or.
divergens, *Blume, Bijdr.* 169.—Java.
diversifolius, *Miq. Fl. Ind Bat. Suppl.* i. 504.—Sumatra.
dysoxyloïdius, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 49.—Burma.
erythrocarpus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 550. —Malacca.
fragrans, *Hiern, l. c.* 551.—Malacca.
glomeratus, *Hiern, l. c.*—Ins. Penang.
grandiflorus, *Kurz, in Journ. As. Soc.* xli (1872) ii. 206.—Burma.
holochrysus, *Hiern. in Hook. f. Fl. Brit. Ind.* i. 551.—Malaya.
Junghuhnii, *Miq. in Ann. Mus. Bot Lugd. Bat.* iv. 30 —As. or.
paniculatus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 552.—Ind. or ; Burma
patens, *Blume, Bijdr.* 169.—Java.
paucijugum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 30.—Sumatra.
pendulifloros, *Planch. ex Hiern, in Hook. f. Fl Brit. Ind.* i. 550.—Malaya.
spectabilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 29. —As. or.
spicatus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 550.—Malacca.
tetrapetalus, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 411.—Java.
Vriesianus, *C. DC. in A. DC. Monog. Phan.* i. 532.—As. or.

CHITHONANTHUS, Lehm. in Ind. Sem. Hort. Hamburg (1842) 4; et Pl. Preiss. ii. 368 (1847) = **Aoacia**, Willd. (Legumin.).
restiaceus, Lehm. ii. cc.—A. restiacea.

CHITONIA, D. Don, in Mem. Wern. Soc. iv. (1823) 317 = **Micontia**, Ruiz & Pav. (Melastomac.).
albicans, D. Don, ex Nand. in Ann. Sc. Nat. Sér. III. xv. (1851) 339 = Tetrazygia albicans.
aurea, D. Don, in Mem. Wern. Soc. iv. (1823) 319 = M. aurea.
bubalina, D. Don, l. c. = M. bubalina.
caudata, D. Don, l. c. = M. Donacena.
Fothergilla, D. Don, l. c. 318 = M. Fothergilla.
macrophylla, D. Don, l. c. 319 = M. macrophylla.
pyramidalis, G. Don, in Loud. Hort. Brit. 174 = M. laevigata.
rudgenosa, Steud. Nom. ed. II. i. 852 = M. Fothergilla.
Tamonia, G. Don, in Loud. Hort. Brit. 174 = M Fothergilla.

CHITONIA, Moç. & Sesse, ex DC. Prod. i. 707 (1824). *ZYGOPHYLLEÆ*; Benth. & Hook. f. i. 266.
mexicana, *Moc. & Sesse, l. c.*—Mexic.

CHITONIA, Saliab. Gen. Pl. Fragm. 51 (1866) = **Zygadenus**, Michx. (Liliac.).

CHIZOCHETON, A. Juss. in Orb. Dict. Sc. viii. 80 (1849) = **Chisocheton**, Blume (Meliac.).

CHLAENANDRA, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 83 (1868–69). *MENISPERMACEÆ*.
ovata, *Miq. l. c.* 84.—N. Guinea ?

CHLAENOBOLUS, Cass. in Dict. Sc. Nat. xlix. 337 (1827) = **Pterocaulon**, Ell. (Compos.).
alopecuroides, Cass. l. c. 339 = P. virgatum.
pycnostachyus, Cass. l. c. 338 = P. pycnostachyum.
rugosus, Cass. l. c. 341 = P. virgatum.
spicatus, Cass. l. c. = P. virgatum.
virgatus, Cass. l. c. 340 = P. virgatum.

CHLAMYDANTHUS, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 355, 359 = **Thymelæa**, Endl.
argenteus, C. A. Mey. l. c. 358 = T. Tartonraira.
canescens, C. A. Mey. l. c. = T. canescens.
elliptieus, C. A. Mey. l. c. 359 = T. elliptica.
hirsutus, Griseb. Spicil. Fl. Rumel. ii. 84 = Passerina hirsuta.

CHLAMYDANTHUS:—
nitidus, C. A. Mey. in Bull. Phys-Math. Acad. Pétersb. i. (1843) 358 = Thymelæa nitida.
Tartonraira, C. A. Mey. l. c. = T. Tartonraira.
thesioides, C. A. Mey. l. c. 359 = T. thesioides.
Thymelæa, C. A. Mey. l. c. = T. Sanamunda.
tinctorius, C. A. Mey. l. c. 358 = T. tinctoria.
villosus, C. A. Mey. l. c. = T. villosa.
virgatus, C. A. Mey l. c. 359 = T. virgata.

CHLAMYDIA, Banks, ex Gaertn. Fruct. i. 71. t. 18 (1788) = **Phormium**, Forst. (Liliac.).
tenacissima, Gaertn. l. c. = P. tenax.

CHLAMYDOCARYA, Baill. Adansonia, x. (1872) 276. *OLACINEAE*.
capitata, *Baill. l. c.* 277.—Afr. trop.
Thomsoniana, *Baill. l. c.* 276.—Afr. trop.

CHLAMYDOPHORA, Ehrenb. ex Less. Syn. Comp. 265, 446 (1832) = **Cotula**, Linn. (Compos.).
pubescens, Cass. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 279 = Cot. pubescens.
tridentata, Ehrenb. ex Less. Syn. Comp. 266 = Cot. tridentata.

CHLAMYDOSPERMA, A. Rich Ess. Fl. Cub. 331 (1845) = **Stegnosperma**, Benth. (Phytolacc.).

CHLAMYDOSTYLUS, Baker, in Journ. Bot. xiv. (1876) 185 = **Nemastylis**, Nutt. (Irid.).
bracteolatus, Baker, in Journ. Linn. Soc. xvi. (1877) 107.—Boliv.
cernuus, Baker, in Journ. Bot. xiv. (1876) 186 = N. trifora.
Medusa, Baker, l. c.—Bras.
multiflorus, Baker, l. c. = N. multiflora.
spathaceus, Griseb. in Goett. Abh. xxiv. (1879) 324 = N. spathacea.
tenuis, Baker, in Journ. Bot. xiv. (1876) 185 = N. tenuis.
triforus, Baker, in Journ. Linn. Soc. xvi. (1877) 107 = N. trifora.

CHLAMYSPERMA, Less. Syn. Comp. 256 (1832) = **Villanova**, Lag. (Compos.).
arenarioides, Hook. & Arn. Bot. Beech Voy. 300. t. 64 = V. pratensis.
polygama, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 40.—N. Granat.
pratense, Less. Syn. Comp. 256 = V. pratensis.

CHLAMYSPORUM, Sallisb. Parad. Lond. t. 108 (1808) = **Rhynanotus**, R. Br. (1810) (Liliac.).
dichotomum, Sallisb. l. c. in nota sub t. 103 = T. dichotomus.
hispidulum, Salisb. ex Steud. Nom. ed. II. i. 352 = T. hispidulus.
juncifolium, Sallisb. Parad. Lond. t. 108 = T. junceus.
multiflorum, Sallisb. ex Steud. Nom. ed. II. i. 352 = T. multiflorus.
pauciflorum, Sallisb. ex Steud. l. c. = T. pauciflorus.
triandrum, Sallisb. ex Steud. l. c. = T. triandrus.

CHLANIS, Klotzsch, in Peters, Reise Mossamb. Bot. 144 (1861) = **Oncoba**, Forsk. (Bixin.).
macrophylla, Klotzsch, l. c. 145 = O. macrophylla.
tettensis, Klotzsch, l. c. = O. tettensis.

CHLAOTRACHELUS, Hook. f. Fl. Brit. Ind. iii. 294 (1881), sphalm. = CLAOTRACHELUS, Zoll. = **Wernonia**, Schreb. (Compos.).

CHLETERUS, Rafin. Princ. Somiol. 25 (1814) = **Boes**, Comm. (Gesneriac.).

CHLIDANTHUS, Herb. App. 46 (1821). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 723.
COLEOPHYLLUM, Klotzsch, in Otto & Dietr. Allg. Gartenz. viii. (1840) 185.
Cumingii, Presl, Bot. Bemerk. 115 = Hippeastrum advenum.
Ehrenbergii, *Kunth, Enum. Pl.* v. 654.—Mexic.
fragrans, *Herb. App.* 46.—Peruv.
luteus, D. Dietr. Syn. Pl. ii. 1175 = Stenomesson luteum.

CHLOAMNIA, Rafin. Neogenyt. 4 (1825) = **Festuca**, Tourn. (Gramin.).

CHLOANTHES, R. Br. Prod. 513 (1810). *VERBENACEAE*, Benth. & Hook. f. ii. 1140.
HEMISTEMON, F. Muell. Fragm. x. 78 (1876).
atriplicina, F. Muell. Fragm. i. 285 = Pityrodia atriplicina.
Bartlingii, Lehm. Ind. Sem. Hort. Hamb. (1844), ex Bartl. in Lehm. Pl. Preiss. i. 352 = Pityrodia Bartlingii.
Bonnemaini, *F. Muell. Fragm.* x. 78.—Austral.
bullata, F. Muell. l. c. vi. 156 = Pityrodia uncinata.
coccinea, *Bartl. in Lehm. Pl. Preiss.* i. 352.—Austral.
cuneata, F. Muell. Census, 103 = Pityrodia cuneata.
dilatata, F. Muell. Fragm. vi. 157 = Pityrodia dilatata.
Elderi, F. Muell. l. c. x. 13 = Hemiphora Elderi.
glandulosa, *R. Br. Prod.* 514.—Austral.
halgnaiacea, *F. Muell. Fragm.* x. 14.—Austral.
hemigymnioides, F. Muell. l. c. vi. 156 = Pityrodia hemigymnioides.
lavandulæfolia, Sieber, ex Spreng. Syst. ii. 756 = Stoechadia.
lepidota, *F. Muell. in Wing, South. Sc. Record*, iii. (1885) 3.—Austral.
Lewellinii, *F. Muell. Fragm.* viii. 50.—Austral.
loricata, *F. Muell. l. c.* x. 14.—Australia.
loxocarpa, F. Muell. l. c. ii. 22 = Pityrodia Drummondii.
Oldfieldii, F. Muell, l. c. i. 284 = Pityrodia Oldfieldii.
paniculata, F. Muell. l. c. iv. 80 = Pityrodia paniculata.
parviflora, *Walp. Rep.* iv. 58.—Australia.
rosmarinifolia, *A. Cunn. ex Steud. Nom.* ed. II. i. 352.—Austral.
salvifolia, F. Muell. Census, 108 = Pityrodia salvifolia.
stachyoides, F. Muell. Fragm. v. 50; vi. 158 = Pityrodia racemosa.
Stoechadia, *R. Br. Prod.* 514.—Austral.
uncinata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 194 = Pityrodia uncinata.
verbascifolia, F. Muell. Fragm. i. 233 = Pityrodia verbascina.

CHLOERUM, Willd. ex Spreng. Jahrb. i. iii. 74 (1833) = **Abolboda**, Humb. & Bonpl. (Xyrid.).
gramineum, Willd. ex Kunth, Enum. Pl. iv. 27 = A. Poeppigii.
imberbe, Willd. ex Steud. Nom. ed. II. i. 352 = A. imberbis.

CHLOIDIA, Lindl. Gen. et Sp. Orch. 484 (1840) = **Corymbis**, Thou. (Orchid.).
decumbens, Lindl. l. c.—Bras.
flava, Rolfe, in Kew Bull. Ann. vi. 244 = Cor. flava.
polystachya, Reichb. f. l. c. = *vernalis*.
vernalis, Lindl. Gen. et Sp. Orch. 484.—Jamaica.

CHLONANTHES, Rafin. New Fl. Am. ii. 20 (1836) = **Chelone**, Linn. (Scrophul.).
tomentosa, Rafin. l. c.—Am. bor.

CHLONANTHUS, Rafin. Prine. Somiol. 26 (1814) = prsec.

CHLOOPSIS, Blume, Enum. Pl. Jav. 14 (1827) = **Ophiopogon**, Ker-Gawl. (Haemodorac.).
acaulis, Blume, l. c. = O. japonicus.
caulescens, Blume, l. c. = O. japonicus.

CHLOOTHAMNUS, Buese, in Miq. Pl. Jungh. 386 (1851–55) = **Schizostachyum**, Nees (Gramin.).
chilianthus, Buese, l. c. 387.—Sumatra.

CHLORA, Rem. ex Adans. Fam. ii. 503 (1763). *GENTIANEAE*, Benth. & Hook. f. ii. 809.
BLACKSTONIA, Huds. Fl. Angl. ed. I. 146 (1762), nomen prius.
CHLOROSTIS, Rafin. Princ. Somiol. 25 (1814).
DIRINSTYLIS, Rafin. Fl. Tellur. iii. 31 (1836).
PLEURIMARIA, Rafin. l. c. (1836).
acuminata, Koch & Zis, Cat. Fl. Palat. 20 = serotina.
affinis, Willk. in Bot. Zeit. v. (1847) 874 = imperfoliata.
citrina, *Boiss. & Reut. Pugill. Pl. Nov.* 77.—Hispan.
cremulata, Salisb. ex Griseb. Gen. et Sp. Gent. 118 = imperfoliata.
dodecandra, Linn. Syst. ed. XII. 367 = Sabbatia chloroides.
dubia, Lam. Tabl. Encyc. ii. 484 = imperfoliata.
emiliata, Griseb. Gen. et Sp. Gent. 116 = Eustoma alienifolium.
grandiflora, Vis. Cors. App. Alt. 4 = perfoliata.

CHLORA:—
hybrida, *Tratt. Arch.* t. 229.—Europ.
imperfoliata, *Linn. f. Suppl.* 218.—Europ.
intermedia, Tenore, Syll. 365 = perfoliata.
lanceolata, Koch & Zis, Cat. Pl. Palat. 20 = imperfoliata.
longifolia, *Lam. Tabl. Encyc.* ii. 434.—Hab.?
mauceriensis, Desf. ex Steud. Nom. ed. I. 190; ed. II. i. 352 = perfoliata.
perfoliata, Gorter, ex Steud. Nom. ed. II. i. 352 = Saponaria Vaccaria.
perfoliata, *Linn. Syst. ed.* XII 267.—Europ.: Asia. Min. quadrifolia, *Linn. l. c.*—Europ. austr.
quadrifolia, F. W. Schmidt, in Roem. Arch. i. t. 18 = quadrifolia.
serotina, *Koch, in Reichb. Pl. Crit.* iii. 6. t. 251.—Europ.; As. Min.
sessilifolia, Desv. in Trav. Soc. Amat. i. (1807) 74. t. 8. f. 2 = imperfoliata.
sessilis, Willd. Sp. Pl. ii. 841 (= Plantagines?).—Reg. Argent.
sicula, Tineo, in Guss Fl. Sic. Syn. ii. 819 = perfoliata.
triphylla, *Badaro, in Brugnat. Giorn. Fis.* II. ix. (1826) 166.—Italia.

CHLORADENIA, Baill. Etud. Gén. Euph. 471. t. 19 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 312.
ADENOGYNUM, Zoll. in Linnaea, xxviii. (1856) 825. discolor, *Baill. l. c.* 472.—Java.

CHLORAEA, Lindl. in Brand. Quart. Journ. Roy. Instit. N. S. i. (1827) 47. *ORCHIDEAE*, Benth. & Hook. f. iii. 518.
ASARCA, Lindl. in Quart. Journ. Roy. Instit. i. (1827) 52.
BIENERIA, Reichb. f. in Bot. Zeit. xl. (1853) 3. t. 1.
GAVILLEA, Poepp. & Endl. ex Steud. Nom. ed. II. i. 666 (1840).
ULANTHIA, Hook. Bot. Mag. lvii. sub t. 2990 (1830).
affinis, *Lindl. in Hook. Journ. Bot.* i. (1834) 4.—Chili.
alaris, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1827) 50.—Chili
alpina, *Poepp. Fragm. Syn. Pl. Chil.* 17.—Chili.
amazonica, *Lindl. Gen. et Sp. Orch.* 403.—Chili.
aures, *Phil. in Linnaea*, xxix. (1857–58) 48.—Chili.
Austinæ, A. Gray, in Proc. Am. Acad. xii. (1877) 88 = Cephalanthera Oregana.
barbata, *Lindl. in Hook. Journ. Bot.* i. (1834) 5.—Chili.
biseriata, *Griseb. in Goett. Abh.* xxiv. (1879) 339.—Reg. Argent.
Bergii, *Hieron. in Bol. Acad. Nac. Cordova*, iii. (1879) 380.—Reg. Argent.
bletioides, Lindl. in Brand. Quart. Journ. Roy. Instit. N. S. i. (1827) 50 = alaniloides.
calopogon, *Phil. in Linnaea*, xxix. (1857–58) 47.—Chili.
campestris, *Poepp. Fragm. Syn. Pl. Chil.* 16.—Chili.
chlorostieta, *Phil. in Linnaea*, xxix. (1857–58) 52.—Chili.
chrysantha, *Poepp. Fragm. Syn. Pl. Chil.* 15.—Chili.
chrysochlora, *Phil. in Linnaea*, xxix. (1857–58) 50.—Chili.
collina, *Phil. l. c.* 54.—Chili.
Commersonii, *Brongn. in Duperr. Voy. Bot.* 189.—Reg. Magell.
crispa, *Lindl. Gen. et Sp. Orch.* 401.—Chili.
cristata, *Lindl. in Hook. Journ. Bot.* i. (1834) 4.—Chili.
crocata, *Phil. in Linnaea*, xxix. (1857–58) 51.—Chili.
cuneata, *Lindl. Gen. et Sp. Orch.* 400.—Chili.
cyrgaea, *Phil. in Linnaea*, xxxii. (1864–63) 241.—Chili.
cylindrostachya, *Poepp. Fragm. Syn. Pl. Chil.* 15.—Chili.
dasyrgon, *Phil. in Linnaea*, xxxiii. (1864–65) 242.—Chili.
decipiens, Poepp. Fragm. Syn. Pl. Chil. 17 = cristata, multiflora, Poeppigiana.
dense, *A. Rich. in C. Gay, Fl. Chil.* v. 464.—Chili.
disclon, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1827) 47.—Chili.
fimbriata, *Phil. in Linnaea*, xxix. (1857–58) 49.—Chili.

CHLORAEA:—
fimbriata, Poepp. Fragm. Syn. Pl. Chil. 15 = Bipinnula unguiculata.
Fonkii, *Phil. in Linnaea*, xxix. (1857–58) 55.—Chili.
galeata, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1857) 48.—Chili.
Gandichaudii, *Brongn. in Duperr. Voy. Bot.* 189.—Reg. Magell.
Gavilu, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1857) 48.—Chili.
Gayana, *A. Rich. in C. Gay, Fl. Chil.* v. 439.—Chili.
grandiflora, *Poepp. Fragm. Syn. Pl. Chil.* 14.—Chili.
gymnoglossa, *Phil. in Linnaea*, xxix. (1857–58) 54.—Chili.
heteroglossa, *Reichb. f. l. c.* xxii. (1849) 863.—Chili.
homopetala, *Phil. l. c.* xxix (1857–58) 48.—Chili.
incisa, *Poepp. Fragm. Syn. Pl. Chil.* 17.—Chili.
inconspicua, *Phil. in Anal. Univ. Chil.* (1865) ii. 321.—Chili.
lamellata, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1857) 49.—Chili.
Lechleri, *Lindl. in Lechl. Berb. Am. Austr.* 53.—Chili.
Lindleyi, *Poepp. Fragm Syn. Pl Chil.* 16.—Chili.
littoralis, *Phil. in Linnaea*, xxxii. (1864–65) 214.—Chili.
longebracteata, *Lindl. in Brand Quart. Journ. Roy. Instit.* N. S. i. (1857) 48.—Peruv.
longipetala, *Lindl. Gen. et Sp. Orch.* 400.—Chili.
magellanica, *Hook. f. Fl. Antarct.* ii. 350.—Reg. Magellan.
membranacea, *Lindl. Gen. et Sp. Orch.* 401.—Reg. Argent.
multiflora, *Lindl. in Brand. Quart. Journ Roy. Instit.* i. (1857) 49.—Chili.
nudilabia, *Poepp. Fragm. Syn. Pl. Chil.* 16.—Chili.
obovata, *Phil. in Linnaea*, xxix. (1857–58) 49.—Chili.
odontoglossa, *A. Rich. in C. Gay, Fl. Chil* v. 455.—Chili.
papillosa, Phil. in Linnaea, xxix. (1857–58) 50 = cylindrostachys.
patagonica, *Phil. l. c.* xxxiii. (1864–65) 245.—Chili.
Pavonii, *Lindl. Gen. et Sp. Orch.* 404.—Chili.
Pearcei, *Phil. in Linnaea*, xxxii. (1864–65) 243.—Chili.
penicillata, *Reichb. f. Otia Bot. Hamb.* 51.—Patagon.
Philippi, *Reichb. f. in Linnaea*, xxii. (1849) 863.—Chili.
Piquichen, *Lindl. Gen. et Sp. Orch.* 400.—Chili.
Poeppigiana, *A. Rich. in C. Gay, Fl. Chil.* v. 443.—Chili.
pogonata, *Phil. in Linnaea*, xxxiii. (1864–65) 240.—Chili.
prodigiosa, *Reichb. f. l. c.* xxii. (1849) 863.—Chili.
reflexa, *Phil. l. c.* xxix. (1857–58) 51.—Chili.
rypalogloss, *A. Rich. in C. Gay, Fl. Chil.* v. 457.—Chili.
Sceptrum, *Reichb. f. in Linnaea*, xxii. (1849) 862.—Chili.
secunda, *Phil. l. c.* xxix. (1857–58) 52.—Chili.
semibarbata, *Lindl. Gen. et Sp. Orch.* 403.—Chili.
speciosa, *Poepp. Fragm. Syn. Pl. Chil.* 14.—Chili.
spectabilis, *Phil. in Linnaea*, xxxiii. (1864–65) 240.—Chili.
suaveolens, *Phil. l. c.* xxix. (1857–58) 46.—Chili.
ulanthoides, *Lindl. Gen. et Sp. Orch.* 404.—Chili.
Unguicati, *Reichb. f. in Linnaea*, xxii. (1849) 864.—Chili.
venosa, *Reichb. f. l. c.*—Chili.
verrucosa, *Phil. l. c.* xxix. (1857–58) 58.—Chili.
virescens, *Lindl. in Brand. Quart. Journ. Roy. Instit.* i. (1857) 51.—Chili.
viridiflora, *Poepp. Fragm. Syn. Pl. Chil.* 14.—Chili.
volucris, *Lindl. in Hook. Journ. Bot.* i. (1834) 4.—Chili.

CHLORANTHUS, Sw. in Phil. Trans. lxxvii. (1787) 259. *CHLORANTHACEAE*, Benth. & Hook. f. iii. 124.
CARODUS, Lour. Fl. Cochinch. 88 (1790).
CRYPHARA, Haum. in Brownt. Edinb. Journ. Sc. ii. (1825) 11.
MOSSINA, Thunb. Nov. Gen. 68 (1782).
PEPERIDIA, Reichb. Consp. 219* (1828).
SAINTLEGERIA, Cordem. in Bull. Adansonia, iii. (1862–63) 260.
SARCANDRA, Garda. in Calc. Journ. Nat. Hist. vi. (1846) 348.
STROPHA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4; ed. II. 66 (1892).
TRICRCANDRA, A. Gray, in Perry, Jap. Exped. ii. 318 (1857).
Blumeanus, Cordem. in Bull. Adansonia, iii. (1862–63) 296 = serratus.
brachystachyus, *Blume, Fl. Jav. Chloranth.* 18. t. 2.—Ind. or.; Malaya; China.
ceylanicus, Miq. Fl. Ind. Bat. i. 1. 802 = brachystachyus.
denticulatus, Cordem in Bull. Adansonia, iii. (1862–63) 296 = brachystachys.
elatior, R. Br. in Bot. Mag. sub t. 2190 = officinalis.
erectus, Sweet, Hort. Suburb. Lond. 28 = officinalis.
Fortunei, *Solms, in DC. Prod.* xvi. 1. 476.—China.
grandifolius, *Miq. Fl. Ind. Bat.* i. 1. 802.—Ind. or
hirsutulus, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 129 = brachystachyus.
inconspicuus, Blanco, Fl. Filip. ed. II. 54 = officinalis.
inconspicuus, *Sw. in Phil. Trans.* lxxvii. (1787) 259. t. 10.—China; Japon.
indicus, Wight, Ic. t. 1945 = inconspicuus.
insignis, *Kurz, in Journ. As. Soc. Beng* xlii. (1873) ii. 108.—Burma.
integrifolius, Schult. f. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 130 = Cornus suecica.
Japonicus, *Siebold, in Nov. Act. Nat. Cur* xiv. ii (1829) 681.—China; Japon.
macranthera, *Schult. f. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 130.—Japon
mandarinus, Kunth, Enum. Pl. Amur. t. 2 = japonicus.
monander, R. Br. in Bot. Mag. sub t. 2190 = brachystachyus.
monostachyus, *R. Br. l. c. t.* 2190.—China.
montanus, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 129 = brachystachyus.
obtusifolius, Miq. Fl. Ind. Bat. i. 1. 802 = inconspicuus ? officinalis; *Blume, Enum. Pl. Jav.* 79.—Ind. or.; Malaya.
Oldhami, *Solms, in DC. Prod.* xvi. t. 476 —Japon.
Perryanus, *Solms, l. c.* 474.—China.
quadrifolius, A. Gray, in Miq. Ann. Mus. Bot. Lugd. Bat. iii. 129 = serratus.
salicifolius, Presl, Epim. Bot. 231 = officinalis.
serratus, *Roem. & Schult. Syst.* iii. 461.—Japon.
sumatranus, Miq. Fl. Ind. Bat. Suppl. 848 = officinalis.

CHLORASTER, Haw. in Phil. Mag. lxiii. 103 (1824) = **Naroissus**, Tourn. (Amaryllid.).
flavus, Haw. Monog. Narcise. 7 = N. viridiflorus.
integer, Haw. l. c. = N. viridiflorus.
obliteratus, M. Roem. Syn. Ensat. 214 = N. Broussonetii.
viridiflorus, M. Roem. l. c. 218 = N. viridiflorus.

CHLORIS, Sw. Prod. Veg. Ind. Occ. 25 (1788). *GRAMINEAE*, Benth. & Hook. f. iii. 1165.
CHLORODES, Flach. ex Regel, in Ind. Sem. Hort. Petrop. (1863) 28.
EUSTACHYS, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188.
HETEROLOPHIS, Ehrenb. ex Boiss. Fl. Orient. v. 554 (1881).
MACROSTACHYA, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 408, in nota (1851).
PHACELLARIA, Willd. ex Steud. Nom. ed. II. i. 358, in syn. (1840); ii. 313 (1841).
SCHULTESIA, Spreng. Pugill. ii. 17 (1815).
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 406.—Abyssin.
acicularis, *Lindl. in Mitch. Journ. Trop. Austral.* 31.—Austral.
acuminata, Trin. Sp. Gram. iii. t. 305 = distichophylla.
adsc., J. & C. Presl, Rel. Haenk. i. 289 = elegans.
Alberti, *Regel, in Act. Hort. Petrop.* vii. (1880) 650.—Mongolia.
amethystea, *Hochst. in Flora*, xxxviii. (1855) 205.—Abyssin.
andropogonoides, *Fourn. Mex. Pl. Enum. Gram.* 143; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 555.—Mexic.
anoplis, *Fourn. l. c.*; *ex Hemsl. l. c.*—Mexic.
Arctidia, Salzm. ex Steud. Syn. Pl. Gram. 218 = Gymnopogon foliosus.
arundinacea, Nees, ex Steud. l. c. 207 = polydactyla.

CHLORIS:—
babiensis, *Steud. Syn. Pl. Gram.* 208.—Bras.
barbata, *Sw. Fl. Ind. Occ.* i. 200.—Reg. trop.
Beyrichiana, Steud. Nom. ed. II. i. 353 = seq.
Beyrichana, Kunth, Rev. Gram. i. 89, 289. t. 56 = radiata.
brachystachys, *Anderss. in Peters, Reise Mossamb. Bot.* 556.—Afr. trop.
brevigluma, *Wright, in Sauv. Fl. Cub.* 191.—Ins. Cuba.
breviseta, *Benth. in Hook. Niger Fl.* 566.—Afr. trop.
campuloides, *Trin. ex Steud. Nom.* ed. II. i. 353, nomen. —Hab. ?
caribaea, Spreng. Syst. i. 295 = barbata.
candata, *Trin. ex Bunge, Enum. Pl. Chin. Bor.* 70.—China.
ciliata, *Sw. Prod. Veg. Ind. Occ.* 25.—Am. trop.
compressa, *DC. Cat. Pl. Hort. Monsp* (1813) 94.—Am. austr.
confertifolia, Trin. in Spreng. Neue Entdeck. ii. 74 = distichophylla.
consanguinea, Kunth, Rev. Gram. i. 89 = polydactyla.
crinita, *Lag Varied. Cienc.* ii. iv. (1805) 143.—Ins. Philipp
crinita, Salzm. ex Doell, in Mart Fl. Bras. ii. iii. 64 = radiata
cruciata, *Sw. Prod. Veg. Ind. Occ.* 25.—Ind. occ.
cryptostachya, *Steud. ex J. A. Schmidt, Beitr. Fl. Cap. Prod. Ins.* 148.—Ins. Cap. Verd.
ctenoides, *Steud Syn. Pl. Gram.* 428.—Madag.
cuculinata, *Buch. in Ann. Sc. Nat. Sér.* III. xix. (1853) 357.—Mexic.
curtipendula, Michx. Fl. Bor. Am. i. 59 = Boutelous racemosa.
Cynodon, Trin. Gram. Unifl 229 = Cynodon Dactylon
cynodontoides, *Balansa, in Bull. Soc. Bot. Fr.* xix. (1872) 318.—N. Caled.
decora, Nees, ex Steud. Syn. Pl. Gram. 205 = barbata.
Digitaria, H. B. & K. Nov. Gen. et Sp. i. 168 = Leptochloa Digitaria.
digitata, *Steud Syn Pl. Gram.* 207.—Ind. or.
distachya, Kunth, Rev. Gram. i. 291. t. 57 = monostachya.
distichophylla, *Lag. Gen. et Sp. Nov.* 4.—Am. austr.
divaricata, *R. Br. Prod.* i. 180.—Austral.
dolichostachya, Lag. Gen. et Sp. Nov. 5 = truncata.
dubia, H. B. & K. Nov. Gen. et Sp. i. 169 = Diplachne dubia.
Durandiana, Schult. Mant. ii. 241 = virgata.
elata, *Desv. Opusc* 73.—Bras.
elegans, *H. B. & K. Nov. Gen. et Sp.* i. 166.—Mexic.
eleusinoides, *Griseb. Fl. Brit. W. Ind.* 539.—Ind. occ.
elongata, Poir. Encyc. Suppl. ii. 286 = truncata.
emarginata, *Beauv. Agrost.* 156, nomen.—Hab. ?
equitans, *Trin. Sp. Gram.* iii. t. 309.—Hab. ?
falcata, Sw in Ges. Naturf. Freunde, Neue Schr. iii. (1801) 160 (= *Campulosus falcatus*).—Afr. austr.
fasciculata, Schrad. in Schult. Mant. ii. 239 = distichophylla.
filiformis, Hort. Par. ex Poir. Encyc. Suppl. ii. 287, n. 8 = Boutelous tenuis.
filiformis, *Poir. l. c.* 288, n. 12.—Hab. ?
floccifolus, Poir. l. c. = Eleusine floccifolia.
floridana, *Vasey, Cat. Gram. Un. St.* 61.—Florida.
foliosa, Willd. Sp. Pl. iv. 924 = Gymnopogon foliosus.
Gayana, *Kunth, Rev. Gram.* i. 89, 293. t. 58.—Afr. trop.
geminata, *Hochst. in Flora*, xxxviii. (1855) 205.—Abyssin.
glabrata, *Anderss. in Peters, Reise Mossamb. Bot.* 557.—Afr. trop.
glauca, *Vasey, Cat. Gram. Un. St.* 61.—Florida.
glaucescens, *Steud. Syn. Pl. Gram.* 206.—Peruv.
gracilis, P. Durand, Chlor. 10 = virgata.
gracilis, H. B. & K. Nov. Gen. et Sp. i. 168 = Leptochloa gracilis.
Gryllus, Houck. Syn. Pl. Germ. i. 437 = Chrysopogon Gryllus.
guineensis, *Schum. & Thonn. Beskr. Guin. Pl.* 55.—Afr. trop.
hispida, *P. Durand, Chlor.* 17.—Ins. Philipp.
Humboldtiana, *Steud. Syn. Pl. Gram.* 205.—Ins. Cuba.

CHLORIS:—
humilis, Kunth, Rev. Gram. i. 39; ii. 531. t. 180 = radiata.
imberbis, Desf. Cat. Hort. Par. ed. III. 15 = subimmutica.
incompleta, *Roth, Nov. Pl. Sp.* 60.—Ind. or.
inermis, *Trin. Gram. Unifl.* 232.—Mexic.
inflata, *Link, Enum. Hort. Berol.* i. 105.—Calif.; Ins. Philipp.
intermedia, *A. Rich. Tent. Fl. Abyss.* ii. 407.—Abyssin.
leptostachya, *Hochst. ex A. Rich. l. c.*—Abyssin.
longibarbis, *Michx. ex Beauv. Agrost.* 158, nomen.—Am. bor.
longifolia, *Steud. Syn. Pl. Gram.* 205.—Ins. Philipp.
macrantha, *Desv. Opusc.* 78.—Arabia.
macrantha, Jaub. & Spach, Illustr. iv. 42 (= Tetrapogonis sp.).—Arabia.
macrostachya, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 408 = Enteropogon macrostachya.
madagascariensis, *Steud. Syn. Pl. Gram.* 206.—Madag.
maritima, Trin. Gram. Unifl. 236 = Cynodon Dactylon.
meccana, *Hochst. ex Steud. Syn. Pl. Gram.* 205.—Aegypt.; Arabia; Beluchist.
megastachya, Schrad. ex Schult. Mant. ii. 340 = truncata.
mendocina, *Phil. Sert. Mend. Alt.* 50.—Chili.
monostachya, Michx. Fl. Bor. Am. i. 59 = Cteniun carolinianum.
monostachya, *Pair. Encyc. Suppl.* ii. 288.—Ind. or.; Ins. Mascar.
montana, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 329.—Ind. or.
Moorei, F. Muell. in Linnaea, xxv. (1852) 444 = acicularis.
mucronata, Michx. Fl. Bor. Am. i. 59 = Eleusine aegyptiaca.
multiradiata, *Hochst. in Flora*, xxxviii. (1855) 204.—Abyssin.
myriostachya, *Hochst. l. c.*—Afr. trop.
notocorna, *Hochst. l. c.*—Abyssin.
obtusifolia, *Desv. Opusc.* 78.—Bras.
obtusifolia, *Trin. Gram. Unifl.* 238.—Ins. Philipp.
orthonoton, *Doell, in Mart. Fl. Bras.* ii. iii. 64.—Bras.
pallida, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 244.—Austral.
pallida, Willd. Sp. Pl. iv. 926 = radiata.
paniea, *Willd. l. c.* 923.—Ind. or.
paraguariensis, Stend. Syn. Pl. Gram. 204 = barbata.
paspaloides, *Hochst. in Flora*, xxxviii. (1855) 206.—Abyssin.
patyensis, *Steud. Syn. Pl. Gram.* 207.—Peruv.
pectinata, *Benth. Fl. Austral.* vii. 612.—Austral.
pendula, Salzm. ex Steud. Syn. Pl. Gram. 208 = bahiensis.
penicillata, Jan, ex Trin. Gram. Unifl. 231 = compressa.
penicillata, *Pers. Syn.* i. 87.—Ind. or.
penicillata, Willd. ex Stend. Nom. ed. II. i. 358 = elegans.
peregrina, *P. Durand, Chlor.* 11.—Ins. Philipp.
petraea, *Sw. Prod. Veg. Ind. Occ.* 25.—Reg. trop.
piloss, *Schum. & Thonn. Beskr. Guin. Pl.* 55.—Afr. trop.
pipertia, Michx. ex Steud. Nom. ed. II. i. 358 = Cteniun caroliniauum.
poaeformis, H. B. & K. Nov. Gen. et Sp. i. 169 = Leptochloa virgata.
polydactyla, P. Durand, Chlor. 14, 22 = virgata.
polydactyla, *Sw. Prod. Veg. Ind. Occ.* 26.—Am. austr.
polystachya, Lag. Gen. et Sp. Nov. 4 = inermis, Leptochloa virgata.
polystachya, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 330.—Ind. or.
Prieurii, *Kunth, Rev. Gram.* i. 39; ii. 541. t. 184.—Afr. trop.
procumbens, P. Durand, Chlor. 16 = Boutelona tenuis.
propinqua, *Steud. Syn. Pl. Gram.* 204.—Guadaloupe.
prostrata, Poir. Encyc. Suppl. ii. 289 = Eleusine aegyptiaca.
pubescens, *Lag. Varied. Cienc.* ii. iv. (1805) 148.—Peruv.
pubescens, *Peyr. in Sitzb. Acad. Wiss, Math. Nat.* xxxviii. (1860) 586.—Am. austr.

CHLORIS:—
pulchra, *Schum. & Thonn. Beskr. Guin. Pl.* 56.—Afr. trop.
Pumilio, *R. Br. Prod.* 186.—Australi.
punctulata, *Hochst. ex Steud. Syn. Pl. Gram.* 205.—Afr. trop.
pycnothrix, Trin. Gram. Unifl. 234 = radiata.
radiata, Heyne, ex Roth, Nov. Pl. Sp. 61 = incompleta.
radiata, *Sw. Prod. Veg. Ind. Occ.* 26.—Am. trop.
regens, Anct. ex Steud. Nom. ed. II. i. 358 = Eleusine indica.
repens, *Hochst. in Flora*, xxxviii. (1855) 204.—Abyssin.
retusa, *Lag. Gen. et Sp. Nov.* 5.—Reg. Argent.
rhachitricha, *Steud. Syn. Pl. Gram.* 205.—Ins. Philipp.
Roxburghii, Edgew. in Journ. As. Soc. Beng. xxi. (1852) 188 = digitata.
Roxburghiana, Schult. Mant. ii. 239 = polystachya.
rufescens, *Lag. Varied. Cienc.* ii. iv. (1805) 148.—Ins. Philipp.
rufescens, *Steud. Syn. Pl. Gram.* 206.—Ins. Mauric.
Sagraeana, *A. Rich. Fl. Cub. Fanerog.* ii. 315.—Ins. Cuba.
Salamanni, Steud. Syn. Pl. Gram. 206 = radiata.
scariosa, *Beauv. Agrost.* 158, nomen.—Hab.?
sciurea, *F. Muell. Fragm.* vi. 86.—Austral.
sclerantha, Lindl. in Mitch. Journ. Trop. Austral. 81 = ventricosa.
scoparia, Desf. Tabl. ed. J. 14 = radiata.
simplex, *Schum. & Thonn. Beskr. Guin. Pl.* 54.—Afr. trop.
spathacea, *Hochst. ex Steud. Syn. Pl. Gram.* 204.—Abyssin.
subbiflora, *Steud. l. c.* 206.—Am. bor.
subdolichostachya, *C. Muell. in Bot. Zeit.* xix. (1861) 341.—Am. bor.
submutica, *H. B. & K. Nov. Gen. et Sp.* i. 167. t. 50.—Mexic.
subtriflora, *Steud. Syn. Pl. Gram.* 208.—Afr. trop.
Swartziana, *Doell, in Mart. Fl. Bras.* ii. iii. 68.—Bras.
Swartzii, C. Muell. in Bot. Zeit. xix. (1861) 341 = pectinata.
tacmensis, *Steud. in Lechl. Bord. Am. Austr.* 56.—Peruv.
tenella, *Koen. ex Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 329.—Ind. or.
tensior, Poir. ex Schult. Mant. ii. 321 = Leptochloa cynosuroides.
tenuis, Poir. Encyc. Suppl. v. 614 = Boutelona tenuis.
tetrameris, Trin. Gram. Unifl. 235 = Cynodon elongatus.
Tetrapogon, Beauv. Agrost. 158 = Tetrapogon villosus.
triangulata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 406.—Abyssin.
truncata, *R. Br. Prod.* 186.—Malaya; Austral.
unispicea, *F. Muell. Fragm.* vii. 118.—Austral.
ventricosa, *R. Br. Prod.* 186.—Australi.
villosa, Pers. Syn. i. 87 = Tetrapogon villosus.
virgata, P. Durand, Chlor. 10 = radiata.
virgata, *Sw. Fl. Ind. Occ.* i. 208.—Am. trop.
Wightiana, Nees, ex Steud. Syn. Pl. Gram. 206 = incompleta.

CHLORIZA, Salisb. Gen. Pl. Fragm. 21 (1866) = **Lachenalia**, Jacq. (Liliac.).
fragrans, Salisb. l. c. = L. versicolor.
lucida, Salisb. l. c. = L. lucida.
media, Salisb. l. c. = L. media.
pallida, Salisb. l. c. = L. pallida.
pustulata, Salisb. l. c. = L. pustulata.
unicolor, Salisb. l. c. = L. versicolor.

CHLOROCAULON, Klotzsch, in Endl. Gen. Suppl. iv. iii. 89 (1850) = **Argithamnia**, Sw. (Euphorbiac.).

CHLOROCHLAMYS, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 149 (1868–69) = **Maesenia**, R. Br. (Asclepiad.).
celebica, Miq. l. c.—Ins. Celebes.

CHLOROCODON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 118 = **Briza**, Tourn.
scoparium, Fourr. l. c. = E. scoparia.

CHLOROCODON, Hook. f. Bot. Mag. t. 5998 (1871). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 745.
Whitsii, *Hook. f. l. c.*—Afr. austr.

CHLOROCREPIS, Griseb. in Goett. Abh. v. (1853) 155 = **Hieracium**, Linn. (Compos.).
staticifolia, Griseb. l. c. = H. staticefolium.

CHLOROGALUM, Kunth, Enum. Pl. iv. 681 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 785.
Laothoe, Raf. Fl. Tellur. iii. 53 (1836), nomen prius.
angustifolium, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 104. t. 90.—Calif.
dinaricetum, Kunth, Enum. Pl. iv. 682 = pomeridianum.
Leichtlinii, *Baker, in Gard. Chron.* (1874) i. 689.—Am. bor. occ.
parviflorum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 249.—Calif.
pomeridianum, *Kunth, Enum. Pl.* iv. 682.—Calif.

CHLOROIDES, Fisch. ex Regel, Ind. Sem. Hort. Petrop. (1863) 28 = **Chloris**, Sw. (Gramin.).
petraeus, Fisch. l. c. = Chloris petraea.

CHLOROMYRON, Pers. Syn. ii. 78 (1807) = **Rhoedia**, Linn. (Guttifer.).
verticillatum, Pers. l. c. = R. acuminata.

CHLOROPHORA, Gaudich. in Freyc. Voy. Bot. 508, in nota (1826). *URTICACEAE*, Benth. & Hook. f. iii. 363.
tinctoria, *Gaudich. l. c.*—Mexic.

CHLOROPHYTUM, [Ker-Gawl.] in Bot. Mag. t. 1071 (1808). *LILIACEAE*, Benth. & Hook. f. iii. 788.
Asphodelopsis, Stend. ex Baker, in Journ. Linn. Soc. xv. (1876) 321.
Hartwegia, Nees, in Nov. Act. Nat. Cur. xv. ii. (1831) 879.
Hollia, Heynh. Nom. ii. 303 (1846).
Schidospermum, Griseb. in Lechl. Berb. Am. Austr. 56 (1857).
abyssinicum, Kotschy & Peyr. Pl. Tinn. 46 = laxiflorum.
acaule, *Baker, in Trans. Linn. Soc.* xv. (1876) 327.—Ind. or.
affine, *Baker, in Trans. Linn. Soc.* xxix. (1875) 160. t. 104.—Afr. trop.
Afzelii, *Baker, in Journ. Linn. Soc.* xv. (1876) 323.—Afr. trop.
allamaefolium, *Baker, l. c.* 324.—Afr. trop.
alpinum, *Benth. ex Baker, l. c.* 328.—Ins. Tasman.
amplexicaule, *Baker, l. c.* 325.—Afr. trop.
andongense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 260.—Afr. trop.
anthericoideum, Dalz. in Hook. Kew Journ. ii. (1850) 141 = tuberosum.
arundinaceum, *Baker, in Journ. Linn. Soc.* xv. (1876) 323.—Ind. or.
attenuatum, *Baker, l. c.* 332.—Ind. or.
blepharophyllum, *Schweinf. ex Baker, l. c.* 327.—Abyssin.
Bowkeri, *Baker, in Saund. Refug. Bot.* t. 352.—Afr. austr.
bracteosum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 260 = stenopetalum.
Brehmeanum, *Schulz. f. Syst.* vii. 454.—Afr. austr.
breviscapum, *Dalz. in Hook. Kew Journ.* ii. (1850) 141.—Ind. or.
Burchellii, *Baker, in Journ. Linn. Soc.* xv. (1876) 330.—Afr. austr.
ciliatum, *Baker, in Journ. Bot.* xvi. (1878) 225.—Afr. trop.
comosum, Baker, in Gard. Chron. (1873) 75 = Sternbergianum.
crispum, *Baker, in Journ. Linn. Soc.* xv. (1876) 321.—Afr. austr.
debile, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1876) 260.—Afr. trop.
decipiens, *Baker, in Journ. Linn. Soc.* xx. (1883) 275.—Madag.
dubium, Schult. f. Syst. vii. 455 = Antherium Subulatum.
elatum, *R. Br. Prod.* 277.—Afr. austr.
falcatum, Baker, in Saund. Refug. Bot. t. 352 = laxiflorum.
filipendulum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1876) 260.—Afr. trop.
Finlaysonianum, Wall. Cat. n. 5021 (Drummondiae sp.).—Siam.

CHLOROPHYTUM:—
galpinianum, Schweinf. ex Baker, in Journ. Linn. Soc. xv. (1876) 325 = Orchidastrum.
glaucum, *Dals. in Hook. Kew Journ.* ii. (1850) 148.—Ind. or.
graminifolium, Kunth, Enum. Pl. iv. 606 = Anthericum undulatum.
Hartmannianum, *Schweinf. Beitr. Fl. Aethiop.* 294, nomen.—Afr. trop.
Heynianum, *Wall. Cat.* n. 6060.—Ind. or.
Heynii, Baker, in Journ. Linn. Soc. xv. (1876) 322 = prec.
inornatum, *Ker-Gawl. in Bot. Mag.* t. 1071.—Afr. trop.
juncifolium, *Baker, in Journ. Linn. Soc.* xv. (1876) 333.—Bras.
Kirkii, *Baker, in Gard. Chron.* (1882) i. 108.—Afr. trop.
lanciflorum, *Wight. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 360.—Afr. trop.
lanciforme, Baker, in Journ. Linn. Soc. xv. (1876) 328 = laxum.
laxum, *R. Br. Prod.* 277.—Trop. As. ; Afr. ; Austral.
leucolepis, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 359 = macrophyllum.
longifolium, *Schweinf. Beitr. Fl. Aethiop.* 294, nomen.—Abyssin.
longipes, *Baker, in Journ. Bot.* xvi. (1878) 325.—Afr. trop.
macrophyllum, *Aschers. in Schweinf. Beitr. Fl. Aethiop.* 294.—Abyssin.
macrosporum, *Baker, in Journ. Linn. Soc.* xv. (1876) 330.—Afr. austr.
madagascariense, *Baker, in Journ. Bot.* xvi. (1878) 326.—Afr. trop.
malabaricum, *Baker, in Journ. Linn. Soc.* xv. (1876) 331.—Ind. or.
micranthum, *Baker, in Journ. Bot.* xvi. (1878) 325.—Afr. trop.
modestum, *Baker, in Journ. Linn. Soc.* xv. (1876) 329.—Afr. austr.
nepalense, *Baker, l. c.* 330.—Reg. Himal.
Nimmonii, Dals. in Hook. Kew Journ. ii. (1850) 142 = Orchidastrum.
Orchidastrum, *Lindl. Bot. Reg.* t. 813.—Ind. or.; Afr. trop.
parviflorum, Dals. in Hook. Kew Journ. ii. (1850) 141 = laxum.
petiolatum, *Baker, in Journ. Linn. Soc.* xv. (1876) 326.—Afr. trop.
polyrhizum, *Baker, in Gard. Chron.* (1878) ii. 396.—Afr. trop.
polystachys, *Baker, in Journ. Bot.* xvi. (1878) 326.—Afr. trop.
pubiflorum, *Baker, in Journ. Linn. Soc.* xv. (1876) 329.—Afr. trop.
pulchellum, Kunth, Enum. Pl. iv. 605 = Anthericum pulchellum.
pusillum, *Schweinf. ex Baker, in Journ. Bot.* xvi. (1878) 325.—Afr. trop.
rhizomatosum, *Baker, in Gard. Chron.* (1885) ii. 230.—Afr. trop.
rigidum, Kunth, Enum. Pl. iv. 604 = Anthericum rigidum.
Rutenbergianum, *Vatke, in Bremen Abh.* ix. (1885) 136.—Ins. Nossibé.
schidospermum, *Baker, in Journ. Linn. Soc.* xv. (1876) 326.—Peruv.
sparsiflorum, *Baker, l. c.* 325.—Afr. trop.
stenopetalum, *Baker, l. c.* 331.—Afr. trop.
Sternbergianum, *Steud. Nom.* ed. II. i. 354.—Afr. austr.
subfruticosum, *Baker, in Journ. Bot.* xvi. (1878) 326.—Afric.
tetraphyllum, *Baker, in Journ. Linn. Soc.* xv. (1876) 328.—Abyssin.
Tisseau, *Baker, l. c.* 328.—Afr. trop.
triflorum, Kunth, Enum. Pl. iv. 606 = Anthericum triflorum.
tuberosum, *Baker, in Journ. Linn. Soc.* xv. (1876) 332.—Ind. or. ; Abyss.
undulatum, Wall. Cat. n. 5059 = nepalense.
ventricosum, *Baker, in Journ. Linn. Soc.* xv. (1876) 326.—Afr. trop.
viscosum, Kunth, Enum. Pl. iv. 605 = Anthericum viscosum.
xerophilum, F. Muell. Fragm. i. 63 = laxum.

CHLORYPHYTUM, Pohl, ex DC. Prod. iv. 340 (1830) = Spermacoce, Dill. (Rubiac.).
racemosum, Pohl, l. c. 548 = S. Poaya.
pratense, Pohl, l. c. = S. Poaya.

CHLOROPYRON, Behr, in Proc. Calif. Acad. i. (1855) 61 = Cordylanthus, Nutt. (Scrophul.).
palustre, Behr, l. c. = Cordylanthus maritimus.

CHLOROSA, Blume, Bijdr. 420. t. 81 (1825). *ORCHIDEÆ*, Benth. & Hook. f. iii. 616.
gracilis, *Blume, l. c.*—Java.
latifolia, *Blume, l. c.*—Java.

CHLOROSPATHA, Engl. in Regel, Gartenfl. (1878) 97. t. 983. *AROIDEÆ*, Benth. & Hook. f. iii. 977.
Kolbii, *Engl. l. c.*—N. Granat.

CHLOROSTEMMA, Fourr. in Ann. Soc. Linn. Lyon, N.S. xvi. (1868) 298 = **Asperula**, Linn. (Rubiac.).
odoratum, Fourr. l. c. = A. odorata.

CHLOROSTIS, Rafin. Princ. Somiol. 26 (1814) = **Chlora**, Ren. (Gentian.).

CHLOROXYLON, Rumph. ex Scop. Introd. 208 (1777) ; DC. Prod. i. 625 (1824). *MELIACEÆ*, Benth. & Hook. f. i. 340.
Dupada, Buch.-Ham. Journ. Mysore, i. 184 = Boswellia serrata.
Swietenia, *DC. Prod.* l. 625.—Ind. or.

CHLOROXYLUM, P. Br. Hist. Jamaic. 187. t. 7. f. 1 (1756) ; Adans. Fam. ii. 308 (1763) (Quid ?).

CHLORYLLIS, E. Mey. Comm. Pl. Afr. Austr. 149 (1835) = **Dolichos**, Linn. (Legumin.).
pratensis, E. Mey. l. c. = D. Chloryllis.

CHNOANTHUS, Phil. in Anal. Univ. Chili (1863) ii. 408 = **Gomphrena**, Linn. (Amarantac.).
meridionalis, Phil. l. c. ; et in Linnæa, xxxiii. (1864–65) 223.—Chili.
ovatus, Phil. Sert. Mend. Alt. 41.—Chili.

CHOCHO, Adans. Fam. ii. 500 (1763) = **Sechium**, P. Br. (Cucurbit.).

CHODONDENDRON, Bosc, Encycl. Agric. vii. 296 (1892), sphalm. = Chondodendron, Ruiz & Pav. (Menispernac.).

CHOENOMELES, Lindl. in Trans. Linn. Soc. xiii. (1822) 97 = **Pyrus**, Linn. (Rosac.).
alba, Carr. in Rev. Hortic. (1874) 410 = P. japonica.
cancellata, Prits. Ic. Ind. i. 250, sphalm. = P. sinensis, Poir.
japonica, Lindl. in Trans. Linn. Soc. xiii. (1822) 97 = P. japonica.

CHOERADODIA, Herb. Amaryll. 87 (1837) = **Strumaria**, Jacq. (Amaryllid.).
chilensis, Herb. l. c. = S. chilensis.

CHOERADOPLECTRON, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 436 = **Habenaria**, Willd. (Orchid.).
Spiranthes, Schau. l. c. = H. lacertifera.

CHOERophyllum, Neck. Delic. Gallo-Belg. i. 147 (1768) = **Chærophyllum**, Hoffm. (Umbell.).

CHOEROSERIS, Link, Handb. i. 798 (1829) = **Picris**, Linn. (Compos.).
heterosperma, Link, l. c. 799 = Crepis Dioscoridis.
phagelioides, Link, l. c. = P. Sprengeriana.
Sprengeriana, Link, l. c. = P. Sprengeriana.

CHOETOPHORA, Fraesch. & Sav. Enum. Pl. Jap. ii. 548 (1879), sphalm. = CHAETOSPORA, R. Br. = **Schœnus**, Linn. (Cyperac.).

CHOISYA, H. B. & K. Nov. Gen. et Sp. vi. 4. t. 518 (1823). *RUTACEÆ*, Benth. & Hook. f. i. 297.
JULIANIA, La Llave, in La Llave & Lex. Nov. Veg. Descr. fasc. ii. 4 (1825).
PLEUSCIA, Mog. & Sesse, ex DC. Prod. i. 724 (1824).
grandiflora, Kegel, Gartenfl. (1876) 357 ; in Act. Hort. Petrop. iv. (1876) 384 = ternata.
ternata, *H. B. & K. Nov. Gen. et Sp.* vi. 6. t. 518.—Mexic.

CHOMELIA, Jacq. Enum. Pl. Carib. 1 (1760). *RUBIACEÆ*, Benth. & Hook. f. ii. 108.
ANISOMERIS, Presl, Symb. Bot. ii. 5. t. 54 (1833).
angustifolia, *Benth. in Hook. Journ. Bot.* iii. (1841) 235.—Guiana.
anisomeris, *Muell. Arg. in Flora*, lviii. (1875) 451.—Bras.
barbinervis, *Moric. ex Benth. in Linnæa*, xxiii. (1850) 446.—Bras.
brasiliana, *A. Rich. l. c.* partim = Vauthieri.
centrillana, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
fasciculata, *Sw. Fl. Ind. Occ.* i. 238.—Jamaica.
filipes, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1855) 41.—Nicaragua.
hirsuta, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 107.—Bras.
intercedens, *Muell. Arg. in Flora*, lviii. (1875) 451.—Bras.
malanoclides, *Muell. Arg. l. c.* 452.—Bras.
Martiana, *Muell. Arg. l. c.* 451.—Bras.
obtusa, *Cham. & Schlecht. in Linnæa*, iv. (1829) 186. —Bras.
obtusa, Mart. in Flora, xxiv. (1841) ii. Beibl. 71 = Martiana.
occidentalis, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
odoratissima, *Tejsm. & Binn. Cat. Hort. Bog.* 884, nomen.—Hab. ?
oligantha, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
parviflora, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 41.—Bras.
pedunculosa, *Benth. in Linnæa*, xxiii. (1850) 445 — Bras.
Pohliana, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
pubescens, *Cham. & Schlecht. in Linnæa*, iv. (1829) 187.—Bras.
ribesioides, *Benth. ex A. Gray, in Proc. Am. Acad.* iv. (1860) 35.—Bras.
sandwicensis, *A. Gray, l. c.*—Ins. Sandvic.
sericea, *Muell. Arg. in Flora*, lviii. (1875) 451.—Bras.
sessilis, *Muell. Arg. l. c.*—Bras.
spinosa, *Jacq. Enum. Pl. Carib.* 13.—Venezuela.
tenuiflora, *Benth. in Hook. Journ. Bot.* iii. (1841) 235. —Guiana.
transiens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 457.—Bras.
triscantha, *Griseb. Cat. Pl. Cub.* 183.—Cuba.
tristis, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
Vauthieri, *Muell. Arg. l. c.*—Bras.
versicolor, Spreng. Syst. i. 410 = Scoloxanthus versicolor.
vulpina, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.

CHOMELIA, Linn. Gen. ed. I. 55 (1737) ; cf. Richter, Codex Linn. 188 = Tarenna, Gaertn. (Rubiac.).

CHOMELIA, Vell. Fl. Flum. i. t. 107 (1827) = **Ilex**, Tourn.
amara, Vell. l. c.—Bras.

CHONA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = **Brica**, Linn.
sanguinea, D. Don, l. c. = E. Nivenia.

CHONAIS, Salisb. Gen. Pl. Fragm. 128 (1866) = **Hippeastrum**, Herb. (Amaryllid.).
vittata, Salisb. l. c. = H. vittatum.

CHONDODENDRON, Benth. & Hook. f. Gen. i. 84 (1863), sphalm. = Odontocarya, Miers (Menispern.).

CHONDODENDRON, Ruiz & Pav. Prod. 128 (1794). *MENISPERMACEÆ*, Benth. & Hook. f. i. 962.
BOTRYOPHIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 43.
acuminum, Miers, l. c. Ser. III. xix. (1867) 192 = platyphyllum.
cinerascens, Miers, l. c. = platyphyllum.
connaroides, Poepp. & Endl. Nov. Gen. et Sp. ii. 65 = tomentosum.

CHONDODENDRON :—
cretosum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xix. (1867) 192.—Peruv.
haderifolium, Miers, l. c. Ser. II. vii. (1851) 44 = tomentosum.
nemophilum, *Miers, Contrib.* iii. 819.—Bras.
obscurum, *Miers, l. c.* 814.—Bras.
ovatum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xix. (1867) 192.—Bras.
platyphyllum, *Miers, l. c.*—Bras.
scabrum, Miers, l. c. Ser. II. vii. (1851) 44 = tomentosum.
tamoides, *Miers, l. c.*—Guiana.
tomentosum, *Ruiz & Pav. Syst. Veg.* 261.—Peruv.

CHONDRACHNE, R. Br. Prod. 220 (1810) = **Lepironia**, Rich. (Cyperac.).
articulata, R. Br. l. c. = *L. mucronata*.

CHONDRACHYRUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836) = **Melica**, Linn. (Gramin.).
scabrum, Nees, l. c. 450.—Bras.

CHONDRILLA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 241 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 594.
CALYCOCORSUS, F. W. Schmidt, Samml. Phys. Aufs. i. 271 (1795).
PELTIDIUM, Zollikof. Nat. Anz. (1820); Bartl. Ord. 145 (1830).
WIBELIA, Roehl. Deutschl. Flora, ii. 426 (1813).
WILLEMETIA, Neck. Elem. i. 60 (1790).
ZOLLIKOFERIA, Nees, in Bluff & Fingerh. Comp. Fl. Germ. ii. 305 (1825).
acanthifolia, Poir. Encyc. Suppl. ii. 330 = Lactuca acanthifolia.
acantholepis, Boiss. Diagn. Ser. I. xi. 48 = juncea.
acanthophylla, Borckh. ex Reichb. Fl. Germ. Excurs. 271 = juncea.
alba, Lam. Encyc. ii. 79 = Prenanthes alba.
altissima, Lam. l. c. 78 = Prenanthes altissima.
ambigua, Fisch. ex Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 398 = pauciflora.
angustissima, Hegetschw. Fl. Schw. 762 = juncea.
aspera, *Poir. Encyc. Suppl.* ii. 329.—Hab.?
asplenifolia, Poir. l. c. 330. = Lactuca saplenifolia.
auriculata, Wall. Cat. n. 3269 = Lactuca dissecta.
brevirostris, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 36.—Sibir. altaic.
caucasica, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 397.—Soongar.
capitata, Sieber, ex DC. Prod. vii. 180 = Lactuca glomerata.
chinensis, Poir. Encyc. Suppl: ii. 331 = Lactuca versicolor.
crepidinea, Poir. l. c. 331 = Prenanthes crepidinea.
crepoides, Lapeyr. Hist. Abr. Pl. Pyr. 462 = Lactuca saligna.
crepoides, *Murr. Syst. Veg.* ed. XIII. 595.—Moet. Pyrm.
crepoides, Reich. Fl. Moeno-Francof. 59 = juncea.
cymosa, Poir. Encyc. Suppl. ii. 330 = Prenanthes cymosa.
debilis, Poir. l. c. 332 = Lactuca debilis.
Decandollei, Spreng. ex DC. Prod. vi. 158 = Crepis foetida.
dentata, Poir. Encyc. Suppl. ii. 328 = Lactuca denticulata.
eryngiifolia, Poir. l. c. 331 = Lactuca muralis.
fontinalis, Wall. Cat. n. 3265 = Lactuca polyorphala.
Gaudini, Hegetschw. in Suter, Fl. Helv. ed. II. ii. 165 = juncea.
glomerata, C. Koch, in Linnaea, xxiii. (1850) 678 = juncea.
gracilis, Wall. Cat. n. 3267 = Lactuca gracilis.
graminea, *Bieb. Fl. Taur. Cauc.* ii. 246.—Reg. Caucas. et Himal.; Sibiria.
graminea, Ledeb. Fl. Alt. iv. 148 = piptocoma.
graminifolia, Fisch. Cat. Hort. Gorenk. (1812) 33 = graminea.
hastata, Poir. Encyc. Suppl. ii. 332 = Lactuca denticulata.
hastata, Wall. Cat. n. 3271 = Lactuca hastata.
hieracioides, Roth, in Usteri, Ann. Bot. x. 50 = Tolpis coronopifolia.
hispida, Desf. Cat. Hort. Par. ed. III. 144 = juncea.
hispida, Poir. Encyc. Suppl. ii. 328 = Lactuca microcephala.
illinoiensis, Poir. l. c. 331 = Prenanthes aspera.

CHONDRILLA:—
indica, Steud. Nom. ed. I. 191 = Lactuca Heyneana.
intybacea, Frivald. in Flora, xviii. (1835) 585 = juncea.
intybacea, Ledeb. Fl. Ross. ii. 809 = pauciflora.
japonica, Lam. Encyc. ii. 79 = Crepis japonica.
juncea, Ledeb. Fl. Alt. iv. 147 = brevirostris.
juncea, *Linn. Sp. Pl.* 796.—Europ.; As. med.
laciniata, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 410 = juncea.
laevigata, Pursh, Fl. Am. Sept. ii. 497 = Pyrrhopappus carolinianus.
lanceolata, Poir. Encyc. Suppl. ii. 329 = Crepis integra.
latifolia, Bieb. Fl. Taur. Cauc. ii. 244 = juncea.
lejourneana, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 456.—Sibir. altaic.
longifolia, Buch.-Ham. in Wall. Cat. sub n. 3274 = Lactuca polyorphala.
longifolia, Wall. Cat. n. 3273 = Lactuca longifolia.
lutea, Dulac, Fl. Hautes-Pyr. 495 = juncea.
lyrata, Poir. Encyc. Suppl. ii. 332 = Crepis japonica.
maracandica, *Bunge, in Mém. Sav. Etr. Petersb.* vii. (1851) 380.—Turkestan.
multiflora, Poir. Encyc. Suppl. ii. 332 = Crepis japonica.
muralis, Lam. Fl. Fr. ii. 105 = Lactuca muralis.
nudicaulis, Linn. Mant. 278 = Launaea nudicaulis.
paniculata, Lam. Encyc. ii. 78 = prenanthoides.
pauciflora, D. Don, in Trans. Linn. Soc. xvi. (1833) 180 = Pyrrhopappus pauciflorus.
pauciflora, *Ledeb. Fl. Alt.* iv. 148.—Rossia; Sibiria.
Pallasiana, *Mortin. Fl. Sarep.* 263.—Europ.
phaeocephala, *Rupr. Sert. Tianach.* 59.—As. centr.
piestocarpa, *Boiss. Fl. Orient.* iii. 793.—Persia.
pinnatifida, Poir. Encyc. Suppl. ii. 330 = Prenanthes cymosa.
piptocoma, *Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop.* viii. 54.—Sibir. altaic.
Prenanthes, Steud. Nom. ed. I. 191 = seq.
prenanthoides, *Vill. Voy.* (1812) 16.—Europ.
pulchra, Lam. Fl. Fr. ii. 106 = Phaecasium lampsanoides.
purpurea, Lam. l. c. 105 = Prenanthes purpurea.
puella, *Boss. Diagn.* Ser. I. vii. 11.—Persia; Turkestan.
racemosa, Poir. Encyc. Suppl. ii. 331 = Lactuca Heyneana.
ramosissima, *Sibth. & Sm. Fl. Graec. Prod.* ii. 128.—Graecia.
ramosissima Zeyher, ex DC. Prod. vii. 176 = Lactuca viminea.
Rauillieri, Walp. Rep. ii. 694, sphalm. = Roullilieri.
repens, Lam. Encyc. ii. 79 = Lactuca repens.
rhombifolia, Humb. ex Willd. Sp. Pl. iii. 1535–36 = Prenanthes rhombifolia.
rigens, Reichb. Fl. Germ. Excurs. 271 = juncea.
Roullieri, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 456.—Sibir. altaic.
rubicunda, Poir. Encyc. Suppl. ii. 329 = Prenanthes alba.
ruderalis, Gaertn. ex Steud. Nom. ed. I. 191, 655 = Lactuca muralis.
runcinata, Wall. Cat. n. 3272 = Crepis japonica.
sagittata, Wall. l. c. n. 3270 = Lactuca sagittarioides.
saxatilis, Poir. Encyc. Suppl. ii. 330 = Launaea pinnatifida.
Schrederi, Steud. Nom. ed. II. ii. 393, in syn. = saxatilis.
secunda, Royle, ex C. B. Clarke, Comp. Ind. 276 = Launaea secunda.
Senaarensa, D. Don, in Trans. Linn. Soc. xvi. (1833) 180 = Pyrrhopappus Senaarensis.
sessiliflora, Lam. Fl. Fr. ii. 104 = Lactuca viminea.
setulosa, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 402.—Himal. bor. occ.
sonchifolia, Poir. Encyc. Suppl. ii. 330 = Lactuca Heyneana.
soongarica, *Stschegl. in Bull. Soc. Nat. Mosc.* xxvii. (1854) i. 179.—Soongar.
squarrosa, Poir. Encyc. Suppl. ii. 332 = Lactuca brevirostris.
stipitata, *Sch. Bip. in Linnaea*, xv. (1841) 558.—Europ.
stricta, *Ledeb. Fl. Alt.* iv. 146.—Sibir. altaic.
tenuifolia, Lam. Encyc. ii. 78 = Prenanthes purpurea.
tenuis, Buch.-Ham. in Wall. Cat. n. 3274 = Lactuca polyorphala.

CHONDRILLA:—
triquetra, Steud. Nom. ed. I. 191; ed. II. i. 354 = Lactuca triquetra.
verricolor, Sch. Bip. in Mus. Senckenb. iii. (1846) 49, in obs. 3 = Lactuca Fischeriana.
viminea, Lam. Encyc. ii. 97 = Lactuca viminea.
virgata, Poir. Encyc. Suppl. ii. 328 = Prenanthes virgata.
virgata, J. & C. Presl, Delic. Prag. 116 = juncea.
viscosa, Gilib. Fl. Lituan. i. 232 = juncea.

CHONDROCARPUS, Nutt. Gen. Am. ii. in corrigend. (1818) = **Hydrocotyle**, Linn. (Umbellif.).
asiaticus, Nutt. l. c.—Cf. i. 177 = H. asiatica.
erectus, Wats. Bibliog. Ind. N. Am. Bot. 425, sphalm. = H. repanda.
repandus, Nutt. Gen. Am. ii. in corrigend.—Cf. i. 177 = H. repanda.
sibthorpioides, Sweet, Hort. Brit. ed. I. 185 = H rotundifolia.
trifurus, Nutt. Gen. Am. ii. in corrigend.—Cf. i. 177 = H. asiatica.

CHONDROCARPUS, Stev. in Bull. Soc. Nat. Mosc. iv (1832) 268 = **Astragalus**, Tourn. (Legumin.).
brachycarpus, Stev. l. c. = A. brachycarpus.
clavatus, Stev. l. c. xxix. (1856) ii. 149 = A. clavatus.
elongatus, Stev. l. c. iv. (1832) 268 = A. elongatus.
incurvus, Stev. l. c. = A. incurvus.
macrorhizus, Stev. l. c. = A. macrorrhizus.
nummularioides, Stev. l. c. xxix. (1856) ii. 149 = A nummularioides.
platyphyllus, Stev. l. c. = A. platyphyllus.

CHONDROCHILUS, Phil. in Linnaea, xxviii. (1856) 711 = **Chaetanthera**, Ruiz & Pav. (Compos.).
crenatus, Phil. l. c.—Chili.
involucratus, Phil. Fl. Atac. 27.—Chili.
parvifolius, Phil. in Linnaea, xxxiii. (1864–65) 114 — Chili.

CHONDROLAENA, Nees, Fl. Afr. Austr. Gram. 133 (1841) = **Prionachne**, Nees (Gramin.).
dentata, Steud. Nom. ed. II. i. 355, sphalm. = seq.
phalaroides, Nees, Fl. Afr. Austr. Gram. 134 — F. Ecklonii.

CHONDRODENDRON, Spreng. Syst. ii. 155, in syn. (1825) = **Chondodendron**, Ruiz & Pav. (Menisp.).

CHONDROLOMIA, Nees, in Mart. Fl. Bras. iii. i. 178 (1842) = **Solaria**, Berg. (Cyperac.).
Sellowiana, Nees, l. c. 174 = S. Sellowiana.

CHONDROPETALUM, Rottb. Progr. 12 (1772); Desc. 10. t. 3 (1778) = **Elegia**, Linn. (Restiac.).
deustum, Rottb. ll. cc. 12, 11 = E. deusta.
nudum, Rottb. ll. cc. 12, 11 = E. nuda.
tectorum, Rafin. Fl. Tellur. iv. 88 = E. deusta.

CHONDROPIS, Rafin. Fl. Tellur. iii. 29 (1836) = **Eraxum**, Linn. (Gentian.).
trinervis, Rafin. l. c. = E. zeylanicum.

CHONDROBRYNCHA, Lindl. Orch. Linden. 12 (1846). *ORCHIDEAE*, Benth. & Hook. f. iii. 548.
Chesterioni, *Reichb. f. in Gard. Chron.* (1879) ii. 648. —N. Granat.
fimbriata, *Reichb. f. in Saund. Refug. Bot.* t. 107.— N. Granat.
rosea, *Lindl. Orch. Linden.* 13.—Venezuela.

CHONDROSEA, Haw. Saxifr. Enum. 10 (1821) = **Saxifraga**, Tourn.
Aizoon, Haw. l. c. 11 = S. Aizoon.
aizoopsis, Jord. & Fourr. Ic. Pl. Eur. iii. 5 = S. Aizoon.
alpicola, Jord. & Fourr. Brev. Pl. Nov. fasc. I. 34 = S. Aizoon.
arcticoides, Haw. Saxifr. Enum. 14 = S. arcticoides.
baugesiana, Jord. & Fourr. Brev. Pl. Nov. fasc. I. 33 = S. Aizoon.
brevicaulis, Jord. & Fourr. Ic. Pl. Eur. ii. 3 = S. Aizoon.
Burseriana, Haw. Saxifr. Enum. 15 = S. Burseriana.
cuneata, Haw. l. c. 14 = S. cuneata.
calcarea, Jord. & Fourr. Ic. Pl. Eur. ii. 8 = S. Aizoon.
calycifera, Haw. Saxifr. Enum. 18 = S. media.

CHONDROSEA:—
carthusiana, Jord. & Fourr. Ic. Pl. Eur. ii. 3 = Saxifraga Aizoon.
cartilaginea, Haw. Saxifr. Enum. 12 = S. cartilaginea.
diapensoides, Haw. l. c. 13 = S. diapensoides.
flexipes, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
glabrata, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 34 = S. Aizoon.
glareosa, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
gracilescens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 35 = S. Aizoon.
intermedia, Haw. Saxifr. Enum. 11 = S. intermedia.
juniperina, Haw. l. c. 15 = S. juniperifolia.
laevigata, Jord. & Fourr. Ic. Pl. Eur. ii. 5 = S. Aizoon.
leptopetala, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 34 = S. Aizoon.
longifolia, Haw. Saxifr. Enum. 11 = S. longifolia.
luteo-purpurea, Haw. l. c. 13 = S. luteopurpures.
mutata, Haw. l. c. = S. mutata.
nitida, Haw. l. c. 16 = S. nitida.
orophila, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 33 = S. Aizoon.
patens, Haw. Saxifr. Enum. 14 = S. caesia.
patulipes, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
petrophila, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 34 = S. Aizoon.
punctillata, Jord. & Fourr. l. c. 33 = S. Aizoon.
pusilla, Jord. & Fourr. Ic. Pl. Eur. ii. 3 = S. Aizoon.
pyramidalis, Haw. Saxifr. Enum. 10 = S. Aizoon.
refracta, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
rosella, Jord. & Fourr. l. c. = S. Aizoon.
rosularis, Haw. Saxifr. Enum. 12 = S. Aizoon.
tenella, Haw. l. c. 15 = S. tenella.
valida, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 32 = S. Aizoon.
Vandellii, Haw. Saxifr. Enum. 15 = S. Vandelli.
virgata, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
viridula, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 33 = S. Aizoon.

CHONDROSIA, Benth. in Journ. Linn. Soc. xix. (1881) 104, err. typ. = seq.

CHONDROSIUM, Desv. Journ. Bot. iii. (1813) 68 = seq.

CHONDROSUM, Desv. in Nouv. Bull. Soc. Philom. ii (1810) 188 = **Bouteloua**, Lag. (Gramin.).
Aschenbornianum, Nees, in Linnaea, xix. (1847) 699 = B. oligostachya.
ciliatum, Beauv. Agrost. 41, nomen = Chloris ciliata.
eriopodum, Torr. in Emory, Notes Mil. Reconnos. 158 = B. eriopoda.
exile, Fourn. Mex. Pl. Enum. Gram. 137, ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
fornewm, Torr. in Emory, Notes Mil. Reconnos. 158. t. 19 = B. hirsuta.
gracile, H. B. & K. Nov. Gen. et. Sp. i. 176. t. 58 = B. oligostachya.
hirsutum, Sweet, Hort. Brit. ed. I. 455, nomen = seq.?
hirtum, H. B. & K. Nov. Gen. et Sp. i. 176. t. 59 = B. hirsuta.
Humboldtianum, Kunth, Rev. Gram. i. 98 = B. ovata.
humile, Beauv. Agrost. 41 = B. prostrata.
Karwinskii, Fourn. Mex. Pl. Enum. Gram. 137, ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
microstachyum, Fourn. l. c. 138, ex Hemsl. l. c. 562 = B. polystachya.
oligostachyum, Torr. in Marcy, Rep. 300 = B. oligostachya.
Parryi, Fourn. Mex. Pl. Enum. Gram. 150, ex Hemsl. Biol. Centr. Am. Bot. iii. 562 = B. polystachya.
polystachyum, Benth. Bot. Voy. Sulph. 56 = B. polystachya.
procumbens, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = B. tenuis.
prostratum, Sweet, Hort. Brit. ed. I. 455 = B. prostrata.
scorpioides, Kunth, Rev. Gram. i. 94 = B. scorpioides.
simplex, Kunth, l. c. = B. prostrata.
subscorpioides, C. Muell. in Bot. Zeit. xiv. (1856) 847. —Calif.
tenue, Beauv. Agrost. 41 = B. tenuis.
Violetii, Fourn. Mex. Pl. Enum. Gram. 136, ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.

CHONDROSPERMUM, Wall. Cat. n. 2827 (1832) = **Myxopyrum**, Blume (Oleac.).
coriaceum, Wall. l. c. n. 2838 = M. smilacifolium.
laurifolium, Voigt, Hort. Suburb. Calc. 648 = M. smilacifolium.
smilacifolium, Wall. Cat. n. 2837 = M. smilacifolium.

CHONEMORPHA, G. Don, Gen. Syst. iv. 76 (1838). *APOCYNACEAE*, Benth. & Hook. f. ii. 720.
EPICHYSIANTHUS, Voigt, Hort. Suburb. Calc. 528 (1845).
antidysenterica, G. Don, l. c. = Holarrhena antidysenterica.
bantamensis, G. Don, l. c. = Echites bantamensis.
convolvuloides, *G. Don, l. c.*—Ins. S. Thomas.
coriacea, G. Don, l. c. = Echites coriacea.
cristata, G. Don, l. c. = Aganosma marginata.
densiflora, G. Don, l. c. = Echites densiflora.
dichotoma, G. Don, l. c. = Aganosma caryophyllata.
*grevei*era, G. Don, l. c. = macrophylla.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 662.—Reg. Himal.
inflata, G. Don, Gen. Syst. iv. 76 = Echites inflata.
macrophylla, *G. Don, l. c.*—Ind. or.; Malaya.
malabarica, *G. Don, l. c.*—Ind. or.
mollis, Miq. Fl Ind. Bat. ii. 444 = macrophylla.
pubescens, G. Don, in Sweet, Hort. Brit. ed. III. 458 = Holarrhena antidysenterica.
reticulata, G. Don, Gen. Syst. iv. 76 = Cryptolepis Buchanani.
wettsiia, G. Don, l. c. = Wrightia tomentosa ?

CHORETIS, Herb. Amaryll. 219. f. 85 (1837) = **Hymenocallis**, Salisb. (Amaryllid.).
galvestonensis, Herb. l. c. 221. t. 41. f. 84, 85 = H. galvestonensis.
glauca, Herb. l. c. 220. t. 85. f. 1 = H. glauca.

CHORETRUM, R. Br. Prod. 354 (1810). *SANTALACEAE*, Benth. & Hook. f. iii. 228.
Candollei, *F. Muell. ex Benth. Fl. Austral.* vi. 219.—Austral.
chrysanthum, F. Muell in Trans. Phil. Soc. Vict. i. (1855) 28 = glomeratum.
glomeratum, *R. Br. Prod.* 354.—Austral.
lateriflorum, A. DC. in DC. Prod. xiv. 675 = Candollei.
lateriflorum, *R. Br. Prod.* 354.—Austral.
oxycladum, *F. Muell. Fragm.* i. 21.—Austral.
pauciflorum, A. DC. in DC. Prod. xiv. 676 = lateriflorum.
Preissianum, Miq. in Lehm. Pl. Preiss. i. 608 = Leptomeria Preissiana.
spicatum, *F. Muell. Fragm.* i. 21.—Austral.
spinosum, Miq. in Lehm. Pl. Preiss. i. 609 = Leptomeria spinosa.

CHORIBENA, Steud. Nom. ed. II. i. 355 (1840), ephalm. = **Chorilaena**, Endl. (Rutac.).

CHORICARPHA, Boeck. in Flora, xli. (1858) 19 = **Leptrionia**, Rich. (Cyperac.).
aphylla, Boeck. l. c. 20 = L. mucronata.

CHORICERAS, Baill. Adansonia, xi. (1873) 119 = **Distillaria**, F. Muell. (Euphorbiac.).
australiana, Baill. l. c. 120 = D. tricornis ?

CHORILAENA, Endl. Enum. Pl. Hueg. 17 (1837). *RUTACEAE*, Benth. & Hook. f. i. 295.
angustifolia, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 10 = Phebalium dicnemum.
hirsuta, *Benth. Fl. Austral.* i. 357.—Austral.
myoporoides, *Endl. ex Heynh. Nom.* ii. 140.—Austral.
quercifolia, *Endl. Enum. Pl. Hueg.* 17.—Austral.

CHORIOPHYLLUM, Benth. in Hook. Ic. Pl. t. 1280 (1859). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 260.
malayanum, *Benth. l. c.*—Malaya.

CHORIOZANDRA, Steud. Nom. ed. II. i. 355 (1840), ephalm. = **Chorizandra**, R. Br. (Cyperac.).

CHORIPETALUM:—
australianum, F. Muell. Fragm. iii. 36 (= *Samara australiana*).—Australi.
Benthamii, Hance, in Walp. Ann. iii. 10 = *obovatum*.
obovatum, Benth. in Hook. Lond. Journ. Bot. i. (1842) 490.—Ins. Hongkong.
Roxburghianum, Wall. ex Voigt, Hort. Suburb. Calc. 332, nomen = *obovatum* ?
undulatum, A. DC. in Trans. Linn. Soc. xvii. (1837) 122 = E. Nagushia.
viridiflorum, A. DC. in DC. Prod. viii. 88 = E. viridiflora.

CHORISANDRA, Benth. & Hook. f. Gen. iii. 1057 (1883) err. typ. = **Chortisandra**, R. Br. (Cyperac.).

CHORISANDRA, Wight, Ic. vi. 13. t. 1994 (1853) = **Fluggea**, Willd. (Euphorbiac.).
pinnata, Wight, l. c. = F. microcarpa.

CHORISANTHERA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 109 = **Pentarhaphia**, Lindl. (Gesnetrac.).
tenera, Oerst. l. c. = P. tenera.

CHORISEMA, Fisch. Ind Sem. Hort. Petrop. viii. 54 (1841) err. typ. = **Chorisema**, Labill. (Legumin.).

CHORISIA, H. B. & K. Nov. Gen. et Sp. v. 295. t. 485 (1821). *MALVACEAE*, Benth. & Hook. f. i. 210.
crispiflora, *H. B. & K. l. c.* 297.—Bras.
insignis, *H. B. & K. l. c.* 485. f. 1.—Peruv.
rosea, *Sem. Bot. Voy. Herald*, 84.—N. Granat.
speciosa, *St. Hil. Pl. Us. Bras.* t. 68.—Bras.
ventricosa, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1828) 103. t. 9 = crispiflora.

CHORISIS, DC. Prod. vii. 177 (1838) = **Lactuca**, Tourn. (Compos.).
repens, DC. l. c. 178 = L. repens.

CHORISMA, D. Don, in Edin. N. Phil. Journ. (1828-29) 208 = CHORISIS, DC. = **Lactuca**, Linn. (Compos.).
repens, D. Don, l. c. = L. repens.

CHORISMA, Lindl in Sweet, Geran. sub t. 79 (1830) = **Pelargonium**, L'Herit. (Geraniac.).
flavescens, Eckl. & Zeyh. Enum. 71 = P. tetragonum.
tetragona, Lindl. in Sweet, Geran. sub t. 79 = P. tetragonum.

CHORISPERMUM, R. Br. in Ait. Hort. Kew. ed. II. iv. 129 (1812) = **Chorispora**, R. Br. (Crucifer.).
arcuatum, Andrz. ex DC. Syst. ii. 486 = Chorispora tenella.
ibericum, Andrz. ex DC. l. c. 487 = Chorispora iberica.
sibiricum, Andrz. ex DC. l. c. = Chorispora sibirica.
strictum, Andrz. ex DC. l. c. 486 = Diptychocarpus strictus.
tenellum, R. Br. in Ait. Hort. Kew. ed. II. iv. 129 = Chorispora tenella.

CHORISPORA, R. Br. ex DC. Syst. ii. 485 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 102.
CHORISPERMUM, R. Br. in Ait. Hort. Kew. ed. II. iv. 129 (1812).
Bungeana, Fisch. & Mey. in Enum. Pl. Nov. Schrenk, 96 = exscapa.
compressa, Hoiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 385 = iberica ?
elegans, Cambess. in Jacquem. Voy. Bot. 15. t. 14 = sabulosa.
exscapa, *Bunge, in Ledeb. Fl. Ross.* i. 169.—Sibir. altaic.
Greigi, *Regel, Gartenfl.* (1879) 257. t. 984.—As. temp.
hispida, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) 1. 266.—As. temp.
iberica, *DC. Syst.* ii. 487.—As. Min.; Persia.
macropoda, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) 1. 109.—Songaria.
persica, *Boiss. in Ann. Sc. Nat. Sér. II.* xvii. (1842) 384.—Persia.
sabulosa, *Cambess. in Jacquem. Voy. Bot.* 15. t. 15.—Reg. Himal.; Afghan.
sibirica, *DC. Syst.* ii. 487.—Sibir. altaic.; Tibet.

CHORISPORA:—
sibirica, Regel & Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) 1. 866 = macropoda.
songanica, *Fisch. & Mey. Enum. Pl. Nov. Schrenk*, ii. 57.—Songar.
stenopetala, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 339.—Tashkent.
stricta, DC. Syst. ii. 426 = Diptychocarpus strictus.
syriaca, *Boiss. in Ann. Sc. Nat.* Sér. iii. xvii. (1842) 354.—Syria.
tenella, *DC. Syst.* ii. 435.—Reg. Caucas. et Himal.; Persia; Afghanist.

CHORISTEA, Thunb. Prod. Pl. Cap. 168 (1800); Nov. Gen. 179 (1801) = Didelta, L'Hérit. (Compos.).
carnosa, Thunb. ll. cc. 168, 180 = D. carnosa.
glabra, Soland. ex Pole. Encyc. v. 507 = D. spinosa.
spinosa, Thunb. Prod. Pl. Cap. 163; Nov. Gen. 180 = D. spinosa.

CHORISTES, Benth. Pl. Hartw. 68 (1840) = Deppea, Cham. & Schlecht. (Rubiac.).
cornifolia, Benth. l. c. 64 = D. cornifolia.
excelsa, Benth. l. c. = Psychotria excelsa.
obtusifora, Benth. l. c. 68 = D. obtusiflora.

CHORISTYLIS, Harv. in Hook. Lond. Journ. Bot. i. (1842) 19. *SAXIFRAGEAE*, Benth. & Hook. f. i. 647.
rhamnoides, *Harv. l. c.*—Afr. austr.

CHORITAENIA, Benth. in Benth. & Hook. f. Gen. i. 907 (1867). *UMBELLIFERAE*, Benth. & Hook. f. i. c.
Pappea, Sond. & Harv. Fl. Cap. ii. 562 (1865).

CHORIZANDRA, Benth. & Hook. f. Gen. iii. 975, in texts (1880) = CHORISANDRA, Wight = Fluggea, Willd. (Euphorb.).

CHORIZANDRA, R. Br. Prod. 321 (1810). *CYPERACEAE*, Benth. & Hook. f. iii. 1057.
cymbaria, *R. Br. l. c.*—Australi.
enodis, *Nees, in Linn. Pl. Preiss.* ii. 78.—Austral.
multiarticulata, *Nees, in Ann. Nat. Hist.* Ser. i. vi. (1841) 48.—Australi.
sphaerocephala, *R. Br. Prod.* 321.—Austral.

CHORIZANTHE, R. Br. ex Benth. in Trans. Linn. Soc. xvii. (1836) 416. t. 19. *POLYGONACEAE*, Benth. & Hook. f. iii. 98.
Acanthogonum, Torr. in Pacif. Rail. Rep. iv. 132 (1856).
Mucronea, Benth. in Trans. Linn. Soc. xvii. (1836) 419. t. 20.
Trigonocarpus, Bert. ex Steud. Nom. ed. II. ii. 716 (1841).
angustifolia, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 167.—Calif.
brevicornis, *Torr. in Bot. Mex. Bound.* 177.—Am. bor. occ.
Breweri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 270. —Calif.
californica, *A. Gray, in Proc. Bost. Soc. Nat. Hist.* vii. (1861) 149.—Calif.
Chevalandi, *Parry, in Proc. Davenp. Acad.* iv. (1884) 68.—Am. bor. occ.
commixta, *Remy, in C. Gay, Fl. Chil.* v. 287.—Chili.
corrugata, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 198.—Calif.
dasyantha, *Phil. in Linnaea*, xxxiii. (1864–65) 237.—Chili.
diffusa, *Benth. Pl. Hartw.* 333.—Calif.
discolor, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 167.—Calif.
Douglasii, *Benth. in Trans. Linn. Soc.* xvii. (1836) 418.—Calif.
fernandina, *S. Wats. Bot. Calif.* ii. 481.—Calif.
fimbriata, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 168.—Calif.
flavescens, *Phil. in Linnaea*, xxxiii. (1864–65) 236.— Chili.
frankeniciides, *Remy, in C. Gay, Fl. Chil.* v. 288.— Chili.
glabrescens, *Benth. in Trans. Linn. Soc.* xvii. (1836) 418.—Chili.
Kingii, *Phil. in Anal. Univ. Chil.* (1873) 536.—Chili.

CHORIZANTHUS:—
laciniata, *Torr. in Pacif. Rail. Rep.* vii. 19.—Calif.
Lasiarrhus, *Parry, in Proc. Davenp. Acad.* iv. (1884) 68.—Am. bor. occ.
leptocerus, *S. Wats. in Proc. Am. Acad.* xii. (1877) 269.—Calif.
Macrotis, *Benth. in Trans. Linn. Soc.* xvii. (1836) 417. —Chili.
membranaceus, *Benth. l. c.* 419.—Chili.
nudicaule, Benth. Pl. Hartw. 333 = Douglasii.
nudicaule, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 166.—Calif.
Orcuttiana, *Parry, in Proc. Davenp. Acad.* iv. (1884) 54.—Am. bor. occ.
Palmeri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 271. —Calif.
pauciseta, *Benth. in Trans. Linn. Soc.* xvii. (1836) 417.—Chili.
Parryi, *S. Wats. in Proc. Am. Acad.* xii. (1877) 271.— Calif.
pedunculata, *Benth. in Trans. Linn. Soc.* xvii. (1836) 416.—Chili.
perfoliata, *A. Gray, in Proc. Bost. Soc. Nat. Hist.* vii. (1861) 146.—Calif.
polygonoides, *A. Gray, in Proc. Am. Acad.* viii. (1870) 197.—Calif.
procumbens, A. Gray, in Proc. Bost. Soc. Nat. Hist. vii. (1861) 148 = Xanti.
procumbens, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 167.—Calif.
pungens, *Benth. in Trans. Linn. Soc.* xvii. (1836) 419. Calif.
ramosissima, *Benth. l. c.* 417.—Chili.
rigida, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 198.—Am. bor. occ.
rosea, *Phil. in Linnaea*, xxxiii. (1864–65) 235.— Chili.
spinosa, *S. Wats. Bot. Calif.* ii. 481.—Calif.
staticoides, *Benth. in Trans. Linn. Soc.* xvii. (1836) 418.—Calif.
stellulata, *Benth. Pl. Hartw.* 333.—Calif.
tenuis, *Phil. in Linnaea*, xxxiii. (1864–65) 237.— Chili.
Thurberi, *S. Wats. in Proc. Am. Acad.* xii. (1877) 269 —Am. bor. occ.
umbellata, *Phil. in Linnaea*, xxxiii. (1864–65) 235.— Chili.
uncinata, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 167.—Calif.
uniaristata, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 196.—Calif.
vaginata, *Benth. in Trans. Linn. Soc.* xvii. (1836) 417. —Chili.
valida, *S. Wats. in Proc. Am. Acad.* xii. (1877) 271.— Calif.
virgata, *Benth. in Trans. Linn. Soc.* xvii. (1836) 416. t. 19. f. 1.—Chili.
Watsoni, *A. Gray, in Proc. Am. Acad.* viii. (1870) 199. —Calif.
Wheeleri, *S. Wats. l. c.* xii. (1877) 272.—Calif.
Xanti, *S. Wats. l. c.*—Calif.

CHORIZEMA, Labill. Voy. i. 408. t. 21 (1799). *LEGUMINOSAE*, Benth. & Hook. f. i. 467.
Orthotropis, Benth. in Lindl. Swan River App. 16 (1839).
angustifolium, *Benth. Enum. Pl. Huegy.* 28.—Austral.
Baueri, Benth. in Ann. Wien. Mus. ii. (1838) 71 = Isotropis pungens.
Baueri, Meisn. in Lehm. Pl. Preiss. i. 94; ii. 309 = Heschmanni.
Bentleyi, R. Grah. in Edinb. Phil. Journ. (Apr.-June 1831) 367 = Oxylobium scandens.
callistachys, F. Muell. Fragm. iv. 18 = Oxylobium callistachys.
capillipes, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 256 = angustifolium.
Chandleri, Hort. ex Nichols. Dict. Gard. i. 318 = varium.
cordatum, *Lindl. Bot. Reg.* (1838) t. 10.—Austral.
cordifolium, F. Muell. Fragm. iv. 17 = Oxylobium cordifolium.
coriaceum, Sm. in Trans. Linn. Soc. ix. (1808) 264 = Oxylobium retusum.
costatum, Meisn. in Lehm. Pl. Preiss. i. 88 = Dickgensii.
cytisoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 256.—Austral.

CHORIZEMA:—
davisioides, Meisn. in Lehm. Pl. Preiss. i. 84 = Mirbelia davisioides.
dentatostatum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 252 = angustifolium.
Dicksoni, *R. Grah. in Mennd, Botanist*, iii. t. 106.— Austral.
diversifolium, *A. DC. Pl. Rar. Jard. Genev.* 7e *Not.* (1856) 44. t. 8.—Austral.
elliptieum, F. Muell. Fl. Vict. ii. 89 = Oxylobium ellipticum.
ericifolium, *Meisn. in Lehm. Pl. Preiss.* ii. 209.— Austral.
ericoides, Hort. Belg. ex Heynh. Nom. ii. 141 = praec.?
Fawcum, Hemsley, in Moore & Ayres, Gard. Mag. Bot. i. 73. t. 7 = cordatum.
gomphocarpum, *Benth. ex A. Gray, in Bot. U. St. Expl. Exped.* i. 380.—Austral.
Henchmanni, *R. br. in Bot. Reg.* t. 986.—Austral.
heterophyllum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 255 = Oxylobium heterophyllum.
humile, *Turcz. l. c.* 254.—Austral.
illicifolium, *Labill. Voy.* i. 406.—Austral.
ilicifolium, Stend. Nom. ed. II. i. 836, sphalm. = Oxylobium trilobatum.
Leichhardtii, F. Muell. Fragm. iv. 20 = Isotropis illicoides.
linearе, F. Muell. Fragm. iv. 17 = Oxylobium lineare.
magnifolium, F. Muell. l. c. 18 = Mirbelia racemosa.
nanum, Sims, Bot. Mag. t. 1083 = ilicifolium.
nervosum, *T. Moore, in Gard. Comp.* (1852) 122.— Austral.
ovatum, Lindl. Bot. Reg. t. 1598 = rhombeum.
parviflorum, *Benth. in Ann. Wien. Mus.* ii. (1838) 71. —Austral.
parvifolium, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 253 = nervosum.
platyloboides, DC. Prod. ii. 108 = Mirbelia grandiflora.
pubescens, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 256 = Oxylobium carinatum.
Pulteneaee, F. Muell. Fragm. iv. 19 = parviflorum.
reticulatum, *Meisn. in Lehm. Pl. Preiss.* i. 84.— Austral.
rhombeum, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 9.— Austral.
rhombeum, Lodd. Bot. Cab. t. 1619 = diversifolium.
rhynchotropis, Meisn. in Lehm. Pl. Preiss. ii. 209 = Henchmanni.
scandens, Sm. in Trans. Linn. Soc. ix. (1808) 258 = Oxylobium scandens.
sericeum, Sm. l. c. = Brachysema undulatum.
spartioides, Lodd. Bot. Cab. t. 1958 = Isotropis striata.
spectabile, Lindl. Bot. Reg. (1841) t. 46 = diversifolium.
superbum, Lem. Illustr. Hortic. i. (1854) t. 29 = cordatum.
triangulare, Lindl. Bot. Reg. t. 1513 = ilicifolium.
trigonum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 254.—Austral.
trilobatum, Sm. in Trans. Linn. Soc. ix. (1808) 258 = Oxylobium trilobatum.
varium, *Benth. ex Lindl. Bot. Reg.* (1839) t. 49; *Misc.* 45.—Austral.

CHORIZOTHECA, Meisl. Arg. in Linnaea, xxxii. (1863) 76 = Pseudantinia, Steud. (Scrophular.).
micranthoides, Meisl. Arg. l. c. = F. virgatus.

CHOROBANCHE, Presl, Bot. Bemerk. 91 (1846) (Chorobane) = Orobanche, Linn.
Indica, Presl, l. c. = O. Indica.

CHOROSEMA, Bronga. Enum. Gen. 126 (1845) = Chorizema, Labill. (Legumin.).

CHOROZEMA, Sm. in Trans. Linn. Soc. ix. (1808) 251, sphalm. = Chorizema, Labill. (Legumin.).

CHORYZEMUM, Rose, Encyc. Agric. vii. 297 (1835) = prase.

CHOTECKIA, Stend. Nom. ed. II. i. 366 (1840) sphalm. = seq.

CHOTEKIA, Opiz & Corda, in Flora, xiii. (1830) 1. 38 = **Dysophylla**, Blume (Labiat.).
sericea, Opiz & Corda, l. c. 38 = D. quadrifolia.

CHOULETTIA, Pomel, Nouv. Mat. Fl. Atl. 61 (1874) = **Gaillonia**, A. Rich. (Rubiac.).
Rebouliana, Pomel, l. c. = G. Rebouliana.

CHRESTA, Vell. Fl. Flum. viii. t. 150, 151 (1827) = **Erexanthus**, Linn. (Compos.).
alpestris, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 289 = Vernonia alpestris.
angustifolia, Gardn. l. c. 240 = E. angustifolius.
cordata, Vell. Fl. Flum. viii. t. 150 = E. sphaeroorphalus.
exsucca, DC. Prod. v. 35 = E. exsuccus.
intermedia, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 236 = E. sphaeroorphalus.
lanceolata, Vell. Fl. Flum. viii. t. 151 = Vanillosmopsis erythropappa.
plantaginifolium, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 241 = E. plantaginifolius.
pycnocephala, DC. Prod. v. 35 = E. pycnocephalus.
scabigera, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 241 = E. scabiger us.
speciosa, Gardn. l. c. 240 = E. speciosus.
sphaerocephala, DC. in Deless. Ic. Sel. iv. 2. t. 6 = E sphaerocephalus.
spathulaefolia, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 241 = E. plantaginifolius.

CHRISANTHEMUM, Neck. Delic. Gallo-Belg. ii. 358 (1768) = **Chrysanthemum**, Tourn. (Compos.).

CHRISOSPLENIUM, Neck. Delic. Gallo-Belg. i. 190 (1768) = **Chrysosplenium**, Tourn. (Saxifrag.).

CHRISTANNIA, Presl, Rel. Haenk. ii. 91. t. 67 (1831) = **Banara**, Aubl. (Samydac.).
salicifolia, Presl, l. c. = B. incana.

CHRISTANNIA, Walp. Rep. i. 860 (1842), sphalm. = **Christiana**, DC. (Tiliac.).

CHRISTIA, Moench, Meth. Suppl. 89 (1802) = **Lourea**, Neck. (Legumin.).
lunata, Moench, l. c. 40 = L. vespertilionis.

CHRISTIANA, DC. Prod. i. 516 (1824). *TILIACEAE*, Benth. & Hook. f. i. 282.
africana, *DC. l. c.*—Afr. trop.
corchifolia, Hook. f. in Niger Fl. 288 = africana.
madagascariensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 543.—Madag.

CHRISTIBONIA, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 153. *OROBANCHACEAE*, Benth. & Hook. f. ii. 982.
CAMPBELLIA, Wight, Ic. iv. 5. t. 1434, 1425 (1850).
OLIGOPHOLIS, Wight, l. c. t. 1422 (1850).
albida, *Thw. ex Hook. f. Fl. Brit. Ind.* iv. 323.—Zeylan.
aurantiaca, Wight, Ic. iv. t. 1436 = bicolor.
bicolor, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 160.—Ind. or.
calcarata, *Wight, Ic.* iv. t. 1426.—Ind. or.
grandiflora, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 155 = tricolor.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit: Ind.* iv. 331.—Reg. Himal.
Lawii, *Wight, Ic.* iv. t. 1427.—Ind. or.
neilgherica, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 157.—Ind. or.
pallida, Gardn. l. c. 159 = bicolor.
Stocksii, Hook. Ic. Pl. t. 336 = calcarata.
subacaulis, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 162.—Ind. or.
Thwaitesii, *Trim. in Journ. Bot.* xxiii. (1885) 240.—Zeylan.
tricolor, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 156.—Zeylan.
tubulosa, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 331.—Ind. or.
unciniae, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 161 = neilgherica.

CHRISTMANNIA, Dennst. Schlüssel. Hort. Malab. 30 (1818). 'Genus delendum,' Benth. & Hook. f. iii. 150. CURCUMA, Rafin. Sylva Tellur. 154 (1838).
Couronii, Dennst. l. c.—Ind. or.

CHRISTOLEA, Cambess. in Jacquem. Voy. Bot. 17. t. 17 (1837). *CRUCIFERAE*, Benth. & Hook. f. i. 80.
crassifolia, *Cambess. l. c.*—Tibet. occ.

CHRISTOPHORIANA, Tourn. ex Rupp. Fl. Jen. ed. Hall. 94 (1745) = **Actaea**, Linn. (Ranunculac.).
spicata, Moench, Meth. 276 = A. spicata.
vulgaris, Rupr. Fl. Caucas. 44 = A. spicata.

CHRISTYA, Ward & Harv. in Hook. Journ. Bot. iv. (1842) 184. t. 21 = **Strophanthus**, DC. (Apocynac.).
speciosa, Ward & Harv. l. c. = S. capensis.

CHRITMUM, Brot. Fl. Lusit. i. 436 (1804), sphalm. = Crithmum, Linn. (Umbell.).

CHROILEMA, Bernh. Del. Sem. Hort. Erf. (1840); et in Linnaea, xv. (1841) Litt. 90 = **Aplopappus**, Cass. † (Compos.).
subcanescens, Bernh. ll. cc.—Chili.

CHROMANTHUS, Phil. Sert. Mend. Alt. 14 (1871). *PORTULACEAE*?
CHROMATOLEPIS, Dulac, Fl. Hautes-Pyr. 526 (1867) = **Carlina**, Linn. (Compos.).
acaulis, Dulac, l. c. 527 = Carl. acaulis.
lanata, Dulac, l. c. = Carl. lanata.
magna, Dulac, l. c. = Carl. acanthifolia.
vulgaris, Dulac, l. c. = Carl. vulgaris.

CHROMATOPOGON, F. W. Schmidt, Samml. Phys. Aufs. i. 275 (1795) = **Soonsros**, Linn. (Compos.).
Jacquini, F. W. Schmidt, l. c. = S. hirsuta.

CHROMOCHITON, Cass. in Dict. St. Nat. lvi. 230 (1828) = **Cassinia**, R. Br. (Compos.).
aculeata, Cass. l. c. = Cassinia aculeata.
affinis, Cass. l. c. = Cassinia aculeata.
aurea, Cass. l. c. = Cassinia aurea.

CHROMOLAENA, DC. Prod. v. 183 (1836) = **Eupatorium**, Linn. (Compos.).
alternifolia, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 465 = E. stachyophyllum.
Candolleana, Gardn. l. c. 464 = E. horminoides.
decumbens, Gardn. l. c. 466 = E. ocoumbens.
opelaecae, Gardn. l. c. vi. (1847) 496 = E. lupulinum.
horminoides, DC. Prod. v. 183 = E. horminoides.
leucocephala, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 466 = E. horminoides.
pratensis, Gardn. l. c. i. (1842) 176 = E. macrocephalum.

CHROMOLEPIS, Benth. Pl. Hartw. 40 (1840). *COMPOSITAE*, Benth. & Hook. f. ii. 366.
heterophylla, *Benth. l. c.*—Mexic.

CHRONE, Dulac, Fl. Hautes-Pyr. 512 (1867) = **Eupatorium**, Linn. (Compos.).
heterophylla, Dulac, l. c. = E. cannabinum.

CHRONOBASIS, DC. ex Benth. & Hook. f. Gen. ii. 457 (1873) = **Urvinia**, Gaertn. (Compos.).

CHRONOPAPPUS, DC. Prod. v. 84 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 336.
bifrons, *DC. l. c.*—Bras.

CHROSPERMA, Raf. Neogenyt. 3 (1825) = **Zygadenus**, Michx. (Liliac.).
lentum, Rafin. l. c. = Z. muscotoxicum.

CHROSOPHORA, Neck. Elem. ii. 837 (1790). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 302.
CROSOPHORA, A. Juss. Tent. Euph. 27 (1824).
LEPIDOCROTON, Presl, Epim. Bot. 213 (1850).
RICINOIDES, Tourn. ex Moench, Meth. 356 (1794).
TOURNEFORTIA, Nimol. ex Scop. Introd. 343 (1777).
Brocchiana, *Schweinf. Pl. Nilot.* 9.—Afr. trop.
Burmanni, Spreng. Syst. iii. 851 = plicata.

CHROSOPHORA:—
gracilis, *Fisch. & Mey. ex Ker. in Bull. Soc. Nat. Mosc.* (1839) 171, nomen; *Boiss. Fl. Orient.* iv. 1140.—As centr.
hierosolymitana, Spreng. Syst. iii. 850 = verbascifolia.
integrifolia, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 490 = verbascifolia.
mollissima, A. Juss. Tent. Euph. 28 = Mallotus viscinoides.
obliqua, *A. Juss. l. c.*—Afr. bor. ; As. occ. ; Ind. or.
obliqua, Schweinf. Pl. Nilot. 18 = plicata.
oblongifolia, A. Juss. Tent. Euph. 28 = obliqua.
parvifolia, Klotzsch, ex Schweinf. Pl. Nilot. 11 = plicata.
peltata, Labill. Sert. Austr.-Caled. 74. t. 75 = Codiaeum inophyllum.
plicata, *A. Juss. Tent. Euph.* 28.—Europ. austr. ; Afr. bor. ; As. occ. ; Ind. or.
prostrata, Dalz. & Gibs. Bomb. Fl. 233 = plicata.
Rottleri, A. Juss. Tent. Euph. 26 = plicata.
subulata, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 446 = gracilis.
senegalensis, *A. Juss. Tent. Euph.* 26.—Afr. trop.
Sieberi, Presl, Bot. Bemerk. 109 = verbascifolia.
tinctoria, *A. Juss. Tent. Euph.* 28. t. 7.—Europ. austr. ; Orient ; Ind. or. ; Afr. bor.
tinctoria, Muell. Arg. in DC. Prod. xv. ii. 749 = obliqua.
verbascifolia, *A. Juss. Tent. Euph.* 28.—Reg. Mediterr.; Orient.

CHROZORRHIZA, Ehrh. Beitr. iv. 147 (1789) = **Asperula**, Linn. (Rubiac.) (A. tinctoria).

CHRYSA, Rafin. in Med. Repos. N. York, II. v. (1809) 350 = Coptis, Salisb. (Ranunculac.).

CHRYSACTINIA, A. Gray, in Mem. Am. Ac. N. S. iv. (1849) (Pl. Fendl.) 93. *COMPOSITAE*, Benth. & Hook. f. ii. 412.
mexicana, *A. Gray, l. c.*—Mexic.

CHRYSACTINIUM, Wedd. Chlor. And. i. 212. t. 39. (1856) = **Lidabum**, Adans. (Compos.).
acaule, Wedd. l. c. = L. acaule.

CHRYSALIDOCARPUS, H. Wendl. in Bot. Zeit. xxxvi. (1878) 171. *PALMAE*, Benth. & Hook. f. iii. 892.
lutescens, *H. Wendl. l. c.*—Madag.

CHRYSANGIA, Link, Handb. i. 632 (1829) = **Musschia**, Dum. (Campanul.).
aurea, Link, l. c. = M. aurea.

CHRYSANTHELLINA, Cass. in Dict. Sc. Nat. xxv. 391 (1822) = Chrysanthellum, Rich. (Compos.).
fasciculata, Cass. l. c. = Chrysanthellum procumbens.
gracilis, Cass. l. c. 392 = Chrysanthellum procumbens.
Swartzii, Cass. l. c. = Chrysanthellum procumbens.

CHRYSANTHELLUM, Rich. in Pers. Syn. ii. 471 (1807). *COMPOSITAE*, Benth. & Hook. f. ii. 389.
ADENOSPERMUM, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 318.
CHRYSANTHELLINA, Cass. in Dict. Sc. Nat. xxv. 391 (1822).
COLLARA, Spreng. Syst. iii. 622 (1826).
HINTERHUBERA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 389 (1873).
NEURACTIS, Cass. in Dict. Sc. Nat. xxxiv. 496 (1825).
SEBASTIANIA, Bertol. Luccubr. (1833) 87.
UCAUOU, Adans. Fam. ii. 131 (1763).
abyssinicum, Sch. Bip. in Flora, xxv. (1842) 440 = Microlecane abyssinica.
americanum, *Vatke, in Bremen Abh.* ix. (1885) 122.—Am. trop.
bolivianse, *Sch. Bip. in Bull. Soc. Fr.* xii. (1865) 82, nomen.—Boliv.
crectum, *Anderss. in Vet. Akad. Handl. Stockh.* 1858 (1855) 188.—Ins. Galapag.
indicum, *DC. Prod.* v. 681.—As et Afr. trop.
integrifolium, *Steetz, in Seem. Bot. Voy. Herald*, 160. —Panama.
procumbens, *A. Rich. in Pers. Syn.* ii. 471.—Am. trop.
pusillum, *Benth. & Hook. f. in Trans. Linn. Soc.* xx. (1851) 214 —Ins. Galapag.
senegalense, DC. Prod. v. 681 = indicum.
Swartzii, DC. l. c. = procumbens.

CHRYSANTHEMOIDES, Tourn. ex Medic. Phil. Bot. i. 159 (1789) = **Osteospermum**, Linn. (Compos.).
coerulea, Moench, Meth. 584 = O. oxarulearum.
pisiformis, Medic. Phil. Bot. i. 159 = O. monilliferum.

CHRYSANTHEMUM, [Tourn.] Linn. Syst. ed. 1 (1735). *COMPOSITÆ*, Benth. & Hook. f. ii. 434.
Ammanthus, Boiss. & Heldr. Diagn. Ser. I. xi. 18 (1849).
Argyranthemum, Webb, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 258. t. 90 ad 96 (1836–50).
Balsamita, Desf. in Act. Soc. Hist. Nat. Par. i. (1795) t. 1.
Brachanthemum, DC. Prod. vi. 44 (1837).
Centrachaena, Schott, Ind. Sem. Hort. Vindob. (1828), ex Reichb. Ic. Pl. Crit. iv. 47 (1827).
Centrospermum, Spreng. Nov. Prov. 19 (1818); Syst iii. 862 (1826).
Coleostephus, Cass. in Dict. Sc. Nat. xli. 43 (1826).
Decaneurum, Sch. Bip. Tanacet. 44 (1844).
Dendranthema, Des Moul. in Act. Soc. Linn. Bord. xx. (1855) 561.
Endopappus, Sch. Bip. in Bonplandia, viii. (1860) 269.
Gleborus, Cass. Dict. Sc. Nat. xli. 41 (1826).
Glossopappus, Kunze, in Flora, xxix. (1846) 748.
Gymnocline, Cass. in Bull. Soc. Philom. (1816) 199; et in Dict. Sc. Nat. xx. 119 (1821).
Heteranthemis, Schott, in Oken, Isis, (1818) 821.
Heteromera, Pomel. Nouv. Mat. Fl. Atl. 60 (1874).
Hymenostemma, Kunze, in Flora, xxix. (1846) 699.
Ismelia, Cass. in Dict. Sc. Nat. xli. 40 (1826).
Krenseria, Dur. in Duch. Rev. Bot. i. (1846) 364.
Leucanthemum, [Tourn.] DC. Prod. vi. 45 (1837).
Matricaria, Hall. ex Scop. Fl. Carn. ed. II. ii. 147 (1772).
Monoptera, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 253 (1836–50).
Myconia, Neck. Elem. i. 22 (1790).
Otospermum, Willk. in Bot. Zeit. xxii. (1864) 251.
Phalacrocarpum, Willk l. c. 252 (1864).
Phalacrodiscus, Less. Syn. Comp. 258 (1832).
Phalacroglossum, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276 (1855–80).
Pinardia, Cass. in Dict. Sc. Nat. xli. 38 (1826).
Plagius, L'Hérit. Dis. ex DC. Prod. vi. 136 (1837).
Pontia, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 215.
Prolixia, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 250 (1836–50).
Prolongoa, Boiss. Voy. Bot. Espagne, 230. t. 29 a (1840).
Pyrethrum, Hall. Enum. Stirp. Helv. ii. 720 (1742).
Richteria, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1846) 196.
Stigmatotheca, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 255 (1836–50).
Tridactylina, Sch. Bip. Tanacet. 48 (1844).
Xanthophthalmum, Sch. Bip. l. c. 17 (1844).
absinthiifolium, Fisch. ex DC. Prod. vi. 46 = sinoatum.
Achillea, *Linn. Syst. ed.* XII. 962.—Italia.
achilleæfolium, *DC. Prod.* vi. 66.—Reg. Caucas.; Sibiria.
adustum, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) i. 415 = arctiocum.
alpestre, DC. Prod. vi. 62 = coronopifolium.
alpinum, Bieb. Casp. 211 = cancasicum.
alpinum, *Linn. Sp. Pl.* 889.—Europ.
anethifolium, *Brouss. ex Willd. Enum. Hort. Berol.* 904, in nota.—Ins. Canar.
angustifolium, *Rafin. Herb.* 25; *Atl. Journ.* 178.—Am. bor.
anomalum, *Leg. Variet. Cienc.* ii. iv. (1805) 40.—Hispan.
apetalum, *Hort. ex Steud. Nom.* ed. II. i. 856.—Europ.
aragonense, Asso, Syn. 132. t. 9. f. 1 = pulverulentum.
arcticum, *Linn. Sp. Pl.* 886.—Reg. arct.
argenteum, *Willd. Sp. Pl.* iii. 2146.—Armenia.
artemisiæfolium, Klatt, in Sitz. Akad. Muench. (1876) 58 = C. Richteria.

CHRYSANTHEMUM:—
arvense, Herb. Mex. ex DC. Prod. v. 311 = Xanthocephalum humile.
asteroides, *Desf. Tabl.* ed. II. 118, nomen.—Hab.?
Atkinsoni, *C. B. Clarke, Comp. Ind.* 147.—Reg. Himal.
atlanticum, *Ball, in Journ. Bot.* xi. (1873) 366.—Marocco.
atratum, Georgi, Beschr. Russ. Reich. iii. iv. 1252 = maximum!
atratum, *Linn. Sp. Pl.* ed.-II. 1252.—Europ.
Balsamita, Baill. Hist. Pl. viii. 311 = Tanacetum Balsamita.
Balsamita, *Linn. Sp. Pl.* ed. II. 1262.—As. occ.
Barrelieri, DC. Prod. vi. 68 = coronopifolium.
Bauhini, Tausch, in Syll. Ratisb. ii. (1828) 251 = atratum.
bellidiflorum, Willd. ex Steud. Nom. ed. II. i. 856 = Bellis dubia.
Biebersteinianum, Adam, in Weber & Mohr, Beitr. i. 70 = Anthemis Biebersteiniana.
bipinnatum, *Linn. Sp. Pl.* 890.—Reg. arct.
Bocconi, *Pourr. ex DC. Prod.* vi. 61.—Hispan.
breviradiatum, Hort. ex DC. Prod. vi. 64 = coronarium.
Broussonetii, *Balb. ex Pers. Syn.* ii. 461.—Ins. Canar.
caespitosum, *Weinm. Cat. Hort. Dorp.* 48.—Hab.?
carinatum, *Schousb. Vestr. Marokko*, 198. t. 6.—Marocco.
carneum, Steud. Nom. ed. I. 192 = coccineum.
carnosulum, DC. Prod. vi. 65 = Phymaspermum carnosulum.
carnosum, Pers. Syn. ii. 460 = glabrum.
carolinianum, Walt. Fl. Carol. 204 = Boltonia Caroliniana.
Cataranche, *Ball, in Journ. Bot.* xi. (1873) 366.—Marocco.
cascasicum, *Pers. Syn.* ii. 462.—Reg. Caucas.
catenarium, DC. ex Steud. Nom. ed. I. 192, 193 = monspeliense.
centrospermum, Bernh. ex Reichb. Ic. Pl. Crit. iv. 47 = viscosum.
ceratophylloides, All. Fl. Pedem. i. 190 = stratum.
Chamomilla, Bernh. Syst. Verz. Erf. 145 = Matricaria Chamomilla.
cinerariæfolium, *Vis. Fl. Dalmat.* ii. 88. t. 8.—Dalmatia.
cinerarium, Steph. in Ledeb. Fl. Ross. ii. 542 = sinuatum.
coccineum, *Willd. Sp. Pl.* iii. 2144.—Reg. Caucas.; Persia.
coronarium, *Linn. Sp. Pl.* 890.—Reg. Mediterr.
coronopifolium, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 208.—Ins. Canar.
coronopifolium, Steud. Nom. ed. II. i. 856 = grandiflorum.
coronopifolium, *Vill. Fl. Delphin.* 98; *Hist. Pl. Dauph.* iii. 202.—Corsica.
coronopifolium, Willd. Herb. ex Boiss. Fl. Orient. iii. 846 = coccineum.
corsicum, Sieber, ex DC. Prod. vi. 47 = coronopifolium, *Vill.*
corymbiflorum, *Linn. Sp. Pl.* ed. II. 1251.—Helv.; Austria.
corymbiferum, Pall Reise, i. 196 = corymbosum.
corymbosum, *Linn. Sp. Pl.* 890.—Europ.; Afr. bor.; Reg. Caucas.
crithmifolium, Brouss. ex Willd. Enum. Hort. Berol. 904, in syn.; Buch, in Berl. Acad. Abh. 1816–17 (1819) 276 = anethifolium.
cuneatum, Rœmb. Hort. Beng. [101]; Fl. Ind. iii. 436 = Boltonia indica.
dasyfolium, *Pers. Syn.* ii. 462.—Hab.?
dentatum, *Steud. Nom.* ed. I. 192.—Syria.
dentatum, Gilib. Fl. Lituan. i. 219 = C. Leucanthemum.
diffusum, *Vell. Fl. Flum.* viii. t. 126.—Bras.
disciforme, C. A. Mey. Verz. Pl. Casc. 75 = Matricaria discoiformis.
discoideum, *All. Fl. Pedem.* i. 190.—Italia.
dissectum, *Lour. in Transa. Camb. Phil. Soc.* vi. (1838) reimpr. 17.—Ins. Canar.
elegans, Steud. Nom. ed. I. 192 = maximum.
ericæfolium, *Burm. f. Fl. Cap. Prod.* 27.—Afr. austr.
floridum, Salisb. Prod. 203 = frutescens.
fosculosum, *Linn. Sp. Pl.* 890.—Reg. Mediterr.

CHRYSANTHEMUM:—
foeniculaceum, Steud. Nom. ed. I. 193; DC. Prod. vi. 66 = anethifolium.
foliosum, Brouss. Hort. Monsp. (1804); ex DC. Prod. vi. 66 = grandiflorum.
frutescens, *Linn. Sp. Pl.* 887.—Ins. Canar.
frutescens, Thunb. Prod. Pl. Cap. 161 = Thunbergii.
fruticosum, Buch, in Abh. Berl. Acad. 1816–17 (1819) 275 = frutescens!
fruticosum, *Vell. Fl. Flum.* viii. t. 130.—Bras.
fruticulosum, *Ledeb. Pl. Alt. iv.* 117.—Sibir. altaic.
fuscatum, Desf. Fl. Atlant. ii. 288 = Matricaria fuscata.
Gayanum, Ball, in Journ. Linn. Soc. xvi. (1878) 509 = Mawii.
glabratum, Thunb. Prod. Pl. Cap. 161 = Matricaria glabrata.
glabrum, DC. Prod. vi. 49, in syn. = paludosum.
glaucum, Pers. Syn. ii. 461 = multifidum.
Gmelini, Turcz. in Bull. Soc. Nat. Mosc. (1838) 94 = arcticum.
gracile, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 207.—Ins. Canar.
gracilicaule, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 306 = montanum.
graminifolium, *Linn. Sp. Pl.* 889.—Europ.
grandiflorum, Brouss. Elench. Pl. Hort. Monsp. (1804) 15 = coronopifolium.
grandiflorum, Hook. in Parry, App. Second Voy. 398 = Matricaria inodora.
grandiflorum, Hort. Par. (1819) ex DC. Prod vi. 47 = lacteum.
grandiflorum, Lapeyr. Hist. Abr. Pl. Pyr. 597 = maximum.
grandiflorum, *Willd. Enum. Hort. Berol.* 904.—Ins. Canar.
Griffithii, C. B. Clarke, Compos. Ind. 148 = Stolickza.
haematomma, *Lowe, in Hook. Kew Journ.* viii. (1856) 284.—Ins. Madera.
Halleri, Suter, Fl. Helv. ii. 193 = coronopifolium, *Vill.*
Hermanni, Hoffngg. & Link, Fl. Port. ii. t. 102 = oppositifolium.
heterophyllum, *Willd. Sp. Pl.* iii. 2142.—Europ.
hirtum, Hort. ex Reichb. Ic. Pl. Crit. iv. 47 = viscidum.
hirtum, Thunb. Prod. Pl. Cap. 161 = Matricaria hirta.
hispidum, Rich. ex DC. Prod. vi. 47 = L. pallens.
humile, Spreng. Syst. iii. 684 = Xanthocephalum humile.
hybridum, Guss. Adnot. Cat. Pl. Bocc. 3 = C. Myconis.
incanum, Thunb. Prod. Pl. Cap. 161 = Pentzia virgata.
indicum, *Linn. Sp. Pl.* 889.—China; Japon.
indicum, Thunb. Fl. Jap. 320 = sinense.
inodorum, Linn. Sp. Pl. ed. II. 1253 = Matricaria inodora.
integrifolium, *Richards. in Frankl. 1st Journ. App.* ed. II. 33.—Reg. arct.
intermedium, Hort. Monsp. ex DC. Prod. vi. 66 = grandiflorum.
Ircutianum, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) ii. 177.—Sibiria.
italicum, *Linn. Mant.* i. 113.—Italia.
japonicum, Thunb. Fl. Jap. 321 = indicum.
lacerum, Auct. ex Desv. in Lam. Encyc. iii. 739, in syn. = pinnatifidum.
laciniatum, Gilib. Fl. Lituan. i. 218 = segetum.
lacustre, *Brot. Fl. Lusit.* i. 370.—Europ.
lanceolatum, *Pers. Syn.* ii. 460.—Europ.
lanceolatum, Vest, in Flora, iii. (1820) 4 = C. Leucanthemum.
lanuginosum, Genev. ex DC. Prod. vi. 57 = corymbosum.
larvatum, *Griseb. ex Pant. in Oestr. Bot. Zeitschr.* xxiii. (1873) 366.—Montenegro.
latifolium, DC. Prod. vi. 69 = grandiflorum.
leptophyllum, DC. l. c. 65 = nodosum.
Leucanthemum, *Linn. Sp. Pl.* 888.—Europ.; Asbor.
lidbeckioides, *Less. Syn. Comp.* 258.—Afr. austr.
macrocephalum, Viv. Fl. Lib. Spec. 56. t. 10. f. 4 = trifurcatum.
macrophyllum, *Waldst. & Kit. Pl. Rar. Hung.* i. 97. —Europ. or.; Reg. Caucas.
maurorum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 509. —Hispan.; Marocco.
maximum, *Anchera. Fl. Brandenb.* 229 = C. Leucanthemum.
Maresii, *Ball, in Journ. Bot.* xi. (1873) 366 (Maresii). —Marocco.
maritimum, *Pers. Syn.* ii. 462.—Europ. occ.

CHRYSANTHEMUM:—
Mawii, *Hook. f. Bot. Mag.* t. 5997.—Marocco.
maximum, *Ramond*, *in Bull. Soc. Philom.* ii. (1800) 140.—Mont. Pyren.
mexicana, *H. B. & K. Nov. Gen. et Sp.* iv. 299.—Mexic.
millefoliatum, *Linn. Syst.* ed. XII. 868.—Reg. Caucas.; As. Min.
minimum, Brot. Fl. Lusit. i. 379 = pulverulentum ?
minimum, Vill. Hist. Pl. Dauph. iii. 202 = alpinum.
monspelliense, *Linn. Sp. Pl.* 880.—Europ.
monspeliense, Schkuhr; Handb. iii. 131. t. 253 = arcticum.
montanum, *Linn. Sp. Pl.* 888.—Europ.
montanum, Willd. Sp. Pl. iii. 2148 = C. Leucanthemum.
moschatum, *Willd. ex Steud. Nom.* ed. I. 192, nomen; ed. II. i. 357.—Hab. ?
multicaule, *Desf. Fl. Atlant.* ii. 282.—Afr. bor.
multifidum, *Desf. Cat. Hort. Par.* ed. III. 170.—Orient.
Myconis, *Linn. Sp. Pl.* ed. II. 1254.—Reg. Mediterr.
nanum, Hook. Fl. Bor. Am. i. 320 = Blennosperma californicum.
nodosum, *DC. Prod.* vi. 65.—Afr. austr.
ochroleucum, *Mauf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 208.—Ins. Canar.
odoratum, Lowe, Man. Fl. Mad. i. 462 = pinnatifidum.
oppositifolium, *Brot. Fl. Lusit.* i. 381.—Lusit.
orcastrum, *Hance, in Journ. Bot.* xi. (1873) 108.—China.
orientale, Pers. Syn. ii. 462 = Anthemis Biebersteiniana.
osmitoides, *Harv. Thes. Cap.* ii. 38. t. 152.—Natal.
palæa, *Y. Gay, ex Perreym. in Guill. Arch. Bot.* ii. (1833) 548.—Europ.
pallens, Pers. Syn. ii. 461, in syn. = pulverulentum.
pallidum, Mill. Gard. Dict. ed. VIII. n. 12 = pulverulentum.
palmatum, Ledeb. Fl. Ross. ii. 552, sphalm. = parthenifolium.
paludosum, *Poir. Voy. Barb.* ii. 241.—Afr. bor.
palustre, *Pers. Syn.* ii. 461.—Armenia.
parthenifolium, Pers. l. c. 462 = præaltum.
Parthenium, *Bernh. Syst. Verz. Erf.* 145.—Europ.
parviflorum, Pers. Syn. ii. 462 = Matricaria præcox.
pectinatum, *Linn. Sp. Pl.* ed. II. 1255.—Hispan.
perpusillum, Loisel. in Desv. Journ. Bot. ii. (1809) 269 = Nananthea perpusilla.
pinnatifidum, *Linn. f. Suppl.* 377.—Ins. Madera.
præaltum, *Vent. Jard. Cels*, t. 43.—Reg. Caucas.; As. Min.; Persia.
præcox, DC. Prod. vi. 69 = Matricaria præcox.
pratense, Sallab. Prod. 203 = C. Leucanthemum.
procumbens, Lour. Fl. Cochinch. ii. 499 = indicum.
prostratum, Pers. Syn. ii. 460 = Egletis domingensis.
ptarmicæfolium, Fischk. ex Willd. Sp. Pl. iii. 2151; Pers. Syn. ii. 460 = Achillea Ptarmica.
pubescens, Hort. ex Reichb. Ic. Fl. Crit. iv. 47 = viscidum.
pulverulentum, *Lag. ex Pers. Syn.* ii. 461.—Hispan.
pumilum, *Willd. Enum. Hort. Berol.* 903.—Hab. ?
punctatum, Pers. Syn. ii. 461, in syn. = palustre.
purpureum, Pers. l. c. = indicum.
pusillum, *Vis. Fl. Lib. Spec.* 56.—Afr. bor.
radicans, *Pers. Syn.* ii. 462.—Hispan.
ramosum, *F. W. Schmidt, Sammel. Phys. Aufs.* i. (1795) 281, nomen.—Europ.
Richteria, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 315.—Sodongaria ? Tibet. occ.
rigidum, Vis. Del. Sem. Hort. Patav. (1825) = cinerariæfolium.
roseum, Adam, in Weber & Mohr, Beitr. i. 70 = coccineum.
rotundifolium, *Waldst. & Kit. Pl. Rar. Hung.* iii. 262. t. 236.—Hungar.
Roxburghii, Desf. Cat. Hort. Par. ed. III. 170 = coronarium.
scandens, *Vell. Fl. Flum.* viii. t. 129.—Bras.
segetale, Sallab. Prod. 203 = segetum.
segetum, Forsk. ex DC. Prod. vi. 64 = coronarium.
segetum, *Linn. Sp. Pl.* 889.—Europ.; Afr. bor.; As. occ.
senecioides, Dun. ex DC. Prod. vi. 64 = coronarium.
serlorum, Adam, in Weber & Mohr, Beitr. i. 69 = segetoides.
serlorum, Hoffmgg. & Link, Fl. Port. ii. t. 101 = oppositifolium.
serotinum, *Linn. Sp. Pl.* 888.—Hungar.

CHRYSANTHEMUM:—
setabense, Dufour, ex DC. Prod. vi. 49 = glabrum.
sibiricum, Turcz. ex DC. l. c. 46 = arcticum.
sinense, *Sabine, in Trans. Linn. Soc.* xiv. (1823) 145.—Japon.
sinuatum, *Ledeb. Fl. Alt.* iv. 116.—Sibir. altaic.
spathulatum, *F. W. Schmidt, Sammel. Phys. Aufs.* i. (1795) 281.—Europ.
speciosum, Brouss. ex Pers. Syn. ii. 463 = coronarium.
speciosum, Steud. Nom. ed. II. i. 357 = grandiflorum.
Stolitzkai, *C. B. Clarke, Comp. Ind.* 147.—Tibet. occ.
suaveolens, *Aschers. Fl. Brandenb.* 382.—Europ.
subcorymbosum, *Schur, in Verh. Siebenb. Ver. Naturw.* x. (1859) 146.—Transsylv.
sylvaticum, Hoffmgg. & Link, ex Link, Handb. i. 755 = C. Leucanthemum ?
sylvaticum, Jacq, ex DC. Prod. vi. 47 = pallens.
sylvestre, Brot. ex Steud. Nom. ed. I. 192; ed. II. i. 357, nomen.—Lusit.
sylvestre, Willd. Enum. Hort. Berol. Suppl. 60 = C. Leucanthemum.
tanacetifolium, Desf. Cat. Hort. Par. (1826); ex DC. Prod. vi. 66 = grandiflorum.
tanacetifolium, *Willd. Sp. Pl.* iii. 2145.—Orient.
Tanacetum, Vis. Fl. Dalmat. ii. 89 = Tanacetum vulgare.
tenuifolium, *Kit. in Schult, Oester. Fl.* ed. II. ii. 498.—Hungaria.
tenuifolium, Tenore, ex Steud. Nom. ed. II. i. 357 = C. Achilleæ.
tenuisectum, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 268.—Syria.
Thunbergii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 162.—Afr. austr.
tibeticum, *C. B. Clarke, Comp. Ind.* 147.—Tibet. occ.
tingitanum, *Bernh. in Linnæa*, x. (1835) *Litt.* 74.—Marocc.
tomentosum, *Loisel. Fl. Gall.* 580. t. 18.—Corsica.
trichophyllum, *Boiss. Diagn.* Ser. I. iv. 10.—As. Min.
trichotomum, Walp. Ann. ii. 891 = trichophyllum.
tricolor, Andr. Bot. Rep. t. 109 = carinatum.
trifurcatum, *Desf. Fl. Atlant.* ii. 281.—Afr. bor.
trspartitum, Sweet, Brit. Flow. Gard. Ser. I. t. 193 = indicum.
Turreanum, Vis. Stirp. Dalm. Spec. 19 = cinerariæfolium.
uliginosum, *Pers. Syn.* ii. 460.—Hungar.
umbellatum, *Vell. Fl. Flum.* viii. t. 127.—Bras.
umbrosum, *Willd. Sp. Pl.* iii. 2149.—Graecia.
variabile, Rochel, Pl. Banat. Rar. Ind. Spec. = atratum.
viscosum, *Desf. Cat. Hort. Par.* ed. III. 170.—Reg. Mediterr.
Webbianum, *Ball. in Journ. Linn. Soc. xvi.* (1878) 509.—Marocco.
Webbii, *Mauf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 207.—Ins. Canar.
Welwitschii, Sch. Bip. ex Nym. Consp. 370 = segetum.
Zawadskii, *Herbich, Add. Fl. Galic.* 48.—Gallicia.

CHRYSEIS, Cass. in Dict. Sc. Nat. vii. 377; ix. 154 (1817) = **Centaurea**, Linn. (Compos.).
Centaurium, Kostel. Allg. Med. Pharm. Fl. ii. 606 = Cent. Centaurium.
glauca, Cass. in Dict. Sc. Nat. xliv. 89 = Cent. moschata.
moschata, Cass. l. c. = Cent. moschata.
odorata, Cass. l. c. ix. 154 = Cent. moschata.
suaveolens, Cass. l. c. xliv. 89 = Cent. moschata.

CHRYSEIS, Lindl. Bot. Reg. t. 1948 (1835) = **Eschscholtzia**, Cham. (Papaverac.).
carnpitosa, Torr. & Gray, Fl. N. Am. i. 64 = E. californica.
californica, Hook. & Arn. Bot. Beech. Voy. 319 = E. californica.
compacta, Lindl. Bot. Reg. t. 1948 = E. californica.
crocea, Lindl. l. c. = E. californica.
Douglasii, Hook. & Arn. Bot. Beech. Voy. 320 = E. californica.
hypocrides, Torr. & Gray, Fl. N. Am. i. 64 = E. californica.
tenuifolia, Torr. & Gray, l. c. = E. californica.

CHRYSION, Spach, Hist. Vég. Phan. v. 509 (1836) = **Viola**, Linn.
Mysorum, Spach, l. c. 510 = V. biflora.

CHRYSIPHIALA, [Ker-Gawl. in] Bot. Reg. t. 778 (1824) = **Stenomesson**, Herb. (Amaryllid.).
aurantiaca, Schult. f. Syst. vii. 904 = S. aurantiacum.
coccinea, [Ker-Gawl. in] Bot. Reg. sub t. 778 = S. coccineum.
curvidentata, Herb. ex Steud. Nom. ed. II. i. 858 = S. flavum.
flava, [Ker-Gawl. in] Bot. Reg. t. 778 = S. flavum.
incarnata, Schult. f. Syst. vii. 908 = S. incarnatum.
latifolia, [Ker-Gawl. in] Bot. Reg. sub t. 778 = Urceolina latifolia.
pauciflora, Hook. Exot. Fl. t. 132 = S. flavum.
recurvata, [Ker-Gawl. in] Bot. Reg. sub t. 778 = S. recurvatum.
trichroma, Schult. f. Syst. vii. 907 = Pancratium trichromum.
variegata, Schult. f. l. c. = Pancratium variegatum.
viridiflora, Schult. f. l. c. 908 = S. viridiflorum.

CHRYSIS, Renealm. ex DC. Prod. v. 585 (1836) = **Hellanthus**, Llan. (Compos.).

CHRYSITRIX, Linn. Mant. ii. 165 (1771). *CYPERACEÆ*, Benth. & Hook. f. iii. 1057.
capensis, *Linn. Mant.* ii. 304; *Linn. f. Gram.* 27. t. 1, 2.—Afr. austr.
junciformis, *Nees, in Linnæa*, x. (1836) 144.—Afr. austr.

CHRYSOBACTRON, Hook. f. Fl. Antarct. i. 72. t. 44 (1845) = **Bulbinella**, Kunth (Liliac.).
Hookeri, Colenso, in Hook. Ic. Pl. t. 817.—N. Zel.
Rossii, Hook. f. Fl. Antarct. i. 72. t. 44–45.—Ins. Auckland.

CHRYSOBALANUS, Linn. Gen. ed. I. 865 (1737). *ROSACEÆ* Benth. & Hook. f. i. 606.
Icaco, Adans. Fam. ii. 305 (1763).
cuspidatus, *Griseb. Fl. Brit. W. Ind.* 711.— Ind. occ.
ellipticus, *Soland. ex Sabine, in Trans. Hort. Soc.* v. (1824) 458.—Afr. trop.
guianensis, Klotzsch, in Schomb. Fauna et Fl. Brit. Guian. 1024 = ellipticus.
Icaco, *Linn. Sp. Pl.* 513.—Afr. et Am. trop.
incanus, Rafin. New Fl. Am. iii. 26 = oblongifolius.
latus, Sabine, in Trans. Hort. Soc. v. (1824) 458 = C. Icaco ?
macrophyllus, *Schott, in Spreng. Syst.* iv. *App.* 407.—Bras.
oblongifolius, *Michx. Fl. Bor. Am.* i. 288.—Am. bor.
oblucularis, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 229 = C. Icaco.
owalifolius, *Schott, in Spreng. Syst.* iv. *App.* 406.—Bras.
pallocarpus, G. F. W. Mey. Prim. Fl. Esseq. 198 = C. Icaco.
prunifolius, Rafin. New Fl. Am. iii. 26 = oblongifolius.
purpureus, Mill. Gard. Dict. ed. VIII. n. 2 = C. Icaco.
racemosus, *Rozb. Hort. Beng.* [92]; *Fl. Ind.* ii. 506.—Ins. Moluce.
retusus, Rafin. New Fl. Am. iii. 26 = oblongifolius.
subundulatus, *Brign. in Mem. Soc. Ital. Mod.* Ser. II. i. (1862) 71.—Bras.

CHRYSOBAPHUS, Wall. Tent. Fl. Nep. 37. t. 27 (1826) = **Anoectochilus**, Blume (Orchid.).
Roxburghii, Wall. l. c. = A. Roxburghii.

CHRYSOBOTRYA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 18 = **Ribes**, Linn. (Saxifrag.).
intermedia, Spach, l. c. 19 = R. aureum.
Lindleyana, Spach, l. c. 20 = R. aureum.
revoluta, Spach, l. c. 19 = R. aureum.

CHRYSOCALYX, Guill. & Perr. Fl. Senegj. Tent. 157. t. 43 (1833) = **Crotalaria**, Linn. (Legumin.).
ebenoides, Guill. & Perr. l. c. 158. t. 43. f. 1 = Crot. ebenoides.
gracilis, Guill. & Perr. l. c. 159 = Crot. gracilis.
Perrottetii, Guill. & Perr. l. c. 160. t. 43. f. 2 = Crot. Perrottetii.
Petitiana, A. Rich. Tent. Fl. Abyss. i. 169 = Crot. macrocalyx.

CHRYSOCALYX :—
platycalyx, Steud. ex A. Rich. Tent. Fl. Abyss. 160 = Crotalaria platycalyx.
Quartinianus, A. Rich. l. c. 150 = Crot. Dillenianæ.
rubiginosus, Guill. & Perr. Fl. Seneg. Tent. 158 = Crot. strorubens.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 151 = Crot. incana.

CHRYSOCEPHALUM, Walp. in Linnæa, xiv. (1840) 508 = Hallchrysum, Vaill. (Compos.).
apiculatum, Steetz, in Lehm. Pl. Preiss. i. 474 = H. apiculatum.
asperum, Steetz, l. c. 473 = H. semipapposum.
Baumannum, Sond. in Linnæa, xxv. (1852) 517 = H. Baxteri.
brevicilium, Walp. Rep. vi. 238 = H. semipapposum.
conferuum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) t. 196 = H. apiculatum.
ciliatum, Steetz, in Lehm. Pl. Preiss. i. 474 = H. semipapposum.
Eminium, Walp. Rep. i. 327, sphalm. = seq.
Eminium, Steetz, in Lehm. Pl. Preiss. i. 472 = H. apiculatum.
glabratum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) t. 197 = H. apiculatum.
helichrysoides, Walp. in Linnæa, xiv. (1840) 508 = H. apiculatum.
odorum, Walp. Rep. vi. 238 = H. apiculatum.
pterochætum, F. Muell. in Linnæa, xxv. (1852) 416 = H. pterochætum.
semipapposum, Steetz, in Lehm. Pl. Preiss. i. 474 = H. semipapposum.
squarrulosum, Sond. in Linnæa, xxi. (1852) 515 = H. semipapposum.
squarrulosum, Steetz, in Lehm. Pl. Preiss. i. 472 = H. apiculatum.
viteliinum, Sond. & F. Muell. in Linnæa, xxv. (1852) 514 = H. apiculatum.

CHRYSOCHAMELA, Boiss. Fl. Orient. i. 813 (1867) = Cochlearia, Tourn. (Cruciferæ).
alliifica, Boiss. l. c. 814.—As Min.
Nodana, Boiss. l. c.—Armenia.
veinina, Boiss. l. c. 813 = Coch. veinina.

CHRYSOCHLAMYS, Poepp. & Endl. Nov. Gen. et Sp. iii. 13. t. 211 (1842). *GUTTIFERÆ*, Benth. & Hook. f. 172.
BERTOLONIA, Spreng. Neue Entdeck. ii. 110. t. 1. f. 1 (1821).
COMMINGSIA, Miers, in Trans. Linn. Soc. xxi. (1855) 252. t. 26.
PORCILOSTEMON, Planch. & Triana, in Ann. Sci. Nat. Sér. IV. xiv. (1860) 266.
TOVOMITOPSIS, Planch. & Triana, l. c. 361 (1860).
cinnicodes, *Griseb. Cat. Pl. Cub.* 37.—Cuba.
costa-ricana, *Hemsl. Biol. Centr. Am. Bot.* i. 37.—Costa Rica.
dependens, *Planch. & Triana*, *in Ann. Sc. Nat.* Ser. IV. xiv. (1860) 259.—Am. trop.
grisea, *Hemsl. Biol. Centr. Am. Bot.* i. 37.—Costa Rica.
Goudotii, *Planch. & Triana*, *in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 259.—N. Granat.
laxa, *Planch. & Triana*, *l. c.* 858.—N. Granat.
membranacea, *Planch. & Triana*, *l. c.* 260.—N. Granat.
multiflora, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 13. t. 211.—Peruv.
myriadea, *Planch. & Triana*, *in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 260.—N. Granat.
nicaraguensis, *Hemsl. Biol. Centr. Am. Bot.* i. 37.—Nicaragua.
pachypoda, *Planch. & Triana*, *in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 257.—Peruv.
Parenti, *Planch. & Triana*, *l. c.* 258.—Peruv.
psychotrisefolia, *Hemsl. Biol. Centr. Am. Bot.* i. 37.—Costa Rica.

CHRYSOCOMA, Linn. Gen. ed. I. 948 (1787), partim. *COMPOSITÆ*, Benth. & Hook. f. ii. 396.
acaulis, Walt. Fl. Carol. 196 = Vernonia oligophylla.
adpressa, Vell. Fl. Flum. 338; viii. t. 35 = Vernonia mosticola.
Altoni, *Spreng. Fl. Hal. Mant.* 50.—Hab.?
alata, *Vell. Fl. Flum.* 335; viii. t. 48 (Eupatorii sp.?).—Bras.
albida, *Vell. l. c.* 338; viii. t. 49.—Bras.
CHRYSOCOMA :—
alterna, *Vell. Fl. Flum.* 338; viii. t. 40 (Vernoniæ sp.?).—Bras.
amara, Schum. & Thonn. Beskr. Guin. Pl. 353 = Vernonia senegalensis.
angustifolia, Steud. Nom. ed. II. i. 359 = Vernonia angustifolia.
aphylla, Vell. Fl. Flum. 324; viii. t. 1 = Baccharis aphylla.
arborea, Vell. l. c. 326; viii. t. 11 = Vernonia Tweediana.
artemisiæfolia, Poepp. ex Steud. Nom. ed. II. i. 356 = Eupatorium coronopifolium.
aspera, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 338 = Inula germanica.
aurea, Boj. ex DC. Prod. vi. 322 = Nidorella Chrysocoma.
aurea, Salisb. Prod. 118 = C. Coma-aureæ.
bifora, Linn. Sp. Pl. 841 = Aster dracunculoides.
camphorata, Rob. & Cass. in Dict. Sc. Nat. xxxiv. 35 = Jasonia glutinosa.
candicans, Delile, Fl. Egypte, 271. t. 46 (= *Varthemia candicans*).—Aegypt.
capillacea, Michx. Fl. Bor. Am. ii. 101 = Eupatorium foeniculaceum.
cernua, Linn. Sp. Pl. 840 = C. Coma-aureæ.
cernua, Thunb. Prod. Fl. Cap. 149, partim = tenuifolia.
ciliaris, Link, Enum. Hort. Berol. ii. 311 = Aster Ecklonis.
ciliaris, Willd. Sp. Pl. iii. 1789 = ciliata.
ciliata, *Linn. Sp. Pl.* 841.—Afr. austr.
ciliata, E. Mey. ex DC. Prod. v. 254 = oblongifolia.
cinerea, Labill. Nov. Holl. Pl. ii. 89. t. 193 = Helichrysum cinereum.
Coma-aurea, *Linn. Sp. Pl.* 840.—Afr. austr.
coronopifolia, Michx. Fl. Bor. Am. ii. 102 = Eupatorium coronopifolium.
crassa, *Vell. Fl. Flum.* 338; viii. t. 8.—Bras.
crenata, *Vell. l. c.* 326; viii. t. 10 (Baccharidis sp.?).—Bras.
cuneifolia, *Jacq. Coll.* iii. 280.—Reg. Magell.
cymosa, Vell. Fl. Flum. 337; viii. t. 16 = Vernonia missionia.
decurrens, *DC. Prod.* v. 254.—Afr. austr.
decurrens, Harv. in Harv. & Sond. Fl. Cap. iii. 95 = Hertzenna decurrens.
decussata, Vell. Fl. Flum. 326; viii. t. 9 = Baccharis camporum.
dentata, Vell. l. c. 324; viii. t. 47 = Baccharis oxyphylla.
denticulata, Jacq. Hort. Schoenb. iii. 62 = Chrysanthemum floseulosum.
dichotoma, Jacq. Ic. Pl. Rar. i. 17. t. 171 = Allagopappus dichotoma.
discaricata, Fisch. Hort. Gorenk. ed II. (1812) 87 = Aster divaricatus.
dracunculoides, Lam. Encyc. ii. 192 = Aster dracunculoides.
dracunculoides, Pursh, Fl. Am. Sept. ii. 517 = Bigelowia dracunculoides.
ferruginea, Spreng. Syst. iii. 494 = Helichrysum ferrugineum.
floribda, *Vell. Fl. Flum.* 334; viii. t. 44.—Bras.
foetida, Lam. Encyc. ii. 198 = Nidorella foliosa.
fruticosa, J. F. Gmel. Syst. 1908 = cuneifolia.
fruticosa, Stend. Nom. ed. II. i. 358 = Vernonia rigida.
glauca, Walt. Fl. Carol. 196 = Vernonia altissima.
graphaloides, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
graminifolia, Bosher, in Pall. N. Nord. Beitr. vi. 261 = Aster Linosyris.
graminifolia, Linn. Sp. Pl. 841 = Solidago lanceolata.
graminifolia, Walt. Fl. Carol. 196 = Vernonia angustifolia.
grandis, *Lepech. It.* iii. 86.—Sibir. ural.
graveolens, Nutt. Gen. Am. ii. 136 = Bigelowia dracunculoides.
graveolens, Torr. in Ann. Lyc. N. York, ii. (1828) 211 = Bigelowia pluriflora.
herbacea, Vell. Fl. Flum. 330; viii. t. 29 = Vernonia oborata.
heterophylla, Raffn. Fl. Ludov. 64 (= Baccharidis sp. ?).—Am. bor.
hirsuta, *Vell. Fl. Flum.* 333; viii. t. 41.—Bras.
horizontalis, Vell. Fl. Flum. 330; viii. t. 26 = Vernonia obscura.
humilis, DC. Prod. vii. 260 = Aster empetrus.
imbarbis, *Vell. Fl. Flum.* 331; viii. t. 34.—Bras.
incana, Burm. f. Fl. Cap. Prod. 28 = Pteronia incana.
CHRYSOCOMA :—
incana, Léveillé in Denkl. Voy. Russ. Mér. ii. 237; iv. 496 (= *Linosyris villosa*).—Russia.
indivisa, Vell. Fl. Flum. 326; viii. t. 12 = Baccharis brachylaenoides.
integerrima, *Vell. l. c.* 328; viii. t. 19.—Bras.
junceus, *Desf. Tabl. ed.* II. 115, nomen.—Hab.?
lappulina, Boj. ex DC. Prod. v. 252 = Nidorella muricronata.
lateralis, Vell. Fl. Flum. 329; viii. t. 21 = Baccharis caniniefolia.
litoralis, Spreng. Syst. iii. 495 = Aster Linosyris.
linearis, *Vell. Fl. Flum.* 330; viii. t. 27.—Bras.
linifolia, Steud. Nom. ed. II. i. 355 = Aster subulatus?
Linosyris, Linn. Sp. Pl. 841 = Aster Linosyris.
longifolia, *DC. Prod.* v. 254.—Afr. austr.
maculata, Vell. Fl. Flum. 325; viii. t. 6 = Eupatorium conyzoides.
maritima, Vell. l. c. 329; viii. t. 22 = Baccharis argyrea.
mediocerrata, *Vell. l. c.* 327; viii. t. 15.—Bras.
megapotamica, Spreng. Syst. iii. 424 = Eupatorium ericoides.
microcephala, DC. Prod. v. 258 = tenuifolia.
microphylla, Thunb. Prod. Fl. Cap. 142 = tenuifolia.
monantha, *Vell. Fl. Flum.* 331; viii. t. 33.—Bras.
montana, Vahl. Symb. Bot. i. 70 (= *Varthemia montana*).—Arabia.
mucronata, Forsk. Fl. Aegypt.-Arab. 147 = Iphiona juniperifolia.
nauseosa, Pursh, Fl. Am. Sept. ii. 517 = Bigelowia dracunculoides.
nives, *Willd. Sp. Pl.* iii. 1790.—Afr. austr.
nouveboracensis, Desf. Tabl. ed. I. 98 = Vernonia noveboracensis.
nuda, Vell. Fl. Flum. 335; viii. t. 50 = Baccharis aphylla.
nudata, Michx. Fl. Bor. Am. ii. 101 = Bigelowia nudata.
nupera, S. F. Gray, Nat. Arr. Brit. Pl. ii. 467 = Aster Linosyris.
oblongifolia, *DC. Prod.* 254.—Afr. austr.
obscura, *Vell. Fl. Flum.* 330; viii. t. 30.—Bras.
odoratissima, Rasusch. Nom. ed. III. 285 = Trilisia odoratissima.
oligophylla, Vell. Fl. Flum. 324; viii. t. 2. = Vernonia Cephalotes.
opposita, *Vell. l. c.* 334; viii. t. 46.—Bras.
oppositifolia, Linn. Pl. Rar. Afr. 18 = Pteronia divaricata.
oppositifolia, *Vell. Fl. Flum.* 338; viii. t. 18.—Bras.
ovata, *Forsk. Fl. Aegypt. Arab.* 147.—Arabia.
palustris, Savi, Pugill. 13; ex Bertol. Fl. Ital. ix. 91 = Aster Linosyris.
paniculata, J. F. Gmel. Syst. 1204 = Trilisia paniculata.
paniculata, Vell. Fl. Flum. 327; viii. t. 14 = Vernonia geminata.
patula, Berg. Desc. Pl. Cap. 234 = C. Coma-aureæ.
pauciflora, Vell. Fl. Flum. 325; viii. t. 5 = Aganetum malleasefolium.
peduncularis, *DC. Prod.* v. 253.—Afr. austr.
pedunculata, Vell. Fl. Flum. 332; viii. t. 37 = Vernonia sanmonophila.
phosphorica, Vell. l. c. 325; viii. t. 4 = Vernonia polyanthus.
pilosa, J. F. Gmel. Syst. 1908 = Llatris pycnostachya.
pilosa, *Vell. l. c.* 338; viii. t. 17.—Bras.
pinifolia, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
pinifolia, Spreng. Syst. iii. 494 (= *Cacchamnus pinifolius*).—N. Zel.
pinnatifida, *DC. Prod.* v. 254.—Afr. austr.
prenanoe, Steud. Nom. ed. II. i. 358 = C. Proscna.
prenanthe, Desf. Tabl. ed. I. 98 = Vernonia altissima!
Prenanes, *Vell. Fl. Flum.* 333; viii. t. 39.—Bras.
pteroconia, *DC. Prod.* v. 254.—Afr. austr.
punctata, Vell. Fl. Flum. 331; viii. t. 32 = Vernonia gracilifora.
punctata, Steud. Nom. ed. II. i. 356 = seq.
punctulata, Vell. Fl. Flum. 335; viii. t. 49 = Eupatorium laevigatum.
purpurascens, *Vell. l. c.* 331; viii. t. 31.—Bras.
purpurea, *Forsk. f. Prod.* 54.—N. Hebrid.
purpurea, Raffn. Fl. Ludov. 64 (= Llatridis sp.).—Am. bor.
purpurea, Vell. l. c. 324; viii. t. 45 = Baccharis caniniefolia.
quadrangularis, Vell. Fl. Flum. 329; viii. t. 23 = Piptocarpha quadrangularis.

CHRYSOCOMA:—
radiata, *Flach. ex Steud. Nom.* ed. II. i. 359.—Hab.?
ramosa, *Rœusch. Nom.* ed. III. 285, nomen.—Carolin.
repanda, Vell. Fl. Flum. 237; viii. t. 18 = Vernonia scorpioides.
reticulata, Labill. Nov. Holl. Pl. ii. 40. t. 183 = Helichrysum reticulatum.
rosmarinifolia, Spreng. Syst. iii. 424 = Helichrysum rosmarinifolium.
sancta, Vell. Fl. Flum. 234; viii. t. 3 = Baccharis Lundii.
saxatilis, DC. Fl. Fr. Suppl. 468 = Jasonia saxatilis.
scabra, Ledeb. in Mém. Acad. Pétersb. v. (1812) 563 = Inula germanica.
scabra, Linn. Sp. Pl. ed. II. 1177 = Felicia rehexa.
scoparia, Linn. Fl. Jam. Pug. 21; Amœn. Acad. v. 381 = Baccharis scoparia.
sericea, Linn. f. Suppl. 360 (= *Schistogyne sericea*).—Ins. Canar.
serrata, Hill, Veg. Syst. iii. 128, sphalm. = C. Comaaurea.
serrata, *Vell. Fl. Flum.* 229; viii. t. 23.—Bras.
sessilis, Vell. l. c. 238; viii. t. 20 = Vernonia macrophylla.
sieberiana, Vell. l. c. 225; viii. t. 7 = Vernonia nitidula et Baccharis ceanothifolia.
spathulata, Forsk. Fl. Aegypt. Arab. 147 = Vernonia atriplicifolia.
sphærotata, Mœnch, Meth. 577 = Senecio paniculatus.
spicata, Forsk. Fl. Aegypt. Arab. p. lxxiii = Iduga Fontanesii.
spinosa, Delile, Fl. Aegypt. 128. t. 46 = Iphiona juniperifolia.
squamata, Labill. Nov. Holl. Pl. ii. 40. t. 184 = Leptorhynchus squamatus.
tatarica, *Less. in Linnæa*, ix. (1834) 186.—Reg. Casp.
tenuifolia, *Berg. Descr. Pl. Cap.* 235.—Afr. austr.
tenuifolia, Salisb. Prod. 188 = Aster Linosyris.
tomentosa, *Vell. Fl. Flum.* 233; viii. t. 58.—Bras.
tomentosa, *Linn. Syst.* ed. XII. 539.—Afr. austr.
tomentosa, Walt. Fl. Carol. 196 = Vernonia noveboracensis.
tridentata, *DC. Prod.* v. 264.—Afr. austr.
trinervis, *Vell. Fl. Flum.* 233; viii. t. 43.—Bras.
truncifolia, J. Shaw, in Journ. Linn. Soc. xiv. (1874) 206, sphalm. = tenuifolia.
tuberosa, *Vell. Fl. Flum.* 232; viii. t. 36.—Bras.
undulata, Thunb. Prod. Pl. Cap. 142 = Nidorella amplexicaulis.
uniflora, *Rœusch. Nom.* ed. III. 285, nomen.—Am. bor.
uniflora, Spreng. Nov. Prov. (1818) 11 (= *Varthemia candicans*).—Aegypt.
verticalis, Lag. Gen. et Sp. Nov. 25 = Jasonia saxatilis.
verticillata, Vell. Fl. Flum. 230; viii. t. 26 = Piptocarpha oxyphylla.
villosa, Linn. Sp. Pl. 841 (= *Linosyris villosa*).—Europ.; Casc.; Sibir.
violacea, Schum. & Thonn. Beskr. Guin. Pl. 334 = Vernonia cinerea.
virgata, Nutt. Gen. Am. ii. 137 = Bigelowia nudata.
virginiana, Hort. ex DC. Prod. v. 341 = Solidago lanceolata.
volubilis, *Vell. Fl. Flum.* 229; viii. t. 24.—Bras.
vulgaris, *Gueldenst. ex Ledeb. Fl. Ross.* ii. 495 = Aster Linosyris.

CHRYSOCOPTIS, Nutt. in Journ. Acad. Philadr vii. (1834) 8. t. 1 = Coptis, Salisb. (Ranunculac.).
asplenifolia, Nutt. l. c. 9, in obs. = Coptis asplenifolia.
occidentalis, Nutt. l. c. 8 = Coptis occidentalis.

CHRYSOCORYNE, Endl. in Bot. Zeit. i. (1843) 487 = Angianthus, Wendl. (Compos.).
anglomicoides, F. Muell. in Linnæa, xxv. (1852) 404, 483 = A. pusillus.
Drummondii, A. Gray, in Hook. Kew Journ. iii. (1851) 153 = A. tenuifus.
Hookerii, A. Gray, l. c. 151 = A. tenuifus.
myosuroides, A. Gray, l. c. 153 = A. myosuroides.
pusilla, Endl. in Bot. Zeit. i. (1843) 488 = A. pusillus.
tenuifus, F. Muell. in Trans. Vict. Inst. (1855) 126 = A. tenuifus.
uniflora, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 188 = A. myosuroides.

CHRYSOCYATHUS, Falc. in Proc. Linn. Soc. i. (1839) 17 = Calathodes, Hook. f. & Thoms. (Ranunculac.).

CHRYSOCYCNIS, Linden & Reichb. f. in Bonplandia, ii. (1854) 280. *ORCHIDEÆ*, Benth. & Hook. f. iii. 583.
Schlimii, *Linden & Reichb. f. l. c.*—N. Granat.

CHRYSODENDRON, Vaill. ex Meisn. in DC. Prod. xiv. 230 (1856-57) = Protea, Linn.

CHRYSODISCUS, Steetz, in Lehm. Pl. Preiss. i. 460 (1845) = Athrixia, Ker-Gawl. (Compos.).
sieversii, Steetz, l. c. = A. stricta.

CHRYSOGLOSSUM, Blume, Bijdr. 887. t. 7 (1825). *ORCHIDEÆ*, Benth. & Hook. f. iii. 507.
DIGLYPHIS, Blume, Fl. Jav. Fragm. p. vii. (1828).
DILYPHOSA, Blume, Bijdr. 336. t. 60 (1825).
latifolium, *Benth. & Hook. f. Gen.* iii. 508.—Java.
ornatum, *Blume, Bijdr.* 336.—Java.
vesicatum, *Reichb. f. in Seem. Fl. Vit.* 304.—Ins. Fiji.
villosum, *Blume, Bijdr.* 338.—Java.

CHRYSOGONUM, Linn. Hort. Cliff. 424 (1737); Gron. 17 (1737). *COMPOSITÆ*, Benth. & Hook. f. ii. 350.
CARGILLA, Adans. Fam. ii. 130 (1763).
DIOTOSTEPHUS, Cass. in Dict. Sc. Nat. xlviii. 548 (1827).
MOONIA, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 848.
PENTALEPIS, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 496.
Arnottianum, *C. B. Clarke, Compos. Ind.* 132.—Ind. or.
dichotomum, *Vahl, in West, Bidr. Ste. Croix*, 303; *DC. Prod.* v. 510.—Ins. S. Crucis.
Diotostephus, DC. l. c. = virginianum.
elliptoides, *Benth. & Hook. f. ex F. Muell. Census*, 88.—Austral.
heterophyllum, *C. B. Clarke, Comp. Ind.* 132.—Ind. or.
pervarianum, Linn. Sp. Pl. 920 = Zinnia multiflora.
procumbens, *Benth. & Hook. f. ex F. Muell. Census*, 88.—Austral.
trichomeroides, *Benth. & Hook. f. l. c.*—Austral.
virginianum, *Linn. Sp. Pl.* 920.—Am. bor.

CHRYSOLIGA, Willd. ex DC. Prod. iii. 89, in syn. (1828) = Mœnes, Comm. (Lythrar.).

CHRYSOLINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349 = Linum, Tourn.
gallicum, Fourr. l. c. = L. gallicum.
strictum, Fourr. l. c. = L. strictum.

CHRYSOLVGA, Willd. ex Steud. Nom. ed. II. i. 359 (1840) = Mœnes, Comm. (Lythrar.).
salicifolia, Willd. ex Steud. l. c. = N. salicifolia.

CHRYSOMA, Nutt. in Journ. Acad. Philad. vii. (1834) 67 = Solidago, Vaill. (Compos.).
pauciflora, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 395 = S. pauciflosculosa.
solidagineides, Nutt. in Journ. Acad. Philad. vii. (1834) 67 = S. pauciflosculosa.
ustilaginoides, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 395 = S. neglecta.

CHRYSOMALLUM, Thou. Gen. Nov. Madag. 8 (1806) = Vitex, Linn. (Verbenac.).
madagascariense, Thou. ex Steud. Nom. ed. I. 194 = V. Chrysomallum.

CHRYSOMELEA, Tausch, Hort. Canal. fasc. i.—Cf. DC. Prod. v. 569 (1836) = Corrosperse, Linn. (Compos.).
auriculata, Tausch, l. c.; ex DC. l. c. 571 = Cor. pubescens.
lanceolata, Tausch, l. c.; ex DC. l. c. 570 = Cor. lanceolata.

CHRYSOMELON, Forst. ex A. Gray, Bot. U. St. Expl. Exped. i. 374, in syn. (1854) = Spondias, Linn. (Anacard.).
pomiferum, Forst, ex A. Gray, l. c. = S. dulcis.

CHRYSONIAS, Benth. ex Steud. Nom. ed. II. i. 359 (1840) err. typ. = Chrysocoma, E. Mey.

CHRYSOPHANIA, Kunth, ex Less. Syn. Comp. 224 (1832) = Baltimoria, Pers. (Compos.).
fastigiata, Kunth, l. c. = *Z. angusta*.

CHRYSOPHORA, Cham. ex Triana, in Trans. Linn. Soc. xxvii. (1871) 94, 95. 147 = Oxymeris, DC. (Melastomac.).
cinnamomifolia, Cham. ex Triana, l. c. 147 = Ossæa cinnamomifolia.
cordigera, Cham. ex Triana, l. c. 94 = Oxym. cordigera.
hirsutior, Cham. ex Triana, l. c. 95 = Oxym. deflexa.
secundifolia, Cham. ex Triana, l. c. = Oxym. secundiflora.

CHRYSOPHTHALMUM, Phil. in Linnæa, xxix. (1857) 9 = Grindelia, Willd. (Compos.).
andinum, Phil. l. c. = G. andina.

CHYSOPHTHALMUM, Sch. Bip. in Walp. Rep. ii. 955 (1843) (Chrysophtalmum). *COMPOSITÆ*. Benth. & Hook. f. ii. 341.
dichotomum, *Boiss. & Heldr. Diagn.* Sér. I. xi. 5.—As. Min.
montanum, Boiss. l. c. 4 = sternutatorium.
sternutatorium, *Sch. Bip. in Walp. Rep.* ii. 955.—As. Min.

CHRYSOPHYLLUM, Linn. Gen. ed. I. 361 (1737). *SAPOTACEÆ*, Benth. & Hook. f. ii. 668.
CAINITO, Plum. ex Adans. Fam. ii. 166 (1763); Tuss. Fl. Antill. iii. 41. t. 9 (1824).
DACTIMALA, Rafin. Sylva Tellur. 158 (1838).
GUERRENTIA, Rafin. l. c. (1838).
NIEMEYERA, F. Muell. Fragm. vii. 114 (1869-71).
NYCTERISITION, Ruiz & Pav. Prod. 80. t. 5 (1794).
SCLEROXYLON, Bertol. in Mem. Acc. Sc. Bologn. viii. (1857) 237 (Misc. Bot. xiii. 15).
acuminatum, Boj. Hort. Maurit. 197 = Sideroxylon ferrugineum.
acuminatum, Lam. Tabl. Encyc. ii. 44 = monopyrenum.
acuminatum, Poir. Encyc. Suppl. ii. 15 = Sideroxylon guyanense.
acuminatum, Roxb. Hort. Beng. 17; Fl. Ind. i. 599 = Roxburghii.
acuminatum, Willd. ex Roem. & Schult. Syst. iv. 818 = Humboldtianum.
africanum, *A. DC. in DC. Prod.* viii. 168.—Afr. trop.
albidum, *G. Don, Gen. Syst.* iv. 93.—Afr. trop.
alnifolium, *Baker, in Oliver, Fl. Trop. Afr.* iii. 499.—Afr. trop.
amplifolium, *A. DC. in DC. Prod.* viii. 159.—Bras.
angustifolium, *Lam. Tabl. Encyc.* ii. 44.—Ins. S. Doming.
aquaticum, H. B. & K. Nov. Gen. et Sp. iii. 286 = C. Cainito.
argenteum, *Jacq. Enum. Pl. Carib.* 15.—Ins. Martinic.
auratum, *Miq. in Mart. Fl. Bras.* vii. 97.—Bras.
bancanum, *Miq. Fl. Ind. Bat. Suppl.* 579.—Ins. Banca.
Barbasco, Loefl. ex DC. Prod. viii. 149 = Jacquinia armillaris.
bicolor, *Poir. Encyc. Suppl.* ii. 15.—Ins. Porto Rico.
Bonplandii, Klotzsch, ex Miq. in Mart. Fl. Bras. vii 94 = C. Cainito.
brasiliense, *A. DC. in DC. Prod.* viii. 156.—Bras.
bumelioides, *Mart. in Flora*, xxii. (1839) 1. *Beibl.* 1.—Bras.
Burenhem, Riedel, in Tausay, Man. Agric. Bras. ed. II. 318 = Lucuma glycyphloes.
caeruleum, Jacq. Stirp. Select. Am. 59 = C. Cainito.
Cainito, *Linn. Sp. Pl.* 198.—Ind. occ.
carolinense, Jacq. Obs. Bot. iii. 3 = Bumelia tenax.
cayennense, *A. DC. in DC. Prod.* viii. 160.—Guiana.
cuneifolium, *A. DC. l. c.*—Guiana.
decandrum, *Montr. in Mém. Acad. Lyon*, x. (1860) 228.—N. Caled.
dubium, *Steud. Nom.* ed. II. i. 359.—Ins. S. Doming.
dubium, *Pauch. & Sebert, Not. Bois. Nouv. Calédd.* 195.—N. Caled.
ebenaceum, *Mart. in Flora*, xxi. (1838) ii. *Beibl.* 96.—Bras.
emarginatum, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 1087.—Guiana.
ferrugineum, Geertn. Fruct. iii. 128 = monopyrenum.
ferrugineum, *Steud. Nom.* ed. II. i. 359.—Peruv.
firmum, *Miq. Fl. Ind. Bat. Suppl.* 579.—Ins. Banca.
flexuosum, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 98.—Bras.
fuscum, *Flach. ex Steud. Nom.* ed. II. i. 359.—Ind. or.

CHRYSOPHYLLUM:—
Gardneri, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 105.—Bras.
glabrum, Vaup. *Resum. Fl. Carib.* 15.—Ins. Martinic.
glabrum, Jusl. ex Lam. Encyc. i. 245 = Bumelia tenax.
glycyphloeum, Casar. in Atti Sci. Ital. Sc. Ital. Firenze (1841) 514; Dec. Stirp. Bras. 13 = Lucuma glycyphloea.
granatense, *Spreng. Syst.* i. 867.—N. Granat.
grandifolium, *Steud. Nom.* ed. II. i. 859.—Jos. Philipp.
guyanense, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 98 = sericeum.
Guillemiinianum, A. DC. in DC. Prod. vii. 161 = ebenaceum.
Hostmannianum, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 98 = sericeum.
Humboldtianum, *Roem. & Schult. Syst.* iv. 613.—Am. austr.
Inophyllum, Mart. ex Miq. in Mart. Fl. Bras. vii. 105 = Lucuma Inophyllum.
inornatum, *Mart. in Flora*, xxi. (1838) ii. *Beibl.* 96.—Bras.
jamaicense, Jacq. Stirp. Select. Am. 53 = C. Cainito.
jamaicense, *Eichl. in Kjoeb. Vidensh. Meddel.* (1870) 206.—Bras.
javanicum, *Steud. Nom.* ed. II. i. 859.—Java.
lanceolatum, Casar. Stirp. Bras. Dec. 64 = incratum.
lanceolatum, *A. DC. in DC. Prod.* viii. 163.—Java.
leptocarpum, Spruce, ex Miq. in Mart. Fl. Bras. vii. 106 = Lucuma leptocarpa.
longifolium, *Mart. & Eichl. ex Miq. l. c.* 97.—Bras.
lucumifolium, *Griseb. in Goett. Abh.* xxiv. (1879) 223.—Reg. Argent.
ludovicianum, *Kuhn. Fl. Ludov.* 58.—Louisiana.
Maconcon, *Aubl. Pl. Gui.* i. 293. t. 92.—Guiana.
macrophyllum, Desf. Cat. Hort. Par. ed. III. 132 = grandifolium.
macrophyllum, Gaertn. f. Fruct. iii. 122. t. 202. f. 1 = dubium.
macrophyllum, Lam. Tabl. Encyc. ii. 44 = Lucuma Rivicoa.
macrophyllum, Mart. in Flora, xxi. (1838) ii. Beibl. 95 = brasiliense.
macrophyllum, Sabine, in Trans. Hort. Soc. v. (1824) 458 = africanum.
magalis-montana, *Sond. in Linnaea*, xxiii. (1850) 72.—Afr. austr.
maliforme, Linn. Syst. ed. X. 987 = C. Cainito.
Mangilila, Lam. Tabl. Encyc. ii. 45 = Myrsine Mangilila.
Martianum, A. DC. in DC. Prod. viii. 161 = ebenaceum.
maximoides, *Mart. in Flora*, xxii. (1839) i. *Beibl.* 1.—Bras.
melanophloeum, Lam. Tabl. Encyc. ii. 45 = Myrsine mitis.
Michino, *H. B. & K. Nov. Gen. et Sp.* iii. 286.—N. Granat.
microcarpum, *Sw. Prod. Veg. Ind. Occ.* 49.—Ins. S. Doming.
microphyllum, A. DC. in DC. Prod. viii. 158 = monopyrenum.
microphyllum, Jacq. Stirp. Select. Am. 53. t. 37. f. 2 = C. Cainito.
Millerianum, Lam. Tabl. Encyc. ii. 45 = Myrsine mitis.
molle, Wall. Cat. n. 4161 = Payena Maingayi.
monopyrenum, Spreng. Syst. i. 866 = C. Cainito.
monopyrenum, *Sw. Prod. Veg. Ind. Occ.* 49.—Ind. occ.
*myrsinoides*enron, F. Muell. Fragm. vi. 178 = Sideroxylon Brownii.
myrtifolium, Mart. in Flora, xxii. (1839) i. Beibl. 1 = maytenoides.
natalense, *Sond. in Linnaea*, xxiii. (1850) 73.—Afr. austr.
nitidissimum, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 98 = sericeum.
nitidum, *G. F. W. Mey. Prim. Fl. Essq.* 116.—Guiana.
obovatum, *Sabine, in Trans. Hort. Soc.* v. (1824) 458.—Afr. trop.
obovatum, Wall. ex A. DC. in DC. Prod. viii. 178 = Sideroxylon ferrugineum.
obtusatum, Willd. ex Roem. & Schult. Syst. iv. 613 = C. Cainito.
oleaefolium, *Spruce, ex Miq. in Mart. Fl. Bras.* vii. 101.—Bras.
Olfersii, Klotzsch, ex Miq. l. c. 91 = splendens.
oliviformis, Lam. Encyc. i. 852 = monopyrenum.
oliviformis, Lian. Syst. ed. X. 987 = C. Cainito.

CHRYSOPHYLLUM:—
Ottonis, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 94 = C. Cainito.
parviflorum, *Casar. Stirp. Bras. Dec.* 63.—Bras.
paucifloram, Lam. Tabl. Encyc. ii. 44 = glabrum.
pentagonum, *Hance, in Journ. Bot.* xx. (1882) 78.—China.
Perdicastrum, *Eichl. in Kjoeb. Vidensh. Meddel.* (1870) 208.—Bras.
pentiformi, *Bert. ex Spreng. Syst.* i. 867.—N. Granat.
Prieurei, *A. DC. in DC. Prod.* viii. 161.—Guiana.
pruniferum, *F. Muell. Fragm.* vi. 26.—Austral.
prunifolium, *Baker, in Oliver, Fl. Trop. Afr.* iii. 499.—Afr. trop.
paliophyllum, *A. DC. in DC. Prod.* viii. 160.—Guiana.
pulcherrimum, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 92.—Bras.
pyriforme, Willd. Sp. Pl. i. 1084 = C. Maconcon.
ramiflorum, A. DC. in DC. Prod. viii. 158 = Ecclinusa ramiflora.
revolutum, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 104.—Bras.
rhodanense, *Hassk. in Flora*, xxxviii. (1855) 579.—Java.
Richardii, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 97 = auratum.
Roxburghii, *G. Don, Gen. Syst.* iv. 83.—As. trop.
rufum, *Mart. in Flora*, xxi. (1838) ii. *Beibl.* 94.—Bras.
rugosum, Sw. Prod. Veg. Ind. Occ. 49 = Sideroxylon rugosum.
Schomburgkianum, *A. DC. in DC. Prod.* viii. 157.—Guiana.
Sebertii, *Panch. in Panch. & Sebert, Not. Bois Nouv. Caléd.* 194.—N. Caled.
sericeum, *A. DC. in DC. Prod.* viii. 158.—Guiana.
sericeum, Salisb. Prod. 138 = C. Cainito.
sessiliflorum, *Poir. Encyc. Suppl.* ii. 16.—Guiana.
sessiliflolium, *Panch. & Sebert, Not. Bois Nouw. Caled.* 193.—N. Caled.
sparsiflorum, *Klotzsch, ex Miq. in Mart. Fl. Bras.* vii. 90.—Guiana.
splendens, *Spreng. Syst.* i. 866.—Bras.; Peruv.
subrotundum, *Baker, in Oliver, Fl. Trop. Afr.* iii. 499.—Afr. trop.
sumatranum, Miq. Fl. Ind. Bat. Suppl. 579 = Roxburghii.
sundaicum, *Miq. l. c.* 578.—Ins. Sunda.
viride, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 102.—Bras.
Wakere, *Panch. & Sebert, Not. Bois Nouw. Calèd.* 193.—N. Caled.

CHRYSOPIA, Noronha, ex Thou. Gen. Nov. Madag. 14 (1806) = **Symphonia**, Linn. (Guttif.).
fasciculata, Thon. l. c. n. 48 = S. fasciculata.
gymnocarpa, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 291.—Madag.
macrophylla, Cambess. Mém. Tern. et Gutt. 55. t. 4.—Madag.
microphylla, Cambess. & Boj. ex Cambess. in Mem. Mus. Par. xvi. (1828) 423. t. 4.—Madag.
urophylla, Decne. ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 291.—Madag.
verrucosa, Hilsenb. & Boj. ex Planch. & Triana, l. c. 290.—Madag.

CHRYSOPOGON, Trin. Fund. Agrost. 187 (1820).
GRAMINEAE, Benth. & Hook. f. iii. 1185.
CENTROPHORUM, Trin. l. c. 106. t. 5 (1820).
PERIANTHERA, Rafin. in Ser. Bull. Bot. i. (1830) 291.
RAPHIS, Beauv. Agrost. 190 (1812).
RHAPHIS, Lour. Fl. Cochinch. 826 (1790).
TALANTHIUM, Desv. Opusc. 69, obs. (1851).
acicularis, *Trin. Fund. Agrost.* 188.—Reg. trop.
alternans, Trin. ex Steud. Nom. ed. II. i. 359 = violaceus!
argutus, *Trin. ex Steud. l. c.* 360.—Austral.
Arnottianus, *Nees, ex Steud. Syn. Pl. Gram.* 396.—Ind. or.
avenaceus, *Benth. in Journ. Linn. Soc. xix.* (1881) 78.—Ind. calid.
ciliolatus, Boiss. Fl. Orient. v. 656 = serrulatus.
filiformis, Volgt, Hort. Suburb. Calc. 704 = Dimeria ornithopoda.
elongatus, *Benth. Fl. Austral.* vii. 686.—Austral.

CHRYSOPOGON:—
Earnhockii, Arn. ex Steud. Syn. Pl. Gram. 395 = serrulatus.
Francavillanus, Hemsl. Biol. Centr. Am. Bot. iii. 530 = avenaceus.
fuscus, Trin. ex Steud. Nom. ed. II. i. 360 = Andropogon serratus.
glaberatus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 318 = C. Gryllus.
Gryllus, *Trin. Fund. Agrost.* 188.—Reg. trop. et subtrop.; Europ. austr.; Afr.
incrassus, Nees, ex Steud. Syn. Pl. Gram. 396 = serrulatus.
Minarum, Benth. in Journ. Linn. Soc. xix. (1881) 78 = stipoides.
monianus, *Trin. in Sprengl. Neue Entdech.* ii. 98.—Ind. or.
nutans, *Benth. in Journ. Linn. Soc. xix.* (1881) 78.—Am. calid.
pallidus, *Trin. ex Steud. Nom.* ed. II. i. 360.—Austral.
parviflorus, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 411 = violaceus.
pauciflorus, Benth. ex Vasey, Gram. U. St. 20 = Sorghum pauciflorum.
pictus, *Hance, in Ann. Sc. Nat.* Ser. V. v. (1866) 252.—As. trop.
quinqueplumis, *A. Rich. Tent. Fl. Abyss.* ii. 450.—Abyssin.
secundum, Benth. ex Vasey, Gram. U. St. 20 = Sorghum secundum.
serrulatus, *Trin. in Mém. Acad. Pétersb.* Ser. VI. ii. (1833) 318.—Ind. or.; Afr. trop.
stipoides, Benth. in Journ. Linn. Soc. xix. (1881) 78 = avenaceus.
stipoides, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 71.—Bras.
strictus, Nees, in Hook. Kew Journ. ii. (1850) 95 = Andropogon intermedius.
subtilis, *Miq. Fl. Ind. Bat.* iii 492.—Java.
subulatus, Trin. ex Steud. Nom. ed. II. i. 360 = acicularis.
tener, *Nees, ex Steud. Syn. Pl. Gram.* 397.—' Andor.'
trivialis, Arn. & Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 171 = acicularis.
verticillatus, *Trin. ex Steud. Nom.* ed. II. i. 360.—Ind. or.
violaceus, *Trin. in Mem. Acad. Pétersb.* Ser. VI. ii. (1833) 319.—As. trop.; Abyssin.; Austral.
Wightianus, *Thw. Enum. Pl. Zeyl.* 366.—Zeylan.
Wrightii, Munro, ex Vasey, Cat. Gram. U. St. 20 = Sorghum pauciflorum.
zeylanicus, Thw. Enum. Pl. Zeyl. 366 = Arnottianus.

CHRYSOPSIS, Ell. Sketch, ii. 338 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 252.
AMMODIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 321.
DIPLOGON, Rafin. in Am. Monthly Mag. (1818) 262.
HECTORIA, DC. Prod. v. 24 (1836).
HEYFELDERA, Sch. Bip. in Flora, xxxvi. (1853) 35.
MACRONEMA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 322.
PITYOPSIS, Nutt. l. c. (1841) 317.
acaulis, Nutt. in Journ. Acad. Philad. vii. (1834) 33 = Aplopappus acaulis.
alba, DC. Prod. v. 284, in syn. = Aster ptarmicoides.
alpina, Nutt. in Journ. Acad. Philad. vii. (1834) 34. t. 2. = Aster scopulorum.
amygdalina, DC. Prod. v. 272, in syn. = Aster umbellatus.
andicola, *Phil. in Anal. Univ. Chil.* (1868) ii. 208; *et in Linnaea*, xxxiii. (1864-65) 121.—Chili.
argentea, Ell. Sketch, ii. 334 = graminifolia.
Belantieri, A. Gray, in Proc. Am. Acad. vi. (1864-65) 548 = villosa.
Bewori, *A. Gray, l. c.* 543.—Calif.
canaglina, Nutt. in Journ. Acad. Philad. vii. (1834) 33 = Aplopappus acaulis.
canescens, DC. Prod. v. 339 = Erigeron filifolius.
canescens, Torr. & Gray, Fl. N. Am. ii. 256 = villosa.
carenophylla, Nutt. in Journ. Acad. Philad. vii. (1834) 34 = Aster tanacetifolius.
decumbens, Chapm. Fl. S. U. St. 317 = gossypina.
dentata, Ell. Sketch, ii. 337 = gossypina.
diversicola, Ell. l. c. 339 = Aster divaricatus.

CHRYSOPSIS:—
schiedeæ, Benth. Bot. Voy. Sulph. 25 = villosa.
ericoides, Torr. & James, ex Eaton, N. Am. Bot. ed. VIII. 196 = Aster ericoëfolius.
falcata, Ell. Sketch, ii. 336 = villosa.
falcata, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 316 = villosa.
gossypina, *Ell. Sketch*, ii. 337.—Am. bor.
graminifolia, *Ell. l. c.* 334.—Am. bor.
hirtella, DC. Prod. v. 337 = Erigeron chrysopsidis.
hispida, DC. l. c. vii. 279 = villosa.
humilis, DC. l. c. v. 278, in syn. = Aster infirmus.
hyssopifolia, *Nutt. in Journ. Acad. Philad.* vii. (1834) 67.—Am. bor.
Lamarckii, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 315 = Aplopappus spinulosus.
linariifolia, DC. Prod. v. 277 = Aster linariifolius.
linifolia, DC. l. c. 255, in syn. = Aster acris.
Mariana, *Ell. Sketch*, ii. 335.—Am. bor.
mollis, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 316 = villosa.
obovata, Nutt. Gen. Am. ii. 152 = Aster reticulatus.
oligantha, Chapm. ex Torr. & Gray, Fl. N. Am. ii. 253 = graminifolia.
oregana, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 548.—Am. bor. occ.
pilosa, *Nutt. in Journ. Acad. Philad.* vii. (1834) 66. —Texas.
pinifolia, *Ell. Sketch*, ii. 335.—Am. bor.
scabra, Ell. l. c. 339 = Heterotheca Lamarckii.
scabra, Hook. & Arn. Bot. Beech. Voy. 484 = Oxypappus scaber.
scabrella, *Torr. & Gray, Fl. N. Am.* ii. 255.—Am. bor.
sessiliflora, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 317 = villosa.
trichophylla, *Ell. Sketch*, ii. 336.—Am. bor.
villosa, *DC. Prod.* v. 287; vii. 279.—Am. bor.
Wrightii, *A. Gray, Syn. Fl. N. Am.* i. ii. 445.—Calif.

CHRYSORHOË, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 357 = Wertordtia, DC. (Myrsac.).
nitens, Lindl. l. c. = V. nitens.
serrata, Lindl. Swan River App. 6 = V. serrata.

CHRYSOSCIAS, E. Mey. Comm. Pl. Afr. Austr. 139 (1835) = Rhynchosia, Lour. (Legumin.).
calycina, E. Mey. l. c. 140 = R. lencoscias.
floribunda, Lem. in Illustrat. Hortic. i. (1854) Misc. 56; ii. (1856) t. 47.—Afr. austr.
grandiflora, E. Mey. Comm. Pl. Afr. Austr. i. 139 = R. Chrysoscias.
media, E. Mey. l. c.—Afr. austr.
parviflora, E. Mey. l. c. = R. microscias.

CHRYSOSPERMUM, Reichb. Consp. 166 (1828), nomen = Anthospermum, Linn. (Rubiac.).
repens, Pritz. Ind. Ic. 263, sphalm. = *Cryphiospermum repens*, Beauv. = Enhydra fluctuans, *Lour.* (Compos.).

CHRYSOSPLENIUM, Tourn. ex Linn. Syst. i. (1735). *SAXIFRAGRAE*, Benth. & Hook. f. i. 633.
adenoides, Hook. f. & Thoms. ex Maxim. in Bull. Acad. Pétersb. xxiii. (1877) 349 = lanuginosum.
altum, *Maxim. l. c.* xvii. (1872) 490.—Japon.
alpinum, Schur, in Verh. Siebenb. Ver. Naturw. v. (1855) 86 = oppositifolium.
alternans, Thunb. Fl. Jap. 183 = alternifolium.
alternifolium, *Linn. Sp. Pl.* 398.—Europ.; As. bor.; Am. bor.
americanum, *Schwein. ex Hook. Fl. Bor. Am.* i. 342.—Am. bor.
sclaocarpum, *Ernst, in Flora*, lvii. (1874) 214.—Venezuela.
auriculatum, Cunn. Inst. i. 139 = oppositifolium.
axillare, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 341.—China.
baicalense, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 91.—Sibir.
carnosulum, Maxim. in Bull. Acad. Pétersb. xxiii. (1877) 340 = carnosum.
carnosum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 78.—Kap. Hlmsl.
Davidianum, *Decne. ex Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 342.—Tibet, &c.
Delavayi, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) [illegible]

CHRYSOSPLENIUM:—
Dickinsii, *Franch. & Sav. Enum. Pl. Jap.* ii. 357.—Japon.
dubium, J. Gay, ex DC. Prod. iv. 48 = macrocarpum.
echimulatum, *Franch. & Sav. Enum. Pl. Jap.* ii. 358.—Japon.
Echinus, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 347.—Japon.
elevatum, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 281 = macranthum.
Fauriæ, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 55.—Japon.
flagelliferum, *F. Schmidt, Reisen Amuri.* 194.—As. or.
flagelliforme, Franch. & Sav. Enum. Pl. Jap. ii. 355, Japan = præc.
grandes, Fisch. ex Ledeb. Fl. Ross. ii. 227 = alternifolium.
glaciale, Fuss, Fl. Transs. 247 = oppositifolium.
glechomæfolium, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 589.—Am. bor. occ.
Grayanum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 346.—Japon.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 74.—Reg. Himal.
kamtschaticum, *Fisch. ex DC. Prod.* iv. 48.—Kamtschat.
kamtschaticum, A. Gray, Bot. Jap. 389 = ramosum.
lanuginosum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 74.—Reg. Himal.
macranthum, *Hook. Lond. Journ. Bot.* i. (1842) 458. t. 16.—Patagon.
macrocarpum, *Cham. in Linnæa*, vi. (1831) 558.—Am. arct.
macrostemon, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* ii. 358.—Japon.
Maximowiczii, *Franch. & Sav. l. c.*—Japon.
multicaule, *Franch. & Sav. l. c.* 361.—Japon.
nepalense, *D. Don, Prod. Fl. Nep.* 210.—Reg. Himal.
nepalense, Maxim. Prim. Fl. Amur. 121, in nota = sulcatum.
nipponicum, *Franch. & Sav. Enum. Pl. Jap.* ii. 356.—Japon.
nivale, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 98 = alternifolium.
nudicaule, *Bunge, in Ledeb. Fl. Alt.* ii. 114.—Sibir.
octandrum, Caqué, ex Steud. Nom. ed. II. i. 360 = oppositifolium.
oppositifolium, Cham. in Linnæa, vi. (1831) 557 = kamtschaticum.
oppositifolium, *Linn. Sp. Pl.* 398.—Europ.; As. bor.; Ind. or.
oppositifolium, Trautv. & Mey. in Middend. Reise (Fl. Ochot). 42 = ramosum.
oppositifolium, Walt. Fl. Carol. 140 = americanum.
ovalifolium, Bertol. Fl. Ital. iv. 449 = macrocarpum.
ovalifolium, *Bieb. ex Ledeb. Fl. Alt.* ii. 115.—Sibir. altaic.
ovalifolium, Miq. Ann. Mus. Lugd. Bat. iii. 87 = macrocarpum.
papillosum, Franch. & Sav. Enum. Pl. Jap. ii. 355 = alternifolium.
peltatum, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 278.—Sibir.
piloseum, *Maxim. Prim. Fl. Amur.* 122.—Manchur.
ramosum, *Maxim. l. c.* 121.—Manchur.
repens, Link, ex Steud. Nom. ed. II. i. 360 = oppositifolium.
rhabdospermum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 347.—As. or.
rosulare, Schott, ex Maxim. l. c. 345 = oppositifolium.
rotundifolium, Schlecht. ex Ledeb. Fl. Ross. ii. 227 = alternifolium.
Sedakowii, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 278.—Sibir.
sibiricum, Steph. ex DC. Prod. iv. 48 = alternifolium.
sinicum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 348.—China.
sphærospermum, *Maxim. l. c.* 349.—Japon.
sulcatum, Maxim. l. c. 347 = nepalense.
tenellum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 73.—Reg. Himal.
tetrandrum, Fries, in Bot. Notiser (1858), 193 = alternifolium.
trachyspermum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 474.—China.
tracuulivalense, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 98.—Transsylvan.

CHRYSOSPLENIUM:—
trichospermum, *Edgew. ex Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 78.—Reg. Himal.
uniflorum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 472.—China.
valdivicum, *Hook. Lond. Journ. Bot.* i. (1842) 459 t. 17.—Chili.
Vidalii, *Franch. & Sav. Enum. Pl. Jap.* ii. 360.—Japon.
Wrightii, *Franch. & Sav. l. c.* 356.—Sibir. or.
yesoense, *Franch. & Sav. l. c.* 355.—Japon.
yunnanense, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 7.—China.

CHRYSOSTACHYS, Pohl, Pl. Bras. ii. 65. t. 148 (1831) = Combretum, Linn.
ovatifolia, Pohl, l. c. 66.—Bras.

CHRYSOSTEMMA, Less. Syn. Comp. 227 (1832) = Coreopsis, Linn. (Compos.).
calendulacea, E. Mey. ex DC. Prod. vi. 501 = Gorteria calendulacea.
senifolium, Shuttl. ex Walp. Ann. ii. 857 = Cor. senifolia.
Tripteris, Less. Syn. Comp. 227 = Cor. Tripteris.

CHRYSOSTEMON, Klotzsch, in Lehm. Pl. Preiss. ii. 282 (1847) = Pseudanthus, Sieber (Compos.).
virgatus, Klotzsch, l. c. = P. virgatus.

CHRYSOSTOMA, Lilja, Fl. Suppl. i. 33 (1840); et in Linnæa, xv. (1841) 365 = Martinella, Linn. (Loas.).
aurea, Lilja, li. cc. 33, 364 = M. oligosperma.

CHRYSOTHAMNUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 323 = Bigelovia, DC. (Compos.).
depressus, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 171 = B. depressa.
dracunculoides, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 324 = B. dracunculoides.
lancolatus, Nutt. l. c. = B. viscidiflora.
pumilus, Nutt. l. c. 323 = B. viscidiflora.
speciosus, Nutt. l. c. = B. viscidiflora.
viscidiflorus, Nutt. l. c. 324 = B. viscidiflora.

CHRYSOTHEMIS, Decne. in Rev. Hortic. Sér. III. iii. (1849) 349 = Tussacia, Reichb. (Gesneriac.).
aurantiaca, Decne. l. c. iv. (1850) 381.—Ind. occ.
melissæfolia, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1409 = T. pulchella.
multifolia, G. Don, ex Loud. l. c. = præc.
ovata, Klotzsch, in Schomb. Fauna et Fl. Gui. 108 — Guiana.
pulchella, Decne. in Rev. Hortic. Sér. III. iii. (1849) 349 = T. pulchella.
venosa, Decne. l. c. = Episcia melittifolia †

CHRYSOTHRIX, Roem. & Schult. Mant. ii. 6. 144 (1824) = Chrysothrix, Linn. (Cyperac.).

CHRYSOXYLON, Cæsar. Nov. Stirp. Bras. Dec. 59 (1843) = Plathymenia, Benth. (Legumin.).
Vinhatico, Cæsar. l. c. = P. reticulata.

CHRYSOXYLON, Wedd. Hist. Nat. Quinquin. 100 (1849) = Pogonopus, Klotzsch (Rubiac.).
febrifugum, Wedd. l. c.—Bolivia.

CHRYSURUS, Pers. Syn. i. 80 (1805) = Lamarckia, Moench (Gramin.).
aurea, Beauv. Agrost. 128 = L. aurea.
cynosuroides, Pers. Syn. i. 80 = L. aurea.
echinatus, Beauv. Agrost. 128 = Cynosurus echinatus.
effusus, Beauv. l. c. = Cynosurus elegans.
elegans, Beauv. l. c. = Cynosurus polybracteatus.
giganteus, Tenore, Fl. Nap. iv. 13. t. 210 = Cynosurus echinatus.
gracilis, Moris, Stirp. Sard. Elench. i. 50 = Cynosurus gracilis.

CHRYTOTHECA, G. Don, Gen. Syst. ii. 706 (1832) = CRYPTOTHECA, Blume = Ammannia, Houst. (Lythrar.).

CHRYZA, Rafin. in Med. Repos. N. York, v. (1808) 353 = Coptis, Linn. (Ranunculac.).
borealis, Rafin. l. c. = Coptis trifolia.

CHTHAMALIA, Decne. in DC. Prod. viii. 605 (1844) = Gonolobus, Michx. (Asclepiad.).
bifora, Decne. l. c.—G. biflorus.
hubsifusa, Fourn. in Mart. Fl. Bras. vi. IV. 318.—Bras.
major, Fourn. l. c. 314.—Bras.
nummularia, Decne. in DC. Prod. viii. 605 = G. nummularius.
pedunculata, Decne. l. c. = G. pedunculatus.
pubiflora, Decne. l. c. = G. pubiflorus.
purpurea, Decne. l. c.—Bras.

CHTHONIA, Cass. in Dict. Sc. Nat. ix. 173 (1817) = Perdix, Linn. (Compos.).
Bonplandiana, Cass. l. c. xxvii 206 = P. Bonplandiana.
canescens, Cass. l. c. = P. canescens.
ciliaris, Cass. l. c. = P. ciliaris.
elongata, Cass. l. c. = P. elongata.
glaucescens, Cass. l. c. ix. 173 = P. ciliaris.
humifusa, Cass. l. c. = P. humifusa.
leptocephala, Cass. l. c. xxvii. 206 = P. Swartziana.
prostrata, Cass. l. c. ix. 173 = P. prostrata.
pygmaea, Cass. l. c. xxvii. 206 = P. pygmaea.
repens, Cass. l. c. = P. serpyllifolia.

CHTHONOCEPHALUS, Steetz, in Lehm. Pl. Preiss. i. 444 (1845). *COMPOSITAE*, Benth. & Hook. f. ii. 392.
CHAMAESPHAERION, A. Gray, in Hook. Kew Journ. iii. (1851) 176.
GYROSTEPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 76.
LACHNOTHALAMUS, F. Muell. Fragm. iii. 156 (1863).
Drummondii, *A. Gray, in Hook. Kew Journ.* iii. (1851) 178.—Austral.
pseudevax, *Steetz, in Lehm. Pl. Preiss.* i. 445.—Austral.
pygmaeus, *Benth. Fl. Austral.* iii. 582.—Austral.
tomentellus, *Benth. l. c.* 581.—Austral.

CHUKRASIA, A. Juss. in Mém. Mus. Par. xix. (1830) 351. t. 23, vide Chickrassia, A. Juss. (Meliac.).

CHULUSIUM, Rafin. Ann. Nat. 12 (1820) = Polygonum, Tourn.
acuminatum, Rafin. Fl. Tellur. iii. 14 = P. acuminatum.
amphibium, Rafin. l. c. = P. amphibium.
filiforme, Rafin. l. c. = P. tenue.
fluitans, Rafin. l. c. = P. amphibium ?
natans, Rafin. l. c. = P. amphibium ?

CHUNCHOA, Pers. Syn. i. 486 (1805) = seq.

CHUNCOA, Pav. ex Juss. Gen. 76 (1789) = Terminalia, Linn. (Combretac.).
actinophylla, Griseb. in Goett. Abh. xxiv. (1879) 132 = T. actinophylla.
amazonia, J. F. Gmel. Syst. 702.—Bras.
arbuscula, Griseb. Fl. Brit. W. Ind. 276 = T. arbuscula.
brasiliensis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 244 = T. brasiliensis, *Eichl.*
Chicharronia, Griseb. Cat. Pl. Cub. 109 = T. Chichacronia.
diptera, F. G. Dietr. Vollst. Lex. Gaertn. Nachtr. ii. 244 = T. oblonga.
eriostachya, Griseb. Cat. Pl. Cub. 109 = T. eriostachya.
flavescens, Presl, Epim. Bot. 215.—Bras.
glabra, Buch.-Ham. ex Wall. Cat. sub n. 3979 = T. Arjuna.
lactiflua, F. Muell. Fragm. ii. 149 = T. bursarina.
oblonga, Pers. Syn. i. 486 = T. oblonga.
obovata, Pers. l. c. = T. obovata.
tomentosa, Buch.-Ham. ex Wall. Cat. sub n. 3978 = T. tomentosa.
trifoxa, Griseb. in Goett. Abh. xxiv. (1879) 132.—Reg. Argent.

CHUPALON Adans. Fam. ii. 164 (1763) = Thibaudia, Pav. (Vacciniac.).

CHUPALONES, Nieremb. ex Steud. Nom. ed. II. i. 361 (1840) = prec.

CHUQUIRAGA, Juss. Gen. 176 (1789). *COMPOSITAE*, Benth. & Hook. f. ii. 488.
DASYPHYLLUM, H. B. & K. Nov. Gen. et Sp. iv. 17. t. 308 (1820).
DIACANTHA, Leg. Amen. Nat. i. 40 (1811).
FLOTOVIA, Spreng. Syst. iii. 359 (1826).
JOANNESIA, Pers. Syn. ii. 388 (1807).
JOHANNIA, Willd. Sp. Pl. iii. 1705 (1801).
PITOCARPA, Hook. & Arn. in Comp. Bot. Mag. i. (1835) 110.
acanthophylla, *Wedd. Chlor. And.* i. 5.—Bolivia.
acicularis, *D. Don, in Phil. Mag.* (1832) 392.—Chili.
alpina, Poepp. ex Less. Syn. Comp. 96 = oppositifolia.
anomala, *D. Don, in Phil. Mag.* (1832) 392.—Chili.
Avellanedae, *Lorentz, in Roca, Exped. Rio-Negro*, ii. 245.—Patagon.
Candolleana, *Baker, in Mart. Fl. Bras.* vi. III. 355.—Bras.
chrysantha, *Field. & Gardn. Sert. Pl.* t. 42.—Chili.
cryptocephala, *Baker, in Mart. Fl. Bras.* vi. III. 355.—Bras.
Douiana, *Baker, l. c.* 358.—Bras.
Echegaryi, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 47.—Reg. Argent.
erinacea, *D. Don, in Phil. Mag.* (1832) 392.—Chili; Patagon.
exonisa, *D. Don, l. c.*—Chili.
floribunda, *Baker, in Mart. Fl. Bras.* vi. III. 362.—Bras.
fodinarum, *Baker, l. c.* 356.—Bras.
fruticosa, Juss. ex Steud. Nom. ed. II. i. 861 = insignis.
glabra, *Baker, in Mart. Fl. Bras.* vi. III. 363.—Bras.
Griseb*achii*, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 359.—Reg. Argent.
hispida, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 287.
Hystrix, *D. Don, in Phil. Mag.* (1832) 392.—Chili; Patagon.
incana, D. Don, l. c. = ulicina.
infundibularis, *Baker, in Mart. Fl. Bras.* vi. III. 357.—Bras.
insignis, Humb. & Bonpl. Pl. Aequin. i. 153 = Jussieni.
jamesonensis, *Decne. in Orbign. Voy. Bot.* t. 11.—Amaustr.
Jussieui, *Y. F. Gmel. Syst.* 1205.—Peruv.; Ecuador.
Kingii, *Bell, in Journ. Linn. Soc.* xxi. (1884) 224.—Patagon.
lanciifolia, *Humb. & Bonpl. Pl. Aequin.* i. 153.—Peruv.; Ecuador.
latifolia, *Baker, in Mart. Fl. Bras.* vi. III. 357.—Bras.
latifolia, D. Don, in Trans. Linn. Soc. xvi. (1830) 288 = glabra.
leptacantha, *Baker, in Mart. Fl. Bras.* vi. III. 360.—Bras.
leucocryon, *Poepp. ex Less. Syn. Comp.* 95, in syn.—Chili.
linearis, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 94.—Chili.
macrantha, *Baker, in Mart. Fl. Bras.* vi. III. 359.—Bras.
microphylla, Humb. & Bonpl. Pl. Aequin. i. 151. t. 43 = Jussieui.
oppositifolia, *D. Don, in Phil. Mag.* (1832) 392.—Chili.
orthacantha, *Baker, in Mart. Fl. Bras.* vi. III. 360.—Bras.
paniculata, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 289.—Bras.
patagonica, *Phil. in Linnaea*, xxxiii. (1864–65) 111.—Patagon.
peruviana, Jaume St. Hil. Fam. Nat. i. 204 = Jussieui.
racemosa, *Baker, in Mart. Fl. Bras.* vi. III. 363.—Bras.
Regnellii, *Baker, l. c.* 360.—Bras.
reticulata, *DC. Prod.* vii. 9.—Bras.
revoluta, *Field. & Gardn. Sert. Pl.* sub t. 42.—Chili.
rotundifolia, *Wedd. Chlor. And.* i. 4. t. 4.—Peruv.
ruscifolia, *D. Don, in Phil. Mag.* (1832) 392.—Chili.
spinescens, *Baker, in Mart. Fl. Bras.* vi. III. 362.—Bras.
spinosa, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 286.—Peruv.
Sprengeliana, *Baker, in Mart. Fl. Bras.* vi. III. 357.—Bras.
synacantha, *Baker, l. c.* 361.—Bras.
tomentosa, Baker, l. c. 360 = paniculata.
trichophylla, *Baker, l. c.* 356.—Bras.
ulicina, *Hook. Comp. Bot. Mag.* i. (1835) 110.—Chili.

CHUQUIRAGA :—
Ungnis-cati, *Cas. in Atti Soc. Sc. Nap.* v. (1878) 11. n. 7.—Hab. ?
vaginata, *Baker, in Mart. Fl. Bras.* vi. III. 361.—Bras.
velutina, *Baker, l. c.* 358.—Bras.
verpreculata, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 290.—Venezuela.

CHURUMAYA, Rafin. Sylva Tellur. 85 (1838) = Piper, Linn.
arborea, Rafin. l. c. = P. Churumaya.

CHUSQUEA, Kunth, Syn. Pl. Aequin. i. 254 (1822). *GRAMINEAE*, Benth. & Hook. f. iii. 1209.
DENDRAGROSTIS, Nees, in Linnaea, ix. (1834) 487.
MUSTELLA, Cav. ex Steud. Nom. ed. II. i. 361, in syn. (1840); ii. 168.
RETTERSIA, Raddi, Agrost. Bras. 17 (1823).
abietifolia, *Griseb. Fl. Brit. W. Ind.* 539.—Ind. occ.
acuminata, *Doell, in Mart. Fl. Bras.* ii. III. 204.—Bras.
amplexipeniculata, Steud. Syn. Pl. Gram. 337 = Dinochloa Tjankorreh.
andina, *Phil. in Linnaea*, xxix. (1857–58) 108.—Chili.
anclythra, *Nees, in Linnaea*, ix. (1834) 491.—Bras.
anclytroides, *Rupr. ex Doell, in Mart. Fl. Bras.* ii. III. 206.—Bras.
aristata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 61.—Ecuador.
Billmeki, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 587.—Mexic.
breviglumis, Phil. in Linnaea, xxix. (1857–58) 108 = C. Coleou.
capitata, *Nees, in Linnaea*, ix. (1834) 489.—Bras.
capituliflora, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. (1835) 318.—Bras.
carinata, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 587.—Mexic.
ciliata, Phil. in Linnaea, xxxiii. (1864–65) 299 = tenuiflora.
Culeou, *E. Desv. in C. Gay, Fl. Chil.* vi. 450.—Chili.
Caminglii, *Nees, in Linnaea*, ix. (1834) 487.—Chili.
Domheyana, *Kunth, Rev. Gram.* ii. 555. t. 191.—Peruv.
fasciculata, *Doell, in Mart. Fl. Bras.* ii. III. 202.—Bras.
Fendleri, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 61.—Venezuela.
Fernandeziana, *Phil. in Anal. Univ. Chil.* (1873) 577.—Chili.
fimbriata, Steud. Syn. Pl. Gram. 338 = Merostachys capitata.
Galeottiana, *Rupr. in Bull. Acad. Brux.* ix. II. (1842) 246.—Mexic.
Gandichaudii, *Kunth, Rev. Gram.* i. 138, 331. t. 78.—Bras.
gionorrhiza, Munro, in Trans. Linn. Soc. xxvi. (1868) 50 = Merostachys capitata.
heterophylla, *Nees, in Linnaea*, ix. (1834) 488.—Bras.
humilis, Lechl. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 60 = spicata.
intermedia, Steud. in Lechl. Berb. Am. Austr. 52, nomen = C. Quila.
Yamesoni, Steud. Syn. Pl. Gram. 337 = scandens.
leptophylla, Nees, in Linnaea, ix. (1834) 489 = Arthrostylidium leptophyllum, Trinii.
Liebmanni, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 587.—Mexic.
ligulata, Munro, in Trans. Linn. Soc. xxvi. (1868) 62 = Fernandeziana.
Lorentziana, *Griseb. in Goett. Abh.* xix. (1874) 249.—Reg. Argent.
Meyeriana, *Rupr. ex Doell, in Mart. Fl. Bras.* ii. III. 208.—Bras.
montana, *Phil. in Linnaea*, xxxiii. (1864–65) 299.—Chili.
Mulleri, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 65.—Mexic.
oligophylla, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1859) 154.—Bras.
nigricans, *Phil. in Anal. Univ. Chil.* (1865) ii. 223.—Chili.
pallida, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 66.—M. Gaumer.
parviflora, Phil. in Linnaea, xxxiii. (1864–65) 299 = Coleou.
pinifolia, *Nees, l. c.* ix. (1834) 490.—Bras.

CHUSQUEA:—
pubescens, Steud. Syn. Pl. Gram. 337 = C. Quila.
Pentlandiana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 56.—N. Granat.
Quila, E. Desv. in C. Gay, Fl. Chil. vi. 447 = tenuiflora.
Quila, *Kunth, Rev. Gram.* i. 138, 329. t. 77.—Chili.
scandens, *Kunth, Syn. Pl. Aequin.* i. 254.—Am. austr.
sclerophylla, *Doell, in Mart. Fl. Bras.* ii. III. 200.—Bras.
Sellowii, *Rupr. in Mém. Acad. Pétersb.* Ser. VI. v. (1839) 126.—Bras.
simpliciflora, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 54.—Panama.
simpliciflora, Hemsl. Biol. Centr. Am. Bot. iii. 487, sphalm. = simplicifloa.
Spencei, *Ernst, in Journ. Bot.* x. (1872) 269.—Venezuela.
spicata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 60.—Peruv.
spinosa, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 487.—Mexic.
tenella, *Nees, in Linnaea*, ix. (1834) 492.—Bras.
tenuiflora, *Phil. in Linnaea*, xxx. (1859–60) 206.—Chili.
tenuigimis, *Doell, in Mart. Fl. Bras.* ii. III. 199.—Bras.
tessellata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 60.—N. Granat.
uliginosa, *Phil. in Linnaea*, xxx. (1859–60) 207.—Chili.
uniflora, *Steud. Syn. Pl. Gram.* 337.—Ins. Cuba.
valdiviensis, E. Desv. in C. Gay, Fl. Chil. vi. 446 = C. Quila.
Venezuelae, *Steud. Syn. Pl. Gram.* 337.—Venezuela.
Wilkesii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 63.—Bras.

CHYDENANTHUS, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 111 = **Barringtonia**, Forst. (Myrtac.).
excelsus, Miers, l. c. 112 = B. excelsa.

CHYLOCALYX, Hassk. in Flora, xxv. (1842) 11. Beibl. 20 = **Polygonum**, Tourn.
perfoliatus, Hassk. l. c. = P. perfoliatum.
senticosus, Meisn. in Miq. Ann. Mus. Bot. Lugd. Bat. ii. 65 = P. senticosum.

CHYLODIA, Rich. ex Cass. in Dict. Sc. Nat. xxix. 491 (1823) = **Wulffia**, Neck. (Compos.).
serrantoides, Rich. l. c. 492 = W. senegalensis.

CHYLOGALA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = **Euphorbia**, Linn.
serrata, Fourr. l. c. = E. serrata.

CHYMOCARPUS. G. Don, in Trans. Linn. Soc. xvii. 13 (1834) = **Tropaeolum**, Linn. (Geraniac.).
brachyceras, Heynh. Nom. 1. 195 = T. brachyceras.
Yaratitii, Heynh. l. c. = T. jaratiti.
pentaphyllus, D. Don, in Trans. Linn. Soc. : vii. (1834) 14 = T. pentaphyllum.
polyphyllus, Heynh. Nom. l. 195 = T. polyphyllum.
speciosus, Walp. Ann. i. 149 = T. speciosum.
tricolor, Heynh. Nom. l. 195 = T. tricolorum.
tuberosus, Heynh. l. c. = T. tuberosum.

CHYMOCOCCA, Meisn. in DC. Prod. xiv. 565 (1857). *THYMELAEACEAE*, Benth. & Hook. f. iii. 194.
empetroides, *Meisn. l. c.*—Afr. austr.

CHYMOCORMUS, Harv. in Hook. Lond. Journ. Bot. i. (1842) 23 = **Fockea**, Endl. (Asclepiad.).
edulis, Harv. l. c. 24.—Afr. austr.

CHYSIS, Lindl. Bot. Reg. xxiii. t. 1937 (1837). *ORCHIDEAE*, Benth. & Hook. f. iii. 514.
aurea, *Lindl. l. c.*—Chili.
bractescens, *Lindl. l. c.* (1840) *Misc.* 61.—Mexic.
bracteoscens, *Reichb. f. & Warsc. in Bot. Zeit.* xv. (1857) 357.—Peruv.
Chelsoni x, Reichb. f. in Gard. Chron. (1874) i. 585 (cf. p. 365).
laevis, *Lindl. Bot. Reg.* (1840) *Misc.* 61.—Mexic.
Limminghii, *Linden & Reichb. f. in C. Koch, Allg. Gartenz.* xxvi. (1858) 205.—Mexic.
Maleyi, *Mesmb. Nom.* ii. 148.—Mexic.
Salmo x, Reichb. f. in Gard. Chron. (1880) i. 616.

CHYTRA, Gaertn. f. Fruct. iii. 184. t. 214 (1805) = **Buchnera**, Linn. ! (Scrophularin.).
anomala, Gaertn. f. l. c.—Hab. ?

CHYTRACULIA, P. Br. Hist. Jamaic. 239. t. 37 (1756) = **Calyptranthes**, Sw. (Myrtac.).

CHYTRALIA, Adans. Fam. ii. 80 (1763) = **Calyptranthes**, Sw. (Myrtac.).

CHYTRANTHUS, Hook. f. in Benth. & Hook. f. Gen. i. 408 (1868). *SAPINDACEAE*, Benth. & Hook. f. i. o.
Mannii, *Hook. f. l. c.*—Afr. trop.
Prieurianus, *Baill. Adansonia*, xi. (1873–76) 241.—Guiana.

CHYTROGLOSSA, Reichb. f. in Hamb. Gartenz. xix. (1863) 546. *ORCHIDEAE*, Benth. & Hook. f. iii. 569.
aurata, *Reichb. f. l. c.*—Hab. ?
Marileoniae, *Reichb. f. l. c.*—Bras.

CHYTROMA, Miers, in Trans. Linn. Soc. xxx. (1874) 229 = **Lecythis**, Loefl. (Myrtac.).
amara, Miers, l. c. 231 = L. amara.
apiculata, Miers, l. c. 245.—Bras.
basilaris, Miers, l. c. 239.—Guiana.
chartacea, Miers, l. c. 231 = L. chartacea.
cinctuata, Miers, l. c. 237.—Bras.
cistella, Miers, l. c. 233.—Bras.
grandifolia, Miers, l. c. 227 = L. grandiflora.
Idatimi, Miers, l. c. 236.—Bras.
Idatimon, Miers, l. c. 239 = L. Idatimon.
incarcerata, Miers, l. c. 232.—Bras.
maranyensis, Miers, l. c. 245 = L. maranyensis.
monosperma, Miers, l. c. 239 = L. monosperma.
ovalifolia, Miers, l. c. 244 = L. ovalifolia.
perspicua, Miers, l. c. 246.—Guiana.
pilocarpa, Miers, l. c. 238.—Bras.
retusa, Miers, l. c. 243 = L. retusa.
rorida, Miers, l. c. = L. rorida.
rosea, Miers, l. c. 242 = L. rosea.
rubriflora, Miers, l. c. 241.—Guiana.
salebrosa, Miers, l. c. 240 = L. salebrosa.
Schomburgkiana, Miers, l. c. 230 = L. Schomburgkii.
Spruceana, Miers, l. c. 235 = L. Spruceana.
turbinata, Miers, l. c. 234 = L. turbinata.
urceolata, Miers, l. c. 237.—Bras.
valida, Miers, l. c. 241.—N. Granat.

CIANITIS, Reinw. in Syll. Ratisb. ii. (1828) 10 = CYANITIS, Reinw. = **Dichroa**, Lour. (Saxifrag.).

CICCA, Linn. Mant. i. 124 (1767) = **Phyllanthus**, Linn. (Euphorbiac.).
acidissima, Blanco, Fl. Filip. ed. I. 700 = P. acidissimus.
alternoides, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 239.—Burma.
anomala, Baill. Etud. Gen. Euph. 619 = P. indicus.
antillana, A. Juss. Tent. Euph. 108. t. 4. f. 13 B = P. nobilis.
arborea, Wright, ex Griseb. in Goett. Nachr. (1865) 166 = P. nobilis.
brasiliensis, Baill. Etud. Gén. Euph. 618, nomen = P. nobilis.
chinensis, Baill. l. c.; Atlas, 45 = P. nobilis.
decandra, Blanco, Fl. Filip. ed. I. 701 = P. reticulatus.
discoidea, Baill. in Adansonia, i. (1860–61) 85 = P. discoideus.
distincta, Linn. Mant. i. 124 = P. distichus.
Emblica, Kurz, For. Fl. Brit. Burma, ii. 352 = P. Emblica.
formosa, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 35 = P. japonicus.
Guerinoviana, Msell. Arg. in DC. Prod. xv. ii. 414, in syn. = P. cyanocarpus.
Leucophyllus, Kurz, For. Fl. Brit. Burma, ii. 353 = Flueggia Leucopyrus.
macrocarpa, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 239.—Burma.
macrostachya, Benth. Bot. Voy. Sulph. 166 = Picrmannia Antidesma.
microcarpa, Benth. Fl. Hongk. 319 = P. reticulatus.
multiflora, Lam. Encyc. ii. 1 = P. distichus.
obovata, Kurz, For. Fl. Brit. Burma, ii. 354 = Flueggia microcarpa.

CICCA :—
Pawoniana, Baill. Etud. Gén. Euph. 618 = Phyllanthus nobilis.
pentandra, Blanco, Fl. Filip. ed. I. 701 = Flueggia microcarpa.
racemosa, Lour. Fl. Cochinch. ii. 556 = P. acidissimus?
reticulata, Kurz, For. Fl. Brit. Burma, ii. 354 = P. reticulatus.
rhomboidalis, Baill. Adansonia, ii. (1861–62) 51 = P. rhomboidalis.
scandens, Griseb. in Goett. Nachr. (1865) 165 = P. scandens.
sinica, Baill. Etud. Gén. Euph. 618. t. 24 = P. nobilis.
surinamensis, Miq. in Linnaea, xxi. (1848) 479 = P. nobilis.
virens, Wright, ex Griseb. in Goett. Nachr. (1865) 166 = P. virens.

CICENDIA, Adans. Fam. ii. 508 (1763). *GENTIANEAE*, Benth. & Hook. f. ii. 810.
campestris, Rafin. Fl. Tellur. iii. 26 = Gentiana campestris.
Candollei, Griseb. Gen. et Sp. Gent. 158 = pusilla.
exalata, Griseb. l. c. 150 = Erythraea Douglasii.
fastigiata, Griseb. l. c. 158 = Hoppea fastigiata.
filiformis, Delarb. Fl. Auv. i. 20 ; ed. II. 30 = Microcala filiformis.
glacialis, Rafin. Fl. Tellur. iii. 26 = Gentiana tenella.
hyssopifolia, Wight & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 249 = Enicostema littorale.
microphylla, Edgew. in Trans. Linn. Soc. xx. (1851) 83 = Schaea khasiana.
Pooppigii, Griseb. in DC. Prod. ix. 62 (Microcalae sp). —Bras.
pulchella, Griseb. Gen. et Sp. Gent. 159 = Erythraea ramosissima.
pusilla, *Griseb. l. c.* 157.—Europ.
quadrangularis, Griseb. l. c. = Microcala quadrangularis.
quitensis, Griseb. in Linnaea, xxii. (1849) 38 = Erythraea quitensis.
Roxburghii, Griseb. Gen. et Sp. Gent. 160 = Hoppea dichotoma.
stricta, *Griseb. in Journ. Linn. Soc.* vi. (1862) 143.—Guatemala.
tenella, Rafin. Fl. Tellur. iii. 26 = Gentiana tenella.

CICER, [Tourn.] Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 524.
NOCHOTTA, S. G. Gmel. Reise, iii. 28. t. 3 (1774).
Adonis, Orph. ex Nym. Consp. 200 = ervoides.
anatolicum, *Alef. in Bonplandia*, ix. (1861) 349.—As. Min.
arietinum, *Linn. Sp. Pl.* 788.—Europ. ; Orient ; Ind. or.
cuneatum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 195 = arietinum.
ervoides, Brign. Fasc. Pl. Forojul. 27 = Lens Lenticula.
ervoides, *Fenzl, in Russegg. Reise*, i. 894, in obs.—Syria ; As. Min. ; Syria.
floribundum, *Fenzl, l. c.* 892. t. 9.—Cilicia.
glutinosum, Alef. in Bonplandia, ix. (1861) 349 = anatolicum.
graecum, *Orph. ex Boiss. Diagn.* Ser. II. ii. 48.—Graecia.
grossum, Sallsb. Prod. 340 = arietinum.
Heyneanensis, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xvii. (1842) 231 = soongaricum.
judaicum, Boiss. Diagn. Ser. I. ix. 180 = pinnatifidum.
Lens, Willd. Sp. Pl. iii. 1114 = Lens esculenta.
microphyllum, Royle, Illustr. Bot. Himal. 200 = soongaricum.
minimum, *Boiss. & Hohen. Diagn.* Ser. I. ix. 180.—Persia.
Monbreti, *Jaub. & Spach, in Ann. Sc. Nat.* Ser. II. xvii. (1842) 228.—As. Min.
nigrum, Hort. ex Steud. Nom. ed. II. i. 561 = Lens esculenta.
nunudariaefolium, Lam. Encyc. ii. 2 = Crotalaria bifora.
oxyodon, *Boiss. & Hohen. Diagn.* Ser. I. ix. 129.—Persia.
physodes, Reichb. Fl. Germ. Excurs. 532 = arietinum.
pumpinellifolium, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xvii. (1842) 228 = ervoides.

CICER :—
pinnatifidum, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. II. xvii. (1842) 327.—As. Min.; Mesopotam.
punctatum, Hort. ex Steud. Nom. ed. II. i. 361 = Lens Lenticula.
punctulatum, Hort. ex Ser. in DC. Prod. ii. 366 = Lens esculenta.
pungens, *Boiss. Diagn.* Ser. II. ii. 44.—Afghanist.
sativum, Schkuhr, Handb. ii. 367. t. 202 = arietinum.
soloniense, Schrank, ex G. Don, Gen. Syst. ii. 312 = Lens Lenticula.
songaricum, Jaub. & Spach, Illustr. i. 84. t. 48 B = anatolicum
songaricum, *Steph. ex DC. Mém. Lég.* viii. 349.—Songar.: Himal. bor. occ.
spirocorres, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. II. xvii. (1842) 328.—Persia.
subsphyllum, *Boiss. Diagn.* Ser. I. vi. 44.—Persia.
tragacanthoides, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. II. xvii. (1842) 324.—Persia; Turkestan.

CICERBITA, Wallr. Sched. Crit. 488 (1822) = Lactuca, Tourn. (Compos.).
acuminata, Wallr. l. c. 494 = L. acuminata.
alpina, Wallr. l. c. = L. leucophaea.
borealis, Wallr. l. c. = L. leucophaea.
canadensis, Wallr. l. c. = L. canadensis.
corymbosa, Wallr. l. c. = L. quercina.
elongata, Wallr. l. c. = L. canadensis.
floridana, Wallr. l. c. = L. floridana.
intybacea, Wallr. l. c. = Brachyramphus intybaceus.
leucophaea, Wallr. l. c. = L. leucophaea.
macrophylla, Wallr. l. c. = L. macrophylla.
muralis, Wallr. l. c. 495 = L. muralis.
Plumieri, Kirschl. in Mém. Soc. Hist. Nat. Strassb. iv. (1855) [n. xv.] t. 1 = L. Plumieri.
quercina, Wallr. Sched. Crit. 495 = L. quercina.

CICERCULA, Medic. in Vorles. Churpf. Phys. Ges. i. (1787) 358 = Lathyrus, Tourn. (Legumin.).
alata, Moench, Meth. 163 = L. sativus.
alba, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = L. sativus.
ancepz, Moench, Meth. 163 = L. Cicera.
cornuta, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = L. sativus.
Cicera, Alef. in Bonplandia, xiv. (1861) 147 = L. Cicera.
sativa, Medic. Phil. Bot. i. 203 = L. sativus.

CICERELLA, DC. Prod. ii. 369 (1825) = præc.

CICHORIUM, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITÆ*, Benth. & Hook. f. ii. 506.
ACANTHOPHYTON, Less. Syn. Comp. 128 (1832).
alatum, Hochst. & Steud. ex DC. Prod. vii. 75 = Geigeria alata.
ambiguum, Schult. Obs. 170 = pumilum.
byzantinum, Clem. in Mem. Acc. Sc. Torin. Ser. II. xvi. (1857) 265 = C. Intybus.
caeruleum, Gilib. Fl. Lituan. i. 294 = C. Intybus.
calliosum, *Pomel, Nouv. Mat. Fl. Atl.* 17.—Afr. bor.
calvum, *Sch. Bip. in Ind. Sem. Hort. Beral.* (1859) 7, nomen ; *Schweinf. Beitr. Fl. Aethiop.* 143.—Afr. trop.
Cassia, C. B. Clarke, Comp. Ind. 250 = C. Intybus.
Cicerea, Dum. Fl. Belg. Prod. 64 = C. Intybus.
commune, Pall. Reise, iii. 655 = C. Intybus.
Cunila, Buch.-Ham. in Wall. Cat. n. 3241 = C. Intybus.
crispum, Mill. Gard. Dict. ed. VIII. n. 4 = C. Endivia.
dichotomum, Link, Handb. i. 811, sphalm. = pumilum.
divaricatum, Heldr. ex Nym. Consp. 372 = C. Intybus.
divaricatum, Schousb. Verz. Marok. 197 = pumilum.
Endivia, *Linn. Sp. Pl.* 813.—Europ. ; Orient.
esculentum, Salisb. Prod. 168 = C. Endivia.
glabratum, Presl, Fl. Sic. i. 83 = C. Intybus.
glandulosum, *Boiss. & Heut, Diagn.* Ser. II. iii. 87.—As. Min. ; Mesopotam.
glaucum, *Hoffmgg. & Link, Fl. Port.* ii. 178.—Hispan.
hirsutum, Gren. in Acad. Sc. Besanç. Séanc. Publ. (1866) 187 = C. Intybus.
Intybus, *Linn. Sp. Pl.* 813.—Europ. ; Orient.

CICHORIUM :—
minimum, *Portenschl. Enum. Pl. Dalm.* 16.—Dalmatia.
nanum, Portenschl. ex Nym. Consp. 372 = pumilum
Noëanum, *Boiss. Fl. Orient.* iii. 737.—Mesopot.
officinale, Gueldenst. ex Ledeb. Fl. Ross. ii. 774 = C. Intybus.
perenne, Moënch, Bot. Mat. Med. 138 = C. Intybus.
polystachyum, *Pomel, Nouv. Mat. Fl. Atl.* 18.—Afr. bor.
pumilum, *Jacq. Obs.* iv. 8. t. 80.—Reg. Mediterr.
rigidum, Sallah. Prod. 168 = C. Intybus.
spinosum, *Linn. Sp. Pl.* 813.—Italia ; Graecia.
stylosum, Lam. Fl. Fr. ii. 120 = C. Intybus.

CICLOSPERMUM, Lag. Amen. Acad. Madr. ii. (1821) 101 = Apium, Linn. (Umbellif.).
Ammi, Lag. l. c. = A. Ammi.

CICONIUM, Sweet, Geran. i. p. ix (1823) = Pelargonium, L'Hérit. (Geraniac.).
acetosellum, Hoffmgg. Verz. Pfl. 48 = P. acetosellum.
acetosum, Hoffmgg. l. c. = P. acetosum.
aequisiflorum, Sweet, ex Steud. Nom. ed. II. i. 362 = P. sequiflorum.
albidum, Sweet, l. c. = P. albidum.
amabile, Hoffmgg. Verz. Pfl. 48 = P. amabile.
Annichianum, Sweet, Hort. Brit. ed. I. 85 = P. Bentinkianum.
Blueckerianum, Hoffmgg. Verz. Pfl. 48 = P. Blueckerianum.
bracteosum, Sweet, Hort. Brit. ed. I. 85 = P. bracteosum.
cerinum, Sweet, Geran. ii. t. 176 = P. inquinans.
clarum, Hoffmgg. Verz. Pfl. 48 = P. zonale.
coccinum, Hoffmgg. l. c. = P. hybridum.
coccinisflorum, Hoffmgg. l. c. = P. zonale.
cornutum, Hoffmgg. l. c. = P. zonale.
crenatum, Hoffmgg. l. c. ; Sweet, Hort. Brit. ed. I. 85 = P. crenatum.
denticulatum, Eckl. & Zeyh. Enum. 75 = P. zonale.
Fothergillii, Sweet, Geran. iii. t. 236 = P. Fothergillii.
fulgens, Hoffmgg. Verz. Pfl. 48 = P. formosissimum.
fulgens, Sweet, Hort. Brit. ed. I. Add. 471 = P. fulgens.
glabrifolium, Sweet, Geran. iv. t. 363 = P. glabrifolium.
heterogamum, Sweet, Hort. Brit. ed. I. 84 = P. heterogamum.
hilare, Hoffmgg. Verz. Pfl. 49 = P. hilare.
homogamum, Hoffmgg. l. c. = P. heterogamum.
humile, Sweet, Hort. Brit. ed. I. 84 = P. scandens.
hybridum, Sweet, Geran. i. Ind. = P. hybridum.
inquinans, Hoffmgg. Verz. Pfl. 49 = P. inquinans.
lanuginosum, Hoffmgg. l. c. = P. lanuginosum.
lateritium, Hoffmgg. l. c. = P. malvaefolium.
lavateraefolium, Eckl. & Zeyh. Enum. 75 = P. lavateraefolium.
leptopetalum, Sweet, Hort. Brit. ed. I. 84 = P. leptopetalum.
leucanthum, Hoffmgg. Verz. Pfl. 49 = P. zonale.
lucidum, Hoffmgg. l. c. = P. lucidum.
malvaefolium, Sweet, Hort. Brit. ed. I. 84 = P. malvaefolium.
marginatum, Hoffmgg. Verz. Pfl. 49 = P. marginatum.
micranthum, Sweet, Geran. iii. t. 195 = P. micranthum.
monstrum, Sweet, Geran. i. Ind. = P. monstrum.
multiflorum, Hoffmgg. Verz. Pfl. 49 = P. multiflorum.
orbiculare, Hoffmgg. l. c. = P. obtusifolium.
oxyphyllum, Sweet, Hort. Brit. ed. III. 190 = P. hybridum.
placidum, Hoffmgg. Verz. Pfl. 49 = P. amoenum.
Porphyrio, Hoffmgg. l. c. 50 = P. Porphyrio.
propinquatum, Hoffmgg. l. c. = P. fulminans.
pumilum, Hoffmgg. l. c. = P. scandens.
puniceum, Hoffmgg. l. c. = P. puniceum.
purpurifloram, Hoffmgg. l. c. = P. purpureum.
reticulatum, Sweet, Geran. ii. t. 148 = P. retinervium.
Rumila, Hoffmgg. Verz. Pfl. 50 = P. zonale.
scandens, Hoffmgg. l. c. = P. scandens.
sicelfolium, Hoffmgg. l. c. = P. sicelfolium.
Skelii, Hoffmgg. l. c. = P. Skelii.
stenopetalum, Hoffmgg. l. c. = P. zonale.
umbelliferum, Hoffmgg. l. c. = P. umbelliferum.
urceolatiflorum, Eckl. & Zeyh. Enum. 76 = P. urceolatiflorum.
volutiosum, Hoffmgg. Verz. Pfl. 50 = P. ovale.
zonale, Hoffmgg. l. c. = P. zonale.

CICUTA, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERÆ*, Benth. & Hook. f. i. 899.
KERASKOMION, Rafin. New Fl. Am. iv. 21 (1836).
africana, Lam. Encyc. ii. 4 = Capnophyllum africanum.
Amomum, Crantz, Cl. Umb. Emend. 96 = Sison Amomum.
angustifolia, Kit. in Schult. Oester. Fl. ed. II. i. 515 = virosa.
Bolanderi, *A. Gray, in Proc. Am. Acad.* xi. (1876) 120. —Am. bor. occ.
bulbifera, *Linn. Sp. Pl.* 255.—Am. bor.
californica, *A. Gray, in Proc. Am. Acad.* vii. (1868) 344.—Calif.
capillacea, Nutt. ex DC. Prod. iv. 107 = Discoplenra Nuttallii.
cynapium, Crantz, Cl. Umb. Emend. 98 = Aethusa Cynapium.
Cynapium, Targ. Ist. Bot. ii. 227 = Aethusa Cynapium.
dalmica, Gilib. Fl. Lituan. ii. 96 = virosa.
dahurica, Fisch. Hort. Gorenk. ed. II. (1812) 45, nomen = Apium lineare.
dichotoma, Foir. Encyc. Suppl. ii. 261 = Capnophyllum filchotomum.
latifolia, Crantz, Cl. Umb. Emend. 97 = Sium latifolium.
maculata, Hook. & Arn. Bot. Beech. Voy. 143 = californica ?
maculata, Lam. Fl. Fr. iii. 104 = Conium maculatum.
maculata, *Linn. Sp. Pl.* 256.—Am. bor.
major, Lam. Fl. Fr. iii. 455 = Conium maculatum.
Nimi, Crantz, Cl. Umb. Emend. 97 = Sium Ninsi.
nipponica, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 86.—Japon.
nodiflora, Crantz, Cl. Umb. Emend. 97 = Apium nodiflorum.
occidentalis, *Dougl. ex Hornk. Nom.* ii. 144, nomen.—Am. bor.
officinalis, Crantz, Cl. Umb. Emend. 98 = Conium maculatum.
peroxnana, Walt. Fl. Carol. 116 = Cryptotaenia canadensis !
rigens, Lam. Encyc. ii. 4 (= *Bunium rigens*).—Afr. austr.
sicula, Crantz, Cl. Umb. Emend. 97 = Kundmannia sicula.
sinensis, Zaccagni, in Roem. Collect. 135 = Sellnum Monnieri.
suffruticosa, Poir. Encyc. Suppl. ii. 261 = Capnophyllum Jacquini.
tenuifolia, Schrank, in Denkschr. Akad. Muench. vii. (1818) 58. t. 4. f. 1 = virosa.
trachyptera, *S. Wats. Bibliog. Ind. N. Am. Bot.* 417. —Am. bor. occ.
venenata, Nutt. ex Steud. Nom. ed. II. i. 362 = Archangelica hirsuta.
venenosa, Greene, in Trans. Am. Phil. Soc. iii. (1793) 234 = Archangelica hirsuta.
virosa, *Linn. Sp. Pl.* 255.—Europ. ; As. bor. ; Am. bor.

CICUTARIA, Moench, Meth. Suppl. 22 (1802) (Umbellif.).
dioica, Moench, l. c. = Trinia vulgaris.
verticillata, Moench, l. c. = Molopospermum cicutarium.

CICUTARIA, [Tourn.] Lam. Fl. Fr. iii. 445 (1778) = Cicuta, Tourn. (Umbellif.).
aquatica, Lam. Fl. Fr. iii. 445 = Cicuta virosa.
bulbifera, Lam. Encyc. ii. 3 = Cicuta bulbifera.
maculata, Lam. l. c. 3 = Cicuta maculata.
virosa, Delarb. Fl. Auv. ed. II. 215 = Cicuta virosa.

CIECA, Adans. Fam. ii. 355 (1763) = Julocroton, Mart. (Euphorb.) (J. argentea).

CIECA, Medic. Malv. 97 (1787) = Passiflora, Linn.
angustifolia, M. Roem. Syn. Pepon. 143 = P. suberosa.
apiculiculata, M. Roem. l. c. 148 = P. cærulea.
cærulea, M. Roem. l. c. 148 = P. cærulea.
lanceolatifolia, M. Roem. l. c. 146 = P. lanceolatifolia.
Serratora, M. Roem. l. c. = P. Serratora.
Sibthorpii, M. Roem. l. c. 146 = P. Sibthorpii.
Cavanillesii, M. Roem. l. c. 146 = P. cupraea.
cinerea, M. Roem. l. c. 148 = P. cinerea.

CIECA :—
colubrina, M. Roem. Syn. Pepon. 143 = Passiflora trilobe.
coriacea, M. Roem. l. c. 148 = P. coriacea.
cuprea, M. Roem. l. c. 139 = P. cuprea.
Dictamo, M. Roem. l. c. 146 = P. Dictamo.
difformis, M. Roem. l. c. 140 = P. coriacea.
discolor, M. Roem. l. c. = P. Maximiliana.
flexuosa, M. Roem. l. c. 146 = P. suberosa.
glabrata, M. Roem. l. c. 143 = P. lanata.
globosa, M. Roem. l. c. 144 = P. suberosa.
gracilis, M. Roem. l. c. 141 = P. gracilis.
guyennensis, Klotzsch, in Schomb. Fauna et Fl. Gui. 986, nomen.—Guian.
hederacea, Klotzsch, l. c. 1090, nomen.—Guian.
hederacea, M. Roem. Syn. Pepon. 141 = P. suberosa.
heterophylla, Moench, Meth. Suppl. 101 = P. angustifolia.
limbata, M. Roem. Syn. Pepon. 148 = P. limbata.
littoralis, M. Roem. l. c. 145 = P. suberosa.
maculata, M. Roem. l. c. = P. maculata.
membranacea, M. Roem. l. c. 140 = P. membranacea.
mexicana, M. Roem. l. c. 146 = P. mexicana.
minima, Moench, Meth. Suppl. 102 = P. minima.
misera, M. Roem. Syn. Pepon. 140 = P. misera.
multiflora, M. Roem. l. c. 148 = P. multiflora.
nigra, Medic. Malv. 97 = P. suberosa.
normalis, M. Roem. Syn. Pepon. 144 = P. perfoliata.
oliveæformis, M. Roem. l. c. = P. suberosa.
pallida, M. Roem. l. c. 142 = P. suberosa.
pannosa, M. Roem. l. c. 148 = P. pannosa.
peltata, M. Roem. l. c. 141 = P. suberosa.
porophylla, M. Roem. l. c. 147 = P. porophylla.
pseudo-suberosa, M. Roem. l. c. 146 = P. suberosa.
pubescens, M. Roem. l. c. 141 = P. capsularis.
suberosa, Moench, Meth. Suppl. 102 = P. suberosa.
Surinuca, M. Roem. Syn. Pepon. 141 = P. setacea.
tristosa, M. Roem. l. c. 147 = P. fœtida.
variolata, M. Roem. l. c. 140 = P. variolata.
Velloxii, M. Roem. l. c. 142 = P. Velloxii.
viridis, Medic. Malv. 97 = P. minima.
Warei, M. Roem. Syn. Pepon. 146 = P. Warei.

CIENFUEGIA, Willd. Sp. Pl. iii. 738 (1800) = seq.

CIENFUEGOSIA, Cav. Diss. 174. t. 72. f. 2 (1787) = **Fugosia**, Juss. (Malvac.).
digitata, Cav. l. c. = F. digitata.
heterophylla, Garcke, in Bonplandia, viii. (1860) 150 = F. heterophylla.
Hildebrandtii, Garcke, in Eichl. Jahrb.Bot. Gard. Berl. ii. (1883) 387.—Afr. trop.
phlomidifolia, Garcke, in Bonplandia, viii. (1860) 150 = F. phlomidifolia.
sulphurea, Garcke, l. c. = F. sulfurea.
Welwitzii, Garcke, in Eichl. Jahrb. Bot. Gard. Berl. ii. (1883) 387.—Arabia.

CIENFUGOSIA, DC. Prod. i. 457 (1824) = præc.

CIENKOWSKIA, Regel & Rach, in Ind. Sem. Hort. Petrop. (1858) 48. 'Gen. dub.' Benth. & Hook. f. i. 997.
nitida, Regel & Rach, l. c.—Hab. ?

CIENKOWSKIA, Solms, in Schweinf. Beitr. Fl. Aethiop. t. 1. (1867) = seq.

CIENKOWSKYA, Solms, in Schweinf. Beitr. Fl. Aethiop. 197 (1867) = **Kaempferia**, Linn. (Scitamin.).
æthiopica, Solms, l. c.—Afr. trop.
Kirkii, Hook. f. Bot. Mag. t. 6964.—Afr. trop.

CILIARIA, Haw. Saxifr. Enum. 41 (1821) = **Saxifraga**, Tourn.
aspera, Haw. l. c. 43 = S. aspera.
bronchialis, Haw. l. c. 43 = S. bronchialis.
hyrcina, Haw. l. c. 42 = S. aspera.

CIMBARIA, Hill, Veg. Syst. xx. 43 (1773) = Cymbaria, Linn. (Scrophul.).

CIMICIFUGA, Linn. Fl. Suec. Comach. 21 (1750). *RANUNCULACEÆ*, Benth. & Hook. f. i. 9.
ACTINOSPORA, Turcz. in Flach. & Mey. Ind. Sem. Hort. Petrop. i. 21 (1838).
BOTRYOSTE, Raf. Med. Fl. i. 26 (1828).
CIMICIFUGA :—
MACROTYS, Rafin. in Med. Repos. N. York. II. v. (1808) 359.
MACROTYS, DC. Syst. i. 288 (1818).
MEGOTYS, Rafin. in Am. Monthly Mag. (1818) 267.
PITYROSPERMA, Sieb. & Zucc. in Abh. Acad. Muench. iii. (1846) 784.
americana, *Michx. Fl. Bor. Am.* i. 316.—Am. bor.
americana, Muhl. Cat. ed. II. 54 = cordifolia.
arisonica, *S. Wats. in Proc. Am. Acad.* xx. (1885) 352. —Arizona.
biternata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 9.— Japon.
cordifolia, *Pursh, Fl. Am. Sept.* ii. 373.—Am. bor.
elata, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 36.—Am. bor. occ.
fœtida, *Linn. Syst. ed.* XII. 659.—Europ. ; Sibir.
fœtida, Pursh, Fl. Am. Sept. ii. 373 = elata.
frigida, Royle, Illustr. Bot. Himal. 57. t. 14 = fœtida.
japonica, Spreng. Syst. ii. 626.—Japon.
laciniata, *S. Wats. in Proc. Am. Acad.* xx. (1885) 352. —Am. bor. occ.
obtusiloba, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 9.— Japon.
palmata, Michx. Fl. Bor. Am. i. 316 = Trautvetteria palmata.
podocarpa, Ell. Sketch, ii. 16 = americana.
racemosa, *Nutt. Gen. Am.* ii. 15.—Am. bor.
Serpentaria, Pursh, Fl. Am. Sept. ii. 373 = racemosa.
simplex, Wormsk. ex DC. Prod. i. 64 = fœtida.

CIMINALIS, Adans. Fam. ii. 504 (1763) = **Gentiana**, Tourn.
acaulis, Borckh. in Roem. Arch. i. 1. 26 = G. acaulis.
algida, Bercht. & Presl, Rostl. i. Gentian. 10 = G. frigida.
decumbens, Bercht. & Presl, l. c. = G. decumbens.
frigida, Bercht. & Presl, l. c. = G. frigida.
glauca, Bercht. & Presl, l. c. 11 = G. glauca.
grandiflora, Mayrh. in Schrank, Fl. Monac. t. 37 = G. acaulis.
linearis, Bercht. & Presl, Rostl. i. Gentian. 11 = G. linearis.
longiflora, Moench, Meth. 514 = G. acaulis.
ochroleuca, Bercht. & Presl, Rostl. i. Gentian. 11 = G. ochroleuca.
Pneumonanthe, Borckh. in Roem. Arch. i. 1. 26 = G. Pneumonanthe.
pseudopneumonanthe, Bercht. & Presl, Rostl. i. Gentian. 11 = G. Pneumonanthe.
saponaria, Bercht. & Presl, l. c. = G. saponaria.
saxosa, Bercht. & Presl, l. c. = G. montana.
septemfida, Bercht. & Presl, l. c. 10 = G. septemfida.
vulgaris, Bercht. & Presl, l. c. = G. Pneumonanthe.

CIMINALIS, Rafin. Fl. Tellur. iii. 19 (1836) = **Voyria**, Aubl. (Gentian.).
alpina, *angustifolia*, *grandiflora*, Rafin. l. c. = V. aphylla.

CINARA, [Tourn.] Linn. Syst. ed. I. (1735) = **Cynara**, Linn. (Compos.).

CINCHONA, Linn. Gen. ed. II. 526 (1742). *RUBIACEÆ*, Benth. & Hook. f. ii. 23.
CALISAYA, Hort. ex Pav. Nuev. Quinol. sub t. 5. p. 6 (1862).
CASCARILLA, Ruiz, ex Steud. Nom. ed. II. i. 165 (1840).
KINKINA, Adans. Fam. ii. 147 (1763).
QUINQUINA, Condam. in 'Act. Acad. Par. (1738) 226.'
acadensium, Gulbourt, Drogues Simples, iii. 98 = officinalis.
acuminata, Polr. Encyc. Suppl. iv. 640 = Cosmibuena acuminata.
acutifolia, Ruiz & Pav. Fl. Per. iii. 1. t. 225 = Cascarilla acutifolia.
affinis, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 8.— Peruv.
afrodata, Willem. in Usteri, Ann. Bot. xviii. (1796) 16 = Danais fragrans.
amygdalifolia, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 6.—Bolivia.
angustifolia, Ruiz, Quin. Suppl. 14. n. 17 = lancifolia.
angustifolia, Sw. Prod. Veg. Ind. Occ. 41 = Exostemma angustifolium.
aspedifolia, *Wedd. in. Ann. Sc. Nat.* Sér. III. x. (1848) 7.—Bolivia.
CINCHONA :—
australis, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 68 = Exostemma australe.
australis, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 7.—Bolivia.
barbacoensis, *Karst. in C. Koch & Fint. Wochensch.* ii. (1859) 31.—N. Granat.
Bergeniana, A. Braun, ex Mart. Reise Bras. iii. 1286 = Remijia Bergeniana.
bicolor, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 68 = Exostemma bicolor.
bogotensis, *Karst. in C. Koch & Fint. Wochenschr.* ii. (1859) 30.—N. Granat.
boliviana, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 7 = C. Calisaya.
Bonplandiana, *Klotzsch, in Hayne, Arzneigew.* vii. t. 37.—Peruv.
brachycarpa, Sw. Prod. Veg. Ind. Occ. 41 = Exostemma brachycarpum.
brasiliensis, Hoffngg. ex Humb. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 219 = Machaonia brasiliensis.
caducifolora, Humb. & Bonpl. Pl. Aequin. i. 168, in obs. = Cascarilla caducifolora.
Calisaya, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 6. —Bolivia.
calopters, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. (1868-69) 278.—Peruv.
Candollii, G. Don, Gen. Syst. iii. 479 = Remijia paniculata.
canescens, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 68 = Exostemma canescens.
capensis, Burm. ex DC. Prod. iv. 369 = Burchellia capensis.
capitata, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 68 = Exostemma capitatum.
carabayensis, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 9.—Peru.
caribæa, Jacq. Enum. Pl. Carib. 16 = Exostemma caribæum.
caribbæana, W. Wright, in Phil. Trans. lxvii. (1778) 506 = prae.
carolinians, Poir. Encyc. vi. 40 = Pinckneya pubens.
Carua, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 275.— Bolivia.
Cattukambar, Koen. ex Retz, in Vet. Acad. Handl. Stockh. (1814) 33. t. 3 = Uncaria acida.
cava, Pav. ex Lamb. Illustr. Cinch. 8 = Cascarilla Pavonii.
Chahuargnero, Pav. ex DC. Prod. iv. 352 = officinalis.
China, DC. Prod. iv. 356, sphalm. = Cosmibuena obtusifolia.
chlorrhiza, Bory, ex DC. l. c. 361 = Danais rotundifolia.
Chomelana, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 9.— Bolivia.
cladorrhiza, Bory, ex Wedd. Hist. Nat. Quinq. 72 = Danais rotundifolia.
coccinea, Pav. ex DC. Prod. iv. 353 = macrocalyx.
colorata, Laubert, in 'Bull. de Pharm. (1810) 294' = officinalis.
Condaminea, Humb. & Bonpl. Pl. Aequin. i. 33 = officinalis.
conglomerata, *Pav. Nuov. Quinol.* t. 15.—Ecuador.
cordifolia, *Mutis, ex Humb. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 117.—N. Granat. ; Peruv.
coriacea, Polr. Encyc. vi. 38 = Exostemma coriaceum.
coronulata, *Miq. in Journ. Bot. Néerl.* i. (1861) 140.— Peruv.
corymbifera, Forst. in Act. Nov. Soc. Sc. Ups. iii. (1780) 176 = Exostemma corymbiferum.
corymbosa, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 68 = Exostemma corymbosum.
corymbosa, Karst. Fl. Columb. i. 19. t. 10 = pitayensis.
crassifolia, Pav. ex DC. Bibl. Univ. xli. (1829) 150 = Cascarilla calyptrata.
crispa, Tafalla, ex Howard, in Pav. Nouv. Quinol. t. 2 = officinalis.
cucumæfolia, Pav. ex Lamb. Illustr. Cinch. 2 = lancifolia.
cujabensis, Manso, ex Klotzsch, in Hayne, Arzneigew. xiv. sub t. 15 = Remijia cujabensis.
cuspidata, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 68 = Exostemma cuspidatum.
decurrentifolia, Pav. Nouv. Quinol. t. 23 = purpurascens.
Delondriana, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 7 = scrobiculata.

CINCHONA:—
densifora, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Remijia densiflora.
dichotoma, Ruiz & Pav. Fl. Per. ii. 58. t. 197 = Ladenbergia dichotoma.
discolor, Klotzsch, in Hayne, Arzneigew. xiv. sub t. 14 = pubescens.
disstmiliflora, Mutis, ex Humb. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 120 = Exostemma dissimiliflorum.
elliptica, *Wedd. in Ann. Sc. Nat.* Sér. V. xii. (1869) 60.—Peruv.
erythrantha, *Pav. Nuev. Quinol.* t. 12.—Ecuador.
erythrodermis, *Wedd. in Ann. Sc. Nat.* Ser. V. xi. (1869) 362.—Peruv.
euneura, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 265.—Bras.
excelsa, Roxb. Fl. Corom. ii. 3. t. 106 = Hymenodictyon excelsum.
ferruginea, St. Hil. in Mém. Mus. Par. ix. (1822) 314 = Remijia ferruginea.
firmula, Mart. ex Klotzsch, in Hayne, Arzneigew. xiv. sub t. 15 = Remijia firmula.
flaccida, Spreng. Syst. iv. Cur. Post. 78 = Hymenodictyon flaccidum.
floribunda, Sw. Prod. Veg. Ind. Occ. 41 = Exostemma floribundum.
Forbesiana, Howard, Quinol. E. Ind. Plant. 37* = lancifolia.
formosa, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma formosum.
fusca, Ruiz, ex Vitm. Summa, Suppl. i. 262 (= *Lasionema roseum*).—Am. trop.
glabra, Ruiz, Quinol. 54 = lancifolia.
glandulifera, *Ruiz & Pav. Fl. Per.* iii. 1. t. 224.—Peruv.
glabrifera, Pav. ex DC. Prod. iv. 345 = Naucles Cinchonae.
govanea, *Miq. in Journ. Bot. Néerl.* i. (1861) 142.—Peruv.
grandiflora, Ruiz & Pav. Fl. Per. ii. 54 = Coamibuena obtusifolia.
grandifolia, Poir. Encyc. vi. 36 = Cascarilla magnifolia.
gratissima, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 154 = Lucilia gratissima.
Haenkeana, Bartl. ex DC. Prod. iv. 530 = Pallcoarea Haenkeana.
Hamkeriana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 266.—Peruv.
Henleana, *Karst. in C. Koch & Fint. Wochenschr.* ii. (1859) 31.—Venezuela.
herbacea, Linn. Mant. ii. 238 = Exostemma caribaeum ?
heterocarpa, *Karst. Fl. Columb.* i. 11. t. 6.—N. Granat.
heterophylla, Pav. ex DC. Prod. iv. 353 = cordifolia.
heterophylla, Pav. Nuev. Quinol. t. 18 = macrocalyx.
hexandra, G. Don, Gen. Syst. iii. 478 = Cascarilla hexandra.
hirsuta, *Ruiz & Pav. Fl. Per.* ii. 51. t. 192.—Peruv.
Howardiana, *O. Kuntze, Monog. Cinch.* (1878). 30.—Peruv.
Humboldtiana, *Lamb. Illustr. Cinch.* 7.—Peruv.
Humboldtiana, Roem. & Schult. Syst. v. 18 (= *Lasionema Humboldtianum*).—Peruv.
Humboldtiana, Wedd. Hist. Nat. Quinq. 67 = villosa.
jamaicensis, W. Wright, in Phil. Trans. lxvii. (1778) 506 = Exostemma caribaeum.
Josephiana, *Wedd. in Ann. Sc. Nat.* Sér. V. xii. (1869) 58.—Bolivia.
Kattuhambar, Koen. ex DC. Prod. iv. 347 = Uncaria acida.
lancifera, Ruiz & Pav. ex Fée, Ess. Crypt. Ecorces Exot. Off. 8; Tafalla, ex Wedd. Hist. Nat. Quinq. 72 = Condamines tinctoria ?
Lamberciana, A. Braun, ex Mart. Reise Bras. iii. 1286 = Cascarilla Lambertiana.
lancolaeta, Ruiz & Pav. Fl. Per. ii. 51 = officinalis.
lancifolia, *Mutis, in Periodico de Santa Fé* (1793) 465, nomen.—Peruv. ; N. Granat.
Lechleriana, *Schlecht. in Linnaea*, xxvi. (1853–55) 738.—Peruv.
Ledgeriana, *Moens, ex Trimen, in Journ. Bot.* xix. (1881) 338.—Bolivia.
legitima, *Ruiz, ex Laubert, in Bull. de Pharm.* (1810) 193.—Peruv. ?

CINCHONA:—
leucocalyx, *Brign. in Mem. Soc. Ital. Mod.* Ser. II. i. (1862) 64.—Bras.
linaeata, Vahl, in Skriv. Naturh.-Selsk. i. (1790) 22 = Exostemma linaetum.
longiflora, Lamb. Desc. Cinch. 38. t. 12 = Exostemma longiflorum.
longiflora, Mutis, ex Steud. Nom. ed. I. 196 = Cosmibuena obtusifolia.
longifolia, Mutis, ex Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 62 = praec. ?
Luciana, Vitm. Summa, Suppl. i. 264 = Exostemma floribundum.
lucumaefolia, Pav. ex Lindl. Fl. Med. 416 = macrocalyx.
lutea, Pav. Nuev. Quinol. t. 14 = cordifolia.
lutescens, Ruiz, ex Vitm. Summa, Suppl. i. 262 = Cascarilla magnifolia.
macrocalyx, *Pav. ex DC. in Bibl. Univ. Sc. et Arts*, xli. (1839) 156.—Am. austr.
macrocarpa, Vahl, in Skriv. Naturh.-Selsk. i. (1790) 20 = Cascarilla macrocarpa.
Macronemia, A. Braun, ex Mart. Reise Bras. iii. 1286 = Remijia Macronemia.
macrophylla, *Karst. in Linnaea*, xxx. (1859–60) 151.—N. Granat.
magnifolora, Pav. ex DC. Prod. iv. 354 = Cascarilla macrocarpa.
magnifolia, Humb. & Bonpl. Pl. Aequin. i. 136 = Cascarilla caducifolia.
magnifolia, Ruiz & Pav. Fl. Per. ii. 53. t. 196 = Cascarilla magnifolia.
mayanensis, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma mayanense.
micrantha, *Ruiz & Pav. Fl. Per.* ii. 52. t. 194.—Peruv.
microphylla, Mutis, ex Lamb. Illustr. Cinch. 9 = Mutisii.
microphylla, *Pav. Nuev. Quinol.* t. 17.—Peruv.
montana, Badier, in Journ. Phys. (1789) 129 = Exostemma floribundum.
Morada, Ruiz, Quinol. 57. n. 5 = purpurea.
Moritziana, *Karst. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 275.—Peruv.
Mutisii, *Lamb. Illustr. Cinch.* 9.—Peruv.
muzonensis, Goudot, in Phil. Mag. (1838) 182 = Cascarilla muzonensis.
myrtifolia, Stokes, Bot. Mat. Med. i. 359 = Exostemma caribaeum.
nitida, Benth. Pl. Hartw. 109 = Cascarilla nitida.
nitida, *Ruiz & Pav. Fl. Per.* ii. 50.—Peruv.
oblongifolia, Lamb. Illustr. Cinch. 13 = Beena Riveroi.
oblongifolia, Mutis, in Periodico de Santa Fé (1793) 466, nomen ; Humb. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 118 = Cascarilla oblongifolia.
obovata, Pav. Nuev. Quinol. sub t. 18. et t. 24 = pubescens.
obovata, Spreng. Syst. iv. Cur. Post. 78 = Hymenodictyon obovatum.
obtusifolia, D. Dietr. Syn. Pl. i. 721 = Cosmibuena obtusifolia.
obtusifolia, Pav. ex DC. Prod. iv. 353 = macrocalyx.
officinalis, Linn. Mant. ii. 238 = cordifolia.
officinalis, *Linn. Syst. ed.* X. 939.—Peruv.
ovalifolia, Humb. & Bonpl. Pl. Aequin. i. 65 (= *Lasionema Humboldtianum*).—Peruv.
ovalifolia, Mutis, ex Humb. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 118 = Cascarilla macrocarpa.
ovata, *Ruiz & Pav. Fl. Per.* ii. 52. t. 195.—Peruv.
Pahudiana, *Howard, in Pav. Nuev. Quinol.* t. 21.—Am. austr.
pallescens, Ruiz, ex DC. Prod. iv. 358 = ovata.
Palton, Pav. Nuev. Quinol. t. 13 = macrocalyx.
paniculata, D. Dietr. Syn. Pl. i. 721 = Remijia paniculata.
parabolica, *Pav. Nuev. Quinol.* t. 16.—Peruv.
parviflora, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma parviflorum.
parviflora, Pohl, Encyc. vi. 36 = micrantha.
pascifices, *Tafalla, ex Steud. Nom.* ed. II. i. 263.—Am. austr.
Pavoniana, *O. Kuntze, Monog. Cinch.* (1878) 26.—Peruv. ; Bolivia.
Pavonii, Lamb. Illustr. Cinch. 8 = Cascarilla Pavonii.
pedunculata, *Karst. in C. Koch & Fint. Wochenschr.* ii. (1859) 30.—N. Granat.
Pelaiba, *Pav. ex DC. Bibl. Univ.* xli. (1839) 153.—Am. austr.

CINCHONA:—
Pelleteriana, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 8 = pubescens.
peruviana, *Howard, in Pav. Nuev. Quinol.* t. 27.—Andes.
peruviana, Poir. Encyc. Suppl. iv. 640 = Exostemma peruvianum.
philippica, *Cav. Ic.* iv. 16 = Exostemma philippicum.
pitayensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xi. (1849) 269 ; et V. xii. (1869) 37.—N. Granat.
platyphylla, *Wedd. l. c.* Sér. V. xi. (1869) 362 ; xii. (1869) 78.—N. Granat.
prismatostylis, *Karst. Fl. Columb.* i. 13. t. 7.—N. Granat.
pubescens, Hort. Mack. ex Walp. Rep. ii. 943 = Rhysocarpus pubescens.
pubescens, Lamb. Illustr. Cinch. 6 = ovata.
pubescens, Vahl, in Skriv. Naturh.-Selsk. i. (1790) 19 = cordifolia.
purpurascens, *Wedd. in Ann. Sc. Nat.* Ser. III. x. (1848) 8.—Bolivia.
purpurea, *Ruiz & Pav. Fl. Per.* ii. 52. t. 193.—Peruv.
quercifolia, Pav. ex Lamb. Illustr. Cinch. 9 = Mutisii.
Quina, Ruiz, in Ruiz & Pav. Fl. Per. iii. 4 = Cosmibuena obtusifolia.
racemosa, Schrank, ex Steud. Nom. ed. II. i. 263 = Exostemma caribaeum.
Reicheliana, Howard, in Pav. Nuev. Quinol. t. 5 = micrantha.
Remijiana, St. Hil. Pl. Us. Bras. sub t. 2 = Remijia Hilarii.
Riedeliana, Casar. in Atti Scia Rium. Scienz. Ital. (1841) 513 = Cascarilla Riedeliana.
robusta, *Howard, in Journ. Linn. Soc.* xx. (1884) 327, nomen.—Hab. ?
Roraimae, Benth. in Hook. Journ. Bot. iii. (1841) 214 = Cascarilla Roraimae.
rosea, Ruiz & Pav. Fl. Per. ii. 54. t. 199 (= *Lasionema roseum*).—Peruv.
roseiecta, *Howard, ex Wedd. in Ann. Sc. Nat.* Sér. V. xi. (1869) 362.—N. Granat.
rotundifolia, Pav. ex Lamb. Illustr. Cinch. 5 = cordifolia.
rubicunda, *Auct. ex Steud. Nom.* ed. II. i. 263.—Am. austr.
rubicunda, Tafalla, ex Wedd. in Ann. Sc. Nat. Sér. V. xi. (1869) 362 = ovata.
rufinervia, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 8.—Peruv.
rugosa, Pav. ex DC. Prod. iv. 353 = cordifolia.
Sanctae-Luciae, Davids, ex Roem. & Schult. Syst. v. 19 = Exostemma floribundum.
scrobiculata, *Humb. & Bonpl. Pl. Aequin.* i. 165. t. 67.—Peruv.
sessilis, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma sessilanum.
spinosa, Vavass. in Journ. Phys. (Oct. 1790) 273 = Exostemma Vavassorii.
stenocarpa, Lamb. Illustr. Cinch. 13 = Cascarilla stenocarpa.
stipea, Pav. ex Lindl. Fl. Med. 416 = lucumaefolia.
subcordata, Pav. Nuev. Quinol. ed. Howard, ad calcem = cordifolia.
suberosa, Pav. l. c. = macrocalyx.
subsessilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 272.—Peruv.
succirubra, *Pav. ex Klotzsch, in Abh. Akad. Berl.* 1857 (1858) 90.—Peruv.
Tarontaron, *Pav. Nuev. Quinol. ed. Howard*, ad calcem.—Peruv.
tenuiflora, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Remijia tenuiflora.
tenuis, Ruiz, ex DC. Prod. iv. 353 = cordifolia.
thyrsiflora, Roxb. Hort. Beng. 15 ; Fl. Ind. i. 530 = Hymenodictyon excelsum.
timorensa, Span. in Linnaea, xv. (1841) 315 = Hymenodictyon timoriense.
Timoriensis, Walp. Rep. ii. 500 = praec.
Trianae, Karst. in C. Koch & Fint. Wochenschr. ii. (1859) 31 = pitayensis.
triflora, W. Wright, in Lond. Med. Journ. viii. (1787) 340 = Exostemma triflorum.
tucujensis, *Karst. Fl. Colomb.* i. 17. t. 9.—N. Granat.
umbellulifera, *Pav. Nuev. Quinol.* t. 23.—Peruv.
undata, Karst. ex Wedd. in Journ. Linn. Soc. xi. (1869) 387 = Cascarilla undata.
undulata, Pav. Nuev. Quinol. t. 7. t. 8 = glandulifera.
Uritusinga, Pav. ex DC. Prod. iv. 356 = macrocalyx.

CINCHONA:—
Uritusinga, Pav. ex Howard, Nuev. Quinol. t. 19 = officinalis.
vanillodora, *Auct. ex Steud. Nom.* ed. II. i. 368.—Am. austr.
Vellozii, St. Hil. Pl. Us. Bras. sub t. 2 = Remijia Vellozii.
villosa, Pav. ex Lamb. Illustr. Cinch. 7 = Humboldtiana.
violacea, *Pav. Nuev. Quinol. ed. Howard, ad calcem.* —Ecuador.
viridiflora, Pav. l. c. = pubescens.
Virtusinga, Pav. ex DC. Prod. iv. 352 = C. Condamines.
Weddelliana, *O. Kuntze, Monog. Cinch.* (1878) 29.—Bolivia.

CINCLIA, Hoffmgg. Prelavers. (1841).—*Cf.* Linnæa, xvi. (1842) Litt. 367 = **Ceropegia**, Linn. (Asclepiad.).

CINCLIDOCARPUS, Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie, iii. (1846) 81 = **Caesalpinia**, Linn. (Legumin.).
nitidus, Zoll. & Mor. l. c. 82 = Caesalpinia Cinclidocarpa.

CINERARIA, Linn. Sp. Pl. ed. II. 1242 (1763). *COMPOSITÆ*, Benth. & Hook. f. ii. 445.
XENOCARPUS, Cass. in Dict. Sc. Nat. liv. 108 (1829).
abrotanifolia, *Berg. Desc. Pl. Cap.* 292.—Afr. austr.
abyssinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 153.—Afr. trop.
acanthifolia, Reichb. Hort. Bot. t. 24 = Senecio Cineraria.
acutangula, Bertol. Fl. Guatimal. 35 = Senecio acutangulus.
Aitoniana, Spreng. Fl. Hal. Mant. 51 = canescens.
alata, Linn. f. Suppl. 374 = Othonna alata.
alchemilloides, *DC. Prod.* vi. 307.—Afr. austr.
allariifolia, *Desf. Cat. Hort. Par.* ed. III. 403.—Hab.?
alpestris, Hoppe, Taschenb. (1806) 130 = Senecio alpestris.
alpestris, Pall. Reise, ii. 38 = Senecio campestris.
alpestris, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 111 = Senecio brachychætus.
alpina, All. Fl. Pedem. i. 208. t. 38 = Senecio auranticus.
alpina, Huds. Fl. Angl. ed. II. 370 = Senecio campestris.
alpina, Linn. Sp. Pl. ed. II. 1243 = Senecio alpinus.
altaica, C. A. Mey. ex DC. Prod. vi. 315 = Senecio altaicus.
ambigua, Bivona, Manip. iii. 5 = Senecio ambiguus.
ambracea, Hornem. Hort. Hafn. Add. 972 = Senecio umbraceus.
amelioides, Linn. Sp. Pl. ed. II. 1245 = Aster rotundifolius.
americana, Linn. f. Suppl. 373 = Senecio denticulatus.
amœna, Sallsb. Prod. 200 = Aster rotundifolius.
angulata, Alam. ex DC. Prod. vi. 431. n. 544 = Senecio sublobatus.
angulata, Mair, ex DC. l. c. n. 547 = Senecio angulifolius.
anguiosa, Lam. Encyc. ii. 6 = humifusa, gelfolia.
anguiosa, E. Mey. ex DC. Prod. vi. 337 = Senecio longiflorus.
angustata, Schur, Enum. Pl. Transs. 346 = *transsilvanica*.
angustifolia, H. B. & K. Nov. Gen. et Sp. iv. 189 = Senecio salignus.
anomala, Sibth. & Sm. Fl. Græc. Prod. ii. 179; Fl. Græc. Ic. 56 = Senecio Othonnæ.
anthemoides, Lam. Encyc. ii. 9 = Senecio gallicus.
anthemoides, Spreng. ex DC. Prod. vi. 365 = Senecio ramosus.
appendiculata, Poir. Encyc. Suppl. ii. 363 = Senecio appendiculatus.
appendiculata, Steud. Nom. ed. II. i. 363 = Senecio pomiliolius.
arachnoidea, Reichb. in Mossel. Handb. ed. II. ii. 1561 = Senecio lanatus.
arachnoidea, Turcz. ex Ledeb. Fl. Ross. ii. 632 = Senecio frigidus.
arctotidea, *DC. Prod.* vi. 306.—Afr. austr.
arctotidea, Drège, ex Harv. & Sond. Fl. Cap. iii. 309 = axillis.
arenosa, Remy, in Bull. Soc. Bot. Fr. xxvii. (1880) 99 = Senecio campestris.

CINERARIA:—
aspera, *Thunb. Prod. Pl. Cap.* 155.—Afr. austr.
atriplicifolia, *DC. Prod.* vi. 308.—Afr. austr.
atropurpurea, Ledeb. in Mém. Acad. Pétersb. v. (1814) 574 = Senecio frigidus.
Auchert, *C. Koch, in Linnæa*, xxiv. (1851) 358.—Orient.
aurantiaca, Bess. Enum. Pl. Volh. 38 = Senecio campestris.
aurantiaca, Hoppe, ex Willd. Sp. Pl. iii. 2081 = Senecio aurantiacus.
aurea, Linn. Sp. Pl. ed. II. 1244 = Senecio auratus.
auriculata, Ledeb. Fl. Alt. iv. 105 = Senecio racemosus.
auritæ, Andr. Bot. Rep. i. 24. t. 24 = Doronicum crnentum.
auritæ, *L'Hérit. Sert. Angl.* 26 = Senecio maderensis.
Balbisiana, *Bertol. Fl. Ital.* ix. 290.—Italia.
Balsamitæ, Lam. Encyc. ii. 7 = Senecio aureus.
balsamitæ, Spreng. Syst. iii. 553 = Senecio balbysius.
Bergerianæ, Spreng. l. c. 551 = Aster Bergerianus.
bicolor, Willd. Sp. Pl. iii. 2085 = Senecio bicolor.
bipinnatæ, Willd. l. c. 2075 = Senecio pinnatifidus.
bracteatæ, Chr. Sm. ex Link, in Buch, Beschr. Canar. Ins. 148 = Senecio malvæfolius.
brasilisensis, Spreng. Neue Entdeck. ii. 149 = Senecio brasiliensis.
cacaliiformis, Lam. Fl. Fr. ii. 124 = Senecio Ligularia.
cacalioides, Linn. f. Suppl. 374 = Othonna carnosa.
Caladonicæ, Spreng. Fl. Hal. Mant. 50 = Brachyglottis repanda.
calvescens, Nym. Syll. Suppl. 1 = Senecio Cineraria.
campestris, DC. Fl. Fr. iv. 164 = Senecio campestris.
campestris, Pellm. ex Ledeb. Fl. Ross. ii. 647 = Senecio aurantiacus.
campestris, Zett. ex Willk. & Lange, Prod. Fl. Hisp. ii. 110 = Senecio pyrenaicus.
canadensis, Linn. Sp. Pl. ed. II. 1244 = Senecio Cineraria.
canadensis, Walt. Fl. Carol. 207 = Erechtites hieracifolia?
candida, J. & C. Presl, Delic. Prag. (1822) 95 = Senecio candidus.
canescens, Spreng. Syst. iii. 549 = Erigeron canescens.
canescens, *Wendl. Obs. ex Link, Enum. Hort. Berol.* ii. 332.—Afr. austr.
cantabrica, Nym. Syll. 2 = Senecio brachychætus.
capillacea, Linn. f. Suppl. 375 = Steirodiscus capillaceus.
capitata, Ledeb. Fl. Alt. iv. 104 = Senecio campestris.
capitata, Wahlenb. Fl. Carp. 271 = Senecio auranticus.
carolinensis, Walt. Fl. Carol. 207 (sp. indeterminab.).—Am. bor.
caspica, Pers. Syn. ii. 439 = Senecio racemosus.
caucasica, Bieb. Fl. Taur. Cauc. ii. 313 = Senecio caucasicus.
causcica, Hoffm. Hort. Mosq. (1808) n. 598 = Senecio platyphyllus.
caucasica, Schlecht. ex Ledeb. Fl. Ross. ii. 558 = Senecio renifolius, *Sch. Bip.*
cerathophila, Tenore, Ind. Sem. Hort. Neap. 1825 = Senecio Cineraria.
cernua, Thore, Chl. Land. 344 = Arnica montana.
cernua, Walt. ex DC. Prod. vi. 314 = Senecio arnicoides.
chamasdrifolia, *Lam. Encyc.* ii. 9.—Afr. austr.
chinensis, Spreng. Syst. iii. 549 = Senecio scandens.
Cineta, Willd. in Webb & Berth. Phyt. Canar. iii. 339 = Doronicum papyraceum.
Clusiana, Host, Fl. Austr. ii. 423 = Senecio abrotifolius.
Cobaea, Nym. Consp. 352 = Senecio brachychætus.
collina, Sallsb. Prod. 200 = Senecio campestris.
conformis, Rochel, Pl. Banat. Rar. Ind. = Senecio auranticus, brachychætus.
congesta, R. Br. in Parry, 1st Voy. App. 279 = Senecio palustris.
cordata, Jacq. Fl. Austr. ii. t. 176 = seq.
cordifolia, Gouan, Illustr. 69 = Senecio alpinus.
cordifolia, Jacq. Fl. Austr. t. 177 = Senecio subalpinus.
cordifolia, Lapeyr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 115 = Senecio Doronicum.
coronata, Thunb. Prod. Pl. Cap. 154 = Senecio lasiochæta.
coronata, Zeyh. ex DC. Prod. vi. 488 = Senecio jacobæa.
corymbosa, Moench, Meth. 586 = Erigeron annuus.

CINERARIA:—
crassiflora, Poir. Encyc. Suppl. ii. 367 = Senecio crassiflorus.
crassifolia, Host, [Fl. Austr. ii. 482] ex DC. Prod. vi. 350.—Europ.
crassifolia, Kit. in Schult. Oestr. Fl. ed. II. ii. 814 = Senecio alpestris.
crenata, Spreng. ex DC. Prod. vi. 388 = Senecio crocus.
crispa, Jacq. Fl. Austr. ii. 48 = Senecio crispatus.
crocea, Trat. Arch. i. 26 = Senecio crispatus.
cruenta, Mass. ex L'Hérit. Sert. Angl. 26 = Doronicum cruentum.
cunaata, E. Mey. ex DC. Prod. vi. 439 = Senecio diversifolius, *Harv.*
cymbalariifolia, Linn. Amœn. Acad. vi. 106 = Senecio cymbalariifolius.
dealbata, Spreng. ex DC. Prod. vi. 369 = Senecio Kunthianus.
decipiens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 312.—Afr. austr.
decipiens, Spreng. ex DC. Prod. vi. 366 = Senecio chrysanthemoides.
degenerans, Kit. ex Kanitz, in Verh. Zool.-Bot. Ges. Wien, xiii. (1863) 623 = Senecio alpestris ? deltoides, *Sond. in Linnæa*, xxiii. (1850) 68.—Afr. austr.
denticulata, DC. Prod. vi. 416, in syn. = Senecio denticulatus.
denticulata, Linn. f. Suppl. 376 (= *Brachyrhynchos cupatorioides*).—Afr. austr.
difformis, Rochel, Pl. Banat. Rar. Ind. = Senecio alpinus.
discoides, *Kit. in Linnæa*, xxxii. (1863) 389.—Europ.
discolor, Sw. Prod. Veg. Ind. Occ. 114 = Senecio discolor.
distans, Kunze, in Linnæa, xvii. (1843) 571 = Senecio gelfolius.
divaricata, Spreng. ex DC. Prod. vi. 370 = Senecio scandens.
Doria, Spreng. ex DC. l. c. 364 = Senecio Wallichii.
draconculoides, Moq. & Sesse, ex DC. l. c. 480 = Senecio salignus.
Dregeana, *DC. Prod.* vi. 305.—Afr. austr.
dubia, Spreng. Syst. iii. 550 = Senecio tomentosus.
elata, Steud. Nom. ed. II. i. 364 = Senecio Ligularia.
elatior, Bouché, in Link, Enum. Hort. Berol. ii. 333 = Senecio multiflorus.
elodes, Nym. Syll. 2 = Senecio elodes.
elongata, Linn. f. Suppl. 374 = Senecio diversifolium, *Harv.*
erodioides, *DC. Prod.* vi. 307.—Afr. austr.
erosa, Harv. in Harv. & Sond. Fl. Cap. iii. 309 = oxyodonta.
erosa, *Willd. Sp. Pl.* iii. 2078.—Afr. austr.
exilis, *DC. Prod.* vi. 305.—Afr. austr.
Farfara, Bernh. Syst. Verz. Erf. 146 = Tussilago Farfara.
filifolia, Thunb. Prod. Pl. Cap. 154 = Senecio longiflorus.
Fischeri, Ledeb. Ind. Sem. Hort. Dorpat. (1820) 17 = Senecio Ligularia.
flammea, Turcz. ex DC. Prod. vi. 362 = Senecio flammeus.
Forsteri, Spreng. Pugill. i. 58 = Brachyglottis Forsteri.
fragrans, Hoffmgg. ex Steud. Nom. ed. II. i. 364 = Erigeron fragrans.
frigida, Richards. in Frankl. Narr. 1st Journ. App. 748 = Senecio frigidus.
fulva, Sm. in Mém. Soc. Nat. Mosc. iv. (1816) 84 = Senecio auranticus.
Fuzoni, Griseb. & Schenk, in Wiegm. Arch. xviii. (1852) 342 = Senecio campestris.
gelfolia, *Linn. Sp. Pl.* ed. II. 1242.—Afr. austr.
gerifolia, C. A. Mey. ex Steud. Nom. ed. II. i. 364 = Senecio gelfolius.
geniculata, Wendl. ex Steud. Nom. ed. II. i. 364 = Senecio geniculatus.
geraniifolia, *DC. Prod.* vi. 306.—Afr. austr.
gibbosa, Gmel. Adnot. Cat. Pl. Hort. Bocc. (1821) 3 = Senecio gibbosus.
gigantea, Sm. Exot. Bot. ii. 11 = Senecio Smithii.
glabella, Turcz. ex DC. Prod. vi. 360 = Senecio campestris.
glaberrima, Spreng. ex DC. l. c. v. 25 = Vernonia cinerea.
glabra, Sw. Prod. Veg. Ind. Occ. 113 = Senecio Swartzii.
glabrata, Ledeb. Ic. Ross. t. 94 = Senecio racemosus.

CINERARIA :—
glabrata, Sw. Fl. Ind. Occ. iii. 1354 = Senecio Swartzii.
glandulifera, Vukot. in Rad Jugos. Akad. Zagreb. ii. (1880) 84 = Senecio crispatus.
glastifolia, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 146 = Senecio glastifolius.
glauca, Linn. Sp. Pl. ed. II. 1249 (= *Senecillis glauca*). —Rom. amstr.
gnaphaloides, *Nym. Consp.* 351.—Europ.
grandiflora, Spreng. ex DC. Prod. vi. 314 = Senecio amicoides.
grandiflora, Vatke, in Linnaea, xxxix. (1875) 508 = abyssinica.
Gualtieria, Gill. ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 340 = Senecio Hualtata.
hastifolia, Linn. f. Suppl. 376 = Senecio cymbalariifolius.
Heldreichii, Nym. Syll. 2 = Senecio Heldreichii.
Heimitis, Georgi, Beschr. Russ. Reich. iii. iv. 1246 = Senecio campris.
Helenitis, Sterd. Nom. ed. II. i. 864 = Senecio brachychaetus.
heterophylla, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 72 = Senecio resedifolius.
heterophylla, Ortog. Hort. Matr. 71 = Senecio minutus.
heterophylla, Pursh, Fl. Am. Sept. ii. 528 = Senecio tomentosus.
Heufleii, Hoppe, ex Nym. Consp. 351 = Senecio lugens.
Hieracium × , Hort. ex Mamock, Floric. Mag. iv. (1839) t. 89.
hirsuta, Vent. Jard. Malm. 95 = Aster hirsutus.
hirta, Vahl, ex DC. Prod. v. 226 = Aster elongatus.
Hualtata, Walp. Rep. ii. 453 = Senecio Hualtata.
humifuse, *L'Hérit. Sert. Angl.* 25.—Afr. austr.
humilis, Poepp. ex DC. Prod. vi. 417 = Senecio glaber.
hybrida, Bernh. Syst. Verz. Erf. 146 = Petasites officinalis.
hybrida, Willd. Enum. Hort. Berol. 898 = Doronicum crantum.
hypoleuca, *Reichb. ex DC. Prod.* vi. 389.—Afr. austr.
hypoleuca, Sieber, ex Harv. & Sond. Fl. Cap. iii. 381 = Senecio verbascifolius.
hypoleuca, Spreng. ex DC. l. c. 360 = Senecio rufinervis.
incana, Sw. Fl. Ind. Occ. iii. 1356 = Senecio jamaicensis.
incisa, Hoffm. ex DC. Prod. vi. 436 = Senecio incisus.
incisa, Thunb. Prod. Pl. Cap. 156 = Senecio incisus.
incisa, Willd. Sp. Pl. iii. 2074 = Senecio crenus.
integrifolia, Benth. ex Willd. & Lange, Prod. Fl. Hisp. ii. 110 = Senecio pyrenaicus.
integrifolia, Hook. & Arn. in Bot. Beech. Voy. 126 = Senecio ingens.
integrifolia, Jacq. ex Willd. Sp. Pl. iii. 2082 = Senecio alpestris.
integrifolia, Pursh, Fl. Am. Sept. ii. 528 = Senecio tomentosus.
integrifolia, Richards. in Frankl. Narr. 1st Journ. ed. II. 21 = Senecio Hookeri.
integrifolia, Schkuhr, Handb. t. 246 = Senecio crispatus.
integrifolia, Vill. ex Steud. Nom. ed. II. i. 864 = Senecio saranticus.
integrifolia, Wallr. Sched. Crit. 474 = Senecio spathulaefolius.
integrifolia, With. Bot. Arr. ed. II. 990 ; Sm. Eng. Bot. t. 159 = Senecio campestris.
japonica, Thunb. Fl. Jap. 317 = Senecio japonicus.
laciniata, Sw. Fl. Ind. Occ. iii. 1353 = Senecio Swartzianus.
lactea, Jacq. f. Eclog. t. 105 = Senecio crantus.
lactea, Willd. Enum. Hort. Berol. Suppl. 59 = Senecio appendiculatus.
laeta, Spreng. ex DC. Prod. vi. 404 = Senecio angulatus.
lanata, Jacq. Coll. iii. 177 = lanosus.
lanata, Lam. Encyc. ii. 7 = Senecio Hertiteri.
lanata, Link, Enum. Hort. Berol. ii. 888 = Doronicum crantum.
lanceolata, Lam. Fl. Fr. ii. 125 = Senecio campestris.
lanosa, *DC. Prod.* vi. 309.—Afr. austr.
laricifolia, Lam. Encyc. ii. 8 = Senecio pinifolius.
leucanthemus, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 316 = Senecio Smithii, scaanthifolius.
Lewisii, Richards. in Frankl. Narr. 1st Journ. App. 748 = Erigeron compositus.

CINERARIA :—
linnaeus, Linn. f. Suppl. 375 = Senecio linnaeus.
linnaeus, Vahl, ex DC. Prod. vi. 396 = Senecio tripli-nervius.
linifolia, Linn. Amoen. Acad. vi. 106 = Euryops linifolius.
linifolia, Zeyh. ex DC. Prod. v. 224 = Aster linifolius.
littoralis, Vest, in Flora, iii. (1820) 2 = Senecio gibbosus.
lobata, *L'Hérit. Sert. Angl.* 26.—Afr. austr.
lobata, Mair. ex DC. Prod. vi. 431 = Senecio sublobatus.
longifolia, All. Fl. Pedem. i. 208 = Senecio Balbisianus.
longifolia, Bess. Enum. Pl. Volh. 33 = Senecio alpestris.
longifolia, Eichw. Casp. Cauc. 22 = Senecio cladobotrys.
longifolia, Jacq. Fl. Austr. ii. 49. t. 181 = Senecio brachychaetus.
longifolia, Lapeyr. Hist. Abr. Pl. Pyr. 521 = Senecio pyrenaicus.
lucida, Sw. Fl. Ind. Occ. iii. 1360 = Senecio lucidus.
lyrata, *DC. Prod.* vi. 308.—Afr. austr.
lyrata, Ledeb. in Mém. Acad. Pétersb. v. (1818) 576 = Senecio resedifolius.
lyrato-pinnatifida, Bluff & Fing. Comp. ii. 377 = Senecio alpinus.
macrophylla, Ledeb. Fl. Alt. iv. 108 = Senecio Ledebourii.
malvaefolia, L'Hérit. Sert. Angl. 26 = Senecio malvaefolius.
maritima, Linn. Sp. Pl. ed. II. 926 = Senecio Cineria.
maritima, Wulf. ex Steud. Nom. ed. II. i. 864 = Senecio gibbosus.
megapotamica, Spreng. Syst. iii. 547 = Senecio pinnatus.
microglossus, *DC. Prod.* vi. 305.—Afr. austr.
microphylla, Vahl, ex DC. l. c. v. 235 = Aster asthiopicus.
microrhiza, Schur, in Verh. Siebenb. Naturw. Ver. (1859) 147 = Senecio campestris.
minuta, Cav. Ic. i. 21. t. 33. f. 3 = Senecio minutus.
mitellifolia, *L'Hérit. Sert. Angl.* 25.—Afr. austr.
mollis, *E. Mey. ex DC. Prod.* vi. 306.—Afr. austr.
mongolica, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 199 = Senecio mongolicus.
monticola, Spreng. *Syst.* iii. 548 = Senecio chilensis.
multiflora, L'Hérit. Sert. Angl. 26 = Senecio multiflorus.
nebrodensis, Guss. Adnot. Cat. Pl. Bocc. (1821) 4 = Senecio candidus.
niveus, Willd. Sp. Pl. iii. 1974 = Senecio niveus.
nuda, Kit. in Linnaea, xxxii. (1863) 389 = Senecio alpestris.
oppositifolia, Moench, Meth. 588 = Aster rotundifolius.
Othonnites, Linn. Sp. Pl. ed. II. 1244 = Othonna frutescens.
othonnoides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 314.—Afr. austr.
ovata, Spreng. ex DC. Prod. v. 344 = Aster asperulus.
ovirensis, Koch, in Flora, vi. (1823) 507 = Senecio alpestris.
oxyodus, *DC. Prod.* vi. 306.—Afr. austr.
palmensis, Nees, Hort. Phys. Berol. 115. t. 24 = Senecio palmensis.
palustris, Linn. Sp. Pl. ed. II. 1243 = Senecio barbareifolius.
pandurata, Thunb. Prod. Pl. Cap. 155 = Senecio panduratus.
papposa, Reichb. in Mossl. Handb. ed. II. ii. 1500 = Senecio papposus.
Parnassi, Nym. Consp. 352 = Senecio Parnassi.
parviflora, Bieb. Fl. Taur. Cauc. ii. 316 = Senecio lampsanoides.
parviflora, [R. Br. in] Ait. Hort. Kew. ed. II. v. 72 = cineseens.
pauciflora, Spreng. ex DC. Prod. vi. 367 = Senecio nudicaulis.
pedunculosus, *DC. l. c.* 305.—Afr. austr.
perfoliatus, Linn. f. Suppl. 378 = Othonna perfoliata.
Petasitis, Sims, Bot. Mag. t. 1286 = Senecio Petasitis.
phylicaefolia, Poepp. ex DC. Prod. vi. 415 = Senecio phylicaefolius.
pinnata, La Llave, in La Llave & Lex. Nov. Veg. Desc. i. 20 = Senecio pinnatisectus.

CINERARIA :—
pinnatifida, Willd. Sp. Pl. iii. 2074 = Senecio incertus.
platanifolia, Schrank, Fl. Rar. Hort. Monac. t. 95 = Senecio Petasites.
platycarpa, *DC. Prod.* vi. 305.—Afr. austr.
polyspathula, *DC. l. c.* 307.—Afr. austr.
polyglossa, DC. l. c. 306 = mollis.
popularifolia, Lam. Encyc. ii. 8 = Senecio appendiculatus.
populifolia, L'Hérit. Sert. Angl. 25 = Senecio populifolius.
praecox, Cav. Ic. iii. 23. t. 244 = Senecio praecox.
pratensis, Herder, in Bull. Soc. Nat. Mosc. xl. (1867) i. 400, partim = Senecio lugens.
pratensis, Hoppe, ex Reichb. Fl. Germ. Exsurs. ii. 241 = Senecio campestris.
procera, *Griseb. Spicil. Fl. Rumel.* ii. 219.—Rumel.
pulchella, E. Mey. ex DC. Prod. vi. 438 = Senecio cymbalariaefolius.
pulchella, *Sweet, Hort. Brit.* ed. I. 282.—Ins. Canar.
pumila, Thunb. Prod. Pl. Cap. 155 = Senecio repandus.
purpurascens, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 318 = Senecio acanthifolius.
purpurata, *Linn. Mant.* 286.—Afr. austr.
pyrenaica, Nym. Syll. 2 = Senecio pyrenaicus.
racemosa, Bieb. Fl. Taur. Cauc. ii. 314 = Senecio racemosus.
ramentosa, L'Hérit. Sert. Angl. 26 = Senecio echinatus.
ramosa, Vukot. in Rad Jugos. Akad. Zagreb. ii. (1880) 84.—Europ.
remifolia, C. A. Mey. Verz. Pfl. Cauc. 82 = Senecio remifolius.
remiformis, Spreng. ex DC. Prod. vi. 315 = Cremanthodium reniforme.
repanda, Lour. Fl. Cochinch. ii. 501 = Senecio scandens.
repanda, Willd. Sp. Pl. iii. 2076 = Senecio Forsteri.
retusa, Wall. ex C. B. Clarke, Comp. Ind. 206 = Senecio retusus.
rivularis, Waldst. & Kit. Pl. Rar. Hung. iii. 265 = Senecio crispatus.
robusta, Ledeb. Fl. Alt. iv. 106 = Senecio robustus.
rotundifolia, J. F. Gmel. Syst. 1240 ; Willd. Sp. Pl. iii. 2074 = Senecio rotundifolius.
rupicola, Nimp. Consp. 351 = *transsilvanica*.
salicifolia, H. B. & K. Nov. Gen. et Sp. iv. 188 = Senecio saligna.
samnitica, Huet, ex Nym. Consp. 352 = Senecio alpinus.
sarmentosa, Lesch. ex DC. Prod. vi. 369 = Senecio candicans.
Saxifraga, *DC. l. c.* 306.—Afr. austr.
scapiflora, L'Hérit. Sert. Angl. 25 = Senecio diversifolius, *Harv.*
scapigera, Banks, ex DC. Prod. vi. 430 = Senecio diversifolius, *Harv.*
Schimperi, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 483.—Abyssin.
Schkuhrii, Reichb. in Mossl. Handb. ed. II. ii. 1499 = Senecio crispatus.
scirpoea, E. Mey. ex DC. Prod. vi. 438 = Senecio juncous.
scordifolia, Link, ex DC. l. c. 429 = Senecio diversifolius, *Harv.*
Sellei, Spreng. Syst. iii. 541 = Senecio Sellei.
senecifolia, Fisch. Enoyc. Suppl. ii. 265 = Senecio alpinus.
Senecionis, *L'Hérit. Sert. Angl.* 26.—Hab. ?
serrata, Willd. Sp. Pl. iii. 2072 = Senecio Leontodontis.
serratifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 490 = Senecio paludosus.
sessilifolia, Jacquem. Voy. Bot. 91 = Wernaria nana.
sibirica, Linn. Sp. Pl. ed. II. 1242 = Senecio Ligularia.
solidaginifolia, Fisch. ex Ledeb. Fl. Ross. ii. 550 = Senecio resedifolius.
sonchifolia, *Linn. Sp. Pl. ed. II.* 1243.—Afr. austr.
spathulaefolia, C. C. Gmel. Fl. Bad. iii. 494 = Senecio spathulaefolius.
spathulaefolia, Lam. Encyc. ii. 8 = Othonna alata.
speciosa, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) ii. 13 (= *Ligularia speciosa*).—Siberia.

CINERARIA:—
speciosa, Schrad. ex Link, Enum. Hort. Berol. ii. 334 = Senecio Ligularia.
spinulosa, *Lam. Encyc.* ii. 9.—Afr. austr.
subdentata, Bunge, Enum. Pl. Chin. Bor. 39 = Senecio campestris.
sudetica, Koch, in Flora, vi. (1823) 505 = Senecio crispatus.
taraxacifolia, Bieb. Fl. Taur. Cauc. ii. 318 = Senecio taraxacifolius.
taygetea, Nym. Syll. 2 = Senecio candidus.
tenella, Link, Enum. Hort. Berol. ii. 334 = Aster tenellus.
tenuifolia, Gaudin, Fl. Helv. v. 306 = Senecio brachychaetus.
thapsoides, Nym. Syll. 2 = Senecio thapsoides.
thyrsoidea, *Icon. Fl. Austr.* ii. 481.—Austria.
thyrsoides, Ledeb. Fl. Alt. iv. 107 = Senecio sibiricus.
tillandsiaefolia, Poepp. ex DC. Prod. vi. 415 = Senecio argenteus.
tomentosa, *Less. Syn. Comp.* 391.—Afr. austr.
tomentosa, Mill. Gard. Dict. ed. VIII. n. 5 = Senecio canadensis.
trachyphylla, Spreng. Neue Entdeck. iii. 48 = Aster Ecklonis.
transilvanica, Schur, Enum. Pl. Transs. 345 = Senecio lugens.
trifurcata, Spreng. Syst. iii. 551 = Senecio trifurcatus.
tuberosa, E. Mey. ex DC. Prod. vi. 438 = Senecio inaequus.
tuxtlaginea, Thunb. Fl. Cap. 671 = Senecio verbascifolius.
Tuxtlaginis, L'Hérit. Sert. Angl. 26 = Senecio Tuxtlaginis.
tuxtlaginoides, Webb, ex Lond. Hort. Brit. Suppl. ii. 693, nomen = Senecio Tuxtlaginis.
uliginosa, Ledeb. in Spreng. Nov. Prov. (1818) 11 = Senecio gallicus.
uncinata, Spreng. ex DC. Prod. vi. 369 = Senecio densiflorus.
unctuosa, Gillb. Fl. Lituan. i. 211 = Senecio palustris.
undulata, Willd. Sp. Pl. iii. 2071 = Senecio petiolaris.
verna, Mair. ex DC. Prod. vi. 430 = Senecio saligna.
vernicifera, Mair. ex DC. l. c. = Senecio Andriezii.
vestita, Haw. in Phil. Mag. (1832) 187 = Othonna vestita.
vestita, Spreng. Syst. iii. 550 = Senecio crassiflorus.
viscosa, *L'Hérit. Sert. Angl.* 25.—Afr. austr.
Vitaliana, Nym. Consp. 351 = Senecio thapsoides.
vulneraria, Alam. ex DC. Prod. vi. 428 = Senecio vulnerarius.
Waterhousiana, Patt. Mag. Bot. iv. (1838) t. 219 = Senecio Tuxtlaginis ?
Webberiana, Patt. l. c. ix. (1843) t. 195.—Hab. ?
Wolffi, Schur, ex Nym. Consp. 359 = Senecio campestris.
Zahlbruckneri, Host, Fl. Austr. ii. 479 = Senecio lyratifolius.

CINGA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid ?).

CINHONA, Linn. Gen. ed. VI. 91 (1764), lapsu = Cinchona, Linn. (Rubiac.).

CINNA, Linn. Sp. Pl. 5 (1753). *GRAMINEAE*. Benth. & Hook. f. iii. 1151.
AGOLA, Adans. Fam. ii. 21 (1763).
BLYTTIA, Fries, Nov. Fl. Suec. Mant. ii. 2 (1839).
agrostidea, Beauv. ex Steud. Nom. ed. I. 20, 196 = arundinacea.
alba, *Nees, ex Steud. Syn. Pl. Gram.* 182.—Chili.
arachnoidea, Kunth, Rev. Gram. i. 67 = arundinacea.
arundinacea, Hook. Fl. Bor. Am. ii. 238 = Colpodium latifolium.
arundinacea, *Linn. Sp. Pl.* 5.—Reg. bor.
arundinacea, Kze. ex Steud. Nom. ed. II. i. 365 = Muehlenbergia mexicana.
Brownii, Roqr. Beitr. Pfl. Russ. Reich. ii. 66 = Colpodium latifolium.
criseta, Trin. Fund. Agrost. 118 = Dichelachne criseta.
duvigneae, Kunth, Rev. Gram. i. 67 = Deyeuxia scabra.
exigua, Link, Hort. Berol. ii. 296 = arundinacea.
filiformis, Link, Enum. Hort. Berol. i. 79 = Muehlenbergia mexicana.
CINNA :—
glomerata, Link, Hort. Berol. ii. 297 = Muehlenbergia glomerata.
glomerata, Walt. Fl. Carol. 59 = Andropogon macrourus.
japonica, Steud. Syn. Pl. Gram. 182 = Sporobolus indicus.
lanata, Kunth, ex Steud. Nom. ed. II. i. 365 = Epicampes lanata.
lateralis, Walt. Fl. Carol. 59 = Andropogon virginicus.
lateriflora, Kunth, Rev. Gram. i. 67 = Muehlenbergia mexicana.
latifolia, Griseb. in Ledeb. Fl. Ross. iv. 435 = arundinacea.
macroura, Kunth, Rev. Gram. i. 67 = Epicampes macroura.
macrura, Thurb. in S. Wats. Bot. Calif. ii. 276 = Epicampes rigens.
mexicana, Beauv. Agrost. 32 = Muehlenbergia mexicana.
ovata, Kunth, Rev. Gram. i. 67 = Echinopogon ovatus.
pendula, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. (1845) ii. 230 = arundinacea.
phleoides, *Kunth, Rev. Gram.* i. 67.—Venezuela.
pubescens, Kunth, l. c. = Epicampes pubescens.
Purshii, Kunth, l. c. = Calamagrostis canadensis.
racemosa, Kunth, l. c. = Muehlenbergia glomerata.
setifolia, Kunth, l. c. 64; et Enum. Pl. i. 208 = Epicampes macroura.
sobolifera, Link, Enum. Hort. Berol. i. 71 = Muehlenbergia sobolifera.
stricta, Kunth, Rev. Gram. i. 67 = Epicampes macroura.
suaveolens, Ruprr. ex Ledeb. Fl. Ross. iv. 435 = arundinacea.
tenuiflora, Link, Enum. Hort. Berol. i. 71 = Muehlenbergia Willdenowii.
valdiviana, *Phil. in Anal. Univ. Chil.* (1873) 563.—Chili.

CINNAGROSTIS, Griseb. in Goett. Abh. xix. (1874) 256. *GRAMINEAE*, Benth. & Hook. f. iii. 1152.
polygama, *Griseb. l. c.* 257.—Reg. Argent.

CINNAMODENDRON, Endl. Gen. 1029 (1840). *CANELLACEAE*, Benth. & Hook. f. i. 121.
axillare, *Endl. ex Walp. Rep.* i. 898.—Bras.
corticosum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 351.—Ind. occ.
macranthum, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 917.—Porto Rico.
rubrum, Griseb. Fl. Brit. W. Ind. 109 = corticosum.

CINNAMOMUM, [Tourn.] Linn. Syst. ed. I. (1735); Blume, Bijdr. 568 (1826). *LAURINEAE*, Benth. & Hook. f. iii. 155.
CAMPHORA, [Banh.] Linn. Syn. ed. I (1735); Nees, in Wall. Pl. As. Rar. ii. 61 (1831); Syst. Laurin. 87 (1836).
CRICODAPHNE, Nees, Syst. Laurin. 202 (1836).
PARTHENOXYLON, Blume, Mus. Bot. Lugd. Bat. i. 292 (1851).
albiflorum, Hook. f. & Thoms. ex Meisn. in DC. Prod. xv. 1. 21 = impressinervium.
albiflorum, Nees, in Wall. Pl. As. Rar. ii. 75 = C. Tamala.
angustifolium, Rafin. Sylva Tellur. 135 = Acrodiclidium salicifolium ?
aromaticum, J. Grah. Cat. Bomb. Pl. 173 = zeylanicum.
aromaticum, Nees, in Wall. Pl. As. Rar. ii. 74 = C. Camphora.
aromaticum, Zoll. Syst. Verz. Ind. Arch. 119 = inera.
Basaseia, Nees, in Wall. Pl. As. Rar. ii. 78 = obtusifolium.
Beijolghota, Sweet, Hort. Brit. ed. I. 344 = obtusifolium.
bonnense, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 260; *Meisn. in DC. Prod.* xv. 1. 19.—Borneo.
brevifolium, *Miq. in Verh. Nederl. Inst.* III. (1852) v. 14.—Japon.
Burmanni, *Blume, Bijdr.* 569.—Java.
calophyllum, Nees, Syst. Laurin. 40 = camphoratum.
calycinulum, *Miq. Fl. Ind. Bat. Suppl.* 359.—Sumatra.
CINNAMOMUM :—
Camphora, *T. Nees & Eberm. Handb. Med.-Pharm. Bot.* ii. 480.—China ; Japon.
camphoratum, *Blume, Bijdr.* 571.—Java.
camphoriferum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = C. Camphora.
Cappara-Coronde, Blume, Rumphia, i. 84. t. 9. f. 2, 3 = inera.
Cassia, *Blume, Bijdr.* 570.—Burma ; China.
Cassia, D. Don, Prod. Fl. Nep. 67 = C. Tamala.
Cassia, Siebold, in Verh. Batav. Gen. xii. (1830) 23 = Burmanni.
Cathia, D. Don, Prod. Fl. Nep. 66 = Phoebe paniculata.
caudatum, *Nees, in Wall. Pl. As. Rar.* ii. 76.—Reg. Himal. ; Burma.
Ceddodaphne, *Meisn. in DC. Prod.* xv. 1. 25.—Reg. Himal.
celebicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 264.—Ins. Celebes.
chinense, Blume, Bijdr. 569 = Burmanni.
Chloroxylon, J. S. Presl, Rostl. ii. 37 = Cenothus Chloroxylon.
citriodorum, *Thw. Enum. Pl. Zeyl.* 258.—Zeylan.
crassinervium, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 264.—Malaya.
Culilawa, Rafin. Sylva Tellur. 135 = seq. ?
Culilawan, *Blume, Bijdr.* 571.—Malaya ; China.
Culitlawan, Nees, in Wall. Pl. As. Rar. ii. 75 = pauc.
curvifolium, *Nees, Syst. Laurin.* 80.—China.
cuspidatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 262.—Sumatra.
cyrtopodum, *Miq. Fl. Ind. Bat.* i. 1. 397.—Sumatra.
daphnoides, *Sieb. & Zucc. in Abh. Acad. Muench. iv.* III. (1846) 202.—Japon.
dasyphyllum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 259.—Sumatra.
Doederleinii, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 57.—Archip. Loo-Choo.
dubium, Nees, in Wall. Pl. As. Rar. ii. 78 = inera.
dulce, Nees, l. c. 75 = Burmanni.
eucalyptoides, Nees, l. c. 73 = inera.
glaucescens, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 264.—Malaya.
glanduliferum, *Meisn. in DC. Prod.* xv. 1. 25.—Reg. Himal.
glaucum, J. S. Presl, Rostl. ii. 37 ; G. Don, ex Loud. Hort. Brit. 160 = Litsaea glauca.
gracile, Hort. ex Sweet, Hort. Brit. ed. I. 344, nomen = C. Camphora.
gracile, Miq. Ann. Mus. Bot. Lugd. Bat. i. 259 = inera.
Griffithii, Meisn. in DC. Prod. xv. 1. 19 = inera.
Heyneanum, *Nees, in Wall. Pl. As. Rar.* ii. 76.—Ind. or.
impressinervium, *Meisn. in DC. Prod.* xv. 1. 21.—Reg. Himal. ; China.
inera, *Reinw. ex Blume, Bijdr.* 570.—Burma ; Malaya.
inera, Wall. Cat. n. 3588 E = nitidum.
inera, Wight, Ic. t. 192, bis = zeylanicum.
inodorum, *Meisn. in DC. Prod.* xv. 1. 26.—Malaya.
inunctum, *Meisn. l. c.* 24.—Burma.
japonicum, Siebold, in Verh. Gen. Batav. xii. (1830) 23 = pedunculatum.
javanicum, *Blume, Bijdr.* 570.—Malaya.
Kiamis, Hassk. Cat. Hort. Bogor. Alt. 87 = pedunculatum.
Kiamis, Nees, in Wall. Pl. As. Rar. ii. 75 = Burmanni.
lampongum, *Miq. Fl. Ind. Bat. Suppl.* i. 356.—Sumatra.
Laubatii, F. Muell. Fragm. v. 165 = C. Tamala.
laxiflorum, *Meisn. in DC. Prod.* xv. 1. 21.—Java.
litseaefolium, *Thw. Enum. Pl. Zeyl.* 253.—Zeylan.
Loureirii, *Nees, Syst. Laurin.* 66.—Cochinch.
lucens, Miq. Ann. Mus. Bot. Lugd. Bat. i. 261 = tavoyanum.
macrophyllum, *Miq. l. c.* 269.—Java.
Malabathrum, Batka, in Nov. Act. Nat. Cur. xvii. (1835) II. 618 = inera.
Malabathrum, Miq. Fl. Jungh. i. 174 = subtetrapterum.
malaccense, Meisn. in DC. Prod. xv. 1. 27 = C. Parthenoxylon.
Marsba, *Meisn. in DC. l. c.* 15.—Malaya.
montanum, J. S. Presl, Rostl. ii. 36 ; Nees, Syst. Laurin. 61 = Phoebe montana.

CINNAMOMUM :—
meltiformis, *Wight, Ic.* t. 181.—Zeylan.
Nessianum, Meisn. in DC. Prod. xv. i. 26 = C. Parthenoxylon.
neglectum, Blume, Rumphia, i. 88, in nota, t. 11. f. 2 = javanicum.
nitidum, *Blume, l. c.* 85. t. 18. f. 2; t. 16. f. 1, 2.—Burma; Malaya.
nitidum, Hook. Exot. Fl. iii. t. 176 = inera.
obscurum, *Meisn. in DC. Prod.* xv. i. 26.—Sumatra.
obtusifolium, *Nees, in Wall. Pl. As. Rar.* ii. 78.—Reg. Himal.; Burma.
ochraceum, Blume, Rumphia, i. 87 = inera.
officinarum, Nees, ex Steud. Nom. ed. II. i. 366 = C. Casia !
ovalifolium, Gardn. ex Meisn. in DC. Prod. xv. i. 11 = Wightii.
ovalifolium, *Wight, Ic.* t. 125.—Zeylan.
paraneuron, *Miq. Fl. Ind. Bat.* i. i. 895.—Sumatra.
Parthenoxylon, *Meisn. in DC. Prod.* xv. i. 26.—Burma; Malaya; China.
pauciflorum, *Nees, in Wall. Pl. As. Rar.* ii. 78.—Reg. Himal.
pedalinervium, *Meisn. in DC. Prod.* xv. i. 15.—Ins. Fiji.
pedunculatum, *J. S. Presl, Rostl.* ii. 37; *Nees, Syst. Laurin.* 79.—Japon.
Perrottetii, *Meisn. in DC. Prod.* xv. i. 22.—Ind. or.
politum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 285.—Malaya.
pseudo-sassafras, Meisn. in DC. Prod. xv. i. 27 = C. Parthenoxylon.
pseudo-zeylanicum, Miq. Fl. Ind. Bat. i. i. 902 = inera.
pyrifolium, *D. Don, Prod. Fl. Nep.* 66 (sp. dubia).—Nepal.
Rauwolfii, Blume, Rumphia, i. 86. t. 9. f. 4, 5 = inera.
recurvatum, Wight, Ic. t. 132 = pauciflorum.
Reinwardti, Miq. Fl. Ind. Bat. i. i. 900 = inera.
Reinwardti, Nees, Syst. Laurin. 70 = C. Tamala.
rhynchophyllum, *Miq. Fl. Ind. Bat.* i. i. 895.—Sumatra.
rubrum, Blume, in Hoev. & De Vriese, Tijdschr. i. (1834) 62 = C. Cuilawan.
sericans, *Hance, in Journ. Bot.* xv. (1877) 336.—China.
sericeum, Siebold, in Batav. Gen. Verh. xii. (1830) 24 = Lauras sericea.
Sieboldii, *Meisn. in* DC. Prod. xv. i. 18 = Burmanni.
Sintok, *Blume, Bijdr.* 571.—Malaya.
subavenium, *Miq. Fl. Ind. Bat.* i. i. 902.—Sumatra.
subenneatum, *Miq. l. c.* 895.—Sumatra.
substriatum, *Miq. l. c.* 902.—Sumatra.
sulphuratum, Kurz, For. Fl. Brit. Burm. ii. 288 = tavoyanum.
sulphuratum, *Nees, in Wall. Pl. As. Rar.* ii. 74.—Ind. or.
sumatranum, *Meisn. in DC. Prod.* xv. i. 25.—Sumatra.
Tamala, *T. Nees & Eberm. Handb. Med.-Pharm. Bot.* ii. 426.—Reg. Himal.
tavoyanum, *Meisn. in DC. Prod.* xv. i. 20.—Burma.
tenuifolium, D. Don, Prod. Fl. Nep. 66 = Phoebe paniculata.
validinerve, *Hance, in Journ. Bot.* xx. (1882) 80.—China.
verum, J. S. Presl, Rostl. ii. 36; Sweet, Hort. Brit. ed. I. 344 = zeylanicum.
villosum, Wight, ex Meisn. in DC. Prod. xv. i. 16 = multiflorum.
villosum, Wight, Ic. t. 127 = inera.
vimineum, *Nees, in Wall. Pl. As. Rar.* ii. 76.—Ins. Penang.
Wightii, *Meisn. in DC. Prod.* xv. i. 11.—Ind. or.
xanthoneurum, *Blume, in Hoev. & De Vriese, Tijdschr.* i. (1834) 68.—N. Guinea.
zeylanicum, *Nees, in Wall. Pl. As. Rar.* ii. 74; ii. 92. —Ind. or.; Malaya.

CINNAMOSMA, Baill. Adansonia, vii. (1866-67) 219. *CANELLACEÆ*, Benth. & Hook. f. i. 970.
fragrans, *Baill. l. c.*—Malaya.

CINNASTRUM, Fourn. Gram. Mex. 90, ex Benth. & Hook. f. Gen. iii. 1158 (1883) = Deyeuxia, Clar. (Gramin.).

CINOGASUM, Neck. Elem. ii. 326 (1790) = Croton, Linn. (Euphorbiac.).

CIONANDRA, Griseb. Fl. Brit. W. Ind. 326 (1860) = **Cayaponia**, Silva Manso (Cucurbitac.).
angustiloba, Ballo, in Anal. Soc. Esp. Hist. Nat. x. (1881) 274.—Porto Rico.
cuspidata, Griseb. Fl. Brit. W. Ind. 327 = Cay. americana.
gracilifora, Griseb. l. c. = Cay. americana.
racemosa, Griseb. l. c. 326 = Cay. racemosa.

CIONIBACCUS, Breda, Orch. Kuhl et Hass. (1827) [t. 8] = **Goodyera**, R. Br. (Orchid.).
lanceolatus, Breda, l. c. = G. procera.

CIONOSICYOS, Benth. & Hook. f. Gen. i. 826 (1867) = sq.

CIONOSICYS, Griseb. Fl. Brit. W. Ind. 288 (1860). *CUCURBITACEÆ*, Benth. & Hook. f. i. 826.
pomiformis, *Griseb. l. c.*—Ind. occ.

CIONURA, Griseb. Spicil. Fl. Rumel. ii. 69 (1844) = **Marsdenia**, R. Br. (Asclepiad.).
erecta, Griseb. l. c. = M. erecta.

CIPADESSA, Blume, Bijdr. 162 (1825). *MELIACEÆ* [Benth. & Hook. f. i. 382].
MALLEA, A. Juss. in Mém. Mus. Par. xix. (1830) 221.
baccifera, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 6 = fruticosa.
Boivinians, *Baill. Adansonia*, xi. (1874) 255.—Madag.
bornensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 6.—Borneo.
depauperata, *Baill. Adansonia*, xi. (1874) 256.—Ins. Comor.
fruticosa, *Blume, Bijdr.* 162.—Java.
subcandens, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 7 = fruticosa.

CIPONIMA, Aubl. Pl. Gui. i. 566. t. 226 (1775) = **Symplocos**, Linn. (Styrac.).
guianensis, Aubl. l. c. 567 = S. Ciponima.
scabridula, Miers, in Journ. Linn. Soc. xvii. (1879) 288 = S. Ciponima.

CIPURA, Aubl. Pl. Gui. i. 28. t. 12 (1775). *IRIDEÆ*, Benth. & Hook. f. iii. 694.
MARICA, Schreb. Gen. Pl. 87 (1789) (non Ker-Gawl.).
aurea, Griseb. & Klatt, in Abh. Naturf. Ges. Halle, xii. (1868) 287 = Calydorea aurea.
caerulea, Heynh. Nom. i. 197 = Marica caerulea.
coelestis, Heynh. l. c. = Nemastylis coelestinus.
cubensis, *Grisb. Cat. Pl. Cub.* 251.—Ins. Cuba.
gracilis, Heynh. Nom. i. 197 = Marica gracilis.
graminea, H. B. & K. Nov. Gen. et Sp. i. 320 = palmdora.
humilis, H. B. & K. l. c. = palmdosa.
limoria, Klotzsch & Klatt, in Abh. Naturf. Ges. Halle, xii. (1868) 287 = Calydorea cipuroides.
longifolia, Heynh. Nom. i. 197 = Marica longifolia.
martinicensis, H. B. & K. Nov. Gen. et Sp. i. 320 = Trimezia lurida.
Northiana, *Bodl. in Harting. Pavad. Vindob.* t. 6.—Bras.
palmdosa, *Aubl. Pl. Gui.* i. 28. t. 13.—Am. trop.
plicata, Griseb. Fl. Brit. W. Ind. 589 = Eleutherine Bulbosa.
Sabini, Heynh. Nom. i. 197 = Northiana.
Sabiniana, Voigt, Hort. Suburb. Calc. 602 = Marica Sabini.
semiaperta, Heynh. l. c. = Trimezia lurida.

CIPURA, Klotzsch, ex Klatt, in Abh. Naturf. Ges. Halle, xii. (1868) 302 = Herbertia, Sweet (Irid.).

CIRCÆA, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 8 (1737). *ONAGRARIEÆ*, Benth. & Hook. f. i. 793.
Reneria, Delac. Fl. Hautes-Pyr. 333 (1867).
alpestris, Schur, Enum. Pl. Transs. 214 = C. lutetiana.
alpestris, Wallr. Beitr. Fl. Hercyn. 167 = alpina.
alpina, *Linn. Sp. Pl.* 9.—Europ.; As. bor.; Am. bor.
canadensis, Michl. Cat. 2 = C. lutetiana.
cordata, *Royle, Illustr. Bot. Himal.* 211. t. 48. f. 1.—Reg. Himal.
cordifolia, Bodard, Bot. Mat. Med. i. 98 = alpina.
decumbens, Gilib. Fl. Litaus. ii. 127 = alpina.
erecta, Gilib. l. c. 126 = C. lutetiana.

CIRCÆA :—
erioctonon, *Mærtr. Fl. Crit. Tarn.* 29.—Gallia.
erubescens, *Franch. & Sav. Enum. Pl. Jap.* ii. 270.—Japon.
intermedia, *Ehrh. Beitr.* iv. 42.—Europ.
intermedia, Wall. Cat. n. 6343 = alpina.
intetiana, Georgi, It. i. 126 = alpina.
intetiana, *Linn. Sp. Pl.* 9.—Reg. bor.
major, Lam. Fl. Fr. iii. 473 = C. lutetiana.
minima, Lam. l. c. = C. alpina.
mollis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. (1847) 184.—Japon.
nemorosa, Salisb. Prod. 276 = C. lutetiana.
ovalifolia, Stokes, Bot. Mat. Med. i. 96 = C. lutetiana.
pacifica, *Aschers. & Magnus, in Bot. Zeit.* xxix. (1871) 392.—Am. bor. occ.
pubescens, Pohl, ex Steud. Nom. ed. II. i. 366 = C. lutetiana.
quadriulcata, *Maxim. in Franch. & Sav. Enum. Pl. Jap.* i. 169.—Japon.
repens, Wall. Cat. n. 6341 = C. lutetiana.
vulgaris, Moench, Meth. 379 = C. lutetiana.

CIRCÆASTER, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 556. *CHLORANTHACEÆ* ?, Benth. & Hook. f. iii. 1220.
agrestis, *Maxim. l. c.* 557.—Reg. Himal.; As. centr.

CIRCINNUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 384 = **Hymenocarpus**, Savi [1798] (Legumin.).
vuinervicides, Medic. l. c. = H. circinnata.

CIRCINUS, Medic. Phil. Bot. i. 208 (1789) = præc.

CIRCIS, Chapm. Fl. S. U. St. 114 (1860), lapsu = **Cercis**, Linn. (Legumin.).

CIRINOSUM, Neck. Elem. ii. 84 (1790) = **Cereus**, Haw. (Cactac.).

CIRIPEDIUM, Zumag. Fl. Pedem. i. 18 (1809) = **Cypripedium**, Linn. (Orchid.).

CIRRHÆA, Lindl. Bot. Reg. sub t. 980 (1825). *ORCHIDEÆ*, Benth. & Hook. f. iii. 585.
SARCOLOBIUM, Beer, Prakt. Orch. (1854).
SCLEROPTERIS, Scheidw. in Allg. Gartenz. (1839) 407 adsperus, *Hoffm. ex Heynh. Nom.* ii. 146, nomen.—Bras.
dependens, Lond. Hort. Brit. 370 = Loddigesii.
dependens, Reichb. f. in Walp. Ann. vi. 496 = viridipurpurea.
fuscolutea, Hook. Bot. Mag. t. 6796 = saccata.
fuscolutea, *Lindl. Bot. Reg.* sub t. 1538.—Bras.
Hoffmannseggii, Heynh. ex Reichb. f. in Walp. Ann. vi. 496 = viridipurpurea.
Loddigesii, *Lindl. Bot. Reg.* t. 1538.—China !
obtusata, *Lindl. l. c.* xxiii. (1837) 2005.—Bras.
pallida, *Lindl. l. c.* sub t. 2005, nomen.—Hab. !
sacosa, *Lindl. l. c.* (1839) *Misc.* 72.—Bras.
tristis, *Lindl. l. c.* t. 1889.—Mexic.
violacea-virens, *Hoffmgg. Verz. Orch.—Cf. Bot. Zeit* i. (1843) 831.—Hab. !
violacea, *Hoffmgg. ll. cc.*—Hab. !
viridipurpurea, *Lindl. Bot. Reg.* t. 1538.—Bras.
viridi-purpurea, Lodd. Bot. Cab. t. 1967 = Loddigesii.
Warreana, *Lindl. ex Reichb. f. in Walp. Ann.* vi. 497. —Bras.

CIRRHOPETALUM, Lindl. Bot. Reg. sub t. 832 (1824). *ORCHIDEÆ*, Benth. & Hook. f. iii. 504.
EPHIPPIUM, Blume, Bijdr. 309. t. 65 (1825).
BOLBOPHYLLOPHIS, Reichb. f. in Bot. Zeit. x. (1852) 993.
HIPPOGLOSSUM, Breda, Orch. Kuhl et Hass. (1827) [t. 14].
Zygoglossum, Retzw. in Syll. Ratisb. ii. (1828) 4, abbreviatum, *Reichb. f. in Gard. Chron.* (1881) ii. 70. —Hab. !
altilobum, *Wight, Ic.* t. 1658.—Ind. or.
Andersonii, *Kurz, Rep. Veg. Andaman.* 51.—Ins. Andaman.
antenniferum, *Lindl. Bot. Reg.* (1842) sub t. 69.—Ins. Philipp.
auratum, *Lindl. l. c.* (1840) *Misc.* 52.—Ins. Philipp.
Blumii, *Lindl. Gen. et Sp. Orch.* 86.—Java.
bootanensis, Griff. Notul. iii. 224.—*cucullatum*.
complanatum, *Wall. ex Lindl. Bot. Reg.* (1838) *Misc.* 26.—Reg. Himal.

CIRRHOPETALUM :—
capitatum, *Lindl. Bot. Reg.* (1843) sub t. 49.—Java.
carinatum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 207.—Java.
caudatum, Wight, Ic. t. 1636 = vaginatum.
chinense, *Lindl. Bot. Reg.* (1842) *Misc.* 87.—China.
clavigerum, *Fitzgerald, in Journ. Bot.* xxi. (1883) 204.—Austral.
compressum, *Lindl. Bot. Reg.* xxix. (1843) sub t. 49.—Java.
cornutum, *Lindl. l. c.* (1886) *Misc.* 75.—Reg. Himal.
Cumingii, *Lindl. l. c.* sub t. 49.—Ins. Philipp.
delitescens, Rolfe, in Gard. Chron. (1892) ii. 461.—Hongkong.
elegans, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 310.—Malaya.
Elisae, F. Muell. Fragm. vi. 120. t. 57 = Bulbophyllum Elisae.
elongatum, *Lindl. Bot. Reg.* (1843) sub t. 49.—Reg. Himal.
fimbriatum, *Lindl. l. c.* (1839) *Misc.* 72.—Ind. or.
flagelliforme, Tejsm. & Binn. ex Reichb. in Walp. Ann. vi 264 = Pahudi.
gamosepalum, *Griff. Notul.* iii. 296.—Burma.
grandiflorum, *Wight, Ic.* v. t. 1658.—Zeylan.
lasiochilum, *Par. & Reichb. f. in Trans. Linn Soc.* xxx. (1874) 158 (Bulbophylli §).—Burma.
leopardinum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 309.—Java.
linearum, *Teijsm. & Binn. l. c.* 311.—Malaya.
longiscapum, *Teijsm. & Binn. l. c.* 310.—Malaya.
Macraei, *Lindl. Gen. et Sp. Orch.* 59.—Zeylan.
Macraei, Wight, Ic. t. 1652 = Wightii.
maculosum, *Lindl. Bot. Reg.* (1841) *Misc.* 81.—Reg. Himal.
Makoyanum, Reichb. f. in Gard. Chron. (1879) i. 284 = Bulbophyllum Makoyanum.
maxillare, *Lindl. Bot. Reg.* (1843) sub t. 49.—Ins. Philipp.
Medusae, *Lindl. l. c.* (1842) t. 12.—Malaya.
neilgherrense, *Wight, Ic.* vi. t. 1954.—Ind. or.
nutans, *Lindl. Bot. Reg.* (1839) *Misc.* 71.—Manilla.
ornatissimum, *Reichb. f. in Gard. Chron.* (1882) ii. 424.—Reg. Himal.
Pahudi, *De Vriese, Epim. Lugd.-Bat.* (1868) 9.—Java.
picturatum, *Lodd. in Bot. Reg.* (1840) *Misc.* 49.—Burma.
Pumilio, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 152 (Bulbophylli §).—Burma.
putidum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 311.—Java.
refractum, *Edl. in Flora*, xxx. (1847) 456.—Reg. Himal.; Malaya.
retusiaculum, *Reichb. f. in Gard. Chron.* (1869) 1182.—Burma.
Roxburghii, *Lindl. Gen. et Sp. Orch.* 58.—Ind. or.
similimum, *Reichb. f. Otia Hamburg.* 48.—Burma.
stramineum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 310.—Java.
Thouarsii, *Lindl. Bot. Reg.* sub t. 882, in texte.—Ins. Masear.
trigonopus, *Reichb. f. in Gard. Chron.* (1861) ii. 71.—Hab.?
tripedians, *Par. & Reichb. f. l. c.* (1876) i. 816.—Burma.
tumidilabium, Linden, Illustr. Hortic. xxviii. (1881) 80. = maculosum?
vaginatum, *Lindl. in Wall. Cat.* n. 1979.—Malaya.
Walkerianum, Wight, Ic. v. t. 1657 = Mecrael.
Wallichii, J. Grah. Cat. Pl. Bomb. 206 = fimbriatum?
Wallichii, Lindl. in Wall. Cat. n. 1980 = refractum.
Wallichii, *Lindl. in Wall. Pl. As. Rar.* i. 58. t. 67.—Reg. Himal.
Wightii, *Thw. Enum. Pl. Zeyl.* 299.—Zeylan.

CIRSELIUM, Brot. Fl. Lusit. i. 346 (1804) = seq.

CIRSELLIUM, Gaertn. Fruct. ii. 454. t. 168 (1791) = Atractylis, Linn. (Compos.).
cancellatum, Gaertn. l. c. = A. cancellata.
humile, Gaertn. l. c. 455 = A. humilis.
gummiferum, Brot. Fl. Lusit. i. 346 = A. gummifera.

CIRSIUM, [Tourn.] Adans. Fam. ii. 116 (1763); DC. Fl. Fr. iv. 110 (1805) = Onicus, Tourn. (Compos.).
abyssinicum, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 456 = Cn. lanceolatus.
acanthifolium, Arv. in Bull. Soc. Dauph. iii. (1876) 78.—Gallia.
Acarna, Moench, Meth. Suppl. 226 = Cn. Acarna.
acaule, [Weber, ex] Wigg. Prim. Fl. Hols. 59 = Cn. acaulis.
acaulon, Scop. Fl. Carn. ed. II. ii. 181 = Cn. acaulis.
aciculare, Tausch, in Flora, xii. (1829) i. Erg. 40.—Hab.?
acinaceum, Fisch. & Mey. ex DC. Prod. vi. 636 = *fallax*.
affine, Tausch, in Flora, xvi. (1833) 228.—Bohem.
afrum, DC. Cat. Hort. Monsp. 96 = Cn. afer.
aggregatum, Ledeb. Fl. Ross. ii. 787.—Reg. Cancas.
Alberti, Regel & Schmalh. in Act. Hort. Petrop. v. (1879) 818.—Turkestan.
albicans, Willk. in Linnaea, xxx. (1859–60) 109.—Hispan.
Allionii, Spenn. Fl. Friburg. iii. 1079 = Cn. acaulis.
Allionii, Thurns. ex Ardoin. Fl. Alpes-Marit. 198 = Cn. pyrenaicus.
alpestre, Nageli, Cirs. Schw. 84 = Cn. acaulis.
alpestre x, Treuinf. in Zeitschr. Ferdinand. Tirol. III. xix. (1875) 228.—Tyrol.
alpinum, All. Fl. Pedem. i. 153 = Saussurea alpina, discolor.
altissimum, Hill, Hort. Kew. 63 = Cn. altissimus.
ambiguum, All. Auct. Pedem. 10 = Cn. heterophyllus.
anglicum, DC. Fl. Fr. iv. 118 = Cn. pratensis.
angulosum, Tausch, in Flora, xix. (1836) 402.—Hab.?
angustifolium, DC. Prod. vi. 648 = Carduus angustifolius.
angustissimum, DC. Prod. vi. 652 = Tricholepis strictophyllum.
apiculatum, DC. l. c. 642 = *libanoticum*.
appendiculatum, Griseb. Spicil. Fl. Rumel. ii. 250.—Rumel.; Herceg.
arachnoideum, Bieb. Fl. Taur. Cauc. iii. 557 = Cn. arachnoideus.
arancense, Stend. Nom. ed. II. i. 367 = Cn. arenoesus.
arctioides, Scop. Fl. Carn. ed. II. ii. 124 = Carduus arctioides.
arctostaphylum, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 144 = Cn. Schimperi.
argentium, Feyer, ex Vest, in Flora, xii. (1829) i. Erg. 87 = Cn. arvensis.
argenteum, DC. Prod. vi. 644.—Sibiria.
argyracanthum, DC. l. c. 640 = Cn. argyracanthus.
aristatum, DC. l. c.—Armenia.
armenum, DC. l. c. 647 = *simplex*.
arvense, Scop. Fl. Carn. ed. II. ii. 126 = Cn. arvensis.
Aschersonii x, Celak. in Arch. Naturfr. Landesd. Bohm. ii. (1873) 268.—Bohem.
atriplicifolium, Fisch. ex Stend. Nom. ed. II. i. 367 = Serratula atriplicifolia.
atticum, Sch. Bip. ex Nym. Consp. 406 = *crinitum*.
Auserdorferi x, Hausm. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 228 = Cn. ochroleucus?
autarcticum, Mutel, Fl. Dauph. ii. 282 = Cn. heterophyllus.
autumnale, Spreng. Syst. iii. 377.—Hispan.
benacense, Treuinf. in Oestr. Bot. Zeitschr. xxiv. (1874) 172.—Tyrol.
bipedunculi, Spreng. Syst. iii. 377 = Cn. Bertolonii.
Biglowii, DC. Prod. vi. 652 = Cn. muticus.
bipontinum, F. Schultz, in Arch. Fl. Fr. et Allem. 84 = Cn. acaulis.
Bonjeani, Sch. Bip. in Oestr. Bot. Wochenbl. vi. (1856) 209 = Cn. ciliatus.
Bourgaeanum, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 191.—Hispan.
brachycephalum, Juratzka, in Verh. Zool. Bot. Ver. Wien, vii. (1857) 99 = *Chailletii*, Koch.
bracteatum, Link, in Linnaea, ix. (1834) 580 = Cn. syriacus.
bracteosum, DC. Prod. vi. 641 = Cn. strigosus.
Braunii, F. Schultz, ex Nym. Consp. 409 = Cn. oleraceus.
breviculo, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 396 = Cn. japonicus.
brevifolium, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 421 = Cn. undulatus.
brevilepium, Sch. Bip. ex Nym. Consp. 410 = Cn. pengens.

CIRSIUM :—
britannicum, Soop. Ann. i). Hist. Nat. (1769) 80 = Cn. pratensis?
Bruggeri x, Kill. ex Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.–xxiv. (1880) 106.—Helvet.
Brunneri, A. Braun, ex Nym. Consp. 407 = Cn. tuberosus.
Buckingeri, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 284, nomen.—Abyss.
Buergeri, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 185 = Cn. Buergeri.
bulbosum, DC. Fl. Fr. iv. 118 = Cn. tuberosus.
bulbosum, Lam. Fl. Fr. ii. 23 = Cn. monspessulanus.
bulbosum, Willk. in Flora, xxxiv. (1851) 760 = *filipendulum*, Lange.
bulgaricum, DC. Prod. vi. 639 = Cn. gigantens.
byzantinum, Stend. Nom. ed. II. i. 367.—Rumel.
californicum, A. Gray, in Pacif. Rail. Rep. iv. 112 = Cn. californicus.
Candelabrum, Griseb. Spicil. Fl. Rumel. ii. 251.—Graecia; Rumel.
Candolleanum, Naeg. Cir. Schweiz. 98 = Cn. ochroleucus?
canescens, A. Gray, Pl. Fendl. 110 = Cn. neo-mexicanus.
canescens, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 420 = Cn. undulatus.
canum, All. Fl. Pedem. i. 151 = Cn. pratensis?
canum, Bieb. Fl. Taur. Cauc. ii. 274 = Cn. canus.
Caput-Medusae, Schur, ex Nym. Consp. 406 = Cn. spinosissimus.
carlinoides, Fisch. in DC. Prod. vi. 661 = Cn. carlinoides.
carniciense, Dum. Fl. Belg. 74.—Belgium.
carnioli cum, All. Fl. Pedem. i. 149 = Cn. rivularis.
carnioli cum, Scop. Fl. Carn. ed. II. ii. 128 = Cn. rufescens.
Carolorum x, C. Jenner, ex Nym. Consp. 407 = Cn. heterophyllus.
carthamoides, Link, Enum. Hort. Berol. ii. 303 = Centaurea carthamoides.
Casabonae, DC. Fl. Fr. iv. 121 = Cn. Casabonae.
castellanum, Willk. in Linnaea, xxx. (1859–60) 109 = Cn. flavispinus.
catanoticum, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 585.—Syria.
Celakovskianum x, Knaf, in Oestr. Bot. Zeitschr. xxii. (1872) 309 = Cn. arvensis.
Cephalotes, Boiss. Diagn. Ser. I. vi. 102 = Cn. horridus.
cernuum, Lag. Gen. et Sp. Nov. 24 = Cn. nivalis.
Cervini, Koch, Syn. Fl. Germ. ed. I. 399 = Cn. spinosissimus.
Chailletii, Gaudin, Fl. Helv. v. 182 = Cn. palustris.
Chailletii, Koch, Syn. Fl. Germ. ed. I. 398.—Europ.
Chamaepeuce, Tenore, Fl. Nap. v. 211 = Cn. gnaphalodes.
cheiranum, Orph. in Boiss. Diagn. Ser. II. iii. 41 = *Candelabrum*.
chinense, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 328 = Cn. chinensis.
chium, DC. Prod. vi. 646 = Carduus chius.
chrysopicum, Boiss. Fl. Orient. iii. 586.—Armenia.
cichoraceum, Spreng. Syst. iii. 370 = Serratula cichoracea.
ciliato-affine, Boss. Enum. Fl. Volh. 33 = Cn. serrulatus.
ciliatum, Bieb. Fl. Taur. Cauc. iii. 556 = Cn. ciliatus.
ciliatum, Moench, Meth. Suppl. 227.—Hab.?
ciliaeum, Tardent, Ess. Hist. Nat. Bessar. 63 = Cn. ciliatus.
compactum, Lam. Fl. Fr. ii. 24 = Cn. monspessulanus.
congestum, Fisch. & Mey. ex DC. Prod. vi. 641.—Persia.
consanguineum, DC. l. c. = *congestum*.
conspicuum, Sch. Bip. in Seem. Bot. Voy. Herald, 312 = Cn. conspicuus.
controversum, DC. Prod. vi. 654 = Cn. spinosissimus.
Czernaii, Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 251 = Cn. lappaceus.
Coulteri, Harv. & Gray, Pl. Fendl. 110 = Cn. occidentalis.
creticum, Urv. in Mém. Soc. Linn. Par. i. (1822) 368.—Archip. Graec.
crinitum, Boiss. ex DC. Prod. vii. 636.—Hispan.
cryptocephalum, Sch. Bip. ex Boiss. Fl. Orient. iii. 592 = Centaurea pygmaea.
cuspidinae, Borb. in Oestr. Bot. Zeitschr. xxviii. (1878) 393.—Europ.
cyanoides, Hill, Hort. Kew. 63 = Jurinea cyanoides.
cyanoides, Griseb. Spicil. Fl. Rumel. ii. 264 = Cn. afer.

CIRSIUM :—
cynaroides, Spreng. Syst. iii. 372 = Cnicus cynaroides.
dealbatum, Bieb. Fl. Taur. Cauc. iii. 560.—Reg. Cancas.
decipiens, Nym. Syll. Suppl. 4 = Cn. acaulis.
decoloratum, Koch, Syn. Fl. Germ. ed. 1. 398 = Cn. acaulis, oleraceus.
decurrens, Janka, in Linnaea, xxx. (1859–60) 582 = Cn. ferox, *Sibth.* & *Sm.*
defloratum, Scop. Fl. Carn. ed. II. ii. 137 = Carduus deflioratus.
dentatum, DC. Prod. vi. 654 = Kentrophyllum dentatum.
depilatum, Bolss. & Bal. in Boiss. Fl. Orient. iii. 648. —As. Min.
desertorum, Fisch. ex Link, Enum. Hort. Berol. ii. 300. —Siberia.
desertorum, Janka, ex Nym. Consp. 405 = Cn. ciliatus.
Diacanthe, DC. Cat. Hort. Monsp. 96 = Cn. Diacantha.
dioicum, Cass. in Dict. Sc. Nat. xxvii. 190 = Cn. arvensis.
discolor, Hort. ex Stend. Nom. ed. II. i. 368 = Cn. ciliatus.
discolor, Spreng. Syst. iii. 373 = Cn. altissimus.
dissectum, Hill, Hort. Kew. 62 = Cn. dissectus.
disserifolium, DC. Prod. vi. 649 = Cn. altissimus.
Dollineri, Sch. Bip. ex Hausm. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 373 = Cn. pannonicus.
Douglasii, DC. Prod. vi. 643 = Cn. undulatus.
Dreieri, Reichb. f. in Fl. Germ. xv. 60 = Cn. lanceolatus.
Drummondii, Torr. & Gray, Fl. N. Am. ii. 459 = Cn. Drummondii.
duriusculum, C. Koch, in Linnaea, xxiv. (1851) 410 = *eiodes*, Cn. palustris.
echinatum, DC. Fl. Fr. Suppl. 466 = Cn. echinatus.
echinocephalum, Bieb. Fl. Taur. Cauc. iii. 559 = Cn. echinocephalus.
echinophyllum, Boiss. ex Tchihat. Asie Min. Bot. ii. 548.—As. Min.
edule, Nutt. in Trans. Am. Phil. Soc. N. S. vi. (1841) 450 = Cn. edulis.
elatius, Link, Enum. Hort. Berol. ii. 299 = Cn. polyanthemum.
elatum, Sauter, in Flora, xxviii. (1845) 190.—Europ.
elatum, Tod. Ind. Sem. Hort. Panorm. (1858) 25; et in Ann. Sc. Nat. Sér. IV. xi. (1859) 378.—Sicilia.
elegans, Willd. ex Spreng. Syst. iii. 375 = *ignitarium*.
eliodes, Bieb. Fl. Taur. Cauc. iii. 555.—Rossia; Reg. Caucas.; Persia.
Empedoclearum, Lojac. in Natural. Sicil. iii. (1884) 284. —Sicil.
eriocephalum, A. Gray, in Proc. Acad. Sc. Philad. 1868 (1864) 69 = Cn. sciocephalus.
eriocephalum, Wallr. Beitr. Fl. Hercyn. 262 = Cn. eriophorus.
eriophorum, Scop. Fl. Carn. ed. II. ii. 130 = Cn. eriophorus.
Erisithales, Scop. Ann. ii. Hist. Nat. 69 = Cn. ochroleucus!
eristhaloides x, Huter, ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 271.—Tyrol.
erucagineum, DC. Fl. Fr. iv. 115 = Cn. rivularis.
erythrolepis, C. Koch, in Linnaea, xvii. (1843) 41.— Reg. Caucas.
esculentum, C. A. Mey. in Beitr. Pfl. Russ. Reich. iv. 48; et in Flora, xxxii. (1849) 747 = Cn. esculentus.
exaltatum x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 109.—Helvet.
exaspera, Fisch. Hort. Gorenk. ed. II. (1812) 85, nomen.—Hab. ?
falcatum, Turcz. ex DC. Prod. vi. 650 = Cn. pendulus.
fallax, Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc. (1838) 260.—Reg. Caucas.
fallax, Franch. in Billot, Annot. 109 = Cn. acaulis.
ferox, DC. Fl. Fr. iv. 120 = Cn. ferox.
ferox, Schur, ex Nym. Consp. 405 = Cn. ciliatus.
Ficfolium, Fisch. in Mém. Soc. Nat. Mosc. til. (1812) 89 = Serratula atriplicifolia.
filipendulum, Engelm. in A. Gray, Man. Bot. N. U. St. ed. V. 272 = Cn. altissimus.
filipendulum, Lange, in Kjoeb. Vidensk. Meddel. (1861) 92.—Hispan.
flabriatum, Spreng. Syst. iii. 373 = Cn. flabriatus.
Flerkeriauum, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 501, nomen.—Rossia.
flavum x, Auserd. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 263.—Tyrol.
flavescens, Koch, Taschenb. 393 = Cn. ochroleucus.
flavescens Lam. Fl. Fr. ii. 26 = Cn. parviflorus.

CIRSIUM :—
flavispinum, Bolss. in DC. Prod. vii. 305 = Cnicus flavispinus.
foliosum, DC. l. c. vi. 654 = Cn. foliosus.
Forsteri, Loud. Hort. Brit. Suppl. i. 585 = Cn. palustris.
fratrum, DC. Prod. vi. 636 = Cn. strigosus.
Freyerianum, Koch, Taschenb. 388; Syn. Fl. Germ. ed. II. 454.—Dalmat.
Frickii, Fisch. & Mey. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. vi. (1849) 47 = *rhizocephalum*.
fulvispinum, Walp. Rep. ii. 676, sphalm. = Cn. flavispinus.
furcatus, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) 1. 348 = Cn. ciliatus.
Gaillardoti, Bolss. Diagn. Ser. II. iii. 42.—Syria.
Galactites, Hill, Hort. Kew. ed. = Galactites tomentosa.
Genderi x, Huter, ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 274.—Tyrol.
Gerhardti, Sch. Bip. in Flora, xxxii. (1849) 547 = Cn. lanceolatus.
gigauteum, Spreng. Syst. iii. 375 = Cn. gigauteus.
glabrum, DC. Fl. Fr. Suppl. 468 = Cn. spinosissimus?
glaucescens x, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 237.—Tyrol.
glutinosum, Lam. Fl. Fr. ii. 27 = Cn. ochroleucus.
Gmelini, Tausch, in Flora, xi. (1828) 483 = Cn. acaulis.
graphalodes, Spreng. Syst. iii. 375 = Cn. graphalodes.
Gudroni, Sch. Bip. ex Nym. Consp. 406 = Cn. lanceolatus.
gracile, Rostr. ex Nym. l. c. = Cn. lanceolatus.
Grahami, Schur, Enum. Pl. Transs. 422 = Cn. rivularis.
Grahami, A. Gray, Pl. Wright. ii. 102 = Cn. Grahami.
grandiflorum x, Ritt. Taschenb. ed. II. 551.—Europ.
graveolens, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 189.—Hispan.
Gremblichii x, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 218.—Tyrol.
Griffithii, Bolss. Fl. Orient. iii. 640.—Afghanist.
grumosum, Fisch. & Mey. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. vi. (1849) 48 = *rhizocephalum*.
grumosum, Willk. & Lange, Prod. Fl. Hisp. ii. 199 = Cn. grumosus.
Gutnichianum x, Loehr, Enum. Fl. Deutsch. 865.— Europ.
Haller, Tausch, i. Flora, xix. (1836) 401.—Helvet.
Hallerianum, Gaudin, Fl. Helv. v. 188 = Cn. spinosissimus.
Hallerianum, Hill, Veg. Syst. iv. 27.—Helvet.
halophilum, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xlii. (1870) i. 83 = Cn. arvensis.
Hausmanni, Bolss. Fl. Orient. iii. 585.—Kurdistan.
helenioides, All. Fl. Pedem. i. 152 = Cn. heterophyllus.
helenioides, Hill, Hort. Kew. 64 = Cn. helenioides.
helleuicum, Bolss. & Orph. ex Bolss. Fl. Orient. iii. 630 = Cn. ferox.
Hermonis, Bolss. Diagn. Ser. II. iii. 40 = Cn. lappaceus.
heterolepis, Bench. Pl. Hartw. 97 = Cn. subcoriaceus.
heterophyllum, Spreng. Syst. iii. 373; Sweet, Hort. Brit. ed. I. 314 = Sanssurea candicans.
heterophylloides x, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 246.—Austr.
heterophyllum, Hill, Hort. Kew. 64; All. Fl. Pedem. i. 152 = Cn. heterophyllus.
heterotrichum, Panč. Elem. Fl. Bulg. 42.—Europ.
Hippolyti, Link, ex Stend. Nom. ed. II. i. 368 = Cn. pungens.
Hookerianum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 418 = Cn. Hookerianus.
horridulum, Michx. Fl. Bor. Am. ii. 90 = Cn. horridulus.
horridum, Fisch. Hort. Gorenk. ed. II. (1812) 85 = Cn. horridus.
horridum, Lag. ex DC. Prod. vi. 660.—Hispan.
Hugueninx x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 109.—Europ.
Hutert x, Hausm. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 231.—Europ.
hybridum, Koch, in DC. Fl. Fr. Suppl. 468 = Cn. echinocephalus.
hygrophilum, Bolss. Diagn. Ser. I. x. 89.—Persia.
hypoleucum, DC. Prod. vi. 648.—Balkan; As. Min.
hypoglaucum, Bolss. & Heldr. Diagn. Ser. I. vi. 101 = Cn. ferox.
ignitarium, Spreng. Syst. iii. 373.—Sibir. alaska.
incurvidens, Bolss. Fl. Orient. iii. 637.—Reg. Caucas.
incurvum, Bieb. Fl. Taur. Cauc. iii. 561 = Cn. arvensis.
inclinatum, Lam. Fl. Fr. ii. 23 = Carduus medius.

CIRSIUM :—
inermo, G. Don, in Lond. Hort. Brit. 330; Reichb. Fl. Germ. Excurs. 387 = Cnicus tuberosus.
inermo, Link, Enum. Hort. Berol. ii. 301 = Cn. cynaroides.
insubricum, Moretti, ex Bertol. Fl. Ital. ix. 26 = Cn. eriophorus.
involucratum, DC. Prod. vi. 639 = Cn. involucratus.
italicum, DC. Cat. Hort. Monsp. 96 = Cn. italicus.
Jaegeri x, F. Schultz, in Flora, xxxii. (1849) 238.— Europ.
japonicum, DC. Prod. vi. 640 = Cn. japonicus.
jorullense, Spreng. Syst. iii. 376 = Cn. jorullensis.
Jurinhi, Pacher, in Jahrb. Landesm. Kaernt. xv. (1882) 136.—Europ.
Jurinhoides x, Reichardt, in Verh. Zool.-Bot. Ges. Wien, xi. (1861) 280.—Austr.
kamtschaticum, Ledeb. ex DC. Prod. vi. 644 = Cn. kamtschaticus.
Kerneri x, Auserd. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 260.—Europ.
Killiani, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 108.—Helvet.
kirbyæ, Porsel, Nouv. Mat. Fl. Act. 274.—Afr. bor.
Kochianum, Loehr, in Flora, xxv. (1842) 111 = Cn. palustris.
Kornulii, Bolss. Fl. Orient. iii. 581, in syn. = Cn. lappaceus.
Kotschyanum, Boiss. Diagn. Ser. I. vi. 108 = *libanoticum*.
Lachenalii, DC. Prod. vi. 648, in syn. = Cn. rigens.
Lachenalii, Koch, Syn. Fl. Germ. ed. I. 397 = Cn. acaulis, oleraceus.
lacinatum, Doell, ex Nym. Consp. 409 = Cn. palustris.
lactum x, Ruhm. in Eichl. Jahrb. Bot. Gart. Berl. i. (1881) 241.—Europ.
lactum, Schleich. Cat. (1821), ex Koch, Syn. Fl. Germ. ed. I. 394 = palustris.
lasor, Moench, Meth. Suppl. 227 = Cn. tataricus.
lasvigatum, Tausch, in Flora, xi. (1828) 483 = Cn. arvensis.
lamprophyllum, Herbich, Stirp. Rar. Bucov. 54.— Bukovina.
lanatum, Spreng. Syst. iii. 373 = Cn. arvensis.
lanceolatum, Hill, Herb. Brit. i. 80; Scop. Fl. Carn. ed. II. ii. 130 = Cn. lanceolatus.
lanifloram, Bieb. Fl. Taur. Cauc. iii. 557 = Cn. lanifloras.
lanigerum, Naegeli, Cirs. Schweiz. 114. t. 3 = Cn. lanceolatus.
lappaceum, Fisch. Hort. Gorenk. ed. II. (1812) 85 = Cn. lappaceus.
lappaceum, Lam. Fl. Fr. ii. 24 = Carduus Personata.
lappoides, Sch. Bip. in Seem. Bot. Voy. Herald, 319 = Cn. lappoides.
latifolium, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 28.—Ins. Madera.
Lecouti, Torr. & Gray, Fl. N. Am. ii. 458 = Cn. Lecouti.
Ledebourii, Steud. Nom. ed. II. i. 368.—Sibiria.
leucanicum, Lojac. in Natural. Sicil. iii. (1884) 283.— Sicilia.
leucocephalum, C. Koch, in Linnaea, xvii. (1851) 401 = *chrysiticum*.
leucocephalum, Sprung. Syst. iii. 377 = Cn. leucocephalus.
leucocrenum, Bolss. & Heldr. in Bolss. Fl. Orient. iii. 584.—Catsonia.
leucophaenum, Schur, Enum. Pl. Transs. 418 = Cn. lanifloras.
leucopsis, DC. Prod. vi. 644.—As. Min.
leucostylum, Mocench, Meth. 556 = Carduus deflonatus.
libanoticum, DC. Prod. vi. 647.—As. Min.; Syria.
lignitare, Bolss. Fl. Orient. iii. 630.—Thracia.
Linnéi, Nym. Syll. 29.—Lusitan.
linearo, Sch. Bip. in Linnaea, xix. (1847) 365 = Cn. linearis.
litorale, Maxim. Prim. Fl. Amur. 178 = Cn. japonicus.
Linkianum x, Loehr, Enum. Fl. Deutsch. 864.—Europ.
Lobelii, Tenore, Fl. Neap. v. 211. t. 189 = Cn. ferox.
longepinnatum, Tod. ex Nym. Consp. 406 = Cn. lanceolatus.
longifolium, Bolss. Fl. Orient. iii. 582, sphalm. (= *Hypochoeris longifolia*).—Reg. Caucas.
lucidum, DC. Prod. vi. 654.—Hab. ?
lutescens, Rafin. in Med. Repos. N. York, II. v. (1806) 354, nomen.—Hab. ?
lycaonicum, Bolss. & Heldr. Diagn. Ser. I. x. 90 = *libanoticum*.

CIRSIUM :—
lyratum, Bunge, Enum. Pl. Chin. Bor. 36 = Saussurea affinis.
Maackii, Maxim. Prim. Fl. Amur. 173 = Cn. japonica.
macracanthum, Sch. Bip. in Flora, xxxv. (1852) 185 = Cn. Wallichii.
macrobotrys, Boiss. Fl. Orient. iii. 582.—Armenia.
macrocephalum, C. A. Mey. Verz. Pfl. Cauc. 70 = Cn. munitus.
macrochaetum, Tausch, in Flora, xix. (1836) 397.—Hab.?
macrostachyum, Boiss. Fl. Orient. iii. 550.—Reg. Caucas.
macrotylon, Reichb. Fl. Germ. Excurs. 856 = Cn. arvensis.
maculatum, Lam. Fl. Fr. ii. 22 = Tyrimnus leucographus.
maculatum, Moench, Meth. 557 = Cn. syriacus.
maculatum, Scop. Fl. Carn. ed. II. ii. 130 = Silybum Marianum.
marocitanum, Spreng. Syst. iii. 377 = Carthamus marocitanus.
matritense, Lag. ex Cut. Fl. Comp. Madrid, 409 = Cn. flavispina.
maximum, Benth. Pl. Hartw. 289 = Cn. suborciaceus.
medium, All. Fl. Pedem. i. 149 = Cn. tuberosum.
megacanthum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 421 = Cn. horridulus.
mexicanum, DC. Prod. vi. 636 = Cn. mexicanus.
micranthum ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 218.—Europ.
microcephalum, Lange, Ind. Sem. Hort. Havn. (1854) 22.—Hispan.
microcephalum, Moris, Fl. Sard. ii. 466 = Cn. microcephalus.
microcephalum, Sch. Bip. in Oestr. Bot. Zeitschr. viii. (1858) 246 = Cn. siculus.
Mielichhoferi, Sauter, in Flora, xxviii. (1845) 180 = Cn. heterophyllus.
misilmerense, Ces. Passer. & Gib. Comp. Fl. Ital. 483 = Cn. misilmerensis.
mite, Fisch. in Spreng. Nov. Prov. 18 = Cn. mitis.
molle, Scop. Fl. Carn. ed. II. ii. 126 = Jurinea mollis.
monspessulanum, Hill, Hort. Kew. 68 ; All. Fl. Pedem. i. 159 = Cn. monspessulanus.
montanum, Hill, Herb. Brit. i. 80 = Cn. montanus.
Moretticanum, Nym. Syll. 24 = Cn. spathulatus.
morinaefolium, Boiss. & Heldr. ex Boiss. Fl. Orient. iii. 580.—Ins. Creta.
Morisianum, Reichb. f. lc. Fl. Germ. xv. 59.—Europ.
munitum, Fisch. Hort. Gorenk, ed. II. (1812) 86 = Cn. munitus.
mutatum, Menyh. Kalocsa Vidék. Növény. 109 = Cn. arvensis.
muticum, Michx. Fl. Bor. Am. ii. 89 = Cn. muticus.
Neapolitanum ×, Loehr, Enum. Fl. Deutsch. 362.—Europ.
neglectum, Fisch. ex Spreng. Syst. iii. 370 = Cn. arvensis.
nemorale, Reichb. Fl. Germ. Excurs. 856 = Cn. lanceolatus.
neo-mexicanum, A. Gray, Pl. Wright, ii. 101 = Cn neo-mexicanus.
nepalense, DC. Prod. vi. 642 = Cn. Wallichii.
nevadense, Willk. in Linnaea, xxx. (1859–60) 111.—Hispan.
nidulans, Regel, in Bull. Soc. Nat. Mosc. xl. (1867) ii. 160.—As. centr.
nivale, Sch. Bip. in Seem. Bot. Voy. Herald, 312 = Cn. rivalis.
nivium, Spreng. Syst. iii. 378 = Cn. nivens.
numidicum, Spreng. l. c. 377.—Afr. bor.
Nuttalii, DC. Prod. vi. 651 = Cn. Nuttallii.
oblongifolium, C. Koch, in Linnaea, xvii. (1843) 41.—Reg. Caucas.
obscurum ×, Kern. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 264.—Europ.
obvallatum, Hab. Fl. Taur. Cauc. iii. 549 = Cn. obvallatus.
ochro-centrum, A. Gray, Pl. Fendl. 110 = Cn. ochrocentrus.
ochroleucum, All. Fl. Pedem. i. 150 = Cn. ochroleucus.
odontolepis, Boiss. ex DC. Prod. vii. 606.—Hispan. ; Baliear.
ommanum ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 215.—Europ.
ompbaloides ×, Treuinf. l. c. 220.—Europ.
oleraceum, Scop. Fl. Carn. ed. II. ii. 124 = Cn. oleraceus.

CIRSIUM :—
onopordioides, Kotschy, ex Boiss. Fl. Orient. iii. 564 = Onopordum polycephalum.
ornithales, Hance, ex Walp. Ann. ii. 944 = Cnicus chinensis.
orgyale, Hort. ex Tausch, in Flora, xii. (1829) 1. Erg. 41 = *aciculare* ?
orgyale, Link, Enum. Hort. Berol. ii. 302 = Cn. orgyalis.
oriens, Spreng. Syst. iii. 378 = Galactites tomentosa.
palatinum, Sch. Bip. ex Nym. Consp. 409 = Cn. palustris.
pallens, DC. Prod. vi. 647 = Cn. oleraceus.
paludosum, G. Don, in Loud. Hort. Brit. 381 = Cn. paludosus.
palustre, Cos. ex Willk. & Lange, Prod. ii. 191 = *Bourgeanum*.
palustre, Scop. Fl. Carn. ed. II. ii. 128 = Cn. palustris.
paniculatum, Boiss. & Hausk. ex Boiss. Fl. Orient. iii. 535 = *catanonicum* !
paniculatum, Lam. Fl. Fr. ii. 25 = Carduus carlinoides.
paniculatum, Spreng. Syst. iii. 371.—Mont. Pyren.
paniculatum, Vahl, ex Willk. & Lange, Prod. Fl. Hisp. ii. 186 = Cn. flavispinus.
pannonicum, Link, Enum. Hort. Berol. ii. 301 = Cn. pannonicus.
paradoxum ×, Wallr. Beitr. Fl. Hercyn. 264.—Europ.
parnifforme, DC. Prod. vi. 647 = Cn. oleraceus.
parviflorum, Lange, ex Nym. Consp. 409 = Cn. palustris.
pauciflorum, Schleich. ex Steud. Nom. ed. II. i. 369 = Cn. ochroleucus.
pauciflorum, Koch, Syn. Fl. Germ. ed. I. 395 = Cn. heterophyllus.
pauciflorum, Lam. Fl. Fr. ii. 22 = Carduus defloratus.
pauciflorum, Spreng. Syst. iii. 375 = Cn. pauciflorus.
pascuorense, Spreng. l. c. 372 = Cn. pascuarensis.
pectinatum, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 395 = Cn. pectinellus.
pendulum, Fisch. ex DC. Prod. vi. 650 = Cn. pendulus.
pentlandii, C. Koch, in Linnaea, xvii. (1843) 41 = Cn. lappaceus.
persicum, Boiss. Diagn. Ser. I. vi. 103 = Cn. strigosus.
phyllocephalum, Boiss. & Blanche, in Boiss. l. c. Ser. II. iii. 89.—Syria.
pinnatifidum, Spreng. Syst. iii. 395 = Serratula pinnatifida.
Pitcheri, Torr. & Gray, Fl. N. Am. ii. 456 = Cn. Pitcheri.
platycephalum, Benth. ex Oerst. in Kjoeb. Videnak. Meddel. (1859) 110 = Cn. suborciaceus.
polyacanthum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 456 = Cn. polyacanthus.
polyacanthum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 460.—Sibir. altaic.
polyanthemum, Spreng. Syst. iii. 371 = Cn. polyanthemus.
polyanthos, Urv. in Mém. Soc. Linn. Par. i. (1822) 368 = Cn. pungens.
polycephalum, DC. Prod. vi. 639 = *lyantinum*.
polycephalum, Willd. ex Spreng. Syst. iii. 371.—Hab.?
polymorphum, Doll. ex Hausm. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 875 = *Linhianum*.
ponticum ×, A. Mey. Ind. Sem. Hort. Petrop. ix. 67 = *polycephalum*.
Portae, Hausm. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 872 = *Linhianum*.
praealtum, Cass. in Dict. Sc. Nat. xli. 333 = Cn. arvensis.
praemorsum, Koch, Syn. Fl., Germ. ed. I. 397 = Cn. ochroleucus ?
pratense, DC. Fl. Fr. iv. 113 = Cn. monspessulanus.
propinticum, Griseb. Spicil. Fl. Rumel. ii. 354 = Cn. eriophorus.
pseudopumilum, Schur, Enum. Pl. Transs. 421 = Cn. pannonicus.
pseudo-lanceolatum, Schur, l. c. 418 = Cn. lanceolatus.
pseudo-oleraceum, Schur, l. c. 423 = Cn. pannonicus.
pseudo-palustre, Schur, l. c. 420 = *Chailletii*, Koch.
pseudo-pannonicum, Schur, l. c. = Cn. canus.
pseudo-pyrenaeum, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 548.—As. Min.
pubigerum, DC. Prod. vi. 650 = Cn. pubigerus.
Pumilio, Tausch, in Syll. Ratisb. ii. (1828) 249.—Austria.
pumilum, Spreng. Syst. iii. 375 = Cn. pumilus.

CIRSIUM :—
pungens, Schur, Enum. Pl. Transs. 420.—Europ.
pungens, Spreng. Syst. iii. 370 = Cnicus polyanthemus.
purpureum, All. Fl. Pedem. i. 150 = Cn. spinosissimus.
Pustaricaum ×, Ausserd. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 258.—Europ.
pygmaeum, Scop. Fl. Carn. ed. II. ii. 128 = Saussurea pygmaea.
pyrenaicum, All. Fl. Pedem. i. 151 = Cn. montanus.
pyrenaicum, DC. Fl. Fr. iv. 113 = C. pyrenaicus.
racemosum, Boiss. in Tchihat. Asie Min. Bot. ii. 388 = *tenuilobum*.
radians, Benth. Pl. Hartw. 77 = Cn. radians.
Reichardtii ×, Jurotzka, in Verh. Zool.-Bot. Ges. Wien, ix. (1859) 317.—Europ.
Reichenbachianum ×, Loehr, Enum. Fl. Deutsch. 364 = Cn. arvensis.
remotifolium, DC. Prod. vi. 655 = Cn. remotifolius.
repandum, Michx. Fl. Bor. Am. iii. 28 = Cn. repandus
replicatum, DC. Prod. vi. 655 = Cn. replicatus.
rhegninum, Spreng. Pugill. ii. 77 = Cn. strictus.
rhizocephalum, C. A. Mey. Verz. Pfl. Cauc. 70.—As Min.; Persia; Afghan.
rhodo-leucanthum, Walp. Ann. i. 453 = Cn. canus.
Richterianum, Gillot, in Bull. Soc. Bot. Fr. xxvii. (1880) p. lii.—Europ.
rigens, Spreng. Neue Entdech. iii. 88 = Cn. oleraceus.
rigens, Wallr. Sched. Crit. 446 = Cn. acaulis.
rigidum, DC. Prod. vi. 640 = Cn. arachnoideus.
rivulare, All. Auct. Pedem. 10 = Cn. montanus.
rivulare, Link, Enum. Hort. Berol. ii. 301 = Cn. rivalis.
Rosani, Tenore, Ind. Sem. Hort. Neap. (1830) 15: ex Syll. 414 = Cn. lanceolatus.
Rosenii, Vill. Hist. Pl. Dauph. iii. 14. t. 21 = Cn. acaulis ?
rufescens, Ramond, ex DC. Fl. Fr. iv. 114 = Cn. rufescens.
ruthenicum, Fisch. Hort. Gorenk. ed. II. (1812) 85 = Cn. arvensis.
sabaudum ×, Loehr, Enum. Fl. Deutsch. 362.—Europ.
sabaudum ×, Ruhm. in Eichl. Jahrb. Bot. Gard. Berl. i. (1881) 240.—Europ.
salamanticum, Hill, Hort. Kew. 64 = Centaurea salmantica.
salinum, Bieb. ex Steud. Nom. ed. II. i. 370 = Centaurea salina.
salisburgense, G. Don, in Loud. Hort. Brit. 381 = Cn. rivularis.
samuticum, Steud. Nom. ed. II. i. 370 = Cn. italicus.
scariosum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 420 = Cn. scariosus.
schanginense, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 58).—Sibir. or.
Schultzianum ×, Loehr, Enum. Fl. Deutsch. 365.
scleranthum, Bieb. Fl. Taur. Cauc. iii. 559.—As. Min. ; Persia.
Scopolianum, Sch. Bip. ex Nym. Consp. 408 = Cn. Eristhales.
segetum, Bunge, Enum. Pl. Chin. Bor. 36 = Cn segetum.
Semenowi, Regel, in Bull. Soc. Nat. Mosc. xl. (1867) ii. 161.—As. centr.
semidecurrens, DC. Prod. vi. 646.—Hab.?
semidecurrens, Richt. Fl. Lips. 378 = Cn. palustris.
semipectinatum, Ledeb. ex Steud. Nom. ed. II. i. 370 = *Ledebourii*.
semipectinatum, Reichb. Fl. Germ. Excurs. 857 = Cn. rivularis.
semipectinatum, Spreng. Syst. iii. 378 = Cn. semipectinatus.
serratuloides, Hill, Hort. Kew. 64 = Cn. serratuloides.
serratuloides, Scop. Fl. Carn. ed. II. ii. 127 = Cn. pannonicus.
serratum, Tausch, in Flora, xii. (1829) 1. Erg. 39.—Europ.
serrulatum, Bieb. Fl. Taur. Cauc. iii. 557 = Cn. serrulatus.
serrulatum, Schur, Enum. Pl. Transs. 419 = Cn. fimbriatus.
sessilifolium, Herbich, Stirp. Rar. Bucov. 54.—Turcia.
setigerum, Ledeb. lc. Pl. Fl. Ross. i. 9.—Sibir. altaic.
setosum, Bieb. Fl. Taur. Cauc. iii. 560 = Cn. arvensis.
siculum, Spreng. Neue Entdeck. iii. 86 = Cn. pungens.
Sieboldi, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 184 = Cn. Sieboldii.
Siegerti, Sch. Bip. ex Nym. Consp. 406 = Cn. rivularis.

CIRSIUM :—
silesiacum, Sch. Bip. ex Nym. Consp. 409 = Cnicus paleacris.
sisauticum, Tausch, in Flora, xii. (1829) 1. Erg. 86 = Cn. lanceolatus.
simplex, Hornem. Hort. Hafn. Suppl. 98 = Centaurea salina.
simplex, C. A. Mey. Verz. Pfl. Cauc. 70.—Reg. Caucas.
sinense, C. B. Clarke, Comp. Ind. 219 = Cn. chinensis.
sinuatum, Bolss. Fl. Orient. iii. 546 = Cn. cynaroides.
sinuatum, Bolss. & Kotschy, ex Bolss. l. c. 542 = *rhiscophalum*.
sordidum, Wallr. Beitr. Fl. Hercyn. 266 = Cn. arvensis.
sorocephalum, DC. Prod. vi. 636 = *fallax*, Fisch. & Mey.
sorocephalum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 38.—Reg. Caucas.
spathulatum, Gandin, Fl. Helv. v. 202 = Cn. eriophorus.
spectabile, DC. Prod. vi. 641.—Persia.
spinifolium x, Beck, in Oestr. Bot. Zeitschr. xxxi. (1881) 312.—Europ.
spinosissimoides x, Auszerd. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 250.—Europ.
spinosissimum, Scop. Fl. Carn. ed. II. ii. 129 = Cn. spinosissimus.
Spitzelii, Sch. Bip. ex Nym. Consp. 408 = Cn. spinosissimus.
spurium, Delastre, in Ann. Sc. Nat. Sér. II. xviii. (1842) 149 = Cn. pratensis.
squarrosum, Fisch. ex Spreng. Syst. iii. 376 = Cn. squarrosus.
stellatum, All. Fl. Pedem. i. 158 = Cn. stellatus.
stenolepidum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 419 = Cn. remotifolius.
stenophyllum, Bolss. Fl. Orient. iii. 534 = *tenuilobum*.
Stschegl, Bolss. Diagn. Ser. II. iii. 40 = Cn. arvensis
strictum, Moris, Stirp. Sard. Elench. i. 27 = Cn. microcephalus.
strictum, Spreng. Syst. iii. 370 = Cn. strictus.
strigosum, Bieb. Fl. Taur. Cauc. iii. 558 = Cn. strigosus.
subalatum, Gaudin, Fl. Helv. v. 180 = Cn. oleraceus.
subalpinum, Gaudin. l. c. 182 = Cn. palustris.
subalpinum, Schleich. ex Reichb. Ic. Fl. Germ. xv. 78. —Helvetia.
subcoriaceum, Sch. Bip. ex Seem. Bot. Voy. Herald, 213 = Cn. subcoriaceus.
subinerme, Fisch. & Mey. in Mém. Acad. Petersb. Ser. VI. Sc. Nat. vi. (1849) 55 = *slodes*.
subnigrifolium, Schur, Enum. Pl. Transs. 421 = Cn. pannonicus.
succinctum, Ledeb. Fl. Ross. ii. 782 = Cn. lappaceus.
superstratum, Tausch, in Flora, xix. (1836) 397.—Hab. ?
sylvaticum, DC. Prod. vi. 656 = Cn. lanceolatus.
syriacum, Gaertn. Fruct. ii. 383. t. 163. f. 3 = Cn. syriacus.
Szovitsii, Bolss. Fl. Orient. iii. 537.—Reg. Caucas.
tartaricum, All. Fl. Pedem. i. 151 = Cn. tataricus.
tataricum, DC. Fl. Fr. iv. 114 = Cn. zigeus.
tauricolum, Bolss. & Bal. ex Bolss. Fl. Orient. iii. 551. —Catacnia.
Tauschii, DC. Prod. vi. 654.—Hab. ?
tchefouense, Debeaux, Fl. Tohef. 86 = Cn. tchefouensis.
tenuilobum, Bolss. Fl. Orient. iii. 534.—Armenia.
tenuinum, Buckl. in Proc. Acad. Sc. Philad. '1861 (1862) 460 = Cn. virginianus.
Thomasii, Nagg. Circ. Schweiz. 93 = Cn. oleraceum.
tiroleuse x, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 243.—Europ.
tomentosum, C. A. Mey. Verz. Pfl. Cauc. 69 = Cn. mucutica.
tomentosum, Moench, Meth. Suppl. 237 = Juricea cynaroides.
trachylepis, Bolss. Fl. Orient. iii. 528.—Armenia.
transsilvanicum, Schur, Enum. Pl. Transs. 419 = Cn. transilvanicus.
tricephalodes, DC. Fl. Fr. iv. 116 = Cn. rivularis.
tricephalum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = prec.
trichotomum, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 251.—Reg. Caucas.
trispicatum, Moench, Meth. 656 = Cn. Casabonae.
tryphillinum x, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 276.—Europ.
tuberosum, All. Fl. Pedem. i. 151 = Cn. tuberosus.

CIRSIUM :—
ukranicum, Bess. ex DC. Prod. vi. 635 = Cnicus serrulatus.
uliginosum, Bieb. Fl. Taur. Cauc. iii. 555 = Cn. uliginosus.
uliginosum, Heldr. ex Bolss. Diagn. Ser. I. x. 90 = *thasaeticum*.
undulatum, Spreng. Syst. iii. 374 = Cn. undulatus.
Vallis-Demonsii, Lojac. in Natural. Sicil. iii. (1884) 267. —Sicilia.
vaviloffii, Moench, Meth. 658 = Cn. oleraceus.
Verutum, Spreng. Syst. iii. 370 = Cn. argyracanthus.
virens, Timb. & Jeanb. ex Nym. Consp. 406 = Cn. lanceolatus.
virgatum, Tausch, in Flora, xix. (1836) 402.—Tauria.
virginianum, Hook. Comp. Bot. Mag. i. (1835) 48 = Cn. Lecontei.
virginianum, Michx. Fl. Bor. Am. ii. 90 = Cn. virginianus.
Vlassovianum, Fisch. ex DC. Prod. vi. 653 = Cn. Vlassovianus.
Wallichii, DC. l. c. 648 = Cn. Wallichii.
Wankelii x, Reichardt, in Verh. Zool.-Bot. Ges. Wien, xi. (1861) 881.—Europ.
Welwitschii, Cos. Notes Crit. 118.—Hispania.
Weyrichii, Maxim. Prim. Fl. Amur. 174 = Cn. Weyrichii.
Wiedemannii, Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop. vi. 48 = *hypoleucum*.
Willdenowii, Steud. Nom. ed. II. i. 370 = Cn. Willdenowii.
Wimmeri, Celak. in Arch. Naturfw. Landesd. Boehm. ii. (1872) 268.—Bohemia.
Winklerianum x, Celak. l. c. 262.—Europ.
Wirtgenii, Lasch, Enum. Fl. Deutsch. 863.—Europ.
Woodwardii, H. C. Wats. ex Nym. Consp. 407 = Cn. tuberosus.
Wrightii, A. Gray, Pl. Wright. ii. 101 = Cn. Wrightii.
Zieianum, Koch, Syn. Fl. Germ. ed. I. 398 = Cn. tuberosus.

CISSAMPELOPSIS, Miq. Fl. Ind. Bat. ii. 102 (1856) = Senecio, Toura. (Compos.).
volubilis, Miq. l. c. = S. araneosus.

CISSAMPELOS, Linn. Gen. ed. I. 368 (1737). *MENISPERMACEAE*, Benth. & Hook. f. i. 37.
ANTIDOTA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 41.
CAAPEBA, Plum. ex Adans. Fam. ii. 357 (1763).
DISSOPETALUM, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 367.
Abutua, Vell. Fl. Flum. x. t. 140 = Chondodendron platyphyllum.
acuminata, Benth. Pl. Hartw. 58 = C. Pareira.
acuminata, *DC. Syst.* i. 538.—Ind. or.
acuta, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 48.—N. Granat.
amazonica, Miers, ex Benth. in Hook. Kew Journ. iii. (1851) 114 = ovalifolia.
andromorpha, *DC. Syst.* i. 539.—Guiana.
angustifolia, *Turcz. Trav.* i. 389.—Afr. austr.
angustifolia, Drège, ex Harv. & Sond. Fl. Cap. i. 13 (= *Antisoma Mierriana*).—Afr. austr.
apiculata, Hochst. in Flora, xxiii. (1845) 93 = C. Pareira.
argentea, H. B. & K. Nov. Gen. et Sp. v. 67 = C. Pareira.
ariteroides, *Casaffola*, Fenzl, in Flora, xxvii. (1844) 312 = C. Pareira.
asiatica, Miers, ex Benth. in Hook. Kew Journ. iii. (1851) 114 = ovalifolia.
auriculata, *Miers, Contrib.* iii. 158.—Reg. Argent.
australis, St. Hil. Fl. Bras. Mer. i. 54 = C. Pareira.
barbata, Wall. Cat. n. 4972 = Cyclea peltata.
Benthamiana, *Miers, Contrib.* iii. 146.—Mexic.
Boivini, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 460. —Madag.
Bojeriana, *Miers, Contrib.* iii. 152.—Ins. Maurit.
Caapeba, Linn. Sp. Pl. 1032 = C. Pareira.
Caapeba, Vell. Fl. Flum. x. t. 139 = fasciculata ?
calcariformis, Buch. Trav. ii. 256 = fruticosa.
canescens, Miq. Sert. Kunt. i. 7. t. 4 = C. Pareira.
capensis, *Thumb. Prod. Pl. Cap.* 116.—Afr. austr.
capitata, *Griff. Notul. iv.* 209.—Ind. or.
clematidea, Presl, Rel. Haenk. 7 = C. Pareira.
Cocculus, Buch.-Ham. ex Steud. Nom. ed. II. i. 372 = C. Pareira.

CISSAMPELOS :—
Cocculus, *Poir. Encyc.* v. 9.—Ind. or.
Cocculus, Steud. Nom. ed. I. 198 = Anamirta paniculata.
comata, Miers, in Hook. Niger Flora, 215 = C. Pareira.
communis, St. Hil. Fl. Bras. Mer. i. 52 = ovalifolia.
consociata, *Miers, Contrib.* iii. 167.—Ind. occ.
convexa, Vell. Fl. Flum. x. t. 142 = Abuta rufescens.
convolvulacea, Willd. Sp. Pl. iv. 868 = C. Pareira.
cordifolia, *Boj. Ann. Sc. Nat.* Sér. II. xx. (1843) 94.—Afr. austr.
crenata, DC. Syst. i. 537 = ovalifolia.
Cumingiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 288.—Ins. Philipp.
delicatula, Miers, Contrib. iii. 197 = C. Pareira.
densiuscula, Miers, ex Benth. in Hook. Kew Journ. iii. (1851) 115 = fasciculata.
diffusa, *Miers, Contrib.* iii. 168.—Ind. occ.
discolor, DC. Syst. i. 534 = Stephania hernandifolia.
discolor, A. Gray, Bot. U. St. Expl. Exped. i. 86 = C. Pareira.
discolor, Wall. Cat. n. 4982 = Cyclea peltata.
diversa, Miers, Contrib. iii. 187 = C. Pareira.
ebracteata, St. Hil. Pl. Us. Bras. t. 35 = ovalifolia.
ciata, Miers, Contrib. iii. 187 = C. Pareira.
eriantha, Miers, l. c. 192 = C. Pareira.
eriocarpa, Triana & Planch. in Ann. Sc. Nat. Ser. IV. xvii. (1862) 41 = C. Pareira.
errabunda, *Miers, Contrib.* iii. 138.—Bras.
fasciculata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 561.—Bras.
floribunda, *Miers, Contrib.* iii. 164.—Peruv.
fluminensis, *Zichi. in Flora*, xlvii. (1864) 593 [*rec. typ* ! *823* ?] *et in Mem. Fl. Bras.* xiii. t. 191.—Bras.
fruticosa, Thunb. Prod. Pl. Cap. 110 = capensis.
glabertima, *St. Hil. Fl. Bras. Mer.* i. 57.—Bras.
glabra, Buch.-Ham. ex Wight & Arn. Prod. i. 14 = Clypea hernandifolia et Cocculus Roxburghianus.
glabra, Roxb. Hort. Beng. 74; Fl. Ind. iii. 840 = Stephania rotunda.
glaucescens, Triana & Planch. in Ann. Sc. Nat. Ser. IV. xvii. (1862) 41 = C. Pareira.
gracilis, St. Hil. Fl. Bras. Mer. i. 56 = C. Pareira.
grallatoria, Miers, Contrib. iii. 188 = C. Pareira.
grandifolia, Triana & Planch. in Ann. Sc. Nat. Ser. IV. xvii. (1862) 44 = C. Pareira.
guapayulensis, H. B. & K. Nov. Gen. et Sp. v. 67 = C. Pareira.
Haenkeana, Presl, Rel. Haenk. ii. 80 = ovalifolia.
hederacea, *Miers, Contrib.* iii. 189.—Reg. Argent.
Hernandia, *Vell. Fl. Flum.* x. t. 136.—Bras.
hernandifolia, Wall. Cat. 4977, partim = C. Pareira.
hernandifolia, Willd. Sp. Pl. iv. 861 = Stephania hernandifolia.
heterophylla, DC. Syst. i. 534 = C. Pareira.
hexandra, Roxb. Hort. Beng. 74; Fl. Ind. iii. 841 = Stephania hernandifolia.
hirsuta, Buch.-Ham. in Trans. Soc. Linn. Syst. i. 535 = C. Pareira.
hirsutissima, Presl, Rel. Haenk. ii. 80 = ovalifolia.
hirta, Klotzsch, in Peters, Reise Mossamb. Bot. 174 = C. Pareira.
humilis, Poir. Encyc. v. 11 = capensis.
hypoglauca, *Schum. in Nov. Act. Acad. Nat. Cur.* xix. Suppl. i. (1843) 478.—Guiana.
Insolita, *Miers, Contrib.* iii. 179.—Afr. trop.
kohautiana, Presl, Rel. Haenk. ii. 81 = C. Pareira.
lanrifolia, *Poir. Encyc.* v. 11.—Ind. occ.
linhata, *Miers, Contrib.* iii. 143.—Afr. trop.
littoralis, St. Hil. Fl. Bras. Mer. i. 54 = C. Pareira.
longipes, *Miers, Contrib.* iii. 188.—Ind. occ.
macrostachya, Klotzsch, in Peters, Reise Mossamb. Bot. 178 = C. Pareira.
madagascariensis, *Miers, Contrib.* iii. 161.—Madag.; Ins. Borb.
mallophylla, *Miers, l. c.* 169.—Am. trop.
mauritiana, Thou. in Desv. Journ. Bot. ii. (1809) 67. t. 8, 4 = C. Pareira.
mauritiana, Wall. Cat. n. 4980 = Pericampylus incanus.
microcarpa, DC. Syst. i. 534 = C. Pareira.
mucuna, St. Hil. Fl. Bras. Mer. i. 55 = C. Pareira.
mucronata, A. Rich. in Guill. & Perr. Fl. Senog. Tent. i. 31 = C. Pareira.
myriantha, Triana & Planch. in Ann. Sc. Nat. Ser. IV. xvii. (1862) 42 = C. Pareira.
nephrophylla, Baj. t. a. Ser. II. xx. (1843) 95 = C. Pareira.

CISSAMPELOS :—
nymphæifolia, R. Br. in Salt, Abyss. App. 65 = Stephania abyssinica.
obtecta, Wall. Cat. n. 4981 = C. Pareira.
olivacea, Wall. l. c. n. 4984 = Parabœna sagittata.
orbiculata, DC. Syst. i. 537 = C. Pareira.
orinocensis, H. B. & K. Nov. Gen. et Sp. v. 68 = C. Pareira.
ovalifolia, *DC. Syst.* i. 537.—Bras.
ovata, Poir. Encyc. v. 10 = Pachygone ovata.
ovata, Vell. Fl. Flum. x. t. 141 = Abuta Selloana ?
ovaviensis, Benav. ex DC. Prod. i. 100 = C. Pareira.
pannosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 388.—Cuba.
Pareira, *Linn. Sp. Pl.* 1031.—Reg. trop.
parvifolia, DC. Enu. Med. Pl. Comp. ed. II. 78 = C. Pareira.
Parriera, Vell. Fl. Flum. x. t. 138 = glaberrima.
Pata, Roxb. ex Wight & Arn. Prod. i. 15 = C. Pareira.
pauciflora, Nutt. in Journ. Acad. Philad. vii. (1834) 114 = Croomia pauciflora.
peltata, Reis, ex Eichl. in Mart. Fl. Bras. xiii. t. 191 = tropæolifolia.
pellophylla, *Presl, Rel. Haenk.* ii. 80.—Ins. Luzon.
ramiflora, *Miers, Contrib.* iii. 168.—Bras.
rotundata, Pohl, ex Eichl. in Mart. Fl. Bras. xiii. t. 187 = ovalifolia.
sagittata, Buch.-Ham. ex Wall. Cat. n. 4983 = Parabœna sagittata.
Salemanni, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 284.—Bras.
scutigera, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 49 = C. Pareira.
senensis, Klotzsch, in Peters, Reise Mossamb. Bot. i. 178 = C. Pareira.
smilacina, Lam. Sp. Pl. 1032 = Menispermum canadense.
subcrenata, *Klotzsch, in Schomb. Fauna, Fl. Gui.* 1162, nomen.—Guiana.
suberbicularis, St. Hil. Fl. Bras. Mer. i. 58 = ovalifolia.
subpeltata, Thw. Enum. Pl. Zeyl. 18, 399 = C. Pareira.
subreniformis, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 41 = C. Pareira.
subtriangularis, St. Hil. Fl. Bras. Mer. i. 51 = ovalifolia.
sympodiales, *Eichl. in Flora*, xlvii. (1864) 382 [err. typ. 302].—Bras.
tamaulipica, *Miers, Contrib.* iii. 185.—Afr. austr.
tamoides, Sagot, ex Eichl. in Mart. Fl. Bras. xiii. t. 194 = fasciculata.
tamoides, *Willd. ex DC. Syst.* i. 536.—Bras.
testudinaria, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 134, nomen = seq.?
tetudinum, *Miers, Contrib.* iii. 148.—Ins. Galapag.
tetrandra, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 842 = C. Pareira.
tomentosa, DC. Syst. i. 535 = C. Pareira.
tomentosa, Vell. Fl. Flum. x. t. 143 = Abuta rufescens.
torulosa, *S. Mey. ex Harv. & Sond. Fl. Cap.* i. 11.—Afr. austr.
triloba, *Sprong. Nouv Entdeck.* ii. 152.—Bras.
tropæolifolia, *DC. Syst.* i. 538.—Am. austr.
umbellata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 10 = Stephania hernandifolia.
velutina, St. Hil. Fl. Bras. Mer. i. 59 = ovalifolia.
ventita, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1863) 44 = ovalifolia.
Vitis, *Vell. Fl. Flum.* x. t. 137.—Bras.
Vogelii, Miers, in Hook. Niger Fl. 214 = C. Pareira.
zahnata, *Miers, Contrib.* iii. 180.—Afr. trop.

CISSARRYON, Meisen. Gen. 68; Comm. 42 (1837) = seq.?

CISSAROBRYON, Poepp. Fragm. Syn. Fl. Chil. 29 (1839) = Wrightia, Cav. (Geraniac.).
elegans, Kunze, l. c. 30.—Chili.
macrophyllum, Phil. in Linnæa, xxxiii. (1864-65) 98. —Chili.
parvifolium, Phil. l. c. 98.—Chili.

CISSOCHYTRON, F. Muell. Cens. 61 (1882) = KIESSOCHYTRON, Spenn. = Kiessera, Tourn. (Arallac.).

CISSUS, Linn. Diss. Dass. 6 (1747); Amoen. Acad. i. 289 (1749) = Vitis, Tourn. (Ampelid.).
acetosa, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 24 = V. acetosa.
acida, Linn. Sp. Pl. ed. II. 170 = V. acida.
acida, Roxb. ex Wight & Arn. Prod. 137 = V. trifolia.
acida, Wall. Cat. n. 6000 B = V. setosa.
acutifolia, Walp. Rep. i. 441 = V. serjaniæfolia.
aculeata, Span. in Linnæa, xv. (1841) 134 = V. aculeata.
acuminata, A. Gray, Bot. U. St. Expl. Exped. i. 378 = V. acuminata.
acuminata, Thw. Enum. Pl. Zeyl. 62 = V. repanda.
acutifolia, Poir. Encyc. Suppl. i. 106 = V. trifolia.
adenantha, Presen. in Mus. Senckenb. ii. (1837) 283 = V. adenantha.
adenocaulis, Stend. ex A. Rich. Tent. Fl. Abyss. i. 111 = V. adenocaulis.
adnata, Roxb. Hort. Beng. 11; Fl. Ind. i. 405 = V. adnata.
adnata, Wight, ex Wall. Cat. sub n. 5988 = V. Heyneana.
ægyrophylla, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 130 = V. ægyrophylla.
alata, Blanco, Fl. Filip. ed. II. 51 = V. geniculata.
alata, Jacq. Select. Am. 28 = V. sulcicaulis ?
albida, Cambess. in St. Hil. Fl. Bras. Mer. i. 844 = V. albida.
altissima, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 74 = V. papuana.
amazonica, Linden, in Belg. Hortic. xv. (1865) 99.—Bras.
amboinensis, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 74 = V. amboinensis.
Ampelopsis, Pers. Syn. i. 142 = V. indivisa.
anamalayana, Wight, in Madr. Journ. Sc. Ser. III. i. (1864) 41 = V. anamalayana.
anemoniifolia, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 80 = V. anemoniifolia.
angulata, Lam. Tabl. Encyc. i. 331 = V. Linnæi.
angustifolia, Roxb. Hort. Beng. 11; Fl. Ind. i. 408 = V. angustifolia.
antarctica, Vent. Choix, 21. t. 21 = V. antarctica.
aquatica, Buch.-Ham. ex Wall. Cat. n. 6000 = V. repanda.
arachnoidea, Hassk. Cat. Hort. Bog. Alt. 166 = V. indica.
arayensis, Miq. Fl. Ind. Bat. Suppl. i. 517 = V. araneosa.
arborea, Blanco, Fl. Filip. ed. II. 51 = Evodia Roxburghiana.
arborea, Desmoul. in Durand, Monog. Vit. 60 = V. arborea.
arborea, Forsk. Fl. Aegypt. Arab. 82 = Salvadora persica.
arborea, Willd. ex Roem. & Schult. Syst. iii. Mant. 248 = Cyblanthus Humboldtii.
argentea, Linden, in Belg. Hortic. xvii. (1867) 108.—Bras.
arguta, Hook. f. in Hook. Niger Fl. 261 = V. arguta.
aristata, Blume, Bijdr. 188 = V. adnata.
articulata, Guill. & Perr. Fl. Seneg. Tent. i. 185, in obs.—Afr. trop.
asimilis, Kurz, ex Laws. in Hook. f. Fl. Brit. Ind. i. 660 = V. lanceolaria.
auriculata, DC. Prod. i. 632 = V. auriculata.
auriculata, Roxb. Hort. Beng. 11; Fl. Ind. i. 411 = V. auriculata.
australasica, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 8 = V. hypoglauca.
Bandiensis, Brouss. ex DC. Prod. i. 629 = V. antarctica.
bifida, Schum. & Thonn. Beskr. Guin. Pl. 80 = V. quadrangularis.
bigemina, Harv. in Harv. & Sond. Fl. Cap. i. 258 = V. gracilis.
bigitanulosa, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 75 = V. pallidora.
bipinnata, Nutt. Gen. Am. i. 144 = V. arborea.
Blumeana, Hassk. in Flora, xxv. (1842) II. Beibl. 39 = V. indica.
Blumeana, Span. in Linnæa, xv. (1841) 134.—Java.
Blumeana, Steud. Nom. ed. II. i. 371 = V. pallida.
bowrtiensis, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 129 = V. striata.
brevipedunculata, Maxim. Prim. Fl. Amur. 68 = V. heterophylla.

CISSUS :—
bryoniifolia, Regel, Fl. Usur. t. 8. f. 3 = Vitis heterophylla.
caesia, Afzel. Rem. Guin. 58 = V. caesia.
canescens, Lam. Tabl. Encyc. i. 331.—Pervv.
cantoniensis, Hook. & Arn. Bot. Beech. Voy. 175 = V. cantoniensis.
capensis, Willd. Sp. Pl. i. 655 = V. capensis.
caprolata, Walp. Rep. i. 439 = V. caprolata.
carnosa, Lam. Encyc. i. 31 = V. trifolia.
caustica, Tussac, Fl. Ant. i. 116. t. 16 = V. trifoliata, *Baker*.
cerifera, Tulasne, & Bhn. in Tijdschr. Nederl. Ind. xxv. (1863) 418 = V. cerifera.
cinerea, Lam. Tabl. Encyc. i. 332 = V. trifolia.
cirrhiflora, Eckl. & Zeyh. Enum. 56 = *punctiflora*.
cirrhosa, Willd. Sp. Pl. i. 667 = V. cirrhosa.
cirrhosa, Willd. ex Roem. & Schult. Syst. iii. Mant. 414 = V. aculeata.
clematifolia, Walp. Rep. i. 439 = V. scuticata.
cochinchinensis, Spreng. Syst. i. 650.—Cochinchin.
compressa, Blume, Bijdr. 183 = V. adnata.
compressicaulis, Kuis & Pav. Fl. Per. i. 64. t. 100.—Peruv.
connivens, Lam. Tabl. Encyc. i. 332.—Madag.
cordata, Roxb. Hort. Beng. 11; Fl. Ind. i. 407 = V. glauca, repens.
cordata, Wall. Cat. n. 6001 = V. adnata.
cordifolia, Linn. Sp. Pl. ed. II. 170.—Am. austr.
coriacea, DC. Prod. i. 633 = V. coriacea.
crenata, Blume, Bijdr. 186 = *Blumeana*.
crenata, Vahl, Symb. Bot. iii. 19 = V. trifolia.
cuneifolia, Eckl. & Zeyh. Enum. 56.—Afr. austr.
Currori, Hook. f. in Hook. Niger Fl. 265 = V. Currori.
cyanocarpa, Miq. Fl. Ind. Bat. i. ii. 608 = V. japonica.
cymosa, Schum. & Thonn. Beskr. Guin. Pl. 82 = V. Thonningii.
cymosa, Steud. Nom. ed. II. i. 372 = V. tenuifolia.
cyphopetala, Fresen. in Mus. Senckenb. ii. (1837) 282 = V. cyphopetala.
Daridi, Miq. Fl. Ind. Bat. Suppl. i. 515.—Sumatra.
Davidiana, Carr. in Rev. Hortic. (1868) 29.—China.
dealbata, Bedd. in Madr. Journ. Sc. Ser. II. xxii. (1861) 74.—Ind. or.
deficiens, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 160 = V. striata.
dendroides, Roem. & Schult. Syst. iii. Mant. 248 = Ardisia acuminata.
denticulata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 591 = V. Afzelii ?
dichotoma, Blume, Bijdr. 186 = V. dichotoma.
dichrostkrix, Miq. Fl. Ind. Bat. Suppl. i. 514.—Sumatra.
Diepenhorstii, Miq. l. c. 515 = V. Diepenhorstii.
digitata, Lam. Tabl. Encyc. i. 332.—Arabia.
dimidiata, Eckl. & Zeyh. Enum. 57.—Afr. austr.
dioica, Roxb. ex DC. Prod. i. 632 = V. lanceolaria.
discolor, Blume, Bijdr. 181 = V. discolor.
distentula, DC. Prod. i. 631 (*= heterophylla*. Link). —Hab. ?
diversifolia, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 314 = V. cantoniensis.
Dregeana, Bernh. in Flora, xxvii. (1844) 397 = V. capensis.
Duarteana, Cambess. in St. Hil. Fl. Bras. Mer. i. 843 = V. Duarteana.
dubia, Steud. Nom. ed. II. i. 379.—China.
edulis, Dals. in Hook. Kew Journ. ix. (1857) 348 = V. quadrangularis.
elegans, Hort. ex C. Koch, Dendrol. i. 555 = V. heterophylla.
elliptica, Cham. & Schlecht. in Linnæa, v. (1830) 221 = V. elliptica.
elongata, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) 110 = V. Miqueliana.
elongata, Roxb. Hort. Beng. 11; Fl. Ind. i. 411 = V. elongata.
emarginata, Sw. in Vet. Akad. Handl. Stockl. (1825) 437 = V. crenata.
Endressii, Hort.—Cf. Gard. Chron. (1876) i. 592.—Costa Rica.
enneaphylla, Vell. Fl. Flum. 40; t. t. 108 = V. enneaphylla.
Epidendron, Eckl. in Mart. Fl. Bras. xiv. II. 217 = prec.

CISSUS:—
arcuata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Vitis arcuata.
erythrodes, Hochst. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 79 = V. erythrodes.
feminea, Roxb. Hort. Beng. 13; Fl. Ind. i. 410 = V. lanceolaria.
ferruginea, E. Mey. Zwei Pfl. Docum. 178 = *Thunbergii*.
ferruginea, Willd. ex Roem. & Schult. Syst. iii. Mant. 348 = Ardisia ferruginea.
Figueiana, Webb, Fragm. Fl. Aethiop. 58 = V. Figueriana.
flexuosa, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 418.—Ins. Lazon.
formina, Steud. Nom. ed. II. i. 372 = V. lanceolaria.
fragariaefolia, Boj. Hort. Maurit. 60.—Zanzibar.
fragilis, E. Mey. Zwei Pfl. Docum. 178, nomen; Harv. & Sond. Fl. Cap. i. 249.—Afr. austr.
frutescens, Blanco, Fl. Filip. ed. i. 70 = Evodia Roxburghiana.
fuliginea, H. B. & K. Nov. Gen. et Sp. v. 294.—Am. austr.
Gardneri, Thw. Enum. Pl. Zeyl. 68 = V. Gardneri.
geniculata, Blume, Bijdr. 184 = V. geniculata.
gigantea, Bedd. in Madr. Journ. Sc. xxii. Ser. II. (1861) 75.—Ind. or.
glaberrima, Steud. Nom. ed. II. i. 372 = V. glaberrima.
glabra, Herb. Madr. ex Wall. Cat. sub n. 6988 = V. Heyneana.
gladra, E. Mey. Zwei Pfl. Docum. 178 = V. orientalis.
glabrata, Blume, Bijdr. 189 = V. serrulata.
glandulosa, J. F. Gmel. Syst. 256.—Arabia.
glandulosa, Fotr. Encyc. Suppl. i. 105 = V. antarctica.
glandulosa, Roxb. Hort. Beng. 11, nomen.—Ind. or.
glauca, Roxb. l. c. 11; Fl. Ind. i. 406 = V. glauca.
glauca, Thw. Enum. Pl. Zeyl. 62 = V. pallida ?
glaucophylla, Hook. f. in Hook. Niger Fl. 363 = V. glaucophylla.
glyptopetala, Thw. Enum. Pl. Zeyl. 62 = V. glyptocarpa.
gongylodes, Burch. ex Baker, in Mart. Fl. Bras. xiv. ii. 209 = V. gongylodes.
gracilis, Guill. & Perr. Fl. Seneg. Tent. i. 184 = V. gracilis, *Baker*.
granulosa, Ruiz & Pav. Fl. Per. i. 62 = V. striata.
haematantha, Miq. in Linnaea, xxvi. (1855) 220.—Guiana.
Hahniana, Ernst, in Journ. Bot. xiv. (1876) 179.—Venezuela.
hastata, Miq. Fl. Ind. Bat. Suppl. i. 517 = V. hastata.
hederacea, Fers. Syn. i. 148 = V. hederacea.
hederacea, Royle, ex Walp. Rep. i. 429 = V. capreolata.
helleborifolia, Zipp. ex Span. in Linnaea, xv. (1841) 184 = V. auriculata.
heptaphylla, Retz. Obs. v. 22 = V. pedata.
heterona, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 591.—Java.
heterophylla, Herb. Madr. ex Wall. Cat. n. 6036 = V. Rheedii.
heterophylla, Link, Enum. Hort. Berol. i. 148 = *diversifolia*, DC.
heterophylla, Poir. Encyc. Suppl. i. 107.—Java.
heterophylla, Walp. Rep. i. 441 = V. Landuk.
heterotoma, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 591.—Java.
Heyneana, Steud. Nom. ed. II. i. 372 = V. Heyneana.
himalayana, Walp. Rep. i. 441 = V. himalayana.
hirsuta, Steud. Nom. ed. I. 190 = V. hederacea.
hirtella, Blume, Bijdr. 185 = V. geniculata.
humulifolia, Regel, Tent. Fl. Ussur. 36 = V. heterophylla.
hydrophora, Gaud. Bot. Voy. Bonite, i. 131.—Hab.?
hypoglauca A. Gray, Bot. U. St. Expl. Exped. i. 373 = V. hypoglauca.
hypoleuca, Harv. in Harv. & Sond. Fl. Cap. i. 252.—Afr. austr.
iboensis, Hook. f. in Hook. Niger Fl. 365 = V. iboensis.
inaequalis, Heyne, ex Wall. Cat. n. 6010 = V. inaequalis.
incarnata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 417 = V. incarnata.
incisa, Dennsod. in Durand, Monog. Vit. 59, ex S. Wats. Bibliog. Ind. N. Am. Bot. 178 = V. incisa.
indica, Kem. ex Steud. Nom. ed. II. i. 371, 372 = V. Linnaei.

CISSUS:—
indica, Rottl. in Ges. Naturf. Freunde, Neue Schr. iv. (1808) 188 = Vitis repanda.
indica, Walp. Rep. i. 441 = V. indica.
indivisa, Desmoul. in Durand, Monog. Vit. 50, ex S. Wats. Bibliog. Ind. N. Am. Bot. 178 = V. indivisa.
intermedia, A. Rich. Ess. Fl. Cub. 311 = V. trifoliata, *Baker*.
involucrata, Spreng. Syst. iv. Cur. Post. 44 = Pterisanthes cissoides.
ipomoeifolia, Webb, Fragm. Fl. Aethiop. 57 = V. ipomoeifolia.
Jacksii, Korth. ex Miq. in Ann. Mus. Bot. Lugd. Bat. i. 81 = V. geniculata.
japonica, Willd. Sp. Pl. i. 689 = V. japonica.
javana, DC. Prod. i. 626.—Java.
juncea, Webb, Fragm. Fl. Aethiop. 57 = V. juncea.
Kleinii, Lawa. in Hook. f. Fl. Brit. Ind. i. 649, in syn. = V. adnata.
lacinata, Linn. Syst. ed. X. 897 = Sicyos laciniata.
laevigata, Blume, Bijdr. 191 = V. auriculata.
Lamarckiana, Roem. & Schult. Syst. iii. Mant. 329 = V. sicyoides.
lanceolaria, Roxb. Hort. Beng. 11; Fl. Ind. i. 412 = V. lanceolaria.
Landuk, Hassk. in Flora, xxv. (1842) ii. Beibl. 89 = V. Landuk.
langsra, Harv. in Harv. & Sond. Fl. Cap. i. 252.—Afr. austr.
latifolia, Descourt. Fl. Med. Ant. v. 29. t. 11 = V. sicyoides.
latifolia, Lam. Encyc. i. 30 = V. glauca.
latifolia, Vahl, Symb. Bot. iii. 18 = V. adnata.
latifolia, Walp. Rep. i. 441 = V. latifolia.
leonensis, Hook. f. in Hook. Niger Fl. 364 = V. leonensis.
leucocarpa, Blume, Bijdr. 189 = V. japonica.
leucostaphyla, Steud. Nom. ed. II. i. 372 = V. lanceolaria.
Lindeni, André, Illustr. Hortic. xvii. (1870) 19. t. 2.—N. Granat.
Linnaeana, Welw. in Journ. Linn. Soc. viii. (1865) 76.—Afr. trop.
lonchiphylla, Thw. Enum. Pl. Zeyl. 62 = V. lonchiphylla.
longipedunculata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 417 = V. longepedunculata.
lucida, Poir. Encyc. Suppl. i. 106 = V. erosa.
macrobotrya, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 369.—Sumatra.
macrophylla, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 306.—Java.
macropus, Welw. in Journ. Linn. Soc. viii. (1865) 77 = V. macropus.
Mappia, Lam. Tabl. Encyc. i. 333 = V. Mappia.
membranacea, Hook. f. in Hook. Niger Fl. 366 = V. gracilis.
mexicana, Moc. & Sesse, ex DC. Prod. i. 631 = V. mexicana.
micrantha, Poir. Encyc. Suppl. i. 105.—Ins. S. Doming.
microantha, Willd. ex Roem. & Schult. Syst. iii. Mant. 348 = *parviflora*.
microcarpa, Vahl, Eclog. Am. i. 16 = V. trifoliata, *Baker*.
microphylla, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 416.—Afr. austr.
mollis, Steud. ex Baker, in Oliver, Fl. Trop. Afr. i. 412 = V. serpens.
macrocarpa, Hook. & Arn. ex Steud. Nom. ed. II. i. 372 = V. lanceolaria.
multibracteata, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 75 = V. pubiflora.
multiflora, Willd. ex Roem. & Schult. Syst. iii. Mant. 349.—Am. austr.
mauritiana, Dals. & Gibs. Bomb. Fl. 40; Thw. Enum. Pl. Zeyl. 62 = V. lanceolaria.
mutabilis, Blume, Bijdr. 190 = V. acutabilis.
napaulensis, DC. Prod. i. 632 = V. capreolata.
natalensis, Bernh. ex Krauss, in Flora, xxvii. (1844) 349 = Rhus natalensis.
nelligherrensis, Wight, Spicil. Neilgh. i. 23. t. 26 = V. himalayana.
nepalensis, Steud. Nom. ed. II. i. 372, vide *napaulensis*.
nicaragirea, Miq. in Linnaea, xxvi. (1853) 221.—Ind. or.
nitida, Vell. Fl. Flum. 39; i. t. 160 = V. sicyoides.

CISSUS:—
nivea, Hochst. ex Schweinf. Beitr. Fl. Aethiop. 32.—Abyssin.
nodosa, Blume, Bijdr. 183 = Vitis nodosa.
obliqua, Ruiz & Pav. Fl. Par. i. 65.—Peruv.
obovata, Vahl, Symb. Bot. iii. 19 = V. trifoliata, *Baker*.
obscura, DC. Prod. i. 629 = V. sicyoides.
obtusata, Benth. Bot. Voy. Sulph. 77.—Panama.
obtusifolia, Lam. Encyc. i. 31 = V. trifolia.
ochracea, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 25 = V. ochracea.
officinalis, Klotzsch, in Bot. Zeit. iv. (1846) 110.—N. Granat.
opaca, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 28 = V. opaca.
orientalis, Lam. Tabl. Encyc. i. 332. t. 84. f. 2 = V. orientalis.
ovata, Lam. l. c. 331 = V. sicyoides.
ovata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106. —Guiana.
oxyodon, Planch. in Catal. Pl. Hort. Donat. 5.—Hab.?
pallida, Salisb. Prod. 66 = V. sicyoides.
pallida, Steud. Nom. ed. II. i. 372 = V. pallida.
palmata, Poir. Encyc. Suppl. i. 107 = V. Commersoni.
paniculata, Heyne, in Wall. Cat. n. 6032 = V. tamnifolia.
papillosa, Blume, Bijdr. 188 = V. pubiflora.
parviflora, Roem. & Schult. Syst. iii. Mant. 346.—Ind. or.
parvifolia, Salisb. Prod. 66 = V. acida.
paucidentata, Ernst, in Journ. Bot. viii. (1870) 374. —Venezuela.
paucidentata, Klotzsch, in Peters, Reise Mossamb. Bot. i. 179 = V. paucidentata.
pauciflora, DC. Prod. i. 630.—Afr. austr.
Pauli-Guilielmi, Schweinf. Rel. Kotsch. 46.—Afr. trop.
paulliniifolia, Vell. Fl. Flum. 40; i. t. 102 = V. ternata ?
pedata, Lam. Encyc. i. 31 = V. pedata.
pedata, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 30 = V. lanceolaria.
pelata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 590.—Java.
pentagona, Roxb. Hort. Beng. 11; Fl. Ind. i. 406 = V. pentagona.
pentandra, Willd. ex Roem. & Schult. Syst. iii. Mant. 348.—Am. austr.
pentaphylla, Buch.-Ham. ex Wall. Cat. sub n. 6016 = V. elongata.
pentaphylla, Herb. Madr. ex Wall. l. c. sub n. 6022 = V. tamnifolia.
pentaphylla, Willd. Sp. Pl. i. 669 = V. pentaphylla.
pergamacea, Blume, Bijdr. 188 = V. pergamacea.
periclata, Hook. f. in Hook. Niger Fl. 362 = V. pallida.
pinnata, [Soland. in] Russell, Aleppo, ed. II. ii. 267. —Cilicia.
poecilophylla, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 79 = V. pinnarpa.
platanifolia, Carr. in Rev. Hortic. (1868) 39.—China.
populnea, Guill. & Perr. Fl. Seneg. Tent. i. 184 = V. pallida.
porphyrophylla, Lindl. in Proc. Hort. Soc. i. (1861) 285 = Piper porphyrophyllum.
productа, Afzel. Rem. Guin. 68 = V. productа.
pruinata, Weinm. in Syll. Ratisb. i. (1824) 224.—Ross.
pruinosa, Desf. Cat. Hort. Par. ed. III. 286.—Hab.?
pubescens, H. B. & K. Nov. Gen. et Sp. v. 296.—Am. austr.
pubescens, Walp. Rep. i. 441 = V. pubescens.
pubiflora, Miq. Fl. Ind. Bat. Suppl. i. 516 = V. pubiflora.
pubiflora, Blume, Bijdr. 185.—Java.
pulcherrima, Vell. Fl. Flum. 40; i. t. 101 = V. pulcherrima.
punctulatae, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = V. sicyoides.
purpurea, Roxb. ex Steud. Nom. ed. II. i. 372 = V. glauca.
purpurascens, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 86 = V. rostrata.
pyrrhadenya, Miq. Fl. Ind. Bat. Suppl. i. 517.—Sumatra.
quadrangularis, Salisb. Prod. 66 = V. quadrangularis.
quadrangularis, Linn. Mant. i. 80 = V. quadrangularis.

CISSUS :—
quadrialata, H. B. & K. Nov. Gen. et Sp. v. 225 = Vitis ecos.
quinata, [Drysad. in] Alt. Hort. Kew. ed. II. i. 260 = V. cirrhosa.
quinquefolia, Desf. Cat. Hort. Par. ed. III. 388 = V. hederacea.
quinquefolia, Seland. in Sims, Bot. Mag. t. 2448 = V. Sinuata.
repanda, Vahl, Symb. Bot. iii. 18 = V. repanda.
repens, Lam. Encyc. i. 31 = V. repens.
repens, Thw. Enum. Pl. Zeyl. 62 = V. repanda.
reticulata, Willd. ex Roem. & Schult. Syst. iii. Mant. 345 = *sanscrena*.
reticulata, Thw. Enum. Pl. Zeyl 62 = V. reticulata.
rhodocarpa, Blume, Bijdr. 185 = V. geniculata.
rhombifolia, Vahl, Eclog. Am. ii. 10 = V. rhombifolia.
rhombicalza, E. Mey. Zwei Pfl. Docum. 173, nomen; Harv. & Sond. Fl. Cap. i. 252.—Afr. austr.
ripuria, Laws. in Hook. f Fl. Brit. Ind. i. 649, in syn. = V. repanda.
Rocheana, Planch. in Comptes-rendus, xxii. (1851) 889.—Afr. trop.
rosea, Walp. Rep. i. 487 = V. repanda.
rotundata, Heyne, ex Wall. Cat. sub n. 6988 = V. Heyneana.
rotundifolia, Blume, Bijdr. 480 = V. pallida.
rotundifolia, Vahl, Symb. Bot. iii. 19.—Arabia.
rubescens, Blanco, Fl. Filip. ed. I. 71 = V. geniculata.
rubricaulis, Carr. in Rev. Hortic. (1868) 89.—China.
rufescens, Guill. & Perr. Fl. Seneg. Tent. i. 133 = V. rufescens.
rugosa, DC. Prod. i. 629.—Ins. Jamaic.
salutaria, H. B. & K. Nov. Gen. et Sp. v. 225 = V. salutaria.
Sandersoni, Harv. in Harv. & Sond. Fl. Cap. i. 253.—Afr. austr.
scariosa, Blume, Bijdr. 191 = V. coriacea.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 114 = V. oxyphylla.
semicordata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 481 = V. himalayana.
semiglabra, Harv. in Harv. & Sond. Fl. Cap. i. 250 = V. semiglabra.
serrata, Eckl. & Zeyh. Enum. 57.—Afr. austr.
serjaniaefolia, Walp. Rep. i. 441 = V. serjaniaefolia.
serpens, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 111 = V. serpens.
serratifolia, Rottl. ex Wight, in Hook. Bot. Misc. ii. (1831) 104 = V. pedata.
serrulata, Roxb. Hort. Beng. 11; Fl. Ind. i. 414 = V. capreolata.
setosa, Roxb. ii. cc. 83; 410 = V. setosa.
sexangularis, Ernst, in Journ. Bot. viii. (1870) 274.—Venezuela.
sicyoides, Klein, ex Stend. Nom. ed. II. i. 372, 373 = V. discolor.
sicyoides, Linn. Syst. ed. X. 897 = V. sicyoides.
simplex, Blanco, Fl. Filip. ed. I. 72 = V. adnata.
Sinuiana, Roem. & Schult. Syst. iii. 346 = V. Sinuana.
smilacina, H. B. & K. Nov. Gen. et Sp. v. 224 = V. sicyoides.
spicifera, Griff. Notul. iv. 698 = V. macrostachya.
spinosa, Cambess. in St. Hil. Fl. Bras. Mer. i. 345 = V. spinosa.
stans, Pers. Syn. i. 149 = V. arborea.
stipulata, Vell. Fl. Flum. 89; i. t. 99 = V. salicifolia?
striata, Ruiz & Pav. Fl. Per. i. 64. t. 100 = V. striata.
stricta, Bert. ex Stend. Nom. ed. II. i. 372, in syn. = V. striata.
subdigitata, Stend. in A. Rich. Tent. Fl. Abyss. i. 110 = V. gracilis.
subcordata, Sedd. in Madr. Journ. Sc. Ser. II. xxii. (1861) 75.—Ind. or.
sylvatica, Cambess. in St. Hil. Fl. Bras. Mer. i. 345 = V. erosa.
tamoides, Cambess. l. c. 343 = V. sicyoides.
tenuicaulis, Hook. f. in Hook. Niger Fl. 266 = V. tamoides.
tenuifolia, Hayne, in Wall. Cat. n. 6062 A = V. tenuifolia.
ternata, J. F. Gmel. Syst. 396.—Arabia.
tetragona, Mavr. in Mavr. & Sond. Fl. Cap. i. 249.—Afr. austr.
tetraptera, Hook. f. in Hook. Niger Fl. 266 = V. quadrangularis.

CISSUS :—
Teysmanni, Miq. Fl. Ind. Bat. Suppl. i. 516.—Sumatra.
Thunbergii, Eckl. & Zeyh. Enum. 56.—Afr. austr.
Thunbergii, Steh. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 195 = Vitis incostans.
thyrsiflora, Blume, Bijdr. 187 = V. thyrsiflora.
tiliacea, H. B. & K. Nov. Gen. et Sp. v. 232 = V. tiliacea.
timoriensis, DC. Prod. i. 630.—Ins. Timor.
tinctoria, Arruda, in Koster, Trav. Bras. 490.—Bras.
tinctoria, Mart. in Spix & Mart. Reise Bras. 368 = V. sicyoides.
tomentosa Lam. Tabl. Encyc. i. 330 = V. capensis.
trianaea, Schum. & Thonn. Beskr. Guin. Pl. 81 = V. quadrangularis.
tricuspidata, Steh. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 196.—Japon.
tricuspis, Burch. ex Baker, in Mart. Fl. Bras. xiv. ii. 309 = V. tricuspis.
tridentata, Eckl. & Zeyh. Enum. 56 = *pauciflora*.
trifoliata, Bech.-Ham. in Wall. Cat. n. 6013 I K = V. angustifolia.
trifoliata, Linn. Sp. Pl. ed. II. 170 = V. trifoliata, *Baker*.
trifoliata, Lour. Fl. Cochinch. 83 = *dubia*.
trigona, Willd. ex Roem. & Schult. Syst. iii. Mant. 348.—Bras.
trilobata, Lam. Encyc. i. 31 = V. Rheedei.
triternata, Miq. Fl. Ind. Bat. Suppl. i. 516.—Sumatra.
truncata, Laws. in Hook. f. Fl. Brit. Ind. i. 660 = V. tenuifolia.
tuberculata, Blume, Bijdr. 189 = V. tuberculata.
tuberculata, Jacq. Hort. Schoenb. i. 14.—Am. austr.
tuberosa, Mog. & Sesse, ex DC. Prod. i. 629 = V. tuberosa.
umbellata, Lour. Fl. Cochinch. 84.—China.
umbrosa, H. B. & K. Nov. Gen. et Sp. v. 223.—Am. austr.
unifoliata, Harv. in Harv. & Sond. Fl. Cap. i. 250.—Afr. austr.
uniflora, Abel. Rem. Guin. 69 = V. uviferi.
vagans, Carr. in Rev. Hortic. (1875) 60 = V. japonica?
Veitchii, Carr. l. c. (1877) 176.—Hab.?
velutina, Linden, ex Hook. Bot. Mag. t. 5207.—Hab.?
venatorum, Descourt. Fl. Méd. Ant. v. 21. t. 309 = V. sicyoides.
verrucosa, Stend. Nom. ed. II. i. 373 = V. tuberculata.
vesicatoria, Blanco, Fl. Filip. ed. II. 50 = V. repens.
villosa, Roxb. Hort. Beng. 11, nomen.—Hab.?
violacea, Meyen, Reise, i. 300.—Chili.
viticifolia, Steh. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 196 = V. serjaniaefolia.
vitiensis, A. Gray, Bot. U. St. Expl. Exped. i. 372 = V. vitiensis.
vitifolia, Boiss. Diagn. Ser. I. vi. 81 = V. persica.
vitifolia, Salisb. Prod. ed. 2 = V. repanda.
vitiginea, Linn. Sp. Pl. 117 = V. repanda.
vitiginea, Roxb. Fl. Ind. i. 406 = V. lanata.
Vogelii, Hook. f. in Hook. Niger Fl. 267 = V. Vogelii.
Wallichiana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 416 = Hedera Helix.
Zollingeri, Turcz. l. c. xxxvi. (1863) i. 591.—Java.

CISTANWCHIA, Hoffmgg. & Link, Fl. Port. i. 318. t. 63 (1806). *OROBANCHACEAE*, Benth. & Hook. f. ii. 983.
Ægyptiaca, Reichb. Consp. 219* (1828).
lutea, *Hoffmgg. & Link, l. c.* 319.—Reg. Mediterr.; Arabia.
lutea, Wight, Illustr. ii. 180. t. 158. b. f. 4 = tubulosa.
tubulosa, *Wight, Ic.* t. 1420 bis.—Ind. bor. occ.

CISTANTHE, Spach, Hist. Vég. Phan. v. 229 (1836) = Calandrinia, Hl. B. & K. (Portulac.).
ancepa, Spach, l. c. 230 = Cal. grandiflora.
discolor, Spach, l. c. 231 = Cal. discolor.
glauca, Spach, ex Heynh. Nom. ii. 146 = Cal. grandiflora.
speciosa, Lilja, ex Heynh. l. c. = Cal. discolor.

CISTELA, Blume, Fl. Jav. praef. p. vi (1828) = seq.

CISTELLA, Blume, Bijdr. 292. t. 55 (1825) = Geodorum, Jack (Orchid.).
cornuta, Blume, l. c. = G. dilatatum.

CISTICAPNOS, Adans. Fam. ii. 431 (1763) = CYSTICAPNOS, Boerh. = Corydalis, Vent. (Papaverac.).

CISTOCARPIUM, Spach, Hist. Vég. Phan. vi. 471 (1838) = Vesicaria, Tourn. (Crucifer.).
utriculatum, Spach, l. c. 472 = V. utricolata.

CISTOCARPUM, Pfeiff. Nom. i. 766 (1874) = seq.

CISTOCARPUS, Kunth, in Mém. Soc. Nat. Hist. Par. iii. (1827) 389 = Balbisia, Cav. (Geraniac.).

CISTOMORPHA, Caley, ex DC. Syst. i. 427, in syn. (1818) = Hibbertia, Andr. (Dilleniac.).
lanceolata, Caley, l. c. = H. saligna.

CISTRUM, Hill, Veg. Syst. iv. 36 (1762) = Centaurea, Linn. (Compos.).
asperum, Hill, l. c. = Cent. aspera.
lenardi, Hill, l. c. = Cent. aspera.
napifolium, Hill, l. c. = Cent. napifolia.

CISTULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

CISTUS, [Tourn.] Linn. Syst. ed. I (1735). *CISTINEAE*, Benth. & Hook. f. i. 118.
HALIMIUM, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 365.
LADANIUM, Spach, l. c. 367 (1836).
LEDONIA, Spach, l. c. 369 (1836).
LIBANOTIS, Rafin. Sylva Tellur. 123 (1836).
RHODOCISTUS, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 367.
STEPHANOCARPUS, Spach, l. c. 366 (1836).
STRABON, Rafin. Sylva Tellur. 122 (1836).
acuminatus, Viv. Ann. Bot. i. ii. (1804) 172 = Helianthemum variabile.
acutifolius, *Sweet, Cist.* t. 75.—Europ.
aegyptiacus, Linn. Sp. Pl. 527 = Helianthemum aegyptiacum.
aegyptius, Pall. ex Bieb. Fl. Taur. Cauc. ii. 8 = Helianthemum salicifolium.
affinis, Bertol. ex Guss. Fl. Sic. Prod. ii. 12 = monspeliensis.
albidus, *Linn. Sp. Pl.* 524.—Reg. Mediterr.
albissimus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 10.—Malaya.
algarvensis, Sims, Bot. Mag. t. 627 = Helianthemum ocymoides.
alpestris, Cranz, Stirp. Austr. ed. I. fasc ii. 78; ed. II. 108 = Helianthemum cinereum.
alterniflolius, Vahl, Symb. Bot. i. 86 = Helianthemum brasiliense.
alyssoides, Lam. Encyc. ii. 20 = Helianthemum alyssoides.
anglicus, Lam. Mant. ii. 245 = Helianthemum cinereum.
angustifolius, Jacq. Hort. Vindob. iii. 29 = Helianthemum variabile.
angustifolius, Lag ex Willk. & Lange, Prod. Fl. Hisp. iii. 784 = Helianthemum leptophyllum.
angustifolius, Salisb. Prod. 368 = Helianthemum umbellatum.
annuus, Lamb. ex Dun. in DC. Prod. i. 272 = Helianthemum villosum.
anthurus, Link, in Schrad. Neues Journ. i. (1805) 112 = Helianthemum umbellatum.
appeninus, Linn. Sp. Pl. 529 = Helianthemum variabile.
arabicus, Linn. Cent. i. 14 = Helianthemum arabicum.
asperifolius, *Pomel, Nouv. Mat. Fl. Atl.* 885.—Afr. bor.
asperifolius, Sweet, Cist. t. 87 = longifolius.
atriplicifolius, Lam. Encyc. ii. 19 = Helianthemum atriplicifolium.
barbatus, Lam. i. c. 24 = Helianthemum vulgare.
Barrelieri, Stend. Nom. ed. I. 199 = Helianthemum glutinosum.
Bourgeanus, *Coss. Notes Crit.* ii. 80.—Hispan.
brasiliensis, Lam. Encyc. ii. 22 = Helianthemum brasiliense.
brevifolius, *Mill. Gard. Dict.* ed. VIII. n. 3.—Hab.?
bupleurifolius, Lam. Encyc. ii. 23 = Helianthemum guttatum.
calycinus, Linn. Mant. ii. 865 = Helianthemum Fumana.
canadensis, Linn. Sp. Pl. 526 = Helianthemum canadense.

CISTUS :—

canariensis, Jacq. Misc. ii. 389 = Helianthemum canariense.

candidissimus, Desf. ex DC. Prod. i. 264 = vaginatus.

canescens, Sweet, Cist. t. 45 = polymorphus.

canus, Gesegl. Bescbr. Russ. Reich. iii. iv. 1047 = Helianthemum oelandicum.

canus, Jacq. Fl. Austr. iii. 43. t. 277 = Helianthemum montanum.

canus, Linn. Sp. Pl. 525 = Helianthemum vineale.

capensis, *Linn. f. c.* ed. II. 736.—*Afr. austr.*

carolinianus, Walt. Fl. Carol. 153 = Helianthemum carolinianum.

carpinifolius, Lam. Encyc. ii. 19 = Helianthemum lasianthum.

ciliatus, Desf. Fl. Atlant. i. 421. t. 109 = Helianthemum virgatum.

ciliatus, Roxburg, ex Dun. in DC. Prod. i. 281 = Helianthemum obtusifolium.

cinereus, Cav. Ic. ii. 38. t. 141 = Helianthemum cinereum.

clathifolius, Steud. Nom. ed. I. 300 = Helianthemum variabile.

Clusii, *Dun. in DC. Prod.* i. 266.—Hispan.; Afr. bor.

collinus, Salisb. Prod. 368 = monspeliensis.

complicatus, Lam. Encyc. ii. 14 = parviflorus.

complicatus, Sprun. ex Nym. Consp. 70 = villosus.

corbariensis, Pourr. in Mém. Acad. Toul. iii. (1788) 312 = salvifolius.

cordifolius, *Mill. Gard. Dict.* ed. VIII. n. 12.—Hab.?

cordifolius, Moench, Meth. 245 = populifolius

coriaceus, Poir. Encyc. Suppl. ii. 278 (= *Hemistemma Aubertii*, *Cremospermeæ*).—Madag.

corsicus, Loisel. Nouv. Not. 34 = villosus.

corymbosus, Poir. Encyc. Suppl. ii. 272 = Helianthemum corymbosum.

crassifolius, Poir. l. c. 277 = Helianthemum cinereum.

creticus, Linn. Sp. Pl. ed. II. 738 = polymorphus.

creticus, Sibth. & Sm. Fl. Græc. Prod. i. 364 = parviflorus.

crispus, *Linn. Sp. Pl.* 524.—Reg. Mediterr.

croceus, Desf. Fl. Atlant. i. 422. t. 110 = Helianthemum glaucum.

Cupanianus, Presl, Fl. Sic. i. 117 = polymorphus.

cymosus, Dun. in DC. Prod. i. 265 = parviflorus.

cyprius, *Lam. Encyc.* ii. 16.—Ins. Cyprus.

denticulatus, *Kit. ex Kanitz, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 529.—Hungar.

denticulatus, Poir. Encyc. Suppl. ii. 278 = Helianthemum salicifolium.

dichotomus, Cav. Ic. iii. 83 = Helianthemum origanifolium.

distachyus, Roth, Catalecta, i. 62 = Helianthemum distachyum.

dubius, *Pourr. in Mém. Acad. Toul.* iii. (1788) 312.—Mont. Pyren.

Dunalianus, Steud. Nom. ed. II. i. 374 = polymorphus.

echioides, Lam. Encyc. i. 21 = Helianthemum variabile.

ellipticus, Desf. Fl. Atlant. i. 418. t. 107 = Helianthemum Lippii.

elongatus, Vahl, Symb. Bot. i. 38 = Helianthemum lasianthum.

ericoides, Cav. Ic. ii. 36. t. 172 = Helianthemum Fumana.

ericoephalus, Viv. Fl. Cors. Sp. Nov. 8 = polymorphus.

fasciculatus, Steud. Nom. ed. I. 300 = Helianthemum fasciculatum.

fastigiatus, Guss. Fl. Sic. Prod. ii. 12 = Clusii.

ferrufinensis, Bertand. in Bull. Soc. Bot. Fr. xxx. (1883) 268.—Afr. bor.

ferrugineus, Lam. Encyc. ii. 25 = Helianthemum umbellatum.

ferrugineus, Lam. l. c. 17 = monspeliensis.

floribundus, *Tausch, in Flora*, xix. (1836) 417.—Gallia.

fontanus, Jacq. Misc. ii. 341 = Helianthemum variabile.

formosissimus, *Schouw. Ic. Pl. Rar.* t. 47.—Lusitan.

formosus, Curt. Bot. Mag. t. 264 = Helianthemum lasianthum.

Fumana, Linn. Sp. Pl. 525 = Helianthemum procumbens.

Fumana, Schouw. ex Ball, in Journ. Linn. Soc. xvi. (1877) 846 = Helianthemum glutinosum.

CISTUS :—

galactites, Schmidel, Ic. Pl. ed. II. 250. t. 85 = Helianthemum vulgare ?

gargasicus, Tenore, Ind. Sem. Hort. Neap. (1829) 15, ex Syll. ed. fol. 74 = villosus.

glaucophyllus, Lam. Fl. Fr. iii. 162 = Helianthemum lævipes.

glaucus, Cav. Ic. iii. 81. t. 261 = Helianthemum glaucum.

glaucus, Desf. Fl. Atlant. i. 418 = Helianthemum crassifolium.

glaucus, *Pourr. in Mém. Acad. Toul.* iii. (1788) 311.—Gallia.

glaucus, Salisb. Prod. 368 = Helianthemum lævipes.

globulariifolius, Lam. Encyc. ii. 23 = Helianthemum Tuberaria.

glomeratus, Lag. Gen. et Sp. Nov. 16 = Helianthemum glomeratum.

glutinosus, Linn. Mant. ii. 246 = Helianthemum glutinosum.

grandiflorus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 713 = ladaniferus.

grandiflorus, Scop. Fl. Carn. ed. II. i. 377 = Helianthemum vulgare.

guttatus, Linn. Sp. Pl. 526 = Helianthemum guttatum.

halimifolius, Linn. l. c. 524 = Helianthemum halimifolium.

helianthemoides, Crantz, Stirp. Austr. ed. I. i. 69; ed. II. 101 = Helianthemum variabile.

helianthemoides, *Desf. Fl. Atlant.* i. 423.—Afr. bor.

Helianthemum, Linn. Sp. Pl. 528 = Helianthemum Fumana.

heterophyllus, Desf. Fl. Atlant. i. 411. t. 104 = polymorphus.

hirsutus, *Lam. Fl. Fr.* iii. 157.—Hispan.; Lusitan.

hirsutus, Lapeyr. Hist. Abr. Pl. Pyr. 303 = Helianthemum polymorphum.

hirtus, Chaub. ex Nym. Consp. 74 = Helianthemum glaucum.

hirtus, Gilib. Fl. Lituan. v. 225 = Helianthemum vulgare.

hirtus, Linn. Sp. Pl. 526 = Helianthemum hirtum.

hirtus, Sibth. & Sm. Fl. Græc. vi. t. t. 501 = Helianthemum obtusifolium.

hirtus, Thlb. ex Dun. in DC. Prod. i. 268 = Helianthemum asperum.

hispanicus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Hispan.

hispidus, Lam. Encyc. ii. 26 = Helianthemum hirtum.

horridus, Lag. in Varied. Cienc. ii. iv. (1805) 214, sphalm. (= *Cistum horridum*).

humilis, Salisb. Prod. 368 = Helianthemum vulgare.

hybridus, Pourr. in Mém. Acad. Toul. iii. (1788) 312 = salvifolius.

hybridus, *Vahl, Symb. Bot.* i. 37.—Hispan.

imbricatus, Poir. Encyc. Suppl. ii. 276 = Helianthemum ledifolium.

immaculatum, Steud. Nom. ed. I. 300 = Helianthemum guttatum.

incanus, Linn. Sp. Pl. 524 = polymorphus.

incanus, Sibth. & Sm. Fl. Græc. v. 76. t. 494 = parviflorus.

inconspicuus, Poir. Encyc. Suppl. ii. 278 = Helianthemum guttatum.

indicus, Linn. Syst. ed. XII. 367, sphalm. vide *Helianthemum*.

intermedius, Poir. Encyc. Suppl. ii. 276 = Helianthemum intermedium.

insolacratus, Lam. Encyc. ii. 20 = Helianthemum lasianthum.

italicus, Linn. Syst. ed. X. 1078 = Helianthemum montanum.

italicus, Pall. ex Bieb. Fl. Taur. Cauc. ii. 8 = Helianthemum oelandicum.

japonicus, *Fortune, in Journ. Agri.-Hort. Soc. Ind.* ix. (1853) 96.—Japon.

kahiricus, Steud. Nom. ed. I. 300 = Helianthemum kahiricum.

ladaniferus, Curt. Bot. Mag. t. 113 = cyprius.

ladaniferus, Gouan, ex Steud. Nom. ed. II. i. 374 = C. Ledon.

ladaniferus, *Linn. Sp. Pl.* 523.—Reg. Mediterr. occ.

ladaniiferus, Stokes, Bot. Mat. Med. iii. 209 = polymorphus.

ladanoides, Hoffingg. Verz. s. = ladaniferus.

laevipes, Linn. Cent. i. 14 = Helianthemum lævipes.

laurifo, Cav. Ic. ii. 35. t. 165. f. 1 = Helianthemum thymifolium.

CISTUS :—

lanceolatus, Vahl, Symb. Bot. ii. 62 = Helianthemum guttatum.

lanuginosus, Viv. Fl. Lib. Spec. 28. t. 14. f. 3 = Helianthemum lanuginosum.

lasianthus, Lam. Encyc. ii. 19 = Helianthemum lasianthum.

latifolius, Sweet, Cist. t. 15 = populifolius.

lazarifolius, *Linn. Sp. Pl.* 523.—Europ. austr. occ.; As. Min.

lavandulæfolius, Lam. Encyc. ii. 25 = Helianthemum lavandulæfolium.

laxus, [Dryand. in] Ait. Hort. Kew. ed. II. iii. 305 = longifolius, *Lam.*

laxus, Brot. ex Nym. Consp. 71 = hirsutus.

ledifolius, Georgi, Beschr. Russ. Reich. Nachtr. 276 = Helianthemum salicifolium.

ledifolius, Linn. Sp. Pl. 527 = Helianthemum ledifolium.

ledonius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = glaucus.

Ledon, Lam. Encyc. ii. 17 = glaucus.

lepidotus, Amo, Fl. Iber. vi. 357 = Helianthemum halimifolium.

Libanotis, Desf. Fl. Atlant. i. 419 = Clusii.

Libanotis, Linn. Syst. ed. X. 1077 = Helianthemum Libanotis.

lilacinus, Hoffmgg. ex Steud. Nom. ed. II. i. 374 = parviflorus.

linearis, Cav. Ic. iii. 8. t. 216 = Helianthemum variabile.

Lippii, Linn. Mant. ii. 245 = Helianthemum Lippii.

longifolius, Hort. ex Steud. Nom. ed. II. i. 374 = grandiflorus.

longifolius, *Lam. Encyc.* ii. 16.—Hispan.; Gallia.

longifolius, *Mill. Gard. Dict.* ed. VIII. n. 18.—Hab.?

lunulatus, All. Auct. Pedem. 80. t. 2. f. 3 = Helianthemum lunulatum.

lusitanicus, Mill. Gard. Dict. ed. VIII. n. 4 = Helianthemum variabile.

lutetus, Gilib. Fl. Lituan. ii. 235 = Helianthemum variabile.

majoranifolius, Gouan, Herbor. 36 = Helianthemum hirtum.

marifolius, Bieb. Fl. Taur. Cauc. ii. 8 = Helianthemum oelandicum.

marifolius, Linn. Sp. Pl. 526 = Helianthemum vineale.

mauritanicus, Thlb. ex Dun. in DC. Prod. i. 276 = Helianthemum glutinosum.

medius, *F. F. Gmel. Syst.* 673.—Italia.

micranthus, Viv. Fl. Lib. Spec. 27 = Helianthemum micranthum.

mollis, Cav. Ic. iii. 82. t. 262. f. 2 = Helianthemum origanifolium.

monspeliensis, *Linn. Sp. Pl.* 524.—Reg. Mediterr. occ.

multiflorus, Amo, Fl. Iber. vi. 359 = Helianthemum halimifolium.

Munbyi, *Pomel, Nouv. Mat. Fl. Atl.* 215.—Afr. bor.

nudabilis, Jacq. Misc. ii. 340 = Helianthemum variabile.

myrtifolius, Lam. Fl. Fr. iii. 161 = Helianthemum oleaster.

nervosus, Lam. l. c. 164 = Helianthemum Tuberaria.

nigricans, Pourr. in Mém. Acad. Toul. iii. (1788) 311 = longifolius.

nilidulus, Lam. Mant. ii. 246 = Helianthemum ledifolium.

nudifolius, Lam. Fl. Fr. iii. 163 = Helianthemum Fumana.

nummularius, Cav. Ic. ii. 34. t. 143 = Helianthemum rubellum.

nummularius, Linn. Sp. Pl. 527 = Helianthemum vulgare.

oblongifolius, Sweet, Cist. t. 67 = longifolius.

obscurus, Steud. Nom. ed. I. 261 = Helianthemum variabile.

obtusifolius, Sweet, Cist. t. 48 = longifolius.

occidentalis, Amo, Fl. Iber. vi. 352 = Helianthemum oleaster.

odoratus, *Chr. Sm., ex Buch, in Berl. Acad. Abh.* 1816-17 (1819) 286.—Ins. Canar.

oxyanthus, Brot.—Cf. Nym. Consp. 74 = Helianthemum cupreum.

oxyanthus, Lam. Encyc. ii. 18 = polymorphus.

CISTUS :—
calandicus, Lam. Encyc. ii. 20 = Helianthemum mon-tanum.
calandicus, Linn. Sp. Pl. 526 = Helianthemum italicum.
clusii, Huet & Hanry, in Bull. Soc. Bot. Fr. vii. (1860) 846 = monspeliensis.
conifolius, Mill. Gard. Dict. ed. VIII. n. 10 = monspeliensis.
corymbosus, Clark, Trav. ii. 462 = Helianthemum oligocephyllum.
crispifolius, Lam. Encyc. ii. 21 = Helianthemum crispifolium.
crotus, Vis. Fl. Ital. Fragm. i. 6 = Helianthemum variabile.
parasiticus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
parvifolius, *Lam. Encyc.* ii. 14.—Reg. Mediterr. or.
Pechii, *Pourr. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* ii. (1874) 90.—Gallia.
pedicularis, *Mœnch. Herbar. Méd. Pr.—Cf. Bull. Soc. Bot. Fr.* ii. (1855) 122.—Gallia.
peloponnesiacus, Poir. Encyc. Suppl. ii. 376 = Helianthemum peloponnesiacum.
pilosus, Gmel. ex Willk. & Lange, Prod. Fl. Hisp. iii. 739 = Helianthemum pulverulentum.
pilosulosides, Lapeyr. Hist. Abr. Pl. Pyr. 301 = Helianthemum montanum.
pilosus, All. Fl. Pedem. ii. 104 = Helianthemum polifolium.
pilosus, Linn. Sp. Pl. 528 = Helianthemum variabile.
*pilulifer*us, Thib. ex Dun. in DC. Prod. i. 282 = Helianthemum variabile.
piperifolius, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
plantagineus, Willd. Sp. Pl. ii. 1197 = Helianthemum guttatum.
plantaginifolius, Sallsb. Prod. 366 = Helianthemum Tuberaria.
platyscpalus, *Sweet, Cist.* t. 47.—Ins. Creta.
polifolius, Linn. Sp. Pl. ed. II. 745 = Helianthemum pulverulentum.
polyfolius, Huds. Fl. Angl. ed. I. 305 = præc.
polyanthos, Desf. Fl. Atlant. i. 490 = Helianthemum polyanthum.
polymorphus, *Willd. Ic. Pl. Hisp.* ii. 19 (sp. aggregat.). —Reg. Mediterr.
populifolius, *Linn. Sp. Pl.* i. 523.—Reg. Mediterr. occ.
porquerollensis, Hanry & Huet, in Bull. Soc. Bot. Fr. vii. (1860) 845 = monspeliensis.
Pouzolzii, Delile, Suppl. Cat. Hort. Monsp. (1839); ex Grm. & Godr. Fl. Fr. i. 169 = albidus.
pulverulentus, Sweet, Cist. t. 38 = hirsutus.
pulverulentus, Pourr. in Mem. Acad. Toul. iii. (1788) 319 = albidus.
punctatus, Willd. Sp. Pl. ii. 1199 = Helianthemum salicifolium.
purpureus, *Lam. Encyc.* ii. 14.—Oriens.
quinquenervis, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
quinquesulnerus, *Vis. in Act. Ist. Ven. Sc.* Ser. III. iv. (1858–59) 185.—Hab. ?
racemosus, Cav. Ic. ii. 31. t. 140 = Helianthemum lavandulæfolium.
racemosus, Lam. Encyc. ii. 25 = Helianthemum virgatum.
racemosus, Linn. Mant. i. 76 = Helianthemum variabile.
ramuliflorus, Poir. Encyc. Suppl. ii. 374 = Helianthemum canadense.
retrofractus, Poir. l. c. 376 = Helianthemum retrofractum.
rorismarinifolius, Pourr. ex Timb. in Bull. Soc. Sc. Phys. & Nat. Toul. ii. (1874) 101 = Clusii.
rosmarinus, All. Fl. Pedem. ii. 105 = Helianthemum vulgare.
rosmarinus, Jacq. Hort. Vindob. iii. 37 = Helianthemum variabile.
rosmarinifolius, All. Fl. Pedem. ii. 105 = Helianthemum rosmarinifolium.
rosmarinifolius, Pourr. in Mém. Acad. Toul. iii. (1788) 313 = Clusii.
rosmarinifolius, Sallsb. Prod. 366 = Helianthemum Libanotis.
rotundifolius, Sweet, Cist. t. 75 = vulgaris.
rufinianus, Viv. Fl. Cors. Sp. Nov. 8 = Helianthemum vulgarissimum.
salicifolius, Cav. Ic. ii. 33. t. 144 = Helianthemum thymifolium.

CISTUS :—
salicifolius, Huds. Fl. Angl. ed. II. 253 = Helianthemum lædifolium.
salicifolius, Linn. Sp. Pl. 527 = Helianthemum salicifolium.
salviæfolius, Boiss. Fl. Orient. iii. 438 = sec.
salvifolius, *Linn. Sp. Pl.* 524.—Reg. Mediterr.
salvifolius, Pall. ex Bieb. Fl. Taur. Cauc. ii. 7 = polymorphus.
sampsucifolius, Cav. Ic. i. 65 = Helianthemum ocymoides.
sampsucifolius, Sims, Bot. Mag. xliii (1816) t. 1808 = Helianthemum variabile.
sanguineus, *Lag. in Varied. Cienc.* ii. IV. (1805) 40.—Austr.
Savi, Bertol. ex Dun. in DC. Prod. i. 275 = Helianthemum arabicum.
scabrosus, [Soland. in] Ait. Hort. Kew. ed. I. ii. 336 = Helianthemum lasianthum.
scorpioides, Desf. ex Steud. Nom. ed. II. i. 375 = Helianthemum glaucum.
sedjera, *Pomel, Nouv. Mat. Fl. Atl.* 216.—Afr. bor.
Segnieri, Cranz, Stirp. Austr. ed. I. fasc. ii. 74, ed. II. 104 = Helianthemum cinereum.
Segnieri, Pourr. ex Dun. in DC. Prod. i. 277 = Helianthemum italicum.
sericea, *Munby, Fl. Algér.* 53.—Algeria.
sericeus, *Vahl, Symb. Bot.* i. 57.—Hispan.
serpillifolius, Balb. ex Dun. in DC. Prod. i. 280 = Helianthemum variabile.
serpyllifolius, Linn. Sp. Pl. 527 = Helianthemum vulgare.
serratifolius, Lamb. ex Dun. in DC. Prod. i. 271 = Helianthemum guttatum.
serratus, Cav. Ic. ii. 97. t. 175 = Helianthemum guttatum.
sessiliflorus, Desf. Fl. Atlant. i. 417 = Helianthemum Lippii.
Sideritis, *Presl, Fl. Sic.* i. 116.—Sicilia.
Skanbergi, *Lojac. in Natural. Sicil.* iv. (1885) 95.—Sicilia.
splendens, Lam. Fl. Fr. iii. 156 = Helianthemum variabile.
squamatus, Linn. Sp. Pl. 1196 = Helianthemum squamatum.
stellulatus, Steud. Nom. ed. I. 201 = Helianthemum stellulatum.
stenophyllus, Link, Enum. Hort. Berol. ii. 74 = Clusii, cyprius.
stipulatus, Forsk. Fl. Aegypt. Arab. 100 = Helianthemum kahiricum.
stoechadifolius, Brot. Fl. Lusit. ii. 270 = Helianthemum glaucum.
stoechadifolius, Hort. ex Dun. in DC. Prod. i. 280 = Helianthemum leptophyllum.
strictus, Cav. Ic. iii. 23. t. 263. f. 2 = Helianthemum virgatum.
strigosus, Fisch. ex Steud. Nom. ed. II. i. 375 = Helianthemum cinereum.
sulphureus, Steud. l. c. ed. I. 201 = Helianthemum variabile.
surrejanus, Linn. Sp. Pl. 527 = Helianthemum Chamæcistus.
symphytifolius, Lam. Encyc. ii. 15 = vaginatus.
syriacus, Jacq. Ic. Pl. Rar. i. 10. t. 96 = Helianthemum lavandulæfolium.
syrticus, Viv. Fl. Lib. Spec. 27 = Helianthemum syrticum.
tauricus, *Y. & C. Prael, Delic. Prag.* 24.—Tauria.
telephiifolius, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
tereifolius, Poir. Encyc. Suppl. ii 379 = Helianthemum hirtum.
tereifolius, Thib. ex Dun. in DC. Prod. i. 281 = Helianthemum variabile.
Thibaudi, Poir. Encyc. Suppl. ii. 377 = Helianthemum lavandulæfolium.
thimifolius, Gouan, ex Vill. Prosp. 49 = Helianthemum thymifolium.
thymifolius, Falk, Beitr. ii. 197 = Helianthemum scoparioides.
thymifolius, Linn. Sp. Pl. 526 = Helianthemum glutinosum.
thymifolius, Sibth. & Sm. Fl. Græc. v. 31. t. 500 = Helianthemum thymifolium.
tomentosus, Lam. Fl. Fr. iii. 156 = polymorphus.
tomentosus, Scop. Ann. II. Hist. Nat. (1769) 58 ; Fl. Cam. ed. II. i. 376 = Helianthemum Chamæcistus.

CISTUS :—
tricuspis, Darwin, Variation Fl. & Anim. Domest. i. 277, ex. typ. = Hibiscus tricuspis.
Tuberaria, Linn. Sp. Pl. 526 = Helianthemum Tuberaria.
umbellatus, Linn. l. c. 525 = Helianthemum umbellatum.
undulatus, Mœnch, Meth. 340 = polymorphus.
vaginatus, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. iii. 504.—Venerifle.
valentinus, Pourr. ex Nym. Consp. 71 = monspeliensis.
varius, Pourr. in Mém. Acad. Toul. iii. (1788) 312 = albidus.
verticillatus, Stend. Nom. ed. II. i. 375 = Helianthemum anthurum.
verticillatus, Brot. Fl. Lusit. ii. 262 = Helianthemum umbellatum.
villosus, *Linn. Gen.* ed. VI. pag ult. (nomen incertum). —Hispan.
vinealis, Willd. Sp. Pl. ii. 1195 = Helianthemum vineale.
violaceus, Cav. Ic. ii. 33. t. 147 = Helianthemum variabile.
virgatus, Desf. Fl. Atlant. i. 422 = Helianthemum virgatum.
virgatus, Thib. ex Dun. in DC. Prod. i. 269 = Helianthemum canadense.
viscosissimus, Tausch, in Flora, xix. (1836) 418.—Hab. ?
viscosus, Stokes, Bot. Mat. Med. iii. 309 = ladaniferus.
vulgaris, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 368 = polymorphus.

CITHARELLA, Norcsha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 = (Quid ?).

CITHAREXYLOM, Bunge, in Cat. Hort. Derp. (1842) 6 ; et in Linnæa, xvii. (1844) 149. *CRUCIFERÆ*, Benth. & Hook. f. i. 67.
canescens, Bunge, in Linnæa, xviii. (1844) 150, in nota = Verbena.
Lehmanni, *Bunge, Cat. Hort. Derp.* (1843) 6 ; *et in Linnæa*, xvii. (1844) 149.—As. med.
vernum, *Bunge, in Linnæa*, xviii. (1844) 150, in nota. —As. med.

CITHAREXYLUM, Mill. Gard. Dict. ed. VI. 175 (1759) ; Linn. Sp. Pl. ed. i. 635 (1753). *VERBENACEÆ*, Benth. & Hook. f. ii. 1149.
POEPPIGIA, Bert. in Bull. Ferussac, xxiii. (Oct. 1830) 109.
RAUWOLFIA, Ruiz & Pav. Fl. Per. ii. 26. t. 152 (1799).
affine, *D. Don, in Edinb. N. Phil. Journ.* (Jan.-March 1831) 238.—Mexic.
affine, Mart. & Gal. in Bull. Acad. Brux. xi. ii. (1844) 328 = reticulatum.
albicaule, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 208.—Cuba.
album, Mill. Gard. Dict. ed. VIII. n. 2 = caudatum.
alpinum, Poepp. ex Schau. in DC. Prod. xi. 545 = Verbena cinerascens.
barbinerve, *Cham. in Linnæa*, vii. (1832) 116.—Bras.
Berteri, *Spreng. Syst.* ii. 768.—Ins. Jamaica.
bracteatum, Hort. ex Walp. Rep. vi. 646 = Baddleia melliodora.
caudatum, *Linn. Sp. Pl.* ed. II. 872.—Ins. Jamaic.
caudatum, Sw. Obs. Bot. 284 = quadrangulare.
Chamissonis, Walp. Rep. iv. 75 = caudatum.
cinereum, *Linn. Sp. Pl.* ed. II. 872.—Am. trop.
cinereum, Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ. (Jan.-March, 1831) 238 = lucidum.
cinereum, Spreng. Syst. ii. 764, partim = myrianthum.
coriaceum, Desf. Cat. Hort. Par. ed. III. 292 = quadrangulare.
cyanocarpum, C. Gay, Fl. Chil. v. 84 = Rhaphithamnus parvifolius.
cyanocarpum, Hook. & Arn. Bot. Beech. Voy. 56. t. 11 = Rhaphithamnus cyanocarpus.
dentatum, *D. Don, in Edinb. N. Phil. Journ.* (Jan.-March, 1831) 237.—Perav.
discolor, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 209.—Cuba.
elegans, Phil. ex Miers, in Trans. Linn. Soc. xxvii. (1870) 98 = Rhaphithamnus longiflorus.
ellipticum, *Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ.* (Jan.-March, 1831) 238.—Mexic.
erectum, Sw. Prod. Veg. Ind. Occ. 91 = caudatum.
flexuosum, *D. Don, in Edinb. N. Phil. Journ.* (Jan.-March 1831) 237.—Peruv.

CITHAREXYLUM:—
fruticosum, Linn. Syst. ed. X. 1115 = cinereum.
ilicifolium, *H. B. & K. Nov. Gen. et Sp.* ii. 256.—Ecuador.
incanum, *Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 286.—Mexic.
laetum, *Horn, in Kjoeb. Vidensk. Meddel.* (1877–78) 104.—Am. austr.
laevigatum, Hostm. ex Griseb. Fl. Brit. W. Ind. 497 = laxiflum.
Lindenii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 208.—Cuba.
longiflorum, *Turcz. l. c.* 207.—Cuba.
lucidum, Cham. & Schlecht. in Linnaea, v. (1830) 97 = caudatum.
lucidum, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 286.—Mexic.
lycioides, *D. Don, l. c.* 287.—Mexic.
macrophyllum, *Poir. Encyc. Suppl.* ii. 367.—Guiana.
melanocarpidium, Sw. Prod. Veg. Ind. Occ. 91 = Petitia domingensis.
mendocinum, *Phil. Sert. Mend. Alt.* 35.—Chili.
Mocinni, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 286.—Mexic.
molle, H. B. & K. Nov. Gen. et Sp. ii. 257 = quitense.
molle, *Yacq. ex Spreng. Syst.* ii. 764.—Am. austr.
molle, Salisb. Prod. 106 = villosum.
myrianthum, *Cham. in Linnaea*, vii. (1832) 117.—Bras.
ovatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 207.—Chili.
paniculatum, *Gaertn. Fruct.* i. 270, t. 56.—Jamaica.
paniculatum, Poir. in Lam. Tabl. Encyc. iii. 95 = Psenna integrifolia.
pentandrum, *Vent. Jard. Cels*, t. 47.—Porto Rico.
perforatum, Forst. f. Prod. 44 = Myoporum laetum.
Poeppigii, *Walp. Rep.* iv. 76.—Bras.
polystachyum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 209.—Cuba.
pulcanthum, *Turcz. l. c.* 207.—Hab. ?
pulverulentum, *Pers. Syn.* ii. 143.—Am. austr.
quadrangulare, Hort. Madr. ex Steud. Nom. ed. II. i. 375 = pulverulentum.
quadrangulare, *Yacq. Enum. Pl. Carib.* 26.—Ind. occ.
quadrangulare, Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ. (Jan.–March, 1831) 288 = Sessei.
quitense, *Spreng. Syst.* ii. 763.—Ecuador.
reticulatum, *H. B. & K. Nov. Gen. et Sp.* ii. 257.—Peruv.
retusum, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 287.—Peruv.
Rugenianum, *Cham. in Linnaea*, vii. (1832) 120.—Bras.
scabrum, *Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 285.—Mexic.
scabrum, Willd. ex Cham. in Linnaea, vii. (1832) 117 = myrianthum.
scurriosum, Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ. (Jan.–March, 1831) 288 = reticulatum.
Sessei, *D. Don, l. c.*—Mexic.
solanaceum, *Cham. in Linnaea*, vii. (1832) 119.—Bras.
spinosum, *H. B. & K. Nov. Gen. et Sp.* ii. 256.—Peruv.
spinosum, *Linn. Sp. Pl.* 625.—Ins. Barbed.
subserratum, *Sw. Prod. Veg. Ind. Occ.* 91.—Ins. S. Doming.
surrectum, *Griseb. Fl. Brit. W. Ind.* 497.—Ind. occ.
tomentosum, *Forst. f. Prod.* 44 = Myoporum tenuifolium.
teres, Jacq. Enum. Pl. Carib. 46 = cinereum.
tomentosum, *H. B. & K. Nov. Gen. et Sp.* ii. 258.—Am. austr.
tomentosum, Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ. (Jan.–March, 1831) 286 = Mocinni.
tomentosum, Poir. Encyc. Suppl. ii. 366 = pantandrum.
tristachyum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 209.—Cuba.
venustum, *Phil. in Bot. Zeit.* xiv. (1856) 646 = Rhaphithamnus longiflorus.
verticillatum, Klotzsch, ex Walp. Rep. iv. 72 = cynocarpum.
villosum, *Yacq. Ic. Pl. Rar.* i. 12, t. 118; *Coll.* i. 72.—Ins. S. Doming.

CITINUS, All. Fl. Pedem. ii. 214 (1785) = Cytinus, Tourn.

CITREUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 141 (1745); Mill. Gard. Dict. ed. VI. (1752) = Citrus, Linn.

CITRIOBATHUS, A. Juss. in Orb. Dict. x. 285 (1849) = seq.

CITRIOBATUS, A. Cunn. in Loud. Hort. Brit. Suppl. i. 585 (1832). *PITTOSPOREAE*, Benth. & Hook. f. i. 122.
Ixosporum, F. Muell. Fragm. ii. 76 (1860).
megacarpus, *F. Muell. Fragm.* ii. 76.—Austral.
multiflorus, *A. Cunn. in Loud. Hort. Brit. Suppl.* i. 585.—Austral.
pauciflorus, *A. Cunn. l. c.*—Austral.

CITRONELLA, D. Don, in Edinb. N. Phil. Journ. [xlii.] (Oct. 1832) 248 = **Villaresia**, Ruiz & Pav. Olacin.
mucronata, D. Don, l. c. = V. mucronata.

CITROSENA, Bosc, ex Steud. Nom. ed. II. i. 375 (1840) = seq.

CITROSMA, Ruiz & Pav. Fl. Per. et Chil. Prod. 134. t. 29 (1794) = **Stiparuna**, Aubl. (Monimiac.).
alternifolia, Spreng. Neue Entdeck. ii. 112 = S. alternifolia.
amazonum, Tul. in Arch. Mus. Par. viii. (1855–56) 329 = S. asperula.
andina, Tul. in Ann. Sc. Nat. Ser. IV. iii. (1855) 36 = S. andina.
apicifera, Tul. l. c. 38 = S. apicifera.
Apiosyce, Mart. ex Tul. l. c. 34 = S. Apiosyce.
aspera, Tul. in Arch. Mus. Par. viii. (1855–56) 324 = S. aspera.
asperula, Tul. in Ann. Sc. Nat. Ser. IV. iii. (1855) 35 = S. asperula.
bifida, Poepp. & Endl. Nov. Gen. et Sp. ii. 48 = S. bifida.
brasiliensis, Spreng. Neue Entdeck. ii. 113 = S. brasiliensis.
buddleiaefolia, Benth. Fl. Hartw. 250 = S. budleiaefolia.
camporum, Tul. in Ann. Sc. Nat. Ser. IV. iii. (1855) 39 = S. camporum.
chiridota, Tul. l. c. 37 = S. chiridota.
cristata, Poepp. & Endl. Nov. Gen. et Sp. ii. 47, t. 164 = S. cristata.
cujabana, Mart. ex Tul. in Ann. Sc. Nat. S.r. IV. iii. (1855) 33 = S. cujabana.
cuspidata, Tul. l. c. 40 = S. cuspidata.
decipiens, Tul. in Arch. Mus. Par viii. (1855–56) 368 = S. decipiens.
dentata, Poepp. ex A. DC. Prod. xvi. ii. 653 = S. mollicoma.
dentata, Ruiz & Pav. Syst. Veg. i. 264 = S. limoniadifolia.
dimidiata, Sello, ex A. DC. Prod. xvi. ii. 656 = S. paniculata.
discolor, Poepp. & Endl. Nov. Gen. et Sp. ii. 47 = S. discolor.
dubia, H. B. & K. Nov. Gen. et Sp. ii. 174 = S. mollis.
echinata, H. B. & K. l. c. 178 = S. echinata.
eriocalyx, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 35 = S. eriocalyx.
erythrocarpa, Mart. ex A. DC. Prod. xvi. ii. 645 = S. estrellensis.
erythracarpa, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 33 = S. erythracarpa.
estrellensis, Tul. l. c. 33 = S. estrellensis.
foliosa, Tul. l. c. 38 = S. foliosa.
gemmiridens, H. B. & K. Nov. Gen. et Sp. ii. 172 = S. glabra.
glabra, Spreng. Syst. ii. 545 = S. glabra.
glabrescens, Presl. Bot. Bemerk. 110 = S. glabrescens.
grandifolia, H.]. & K. Nov. Gen. et Sp. ii. 172 = S. Kunthii.
Gudesiana, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 38 = S. Gondesiana.
guianensis, Tul. in Arch. Mus. Par. viii. (1855–56) 361 = S. guianensis.
Kunthii, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 37 = S. Kunthii.
Lagopus, Tul. l. c. = S. Lagopus.
lanceolata, Tul. l. c. 35 = S. lanceolata.

CITROSMA:—
laurifolia, H. B. & K. Nov. Gen. et Sp. ii. 171 = Stiparuna laurifolia.
lepidota, H. B. & K. l. c. 170 = S. lepidota.
limoniadora, Pav. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 35 = S. limoniadora.
Lindeni, Seem. Journ. Bot. ii. (1864) 343 = S. Lindeni.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 172 = S. macrophylla.
macrophylla, Mart. ex A. DC. Prod. xvi. ii. 655 = S. cristata.
mollicoma, Mart. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 36 = S. mollicoma.
mollis, H. B. & K. Nov. Gen. et Sp. ii. 178 = S. mollis.
muricata, Ruiz & Pav. Syst. Veg. 265 = S. muricata.
Mutisii, H. B. & K. Nov. Gen. et Sp. ii. 170 = S. Mutisii.
myristicoides, Spruce, in Journ. Linn. Soc. v. (1861) 5.—Bras.
neglecta, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 36 = S. neglecta.
oblongifolia, Ruiz & Pav. Syst. Veg. i. 266 = S. laurifolia, oblongifolia.
obovata, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 343 = S. obovata.
oligandra, Tul. in Ann. Sc. Nat. Ser. IV. iii. (1855) 39 = S. obovata.
oligocarpa, Mart. ex A. DC. in DC. Prod. xvi. ii. 654 = S. guianensis.
ovalis, Ruiz & Pav. Syst. Veg. i. 266 = S. ovalis.
paniculata, Spreng. Syst. ii. 545 = S. paniculata.
pauciflora, Beurling, in Vet. Akad. Handl. Stockh. 1854 (1856) 144 = S. pancifiora.
pellita, Tul. in Ann. Sc. Nat. Ser. IV. iii. (1855) 34 = S. pellita.
petiolaris, H. B. & K. Nov. Gen. et Sp. ii. 171 = S. petiolaris.
pleuria, Tul. in Ann. Sc. Nat. Ser. IV. iii. (1855) 33 = S. cujabana.
Poeppigii, Tul. l. c. 39 = S. Poeppigii.
polyantha, Tul. l. c. 31 = S. polyantha.
pyrifera, Ruiz & Pav. Syst. Veg. i. 264 = S. macrophylla.
pyricarpa, Willd. ex Tul. in Arch. Mus. Par. viii. (1855–56) 358 = S. mollis.
radiata, Poepp. & Endl. Nov. Gen. et Sp. ii. 48 = S. radiata.
reginae, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 39 = S. reginae.
riparia, Tul. l. c. 36 = S. riparia.
ruficeps, Tul. l. c. 34 = S. ruficeps.
Schottiana, Spreng. Syst. iv. Cur. Post. 407 = Mollinedia lanceolata.
Selloi, Spreng. Syst. ii. 545 = Mollinedia cinerea.
sessilifiora, H. B. & K. Nov. Gen. et Sp. ii. 174 = S. sessilifiora.
sinuosolens, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 36 = S. sinuosolens.
subinodora, Ruiz & Pav. Syst. Veg. i. 265 = S. subinodora.
Thea, Seem. Journ. Bot. ii. (1864) 343 = S. Thea.
theophora, Poepp. & Endl. Nov. Gen. et Sp. ii. 48 = S. theophora.
tomentosa, Ruiz & Pav. Syst. Veg. i. 266 = S. aspera, tomentosa.
tomentosa, Willd. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 36 = S. neglecta.
trifora, Spreng. Syst. ii. 544 = Mollinedia triflora.
umbellata, Spreng. l. c. 545 = Mollinedia umbellata.

CITRULLUS, Forsk. Fl. Aegypt. Arab. 167 (1775); Neck. Elem. i. 340 (1790). *CUCURBITACEAE*, Benth. & Hook. f. i. 826.
amarus, Schrad. in Eckl. & Zeyh. Enum. 279 = vulgaris.
aqueous, Schur, Enum. Pl. Transs. 221 = vulgaris.
Battichi, *Forsk. Fl. Aegypt. Arab.* 167.—Arabia.
caffer, Schrad. Ind. Sem. Hort. Gotting. (1834); ex Eckl. & Zeyh. Enum. 279 = vulgaris.
caffrorum, Schrad. in Linnaea, xii. (1838) 413 = vulgaris.
chodospermus, Fal. & Dun. in Bull. Soc. Agr. Hérault (1856), 264 = vulgaris.
Colocynthis, *Schrad. in Linnaea*, xii. (1838) 414.—Afr. subt.
edulis, Spach, Hist. Vég. Phan. vi. 214 = vulgaris.
fistulosus, Stocks, in Hook. Kew Journ. iii. (1851) 74, t. 5 = vulgaris.
Naudinianus, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 340.—Afr. trop.

CITRULLUS:—
Pastece, Sagaret, in Ann. Sc. Nat. Sér. I. viii. (1826) 312 = vulgaris.
Pseudo-Colocynthis, Moench. Meth. Suppl. 50 = C. Colocynthis.
variegatus, Schrad. ex M. Roem. I. c. 51 = Cucurbita, Pepo.
vulgaris, *Schrad. ex Echl. & Zeyh. Enum.* 279.—Afr. trop.

CITRUS, Linn. Syst. ed. I (1735). *RUTACEAE*, Benth. & Hook. f. i. 305.
CITREUM, Toorn. ex Rupp. Fl. jen. ed. Hall. 141 (1745); Mill. Gard. Dict. ed. VI (1752).
CITROPHORUM, Neck. Elem. ii. 401 (1790).
LIMON, Tourn. ex Mill. Gard. Dict. ed. VI (1752).
CHAETHERA, Monbrona. in Mém. Acad. Lyon, x. (1860) 186.
PAPEDA, Hassk. in Flora, xxv. (1842) ii. Beibl. 42.
PONCIRUS, Rafin. Sylva Tellur. 148 (1838).
PSEUDAEGLE, Miq. Ann. Mus. Lugd. Bat. ii. 88 (1865–66).
SARCODACTILIS, Gaertn. f. Fruct. iii. 39. t. 185 (1805).
acida, Pers. Syn. ii. 78 = C. Medica.
albida, *Boj. Hort. Maurit.* 49.—Ins. Maurit.
amara, *Hassk. in Flora*, xxv. (1842) *Beibl.* ii. 43.—Java.
amara, Link, Handb. ii. 346 = C. Aurantium.
angulata, *Willd. Sp. Pl.* iii. 1436.—Ins. Amboin.
articulata, *Willd. ex Spreng. Syst.* iii. 834.—Afr. trop.
Aurantium, *Linn. Sp. Pl.* 782.—As. trop.
aurata, Risso, Hist. Nat. Or. Europ. Mérid. i. 409 = C. Aurantium.
australasica, *F. Muell. Fragm.* i. 26.—Austral.
australis, *Planch. in Hort. Donat.* 18.—Austral.
Bergamia, Risso, Hist. Nat. Or. Europ. Mérid. t. 58–65 = C. Aurantium.
Bergamota, Rafin. Sylva Tellur. 141 = præc.
Bigaradia, Loisel. in Duham. Arb. ed. nov. vii. 99 = C. Aurantium.
bonifolia, Hort. ex Ten. Atti 8ª Reun. Sc. Ital. (1841) 508 = Atalantia buxifolia.
buxifolia, Poir. in Lam. Encyc. iv. 580 = C. Aurantium.
Calot, Lag. Gen. et Sp. Nov. 17 = C. Aurantium.
cedra, Link, Handb. ii. 346 = C. Medica.
cedrata, Rafin. Sylva Tellur. 141 = C. Medica.
chilensis, Molina, Sagg. Chile, 171 = Villaresia mucronata.
Combora, Rafin. Sylva Tellur. 142.—Malaya.
costata, Rafin. l. c. = decumana ?
crassa, *Hassk. Cat. Pl. Hort. Bog. Alt.* 217.—Java.
Daidai, *Bieb. ex Van Houtte, Cat.* n. 196 (1849).—Japon.
Decumana, *Murr. Syst.* ed. XIII. 508.—As. trop.
delicious, Tenore, Sull' Arancio Mandarino, 11 = C. Aurantium.
dulcimedulla, Prits. Ic. Bot. Ind. 375 = C. Aurantium ?
dulcis, Pers. Syn. ii. 74 = C. Aurantium.
echinata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 193 = C. Hystrix.
emarginata, *Dcsf. Cat. Hort. Par.* ed. III. 406.—Hab. ?
florida, Salisb. Prod. 378 = C. Aurantium.
fragrans, Salisb. l. c. = C. Medica.
fusca, Lour. Fl. Cochinch. 467 = C. Aurantium.
fusiformis, Rafin. Sylva Tellur. 142.—Hab. ?
Gongra, Rafin. l. c. = C. Medica ?
grandis, *Hassk. in Flora*, xxv. (1842) *Beibl.* ii. 44.—Java.
granulata, Rafin. Sylva Tellur. 142.—Hab. ?
heterophylla, Rafin. l. c. 142.—Hab. ?
Hillii, *F. Muell.* i. 26.—Austral.
humilis, *Poir. in Lam. Encyc.* iv. 580.—Ind. or.
Hystrix, *DC. Cat. Hort. Monsp.* 97.— Ind. or. ; Malaya.
inermis, Raxb. Hort. Beng. 39 ; Fl. Ind. iii. 393 = japonica.
japonica, *Thunb. Fl. Jap.* 292.—Japon ; China.
javanica, *Blume, Bijdr.* 140.—Java.
Kawna, Rafin. Sylva Tellur. 143.—Hab. ?
lacuca, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. i. 515 = C. Hystrix.
Limo, Mandel. in Hook. Bot. Misc. i. (1830) 306 = C. Medica.
Limonia, Risso, in Ann. Mus. Par. xx. (1813) 296. t. 2. f. 1 = C. Medica.
Limonellus, Buch.-Ham. ex Wall. Cat. n. 6386 = C. Medica.
Limonum, Risso, in Ann. Mus. Par. xx. (1813) 301 = C. Medica.
lumia, Risso, Hist. Nat. Or. Europ. Mérid. i. 414 = C. Medica.
macrocantha, *Hassk. Cat. Pl. Hort. Bog. Alt.* 218.—Java ?
macroptera, *Montrous. in Mém. Acad. Lyon*, x. (1860) 137.—N. Caled.
maderensis, Lour. Fl. Cochinch. 467 = japonica.
Margarita, Lour. l. c. = japonica.
marginata, Stend. Nom. ed. II. i. 376 = japonica.
Medica, *Linn. Sp. Pl.* 782.—As. trop.
microcarpa, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 84 = japonica.
mitis, Blanco, Fl. Filip. ed. I. 610 = C. Medica.
myrtifolia, Rafin. Sylva Tellur. 141.—Hab. ?
nobilis, Lour. Fl. Cochinch. 466 = C. Aurantium.
notissima, Blanco, Fl. Filip. ed. I. 607 = C. Medica.
obovata, Rafin. Sylva Tellur. 142.—Hab. ?
obversa, *Hassk. Cat. Pl. Hort. Bog. Alt.* 218.—Java.
ovata, *Hassk. in Flora*, xxv. (1842) *Beibl.* ii. 43.—Java.
paniculata, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* iv. (1828) 152.—Afr. trop.
Papaya, *Hassk. in Cat. Pl. Hort. Bog. Alt.* 218.
Pepedia, Mig. Fl. Ind. Bat. ii. 580 = C. Hystrix.
papillaris, Blanco, Fl. Filip. ed. I. 610 = C. Aurantium.
Paradisi, Macfad. Fl. Jamaic. i. 181 = C. Aurantium.
Pleroma, *F. Muell. Sel. Pl. Indust. Cult.* 84.—Austral.
pyriformis, *Hassk. in Flora*, xxv. (1842) *Beibl.* ii. 44.—Java.
reticulata, Blanco, Fl. Filip. ed. I. 610 = C. Aurantium.
rotundifolia, Rafin. Sylva Tellur. 141.—Hab. ?
salicifolia, Rafin. l. c.—Hab. ?
scaveola, Griff. Notul. iv. 495 = Paramignya Griffithii.
sinensis, Pers. Syn. ii. 74 = C. Aurantium.
spinosa, S. G. Gmel. Reise Russl. iii. 278. t. 22.—Persia.
spinosissima, G. F. W. Mey. Prim. Fl. Esseq. 247 = C. Medica.
torosa, *Blanco, Fl. Filip.* ed. I. 609.—Ins. Philipp.
trifoliata, Thunb. Fl. Jap. 294 = trifoliata.
trifoliata, Hort. ex M. Roem. Syn. Hesper. 89 = Glycosmis pentaphylla.
trifoliata, *Linn. Sp. Pl.* ed. II. 1101.—Japon.
triptera, Desf. Cat. Hort. Par. ed. III. 406 = trifoliata.
tuberosa, Mill. Gard. Dict. ed. VIII. n. 2 = C. Medica.
Vangasay, *Boj. Hort. Maurit.* 49.—Hab. ?
Volkameriana, *Pasq. Cat. Orto Bot. Nap.* 29. — Hab. ?
vulgaris, Risso, in Ann. Mus. Par. xx. (1813) 190 = C. Aurantium.

CITTA, Lour. Fl. Cochinch. 456 (1790) = **Mucuna**, Adans. (Legumin.).
nigricans, Lour. l. c. = M. nigricans.

CITTARONIUM, Reichb. Nom. 186 (1841) = **Viola**, Tourn.

CITTORHINCHUS, Willd. ex H. B. & K. Nov. Gen. et Sp. vi. 14 (1823) = **Gomphia**, Schreb. (Ochnac.).
javitensis, Willd. l. c. = G. aquatica.

CLADANTHUS, Cass. in Bull. Soc. Philom. (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 491.
arabicus, *Cass. l. c.; et in Dict. Sc. Nat.* ix. 342. t. 37.—Arab. (Quid?).
canescens, Sweet, Hort. Brit. ed. III. 886.—Ins. Canar.
Gaubini, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 89 = Mecomischus Cealbi.
hallmifolius, Coss. & Dur. l. c. = Mecomischus Cealbi.
pedunculatus, Benth. & Hook. f. Gen. ii. 422, err. typ. = sec.
pedunculatus, Dur. in Bull. Soc. Bot. Fr. iv. (1857) 14 = Anthemis pedunculata.
proliferus, *DC. Prod.* vi. 15.—Afr. bor.

CLADERIA, Rafin. Sylva Tellur. 12 (1838) = **Malta**, Linn. ?
parviflora, Rafin. l. c. = Hab. ?

CLADIUM, P. Br. Hist. Jamaic. 114 (1756). *CYPERACEAE*, Benth. & Hook. f. iii. 1068.

CLADIUM:—
AGVILLA, Phil. in Anal. Univ. Chil. (1865) i. 643.
BAUMEA, Gaudich. in Freyc. Voy. Bot. 416. t. 29 (1826).
CHAPELLIERA, Nees, in Linnæa, ix. (1834) 298
MACHAERINA, Vahl, Enum. ii. 228 (1806).
SCHOENOPSIS, Beauv. in Lestib. Essai Fam. Cyp. 241 (1819).
TEROSERA, Stend. Syn. Pl. Cyp. 164 (1855).
TRAIL, Beauv. in Lestib. Essai Fam. Cyp. 32 (1819).
TRACHYRHYNCHIUM, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 118.
TRACHYRHYNGIUM, Nees, ex Kunth, Enum. Pl. ii. 214 (1837).
VINCENTIA, Gaudich. in Freyc. Voy. Bot. 417 (1826).
acutum, Poir. in Dict. Sc. Nat. ix. 344 = schoenoides.
ancepe, Hook. f. Handb. New Zeal. Fl. 305, in obs. = trifidiflorum.
ancistrophyllum, F. Muell. ex Benth. Fl. Austral. vii. 415 = Gahnia ancistrophylla.
aristatum, *F. Br. Prod.* 227.—Austral.
asperum, F. Muell. Fragm. ix. 12 = Gahnia aspera.
athrophyllum, *F. Muell. l. c.* 49.—Austral.
baldense, Nees, in Linnæa, ix. (1834) 301 = C. germanicum.
chinense, Nees, l. c.; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 116 = germanicum.
complanatum, *Bergg. in Minnesbr. Fisiog. Sallsk. Lund*, (1877) n. 8. p. 23.—N. Zel.
decompositum, R. Br. Prod. 227 = Gahnia decomposita.
deustum, R. Br. l. c. = Gahnia deusta.
dubium, Spreng. Syst. iv. Cur. Post. 21 = glomeratum
effusum, Torr. in Ann. Lyc. N. York, iii. (1836) 274 = germanicum.
elynanthoides, F. Muell. Fragm. ix. 51.—Austral.
ensifolium, *Benth. & Hook. f. Gen.* iii. 1066.—Sess.
ensigerum, *Hance, in Journ. Bot.* xxiii. (1885) 60.—China.
ficticum, *Hemsl. Bot. Challenger, Voy.* iii. 59, in obs. t. 60. f. 5–7.—Am. austr.
filifolium, *Benth. & Hook. f. Gen.* iii. 1066.—Cuba
Filum, R. Br. Prod. 227, partim = Gahnia trifida.
floribundum, Presl, in Okeu, Isis, xxi. (1828) 270 = germanicum.
gahnoides, Colenso, in Trans. N. Z. Inst. xvi. (1884) 340 = Sinclairi.
germanicum, *Sond. ex Fl. Germ.* i. 75. t. 5. f. 7.—Reg. temp. et subtrop.
giganteum, Willd. in Willk. & Lange, Prod. Fl. Hisp. ii. 136 = germanicum.
glomeratum, *R. Br. Prod.* 227.—As. or. ; Australas.
Gunnii, *Hook. f. Fl. Tasm.* ii. 95. t. 148 B.—Austral. ; N. Zeland.
Hattoni, *T. Kirk. in Trans. N. Z. Inst.* ix. (1877) 551. —N. Zeland.
insulare, *Benth. Fl. Austral.* vii. 403.—Austral.
iridifolium, *Baker, Fl. Maurit.* 434.—Ins. Mascar. ; Bras.
jamaicense, Crantz, Inst. i. 362 = germanicum.
japonicum, Steud. Syn. Pl. Cyp. 152 = germanicum.
junceum, *R. Br. Prod.* 227.—Austral.
lanigerum, R. Br. l. c. = Gahnia lanigera.
latissimum, *F. Muell. Fragm.* ix. 15.—Austral.
lanciforme, Hook. f. Fl. Tasm. ii. 95. t. 148 A = Gunnii.
laxum, *Benth. Fl. Austral.* vii. 405.—Austral.
leptostachyum, [Nees, in] Meyen, Reise, ii. 119 ; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 115 = germanicum.
mariscoides, Torr. in Ann. Lyc. N. York, iii. (1836) 272 = triglomeratum.
mariscoides, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 209.—Ins. Philipp.
Mariscus, R. Br. Prod. 226 = germanicum.
medium, R. Br. l. c. 227 = Gahnia lanigera.
melanocarpum, F. Muell. Fragm. ix. 12 = Gahnia melanocarpa.
Mellei, *Baker, in Journ. Linn. Soc.* xxi. (1885) 451. —Madag.
microstachyum, F. Muell. Census, 129 = Gahnia microstachya.
nodosum, Boeck. in Linnæa, xxxviii. (1874) 226 = Gunnii.

CLADIUM :—
occidentale, Schrad. Fl. Germ. i. 76 = germanicum.
palustre, Poir. in Dict. Sc. Nat. ix. 344 = germanicum.
pantopodum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 457.—Madag.
pauciflorum, R. Br. Prod. 237 = Tricostularia pauciflora.
polyphyllum, F. Muell. Census, 129 = Gahnia polyphylla.
Preissii, F. Muell. ex Benth. Fl. Austral. vii. 405 = latisamum.
psittacorum, F. Muell. Fragm. ix. 13 = Gahnia psittacorum.
quadrangulare, Nees, in Linnaea, ix. (1834) 301 = Lamposoma Gaudichaudii.
Radula, R. Br. Prod. 237 = Gahnia Radula.
rigidifolium, Presl, in Oken, Isis, xxi. (1828) 270 = teretifolium.
riparium, *Benth. Fl. Australi.* vii. 405.—Austral.
schoenoides, *R. Br. Prod.* 237.—Austral.
scirpoideum, *Benth. & Hook. f. ex Hemsl. Bot. Challenger Voy.* iii. 59. t. 60, f. 1–4.—Ins. Juan Fernand.
sclerioides, F. Muell. Fragm. ix. 12 = Exocarya sclerioides.
Siebori, F. Muell. l. c. 14 = Gahnia Sieberi.
Sinclairi, *Hook. f. Handb. N. Zeal. Fl.* 304.—N. Zel.
teretifolium, *R. Br. Prod.* 237.—Austral.
tetragonocarpum, F. Muell. Census, 130 = Gahnia tetragonocarpa.
tetraquetrum, *Hook. f. Fl. Tasm.* ii. 95. t. 149.—Tasmania.
trifidum, F. Muell. in Papers Roy. Soc. Tasm. (1878) 117 = Gahnia trifida.
triglochinatum, *Nees, in Linnaea*, ix. (1834) 301.—Am. bor.
undulatum, *Thw. Enum. Pl. Zeyl.* 353.—Malaya; Austral.
vaginale, *Benth. Fl. Austral.* vii. 406.—Austral.
xanthocarpum, F. Muell. Fragm. ix. 13 = Gahnia xanthocarpa.
xiphalepis, Baker, Fl. Maurit. 434 = Schoenus elongatus.

CLADOBIUM, Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = Scaphyglottis, Pospp. (Orchid.).
violaceum, Lindl. l. c. = S. violacea.

CLADOCALON, Gardn. in Hook. Ic. Pl. t. 528 (1848) = Paepalanthus, Mart. (Eriocaulacee).
brasiliense, Gardn. l. c. = P. Gardnerianus.

CLADOCHAETA, DC. Prod. vi. 245 (1837) = Helichrysum, Vaill. (Compos.).
candidissima, DC. l. c. = H. candidissimum.

CLADODA et CLADODEA, Poir. in Dict. Sc. Nat. ix. 345 (1817) = acq.

CLADODES, Lour. Fl. Cochinch. ii. 574 (1790) = Alchornea, Sw. (Euphorbiac.).
ilicifolia, Baill. Adansonia, vi. (1866) 1 = A. Ilicifolia.
rugosa, Lour. Fl. Cochinch. ii. 574 = A. rugosa.
Thouviana, Baill. Adansonia, vi. (1866) 321 = A. Thouviana.

CLADOGYNOS, Zipp. ex Span. in Linnaea, xv. (1841) 349. *EUPHORBIACEAE*, Benth. & Hook. *f.* iii. 323.
orientalis, *Zipp. l. c.*—Malaya.

CLADOLEPIS, Moq. in DC. Prod. xiv. ii. 268, in syn. (1849) = Obionton, Raf. (Chenopod.).

CLADOMISCHUS, Klotzsch, ex A. DC. in DC. Prod. xv. i. 278 (1864) = Begonia, Linn.

CLADOPOGON, Sch. Bip. in Ind. Sem. Hort. Hamb. (1852) 8, ex Lehm. in Ann. Sc. Nat. Sér. III. xix. (1853) 359 = Senecio, Tourn. (Compos.).
aurantiacus, Lehm. ll. cc. 8; 360.—Mexic.?

CLADOSERIS, Spach, Hist. Vég. Phan. x. 35 (1841) = Onoseris, DC. (Compos.).

CLADOSICYOS, Hook. f. in Oliver, Fl. Trop. Afr. ii. 584 (1871) = Cucumeropsis, Naud. (Cucurbitac.).
edulis, Hook. f. l. c. = Cucum. edulis.

CLADOSTACHYS, D. Don, Prod. Fl. Nep. 76 (1825) = Digera, R. Br. (Amarantac.).
alternifolia, *Sweet*, Hort. Brit. ed. I. 326 = Digera arvensis.
arborescens, D. Don, Syn. Pl. i. 861 = Digera arvensis.
frutescens, D. Don, Prod. Fl. Nep. 76 = Deeringia celosioides.
mauricata, Moq. in DC. Prod. xiii. ii. 235 = Digera arvensis.

CLADOSTEMON, Ad. B. & Vathe, in Monatsb. Acad. Berl. (1876) 866. *CAPPARIDEAE*.
paradoxus, *A. Br. & Vathe, l. c.*—Afr. trop.

CLADOSTIGMA, Radlk. in Bremen Abh. viii. (1883) 413. *CONVOLVULACEAE*.
dioicum, *Radlk. l. c.*—Abyssin.

CLADOSTYLES, Humb. & Bonpl. Pl. Aequin. i. 302. t. 57 (1808) = Ipomoea, Linn. (Convolvulac.).
ericoides, Nees, ex Steud. Nom. ed. II. i. 577 = E. glomerata.
glomerulata, H. B. & K. Pl. Aequin. i. 302 = E. paniculata.

CLADOTHAMNUS, Bong. Veg. Sitcha, 37. t. 1, in Mém. Acad. Pétersb. Sér. VI. ii. (1833). *ERICACEAE*, Benth. & Hook. f. ii. 598.
pyroliflorus, *Bong. ll. cc.*—Am. bor. occ.

CLADOTHECA, Steud. Syn. Pl. Cyp. 178 (1855) = Cryptangium, Schrad. (Cyperac.).
vaginata, Steud. l. c. = Crypt. leptocladium.

CLADOTHRIX, Nutt. ex Moq. in DC. Prod. xiii. ii. 359 (1849). *AMARANTACEAE*, Benth. & Hook. f. iii. 67.
lanuginosa, *Nutt. ex Moq. l. c.* 360.—Calif.
oblongifolia, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 376.—Am. bor. occ.

CLADOTRICHIUM, Vog. in Linnaea, xi. (1837) 401 = Cassipinnia, Linn. (Legumin.).
spunctatum, Vog. l. c. 402 = Cass. spunctata.
villosum, Vog. l. c. = Cass. pilosa.
rubicundum, Vog. l. c. 401 = Cass. rubicunda.
stipulare, Vog. l. c. 402 = Cass. stipulata.

CLADRASTIS, Rafin. Neogenyt. 1 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 554.
BURSERIA, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 53 (1867).
MAACLA, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 188, 143.
alnifolia, *Rafin. New Fl. Am.* iii. 83.—Am. bor.
amurensis, *Benth. ex Maxim. in Bull. Acad. Pétersb. xviii.* (1873) 400.—Reg. Amur.
bentuchensis, Rafin. Neogenyt. (1825) 1 = cinctoria?
lutea, Rafin. l. c. = tinctoria.
secundiflora, Rafin. l. c. = Sophora secundiflora.
tinctoria, *Rafin. l. c.*; *New Fl. Am.* iii. 83.—Am. bor.

CLAIRINIA, Abat, ex Benth. & Hook. f. Gen. iii. 76 (1878) = CLARISIA, Abat = Anredera, Juss. (Chenopodiac.).

CLAIRVILLEA, DC. Prod. v. 636 (1836) = Cacosmia, R. Br. & K. (Compos.).
quinquenervia, DC. l. c. = Cacosmia rugosa.

CLAMBUS, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 16. *MENISPERMACEAE*, Benth. & Hook. f. i. 92.
araucanus, *Miers, l. c.* 17.—Mexic.

CLAMYDANTHUS, *Fenzl, in Ann. Sc. Nat. Linn. Lyon, N. S.* xvii. (1869) 147 = CHLAMYDANTHUS, C. A. Mey. = Thymelaea, Linn.

CLANDESTINA, Tourn. ex Adans. Fam. ii. 207 (1763) = Lathraea, Linn. (Orobanch.).
japonica, Miq. in Verzl. en Med. Kon. Akad. Wetensch. II. ii. (1866) 58 = L. Miqueliana.
penduliflora, Lam. Fl. Fr. ii. 329 = L. Clandestina.
rectiflora, Lam. l. c. 329 = L. Squamaria.

CLANDESTINARIA, Spach, Hist. Vég. Phan. vi. 437 (1838) = Nasturtium, R. Br. (Crucif.).
indica, Spach, l. c. 439 = N. indicum.

CLAOTRACHELUS, Zoll. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 566 = Vernonia, Schreb. (Compos.).
rupestris, Zoll. & Mor. l. c.—Java.

CLAOXYLON, A. Juss. Euph. Tent. 48. t. 14 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 309.
ERYTHROCHILUS, Reinw. ex Flora, viii. (1825) 1. 109.
affine, Baill. Adansonia, ii. (1861–62) 237 = insulanum.
affine, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie*, ii. (1846) 582.—Java.
africanum, Muell. Arg. in DC. Prod. xv. ii. 777 = Barteri.
angolense, *Muell. Arg. in Journ. Bot.* ii. (1864) 388. —Afr. occ.
angustifolium, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 165.—Austral.
australe, *Baill. Etud. Gén. Euph.* 493.—Austral.
Barteri, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 21.—Afr. occ.
capense, *Baill. Etud. Gén. Euph.* 492.—Afr. austr.
columnare, *Muell. Arg. in Flora*, xlvii. (1864) 487.—Afr. trop.
corsifolium, Benth. in Hook. Niger Fl. 506 = Mallotus oppositifolius.
coriaceum, Baill. Etud. Gén. Euph. 493 = erythrophyllum.
crassifolium, *Baill. Adansonia*, i. (1860–61) 279.—Afr. trop.
digynum, Wight, Ic. t. 1884 = Macaranga digyna.
echinocarpum, *Muell. Arg. in DC. Prod.* xv. ii. 787. —Ins. Pacif.
erythrophyllum, *Miq. Fl. Ind. Bat.* i. ii. 587.—Java.
fallax, *Muell. Arg. in DC. Prod.* xv. ii. 780.—Ins. Pacif.
glandifolium, *Miq. Fl. Ind. Bat.* i. ii. 587.—Java.
glanduliferum, *Baill. Adansonia*, i. (1860–61) 281.—Ins. Mascar.
graciliflorum, Miq. Fl. Ind. Bat. i. ii. 586 = affine.
grandidentatum, Boiv. ex Baill. in Adansonia, i. (1860–61) 283 = racemiflorum.
grandifolium, Muell. Arg. in Linnaea, xxxiv. (1865–66) 168 = crassifolium.
hexandrum, *Muell. Arg. in Flora*, xlvii. (1864) 488.—Afr. occ.
Hillii, *Benth. Fl. Austral.* vi. 181.—Austral.
hirtellum, *Baill. Adansonia*, i. (1860–61) 284.—Madag.
indicum, *Hassk. Cat. Hort. Bogor. Alt.* 235.—As. trop.
insulanum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 164.—N. Caledon.
Kirkii, *Muell. Arg. in Flora*, xlvii. (1864) 486.—Afr. occ.
leucocarpum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) 11. 244.—Burma.
lineatostyle, *Baill. Adansonia*, i. (1860–61) 281.—Ins. Mascar.
longifolium, Baill. Etud. Gén. Euph. 493, pro parte = oligandrum.
longifolium, *Muell. Arg. in DC. Prod.* xv. ii. 781.—Malaya.
longipetiolatum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) 11. 244.—Burma.
macranthum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 164.—Ins. Madag.
macrophyllum, Boj. Hort. Maurit. 284 = indicum.
Maingayi, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 20.—Afr. trop.
marianum, *Muell. Arg. in DC. Prod.* xv. ii. 782.—Ins. Marian.
medullosum, *Baill. Adansonia*, i. (1860–61) 283.—Madag.
micrandrum, *Muell. Arg. in Flora*, xlvii. (1864) 487.—Afr. trop.

CLAOXYLON :—
Marcanzilis, *Thw. Enum. Pl. Zeyl.* 271.—Ae. trop.; Afr. trop.
Metsérii, *Soulou, ex Boj. Hort. Maurit.* 285.—Ins. Maurit.
minus, Hamk. Pl. Jav. Rar. 251 = indicum.
molle, Miq. Fl. Ind. Bat. i. ii. 386 = indicum.
monoicum, *Baill. Adansonia*, i. (1860–61) 283.—Madag.
mauritanum, Wight, Ic. t. 1886 = Mallotus muricatus.
Nevadianum, *Baill. Adansonia*, i. (1860–61) 280.—Afr. trop.
occidentale, *Muell. Arg. in Flora*, xlvii. (1864) 438.—Afr. trop.
oligandrum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 164.—Afr. trop.
parviflorum, Hook. & Arn. Bot. Beech. Voy. 212 = indicum.
parviflorum, *A. Juss. Tent. Euph.* 114. t. 14. f. 48.—Ins. Mascar.
parviflorum, Scem. in Bonplandia, ix. (1861) 258 = fallax.
pauciflorum, *Muell. Arg. in Journ. Bot.* ii. (1864) 388.—Afr. trop.
pedicellare, *Muell. Arg. in Flora*, xlvii. (1864) 437.—Afr. trop.
racemiflorum, *Baill. Etud. Gén. Euph.* 492.—Ins. Borbon.
reticulatum, *Boj. Hort. Maurit.* 285.—Ins. Maurit.
rivulare, *Muell. Arg. in Flora*, xlvii. (1864) 513.—Afr. trop.
rubescens, *Miq. Fl. Ind. Bat.* i. ii. 387.—Ins. Pacif.
rubriverve, Baill. Etud. Gén. Euph. 493, nomen = rubescens.
sandwicense, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 166.—Ins. Sandvic.
scabrum, Boj. Hort. Maurit. 285, nomen = Macostachys.
spiciflorum, Baill. in Adansonia, i. (1860–61) 285 = indicum.
spiciflorum, A. Juss. Tent. Euph. 43 = Acalypha spiciflora.
stipulosum, Reichb. [f.] ex Zoll. in Verh. Naturk. Ver. Nederl. Indie, i. (1856) 30; et in Linnaea, xxviii. (1856) 938 = Mallotus stipulosus.
taitense, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 166.—Ins. Tahiti.
tenerifolium, *F. Muell. ex Baill. Adansonia*, vi. (1866) 295.—Austral.
trichogyne, *Muell. Arg. in Journ. Bot.* ii. (1864) 384.—Afr. trop.
triste, *Muell. Arg. l. c.*—Afr. trop.
Wallichianum, *Muell. Arg. in DC. Prod.* xv. ii. 781.—Malaya.
Weinwichianum, *Muell. Arg. in Journ. Bot.* ii. (1864) 383.—Afr. trop.

CLAPPERTONIA, Meisn. Gen. Pl. 86 [98] (1837) = **Honckenya**, Willd. (Tiliac.).
ficifolia, Decne. in Deless. Ic. Select. Pl. v. t. 1 = H. ficifolia.

CLAPTIA, A. Gray, in Torr. Bot. Mex. Bound. 98. (1858). *COMPOSITAE*, Benth. & Hook. f. ii. 413.
aurantiaca, Benth. in Hook. Ic. Pl. t. 1104 (= Dysodiae sp.).—Mexic.
suaedaefolia, *A. Gray, in Torr. Bot. Mex. Bound.* 98.—Mexic.

CLARA, Kunth, in Abh. Akad. Berl. (1848) 44 = **Herrania**, Ruiz & Pav. (Liliac.).
sphinganoides, Kunth, Enum. Pl. v. 296.—Bras.

CLARCKIA, Pursh, Fl. Am. Sept. i. 260 (1814) = **Clarkia**, Pursh (Onagrar.).

CLARIONA, Spreng. Syst. fil. 504 (1826) = seq.

CLARIONEA, Lag. ex DC. in Ann. Mus. Par. xix. (1812) 95 = **Perezia**, Lag. (Compos.).
acanthoides, D. Don, ex Hook. & Arn. in Comp. Bot. Mag. i. (1835) 94 = P. multiflora.
affinis, Phil. in Anal. Univ. Chil. (1873) 479.—Chili.
carthamoides, D. Don, in Phil. Mag. (1832) 393 = P. Hieracifolia.
ciliaris, D. Don, ex Hook. & Arn. in Comp. Bot. Mag. i. (1836) 94 = P. ciliaris.
dicephala, DC. Prod. vii. 93 = P. dicephala.
elegans, Phil. in Linnaea, xxviii. (1856) 717.—Chili.
Feuch, Phil. t. c. 716.—Chili.
glaberrima, Cass. Opusc. ii. 165 = Perezia lactucoides.
humilis, Phil. in Linnaea, xxviii. (1856) 717.—Chili.
laciniata, Phil. in Anal. Univ. Chil. (1873) 479.—Chili.
lactucoides, D. Don, in Trans. Linn. Soc. xvi. (1833) 206 = P. lactucoides.
Lechleri, Sch. Bip. in Flora, xxxviii. (1855) 129.—Peruv.
macrocephala, Sch. Bip. in Lechl. Berb. Am. Austr. 57.—Peruv.
magellanica, DC. in Ann. Mus. Par. xix. (1812) 66. t. 3 = P. magellanica.
multicapitata, Remy, in C. Gay, Fl. Chil. ii. 410 = P. multicapitata.
parvifolia, Phil. in Anal. Univ. Chil. (1873) 480.—Chili.
pedicularifolia, DC. Prod. vii. 61 = P. pedicularidifolia.
pilifera, D. Don, in Phil. Mag. (1832) 388 = P. pilifera.
pinnata, Phil. in Linnaea, xxviii. (1856) 718.—Chili.
pinnatifida, DC. Prod. vii. 62 = P. pinnatifida.
polycephala, Cass. Opusc. ii. 167 = P. multiflora.
pungens, DC. Prod. vii. 62 = P. pungens.
recurvata, D. Don, in Trans. Linn. Soc. xvi. (1830) 206 = P. Doniana.
runcinata, Lag. ex D. Don, l. c. 207 = P. runcinata.
spathulata, Lag. ex D. Don, l. c. 205 = P. viscosa.
variabilis, Phil. in Linnaea, xxxiii. (1864–65) 195.—Chili.
virens, D. Don, in Trans. Linn. Soc. xvi. (1833) 208 = P. Poeppigii.
Volchmanni, Phil. in Anal. Univ. Chil. (1873) 480.—Chili.

CLARIONELLA, DC. ex Steud. Nom. ed. II. i. 377 (1840) = CLARIONEA, Lag. = **Perezia**, Lag. (Compos.).
magellanica, Hombr. & Jacquin. ex Decne. Bot. Voy. Astrol. & Zél. 45 = P. magellanica.

CLARIONEMA, Phil. in Linnaea, xxviii. (1856) 717 = prsec.

CLARIONIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 204 = prsec.

CLARISIA, Abat, in Mem. Soc. Med. Sev. x. (1792) 418.—Cf. Spreng. Gen. 202 (1830) = **Antedere**, Juss. (Chenopod.).
volubilis, Abat, l. c.; ex Moq. in DC. Prod. xiii. ii. 230 = A. spicata.

CLARISIA, Ruiz & Pav. Prod. Fl. Per. & Chil. 158. t. 28 (1794). *URTICACEAE*, Benth. & Hook. f. iii. 377.
SOARESIA, Fr. Allem. in Palestr. Scient. Rio Jan. (1855) 142.
biflora, *Ruiz & Pav. Syst. Veg.* 256.—Peruv.
racemosa, *Ruiz & Pav. l. c.* 255.—Peruv.

CLARKELLIA, Hook. f. Fl. Brit. Ind. iii. 46 (1880). *RUBIACEAE*.
nana, *Hook. f. l. c.*—Reg. Himal.

CLARKIA, Pursh, Fl. Am. Sept. i. t. 11 (1814). *ONAGRARIACEAE*, Benth. & Hook. f. i. 789.
QUARONIA, Presl, Epim. Bot. 219 (1849).
OPISANTHES, Lilja, Fl. Sverig. Suppl. 35 (1840).
PHAROSTOMA, Spach, Hist. Vég. Phan. iv. 392 (1835).
Elegans, *Kellogg, in Proc. Calif. Acad.* vii. (1877) 94.—Calif.
elegans, *Dougl. in Lindl. Bot. Reg.* t. 1575.—Am. bor. occ.
elegans, Poir. in Dict. Sc. Nat. ix. 255 = pulchella.
gracilens, Dougl. ex Sweet, Brit. Flow. Gard. Ser. II. t. 279 = rhomboides.
pulchella, *Pursh, Fl. Am. Sept.* i. t. 11.—Am. bor. occ.
pulcherrima, Hort. ex L'Horticol. Franç. (1861) 7 = pulchella.
rhomboides, *Dougl. in Hook. Fl. Bor. Am.* i. 214.—Am. bor. occ.
unguiculata, Lindl. Bot. Reg. t. 1921 = elegans.
Xantiana, *A. Gray, in Proc. Bost. Soc. Nat. Hist.* vii. (1858–61) 145.—Calif.

CLASTA, Comm. ex Vent. Choix, sub t. 47 (1808) = **Casearia**, Jacq. (Samydac.).
fragilis, Comm. ex Vent. l. c. = Cas. fragilis.

CLASTILIX, Rafin. Sylva Tellur. 99 (1838) = **Miconia**, Ruiz & Pav. (Melast.).
tunicata, Rafin. l. c. = M. mexicana.
tunicata, Rafin. l. c. = M. tunicata.

CLASTOPUS, Bunge, ex Boiss. Fl. Orient. i. 261 (1867) = **Vesicaria**, Tourn. (Crucif.).
purpureus, Bunge, l. c.—Persia.
vesitus, Boiss. l. c.—Persia.

CLATHROSPERMUM, Planch. ex Benth. & Hook. f. Gen. i. 29 (1862). *ANONACEAE*, Benth. & Hook. f. i. c.
biovulatum, *S. Moore, in Journ. Bot.* xv. (1877) 65.—Afr. trop.
Mannii, *Oliver, Fl. Trop. Afr.* i. 26.—Afr. trop.
Vogelii, *Planch. ex Benth. in Trans. Linn. Soc.* xxiii. (1862) 479.—Afr. trop.

CLAUCENA, Barm. f. Fl. Ind. 87 (1768) = **Clausena**, Burm. f. (Rutac.).

CLAUDIA, Opiz, ex Pans. in Denkschr. Rot. Ges. (1808) ii. 177 = **Malica**, Linn. (Gramin.).
ciliata, Opiz, l. c. = M. ciliata.

CLAUSENA, Burm. f. Fl. Ind. 248 (1768). *RUTACEAE*, Benth. & Hook. f. i. 304.
AULACIA, Lour. Fl. Cochinch. 273 (1790).
COOKIA, Sonner. Voy. ii. 130. t. 131 (1782).
FAGARASTRUM, C. Don, Gen. Syn. i. 87 (1832).
GALLESIOA, M. Roem. Syn. Hesper. 45 (1846).
GLAUCENA, Vitm. Summa, ii. 441 (1789).
MYARIS, Presl, Bot. Bemerk. 40 (1844).
QUINARIA, Lour. Fl. Cochinch. 272 (1790).
PIPTOSTYLIS, Dalz. in Hook. Kew Journ. iii. (1851) 33. t. 2.
POLYCYEMA, Volgt, Hort. Suburb. Calc. 141 (1845).
anisata, *Hook. f. in Hook. Niger Fl.* 256.—Afr. trop.
brevistyla, *Oliver, in Journ. Linn. Soc. v. Suppl.* ii. (1861) 31.—Austral.
chrysogyne, *Miq. Fl. Ind. Bat. Suppl.* i. 502.—Malaya.
corymbifolia, *Linden, Cat.* n. 98 (1878), 3.—Ins. Loyalt.
crenulata, F. Muell. Phyt. New Hebr. 7 = Murraya crenulata.
dentata, M. Roem. Syn. Hesper. 44 = Willdenowii.
excavata, *Burm. f. Fl. Ind.* 89. t. 29.—Ind. or.; Malaya.
heptaphylla, *Wight & Arn. Prod.* 95, in nota.—Ind. or.; Burma.
inequalis, *Benth. in Hook. Niger Fl.* 257.—Afr. trop.
indica, *Oliver, in Journ. Linn. Soc. v. Suppl.* ii. (1861) 36.—Ind. or.
javanensis, Rœusch. ex DC. Prod. i. 538 = excavata.
javanica, *M. Roem. Syn. Hesper.* 45.—Java.
javensis, J. F. Gmel. Syst. 610 = excavata?
macrophylla, *Hook. f. Fl. Brit. Ind.* i. 504.—Burma.
nana, Wight & Arn. Prod. 96 = Willdenowii.
pentaphylla, *DC. Prod.* i. 538.—Reg. Himal.
pubescens, Wight & Arn. Prod. 96 = Willdenowii.
punctata, Wight & Arn. l. c. 95 = excavata.
simplicifolia, Dalz. in Hook. Kew Journ. iii. (1851) 180 = Acronychia lanrifolia.
suffruticosa, *Wight & Arn. Prod.* 96, in nota.—Reg. Himal.
sumatrana, Wight & Arn. l. c. 95 = excavata.
timorensis, *M. Roem. Syn. Hesper.* 44.—Ins. Timor.
Wallichii, *Oliver, in Journ. Linn. Soc. v. Suppl.* ii. (1861) 35.—Burma.
Wampi, *Oliver, l. c.* 34.—China?
Wightii, M. Roem. Syn. Hesper. 44 = Willdenowii?
Willdenowii, *Wight & Arn. Prod.* 96.—Ind. or.; Malaya.

CLAUSIA, Trotsky, Ind. Sem. Hort. Casan. (1899) = **Hesperis**, Linn. (Crucif.).
aprica, Trotsky, l. c. = H. aprica.

CLAUSONIA, Pomel, Mat. Fl. Atl. i (1860) = **Asphodelus**, Tourn. (Liliac.).
acaulis, Pomel, l. c. = A. acaulis.

CLAVARIA, Steud. Nom. ed. I. 204 (1821), sphalm. = **Calvaria**, Gaertn. (Gen. dub.).

CLAVENA, DC. Prod. vi. 638 (1837) = Carduus, Tourn. (Compos.).
leucophaula, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276 = Card. baccouphalus.
canariensis, DC. Prod. vi. 638 = Card. clavulatus.
squarrosa, DC. l. c. = Card. squarrosus.

CLAVENNA, Neck. Elem. ii. 145 (1790) = Ammannia, Houst. † (Lythrar.).

CLAVIGERA, DC. Prod. v. 127 (1836) = Brickellia, Ell. (Compos.).
brachyphylla, A. Gray, Pl. Fendl. 68 = B. brachyphylla.
corymbosa, DC. Prod. v. 128 = B. corymbosa.
dentata, DC. l. c. = B. Riddellii.
pinifolia, Carda. in Hook. Lond. Journ. Bot. v. (1846) 461 = B. pinifolia.
Riddellii, Torr. & Gray, Fl. N. Am. ii. 77 = B. Riddellii.
scabra, Benth. Pl. Hartw. 19 = B. corymbosa.
scoparia, DC. Prod. v. 128 = B. scoparia.
spinulosa, A. Gray, Pl. Fendl. 68 = B. spinulosa.

CLAVIJA, Ruiz & Pav. Prod. 142. t. 20 (1794).
MYRSINEAE, Benth. & Hook. f. ii. 649.
Horta, Vell. Fl. Flum. 48 (1825); i. t. 194.
Theophrasta, Linn. Gen. ed. 1. 45 (1737).
Zacintha, Vell. Fl. Flum. 276 (1825); viii. t. 9.
biborrana, *Orrst. in Kjoeb. Vidensk. Meddel.* (1861) 117.—Costa Rica.
calonoura, *Mart. ex Miq. in Mart. Fl. Bras.* x. 277.—Bras.
clavata, *Decne. in Ann. Sc. Nat.* Sér. VI. iii. (1876) 144.—Hab.!
grandis, *Decne. l. c.*—Hab.!
fulgens, *Hook. f. Bot. Mag.* t. 5696.—Peruv.
Hookeri, *DC. Prod.* viii. 148.—Peruv.
integrifolia, *Mart. ex Miq. in Mart. Fl. Bras.* x. 277.—Bras.
lancifolia, *Desf. in Nouv. Ann. Mus. Par.* i. (1832) 402. t. 14.—Guiana.
latifolia, *C. Koch, in C. Koch & Fint. Wochenschr.* (1859) 108.—Ind. occ.
longifolia, *Ruiz & Pav. Syst. Veg.* i. 284.—Peruv.
macrocarpa, *Ruiz & Pav. l. c.*—Am. austr.
macrophylla, *Miq. in Mart. Fl. Bras.* x. 275.—Bras.
ornata, *D. Don, in Edinb. Phil. Journ.* (Jan. 1831) 286.—Am. austr.
pendula, *Ruiz & Pav. Syst. Veg.* i. 285.—Am. austr.
Riedeliana, *Regel, Gartenfl.* (1859) 245.—Bras.
Rodolphiana, *Linden & André, Illustr. Hortic.* xxi. (1874) t. 188.—N. Granat.
sparsiflora, Miq. in Mart. Fl. Bras. x. 276 = Leonia glycycarpa.
spathicalata, Hook. Ic. Pl. t. 140 = Hookeri.
spathulata, *Ruiz & Pav. Syst. Veg.* i. 285.—Am. austr.
speciosa, *C. Koch & Linden, Ind. Sem. Hort. Berol.* (1867) *App.* i. 8.—Am. trop.
umbrosa, *Regel, Ind. Sem. Hort. Petrop.* (1868) 86.—Bras.
undulata, *D. Don, in Edinb. Phil. Journ.* (Jan. 1831) 284.—Am. austr.

CLAVIMYRTUS, Blume, Mus. Bot. Lugd. Bat. i. 118. t. 49 (1849) = Eugenia, Linn. (Myrtac.).
clavifiora, Blume, l. c. 114, in obs. = E. clavifiora.
cylindrica, Blume, l. c. 115 = E. cylindrica.
firma, Blume, l. c. 116.—Java.
glabrata, Blume, l. c. 114.—Java.
latifolia, Blume, l. c. 117.—Borneo.
linnata, Blume, l. c. 116 = E. linnata, *Duthie*.
marginata, Blume, l. c. 115.—Java.
punctifera, Blume, l. c. 114, in obs. = E. lacta.
ramosissima, Blume, l. c. 115 = E. ramosissima.
symphytocarpa, Blume, l. c. 117.—Sumatra.
virens, Blume, l. c. 114.—Java.

CLAVULA, Dum. Fl. Belg. 148 (1827) = Blechcharis, R. Br. & Sabrpus, Linn. (Cyperac.).
acicularis, Dum. l. c. = E. acicularis.
Boscoyon, Dum. l. c. = S. pauciflorus.
caespitosa, Dum. l. c. = S. caespitosus.
comosa, Dum. l. c. = S. nanus?
fluitans, Dum. l. c. = S. fluitans.
multiculmis, Dum. l. c. = E. multiculmis.
ovata, Dum. l. c. = E. ovata.
palustris, Dum. l. c. = E. palustris.
uniglumis, Dum. l. c. = E. palustris.

CLAVULIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 407 = Crotalaria, Dill. (Legumin.).
sericeoides, G. Don, Gen. Syst. ii. 142 = Crot. mucronata.
pedunculatum, Steud. Nom. ed. II. i. 376 = seq.
pedunculosum, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 407 = Crot. labumifolia.

CLAVULUM, G. Don. Gen. Syst. ii. 96, 142 (1832) = prec.

CLAYTONA, Linn. Hort. Ups. 52 (1745) = seq.

CLAYTONIA, Gronov. ex Linn. Gen. ed. I. 339 (1737). *PORTULACEAE*, Benth. & Hook. f. i. 158.
Beija, Steller, ex S. G. Gmel. Fl. Sibir. iv. 88 (1769).
acutifolia, Ledeb. Fl. Alt. i. 353; Ic. t. 373 = arctica.
acutifolia, *Pall. ex Willd. in Roem. & Schult. Syst.* v. 426.—Sibir. or.
cautifiora, Sweet, Hort. Brit. ed. II. 220 = virginica.
alsinoides, Sims, Bot. Mag. t. 1309 = sibirica.
ambigua, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 365.—Am. bor. occ.
aquatica, Nutt. in Torr. & Gray, Fl. N. Am. i. 201 = Chamissonis.
arctica, *Adam, in Mem. Soc. Nat. Mosc.* v. (1817) 94. —As. et Am. arct.
asarifolia, Bong. Veg. Sitcha, 136 = sibirica?
australiasica, *Hook. f. in Hook. Ic. Pl.* t. 298.—Austral.
balcomensis, F. Muell. Census, 27 = Calandrinia balcomensis.
bifida, Willd. ex Roem. & Schult. Syst. v. 434 = sibirica.
brevipedata, F. Muell. Census, 27 = Calandrinia brevipedata.
bulbifera, *A. Gray, in Proc. Am. Acad.* xii. (1877) 54. —Am. bor. occ.
californica, *Hort. Petrop. ex Loud. Hort. Brit. Suppl.* iii. 21.—Calif.
calyptrata, F. Muell. Fragm. iii. 89 = Calandrinia calyptrata.
carolinians, *Michx. Fl. Bor. Am.* i. 160.—Am. bor.
Chamissoi, Ledeb. ex Spreng. Syst. i. 790 = sarmentosa.
Chamissonis, *Eschscholtz, ex Cham. in Linnaea*, vi. (1831) 562.—Am. bor. occ.
compositi, F. Muell. Census, 27 = Calandrinia compositi.
corrigioloides, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 365.—Am. bor. occ.
corrigioloides, F. Muell. Census, 27 = Calandrinia corrigioloides.
cubensis, Bonpl. in Ann. Mus. Par. vii. (1806) 82. t. 6 = perfoliata.
dichotoma, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 202. —Am. bor. occ.
diffusa, *Nutt. l. c.*—Calif.
Eschscholtzii, *Cham. in Linnaea*, vi. (1831) 561.—Am. arct.
exigua, *Torr. & Gray, Fl. N. Am.* i. 200.—Calif.
Alicanilie, Dougl. in Hook. Fl. Bor. Am. i. 294. t. 72 = parvifolia.
flagellaris, Bong. Veg. Sitcha, 136 = Chamissonis.
gracilis, F. Muell. Census, 27 = Calandrinia gracilis.
grandiflora, Sweet, Brit. Flow. Gard. t. 216 = virginica.
granuliflora, F. Muell. Census, 27 = Calandrinia granuliflora.
Gynophiloides, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 33 = perfoliata.
Howardiana, Roem. & Schult. Syst. v. 434 = arctica.
lanceolata, Hook. & Arn. Bot. Beech. Voy. 138, 344 = sarmentosa.
lanceolata, Pursh, Fl. Am. Sept. i. 176. t. 8 = caroliniana.
Lehmanni, F. Muell. Census, 27 = Calandrinia Lehmanni.
linearis, *Dougl. in Hook. Fl. Bor. Am.* i. 294. t. 71.—Am. bor. occ.
linifiora, F. Muell. Census, 27 = Calandrinia linifiora.
media, *Lieb. Hamth.* ii. 44 = virginica.
megarrhiza, *Parry, ex S. Wats. Bibliog. Ind. N. Am. Bot.* i. 118.—Am. bor. occ.
novarosea, Willd. ex Roem. & Schult. Syst. v. 426 (= *Iribachia Bonplandiana*).—N. Granat.
novoalensis, *S. Wats. Bot. Calif.* i. 77.—Calif.

CLAYTONIA—
parviflora, Dougl. ex Hook. Fl. Bor. Am. i. 295. t. 73 = perfoliata.
parvifolia, *Mog. ex DC. Prod.* iii. 361.—Am. bor. occ.
perfoliata, *Denn. Ind. Hort. Cant.* 25, *ex Willd. Sp. Pl.* i. 1186.—Am. bor.
Pickeringi, F. Muell. Census, 27 = Calandrinia Pickeringii.
pleiopetala, F. Muell. l. c. = Calandrinia pleiopetala.
pogonophora, F. Muell. l. c. = Calandrinia pogonophora.
polyandra, F. Muell. l. c. = Calandrinia polyandra.
polypetala, F. Muell. l. c. = Calandrinia polypetala.
Portulacaria, Linn. Mant. ii. 211 = Portulacaria afra.
ptychosperma, F. Muell. Census, 27 = Calandrinia ptychosperma.
pumila, F. Muell. l. c. = Calandrinia pumila.
quadrivalvis, F. Muell. l. c. = Calandrinia quadrivalvis.
sarmentosa, *C. A. Mey. in Nouv. Mém. Soc. Nat. Mosc.* vii. (1839) 137. t. 3.—As. et Am. arct.
sibirica, *Linn. Sp. Pl.* 204.—As. bor.; Am. bor. occ.
sibirica, Pall. ex Stend. Nom. ed. II. i. 376 = arctica.
Simsii, Sweet, Brit. Flow. Gard. sub t. 216 = virginica.
spathulaefolia, Salisb. Parad. Lond. t. 71 = caroliniana.
spathulata, Dougl. in Hook. Fl. Bor. Am. i. 296. t. 74 = perfoliata.
spatulata, Hook. & Arn. Bot. Beech. Voy. 344 = exigua.
spatulata, Eaton, Man. ed. IV. 263 = caroliniana.
spergularina, F. Muell. Census, 27 = Calandrinia spergularina.
stolonifera, C. A. Mey. in Nouv. Mem. Soc. Nat. Mosc. vii. (1829) 139 = Chamissonis.
strophiolata, F. Muell. Fragm. xi. 82 = Calandrinia strophiolata.
tennifolia, Torr. & Gray, Fl. N. Am. i. 201 = exigua.
triphylla, *S. Wats. in Proc. Am. Acad.* x. (1875) 345. —Am. bor. occ.
tuberosa, Mog. & Sesse, ex DC. Prod. iii. 357 = Talinum napiforme.
tuberosa, *Pall. ex Willd. in Roem. & Schult. Syst.* v. 426.—Sibir. or.
umbellata, *S. Wats. in Bot. King, Exp.* 43. t. 6.—Am. bor. occ.
undulatifolia, Ftsch. Hort. Gorenk. ed. II. (1812) 62; et in Roem. & Schult. Syst. v. 434 = sibirica.
uniflora, F. Muell. Census, 27 = Calandrinia uniflora.
Vancouverii, *A. Murr. Oreg. Circ.* 1.—Am. occ.
Verlaine, Fisch. ex G. Don, Gen. Syst. iii. 89 = sarmentosa.
virginica, *Linn. Sp. Pl.* 204.—Am. bor.
volubilis, F. Muell. Census, 27 = Calandrinia volubilis.

CLEACHNE, Roland. ex Steud. Nom. ed. II. i. 378 (1840) = Panpalum, Linn. (Gramin.).

CLEANTHES, Salisb. in Trans. Hort. Soc. i. (1812) 313. *IRIDEAE*, Benth. & Hook. f. iii. 701.
bicolor, *Salisb. l. c.*—Ab. austr.
metelanea, Baker, in Journ. Linn. Soc. xvi. (1877) 112 = bicolor.

CLEANTHES, D. Don, in Trans. Linn. Soc. xvi. (1830) 194 = Trixis, P. Br. (Compos.).
brasilienais, D. Don, l. c. 195 = T. brasilienais.
hieraciodes, D. Don, l. c. 197 = T. ochroleuca.
ochroleuca, Griseb. in Goett. Abh. xxiv. (1879) 217 = T. brasilienais.

CLEGHORNIA, Wight, Ic. t. 1810, 1812 (1850) = Balsamea, DC. (Apocyn.).
acuminata, Wight, l. c. t. 1810 = B. acuminata.
cymosa, Wight, l. c. t. 1811 = B. acuminata.

CLEIDION, Blume, Bijdr. 619 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 290.
Lasiostylis, Presl, Bot. Bemerk. 148 (1844).
Prlostachys, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 98.
Rebia, Comm. Nov. Rdrgs. Bens. Don 61 (1849).
Thylacoma, Bedd. in Madr. Journ. Sc. Ser. II. xiii. (1861) 79.
castaneifolium, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 194.—Penne.
chrysolepidum, *Muell. Arg. l. c.*—N. Calad.

CLEIDION [—]
coriaceum, Baill. Adansonia, ii. (1861–62) 218 = Macaranga coriacea.
gaboniense, *Baill. l. c.* xi. (1874) 120.—Afr. trop.
javanicum, *Blume, Bijdr.* 613.—Ind. or. ; Malaya.
macrophyllum, Baill. Adansonia, ii. (1861–62) 219 = Vitellariâ.
nicaraguense, *Hemsl. Biol. Centr. Am. Bot.* iii. 120.—Nicaragua.
nicobar, *Thw. ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 245.—Ins. Andaman.
populifolium, *Zipp. ex Span. in Linnaea*, xv. (1841) 849, nomen.—Malaya.
spathulatum, *Baill. Adansonia*, ii. (1861–62) 221.—N. Caled.
trioccoum, *Baill. l. c.* iv. (1863–64) 370.—Brasil.
ulmifolium, *Muell. Arg. in Flora*, xlvii. (1864) 481.—Ins. Loo-Choo.
verticillatum, *Baill. Adansonia*, ii. (1861–62) 221.—N. Caled.
Vieillardii, *Baill. l. c.* 220.—N. Caled.

CLEIEMERA; Rafin. Fl. Tellur. iv. 77 (1836) = Ipomœa, Linn. (Convolvulac.).
cuspidata, Rafin. l. c. 78 = Convolvulus Meyeri.
guineensis, Rafin. l. c. = I. capitata.
hederacea, Rafin. l. c. 77 = I. Nil.
hirsuta, Rafin. l. c. 78 = I. hederacea.
indica, Rafin. l. c. (= *Pharbitis violacea*).—As. trop. ?
peruviana, Rafin. l. c. = I. longicupis.

CLEISOCRATERA, Korth. Verh. Nat. Geschied. 256. t. 69 (1843). *RUBIACEÆ*, Benth. & Hook. f. ii. 128.
elegans, Korth. l. c. 257. t. 62.—Borneo.

CLEISOSTOMA, Blume, Bijdr. 362 (1825). *ORCHIDEÆ*, Benth. & Hook. f. iii. 580.
ECHINOGLOSSUM, Reichb. Nom. 54 (1841).
ECHIOGLOSSUM, Blume, Bijdr. 364. t. 98 (1825).
POMATOCALPA, Breda, Orch. Kabl & Hass. (1837).
SYMPTERA, Llanos, Fragm. Fl. Filip. 98 (1851).
acaule, Lindl. Gen. et Sp. Orch. 327 (= Saccolabii sp.). —Zeylan.
amabile, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* v. (1853) 493.—Malaya.
appendiculatum, *Benth. & Hook. f. Gen.* iii. 577.—Ind. or.
Armitii, F. Muell. Fragm. ix. 49 = Sarcochilus Armitii.
Beckleri, *F. Muell. ex Benth. Fl. Austral.* vi. 296.—Austral.
bicolor, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 99. —Manilla.
bifolium, *Teijsm. & Binn. Cat. Hort. Bog.* 49, nomen. —Java.
brevilabre, *F. Muell. Fragm.* xi. 87.—Austral.
callosum, *Blume, Bijdr.* 364.—Java.
carinum, *Hance, in Journ. Bot.* xx. (1882) 359.—China.
conspicuatum, *Lindl. Or. Gen. et Sp. Orch.* 327. n. 8) ; *Slensd. Nem.* ed. II. t. 278.—Ind. or.
crassifolium, *Lindl. & Paxt. Flow. Gard.* iii. (1852–53) t. 99, 126.—Burma ?
cryptochilum, *F. Muell. in Wing, South Sc. Record*, N. S. t. (1885).—N. Guinea.
Cumingii, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 214.—Hab. ?
Dawsoniensis, Reichb. f. in Gard. Chron. (1868) 815 = Trichoglottis Dawsoniana.
deshlatum, *Lindl. Bot. Reg.* (1843) *Misc.* 6.—Manilla.
decipiens, *Lindl. l. c.* (1844) *Misc.* 11.—Zeylan.
discolor, *Lindl. l. c.* (1845) *Misc.* 59.—Ind. or.
emarginatum, *Blume, ex Teijsm. & Binn. Cat. Hort. Bog.* 49, nomen.—Java.
erectum, *Bklgr. Austral. Orch.* t. iv.—Austral.
expansum, *Reichb. f. Otto Bot. Hamb.* 52.—Ins. Mindanao.
Fordii, *Hance, in Journ. Bot.* xiv. (1876) 45.—Ins. Hongkong.
formosanum, *Hance, l. c.* xxii. (1884) 264.—China.
fuscum, Lindl. in Journ. Hort. Soc. v. (1850) 80 = latifolium.
guineense, Thw. Enum. Fl. Zeyl. 306 = maculosum, *Lindl.*

CLEISOSTOMA—
Guiberti, *Linden & Reichb. in Bot. Zeit.* xx. (1862) 375 ; *in Hamb. Gartenz.* xviii. (1862) 529.—Hab. ?
loncostum, *Lindl. Bot. Reg.* (1847) t. 43.—Manilla.
Keffordii, *F. M. Bailey, in Rep. Queensl. Acclim. Soc.* (Apr. 1884) 9.—Austral.
lanatum, *Lindl. in Journ. Hort. Soc.* iv. (1849) 164.—Burma.
latifolium, *Lindl. Bot. Reg.* (1840) *Misc.* 60.—Manilla.
longifolium, *Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind.* v. (1853) 494.—Java.
loratum, *Reichb. f. in Flora*, lv. (1872) 273.—Reg. Himal.
Macphersoni, *F. Muell. ex Benth. Fl. Austral.* vi. 297. —Austral.
macrostachyum, *Teijsm. & Binn. Cat. Hort. Bog.* 49. —Java.
maculosum, *Blume, ex Teijsm. & Binn. Cat. Hort. Bog.* 49, nomen.—Java.
maculosum, *Lindl. Gen. et Sp. Orch.* 227.—Zeylan.
maculosum, Thw. Enum. Fl. Zeyl. 304 = decipiens.
Mannii, *Reichb. f. in Flora*, lv. (1872) 274.—Reg. Himal.
roseum, *Lindl. Bot. Reg.* (1838) *Misc.* 80.—Manilla.
sagittatum, *Blume, Bijdr.* 363.—Java.
spathulatum, *Blume, l. c.* 364.—Java.
spicatum, *Lindl. Bot. Reg.* (1847) sub t. 82.—Malaya.
suaveolens, *Blume, Bijdr.* 363.—Java.
subulatum, *Blume, l. c.*—Java.
subviolaceum, *Reichb. f. in Bonplandia*, x. (1862) 335. —Philipp.
teretifolium, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 20.—Sumatra.
Thwaitesianum, Trim. in Journ. Bot. xxiii. (1885) 244 = decipiens.
tridentatum, *Lindl. Bot. Reg.* (1838) *Misc.* 83.—Austral.
undulatum, *Reichb. f. in Flora*, lv. (1872) 274.—Reg. Himal.
virginale, *Hance, in Journ. Bot.* xv. (1877) 38.—Ins. Hongkong.
vitellinum, *Reichb. f. in Linnæa*, xli. (1877) 76.—Ins. Philipp.
Wendlandorum, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 219.—Reg. Himal. ; Burma.

CLEISOSTOMA, Rafin. Fl. Tellur. iv. 80 (1836) = Convolvulus, Tourn.
villosum, Rafin. l. c. = Conv. aggregatus.

CLEISSOCRATERA, Miq. Fl. Ind. Bat. ii. 383 (1856) = Cleisocrastera, Korth. (Rubiac.).

CLEISTACHNE, Benth. in Journ. Linn. Soc. xix. (1881) 61. *GRAMINEÆ*, Benth. & Hook. f. iii. 1190.
sorgholdes, *Benth. in Hook. Ic. Pl.* t. 1379.—Afr. trop. ; Ind. or.

CLEISTANTHIUM, Kunze, in Bot. Zeit. ix. (1851) 850 = Garbera, Gronov. (Compos.).
nepalense, Kunze, l. c. = G. Kunzeana.

CLEISTANTHUS, Hook. f. ex Planch. in Hook. Ic. Pl. t. 779 (1848). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 368.
LEBIDIERA, Baill. Etud. Gén. Euph. Atlas, 50. t. 97 (1858).
LEBIDIEROPSIS, Muell. Arg. in Linnæa, xxxii. (1863) 79.
LEIOPYXIS, Miq. Fl. Ind. Bat. Suppl. 445 (1860).
NANOPETALUM, Hemsl. in Verh. Kon. Akad. Wetensch iv. (1855) 140.
acuminatus, *Muell. Arg. in DC. Prod.* xv. ii. 506.—Zeylan.
angulensis, *Weiw. ex Muell. Arg. in Journ. Bot.* ii. (1864) 339.—Afr. occ.
apodus, *Benth. Fl. Austral.* vi. 122.—Austral.
Blancoi, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 315.— Ins. Philipp.
Bolvinianus, *Muell. Arg. in DC. Prod.* xv. ii. 506.— Madag.
chartaceus, *Muell. Arg. l. c.* 507.—Reg. Himal. ; Malaya.
collinus, *Benth. & Hook. f. Gen.* iii. 368.—Ind. or.

CLEISTANTHUS—
Cunninghamii, *Muell. Arg. in DC. Prod.* xv. ii. 506.— Austral.
Dallachyanus, *Benth. ex Benth. Fl. Austral.* vi. 122. —Austral.
diversifolius, *Muell. Arg. in DC. Prod.* xv. ii. 509.— Madag.
ferrugineus, *Muell. Arg. in DC. Prod.* xv. ii. 507.— Zeylan.
ferrugineus, F. Vill. in Blanco, Fl. Philipp. ed. III. Nov. App. 187. t. 358 = Blancoi.
lanceolatus, *Muell. Arg. l. c.*—Java.
malabaricus, *Muell. Arg. l. c.* 508.—Ind. or.
monoicus, *Muell. Arg. l. c.*—China.
myrianthus, *Kurz, For. Fl. Br. Burm.* ii. 370.— Burma ; Malaya.
oblongifolius, Muell. Arg. in DC. Prod. xv. ii. 506, pro parte = chartaceus.
pallidus, *Muell. Arg. l. c.* 508.—Zeylan.
patulus, *Muell. Arg. l. c.* 508.—Ind. or.
polystachyus, *Hook. f. ex Planch. in Hook. Ic. Pl.* t. 779.—Afr. trop.
robustus, *Muell. Arg. in DC. Prod.* xv. ii. 504.— Zeylan.
semiopacus, *F. Muell. ex Benth. Fl. Austral.* vi. 123. —Austral.
stenophyllus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 242.—Burma.
stipitatus, *Muell. Arg. in DC. Prod.* xv. ii. 506.—N. Caled.
stipularis, *Muell. Arg. l. c.* 508.—Ind. or.
subglaucus, Thw. ex Trimen, Cat. Pl. Ceyl. 78 = pallidus.
sumatranus, *Muell. Arg. in DC. Prod.* xv. ii. 504.— Sumatra.
tomentosus, *Hance, in Journ. Bot.* xv. (1877) 337.— China.

CLEISTES, Rich. in Mém. Mus. Par. iv. (1818) 31 = Pogonia, Juss. (Orchid.).
Liliastrum, Reichb. f. in Linnæa, xxii. (1849) 816 = P. Liliastrum.
lutea, Lindl. Gen. et Sp. Orch. 409.—Guiana.
Miersii, Gardn. in Hook. Ic. Pl. sub t. 478–474 = P. Miersii.
montana, Gardn. l. c. = P. montana.
paludosa, Reichb. f. in Linnæa, xxii. (1849) 816 = P. paludosa.
parviflora, Lindl. Gen. et Sp. Orch. 410.—Guiana.
rosea, Lindl. l. c. = P. rosea.
speciosa, Gardn. in Hook. Ic. Pl. t. 478–474 = P. speciosa.

CLEISTOCACTUS, Lem. Illustr. Hortic. viii. (1861) Misc. 32 = Cereus, Mill. (Cact.).
Baumannii, Lem. l. c. = Cereus Baumannii.
colubrinus, Lem. l. c. = Cereus colubrinus.
rhodacanthus, Lem. l. c. = Echinopsis rhodacantha.

CLEISTOCALYX, Blume, Mus. Bot. Lugd. Bat. i. 84. t. 56 (1849) = Eugenia, Linn. (Myrtac.).
nervosus, Blume, l. c. 85 = E. nervosa.
nitidus, Blume, l. c. 84.—Java.

CLEISTOCALYX, Steud. Syn. Pl. Cyp. 151 (1855) = Eyncholepis, Vahl (Cyperac.).
junciformis, Steud. l. c. = E. oligantha.

CLEISTOCHLAMYS, Oliver, in Journ. Linn. Soc. ix. (1867) 175. *ANONACEÆ*, Benth. & Hook. f. i. 256.
Kirkii, *Oliver, l. c.*—Afr. trop.

CLEITHRIA, Steud. Nom. ed. II. ii. 746 (1841) = seq.

CLEITRIA, Schrad. Ind. Sem. Hort. Gotting. (1831) ; ex Linnæa, viii. (1833) Litt. 28 = Wendtiium, Less. (Compos.).
hirta, Schrad. ll. cc. = V. Schraderi.

CLELIA, Cassr. Nov. Stirp. Bras. Dec. 83 (1843) = Calliandra, Benth. (Legumin.).
ornata, Cassr. l. c. 84 = Call. Harrisii.

CLEMANTHUS, Klotzsch, in Peters, Reise Mossamb. Bot. 148 (1861) = Modesca, Lem. (Passifior.).
serrulata Klotzsch, l. c. = M. serrulata.

CLEMATIS, Dill. ex Linn. Gen. ed. I. 168 (1787).
RANUNCULACEÆ, Benth. & Hook. f. i. 3.
ATRAGENA, Linn. Syst. ed. I (1735); et Dis. Dans. 21 (1767); in Amoen. Acad. I. 182 (1749) (Atragene).
CHEIROPSIS, Spach, Hist. Vég. Phan. vii. 260 (1839).
CLEMATITIS, [Tourn.] Linn. Syst. ed. I (1735).
MACLATIA, Spach, Hist. Vég. Phan. vii. 273 (1839).
MURALTA, Adans. Fam. ii. 460 (1763).
SIEBOLDIA, Hoffmgg. ex Heynh. Nom. ii. 665 (1846).
TRIGULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4; ed. II. 67 (1827).
VIORNA, Reichb. Handb. 277 (1837).
VITICELLA, Dill. ex Moench, Meth. 296 (1794).
acapulcensis, *Hook. & Arn. Bot. Beech. Voy.* 410.—Mexic
acerifolia, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 2.—China.
acuminata, *DC. Syst.* i. 148.—Reg. Himal.
aemiangula, *Hook. f. & Thoms. Fl. Ind.* 5.—Reg. Himal.
aethusæfolia, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 181.—Mongolia.
affinis, *St. Hil. Fl. Bras. Mer.* i. 8.—Bras.
affinis, *Wight, Illustr.* i. 5.—Ind. or.
afoliata, *F. Buch. in Trans. N. Z. Inst.* iii. (1871) 211. —N. Zel.
ajanensis, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1886) 176.—Aa. bor.
alpina, Klotzsch, in Bot. Ergeb. Waldem. Reise 181 = orientalis.
alpina, *Mill. Gard. Dict.* ed. VIII. n. 9.—Europ.; Am. bor.
americana, Mill. i. c. n. 14 = dioica.
amplexicaulis, Edgew. in Trans. Linn. Soc. xx. (1851) 44 = connata.
anemonifera, D. Don, Prod. Fl. Nep. 192 = montana.
anemonoides, O. Kuntze, in Verh. Bot. Ver. Brand. xxvi. (1885) 149 = florida.
anethifolia, *Hook. fc. i.* t. 78.—Madag.
angustifolia, Jacq. Enum. Vindob. 310 = C. Flammula.
aphylla, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 148.—N. Zel.
apiculata, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 4.— Reg. Himal.
apiifolia, *DC. Syst.* i. 149.—Japon.
aquifolia, Stend. Nom. ed. II. i. 278, sphalm. = apiifolia.
aristata, *R. Br. in Bot. Reg.* t. 258.—Austral.
Armandi, *Franch. in Nouv. Arch. Mus. Par.* Sér. II. viii. (1885) 184.—Aa. or.
aromatica, *Lemd & C. Koch, in App. Ind. Sem. Hort. Berol.* (1856) 15; *ex Walp. Ann.* iv. 4.—Hab.?
asiatica, *Schrenk, Enum. Pl. Nov. Alt.* 68.— Scongaria; Afghanist.
asiæa, *Hort. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 272 [sic]—Tauria (calt.).
asræa, Siebold, ex Steud. Nom. ed. II. i. 278 = florida.
baccata, Pers. Syn. ii. 99 = C. Viticella.
Baldwini, *Torr. & Gray, Fl. N. Am.* i. 8.—Am. bor.
balearica, Rich. in Journ. Phys. (Févr. 1779) 137 = cirrhosa.
bannatica, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 7.—Europ.
barbellata, *Edgew. in Trans. Linn. Soc.* xx. (1851) 25. —Reg. Himal.
Bergeroni, *A. Lowell. Clemat.* 35. t. 10.—Hab.?
bicolor, Stend. Nom. ed. II. i. 278 = florida.
Bigelovi, *Torr. in Pacif. Rail. Rep.* iv. 61.—Am. bor. occ.
biloba, Stend. Nom. ed. II. i. 279, vide *triloba*, St. Hil.
biternata, *DC. Syst.* i. 149.—Japon.; China.
blanda, Hook. Journ. Bot. i. (1834) 241 = aristata.
Bojeri, *Hook. fc. Pl.* t. 10.—Madag.
bosniensis, *Vuss. ex DC. Syst.* i. 145.—Reg. Argent.
brachiata, Ker-Gawl. in Bot. Reg. t. 97 = Kerrii.
brachiata, *Thunb. Prod. Pl. Cap.* 94.—Afr. austr.
brachyura, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 221.—Corea.
bracteata, DC. Syst. i. 149 = virginiana.
bracteata, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 43.—Ind. or.
bracteosa, Banks, ex Stend. Nom. ed. I. 204, 205 = recta.
brasiliana, *DC. Syst.* i. 148.—Bras.
brevicaudata, *DC. l. c.* 138.—China.

CLEMATIS :—
Bucamara, Buch.-Ham. ex DC. Syst. i. 140 = Buchaniana.
Buchanani, D. Don, Prod. Fl. Nep. 191 = seq.
Buchananiana, Wall. Cat. n. 4677 = seq.
Buchaniana, *DC. Syst.* i. 140.—Reg. Himal.
Cadmia, *Buch.-Ham. ex Wall. Cat.* n. 4669.—Reg. Himal.; Burma.
caerulea, Lindl. Bot. Reg. t. 1955 = patens.
casmiriata, *Hance, in Journ. Bot.* viii. (1870) 71.— China.
caspitosa, Scop. Fl. Carn. ed. II. i. 389 = C. Flammula.
calycina, [Soland. in] Ait. Hort. Kew. ed. I. ii. 259 = cirrhosa.
campanifiora, *Brot. Fl. Lusit.* ii. 350.—Lusitan.
campaniflora, Lodd. ex Steud. Nom. ed. II. i. 279 = parviflora.
campestris, *St. Hil. Fl. Bras. Mer.* i. 4.—Bras.
cana, Wall. Cat. n. 4672 = Gouriana.
canadensis, Mill. Gard. Dict. ed. VIII. n. 5 = virginiana.
canaliculata, Lag. Gen. et Sp. Nov. 17 = C. Flammula.
capensis, Poir. Encyc. Suppl. ii. 296 = Anemone capensis.
caracassana, H. B. & K. ex DC. Syst. i. 141 = caripensis.
caripensis, *H. B. & K. Nov. Gen. et Sp.* v. 36.—Mexic Caiethyana, *Pursh, Fl. Am. Sept.* ii. 736.—Am. bor.
caudata, Hook. ex Hemsl. Biol. Centr. Am. Bot. i. 2 = Drummondii.
Chandleri, Hort. ex Lavall. Clemat. 40 = eriostemon.
chinensis, Osb. Itr, ed. Angl. i. 398 = recta.
chinensis, *Retz. Obs.* ii. 18. t. 2.—China.
chirsutissima, Lindl. Bot. Reg. t. 1924 = grandiflora.
chrysocarpa, *Welw. ex Oliver, Fl. Trop. Afr.* i. 5.— Afr. trop.
cirrhosa, *Linn. Sp. Pl.* 544.—Reg. Mediterr.
citalcarica, DC. Syst. i. 158 = aristata.
coccinea, Engelm. ex A. Gray, Pl. Wright. ii. 7 = C. Viorna.
coccinifiora, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 260.—N. Zel.
cognata, Steud. in Lehm. Pl. Preiss. i. 363 = aristata.
Colensoi, *Hook. f. Fl. N. Zelland.* i. 6.—N. Zel.
colorata, Lindl. Bot. Reg. t. 1924 = grandiflora.
columbiana, Torr. & Gray, Fl. N. Am. i. 11 = verticillaris.
communata, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 138.—Afr. trop.
comosa, Hook. f. Syst. i. 157 = integrifolia?
connata, *DC. Prod.* i. 10.—Reg. Himal.
contorta ×, Carr. in Rev. Hortic. (1879) 380.
cordata, Pursh, Fl. Am. Sept. ii. 354 = virginiana.
cordata, *Royle, Illustr. Bot. Himal.* 51.—Ind. or.
cordata, Sims, Bot. Mag. t. 1316 = crispa.
cordifolia, DC. Syst. i. 142 = virginiana.
coriacea, DC. l. c. 146 = aristata.
coriacea, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 208.—Java.
corymbosa, Poir. Encyc. Suppl. ii. 297 = recta.
crassifolia, *Benth. Pl. Hongk.* 7.—China.
cronata, Jord. in Billot, Annot. Fl. Fr. Allem. 12 = C. Vitalba.
crispa, Encyc. ii. 44 = cylindrica.
crispa, *Linn. Sp. Pl.* 543.—Am. bor.
crispa, Thunb. Fl. Jap. 229 = paniculata.
cylindrica, Sims, Bot. Mag. t. 1160 = crispa.
dahurica, DC. Prod. i. 7 = daurica.
dasycnema, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 138.—Borneo.
daurica, *Pers. Syn.* ii. 99.—Aa. temp.
Davidiana, Decne. ex Verlot, in Rev. Hortic. (1867) 90 = heracleæfolia.
davurica, Ledeb. Fl. Ross. i. 4 = daurica.
denticulata, Vell. Fl. Flum. 240; v. t. 134 = Hilarii.
depauperata, *Colenso, in Trans. N. Z. Inst.* i. (1868) 248, nomen.—N. Zel.
dioica, *Linn. Syst. ed.* X. 1084.—Jamaica.
dioica, Lour. Fl. Cochinch. 344 = Loureiriana.
dioica, *Sieber, ex Presl, in Oken, Isis, xxi.* (1828) 373 = caripensis.
discolor, *Garbin. in Hook. Lond. Journ. Bot.* ii. (1843) 320.—Bras.
discolor, Steud. in Lehm. Pl. Preiss. i. 363 = aristata.
dissecta, *Baker, in Journ. Linn. Soc.* xx. (1883) 67.— Madag.

CLEMATIS :—
distorta, *Lavall. Clemat.* 37. t. 11.—Hab.?
divaricata, Jacq. Eclog. Fl. i. 51. t. 33 = crispa.
divaricifolia, *DC. Syst.* i. 158.—Hab.?
diversifolia, Gilib. Exercit. i. 338 = C. Flammula.
dominica, *Lam. Encyc.* ii. 45.—Ind. occ.
Douglasi, *Hook. Fl. Bor. Am.* i. t. 1.—Am. bor. occ.
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 9.—Am. bor.
dumosa, Salisb. Prod. 371 = C. Vitalba.
edentata, *Baker, in Journ. Linn. Soc.* xxi. (1884) 316. —Madag.
elliptica, Endl. in Enum. Pl. Huegl. 1 = aristata.
elongata, Trait. Archiv, t. 149 = integrifolia.
erecta, Linn. Syst. ed. XII. 477; All. Fl. Pedem. i. 296 = recta.
eriopoda, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 238.—Japon.
eriostemon, *Decne. in Rev. Hortic.* Sér. IV. i. (1852) 341.—Hab.?
excelsior ×, Van Houtte, in Fl. des Serres, Sér. II. x. (1874) 24. t. 2054-55.
Fawcettii, *F. Muell. Fragm.* x. 1.—Australi.
filifera, Benth. Pl. Hartw. 285 = reticulata.
Flammula, *Linn. Sp. Pl.* 544.—Reg. Mediterr.
Flammulastrum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 158.—Cuba.
flava, DC. Syst. i. 135 = orientalis.
flavescens, *Regin. Fl. Ludov.* 81.—Am. bor.
floribunda, *Kurz, in Journ. Bot.* v. (1867) 240.—Ind. or.
floribunda, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 9.—N. Granat.
florida, *Thunb. Fl. Jap.* 240.—Japon.
fluminensis, *Vell. Fl. Flum.* 240; v. t. 133.—Bras.
foetida, *Raoul, Choix Pl. Nouv. Zél.* 28.—N. Zel.
forsteri, J. F. Gmel. Syst. 873 = hexasepala.
Fortunei, T. Moore, in Gard. Chron. (1868) 460, 676 = Williamsii.
fragrans, Salisb. Prod. 371 = virginiana.
fragrans, Tenore, Prod. Fl. Nap. p. xxxii = C. Flammula.
francofurtensis, Rinz. ex C. Koch, Dendrol. i. 491 = C. Viticella.
Fremonti, *S. Wats. in Proc. Am. Acad.* x. (1875) 339. —Am. bor. occ.
fruticosa, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 180.—China.
fulva, *Zoll. & Moritzi, Syst. Verz. Zoll.* 35.—Java.
fusca, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 60.— China; Japon.
Gebleriana, Bong. in Bull. Sc. Acad. Pétersb. viii. (1841) 382 = soongarica.
guntiacoides, *DC. Syst.* i. 159.—Austral.
Gilbertiana, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 278 [bis] = aristata.
glabra, *DC. Syst.* i. 143.—Ins. S. Doming.
glandulosa, Blume, Bijdr. i. = smilacifolia.
glauca, Class. in Goebel, Reise, ii. 347 = orientalis.
glauca, Willd. Berl. Baumz. 65 = orientalis.
glaucescens, Presl, in Mss. Senckenb. ii. (1837) 268 = Thunbergii.
globosa, Royle, Illustr. Bot. Himal. 51 = orientalis.
glychoides, *DC. Syst.* i. 145.—Austral.
Gondotiana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 10.—N. Granada.
Gouriana, *Roxb. Hort. Beng.* 43; *Fl. Ind.* ii. 570.— Ind. or.; Malaya.
gracilis, Edgew. in Trans. Linn. Soc. xx. (1851) 24 = connata.
Grahami, *Benth. Pl. Hartw.* 8.—Mexic.
grandiflora, *DC. Syst.* i. 151.—Afr. trop.
grata, O. Hoffm. ex Baker, in Journ. Linn. Soc. xx. (1883) 88 = Wightiana.
grata, *Wall. Pl. As. Rar.* i. 53. t. 98.—Reg. Himal.; China; Afr. trop.
grewioides, Lindl. Bot. Mag. t. 4495 = orientalis.
grewioides, *Lindl. in Journ. Hort. Soc.* i. (1846) 267. —Reg. Himal.
gueriana, *DC. Syst.* i. 140.—Reg. Himal.
grosa, *Benth. Pl. Hartw.* 33.—Mexic.
grosa, *Wall. Cat.* n. 4671.—Sumatra.
Guadalupæa, Pers. Syn. ii. 99 = acapulcensis.
Guaraná ×, Lem. & Verlot, in Illustr. Horticole, iv. (1857) 117.
Hookeriana, *Presl, Rel. Haenk.* ii. 68.—Peruv.
hakoneusis, *Franch. & Sav. Enum. Pl. Jap.* ii. 228.— Japon.

CLEMATIS :—
Hancockiana, *Maxim. in Bull. Soc. Nat. Moac.* liv. (1879) i. 1.—China.
Hartwegi, Hort. ex C. Koch, Dendrol. i. 434 = reticulata.
havanensis, *H. B. & K. ex DC. Syst.* i. 152.—Cuba.
hedysarifolia, *DC. l. c.* 148.—Ind. or.
Henderaonii, Hort. ex Steud. Nom. ed. II. i. 379 = reticulata.
Henryi, *Hort.—Cf. Gard. Chron.* (1878) ii. 489.—Hab.?
heracleaefolia, *DC. Syst.* i. 126.—China.
hexapetala, Linn. f. Suppl. 271 = hexasepala.
hexapetala, Pall. Reise, iii. 735 = C. Flammula.
hexasepala, *DC. Syst.* i. 146.—N. Zel.
Hilarii, *Spreng. Syst.* v. 177.—Bras.
hirsuta, Guill. & Perr. Fl. Seneg. Tent. i. 1 = Thunbergii.
hirsutissima, Pursh, Fl. Am. Sept. ii. 385 = Anemone patens.
hispanica, Mill. Gard. Dict. ed. VIII. n. 3 = recta.
holosericea, *Pursh, Fl. Am. Sept.* ii. 384.—Am. bor.
Hookeri, Decne. in Fl. des Serres, Sér. II. xiii. (1860) 389 = heracleaefolia.
Hothia, *Kurz, For. Fl. Brit. Burma*, i. 17.—Burma.
hyemalis, Munro, ex Hook. f. & Thoms. Fl. Ind. 9 = orientalis.
Ibarensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 264.—Madag.
incisodentata, A. Rich. Tent. Fl. Abyss. i. 2 = grata.
inclinata, Scop. Fl. Carn. ed. II. i. 389 = integrifolia.
indica, Heyne, ex Roth, Nov. Pl. Sp. 252 = Gouriana.
indivisa, Steud. in Lehm. Pl. Preiss. i. 262 = aristata.
indivisa, *Willd. Sp. Pl.* ii. 1291.—N. Zel.
indivisa, *Banks. in Bull. Soc. Linn. Par.* i. (1883) 831.—Madag.
insulensis, Hort. ex C. Koch, Dendrol. i. 433 = crispa.
integra, *Vell. Fl. Flum.* 241 ; v. t. 185.—Bras.
integrifolia, Forst. f. Prod. 49 = indivisa.
integrifolia, *Linn. Sp. Pl.* 544.—Eur. asstr. ; As. bor.
intermedia, *Carr. in Rev. Hort.* (1866) 339.—Hab.?
Intricata, *Bunge, in Mém. Sav. Etr. Acad. Pétersb.* vi. (1855) 75.—China.
inversa, *Grig. Notul.* iv. 700.—Ind. or.
japonica, Boiss. Diagn. Ser. I. vi. 3 = orientalis.
italica, *Schmid. Icon. Ill. Ind.* 2.—Europ.
Jackmanni x, Van Houtte, in Fl. des Serres, xvi. (1865) 89.
japonica, *Thunb. Fl. Jap.* 240.—Japon.
javana, DC. Syst. i. 153 = Gouriana.
javanica, Steud. Nom. ed. II. i. 379 = pmec.
Jubata, *Bisch. Del. Sem. Hort. Heidelb.* (1849) 4 ; *ex Walp. Ann.* ii. 4.—Hab.?
Junghuniana, *De Vriese, in Mig. Pl. Jungh.* 75.—Java.
kamtschatica, *Bong. Vers. Saisang-Nor Pf.* 10 = Kamtchatka.
Kerii, *Steud. Nom.* ed. II. i. 379.—Afr. austr.
Kirilowi, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 210.—China.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 5.—Afr. trop.
Koumbotsa, *Decne. in Fl. des Serres*, xxiii (1880) 284.—Japon.
laeta, Salisb. Prod. 370 = cirrhosa.
lanuginosa, *Lindl. & Paxt. Flow. Gard.* iii. (1852-53) 107.—China.
lasiandra, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 218.—Japon.
lasiandra, Hort. Gorenk. ed. II. (1812) 47 = C. Flammula.
lasiantha, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 9.—Am. bor. occ.
lathyrifolia, Bess. ex Reichb. Fl. Germ. Excurs. 734 = recta.
Lavallei, *Decne. in Fl. des Serres*, xxiii. (1880) 284.—Japon.
lazulica, *Baker, in Journ. Linn. Soc.* xxi. (1884) 317.—Madag.
Lechenaultiana, *DC. Syst.* i. 151.—Java ; Ind. or.
leptomera, *Hance, in Journ. Bot. xvii.* (1880) 357.—China.
leptophylla, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 9.—Am. bor.
Hexuosifolia, Steud. in Lehm. Pl. Preiss. i. 262 = microphylla.
Hindsiana, DC. Syst. i. 138 = crispa.
Hookeriana, DC. l. c. 140 = grewiaefolia ?
Hookeriana, D. Don, Prod. Fl. Nep. 191 = Buchaniana ?

CLEMATIS :—
longicaudata, Ledeb. Fl. Ross. i. 3 = orientalis.
longicauda, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 2.—Abyssin.
longiloba, *DC. Syst.* i. 136.—Chili.
longipes, *Freyn, in Bremen Abh.* viii. (1880) 5.—Madag.
Loureiriana, *DC. Syst.* i. 144.—China.
lugubris, Salisb. Prod. 371 = C. Viticella.
Lulovi, Hort. ex C. Koch, Dendrol. i. 435 = patens.
macropetala, *Ledeb. Ic. Pl. Fl. Ross.* i. 5. t. 11.—Siberia.
magniflora, *Hoffegg. Vers. Pft. Nachtr.* i. 167.—Hab.?
mandschurensis, Mort. ex C. Koch, Dendrol. i. 439 = fusca.
mandshurica, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 388 = recta.
marata, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 385.—N. Zel.
maritima, Lam. Encyc. ii. 174 = C. Flammula.
maritima, Linn. Sp. Pl. 767 = recta.
Massoniana, DC. Syst. i. 135 = brachiata.
Massoniana, Maxim. Ind. Fl. Pekin. in Prim. Fl. Amur. 466 = Kirilowi.
mauritiana, *Lam. Encyc.* ii. 42.—Ins. Mauric. ; Madag.
Maximowiciana, *Franch. & Sav. Enum. Pl. Jap.* ii. 281.—Japon.
Mechowiana, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 171.—Afr. trop.
meduseae, *Planch. & Linden, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 11.—N. Granat.
mendocina, *Phil. in Anal. Univ. Chil.* (1862) ii. 889.—Chili.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 297.—China.
microcuspis, *Baker, in Journ. Linn. Soc.* xxi. (1884) 317.—Madag.
microphylla, *DC. Syst.* i. 147.—Austral.
millefoliata, *Eichl. in Mart. Fl. Bras.* xiii. t. 150.—Peruv.
minor, *Curt. Fl. Cochinch.* i. 845.—China.
Mociniana, *G. Don, Gen. Syst.* i. 5.—Mexic.
monstrosa, Hort. ex C. Koch, Dendrol. i. 436 = patens.
montana, *Buch.-Ham. ex DC. Syst.* i. 164.—Reg. Himal.
montana, D. Don, Prod. Fl. Nep. 192 = napaulensis.
montevidensis, *Spreng. Syst.* iii. 867.—Reg. Argent.
Munroiana, Wight, Illustr. Ind. Bot. i. t. 1 = smilacifolia.
nannophylla, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 219.—Ind. or.
napaulensis, *DC. Syst.* i. 164.—Nepal.
naravelioides, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 119.—Ind. or.
nepalensis, Koyle, Illustr. Bot. Himal. 51 = barbellata.
nervata, Benth. Pl. Hartw. 5 = Drummondii.
nutans, Grant, Stirp. Austr. ed. I. fasc. ii. 110 ; ed. II. 127 = integrifolia.
nutans, *Royle, Illustr. Bot. Himal.* 51.—Reg. Himal.
occidentalis, A. Cunn. ex Hook. in Mitch. Journ. Trop. Austral. 366 = microphylla.
occidentalis, *DC. Prod.* i. 10.—Hab.?
ochotensis, *Pair. Encyc. Suppl.* ii. 298.—Sibir. or.
ochroleuca, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 260.—Am. bor.
odora, *Wendor. ! ind. Sem. Hort. Herb. App.* n. 4.'—*ex C. Steud. Nom.* ed II. i. 380.—Hab.?
odontia, *Wall. ex Steud. Nom.* ed. II. i. 380.—Ind. or.
oligophylla, *Hook. Ic. Pl.* t. 80.—Madag.
Olivieri, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 165.—Abyssin.
oreophila, Hance, in Walp. Ann. ii. 8 = Meyeniana.
orientalis, *Linn. Sp. Pl.* 543.—Reg. Himal. ; Orient.
orientalis, Sieb. ex Steud. Nom. ed. II. i. 380 = C. Flammula.
ovata, *Pursh, Fl. Am. Sept.* ii. 386.—Am. bor.
Ovenizae, *Hare. in Hare. & Sond, Fl. Cap.* i. 2.—Afr. trop.
Pallasii, J. F. Gmel. Syst. 573 = C. Flammula.
pallida, *A. Rich. Esc. Fl. Cub.* 10.—Cuba.
pandurata, Steud. Nom. ed. II. i. 380, sphalm. = montana.
paniculata, J. F. Gmel. Syst. 573 = indivisa.
paniculata, *Thumb. in Trans. Linn. Soc.* ii. (1794) 337.—Japon.
Parkinsoniana, *Colenso, in Trans. N. Z. Inst.* xii. (1880) 398.—N. Zel.

CLEMATIS :—
parviflora, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 289.—N. Zel.
parviflora, *DC. in Mém. Soc. Phys. Genev.* i. (1823) 493.—Hab.?
parviflora, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 9.—Am. bor.
parvifolia, Edgew. in Trans. Linn. Soc. xx. (1846) 25 = orientalis.
parvifolia, Hook. f. & Thoms. Fl. Ind. 9 = gravelens.
parviloba, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 241.—China.
patens, *Morr. & Decne. in Bull. Acad. Brux.* iii. (1836) 178.—Japon.
pauciflora, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 9.—Calif.
pedicellata, Sweet, Hort. Brit. ed. I. 2 ; Loud. Hort. Brit. 1 = cirrhosa.
pensylvanica, *Donn, Hort. Cantab.* ed. IV. 129 ; *Hort. ex Turca. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 278 [sic].—Am. bor.
perulata, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 158.—Bras.
peruviana, *DC. Syst.* i. 141.—Peruv.
Petersiana, Klotzsch, in Peters, Reise Mossamb. Bot. i. 170 = grata.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 1.—Ins. Fiji.
Pierotii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 1.—Japon.
pilosa, Dulac, Fl. Hautes-Pyr. 211 = C. Vitalba.
pimpinellifolia, *Hook. Ic. Pl.* t. 77.—Madag.
pinnata, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 218.—China.
Pitcheri, Sarg. ex Lavall. Clemat. 60 = Sargenti.
Pitcheri, *Torr. & Gray, Fl. N. Am.* i. 10.—Am. bor. occ.
Pluknetii, DC. Syst. i. 158 = Catesbyana.
Poiretii, Hort. ex Lavall. Clemat. 40 = eriostemon.
polyopetala, *Bertol. Fl. Guat.* 23.—Guatemala.
polygama, *Vacq. Bull. Fl. Carib.* 24.—Ind. occ.
polymorpha, Viv. Fl. Cors. Sp. Nov. 9 = cirrhosa.
polypetala, DC. Syst. i. 167 = Anemone japonica.
polypetala, *Poir. Encyc. Suppl.* ii. 296.—Japon.
popaliticola, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 272.—Venezuela.
pseudoatragene, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 160.—Am. bor.
pseudograndiflora, *O. Kuntze, l. c.* 138.—Afr. trop.
pseudo-orientalis, *O. Kuntze, l. c.* 171.—Persia.
puberula, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 4.—Reg. Himal.
pubescens, *Benth. Pl. Hartw.* 5.—Mexic.
pubescens, Haeg. Enum. Pl. Hoeg. ? = aristata.
pulchella, Pers. Syn. ii. 99 = C. Viticella.
punduana, Wall. Cat. n. 4682 = montana.
Purshii, D. Dietr. Syn. Pl. iii. 345 = virginiana.
quadribracteolata, *Colenso, in Trans. N. Z. Inst.* xiv. (1882) 329.—N. Zel.
recta, *Linn. Sp. Pl.* 544.—Europ. austr.
reticulata, *Walt. Fl. Carol.* 156.—Am. bor.
revoluta, *Hort. Par. ex Steud. Nom.* ed. II. i. 380 = parviflora.
Robertsiana, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 39.—Afghan.
rosea, Hort. ex Lavall. Clemat. 35 = Bergeroni.
rosea, Sm. in Abbot, Nat. Hist. Lepidopt. 201. t. 101 = reticulata.
rubella, Pers. Syn. ii. 100 = C. Flammula.
rubro-violacea x, Van Houtte, in Fl. des Serres, xvi. (1865) 87 = lanuginosa.
Sargenti, *Lavall. Clemat.* 60. t. 18.—Am. bor.
Savatieri, *Decne. in Fl. des Serres*, xxiii. (1880) 286.—Japon.
scabiosa, *Hilz. & Bej. ex Bull. Bull. Soc. Linn. Par.* i. (1885) 831, nomen.—Madag.
scabiosaefolia, *DC. Syst.* i. 154.—Ind. or.
sawaiana, Bentl. Handb. Forstb. ii. 1199 = C. Vitalba.
Schlechteri, C. Koch, in App. Ind. Sem. Hort. Berol. (1856) 16 = cylindrica.
Scottii, *Porter, Syn. Fl. Colorado*, 1.—Am. bor. occ.
Seemanni, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 144.—Peruv.
semitriloba, Lag. Gen. et Sp. Nov. 17 = cirrhosa.
sepium, Lam. Fl. Fr. iii. 306 = C. Vitalba.
sericea, *H. B. & K. ex DC. Syst.* i. 144.—Am. austr.
sericea, Michx. Fl. Bor. Am. i. 319 = ochroleuca.

CLEMATIS:—
sibirica, Mill. Gard. Dict. ed. VIII. n. 12 = angustifolia.
Sieboldii, D. Don, ex Steud. Nom. ed. II. i. 380 = florida.
simensis, *Fresen. in Mus. Senckenb.* ii. (1837) 267.—Abyssin.
sinensis, Lour. Fl. Cochinch. 844 = chinensis.
Simsii, Sweet, Hort. Brit. ed. I. 1 = crispa.
smilacifolia, *Wall. in As. Res.* xiii. (1820) 402.—Ind. or.; Malaya.
smilacina, Blume, Bijdr. 1 = smilacifolia.
songarica, Siev. ex Steud. Nom. ed. II. i. 380 = glauca.
songorica, *Bunge, Del. Sem. Hort. Dorp.* (1839) 8.—*Cf. Linnaea*, xiv. (1840) *Litt.* 119.—Sibiria.
Sonnerati, Pers. Syn. ii. 99 = mauritiana.
Standishii, Van Houtte, in Fl. des Serres, xvi. (1865) 89 = patens.
Stanleyi, *Hook. Ic. Pl.* t. 589.—Afr. austr.
stans, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. (1846) II. 177.—Japon.
stenopetala, R. Br. ex DC. Syst. i. 147 = glycinoides.
stenophylla, Fraser, ex Hook. in Mitch. Trop. Austr. 364 = microphylla.
stenopsis, DC. Syst. i. 147 = glycinoides.
stipulata, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 164.—Mexic.
strata, *Eagln. Fl. Ludov.* 83.—Am. bor.
stricta, *Wender. in Ind. Sem. Marb.* (1836).—*Cf. Linnaea*, xii. (1838) *Litt.* 85.—Hab.?
strigillosa, *Baker, in Journ. Linn. Soc.* xviii. (1881) 245.—Madag.
Strobeliana, *Ces. in Atti Soc. Acc. Sc. Nap.* v. (1873) 8. n. 7.—Europ.
Stronachii, *Hance, in Journ. Bot.* xvi. (1878) 103.—China.
suaveolens, Salisb. Prod. 371 = C. Flammula.
subpeltata, Wall. Pl. As. Rar. i. 19. t. 20 = smilacifolia.
substipulata, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 147.—Ind. or.
subtriflora, Walp. Rep. i. 8, sphalm. = seq.
subtriloba, *Nees, ex G. Don, Gen. Syst.* i. 7.—Mexic.
subumbellata, *Kurz, For. Fl. Brit. Burma*, i. 16.—Burma.
sulcata, Wall. Cat. n. 4667 = C. Cadmia.
Tatarinowii, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 214.—China.
tenuiloba, Bess. ex Nym. Consp. 1 = C. Vitalba.
tenuiflora, DC. Prod. i. 3 = recta.
tenuifolia, Poir. Encyc. Suppl. ii. 298 = Anemone tenuifolia.
tenuifolia, Royle, Illustr. Bot. Himal. 51 = orientalis.
ternifiora, DC. Syst. i. 138 = recta.
Tesane, Hamilt. Cat. Hort. Bog. Alt. 179 = Norvehiana.
texensis, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 446 = C. Viorna.
thalictroides, *Steud. in Flora*, xxxix. (1856) 407.—Peruv.
Thunbergii, *Steud. Nom.* ed. II. i. 380.—Afr. austr.
tibetana, *O. Kuntze, in Verh. Bot. Ver Brand.* xxvi. (1885) 179.—Tibet.
tortuosa, Wall. Cat. n. 4675 = Buchaniana.
Tourneforti, *DC. Syst.* i. 167.—Oriens.
trifida, *Host. Ic. Pl.* t. 79.—Madag.
trifoca, Vahl, Symb. Bot. iii. 74 = mauritiana.
trifoliata, *Thunb. in Trans. Linn. Soc. ii.* (1794) 337.—Japon.
trifolia, *Heyne, ex Roth, Nov. Pl. Sp.* 251.—Ind. or.
trifolia, Sc. Hil. Fl. Bras. Mer. i. 3 = montevidensis.
triloba, Thunb. Prod. Pl. Cap. 94 = Thunbergii.
trinervia, Buch.-Ham. ex DC. Syst. i. 148 = acuminata.
trisecta, *DC. l. c.* 156.—Hab.?
tubulosa, Hook. Bot. Mag. t. 6601 = heracleæfolia.
tuberosa, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 146 = heracleæfolia.
tunbridgensis ×, Hort. ex Lavall. Clemat. 13.
umbraticola, Schur, in Verh. Naturf. Ver. Brunna, xv. 11. (1877) 8 = uncin.
uncinata, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 255.—Hongkong.
uniflora, Balb. Cat. Hort. Taur. (1810) 21 = ochroleuca.
urophylla, *Franch. in Bull. Soc. Linn. Par.* i. (1884) 435.—China.
velutina, Edgew. in Trans. Linn. Soc. xx. (1859) 24 = connata.

CLEMATIS:—
venosa, Royle, Illustr. Bot. Himal. 51 = connata.
verticillaris, *DC. Syst.* i. 166.—Am. bor.
villosa, *DC. l. c.* 154.—Ind. or.
violacea, *A. DC. in Mem. Soc. Phys. Genev.* xi. (1845) 67.—Hab.?
Viorna, Andr. Bot. Rep. i. t. 71 = crispa.
Viorna, *Linn. Sp. Pl.* 543.—Am. bor.
viorndides, *Schrad. ex Steud. Nom.* ed. II. i. 380.—Am. bor.
virginiana, *Linn. Cent.* i. 15; *Amœn. Acad.* iv. 275.—Am. bor.
virginiana, Lour. Fl. Cochinch. 845 = apiifolia.
virginica, Thunb. Fl. Jap. 240 = biternata.
viridiflora, Bert. Misc. Bot. xix. 7. t. 3 = grata.
vitæa, *Sterl. Hort. Nymph. ex Steud. Nom.* ed. II. i. 380.—Hab.?
Vitalba, Buch.-Ham. ex Steud. l. c. 379, 380 = grewiæfolia.
Vitalba, *Linn. Sp. Pl.* 544.—Europ.; Afr. bor.; Reg. Caucas.
Vitalella, *Linn. l. c.* 543.—Europ. or.; As. Min.; Persia.
vitifolia, Wall. Cat. n. 4676 = Buchaniana.
Walteri, Pursh, Fl. Am. Sept. ii. 384 = crispa.
Welwitschii, *Hiern, ex O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 171.—Afr. trop.
Wenderothii, *Steud. Nom.* ed. II. i. 380.—Europ.
Wightiana, *Wall. Cat.* n. 4674.—Ind. or.
Williamsii, *A. Gray, ex Perry, Jap. Exp.* ii. 306.—Japon.
Wyethii, Nutt. ex Torr. & Gray, Fl. N. Am. i. 8 = Douglasii.
xanthoricæ, Sweet, Hort. Brit. ed. II. 1 = sec.
zanzibarensis, *Boj. ex Loud. Hort. Brit.* 228.—Ins. Zanzibar.
zeylanica, Poir. Encyc. Suppl. ii. 296 = Naravelia zeylanica.
Zollingeri, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 11. 374 [bis].—Java.

CLEMATITARIA, Bur. Monog. Big. 44 (1864) = Bignonia, Linn.

CLEMATITIS, [Tourn.] Linn. Syst. ed. I (1735) = Clematis, Dill. (Ranunculac.).
brachiata, Moench, Meth. Suppl. 103 = Clematis virginiana.
cordifolia, Moench, l. c. 104 = Clematis virginiana.
crispa, Moench, Meth. 296 = Clematis crispa.
erecta, Steud. Nom. ed. I. 206 = Clematis recta.
flava, Moench, Meth. 296 = Clematis orientalis.
integrifolia, Moench, l. c. = Clematis integrifolia.
Vitalba, Moench, l. c. = Clematis Vitalba.

CLEMATOPSIS, Boj. ex Hook. Ic. Pl. t. t. 10 (1887) = Clematis, Dill. (Ranunculac.).
sumaculana, Boj. l. c. = Clematis Bojeri.

CLEMENTEA, Cav. in Anal. Cienc. Nat. vii. (1804) 68 = Cæsalvinia, Adans. (Legumin.).
nitida, Cav. l. c. t. 67 = Cæs. emiliformis.

CLEOBULA, Vell. Fl. Flum. 187 (1825). *INC. SED.*
CHEOBULA, Vell. Fl. Flum. iv. t. 146 (1827).
pinnata, *Vell. l. c.*—Bras.

CLEOBULLA, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 181. *LEGUMINOSÆ*, Benth. & Hook. f. i. 587.
diodecidden, *Benth. in Mart. Fl. Bras.* xv. i. 168.—Am. trop.
leiantha, *Benth. l. c.*—Bras.
mutisiflora, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 181.—Bras.

CLEOCHROMA, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 340 = Lophocosta, Benth. (Solanac.).
calycina, Miers, l. c. 350 = I. calycina.
grandiflora, Miers, l. c. 351 = I. grandiflora.
macrocalyx, Miers, l. c. 350 = I. macrocalyx.

CLEODORA, Klotzsch, in Wiegm. Archiv, vii. (1841) 196 = Croton, Linn. (Euphorbiac.).
Sellowiana, Klotzsch, l. c. 197 = Croton sphærogynus.

CLEOME, Linn. Syst. ed. I (1735). *CAPPARIDEÆ*, Benth. & Hook. f. i. 106.
ALEOME, Neck. Elem. iii. 68 (1790).
ANOMALOSTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 169 (1861).
ARIVELA, Rafin. Sylva Tellur. 110 (1838).
ATALANTA, Nutt. Gen. Am. ii. 78 (1818).
AUBLON, Rafin. Sylva Tellur. 110 (1838).
BUSBIA, Bunge, Del. Sem. Hort. Dorpat. (1859) 4.
CHILOCALYX, Klotzsch, in Peters, Reise Mossamb. Bot. 164 (1861).
COALISIA, Rafin. Sylva Tellur. 114 (1838).
CORYMANDRA, Schrad. Cat. Sem. Hort. Gotting. (1846).
DECASTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 167 (1861).
DIANTHERA, Klotzsch, l. c. 160. t. 27 (1861).
DIORIMASPERMA, Rafin. Sylva Tellur. 111 (1838).
HEMICOLA, Rafin. l. c. (1838).
ISEXINA, Rafin. l. c. 112 (1838).
LAGANIA, Ruiopf, ex Rafin. l. c. 110 (1838).
MELANOSTIGMA, Rafin. l. c. (1838).
MICAMEE, Adans. Fam. ii. 407 (1763).
MITOSTYLIS, Rafin. Sylva Tellur. 114 (1838).
MOZAMBE, Rafin. l. c. 113 (1838).
ONCUPIS, Rafin. l. c. 114 (1838).
PERICLA, Rafin. l. c. 112 (1838).
PERITOMA, DC. Prod. i. 237 (1824).
PHYSOSTEMON, Mart. Nov. Gen. et Sp. i. 78. t. 45 (1824).
PODOGYNE, Hoffngg. Verz. Pfl. 185 (1840).
POLANISIA, Rafin. in Am. Monthly Mag. (1818) 367; Journ. Phys. (Août 1819) 93.
RORIDA, J. F. Gmel. Syst. 960 (1791).
RORIDULA, Forsk. Fl. Aegypt. Arab. 85 (1775).
SCOLOPERMA, Rafin. Sylva Tellur. 111 (1838).
SIERULA, Rafin. l. c. 112 (1838).
SILIQUARIA, Forsk. Fl. Aegypt. Arab. 78 (1775).
SINAPISPERMUM, Tourn. ex Hall. in Rupp. Fl. Jen. 269 (1745).
STYLISTA, Rafin. Sylva Tellur. 112 (1838).
SYMPHOSTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 159 (1861).
TARENAYA, Rafin. Sylva Tellur. 111 (1838).
TETRATELEIA, Sond. in Harv. & Sond. Fl. Cap. i. 58 (1859).
aculeata, *Linn. Syst.* ed. XII. iii. 232.—Am. trop.
acuta, Schum. & Thonn. in Dansk. Vidensk. Selsk. Afh. iv. (1828) 67 = Gynandropsis pentaphylla.
affinis, *DC. Prod.* i. 241.—Bras.
affinis, Sprang. Syst. iv. Cur. Post. 188 = Blumeana.
albescens, *Franch. in Revoil, Comalli*, 11.—Afr. trop.
alticaea, Blanco, Fl. Filip. ed. I. 592 = Gynandropsis pentaphylla.
alticolora, Blanco, l. c. ed. II. 363 = Gynandropsis pentaphylla.
angulata, *Schult. f. Syst.* vii. 45.—Java.
angustifolia, Forsk. Fl. Aegypt. Arab. 120 = tenella.
anomala, *H. B. & K. Nov. Gen. et Sp.* v. 85.—N. Granad.
aphylla, Thunb. Prod. Pl. Cap. 109 = Schepperia juncea.
arabica, *Linn. Cent. Pl.* i. 20.—Afr. bor.; Arab.
arborea, *H. B. & K. Nov. Gen. et Sp.* v. 86.—Venezuela.
arborea, Schrad. in Goett. Gel. Anz. i. (1821) 707 = Cratæva Tapia.
arborea, Wetsm. in Syll. Retzab. i. (1824) 227 = dendroides.
arnuta, *Thunb. Prod. Pl. Cap.* 109.—Afr. austr.
Aschersoniana, Pfund, in Flora, lvii. (1874) 413 = trinervia.
aspera, *Korn. in DC. Prod.* i. 241.—Ind. or.
atriplicifolia, Schott, in Schreth Neuhr. Oestr. Naturf. i. 239 = dendroides.
aurea, *Celak. in Oestr. Bot. Zeitschr.* xxxiv. (1884) 113.—Graecia.
aurea, Nutt. ex Torr. & Gray, Fl. N. Am. i. 133 = lutea.
bicolor, *Garcin. in Hook. Lond. Journ. Bot.* ii. (1843) 226.—Bras.
Blumeana, *D. Dietr. Syn. Pl.* ii. 1085.—Java.
bonariensis, *Oliver, Fl. Trop. Afr.* i. 81.—Afr. trop.
brachycarpa, *Vahl, ex DC. Prod.* i. 240.—Afr. bor.; Arab.; Ind. bor. occ.
brasiliensis, *Willem. in Syll. Retzab.* i. (1824) 228.—Bras.
brevifolia, *Schum. & Thonn. in Dansk. Vidensk. Selsk.* Afh.

CLEOME:—
Bungei, *Stend. Nom.* ed. II. i. 321.—China.
Burmanni, *Wight & Arn. Prod.* 25.—Ind. or.
canadensis, Hort. ex Schult. f. Syst. vii. 38 = cardaminoides.
Candelabrum, *Sims, Bot. Mag.* t. 2656.—Hab.?
cannacrus, Stev. in DC. Prod. i. 241 = omithopodioides.
capensis, Linn. Sp. Pl. ed. II. 940 = Heliophila callosa.
capparoides, Willd. ex Eichl. in Mart. Fl. Bras. xiii. i. 247 = anomala.
cardaminoides, *Hoffmgg. Verz. Pfl.* 99.—Afr. austr.
cardinalis, *DC. Prod.* i. 238.—Mexic.
Chelidonii, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 56 = strigosa.
Cheiidonii, *Linh. f. Suppl.* 300.—Ind. or.; Java.
chilensis, *DC. Prod.* i. 238.—Chili.
chiloeiisis, *Oliver, Fl. Trop. Afr.* i. 81.—Afr. trop.
chrysantha, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 274.—Arab.; Nubia.
ciliata, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* iv. (1828) 67.—Afr. trop.
colutaeoides, *Botan. Diagn.* Ser. I. i. 3.—Persia.
cordata, Burch. ex DC. Prod. i. 239 = monophylla.
cordata, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 68 = Ehrenbergiana.
cordobensis, *Eichl. ex Griseb. in Goett. Abh.* xix. (1874) 78.—Am. austr.
cremolobe, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 318.—Mexic.
crenopetala, *DC. in Mém. Soc. Phys. Genèv.* vi. (1836) 220. t. 2.—Am. austr.
cubensis, A. Rich. Ess. Fl. Cub. 74 = Houstoni.
cuneifolia, *Muhl. ex Nutt. Gen. Am.* ii. 78.—Georgia.
cypria, *Colak. in Oestr. Bot. Zeitschr.* xxxiv. (1884) 114.—Ins. Cyprus.
decipiens, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 74.—N. Granat.
deflexa, *Ruiz & Pav. ex DC. Prod.* i. 240.—Peruv.
dendroides, *Schult. f. Syst.* vii. 38.—Bras.
denticulata, *Schult. f. l. c.* 36.—Afr. trop.
diandra, *Burch. Trav.* i. 548.—Afr. austr.
dichotoma, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 318.—Bras.
didymana, *Hochst. ex Oliver, Fl. Trop. Afr.* i. 79.—Abyssin.
diffusa, *Banks, ex DC. Prod.* i. 241.—Bras.
diffusa, Koxb. Hort. Beng. [96]; Fl. Ind. iii. 129 = aspera.
digitata, *Forsk. Fl. Aegypt. Arab.* 120.—Arabia.
Dilleniana, *DC. Prod.* i. 240 = ornithopodioides.
diversifolia, Hochst. & Steud. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 5 = brachycarpa.
dodecandra, Banks, ex Wight & Arn. Prod. 23, partim = Burmanni.
dodecandra, Linn. Sp. Pl. ed. II. 939 = graveolens.
dodecaphylla, *Vell. Fl. Flum.* 271; vi. t. 110.—Bras.
drocrifolia, *Delile, Fl. Egypt.* 250.—Aegypt.; Arab.
Eclkloniana, Schrad. in Linnaea, x. (1836) Lit. 109 = Gynandropsis pentaphylla.
edulis, *Rafin. Fl. Ludov.* 86.—Am. bor.
Ehrenbergiana, *Schweinf. Beitr. Fl. Aethiop.* i. 66.—Nubia.
epilobioides, Jan, Elench. fide Schult. f. Syst. vii. 47 = gravoleens.
Erucago, Mill. Gard. Dict. ed. VIII. n. 6 = heptaphylla, sphoeta.
fetisa, *Linn. f. Suppl.* 300.—Ind. or.
filifolia, Vahl, Symb. Bot. i. 48 = tenella.
fimbriata, *Vicary, in Ann. Nat. Hist.* Ser. II. i. (1848) 495.—Scinde.
flava, Banks, ex DC. Prod. i. 241 = gravoleens.
flavescens, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 396; ex Schult. f. Syst. vii. 37 = Gynandropsis pentaphylla
flexuosa, *Griseb. in Goett. Abh.* xix. (1874) 78.—Am. austr.
foliolosa, *DC. Prod.* i. 240.—Persia.
foliosa, *Hook. f. in Hook. Niger Fl.* 219.—Afr. trop.
fruticosa, *Ambl. Pl. Gui.* ii. 277.—Guiana.
frutescens, Linn. Sp. Pl. 671 = Cadaba indica.
fugax, *Schrad. Ind. Sem. Hort. Gotting.* (1830).—*Cf. Linnaea*, vi. (1831) *Lit.* 72.—Hab.?
gigantea, Nimmo, Fl. Filip. ed. II. 356 = Gynandropsis pentaphylla.
glaberrima, *Edgw. Mssot.* 400.—Am. austr.
glandulosa, *Ruiz & Pav. ex DC. Prod.* i. 238.—Peruv.

CLEOME:—
glauca, *DC. Prod.* i. 239.—Mesopotam.
glaucescens, *DC. l. c.*—Mesopotam. ; Persia.
gracilis, Edgew. in Journ. As. Soc. Beng. xvi. (1847) ii. 1213 = papillosa.
gracilis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 74.—N. Granat.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 462.—Madag.
grandiflora, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 71.—Afr. trop.
gravoleens, *Rafin. in Journ. Phys. (Août* 1819) 98.—Am. bor.
guianensis, *Aubl. Pl. Gui.* ii. 675.—Guiana.
guineensis, Hook. f. in Hook. Niger Fl. 216 = ciliata.
gynandra, Linn. Sp. Pl. 671 = pentaphylla.
heptaphylla, *Linn. f. c.* ed. II. 967.—Am. trop.
heptaphylla, G. F. W. Mey. Prim. Fl. Esseq. 222 = Houstoni.
heptaphylla, Sw. Obs. 252 = pungens.
herbacea, *Bunge & Bien. ex Boiss. Fl. Orient.* i. 412. —Afghanist.
heterotricha, Burch. Trav. i. 547 = Gynandropsis heterotricha.
hexandra, Poir. ex Steud. Nom. ed. I. 206 = arabica.
hirsuta, *Ruiz & Pav. ex DC. Prod.* i. 238.—Peruv.
hirta, *Oliver, Fl. Trop. Afr.* i. 81.—Afr. trop.
horrida, Mart. in Schult. f. Syst. vii. 32 = spinosa.
Houtmanni, Miq. in Linnaea, xxii. (1849) 470 = latifolia.
Houstoni, *R. Br. in Ait. Hort. Kew.* iv. 181.—Jamaica.
Houstoni, Miq. l. c. xix. (1847) 483 = diffusa.
Houstoniana, *Schlecht. in Linnaea*, xxiv. (1851) 689.—Am. austr.
Humboldtii, *DC. Prod.* i. 241.—Am. trop.
iberica, DC. l. c. 240 = ornithopodioides.
iberidella, *Welw. in Oliver, Fl. Trop. Afr.* i. 79.—Afr. trop.
icosandra, Linn. Sp. Pl. 672 = gravoleens.
indecora, *Hornem. Ind. Sem. Hort. Hafn.* (1846) ; *ex Walp. Ann.* i. 50.—Mexic.
integrifolia, *Torr. & Gray, Fl. N. Am.* i. 122.—Am. bor.
jamesa, Berg. Desc. Pl. Cap. 164 = Heliophila callosa.
junceo, Sparm. in Nov. Act. Soc. Sc. Upsal. iii. (1780) 192 = Schepperia juncea.
khorassanica, *Bunge & Bien. ex Boiss. Fl. Orient.* i. 413.—Persia.
Kotschyana, *Boiss. l. c.* 413.—Assyria.
laevigata, Soland. ex DC. Prod. i. 200 = Stanleya grandiflora.
lateralis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1865) 73.—N. Granat.
latifolia, *Vahl, ex DC. Prod.* i. 239.—Guiana.
laxa, *Thunb. Prod. Pl. Cap.* 109.—Afr. austr.
Lechleri, *Eichl. in Mart. Fl. Bras.* xiii. i. 250.—Peruv.
Leschenaultii, Schult. f. Syst. vii. 49 = Cheiidonii.
linearis, Stocks, ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 3 = papillosa.
loughlinii, *Prod. Rel. Haenk.* ii. 84.—Peruv.
longifolia, Vahl, ex DC. Prod. i. 240 = Wormskioldia heterophylla.
longifolia, Willd. ex Schult. f. Syst. vii. 1617 = speciosa.
longipes, *DC. Prod.* i. 239.—Ecuador.
lusitanica, Mill. Gard. Dict. ed. VIII. n. 3 = gravoleens.
lutea, *Hook. Fl. Bor. Am.* i. 70. t. 25.—Am. bor. occ.
lutea, *E. Mey. Exsi Pfl. Docum.* 173.—Afr. austr.
macrocarpa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 319.—Mexic.
macrophylla, Rezb. Hort. Beng. [96], nomen (quid?). —Ind. or.
macrostachya, *Sauv. Fl. Cub.* 4.—Cuba.
macrothyrsa, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 73.—N. Granat.
maderensis, Mart. ex Schult. f. Syst. vii. 48 = aculeata.
mollistoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 320.—Perlia.
mexicana, *D. Dietr. Syn. Pl.* ii. 1068.—Mexic.
mexicana, *Hemsl. Diag. Pl. Nov.* 20.—Mexic.
microantha, *Desv. in Hum. Prod. Fl. Ind. Occ.* 68.—Guiana.
microcarpa, Schrad. ex Schult. f. Syst. vii. 30 = Dactylaena microantha.

CLEOME:—
monandra, DC. Pl. Rar. Hort. Genèv. 54. t. 15 = Dactylaena microantha.
monophylla, Law, in J. Grah. Cat. Bomb. Pl. 7 = simplicifolia.
monophylla, *Linn. Sp. Pl.* 672.—Ind. or.; Afr. trop.
Mortiziana, *Klotzsch, ex Eichl. in Mart. Fl. Bras.* xiii. i. 250.—Venezuela.
muschata, Stocks, ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 4 = brachycarpa.
multicaulis, *Moç. & Sesse, ex DC. Prod.* i. 240.—Mexic.
mutisea, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) ii. 1212 = paradoxa.
muricata, *Schult. f. Syst.* vii. 35.—Ind. or.
nivalis, Vahl, ex DC. Syst. ii. 245 = Macropodium nivale.
Notana, Boiss. Diagn. Ser. II. i. 48 = quinquenervia.
nummularia, DC. Prod. i. 239 = rosea.
ofiicacyra, Balb. Hort. Lagd. Alob. (1826); ex Schult. f. Syst. vii. 51 = Gynandropsis muricata.
oleracea, Welw. Apont. 556 = pentaphylla.
ornithopodioides, Forsk. Fl. Aegypt. Arab. n. 402 = brevistiliosa.
ornithopodioides, *Linn. Sp. Pl.* 672.—As. Min. ; Persia.
oxalides, *F. Muell. Fragm.* i. 69.—Austral.
oxypetala, *Boiss. Diagn.* Ser. I. vi. 30.—Persia; Arab.
oxyphylla, *Burch. Trav.* ii. 230.—Afr. austr.
pallida, *Kotschy, in Sitzb. Acad. Wiem. Math. Nat.* liii. *Abth.* i. (1866) 302.—Arabia.
palmipes, *Spreng. Syst.* iii. 122.—Guiana.
paludosa, *Willd. ex Eichl. in Mart. Fl. Bras.* xiii. i. 256.—Bras.
palustris, *Salzm. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 317.—Bras.
papillosa, *Steud. Nom.* ed. II. i. 362.—Aegypt.; Amb. ; Ind. bor. occ.
paradoxa, *R. Br. in Salt, Abyss. App.* 65.—Abyssin.
parviflora, R. Br. in Salt, Abyss. App. 65 = brachycarpa.
parvifiora, H. B. & K. Nov. Gen. et Sp. v. 68 = psoraleoides.
pedunculata, Vell. Fl. Flum. 272; vi. t. 112 = gigantea.
pentaphylla, Blanco, Fl. Filip. ed. I. 252 = Gynandropsis speciosa.
pentaphylla, Hort. Van Houtte, ex Eichl. in Mart. Fl. Bras. xiii. i. 251 = Houtteana.
pentaphylla, Linn. Sp. Pl. ed. II. 938 = Gynandropsis pentaphylla.
pentaphylla, Vell. Fl. Flum. 272; vi. t. 111 = rosea.
pilosa, *Benth. Bot. Voy. Sulph.* 65.—Am. austr.
pinnata, Pursh, Fl. Am. Sept. ii. 739 = Stanleya pinnatifida.
platycarpa, *Torr. in Bot. U. St. Expl. Exped.* ii. 235. t. 2.—Am. bor. occ.
Poissoniana, *Pacquet, in Journ. Hort. Prat.* i. (1843) 349.—Hab.?
polygama, Linn. Sp. Pl. ed. II. 939; DC. Prod. i. 241 = serrata.
porphyrantha, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 71.—N. Granat.
procumbens, *Vacq. Enum. Pl. Caris.* 361; *Linn. Syst.* ed. XII. 443.—Ind. occ.
prostrata, Hort. ex DC. Prod. ii. 362 = Rothia trifoliata.
prostrata, Koen. ex Steud. Nom. ed. II. i. 362 = aspera.
pruinose, *T. Anders, in Journ. Linn. Soc. v. Suppl.* i. (1860) 3.—Arabia.
pruricea, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 68.—N. Granat.
psoraleoefolia, *DC. Prod.* i. 239.—Bras.
puberula, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 71.—N. Granat.
pubescens, Sieber, ex Steud. Nom. ed. II. i. 362 = Gynandropsis pentaphylla.
pubescens, *Sims, Bot. Mag.* t. 1857.—Panama.
pulchella, *Schult. f. Syst.* vii. 1617.—Bras.
pungens, Willd. Enum. Hort. Berol. 639 = spinosa.
purpurea, *DC. Prod.* i. 239.—Peruv.
quinquenervia, *DC. l. c.*—Persia ; Afghanist. ; Ind. bor. occ.
Raddeana, *Trautv. in Act. Hort. Petrop.* i. (1871) 19. —Turcoman.
Radula, Fenzl, in Flora, xxvii. (1844) 312 = papillosa.

CLEOME :—
ramosissima, *Parl. ex Webb, Progm. Fl. Aethiop.* 23. —Afr. trop.
rephaeoides, DC. Syst. ii. 662 = Wormskioldia heterophylla.
rocia, *Yacquem. Voy. Bot.* t0.—Ind. or.
Ragonlii, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 190.—Am. austr.
Richii, *A. Gray, in Bot. U. St. Exped.* i. 67.—Bras.
ridibula, R. Br. in Salt, Abyss. App. 65 = droseriifolia.
rosea, Eckl. ex Steud. Nom. ed. II. i. 582 = Gynandropsis pentaphylla.
rosea, *Poit. in Ker. Hortic. Ser.* II. i. (1841-48) 69.—Texas.
roseo, *Vahl, ex DC. Prod.* i. 239.—Bras.
rubella, *Burch. Trav.* i. 543.—Afr. austr.
rubiginosa, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xvii. (1862) 69.—N. Granat.
rupestris, *Sond. in Linnaea*, xxiii. (1850) 6.—Afr. austr.
rupicola, *Vicary, in Ann. Nat. Hist.* Ser. II. i. (1848) 455.—Scinde.
Rutea, Cambess. in Jacquem. Voy. Bot. 19. t. 19 = brachycarpa.
rutidosperma, *DC. Prod.* i. 241.—Ind. occ.?
sandwicensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 65. —Ins. Sandvic.
scabrella, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 188.—Am. trop.
scandens, Rudx, ex Eichl. in Mart. Fl. Bras. xiii. i. 251 = longipes.
scoposa, DC. Prod. i. 239 = papillosa.
Schraderi, Schult. f. Syst. vii. 45 = Cheiidonii.
secundiflora, *Schrad. in Schult. f. l. c.* 37.—Hab.?
Sellowia, *Klotzsch, ex Eichl. in Mart. Fl. Bras.* xiii. i. 253.—Bras.
semibetandra, *Send. in Linnaea*, xxiii. (1850) 5.—Afr. austr.
serrata, *Yacq. Ensum. Pl. Carib.* 26.—Ind. occ.
serrulata, Forsk, Fl. Bor. Am. ii. 441 = integrifolia.
sessiliflora, *Schrank. f. Syst.* vii. 28.—Ind. or. occ.
silicuifera, *Eichl. in Mart. Fl. Bras.* xiii. i. 260.—Bras.
siliquaria, R. Br. in Salt, Abyss. App. 65 = arabica.
simplicifolia, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 159.—Ind. or.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 501.—Ins. Socotra.
Sonora, *A. Gray, Pl. Wright.* ii. 16.—N. Mexic.
sparsiflia, *S. Wats. in Bot. King, Exp.* 32. t. 5.—Am. bor. occ.
speciosa, H. B. & K. Nov. Gen. et Sp. v. 84. t. 436 = Gynandropsis speciosa.
speciosa, *Rafin. Fl. Ludov.* 86.—Am. bor.
speciosissima, *Deppe, ex Lindl. Bot. Reg.* t. 1812.—Mexic.
spinosa, *Yacq. Ensum. Pl. Carib.* 26; *Linn. Sp. Pl.* ed. II. 939.—Am. austr.
spinosa, Sw. Obs. 289 = pungens.
stenophylla, Klotzsch, in Rich. Schomb. Reise, iii. 1104 = Physostemon lanceolatum.
Steveniana, Schult. f. Syst. vii. 40 = cretinifolium.
Stockiana, *Botan. Magaz. Ser. II. i.* 47.—Baluchistan.
strigosa, *Oliver, Fl. Trop. Afr.* i. 80.—Afr. trop.
stylosa, *Eichl. in Mart. Fl. Bras.* xiii. i. 250.—Venezuela.
subcordata, Steud. ex Oliver, in Trans. Linn. Soc. xxix. (1873) 58 = monophylla.
suchanezensis, *Miq. in Linnaea*, xviii. (1844) 580.—Guiana.
tenella, *Linn. f. Suppl.* 300.—Ind. or.; Afr. trop.
tenuifolia, *Le Conte, ex Torr. & Gray, Fl. N. Am.* i. 138.—Am. bor.
ternata, Willd. ex Eichl. in Mart. Fl. Bras. xiii. i. 257 = polygama.
tetrandra, *Benth, ex DC. Prod.* i. 240.—Austral.
tetraphylla, Willd. ex Eichl. in Mart. Fl. Bras. xiii. i. 257 = psoraleoides.
tinnaea, *Spag. in Anal. Soc. Cient. Argent.* xv. (1883) 97.—Reg. Argent.
trachycarpa, *Klotzsch, ex Eichl. in Mart. Fl. Bras.* xiii. i. 256.—Bras.
trianæi, *Pranco. in Mus. Seuchenb.* i. (1884) 177.—Angyst.; Arabia.
trifoliata, Descourt. Fl. Med. Ant. i. t. 44 = polygama.

CLEOME :—
tripfylla, James, in Long, Exped. ii. 289 = integrifolia.
tripfylla, Linn. Sp. Pl. ed. II. 938 = serrata.
tripfylla, Vell. Fl. Flum. 279; vi. t. 118 = affinis.
uniglandulosa, *Cav. Ic.* iv. 8.—Mexic.
usana, *Brot. ex Heynh. Nom.* ii. 149, nomen.—Hab.?
Vahliana, Presen. in Mus. Senckenb. ii. (1837) 110 = brachycarpa.
venusta, Fenzl, in Flora, xxvii. (1844) 812 = paradoxa.
villosa, *Garsn. in Hook. Lond. Journ. Bot.* i. (1842) 166.—Bras.
viminea, *Schrad. in Linnaea*, x. (1836) *Lit.* 109.—Hab.?
violacea, Linn. Sp. Pl. 672 = gravesiana.
virgata, Ster. in DC. Prod. i. 240 = ornithopodioides.
virgata, *Thumb. Prod. Pl. Cap.* 109.—Afr. austr.
viridiflora, Schreb. in Nov. Act. Nat. Cur. iv. (1770) 156. t. 3 = gigantea.
viscosa, *Linn. Sp. Pl.* 672.—Geront. trop.

CLEOMELLA, DC. Prod. i. 237 (1824). *CAPPARIDEAE*, Benth. & Hook. f. i. 105.
ISOPARA, Rafin. Acti. Journ. 144 (1832).
angustifolia, *Torr. in Hook. Kew Journ.* ii. (1850) 255. —Mexic.
brevipes, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 368.—Calif.
Caulfieri, Herv. ex A. Gray, Pl. Wright. i. 12, 139 = Wislizenia refracta.
Fremontii, *Torr. in Smithson, Contrib.* iii. (1852) 12, in note.—Am. bor.
longifora, *Torr. in Hook. Kew Journ.* ii. (1850) 255.—Mexic.
medicaginea, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 318.—Mexic.
mexicana, *Moç. & Sesse, ex DC. Prod.* i. 237.—Mexic.
Wislizenia, Torr. in Ann. Lyc. N. York, ii. (1828) 167 = angustifolia.
obtusifolia, *Torr. & Frém. in Frem. Second Report*, 211.—Am. bor. occ.
occarpa, *A. Gray, in Proc. Am. Acad.* xi. (1876) 72.—Am. bor. occ.
parviflora, *A. Gray, l. c.* vi. (1864-65) 520.—Calif.
plocasperma, *S. Wats. in Bot. King, Exp.* 33.—Am. bor. occ.

CLEOMENA, Roem. & Schult. Syst. ii. 18, 888 (1817) = Mushlenbergia, Schreb. (Gramin.).

CLIBONIA, Linn. Sp. Pl. ed. II. 587 (1762). *LABIATAE*, Benth. & Hook. f. ii. 1208.
Isatisaica, *Linn. l. c.*—Europ. austr.

CLEOPHORA, Gaertn. Fruct. ii. 285. t. 120 (1791) = Lasiandra, Comm. (Palm.).
dendriformis, Lodd. ex Baker, Fl. Maurit. 281 = L. Loddigesii.
lonicerodes, Gaertn. Fruct. ii. 185. t. 120 = L. Commersonii.

CLERCIA, Vell. Fl. Flum. 29; i. t. 78, 74 (1825) = Salacia, Linn. (Celastr.).
cinnicnofolia, Miers, in Trans. Linn. Soc. xxvii. (1872) 381 = S. spivestris.
dispersa, Miers, l. c. 379.—Bras.
guianensis, Miers, l. c. 378.—Guiana.
lanceolata, Miers, l. c. 377.—Bras.
microcarpa, Miers, l. c. 379 = S. Miersii.
ovata, Vell. Fl. Flum. 30; i. t. 78 = S. arborea.
Passiflora, Vell. l. c. t. 74.—Bras.
retusa, Miers, in Trans. Linn. Soc. xxvii. (1872) 380 = S. spivestris.
Tweediana, Miers, l. c.—Bras.

CLERKIA, Neck. Elem. i. 251 (1790) = Tabernaemontana, Plum. (Apocyn.).

CLERMONTIA, Gaudich. in Freyc. Voy. Bot. 459. t. 71 ad 78 (1826). *CAMPANULACEAE*, Benth. & Hook. f. ii. 546.
grandiflora, *Gaudich. l. c.* t. 78.—Ins. Sandvic.
Kakoana, *Meyen, Reise*, ii. 139.—Ins. Sandvic.
macrocarpa, Gaudich. Bot. Voy. Bonite, t. 49 = Kakoana.
macrophylla, Nutt. Trans. Am. Phil. Soc. N. S. viii. (1842) 251 = Kakoana.
oblongifolia, *Gaudich. in Freyc. Voy. Bot.* 459. t. 71. —Ins. Sandvic.

CLERMONTIA :—
parviflora, *Gaudich. ex A. Gray, in Proc. Am. Acad.* v. (1861) 150.—Ins. Sandvic.
paroalcalis, *Gaudich. in Freyc. Voy. Bot.* 459. t. 72.—Ins. Sandvic.

CLERODENDRON, Linn. Gen. ed. I. 186 (1737) *VERBENACEAE*, Benth. & Hook. f. ii. 1156.
AGRICOLAEA, Schrank, in Denkschr. Akad. Munch. (1806) 98.
CORMACCHINIA, Savi, in Mem. Soc. Ital. Mod. xxi. (1837) 184. t. 7.
CYCLONEMA, Hochst. in Flora, xxv. (1842) 235.
CYRTOSTEMMA, Kunze, in Bot. Zeit. i. (1843) 272.
DOUGLASIA, Adans. Fam. ii. 300 (1763).
EGENA, Rafin. Fl. Tellur. ii. 85 (1836).
OVIEDA, Linn. Gen. ed. I. 79 (1737).
ROTHEEA, Rafin. Fl. Tellur. iv. 69 (1836).
SIPHONANTHUS, Linn. Gen. ed. II. 236 (1742).
SIPHONEMA, Hochst. in Flora, xxv. (1842) 236.
TORREYA, Spreng. Neue Entdeck. ii. 231 (1821).
VALDIA, Flum. ex Adans. Fam. ii. 157 (1763).
VOLKAMERIA, Linn. Syst. ed. I (1735); Gen. ed. I. 847 (1737).
VOLKAMANIA, Jacq. Hort. Schoenbr. iii. 48. t. 338 (1798).
Acerbiana, *Benth. & Hook. f. Gen.* ii. 1156.—Nubia.
aculeatum, *Griseb. Fl. Brit. W. Ind.* 500.—Jamaic.
acuminatum, Wall. Cat. n. 1792 = disparifolium.
affine, *Griff. Notul. iv.* 170.—Burma.
amicorum, *Seem. in Bonplandia*, x. (1862) 249.—Ins. Fasif.
amplius, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 238 = cyrtophyllum.
angustifolium, Salisb. Prod. 108 = fortunatum.
angustifolium, *Spreng. Syst.* ii. 758.—Jamaic.
atlernatum, R. Br. Prod. 511 = floribundum.
aurantium, *G. Don, in Edinb. Phil. Journ.* xi. (Oct. 1824) 349.—Afr. trop.
Arthur-Gordoni, Horne, ex Baker, in Journ. Linn. Soc. xx. (1883) 370 = Gordoni.
Balfouri, Hort. ex Flor. Mag. viii. (1869) sub t. 432 = Thomsonae.
Bethenecanum, *Low, Sarawak*, 378; *et in Bot. Mag.* t. 6485.—Borneo.
Blancoanum, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 161 = Nauseosum.
Blancoi, *Nowes, ex Villar, l. c.*—Ins. Philipp.
Blumeanum, *Schau. in DC. Prod.* xi. 669.—Java.
brachyanthum, *Schau. l. c.* 666.—Ins. Philipp.
bracteatum, *Wall. Cat.* n. 1800.—Reg. Himal.
brunsvigioides, *Baker, in Journ. Linn. Soc.* xxi. (1885) 455.—Madag.
Buchananii, *Herb. Roxb. ex Wall. Cat.* 2658 (sp. dub.). —Ind. or.
Bungei, Steud. Nom. ed. II. i. 382 = foetidum.
burmanum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 252. —Malaya.
buzifolium, *Spreng. Syst.* ii. 758.—Hab.?
calamitrotum, Hort. Belg. ex Lem. Jard. Fleur. iv. (1854) Misc. 48 = nutans.
calamitosum, *Linn. Mant.* i. 80.—Malaya.
calycinum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 232 = infortunatum.
canescens, *Wall. Cat.* n. 1806.—China.
capense, *D. Don, ex Steud. Nom.* ed. II. i. 382.—Afr. austr.
capreae, Eckl. & Zeyh. ex Schau. in DC. Prod. xi. 661 = glabrum.
capitatum, Hook. Bot. Mag. t. 4355 = Whitfieldii.
capitatum, *Schnee & Thou. in Dan. Vid. Selsk. Afh.* iv. (1826) 21.—Afr. trop.
capsulare, Blanco, Fl. Filip. ed. I. 509 = inerme.
cardiophyllum, F. Muell. Fragm. iii. 144 = floribundum.
castaneaefolium, Kintzsch, in Bot. Ergeb. Waldem. Reise, 109 = infortunatum.
castaneifolium, Hook. & Arn. Bot. Beech. Voy. 205 = fortunatum.
cauliflorum, *Vathe, in Linnaea*, xliii. (1880-82) 538. —Ins. Nossibé.
cereum, *Wall. ex Steud. Nom.* ed. II. i. 382.—Ind. or.
coccineum, D. Dietr. Syn. Fl. iii. 216 = squamatum.
Colebrookianum, *Walp. Rep.* iv. 114.—Ind. or.; Malaya.
Commersonii, *Spreng. Syst.* ii. 736.—Ins. Philipp.
condensatum, *Miq. Fl. Ind. Bat.* ii. 976.—Malaya.

CLERODENDRON:—
cordatum, D. Don, Prod. Fl. Nep. 103 = infortunatum.
cordifolium, *A. Rich. Tent. Fl. Abyss.* ii. 170.—Abyssin.
coriaceum, R. Br. Prod. 511 = floribundum.
coriaceum, Poir. Encyc. Suppl. iv. 353 = javanicum.
coromandelianum, Spreng. Syst. ii. 758 = ovatum.
costatum, *R. Br. Prod.* 511.—Austral.
crenatum, *Lindl. in Gard. Chron.* (1860) 456.—Aa. trop.
cubense, *Schau. in DC. Prod.* xi. 668.—Cuba.
Cumingianum, *Schau. l. c.* 666.—Ins. Philipp.
cuneatum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 221 = serratum.
Cunninghamii, *Benth. Fl. Austral.* v. 54.—Austral.
cupidatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 221.—N. Granat.
cyrtophyllum, *Turcz. l. c.* 229.—China.
deflexum, *Wall. Pl. As. Rar.* iii. 10. t. 215.—Malaya.
densiflorum, *Grisf. Notul.* iv. 170.—Burma.
dentatum, Wall. Cat. n. 1799 = squamatum.
depauperatum, Wall. ex Steud. Nom. ed. II. i. 882 = infortunatum.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 568.—Malaya.
discolor, *Vatke, in Linnaea*, xliii. (1880–82) 536.—Afr. trop.
disparifolium, *Blume, Bijdr.* 809.—Malaya.
divaricatum, Jack, in Malay. Misc. i. (1820) n. v. p. 43 = serratum.
divaricatum, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 154 = Caryopteris divaricata.
diversifolium, Vahl, Symb. Bot. ii. 75 = emirnense!
elegans, *Manetti, ex Lem. Jard. Fleur.* iv. (1854) *Misc.* 47.—Hab.!
ellipticum, Zipp. ex Span. in Linnaea, xv. (1841) 329 = laevifolium.
emirnense, *Boj. ex Hook. Bot. Mag.* t. 2925.—Madag.
ericolphon, *Schau. in DC. Prod.* xi. 662.—Java.
fallax, *Lindl. Bot. Reg.* (1844) sub t. 19.—Java.
farinosum, Wall. Cat. n. 1810 = serratum.
ferruginosum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 221 = villosum.
fœtulosum, *Exc. Malacca*, ii. (1884) 48.—Malaya.
floribundum, *R. Br. Prod.* 511.—Austral.
floribundum, Hort. Angl. ex Walp. Rep. iv. 111 = emirnense.
fœtidum, *Bunge, Enum. Pl. Chin. Bor.* 52.—China.
fœtidum, D. Don, Prod. Fl. Nep. 103 = Caryopteris grata.
foetidum, Hort. Par. ex Planch. in Fl. des Serres, Sér. I. ix. (1853–54) 17 = fragrans, *Vent.*
fortunatum, Bach.-Ham. ex Wall. Cat. n. 3652 (sp. dub.)—Assam.
fortunatum, *Linn. Cent. Pl.* ii. 28.—China.
fragrans, *Vent. Jard. Malm.* t. 70.—China; Sumatra; Java.
fragrans, *Willd. Enum. Hort. Berol.* 659.—Am. trop.
galeatum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 91.—Ins. Socotra.
glabrum, *E. Mey. Comm. Pl. Afr. Austr.* i. 278.—Afr. austr.
glandulosum, Colebr. ex Wall. Cat. n. 1806 = Colebrookianum.
glandulosum, Lindl. Bot. Reg. (1844) sub t. 10 = prænc.!
glaucum, *Wall. ex Steud. Nom.* ed. II. i. 883.—Ind. or.
Gordoni, *Baker, in Journ. Linn. Soc.* xx. (1883) 370.—Ins. Fiji.
grandiflorum, *Schau. in DC. Prod.* xi. 659.—Hab.!
grandifolium, Salisb. Prod. 106 = serrata.
grandiose, Kurz, For. Fl. Brit. Burma, ii. 366 = Caryopteris paniculata.
grandiose, Wall. Cat. n. 1813 = Caryopteris grata.
Griffithianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 596.—Assam; Burma.
hamiltoniense, Hance, in Walp. Ann. iii. 386 = canescens.
Hanceanum, *Schauinsl. Bot. Fl. Asthiop.* 119.—Afr. trop.
hastatum, *Lindl. Bot. Reg.* t. 1207.—Reg. Himal.
heterophyllum, Wall. ex Steud. Nom. ed. II. i. 886, sphalm. = Caryopteris Wallichiana.

CLERODENDRON:—
hemidermo, *F. Muell. ex Benth. Fl. Austral.* v. 61.—Austral.
herbaceum, Wall. Cat. n. 1815 = serratum.
heterophyllum, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 64.—Ins. Mascar.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 536.—Afr. trop.
hircinum, *Schau. in DC. Prod.* xi. 661.—Madag.
hirsutum, G. Don, in Edinb. Phil. Journ. xi. (1824) 349 = umbellatum.
Horsfieldii, *Miq. Fl. Ind. Bat.* ii. 880.—Java.
Huegelii, *Hort. ex Regel, Gartenfl.* (1861) 51.—Austral.
illustre, *N. E. Br. in Gard. Chron.* (1884) ii. 424.—Ins. Celebes.
imperiale, *Carr. in Rev. Hortic.* (1874) 110.—Hab.!
inclusum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 257.—Afr. trop.
inermum, R. Br. Prod. 511 = nerifolium.
inerme, *Guerin. Pruct.* i. 271.—Ind. or.
infortunatum, *Guerin. l. c.*—Ind. or.; Malaya.
infortunatum, Dennst. Schlüssel. Hort. Malab. 27 = villosum.
intermedium, *Cham. in Linnaea*, vii. (1832) 150.—Ins. Luzon.
involucratum, *Vatke, in Linnaea*, xliii. (1880–82) 537.—Afr. trop.
ixoraefolium, *Hassk. in Retzia*, 60.—Singapore.
Jacksonianum, Wall. in Hook. Bot. Misc. i. (1830) 284; Cat. n. 1794 = disparifolium.
japonicum, *Sweet, Hort. Brit.* ed. I. 393.—Japon.
javanicum, *Spreng. Syst.* ii. 759.—Java.
jenense, Walp. Rep. iv. 113 = serratum.
Kaempferi, Fisch. ex Morr. Ann. Soc. Gand, i. (1845) 17, in syn. = squamatum.
laciniatum, *Balf. f. in Journ. Linn. Soc.* xvii. 1877) 19.—Ins. Mascar.
laevifolium, *Blume, Bijdr.* 808.—Malaya.
lanceolatum, *F. Muell. Fragm.* iii. 145.—Austral.
lasiogeneum, *Blume, Bijdr.* 810.—Ins. Molucc.
lasiocephalum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 594.—Reg. Himal.
lasiophyllum, *Baker, in Journ. Linn. Soc.* xx. (1883) 229.—Ins. Fiji.
Lehmtel, *Horne, ex Baker, l. c.* 369.—Ins. Fiji.
leucophloeum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 91.—Ins. Socotra.
Leucosceptrum, D. Don, Prod. Fl. Nep. 103 = Leucosceptrum canum.
ligustrifolium, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 64.—Mexic.
Lindemuthianum, *Vatke, in Linnaea*, xliii. (1880–82) 537.—Afr. trop.
Lindenianum, *A. Rich. Fl. Cub. Fanerog.* ii. 147.—Cuba.
Lindleyi, Decne. ex Planch. in Fl. des Serres, Sér. I. ix. (1853–54) 17 = fragrans, *Vent.*
Linnæanum, F. Muell. Fragm. vi. 151 = hemidermo.
Linnæi, Thw. Enum. Pl. Zeyl. 243 = Glossocarya Linnæi.
lividum, Lindl. Bot. Reg. t. 945 = fortunatum.
Lobbii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 590.—Ins. Penang.
longiflorum, *G. F. W. Mey. Prim. Fl. Esseq.* 217.—Guiana.
longiflorum, *Decne. in Nov. Ann. Mus. Par.* iii. (1834) 400.—Ins. Timor.
macrobotrium, *Miq. Fl. Ind. Bat.* ii. 874.—Java.
macrocalycinum, *Baker, in Journ. Linn. Soc.* xvii. (1881) 276.—Madag.
macrophyllum, *Blume, Bijdr.* 808.—Java.
macrophyllum, Sims, Bot. Mag. t. 2536 = serratum.
macrolophon, *Hook. f. Bot. Mag.* t. 6595.—Afr. trop.
macrostachyum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 220.—Reg. Himal.; Burma.
macrostegium, *Schau. in DC. Prod.* xi. 666.—Ins. Philipp.
Madagascariense, *Voigt, Hort. Suburb. Calc.* 467.—Ind. or.!
magnolizefolium, *Baker, in Journ. Bot.* xx. (1882) 343.—Madag.
Manetti, *Vis. Sem. Hort. Patav.* (1846–49) n. 2.—Hab.!
medium, R. Br. Prod. 510 = floribundum.
Minahassæ, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 409.—Ins. Celebes.
mite, *Vatke, in Linnaea*, xliii. (1880–82) 537.—Afr. trop.
molle, *R. Br. & K. Nov. Gen. et Sp.* ii. 244.—Ecuador.

CLERODENDRON:—
molle, Jack, in Malay. Misc. i. (1820) n. i. 15 = villosum.
mossambicense, *Klotzsch, in Peters, Reise Mossamb. Bot.* 256.—Afr. trop.
multiflorum, G. Don, in Edinb. Phil. Journ. xi. (1824) 350 = volubile!
myricoides, *R. Br. in Salt, Abyss. App.* p. lxv, nomen.—Abyssin.
Navesianum, *Vidal, Cat. Prov. Manila*, n. 398.—Ins. Philipp.
nerelfolium, *Wall. Cat.* n. 1789.—Malaya.
Neumayeri, *Vatke, in Linnaea*, xliii. (1880–82) 535.—Afr. trop.
nutans, Jack, in Malay. Misc. i. (1820) n. i. 17 = Jackianum.
nutans, *Wall. Cat.* n. 1793.—Reg. Himal.; Burma.
obovatum, *Walp. Rep.* iv. 113.—Ins. Molucc.
obtusidens, *Miq. Fl. Ind. Bat.* ii. 870.—Malaya.
odoratum, D. Don, Prod. Fl. Nep. 102 = Caryopteris Wallichiana.
ornatum, Wall. Cat. n. 1811 = serratum.
ovale, *Klotzsch, in Peters, Reise Mossamb. Bot.* 257.—Afr. trop.
ovalifolium, *A. Gray, in Proc. Am. Acad.* vi. (1862) 50.—Ins. Fiji.
ovatum, Poir. Encyc. Suppl. iv. 352 = coromandelinum.
ovatum, R. Br. Prod. 511 = floribundum.
oxyscpalum, *Miq. in Journ. Bot. Ntcrl.* i. (1851) 114.—China.
paniculatum, *Linn. Mant.* i. 90.—Burma; Malaya; China.
papanaun, *Schef. in Ann. Jard. Buitenz.* i. (1876) 41.—N. Guin.
pauciflorum, *Wall. Cat.* n. 1795.—Burma.
pentagonum, Hance, in Walp. Ann. iii. 388 = fortunatum.
petunioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 230.—Madag.
philippinum, *Schau. in DC. Prod.* xi. 667.—Malaya.
phlomoides, Hort. Ital. ex DC. l. c. v. 618 = Montanoa arborescens.
phlomoides, *Linn. f. Suppl.* 292.—Ind. or.
phyllomega, Stend. Nom. ed. II. i. 883 = macrophyllum
pubescens, *Lindl. in Bot. Reg.* t. 1205.—Ind. or.
pubescens, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 380, lapsu = canescens.
pulchrum, *Forsc. in O. Forbes, Nat. Wand. East. Archip.* 514.—Ins. Timor.
pumilum, *Spreng. Syst.* ii. 759.—China.
putre, *Schau. in DC. Prod.* xi. 661.—Madag.
pyramidale, Andr. Bot. Rep. t. 638 = paniculatum.
pyrifolium, *Baker, in Journ. Linn. Soc.* xx. (1883) 238.—Madag.
ramosissimum, *Baker, l. c.*—Madag.
Riedelii, *Oliver, in Journ. Linn. Soc.* xv. (1876) 100.—Ins. Celebes.
robustum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 259.—Afr. trop.
roseum, *Poit. in Rev. Hortic.* (1876) 80.—Hab.!
rotundifolium, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 128.—Afr. trop.
rubellum, *Baker, in Journ. Linn. Soc.* xx. (1883) 229.—Madag.
Rumphianum, *De Vriese, in Mig. Ann. Mus. Bot. Lugd. Bat.* iii. 252.—Ins. Amboina.
sagittatum, Wall. Cat. n. 1786 = hastatum.
Sagraei, *Schau. in DC. Prod.* xi. 659.—Ins. Cuba.
scandens, Beauv. Fl. Owar. i. 52. t. 28 = umbellatum.
scopiferum, *Miq. Fl. Ind. Bat.* ii. 881.—Malaya.
semiserratum, *Wall. Cat.* n. 1785.—Burma.
sericeum, Wall. l. c. n. 1814 = Hipteage sericea.
serotinum, Carr. in Rev. Hortic. (1867) 351 = trichotomum.
serratum, *Spreng. Syst.* ii. 758.—Ind. or.; Burma.
simplex, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 349.—Afr. trop.
singalense, *Miq. Fl. Ind. Bat. Suppl.* 568.—Sumatra.
sinuatum, *Hook. Bot. Mag.* t. 2359.—Afr. trop.
Siphonanthus, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 65.—Ind. or.; Malaya.
speciosissimum, Hort. Angl. ex Schau. in DC. Prod. xi. 673 = glandulosum.
speciosissimum, Fenz. Mag. Bot. iii. (1837) 217 et 271 = squamatum.
speciosum ×, Teijsm. & Binn. Cat. Hort. Bog. 386; et Lam. Illustr. Hortic. xvi. (1869) t. 592.
spicatum, *Thunb. Fl. Yaponn.* 22.—Java.
sphæcum, *Spreng. Syst.* ii. 760.—Ins. Dominic.

CLERODENDRON :—
splendens, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 349.—Ab. trop.
splendidum, Wall. Cat. n. 1803 = paniculatum.
squamatum, *Vahl, Symb. Bot.* ii. 74.—Reg. Himal.; Burma; China.
stenanthum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 258.—Afr. trop.
tenuifolium, *Baker, in Journ. Linn. Soc.* xx. (1883) 259.—Madag.
ternifolium, D. Don, Prod. Fl. Nep. 108 = serratum.
ternifolium, *H. B. & K. Nov. Gen. et Sp.* ii. 244.—Fl. Orinoco.
Thomasiae, *Balf. in Edinb. New Phil. Journ.* N. S. xv. (Jan.–Apr. 1862) 333.—Afr. trop.
tomentosum, *R. Br. Prod.* 510.—Austral.
Tracyanum, *F. Muell. ex Benth. Fl. Austri.* v. 62.—Austral.
trichotomum, *Thunb. Fl. Jap.* 256.—Japon.
trichotomum, Wall. Cat. n. 6815 B = villosum.
trifolium, *Vis. Illustr. Plants Nov. Ort Padov.* 19.—Nubia.
trifoliatum, *Steud. Nom.* ed. II. i. 883.—Ind. or.
tuberculatum, *A. Rich. Fl. Cub. Fanerog.* ii. 147.—Cuba.
umbellatum, *Pair. Encyc.* v. 166.—Afr. trop.
urticifolium, Wall. Cat. n. 1801 = squamatum.
velutinum, Wall. l. c. n. 1797 = villosum.
venosum, *Wall. l. c.* n. 1807.—Reg. Himal.
verrucosum, *Spitig. ex De Vriese, in Nederl. Kruidk. Arch.* i. (1848) 851.—Guiana.
verticillatum, D. Don, Prod. Fl. Nep. 102 = C. Siphonanthus.
vestitum, Wall. ex Steud. Nom. ed. II. i. 888 = infortunatum.
villosum, *Blume, Bijdr.*—Malaya.
viscosum, Vent. Jard. Malm. t. 25 = infortunatum.
volubile, *Beauv. Fl. Owar.* i. 92. t. 89.—Afr. trop.
Whitfieldii, *Sweet. in Bonplandia*, x. (1862) 250.—Afr. trop.

CLERODENDRUM, Linn. = prec.

CLETHRA, Bert. ex Steud. Nom. ed. II. i. 884 (1840) = Vivienia, *Cav.*
mollis, Bert. ex Steud. l. c. = V. parvifolia.

CLETHRA, Gronov. ex Linn. Syst. ed. I (1735).
ERICACEAE, Benth. & Hook. f. ii. 603.
Crossopheya, Klotzsch, in Linnaea, xxiv. (1851) 8.
Cuellaria, Ruiz & Pav. Prod. 59. t. 10 (1794); Syst. Veg. 108 (1798).
Guliena, Adans. Fam. ii. 166 (1763).
Junia, Adans. l. c. 165 (1763).
Kowalewskia, Turcz. in Bull. Soc. Nat. Mosc. (1859) i. 268.
Tinta, Linn. Syst. ed. X. 1010 (1759).
Volkameria, F. Br. Hist. Jamaic. 214. t. 21. f. 1 (1756).
acuminata, *Micks. Fl. Ber. Am.* i. 260.—Am. bor.
Alexandri, *Griseb. Fl. Brit. W. Ind.* 142.—Ind. occ.
alnifolia, Blanco, Fl. Filip. ed. II. 259 = canescens.
alnifolia, *Linn. Sp. Pl.* 396.—Am. bor.
angustifolia, *DC. Prod.* vii. 589.—Chili.
arborea, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 78.—Ins. Madera.
arborea, Vent. Jard. Malm. t. 40 = secundiflora.
barbinervis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 138.—Japon.
bicolor, *H. B. & K. Nov. Gen. et Sp.* iii. 289.—N. Granat.
bracteata, *Griseb. Fl. Brit. W. Ind.* 141.—Ind. occ.
brasiliensis, *Cham. & Schlecht. in Linnaea*, viii. (1833) 510.—Bras.
brevifolia, *Benth. Pl. Hartw.* 148.—N. Granat.
bynsonioides, *Griseb. Fl. Wright. Cub.* 170.—Cuba.
canescens, *Herb. Retzius. ex Blume, Bijdr.* 868.—Java.
castaneaefolia, *Meisn. in Mart. Fl. Bras.* vii. 169.—Peruv.
comitifolia, *Brust, in Flora*, lvii. (1874) 212.—Venezuela.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 75.—Cuba.
deutzia, [Soland. in] Ait. Hort. Kew. ed. I. ii. 78 = alnifolia.
divaricata, *Loisel. in Duham. Arb. ed. Nov.* v. 134.—Am. austr.
Fabri, *Hance, in Journ. Bot.* xxi. (1883) 130.—China.
fagifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 289.—Am. austr.
ferruginea, *Link, Enum. Hort. Berol.* i. 296, nomen.—Hab. ?
CLETHRA :—
ferruginea, *Ruiz & Pav. Fl. Per.* iv. t. 380.—Peruv.
fimbriata, *H. B. & K. Nov. Gen. et Sp.* iii. 290. t. 264.—N. Granat.
Gardneri, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 338.—Bras.
glauca, Hort. ex Steud. Nom. ed. II. i. 883 = pubescens.
glauca, Pers. Syn. i. 483 = acuminata.
guianensis, *Klotzsch, ex Meisn. in Mart. Fl. Bras.* vii. 170.—Guiana.
incana, Pers. Syn. i. 483 = pubescens.
integrifolia, Moç. & Sesse, ex DC. Prod. vii. 590 = mexicana.
japonica, Thunb. ex Steud. Nom. ed. II. i. 883 = barbinervis.
javanica, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 339.—Java.
Kewaleskii, Turcz. l. c. 338 = mexicana.
laevigata, *Meisn. in Mart. Fl. Bras.* vii. 168.—Bras.
lanata, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 538.—Mexic.
lancifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 331.—Singapore.
macrophylla, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 539.—Mexic.
macrostachya, *Loisel. in Duham. Arb. ed. Nov.* v. 155.—Hab. ?
Matherugii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 332.—Peruv.
mexicana, *DC. Prod.* vii. 590.—Mexic.
Michauxii, *St. Court. in Mag. Hortic.* i. (1838) 249.—Am. bor.
micrantha, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 238.—N. Granat.
monticola, Fras. ex Loisel. in Duham. Arb. ed. Nov. v. 130 = acuminata.
nervosa, *Loisel. in Duham. l. c.* 138.—Peruv.
obovata, Hook. & Arn. Bot. Beech. Voy. 302 = quercifolia.
odontata, *Ruiz & Pav. Fl. Per.* iv. t. 381.—Peruv.
oxanensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 331.—N. Granat.
occidentalis, Asct. ex Steud. Nom. ed. II. i. 883 = tinifolia.
ovalifolia, Moç. & Sesse, ex DC. Prod. vii. 590 = mexicana.
ovalifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 336.—Reg. Andin.
paniculata, [Soland. in] Ait. Hort. Kew. ed. I. ii. 78 = alnifolia.
pedicellaris, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 334.—Peruv.
pubescens, Willd. Enum. Hort. Berol. 455 = alnifolia.
punctata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 335.—N. Granat.
quercifolia, *Schlecht. in Linnaea*, viii. (1833) 594; *Lindl. Bot. Reg.* (1842) t. 23.—Mexic.
repanda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 331.—Venezuela.
revoluta, *Ruiz & Pav. Fl. Per.* iv. t. 380.—Peruv.
scabra, Loisel. in Duham. Arb. ed. Nov. v. 132 = brasiliensis.
scabra, Pers. Syn. i. 483 = alnifolia.
secundiflora, *Decne. in Fl. des Serres*, Sér. II. xii. (1877) 171, in nota.—Hab. ?
serrulata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 338 = mexicana.
spicigera, *Meisn. in Mart. Fl. Bras.* vii. 168.—Bras.
Spruceï, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 332.—Peruv.
suaveolens, *Turcz. l. c.* 330.—Mexic.
tinifolia, Schlecht. in Linnaea, viii. (1833) 594 = quercifolia.
tinifolia, *Sw. Prod. Veg. Ind. Occ.* 76.—Jamaic.
tomentosa, Lam. Encyc. ii. 46 = alnifolia.
virgata, *Ruiz & Pav. Fl. Per.* iv. t. 381.—Peruv.

CLETHROPSIS, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 301 = Alnus, Tourn. (Capelli?).
nepalensis, Spach, l. c. 302 = A. nepalensis.
nitida, Spach, l. c. = A. nitida.

CLETHROSPERUM, Planch. in Hook. Ic. Pl. sub t. 767 (1848), sphalm. = Clethrospermum, Planch. (Asclep.).

CLEVERA, Adans. Fam. ii. 234 (1763) = Polypremum, Linn. (Loganiac.).

CLEVERA, DC. in Mém. Soc. Phys. Genèv. i. (1822) 419; Prod. i. 594. (1824). *TERNSTROMIACEAE*, Benth. & Hook. f. i. 253.
Mocoy, Adans. Fam. ii. 50 (1763).
Trittylium, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 547.
elegans, *Choisy, in Mém. Soc. Phys. Genève.* xiv. (1856) 119.—Ins. Guadelup.
grandiflora, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 284.—Reg. Himal.
grandiflora, Wall. Cat. n. 1461 = ochnacea.
integrifolia, *Choisy, in Mém. Soc. Phys. Genève.* xiv. (1855) 119.—Mexic.
japonica, Sieb. & Zucc. Fl. Jap. i. 158. t. 81 = ochnacea.
Lushia, G. Don, Gen. Syst. i. 506 = ochnacea.
Merteniana, *Sieb. & Zucc. Fl. Jap.* i. 154.—Japon.
mexicana, *Planch. ex Hemsl. Biol. Centr. Am. Bot.* i. 93.—Mexic.
ochnacea, *DC. in Mém. Soc. Phys. Genève.* i. (1822) 418.—Reg. Himal.
ochnoides, Wall. ex G. Don, Gen. Syst. i. 506 = ochnacea.
serrulata, *Choisy, in Mém. Soc. Phys. Genève.* xiv. (1855) 119.—Mexic.
siphilitica, *Choisy, l. c.*—Mexic.
theoides, *Choisy, l. c.*—Ind. occ.
Wallichiana, Steb. & Zucc. Fl. Jap. i. 154 = ochnacea.

CLEVERA, Thunb. Nov. Gen. 68 (1783) = Ternstroemia, Linn. f.
dubia, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 308 = T. japonica.
emarginata, Garth. in Calc. Journ. Nat. Hist. vii. (1847) 447 = T. emarginata.
fragrans, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 307 = T. japonica.
gymnanthera, Wight & Arn. Prod. 87 = T. japonica.
japonica, Thunb. Nov. Gen. 69 = T. japonica.
laciopetala, Wight, Illustr. i. 99 = Adinandra lasiopetala.
Millettii, Hook. & Arn. Bot. Beech. Voy. 171 = Adinandra Millettii.
pentapetala, Spreng. Syst. ii. 596 = Saurauja tristyla.
rubiginosa, Spreng. l. c.—Sumatra.

CLEVRIA, Neck. Elem. ii. 188 (1790) = Dialium, Linn. (Leguminos.).

CLIANTHUS, Banks & Soland. ex G. Don, Syst. ii. 468 (1832). *LEGUMINOSAE*, Benth. & Hook. f. i. 503.
Donia, D. & G. Don, Gen. Syst. ii. 467 (1832).
Eremocharis, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 72 (1849).
Binnendykianus, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) 51.—Ins. Ceram.
cornutus, Lindl. Bot. Reg. (1841) Misc. 2 = Streblorrhiza speciosa.
Dampieri, *A. Cunn. ex Lindl. in Trans. Hort. Soc. Ser. I.* (1826) 522.—Austral.
Dampieri, R. Br. in Sturt, Exped. Centr. Austral. ii. 71 = Oxleyi.
Oxleyi, *A. Cunn. ex Lindl. in Trans. Hort. Soc. Ser. II.* i. (1835) 523.—Austral.
plicatus, *Endl. in Ann. Wien. Mus.* i. (1836) 185 = Streblorrhiza speciosa.
puniceus, *Banks & Soland. ex Lindl. in Trans. Hort. Soc. Ser. II.* i. (1835) 521.—N. Zel.
speciosus, Steud. Nom. ed. II. i. 381 = Streblorrhiza speciosa.

CLIBADIUM, Linn. Mant. ii. 161 (1771). *COMPOSITAE*, Benth. & Hook. f. ii. 345.
Baillieria, Arabl. Pl. Cel. ii. 994. t. 317 (1778).
Ormenia, DC. Prod. v. 104 (1836).
Oswalda, Cass. in Dict. Sc. Nat. liv. 323 (1828).
Thinia, Sw. Prod. Veg. Ind. Occ. 115 (1788).
acuminatum, *Benth. Bot. Voy. Sulph.* 114.—Ins. Cocos.
agreste, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 157 = Ichthyothera agreste.
Alexandri, *Griseb. Fl. Brit. W. Ind.* 368.—Jamaic.
angustifolium, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 154 = Sauvagesia.
Arouca, Sch. Bip. ex Baker, l. c. 155 = Scaudellaëa.
asperum, DC. Prod. v. 566 = quinquenerve.
Baderi, Griseb. Fl. Brit. W. Ind. 367 = quinquenerve.

CLIBADIUM :—
Barbasco, *DC. Prod.* v. 507.—Venezuela.
carrasanum, *DC. l. c.* 506.—Venezuela.
commelinoides, DC. l. c. = Lethyothera Canabi.
crosum, *DC. l. c.*—Ind. occ.
foetidum, Aiemand, ex Linn. Mant. ii. 294 = surinamense.
fragiferum, *Griseb. Fl. Brit. W. Ind.* 268.—Ind. occ. havanense, *DC. Prod.* v. 506.—Cuba.
leiocarpum, *Steetz, in Sem. Bot. Voy. Herald*, 152.—Panama.
neriifolium, *DC. Prod.* v. 507.—Venezuela.
pedunculosum, *DC. l. c.*—Guiana.
pervicinum, Poepp. ex DC. l. c. 506 = surinamense.
rotundifolium, *DC. l. c.*—Bras.
Schomburgkii, *Sch. Bip. in Schomb. Fauna & Fl. Guian.* noviss. 940.—Guiana.
surinamense, *Linn. Mant.* ii. 294.—Am. austr.
sylvestre, Baill. Hist. Pl. viii. 307 = surinamense.
trichinthinaeum, *DC. Prod.* v. 506.—Ind. occ.
Trinitatis, DC. l. c. 508 = surinamense.
Vargasi, *DC. l. c.* 506.—Venezuela.
villosum, Benth. Pl. Hartw. 205 = surinamense.

CLIDANTHERA, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 73 (1849) = Glycyrrhiza, Linn. (Legumin.).
paucifoliolata, R. Br. l. c. 74 = G. pacraloides.

CLIDEMIA, D. Don, in Mem. Wern. Soc. iv. (1823) 306. *MELASTOMACEAE*, Benth. & Hook. f. i. 766.
DANCERA, Rafin. Sylva Tellur. 93 (1838).
STAPHIDIUM, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 306.
STEPHANOTRICHUM, Naud. l. c. iv. (1846) 54.
acutiflora, Naud. l. c. xvii. (1852) 371 = Oxymeris acutiflora.
aggregata, D. Don, in Mem. Wern. Soc. iv. (1823) 309 = Henriettella aggregata.
agrestis, D. Don, l. c. = Oxymeris agrestis.
alata, G. Don, ex Lond. Hort. Brit. 174 = Miconia alata.
alpestris, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 103 = Oxymeris alpestris.
althaeoides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 368 = Octopleura confertiflora.
amblyandra, Naud. l. c. 358 = Oxymeris subseriata.
amygdaloides, DC. Prod. iii. 156 = Ossaea amygdaloides.
angustifolia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 358 = Oxymeris angustifolia.
aniophylla, DC. Prod. iii. 161 = Oxymeris aniophylla.
antostrichum, Schlecht. in Linnaea, xiii. (1839) 497 = Miconia Berghestana.
aphananthe, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 227.—Guiana.
aptera, DC. Prod. iii. 161 = Oxymeris aptera.
aristigera, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 366 = Oxymeris aristigera.
asperifolia, Naud. l. c. 343 = Oxymeris asperifolia, Ossaea asperifolia.
astrotricha, DC. Prod. iii. 161 = Miconia astrotricha.
atrata, Spreng. in Flora, xx. (1837) Beibl. ii. 78 = Miconia atrata.
aurea, Cham. in Linnaea, x. (1836) 47 = Oxymeris aurea.
auricoma, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 375 = Leandra aurea.
australis, Cham. in Linnaea, x. (1836) 44 = Oxymeris australis.
barbata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 186.—N. Granat.
barbinervis, Benth. Bot. Voy. Sulph. 96 = Miconia barbinervis.
Beathamiana, Miq. in Linnaea, xviii. (1844) 576 = hirta.
berbicensis, DC. Prod. iii. 169 = Miconia uniopoda.
Bertoni, Griseb. in Goett. Abh. ix. (1861) 47 = Sagraea plumosa.
bifaria, Rich. ex Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 374 = Oxymeris hexandrula.
bisglauca, DC. Prod. iii. 164 = Oxymeris dichotoma.
blepharoides, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 121 = Oxymeris Mitscherlichii.
blumana, *DC. Prod.* iii. 166.—Bras.
blepharoides, *DC. l. c.*—Bras.

CLIDEMIA :—
botryophora, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 350 = Oxymeris Nianga.
brachystachya, DC. Prod. iii. 156 = Ossaea brachystachya.
Brachenridgei, A. Gray, Bot. U. St. Expl. Exped. i. 591 = Oxymeris Brackenridgei.
bracteata, DC. Prod. iii. 162 = Miconia bracteata.
bullosa, *DC. l. c.* 158.—Bras.
calvescens, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 92 = Oxymeris calvescens.
camptoneura, Benth. in Hook. Journ. Bot. ii. (1840) 308 = Miconia campestris.
capillaris, D. Don, in Mem. Wern. Soc. iv. (1823) 309 = Octopleura leptopoa.
capillaris, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 184 = Sagraea capillaris.
capitata, Benth. in Hook. Journ. Bot. ii. (1840) 307 = Sagraea capitata.
capitellata, D. Don, in Mem. Wern. Soc. iv. (1823) 310 = dependens.
carassana, DC. Prod. iii. 162 = Oxymeris carassana.
carassana, DC. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 111 = Miconia spondylantha.
cephalophora, Steud. ex Triana, l. c. = Miconia bracteata.
cervanicarpa, DC. Prod. iii. 160 = Miconia spondylantha.
chantoni, *D. Don, in Lond. Hort. Brit.* 174.—Peruv.
chaetocalyx, A. Gray, Bot. U. S. Expl. Exped. i. 589. t. 74 = Oxymeris chaetocalyx.
Chantoni, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 360 = Oxymeris Chaetodon.
Chamissois, Naud. l. c. 375 = Miconia ibaguensis.
chinatiana, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 135.—Mexic.
chrysogogon, Benri. in Vet. Akad. Handl. Stockh. 1854 (1856) 127 = Miconia barbinervis.
ciliata, D. Don, in Mem. Wern. Soc. iv. (1824) 309 = Calophysa ciliata.
cinerea, Griseb. Cat. Pl. Cub. 97 = Oxymeris cinerea.
cinnamomifolia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 359 = Ossaea cinnamomifolia.
Claussenii, Naud. l. c. 357 = Oxymeris purpurascens.
coccinea, DC. Prod. iii. 162 = Miconia spondylantha.
cognata, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 308 = hirta.
confertiflora, DC. Prod. iii. 156 = Ossaea confertiflora.
congestiflora, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 344 = Oxymeris congestiflora.
conglomerata, *DC. Prod.* iii. 156.—Guiana.
cordifolia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 361 = Oxymeris cordifolia.
cordigera, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 94 = Oxymeris cordigera.
coriacea, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 356 = Ossaea coriacea.
cornoides, *Griseb. in Goett. Abh.* ix. (1861) 50.—Reg. Argent.
crenata, DC. Prod. iii. 157 = hirta.
crenata, D. Don, in Mem. Wern. Soc. iv. (1823) 308 = Oxymeris crenata.
crenicalpala, Griseb. Fl. Brit. W. Ind. 248 = Sagraea heteromera, crenopetala.
Cruegeriana, Griseb. l. c. 249 = Sagraea salicifolia.
cuneata, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 376 = Oxymeris cuneata.
cyanocarpa, Benth. Bot. Voy. Sulph. 94 = Octopleura quinquenervia.
dasytricha, A. Gray, Bot. U. S. Expl. Exped. i. 588 = Oxymeris carassana.
debilis, Crueg. in Linnaea, xx. (1847) 104 = Sagraea rubra.
debilis, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 361 = Oxymeris debilis.
decurrens, Benri. in Vet. Akad. Handl. Stockh. 1854 (1856) 127 = Octopleura quinquenervia.
deflexa, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 94 = Oxymeris deflexa.
deflexa, *Benth. in Proc. Linn. Soc. i.* (1841) 108.—Bras.
dendroides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 367 = Oxymeris dendroides.

CLIDEMIA :—
dentata, *D. Don, in Mem. Wern. Soc.* iv. (1823) 308.—Peruv.
depauperata, DC. Prod. iii. 161 = Oxymeris agrestis.
dependens, *D. Don, in Mem. Wern. Soc.* iv. (1823) 307.—Peruv.
Deppeana, *Steud. Nom.* ed. II. i. 884.—Mexic.
desmantha, Benth. in Hook. Journ. Bot. ii. (1840) 309 = Miconia rhytidophylla.
dichotoma, D. Don, in Mem. Wern. Soc. iv. (1823) 307 = Oxymeris dichotoma.
dispar, Gardn. in Hook. Lond. Journ. Bot. iii. (1843) 545 = Oxymeris dispar.
divaricata, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 378 = Oxymeris secundiflora.
diversifolia, DC. Prod. iii. 159 = Octopleura diversifolia.
dolichostachya, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 351 = Oxymeris dolichantha.
dolichostachya, Naud. l. c. 359 = Oxymeris dolichostachya.
Drosera, Sagot, l. c. Sér. VI. xv. (1883) 238 = aphananthe.
elegans, *DC. Prod.* iii. 157.—Am. trop.
elegans, D. Don, in Mem. Wern. Soc. iv. (1823) 309 = hirta.
Epibaterium, DC. Prod. iii. 167 = Octopleura Epibaterium.
erostrata, DC. l. c. 160 = Oxymeris erostrata.
erythropogon, *DC. l. c.* 157.—Jamaica.
euphorbioides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 370.—Ossaea euphorbioides.
fallax, Cham. in Linnaea, x. (1836) 41 = Oxymeris fallax.
fallax, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 344 = Oxymeris xanthoiasa.
fenestrata, Benth. Bot. Voy. Sulph. 94 = umbonata.
fenestrata, Seem. Bot. Voy. Herald, 154 = neglecta.
floribunda, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 349 = Oxymeris floribunda.
foveolata, DC. Prod. iii. 163 = Oxymeris foveolata.
fulva, *Benth. in Proc. Linn. Soc.* i. (1841) 108.—Bras.
glabrata, Steud. in Flora, xxvii. (1844) ii. 721 = Miconia articulata.
glabriflora, Presl, Bot. Bemerk. 70 = Miconia decussata.
glomerata, DC. Prod. iii. 168 = Oxymeris Nianga.
guadalupensis, Griseb. in Goett. Abh. vii. (1857) 218 = Sagraea guadalupensis.
hematostemon, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 348 = Oxymeris aurea.
herbacea, Steud. ex Cogn. in Mart. Fl. Bras. xiv. iii. 468 = Acletis herbacea.
heterobasis, DC. Prod. iii. 164 = Oxymeris heterobasis.
heterolcida, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 365 = japurensis.
heteromalla, D. Don, in Mem. Wern. Soc. iv. (1823) 310 = Sagraea rubra.
heterophylla, *Steud. in Flora*, xxvii. (1844) ii. 721.—Bras.
heterotricha, Griseb. Cat. Pl. Cub. 98 = Calycogonium heterotrichum.
hirsuta, *Griseb. in Mem. Am. Acad. N. S.* viii. (1861) 185.—Ind. occ.
hirsuta, Griseb. Fl. Brit. W. Ind. 248 = Ossaea hirsuta.
hirsuta, Macf. Fl. Jamaic. ii. 45 = Ossaea appendiculata.
hirta, *D. Don, in Mem. Wern. Soc.* iv. (1823) 309.—Jamaica.
hirtella, Griseb. in Goett. Abh. ix. (1861) 50 = Ossaea hirtella, Wrightii.
humilis, Miq. in Linnaea, xxii. (1849) 543 = Oxymeris Nianga.
holosericea, *D. Don, ex Loud. Hort. Brit.* 174.—Bras.
inaequalifolia; DC. Prod. iii. 164 = Oxymeris inaequalifolia.
inopogon, Mart. in Flora, xxiv. (1841) Beibl. ii. 87 = Oxymeris inopogon.
intermedia, DC. Prod. iii. 164 = Oxymeris intermedia.
interrupta, Poepp. ex Triana, in Trans. Linn. Soc. xxvii. (1871) 111 = Miconia bractosta.
hyselnerata, *DC. Prod.* iii. 168.—Guiana.
japurensis, *DC. l. c.* 159.—Bras.
lacera, DC. l. c. 164 = Miconia lacera.

CLIDEMIA :—

laevigata, O. Berg, ex Triana, in Trans. Linn. Soc. xxvii. (1871) 94 = Osneria laevigata.

Lamberiana, DC. Prod. iii. 164 = Clyneria crenata.

lanata, DC. l. c. 163 = Miconia lanata.

lanata, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 184 = Ossæa lanata.

lappacea, DC. Prod. iii. 160 = Miconia lappacea.

latifolia, DC. l. c. 159 = Sagræa latifolia.

latifolia, Griseb. Fl. Brit. W. Ind. 247 = Sagræa umbrosa.

laxiflora, *Schlecht. in Linnæa*, xxii. (1839) 426.—Mexic.

laxinervioides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 355 = Clyneria purpurascens.

leptocladoides, H. Crueg. in Linnæa, xx. (1847) 103 = hirta.

leptopus, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 147 = Ossæa leptopus.

leptostachya, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 172 = Ossæa brachystachya.

leucandra, Wright, ex Griseb. Cat. Pl. Cub. 96 = Sagræa domingensis.

Lima, DC. Prod. iii. 161 = Ossæa Lima.

linearis, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 356 = Clyneria purpurascens.

longiberbis, DC. Prod. iii. 168 = Oxymeris longiberbis.

lugubris, Benth. in Vet. Akad. Handl. Stockh. 1854 (1856) 127 = Oxymeris heterobasis.

lutescens, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 347 = Oxymeris crenata.

macrophylla, *Naud. l. c.* 375.—Ins. Cocos.

macrophylla, Sesm. Bot. Voy. Herald, 124 = Miconia simplex.

macropora, O. Berg, ex Triana, in Trans. Linn. Soc. xxvii. (1871) 94 = Oxymeris macropora.

maculata, Benth. in Hook. Journ. Bot. ii. (1840) 310 = Miconia auriculata.

marginata, DC. Prod. iii. 156 = Ossæa marginata.

Martiana, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 340 = Oxymeris cuneata.

Martii, Naud. l. c. 375 = Oxymeris ciliata.

melanoderma, Naud. l. c. 358 = Oxymeris melanodesma.

melanotricha, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 186.—Costa Rica.

Microcistrum, Naud. in Ann. Sc. Nat. Sér. III. xvii (1852) 359 = Oxymeris Microcistrum.

microcladoides, Benth. in Hook. Journ. Bot. ii. (1840) 310 = Miconia auriculata.

microcrantha, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 397.—Guiana.

microphylla, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 367 = Oxymeris micropetala.

microphylla, *D. Don, in Loud. Hort. Brit.* 174.—Ann. austr.

microphylla, Griseb. Fl. Brit. W. Ind. 248 = Ossæa microphylla.

microstachya, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 357 = Oxymeris purpurascens.

monticola, Naud. l. c. 341 = Miconia erythrantha.

multiplinervis, Naud. l. c. 358 = Oxymeris multiplinervis.

mutabilis, DC. Prod. iii. 162 = Miconia mutabilis.

naevula, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 187.—Guiana.

neglecta, *D. Don, in Mem. Wern. Soc.* iv. (1823) 307. —Peruv.

nervosa, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 341 = Oxymeris nervosa.

Nianga, DC. Prod. iii. 168 = Oxymeris Nianga.

Nianga, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 344 = Oxymeris xantholoata.

nivea, *D. Don, in Loud. Hort. Brit.* 174.—Am. austr.

novemnervia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 186.—Am. trop.

nunezensis, Steud. ex Triana, l. c. 92 = Oxymeris subsessilis.

obscura, DC. Prod. iii. 162 = Miconia obscura.

octona, *Griseb. in Goett. Abh.* ix. (1861) 50.—Reg. Argent.

oligocheata, Cham. in Linnæa, x. (1836) 45 = Oxymeris foveolata.

oocarpa, A. Gray, Bot. U. S. Expl. Exped. i. 590 = Oxymeris oocarpa.

ozanonioides, Naud. in Ann. Sc. Nat. Sér. III. xvii (1852) 360 = Oxymeris ozanonioides.

CLIDEMIA :—

oxyura, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 185.—Guiana.

paraguayensis, Steud. in Flora, xxvii. (1844) ii. 721 = Oxymeris paraguayensis.

parasitica, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxvii. (1871) 185.—Bras.

pauciflora, DC. Prod. iii. 157 = hirta.

pendulifiora, Naud. in Ann. Sc Nat. Sér. III. xvii. (1852) 362 = Oxymeris pendulifiora.

penninervis, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 185 = Sagræa penninervis.

pennipilis, O. Berg, ex Triana, in Trans. Linn. Soc. xxvii. (1871) 92 = Oxymeris pennipilis.

petiolaris, Triana, l. c. 125 = Deppeana.

petiolata, DC. Prod. iii. 157 = Sagræa petiolata.

phaeostephia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 342 = Oxymeris dichotoma.

phasotricha, Naud. l. c. 35 = Oxymeris australis.

Pickeringii, A. Gray, Bot. U. S. Expl. Exped. i. 591 = Oxymeris Pickeringii.

pilosa, D. Don, in Mem. Wern. Soc. iv. (1823) 308 = Calophyxe pilosa.

plumosa, DC. Prod. iii. 159 = Sagræa plumosa.

polyandra, Benth. Bot. Voy. Sulph. 95 = Miconia Benthamiana.

polystephia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 347 = Oxymeris polystachya.

pulchra, A. Gray, Bot. U. St. Expl. Exped. i. 591 = Oxymeris pulchra.

purpurascens, DC. Prod. iii. 161 = Oxymeris purpurascens.

purpurea, DC. l. c. 158 = dentata.

purpurea, *D. Don, in Mem. Wern. Soc.* iv. (1823) 308. —Peruv.

pustulata, DC. Prod. iii. 159 = umbonata.

quadrinodis, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 364 = Oxymeris secundiflora.

quinquedentata, Naud. l. c. 375 = Oxymeris quinquedentata.

quinquenodis, Naud. l. c. 370 = Oxymeris quinquenodis.

quintuplinervia, Steud. ex Triana, in Trans. Linn. Soc. xxvii. (1871) 187 = involucrata.

radulifolia, Benth. in Hook. Journ. Bot. ii. (1840) 309 = Miconia radulæfolia.

rariflora, *Benth. l. c.* 308.—Guiana.

reticulata, Benzl. in Vet. Akad. Handl. Stockh. 1854 (1856) 125 = petiolaris.

Regnellii, O. Berg, ex Triana, in Trans. Linn. Soc. xxvii. (1871) 91 = Oxymeris Regnellii.

Rengerri, Steud. in Flora, xxvii. (1844) 721 = Oxymeris inopogon.

repens, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 186.—N. Granad.

retropila, DC. Prod. iii. 156 = Ossæa retropila.

reversa, DC. l. c. 161 = Oxymeris reversa.

rhomboidea, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 378 = Oxymeris secundiflora.

rhodolaxia, Benzl. in Vet. Akad. Handl. Stockh. 1854 (1856) 126 = Miconia laxa.

rhodopogon, DC. Prod. iii. 163 = Oxymeris rhodopogon.

ribesifiora, Cham. in Linnæa, x. (1836) 38 = Oxymeris ribesiflora.

Riedeliana, O. Berg, ex Triana, in Trans. Linn. Soc. xxvii. (1871) 92 = Oxymeris Riedeliana.

rubra, *Mart. Nov. Gen. et Sp.* iii. 153. t. 281.—Am. austr.

rubrinervis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 183.—Cuba.

rufescens, Griseb. Cat. Pl. Cub. 97 = Ossæa rufescens.

sagittata, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 359 = Oxymeris subsessilis.

salicifolia, Naud. l. c. 364 = Ossæa brachystachya.

scabrosa, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 184.—Ind. occ.

secundiflora, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 346 = bipharochoides.

secunda, D. Don, in Mem. Wern. Soc. iv. (1823) 308 = Oxymeris secunda.

secundiflora, DC. Prod. iii. 164 = Oxymeris secundiflora.

serrata, D. Don, in Mem. Wern. Soc. iv. (1823) 309 = Sagræa rubra.

serrulata, *Schlecht. in Linnæa*, xiii. (1839) 495.—Mexic.

sibundoyia, C. Muell. in Walp. Ann. iv. 709 = Oxymeris sylvestris.

CLIDEMIA :—

simplicicaulis, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 346 = Oxymeris simplicicaulis.

solearis, Naud. l. c. 339 = Octoplaea macrophylla.

spanidlora, *Griseb. in Goett. Abh.* vii. (1857) 218.—Reg. Argent.

spicaeformis, DC. Prod. iii. 160 = Miconia spondylantha.

spicata, DC. l. c. 159 = dependens.

spicata, D. Don, in Mem. Wern. Soc. iv. (1823) 308 = Miconia spondylantha.

spicata, Griseb. Cat. Pl. Cub. 96 = neglecta, Heterotrichum octonum.

spondylantha, Mart. ex Triana, in Trans. Linn. Soc xxviii. (1871) 111 = Miconia spondylantha.

staphidioides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 344 = Miconia staphidioides.

stenopetala, DC. Prod. iii. 168 = Oxymeris dichotoma.

strigilifiora, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 372 = Oxymeris strigillosa.

strigillosa, *DC. Prod.* iii. 159.—Am. trop.

subsessilis, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 354 = Oxymeris subsessilis.

subspicata, Benzl. in Vet. Akad. Handl. Stockh. 1854 (1856) 126 = dependens.

suffruticosa, O. Berg, ex Triana, in Trans. Linn. Soc. xxvii. (1871) 185.—Bras.

sulcicaulis, Poepp. ex Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 343 = Oxymeris dichotoma.

sulphurea, Naud. l. c. 371 = Oxymeris velutina.

surinamensis, Miq. Comm. Phytog. ii. 82 = umbonata

Swartzii, Griseb. Fl. Brit. W. Ind. 248, pro parte = Sagræa pilosa.

sylvestris, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 356 = Oxymeris sylvestris.

tetraquetra, Cham. in Linnæa, x. (1836) 42 = Oxymeris tetraquetra.

tiliaefolia, *DC. Prod.* iii. 158.—Bras.

trichodes, DC. l. c. 161 = Miconia trichodes.

trichotoma, Wright, ex Griseb. Cat. Pl. Cub. 96 = Sagræa trichotoma.

trifora, Scheele, in Linnæa, xvii. (1843) 388 = Tibouchina viminea.

trinitensis, Griseb. Fl. Brit. W. Ind. 249 = Sagræa trinitensis.

tristis, D. Don, in Mem. Wern. Soc. iv. (1823) 307 = strigillosa.

umbonata, *DC. Prod.* iii. 158.—Bras.

urceolata, DC. l. c. = neglecta.

urticaefolia, Naud. in Ann. Sc. Nat. Sér. III. xvii (1852) 384 = Miconia Burchellana.

Valenzuelana, A. Rich. Ess. Fl. Cub. 559 = Heterotrichum octonum.

velutina, *D. Don, in Loud. Hort. Brit.* 174.—Ind: occ.

verticillata, DC. Prod. iii. 160 = Ossæa verticillata.

violacea, DC. l. c. 162 = Miconia spondylantha.

vittata, *Linden & André, Illustr. Hortic.* xxii. (1875) 152. t. 219.—Peruv.

Weddelliii, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 346 = Oxymeris crostrata.

Wrightii, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 185 = Sagræa Wrightii.

xanthocoma, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 353 = Oxymeris xanthocoma.

xantholoata, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 344 = Oxymeris cuneata.

xanthopogon, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 349 = Oxymeris xanthopogon.

CLIDEMIASTRUM, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 87 = Oxymeris, DC. (Melast.).

Klugianum, Naud. l. c. 88 = Leandra reversa.

mexicanum, Naud. l. c. 87 = O. heterobasis.

CLIFFORTIA, Linn. Gen. ed. I. 311 (1737). *ROSACEAE*, Benth. & Hook. f. i. 624.

MONOGRAPHIDIUM, Presl, Epim. Bot. 208 (1850).

MORLANDIA, Neck. Elem. ii. 99 (1790).

acerosa, Banks, ex Gaertn. Fruct. i. 265 = Nenax acerosa.

arachnoidea, Ludd. Bot. Cab. t. 900 = rucifolia.

bancana, *Horn. in Horn. & Sond. Fl. Cap.* ii. 308.—Ad. austr.

berberidifolia, Linn. Encyc. ii. 46 = ferruginea.

cinerea, Thunb. Prod. Fl. Cap. 93 (= *Ambrosia hirta*). —Ad. austr.

CLIFFORTIA:—
complanata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 593 = serpyllifolia.
concavifolia, *Eckl. & Zeyh. Enum.* 268.—Afr. austr.
cordifolia, Lam. Encyc. ii. 47 = illicifolia.
cordifolia, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 595. —Afr. austr.
crenata, *Linn. f. Suppl.* 430.—Afr. austr.
cuneata, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 413. Afr. austr.
dentata, Eckl. & Zeyh. Enum. 268 = trilobe.
dentata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 599 = gracilis.
dentata, *Willd. Sp. Pl.* iv. 845.— Afr. austr.
Dregeana, *Presl, Bot. Bemerk.* 141.—Afr. austr.
drepanoides, *Eckl. & Zeyh. Enum.* 269.—Afr. austr.
ericaefolia, *Linn. f. Suppl.* 430.—Afr. austr.
eriocephalina, *Cham. in Linnaea*, vi. (1831) 349.— Afr. austr.
falcata, *Linn. f. Suppl.* 431.—Afr. austr.
falcata, Spreng. ex Eckl. & Zeyh. Enum. 269 = drepanoides.
ferruginea, *Linn. f. Suppl.* 429.—Afr. austr.
filicaulis, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 599 = tricuspidata.
filicaulis, *Schlecht. in Linnaea*, ii. (1827) 33.—Afr. austr.
filifolia, *Linn. f. Suppl.* 430.—Afr. austr.
flabellifolia, Sond. ex Harv. & Sond. Fl. Cap. ii. 597 = Myrothamnus flabellifolia.
flexuosa, E. Mey. ex Harv. & Sond. l. c. ii. 596 = ferruginea.
gracilis, *Harv. l. c.* 599.—Afr. austr.
graminea, *Linn. f. Suppl.* 429.—Afr. austr.
grandifolia, *Eckl. & Zeyh. Enum.* 266.—Afr. austr.
hirsuta, Eckl. & Zeyh. l. c. = odorata.
hirta, *Burm. f. Fl. Cap. Prod.* 31 (err. typ. 27).—Afr. austr.
illicifolia, *Linn. Sp. Pl.* 1088.—Afr. austr.
internedia, *Eckl. & Zeyh. Enum.* 266.—Afr. austr.
juniperina, *Linn. f. Suppl.* 430.—Afr. austr.
laricina, E. Mey. ex Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 528 = juniperina.
leptophylla, Eckl. & Zeyh. Enum. 271 = filifolia.
linearifolia, *Eckl. & Zeyh. l. c.* 270.—Afr. austr.
marginata, *Eckl. & Zeyh. l. c.* 269.—Afr. austr.
Meyeriana, *Presl, Epim. Bot.* 202.—Afr. austr.
obcordata, *Linn. f. Suppl.* 428.—Afr. austr.
obtusa, Spreng. Neue Entdeck. ii. 174 = obcordata.
obovata, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 597.— Afr. austr.
ocinoides, *Cham. in Linnaea*, vi. (1831) 350.—Afr. austr.
odorata, *Linn. f. Suppl.* 431.—Afr. austr.
phylicoides, Eckl. & Zeyh. Enum. 269 = eriocephalina.
polycephala, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 602.—Afr. austr.
polygonifolia, *Linn. Sp. Pl.* 1038.—Afr. austr.
polyphylla, Eckl. & Zeyh. Enum. 268 = serpyllifolia.
propinqua, Eckl. & Zeyh. l. c. = serpyllifolia.
pulchella, *Linn f. Suppl.* 430.—Afr. austr.
pungens, *Presl, Epim. Bot.* 202.—Afr. austr.
reticulata, Eckl. & Zeyh. Enum. 266 = odorata.
rubricaulis, Presl, Bot. Bemerk. 141 = illicifolia.
ruscifolia, *Linn. Sp. Pl.* 1038.—Afr. austr.
sarmentosa, *Linn. Mant.* ii. 299.—Afr. austr.
serrata, *Eckl & Zeyh. Enum.* 269.—Afr. austr.
serpyllifolia, *Cham. & Schlecht. in Linnaea*, ii. (1827) 24.—Afr. austr.
serpyllifolia, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 599 = odorata.
serrata, *Reichb. ex Spreng. Syst.* iv. *Cur. Post.* 209.— Afr. austr.
serrata, Thunb. Prod. Fl. Cap. 98 = ferruginea.
spicata, *Reichb. ex Spreng. Syst.* iv. *Cur. Post.* 209.— Afr. austr.
stirbitiflora, *Murr. Syst. ed.* XIII. 749.—Afr. austr.
teretifolia, *Linn. f. Suppl.* 430.—Afr. austr.
ternata, *Linn. f.* l. c. = polygonifolia.
tricuspidata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 599. —Afr. austr.
tridentata, Willd. Sp. Pl. iv. 836 = ruscifolia.
trifoliata, Linn. Sp. Pl. 1038 = polygonifolia.

CLIFFORTIA:—
trilobe, *Harv. in Harv. & Sond. Fl. Cap.* ii. 598.— Afr. austr.

CLIFFORTIODES, Dryand. ex Hook. Ic. Pl. t. 689 (1844) = **Fagus**, Tourn. (Cupulif.).
oblongata, Dryand. l. c. 689, 678 = F. Solandri.

CLIPTONIA, Banks, ex Gaertn. f. Fruct. lii. 246. t. 225 (1805). *CYRILLEAE*, Benth. & Hook. f. ii. 1226.
MYLOCARYUM, Willd. Enum. Hort. Berol. 454 (1809).
WALTERIANA, Fras. ex Endl. Gen. 1413 (1841).
ligustrina, Sims, ex Spreng. Syst. ii. 816 = nitida.
nitida, *Gaertn. f. l. c.* 247.—Am. bor.

CLIMACANDRA, Miq. Pl. Jungh. 199 (1851–55) = **Actela**, Sw. (Myrsin.).
littoralis, Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 68 = A. humilis.
multiflora, Miq. Pl. Jungh. 200 = A. humilis.
obovata, Miq. l. c. 199 = A. humilis.
pergamacea, Miq. Fl. Ind. Bat. Suppl. 578 = Embelia pergamacea.
salicifolia, Miq. Pl. Jungh. 200 = A. humilis.

CLIMEDIA, Rafarin, in Rev. Hortic. (1877) 88, sphalm. = **Clidemia**, D. Don (Melast.).

CLINACANTHUS, Nees, in DC. Prod. xi. 511 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1105.
angustus, Nees, l. c. = lagiosma angusta.
Burmanni, *Nees, l. c.*—Java.

CLINANTHUS, Herb. App. 40 (1821) = **Stenomesson**, Herb. (Amaryll.).
luteus, Herb. l. c. 40 = S. luteum.

CLINHYMENIA, A. Rich. & Gal. in Comptes-Rendus, Acad. Sc. Par. xviii (1844) 512 = **Cryptarrhena**, R. Br. (Orchid.).
pallidiflora, A. Rich. & Gal. l. c. = Crypt. pallidiflora.

CLINOGYNE, Salisb. in Trans. Hort. Soc. i. (1812) 276. *SCITAMINEAE*, Benth. & Hook. f. iii. 653.
DONAX, Lour. Fl. Cochinch. ii (1790).
dichotoma, *Salisb. l. c.*—Ind. or.; Malaya.
filipes, *Benth. & Hook. f. Gen.* iii. 654.—Afr. trop.
grandis, *Benth. & Hook. f. l. c.*—Burma; Malaya.
virgata, *Benth. & Hook. f. l. c.*—Ind. or.

CLINOPODIUM, Linn. Syst. ed. I (1735) = **Calamintha**, Tourn. (Labiat.).
aegyptiacum, Mill. Gard. Dict. ed. VIII. n. 6 = Calamintha Clinopodium.
albidum, Vell. Fl. Flum. 242; vi. t. 6 = Hyptis paludosa.
angustifolium, Roxb. Hort. Beng. 44, nomen (quid ?). —Hab.?
arundanum, Nym. Syll. 101 = Calamintha Clinopodium.
arvense, Vell. Fl. Flum. 242; vi. t. 9.—Bras.
asiaticum, Lour. Fl. Cochinch. ii. 374 (= *Melissa asiatica*) (= Anisochili sp.).—Cochinch.
atropurpureum, Hort. Hafn. ex Steud. Nom. ed. II. i. 256 = Calamintha Clinopodium.
brasiliense, Vell. Fl. Flum. 241; vi. t. 8.—Bras.
capitatum, Sw. Prod. Veg. Ind. Occ. 88 = Hyptis capitata.
carolinianum, Mill. Gard. Dict. ed. VIII. n. 5 = Calamintha caroliniana.
Chamaedrys, Vahl, Symb. Bot. iii. 77 = Marsypianthes hyptoides.
cordatum, Vell. Fl. Flum. 243; vi. t. 5.—Bras.
fruticosum, Forsk. Fl. Aegypt. Arab. 107 = Otostegia sericea.
glabrescens, Pomel, Nouv. Mat. Fl. Atl. 128.— Algeria.
grandiflorum, Gouldest. ex Bieb. Fl. Taur. Cauc. iii. 299 = Stachys grandiflora.
humile, Mill. Gard. Dict. ed. VIII. n. 4.—Am. bor.

CLINOPODIUM:—
imbricatum, Vell. Fl. Flum. 242; vi. t. 5 = Hyptis pectinata.
incanum, Linn. Sp. Pl. 588 = Pycnanthemum incanum.
longicaule, Benth. in Wall. Pl. As. Rar. i. 66 = Calamintha longicaulis.
martinicense, Jacq. Enum. Pl. Carib. 25 = Leucas martinicensis.
origanifacie, Gillib. Fl. Lituan. i. 76 = Calamintha Clinopodium.
origanifolium, Labill. Ic. Pl. Syr. dec. iv. 14. t. 9 = Calamintha origanifolia.
plumosum, Sieber, in Flora, v. (1822) 242 = Calamintha Clinopodium.
repens, Roxb. Hort. Beng. 44; Fl. Ind. iii. 13 = Calamintha umbrosa.
repens, Vell. Fl. Flum. 242; vi. t. 7 = Peltodon radicans.
rugosum, Linn. Sp. Pl. 588 = Hyptis radiata.
variegatum, Hort. ex Steud. Nom. ed. II. i. 886 = Calamintha Clinopodium.
verticillatum, Vell. Fl. Flum. 242; vi. t. 4.— Bras.
villosum, Noé, in Bull. Soc. Bot. Fr. ii. (1855) 580 = Calamintha Clinopodium.
vulgare, Linn. Sp. Pl. 587 = Calamintha Clinopodium.

CLINOSTIGMA, H. Wendl. in Bonplandia, x. (1862) 196. *PALMAE*, Benth. & Hook. f. iii. 894.
Billardierii, Becc. Malesia, i. (1877) 41 = Microkentia Billardierii.
bractealis, Becc. l. c. 40 = Cyphokentia bractealis.
Deplanchei, Becc. l. c. 41 = Microkentia Deplanchei.
eriostachys, Becc. l. c. = Microkentia eriostachys.
Humboldtiana, Becc. l. c. 40 = Cyphokentia Humboldtiana.
macrostachya, Becc. l. c. = Cyphokentia macrostachya.
Moortanum, *F. Muell. Fragm.* viii. 265.—Austral.
Pancheri, Becc. Malesia, i. (1877) 40 = Microkentia Pancheri.
robusta, Becc. l. c = Cyphokentia robusta.
samoëne, *H. Wendl. in Bonplandia*, x. (1862) 196.— Ins. Samoa.
surculosa, Becc. Malesia, i. (1877) 41 = Microkentia surculosa.
vaginata, Becc. l. c. = Cyphokentia vaginata.

CLINOSTYLIS, Hochst. in Flora, xxvii. (1844) 26 = **Gloriosa**, Linn. (Liliac.).
speciosa, Hochst. l. c. = G. abyssinica.

CLINTONIA, Dougl. ex Lindl. Bot. Reg. t. 1241 (1829) = **Downingia**, Torr. (Campanulac.).
Bergiana, Cham. in Linnaea, vii. (1832) 217 = Lobelia Bergiana.
corymbosa, DC. Prod. vii. 347 = D. elegans.
elegans, Dougl. ex Lindl. Bot. Reg. t. 1241 = D. elegans.
pulchella, Lindl. l. c. t. 1909 = D. pulchella.
pusilla, G. Don, Gen. Syst. iii. 716.—Chili.

CLINTONIA, Rafin. in Am. Monthly Mag. (1818) 266; Journ. Phys. lxxxix. (1819) 109. *LILIACEAE*, Benth. & Hook. f. iii. 832.
HYLOCHARIS, Spreng. ex Regel & Tiling, in Nouv. Mém. Soc. Nat. Hosc. xi. (1859) 123.
XENIATRUM, Salisb. Gen. Pl. Fragm. 58 (1866).
Aitoni, Rafin. Ant. Journ. 120 = borealis.
alpina, *Kunth, Enum. Pl.* v. 156.—Reg. Himal.
Andrewsiana, *Torr. in Pacif. Rail. Rep.* iv. 150.— Calif.
Andrewsii, Wood, in Proc. Acad. Philad. (1868) 174 = Andrewsiana.
angustifolia, *Rafin. New Fl. Am.* ii. 79.—Am. bor.
bifora, Rafin. l. c. 78.—Am. bor.
bisumbella, Rafin. l. c. 81 = borealis.
borealis, *Rafin. Atl. Journ.* 120.—Am. bor.
ciliata, Rafin. in Am. Monthly Mag. (1819) 357 = borealis.
decandra, *Rafin. Atl. Journ.* 120.—Am. bor.
falcata, *Rafin. New Fl. Am.* ii. 79.—Am. bor.
fulva, *Rafin. l. c.* 82.—Am. bor.
glomerata, *Rafin. l. c.* 83.—Am. bor.
latifolia, *Rafin. l. c.* 81.—Am. bor.
multiflora, Beck, Bot. 358 = umbellata.

CLINTONIA:—
multiflora, *Rafin. Atl. Journ.* 190.—Am. bor.
mutans, Rafin. Ann. Nat. 16 = borealis.
nutans, *Rafin. Med. Fl.* ii. 58 = præc.
odorata, Rafin. Ann. Nat. 16 = umbellata.
ophioglossoides, *Rafin. New Fl. Am.* ii. 80.—Am. bor.
parviflora, *Rafin. Ann. Nat.* 16.—Am. bor.
podagrista, *Rafin. l. c.*—Am. bor.
trifora, *Rafin. New Fl. Am.* ii. 79.—Am. bor.
udensis, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 99).—Sibiria; Japon.
umbellata, *Torr. Fl. N. York*, ii. 201.—Am. bor.
undulata, Rafin. New Fl. Am. ii. 81 = borealis.
uniflora, *Kunth, Enum. Pl.* v. 159.—Am. bor. occ.

CLIOCARPUS, Miers, in Ann. Nat. Hist. Ser. II. iv. (1849) 141 = **Solanum**, Tourn.
didymus, Miers, l. c. xv. (1855) 204 = S. didymum.
eriocalyx, Miers, l. c. = S. trichocalyx.
Gardneri, Miers, l. c. iv. (1849) 141.—Bras.
megalochiton, Miers, l. c. xv. (1855) 203 = S. megalochiton.

CLIOCOCCA, Bab. in Proc. Linn. Soc. i. (1841) 90; et Trans. Linn. Soc. xix. (1842) 33. t. 3 = **Linum**, Tourn.
tenuifolia, Bab. in Trans. Linn. Soc. xix. (1845) 33, 34 = l. scleginoides.

CLIPTERIA, Rafin. New Fl. Am. ii. 44 (1838) = **Bolipta** †, Linn. (Compos.).
dichotoma, Rafin. l. c.—Am. bor.

CLISTANTHUS, Musil. Arg. in DC. Prod. xv. ii. 1025 (1866) = CLISTRANTHUS, Polt. = **Pera**, Mutis (Euphorb.).

CLISTAX, Mart. Nov. Gen. et Sp. iii. 26 (1829).
ACANTHACEAE, Benth. & Hook. f. ii. 1100.
CORYTHACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836).
brasiliensis, *Mart. l. c.*—Bras.
speciosus, *Nees, in Mart. Fl. Bras.* ix. 14.—Bras.

CLISTRANTHUS, Polt. ex Baill. Etud. Gén. Euph. 434 (1858) = **Pera**, Mutis (Euphorb.).

CLITANDRA, Benth. in Hook. Niger Fl. 445 (1849).
APOCYNACEAE, Benth. & Hook. f. ii. 692.
cirrhosa, *Radlk. in Bremen Abh.* viii. (1883) 400.—Afr. trop.
cymulosa, *Benth. in Hook. Niger Fl.* 445.—Afr. trop.

CLITANTHES, Herb. in Bot. Reg. (1839) Misc. 87 = **Childanthus**, Herb. & **Stenomesson**, Herb. (Amaryll.).
humilis, Herb. l. c. = Stenomesson humile.
luteo, Herb. l. c. = Childanthus fragrans.
Macleanicus, Herb. l. c.—Peruv.

CLITANTHUM, Benth. & Hook. f. Gen. iii. 738 (1880–83) = **Childanthus**, Herb. (Amaryll.).

CLITOCYAMOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128, 133 = **Ipomoea**, Linn. (Convolvulac.).
pinnatifidus, St. Lag. ll. cc.=I. Quamoclit.

CLITORIA, Linn. Gen. Pl. ed. I. 216 (1737).
LEGUMINOSAE, Benth. & Hook. f. i. 527.
MARTIA, Leandr. Secr. in Denkschr. Akad. Wissensch. vii. (1821) 228. t. 12.
MARTIUSIA, Schult. Mant. i. 69 (1822).
NAUCHEA, Desvaux. in Mem. Soc. Linn. Par. iv. (1826) 7. t. 1.
NEUROCARPUM, Desv. Journ. Bot. i. (1813) 119; iii. (1814) 75.
RHOMBIFOLIUM, Rich. ex DC. Prod. ii. 235 (1825).
TERNATEA, Tourn. ex Linn. Syst. ed. I. (1735); N. B. & K. Nov. Gen. et Sp. vi. 415 (1823).
VEXILLARIA, Rafin. in Am. Monthly Mag. (1818) 366.
acuminata, Benth. in Ann. Wien. Mus. i. (1838) 115 = RHOMBIFOLIUM.

CLITORIA:—
acuminata, R. Grah. in Wall. Cat. n. 5946 = Martiana.
alabamensis, Bertol. Misc. Bot. ix. 12 = virginiana.
alba, G. Don, Gen. Syst. ii. 216.—Ins. S. Thomas.
amazonum, *Mart. in Ann. Wien. Mus.* ii. (1838) 115.—Bras.
amœna, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) 24 = arborescens.
amœna, Roth, in Roem. Arch. i. III. 42 = Centrosema virginianum.
angustifolia, H. B. & K. Nov. Gen. et Sp. vi. 417 = Centrosema angustifolium.
angustifolia, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 29 = Galactia Martii.
aquilupinoïs, Raeusch. Nom. ed. III. 210 = quadalupeïdis.
arborea, Benth. in Ann. Wien. Mus. ii. (1838) 115 = Hoffmanseggii.
arborescens, *[R. Br. in] Ait. Hort. Kew.* ed. II. iv. 302.—Ins. Trinitat.
australis, *Benth. Fl. Austral.* ii. 242.—Austral.
Berteriana, DC. Mém. Lég. vi. 289 = Feriandra Berteriana.
bifora, *Dals. in Hook. Kew Journ.* ii. (1850) 35.—Ind. or.
brachystegia, *Benth. Bot. Voy. Sulph.* 84.—Ecuador.
bracteata, Poir. Encyc. Suppl. ii. 801 = C. Ternatea.
brasiliana, Linn. Sp. Pl. 753 = Centrosema brasilianum.
brasiliana, Vell. Fl. Flum. 312; vii. t. 129 = Canavalia ensiformis.
Broussonetii, Balb. Cat. Taur. (1813) 26 = Cologania Broussonetii.
cajanifolia, *Benth. in Mart. Fl. Bras.* xv. i. 121.—Reg. trop.
calcarata, L'Hérit. ex DC. Prod. ii. 235 = Centrosema Plumieri.
calcarigera, Salisb. Parad. Lond. t. 51 = Centrosema virginianum.
capitata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Ind. occ.
coccinea, Schrad. in Goett. Gel. Ans. i. (1821) 717 = Periandra coccinea.
coccinea, *Benth. in Journ. Linn. Soc.* ii. (1858) 41.—Am. trop.
erecta, *Raxb. Hort. Beng.* 56.—Am. trop.
falcata, Lam. Encyc. ii. 51 = glycinoides.
falcata, Nees, in Flora, iv. (1821) 329 = Feriandra coccinea.
flagellaris, *Benth. in Journ. Linn. Soc.* ii. (1858) 39.—Am. trop.
fluminensis, Vell. Fl. Flum. 312; vii. t. 128 = Centrosema Plumieri.
formosa, H. B. & K. Nov. Gen. et Sp. vi. 417 = Centrosema brasilianum.
fulgens, Paxt. Mag. Bot. xi. (1844) 121 (= *Collaea scarlatina*).—Bras.
Galactia, Linn. Sp. Pl. ed. II. 1026 = Galactia pendula.
gemina, Vell. Fl. Flum. 312; vii. t. 126 = Centrosema virginianum.
glabella, Desf. Tabl. ed. I. 192 = Galactia glabella.
gladiata, Schrank, in Syll. Ratisb. i. (1824) 229 = Centrosema virginianum.
glomerata, *Griseb. Cat. Pl. Cub.* 74.—Cuba.
glycinoides, *DC. Prod.* ii. 234.—Bras.
Grahami, Steud. Nom. ed. II. i. 886 = Mariana.
grandiflora, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) ii. 189.—Mexic.
guadelupensis, vide quadelupeensis.
guianensis, *Benth. in Journ. Linn. Soc.* ii. (1858) 40.—Am. trop.
heterophylla, *Lam. Encyc.* ii. 51.—Ind. or.
Hoffmanseggii, *Benth. in Journ. Linn. Soc.* ii. (1858) 42.—Am. trop.
insulana, Vell. Fl. Flum. 313; vii. t. 131 = Centrosema brasilianum.
javanica, Miq. Fl. Ind. Bat. i. t. 226 = macrophylla.
javitensis, *Benth. in Journ. Linn. Soc.* ii. (1858) 42.—Mexico.
lactescens, Linn. Syst. ed. X. 1173 = Galactia pendula.
lascive, *Bojr. Hort. Maurit.* 93, nomen; *Benth. in Ann. Wien. Mus.* ii. (1838) 114.—Madag.
lasciflora, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 411 = Centrosema vexillatum.
laurifolia, Poir. Encyc. Suppl. ii. 801 = cajanifolia.
leptostachya, *Benth. in Journ. Linn. Soc.* ii. (1858) 43.—Am. trop.
macrophylla, *Wall. Cat.* n. 5345.—Burma.

CLITORIA:—
Mariana, *Linn. Sp. Pl.* 753.—Ind. or.; Burma; Am. bor.
Mariana, Moç. & Sesse, ex DC. Prod. ii. 238 = Galactia radicata.
mexicana, Link, Enum. Hort. Berol. ii. 235 = Mariana.
micrantha, *Guill. & Perr. Fl. Senegr. Tent.* i. 190.—Afr. trop.
micrantha, Spreng. Delic. Insub. ii. 1 = Galactia longifolia.
micrantha, Sm. ex Steud. Nom. ed. II. i. 886 = Centrosema virginianum.
mucronulata, *Benth. in Ann. Nat. Hist.* (1839) 435.—Bras.
multiflora, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 188.—Mexic.
multiflora, Sw. Prod. Veg. Ind. Occ. 106 = Vilmorinia multiflora.
nana, *Benth. in Journ. Linn. Soc.* ii. (1858) 40.—Am. trop.
oblonga, Hamk. in Flora, xxv. (1842) ii. Beibl. 48 = Centrosema virginianum.
occidentalis, *Sweet, Hort. Brit.* ed. II. 140.—Ind. occ.
parviflora, *Rafin. Atl. Journ.* 147.—Am. bor.
pedunculata, *Bojr. ex Benth. in Ann. Wien. Mus.* ii. (1838) 114.—Madag.
pedunculata, *Michels, in Kjoeb. Vidensk. Meddel.* (1875) 77.—Bras.
Pyrrse, Steud. Nom. ed. I. 208, 349 = Galactia sericea.
pilosula, Wall. Cat. n. 5347 = C. Ternatea.
Plumieri, *Turp. in Perr. Syn.* ii. 308 = Centrosema Plumieri.
Poitaei, Benth. in Ann. Nat. Hist. ii. (1839) 434 = arborescens.
Poiteaui, DC. Prod. ii. 234 = arborescens.
polyphylla, Poir. Encyc. Suppl. ii. 800 = Barbieria polyphylla.
polystachya, *Benth. Pl. Hartw.* 60.—Mexic.
portobellensis, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 119.—Panama.
pudica, *Steud. Nom.* ed. II. i. 886.—Cuba.
quadacensis, *Scop. Delic. Insub.* ii. 4.—Ins. Guadalup.
racemosa, *G. Don, Gen. Syst.* ii. 215; *Benth. in Ann Wien. Mus.* ii. (1838) 116.—Bras.
rubiginosa, Juss. ex Pers. Syn. ii. 313 = glycinoides.
rubiginosa, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 29 = Galactia Neesii.
rufescens, *Benth. in Journ. Linn. Soc.* ii. (1858) 39.—Bras.
Schiedeana, Schlecht. in Linnæa, xii. (1838) 284 = Centrosema pubescens.
Sellei, *Benth. in Journ. Linn. Soc.* ii. (1858) 42.—Am. trop.
simplicifolia, *Benth. l. c.* 40.—Am. trop.
sinuata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 120 = Centrosema sinuatum.
speciosa, *Cav. Descr.* 183.—Mexic.
spectabilis, Salisb. Prod. 336 = C. Ternatea.
stipularis, *Benth. in Journ. Linn. Soc.* ii. (1858) 41.—Am. trop.
Ternatea, *Linn. Sp. Pl.* 753.—Reg. trop.
ternatensium, Cranz, Inst. ii. 59 = C. Ternatea.
tetragona, Poir. ex DC. Prod. ii. 236 = glycinoides.
tomentosa, Mart. in Nov. Act. Nat. Cur. xii. (1824) 30 = Galactia tomentosa.
tristis, *Tenore, Cat. Hort. Neap.* (1845) 32.—Hab.?
viciodes, Nees & Mart. in Nor. Act. Nat. Cur. xii. (1824) 28 = Cologania viciodes.
virginiana, Linn. Sp. Pl. 753 = Centrosema virginianum.
viridiflora, Boston, ex Hook. Ic. Pl. t. 152 = Dolichos axillaris.
xanthorrensis, *Vathe, in Oestr. Bot. Zeitschr.* xxviii. (1878) 261.—Afr. trop.
xeuphthalmum, Linn. Syst. ed. X. 1172 = Dolichos uncus?

CLITORIASTRUM, Heist. Syst. 9 (1748) = **Clitoria**, Linn. † (Leguminos.).

CLIVIA, Lindl. Bot. Reg. t. 1128 (1828). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 729.
IMATOPHYLLUM, Hook. Bot. Mag. t. 2826 (1836).
Gardeni, *Hook. Bot. Mag.* t. 4826.—Afr. austr.
miniata, *Regel, Gartenfl.* (1864) 181.—Afr. austr.
nobilis, *Lindl. Bot. Reg.* t. 1183.—Afr. austr.

CLOANTHE, Nees, in R. Br. Verm. Schr. i. 78 (1825) = **Chloanthes**, R. Br. (Verbenac.).

CLOEZIA, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 578. *MYRTACEÆ*, Benth. & Hook. f. i. 709.
BALLARDIA, Montr. in Mém. Acad. Lyon, x. (1860) 204.
MOORIA, Montr. l. c. (1860) 207.
buxifolia, *Brongn. & Gris, l. c.* 577.—N. Caled.
canescens, *Brongn. & Gris, l. c.*—N. Caled.
Deplanchei, *Brongn. & Gris, l. c.*—N. Caled.
floribunda, *Brongn. & Gris, l. c.*—N. Caled.
ligustrina, *Brongn. & Gris, l. c.*—N. Caled.
semilifolia, *Brongn. & Gris, l. c.*—N. Caled.

CLOMENA, Beauv. Agrost. 98. t. 7. f. 10 (1812) = Mœhlenbergia, Schreb. (Gramin.).
perwviana, Beauv. l. c. = M. Clomena.

CLOMENOCOMA, Cass. in Dict. Sc. Nat. lx. 416 (1817) = Dysodia, Cav. (Compos.).
aurantia, Cass. l. c. = D. appendiculata.
montana, Benth. Pl. Hartw. 86 = D. grandiflora.
pinnata, DC. Prod. v. 641 = D. pubescens.

CLOMENOLEPIS, Cass. in Dict. Sc. Nat. iii. Suppl. 64 (1816), nomen (Compos.).

CLOMIUM, Adans. Fam. ii. 116 (1763) = Carduus, Tourn. (Compos.).

CLOMOPANUS, Stend. Nom. ed. II. i. 386 (1840) = seq.

CLOMPANUS, Aubl. Pl. Gui. ii. 778 (1775) = Lonchocarpus, Humb. & Bonpl. (Legumin.).
paniculata, Aubl. l. c.—Guiana.

CLOMPANUS, Rafin. Sylva Tellur. 78 (1838) = Sterculia, Linn.
molucanus, Rafin. l. c.—Ins. Molucc.

CLONODIA, Griseb. in Mart. Fl. Bras. xii. 1. 26 (1858). *MALPIGHIACEÆ*, Benth. & Hook. f. i. 258.
verruosa, *Griseb. l. c.*—Bras.

CLONOSTACHYS, Klotzsch, in Wiegm. Archiv, vii. (1841) 185 = Sebastiania, Spreng. (Euphorbiac.).

CLOSASCHIMA, Korth. Verh. Nat. Gesch. Bot. 139 (1842) = Lasplacea, H. B. & K. (Ternstroemiac.).
marginata, Korth. l. c. 141 = L. marginata.
ovalis, Korth. l. c. 140 = L. ovalis.

CLOSIA, Remy, in C. Gay, Fl. Chil. iv. 119. t. 46. f. 2 (1849). *COMPOSITÆ*, Benth. & Hook. f. ii. 404.
anthemoides, *Phil. Fl. Atac.* 31.—Chili.
catnia, *Remy, in C. Gay, Fl. Chil.* iv. 120.—Chili.
chilensis, *Regel & Korrn. Ind. Sem. Hort. Petrop.* (1857) 39.—Chili.
discoides, *Phil. Fl. Atac.* 31.—Chili.
elata, *Phil. l. c.*—Chili.
pusilla, *Phil. l. c.*—Chili.

CLOSIROSPERMUM, Neck. Elem. i. 54 (1790) = Crepis, Linn. (Compos.).

CLOSTERANDRA, Boiv. ex Belang. Voy. Indes Or. t. absque num. (1839) = Papaver, Tourn.
minor, Boiv. l. c. = P. Belangeri.

CLOSTERANTHERA, Walp. Rep. i. 115 (1842) = præc.

CLOWESIA, Lindl. Bot. Reg. xxix. (1843) Misc. 25. t. 39 = Catasetum, Rich. (Orchid.).
rosea, Lindl. l. c. = Catasetum roseum, *Reichb. f.*

CLUACENA, Rafin. Sylva Tellur. 104 (1838) = Myrtus, Tourn.
myrsinoides, Rafin. l. c. = M. myrsinoides.
vaccinioides, Rafin. l. c. = M. vaccinioides, *H. B. & K.*

CLUERIA, Rafin. Fl. Tellur. iii. 87 (1836) = Bromostachys, Bunge (Labiat.).
laciniata, Rafin. l. c. = E. laciniata.

CLUGNIA, Comm. ex DC. Syst. i. 433 (1818) = Wormia, Rottb. (Dilleniac.).
volubilis, Comm. ex Steud. Nom. ed. I. 906 = W. madagascariensis.
volubilis, Comm. ex DC. Syst. i. 433 = W. madagascariensis.

CLUSIA, Linn. Gen. ed. I. 344 (1737). *GUTTIFERÆ*, Benth. & Hook. f. i. 170.
ANDROSTYLIUM, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) 93.
ARRUDEA, St. Hil. Fl. Bras. Mer. i. 318. t. 66 (1827).
ASTROTHRICA, Miers, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 254.
BIROLIA, Rafin. Sylva Tellur. 103 (1838).
CAHOTIA, Karst. in Linnæa, xxvii. (1856) 448.
COCHLANTHERA, Choisy, Coll. Ind. 46. t. 3 (1823).
ELWERTIA, Rafin. Sylva Tellur. 103 (1838).
FIRKEA, Rafin. l. c. (1838).
HAVETIA, H. B. & K. Nov. Gen. et Sp. v. 208 (1821).
ICOSTEGIA, Rafin. Sylva Tellur. 103 (1838).
LIPOSTYLLUM, Miers, in Trans. Linn. Soc. xxi. (1855) 251. t. 26.
OXYSTEMON, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 314.
QUAPOYA, Aubl. Pl. Gui. ii. 897 (1775).
SMITHIA, Scop. Introd. 1272 (1777..
TRIPLANDRON, Benth. Bot. Voy. Sulph. 78. t. 36 (1844).
XANTHE, Schreb. Gen. 710 (1791).
acuminata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 358.—Panama.
acuminata, *Spreng. Syst.* ii. 599.—Porto Rico.
alata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 361.—N. Granat.
alba, Choisy, in Planch. & Triana, l. c. 326 = palmicida.
alba, *Yacq. Enum. Pl. Carib.* 34.—Ind. occ.
alba, Kunth, ex Griseb. Fl. Brit. W. Ind. 107 = rosea.
amazonica, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 358.—Bras.
bicolor, *Nort. Nov Gen. et Sp.* iii. 164.—Bras.
Brongniartiana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 355.—Ind. occ.
Cambessedii, *Planch. & Triana, l. c.* 360.—Bras.
Candelariana, *Planch. & Triana, l. c.* 348.—Am. trop.
casuinioides, *Planch. & Triana, l. c.* 369.—Peruv.
Coulteri, Duchass. ex Planch. & Triana, l. c. 333 = minor.
crassifolia, *Planch. & Triana, l. c.* 351.—Am. trop.
Criuva, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 317.—Bras.
cuneata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 368.—Guiana.
decussata, *Ruiz & Pav. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 320, 345.—Peruv.
Ducu, *Benth. Pl. Hartw.* 126.—Peruv.
elliptica, *H. B. & K. Nov. Gen. et Sp.* v. 199.—Peruv.
eugenioides, *Planch. & Linden, ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 319, 328.—N. Granat.
flava, *Yacq. Enum. Pl. Carib.* 34.—Jamaica.
flava, Willd. ex Steud. Nom. ed. II. i. 387 = Hoffmannseggiana.
flumineusis, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 349.—Bras.
Fockeana, *Miq. in Hoev. & De Vriese, Tijdschr.* x. (1843) 89.—Guiana.
fragrans, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 334.—Bras.
ganabarica, Casar. Nov. Stirp. Bras. Decad. 60 = Sellowiana.
Gardneri, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 319, 328.—Bras.
Gaudichaudii, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 316.—Bras.
Gaudichaudii, Choisy, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 359 = Sellowiana.
grandiflora, *Splitg. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 101.—Guiana.
guatemalensis, *Hemsl. Diag. Pl. Nov.* 2.—Am. centr.
Gundlachi, *Stahl, Enbod. Pl. Puerto-Rico, foll.* ii. 123.—Ind. occ.
havetioides, *Planch. & Triana, in Ann. Sc. Nat. Sér.* IV. xiii. (1860) 368.—Ind. occ.
Hilariana, *Schlecht. in Linnæa*, viii. (1833) 161.—Bras.
Hoffmannseggiana, *Schlecht. l. c.* 165.—Bras.
Hookeriana, A. Rich. ex Planch. & Triana, in Ann. Sc. Nat. Ser. IV. xiii. (1860) 359 = Sellowiana.

CLUSIA:—
insignis, *Mart. Nov. Gen. et Sp.* iii. 164.—Bras.
lanceolata, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 318.—Bras.
latipes, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 365.—N. Granat.
laurifolia, *Planch. & Triana, l. c.* 344.—N. Granat.
laxiflora, *D. Dietr Syn. Pl.* iv. 866.—Peruv.
leprantha, *Mart. Nov. Gen. et Sp.* iii. 165.—Bras.
leprata, *Mart. l. c.*—Bras.
lencantha, *Schlecht. in Linnæa*, viii. (1833) 186.—Bras.
Lhotskyana, *Schlecht. in Linnæa*, viii. (1833) 184.—Bras.
Lhotzkyana, Choisy, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 319, pro parte = Hilariana.
linesta, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 345.—N. Granat.
longifolia, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 113 = Micranthera clusioides.
loranthacea, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 346.—N. Granat.
macrocarpa, *Spreng. Syst.* ii. 599.—Guiana.
major, Linn. Sp. Pl. 509 = alba.
mammosa, *Casar. Nov. Stirp. Bras. Decad.* 60.—Bras.
Mangle, *Rich. ex Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 369.—Ins. Guadelup.
maxima, Rich. ex Planch. & Triana, l. c. 325 = grandiflora.
micrantha, Choisy, ex Planch. & Triana, l. c. 329 = Sellowiana.
microcarpa, *Spreng. Syst.* ii. 598.—Guiana.
microstemon, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 331.—Bras.
minor, *Linn. Sp. Pl.* 510.—Guatemala.
multiflora, *H. B. & K. Nov. Gen. et Sp.* v. 200.—N Granat.
myriandra, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 332.—Guiana.
nemorosa, *G. F. W. Mey. Prim. Fl. Esseq.* 203.—Guiana.
nutans, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 373.—N. Granat.
odorata, *Sesm. Bot. Voy. Herald*, 89.—Am. centr.
organensis, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 349.—Bras.
Orizabæ, *Hemsl. Diag. Pl. Nov.* 3.—Mexic.
ovigera, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 354.—Mexic.
palmicida, *Rich. ex Planch. & Triana, l. c.* 326.—Guian.
Panapanari, Choisy, in DC. Prod. i. 559 = macrocarpa.
parviflora, *Humb & Bonpl. ex Willd. Sp. Pl.* iv. 976.—Ins. Margarita.
Pavonii, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV xiii. (1860) 363.—Peruv.
pedicellata, *Forst. f. Prod.* 74. n. 390.—N. Caled.
pentarhyncha, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 365.—N. Granat.
petiolaris, *Planch. & Triana, l. c.* 373.—N. Granat.
Plumerii, *Planch. & Triana, l. c.* 340.—Ina. S. Doming.
popayenensis, *Planch. & Triana, l. c.* 363.—N. Granat.
pratensis, Seem. Bot. Voy. Herald, 89 = minor.
pseudochina, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 12.—Peruv.
pseudo-haveti, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 366.—Peruv.
pseudo-mangle, *Planch. & Triana, l. c.* 370.—Peruv.
Puat, *Montr. in Mém. Acad. Lyon*, x. (1860) 188.—N. Caled.
Quapoya, Choisy, in DC. Prod. i. 559 = microcarpa.
radicans, *Pav. ex Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 374.—Peruv.
ranggeriolides, *Planch. & Triana, l. c.* 350.—Bras.
retusa, Poir. Encyc. v. 138 = rosea.
rosæflora, *Planch. & Linden, ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 343.—Venezuela.
rosea, *Yacq. Enum. Pl. Carib.* 34.—Ins. S. Doming.
rosea, Cambess. in St. Hil. Fl. Bras. Mer. i. 316 = Hilariana.
rupicola, *Casar. Nov. Stirp. Bras. Decad.* 60.—Bras.
Seemanii, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 347.—Panama.

CLUSIA :—
Sellowiana, *Schlecht. in Linnaea*, viii. (1833) 163.—Bras.
sessilifora, *Pair. Encyc.* v. 133.—Madag.
sessilis, *Forst. f. Prod.* 74. n. 391.—Ins. Amicor.
Sieboldii, *Regel, Cat. Pl. Hort. Absukor.* 36.—Hab.?
sphaerocarpa, *Planch. & Triana, in Ann. Sc. Nat.* Sér IV. xiii. (1860) 362.—Peruv.
Spruceana, *Planch. & Triana, l. c.* 346.—Peruv.
subsessilis, *Benth. Bot. Voy. Sulph.* 75.—N. Granat.
sulphurea, *D. Dietr. Syn. Pl.* iv. 866.—Peruv.
tetrandra, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 978. —Ecuador.
tharifera, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 364.—Peruv.?
venosa, *Yacq. Enum. Pl. Carib.* 24.—Ind. occ.
venosa, Linn. Sp. Pl. ed. II. 1495 = minor.
volubilis, *H. B. & K. Nov. Gen. et Sp.* v. 200.— N. Granat.
Weddelliana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 283.—Bras.

CLUSIANTHEMUM, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 333 = Garcinia, Linn. (Gustifer.).
coriaceum, Vieill. l. c. 339 = *pedicellatum*.
densiflorum, Vieill. ex Pierre, Fl. For. Cochinch. fasc. v. 25 = *D. densiflora*.
pedicellatum, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 339.—N. Caled.

CLUSIELLA, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 253. *GUTTIFERAE*, Benth. & Hook. f. i. 172.
elegans, *Planch. & Triana, l. c.*—N. Granat.

CLUSIOPHYLLEA, Baill. Adansonia, xii. (1878) 197 = **Plectronia**, Linn. (Rubiac.).
Perrieriana, Baill. l. c. (= *Canthium Perrillenanum*). —Madag.

CLUSIOPHYLLUM, Muell. Arg. in Flora, xlvii. (1864) 518 = Cunuria, Baill. (Euphorb.).
Spruceri, Muell. l. c. 519 = Cun. Spruceana.

CLUYTIA, Boerh. ex Linn. Syst. ed. I (1736) (Clutia). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 302.
ALTOR, Adans. Fam. ii. 356 (1763).
CRATOCHWIZLA, Neck. Elem. ii. 339 (1790).
abyssinica, *Jaub. & Spach, Illustr.* v. 77. t. 468.— Abyssin.
acuminata, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 135 = polifolia.
acuminata, *Linn. f. Suppl.* 432.—Afr. austr.
acuminata, Thunb. Prod. Pl. Cap. 58 = Lachnostylis hirta.
affinis, *Sond. in Linnaea*, xxiii. (1850) 126.—Afr. austr.
africana, Poir. Encyc. Suppl. ii. 802 = daphnoides.
alaternoides, *Linn. Sp. Pl.* 1042.—Afr. austr.
androgyna, Linn. Mant. i. 138 = Sauropus albicans.
bengalensis, *Mudl. Arg. in Journ. Bot.* ii. (1864) 357.—Afr. trop.
brevifolia, Sond. in Linnaea, xxiii. (1850) 125 = polifolia.
Cascarilla, Linn. Sp. Pl. 1042 = Croton Cascarilla.
collina, Roxb. Pl. Corom. ii. 37 = Cleistanthus collinus.
cordata, *Bernh. in Flora*, xxviii. (1845) 81.—Afr. austr.
cotinifolia, Salisb. Prod. 396 = pulchella.
curvata, C. Mey. ex Sond. in Linnaea, xxiii. (1850) 121 = polygonoides.
daphnoides, Eckl. & Zeyh. ex Muell. Arg. in DC. Prod. xv. ii. 1064 = polygonoides.
daphnoides, *Lam. Encyc.* ii. 54.—Afr. austr.
decandra, Crantz, Inst. i. 201 = Croton Eluteria.
dicmoides, *Sond. in Linnaea*, xxiii. (1850) 122.—Afr. austr.
diversifolia, Roxb. Hort. Beng. 104; Fl. Ind. iii. 781 = Cleistanthus diversifolius.
Dongeana, *Scheele, in Linnaea*, xxv. (1852) 568.—Afr. austr.
Eckloniana, *Muell. Arg. in DC. Prod.* xv. ii. 1064.— Afr. austr.
Eluteria, Linn. Sp. Pl. 1043 = Croton Eluteria.

CLUYTIA :—
ericoides, Krebs, in Flora, xxviii. (1845) 82 = polygonoides.
ericoides, *Thunb. Prod. Pl. Cap.* 58.—Afr. austr.
floribunda, Baill. Etud. Gen. Euph. Atlas, 20. t. 16. f. 1–8 = alaternoides.
gracilis, Baill. Adansonia, iii. (1862–63) 151 = tenuifolia.
heterophylla, Sond. in Linnaea, xxiii. (1850) 128 = Sonderiana.
heterophylla, *Thunb. Prod. Pl. Cap.* 58.—Afr. austr.
hirsuta, Eckl. & Zeyh. ex Sond. in Linnaea, xxiii. (1850) 128 = daphnoides.
hirsuta, *Muell. Arg. in DC. Prod.* xv. ii. 1046.—Afr. austr.
hirta, Linn. f. Suppl. 432 = Lachnostylis hirta.
hirta, *Vahl, Symb. Bot.* ii. 101.—Afr. austr.
humilis, Bernh. in Flora, xxvii. (1845) 81 = pubescens.
Jaubertiana, *Muell. Arg. in DC. Prod.* xv. ii. 1044.— Afr. trop.
lanceolata, *Forsk. Fl. Aegypt. Arab.* 170.—Arabia.
lanceolata, Jaub. & Spach, Illustr. Pl. Or. v. t. 467 = Jaubertiana.
marginata, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 130 = tomentosa.
Meyeriana, *Muell. Arg. in DC. Prod.* xv. ii. 1055.— Afr. austr.
monoica, Lour. Fl. Cochinch. ii. 638 = Cleistanthus monoicus.
montana, Roxb. Pl. Corom. iii. 38. t. 271 = Briedelia montana.
myricoides, *Jaub. & Spach, Illustr. Pl. Or.* v. 78. t. 456, 466.—Arabia.
natalensis, *Bernh. in Flora*, xxviii. (1845) 81.—Afr. austr.
oblongifolia, Roxb. Hort. Beng. 70; Fl. Ind. iii. 780 = Cleistanthus obovatus.
oblongifolia, Wall. Cat. n. 7886 = Trigonostemon semperflorens.
ovalis, Scheele, in Linnaea, xxv. (1852) 568 = Andrachne ovalis.
ovalis, *Sond. in Linnaea*, xxiii. (1850) 129.—Afr. austr.
patula, Herb. Roxb. ex Wall. Cat. n. 7877 = Cleistanthus collinus.
patula, Roxb. Pl. Corom. ii. 37. t. 170 = Cleistanthus patulus.
phyllanthifolia, Baill. Adansonia, iii. (1862–63) 153 = affinis.
polifolia, *Yacq. Hort. Schoenb.* iii. 67. t. 250.—Afr. austr.
polifolia, Sond. in Linnaea, xxiii. (1850) 124 = pterogona.
polygaloefolia, Salisb. Prod. 396 = alaternoides.
polygonoides, *Burm. f. Fl. Cap. Prod.* 81 (err. typ. 97). —Afr. austr.
polygonoides, *Linn. Sp. Pl.* ed. II. 1475.—Afr. austr.
polygonoides, Thunb. Prod. Pl. Cap. 58 = daphnoides.
polygonoides, Willd. Hort. Berol. t. 51 = alaternoides.
pterogona, *Muell. Arg. in DC. Prod.* xv. ii. 1048.— Afr. austr.
pubescens, Eckl. & Zeyh. ex Sond. in Linnaea, xxiii. (1850) 126 = affinis.
pubescens, *Thunb. Prod. Pl. Cap.* 58.—Afr. austr.
pubescens, Willd. Sp. Pl. iv. 581 = daphnoides.
pulchella, *Linn. Sp. Pl.* 1043.—Afr. austr.
pulchella, Sperm. ex Sond. in Linnaea, xxiii. (1850) 126 = daphnoides.
retusa, Herb. Wight, ex Wall. Cat. n. 7877 = Cleistanthus collinus.
retusa, Linn. Sp. Pl. 1475 = Bridelia retusa.
retusa, Moon, Cat. Pl. Ceyl. 71 = Bridelia Moonii.
Richardiana, *Muell. Arg. in DC. Prod.* xv. ii. 1044.— Afr. trop.
rubricaulis, Eckl. ex Sond. in Linnaea, xxiii. (1850) 128 = alaternoides.
scandens, Roxb. Pl. Corom. ii. 39. t. 173 = Bridelia stipularis.
semperflorens, Roxb. Hort. Beng. 104; Fl. Ind. iii. 780 = Trigonostemon semperflorens.
semperflorens, Wall. Cat. n. 7852 B, ex Baill. Etud. Gén. Euph. 474 = Symphylia silhetiana.
sessiliflora, Muell. Arg. in DC. Prod. xv. ii. 764 = Symphylia silhetiana, Trigonostemon Hookerianus.
serices, *Muell. Arg. l. c.* 1063.—Afr. austr.
similis, *Muell. Arg. l. c.* 1046.—Afr. austr.
Sonderiana, *Muell. Arg. l. c.* 1061.—Afr. austr.

CLUYTIA :—
spinosa, Roxb. Fl. Corom. ii. 38. t. 172 = Briedelia retusa.
squarrosa, Lam. Encyc. ii. 54 = Briedelia retusa.
stipularis, Linn. Mant. 137 = Briedelia stipularis.
tabularis, Eckl. & Zeyh. ex Sond. in Linnaea, xxiii. (1850) 123 = polygonoides.
tenuifolia, *Willd. Sp. Pl.* iv. 580.—Afr. austr.
tenuifolia, Sond. in Linnaea, xxiii. (1850) 124 = polifolia.
Thunbergii, Sond. l. c. 130 = daphnoides.
tomentosa, *Linn. Mant.* 299.—Afr. austr.
tomentosa, Thunb. Fl. Cap. ed. Schult. 971 = daphnoides.

CLUYTIA, Roxb. ex Steud. Nom. ed. II. i. 225 (1840) = **Briedelia**, Willd. (Euphorb.).

CLUYTIANDRA, Muell. Arg. in Journ. Bot. ii (1864) 328. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 272.
trichopoda, *Muell. Arg. l. c.*—Afr. trop.

CLYBATIS, Phil. in Anal. Univ. Chil. (1873) 742. *COMPOSITAE*, Benth. & Hook. f. ii. 1237.
Volckmanni, *Phil. l. c.* 748.—Chili.

CLYMENUM, [Tourn.] Linn. Syst. ed. I (1735) = **Lathyrus**, Tourn. (Legumin.).
alatum, Link, Handb. ii. 188 = L. alatus.
articulatum, Medic. Phil. Bot. i. 204 = L. articulatus.
bicolor, Moench, Meth. 150 = L. articulatus.
dalichoides, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = L. articulatus.
Ochrus, Link, Handb. ii. 188; Alef. in Bonplandia, ix. (1861) 127 = L. Ochrus.
tenuifolium, Link, l. c. 494; Alef. l. c. 138 = L. tenuifolius.
tingitanum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = L. tingitanus.
uncinatum, Moench, Meth. 150 = L. Clymenum.

CLYNHYMENIA, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 24, in syn. = CLINHYMENIA, A. Rich. & Gal.—Cryptarrhena, R. Br. (Orchid.).

CLYPEA, Blume, Bijdr. 26 (1825) = **Stephania**, Lour. (Menispermac.).
abyssinica, Dillon & A. Rich. in Ann. Sc. Nat. Ser. II. xiv. (1840) 268 = S. abyssinica.
acuminatissima, Blume, Bijdr. 28 = S. acuminata.
Burmanni, Wight & Arn. Prod. 14 = Cyclea Burmanni.
capitata, Blume, Bijdr. 28 = S. capitata.
consummata, Miers, Contrib. iii. 209.—Japon.
corymbosa, Blume, Bijdr. 27 = S. rotunda.
discolor, Blume, l. c. 26 = S. discolor.
effusa, Miers, Contrib. iii. 207.—Malabar.
Forsteri, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 270 = S. Forsteri.
glaucocarpa, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 493 t. 18 = S. glaucocarpa.
hernandifolia, Wight & Arn. Prod. 14 = S. hernandifolia.
longa, G. Don, Gen. Syst. i. 113 = S. hernandifolia.
oppfylla, Miers, Contrib. iii. 206.—Nepal.
rotunda, Steud. Nom. ed. II. i. 387 = S. rotunda.
subcostata, Miers, Contrib. iii. 209.—Japon.
tomentosa, Blume, Bijdr. 27 = S. tomentosa.
venosa, Blume, l. c. = S. venosa.

CLYPEOLA, Burm. ex DC. Prod. ii. 430 (1825) = **Pterocarpus**, Linn. (Legumin.).

CLYPEOLA, Linn. Gen. ed. I. 198 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 68.
BERGERETIA, Desv. Journ. Bot. iii. (1813) 161.
FOSSELINIA, Scop. Introd. 318 (1777).
IONTHLASPI, Adans. Fam. ii. 493 (1763).
ORIUM, Desv. Journ. Bot. iii. (1813) 161. t. 35. f. 10.
alliacea, Lam. Tabl. Encyc. iii. 118 = Peltaria alliacea.
alliacea, Schaeglin, ex Ledeb. Fl. Ross. i. 135 = Alyssum spathulatum.
alpestris, Linn. Sp. Pl. 953 = Alyssum calycinum.
amalgma, Jord. & Fourr. Rev. Pl. Nov. ii. 15 = microcarpa.

CLYPEOLA:—
auriculata, Cranta, Class. Crucif. 98 = Biscutella auriculata.
Brahnii, *Gruner, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 206.—Reg. Casp.
calycina, All. ex Stend. Nom. ed. II. i. 387 = Alyssum calycinum.
capensis, Burm. ex DC. Prod. ii. 490 = Wiborgia fusca.
capensis, Poir. Encyc. Suppl. ii. 304 = Heliophila diffusa.
carolinana, Walt. Fl. Carol. 178 = Lepidium virginicum.
chaetocarpa, Jaub. & Spach, Illustr. iii. 8. t. 206 = echinata.
ciliata, *Boiss. Fl. Orient.* i. 309.—Lycia.
edentula, Link, Handb. ii. 296 = Alyssum gemonense.
cyclostoma, *Delile, in Bull. Soc. Agr. Hérault*, (1880) 268.—Gallia.
dichotoma, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 176.—Persia; Armen.
didyma, Cranta, Stirp. Austr. ed. I. fasc. i. 20; ed. II. 22 = Biscutella laevigata.
echinata, Boiss. Fl. Orient. i. 310 = lappacea.
echinata, *DC. Syst.* ii. 338.—Syria; Persia; Afghan.
elegans, *Boiss. & Huet, Diagn.* Ser. II. v. 38.—Armenia.
eriocarpa, *Cav. Descr.* 401.—Hispan.
eriophora, DC. Syst. ii. 337 = eriocarpa.
Gaudini, Trachsel, in Flora, xiv. (1831) 748 = microcarpa.
glabra, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 178 = microcarpa.
gracilis, Planch. in Bull. Soc. Bot. Fr. v. (1858) 494 = microcarpa.
halimifolia, Link, Handb. ii. 295 = Alyssum halimifolium.
hispida, *Presl, Bot. Bemerk*, 9.—Arabia.
hispidula, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 15 = microcarpa.
indica, Poir. Encyc. Suppl. ii. 304 = Fortuynia Garcini.
Jonthlaspi, *Linn. Sp. Pl.* 652.—Reg. Mediterr.; Oriens.
laevigata, Jord. Brev. Pl. Nov. fasc. ii. 16 = microcarpa.
lapidicola, Jord. & Fourr. l. c. 14 = C. Jonthlaspi.
lappacea, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 174.—Mesopotam.; Persia.
lasiocarpa, Jaub. & Spach, Illustr. iii. 6. t. 205 = lappacea.
lasiocarpa, Jusa. ex Pers. Syn. ii. 198 = echinata.
lomatostricta, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 194 = microcarpa.
macrocarpa, Link, Handb. ii. 295 = Alyssum macrocarpum.
maritima, Georgi, Beschr. Russ. Reich. iii. iv. 1196 = Alyssum canescens.
maritima, Linn. Sp. Pl. 653 = Alyssum maritimum.
microcarpa, *Boiss. Diagn.* Ser. I. i. 74.—Graecia; Asia occ.
monosperma, Lam. Fl. Fr. ii. 462; 484 = C. Jonthlaspi.
montana, Cranta, Stirp. Austr. ed. I. fasc. i. 19; ed. II. 19 = Alyssum montanum.
perennis, Ard. Animad. Spec. 16. t. 6 = Peltaria alliacea.
petraea, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 14 = C. Jonthlaspi.
psilocarpa, Jord. & Fourr. l. c. = C. Jonthlaspi.
pyrenaica, Bordère, ex Dur. in Act. Soc. Linn. Bord. xxvi. (1866) 87 = C. Jonthlaspi
pyrenaica, Link, Handb. ii. 295 = Alyssum pyrenaicum.
rupestris, Link, l. c. = Alyssum rupestre.
Schengrini, Steven, in Pall. N. Nord. Beitr. vi. 274 = Alyssum spathulatum.
semiglabra, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 194 = C. Jonthlaspi.
spathulaefolia, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 15 = microcarpa.
spinosa, Link, Handb. ii. 295 = Alyssum spinosum.
tomentosa, *Linn.* Mant. 92 = Alyssum orientale.

CLYTIA, Stokes, Bot. Mat. Med. iv. 548 (1812) = Chrysita, Bourb. (Euphorb.).

CLYTORIA, J. S. Presl, Rostl. iii. 196 (1835) = Gliricidia, Linn. (Leguman.).

CLYTOSTOMA, Miers, ex Bur. in Adansonia, viii. (1868) 358 = Bignoniacostomum, Mast. (Bignonice.).
cuspidatolobatum, Bur.—Cf. Adansonia, viii. (1868) 358.—Sum. austr.

CNEMIDIA, Lindl. Bot. Reg. sub t. 1618 (1833) = Tropidia, Lindl. (Orchid.).
anguicoa, Lindl. l. c. = T. anguicoa.
bambusifolia, Thw. Enum. Pl. Zeyl. 314.—Zeylan.
ctenophora, Reichb. f. Otia Bot. Hamb. 51.—Ins. Fiji.
semilibera, Lindl. Bot. Reg. sub t. 1618 = T. anguicoa.

CNEMIDOSTACHYS, Mart. Nov. Gen. et Sp. i. 66. t. 40–44 (1824) = Sebastiana, Spreng. (Euphorb.).
acalyphoides, Mart. l. c. 71 = S. corniculata.
bidentata, Mart. l. c. 69 = S. virgata.
campestris, Mart. ex Baill. Etud. Gén. Euph. 516 = S. corniculata.
Chamaeleae, Spreng. Syst. iii. 885 = S. Chamaeleа.
coriacea, Mart. Nov. Gen. et Sp. i. 68 = S. marginata.
crotonoides, Mart. l. c. = S. corniculata.
daphnoides, Mart. l. c. = S. daphnoides.
dilasioides, F. Didr. in Kjoeb. Vidensk. Meddel. (1853) 88 = S. ditassoides.
dubia, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 241 = S. corniculata.
glabra, Mart. ex Baill. in Adansonia, v. (1864–65) 325 = S. multiramea.
glabrata, Mart. Nov. Gen. et Sp. i. 70 = S. corniculata.
glandulosa, Mart. l. c. 71 = S. corniculata.
hispida, Mart. l. c. = S. corniculata.
linearifolia, Miq. Fl. Ind. Bat. Suppl. 460 = S. Chamaeleа.
longifolia, Mart. Nov. Gen. et Sp. i. 71 = S. corniculata.
madagascariensis, Boj. Hort. Maurit. 284 = Excoecaria melanostricta.
marginata, Mart. Nov. Gen. et Sp. i. 68. t. 41 = S. marginata.
myrtilloides, Mart. l. c. 67. t. 40 = S. daphnoides.
oleoides, Mart. l. c. 71 = S. oleoides.
patula, Mart. in Flora, xxiv. (1841) ii. Beibl. 8 = S. corniculata.
prostrata, Mart. Nov. Gen. et Sp. i. 70 = S. corniculata.
salicifolia, Mart. l. c. = S. corniculata.
scoparia, Mart. l. c. = S. virgata.
Sellowiana, Klotzsch, ex Baill. Etud. Euph. 516 = S. corniculata.
serrulata, Mart. Nov. Gen. et Sp. i. 66 = S. serrulata.
stipulacea, Klotzsch, ex Baill. Etud. Gén. Euph. 516 = S. stipulacea.
tragioides, Mart. Nov. Gen. et Sp. i. 70 = S. corniculata.
Pohlii, Spreng. Syst. iii. 834 = S. corniculata.
velutina, Klotzsch, ex Baill. Etud. Gén. Euph. 516 = S. corniculata.

CNENAMUM, Teusch, in Flora, xi. (1828) 1. Eg. 80 = CRENAMUM, Adans. = Crepis, Linn. (Compos.).

CNEORIDIUM, Hook. f. in Benth. & Hook. f. Gen. i. 812 (1863). *SIMARUBEAE*, Benth. & Hook. f. i. 0.
PITAVIA, Nutt. ex Torr. & Gray, Fl. N. Am. 215 (1840).
dumosum, *Hook. f. l. c.*—Am. bor. occ.

CNEORUM, Linn. Syst. ed. I (1735). *SIMARUBEAE*, Benth. & Hook. f. i. 811.
pulverulentum, *Vent. Jard. Cels*, t. 77.—Ins. Canar.
tricoccon, Gmelid. ex Bieb. Fl. Taur. Casc. iii. 160 = Evonymus nanus.
tricoccon, *Linn. Sp. Pl.* 34.—Europ. austr.

CNESMOCARPUS, Zipp. ex Blume, Rumphia, iii. 115 (1847) = Pomatia, Forst. (Sapind.).
excelsus, Zipp. ex Blume, l. c. (= *Irina Diplocardia*).—N. Guinea.

CNESMONE, Blume, Bijdr. 680 (1825) (Cnesmose); Fl. Jav. Praef. p. vi (1828). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 320.
glabrata, *Kurz, in Flora*, lviii. (1875) 81.—Singapore.
javanica, *Blume, Bijdr.* 680.—Java.

CNESTIDIUM, Planch. in Linnaea, xxiii. (1850) 489. *CONNARACEAE*, Benth. & Hook. f. i. 428.
laxiconcarpum, *Baker, in Mart. Fl. Bras.* xiv. ii. 195.—Bras.
rubescens, *Planch. in Linnaea*, xxiii. (1850) 440.—Mexic.; Panama.

CNESTIS, Juss. Gen. 374 (1789). *CONNARACEAE*, Benth. & Hook. f. i. 428.
SPONDIOIDES, Sneathm. ex Lam. Encyc. iii. 28 (1789).
acuminata, Wall. Cat. n. 8838 = Rourea acuminata.
borbonientis, Rœusch. Nom. ed. III. 132 = glabra.
corymbulata, Blanco, Fl. Filip. ed. I. 386 = Connarus ferrugineus.
cornicalata, *Lam. Encyc.* iii. 28.—Afr. trop.
diffusa, Blanco, Fl. Filip. ed. I. 386 = Rourea rugosa.
emarginata, *Yach, in Malay. Misc.* ii. (1822) vii. 43.—Malaya.
ferruginea, *DC. Prod.* ii. 87.—Afr. trop.
flaminea, Griff. Notul. iv. 488 = ramiflora.
flammea, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 76 = prace.
florida, *Yach, in Malay. Misc.* ii. (1822) vii. 43.—Sumatra.
foliosa, Planch. ex Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 216 = ramiflora.
fraterna, *Planch. in Linnaea*, xxiii. (1850) 440.—Afr. trop.
glabra, Blanco, Fl. Filip. ed. I. 387 = Connarus paniculatus.
glabra, *Lam. Encyc.* iii. 28.—Ins. Mascar.
glabra, Naves, in Blanco, Fl. Philipp. ed. III. t. 146 = Rourea multiflora.
gracea, *Baker, in Oliver, Fl. Trop. Afr.* i. 461.—Afr. trop.
ignea, Planch. ex Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 216 = ramiflora.
laxiconcarpa, Mart. ex Baker, in Mart. Fl. Bras. xiv. ii. 195 = Cnestidium laxiconcarpum.
lucida, *Baill. Adansonia*, vii. (1866–67) 244.—Ins. Nossibé.
macrantha, *Baill. l. c.* 242.—Afr. trop.
madagascariensis, Raeusch. Nom. ed. III. 132 = polyphylla.
mimosoides, *Yach, in Malay. Misc.* ii. (1822) vii. 44.—Malaya.
monadelpha, Roxb. Hort. Beng. 34; Fl. Ind. ii. 454 = Rourea commutata.
natalensis, *Planch. & Sond. ex Harv. & Sond. Fl. Cap.* i. 528.—Afr. austr.
oblitqua, *Benth. Fl. Oman.* i. 97. t. 59.—Afr. trop.
obliqua, Boj. Hort. Maurit. 84 = Agelaea Lamarckii.
oblongifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 462.—Afr. trop.
pentaphylla, *Spen. in Linnaea*, xv. (1841) 189.—Ins. Timor.
pinnata, Beauv. Fl. Owar. i. 98. t. 60 = Rourea Palisotii.
platantha, Griff. Notul. iv. 484 = ramiflora.
polyphylla, Blanco, Fl. Filip. ed. II. 270 = Rourea rugosa.
polyphylla, *Lam. Encyc.* iii. 28.—Maurit.; Madag.
racemosa, *G. Don, Gen. Syst.* ii. 91.—Afr. trop.
ramiflora, *Griff. Notul. iv.* 483.—Ind. or.
scandens, J. F. Gmel. Syst. 729 = glabra ?
stenopetala, Griff. Notul. iv. 488 = Rourea stenopetala.
trifolia, Lam. Encyc. iii. 24 = Agelaea villosa.
westiis, Wall. ex Hook. f. Fl. Brit. Ind. ii. 47 = Agelaea vestita.
volubilis, Blanco, Fl. Filip. ed. I. 385 = Rourea heterophylla.

CNICOTHAMNUS, Griseb. in Goett. Abh. xix. (1874) 196. *COMPOSITAE*, Benth. & Hook. f. ii. 1387.
Lorentzii, *Griseb. l. c.* 197.—Reg. Argent.

CNICUS, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 466.
ACARNA, Hill, Veg. Syst. iv. 18 (1762).
ANCATHIA, DC. in Guill. Arch. Bot. ii. (1833) 331.
BERSEA, Lam. Syn. Comp. 9 (1888).
CEPHANANOPLOS, Neck. Elem. i. 46 (1790).
CHAMAEPEUCE, DC. Prod. vi. 657 (1837).
CIRSIUM, [Tourn.] Adans. Fam. ii. 116 (1763); DC. Fl. Fr. iv. 110 (1805).
CLOMIUM, Adans. Fam. ii. 116 (1763).
ECHENAIS, Cass. in Bull. Soc. Philom. (1816) 83.
ERYTHRACHYS, C. Koch, in Linnaea, xxiv. (1851) 396.
ERIOLEPIS, Cass. in Dict. Sc. Nat. xxiv. 173 (1825); xli. 331 (1826).
ERYTHROLAENA, Sweet, Brit. Flow. Gard. Ser. I. t. 134 (1825).

CNICUS:—
Ixine, Hill, Veg. Syst. iv. 17 (1762).
Lamyra, Cass. in Dict. Sc. Nat. xxv. 216 (1822).
Lophiolepis, Cass. l. c. xli. 218 (1826).
Notobasis, Cass. l. c. xxxv. 170 (1825).
Onotrophe, Cass. l. c. xxxvi. 146 (1825).
Orthocentron, Cass. l. c. xlix. 429 (1826).
Picnomon, Adans. Fam. ii. 116 (1763).
Platyraphium, Cass. in Dict. Sc. Nat. xxxv. 173 (1825).
Polyacantha, Hill, Veg. Syst. iv. 12 (1762).
Ptilostemon, Cass. in Dict. Sc. Nat. xxxv. 173 (1825).
Pycnocomon, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 78, 84.
Spanioptilon, Less. Syn. Comp. 10 (1832).
Tetralix, Hill, Veg. Syst. iv. 13 (1762).
Xylanthema, Neck. Elem. i. 67 (1790).

acanthoclepis, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.

Acarna, *Linn. Syst.* ed. X. 1201.—Europ.

acaulis, *Willd. Prod. Fl. Berl.* 260.—Europ.

acer, *Sibth. & Sm. Fl. Graec.* ix. 20. t. 839.—Graecia; As. Min.

acer, Tenore, Prod. p. lxiii = nivens.

acer, Willd. Sp. Pl iii. 1682 = Cn. Diacantha.

alpinus, Lobel. Fl. Gall. 542 = Saussurea alpina.

alsophilus, *Pollini, Fl. Veron.* ii. 620. t. 5.—Italia.

altaicus, Steph. ex DC. Prod. vi. 538 = Saussurea Frolowii.

altissimus, *Willd. Sp. Pl.* iii. 1671.—Am. bor.

ambiguus, Loisel. Fl. Gall. 540 = heterophyllus.

americanus, *A. Gray, in Proc. Am. Acad.* xix. (1884) 56.—Am. bor. occ.

Andersonii, *A. Gray, l. c.* x. (1874) 44.—Am. bor. occ.

Andrewsii, *A. Gray, l. c.* 45.—Calif.

anglicus, C. C. Gmel. Fl. Bad. iii. 872 = pratensis.

angustifolius, Buch.-Ham. in Wall. Cat. n. 2915 = Tricholepis Stebbingiana.

arabicus, *Roth, in Usteri, Ann. Bot.* viii. 8.—Arabia.

arachnoides, Hort. Par. ex DC. Prod. vi. 635 = fimbriatus.

arachnoides, Wall. Cat. n. 3882 = Wallichii.

arachnoideus, *Bieb. Fl. Taur. Cauc.* ii. 277.—Reg. Caucas.

arancosus, *Desf. ex DC. Prod.* vi. 635.—Reg. Caucas.

arenarius, Willd. Sp. Pl. iii. 1663 = Carduus arenarius.

Argemone, Lapeyr. Hist. Abr. Pl. Pyr. ii. 493 = medius.

argentatus, Roth, Catalect. i. 111 = Carduus argentatus.

argyracanthus, *C. B. Clarke, Comp. Ind.* 218.—Reg. Himal.

arisonicus, *A. Gray, in Proc. Am. Acad.* x. (1874) 44. —Arizona.

armatus, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 262.—Syria.

arvensis, *Hoffm. Deutschl. Fl.* ed. II. i. 11. 130.—Europ.; As. temp.

arvensis, Hook. Comp. Bot. Mag. i. (1835) 48 = virginianus.

asiaticus, *Sievers, in Pall. N. Nord. Beitr.* vii. 260. —Sibir. altaic.

auriculatus, Wall. Cat. n. 3889 = Saussurea hypoleuca.

autariscus, Chaix, in Vill. Hist. Pl. Dauph. i. 381; Willd. Sp. Pl. iii. 1676 = heterophyllus.

autariscus, Lobel. Fl. Gall. 541 = rivularis.

Bertolonii, *Bertol. Fl. Ital.* ix. 28.—Italia.

Boujarti, DC. Prod. vi. 635 = fimbriatus vel arachnoideus.

bracteatus, Cilib. Fl. Lituan. i. 184 = oleraceus.

Breweri, *A. Gray, in Proc. Am. Acad.* x. (1874) 43.—Calif.

Buergeri, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 502.—Japon.

californicus, *A. Gray, in Proc. Am. Acad.* x. (1874) 46.—Calif.

callosus, Roth, Catalect. iii. 76 = Carduus callosus.

candicans, Wall. Cat. n. 3893 = arvensis.

canus, *Roth, Tent. Fl. Germ.* i. 345.—Europ.

capitatus, *Kroch. Fl. Sil. Suppl.* ii. 290.—Europ.

carlinoides, *Fisch. ex Schrank, Pl. Rar. Hort. Monac.* t. 11.—Reg. Caucas.

carnolicus, *Willd. Sp. Pl.* iii. 1677.—Europ.

carthamoides, Wall. Cat. n. 3896 = Saussurea affinis.

carthamoides, Willd. Sp. Pl. iii. 1686 = Centaurea carthamoides.

CNICUS:—
Casabonae, *Roth, in Usteri, Ann. Bot.* viii. 9.—Ins. Ital.; Macrocos.

centauroides, *Linn. Sp. Pl.* 826.—Mont. Pyren.

centauroides, Willd. ex Ledeb. Fl. Ross. ii. 753 = Lemnea salina.

cernuus, A. Gray, in Proc. Am. Acad. x. (1874) 44 = subcoriaceus.

cernuus, Linn. Sp. Pl. 826 = Carduus cernuus.

cernuus, Wall. Cat. n. 3892 = Wallichii.

chamaecephalus, *Vahl, in Linnaea*, xxxix. (1875) 511. —Abyssin.

Chamaepence, *Desf. Arb.* i. 280.—Graecia; As. Min.

Chamaepence, Tenore, Prod. p. xlviii = gnaphalodes.

chinensis, *Benth. ex Maxim. in Bull. Acad. Pétersb.* xix. (1874) 509.—China.

chrysacanthus, *Ball, in Journ. Bot.* xi. (1873) 369.—Marocco.

ciliatus, *Roth, Catalect.* ii. 104.—Europ.

ciliatus, Samuela, iv. 447 = ochroleucus.

collinus, Hornem. Hort. Hafn. Suppl. 92 = Carduus collinus.

communis, Lam. Fl. Fr. ii. 25 = Cynara Cardunculus.

connexus, *Franch. & Sav. Enum. Pl. Jap.* ii. 409.—Japon.

conspicuus, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.

costaricensis, *Polak. in Linnaea*, xli. (1879) 561.—Costa Rica.

cyanoides, *Roth, in Usteri, Ann. Bot.* viii. 9.—Europ. *Cynara*, Lam. Fl. Fr. ii. 14 = Centaurea cynaroides.

cynaroides, Bieb. Fl. Taur. Cauc. ii. 280 (= *Cirsium sinuatum*).—Reg. Caucas.

cynaroides, Sibth. & Sm. Fl. Graec. ix. 20. t. 838 = eriophorus.

cynaroides, *Willd. Sp. Pl.* iii. 1670.—Ins. Creta.

dalidaeus, Wall. ex DC. Prod. vi. 541 = Saussurea deltoidea.

dentatus, Forsk. Fl. Aegypt. Arab. 217 = Kentrophyllum dentatum.

dentatus, Willd. Sp. Pl. iii. 1672 = pratensis.

Diacantha, *Desf. Cat. Hort. Par.* ed. III. 252.—As. Min.; Syria.

digynocephala, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 497.—Japon.

discolor, *Mushl. ex Willd. Sp. Pl.* iii. 1670.—Am. bor.

discolor, Roth, ex Steud. Nom. ed. II. i. 388, 388 = Carduus callosus.

dissectus, *Willd. Sp. Pl.* iii. 1665.—Europ.

Drummondii, *A. Gray, in Proc. Am. Acad.* x. (1874) 40.—Am. bor. occ.

dubius, Willd. Prod. Fl. Berl. 260 = acaulis.

Eatoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 56.—Am. bor. occ.

echinatus, *Willd. Sp. Pl.* iii. 1668.—Reg. Mediterr. occ.

echinocephalus, *Willd. l. c.* 1665.—Reg. Caucas.

edulis, *A. Gray, in Proc. Am. Acad.* x. (1874) 47.—Am. bor. occ.

effusus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 500.—Japon.

elongatus, Wall. ex C. B. Clarke, Comp. Ind. 239 = Tricholepis elongata.

eriocephalus, *A. Gray, in Proc. Am. Acad.* x. (1874) 46.—Am. bor. occ.

eriophorodes, *Hook. f. Fl. Brit. Ind.* iii. 363.—Reg. Himal.

eriophorus, C. B. Clarke, Comp. Ind. 217 = eriophoroides.

eriophorus, Juss. ex DC. Prod. vi. 635 = spathulatus.

eriophorus, *Roth, Tent. Fl. Germ.* i. 345.—Europ.; As. Min.

Erisithales, Linn. Sp. Pl. ed. II. 1157 = ochroleucus?

esculentus, *Sievers, in Pall. N. Nord. Beitr.* vii. 262.—Rossia.

Falconeri, *Hook. f. Fl. Brit. Ind.* iii. 363.—Reg. Himal.

ferox, *Linn. Mant.* i. 109; *Sibth. & Sm. Fl. Graec. Prod.* ii. 152.—Europ. or.; As. Min.

fimbriatus, *Bieb. Fl. Taur. Cauc.* ii. 276.—Reg. Caucas.

firmus, *Ÿ. & C. Presl, Delic. Prag.* 107.—Europ.

flavescens, Willd. Sp. Pl. iii. 1688 = Serratula flavescens.

flavispinus, *Ball, in Journ. Linn. Soc.* xvi. (1878) 522.—Hispan.; Marocco.

foliosus, *A. Gray, in Proc. Am. Acad.* x. (1874) 40.—Am. bor. occ.

CNICUS:—
fruticosus, *Desf. Arb.* i. 280.—Graecia.

Galactites, Loisel. Fl. Gall. ed. I. 538 = Galactites tomentosa.

gigantea, *Willd. Sp. Pl.* iii. 1671.—Reg. Mediterr.

glabor, Ell. Sketch. ii. 270 = pitcheilis.

glabor, Nutt. Gen. Am. ii. 129 = muticus?

glutinosus, Bigel. Fl. Bost. ed. II. 291 = muticus.

glutinosus, Gilib. Fl. Lituan. i. 185 = ochroleucus?

Gmelini, Spreng. Hist. Rei Herb. ii. 270 = acaulis.

gnaphalodes, *Willd. ex Spreng. Syst.* iii. 675.—Italia.

gossypinus, Wall. ex DC. Prod. vi. 541, 643 = Saussurea gossypiphora.

Gouanii, Willd. Sp. Pl. iii. 1665 = Carduus medius.

Grahami, *A. Gray, in Proc. Am. Acad.* xix. (1883) 57.—Arizona.

Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 363.—Reg. Himal.

grumosus, *Hoffmgg. & Link, Fl. Port.* ii. 196.—Lusitan.

Hallii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 56.—Am. bor. occ.

helenioides, Willd. Sp. Pl. iii. 1674 = heterophyllus.

heterolepis, A. Gray, in Proc. Am. Acad. x. (1874) 44 = subcoriaceus.

heteromallus, Steud. Nom. ed. II. i. 388 = Saussurea candicans.

heterophyllus, *Roth, Catalect.* i. 114.—Europ.

Hilgendorfi, *Franch. & Sav. Enum. Pl. Jap.* ii. 410. —Japon.

hispanicus, *Ball, in Journ. Linn. Soc.* xvi. (1878) 522. —Hispan.

Hookerianus, *A. Gray, in Proc. Am. Acad.* x. (1874) 46.—Am. bor. occ.

horridulus, *Pursh, Fl. Am. Sept.* ii. 507.—Am. bor.

horridus, Bertol. Amoen. Ital. 40 = Bertolonii.

horridus, *Bieb. Fl. Taur. Cauc.* ii. 279.—Reg. Caucas.

horridus, C. B. Clarke, Comp. Ind. 220 = falconeri.

horridus, Forsk. Fl. Aegypt. Arab. 217 = Cardopatium corymbosum.

hybridus, Schleich. Cat. Pl. Helv. ed. III. 12 = ochroleucus.

igniarius, *Benth. & Hook. f. Gen.* ii. 469.—Sibir. altaic. incomptus, *Franch. & Sav. Enum. Pl. Jap.* ii. 410.—Japon.

inermis, Willd. Sp. Pl. iii. 1672 = centauroides.

intermedius, *Heller, Pl. Wisc.* ii. 256.—Europ.

involucratus, *Wall. ex DC. Prod.* vi. 639.—Reg. Himal.

italicus, *Seb. & Mauri, Fl. Rom. Prodr.* ii. 282 — Italia.

japonicus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 503.—Japon.

jordanels, *H. B. & K. Nov. Gen. et Sp.* iv. 22.—Mexic.

kamtschaticus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 498.—Japon.; Alaska.

Kermesina, Willd. ex Ledeb. Fl. Ross. ii. 739 = lappaceus.

Kotzkyi, Sch. Bip. in Ind. Sem. Hort. Berol. (1857) App. 24 = Carbenia benedicta.

Lachenalii, *C. C. Gmel. Fl. Bad.* iii. 880.—Europ.

lactens, Schleich. Cat. Pl. Helv. ed. IV. 18 = palustris.

lanatus, Willd. Sp. Pl. iii. 1671 = arvensis.

lanceolatus, *Willd. Prod. Fl. Berl.* 259.—Europ.

lanifolius, *Bieb. Fl. Taur. Cauc.* ii. 276.—Reg. Caucas.

lappaceus, *Bieb. l. c.* 277.—Reg. Caucas.

lappaceus, Hort. Par. ex DC. Prod. vi. 639 = nivalis.

lappoides, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.

Lecontei, *A. Gray, in Proc. Am. Acad.* x. (1874) 39.—Am. bor.

leucocephalus, *Willd. Sp. Pl.* iii. 1668.—Creta; As. Min.

leucographus, Roth, in Usteri, Ann. Bot. viii. 8 = Tyrimnus leucographus.

leucophaeus, Baumg. ex Schur, Enum. Pl. Transs. 418 = lanceolatus.

lineatus, *Benth. & Hook. f. ex Franch. & Sav. Enum. Pl. Jap.* i. 261.—Japon.

lomatolepis, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.

longifolius, Lam. Fl. Fr. ii. 12 = Carduncellus mospelianus.

lucidus, Wall. ex DC. Prod. vi. 639 = Cardnus setans.

macrocephalus, *Moretti, in Brugnat. Giorn. Fis.* II. (1824) 162.—Italia.

CNICUS :—
medius, *Kroch. Fl. Sil. Suppl.* ii. 276.—Silesia.
mexicanus, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.
microcephalus, *Bertol. Fl. Ital.* ix. 36.—Ital.; Sardinia.
microcephalus, Boiss. Fl. Orient. iii. 706 = Carbenia benedicta.
mollmeresis, *Tineo, ex Ces. Passer. & Gib. Comp. Fl. Ital.* 488.—Ital.
mitis, *Hornem. Hort. Hafn. Suppl.* 98.—Sibir.
mitissimus, Lam. Fl. Fr. ii. 13 = Carduncellus mitissimus.
mollis, DC. Prod. vi. 652, sphalm. = acaulis.
mollis, Roth, Tent. Fl. Germ. i. 346 = Juncea mollis.
monspessulanus, *Koch, in Usteri, Ann. Bot.* viii. 9.—Gallia.
montanus, *Waldst. & Kit. ex Willd. Sp. Pl.* iii. 1676.—Europ.
multiceps, Wall. ex DC. Prod. vi. 540 = Saussurea affinis.
munitus, *Bieb. Fl. Taur. Cauc.* ii. 279.—Reg. Caucas.
muticus, *Ell. Sketch.*, ii. 268.—Am. bor.
neo-mexicanus, *A. Gray, in Proc. Am. Acad.* x. (1874) 45.—Am. bor. occ.
nipponicus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 496.—Japon.
nivalis, *H. B. & K. Nov. Gen. et Sp.* iv. 28.—Mexic.
niveus, *F. & C. Presl, Delic. Prag.* 108.—Europ.
niveus, Wall. Cat. n. 2898 = Saussurea deltoides.
nudiflorus, *Schleich. Cat. Pl. Helv.* ed. IV. 14, nomen.—Helvet.
Nuttallii, *A. Gray, in Proc. Am. Acad.* x. (1874) 41.—Am. bor.
obvallatus, *Bieb. Fl. Taur.-Cauc.* ii. 279.—Reg. Caucas.
oboallatus, Salzm. ex DC. Prod. vi. 660 = Notobasis syriaca.
occidentalis, *A. Gray, in Proc. Am. Acad.* x. (1874) 45.—Am. bor. occ.
ochroocentrus, *A. Gray, l. c.* xix. (1883) 57.—Am. bor. occ.
ochroleucus, *Spreng. Nachr.* i. *Bot. Gart. Halle*, 19.—Europ.
odoratus, Muhl. ex Steud. Nom. ed. II. i. 588 = pumilus.
oleraceus, *Linn. Sp. Pl.* 826.—Europ.
oligophyllus, *Franch. & Sav. Enum. Pl. Jap.* ii. 412.—Japon.
orgyalis, *Willd. Enum. Hort. Berol. Suppl.* 56.—Hab.?
orientalis, Willd. Sp. Pl. iii. 1670 = Galactites tomentosa.
ornatus, *Ball, in Journ. Bot.* xi. (1873) 868.—Marocco.
ovalifolius, *Franch. & Sav. Enum. Pl. Jap.* ii. 412.—Japon.
paludosus, *Loisel. Fl. Gall.* 542.—Europ.
palustris, Sibth. & Sm. Fl. Graec. Prod. ii. 151 = pungens.
palustris, *Willd. Prod. Fl. Berl.* 260.—Europ.; As. bor.
pannonicus, *Host, Fl. Austriac.* ii. 444.—Europ.
Parryi, *A. Gray, in Proc. Am. Acad.* x. (1874) 47.—Am. bor. occ.
parviflorus, *Heller, Fl. Wirc. Suppl.* 76.—Europ.
pauciflorus, *Waldst. & Kit. Pl. Rar. Hung.* ii. 176.—Europ. or.
pascuarensis, *H. B. & K. Nov. Gen. et Sp.* iv. 22. t. 310.—Mexic.
pectinellus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 494.—Japon.
pedemontanus, *Willd. Sp. Pl.* iii. 1675.—Europ.
pendulus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 510.—Sibir. or.
pexus, *Franch. & Sav. Enum. Pl. Jap.* ii. 410.—Japon.
pinnatifidus, Willd. Sp. Pl. iii. 1684 = Serratula pinnatifida.
Pitcheri, *Torr. in Eaton, Man.* ed. V. 180.—Am. bor.
polyacanthus, *Vathe, in Linnaea*, xxxix. (1875) 510.—Afr. trop.
polyanthemus, *Bertol. Amoen. Ital.* 41.—Italia.
Pontadorae, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 96 = rivularis.
praemorsus, Michl, in Flora, iii. (1820) 317 = ochroleucus ?
prostratus, Lam. Fl. Fr. ii. 14 = oleraceus.
pratensis, *Willd. Sp. Pl.* iii. 1679.—Europ.

CNICUS :—
pseudo-benedictus, Hort. Dorpat. ex Aschers. Ind. Sem. Hort. Berol. (1861) 5 = Carbenia benedicta.
pubigerus, *Desf. Cat. Hort. Par.* ed. III. 299.—As. Min.
pumilus, *Torr. Compend.* 282.—Am. bor.
pungens, Bivona, Sic. Pl. Cent. i. 26 = siculus.
pungens, Sebast. Fl. Rom. fasc. i. 9. t. 2 = polyanthemus.
pungens, *Willd. Sp. Pl.* iii. 1668.—Europ. or.; As. Min.
purpuratus, *Maxim. in Bull. Acad. Pétersb.* xix. (1879) 491.—Japon.
purpureus, Bertol. Fl. Ital. ix. 21 = spinosissimus.
pycnocephalus, Roth, in Usteri, Ann. Bot. viii. 8 = Carduus pycnocephalus ?
pygmaeus, Linn. Sp. Pl. ed. II. 1156 = Saussurea pygmaea.
pyrenaicus, *Willd. Sp. Pl.* iii. 1664.—Europ.
quercetorum, *A. Gray, in Proc. Am. Acad.* x. (1874) 44.—Calif.
radians, *Hemsl. Biol. Centr. Am. Bot.* ii. 252.—Guatemala.
rhaphilepis, *Hemsl. l. c.*—Mexic.
rectus, Hort. Par. ex DC. Prod. vi. 659 = stictus.
Reinii, *Franch. & Sav. Enum. Pl. Jap.* ii. 413.—Japon.
remotilobus, *A. Gray, in Proc. Am. Acad.* x. (1874) 47.—Am. bor. occ.
repandus, *Ell. Sketch*, ii. 269.—Am. bor.
replicatus, *Hornem. Hort. Hafn. Suppl.* 98.—Hungar.—Calif.
rigens, *Willd. Sp. Pl.* iii. 1675.—Europ.
rivularis, *Willd. l. c.* 1676.—Europ.
Rothrockii, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 220.—Arizona.
rufescens, *Loisel. Fl. Gall.* 543.—Gallia.
ruthenicus, Henning, in Mém. Soc. Nat. Mosc. vi. (1932) 78 = arvensis.
saanensis, Schlecht. ex DC. Prod. vi. 648 = ochroleucus.
salinus, Willd. ex Ledeb. Fl. Ross. ii. 752 = Leuzea serratuloides
salisburgensis, Willd. Sp. Pl. iii. 1675 = rivularis.
sammiticus, Tenore, ex Steud. Nom. ed. II. i. 589 = lanceolatus.
scariosus, *A. Gray, Syn. Fl. N. Am.* i. ii. 402.—Am. bor. occ.
Schimperi, *Vathe, in Linnaea*, xxxix. (1875) 511.—Afr. trop.
segetum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 511.—China.
seminudus, Schleich. ex DC. Prod. vi. 650 = rivularis.
semipectinatus, *Desf. Cat. Hort. Par.* ed. III. 151.—As. centr.
semipectinatus, *Hornem. & Ledeb. ex DC. Prod.* vi. 647.—Hab.?
semipectinatus, Thom. ex Steud. Nom. ed. II. i. 589 (= *Cirsium subalpinum*).—Europ.
serratuloides, *Roth, Tent. Fl. Germ.* i. 346.—Silseria.
serrulatus, *Bieb. Fl. Taur.-Cauc.* ii. 275.—Reg. Caucas.
setosus, Bess. Prim. Fl. Galic. ii. 172 = arvensis ?
siculus, Lam. Fl. Sic. Syn. ii. 442 = pungens.
Sieboldii, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 508.—Japon.
sinensis, C. B. Clarke, Comp. Ind. 219 = chinensis.
spathulatus, *Moretti, in Brugnat. Giorn. Fis.* II. v. (1822) 111.—Italia.
spicatus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 501.—Japon.
spinosissimus, Darling, Fl. Cestr. ed. II. 438 = horridulus.
spinosissimus, Habl. Tanr. 158, ex Bieb. Fl. Taur.-Cauc. iii. 261 = echinocephalus.
spinosissimus, Lapeyr. Hist. Abr. Pl. Pyr. ii. 496 = seq. ?
spinosissimus, *Linn. Sp. Pl.* 826.—Europ.
spurius, Hoffm. Deutschl. Fl. ed. II. ii 198 = pratensis.
squarrosus, *Hornem. Hort. Hafn.* ii. 782.—Sibiria.
stellatus, *Roth, in Usteri, Ann. Bot.* viii. 9.—Europ.
strictus, *Tenore, Prod.* p. xlviii.—Italia.
strigosus, *Bieb. Fl. Taur.-Cauc.* ii. 277.—Reg. Caucas.; Persia.
suboriaceus, *Hemsl. Biol. Centr. Am. Bot.* ii. 252.—Mexic.
suffultus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 498.—Japon.

CNICUS :—
syriacus, *Roth, in Usteri, Ann. Bot.* viii. 10.—Reg. Mediterr.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* ii. 411.—Japon.
tartaricus, *Willd. Sp. Pl.* iii. 1677.—As. temp.
tchefouensis, *Franch. in Mém Soc. Sc. Nat. Cherbourg*, xxiv. (1884) 229.—China.
transsylvanicus, *Schur, in Oestr. Bot. Zeitschr.* x. (1860) 179.—Transsylvan.
tuberosus, *Roth, Tent. Fl. Germ.* i. 345.—Europ.
uliginosus, *Bieb. Fl. Taur.-Cauc.* ii. 274.—Reg Caucas.
undulatus, *A Gray, in Proc. Am. Acad.* x. (1874) 42.—Am. bor.
uniflorus, Linn. Mant. ii. 572 = Centaurea monanthos
uniflorus, Sievers, in Pall. N. Nord. Beitr. vii. 346 = Leuzea carthamoides.
Vervutus, D. Don, Prod. Fl. Nep. 167 = argyracanthus.
virginianus, Hook. Comp. Bot. Mag. i. (1836) 48 = Lecontei.
virginianus, *Pursh, Fl. Am. Sept.* ii. 506.—Am. bor.
Vlassovianus, *Maxim. in Bull. Acad. Pétersb.* xix (1874) 509.—As. temp.
Wallichii, *Hook. f. Fl. Brit. Ind.* iii. 363.—Reg. Himal.
Weyrichii, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 495.—Jap. Sachalin.
Wheeleri, *A. Gray, in Proc. Am. Acad.* xix. (1883) 56.—Arizona.
Wildenowii, *F. G. Dieter. Vollst. Lexik. Gaertn. Nachtr* ii. 231.—Hab.?
Wrightii, *A. Gray, in Proc. Am. Acad.* x. (1874) 41.—Texas; Arizona.
yezoensis, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 508.—Japon.

CNICUS, [Vaill.] Gaertn. Fruct. ii. 385. t. 162 (1791) = Carbenia, Adans. (Compos.).
benedictus, Linn. Sp. Pl. 826; Gaertn. l. c. = Carbenia benedicta.

CNIDIUM, Cusson, in Mém. Soc. Méd. Paris (1782) 280 = Bellnum, Linn. (Umbell.).
affine, C. A. Mey. in Beitr. Pfl. Russ. Reich. vi. 88.—Reg. Caucas.
alatum, Spreng. ex Link, Handb. i. 329 = Lugasticum alatum.
alsaticum, Spreng. Pl. Umbellif. Prod. 40 = Peucedanum alsaticum.
anomalum, Ledeb. Fl. Alt. i. 330; Ic. Fl. Ross. t. 311.—Sibir. altaic.
apioides, Spreng. Pl. Umbellif. Prod. 40 = Carum rigidulum.
argenteum, Cesati, in Linnaea, xi. (1837) 324 = Peucedanum involucratum ?
athoum, Griseb. Spicil. Fl. Rumel. i. 363 = Carum rigidulum.
atropurpureum, Spreng. in Roem. & Schult. Syst. vi. 418 = Thaspium trifoliatum.
canadense, Spreng. l. c. 415 = S. canadense.
carvifolium, Bieb. Fl. Taur.-Cauc. iii. 212 = Carum caucasicum.
cenolophioides, Benth. & Hook. f. Gen. i. 914 = S. sylvestre.
chinense, Spreng. ex Steud. Nom. ed. II. i. 389 = S. canadense.
confertum, Moench, Meth. 98 = S. Monnieri.
conisifolium, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 299.—As. Min.
cuneatum, Ledeb. Fl. Alt. i. 331; et Ic. Fl. Ross. t. 312.—Sibir. altaic.
cuneifolium, Griseb. Spicil. Fl. Rumel. i. 363, in obs. = Ligusticum cuneifolium.
dahuricum, Fisch. & Mey. in Ann. Sc. Nat. Sér. II. v. (1836) 186.—Dahuria.
diffusum, DC. Prod. iv. 158 = Seseli indicum.
divaricatum, Ledeb. Fl. Alt. i. 324 = Silaus longifolius.
Fischeri, Spreng. Nov. Prov. 12 = Silaus longifolius.
Fontanesii, Spreng. Sp. Umbellif. Prod. 41.—Numidia.
intermedium, Tchihat. Asie Min. Bot. i. 430.—As. Min.
japonicum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 60.—Japon.
Krauseianum, Sond. in Harv. & Sond. Fl. Cap. ii. 552 = Pimpinella caffra.

CNIDIUM :—
meifolium, Bieb. Fl. Taur. Casc. iii. 213 = Carum asperum.
microcarpum, Turcz. ex Steud. Nom. ed. II. i. 389.—Sibir. Baical.
Monnieri, Cusson, in Mém. Soc. Méd. Par. (1782) 280 = Selinum Monnieri.
multicaule, Ledeb. Fl. Ross. ii. 284.—Sibiria.
myrrhidifolium, Bieb. Fl. Taur. Casc. iii. 212 = Ligusticum alatum.
orientale, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 299 = S. orientale.
palustre, Reichb. Fl. Germ. Excurs. 469 = S. sylvestre.
Petroselinum, DC. Prod. iv. 152 = Carum Petroselinum.
peucedanoides, H. B. & K. Nov. Gen. et Sp. v. 15 = Eulophus peucedanoides.
pyrenæum, Spreng. in Roem. & Schult. Syst. vi. 416 = Ligusticum pyrenæum.
selinum, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 783.—Dahuria.
Silaus, Bieb. Fl. Taur. Casc. iii. 212 = Silaus Besseri.
Silaus, Spreng. Pl. Umbellif. Prod. 40 = Silaus pratensis.
striatum, Turcz. ex Fisch. & Mey. in Ann. Sc. Nat. Sér. II. v. (1836) 188 = *dahuricum*.
suffruticosum, Cham. & Schlecht. in Linnæa, i. (1826) 387 = S. suffruticosum.
tenuifolium, Moench, Meth. 98 = Apium leptophyllum.
toluense, Spreng. Syst. i. 888 = Arracacia toluensis.
venosum, Koch, Umbellif. 109 = S. sylvestre.

CNIDONE, E. Mey. ex Endl. Gen. Suppl. ii. 76 (1842) = **Kissenia**, R. Br. (Lossac.).
muricatissima, E. Mey. ex Presl, Bot. Bemerk. 78 = K. spathulata.

CNIDOSCOLUS, Pohl, Pl. Bras. Ic. i. 56. t. 49 (1827) = **Jatropha**, Linn. (Euphorbiac.).
angustidens, Torr. Bot. Mex. Bound. 198 = J. angustidens.
cnicodendron, Griseb. in Goett. Abh. xxiv. (1879) 58. —Reg. Argent.
fragrans, Pohl, Pl. Bras. i. 63 = J. fragrans.
hamosus, Pohl, l. c. 57 = J. hamosa.
lobatus, Pohl, l. c. 62 = J. phyllacantha.
Marcgravii, Pohl, l. c. 58. t. 50 = J. urens.
Michauxii, Pohl, l. c. 58 = J. urens.
napæifolius, Pohl, l. c. 62 = J. aconitifolia.
neglectus, Pohl, l. c. 60. t. 51 = J. urens.
obtusifolius, Pohl, ex Baill. in Adansonia, iv. (1863–64) 275 = J. obtusifolia.
osteocarpus, Pohl, Pl. Bras. i. 63 = J. osteocarpa.
palmatus, Pohl, l. c. = J. palmata.
pubescens, Pohl, l. c. 62 = J. obtusifolia.
quercifolius, Pohl, ex Baill. in Adansonia, iv. (1863–64) 276 = J. phyllacantha.
quinquelobus, Pohl, Pl. Bras. i. 63 = J. urens.
repandus, Pohl, l. c. 62 = J. phyllacantha.
stimulosus, Ræusm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1843) 324 = J. texana.
stimulosus, A. Gray, Man. Bot. 389 = J. urens.
surinamensis, Miq. in Linnæa, xvii. (1844) 749 = Croton lobatus.
vitifolius, Pohl, Pl. Bras. i. 61 = J. vitifolia.

CNOPOS, Raf. Fl. Tellur. iii. 18 (1836) = **Polygonum**, Tourn.
ramosissimum, Raf. l. c. = P. ramosissimum.

COA, Adans. Fam. ii. 445 (1763) = **Hippocratea**, Linn. (Celastr.).

COALISIA, Raf. Sylva Tellur. 114 (1838) = **Cleome**, Linn. (Capparid.).
angustifolia, Raf. l. c. = Cleome angustifolia.

COATESIA, F. Muell. Fragm. iii. 26 (1862) = **Goetzea**, Schott (Rutac.).
paniculata, F. Muell. l. c. = G. Muelleri.

COBÆA, Cav. Ic. i. 11. t. 16, 17 (1791). *POLEMONIACEÆ*, Benth. & Hook. f. ii. 894.
ROSENBERGIA, Cent. in Kjob. Vidensk. Meddel. (1856) 30.
grandiflora, *Hemsl. in The Garden*, xvii. (1880) 329.—Costa Rica.

COBÆA :—
lutea, D. Don, in Edinb. Phil. Journ. x. (1824) 4 = macrostemma.
macrostemma, *Pav. ex D. Don, l. c.* 112.—Guatemala.
minor, *Mart. & Gal. in Bull. Acad. Brux.* xii. (1845) 11. 376.—Mexic.
pendulifora, *Hook. f. Bot. Mag.* t. 5757.—Venezuela.
scandens, *Cav. Ic.* i. 11. t. 16, 17.—Mexic.
stipularis, *Benth. Pl. Hartw.* 45.—Mexic.

COBAEA, Neck. Elem. i. 129 (1790) = **Lonicera**, Linn. (Caprifol.).

COBAMBA, Blanco, Fl. Filip. ed. I. 510 (1837) = **Cansoora**, Lam. (Gentian.).
Blanci, Blanco, l. c. ed. II. Suppl. 591 = Cans. decussata.
dichotoma, Blanco, l. c. ed. I. 510 = Cans. diffusa.

COBRESIA, Pers. Syn. ii. 534 (1807) = **Kobresia**, Willd. (Cyperac.).

COBURGIA, Sweet, Brit. Flow. Gard. Ser. II. t. 17 (1838) = **Stenomesson**, Herb. (Amaryllid.).
acuta, Herb. in Bot. Mag. t. 3865 = S. incarnatum.
angusta, Herb. in Bot. Reg. (1842) Misc. 55 = S. recurvatum.
Belladonna, Herb. in Bot. Mag. sub t. 2113, 2114 = Amaryllis Belladonna.
blanda, Herb. l. c. = Amaryllis Belladonna.
chachapoyensis, Herb. in Bot. Reg. (1842) Misc. 53 = S. incarnatum.
ciliaris, Herb. in Trans. Hort. Soc. iv. (1822) 181 = Buphane disticha.
coccinea, Herb. in Bot. Reg. (1842) Misc. 54 = S. coccineum.
discolor, Herb. Amaryll. 197 = S. recurvatum.
flava, Hort. Belg. ex Heynh. Nom. ii. 151 = seq.
fulva, Herb. in Bot. Reg. t. 1497 = S. incarnatum.
humilis, Herb. l. c. (1842) Misc. 55 = S. humile.
incarnata, Sweet, Brit. Flow. Gard. Ser. II. t. 17 = S. incarnatum.
Josephinæ, Herb. in Trans. Hort. Soc. iv. (1822) 181 = Brunsvigia Josephinæ.
lacta, Herb. in Bot. Reg. (1842) Misc. 53 = S. incarnatum.
longæva, Herb. l. c. = S. incarnatum.
lutea, Herb. l. c. 55 = S. recurvatum.
Macleanica, Herb. l. c. = S. recurvatum.
miniata, Lindl. Bot. Reg. (1844) Misc. 13.—Peruv.
multiflora, Herb. in Bot. Mag. sub tt. 2113, 2114 = Brunsvigia multiflora.
obvallaris, Herb. in Bot. Reg. (1842) Misc. 54 = S. recurvatum.
pallida, Herb. in Trans. Hort. Soc. iv. (1822) 181 = Amaryllis Belladonna.
præcipitata, Herb. in Bot. Reg. (1842) Misc. 54 = S. coccineum.
pudica, Herb. in Bot. Mag. sub tt. 2113, 2114 = Amaryllis Belladonna.
quitensis, M. Roem. Syn. Enest. 165.—Ecuador.
Radula, Herb. ex Steud. Nom. ed. II. i. 390 = Brunsvigia Radula.
recurvata, Herb. in Bot. Reg. (1842) Misc. 54 = S. recurvatum.
reticulata, Herb. App. 34 = Hippeastrum reticulatum.
splendens, Herb. Amaryll. 400. t. 47. f. 5 = S. incarnatum.
striatifolia, Herb. App. 34 = Hippeastrum reticulatum.
stylosa, Lindl. in Journ. Hort. Soc. ii. (1847) 210.—Peruv.
taeniophylla, Salisb. Gen. Pl. Fragm. 134.—Hab.?
trichroma, Herb. Amaryll. 196 = S. incarnatum.
variegata, Herb. l. c. = S. incarnatum.
venusta, Herb. in Bot. Reg. (1842) Misc. 54 = S. coccineum.
versicolor, Herb. l. c. 53. t. 66 = S. incarnatum.

COCCANTHERA, C. Koch & Hanst. Ind. Sem. Hort. Berol. (1855) 17 = **Cocconanthe**, Hanst. (Gesneriac.).
Descoingsii, C. Koch & Hanst. l. c. = Cocl. Devosiana.
Hookeriana, C. Koch & Hanst. l. c. = Cocl. Hookeriii.

COCCINIA, Wight & Arn. Prod. 347 (1834). *CUCURBITACEÆ*, Benth. & Hook. f. i. 827.
CEPHALANDRA, Schrad. in Eckl. & Zeyh. Enum. ii. 280 (1836).
abyssinica, *Cogn. in DC. Monog. Phan.* iii. 536.—Abyssin.
adolfensis, *Cogn. l. c.* 538.—Afr. trop. et austr.
cordifolia, Cogn. l. c. 529 = indica.
decipiens, *Cogn. l. c.* 530.—Afr. trop.
discrepiflora, Cogn. l. c. 536 = C. Hartmanniana.
grandis, Voigt, Hort. Suburb. Calc. 59 = indica.
Hartmanniana, *Schweinf. Reliq. Kotschy.* 93. t. 27.—Abyssin.
indica, *Wight & Arn. Prod.* 347.—As. trop.
jatrophæfolia, *Cogn. in DC. Monog. Phan.* iii. 533.—Abyssin.
Leuweriana, M. Roem. Syn. Pepon. 93 = indica.
Mac-Kennii, *Cogn. in DC. Monog. Phan.* iii. 541.—Afr. austr.
Maghadi, Aschers. in Schweinf. Beitr. Fl. Aethiop. 267 = C. Motmoi.
Motmoi, *M. Roem. Syn. Pepon.* 93.—Afr. trop.; Arabia.
Monteiri, *Hort.*—*Cf. Cat. Gr. Yard. Bot. Bordeaux* (1866).—Afr. trop.?
palmata, *Cogn. in DC. Monog. Phan.* iii. 540.—Afr. austr.
palmata, M. Roem. Syn. Pepon. 93 = Bryonopsis lacinicea.
palmatisecta, *Kotschy, in Sitzb. Acad. Wien, Math. Nat.* ii. (1865) 300. t. 4.—Abyssin.
quinqueloba, *Cogn. in DC. Monog. Phan.* iii. 533.—Afr. austr.
Schimperi, Naud. in Ann. Sc. Nat. Sér. IV. viii. (1857) 366 = C. Motmoi.
senensis, *Cogn. in DC. Monog. Phan.* iii. 535.—Afr. trop.
sessilifolia, *Cogn. l. c.* 534.—Afr. austr.
Wightiana, M. Roem. Syn. Pepon. 93 = indica.

COCCOBRYON, Klotzsch, in Miq. Syst. Pip. 843 (1843) = **Piper**, Linn.
capense, Klotzsch, l. c. = P. capensis, radicans.

COCCOCERAS, Miq. Fl. Ind. Bat. Suppl. 455 (1860). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 318.
muticum, *Muell. Arg. in Flora*, xlvii. (1864) 470.—Malaya.
plicatum, *Muell. Arg.* l. c. 539.—Burma.
samatranum, *Miq. Fl. Ind. Bat. Suppl.* 456.—Sumatra.

COCCOCYPSELUM, P. Br. Hist. Jamaic. 144 (1756) (Coccocipalum). *RUBIACEÆ*, Benth. & Hook. f. ii. 73.
BELLARDIA, Schreb. Gen. i. 790 (1789).
CONDALIA, Rnia & Pav. Prod. 11. t. 2 (1794).
LITOSTOMA, D. Don, in Edinb. Phil. Journ. (1830) i. 168.
SICELIUM, P. Br. Hist. Jamaic. 144 (1756).
TONTANEA, Aubl. Pl. Gui. i. 108. t. 43 (1775).
aureum, *Cham. & Schlecht. in Linnæa*, iv. (1829) 139.—Bras.
bifarium, Willd. Sp. Pl. i. 618 = Manettia Lygistum.
buxifolium, Spreng. Syst. i. 416 = Fernelia buxifolia.
campanaliflorum, *Cham. & Schlecht. in Linnæa*, iv. (1829) 140.—Bras.
canescens, *Willd. ex Roem. & Schult. Syst. iii. Mant.* 190.—Am. austr.
capitatum, Willd. ex DC. Prod. iv. 396 = umbellatum.
ciliatum, *Cham. & Schlecht. in Linnæa*, vi. (1831) 414.—Mexic.
Condalia, Pers. Syn. i. 183 = umbellatum.
cordifolium, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 14.—Bras.
dichroolasium, *Mart. in Flora*, xxiv. (1841) 11. *Beibl.* 76.—Bras.
discolor, *Van Houtte, ex Kunze, in Linnæa*, xxiv. (1851) 192.—Bras.?
erythrocospalum, *Cham. & Schlecht. in Linnæa*, iv. (1829) 144.—Bras.
gaophilotdes, *Wawra, in Oestr. Bot. Zeitschr.* xxxi. (1881) 70.—Bras.
glabrum, *Benth. ex DC. Prod.* iv. 397.—Panama.
herbaceum, Aubl. Pl. Gui. i. 98 = repens.
hirsutum, *Benth. ex DC. Prod.* iv. 396.—Hab.?
lanceolatum, *Pers. Syn.* i. 183.—Peruv.

COCCOCYPSELUM:—
mentalicum, *Linden, in Belg. Hortic.* xv. (1888) 99.—Guiana.
microphyllum, *Willd. ex Roem. & Schult. Syst.* iii. *Mant.* 120.—Ins. Mauriti.
montanum, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 76.—Bras.
nummularifolium, *Cham. & Schlecht. in Linnaea*, iv. (1829) 145.—Bras.; Ind. occ.; Mexic.
obovatum, *Pers. Syn.* i. 183.—Peruv.
ovatum, *Cham. & Schlecht. in Linnaea*, iv. (1829) 141.—Bras.
pedunculare, *Cham. & Schlecht. l. c.* 142.—Bras.
Pseudo-Tentanea, *Griseb. Fl. Brit. W. Ind.* 932.—Ind. occ.
pulchellum, *Cham. in Linnaea*, ix. (1834) 288.—Bras.
repens, H. B. & K. Nov. Gen. et Sp. iii. 408 = canescens.
repens, *Sw. Prod. Veg. Ind. Occ.* 81.—Ind. occ.
sessile, *Pers. Syn.* i. 193.—Peruv.
spicatum, H. B. & K. Nov. Gen. et Sp. iii. 406 = Gonzales spicata.
Tentanea, *H. B. & K. l. c.*—Am. austr.
umbellatum, *Poir. Encyc. Suppl.* ii. 807.—Peruv.
uniflorum, Willd. Sp. Pl. i. 618 = Fernelia obovata veronicoides, *Presl, Bot. Bemerk.* 87.—Am. trop.
virgatum, *Lam. Illustr.* i. 270. n. 1457 = Manettia picta.
violaceum, Desf. Cat. Hort. Par. ed. III. 404 = capitulifloruin.

COCCODERMA, Miers, in Lindl. Veg. Kingd. 309 b (1847), nomen (Menispermn.).

COCCOLOBA, Linn. Syst. ed. X. 1007 (1759). *POLYGONACEAE*, Benth. & Hook. f. iii. 102.
LYPERODENDRON, Willd. ex Meisn. in DC. Prod. xiv. 146, in syn. (1857).
NAUCORRHES, Raïn. Fl. Tellur. ii. 84 (1836).
SCHLOSSERIA, Mill. ex Steud. Nom. ed. II. ii. 531 (1841).
acrostichoides, *Cham. in Linnaea*, viii. (1833) 132.—Bras.
acuminata, *H. B. & K. Nov. Gen. et Sp.* ii. 176.—N. Granat.
alagoënsis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 260.—Bras.
alnifolia, Casar. Nov. Stirp. Bras. Dec. viii. 71 = populifolia.
appressa, Meisn. ex Steud. Nom. ed. II. i. 390 = Polygonum appressum.
armata, *Wright, ex Griseb. Cat. Pl. Cub.* 288.—Cuba.
asiatica, *Lour.* Fl. Cochinch. 289 = Polygonum asiaticum.
australis, Forst. f. Prod. 29. n. 176 = Muehlenbeckia australis.
barbadensis, *Jacq. Enum. Pl. Carib.* 87; *Linn. Syst.* ed. XII. 277.—Am. trop.
barbadensis, Willd. Sp. Pl. ii. 459 = caracasana.
Blanchetiana, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 257 = ochreolata.
brasiliensis, *Meisn. in Mart. Fl. Bras.* v. i. 80.—Bras.
brasiliensis, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) i. 80.—Bras.
brasiliensis, Spreng. Syst. ii. 262 (= Hedyosmi sp.).—Bras.
bullata, Hort. ex Meisn. in DC. Prod. xiv. 152 = rugosa.
calobotrya, Meisn. l. c. 157 = coriacea.
Candolleana, *Meisn. in Mart. Fl. Bras.* v. i. 41.—Bras.
caracasana, *Meisn. in DC. Prod.* xiv. 137.—Venezuela.
coriacea, Ruiz, ex Meisn. l. c. 150 = Muehlenbeckia leptobotrya.
costata, *Cham. in Linnaea*, viii. (1833) 132.—Bras.
cordifolia, Meisn. in Mart. Fl. Bras. v. i. 37 = laevis.
coriacea, *A. Rich. Fl. Cub. Faxergr.* ii. 184.—Cuba.
coriacea, Willd. ex Meisn. in DC. Prod. xiv. 155 = obovata.
coronata, *Jacq. Enum. Pl. Carib.* 19.—Ind. occ.
cuneata, Mart. ex Meisn. in DC. Prod. xiv. 153 = populifolia.
cubata, *Wright, in Sauv. Fl. Cub.* 189.—Cuba.
cymosinflorida, *Cham. in Linnaea*, viii. (1833) 134.—Bras.
cymosa, Roxb. Fl. Ind. ii. 392 = Polygonum chinense.

COCCOLOBA:—
Cruegerii, *Griseb. Fl. Brit. W. Ind.* 710.—Ind. occ.
cubensis, *Meisn. in DC. Prod.* xiv. 139.—Cuba.
cujabensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 259.—Bras.
cymosa, Lour. Fl. Cochinch. 240 = Polygonum cymosum.
declinata, *Mart. ex Meisn. in Mart. Fl. Bras.* v. i. 80.—Bras.
densiflora, *Mart. ex Meisn. l. c.* 26.—Bras.
dioica, Sieud. Nom. ed. II. i. 390 = Muehlenbeckia sagittifolia.
diversifolia, *Jacq. Enum. Pl. Carib.* 19.—Ins. S. Doming.
emarginata, *Jacq. l. c.* 87 (sp. dub.).—Am. austr.
excelsa, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 694.—Am. trop.
excoriata, *Linn. Syst.* ed. X. 1007.—Ind. occ.
fagifolia, *Jacq. Hort. Schoenb.* iii. 55. t. 359.—Am. austr.
fasciculata, Meisn. in Kjoeb. Vidensk. Meddel. (1870) 138 = longependula.
fasciculata, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 255 = crescentiaefolia.
fastigiata, *Meisn. in Mart. Fl. Bras.* v. i. 34.—Bras.
ferruginea, Endl. Cat. i. 274 = excoriata.
flavescens, *Jacq. Enum. Pl. Carib.* 19.—Ins. S. Doming.
floridana, Meisn. in DC. Prod. xiv. 165 = laurifolia.
Gardneri, Meisn. in Mart. Fl. Bras. v. i. 36 = crescentiaefolia.
Gaudichiana, Meisn. l. c. 35 = fastigiata.
Gaudotiana, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 260 = obovata.
gracilis, *H. B. & K. Nov. Gen. et Sp.* ii. 176.—Peruv.
grandifolia, *Jacq. Enum. Pl. Carib.* 19.—Am. trop.
grandis, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 694 = latifolia.
guatemalensis, *Teijsm. & Binn. Cat. Hort. Bog.* 383, nomen.—Hab.?
guianensis, *Meisn. in Linnaea*, xxi. (1848) 254.—Guiana.
Humboldtii, *Meisn. in DC. Prod.* xiv. 168.—Bras.
ilbesmais, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 256.—Bras.
indica, Wight, ex Meisn. in DC. Prod. xiv. 131 = Polygonum chinense.
japurana, Meisn. in Mart. Fl. Bras. v. i. 25 = (Ancheles sp.).—Bras.
Klotzschiana, *Meisn. in DC. Prod.* xiv. 155.—Ins. S. Thomas.
Kunthiana, Meisn. l. c. 166 = pyrifolia.
laevis, *Casar. Nov. Stirp. Bras. Dec.* viii. 71.—Bras.
latifolia, *Lam. Encyc.* vi. 61.—Am. austr.
latifolia, Poepp. ex Meisn. in DC. Prod. xiv. 151 = polystachya.
laurifolia, *Jacq. Hort. Schoenbr.* iii. 9. t. 267.—Am. trop.
laxiflora, Desf. Cat. Hort. Par. ed. III. 46 = latifolia.
leptostachya, *Benth. Bot. Voy. Sulph.* 159.—Am. austr.
longependula, *Mart. ex Meisn. in Mart. Fl. Bras.* v. i. 27.—Bras.
longifolia, *Flach. Hort. Gorensk.* 25.—Ind. occ.
luciduda, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 697.—Guiana.
macrantha, Hook. Bot. Mag. sub t. 4536 = sec.
macrophylla, Desf. Tabl. ed. I. 88 = rugosa.
magnifolia, Hort. ex Meisn. in DC. Prod. xiv. 152 = rugosa.
maximiliensis, *Bouché in Vet. Akad. Handl. Stockh.* 1854 (1856) 149.—Panama.
margina, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 696.—Bras.
Martii, Meisn. in Mart. Fl. Bras. v. i. 37 = nitida.
membranacea, Klotzsch, in Linnaea, xiv. (1841) 389, nomen = ilbesmais.
microneura, *Meisn. in DC. Prod.* xiv. 168.—Demerara.
microphylla, *Griseb. Cat. Pl. Cub.* 62.—Cuba.
microstachya, *Willd. Sp. Pl.* ii. 459.—Ind. occ.
mollis, Casar. Nov. Stirp. Bras. Dec. viii. 72 = polystachya.
monoica, Ruiz, ex Meisn. in DC. Prod. xiv. 149 = Muehlenbeckia tamnifolia.
Moritzii, Klotzsch, ex Meisn. in Mart. Fl. Bras. v. i. 28 = ovata.

COCCOLOBA:—
nitida, Desf. Tabl. ed. II. 46 = fagifolia.
nitida, *H. B. & K. Nov. Gen. et Sp.* ii. 176.—N. Granat.
nitida, Mart. ex Meisn. in DC. Prod. xiv. 164 = ovata.
nitens, Jacq. Enum. Pl. Carib. 19 = excoriata.
nutans, *H. B. & K. Nov. Gen. & Sp.* ii. 176.—Peruv.
nymphaeifolia, Linden, Cat. n. 18 (1858) = peltata.
obovata, *H. B. & K. Nov. Gen. et Sp.* ii. 175.—N. Granat.
obtusifolia, Jacq. Enum. Pl. Carib. 19 = microstachya.
ochreolata, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii (1849) 259.—Am. austr.
orbicularis, Lodd. Cat. ex Loud. Hort. Brit. 159 = Muehlenbeckia orbicularis.
ovata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 697.—Guiana.
padiformis, *Meisn. in DC. Prod.* xiv. 166.—Venezuela.
pallida, *Wright, ex Griseb. Cat. Pl. Cub.* 61.—Cuba.
paniculata, Meisn. in Mart. Fl. Bras. v. i. 43 = polystachya.
parensis, *Meisn. l. c.* 38.—Bras.
parvimontii, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 696 = excelsa.
parvifolia, *Nutt. N. Am. Sylv.* iii. 95. t. 89 (sp. dub.).—Am. bor.
parvifolia, Poir. Encyc. vi. 64 = microstachya.
peltata, *Schott, in Spreng. Syst. iv. Curr. Post.* 605.—Bras.
peltigera, Meisn. in Mart. Fl. Bras. v. i. 39 = peltata.
pendula, Salzm. ex Meisn. in DC. Prod. xiv. 159 = nitida.
Persicaria, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 256.—Bolivia.
peruviana, Willd. ex Meisn. in DC. Prod. xiv. 163 = gracilis.
piperifolia, *Mart. ex Meisn. l. c.* 161.—Bras.
plantaginea, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 257.—Bras.
platyclada, F. Muell. ex Hook. Bot. Mag. t. 5883 = Muehlenbeckia platyclada.
Plumieri, *Griseb. Fl. Brit. W. Ind.* 162.—Ind. occ.
polystachya, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 261.—Bras.
populifolia, *Wedd. l. c.* 257.—Bras.
pubescens, Linn. Syst. ed. X. 1007 = grandifolia.
punctata, Linn. Sp. Pl. ed. II. 523 = coronata.
pyrifolia, *Desf. Tabl.* ed. II. 46.—Ind. occ.
racemulosa, *Meisn. in Mart. Fl. Bras.* v. i. 30.—Bras.
ramosissima, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 258.—Bras.
retusa, *Griseb. Cat. Pl. Cub.* 61.—Ind. occ.
rhofolia, Desf. Tabl. ed. II. 46 = latifolia.
rigida, Meisn. in Mart. Fl. Bras. v. i. 29 = parvifolia.
rigida, Willd. ex Meisn. in DC. Prod. xiv. 163 = Humboldtii.
roses, *Meisn. in Mart. Fl. Bras.* v. i. 33.—Bras.
rotundifolia, *Meisn. in DC. Prod.* xiv. 154.—Ins. S. Doming.
rubescens, Linn. Sp. Pl. ed. II. 523 = grandifolia.
rubiginosa, Mart. ex Meisn. in DC. Prod. xiv. 160 = acrostichoides.
rufescens, *Wright, in Sauv. Fl. Cub.* 188.—Cuba.
rugosa, *Desf. Cat. Hort. Par.* ed. III. 389.—Ind. occ.
sagittata, Poir. Encyc. vi. 64 = Muehlenbeckia sagittifolia.
sagittifolia, Ortega, Hort. Matr. v. 80 = Muehlenbeckia sagittifolia.
salicifolia, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 259.—Bras.
scandens, Casar. Nov. Stirp. Bras. Dec. viii. 70 = sticticaulis.
scandens *Forsyth. ex Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 695.—Ind. occ.
scandens, Poepp. ex Meisn. in DC. Prod. xiv. 154 = peltata.
Schomburgkiana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 698.—Guiana.
Schomburgkii, *Meisn. in Linnaea*, xxi. (1848) 265.—Guiana.

COCCOLOBA:—
sticticaulis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 361.—Bras.
striata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 626.—Bras.
stricta, Klotzsch, in Schomb. Faun. et Fl. Guian. 934 = laxiflda.
stroblulifera, *Meisn. in Mart. Fl. Bras.* v. 1. 95.—N. Granat.
Swartzii, *Meisn. in DC. Prod.* xiv. 159.—Ind. occ.
tenuifolia, Grisb. Fl. Brit. W. Ind. 163 = retusa.
tenuifolia, *Linn. Syst.* ed. X. 1007; *Amoen. Acad.* v. 397 (sp. dub.).—Jamaic.
Tetrace, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 74 = Polygonum molle.
uvifera, *Linn. Syst.* ed. X. 1007.—Am. trop.
Vellosiana, Casar. Nov. Stirp. Bras. Dec. viii. 70 = crassensiefolia.
venosa, Linn. Syst. ed. X. 1007 = coronata.
virens, Lindl. Bot. Reg. t. 1816 = coronata.
Warmingii, *Meisn. in Kjoeb. Vidensk. Meddel.* (1870) 128.—Bras.
Zebra, *Griseb. Fl. Brit. W. Ind.* 163.—Jamaic.

COCCOLOBIS, P. Br. Hist. Jamaic. 209 (1756) = Coccoloba, Linn.

COCCOMELIA, Reinw. Cat. Gew. Buitenz. 110 (1823); et in Syll. Ratisb. ii. (1828) 5 = Baccaurea, Lour. (Euphorbiac.).
racemosa, Reinw. ll. cc. 110, 6 = B. racemosa.

COCCONERION, Baill. Adansonia, xi. (1878) 87. *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 323.
Balansae, *Baill. l. c.* 88.—N. Caled.
minus, *Baill. l. c.*—N. Caled.

COCCOS, Gaertn. Fruct. i. 15 (1788) = Cocos, Linn. (Palm.).

COCCOSIPSILUM, Sw. Prod. Veg. Ind. Occ. 31 (1788) = Coccocypselum, P. Br. (Rubiac.).

COCCOSPERMA, Klotzsch, in Linnaea, xii. (1838) 215 = Balanza, Salisb. (Ericac.).
Forbesianum, Klotzsch, l. c. = S. hexandra.
rugosum, Klotzsch, l. c. = S. rugosa.

COCCULIDIUM, Spach, Hist. Vég. Phan. viii. 16 (1839) = Cocculus, DC. (Menisp.).
populifolium, Spach, l. c. 17 = Cocculus carolinus.

COCCULUS, DC. Syst. i. 515 (1818). *MENISPERMACEÆ*, Benth. & Hook. f. i. 86.
ADENOCHETON, Fenzl, in Flora, xxv. (1842) 312.
ANDROPHYLAX, Wendl. Beobacht. 38 (1798).
BAUMGARTIA, Moench, Meth. 650 (1794).
CERATHA, Forsk. Fl. Aegypt. Arab. 171 (1775).
COCCULIDIUM, Spach, Hist. Vég. Phan. viii. 16 (1839).
COLUMBA, Comm. ex Endl. Gen. 826 (1840).
DIPLOCLISIA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42.
EPIBATERIUM, Forst. Char. Gen. 107. t. 54 (1776).
EPIBLASTRUM, Scop. Introd. 258 (1777).
FERRANDIA, Gardich. Fragv. Voy. Bot. 477 (1826).
GALLOA, Hassk. Cat. Hort. Bog. Alt. 172 (1844).
HOLOPELEA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42.
LEAENA, Forsk. Fl. Aegypt. Arab. 173 (1775).
LEONTHEMA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 58.
NEPHROIA, Lour. Fl. Cochinch. 566 (1790).
QUIMO, Schlecht. in Linnaea, xxvi. (1854) 792.
SELWYNIA, F. Muell. Fragm. iv. 158 (1864).
WENDLANDIA, Willd. Sp. Pl. ii. 876 (1799).
Abata, Kostel. Allg. Med.-Pharm. Fl. ii. 501 = Abuta rufescens.
acuminatus, DC. Syst. i. 527 = Tiliacora racemosa.
amazonum, *Mart. ex Herberger. in Buchn. Rep. Pharm.* xxxvi. (1830) 253.—Bras.
angustifolius, Hassk. Cat. Fl. Hort. Bog. Alt. 172 = laurifolius.
Aristolochiæ, DC. Syst. i. 520 = villosus.
Bakis, A. Rich. in Guill. & Perr. Fl. Senegamb. Tent. i. 12. t. 4 = Tinospora Bakis.
COCCULUS:—
Balfourii, *Schweinf. ex Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 500.—Ins. Socotra.
bracteiferifolius, A. Rich. ex St. Hil. & Tul. in Ann. Sc. Nat. Sér. II. xvii. (1842) 186 = Pachygone oblongifolia.
bustamensis, *Blume, Bijdr.* 26.—Java.
Blumeanus, Wall. Cat. n. 4971, partim = Cocsinium fenestratum.
brachystachyus, *DC. Syst.* i. 528.—Ins. Timor.
Burmanni, DC. l. c. 517 = Cyclea Burmanni.
calophyllus, Wall. ex Voigt, Hort. Suburb. Calc. 331, nomen; et ex Mast. in Hook. f. Fl. Brit. Ind. i. 578 = Erythropalum scandens.
carolinus, *DC. Syst.* i. 524.—Am. bor.
Cebatha, *DC. l. c.* 526.—Arabia.
Chondodendrum, *DC. l. c.* 529.—Am. austr.
cinerascens, St. Hil. Fl. Bras. Mer. i. 50 = Chondodendron platyphyllum.
cinereus, *Zoll. & Moritzi, Syst. Verz. Zoll.* 38.—Java.
convolvulaceus, DC. Syst. i. 519 = Tinospora cordifolia.
convolvulaceus, *Rudl. ex Walp. Rep.* i. 95.—Peruv.
cordifolius, DC. Syst. i. 517 = Tinospora cordifolia.
coriaceus, Blume, Bijdr. 25.—Java.
corymbosus, *Blume, l. c.* 24.—Java.
Cotoneaster, *DC. Syst.* i. 525.—Am. austr.
crispus, DC. l. c. 521 = Tinospora crispa.
cumanus, *Benth. in Journ. Linn. Soc. v. Suppl.* ii. (1861) 50.—China.
cuspidatus, Wall. Cat. n. 4960 = Limacia cuspidata.
cyanoclados, *Presl, Rel. Haenk.* ii. 79.—Zeylan.
dentatus, Hook. & Arn. Bot. Beech. Voy. 167 = Thanbergii.
dichopetalus, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 280.—Japon.
dichrous, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 46.—Bras.
diversifolius, *DC. Syst.* i. 523.—Mexic.
domingensis, DC. l. c. 528 = Pachygone domingensis.
ellipticus, DC. l. c. 526 = C. Leaeba.
emeuadus, Eichl. in Flora, xlvii. (1864) 391.—Peruv.
Epibasterium, *DC. Syst.* i. 520.—Ins. Pacif.
Ferrandianus, *Gaudich. in Freyc. Voy. Bot.* 477.—Ins. Sandvic.
Fibraureus, DC. Syst. i. 525 = Fibraurea tinctoria.
filipendulus, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 43.—Bras.
Finlaysonianus, Wall. Cat. n. 4974 = Stephania rotunda.
flavescens, *DC. Syst.* i. 520.—Ins. Molucc.
floridanus, *Wall. Cat.* n. 4976.—Ins. Penang.
Forsteri, *DC. Syst.* i. 517.—Hab. ?
glaber, Wight & Arn. Prod. i. 13 = C. Leaeba.
glaucescens, *Blume, Bijdr.* 26.—Borneo; Java.
glaucus, *DC. Syst.* i. 521.—Ins. Moluce.
gomphioides, *DC. l. c.* 530.—Madag.
hastatus, DC. l. c. 522 = villosus.
hexagynus, *Colebr. in Trans. Linn. Soc.* xiii. (1822) 68.—China.
Hookerianus, *F. Muell. ex Benth. in Journ. Linn. Soc. v. Suppl.* ii. (1861) 50.—Austral.
imenus, Mart. in Flora, xxiv. (1841) ii. *Beibl.* 44.—Anomospermum japonense.
incanus, Colebr. in Trans. Linn. Soc. xiii. (1822) 67. t. 17 = Anomospermum incanum.
japonicus, *DC. Syst.* i. 516.—Japon.
japurensis, Mart. in Flora, xxiv. (1841) ii. Beibl. 44 = Anomospermum japonense.
lacunosus, DC. Syst. i. 519 = Anamirta Cocculus.
laevigatus, Mart. in Flora, xxiv. (1841) ii. Beibl. 45 = Abuta sancelon.
laevis, Wall. Cat. n. 4975 = C. Leaeba.
lanuginosus, *Blume, Bijdr.* 24.—Java.
laurifolius, *DC. Syst.* i. 530.—Reg. Himal. ; Japon.
Leaeba, *DC. l. c.* i. 526.—As. trop. occ. ; Afr. trop.
leptostachyus, *DC. l. c.* 526.—Ins. Timor.
Limacia, *DC. l. c.* 526.—Cochinch.
Limacianus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 62 (err. typ. 61).—Burma.
lobata, Mart. ex Eichl. in Mart. Fl. Bras. xiii. i. 189 = Disciphania lobata.
longifolius, *Decne. in Mig. Ann. Mus. Bot. Lugd. Bat.* iv. 84.—Ins. Timor.
lucidus, Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind. iv. (1856) 397 = Antitaxis fasciculata.
macrocarpus, Hook. f. in Hook. Ic. Pl. t. 750 = Jateorhiza strigosa.
COCCULUS:—
macrocarpus, *Wight & Arn. Prod.* 13.—Ind. or. ; Burma ; China.
macrophyllus, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 184.—Bras.
malabaricus, DC. Syst. i. 518 = Tinospora malabarica.
Martii, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 185.—Bras.
megaspermus, *Voigt, Hort. Suburb. Calc.* 331.—Silhet.
membranaceus, Wall. Cat. n. 4967 = Cissampelos Pareira.
milleflorus, *DC. Syst.* i. 530.—Madag.
mollis, *Wall. Cat.* n. 4973.—Reg. Himal.
Moorei, F. Muell. Fragm. i. 162 = Pericampylus incanus.
Nephroia, *DC. Syst.* i. 531.—Cochinch.
nudiflorus, *Griff. Notul.* iv. 307.—Burma.
oblongifolius, *DC. Syst.* i. 529.—Am. bor. occ.
oblongifolius, Mart. in Flora, xxiv. (1841) ii. Beibl. 43 = Pachygone oblongifolia.
oblongus, Wall. Cat. n. 4968 = Limacia oblonga.
orbiculatus, DC. Syst. i. 523 = Cissampelos Pareira.
ovalifolius, Blume, Bijdr. 26 = umbellatus.
ovalifolius, DC. Syst. i. 526 = Thunbergii.
Pahni, Mart. in Flora, xxiv. (1841) ii. Beibl. 45 = Anomospermum japurense.
palmatus, DC. Syst. i. 522 = Jateorhiza Columba.
palmatus, Hook. Bot. Mag. t. 2970–71 = Jateorhiza Miersii.
paniculiger, Mart. in Flora, xxiv. (1841) ii. Beibl. 48 = Pachygone domingensis.
panper, *Griseb. in Gottt. Abh.* vii. (1856–57) 162.—Am. trop.
peltatus, DC. Syst. i. 516 = Cyclea peltata.
petiolaris, Wall. Cat. n. 4964 = Tinomiscium petiolare.
planiflorus, Wall. l. c. n. 4961 = Pycnarrhena planiflora.
platyphyllus, St. Hil. Fl. Bras. Mer. i. 59 = Chondodendron platyphyllum.
Pluknetii, DC. Syst. i. 520 = Pachygone ovata.
polycarpus, Wall. Cat. n. 4958 = Tiliacora racemosa.
populifolius, DC. Syst. i. 519 = Anamirta Cocculus.
radiatus, DC. l. c. 527 = Tiliacora racemosa.
racinus, Miers, Contrib. iii. 258 = C. Leaeba.
reticulatus, Mart. in Flora, xxiv. (1841) ii. Beibl. 44 = Anomospermum reticulatum.
rimosus, *Blume, Bijdr.* 24.—Java.
rotundifolius, *DC. Syst.* i. 517.—Hab. ?
Rosthurghianus, DC. l. c. 516 = Stephania rotunda.
sagittaefolius, *Meisn., Contrib.* iii. 255.—Am. bor.
Selwynii, F. Muell. Fragm. iv. 158 = Selwynia laurina.
sepium, Colebr. in Trans. Linn. Soc. xiii. (1822) 58. t. 6 = villosus.
smilacinus, F. Muell. ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 59 = Tinospora smilacina.
suberosus, DC. Syst. i. 519 = Anamirta paniculata.
tamoides, DC. l. c. 521 = Chondodendron tamoides.
tetrandrus, *Zipp. ex Span. in Linnaea*, xv. (1841) 164, nomen.—Malaya.
Thunbergii, *DC. Syst.* i. 524.—Japon.
tomentosus, Colebr. in Trans. Linn. Soc. xiii. (1822) 59 = Tinospora tomentosa.
tomentosus, Mart. ex Eichl. in Mart. Fl. Bras. xiii. i. 178 = Abuta rufescens.
triandrus, Colebr. in Trans. Linn. Soc. xiii. (1822) 64 = Limacia triandra.
triangularis, *Zipp. ex Span. in Linnaea*, xv. (1841) 164, nomen.—Malaya.
trilobus, *DC. Syst.* i. 529.—Java.
trilobus, *DC. l. c.* 522.—Japon.
umbellatus, *Steud. Nom. ed.* II. i. 802.—Java.
urophyllus, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 45.—Bras.
velutinus, Wall. Cat. n. 4970 = Limacia velutina.
verrucosus, Wall. l. c. n. 4966 A, B = Tinospora crispa.
verrucosus, Wall. l. c. n. 4966 C, D, E = Tinospora cordifolia.
villosus, *DC. Syst.* i. 525.—Ind. or. ; Afr. trop.
villosus, Wall. Cat. n. 4967, partim = Cissampelos Pareira.
Wightianus, Wall. l. c. n. 4959 A = Pachygone ovata.

COCCUS, Linn. Mus. Cliff. 11 (1786); Gen. ed. I. 336 (1737) = Cocos, Linn. (Palm.).

COCCYGANTHE, Reichb. Fl. Germ. Excurs. xvi. 55 (1844) = **Lychnis**, Linn. (Caryophyll.).
Flos-cuculi, Reichb. l. c. = L. Flos-cuculi.
pratensis, Schur, Enum. Fl. Transs. 107 = L. Flos-cuculi.

COCHLANTHERA, Choisy, in Mém. Soc. Phys. Genèv. xii. (1851) 426 = **Clusia**, Linn. (Guttifer.).
lanceolata, Choisy, l. c.—Venezuela.

COCHLANTHUS, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 78. *ASCLEPIADACEAE.*
socotranus, *Balf. f. l. c.* 79.—Ins. Socotra.

COCHLEANTHES, Rafin. Fl. Tellur. 45 (1836) = **Zygopetalum**, Hook. (Orchid.).
fragrans, Rafin. l. c. = Z. cochleare.

COCHLEARIA, Tourn. ex Linn. Syst. ed. 1 (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 75.
ARMORACIA, Rtv. ex Gaertn. Mey. & Scherb. Fl. Wett. ii. 426 (1800).
CHRYSOCHAMELA, Boiss. Fl. Orient. i. 813 (1867).
GONYCLISIA, Dulac, Fl. Hautes-Pyr. 191 (1867).
IONOPSIDIUM, DC. ex Steud. Nom. ed. II. i. 392 (1840).
KERNERA, Medic. Pflanzeng. 71 (1792).
RAPHANIS, Dod. ex Moench, Meth. 267 (1794).
RHIZOBOTRYA, Tausch, in Flora, xix (1836) 33.
TAPHROSPERMUM, C. A. Mey. in Ledeb. Fl. Alt. iii. 179 (1831).
TROPHOSPERMUM, Walp. Rep. i. 171 (1842).
acaulis, *Desf. Fl. Atlant.* ii. 69.—Lusitan. ; Afr. bor.
alpina, Sweet, Hort. Brit. ed. I. 22 = officinalis.
alsacica, Schlecht. ex DC. Prod. i. 178 = microcarpa.
alyssoides, DC. l. c. 172 = flava.
amphibia, *Ledeb. Fl. Ross.* i. 160.—Soongaria.
anglica, *Linn. Syst. ed.* X. 1136.—Reg. bor. et arct.
aphragmodes, *Boiss. Diagn.* Ser. I. vi. 15.—Persia.
aquatica, Kaston, Man. ed. V. 181 = Nasturtium lacustre !
aquatica, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 88.—As. bor.
arctica, Schlecht. in DC. Syst. ii. 367 = anglica.
Armoracia, *Linn. Sp. Pl.* i. 643.—Europ.
Armoracia, Pall. ex DC. Syst. ii. 367 = microcarpa.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 168.—Armenia.
auriculata, *Lam. Encyc.* ii. 165.—Europ.
austriaca, *Ledeb. Fl. Ross.* i. 160.—Rossia.
austriaca, Vis. Fl. Dalm. iii. 122 = Nasturtium austriacum.
batava, *Dum. Fl. Belg. Prod.* 120.—Batavia.
brevicaulis, *Facch. in Zeitschr. Ferdin. Tirol.* III. (1854) 79.—Tyrol.
Cakile, Wall. Cat. n. 4802 = flava.
Camelinae, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 173.—Persia.
Camelinae, *Ledeb. Fl. Ross.* i. 160, in nota.—Manchuria.
campylocarpa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 171.—Persia.
cardiolla, *Turcz. ex Bunge, in Mém. Sav. Etr. Acad. Pétersb.* ii. (1835) 570.—Sibir. balcal.
Coronopus, Linn. Sp. Pl. 648 = Senebiera Coronopus.
Crantzii, *Crantz, Class. Crucif. Emend.* 100.—Europ.
crassifolia, Willd. ex DC. Prod. i. 173 = macrocarpa.
danica, Gunn. Fl. Norv. 72. n. 197 = officinalis.
danica, *Linn. Sp. Pl.* 647.—Reg. bor. et arct.
decipiens, *Willk Sert. Fl. Hispan.* 13.—Mont. Pyren.
Draba, Linn. Syst. ed. X. 1129 = Lepidium Draba.
drabaecarpa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 169.—Persia.
fenestrata, R. Br. in App. Voy. Ross, (1819) p. cxliii = officinalis.
fragrans, Gilib. Fl. Lituan. ii. 51 = officinalis.
flava, *Buch.-Ham. ex Roxb. Hort. Beng.* 48, nomen.—Ind. or.
fortida, Schkuhr, Beitr. Fl. ii. 105 = Camelina foetida.
glastifolia, Falk, Beitr. ii. 215 = Hymenophysa pubescens.
glastifolia, *Linn. Sp. Pl.* 646.—Europ.
glastifolia, Pall. ex DC. Syst. ii. 369 = Draba aizoa.
glaucophylla, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 168.—Persia.
globosa, *Ledeb. Fl. Ross.* i. 159.—Dahuria.
grandiflora, *DC. Syst.* ii. 368.—Sibiria.
groenlandica, Gunn. Fl. Norv. 72. n. 196 = anglica.

COCHLEARIA:—
groenlandica, Linn. Sp. Pl. 647 = officinalis.
groenlandica, Willd. ex DC. Prod. i. 173 = lenensis.
hastata, Moench, Meth. Suppl. 87 = danica.
hederacea, S. F. Gray, Nat. Arr. Brit. Pl. ii. 696 = danica.
heterophylla, Cav. Descr. 405 = Camelina dentata.
heterophylla, Fisch. Hort. Gorenk. ed. II. (1812) 50, nomen = seq.
heterophylla, Schlecht. in DC. Syst. ii. 372 = sisymbrioides.
himalaica, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 154.—Reg. Himal.
humifusa, Michx. Fl. Bor. Am. ii. 27 = Senebiera didyma.
integrifolia, DC. Syst. ii. 368 = Eutrema alpestre.
kamtschatica, Schlecht. ex DC. Prod. i. 173 = officinalis.
lancifolia, Stokes, Bot. Mat. Med. iii. 437 = C. Armoracia.
lapathifolia, Gilib. Fl. Lituan. iv. 53 = C. Armoracia.
lenensis, *Adams, ex Fisch. in DC. Syst.* ii. 367.—Sibiria.
Linnaei, Grlewank, ex Aschers. in Fl. Prov. Brandenb. 54 = officinalis.
longifolia, *Medic. in Usteri, Neu. Ann.* ii. 41.—Europ.
lyrata, *Sibth. & Sm. Fl. Graec. Prod.* ii 8 ; *Fl. Graec.* t. 619.—Sicilia.
macrocarpa, *Waldst. & Kit. Pl. Rar. Hung.* ii. 101. t. 184.—Hungar : Transsylv.
mexicana, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 220.—Mexic.
microcarpa, *DC. Syst.* ii. 362.—Sibir.
minuta, Steud. Nom. ed. II. i. 392 = danica.
natans, C. A. Mey. in Ledeb. Fl. Alt. iii. 90 = amphibia.
nilotica, Delile, Fl. Egypte, 245 = Senebiera nilotica.
oblongifolia, DC. Syst. ii. 368 = officinalis.
officinalis, Lapeyr. Hist. Abr. Pl. Pyr. 368 = pyrenaica.
officinalis, *Linn. Sp. Pl.* 647.—Reg. bor. et arct.
olyssoides, Brot. Fl. Lusit. i. 571 = acaulis.
ovalifolia, Stokes, Bot. Mat. Med. iii. 436 = anglica.
planisiliqua, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 169.—Persia.
polysperma, Costa, ex Willk. in Linnaea, xxx. (1859-60) 56 = Kernera polysperma.
Pseud-Armoracia, Czern. ex Trautv. in Act. Hort. Petrop. viii. (1883) 93 = macrocarpa !
pusilla, Brot. Phyt. 100 = acaulis.
pyrenaica, DC. Syst. ii. 365 = officinalis.
renifolia, Stokes, Bot. Mat. Med. iii. 435 = officinalis.
repanda, *Medic. in Usteri. Neu. Ann.* ii. 42.—Europ.
repens, Lam. Fl. Fr. ii. 473 = Senebiera Coronopus.
rhizobotrya, *Walp. Rep.* i. 158.—Austria.
rotundifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 695 = officinalis.
rusticana, Lam. Fl. Fr. ii. 471 = C. Armoracia.
sagittaefolia, *Desv. Journ. Bot.* iii. (1812) 182.—Hab. ?
sagittata, Crantz, Class. Crucif. Emend. 99 = Neslia paniculata.
salsa, *Schlecht. ex DC. Syst.* ii. 369.—Sibiria ?
sativa, Cav. Descr. 405 = Camelina sativa.
saxatilis, *Linn. Sp. Pl.* 648.—Europ.
saxifragaefolia, DC. Syst. ii. 370 = Graellsia saxifragaefolia.
scapiflora, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 154.—Reg. Himal.
Sempervivum, *Boiss. & Bal. Diagn.* Ser. II. v. 28.—As. Min.
septentrionalis, DC. Prod. i. 174 = Draba hyperborea.
septentrionalis, Schlecht. in DC. Syst. ii. 365 = officinalis.
sibirica, Willd. Sp. Pl. iii. 450 = Sobolewskia lithophila.
siliquosa, Schlecht. in DC. Syst. ii. 369 = Draba hyperborea.
sisymbroides, *DC. l. c.* 366.—Sibir. arct.
spathulata, *Schlecht. in DC. l. c.* 369 = Draba hyperborea.
Szovitzii, *Boiss. Fl. Orient.* i. 248.—Persia.
tenuifolia, Saliab. Prod. 267 = Senebiera Coronopus.
tridactylites, *Bunbe, ex DC. Syst.* ii. 367.—Am. arct.
turritifolia, Saliab. Prod. 267 = C. Armoracia.
velutina, *DC. Syst.* ii. 370.—Mesopot.
violacea, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 171.—Persia.

COCHLEARIA:—
Wahlenbergii, Rupr. Beitr. Pfl. Russ. Reich. ii. 21 = anglica.
Wunderlichii, *C. A. Mey. ex Claus, in Beitr. Pfl. Russ. Reich.* viii. 208.—Rossia.

COCHLEATA, Rtv. ex Rupp. Fl. Jen. ed. Hall. 264 (1745) = **Medicago**, Linn. (Legum.).
intertexta, Medic. in Vorles. Churf. Ges. ii. (1787) 385 = M. intertexta.
scutellata, Medic. Phil. Bot. i. 210 = M. scutellata.

COCHLIA, Blume, Bijdr. 320. t. 59 (1825) = **Bulbophyllum**, Thou. (Orchid.).
violacea, Blume, l. c. = B. violaceum.

COCHLIANTHUS, Benth. in Miq. Pl. Jungh. i. 284, in note (1851 ?). *LEGUMINOSAE*, Benth. & Hook. f. i. 532.
graeilis, *Benth. l. c.*—Reg. Himal.

COCHLIDIOSPERMUM, Reichb. Consp. 121 (1828) = **Veronica**, Linn. (Scrophularin.).
agrestis, Opiz, Seznam, 31 = V. agrestis.
Buchaumii, Opiz, l. c. = V. Tournefortii.
digitatum, Opiz, l. c. = V triphyllos.
Fresenanum, Opiz, l. c. = V. opaca.
hederaefolium, Opiz, l. c. = V. hederaefolia.
Lappago, Opiz, in Lotos, iv. (1854) 168 = V. hederaefolia.
opacum, Opiz, Seznam, 31 = V. opaca.
praecox, Opiz, l. c. = V. praecox.

COCHLIODA, Lindl. Fol. Orchid. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 560.
densiflora, *Lindl. l. c.*—Reg. Andin.

COCHLIOPETALUM, Beer, in Flora, xxxvii. (1854) 347 ; Bromel. 68 (1857) = **Pitcairnia**, L'Hérit. (Bromeliac.).
albiflos, Beer, Bromel. 68 = P. albiflos.
flavescens, Beer, l. c. 69 = P. xanthocalyx.
odoratum, Hemsl. Biol. Centr. Am. Bot. iii. 317 = P. odorata.
Schuchii, Beer, Bromel. 69 = P. albiflos.
staminosum, Beer, in Flora, xxxvii. (1854) 347 = P. staminea.

COCHLIOSPERMUM, Lag. Mem. Pl. Barill. 58 (1817) = **Bunaea**, Forsk. (Chenopod.).
altissimum, Lag. l. c. 57 = S. altissima.
Covanillesii, Lag. l. c. 60 = S. maritima.
Clementei, Lag. l. c. 62 = S. splendens.
fruticosum, Lag. l. c. 59 = S. fruticosa.
hispanicum, Lag. l. c. 58 = S. altissima.
salsum, Lag. l. c. 55 = S. maritima.

COCHLIOSTEMA, Lem. Illustr. Hortic. vi. (1859) Misc. 70 et t. 217. *COMMELINACEAE*, Benth. & Hook. f. iii. 850.
Jacobianum, *C. Koch & Linden, in C. Koch & Fint. Wochenschr.* x. (1867) 322.—Ecuador.
odoratissimum, *Lem. Illustr. Hortic.* vi. (1859) *Misc.* 70. t. 217.—Ecuador.
odoratum, C. Koch, in C. Koch & Fint. Wochenschr. ii. (1859) 340 = odoratissimum.

COCHLOSPERMUM, Kunth, Malv. 6 (1822). *BIXINEAE*, Benth. & Hook. f. i. 124.
AZEREDIA, Arruda, ex Allem. Denemb. (1846).
MAXIMILIANA, Mart. in Flora, ii. (1819) 481.
WITTELSBACHIA, Mart. Nov. Gen. et Sp. i. 80. t. 55 (1824).
anglense, *Welw. Apont.* 566 ; *Oliver, Fl. Trop. Afr.* i. 113.—Afr. trop.
Cadinae, *Eichl. in Mart. Fl. Bras.* xiii. i. 481.—Bras.
Fraseri, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 307.—Austral.
Gillivraei, *Benth. Fl. Austral.* i. 106.—Austral.
Gossypium, *DC. Prod.* i. 527.—Ind. or.
Gregorii, *F. Muell. Fragm.* i. 71.—Austral.
heteronemum, *F. Muell. in Hook. Kew Journ.* ix. (1857) 15 (= seq. ?).—Austral.
heteronervum, *F. Muell. ex Benth. Fl. Austral.* i. 106.—Austral.
hibiscoides, *Kunth, Syn. Pl. Aeq.* iii. 214.—Mexic.
indigne, *St. Hil. Pl. Or. Bras.* t. 57.—Bras.
nilotieum, *Oliver, Fl. Trop. Afr.* i. 113.—Afr. trop.

COCHLOSPERMUM :—
orinocoense, *Steud. Nom.* ed. II. i. 393.—Venezuela.
Parkeri, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 210.—Guiana.
periticifolium, *Planch. l. c.* 211.—Guiana.
Planchoni, *Hook. f. ex Planch. l. c.* 209.—Afr. trop.
serratifolium, DC. Prod. i. 537 = hibiscoides.
tinctorium, *Perr. in Ann. de la Marine et des Colonies* (1830); *ex A. Rich. in Guill. & Perr. Fl. Senieg. Tent.* i. 99.—Afr. trop.
vitifolium, Spreng. Syst. ii. 596 = serratifolium.
vitifolium, *Spreng. l. c.* iv. *Cur. Post.* 206.—Mexic.

COCHRANIA, Miers, Trav. Chile, ii. 529 (1826) nomen; et in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 125. *BORAGINEÆ*, Benth. & Hook. f. ii. 845.
chenopodioides, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868) 132.—Chili.
conferta, *Miers, Trav. Chile*, ii. 529, nomen; *et i. c.* 125. —Chili.
corymbosa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868) 126.—Chili.
ericoides, *Miers, l. c.* 130.—Chili.
filifolia, *Miers, l. c.* 131.—Chili.
floribunda, *Miers, l. c.* 129.—Chili.
hebeculis, *Miers, l. c.* 130.—Chili.
hispidula, *Miers, l. c.* 132.—Chili.
myosotifolia, *Miers, l. c.* 128.—Chili.
sinuata, *Miers, l. c.* 127.—Chili.
stenophylla, *Miers, l. c.* 128.—Chili.

COCKEBURNIA, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 90. *SELAGINEÆ*.
scoparia, *Balf. f. l. c.*—Ins. Socotra.

COCOCIPSILUM, Jaume St. Hil. Expos. i. 434 (1805) = Cocococypselum, P. Br. (Rubiac.).

COCOLOBA, Rafin. Fl. Tellur. ii. 84 (1836) = Coccoloba, Linn. (Polygal.).

COCOS, Linn. Sp. Pl. ed. I. 1188 (1753). *PALMAE*, Benth. & Hook. f. iii. 945.
GLAZIOVA, Mart. ex H. Wendl. in Flor. et Pom. (1871) 114.
LARDIZOBALIA, Raddi, in Mem. Soc. Ital. Mod. xviii. (1820) 347. t. 1.
LITHOCARPUS, Targ. Tozz. ex Stend. Nom. ed. II. i. 178, in syn. (1840).
PLATENIA, Karst. in Linnæa, xxviii. (1856) 250.
SYAGRUS, Mart. Hist. Nat. Palm. ii. 129 (1824).
acaulis, *Drude, in Mart. Fl. Bras.* iii. ii. 426.—Bras.
arioulevis, Sw. Prod. Veg. Ind. Occ. 58 = Bactris minor.
acrocomioides, *Drude, in Mart. Fl. Bras.* iii. ii. 409. —Bras.
aculeata, Jacq. Select. Am. 278 = Acrocomia sclerocarpa.
aequatorialis, Drude, in Mart. Fl. Bras. iii. ii. 407, sphalm. = C. lunjai.
amara, *Jacq. Select. Am.* 277.—Ind. occ.
arenaria, Gomes, in Mem. Ac. Lisboa, iii. (1812) Mem. dos Corresp. 61 = Diplothemium littorale.
argentea, *Engel, in Linnæa*, xxxiii. (1864–65) 690.— N. Granat.
Ariouï, Neuwied, Reise Bras. i. 278 = schizophylla.
australis, *Mart. Palmet. Orbign.* 95.—Paraguay.
Bonnetti, *Hort. ex H. Wendl. in Kerch. Palm.* 240, nomen.—Hab. ?
botryophora, *Mart. Hist Nat. Palm.* ii. 118. t. 83, 84. —Bras.
butyracea, *Linn. f. Suppl.* 454.—Am. austr.
butyracea, Mart. ex H. Wendl. in Kerch. Palm. 140 = Scheelea butyracea.
butyrosa, H. Wendl. l. c. = Attalea humilis.
campestris, *Mart. Hist. Nat. Palm.* ii. 121.—Bras.
capitata, *Mart. l. c.* 114.—Bras.
chiliensis, Molina, ex H. B. & K. Nov. Gen. et Sp. i. 309, in obs. = Jubæa spectabilis.
comosa, *Mart. Hist. Nat. Palm.* ii. 121. t. 88. f. 1–2.— Bras.
coronata, *Mart. l. c.* 115. t. 80, 81.—Bras.
crispa, *H. B. & K. Nov. Gen. et Sp.* i. 309.—Cuba.
Datil, *Drude & Griseb. in Gœtt. Abh.* xxiv. (1879) 285.—Reg. Argent.
elegantissima, Hort. ex H. Wendl. in Kerch. Palm. 241 = Weddelliana.
COCOS :—
equatorialis, Barb.-Rodr. Enum. Palm. Nov. 38 = C. lunjai.
eriospatha, *Mart. ex Drude, in Mart. Fl. Bras.* iii. ii. 424.—Bras.
flexuosa, *Mart. Hist. Nat. Palm.* ii. 120.—Bras.
frigida, *Lindm. in Illustr. Hortic.* xxviii. (1881) 16, nomen.—Bras.
fusiformis, Sw. Fl. Ind. Occ. i. 616 = Acrocomia sclerocarpa.
Gerdia, Barb.-Rodr. Prot. App. 48 = botryophora.
graminifolia, *Drude, in Mart. Fl. Bras.* iii. ii. 415.— Bras.
Gnacaryale, *Lisbm. ex Mart. Hist. Nat. Palm.* iii. 233. —Mexic.
guineensis, Linn. Mant. i. 137 = Bactris minor.
lunjai, *Spruce, ex Trail, in Journ. Bot.* xv. (1877) 79. —Bras.
insignis, *Mart. ex H. Wendl. in Kerch. Palm.* 241.— Bras.
Kotchoubeyi, *Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Bras.
lapidea, *Gaertn. Fruct.* i. 16. t. 6. f. 1.—Bras. ?
leiophylla, *Barb.-Rodr. Prot. App.* 44.—Bras.
maldivica, J. F. Gmel. Syst. 969 = Lodoicea seschellarum.
mamillaris, *Blanco, Fl. Filip.* ed. I. 722.—Ins. Philipp.
maritima, Comm. ex H. Wendl. in Kerch. Palm. 241 = Lodoicea sp.
Martiana, *Drude & Glaz. ex Drude, in Mart. Fl. Bras.* iii. ii. 418.—Bras.
melanococca, Hort. Belg. ex Heynh. Nom. ii. 152, nomen.—Hab. ?
Mikaniana, *Mart. Hist. Nat. Palm.* iii. 138.—Bras.
Naja, *Arruda, ex Kunth, Enum. Pl.* iii. 288.— Am. trop.
nana, Griff. Notul. iii. 166 = nucifera.
Nolala, *Neuwied, Reise Bras.* i. 271.—Bras.
Normandyi, W. Hill, Rep. Brsb. Bot. (1876) 6; ex F. Muell. Fragm. viii. 235 = Areca Normanbyi.
nucifera, *Sp. Pl.* 1188.—Reg. trop.
Nypa, Lour. Fl. Cochinch. 567 = Nipa fruticans.
oleracea, *Mart. Hist. Nat. Palm.* ii. 117. t. 82.—Bras.
oriocomoides, *Spruce, in Journ. Linn. Soc.* xi. (1871) 161.—Am. trop.
ovata, *Hort. Belg. ex Heynh. Nom.* ii. 152, nomen.— Hab. ?
petræa, *Mart. Palmet. Orbign.* 100.—Bolivia.
pityrophylla, *Mart. l. c.* 99.—Bolivia.
plumosa, *Hook. Bot. Mag.* t. 5180.—Bras.
plumosa, Lodd. ex Lond. Hort. Brit. 881 = comosa.
Procopiana, *Glaz. ex Drude, in Mart. Fl. Bras.* iii. ii. 412.—Bras.
regia, *Lisbm. in Mart. Hist. Nat. Palm.* iii. 228.— Mexic.
Romanzoffiana, *Cham. in Choris, Voy. Pittor. (Chili)* 5. t. 6 & 6 [bis].—Bras.
rupestris, Barb.-Rodr. Prot. App. 45 = petræa.
schizophylla, *Mart. Hist. Nat. Palm.* ii. 119. t. 84, 85. —Bras.
speciosa, *Barb.-Rodr. Enum. Palm. Nov.* 38.—Bras.
simplicifolia, *Hort. Belg. ex Heynh. Nom.* ii. 152.— Hab. ?
Syagrus, *Drude, in Mart. Fl. Bras.* iii. ii. 406.—Bras.
syivestris, *Hort. ex H. Wendl. in Kerch. Palm.* 241, nomen.—Hab. ?
Uracuru, *Lindem, in Illustr. Hortic.* xxviii. (1881) 16, nomen.—Bras.
ventricosa, *Arruda, in Koster, Trav. Bras.* 485.—Bras.
venustorum, Poepp. ex H. Wendl. in Kerch. Palm. 241 = Maximiliana venatorum.
viridens, Mart. Hist. Nat. Palm. iii. 224 = Gaussia viridens.
Weddelliana, *H. Wendl. in Flor. et Pom.* (1871) 114.
Weddelliî, *Drude, in Mart. Fl. Bras.* iii. ii. 411.— Bras.
Yatay, *Mart. Palmet. Orb.* 93.—Reg. Argent.
Yarumaguas, *Mart. ex H. Wendl. in Kerch. Palm.* 241. —Peruv.

CODANTHERA, Rafin. Fl. Tellur. iii. 90 (1836) = Salvia, Tourn. (Labiat.).
bifora, Rafin. l. c. = S. bifolora.
glabra, Rafin. l. c. = S. strictiflora.

CODARIA, Linn. ex Benn. Pl. Jav. Rar. 99 (1838) = Lorchhea, Kalm. (Rubiac.).

CODARIOCALYX, Hask. in Flora, xxv. (1842) ii. Beibl. 48 = Desmodium, Desv. (Legumin.).
capitatus, Hask. l. c. 49 = D. capitatum.
conicus, Hask. l. c. 49, 62 = D. gyroides.
gyrans, Hask. l. c. = D. gyrans.
gyroides, Hask. l. c. 49 = D. gyroides.

CODARIUM, Soland. ex Vahl, Enum. i. 302 (1806) = Dialium, Linn. (Legumin.).
acutifolium, Afzel. in Schrad. N. Journ. ii. (1807) 233 = D. guineense.
discolor, DC. Prod. ii. 520 = D. guineense.
nitidum, Vahl, Enum. i. 302 ; ii. 400 = D. guineense.
obtusifolium, Afzel. in Schrad. N. Journ. ii. (1807) 233 = D. guineense.
Solandri, Vahl, Enum. i. 302 = D. guineense.

CODAZZIA, Karst. & Triana, in Linnæa, xxviii. (1856) 426 = Dalostoma, D. Don (Bignoniac.).
dentata, Karst. Fl. Colomb. ii. 87. t. 118 = D. roseo.
rosea, Karst. in Linnæa, xxviii. (1856) 427 = D. roseo.
speciosa, Karst. l. c.—N. Granat.

CODDAMPULI, Adans. Fam. ii. 445 (1763) = Garcinia, Linn. (Guttif.).

CODDA-PANA, Adans. Fam. ii. 25 (1763) = Bentinckia, Berry (Palm.).

CODIA, Forst. Char. Gen. 59. t. 30 (1776). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 848.
ferruginea, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 77.—N. Caled.
floribunda, *Brongn. & Gris, l. c.*—N. Caled.
montana, *Forst. Char. Gen.* 68.—N. Caled.
montana, Labill. ex D. Don, in Edinb. N. Phil. Journ. (Apr.–June 1830) 94 = Callicoma Billardieri.
obcordata, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 77.—N. Caled.
serratifolia, Ser. ex DC. Prod. iv. 7 = Callicoma serratifolia.
spathulata, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 77.—N. Caled.

CODIAEUM, Rumph. ex A. Juss. Euph. Tent. 23. t. 9 (1824). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 299.
CROSOPHYLA, Rafin. Sylva Tellur. 64 (1838).
JUNGHUHNIA, Miq. Fl. Ind. Bat. i. 11. 412 (1855).
PHYLLAURea, Lour. Fl. Cochinch. 575 (1790).
SYNASPISMA, Endl. Gen. 1110 (1840).
alternifolium, *Muell. Arg. in DC. Prod.* xv. ii. 1117. —N. Caled.
andamanicum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 246 (= Binchiae sp.).—Ins. Andaman.
aurantiacum, *Muell. Arg. in DC. Prod.* xv. ii. 1118.— Ins. Banca.
Balansae, *Baill. Adansonia*, xi. (1873–76) 77.—N. Caled.
Brongniartii, *Baill. l. c.* 76.—N. Caled.
Bureavi, *Baill. l. c.* 74.—N. Caled.
carunculatum, *Muell. Arg. in DC. Prod.* xv. ii. 1117.— N. Caled.
chrysosticton, Spreng. Syst. iii. 886 = variegatum.
cuneifolium, Zipp. ex Span. in Linnæa, xv. (1841) 345 = variegatum.
Deplanchei, *Baill. Adansonia*, xi. (1873–76) 75.—N. Caled.
drinciflorum, *Baill. l. c.*—N. Caled.
ingobyliense, *Muell. Arg. in DC. Prod.* xv. ii. 1120.— N. Caled.
lutescens, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 246 = Sphyranthera cephalitata.
medium, Baill. Adansonia, i. (1860–61) 251 = variegatum.
meleagrinum, Decne, in Nouv. Ann. Mus. Par. iii. (1834) 488 = variegatum.
obovatum, Zoll. & Mor. in Nat. en Genees. Arch. Nederl. Indie, ii. (1845) 868 = variegatum.
Pancheri, Muell. Arg. in DC. Prod. xv. ii. 1117 = Baloghia Pancheri.
Pentoti, Muell. Arg. l. c. 1118 = Blachia Pentoti.
pictum, Hook. Bot. Mag. t. 3061 = variegatum.
roseo-pictum, *André, Illustr. Hortic.* xxvi. (1879) 170. t. 286.—Hab. ?
timorense, A. Juss. Tent. Euph. 24, nomen = variegatum.
umbellatum, Muell. Arg. in DC. Prod. xv. ii. 1118 = Blachia umbellata.
variegatum, *Blume, Bijdr.* 606.—Malaya.

CODIAMINUM, Rafin. Fl. Tellur. iv. 21 (1836) = **Narcissus**, Tourn. (Amaryllid.) (N. Bulbocodium).

CODIPHUS, Rafin. Fl. Tellur. ii. 77 (1836) = **Prismatocarpus**, L'Herit. (Campanulac.). *nitidus*, Rafin. l. c. = P. nitidus.

CODIVALIA, Rafin. Fl. Tellur. iii. 40 (1836) = **Pupalia**, Juss. (Amarant.). *lappacea*, Rafin. l. c. = P. lappacea. *patula*, Rafin. l. c. = P. lappacea.

CODOCHONIA, Dun. in DC. Prod. xiii. 1. 482 (1852) = **Acnistus**, Schott (Solanac.). *solanoides*, Dun. l. c. = A. umbellatus.

CODON, Linn. Syst. ed. XII. 292 (1767). *HYDROPHYLLACEÆ*, Benth. & Hook. f. ii. 880. *aculeatum*, Gaertn. Fruct. ii. 88 = Royeni. Dregei, *E. Mey. Zwei Pfl. Docum.* 174, nomen.—Afr. austr. Royeni. *Linn. Syst.* ed. XII. 292.—Afr. austr.

CODONACANTHUS, Nees, in DC. Prod. xi. 108 (1847). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1098. *acuminatus*, Nees, l. c. = pauciflorus. pauciflorus, *Nees, l. c.*—Hongkong.

CODONACHNE, Wight & Arn. ex Steud. Nom. ed. II. i. 393 (1840) = **Chloris**, Sw. (Gramin.). *Neesiana*, Wight & Arn. l. c. = Chloris tenella.

CODONANDRA, Karst. Fl. Columb. ii. 48. t. 122 (1869 ?) = **Calliandra**, Benth. (Legumin.). *corolnata*, C. Muell. in Walp. Ann. vii. 841, sphalm. = Call. purpurea. *purpurea*, Karst. Fl. Columb. ii. 48. t. 122 = Call. Codonandra.

CODONANTHE, Hanst. in Linnæa, xxvi. (1853) 209. *GESNERACEÆ*, Benth. & Hook. f. ii. 1011.
COCCANTHERA, C. Koch & Hanst. Ind. Sem. Hort. Berol. (1855) 17.
aggregata, *Hanst. in Linnæa*, xxvi. (1853–55) 209.—Bras.
calcarata, *Hanst. l. c.* xxxiv. (1865–66) 416.—Am. trop.
carnosa, *Hanst. in Mart. Fl. Bras.* viii. 418.—Bras.
elliptica, *Lem. in Illustr. Hortic.* ii. (1855) sub t. 56.—Bras.
Devosiana, *Lem. l. c.* t. 56.—Bras.
gracilis, *Hanst. in Linnæa*, xxvi. (1853–55) 209.—Bras.
Hookertii, *Lem. in Illustr. Hortic.* ii. (1855) sub t. 56.—Bras.
picta, Lem. l. c. iv. (1857) t. 144 = Devosiana.

CODONANTHE, Mart. ex Steud. Nom. ed. II. i. 791, in syn. (1840) = **Hypocyrta**, Mart. (Gesnerac.).

CODONANTHEMUM, Klotzsch, in Linnæa, xii. (1838) 240 = **Sinochetus**, Klotzsch (Ericac.). *discolor*, Benth. in DC. Prod. vii. 708.—Afr. austr. *parviflorum*, Klotzsch, in Linnæa, xii. (1838) 240.—Afr. austr.
puberulum, Klotzsch, l. c.—Afr. austr.
tenue, Benth. in DC. Prod. vii. 708.—Afr. austr.

CODONANTHES, Rafin. Fl. Tellur. iv. 24 (1836) = **Pitcairnia**, L'Hérit. (Bromeliac.). *albiflos*, Rafin. l. c. = P. albiflos.

CODONANTHUS, G. Don, Gen. Syst. iv. 166 (1838) = **Erweseia**, R. Br. (Convolvulac.).
africana, G. Don, l. c. = R. africana.
alternifolia, Planch. in Hook. Ic. Pl. t. 796.—Afr. trop.

CODONANTHUS, Hassk. in Flora, xxv. (1842) ii. Beibl. 24 = **Physostelma**, Wight (Asclepiad.).
campanulatus, Hassk. l. c. = P. Wallichii.

CODONEMMA, Miers, Apocyn. S. Am. 72 (1878) = **Tabernæmontana**, Plum. (Apocynac.).
calycinum, Miers, l. c. 78 = T. Muelleriana.
macrocalyx, Miers, l. c. = T. macrocalyx.

CODONIUM, Vahl, in Skriv. Naturh. Selsk. ii. (1793) t. 206; Symb. Bot. iii. 36 (1794) = **Schoepfia**, Schreb. (Olacin.).
arborescens, Vahl, ii. oc. 207; 36 = S. arborescens.
flexuosum, M. Roem. Syn. Hesper. 21 = S. flexuosa.

CODONOCALYX, Klotzsch, ex Baill. Etud. Gen. Euph. 369 (1858) = **Croton**, Linn. (Euphorb.).
discoricatus, Klotzsch, l. c. 370 = Crot. nitrariæfolius.
lanatus, Klotzsch, l. c. = Crot. lanuginosus.
longifolius, Klotzsch, l. c. = Crot. montevidensis.
montevidensis, Klotzsch, l. c. = Crot. montevidensis.
polymorphus, Klotzsch, l. c. = Crot. nitrariæfolius.
wellerifloros, Klotzsch, l. c. = Crot. nitrariæfolius.

CODONOCALYX, Miers, in Lindl. Veg. Kingd. 764 (1847) = **Buteria**, DC. (Rubiac.).

CODONOCARPUS, A. Cunn. ex Hook. Bot. Misc. i. (1830) 244, in syn. *PHYTOLACCACEÆ*, Benth. & Hook. f. iii. 86.
HYMENOTHECA, F. Muell. Fragm. i. 201 (1859). australis, *A. Cunn. l. c.*—Austral.
cotinifolia, *F. Muell. Pl. Vict.* i. 200.—Austral.
pyramidalis, *F. Muell. l. c.* 201.—Austral.

CODONOCEPHALUM, Fenzl, in Flora, xxvi. (1843) 897. *COMPOSITÆ*, Benth. & Hook. f. ii. 520.
SPRUNNERA, Sch. Bip. in Walp. Rep. ii. 954 (1843). inuloides, *Fenzl, l. c.*—Kurdistan.

CODONOCRINUM, Willd. ex Schult. f. Syst. vii. 718 (1829) = **Yucca**. Dill. (Liliac.).
agavoides, Willd. ex Schult. f. l. c. = Y. acaulis.

CODONOPHORA, Lindl. Bot. Reg. sub t. 1110 (1827) = **Paliavana**, Vand. (Gesnerac.).
grandiflora, Lindl. l. c. = P. prasinata.
lanceolata, Lindl. l. c. = Rytidophyllum tomentosum.
prasinata, Lindl. l. c. = P. sericiflora.
Selloana, Hanst. in Mart. Fl. Bras. viii. 396 = P. prasinata.

CODONOPRASUM, Reichb. Consp. 66 (1828) = **Allium**, Tourn. (Liliac.).
alpicolum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 127 = A. oleraceum.
carinatum, Reichb. Fl. Germ. Excurs. 114 = A. carinatum.
complanatum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 159 = A. oleraceum.
consimile, Fourr. l. c. = A. carinatum.
flavum, Reichb. Fl. Germ. Excurs. 115 = A. flavum.
flexifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = A. carinatum.
flexum, Reichb. Fl. Germ. Excurs. 114 = A. flavum.
fuscum, Reichb. l. c. 115 = A. paniculatum.
intermedium, Reichb. l. c. 114 = A. oleraceum.
longispathum, Reichb. l. c. 115 = A. paniculatum.
oleraceum, Reichb. l. c. 114 = A. oleraceum.
pallens, Reichb. l. c. 115 = A. flavum.
paniculatum, Reichb. l. c. = A. pulchellum.
pulchellum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = A. pulchellum.
viridiflorum, Schur, Enum. Pl. Transs. 671 = A. oleraceum.

CODONOPSIS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 108 (1824). *CAMPANULACEÆ*, Benth. & Hook. f. ii. 557.
CLEMATOCOMIA, D. Don. Prod. Fl. Nep. 158 (1824).
affinis, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 12.—Reg. Himal.
albiflora, Griff. Notul. iv. 279 = Campanumœa celebica.
Benthamii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 14.—Reg. Himal.
celebica, Miq. Fl. Ind. Bat. ii. 566 = Campanumœa celebica.
clematidea, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 433 = gracilis.
convolvulacea, *Kurz, in Journ. Bot.* xi. (1873) 195.—Burma.
cordata, Hassk. in Retzia, i. 9 = Campanumœa javanica.
foetens, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 16.—Reg. Himal.
gracilis, Hook. f. & Thoms. in Illustr. Himal. Pl. t. 16 A = Leptocodon gracilis.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 431.—Reg. Himal.
inflata, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 13 = Campanumœa inflata.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 192.—Japon.

CODONOPSIS:—
javanica, *Hook. f. & Thoms. in Illustr. Himal. Pl.* t. 16 B = Campanumœa javanica.
lanceolata, *Benth. & Hook. f. Gen.* ii. 558; *Trans. in Act. Hort. Petrop.* vi. (1879) 46.—As. temp.
leucocarpa, Miq. Fl. Ind. Bat. ii. 566 = Campanumœa celebica.
luridа, Lindl. Bot. Reg. (1839) Misc. 82 = rotundifolia.
ovata, *Benth. in Royle, Illustr. Bot. Himal.* 253. t. 60. f. 3.—Reg. Himal.
parviflora, Wall. Cat. n. 1800 = Campanumœa parviflora.
purpurea, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 105.—Reg. Himal.
rotundifolia, *Royle, Illustr. Bot. Himal.* 254. t. 62.—Reg. Himal.
subsimplex, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 16.—Reg. Himal.
thalictrifolia, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 106.—Reg. Himal.
truncata, Wall. Cat. n. 1801 = Campanumœa celebica.
viridiflora, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 496.—As. or.
virides, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 103.—Reg. Himal.

CODONORAPHIA, Oerst. in Videnak. Selsk. Skr. V. v. (1859) 82; (Gen. Centr. Am. 8) = **Pentaraphia**, Lindl. (Gesnerac.).

CODONORCHIS, Lindl. Gen. et Sp. Orch. 410 (1840) = **Pogonia**, Juss. (Orchid.).
Lessonii, Lindl. l. c. 411 = P. tetraphylla.
Poeppigii, Lindl. l. c. 410 = P. tetraphylla.

CODONOSTIGMA, Klotzsch, ex Benth. in DC. Prod. vii. 709 (1839) = **Scyphogyne**, Brongn. (Ericac.). *Erinus*, Klotzsch, ex Benth. l. c.—Afr. austr.

CODORIOCALYX, Hassk.—Cf. Linnæa, xv. (1841) Lit. 80, 81 = CODARIOCALYX, Hassk. = **Desmodium**, Desv. (Legumin.).

CODVLIS, Rafin. in Am. Monthly Mag. (1819) 191 = **Nicotiana**, Linn. (Solanac.).

COELACHNE, R. Br. Prod. 187 (1810). *GRAMINEÆ*, Benth. & Hook. f. iii. 1156.
brachiata, *Munro, ex Benth. Fl. Austral.* vii. 626, in obs.—Zeylan.; China.
infirma, *Buese, in Miq. Pl. Jungh.* 350.—Java.
perpusilla, *Thw. Enum. Pl. Zeyl.* 373.—Zeylan.
pulchella, *R. Br. Prod.* 187.—Burma; Austral.
simplicissacha, Munro, ex Benth. in Journ. Linn. Soc. xix. (1881) 98 = brachiata.
subulifolia, F. Muell. Fragm. v. 208 = Micraira subulifolia.

COELACHYRUM, Nees, in Linnæa, xvi. (1842) 221 = **Eragrostis**, Beauv. (Gramin.).
brevifolium, Nees, l. c. = E. Coelachyrum.

COELANDRIA, Fitzg. Austral. Orch. i. pt. vii. t. 2 (1889) = **Dendrobium**, Sw. (Orchid.).
Smilliæ, Fitzg. l. c. = D. Smillieæ.

COELANTHE, Griseb. Gen. et Sp. Gent. 297, in syn. (1839) = COILANTHA, Borckh. = **Gentiana**, Linn.

COELANTHIUM, Sond. in Harv. & Sond. Fl. Cap. i. 147 (1859) = seq.

COELANTHIUM, E. Mey. ex Fenzl, in Ann. Wien. Mus. i. (1836) 358; iii. (1838) 367. *FICOIDEÆ*, Benth. & Hook. f. i. 857.
grandiflorum, *E. Mey. ex Fenzl, l. c.*—Afr. austr.
parviflorum, *Fenzl, l. c.* 366.—Afr. austr.

COELANTHUS, Willd. ex Schult. f. Syst. vii. p. xlvi (1830) = **Lachenalia**, Jacq. (Liliac.).
complicatus, Willd. l. c. in admot. = L. reflexa.

COELAS, Dulac, Fl. Hautes-Pyr. 308 (1867) = **Potentilla**, Tourn. (Rosac.).
procumbens, Dulac, l. c. = P. Sibbaldi.

COELEBOGYNE, J. Sm. in Proc. Linn. Soc. i. (1839) 41; Trans. Linn. Soc. xviii. (1841) 511. t. 26 = **Alchornea**, Sw. (Euphorb.).
ilicifolia, J. Sm. l. c. 513. t. 26 = A. ilicifolia.

COELESTINA, Cass. in Bull. Soc. Philom. (Jan. 1817) 10 = CAELESTINA, Cass. = **Ageratum**, Linn. (Compos.).

COELESTINA, Hill, Hort. Kew. 3 (1769) ; Veg. Syst. ii. 60** = **Aster**, Tourn. (Compos.).

COELESTINIA, Endl. Gen. 866 (1838) = CAELESTINA, Cass. = **Ageratum**, Linn. (Compos.).

COELIA, Lindl. Gen. et Sp. Orch. 86 (1830). *ORCHIDEAE*, Benth. & Hook. f. iii. 508.
BOTHRIOCHILUS, Lem. Illustr. Hortic. iii. (1856) Misc. 30.
Baueriana, *Lindl. Gen. et Sp. Orch.* 86.—Mexic.
bella, *Reichb. f. in Walp. Ann* vi. 218.—Guatemala.
Galeottiana, *Van Houtte, ex Heynh. Nom.* ii. 152, nomen.—Mexic.
glicinila, *Van Houtte, l. c.*, nomen.—Am. austr.
guatemalensis, *Reichb. f. in Walp. Ann.* vi. 219.—Guatemala.
macrostachya, *Lindl. in Benth. Pl. Hartw.* 92.—Mexic.
triptera, G. Don, ex Steud. Nom. ed. II. i. 394 = Baueriana.

COELIDIUM, Vog. ex Walp. in Linnaea, xiii. (1839) 472. *LEGUMINOSAE*, Benth. & Hook. f. i. 473.
Bowei, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 454.—Afr. austr.
bullatum, *Benth. l. c.* 453.—Afr. austr.
ciliare, *Vog. ex Walp. in Linnaea*, xiii. (1839) 472 — Afr. austr.
muriltioides, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 455.—Afr. austr.
roseum, *Benth. l. c.* 454.—Afr. austr.
Thunbergii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 24. —Afr. austr.
Vogelii, *Walp. in Linnaea*, xiii. (1839) 472.—Afr. austr.

COELINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid ?).

COELIOPSIS, Reichb. f. in Gard. Chron. (1872) 9 *ORCHIDEAE*, Benth. & Hook. f. iii. 477.
hyacinthosma, *Reichb. f. l. c.*—Panama.

COELOCARPUM, Balf. f. in Proc. Roy. Soc. Edinb xii. (1884) 90. *VERBENACEAE*.
socotranum, *Balf. f. l. c.* 91.—Ins. Socotra.

COELOCHLOA, Hochst. ex Steud. Nom. ed. II. i. 394 (1840) = COELACHYRUM, Nees = **Eragrostis**, Beauv. (Gramin.).
brevifolia, Hochst l. c. = E. Coelachyrum.

COELOCLINE, A. DC. in Mem. Soc. Phys. Genev. v. (1832) 208 = **Xylopia**, Linn. (Anonac.).
acutiflora, A. DC. l. c. = X. scutiflora.
lucida, A. DC. l. c. 209 = X. longifolia.
oxypetala, A. DC. l. c. = X. scutiflora ?
parviflora, A. DC. l. c. = X. parviflora.
polycarpa, A. DC. l. c. = X. polycarpa.

COELOCOCCUS, H. Wendl. in Bonplandia, x. (1862) 199 = **Metroxylon**, Rottb. (Palm.).
vitiensis, H. Wendl. l. c. = M. vitiense.

COELODEPAS, Hassk. in Flora, xl. (1857) 581. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 313.
KOILODEPAS, Hassk. in Versl. en Med. Kon. Akad. Wetensch. iv. (1855) 139.
bantamense, *Hassk. in Versl. en Med. Kon. Acad. Wetensch.* iv. (1855) 140 (Koilodepas).—Ins. Bantam.
calycinum, *Bedd. Forest. Man. Bot.* 207.—Ind. or.
Wallichianum, *Benth. in Hook. Ic. Pl.* t. 1288.—Ins. Penang.

COELODISCUS, Baill. Etud. Gén. Euph. 298 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 317.
dispar, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 244.—Ind. or.
eriocarpoides, Kurz, l. c. = Mallotus decipiens.
eriocarpus, *Kurz, l. c.*—Ind. or.
glabriusculus, *Kurz, For. Fl. Brit. Burma*, ii. 393.—Burma.

COELODISCUS:—
hirsutulus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 248.—Burma.
lappaceus, *Kurz, l. c.* 244 ; *et For. Fl. Brit. Burma*, ii. 392.—Burma.
loogipes, *Kurz, ll. cc.* 244, 393.—Burma.
montanus, *Muell. Arg. in DC. Prod.* xv. ii. 759.—Ins. Penang.
speciosus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 164.—Ind. or.
Thunbergianus, *Muell. Arg. in DC. Prod.* xv. ii. 758. —Zeylan.

COELOGLOSSUM, Hartm. Handb. Skand. Fl. ed. I. 329 (1820) = **Habenaria**, Willd (Orchid.).
acuminatum, Lindl. Gen. et Sp. Orch. 302 = H. lacertifera.
albidum, Hartm. Handb. Skand. Fl. ed. I. 329 = H. albida.
alpinum, Schur. in Verh. Siebenb. Ver. Naturw. ii. (1851) 199 ; iv. (1853) 72 = H. cucullata.
bracteatum, Parl. Fl. Ital iii. 409 = H. viridis.
brevifolium, Lindl. Gen. et Sp. Orch. 302 = H. cubitalis.
cernuum, Reichb. f. in Bonplandia, iii. (1855) 250 = H. stenostachya.
cordatum, Nym. Syll. 359 = H. cordata.
densiflorum, Hartm. ex Willk. & Lange, Prod. Fl. Hisp. i. 164 = H. intacta
densum, Lindl. Gen. et Sp. Orch. 302 = H. stenostachya.
Erdingeri, Kerner, in Oestr. Bot. Zeitschr. xiv. (1864) 140 = H. viridis.
lacertiferum, Lindl. Gen. et Sp. Orch. 302 = H. lacertifera.
luteum, Dalz. in Hook. Kew Journ. ii. (1850) 263.—Ind. or.
Manerii, Reichb. f. in Linnaea, xli. (1877) 54.—Reg. Himal.
persityloides, Reichb. f. in Bonplandia, iv. (1856) 321 — H. stenostachya.
purpureum, Schur, Enum. Pl. Transs 646 = H. cucullata.
satyrioides, Nym. Syll 359.—Tauria.
secundum, Lindl. Bot. Reg. sub t. 1701.—Ind. or.
tenue, Lindl. l. c. = H. stenostachya.
Vaillantii, Schur, Enum. Pl. Transs. 645 = H. viridis.
viride, Hartm. Handb. Skand. Fl. ed. I. 329 = H. viridis.

COELOGYNE, Lindl. Collect. sub t. 38 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 518
ACANTHOGLOSSUM, Blume, Bijdr. 381 (1825).
BOLBORCHIS, Zoll. & Mor. Verz. Pfl. 89 (1846).
GOMPHOTRIUS, Wall. ex Lindl. Gen. et Sp. Orch. 43 (1830).
NEOGYNA, Reichb. f. in Bot. Zeit. x. (1852) 981.
PLEIONE, D. Don, Prod. Fl. Nep. 36 (1825).
advena, *Par. & Reichb. f. Otia Bot. Hamb.* 47.—Burma.
alba, Reichb. f. in Walp. Ann. vi. 238 = Otochilus alba.
angustifolia, A. Rich. in Ann. Sc. Nat. Ser. II. xv. (1841) 16. t. 6 = odoratissima.
angustifolia, Wight, Ic. t. 1641 = breviscapa.
apiculata, Reichb. f. in Walp. Ann. vi. 225 = Panisea tenuilala.
Arthuriana, Reichb. f. in Gard. Chron. (1881) i. 40 = maculata.
articulata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota articulata.
asperata, *Lindl. in Journ. Hort. Soc.* iv. (1849) 221.—Malaya.
assamica, Linden & Reichb. f. in C. Koch, Allg. Gartenz. xxv. (1857) 403 = fuscescens.
barbata, *Lindl. ex Grif. Itin. Not.* 73.—Reg. Himal.
biflora, *Parish, ex Reichb. f. in Gard. Chron.* (1865) 1086.—Burma.
bilamellata, *Lindl. Fol. Orchid. Coelog.* 14.—Ins. Philipp.
birmanica, *Reichb. f. in Gard. Chron.* (1882) ii. 840. —Burma.
brachyptera, *Reichb. f. l. c.* (1881) ii. 6.—Burma.
breviscila, *Lindl. Fol. Orchid. Coelog.* 7.—Reg. Himal.
breviscapa, *Lindl. l. c.* 4.—Ind. or.
brunnea, Lindl. in Gard. Chron. (1848) 71 = fuscescens.
calcarata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota calcarata.

COELOGYNE:—
camelostalix, Reichb. f. in Walp. Ann. vi. 238 = Pholidota camelostalix.
carnea, Reichb. f. l. c. 237 = Pholidota carnea.
caulescens, *Grif. Notul.* iii. 282.—Malacca.
chinensis, Reichb. f. in Walp. Ann. vi. 237 = Pholidota chinensis.
chloroptera, *Reichb. f. in Gard. Chron.* (1883) i. 466. —Ins. Philipp.
ciliata, *Bonall, ex Naves, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 237.—Ins. Philipp.
clypeata, Reichb. f. in Walp. Ann. vi. 237 = Pholidota clypeata.
conchoidea, Reichb. f. l. c. = Pholidota conchoidea.
conferta, *Hort.—Cf. Gard. Chron.* (1875) i. 314.— Ind. or.
Convallariae, *Par. & Reichb. f. in Flora*, lv. (1872) 277.—Reg. Himal. ; Burma.
corniculata, *Reichb. f. in Gard. Chron.* (1865) 746.— Assam !
cornuta, Lindl. Bot. Reg. (1841) Misc. 83 = Trichosma suavis.
corrugata, *Wight, Ic.* t. 1630.—Ind. or.
corymbosa, *Lindl. Fol. Orchid. Coelog.* 7.—Reg. Himal.
cristata, *Lindl. Collect. Bot.* sub t. 38.—Reg. Himal.
Croockewitii, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1858) 488.—Java.
crotalina, Reichb. f. in Walp. Ann. vi. 238 = Pholidota crotalina.
Cumingii, *Lindl. Bot. Reg.* (1843) *Misc.* 76.—Malaya.
Cyenoches, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 147.—Burma.
cymbidioides, *Reichb. f. in Walp. Ann.* vi. 239.—Ins. Moluce.
Dayana, *Reichb. f. in Gard. Chron.* (1884) i. 826.— Borneo.
decora, *Wall. ex Voigt, Hort. Suburb. Calc.* 621, nomen.—Hab. ?
diphylla, *Lindl. Fol. Orchid. Coelog.* 15.—Ind. or.
ciata, *Lindl. in Wall. Cat. n.* 1959.—Reg. Himal.
fimbriata, *Lindl. Bot. Reg.* t. 986.—Reg. Himal.
flaccida, *Lindl. in Wall. Cat. n.* 1961.—Reg. Himal.
flavida, *Hook. f. ex Lindl. Fol. Orchid. Coelog.* 10.— Reg. Himal.
fuligincsa, *Lodd. ex Hook. Bot. Mag.* t. 4440.—Reg. Himal.
fusca, Reichb. f. in Walp. Ann. vi. 236 = Otochilus fusca.
fuscescens, *Lindl. in Wall. Cat.* n. 1962.—Malaya.
fusco-lutea, *Teijsm. & Binn. Cat. Hort. Bog.* 378, nomen.—Hab. !
Gardneriana, *Lindl. in Wall. Pl. As. Rar.* i. 88. t. 88. —Reg. Himal.
gibbosa, Reichb. f. in Walp. Ann. vi. 237 = Pholidota gibbosa.
glandulosa, *Lindl. Fol. Orchid. Coelog.* 6.—Ind. or.
globosa, Reichb. f. in Walp. Ann. vi. 236 = Pholidota globosa.
Goweri, *Reichb. f. in Gard. Chron.* (1869) 448.— Reg. Himal.
graminifolia, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 146.—Burma.
Hookeriana, *Lindl. Fol. Orchid. Coelog.* 14.—Reg. Himal.
Huettneriana, *Reichb. f. in Flora*, lv. (1872) 277.— Burma.
humilis, *Lindl. Collect. Bot.* sub t. 37.—Reg. Himal.
imbricata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota imbricata.
incrassata, *Lindl. Gen. et Sp. Orch.* 40.—Java.
interrupta, *Lindl. ex Heynh. Nom.* ii. 153, nomen.— Ind. or.
javanica, *Lindl. Fol. Orchid. Coelog.* 17.—Java.
khasyana, Reichb. f. in Walp. Ann. vi. 238 = Pholidota articulata.
lactea, *Reichb. f. in Gard. Chron.* (1885) i. 692.— Burma.
Lagenaria, *Lindl. Fol. Orchid. Coelog.* 15.— Reg. Himal.
lentiginosa, *Lindl. l. c.* 3.—Burma.
ligulata, *Teijsm. & Binn. Cat. Hort. Bog.* 46.— Java.
longecillata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 16.—Ind. or.
longifolia, *Lindl. Gen. et Sp. Orch.* 42.—Java.
longipes, *Lindl. Fol. Orchid. Coelog.* 10.— Reg. Himal.

COELOGYNE:—
loricata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota loricata.
Lowii, Paxt. Mag. Bot. xvi. (1849) 225 = asperata.
macrophylla, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 241.—Sumatra.
maculata, *Lindl. Gen. et Sp. Orch.* 43.—Reg. Himal.
marmorata, *Reichb. f. in Linnaea*, xli. (1877) 116.—Ins. Philipp.
Massangeana, *Reichb. f. in Gard. Chron.* (1878) ii. 684.—Malay.
Mayeriana, *Reichb. f. l. c.* (1877) ii. 184.—Hab.?
media, *Wall. ex Voigt, Hort. Suburb. Calc.* 621, nomen.—Ind. or.
micrantha, *Lindl. in Gard. Chron.* (1855) 173.—Reg. Himal.; Burma.
minista, *Lindl. Gen. et Sp. Orch.* 42.—Java.
nervulosa, Reichb. f. in Walp. Ann. vi. 236 = Pholidota nervosa.
nervosa, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 16 = corrugata.
nervosa, Wight, Ic. t. 1688 = glandulosa.
nigrescens, *P. Don, ex Loud. Hort. Brit. Suppl.* iii. 590, nomen.—Ind. or.
nitida, *Lindl. in Wall. Cat.* n. 1954.—Reg. Himal.
ocellata, *Lindl. Gen. et Sp. Orch.* 40.—Ind. or.
ocellata, Lindl. in Wall. Cat. n. 1958 = nitida, ochracea, cristata.
ochracea, *Lindl. Bot. Reg.* (1846) t. 69.—Reg. Himal.
odoratissima, *Lindl. in Wall. Cat.* n. 1960.—Ind. or.
ovalis, *Lindl. Bot. Reg.* (1838) *Misc.* 91.—Reg. Himal.
patiala, Reichb. f. in Walp. Ann. vi. 238 = Pholidota imbricata.
pandurata, *Lindl. in Gard. Chron.* (1853) 791.—Malaya.
Papagena, Reichb. f. in Bot. Zeit. xx. (1862) 214 = micrantha.
Parishii, *Hook. f. in Bot. Mag.* t. 5823.—Burma.
parviflora, Lindl. Gen. et Sp. Orch. 44 = Panisea parviflora.
peltastes, *Reichb. f. in Gard. Chron.* (1880) ii. 296.—Borneo.
Pholas, Reichb. f. in Walp. Ann. vi. 237 = Pholidota Pholas.
pilosissima, Planch. Hort. Donat. 144 = ovalis.
plantaginea, *Lindl. in Gard. Chron.* (1855) 20.—Ind. or.
porrecta, Reichb. f. in Walp. Ann. vi. 236 = Otochilus porrecta.
praecox, *Lindl. Collect. Bot.* sub t. 37.—Reg. Himal.
prolifera, *Lindl. in Wall. Cat.* n. 1956.—Reg. Himal.
psittacina, *Reichb. f. Xenia Orch.* ii. 141.—Ins. Moluce.
pulverula, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 306.—Sumatra.
pumila, Reichb. f. in Walp. Ann. vi. 236 = Dendrochilum pumilum.
punctulata, Lindl Collect. Bot. sub t. 33 = nitida.
recurva, Reichb. f. in Walp. Ann. vi. 237 = Pholidota recurva.
Reichenbachiana, *T. Moore, in Gard. Chron.* (1868) 1210.—Burma.
Rhodoana, Reichb. f. l. c. (1867) 901 = Rossiana.
rhombophora, *Reichb. f. in Linnaea*, xli. (1877) 116.—Ins. Manilla.
rigida, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 146.—Burma.
Rochussenii, *De Vriese, Illustr. Orch.* [t. 2. et t. 11. f. 6].—Java.
Rossiana, *Reichb. f. in Gard. Chron.* (1884) ii. 808.—Burma.
rubra, Reichb. f. in Walp. Ann. vi. 238 = Pholidota rubra.
Rumphii, *Lindl. Fol. Orchid. Coelog.* 14.—Malaya.
salmonicolor, *Reichb. f. in Gard. Chron.* (1888) ii. 328.—Ins. Sundaic.
Schilleriana, *Reichb. f. in C. Koch, Allg. Gartenz.* xxvi. (1858) 189.—Burma.
simplex, *Lindl. Fol. Orchid. Coelog.* 18.—Ind. or.
speciosa, *Reichb. f. in Gard. Chron.* (1888) i. 806.—Ins. Philipp.
speciosa, *Lindl. Gen. et Sp. Orch.* 39.—Malaya.
sulphurea, *Reichb. f. in Bonplandia*, v. (1857) 48.—Malaya.

COELOGYNE:—
testacea, *Lindl. Bot. Reg.* (1842) *Misc.* 88.—Malaya. Thuniana, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 145.—Nepal.
tomentosa, *Lindl. Fol. Orchid. Coelog.* 8.—Malaya.
trifida, Reichb. f. in Hamb. Gartenz. xix. (1863) 646 = odoratissima.
trinervis, *Lindl. in Wall. Cat.* n. 1955.—Burma.
triotes, *Reichb. f. in Hamb. Gartenz.* xviii. (1862) 84.—Ins. Philipp.
triplicatula, Reichb. f. in Bot. Zeit. xxii. (1864) 415 = fuligmosum?
tristera, A. Brongn. in Duperr. Voy. Coq. Bot. 201 = Epidendrum cacspitosum, *Poepp. & Endl.*
trisaccata, Griff. Notul. iii. 400 = Gardneriana.
undulata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota rubra.
undulata, *Wall. ex Voigt, Hort. Suburb. Calc.* 621, nomen.—Ind. or
uniflora, *Lindl. in Wall. Cat.* n. 1966.—Reg. Himal.; Burma.
ustulata, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 146.—Burma.
ventricosa, Reichb. f. in Walp. Ann. vi. 237 = Pholidota ventricosa.
viscosa, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 218.—Reg. Himal
Wallichiana, Lindl. Gen. et Sp. Orch. 43 = praecox.

COELONEMA, Maxim. in Bull. Acad. Pétersb. xxvi. (1880) 428. *CRUCIFERAE.*
draboides, *Maxim. l. c.* 424.—China.

COELOPLEURUM, Ledeb. Fl. Ross. ii. 861 (1844) = **Archangelica**, Hoffm. (Umbellul.).
Gmelini, Ledeb. l. c. = A. Gmelini.

COELOPYRUM, Jack, in Malay. Misc. ii. (1822) vii. 65. *INC. SED.*
coriaceum, *Jack, l. c.*—Malaya.

COELORACHIS, Brongn. in Duperr. Voy. Coq. Bot. 64. t. 14 (1829) = **Rottboellia**, Linn. (Gramin.).
hirsuta, Brongn. l. c. 65, in obs. = R. hirsuta.
muricata, Brongn. l. c. 65. t. 14 = R. glandulosa.

COELOSPERMUM, Blume, Bydr. 994 (1826) (Coelospermum). *RUBIACEAE*, Benth. & Hook. f. ii. 119.
POGONOLOBUS, F. Muell. Fragm. i. 55 (1858).
Balansaeanum, *Baill. Adansonia*, xii. (1876–79) 236.—N. Caled.
barbatum, *Span. in Linnaea*, xv. (1841) 318.—Ins. Timor.
corymbosum, *Blume, ex DC. Prod.* iv. 468.—Java.
decipiens, *Baill. in Bull. Soc. Linn. Par* i. (1879) 218.—Austral.
Gmelini, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 62, sphalm. = Archangelica Gmelini.
paniculatum, *F. Muell. Fragm.* v. 19.—Austral.
reticulatum, *Benth. Fl. Austral.* iii. 425.—Austral.
scandens, *Blume, Bydr.* 994.—Malaya.

COELOSTEGIA, Benth. in Benth. & Hook. f. Gen i. 918 (1862). *MALVACEAE*, Benth. & Hook. f. i. c. Griffithii, *Benth. l. c.*—Malacca.

COELOSTELMA, Fourn. in Mart. Fl. Bras. vi. iv. 320 (1885). *ASCLEPIADEAE.*
refractum, *Fourn. l. c.*—Bras.

COLOSTYLIS, Torr. & Gray, ex Endl. Nov Stirp. Dec. 33 (1839) = **Spigelia**, Linn. (Loganiac.).
loganioides, Torr. & Gray, l. c. = S. loganioides.
texana, Torr. & Gray, Fl. N. Am. ii. 46 = S. texana.

COEMANSIA, March. in Bull. Acad. Belg. Sér. II. xlvii. (1879) 94. *ARALIACEAE.*
COUDENBERGIA, March. l. c. 514 (1879).
Warmingiana, *March. l. c.* 95.—Am. austr.

COENOLOPHIUM, Reichb. Consp. 143 (1828) = CENOLOPHIUM, Koch = **Selinum**, Linn. (Umbell.).

COENOTUS, Benth. & Hook. f. Gen. ii. 981 (1873), lapsu = CARNOTUS, Rafin. = **Erigeron**, Linn. (Compos.).

COERULINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophular.).
montana, Fourr. l. c. = V. montana.

COETOCAPNIA, Link & Otto, Ic. Pl. Rar. 35. t. 18 (1828) = **Bravoa**, La Llav. (Amaryllid.).
geminiflora, Link & Otto, l. c. = B. geminiflora.

COFFEA, Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth. & Hook. f. ii. 114.
HEXEPTA, Rafin. Sylva Tellur. 164 (1838).
LACHNASTOMA, Korth. in Ned. Arch. Kruidk. ii. ii. (1851) 201.
acuminata, *Ruiz & Pav. Fl. Per.* ii. 64.—Peruv.
Afzelii, *Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 174.—Afr. trop.
alpestris, Wight, Ic. t. 1040 = Webera lucens.
angustifolia, *Roxb. Hort. Beng.* 86; *Fl. Ind. i.* 541.—Malaya.
arabica, Benth in Hook. Niger Fl. 413 = liberica, stenophylla.
arabica, *Linn. Sp. Pl.* 172.—Arabia; Afr. trop.
australis, Vell. Fl. Flum. 62; ii. t. 13 = Faramea australis.
bengalensis, *Roxb. Hort. Beng.* 15; *Fl. Ind. i.* 540.—Ind. or.; Malaya.
bidentata, D. Dietr. Syn. Pl. i. 776 = microcarpa.
biflora, Vell. Fl. Flum. 63; ii. t. 16 = Coussarea biflora.
Boryana, D. Dietr. Syn. Pl i. 777 = Chasalia Boryana.
brachyphylla, *Endlk. in Bremen Abh.* viii. (1883) 390.—Ins. Nosaibé.
brasiliensis, Walp. Rep. ii. 477 = Rudgea lanceolata.
brevipes, *Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 172.—Afr. trop.
calycina, *Benth. in Hook. Journ. Bot.* iii. (1841) 282.—Guiana.
capitata, D. Dietr. Syn. Pl. i. 777 = Chasalia capitata.
capitata, Sieber, ex DC. Prod. iv. 531 = Chasalia divaricata?
Chamissonis, Hook. & Arn. Bot. Beech. Voy. 86 = Strumpfia haduana.
chasaliodes, D. Dietr. Syn. Pl. i. 777 = Chasalia divaricata?
ciliata, *Ruiz & Pav. Fl. Per.* ii. 65. t. 216.—Peruv.
clusiaefolia, D. Dietr. Syn. Pl. i. 777 = Chasalia clusiaefolia.
coriacea, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 201 = laurifolia.
corymbulosa, *Bertol. Fl. Guatimal.* 10.—Guatemala.
crassiloba, Benth. in Hook. Journ. Bot. iii. (1841) 233 = Rudgea Schomburgkiana.
cymosa, Willd. ex Roem. & Schult. Syst. v. 201 = Chasalia Fontanesii.
densiflora, *Blume, Bydr.* 965.—Java.
Deppeana, *Steud. Nom.* ed. II. i. 394.—Mexic.
didymocarpa, Hartl. ex DC. Prod. iv. 481 = Declieuxia psychotroides.
divaricata, Tausch, ex DC. l. c. 532 = Chasalia divaricata.
elliptica, Thw. Enum. Pl. Zeyl 154 = Byrsophyllum ellipticum.
elongata, *Korth. in Nederl Kruidk. Arch.* ii. ii. (1851) 254.—Sumatra.
erianthа, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 534 = Rudgea erianthа.
flavicans, Humb & Bonpl. ex Roem. & Schult. Syst. v. 201 = Faramea jasminoides.
floribunda, Mart. in Flora, xxiv. (1841) ii. Beibl. 85 = Ixora densiflora.
floribunda, *Miq. in Linnaea*, xxii. (1849) 804.—Bras.
Fontanesii, D. Dietr. Syn. Pl. i. 777 = Chasalia Fontanesii.
foveolata, *Ruiz & Pav. Fl. Per.* ii. 65. t. 216.—Peruv.
fragrans, *Wall. ex Hook. f. Fl. Brit. Ind.* iii. 154.—Ind. or.; Burma.
gardneriodes, Cham. in Linnaea, ix. (1834) 280 = Rudgea gardenioides.
glabra, *Korth. in Nederl. Kruidk. Arch.* ii. ii. (1851) 254.—Borneo.
grandifolia, Boj. ex Baker, Fl. Maurit. 152 = macrocarpa.
grandifolia, D. Dietr. Syn. Pl. i. 777 = Chasalia grandifolia.
grumeliodes, Wight, Ic. t. 1041 = Webera lucens.
guianensis, *Aubl. Pl. Gui.* i. 150. t. 57.—Guiana.
hirsuta, G. Don, Gen. Syst. iii. 581 = Cremaspora africana.

COFFEA :—
Horsfieldiana, Miq. Fl. Ind. Bat. ii. 306 = bengalensis.
Humblotiana, *Baill. in Bull. Soc. Linn. Par.* i. (1886) 514.—Ins. Comor.
hypoglauca, *Welw. ex Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 173.—Afr. trop.
indica, *Pair. Encyc. Suppl.* ii. 14.—Java.
jasminoides, Cham. in Linnæa, ix. (1834) 222 = Rudgea jasminoides.
jasminoides, *Welw. ex Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 175.—Afr. trop.
javanica, Blume, Cat. Gew. Buitenz. 46 = Chiococca javanica.
Jenkinsii, *Hook. f. Fl. Brit. Ind.* iii. 155.—Reg. Himal.
kaduana, Cham. & Schlecht. in Linnæa, iv. (1829) 33 = Strausia kaduana.
kisantuana, *Hook. f. Fl. Brit. Ind.* iii. 154.—Reg. Himal.
Krausiana, Hochst. in Flora, xxv (1842) 287 = Krausia floribunda.
lanceolata, Cham. in Linnæa, ix. (1834) 232 = Rudgea lanceolata.
lanceolata, Cham. & Schlecht. l. c. vi. (1831) 412 = Deppeana.
laurifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 372.—Venezuela.
laurifolia, Salisb. Prod. 62 = arabica.
laurina, Poir. in Lam. Encyc. Suppl. ii. 14 = Craterispermum laurinum.
laurina, *Smeathm. ex DC. Prod.* iv. 499.—Afr. trop.
lepidophloia, *Miq. Fl. Ind. Bat. Suppl.* 548.—Ins. Banca.
liberica, *Hiern*, in *Trans. Linn. Soc.* Ser. II. i. (1876) 171. t. 24.—Afr. trop.
longiloba, *Ruiz & Pav. Fl. Per.* ii. 66. t. 218.—Peruv.
luxoniensis, Cham. & Schlecht. in Linnæa, iv. (1829) 32 = Psychotria luxoniensis.
macrocarpa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 168.—Ins. Mascar.
macrophylla, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 344 = subsessilis.
madrensis, *Teijsm. & Binn. Cat. Hort. Bog.* 112.—Ins. Madur.
magnoliæfolia, Cham. in Linnæa, ix. (1834) 225 = Rudgea magnoliæfolia.
major, Cham. l. c. 226 = Rudgea major.
marginata, *Benth. Pl. Hartw.* 198.—N. Granat.
Mariniana, Cham. & Schlecht. in Linnæa, iv. (1829) 35 = Strausia Mariniana.
mauritiana, *Lam. Encyc.* i. 550.—Ins. Mascar.
melanocarpa, *Welw. ex Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 173.—Afr. trop.
meridionalis, Vell. Fl. Flum. 62; it. t. 14 = Conssarea meridionalis.
mexicana, *DC. Prod.* iv. 501.—Mexic.
microcarpa, DC. l. c. 499 = Perrottetii.
microcarpa, *Ruiz & Pav. Fl. Per.* ii. 66. t. 218.—Peruv.
minor, Cham. in Linnæa, ix. (1834) 227 = Rudgea Clausseniana.
Moka, Hort. ex Heynh. Nom. ii. 153 = arabica.
mosambicana, DC. Prod. iv. 500 = racemosa, *Lour.*
myrtifolia, *Roxb. Hort. Beng.* 15, nomen.—Ins. Maurit.
neurophylla, *Miq. Fl. Ind. Bat.* ii. 1078.—Java.
nitida, *Ruiz & Pav. Fl. Per.* ii. 65. t. 217.—Peruv.
nodosa, Cham. in Linnæa, ix. (1834) 228 = Rudgea nodosa.
novoguineensis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 259.—N. Guinea.
obovata, *Cham. & Schlecht. in Linnæa*, vii. (1832) 412.—Mexic.
obovata, D. Dietr. Syn. Pl. i. 777 = Chasalia divaricata.
occidentalis, Jacq. Enum. Pl. Carib. 16 = Faramea odoratissima.
occidentalis, Vell. Fl. Flum. 62; it. t. 12 = Faramea occidentalis.
odorata, *Forst. f. Prod.* 16.—Ins. Amicor.
oleifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 372.—N. Granat.
opulens, Forst. f. Prod. 16 = Pavetta opulina.
paniculata, *Aubl. Pl. Gui.* i. 152. t. 58.—Guiana.
parquioides, Cham. in Linnæa, ix. (1834) 224 = Rudgea parquioides.
parvifolia, Cham. l. c. 231 = Rudgea parvifolia.
pedunculata, *Roxb. Hort. Beng.* 86 ; *Fl. Ind.* i. 541.—Ins. Molucc.
Perrottetii, *Steud. ex Bush, DC. Prod. Index*, i. præf. p. ix.—Afr. trop.
porophylla, Vell. Fl. Flum. 62; it. t. 11 = Conssarea porophylla.

COFFEA :—
psychotrioides, D. Dietr. Syn. Pl. i. 777 = Gaertnera psychotrioides.
racemosa, *Lour. Fl. Cochinch.* 145.—Afr. trop.
racemosa, *Ruiz & Pav. Fl. Per.* ii. 64. t. 214.—Peruv.
rachiflorula, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 514.—Ins. Comor.
ramosa, Roem. & Schult. Syst. v. 108 = racemosa, *Lour.*
roscs, *Mor. & Sesse, ex DC. Prod.* iv. 499.—Mexic.
rupestris, *Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 174.—Afr. trop.
salicifolia, *Miq. Fl. Ind. Bat.* ii 307.—Java.
sambucina, Forst. f. Prod. 16 = Pavetta sambucina.
semisericea, *Celebr. ex Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 195.—Ind. or.
sessilis, Vell. Fl. Flum. 64 ; ii. t. 20 = Psychotria axillaris.
spicata, *H. B. & K. Nov. Gen. et Sp.* iii. 371. t. 286.—N. Granat.
stenophylla, *G. Don, Gen. Syst* iii. 581.—Abyssin.
stipulacea, *DC. Prod.* iv. 498.—Guiana.
stipulacea, Steud. Nom ed. II. i. 895 = seq.
stipulata, Vell. Fl. Flum. 63 ; ii. t. 17 = Bathysa stipulata.
subcordata, *Hiern*, in *Trans. Linn. Soc.* Ser. II. i. (1876) 174.—Afr. trop.
subsessilis, Benth. in Hook. Journ. Bot. iii. (1841) 282 = Mapouria subsessilis.
subsessilis, *Ruiz & Pav. Fl. Per.* ii. 64. t. 215.—Peruv.
sundana, *Miq. Fl. Ind. Bat.* ii. 306.—Java.
sylvestris, Willd. ex Roem. & Schult. Syst. v. 201 = mauritiana.
tenuiflora, *Benth. in Hook. Journ. Bot.* iii. (1841) 282.—Guiana.
tetrandra, Roxb. Hort. Beng. 85; Fl. Ind. i. 538 = Priamatomeris albidiflora.
travancorensis, *Wight & Arn. Prod.* 435.—Ind. or.
trifora, Vell. Fl. Flum. 16 = Pavetta trifora.
trifora, Moon, Cat. 15 = travancorensis.
truncata, Vell. Fl. Flum. 63 ; it. t. 15 = Faramea truncata.
umbellata, *Ruiz & Pav. Fl. Per.* ii. 64. t. 215.—Peruv.
umbellata, Vell. Fl. Flum. 63 ; it. t. 19 = Faramea salicifolia.
verticillata, *Ruiz & Pav. Fl. Per* ii. 66. t. 217.—Peruv.
verticillata, Vell. Fl. Flum. 63 ; it. t. 18 = Ixora verticillata.
viburnoides, Cham. in Linnæa, ix. (1834) 228 = Rudgea viburnoides.
volubilis, Blanco, Fl. Filip. ed. I. 157 = Morinda tinctoria.
vulgaris, Mœnch, Meth 504 = arabica.
Wightiana, *Wall. Cat.* n. 6246 ; *Wight & Arn. Prod.* 436.—Ind. or.
Zanguebarica, *Lour. Fl. Cochinch.* i. 145.—Afr. trop.

COGNIAUXELLA, Baill. in Bull. Soc. Linn. Par. i. (1884) 494 = **Cogniauxia**, Baill. (Cucurb.).

COGNIAUXIA, Baill. in Bull. Soc. Linn. Par. i. (1884) 493. *CUCURBITACEÆ.*
(COGNIAUXELLA, Baill. l. c. 494.
podolaema, *Baill. l. c.* 494.—Madag. ?

COGSWELLIA, Schult. in Roem. & Schult. Syst. vi. p. xlviii (1820) = **Peucedanum**, Tourn. (Umbellif.).
villosa, Schult. l. c. 588 = P. foeniculaceum.

COGYLIA, Molina, Sagg. Chil. ed. II. 300 (1810) = **Lavradia**, Ruiz & Pav. (Berberid.).
bitternata, Molina, l. c. = L. bitternata.
ternata, Molina, l. c. = L. trifoliata.
triternata, Molina, l. c. = L. triternata.

COHAUTIA, Endl. Ench. 278 (1841) = KOHAUTIA, Cham. & Schlecht. = **Oldenlandia**, Linn. (Rubiac.).

COHIBA, Rafin. Fl. Tellur. iii. 75 (1836) = **Wigandia**, H. B. & K. (Hydrophyllac.).
urens, Rafin. l. c. = W. pruriticea.

COHNIA, Kunth, Enum. Pl. v. 85 (1850) = **Cordyline**, Comm. (Lilieæ.).
floribunda, Kunth, l. c. 86 = Cord. floribunda,
macrophylla, Kunth, l. c. 87 = Cord. macrophylla.
mauritiana, Baker, in Journ. Linn. Soc. xiv. (1876) 546 = Cord. macrophylla.
neocaledonica, Baker, l. c. = Cord. neocaledonica.
parviflora, Kunth, Enum. Pl. v. 87 = Cord. floribunda.

COHNIA, Reichb. f. in Bot. Zeit. x. (1852) 926. *ORCHIDEÆ*, Benth. & Hook. f. iii. 477.
quekettioides, *Reichb. f. l. c.*—Nicaragua.

COILADENA, Rafin. Fl. Tellur. ii. 12 (1836) = **Ipomœa**, Linn. (Convolvulac.).
hyemalis, Rafin. l. c. = I. Horsfalliae.

COILANTHA, Borckh. in Roem. Arch. i. 1. (1796) 25 = **Gentiana**, Tourn.
asclepiadea, G. Don, Gen. Syst. iv. 186 = G. asclepiadea.
biloba, Bercht. & Presl, Rostl. i. Gentian. 6 = G. lutea.
Bursseri, Bercht. & Presl, l. c. = G. Burseri.
campanulata, G. Don, Gen. Syst. iv. 185 = G. punctata.
glauca, G. Don, l. c. = G. glauca.
Macinni, D. Don, ex G. Don, l. c. = G. calyculata.
pannonica, G. Don, l. c. = G. pannonica.
punctata, G. Don, l. c. = G. punctata.
purpurea, Borckh. in Roem. Arch. i. 1. (1796) 25 = G. purpurea.
Sessaei, D. Don, ex G. Don, Gen. Syst. iv. 185 = G. Sessaei.

COILANTHERA, Rafin. Sylva Tellur. 38 (1838) = **Cordia**, Linn. (Boragin.).
rotundifolia, Rafin. l. c. = Cordia rotundifolia.

COILMEROA, Endl. Gen. Suppl. iii. 99 (1848) = COLMEIROA, Reut. = **Seguriniega**, Juss. (Euphorbiac.).

COILONOX, Rafin. Fl. Tellur. ii. 98 (1836) = **Ornithogalum**, Tourn. (Liliac.).
albucoides, Rafin. l. c. = O. suaveolens.

COILOSPERMA, Rafin. Fl. Tellur. iii. 48 (1836) = **Desringia**, R. Br. (Amarantac.).
cordata, Rafin. l. c. = D. baccata.

COILOSTIGMA, Klotzsch, in Linnæa, xii. (1838) 284 = **Scyphopyne**, Bronghn. (Eriac.).
Dregeanum, Klotzsch, l. c. 285.—Afr. austr.
glabrum, Benth. in DC. Prod. vii. 708.—Afr. austr.
puberulum, Benth. l. c. = S. puberula.
tenuifolium, Klotzsch, in Linnæa, xii. (1838) 284.—Afr. austr.
Zeyherianum, Klotzsch, l. c.—Afr. austr.

COILOSTYLIS, Rafin. Fl. Tellur. iv. 37 (1836) = **Epidendrum**, Linn. (Orchid.).
emarginata, Rafin. l. c. = E. ciliare.
obtusifolia, Rafin. l. c. = E. lineare !

COILOTAPALUS, P. Br. Hist. Jamaic. 111 (1756) = **Cecropia**, Loefl. (Urticac.).

COINOCHLAMYS, T. Anders. ex Benth. & Hook. f. Gen. ii. 1091 (1876). *ACANTHACEÆ*, Benth. & Hook. f. l. c.
angolana, *S. Moore, in Journ. Bot.* xiv. (1876) 322.—Afr. trop.
hirsuta, *T. Anders. ex Benth. & Hook. f. Gen.* ii. 1091.—Afr. trop.

COINOGYNE, Less. in Linnæa, vi. (1831) 520 = **Jaumea**, Pers. (Compos.).
carnosa, Less. l. c. 521 = J. carnosa.

COIX, Linn. Gen. ed. I. 980 (1737). *GRAMINEÆ*, Benth. & Hook. f. iii. 1119.
LACRYMA, Medic. Phil. Bot. i. 177 (1789).
LACRYMA-JOB, Tourn. ex Linn. Gen. ed. I 980 (1737).
LACRYSMARIA, Heist. Syst. 12 (1748).
LITHAGROSTIS, Gaertn. Fruct. t. 7. t. 1 (1788).
SPHÆRIUM, Linn. Syst. ed. I (1735).
agrestis, *Lour. Fl. Cochinch.* 551.—Amboina ; China.
anguiste, Mill. Gard. Dict. ed. VIII. n. 2 = Tripsacum dactyloides.
aquatica, *Roxb. Hort. Beng.* 66 ; *Fl. Ind.* iii. 571.—Ind. or.
arundinacea, Koen. ex Willd. Sp. Pl. iv. 908 = Chionachne barbata.
arundinacea, Lam. Encyc. iii. 423 = C. Lacryma-Jobi.
barbata, Roxb. Hort. Beng. 66 ; Fl. Ind. iii. 569 = Chionachne barbata.

COIX :—
cypsoides, *C. Muell. in Bot. Zeit.* xix. (1861) 334.—Ind. or.
dactyloides, Linn. Sp. Pl. 972 = Tripsacum dactyloides.
exaltata, Jacq. ex Spreng. Syst. i. 228 = C. Lachryma-Jobi.
gigantea, Koen. ex Roxb. Hort. Beng. 66; Fl. Ind. iii. 570 = C. Lachryma-Jobi.
heterocilita, Roxb. l. c. 66; 572 = Polytoca bractesta.
Koenigii, Spreng. Syst. i. 228 = Chionache barbata.
Lacryma, Linn. Syst. ed. X. 1361 = C. Lachryma-Jobi. Lacryma-Jobi, *Linn. Sp. Pl.* 972 —As. trop.
Ma-Yuén, *Romanet, in Bull. Soc. Acclimat.* Sér. II. viii. (1881) 442.—Cochinch.; China; Burma.
ovata, Stokes, Bot. Mat. Med. iv. 842 = C. Lachryma-Jobi.
perdula, Saliab. Prod. 28 = C. Lachryma-Jobi.
pumila, *Roxb. Hort. Beng.* 66; *Fl. Ind.* iii. 572.—Ins. Maurit.
stigmatosa, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1855) 9.—Hab. ?

COLA, Schott & Endl. Meletem. 33 (1832). *STERCULIACEÆ*, Benth. & Hook. f. i. 218.
BICHEA, Stokes, Bot. Mat. Med. ii. 564 (1812).
COLARIA, Rafin. Sylva Tellur. 78 (1838).
COURTENIA, R. Br. in Denn. Pl. Jav. Rar. 236 (1844).
EDWARDIA, Rafin. Specch. i. 158 (1824); ex DC. Prod. ii. 92 (1825).
LUNANEA, DC. Prod. ii. 92 (1825).
SIPHONIOPSIS, Karst. Fl. Columb. i. 139. t. 69 (1858).
acuminata, *Schott & Endl. Meletem.* 33.—Afr. trop.
Afzelii, *Mast. in Oliver, Fl. Trop. Afr.* i. 223.—Afr. trop.
argutea, *Mast. l. c.* 224.—Afr. trop.
cauliflora, *Mast. l. c.* 221.—Afr. trop.
clavata, *Mast. l. c.* 222.—Afr. trop.
cordifolia, *[R. Br. in] Benn Pl. Jav. Rar.* 237.—Java.
digitata, *Mast. in Oliver, Fl. Trop. Afr.* i. 224.—Afr. trop.
ficifolia, *Mast. l. c.* 223.—Afr. trop.
gabonensis, *Mast. l. c.* 222.—Afr. trop.
grandiflora, *Schott & Endl. Meletem.* 33.—Hab. ?
heterophylla, *Schott & Endl. l. c.*; *Mast. in Oliver, Fl. Trop. Afr.* i. 223.—Afr. trop.
lanifolia, *Mast. in Oliver, l. c.* 222.—Afr. trop.
macrocarpa, Schott & Endl. Meletem. 33 = acuminata.
natalensis, *Oliver, in Hook. Ic. Pl.* t. 1390.—Afr. austr.
nitida, Schott & Endl. Meletem. 33 = acuminata.
quinqueloba, *Garcke, in Peters; Reise Mossamb. Bot.* 180.—Afr. trop.
tomentosa, Schott & Endl. Meletem. 33 = Sterculia tomentosa.

COLARIA, Rafin. Sylva Tellur. 78 (1838) = **Cola**, Schott & Endl. (Stercul.).
acuminata, Rafin. l. c. = Cola acuminata.
heterophyla, Rafin. l. c. = Cola heterophylla.

COLAX, Lindl. ex Spreng. Syst. iii. 737 (1826); et Bot. Reg. (1843) Misc. 50 = **Lycaste**, Lindl. (Orchid.).
aromaticus, Spreng. Syst. iv. Cur. Post. 807 = L. aromatica.
Barringtoniæ, Lindl. ex Spreng. Syst. iii. 737 = L. Barringtoniæ.
grandiflorus, Rafin. Fl. Tellur. ii. 85 = Bifrenaria Harrisoniæ.
Harrisoniæ, Lindl. ex Spreng. Syst. iii. 737 = Bifrenaria Harrisoniæ.
inconus, Lindl. Bot. Reg. (1843) Misc. 51.—Bras.
modestior, Reichb. f. in Hamb. Gartens. xvi. (1860) 14.—Bras.
pallidiflorus, Spreng. *l.* Tent. 29 = Maxillaria pallidiflora.
palmifolius, Lindl. ex Spreng. Syst. iii. 737 = Maxillaria palmifolia.
Parkeri, Spreng. f. Tent. 28 = Maxillaria Parkeri.
parvulus, Spreng. Syst. iv. Cur. Post. 807 = Bifrenaria parvula.
planiscapus, Lindl. Bot. Reg. (1843) Misc. 50.—Bras.
Pupulii, Linden & André, in Illustr. Hortic. xxvii. (1880) t. 959.—Bras.
racemosus, Spreng. f. Tent. 29 = Bifrenaria racemosa.
viridis, Lindl. Bot. Reg. (1843) Misc. 50 = L. viridis.

COLBERTIA, Salisb. Parad. Lond. sub t. 78 (1807) = **Dillenia**, Linn.
angustata, Wall. Cat. n. 948 = D. pentagyna.
coromandelina, DC. Syst. i. 435 = D. pentagyna.
floribunda, Wall. Cat. n. 950 = D. floribunda.
minor, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 579.—Java.
obovata, Blume, Bijdr. 6 = D. aurea.
scabrella, D. Don, Prod. Fl. Nep. 226 = D. scabrella.

COLCHICUM, Linn. Syst. ed. I (1735); Gen. ed. I. 107 (1737). *LILIACEÆ*, Benth. & Hook. f. iii. 821.
FOLINA, Pomel, Mat. Fl. All. 9 (1860).
HERMODACTYLOS, Reichb. Consp. 64 (1828).
MONOCARYUM, R. Br. in App. Voy. Denh. et Clapp. 248 (1826).
PALIDARIA, Salisb. Gen. Pl. Fragm. 53 (1865).
SYNSIPHON, Regel, in Act. Hort. Petrop. vi. (1879) 490.
aegyptiacum, Boiss. Diagn. Ser. I. v. 66 = montanum
aestivale, Bor. Fl. Centr. Fr. ed. III. ii. 612 = byzantinum.
aetnense, Tineo, in Guss. Fl. Sic. Syn. ii. 818 = neapolitanum.
agrippinum, *Baker, in Journ. Linn. Soc.* xvii. (1879) 425.—Hab. ?
Alberti, *Regel, in Act. Hort. Petrop.* viii. (1884) 647.—As. centr.
alpinum, *DC. Fl. Fr.* iii. 195.—Europ.
alpinum, Schimp. ex Nym. Consp. 742 = montanum.
amabile, *Heldr. in Atti Congr. Bot. Firense*, 1874 (1876) 27.—Graecia.
arenarium, Koch, Syn. Fl. Germ. ed. I. 836 = neapolitanum.
asturicum, Boiss. & Reut. Pugill. 97 = montanum.
atticum, Sprun. ex Tommas. in Flora, xxiii. (1840) 780 = Merendera attica
autumnale, *Linn. Sp. Pl.* 341.—Europ.
Balansae, Planch. in Ann. Sc. Nat. Sér. IV. iv. (1855) 146 = laetum.
Bertolonii, Stev. in Nouv. Mém. Soc. Nat. Mosc. vii. (1829) 268 = montanum.
Bisignani, Tenore, fide Janka, ex Baker, in Journ. Linn. Soc. xvii. (1879) 427 = Tenorii.
Bivonae, *Guss. Adnot. Cat. Pl. Bocc.* (1821) 4.—Europ. austr.
Bivonae, Tenore, Fl. Nap. Prod. App. v. 11 = Tenorii.
Bivonae, Willk. & Lange, Prod. Fl. Hisp. i. 194, ex parte = lusitanum.
Boissieri, *Orph. in Atti Cong. Bot. Firense*, 1874 (1876) 31.—Graecia.
brachyphyllum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 164.—Syria.
bulbocodioides, Bieb. Fl. Taur. Cauc. i. 293; iii. 581 = montanum.
bulbocodioides, Brot. Fl. Lusit. i. 597 = Merendera Bulbocodium.
Bulbocodium, Ker-Gawl. in Bot. Mag. sub t. 1028 = Bulbocodium vernum.
byzantinum, *Ker-Gawl. l. c.*—Graecia; As. Min.
byzantinum, Tenore, Fl. Nap. iii. 397 = Tenorii.
castroviejoi, Lazamb. in Bull. Soc. Bot. Fr. ii. (1855) 688 = neapolitanum.
candidum, Schott & Kotschy, ex Baker, in Journ. Linn. Soc. xvii. (1879) 429 = laetum.
catacuzenum, Heldr. ex Boiss. Fl. Orient. v. 165 = montanum.
caucasicum, Spreng. Syst. i. 149 = Merenders caucasica.
chinense, Haw. ex Kunth, Enum. Pl. iv. 139 = variegatum?
Clementei, Graells, in Mem. Acad. Cienc. Madr. ii. [1854] 688 = bulbocodioides.
communis, Neck. Delic. Gallo-Belg. i. 176 = autumnale.
corsicum, *Baker, in Journ. Linn. Soc.* xvii. (1879) 431.—Ins. Corsica.
crociflorum, *Sims, Bot. Mag.* t. 2673.—Europ. austr.
crociflorum, *Regel, in Act. Hort. Petrop.* vii. (1880) 385.—Kokan.
crociflorum, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 97 = montanum.
crocifolium, Boiss. Diagn. Ser. I. v. 67 = montanum.
Cupani, Guss. Fl. Sic. Prod. i. 452 = montanum.
Decaisnei, *Boiss. Fl. Orient.* v. 157.—Syria.
euboeum, Orph. in Atti Cong. Bot. Firenze, 1874 (1876) 28 = Bivonae.
fasciculare, *Boiss. Fl. Orient.* v. 163.—Syria.
fasciculare, R. Br. in Denh. & Clapp. Trav. 248 = montanum.

COLCHICUM :—
falcifolium, *Stapf, in Denkschr. Acad. Wien*, I. (1885) 19.—Persia.
floribundum, Lawson, Cat. 6; ex Saliab. in Trans. Hort. Soc. i. (1812) 329 = byzantinum.
Hausknechtii, *Boiss. Fl. Orient.* v. 157.—Persia.
Haynaldi, Heuff. in Verh. Zool.-Bot. Ges. Wien, viii (1858) 213 = autumnale.
hexapetalum, Pourr. ex Schult. f. Syst vii. 1521 = Merendera Bulbocodium.
hololophum, Coss. & Dur. in Bull. Soc. Bot. Fr. iv (1857) 489 = montanum.
illyricum, Frivald. ex Kunth, Enum. Pl. iv. 140 = speciosum.
illyricum, Stokes, *Bot. Mat. Med.* ii. 331.—As. Min.
Jankae, Freyn, in Oestr. Bot. Zeitschr. xxvii. (1877) 361 = neapolitanum.
Kochii, Parl. Fl. Ital. iii. 188 = neapolitanum.
Kotschyi, Boiss. Diagn. Ser. I. xiii. 28 = laetum.
laetum, *Sm. in Nouv. Mém. Soc. Nat. Mosc.* vii. (1829) 262. t. 13.—Orient.
latifolium, Graeb. Spicil. Fl. Rumel. ii. 378 = speciosum.
latifolium, Sibth. & Sm. Fl. Graec. iv. 43. t. 350 = Sibthorpii, byzantinum.
Levieri, *Janka, in Oestr. Bot. Zeitschr.* xxv. (1875) 82.—Italia.
libanoticum, *Ehrenb. ex Boiss. Fl. Orsent.* iii. 166.—Syria.
linguiatum, *Boiss. & Sprun. Diagn.* Ser. I. v. 66.—Europ.
longifolium, Castag. Cat. Pl. Marseille, 135 = neapolitanum.
lusitanum, *Brot. Phyt. Lusit.* ii. 211. t. 178, 174.—Lusitania.
luteum, *Baker, in Gard. Chron.* (1874) 33.—Reg. Himal.
Merendera, Ker-Gawl. in Bot. Reg. sub t. 541 = Merendera Bulbocodium.
micranthum, *Boiss. Fl. Orient.* v. 162.—Byzant.
minimum, Elwes, ex Baker, in Journ. Linn. Soc. xvii. (1879) 433 = montanum.
monogynum, Bieb. ex Schult. f. Syst. vii. 796 = Sternbergia caucasica.
montanum, Ali. Fl. Pedem. i. 117 = alpinum.
montanum, *Linn. Sp. Pl.* 342.—Europ austr.
montanum, Linn. l. c. partim = Merendera Bulbocodium.
montanum, Schousb. Vextr. Marok. 185 = autumnale.
multiflorum, Brot. Fl. Lusit. i. 597 = autumnale.
neapolitanum, Orph. ex Nym. Consp. 743 = parnassicum.
neapolitanum, *Tenore, Fl. Neap. Prod. App.* v. 11.—Italia.
nivale, Boiss. & Huet, ex Baker, in Journ. Linn. Soc. xvii. (1879) 433 = montanum.
orientale, Frivald. ex Kunth, Enum. Pl. iv. 143 = neapolitanum.
pannonicum, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 359 = autumnale.
Parkinsoni, Hook. f. Bot. Mag. t. 6090 = variegatum.
Parlatoris, *Orph. in Atti Congr. Bot. Firenze*, 1874 (1876) 32.—Graecia.
parnassicum, *Sart. Orph. & Heldr. in Boiss. Diagn.* Ser. II. iv. 122.—Graecia.
parviflorum, Bivona, ex Parl. Fl. Ital. iii. 101 = montanum.
parvulum, Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 343 = micranthum.
parvulum, Tenore, Viag. in Basilic. 120 = alpinum.
patens, Schultz, in Flora, ix. (1826) 132 = autumnale.
persicum, *Baker, in Journ. Linn. Soc.* xvii. (1879) 430.—Persia.
polyanthon, Ker-Gawl. in Bot. Mag. sub t. 1028 = autumnale.
polymorphum, Orph. in Atti Congr. Bot. Firenze, 1874 (1876) 29 = neapolitanum.
polyphyllum, Boiss. & Heldr. Diagn. Ser. II. iv. 121 = Stevenii.
procox, Spenn. Fl. Friburg. i. 215 = autumnale.
provinciale, H. Loret, in Bull. Soc. Bot. Fr. vi. (1859) 659 = neapolitanum.
pulchrum, *Herb. ex Baker, in Journ. Linn. Soc.* xvii. (1879) 425.—Graecia.
pusillum, Sieber, in Flora, v. (1822) 248 = montanum.
pyrenaicum, Pourr. in Mém. Acad. Toul. iii. (1788) 316 = Merendera Bulbocodium.

COLCHICUM :—
Ritchii, R. Br. in App. Denh. & Clapp. Trav. 241 = montanum.
Sibthorpii, *Baker, in Journ. Linn. Soc.* xvii. (1879) 427.—Graecia.
speciosum, *Stev. in Nouv. Mém. Soc. Nat. Mosc.* vii. (1839) 265. t. 15.—Reg. Caucas.
Steveni, *Kunth, Enum. Pl.* iv. 144.—Syria; Arab.
Szovitsii, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 24. —Reg. Caucas.
Tenorii, *Parl. Fl. Ital.* iii. 176.—Italia.
tessellatum, Hort. ex Baker, in Journ. Linn. Soc. xvii. (1879) 425 = agrippinum.
tessalatum, Link.—Cf. Kunth, Enum. Pl. iv. 139 = Bivonae.
tessellatum, Saltab. Prod. 286 = variegatum.
tessulatum, Mill. Gard. Dict. ed. VIII. n. 4 = variegatum.
Todarii, Parl. Fl. Ital. iii 178 = lusitanum.
trapezunticum, Boiss ex Baker, in Journ. Linn. Soc. xvii. (1879) 430 = polyphyllum.
transsilvanicum, Schur, Enum. Pl. Transs. 679 = byzantinum.
triphyllum, Kunze, in Flora, xxix. (1846) 755 = montanum.
Troodi, *Kotschy, in Unger & Kotschy, Cyprus*, 190.—Ins. Cyprus.
turcicum, *Yanka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 242.—Europ. or.
umbrosum, *Stev. in Nouv. Mem. Soc. Nat. Mosc.* vii. (1839) 268. t. 14.—Tauria.
Valesy, Tineo, in Guss. Fl. Sic. Syn. ii. 818 = montanum.
variegatum, Bivona, Sic. Pl. Cent. i. 27 = Bivonae.
variegatum, *Linn. Sp. Pl.* 342.—Europ. austr.
variegatum, Sieber, ex Baker, in Journ. Linn. Soc. xvii. (1879) 427 = Sibthorpii.
variepictum, Janka, in Oestr. Bot. Zeitschr. xxv. (1875) 83 = amabile.
vernale, Hoffm. Deutschl. Fl. ed. II. i. 174 = autumnale.
vernum, Kunth, Enum. Pl. iv. 141 = autumnale.
vernum, Kunth, l. c. 146 = Bulbocodium vernum.
vernum, Pall. ex Bieb. Fl. Taur. Cauc. i. 298 = montanum.
versicolor, Ker-Gawl. in Bot. Reg. t. 571 = Bulbocodium versicolor.
Visianii, Parl. Fl. Ital. iii. 175 = Bivonae.

COLDENIA, Linn. Dis. Dasa. 9 (1747); Amoen. Acad. i. 393 (1749). *BORAGINEAE*, Benth. & Hook. f. ii. 841
EDDYA, Torr. & A. Gray, in Pacif. Rail. Rep. ii. 170 (1855).
GALAPAGOA, Hook. f. in Trans. Linn. Soc. xx. (1847) 196.
GOLDENIA, Rasnach. Nom. ed. III. 43 (1797).
LOPOPHYLLUM, F. Muell. in Hook. Kew Journ. ix. (1857) 21.
MONOMESIA, Rafin. Fl. Tellur. iv. 87 (1836).
PTILOCALYX, Torr. & A. Gray, in Pacif. Rail. Rep. ii. 170. t. 8 (1855).
STROBOCARPUS, Torr. l. c. 169 (1855).
TIQUILIA, Pers. Syn. i. 157 (1805).
angolensis, *Welw. Apont.* 591.—Afr. trop.
atacamensis, *Phil. Fl. Atac.* 37.—Chili.
canescens, *DC. Prod.* ix. 559.—Am. bor. occ.
dichotoma, *Lehm. Asperifol.* 9.—Peruv.
Dombeyana, Benk, Ind. DC. Prod. iii. 108 = dichotoma.
Greggii, *A. Gray, Syn. Fl. N. Am.* ii. i. 182.—Am. bor. occ.
hispidissima, *A. Gray, in Proc. Am. Acad.* v. (1862) 340.—Am. bor. occ.
littoralis, *Phil. Fl. Atac.* 37.—Chili.
mexicana, *S. Wats. in Proc. Am. Acad.* xvii. (1882–83) 119.—Mexic.
Nuttallii, *Hook. in Hook. Kew Journ.* iii. (1851) 296.—Am. bor. occ.
Palmeri, *A. Gray, in Proc. Am. Acad.* viii. (1870) 292. —Am. bor. occ.
pentandra, Juss. ex Steud. Nom. ed. II. i. 396 = dichotoma.
procumbens, *Linn. Sp. Pl.* 125.—Ind. or.
tomentosa, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 120.—Am. bor. occ.

COLEA, Boj. Hort. Maurit. 220 (1837). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1051.

COLEA :—
TRIPINNA, Lour. Fl. Cochinch. 391 (1790).
TRIPINNARIA, Pers. Syn. ii. 178 (1807).
ULOMA, Rafin. Fl. Tellur. ii. 62 (1836).
caulisflora, DC. Prod. ix. 241 = floribunda.
Chapelieri, DC. l. c. 242 = decora.
Commersonii, DC. l. c. = floribunda.
decora, *Boj. Hort. Maurit.* 220, nomen; *DC. l. c.* 241. —Madag.
diacolor, *Seem. in Trans. Linn. Soc.* xxiii. (1860) 8.—Madag.
floribunda, *Boj. Hort. Maurit.* 220, nomen; *Lindl. Bot. Reg.* (1841) t. 19.—Madag.
hispidissima, *Seem. in Trans. Linn. Soc.* xxiii. (1860) 9.—Madag.
involucrata, *Boj. Hort. Maurit.* 221, nomen; *DC. Prod.* ix. 242.—Madag.
mauritiana, *Boj. l. c.* 220.—Ins. Mascar.
nitida, DC. Prod. ix. 242 = decora.
obtusifolia, DC. l. c. 241 = mauritiana.
parviflora, *Baker, in Journ. Linn. Soc.* xxi. (1885) 428. —Madag.
pedunculata, *Baker, Fl. Maurit.* 244.—Ins. Seychelles.
purpurascens, *Seem. in Trans. Linn. Soc.* xxiii. (1860) 8.—Madag.
ramiflora, DC. Prod. ix. 241 = mauritiana.
Seychellarum, *Seem. in Trans. Linn. Soc.* xxiii. (1860) 8.—Ins. Seychelles.
Sieberi, *A. DC. in DC. Prod.* ix. 240, in nota.—Hab.?
Telfairii, *Boj. Hort. Maurit.* 220.—Madag.
tetragona, *DC. Prod.* ix. 242.—Madag.
tripennata, *Seem. in Bonplandia*, iv. (1856) 128.—Guinea.
undulata, *Regel, Gortenfl.* (1870) 322.—Madag.?

COLEACHYRON, J. Gay, ex Boiss. Fl. Orient. v. 408 (1881) = **Carex**, Dill. (Cyperac.).
Oliverianum, J. Gay, l. c. = Carex illegitima.

COLEANTHERA, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) t. 4. *SPACRIDEAE*, Benth. & Hook. f. ii. 611.
MICRAEA, F. Muell. Fragm. iv. 95. t. 37 (1864).
caelophylla, *Benth. Fl. Austral.* iv. 150.—Austral.
myrtioides, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) t. 4.—Austral.
virgata, *Stschegl. l. c.* 5.—Austral.

COLEANTHUS, Seid. in Roem. & Schult. Syst. ii. 11, 376 (1817). *GRAMINEAE*, Benth. & Hook. f. iii. 1147.
SCHMIDTIA, Tratt. Fl. Austr. i. 12. t. 451 (1811).
WILIBALDA, Stemb. in Flora, ii. (1819) 6.
subtilis, *Seid. in Roem. & Schult. Syst.* ii. 376.—Europ.

COLEATAENIA, Griseb. in Goett. Abh. xxiv. (1879) 308 = **Panicum**, Linn. (Gramin.).
gynerioides, Griseb. l. c.—Reg. Argent.

COLEBROOKEA, Sm. Exot. Bot. ii. 111. t. 115 (1806). *LABIATAE*, Benth. & Hook. f. ii. 1180.
oppositifolia, Lodd. Bot. Cab. t. 487 = Ehholtzia polystachya.
oppositifolia, *Sm. Exot. Bot.* ii. 111. t. 115.—Ind. or.; Burma.
ternifolia, Roxb. Hort. Beng. 45; Fl. Ind. iii. 25 = oppositifolia.

COLEBROOKIA, Donn, Cat. Hort. Cantab. i. (1796) = **Globbe**, Linn. (Zingib.).
bulbifera, Donn, l. c. = G. marantina.

COLEBROOKIA, Spreng. Syst. ii. 713 = **Colebrookea**, Sm. (Labiat.).

COLEMA, Rafin. New Fl. Am. iii. 49 (1836) = **Corema**, D. Don (Empetrac.).
arenaria, Rafin. l. c. = Cor. Conradii.

COLENSOA, Hook. f. Fl. N. Zel. i. 156 (1853) = **Pratia**, Gaudich. (Campanulac.).
physaloides, Hook. f. l. c. 157.—N. Zel.

COLEOCOMA, F. Muell. in Hook. Kew Journ. ix. (1857) 19. *COMPOSITAE*, Benth. & Hook. f. ii. 393.
Centaurea, *F. Muell. l. c.*—Austral.

COLEOGYNE, Torr. in Smithson. Contrib. vi. (Plant. Fremont.) 8. t. iv. (1854). *ROSACEAE*, Benth. & Hook. f. i. 617.
ramosissima, *Torr. l. c.*—Calif.

COLEONEMA, Bartl. & Wendl. f. Diosm. 55. t. A (1824). *RUTACEAE*, Benth. & Hook. f. i. 289.
album, *Bartl. & Wendl. f. l. c.* 56.—Afr. austr.
album, E. Mey. Zwei Pfl. Docum. 174 = Diosma vulgaris.
capalatholdes, *A. Juss. in Mém. Mus. Par.* xii. (1825) 471.—Afr. austr.
Dregeanum, Presl, Bot. Bemerk. 31 = pulchrum.
filiformis, A. Juss. in Mém. Mus. Par. xii. (1825) 471 = pulchrum.
gracile, Eckl. & Zeyh. Enum. 106 = pulchrum.
juniperifolium, *Eckl. & Zeyh. l. c.*—Afr. austr.
juniperinum, Sond. in Harv. & Sond. Fl. Cap. i. 379 = juniperifolium.
pulchrum, *Hook. Bot. Mag.* t. 3840.—Afr. austr.
virgatum, Eckl. & Zeyh. Enum. 106 = pulchrum.
virgnianum, Eckl. & Zeyh. l. c. = juniperifolium.

COLEOPHORA, Miers, in Ann. Nat. Hist. Ser. II. vii (1851) 197 = **Daphnopsis**, Mart. (Thymel.).
gemmiflora, Miers, l. c. = D. Martii.

COLEOPHYLLUM, Klotzsch, in Otto & Dietr. Allg. Gartenz. viii. (1840) 185 = **Childanthus**, Herb. (Amaryll.).
Ehrenbergii, Klotzsch, l. c. = Childanthus Ehrenbergii

COLEOSANTHUS, Cass. in Bull. Soc. Philom. (1817) 67; et Dict. Sc. Nat. x. (1818) 36 = **Britokallia**, Ell (Compos.).
Cavanillesii, Cass. in Dict. Sc. Nat. x. 37 = B Cavanillesii.
tiliaefolius, Cass. l. c. xxiv. 519 = Eupatorium macrophyllum.

COLEOSPADIX, Becc. in Ann. Jard. Buitenz. ii. (1885) 90. *PALMAE*.
litigiosa, *Becc. l. c.*—N. Guinea.
onanensis, *Becc. l. c.*—N. Guinea.

COLEOSTACHYS, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 329. *MALPIGHIACEAE*, Benth. & Hook. f. i. 258.
genipaefolia, *A. Juss. l. c.*—Bras.
hypoleuca, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 125 = Blepharandra hypoleuca.
spicata, Juss. in Arch. Mus. Par. iii. (1843) t. 5 = genipaefolia.
vestita, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 124 = Diacidia vestita.

COLEOSTEPHUS, Cass. in Dict. Sc. Nat. xli. 43 (1826) = **Chrysanthemum**, Linn. (Compos.).
Clausonis, Pomel, Nouv. Mat. Fl. Atl. 39.—Algeria.
hybridus, Lange, in Kjoeb. Vidensk. Meddel. (1861) 77 (Pugill. 127) = Chrys. Myconis.
macrotus, Dur. in Duch. Rev. Bot. i. (1846–46) 368 = Chrys. macrotum.
multicaulis, Dur. l. c. 364 = Chrys. multicaule.
Myconis, Cass. in Dict. Sc. Nat. xli. 43 = Chrys. Myconis.

COLEOSTYLES, Benth. & Hook. f. Gen. ii. 535 [bis] (1873) = seq.

COLEOSTYLIS, Sond. in Lehm. Pl. Preiss. i. 391 (1845) = **Levenhookia**, R. Br. (Stylid.).
Preissii, Sond. l. c. = L. Preissii.
Sonderi, F. Muell. in Trans. Phil. Soc. Vict. i. (1856) 46 = L. Sonderi.
umbellulata, Sond. in Lehm. Pl. Preiss. i. 391 = L. stipitata.

COLEOTRYPE, C. B. Clarke, in DC. Monog. Phan. iii. 288. t. 8 (1881). *COMMELINACEAE*, Benth. & Hook. f. iii. 851.
Goudotii, *C. B. Clarke, l. c.* 240.—Madag.
madagascarica, *C. B. Clarke, l. c.* 289.—Madag.
natalensis, *C. B. Clarke, l. c.*—Afr. austr.

COLETIA, Vell. Fl. Flum. 83 (1829) = **Mayaca**, Aubl. *maidida*, Vell. l. c. 83; t. t. 70 = M. Sellowiana.

COLEUS, Lour. Fl. Cochinch. ii. 372 (1790). *LABIATAE*, Benth. & Hook. f. ii. 1176.
Mitha, Chap. ex Benth. Lab. Gen. et Sp. 52 (1832).
scuminatus, *Benth. in Linnaea*, vi. (1831) 81.—Ins. Philipp.
africanus, *Benth. Lab. Gen. et Sp.* 54.—Afr. trop.
albidus, *Vatke, in Linnaea*, xxxvii. (1871-72) 221.—Abyss.
alpinus, *Vatke, l. c.* 232.—Abyss.
amboinicus, Lour. Fl. Cochinch. ii. 372 = aromaticus.
anamallayensis, *Bedd. in Madr. Journ. Sc.* Ser. II. xxii. (1861) 78.—Ind. or.
arabicus, *Benth. in DC. Prod.* xii. 79.—Arab.
aromaticus, *Benth. in Wall. Pl. As. Rar.* ii. 16.—Ind. or.
atropurpureus, *Benth. l. c.*—Malaya; Pacific.
barbatus, *Benth. l. c.* 15.—Ind. or.; Afr. trop.
Benthamianus, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 354 = inflatus.
bicolor, *Benth. Lab. Gen. et Sp.* 55.—Java.
Blancoi, *Benth. in DC. Prod.* xii. 79.—Ins. Philipp.
Blumei, *Benth. Lab. Gen. et Sp.* 56.—Java.
Bojeri, *Benth. l. c.* 52.—Madag.
canianus, *Vatke, in Linnaea*, xxxvii. (1871-72) 218.—Abyss.
carnosus, *Hassk. in Flora*, xxv. (1842) ii. *Beibl.* 25.—Java.
comosus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 183 = spicatus.
crassifolius, Benth. in Wall. Pl. As. Rar. ii. 15 = aromaticus.
edulis, *Vatke, in Linnaea*, xxxvii. (1871-72) 219.—Abyss.
flaccidus, *Vatke, in Linnaea*, xliii. (1880-82) 90.—Afr. trop.
fruticosus, Benth. in DC. Prod. xii. 78 = Plectranthus fruticosus.
fruticosus, *Wight, ex Benth. in DC. Prod.* xii. 78.—Ind. or.
galeatus, *Benth. Lab. Gen. et Sp.* 56.—Java.
Garckeanus, *Vatke, in Linnaea*, xxxvii. (1871-72) 228.—Abyss.
Gibsonii, *Verlot, in Rev. Hortic.* (1866) 279.—Hab.?
glabratus, Benth. Lab. Gen. et Sp. 58 = Plectranthus coleoides.
glandulosus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 211.—Afr. trop.
grandifolius, *Benth. Lab. Gen. et Sp.* 58.—Ins Timor.
grandifolius, Blanco, Fl. Filip. ed. I. 482 = Blancoi.
Heynei, Benth. Lab. Gen. et Sp. 50 = spicatus.
Huberi, *Regel, in Act. Hort. Petrop.* vii. (1880) 543.—Abyss.
igniarius, *Schweinf. Beitr. Fl. Aethiop.* 121.—Afr. trop.
inflatus, *Benth. Lab. Gen. et Sp.* 58.—Zeylan.
ingratus, *Benth. l. c.* 53.—Java.
laciniatus, *Benth. l. c.* 56.—Java.
lactiflorus, *Vatke, in Linnaea*, xliii. (1880-82) 89.—Afr. trop.
lanuginosus, *Hochst. ex Benth. in DC. Prod.* xii. 79.—Abyss.
latiflolius, *Hochst. ex Benth. l. c.* 74.—Abyss.
leptostachys, *Benth. l. c.* 77.—Zeylan.
Macvari, Benth. Lab. Gen. et Sp. 58 = malabaricus.
macrophyllus, *Benth. l. c.* 55.—Java.
macropus, *Miq. Fl. Ind. Bat.* ii. 956.—Java.
macrostachys, *Benth. Lab. Gen. et Sp.* 57.—Java.
malabaricus, *Benth. in Wall. Pl. As. Rar.* ii. 16.—Ind. or.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 211.—Afr. trop.
micranthus, *Maxim. in Act. Hort. Petrop.* iii. (1875) 108.—Abyss.
mollis, Benth. in DC. Prod. xii. 77 = malabaricus.
multiflorus, *Benth. Lab. Gen. et Sp.* 55.—Ins. Philipp.
ovatus, Benth. l. c. 57 = malabaricus.
palustris, *Vatke, in Linnaea*, xxxvii. (1871-72) 219.—Abyss.
paniculatus, Benth. in Wall. Pl. As. Rar. ii. 16 = Plectranthus coleoides.
parviflorus, *Benth. in DC. Prod.* xii. 72.—Ind. or.
Perrottii, *Benth. Lab. Gen. et Sp.* 56.—Madag.
Petersianus, *Vatke, in Linnaea*, xl. (1876) 180.—Afr. trop.
peberatus, *Miq. Fl. Ind. Bat.* ii. 958.—Malaya.
pumilius, Blanco, Fl. Filip. ed. I. 482 = scuminatus.
remotiflorus, *Miq. Fl. Ind. Bat.* ii. 954.—Java.
rivularis, *Vatke, in Linnaea*, xxxvii. (1871-72) 230.—Abyss.

COLEUS:—
rugosus, Benth. in Wall. Pl. As. Rar. ii. 15 = Plectranthus rotundifolius.
Schimperi, Vatke, in Linnaea, xxxvii. (1871-72) 230 = barbatus.
Schweinfurthii, *Vatke, l. c.* 232, in texte —Afr. trop.
scutellarioides, *Benth. in Wall. Pl. As. Rar.* ii. 16.—Malaya; Austral.
secundiflorus, *Benth. Lab. Gen. et Sp.* 55.—Ins. Timor.
spectabilis, *Miq. Fl. Ind. Bat.* ii. 955.—Java.
spicatus, *Benth. in Wall. Pl. As Rar.* ii. 15.—Ind. or.
suborbicularis, *Zoll. & M oritz, Syst. Verz. Zoll.* 4.—Java.
subspicatus, Walp. Rep. vi. 658 = Plectranthus spicatus.
Sugeroba, Blanco, Fl. Filip. ed. I. 488 = aromaticus.
tenuicaulis, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 211.—Afr. trop.
tenuiflorus, *Vatke, in Linnaea*, xliii. (1880-82) 92.—Afr. trop.
Tryoni x, *Hort. ex Flor. Mag.* N. S. i. (1872) t. 84.
tuberosus, Benth. Lab. Gen. et Sp. 59 = parviflorus.
tuberosus, *A. Rich. Tent. Fl. Abyss.* ii. 185.—Abyss.
umbrosus, *Vatke, in Linnaea*, xliii. (1880-82) 91.—Afr. trop.
urticifolius, Benth. in DC. Prod. xii. 78 = Plectranthus urticifolius.
Verschaffeltii, *Lem. Illustr. Hortic.* viii. (1861) t. 298.—Java.
Wallichii, Benth. in DC. Prod. xii. 77 = malabaricus.
Wightii, Benth. Lab. Gen. et Sp. 58 = Plectranthus coleoides.
Zatarhendi, Benth. l. c. 50 = spicatus?

COLICODENDRON, Mart. in Flora, xxii. (1839) i. Beibl. 25 = **Capparis**, Linn.
anceps, Retz. ex Chapm. Fl. S. U. St. 32 = Capp. jamaicensis.
avicenniaefolium, Seem. Bot. Voy. Herald, 78 = Capp. avicenniifolia.
Breynia, Endl. ex Heynh. Nom. ii. 154 = Capp. Breynia.
lepidotum, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 237.—Venezuela.
longifolium, Mart. in Flora, xxii. (1839) i. Beibl. 26.—Bras.
oblitquifolium, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 238.—N. Granat.
obovatum, Turcz. l. c. 237.—Venezuela.
pulchellum, Seem. Bot. Voy. Herald, 78 = Capp. pulcherrima.
scabridum, Seem. l. c. = Capp. scabrida.
subbiloburn, Seem. l. c. = Capp. subbiloba.
Yco, Mart. in Flora, xxii. (1839) i. Beibl. 25 = Capp. Yco.

COLIGNONIA, Endl. Gen. 311 (1837). *NYCTAGINEAE*, Benth. & Hook. f. iii. 5.
COLLIGNONIA, Endl. Ench. 193 (1841).
biumbellata, *Ball, in Journ. Linn. Soc.* xxii. (1885) 54.—Peruv.
glomerata, *Griseb. in Gentt. Abh.* xix. (1874) 87.—Reg. Argent.
parviflora, *Endl. Gen.* 311; *Choisy, in DC. Prod.* xiii. ii. 428.—Peruv.
scandens, *Benth. Pl. Hartw.* 148.—Peruv.

COLIQUEA, Steud. ex Bibra, in Denkschr. Wien. Acad. v. (1853) 115 = **Chusquea**, Kunth (Gramin.).
Quila, Steud. ex Bibra, l. c. = Chus. Cumingii.

COLLABIUM, Blume, Bijdr. 357 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 506.
nebulosum, *Blume, l. c.* 358.—Java.
simplex, *Reichb. f. in Gard. Chron.* (1881) i. 462.—Borneo.

COLLADEA, Pers. Syn. i. 107 (1805) = seq.

COLLADOA, Cav. Ic. v. 37. t. 460 (1799) = **Tschaemum**, Linn. (Gramin.).
distachia, Cav. l. c. = I. rugosum.
monostachya, Pers. Syn. i. 107 = Anthephora elegans.

COLLADONIA, Spreng. Syst. i. 516 (1825) = **Pallicourea**, Aubl. (Rubiac.).
timifolia, Spreng. l. c. 767 = P. timifolia.

COLLADONIA, DC. Prod. iv. 240 (1830) = **Prangos**, Lindl. (Umbellif.).
alata, Boiss. Fl. Orient. ii. 945.—Syria.
anatolica, Boiss. l. c.—As. Min.
angustifolia, Bertol. Fl. Ital. iii. 408 = P. angustifolia.
anisoptera, Boiss. Fl. Orient. ii. 946.—Syria.
cilicica, Boiss. l. c. 947.—Cilicia.
crenata, Boiss. l. c. 946.—Syria.
heptoptera, Boiss. l. c. 945 = P. colladonioides.
microcarpa, Boiss. l. c. 946 = P. microcarpa.
syriaca, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 86 = Smyrniopsis cachroides.
triquetra, DC. Prod. iv. 240 = P. triquetra.

COLLAEA, DC. Mém. Leg. vi. 244 (1825); et in Ann. Sc. Nat. Sér. I. iv. (1825) 96 = **Galactia**, P. Br. (Legumin.).
angustifolia, Benth. in Ann. Wien. Mus. ii. (1838) 129 = G. angustifolia.
argentina, Griseb. in Goett. Abh. xix. (1874) 125.—Reg. Argent.
bullata, Benth. in Mart. Fl. Bras. xv. i. 149.—Bras.
cinerascens, R. Grah. in Wall. Cat. n. 5575 = Atylosia mollis.
crassifolia, Benth. in Ann. Wien. Mus. ii. (1838) 128.—Bras.
decumbens, Benth. in Linnaea, xxii. (1849) 515.—Bras.
diversifolia, Benth. in Mart. Fl. Bras. xv. i. 147.—Am. trop.
formosa, Griseb. in Goett. Abh. xix. (1874) 126.—Reg. Argent.
gibba, R. Grah. in Wall. Cat. n. 5572 = Dunbaria Heynei, ferruginea.
glaucescens, Benth. in Ann. Wien. Mus. ii. (1838) 128 = G. glaucescens.
grewiaefolia, Benth. l. c. 132.—Bras.
guianensis, Klotzsch, ex Schomb. Faun. & Fl. Gui. 1208.—Guiana.
lancifolia, Benth. in Mart. Fl. Bras. xv. i. 150.—Bras.
latisiliqua, Benth. in Ann. Wien. Mus. ii. (1838) 129 = G. latisiliqua.
longiflora, Benth. l. c. 128.—Ins. Trinit
longifolia, Benth. in Mart. Fl. Bras. xv. i. 151.—Am. trop.
macrophylla, Benth. in Ann. Wien. Mus. ii. (1838) 128.—Bras.
Martii, Benth. l. c. 129 = G. Martii.
mollis, R. Grah. in Wall. Cat. n. 5574 = Atylosia mollis.
Neesii, Benth. in Ann. Wien. Mus. ii. (1838) 129 = G. Neesii.
obtusa, Benth. l. c. 128.—Bras.
pauciuscula, Mart. ex Benth. l. c.—Bras.
pedicellatum, Benth. in Hook. Journ. Bot. ii. (1840) 60.—Bras.
peduncularis, Benth. in Ann. Wien. Mus. ii. (1838) 128.—Bras.
pendula, Benth. l. c. = G. pendula.
rosea, Benth. in Ann. Nat. Hist. iii. (1839) 487 = G. Jussiaeana.
rotundifolia, Benth. in Mart. Fl. Bras. xv. i. 149.—Bras.
rugosa, Benth. in Ann. Wien. Mus. ii. (1838) 128.—Bras.
scarlatina, Mart. ex Benth. l. c. 129.—Bras.
speciosa, DC. in Mém. Lég. vi. 245.—Peruv.
stenophylla, Benth. in Mart. Fl. Bras. xv. i. 146 = G. stenophylla.
trinervia, DC. Mém. Lég. vi. 247 = Atylosia Candollei.
tomentosa, Benth. in Mart. Fl. Bras. xv. i. 149 = G. tomentosa.
velutina, Benth. in Ann. Wien. Mus. ii. (1838) 129 = G. velutina.
venosa, R. Grah. in Wall. Cat. n. 5573 = Dunbaria ferrugines.
virgata, Benth. in Mart. Fl. Bras. xv. i. 146.—Bras.

COLLAEA, Endl. Gen. Suppl. ii. 20 (1842) in syn. = COLLEA, Lindl. = **Polexia**, Lindl. (Orchid.).

COLLAEA, Spreng. Syst. iii. 622 (1826) = **Chrysanthellum**, Rich. (Compos.).
procumbens, Spreng. l. c. = Chrys. procumbens.

COLLANDRA, Lem. in Fl. des Serres, iii. (1847) sub t. 223 = **Columnea**, Linn. (Gesnetrac.).
atrosiriana, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1402 = Columnea aureo-nitens.
aureo-nitens, Hanst. in Linnaea, xxvi. (1853) 209 = Columnea aureo-nitens.
fortolaris, Griseb. Fl. Brit. W. Ind. 468.—Ind. occ.
phoenicea, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1402.—N. Granat.
picta, Lem. Jard. Fleur. ii. (1852) t. 214 = Columnea picta.
pilosa, Lem. in Fl. des Serres, iii. (1847) sub t. 223 = Columnea aureo-nitens.
sanguinea, Griseb. in Mem. Am. Acad. N. S. viii. (1868) 526 = Columnea sanguinea.

COLLANIA, Herb. Amaryll. 108. t. 8 (1837) = **Bomarea**, Mirb. (Amaryllid.).
andimarcana, Herb. l. c. 105 = B. andimarcana.
dulcis, Herb. l. c. 104 = B. glaucescens.
glaucescens, Herb. l. c. = B. glaucescens.
involucrosa, Herb. l. c. 103. t. 9 = B. glaucescens.
puberula, Herb. l. c. 105. t. 11. f. 1 = B. glaucescens.

COLLANIA, Schult. f. Syst. vii. ii. p. lii, 898 (1830) = **Urocoltina**, Reichb. (Amaryllid.).
dubia, Schult. f. l. c.—Ecuador.
fulva, M. Roem. Syn Enast. 55 = U. fulva.
urceolata. Schult. f. Syst. vii. 898 = U. pendula.

COLLEA, Lindl. Bot. Reg. sub t. 760 (1823) = **Palexia**, Lindl. (Orchid.).
adnata, Lindl. l. c. = P. spiranthoides.
calcarata, Lindl. l. c. = P. setacea.

COLLEMA, Anders. ex DC. Prod. vii. 512 (1839) = **Goodania**, Sm.

COLLETIA, Comm. ex Juss. Gen. 380 (1789). *RHAMNEAE*, Benth. & Hook. f. i. 383.
SCYPHARIA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 8.
asiciulata, *Miers*, *in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 214.—Peruv.
armata, Miers, l. c. 211 = spinosa.
articulata, *Phil. in Linnaea*, xxviii. (1856) 679.—Chili.
atrox, *Miers*, *in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 206.—Reg. Argent.
bictonensis, Lindl. in Journ. Hort. Soc. v. (1850) 81 = cruciata.
brevispina, *Phil. in Anal. Univ. Chil.* (1872) 690.—Chili.
cataphracta, *Miers*, *in Ann. & Mag. Mat. Hist.* Ser. III. v. (1860) 210.—Chili.
Chacaye, G. Don, Gen. Syst. ii. 85 = Discaria serratifolia.
crenata, *Clos*, *in C. Gay, Fl. Chil.* ii. 35.—Chili.
crenata, Regel, Ind. Sem. Hort. Petrop. (1856) 37 = Discaria crenata.
cruciata, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 152. t. 48.—Chili.
Cruzillo, *Bert. in 'Merc. Chil.* 1839'.—*Cf. Bull. Férussac*, xx. (1830) 100, nomen.—Chili.
Cunninghamii, Penzl, in Enum. Pl. Hueg. 23 = Discaria australis.
discolor, *Hook. Ic. Pl.* t. 538.—Patagon.
diasperma, *Miq. & Sesse, ex DC. Prod.* ii. 29.—Mexic.
Doniana, *Clos, in C. Gay, Fl. Chil.* ii. 36.—Chili.
dumosa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 212.—Chili.
Ephedra, Bert. ex Steud. Nom. ed. II. i. 397 = Trevoa Trutilla.
Ephedra, *Vent. Choix*, 15. t. 16.—Peruv.
exserta, *Klotzsch, ex Reiss. in Mart. Fl. Bras.* xi. t. 100.—Bras.
ferox, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 154.—Chili.
horrida, Brongn. ex Drap. Encyc. & Fl. des Serres Angl. iii. (Jull. 1835) f. 6 = ferox.
horrida, Willd. Sp. Pl. i. 1113 = spinosa.
Hystrix, *Clos, in C. Gay, Fl. Chil.* ii. 33.—Chili.
infesta, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 366 = Adolphia infesta.
lanuginosa, *Reiss. in Mart. Fl. Bras.* xi. t. 100.—Bras.
intrinsa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 204.—Chili.
invicta, *Miers*, *l. c.* 205.—Chili.
Kunthiana, *Miers*, *l. c.* 206.—Peruv.

COLLETIA :—
longispina, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 178.—Chili.
longissima, Steud. Nom. ed. II. i. 397 = praec.
maytenoides, *Griseb. in Gottt. Abh.* vi. (1854) 117.—Reg. Argent.
montana, *Hist. in Linnaea*, xxviii. (1856–65) 37.—Chili.
multiflora, Moç. & Sesse, ex DC. Prod. ii. 29 = Colubrina nana, *Clos, in C. Gay, Fl. Chil.* ii. 37.—Chili.
obcordata, *Vent. Jard. Cels*, t. 92.—Peruv.
polyacantha, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 513 = obcordata, spinosa.
pubescens, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 366 = Discaria australis.
pungens, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 209.—Chili.
serratifolia, Vent. Choix, 15. t. 15 = Discaria serratifolia.
spartioides, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 52.—Ins. J. Fernand.
spicata, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 513.—Am. austr.
spinosa, *Lam. Tabl. Encyc.* ii. 91. t. 129.—Am. austr.
spinosissima, J. F. Gmel. Syst. 408 = praec.
stipellacea, *Phil. in Linnaea*, xxxiii. (1864–65) 36.—Chili.
tenuicola, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 209.—Reg. Argent.!
tetragona, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 366.—Peruv.
tetrandra, *Clos, in C. Gay, Fl. Chil.* ii. 31.—Chili.
tomentosa, *Phil. in Linnaea*, xxviii. (1856) 618.—Chili.
Trailliana, Bert. ex Colla, in Mem. Acc. Torin. xxxvii. (1834) 53 = Trevoa quinquenervia.
Trella, Bert. ex Colla, l. c. = Trevoa trinervia.
ulicina, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 155.—Chili.
valdiviana, *Phil. in Linnaea*, xxxiii. (1864–65) 35.—Chili.
Valensuela, Bert. ex Steud Nom. ed. II. i. 397 = Colubrina.
velutina, *Spreng. Syst.* i. 771.—Bras.
veprescula, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 213.—Chili.
Weddelliana, *Miers, l. c.* 207.—Bolivia.

COLLETIA, Scop. Introd. 207 (1777) = **Coltis**, Linn. (Urticac.).

COLLIGNONIA, Endl. Ench. 198 (1841) = **Colignonia**, Endl. (Nyctag.).

COLLIGUAJA, Molina, Sagg. Chil. 158, 354 (1782). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 338.
brasiliensis, *Klotzsch, ex Baill. Etud. Gén. Euph.* 535.—Bras.
Bridgesii, *Muell. Arg. in Linnaea*, xxxii. (1863) 125.—Chili.
Dombeyana, C. Gay, Fl. Chil. 343 = salicifolia.
Dombeyana, *A. Juss. in Ann. Sc. Nat.* Sér. I. xxv. (1832) 23.—Chili.
integerrima, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 140.—Chili.
obtusa, *Regel, Ind. Sem. Hort. Petrop.* (1855) 18.—Chili.
odorifera, *Molina, Sagg. Chil.* 158, 354.—Chili.
salicifolia, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 141.—Chili.
triquetra, *Gill. & Hook. l. c.* 142.—Chili.

COLLINARIA, Ehrh. Beitr. iv. (1780) 147 = **Koeleria**, Pers. (Gramin.).

COLLINIA, Liebm. in Oefers. Danske Vid. Selsk. 1845 (1846) 8 = **Chamaedorea**, Willd. (Palm.).
elatior, Liebm. l. c. = Cham. Liebmanni.
elegans, Liebm. l. c. = Cham. elegans.
fibrosa, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 5 = Synechanthus fibrosus.
humilis, Liebm. in Oefers. Danske Vid. Selsk. 1845 (1846) 8 = Cham. humilis.

COLLINIA, & COLLINIANA, Rafin. in Am. Monthly Mag. (1819) 194 = seq.

COLLINSIA, Nutt. in Journ. Acad. Philad. i. (1817) 190. t. 9. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 941.
barbata, Bosse, in Verh. Berl. Gartenh. Ver. N. Reihe, i. (1858) 299 = tinctoria.
bartsiaefolia, *Benth. in DC. Prod.* x. 318.—Calif.
bicolor, *Benth. in Trans. Hort. Soc.* N. S. i. (1831–35) 480.—Calif.
Childii, *Parry, ex A. Gray, Syn. Fl. N. Am.* ii. 1. 287.—Calif.
corymbosa, *Herder, Ind. Sem. Hort. Petrop.* (1857) 32; *Regel, Gartenfl.* (1858) 33. t. 568.—Calif.
divaricata, *Kellogg, in Proc. Calif. Acad.* iii. (1863) 26.—Calif.
grandiflora, *Dougl. ex Lindl. Bot. Reg.* sub t. 1166 et t. 1107.—Am. bor. occ.
Greenii, *A. Gray, in Proc. Am. Acad.* x. (1874) 75.—Calif.
heterophylla, R. Grah. in Hook. Bot. Mag. t. 3695 = bicolor.
hirsuta, Kellogg, in Proc. Calif. Acad. ii. (1863) 110 t. 33 = bartsiaefolia.
linearis, *A. Gray, in Proc. Am. Acad.* xv. (1880) 50.—Am. bor. occ.
minima, Nutt. in Journ. Acad. Philad. vii. (1834) 47 = parviflora.
multicolor, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 89. t. 55.—Calif.
Parryi, *A. Gray, Syn. Fl. N. Am.* ii. t. 257.—Calif.
parviflora, *Lindl. Bot. Reg.* t. 1082.—Am. bor.
pauciflora, Hook. Fl. Bor. Am. ii. 94, sphalm. = parviflora.
purpurea, *Rafin. Atl. Journ.* 149.—Am. bor.
Rattani, *A. Gray, in Proc. Am. Acad.* xv. (1880) 50.—Am. bor. occ.
septemnervia, Kellogg, in Proc. Calif. Acad. ii. (1863) 224 = tinctoria.
solitaria, Kellogg, l. c. 10 = sparsiflora.
sparsiflora, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1836) 33.—Calif.
tenella, Benth. in DC. Prod. x. 598 = Tonella collinsioides.
tinctoria, *Hartw. ex Benth. Pl. Hartw.* 328.—Calif.
Torreyi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 378.—Calif.
trycolor, Rafin. Cat. 18 = verna.
verna, *Nutt. in Journ. Acad. Philad.* i. (1817) 190. t. 9.—Am. bor.
violacea, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 170 —Am. bor.

COLLINSONIA, Linn. Gen. ed. I. 884 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1181.
DIALLOSTIRA, Rafin. Neogenyt. 3 (1825).
HYPOGON, Rafin. Fl. Ludov. 149 (1817).
PLEURADENIA, Rafin. Neogenyt. 2 (1825).
acinata, *Sims, Bot. Mag.* t. 1213.—Am. bor.
canadensis, *Linn. Sp. Pl.* 26.—Am. bor.
cuneata, Wender. in Schr. Nat. Ges. Marb. † ex Benth. in DC. Prod. xii. 558 = canadensis.
decussata, Moench, Meth. 379 = canadensis.
dentis, Pursh, Fl. Am. Sept. i. 21 = scabriuscula.
grasces, Walt. Fl. Carol. 65 = scabriuscula.
punctata, Ell. Sketch, i. 36 = canadensis.
scabra, Pers. Syn. i. 29 = scabriuscula.
scabriuscula, *Ait. Hort. Kew.* ed. I. i. 47.—Am. bor.
serotina, Walt. Fl. Carol. 65 = canadensis.
tuberosa, Michx. Fl. Bor. Am. i. 17 = scabriuscula.
verticillata, Salisb. Prod. 78 = canadensis.
verticillaris, Rafin. Fl. Ludov. 41 = seq. ?
verticillata, *Baldw. ex Ell. Sketch*, i. 36.—Am. bor.

COLLOCOCCUS, P. Br. Hist. Jamaic. 167 (1756) (Gen. dub.).

COLLOMIA, Nutt. Gen. Am. i. 126 (1818). *POLEMONIACEAE*, Benth. & Hook. f. ii. 822.
COURTOSIA, Reichb. Io. Exot. iii. 4. t. 208 (1837).
aggregata, T. C. Porter, ex A. Gray, Syn. Fl. N. Am. ii. 1. 294 = Gilia aggregata.
Cavanillesiana, *G. Don, Gen. Syst.* iv. 347.—Mexic.; Texas.
Cavanillesii, Hook. & Arn. Bot. Beech. Voy. 84 = coccinea.
coccinea, *Lehm. ex Benth. in Bot. Reg.* t. 1032.—Chili.
orthocladon, *Griseb. in Gottt. Abh.* vi. (1853–56) 126.—Chili.
erythrocodon, *Griseb. l. c.*—Chili.

COLLOMIA:—
gilloides, *Benth. in Bot. Reg.* sub t. 1622.—Calif.
glutinosa, Benth. l. c. = gilloides.
gracilis, *Dougl. ex Benth. l. c.*—Am. bor. occ.
grandiflora, *Dougl. in Lindl. Bot. Reg.* t. 1174.—Am. bor. occ.
heterophylla, *Hook. Bot. Mag.* t. 2895.—Am. bor. occ.
laterifia, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. iii. t. 206 = coccinea.
leptalea, *A. Gray, in Proc. Am. Acad.* viii. (1870) 261. —Calif.
linearis, *Nutt. Gen. Am.* i. 136.—Am. bor. occ.
linoides, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 159 = Gilia minutiflora.
longiflora, *A. Gray, in Proc. Am. Acad.* viii. (1870) 261.—Am. bor. occ.
micrantha, Kellogg, in Proc. Calif. Acad. iii. (1868) 18. f. B = gracilis.
myotica, *Griseb. in Gott. Abh.* vi. (1853-55) 129.—Chili.
Navarretia, G. Don, Gen. Syst. iv. 247 = Gilia Navarretia
nudicaulis, *Hook. & Arn. Bot. Beech. Voy.* 368.—Am. bor. occ.
stenothyrsa, *Kunze, in Linnæa*, xxiv. (1851) 228.—Chili.
tenella, *A. Gray, in Proc. Am. Acad.* viii. (1870) 259. —Am. bor. occ.
Thurberi, *A. Gray, l. c.* 261.—N. Mexic.
tinctoria, Kellogg, in Proc. Calif. Acad. iii. (1868) 17. f. 2 = linearis.

COLLOMIA, Sieber, ex Steud. Nom. ed. II. i. 398 (1840) = **Felicia**, Cass. (Compos.).
setosa, Sieber, ex Steud. l. c. = F. reflexa.

COLLOPHORA, Mart. in Buchn. Rep. Pharm. (1828).—Cf. Bull. Férussac (Jan. 1831) 64 = **Couma**, Aubl. (Apocyn.).
utilis, Mart. l. c. = Couma utilis.

COLLYRIS, Vahl, in Skrift. Nat. Selsk. Kiobenh. vi. (1810) 110 = **Dischidia**, R. Br. (Asclepiad.).
major, Vahl, l. c. = D. Rafflesiana, D. Collyris.
minor, Vahl, l. c. 111 = D. nummularia.

COLMEIROA, F. Muell. Fragm. vii. 149 (1871). *SAXIFRAGEAE.*
carpodetoides, *F. Muell. l. c.*—Ins. D. Howe.

COLMEIROA, Reut. in Mém. Soc. Phys. Genèv. x. (1843) 240. t. 6 = **Scourinoga**, Comm. (Euphorb.).
buxifolia, Reut. l. c. = S. buxifolia.

COLOBACHNE, Beauv. Agrost. 22. t. 6. f. 9 (1812) = **Alopecurus**, Linn. (Gramin.).
brachystachys, Nym. in Bot. Notiser (1851) 69 = A. brachystachys.
Gerardi, Link, Hort. Berol. i. 74 = Phleum Gerardi
jonitica, Nym. in Bot. Notiser (1851) 69 = A. glacialis.
vaginata, Beauv. Agrost. 22 = A. vaginatus.

COLOBANDRA, Bartl. in Lehm. Pl. Preiss. i. 357 (1845) = **Hemigenia**, R. Br. (Labiat.).
canescens, Bartl. l. c. 358 = H. canescens.
lanata, Bartl. l. c. 358 = H. canescens ?
mollis, Bartl. l. c. 358 = H. canescens.
platyphylla, Bartl. l. c. = H. platyphylla.
robusta, Bartl. l. c. 357 = H. polystachya.
subcillosa, Bartl. l. c. 359 = H. platyphylla ?

COLOBANTHUS, Bartl. in Presl, Reliq. Haenk. ii. 18. t. 49 (1830). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 151.
acicularis, *Hook. f. Handb. New Zeal. Fl.* 25.—N. Zel.
affinis, Hook. f. in Hook. Journ. Bot. ii. (1840) 410 = Billardieri.
arcticoides, *Gill. ex Hook. Bot. Misc.* iii. (1833) 336.—Chili.
Benthamianus, Fenzl, in Ann. Wien. Mus. i. (1836) 49 = subulatus.
Billardieri, *Fenzl, l. c.*—Austral.
charlottiodes, *Hook. f. Fl. Antarct.* 349.—Patagon.
crassifolius, *Hook. f. l. c.* 348.—Chili; Patagon.
diffusus, *Hook. f. l. c.* 349.—Ins. Amsterdam.
hartwegianus, *Hook. f. l. c.*—Ins. Kerguel.
Lechleri, *Phil. in Linnæa*, xxviii. (1856) 659.—Chili.

COLOBANTHUS:—
lycopodioides, *Griseb. in Gott. Abh.* vi. (1853-55) 116.—Reg. Magellan.
muscoides, *Hook. f. Fl. Antarct.* 14.—Ins. Antarct.
polycnemoides, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 334.—Reg. Argent.
pulvinatus, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 101 = subulatus.
quitensis, *Bartl. in Presl, Reliq. Haenk.* ii. 18. t. 49. f. 2.—Ecuador.
saginoides, *Bartl. l. c.*—Chili.
subulatus, *Hook. f. Fl. Antarct.* 13.—Ins. Antarct.

COLOBANTHUS, Trin. in Mém. Acad. Pétersb. Sér. VI i. (1831) 66 = **Avellinia**, Parl. (Gramin.).

COLOBATUS, Walp. in Linnæa, xiii. (1839) 478 = COLOBOTUS, E. Mey. = **Buchenroedera**, Eckl. & Zeyh. (Legumin.).

COLOBIUM, Roth, in Roem. Archiv, i. (1796) i. 86 = **Leontodon**, Linn. (Compos.).
hirtum, Roth, l. c. = L. hirtum.
hispidum, Roth, l. c. 87 = L. Rothii.

COLOBOGYNIUM, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 84 = **Schismatoglottis**, Zoll. (Aroid.).
lectuarium, Schott, l. c. = S. variegata.

COLOBOTUS, E. Mey. Comm. Pl. Afr. Austr. 156 (1835) = **Buchenroedera**, Eckl. & Zeyh. (Legumin.).
ochreatus, E. Mey. l. c.—Afr. austr.

COLOCASIA, Link, Diss. Bot. Suerin (1795) 77; et Handb. i. 267 = **Richardia**, Kunth (Aroid.).
aethiopica, Link, ll. cc. = R. aethiopica.

COLOCASIA, Neck. Elem. iii. 289 (1790) (Gen. dub. Aroidearum).

COLOCASIA, Schott, Meletem. i. 18 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 974.
LEUCOCASIA, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 34.
acris, Schott, Meletem. i. 18 = antiquorum.
affinis, *Schott, in Bonplandia*, vii. (1859) 28.—Reg. Himal.
alba, Hort. ex Engl. in DC. Monog. Phan. ii. 501 = Alocasia alba.
argyrenerura, Hort.—Cf Gard. Chron. (1875) i. 589 = Caladium Schomburgkii.
antiquorum, *Schott, Meletem.* i. 18.—As. trop.
Boryi, Kunth, Enum. Pl. iii. 41 = Alocasia indica.
cochleata, Miq. Epim. Sem. Hort. Amstelod. (1858) = Alocasia cucullata.
cucullata, Schott, Meletem. i. 18 = Alocasia cucullata.
esculenta, Schott, l. c. = antiquorum.
euchlora, C. Koch & Lladen, in App. Sem. Hort. Bot. Berol. (1854) 4 = antiquorum.
fallax, *Schott, in Bonplandia*, vii. (1859) 28.—Ind. or *Fontanesii*, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 409 = antiquorum.
formicaria, Kunth, Enum. Pl. iii. 41 = Alocasia fornicata.
gracilis, *Engl. Bot. Jahrb.* i. (1881) 185.—Sumatra.
heterophylla, *Kunth, Enum. Pl.* iii. 40.—Ins. Philipp.
humilis, *Hassk. Cat. Hort. Bogor. Alt.* 86.—Java.
indica, *Hassk. in Hoov. & De Vriese, Tijdschr.* ix. (1842) 160.—Java.
indica, Kunth, Enum. Pl. iii. 39 = Alocasia indica.
macrorrhiza, Schott, Meletem. i. 18 = Alocasia macrorrhiza.
Marchalii, *Engl. in DC. Monog. Phan.* ii. 494.—Hab.?
montana, Kunth, Enum. Pl. iii. 40 = Alocasia montana.
mucronata, Kunth, l. c. = Alocasia macrorrhiza.
neocaledonica, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1853) 13 = Alocasia navicularia.
neo-guineana, *André, in Illustr. Hortic.* xxvii. (1880) 68.—N. Guinea.
nitida, Walp. Ann. i. 1041, sphalm. = Coprosma nitida.
nymphaefolia, Kunth, Enum. Pl. iii. 37 = antiquorum.
obtusiloba, *Kunth, l. c.* 41.—Hab.?
odora, Brongn. in Nouv. Ann. Mus. Par. iii. (1834) 145. t. 7 = Alocasia odora.
peregrina, *Engl. in Fl. Tellur.* iii. 65.—Hab.?
pruinifera, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1854) 4 = indica.
puberula, Hassk. Cat. Hort. Bogor. Alt. 302 = Alocasia puberula.

COLOCASIA:—
pumila, Kunth, Enum. Pl. iii. 40 = Gonstanthus sarmentosus.
ramosa, Schott, ex Heynh. Nom. ii. 154 = Arum ramosum.
rapiformis, *Kunth, Enum. Pl.* iii. 40.—Burma.
rupestris, Kunth, l. c. 41 = Alocasia cucullata.
virosa, *Kunth, l. c.*—Ind. or.
vivipara, *Thw. Enum. Pl. Zeyl.* 326.—Zeylan.
vulgaris, Rafin. Fl. Tellur. iii. 66 = antiquorum.

COLOCOCCA, Rafin. Sylva Tellur. 40 (1838) = **Cordia**, Linn. (Boragin.).
macrophyla, Rafin. l. c. = Cordia macrophylla.

COLOCYNTHIS, [Tourn.] Linn. Syst. ed. I (1735); Schrad. Ind. Sem. Hort. Gotting. (1832) 2; Linnæa, x. (1836) Lit. 70 = **Citrullus**, Neck. (Cucurbit.).
amarissima, Schrad. ll. cc. = Cit. vulgaris.
officinalis, Schrad. in Linnæa, xii. (1838) 421 = Cit. Colocynthis.
officinarum, Cogn. in DC. Monog. Phan. iii. 511, sphalm. = præc.
vulgaris, Schrad. Ind. Sem. Hort. Gotting. (1832) 2 = Cit. Colocynthis.

COLOGANIA, Kunth, Mim. 205. t. 57 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 539.
affinis, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) ii. 186.—Mexic.
angustifolia, *Kunth, Mim.* 209. t. 58.—Am. bor. occ.
australis, *Griseb. in Gott. Abh.* xix. (1874) 124.—Reg. Argent.
bliloba, *Nichols. Dict. Gard.* i. 363.—Mexic.
Broussonetii, *DC. Prod.* ii. 257.—Mexic.
heterophylla, Gill. ex Hook. Bot. Misc. iii. (1833) 181 = Calictis marginalis.
humifusa, *Hemsl. Diagn. Pl Nov.* 47.—Mexic.
intermedia, *H. B. & K. Nov. Gen. et Sp.* vi. 414.—Mexic.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 74.—Arizona.
longifolia, *A. Gray, Pl. Wright.* ii. 35.—Am. bor. occ.
Martii, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 345.—Mexic.
obovata, *Schlecht. in Linnæa*, xii. (1838) 287.—Mexic.
ovalifolia, H. B. & K. Nov. Gen. et Sp. vi. 412 = pulchella.
procumbens, *Kunth, Mim.* 205. t. 57.—N. Granat.
pulchella, *H. B. & K. Nov. Gen. et Sp.* vi. 413.—Am. bor. occ.

COLOGYNE, Griff. Notul. iii. 280 (1851) err. typ. = **Coelogyne**, Lindl. (Orchid.).

COLOMANDRA, Neck. Elem. ii. 142 (1790) = **Alouea**, Aubl. (Laurin.).

COLONA, Cav. Ic. iv. 47. t. 370 (1797) = **Columbia**, Pers. (Tiliac.).
serratifolia, Cav. l. c. = Columbia americana.

COLONNA, Jaume St. Hil. Expos. ii. 95 (1806) = præc.

COLONNEA, Endl. Gen. 421, in syn. (1838) = CALONNEA, Buchoz = **Gaillardia**, Foug. (Compos.).

COLOPHONIA, Comm. ex Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 353 = **Canarium**, Linn. (Burserac.).
mauritiana, DC. Prod. ii. 79 = Canarium mauritianum.
tetramera, Boiv. ex Engl. in DC. Monog. Phan. iv. 27 = Commiphora tetramera.

COLPIAS, E. Mey. ex Benth. in Hook. Comp. Bot. Mag. ii. (1836) 58. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 981.
affinis, *E. Mey. l. c.*—Afr. austr.

COLPODIUM, Trin. Fund. Agrost. 119 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1196.
altaicum, *Trin. in Ledeb. Fl. Alt.* i. 100.—Sibir. alt.
aquaticum, Trin. Fund. Agrost. 138 = Catabrosa aquatica.
arundinaceum, *Hook. Fl. Bor. Am.* ii. 238.—Am. bor.
bulbosum, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 294 = Catabrosa humilis.
Calderi, Boiss. Diagn. Sér. II. iv. 138 = Catabrosa humilis.

COLPODIUM:—
compressum, Trin. in Spreng. Neue Entdeck. ii. 58 = Sporobolus compressus.
fibrosum, Trautv. in Act. Hort. Petrop. ii. (1873) 486 = Catabrosa fibrosa.
filiifolium, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 69 = Catabrosa humilis.
fulvum, Griseb. in Ledeb. Fl. Ross. iv. 385 = Poa fulva.
humile, Griseb. l. c. 384 = Catabrosa humilis.
junceum, Trin. in Spreng. Neue Entdeck. ii. 57 = Sporobolus junceus.
latifolium, *R. Br. in Parry, 1st Voy. Suppl.* p. clxxvi. —Reg. arct.
Malmgrenii, *Anderss. in Öfvers. Vet. Akad. Stockh.* xxiii. (1866) 121.—Ins. Spitzberg.
minutum, *Steud. Syn. Pl. Gram.* 177.—Hispan.
monandrum, Trin. Fund. Agrost. 119 = Phippsia algida.
parviflorum, Boiss. & Buhse, Diagn. Ser. II. iv. 133 = Catabrosa parviflora.
pauciflorum, *Hook. Fl. Bor. Am.* ii. 238.—Am. bor.
pendulinum, Griseb. in Ledeb. Fl. Ross. iv. 386 = Poa Lasiadii.
pusillum, *Nees, Fl. Afr. Austr.* 149.—Afr. austr.
Stevani, Hoben. in Bull. Soc. Nat. Mosc. vi. (1833) 218 = Catabrosa humilis.
Stevani, *Trin. Fund. Agrost.* 119. t. 7.—Reg. Caucas.
Tilesii, *Griseb. in Ledeb. Fl. Ross.* iv. 385.—Sibir. or.
variegatum, *Boiss. ex Griseb. l. c.* 386, in textu.—Aa. Min.
violaceum, *Griseb. l. c.*—Persia.

COLPOON, Berg. Desc. Pl. Cap. 88. t. 1 (1767).
SANTALACEÆ, Benth. & Hook. f. iii. 225.
FUSANUS, Murr. Syst. ed. XIII. 765 (1774).
HAMILTONIA, Hartt. Gen. Pl. Cap. ed. I. 298 (1888).
RHOICARPOS, A. DC. Prod. xiv. 634 (1857).
compressum, *Berg. Desc. Pl. Cap.* 88.—Afr. austr.

COLPOTHRINAX, Griseb. & H. Wendl. in Bot. Zeit. xxxvii. (1879) 147. *PALMAE*, Benth. & Hook. f. iii. 937.
Wrightii, *H. Wendl. in Kerch. Palm.* 241.—Cuba.

COLQUHOUNIA, Wall. in Trans. Linn. Soc. xiii. (1822) 606. *LABIATAE*, Benth. & Hook. f. ii. 1208.
coccinea, *Wall. l. c.* 606.—Reg. Himal.
elegans, *Wall. Pl. As. Rar.* i. 85.—Burma.
mollis, *Schlecht. in Linnaea*, xxiv. (1851) 681.—Reg. Himal. ?
tenuiflora, *Hook. f. Fl. Brit. Ind.* iv. 674.—Burma.
tomentosa, Houtt. in Nov. Hortic. (1778) 131 = coccinea.
vestita, *Wall. Tent. Fl. Nep.* i. 14.—Reg. Himal.

COLSMANNIA, Lehm. in Ges. Nat. Fr. Berl. Mag. viii. (1814) 92. t. 4 = Oncosma, Linn. (Boraginˑ.).
flava, Lehm. l. c. = O. flavum.

COLUBRINA, Rich. ex Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 366. t. 15. f. B. *RHAMNEAE*, Benth. & Hook. f. i. 379.
DIPLISCA, Rafin. Sylva Tellur. 31 (1838).
MARCORELLA, Neck. Elem. iii. 122 (1790).
TUBANTHERA, Comm. ex DC. Prod. ii. 30 (1825).
acuminata, *Griseb. Cat. Pl. Cub.* 84.—Cuba.
Alamani, *G. Don, Gen. Syst.* ii. 37.—Mexic.
americana, Nutt. Sylva, ii. 58 = ferruginea.
asiatica, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 369.—As. et Afr. trop.
buxifolia, Schlecht. in Linnaea, xv. (1841) 469 = Canothus buxifolius.
celtidifolia, *Schlecht. l. c.* 471.—Mexic.
cordifolia, *Reiss. in Mart. Fl. Bras.* xi. 1. 98.—Bras.
cubensis, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 369 = Canothus cubensis.
Ehrenbergii, *Schlecht. in Linnaea*, xv. (1841) 469.—Mexic.
excelsa, Fenzl, in Enum. Pl. Huegˑ. 20 = Alphitonia excelsa.
fermenta, *Rich. ex Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 369.—Guiana.
ferruginosa, *Brongn. l. c.*—Am. bor.
glomerata, *Hemsl. Biol. Centr.-Am. Bot.* i. 200.—Mexic.
granulosa, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 369.—Mexic.
Greggii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 336.—Am. bor. occ.
guineensis, G. Don, Gen. Syst. ii. 37 = Chailletia toxicaria ?

COLUBRINA:—
infesta, Schlecht. in Linnaea, xv. (1841) 468 = Adolphia infesta.
jamaicensis, Miq. Fl. Ind. Bat. i. 1. 646 = asiatica.
Leschenaultii, G. Don, Gen. Syst. ii. 37 = asiatica ?
macrocarpa, G. Don, l. c. 36 = Ceanothus macrocarpus.
Mocinianna, G. Don, l. c. = Ceanothus Mocinianus.
mystacina, G. Don, l. c. 37 = Helinus scandens.
nepaulensis, G. Don, l. c. 36 = Rhamnus nepaulensis.
oppositifolia, *Brongn. ex H. Mann, in Proc. Am. Acad.* vii. (1867) 161.—Ins. Sandvic.
pubescens, G. Don, Gen. Syst. ii. 36 = Ceanothus pubescens.
pubescens, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 301.—Burma.
pubiflora, G. Don, Gen. Syst. ii. 37 = Zizyphus pubiflorus.
reclinata, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 369 = Ceanothus reclinatus.
rufa, *Reiss. in Mart. Fl. Bras.* xi. 1. 98.—Bras.
texensis, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 169.—Texas.
travancorica, *Bedd. Ic. Pl. Ind. Or.* i. 48.—Ind. or.
trifora, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 369.—Mexic.
triquetra, G. Don, Gen. Syst. ii. 37 = Rhamnus triquetra.
vitiensis, Seem. Syst. List Vit. Pl. 4 = Rhamnus vitiensis.

COLUMBARIA, J. & C. Presl, Fl. Čech. 32 (1819) = Scandixˑ, Tourn. (Dipsac.).
canescens, J. & C. Presl, l. c. = S. suaveolens.
crenata, Presl, ex Steud. Nom. ed. II. i. 398 = S. crenata.
glabrescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = S. lucida.
graminifolia, Fourr. l. c. = S. graminifolia.
minor, S. F. Gray, Nat. Arr. Brit. Pl. ii. 476 = S. Columbaria.
norica, J. & C. Presl, Fl. Čech. 32 = S. lucida.
ochroleuca, J. & C. Presl, l. c. = S. ochroleuca.
patens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = S. Columbaria.
pratensis, Fourr. l. c. = Knautia arvensis.
rubella, Opiz, ex Steud. Nom. ed. II. i. 398 = S. Columbaria.
suaveolens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = S. suaveolens.
tenuifolia, Fourr. l. c. = S. graminifolia.
vulgaris, J. & C. Presl, Fl. Čech. 32 = S. Colombaria.

COLUMBEA, Salisb. in Trans. Linn. Soc. viii. (1807) 317 = Tradescantia, Juss. (Conifer.).
angustifolia, Bertol. in Bologn. Opusc. Sc. iii. (1819) 411 = excelsa.
angustifolia, Raddi, ex Steud. Nom. ed. II. i. 398 = A. brasiliana.
Bidwilli, Carr. Conif. ed. II. 601 = A. Bidwillii.
brasiliensis, Carr. l. c. 596 = A. brasiliana.
excelsa, Spreng. Syst. iv. Cur. Post. 215 = A. excelsa.
imbricata, Carr. Conif. ed. II. 598 = A. imbricata.
quadrifaria, Salisb. in Trans. Linn. Soc. viii. (1807) 317 = A. imbricata.

COLUMBIA, Pers. Syn. ii. 66 (1807). *TILIACEAE*, Benth. & Hook. f. i. 232.
COLONA, Cav. Ic. iv. 47. t. 370 (1797).
americana, *Pers. Syn. ii.* 66.—Ins. Philipp.
Anilao, *Blanco, Fl. Filip.* ed. I. 654.—Ins. Philipp.
Blancoi, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 308. —Ins. Philipp.
celebica, *Bisma, Bijdr.* 117.—Ins. Celebes.
floribunda, Naves, in Blanco, Fl. Philipp. ed. III. t. 513 = Blancoi.
floribunda, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 93.—Burma.
gramosepala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 575.—Hab. ?
inaequilatera, Turcz. l. c. xxxi. (1858) i. 233 = serratifolia.
javanica, *Blume, Bijdr.* 117.—Java.
marguensis, *Planch. ex Mast. in Hook. f. Fl. Brit. Ind.* i. 394.—Burma.
serratifolia, Blanco, Fl. Filip. ed. II. 427 = Grewia columnaris.
serratifolia, *DC. Prod.* i. 512.—Ins. Philipp.

COLUMELLA, Comm. ex DC. Prod. i. 443, in syn. (1824) = Pavonia, Cav. (Malvac.).
Malvinda, Comm. ex DC. l. c. = P. Columella.

COLUMELLA, Lour. Fl. Cochinch. i. 85 (1790) = Vitis, Linn. (Ampelid.).
pedata, Lour. l. c. 86 = V. cochinchinensis.

COLUMELLA, Vahl, Enum. i. 300 (1805) = Columellia, Ruiz & Pav.

COLUMELLA, Vell. Fl. Flum. 155 (1825); iv. t. 17 = Pisonia, Linn. (Nyctagin.).
rustica, Vell. l. c. = P. subcordata.

COLUMELLEA, Jacq. Hort. Schoenbr. iii. 28. t. 301 (1798) = Nestlera, Spreng. (Compos.).
hernna, Jacq. l. c. 29 = N. bicaulis.

COLUMELLIA, Ruiz & Pav. Prod. 3. t. 1 (1794). *COLUMELLIACEAE*, Benth. & Hook. f. iii. 989.
ULIXIA, Juss. in Dict. Sc. Nat. x. 103 (1818).
arborescens, Pers. Syn. i. 13 = oblonga.
frutescens, Pers. l. c. = oblonga.
oblonga, *Ruiz & Pav. Fl. Per.* i. 28.—Peruv.
obovata, *Ruiz & Pav. l. c.*—Peruv.
serices, *H. B. & K. Nov. Gen. et Sp.* ii. 288.—Ecuador.

COLUMNEA, Plum. ex Linn. Gen. ed. I. 278 (1737). *GESNERACEAE*, Benth. & Hook. f. ii. 1009.
APONOA, Rafin Sylva Tellur. 84 (1838).
COLLANDRA, Lem. in Fl. des Serres, iii. (1847) sub t. 223.
DALBERGARIA, Tussac, Fl. Antill. i. 141. t. 19 (1808).
EUSYNETRA, Rafin. Fl. Tellur. ii. 57 (1836).
GLYCANTHES, Rafin. Sylva Tellur. 88 (1838).
ORTHOLOMA, Hanst. in Linnaea, xxvi. (1853) 909.
PENTADESIA, Hanst. l. c. 211 (1853).
STENANTHUS, Oerst. ex Hanst. l. c. 209 (1858).
STYGNANTHE, Hanst. l. c. (1853).
acuminata, *Benth. Pl. Hartw.* 281.—N. Granat.
anisophylla, *DC. Prod.* vii. 543.—Am. austr.
argentea, *Griseb. Fl. Brit. W. Ind.* 465.—Ind. occ.
aurantiaca, *Decne. ex Planch. in Fl. des Serres*, vi. (1850–51) 45.—Reg. Andin.
aures, *Warcz. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 258.—N. Granat. ?
aureoalbica, *Hook. Bot. Mag.* t. 4294.—N. Granat.
balsamea, Roxb. Hort. Beng. 47; Fl. Ind. iii. 97 = Limnophila grattioloides.
bilabiata, *Seem. Bot. Voy. Herald*, 186.—Panama.
Billbergiana, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 185.—Panama.
campanulata, *Benth. Pl. Hartw.* 282.—N. Granat.
capitata, Walp. Rep. vi. 405, sphalm. = Besleria capitata.
coccinea, Hort. ex Hanst. in Linnaea, xxvii. (1855–56) 405 = rotundifolia.
cochinchinensis, Poir. Encyc Suppl. iii. 815 (= *Dicrus cochinchinensis*).—Cochinch.
consanguinea, *Hanst. in Linnaea*, xxxiv. (1865–66) 368.—Costa Rica.
crassifolia, *Hook. Bot. Mag.* t. 4330.—Mexic.
divaricata, Walp. Rep. vi. 405, sphalm. = Besleria divaricata.
diffusa, Roxb. ex Hook. f. Fl. Brit. Ind. iv. 277 = Torenia vagans.
erecta, Lam. Encyc. ii. 66 = Achimenes coccinea.
erianthæ, *Hanst. in Linnaea*, xxxiv. (1865–66) 891.—Ecuador.
erythrocalyx, Oerst. in Videnak. Selsk. Skr. V. v. (1861) 188 = querceti.
erythrophæa, *Decne. ex Houtt. in Nov. Hortic.* (1867) 179.—Mexic.
erythrophylla, *Hanst. in Linnaea*, xxxiv. (1865–66) 389.—N. Granat.
flaccida, *Seem. Bot. Voy. Herald*, 186.—Panama.
flava, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) ii. 39.—Mexic.
glabra, *Oerst. in Videnak. Selsk. Skr.* V. v. (1861) 186. —Costa Rica.
grandiflora, Hort. ex Hanst. in Mart. Fl. Bras. viii. 414 = Nematanthus longipes.
guttata, *Prepp. & Endl. Nov. Gen. et Sp.* iii. 1.—Peruv.
hondeloides, Spreng. ex Hayne, Nom. ii. 154 = Streptocarpus Rexii.

COLUMNEA:—
heterophylla, *Hanst. in Linnæa*, xxxiv. (1865–66) 390.—Costa Rica.
heterophylla, Roxb. Hort. Beng. 47; Fl. Ind. iii. 97 = Linnophila heterophylla.
hirsuta, Klotzsch, ex Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 135 = hirta.
hirsuta, *Sw. Prod. Veg. Ind. Occ.* 94.—Jamaica.
hirta, *Klotzsch & Hanst. in Linnæa*, xxxiv. (1865–66) 403.—Costa Rica.
hispida, *Sw. Prod. Veg. Ind. Occ.* 94.—Jamaica.
humilis, *Law. ex Steud. Nom.* ed. I. 213; ed. II. i. 899.—Hab.?
inæqualistera, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 1.—Peruv.
Kalbreyeri, Hook. f. Bot. Mag. t. 6638 = Kalbreyeriana.
Kalbreyeriana, *Mast. in Gard. Chron.* (1882) i. 44 217.—N. Granat.
Kienastiana, *Regel, in Act. Hort. Petrop.* viii. (1883) 374.—N. Granat.
labellosa, *Karst. Fl. Columb.* ii. 105. t. 154.—N. Granat.
lepidocaula, *Hanst. in Linnæa*, xxxiv. (1865–66) 411.—Costa Rica.
Lindleyana, *Pasq. Cat. Orto Bot. Nap.* 31.—Hab.?
linearis, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 133.—Costa Rica.
longepedunculata, Hort. ex Hanst. in Mart. Fl. Bras. viii. 414 = Nematanthus longipes.
longifolia, Linn. Mant. 90 = Artanema sesamoides.
lucida, Walp. Rep. vi. 405. sphalm. = Besleria lucida.
macrantha, *Benth. Pl. Hartw.* 282.—Ecuador.
magnifica, *Klotzsch & Hanst. ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 134.—Costa Rica.
major, *Hanst. in Linnæa*, xxxiv. (1865–66) 388.—N. Granat.
microcalyx, *Hanst. l. c.* 408.—Costa Rica.
microphylla, *Klotzsch & Hanst. ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 137.—Costa Rica.
minor, *Hanst. in Linnæa*, xxxiv. (1865–66) 387.—N. Granat.
minuta, Roxb. Hort. Beng. 47; Fl. Ind. iii. 98 = Vandellia scabra.
moesta, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 1. t. 206.—Peruv.
nervosa, *Klotzsch & Hanst. in Linnæa*, xxxiv. (1865–66) 400.—Costa Rica.
nicaraguensis, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 136.—Nicaragua.
ochroleuca, *Klotzsch & Hanst. in Linnæa*, xxxiv. (1865–66) 393.—Costa Rica.
Oerstediana, *Klotzsch, ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 135.—Costa Rica.
ovata, *Cav. Ic.* iv. 62. t. 391.—Ins. Chiloe.
oxyphylla, *Hanst. in Linnæa*, xxxiv. (1865–66) 405.—Costa Rica.
pendula, *Klotzsch & Hanst. l. c.* 397.—Costa Rica.
pichinchensis, *Hanst. l. c.* 393.—Ecuador.
picta, *Karst. Fl. Columb.* ii. 105. t. 154.—N Granat.
pilosa, Lem. in Fl. des Serres, iii. (1847) t. 223 = aureonitens.
praetexta, *Hanst. in Linnæa*, xxxiv. (1865–66) 394.—Costa Rica.
purpurata, *Hanst. l. c.* 386.—Costa Rica.
querceti, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 132.—Costa Rica.
repens, *Hanst. in Linnæa*, xxxiv. (1865–66) 396.—Costa Rica.
repens, Rafin. Sylva Tellur. 84 = scandens?
ringens, *Regel, in Act. Hort. Petrop.* viii. (1883) 373.—N. Granat.
rotundifolia, *Salisb. Parad. Lond.* t. 22.—Ins. Trinit.
rutilans, *Sw. Prod. Veg. Ind. Occ.* 94.—Jamaica.
sanguinea, *Hanst. in Linnæa*, xxxiv. (1865–66) 384.—Ins. S. Doming.
sanguinolenta, *Hanst. l. c.* 389.—Costa Rica.
scabrosa, Soland. ex DC. Prod. ix. 277 = Rhabdothamnus scabrosus.
scandens, Billh. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 126 = Billbergiana.
scandens, *Linn. Sp. Pl.* 988.—Am. austr.
scandens-hirsuta, Schott, ex Hanst. in Mart. Fl. Bras. viii. 417 = Codonanthe Hookeril.
Schiedeana, *Schlecht. in Linnæa*, viii. (1833) 849.—Mexic.
serrata, *Hanst. in Linnæa*, xxxiv. (1865–66) 390.—Costa Rica.

COLUMNEA:—
speciosa, Presl, Bot. Bemerk. 145 = rotundifolia.
speciosa, Wendl. ex Steud. Nom. ed. II. i. 899 = Rytidophyllum praunatum.
splendens, Part. Mag. Bot. x. (1843) 5 = Nematanthus longipes.
stellata, *Lour. Fl. Cochinch.* 384.—Cochinch.
strigosa, *Benth. Pl. Hartw.* 282.—N. Granat.
tenuis, *Klotzsch, ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 137.—Costa Rica.
tincta, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 526.—Ins. Cuba.
tomentosa, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 138.—Nicaragua.
tomentosa, Roxb. Hort. Beng. 47; Fl. Ind. iii. 98 = Mazus rugosus.
trifoliata, Link, Enum Hort. Berol. ii. 145 = Stemodia trifoliata.
umbellata, *Bertol. Fl. Guatim.* 26.—Guatemala.
vestita, Hanst. in Linnæa, xxxvi. (1865–66) 392 = Warszewicziana.
villosa, Willd. Enum Hort. Berol. Suppl. 43 = scandens.
violacea, Jacq. f. ex Steud. Nom. ed. II. i. 899 = Stemodia trifoliata.
Warszewicziana, *Klotzsch & Hanst. in Linnæa*, xxxiv. (1865–66) 892.—Costa Rica.
Wendlandiana, *Hanst. l. c.* 402.—Costa Rica.
Wullschlaegeliana, *Hanst. l. c.* 414.—Jamaica.
zebrina, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1402 = Alloplectus dichrous.

COLUPPA, Adans. Fam. ii. 268 (1763) = Gomphrena, Linn. (Amarantac.).

COLURIA, R. Br. in Parry, Voy. App. 276 (1823). *ROSACEÆ*, Benth. & Hook. f. i. 619.
LAMARNIA, Flesch. ex Ledeb. Fl. Alt. ii. 262 (1830).
groidez, Ledeb. Fl. Alt. ii. 263 = potentilloides.
mongolica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 466.—As. or.
potentilloides, *R. Br. in Parry, 1st Voy. Suppl.* p. cclxxvi.—Sibir.

COLUS, Raeusch. Nom. ed. III. 171 (1797) = Coleus, Lour. (Labiat.).

COLUTEA, [Tourn.] Linn. Syst. ed. I (1735) *LEGUMINOSÆ*, Benth. & Hook. f. i. 505.
acachmenides, *Sosp. Delic. Insub.* iii. 22. t. 12.—Ins. Baham.
alpina, Lam. Encyc. i. 354 = Astragalus alpinus, membranaceus.
americana, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
annua, Murr. in Nov. Comm. Gotting. 1774 (1775) 40 = halepica.
aperta, Mœnch, Vers. 24 = orientalis.
arborescens, *Linn. Sp. Pl.* 723.—Europ.; Orient.
arenaria, Poir Encyc. Suppl. i. 562 = Astragalus chorinensis.
armena, *Boiss. & Huet. Diagn.* Sér. II. v. 83.—Armenia.
astragalina, Poir. Encyc. Suppl. i. 561 = Astragalus alpinus.
australis, Lam. Encyc. i. 354 = Astragalus australis.
baetica, Poir. Encyc. Suppl. i. 561 = Astragalus lusitanicus.
brevialata, *Lange, Ind. Sem. Hort. Haun.* (1861) 30.—Hab.?
cuspica, Bieb. Fl. Taur. Cauc. ii. 169 = Sphaerophysa salsula.
cilicica, *Boiss. & Bal. Diagn.* Sér. II. v. 83.—Cilicia.
cruenta, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. ii. 55.—Reg. Caucas.
daurica, Spreng. Syst. iii. 242 = Sphaerophysa salsula.
sacica, Thunb. Prod. Pl. Cap. 134 = Lessertia sacica.
fistulosa, Retz. Obs. iii. 40 = Lessertia perennans.
floribunda, Poir. Encyc. Suppl. i. 562 = Sesbania vesicaria.
florida, Salisb. Prod. 337 = arborescens.
frigida, Poir. Encyc. Suppl. i. 561 = Astragalus frigidus.

COLUTEA:—
frutescens, Linn. Sp. Pl. 723 = Sutherlandia frutescens.
galegifolia, Sims, Bot. Mag. t. 792 = Swainsona galegifolia.
glabra, Poir. Encyc. Suppl. i. 561 = Astragalus australis.
grandiflora, Salisb. Prod. 338 = Sutherlandia frutescens
halepica, *Lam. Encyc.* i. 353.—Arab.; Abyssin.
halicacaba, Poir. Encyc. Suppl. i. 562 = Astragalus Valmeranis.
herbacea, Linn. Sp. Pl. 723 = Lessertia linearis.
hirsuta, Roth, Tent. Fl. Germ. i. 305 = arborescens.
humilis, Scop. Delic. Insub. ii. 23 = orientalis.
incana, Poir. Encyc. Suppl. i. 562 = Astragalus anthylloides.
istria, Mill. Gard. Dict. ed. VIII. n. 2 = halepica.
linearis, Thunb. Prod. Pl. Cap. 135 = Lessertia linearis.
media, *Willd. Enum. Hort. Berol.* 771.—Hab.?
melanocalyx, Boiss. & Heldr. Diagn. Ser. I. ix. 35 = arborescens.
microphylla, Delile, Ind. Sem. Hort. Monsp. (1847) 7 = halepica.
nepalensis, Sims, Bot. Mag. t. 2622 = arborescens.
Novæ-Hollandiæ, Walp. Ann. ii. 868 = Clianthus Dampieri.
obtusata, Thunb. Prod. Pl. Cap. 134 = Lessertia obtusata.
orientalis, Lam. Encyc. i. 353 = cruenta.
pallida, Salisb. Prod. 337 = halepica.
perennans, Jacq. Enum. Vindob. App. 31 = Lessertia perennans.
persica, *Boiss. Diagn.* Ser. I. vi. 33.—Persia.
Peucedi, [*Dryand. in*] Ait. Hort. Kew. ed. I. iii. 55 = halepica.
procumbens, Mill. Gard. Dict. ed. VIII. n. 7 = Lessertia procumbens.
prostrata, Thunb. Prod. Pl. Cap. 134 = Lessertia prostrata.
pubescens, Thunb. l. c. = Lessertia pubescens.
rapida, Thunb. l. c. = Lessertia frutiosa.
rubra, Medic. Beobacht. (1782) 359 = arborescens.
salsula, Poir. Encyc. Suppl. i. 562 = Sphaerophysa salsula.
sanguinea, Pall. Fl. Ross. ii. 88 = cruenta.
spinosa, Forsk. Fl. Aegypt. Arab. 181 = Astragalus Russellii.
tomentosa, Thunb. Prod. Pl. Cap. 135 = Lessertia tomentosa.
trifora, Poir. Encyc. Suppl. i. 562 = Astragalus triflorus.
trifoliata, Poir. l. c. (= *Phaca trifoliata*).—China.
triphylla, *Bunge, ex Boiss. Fl. Orient.* ii. 196.—Persia.
vesicaria, Thunb. Prod. Pl. Cap. 135 = Lessertia vesicaria.
versicolor, Salisb. Prod. 337 = cruenta.
Wightiana, Wall. ex Steud. Nom. ed. II. i. 899 = Sutherlandia frutescens.
wolgarica, Lam. Encyc. i. 353 = Calophaca wolgarica.

COLUTEOCARPUS, Boiss. in Ann. Sc. Nat. Sér. II. xxii. (1842) 162. *CRUCIFERÆ*, Benth. & Hook. f. i. 78.
LAGOWSKIA, Trautv. in Bull. Phys.-Math. Acad. Pétersb. xvi. (1858) 321.
reticulatus, *Boiss. l. c.*—As. Min.

COLUTIA, Medic. in Vorles. Cherpf. Phys. Ges. ii. (1787) 366 = Sutherlandia, R. Br. (1812) (Legumin.).
amœna, Medic. l. c. = S. frutescens.
frutescens, Medic. Phil. Bot. i. 210 = S. frutescens.
rubra, Mœnch, Meth. 160 = S. frutescens.

COLVILLEA, Boj. ex Hook. Bot. Mag. tt. 3225, 3226 (1834). *LEGUMINOSÆ*, Benth. & Hook. f. i. 569.
racemosa, *Boj. l. c.*—Madag.

COLYMBADA, Hill, Veg. Syst. iv. 31 (1762) = Centaurea, Linn. (Compos.).
centauroides, Hill, Hort. Kew. 65 = Cent. Rhapontica.
sicula, Hill, l. c. = Cent. sicula.

COLYMBEA, Stend. Nom. Bot. ed. II. i. 399 (1840) = **COLUMBEA**, Salisb. = **Araucaria**, juss. (Conif.).

COLYRIS, Endl. Gen. 596 (1838) = **COLLYRIS**, Vahl = **Dischidia**, R. Br. (Asclepiad.).

COLYTHRUM, Schott, Rutac. 13. t. 7 (1834) = **Boenbeckia**, Knnth (Rutac.).
febrifugum, Schott, l. c. 14 = E. febrifuga.
grandiflorum, Stend. Nom. ed. II. i. 599 = E. grandiflora.
latifolium, Schott, Rutac. 14 = E. pumila.
mauritiodes, Schott, l. c. = E. mauritiodes.
puberulum, Schott, l. c. 13. t. 7.—Bras.
pumilum, Schott, l. c. 14 = E. pumila.

COMACEPHALUS, Klotzsch, in Linnæa, xii. (1838) 324 = **Griesbachia**, Klotzsch (Ericac.).
incurvus, Klotzsch, l. c.—Afr. austr.

COMACLINIUM, Scheidw. & Planch. in Fl. des Serres, viii. (1852) 19. t. 756 = **Dysodia**, Cav. (Compos.).
aurantiacum, Scheidw. & Planch. l. c. = D. grandiflora.

COMACUM, Adans. Fam. ii. 845 (1763) = **Myristica**, Linn.

COMANDRA, Nutt. Gen. Am. i. 157 (1818).
SANTALACEÆ, Benth. & Hook. f. iii. 224.
HAMILTONIA, Spreng. Syst. i. 581 (1825), pro parte.
cuneifolia, Rafin. New Fl. Am. ii. 33 = umbellata.
Darbya, *A. DC. in DC. Prod.* xiv. 687.—Am. bor.
elegans, *Reichb. f. Ic. Fl. Germ.* xi. 11. t. 547. f. 1162. —Europ. or. austr.
elliptica, Rafin. New Fl. Am. ii. 33 = umbellata.
livida, *Richards. in Frankl. Narr. 1st Journ. App.* 734. —Am. bor.
media, Rafin. New Fl. Am. ii. 33 = umbellata.
obovata, Rafin. l. c. = umbellata.
obtusifolia, Rafin. l. c. = umbellata.
pallida, *A. DC. in DC. Prod.* xiv. 686 —Am. bor.
rigida, Stend. Nom. ed. II. i. 399 = livida.
umbellata, *Nutt. Gen. Am.* i. 157.—Am. bor.
umbellulata, Rafin. New Fl. Am. ii. 33 = præc.

COMANTHOSPHACE, S. Moore, in Journ. Bot xv. (1877) 293. *LABIATÆ.*
barbinervis, *S. Moore, l. c.*—Japon.
japonica, *S. Moore, l. c.*—Japon.
stellipila, *S. Moore, l. c.*—Japon.
sublanceolata, *S. Moore, l. c.*—Japon.

COMAROPSIS, Rich. in Nestl. Monog. Potent. 16. t. 1 (1816) = **Waldsteinia**, Willd. (Rosac.).
Donians, Ser. in DC. Prod. ii. 558 = W. fragarioides.
fragarioides, Ser. l. c. = W. fragarioides.
pedata, Ser. l. c. = Rubus pedatus.
radicans, Ser. l. c. = Rubus radicans.
sibirica, Ser. l. c. = W. sibirica.

COMAROSTAPHYLIS, Zucc. in Abh. Akad. Muench. ii. (1837) 331 = **Arctostaphylos**, Adans. (Ericac.).
angustifolia, Klotzsch, in Linnæa, xxiv. (1851) 74 = A. angustifolia.
arbutoides, Lindl. Bot. Reg. (1843) t. 30 = A. arbutoides.
arguta, Zucc. in Linnæa, xii. (1838) Litt. 86 = A. arguta.
attenuata, Klotzsch, l. c. xxiv. (1851) 75 = A. attenuata.
bicolor, Klotzsch, l. c. 78 = A. bicolor.
formosa, Lem. in Illustr. Hortic. v. (1858) sub t. 162 = Pieris formosa.
glauca, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 461 = Arbutus Menziesii.
glaucescens, Zucc. ex Klotzsch, in Linnæa, xxiv. (1851) 76 = A. glaucescens.
Hartwegiana, Klotzsch, in Linnæa, xxiv. (1851) 77 = A. Hartwegiana.
latifolia, Klotzsch, l. c. 76.—Mexic.
longifolia, Klotzsch, l. c. 75 = A. longifolia.
mucronata, Klotzsch, l. c. 77 = A. mucronata.
mucronifera, Klotzsch, l. c. 79 = A. mucronifera.
oaxacana, Klotzsch, l. c. 77 = A. oaxacana.
polifolia, Zucc. ex Klotzsch, l. c. = A. polifolia.
rubescens, Klotzsch, l. c. = A. rubescens.

COMAROUNA, Carr. in Rev. Hortic. (1873) 282 = COUMAROUNA, Aubl. = **Dipteryx**, Schreb. (Legumin.).

COMARUM, Linn. Syst. ed. I (1735) = **Potentilla**, Linn. (Rosac.).
angustifolium, Rafin. Fl. Tellur. ii. 56 = P. Comarum.
digitatum, Rafin. l. c. = P. Comarum.
flavum, Bach.-Ham. ex Roxb. Hort. Beng. 39 ; Fl. Ind. ii. 321 = P. supina.
fragarioides, Roth, Tent. Fl. Germ. ii. 377 = P. fragariastrum.
palustre, Linn. Sp. Pl. 502 = P. Comarum.
rubrum, Gilib. Fl. Lituan. ii. 265 = P. Comarum.
Salessowii, Bunge, Del. Sem. Dorp. (1839) 8.—Cf. Linnæa, xiv. (1840) Litt. 119 = P. Salesoviana.
supinum, Alef. in Bot. Zeit. xxiv. (1866) 262 = P. supina ?

COMASPERMUM, Pers. Syn. ii. 275 (1807) = **Comesperma**, Labill. (Polygal.).

COMATOCROTON, Karst. in C. Koch & Pint. Wochenschr. (1859) 6 = Croton, Linn. (Euphorbiac.)
ovalifolia, Karst. l. c. = C. ovalifolius.

COMATOGLOSSUM, Karst. & Triana, in Linnæa, xxviii. (1856) 486 = **Talisia**, Aubl.? (Sapindac.
strictum, Karst. & Triana, l. c. 497.—N. Granat.

COMBESIA, A. Rich. Tent. Fl. Abyss. i. 307 (1847) = **Tillæa**, Linn. (Crassulac.).
abyssinica, A. Rich. l. c. = T. pharnaceoides.
Schimperi, Schweinf. in Beitr. Fl. Aethiop. 80 = T. pentandra.

COMBRETOCARPUS, Hook. f. in Benth. & Hook. f. Gen. i. 683 (1865). *RHIZOPHOREÆ*, Benth. & Hook. f. l. c.
Motleyi, *Hook. f. l. c.*—Borneo.

COMBRETUM, Linn. Gen. ed. I. App. 306 (1737). *COMBRETACEÆ*, Benth. & Hook. f. i. 688.
ÆTIA, Adans. Fam. ii. 84 (1763).
BURRAVA, Baill. Adansonia, i. 71 (1860).
CALOPYXIS, Tul. in Ann. Sc. Nat. Ser. IV. vi. (1856) 86.
CHRYSOSTACHYS, Pohl, Pl. Bras. ii. 65. t. 143 (1831).
CRISTARIA, Sonner. Voy. Ind. iii. 284 (1782).
EMBRYOGONIUM, Blume, Mus. Bot. Lugd. Bat. ii. 122 (1854).
FORSGARDIA, Vell. Fl. Flum. 152 (1825) ; iv. t. 13 (1827).
GONOCARPUS, Ham. Prod. Fl. Ind. Occ. 89 (1825).
PEVRAEA, Comm. ex Juss. Gen. 390 (1789).
POIVREA, Comm. ex Thou. Obs. Pl. Afr. Austr. 28 (1806).
SEGUIERA, Reichb. ex Oliver, Fl. Trop. Afr. ii. 424 (1871).
SCHREADENDRON, Bertol. f. in Mem. Acc. Sc. Bologn. ii. (1850) 572.
SHEADENDRON, Bertol. f. l. c. t. 40 (1850).
THILOA, Eichl. in Flora, xlix. (1866) 149.
accedens, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 284.—Guiana.
acuelestum, *Vent. Choix*, 58.—Afr. trop.
acuminatum, *Roxb. Hort. Beng.* [38] ; *Fl. Ind.* ii. 238. —Ind. or. ; Malaya.
acutum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 424.—Afr. trop.
adenogonium, Steud. ex A. Rich. Tent. Fl. Abyss. i. 266 = reticulatum.
adenophyllum, Mart. in Flora, xxiv. (1841) ii. Beibl. 1 = Jacquini.
Afzelii, G. Don, in Trans. Linn. Soc. xv. (1827) 487 = grandiflorum.
Afzelii, Hort. ex A. Juss. in Walp. Rep. v. 338 = Hirsea Wiedeana.
albidum, G. Don, in Trans. Linn. Soc. xv. (1827) 429 = ovalifolium.
album, Pers. Syn. i. 411 = decandrum.
Alfredi, *Hance, in Journ. Bot.* ix. (1871) 131.—China.
alternifolium, Herb. Madr. ex Wight & Arn. Prod. 317 = Lumnitzera racemosa.
alternifolium, *Pers. Syn.* ii. 213.—Panama.
alternifolium, Spreng. Syst. ii. 331 = acuelestum.

COMBRETUM :—
altum, *Guill. & Perr. ex DC. Prod.* iii. 20.—Afr. trop.
anfractuosum, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 3. —Bras.
angolense, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 423.—Afr. trop.
anisopterum, *Welw. ex Laws. l. c.* 429.—Afr. trop.
apetalum, *Wall. Cat.* n. 3990.—Burma.
apiculatum, *Sond. in Linnæa*, xxiii. (1850) 45.—Afr. austr.
arboreum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 118.— Malaya.
argenteum, Bertol. Fl. Guatimal. 19 = erianthum.
argyrotrichum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 423.—Afr. trop.
asiaticum, *Eichl. in Mart. Fl. Bras.* xiv. ii. 109.— Bras.
attenuatum, Wall. Cat. n. 3989 = quadrangulare.
Aubletii, *DC. Prod.* i. 18.—Guiana.
aurantiacum, *Benth. in Hook. Journ. Bot.* ii. (1840) 232.—Guiana.
auratum, Stend. Nom. ed. II. i. 400 = seq.
aureum, *Wall. Cat.* n. 3988.—Burma.
barbatum, *Neum. ex G. Don, in Loud. Hort. Brit.* 157. —Bras.
bellum, Stend. Nom. ed. II. i. 400 = trifoliatum.
Benthamianum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 220.—N. Granat.
Binderianum, *Kotschy, in Sitzb. Wien. Akad.* ii. *Abth.* ii. (1865) 363.—Afr. trop.
Blanchetti, *Eichl. in Mart. Fl. Bras.* xiv. ii. 114.— Bras.
bracteatum, Herb. Madr. ex Wall. Cat. n. 4009 = decandrum.
brevistylum, *Eichl. in Mart. Fl. Bras.* xiv. ii. 118.— Bras.
Bugi, Cambess. in St. Hil. Fl. Bras. Mer. ii. 247. t. 130 = Jacquini.
butyrosum, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 87, in nota.—Afr. trop.
capitiliflorum, *Fenzl. ex Schweinf. in Reliq. Kotsch* 28.—Afr. trop.
celastroides, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 422.—Afr. trop.
chinense, *Roxb. Hort. Beng.* [38] ; *Fl. Ind.* ii. 230.— As. trop.
chrysophyllum, Guill. & Perr. Fl. Seneg. Tent. i. 289 = tomentosum.
coccineum, *Lam. Encyc.* i. 734.—Madag. ; Ins. Maurit.
collinum, *Fresen. in Mus. Senckenb.* ii. (1837) 158.— Abyssin.
comosum, *G. Don, in Edinb. Phil. Journ.* (1834) 344. —Afr. trop.
confertum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 422.— Afr. trop.
constrictum, *Laws. l. c.* 422.—Afr. trop.
cordatum, *G. Don, in Trans. Linn. Soc.* xv. (1827) 440.—Ins. S. Doming.
corylifolium, *Griseb. Fl. Brit. W. Ind.* 711.—Ind. occ.
corymbosum, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 135 = racemosum.
costatum, Roxb. Hort. Beng. [38] ; Fl. Ind. ii. 237 = acuminatum.
cuspidatum, *Planch. ex Benth. in Hook. Niger Fl.* 340.—Afr. trop.
cyclophyllum, Stend. Nom. ed. II. i. 400 = extensum.
dasystachyum, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 187.—Reg. Himal. ; Burma.
decandrum, Jacq. Enum. Pl. Carib. 19 = alternifolium.
decandrum, Knis & Pav. ex G. Don, Gen. Syst. ii. 665 = Parcoli.
decandrum, *Roxb. Pl. Corom.* i. 43. t. 59.—Ind. or. ; Burma.
dipterum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 420.—Afr. trop.
distillatorium, Blanco, Fl. Filip. ed. I. 295 = ovalifolium.
Duarteanum, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 248.—Bras.
elæagnifolium, Planch. in App. Speke, Journ. Disc. Nile, 534 = collinum.
elæagnoides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 78.—Afr. trop.
elegans, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 247.— Bras.
elegans, H. B. & K. Nov. Gen. et Sp. vi. 100. t. 558 = Aubletii.

COMBRETUM :—
elegans, Wall. Cat. n. 4003 = trifoliatum.
erianthum, *Benth. Pl. Hartw.* 72.—Guatemala.
eriopetalum, *G. Don, Gen. Syst.* ii. 866.—Ins. S. Doming.
erythrophyllum, *Sond. in Linnaea*, xxiii. (1850) 43.—Afr. austr.
extensum, *Roxb. Hort. Beng.* [38] ; *Fl. Ind.* ii. 229.—Ind. or. ; Malaya.
farinosum, *H. B. & K. Nov. Gen. et Sp.* vi. 110.—Mexic.
ferrugineum, *G. Don, in Trans. Linn. Soc.* xv. (1827) 440.—Am. austr.
ferrugineum, Hoffmgg. ex Walp. Rep. ii. 65 = Jacquini.
ferrugineum, A. Rich. Tent. Fl. Abyss. i. 267 = trichanthum.
flagrocarpum, *Herb. Calc. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 455.—Ind. or. ; Burma.
floccosum, *Eichl. in Mart. Fl. Bras.* xiv. ii. 119.—Bras.
formosum, *G. Don, in Trans. Linn. Soc* xv. (1827) 420.—Bras.
formosum, Griff. Notul. iv. 683 = extensum.
frangulæfolium, *H. B. & K. Nov. Gen. et Sp.* vi. 109.—Venezuela.
fuscum, *Planch. ex Benth. in Hook. Niger Fl.* 339.—Afr. trop.
gallaetense, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 664.—Afr. trop.
Gehesebii, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 232.—Bras.
glabrum, *DC. Prod.* iii. 19.—Guiana.
glaucocarpum, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 8.—Bras.
glomeruliflorum, *Sond. in Linnaea*, xxiii. (1850) 42.—Afr. austr.
glutinosum, *Guill. & Perr. Fl. Seneg. Tent.* 288. t. 68.—Afr. trop.
glutinosum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 268 = lepidotum.
Goldieanum, *F. Muell. Desc. Papuan Pl. iv.* 66.—N. Guinea.
gracile, *Schott, in Spreng. Syst. iv. App.* 406.—Bras.
grandiflorum, *G. Don, in Edinb. Phil. Journ.* (1824) 346.—Afr. trop.
Griffithii, Heurck & Muell. Arg. in Heurck, Obs. Bot. 231 = chinense.
Gueinzii, *Sond. in Linnaea*, xxiii. (1850) 43.—Afr. austr.
guianense, Miq. in Ann. Mag. Nat. Hist. xi. (1843) 13 = Aubletii.
Hartmannianum, *Schweinf. Beitr. Fl. Aethiop.* 24. t. 3.—Afr. trop.
herbaceum, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 347.—Afr. trop.
Heyneanum, Wall. Cat. n. 4001 = ovalifolium.
Hilariana, *D. Dietr. Syn. Pl.* ii. 1303.—Bras.
hispidum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 421.—Afr. trop.
holosericeum, Sond. in Linnaea, xxiii. (1850) 44.—Afr. austr.
Hostyldii, Miq. Fl. Ind. Bat. i. 1. 609 = extensum.
imberbe, *Wawra, in Sitzb. Acad. Wien, Math.-Nat.* xxxviii. (1860) 556.—Afr. trop.
insigne, Heurck & Muell. Arg. in Heurck, Obs. Bot. 247 = pilosum.
intermedium, G. Don, in Edinb. Phil. Journ. xi. (1824) 346 = comosum.
Jacquini, *Gritsch. Fl. Brit. W. Ind.* 276.—Ind. occ.
Kirkii, *Laws. in Oliver, Fl. Trop. Afr.* ii. 429.—Afr. trop.
Klotzschii, *Welw. ex Laws. l. c.* 422.—Afr. trop.
Kraussii, *Hochst. in Flora*, xxvii. (1844) ii. 484.—Afr. austr.
laetum, Wall. Cat. n. 4004 = pilosum.
lanceolatum, *Pohl, ex Eichl. in Mart. Fl. Bras.* xiv. ii. 110.—Bras.
lanuginosum, *G. Don, in Trans. Linn. Soc.* xv. (1827) 423.—Abyss.
latifolium, Bazme, Bijdr. 641 = extensum.
latifolium, *G. Don, in Trans. Linn. Soc.* xv. (1827) 432.—Ind. or.
laurifolium, *Mart. in Flora*, xxii. (1839) i. *Beibl.* 64.—Bras.
laxiflorum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 428.—Afr. trop.
leaeum, Arabl. Fl. Gui. 351 = Aubletii.
leaeum, Jacq. Enum. Pl. Carib. 19 = Jacquini.

COMBRETUM :—
laxum, Loefl. it. 248, 308 = Loeflingii.
laxum, Roxb. Hort. Beng. [38] ; Fl. Ind. ii. 231 = ovalifolium.
lepidotum, Presl, Bot. Bemerk. 142 = squamosum.
lepidotum, *A. Rich. Tent. Fl. Abyss.* i. 268.—Abyss.
leprosum, *Mart. in Flora*, xxiv. (1841) ii. *Beibl* 1.—Bras.
leptostachyum, *Mart. l. c.* 2.—Bras.
leucanthum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 240 = extensum.
leucophyllum, G. Don, in Edinb. Phil. Journ. xi. (1824) 346 = racemosum.
Loeflingii, *Eichl. in Mart. Fl. Bras* xiv. ii. 110.—Am trop.
lucidum, Blume, Bijdr. 641 = trifoliatum.
macrocarpum, Beauv. Fl. Owar. ii. 90. t. 118 = racemosum.
macrophyllum, Roxb. Hort. Beng. [38] ; Fl. Ind. ii. 231 = latifolium.
macrostachyum, Wall. Cat. n. 3997 = extensum.
Madolata, Buch.-Ham. ex Wall. Cat. n. 4009 = decandrum.
magnificum, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 4.—Bras.
Malaiuca, Wall. Cat. n. 3991 = squamosum.
Mechowianum, *O. Hoffm. in Linnaea*, xliii. (1880–82) 151.—Afr. trop.
mellifluum, *Eichl. in Mart. Fl. Bras.* xiv. ii. 113.—Bras.
mexicanum, *Humb. & Bonpl. Pl. Aequin.* ii. 159. t. 132.—Mexic.
micranthum, G. Don, in Edinb. Phil. Journ. xi. (1824) 347 = altum.
microphyllum, DC. Prod. iii. 19 = Loeflingii.
microphyllum, Llanos, in Mem. Acad. Cienc. Madr. iv. (1856) 502 = extensum.
microphyllum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 74.—Afr. trop.
molle, *R. Br. ex G. Don, in Trans. Linn. Soc.* xv. (1827) 431.—Abyss.
Moneraia, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 2.—Bras.
mucronatum, *Schum. & Thonn. Beskr. Guin. Pl.* 184.—Afr. trop.
myrtifolium, *Laws. in Oliver, Fl. Trop. Afr.* ii. 431.—Afr. trop.
nanum, *Lech.-Ham. in D. Don, Prod. Fl. Nep.* 219.—Reg. Himal.
neurophyllum, Miq. Fl. Ind. Bat. i. 1. 608 = acuminatum.
nigricans, *Lepriour, ex Guill. & Perr. Fl. Senog. Tent.* i. 290.—Afr. trop.
nitidum, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiv. ii. 118.—Bras.
obscurum, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 88.—Madag.
obtusifolium, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 106.—Guian.
obtusum, Walp. Rep. ii. 67 = prunei.?
occidentalis, *Linn. Syst. ed.* X. 999.—Am. austr.
odoratum, *Pers. ex G. Don, in Trans. Linn. Soc.* xv. (1827) 430.—Ecuador.
ovale, R. Br. ex G. Don, l. c. 434 = aculeatum.
ovalifolium, *Roxb. Hort. Beng.* [38] ; *Fl. Ind.* ii. 226.—Ind. or.
oxypetalum, G. Don, in Trans. Linn. Soc. xv. (1827) 420 = Loeflingii.
oxystachyum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 422.—Afr. trop.
pachycladum, *Baker, in Journ. Linn. Soc.* xviii. (1881) 270.—Madag.
paniculatum, *Vent. Choix*, t. 58.—Afr. trop.
paradoxum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 430.—Afr. trop.
parviflorum, *Eichl. in Mart. Fl. Bras.* xiv. ii. 114.—Bras.
parviflorum, Reichb. ex DC. Prod. iii. 21 = altum.
Pavoeii, *G. Don, in Trans. Linn. Soc.* xv. (1827) 436.—Ecuador.
pedicellatum, *DC. Prod.* iii. 20.—Bras.
pentagonum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 424.—Afr. trop.
Petitianum, *A. Rich. Tent. Fl. Abyss.* i. 268.—Abyss.
phaeocarpum, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 1.—Bras.
pilosum, *Roxb. Hort. Beng.* [38] ; *Fl. Ind.* ii. 231.—Ind. or. ; Burma.

COMBRETUM :—
Pinciauum, *Hook. Bot. Mag.* (1846) sub t. 4262.—Afr. trop.
pisoniæflorum, *Klotzsch, in Peters, Reise Mossamb. Bot.* t. 14.—Afr. trop.
platypetalum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 433.—Afr. trop.
platyphyllum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 242 = extensum.
polyemdrum, Hoffmgg. ex Walp. Rep. ii. 6 = Moquilia parecnia.
Poriersanum, Wall. Cat. n. 4000 = chinense.
puberum, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 108.—Guiana.
pulchellum, Mart. in Flora, xxii. (1839) i. Beibl. 64 = Jacquini.
punctatum, A. Rich. Tent. Fl. Abyss. i. 266 = lepidotum.
punctatum, *Blume, Bijdr.* 640.—Java.
punctatum, Steud. in Flora, xxvi. (1843) ii. 761 = Aubletii.
purpureum, Vahl, Symb. Bot. iii. 51 = coccineum.
pyramidatum, *Desv. in Ham. Prod. Fl. Ind. Occ.* 85.—Guiana.
pyrifolium, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 188.—Burma.
quadrangulare, *Kurz, l. c.*—Burma.
Quartinianum, A. Rich Tent. Fl. Abyss. i. 266 = lepidotum.
racemosum, *Beauv. Fl. Owar.* 87. t. 118.—Afr. trop.
reticulatum, *Fresen. in Mus. Senckenb.* ii. (1837) 154.—Abyss.
reticulatum, *Presl, Reliq. Haenk.* ii. 25.—Mexic.
Richardsianum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 234.—Abyss.
riparium, *Sond. in Linnaea*, xxiii. (1850) 47.—Afr. austr.
Rochellanum, *A. Rich. ex A. Juss. in Ann. Sc. Nat.* Sér. III. xv. (1851) 369.—Abyss.
rotundifolium, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Aubletii.
rotundifolium, Roxb. Hort. Beng. [38] ; Fl. Ind. ii. 226 = extensum.
Roxburghii, G. Don, in Trans. Linn. Soc. xv. (1827) 429 = ovalifolium.
Roxburghii, Spreng. Syst. ii. 331 = decandrum.
rubiginosum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 428.—Afr. trop.
Rueppellianum, *A. Rich. Tent. Fl. Abyss.* i. 267.—Abyss.
salicifolium, *E. Mey. ex Hook. fc. t.* 592.—Afr. austr.
sarcopterum, Thw. Enum. Pl. Zeyl. 115 = acuminatum.
Schimperianum, *A. Rich. Tent. Fl. Abyss.* i. 265.—Abyss.
secundum, G. Don, in Trans. Linn. Soc. xv. (1827) 419 = Aubletii.
secundum, Jacq. Enum. Pl. Carib. 19 = Jacquini.
semiadnatum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 244.—Ind. or.
sericeum, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 347.—Afr. trop.
sericeum, Wall. ex C B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 450 = Calycopteris floribunda.
Smeathmanni, G. Don, in Trans. Linn. Soc. xv. (1827) 425 = racemosum.
Sonderi, *Germ. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 511.—Afr austr.
spinescens, Wall. Cat. n. 4006 = pilosum.
spinosum, G. Don, in Edinb. Phil. Journ. xi. (1824) 345 = paniculatum.
spinosum, Humb. & Bonpl. Pl. Aequin. ii. 161 = alternifolium.
Sprucei, *Eichl. in Mart. Fl. Bras.* xiv. ii. 115.—Bras.
squamosum, *Roxb. Hort. Beng.* [38] ; *Fl. Ind.* ii. 231. —Ind. or. ; Malaya.
stelligerum, *Presl, Epim. Bot.* 215.—Ind. or.
stenopetalum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 225 = acuminatum.
stigmarium, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 3.—Bras.
subalternans, Wall. Cat. n. 4005 = trifoliatum.
sundaicum, *Miq. Fl. Ind. Bat. Suppl.* 327.—Sumatra.
terminalioides, *Steud. in Flora*, xxvi. (1843) ii. 762.—Guiana.
ternatum, Wall. Cat. n. 4002 = pilosum.
tetragonocarpum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 306.—Burma.

COMBRETUM, L.

tetragonum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 480.—Afr. trop.

tetragonum, *Presl, ex Steud. Nom.* ed. II. i. 400.—Mexic.

tetralophum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 464.—Malaya.

Themehli, *D. Hoffm. in Linnaea*, xliii. (1880–82) 182.—Afr. trop.

Thouarsianum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 238 = ovalifolium.

tinctorum, Walw. ex Laws. in Oliver, Fl. Trop. Afr. ii. 480 = Kirkii.

tomentosum, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 346.—Afr. trop.

tomentosum, Willd. ex Eichl. in Mart. Fl. Bras. xiv. ii. 111 = frangulæfolium.

trichanthum, *Fresen. in Mus. Senckenb.* ii. (1837) 155.—Abyssin.

trifoliatum, *Vent. Choix*, t. 58.—Hab. ?

trigonoides, Perr. ex DC. Prod. iii. 20 = racemosum.

truncatum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 427.—Afr. trop.

undulatum, Wall. Cat. n. 3993 = trifoliatum.

Vahlenckii, *Muell. Arg. in Heurck, Obs. Bot.* 227.—Ind. or.

variabile, *Mart. in Flora*, xxii. (1839) i. *Beibl.* 52.—Bras.

velutinum, *DC. Prod.* iii. 20.—Bras.

vernicosum, Fenzl, in Flora, xxvii. (1844) 312 = lepidotum.

villosum, *Boj. ex Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 80, in syn.—Madag.

violaceum, *Boj. ex Tul. l. c.* 79.—Madag.

virgatum, Wall. Cat. n. 3992 = apetalum.

virgatum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 435.—Afr. trop.

viscidum, *Griseb. Cat. Pl. Cub.* 109.—Cuba.

Wallichii, *DC. Prod.* iii. 21.—Reg. Himal.

Wallichii, Kurz, For. Fl. Burm. i. 465 = flagrocarpum.

Wightianum, Thw. Enum. Pl. Zeyl. 103, partim = ovalifolium.

Wightianum, Wall. Cat. n. 4007 = extensum.

Zeyheri, *Sond. in Linnaea*, xxiii. (1850) 46.—Afr. austr.

COMBSPERMA, Labill. Nov. Holl. Pl. ii. 21. t. 159—163 (1806). *POLYGALEAE*, Benth. & Hook. f. i. 136.

acerosum, *Steetz, in Lehm. Pl. Preiss.* ii. 299.—Austral.

acutifolium, Steetz, l. c. 296 = ericinum.

aemulum, Steud. l. c. i. 308 = virgatum.

anthocarpum, Walp. Rep. v. 66, sphalm. = *xanthocarpum*.

aphyllum, *R. Br. ex Benth. Fl. Austral.* i. 148.—Austral.

calycoptera, *Labill. Nov. Holl. Pl.* ii. 23.—Austral.

ciliatum, *Steetz, in Lehm. Pl. Preiss.* ii. 304.—Austral.

compactum, Sieber, ex Steud. Nom. ed. II. i. 400 = retusum.

confertum, *Labill. Nov. Holl. Pl.* ii. 23, t. 161.—Austral.

contractum, Steud. in Lehm. Pl. Preiss. i. 298 = virgatum.

cordifolium, A. Cunn. in Field, N. S. Wales, 337 = ericinum.

corniculatum, Steud. in Lehm. Pl. Preiss. i. 306 = virgatum.

defoliatum, *F. Muell. Vict. Pl.* i. 189.—Austral.

Drummondii, *Steetz, in Lehm. Pl. Preiss.* ii. 301.—Austral.

ericinum, *DC. Prod.* i. 394.—Austral.

flavum, *DC. l. c.*—Austral.

floribundum, St. Hil. Fl. Bras. Mer. ii. 55. t. 91 = Bredemeyera floribunda.

gracile, Pax. Mag. Bot. v. (1838) 145 = volubile.

herbaceum, Steud. in Lehm. Pl. Preiss. i. 311 = calymega.

hirtulum, Steud. l. c. 309 = confertum.

integerrimum, *Endl. in Enum. Pl. Hueg.* 7.—Austral.

leucalys, Spreng. Syst. iii. 173 = calymega.

Xanthianum, St. Hil. Fl. Bras. Mer. ii. 54. t. 90 = Bredemeyera Xanthiana.

lanceolatum, *R. Br. ex Benth. Fl. Austral.* i. 148.—Austral.

latifolium, Steetz, in Lehm. Pl. Preiss. ii. 295 = ericinum.

COMBSPERMA:—

laurifolium, St. Hil. Fl. Bras. Mer. ii. 56 = Bredemeyera laurifolia.

laxiusculum, Steud. in Lehm. Pl. Preiss. i. 310 = virgatum.

Lhotskyanum, Manso, ex Klotzsch, in Peters, Reise Mossamb. Bot. 117, nomen (= Bredemeyerae sp.).—Afr. trop.

linariaefolium, A. Cunn. ex Steetz, in Lehm. Pl. Preiss. ii. 297 = ericinum.

longebracteatum, Steud. l. c. i. 307 = virgatum.

longifolium, Steud. l. c. 306 = confertum.

lucidum, Baill. Hist. Pl. v. 82 = Bredemeyera lucida.

Mansonii, Lhotsky, ex Klotzsch, in Peters, Reise Mossamb. Bot. 117 = Bredemeyera lucida.

megapteryga, Steud. in Lehm. Pl. Preiss. i. 307 = undulosum.

nudiusculum, *DC. Prod.* i. 394.—Austral.

nudiusculum, Steetz, in Lehm. Pl. Preiss. ii. 308 = defoliatum.

parviflorum, Steud. l. c. i. 310 = calymega.

patentifolium, F. Muell. Fragm. i. 48 = ericinum.

pancifolium, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 352 = volubile.

planifolium, *Hassk. in Mig. Ann. Mus. Bot. Lugd. Bat.* i. 185.—Austral.

polygaloides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 7.—Austral.

praecelsum, *F. Muell. Fragm.* xi. 2.—Austral.

pubescens, Gardn. ex A. W. Benn. in Mart. Fl. Bras. xiii. 111. 49 = Bredemeyera brevifolia.

ramentiferum, Steud. in Lehm. Pl. Preiss. i. 309 = undulosum.

retusum, *Labill. Nov. Holl. Pl.* ii. 22. t. 160.—Austral.

rhadinocarpum, *F. Muell. Fragm.* xi. 1.—Austral.

roseum, Steud. in Lehm. Pl. Preiss. i. 307 = virgatum.

scandens, Steud. l. c. 311 = integerrimum.

scoparium, *Drumm. in Hook. Journ. Bot.* ii. (1840) 209; *Steetz, in Lehm. Pl. Preiss.* ii. 309.—Austral.

secundum, *Banks, ex DC. Prod.* i. 394.—Austral.

telagiroides, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 352 = virgatum.

silvestre, F. Muell. Fragm. i. 49 = sylvestre.

simplex, Endl. in Enum. Pl. Hueg. 7 = virgatum.

spathulatum, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 352 = calymega.

sphaerocarpum, *Steetz, in Lehm. Pl. Preiss.* ii. 314.—Austral.

spinosum, *F. Muell. Fragm.* i. 144.—Austral.

strictum, Endl. in Enum. Pl. Hueg. 7 = calymega.

subbracteatum, Walp. Ann. ii. 84, sphalm. = calymega.

sylvestre, *Lindl. in Mitch. Journ. Trop. Austral.* 342.—Austral.

tenue, Steud. in Lehm. Pl. Preiss. i. 308 = calymega.

tortuosum, Steetz, l. c. ii. 308 = volubile.

variabile, Mart. ex Steud. Nom. ed. II. i. 400 (= Bredemeyerae sp.).—Bras.

varians, Steud. in Lehm. Pl. Preiss. i. 310 = calymega.

virgatum, *Labill. Nov. Holl. Pl.* ii. 21. t. 159.—Austral.

viscidulum, *F. Muell. Fragm.* x. 4.—Austral.

volubile, *Labill. Nov. Holl. Pl.* ii. 24.—Austral.

xanthocarpum, Steud. in Lehm. Pl. Preiss. i. 309 = flavum.

COMETES, Linn. Mant. i. 4 (1767). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 18.

CERATONYCHIA, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1215.

SALTIA, R. Br. in Salt, Abyss. App. 64 (1814), nomen. abyssinica, *R. Br. in Wall. Pl. As. Rar.* i. 18.—Abyssin.

altermifolia, Linn. Syst. ed. XII. 137 = aurattensis.

aphylla, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 344 = aurattensis.

aurattensis, *Linn. Mant.* i. 39; *Burm. f. Fl. Ind.* 39. t. 15.—Ind. bor. occ. ; Arab.

COMETIA, Thou. ex Baill. Etud. Gén. Euph. 642 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 353.

lucida, *Baill. Adansonia*, ii. (1861) 55.—Madag.

Thouarsii, *Baill. Etud. Gén. Euph.* 643.—Madag.

COMEURYA, Baill. Adansonia, x. (1872) 329 = Dracontomelum, Blume (Account.).

Cumingiana, Baill. l. c. 330 = D. Cumingianum.

COMMARUM, Schrank, Prim. Fl. Salish. 132 (1792) = COMARUM, Linn. = Potentilla, Linn. (Rosac.).

COMMELINA, Plum. ex Linn. Syst. ed. I. (1735). *COMMELINACEAE*, Benth. & Hook. f. iii. 647.

ANANTHOPUS, Raﬁn. Fl. Ludov. 20 (1817).

DICHORAPHIS, Hassk. in Flora, xlix. (1866) 211.

EELEMIA, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 494.

HEDWIGIA, Medic. l. c. 495 (1790).

HETEROCARPUS, Wight, Ic. vi. 29. t. 2067 (1853).

LECHEA, Lour. Fl. Cochinch. 60 (1790).

OMPHALOTHECA, Hassk. in Bull. Congr. Bot. Amst. (1865) 109.

SPATHODITHYROS, Hassk. in Flora, xlix. (1866) 211.

TRITHYROCARPUS, Hassk. l. c. (1866).

ALLOSPERMA, ALLOTRIA, DITELESIA, DIETEA, EUDIPETALA, LARNALIES, NEPHRALES, et OVIDIA, Raﬁn. Fl. Tellur. iii. 68–70, 129 (1836).

acuminata, *H. B. & K. Nov. Gen. et Sp.* i. 258.—Mexic.

acuminata, Poir. Encyc. Suppl. ii. 324 = Aneilema acuminatum.

acuminata, R. Br. in Salt, Abyss. App. 62 = pyrrhobiephantis.

aequinoctialis, Roem. & Schult. Syst. i. 538 = Aneilema aequinoctiale.

aequinoctialis, Beauv. Fl. Owar. i. t. 38 = prasc.

aethiopica, *C. B. Clarke, in DC. Monog. Phan.* iii. 189.—Afr. trop.

affinis, Poir. Encyc. Suppl. ii. 324 = Aneilema gramineum.

africana, *Linn. Sp. Pl.* 41.—Afr. austr.

africana, Mirb. in Ann. Mus. Par. xvi. (1810) 456 = communis.

agraria, Kunth, Enum. Pl. iv. 38 = nudiflora.

agrostophylla, *F. Muell. Fragm.* viii. 59.—Austral

alba, Buch.-Ham. ex C. B. Clarke, in DC. Monog. Phan. iii. 186 = appendiculata.

albescens, *Hassk. in Schweinf. Beitr. Fl. Aethiop.* 210.—Afr. trop.

ambigua, Beauv. Fl. Owar. i. 26. t. 15 = Pallasia ambigua.

amplexicaulis, *Hassk. in Schweinf. Beitr. Fl. Aethiop.* 208.—Afr. trop.

angolensis, *C. B. Clarke, in DC. Monog. Phan.* iii. 167.—Afr. trop.

angustifolia, Hassk. in Peters, Reise Mossamb. Bot. 538 = Gerrardi.

angustifolia, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = graminifolia.

angustifolia, Michx. Fl. Bor. Am. i. 24 = virginica.

angustifolia, Reichb. Ic. Bot. Exot. ii. 17. t. 148 = graminifolia.

anthericoides, Poir. Encyc. Suppl. ii. 324 = Aneilema graminosum.

appendiculata, *C. B. Clarke, Commel. et Cyrt. Beng.* t. 18.—Ind. or.

aspera, *G. Don, ex Benth. in Hook. Niger Fl.* 562.—Afr. trop.

attenuata, Heyne, ex Steud. Nom. ed. II. i. 401 = subulata.

attenuata, *Kunth, ex Vahl, Enum. Hell. Pl.* ii. 168.—Ind. or.

auriculata, *Blume, Enum. Pl. Jav.* i. 2.—Sumatra.

auriculata, E. Mey. in Presl, Rel. Haenk. i. 137 = virginica.

avenacifolia, *Y. Grah. Cat. Pl. Bombay*, 234.—Ind. or.

axillaris, Linn. Sp. Pl. 42 = Cyanotis axillaris.

bahiensis, Hoffmgg. ex Sprog. Jahrb. i. 111. 73 = virginica.

Balnedi, *C. B. Clarke, in DC. Monog. Phan.* iii. 184.—Afr. austr.

barbata, Boj. Hort. Maurit. 259 = communis.

barbata, *Lam. Tabl. Encyc.* i. 129.—Am. austr.

barbata, Sieber, ex C. B. Clarke, in DC. Monog. Phan. iii. 247 = Cyanotis cristata.

benghalensis, Forsk. Fl. Aegypt. Arab. 12 = divaricata.

benghalensis, *Linn. Sp. Pl.* 41.—Geront. trop.

benghalensis, Wall. Cat. n. 8980 = persicariaefolia.

bracteolensis, Beauv. Fl. Owar. ii. 49. t. 87 = Aneilema bambusae.

bicolor, Poepp. ex Kunth, Enum. Pl. iv. 89 = Tradescantia gracilis.

biflora, Poir. Encyc. Suppl. ii. 324 = Aneilema biflorum.

Blumei, A. Dietr. Sp. Pl. ii. 400 = sarcinulata.

Boissieriana, *C. B. Clarke, in DC. Monog. Phan.* iii. 161.—Afr. trop.

brachiolata, Linn. Encyc. ii. 49 = Aneilema spicatum.

COMMELINA :—
bracteosa, *Hassk. in Flora*, xlvi. (1863) 386 —Afr. trop.
Brownii, A. Dietr. Sp. Pl. ii. 415 = Anellema acuminatum.
bulbosa, Heyne, ex Steud. Nom. ed. II. i. 401 = Anellema dimorphum.
caerulea, Salisb. Prod. 216 = virginica.
caespitosa, Roxb. Hort. Beng. 5; Fl. Ind. i. 174 = nudiflora.
cajanonala, *Sieber, ex Steud. Nom.* ed. II. i. 401.—Ins. Martinic.
cajennensis, Kunth, Enum. Pl. iv. 38, in syn. = nudiflora.
Callisia, Steud. Nom. ed. II. i. 401 = Anellema Hamiltonianum.
canariensis, Ch. Sm. in Buch, Canar. Ins. 149 = nudiflora.
canescens, Vahl, Enum. ii. 178 = benghalensis.
capitata, *Benth. in Hook Niger Fl.* 541.—Afr. trop.
capitata, Klotzsch, ex C. B. Clarke, in DC. Monog. Phan. iii. 178 = obliqua.
cardiocephala, Kunze, ex C. B. Clarke, l. c. 157 = quitensis.
cardiospala, Kunze, in Linnaea, xx. (1847) 9 = quitensis.
caripensis, H. B. & K. Nov. Gen. et Sp. i. 260 = virginica.
carnea, Schlecht. Ind. Sem. Hort. Halens. (1839) adn. 9 = scabra.
caroliniana, Herb. Berol. (1841) ex Kunth, Enum. Pl. iv. 59 = stricta.
caroliniana, Walt. Fl. Carol. 68 = nudiflora.
caroliniana, Willd. ex Kunth, Enum. Pl. iv. 54 = angustifolia.
cayennensis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 105 = nudiflora.
Chamissonis, *Klotzsch, ex C. B. Clarke, in DC. Monog. Phan.* iii. 186.—Austral.
Chantransia, Roem. & Schult. Mant. i. Add. i. 376 = Floscopa africana.
clandestina, Hort. Berol. (1832) ex Kunth, Enum. Pl. iv. 46 = graminifolia.
clandestina, *Mart. ex Steud. Nom.* ed. II. i. 401.—Bras.
clavata, *C. B. Clarke, Commel. et Cyrt. Beng.* t. 5.—Ind. or. ; Malaya.
coelestis, *Willd. Enum. Hort. Berol.* 69.—Mexic.
commelinoides, *Forsk. Fl. Aegypt. Arab.* 12.—Arab.
communis, Engelm. ex Kunth, Enum. Pl. iv. 38 = agnuia.
communis, F. Muell. Fragm. vill. 59 = cyanea.
communis, *Linn. Sp. Pl.* 40.—China.
communis, Roxb. Hort. Beng. 5; Fl. Ind. i. 171 = obliqua.
communis, Wall. Cat. n. 8978 = sallcifolia, Haskarlii.
communis, Walt. Fl. Carol. 68 = nudiflora.
comosa, E. Mey. Zwei Pfl Docum. 174 = Anellema Dregeanum.
condensata, *C. B. Clarke, in DC. Monog. Phan.* iii. 190.—Afr. trop.
congesta, *C. B. Clarke, l. c.* 160.—Afr. trop.
conspicua, Blume, Enum. Pl. Jav. i. 4 = Anellema conspicuum.
conspicua, Zoll. Syst. Verz. Ind. Arch. 64 = Anellema monadelphum.
cordifolia, *A. Rich. Tent. Fl. Abyss.* ii. 341.—Abyss.
crassicaulis, *C. B. Clarke, in DC. Monog. Phan.* iii. 149.—Afr. trop.
crispata, Poir. Encyc. Suppl. ii. 324 = Pollia crispata.
cristata, Linn. Sp. Pl. 42 = Cyanotis cristata.
cucullata, Linn. Mant. ii. 178 = benghalensis.
cyanea, *R. Br. Prod.* 269.—Austral.
cymosa, Blume, Enum. Pl. Jav. i. 4 = Floscopa scandens.
debilis, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 59 = graminifolia.
debilis, Ledeb. in Roem. & Schult. Mant. i. 349 = virginica.
decumbens, E. Mey. in Presl, Rel. Haenk. i. 136 = pallida.
dichotoma, *Herb. ex Steud. Nom.* ed. II. i. 401.—Bras.
diffusa, Hook. Bot. Mag. t. 3844 = virginica.
difficiens, Van Houtte, Fl. des Serres, xvill. (1862) t. 1824 = nudiflora.
dolichotula, Schlecht. in Gartenfl. (1856) 349. t. 104 = benghalensis.
dregeiana, Blume, Enum. Pl. Jav. i. 4 = Floscopa scandens.
dicandra, Steud. Nom. ed. II. i. 401 = Anellema nudiflorum.
dianthifolia, *[Delile, in] Red. Lil.* vii. t. 390.—Mexic.

COMMELINA :—
didyma, Steud. Nom. ed. II. i. 401 = Pollia sorsogonensis.
diffusa, Barm. f. Fl. Ind. 18. t. 7. f. 2 = nudiflora.
diffusa, Willd. ex Kunth, Enum. Pl. iv. 39, 40 = salicifolia, barbata.
diffusa, Zoll. ex C. B. Clarke, in DC. Monog. Phan. iii. 171 = clavata.
divaricata, *Vahl, Enum.* ii. 169.—Arab.
Dorii, A. Dietr. Sp. Pl. ii. 396 = obliqua.
dubia, *[Delile, in] Red. Lil.* t. 359.—Mexic.
dubia, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = coelestis.
dubia, Jacq. Obs. Bot. iii. 9 = Heteranthera graminea.
ebracteata, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 210 = Anellema Ehrenbergii.
Eckloniana, *Kunth, Enum. Pl.* iv. 57.—Austral.
edulis, *A. Rich. Tent. Fl. Abyss.* ii. 341.—Abyss.
edulis, Stokes, Bot. Mat. Med. i. 184 = tuberosa.
Ehrenbergiana, Link, Klotzsch & Otto, Ic. Pl. Rar. 75. t. 30 = scabra.
elata, Vahl, Enum. ii. 178 = Anellema herbaceum.
elegans, H. B. & K. Nov. Gen. et Sp. i. 259 = virginica.
elliptica, *R. Br. & K. l. c.*—Am. trop.
ensifolia, F. Muell. Fragm. viii. 60 = undulata.
ensifolia, *R. Br. Prod.* 269.—Austral.
equinoctialis, Beauv. Fl. Owar. i. 65 = Anellema aequinoctiale.
erecta, Chapm. Fl. S. U. St. 497 = hirtella.
erecta, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = coelestis.
erecta, *Linn. Sp. Pl.* 41.—Am. bor.
esculenta, Herb. Madr. ex Wall. Cat. sub n. 5208 = Anellema esculentum.
esilis, Steud. Nom. ed. II. i. 401 = Anellema nudiflorum.
falcata, Hassk. in Flora, xlvi. (1863) 387 = Forskalaei.
fasciculata, *Ruiz & Pav. Fl. Per.* i. 44. t. 72. f. 6.—Peru.
filiformis, Steud. Nom. ed. II. i. 401 = Anellema vaginatum.
firma, Welw. ex C. B. Clarke, in DC. Monog. Phan. iii. 180 = aspera.
flava, Salisb. Prod. 215 = africana.
flexuosa, Welw. ex C. B. Clarke, in DC. Monog. Phan. iii. 208 = Cyanotis nudiflora.
floribunda, H. B. & K. Nov. Gen. et Sp. i. 260 = Tradescantia cumanensis.
forcandia, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 255 = Cyanotis foecunda.
formosa, R. Grah. in Edinb. N. Phil. Journ. (July-Sept. 1830) 386 = nudiflora.
Forskalii, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 173 = Kotschyi.
Forskâlaei, *Vahl, Enum.* ii. 172.—Geront. trop.
Gambiae, *C. B. Clarke, in DC. Monog. Phan.* iii. 146. —Afr. trop.
gelatinosa, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 177.—Ind. or.
genienlata, *Hem. Prod. Fl. Ind. Occ.* 4.—Ins. S. Doming.
Gerrardi, *C. B. Clarke, in DC. Monog. Phan.* iii. 153. —Afr. austr.
gigantea, Vahl, Enum. ii. 177 = Anellema giganteum.
glabra, Baker, in Saund. Refug. Bot. ii. t. 166 = flaccida.
glabra, *C. B. Clarke, in DC. Monog. Phan.* iii. 168.—Ind. or.
glabra, G. F. W. Mey. Prim. Fl. Esseq. 22 = nudiflora.
gracilis, Hook. Bot. Mag. t. 3047 = flaccidula.
gracilis, Ruiz & Pav. Fl. Per. i. 44. t. 72. f. a = nudiflora.
graminea, Poir. Encyc. Suppl. ii. 324 = Anellema gramineum.
graminifolia, *H. B. & K. Nov. Gen. et Sp.* i. 258.—Mexic.
graminifolia, Pav. ex C. B. Clarke, in DC. Monog. Phan. iii. 151 = dianthifolia.
grandiflora, *Tenore, Cat. Hort. Bot. Nap.* (1845) 22.—Hab. ?
guianensis, Klotzsch, in Schomb. Faun. et Fl. Guian. 1064 = Phaeosphaerion persicariaefolium.
Hamiltonii, Spreng. Syst. v. 186 = Floscopa scandens.
hamiltis, *Wright, in Sauv. Fl. Cub.* 157.—Cuba.
Haroocmskii, *Hort. Pillnitz. ex Tenore, Cat. Ort. Bot. Nap.* (1845) 100.—Hab. ?
Haskarlii, *C. B. Clarke, Commel. et Cyrt. Beng.* t. 8. —Ind. or.
herbacea, Roxb. Hort. Beng. [5]; Fl. Ind. i. 175 = Anellema herbaceum.

COMMELINA :—
Hendelotii, *C. B. Clarke, in DC. Monog. Phan.* iii. 154.—Afr. trop.
hexandra, Aubl. Pl. Gui. i. 35. t. 12 = Dichorisandra Aubletiana.
hirsuta, R. Br. in Salt, Abyss. App. 62 = benghalensis.
hirsuta, *C. B. Clarke, in DC. Monog. Phan.* iii. 163.—Ind. or.
hirsuta, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 344 = Cyanotis hirsuta.
hirsuta, Hort. Berol. (1841) ex Kunth, Enu n. Pl. iv. 660 = elliptica.
hirsuta, Willd. in Spreng. & Link, Jahrb. i. III. 78 = virginica.
hirtella, *Vahl, Enum.* ii. 166.—Am. bor.
hispida, Ham. ex Spreng. Syst. iv. Cur. Post. 25 = Floscopa scandens.
hispida, *Ruiz & Pav. Fl. Per.* i. 43. t. 73. f. a.—Peruv.
Hookeri, A. Dietr. Sp. Pl. ii. 404 = Anellema giganteum.
Hookeriana, Spreng. Syst. v. 186 = Anellema giganteum.
huillensis, *Welw. ex C. B. Clarke, in DC. Monog. Phan.* iii. 178.—Afr. trop.
humilis, Klotzsch, ex C. B. Clarke, l. c. 154 = C. elliptica.
ignorata, *Kunth, Enum. Pl.* iv. 60.—Am. bor.
imberbis, Ehrenb. ex Hassk. in Schweinf. Beitr. Fl. Aethiop. 309 = latifolia.
intermedia, Schlecht. in Linnaea, xxiv. (1851) 655 = coelestis.
involucrosa, *A. Rich. Tent. Fl. Abyss.* ii. 342.—Abyss.
Jamesoni, *C. B. Clarke, in DC. Monog. Phan.* iii. 155. —Ecuador.
japonica, Hort. Berol. (1841) ex Kunth, Enum. iv. 659 = coelestis.
japonica, Thunb. in Trans. Linn. Soc. ii. (1794) 332 = Anellema japonicum.
karooica, *C. B. Clarke, in DC. Monog. Phan.* iii. 166. —Afr. austr.
Karwinskiana, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 660 = seq.
Karwinskii, Mart. ex Kunth, l. c. = elliptica.
Kirkii, *C. B. Clarke, in DC. Monog. Phan.* iii. 167.—Afr. trop.
Koenigii, Steud. Nom. ed. II. i. 401 = Anellema herbaceum, Koenigii.
kotschyi, Hassk. in Schweinf. Beitr. Fl. Aethiop. 207. —Aegypt.
Kretchmara, Kunth, Enum. Pl. iv. 40 = africana.
Kurzii, *C. B. Clarke, in Journ. Linn. Soc.* xi. (1871) 444.—Ind. or.
lanceolata, Benth. Fl. Austral. vii. 84 = agrostophylla.
lanceolata, *R. Br. Prod.* 269.—Austral.
lanuginosa, Heyne, ex Wall. Cat. n. 5221 = Anellema lanuginosum.
latifolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 340. —Abyss.
latifolia, Hoobst. ex C. B. Clarke, in DC. Monog. Phan. iii. 160, 169 = benghalensis, uncata.
laxa, Poir. Encyc. Suppl. ii. 324 = Anellema laxum.
leiandra, *C. B. Clarke, in DC. Monog. Phan.* iii. 150.—Mexic.
leiocarpa, Benth. in Bot. Voy. Sulph. 176 = Athyrocarpus leiocarpus.
linearifolia, Kunth, Enum. Pl. iv. 48 = subulata.
linearis, Benth. Fl. Hartw. 27 = dianthifolia.
linoides, Blume, Enum. Pl. Jav. i. 8 = Anellema herbaceum.
Livingstoni, *C. B. Clarke, in DC. Monog. Phan.* iii. 190. —Afr. austr.
Loddigesii, Steud. Nom. ed. II. i. 401 = tuberosa.
longicapsa, *C. B. Clarke, in DC. Monog. Phan.* iii. 176. —Afr. occ.
longicaulis, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = coelestis.
longicaulis, Jacq. Coll. iii. 284 = nudiflora.
longifolia, *Lam. Tabl. Encyc.* i. 139.—Java.
longifolia, Micha. Fl. Am. Bor. i. 58 = hirtella.
longifolia, Spreng. Syst. iv. Cur. Post. 25 = Anellema giganteum.
longifolia, Thw. Enum. Pl. Zeyl. 322 = Kurzii.
Lourcirii, *Kunth, Enum. Pl.* iv. 60.—China.
ludens, Miq. in Journ. Bot. Néerl. i. (1861) 88 = communis.
lunata, Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 138 = ensifolia.
lutea, Moench, Meth. 287 = africana.
macrophylla, Poir. Encyc. Suppl. ii. 324 = Pollia macrophylla.

COMMELINA :—

maculata, Edgew. in Trans. Linn. Soc. xx. (1851) 89 = obliqua.
madagascarica, *C. B. Clarke, in DC. Monog. Phan.* iii. 174.—Madag.
Maonli, *C. B. Clarke, l. c.* 167.—Afr. trop.
Martiana, Seub. in Mart. Fl. Bras. iii. 1. 265.·t. 87. f. 1 = virginica.
mascarenica, *C. B. Clarke, in DC. Monog. Phan.* iii. 174.—Ins. Maurit.
medica, Lour. Fl. Cochinch. 40 = Aneilema medicum.
mexicana, E. Mey. in Presl, Rel. Haenk. i. 137 = Tradescantia elongata.
micrantha, Vahl, Enum. ii. 178 = Aneilema micranthum.
minuta, Blume, Cat. Gew. Buitenz. 34 = Aneilema nudiflorum.
mollis, Jacq. Coll. iii. 235 = benghalensis.
moluccana, Roxb. Hort. Beng. (81); Fl. Ind. i. 172 = Cyanotis uniflora.
monadelpha, Blume, Enum. Pl. Jav. i. 4 = Aneilema monadelphum.
montana, Stend. Nom. ed. II. i. 401 = Aneilema montanum.
monticola, *Sonb. in Mart. Fl. Bras.* iii. 1. 264.—Bras.
multicaulis, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 164 = albescens.
multiflora, Martens & Gal. in Bull. Acad. Brux. ix. (1842) ii. 374 = Callisia Martensiana.
nana, Roxb. Hort. Beng. 5; Fl. Ind. i. 173 = Aneilema spiratum.
nervosa, Burm. f. Fl. Ind. 18 = benghalensis.
nervosa, Ruiz & Pav. Fl. Per. i. 44 = fasciculata.
nigritana, Baker, in Trans. Linn. Soc. xxix. (1875) 163 = nubulata.
nigritana, Benth. in Hook. Niger Fl. 541 = umbellata.
nilagirica, Steud. ex C. B. Clarke, in DC. Monog. Phan. iii. 163, 171 = hirsuta, clavata.
Nimmoniana, *J. Grah. Cat. Pl. Bombay*, 224.—Ind. or.
nitens, *Hort. Pillnitz. ex Tenore, Cat. Ort. Bot. Nap.* (1845) 100.—Hab. ?
nudicaulis, Burm. f. Fl. Ind. 17. t. 8. f. 1 = Aneilema nudiflorum.
nudiflora, Linn. Mant. ii. 177 = Aneilema nudiflorum.
nudiflora, *Linn. Sp. Pl.* 4.—Reg. trop.
obliqua, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 45.—Ind. or. ; Malaya.
obliqua, Vahl, Enum. ii. 173 = erecta.
oblongifolia, Hassk. Commel. Ind. 27 = auriculata.
obtusifolia, Vahl, Enum. ii. 168 = nudiflora.
ochrosta, Schan. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 447 = nudiflora.
oligotricha, *Mig. Fl. Ind. Bat.* iii. 532.—Malaya.
orchioides, Booth, ex Lindl. Bot. Reg. (1838) Misc. 58 = elliptica.
orchioides, *Roth, ex Steud. Nom.* ed. II. i. 402.—Mexic.
ovata, Stend. l. c. = Aneilema ovatum.
ovato-oblonga, Roem. & Schult. Mant. i. Add. t. 876 = Aneilema ovato-oblongum.
pacifica, Vahl, Enum. ii. 168 = nudiflora.
palasta, *Hassk. in Mig. Fl. Jungh.* 189.—Ind. or. ; Malaya.
Palisoti, A. Dietr. Sp. Pl. ii. 414 = Aneilema ovato-oblongum.
pallida, *Willd. Hort. Berol.* ii. 87. t. 87.—Mexic.
pallida, Schlecht. in Linnaea, xxvi. (1853-55) 454 = Athyrocarpus leiocarpus.
paludosa, Blume, Enum. Pl. Jav. i. 2 = obliqua.
paludosa, Stend. Nom. ed. II. i. 402 = persicariaefolia.
paniculata, *Hill, Veg. Syst.* xxiii. 4.—Ind. or.
paniculata, Nort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = coelestis.
paniculata, Soland. ex C. B. Clarke, in DC. Monog. Phan. iii. 228 = Aneilema laxum.
paniculata, Vahl, Enum. ii. 179 = Aneilema Forskalei.
papillonacea, Burm. f. Fl. Ind. 17 = Cyanotis papilionacea.
parviflora, Link, Enum. Hort. Berol. i. 61 = gramuifolia.
parviflora, Reichb. Fl. Exot. ii. 17. t. 146 = tuberosa.
paucinervis, Spreng. & Link, Jahrb. i. 241. 74 = salicifolia.
persicariaefolia, [Delile, in] Red. Lil. t. 472 = Athyrocarpus persicariaefolius.
persicariaefolia, *Wight, in Wall. Cat.* n. 8984.—Burma.
Peterii, *Hassk. in Flora*, xlvi. (1863) 255.—Afr. trop.

COMMELINA :—

phalangoides, *Cranis, Inst.* i. 459.—Hab. ?
pilosa, Pers. Syn. i. 55 = nudiflora.
pilosa, Stend. Nom. ed. II. i. 402 = Aneilema ochraceum.
pilosula, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = nudiflora.
platyphylla, *Klotzsch, in Schomb. Faun. et Fl. Guiana*, 897.—Guiana.
Poiteana, Seub. in Mart. Fl. Bras. iii. 1. 265. t. 87. f. 1 = erecta.
polygama, Blanco, Fl. Filip. ed. Naves (Ic. sine num.) = benghalensis.
polycloda, Welw. ex C. B. Clarke, in DC. Monog. Phan. iii. 165 = africana.
polygama, Roth, in Roem. & Ust. Mag. iv. x. (1790) 14 = communis.
polygastha, Wight, Ic. t. 2066 = obliqua.
procurrens, Schlecht. in Linnaea, xxv. (1852) 183 = benghalensis.
prostrata, H. B. & K. Nov. Gen. et Sp. i. 259 = nudiflora.
prostrata, Poepp. ex Kunth, Enum. Pl. iv. 38 = nudiflora.
prostrata, Regel, Gartenfl. (1868) 289. t. 592. f. 1 = benghalensis.
protensa, Steud. Nom. ed. II. i. 402 = Aneilema protensum.
pubernla, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv 659 ; et in Linnaea, xxiv. (1851) 197 = dianthifolia.
pubescens, Klotzsch, ex C. B. Clarke, in DC. Monog. Phan. iii. 159 = acuminata.
pubinervis, *Hassk. in Flora*, xxv. (1842) ii. *Beibl.* 4.—Java.
pumila, Royle, ex C. B. Clarke, in DC. Monog. Phan. iii. 207 = Aneilema spiratum.
pyrrho-blephara, *Hassk. in Schweinf. Beitr. Fl. Aethiop.* 309.—Abyssin.
quitensis, *Benth. Pl. Hartw.* 258.—Ecuador.
radicans, Spreng. Syst. iv. Cur. Post. 25 = Aneilema nudiflorum.
radicifora, R. Br. ex C. B. Clarke, in DC. Monog. Phan. iii. 159 = benghalensis.
Rahnesquet, *Kunth. Enum. Pl.* iv. 40.—Am. bor.
rajmahalensis, C. B. Clarke, in Journ. Linn. Soc. xi. (1871) 444 = attenuata.
rara, *Roxb. ex Steud. Nom.* ed. II. i. 402.—Ind. or.
reniformis, Steud. l. c. = Pollia subumbellata.
repens, Roxb. & Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 172 = attenuata.
rhizocarpa, Afzel. ex C. B. Clarke, l. c. 159 = benghalensis.
robusta, *Kunth, Enum. Pl.* iv. 59.—Bras.
rorigera, *Hsch. ex Tenore, Cat. Ort. Bot. Nap.* (1845) 100.—Hab. ?
rosea, *Schlecht. Ind. Sem. Hort. Hal.* (1838) 7.—Mexic.
rubens, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 639 = gramuifolia.
rubens, [Delile, in] Red. Lil. vii. t. 367 = pallida.
rufipes, Seub. in Mart. Fl. Bras. iii. 1. 265 = Phaeospherion persicariaefolium.
rugulosa, C. B. Clarke, in Journ. Linn. Soc. xi. (1871) 446 = suffruticosa.
Ramphii, *Kostel. Allg. Med.-Pharm. Fl.* i. 126.—As. trop.
Sabatieri, *C. B. Clarke, in DC. Monog. Phan.* iii. 146. —Afr. trop.
sagittifolia, Hassk. in Schweinf. Beitr. Fl. Aethiop. 306 = latifolia.
salicifolia, Boj. Hort. Maurit. 260 = communis.
salicifolia, *Roxb. Hort. Beng.* 5 ; *Fl. Ind.* i. 172.—Ind. or.
salicifolia, Thw. Enum. Pl. Zeyl. 321 = clavata.
Salziana, Stend. Nom. ed. II. i. 402 = pyrrho-blepharis.
scabra, Blume, Enum. Pl. Jav. i. 4 = Aneilema protensum.
scabra, *Benth. Pl. Hartw.* 26.—Mexic.
scabrata, Seub. in Mart. Fl. Bras. iii. 1. 266 = Phaeospherion persicariaefolium.
scandens, *Welw. ex C. B. Clarke, in DC. Monog. Phan.* iii. 146.—Afr. trop.
scapifera, Roxb. Hort. Beng. [8]; Fl. Ind. i. 175 = Aneilema scapiflorum.
scapiflora, Hort. Berol. (1833) ex Kunth, Enum. Pl. iv. 46 = elliptica.
Schimperiana, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 164 = albescens.
Schomburgkiana, Klotzsch, in Schomb. Faun. et Fl. Guiana, 1064 = undiflora.

COMMELINA :—

Schouberti, Zucc. ex Tenore, Cat. Hort. Bot. Nap. (1845) 22 = hispida.
Schweinfurthii, *C. B. Clarke, in DC. Monog. Phan.* iii. 158.—Afr. trop.
secundiflora, Blume, Enum. Pl. Jav. i. 3 = Pollia secundiflora.
Sellowiana, Kunth, Enum. Pl. iv. 44 = nudiflora.
Sellowii, Schlecht, Ind. Sem. Hort. Hal. (1850) 7 = nudiflora.
semi-ovata, Buch.-Ham. in Wall. Cat. n. 8985 = obliqua, suffruticosa.
senegalensis, Tenore, Cat. Ort. Bot. Nap. (1845) 22 = benghalensis.
serrulata, *Vahl, Eclog. Am.* ii. 4.—Guiana.
setosa, Wight, in Wall. Cat. n. 8981 = undulata.
sikkimensis, *C. B. Clarke, Commel. & Cyrt. Beng.* t. 6 —Reg. Himal.
silicuiosa, Poir. Encyc. Suppl. ii. 324 = Aneilema silicuiosum.
simplex, Vahl, Enum. ii. 177 = Aneilema sinicum.
Simsoni, C. B. Clarke, in Journ. Linn. Soc. xi. (1871) 446 = suffruticosa.
singularis, *Vell. Fl. Flum.* 31 ; i. t. 76.—Bras.
sinica, Roem. & Schult. Mant. i. App. t. 876 = Aneilema sinicum.
speciosa, Thunb. Prod. Pl. Cap. 58 = Cyanotis nodiflora.
spectabilis, *C. B. Clarke, in DC. Monog. Phan.* iii. 175.—Afr. trop.
spicata, Stend. Nom. ed. II. i. 402 = Aneilema Loureiri.
spirata, Koen. ex Steud. l. c. = Aneilema herbaceum.
spirata, Linn. Mant. ii. 176 = Aneilema spiratum.
striata, Edgew. in Trans. Linn. Soc. xx. (1851) 89 = obliqua.
striata, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 340 = subulata.
striata, Wall. Cat. n. 8981 = albescens, Kurzii, emifolia *stricta*, Desf. Cat. Hort. Par. ed. III. 383 = graminifolia.
subamplectens, Hasak. Commel. Ind. 27 = africana.
subaxillaris, Hochst. in Kunth, Enum. Pl. iv. 658 = subulata.
subulata, *Roth, Nov. Pl. Sp.* 28.—Geront. trop.
suffruticosa, *Blume, Enum. Pl. Jav.* i. 3.—Java.
sulcata, Benth. in Hook. Niger Fl. 541 = Vogelii.
sulcata, Hoffmgg. in Spreng. & Link, Jahrb. i. iii. 74 = virginica.
tennis, Roth, Nov. Pl. Sp. 22 = Aneilema acuminatum.
tinctoria, Nees, in Linnaea, xix. (1847) 699 = dianthifolia.
triquetra, Stend. Nom. ed. II. i. 402 = Aneilema triquetrum.
truncata, Willd. Enum. Hort. Berol. Suppl. 4 = virginica.
tuberosa, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1824 (1826) 110 = virginica.
tuberosa, Burm. ex Steud. Nom. ed. II. i. 402 = salicifolia.
tuberosa, [DC. in] Red. Lil. ii. t. 108 = coelestis.
tuberosa, Forsk. Fl. Aegypt. Arab. 12 = Aneilema Forskalei.
tuberosa, *Linn. Sp. Pl.* 41.—Mexic.
tuberosa, Lour. Fl. Cochinch. 40 = Aneilema Loureiri.
turbinata, Vahl, Enum. ii. 171 = benghalensis.
umbellata, *Schum. & Thonn. Beskr. Guin. Pl.* 21.—Afr. trop.
umbrosa, Vahl, Enum. ii. 179 = Aneilema ovato-oblongum.
uncta, *C. B. Clarke, in DC. Monog. Phan.* iii. 169. — Abyssin.
undulata, Benth. Fl. Austral. vii. 88 = Chamtscania.
undulata, Lodd. Bot. Cab. t. 1858 = tuberosa.
undulata, *R. Br. Prod.* 270.—Austral.
vaginata, Heyne, ex Steud. Nom. ed. II. i. 402 = Aneilema cucullatum.
vaginata, Linn. Mant. 177 = Aneilema vaginatum.
vaginata, Sonn. ex C. B. Clarke, in DC. Monog. Phan. iii. 166 = emifolia.
variabilis, Schlecht. Ind. Sem. Hort. Hal. (1836) 7 = elliptica.
ventricosa, Seub. in Mart. Fl. Bras. iii. 1. 264 = montioides.
villosicaulis, Soland. ex C. B. Clarke, in DC. Monog. Phan. iii. 159 = benghalensis.
virginica, Vent. f. Prod. 6 = nudiflora.
virginica, *Linn. Sp. Pl.* ed. II. 61.—Am. bor.
Vogelii, *C. B. Clarke, in DC. Monog. Phan.* iii. 189.—Afr. trop.

COMMELINA:—
vulgaris, [DC. in] Red. Lil. tv. t. 306 = communis.
Wallichiana, Steud. Nom. ed. II. i. 409 = Anellema divergens.
Welwitschii, *C. B. Clarke, in DC. Monog. Phan.* iii. 178.—Afr. trop.
Wernerana, Hassk. in Schweinf. Beitr. Fl. Aethiop. 206 = undiflora.
Willdenowii, Kunth, Enum. Pl. iv. 37 = communis.
zambesica, *C. B. Clarke, in DC. Monog. Phan.* iii. 161.—Afr. trop.
Zanonia, Linn. Sp. Pl. 41 = Campelia Zanonia.
Zebrina, Hort. ex C. B. Clarke, in DC. Monog. Phan. iii. 818 = Zebrina pendula.
zeylanica, *Ralfsenberg, in Yust, Jahresb.* iv. (1876) 406.—Zeylan.

COMMELYNA, Hoffmgg. ex Endl. Gen. 125 (1837) = Commelina, Linn.

COMMERÇONIA, F. Muell. Census, 16 (1882) = Commersonia, Forst. (Sterculi.).

COMMERSONA, Sonner. Voy. t. 89 (1782) = Barringtonia, Forst. (Myrtac.).
speciosa, Salisb. Prod. 356 = B. Butonica.

COMMERSONIA, Comm. ex DC. Prod. ii. 10, in syn. (1825) = Polycardia, Juss. (Lecythidear.).

COMMERSONIA, Forst. Char. Gen. 43. t. 22 (1776). *STERCULIACEAE*, Benth. & Hook. f. i. 226.
aspera, G. Don, Gen. Syst. i. 523 = Buettneria aspera.
cinerea, Steud. in Lehm. Pl. Preiss. i. 288 = Rulingia grandiflora.
crassophylla, F. Muell. Fragm. ix. 59 = Rulingia crassophylla.
crispa, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) i. 501.—Austral.
cuneata, F. Muell. Census, 16 = Rulingia cuneata.
cygnorum, Steud. in Lehm. Pl. Preiss. i. 287 = Rulingia malvaefolia.
dasyphylla, Andr. Bot. Rep. t. 608 = Rulingia pannosa.
dasyphylla, Hort. ex Steud. Nom. ed. II. i. 402 = Rulingia pannosa.
densiflora, F. Muell. Fragm. x. 21 = Rulingia densiflora.
diphylla, Andr. Bot. Rep. t. 608 = Rulingia pannosa.
echinata, Blume, Bijdr. 86 = platyphylla.
echinata, *Forst. Char. Gen.* 43. t. 22.—Malaya.
Fraseri, *Y. Gay, in Mém. Mus. Par.* x. (1823) 215. t. 15.—Austral.
Fraseri, Sieber, ex Steets, in Lehm. Pl. Preiss. ii. 359 = Rulingia pannosa.
Gaudichaudii, *Y. Gay, in Mém. Mus. Par.* x. (1823) 213. t. 14.—Austral.
herbacea, G. Don, Gen. Syst. i. 523 = Buettneria herbacea.
hermannaefolia, *Y. Gay, in H. B. & K. Nov. Gen. et Sp.* v. 311, in nota.—Hab. ?
hermanniaefolia, F. Muell. Census, 16 = Rulingia hermanniaefolia.
Jacksiana, G. Don, Gen. Syst. i. 523 = Buettneria Jacksiana.
juvenis, G. Don, l. c. = platyphylla.
Kempeana, F. Muell. Fragm. xi. 118 = Rulingia Kempeana.
laxophylla, F. Muell. l. c. x. 29 = Rulingia laxophylla.
Leichardtii, *Benth. Fl. Austr.* i. 246.—Austral.
magnifera, F. Muell. Fragm. viii. 233 = Rulingia magnifera.
melanopetala, *F. Muell. l. c.* x. 21.—Austral.
microphylla, *Benth. Fl. Australas.* i. 244.—Austral.
parviflora, F. Muell. Census, 16 = Rulingia parviflora.
pilosa, G. Don, Gen. Syst. ii. 594 = Buettneria pilosa.
platycalyx, F. Muell. Census, 16 = Rulingia platycalyx.
platyphylla, *Andr. Bot. Rep.* t. 608 adn. et t. 518.—Malaya.
Preissii, Steud. in Lehm. Pl. Preiss. i. 288 = Rulingia corylifolia.
pulchella, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) i. 502.—Austral
retusifolia, F. Muell. Census, 16 = Rulingia retusifolia.
regnans, F. Muell. l. c. = Rulingia regnans.
salicifolia, F. Muell. l. c. = Rulingia salicifolia.
COMMIA, Lour. Fl. Cochin. 605 (1790) = Excoecaria, Linn. (Euphorb.).
cochinchinensis, Lour. l. c. 606 = E. Agallocha.

COMMIANTHUS, Benth. in Hook. Journ. Bot. iii. (1841) 328 = Retiniphyllum, Humb. & Bonpl. (Rubiac.).
concolor, Spruce, ex Benth. in Hook. Kew Journ. v. (1853) 235 = R. concolor.
discolor, Spruce, ex Benth. l. c. 234 = R. discolor.
pilosus, Spruce, ex Benth. l. c. 233 = R. pilosum.
Schomburgkii, Benth. in Hook. Journ. Bot. iii. (1841) 328 = R. Schomburgkii.
speciosus, Spruce, ex Benth. in Hook. Kew Journ. v. (1853) 234 = R. speciosum.

COMMIDENDRON, Lem. in Orb. Dict. iv. 147 (1849) = seq.

COMMIDENDRUM, Barch. ex DC. in Guill. Archiv. Bot. ii. (1833) 384. *COMPOSITAE*, Benth. & Hook. ii. 277.
Burchellii, *Benth. & Hook. f. ex Hemsl. Bot. Chall Voy.* i. ii. 71.—Ins. S. Helena.
gummiferum, DC. Prod. v. 344 = robustum.
robustum, *DC. l. c.*—Ins. S. Helena.
rotundifolium, DC. l. c. = Psiadia rotundifolia.
rugosum, *DC. l. c.* 345.—Ins. S. Helena.
sparsum, *DC. l. c.* 344.—Ins. S. Helena.

COMMILOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 110 = Psorodon, Vog. (Legumin.).
abruptum, Moric. Fl. Rar. Nouv. Am. 98. t. 62 = P. abruptus.
polygonifolium, Benth. in Ann. Wien. Mus. ii. (1838) 111 = P. polygalaefolius.
pubescens, Benth. l. c. = P. pubescens.

COMMIPHORA, Jacq. Hort. Schoenb. ii. 66 (1797). *BURSERACEAE*.
BALSAMEA, Gled. in Berl. Ges. Naturf. Fr. Schr. iii. (1782) 197.
BALSAMODENDRON, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 348.
BALSAMOPHLOEOS, O. Berg, in Bot. Zeit. xx. (1862) 168.
BALSAMUS, Stackh. De Liban. 11 (1814).
HEMPRICHIA, Ehrenb. in Linnaea, iv. (1829) 396.
HEUDELOTIA, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 160. t. 39 (1832).
HITTERIA, Klotzsch, in Peters, Reise Mossamb. Bot. 89 (1861).
NIOTOUTT, Adans. Voy. 162 (1759).
PROTIONOPSIS, Blume, Mus. Bot. Lugd. Bat. i. 229 (1850).
PROTIUM, Wight & Arn. Prod. 176 (1834).
abyssinica, *Engl. in DC. Monog. Phan.* iv. 10.—Abyss.
africanum, *Endl. l. c.* 14.—Abyss.
Agallocha, *Engl. l. c.* 11.—Ind. or.
angolensis, *Engl. l. c.* 34.—Afr. trop.
Berryi, *Engl. l. c.* 17.—Ind. or.
Botviniana, *Engl. l. c.* 21.—Afr. trop.
capensis, *Engl. l. c.* 18.—Afr. austr.
caudata, *Engl. l. c.* 27.—Ind. or.
edulis, *Engl. l. c.* 22.—Afr. trop.
erythraea, *Engl. l. c.* 30.—Afr. trop.
grandifolia, *Engl. l. c.* 25.—Madag.
Harveyi, *Engl. l. c.*—Afr. austr.
Hildebrandtii, *Engl. l. c.* 15.—Afr. trop.
Kirkii, *Engl. l. c.* 19.—Ind. or.
longebractesta, *Engl. l. c.*—Afr. trop.
madagascariensis, *Jacq. Hort. Schoenb.* ii. 66.—Madag.
Marchandii, *Engl. in DC. Monog. Phan.* iv. 26.—Madag.
mollis, *Engl. l. c.* 28.—Afr. trop.
mossambicensis, *Engl. l. c.* 26.—Afr. trop.
Mukul, *Engl. l. c.* 13.—Ind. or.
Myrrha, *Engl. l. c.* 10.—Arab.
Olivieri, *Engl. l. c.* 24.—Afr. trop.
Opobalsamum, *Engl. l. c.* 15.—Arab.
orbicularis, *Engl. l. c.* 18.—Madag.
parvifolia, *Engl. l. c.* 536.—Ins. Socotra.
pedunculata, *Engl. l. c.* 23.—Afrca.
Perrieliana, *Engl. l. c.* 29.—Madag.
pilosa, *Engl. l. c.* 12.—Afr. trop.
planifrons, *Engl. l. c.* 536.—Ins. Socotra
pubescens, *Engl. l. c.* 31.—Ind. or.
Rahmanni, *Engl. l. c.* 15.—Afr. austr.
Schimperi, *Engl. l. c.* 12.—Abyssin.

COMMIPHORA:—
serrata, *Engl. in DC. Monog. Phan.* iv. 24.—Afr. trop.
socotrana, *Engl. l. c.* 536.—Ins. Socotra.
Stocksiana, *Engl. l. c.* 17.—Beluchist.
tetramera, *Engl. l. c.* 37.—Madag.
Welwitschii, *Engl. l. c.* 22.—Afr. trop.
zanzibarica, *Engl. l. c.* 28.—Afr. trop.

COMMIRHOEA, Miers, in Trans. Linn. Soc. xxi. (1855) 252 = Chrysochlamys, Poepp. † (Guttif.).
mexicana, Miers, l. c. 258.—Mex. ?

COMOCLADIA, P. Br. Hist. Jamaic 124 (1756). *ANACARDIACEAE*, Benth. & Hook. f. i. 419.
DODONAEA, Plum. ex Adans. Fam. ii. 342 (1763).
acuminata, *Moç. & Sesse, ex DC. Prod.* ii. 65.—Mexic.
anguiosa, Willd. Sp. Pl. i. 1483 = Ilicifolia.
Brasiliastrum, *Pohr. Encyc. Suppl.* ii. 226.—Am. austr.
dentata, *Jacq. Enum. Pl. Carib.* 12.—Ind. occ.
Ehrenbergii, *Engl. in Bot. Jahrb.* i. (1881) 420.—Ins. S. Doming.
glabra, *Sprng. Syst.* i. 1176.—Cuba.
ilicifolia, *Sw. Prod. Veg. Ind. Occ.* 17.—Ind. occ.
integrifolia, *Jacq. Enum. Pl. Carib.* 12.—Am. trop.
internecilis, *Wright, ex Engl. in DC. Monog. Phan.* iv. 366.—Cuba.
loxensis, *H. B. & K. Nov. Gen. et Sp.* vii. 16.—N. Granat.
mollissima, *H. B. & K. l. c.* 16. t. 607.—Mexic.
pinnatifida, Linn. Syst. ed. X. 861 = integrifolia.
platyphylla, *A. Rich. ex Griseb. Cat. Pl. Cub.* 68.—Cuba.
propinqua, *H. B & K. Nov. Gen. et Sp.* vii. 16.—Cuba.
pubescens, *Engl. in Bot. Jahrb.* i. (1881) 420.—Jamaic.
serrata, Blanco, Fl. Filip. ed. I. 80 = Salacia oblonga.
Tapaculo, H. B. & K. Nov. Gen. et Sp. vii. 16 = Tapiria guianensis.
tomentosa, Willd. ex Schult. Mant. i. 350 = mollissima.
tricuspidata, Lam. in Mem. Acad. Sc. Par. 1784 (1787) 347 = ilicifolia.

COMOLIA, DC. Prod. iii. 114 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 741.
HOSTMANNIA, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 25.
LEIOSTEGIA, Benth. in Hook. Journ. Bot. ii. (1840) 294.
PACHYLOMA, DC. Prod. iii. 120 (1828).
TETRAMERIS, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 120. t. 4.
TRICENTRUM, DC. Prod. iii. 123 (1828).
amazonica, *Cogn. in Mart. Fl. Bras.* xiv. III. 422.—Bras.
Aubletii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 87.—Guiana.
berbertiolia, *DC. Prod.* iii. 115.—Bras.
denudata, *Naud, in Ann. Sc. Nat.* Ser. III. xiii. (1850) 25.—Bras.
birtella, *Naud. l. c.* 26 = microphylla.
lanceolata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 87.—Bras.
latifolia, Cogn. in Mart. Fl. Bras. xiv. III. 433 = Aubletii.
leptophylla, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1850) 27.—Guiana.
lythrarioides, *Naud. l. c.*—Guiana.
microphylla, *Benth. in Hook. Journ. Bot.* ii. (1840) 290.—Guiana.
neglecta, *Cogn. in Mart. Fl. Bras.* xiv. III. 423.—Guiana.
nummularioides, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1850) 27.—Bras.
ovalifolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 87. t. 2 = denudata.
purpurea, Miq. in Linnaea, xviii. (1844) 617 = veronicaefolia.
Sertularia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 87. t. 2. f. 7.—Bras.
sessilis, *Triana, l. c.* f. 27 c.—Bras.
stenodon, *Triana, l. c.* 87.—Bras.
surinamensis, Miq. Stirp. Surin. 55 = lythrarioides.
tetraptera, *Cogn. in Mart. Fl. Bras.* xiv. III. 426.—Bras.

COMOLIA:—
verniciose, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 87. t. 3. f. 37 g.—Bras.
veronicæfolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 295.—Guiana.
veronicifolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 86 = purpurea.
villosa, *Triana, l. c.* 87.—Guiana.
violacea, *Triana, l. c.*—Bras.

COMOMYRSINE, Hook. f. in Benth. & Hook. f. Gen. ii. 648 (1876). *MYRSINEÆ*, Benth. & Hook. f. l. c.
Schlimii, *Hook. f. l. c.* 644.—Am. austr.
simplex, *Hook. f. l. c.*—Am. austr.
Sprucei, *Hook. f. l. c.*—Am. austr.

COMOSPERMA, Poir. in Dict. Sc. Nat. x. (1818) 128 = Comesperma, Labill. (Polygal.).

COMOSTEMUM, Nees, in Linnæa, ix. (1834) 288 = Androtrichum, Brongn. (Cyperac.).
globuliferum, Nees, in Mart. Fl. Bras. ii. t. 5 = A. polycephalum.
impolitum, Nees, in Hook. Journ. Bot. ii. (1840) 297 = Cyperus impolitus.
laetum, Nees, l. c. = Cyperus laetus, *Presl.*
montevidense, Nees, in Linnæa, ix. (1834) 288 = A. polycephalum.
prolixum, Nees, in Hook. Journ. Bot. ii. (1840) 297 = Cyperus prolixus.
Schottii, Nees, in Linnæa, ix. (1834) 288 = Cyperus prolixus.

COMPARETTIA, Poepp. & Endl. Nov. Gen. et Sp. i. 42. t. 78 (1835). *ORCHIDEÆ*, Benth. & Hook. f. iii. 586.
PLECTROPHORA, H. C. Focke, in Tijdschr. Natuurk. Wetensch. i. (1848) 212.
coccinea, *Lindl. Bot. Reg.* xxiv. (1838) t. 68.—Bras.
cryptocera, *Morr. in Belg. Hortic.* ii. (1852) 310.—Hab. ?
falcata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 42. t. 78.—Peruv.
macroplectron, *Reichb. f. & Triana, in Gard. Chron.* (1876) ii. 524; (1876) i. 293.—N. Granat.
rosea, Lindl. Bot. Reg. (1840) Misc. 78 = falcata.
saccata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 42. t. 72.—Peruv.
speciosa, *Reichb. f. in Gard. Chron.* (1878) ii. 524.—Ecuador.

COMPERIA, C. Koch, in Linnæa, xxii. (1849) 287 = Orchis, Linn.
taurica, C. Koch, l. c. 288 = O. Comperiana.

COMPHOROPSIS, Meq. in DC. Prod. xiii. ii. 900, in syn. (1849) = CAMPHOROPSIS, Meq. = Panophytum, Less. (Chenopod.).

COMPHRENA, Aubl. Pl. Gui. i. 280 (1775), sphalm. = Gomphrena, Linn. (Amarant.).

COMPSANTHUS, Spreng. Syst. iv. Cur. Post. 187 (1827) = Trigyrtis, Wall. (Liliac.).
maculatus, Spreng. l. c. = T. pilosa.

COMPSOA, D. Don, Prod. Fl. Nep. 50 (1825) = Trigyrtis, Wall. (1826) (Liliac.).
maculata, D. Don, l. c. 51 = T. pilosa.

COMPTONIA, Banks, ex Gaertn. Fruct. ii. 58. t. 90 (1791) = Myrica, Linn.
asplenifolia, Banks, l. c. = M. asplenifolia.
Catesbæi, Mirb. in Duham. Arb. ed. nov. ii. t. 11 = M. asplenifolia.

COMUS, Salisb. Gen. Pl. Fragm. 24 (1866) = Muscari, Tourn. (Liliac.).
carnosus, Salisb. l. c. 25 = M. comosum.

CONAMI, Aubl. Pl. Gui. ii. 926. t. 354 (1775) = Phyllanthus, Linn. (Euphorb.).
brasiliensis, Aubl. l. c. 927 = P. Conami.

CONANDRON, Sieb. & Zucc. in Abh. Akad. Wiss. Munich. iii. iii. (1844) 739. t. 3. *GESNERACEÆ*, Benth. & Hook. f. ii. 1094.
ramondioides, *Sieb. & Zucc. l. c.*—Japon.

CONANTHERA, Ruiz & Pav. Fl. Per. iii. 68. t. 301 (1802). *HÆMODORACEÆ*, Benth. & Hook. f. iii. 679.
CUMMINGIA, D. Don, in Sweet, Brit. Flow. Gard. 2nd. ii. t. 207 (1828).
albiflora, Cham. & Schlecht. in Linnæa, vi. (1831) 80 = Echeandia terniflora.
bifolia, *Ruiz & Pav. Fl. Per.* iii. 68. t. 301.—Chili.
bifolia, Sims, in Bot. Mag. t. 2496 = Simsii.
campanulata, Lindl. in Trans. Hort. Soc. vi. (1826) 258 = Simsii.
Echeandia, Pers. Syn. i. 370 = Echeandia terniflora.
Forsteri, *Spreng. Syst.* ii. 91.—N. Caled.
Simsii, *Sweet, Hort. Brit.* ed. I. 428.—Chili.

CONANTHODIUM, A. Gray, in Hook. Kew Journ. iv. (1852) 272 = Hellohrysum, Vaill. (Compos.).
Drummondii, A. Gray, l. c. = H. argyroglottis.

CONANTHUS, S. Wats. Bot. King's Exped. 256 (1871). *HYDROPHYLLACEÆ*, Benth. & Hook. f. ii. 896.
arcuiides, *S. Wats. l. c.*—Am. bor. occ.

CONCHEVERIA, Aubl. Pl. Gui. ii. 928. t. 353 (1775). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 316.
africana, *Muell. Arg. in Flora*, xxxiii. (1864–65) 530.—Afr. trop.
cordata, A. Juss. Tent. Euph. 48. t. 18 = Alchornea cordata.
guianensis, *Aubl. Pl. Gui.* ii. 924. t. 353.—Guiana.
Hostmani, *Benth. in Hook. Kew Journ.* vi. (1854) 332.—Guiana.
javanensis, Blume, Bijdr. 614 = Alchornea rugosa.
latifolia, Zipp. ex Span. in Linnæa, xv. (1841) 349 = Alchornea rugosa.
latifolia, *Benth. in Hook. Kew Journ.* vi. (1854) 332.—Bras.
macrophylla, Klotzsch, ex Benth. l. c. 333 = Alchornea cordata.
Martiana, *Baill. in Adansonia*, v. (1864–65) 221.—Bras.
magnifolia, Muell. Arg. in Linnæa, xxxiv. (1865–66) 167 = Alchornea Martiana.
ovata, Rich. ex A. Juss. Tent. Euph. 48 = guyanensis.
rhytidocarpa, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 273.—Peruv.
terminalis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 167.—Bras.
tomentosa, Span. l. c. xv. (1841) 349 = Cladogynos orientalis.
trigonocarpa, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 272.—Bras.

CONCEVEIBUM, Rich. ex A. Juss. Tent. Euph. 42 (1824) = prace.

CONCHIDIUM, Griff. Notul. iii. 321. t. 310 (1851) = Bria, Lindl. (Orchid.).
pusillum, Griff. l. c. = E. pusilla.
ninicum, Lindl. ex Benth. in Hook. Kew Journ. vii. (1855) 94 = E. stolon.

CONCHIUM, Sm. in Trans. Linn. Soc. iv. (1798) 215 = Hakea, Schrad. (Proteac.).
aciculare, Sm. ex Vent. Jard. Malm. t. 111 = H. sclenlaria.
ceratophyllum, Sm. in Trans. Linn. Soc. ix. (1808) 124 = H. ceratophylla.
clavatum, Willd. Enum. Hort. Berol. 141 = H. clavata.
compressum, Sm. in Trans. Linn. Soc. ix. (1808) 121 = H. scienlaria.
cornicalatum, Sm. ex Willd. Enum. Hort. Berol. 141 = H. pugioniformis.
cornutum, Gaertn. f. Fruct. iii. 216. t. 219 = H. gibbosa.
dactyloides, Vent. Jard. Malm. t. 110 = H. dactyloides.
drupaceum, Gaertn. f. Fruct. iii. 217. t. 219 = H. leucopteta.
ellipticum, Sm. in Trans. Linn. Soc. ix. (1808) 122 = H. ellipticu.

CONCHIUM:—
epiglottis, Willd. Enum. Hort. Berol. 141 = H. epiglottis.
gibbosum, Sm. in Trans. Linn. Soc. ix. (1808) 119 = H. gibbosa.
longifolium, Sm. l. c. 121 = H. pugioniformis.
mucronatum, Cels, ex Spreng. Syst. i. 680 = H. varia.
myrsinum, Sm. ex Willd. Enum. Hort. Berol. 141 = H. dactyloides.
oleifolium, Sm. in Trans. Linn. Soc. ix. (1808) 124 = H. oleifolia.
pectinatum, Willd. Enum. Hort. Berol. 141 = H. gibbosa.
pugioniforme, Sm. in Trans. Linn. Soc. ix. (1808) 122 = H. pugioniformis.
pyriforme, Willd. Enum. Hort. Berol. 141 = Xylomelum pyriforme.
ruscifolium, Willd. l. c. = H. ruscifolia.
salicifolium, Gaertn. f. Fruct. iii. 217 = H. saligna.
saligneum, Sm. in Trans. Linn. Soc. ix. (1808) 124 = H. saligna.
sphaeroideum, Sm. l. c. 120 = H. gibbosa.
terretifolium, Gaertn. f. Fruct. iii. 217. t. 219 = E. epiglottis.
trifurcatum, Sm. in Trans. Linn. Soc. ix. (1808) 122 = H. trifurcata.

CONCHOCARPUS, Mikan, Delect. fasc. i. t. 2 (1820) = Casparia, Humb. (Rutac.).
cuneifolius, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 102 = Cusp. cuneifolia.
macrophyllus, Mikan, Delect. fasc. i. t. 2 = Cusp. macrophylla.

CONCHOCHILUS, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 146 = Appendicula, Blume, (Orchid.).
distichus, Hassk. l. c. = A. pendula.
opposiflorus, Hassk. l. c. 147 = A. cristata.

CONCHOPHYLLUM, Blume, Bijdr. 1060 (1826) = Dischidia, R. Br. (Asclepiad.).
imbricatum, Blume, l. c. 1061 = D. Collyris.

CONCILIUM, Rafin. Fl. Tellur. ii. 78 (1836) = Lightfootia, L'Hérit. (Campan.).
pedunculare, Rafin. l. c. = L. subulata.

CONDAEA, Stend. Nom. ed. II. i. 403 (1840) = CONDEA, Adans. = Hyptis, Jacq. (Labiat.).

CONDALIA, Cav. in Anal. Cienc. Nat. i. (1799) 89. t. 4. *RHAMNEÆ*, Benth. & Hook. f. i. 376.
buxifolia, *Reiss. in Mart. Fl. Bras.* xi. i. 89.—Bras.
coriacea, Reichb. in Flora, xiii. (1830) i. 130 = Rhamnus integrifolius.
ferrea, Griseb. Fl. Brit. W. Ind. 100 = Scutia ferrea.
infectoria, *Reiss. in Mart. Fl. Bras.* xi. i. 90.—Bras.
linearis, *A. Gray, Bot. U. St. Expl. Exped.* i. 275.—Patagon.
maytenoides, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 14.—Chili.
mexicana, *Schlecht. in Linnæa*, xv. (1841) 471.—Mexic.
microphylla, *Cav. in Anal. Cienc. Nat.* i. (1799) 40.—Chili.
oborata, *Hook. Ic. Pl.* t. 287.—Texas.
paradoxa, Spreng. Syst. i. 835 = Colletia cruciata.
spathulata, *A. Gray, Pl. Wright.* i. 82.—Am. bor. occ.
spinosa, *Spreng. Syst. iv. Cur. Post.* 106.—Bras.

CONDALIA, Ruiz & Pav. Prod. 11. t. 2 (1794) = Cococcypselum, P. Br. (Rubiac.).
lanceolata, Ruiz & Pav. Fl. Per. i. 54 = Cocc. lanceolatum.
obovata, Ruiz & Pav. l. c. = Cocc. obovatum.
repens, Ruiz & Pav. l. c. = Cocc. umbellatum.
ovatifolia, Ruiz & Pav. l. c. = Cocc. ovatifolia.

CONDAMINEA, DC. Prod. iv. 402 (1830). *RUBIACEÆ*, Benth. & Hook. f. ii. 43.
corymbosa, *DC. l. c.*—Am. austr.
glabrata, *DC. l. c.*—Peruv.
macrophylla, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 20. —Bras.
microcarpa, *DC. Prod.* iv. 402.—Peruv.
tinctoria, *DC. l. c.*—Venezuela.
utilis, *Goudot, in Comptes Rendus*, xviii. (1844) 999.—N. Granat.
venosa, *DC. Prod.* iv. 402.—Peruv.

CONDEA, Adans. Fam. ii. 504 (1763) = **Hyptis**, Jacq. (Labiat.).

CONDYLICARPUS, Steud. Nom. ed. II. ii. 491 (1841) err. typ. = CONDYLOCARPUS, Hoffm. = **Tordylium**, Tourn. (Umbellif.).

OONDYLOCARPON, Desf. in Mém. Mus. Par. viii. (1822) 119. t. 11. *APOCYNACEAE*, Benth. & Hook. f. ii. 701.
HOSTINIANIA, Miq. in Nat. Verh. Wet. Haarl. vii. 107 (1850).
MAYCOCKIA, A. DC. Prod. viii. 324 (1844).
brasiliense, Mart. ex Muell. Arg. in Mart. Fl. Bras vi. t. 64 = Rauwolfiae.
brevistipulatum, *Muell. Arg. l. c.* 65.—Bras.
ciliatum, *Muell. Arg. l. c.* 66.—Bras.
gracile, *Miers, Apocyn. S. Am.* 33.—Am. austr.
guyanense, *Desf. in Mém. Mus. Par.* viii. (1822) 120. t. 8.—Guiana.
intermedium, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 65.—Bras.
isthmicum, *A. DC. Prod.* viii. 361.—Bras.
laxum, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 66.—Bras.
myrtifolium, *Muell. Arg. l. c.* 65, in obs.—Guiana.
obtusiusculum, *Muell. Arg. l. c.*—Bras.
pubiflorum, *Muell. Arg. l. c.* 67.—Bras.
Rauwolfiae, *Muell. Arg. l. c.* 64.—Bras.

CONDYLOCARPUS, Hoffm. Gen Umb. ed. II. 202 (1816) = **Tordylium**, Tourn. (Umbellif.).
apulus, Hoffm. l. c. 208 = T. apulum.
humilis, Koch, in Nov. Act. Nat. Cur. xii. t. (1824) 87 = T. apulum.
officinalis, Koch, l. c. = T. officinale.

CONDYLOCARPUS, Salisb ex Lamb. Pinus, ed. 8vo, 120, sub t. 64 (1825) = **Sequoia**, Endl. (Conif.).
sempervirens, Salisb. ex Carr. Conif. ed. I. 162 = S. sempervirens.

CONFORATA, Cassalp. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 91 = **Achillea**, Linn. (Compos.).
Ageratum, Fourr. l. c. = A. Ageratum.

CONGDONIA, Muell. Arg. in Flora, lix. (1876) 437.
RUBIACEAE.
caerulea, *Muell. Arg. l. c.*—Bras.

CONGEA, Roxb. Fl. Corom. iii. 90. t. 293 (1819).
VERBENACEAE, Benth. & Hook. f. ii. 1159.
CALOCHLAMYS, Presl, Bot. Bemerk. 143 (1844).
aurea, Wall. Cat. n. 1788 = Sphenodesma pentandra.
barbata, Wall. l. c. n. 1788 = Sphenodesma barbata.
ferruginea, Wall. l. c. n. 1787 = Sphenodesma unguiculata.
involucrata, Wall. l. c. n. 1740 = Symphorema polyandrum.
Jacksiana, Wall. l. c. n. 1786 = Sphenodesma pentandra, trifora.
paniculata, Wall. l. c. n. 1739 = Symphorema involucratum, Sphenodesma paniculata.
pentandra, Wall. l. c. n. 1784 = Sphenodesma pentandra.
tomentosa, *Roxb. Fl. Corom.* iii. 90.—Burma.
unguiculata, Wall. Cat. n. 1786 = Sphenodesma unguiculata.
veinina, *Wight, Ic.* t. 1479/2.—Ind. or.
ventia, *Griff. Notul. iv.* 174.—Burma.
villosa, *Wight, Ic.* t. 1479 B.—Burma.

CONGHAS, Wall. Cat. n. 3080; ex Hiern, in Hook. f. Fl. Intr. Ind. I. 681 (1875) = **Schlatterera**, Willd. (Sapindac.).
scylensuela, Wall. l. c. = S. trijuga.

CONIANDRA, Schrad. in Eckl. & Zeyh. Enum. 275 (1836) = **Kedrostis**, Medic. (Cucurb.).
africana, Sond. in Harv. & Sond. Fl. Cap. ii. 483 = K. obtusa.
cordifolia, Fenzl, ex Nand. in Ann. Sc. Nat. Ser. IV. xvi. (1862) 180 = Cardiocarpus Fenzlii.
digitata, Sond. in Harv. & Sond. Fl. Cap. ii. 483 = K. digitata.
discoidea, Schrad. in Eckl. & Zeyh. Enum. 276 = K. discoidea.
glauca, Schrad. l. c. 275 = K. grossulariaefolia.

CONIANDRA:—
grossulariaefolia, Arn. in Hook. Journ. Bot. iii. (1841) 273 = Kedrostis grossulariaefolia.
mollis, Sond. in Harv. & Sond. Fl. Cap. ii. 485 = K. mollis.
pinnatisecta, Schrad. in Eckl. & Zeyh. Enum. 276 = K. africana.
punctulata, Sond. in Harv. & Sond. Fl. Cap. ii. 484 = K. punctulata.
Thunbergi, Sond. l. c. = K. nana.
Zeyheri, Schrad. in Eckl. & Zeyh. Enum. 275 = K. Zeyheri.

CONIOGETON, Blume, Bijdr. 1156 (1826) = **Buchanania**, Roxb. (Anacard.).
arborescens, Blume, l. c. = B. florida.

CONIOSELINUM, Fisch. ex Hoffm. Gen. Umb. ed. I. p. 180 et xxviii. (1814); ed. II. 185 (1816) = **Ligusticum**, Linn. (Umbellif.).
altaicum, Rupr. in Beitr. Pfl. Russ. Reich. xi. 22.—Sibir.
canadense, Torr. & Gray, Fl. N. Am. i. 619 = Selinum canadense.
cenolophioides, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 736 = Selinum cenolophioides.
Czernaevia, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 53 = Pencedanum latifolium.
Fischeri, Wimm. & Grab. Fl. Siles. (1827) i. 366 = Selinum Benthami.
gayoides, Less. in Linnaea, ix. (1834) 178.—Sibir.
Gmelini, Stend. Nom. ed. II. i. 408 = Ligusticum Gmelini.
humile, Turcz. ex Ledeb. Fl. Ross. ii. 818 = Peucedanum salsum.
ivanense, Fisch. Ind. Sem. Hort. Petrop. (1834) ex Ledeb. l. c. 290 = Selinum Benthami.
kamtschaticum, Rupr. in Beitr. Pfl. Russ. Reich. xi. 22.—Kamtsch.
latifolium, Rupr. Sert. Tianach. 48.—Am. centr.
longifolium, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 736.—Sibir.
neglectum, Fisch. ex Steud. Nom. ed. II. i. 408, 404 = Selinum Benthami.
tataricum, Hoffm. Gen. Umb. ed. II. 185 = Selinum Benthami.
uniwittatum, Turcz. ex Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 368 = Selinum Benthami.

CONIOTHELE, DC. Prod. v. 531 (1836) = **Blennosperma**, Less. (Compos.).
californica, DC. l. c. = B. californicum.

CONIROSTRUM, Dulac, Fl. Hautes-Pyr. 195 (1867) = **Bromidium**, Tourn. (Crucif.).
obtusangulum, Dulac, l. c. = B. Erucastrum.

CONISA, Desf. ex Steud. Nom. ed. II. i. 404 (1840) = **Conyza**, Linn. (Compos.).

CONIUM, Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 893.
africanum, Hort. ex DC. Prod. iv. 108 = Carum copticum.
africanum, Jacq. Hort. Vind. ii. 91. t. 194 = Capnophyllum Jacquini.
africanum, Linn. Sp. Pl. 248 = Capnophyllum africanum.
Arracacia, Hook. Exot. Bot. t. 152 = Arracacia xanthorrhiza.
chaerophylloides, Eckl. & Zeyh. Enum. 255 = Sceli chaerophylloides.
Cicuta, Neck. Delic. Gallo-Belg. i. 149 = maculatum.
crosticum, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 305 = maculatum.
dichotomum, Desf. Fl. Atlant. i. 246. t. 66 = Capnophyllum dichotomum.
divaricatum, Boiss. & Orph. Diagn. Ser. II. v. 103 = maculatum.
filifolium, Vahl, Symb. Bot. iii. 49 = Oenanthe filiformis.
Jacquini, D. Dietr. Syn. Pl. ii. 968 = Capnophyllum Jacquini.
maculatum, *Linn. Sp. Pl.* 243.—Europ.; Orient.
maculosum, Pall. Reis. Statth. Russ. Reich. i. 478 = maculatum.
Monbret, *Hort. ex Opiz, Natural.* x. (1826) 260, nomen.—Europ.

CONIUM:—
moschatum, H. B. & K. Nov. Gen. et Sp. v. 14. t. 420 = Arracacia moschata.
nepalense, D. Dietr. Syn. Pl. ii. 988 = Vicatia conifolia.
nodosum, Fisch. ex Steud. Nom. ed. I. 215 = maculatum.
rigens, Linn. Mant. iii. 252, 512 = Bunium rigens.
rigidum, Mert. ex DC. Prod. iv. 140 (= *Sclerocciadium punctilus*).—Marocco.
Royeni, Linn. Sp. Pl. 243 = Caucalis daucoides.
rugosum, Thunb. Prod. Pl. Cap. 50 = Capnophyllum Jacquini.
sibiricum, Hort. ex Steud. Nom. ed. II. i. 404 = maculatum.
strictum, Trait. Arch. Gew. i. t. 21 = maculatum.
suffruticosum, Berg. Descr. Pl. Cap. 77 = Selinum suffruticosum.
tenuifolium, Mill. Gard. Dict. ed. VIII. n. 2 = maculatum.
tenuifolium, Vahl, Symb. Bot. iii. 49 = Oenanthe filiformis.
verrucosum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 324 = Pimpinella simensis.

CONIZA, Neck. Delic. Gallo-Belg. ii. 848 (1768) = **Conyza**, Linn. (Compos.).

CONNAROPSIS, Planch. ex Hook. f. in Trans. Linn. Soc. xxiii. (1860) 166. *GERANIACEAE*, Benth. & Hook. f. i. 377.
diversifolia, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) 11. 69 = Griffithii.
glauca, *Hook. f. in Linn. Soc. Trans.* xxiii. (1860) 166.—Borneo.
Griffithii, *Planch. l. c. et ex Hook. f. Fl. Brit. Ind.* i. 440.—Malaya.
monophylla, *Planch. il. cc.*—Malacca.
philippica, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 98.—Ins. Philipp.

CONNARUS, Linn. Diss. Dass. 25 (1747); in Amoen. Acad. i. 406 (1749). *CONNARACEAE*, Benth. & Hook. f. i. 482.
ANISOSTEMON, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 152.
CAMICIDIA, Vell. Fl. Flum. 184 (1825); iv. t. 139 (1827).
CASTANOLA, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 503.
ERYTHROSTIGMA, Hassk. in Flora, xxv. (1842) ii. Beibl. 45.
KALAWARI, Adans. Fam. ii. 344 (1763).
MATHERANCIA, Stend. Nom. ed. II. i. 404, in syn. (1840).
OMPHALOBIUM, Gaertn. Fruct. i. 217. t. 46 (1788).
TALI, Adans. Fam. ii. 312 (1763).
TAPOMANA, Adans. Fam. ii. 344 (1763).
africanus, *Lam. Encyc.* ii. 95.—Afr. trop.
africanus, G. F. W. Mey. Prim. Fl. Essequ. 228 = guianensis.
Afzelii, R. Br. ex Planch. in Linnaea, xxiii. (1850) 416 = Roures santaloides.
asiaticus, Willd. Sp. Pl. iii. 692 = monocarpus.
Beyrichii, *Planch. in Linnaea*, xxiii. (1850) 480.—Bras.
Blanchetii, *Planch. l. c.* 481.—Bras.
Championi, *Thw. Enum. Pl. Zeyl.* 80.—Zeylan.
conchocarpus, *F. Muell. Fragm.* v. 108.—Austral.
confertiflorus, *Baker, in Mart. Fl. Bras.* xiv. ii. 198.—Guiana.
crassifolius, Benth. ex Baker, l. c. 186 = favosus.
cuneifolius, *Baker, l. c.* 194.—Bras.
cymosus, *Planch. in Linnaea*, xxiii. (1850) 430.—Bras.
decumbens, *Thumb. in Roem. Arch.* i. t. (1796) 1.—Afr. austr.
deustus, *Planch. in Linnaea*, xxiii. (1850) 485.—Bras.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* i. 539.—Sumatra.
Duparquetianus, *Baill. Adansonia*, vii. (1866–67) 236.—Afr. trop.
erianthus, *Benth. ex Baker, in Mart. Fl. Bras.* xiv. ii. 191.—Bras.
falcatus, *Blume, Mus. Bot. Lugd. Bat.* i. 266.—Malaya.
fasciculatus, Planch. in Linnaea, xxiii. (1850) 492 = erianthus.
favosus, *Planch. l. c.* 484.—Bras.

CONNARUS:—
fecundus, *Baker, in Mart. Fl. Bras.* xiv. ii. 185.—Bras.
ferrugineus, *Yach, in Malay. Misc.* ii. (1822) n. vii. 37. —Malaya.
floribundus, Planch. in Linnæa, xxiii. (1850) 435 = Perrottetii.
floribundus, *Schum. & Thonn. Beskr. Guin. Pl.* 299.— Afr. trop.
floribundus, Wall. ex Hook. f. Fl. Brit. Ind. ii. 52 = semidecandras.
floridanus, Thonn. ex DC. Prod. ii. 86 = floribundus.
fluminensis, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 529 = Bernardinia fluminensis.
fortes, Blanco, Fl. Filip. ed. I. 253 = Murraya exotica.
foliolosus, Jack, in Wall. Cat. n. 8529 = Cnestis ramiflora.
fulgens, Wall. Cat. n. 8524 = Roures fulgens.
fulvus, *Planch. in Linnæa*, xxiii. (1850) 434.—Bras.
furfuraceus, *Blume, Mus. Bot. Lugd. Bat.* i. 268.— Sumatra.
Gaudichaudii, *Planch. in Linnæa*, xxiii. (1850) 429.— Ins. Moluce.
gibbosus, *Wall. Cat.* n. 8541, *ex Planch. l. c.* 436.— Burma; Malaya.
glaber, DC. Prod. ii. 85 = Roures glabra.
grandifolius, *Planch. in Linnæa*, xxiii. (1850) 429.— Ins. S. Doming.
grandis, *Yach, in Malay. Misc.* ii. (1822) n. vii. 40. —Burma; Malaya.
Griffithii, *Hook. f. Fl. Brit. Ind.* ii. 53.—Burma.
Griffonianus, *Baill. Adansonia*, vii. (1866–67) 285.— Afr. trop.
guianensis, *Lamb. ex DC. Prod.* ii. 85.—Am. trop.
hæmorrhæus, *Kurst. Fl. Columb.* ii. 73. t. 127.— N. Granat.
Hasseltii, *Blume, Mus. Bot. Lugd. Bat.* i. 266.—Java.
ignaus, Wall. Cat. n. 8528 = Cnestis ramiflora.
incomptus, *Planch. in Linnæa*, xxiii. (1850) 433.— Guiana.
Jachienus, Wall. Cat. n. 8552 = Cupania Jackiana.
javanicus, *Blume, Bijdr.* 1166.—Java.
juglandifolius, Hook. & Arn. Bot. Beech. Voy. 179 = Rhus succedanea.
kavaiensis, *H. Mann, in Proc. Am. Acad.* vii. (1867) 163.—Ins. Sandvic.
latifolius, *Wall. Cat.* n. 8537.—Burma.
laurifolius, *Baker, in Mart. Fl. Bras.* xiv. ii. 186.— Bras.
luckiae, *Hassk. in Hort. & De Vriese, Tijdschr.* x. (1843) 144.—Malaya.
luckius, *Yach, in Malay. Misc.* ii. (1822) n. vii. 41. —Malaya.
macrophyllus, Poepp. & Endl Nov. Gen. et Sp. iii. 76 = Roures macrophylla.
Maingayi, *Hook. f. Fl. Brit. Ind.* ii. 53.—Malaya.
Mann, *Baker, in Oliver, Fl. Trop. Afr.* i. 455.—Afr. trop.
marginatus, *Planch. in Linnæa*, xxiii. (1850) 429.— Bras.
microphyllus, Hook. & Arn. Bot. Beech. Voy. 179 = Roures microphylla.
mimosoides, Jack, in Malay. Misc. ii. (1822) n. vii. 44 = Roures concolor.
mimosoides, *Vahl, Symb. Bot.* iii. 87 (sp. dubia).—Ins. Nicobar.
mimosoides, Wall. Cat. n. 8526 = Roures Wallichiana, similis.
moluccanus, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 266 = Gaudichaudii.
monadelphus, *Wall. Cat.* n. 8547 B.—Ind. or.
monocarpus, *Linn. Sp. Pl.* 675.—Ind. or.
monocarpus, Wight & Arn. Prod. 143 = Roures santaloides.
monophyllus, Wall. Cat. n. 8551 = Ellipanthus tomentosulus.
mustalifa, *Blume, Mus. Bot. Lugd. Bat.* i. 269.— Malaya.
nemorosus, *Vahl, ex Schum. & Thonn. Beskr. Guin. Pl.* 299.—Afr. trop. ?
neurocalyx, *Planch. in Linnæa*, xxiii. (1850) 345.— Ins. Philipp.
nicicus, *Hassk. in Flora*, xxvii. (1844) ii. 616.—Java.
nitidus, *Poxb. Hort. Beng.* 49; *Wall. Cat.* n. 8540 C. —Ind. or.
nodosus, *Baker, in Mart. Fl. Bras.* xiv. ii. 190.—Bras.
obliquus, *Presl, in Walp. Ann.* iii. 844.—Ins. Luzon.
oblongifolius, Mart. Herb. Fl. Bras. n. 1267 = Roures glabra.

CONNARUS:—
obtusifolius, *Planch. in Linnæa*, xxiii. (1850) 428.— Ins. Luzon.
odoratus, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 172.—Borneo.
oligophyllus, *Planch. in Linnæa*, xxiii. (1850) 437.— Penang.
panamensis, *Griseb. in Bonplandia*, vi. (1858) 6.— Panama.
panamensis, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 277 = Turczaninowii.
paniculatus, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 139.— Ind. or.
Parisii, *Planch. in Linnæa*, xxiii. (1850) 432.— Guiana.
pentagynus, Lam. Encyc. ii. 95 = Agelæa Lamarckii.
pentandrus, Roxb. Hort. Beng. 49; Fl. Ind. iii. 140 = paniculatus.
Perrottetii, *Planch. in Linnæa*, xxiii. (1850) 432.— Guiana.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 375. —Ins. Fiji.
pinnatus, Hasselt, ex Blume, Mus. Bot. Lugd. Bat. i. 266 = Hasseltii.
pinnatus, Lam. Encyc. ii. 95 = monocarpus.
polyanthus, *Planch. in Linnæa*, xxiii. (1850) 428.— Ins. Luzon.
polyphyllus, *Miq. Fl. Ind. Bat. Suppl.* 239.— Sumatra.
pubescens, *Baker, in Oliver, Fl. Trop. Afr.* i. 455.— Afr. trop.
pubescens, DC. Prod. ii. 85 = Roures revoluta.
punctatus, *Planch. in Linnæa*, xxiii. (1850) 433.— Guiana.
pyrrhocarpus, *Miq. Fl. Ind. Bat. Suppl.* i. 530.— Sumatra.
reticulatus, *Griseb. Cat. Pl. Cub.* 84.—Cuba.
Ritchiei, *Hook. f. Fl. Brit. Ind.* ii. 51.—Ind. or.
Rolfei, *Vidal, Phan. Cuming. Philipp.* 32, 106.—Ins. Philipp.
Roxburghii, Hook. & Arn. Bot. Beech. Voy. 179 = Roures Milletii.
ruber, *Planch. in Linnæa*, xxiii. (1850) 436.—Am. austr.
rugosus, Wall. Cat. n. 8527 = Roures rugosa.
sanctalioides, Blanco, Fl. Filip. ed. II. 266 = Murraya exotica.
santaloides, Vahl, Sym. Bot. iii. 87 = Roures santaloides.
Schomburgkii, *Planch. in Linnæa*, xxiii. (1850) 431. —Guiana.
semidecandrus, *Yach, in Malay. Misc.* ii. (1822) n. vii. 39.—Burma; Malaya.
Smeathmanni, *Planch. in Linnæa*, xxiii. (1850) 426; *Baker, in Oliver, Fl. Trop. Afr.* i. 456.—Afr. trop.
Spanoghei, *Blume, Mus. Bot. Lugd. Bat.* i. 267.—Ins. Timor.
Spruceï, *Baker, in Mart. Fl. Bras.* xiv. ii. 187.—Bras.
stictophyllus, *Kurz, For. Fl. Brit. Burma*, i. 546.— Burma.
suberosus, *Planch. in Linnæa*, xxiii. (1850) 433.— Bras.
Thomsoni, *Baker, in Oliver, Fl. Trop. Afr.* i. 456.— Afr. trop.
Turczaninowii, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 364.—Panama.
turgidus, *Wall. Cat.* n. 8545, *ex Planch. in Linnæa*, xxiii. (1850) 436.—Ind. or.
umbifoliolatus, Thw. Enum. Pl. Zeyl. 86 = Ellipanthus Thwaitesii.
velutinosus, Wall. Cat. n. 8536 = Ellipanthus tomentosus.
venezuelanus, *Baill. Adansonia*, ix. (1868–70) 151.— Venezuela.
venosus, Bennthm. ex DC. Prod. ii. 85 = africanus.
villosus, *Yach, in Malay. Misc.* ii. (1822) n. vii. 38. —Sumatra.
villosus, Planch. in Linnæa, xxiii. (1850) 429 = grandis.
Wallichii, Planch. l. c. 426 = grandis.
Wightii, *Hook. f. Fl. Brit. Ind.* ii. 51.—Ind. or.

CONOBAEA, Bert. ex Steud. Nom. ed II. i. 404 (1840) = Mukhlenbeckia, Meisn. (Polygon.).
sagittifolia, Bert. ex Steud. l. c. = M. sagittifolia.

CONOCEPHALUS, Adsh. Pl. Gui. ii. 689. t. 256 (1775). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 951.
BLANCKIA, Neck. Elem. i. 247 (1790).
LEUCOPHYLLA, Nutt. in Journ. Acad. Philad. vii. (1834) 87.

CONOPEA.—
SCHISTOPHRAGMA, Benth. in Endl. Gen. 679 (1836).
SPHAEROTHECA, Cham. & Schlecht. in Linnæa, ii. (1827) 606.
alata, R. Grah. in Edinb. N. Phil. Journ. (1836) 3 = Leucocarpon alatum.
aquatica, *Aubl. Pl. Gui.* ii. 689. t. 256.—Guiana.
aquatica, Willd. ex Benzl. in Vet. Akad. Handl. Stockh. 1854 (1856) 141 = Herpestis lanigera.
borealis, Spreng. Neue Entdeck. iii. 96 = Gratiola virginiana.
indica, Spreng. Syst. ii. 771 = Limnophila polystachya.
intermedia, *A. Gray, in Torr. Bot. Mex. Bound.* 117. —Mexic.
multifida, *Benth. in DC. Prod.* x. 391.—Am. bor.
ovata, Spreng. ex Schrank, in Syll. Ratisb. ii. (1828) 69 = Stemodia parviflora !
pumila, Spreng. Nov. Prov. 13 = Stemodia parviflora.
punctata, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 143.—Bras.
pusilla, *Benth. & Hook. f. Gen.* ii. 951.—Mexic.
scoparioides, *Benth. in DC. Prod.* x. 391.—Mexic.
vandelioides, *Benth. l. c.*—Bras.
verticillaris, Hort. Berol. ex Spreng. Nov. Prov. 13 = Stemodia durantifolia.
viscosa, Schrank, in Syll. Ratisb. ii. (1828) 61 = Stemodia durantifolia.
viscosa, *Spreng. Neue Entdeck.* i. 263.—Bras.

CONOCALPIS, Boj. ex Decne. in DC. Prod. viii. 621 (1844) = Gymnema, R. Br. (Asclepiad.).
umbellata, Boj. ex Decne. l. c. = G. rufescens.

CONOCARPUS, Adans. Fam. ii. 384 (1763) = Leucadendron, R. Br. (Proteac.).

CONOCARPUS, Linn. Gen. ed. I. 376 (1737); Geertn. Fruct. ii. 470. t. 177 (1791). *COMBRETACEAE*, Benth. & Hook. f. i. 686.
RHIZARIA, Rafin. Sylva Tellur. 90 (1838).
RUDBECKIA, Adans. Fam. ii. 80 (1763).
acuminatus, Buch.-Ham. ex Wall. Cat. n. 4014, 4015 = Anogeissus latifolia.
acuminatus, Roxb. Hort. Beng. 84; Fl. Ind. ii. 443 = Anogeissus acuminata.
acutifolius, Willd. in Roem. & Schult. Syst. v. 574 = erectus.
erectus, *Linn. Sp. Pl.* 147; *Jacq. Select. Am.* 78. t. 52. —Ind. occ.
hirtus, Buch.-Ham. ex Wall. Cat. n. 4016 = Anogeissus acuminata.
lanceolatus, Heyne, ex Wall. l. c. n. 4014 = Anogeissus acuminata.
latifolius, Roxb. Hort. Beng. 84; Fl. Ind. ii. 443 = Anogeissus latifolia.
leiocarpus, DC. Prod. iii. 16 = Anogeissus leiocarpa.
mucronifera, Steud. Nom. ed. I. 218 = Rhus Rhadsellavel.
myrtifolius, Buch.-Ham. ex Wall. Cat. n. 4017 = Anogeissus pendula.
nitidicus, *Dehlle, ex Steud. Nom.* ed. II. i. 404.— Aegypt.
Novæ-Zelandiæ, *Cels, ex Tenore, Cat. Hort. Neap.* (1846) 98.—N. Zel.
parvifolius, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 269 = Anogeissus leiocarpa.
procumbens, Linn. Sp. Pl. 148 = erectus.
pubescens, *Schum. & Thonn. Beskr. Guin. Pl.* 115.— Afr. trop.
racemosus, Linn. Syst. ed. X. 930; Jacq. Select. Am. 80 = Laguncularia racemosa.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 269 = Anogeissus leiocarpa.
sericeus, Forst. ex G. Don, Gen. Syst. ii. 662 = erectus.
sulcius, Cassin, Ann. i. 256 = erectus.

CONOCEPHALUS, Blume, Bijdr. 489 (1825). *URTICACEAE*, Benth. & Hook. f. iii. 390.
acuminatus, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 91.—Ins. Philipp.
asperrimus, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvi. (1864) 36.—Java; Sumatra.
Blumei, *Gaudich. Bot. Voy. Bonite*, t. 96.—Malaya.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 210. —Borneo.
concolor, *Dalz. in Dalz. & Gibs. Bomb. Pl.* 220.— Ind. or.
ellipticus, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 91.—Penang.

CONOCEPHALUS:—
erectus, *Villar. in Bianco, Fl. Philipp.* ed. III. *Nov. App.* 908.—Ins. Philipp.
gratus, Miq. Fl. Jungh. I. 43 = suaveolens.
lanceolatus, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 68.—Ind. or.
micranthus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 210. —Borneo.
microphyllus, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 92.—Ins. Philipp.
nasciformis, Lindl. Bot. Reg. t. 1308 = suaveolens.
nivens, Wight, Ic. vi. 7. t. 1959 = Debregeasia velutina.
ovatus, Trec. in Ann. Sc. Nat. Sér. III. viii. (1847) 88 = suaveolens.
pubescens, Trec. l. c. 90 = suaveolens.
Roxburghii, Trec. l. c. 88 = suaveolens.
suaveolens, *Blume, Bijdr.* 483.—Ind. or.; Malaya.
subrinervius, *Miq. Fl. Ind. Bat. Suppl.* 617.—Malaya.

CONOCLINIUM, DC. Prod. v. 135 (1836) = Eupatorium, Tourn. (Compos.).
acuminatum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 361 = E. organense.
adenolepis, Sch. Bip. ex Baker, l. c. = E. Selloi.
affine, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 466 = E. betonicæforme.
album, Mart. Del. Sem. Hort. Monac. (1842) 4.—Mexic.
ballotæfolium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 360 = E. ballotæfolium.
betonicæforme, DC. Prod. v. 135 = E. betonicæforme.
betonicum, DC. Prod. v. 135 = E. betonicum.
crenæfolium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 358 = E. cerasifolium.
coelestinum, DC. Prod. v. 135 = E. coelestinum.
dichotomum, Chapm. in Coult. Bot. Gaz. iii. (1878) 5 = E. coelestinum.
dissectum, A. Gray, Pl. Wright. I. 88 = E. Greggii.
hirsutissimum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 311 = E. hirsutissimum.
humile, Benth. Pl. Hartw. 199.—N. Granat.
ianthinum, Mart. in Ann. Soc. Gand, v. (1849) 178 = E. ianthinum.
Lassauxii, Dur. Ind. Sem. Hort. Burdig. (1872) 15.—Uruguay.
maritimum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 362 = E. apiculatum.
microcephalum, Sch. Bip. ex Baker, l. c. 364 = E. Candollianum.
microphyllum, DC. Prod. v. 135 = E. microphyllum.
mikanioides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 306 = E. sordescens.
oligolepis, Kunze, Del. Sem. Hort. Lips. (1840) iv. 2 = E. oligolepis.
organense, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 361 = E. organense.
palustre, Sch. Bip. ex Baker, l. c. 363 = E. palustre.
platylepis, Sch. Bip. ex Baker, l. c. 356 = E. platylepis.
prasifolium, DC. in Deless. Ic. Sel. iv. t. 15 = E. nepetoides.
retinaculum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 306 = E. sordescens.
rigidum, Chapm. in Coult. Bot. Gaz. iii. (1878) 6 = E. heterocliniun.
rugosum, Torrs. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 188.—Ecuador.
scandens, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 487.—Bras.
semidentatum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 316 = E. semistriatum.
subglutinosum, A. Gray, in Proc. Am. Acad. v. (1861) 318; Sch. Bip. ex Baker, l. c. 362 = E. apiculatum.
subviscosum, Baker, l. c., spbalm. = E. apiculatum.
Viradus, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 313 = E. Virdus.

CONOHORIA, Aubl. Pl. Gui. i. 389. t. 96 (1775) = Alsodeia, Thou. (Violar.).
alternifolia, Spreng. Neue Entdeck. ii. 151 = A. integrifolia.
Aublet, D. Dietr. Syn. Pl. i. 371 = Casearia decandra.
cauliflora, Sw. Fl. Ind. Occ. iii. (1806) 1267 = A. guianensis.
CONOHORIA:—
Cuspa, H. B. & K. Nov. Gen. et Sp. vii. 242 = Alsodeia Cuspa.
dentata, D. Dietr. Syn. Pl. i. 381 = A. guianensis.
flavescens, Aubl. Pl. Gui. i. 389 = A. flavescens.
Lobelioba, St. Hil. in Mém. Mus. Par. xi. (1824) 494 = A. physiphora.
megapotamica, G. Don, Gen. Syst. i. 841 = A. megapotamica.
ovariensis, D. Dietr. Syn. Pl. i. 381 = A. ovariensis.
Pazsowa, DC. Prod. i. 312 = A. flavescens.
Perrini, D. Dietr. Syn. Pl. i. 382 = A. Perrini.
Regnellii, Miq. in Linnæa, xxii. (1849) 556.—Bras.
Riana, DC. Prod. i. 312 = A. prunifolia.
Rimosa, St. Hil. Pl. Rem. 321 = A. paniculata.
ulmifolia, H. B. & K. Nov. Gen. et Sp. v. 287. t. 491 = A. ulmifolia.

CONOMITRA, Fenzl, in Endl. Nov. Stirp. Dec. 85 (1839) = Glossonema, Decne. (Asclepiad.).
linearis, Fenzl, l. c. = G. lineare.

CONOMORPHA, A. DC. in Trans. Linn. Soc. xvii. (1834) 102. *MYRSINEÆ*, Benth. & Hook. f. ii. 644.
CONOSTYLUS, Pohl, ex A. DC. l. c. (1834).
buxoides, *Griseb. Cat. Pl. Cub.* 168.—Cuba.
guianensis, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 92 = Cybianthus guyanensis.
heteoclada, *Benth. ex Miq. in Mart. Fl. Bras.* x. 304. —Bras.
jacquinioides, *Griseb. Cat. Pl. Cub.* 163.—Cuba.
laxiflora, *DC. in Trans. Linn. Soc.* xvii. (1837) 103.—Bras.
macrophylla, *Mart. Herb. Fl. Bras.* n. 260.—Bras.
multipunctata, *Miq. Stirp. Surin. Sel.* 111. t. 54.—Guiana.
nemoralis, *Mart. & Miq. in Mart. Fl. Bras.* x. 304.—Bras.
oblongifolia, *A. DC. in Trans. Linn. Soc.* xvii. (1837) 103.—Bras.
peruviana, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 92.—Peruv.
reticulata, *Benth. ex Miq. in Mart. Fl. Bras.* x. 303.—Bras.
robusta, *Klotzsch, in Faun. & Fl. Gui.* 974.—Guiana.

CONOPHALLUS, Schott, Syn. Aroid. 84 (1856) = Amorphophallus, Blume (Aroid.).
Blumei, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 124.—Java.
bulbifer, Schott, Syn. Aroid. 84 = A. bulbifer.
commutatus, Schott, in Bonplandia, vii. (1859) 28 = A. commutatus.
giganteus, Schott, ex Miq. Fl. Ind. Bat. iii. 198 = A. giganteus.
Gigas, Miq. Ann. Mus. Bot. Lugd. Bat. i. 286 = A. Gigas.
Konjak, Schott, Syn. Aroid. 84 = A. Rivieri.
lyratus, Schott, l. c. 85 = A. lyratus.
Muelleri, Miq. Ann. Mus. Bot. Lugd. Bat. 286, in obs. = A. Mulleri.
planus, Miq. l. c. 286 = A. planus.
punctulatus, Schott, Prod. Aroid. 35 = A. punctulatus.
sativus, Schott, l. c. = A. sativus.
spectabilis, Miq. Fl. Ind. Bat. iii. 199 = A. spectabilis.
Titanum, Becc. in Bull. Soc. Tosc. Ortic. iii. (1878) 271, 291 = A. Titanum.
tuberculiger, Schott, in Bonplandia, vii. (1859) 28 = A. tuberculiger.

CONOPHARYNGIA, G. Don, Gen. Syst. iv. 94 (1837) = Fiumaria, Tourn. (Apocyn.).
longifolia, G. Don, l. c. 95 = F. longifolia.
retusa, G. Don, l. c. = F. retusa.

CONOPHOLIS, Wallr. Oroh. Diask. 78 (1825). *OROBANCHACEÆ*, Benth. & Hook. f. ii. 985.
americana, *Wallr. l. c.*—Am. bor.
americana, Endl. Iconogr. t. 81 = mexicana.
mexicana, *A. Gray, ex S. Watz. in Proc. Am. Acad.* xvii. (1882–83) 131.—N. Mexic.

CONOPHYTA, Scham. ex Hook. f. in Trans. Linn. Soc. xxii. (1856) 42 = Thouinigia, Vahl (Balsamph.).

CONOPODIUM, Koch, in Nov. Act. Nat. Cur. xii. (1824) 116. *UMBELLIFERÆ*, Benth. & Hook. f. ii. 896.
BULBOCASTANUM, Lag. Amen. Acad. Madr. 99 (1821).
BUTINIA, Boiss. Elench. 58 (1838).
ELAEOSTICTA, Fenzl, in Flora, xxvi. (1843) 458.
HETEROTAENIA, Boiss. Voy. Bot. Espagne, 267. t. 80 A (1840).
SCALIGERIA, DC. Prod. iv. 248 (1830).
SPHALLEROCARPUS, Bess. in DC. Prod. iv. 230 (1830).
alliodes, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 588.—Turkestan.
boetics, *Boiss. Voy. Bot. Espagne*, ii. 786.—Hispan.
Bourgaei, *Coss. Notes Crit.* 110.—Hispan.
brachycarpum, *Boiss. ex Lange, in Kjoeb. Vidensk. Meddel.* (1865) 44.—Hispan.
bunioides, *Boiss. & Reut. ex Boiss. Voy. Bot. Espagne*, ii. 786.—Hispan.
canadense, Koch, in Nov. Act. Nat. Cur. xii. (1824) 119 = Cryptotaenia canadensis.
capillifolium, *Boiss. Voy. Bot. Espagne*, ii. 786.—Lusitan.
Cyminum, *Benth. & Hook. f. Gen.* i. 896.—Dahuria; Volh.
denudatum, *Koch, in Nov. Act. Nat. Cur.* xii. (1824) 118.—Europ.
hirtulum, *Regel & Schmalh. in Pl. Nov. Fedsch.* 28.—Alt. centr.
Marianum, *Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 282.—Hispan.
mutabile, Miégev. in Bull. Soc. Bot. Fr. xxi. (1874) p. xxxii = denudatum.
pumilum, Nym. Consp. 308 = Bourgaei.
pyrenaicum, Miégev. in Bull. Soc. Bot. Fr. xxi. (1874) p. xxxii = denudatum.
ramosum, *Costa, Cat. Catal.* 105.—Hispan.
subcarneum, *Boiss. & Reut. ex Boiss. Voy. Bot. Espagne*, ii. 786.—Hispan.
subcarneum, Bourg. in Bull. Soc. Bot. Fr. ii. (1855) 648 = Bourgaei.

CONOPSIDIUM, Wallr. in Linnæa, xiv. (1840) 147 = Habenaria, Willd. (Orchid.).
platanthereum, Wallr. l. c. = H. bifolia.
sthenanthereum, Wallr. l. c. = H. bifolia.

CONORIA, Juss. Gen. 287 (1789) = Alsodeia, Thou. (Violar.).

CONOSAPIUM, Muell. Arg. in Linnæa, xxxii. (1863) 87 = Baptium, P. Br. (Euphorb.).
madagascariense, Muell. Arg. l. c.—Madag.

CONOSILENE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 244 = Silene, Linn. (Caryophyll.).
conica, Fourr. l. c. = S. conica.
conoidea, Fourr. l. c. = S. conoidea.

CONOSPIPHON, Poepp. & Endl. Nov. Gen. et Sp. iii. 27 (1842); Endl. Gen. Pl. Suppl. ii. 54 (1842) = Sphinctanthus, Benth. (Rubiac.).
aureus, Poepp. & Endl. l. c. = S. rupestris.
polycarpus, Karst. in Linnæa, xxx. (1859–60) 155.—N. Granat.
strictiflorus, Walp. Rep. ii. 520 = S. rupestris.

CONOSPERMUM, Sm. in Trans. Linn. Soc. iv. (1798) 213. *PROTEACEÆ*, Benth. & Hook. f. iii. 174.
acerosum, *Lindl. Swan River App.* 80.—Australasiacifolium, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.–July, 1836) 171.—Austral.
affine, *Room. & Schult. Syst.* iii. *Mant.* 274.—Austral.
amœnum, *Meisn. in Lehm. Pl. Preiss.* i. 522.—Austral.
brachyphyllum, *Lindl. Swan River App.* 81.—Austral.
bracteosum, *Meisn. in Lehm. Pl. Preiss* i. 518; ii. 248.—Austral.
Brownii, *Meisn. l. c.* ii. 248.—Austral.
canaliculatum, Meisn. l. c. 250 = Stoechadis.
capitatum, *R. Br. in Trans. Linn. Soc.* x. (1811) 155. —Austral.
corruiscens, F. Muell. Fragm. i. 157 = amœnum.
coeruleum, *R. Br. in Trans. Linn. Soc.* x. (1811) 154. —Austral.
commutatum, *Room. & Schult. Syst.* iii. *Mant.* 276.—Austral.

CONOSPERMUM, —
crassinervium, *Meisn. in Hook. Kew Journ.* iv. (1852) 184.—Austral.
Dallachyi, F. Muell. Ann. Rep. (1858) 8de Benth. Fl. Austral. v. 371 = Mitchellii.
debile, *Kipp. ex Meisn. in Hook. Kew Journ.* vii. (1855) 76.—Austral.
densiflorum, *Lindl. Swan River App.* 32.—Austral.
diffusum, *Benth Fl. Austral.* v. 367.—Austral.
distichum, Meisn. in Lehm. Pl. Preiss. i. 533 = floribundum.
distichum, *R. Br. in Trans. Linn. Soc.* x. (1811) 155. —Austral.
ellipticum, *Sm. in Rees, Cycl.* ix. n. 2.—Austral.
ephedroides, *Kipp. ex Meisn. in Hook. Kew Journ.* vii. (1855) 70.—Austral.
erectum, Hort. ex R. Grah. in Edinb. N. Phil. Journ. (Apr.-June 1838) 178 = ericifolium.
ericifolium, *Sm. in Rees, Cycl.* ix. n. 4.—Austral.
falcifolium, [Salisb. in] Knight, Prot. 95 = taxifolium.
filifolium, Meisn. in Lehm. Pl. Preiss. i. 533 = brachyphyllum.
flexuosum, *R. Br. Prod. Suppl.* 11.—Austral.
floribundum, *Benth. Fl. Austral.* v. 378.—Austral.
glumaceum, *Lindl. Swan River App.* 30.—Austral.
Huegelii, *R. Br. ex Endl. Nov. Stirp. Dec.* 58.— Austral.
imbricatum, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 46. —Austral.
incurvum, *Lindl. Swan River App.* 30.—Austral.
intricatum, Lindl. ex Meisn. in DC. Prod. xiv. 228 = Huegelii.
lanceolatum, *R. Br. Prod. Suppl.* 10.—Austral.
lanifolium, *Endl. in Ann. Wien. Mus.* ii. (1838) 308. —Austral.
lavaudulaefolium, *A. Cunn. ex Meisn. in DC. Prod.* xiv. 218.—Austral.
linifolium, *A. Cunn. l. c.* 220.—Austral.
longifolium, *Sm. Exot. Bot.* ii. 45. t. 92.—Austral.
lupulinum, Endl. ex Meisn. in Lehm. Pl. Preiss. ii. 249 = glumaceum.
marginatum, *Meisn. l. c.* 248.—Austral.
Mitchellii, *Meisn. in DC. Prod.* xiv. 220.—Austral.
nervosum, *Meisn. in Hook. Kew Journ.* vii. (1855) 71.—Austral.
patens, *Schlecht. in Linnaea*, xx. (1847) 587.—Austral.
petiolare, *R. Br. Prod. Suppl.* 11.—Austral.
polyophallum, *Meisn. in Lehm. Pl. Preiss.* ii. 249.— Austral.
procerum, F. Muell. Fragm. i. 157 = distichum.
propinquum, R. Br. Prod. Suppl. 10 = taxifolium.
repens, Sieber, ex Roem. & Schult. Syst. iii. Mant. 276 = tenuifolium.
reticulatum, Sm. in Rees, Cycl. ix. n. 5 = Synaphea petiolaris.
rigidum, [Salisb. in] Knight, Prot. 95 = ellipticum.
sonposum, *Benth. Fl. Austral.* v. 369.—Austral.
sclerophyllum, *Lindl. Swan River App.* 30.—Austral.
Smithii, Pers. Syn. i. 113 = longifolium.
sphacelatum, *Hook. in Mitch. Trop. Austral.* 349. —Austral.
spicatum, R. Br. Prod. Suppl. 10 = taxifolium.
Stoechadis, *Endl. in Ann. Wien. Mus.* ii. (1838) 308. —Austral.
taxifolium, Sieber, ex Roem. & Schult. Syst. iii. Mant. 274 = affine.
taxifolium, *Sm. in Rees, Cycl.* ix. n. 3.—Austral.
tenuifolium, *R. Br. in Trans. Linn. Soc.* x. (1811) 154. —Austral.
tenuifolium, Sieber, ex Roem. & Schult. Syst. iii. Mant. 275 = commutatum.
tenuifolium, *R. Br. in Trans. Linn. Soc.* x. (1811) 155. —Austral.
Toddii, *F. Muell. Fragm.* x. 20.—Austral.
tripinervium, *R. Br. Prod. Suppl.* 11.—Austral.
undulatum, *Lindl. Swan River App.* 31.—Austral.

CONOSTEGIA, D. Don, in Mem. Wern. Soc. iv. (1823) 316. *MELASTOMACEAE*, Benth. & Hook. f. i. 762.
BAUGUIERA, Rich. ex DC. Prod. iii. 178, in syn. (1828).
CALYCOTOMUS, Rich. ex DC. l. c. (1828).
SYNODON, Raf. in Sylva Tellur. 95 (1838).
acuminata, Steud. in Flora, xxvii. (1844) 722 = Miconia axillifera.
acutidentata, A. Rich. Ess. Fl. Cub. 556 = salapensis.
alpina, Macf. Fl. Jamaic. ii. 78 = montana.
alternifolia, Macf. Fl. Jamaic. ii. 71 = superba.
arborea, *Steud. Nom. ed.* II. i. 408; *Schaus. in Linnaea*, xx. (1847) 738.—Mexic.
attenuata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 98.—N. Granat.
Balbisiana, Ser. ex DC. Prod. iii. 174 = procera.
bractesta, *Triana, in Journ. Bot.* v. (1867) 309.— Nicaragua.
calyptrata, *D. Don, in Mem. Wern. Soc. iv.* (1823) 317. —Ind. occ.
clidemicoides, *Wright, ex Griseb. Cat. Pl. Cub.* 96.— Cuba.
cornifolia, Ser. ex DC. Prod. iii. 175 = Miconia cinnamomifolia.
cucullata, D. Don, in Mem. Wern. Soc. iv. (1823) 317 = Calyptrella cucullata.
dentata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 99. —N. Granat.
discolor, D. C. Prod. iii. 174 = Charianthus coccinea.
extinctoria, *D. Don, in Mem. Wern. Soc. iv.* (1823) 317.—N. Granat.
formosa, Macf. Fl. Jamaic. ii. 70 = rufescens.
Galeottii, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 107 = arbores.
glabra, D. Don, in Mem. Wern. Soc. iv. (1823) 317 = Antonia Forsteri.
gloriosa, Macf. Fl. Jamaic. ii. 68 = procera.
holosericea, D. Don, in Mem. Wern. Soc. iv. (1823) 317 = salapensis.
lasiopoda, *Benth. Bot. Voy. Sulph.* 96.—N. Granat.
lutescens, Ser. ex DC. Prod. iii. 175 = Miconia cinnamomifolia.
macrantha, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 97.—Costa Rica.
macrophylla, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 113 = superba.
mexicana, Ser. ex DC. Prod. iii. 175 = Miconia mexicana.
montana, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317. —Jamaic.
Mutisii, Ser. ex DC. Prod. iii. 174 = Centronia Mutisii.
myriocarpoides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 99.—N. Granat.
Oerstediana, *O. Berg, ex Triana, l. c.* 98.—Costa Rica.
parvifiora, *DC. Prod.* iii. 175.—Ins. S. Doming.
polyandra, *Benth. Bot. Voy. Sulph.* 96. t. 35.—N. Granat.
procera, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317. —Jamaic.
pulverulenta, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 110 = extinctoria.
purpurea, *Griseb. in Bonplandia*, vi. (1858) 6.— Panama.
quadrangularis, *Schlecht. ex Steud. Nom.* ed. II. i. 406.—Mexic.
rufescens, *Naud. in Ann. Sc. Nat. Sér.* III. xvi. (1851) 107.—Ind. occ.
semivernans, Ser. ex DC. Prod. iii. 175 = Miconia semivernans.
setosa, *Triana, in Trans. Linn. Soc. xxviii.* (1871) 99. —N. Granat.
speciosa, *Naud. in Ann. Sc. Nat. Sér.* III. xvi. (1851) 109.—Panama.
sphaerica, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 98.—Mexic.
suberostulosa, Triana, l. c. = purpurea.
subhirsuta, *DC. Prod.* iii. 174.—Cuba.
superba, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317.— Ins. exter.
tunicata, Ser. ex DC. Prod. iii. 175 = Miconia tunicata.
salapensis, *D. Don, in Mem. Wern. Soc. iv.* (1823) 317.—Mexic.

CONOSTEMUM, Lindl. Swan River App. 31. (1839). *MELASTOMACEAE*, Benth. & Hook. f. i. 757. SPHAIM. = COMOSTEMUM, Nees = Androtoechium, Bronga. (Cyperac.).

CONOSTEPHIOPSIS, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) 1. 5 = Conostephium, Benth. (Epacrid.).
Drummondi, Stschegl. l. c. 6 = Conostephium planifolium.
minor, Stschegl. l. c. = Conostephium minus.
Preissii, Stschegl. l. c. = Conostephium Preissii.

CONOSTEPHIUM, Benth. in Enum. Pl. Hueg. 76 (1837). *EPACRIDEAE*, Benth. & Hook. f. ii. 611. CONOSTEPHIOPSIS, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) 1. 5.
minus, *Lindl. Swan River App.* 25.—Austral.
nitens, *Lindl. ex Sond. in Lehm. Pl. Preiss.* i. 208.— Austral.
pendulum, *Benth. in Enum. Pl. Hueg.* 76.—Austral. *pendulum*, Deless. Ic. Sel. v. t. 28 = minus.
planifolium, *F. Muell. Fragm.* vi. 90.—Austral.
Preissii, *Sond. in Lehm. Pl. Preiss.* i. 204.—Austral.
Roei, *Benth. Fl. Austral.* iv. 160.—Austral.

CONOSTYLIS, R. Br. Prod. 300 (1810). *HAEMODORACEAE*, Benth. & Hook. f. iii. 676.
ANDROSTEMMA, Lindl. Swan River App. 46 (1839).
aculeata, *R. Br. Prod.* 300.—Austral.
aemula, Lindl. Swan River App. 45 = setigera.
albicans, A. Cunn. ex Benth. Fl. Austral. vi. 436 = candicans.
americana, Pursh, Fl. Am. Sept. i. 224. t. 6 = Lophiola aurea.
androstemma, *F. Muell. Fragm.* viii. 19.—Austral.
asswinitis, Endl. in Lehm. Pl. Preiss. ii. 20 = setigera.
aurea, *Lindl. Swan River App.* 44.—Austral.
Beallana, *F. Muell. Fragm.* ix. 50.—Austral.
bracteata, *Endl. in Lehm. Pl. Preiss.* ii. 16.— Austral.
bracteata, Lindl. Swan River App. 45 = dealbata.
breviscapa, *R. Br. Prod.* 301.—Austral.
bromelioides, *Endl. in Lehm. Pl. Preiss.* ii. 18.— Austral.
candicans, *Endl. Nov. Stirp. Dec.* 20.—Austral.
canescens, F. Muell. Fragm. viii. 19 = Blancoa canescens.
caricina, *Lindl. Swan River App.* 45.—Austral.
cymose, *F. Muell. ex Benth. Fl. Austral.* vi. 439.— Austral.
dealbata, *Lindl. Swan River App.* 45.—Austral.
discolor, Endl. in Lehm. Pl. Preiss. ii. 20 = setigera.
Drummondi, *Benth. Fl. Austral.* vi. 433.—Austral.
ensifolia, Endl. in Lehm. Pl. Preiss. ii. 21 = serrulata.
festucacea, Endl. l. c. 18 = Preissii.
filifolia, *F. Muell. Fragm.* viii. 18.—Austral.
gladiata, *Benth. Fl. Austral.* vi. 434.—Austral.
graminea, Endl. in Lehm. Pl. Preiss. ii. 19 = caricina.
involucrata, *Endl. l. c.* ii. 23.—Austral.
juncea, *Endl. Nov. Stirp. Dec.* 19.—Austral.
juncea, *Lindl. Bot. Reg.* (1839) App. 54.—Austral.
laxiflora, *Benth. Fl. Austral.* vi. 439.—Austral.
longifolia, Endl. in Lehm. Pl. Preiss. ii. 22 = serrulata.
melanopogon, *Endl. l. c.* ii. 18.—Austral.
minima, Endl. l. c. ii. 21 = C. Pyrillum.
misera, Endl. l. c. 22 = serrulata.
occulta, Endl. l. c. = serrulata.
petrophiloides, *F. Muell. ex Benth. Fl. Austral.* vi. 431.—Austral.
Preissii, *Endl. in Lehm. Pl. Preiss.* ii. 18.—Austral.
prolifera, *Benth. Fl. Austral.* vi. 436.—Austral.
propinqua, Endl. in Lehm. Pl. Preiss. ii. 17 = candicans.
Pyrillum, *Endl. l. c.* 21.—Austral.
pusilla, Endl. l. c. 20 = pyrillum.
racemosa, *Benth. Fl. Austral.* vi. 436.—Austral.
recurvifora, *F. Muell. Fragm.* i. 158.—Austral.
serrulata, *R. Br. Prod.* 300.—Austral.
setigera, *R. Br. l. c.*—Austral.
setosa, *Lindl. Swan River App.* 44. t. 6 A.—Austral.
spathacea, Endl. in Lehm. Pl. Preiss. ii. 22 = serrulata.
spinuligera, *F. Muell. ex Benth. Fl. Austral.* vi. 438.— Austral.
stylidioides, *F. Muell. Fragm.* viii. 17.—Austral.
subpaniculata, Endl. in Lehm. Pl. Preiss. ii. 17 = setosa.
tenuissima, *F. Muell. Fragm.* viii. 18.—Austral.
vaginata, *Endl. in Lehm. Pl. Preiss.* ii. 22.—Austral.
villosa, *Benth. Fl. Austral.* vi. 432.—Austral.

CONOSTYLUS, Pohl, ex A. DC. in Trans. Linn. Soc. xvii. (1834) 103 = Commocrinphe, A. DC. (Myrtac.). *Stenophalus*, Pohl, ex A. DC. l. c. 102 = Cuspa. ob-longifolia.

CONOTHAMNUS, *Lindl. Swan River App.* 9 (1839). *MYRTACEAE*, Benth. & Hook. f. i. 765.
divaricatus, *Benth. Fl. Austral.* iii. 144.—Austral.
tuberculis, *Lindl. Swan River App.* 9.—Austral.

CONOTRICHIA, A Rich. in Mém. Soc. Hist. Nat. Par. (1830) 277. t. 24. f. 1 = **Manettia**, Mutis (Rubiac.).
alba, A. Rich. l. c. = M. picta.

CONRADIA, Mart. Nov. Gen. et Sp. iii. 88 (1829) = **Pentarhaphia**, Lindl. (Generac.).
calycina, Mart. l. c. 88 = P. calycina.
calycosa, Hook. Ic. Pl. t. 889–890 = P. calycosa.
cassioides, Benth. in DC. Prod. x. 511, lapsu, in syn. = Seymeria tenuifolia.
cetioides, Griseb. Cat. Pl. Cub. 200.—Cuba.
clandestina, Griseb. Fl. Brit. W. Ind. 461.—Jamaica.
corrugata, Griseb. in Mem. Am. Acad. N. S. vii. (1863) 596.—Cuba.
Cranciaria, Mart. Nov. Gen. et Sp. iii. 88 = P. Cranciaria.
cuneifolia, DC. Prod. vii. 526 = P. cuneifolia.
deppeana, Griseb. Cat. Pl. Cub. 200.—Cuba.
exserta, Mart. Nov. Gen. et Sp. iii. 88 = P. exserta.
floribunda, Pæpp. Mag. Bot. xv. (1849) 99 = P. libanensis.
glaucinoides, Griseb. Cat. Pl. Cub. 200.—Cuba.
hispida, Benth. Pl. Hartw. 264 = P. hispida.
humilis, Mart. Nov. Gen. et Sp. iii. 88 = P. humilis.
libanensis, Griseb. Fl. Brit. W. Ind. 463 = P. libanensis.
mimuloides, Griseb. l. c. 461.—Jamaica.
neglecta, Hook. Ic. Pl. t. 874 = P. neglecta.
pedunculosa, DC. Prod. vii. 526 = P. pedunculosa.
pumila, Mart. Nov. Gen. et Sp. iii. 88 = P. pumila.
reticulata, Griseb. Cat. Pl. Cub. 201.—Cuba.
scabra, Mart. Nov. Gen. et Sp. iii. 88 = P. scabra.
Sloanei, DC. Prod. vii. 526 = P. Sloanei.
ventricosa, Mart. Nov. Gen. et Sp. iii. 88 = P. longiflora.
verrucosa, Scheidw. in Otto & Dietr. Allg. Gartenz. xv. (1847) 226 = P. verrucosa.

CONRADIA, Nutt. in Journ. Acad. Philad. vii. (1834) 85. t. 13 = **Macranthera**, Torr. (Scrophul.).
fuchsioides, Nutt. l. c. = M. fuchsioides.

CONRADIA, Rafin. Neog. 3 (1825) = **Tofieldia**, Huds. (Liliac.).

CONRADINA, A. Gray, in Proc. Am. Acad. viii (1870) 394. *LABIATAE*, Benth. & Hook. f. ii. 1191.
canescens, *A. Gray, l. c.* 395.—Am. bor.

CONRINGIA, Heist. ex Linn. Syst. ed. I (1735); Link, Enum. Hort. Berol. ii. 172 (1822). *CRUCIFERAE*, Benth. and Hook. f. i. 78.
CONRINGIA, J. & C. Presl, Fl. Čech. 140 (1819).
alpina, *Link, Enum. Hort. Berol.* ii. 173.—Europ.
austriaca, *Sweet, Hort. Brit.* ed. 1. 25.—Austria; Reg. Caucas.
clavata, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii (1842) 84.—As. Min.; Syria.
grandiflora, *Boiss. & Heldr. Diagn.* Ser. I. viii. 25.—As. Min.
nana, Heist. l. c. Ser. II. v. 25 = Malcolmia conringioides.
orientalis, *Dum. Fl. Belg.* 123.—Europ.; Orient.
perfoliata, Link, Enum. Hort. Berol. ii. 173 = orientalis.
persica, *Boiss. Diagn.* Ser. I. vi. 12.—Persia.
plantitiqua, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 25.—As. occ.
Thaliana, Reichb. Ic. Fl. Germ. ii. t. 60. fig. 4380 = Sisymbrium Thalianum.

CONSANA, Adans. Fam. ii. 420 (1763) = **Drabe**, Linn. (Crucif.).

CONSOLEA, Lem. Rev. Hortic. (1862) 174 = **Opuntia**, Tourn. (Cactese).
catacanthe, Lem. l. c. = O. catacantha.
ferox, Lem. l. c. = O. ferox.
leucacantha, Lem. l. c. = O. leucacantha.
rubescens, Lem. l. c. = O. rubescens.
spinosissima, Lem. l. c. = O. spinosissima.

CONSOLIDA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 711 (1821) = **Delphinium**, Tourn. (Ranunculac.).
Ajacis, Lindl. in Journ. Hort. Soc. vi. (1851) 55 = D. Ajacis.
Ajacis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = D. Ajacis.
arvensis, Opiz, Seznam, 32 = D. Consolida.
CONSOLIDA :—
paniculata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = Delphinium paniculatum.
regalis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 711 = D. Consolida.
segetum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = D. Consolida.

CONSOLIDA, Riv. ex Rupp. Fl. Jen. ed. Hall. 8 (1745) = **Symphytum**, Tourn. (Boragin.).
major, Gilib. Fl. Lituan. i. 34 = S. officinale.

CONSTANTIA, Rodrig. Orch. Nov. i. 78 (1877). *ORCHIDEAE*.
rupestris, *Rodrig. l. c.* 79.—Bras.

CONSUEGRA, Spreng. Neue Entdeck. ii. 808 (1821) = seq.

CONSUEGRIA, Mutis, ex Caldas, in Nuev. Seman. Gran. III. (1810) 22 (Quid?).

CONTARENA, Adans. Fam. ii. 120 (1763) = Corynmbium, Linn. (Compos.).

CONTARENIA, Vand. Fl. Lusit. et Bras. Spec. 42. t. 8. f. 20 (1788) (Gen. delend.). Benth. & Hook. f. ii. 1187.
trinervia, Vand. ex Roem. & Schult. Syst. iii. 294.—Bras.

CONTORTUPLICATA, Medic. in Vorles. Charpf. Phys. Ges. ii. (1787) 378 = **Astragalus**, Tourn. (Legumin.).
astragaloides, Medic. l. c. = A. contortuplicatus.

CONTRAKENIA, Jaume St. Hil. Expos. ii. 846 (1805) = CONTARENIA, Vand.

CONULEUM, A. Rich. in Mém. Soc. Hist. Par. i. (1823) 391. t. 25. *MONIMIACEAE*, Benth. & Hook. f. iii. 142.
guianense, *A. Rich. l. c.*—Guiana.

CONVALLARIA, Linn. Syst. ed. I (1735); Gen. ed. I. 96 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 770.
AXILLARIA, Rafin. in Journ. Phys. lxxix. (1819) 102, 261.
GLOMERIA, Rafin. Med. Fl. ii. 84 (1830).
LILIUM-CONVALLIUM, Tourn. ex Moench, Meth. 686 (1794).
ambigua, Schrank, Baier. Fl. i. 606 = Polygonatum multiflorum.
angulosa, Lam. Fl. Fr. iii. 266 = Polygonatum officinale.
angustifolia, Poir. Encyc. Suppl. v. 787 = Polygonatum biflorum.
biflora, Walt. Fl. Carol. 122 = Polygonatum biflorum.
bifolia, Linn. Sp. Pl. 316 = Smilacina bifolia.
borealis, Poir. Encyc. Suppl. v. 787 = Clintonia borealis.
bracteata, Dulac, Fl. Hautes-Pyr. 135 = majalis.
bracteata, Thowas, in Gand. Fl. Helv. ii. 581 = Polygonatum multiflorum.
Broteri, Gus. Fl. Sic. Syn. i. 416 = Polygonatum Gussonii.
canadensis, Poir. Encyc. Suppl. iv. 80 = Smilacina bifolia.
canaliculata, Willd. Enum. Hort. Berol. 45 = Polygonatum biflorum.
ciliata, Poir. Encyc. Suppl. iv. 80 = Smilacina racemosa.
cirrhifolia, Wall. in As. Res. xiii. (1820) 382 = Polygonatum sibiricum.
communata, Schult. f. Syst. vii. 1671 = Polygonatum latifolium.
compressa, Steud. Nom. ed. II. i. 406 = Polygonatum japonicum.
cordifolia, Stokes, Bot. Mat. Med. ii. 265 = Smilacina bifolia.
dichotoma, Thib. ex Pers. Syn. i. 372 = Streptopus distortus.
diffusa, Veil. Fl. Flum. 143; ill. t. 160 = Dichorisandra pubescens.
fragrans, Salisb. Prod. 254 = majalis.
fruticosa, Linn. Syst. Nat. ed. X. 984 = Cordyline terminalis.
gigantea, Hort. ex Kunth, Enum. Pl. v. 186 = Polygonatum latifolium.
glabra, *Kit. in Linnæa*, xxxii. (1863) 247.—Europ.
Gouaniana, Wall. Cat. n. 5187 = Polygonatum multiflorum.

CONVALLARIA:—
graminifolia, Salisb. Prod. 254 = Ophiopogon japonicus.
hirta, Bosc, ex Poir. Encyc. iv. 369 = Polygonatum latifolium.
hybrida, March. in Van Hall, Bijdr. iv. 843 = Smilacina stellata.
japonica, Linn. f. Suppl. 204 = Ophiopogon japonicus.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 148.—Japon.
latifolia, Jacq. Fl. Austr. iii. 18. t. 232 = Polygonatum latifolium.
latifolia, Lam. ex Steud. Nom. ed. II. i. 406 = Polygonatum multiflorum.
latifolia, Mill. Gard. Dict. ed. VIII. n. 2 = majalis.
latifolia, Savi, Bot. Etrusc. ii. 222 = Polygonatum officinale.
leptophylla, D. Don, Prod. Fl. Nep. 47 = Polygonatum verticillatum.
majalis, *Linn. Sp. Pl.* 314.—Reg. bor. temp.
Mappii, C. C. Gmel. Fl. Bad. ii. 52 = majalis.
monophylla, Oed. in Fl. Dan. t. 231 = Smilacina bifolia.
multiflora, Linn. Sp. Pl. 315 = Polygonatum multiflorum.
obtusifolia, Guenth. ex Steud. Nom. ed. II. i. 406 = Polygonum officinale.
odorata, Mill. Gard. Dict. ed. VIII. n. 4 = Polygonatum officinale.
oppositifolia, Wall. in As. Res. xiii. (1820) 380 = Polygonatum oppositifolium.
orientalis, Poir. Encyc. Suppl. iv. 80 = Polygonatum polyanthemum.
parviflora, Poir. l. c. 29 = Polygonatum officinale.
polyanthema, Bieb. Fl. Taur. Cauc. iii 272 = Polygonatum polyanthemum.
Polygonatum, Linn. Sp. Pl. 315 = Polygonatum officinale.
Polygonatum, Mnhl. Cat. 35 = Polygonatum latifolium.
Polygonatum, Thunb. Fl. Jap. 142 = Polygonatum japonicum.
pubescens, Willd. Hort. Berol. i. 45 = Polygonatum biflorum.
pulchella, Salisb. Prod. 255 = Smilacina bifolia.
punctata, Wall. Cat. n. 5133 = Polygonatum punctatum.
quadrifida, Lam. Fl. Fr. iii. 269 = Smilacina bifolia.
racemosa, Forsk. Fl. Aegypt. Arab. 73 = Smilacina Forskaliana.
racemosa, Linn. Sp. Pl. 315 = Smilacina racemosa.
racemosa, Vell. Fl. Flum. 142; ill. t. 159 = Dichorisandra Gaudichaudiana.
rosea, Leclerc, Fl. Alt. ii. 41 = Polygonatum roseum.
rupestris, Salisb. Prod. 254 = Polygonatum officinale.
scaposa, Gilib. Exercit. ii. 460 = majalis.
scopulosensis, Kit. in Linnæa, xxxii. (1863) 247 = glabra.
sibirica, Schult. f. Syst. vii. 1, 297 = Polygonatum sibiricum.
spicata, Thunb. Fl. Jap. 141 = Liriope graminifolia.
stellata, Linn. Sp. Pl. 316 = Smilacina stellata.
stolifolia, Peterm. in Flora, xxvii. (1844) 863 = Polygonatum verticillatum.
tetrapetala, Gilib. Exercit. ii. 461 = Smilacina bifolia.
Thunbergii, Steud. Nom. ed. II. i. 406 = Polygonatum latifolium.
trifolia, Linn. Sp. Pl. 316 = Smilacina trifolia.
umbellata, Bunge, ex Baker, in Joarn. Linn. Soc. xiv. (1875) 558 = Polygonatum umbellatum.
umbellulata, Michx. Fl. Bor. Am. i. 202 = Clintonia umbellata.
verticillata, Linn. Sp. Pl. 315 = Polygonatum verticillatum.

CONVOLVULOIDES, Moench, Meth. 451 (1794) = **Ipomœa**, Linn. (Convolv.).
alnifolia, Moench, l. c. 452 = I. sibirica.
leucosperma, Moench, l. c. = I. purpurea.
palmata, Moench, l. c. = I. Pes-tigridis.
pilosa, Moench, l. c. = I. pubescens.
purpurea, Moench, l. c. = I. purpurea.
trilobe, Moench, l. c. = I. hederacea.

CONVOLVULUS, [Tourn.]. Linn. Syst. ed. I (1735). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 874.
BUCHARRA, CALIXNOS, CAULOTULIS, CLEISOSTOMA, DITERRIA, EMULINA, EXALLOSIS, EXOCRIUS, IDALIA, LATRIENDA, LIZERON, NEMOSTIMA, PENTANTHUS, PERIPHAS, PODALETRA, RHODOXYLON, SAMIUM, SCIADIARA, SYMETHUS, THYELLA, Rafin. Fl. Tellur. 70–84 (1836).

CONVOLVULUS:—
Meresmia, Dennst. Schluss. Hort. Malab. 34 (1818).
Pantoceskra, Grimb. in Oestr. Bot. Zeitschr. (1878) 267.
Rhodoriza, Webb, in Bot. Reg. (1841) Misc. 69.
Syringotia, Neck. Elem. ii. 28 (1790).
abruptus, Spreng. Syst. i. 596 = Ipomœa abrupta.
abutiloides, H. B. & K. Nov. Gen. et Sp. iii. 106 = Ipomœa abutiloides.
acanthocladus, *Boiss. & Kotschy, Diagn.* Ser. I. vii. 27. —Persia.
acaulis, Choisy, in DC. Prod. ix. 406 = erubescens.
acetosæfolius, Steud. Nom. ed. II. i. 400 = Ipomœa martinicensis.
acetosæfolius, Turcz. in Bull. Soc. Nat. Mosc. (1840) 78 = Calystegia hederacea.
acetosæfolius, Vahl, Eclog. Am. i. 18 = Ipomœa carnosa.
acetosellæfolius, Desr. in Lam. Encyc. iii. 564 = Ipomœa acetosellæfolia.
acicularis, *Vatke, in Linnæa*, xliii. (1880–82) 518.—Afr. trop.
aculeatus, D. Dietr. Syn. Pl. i. 667 = Ipomœa Bonanox.
aculeatus, *Hill, Veg. Syst.* vii. 10 (= seq. ?).—Am. merid.
cuiensis, *Linn. Sp. Pl.* 155.—Am. trop.
acuminatus, Salzm. ex Meisn. in Mart. Fl. Bras. vii. 267 = Ipomœa fastigiata.
acuminatus, Vahl, Symb. Bot. iii. 26 = Ipomœa acuminata.
acutangulus, Spreng. Syst. i. 606 = Ipomœa coccinea.
Adansonii, Desr. in Lam. Encyc. iii. 560 = Ipomœa aquatica.
adpressus, Wall. Cat. n. 1424 = Lettsomia adpressa.
adscendens, De Vriese, in Lehm. Pl. Preiss. i. 346 = erubescens.
adscendens, Steud. ex Boiss. Fl. Orient. iv. 100 = reticulatus.
aegyptius, *Linn. Syst. ed.* X. 923.—Aegypt.
afghanus, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 168.—Afghan.
africh, *Schoele, in Linnæa*, xxi. (1848) 749.—Texas.
affinis, C. Sm. ex Steud. Nom. ed. II. i. 407 = Tuckeyanus.
affinis, Wall. Cat. n. 1385 = Ipomœa polysantha.
africanus, Nickols, Hort. St. Doming. 360 = Ipomœa cathartica.
aggregatus, Lour. Fl. Cochinch. 109 = Ipomœa aggregata.
aggregatus, Steud. Nom. ed. II. i. 407 = Lettsomia aggregata.
agrestis, Mart. ex Choisy, in DC. Prod. ix. 405 = Jacquemontia agrestis.
agrophilus, C. Koch, in Linnæa, xxii. (1849) 745 = galaticus.
agrostophyllus, Vell. Fl. Flum. 71; ii. t. 51 = Breweria Burchellii.
Aitchisoni, *C. B. Clarke, in Journ. Linn. Soc.* xix. (1883) 179.—Afghan.
alatus,Ham. Prod. Fl. Ind. Occ. 94 = Ipomœa Turpethum.
alatus, Spreng. Syst. i. 596 = Ipomœa alata.
albicans, Wall. Cat. n. 1401 = Ipomœa pilosa.
albiflorus, West, Bidr. Ste Croix, 271 = nodiflorus.
albivenius, Lindl. Bot. Reg. t. 1116 = Ipomœa albivenia.
albus, [*Nacca*] *Onom. Hort. Ticin.* (1812) 19.—Hab.?
alcæafolius, Bory & Chaub. Fl. Pélop. 14 = althæoides.
alceæfolius, *Lam. Tabl. Encyc.* i. 461.—Afr. austr.
alcinoides, Linn. Sp. Pl. 157 = Evolvulus alsinoides.
althæoides, *Linn. f. c.* 156.—Reg. Mediterr.
althæoides, Tenore, Syll. Fl. Neap. 92 = tuberosus.
althæoides, Thunb. Fl. Cap. ii. 18 = Thunbergii.
altissimus, Spreng. Syst. i. 613 = Ipomœa staphylina?
amblyocalyx, Steud. Nom. ed. II. i. 407 = fatmensis.
Ammannii, *Desr. in Lam. Encyc.* iii. 549.—Sibir.
ammocharis, *Boiss. & Hausk. Pl. Or. Nov. Dec.* i. 6. —Persia.
anamophilus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = ammocharis.
amœnus, D. Dietr. Syn. Pl. i. 670 = Ipomœa amœna.
amœnus, C. Koch, in Linnæa, xix. (1846) 19 = hirsutus.
amplexifolius, Cham. & Schlecht. i. c. v. (1830) 118 = Ipomœa quinquefolia.

CONVOLVULUS:—
amplicaulis, Herb. Brit. Mus. ex Choisy, in DC. Prod. ix. 361 = Ipomœa amplicaulis.
amplus, Wall. Cat. n. 1490 = Argyreia Roxburghii.
ampullaceus, Vahl, ex Choisy, Conv. Or. 61 = Lepistemon Wallichii.
anceps, Linn. Mant. i. 48 = Ipomœa Turpethum.
andinus, *Phil. in Linnæa*, xxxiii. (1864–65) 184.—Chili.
angularis, Burm. f. Fl. Ind. 46 = Ipomœa vitifolia.
angulatus, Buch.-Ham. ex Choisy, Conv. Or. 58 = Ipomœa coccinea.
angulatus, Salzm. ex Choisy, in Mém. Soc. Phys. Genèv. ix. (1837) 137 = Ipomœa bahiensis.
angulatus, Spreng. Syst. i. 594 = Ipomœa angulata.
angustifolius, Desr. in Lam. Encyc. iii. 547 = Ipomœa angustifolia.
angustatissimus, R. Br. Prod. 482 = erubescens.
aphyllus, Viv. Ann. Bot. i. 95 = Ipomœa pentaphylla.
apocynoides, Cham. & Schlecht. in Linnæa, v. (1830) 117 = Jacquemontia hirsuta.
aquaticus, Walt. Fl. Carol. 94 = Breweria aquatica.
arborescens, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 284 = Ipomœa arborescens.
arborens, Balb. ex Steud. Nom. ed. II. i. 407 = præc.
arcuatus, Presl, Fl. Sic. p. xxxiii = pentapetaloides.
arenarius, Vahl, Symb. Bot. i. 18 = Ipomœa carnosa.
argenteus, Heyne, ex Wall. Cat. n. 1394 = Argyreia fulgens.
argenteus, Lam. Fl. Fr. ii. 266; Desr. in Lam. Encyc. iii. 553 = C. Cneorum.
argenteus, Fourr. in Mém. Acad. Toul. iii. (1788) 316 = lanuginosus.
argenteus, Wall. Cat. n. 1364 = Argyreia argentea.
argyreus, DC. Fl. Fr. Suppl. 423 = althæoides.
argyrophyllus, Hoffmgg. Verz. Pfl. 53 = althæoides.
argyrophyllus, Wall. Cat. n. 1395 = Argyreia lanceolata.
aristolochiæfolius, H. B. & K. Nov. Gen. et Sp. iii. 102 = Ipomœa aristolochiæfolia.
aristolochiæfolius, Mill. Gard. Dict. ed. VIII. n. 9 = Ipomœa umbellata.
aristolochiæfolius, Willd. ex Roem. & Schult. Syst. iv. 802 = Humboldtianus.
armatus, Delile, Fl. Egypte, 189. t. 18 = C. Hystrix.
armenus, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 105. —Armenia; Syria.
arvensis, *Linn. Sp. Pl.* 153.—Geront. temp.
asarifolius, Desr. in Lam. Encyc. iii. 562 = Ipomœa asarifolia.
asarifolius, Sallab. Prod. 125 = Calystegia Soldanella.
ascendens, Steud. ex Choisy, in DC. Prod. ix. 400 = reticulatus.
ascendens, Walp. Rep. vi. 540 = erubescens.
asclepiadeus, Wall. Cat. n. 1396 = parviflorus.
asper, Wall. l. c. n. 1398 = Ipomœa grandiflora.
assyricus, *Griseb. Spicil. Fl. Rumel.* ii. 75, in nota.—Asa Min.; Mesopotam.
astrensis, *Kotschy, in Sitzb. Acad. Wien, Math. Nat. lii. Abth.* i. (1866) 260.—Oriens.
atriplicifolius, Poir. Encyc. Suppl. iii. 467 = hirsutus.
atropurpureus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 54 = Lettsomia atropurpurea.
attenuatus, *Mart. & Gal. in Bull. Acad. Brux.* xii. ii. (1845) 261.—Mexic.
aturensis, H. B. & K. Nov. Gen. et Sp. iii. 96 = Ipomœa aturensis.
Auscheri, *Choisy, in DC. Prod.* ix. 402.—Syria.
aureus, D. Dietr. Syn. Pl. i. 672 = Jacquemontia hirsuta.
aureus, *Thunb. Mus. Nat. Ups.* 185.—Hab.?
auriculatus, Desr. in Lam. Encyc. iii. 540 = arvensis.
auritus, *Willd. ex Roem. & Schult. Syst.* iv. 801.—Ins. Porto Rico.
aurosus, Desr. in Lam. Encyc. iii. 554; Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 107 = Jacquemontia hirsuta.
babylonicus, *Tenore, Ind. Sem. Hort. Nap.* (1839) 16. —? Afr.
bahiensis, Spreng. Syst. i. 596 = Ipomœa bahiensis.
barbeyi, *Pomel, Nouv. Mat. Fl. Atl.* 294.—Afr. bor.
barbatus, Wall. Cat. n. 1391; Pl. As. Rar. ii. 45. t. 155 = Lettsomia barbata.
barbiger, Wall. Cat. n. 1404 = Lettsomia barbigera.
barlerioides, Buch.-Ham. ex Wall. l. c. n. 1396 = Ipomœa barlerioides.
Batata, Veil. Fl. Flum. 76; ii. t. 89(= seq. ?).—Bras.
Batatas, Linn. Sp. Pl. 154; *Amœn. Acad.* vi. 121 = Ipomœa Batatas.

CONVOLVULUS:—
batatæfolius, Boiss. & Hausk. ex Boiss. Fl. Orient. iv. 106 = armenus.
Batatilla, H. B. & K. Nov. Gen. et Sp. iii. 106 = Ipomœa Batatilla.
bauhiniæfolius, Salisb. Prod. 125 = Ipomœa Pescapræ.
Beladambu, Spreng. Syst. i. 608 = Ipomœa Beladambœa.
bellus, Spreng. l. c. 590 = Ipomœa pulchella.
benehoavensis, *C. Bolle, in Bonplandia*, ix. (1861) 54. —Ins. Canar.
Benthami, Wall. Cat. n. 1393 = Ipomœa hirsutula.
Bentzii, Buch.-Ham. ex Wall. l. c. sub n. 1388 = Ipomœa martinicensis.
Besseri, Spreng. Syst. i. 610 = lineatus.
betonicæfolius, Mill. Gard. Dict. ed. VIII. n. 20 = hirsutus.
bibractiatus, Poepp. ex Meisn. in Mart. Fl. Bras. vii. 270 = Ipomœa luxurians.
bicolor, Desr. in Lam. Encyc. iii. 564 = Ipomœa pilosa.
bicolor, Vahl, Symb. Bot. iii. 25 = Hewittia bicolor.
bicuspidatus, Fisch. ex Link, Enum. Hort. Berol. i. 201 = arvensis.
bidentatus, *Bernh. in Flora*, xxvii. (1844) ii. 629.—Afr. austr.
bifidus, Vahl, Symb. Bot. iii. 30 = Ipomœa cymosa.
bifidus, Vell. Fl. Flum. 71; ii. t. 59 (= *Aniseia Velloziana*).—Bras.
biflorus, Forsk. Fl. Aegypt. Arab. 908 = Ipomœa acetosæfolia.
biflorus, *Georgi, Beschr. Russ. Reich. Nachtr.* 261 (quid?).—Tauria.
biflorus, Linn. Sp. Pl. ed. II. 1668 = Ipomœa biflora.
bigelandulosus, Stokes, Bot. Mat. Med. i. 326 = Ipomœa biloba.
bignonioides, Spreng. Syst. iv. Cur. Post. 60 = Ipomœa bignoniodes.
bilobatus, Roxb. Hort. Beng. 14; Fl. Ind. i. 485 = Ipomœa biloba.
binecturiferus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 47 = Lepistemon Wallichii.
Blanchetii, Steud. Nom. ed. II. i. 407 = Jacquemontia Blanchetii.
blandus, Roxb. Hort. Beng. [13]; Fl. Ind. i. 470 = Ipomœa cymosa.
Blumii, D. Dietr. Syn. Pl. i. 675 = Ipomœa angustifolia.
Boerhavia, [*Spreng.*] *Bot. Cent. Halle*, 38.—Hab.?
boerhavioides, Blanco, Fl. Filip. ed. II. 67 = Breweria valerianoides.
bogotensis, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 203 = Ipomœa bogotensis.
Boissieri, Steud. Nom. ed. II. i. 407 = nitidus.
Bona-nox, Spreng. Syst. i. 600 = Ipomœa Bona-nox.
bonariensis, Cav. Ic. v. 54. t. 480 = incanus.
Bonplandianus, Roem. & Schult. Syst. iv. 884 = Ipomœa sidæfolia.
Bonplandii, Meisn. in Mart. Fl. Bras. vii. 235 = præc.
Bourgæi, *C. Bolle, in Bonplandia*, ix. (1861) 54.—Ins. Teneriffæ.
brachyphyllus, Boiss. Diagn. Ser. II. iii. 122 = scindicus.
bracteatus, Bert. & Balb. ex Choisy, in DC. Prod. ix. 359 = Ipomœa setifera.
bractesatus, Vahl, Symb. Bot. iii. 25 = Hewittia bicolor.
brasiliensis, L. Sp. Pl. 159 = Ipomœa Pes-capræ.
brevicuspinatus, Mart. & Choisy, in DC. Prod. ix. 409 = Jacquemontia racemosa.
brevifforus, Spreng. Syst. i. 606 = Ipomœa setifera.
brevipes, *Pomel, Nouv. Mat. Fl. Atl.* 36.—Afr. bor.
breyniphyllus, Jaub. & Spach, Illustr. iv. 106. t. 369 = fruticosus.
Brownii, Spreng. Syst. i. 590 = Ipomœa quinata.
Brownii, Spreng. l. c. 612 = Ipomœa heptaphylla.
bryoniæfolius, Reichb. Fl. Germ. Excurs. 848 = althæoides.
bryoniæfolius, Salisb. Prod. 125 = Ipomœa Pes-tigridii.
bryoniæfolius, *Sims, Bot. Mag.* t. 943.—Reg. Mediterr.
bufalinus, Lour. Fl. Cochinch. 109 = Ipomœa bufalina.
Burmanni, *Choisy, in DC. Prod.* ix. 406.—Afr. austr.
caballeroides, Boiss. in Prod. Acad. Sc. Phlad. 1862 (1863) 6 = Ipomœa leptophylla.
cærulescens, Auct. vide convolvens.
cæspitosus, Roxb. Hort. Beng. 14; Fl. Ind. i. 488 = Hewittia bicolor.
cairicus, Linn. Syst. ed. X. 923 = Ipomœa palmata.

CONVOLVULUS :—

californicus, Benth. Pl. Hartw. 326 = luteolus.
californicus, *Choisy, in DC. Prod.* ix. 406.—Am. bor. occ.
caliginosus, Koen. ex Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 461 = Ipomœa cymosa.
Calvertî, *Boiss. Diagn.* Ser. II. iii. 154.—Tauria ; As. Min.
calycinus, *Drége, ex Choisy, in DC. Prod.* ix. 408.—Afr. austr.
calycinus, H. B. & K. Nov. Gen. et Sp. iii. 109 = Ipomœa cissoides.
calycinus, Roxb. Hort. Beng. [13] ; Fl. Ind. i. 471 = Ipomœa calycina.
calyculatus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 10.—Malaya.
calystegioides, *Choisy, Conv. Or.* 98.—China.
campanulatus, Spreng. Syst. i. 608 = Ipomœa campanulata.
campestris, Vell. Fl. Flum. 74 ; ii. t. 66 = Ipomœa polymorpha.
canariensis, *Linn. Sp. Pl.* 155.—Ins. Canar.
candicans, Roem. & Schult. Syst. iv. 302 = Rivea hypocrateriformis.
candicans, Roth, Nov. Pl. Sp. 106 = Rivea ornata.
candicans, Soland. ex Sims, in Bot. Mag. t. 1608 = Ipomœa fastigiata.
canescens, H. B. & K. Nov. Gen. et Sp. iii. 99 = Jacquemontia violacea.
canescens, *Phil. in Linnæa*, xxxiii. (1864–65) 182.—Chili.
canescens, Roth, Nov. Pl. Sp. 107 = Lettsomia aggregata.
cantaber, Pall. Phyt. Topogr. Gem. Taur. 99 = C. Cantabrica !
Cantabrica, *Linn. Sp. Pl.* 158.—Europ. ; Orient.
capensis, *Burm. f. Fl. Cap. Prod.* 5.—Afr. austr.
capensis, Willd. Sp. Pl. i. 877 = hastata.
capillaceus, H. B. & K. Nov. Gen. et Sp. iv. 260 = Ipomœa muricata.
capitatus, Cav. Ic. ii. 72. t. 189 = lanuginosa.
capitatus, Desr. in Lam. Encyc. iii. 554 = Jacquemontia tamnifolia.
capitatus, Schimp. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 25 = glomerulus.
capitatus, Vahl, Symb. Bot. iii. 28 = Lettsomia strigosa.
capitellatus, Buch.-Ham. ex Wall. Cat. n. 2250 = Ipomœa Pes-tigridis.
capitiformis, Poir. Encyc. Suppl. iii. 469 = Lettsomia strigosa.
capituliferus, *Franch. in Rèvoil, Çomali*, 41.—Afr. or. trop.
capripes, Stokes, Bot. Mat. Med. i. 327 = Ipomœa Pes-capræ.
Caput-Medusæ, *Lowe, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 155.—Ins. Madera.
caracasanus, Willd. ex Roem. & Schult. Syst. iv. 301 = Ipomœa umbellata.
cardiosepalus, *Boiss. Fl. Orient.* iv. 96.—Cilicia.
carneus, Spreng. Syst. i. 602 = Ipomœa carnea.
carnosus, Spreng. l. c. 608 = Ipomœa carnosa.
carolinus, Linn. Sp. Pl. 154 = Ipomœa commutata.
catesæicus, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. dec.* i. 5.—As. Min.
Catesbæi, Spreng. Syst. i. 608 = Calystegia sepium.
catharticus, Blanco, Fl. Filip. ed. I. 94 = Ipomœa grandiflora.
catharticus, *Cranto, Inst.* ii. 885.—Am. trop. ?
Cavanillesii, Spreng. Syst. i. 590 = Ipomœa Cavanillesii.
cephalanthus, Mart. ex Choisy, in DC. Prod. ix. 406 = Ipomœa muricata.
cephalophorus, *Boiss. Diagn.* Ser. I. vii. 92.—Persia.
cephalopodus, Boiss. l. c. 94 = sericeus.
cerpitense, Ehrenb. ex Boiss. Fl. Orient. iv. 97 = Blanoticus.
chenopodioides, *Desr. in Lam. Encyc.* iii. 544.—Hab. ?
Chevleri, Agardh, ex Roem. & Schult. Syst. iv. 261 = arvensis.
chilensis, Pers. Syn. i. 180 = dissectus.
chinensis, Ker-Gawl. in Bot. Reg. t. 223 = arvensis.
Chodjamuss, Wall. Cat. n. 2250 = Argyreia Lascheanaidii.
choisiocola, Spreng. Syst. i. 599 = Ipomœa coccinea.
chorassifolius, *Boiss. Diagn.* Ser. I. vii. 93.—Persia.
Chrysidesmi, Steud. Nom. ed. II. i. 466, lapsu = *Chrysidesmus*.

CONVOLVULUS :—

chrysoides, Spreng. Syst. i. 598 = Ipomœa chrysoides.
chrysorrhizus, Schrad. (vel Soland.) ex Steud. Nom. ed. II. i. 408, 109 = Ipomœa mammosa !
ciliatus, Roth, Catalect. fasc. i. 29 = undulatus.
ciliatus, Vahl, Eclog. Am. ii. 18 = Jacquemontia tamnifolia.
ciliolatus, Michx. Fl. Bor. Am. i. 137 = Ipomœa pandurata.
cinerosus, Spreng. Syst. i. 609 = Ipomœa incisa.
circinnatus, Willd. ex Roem. & Schult. Syst. iv. 302 = Ipomœa parasitica.
cirrhosus, *R. Br. in Salt, Abyss. App.* p. lxiv ; *ed. Germ.* 475.—Abyss.
cissoides, Lam. Tabl. Encyc. i. 462 ; Vahl, Eclog. Am. ii. 18 = Ipomœa cissoides.
cladotrichus, *Mart. Del. Sem. Hort. Monac.* (1846) 4. —Mexic.
Clappertonii, Spreng. Syst. iv. 62 = Ipomœa Clappertonii.
clavatus, Pav. ex Choisy, in DC. Prod. ix. 46 (= *Calonyction clavatum*).—Bras.
Cneorum, Forsk. Fl. Aegypt. Arab. p. lxiii = lanatus.
Cneorum, Habl. Taur. 147, ex Bieb. Fl. Taur. Casc. i. 146 = Calvertî.
Cneorum, *Linn. Sp. Pl.* 157.—Europ. austr.
Cneorum, Sprun. ex Nym. Consp. 505 = oleæfolius.
coccineus, Salisb. Prod. 124 = Ipomœa coccinea.
cochlearis, *Griseb. Spicil. Fl. Rumel.* ii. 76, in nota.—Græc. ; As. Min.
coelestis, Forst. f. Prod. 14 = Ipomœa hederacea.
coelesyriacus, *Boiss. Diagn.* Ser. I. xi. 85.—Syria.
coeruleus, Mart. & Gal. in Bull. Acad. Brux. xii. 11. (1845) 359 = Jacquemontia pavifiora.
coeruleus, Schum. & Thonn. Beskr. Guin. Pl. 101 = Ipomœa ovalifolia.
coeruleus, Sprengel, Syst. i. 593 = Ipomœa hederacea.
colubrinus, Blanco, Fl. Filip. ed. II. 66 = Ipomœa muricata.
commutatipilus, Steud. ex Choisy, in DC. Prod. ix. 354 = Ipomœa commutatipila.
Commersonii, Steud. Nom. ed. II. i. 408 = Evolvulus incanus.
communatus, *Boiss. Diagn.* Ser. I. xi. 81.—Mesopot.
compactus, *Boiss. l. c.* iv. 40.—As. Min.
condensatus, Bertol. Misc. Bot. xv. 18 = Jacquemontia tamnifolia.
confertus, Hort. Hamb. ex Steud. Nom. ed. II. i. 408 = Argyreia setosa.
congestus, *R. Br. in Salt, Abyss. App.* p. lxiv.—Abyss.
congestus, Spreng. Syst. i. 601 = Ipomœa congesta.
contortus, Vell. Fl. Flum. 70 ; ii. t. 48 = Ipomœa glabra.
copticus, Linn. Mant. ii. 559 = Ipomœa dissecta.
cordatifolius, Vell. Fl. Flum. 78 ; ii. t. 60 = Ipomœa Batatas.
cordifolius, Bucle, ex Choisy, in DC. Prod. ix. 381 = Ipomœa Bucli.
cordifolius, Jus. ex Choisy, l. c. 412 = farinosus.
cordifolius, Salzm. ex Meisn. in Mart. Fl. Bras. vii. 267 = Ipomœa fastigiata.
cordifolius, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
coriaceus, Herb. Brit. Mus. ex Choisy, in DC. Prod. ix. 386 = Ipomœa pentaphylla.
Cornuti, *Steud. Nom.* ed. II. i. 408.—Chili.
corsicus, Roem. & Schult. Syst. iv. 256 = arvensis.
corymbosus, Forst. f. Prod. 14 = parviflorus.
corymbosus, Linn. Syst. ed. X. 923 = Ipomœa Burmanni.
costatus, Meyen, Reise, i. 364 (= *Aniseia costata*).—Chili.
crassifolius, *Schenle, in Linnæa*, xxi. (1848) 752.—Bras.
crassipes, Kunze, ex Choisy, in DC. Prod. ix. 433 = Calystegia sepium.
crenatifolius, *Ruiz & Pav. Fl. Per.* ii. 10.—Peruv.
crenatus, Jacq. Coll. ii. 277 = Hermannise.
crenatus, Vahl, Symb. Bot. iii. 31 = Vahlii.
crinitus, Desr. in Lam. Encyc. iii. 568 = Jacquemontia tamnifolia.
crispatulus, Wall. Cat. n. 1408 = Ipomœa petaloidea.
crispifolius, *F. Muell. in Linnæa*, xxv. (1852) 422.—Austral.
crispus, *Thunb. Fl. Cap.* ii. 15.—Afr. austr.
cumanensis, H. B. & K. Nov. Gen. et Sp. iii. 99 = Ipomœa farinosa.

CONVOLVULUS :—

cunoatus, Willd. Sp. Pl. i. 872 = Argyreia cuneata.
cuneiformis, Buch.-Ham. ex Wall. Cat. p. 84. n. 1366 = Argyreia cuneata.
Cupanianus, Tod. in Ann. Sc. Nat. Sér. IV. xx. (1863) 304 = tricolor.
Curtisus, Buch.-Ham. ex Wall. Cat. p. 83, n. 1344 = Ipomœa eriocarpa.
cuspidatus, Willd. ex Spreng. Syst. i. 697 = Meyeri.
cymosus, Desr. in Lam. Encyc. iii. 556 = Ipomœa cymosa.
cymosus, Herb. Madr. ex Wall. Cat. p. 84, n. 1366 = Argyreia cuneata.
cymosus, Roem. & Schult. Syst. iv. 303 = Senegambiæ.
cymosus, Ruiz & Pav. Fl. Per. ii. 9 = Ipomœa ramosissima.
cynanchifolius, Wall. Cat. n. 1399 = Ipomœa cynanchifolia.
cynanchifolius, Willd. ex Roem. & Schult. Syst. iv. 301 = mollis, Ipomœa Bona-nox.
cyprius, *Boiss. Fl. Orient.* iv. 98.—Ins. Cyprus.
dahuricus, Herb. in Bot. Mag. t. 2609 = Calystegia dahurica.
damascenus, Boiss. & Gaill. Diagn. Ser. II. vi. 122 = stachydifolius.
dasycephalus, Pall. ex Choisy, in DC. Prod. ix. 401 = lanuginosus.
dasyspermus, Spreng. Syst. i. 591 = Ipomœa dasysperma.
decumbens, Orteg. ex Steud. Nom. ed. II. i. 408 = undulatus.
defloratus, *Choisy, in Zoll. Syst. Vers. Ind. Archip.* 130, 193.—Malaya.
demissus, *Choisy, in DC. Prod.* ix. 404.—Chili.
dentatus, Blanco, Fl. Filip. ed. I. 89 = Ipomœa Blanconi.
dentatus, Vahl, Symb. Bot. iii. 25 = Ipomœa chrysoides.
denticulatus, Desr. in Lam. Encyc. iii. 540 = Ipomœa denticulata.
denticulatus, Spreng. Syst. i. 603 = Ipomœa angustifolia.
denticulatus, Steud. Nom. ed. II. i. 408 = Ipomœa chrysoides.
deserti, Hochst. & Steud. ex Steud. l. c. = microphyllus.
dianthoides, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 708 = C. Pseudo-Cantabrica.
Dianthus, J. F. Gmel. Syst. 348 = Ipomœa littoralis.
dichrous, Koem. & Schult. Syst. iv. 263 = Ipomœa pilosa.
digitatus, Buch.-Ham. ex Wall. Cat. p. 85. n. 1423 = Ipomœa cissoides.
digitatus, Roxb. Hort. Beng. 14 ; Fl. Ind. i. 479 = Ipomœa palmata.
digitatus, Spreng. Syst. i. 592 = Ipomœa digitata.
Dillenii, Desr. in Lam. Encyc. iii. 544 = Ipomœa hederacea.
discolor, H. B. & K. Nov. Gen. et Sp. iii. 105 = Ipomœa discolor.
dissectus, Cav. Ic. v. 54 = incanus.
dissectus, Herb. Madr. ex Wall. Cat. p. 84, n. 1351 = Ipomœa dissecta.
dissectus, Jacq. Obs. Bot. ii. 4 = Ipomœa sinuata.
distillatorius, Blanco, Fl. Filip. ed. I. 96 = Ipomœa campanulata.
divaricatus, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 537, 588.—Turkestan.
divaricatus, Wall. Cat. n. 1402 = arvensis.
diversifolius, Kunze, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 867 = incanus.
diversifolius, Schum. & Thonn. Beskr. Guin. Pl. 94 = Ipomœa sagittata.
diversifolius, Spreng. Syst. i. 592 = Ipomœa diversifolia.
divinus, Vell. Fl. Flum. 71 ; ii. t. 49 = Jacquemontia membranacea.
domingensis, Desr. in Lam. Encyc. iii. 554 = Ipomœa tuberosa.
domingensis, Spreng. Nova Enchirid. iii. 29 = Ipomœa Hispaniolæ.
doryoniclides, *Den. Rep. Lig.* 282.—Ligur. ; Ins. Palm. Doryoniclides, *Linn. Syst.* ed. X. 923.—Græcia.
Doryonium, Pall. ex Steud. Nom. ed. II. i. 408 = lanuginosus !

CONVOLVULUS:—
Dregeanus, *Choisy, in DC. Prod.* ix. 411.—Afr. austr.
Deyandri, Spreng. Syst. i. 597 = Massoni.
Dudia, Buch.-Ham. ex Wall. Cat. p. 88 n. 1238 = Calystegia hederacea.
dumetorum, H. B. & K. Nov. Gen. et Sp. iii. 101 = Ipomoea dumetorum.
Durandoi, *Pomel, Nouv. Mat. Fl. Atl.* 85.—Afr. bor.
ebracteatus, Desr. in Lam. Encyc. iii. 541 = Ipomoea ebracteata.
echioides, Hochst. ex Choisy, in DC. Prod. ix. 382 = Argyreia abyssinica.
edulis, Thunb. Fl. Jap. 84 = Ipomoea Batatas.
edulis, *Vell. Fl. Flum.* 73; ii. t. 55.—Bras.
Eichwaldi, Boiss. Diagn. Ser. I. vii. 27, in nota = fruticosa.
elegans, Wall. Cat. n. 1892 = Breweria elegans.
elegantissimus, Mill. Gard. Dict. ed. VIII. n. 22 = silhasoides.
ellipticus, Spreng. Syst. i. 618 = Lettsomia elliptica.
elongatus, Salisb. Prod. 128 = C. Scammonia.
elongatus, *Willd. Enum. Hort. Berol.* 205.—Ins. Canar.
emarginatus, *Link, Enum. Hort. Berol.* i. 201.—Hab.?
emarginatus, Vahl, Symb. Bot. iii. 23 = Ipomoea maritimoides.
equitans, Benth. Pl. Hartw. 16 = incanus.
erectus, Spreng. Syst. i. 612 = Ipomoea erecta.
eremophilus, *Boiss. & Buhse, in Nouv. Mem. Soc. Nat. Mosc.* xii. (1860) 148.—Persia.
erianthus, Willd. ex Spreng. Syst. i. 610 = Breweria aquatica.
erinaceus, *Ledeb. in Eichw. Casp. Cauc.* 11. t. 7.—Reg. Casp.
erinaceus, C. A. Mey. Verz. Pfl. Cauc. 102 = eremophilus.
eriocarpus, Spreng. Syst. i. 598 = Ipomoea eriocarpa.
eriocaulos, Willd. ex Roem. & Schult. Syst. iv. 301 = Ipomoea purpurea.
eriospermus, Desr. in Lam. Encyc. iii. 567 = Ipomoea Desrousseauxii.
erosus, Desr. l. c. 558 = Hermanniae.
erubescens, *Sims, Bot. Mag.* t. 1067.—Austral.
erythrocarpus, Wall. Cat. n. 1413 = Argyreia lanceolata.
esculentus, Salisb. Prod. 132 = Ipomoea Batatas.
esculentus, Vell. Fl. Flum. 73; ii. t. 58 = septangularis.
exaequebenetis, Spreng. Syst. i. 600 = Ipomoea fastigiata.
evolvuloides, *Boiss. Diagn.* Ser. I. vii. 25.—Persia.
evolvuloides, Desf. Fl. Allant. i. 178. t. 49 = undulatus.
excisus, Zipp. ex Span. in Linnaea, xv. (1841) 388 = Ipomoea reniformis.
Falckia, Thunb. Prod. Pl. Cap. 35 = Falkia repens.
Falkia, *Spicq. Hort. Schoenh.* ii. 38. t. 198.—Afr. austr. farinosus *Linn. Mant.* ii. 208.—Afr. trop.
fastigiatus, Herb. Roxb. ex Wall. Cat. n. 2258 = Argyreia populifolia.
fastigiatus, Roxb. Hort. Beng. [18]; Fl. Ind. i. 468 = Ipomoea fastigiata.
fatmensis, *Kunze, in Flora*, xxiii. (1840) i. 172.—Afr. bor.
ferrugineus, Stend. Nom. ed. II. i. 408 = Jacquemontia ferruginea.
ferrugineus, Vahl, Eclog. Am. i. 17 = Ipomoea ferrugines.
festivus, Wall. Cat. n. 1414 = Argyreia argentea, populifolia.
filicaulis, Vahl, Symb. Bot. iii. 24 = Ipomoea angustifolia.
filifolius, Hook. & Arn. Bot. Beech. Voy. 35 = Ipomoea Cruckshanksii.
filiformis, Desr. in Lam. Encyc. iii. 555 = Ipomoea filiformis.
filiformis, Thunb. Fl. Cap. ii. 16 = Ipomoea angustifolia.
filipes, *Bailf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 62.—Ins. Socotr.
Finlaysonianus, Wall. Cat. n. 6807 (Argyreiae sp.).—Cochinch.
firmus, Spreng. Syst. i. 613 = Ipomoea stans.
flagelliformis, Roxb. Hort. Beng. 14; Fl. Ind. i. 481 = Ipomoea Beladamboo.
flavescens, D. Dietr. Syn. Pl. i. 680 = Ipomoea flavescens.
flavus, Heyne, ex Wall. Cat. sub n. 1849 = Ipomoea chrysides.
flavus, Salisb. Prod. 134 = Ipomoea umbellata.

CONVOLVULUS:—
flavus, *Willd. Sp. Pl.* i. 859.—Ind. or.
flexuosus, *Pomel, Nouv. Mat. Fl. Atl.* 84.—Algeria.
flexuosus, Spreng. Syst. i. 612 = Ipomoea erecta.
floribundus, *H. B. & K. Nov. Gen. et Sp.* iii. 96.—Am. austr.
floridus, *Linn. f. Suppl.* 136.—Ins. Teneriff.
formosus, J. F. Gmel Syst. 343 = Ipomoea sagittata.
formosus. Roxb. ex Hook. f. Fl. Brit. Ind. iv. 214 = Ipomoea kentrocaulos.
Forskalei, Delile, Fl. Egypte, 190. t. 18. f. 3 = sericeus.
Forskolei, Spreng. Syst. i. 596 = Ipomoea verbellacea.
Frankeanus, Schlecht. in Linnaea, vi. (1831) 613 = Jacquemontia mexicanoides.
frondosus, Willd. ex Roem. & Schult. Syst. iv. 303 = havanensis.
fruticosus, Mill. Gard. Dict. ed. VIII. n. 13 = Ipomoea operculata?
frutescens, Pall. ex Ledeb. Fl. Ross. iii. 33 = fruticosus.
frutescens, Pall. Reise, ii. 474 = seq.
fruticosus, *Pall. l. c.* ii. 784.—Oricns.
fruticulosus, *Desr. in Lam. Encyc.* iii. 541.—Ins. Canar.
fugacissimus, Hochst. ex Choisy, in DC. Prod. ix. 449 = Evolvulus alsinoides.
fulvatus, Greene, in Bull. Calif. Acad. i. (1885) 208 = *fulgens*, Wall. Cat. n. 1394 = Argyreia fulgens.
galapagensis, *Andersz. in Vet. Acad. Handl. Stockh.* 1852 (1855) 213.—Ins. Galap.
galaticus, *Rost. ex Choisy, in DC. Prod.* ix. 408.—As. Min.
gangeticus, Linn. Cent. Pl. ii. 9; Amoen. Acad. iv. 306 = Evolvulus gangeticus.
gangeticus, Roxb. Hort. Beng. [18]; Fl. Ind. i. 467 = Argyreia tiliaefolia.
gangeticus, Wall. Cat. n. 1340 = Ipomoea staphylina.
Garberi, Chapm. in Coult. Bot. Gaz. iii. (1878) 11 = havanensis.
Gaudichaudii, Choisy, Conv. Or. 68 = Ipomoea Turpethum.
gemellus, Burm. f. Fl. Ind. 46; Vahl, Symb. Bot. iii. 27 = Ipomoea obscura.
generalis, Lehm. Sem. Hort. Hamb. (1826) 17 = erubescens.
geniculatus, *Munby, ex Pomel, Nouv. Mat. Fl. Atl.* 85.—Afr. bor.
genistoides, Jaub. & Spach, Illustr. iv. 107. t. 370 = spinosus.
geranioides, *Phil. in Anal. Univ. Chil.* (1884) reimp. 7.—Chili.
Gerardi, Roem. & Schult. Syst. iv. 294 = linastus.
Germaniceae, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 1.—As. Min.
giganteus, Silva Manso, Enum. Subst. Bras. 18 = Calystegia palmato-pinnata.
glaber, Aubl. Pl. Gui. i. 138 = Ipomoea glabra.
glaber, *Mill. Gard. Dict.* ed. VIII. n. 11.—Ind. occ.
glandulifer, Spreng. Syst. i. 599 = Ipomoea purpurea.
glandulosus, Herb. Buch.-Ham. ex Wall. Cat. n. 2252 = Lettsomia rubens.
glaucescens, H. B. & K. Nov. Gen. et Sp. iii. 101 = Ipomoea dumetorum.
glaucifolius, Choisy, in DC. Prod. ix. 419 = incanus.
glaucifolius, Spreng. Syst. i. 604 = Ipomoea glaucifolia.
glomeratus, *Choisy, in DC. Prod.* ix. 401.—Arab.
glomeratus, *Thunb. in Mus. Nat. Upс.* 168.—Hab.?
glomeratus, Wall. Cat. n. 2257 = Argyreia cymosa.
gonatodes, Stend. ex A. Rich. Tent. Fl. Abyss. ii. 69 = Ipomoea obscura.
gonocladus, Boiss. Diagn. Ser. I. vii. 22 = Hausknechtii.
Gortschakovii, *Schrenk, Enum. Pl. Nov. Song.* 18.—Soongaria.
gossypifolius, H. B. & K. Nov. Gen. et Sp. iii. 107 = Ipomoea tuberosa.
gossypinus, Wall. Cat. n. 1407 = Ipomoea Wightii.
gracilis.—Salisb. Prod. 134 = Argyreia mollis.
gracilis, Spreng. Syst. i. 604 = Ipomoea gracilis.
graminosus, Spreng. l. c. 607 = Ipomoea graminea.
grandiflorus, Desr. in Lam. Encyc. iii. 548 = Ipomoea Tuba.
grandiflorus, Hort. Brit. ex Stend. Nom. ed. II. i. 409 = Argyreia tiliaefolia.
grandiflorus, Jacq. Hort. Vindob. iii. 39 = Ipomoea latifolia.

CONVOLVULUS:—
grandiflorus, Linn. f. Suppl. 136 = Ipomoea grandiflora.
grandiflorus, Wall. Cat. n. 1370 = Ipomoea Bona-nox.
griseus, *Raffn. Act. Sonora.* 177.—Am. bor.
guadaloupensis, *Steud. Nom.* ed. II. i. 409.—Ins. Guadalup.
guianensis, Aubl. Pl. Gui. i. 136 = Jacquemontia guyanensis.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 90 = Ipomoea capillacea.
guineensis, D. Dietr. Syn. Pl. i. 678 = Ipomoea pileata.
Hamiltonii, Spreng. Syst. iv. Cur. Post. 61 = Ipomoea Hamiltonii.
Hardwickii, Spreng. l. c. 60 = Ipomoea calycina.
hastatus, Desr. in Lam. Encyc. iii. 548 = Ipomoea angustifolia.
hastatus, Forsk. Fl. Aegypt. Arab. 308 = arvensis.
hastatus, Heyne, ex Wall. Cat. n. 1854 = Ipomoea sepiaria.
hastatus, Nutt. in Am. Phil. Trans. N. S. v. (1837) 194 = incanus.
hastatus, Sieber, ex Choisy, in DC. Prod. ix. 413 = fatmensis.
hastatus, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
hastifolius, Poir. Encyc. Suppl. iii. 467 = arvensis.
hastigerus, Spreng. Syst. i. 594 = Ipomoea coccinea.
Hausknechtii, *Boiss. Fl. Orient.* iv. 109.—Persia.
havanensis, *Jacq. Obs.* iv. 6.—Ind. occ.
hawanensis, Willd. ex Steud. Nom. ed. II. i. 409 = Ipomoea umbellata.
hederaceus, Blanco, Fl. Filip. ed. I. 90 = Hewittia bicolor.
hederaceus, Linn. Sp. Pl. 154 = Ipomoea hederacea.
hederaceus, Mill. Gard. Dict. ed. VIII. n. 17 = Ipomoea commutata.
hederaefolius, Salisb. Prod. 128 = Ipomoea hederacea.
helianthemifolius, *Poir. Encyc. Suppl.* iii. 466.—Am. austr.
hepaticaefolius, Spreng. Syst. i. 594 = Ipomoea Pestigridis.
hepaticaefolius, Willd ex Roem & Schult. Syst. iv. 306 = Ipomoea trifida.
heptaphyllus, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 196 = Ipomoea palmata.
Hermanniae, Choisy, in DC. Prod. ix. 408 = hermannioides.
Hermanniae, *L'Hérit. Stirp. Nov.* 67. t. 33.—Peruv.
hermannioides, *A. Gray, Syn. Fl. N. Am.* ii. i. 216.—Texas.
heterophyllus, Ehrenb. ex Boiss. Fl. Orient. iv. 109 = coelesyriacus.
heterophyllus, Spreng. Syst. i. 592 = Ipomoea heterophylla.
Heynii, Spreng. l. c. 699 = Lettsomia aggregata.
Heynii, Wall. Cat. n. 1387 = Ipomoea sepiaria.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 519.—Afr. trop.
hirsutissimus, Wall. Cat. n. 1400 = Lettsomia hirsutissima.
hirsutus, *Bieb. Fl. Taur. Cauc.* i. 432.—Graecia; As. Min.
hirsutus, Roem. ex Choisy, in DC. Prod. ix. 344 = Ipomoea hederacea.
hirsutus, Roxb. Hort. Beng. 14; Fl. Ind. i. 479 = Ipomoea pentaphylla.
hirsutus, Tenore, Fl. Nap. i. 60. t. 15 = althaeoides.
hirtus, Herb. Madr. ex Wall. Cat. n. 2254 = Ipomoea rumicifolia.
hirtus, *Linn. Sp. Pl.* 159.—Ind. or.
hirtus, Wight, ex Stend. Nom. ed. II. i. 409 = Ipomoea eriocarpa.
Hispaniolae, Spreng. Syst. i. 590 = Ipomoea Hispaniolae.
hispidus, Vahl, Symb. Bot. iii. 29 = Ipomoea eriocarpa.
holosericeus, Mönch, ex Choisy, in DC. Prod. ix. 401 = Anchei.
holosericeus, *Bieb. Fl. Taur. Cauc.* i. 147.—As. Min.
Hookerianus, D. Dietr. Syn. Pl. i. 666 = Ipomoea fistulosa.
Horsfalliae, D. Dietr. l. c. 664 = Ipomoea Horsfalliae.
Houstoni, Stend. Nom. ed. II. i. 408, in syn. = glaber.
Howgiei, De Vriese, in Lehm. Pl. Preiss. i. 346 = orthocaulis.
Humboldtianus, *Roem. & Schult. Syst.* iv. 284.—Venezuela.
humilis, Jacq. Coll. iv. 280. t. 20 = undulatus.
humilis, Sadlisb. Prod. 126 = Hiscidus.
hunteriensis, Walk. Fl. Carol. 84 = Stenosis hunteriana.

CONVOLVULUS :—

hybridus, *Zeil. & Morr. in Nat. Genesch. Arch. Nobrl. Ind.* ii. (1846) 6.—Malaya.
hyoscyamoides, *Vahl, in Linnæa*, xiii. (1839–39) 590.—Afr. trop.
hypocrateriformis, Desr. in Lam. Encyc. iii. 561 = Rivea hypocrateriformis.
hypoleucus, Roem. & Schult. Syst. iv. 884 = Rivea hypocrateriformis.
Hystrix, *Vahl, Symb. Bot.* i. 16.—Aegypt.; Arab.
ianthinus, Hance, ex Walp. Ann. iii. 118 = parviflorus.
ignavus, Schrad. in Goett. Gel. Anz. i. (1821) 716 = Ipomœa flammea.
imbricatus, Spreng. Syst. i. 613 = Lettsomia aggregata.
imperati, Vahl, Symb. Bot. i. 17 = Ipomœa littoralis.
incanus, *Vahl, l. c.* iii. 23.—Am. bor. et austr.
incarnatus, Vahl, Eclog. Am. ii. 13 = Ipomœa incarnata.
incisus, *Herb. Mus. Par. ex Choisy, in DC. Prod.* ix. 409.—Peruv.†
incisus, Spreng. Syst. i. 609 = Ipomœa incisa.
incrassatus, Wall. Cat. n. 1386 = Ipomœa sepiaria.
incurvus, Schum. & Thonn. Beskr. Guin. Pl. 99 = Ipomœa Clappertonii.
indicus, Mill. Gard. Dict. ed. VIII. n. 5 = Ipomœa violacea.
indivisus, Vell. Fl. Flum. 71; ii. t. 50 = Ipomœa coccinea.
inflatus, Desf. Tabl. ed. I. 74 = Calystegia silvatica.
insignis, Spreng. Syst. i. 592 = Ipomœa digitata.
insularis, D. Dietr. Syn. Pl. i. 668 = Ipomœa obscura.
intermedius, Loisel. in Journ. Bot. ii. (1809) 264 (Not. 40) = linostus.
intermedius, Roem. & Schult. Syst. iv. 284 = Ipomœa purpurea.
involucellatus, *Klotzsch, in Peters, Reise Mossamb. Bot.* 241.—Afr. trop.
involucratus, Ker-Gawl. in Bot. Reg. t. 318 = Hewittia bicolor.
involucratus, Spreng. Syst. i. 690 = Ipomœa pilosta.
involucratus, Willd. Sp. Pl. i. 845 = Ipomœa teretistigma.
ipomœa, Vell. Fl. Flum. 72; ii. t. 56 = Ipomœa trichoba.
italicus, Roem. & Schult. Syst. iv. 286 = althæoides.
jacobseus, *C. Sm. in Tuckey, Congo*, 249, nomen.—Ins. Cap. Verd.
Jalapa, Linn. Mant. i. 42 = Ipomœa Jalapa.
jamaicensis, Jacq. Obs. iii. 6 = havanensis†
jamaicensis, Spreng. Syst. i. 595 = Ipomœa jamaicensis.
japonicus, Thunb. Fl. Jap. 85 = Ipomœa angustifolia.
fatiauca, J. F. Gmel. Syst. 239 = Ipomœa Jalapa.
javanensis, Burm. f. Fl. Ind. 48 = Ipomœa denticulata.
javanicus, Burm. f. l. c. 50 = Ipomœa sagittæfolia.
jenamicus, Spreng. Syst. i. 605 = Ipomœa hastata.
jenensis, *Kotschy, in Sitzb. Wien. Acad.* liii. (1866) t. 2.—Arab.
kentrocaulos, Steud. ex Choisy, in DC. Prod. ix. 362 = Ipomœa kentrocaulos.
Kleinii, Spreng. Syst. i. 606 = Ipomœa staphylina.
Kleinii, Wall. Cat. n. 1377 = Ipomœa sinuata.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 837, 336.—Turkestan.
Kotschyanus, *Boiss. Diagn.* Ser. I. vii. 93.—Persia.
Kraussianus, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 332.—Turkestan.
lachnaeus, Spreng. Syst. i. 606 = Ipomœa lachnaea.
lachnosperma, Hochst. in Ind. Sem. Hort. Monsp. (1847) 7 = Ipomœa lachnosperma.
lacinulatus, *Desr. in Lam. Encyc.* iii. 546.—Reg. Argent.
lactescens, Grenier, ex Choisy, in DC. Prod. ix. 483 = Calystegia sepium.
lactens, Wall. in J. Geol. Cat. Bomb. Pl. 133 = Ipomœa lactea.
lacunosus, Spreng. Syst. i. 597 = Ipomœa lacunosa.
lactus, Naked. Prod. 128 = Calystegia sepium.
laetissimus, Willd. ex Roem. & Schult. Syst. iv. 303 = Ipomœa dissecta.
lævigatus, Herb. Mus. Brit. ex Steud. Nom. ed. II. i. 898 = Ipomœa laxigata.
lanatus, Vahl, Symb. Bot. i. 26 = setosus.
lanigus, *Willd. ex Spreng. Syst.* i. 598.—Hab.†
lanuginosus, Alleh. Cat. Punjab Pl. 98 = Allolobium lanuginosum, *Desr. in Lam. Encyc.* iii. 551.—Hispan.

CONVOLVULUS :—

lanuginosus, Heyne, ex Wall. Cat. n. 1407 = Ipomœa Wightii.
lanuginosus, Ledeb. Fl. Ross. iii. 88 = Calvertii.
lanuginosus, Vahl, Symb. Bot. iii. 28 = hirsutus.
lapathifolius, *Schlecht. in Linnæa*, xix. (1847) 786.—Mexic.
lapathifolius, Spreng. Syst. i. 604 = Ipomœa acetoselleæfolia.
lasiantha, *Cav. Ic.* v. 58. t. 479.—Chili.
laxiophlœus, Janb. & Spach, Illustr. iv. 105. t. 368 = lobatus.
laxiosperma, Vis. Pl. Aeg. Nub. 13. t. 1. f. 2 = Ipomœa rumicifolia.
latifolius, Desr. in Lam. Encyc. iii. 561 = Ipomœa ilosa-nox.
laurifolius, Roxb. Hort. Beng. [13]; Fl. Ind. i. 470 = Lettsomia elliptica.
laurifolius, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 186 = Argyreia obtecta.
leiocalycinus, *Boiss. Diagn.* Ser. I. vii. 28.—Persia.
leptocladus, *Boiss. l. c.* 25.—Persia.
leucanthus, Desr. in Lam. Encyc. iii. 541 = Ipomœa leucantha.
leucanthus, Herb Madr. ex Wall. Cat. p. 83, n. 1340 = Argyreia tiliæfolia.
leucophyllus, Willd. ex Roem. & Schult. Syst. iv. 301 = Ipomœa discolor.
leucotrichus, *Pomel, Nouv. Mat. Fl. Atl.* 87.—Afr. bor.
liliacinus, *Boiss. Diagn.* Ser. I. xi. 89.—Syria.
liberalis, Lhotsky, ex Choisy, in Mém. Soc. Phys. Genèv. ix. (1837) 132 = Ipomœa echioides.
lilacinus, D. Dietr. Syn. Pl. i. 669 = Ipomœa lilacina.
limphaticus, Vell. Fl. Flum. 70 = Ipomœa palmata†
Lindleyi, Steud. Nom. ed. II. i. 409 = Ipomœa hederacea.
linearifolius, Mill. Gard. Dict. ed. VIII. n. 28 = pilosellæfolius.
linearis, Curt. Bot. Mag. t. 289 = oleæfolius.
linearis, DC. Fl. Fr. Suppl. 424 = lanuginosus.
linearis, Lam. Fl. Fr. ii. 267 = cantabrica.
linearis, *Linn. Syst. ed.* X. 923.—Europ. austr.; As. temp.
linearis, Sibth. & Sm. Fl. Græc. ii. 81. t. 199 = oleæfolius.
limifolius, D. Dietr. Syn. Pl. i. 675 = Ipomœa linifolia.
linifolius, Linn. Cent. Pl. ii. 10 = Evolvulus alsinoides.
linifolius, Wall. Cat. n. 1389 = Ipomœa angustifolia.
Linkii, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 111. 94 = cœruleus.
littoralis, Linn. Syst. ed. X. 924 = Ipomœa carnea.
littoralis, *Vahl, in Linnæa*, xiii. (1839–39) 519.—Afr. trop.
lividus, Moq. ex Steud. Nom. ed. II. i. 409 = Ipomœa Jalapa.
lobatus, Engelm. & Gray, Pl. Lindheim. i. 44 = incanus.
longiflorus, Heyne, ex Steud. Nom. ed. II. i. 409 = Ipomœa palmata.
longiflorus, Spreng. Syst. i. 595 = Ipomœa grandiflora†
longipes, *S. Wats. in Am. Natural.* vii. (1873) 302.—Am. bor. occ.
Lotuka, Buch.-Ham. ex Wall. Cat. p. 83, n. 1342 = Ipomœa cymosa.
Lourieri, G. Don, Gen. Syst. iv. 290 = calystegioides.
lucanus, Tenore, Fl. Neap. Prod. App. v. 9 = Calystegia sepium.
lupulifolia, Gmel. Noteal. iv. 284 = Ipomœa palmata.
luteolus, *A. Gray, in Proc. Am. Acad.* xi. (1876) 90.—Calif.
luteolus, Spreng. Syst. i. 599 = Ipomœa coccinea.
luteolus, Wight, ex Wall. Cat. p. 84, n. 1349 = Ipomœa chrysoides.
lutus, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 280.—Mexic.
lycioides, Boiss. Diagn. Ser. I. vii. 28 = leiocalycinus.
lycioides, Wall. Cat. n. 1390 = Blinkworthia lycioides.
lymphaticus, Vell. Fl. Flum. ii. t. 47 = Ipomœa palmata.
macranthus, H. B. & K. Nov. Gen. et Sp. iii. 95 = Ipomœa muricoides.
macrocalyx, Ruiz & Pav. Fl. Per. ii. 10 = Ipomœa glabra.
macrocarpus, *Linn. Syst.* ed. X. 923.—Ind. occ.

CONVOLVULUS :—

macrocarpus, Spreng. Syst. i. 592 = Ipomœa tuberosa.
macrorhizus, Ell. Sketch. i. 253 = Ipomœa Jalapa.
macrorhizus, A. Gray, ex Choisy, in DC. Prod. ix. 381 = Ipomœa fastigiata.
macrorhizus, Linn. Syst. ed. X. 923 = Ipomœa Jalapa. macrorrhizus.
macrorrhizus, Heyne, ex Wall. Cat. n. 1350 = Ipomœa paniculata.
macrosiphon, Spreng. Syst. i. 608 = Ipomœa Bona-nox.
macrospermus, Willd. Sp. Pl. i. 860 = Maripa paniculoides.
maroccanus, *Greene, in Bull. Calif. Acad.* i. (1885) 208.—Calif.
Mahur, Buch.-Ham. ex Wall. Cat. p. 85, n. 1422 = arvensis.
major, Gilib. Fl. Lituan. i. 48 = Calystegia sepium.
major, Hort. ex Nichols, Dict. Gard. i. 870 = Ipomœa purpurea.
malabaricus, Linn. Sp. Pl. 155 = Argyreia malabarica.
malabaricus, Roxb. Hort. Beng. [13]; Fl. Ind. i. 469 = Ipomœa staphylina.
malabaricus, Wall. Cat. n. 1414 B = Argyreia Hookeri.
Malcolmi, Roxb. Hort. Beng. [14]; Fl. Ind. i. 474 = arvensis.
malvaceus, *Oliver, in Trans. Linn. Soc.* xxix. (1874) 117.—Afr. trop.
mammosus, Lour. Fl. Cochinch. 108 = Ipomœa mammosa.
marginalis, Buch.-Ham. ex Wall. Cat. p. 83, n. 1340 = Argyreia tiliæfolia.
marginatus, Desr. in Lam. Encyc. iii. 558 = Ipomœa sepiaria.
marginatus, Spreng. Syst. i. 603 = Calystegia margaritata.
maritimus, Desr. in Lam. Encyc. iii. 550 = Ipomœa biloba.
maritimus, D. Dietr. Syn. Pl. i. 680 = Ipomœa littoralis.
maritimus, Gouan, Fl. Monsp. 27 = Calystegia sepium.
maritimus, Lam. Fl. Fr. ii. 265 = Calystegia Soldanella.
maritimus, Pall. ex Ledeb. Fl. Ross. iii. 89 = C. Cantabrica.
Martii, Spreng. Syst. i. 598 = Calystegia heterantha.
martinicensis, Jacq. Select. Am. 26 = Ipomœa martinicensis.
mascatensis, Boiss. Diagn. Ser. I. vii. 25 = virgatus.
Massoni, *A. Distr. Gaert. Nachr.* ii. 377.—Ins. Canar. maxacocarpus, *Spreng. Syst.* i. 606.—N. Granat.
matutinus, *Wend. Bubrag. Ste-Croix*, 271.—Ind. occ.
mauritanicus, *Boiss. Voy. Esp.* 418. t. 122.—Afr. bor.
Maximiliani, Nees, in Pr. Neuwied, Reise, 17 = Calystegia sepium.
maximus, Buch.-Ham. ex Wall. Cat. n. 1341 = Ipomœa staphylina.
maximus, Linn. f. Suppl. 137; Vahl, Symb. Bot. iii. 26 = Ipomœa sepiaria.
Mechoacana, Vitm. Summa Pl. i. 434 = Ipomœa Jalapa.
Medium, Linn. Sp. Pl. 156 = Ipomœa denticulata.
Medium, Roxb. Hort. Beng. 14; Fl. Ind. i. 474 = Ipomœa angustifolia.
megapotamicus, Spreng. ex Meisn. in Mart. Fl. Bras. vii. 259 = Ipomœa megapotamica.
melanostictus, Schlecht. in Linnæa, vi. (1831) 737 = Argyreia tiliæfolia.
memanthus, Hoffmgg. & Link, Fl. Port. i. 369 = tricolor.
menistanus, *Thunb. Mus. Nat Upс.* 168.—Hab.†
Meyeri, *Spreng. Syst.* i. 597.—Hab.†
micranthus, Riddell, Syn. Fl. W. St. 70 = Ipomœa lacunosa.
micranthus, *Roem. & Schult. Syst.* iv. 276.—Ind. occ.; Mexic.
micranthus, Willd ex Spreng. Syst. i. 601 = Sprengelii.
microcalyx, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 230.—Assam.
microcalyx, *Polletan, ex Steud. Nom.* ed. II. i. 410.—Mexic.
microphyllus, *Sieber, ex Spreng. Syst.* i. 611.—Aegypt.
Millerianus, Roem. & Schult. Syst. iv. Index, 831 = Ipomœa umbellata.
minimus, *Asch. Pl. Gui.* i. 141.—Guiana.
minimus, Vitm. Summa Pl. i. 434 = Evolvulus incanus.

CONVOLVULUS:—
minor, Gilib. Fl. Lituan. i. 43 = arvensis.
minor, Host. ex Mill. Gard. Dict. ed. VIII. sub n. 24 = tricolor.
minutiflora, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 269.—Mexic.
modestus, Bois. Diagn. Ser. I. xi. 82 = commutatus.
mollis, Burm. f. Fl. Ind. 44 = Argyreia mollis.
mollis, H. B. & K. Nov. Gen. et Sp. iii. 104 = Ipomoea mollis.
mollis, *Meisn. in Mart. Fl. Bras.* vii. 314.—Bras.
mollissimus, *Bertol. Misc. Bot.* xx. 11. t. 3.—Guatemala.
mollissimus, Pall. ex Choisy, in DC. Prod. ix. 408 = holosericeus.
monospermus, [*Tin.*] *Enech. Pl. Hort. Parm.* (1802) 13.—Hab. ?
montevidensis, *Spreng. Syst.* i. 604.—Reg. Argent.
mucronatus, Benth. Pl. Hartw. 130 = nodiflorus.
mucronatus, Forst. f. Prod. 14 = Ipomoea palmata.
mucronifer, Choisy, in DC. Prod. ix. 414 = Jacquemontia guyanensis.
multibracteatus, Wall. Cat. n. 1408 = Argyreia tomentosa.
multifidus, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
multiflorus, Banks, ex Stend. Nom. ed. II. i. 410 = Ipomoea congesta.
multiflorus, Bess. ex Choisy, in DC. Prod. ix. 407 = arvensis.
multiflorus, H. B. & K. Nov. Gen. et Sp. iii. 100 = Ipomoea tubata.
multiflorus, Mill. Gard. Dict. ed. VIII. n. 15 = Ipomoea multiflora.
multiflorus, Poit. ex Choisy, in DC. Prod. ix. 367 = Ipomoea quinquepartita.
multivalvis, R. Br. Prod. 483 = parviflorus.
munitiflorus, Mart. & Gal. in Bull. Acad. Brux. xii. 11. (1845) 262, sphalm. = minutiflorus.
munitus, Wall. Cat. n. 1354 = Ipomoea quinata, pentaphylla.
muricatus, Linn. Mant. i. 44 = Ipomoea muricata.
mutabilis, Blanco, Fl. Filip. ed. I. 92 = Ipomoea Bona-nox.
mutabilis, Sallab. Prod. 132 = Ipomoea purpurea.
mutabilis, Spreng. Syst. i. 599 = Ipomoea acuminata.
muticus, Stend. Nom. ed. II. i. 410 = Ipomoea grandiflora.
natalensis, *Bernh. in Flora*, xxvii. (1844) ii. 829.—Afr. austr.
Nevi, Spreng. Syst. i. 593 = Ipomoea pedunculatis.
nemorosus, Willd. ex Roem. & Schult. Syst. iv. 303 = Ipomoea pentaphylla.
nervosus, Burm. f. Fl. Ind. 48 = Argyreia speciosa.
nervosus, *Rafin. Fl. Ludov.* 46.—Am. bor.
nevinervis, Ham. Prod. Fl. Ind. Occ. 25 = Jacquemontia hirsuta.
Nil, Linn. Sp. Pl. ed. II. 219 = Ipomoea hederacea.
nitens, C. Koch, in Linnæa, xxii. (1849) 743 = lineatus.
nitidus, *Botan. Mexican.* 65.—Hispan.
nitidus, Desr. in Lam. Encyc. iii. 544 = Argyreia nitida.
nodiflorus, *Desr. l. c.* 557.—Mexic.
nodosanthera, Zipp. ex Span. in Linnæa, xv. (1841) 341 = Ipomoea pamila.
nummularius, Linn. Sp. Pl. 157 = Evolvulus alsinoides.
nummularius, Vahl, Eclog. Am. ii. 18 = Jacquemontia nummularia.
nymphæifolius, D. Dietr. Syn. Pl. i. 670 = Ipomoea peltata.
obcrenatus, Stend. Nom. ed. II. i. 410 = Ipomoea chrysoides.
oblongus, Wall. Cat. n. 1389 = Ipomoea barlerioides.
obscurus, Wall. l. c. n. 1381 = Lettsomia setosa.
obscurus, Linn. Sp. Pl. ed. II. 220 = Ipomoea obscura.
obtectus, Wall. Cat. n. 1416 = Argyreia obtecta.
obtusilobus, Michx. Fl. Bor. Am. i. 139 = Ipomoea commutabilis.
obsullatus, Spreng. Syst. i. 595 = Ipomoea bracteata.
occidentalis, *A. Gray, in Proc. Am. Acad.* xi. (1876) 89.—Am. bor. occ.
occidentalis, S. Wats. l. c. 116 = macrostegia.
ocellatus, *Hook. Bot. Mag.* t. 4065.—Afr. austr.
ochraceus, Lindl. Bot. Reg. t. 1060 = Ipomoea ochracea.
ochraceus, Vahl, ex Choisy, in DC. Prod. ix. 380 = Ipomoea afra.
Oenotheræ, *Vatke, in Linnæa*, xliii. (1880–81) 520.—Afr. trop.
oenotheroides, Linn. f. Suppl. 137 = Ipomoea argyreioides ?

CONVOLVULUS:—
officinalis, Pelletan, ex Stend. Nom. ed. II. i. 410 = Ipomoea Purga.
oleaefolius, *Desr. in Lam. Encyc.* iii. 552.—Graecis.
Olgæ, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 25.—An. cantr.
oligodontus, *Baker, in Journ. Linn. Soc.* xx. (1883) 213.—Madag.
operculatus, Gomes, in Mem. Ac. Lisbos, iii. (1812) Mem. Corresp. 27 = Ipomoea operculata.
orinocensis, Willd. ex Stend. Nom. ed. II. i. 410 = Ipomoea clasoides.
orixensis, *Wall. Cat.* n. 1362 = Lettsomia aggregata.
orixæbotrys, Pelletan, ex Stend. Nom. ed. II. i. 410 = Ipomoea oriæbensis.
ornatus, Wall. Cat. n. 1369 = Rivea ornata.
oronocensis, Willd. ex Roem. & Schult. Syst. iv. 308, vide *orinocensis*.
oxyrensis, Roth, Nov. Pl. Sp. 405 = Lettsomia aggregata.
Oüles, *Willd. ex Fisch. in Cat. Hort. Gorenk.* 28.—Hab. ?
Ottonis, *Meisn. in Mart. Fl. Bras.* vii. 311.—Bras.
ovalifolius, Vahl, Eclog. Am. ii. 16 = Ipomoea ovalifolia.
ovalifolius, West, Bidrag. Ste-Croix, 271 = Ipomoea quinquepartita.
ovatus, Moench, Meth. 450 = sicolus.
ovatus, Reinw. ex Choisy, in DC. Prod. ix. 382 = Argyreia Gutchenotii.
ovariensis, Spreng. Syst. i. 594 = Ipomoea ovariensis.
Owsjanikowii, *Smirnow, ex Trautv. in Act. Hort. Petrop.* ix. (1884) 25, nomen.—Rossia.
oxyphyllus, *Boiss. Diagn.* Ser. I. vi. 26.—Persia.
oxyrepus, *Boiss. l. c.* 28.—Persia.
palæstinus, *Boiss. l. c.* xi. 84.—Palaestin.
pallidus, Sallieb. Prod. 128 = canariensis.
pallidus, Wall. Cat. n. 1418 = Argyreia pallida.
palmatus, Mill. Gard. Dict. ed. VIII. n. 8 = Ipomoea quinquefolia.
palustris, *Cav. Ic.* vi. 80. t. 600. f. 2.—Mexic.
palustris, Herb. Ham. ex Wall. Cat. n. 85, n. 1422 = arvensis.
panduratus, Linn. Sp. Pl. 158 = Ipomoea fastigiata.
panduratus, Lour. Fl. Cochinch. 107 = Ipomoea pandurieformis.
paniculatus, Blanco, Fl. Filip. ed. I. 96 = Ipomoea tuberosa.
paniculatus, Linn. Sp. Pl. 156 = Ipomoea digitata.
paniculatus, Wall. Cat. n. 1350 = Ipomoea kentrocaulos.
pannifolius, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. i. 338.—Ins. Canar.
pannosus, Spreng. Syst. i. 612 = Ipomoea erecta.
Papiria, Spreng. l. c. 595 = Ipomoea pubescens.
paradoxus, Foir. Encyc. Suppl. v. 730 = sagittatus.
parasiticus, H. B. & K. Nov. Gen. et Sp. iii. 169 = Ipomoea parasitica.
parviagrifolius, Bert. ex Spreng. Syst. i. 613 = Ipomoea paeriresifolia.
parnassicus, Boiss. & Orph. Diagn. Ser. II. iii. 125 = cochlearis.
parviflorus, Arn. ex Stend. Nom. ed. II. i. 410 = C. Retzii.
parviflorus, Desr. in Lam. Encyc. iii. 556 = micranthus.
parviflorus, Sallab. Prod. 126 = sicolus.
parvifolius, *Vahl, Symb. Bot.* iii. 29.—Geront. trop.
parvifolius, Spreng. Syst. i. 611 = pluricaulis.
patens, *Clarke, Trav.* i. ed. II. *Add.* p. xxii.—Orient.
patens, Desr. in Lam. Encyc. iii. 547 = Breweria humilstrata.
pauciflorus, Willd. ex Roem. & Schult. Syst. iv. 302 = Ipomoea fastigiata.
paulistanus, Silva Manso, Enum. Subst. Bras. 17 = Ipomoea schiedeana.
pedatus, Forsk. ex Choisy, in DC. Prod. ix. 410 = sicolus.
pedatus, Roxb. Hort. Beng. 14; Fl. Ind. i. 478 = Ipomoea cheiophylla.
pedunculiferta, Boiss. Diagn. Ser. I. xi. 84 = hirsutus.
pedunculosus, Wall. Cat. n. 1417 = Argyreia elliptica.
pedunculatus, Walp. Ann. iii. 112, sphalm. = hirsutus.
pellitus, Lasch. Fl. Alt. i. 282 = Calystegia pellita.
peltatus, Linn. Sp. Pl. 1194 = Ipomoea peltata.
penangiensis, Wall. Cat. n. 1425 = Lettsomia penangiana.
pendulus, Silva Manso, Enum. Subst. Bras. 17 = Ipomoea utilis.

CONVOLVULUS:—
pendulus, Spreng. Syst. i. 590 = Ipomoea palmata.
penicillatus, *A. Rich. Tent. Fl. Abyss.* ii. 74.—Abyssin.
peronatifolius, Sallab. Prod. 134 = Ipomoea Quamoclit.
pennatus, Desr. in Lam. Encyc. iii. 567 = prase.
pennatifidus, Drapiez, Encycl. iii. (1886) = prase.
pensylvanicus, Schrank, in Syll. Ratisb. i. (1824) 197 = spithamaeus.
pentadactylis, Wall. Cat. n. 1367 = Ipomoea quinata.
pentagonus, Roxb. Hort. Beng. 14; Fl. Ind. i. 485 = Ipomoea cymosa.
pentanthus, Jacq. Coll. iv. 210 = Jacquemontia violacea.
pentapetaloides, *Linn. Syst.* ed. XII. 229.—Europ. austr.; As. Min.
pentaphylloides, *Schrank, in Flora*, vii. (1824) ii. *Beil.* 50.—Bras.
pentaphyllus, Linn. Sp. Pl. ed. II. 223 = Ipomoea pentaphylla.
pentaphyllus, Salzm. ex Meisn. in Mart. Fl. Bras. vii. 287 = Ipomoea glabra.
perfoliatus, Schum. & Thonn. Beskr. Guin. Pl. 89 = Ipomoea involucrata.
persicus, *Linn. Sp. Pl.* 158.—As. Min.; Persia.
peruvianus, Spreng. Syst. i. 598 = Ipomoea cuspidata.
Pes-caprae, Linn. Sp. Pl. 159 = Ipomoea Pes-caprae.
Pes-tigridis, Spreng. Syst. i. 602 = Ipomoea Pestigridis.
petiolaris, Kunth, ex Spreng. l. c. 598 = Ipomoea Bona-nox.
petraeus, *Lee, ex Choisy, in DC. Prod* ix. 418.—Afr. austr.
philomega, Vell. Fl. Flum. 74; ii. t. 63 = Ipomoea capparoides.
phoenicus, Prav. ex Meisn. in Mart. Fl. Bras. vii. 214 (= *Operculina pteropus*).—Bras.
phoenicus, Spreng. Syst. i. 596 = Ipomoea coccinea.
physcides, *Presl, Novem. Mat. Fl. Sic.* 294.—Algeria.
Pickeringii, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 130 = Breweria Pickeringii.
pilcatus, Spreng. Syst. iv. 61 = Ipomoea pilcata.
pilifera, *Stend. Nov. ed.* II. i. 410.—Abyssin.
pilosella, *Liebm. Ind. Sem. Hort. Haun.* (1853) 22; in *Ann. Sc. Nat.* Sér. IV. i. (1854) 33.—Hab. ?
pilosellaefolius, *Desr. in Lam. Encyc.* iii. 551.—As. Min.; Persia.
pilosellioides, D. Dietr. Syn. Pl. i. 676 = pilosellaefolius.
pilosus, R. Br. in Salt, Abyss. App. p. lxiv = pilifera.
pilosus, *Rottl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 196.—Ind. or.
pilosus, Roxb. Hort. Beng. 14; Fl. Ind. i. 478 = Ipomoea pilosa.
pilosus, Wickstr. in Vet. Acad. Handb. Stockh. (1837) 60 = Ipomoea clasoides.
pinnatifidus, *H. B. & K. Nov. Gen. et Sp.* iii. 168.—Cuba.
pinnatus, Baker, Fl. Maurit. 212, sphalm. vide pennatus.
Pinto, *Vand. Fasc. Pl.* 9.—Bras.
plantagineus, Choisy, in DC. Prod. ix. 405 = Ipomoea simplex.
platanifolius, Vahl, Symb. Bot. iii. 26 = Ipomoea fastigiata.
platensis, Spreng. Syst. i. 591 = Ipomoea digitata.
platycarpus, Cav. Ic. v. 56 = Bracteila platycarpa.
platypoltis, Zipp. ex Span. in Linnæa, xv. (1841) 338 = Ipomoea campanulata.
platyphyllus, Don, ex Stend. Nom. ed. II. i. 410; ii. 460 = Rivea tiliaefolia.
plebejus, Spreng. Syst. i. 604 = Ipomoea plebeja.
plicatus, *Desr. in Lam. Encyc.* iii. 558.—Afr. austr.
Plumeri, *Spreng. Syst.* i. 603.—Ins. S. Doming.
pluricaulis, *Choisy, Conv. Or.* 95.—Ind. or.
polyanthus, Chan. & Schlecht. in Linnæa, v. (1830) 117 = Jacquemontia violacea.
polyanthus, Wall. Cat. n. 1378 = Ipomoea polyantha, xanthyllina.
polyanthus, H. B. & K. Nov. Gen. et Sp. iii. 98 = mikranthus.
polybrizus, Silva Manso, Enum. Subst. Bras. 16 = Ipomoea polybrizos.
pomeensis, Wall. Cat. n. 1429 = Argyreia pomeensis, sulfyghetta.
portoricensis, Spreng. Syst. i. 605 = Ipomoea sagittifolia.
posduensis, Buch.-Ham. ex Choisy, Convr. Or. 105 = Ipomoea bimbifolia.

CONVOLVULUS:—

Preissii, De Vriese, in Lehm. Pl. Preiss. i. 346 = erubescens.
procumbens, Lund, ex Choisy, in DC. Prod. ix. 352 = Ipomoea polymorpha.
procumbens, Pall. ex Roem. & Schult. Syst. iv. 295 = holosericea.
prolifera, *Vahl, Eclog. Am.* i. 18.—Am. austr.
prolifera, Willd. ex Roem. & Schult. Syst. iv. 302 = Ipomoea tubata.
prostrata, Forsk. Fl. Aegypt. Arab. 203 = pentapetaloides.
prostrata, F. W. Schmidt, Fl. Boëm. ii. 93 = arvensis.
pruinosa, Stend. non ed. II. i. 411 = Jacquemontia glaucescens.
Pseudo-Cantabrica, Schrank, Enum. Pl. Nov. i. 21 = dianthoides.
Pseudo-Scammonia, C. Koch, in Linnaea, xxii. (1849) 746 = C. Scammonia.
Pseudo-siculus, Cav. Descr. 97 = elongatus.
Pseudo-Soldanella, Mer. in App. Loisel. ex Bull. Férussac (1825) 111 = Calystegia Soldanella.
Pseudo-tricolor, Viv. Ann. Bot. i. ii. 184 = tricolor.
pterocarpus, Bert. ex Coll. Hort. Ripul. 87 = Ipomoea martinicensis.
pteygocaulos, Steud. ex Choisy, in DC. Prod. ix. 351 = Ipomoea pteygocaulos.
pubescens, [Soland. in] Russ. Aleppo, ed. II. ii. 246 = hirsutus.
pubescens, Willd. Enum. Hort. Berol. 203 = Ipomoea pubescens.
pudibundus, Lindl. Bot. Reg. t. 999 = Ipomoea cathartica.
puellaris, *Thunb. Mus. Nat. Ups.* 168.—Hab.?
pulchellus, H. B. & K. Nov. Gen. et Sp. iii. 101 = Ipomoea fastigiata.
pulcher, D. Dietr. Syn. Pl. i. 667 = Argyreia Guichenotii.
pulcherrimus, Vell. Fl. Flum. 72; ii. t. 54 = Ipomoea Bona-nox.
pungens, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 709 = Gortschakovii.
punicus, Silva Manso, Enum. Subst. Bras. 18 = Ipomoea pruinosa.
Purga, Wender. in Pharm. Centralb. i. (1880) 457 = Ipomoea Purga.
purpureus, Linn. Sp. Pl. ed. II. 219 = Ipomoea purpurea.
pusillus, Pall. ex Roem. & Schult. Syst. iv. 295 = Calvertii.
pycnanthus, Hochst. ex Choisy, in DC. Prod. ix. 365 Ipomoea capitata.
pyrrhotrichus, Boiss. Diagn. Ser. II. iii. 122 = Hausknechtii.
quadrifolius, *Hochst. in Lorent, Wanderungen*, 235.—Ax. Min.
Quamoclit, Spreng. Syst. i. 591 = Ipomoea Quamoclit.
quinatus, Spreng. l. c. 590 = Ipomoea quinata.
quinquefolius, *Vahl, Symb. Bot.* iii. 31.—Ins. Mascar.
quinquefolius, Linn. Syst. ed. X. 928 = Ipomoea quinquefolia.
quinquelobus, Lindem. in Bull. Soc. Nat. Mosc. xxiii. (1850) ii. 508 = arvensis.
quinquelobus, Vahl, Symb. Bot. iii. 32 = Ipomoea palmata.
quinquenervis, Buch.-Ham. ex Wall. Cat. p. 85, n. 1396 = parviflorus.
quinquepartitus, Vahl, Eclog. Am. ii. 14 = Ipomoea quinquepartita.
racemosus, Roem. & Schult. Syst. iv. 302 = Ipomoea staphylina.
radicans, *Thunb. Fl. Cap.* ii. 15.—Afr. austr.
radiosus, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. iii. 134.—*Graecia.*
ramosissimus, Poir. Encyc. Suppl. iii. 468 = Ipomoea ramosissima.
Rampanis, Buch.-Ham. ex Wall. Cat. n. 2251 = Ipomoea stolonifera.
reflexus, *Pomel, Nouv. Mat. Fl. Atl.* 84.—Afr. bor.
Reinwardtianus, D. Dietr. Syn. Pl. i. 675 = Argyreia Guichenotii.
remotus, R. Br. Prod. 488 = erubescens.
repifornis, Cat. Hort. Cantab. (1851) 76 (Quid?).
repiformis, Poir. Encyc. Suppl. iii. 475 = Calystegia Soldanella.
repiformis, Roxb. Hort. Beng. 34; Fl. Ind. i. 482 = Ipomoea muricata.

CONVOLVULUS:—

repandus, Desr. in Lam. Encyc. iii. 555 = Ipomoea repanda.
repens, Linn. Sp. Pl. 158 = Calystegia sepium.
repens, Sw. Obs. 64 = Ipomoea acetosaefolia.
repens, Vahl, Symb. Bot. i. 17 = Ipomoea equatica.
repens, Willd. Sp. Pl. i. 874 = Ipomoea reptans.
reptans, Linn. Sp. Pl. 158 = Ipomoea aquatica.
reticulatus, Buch.-Ham. ex Stend. Nom. ed. II. i. 411 = Ipomoea petaloidea.
reticulatus, *Choisy, in DC. Prod.* ix. 399.—Syria.
retusus, Colla, App. Hort. Ripul. iii. 31 = Ipomoea bilobata.
Retzii, *Steud. Nom.* ed. II. i. 410.—Ind. or.
Rheedii, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 70 = Ipomoea uniflora.
rhytidosperma, *Hochst. ex Choisy, in DC. Prod.* ix. 405.—Afr. trop.
Riedlei, Choisy, Conv. Or. 68 = Ipomoea Turpethum.
riparius, H. B. & K. Nov. Gen. et Sp. iii. 109 = Ipomoea cissoides.
Robertianus, Spreng. Syst. v. 192 = Ipomoea heterophylla.
roseus, H. B. & K. Nov. Gen. Sp. iii. 108 = Ipomoea digitata.
roseus, Mill. Gard. Dict. ed. VIII. n. 18 = Ipomoea fastigiata.
roseus, Pall. Reise, iii. 317 = lineatus.
rostratus, Zipp. ex Span. in Linnaea, xv. (1841) 339 = Ipomoea aquatica.
Rotkii, Spreng. Syst. i. 600 = Ipomaea cymosa.
Rottleri, Spreng. Syst. iv. Cur. Post. 61 = Argyreia malabarica.
Rottlerianus, *Choisy, Convr. Or.* 95.—Ind. or.
rotundifolius, Schum. & Thonn. Fl. Beskr. Guin. Pl. 102 = Ipomoea biloba.
Roxburghii, Wall. Cat. n. 1415 = Argyreia Roxburghii.
Royeni, Steud. Nom. ed. II. i. 410, in syn. = Ipomoea pentaphylla.
ruber, Wall. Cat. n. 1491 = Lettsomia rubens.
ruber, Vahl, Eclog. Am. ii. 12 = Calystegia setifera.
rubescens, Poir. Encyc. Suppl. iii. 466 = erubescens.
rubicundus, Wall. Cat. n. 1409 = Lettsomia rubicunda.
rubrocoruleus, D. Dietr. Syn. Pl. i. 670 = Ipomoea rubrocoerulea.
ruderatus, H. B. & K. Nov. Gen. et Sp. iii. 96 = havanensis.
rufescens, Choisy, Conv. Or. 97 = flavus.
rugosus, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 198 = Ipomoea Beladamboe.
Ruiwii, Spreng. Syst. i. 594 = Ipomoea villosa.
rupestris, Buch, Reschr. Canar. Ins. 198 = Massoni.
rupestris, Pall. Reise, iii. 723 = Ipomoea sibirica.
rupestris, *Willd. Sp. Pl.* i. 846.—Sibir.
Ruprechti, *Boiss. Fl. Orient.* iv. 96.—Reg. Caucas.
sabatius, Viv. Fl. Lib. Spec. 67 = elongatus.
sagittaefolius, Salisb. Prod. 128 = arvensis.
sagittatus, Dulac, Fl. Hautes-Pyr. 489 = arvensis.
sagittatus, A. Gray, in Syn. Fl. N. Am. ii. i. 484, = hirsutus?
sagittatus, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
sagittifer, H. B. & K. Nov. Gen. et Sp. iii. 100 = Ipomoea umbellata.
sagittifolius, Flesch. ex Choisy, in DC. Prod. ix. 407 = arvensis.
sagittifolius, Micha. Fl. Bor. Am. i. 138 = Ipomoea sagittata.
sagittifolius, *Schlecht. in Linnaea*, xxi. (1848) 747.—Texas.
sagittifolius, Silbth. & Sm. Fl. Graec. Prod. i. 138; Fl. Graec. t. 198 = hirsutus.
salicifolius, Desr. in Lam. Encyc. iii. 548 = Ipomoea martinicensis.
salicifolius, Roxb. ex Steud. Nom. ed. II. i. 411 = Ipomoea salicifolia.
Saltii, *Steud. l. c.* = congestus.
salsifolius, Sieber, ex Link, Enum. Hort. Berol. i. 203 = sepiaria.
sanguineus, Spreng. Syst. i. 594 = Ipomoea hederifolia.
sanguineus, Willd. ex Roem. & Schult. Syst. iv. 302 = Ipomoea superba.
sarcocarpus, Silva Manso, Enum. Subst. Bras. 49 = Ipomoea operculata.
sarmentosus, *Bolf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 83.—Ins. Socotra.

CONVOLVULUS:—

saxothrocladus, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* iv. 93.—Assyria.
saxatilis, Bieb. Fl. Taur. Cauc. i. 146 = Calvertii.
saxatilis, Salisb. Prod. 124 = Massoni.
saxatilis, Vahl, Symb. Bot. iii. 32 = lanuginosus.
scammonia, Wall. ex Stend. Nom. ed. II. i. 411 = stamensis.
Scammonia, *Linn. Sp. Pl.* 153.—An. Min.
Scammonia, Lour. Fl. Cochinch. 106 = calystegioides.
scandens, *Delile, Ind. Sem. Hort. Monsp.* (1847) ?.—Aegypt.
Schiedeanus, D. Dietr. Syn. Pl. i. 670 = Ipomoea Schiedeana.
Schimperi, *Boiss. Diagn.* Ser. I. xi. 61.—Arabia.
schirasianus, *Boiss. l. c.* 95.—Persia.
Schottianus, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 19. —Hab.?
Schultesii, Roem. & Schult. ex Steud. Nom. ed. II. i. 411 = Ipomoea purpurea.
Schultesii, Weinm. ex Stend. l. c. = Hewittia bicolor.
scindicus, Bolss. Diagn. Ser. II. iii. 122 = microphylla.
scindicus, *Stocks, in Hook. Kew Journ. iv.* (1852) 178. —Ind. bor. occ.
scoparius, *Linn. f. Suppl.* 135.—Ins. Teneriff.
scrobiculatus, Lindl. Bot. Reg. t. 1076 = Ipomoea scrobiculata.
secundus, *Desr. in Lam. Encyc.* iii. 553.—Aegypt.; Syria.
secundus, Ruiz & Pav. Fl. Per. ii. 10 = Jacquemontia secunda.
Selloi, *Meisn. in Mart. Fl. Bras.* vii. 314.—Bras.
Semenowii, *Herder, in Bull. Soc. Nat. Mosc.* xii. (1868) t. 74.—Sibir.
semidigynus, Roxb. Hort. Beng. [18]; Fl. Ind. i. 468 = Breweria cordata.
Senegambiae, Spreng. Syst. i. 610 = Ipomoea Senegambiae.
sepiarius, Wall. Cat. n. 1384 = Ipomoea sepiaria.
sepincola, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1880) 128 = Calystegia sepium.
sepium, Linn. Sp. Pl. 153 = Calystegia sepium.
sepium, Pall. Reise, iii. 270 = arvensis.
septangularis, Stend. Nom. ed. II. i. 411 = Ipomoea Batatas.
septenatus, Spreng. Syst. i. 591 = Ipomoea umbellifera.
Ser, Spreng. l. c. 598 = Ipomoea sinensis.
sericeus, *Burm. f. Fl. Ind.* 49. t. 19. f. 3.—Persia.
sericeus, Gholsy, in DC. Prod. ix. 400 = lanatus.
sericeus, Forsk. Fl. Aegypt. Arab. 208 = tenuissimus.
sericeus, Linn. Mant. 43 = Argyreia mollis.
sericophyllus, *T. Anders. in Journ. Linn. Soc.* v. *Suppl.* i. (1860) 25.—Arab.
serotinus, DC. Cat. Hort. Monsp. 97 = Ipomoea serotina.
serpens, Linn. Syst. ed. X. 923 = Ipomoea pentaphylla.
serpiculatus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I, Art. iv. 10.—Malaya!
serpyllifolius, H. B. & K. Nov. Gen. et Sp. iii. 95 = Ipomoea serpyllifolia.
serpyllifolius, *Griseb. Cat. Pl. Cub.* 207.—Cuba.
sessiliflorus, Spreng. Syst. i. 599 = Ipomoea eriocarpa.
setifer, Spreng. l. c. 597 = Calystegia setifera.
setosus, Spreng. l. c. 594 = Argyreia setosa.
setosus, Wall. Cat. n. 1412 = Lettsomia setosa.
Sherardi, Pursh, Fl. Am. Sept. ii. 80 = micranthus.
Shumardianus, *Torr. Bot. Expl. Red River* (1853) 291. —Am. bor.
stamensis, *Wall. Cat.* n. 6868.—Siam.
sibiricus, Linn. Mant. ii. 302 = Ipomoea sibirica.
Sibthorpii, Roem. & Schult. Syst. iv. 285 = hirsutus.
sicolus, *Linn. Sp. Pl.* 156.—Reg. Mediterr.
sidaefolius, H. B. & K. Nov. Gen. et Sp. iii. 98 = Ipomoea tubata.
simplex, Busk, Ind. DC. Prod. iii. 118 = plantagineus.
simplex, Pers. Syn. i. 178 = Ipomoea angustifolia.
simplex, Spreng. Syst. i. 607 = Ipomoea simplex.
sinensis, Desr. in Lam. Encyc. iii. 557 = Ipomoea sinensis.
sinensis, Nees, ex G. Don, Gen. Syst. iv. 276 = Ipomoea Desponna.
sinuatus, Petagna, ex Steud. Nom. ed. II. i. 411 = Ipomoea carnea.
sinuatus, *Schenk, in Linnaea*, xxi. (1848) 748.—Texas.
Sloanei, Spreng. Syst. i. 606 = Ipomoea triloba.

CONVOLVULUS:—
cmilacifolius, Salisb. Prod. 124 = Ipomoea maricata.
cogelianus, Bunge, Reliq. Lehm. Bot. 219 = pilosellaefolius.
solanifolius, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 93 = Madeira.
solanifolius, Spreng. Syst. i. 598 = Ipomoea solanifolia.
Soldanella, Despr. ex Choisy, in DC. Prod. ix. 345 (= *Pharbitis Pravaziana*).—Jap. Cenar.
Soldanella, Linn. Sp. Pl. 159 = Calystegia Soldanella.
somalensis, *Franch. in Rhodl, Cosmati*, 43.—Afr. trop.
somalensis, *Vatke, in Linnaea*, xliii. (1880–82) 517.—Afr. trop.
speciosus, Linn. f. Suppl. 137 = Argyreia speciosa.
speciosus, Wall. Fl. Carol. 93 = Ipomoea sagittata.
spectabilis, Hort. Brit. ex Steud. Nom. ed. II. i. 411 = Rivea tiliacfolia.
sphaerocephalus, Roxb. Hort. Beng. 14; Fl. Ind. i. 472 = eriocarpa.
sphaerostigma, Cav. Ic. v. 54. t. 481 = Jacquemontia hirsuta.
spicaefolius, Desr. in Lam. Encyc. iii. 549 = linearius.
spicaefolius, Buch.-Ham. ex Wall. Cat. p. 58. n. 1816 = pluricaulis.
spinosus, *Burm. f. Fl. Ind.* 47. t. 19. f. 4.—Persia.
spinosus, Desr. in Lam. Encyc. iii. 546 = fruticosus.
spinosus, Forsk. Fl. Aegypt. Arab. p. cvi = Hystrix.
spiralis, Willd. ex Roem. & Schult. Syst. iv. 302 = nodiflorus.
spithamaeus, *Linn. Sp. Pl.* 158.—Am. bor.
splendens, Horsem. Hort. Hafn. Suppl. 128 = Argyreia splendens.
Sprengelii, *Choisy, in DC. Prod.* ix. 416.—Hab. ?
stachydifolius, *Choisy, l. c.* 408.—Syria; Mesopotam.
stans, H. B. & K. Nov. Gen. et Sp. iii. 96 = Ipomoea stans.
stans, Michx. Fl. Bor. Am. i. 136 = spithamaeus.
stellaris, Salisb. Prod. 124 = Hermanniae.
stellata, *Brot. ex Heynh. Nom.* ii. 158, nomen.—Hab. ?
stenophyllus, *Boiss. Fl. Orient.* iv. 106.—Syria.
stipulaceus, Roxb. Hort. Beng. 14; Fl. Ind. i. 484 = Ipomoea sepiaria.
stipulatus, Desr. in Lam. Encyc. iii. 546 = Ipomoea dissecta.
Stockii, *Boiss. Fl. Orient.* iv. 110.—Beluchist.
stolonifernus, Cyrilli, Pl. Rar. Neap. i. 14. t. 5 = Ipomoea carnea.
striatus, Vahl, Symb. Bot. iii. 28 = Ipomoea aspleria.
strictus, Lehm. Ind. Sem. Hort. Hamb. (1852) 17 = undulatus.
strictus, Willd. ex Steud. Nom. ed. II. i. 412 = Ipomoea murucoides.
strigosus, Russell, ex Wall. Cat. n. 2259 = Argyreia Leschenaultii.
strigosus, Spreng. Syst. i. 600 = Lettsomia setosa.
strigosus, Wall. Cat. n. 1365 = Lettsomia strigosa.
strigulosus, Boiss. Diagn. Ser. I. xi. 89 = assyricus.
suaveolens, Mart. & Gal. in Bull. Acad. Brux. xii. 11. (1845) 261 = Ipomoea suaveolens.
subacaulis, Buch.-Ham. ex Wall. Cat. sub n. 2256 = Ipomoea barlerioides.
subangularis, Buch.-Ham. ex Steud. Nom. ed. II. i. 413 = Ipomoea obscura.
subcreus, Willd. Enum. Hort. Berol. Suppl. 10 = Mexicani.
subglobosus, *G. Don, Gen. Syst.* iv. 298.—Cochinch.
subhirsutus, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 338, 339.—Turkestan.
subhastatus, Linn. f. Suppl. 135 = Hewittia bicolor.
subpinnatifidus, De Vriese, in Lehm. Pl. Preiss. i. 347 = erubescens.
subpubescens, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 357 = incanus.
subsericeus, *Schrenk, Enum. Pl. Nov.* i. 19.—Sibir. altaic.
subsessilis, Steud. Nom. ed. II. i. 413 = Jacquemontia subsessilis.
subspathulatus, *Vatke, in Linnaea*, xliii. (1880–82) 520.—Afr. trop.
subulatifolius, *Neronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 10.—Malaya.
subvaginalis, *Steud. Nom.* ed. II. i. 412.—Chili.
subvolubilis, Ledeb. Fl. Alt. i. 225, in note = Calystegia subvolubilis.
sufruticosus, [Dryand. in] Ait. Hort. Kew. ed. II. i. 331 = Masseni.

CONVOLVULUS:—
suffruticosus, *Desf. Fl. Atlant.* i. 175. t. 48.—Afr. bor.
sufjultus, H. B. & K. Nov. Gen. et Sp. iii. 103. t. 211 = Ipomoea suffulta.
sufjuburnus, La Llave, in Nov. Veg. Desc. fasc. i. 17 = Ipomoea convalescentis.
superbus, H. B. & K. Nov. Gen. et Sp. iii. 108 = Ipomoea superba.
supinus, *Cass. & Kral, in Bull. Soc. Bot. Fr.* iv. (1857) 397, 487, nomen.—Algeria.
sylvaticus, Spreng. Syst. i. 602 = Calystegia sylvatica.
sylvestris, Waldst. & Kit. Pl. Rar. Hung. iii. 290 = Calystegia sylvatica.
tamnifolius, G. F. W. Mey. Prim. Fl. Esseq. 95 = Jacquemontia tamnifolia.
tamnifolius, Willd. ex Roem. & Schult. Syst. iv. 301 = Ipomoea tricocarpa.
Tarita, Buch.-Ham. ex Wall. Cat. n. 2253 = Rivea ornata.
tenuifus, Desr. in Lam. Encyc. iii. 559 = Breweria hirsutrata.
tenuifus, Stocks, in Hook. Kew Journ. iv. (1852) 172 = Stocksii.
tenuiflorus, Steud. Nom. ed. II. i. 412 = Jacquemontia parviflora.
tenuifolius, Buch.-Ham. ex Wall. Cat. p. 84, n. 1854 = Ipomoea pentaphylla.
tenuifolius, Heyne, ex Wall. l. c. n. 1858 = Ipomoea palmata.
tenuifolius, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 260.—Mexic.
tenuifolius, Vahl, Symb. Bot. iii. 33 = Ipomoea quinata.
tenuiscapus, Steud. ex Choisy, in DC. Prod. ix. 380 = Ipomoea tenuiscapa.
tenuissimus, *Sibth. & Sm. Fl. Graec. Prod.* i. 134; *Fl. Graec.* t. 195.—Europ. austr. ; Afr. bor.
tenuistachys, Salisb. Prod. 128 = cantabrica.
ternatus, Spreng. Syst. i. 590 = Ipomoea ternata.
terrifolius, Spreng. l. c. 613 = Ipomoea terrifolia.
terrestris, Linn. Sp. Pl. ed. II. 334 = cantabrica.
thalassicus, Steud. Nom. ed. II. i. 412 = Ipomoea littoralis.
Thonningii Schum. & Thonn. Beskr. Guin. Pl. 98 = Ipomoea dissecta.
Thunbergii, *Roem. & Schult. Syst.* iv. 268.—Afr. austr.
tiliaceus, Willd. Enum. Hort. Berol. 208 = Ipomoea fastigiata.
tiliaefolius, Desr. in Lam. Encyc. iii. 544 = Argyreia tiliaefolia.
timornsis, D. Dietr. Syn. Pl. i. 667 = Hewittia bicolor.
tomentosus, Choisy, in DC. Prod. ix. 418 = hirsutus.
tomentosus, *Linn. Sp. Pl.* 156.—Jamaica.
tomentosus, Vell. Fl. Flum. 74; ii. t. 65 = Ipomoea acuminata.
Tourneforti, Sieber, ex Spreng. Syst. i. 611 = oleaefolius.
tragacanthoides, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 31.—Mongolia.
translucens, *Sorm. in Journ. Bot.* vii. (1869) 165.—Mongolia.
trianthus, Spreng. Syst. i. 598 = Ipomoea sepiaria ?
trichocalyx, Schum. & Thonn. Beskr. Guin. Pl. 91 = Ipomoea ochracea.
trichocalyx, Zoll. & Mor. in Nat. en Geneesk. Arch. Ned. Ind. ii. (1845) 6 = Ipomoea pilosa.
trichocarpus, Spreng. Syst. i. 598 = Ipomoea commutata.
trichophorus, Wall. Cat. n. 1433 = Ipomoea cissoides.
trichosanthes, Michx. Fl. Bor. Am. i. 137 = Breweria humistrata.
trichostoma, D. Dietr. Syn. Pl. i. 670 = Argyreia capitata.
tricolor, *Linn. Sp. Pl.* 158.—Europ. austr.
tricornutus, *Neronha, in Verh. Batav. Gen.* v. (1790) ed. I. art. iv. 10.—Malaya.
tridentatus, Linn. Sp. Pl. 157 = Ipomoea tridentata.
trifidus, H. B. & K. Nov. Gen. et Sp. iii. 107 = Ipomoea trifida.
triflorus, Pav. ex Choisy, in DC. Prod. ix. 340 = Ipomoea quinquefolia.
trifloros, *Phil. in Linnaea*, xxxiii. (1864–65) 183.—Chili.
trifurus, Vahl, Symb. Bot. iii. 30 = Ipomoea aspleria.
trigonus, Buch.-Ham. ex Wall. Cat. p. 84, n. 1853 = Ipomoea obscura.

CONVOLVULUS:—
trilobus, Desr. in Lam. Encyc. iii. 564 = Ipomoea triloba.
triloba, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
trinervis, *Thunb. Fl. Jap.* 85.—Japon.
triquetter, Rehm. ex Boiss. Fl. Orient. iv. 94 = Calvertii.
triquetter, Vahl, Symb. Bot. iii. 30 = Ipomoea Turpethum.
trigustrosus, Vell. Fl. Flum. 71; ii. t. 53 = Ipomoea batatoides.
truncatus, *Vell. l. c.* 74; ii. t. 74.—Bras.
Tuba, Schlecht. in Linnaea, vi. (1831) 735 = Ipomoea grandiflora.
tuberculatus, Desr. in Lam. Encyc. iii. 545 = Ipomoea palmata.
tuberifer, Steud. Nom. ed. II. i. 412 = Ipomoea Batatas.
tuberosus, Spreng. Syst. i. 591 = Ipomoea tuberosa.
tuberosus, Vell. Fl. Flum. 73; ii. t. 57 = Ipomoea Batatas.
Tuckeyanus, *Steud. Nom.* ed. II. i. 407. lin. 1.—Afr. trop.
tuguriorum, Forst. f. Prod. 14 = Calystegia tuguriorum.
turbinifolius, *Hort. Bonon. ex Pasq. Cat. Orto Bot Nap.* 81, nomen.—Hab. ?
Turpethum, Linn. Sp. Pl. 155 = Ipomoea Turpethum.
uliginosus, *Beiss. Diagn.* Ser. I. vii. 36.—Arabia.
umbellatus, Linn. Sp. Pl. 155 = Ipomoea umbellata.
umbellatus, Wall. Cat. n. 2329 = Ipomoea cymosa.
undulatus, *Cav. Ic.* iii. 39. t. 277. f. 1.—Reg. Mediterr.
uniflorus, Burm. f. Fl. Ind. 47 = Ipomoea unifora.
unilateralis, Roem. & Schult. Syst. iv. 284 = Jacquemontia secunda.
urbicus, Salzm. ex Choisy, in DC. Prod. ix. 350 = Ipomoea asarifolia.
urceolatus, Spreng. Syst. i. 601 = Ipomoea urceolata.
Vahlii, *Roem. & Schult. Syst.* iv. 260.—Bras.
Vahlii, Spreng. Syst. i. 611 = Ipomoea chryseides.
valentinus, *Cav. Ic.* ii. 65. t. 180. f. 2.—Hispan.
valentinus, Hort. ex Steud. Nom. ed. II. i. 412 = Ipomoea sagittata.
valensuelanus, A. Rich. Fl. Cub. Fanerog. ii. 124 = havanensis.
valerianoides, Blanco, Fl. Filip. ed. I. 90 = Evolvulus alsinoides.
variabilis, Cham. & Schlecht. in Linnaea, v. (1830) 116 = Ipomoea variabilis.
variabilis, Weinm. ex Steud. Nom. ed. II. i. 412 dimactus.
varifolius, Steud. l. c. = Ipomoea hederacea.
varius, Vell. Fl. Flum. 73; ii. t. 61 = Ipomoea Batatas.
velutinus, Spreng. Syst. i. 601 = Ipomoea velutina.
venenatus, *West. Bidr. Sto-Croix*, 371.—Ins. S. Crux.
venosus, Desr. in Lam. Encyc. iii. 566 = Ipomoea venosa.
venosus, Herb. Madr. ex Wall. Cat. p. 84, n. 1853 = Ipomoea palmata.
ventricosus, Bert. ex Colla, Hort. Ripul. 87 = Ipomoea ventricosa.
ventricosus, Silva Manso, Enum. Subst. Bras. 20 = Ipomoea silvana.
venulosus, Ehrenb. ex Boiss. Fl. Orient. iv. 107 = stachydifolius.
venustus, Spreng. Syst. i. 600 = Ipomoea tricolor.
verrucipes, Tenore, Ind. Sem. Hort. Neap. (1840) = Ipomoea verrucipes.
verrucosus, D. Dietr. Syn. Pl. i. 670 = Ipomoea verrucosa.
versicolor, Sallab. Prod. 128 = tricolor.
vertebellatus, Linn. Sp. Pl. ed. II. 220 = Jacquemontia hirsuta.
vestitus, Wall. Cat. n. 1411 = Lettsomia bella ?
villosulus, D. Dietr. Syn. Pl. i. 667 = Ipomoea villosula.
villosus, *A. Gray, in Proc. Am. Acad.* xi. (1876) 90.—Calif.
villosus, Pers. Syn. i. 177 = Jacquemontia tamnifolia.
violaceus, Spreng. Syst. i. 599 = Ipomoea violacea.
violaceus, Vahl, Symb. Bot. iii. 29 = Jacquemontia violacea.
virgatus, *Beiss. Diagn.* Ser. I. vii. 34.—Persia; Arab.
viridis, Roxb. Hort. Beng. 14, nomen = Ipomoea tridentata.
viridiflorus, Burm. f. Fl. Ind. 46 = Ipomoea viridiflora.
viridulus, Zipp. ex Span. in Linnaea, xv. (1841) 640 = Ipomoea palmata.

CONVOLVULUS :—
volubilis, Brouss. ex Link, in Bach, Beschr. Canar. Ins. 145 = Mannoni.
Wallichianus, Spreng. Syst. iv. Cur. Post. 61 = Calystegia hederacea.
walthenioides, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov.* dec. i. 6.—Persia.
Warleri, Vahl, Symb. Bot. ii. 36 = Ipomœa sagittata.
Wightii, Wall. Pl. As. Rar. ii. 55. t. 171 = Ipomœa Wightii.
Willdenowii, Steud. Nom. ed. II. i. 412 = Ipomœa heterophylla.
seylanicus, Hook. ex Steud. l. c. = Argyreia populifolia.
seylanicus, Moon, ex Wall. Cat. n. 1879 = Ipomœa sidæfolia.
Zollingeri, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 180, 192.—Ins. Saraja.

CONYSA, Burm. f. Fl. Ind. 180 (1768) = **Conyza**, Linn. (Compos.).

CONVSTVLUS, Pritz. Ic. Index, ii. 90 (1866) = **Conystylus**, Teijsm. & Binn. (Thymel.).

CONYZA, Linn. Gen. ed. I. 251 (1737); Less. Syn. Comp. 209 (1832). *COMPOSITÆ*, Benth. & Hook. f. ii. 288.
DIMORPHANTHES, Cass. in Bull. Soc. Philom. (1818) 30; et in Dict. Sc. Nat. xiii. 354 (1819).
ENEMIAS, Rafin. Fl. Tellur. ii. 49 (1836).
ESCHENBACHIA, Moench, Meth. 573 (1794).
FIMBRILLARIA, Cass. in Bull. Soc. Philom. (1818) 31; et in Dict. Sc. Nat. xvii. 54 (1820).
FRAGMORA, Rafin. Fl. Tellur. ii. 50 (1836).
LAENNECIA, Cass. in Dict. Sc. Nat. xxv. 91 (1822).
MARESIA, Arbass. Fam. ii. 122 (1768).
MOSTERION, Rafin. Fl. Tellur. ii. 50 (1836).
PINARDIA, Neck. ex Steud. Nom. ed. II. i. 412 (1840).
TURIFERA, DC. ex Steud. l. c. (1840).
WEBBIA, Sch. Bip. in Walp. Rep. ii. 970 (1843).
abbreviata, Wall. Cat. n. 3048 = Vernonia cinerea.
abnutisifolia, DC. in Wight, Contrib. 16 = stricta.
abyssinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 390.—Geront. trop.
acrocephala, *Miq. Fl. Ind. Bat.* ii. 48.—Malaya.
acuminata, Wall. Cat. n. 3084 = Vernonia arborea.
acutata, Miq. Fl. Ind. Bat. ii. 48 = Blumea laciniata.
adenocarpa, *Dals. & Gibs. Bomb. Fl.* 125.—Ind. or.
adnata, *H. B. & K. Nov. Gen. et Sp.* iv. 74.—Mexic.
ægyptiaca, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 188.—Aegypt.
ægyptiaca, Hook. f. & Thoms. ex Fl. Brit. Ind. iii. 354 = Erigeron asteroides.
ægerantoides, *DC. Prod.* v. 385.—Madag.
alata, Buch. Enum. Transyl. iii. 102 = Inula bifrons.
alata, Roxb. Hort. Beng. 61; Fl. Ind. iii. 430 = Laggera alata.
albicans, Wall. Cat. n. 3079 = Vernonia albicans.
albida, Willd. ex Spreng. Syst. iii. 514 = Erigeron bonariensis.
alopecuroides, Lam. Encyc. ii. 98; Sw. Fl. Ind. Occ. iii. 1314 = Pterocaulon virgatum.
altaica, DC. Prod. v. 280 = Aster angustus.
ambigua, DC. Fl. Fr. Suppl. 468 = Erigeron unifolius.
ambigua, Hook. & Arn. in Bot. Beech. Voy. 57 = chilensis.
ambotnica, *Crantz, Inst.* i. 273.—Amboina.
amœna, Link, Enum. Hort. Berol. ii. 232 = Blumea lacera.
amorphoglossa, Boiss. Diagn. Ser. I. vi. 81 = Erigeron amorphoglossus.
amplectens, Wall. Cat. n. 3096 = Blumea amplectens.
amplexicaulis, Lam. Encyc. ii. 84 = Blumea amplectens.
amplexicaulis, Michx. Fl. Bor. Am. ii. 126 = Pluchea bifrons.
amplexicaulis, Wall. Cat. n. 3104 = Blumea bifoliata.
amygdalina, *Lam. Encyc.* ii. 90.—Ins. Mascar.
amygdalina, Spreng. ex DC. Prod. vi. 446 = Blumea densiflora.
anagallidifolia, Wall. Cat. n. 3066 = Blumea bifoliata.
anchonæfolia, *Poir. Encyc. Suppl.* ii. 330.—Ins. Mascar.
andina, *Remy, in C. Gay, Fl. Chil.* iv. 72.—Chili.

CONYZA :—
andryaloides, DC. Prod. v. 377 = Erigeron andryaloides.
angustifolia, Hort. Par. ex DC. l. c. 420 = Baccharis Pingraea.
angustifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 109 = Pluchea bifrons?
angustifolia, *Roxb. Hort. Beng.* 61; *Fl. Ind.* iii. 429.—Reg. Himal.
angustifolia, Steud. Nom. ed. I. 221 = Monarrhenus salicifolius.
angustifolia, Thw. ex Hook. f. Fl. Brit. Ind. iii. 254 = Erigeron linifolius.
anthelmintica, Linn. Sp. Pl. ed. II. 1207 = Vernonia anthelmintica.
appendiculata, Blume, Bijdr. 895 = Blumea balsamifera.
appendiculata, Lam. Encyc. ii. 88 = Senecio appendiculatus.
apurensis, H. B. & K. Nov. Gen. et Sp. iv. 73 = Erigeron apurensis.
arabica, *Willd. Sp. Pl.* iii. 1949.—Arab.
arabidæfolia, *Remy, in C. Gay, Fl. Chil.* iv. 76.—Chili.
arborea, Wall. Cat. n. 3074 = Vernonia arborea.
arborescens, DC. Prod. v. 344, in syn. cum cit. falsa = Commidendron spurium!
arborescens, Linn. Syst. ed. X. 1213 = Vernonia arborescens.
arbutifolia, Lam. Encyc. ii. 92 = Baccharis arbutifolia.
argentea, *Lam. l. c.* 87.—Ins. Mascar.
argentea, Perr. & Lepyr. ex DC. Prod. v. 449 = Laggera aurita.
argentea, Wall. Cat. n. 3014 = Inula Cappa.
argyrea, *Less. in Linnæa*, vi. (1831) 138.—Bras.
argutidens, *Miq. Fl. Ind. Bat.* ii. 58.—Malaya.
aromatica, Wall. Cat. n. 3054 = Blumea aromatica.
artemisiæfolia, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 202.—Peruv.
artemisioides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 111. 68 = Baccharis rufescens.
articulata, Lam. Encyc. ii. 94 = Baccharis articulata.
arvensis, *Phil. in Linnæa*, xxxiii. (1864-65) 143.—Chili.
aspera, *Rory, Vy.* iii. 174 = melastomoides.
aspera, Wall. ex Steud. Nom. ed. II. i. 413 = Vernonia Roxburghii.
asperifolia, *Benth. & Hook. f. Gen.* ii. 284.—Guatemala.
asperula, *Phil. in Linnæa*, xxviii. (1856) 736.—Chili.
asteroides, Desf. Tabl. ed. II. (1815) 114 = Pluchea sebdecurrens.
asteroides, Host, ex Beck, in Denkschr. Acad. Wien, Math. Nat xliv. (1882) 332 = Inula bifrons.
asteroides, Lam. Sp. Pl. 861 = Sericocarpus conyzoides.
asteroides, Wall. Cat. n. 3052 = aegyptiaca.
asteroides, Wall. ex Steud. Nom. ed. II. i. 413 = Chrysopsis gossypina.
attenuata, *DC. Prod.* v. 384.—Madag.
attenuata, Wall. Cat. n. 3020 = Vernonia attenuata.
Aucheri, DC. Prod. vii. 281 = Erigeron Aucheri.
aurea, Perr. ex DC. l. c. v. 68 = Vernonia senegalensis.
auriculata, Moench, Meth. Suppl. 235 = Phagnalon rupestre.
aurita, Heyne, ex DC. Prod. v. 441 = Blumea lacera.
aurita, *Linn. f. Suppl.* 367.—Nutria.
axillaris, *Hornem. ex Willd. Herb. Hort. Berol.* 871. —Afr. occ.
axillaris, Lam. Encyc. ii. 84 = Blumea lacera.
Baccharis, *Mill. Gard. Dict.* ed. VIII. n. 16.—Honduras.
baccharoides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 390 = Pluchea Dioscoridis.
balsamica, Wall. ex DC. Prod. v. 319 = Priadia balsamica.
balsamifera, Linn. Sp. Pl. ed. II. 1206 = Blumea balsamifera.
barbata, Wall. Cat. n. 3099 = Blumea barbata.
bellidifolia, *Baker, in Journ. Bot.* xx. (1882) 169.—Madag.
bellidifolia, Wall. Cat. n. 3048 = Vernonia cinerea.
Bertoniana, *Phil. in Linnæa*, xxviii. (1856) 737.—Chili.
bidentata, Steud. Nom. ed. I. 222 = calocephala.
bifoliata, Cham. & Less. in Linnæa, vi. (1831) 135 = Blumea Wightiana.
bifoliata, Linn. Sp. Pl. 862 = Blumea bifoliata.

CONYZA :—
bifoliata, Walt. Fl. Carol. 204 = Sericocarpus tortifolius.
bifoliolata, DC. Prod. v. 435 = Blumea Wightiana.
bifrons, Linn. Sp. Pl. 861; Gouan, Hort. Reg. 436 = Inula bifrons.
bifrons, Pursh, Fl. Am. Sept. ii. 524 = Pluchea bifrons.
Bilbaoana, Remy, in C. Gay, Fl. Chil. iv. 76 = Erigeron canadensis.
Blanchetii, *Baker, in Mart. Fl. Bras.* vi. 111. 86.—Bras.
blanda, Wall. Cat. n. 3033 = Vernonia blanda.
Bojeri, *DC. Prod.* v. 384.—Madag.
Bovei, DC. in Ann. Sc. Nat. Sér. II. i. (1834) 261 = Erigeron Bovei.
bovina, Buch.-Ham. ex DC. Prod. v. 444 = Blumea oxyodontis.
bracteolata, Rottl. ex DC. l. c. 434 = Blumea bifoliata.
bractealata, Wall. Cat. n. 3036 = Vernonia subsessilis
brasiliensis, Steetz, in Peters, Reise Mossamb. Bot. 455 = Epaltes brasiliensis.
britannica, Rupr. Fl. Ingric. 569 = Inula britannica.
bryoides, Lam. Encyc. ii. 91 = Baccharis bryoides.
burmeana, *Miq. Fl. Ind. Bat.* ii. 44.—Burma.
Buxifoliana, *Remy, in C. Gay, Fl. Chil.* iv. 72.—Chili.
buxifolia, Lam. Encyc. ii. 92 = Baccharis buxifolia.
cæfra, DC. Prod. v. 351 = Blumea laciniata.
calocephala, *Bory, Voy.* ii. 395.—Ins. Borbon.
callosa, *Klatt, in Decken, Reise Ost-Afr.* iii. 76. — Afr. trop.
calycina, Cav. in Ansl. Cienc. Nat. iv. (1801) 87 = Phagnalon rupestre.
camphorata, Pursh, Fl. Am. Sept. ii. 523 = Pluchea camphorata.
canarensis, Willd. Sp. Pl. iii. 1957 = Inula candida
candicans, Wall. Cat. n. 2989 = Leucomeris spectabilis.
candida, Linn. Sp. Pl. 863 = Inula candida.
Candolleana, Boiss. Fl. Orient. iii. 317 = Pluchea Wallichiana.
canescens, Bory, ex DC. Prod. v. 478 (= *Schizogyne sericea*).—Ins. Canar.
canescens, Linn. f. Suppl. 367 = Vernonia pinifolia.
canescens, *Phil. in Linnæa*, xxviii. (1856) 734.—Chili.
canescens, Thunb. Fl. Cap. 655 = Vernonia pinifolia.
capitata, Spreng. Neue Entdeck. ii. 141 = Vanillosmopsis capitata.
Cappa, Buch.-Ham. in D. Don, Prod. Fl. Nep. 176 = Inula Cappa.
caprariæfolia, *Willd. ex Steud. Nom.* ed. II. i. 413.—Hab.?
cardaminæfolia, *H. B. & K. Nov. Gen. et Sp.* iv. 71.—Ecuador.
carolinensis, Jacq. Ic. Rar. iii. t. 585 = Pluchea odorata.
cataphracta, Willd. ex DC. Prod. v. 427 = Baccharis cataphracta.
cayennensis, *Spreng. Syst.* iii. 512.—Guian.
celebica, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 122.—Ins. Celebes.
cernua, Wall. Cat. n. 3012 = Laggera alata.
chenopodiifolia, Lam. Encyc. ii. 85 = Gynura mauritiana.
chenopodioides, DC. Prod. v. 379 = Erigeron bonariensis.
chilensis, *Spreng. Nov. Prov.* 14.—Chili.
chinensis, Lam. Encyc. ii. 88 = Vernonia chinensis.
chinensis, Linn. Sp. Pl. 862 = Blumea chinensis.
chinensis, Pers. Syn. ii. 426 = Lourieria.
Chingoyo, H. B. & K. Nov. Gen. et Sp. iv. 76. t. 398 = Pluchea Chingoyo.
chrysocoma, *Vatke, in Bremen Abh.* ix. (1885) 190.—Madag.
chrysocomoides, Desf. Fl. Atlant. ii. 269. t. 232 = Nolletia chrysocomoides.
chrysocomoides, Desf. Hist. Arb. i. 292 = Felicia angustifolia.
chrysocomoides, Tenore, ex Steud. Nom. ed. II. i. 413 = procumbens.
cichoriifolia, Miq. Fl. Ind. Bat. ii. 47 = Blumea laciniata.
ciliaris, *Spreng. Nov. Prov.* 14.—Hab.?
cinerariæfolia, Boj. ex DC. Prod. v. 345 = Rochonia cinerariæfolia.
cinerascens, Wall. Cat. n. 3062 = Vernonia cinerea.
cinerea, Linn. Sp. Pl. 862 = Vernonia cinerea.
ciliarucnsna, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 316.—Afr. trop.
clathrafolia, Willd. ex Baker, in Mart. Fl. Bras. vi. 11. 100 = Vernonia scabra.

CONYZA:—
collina, *Phil. in Linnaea*, xxxiii. (1864–65) 144.—Chili.
communis, Wall. Cat. n. 3018 = Blumea hieracifolia.
concava, Spreng. ex Steud. Nom. ed. II. i. 418 = Baccharis concava.
congesta, Spreng. ex DC. Prod. v. 333 = japonica.
conica, *Poir. Encyc. Suppl.* ii. 341.—Hab.?
consanguinea, *Phil. in Linnaea*, xxxiii. (1864–65) 142.—Chili.
conspicua, Wall. Cat. n. 3066 = Microglossa albescens.
corcovipifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 70.—Mexic.
Coronopus, Lam. Encyc. ii. 89 = Psiadia Coronopus.
Cortesii, *H. B. & K. Nov. Gen. et Sp.* iv. 75.—Mexic.
corymbosa, *Mill. Gard. Dict. ed.* VIII. n. 7.—Mexic.
corymbosa, Roxb. Hort. Beng. 61; Fl. Ind. iii. 436 = Pluchea indica.
costata, *Herv. in Herv. & Sond. Fl. Cap.* iii. 114.—Afr. austr.
Cotoneaster, Willd. ex Spreng. Syst. iii. 509 = Vernonia Cotoneaster.
Coulteri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 355.—Am. bor. occ.
Coulteri, Rothr. in Wheeler, Rep. 155 = Erigeron subdecurrens.
crispa, Rapr. in Bull. Phys.-Math. Acad. Pétersb. xiv. (1855) 225 = Erigeron linifolius.
crispata, *Vahl, Symb. Bot.* i. 71.—Arabia.
ctenoptera, Kunth, Ind. Sem. Hort. Berol. (1845) 12 = Laggera pterodonta.
cuneata, Boj. ex DC. Prod. v. 345 = Rochonia cuneata.
cuneata, Buch.-Ham. ex DC. l. c. 384 = absinthifolia.
cuneifolia, Lam. Encyc. ii. 91 = Baccharis cuneifolia.
cuneifolia, Rastach. Nom. ed. III. 240 = Commicendrum spurium.
cuneifolia, Wall. Cat. n. 3101 = Blumea lacera.
cuspreæformis, Lam. Encyc. ii. 91; Illustr. t. 697 = Leptolophyllum cupressiforme.
Dampieri, A. Cunn. ex DC. Prod. v. 266 = Olearia axillaris.
dasycoma, Miq. Fl. Ind. Bat. ii. 56 = Blumea densiflora.
decurrens, *Linn. Sp. Pl.* ed. II. 1206.—Ind. or.
densiflora, Heyne, in Wall. Cat. n. 3907 = Blumea densiflora.
dentata, Blanco, Fl. Filip. ed. I. 629 = Inula Cappa.
dentata, *Wild. Sp. Pl. iii.* 1928.—Afr. trop.
dichotoma, Pers. Syn. ii. 428 = Jasonia dichotoma.
diffusa, Poir. Encyc. Suppl. ii. 338 = Gouani.
diffusa, Roxb. Hort. Beng. 61; Fl. Ind. iii. 429 = Blumea membranacea.
dilatata, Steud. Nom. ed. I. 229 = laurifolia.
Discoroïdis, Desf. Tabl. ed. II. 114 = PluchеaDioscoridis.
discolor, Bory, ex DC. Prod. vi. 276 = Senecio appendiculatus.
distichа, Heyne, in Wall. Cat. n. 3024 = viscidula.
distichа, *Spreng. Syst. iii.* 509.—Am. austr.
divaricata, Spreng. l. c. 508 = Vernonia diffusa.
divergens, Wall. Cat. n. 3027 = Vernonia divergens.
diversifolia, Weinm. in Flora, iii. (1820) 611 = chilensis.
dominguensis, Willd. ex DC. Prod. v. 289 = Incens. domgolensis, *Delile, Pl. Afr. Voy. Méred*, i. 43.—Nubia.
dumetorum, *Phil. in Linnaea*, xxxiii. (1864–65) 144.—Chili.
echinoides, A. Rich. Tent. Fl. Abyss. i. 388 = aegyptiaca.
elaeagnifolia, Wall. Cat. n. 3041 = Vernonia elaeagnifolia.
elata, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 14.—Bras.
elegantula, Wall. Cat. n. 3049 = Vernonia cinerea.
elongata, Wall. l. c. n. 3078 = Blumea elongata.
ericoides, Lam. Encyc. ii. 92 = Liebum ericoides.
erigeroides, DC. Prod. v. 273 = Erigeron bonariensis.
eriophora, Wall. Cat. n. 3018 = Inula Cappa.
erosa, Blanco, Fl. Filip. ed. II. 489 = Blumea mannlensis.
esoona, *Miq. Pl. Jungh.* 500.—Java.
eupatorioides, Wall. Cat. n. 3998 = Inula eupatorioides.
excisa, Wall. l. c. n. 3011 = Blumea densiflora.
extensa, Wall. l. c. n. 3016 = Vernonia extensa.
falcata, *Buch.-Ham. ex Spreng. Syst.* ii. 513.—Reg. Himal.
falcigera, Wall. Cat. n. 3046 = Blumea densiflora.
fasciculata, Wall. l. c. n. 3019 = Laggera flava.
fasciculata, Wall. l. c. n. 3094 (ex Steud.) = Blumea glomerata.

CONYZA:—
fastigiata, *Willd. Enum. Hort. Berol. Suppl.* 57.—Afr. bor.
fatuiosa, Roxb. Hort. Beng. [101]; Fl. Ind. iii. 429 = Blumea glomerata.
fecifilia, *DC. Prod.* v. 285.—Madag.
floresiana, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 121.—Malaya.
floribunda, H. B. & K. Nov. Gen. et Sp. iv. 73 = Erigeron bonariensis.
foetida, Lam. Encyc. ii. 85 = Pluchea camphorata.
foetida, Willd. Sp. Pl. iii. 1952 = Nidorella foliosa.
foliolosa, Buch.-Ham. in Wall. Cat. n. 3981 = Pulicaria folliolosa.
foliolosa, Wall. ex DC. Prod. v. 451 = Pluchea indica.
foliolosa, Wight, in Wall. Cat. n. 3092 = Blumea bifoliata.
folliolosa, *Willd. Sp. Pl.* iii. 1919.—Ind. or.
fruticosa, Linn. Sp. Pl. ed. II. 1209 = Vernonia rigida.
Garnieri, *Klatt, in Linnaea*, xxxvii. (1871–73) 566.—Madag.
Gayana, *Phil. i. c.* xxvii. (1856) 737.—Chili.
geminiflora, Tenore, Cat. Hort. Neap. (1819) 75 = Phagnalon rupestre.
genistelloides, Lam. Encyc. ii. 93 = Baccharis genistelloides.
glabra, Willd. Sp. Pl. iii. 1940 = Vernonia Vahliana.
glabrata, Wall. ex Steud. Nom. ed. II. i. 414 = Eupatorium glabrum.
glomerata, Link, Enum. Hort. Berol. ii. 322 = decurrens.
glomerata, Wall. Cat. n. 3055 = Blumea glomerata.
glutinosa, Lam. Encyc. ii. 89 = Psiadia glutinosa.
gnaphaloides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 389 = Gouani.
gnaphaloides, *H. B. & K. Nov. Gen. et Sp.* iv. 73. t. 337.—Mexic.
gnaphaloides, Sieber, Reise Kreta, i. 1853; ii. 322 = Senecio gnaphaloides.
Gouani, Blanco, Fl. Filip. ed. I. 627 = Blumea mannlensis.
Gouani, *Willd. Sp. Pl.* iii. 1928.—Ins. Canar.
gracilis, Heyne, in Wall. Cat. n. 3064 = Blumea membranacea.
gracilis, Hoffmgg. & Link, Fl. Port. ii. 253 = Pulicaria gracilis.
gracilis, Wall. Cat. n. 3064 = Blumea glomerata.
graeca, Sm. Sym. Sull. 16 = Phagnalon graecum.
graminifolia, Spreng. Syst. iii. 515 = Aster divaricatus.
Grahami, *DC. Prod.* v. 280.—Reg. Argent.
grandiflora, Wall. Cat. n. 3081 = Blumea densiflora.
graveolens, Steud. Nom. ed. I. 222 = Inula graveolens.
graveolens, Wall. ex DC. Prod. v. 292 = Erigeron asteroides.
guineensis, Willd. Sp. Pl. iii. 1980 = Laggera aurita.
gummifera, Roxb. in Beatn. St. Hel. Tracts, App. 804 = Commicendrum robustum.
halimifolia, Desf. Tabl. ed. II. 114 = Baccharis halimifolia.
Hamiltoni, Steud. Nom. ed. II. i. 414 = Blumea hieracifolia.
heterophylla, Lam. Encyc. ii. 90 = Monarrhenus rufescens.
heterophylla, Lam. Encyc. ii. 84 = Vernonia cinerea.
heterotricha, Wall. Cat. n. 3235 = Inula eupatorioides.
Hemdelotii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 317.—Afr. trop.
Heynеana, Wall. Cat. n. 3069 = Blumea lacera.
hieracifolia, Spreng. Syst. iii. 514 = Blumea hieracifolia.
hirsuta, Comm. ex DC. Prod. v. 458 = Cylindrocline Commersonii.
hirsuta, Linn. Sp. Pl. i. 863 = Pluchea hirsuta.
hirsuta, Wall. ex DC. Prod. v. 292; et Cat. n. 3007 B, partim = aegyptiaca.
hirtella, *DC. Prod.* v. 285.—Madag.
hispida, *H. B. & K. Nov. Gen. et Sp.* iv. 71.—Mexic.
Hochstetteri, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 387.—Abyssin.
holcoricus, Wall. Cat. n. 3108 = Blumea glomerata.
Huberi, *Boj. ex Sweet, Hort. Brit.* ed. III. 349.—Madag.
hamiltonia, *Miq. Fl. Ind. Bat.* ii. 41.—Java.
hymenophylla, Wall. Cat. n. 3085 = Blumea membranacea.
hypoleuca, *A. Rich. Tent. Fl. Abyss.* i. 391.—Abyssin.
Illinois, *Tenore, in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 249.—Am. centr.

CONYZA:—
incana, Heyne, ex DC. Prod. v. 24 = Vernonia cinerea?
incana, Wall. Cat. n. 3046 = Vernonia cinerea.
incana, *Willd. Sp. Pl.* iii. 1987.—Arabia.
incisa, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 184.—Afr. austr.
indica, Blume, ex DC. Prod. v. 451 = Pluchea indica.
indica, Wall. Cat. n. 3042 = Vernonia indica.
intermedia, Lag. Gen. et Sp. Nov. 38 = Phagnalon saxatile.
inuloides, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 182 = Jasonia dichotoma.
ivaefolia, *Burm. f. Fl. Ind.* 180.—Ind. or.
ivaefolia, *Less. in Linnaea*, vi. (1831) 138.—Afr austr.
japonica, *Less. Syn. Comp.* 204.—Reg. Himal.; China; Japon.
javanica, Blume, Bijdr. 897 = Blumea lacera.
Jerdoni, C. B. Clarke, Comp. Ind. 62 = Erigeron asteroides.
Junghuhniana, *Miq. Fl. Ind. Bat.* ii. 55.—Java.
Kotschyi, Boiss. Diagn. Ser. I. vi. 85 = Erigeron Aucheri.
Kotschyi, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop 307 = Vicoa auriculata.
Korthalsiana, *Miq. Fl. Ind. Bat.* ii. 51.—Malaya.
Kraussеana, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 616.—Turkestan.
Kraussii, *Sch. Bip. ex Walp. Rep.* ii. 972.—Afr. austr.
Kunthiana, *DC. Prod.* v. 279.—Ecuador.
lacera, Burm. f. Fl. Ind. 180 = Blumea lacera.
lacinista, Roxb. Hort. Beng. 61; Fl. Ind. iii. 437 = Blumea laciniata.
lactucaefolia, Wall. Cat. n. 3088 = Blumea Wightiana
laeta, Hort. Calc. ex C. B. Clarke, Comp. Ind. 14 = Vernonia saliga.
lanceolaria, Roxb. Hort. Beng. 61; Fl. Ind. iii. 432 = Blumea myriocephala.
lanceolata, Wall. l. c. n. 2991 = Pluchea lanceolata.
lanceolata, Wall. Cat. n. 3059 = Vernonia divergens.
lanuginosa, Wall. l. c. n. 2923 = Inula Cappa.
Larrañiana, *Remy, in C. Gay, Fl. Chil.* iv. 71.—Chili.
laurifolia, *Lam. Encyc.* ii. 89.—Ins. Mascar.
laxa, *Spreng. Syst.* iii. 518.—Bras.
laxiflora, *Desf. Cat. Hort. Par.* ed. III. 400.—Hab.?
leucanthema, Buch.-Ham. ex Steud. Nom. ed. II. i. 414 = Blumea leucanthema.
leucodasys, *Miq. in Journ. Bot. Néerl.* i. (1860) 108.—Malaya.
leucophylla, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 387 = incana.
limonifolia, Sibth. & Sm. Fl. Graec. Prod. ii. 174 = Inula candida.
linearifolia, Lam. Encyc. ii. 92 = Psiadia linearifolia.
linearifolia, DC. Prod. v. 285 = aegyptiaca.
linearis, *DC. l. c.* 278.—Chili.
linifolia, *Burm. f. Prod. Fl. Cap.* 26.—Afr. austr.
linifolia, Linn. Sp. Pl. 861 = Sericocarpus solidagineus.
linifolia, *Phil. in Linnaea*, xxviii. (1856) 784.—Chili.
linifolia, Wall. Cat. n. 3069 = Vernonia cinerea.
lithospermifolia, *Lam. Encyc.* ii. 90.—Ins. Mascar.
lobata, Juss. ex Steud. Nom. ed. II. i. 414, 521 = Verbesina diversifolia.
lobata, Lam. Sp. Pl. 863 = Neurolaena lobata.
longicaulis, Wall. Cat. n. 3078 = Eupatorium Reevesii.
longifolia, Desf. Cat. Hort. Par. ed. III. 164 = chilensis.
longifolia, Heyne, in Wall. Cat. n. 3026 = Blumea spectabilis.
longipina, Hook. f. Fl. Brit. Ind. iii. 268, sphalm. = ses.
longissima, Zoll. & Moritzi, Syst. Verz. Zoll. 58 = Blumea chinensis.
Lorentzii, *Griseb. in Gott. Abh.* xxiv. (1879) 176.—Reg. Argent.
Lorentzii, *Poir. Encyc. Suppl.* ii. 338.—China.
lucens, *Pohr. l. c.* 341.—Ins. S. Doming.
lucida, Spreng. Neue Entdeck. ii. 143 = Piptocarpha lucida.
lucida, *Y. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 196.—Ins. Madeira.
lyoopsoloides, Lam. Encyc. ii. 93 = Erichrix junciformis.
lyrata, *H. B. & K. Nov. Gen. et Sp.* iv. 70.—Ecuador.
macrocephala, *Spreng. Syst.* iii. 512.—Reg. Argent.
macrolepis, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 387.—Abyssin.

CONYZA :—
macrostachya, Wall. Cat. n. 8063 = Blumea hieracifolia.
madagascariensis, Lam. Encyc. ii. 85 = Paladia madagascariensis.
magellanica, Lam. i. c. 91 = Baccharis magellanica.
manillensis, Less. in Linnæa, vi. (1831) 135 = Blumea manillensis.
marginata, Edgew. in Trans. Linn. Soc. xx. (1851) 67 = rufa.
marylandica, Michx. Fl. Bor. Am. ii. 126 = Pluchea camphorata.
mauritiana, *Bcj. ex Sweet, Hort. Brit.* ed. III. 349.—Ins. Maurit.
mauritiana, Mérat, ex Steud. Nom. ed. II. i. 414 = Nidorella compressa.
mauritiana, Willd. ex Steud. l. c. = Nolletia chrysocomoides.
maxima, *Zoll. & Mor. in Nat. en Genesh. Arch. Nederl. Ind.* ii. (1845) 236.—Java.
megapotamica, Spreng. Syst. iii. 516 = Stenachaenium megapotamicum.
melastomoides, *Lam. Encyc.* ii. 90.—Ins. Mascar.
membranacea, Wall. ex DC. Prod. v. 441, Cat. n. 8019 = Blumea membranacea.
membranifolia, *Klatt, in Leopoldina*, xx. (1884) 94.—Bras.
micrantha, Spreng. ex DC. Prod. v. 384 = stricta.
microcephala, *Hemsl. Biol. Centr. Am. Bot.* 126.—Mexic.
modatensis, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 144 = Pluchea Dioscoridis.
modesta, *Phil. in Linnæa*, xxxiii. (1864–65) 142.—Chili.
modesta, *Kunth, Ind. Sem. Hort. Berol.* (1845) 11.—Hab. ?
molitis, Willd. Sp. Pl. iii. 1924 = Vernonia cinerea.
moluccana, *Gaudich. Bot. Voy. Bonite*, 485.—Ins. Molucc.
monocephala, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 122.—Malaya.
montana, Heyne, ex DC. Prod. v. 384 = stricta.
montevidensis, *Spreng. Syst.* iii. 510.—Reg. Argent.
multicaulis, DC. Prod. v. 383 = japonica.
multiflora, Wall. Cat. n. 8022 = Vernonia divergens.
murralis, Buch.-Ham. in Wall. Cat. n. 8077 = Blumea membranacea.
Murrea, Buch.-Ham. l. c. n. 8095 = Blumea lacera.
myosotifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 69.—N. Granat.
myricephalos, Remy, in C. Gay, Fl. Chil. iv. 76 = Erigeron canadensis.
myrsinites, Lam. Encyc. ii. 92 = Baccharis myrsinites.
nana, *Sch. Bip. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 315.—Abyss.
napifolia, Wall. Cat. n. 8076 = Blumea lacera.
natalensis, *Sch. Bip. ex Walp. Rep.* ii. 971.—Afr. austr.
natans, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 263 = Blumea hieracifolia.
Nandini, *Bennet, in Bull. Soc. Bot. Fr.* xxv. (1878) 208.—Hab. ?
necessaria, Buch.-Ham. ex DC. Prod. v. 439 = Blumea hieracifolia.
nereifolia, L'Hérit. ex DC. l. c. 430 = Brachylaena nereifolia.
Neuri, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 317.—Afr. trop.
nitida, Miq. Fl. Ind. Bat. ii. 54 = Blumea Wallichii.
nitida, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 121.—Malaya.
nivalis, Boiss. Diagn. Ser. I. vi. 83 = Erigeron chionophilus.
notobullidiastrum, *Griseb. in Goett. Abh.* xxiv. (1879) 177.—Reg. Argent.
nudans, Blume, Bijdr. 896 = Laggera alata.
nudans, Buch.-Ham. ex DC. Prod. v. 436 = Blumea cernua.
nutans, Wall. Cat. n. 8080 = Blumea hieracifolia.
obliqua, Buch.-Ham. ex DC. Prod. v. 439 = Blumea lacera.
obliqua, Willd. Sp. Pl. iii. 2880 = Blumea axillaceps.
obovata, Wall. ex DC. Prod. v. 449 = Inula Cappa.
oblongifolia, *Spreng. Syst. iii.* 513.—Bras.
obovata, Wall. Cat. n. 8083 = Blumea obovata.
obovata, *DC. Prod.* v. 387.—Ann. austr.
obtusa, *H. B. & K. Nov. Gen. et Sp.* iv. 71.—Mexic.

CONYZA :—
odontophylla, Boiss. Fl. Orient. iii. 217 = Pluchea arguta.
odontoptera, *Webb, in Hook. Niger Fl.* 135.—Ins. Cap. Verd.
odora, *Forsk. Fl. Aegypt. Arab.* 148.—Aegypt
odorata, Linn. S.jst. ed. X. 1218 = Pluchea odorata.
odorata, Willd. ex Steud. Nom. ed. II. i. 414 = Vernonia chinensis.
oleaefolia, Lam. Encyc. ii. 86 = Linosyris villosa.
oligandra, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 1. 179.—Java.
oligocephala, Miq. Fl. Ind. Bat. ii. 41 = Blumea bifoliata.
orientalis, *Willd. Sp. Pl.* iii. 1931.—Armen.
ovata, Wall. Cat. n. 8097 = Blumea cinerea.
oxyodonta, Wall. ex DC. Prod. v. 444 = Blumea oxyodonta.
pallida, *Zipp. ex Span. in Linnæa*, xv. (1841) 822 nomen.—Malaya.
panamensis, *Willd. Sp. Pl.* iii. 1942.—Panama.
paniculata, Desf. Tabl. ed. II. 114 = fastigiata.
paniculata, Wall. Cat. n. 8090 = Blumea Wightiana.
paniculata, Willd. Sp. Pl. iii. 1920 = Blumea membranacea.
pannosa, *Webb, in Hook. Niger Fl.* 135.—Afr. trop.
parviflora, Wall. Cat. n. 8004 = Blumea Wightiana.
patula, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 184 = Vernonia chinensis.
pauciflora, Willd. Sp. Pl. iii. 1927 = Vernonia pauciflora.
paucifolia, Wall. Cat. n. 8037 = Blumea membranacea.
pectinata, *Sch. Bip. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 315.—Afr. trop.
pedunculata, *Mill. Gard. Dict.* ed. VIII. n. 15.—Hab. ?
persica, Boiss. Diagn. Ser. I. vi. 81 = Erigeron persicus.
persicifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 312.—Afr. trop.
petiolaris, Wall. Cat. n. 8382 = japonica.
pinifolia, Bory, Voy. ii. 326 = Faujasia pinifolia.
pinifolia, Lam. Encyc. ii. 86 = Vernonia pinifolia.
pinnatifida, Buch.-Ham. ex Roxb. Hort. Beng. 61; Fl. Ind. iii. 430 = stricta.
pinnatifida, *Less. Syn. Comp.* 204.—Afr. austr.
pinnatilobata, *DC. Prod.* v. 387.—Afr. austr.
platensis, Spreng. Syst. iii. 509 = Vernonia platensis.
platylepis, Spreng. ex DC. Prod. v. 81 = Vernonia floribunda.
podocephala, *DC. l. c.* 387.—Afr. austr.
polycephala, Edgew. in Trans. Linn. Soc. xx. (1851) 68 = viscidula.
polygonata, Wall. Cat. n. 8010 = Inula polygonata.
Ponæa, Sieber, ex Tausch, in Flora, xii. (1829) 1. Erg. 87 = Cylindrocline Commersonii.
populifolia, Lam. Encyc. ii. 87 = Vernonia populifolia.
portoricensis, Bert. ex DC. Prod. v. 52 = Vernonia remotiflora ?
prunnea, *Phil. in Linnæa*, xxviii. (1856) 786.—Chili.
procera, Desf. Cat. Hort. Par. ed. III. 164 = chilensis.
procera, Wall. Cat. n. 8060 = Blumea procera.
procumbens, *Balb. Cat. Hort. Taur.* 36.—Ins. Creta.
proiifera, Lam. Encyc. ii. 84 = Vernonia cinerea.
prostrata, Tenore, Cat. Hort. Neap. (1819) 75 = procumbens.
protistiformis, Perr. ex DC. Prod. v. 376 = Pluchea lanceolata.
pterocaula, *C. Bolle, in Bonplandia*, vii. (1859) 295.—Ins. Canar.
pubescens, DC. Prod. v. 381 = Brachyactis robusta.
pubigera, *Linn. Mant.* i. 118.—Amboina.
pulchella, H. B. & K. Nov. Gen. et Sp. iv. 78 = sophiaefolia.
pulchra, Lam. ex Steud. Nom. ed. II. i. 415 = scorpioides.
pumila, *Lam. ex Pers. Syn.* ii. 427.—Hab. ?
pumila, Sibth. & Sm. Fl. Græc. Prod. ii. 173 = Phagnalon pumilum.
punctata, Willd. Sp. Pl. iii. 1937 = Aplopappus Chamissonis.
punctulata, Wall. Cat. n. 8095 = Vernonia chinensis.
punduana, Wall. l. c. n. 8085 = Vernonia volkameriæfolia.
pungens, Lam. Encyc. ii. 86 = Iphiona juniperifolia.
purpurascens, Sw. Prod. Veg. Ind. Occ. 112 = Pluchea purpurascens.
purpurea, Buch.-Ham. in Wall. Cat. n. 8076 = Blumea glomerata.

CONYZA :—
pusilla, *Houtt. Handlexid.* x. 616. t. 69.—Afr. austr.
pusilla, H. B. & K. Nov. Gen. et Sp. iv. 69 = Kunthiana.
pycnostachya, Michx. Fl. Bor. Am. ii. 126 = Pterocaulon pycnostachyum.
pygmæa, Sieber, Reise Kreta, ii. 322 = Phagnalon pumilum.
pyrifolia, Lam. Encyc. ii. 89 = Microglossa volubilis.
pyrrhopappa, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 389.—Abyss.
racemosa, Spreng. Syst. iii. 508 = Baccharis aviceniæfolia.
racemosa, Wall. Cat. n. 8092 = Blumea glomerata.
ramiflora, Spreng. Syst. iii. 509 = Piptocarpha ramiflora.
redolens, Willd. Sp. Pl. iii. 1951 (= *Monenteles redolens*).—N. Caled.
repanda, Roxb. Hort. Beng. 61; Fl. Ind. iii. 431 = Blumea procera.
retusa, *Lam. Encyc.* ii. 90.—Ins. Mascar.
rigida, Sw. Prod. Veg. Ind. Occ. 113 = Vernonia rigida.
riparia, Blume, Bijdr. 899 = Blumea chinensis.
riparia, H. B. & K. Nov. Gen. et Sp. iv. 76 = Tessaria integrifolia.
rivularis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 124.—Bras.
robusta, Roxb. in Beats. St. Hel. Tracts, App. 804 = Commidendrum robustum.
Roylei, DC. Prod. v. 381 = Brachyactis menthodora.
rubra, Buch.-Ham. ex DC. l. c. 376 = Pluchea lanceolata.
rudis, Willd. ex Steud. Nom. ed. II. i. 415 = lyrata.
rufa, *Wall. Cat.* n. 8079.—Ind. or.
rufescens, Hoffmgg. & Link, Fl. Port. ii. 253 = Pallasia rufescens.
rufidula, *Spreng. Neue Entdeck.* ii. 141.—Bras.
rugosa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 184 = Commidendrum rugosum.
rugosa, Vahl, Symb. Bot. i. 71 = Pterocaulon virgatum.
runcinata, Wall. Cat. n. 8087 = Blumea laciniata.
ruscifolia, Linn. Mant. 113 = Phagnalon rupestre.
rutilans, Poir. Encyc. Suppl. ii. 841 = Vernonia senegalensis.
sagittalis, Lam. Encyc. ii. 94 = Baccharis sagittalis.
salicifolia, Lam. Encyc. ii. 89 = Monarrhenus salicifolius.
salicifolia, *Mill. Gard. Dict.* ed. VIII. n. 6.—Ind. occ.
salicina, *Kellogg, in Proc. Calif. Acad.* iii. (1868) 36.—Calif.
salicina, Rapr. Fl. Ingric. 568 = Inula salicina.
saligna, Wall. Cat. n. 8061 = Vernonia saligna.
salsoloides, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 197 = Inula ammophila.
salvifolia, Bory, Voy. iii. 176 = Blumea salvifolia.
sanctilis, Linn. Sp. Pl ed. II. 1206 = Phagnalon saxatile.
saxatilis, Sibth. & Sm. Fl. Græc. t. 862 = Phagnalon græcum.
saxatilis, Tenore, Fl. Neap. ii. 213 = Phagnalon rupestre.
saxatilis, Zoll. ex C. B. Clarke, Comp. Ind. 90 = Blumea balsamifera.
scabiosæfolia, *Remy, in C. Gay, Fl. Chil.* iv. 72.—Chili.
scabra, *Linn. Mant.* 113.—Ind. or.
scabrida, *DC. Prod.* v. 387.—Afr. austr.
scandens, *Mill. Gard. Dict.* ed. VIII. n. 11.—Mexic.
scandens, Wall. Cat. n. 8060 = Vernonia scandens.
Schimperi, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 386 = stricta.
Schlechtendalii, *C. Bolle, in Bonplandia*, vii. (1859) 294.—Ins. Canar.
scorpioides, Lam. Encyc. ii. 88 = Vernonia scorpioides.
semipinnatifida, *Wall. Cat.* n. 8058.—Ind. or.
seminestita, Wall. l. c. n. 8096 = Blumea procera.
senegalensis, Sieber, ex DC. Prod. v. 449 = Laggera aurita.
senegalensis, *Willd. Sp. Pl.* iii. 1928.—Afr. trop.
sericea, *Bory, Voy.* ii. 376.—Ins. Borbon.
sericea, (Soland. in] Ait. Hort. Kew. ed. I. iii. 183 (= *Schizogyne sericea*).—Ins. Canar.
sericea, Wall. Cat. n. 8021 = Blumea glomerata.
serici-nestita, Wall. l. c. n. 8086 = Blumea chinensis.
serpentaria, *Griseb. in Goett. Abh.* xxiv. (1879) 176.—Reg. Argent.

CONYZA:—
arvensis, Bory, ex DC. Prod. vi. 376 = Senecio salicifolius.
serrulata, *C. B. Clarke, Comp. Ind.* 64.—Ind. or.
serrulata, Lam. Encyc. ii. 55 = Baccharis serrulata.
serrulata, Wall. Cat. sub n. 3094 = vicidula.
sessiflora, Blume, Bijdr. 897 = Pluchea sessiflora.
sessiliflora, *Mig. Fl. Ind. Bat.* ii. 48.—Ins. Timor.
sessiliflora, *Salisb. Prod.* 195.—Ins. Bourbon.
sessilifolia, Steud. Nom. ed. II. i. 415 = Pluchea sessilifolia.
sessilifolia, Wall. Cat. n. 3029 = Pluchea Wallichiana.
sicula, Willd. Sp. Pl. iii. 1981 = Pulicaria sicula.
sinensis, J. F. Gmel. Syst. 1221 = Blumea chinensis.
sinuata, Ell. Sketch, ii. 328 = Erigeron latifolius.
solidaginoides, Wall. Cat. n. 3056 = Aster ignoratus.
sonchifolia, Buch.-Ham. in Wall. i. c. n. 3068 = Blumea laciniata.
sophiæfolia, *H. B. & K. Nov. Gen. et Sp.* iv. 78. t. 326.—Mexic.
sordida, Linn. Mant. 466 = Phagnalon sordidum.
spathulata, *Hornem. Hort. Hafn.* ii. 806.— Hab. ?
spathulifolia, Maxim. ex Regel, in Act. Hort. Petrop. iii. (1874) 109 = stricta.
spatulata, Link, Enum. Hort. Berol. ii. 322 = Vernonia pauciflora.
spicata, Lam. Encyc. ii. 98 = Pterocaulon virgatum.
spicata, *Mill. Gard. Dict. ed.* VIII. n. 14.—Hab. ?
spinulosa, *Mig. Fl. Ind. Bat.* ii. 45.—Burma.
spinulosa, Miq. l. c. 44 = Blumea membranacea.
spinosa, *Sch. Bip. ex Oliver & Hiern, in Fl. Trop. Afr.* iii. 312.—Afr. trop.
spinulosa, Spreng. Syst. iii. 510 = Sommerfeldtia spinulosa.
splendens, Spreng. l. c. 509 = Piptocarpa lucida.
Sprengelii, *Sch. Bip. ex Walp. Rep.* ii. 956.—Afr. austr.
squamata, Spreng. Syst. iii. 515 = Aster divaricatus.
squamosa, Bory, Voy. iii. 110 = Senecio squamosus.
squarrosa, *Georgi, It.* i. 239.—Sibir. baical.
squarrosa, Linn. Sp. Pl. 863 = Inula vulgaris.
squarrosa, Wall. Cat. n. 3092 = Blumea myriocephala.
stachyoides, Spreng. ex DC. Prod. v. 442 = Blumea glomerata.
Steetzii, *Griseb. App. Ind. Sem. Hort. Gotting.* (1878) 8.—Afr. austr.
stellata, Spreng. Neue Entdeck. ii. 142 = Vernonia oppositifolia.
Steudelii, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 388.—Abyssin.
stipulacea, *Decf. Cat. Hort. Par.* ed. III. 164.—Hab. ?
stipulata, Vahl, ex DC. Prod. v. 97 = Liabum Jussieui.
striata, Wall. Cat. n. 3065 = vicidula.
stricta, Wall. l. c. n. 3044 = japonica.
stricta, *Willd. Sp. Pl.* iii. 1952.—Ind. or.; Afr. trop.
subcapitata, Wall. Cat. n. 3056 = Blumea subcapitata.
subdecurrens, DC. Prod. v. 379 = Erigeron subdecurrens.
subdecurrens, A. Gray, Pl. Fendl. 78; Pl. Wright. i. 102 = Coulteri.
subracemosa, *Miq. Fl. Ind. Bat.* ii. 41.—Java.
subscandiis, Wall. Cat. n. 3000 = Vernonia subscandiis.
subsimplex, Wall. l. c. n. 3003 = Vernonia cinerea.
subspicata, *Phil. in Linnæa*, xxviii. (1856) 735.—Chili.
suffruticosa, *Phil. l. c.*—Chili.
sumbavensis, *Miq. Fl. Ind. Bat.* ii. 46.—Malaya.
sylvatica, Blume, Bijdr. 896 = Blumea sylvatica.
symphytifolia, *Mill. Gard. Dict.* ed. VIII. n. 10.—Hab. ?
syringæfolia, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 283.—China.
tenella, *Miq. Fl. Ind. Bat.* ii. 40.—Ins. Timor.
tenera, Wall. Cat. n. 3023 = Blumea oxyodonta.
Tenorii, Presl, ex Spreng. Pugill. i. 55 = Phagnalon rupestre.
tenuifolia, Salisb. Prod. 194 (= *Schkuhria serrata*).—Ins. Canar.
ternatensis, *Miq. Fl. Ind. Bat.* ii. 49.—Ins. Ternate.
tetragona, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 178 = Blumea tetraptera.
thapsoides, Bieb. Fl. Taur. Cauc. ii. 304 = Inula thapsoides.
thodiifolia, *Phil. in Anal. Univ. Chil.* (1862) ii. 381.—Chili.
thodiifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 75.—Mexic.

CONYZA:—
thapsoides, Lam. Encyc. ii. 90 = Baccharis thyoides.
thyrsiflora, C. B. Clarke, Comp. Ind. 76, sphalm. = sev.
thyrsiflora, Wall. Cat. n. 3082 = Blumea lacera.
thyrsoidea, Pers. Syn. ii. 426. n. 8 = Blumea lacera.
tigreana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 316.—Afr. trop.
timorensis, *Miq. Fl. Ind. Bat.* ii. 47.—Ins. Timor.
tomentosa, Burm. f. Fl. Cap. Prod. 26 = Alcicope lanosa.
tomentosa, Forsk. Fl. Aegypt. Arab. 148 = Phagnalon rupestre.
tomentosa, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
tomentosa, Wall. Cat. n. 3031 = Pluchea tomentosa.
tortuosa, *Linn. Sp. Pl.* 862.—Am. austr.
trasoncorensis, Wall. Cat. n. 3067 = vicidula.
tricuneata, Willd. Sp. Pl. iii. 1946 = Baccharis confertæ.
tridentata, Poir. Encyc. Suppl. ii. 842 = Baccharis tridentata ?
trilobæ, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 261 = Erigeron trilobus.
trinervis, *Mill. Gard. Dict.* ed. VIII. n. 12.—Mexic.
trinervis, Lam. Encyc. ii. 55 = Baccharis trinervis.
triplinervia, *Less. in Linnæa*, vi. (1831) 127.—Bras.
trinervis, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 172 = sativa.
tuberosa, Lam. ex Steud. Nom. ed. II. i. 415 = Jasonia radiata.
uliginosa, Pers. Syn. ii. 426 = Pluchea bifrons.
umbrosa, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 879.—Sibiria.
undulata, *Poir. Encyc. Suppl.* ii. 839.—Afr. trop.
uniflora, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 140 = Helichrysum filicaule.
uniflora, *Mill. Gard. Dict.* ed. VIII. n. 18.—Mexic.
vagans, Wall. Cat. n. 3040 = Vernonia vagans.
variegata, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 389.—Abyssin.
Vatikeana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 311.—Afr. trop.
verbascifolia, Bieb. ex Georgi, Beschr. Russ. Reich. iv. 1239 = Inula thapsoides.
verbascifolia, Bory, Voy. ii. 377 = anchusæfolia.
verbascifolia, Frivald. ex Nym. Consp. 398 = Inula Aschersoniana
verbascifolia, Willd. Sp. Pl. iii. 1924 = Inula candida.
veronicæfolia, Wall. Cat. n. 3002 = japonica.
verrucosa, Houtt. ex DC. Prod. vii. 18 = Printzia Bergii.
vestita, Wall. Cat. n. 3008 = Blumea balsamifera.
villosa, Herb. Madr. ex Wall. i. c. n. 3108 = Blumea amplectens.
villosa, Willd. Sp. Pl. iii. 1929 = Laggera aurita.
viminea, *Mig. Fl. Ind. Bat.* ii. 49.—Malaya.
virens, Wall. Cat. n. 3037 = Blumea virens.
virgata, *DC. Prod.* v. 385.—Madag.
virgata, Linn. Sp. Pl. ed. II. 1206 = Pterocaulon virgatum.
vicidula, *Wall. Cat.* n. 3006.—As. trop.; Austral.
viscosa, *Soy.-Will. & Godr. in Mem. Soc. Nancy*, ii. (1851) 349.—Ins. Maurit.
viscosa, Hort. ex Steud. Nom. ed. II. i. 415 = Inula viscosa.
viscosa, *Mill. Gard. Dict.* ed. VIII. n. 8.—Mexic.
viscosula, Wall. ex DC. Prod. v. 441 = Blumea membranacea.
volkameriifolia, Wall. Cat. n. 3001 = Vernonia volkameriifolia.
volubilis, Wall. l. c. n. 3057 = Microglossa volubilis.
vulgaris, Lam. Fl. Fr. ii. 78 = Inula vulgaris.
vulgaris, *Phil. in Linnæa*, xxviii. (1856) 735.—Chili.
Wallichii, DC. Prod. v. 384 = vicidula.
wangtsueuis, *C. B. Clarke, Comp. Ind.* 61.—Reg. Himal.
Wightiana, Wall. Cat. n. 3093 = Vernonia arborea.
Wightiana, Wall. ex DC. Prod. v. 426 = Blumea Wightiana.
Wulfeni, Host, ex Beck, in Denkschr. Akad. Wien, Math. Nat. xliv. (1882) 329 = Inula bifrons.
Zollingeriana, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 179.—Java.

CONYZELLA, Rupr. in Mém. Acad. Pétersb. Sér. VII. xiv. n. 6 (1869); Sert. Tianschi. 51 = Erigeron, Linn. (Compos.).
canadensis, Rupr. l. c. = E. canadensis.

CONYZOIDES, Tourn. ex DC. Prod. vi. 281 (1837) = Carpesium, Linn. (Compos.).

COOCKIA, Batsch, Tab. 58 (1802) = COOKIA, Sonner.

COOKIA, J. F. Gmel. Syst. 24 (1791) = Pimelea, Banks & Soland. (Thymel.).
Gnidia, J. F. Gmel. l. c. = P. Gnidia.
pilosa, J. F. Gmel. l. c. = P. pilosa.
prostrata, J. F. Gmel. l. c. = P. prostrata.

COOKIA, Sonner. Voy. Ind. iii. 258. t. 130 (1782) = Clausena, Burm. f. (Rutac.).
anisata, Desf. Cat. Hort. Par. ed. III. 415, 406.—Hab. ?
anisodora, Blanco, Fl. Filip. ed. II. 253 = Clausena excavata.
Anisum-olens, Blanco, l. c. ed. I. 359 = Clausena excavata.
australis, F. Muell. Fragm. i. 25 = Pentaceras australis.
chlorosperma, Blume, Bijdr. 135 = Glycosmis chlorosperma.
cyanocarpa, Blume, l. c. 136 = Glycosmis cyanocarpa.
dulcis, Bedd. in Madr. Journ. Sc. Ser. II. xxii. (1861) 73 = Clausena Willdenowii.
falcata, DC. Prod. i. 537.—China.
glabrescens, Miq. Fl. Ind. Bat. i. ii. 534.—Ins. Philipp.
gravcolens, Wight & Arn. Prod. 95 = Clausena excavata.
macrophylla, Lindl. in Wall. Cat. n. 6367 = Clausena heptaphylla.
pentaphylla, Wight & Arn. ex M. Roem. Syn. Hesper. 45 = Clausena pentaphylla.
punctata, Hassk. Cat. Hort. Bog. Alt. 215 = Micromelum pubescens.
punctata, Ket. Obs. vi. 29 = Clausena excavata.
punctata, Sonner. Voy. Ind. iii. 258. t. 130 = Clausena Wampi.
Sonnerotii, Montr. in Mém. Acad. Lyon, x. (1860) 184. —N. Caled. ?
Wampi, Blanco, Fl. Filip. ed. I. 358 = Clausena Wampi.

COOPERIA, Herb. in Bot. Reg. t. 1835 (1836). *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 732.
SCEPTRANTHUS, R. Grah. in Edinb. N. Phil. Journ. xx. (1836) 413.
chlorosolen, Herb. in Bot. Reg. t. 1835 = Drummondi.
Drummondii, *Herb. l. c.*—Texas.
mexicana, Herb. Amaryll. 183 = Drummondii.
pedunculata, *Herb. l. c.* 179. t. 42. ff. 8–5.—Texas.

COPAIBA, Adans. Fam. ii. 341 (1763) = seq.

COPAIFERA, Linn. Sp. Pl. ed. II. 557 (1762). *LEGUMINOSÆ*, Benth. & Hook. f. i. 565.
COPAIBA, Adans. Fam. ii. 341 (1763).
COPAIVA, Jacq. Enum. Pl. Carib. 4 (1760).
GORSKIA, Bolla, in Peters, Reise Mossamb. Bot. i. 15. t. 8 (1861).
GUIBOURTIA, J. J. Benn. in Journ. Linn. Soc. i. (1857) 149.
Beyrichii, *Hayne, Arzneig.* x. t. 12.—Bras.
bijuga, *Willd. ex Hayne, l. c.* t. 16.—Bras.
bracteata, *Benth. in Hook. Journ. Bot.* ii. (1840) 101. —Guiana.
coleosperma, *Benth. in Trans. Linn. Soc.* xxv. (1866) 316. t. 48.—Afr. trop.
confertiflora, *Benth. in Mart. Fl. Bras.* xv. ii. 241.—Am. trop.
copaifera, Baill. Hist. Pl. ii. 142, 163 = Guibourtiana.
cordifolia, *Hayne, Arzneig.* x. t. 21.—Bras.
coriacea, *Mart. Reise Bras.* i. 365; *et in Hayne, l. c.* t. 20.—Bras.
disperma, *Willem. in Usteri, Ann. Bot.* xviii. (1796) 54.—Ins. Maurit.
elliptica, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 127.—Bras.
glabra, *Vogel, in Linnæa*, xi. (1837) 410.—Bras.
Gorskiana, *Benth. in Trans. Linn. Soc.* xxv. (1866) 317.—Afr. trop.
guyanensis, *Decf. in Mém. Mus. Par.* vii. (1821) 378.—Guiana.
Guibourtiana, *Benth. in Trans. Linn. Soc.* xxv. (1866) 317.—Afr. trop.

COPAIFERA :—
hymenæifolia, *Moric. in Mem. Soc. Phys. Genev.* vi. (1833) 529.—Cuba.
Jacquini, Desf. in Mem. Mus. Par. vi. (1821) 376 = officinalis.
Jacquiniana, G. Don, Gen. Syst. ii. 455 = officinalis.
Jamesii, *Hayne, Arzneig.* x. sub t. 17.—Peruv.
Lansdorii, *Desf. in Mem. Mus. Par.* vii. (1821) 377. —Bras.
laxa, *Hayne, Arzneig.* x. t. 16.—Bras.
Mannii, Baill. Adansonia, vi. (1866) 202 = Hardwickia Mannii.
marginata, *Benth. in Mart. Fl. Bras.* xv. ii. 243.—Bras.
Martii, *Hayne, Arzneig.* x. t. 15.—Bras.
Mopane, J. Kirk, ex Benth. in Trans. Linn. Soc. xxv. (1866) 317.—Afr. trop.
multijuga, *Hayne, Arzneig.* x. sub t. 17.—Bras.
nitida, *Mart. ex Hayne, l. c.* x. t. 17.—Bras.
oblongifolia, *Mart. ex Hayne, l. c.* t. 23; *et in Mart. Reise Bras.* i. 548.—Bras.
officinalis, *Linn. Sp. Pl.* ed. II. 557.—Am. trop.
pubiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 101. —Guiana.
rigida, *Benth. in Mart. Fl. Bras.* xv. ii. 248.—Bras.
Sellowii, *Hayne, Arzneig.* x. t. 22.—Bras.
trapezifolia, *Hayne, l. c.* sub t. 23.—Bras.

COPAIVA, Jacq. Enum. Pl. Carib. 4 (1760) = Copaifera, Linn. (1762).
officinalis, Jacq. l. c. 31 = Copaifera officinalis.

COPERNICIA, Mart. Hist. Nat. Palm. ii. 151. t. 50 A; iii. 242 (1837 ?). *PALMÆ*, Benth. & Hook. f. iii. 927.
CRYOSOPHILA, Blume, Rumphia. ii. 53, in note (1836).
CRYOSOPHILA, Benth. & Hook. f. Gen. iii. 928 (1883).
cerifera, *Mart. Hist. Nat. Palm.* iii. 242. t. 50 A.—Am. trop.
hospita, *Mart. l. c.*—Cuba.
macroglossa, *H. Wendl. in Kerch. Palm.* 241, nomen. —Hab. ?
maritima, *Mart. Hist. Nat. Palm.* iii. 243. t. 50 A. —Cuba.
Miraguana, Mart. l. c. = Thrinax Miraguana.
nana, *Lodd. ex Hemsl. Biol. Centr. Am. Bot.* iii. 411. —Mexic.
Pumos, *Mart. Hist. Nat. Palm.* iii. 243,—Cuba.
robusta, *Hort. ex Linden, Illustr. Hortic.* xxviii. (1881) 16.—Cuba.
tectorum, *Mart. Hist. Nat. Palm.* iii. 243.—N. Granat. Wrightii, *Griseb., Cat. Pl. Cub.* 230.—Cuba.

COPIANTHUS, Hill, Decade t. 1 (1778). Amaranthac. ? monstr.
indica, Hill, l. c.—Ind. or.

COPIOGLOSSA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 194 = Ruellia, Linn. (Acanthac.).
pilosa, Miers, l. c. (= *Dipteracanthus Neesianus*).—Bras.

COPISMA, E. Mey. Comm. Pl. Afr. Austr. 189 (1835) = Rhynchosida, Lour. (Legumin.).
caribæum, E. Mey. l. c. 188 = R. caribæa.
diversifolium, E. Mey. l. c. 185 = R. hirsuta.
effusum, E. Mey. l. c. = R. sclerodes.
falcatum, E. Mey. l. c. 186 = R. minima.
gibbum, E. Mey. l. c. 187 = R. caribæa.
glabrum, E. Mey. l. c. 184 = R. glandulosa.
glandulosum, E. Mey. l. c. 183 = R. glandulosa.
grandifolium, E. Mey. l. c. 188 = R. grandifolia.
nitidum, E. Mey. l. c. 186 = R. nitida.
paniculatum, E. Mey. l. c. 184 = R. totta.
pictum, E. Mey. l. c. 185 = R. caribæa.
pilosum, E. Mey. l. c. 183 = R. pilosum.
rotundifolium, E. Mey. l. c. 187 = R. rotundifolia.
secundum, Walp. in Linnæa, xiii. (1839) 587 = R. secunda.
simplicifolium, E. Mey. Comm. Pl. Afr. Austr. i. 186 = R. stagnifolia.
stagnifolium, Steud. Nom. ed. II. ii. 454, sphalm. = R. stagnifolia.
subcordatum, Bond. in Linnæa, xxiii. (1850) 34 = Calastia tenuiflora.
tenue, E. Mey. Comm. Pl. Afr. Austr. 186 = R. minima.
tetinum, E. Mey. l. c. 183 = R. puberula.
trichodes, E. Mey. l. c. 184 = R. tetra.
viscidulum, E. Mey. l. c. = R. viscidula.

COPPENSIA, Dum. in Mém. Acad. Brux. ix. (1835) (Sur Maelen.) 10, in nota. = Oncidium, Sw. (Orchid.).
bifolia, Dum. l. c. = O. bifolium.

COPPOLERIA, Todaro, in Atti Acc. Sc. Palermo, N. S. i. (1845) 14 = Vicia, Tourn. (Legumin.).
monantha, Todaro. l. c. = V. monantha.

COPROSMA, Forst. Char. Gen. 137. t. 69 (1776). *RUBIACEÆ*, Benth. & Hook. f. ii. 139.
EDARTHRONIA, Nutt. ex A. Gray, in Proc. Am. Acad. iv. (1860) 49, in syn.
EURYNOME, DC. Prod. iv. 475 (1830).
MARQUISIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 192.
PELAPHIA, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. ii. (1839) 306.
acerosa, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 207. —N. Zel.
acutifolia, *Hook. f. in Journ. Linn. Soc.* i. (1857) 128. —N. Zel.
acutifolia, F. Muell. ex Benth. Fl. Austral. iii. 429 = Morinda acutifolia.
affinis, Hook. f. Fl. Antarct. i. 21. t. 4 = foetidissima.
arborea, *T. Kirk, in Trans. N. Z. Inst.* x. (1878) 430. —N. Zel.
arcuata, *Colenso, Excurs. North. ist.* 84.—N. Zel.
Baueri, *Endl. Iconogr.* t. 111.—Ins. Norfolc.; N. Zel.
Baueriana, Hook. f. Fl. N. Zel. i. 104 = Baueri.
Billardieri, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 465 [bis].—Austral.
calycina, A. Gray, in Proc. Am. Acad. iv. (1860) 306 (= *Leptostigma Arnottianum*).—Chili
ciliata, *Hook. f. Fl. Antarct.* i. 22.—Ins. Auckland.
Colensoi, *Hook. f. Handb. New Zeal. Fl.* 117.—N. Zel.
concinna, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 380.—N. Zel.
conferta, A. Cunn. in Ann. Nat. Hist. ii. (1839) 206 = Cunninghamii.
crassifolia, *Colenso, Excurs. North. Isl.* 75.—N. Zel.
cuneata, *Hook. f. Fl. Antarct.* i. 21. t. 15.—N. Zel.
Cunninghamii, *Hook. f. Handb. New Zeal. Fl.* 113.—N. Zel.
cuspidifolia, DC. Prod. iv. 578 = hirtella.
depressa, *Colenso, ex Hook. f. Fl. N. Zel.* 110.—N. Zel.
divaricata, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 207. —N. Zel.
ernodeoides, *A. Gray, in Proc. Am. Acad.* iv. (1860) 49.—Ins. Pacif.
foetidissima, A. Cunn. in Ann. Nat. Hist. ii. (1839) 206 = Cunninghamii.
foetidissima, *Forst. Char. Gen.* 138.—N. Zel.
foliosa, *A. Gray, in Proc. Am. Acad.* iv. (1860) 48.—Ins. Sandvic.
graveolens, A. Cunn. in Ann. Nat. Hist. ii. (1839) 206 = rhamnoides ?
grandifolia, *Hook. f. Fl. N. Zel.* 104.—N. Zel.
hirtella, *Labill. Nov. Holl. Pl.* i. 70. t. 95.—Austral.
lanceolaria, *F. Muell.* Fragm. ix. 70.—Austral.
ligustrina, *Kunth, Ind. Sem. Hort. Berol.* (1846) 13. —Hab. ?
linariifolia, *Hook. f. Handb. New Zeal. Fl.* 118.—N. Zel.
longifolia, *A. Gray, in Proc. Am. Acad.* iv. (1860) 48. —Ins. Sandvic.
lucida, *Forst. Char. Gen.* 138.—N. Zel.
Menziesii, *A. Gray, in Proc. Am. Acad.* iv. (1860) 49.—Ins. Pacific.
microcarpa, *Hook. f. Fl. N. Zel.* 110.—N. Zel.
microphylla, A. Cunn. ex Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 465 [bis] = Billardieri.
myrtifolia, *Naronha, in Ver. Batav. Gen.* v. (1790) ed. I. Art. iv. 10.—Java.
myrtilliifolia, Hook. f. Fl. Antarct. i. 21 = parviflora.
Nertera, F. Muell. Fragm. ix. 186 = Nertera depressa.
nitida, *Forst. f. in Hook. Lond. Journ. Bot.* vi. (1847) 465 [bis].—Austral.
parviflora, *Hook. f. Fl. N. Zel.* 108.—N. Zel.
pedicellata, *A. Gray, in Proc. Am. Acad.* iv. (1860) 50.—Ins. Fiji.
petiolata, *Hook. f. in Journ. Linn. Soc.* i. (1857) 128. —Ins. Kermadec.
pilosa, *Endl. Fl. Ins. Norf. Prod.* 60.—Ins. Norfolc.
propinqua, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 206.—N. Zel.
pubens, *A. Gray, in Proc. Am. Acad.* iv. (1860) 49.—Ins. Sandvic.

COPROSMA :—
pumila, *Hook. f. Fl. N. Zel.* 111.—Austral.; N. Zel.
pusilla, Forst. t. Prod. 90; A. Rich. Voy. Astrol. Bot. i. 365 = foetidissima.
putida, *C. Moore & F. Muell. Fragm.* vii. 45.—Ins. D. Hown.
repens, A. Rich. Voy. Astrol. Bot. i. 264 = foetidissima.
repens, *Hook. f. Fl. Antarct.* i. 22. t. 16.—N. Zel.
reptans, *F. Muell. Fragm.* ix. 186.—Austral.
retusa, Banks & Soland. ex Hook. f. in Hook. Lond. Journ. Bot. iii. (1844) 415 = Baueri.
rhamnoides, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 206.—N. Zel.
rhynchocarpa, *A. Gray, in Proc. Am. Acad.* iv. (1860) 48.—Ins. Sandvic.
robusta, *Raoul, in Ann. Sc. Nat.* Ser. III. ii. (1844) 121.—N. Zel.
rotundifolia, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 206.—N. Zel.
rubra, *Petrie, in Trans. N. Z. Inst.* xvii. (1885) 269.—N. Zel.
serrulata, *Hook. f. ex Y. Buch. l. c.* iii. (1871) 212.—N. Zel.
spathulata, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 207.—N. Zel.
Stochi Hort. ex J. Barbier, in Rev. Hortic. Belg. iii (1877) t. 12 = Baueri.
sundana, *Miq. Fl. Ind. Bat.* ii. 327.—Malaya.
taitensis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 49. —Ins. Tahiti.
tenuicaulis, *Hook. f. Fl. N. Zel.* 106.—N. Zel.
triflora, *Benth. & Hook. f. Gen.* ii. 139.—Ins. Juan Fernand.
ovnata, A. Cunn. ex Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 465 [bis] = hirtella.
virescens, *Petrie, in Trans. N. Z. Inst.* xi. (1879) 496. —N. Zel.
Weinmannioides, *Wawra, in Flora*, lvii. (1874) 327.—Ins. Pacif.

COPROSMANTHUS, Kunth, in Abh. Akad. Berl. (1846) 85; et Enum. Pl. v. 365 (1850) = Smilax, Linn. (Liliac.).
consanguineus, Kunth, Enum. Pl. v. 366 = S. herbacea.
herbacea, Kunth, l. c. 365 = S. herbacea.
japonicus, Kunth, l. c. 368 = S. China.
lasioneuron, Kunth, l. c. 365 = S. herbacea.
peduncularis, Kunth, l. c. 364 = S. herbacea.
tamnifolius, Kunth, l. c. 367 = S. tamnifolia.

COPTIDIUM, Nym. Consp. 13 (1878) = Ranunculus, Tourn.
ranunculoides, Nym. l. c. = R. lapponicus.

COPTIS, Salisb. in Trans. Linn. Soc. viii. (1807) 305 *RANUNCULACEÆ*, Benth. & Hook. f. i. 8.
CHRYSOCOPTIS, Nutt. in Journ. Acad. Philad. vii. (1834) 9. t. 1.
anemonæfolia, *Sieb. & Zucc. in Abh. Math. Phys. Kl. Munch.* iv. ii. (1848) 180.—Japon.
asplenifolia, *Salisb. in Trans. Linn. Soc.* viii. (1807) 306.—Am. bor.
brachypetala, *Sieb. & Zucc. in Abh. Math. Phys. Kl. Munch* iv. ii. (1848) 180.—Japon.
occidentalis, *Torr. & Gray, Fl. N. Am.* i. 38.—Am. bor. occ.
orientalis, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 80.—Japon.
quinquefolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 7.—Japon.
Teeta, *Wall. in Linnæa*, xii. (1838) *Litt.* 227.—Reg. Himal.
trifolia, *Salisb. in Trans. Linn. Soc.* viii. (1807) 305.—Reg. bor. temp. et arct.

COPTOCHEILE, Hoffingg. Vers. Pfl. Nachtr. (1841) ex Linnæa, xvi. (1842) Litt. 279. *GESNERACEÆ* ?
macrorhiza, *Hoffingg. l. c.*—Bras. ?

COPTOPHYLLUM, Korth. in Ned. Kruidk. Arch. ii. (1851) 161. *RUBIACEÆ*, Benth. & Hook. f. ii. 68.
bracteatum, *Korth. l. c.* 162.—Sumatra.
capitatum, *Miq. Fl. Ind. Bat.* ii. 845.—Malaya.
pilosum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 230.—Java.

COPTOSAPELTA, Korth. in Ned. Kruidk. Arch. ii. 11. (1851) 112. *RUBIACEÆ*, Benth. & Hook. f. ii. 35.
flavescens, *Korth. l. c.* 113.—Malaya.
Griffithii, *Hook. f. Ic. Pl.* t. 1069.—Malaya

COPTOSPERMA, Hook. f. in Benth. & Hook. f. Gen. ii. 86 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
nigrescens, *Hook. f. l. c.* 87.—Afr. trop.

COQUEBERTIA, Brongn. in Ann. Sc. Nat. Sér. I. xxv. (1832) 111 = **Sollerntia**, Maximil. & Nees (Legumin.).
ilicifolia, Brongn. l. c. = Z. ilicifolia.

CORALLIORRHIZA, Aschers. Fl. Brandenb. 697 (1864) = **Corallorrhiza**, Hall. (Orchid.).

CORALLOBOTRYS, Hook. f. in Benth. & Hook. f. Gen. ii. 575 (1876). *VACCINIACEAE*, Benth. & Hook. f. l. c.
acuminata, *Hook. f. Fl. Brit. Ind.* iii. 455.—Reg. Himal.

CORALLOCARPUS, Welw. ex Benth. & Hook. f. Gen. i. 831 (1867). *CUCURBITACEAE*, Benth. & Hook. f. l. c.
conocarpus, *Benth. & Hook. f. l. c.*—Ind. or.; Afr. trop.
corallinus, Cogn. in DC. Monog. Phan. iii. 647 = Fenzlii.
Courbonii, Cogn. l. c. 655 = velutinus.
Ehrenbergii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 566.—Nubia.
epigaeus, *Benth. & Hook. f. Gen.* i. 831.—Ind. or.
erostris, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 567.—Afr. trop.
ethaicus, Hook. f. l. c. = velutinus.
Fenzlii, *Hook. f. l. c.* 565.—Afr. trop.
Gijef, *Hook. f. l. c.* 566.—Nubia.
gloneratus, *Cogn. in DC. Monog. Phan.* iii. 658.—Venezuela.
gracilipes, *Cogn. l. c.* 650.—Ind. or.
palmatus, *Cogn. l. c.* 648.—Ind. or.
parvifolius, *Cogn. l. c.* 653.—Arabia.
pedunculosus, *Cogn. l. c.* 654.—Abyssin.
Poissonii, *Cogn. l. c.* 651.—Madag.
Schimperi, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 567.—Abyssin.
velutinus, *Benth. & Hook. f. Gen.* i. 831.—Ind. or. Welwitschii, *Benth. & Hook. f. l. c.*—Afr. trop.

CORALLODENDRON, Tourn. ex Rupp. Fl. Jen. ed. Hall. 237 (1745) = **Erythrina**, Linn. (Legumin.).
aculeatum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 341 = E. Corallodendron.

CORALLOPHYLLUM, H. B. & K. Nov. Gen. et Sp. vii. 275. t. 660 bis (1825) = **Lennoe**, La Llave & Lex.
coerulenm, H. B. & K. l. c. 276 = L. caerulea.

CORALLORRHIZA, Hall. Enum. Stirp. Helv. i. 278 (1743) (Corallorhiza); R. Br. in Ait. Hort. Kew. ed. II. v. 209 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 497.
RHIZOCORALLON, Hall. in Rupp. Fl. Jen. 501 (1745).
arizonica, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 379.—Arizona.
Bigelovii, *Wats. l. c.* xii. (1877) 275.—Calif.
bulbosa, A. Rich. & Gal. in Ann. Sc. Nat. Ser. III. iii. (1845) 19 = mexicana.
dentata, Host, Fl. Austr. ii. 547 = innata.
Ehrenbergii, *Reichb. f. in Linnaea*, xxii. (1849) 883.—Mexic.
erictorum, Drej. in Fl. Dan. t. 2264 = innata.
follosa, *Lindl. Gen. et Sp. Orch.* 535.—Ind. or.
grandiflora, *A. Rich. & Gal. in Ann. Sc. Nat. Sér. III.* iii. (1845) 19.—Mexic.
Hallerí, Rich. in Mém. Mus. Par. iv. (1818) 61 = innata.
hiemalis, *Nutt. Gen. Am.* ii. 198.—Am. bor.
indica, *Lindl. in Journ. Linn. Soc.* iii. (1859) 26.—Ind. or.
innata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 209.—Europ.; Am. bor.
intacta, Cham. & Schlecht. in Linnaea, iii. (1828) 85 = innata.
Jacquemontii, *Decne. in Jacquem. Voy. Bot.* 166. t. 163.—Reg. Himal.
Macraei, A. Gray, Man. Bot. N. U. St. (1856) 453 = stricta.
CORALLORHIZA:—
Mortoniana, *Bong. Veg. Ins. Sitcha*, 166.—Alaska.
mexicana, *Lindl. Gen. et Sp. Orch.* 534.—Mexic.
micrantha, *Chapm. Fl. S. U. St.* 454.—Am. bor.
multiflora, *Nutt. in Journ. Acad. Philad.* iii. (1823) 138. t. 7.—Am. bor.
nemoralis, Sw. ex Nym. Consp. 686 = innata.
Neottia, Scop. Fl. Carn. ed. II. ii. 207 = innata.
odontorhiza, *Nutt. Gen. Am.* ii. 197.—Am. bor.
patens, Lindl. Gen. et Sp. Orch. 535 = Oreorchis patena.
squamata, Poir. in Dict. Sc. Nat. x. 276 = Dipodium squamatum.
striata, *Lindl. Gen. et Sp. Orch.* 534.—Am. bor.
verna, Nutt. in Journ. Acad. Philad. iii. (1823) 136 = innata.
virescens, Drej. ex Heynh. Nom. ii. 159, nomen = innata.
Wisteriana, Conrad, in Journ. Acad. Philad. vi. (1829) 145 = odontorhiza.

CORALLOSPARTIUM, Armst. in Trans. N. Z. Inst. xiii. (1881) 333. *LEGUMINOSAE*.
crassicaule, *Armst. l. c.*—N. Zel.

CORALLUMA. Schrank, ex Haw. Syn. Pl. Succ. ed. II. 47 (1819) = **Caralluma**, R. Br. (Asclep.).

CORBICHONIA, Scop. Introd. 264 (1777) = **Orygia**, Forsk. (Ficoid.).

CORBULARIA, Salisb. in Trans. Hort. Soc. i. (1812) 349 = **Narcissus**, Tourn. (Amaryllid.).
albicans, Haw. Revis. 122 = N. Bulbocodium.
aurea, Haw. Monog. (in Sweet, Brit. Flow. Gard. iv.) t. n. 5 = N. Bulbocodium.
Bulbocodium, Haw. Revis. 123 = N. Bulbocodium.
cantabrica, Haw. Monog. l. n. 1 = N. Bulbocodium.
conspicua, Haw. l. c. n. 8 = N. Bulbocodium.
filifolia, M. Roem. Syn. Ensat. 190 = N. Bulbocodium.
Gigas, Haw. Monog. l. n. 10 = N. Bulbocodium.
Gratisii, Webb, ex Willk. & Lange, Prod. Fl. Hisp. i. 150 = N. Bulbocodium.
hedraeantha, Webb & Heldr. Cat. Pl. Hisp. 220 = N. Bulbocodium.
lobulata, Haw. Revis. 122 = N. Bulbocodium.
monophylla, Dur. in Duch. Rev. Bot. ii. (1846–47) 425 = N. Bulbocodium.
ninalis, Nym. Consp. 718 = N. Bulbocodium.
obesa, Salisb. in Trans. Hort. Soc. i. (1812) 350 = N. Bulbocodium.
Schultesii, M. Roem. Syn. Ensat. 190 = N. Bulbocodium.
serotina, Haw. Monog. l. n. 9 = N. Bulbocodium.
tenuifolia, Salisb. in Trans. Hort. Soc. i. (1812) 349 = N. Bulbocodium.
turgida, Salisb. l. c. 351 = N. Bulbocodium.

CORCHOROPSIS, Sieb. & Zucc. in Abh. Akad. Muench. iii. (1843) 737. t. 4. *TILIACEAE*, Benth. & Hook. f. i. 235.
crenata, *Sieb. & Zucc. l. c.*—Japon.

CORCHORUS, [Tourn.] Linn. Syst. ed. I (1735). *TILIACEAE*, Benth. & Hook. f. i. 235.
ANTICHORUS, Linn. Mant. 9. n. 1357 (1767).
CARICHERIA, Scop. Introd. 104 (1777).
GANJA, Reichb. Handb. 302 (1837).
MARELENSIA, Vell. Fl. Flum. v. t. 112 (1827).
PALLADIA, Lam. Illustr. t. 385 (1792).
acutangulus, *Lam. Encyc.* ii. 104.—Cosmop. trop.
acutiformis, Hill, Veg. Syst. xiv. 28 = asiaticus.
aestuans, Blanco, Fl. Filip. ed. I. 441 = acutangulus.
aestuans, Forsk. Fl. Aegypt. Arab. 101 = trilocularis.
aestuans, Herb. Madr. ex Wall. Cat. p. 237. n. 1074 = urticaefolius.
aestuans, *Linn. Syst.* ed. X. 1079.—Am. trop.
alatus, *G. Don, Gen. Syst.* i. 545.—Afr. trop.
americanus, Schrank, ex Steud. Nom. ed. II. i. 416 = Sida erubescens.
angustifolius, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* x. (1828) 18.—Afr. trop.
Antichorus, *Raeusch. Nom.* ed. III. 158.—Geront. trop.
arcuatus, *H. B. & K. Nov. Gen. et Sp.* v. 239.—Mexic.
CORCHORUS:—
argutus, *H. B. & K. Nov. Gen. et Sp.* v. 237.—N. Granat.
asplenifolius, *Burch. Trav.* i. 400.—Afr. austr.
asplenifolius, E. Mey. ex Harv. & Sond. Fl. Cap. i. 229 = trilocularis.
australis, A. Cunn. ex F. Muell. Fragm. iii. 8 = Cunninghamii.
bifurcatus, *Mill. Gard. Dict.* ed VIII. n. 5.—Jamaic.
brachycarpus, *Rich. ex Steud. Nom.* ed. II. i. 416.—Afr. trop.
brachycarpus, Guill. & Perr. Fl. Seneg. Tent. i. 89 = fascicularis.
bullatus, Jacq. Eclog. Pl. t. 168 = pilolobus.
Burmanni, DC. Prod. i. 505 = tridens.
campestris, Macfad. Fl. Jamaic. i. 107 = acutangulus.
capsularis, *Linn. Sp. Pl.* 529.—Cosmop. trop.
carophylloides, Linn. Syst. ed. X. 1079 = siliquosus ?
catharticus, Blanco, Fl. Filip. ed. I. 442 = olitorius.
cordifolius, Salisb. Prod. 380 = capsularis.
Corsta, Linn. Pl Jam. Pugill. 14 = siliquosus.
Cuninghami, *F. Muell. Fragm.* iii. 8.—Austral.
decumanguloris, Roxb. Hort. Beng 42; Fl. Ind. ii. 582 = olitorius.
depressus, Stocks, in Proc. Linn. Soc. i. (1848) 367 = C. Antichorus.
echinatus, *Benth. Fl. Australiens.* i. 276.—Austral.
elachocarpus, *F. Muell. Fragm.* viii. 6.—Austral.
erodioides, *Balf. f in Proc. Roy. Soc. Edinb.* xi. (1882) 504.—Ins. Socotra.
fascicularis, *Lam. Encyc.* ii. 104.—Geront. trop.
flexuosus, *Thunb. in Trans. Linn. Soc.* ii. (1794) 335.—Japon.
follosus, *Sprang. Syst.* ii. 585.—Cuba.
frutescens, Lam. Encyc. ii. 105 = hirsutus.
fruticulosus, Vis. in Flora, xx. (1837) 448 = trilocularis.
furcatus, G. Don, Gen. Syst. i. 544 = seq.
fuscus, Roxb. Hort. Beng. 42; Fl. Ind. ii. 582 = acutangulus.
gracilis, *R. Br. in Salt, Abyss. App.* 65.—Abyssin.
grandiflorus, Spring, ex Mart. in Flora, xx. (1837) 11. Beibl. 102 = Turnera serrata.
Greveanus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 548.—Madag.
guadalupensis, *Sprang. Syst.* ii. 584.—Ins. Guadalup.
hirsutus, *Linn. Sp. Pl.* 530.—Ind. occ.
hirtus, *Linn. f. c. el.* II. 747.—Am. calid.
hirtus, Thunb. Fl. Jap. 228 = Zehkova acuminata.
humilis, Munro, ex Stocks, in Proc. Linn. Soc. i. (1848) 367 = C. Antichorus.
humilis, *A. St. Hil. Fl. Bras. Merid.* i. 280.—Bras.
hygrophilus, *A. Cunn. ex Benth. Fl. Australiens.* i. 276.—Austral.
japonicus, Thunb. Fl. Jap. 227 = Kerria japonica.
javanicus, *Burm. f. Fl. Ind.* 123. t. 36. f. 3.—Ind. or.
lanceolatus, *G. Don, Gen. Syst.* i. 543.—Guinea.
lanuginosus, Macfad. Fl. Jamaic. i. 106 = hirsutus.
laxiflorus, Sprang. Syst. ii. 585 = pilolobus.
leptocarpus, *A. Cunn. ex Benth. Fl. Australiens.* i. 278.—Austral.
linearis, *Mill. Gard. Dict.* ed. VIII. n. 8.—Mexic.
linearis, Wall. Cat. n. 1070 = tridens.
longicarpus, *G. Don, Gen. Syst.* i. 543.—Guinea.
longipedunculatus, *Mast. in Oliver, Fl. Trop. Afr.* i. 262.—Afr. trop.
Marwa, Buch.-Ham. in Wall. Cat. n. 6311 = capsularis.
microphyllus, Fresen. in Mus. Senckenb. ii. (1837) 156 = C. Antichorus.
monopetalus, *H. B. & K. Nov. Gen. et Sp.* v. 238.—N. Granat.
muricatus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 31.—Abyssin.
muricatus, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* x. (1828) 20.—Afr. trop.
olitorius, *Linn. Sp. Pl.* 529.—Cosmop. trop.
oppositifolius, Hasskarl, in Hoev. & De Vriese, Tijdschr. xii. (1845) 126 = acutangulus.
ortocoensis, *H. B. & K. Nov. Gen. et Sp.* v. 237.—Venezuela.
patens, Lehm. Sem. Hort. Hamb. (1852) ex Linnaea, iii. (1852) Lit. 9 = tridens.
pilolobus, *Link, Enum. Hort. Berol.* ii. 72.—Am. calid.
pilosus, *H. B. & K. Nov. Gen. et Sp.* v. 236. t. 457.—N. Granat.
polygonus, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* x. (1828) 20.—Afr. trop.

CORCHORUS:—
prismatocarpus, *St. Hil. Fl. Bras. Mer.* i. 382.—Bras.
procumbens, *Boj. Hort. Maurit.* 43.—Afr. trop.
pseudocapsularis, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 670.—Afr. trop.
Pumilio, *R. Br. ex Benth. Fl. Austral.* i. 277.—Austral.
pumilus, *Buch.-Ham. ex Wall. Cat.* n. 6310.—Ind. or.
quadrangularis, *Y. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 298.—Ins. Cap. Verd.
quinquelocularis, Moench, Meth. 348 = olitorius.
quinquenervis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 80 = urticifolius.
scandens, *Thunb. in Trans. Linn. Soc. ii.* (1794) 335.—Japon.
secundiflorus, Moç. & Sesse, ex DC. Prod. i. 504 = siliquosus.
senegalensis, Juss. ex Steud. Nom. ed. II. i. 417 = tridens.
serrasfolius, *Burch. Trav.* i. 587.—Afr. austr.
serrasfolius, DC. Prod. i. 504 = trilocularis.
serratus, *Thunb. in Trans. Linn. Soc.* ii. (1794) 335.—Japon.
sidoides, *F. Muell. Fragm.* iii. 9.—Austral.
siliquosus, *Linn. Sp. Pl.* 529.—Am. calid.
siliquosus, Poepp. ex Steud. Nom. ed. II. i. 417 = foliosus.
siliquosus, Torr. & Gray, Fl. N. Am. i. 229 = pilolobus.
spinosus, Sieber, ex Steud. Nom. ed. II. i. 417 = Pavonia spinifex.
subscandens, Wall. Cat. n. 1074 = urticæfolius.
tetragona, *Mill. Gard. Dict.* ed. VIII. n. 4.—Hab. ?
tetragonulus, Langsd. in Sylt. Ratisb. ii. (1828) 168 = pilolobus
tomentillus, *F. Muell. Fragm.* iii. 10.—Austral.
tomentosus, *Thunb. Fl. Jap.* 238.—Japon.
Torresianus, *Gaudich. Bot. Voy. Bonite*, 477.—Ins. Marian.
tortipes, A. St. Hil. Fl. Bras. Mer. i. 281 = hirtus.
tridens, *Linn. Mant.* 566.—Cosmop. trop.
triflorus, Boj. Hort. Maurit. 43 = trilocularis.
trilocularis, Burm. f. Fl. Ind. 123. t. 37. f. 2 = tridens.
trilocularis, *Linn. Mant.* 77.—Geront. trop.
urticæfolius, *Wight & Arn. Prod.* 73.—Ind. or.; Afr. trop.
vermicularis, *F. Muell. Fragm.* iii. 10.—Austral.
villosissimus, *A. St. Hil. Fl. Bras. Mer.* i. 280.—Bras.
Walcottii, *F. Muell. Fragm.* iii. 9.—Austral.

CORDA, St. Leg. in Ann. Soc. Bot. Lyon, viii. (1881) 177 = Cordia, Linn.

CORDAEA, Spreng. Gen. ii. 581 (1831) = Cyamopsis, DC. (Legumin.).
fabaeformis, Spreng. l. c. = Cyamopsis psoraloides.

CORDEMOYA, Baill. Adansonia, i. 255 (1861) = Mallotus, Lour. (Euphorb.).
acuminata, Baill. l. c. = M. Baillonianus.
integrifolia, Baill. l. c. = M. integrifolius.

CORDIA, Linn. Gen. ed. I. 89 (1737). *BORAGINEAE*, Benth. & Hook. f. ii. 883.
ACNADENA, CARPIPHRA, COILANTHERA, COLOCOCCA, ECTEMIS, NOVELLA, PILOISA, QUARENA, TOPIARIS, TOQUERA, Rafin. Sylva Tellur. 39-48 (1838).
BOBELLIA, Neck. Elem. i. 275, 276 (1790).
CORDIADA, Vell. Fl. Flum. 98 (1825); ii. t. 156.
CERDANA, Ruiz & Pav. Prod. 37. t. 6 (1794).
FIRENSIA, Neck. Elem. i. 275, 276 (1790).
GERASCANTHUS, P. Br. Hist. Jamaic. 170. t. 29. f. 3 (1756).
GYMALON, A. DC. Prod. ix. 468 (1845).
HEMIOTMENIA, Griff. in Calc. Journ. Nat. Hist. iii. (1842) 366.
MACHILIA, Vand. Fl. Lusit. et Bras. Spec. 14 (1788).
MACRIA, Tenore, in Mem. Soc. Ital. Modem. xxiv. (1845) 366.
MYXIA, Cosmlp. ex Linn. Hort. Cliff. 68 (1737) in syn.
PARADIGMA, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 96. t. 8.
PAVONIA, Domb. ex Lam. Tabl. Encyc. i. 421 (1800).

CORDIA:—
PLETHOSTEPHIA, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 92. t. 8.
RHARDOCALYX, Lindl. Veg. Kingd. 629 (1847).
SERESTENA, [Dill.] Gaertn. Fruct. i. 264. t. 76. f. 1 (1788).
VARRONIA, P. Br. Hist. Jamaic. 172 (1756); L. n. Syst. ed. X. 916 (1759).
abyssinica, R. Br. in Salt, Abyss. App. 65; ed. Germ. 473 = C. Myxa.
acuminata, Wall. Cat. n. 896; et in Roxb. Fl. Ind. ed. Carey, ii. 839 = Ehretia Wallichiana.
acutifolia, *Fresen. in Mart. Fl. Bras.* viii. 11.—Am. trop.
adnata, *DC. Prod.* ix. 493.—Cuba!
affinis, *Fresen. in Mart. Fl. Bras.* viii. 22.—Am. trop.
africana, Lam. Illustr. n. 1286 = C. Myxa.
alba, *Roem. & Schult. Syst.* iv. 466.—Ind. occ.
alliodora, *Cham. in Linnaea*, viii. (1833) 121.—Am. trop.
alnifolia, *Hornem. Hort. Hafn.* i. 227.—Hab. ?
alnifolia, Roem & Schult. Syst. iv. 801 = serratifolia.
ambigua, *Cham. in Linnaea*, v. (1830) 115.—Mexic.
ambigua, Cham. l. c. viii. (1833) 125 = anabaptista.
amplifolia, *A. DC. in DC. Prod.* ix. 481.—Maurit. ?
anabaptista, *Cham. in Linnaea*, viii. (1833) 512.—Bras. angiocarpa, *A. Rich. Fl. Cub. Fanerog.* ii. 110.—Cuba
angustifolia, *Roem. & Schult. Syst.* iv. 460.—Ind. occ.
angustifolia, Roxb. Hort. Beng. 17; Fl. Ind. i. 595 = Rothii.
aspera, *Forst. f. Prod.* 18. n. 109.—Austral.; Ins. Pacif.
asperrima, *DC. Prod.* ix. 498.—Jamaic.
asperrima, Spreng. Syst. i. 649 = Hyptis membranacea.
asteroptera, *Mart. ex Fresen. in Mart. Fl. Bras.* viii. 5.—Am. trop.
auricularis, Salisb. Prod. 111 = nodosa.
Azuleti, *DC. Prod.* ix. 490.—Guiana.
bahiensis, DC. l. c. ix. 489 = cylindristachya.
Banalo, Blanco, Fl. Filip. ed. I. 124 = subcordata.
bantamensis, *Blume, Bijdr.* 843.—Java.
Bertero, Spreng. ex Gœrdt. Fl. Brit. W. Ind. 480 = salviifolia.
Beurreria, Willd. Sp. Pl. i. 1078, in syn. = Bourreria succulenta.
bicolor, *A. DC. ex DC. Prod.* ix. 486.—Guiana.
bifurcata, *Roem. & Schult. Syst.* iv. 466.—Peruv.
Blanchetii, *DC. Prod.* ix. 477.—Am. trop.
bogotensis, *Benth. Pl. Hartw.* 340.—N. Granat.
Bonsieri, *A. DC. in DC. Prod* ix. 478.—Mexic.
Bonplandiana, H. B. & K. Nov. Gen et Sp. iii. 77 = sco.
Bonplandii, *Roem. & Schult. Syst.* iv. 462.—Am. austr.
Bourreria, Linn. Syst. ed. X. 936 = Bourreria succulenta.
brachypoda, *DC. Prod.* ix. 487.—Am. trop.
brachytricha, *Fresen. in Mart. Fl. Bras.* viii. 18.—Am. trop.
bracteata, *DC. Prod.* ix. 472.—Cuba.
brevispicata, *Mart. & Gal. in Bull. Acad. Brux.* xi. ii. (1844) 331.—Mexic.
Brownii, DC. Prod. ix. 499 = C. Myxa.
bullata, Coem. & Schult. Syst. iv. 462 = globosa.
buxifolia, *Juss. in Poir. Encyc.* vii. 47.—Ind. occ.
caffra, *Sond. in Linnaea*, xxiii. (1850) 81.—Afr. austr.
Calocoma, Colloccoca.
calocephala, *Cham. in Linnaea*, iv. (1829) 488.—Bras.
calocoma, *Miq. Fl. Ind. Bat.* ii. 916.—Java.
calophylla, *Vahl, Eclog. Am.* iii. 8.—Guiana.
calyptrata, Bert. ex Spreng. Syst. i. 649 = alba.
campanulata, Roxb. Hort. Beng. 17; Fl. Ind. i. 593 = subcordata.
campestris, *Warm. in Kjoeb. Vidensh. Meddel.* (1867) 12.—Bras.
cana, *Mart. & Gal. in Bull. Acad. Brux.* xi. ii. (1844) 331.—Mexic.
candida, *Vell. Fl. Flum.* 98; ii. t. 155.—Bras.
canescens, *H. B. & K. Nov. Gen. et Sp.* iii. 73.—N. Granat.
canescens, Willd. ex Roem. & Schult. Syst. iv. 799 = polystachya.
carassanana, *DC. Prod.* ix. 489.—Venezuela.
Cerdana, Roem. & Schult. Syst. iv. 467 = alliodora.
chamædrifolia, DC. Prod. ix. 499, in syn. = mirabilis flora.
chamædryoides, *Roem. & Schult. Syst.* iv. 801.—Hab. ?
Chamissoniana, *Steud. Nom.* ed. II. i. 417.—Bras.
cinerascens, DC. Prod. ix. 492 = cylindristachya.

CORDIA:—
collocoides, *Warm. in Kjoeb. Vidensh. Meddel.* (1867) 4.—Bras.
Collococa, Aubl. Pl. Gui. i. 219 = nodosa.
Collococca, *Linn. Sp. Pl.* ed. II. 274.—Jamaic.
Collococca, Sw. ex Griseb. Fl. Br. Wist. Ind. 479 = nitida.
copulata, *Poir. in Dict. Sc. Nat.* x. 410.—Java.
corchorifolia, *A. DC. in DC. Prod.* ix. 496.—Am. trop.
cordata, Steud. Nom. ed. II. i. 417 = grandis.
cordifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 70.—N. Granat.
cordifolia, Wall. Cat. n. 894 = grandis.
cormandeliana, Retz. ex DC. Prod. ix. 509 = Ehretia buxifolia.
coryifolia, Willd. ex Roem. & Schult. Syst. iv. 461 = alba.
corymbosa, G. Don, Gen. Syst. iv. 383 = monosperma.
corymbosa, Willd. ex Steud. Nom. ed. II. i. 417 = alba
crenata, *Delile, Fl. Egypte*, 195. t. 20.—Aegypt.
crenata, Roem. & Schult. Syst. iv. 468 = lantanoides.
crenulata, *A. DC. in DC. Prod.* ix. 492.—Mexic.
crispiflora, *DC. l. c.* 476.—Mexic.
cujabensis, *Menso & Lhotsky, ex Cham. in Linnaea*, viii. (1833) 12.—Bras.
Cuminginna, *Vidal, Phan. Cuming. Philipp.* 187.—Ins. Philipp.
cuneata, Heyne, ex DC. Prod. ix. 480 = Rothii.
cuneiformis, *DC. l. c.* 492.—Venezuela.
curassavica, *Roem. & Schult Syst.* iv. 460.—Am. austr.
cylindristachya, *Roem. & Schult. l. c.* 459.—Am. trop.
dasycephala, *H. B. & K. Nov. Gen. et Sp.* iii. 76.—Mexic.
decandra, *Hook. & Arn Bot. Beech. Voy.* i. 38.—Chili.
dentata, Vahl, Eclog. Am. iii. 8 = alba.
dichotoma, Forst. f. Prod. 18. n 110 = C. Myxa.
digynia, *Vell. Fl. Flum.* 97; ii. t. 153.—Bras.
Dillenii, *Spreng. Syst.* i. 645.—Ins. Bahama.
diodea, *Roy. ex DC. Prod.* ix. 481.—Madag.
diosca, Thonn. ex DC. l. c. 480 = guineensis.
diospyrifolia, *Cham. in Linnaea*, iv. (1829) 477.—Bras.
discolor, *Cham. l. c.* 483.—Bras.
divaricata, *H. B. & K. Nov. Gen. et Sp.* iii. 74.—Am. austr.
diversa, Thw. Enum. Pl. Zeyl. 214 = monoica.
diversifolia, *Pav. ex DC. Prod.* ix. 474.—Mexic.
dodecandra, *DC. l. c.* 478.—Mexic.
domestica, Boutou, ex Boj. Hort. Maurit. 225 = amplifolia.
domestica, Roth, Nov. Sp. Pl. 128 = C. Myxa.
domingensis, *Lam. Illustr.* n. 1900.—Ins. S. Doming.
dubiosa, *Blume, Bijdr.* 844.—Java.
ecalyculata, *Vell. Fl. Flum.* 96; ii. t. 149.—Bras.
egyptiaca, *Rafin. Sylva Tellur.* 37.—Aegypt.
ehretioides, Lam. Illustr. n. 1902 = C. Collococca vel nitida.
elæagnoides, *DC. Prod.* ix. 474.—Mexic.
elliptica, Sieber, ex Griseb. Fl. Brit. W. Ind. 479 = laevigata.
elliptica, *Sw. Prod. Veg. Ind. Occ.* 47.—Ind. occ.
erythrococca, *Griseb. Cat. Pl. Cub.* 208.—Cuba.
exaltata, *Lam. Illustr.* n. 1919.—Guiana.
excelsa, *A. DC. in DC. Prod.* ix. 473.—Am. trop.
ferruginea, *Roem. & Schult. Syst.* iv. 458; *H. B. & K. Nov. Gen. et Sp.* iii. 71.—Mexic.; N. Granat.
Finlaysoniana, *Wall. Cat.* n. 7010.—Ind. or.
flavescens, *Aubl. Pl. Gui.* i. 226. t. 89.—Guiana.
floribunda, *Spreng. Syst.* i. 652.—Am. austr.
floridana, Nutt. Gen. ii. 107 = Bourreria havanensis.
foliosa, *Mart. & Gal. in Bull. Acad. Brux.* xi. ii. (1844) 330.—Mexic.
formicarum, Roem. & Schult. Syst. iv. 800 = nodosa.
fragrantissima, *Kurz, For. Fl. Brit. Burma*, ii. 207.—Ind. or.; Burma.
Francisci, *Tenore, Cat. Orto Nap.* (1845) 88.—Hab. ?
frondosa, Schott, in Spreng. Syst. iv. App. 403 = alliodora.
fulvosa, *Wight, Ic.* iv. t. 1260.—Ind. or.
Galeottiana, *A. Rich. Fl. Cub. Fanerog.* ii. 109.—Cuba.
Gerascanthus, H. B. & K. Nov. Gen. et Sp. iii. 69 = tinifolia.
Gerascanthus, *Linn. Syst.* ed. X. 936.—Mexic.
Gerascanthus, Sw. ex Griseb. Fl. Brit. W. Ind. 478 = gerascanthoides.
gerascanthoides, *H. B. & K. Nov. Gen. et Sp.* iii. 69.—Cuba.
gerascanthoides, Rich. ex Griseb. Fl. Brit. W. Ind. 478 = A. Gerascanthus.

CORDIA :—
Gerascanthus, Jacq. Select. Am. 49. t. 175 = A. Gerascanthus.
glabra, *Chem. in Linnaea*, viii. (1833) 194.—Bras.
glabra, Lion. Sp. Pl. 191 = C. Collococca.
glabrata, *A. DC. Prod.* ix. 478.—Am. trop.
glandulosa, *Presen. in Mart. Fl. Bras.* viii. 19.—Am. trop.
globosa, Andr. ex DC. Prod. ix. 497 = oxyacuna.
globosa, *H. B. & K. Nov. Gen. et Sp.* iii. 76.—Ind. occ.; Am. centr.
glomerata, *Lem. Yard. Fleur.* iv. (1854) t. 326.—Bras.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 77.—Am. austr.
grandifolia, *DC. Prod.* ix. 475.—Am. trop.
grandis, Cham. in Linnæa, iv. (1829) 475 = grandifolia.
grandis, *Roxb. Hort. Beng.* 17; *Fl. Ind.* i. 593.—Ind. or.
grandis, Wall. Cat. n. 892 B = obliqua.
graveolens, *H. B. & K. Nov. Gen. et Sp.* iii. 74.—Am. austr.
Greggii, *Torr. in Bot. Mex. Bound.* 135.—Calif.; Mexic.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 139.—Malaya.
guayaquilensis, *DC. Prod.* ix. 496.—Ecuador.
guazumæfolia, Roem. & Schult. Syst. iv. 468 = urticifolia.
guianensis, *Roem. & Schult. l. c.* 460.—Guiana.
guineensis, *Thonn. in Schum. Beskr. Guin. Pl.* 128.—Afr. trop.
Hartwisiana, *Regel, Gartenfl.* (1857) 242.—Mexic.
bebecarpa, *DC. Prod.* ix. 488.—Am. trop.
hermanniæfolia, *Cham. in Linnæa*, iv. (1829) 484.—Am. trop.
heterophylla, *Poir. in Dict. Sc. Nat.* x. 409; *Roem. & Schult. Syst.* iv. 800.—Guiana.
hexandra, Willd. ex Roem. & Schult. l. c. 799 = subcordata.
hirsuta, Willd. Sp. Pl. i. 1076 = nodosa.
hispida, *Benth. Bot. Voy. Sulph.* 139.—Mexic.
hispidissima, *DC. Prod.* ix. 475.—Am. trop.
Humboldtii, Willd. ex Roem. & Schult. Syst. iv. 801 = floribunda.
hypoleuca, *DC. Prod.* ix. 479.—Am. trop.
Ignotæ, Blanco, Fl. Filip. ed. II. 88 = subcordata.
incana, Royle, Illustr. Bot. Himal. 306, nom en = vestita.
indica, Lam. Encyc. vii. 49; Illustr. n. 1913 = C. Myxa.
insignis, *Cham. in Linnæa*, viii. (1833) 123.—Bras.
integrifolia, *Roem. & Schult. Syst.* iv. 461.—Ins. S. Doming.
intermedia, *Presen. in Mart. Fl. Bras.* viii. 8.—Am. trop.
interrupta, *DC. Prod.* ix. 491.—Guiana.
intricata, *Wright, in Sauv. Fl. Cub.* 109.—Cuba.
ipomœæfolia, *Hook. Bot. Mag.* t. 5027.—Hab.?
isiocarpa, F. Muell. Fragm. i. 59 = C. Myxa.
juglandifolia, *Jacq. Enum. Pl. Carib.* 14.—Ins. S. Doming.
lacerata, F. Muell. Fragm. v. 198 = aspera.
laevigata, *Lam. Illustr.* n. 1912.—Ind. occ.
laevis, *Jacq. Hort. Schœnb.* i. t. 40.—N. Granat.
lanata, H. B. & K. Nov. Gen. et Sp. iii. 73 = tomentosa.
lanceolata, *H. B. & K. l. c.* 75.—N. Granat.
lantanoides, *Spreng. Syst.* i. 858.—Peruv.
lapensis, *Warm. in Kjoesb. Vidensk. Meddel.* (1867) 9.—Bras.
latifolia, Cham. in *Linnæa*, viii. (1833) 126.—Bras.
latifolia, Roxb. Hort. Beng. 17; Fl. Ind. i. 588 = obliqua.
latifolia, Wall. Cat. n. 893 = C. Myxa.
Latora, *Buch.-Ham. ex Royle, Illustr. Bot. Himal.* 306.—Ind. or.
laxiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 73.—N. Granat.
Leschenaultii, DC. Prod. ix. 483 = oblongifolia.
leucocalyx, *Presen. in Mart. Fl. Bras.* viii. 23.—Am. trop.
leucocephala, *Moric. Pl. Nouv. Am.* 148. t. 88.—Bras.
leucocoma, *Miq. Fl. Ind. Bat.* ii. 916.—Malaya.
leucophlyctis, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 199.—Ins. Galap.
leucosebestera, *Griseb. Cat. Pl. Cub.* 208.—Cuba.
Lima, *Roem. & Schult. Syst.* iv. 475.—Ins. S. Doming.
linearis, *A. DC. in DC. Prod.* ix. 493.—Mexic.
linearis, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 199.—Ins. Galap.

CORDIA :—
linnæta, Roem. & Schult. Syst. iv. 464 = ulmifolia.
longifolia, *A. DC. in DC. Prod.* ix. 485.—Am. trop.
Lourairi, *Roem. & Schult. Syst.* iv. 465.—China.
lutea, *Lam. Tabl. Encyc.* i. 431.—Peruv.
Macleodii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 128.—Ind. or.
macrocephala, *H. B. & K. Nov. Gen. et Sp.* iii. 77.—Peruv.
macrophylla, *Linn. Sp. Pl.* ed. II. 378.—Ind. occ.
macrophylla, Vahl, ex Griseb. Fl. Brit. W. Ind. 480 = sulcata.
macrostachya, *Roem. & Schult. Syst.* iv. 461.—N. Granat.
macrostachya, Spreng. Syst. i. 852 = cylindristachya.
magnoliæfolia, *Cham. in Linnæa*, iv. (1829) 476.—Bras.
mariquitensis, *H. B. & K. Nov. Gen. et Sp.* iii. 75.—N. Granat.
Martii, DC. Prod. ix. 471 = insignis.
martinicensis, Link, Enum. Hort. Berol. i. 162 = curassavica.
martinicensis, *Roem. & Schult. Syst.* iv. 461.—Ins. Martinic.
martinicensis, Sieber, ex Griseb. Fl. Brit. W. Ind. 480 = cylindristachya.
McLeodii, vide Macleodii.
melanocarpa, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 960.—Guiana.
membranacea, *A. DC. ex DC. Prod.* ix. 484.—Ecuador.
micrantha, *Sw. Prod. Veg. Ind. Occ.* 47.—Jamaica.
micrantha, Sw. l. c. pro parte, ex Griseb. Fl. Brit. W. Ind. 479 = C. Collococca.
microcephala, *Willd. in Roem. & Schult. Syst.* iv. 801.—Venezuela.
microphylla, *Roem. & Schult. l. c.* 468.—Ins. S. Doming.
mirabilifolia, *A. DC. in DC. Prod.* ix. 499.—Cuba.
mirabiloides, *Roem. & Schult. Syst.* iv. 465.—Ins. S. Doming.
miranda, *DC. Prod.* ix. 475.—Am. trop.
molinæfolia, *Kosh. Hort. Beng.* 84; *Fl. Ind.* ii. 594.—Ins. Moluce.
monoica, Boj. Hort. Maurit. 235 = ovalis.
monoica, *Roxb. Fl. Corom.* i. 48. t. 58.—Ind. or.
monosperma, *Roem. & Schult. Syst.* iv. 468.—Venezuela.
mucronata, *Presen. in Mart. Fl. Bras.* viii. 9.—Am. trop.
multiplicata, *Cham. in Linnæa*, iv. (1829) 499.—Bras.
Munneco, *H. B. & K. Nov. Gen. et Sp.* vii. 207.—Venezuela.
Myxa, Forsk. Fl. Aegypt. Arab. p. lxiii = crenata.
Myxa, *Linn. Sp. Pl.* 190.—As. trop.; Austral.
Myxa, Tiw. Enum. Pl. Zeyl. 218 = obliqua.
Neowedlana, *A. DC. in DC. Prod.* ix. 495.—Am. trop.
nervosa, *Lam. Illustr.* n. 1906.—Guiana.
nitida, *Vahl, in West, Bidrag Ste-Croix*, 275.—Ind. occ.
nitida, *Wild. in Roem. & Schult. Syst.* iv. 800.—Hab.?
nives, *Presen. in Mart. Fl. Bras.* viii. 26.—Am. trop.
nodosa, *Lam. Illustr.* n. 1905.—Guiana.
oaxacana, *DC. Prod.* ix. 497.—Mexic.
obliqua, Heyne, ex DC. l. c. 479 = Wallichii.
obliqua, H. B. & K. Nov. Gen. et Sp. iii. 74 = peruviana.
oblique, *Vell. Fl. Flum.* 97; ii. t. 150.—Bras.
obliqua, *Willd. Sp. Pl.* i. 1073.—Ind. or.
oblongifolia, *Hochst. ex DC. Prod.* ix. 480, nota.—Abyssin.
oblongifolia, *Thw. Enum. Pl. Zeyl.* 214.—Zeylan.
obovata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 80.—Ins. Socotra.
obscura, *Cham. in Linnæa*, xiii. (1839) 480.—Bras.
obtusa, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 80.—Ins. Socotra.
ochnacea, *DC. Prod.* ix. 477.—Am. trop.
ocnodes, *A. DC. l. c.*—Ind. or.
officinalis, Lam. Illustr. ii. 490 = C. Myxa.
oliteria, *Blanco, Fl. Filip.* ed. I. 128.—Ins. Philipp.
oncocalyx, *F. Allem. in Trab. Comm. Sc. Expl. Bot.* i. (1862) 11.—Bras.
orientalis, R. Br. Prod. 496 = subcordata.
ovalis, *R. Br. in Salt, Abyss. App.* 66.—Abyssin.
oxyphylla, *DC. Prod.* ix. 492.—Guiana.
paniculaire, *Rudge, Pl. Guian.* 30. t. 46.—Guiana.
paniculata, Roth, Nov. Pl. Sp. 138 = C. Myxa.

CORDIA :—
paniculata, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 278 = Wilbrandia paniculata.
parvifolia, Orteg. Hort. Matr. 85 = serratifolia.
parvifolia, *A. DC. in DC. Prod.* ix. 498.—Mexic.
parvifolia, Hemsl. Biol. Centr. Am. Bot. ii. 369, in syn. = serratifolia.
Patagonula, Ait. Hort. Kew. i. 259 = Patagonula americana.
patens, *H. B. & K. Nov. Gen. et Sp.* iii. 75.—Am. austr.
patens, Miq. ex Griseb. Fl. Brit. W. Ind. 480 = ulmifolia.
paucident ta, *Presen. in Mart. Fl. Bras.* viii. 25.—Bras.
pedunculosa, *Griseb. Cat. Pl. Cub.* 208.—Cuba.
Pelida, Perr. ex DC. Prod. ix. 480 = senegalensis.
pellita, Willd. ex Roem. & Schult. Syst. iv. 800 = Tournefortia hirsutissima.
Perrottetii, *A. DC. Prod.* ix. 482.—Ind. or.
peruviana, *Roem. & Schult. Syst.* iv. 459.—Peruv.
petiolata, Hort. ex Steud. Nom. ed. II. i. 416 = Ehretia petiolaris.
Petta-poligoret, Heyne, ex Roth, Nov. Pl. Sp. 125 = paniculata.
plushieuis, *Presen. in Mart. Fl. Bras.* viii. 9.—Bras.
pilulifera, Willd. ex Roem. & Schult. Syst. iv. 801 = globosa.
platyphylla, Steud. Nom. ed. II. i. 418 = latifolia.
podocephala, *Torr. in Bot. Mex. Bound.* 135.—Texas; Mexic.
Poeppigii, *DC. Prod.* ix. 492.—Am. austr.
poliophylla, *Presen. in Mart. Fl. Bras.* viii. 26.—Am. trop.
polyantha, *Benth. Pl. Hartw.* 121.—Ecuador.
polygama, Roxb. Hort. Beng. 17; Fl. Ind. i. 594 = monoica.
polygama, Wall. Cat. n. 892, partim = fulvosa.
polystachya, *H. B. & K. Nov. Gen. et Sp.* iii. 78.—Venezuela.
portoricensis, Spreng. Neue Entdeck. ii. 127 = cylindristachya.
prionodes, Voigt, Hort. Suburb. Calc. 441 = octandra.
pubescens, *Willd. ex Roem. & Schult. Syst.* iv. 808.—Venezuela.
pyrifolia, Willd. ex Roem. & Schult. l. c. 802 = ferruginea.
querciifolia, *Klotzsch, in Peters, Reise Mossamb. Bot.* 247.—Afr. trop.
Radula, Spreng. Syst. i. 854 = Lima.
ramiflora, *A. Rich. Fl. Cub. Fanerog.* ii. 110.—Cuba.
reflexa, *Willd. ex Roem. & Schult. Syst.* iv. 800.—Ind. or.
reticulata, Roth, Nov. Pl. Sp. 124 = Rothii.
reticulata, *Vahl, Eclog. Am.* iii. 5.—Ins. Montserrat.
retusa, Vahl, Symb. Bot. ii. 43 = Ehretia buxifolia.
revoluta, *Hook. f. in Trans. Linn. Soc. xx.* (1851) 199.—Ins. Galap.
riparia, *H. B. & K. Nov. Gen. et Sp.* iii. 71.—N. Granat.
rotata, *Miq. ex DC. Prod.* ix. 488.—Mexic.
Rothii, *Roem. & Schult. Syst.* iv. 796.—Ind. or.
rotundifolia, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 139 = fragrantissima.
rotundifolia, *Ruiz & Pav. Fl. Per.* ii. 24. t. 148.—Panama.
Roxburghii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 138.—Zeylan.
rubra, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 92 = C. Myxa.
rufa, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 960.—Guiana.
rufescens, *A. DC. in DC. Prod.* ix. 476.—Am. trop.
rugosa, Willd. ex Roem. & Schult. Syst. iv. 801 = cumanensis.
Rumphii, Brume, Bijdr. 846 = subcordata.
salicifolia, *Cham. in Linnæa*, iv. (1829) 481.—Bras.
salicina, *DC. Prod.* ix. 492.—Am. trop.
sebestenoides, H. B. & K. Nov. Gen. et Sp. iii. 72 = sylvestris.
sebestilefa, *Juss. in Poir. Suppl.* vii. 46.—Ins. Guadaloup.
Selowayana, [Benth. ex] Griseb. Fl. Brit. W. Ind. 480 = ulmifolia.
Sellowiana, *DC. Prod.* ix. 494.—Bras.
serrulata, Lam. Illustr. n. 1907 = Sebestena.
sessilifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 71.—Peruv.

CORDIA :—
scabra, *Desf. Tabl.* ed. I. 71.—Ind. or.
scabrida, *Mart. ex Fresen. in Mart. Fl. Bras.* viii. 11. —Bras.
scabrifolia, [Benth. ex] Griseb. Fl. Brit. W. Ind. 479 = paniculata.
scabrifolia, *A. DC. in DC. Prod.* ix. 485.—Guiana.
scandens, *Martin, ex Poir. in Dict. Sc. Nat.* x. 410.— Guiana.
Schomburgkii, *DC. Prod.* ix. 490.—Guiana.
Schottiana, *Fresen. in Mart. Fl. Bras.* viii. 7.—Am. trop.
Sconleri, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 200. —Ins. Galap.
Sebestena, Andr. Bot. Rep. iii. t. 157 = scabra.
Sebestena, Forsk. Fl. Aegypt. Arab. p. lxiii = C. Myxa.
Sebestena, Forst. f. Prod. 15 = subcordata.
Sebestena, *Linn. Sp. Pl.* 190.—Am. trop.
Sebestena, Willd. ex Spreng. Syst. I. 649 = Dillenii.
Sellowiana, *Cham. in Linnaea*, iv. (1829) 479.—Bras.
serrgelensis, *Yuss. in Petr. Encyc.* vii. 47.—Afr. occ.
sericicalyx, *A. DC. in DC. Prod.* ix. 485.—Guiana.
serrata, *Yuss. ex Petr. Encyc.* vii. 41.—Ind. or.
serrata, Roxb. Hort. Beng. 17; Fl. Ind. i. 591 = octandra.
serratifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 76.— Mexic.
sessilifolia. *Cham. in Linnaea*, iv. (1829) 488.—Bras.
sinensis, *Lam. Illustr.* n. 1914.—China.
speciosa. Saliab. Prod. 111 = C. Sebestena.
spectabilis, *Steud. Nom.* ed. II. i. 419.—Ind. or.
sphaerocephala, Willd. ex Roem. & Schult. Syst. iv. 801 = dasycephala.
spicata, Willd. ex Roem. & Schult. l. c. 799 = graveolens.
spinescens, *Linn. Mant.* 206.—Ind. or.
Sprengelii, *DC. Prod.* iv. 473.—Ins. Fiji.
striata, *Fresen. in Mart. Fl. Bras.* viii. 28.—Am. trop.
strigosa, *Spreng. Neue Entdeck.* iii. 80.—Ind. occ.
suaveolens, *Blume, Bijdr.* 843.—Java.
subcordata, *Lam. Illustr.* n. 1899.—As. et Austral. trop.
subdentata, *Miq. Fl. Ind. Bat. Suppl.* 571.—Sumatra.
subopposita, DC. Prod. ix. 480 = Rothii.
subpubescens, *Decne. Herb. Timor.* 67.—Ins. Timor.
sulcata, *DC. Prod.* ix. 488.—Ind. or.
superba, *Cham. in Linnaea*, iv. (1829) 474.—Bras.
taguahyensis, *Vell. Fl. Flum.* 98; ii. t. 154.—Bras.
tenuifolia, *Bertol. Misc. Bot. Suri.* 13. t. 1.—Guatemala.
tetrandra. *Aubl. Pl. Gui.* i. 222. t. 87.—Guiana.
tetraphylla, *Aubl. l. c.* 224. t. 88.—Guiana.
Thibaudiana, *DC. Prod.* ix. 489.—Am. austr.
thyrsiflora, Sieb. & Zucc. Fl. jap. Fam. Nat. ii. 26 = Ehretia acuminata.
timorensis, Span. in Linnaea, xv. (1841) 334 = subpubescens.
tinifolia, *Willd. ex Roem. & Schult. Syst.* iv. 800.—Am. austr.
tomentosa, Cham. & Schlecht. in Linnaea, iv. (1829) 479 = Chamissoniana.
tomentosa, *Roem. & Schult. Syst.* iv. 459.—Am. austr.
tomentosa, Wall. in Roxb. Fl. Ind. ed Carey, ii. 339 = obliqua.
Toqueve, *Aubl. Pl. Gui.* i. 228. t. 90.—Am. trop.
Toqueve, Sieber, ex Griseb. Fl. Brit. W. Ind. 480 = sulcata.
trachyphylla, *Mart. Herb. Fl. Bras.* n. 412.—Bras.
tremula, *Griseb. Fl. Brit. W. Ind.* 479.—Ind. occ.
trichocladea, *DC. Prod.* ix. 474.—Am. trop.
trichostemon, *DC. l. c.* 482.—Ins. Timor.
trichotoma, *Vell. ex Steud. Nom.* ed. II. i. 419.— Bras.
truncata, *Fresen. in Mart. Fl. Bras.* viii. 25.—Am. trop.
ulmifolia, *Yuss. in Dem.-Cours. Bot. Cult.* ed. I. ii. 146.—Panama; Ind. occ.
ulmifolia, Spreng. Syst. i. 653 = monosperma.
umbraculifera, *DC. Prod.* ix. 484.—Am. trop.
urticifolia, *Cham. in Linnaea*, iv. (1829) 483.—Bras.
Valenzuelana, *A. Rich. Fl. Cub. Fanerog.* ii. 110.— Cuba.
velutina, Mart. Herb. Fl. Bras. n. 268 = sujabensis.
verbenacea, *DC. Prod.* ix. 491.—Am. trop.
vestita, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 128.—Reg. Himal.
vilmoriniana, *Zipp. ex Span. in Linnaea*, xv. (1841) 336, nomen.—Malaya.
villicaulis, *Fresen. in Mart. Fl. Bras.* viii. 24.—Am. trop.
villosa, *Spreng. Neue Entdeck.* iii. 81.—Ins. S. Doming.
Wallichii, G. Don, Gen. Syst. iv. 379 = obliqua.
Wickstroemii, *Steud. Nom.* ed. II. i. 419.—Ins. Guadalup.

CORDIADA, Vell. Fl. Flum. 98 (1825); ii. t. 156 = **Cordia**, Linn. (Boragin.).
trichotoma, Vell. li. cc. = Cordia trichotoma.

CORDIERA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 222. t. 20 = **Alibertia**, A. Rich. (Rubiac.).
acuminata, Benth. in Hook. Journ. Bot. iii. (1841) 291.—Guiana.
latifolia, Benth. l. c.—Bras.
trifiora, DC. Prod. iv. 445.—Guiana.
trifolia, Steud. Nom. ed. II. i. 419, sphalm. = praec.

CORDIOPSIS, Desv. in Ham. Prod. Fl. Ind. Occ. 23 (1825) = **Cordia**, Linn. (Boragin.).
mirabilifiora, DC. Prod. ix. 499 = Cordia mirabilifiora.
mirabilioides, Desv. in Ham. Prod. Fl. Ind. Occ. 23 = Cordia mirabilifiora.
monosperma, Ham. ex Heynh. Nom. ii. 160 = Cordia monosperma.

CORDULA, Rafin. Fl. Tellur. iv. 46 (1836) = **Cypripedium**, Linn. (Orchid.).
insignis, Rafin. l. c. = Cyprip. insigne.

CORDYLA, Blume, Bijdr. 416 (1825) = **Pogonia**, Juss. (Orchid.).
concolor, Blume, l. c. = P. concolor.
discolor, Blume, l. c. = P. discolor.

CORDYLA, Lour. Fl. Cochinch. 411 (1790). *LEGUMINOSAE*, Benth. & Hook. f. i. 562.
CALYCANDRA, Lepr. ex A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 30; et 232. t. 9 (1832).
africana, *Lour. l. c.* 412.—Afr. trop.
Richardi, Planch. in App. Speke, Journ. Disc. Nile, 632 = africana.

CORDYLANTHUS, Blume, Mus. Bot. Lugd. Bat. ii. 27. t. 3. 1852 [1856] = **Homalium**, Jacq. (Samydac.).
frutescens, Blume, l. c. 28 = H. Cordylanthus.

CORDYLANTHUS, Nutt. ex Benth. in DC. Prod. x. 597 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 974.
ADENOSTEGIA, Benth. in Lindl. Nat. Syst. ed. II. 445 (1826).
CHLOROPYRON, Behr, in Proc. Calif. Acad. i. (1855) 61.
canescens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 383. —Am. bor. occ.
capitatus, *Nutt. ex Benth. in DC. Prod.* x. 597.—Am. bor.
filifolius, *Nutt. ex Benth. l. c.*—Calif.
Kingii, *S. Wats. in Bot. King's Exp.* 233, 459. t. 22.— Am. bor. occ.
laxiflorus, *A. Gray, in Torr. Bot. Mex. Bound.* 120.— Arizona.
maritimus, *Nutt. ex Benth. in DC. Prod.* x. 598.— Calif.
mollis, *A. Gray, in Proc. Am. Acad.* vii. (1868) 384.— Calif.
Nevinii, *A. Gray, l. c.* xvii. (1881-82) 229.—Calif.
Orcuttianus, *A. Gray, l. c.* xix. (1883) 95.—Calif.
Parryi, S. Wats. in Am. Natural. ix. (1875) 846 = canescens.
pilosus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 382. —Calif.
Pringlei, *A. Gray, l. c.* xix. (1883) 94.—Calif.
ramosus, *Nutt. ex Benth. in DC. Prod.* x. 597.—Am. bor. occ.
tenuis, *A. Gray, in Proc. Am. Acad.* vii. (1868) 383.— Am. bor. occ.
Wrightii, *A. Gray, in Torr. Bot. Mex. Bound.* 120.— Texas; N. Mexic.

CORDYLESTYLIS, Falc. in Hook. Journ. Bot. iv. (1842) 74 = **Goodyera**, R. Br. (Orchid.).
foliosa, Falc. l. c. 75 = G. procera.

CORDYLIA, Pers. Syn. ii. 260 (1807) = **Coryda**, Lour. (Legumin.).

CORDYLINE, Royen, ex Adans. Fam. ii. 54 (1763). *LILIACEAE*, Benth. & Hook. f. iii. 799.
CALODRACON, Planch. in Fl. des Serres, Sér. I. vi. (1850-51) 137.
CHARLWOODIA, Sweet, Fl. Austral. t. 18 (1827).
CONNIA, Kunth, Enum. Pl. v. 85 (1850).
DRACAENUPSIS, Planch. in Fl des Serres, Sér. I. vi. (1850-51) 110.
TAETSIA, Medic. Theodora, 82 (1786).
angustifolia, Kunth, Enum. Pl. v. 32 = stricta.
arborea, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55. —Afr. trop.
atropurpurea, Planch. in Fl. des Serres, Sér. I. vi. (1850-51) 139 = Dracaena elliptica.
australis, Endl. Prod. Fl. Ins. Norf. 29 = Baueri.
australis, *Hook. f. in Gard. Chron.* (1860) 792.— N. Zel.
australis, Nadeaud & Josan, in Mém. Soc. Sc. Nat. Cherb. xi. (1865) 127 = terminalis.
Banksii, *Hook. f. in Gard. Chron.* (1860) 792.— N. Zel.
Baueri, *Hook. f. l. c.*—Ins. Norfolc.
Botschieriana, Goepp. in Nov. Act. Nat. Cur. xxv. (1855) 55 = Dracaena concinna.
Brackenridgei, C. Koch, Wochenschr. viii. (1868) 21 = Banksii.
brasiliensis, *Planch. in Fl. des Serres*, Sér. I. vi. (1850-51) 136.—Bras.
calocoma, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 542 = australis.
cannaefolia, R. Br. Prod. 280 = terminalis.
Casanovae, *Linden. ex André, Illustr. Hortic.* xxi. (1874) 137. t. 181.—N. Hebrid.
cernua, Planch. in Fl. des Serres, Sér. I. vi. (1850-51) 111 = Dracaena reflexa.
coerulea, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55. —Hab.?
congesta, Endl. Gen. 151 = stricta.
congesta, *Steud. Nom.* ed. II. i. 419.—Austral.
Denniisoni, *Hort. ex André, Illustr. Hortic.* xix. (1872) 260.—Hab.?
dennisoniana, *Linden & André, l. c.* (1874) 185. t. 190.— Austral.
diffusa, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 330. —N. Zel.
dracaenoides, *Kunth, in Abh. Acad. Berl.* (1842) 80.— Bras.
ensifolia, Planch. in Fl. des Serres, Sér. I. vi. (1850-51) 139 = Dianella ensifolia.
erythrorachis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 541 = Baueri.
Eschscholziana, Mart. in Schult. f. Syst. vii. 847, 1677 = terminalis.
ferrea, C. Koch, Wochenschr. xiv. (1871) 12 = terminalis.
flabelliformis, *C. Koch, Allg. Gartenz.* (1858) 254.—Ins. Maurit.
flexuosa, *Miq. Fl. Ind. Bat.* iii. 558.—Malaya.
fruticosa, *C. Koch, Allg. Gartenz.* (1858) 254.— Ins. Maurit.
Fontanesiana, Goepp. in Nov. Act. Nat. Cur. xxv. (1855) 54. t. 3 = Dracaena Fontanesiana.
Forsteri, F. Muell. Sel. Pl. Indust. Cult. 58 = australis.
fragrans, *Planch. in Fl. des Serres*, Sér. I. vi. (1850-51) 111, 136.—Afr. trop.
fruticosa, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 53.—Java.
glorioso, Linden & André, Illustr. Hortic. xx. (1873) t. 125-6 = terminalis.
Guilfoylei, Hort. Linden, ex Lem. Illustr. Hortic. xvi. (1869) t. 600 = terminalis.
Hasgeana, *C. Koch, Wochenschr.* (1867) 195.— Austral.
hedychioides, F. Muell. Fragm. v. 196 = terminalis.
heliconiaefolia, Otto & Dietr. in Allg. Gartenz. (1836) 33 = terminalis.
hemichrysa, *Comm. ex Lam. Encyc.* ii. 276.—Ins. Maurit.
Hoolbrenkeana, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55.—N. Zel.
Hookeri, *T. Kirk, in Trans. N. Z. Inst.* vi. (1874) 346.—N. Zel.
humilis, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55. —Hab.?

CORDYLINE:—
indivisa, Regel, Gartenfl. (1859) 331 = australis.
indivisa, *Steud. Nom.* ed. II. i. 419.—N. Zel.
Youngwinii, Kunth, in Abh. Acad. Berl. (1843) 80 = terminalis.
javanica, Klotzsch, ex Kunth, Enum. Pl. v. 26 = terminalis.
lentiginosa, Linden & André, Illustr. Hortic. xvii. (1870) t. 25 = australis.
lineariflola, *Miq. Fl. Ind. Bat. Suppl.* 610.—Sumatra.
longifolia, Benth. Pl. Hartw. 53 = Nolina Hartwegiana.
macrophylla, *C. Koch, Allg. Gartenz.* (1858) 254.—Ins. Mascar.
maculata, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 152 = Dracæna elliptica.
madagascariensis, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 58.—Madag.
Mannerr-Suttoniæ, F. Muell. Fragm. v. 195 = terminalis.
marginata, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55.—Madag.; Ins. Borb.
metallica, Hort. Dallière, in Belg. Hortic. xxii. (1872) 358. t. 26 = terminalis.
micrantha, Baker, in Gard. Chron. (1880) ii. 243 = Brocchinia cordylinoides.
Murchisoniæ, F. Muell. Fragm. v. 195 = Haageana.
neo-caledonica, *Linden, Cat.* n. 97 (1877) 4.—N. Caled.
nobilis, C. Koch, Allg. Gartenz. (1858) 254 = terminalis.
nutans, *Hort. ex Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55, nomen.—Hab.?
obtecta, Baker, in Journ. Linn. Soc. xiv. (1875) 548 = Baueri.
odorata, C. Koch, in C. Koch & Fint. Wochenschr. (1858) 97 = dracænoides.
ovata, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 111, 156 = Dracæna ovata.
oxyphylla, *Miq. Fl. Ind. Bat. Suppl.* 609.—Sumatra.
parvifolia, H. B. & K. Nov. Gen. et Sp. i. 259 = Nolina parvifolia.
Pumilio, *Hook. f. in Gard. Chron.* (1860) 792.—N. Zel.
Reali, *Nichols. Dict. Gard.* i. 374.—Hab.?
reflexa, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 109.—Afr. trop.
Regelii, *C. Koch, Wochenschr.* viii. (1865) 90.—Hab.?
Regina, Veitch, ex Regel, Gartenfl. (1871) 149 = terminalis.
rigida, *Miq. Fl. Ind. Bat. Suppl.* 610.—Sumatra.
rigidifolia, C. Koch, in C. Koch & Fint. Wochenschr. (1858) 389 = stricta.
rubra, *Hoog. ex Kunth, Enum. Pl.* v. 34.—N. Zel.
Rumphii, *Hook. Bot. Mag.* t. 4297 = Dracæna angustifolia, Hookeriana.
Rumphii, F. Muell. Fragm. v. 194 = Dracæna angustifolia.
salicifolia, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 59.—Java.
Sellowiana, Kunth, in Abh. Acad. Berl. (1842) 80 = dracænoides.
Sepiaria, Sesm. Fl. Vit. 311. t. 94 = terminalis.
Siebori, Kunth, in Abh. Acad. Berl. (1843) 80 = terminalis.
Sieboldii, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 109. sub t. 569 = Dracæna elliptica.
spectabilis, Kunth & Bouché, in Ind. Sem. Hort. Berol. (1848) = dracænoides.
spicata, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 152.—Hab.?
stricta, *Endl. in Ann. Wien. Mus.* i. (1836) 162.—Austral.
stricta, Hook. f. in Fl. N. Zel. i. 257. t. 80 = Pumilio.
Sturmii, *Colenso, in Transz. N. Z. Inst.* xv. (1883) 331.—N. Zel.
superbiens, C. Koch, in C. Koch & Fint. Wochenschr. (1858) 381 = australis.
terminalis, *Kunth, in Abh. Acad. Berl.* (1850) 80.—Ins. trop.; Austral.
Ti, Schott, in Flora, xi. (1828) 578 = terminalis.
timoriensis, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 153.—Ins. Timor.
umbraculifera, Planch. l. c. t. 111 = Dracæna umbraculifera.

CORDYLINE:—
Veitchii, Regel, Gartenfl. (1871) 149 = australis.
violacarna, Regel, Ind. Sem. Hort. Petrop. (1858) 25; et Gartenfl. (1859) 331 = rubra.
viridiflora, Hort. ex Steud. Nom. ed. II. i. 419 = Chlorophytum Sternbergianum.
Wallichii, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 152 (Dracænæ sp.).—Ind. or.

CORDYLOBLASTE, Hensch. ex Moritzi, in Bot. Zeit. v. (1847) 604 = Symplocos, Linn. (Styrac.).
Henschelii, Moritzi, l. c. = S. Henschelii.

CORDYLOCARPUS, Desf. Fl. Atlant. ii. 79. t. 182 (1800) = Rapistrum, Tourn. (Crucif.).
crassifolius, Steud. Nom. ed. I. 225 = Erucaria crassifolia.
glaber, Dietr. ex Steud. l. c. = R. glabrum.
laevigatus, Willd. Sp. Pl. iii. 563 = Erucaria aleppica.
muricatus, Desf. Fl. Atlant. ii. 79. t. 152.—Algeria.
pubescens, Sibth. & Sm. Fl. Græc. Prod. ii. 33 = Brassica adpressa.
ramosissimus, Pomel, Nouv. Mat. Fl. Atl. 357.—Algeria.
tenuisiliqua, Sibth. & Sm. Fl. Græc. Prod. ii. 33 = Erucaria tenuifolia.

CORDYLOGNE, Lindl. Veg. Kingd. 626 (1848), sphalm. = seq.

CORDYLOGYNE, E. Mey. Comm. Pl. Afr. Austr. 218 (1837). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 759.
Fellicaulum, Decne. in DC. Prod. viii. 590 (1844).
globosa, *E. Mey. Comm. Pl. Afr. Austr.* 218.—Afr. austr.

COREMA, D. Don, in Edinb. N. Phil. Journ. ii. (1826–27) 68. *EMPETRACEÆ*, Benth. & Hook. f. iii. 414.
Endamia, Rafin. New Fl. Am. iii. 49 (1836).
Ericulum, Rafin. Fl. Tellur. iii. 56 (1836).
Oaksia, Tuckerm. in Hook. Lond. Journ. Bot. i. (1842) 645.
Tuckermannia, Klotzsch, in Wiegm. Archiv, vii. (1841) 248.
album, *D. Don, in Edinb. N. Phil. Journ.* ii. (1827) 68.—Lusitan.
Conradi, *Torr. ex Loud. Encyc. Trees*, 1092.—Am. bor.
fobrifugum, Bolss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 312 = album.

COREOCARPUS, Benth. Bot. Voy. Sulph. 28. t. 16 (1844). *COMPOSITÆ*, Benth. & Hook. f. ii. 324.
Acoma, Benth. Bot. Voy. Sulph. 29. t. 17 (1844).
heterocarpus, *A. Gray, in Proc. Am. Acad.* v. (1861) 163.—Calif.
parthenioides, *Benth. Bot. Voy. Sulph.* 28. t. 16.—Calif.

COREOPIS, Gunn. Fl. Norveg. ii. 87 (1773), sphalm. = seq.

COREOPSIS, Linn. Gen. ed. I. 268 (1737). *COMPOSITÆ*, Benth. & Hook. f. ii. 325.
Achetum, Neck. Elem. i. 34 (1790).
Agarista, DC. Prod. v. 569 (1836).
Anacis, Schrank, in Denkschr. Acad. Muench. v. (1817) 8.
Calliopsis, Reichb. Ic. et Descr. Pl. t. 10 (1824).
Campylotheca, Cass. Dict. Sc. Nat. ii. 476 (1817).
Chrysanthemella, Tausch, Hort. Canal. fasc. i. (1832).
Chrysostemma, Less. Syn. Comp. 237 (1832).
Coreopsoides, Moench, Meth. 594 (1794).
Diatonta, Walp. Rep. ii. 614 (1843).
Diodonta, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 360.
Diplosastera, Tausch, Hort. Canal. fasc. i. t. 4 (1832).
Ecetea, DC. Prod. v. 569 (1836).
Epilepis, Benth. Fl. Hastw. 17 (1839).

COREOPSIS:—
Heterodonta, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 361.
Leachia, Cass. in Dict. Sc. Nat. xxv. 388 (1822).
Lerchia, Reichb. Consp. 109 (1828).
Peramiuus, Rafin. ex DC. Prod. v. 568 (1836).
Prestinaria, Sch. Bip. ex Steud. Nom. ed. II. ii. 298 (1841).
abyssinica, *Sch. Bip. ex Walp. Rep.* vi. 168.—Abyssin.
acuta, Pursh, Fl. Am. Sept. ii. 569 = Actinomeris squarrosa.
alata, Cav. Ic. iii. 30. t. 260 = Actinomeris tetraptera.
alata, Pursh, Fl. Am. Sept. ii. 567 = Verbesina occidentalis.
alba, Linn. Sp. Pl. 908 = Bidens pilosa.
alternifolia, Linn. f. c. 909 = Actinomeris squarrosa.
ambigua, Nutt. in Journ. Acad. Philad. vii. (1834) 75 = aurea.
amplexicaulis, Cav. Desc. 226 = Encelia amplexicaulis.
angustifolia, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 252.—Am. bor.
angustifolia, Linn. Sp. Pl. 908 = Helianthus angustifolius.
angustifolia, Pav. ex DC. Prod. v. 603 = Bidens prosera.
anthemoides, *DC. Prod.* v. 578.—Mexic.
arguta, Pursh, Fl. Am. Sept. ii. 517 = aurea.
aristata, Willd. Sp. Pl. iii. 2258 = aristosa.
aristosa, *Michx. Fl. Bor. Am.* ii. 140.—Am. bor.
artemisiæfolia, Jacq. Coll. Suppl. 135 = Cosmus sulphureus.
aspera, *Pursh, Fl. Am. Sept.* ii. 570.—Am. bor.
Atkinsoniana, *Dougl. in Lindl. Bot. Reg.* t. 1876.—Am. bor. occ.
aurea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 252.—Am. bor.
aurea, Lindl. Bot. Reg. t. 1228 = trichosperma.
auriculata, *Linn. Sp. Pl.* 908.—Am. bor.
auriculata, Schkuhr, Handb. t. 260 = pubescens.
Bacana, Heyne, ex DC. Prod. v. 582 = Glossogyne pinnatifida.
baccata, Linn. Pl. Surin. n. 111; Amoen. Acad. viii. 262 = Wulfia stenoglossa.
baccata, *Linn. f. Suppl.* 350.—Guiana.
Bartavi, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
bicolor, *Bosse, ex Buchenau, in Linnaea*, xxv. (1852) 680.—Hab.?
Bidens, Linn. Sp. Pl. 908 = Bidens cernua.
Bidens, Walt. Fl. Carol. 215 = Bidens chrysanthemoides.
bidentoides, *Nutt. ex Torr. & Gray, Fl. N. Am.* ii. 389.—Am. bor.
biternata, *Lour. Fl. Cochinch.* 508.—China.
bituminosa, *Naronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 11.—Malaya.
Borianiana, *Sch. Bip. ex Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 984.—Afr. trop.
Boyhiniana, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 356 = grandiflora.
calliopsidea, A. Gray, in Torr. Bot. Mex. Bound. 90 = Lagascera calliopsidea.
callosa, *Bertol. Misc. Bot.* vii. 42.—Am. bor.
capillacea, *H. B. & K. Nov. Gen. et Sp.* iv. 230.—Peruv.
cardaminefolia, *Torr. & Gray, Fl. N. Am.* ii. 346.—Am. bor. occ.
chrysanthe, Linn. Sp. Pl. ed. II. 1262 = Bidens chrysantha.
chrysanthe, Spreng. Syst. iii. 674 = Bidens rubifolia.
chrysantha, *Vatke, in Linnaea*, xxxix. (1875) 499.—Afr. trop.
coronata, Bert. ex Steud. Nom. ed. II. i. 420 = Cosmus candatus.
coronata, *Hook. Bot. Mag.* t. 3460.—Texas.
coronata, Linn. Sp. Pl. ed. II. 1263 (ex syn. Plum.) = Bidens pilosa.
coronata, Walt. Fl. Carol. ii. 16 = aurea.
corymbifolia, Buch.-Ham. ex DC. Prod. v. 602 = Bidens pilosa.
cosmoides, *A. Gray, in Proc. Am. Acad.* v. (1861) 158.—Ins. Bonvic.
crassifolia, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 252 = lanceolata.
cuspidata, *Bertol. Misc. Bot.* vii. 44.—Am. bor.
delphinifolia, *Lam. Encyc.* ii. 126.—Am. bor.
dichotoma, Michx. Fl. Bor. Am. ii. 137 = gladiata.

COREOPSIS:—
discoidea, *Torr. & Gray, Fl. N. Am.* ii. 339.—Am. bor.
discolor, *Linn. Enum. Hort. Berol.* ii. 858.—Am. bor.
diversifolia, DC. Prod. v. 571. (excl. syn.) = auriculata.
diversifolia, Hook. Bot. Mag. t. 3474 = Drummondii.
diversifolia, Jacq. Eclog. 80. t. 54 = Bidens grandiflora. Drummondii, *Torr. & Gray, Fl. N. Am.* ii. 345.—Texas.
emarcida, *Noronha, in Verh. Batavo. Gen.* v. (1790) ed. I. Art. IV. 11.—Malaya.
fasciculata, *Wedd. Chlor. And.* i. 71.—Peruv.
ferulaefolia, Jacq. Hort. Schoenb. iii. 65 = Bidens ferulaefolia.
filifolia, Hook. Bot. Mag. t. 3505 = Thelesperma filifolium.
Flammula, Banks, ex Steud. Nom. ed. I. 926, 108 = Bidens chrysanthemoides.
flexicaulis, *Rafin. in Med. Repos. N. York*, v. (1808) 361.—Am. bor.
foeniculacea, Moc. & Sesse, ex DC. Prod. v. 608 = Bidens foeniculifolia.
fœtida, Cav. Ic. i. 55 = Encelia fœtida.
foliosa, *A. Gray, in Proc. Am. Acad.* v. (1861) 125.—Peruv.
formosa, *Bonato, Memoria 9*; *ex Prits. Ic. Ind.* i. 301. —Hab. ?
formosa, Hort. ex Steud. Nom. ed. II. i. 620 = Cosmos bipinnatus.
fruticosa, *Forst. f. Prod.* 91.—Ins. Sandvic.
fruticosa, *Soland. ex Seem. Fl. Vit.* 148.—Ins. Pacif.
frutescens, Vest, in Flora, iii. (1820) 7 = Bidens frutescens.
Galpinii, *Hemsl. Biol. Centr. Am. Bot.* ii. 195.—Mexic.
Georgina, Cass. in Dict. Sc. Nat. xviii. 441 = Dahlia variabilis.
gladiata, *Walt. Fl. Carol.* 215.—Am. bor.
glaucescens, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 389.—Afr. trop.
grandiflora, Blanco, Fl. Filip. ed. II. Suppl. 591 = Cosmos caudatus.
grandiflora, *Nutt. ex Chapm. Fl. S. U. St.* 385.—Am. bor.
Grantii, *Oliver, in Trans. Linn. Soc.* xxix. (1874) 98. t. 65.—Afr. trop.
guineensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
Harveyana, *A. Gray, Syn. Fl. N. Am.* i. 11. 292.—Am. bor. occ.
heterophylla, *Bertol. Misc. Bot.* vii. 43.—Am. bor.
heterophylla, Cav. Ic. iii. 34. t. 268 = Iostephane heterophylla.
heterophylla, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 388 = grandiflora.
hirta, *Rafin. Ann. Nat.* 15.—Am. bor.
incisa, Ker-Gawl. in Bot. Reg. t. 7 = Bidens rubifolia.
incurvoa, Moench, Meth. Suppl. 245 = Bidens ferulaefolia.
integrifolia, *Poir. Encyc. Suppl.* ii. 353 (1811).—Am. bor.
involucrata, *Nutt. in Journ. Acad. Philad.* vii. (1834) 74.—Am. bor.
involucrata, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyss.
jasminifolia, *Bertol. Misc. Bot.* vii. 44.—Am. bor.
Kirkii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
lanceolata, *Linn. Sp. Pl.* 908.—Am. bor.
latifolia, *Michx. Fl. Bor. Am.* ii. 127.—Am. bor.
Leavenworthii, *Torr. & Gray, Fl. N. Am.* ii. 346.—Am. bor.
leucantha, Linn. Sp. Pl. ed. II. 1282 = Bidens pilosa.
leucanthema, Linn. Cent. Pl. i. 29 = prænc.
leucorhiza, Lour. Fl. Cochinch. ii. 508 = Bidens leucorrhiza.
Liebmanni, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 196.—Mexic.
limonoia, Jacq. Coll. ii. 299 = Encelia canescens.
linearifolia, *DC. Prod.* v. 570.—Mexic.
linearifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
linifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 75 = angustifolia.
longipes, Hook. Bot. Mag. t. 3568 = grandiflora.
lucida, Cav. Desc. 226 = Bidens heterophylla.
Macmei, *A. Gray, in Proc. Am. Acad.* v. (1861) 126.—Ins. Sandvic.
macrantha, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyss.

COREOPSIS:—
macrocarpa, *A. Gray, in Proc. Am. Acad.* v. (1861) 126.—Ins. Sandvic.
major, Walt. Fl. Carol. 214 = senifolia.
maritima, *Hook. f. Bot. Mag.* t. 6241.—Calif.
Mentzelii, *A. Gray, in Proc. Am. Acad.* v. (1861) 127. —Ins. Sandvic.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* ii. 196.—Mexic.
micrantha, *A. Gray, in Proc. Am. Acad.* v. (1861) 127. —Ins. Sandvic.
minima, *Hort. ex Steud. Nom.* ed. II. i. 620.—Am. bor.
mitis, Michx. Fl. Am. Bor. ii. 140 = aurea.
monticola, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
mutisiifolia, *DC. Prod.* v. 578.—Peruv.
mutica, *DC. l. c.* 571.—Mexic.
Nesei, *DC. l. c.* 573.—Am. austr.
nudata, *Nutt. Gen. Am.* ii. 180.—Florida.
oblongifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 76 = lanceolata.
odorata, Lam. Illustr. t. 704. f. 1 = Bidens odorata.
odoratissima, Cav. ex Pers. Syn. ii. 477 = Cosmos tenellus.
Oemleri, Ell. Sketch, ii. 435 = senifolia.
Oerstedtiana, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 93.—Bras.
ovata, Cav. Ic. iii. 41. t. 280 = Actinomeris ovata.
pachyloma, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 391.—Afr. trop.
palmata, *Nutt. Gen. Am.* ii. 180.—Am. bor.
parviflora, Jacq. Hort. Schoenb. iii. 65 = Cosmos parviflorus.
pauciflora, Lehm. Ind. Sem. Hort. Hamb. (1838); Linnæa, x. (1836) Litt. 76 = palmata.
perfoliata, Walt. Fl. Carol. 214 = Bidens chrysanthemoides?
Pickeringii, *A. Gray, in Proc. Am. Acad.* v. (1861) 124.—Peruv.
prærcox, Presen. in Linnæa, xiii. (1839) Litt. 93 = palmata.
Prestinaria, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyss.
prestinariæformis, *Vatke, in Linnæa*, xxxix. (1875) 499.—Afr. trop.
procera, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 353 = Actinomeris squarrosa.
pubescens, *Ell. Sketch*, ii. 441.—Am. bor.
purpurea, Moç. & Sesse, ex DC. Prod. v. 604 = Cosmos purpureus.
quadricornis, Krock. Fl. Siles. ii. 11. 481 = Bidens cernua.
radiata, *Mill. Gard. Dict. ed.* VIII. n. 5.—Am. bor.
reptans, Linn. Syst. ed. X. 1228; Amoen. Acad. v. 507 = Bidens rubifolia.
ridens, *Gunn. Fl. Norveg.* ii. 87.—Norvegia.
rigida, Nutt. Gen. Am. ii. 180 = delphinifolia.
rosea, *Nutt. l. c.* 179.—Am. bor.
rudis, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 196.—Mexic.
Ruppellii, Sch. Bip. ex Walp. Rep. vi. 163 = chrysantha.
sambucifolia, Cav. Desc. 225 = Bidens sambucifolia.
scabra, *Rafin. Fl. Ludov.* 72.—Louisiana.
Schaffneri, *A. Gray, in Proc. Am. Acad.* xix. (1883) 15.—Mexic.
senifolia, *Michx. Fl. Bor. Am.* ii. 138.—Am. bor.
serrulata, Poir. Encyc. Suppl. iii. 352 = Bidens grandiflora.
setigera, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyss.
spectabilis, *A. Gray, in Proc. Am. Acad.* v. (1861) 125.—Peruv.
stellata, Banks, ex DC. Prod. v. 572 = senifolia.
stellata, *Nutt. in Journ. Acad. Philad.* vii. (1834) 76. —Am. bor.
Steppia, *Steetz, in Peters, Reise Mossamb. Bot.* 496.—Afr. trop.
Tausa, Steud. Nom. ed. II. i. 620 = Glossogyne tenuifolia.
tenuicula, Spreng. Fl. Hal. Mant. 53 = Glossogyne tenuifolia.
tenuifolia, Ehrh. Beitr. vii. 168 = verticillata.
tetragona, Cerv. ex La Llave & Lex. Nov. Veg. Desc. fasc. i. 21 = Bidens tetragona.
tinctoria, *Nutt. in Journ. Acad. Philad.* ii. (1821) 116.—Am. bor.
trichosperma, *Michx. Fl. Bor. Am.* ii. 139.—Am. bor.
trifida, Poir. Encyc. Suppl. iii. 353 = Thelesperma filifolium.

COREOPSIS:—
trifoliata, *Bertol. Fl. Guatimal.* 26.—Guatemala.
Tripteris, *Linn. Sp. Pl.* 908.—Am. bor.
verticillata, Salisb. Prod. 206 = Bidens rubifolia.
venusta, *H. B. & K. Nov. Gen. et Sp.* iv. 239.—Peruv.
verticillata, Ehrh. Beitr. vii. 168 = delphinifolia.
verticillata, Lam. Encyc. ii. 108 = tenuifolia.
verticillata, *Linn. Sp. Pl.* 907.—Am. bor.
Wrayi, Nutt. in Journ. Acad. Philad. vii. (1834) 76 = senifolia.

COREOPSOIDES, Moench, Meth. 594 (1794) = Coreopsis, Linn. (Compos.).
lanceolata, Moench, l. c. = Coreopsis lanceolata.

COREOSMA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 2 = Ribes, Linn. (Saxifrag.).
aphylla, Spach, l. c. 24 = R. campanulatum.
Bombeyana, Spach, l. c. 25 = R. viscosum.
florida, Spach, l. c. 22 = R. floridum.
multiflora, Spach, l. c. 24 = R. Kunthii.
sanguinea, Spach, Hist. Vég. Phan. vi. 155 = R. sanguineum.
viscosissima, Spach, in Ann. Sc. Nat. Ser. II. iv. (1835) 23 = R. viscosissimum.

CORESANTHA, Alef. in Bot. Zeit. xxi. (1863) 298 = Iris, Tourn.
alata, Klatt, in Linnæa, xxxiv. (1865–66) 575 = I. stylosa.
caucasica, Klatt, l. c. = I. caucasica.
persica, Alef. in Bot. Zeit. xxi. (1863) 298 = I. persica.

CORESANTHE, Baker, in Journ. Linn. Soc. xvi. (1877) 123 = præc.

CORETHRODENDRON, Fisch. & Basin. in Mem. Sav. Etr. Petersb. vi. (1851) 90 (Monogr. Hedys. 46. t. 2). *LEGUMINOSÆ*, Benth. & Hook. f. i. 512.
scoparium, *Basin. l. c.* 91.—Scoogaria.

CORETHROGYNE, DC. Prod. v. 215 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 370.
californica, *DC. l. c.*—Calif.
detonsa, Greene, in Bull. Torrey Club, x. (1883) 41 = Diplostephium canum.
filaginifolia, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 390.—Calif.
incana, Nutt. l. c. = californica.
obovata, *Benth. Bot. Voy. Sulph.* 22.—Calif.
spathulata, A. Gray, in Proc. Am. Acad. vii. (1868) 351 = obovata.
tomentella, Torr. & Gray, Fl. N. Am. ii. 99 = filaginifolia.
virgata, Benth. Bot. Voy. Sulph. 23 = filaginifolia.

CORETHROSTYLES, Benth. & Hook. f. Gen. i. 228 (1862), sphalm. = seq.

CORETHROSTYLIS, Endl. Nov. Stirp. Dec. i. (1839) 1 = Lasiopetalum, Sm. (Stercul.).
bracteata, Endl. l. c. i. 2 = L. bracteatum.
cordifolia, Steetz, in Lehm. Pl. Preiss. ii. 344 = L. cordifolium.
coriacea, Steud. l. c. i. 226 = L. bracteatum.
membranacea, Steud. l. c. = L. membranaceum.
microphylla, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 148 = L. cordifolium.
oppositifolia, F. Muell. Fragm. ii. 6 = L. floribundum.
parvifora, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 174 = L. floribundum.
Schulzenii, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 86 = L. Schulzenii.

CORETHRUM, Vahl, in Skr. Naturk. Selsk. Kiobenh. vi. (1810) 85 = Boutelouа, Lag. (Gramin.).
bromoides, Vahl, l. c. = B. bromoides.

CORIANDRUM, [Tourn.] Linn. Syst ed. I (1735). *UMBELLIFERÆ*, Benth. & Hook. f. i. 926.
KERAMOCARPUS, Fenzl, in Russegg. Reise ii. 966 (1848).
americanum, Nutt. ex DC. Prod. iv. 250 = Bifora americana.
Cicuta, Crantz, Stirp. Austr. ed. I. fasc. iii. 100; ed. II. 211 = Conium maculatum.
Cicuta, Roth, Tent. Fl. Germ. i. 180 = Cicuta virosa.

CORIANDRUM:—
Cynapium, Crantz, Stirp. Austr. ed. I. fasc. iii. 99; ed. II. 211 = Aethusa Cynapium.
didymum, Stoken, Bot. Mat. Med. ii. 121 = Bifora testiculata.
diversifolium, Gilib. Fl. Lituan. ii. 26 = sativum.
globosum, Salisb. Prod. 166 = sativum.
latifolium, Crantz, Stirp. Austr. ed. I. fasc. iii. 101; ed. II. 211 = Sium latifolium.
maculatum, Roth, Tent. Fl. Germ. i. 130 = Conium maculatum.
majus, Goüan, Hort. Monsp. 145 = sativum.
melphitense, *Tenore & Guss. Ind. Sem. Hort. Neap.* (1837) 8.—Italia.
sativum, *Linn. Sp. Pl.* 256.—Europ. austr.; Oriens.
sessilifolium, DC. ex DC. Prod. iv. 250 = Astoma sessilifolium.
tordylioides, *Boiss. Fl. Orient.* ii. 921.—Orient.
testiculare, Salisb. Prod. 166 = Bifora testiculata.
testiculatum, Bieb. Fl. Taur. Cauc. i. 238 = Bifora radians.
testiculatum, Linn. Sp. Pl. 256 = Bifora testiculata.
testiculatum, Lour. Fl. Cochinch. 225 = Bifora Loureiri.

CORIARIA, Nins. ex Linn. Gen. ed. I. 310 (1737). *CORIARIEÆ*, Benth. & Hook. f. i. 429.
HETEROCLADUS, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 152.
HETEROPHYLLARIA, Turcz. l. c. (1848) i. 591.
angustisima, *Hook. f. Handb. New Zeal. Fl.* 47.—N. Zel.
arborea, *Lindsay, Contrib. N. Z. Bot.* 84.—N. Zel.
atropurpurea, Moç. & Sesse, ex DC. Prod. i. 740 = thymifolia.
hermaphrodita, Turcz, Faraet. Nov. Gen. 18 = myrtifolia.
japonica, *A. Gray, in Mem. Acad. Am.* N. S. vi. (1858–59) 383.—Japon.
microphylla, *Poir. Encyc.* vi. 87.—? Peruv.
myrtifolia, *Linn. Sp. Pl.* 1057.—Reg. Mediterr.
nepalensis, *Wall. Pl. As. Rar.* iii. 67.—Reg. Himal.; China.
phylicifolia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 819. —Peruv.
procumbens, *Whitl. ex Hoffmgg. Vers. Pfl. Nachtr.* iii. 9, nomen.—Hab.?
ruscifolia, *Linn. Sp. Pl.* 1057.—Peruv.
sarmentosa, *Forst. f. Prod.* 71. n. 377.—N. Zel.
sinica, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. (1881) n. 111. 9.—China.
thymifolia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 819. —Peruv.
tinctoria, Dulac, Fl. Hautes-Pyr. 248 = myrtifolia.
Tutu, *Lindsay, Contrib. N. Z. Bot.* 84.—N. Zel.

CORIDOCHLOA, Nees, in Edinb. N. Phil. Journ. xv. (1833) 361, in nota = Panicum, Linn. (Gramin.).
cimicina, Nees, l. c. = P. cimicinum.
semialata, Nees, ex Benth. Fl. Austral. vii. 478 = P. semialatum.

CORIDOTHYMUS, Reichb. f. in Oestr. Bot. Wochenbl. vi. (1857) 160 = Thymus, Tourn. (Labiat.).
capitatus, Reichb. f. l. c. 161 = T. capitatus.

CORILUS, Nocca, in Ustari, Ann. Bot. ii. (1793) 50 = Corylus, Tourn. (Cupul.).

CORINDUM, Adans. Fam. ii. 388 (1763) = Paullinia, Linn. (Sapind.).

CORINDUM, Tourn. ex Medic. Malv. 110 (1787) = Cardiospermum, Linn. (Sapind.).
Halicacabum, Medic. l. c. = Card. Halicacabum.

CORINGIA, J. & C. Presl, 'Fl. Čech. 141'; Fl. Sicula, 79 (1826) = Conringia, Heist. (Crucif.).

CORINOCARPUS, Forr. in Dict. Sc. Nat. x. 441 (1818) = Corynocarpus, Forst. (Anacard.).

CORION, Hoffmgg. & Link, Fl. Port. ii. 457 (1820) = Bifora, Hoffm. (Umbellif.).
testiculatum, Hoffmgg. & Link, l. c. = B. testiculata.

CORION, Mitch. in Act. Nat. Cur. viii. (1748) 208, App. = Spergularia, J. & C. Presl (Caryophyll.).

CORIS, Tourn. ex Linn. Gen. ed. I. 336 (1737). *PRIMULACEÆ*, Benth. & Hook. f. ii. 636.
hispanica, *Lange, in Kjoeb. Vidensk. Meddel.* (1866) 69 (*Pugill.* 219).—Hispan.
monspeliensis, *Linn. Sp. Pl.* 177.—Europ. austr.

CORISANTHERA, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 373 (1884) = CORYSANTHERA, Wall. = Rhynchotechum, Blume (Gesnetrac.).

CORISANTHES, Stend. Nom. ed. II. i. 474, in syn. (1840) = CRIOSANTHES, Rafin. = Cypripedium, Linn. (Orchid.).

CORISPERMUM, B. Juss. ex Linn. Syst. ed. I (1735). *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 57.
apetalum, *Hort. ex Steud. Nom.* ed. II. i. 420.—Hab.?
borysthenicum, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 129, nomen.—Rossia.
bracteatum, Vitv. Lett. Corisp. 2 = Marschallii.
caucasicum, *Kit. ex Schult. Oestr. Fl.* ed. II. i. 7.—Europ. occ.
coloratum, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 130, nomen.—Rossia.
confertum, *Bunge, in Maxim. Prim. Fl. Amur.* 225.—Reg. Amur.
crassifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) 428.—Sibiria.
declinatum, Steph. ex Stev. in Mém. Soc. Nat. Mosc. v. (1814) 334 = hyssopifolium.
elatum, Host, Fl. Austr. i. 319 = Marschallii.
elongatum, *Bunge, ex Maxim. Prim. Fl. Amur.* 224. —Manchur.
filifolium, *Becker, in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 13.—Rossia.
glomeratum, *Lag. Gen. et Sp. Nov.* 1.—Hab.?
Gmelini, *Bunge, in Bull. Acad. Pétersb.* xxv. (1879) 187.—Persia.
heteracanthum, *Ledeb. ex Heynh. Nom.* iii. 161, nomen. —Rossia.
hybridum, Bess. ex Moq. in DC. Prod. xiii. ii. 141 = Marschallii.
hyssopifolium, *Linn. Sp. Pl.* 4.—Europ.; As. temp.; Am. bor.
hyssopifolium, Pall. Reise, i. 484; Fl. Ross. t. 98 = orientale.
intermedium, Reichb. Fl. Germ. Excurs. 584 = Marschallii.
intermedium, *Schweig. in Koenigsb. Arch. Naturw.* i. (1812) 211.—Europ.
latifolium, Link, Handb. ii. 408 = orientale.
laxiflorum, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 361 = Marschallii.
Lehmannianum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 468.—Turkestan.
macrocarpum, *Bunge, in Maxim. Prim. Fl. Amur.* 226.—Manchur.
macrocarpum, Trautv. in Act. Hort. Petrop. ix. (1884) 130 = macrocarpum.
marginatum, Steph. ex Moq. in DC. Prod. xiii. ii. 140 = Marschallii.
Marschallii, *Stev. in Mém. Soc. Nat. Mosc.* v. (1814) 336.—Europ.
microspermum, Host, Fl. Austr. i. 319 = nitidum.
nitidum, *Kit. ex Schult, Oestr. Fl.* ed. II. i. 7.—Europ.; As. bor.
orientale, *Lam. Encyc.* ii. 111; *Illustr.* i. 15.—Rossia; As. bor.; Persia.
Pallasii, Stev. in Mém. Soc. Nat. Mosc. v. (1814) 336 = orientale.
patens, Fisch. in Roem. & Schult. Syst. i. 579 = hyssopifolium.
patens, Link, in Spreng. Jahrb. Gewaechsk. i. 111. 28 = Redowskii.
patens, *Roem. & Schult. Syst.* i. 579.—Sibir.
pendulum, Willd. ex Link, in Spreng. Jahrb. Gewaechsk. i. 111. 28 = orientale.
pilosum, Rafin. New Fl. Am. iv. 46 = hyssopifolium?
polygaloides, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 34 = Anthochlamys polygaloides.
pungens, Savi, sec. Uechtr. ex Steud. Nom. ed. II. i. 421 = squarrosum.
pungens, Vahl, Enum. i. 117 = Agriophyllum arenarium, Bieb.
purpurascens, Host, Fl. Austr. i. 319 = nitidum.
Redowskii, Fisch. & Stev. in Mém. Soc. Nat. Mosc. v. (1814) 336 = sq.
rigidum, Fisch. & Stev. in Mém. Soc. Nat. Mosc. v. (1814) 336 = sq.
salsoloides, Host. ex Steud. Nom. ed. II. i. 420 = glomeratum.
spicatum, Willd. Enum. Hort. Berol. Suppl. 1 = orientale.
squarrosum, Bieb. Fl. Taur. Cauc. i. 8 = Marschallii.
squarrosum, Link, Enum. Hort. Berol. i. 7 = orientale.
squarrosum, Linn. Sp. Pl. 4 = hyssopifolium.
squarrosum, Pall. Fl. Ross. i. ii. 113. t. 99 = Agriophyllum arenarium, Bieb.
squarrosum, *Vahl, Enum.* i. 16.—Europ.; As. temp.
squarrosum, Willd. Enum. Hort. Berol. 8 = Marschallii.
Stauntoni, *Moq. Chenop. Enum.* 104.—China.
subpetandrum, Pall. Illustr. i. 59 = orientale.
tenue, Link, in Spreng. Jahrb. Gewaechsk. i. 111. 29 = nitidum.
tylocarpum, *Hance, in Journ. Bot.* vi. (1868) 47.—China.
ulopterum, *Fenzl, ex Ledeb. Fl. Ross.* iii. 765.—Sibir.

CORISTOSPERMUM, Bertol. Fl. Ital. iii. 466 (1837) = Ligusticum, Linn. (Umbellif.).
cuneifolium, Bertol. l. c. 467 = L. cuneifolium.

CORIZOSPERMUM, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 255 (1850) = Cassaria, Jacq. (Samyd.).
clutiaefolium, Zipp. ex Blume, l. c. = Cas. clutiaefolia.

CORMIGONUS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 83 = Bikkia, Reinw. (Rubiac.).

CORMONEMA, Reissek, ex Endl. Gen. 1098 (1840). *RHAMNEÆ*, Benth. & Hook. f. i. 379.
CAESIA, Vell. Fl. Flum. iii. t. 23 (1827).
spinosum, *Reissek, in Mart. Fl. Bras.* xi. i. 96.—Bras.

CORMUS, Spach, Hist. Vég. Phan. ii. 96 (1834) = Pyrus, Linn. (Rosac.).
domestica, Spach, l. c. 97 = P. Sorbus.

CORNA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (quid?).

CORNACCHINIA, Savi, in Mem. Soc. Ital. Moden. xxi. (1827) 184. t. 7 = Clerodendron, Linn. (Verbenac.).
fragiformis, Savi, l. c. 185 = Clerod. Acerbianum.

CORNELIA, Ard. Spec. ii. 9 (1764) = Ammannia, Houst. (Lythr.).
verticillata, Ard. l. c. = A. salicifolia.

CORNICINA, Boiss. Voy. Bot. Espagne, ii. 162 (1840?) = Anthyllis, Linn. (Legumin.).
cicrinata, Boiss. l. c. 163 = Hymenocarpus circinnatus.
hamosa, Boiss. l. c. 163 = A. hamosa.
Loeflingii, Boiss. l. c. = A. Cornicina.
lotoides, Boiss. l. c. = A. lotoides.
nummularia, Boiss. l. c. 163 = Hymenocarpos nummularius.
pseudocytisus, Boiss. Diagn. Ser. I. ii. 15 = Cytisopsis dorycnifolia.

CORNIDIA, Ruiz & Pav. Prod. 68. t. 25 (1794) = Hydrangea, Linn. (Saxifrag.).
integerrima, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 344 = H. scandens, *Poepp.*
radiata, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 42 = H. peruviana.
serratifolia, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 344 = H. serratifolia.
umbellata, Ruiz & Pav. Syst. Veg. 91.—Peruv.

CORNIOLA, Adans. Fam. ii. 321 (1763) = Genista, Linn. (Legumin.).
emarginata, Presl, Bot. Bemerk. 127 = G. sessilifolia.
cavalica, Presl, l. c. 126 = G. cavalica.
mantica, Presl, l. c. = G. mantica.
microphylla, Presl, l. c. 127 = G. microphylla.
ovata, Presl, l. c. 126 = G. ovata.
scariosa, Presl, l. c. = G. triangularis.
Scorpius, Presl, l. c. 127 = G. Scorpius.
sibirica, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 343 = G. tinctoria.
tinctoria, Medic. l. c. = G. tinctoria.
triangularis, Presl, Bot. Bemerk. 126 = G. triangularis.
virgata, Presl, l. c. = G. virgata.

CORNUCOPIÆ, Linn. Gen. ed. i. 13 (1737). *GRAMINEÆ*, Benth. & Hook. f. iii. 1189.
alopecuroides, Linn. Mant. i. 28 = Alopecurus striculatus.
altissimum, Walt. Fl. Carol. 74 = Agrostis elata ?
cacullatum, *Linn. Sp. Pl.* 84.—As. Min.; Mesopot.
hyemale, Walt. Fl. Carol. 73 = Agrostis scabra.
perennans, Walt. l. c. 74 = Agrostis scabra.

CORNULACA, Delile, Fl. Egypte, 206. t. 22. f. 3 (1813). *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 69.
amblyacantha, *Bunge, Anahas. Rev.* 68.—Persia.
Ancheri, *Moq. Chenop. Enum.* 163.—Persia.
Ehrenbergii, *Aschers. in Schweinf. Beitr. Fl. Aethiop.* 184.—Afr. trop.
monacantha, *Delile, Fl. Egypte*, 206.—Aegypt.; Persia.
muricata, Delile, Fl. Egypte Illustr. 57 = Kochia muricata.
setifera, Moq. in DC. Prod. xiii. ii. 218 = tragacanthoides.
tragacanthoides, *Moq. Chenop. Enum.* 163.—Syria; Mesopot.

CORNUS, [Tourn.] Linn. Syst. ed. i (1735). *CORNACEÆ*, Benth. & Hook. f. i. 950.
BENTHAMIA, Lindl. Bot. Reg. t. 1579 (1833).
CYNOXYLON, Rafin. Alsog. Am. 59 (1838).
EUKRANIA, Rafin. l. c. (1838).
SVIDA, Opiz, Seznam, 94 (1852).
TELUKRAMA, Rafin. Alsog. Am. 64 (1838).
alba, Hook. & Arn. Bot. Beech. Voy. 142 = californica.
alba, Lam. Encyc. ii. 115, partim (et auct. Amer.) = stolonifera.
alba, *Linn. Mant.* i. 40.—Sibir.
alba, Thunb. Fl. Jap. 69 = macrophylla.
alba, Walt. Fl. Carol. 88 = C. Amomum.
albida, Ehrh. Beitr. iv. 16 = candidissima.
alterna, Marsh. Arbust. 85 = alternifolia.
alternifolia, *Linn. f. Suppl.* 136.—Am. bor.
Amomum, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. bor.
asperifolia, *Michx. Fl. Bor. Am.* i. 98.—Am. bor.
strata, *Rafin. Alsog. Am.* 61.—Am. bor.
australis, C. A. Mey. in Bull. Phys.-Math. Acad. Petersb. iii. (1845) 372 = sanguinea.
biramis, Stokes, Bot. Mat. Med. i. 291 = suecica.
bovalis, Krasch. Fl. Ingr. 94 = suecica.
brachypoda, C. A. Mey. in Mém. Acad. Pétersb. vii. (1846) ii. 228 = macrophylla.
caerulea, Auct. vide *coerulea*.
californica, *C. A. Mey. in Bull. Phys.-Math. Acad. Petersb.* iii. (1845) 278.—Am. bor. occ.
canadensis, Cham. & Schlecht. in Linnæa, iii. (1828) 139 = unalaschensis.
canadensis, *Linn. Sp. Pl.* 118.—Am. bor.
canadensis, Willd. ex Ledeb. Fl. Ross. ii. 877 = suecica.
candidissima, Bisch. Ind. Sem. Hort. Heidelb. (1851) = stoloníferi.
candidissima, *Mill. Gard. Dict.* ed. VIII. n. 6.—Am. bor.
capitata, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 484.—Reg. Himal.
candata, Zoll. in Flora, xxx. (1847) 603 = Polyosma integrifolia.
chinensis, Molina, Sagg. Chil. 178 = Aristotelia Macqui.
cinerea, *Rafin. Alsog. Am.* 59.—Am. bor.
circinata, *L'Hérit. Cornus*, 7.—Am. bor.
circinnata, Cham. & Schlecht. in Linnæa, iii. (1828) 139 = californica.
circinnata, Hort. ex Lam. Encyc. ii. 116 = candidissima.
circinnata, Wahlenb. in Isis, xxi. (1828) 983 = sanguinea.
coerulea, Lam. Encyc. ii. 116 = C. Amomum.
coerulea, Mœnch. Fl. Sel. i. 8 = sericea.
comosa, *Rafin. Alsog. Am.* 62.—Am. bor.
cricquia, Hance, in Journ. Bot. xix. (1881) 216 = macrophylla.
cynanthus, *Rafin. Atl. Journ.* 151.—Am. bor.
cynoxarpus, J. F. Gmel. Syst. 367 = sericea.
cynoxarpus, Mœnch, Verz. 27 = C. Amomum.
dauurica, Lamm. ex Ledeb. Fl. Ross. ii. 866 = Wilsoniana dauurica.
dichotoma, *Rafin. Alsog. Am.* 60.—Am. bor.
disciflora, *Mey. & Sesse, ex DC. Prod.* iv. 278.—Mexic.
Drummondi, *C. A. Mey. in Bull. Phys.-Math. Acad. Petersb.* iii. (1846) 278.—Am. bor.

CORNUS:—
asythracantha, St. Lager, ex Bull. Soc. Bot. Fr. xxxi. (1883) Bibl. 201 = C. Mas.
excelsa, *H. B. & K. Nov. Gen. et Sp.* iii. 430.—Mexic.
fastigiata, Michx. Fl. Bor. Am. i. 92 = stricta.
femina, Anct. vide *foemina*.
ferruginea, Hort. ex Lam. Encyc. ii. 116 = C. Amomum.
ferruginea, Jacq. ex Roem. & Schult. Syst. iii. 325 = ferulaefolia.
ferulaefolia, *Necca, Syn. Pl.* 38.—Hab. ?
flava, Stend. Nom. ed. I. 357 = C. Mas.
florentina, *Decne. in Nouv. Arch. Mus. Par.* ser. I. x. (1874) 157.—Hab. ?
florida, Hook. Fl. Bor. Am. i. 277, partim = Nuttallii.
florida, *Linn. Sp. Pl.* 117.—Am. bor.
foemina, Mill. Gard. Dict. ed. VIII. n. 4 = sanguinea.
glabrata, *Benth. Bot. Voy. Sulph.* 18.—Calif.
grandis, *Cham. & Schlecht. in Linnæa*, v. (1830) 171.—Mexic.
herbacea, Steller, in Pall. N. Nord. Beitr. ii. 300 = suecica.
ignorata, *C. Koch, Ind. Sem. Hort. Berol.* (1867) *App.* i. 8.—Japon.
illicifolia, Hassk. & Zoll. ex Hassk. Cat. Hort. Bog. Alt. 168, 211 = Polyosma illicifolia.
japonica, *Thunb. Fl. Jap.* 68.—Japon.
Kousa, *Buerger, ex Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 169, in syn.—Japon.
lanceolata, *Rafin. Alsog. Am.* 60.—Am. bor.
lanuginosa, Michx. Fl. Bor. Am. i. 92 = C. Amomum.
latifolia, Bray, in Denkschr. Bot. Ges. Regensb. i. 11. (1818) 85 = sanguinea.
longifolia, *Medic. Beobachtt.* (1782) 308.—Hab. ?
macrophylla, *Wall. in Roxb. Fl. Ind. ed Carey*, i. 483. —Reg. Himal.; China; Japon.
Mas, *Linn. Sp. Pl.* 117.—Europ.; As. Min.
mascula, Linn. Syst. ed. X. 897 = C. Mas.
mexicana, Carr. in Rev. Hortic. (1875) 460 = C. Amomum.
minor, Hort. ex Stend. Nom. ed. II. i. 421 = sericea.
nudiflora, Dum. Fl. Belg. 88 = C. Mas.
Nuttallii, *Audubon, Birds Am.* t. 367; *ex Torr. & Gray, Fl. N. Am.* i. 651.—Am. bor.
obliqua, Rafin. Ann. Nat. (1820) 13 = C. Amomum.
oblonga, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 482.—Reg. Himal.
oblongifolia, Rafin. ex DC. Prod. iv. 272 = sericea.
officinalis, *Sieb. & Zucc. Fl. Jap.* i. 100. t. 50.—Japon.
paniculata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 140 = oblonga.
paniculata, L'Hérit. Cornus, 9. t. 5 = candidissima.
parvifolia, *Rafin. Alsog. Am.* 61.—Am. bor.
paucinervis, *Hance, in Journ. Bot.* xix. (1881) 216.—China.
plicata, Tausch, in Flora, xxi. (1838) 733 = alternifolia.
polygama, Rafin. Fl. Ludov. 78 = C. Amomum ?
praecox, Stokes, Bot. Mat. Med. i. 292 = C. Mas.
pubescens, Nutt. Sylva, iii. 84 = C. Amomum.
pubescens, Torr. Bot. Mex. Bound. 71 = californica.
pubescens, Willd. ex Roem. & Schult. Syst. iii. 252 = excelsa.
punctata, *Rafin. Alsog. Am.* 62.—Am. bor.
purpurea, Tausch, in Flora, xxi. (1838) 731 = tartarica.
Purshii, G. Don, Gen. Syst. iii. 399 = stoloníferi.
racemosa, Lam. Encyc. ii. 116 = candidissima.
riparia, *Rafin. Alsog. Am.* 62.—Am. bor.
rotundifolia, *Rafin. l. c.*—Am. bor.
rubiginosa, Ehrh. Beitr. iv. 15 = C. Amomum.
rugosa, Lam. Encyc. ii. 115 = circinata.
sanguinea, Forsk. Fl. Aegypt. Arab. 88 = Cordia Myxa.
sanguinea, *Linn. Sp. Pl.* 117.—Europ.; As. bor.
sanguinea, Thunb. Fl. Jap. 68 = macrophylla.
sanguinea, Walt. Fl. Carol. 88 = stricta.
sericea, Linn. Mant. ii. 199; L'Hérit. Cornus, 8. t. 2 = C. Amomum.
sericea, *Rafin. Alsog. Am.* 60.—Am. bor.
serrulata, Hamk. & Zoll. ex Hassk. Cat. Bog. Alt. 168 = Polyosma serrulata.
sessilis, *Torr. ex Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 86.—Am. bor. occ.
sibirica, Lodd. in Lond. Hort. Brit. 50 = tartarica.
stolonifera, *Michx. Fl. Bor. Am.* i. 92.—Am. bor.

CORNUS:—
stricta, DC. Prod. iv. 272 = foemina.
stricta, A. Gray, in Hall. Pl. Texas, 11, ex S. Watts. Bibliog. Ind. N. Am. Bot. i. 488 = Drummondii.
stricta, L'Hérit. Cornus, 6 = foemina.
stricta, *Lam. Encyc.* ii. 116.—Am. bor.
stricta, Zoll. & Morr. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 10 = Polyosma integrifolia.
suecica, A. Gray, in Proc. Am. Acad. viii. (1878) 367 = canadensis.
suecica, *Linn. Sp. Pl.* 118.—Reg. bor. et arct.
suffruticosa, *Rafin. Atl. Journ.* 151.—Am. bor.
tartarica, *Mill. Gard. Dict* ed. VIII. n. 7.—Sibir.
toluccnsis, *H. B. & K. Nov. Gen. et Sp.* iii. 430.—Mexic.
tomentosa, Stend. Nom. ed. II. i. 432 = circinata.
tomentosula, DC. Prod. iv. 272 = circinata.
tomentulosa, Michx. Fl. Bor. Am. i. 91 = circinata.
Torreyi, *S. Wats. in Proc. Am. Acad.* xi. (1876) 145.—Calif.
trilobata, *Decne in Nouv. Arch. Mus. Par.* Ser. I. x. (1874) 157.—Hab. ?
unalaschkensis, *Ledeb. Fl. Ross.* ii. 878.—Alaska.
undulata, *Rafin. Alsog. Am.* 61.—Am. bor.
vernalis, Salisb. Prod. 66 = C. Mas.
virginiana, Hort ex Lam. Encyc. ii. 115 = circinata.

CORNUTA, Linn. Gen. ed. VI. 316 (1764) = **Cornutia**, Plum. (Verbenac.).

CORNUTIA, Plum. ex Linn. Gen. ed. I. 366 (1737). *VERBENACEÆ*, Benth. & Hook. f. ii. 1153.
AGNANTHUS, Vaill. ex Linn. Gen. ed. I. 366 (1737).
CORNUTA, Linn. l. c. ed. VI. 316 (1764).
HOSTA, Jacq. Hort. Schœnb. i. 10. t. 114 (1797).
HOSTANA, Pers. Syn. i. 143 (1805).
cayennensis, *Schau. in DC. Prod.* xi. 681.—Guiana.
corymbosa, Burm. f. Fl. Ind. 132 = Premna corymbosa.
corymbosa, Lam. Encyc. i. 54 = Callicarpa lanata et Premna tomentosa.
crenata, Medic. Beobachtt. (1782) 119 = pyramidata.
grandiflora, Stend. Nom. ed. II. i. 422 = acq. grandiflolia, *Schau. in DC. Prod.* xi. 682.—Mexic.
longifolia, *Sprengl. Syst.* i. 89.—Mexic.
odorata, *Poepp. ex Schau. in DC. Prod.* xi. 681.—Am. trop.
pubescens, *Guertn. f. Fruct.* iii. 172. t. 218.—Guiana.
punctata, *Willd. Sp. Pl.* iii. 322.—Am. trop.
pyramidata, Ait. Hort. Kew. ii. 358 = punctata.
pyramidata, *Linn. Sp. Pl.* 628.—Mexic.
quinata, Lour. Fl. Cochinch. ii. 387 = Vitex Loureiri.

CORNUTIOIDES, Linn. Fl. Zeyl. 195 (1747) (Gen. dub.).

CORDCEPHALUS, D. Dietr. Syn. Pl. i. 424, 540 (1839), sphalm. = **Concocephalus**, Blume (Urtic.).

CORONA-IMPERIALIS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 148 (1748) = **Fritillaria**, Linn. (Liliac.).

COROKIA, A. Cunn. in Ann. Nat. Hist. iii. (1839) 249. *CORIACEÆ*, Benth. & Hook. f. i. 949.
buddleioides, *A. Cunn. l. c.*—N. Zel.
Cotoneaster, *Raoul, Choix Pl. N. Zél.* 22.—N. Zel.

COROWANTHERA, Vieill. ex C. B. Clarke, in DC. Monog. Phan. v. 170 (1883). *GESNERACEÆ*.
aspera, *C. B. Clarke, l. c.* 172.—N. Caled.
barbata, *C. B. Clarke, l. c.*—N. Caled.
deltoidifolia, *Vieill. ex C. B. Clarke, l. c.* 171.—N. Caled.
glabra, *C. B. Clarke, l c.* 174.—N. Caled.
Pancheri, *C. B. Clarke, l. c.*—N. Caled.
pedunculosa, *C. B. Clarke, l. c.* 173.—N. Caled.
pinguior, *C. B. Clarke, l. c.* 171.—N. Caled.
pulchra, *C. B. Clarke, l. c.* 173.—N. Caled.
sertosa, *C. B. Clarke, l. c.* 174.—N. Caled.

CORONARIA, Linn. Gen. ed. I. 185 (1737) = **Lychnis**, Linn. (Caryophyll.).
Coeli-rosa, Fries, ex Heynh. Nom. ii. 161 = L. Coeli-rosa.
Flos-cuculi, A. Br. l. c. = L. Flos-cuculi.
Flos-Jovis, A. Br. l. c. = L. Flos-Jovis.
tomentosa, A. Br. l. c. = L. Coronaria.

CORONE, Hoffmgg. ex Steud. Nom. ed. II. i. 492 (1840) = **Silene**, Linn. (Caryophyll.).
agrestina, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = S. sylvestris.
gallica, Fourr. l. c. = S. gallica.
nocturna, Fourr. l. c. = S. nocturna.
quinquevulnera, Hoffmgg. ex Steud. Nom. ed. II. i. 492 = S. quinquevulnera.
vespertina, Hoffmgg. ex Steud. l. c. = S. vespertina.

CORONILLA, Tourn. ex Linn. Gen. ed. I. 327 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 509.
AVIUNCULITA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 367.
CALLISTEPHANA, Fourr. l. c. (1868).
EMERUS, Tourn. ex Mill. Gard. Dict. ed. VI. (1759).
SCORPUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 369.
aculeata, Willd. Sp. Pl. iii. 1147 = Sesbania aculeata.
argentea, Burm. f. Cap. Prod. 73; Thunb. Prod. ii. 181 = Lessertia argentea.
argentea, *Linn. Sp. Pl.* 743.—Ins. Creta.
atlantica, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 42, in syn.—Marocco.
Balansae, Boiss. & Bal. in Boiss. Fl. Orient. ii. 181 = cappadocica.
boxiifolia, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 319.—China.
cannabina, Willd. Sp. Pl. iii. 1147 = Sesbania aculeata.
cappadocica, *Willd. l. c.* 1153.—As. Min.
chia, Orph. ex Boiss. Fl. Orient. ii. 183 = parviflora.
Clusii, Dufour, in Ann. Sc. Gén. Phys. viii. (1820) 307 = minima.
coccinea, Willd. Sp. Pl. iii. 1145 = Sesbania grandiflora.
cochinchinensis, *Lour. Fl. Cochinch.* ii. 492.—Cochinch.
coronata, DC. Prod. ii. 310 = minima.
coronata, Jacq. Fl. Austr. i. 59. t. 95 = montana.
coronata, *Linn. Syst. ed.* X. 1168.—Europ. austr.
cretica, *Linn. Sp. Pl.* 743.—Europ. or.; As. Min.
dura, *Boiss. Diagn.* Ser. II. ii. 34.—Lusitan.
elegans, Pand. ex Nym. Consp. 185 = varia ?
emerus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 123 = C. Emerus.
emeroides, *Boiss. & Sprun. Diagn.* Ser. I. ii. 100.—Graecia; As. Min.
Emerus, *Linn. Sp. Pl.* 743.—Europ. austr.
Emerus, Sibth. & Sm. Fl. Graec. viii. 7. t. 710 = emeroides.
eriocarpa, Boiss. Elench. 37 = Hippocrepis squamata.
extensa, Jord. Pugill. 61 = minima.
floridа, Salisb. Prod. 341 = C. Emerus.
fruticana, *Jord. Pugill.* 62.—Europ.
gibraltarica, *Brouss. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 428.—Hispan.; Afr. bor.
glauca, Asso, ex Nym. Consp. 185 = coronata.
glauca, *Linn. Cent. Pl.* i. 28.—Reg. Mediterr.
globosa, *Lam. Encyc.* ii. 122.—Ins. Creta.
Glycyrrhiza, Spreng. Syst. iii. 324 = Glycyrrhiza asperrima.
grandiflora, *Boiss. Diagn.* Ser. I. v. 88.—As. Min.
grandiflora, Willd. Sp. Pl. iii. 1145 = Sesbania grandiflora.
Haussknechtii, Reut. ex Boiss. Fl. Orient. ii. 182 = varia.
Heinei, *Regel, Cat. Pl. Hort. Asakhor.* 41.—As. temp.
hirsuta, *DC. Prod.* ii. 210.—Afr. austr.
hirsuta, Vell. Fl. Flum. 311; vii. t. 122 = Isodesmia tomentosa.
hirta, Bunge, ex Boiss. Fl. Orient. ii. 183 = varia.
hispanica, Mill. Gard. Dict. ed. VIII. n. 4 = glauca.
iberica, Bieb. Fl. Taur. Cauc. ii. 171 = cappadocica.
juncea, *Linn. Sp. Pl.* 743.—Europ. austr.
libanotica, Boiss. Diagn. Ser. I. ix. 104 = varia.
lotoides, Koch, in Roehl. Deutschl. Fl. ed. Mert. & Koch, v. 199 = minima.
minima, Jacq. Fl. Austr. Mill. t. 271 = virginalis.
minima, *Linn. Cent. Pl.* ii. 28.—Europ.
monilis, Linn. Pl. Surin. n. 102; Amoen. Acad. viii. 261 = Moellera monilioformis ?
montana, *Scop. Pl. Carn.* ed. II. ii. 73.—Reg. Caucas.; Persia.
moniliformis, *DC. Prod.* ii. 210.—Hispan.
myrtifolia, Burm. f. Fl. Cap. Prod. 32 = Tephrosia myrtifolia.
occidentalis, Willd. Sp. Pl. iii. 1147 = Sesbania occidentalis.
orbiculata, Moench, Meth. Suppl. 40 = valentina.

CORONILLA:—
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 8.—Iberia; As. Min.
pallida, Salisb. Prod. 341 = glauca.
parviflora, Moench, Meth. 121 = cretica.
parviflora, *Willd. Sp. Pl.* iii. 1135.—Graecia; As. Min.
pauciflora, Lam. Fl. Fr. ii. 661 = C. Emerus.
pendula, *Kit. in Linnaea*, xxxii. (1863) 684.—Europ.
penuphylla, *Desf. Pl. Atlant.* ii. 171.—Afr. bor.
picta, Willd. Sp. Pl. iii. 1148 = Sesbania aegyptiaca.
pubescens, *Schum. & Thonn. in Kon. Dan. Vid. Sel. Afh.* iv. (1828) 138.—Afr. trop.
pulchra, Bell, in Journ. Bot. xi. (1873) 307 = viminalis.
ramosissima, *Ball, l. c.* 306.—Marocco.
repanda, *Boiss. Diagn.* Ser. II. ii. 86.—Hispan.
rostrata, Boiss. & Sprun. Diagn. Ser. I. ii. 100 = parviflora.
sanguinea, *Roxb. Hort. Beng.* 56, nomen.—Hab. ?
scandens, *Linn. Sp. Pl.* 743.—Ind. occ.
scandens, Vell. Fl. Flum. 310; vii. t. 121 = Chaetocalyx brasiliensis.
scorpioides, *Koch, in Roehl. Deutschl. Fl. ed Mert. & Koch*, v. 201.—Reg. Mediterr.; Persia.
Securidaca, Linn. Sp. Pl. 743 = Securigera Coronilla.
Securigera, Salisb. Prod. 341 = Securigera Securidaca.
sericea, Willd. Enum. Hort. Berol. 778 = Sesbania sericea.
Sesban, Moench, Meth. 120 = Sesbania aegyptiaca.
squamata, Cav. Ic. ii. 48 = Hippocrepis squamata.
stipularis, Lam. Encyc. ii. 120 = valentina.
stipulata, *Buch.-Ham. ex Don, Prod. Fl. Nep.* 246. —Reg. Himal.
suaveolens, Salisb. Prod. 340 = valentina.
suratiensis, Garc. ex DC. Prod. ii. 323 = Smithia thymifolia.
thymifolia, *Burm. f. Fl. Cap. Prod.* 22.—Afr. austr.
vaginalis, *Lam. Encyc.* ii. 121.—Europ.
virginalis, Ledeb. Fl. Ross. i. 652 = sulcorhiza.
valentina, Lam. Fl. Fr. ii. 663 = coronata.
valentina, *Linn. Sp. Pl.* 743.—Europ. austr.
valentina, Pall. ex Bieb. Fl. Taur. Cauc. ii. 178 = parviflora.
varia, *Linn. Sp. Pl.* 743.—Europ.
viminalis, *Salisb. Parad. Lond.* t. 18.—Afr. bor.
virgata, Willd. Sp. Pl. iii. 1148 = Courtoisia virgata.

CORONOCARPUS, Schum. & Thonn. Beskr. Guin. Pl. 303 (1827) = **Aspilia**, Thou. (Compos.).
Gayanus, Benth. in Hook. Niger Fl. 484 = A. helianthoides.
helianthoides, Benth. l. c. = A. latifolia.
helianthoides, Schum. & Thonn. Beskr. Guin. Pl. 303 = A. helianthoides.
Kotschyi, Benth. in Hook. Niger Fl. 483 = A. Kotschyana.
Prieurianus, Benth. l. c. = A. Kotschyana.

CORONOPUS, Reichb. Handb. 202 (1837) = **Plantago**, Tourn.
vulgaris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 140 = P. Coronopus.

CORONOPUS, Rupp. ex Linn. Syst. ed. I (1735); Gaertn. Fruct. ii. 398 (1791) = **Senebiera**, DC. (1799) (Crucif.).
anomalus, Spreng. Syst. ii. 853 = Brachycarpaea variana.
coadunatus, Stokes, Bot. Mat. Med. iii. 440 = S. Coronopus.
cristatus, [Clairv.] Man. Herb. 219 = S. Coronopus.
depressus, Moench, Meth. 290 = S. Coronopus.
didymus, Sm. Fl. Brit. ii. 691 = S. didyma.
fungosus, Sm. ex DC. Syst. ii. 592 = S. integrifolia.
Helenae, Spreng. Syst. ii. 853 = *S. Helenianus.*
incisus, Hornem. Hort. Hafn. 599 = S. didyma.
integrifolius, Spreng. Syst. ii. 853 = S. integrifolia.
linoides, Spreng. l. c. = S. integrifolia.
myricoides, Salisb., Sm. ex DC. Syst. ii. 592 = S. serrata.
niloticus, Spreng. Syst. ii. 853 = S. niloticus.
pinnatifidus, Dulac, Fl. Hautes-Pyr. [186] = S. didyma.
pinnatus, Hornem. Hort. Hafn. 599 = S. didyma.
procumbens, Gilib. Fl. Lituan. ii. 52 = S. Coronopus.
Raddii, Savi, in Mem. G. Raddi, (1820) 88.—Aegypt.
Ruellii, All. Fl. Pedem. i. 256 = S. Coronopus.
serratus, Desv. Journ. Bot. iii. (1814) 169 = S. serrata.
squamatus, Arcbers. Fl. Brandenb. 69 = S. Coronopus.
vulgaris, Desf. Tabl. ed. I. 122 = S. Coronopus.

COROTHAMNUS, Presl (Bot. Bemerk. 187); in Abh. Boehm. Ges. V. iii. (1844) 567 = **Genista**, Linn. (Legumin.).
decumbens, C. Koch, Dendrol. i. 44 = Cytisus decumbens.
diffusus, Presl, Bot. Bemerk. 187 = Cytisus decumbens.
Halleri, Presl, l. c. = Cytisus decumbens.
myrtifolius, Presl, l. c.—Hungar.
procumbens, Presl, l. c. = G. procumbens, *Waldst.*

COROZO, Jacq. ex Gaertn. Fruct. 42 (1792) = **Elaeis**, Jacq. (Palm.).

CORPODETES, Reichb. Consp. 61 (1828) = CARPODETES, Herb. = **Stenomesson**, Herb. (Amaryll.).

CORRAEA, Sm. in Trans. Linn. Soc. iv. (1798) 219 = seq.

CORREA, Andr. Bot. Rep. t. 18 (1798). *RUTACEAE*, Benth. & Hook. f. i. 384.
ANTOMMARCHIA, Colla, Hort. Ripul. App. ii. 345 (1836).
DIDIMERIA, Lindl. in Mitch. Three Exp. Austral. ed. II. ii. 198 (1889).
MAZEUTOXERON, Labill. Voy. ii. 11 (1798).
acmula, *F. Muell. Fragm.* i. 8.—Austral.
alba, *Andr. Bot. Rep.* t. 18.—Austral.
Backhousiana, Hook. Journ. Bot. i. (1834) 353 = speciosa.
Bauerlenii, *F. Muell. in Proc. Linn. Soc. N. S. W.* ix. (1885) 960.—Austral.
bicolor, Paxt. Mag. Bot. ix (1842) 267 = speciosa.
cardinalis, F. Muell. ex Hook. Bot. Mag. t. 4912 = speciosa.
cordifolia, Lindl. in Mitch. Three Exp. Austral. ii. 233 = speciosa.
cotinifolia, Salisb. Parad. Lond. t. 100 = alba.
decumbens, *F. Muell. in Trans. Vict. Inst.* i. (1855) 30.—Austral.
ferruginea, Hook. Comp. Bot. Mag. i. (1835) 276 = Lawrenciana.
glabra, Lindl. in Mitch. Three Exp. Austral. ii. 48 = speciosa.
Latrobeana, F. Muell. in Dietr. Fl. Univ. N. Ser. t. 11 = Lawrenciana.
Lawrenciana, *Hook. Journ. Bot.* i. (1834) 254.—Austral.
leucocladda, Lindl. in Mitch. Three Exp. Austral. ii. 39 = speciosa.
pulchella, Mackay, ex Sweet, Fl. Austral. t. 1 = speciosa.
reflexa, Vent. Jard. Malm. sub t. 13 = speciosa.
revoluta, Vent. l. c. sub t. 13 = speciosa.
rotundifolia, Lindl. in Mitch. Three Exp. Austral. ii. 219 = alba.
rubra, Sm. Exot. Bot. ii. 26 = speciosa.
rufa, Vent. Jard. Malm. t. 13 adnot. = alba.
Schlechtendalii, Behr, in Linnaea, xx. (1847) 680 = speciosa.
speciosa, *Ait. Epit. Hort. Kew.* 266.—Austral.
virens, *Hook. Journ. Bot.* i. (1834) 253.—Austral.
virens, Sm. Exot. Bot. ii. 25. t. 72 = speciosa.
viridiflora, Andr. Bot. Rep. t. 480 = speciosa.

CORREAS, Hoffmgg. Verz. Pfl. Nachtr. i. 168 (1824) = Correa, Andr. (Rutac.).

CORREIA, Vell. in Roem. Script. 196. t. 6 (1796) = **Gomphia**, Schreb. (Ochnac.).

CORRIGIOLA, Linn. Gen. ed. I. 340 (1737). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 17.
POLYGONARIA, Brot. Symbol. Phyt. 6 (1765).
POLYGONIPOLIA, Adans. Fam. ii. 272 (1763).
albella, Forsk. Fl. Aegypt. Arab. 207 = Paronychia arabica.
andina, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 146.—N. Granada.
capensis, *Willd. Sp. Pl.* i. 1507.—Afr. austr.
daltoidea, Hook. & Arn. Bot. Beech. Voy. 24 = Monocosmia corrigioloides.
glomerulifera, *Steud. Nom.* ed. II. i. 428.—Chili.
imbricata, Lapeyr. Fl. Pyr. i. 33; Abr. 169 = littoralis.
latifolia, *C. Gay, Fl. Chil.* ii. 217.—Chili.
littoralis, *Linn. Sp. Pl.* 271.—Europ.
littoralis, Thunb. Prod. Pl. Cap. 28 = capensis.
propinqua, *C. Gay, Fl. Chil.* ii. 217.—Chili.
pseudomorphoides, *Baker, in Journ. Linn. Soc.* xx. (1883) 233.—Madag.

CORRIGIOLA :—
repens, Forsk. Fl. Aegypt. Arab. 207 = Polycarpaea fragilis.
squamosa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 337.—Chili.
telephiifolia, Pourr. in Mém. Acad. Toul. iii. (1788) 316 ; (Chl. Narb. 20) = littorella.

CORSIA, Becc. Malesia, i. 288. t. 9 (1878). *BURMANNIACEAE*, Benth. & Hook. f. iii. 660.
ornata, *Becc. l. c.* 289.—N. Guin.

CORTESIA, Cav. Ic. iv. 58. t. 377 (1797). *BORAGINEAE*, Benth & Hook. f. ii. 941.
cuneata, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 439 = cuneifolia.
cuneifolia, *Cav. Ic.* iv. 58. t. 377.—Reg. Argent.
cuneiformis, Poir. in Dict. Sc. Nat. x. 568 = cuneifolia.
microphylla, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868) 430.—Am. austr.

CORTHUMIA, Reichb. Nom. 201 (1841) = **Pelargonium**, L'Hérit. (Geran.).

CORTHUSA, Reichb. Consp. 128 (1828) = **Cortusa**, Linn. (Primul.).

CORTIA, DC. Prod. iv. 186 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 915.
SCHULTZIA, Wall. Cat. n. 589 (1830).
elata, Edgew. in Trans. Linn. Soc. xx. (1851) 55 = Ligusticum elatum.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 709.—Reg. Himal.
Lindleyi, *DC. Prod.* iv. 187.—Reg. Himal.
vaginata, Edgew. in Trans. Linn. Soc. xx. (1851) 55 = Selinum vaginatum.

CORTUSA, Linn. Gen. ed. I. 40 (1737). *PRIMULACEAE*, Benth. & Hook. f. ii. 633.
Gmelini, Linn. Sp. Pl. 144 = Androsace Gmelini.
grandiflora, Schur. Enum. Pl. Transs. 556 = Matthioli.
Matthioli, *Linn. Sp. Pl.* 144.—Europ. ; As. bor.
pubens, *Schott, Nyman & Kotschy, Analect. Bot.* 17. —Transylv.
Semenovii, *Herd. in Bull. Soc. Nat. Mosc.* xli. (1868) i. 68.—As. centr.
sibirica, *Andrz. ex Steud. Nom.* ed. II. i. 423.—Sibir. baical.

CORTUSINA, Eckl. & Zeyh. Enum. 77 (1835) = **Pelargonium**, L'Hérit. (Geraniac.).
odoratissima, Eckl. & Zeyh. l. c. = P. odoratissimum.
reniformis, Eckl. & Zeyh. l. c. = P. reniforme.
rubropurpurea, Eckl. & Zeyh. l. c. = P. echinatum.
sidaefolia, Eckl. & Zeyh. l. c. = P. sidoides.
volubina, Eckl. & Zeyh. l. c. = P. volutinum.

CORVINA, Steud. Nom. ed. II. i. 356 (1840) sphalm. = **Muricaria**, Desv. (Crucif.).
prostrata, Steud. l. c. = M. prostrata.

CORVINIA, Stadtm. ex Willem. in Usteri, Ann. Bot. xviii. 28 (1796) = **Nephelium**, Linn. (Sapindac.).
Litschi, Stadtm. ex Willem. l. c. = N. Litchi.

CORVISARTIA, Mérat, Fl. Par. ed. I. 238 (1812) = Inula, Linn. (Compos.).
caucasica, G. Don, ex Loud. Hort. Brit. 350 = Buphthalmum speciosum.
Helenium, Mérat, Fl. Par. ed. I. 238 = I. Helenium.
indica, Lindl. Bot. Reg. (1842) Misc. 58 = I. Royleana.

CORYANTHES, Hook. Bot. Mag. t. 3102 (1831). *ORCHIDEAE*, Benth. & Hook. f. iii. 549.
CORYTHANTHES, Lem. in Orth. Diet. iv. 259 (1849).
PANSTRIPH, Rafin. Fl. Tellur. iv. 41 (1836).
MELICLIS, Rafin. l. c. ii. 99 (1836).
Albertinae, Karst. Answ. Neuer Gew. Venes. 5 = maculata.
Barberii, Beer, Prakt. Orch. 223, sphalm. = *Parberi*.
biflora, *Rodrig. Orch. Nov.* i. 102.—Bras.
Bruchmuelleri, *Reichb. f. in Linnaea*, xli. (1877) 74.—N. Granad.
cinguettissima, Mast. in Gard. Chron. (1885) i. 144 = elegantissima.
elegantissima, *Linden & Reichb. f. in Gard. Chron.* (1868) 1114.—Bras.
fimbriata, Reichb. f. Beitr. Syst. Pfl. 42 = Coryanthes fimbriata.
fornicata, Reichb. f. l. c. 67 = Coryanthes fornicata.
macrantha, Reichb. f. l. c. = Coryanthes macrantha.
mucronata, Reichb. f. l. c. = Coryanthes mucronata.
oblonga, Reichb. f. l. c. = Coryanthes oblonga.
picta, Reichb. f. l. c. = Coryanthes picta.
pruinosa, Reichb. f. l. c. 43 = Coryanthes fimbriata.
rivularis, Reichb. f. l. c. 67 = Coryanthes rivularis.
rotundifolius, Reichb. f. l. c. = Coryanthes rotundifolia.
trilobus, Reichb. f. l. c. = Coryanthes trilobs.
unguiculatus, Reichb. f. l. c. 43 = Coryanthes unguiculata.

CORVCARPUS, *Zea*, ex Spreng. Syst. i. 128 (1825) = DIACHASMA, Rafin. (Gramin.).
arundinaceus, *Zea*, ex Spreng. l. c. = D. americana.

CORYCIUM, Sw. in Vet. Acad. Nya Handl. Stockh. xxi. (1800) 220. *ORCHIDEAE*, Benth. & Hook. f. iii. 533.
bicolorum, *Sw. l. c.* 222.—Afr. austr.
bifidum, *Sond. in Linnaea*, xix. (1847) 111.—Afr. austr.
crispum, *Sw. in Vet. Acad. Nya Handl. Stockh.* xxi. (1800) 222.—Afr. austr.
exclusum, *Lindl. Gen. et Sp. Orch.* 368.—Afr. austr.
lignlatum, Reichb. f. in Linnaea, xix. (1847) 875 = bifidum.
microglossum, *Lindl. Gen. et Sp. Orch.* 369.—Afr. austr.
nigrescens, *Sond. in Linnaea*, xix. (1847) 110.—Afr. austr.
orobanchoides, *Sw. in Vet. Acad. Nya Handl. Stockh.* xxi. (1800) 222.—Afr. austr.
vestitum, *Sw. l. c.*—Afr. austr.

CORYDALIS, Vent. Choix, 19 (1803), in nota ; DC. Fl. Fr. iv. 636 (1805). *PAPAVERACEAE*, Benth. & Hook. f. i. 55.
BORCKHAUSENIA, Gaertn. Mey. & Scherb. Fl. Wett. iii. 4 (1801).
BULBOCAPNOS, Bernh. in Linnaea, viii. (1833) 469.
CAPNOIDES, Adans. Fam. ii. 431 (1763).
CERATOCAPNOS, Dur. in Parl. Giorn. Bot. Ital. i. 1. (1844) 326.
CRYPTOCERAS, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 121.
CYSTICAPNOS, Boerh. ex Adans. Fam. ii. 431 (1763).
NECKERIA, Scop. Introd. 313 (1777).
PHACOCAPNOS, Bernh. in Linnaea, xii. (1838) 664.
PISTOLOCHIA, Bernh. Syst. Verz. Erf. 57 (1800).
PSEUDOFUMARIA, Medic. Phil. Bot. I. 110 (1789).
SOPHOROCAPNOS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 570.
acaulis, Pers. Syn. ii. 270 = ochroleuca.
adiantifolia, *Hook. f. & Thoms. Fl. Ind.* i. 271.—Reg. Himal.
adnata, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1879) 29.—China.
albicaulis, *Franch. Pl. David.* i. (1884) 86.—As. or.
albiflora, *Kit. in Linnaea*, xxxii. (1863) 491.—Hungar.
alpestris, C. A. Mey. Vers. Pfl. Casc. 176 = pauciflora.
alpina, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 259 = bracteata.
alpina, C. Koch, in Linnaea, xv. (1841) 252 = rutaefolia.
ambigua, *Cham. & Schlecht. in Linnaea*, i. (1826) 558.—Kamtschat.
angustifolia, *DC. Syst.* ii. 120.—Reg. Caucas. ; Persia.
annua, Hoffmgg. ex Steud. Nom. ed. II. i. 423 = glauca.
astragalina, Hook. f. & Thoms. Fl. Ind. i. 270 = stricta.
aurea, *Willd. Enum. Hort. Berol.* 740.—Am. bor.
biauritia, Hornem. Hort. Hafn. ii. 668 = Dicentra formosa.
bicalcarata, Haw. ex G. Don, in Loud. Hort. Brit. Suppl. i. 186 = rutaefolia.
Bidwelliae, *S. Wats. in Bot. Calif.* ii. 429.—Calif.
blanda, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 149 = tuberosa.
bracteata, *Pers. Syn.* ii. 269.—Sibiria.
bracteosa, Spreng Syst. iii. 162 (= *Dicentra bracteosa*). —Am. bor. ?
Brandegei, *S. Wats. in Bot. Calif.* ii. 430.—Calif.
breviflora, *DC. Syst.* ii. 127.—Kamtschat.
bulbosa, *DC. Fl. Fr.* iv. 637.—Europ.
bulbosa, Pers. Syn. ii. 269 = tuberosa.
bulbosa, Turcz. ex Steud. Nom. ed. II. i. 423 = Turtschaninovii.
Bungeana, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 62. —Sibir.
Burmanni, *Eckl. & Zeyh. Enum.* 4.—Afr. austr.
canadensis, Goldb. in Edinb. Phil. Journ. vi. (1822) 259 = Dicentra canadensis.
capnoides, Pers. Syn. ii. 270 = lutea.
Cascana, *A. Gray, in Proc. Am Acad.* x. (1874) 69.—Calif.
cashmeriana, *Royle, Illustr. Bot. Himal.* 69. t. 16. f. 1. —Reg. Himal.
caucasica, Adams, ex Steud. Nom. ed. II. i. 423 = angustifolia.
caucasica, *DC. Syst.* ii. 119.—Reg. Caucas.
caudata, Pers. Syn. ii. 269 = longiflora.
cava, Schweigg. & Kort. Fl Erlang. ii. 44 ; Wahlenb. Veg. Helv. 126 [bis] = tuberosa.
chaerophylla, *DC. Prod.* i. 128.—Reg. Himal.
chinensis, Franch. Fl. David i. 28 = edulis.
clavicula, S. F. Gray, Nat. Arr. Brit. Pl. ii. 700 = seq.
claviculata, *DC. Fl. Fr.* iv. 638.—Europ.
conorhiza, *Ledeb. Fl. Ross.* i. 99.—Reg. Caucas.
cornuta, *Royle, Illustr. Bot. Himal.* 69.—Reg. Himal.
Cracca, *Chum. & Schlecht. in Linnaea*, i. (1826) 567. —Afr. austr.
crassifolia, *Royle, Illustr. Bot. Himal.* 68.—Reg. Himal.
crassissima, Jacquem. Voy. Bot. 12. t. 11 = crassifolia.
crithmifolia, *Royle, Illustr. Bot. Himal.* 68.—Reg. Himal.
crystallina, *Engelm. ex A. Gray, Man. U. St. Bot.* ed. V. 62.—Am. bor.
Cucullaria, Pers. Syn. ii. 269 = Dicentra Cucullaria.
curviatiloua, *Engelm. ex A. Gray, in Mem. Bot. U. St.* ed. V. 62.—Am. bor. occ.
Casicki, *S. Wats. in Bot. Calif.* ii. 430.—Calif.
dasypteva, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 28.—China.
debilis, Edgew. in Trans. Linn. Soc. xx. (1851) 30 = cornuta.
decipiens, Schott, Nyman & Kotschy, Analect. Bot. 42 = pumila.
decumbens, *Pers. Syn.* ii. 269.—Japon.
densiflora, J. & C. Presl, Delic. Prag. 10 = bulbosa.
depauperata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 51 = bulbosa.
diphylla, Pers. Syn. ii. 269 = bulbosa.
diphylla, Wall. Tent. Fl. Nep. 54 = rutaefolia.
edulis, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 30. —China.
elegans, *Wall. Cat.* n. 1485.—Reg. Himal.
Emanueli, C. A. Mey. Vers. Pfl. Casc. 176 = pauciflora.
enneaphylla, DC. Fl. Fr. Suppl. 587 = Sarcocapnos enneaphylla.
epithymifolia, Walp. Rep. i. 120, sphalm. = orithmifolia.
Erdelii, Zucc. in Abh. Akad. Muench. iii. (1848) 251 = rutaefolia.
eximia, Link, Enum. Hort. Berol. ii. 218 = Dicentra eximia.
fabacea, *Pers. Syn.* ii. 269.—Europ.

CORYDALIS:—
Falconeri, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 124 —Reg. Himal.
Fedtschenkoana, *Regel, Pl. Nov. Fedsch.* 3.—As. centr.
fibrosa, *Maxim. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1857) 397, nomen.—Manchur.
filiformis, Royle, Illustr. Bot. Himal. 68 = sibirica.
fimbellata, *Edgew. in Trans. Linn. Soc.* xx. (1851) 80.—Reg. Himal.
flaccida, *Hook. f. & Thoms. Fl. Ind.* i. 260.→Reg. Himal.
flavidula, Chapm. Fl. S. U. St. Suppl. 604 = flavula.
flavula, *DC. Prod.* i. 129.—Am. bor.
formosa, Pursh, Fl. Am. Sept. ii. 462 = Dicentra canadensis, eximia.
formosa, Spreng. Syst. iii. 162 = Dicentra formosa.
fumariaefolia, *Maxim. in Prim. Fl. Amur.* 89.—Manchur.
fungosa, Vent. Choix, t. 19 = Adlumia cirrhosa.
gamosepala, Maxim. Prim. Fl. Amur. 88 = bulbosa.
Gebleri, Ledeb. Ind. Hort. Dorpat. (1828) 3 = capnoides.
geranifolia, Hook. f. & Thoms. Fl. Ind. i. 269 = chaerophylla.
gigantea, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 15).—Manchur.
glauca, *Pursh, Fl. Am. Sept.* ii. 462.—Am. bor.
glaucescens, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) i. 253.—As. temp.
Gortschakovii, *Schrenk, Enum. Pl. Nov.* i. 100.—Reg. Himal.; Soongar.
Govaniana, *Wall. Tent. Fl. Nep.* 55.—Reg. Himal.
gracilipes, S. Moore, in journ. Bot. xiii. (1875) 226 = decumbens.
gracilis, *Ledeb. Fl. Ross.* i. 101.—Reg. Caucas.
Griffithii, Boiss. Diagn. Ser. II. i. 15 (*Griffithsii*) = rutaefolia.
Halleri, Willd. Enum. Hort. Berol. 740 = bulbosa.
Hamiltoniana, G. Don, Gen. Syst. i. 148 = rutaefolia.
heterocarpa, *Bull, in Journ. Linn. Soc.* xvi. (1877) 314.—Afr. bor. occ.
heterocarpa, Sieb. & Zucc. in Abh. Akad. Muench. iv. (1843) 11. 173 = pallida.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 139 = mucronifolia.
Impatiens, Fisch. in DC. Syst. ii. 124 = sibirica.
incisa, *Pers. Syn.* ii. 269.—Japon.
inconspicua, *Bunge, ex Ledeb. Fl. Ross.* i. 104.—Sibiria.
intermedia, Mérat, Fl. Par. 273 = lutacea.
juncea, *Wall. Tent. Fl. Nep.* 54. t. 42.—Reg. Himal.
Karelliana, *Prita. ex Walp. Rep.* ii. 750.—As. centr.
kashmiriana, Anct.=cashmeriana.
kashgarica, *Rupr. Sert. Tianach.* 38.—As. centr.
Kolpakowskiana, *Regel, in Act. Hort. Petrop.* v. (1877) 633.—Turkestan.
lachenaliaeflora, Fisch. ex DC. Syst. ii. 111 = Dicentra lachenaliaeflora.
laevigata, E. Mey. ex Bernh. in Linnaea, xii. (1838) 264 = C. Cracca.
latifiora, *Hook. f. & Thoms. Fl. Ind.* i. 270.—Reg. Himal.
laxa, *Fraenck. & Sav. Enum. Pl. Jap.* ii. 274.—Japon.
laxa, *Fries, Summa Veg. Scand.* 145.—Fennia.
Leiobotrya, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 377.—Sibiria.
leptocarpa, *Hook. f. & Thoms. Fl. Ind.* i. 260.—Reg. Himal.
libanotica, Hochst. in Lorent, Wanderungen, 339 = rutaefolia.
linarioides, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 27.—China.
linearifolia, Sieb. & Zucc. in Abh. Akad. Muench. iv. 11. (1843) 174 = bulbosa.
Lobelii, Tausch, ex Hoppe, in Flora, xxii. (1839) 279 = pumila.
longiflora, *Pers. Syn.* ii. 269.—Reg. Casp.; Sibir.
longipes, DC. Prod. i. 128 = sibirica.
longipes, D. Don, Prod. Fl. Nep. 196 = rutaefolia.
lutea, *DC. Fl. Fr.* iv. 938.—Europ.
Maackii, *Rupr. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 71.—Manchur.
macrocentra, *Regel, l. c.* viii. (1884) 694.—As. centr.
macrophylla, *Mutt. in Terr. & Gray, Fl. N. Am.* i. 69. —Am. bor. occ.
macrosepala, *Sered. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 71.—Reg. Caucas.
Marschalliana, *Pers. Syn.* ii. 269.—Russia; As. Min.

CORYDALIS:—
media, *Wall. Tent. Fl. Nep.* 55.—Reg. Himal.
melanochlora, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 26.—China.
micropoda, *Franch. Pl. Davidian.* i. (1884) 29.—As. or.
mucronifera, Engelm. ex A. Gray, Man. U. St. Bot. ed. V. 62 = aurea.
Moorcroftiana, *Wall. Cat.* n. 1492.—Reg. Himal.
nana, *Royle, Illustr. Bot. Himal.* 68.—Reg. Himal.
nivalis, Boiss. & Hsut, in Diagn. Ser. II. v. 16 = pauciflora.
nobilis, *Pers. Syn.* ii. 269.—Sibir.
nudicaulis, *Regel, in Act. Hort. Petrop.* viii. (1884) 695.—As. centr.
ochotensis, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 68. —Sibir.
ochroleuca, *Koch, in Flora*, xiv. (1831) 708.—Italia.
ophiocarpa, *Hook. f. & Thoms. Fl. Ind.* i. 259.—Reg. Himal.
oppositifolia, DC. Syst. ii. 114 = rutaefolia.
orthoceras, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. (1843) 11. 175.—Japon.
pauciflora, *Pers. Syn.* ii. 269.—Sibir.; Am. bor. occ.
pallida, *Pers. l. c.* 270.—China; Japon.
paniculigera, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 3.—As. centr.
parnassica, *Orph. & Heldr. in Boiss. Diagn.* Ser. II. vi. 9.—Graecia.
pauciflora, Edgew. in Trans. Linn. Soc. xx. (1851) 80 = rutaefolia.
pauciflora, *Pers. Syn.* ii. 269.—As. bor.; Am. bor. occ.
persica, *Chasm. & Schlecht. in Linnaea*, i. (1826) 567. —Persia.
phyzocarpa, Cambess. in Jacquem. Voy. Bot. 18. t. 12 = crassifolia.
polygalina, *Hook. f. & Thoms. Fl. Ind.* i. 263.—Reg. Himal.
prainosa, *E. Mey. ex Bernh. in Linnaea*, xii. (1838) 264, in syn.—Afr. austr.
pseudo-cava, Pantoč. in Verh. Ver. Natur- u. Heilk. Preßb. ii. (1871-72) 88 = tuberosa.
pseudo-pumila, Schur, Enum. Pl. Transs. 37 = pumila.
pulchella, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1881) 131.—Afghan.
pumila, *Koch, Syn. Fl. Germ.* ed. II. 34.—Europ.
racemosa, Bunge, Enum. Pl. Chin. Bor. 5 = Bungeana.
racemosa, *Pers. Syn.* ii. 270.—Japon.
Raddeana, *Fisch. ex Bull. Soc. Nat. Mosc.* xxxiv. (1861) 11. 143.—Manchur.
ramosa, *Wall. Cat.* n. 1494.—Reg. Himal.
remota, Fisch. ex Maxim. Prim. Fl. Amur. 87 = bulbosa.
rosea, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 28. —China.
rosea, *Zeph. ex Steud. Nom.* ed. II. i. 424.—Afr. austr.
rupestris, *Kotschy, ex Boiss. Diagn.* Ser. I. vi. 8.— Persia.
rutaceo, *T. M. Fr. in Bot. Notizer* (1854) 89.—Sibir.
rutaefolia, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 13 = persica.
rutaefolia, *DC. Syst.* ii. 115.—As. Min.; Persia; Reg. Himal.
scandens, Spreng. Syst. iv. Cur. Post. 365 = Dicentra spectabilis.
Schanginiana, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 4.—As. centr.
Scouleri, *Hook. Fl. Bor. Am.* i. 86. t. 14.—Am. bor. occ.
Semenowii, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxvi. (1864) i. 407.—Sibir.
semperrirens, Pers. Syn. ii. 269 = glauca.
senanensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 273.— Japon.
Seroszowii, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) i. 256.—As. centr.
Shearsi, *S. Moore, in Journ. Bot.* xiii. (1875) 226.— China.
sibirica, *Pers. Syn.* ii. 270.—Sibir.; Reg. Himal.
solida, [Scr. in] Svensk Bot. viii. 561 (1819) = bulbosa.
speciosa, Maxim. Prim. Fl. Amur. 89 = pallida.
spectabilis, Pers. Syn. ii. 269 = Dicentra spectabilis.
stenopetala, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 26.—China.
stricta, *Steph. ex DC. Syst.* ii. 128.—Sibir.; Reg. Himal.
Sommieri, *Pantoč. in Magyar Növénz. Lapok*, v. (1881) 166.—Hungar.

CORYDALIS:—
suaveolens, *Hance, in Journ. Bot.* xviii. (1880) 268.— China.
tenella, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 148.—As. centr.
tenella, *Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb.* ii. (1837) 313.—Reg. Caucas.
tenuifolia, Pursh, Fl. Am. Sept. ii. 462 = Dicentra lachenaliaeflora.
tenuis, Schott, Nyman & Kotschy, Analect. Bot. 43 = bulbosa.
thalictrifolia, *Jamesion, ex Regel, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) 11. 148.—China.
tibetica, *Hook. f. & Thoms. Fl. Ind.* i. 266.—Reg. Himal.
trachycarpa, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 27.—China.
triternata, Cav. in Abh. Akad. Muench. iii. (1843) 251 = bulbosa.
tuberosa, *DC. Fl. Fr.* iv. 627.—Europ.
Turczaninowii, *Bess. in Flora*, xvii. (1834) i. *Beibl.* 6. —Russia.
uniflora, Nym. Syll. 185 = rutaefolia.
urulensis, *Fisch. ex DC. Syst.* ii. 127.—Mont. Ural.
vaginans, *Royle, Illustr. Bot. Himal.* 68.—Reg. Himal.
Vernyi, *Franch. & Sav. Enum. Pl. Jap.* ii. 273.— Japon.
verticillaris, DC. Syst. ii. 114 = rutaefolia.
vesicaria, *Pers. Syn.* ii. 269.—Afr. austr.
Wilfordi, Regel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) ii. 148 = pallida.

CORYDALLIS, Aschers. Fl. Brandenb. 28 (1864) = praec.

CORYDANDRA, Reichb. Nom. 58 (1841) = Galeandra, Lindl. (Orchid.).

CORYLOPSIS, Sieb. & Zucc. Fl. Jap. i. 46. tt. 19, 20 (1835). *HAMAMELIDEAE*, Benth. & Hook. f. i. 667.
glabrescens, *Franch. & Sav. Enum. Pl. Jap.* ii. 367.— Japon.
himalayana, *Grif. in Journ. As. Soc. Beng.* xxiii. (1854) 641.—Reg. Himal.
Kesakii, *Sieb. & Zucc. Fl. Jap.* i. 49.—Japon.
multiflora, *Hance, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 224.—China.
pauciflora, *Sieb. & Zucc. Fl. Jap.* i. 48. t. 20.—Japon.
spicata, *Sieb. & Zucc. l. c.* 45. t. 19.—Japon.

CORYLUS, [Tourn.] Linn. Syst. ed. I (1735). *CUPULIFERAE*, Benth. & Hook. f. iii. 406.
alba, Ait. ex Steud. Nom. ed. II. i. 421 = C. Avellana.
americana, Siebold, in Verh. Bat. Gen. xii. (1830) 36 = heterophylla.
americana, *Walt. Fl. Carol.* 236.—Am. bor.
americana, Willd. Sp. Pl. iv. 471 = humilis.
arborea, Hort. ex Steud. Nom. ed. II. i. 424 = C. Avellana.
arborescens, Gaertn. Mey. & Scherb. Fl. Wett. iii. 276 = tubulosa.
arborescens, Muench. Hesse. iii. 236 = C. Colurna.
atropurpurea, Hort. ex A. DC. in DC. Prod. xvi. 11. 182 = tubulosa.
Avellana, *Linn. Sp. Pl.* 996.—Europ.; As. Min.
Avellana, Thunb. Fl. Jap. 160 = heterophylla.
Bianntina, Desf. Tabl. ed. II. 245 = C. Colurna.
californica, Hort. ex C. Koch, Dendrol. ii. 11. 15 = rostrata.
Colurna, *Linn. Sp. Pl.* 999.—Europ. or; As. Min.; Himal.
corylodes, Desrol, ex Steud. Nom. ed. I. 229 = rostrata.
Davidiana, *Baill. Hist. Pl.* vi. 234.—China.
ferox, *Wall. Cat.* n. 2797.—Reg. Himal.
filiciphila, Hort. ex A. DC. in DC. Prod. xvi. 11. 181 = C. Avellana.
grandis, [Drymod. in] Ait. Hort. Kew. ed. I. iii. 368 = C. Avellana.
Heterophylla, *Siebold, Kruidk. Naamsl.* 27.—Japon.
heterophylla, Fisch. ex Rom. in Flora, xvii. (1834) i. *Beibl.* 34 = C. Avellana.
humilis, Willd. Enum. Hort. Berol. 988 = americana.
intermedia, Fingerh. in Linnaea, iv. (1829) 365 = C. Colurna.
Jacquemontii, Decne. in Jacquem. Voy. Bot. 200. t. 169 = C. Colurna.
lacera, Wall. Cat. n. 2796 = C. Colurna.

CORYLUS;—
laciniata, Hort. ex A. DC. in DC. Prod. xvi. ii. 181 = C. Avellana.
mandshurica, Maxim. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 187 = rostrata.
maxima, Mill. Gard. Dict. ed. VIII. n. 2 = C. Avellana.
mexicana, Hort. ex C. Koch, Dendrol. ii. 11. 15 = rostrata.
mongolica, Hort. ex C. Koch, l. c. 13 = C. Avellana.
oweta, Lam. ex Steud. Nom. ed. II. i. 424 = C. Avellana.
pontica, Dochmahl, Fuehr. Obstkunde, iv. 38 = C. Avellana.
pontica, C. Koch, in Linnæa, xxii. (1849) 329 = C. Colurna.
pumila, Lodd. ex Loud. Hort. Brit. Suppl. iii. 538 = C. Avellana.
purpurea, Hort. ex A. DC. in DC. Prod. xvi. ii. 182 = tubulosa.
quercifolia, Hort. ex A. DC. l. c. 181 = C. Avellana.
rostrata, *Ait. Hort. Kew.* iii. 364.—Am. bor.
Serényiana, Pinak. in Oestr. Bot. Wochenbl. ii. (1852) 394 = C. Avellana.
Sieboldiana, Blume, Mus. Bot. Lugd. Bat. i. 310 = heterophylla.
sylvestris, Salisb. Prod. 393 = C. Avellana.
tetraphylla, Lodeb. in Denkschr. Baier. Bot. Ges. iii. (1841) ii. 56, sphalm. = C. Avellana.
Thunbergii, Hort. ex C. Koch, Dendrol. ii. 11. 13 = heterophylla.
tiliacea, Jacquem. Voy. Bot. 160 = C. Colurna.
tubulosa, *Willd. Sp. Pl.* iv. 470.—Europ. austr.; As. Min.
Turczaninowii, Steud. Nom. ed. II. i. 425, sphalm. = Corydalis Turczaninowii.

CORYMBIS, Thou. Orch. Iles Afr. t. 87, 88 (1822). *ORCHIDEÆ*, Benth. & Hook. f. iii. 591.
CALOIDIA, Lindl. Gen. et Sp. Orch. 494 (1840).
CORYMBORCHIS, Thou. ex Blume, Orch. Arch. Ind. 125 (1855).
HYSTERIA, Reinw. in Blume, Cat. Gew. Buitenz. 99 (1839).
MACROSTYLIS, Breda, Orch. Kubl. et Hass. (1827 ?).
RHYNCHANTHERA, Blume, Bijdr. Tabell. 78 (1826).
TOMOTRIS, Rafin. Fl. Tellur. ii. 89 (1836).
angustifolia, *Miq. Fl. Ind. Bat. Suppl.* 617.—Sumatra.
corymbosa, *Thou. Orch. Iles Afr.* ii. 87, 88.—Madag.
disticha, Lindl. Fol. Orchid. Corymbis = corymbosa.
flava, *Hensl. Biol. Centr. Am. Bot.* iii. 297.—Mexic.
galipanensis, *Reichb. f. in Flora*, xlviii. (1865) 184.—Am. trop.
Thouarsii, Reichb. f. in Bot. Zeit. vii. (1849) 886 = corymbosa.
verstrifolia, *Reichb. f. in Flora*, xlviii. (1865) 184.—Ind. or.; Malaya.
Welwitschii, *Reichb. f. l. c.* 183.—Afr. occ.

CORYMBIUM, Linn. Coroll. Gen. 14 (1787). *COMPOSITÆ*, Benth. & Hook. f. ii. 324.
CONTARENA, Adans. Fam. ii. 120 (1763).
africanum, *Linn. Sp. Pl.* 958.—Afr. austr.
congestum, *E. Mey. ex Steud. Nom.* ed. II. i. 425.—Afr. austr.
cymosum, *E. Mey. ex Steud. l. c.*—Afr. austr.
filiforme, *Linn. f. Suppl.* 293.—Afr. austr.
glabrum, *Linn. Syst.* ed. XII. 582.—Afr. austr.
glabrum, Thunb. Prod. Pl. Cap. 170 = filiforme ?
graminifolium, Burm. f. Fl. Cap. Prod. 29 = filiforme.
graminifolium, Lam. Encyc. ii. 129 = filiforme, glabrum.
hirsutum, Eckl. ex DC. Prod. v. 89 = villosum.
hirsutum, Thunb. Fl. Cap. 729, 730 = seq.
hirtum, *Thunb. Prod. Pl. Cap.* 170.—Afr. austr.
latifolium, *Harv. in Harv. & Sond. Fl. Cap.* iii. 55.—Afr. austr.
listeum, *E. Mey. ex Steud. Nom.* ed. II. i. 425.—Afr. austr.
nervosum, *Thunb. Prod. Pl. Cap.* 170.—Afr. austr.
scabridum, Berg. Descr. Pl. Cap. 241 = africanum.
sinuatum, Lam. Mant. i. 130 = præne.
villosum, *Linn. f. Suppl.* 292.—Afr. austr.

CORYMBORCHIS, Thou. ex Blume, Orch. Ind. Arch. 125 (1855) = Corycrabis, Thou. (Orchid.).
angustica, Blume, Orchid. Archip. Ind. 126. t. 48 = Corymbis ventricifolia.
Thouarsii, Blume, l. c. 126. t. 44. f. 1. a–c = Corymbis corymbosa.
ventricifolia, Blume, l. c. 125. t. 43 = Corymbis ventricifolia.

CORYMBULA, Rafin. New Fl. Am. iv. 88 (1836) = Polygala, Linn.

CORYNABA, Hook. f. in Trans. Linn. Soc. xxii. (1856) 31. t. 18, 14. *BALANOPHOREÆ*, Benth. & Hook. f. iii. 237.
crassa, *Hook. f. l. c.* 31, 54. t. 18.—N. Granat.
Purdiei, *Hook. f. l. c.* 31, 55.—N. Granat.
sphærica, *Hook. f. l. c.* 31, 54. t. 14.—N. Granat.
Spruceï, *Eichl. in DC. Prod.* xvii. 187.—N. Granat.

CORYNANDRA, Schrad. ex Spreng. Syst. iv. Cur. Post. 201 (1827) = Cleome, Linn. (Capparr.).
pulchella, Schrad. ex Spreng. l. c. 204 = Cleome Chelidonii.

CORYNANTHE, Welw. in Trans. Linn. Soc. xxvii. (1869) 87. t. 14. *RUBIACEÆ*, Benth. & Hook. f. ii. 96.
paniculata, *Welw. l. c.*—Afr. trop.

CORYNANTHELIUM, Kunze, in Linnæa, xx. (1847) 19 = Mikania, Willd. (Compos.).
Moronoa, Kunze, l. c.—Bras.

CORYNANTHES, Schlecht. in Bot. Zeit. vi. (1848) 65 = Coryanthes, Hook. (Orchid.).

CORYNELLIA, Reichb. Nom. 148 (1841) = seq.

CORYNELLA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 93. *LEGUMINOSÆ*, Benth. & Hook. f. i. 500.
CORYNITIS, Spreng. Syst. iv. Cur. Post. 268 (1827).
TOXOTROPIS, Turcz. in Bull. Soc. Nat Mosc. xix. (1846) 326.
gracilis, *Grseb. Cat. Pl. Cub.* 71.—Cuba.
immarginata, *Wright, in Sauv. Fl. Cub.* 96.—Cuba.
pauciflora, *DC. in Ann. Sc. Nat. Sér.* I. iv. (1825) 93. —St. Domingo.
polyantha, *DC. l. c.*—Ind. occ.

CORYNELOBOS, R. Roem. ex Willk. in Linnæa, xxv. (1852) 7 = Brassica, Linn. (Cruciff.).
boeticus, R. Roem. ex Willk. l. c. = Brassica cornelobo.

CORYNEPHORUS, Beauv. Agrost. 90. t. 18. f. 2 (1812). *GRAMINEÆ*, Benth. & Hook. f. iii. 1187.
WEINGÆRTNERIA, Bernh. Syst. Verz. Erf. 28, 51 (1800), nomen prius.
aetnensis, Schult. Mant. iii. 641 = articulatus.
articulatus, *Beauv. Agrost.* 90.—Europ.; As. Min. canescens, *Beauv. l. c.* t. 18. f. 2.—Europ. austr.
fasciculatus, *Boiss. & Reut. Pugill. Pl. Nov.* 123.—Algieria.
macranthorus, Boiss. & Reut. l. c. 124 = fasciculatus.

CORYNITIS, Spreng. Syst. iv. Cur. Post. 268, 280 (1827) = Corynella, DC. (Legum.).
dominguensis, Spreng. l. c. 280 = Corynella paucifolia.
polyantha, Spreng. l. c. = Corynella polyantha.

CORYNOCARPUS, Forst. Char. Gen. 31. t. 16 (1776). *ANACARDIACEÆ*, Benth. & Hook. f. i. 423.
laevigata, *Forst. l. c.* 32.—N. Zel.

CORYNOPHALLUS, Schott, in Oestr. Bot. Wochenbl. (1857) 289 = Amorphophallus, Blume (Aroid.).
Afzelii, Schott, l. c. (= *Hydrosme leonensis*).—Afr. trop.
leonensis, Engl. in DC. Monog. Phan. ii. 396 = A. leonensis.

CORYNOPHORUS, Kunth, in Mém. Mus. Par. ii. (1815) 72 = Corynephorus, Beauv. (Gramin.)

CORYNOSICYOS, F. Muell. Sel. Pl. Indust. Cult. 59 (1876) = Cucumeropsis, Naud. (Cucurbit.).
edulis, F. Muell. l. c. = Cucum. edulis.

CORYNOSTIGMA, Presl, Epim. Bot. 218 (1849) = Jussiæna, Linn. (Onagrar.).
fasciculatum, Presl, l. c. = J. nervosa.

CORYNOSTYLIS, Mart. Nov. Gen. et Sp. i. 25. t. 17, 18 (1824). *VIOLARIEÆ*, Benth. & Hook. f. i. 116.
CALYPTERION, Gieg. in DC. Prod. i. 288 (1824).
albiflora, Linden, ex T. Moore, in Florist (1872) 9.—Bras.
Benthami, Walp. Rep. i. 223 = C. Hybanthus.
Berterii, *Sprong. Syst.* i. 806.—Mexic.
carthagenensis, Karst. Fl. Columb. iii. 58. t. 127. f. 9–14 = C. Hybanthus.
diandra, *Sprong. Syst.* i. 805 — Am. austr.
guayanensis, Karst. Fl. Columb. iii. 53. t. 127. f. 1–8 = C. Hybanthus.
Hybanthus, *Mart. & Zucc. Nov. Gen. et Sp.* i. 26. t. 17.—Mexic.
Loeflingii, *Sprong. Syst.* i. 805.—Am. austr.

CORYNOTHECA, F. Muell. Fragm. vii. 68. (1870). *LILIACEÆ*, Benth. & Hook. f. iii. 792.
scaubockeda, *F. Muell. l. c.*—Austral.
dichotoma, *F. Muell. l. c.*—Austral.
interiflora, *F. Muell. l. c.*—Austral.

CORYNULA, Hook. f. Ic. Pl. t. 1123 (1872). *RUBIACEÆ*, Benth. & Hook. f. iii. 138.
pilosa, *Hook. f. l. c.*—Am. austr.

CORYPHA, Linn. Mus. Cliff. 11 (1736); Gen. ed. I. 354 (1737). *PALMÆ*, Benth. & Hook. f. iii. 922.
BEDDA, Rafin. Sylva Tellur. 13 (1838).
CODDA-PANA, Adans. Fam. ii. 25 (1763).
DENDREMA, Rafin. Sylva Tellur. 14 (1838).
GEMBANGA, Blume, in Flora, viii. (1825) 580, 672.
TALIERA, Mart. Progr. Palm. Fam. et Gen. 10 (1824).
africana, Lour. Fl. Cochinch. 213 = Hyphæne coriacea.
australis, R. Br. Prod. 267 — Livistona australis.
cerifera, Arruda, in Koster, Trav. Braz. 494; Mart. Hist. Nat. Palm. ii. 56. t. 49 = Copernicia cerifera.
dulcis, H. B. & K. Nov. Gen. et Sp. i. 300 = Brahea dulcis.
elata, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 176.—Ind. or. Gebang, *Mart. Hist. Nat. Palm.* iii. 233.—Bras.
Gebanga, *Blume, Rumphia*, ii. 59. t. 97, 98, 105.—Malaya.
glaucescens, Lodd. ex Loud. Hort. Brit. 125 = Sabal glaucescens ?
guineensis, Linn. Mant. i. 187 = umbraculifera ?
Hystrix, Desf. Tabl. ed. I. 19 = Rhapidophyllum Hystrix.
Licuala, Lam. Encyc. ii. 181 = Licuala Rumphii.
macropoda, *Kurz, ex Linden, Cat.* n. 87 (1873).—Ins. Andam.
maritima, H. B. & K. Nov. Gen. et Sp. i. 298 = Copernicia maritima.
minor, *Blanco, Fl. Filip.* ed. I. 229.—Ins. Philipp.
minor, Jacq. Hort. Vindob. iii. 8 = Sabal Adansoni.
Miraguama, H. B. & K. Nov. Gen. et Sp. i. 296 = Thrinax Miraguama.
nana, H. B. & K. l. c. 299 = Copernicia nana.
palmacea, Sterd. Nom. ed. I. 225, 334 = Thrinax parviflora.
Palmetto, Walt. Fl. Carol. 119 = Sabal Palmetto.
pilearia, Lour. Fl. Cochinch. 213 = Licuala pilearia.
pumila, Walt. Fl. Carol. 119 = Sabal Adansoni.
Pumos, H. B. & K. Nov. Gen. et Sp. i. 298 = Copernicia Pumos.
repens, Bertero, Trav. 61 = Rhapidophyllum Hystrix.
rotundifolia, Lam. Encyc. ii. 181 = Livistona rotundifolia.
Saribus, Lour. Fl. Cochinch. 212 = Livistona cochinchinensis.
sylvestris, *Mart. Hist. Nat. Palm.* iii. 233.—Ins. Molucc.
Taliera, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 174.—Ind. or.
tectorum, H. B. & K. Nov. Gen. et Sp. i. 299 = Copernicia tectorum.
thebaica, Linn. Sp. Pl. 1187 = Hyphæne crudeta.
umbraculifera, Jacq. Fragm. Bot. 13 = Sabal Blackburniana.
umbraculifera, *Linn. Sp. Pl.* 1178.—Afr. trop.; Ind. or.
Utan, Lam. Encyc. ii. 181 = sylvestris.

CORYPHANTHA, Lem. Cact. 32 (1868) = Mammillaria, Haw. (Cact.).
acanthostephes, Lem. l. c. 55 = M. acanthostepha.
ancistracanthia, Lem. l. c. 34 = M. raphidacantha.

CORYPHANTHA :—
sulacothele, Lem. Cact. 34 = Mammillaria sulacothele.
brevimamma, Lem. ex Foerst. Handb. Cact. ed. II. 394 = M. brevimamma.
calcarata, Lem. Cact. 35 = M. calcarata.
Clava, Lem. l. c. 34 = M. Clava.
compacta, Lem. l. c.—Hab.!
cornifera, Lem. l. c. 35 = M. cornifera.
daimonoceras, Lem. l. c. = M. scolymoides.
elephantidens, Lem. l. c. = M. elephantidens.
Engelmanni, Lem. l. c. 34.—Hab.!
erecta, Lem. l. c. = M. erecta.
exsudans, Lem. ex Foerst. Handb. Cact. ed. II. 395 = M. exsudans.
glandulifera, Lem. Cact. 34 = M. glandulifera!
heterophylla, Lem. l. c.—Hab.!
Hookeri, Lem. l. c.—Hab.!
impexicoma, Lem. ex Foerst. Handb. Cact. ed. II. 414 = M. cornifera.
Lehmanni, Lem. Cact. 34 = M. Lehmanni.
loricata, Lem. l. c. 35 = M. loricata.
macromeris, Lem. l. c. = M. macromeris.
Nuttalii, Engelm. ex Foerst. Handb. Cact. ed. II. 407 = M. Nuttalii.
Ottonis, Lem. Cact. 34 = M. Ottonis.
pycnacantha, Lem. l. c. 35 = M. pycnacantha.
raphidacantha, Lem. l. c. 34 = M. raphidacantha.
Scheeri, Lem. l. c. 35 = M. Scheeri.
Schiechtendalii, Lem. l. c. 34 = M. Schiechtendalii.
sublanata, Lem. l. c. 35.—Hab.!
sulcolanata, Lem. l. c. = M. sulcolanata.

CORYSADENIA, Griff. in Proc. Linn. Soc. i. (1846) 281 = **Illigera**, Blume (Combret.).

CORYSANTHERA, Wall. Cat. n. 6411 (1832) = Rhynchotechoum, Blume (Gesnerac.).
elliptica, Wall. l. c. = R. elliptieum.
vestita, Griff. Itin. Notes, 124 = R. vestitum.

CORYSANTHES, R. Br. Prod. 328 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 509.
CALCEARIA, Blume, Bijdr. 417. t. 83 (1825).
CORYBAS, Salisb. Parad. Lond. t. 83 (1806), nomen prius.
NEMATOSCERAS, Hook. f. Fl. N. Zel. i. 249. t. 57 (1853).
Betchel, *R. Mustl. in Wing, South Sc. Record*, i. (1881) 171.—Ins. Sandwich.
bicalcarata. *R. Br. Prod.* 328.—Austral.
calloza, *Blume, Orch. Archip. Ind.* 128. t. 68.—Malaya.
Cheesemanii, *Hook. f. ex T. Kirk, in Trans. N. Z. Inst.* iii. (1871) 180.—N. Zel.
dienemica, Lindl. Gen. et Sp. Orch. 398 = fimbriata.
fimbriata, *R. Br. Prod.* 328.—Austral.
fornicata, *Lindl. Gen. et Sp. Orch.* 394.—Malaya.
hypogea, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 336.—N. Zel.
limbata, *Hook. f. Bot. Mag.* t. 5357.—Java.
macrantha, *Hook. f. Handb. N. Zeal. Fl.* 266.—N. Zel.
maculata, Heynh. Nom. i. 232 = Coryanthes maculata.
macrosata, *Blume, Orch. Archip. Ind.* 175.—Malaya.
oblonga, *Hook f. Handb. N. Zeal. Fl.* 266.—N. Zel.
papillosa, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 337.—N. Zel.
picta, *Lindl. Gen. et Sp. Orch.* 394.—Malaya.
prudnosa, R. Cunn. in N. S. Wales Mag. (1833) n. 1 = fimbriata.
rivularis, *Hook. f. Handb. N. Zeal. Fl.* 266.—N. Zel.
rotundifolia, *Hook. f. l. c.*—N. Zel.
triloba, *Hook. f. l. c.* 265.—N. Zel.
undulata, *R. Cunn. in N. S. Wales Mag.* (1833), nomen.—Austral.
unguiculata, *R. Br. Prod.* 328.—Austral.

CORYTHACANTHUS, *Nees* in Lindl. Introd. Nat. Syst. ed II. 444 (1836) = Chiatax, Mart. (Acanthac.).
speciosus, Nees, l. c.—Bras.

CORYTHANTHES, Lem. in Orb. Dict. iv. 259 (1849) = Coryanthes, Hook. (Orchid.).

CORYTHOLOBIUM, Mart. ex Benth. in Ann. Wien. Mus. ii (1838) 93 = Securidaca, Linn. (Polygal.).
macrophyllum, Benth. l. c. = S. Corytholobium.

CORYTHOLOMA, Decne. in Rev. Hortic. Sér. III. ii. (1848) 486 = Gesneria, Mart.
aggregatum, Decne. l. c. 487 = G. aggregata.
carcaeaganum, Decne. l. c. = G. caracasana.
chelonoides, Regel, Gartenfl. vii. (1858) 374 = G. chelonoides.
elliptieum, Decne. in Rev. Hortic. Sér. III. ii. (1848) 486 = G. elliptica.
fragile, Decne. l. c. 487 = G. Sceptrum.
gracile, Bronge. ex Decne. l. c. 486.—Hab.!
ignneum, Decne. l. c. 487 = G. Sceptrum.
Lindleyi, Decne. l. c. = G. Lindleyi.
Marchii, Decne. l. c. = G. Marchii.
melittifolium, Brongn. ex Decne. l. c. = G. melittifolia.
Morchii, Decne. ex Regel, Gartenfl. iv. (1855) 247 = G. Merckii.
pendulinum, Decne. in Rev. Hortic. Sér. III. ii. (1848) 487 = G. pendulina.
rupicolum, Decne. l. c. = G. rupicola.
rutilum, Decne. l. c. = G. rutila.
Sceptrum, Decne. l. c. = G. Sceptrum.
strictum, Decne. l. c. 486 = G. stricta.
tubiflorum, Decne. l. c. 487 = Isolema tubiflorum.

CORYTOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 119 = Alloplectus, Mart. (Gesneruc.).
capitatus, Oerst. l. c. 120 = A. capitatus!

CORYZADENIA, Griff. Notul. iv. 256 (1854) = Illigera, Blume (Combretac.).
trifoliata, Griff. l. c. = I. Coryzadenia.

COSARIA, J. F. Gmel. Syst. 29, 71 (1791) = KOSARIA, Forsk. = Dioscorea, Linn. (Uticac.).
Forskålii, J. F. Gmel. l. c. = D. radiata.

COSBAEA, Lem. Illustr. Hortic. ii. (1855) Misc. 71 = Schisandra, Michx. (Magnoliac.)
coccinea, Lem. l. c. = S. Hanosana.

COSCINIUM, Colebr. in Trans. Linn. Soc. xiii. (1822) 51. *MENISPERMACEAE*, Benth. & Hook. f. i. 85.
PEREIRIA, Lindl. Fl. Med. 370 (1838).
Blumeanum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. vii. (1851) 37.—Ind. or.
fenestratum, *Colebr. in Trans. Linn. Soc.* xiii. (1822) 55.—Ind. or.
Maingayi, *Pierre, Fl. For. Cochin.* sub t. 112 (err. Mangay).—Malaya.
usitatum, *Pierre, l. c.*—Cambodia.
Wallichianum, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 37 = fenestratum.
Wightianum, Miers, l. c. = fenestratum.

COSMANTHUS, Nolte, ex A. DC. Prod. ix. 296 (1838) = Phacelia, Juss. (Hydrophyll.).
fimbriatus, Nolte, ex A. DC. l. c. 297 = P. fimbriata.
grandiflorus, A. DC. l. c. = P. grandiflora.
mexicanus, A. DC. l. c. = P. pimpinelloides.
nemophiloides, Kunth, Ind. Sem. Hort. Berol. (1846) 12.—Texas.
parviflorus, A. DC. in DC. Prod. ix. 297 = P. parviflora.
pectinatus, E. Mey. in Ann. Sc. Nat. Sér. III. v. (1846) 342.—Am. bor.
viscidus, A. DC. in DC. Prod. ix. 296 = P. grandiflora.

COSMARIUM, Dulac, Fl. Hautes-Pyr. 215 (1867) [non Corda, quod Algar. gen.] = Adonis, Linn. (Ranunc.).
aestivale, Dulac, l. c. = A. aestivalis.
autumnale, Dulac, l. c. = A. autumnalis.
pyrenaicum, Dulac, l. c. = A. pyrenaica.

COSMEA, Willd. Sp. Pl. iii. 2350 (1803) = Cosmos, Cav. (Compos.).
bipinnata, Willd. l. c. = Cosmos bipinnatus.
candata, Spreng. Syst. iii. 615 = Cosmos candatus.
chrysanthemifolia, Spreng. l. c. = Cosmos chrysanthemifolius.
crithmifolia, Spreng. l. c. = Cosmos crithmifolius.
diversifolia, Heynh. Nom. ii. 163 = Cosmos diversifolius.
intea, Sims, Bot. Mag. t. 1689 = Bidens grandiflora.

COSMEA :—
parviflora, Willd. Sp. Pl. iii. 2350 = Cosmos parviflorus.
pilosa, Spreng. Syst. iii. 615 = Cosmos pilosus.
scabiosoides, Spreng. l. c. = Cosmos scabiosoides.
sulphurea, Willd. Sp. Pl. iii. 2350 = Cosmos sulphureus.
tenella, Spreng. Syst. iii. 615 = Cosmos tenellus.

COSMELIA, R. Br. Prod. 558 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 617.
angustifolia, DC. Prod. vii. 766 = rubra.
rubra, *R. Br. Prod.* 558.—Austral.

COSMIA, Domb. ex Juss. Gen. 212 (1789) = Calandrinia, H. B. & K. (Portulac.).
caulescens, Domb. ex Spreng. Syst. ii. 458 = Calandrinia caulescens.
montana, Domb. ex DC. Prod. iii. 359 = Calandrinia caulescens.
prostrata, Domb. l. c. = Calandrinia acaulis.

COSMIBUENA, Ruiz & Pav. Fl. Per. iii. 8. t. 226 (1802). *RUBIACEAE*, Benth. & Hook. f. ii. 40.
BUENA, Pohl, Pl. Bras. Ic. i. 8, partim (1827).
acuminata, *Ruiz & Pav. Fl. Per.* iii. 8. t. 226.—Peruv.
dichotoma, G. Don, Gen. Syst. iii. 479 = Ladenbergia dichotoma.
latifolia, *Klotzsch, ex Walp. Rep.* vi. 69.—N. Granat.
macrocarpa, *Klotzsch, ex Walp. l. c.*—Ins. Gorgona.
obtusifolia, *Ruiz & Pav. Fl. Per.* iii. 8.—Peruv.
ochracea, *Endl. Icongr.* t. 90.—Hab.!
quinquenervia, *Klotzsch, in Hayne, Arzneig.* xiv. sub t. 15.—Venezuela.
Skinneri, *Hemsl. Biol. Centr. Am. Bo* ii. 12.—Nicaragua.
trifiora, *Klotzsch, in Hayne, Arzneig.* xiv. sub t. 15.—Guiana.

COSMIBUENA, Ruiz & Pav. Prod. Fl. Per. 10. t. 2 (1794) = Hirtella, Linn. (Rosac.).

COSMIDIUM, Nutt. in Trans. Am. Phil. Soc. Ser. II vii. (1841) 361 = Thalesperma, Less. (Compos.).
Burridgeanum ×, Hort. ex Regel, Ind. Sem. Hort Petrop. (1857) 40 = T. filifolium.
filifolium, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 362 = T. filifolium.
gracile, Nutt. l. c. = T. gracile.
simplicifolium, A. Gray, Fl. Fendl. 86 = T. subsimplicifolium.

COSMIUSA, Alef. in Bot. Zeit. xxiv. (1866) 146 = Parochaetus, Buch.-Ham. (Legumin.).
repens, Alef. l. c. = P. communis!

COSMIZA, Rafin. Fl. Tellur. iv. 110 (1836) = Polypompholyx, Lehm. (Lentibul.).
coccinea, Rafin. l. c. = P. multifida.

COSMOPHYLLUM, C. Koch, Ind. Sem. Hort. Berol (1854) 12 = Podachaenium, Benth. (Compos.).
cacaliaefolium, C. Koch, l. c.—Guatemala.

COSMOS, Cav. Ic. i. 9. tt. 14, 79 (1791). *CONPOSITAE*, Benth. & Hook. f. ii. 287.
ADENOLEPIS, Less. in Linnaea. vi. (1831) 510.
COSMEA, Willd. Sp. Pl. iii. 2350 (1803).
COSMUS, Pers. Syn. ii. 477 (1807).
bipinnatus, *Cav. l. i. 10. t. 14.*—Am. bor. ; Mexic.
cervicolus, *Benth. Bot. Voy. Sulph.* 117.—Mexic.
candatus, *H. B. & K. Nov. Gen. et Sp.* v. 240.—Ind. occ.
chrysanthemifolius, *H. B. & K. l. c.* 239. t. 462.—Mexic.
crithmifolius, *H. B. & K. l. c.* 242.—Mexic.
diversifolius, *Otto, in Knowles & Westc. Flor. Cab.* ii. 3. t. 97.—Am. bor.
integrifolius, *Weald. Chlor. Amd.* t. 70, bota.—Bolivia.
linearifolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 200.—Mexic.
marginatus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 253.—Peruv.
parviflorus, *Pers. Syn.* ii. 477 ; *H. B. & K. Nov. Gen. et Sp.* iv. 241.—N. Mexic.
pseudosulphureus, *Weald. Chlor. Amd.* t. 70.—Bolivia.
pilosus, *H. B. & K. Nov. Gen. et Sp.* iv. 243.—Mexic.
pinnatus, Jacq. ex Steud. Nom. ed. II. i. 495 = Conopsis coronata.

COSMOS :—
pulcherrimus, Sch. Bip. in Ball. Soc. Bot. Fr. xii. (1865) 79 = diversifolius.
purpureus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 200.—Mexic.
reptans, Benth. Pl. Hartw. 40 = diversifolius.
rosens, *Mart. ex Steud. Nom.* ed. II. i. 426.—Bras.
scabiosoides, *H. B. & K. Nov. Gen. et Sp.* iv. 242.—Mexic.
subpubescens, *Wedd. Chlor. And.* i. 70.—Peruv.
sulphureus, *Cav. Ic.* i. 58. t. 79.—Mexic.
tenellus, *H. B. & K. Nov. Gen. et Sp.* iv. 240.—Mexic.
tenuifolius, *Lindl. Bot. Reg.* t. 2007.—Mexic.
Uhdeanus, *Kunth, Ind. Sem. Hort. Berol.* (1846) 12.—Mexic.

COSMOSTIGMA, Wight, Contrib. Bot. Ind. 41 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 774.
acuminata, Wight, Ic. t. 1270 = pinnata.
racemosa, *Wight, Contrib. Bot. Ind.* 43.—Ind. or.

COSSIGNEA, Willd. Sp. Pl. ii. 226 (1799) = seq.

COSSIGNIA, Comm. ex Juss. Gen. 248 (1789) = seq.

COSSIGNYA, Baker, Fl. Maurit. 53 (1877) = seq.

COSSINIA, Comm. ex Lam. Encyc. ii. 182 (1786). *SAPINDACEAE*, Benth. & Hook. f. i. 397.
MELICOPSIDIUM, Baill. Adansonia, xi. (1874) 248.
borbonica, DC. Prod. i. 614 = pinnata.
Commersonii, Raeusch. Nom. ed. III. 108 = pinnata.
madagascariensis, *Baill. Adansonia*, xi. (1874) 247.—Madag.
pinnata, *Comm. ex Lam. Encyc.* ii. 182. t. 256.—Ins. Maurit.
splendens, Sieber, ex Steud. Nom. ed. II. i. 426 = pinnata.
ternata, Baker, Fl. Maurit. 59 = pinnata.
trifoliata, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 372.—N. Zeal.
triphylla, Comm. ex Lam. Encyc. ii. 182 = pinnata.

COSSONIA, Dur. in Ann. Sc. Nat. Sér. III. xx. (1853) 82. t. 6 = **Raffenaldia**, Godr. (Crucifer.).
africana, Dur. l. c. = R. primuloides.

COSTA, Vell. Fl. Flum. 19 (1825); t. t. 48 (1827) = **Galipea**, Aubl. (Rutac.).
aromatica, Vell. l. c. 20; t. t. 48 = Galipea Arrabidae.

COSTAEA, A. Rich. Fl. Cub. Fanerog. ii. 75 (1858) = **Purdiaea**, Planch. (Cyrill.).
cubensis, A. Rich. l. c. 78 = P. monodynama.

COSTEA, A. Rich. Fl. Cub. Fanerog. ii. 328 (1858) = prec.

COSTIA, Willk. in Bot. Zeit. xvi. (1858) 377 = **Agropyron**, J. Gaertn. (Gramin.).
cristata, Willk. l. c. = A. cristatum.
imbricata, Willk. l. c. = A. imbricatum.
orientalis, Willk. l. c. = A. orientale.
squarrosa, Willk. l. c. = A. patulum.

COSTIA, Willk. in Bot. Zeit. xviii. (1860) 181 = **Iris**, Tourn.
caucasica, Willk. l. c. = I. caucasica.
persica, Willk. l. c. = I. persica.
scorpioides, Willk. l. c. = I. alata.

COSTUS, Linn. Mus. Cliff. 9 (1736); Gen. ed I. 331 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 646.
ACINAX, Rafin. Fl. Tellur. iv. 53 (1838).
BANKSIA, Koen. in Retz. Obs. iii. 75 (1788).
GISSANTHUS, Salisb. in Trans. Hort. Soc. i. (1812) 279.
HELLENIA, Retz. Obs. vi. 18 (1791).
TRIANA, J. F. Gmel. Syst. 9 (1791).
JAGUANGA, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 262, 343.
PLANERA, Gaertn., Fasel. 205 (1792).
abe, Ker-Gawl. in Bot. Reg. t. 633.—*Afr.* trop.
Anachiri, Jacq. Fragm. 55 = spicatus.
angustifolius, Ker-Gawl. in Bot. Reg. t. 965 = speciosus.
appendiculatus, *Humb. & Bonpl. ex Rorm. & Schult. Mant.* i. 32.—Am. austr.
arabicus, Asbl. Fl. Gui. i. 2 = spicatus.
arabicus, Linn. Sp. Pl. 2 = speciosus.
arabicus, Rosc. ex Spreng. Syst. i. 13 = glabratus.
arabicus, Vell. Fl. Flum. 2; i. t. 5 = Arrabidae.
argenteus, *Ruiz & Pav. Fl. Per.* i. 3. t. 4.—Peruv.
argyrophyllus, Wall. Cat. n. 6535 G, H = speciosus.
Arrabidae, *Steud. Nom.* ed. II. i. 426.—Bras.
cernuus, *Sw. ex Rorm. & Schult. Syst.* i. 25.—Am. trop.
ciliatus, *Humb. & Bonpl. ex Rorm. & Schult Mant.* i. 32.—Am. trop.
ciliatus, *Mog. in Linnaea*, xviii. (1844) 78.—Guian.
cosmosus, *Rosc. in Trans. Linn. Soc.* viii. (1807) 350.—Am. austr.
conicus, Stokes, Bot. Mat. Med. i. 75 = spicatus.
cylindricus, Stokes, l. c. = speciosus.
cylindricus, Jacq. Fragm. 54 = spiralis.
cylindricus, *Rosc. Scitam.* t. 78.—Ins. Trinit.
discolor, *Rosc. l. c.* t. 81.—Bras.
echinatus, Pers. Syn. i. 3 = Amomum echinatum.
foeniculaceus, *Noronha, in Verh. Batav. Gen.* v (1790) ed. I. Art. IV. 12.—Malaya.
glabratus, Reichb. in Moesel. Handb. ed. II. i. 8 = speciosus.
glabratus, *Sw. Prod. Veg. Ind. Occ.* 11.—Ind. occ.
globosus, *Blume, Enum. Pl. Jav.* 62.—Java.
hirsutus, *Presl, Rel. Haenk.* i. 112.—Mexic.
igneus, *N. E. Br. in Illustr. Hortic.* xxxi. (1884) 25. t. 611.—Bras.
laevis, *Ruiz & Pav. Fl. Per.* i. 3.—Peruv.
linearis, Spreng. Syst. i. 13 = Kaempferia secunda.
Loureiri, *Horan. Prod. Scitam.* 38.—Sumatra.
Malmei, Blanco, Fl. Filip. ed. 2. 4 = Roscoea lutea.
maculatus, *Rosc. Scitam.* t. 82.—Afr. trop.
malacensis, Retz. Obs. iii. 71 = Alpinia malaccensis.
Malortieanus, *H. Wendl. in Hamb. Gartenz.* xix. (1863) 30.—Am. centr.
nepalensis, Rosc. Scitam. t. 80 = speciosus.
nigricans, Blanco, Fl. Filip. ed. II. 3 = Roscoea nigrolabra.
nivosus, *Jacq. Fragm.* 55.—Ins. Martinic.
niveus, *G. F. W. Mey. Prim. Fl. Essq.* 1.—Guian.
pictus, *D. Don, ex Lindl. Bot. Reg.* t. 1094.—Mexic.
Pisonis, Lindl. l. c. t. 899 = spiralis.
Potierae, *F. Muell. Fragm.* iv. 164.—Austral.
pulverulentus, *Presl, Rel. Haenk.* i. 111.—Mexic.
pungens, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 244 = Tapeinochelos pungens
quartus, Roem. & Schult. Syst. i. 26 = cylindricus.
quintus, Roem. & Schult. l. c. = spicatus.
ruber, *Wright, ex Griseb. Cat. Pl. Cub.* 256.—Cuba.
sarmentosus, *Bej. in Ann. Sc. Nat. Sér.* II. iv. (1835) 262. t. 8.—Ins. Maurit.
scaber, *Ruiz & Pav. Fl. Per.* i. 2. t. 3.—Peruv.
secundus, Roem. & Schult. Syst. i. 22 = Renealmia Prasco-Secora.
secundus, Spreng. Syst. i. 13 = Kaempféria secunda.
septimus, Roem. & Schult. Syst. i. 26 = villosissimus.
setiger, Blume, Enum. Pl. Jav. 62 = speciosus.
sextus, Roem. & Schult. Syst. i. 26 = niveo-purpureus.
speciosus, *Sm. in Trans. Linn. Soc.* i. (1791) 249.—Ind. or.; Malaya.
spicatus, *Sw. Prod. Veg. Ind. Occ.* 11.—Ind. occ.
spiralis, *in Trans. Linn. Soc.* viii. (1807) 350.—Venezuela.
vaginatus, Salisb. in Trans. Hort. Soc. i. (1812) 277 = speciosus.
Verschaffeltianus, *Lem. in Ygard. Fleur.* iv. (1854) t. 381.—Bras.
villosissimus, *Jacq. Fragm.* 55.—Ins. S. Vincent.
Zerumbet, Pers. Syn. i. 3 = Alpinia zutans.

COTA, J. Gay, in Guss. Fl. Sic. Syn. ii. 866 (1844) = **Anthemis**, Mich. (Compos.).
altissima, J. Gay, l. c. 867 = A. Cota.
austriaca, Sch. Bip. in Oestr. Bot. Wochenbl. iv. (1854) 185 = A. austriaca.
Bourgaei, Boiss. Fl. Orient. iii. 284 = A. coelopoda.
brachycentros, J. Gay, in Guss. Fl. Sic. Syn. ii. 867 = A. brachycentros.
Brachmanni, Boiss. Diagn. Ser. II. iii. 21 = A. Brachmanni.
caelopoda, Boiss. l. c. = A. coelopoda.
Commersonii, Willd. & Lange, Prod. Fl. Hisp. ii. 35 = A. austriaca.
COTA :—
Iyonnetioides, Boiss. & Kotschy, Diagn Ser. II. v 109 = A. lyonnetioides.
ormenioides, Boiss. l. c. iii 21 = A Wiedemanniana.
oxylepis, Boiss l. c. v. 109 = A. oxylepis
palaestina, Rent. ex Unger & Kotschy, Ins. Cyp. 240 = A. palestina!
parnassica, Boiss. & Heldr. Diagn. Ser. II iii. 20 = A tinctoria.
Pestalozzae, Boiss. l. c. 21 = A. Pestalozzae.
Thirkei, Sch. Bip. in Oestr. Bot. Wochenbl. iv. (1854) 183 = A. brachycentros.
tinctoria, J. Gay, in Guss. Fl. Sic. Syn. ii. 867 = A tinctoria.
Trivmfetti, J. Gay, l. c. = A. austriaca.

COTINUS, [Tourn.] Linn. Syst. ed. I (1735) = **Rhus**, Linn. (Anacard.).
americanus, Nutt. Sylva, ii. t. 81 = R. cotinoides
coggygria, Scop. Fl. Carn. ed. II. i. 220 = R. Cotinus.
Coriaria, Duham. Arb. ed. I. i. 191 = R. Cotinus.

COTONEASTER, Rupp. Fl. Jen. ed. Hall. 137 (1745); Medik. Phil. Bot. i. 155 (1789). *ROSACEAE*. Benth. & Hook. f. i. 627.
GYMNOPYRENIUM, Dulac, Fl. Hautes-Pyr. 316 (1867).
NARGELIA, Lindl. Veg. Kingd. 560 (1847).
acuminata, *Lindl. in Trans. Linn. Soc.* xiii. (1822) 101.—Reg. Himal.
acutifolia, Maxim. Prim. Fl. Amur. 171 = integerrima.
acutifolia, *Turcz. in Bull. Soc. Nat. Mosc.* iv. (1832) 190.—Mongolia.
aestivalis, Wenzig, in Linnaea, xxxviii. (1874) 202 = Crataegus aestivalis.
affinis, DC. Prod. ii. 632 = buxifolia.
affinis, Hohen. ex Hook. f. Fl. Brit. Ind. ii. 385 = frigida!
affinis, Lindl. in Trans. Linn. Soc. xiii. (1822) 101 = Stranvaesia glaucescens.
arborescens, Wenzig, in Linnaea, xxxviii. (1874) 208 = Crataegus arborescens.
bacillaris, *Wall. ex Lindl. Bot. Reg.* sub t. 1229.—Reg Himal.
buxifolia, Baker, in Saund. Refug. Bot. i. t. 52 = microphylla.
buxifolia, *Wall. Cat.* n. 661; *et Lindl. Bot. Reg.* sub t. 1229.—Reg. Himal.
californica, *A. Murr. Oreg. Circ.* 1, nomen.—Calif
coccinea, Steud. Nom. ed. II. i. 426 = tomentosa.
comptus, *Lem. in Fl. des Serres*, Sér. I. iv. (1848) 388 b.—Mexic.
congesta, Baker, in Saund. Refug. Bot. i. t. 51 = microphylla.
crenulata, C. Koch, Dendrol. i. 175 = Crataegus crenulata.
Cuila, Lee, ex C. Koch, l. c. 172 = nummularia.
denticulata, *H. B. & K. Nov. Gen. et Sp.* vi. 214. t. 556. —Mexic.
distichea, *Lange, in Bot. Tidsk.* xiii. (1882) 19.—Nepal!
ellipsoidea, Hort. ex Loud. Encyc. Pl. 1208 = nummularia.
emarginata, Hort. ex C. Koch, Dendrol. i. 178 = microphylla.
Fontanesii, *Spach, Hist. Vég. Phan.* ii. 77.—Hab.!
Fortunei, Wenzig, in Linnaea, xxxviii. (1874) 200 = Photinia Fortunea.
frigida, *Wall. Cat.* n. 657; *et Lindl. Bot. Reg.* t. 1229.—Reg. Himal.
Grammontii, Hort. ex C. Koch, Dendrol. i. 172 = nummularia.
granatensis, *Boiss. Elench.* 41.—Hispan.
horizontalis, *Decne. in Fl. des Serres*, Sér. II. xii. (1877) 168.—China.
hymalaica, *Carr. in Rev. Hortic.* (1878) 180.—Reg. Himal.
integerrima, *Medic. Geech.* 85.—Europ.; As. temp.
intermedia, Medic. ex Trautv. in Act. Hort. Petrop. viii. (1883) 387, sphalm. = prsec.
laevis, Hort. ex Steud. Nom. ed. II. i. 426 = nummularia.
lanata, Hort. ex Ragel, Gartenfl. ix. (1860) 59 = buxifolia.
lanifera, Jacq. ex Lindl. Bot. Reg. t. 1205 = integerrima.
Lindleyi, Steud. Nom. ed. II. i. 426 = nummularia.

COTONEASTER:—
lucida, Schlecht. in Linnaea, xxvii. (1856) 641 = integerrima.
marginata, Hort. ex C. Koch, Dendrol. i. 177 = buxifolia.
melanocarpa, Fisch. ex Loud. Hort. Brit. Suppl. ii. 624 = integerrima.
microphylla, *Wall. ex Lindl. Bot. Reg.* t. 1114; *Cat.* n. 662 A.—Reg. Himal.
microphylla, Wall. Cat. n. 662 B = thymifolia.
multiflora, *Bunge, in Ledeb. Fl. Alt.* ii. 290.—Sibir. alt.
multiflora, C. A. Mey. Vers. Pfl. Casc. 171 = nummularia.
nebrodensis, *Nym. Syll.* 367.—Sicil.
nepalensis, Hort. ex C. Koch, Dendrol. i. 177 = frigida.
nervosa, *Decne. in Nouv. Arch. Mus. Par.* Ser. I. x. (1874) 177.—Hab.!
nigra, Wahlenb. Fl. Gothob. 58 = integerrima.
nummularia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 34.—Afr. bor.; Orient.
obtusa, Wall. Cat. n. 659; et Lindl. Bot. Reg. sub t. 1229 = bacillaris.
orientalis, *Kerner, in Oestr. Bot. Zeitschr.* xix. (1869) 270.—Europ.
parnassica, *Boiss. & Heldr. Diagn.* Ser. II. ii. 48.—Graecia.
peduncularis, Boiss. l. c. Ser. I. iii. 8 = integerrima.
polonica, *Jastrz. ex Rostaf. Fl. Polon. Prod.* 121.—Polonia.
prostrata, Baker, in Saund. Refug. Bot. i. t. 53 = rotundifolia.
Pyracantha, Spach, Hist. Vég. Phan. ii. 78 = Crataegus Pyracantha.
racemiflora, C. Koch, Dendrol. i. 170 = nummularia.
reflexa, *Carr. in Rev. Hortic.* (1870-71) 230.—China.
rosea, Edgew. in Trans. Linn. Soc. xx. (1851) 46 = bacillaris.
rotundifolia, *Wall. Cat.* n. 668; *et Lindl. Bot. Reg.* sub t. 1229.—Reg. Himal.
Royleana, Hort. ex C. Koch, Dendrol. i. 171 = acuminata.
Simonsii, *Hort. ex Baker, in Saund. Refug. Bot.* i. t. 55.—Reg. Himal.
spathulata, *Wenzig, in Linnaea*, xxxviii. (1874) 201.—Am. bor.
spicata, Hort. ex C. Koch, Dendrol. i. 166 = tomentosa.
thymaefolia, *Hort. ex Baker, in Saund. Refug. Bot.* i. t. 50.—Reg. Himal.
thymifolia, Hort. ex Loud. Encyc. Pl. 1384 = microphylla.
tomentosa, *Lindl. in Trans. Linn. Soc.* xiii. (1822) 101.—Europ.
tomentosa, C. A. Mey. Vers. Pfl. Casc. 171 = nummularia.
uniflora, *Bunge, in Ledeb. Fl. Alt.* ii. 290.—Sibir. alt.
Uva-ursi, G. Don, in Loud. Hort. Brit. 480 = rotundifolia.
vulgaris, Lindl. in Trans. Linn. Soc. xiii. (1822) 101 = integerrima.

COTTAEA, Endl. Gen. 92 (1841) = seq.

COTTEA, Kunth, Rév. Gram. i. 84, 281. t. 52 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1174.
pappophoroides, *Kunth, l. c.*—Peruv.
tarmentosa, *Nees, ex Steud. Syn. Pl. Gram.* 201.—Ins. S. Vinc.

COTTENDORFIA, Schult. f. Syst. vii. 64, 1193 (1830). *BROMELIACEAE* [Benth. & Hook. f. iii. 667].
albicans, *Griseb. in Gott. Abh.* xxiv. (1879) 230.—Reg. Argent.
floribs, *Schult. f. Syst.* vii. 11. 1193.—Bras.
guianensis, *Klotzsch, ex Beer, Bromel.* 44.—Guianae.

COTTONIA, Wight, Ic. v. t. 21. t. 1755 (1852). *ORCHIDEAE*, Benth. & Hook. f. iii. 573.
Championii, *Lindl. ex Benth. in Hook. Kew Journ.* vii. (1855) 35.—Hong Kong.
macrostachya, *Wight, Ic.* t. 1755.—Ind or.
pedunculata, Thw. Enum. Pl. Zeyl. 308 = macrostachya.

COTULA, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 438.
BALDISORIA, Neck. Elem. i. 281 (1790).
BROCCHIA, Vis. Pl. Aeg. et Nub. 35 (1856).
CENOCLINE, C. Koch, in Bot. Zeit. i. (1843) 41.
CHLAMYDOPHORA, Ehrenb. ex Less. Syn. Comp. 265, 448 (1830).
COTULINA, Pomel, Nouv. Mat. Fl. Atl. 56 (1874).
CTENOSPERMA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 116.
GYMNOGYNE, Steetz, in Lehm. Pl. Preiss. i. 431 (1845).
LANCISIA, Adans. Fam. ii. 126 (1763).
LEPTINELLA, Cass. in Bull. Soc. Philom. (1822) 127.
LEPTOGYNE, Less. in Linnaea, vi. (1831) 151.
MACHLIS, DC. Prod. vi. 140 (1837).
OTOGLYPHIS, Pomel, Nouv. Mat. Fl. Atl. 56 (1874).
PLEIOGYNE, C. Koch, in Bot. Zeit. i. (1843) 40.
STRONGYLOSPERMA, Less. Syn. Comp. 261 (1832).
SYMPHYOMERA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 116.
abrotanifolia, Willd. Sp. Pl. iii. 2167 = Anacyclus radiatus.
abyssinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 419.—Afr. trop.
africana, Steud. Nom. ed. I. 230 = Matricaria capensis.
alba, Linn. Syst. ed. XII. ii. 564 = Eclipta erecta.
alpina, *Hook. f. Fl. Tasm.* i. 192. t. 51 A.—Austral. anthemoides, *Linn. Sp. Pl.* 891.—Hisp.; Afr. bor. et austr.; Ind. or.
anthemoides, Willd. Sp. Pl. iii. 2165 = microcephala.
atrata, *Hook. f. Handb. N. Zeal. Fl.* 142.—N. Zel.
aurea, *Ser.* 168.—Reg. Mediterr.; Orient.
aurea, Sibth. & Sm. Fl. Graec. t. 61 = Matricaria decipiens.
australis, *Hook. f. Fl. Nov. Zel.* 128.—Austral.; N. Zel.
barbata, DC. Prod. vi. 79 = Cenia barbata.
bicolor, Roth, Catalect. fasc. ii. 116 = Dichrocephala latifolia.
bipinnata, *Thunb. Prod. Pl. Cap.* 162.—Afr. austr.
bracteolata, *E. Mey. ex DC. Prod.* vi. 78.—Afr. austr.
capensis, Linn. Mant. ii. 287 = Matricaria capensis.
cenioefolia, *DC. Prod.* vi. 79.—Afr. austr.
chamaemelifolia, *Ehrenb. ex C. Koch, in Bot. Zeit.* i. (1843) 41.—Africa!
chrysanthemifolia, Blume, Bijdr. 918 = Dichrocephala chrysanthemifolia.
cinerea, *Delile, Fl. Egypte*, 275. t. 47. f. 4.—Afr. bor.; Arab.
cinerea, Kotschy, ex Benth. & Hook. f. Gen. ii. 420 = Kotschyi.
complanata, Sibth. & Sm. Fl. Graec. Prod. ii. 187 = Anthemis pectinata.
corica, Wall. Cat. n. 3185 = Spilanthes Acmella.
coronopifolia, *Linn. Sp. Pl.* 892.—Afr. austr. (Europ. alien.).
crinita, E. Mey. ex DC. Prod. vi. 144 = Hippia gracilis.
cryptocephala, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 420.—Abyss.
cunifolia, Willd. Sp. Pl. iii. 2169 = Centipeda orbicularis.
Cunninghamii, F. Muell. Fragm. viii. 149 = Centipeda Cunninghamii.
debilis, Balb. ex Spreng. Neue Entdeck. ii. 187 = Eclipta pusilla.
decumbens, Pers. Syn. ii. 464 = Centipeda orbicularis.
dichotoma, Pers. l. c. = C. Pyrethrata.
dichrocephala, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 419 = anthemoides.
dichrocephaloides, C. B. Clarke, Comp. Ind. 150 = Dichrocephala Hamiltoni.
dioica, *Hook. f. Handb. N. Zeal. Fl.* 148.—N. Zel.
divaricata, Wall. ex DC. Prod. v. 576 = Thespis divaricata.
Drummondii, *Benth. Fl. Australiens.* iii. 550.—Austral.
abractenta, Walp. Rep. ii. 991 = bracteolata.
abractenlata, C. Koch, in Bot. Zeit. i. (1843) 39 = bracteolata.
elongata, *C. B. Vogel, in Trew, Pl. Rar.* iii. 11. t. 26. —Java.
Featherstonei, *F. Muell. ex J. Buch. in Trans. N. Z. Inst.* vii. (1875) 267.—Ins. Chatham.
Filicula, *Hook. f. ex Benth. Fl. Australiens.* iii. 551.—Austral.
filifolia, *Thunb. Prod. Pl. Cap.* 161.—Afr. austr.
filiformis, *Hook. f. Handb. N. Zeal. Fl.* 148.—N. Zel.

COTULA:—
fimbriata, Spreng. Neue Entdeck. iii. 41 = Hippia gracilis.
fontida, Poepp. ex DC. Prod. vi. 130 (= *Myriogyne clatinoides*).—Chili.
globifera, Licht. ex Steud. Nom. ed. II. i. 637 = Fentzia globosa.
globifera, Thunb. Prod. Pl. Cap. 162 = Matricaria globifera.
grandis, Jacq. Obs. Bot. iv. 4. t. 81 = Chrysanthemum floccolosum.
grandis, Linn. Sp. Pl. ed. II. 1257 (= *Plagius grandiflorus*).—Afr. bor.
gymnogyne, *F. Muell. ex Benth. Fl. Australiens.* iii. 549. —Austral.
hemisphaerica, *Wall. Cat.* n. 3236.—Ind. or.
heterocarpa, *DC. Prod.* vi. 80.—Afr. austr.
hispida, Harv. in Harv. & Sond. Fl. Cap. iii. 183 = Cenia hispida.
humifusa, Willd. ex Steud. Nom. ed. II. i. 637 = Egletes humifusa.
integrifolia, Burch. Trav. i. 51 = bracteolata.
integrifolia, *Hook. f. Fl. Tasm.* i. 192. t. 50 B.—Tasmania.
Kotschyi, *Benth. & Hook. f. Gen.* ii. 429.—Afr. trop.
lanata, *Hook. f. Handb. N. Zeal. Fl.* 141.—N. Zel.
latifolia, Forsk. Fl. Aegypt. Arab. 164 = Dichrocephala latifolia.
lasa, *DC. Prod.* vi. 78.—Afr. austr.
leptalea, *DC. l. c.* 80.—Afr. austr.
linearifolia, *Cheesem. in Trans. N. Z. Inst.* xv. (1883) 399.—N. Zel.
linifolia, Burm. f. Fl. Cap. Prod. 27 = Felicia fragilis.
lyrata, Boj. ex DC. Prod. v. 372 = Dichrocephala lyrata.
madagascarica, Willd. Sp. Pl. iii. 2170 = Grangea madagascarica.
Manlotota, *Petrie, in Trans. N. Z. Inst.* xiv. (1882) 362.—N. Zel.
matricarioides, Bong. Veg. Ins. Sitcha, 29 = Matricaria discoidea.
microcephala, *DC. Prod.* vi. 79.—Afr. austr.
minima, *DC. l. c.* 80.—Hab.!
minima, Willd. Sp. Pl. iii. 2170 = Centipeda orbicularis.
minor, Caruel, ex Boiss. Fl. Orient. iii. 358 = cinerea.
minor, *Hook. f. Handb. N. Zeal. Fl.* 143.—N. Zel.
minuta, Forst. f. Prod. 57 = Centipeda orbicularis.
montevidensis, Spreng. Syst. iii. 497 = coronopifolia.
Moseleyi, *Hemsl. Bot. Voy. Challenger*, i. 11. 152.—Tristan d'Acugna.
multifida, DC. Prod. vi. 80 = villosa.
myriophylloides, *Hook. Ic. Pl.* t. 835.—Afr. austr.
nudicaulis, *Thunb. Prod. Pl. Cap.* 162.—Afr. austr.
Oederi, *Murr. Hort. Bot. Stirp.* 228.—Hab.!
oxydonta, DC. Prod. vi. 78 = bipinnata.
pectinata, *Hook. f. Handb. N. Zeal. Fl.* 142.—N. Zel.
perpusilla, *Hook. f. l. c.* 148.—N. Zel.
pilulifera, Linn. f. Suppl. 378 = Matricaria globifera.
Piper, *Vell. Fl. Flum.* viii. t. 136.—Bras.
plumosa, *Hook. f. Handb. N. Zeal. Fl.* 141.—N. Zel.
poskilophylla, *C. Koch, in Bot. Zeit.* i. (1843) 39.—Abyss.
prostrata, Linn. Syst. ed. XII. 564 = Eclipta erecta.
pterocarpa, *DC. Prod.* vi. 80.—Afr. austr.
pubescens, *Desf. Fl. Atlant.* ii. 264.—Afr. bor.
pubescens, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 278 = setacea.
pumilio, Hoott. Handleid. z. 278 = Cenia turbinata.
pusilla, *Thunb. Prod. Pl. Cap.* 162.—Afr. austr.
pygmaea, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 330.—Mexico.
pygmaea, Poir. Encyc. Suppl. ii. (1810) 371 = Nananthea perpusilla.
Pyrethraria, *Linn. Mant.* i. 116.—Am. austr.
pyrethrifolia, *Hook. f. Handb. N. Zeal. Fl.* 149.—N. Zel.
quinqueloba, Thunb. Prod. Fl. Cap. 161 = Fentzia quinquefida.
quinqueloba, Blanco, Fl. Filip. ed. I. 636 = Centipeda orbicularis.
quinqueloba, Linn. f. Suppl. 377 = Lidbeckia lobata.
reptans, *Benth. Fl. Australiens.* iii. 551.—Austral.
reptans, Boj. ex DC. Prod. v. 371 = Sphaeranthus ecutellatus.
rosea, *Boj. ex Less. in Linnaea*, vi. (1831) 152.—Madag.
Russeliana, Wall. Cat. n. 3240 = Sphaeranthus Russeliana.
setacea, Linn. f. Suppl. 377 = Cenia setacea.

COTULA :—
sinapifolia, Buch.-Ham. ex Roxb Hort. Beng. 69 ; Fl. Ind. iii. 427 = Dichrocephala latifolia.
sinapifolia, Wall. ex DC. Prod. v. 575 = Thespis divaricata.
sonchifolia, Bieb. Fl. Taur. Cauc. iii. 328 = Dichrocephala latifolia.
sororia, *DC. Prod.* vi. 79.—Afr. austr.
Sphaeranthus, Link, Enum. Hort. Berol. ii. 344 = Grangea madraspatana.
Spilanthus, Linn. Mant. i. 116 = Spilanthes urens.
squalida, *Hook. f. Handb. N. Zeal. Fl.* 148.—N. Zel.
stenophylla, *C. Koch, in Bot. Zeit.* i. (1848) 40.—Afr. austr.
sternutatoria, Wall. Cat. n. 3259 = Centipeda orbicularis.
stolonifera, Harv. in Harv. & Sond. Fl. Cap. iii. 182 = caniasfolia.
stricta, Linn. Mant. 287 = Lidbeckia pectinata.
tanacetifolia, Linn. Syst. ed. XII ; Murr. Syst. Veg. ed. XIII. 844 n. 10 = Matricaria tanacetifolia.
Teesdaliae, *DC. Prod.* vi. 80.—Afr. austr.
tenella, *E. Mey. ex DC. l. c.* 80.—Afr. austr.
thespidioides, F. Muell. Fragm. viii. 143 = Centipeda thespidioides.
Thunbergii, Harv. in Harv. & Sond. Fl. Cap. iii. 183 = Cenia Thunbergii.
tridentata, *Benth. & Hook. f. Gen.* ii. 430.—Aegypt.
tripinnata, Thunb. Prod. Pl. Cap. 162 = Matricaria grandiflora.
turbinata, Linn. Sp. Pl. 892 = Cenia turbinata.
umbellata, Linn. f. Suppl. 378 = Peyrousea calycina.
vaiparadisea, *Phil. in Anal. Univ. Chil.* (1872) 508.—Chili.
Verbesina, *Linn. Amoen. Acad.* v. 407 = Adenostemma Swartzii.
villosa, *DC. Prod.* vi. 79.—Afr. austr.
viscosa, Linn. Sp. Pl. 892 = Egletes viscosa.
Zeyheri, *Fenzl, ex Harv. & Sond. Fl. Cap.* iii. 180.—Afr. austr.

COTULINA, Pomel, Nouv. Mat. Fl. Atl. 56 (1874) = Cotula, Tourn. (Compos.).
aurea, Pomel, l. c. = Cotula aurea.

COTYLANTHERA, Blume, Bijdr. 707 (1825).
GENTIANEAE, Benth. & Hook. f. ii. 808.
EOPHYLON, A. Gray, in Journ. Linn. Soc. xi. (1871) 22.
paniculosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 94.—Reg. Himal.
tenuis, *Blume, Bijdr.* 708.—Java.

COTYLARIA, Raïn. Princ. Somiol. 25 (1814) = Cotyledon, Tourn. (Crassulac.).

COTYLEDON, Tourn. ex Linn. Syst. ed. I (1735).
CRASSULACEAE, Benth. & Hook. f. i. 659.
COTYLARIA, Raïn. Princ. Somiol. 25 (1814).
COTYLIPHYLLUM, Link, Handb. ii. 22 (1831).
COURANTIA, Lam. Encyl. Fleur. i. (1851) Misc. 98.
ECHEVERIA, DC. Prod. iii. 401 (1828).
OROSTACHYS, [Fisch.] Hort. Gorenk. ed. I. 99 (1808).
ORTHOSTACHYS, Spach, Hist. Vég. Phan. ix. 32 (1840).
PACHYPHYTUM, Klotzsch, in Otto & Dietr. Allg. Gartenz. ix. (1841) 9.
PISTORINIA, DC. Prod. iii. 399 (1828).
UMBILICARIA, Pers. Syn. i. 510 (1805).
UMBILICUS, DC. in Bull. Soc. Philom. (1801) 49.
acaulon, Wall. ex Steud. Nom. ed. I. 280 = spuria.
acutifolia, *Baker, in Saund. Refug. Bot.* i. sub t. 71. n. 34.—Mexic.
adnata, *Baker, l. c.* t. 90.—Mexic.
aegyptiaca, Lam. Encyc. ii. 142 = Kalanchoë crenata.
africana, Burm. f. ex Steud. Nom. ed. I. 280 = spuria.
agavoides, *Baker, in Saund. Refug. Bot.* t. 67.—Mexic.
albiflora, *Hemsl. Diagn. Pl. Nov.* 9.—Mexic.
alborosea, Haw. Suppl. Pl. Succ. 26 = mamillata.
alborosea, Vahl, Symb. Bot. ii. 51 = Kalanchoë alborosea.
amblyon, Schlech. Prod. 367 = orbiculata.
amblyophylla, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 452 = Kalanchoë floribunda.

COTYLEDON :—
arborescens, Mill. Gard. Dict. ed. VIII. n. 7 = Crassula arborescens.
ariculata, Walp. Rep. ii. 258, sphalm. = cristata.
atropurpurea, *Baker, in Saund. Refug. Bot.* t. 198.—Mexic.
Batesii, *Hemsl. Diagn. Pl. Nov.* 9.—Mexic.
bifida, *Hemsl. Biol. Centr. Am. Bot.* i. 388.—Mexic.
bracteolata, *Baker, in Saund. Refug. Bot.* sub t. 62. —N. transat.
brasilica, Vell. Fl. Flum. 197 ; iv. t. 184 = Kalanchoë brasiliensis.
cacalioides, Eckl. & Zeyh. Enum. 307 = Ecklonianа.
cacalioides, *Linn. f. Suppl.* 243.—Afr. austr.
cacalioides, *How. Syn. Pl. Succ.* 107.—Am. bor. occ.
californica, *Baker, in Saund. Refug. Bot.* 70.—Am. bor. occ.
calycina, Roth, Nov. Pl. Sp. 217 = Bryophyllum calycinum.
canaliculata, *Baker, in Saund. Refug. Bot.* i. sub t. 58.—Mexic.
canaliculata, Haw. Suppl. Pl. Succ. 29 = ungulata.
canaliculata, Haw. in Phil. Mag. (1826) 327 = coruscans.
carnicolor, *Baker, in Saund. Refug. Bot.* iii. t. 190.—Mexic.
caryophyllacea, *Burm. f. Prod. Fl. Cap.* 18.—Afr. austr.
cespitosa, Haw. Misc. 180 = caespitosa.
chilensis, *Bull., in Journ. Linn. Soc.* xxii. (1885) 38. —Peruv.
clavifolia, *Haw. in Phil. Mag.* (1827) 274.—Afr. austr.
coccinea, *Cav. Ic.* ii. 54. t. 170.—Mexic.
Cooperi, *Baker, in Saund. Refug. Bot.* t. 72.—Afr. austr.
Cornuti, *Baker, in Gard. Chron.* (1874) 599.—Mexic.
corsaensa, *Haw. Suppl. Pl. Succ.* 21.—Afr. austr.
corymbosa, Rottl. ex Wight & Arn. Prod. 360 = Kalanchoë floribunda.
Cossoniana, *Ball, in Journ. Bot.* xi. (1873) 332.—Marocc.
crassifolia, Haw. in Phil. Mag. (1827) 278 = orbiculata ?
crassifolia, Salisb. Prod. 307 = hemisphaerica.
crenata, Vent. Jard. Malm. 49 = Kalanchoë crenata.
crispa, *How. in Phil. Mag.* (1837) 274.—Afr. austr.
cunnata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 372 = Meyeri.
cunnata, *Thunb. Prod. Pl. Cap.* 83.—Afr. austr.
cuneiformis, *How. in Phil. Mag.* (1826) 185.—Afr. austr.
curviflora, *Sims, Bot. Mag.* t. 2044.—Afr. austr.
cymosa, *Baker, in Saund. Refug. Bot.* t. 68.—Mexic.
decipiens, *Baker, l. c.* 200.—Peruv.
decipiens, *Sims, Bot. Mag.* t. 2518.—Afr. austr.
deficiens, Forsk. Fl. Aegypt. Arab. 89 = Kalanchoë crenata.
deficiens, Hochst. & Steud. ex A. Rich. Tent. Fl. Abyss. i. 310 = Kalanchoë Schimperiana.
Desmetiana, *Hemsl. Biol. Centr. Am. Bot.* i. 389.—Mexic.
dichotoma, *Hemsl. Suppl. Pl. Succ.* 27.—Afr. austr.
Ecklon iana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 374. —Afr. austr.
edulis, *Brenner, in Bot. Calif.* i. 211.—Calif.
elata, Haw. Suppl. Pl. Succ. 20 = orbiculata, trifora.
erecta, D. Dietr. Syn. Fl. ii. 1626 = lustanica.
farinosa, *Baker, in Saund. Refug. Bot.* i. t. 71.—Calif.
farinulenta, *Hemsl. Biol. Centr. Am. Bot.* i. 389.—Mexic.
fasciculata, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 106 ; *Bot. Mag.* t. 5605.—Afr. austr.
filicaulis, Eckl. & Zeyh. Enum. 307 = mamillaris.
fimbriata, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 241.—China ; Au. bor.
fulgens, *Baker, in Saund. Refug. Bot.* t. 64.—Mexic.
Galeottiana, *Hemsl. Diagn. Pl. Nov.* i. 9.—Mexic.
gibbiflora, *Moç. & Sesse, ex DC. Prod.* iii. 401.—Mexic.
glauca, *Baker, in Saund. Refug. Bot.* t. 61.—Mexic.

COTYLEDON :—
globulariaefolia, *Baker, in Saund. Refug. Bot.* t. 201 —Syria ; As. Min.
gracilis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 378.—Afr. austr.
gracilis, *How. Suppl. Pl. Succ.* 26.—Afr. austr.
grandiflora, Burm. f. Fl. Cap. Prod. 13 = tuberculosa.
Grayii, *Baker, in Saund. Refug. Bot.* sub t. 71. n. 33. —N. Mexic.
hemisphaerica, *Linn. Sp. Pl.* 429.—Afr. austr.
heterophylla, Roxb. Hort. Beng. 34 ; Fl. Ind. ii. 456 = Kalanchoë floribunda.
hirsuta, Herb. Heyne, ex C. B. Clarke, in Fl. Brit. Ind. 414 = Kalanchoë glandulosa.
hispanica, *Linn. Sp. Pl.* 1196.—Hispan. ; Marocc.
hispida, Lam. Encyc. ii. 141 = C. Mucizonia.
horizontalis, Guss. ind. Sem. Hort. Boccad. (1826) 4 = C. Umbilicus.
hybrida, Hort. Par. ex DC. Prod. iii. 395 = Kalanchoë spathulata.
incarnata, *Bull, in Journ. Linn. Soc.* xxii. (1885) 37 (err. typ. *incanum*).—Peruv.
integra, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 200. t. 9 = Kalanchoë crenata.
interjecta, *Haw. in Phil. Mag.* (1826) 185.—Afr. austr.
japonica, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 122.—Japon.
jasminiflora, *Salm-Dyck, Obs.* 5.—Afr. austr.
Jurgensenii, *Hemsl. Diag. Pl. Nov.* 9.—Mexic.
lacinista, Linn. Sp. Pl. 430 = Kalanchoe lacinista.
lanceolata, *Benth. & Hook. f. ex S. Wats. Bot. Calif.* i. 211.—Calif.
lanceolata, Blanco, Fl. Filip. ed. I. 282 = spathulata.
lanceolata, Forsk. Fl. Aegypt. Arab. 89 = Kalanchoë lanceolata.
laxa, Benth. & Hook. f. ex Bot. Calif. i. 212 = californica.
leucantha, *Ledeb. Fl. Alt.* ii. 198.—As. bor.; Sibir. altaic.
libanotica, Labill. Ic. Pl. Syr. Dec. iii. 3.—Kurdist. ; Syria.
Lieveni, *Ledeb. Fl. Alt.* ii. 197.—Persia ; Sibiria.
linguaefolia, *Lem. Illustr. Hortic. x.* (1863) *Misc.* 81, in syn.—Mexic.
linguaeformis, [R. Br. in] Ait. Hort. Kew. ed. II. iii. 109 = caespitosa.
Lingula, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 293.—Am. bor. occ.
lurida, *Baker, in Saund. Refug. Bot.* t. 59.—Mexic.
lusitanica, *Lam. Encyc.* ii. 140.—Europ.
lutea, Hort. ex Lam. l. c. 172 = Crassula portulacea.
lutea, Huds. Fl. Angl. ed. II. 194 = lusitanica.
maculata, *Salm-Dyck, Obs.* 5.—Afr. austr.
malacophylla, *Pall. Reise*, iii. 728.—Sibiria.
mamillaris, *Linn. f. Suppl.* 242.—Afr. austr.
marginalum, *Regin. Afr. Journ.* 79.—Hab. ?
mexicana, *Hemsl. Biol. Centr. Am. Bot.* i. 390.—Mexic.
Meyeri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 372.—Afr. austr.
Mucizonia, *Ortega, Mucis.* (1772).—Hispan. ; Marocc.
mucronata, *Baker, in Saund. Refug. Bot.* sub t. 55. n. 4.—Mexic.
mucronata, *Lam. Encyc.* ii. 142.—Afr. austr.
nevadensis, *S. Wats. Bot. Calif.* i. 212.—Am. bor. occ.
nodulosa, *Baker, in Saund. Refug. Bot.* t. 56.—Mexic.
nuda, *Baker, l. c.* t. 57.—Mexic.
nudicaulis, Lam. Encyc. ii. 142 ; Vahl, Symb. Bot. ii. 51 = Kalanchoë crenata.
oblonga, Haw. Syn. Pl. Succ. 106 = orbiculata.
ombilicus, Lam. Encyc. ii. 140 = C. Umbilicus.
oppositifolia, *Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb.* ii. (1837) 818.—Reg. Caucas.
orbiculata, Forsk. Fl. Aegypt. Arab. p. cxii = Kalanchoë attenuata.
orbiculata, *Linn. Sp. Pl.* 429.—Afr. austr.
Oreades, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 416.—Ind. or.
oregonensis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 372.—Am. bor. occ.
ovata, Haw. Syn. Pl. Succ. 105 = orbiculata.
ovata, Mill. Gard. Dict. ed. VIII. n. 8 = Crassula portulacea.
Pachyphytum, *Baker, in Saund. Refug. Bot.* i. sub t. 59. n. 12.—Mexic.
Palmeri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 292. —Am. bor. occ.

COTYLEDON :—
paniculata, Linn. f. Suppl. 242 = orbiculata.
paniculata, Retzl. ex Wight & Arn. Prod. 360 = Kalanchoë floribunda.
paniculata, Thunb. Prod. Pl. Cap. 58 = fasciularis.
pannosa, *Baker, in Journ. Linn. Soc.* xvii. (1881) 269. —Madag.
papillaria, Haw. Suppl. Pl. Succ. 21 = decussata.
papillaris, *Linn. f. Suppl.* 243.—Afr. austr.
papillosa, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 58.—Afghan.
parviflora, *Desf. in Ann. Mus. Par.* xi. (1808) 444 ; *Sibth. & Sm. Fl. Graec. Prod.* i. 303.—Europ.
parviflora, *Hemsl. Diag. Pl. Nov.* 9.—Mexic.
parvula, *Burch. Trav.* i. 219.—Afr. austr.
Peacocki, *Baker, in Gard. Chron.* (1874) ii. 258.—Am. hor.
peltata, Wendl. Beobacht. 49 = C. Umbilicus.
peruviana, *Baker, in Saund. Refug. Bot.* sub t. 58.—Peruv.
Pestalozzae, *Mast. in Gard. Chron.* (1877) ii. 456.—Oriens.
pinnata, Lam. Encyc. ii. 141 = Bryophyllum calycinum.
Pistorinia, *Ortega, Mucis.* (1773).—Hispan.
pubescens, *Baker, in Saund. Refug. Bot.* i. sub t. 55. —Mexic.
pubescens, C. A. Mey. Vers. Pfl. Cauc. 150 = Sedum pilosum.
pulverulenta, *Baker, in Saund. Refug. Bot.* i. t. 66.—Calif.
pumila, *Baker, l. c.* t. 62.—Mexic.
punctata, Hort. ex Lam. Encyc. ii. 173 = Crassula arborescens.
purpurea, Haw. Suppl. Pl. Succ. 28 = curvifiora.
purpurea, *Thunb. Prod. Pl. Cap.* 88.—Afr. austr.
quitensis, *Baker, in Saund. Refug. Bot.* sub t. 55.—Ecuador.
racemosa, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 375. —Afr. austr.
ramosa, Haw. Suppl. Pl. Succ. 24 = orbiculata.
ramosissima, Mill. Gard. Dict. ed. VIII, n. 6 = orbiculata.
ramosissima, *Salm-Dyck, ex Hem. l. c.* 26.—Afr. austr.
reflexa, Willd. Enum. Hort. Berol. Suppl. 24 = caespitosa.
reticulata, *Thunb. Prod. Pl. Cap.* 88.—Afr. austr.
retusa, *Baker, in Saund. Refug. Bot.* sub t. 64. n. 23.—Mexic.
rhizophylla, Roxb. Hort. Beng. 34 ; Fl. Ind. ii. 456 = Bryophyllum calycinum.
rhomboidea, Eckl. & Zeyh. Enum. 307 = hemisphaerica.
rhombifolia, *Haw. in Phil. Mag.* (1825) 33.—Afr. austr.
rosea, *Lezs. in Linnaea*, ix. (1834) 177.—Sibiria.
rosseia, *Baker, in Saund. Refug. Bot.* sub t. 55. n. 3. —Mexic.
rotundifolia, Eckl. & Zeyh. Enum. 307 = hemisphaerica.
rotundifolia, Haw. in Phil. Mag. (1827) 278 = orbiculata.
rupestris, Salisb. Prod. 307 = C. Umbilicus.
Rusbyi, *Greene, in Bull. Torrey Club*, x. (1883) 125. —Am. bor. occ.
Salzmanni, Hook. f. Bot. Mag. t. 5801 = hispanica.
Samium, *Urv. in Mem. Soc. Linn. Par.* i. (1822) 306. —Ins. Min.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 354.—Am. bor. occ.
Schaerii, *Baker, in Saund. Refug. Bot.* sub t. 62. n. 19.—Mexic.
secunda, *Baker, l. c.* sub t. 60. n. 14.—Mexic.
sediformis, Lapeyr. Hist. Abr. Pl. Pyr. 267 = sedoides.
sedoides, *DC. Regn. Veget. Bot. Gen.*ès. ii. 79.—Europ.
Semperivum, *Rich. Bourbr. Cap.* 176.—Reg. Cap.
serrata, *Linn. Sp. Pl.* 429.—Ins. Min.
simensis, *Britten, in Oliver, Fl. Trop. Afr.* ii. 399.—Afr. trop.
spathulata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 416.—Reg. Himal.
spathulata, Poir. Encyc. Suppl. ii. 372 = Kalanchoë spathulata.
spinosa, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 416 = leucantha.
spinosa, *Linn. Sp. Pl.* 429.—Ins. Min.
Sprengeli, *Baker, in Saund. Refug. Bot.* sub t. 71.—Ecuador.

COTYLEDON :—
spuria, *Linn. Sp. Pl.* ed. II. 614.—Afr. austr.
stolonifera, *Baker, in Saund. Refug. Bot.* t. 63.—Mexic.
strictiflora, *Baker, l. c.* sub t. 69. n. 17.—N. Mexic.
subspicata, *Baker, l. c.* sub t. 71.—N. Granat.
subulata, C. A. Mey. Vers. Pfl. Cauc. 150 = Sedum subulatum.
subulifolia, *Baker, in Saund. Refug. Bot.* sub t. 71. n. 12.—Mexic.
tardiflora, Bonpl. Pl. Rar. Malm. 98. t. 37 = fasciularis.
tenuicaulis, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 57.—Afghan.
terrifolia, Lam. Encyc. ii. 139 = spuria.
tetrafolia, *Thunb. Prod. Pl. Cap.* 88.—Afr. austr.
thyrsiflora, *D. Dietr. Syn. Pl.* ii. 1037.—Sibir.
thyrsiflora, Maxim. in Bull. Acad. Petersb. xxix. (1884) 123 = leucantha.
tomentosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 378.—Afr. austr.
tricuspidata, Harv. in Phil. Mag. (1825) 33 = papillaris.
trifida, *Thunb. Prod. Pl. Cap.* 88.—Afr. austr.
trigyna, *Burch. Trav.* ii. 236.—Afr. austr.
tuberculosa, *Lam. Encyc.* iii. 139.—Afr. austr.
tuberosa, *Hort. Par. ex Webb & Berth. Phyt. Canar.* i. 177.—Ins. Canar.
umbilicata, Lam. Fl. Fr. iii. 59 = C. Umbilicus.
umbilicifolia, Stokes, Bot. Mat. Med. ii. 546 = C. Umbilicus.
Umbilicus, *Linn. Sp. Pl.* 429.—Europ. ; As. Min.
undulata, Eckl. & Zeyh. Enum. 306 = cuneata.
undulata, *Haw. Suppl. Pl.* 30.—Afr. austr.
ungulata, *Lam. Encyc.* ii. 139.—Afr. austr.
valentina, *Hook. f. Bot. Mag.* t. 5802.—Afr. austr.
ventricosa, *Burm. f. Pl. Cap. Prod.* 18.—Afr. austr.
Verae, Jacq. Hort. Schoenb. iv. 17 = Kalanchoë crenata.
viridis, Haw. in Phil. Mag. (1827) 278 = orbiculata ?
viscida, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 372.—Am. bor. occ.
viscosa, Vahl, Symb. Bot. ii. 51 = C. Muciosa.
Wallichii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 374.—Afr. austr.
Zeyheri, *Harv. l. c.* 377.—Afr. austr.

COTYLEPHORA, Meisn. Gen. Comm. 36 (1837) = **Nesaea**, Blume (Malvac.).
altissima, Meisn. l. c. = N. altissima.

COTYLIPHYLLUM, Link, Handb. ii. 28 (1831) = **Cotyledon**, Tourn. (Crassulac.).
arcuatum, Link, l. c. = Cotyledon Umbilicus.
horizontale, Link, l. c. = Cotyledon Umbilicus.
Muciconia, Link, l. c. = Cotyledon Muciconia.
Umbilicus, Link, l. c. = Cotyledon Umbilicus.

COTYLISCUS, Desv. Journ. Bot. iii. (1813) 164, 175 = **Senebiera**, DC. (Cracif.).
siliculosa, Desv. l. c. = S. siliculosa.

COTYLODISCUS, Radlk. in Sitzb. Math.-Phys. Acad. Muench. vii. (1878) 324. *SAPINDACEAE*. Melachanthus, *Radlk. l. c.*—Madag.

COTYLOPLECTA, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 14 = **Hibiscus**, Linn. (Malvac.).
corymbosa, *Alef. l. c.* 15 = H. corymbosus.
dongolensis, Alef. l. c. = H. lunariifolius.
macranthe, Alef. l. c. = H. lunariifolius.

COUELANDIA, Aubl. Pl. Gui. ii. 937. t. 356 (1775) = **Muellera**, Linn. (Legumin.).
frutescens, Aubl. l. c. = M. monilisformis.
mexicana, Benth. Adansonia, ix. (1870) 288.—Mexic.

COUDENBERGIA, March. in Bull. Acad. Belg. Sér. II. xlvii. (1879) 514 = **Cosmaneia**, March. (Araliac.).

COUEPIA, Aubl. Pl. Gui. i. 349. t. 321 (1775). *ROSACEAE*, Benth. & Hook. f. i. 606.
Acra, Willd. Sp. Pl. iii. 717 (1800).
Acuoa, Aubl. Pl. Gui. ii. 696. t. 280 (1775).
Dolacia, Neck. Elem. ii. 414 (1790).
brasiensis, *Benth. in Hook. Journ. Bot.* ii. (1840) 215. —Guiana.

COUEPIA :—
canomensis, *Benth. in Hook. Journ. Bot.* ii. (1840) 215. —Bras.
chrysocalyx, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 42.—Bras.
comosa, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Guiana.
crianths, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 45.—Bras.
glandulosa, *Miq. Stirp. Surin. Sel.* 38.—Guian.
glaucescens, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 49.—Bras.
grandiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 519, 307.—Guiana.
hypoleuca, *Miq. in Linnaea*, xix. (1847) 438.—Mexic.
Kunthiana, *Benth. in Hook. Journ. Bot.* ii. (1840) 216.—Mexic.
leptostachya, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 44.—Bras.
macrophylla, *Spruce, ex Hook. f. l. c.* 41.—Peruv.
magnoliaefolia, *Benth. ex Hook. f. l. c.* 43.—Bras.
Martiana, *Hook. f. l. c.* 47.—Bras.
multiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 215. —Guiana.
myrtifolia, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 44.—Bras.
ovatifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 216. —Bras.
parasitica, *Benth. l. c.*—Bras.
Parillo, *DC. Prod.* ii. 536.—Guiana.
racemosa, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 43.—Bras.
Steudeliana, *Miq. Stirp. Surin. Sel.* 38.—Guianas.
subcordata, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 46.—Bras.
thyrsiflora, *Benth. f. l. c.* 45.—Bras.
trimioides, Klotzsch, ex Hook. f. in Mart. Fl. Bras. xiv. ii. 48 = parasitica.
Uiti, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Bras.

COULA, Baill. Adansonia, iii. 61, 64. t. 3 (1862). *OLACINEAE*, Benth. & Hook. f. i. 995.
coulejia, *Baill. l. c.* 64.—Afr. trop.

COULEJIA, Dennst. Schluss. Hort. Malab. xi (1818) = **Antidesma**, Linn. (Euphorb.).
amentacea, Dennst. l. c.—Ind. or.

COULTERIA, H. B. & K. Nov. Gen. et Sp. vi. 328. t. 568 = **Cassalpinia**, Linn. (Legumin.).
africana, Guill. & Perr. Fl. Seneg. Tent. i. 256 = Propsopis oblonga.
chilensis, DC. Prod. ii. 481 = Cassalp. tinctoria.
horrida, H. B. & K. Nov. Gen. et Sp. vi. 330 = Cassalp. horrida.
mexicana, DC. Prod. ii. 481 = Cassalp. vesicaria.
mollis, H. B. & K. Nov. Gen. et Sp. vi. 330 = Cassalp. mollis.
tinctoria, H. B. & K. l. c. 331 = Cassalp. tinctoria.

COUMA, Aubl. Pl. Gui. Suppl. 39. t. 392 (1775). *APOCYNACEAE*, Benth. & Hook. f. ii. 693.
Collophora, Mart. in Buchn. Nep. Pharm. xxxv. (1836) 196, ex Bull. Férussac (Janv. 1831), 64.
dulcis, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. i. 19 = utilis.
guianensis, *Aubl. Pl. Gui. Suppl.* 39. t. 392.—Guiana.
rigida, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 20.—Bras.
utilis, *Muell. Arg. l. c.* 19.—Bras.

COUMAROUNA, Aubl. Pl. Gui. ii. 740. t. 296 (1775) = **Dipteryx**, Schreb. (1791) (Legumin.).
odora, Aubl. l. c. t. 296 = sp.
odorata, Aubl. l. c. 740 = D. odorata.

COUPIA, G. Don, Gen. Syst. ii. 44 (1832) sphalm. = Goupia, Aubl. (Celastrac.).

COUPOUI, Aubl. Pl. Gui. Suppl. 16. t. 377 (1775). *APOCYNACEAE* [Benth. & Hook. f. i. 696].
Cupuliana, Miers, Apocyn. S. Am. 16 (1878).
Cupuia, Radlk. Princ. Sembuk. 39 (1879).
aquatica, *Aubl. l. c.*—Guiana.

COURALIA, Spltg. in Hoev. & De Vriese, Tijdschr. ix. (1842) 14. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1049.
fluviatilis, *Spltg. l. c.* 15.—Guiana.

COURANTIA, Lem. Jard. Fleur. i. (1851) Misc. 92 = Octyladon, Tourn. (Crasulac.).
achroeriodes, Lem. l. c. = Cotyl. rosea.
rosea, Lem. l. c. ill. (1858) 344, in nota = Cotyl. rosea.

COURATARI, Aubl. Pl. Gui. ii. 723. t. 290 (1775). *MYRTACEAE*, Benth. & Hook. f. i. 722.
AMPHORICARPUS, Syrace, ex Miers, in Trans. Linn. Soc. xxx. (1874) 288.
CARINIANA, Casar. Nov. Stirp. Bras. 85 (1842).
asiaxocarpa, *Mart. ex Berg*, in *Mart. Fl. Bras.* xiv. 1. 509.—Bras.
coriacea, *Mart. ex Berg*, l. c. 510.—Bras.
dictyocarpa, *Mart. ex Berg*, l. c.—Bras.
domestica, *Mart. in Flora*, xx. (1837) 11. *Beibl.* 127, in textu.—Bras.
estrellensis, *Raddi*, in *Mem. Mod.* xviii. *Fis.* (1820) 408.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 724. t. 290.—Am. trop.
Lecythopsis, *Mart. in Flora*, xx. (1837) 11. *Beibl.* 89.—Bras.
legalis, *Mart. l. c.* 88.—Bras.
lineata, *Mart. ex Berg*, in *Mart. Fl. Bras.* xiv. 1. 508.—Bras.
macrocarpa, *Mart. ex Berg*, l. c. 509.—Bras.
Martiana, *Miers*, in *Trans. Linn. Soc.* xxx. (1874) 282.—Bras.
paraënsis, *Mart. ex Berg*, in *Mart. Fl. Bras.* xiv. 1. 510.—Bras.
rubra, *Gardn. ex Miers*, in *Trans. Linn. Soc.* xxx. (1874) 288.—Brsa.
Tauari, *Berg*, in *Mart. Fl. Bras.* xiv. 1. 509.—Bras.
usupensis, *Spruce, ex Berg*, l. c. 508.—Bras.
Vriesii, *Miers*, in *Trans. Linn. Soc.* xxx. (1874) 284.—Guiana.

COURATARI, Cambess. in St. Hil. Fl. Bras. Mer. ii. 379 (1829) = **Lecythopsis**, Schrank (Myrtac.).
glabra, Cambess. l. c. = L. glabra.
rufescens, Cambess. l. c. 380 = L. rufescens.

COURBARI, Adans. Fam. ii. 317 (1763) = **Hymenaea**, Linn. (Legumin.).

COURBARIL, Plum. ex Endl. Gen. 1317 (1841) = praec.

COURBONIA, Bronga. in Bull. Soc. Bot. Fr. vii. (1860) 901. *CAPPARIDEAE*, Benth. & Hook. f. i. 969.
PHYSANTHEMUM, Klotzsch, in Peters, Reise Mosamb. Bot. 167. t. 29 (1861).
decumbens, *Brongn. l. c.*—Afr. trop.
virgata, *Brongn. l. c.*—Afr. trop.

COURIMARI, Aubl. Pl. Gui. Suppl. 27 (1775) (Gen. dub. affin.).
guianensis, *Aubl. l. c.*—Guiana.

COURINGIA, Adans. Fam. ii. 418 (1763) = **Conringia**, Heist. (Crucif.).

COUROUPITA, Aubl. Pl. Gui. ii. 706. t. 282 (1775). *MYRTACEAE*, Benth. & Hook. f. i. 725.
ELSHOLTHIA, Neck. Elem. ii. 256 (1790).
PONTOPIDANA, Scop. Introd. 195 (1777).
antillana, *Miers*, in *Trans. Linn. Soc.* xxx. (1874) 191.—Ind. occ.
crenulata, *Miers*, l. c. 195.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 708. t. 282.—Guian.
guianensis, Hook. Bot. Mag. t. 6158 = antillana.
lentula, *Miers*, in *Trans. Linn. Soc.* xxx. (1874) 194.—Bras.
membranacea, *Miers*, l. c.—N. Granat.
nicaraguensis, *DC. Prod.* iii. 294.—Am. centr.
odoratissima, *Seem. Bot. Voy. Herald*, 126.—Panama.
peruviana, *Berg*, in *Linnaea*, xxxi. (1861–62) 961.—Peruv.
surinamensis, *Mart. ex Berg*, in *Mart. Fl. Bras.* xiv. 1. 478.—Guian.

COURRANTIA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276. t. 89 (dr. 1845) = **Mantisalca**, Linn. (Compos.).
dendroïdes, Sch. Bip. l. c. 278 = M. Courrantia.

COURSETIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 92. *LEGUMINOSAE*, Benth. & Hook. f. i. 501.
POISSONIA, Baill. Adansonia, ix. (1870) 295.
arborea, *Grisb. Fl. Brit. W. Ind.* 183.—Ins. Trinit.
dubia, *DC. Prod.* ii. 264.—Am. austr.
erianthe, *Benth. in Hook. Ic. Pl.* sub t. 1065.—Peruv.
glandulosa, *A. Gray*, in *Proc. Am. Acad.* v. (1861) 156.—Calif.
grandiflora, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 10.—Peruv.
microphylla, *A. Gray*, in *Proc. Am. Acad.* xvii. (1881–82) 201.—Arizona.
orbicularis, *Benth. in Hook. Ic. Pl.* t. 1065.—Peruv.
rosensis, *Benth. in Mart. Fl. Bras.* xv. 1. 44.—Am. trop.
tomentosa, *DC. in Ann. Sc. Nat. Sér.* I. iv. (1825) 92.—Peruv.
viciodes, *Benth. in Mart. Fl. Bras.* xv. 1. 44.—Am. trop.
virgata, *DC. in Ann. Sc. Nat. Sér.* I. iv. (1825) 92.—Mexic.

COURTENIA, R. Br. in Benn. Pl. Jav. Rar. 236 (1844) = Cola, Schott (Sterculiac.).
Afzelii, R. Br. l. c. = Cola Afzelii.
heterophylla, R. Br. l. c. = Cola heterophylla.
triloba, R. Br. l. c. = Cola heterophylla.

COURTOISIA, Nees, in Linnaea, ix. (1834) 286. *CYPERACEAE*, Benth. & Hook. f. iii. 1046.
cyperoides, *Nees*, l. c.—Ind. or.
olivacea, *Boeck. in Flora*, xliv. (1861) 331.—Guiana.

COUSINIA, Cass. in Dict. Sc. Nat. xlvii. 508 (1827). *COMPOSITAE*, Benth. & Hook. f. ii. 467.
AUCHERA, DC. Prod. vi. 557 (1837).
acicularis, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 321.—Turkestan.
Actinia, *Boiss. Diagn.* Ser. II. iii. 55.—Afghan.
actinocephala, Jaub. & Spach, Illustr. ii. 98. t. 175 = xiphilolepis.
acinaca, *Bunge, Gatt. Cous.* 80.—Persia.
affinis, *Schrenk, Enum. Pl. Nov.* i. 41.—Soongaria.
aggregata, *DC. Prod.* vi. 555.—Persia.
Aintabensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 508.—Syria.
alata, *Schrenk, Enum. Pl. Nov.* i. 40.—Reg. Casp.; Soongaria.
Alberti, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 315.—Turkestan.
albicaulis, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 128.—Persia.
albida, *DC. Prod.* vi. 552.—Persia.
Alepidea, *Boiss. Diagn.* Ser. II. iii. 53.—Beluchist.
alepplca, *Boiss. l. c.* Ser. I. x. 101.—Syria.
alpicola, Borszczow, ex Trautv. in Act. Hort. Petrop. viii. (1883) 484 = seq.
alpina, *Bunge, Del. Sem. Hort. Dorp.* (1845) 3.—Turkest.
amplissima, *Boiss. Fl. Orient.* iii. 462.—Persia.
anisoptera, Jaub. & Spach, Illustr. ii. 88. t. 170 = calolepis.
anomala, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 316.—Turkestan.
anthacantha, Regel, in Bull. Soc. Nat. Mosc. xl. (1867) ii. 156, sphalm. *vide* xanthacantha.
aptera, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 170.—Afghan.
arachnoides, *Fisch. & Mey. ex DC. Prod.* vi. 558.—Soongar.
aralensis, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 554 = minuta.
arenosa, *DC. Prod.* vi. 556.—Persia.
arctioides, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 115.—Soongaria.
arctoidifolia, *Bunge, Gatt. Cous.* 42.—Persia.
arenaria, *Bunge*, l. c. 39.—Persia.
assyriaca, Jaub. & Spach, Illustr. ii. 87. t. 179 = leucophlora.
atropatana, *Bunge, Gatt. Cous.* 17.—Persia.
Auchert, *DC. Prod.* vi. 555.—Cappadoc.
auriculata, *Boiss. Diagn.* Ser. II. iii. 58.—Ind. bor. occ.
auriculata, Hook. f. Fl. Brit. Ind. iii. 360 = buphthalmoides.
Balsmi, *Boiss. Diagn.* Ser. II. iii. 53.—Afghan.
baselsiana, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 476.—Persia.

COUSINIA:—
Belangeri, *DC. Prod.* vi. 553.—Persia.
Bienerti, *Bunge, Gatt. Cous.* 27.—Persia.
bipinnata, *Boiss. Diagn.* Ser. II. iii. 52.—Afghan.
Boissieri, *Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 127.—Persia.
Bonvalotii, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 320.—Turkestan.
brachyptera, *DC. Prod.* vi. 554.—Armenia.
Brantsii, C. B. Clarke, Comp. Ind. 214 = Thomsoni.
Bukiei, Bunge, Gatt. Cous. 22 = hamosa.
bulgarica, *C. Koch, in Linnaea*, xxiv. (1851) 385.—Europ. or.
Bungeana, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 318.—Turkestan.
buphthalmoides, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 155.—Afghan.; Ind. bor. occ.
cacaavra, *Boiss. & Bal. Diagn.* Ser. II. vi. 116.—Cappadoc.
calcitrapaeformis, Jaub. & Spach, Illustr. ii. 76. t. 158 = minuta.
Calcitrapa, *Boiss Fl. Orient.* iii. 490.—Persia.
calocephala, *Yaub. & Spach, Illustr.* ii. 96. t. 178.—Persia.
calolepis, *Boiss. Diagn.* Ser. I. vi. 116.—Persia.
calolepis, Boiss. Fl. Orient. iii. 480 = commutata.
Candolleana, *Yaub. & Spach, Illustr.* ii. 74. t. 156.—Persia.
canescens, *DC. Prod.* vi. 556.—Armenia.
canescens, *Franch. in Ann. Sc. Nat. Sér.* VI. xvi. (1883) 325.—Turkestan.
Capusi, *Franch. l. c.* 317.—Turkestan.
carduiformis, *Cass. in Dict. Sc. Nat.* xlvii. 508.—Reg. Caucas.
carduncelloides, *Regel & Schmalh. in Pl. Nov. Fedsch.* 50.—As. centr.
carlinoides, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 126 = olivacea.
carlinoides, *DC. Prod.* vi. 554.—Persia.
carthamoides, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 171.—Afghan.
catanicu, *Boiss. & Haussk. Fl. Orient.* iii. 488.—As. Min.
centauroides, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 150.—Persia.
centauroides, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 152.—As centr.
Chamaepeuce, *Boiss. Diagn.* Ser. I. vi. 120.—Persia.
chlorocephala, *C. A. Mey. ex DC. Prod.* vi. 554.—Armenia.
chrysacantha, *Yaub. & Spach, Illustr.* ii. 78. t. 160.—Persia.
cirsioides, *Boiss. & Bal. Diagn.* Ser. II. vi. 117.—Cappadoc.
commutata, *Bunge, Gatt. Cous.* 24.—Persia.
concinna, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 513.—Persia.
concolor, *Bunge, Gatt. Cous.* 33.—Persia.
congesta, *Bunge*, l. c. 19.—Persia.
coronata, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 321.—Turkestan.
crispa, *Yaub. & Spach, Illustr.* ii. 81. t. 162.—Persia.
cylindracea, *Boiss. Diagn.* Ser. I. vi. 115.—Persia.
cylindrocephala, *Yaub. & Spach, Illustr.* ii. 84. t. 166.—Persia.
cynaloeoides, *Boiss. Diagn.* Ser. I. vi. 119.—Kurdist.
cynaroides, *C. A. Mey. Verz. Pfl. Casc.* 230.—Reg. Caucas.
decipiens, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 125.—Persia.
decurrens, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 154.—Afghan.
deserti, *Bunge, Gatt. Cous.* 39.—Persia.
dichotoma, *Bunge, Del. Sem. Hort. Dorpat.* (1845) 3.—Turkest.
discolor, *Bunge, Gatt. Cous.* 34.—Persia.
dissecta, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 391.—Turcoman.
dolichoclada, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 107.—Soongaria.
elata, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 126.—Persia.
elegans, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 173.—Afghan.
erinacea, *Yaub. & Spach, Illustr.* ii. 83. t. 165.—Persia.
eriobasis, *Bunge, Gatt. Cous.* 37.—Persia.
eriocephala, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 504.—As. Min.

COUSINIA :—
- eriophora, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 318.—Turkestan.
- eryngioides, *Boiss. Diagn.* Ser. I. x. 101.—Persia.
- Falconeri, *Hook. f. Fl. Brit. Ind.* iii. 360.—Reg. Himal.
- flaviplena, *Franch. in Ann. Sc. Nat. Sér.* VI. xvi. (1883) 313.—Turkestan.
- foliosa, *Boiss. & Bal. Diagn.* Ser. II. vi. 116.—Cappadoc.
- gracilis, *Boiss. l. c.* Ser. I. vi. 117.—Persia.
- grandiceps, *Bunge, Gatt. Cous.* 54.—Persia.
- grandis, *C. A. Mey. ex DC. Prod.* vi. 557.—Persia.
- Griffithiana, *Boiss. Diagn.* Ser. II. iii. 55.—Afghan.
- *Gundelia*, C. Koch, in Linnæa, xxiv. (1851) 387 = brachyptera.
- Habilitzii, *C. A. Mey. ex DC. Prod.* vi. 554.—Persia.
- hamosa, *C. A. Mey. l. c.* 558.—Persia.
- hamulosa, *C. A. Mey. ex Boiss. Fl. Orient.* vi. 483 = hamosa.
- helianthæ, *Bunge, Gatt. Cous.* 89.—Persia.
- Hermonis, *Boiss. Diagn.* Ser. I. x. 102.—Syria.
- *heterolepis*, C. Koch, in Linnæa, xxiv. (1851) 386 = brachyptera.
- heterophylla, *Boiss. Diagn.* Ser. II. iii. 54.—Afghan.
- Hohenackeri, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 84.—Persia.
- hololenca, *Bunge, Gatt. Cous.* 27.—Persia.
- humilis, *Boiss. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 248.—Cappadoc.
- hypoleuca, *Boiss. Diagn.* Ser. I. x. 103 ; *Fl. Orient.* iii. 464.—Persia.
- *Hystrix*, C. B. Clarke, Comp. Ind. 214 = microcarpa.
- Hystrix, *C. A. Mey. Ind. Sem. Hort. Petrop.* ii. 84.—Persia.
- ilicifolia, *Jaub. & Spach, Illustr.* ii. 89. t. 164.—Persia.
- incompta, *DC. Prod.* vi. 553.—Persia.
- inflata, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* iii. 512. —Kurdistan.
- integrifolia, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 329.—Turkestan.
- intermedia, *C. A. Mey. ex DC. Prod.* vi. 553.—Armenia.
- involucrata, *Boiss. Diagn.* Ser. I. vi. 114.—Persia.
- karnavica, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 317.—Turkestan.
- *Karelini*, Less. ex Kar. in Bull. Soc. Nat. Mosc. (1839) 159 = onopordioides.
- kokanica, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 49.—As. centr.
- Kornhuberi, *Heimerl, in Denkschr. Acad. Wien*, l. (1885) 69.—Persia.
- Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 315.—Turkestan.
- Kotschyi, *Boiss. Diagn.* Ser. I. vi. 120.—Persia.
- Krauseana, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 48.—As. centr.
- lachnosphæra, *Bunge, Gatt. Cous.* 86.—Persia.
- lappacea, *Schrenk, Enum. Pl. Nov.* i. 42.—Soongaria.
- lasiandra, *Bunge, Gatt. Cous.* 40.—Persia.
- lasiolepis, *Boiss. Diagn.* Ser. I. vi. 113.—Persia.
- lepida, *Bunge, Gatt. Cous.* 98, 99.—Persia.
- leptocephala, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 159.—Turcoman.
- leucochlora, *Bunge, Gatt. Cous.* 24, in nota.—Persia.
- libanotica, *DC. Prod.* vi. 556.—Syria.
- lucida, *DC. l. c.* 553.—Assyria.
- lyrata, *Bunge, Gatt. Cous.* 53.—Persia.
- macrocephala, *C. A. Mey. Verz. Pfl. Cauc.* 231.—Reg. Caucas.
- macrolepis, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* iii. 505.—Persia.
- macroptera, *C. A. Mey. ex DC. Prod.* vi. 555.—Armenia.
- microcarpa, *Boiss. Diagn.* Ser. II. iii. 59.—Persia ; Reg. Himal.
- microcephala, *C. A. Mey. ex DC. Prod.* vi. 552.—Persia.
- minuta, *Boiss. Diagn.* Ser. I. vi. 117.—Persia ; Ind. bor. occ.
- mollis, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1846) 106.—Soongaria.
- monocephala, *Bunge, Gatt. Cous.* 59.—Persia.
- multiloba, *DC. Prod.* vi. 554.—Persia.
- neurocentra, *Bunge, Gatt. Cous.* 29.—Persia.
- Notzeana, *Boiss. Diagn.* Ser. II. iii. 57.—Persia.
- oberrata, *Boiss. Fl. Orient.* iii. 491.—Persia.
- odontolepis, *DC. Prod.* vi. 556.—Assyria.

COUSINIA :—
- *odontolepis*, Sch. Bip. ex Boiss. l. c. 512 = symbolepis.
- Olgæ, *Regel & Schmalh. in Pl. Nov. Fedsch.* 48.—As. centr.
- oligocephala, *Boiss. Diagn.* Ser. I. vi. 116.—Persia.
- *oligocephala*, Hausskn. ex Boiss. Fl. Orient. iii. 475 = bachtiarica.
- Olivieri, *DC. Prod.* vi. 552.—Persia !
- *Olivieri*, Jaub. & Spach, Illustr. t. 178 = serratuloides.
- onopordioides, *Ledeb. in Eichw. Pl. Casp. Cauc.* 60. t. 27.—Turcoman.
- orientalis, *C. Koch, in Linnæa*, xxiv. (1851) 386.—Orient.
- ovtchachemxia, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 322.—Turkestan.
- *paimatiloba*, Jaub. & Spach, Illustr. ii. 80. t. 162 = multiloba.
- *pancifolva*, Bunge, Gatt. Cous. 19 = aggregata.
- pentacantha, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 315.—As. centr.
- pergamacea, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* iii. 513.—Persia.
- Pestalozzæ, *Boiss. l. c.* 471.—Syria.
- pinnatocephala, *Boiss. Diagn.* Ser. I. vi. 121.—Persia.
- pipirocephala, *Bunge, Gatt. Cous.* 26.—Persia.
- *platyacantha*, Bunge, l. c. 87 = centauroides.
- platyleps, *Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop. Ins. Suppl.* 10.—Turkestan.
- polycephala, *Rupr. Sert. Tianschi.* 54.—As. centr.
- prasina, *Jaub. & Spach, Illustr.* ii. 86. t. 168.—Persia.
- Princeps, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 324.—Turkestan.
- prolifera, *Jaub. & Spach, Illustr.* ii. 74. t. 157.—Persia.
- pterocarpa, *Boiss. Fl. Orient.* iii. 501.—Afghan.
- pugionifera, *Jaub. & Spach, Illustr.* ii. 92. t. 174.—Persia.
- pulchella, *Bunge, Del. Sem. Hort. Dorpat.* (1845) 5.—As. centr.
- purpurea, *C. A. Mey. ex DC. Prod.* vi. 555.—Persia.
- *pycnoloba*, Boiss. Diagn. Ser. II. iii. 57 = multiloba.
- *pyrrhophaka*, Jaub. & Spach, Illustr. ii. 85. t. 167 = Olivieri.
- racemosa, *Boiss. Diagn.* Ser. II. iii. 56.—Afghan.
- radiata, *Bunge, Del. Sem. Hort. Dorpat.* (1846) 3.—As. centr.
- ramosissima, *DC. Prod.* vi. 552.—Cappadoc.
- recurvata, *DC. l. c.* 555.—Persia.
- sahewensis, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* iii. 488.—Persia.
- Scala, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 131.—Afghan.
- scariosa, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 154.—Afghan.
- Schischurowskiana; *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 50.—As. centr.
- *Schultziana*, Jaub. & Spach, Illustr. ii. 95. t. 177 = cymbolepis.
- Seidlitzii, *Bunge, Gatt. Cous.* 90.—Persia.
- Semenovii, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 147.—As. centr.
- serratuloides, *Boiss. Diagn.* Ser. I. vi. 118.—Persia.
- Severzowii, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 151.—Turkestan.
- *silybioides*, Jaub. & Spach, Illustr. ii. 94. t. 176 = brachyptera.
- Smirnowii, *Trautv. in Act. Hort. Petrop.* viii. (1883) 488.—As. centr.
- *sphærocephala*, Jaub. & Spach, Illustr. ii. 79. t. 161 = C. Chamæpersea.
- squarrosa, *Boiss. Diagn.* Ser. I. x. 103.—Persia.
- *squarrosa*, C. Koch, in Linnæa, xxiv. (1851) 387 = brachyptera.
- stenocephala, *Boiss. Diagn.* Ser. I. vi. 115.—Assyria.
- stenolepis, *Regel & Schmalh. in Pl. Nov. Fedsch.* 49. —As. centr.
- subnutans, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 314.—Turkestan.
- *sylvicola*, Bunge, Del. Sem. Hort. Dorp. (1845) 5 = dichotoma.
- tabrisiana, *Bunge, Gatt. Cous.* 16.—Persia.
- tenella, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 25.—Armenia.
- tenuifolia, *C. A. Mey. ex DC. Prod.* vi. 554.—Persia.
- thamnodes, *Boiss. & Hausskn. in Boiss. Fl. Orient.* iii. 471.—Persia.
- Thomsoni, *C. B. Clarke, Comp. Ind.* 213.—Reg. Himal.

COUSINIA :—
- trachylepis, *Bunge, Gatt. Cous.* 22.—Persia.
- Trautvetteri, *Regel, in Act. Hort. Petrop.* v. (1877) 327.—Turkestan.
- trifoca, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 106.—Soongaria.
- umbrosa, *Bunge, Gatt. Cous.* 10.—Persia.
- uncinata, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 145.—Soongaria.
- verbascifolia, *Bunge, Gatt. Cous.* 59.—Persia.
- verticillaris, *Bunge, in Mém. Sav. Etr. Pétersb.* vi. (1851) 857.—As. centr.
- Verutum, *Bunge, Gatt. Cous.* 31.—Persia.
- wolgensis, *C. A. Mey. ex DC. Prod.* vi. 555.—Rossia.
- xanthacantha, *Regel, in Bull. Soc. Nat. Mosc.* xl (1867) ii. 156 (*sphalm.* anthacantha).—Orient.
- *xeranthemoides*, Bunge, ex Boiss. Fl. Orient. iii. 497 = microcarpa.
- siphiloides, *Boiss. Diagn.* Ser. I. vi. 121.—Persia.

COUSSAPOA, Aubl. Pl. Gui. ii. 955. t. 369, 370 (1775). *URTICACEÆ*, Benth. & Hook. f. iii. 379.
- acutifolia, *Klotzsch, in Linnæa*, xx. (1847) 529.—Peruv.
- angustifolia, *Aubl. Pl. Gui.* ii. 956. t. 368.—Guiana.
- asperifolia, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 96.—Guiana.
- cuneata, *Miq. in Mart. Fl. Bras.* iv. i. 188.—Guiana.
- dealbata, *Andrè, Illustr. Hortic.* xvii. (1870) 16.—Bras.
- fagifolia, *Klotzsch, in Linnæa*, xx. (1847) 528.—Guiana.
- ferruginea, *Trec. in Ann. Sc. Nat.* Ser. III. viii. (1847) 98.—Guiana.
- Fonsinasiana, *A. Brongn. ex Trec. l. c.* 94.—Bras.
- hirsuta, *Trec. l. c.* 97.—Peruv.
- intermedia, *Mart. ex Miq. in Mart. Fl. Bras.* iv. i. 188.—Bras.
- laevigata, *Poepp. & Endl. Gen. et Sp.* ii. 33.—Bras.
- latifolia, *Aubl. Pl. Gui.* ii. 955. t. 362.—Guiana.
- magnifolia, *Trec. in Ann. Sc. Nat.* Ser. III. viii. (1847) 98.—Peruv.
- Martiana, *Miq. in Mart. Fl. Bras.* iv. i. 192.—Bras.
- microcephala, *Trec. in Ann. Sc. Nat.* Ser. III. viii. (1847) 96.—Guiana.
- nitida, *Miq. in Mart. Fl. Bras.* iv. i. 188.—Bras.
- obovata, *Miq. in Verol. en Meded. Kon. Nederl. Inst.* 1843 (1848) 200.—Hab. ?
- ovalifolia, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 95.—Peruv.
- paberula, *Klotzsch, in Linnæa*, xx. (1847) 529.—Peruv.
- Ruiziî, *Klotzsch, l. c.* 529.—Peruv.
- Schottii, *Miq. in Mart. Fl. Bras.* iv. i. 187.—Bras.
- setosa, *Klotzsch, in Linnæa*, xx. (1847) 528.—Peruv.
- sublanosa, *Mart. ex Miq. in Mart. Fl. Bras.* iv. i. 184. —Bras.
- velleræa, *Klotzsch, in Linnæa*, xx. (1847) 527.—Peruv.
- villosa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 33. t. 147. —Peruv.

COUSSAREA, Aubl. Pl. Gui. i. 93. t. 36 (1775). *RUBIACEÆ*, Benth. & Hook. f. ii. 120.
- BILLARDIERIA, Vahl, Eclog. Am. i. 18. t. 10 (1796).
- FRORIEPSIA, Vahl, l. c. Praef. 9 (1796).
- PECHEYA, Scop. Introd. 148 (1777).
- acuminata, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
- albescens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 87, in nota.—Guiana.
- amœna, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
- bahiensis, *Muell. Arg. l. c.* 467.—Bras.
- bidora, *Muell. Arg. l. c.* 466.—Bras.
- capitata, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
- Catingana, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
- coriacea, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 89.—Bras.
- congestiflora, *Muell. Arg. in Flora*, lviii. (1875) 466. —Bras.
- contracta, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
- corcovadensis, *Muell. Arg. in Flora*, lviii. (1875) 469. —Bras.
- corniflora, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
- flava, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 26. t. 231. —Peruv.

COUSSAREA :—
Froelichia, *A. Rich. in Mem. Soc. Hist. Nat. Par.* v. (1834) 177.—Ins. Trinit.
gracillifora, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
grandis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 100.—Bras.
hydrangeaefolia, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
ilheotica, *Muell. Arg. in Flora*, lviii. (1876) 466.—Bras.
lagoënsis, *Muell. Arg. l. c.*—Bras.
lanceolata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 104.—Bras.
leptoloba, *Muell. Arg. in Flora*, lviii. (1876) 468.—Bras.
leptophragma, *Muell. Arg. l. c.* 466.—Bras.
leptopus, *Muell. Arg. l. c.* 465.—Bras.
longiflora, *Muell. Arg. l. c.* 468.—Bras.
longifolia, *Muell. Arg. l. c.*—Bras.
macrocarpa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 177.—Guiana.
macrophylla, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
Martini, *Benth. & Hook. f. Gen.* ii. 121.—Am. trop.
meridionalis, Muell. Arg. in Mart. Fl. Bras. vi. v. 85 = nemoralis.
nemoralis, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
nodosa, *Benth. & Hook. f. l. c.*—Bras.
obscura, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
pentanervia, *Karst. Fl. Columb.* ii. 13. t. 107.—Costa Rica.
platyphylla, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
porophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 87.—Bras.
procumbens, *Muell. Arg. l. c.* 104.—Bras.
racemosa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 177.—Guiana.
Regnelliana, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
Schomburgkiana, Benth. & Hook. f. Gen. ii. 121 = violacea.
squamosa, *Lam. Tabl. Encyc.* i. 281.—Ind. occ.
strigosa, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
trifora, *Muell. Arg. l. c.* 468.—Bras.
unifora, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 107.—Bras.
verticillata, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
violacea, *Aubl. Pl. Gui.* i. 98. t. 38.—Guiana, virens, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
viridis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 104.—Bras.

COUTARREA, Aubl. Pl. Gui. i. 314. t. 122 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 42.
alba, *Griseb. in Gott. Abh.* xxiv. (1879) 158.—Reg. Argent.
Campanilla, *DC. Prod.* iv. 350.—Venezuela.
corymbosa, *Brign. in Mem. Soc. Ital. Mod.* Ser. II. i. (1882) 69.—Bras.
diervilloides, *Planch. & Linden, Cat.* n. 9 (1854) 3.—N. Granat.
flavescens, *Mog. & Sesse, ex DC. Prod.* iv. 350.—Mexic.
latifolia, *Mog. & Sesse, ex DC. l. c.*—Mexic.
Lindeniana, *Baill. Adansonia*, xii. (1879) 300.—Venezuela.
Lemaeana, *Baill. l. c.* 301.—Guatemala.
mexicana, *Zucc. & Mart. ex DC. Prod.* iv. 350.—Mexic.
mollis, *Cham. in Linnaea*, iv. (1834) 359.—Bras.
pubescens, *Pohl, Pl. Bras. Descr.* ii. 146. t. 200.—Bras.
Schottiana, *André, in Illustr. Hortic.* xxv. (1878) 190. t. 321.—N. Granat.
speciosa, *Aubl. Pl. Gui.* i. 314. t. 122.—Guiana.

COUTHOVIA, A. Gray, in Proc. Am. Acad. iv. (1858) 294; v. (1861) 330. *LOGANIACEAE*, Benth. & Hook. f. ii. 797.
corymbosa, *A. Gray, l. c.* iv. (1858) 294.—Ins. Sandvic.
Seemanni, *A. Gray, l. c.* v. (1861) 330.—Ins. Fiji.

COUTOUBEA, Mem. Pl. Ind. Occ. 18 (1825) = seq.

COUTOUBEA, Aubl. Pl. Gui. i. 72. t. 27, 28 (1775) *GENTIANEAE*, Benth. & Hook. f. ii. 812.
LISBORCHIA, Soop. Introd. 139 (1777).
PICRIUM, Schreb. Gen. 791 (1791).
alba, Lam. Encyc. ii. 262 = spicata.
arenaria, Willd. ex Griseb. Gen. et Sp. Gent. 182 = racemosa.
densiflora, Mart. Nov. Gen. et Sp. ii. 111. t. 185 = spicata.
laxa, Spreng. in Flora, xxvi. (1843) 765 = spicata.
minor, H. B. & K. Nov. Gen. et Sp. iii. 179 = spicata.
purpurea, Lam. Encyc. ii. 262 = ramosa.
racemosa, *G. F. W. Mey. Prim. Fl. Esseq.* 86.—Guiana.
ramosa, *Aubl. Pl. Gui.* i. 74. t. 28.—Am. trop.
reflexa, *Benth. in Ann. Nat. Hist.* ii. (1839) 442.—Guiana.
spicata, *Aubl. Pl. Gui.* i. 72.—Am. trop.
spicata, Griseb. in Journ. Linn. Soc. vi. (1862) 141 = reflexa.
spicata, H. B. & K. Nov. Gen. et Sp. iii. 179 = densiflora.
ternifolia, Cav. Ic. iv. 14 = densiflora.
verticillata, G. Don, in Loud. Hort. Brit. 48 = Enicostema littorale.
volubilis, *Mart. Nov. Gen. et Sp.* ii. 112.—Ins. S. Doming.

COUTUBEA, Steud. Nom. ed. II. i. 429 (1840) = Coutoubea, Aubl. (Gentian.).

COVALIA, Reichb. Consp. 94 (1828) = COVOLIA, Neck. = Spermacoce, Linn. (Rubiac.).

COVELIA, Endl. Gen. 527, in syn. (1838) = praec.

COVELLIA, Gasp. Nov. Gen. Fic. 10 (1844) = **Ficus**, Tourn. (Urticac.).
elegia, Miq. Fl. Ind. Bat. Suppl. 434 = F. mollis.
assamica, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 464 = F. hispida.
aurata, Miq. Fl. Ind. Bat. Suppl. 433 = F. chrysocarpa.
Barclayana, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 461. t. 7 B = F. Barclayana.
compacta, Miq. Fl. Ind. Bat. i. II. 324 = F. rapiformis.
congesta, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 468 = F. congesta.
conglomerata, Miq. l. c. 459 = F. Cunia.
costata, Miq. l. c. 468 = F. costigera.
courtallensis, Miq. l. c. 463 = F. hispida.
cunesta, Miq. l. c. 466. t. 8 B = F. cunesta.
Cunia, Miq. l. c. 459 = F. Cunia.
cyrtophylla, Miq. l. c. 460 = F. cyrtophylla.
dasmonum, Miq. l. c. 463 = F. hispida.
dasycarpa, Miq. l. c. 465 = F. hispida.
decipiens, Miq. l. c. 460 = F. obscura.
didymana, Miq. Pl. Jungh. i. 65 = F. lepicarpa.
glomerata, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 465 = F. glomerata.
grandifolia, Miq. Fl. Ind. Bat. Suppl. 434 = F. rapiformis.
Griffithii, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 462 = F. heterophylla.
guttata, Wight, Ic. t. 1966 = F. guttata.
hispida, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 463 = F. hispida.
inaequiloba, Miq. l. c. 459 = F. Cunia.
lanceolata, Miq. l. c. 465 = F. lanceolata.
lepicarpa, Miq. Fl. Ind. Bat. i. II. 326 = F. lepicarpa.
leucophaea, Miq. l. c. 326 = F. leucatoma.
macrophylla, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 465 = F. Roxburghii.
microcarpa, Miq. l. c. 466. t. 9 A = F. Ribes.
mollis, Miq. l. c. 466 = F. glomerata.
oppositifolia, Gasp. Nov. Gen. Fic. 35 = F. hispida.
paniculata, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 467 = F. Ribes.
prostrata, Miq. l. c. 465 = F. prostrata.
racemifera, Miq. l. c. = F. variegata.
radicata, Miq. Fl. Ind. Bat. i. II. 326 = F. septica.
rapiformis, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 464 = F. rapiformis.
Ribes, Miq. Fl. Ind. Bat. i. II. 325 = F. Ribes.
rufescens, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 36 = F. botryocarpa.
setulosa, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 463 = F. hispida.
stictocarpa, Miq. Fl. Jungh. 65 = F. leucantatoma.

COVELLIA :—
subdenticulata, Miq. Fl. Ind. Bat. i. II. 324 = Ficus subdenticulata.
subopposita, Miq. Pl. Jungh. 66 = F. fistulosa.
ternatana, Miq. Fl. Ind. Bat. i. II. 323 = F. ternatana.
tuberculata, Miq. in Zoll. Syst. Verz. Ind. Archip. 94, 99 = F. fistulosa.
umbonata, Miq. Fl. Ind. Bat. i. II. 328 = F. umbonata.
venose, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 468 = F. lascatoma.
wolkenariaefolia, Miq. in Hook. l. c. 464. t. 8 A = F. lepicarpa.
Webbiana, Miq. l. c. 467 = F. Webbiana.
Wightiana, Miq. l. c. 468 = F. hispida.
Zollingeriana, Miq. l. c. 460 = F. coronata.

COVILHAMIA, Korth. in Nederl. Kruidk. Arch. i. (1848) 307. *STERCULIACEAE*?
ovata, *Korth. l. c.*—Borneo.

COVOLA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 227 = **Salvia**, Tourn. (Labiat.).
napifolia, Medic. Phil. Bot. ii. 67 = S. napifolia.
verticillata, Medic. l. c. = S. verticillata.

COVOLIA, Neck. Elem. i. 201 (1790) = **Spermacoce**, Linn. (Rubiac.).

COWANIA, D. Don, in Trans. Linn. Soc. xiv. (1825) 574. t. 22. *ROSACEAE*, Benth. & Hook. f. i. 618.
GERGIA, Engelm. in Bot. Wisliz. Exped. 20 (1848).
ericaefolia, *Torr. ex A. Gray, Pl. Wright.* ii. 106.—Calif.
Havardia, *S. Wats. in Proc. Am. Acad.* xx. (1885) 364.—Texas.
mexicana, *D. Don, in Trans. Linn. Soc.* xiv. (1825) 575. t. 22. f. 1–6.—Am. bor. occ.
plicata, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 400.—Am. bor. occ.
plicata, Torr. in Frém. Rep. 314 = mexicana.
purpurea, Zucc. in Abh. Akad. Muench. iv. ii. (1848) 7. t. 2 = plicata.
Stansburiana, Torr. in Stansb. Exped. Utah, 386 = mexicana.

COXIA, Endl. Gen. 738 (1839) = **Lysimachia**, Linn. (Primulac.).

CRABBEA, Harv. Gen. S. Afr. Pl. 276 (1838) = **Barleria**, Linn. (Acanthac.).
pungens, Harv. l. c. = B. pungens.

CRABBEA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 26. *ACANTHACEAE*, Benth. & Hook. f. ii. 1092.
angustifolia, *Nees, in DC. Prod.* xi. 163.—Afr. austr.
cirsioides, Nees, l. c. 163 = nana.
hirsuta, *Harv. in Hook. Lond. Journ. Bot.* i. (1842) 27.—Afr. austr.
nana, *Nees, in DC. Prod.* xi. 162.—Afr. austr.
ovalifolia, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 34.—Afr. trop.

CRABOWSKIA, G. Don, Gen. Syst. iv. 480 (1836), sphalm. = **Grabowskia**, Schlecht. (Solan.).

CRACCA, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1855) 8. *LEGUMINOSAE*, Benth. & Hook. f. i. 501.
caribaea, *Benth. ex Oerst. l. c.* 9.—Ind. occ.
Edwardsii, *A. Gray, Pl. Wright.* ii. 35.—Mexic.
glabrescens, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1855) 9.—Ecuador.
glandulifera, *Benth. l. c.* 8.—Ecuador.
mollis, *Benth. l. c.* 9.—Peruv.
oehrolenca, *Benth. l. c.*—Peruv.
sericea, *A. Gray, in Proc. Am. Acad.* xix. (1883) 74.—N. Mexic.

CRACCA, Linn. Fl. Zeyl. 139 (1747) = **Tephrosia**, Pers. (1807) (Legumin.).
maxima, Linn. Sp. Pl. ed. I. 752 = T. purpurea.
purpurea, Linn. l. c. = T. purpurea.
senticosa, Linn. l. c. = T. senticosa.
tinctoria, Linn. l. c. = T. tinctoria.
villosa, Linn. l. c. = T. villosa.
virginiana, Linn. l. c. = T. virginiana.

CRACCA, [Riv.] Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359 = **Vicia**, Tourn. (Legum.).
acutifolia, Alef. in Bonplandia, ix. (1861) 118 = Vicia acutifolia.
agrigentina, Alef. l. c. 119 = V. Bivonea.
albicans, Alef. l. c. 120 = V. atropurpurea.
atropurpurea, Gren. & Godr. Fl. Fr. i. 471 = V. atropurpurea.
benghalensis, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359 = V. benghalensis.
Bertaloni, Gren. & Godr. Fl. Fr. i. 470 = V. Pseudocracca.
Bivonea, Alef. in Bonplandia, ix. (1861) 121 = V. Bivonea.
Boissieri, Alef. l. c. = V. villosa.
Broteriana, Alef. l. c. 120 = V. atropurpurea.
calcarata, Gren. & Godr. Fl. Fr. i. 472 = V. calcarata.
carolinana, Alef. in Bonplandia, ix. (1861) 124 = V. carolinana.
cassubica, Opiz, Seznam, 34 = V. cassubica.
ciceroidea, Alef. in Bonplandia, ix. (1861) 121 = V. cicerioides.
cirrhosa, Alef. l. c. 118 = V. cirrhosa.
corsica, Gren. & Godr. Fl. Fr. i. 473 (= *Ervum corsicum*).—Corsica.
dasycarpa, Alef. in Bonplandia, ix. (1861) 121 = V. varia.
disperma, Gren. & Godr. Fl. Fr. i. 472 = V. disperma.
dumetorum, Opiz, Seznam, 34 = V. dumetorum.
elegans, Alef. in Bonplandia, ix. (1861) 124 = V. tenuifolia.
erotanthos, Alef. l. c. 118 = V. Leavenworthii.
exigua, Alef. l. c. 119 = V. exigua.
Gerardi, Gren. & Godr. Fl. Fr. i. 469 = V. Gerardi.
Gerhardi, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359 = V. Gerardi.
hybridesca, Alef. in Bonplandia, ix. (1861) 123 = V. Kotschyana.
Kotschyana, Alef. l. c. 119 = V. Kotschyana.
lusitanica, Alef. l. c. 118 = V. atropurpurea.
longifolia, Alef. l. c. 123 = V. longifolia.
ludoviciana, Alef. l. c. 119 = V. ludoviciana.
major, Alef. l. c. 125 = V. Cracca.
microphylla, Alef. l. c. 121 = V. microphylla.
minor, Gren. & Godr. Fl. Fr. i. 473 = V. hirsuta.
monanthos, Gren. & Godr. l. c. 471 = V. monanthos.
Monardi, Alef. in Bonplandia, ix. (1861) 119 = V. Monardi.
ochroleuca, Alef. l. c. 123 = V. ochroleuca.
pauciflora, Gennari, in Nuov. Giorn. Bot. ii. (1870) 136 = V. pauciflora!
pinata, Alef. in Bonplandia, ix. (1861) 123 = V. pinetorum.
pisiformis, Opiz, Seznam, 34 = V. pisiformis.
Plumnea, Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, iii. (1875–76) 148 = V. Cracca.
polyphylla, Alef. in Bonplandia, ix. (1861) 124 = V. tenuifolia!
Pseudocracca, Alef. l. c. 121 = V. Pseudocracca.
pulchella, Alef. l. c. 119 = V. pulchella.
Sibthorpii, Alef. l. c. 120 = V. Sibthorpii.
sylvatica, Opiz, Seznam, 34 = V. sylvatica.
syriaca, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359.—Syria.
tenuifolia, Opiz, Seznam, 34; Gren. & Godr. Fl. Fr. i. 469 = V. tenuifolia.
tenuissima, Alef. in Bonplandia, ix. (1861) 120 = V. gracilis.
Timbaliana, Debeaux.—Cf. Bull. Soc. Bot. Fr. xxviii. (1881) Bibl. 73 = V. cuneata.
tridentata, Alef. in Bonplandia, ix. (1861) 119, 125 = V. ludoviciana.
varia, Gren. & Godr. Fl. Fr. i. 469 = V. varia.
villosa, Gren. & Godr. l. c. 470 = V. villosa.
vulgaris, Opiz, Seznam, 34 = V. Cracca.

CRACCINA, Stev. in Ball. Soc. Nat. Mosc. iv. (1832) 266 = Astragalus, Tourn. (Legumin.).
austriaca, Stev. l. c. = A. austriacus.
bifidus, Stev. l. c. xxix. (1856) ii. 145 = A. multicaulis.
Cracca, Stev. l. c. (1832) 266 = A. Cracca.
depressa, Stev. l. c. = A. depressus.
erroides, Stev. l. c. xxix. (1856) ii. 145 = A. erroides.
falcata, Stev. l. c. iv. (1832) 266 = A. falcatus.
gracilis, Stev. l. c. = A. gracilis.
leptocaulis, Stev. l. c. xxix. (1856) ii. 145 = A. leptocaulis.
CRACCINA:—
leptostachya, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 266 = Astragalus sulcatus.
leucophaea, Stev. l. c. = A. leucophaeus.
longipes, Stev. l. c. xxix. (1856) ii. 145 = A. macropterus.
multicaulis, Stev. l. c. = A. multicaulis.
puberula, Stev. l. c. = A. puberulus.
racemosa, Stev. l. c. iv. (1832) 266 = A. racemosus.
reptans, Stev. l. c. = A. reptans.
scopaeformis, Stev. l. c. xxix. (1856) ii. 145 = A. tauricus.
sulcata, Stev. l. c. iv. (1832) 266 = A. sulcatus.
taurica, Stev. l. c. = A. tauricus.
triangularis, Stev. l. c. = A. triangularis.
tunetana, Stev. l. c. = A. tunetanus.
vaginata, Stev. l. c. = A. vaginatus.
versicolor, Stev. l. c. = A. versicolor.
vicioides, Stev. l. c. xxix. (1856) ii. 145 = A. macropterus.

CRAEPALIA, Schrank, Baier. Fl. i. 109, 382 (1789) = **Lolium**, Linn. (Gramin.).
temulenta, Schrank, l. c. 382 = L. temulentum.

CRAEPALOPRUMNON, Karst. Fl. Columb. i. 123. t. 61, 62 (1859?) = **Xylosma**, Forst. (Bixin.).
heterophyllum, Karst. l. c. 123. t. 61.—N. Granat.
cordatum, Karst. l. c. 124 = Flacourtia cordata.
obovatum, Karst. l. c. 123. t. 61.—N. Granat.
prunifolium, Karst. l. c. 124 = Flacourtia prunifolia.
racemosum, Karst. l. c. = Flacourtia racemosa.
rubicundum, Karst. l. c. 125. t. 62.—Venezuela.

CRAFORDIA, Rafin. Specch. i. 156 (1814) = **Tephrosia**, Pers.! (Legumin.).
bracteata, Rafin. l. c. = *T. spicata*!

CRAMBE, Tourn. ex Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 98.
amplexicaulis, [Soland. in] Russell, Aleppo, ed. II. ii. 258 = Calepina Corvini.
aspera, Bieb. Fl. Taur. Cauc. ii. 90 = tataria.
Aucheri, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 388 = orientalis.
Bieberstaini, *Yanka, in Termész. Füzetek*, viii. (1884) 36.—Europ.
bursaefolia, L'Hérit. ex DC. Syst. ii. 649 = Calepina Corvini.
Busbaumii, Willd. ex Ledeb. Fl. Ross. i. 292 = orientalis.
caspica, Raeusch. Nom. ed. III. 183 = tataria.
chlorocarpa, *Kit. in Linnaea*, xxxii. (1863) 502.—Austria.
cordata, Willd. Enum. Hort. Berol. Suppl. 43 = cordifolia.
cordifolia, Duf. in Ann. Gén. Sc. Phys. vii. (1820) 308 = glabrata.
cordifolia, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 287.—Reg. Caucas.
Corvini, All. Fl. Pedem. i. 256 = Calepina Corvini.
edentula, *Pisch. & Mey. ex Ker. in Bull. Soc. Nat. Mosc.* (1839) 145, nomen.—Turcoman.
filiformis, Boiss. Voy. Bot. Espagne, 43 = hispanica.
filiformis, *Jacq. Coll. Suppl.* 130.—Patagon.; Chili.
fruticosa, *Linn. f. Suppl.* 289.—Ins. Madera.
gibberosa, *Rupr. Fl. Cauc.* 136, in textu.—Reg. Caucas.
gigantea, Kit. ex Janka, in Oestr. Bot. Zeitschr. xxi. (1871) 95 = maritima.
glabrata, *DC. Prod.* i. 236.—Hispan.
grandiflora, *DC. Syst.* ii. 652.—As. Min.
hispanica, *Linn. Sp. Pl.* 671.—Reg. Mediterr.
juncea, *Bieb. Fl. Taur. Cauc.* iii. 431.—Reg. Cauc. et Himal.
Kotschyana, Boiss. Diagn. Ser. I. vi. 19 = cordifolia.
Kralikii, *Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 496, nomen.—Afr. bor.!
laciniata, Desf. ex Steud. Nom. ed. II. i. 429 = pinnatifida.
lacrimata, Lam. Encyc. ii. 162 = tataria.
laxata, Kit. in Linnaea, xxiii. (1850) 502 = chlorocarpa.
maracandica, Bieb. Fl. Taur. Cauc. ii. 90 = Sobolewskia lithophila.
maritima, *Linn. Sp. Pl.* 671 (Gramin.).—Europ.
orientalis, Jacq. ic. Rar. i. t. 125 = pinnatifida.
orientalis, *Linn. Sp. Pl.* 671.—Ins. Eils.; Persia.
orientalis, Retch. Syst. iii. 220 = tataria.

CRAMBE:—
palmatifida, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 548.—Turkestan.
paniculata, All. Fl. Pedem. i. 256 = Neslia paniculata.
pannonica, Hort. Par. ex Lam. Encyc. ii. 162 = tataria.
persica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 389.—Persia.
pinnatifida, [*F. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 72. —Reg. Caucas.
pontica, *Stev. ex Rupr. Fl. Cauc.* 393.—Tauria.
Pritzelii, *C. Bolle, Ind. Sem. Hort Berol.* (1861) 10.— Ins. Canar.
quadricostata, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 388 = orientalis.
reniformis, Desf. Fl. Atlant. ii. 78. t. 161 = hispanica.
scabra, Lam. Encyc. ii. 162 = strigosa.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) 1. 282.—Turkestan.
sinuato-dentata, *Hochst. & Schimp. Beitr. Fl. Aethiop.* 288.—Abyss.
Steveniana, *Rupr. Fl. Cauc.* 136, in textu.—Reg. Caucas.
strigosa, *L'Hérit. Stirp. Nov.* 151. t. 72.—Ins. Canar.
suecica, Mill. Gard. Dict. ed. VIII. n. 2 = pinnatifida.
tatarica, *Jacq. Misc.* ii. 274.—Europ. or.; As. bor.

CRAMERIA, Murr. Syst. ed. XIII. 138 (1774) = **Krameria**, Linn. (Polygal.).

CRANICHIS, Sw. Prod. Veg. Ind. Occ. 120 (1788) *ORCHIDEAE*, Benth. & Hook. f. iii. 598.
OCAMPOA, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 31.
aphylla, *Sw. l. c.*—Jamaica.
apiculata, *Lindl. in Benth. Pl. Hartw.* 92.—Guatemala.
ciliata, *Kunth, Syn. Pl. Aeq.* i. 324.—Venezuela.
diphylla, *Sw. Prod. Veg. Ind. Occ.* 120.—Jamaica.
Engelii, *Reichb. f. in Linnaea*, xli. (1877) 19.— Venezuela.
foliosa, *Lindl. Gen. et Sp. Orch.* 451.—Peruv.
gibbosa, *Lindl. in Ann. Nat. Hist.* xv. (1845) 385.— Jamaica.
glandulosa, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 30.—Mexic.
Lehmanni, *Reichb. f. Otia Bot. Hamb.* 4.—Ecuador.
luteola, Sw. Fl. Ind. Occ. iii. 1438 = Polystachya luteola
macroblepharis, *Reichb. f. Otia Bot. Hamb.* 4.— Ecuador.
micrantha, *Griseb. in Goett. Abh.* xxiv. (1879) 337.— Reg. Argent.
micrantha, Spreng. Syst. iii. 700 = Prescottia Myosurus
monophylla, *Lindl. Orchid. Linden*, 27.—N. Granat.
multiflora, Ell. ex Nutt. Gen. Am. ii. 191 = Ponthieva glandulosa.
muscosa, *Sw. Prod. Veg. Ind. Occ.* 120.—Jamaic.
nudifolia, *Forst. Syn.* ii. 511.—China.
oligantha, Sw. Prod. Veg. Ind. Occ. 120 = Prescottia Myosurus.
ovata, *Wickstr. in Vet. Akad. Handl. Stockh.* (1837) 73.—Ins. Gandalup.
parvilabris, *Lindl. Orchid. Linden*, 27.—N. Granat.
pauciflora, *Sw. Fl. Ind. Occ.* iii. 1431.—Jamaic.
picta, *Reichb. f. in Linnaea*, xli. (1877) 53.—Ecuador.
reticulata, *Reichb. f. Beitr. Orch. Centr. Am.* 62.— Costa Rica.
Schaffneri, *Reichb. f. in Bonplandia*, iii. (1855) 298.— Mexic.
Schlimii, *Reichb. f. in Linnaea*, xli. (1877) 19.— Venezuela.
similis, *Reichb. f. Otia Bot. Hamb.* 62.—Bras.
spectosa, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 2.—Mexic.
stachyoides, Sw. Prod. Veg. Ind. Occ. 120 = Ponthieva glandulosa.
subumbellata, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 30.—Mexic.
sylvestris, *A. Rich. & Gal. l. c.*—Mexic.
tuberifera, *Griseb. Cat. Pl. Cub.* 228.—Cuba.
tuscia, *Reichb. f. in Flora*, xlviii. (1865) 276.—Ind. occ.
tuberosa, *La Llave & Lex Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 1.—Mexic.
tubulosa, *Lindl. Gen. et Sp Orch.* 451.—Mexic.
ventricosa, *Griseb. Fl. Brit. W. Ind.* 629.—Ind. occ.
Wagneri, *Reichb. f. in Linnaea*, xli. (1877) 19.— Venezuela.

CRANIOLARIA, Linn. Gen. ed. I. 346 (1737). *PEDALINEÆ*, Benth. & Hook. f. ii. 1066.
PETRAMNIA, Raffin. Fl. Tellur. iv. 68 (1836).
annua, *Linn. Sp. Pl.* 618.—N. Granat.
fallax, *A. DC. ex DC. Prod.* ix. 354.—Hab.†
fragrans, Schenk, Ind. Sem. Hort. Wirceb. (1861) = Martynia fragrans.
fruticosa, Linn. Sp. Pl. 618 = Pentarhaphia Craniolaria.
integrifolia, *Chaw. in Linnæa*, vii. (1832) 725.—Bras.
unibracteata, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1826) 67 (= *Proboscidea unibracteata*).—Bras.

CRANIOSPERMUM, Lehm. Asperif. 336; lc. t. 50 (1818). *BORAGINEÆ*, Benth. & Hook. f. ii. 852.
DIPLOLOMA, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 195.
canescens, *DC. Prod.* x. 176.—Sibir.
hirsutum, DC. l. c. = subvillosum.
parviflorum, Decne. in Jacquem. Voy. Bot. 196. t. 130 = Moltkia parviflora.
subvillosum, Bunge, Verz. Alt. 8 = canescens.
subvillosum, *Lehm. Asperif.* 327.—Sibir.

CRANIOTOME, Reichb. lc. Exot. i. 39. t. 54 (1824). *LABIATÆ*, Benth. & Hook. f. ii. 1207.
mauritiensem, Boj. Hort. Maurit. 249 = Anisomeles malabarica.
versicolor, *Reichb. lc. Exot.* i. 39.—Reg. Himal.

CRAWOCARPUS, Benth. in Mart. Fl. Bras. xv. i. 106. t. 28 (1859). *LEGUMINOSÆ*, Benth. & Hook. f. i. 523.
Mertii, *Benth. l. c.*—Bras.

CRANTZIA, Lag. ex DC. Syst. ii. 509, 627, in syn. (1821).
frutescens, Lag. ex DC. l. c. 627 = Moricandia arvensis.
ochroleuca, Lag. ex DC. l. c. 509 = Coringia austriaca.

CRANTZIA, Nutt. Gen. Am. i. 177 (1818). *UMBELLIFERÆ*, Benth. & Hook. f. i. 906.
attenuata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 348 = lineata.
australica, F. Muell. First Gen. Rep. 13, ex Klatt, in Linnæa, xxix. (1857–58) 714 = lineata.
lineata, *Nutt. Gen. Am.* i. 177.—Am. bor. et austr. ; Austral.
Schaffneri, Hemsl. Biol. Centr. Am. Bot. i. 569 = lineata.
Schaffneriana, Schlecht. in Linnæa, xxvi. (1853–55) 870 = lineata.

CRANTZIA, Pohl, in Flora, viii. (1825) 188, nomen (Quid?).

CRANTZIA, Scop. Introd. 173 (1777) = **Alloplectus**, Mart. (Gesneriac.).

CRANTZIA, Sw. Prod. Veg. Ind. Occ. 38 (1788) = **Buxus**, Linn.(Euphorb.).
laevigata, Sw. l. c. = B. laevigata.
laevigata, Vahl, Symb. Bot. ii. 99 = B. Vahlii.
Salandri, Vahl, ex Baill. Mon. Bux. 67 = B. Vahlii.

CRANTZIA, Vell. Fl. Flum. viii. t. 153 (1827) = Centratherum, Cass. (Compos.).
ovata, Vell. l. c.—Bras.

CRANZIA, J. F. Gmel. Syst. 386 (1791) = CRANTZIA, Sw. = **Buxus**, Linn. (Euphorb.).
laevigata, J. F. Gmel. l. c. = B. laevigata.

CRANZIA, Schreb. Gen. i. 148 (1789) = **Toddalia**, Juss. (Rutac.).
aculeata, Steud. Nom. ed. II. i. 480 = T. aculeata.

CRASPEDIA, Forst. f. Prod. 58 (1786). *COMPOSITÆ*, Benth. & Hook. f. ii. 333.
CARTODIUM, Soland. ex R. Br. in Trans. Linn. Soc. xii. (1818) 106.
PYENOSORUS, Benth. in Ennm. Pl. Hueg. 68 (1837).
RICHEA, Labill. Voy. i. 186. t. 16 (1792).
alpina, Banks in Hook. Lond. Journ. Bot. vi. (1847) 159 = uniflora.
chrysantha, *Benth. Fl. Austral.* iii. 580.—Austral.
fimbriata, DC. Prod. vi. 163 = uniflora.
glauca, Spreng. Syst. iii. 441 = C. Richea.
globosa, *Benth. Fl. Australi.* iii. 580.—Austral.

CRASPEDIA :—
gracilis, Hook. f. in Lond. Journ. Bot. vi. (1847) 118 = C. Richea.
macrocephala, Hook. Bot. Mag. t. 3415 = C. Richea.
pilosa, *Sprengel. Syst.* iii. 441.—Austral.
plebeia, Cunn. ex Steud. Nom. ed. II. i. 480 = C. Richea ?
pleiocephala, *F. Muell. in Linnæa*, xxv. (1852) 404. —Austral.
Richea, *Cass. in Dict. Sc. Nat.* xi. 358.—Austral. ; N. Zel.
uniflora, *Forst. f. Prod.* 58.—N. Zel.
viscosa, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 333. —N. Zel.

CRASPEDOLEPIS, Steud. Syn. Pl. Gram. 364 (1855) = **Bestia**, Linn.
Ambrosia, Steud. l. c. = R. Garnottianus.
Verreauxii, Steud. ex Mant. in DC. Monog. Phan. i. 280 = R. Garnottianus.

CRASPEDORHACHIS, Benth. in Hook. Ic. Pl. t. 1377 (1881). *GRAMINEÆ*, Benth. & Hook. f. iii. 1168.
africana, *Benth. l. c.*—Afr. trop.

CRASPEDUM, Lour. Fl. Cochinch. 336 (1790) = **Elæocarpus**, Burm. (Tiliac.).
lectorius, Lour. l. c. = E. lectorius.
tinctorium, Steud. Nom. ed. II. i. 480, sphalm. = præc.

CRASPIDOSPERMUM, Boj. ex DC. Prod. viii. 323 (1844). *APOCYNACEÆ*. Benth. & Hook. f. ii. 695.
verticillatum, *Boj. l. c.*—Madag.

CRASSA, Rupp. Fl. Jen. ed. Hall. 26 (1745) = **Asteoliptas**, Linn.

CRASSINA, Scopin, Schediasma de Acido Veget. 42 (1758) = **Zinnia**, Linn. (Compos.).

CRASSOCEPHALUM, Moench, Meth. 516 (1794) = **Gynura**, Cass. (1825) (Compos.).
cernuum, Moench, l. c. = G. cernua.
flatmense, Hochst. & Steud. ex DC. Prod. vii. 263 = Vernonia cinerea.
flavum, Decne. in Ann. Sc. Nat. Ser. II. ii. (1834) 265 = Senecio Decaisnei.
sonchifolium, Less. Syn. Comp. 395 = Emilia sonchifolia.
valerianæfolium, Less. in Linnæa, v. (1830) 168 = Erechtites valerianæfolia.

CRASSOUVIA, Comm. ex DC. Prod. iii. 395 (1828) = CRASSUVIA, Comm. = **Bryophyllum**, Salisb. (Crassulac.).

CRASSULA, Dill. ex Linn. Syst. ed. I (1735). *CRASSULACEÆ*, Benth. & Hook. f. i. 657.
CRASSULARIA, Hochst. ex Schweinf. Bestr. Fl. Aethiop. 50 (1867).
CURTOGYNE, Haw. Revis. Pl. Succ. 8 (1821).
DASYSTEMON, DC. Prod. iii. 393 (1828).
GLOBULEA, Haw. Syn. Pl. Succ. 60 (1821).
GOMARA, Adans. Fam. ii. 346 (1763).
LAROCHEA, Pers. Syn. i. 387 (1805).
PETROGYTON, Eckl. & Zeyh. Enum. 291 (1837).
TURGOSIA, Haw. in Phil. Mag. iii. (1828) 184.
ROCHEA, DC. Prod. iii. 393 (1828) partim.
SARCOLIPES, Eckl. & Zeyh. Enum. 290 (1837).
SEPTAS, Linn. Pl. Afr. Rar. 10 (1760); Amoen. Acad. vi. 87 (1764).
SPHÆROTIS, Eckl. & Zeyh. Enum. 299 (1837).
TETRAPHYLE, Eckl. & Zeyh. l. c. 292 (1837).
TRIBASTHA, Eckl. & Zeyh. l. c. 302 (1837).
abyssinica, *A. Rich. Tent. Fl. Abyss.* i. 209.—Afr. trop.
acuminata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 349 = corymbulosa.
acutifolia, *Less. Encyc.* ii. 176.—Afr. austr.
adscendens, *Thunb. in Nov. Act. Nat. Cur.* vi. (1778) 241.—Afr. austr.
Agardhiana, *Fenzl. & Schult. Syst.* vii. 718.—Hispan.
alba, *Forsk. Fl. Aegypt. Arab.* 80.—Arabia.
alba, Hort. ex Steud. Nom. ed. II. i. 480 = albiflora.
albicaulis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 353.—Afr. austr.
albiflora, *Sims, Bot. Mag.* t. 2391.—Afr. austr.
alooides, [*Soland. in*] *Ait. Hort. Kew.* ed. i. i. 394.—Afr. austr.

CRASSULA :—
alpestris, *Linn. f. Suppl.* 189.—Afr. austr.
alpina, *Bmdl. in Walp. Rep.* ii. 253.—Afr. austr.
alternifolia, *Linn. Sp. Pl.* 383.—Afr. austr.
andegavensis, DC. Fl. Fr. vi. 522 = Sedum andegavense.
anguina, *Harv. in Harv. & Sond. Fl. Cap.* ii. 348.—Afr. austr.
anthurus, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 338 = perforata.
arborea, *Medic. Beobacht.* (1782) 304.—Afr. austr
arborescens, *Willd. Sp. Pl.* i. 1554.—Afr. austr.
argentea, *Linn. f. Suppl.* 188.—Afr. austr.
articulata, Zuccagni, in Koem. Collect. 136 = portulacea.
ascendens, Harv. in Harv. & Sond. Fl. Cap. ii. 368 = adscendens.
atropurpurea, D. Dietr. Syn. Pl. ii. 1081 = portulacea.
barbata, *Linn. f. Suppl.* 188.—Afr. austr.
bergioides, *Harv. in Harv. & Sond. Fl. Cap.* ii. 352.—Afr. austr.
bibracteata, Eckl. & Zeyh. Enum. 295 = densifolia.
bibracteata, *Haw. in Phil. Mag.* (1824) 186.—Afr. austr.
biconvexa, *Haw. Misc.* 176.—Afr. austr.
bipinnata, *Haw. in. Phil. Mag.* (1824) 186.—Afr. austr.
Bolusii, *Hook. f. Bot. Mag.* t. 8194.—Afr. austr.
bonariensis, Cambess. in St. Hil. Fl. Bras. Mer. iii. 140 = Tillaea pedunculariis.
brachypetala, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 354.—Afr. austr.
brevifolia, *Harv. l. c.* 339.—Afr. austr.
bullata, *Haw Rev. Pl. Succ.* 11.—Afr. austr.
bullulata, *Haw. l. c.*—Afr. austr.
Burmanniana, *D. Dietr. Syn. Pl.* ii. 1082.—Afr. austr.
caerulata, J. F. Gmel. Syst. 518 = crenulata ?
caespitosa, Auct. vide *cespitosa*.
caffra, Linn. Mant. ii. 252 = fruticulosa.
caffraria, Hunnem. ex Link & Otto, Ic. Pl. Sel. i. 40 = corymbulosa.
calicina, Desf. Tabl. ed. II. 167.—Austral.
campestris, Harv. in Harv. & Sond. Fl. Cap. ii. 351 = Tillaea pharnaceoides.
campestris, *Bmdl. in Walp. Rep.* ii. 258.—Afr. austr.
canescens, *Kurm. & Schault. Syst.* vi. 734.—Afr. austr.
capensis, Baill. Hist. Pl. ii. 312 = C. Septas.
capillacea, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 331. —Afr. austr.
capitata, Lam. Encyc. ii. 171 = cymosa.
capitata, *Salm-Dyck, Cat.* (1820) 14.—Afr. austr.
capitella, *Linn. f. Suppl.* 190.—Afr. austr.
capitellata, DC. Prod. iii. 397 = capitella.
capitellata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 344 = lasiantha.
cemaurioides, *Linn. Pl. Afr. Rar.* 9.—Afr. austr.
cephalophora, *Linn. f. Suppl.* 190.—Afr. austr.
cespitosa, Cav. Ic. i. 50 = Sedum caespitosum.
chloraefolia, D. Dietr. Syn. Pl. ii. 1089 = Grammanthes gentianoides.
ciliata, *Linn. Sp. Pl.* 283.—Afr. austr.
clavifolia, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 360. —Afr. austr.
coccinea, Linn. Sp. Pl. 282 = Rochea coccinea.
columnaris, *Linn. f. Suppl.* 191.—Afr. austr.
concinna, *Haw. Rev. Pl. Succ.* 199.—Afr. austr.
concinnella, *Haw. in Phil. Mag.* (1833) 383.—Afr. austr.
connata, Donn, Hort. Cantab. ed. III. 52 = perfossa.
Cooperi, *Regel, Gartenfl.* (1874) 88. t. 788.—Afr. austr.†
corallina, *Linn. f. Suppl.* 188.—Afr. austr.
cordata, Lodd. Bot. Cab. t. 359 = spathulata.
cordata, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 396.—Afr. austr.
cordata, Thunb. Prod. Pl. Cap. 57 = neglecta.
coronata, Heynh. Nom. ii. 165 = perfossa.
corymbulosa, *Link*, Enum. *Hort. Berol.* i. 301.—Afr. austr.
Cotyledon, Haw. Syn. Pl. Succ. 56 = C. Cotyledonis.
Cotyledon, Jacq. Misc. ii. 296 = arborescens.
cotyledonifolia, Salisb. Prod. 309 = arborescens.
Cotyledonis, *Linn. f. Suppl.* 190.—Afr. austr.
crenata, Desf. Choix, 76. t. 58 = Sedum crenatum.
crenulata, *Linn. f. Suppl.* 189.—Afr. austr.
cultrata, *Linn. Sp. Pl.* 283.—Afr. austr.
cymosa, *Berg. Descr. Pl. Cap.* 84.—Afr. austr.
dasyphylla, *Harv. in Harv. & Sond. Fl. Cap.* ii. 365.—Afr. austr.

CRASSULA :—
deblis, *Thunb. Fl. Cap.* 380.—Afr. austr.
decumbens, *Eckl. & Zeyh. Enum.* 303.—Afr. austr.
decumbens, Thunb. Prod. Pl. Cap. 54 = Tillaea decumbens.
decussata, Hart. ex DC. Prod. iii. 398 = falcata.
decussata, Salisb. Prod. 309 = tetragona.
decussata, *Spreng. Syst.* i. 966.—Afr. austr.
dejecta, *Jacq. Hort. Schoenb.* iv. 16. t. 433.—Afr. austr.
deltoides, *Linn. f. Suppl.* 189.—Afr. austr.
densifolia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 340.—Afr. austr.
dentata, *Thunb. Prod. Pl. Cap.* 57.—Afr. austr.
dependens, *Bolus, in Journ. Linn. Soc.* xviii. (1881) 291.—Afr. austr.
diaphana, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 355.—Afr. austr.
dichotoma, Linn. Pl. Afr. Rar. 9 ; Amoen. Acad. vi. 86 = Grammanthes gentianoides.
dichotoma, Linn. f. Suppl. 188 = ramosa.
diffusa, Lam. Fl. Fr. iii. 80 = Sedum caespitosum.
diffusa, [*Soland. in*] *Ait. Hort. Kew.* ed. 1. i. 395.—Afr. austr.
divaricata, *Eckl. & Zeyh. Enum.* 296.—Afr. austr.
Dregeana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 346.—Afr. austr.
Echlonii, D. Dietr. Syn. Pl. ii. 1032 = Grammanthes gentianoides.
ericoides, *Haw. in Phil. Mag.* (1825) 30.—Afr. austr.
exilis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 347.—Afr. austr.
expansa, [*Soland. in*] *Ait. Hort. Kew.* ed. 1. i. 390.—Afr. austr.
falcata, *Wendl. Beobacht.* 44.—Afr. austr.
fascicularis, Lam. Encyc. ii. 171 = Roches fasciularis.
filicaulis, *Haw. in Phil. Mag.* (1824) 186.—Afr. austr.
filiformis, D. Dietr. Syn. Pl. ii. 1029 = Grammanthes gentianoides.
flabelliflia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 357.—Afr. austr.
flava, *Linn. Mant.* i. 60.—Afr. austr.
fruticosa, Mill. Gard. Dict. ed. VIII a. 8 = Euryops tenuissimus.
fruticulosa, *Linn. Mant.* i. 61.—Afr. austr.
furcata, *Endl. in Walp. Rep.* ii. 253.—Afr. austr.
gentianoides, Lam. Encyc. ii. 175 = Grammanthes gentianoides.
glabra, *How. Syn. Pl. Succ.* 58.—Afr. austr.
glabrifolia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 361.—Afr. austr.
globiflora, Spreng. Syst. i. 971 = C. Septas.
globularioides, *Britten, in Oliver, Fl. Trop. Afr.* ii. 389.—Afr. trop.
globulifolia, Stend. Nom. ed. II. i. 481 = Sedum andegavense.
glomerata, *Berg. Descr. Pl. Cap.* 85 ; *Linn. Mant.* i. 60.—Afr. austr.
hemisphaerica, *Thunb. Prod. Pl. Cap.* 57.—Afr. austr.
hirta, *Thunb. Fl. Cap.* 284.—Afr. austr.
hirtipes, *Harv. in Harv. & Sond. Fl. Cap.* ii. 361.—Afr. austr.
hispida, *D. Dietr. Syn. Pl.* ii. 1031.—Afr. austr.
imbricata, *Burm. f. Fl. Cap. Prod.* 8.—Afr. austr.
imbricata, [Soland. in] Ait. Hort. Kew. ed. I. i. 388 = lycopodioides.
impressa, *N. E. Br. in Gard. Chron.* (1879) ii. 238.—Afr. austr.
impressa, *D. Dietr. Syn. Pl.* ii. 1031.—Afr. austr.
inanis, Thunb. Prod. Pl. Cap. 54 = Tillaea capensis.
incana, *Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap.* ii. 355.—Afr. austr.
indica, *Decne. in Jacquem. Voy. Bot.* 61.—Reg. Himal.
interrupta, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 361.—Afr. austr.
jasminea, Ker-Gawl. in Bot. Mag. t. 2178 = Rochea jasminea.
jasminiflora, Haw. in Bot. Mag. sub t. 2178 = Rochea jasminea.
lactea, [*Soland. in*] *Ait. Hort. Kew.* ed. 1. i. 396.—Afr. austr.
lanceolata, *Endl. in Walp. Rep.* ii. 254.—Afr. austr.
lanuginosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 347.—Afr. austr.
lasiantha, *E. Mey. ex Harv. l. c.* 344.—Afr. austr.
lepidosfolia, *Haw. Misc.* 175.—Afr. austr.

CRASSULA :—
lineolata, [*Soland. in*] *Ait. Hort. Kew.* ed. 1. i. 390.—Afr. austr.
Lingua, *D. Dietr. Syn. Pl.* ii. 1031.—Afr. austr.
linguaefolia, *Haw. Misc.* 176.—Afr. austr.
Lingula, *D. Dietr. Syn. Pl.* ii. 1031.—Afr. austr.
littorella, *Endl. in Walp. Rep.* ii. 253.—Afr. austr.
lucida, Lam. Encyc. ii. 173 = spathulata.
lycioides, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 351 = lycopodioides.
lycopodioides, *Lam. Encyc.* ii. 178.—Afr. austr.
Magnolii, DC. Fl. Fr. Suppl. 522 = Sedum caespitosum.
Mannii, Hook. f. in Journ. Linn. Soc. vii. (1864) 198 = abyssinica.
margaritifera, *Harv. in Harv. & Sond. Fl. Cap.* ii. 362.—Afr. austr.
marginalis, [*Soland. in*] *Ait. Hort. Kew.* ed. 1. i. 396.—Afr. austr.
marginata, Thunb. Prod. Pl. Cap. 56 = marginalis.
media, D. Dietr. Syn. Pl. ii. 1033 = Rochea versicolor.
mesembryanthemoides, *D. Dietr. l. c.* 1031.—Afr. austr.
Meyeri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 344.—Afr. austr.
micans, *Vahl, ex Baill. in Bull. Soc. Linn. Par.* i. (1885) 459.—Madag.
microcarpa, *Sibth. & Sm. Fl. Graec. Prod.* i. 217.—Syria.
minima, Harv. in Harv. & Sond. Fl. Cap. ii. 354 = centaurioides.
minima, *Thunb. Prod. Pl. Cap.* 57.—Afr. austr.
mollis, *Linn. f. Suppl.* 189.—Afr. austr.
mongolica, *Franch. Pl. David.* i. (1884) 127.—Mongolia.
monticola, *Linn. f. Suppl.* 189.—Afr. austr.
monticola, *N. E. Br. in Gard. Chron.* (1882) ii. 264.—Afr. austr.
mucronata, Forst. in Comm. Gotting. ix. (1789) 26 = Tillaea moschata.
multicava, *Lem. Illustr. Hortic.* ix. (1861) *Misc.* 40.—Afr. austr.
multiceps, *Haw. in Haw. & Sond. Fl. Cap.* ii. 359.—Afr. austr.
muricata, *Thunb. Prod. Pl. Cap.* 55.—Afr. austr.
muscosa, *Linn. Pl. Afr. Rar.* 10.—Afr. austr.
muscosa, Roth, Enum. Pl. Phan. Germ. i. 994 = Tillaea muscosa.
natans, Thunb. Prod. Pl. Cap. 54 = Tillaea capensis.
neglecta, *Roem. & Schult. Syst.* vi. 722.—Afr. austr.
nemorosa, *Endl. in Walp. Rep.* ii. 253.—Afr. austr.
nivalis, *Endl. i. c.*—Afr. austr.
nudicaulis, *Linn. Sp. Pl.* 388.—Afr. austr.
nummularisefolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 152.—Madag.
obtalcata, Hort. ex Willd. Enum. Hort. Berol. 342 = obtalcata.
obliqua, Haw. in Andr. Bot. Rep. t. 414 = falcata.
obliqua, [Soland. in] Ait. Hort. Kew. ed. 1. i. 393 = portulacea.
obovata, *Haw. Suppl. Pl. Succ.* 17.—Afr. austr.
obtusa, *Haw. l. c.* 16.—Afr. austr.
obvallaria, Hort. ex DC. Prod. iii. 391 = teq.
obvallata, *Linn. Mant.* i. 61.—Afr. austr.
odoratissima, Andr. Bot. Rep. t. 26 = Rochea odoratissima.
orbicularis, *Linn. Sp. Pl.* 388.—Afr. austr.
ovata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 348 = sarmentosa.
pallida, *Baker, in Gard. Chron.* (1874) i. 786.—Afr. austr.
paniculata, *D. Dietr. Syn. Pl.* ii. 1031.—Afr. austr.
parviflora, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 354 = expansa.
parvula, *Endl. in Walp. Rep.* ii. 253.—Afr. austr.
patens, *Endl. l. c.*—Afr. austr.
pellucida, Soq. Fragm. t. 44. f. 2 = centaurioides.
pellucida, *Linn. Sp. Pl.* 388.—Afr. austr.
pepicides, *Harv. in Harv. & Sond. Fl. Cap.* ii. 355.—Afr. austr.
perfoliata, Soop. Delic. Insub. iii. 12 = perfossa.
perfoliata, *Linn. Sp. Pl.* 392.—Afr. austr.
perforata, *Linn. f. Suppl.* 190.—Afr. austr.
perforata, E. Mey. ex Harv. & Sond. Fl. Cap ii. 344 = stachyura.
perfossa, *Lam. Encyc.* ii. 172.—Afr. austr.
perfossa, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 347 = cordata.
Petrogotae, *Walp. Rep.* ii. 262.—Afr. austr.

CRASSULA :—
pharmacoides, C. A. Mey. in Ind. Sem. Hort. Petrop. viii. 56 = Tillaea pharmacoides.
pinnata, Linn. f. Suppl. 191 = Bryophyllum calycinum.
pinnata, *Lour. Fl. Cochinch.* i. 281.—Cochinch.
platyphylla, *Haw. in Haw. & Sond. Fl. Cap.* ii. 368.—Afr. austr.
polpodacea, *Endl. in Walp. Rep.* ii. 253.—Afr. austr.
Portulacaria, Linn. Sp. Pl. ed. II. 606 = Portulacaria afra.
Portulacastrum, Harv. in Harv. & Sond. Fl. Cap. ii. 336 = Portulacaria afra.
portulacea, *Lam. Encyc.* ii. 172.—Afr. austr
profusa, Hook. f. Bot. Mag. t. 6044 = marginalis.
propinqua, *Endl. in Walp. Rep.* ii. 253.—Afr. austr.
prostrata, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 354.—Afr. austr.
prostrata, *Thunb. Prod. Pl. Cap.* 54.—Afr. austr.
pruinosa, *Linn. Mant.* i. 60.—Afr. austr.
pubernla, R. Br. in Salt, Abyss. App. 64 = abyssinica.
pubescens, Endl. in Walp. Rep. ii. 252 = C. Sarcolipes.
pubescens, *Linn. f. Suppl.* 190.—Afr. austr.
pubescens, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 360 = incana.
pulchella, [*Soland. in*] *Ait. Hort. Kew.* ed. 1 i. 392.—Afr. austr.
punctata, *Linn. Syst. Nat.* ed. X. 969.—Afr. austr.
punctata, Mill. Gard. Dict. ed. VIII. n. 7 = perfossa.
pyramidalis, *Linn. f. Suppl.* 189.—Afr. austr.
quadrangula, *Endl. in Walp. Rep.* ii. 253.—Afr. austr.
quadrifida, *Baker, in Saund. Refug. Bot.* t. 298.—Afr. austr.
radicans, *D. Dietr. Syn. Pl.* ii. 1031.—Afr. austr.
ramosa, [*Soland. in*] *Ait. Hort. Kew.* ed. 1. i. 390.—Afr. austr.
ramulifora, *Link, Enum. Hort. Berol.* i. 301.—Afr. austr.
retroflexa, *Ait. Hort. Kew.* ed. II. ii. 194.—Afr. austr.
retroflexa, Linn. f. Suppl. 188 ; Thunb. Prod. Pl. Cap. 53 = Grammanthes gentianoides.
retroflexa, Meerb Pl. Select. t. 16 = falcata.
revolvens, *Haw. in Phil. Mag.* (1824) 188.—Afr. austr.
rosularis, *Haw. Rev. Pl. Succ.* 13.—Afr. austr.
rotundifolia, *Haw. in Phil. Mag.* (1824) 188.—Afr. austr.
rubens, Linn. Syst. Nat. ed. X. 969 = Sedum rubens.
rubicunda, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 341.—Afr. austr.
rubricaulis, *Eckl. & Zeyh. Enum.* 296.—Afr. austr.
rupestris, *Linn. f. Suppl.* 189.—Afr. austr.
sarcocaulis, *Eckl. & Zeyh. Enum.* 295.—Afr. austr.
Sarcolipes, *Haw. in Haw. & Sond. Fl. Cap.* ii. 355.—Afr. austr.
sarmentosa, *Harv. l. c.* 348.—Afr. austr.
Saxifraga, *Haw. l. c.* 357.—Afr. austr.
scabra, *Linn. Sp. Pl.* 388.—Afr. austr.
scabrella, *Haw. Rev. Pl. Succ.* 11.—Afr. austr.
Schimperi, C. A. Mey. Ind. Sem. Hort. Petrop. viii. 56 = Tillaea pentandra.
Schinzii, Roth, Enum. Pl. Phan. Germ. i. 993 = Tillaea aquatica.
sclerantholdes, Burm. f. Fl. Cap. Prod. 8 = glomerata.
Scutellaria, *Burm. f. Fl. Ind.* 78.—Ind. or.
sedifiora, *Endl. in Walp. Rep.* ii. 254.—Afr. austr.
sedifformis, *Schweinf. Beitr. Fl. Aethiop.* 80.—Afr. trop.
sedioides, Mill. Gard. Dict. ed. VIII. n. 9 = orbicularis.
semiorbicularis, *Eckl. & Zeyh. Enum.* 296.—Afr. austr.
Septas, *Thunb. Prod. Pl. Cap.* 57.—Afr. austr.
setacea, *Harv. in Harv. & Sond. Fl. Cap.* ii. 347.—Afr. austr.
sparsa, [*Soland. in*] *Ait. Hort. Kew.* ed. 1. i. 395.—Afr. austr.
spathulata, *Thunb. Prod. Pl. Cap.* 57.—Afr. austr.
Sphaerella, Harv. in Harv. & Sond. Fl. Cap. ii. 339 = cymosa.
spicata, *Linn. f. Suppl.* 189.—Afr. austr.
spinosa, Linn. Mant. ii. 268 = Cotyledon spinosa.
squamosa, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 346 = Dregeana.
squamulosa, *Willd. Enum. Hort. Berol. Suppl.* 15.—Afr. austr.
stachyura, *Eckl. & Zeyh. Enum.* 297.—Afr. austr.
strigosa, Delap. ex Harv. & Sond. Fl. Cap. ii. 347 = lanuginosa.
strigosa, Linn. Fl. Afr. Rar. 10 ; Amoen. Acad. vi. 86 = glomerata ?

CRASSULA :—
subaphylla, *Hovr. in Harv. & Sond. Fl. Cap.* ii. 362. —Afr. austr.
sublacuna, *D. Dietr. Syn. Pl.* ii. 1081.—Afr. austr.
subulata, Eckl. & Zeyh. Enum. 298 = cymosa.
subulata, Harv. in Harv. & Sond. Fl. Cap. ii. 352 = Tillæa subulata.
subulata, *Linn. Syst. Nat.* ed. X. 969.—Afr. austr.
sulcata, *D. Dietr. Syn. Pl.* ii. 1081.—Afr. austr.
sulphurea, *Kunze, in Linnæa*, xvii. (1843) 378.—Afr. austr.
swellingreibiana, Hort. ex DC. Prod. iii. 393 = falcata.
sylvatica, *Licht. in Roem. & Schult. Syst.* vi. 736.— Afr. austr.
tecta, *Linn. f. Suppl.* 190.—Afr. austr.
telephioides, *Haw. Rev. Pl. Succ.* 9.—Afr. austr.
tetragona, *Linn. Sp. Pl.* 353.—Afr. austr.
Thunbergiana, Roem. & Schult. Syst. vi. 738 = Tillæa decumbens.
thyrsiflora, *Linn. f. Suppl.* 190.—Afr. austr.
tomentosa, *Linn. f. l. c.*—Afr. austr.
torquata, *Baker, in Saund. Refug. Bot.* t. 154.—Afr. austr.
trachyantha, *Haw. in Harv. & Sond. Fl. Cap.* ii. 362. —Afr. austr.
turrita, *Thunb. Prod. Pl. Cap.* 56.—Afr. austr.
Umbella, *Pacq. Coll. iv.* 172.—Afr. austr.
umbellata, *Thunb. Prod. Pl. Cap.* 54.—Afr. austr.
undata, DC. Prod. iii. 392, in syn. = dejecta.
undulata, *Haw. Misc.* 176.—Afr. austr.
vaginata, *Eckl. & Zeyh. Enum.* 298.—Afr. austr.
Vaillantii, Roth, Enum. Pl. Phan. Germ. i. 992 = Tillæa Vaillantii.
verticolor, Burch. ex Bot. Reg. t. 320 = Rochea versicolor.
verticillaris, Linn. Syst. Nat. ed. XII. 280 = Sedum caespitosum.
vestita, *Linn. f. Suppl.* 188.—Afr. austr.
virgata, *Haw. in Harv. & Sond. Fl. Cap.* ii. 360.— Afr. austr.
Whiteheadii, *Harv. l. c.* 346.—Afr. austr.
Zeyheri, *Harv. l. c.* 377.—Afr. austr.

CRASSULARIA, Hochst. ex Schweinf. Beitr. Fl. Aethiop. 80 (1867) = Crassula, Dill.
sediformis, Hochst. l. c. = Crassula sediformis.

CRASSUVIA, Comm. ex Lam. Encyc. ii. 141 (1786) = Bryophyllum, Salisb. (1805) (Crassul.).
floripendula, Comm. ex Lam. l. c. = B. calycinum.
floripendula, Sims, Bot. Mag. t. 1409 = præc.

CRATÆGUS, Tourn. ex Linn. Syst. ed. I (1735).
ROSACEÆ, Benth. & Hook. f. i. 626.
PYRACANTHA, M. Roem. Syn. fasc. iii. (Rosifl.) 104, 219 (1847).
PHALACROS, Wenzig, in Linnæa, xxxviii. (1874) 164.
SPORTELLA, Hance, in Journ. Bot. xv. (1877) 207.
TIMBALIA, Clos, in Bull. Soc. Bot. Fr. xviii. (1871) 178.
acerifolia, Mœnch, Vers. Anal. Baeume, 31 = cordata.
acuminata, Desf. ex Steud. Nom. ed. II. i. 431 = Cotoneaster acuminata.
aestivalis, *Torr. & Gray, Fl. N. Am.* i. 468.—Am. bor.
alnifolia, Hort. ex Lavall. Arbor. Segrez. 18 = cuneata.
alnifolia, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1846) 120.—Japon.
alpestris, Hort. ex C. Koch, Dendrol. i. 144 = C. Crus-galli.
alpina, E. F. Gray, Nat. Arr. Brit. Pl. ii. 564 = Pyrus Aria.
alpina, Mill. Gard. Dict. ed. VIII. n. 3 = Pyrus Chamaemespilus.
altaica, Ledeb. ex Loud. Encyc. Trees, 868 = sanguinea.
amblyon, *Becker, in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 13.—Rossia.
Amelanchier, Desf. Hist. Arbr. i. 149 = Amelanchier canadensis.
amenaea, Salisb. Prod. 357 = Amelanchier canadensis.
angustifolia, Benth. in Roem. Arch. i. iii. 86 = C. Crus-galli.
apiifolia, Mœnch, Gmelb. 88 = C. Oxyacantha.
apiifolia, *Michx. Fl. Bor. Am.* i. 287.—Am. bor.
arbuscorum, *Ell. Sketch*, i. 550.—Am. bor.
arbutifolia, Hort. ex Steud. Nom. ed. II. i. 431 = Pyrus Aucuparia.
arbutifolia, [Nayrand. in] Ait. Hort. Kew. ed. II. iii. 268 = Photinia arbutifolia.

CRATÆGUS :—
arbutifolia, Lam. Encyc. i. 83 = Pyrus arbutifolia.
Aria, Linn. Sp. Pl. 475 = Pyrus Aria.
arnoldiana, Sarg. in Bot. Gaz. xvi. (1891) 229.—Mass.
Aronia, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 264 = sinaica.
atrophaea, Stev. ex Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1839) 866 = pentagyna.
atropurpurea, Stev. ex Nym. Consp. 244 = C. Oxyacantha.
Aucuparia, Salisb. Prod. 357 = Pyrus Aucuparia.
ustera, Salisb. l. c. = Pyrus domestica.
axillaris, Pers. Syn. ii. 39 = C. Pyracantha.
Azarolus, Griseb. Spicil. Fl. Rumel. i. 88 = C. Oxyacantha.
Azarolus, *Linn. Sp. Pl.* 477.—As. Min.; Persia.
berberifolia, *Torr. & Gray, Fl. N. Am.* i. 469.—Am. bor.
Bibas, Lour. Fl. Cochinch. 319 = Eriobotrya fragrans.
Boscana, M. Roem. Syn. Rosifl. iii. 119 = C. Crus-galli.
brachyacantha, *Engelm. in Coult. Bot. Gas.* vii. (1882) 128.—Am. bor.
brevispina, Dougl. ex Steud. Nom. ed. II. i. 432 = Douglasii.
brevispina, *Kunze, in Flora*, xxix. (1846) 737.— Hispan.
Calpodendron, Borckh. in Roem. Archiv. i. iii. 87 = tomentosa.
calycina, Peterm. ex Nym. Consp. 244 = C. Oxyacantha.
carolviana, Pers. Syn. ii. 36 = flava.
Celsiana, Bosc, in Nouv. Cours Agr. ii. 223 = nigra.
Chamaemespilus, Jacq. Enum. Vindob. 36; 243 = Pyrus Chamaemespilus.
chinensis, Steud. Nom. ed. II. i. 432 = Raphiolepis indica.
chlorocarpa, C. Koch, Ind. Sem. Hort. Berol. (1856) 17 = coccinea.
chlorosarca, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 30.—Manchur.
Clarkei, *Hook. f. Fl. Brit. Ind.* ii. 383.—Reg. Himal.
coccinea, *Linn. Sp. Pl.* 476.—Am. bor.
coralinus, Hort. Par. ex Pers. Syn. ii. 36 = cordata.
cordata, *Soland. in] Ait. Hort. Kew.* ed. I. ii. 198.— Am. bor.
coriacea, Soland. in Trans. Linn. Soc. xiii. (1822) 104 = Chamaemeles coriacea.
coronaria, Salisb. Prod. 357 = Pyrus coronaria.
Cotoneaster, Borck. in Roem. Arch. i. iii. 87 = Cotoneaster integerrima.
Courretiana, M. Roem. Syn. Rosifl. 117 = C. Crus-galli.
crenulata, *Roxb. Hort. Beng.* 38; *Fl. Ind.* ii. 509.— Reg. Himal.
cretica, Desf. Choix, 79 = Amelanchier vulgaris.
Crus-galli, Bigel. Fl. Bost. ed. II. 194 = coccinea.
Crus-galli, Duroi, Harbk. Baums. ed. I. i. 183 = tomentosa.
Crus-galli, *Linn. Sp. Pl.* 476.—Am. bor.
cuneata, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1846) 120.—Japon.
cuneifolia, Borckh. in Roem. Arch. i. iii. 86 = tomentosa.
cuspidata, Spach, Hist. Vég. Phan. ii. 106 = Pyrus cuspidata.
dentata, Tmtl. Fl. Par. 246 = Pyrus rotundifolia.
depressa, *Presl, Epim. Bot.* 204.—Peruv.
digyna, Pall. ex Ledeb. Fl. Ross. ii. 88 = C. Oxyacantha.
discolor, Borckh. in Roem. Arch. i. iii. 86 = C. Oxyacantha.
diversifolia, Steud. Nom. ed. I. 284 = C. Oxyacantha.
Douglasii, *Lindl. Bot. Reg.* t. 1810.—Am. bor. occ.
dumosa, Salisb. Prod. 357 = C. Pyracantha.
edulis, Bodd. ex Mœnch, Vers. Anal. Baeume, 30 = Pyrus edulis.
elegans, Pers. Encyc. iv. 439 = C. Oxyacantha.
elliptica, Ell. Sketch, i. 549 = æstivalis.
elliptica, [Soland. in] Ait. Hort. Kew. ed. I. ii. 198 = arborescens.
eriocarpa, *Pomel, Nouv. Mat. Fl. Atl.* 157.—Afr. bor.
escallonifolia, *Schlecht. in Linnæa*, xxviii. (1856) 668.—Peruv.
esculea, Salisb. Prod. 357 = Pyrus communis.

CRATÆGUS :—
fennica, Kalm, in Linn. Fl. Suec. ed. II. 167 = Pyrus intermedia.
ferruginea, Pers. Syn. ii. 37 = Eriobotrya cordata.
fissa, Bosc, ex DC. Prod. ii. 633 = C. Oxyacantha.
flabellata, Heldr. ex Regel, in Act. Hort. Petrop. i. (1871) 114 = orientalis.
flabelliformis, Spach, Hist. Vég. Phan. ii. 108 = Pyrus intermedia.
flava, Hook. Fl. Bor. Am. i. 202 (excl. syn.) = tomentosa.
flava, [*Soland. in*] *Ait. Hort. Kew.* ed. II. ii. 19.—Am. bor.
flavescens, *Bosc, in DC. Prod.* ii. 630.—Hab. ?
flexuosa, Poir. Encyc. Suppl. iv. 78 = tomentosa.
flexuosa, Schwein. in Long, Exped. Winnep. ii. 387 = tomentosa ?
florentina, Zuccagni, in Roem. Collect. 142 = Pyrus cratægifolia.
floribunda, C. Koch, Dendrol. i. 149 = coccinea.
Fontanesiana, Steud. Nom. ed. II. i. 432 = C. Crus-galli.
glabra, Thunb. Fl. Jap. 205 = Photinia serrulata.
glandulosa, Georgi, Beschr. Russ. Reich. iii. Nachtr. 278 = C. Oxyacantha.
glandulosa, *Moench, Vers. Anal. Baeume*, 31.—Am. bor.
glandulosa, [Soland. in] Ait. Hort. Kew. ed. I. ii. 199 = flava.
glandulosa, Willd. Sp. Pl. ii. 1002, partim = coccinea.
glauca, Wall. Cat. n. 672 = Strasvæsia glaucescens.
glomerata, *Booth, in Linnæa*, xxiv. (1851) 185.— Hab. ?
grasca, Spach. Hist. Vég. Phan. ii. 102 = Pyrus Aria.
granatensis, *Boiss. Elench.* 41.—Hispan.
grandiflora, *Hort. Belg. ex Hornem. Enum. Pl. Hort. Hafn.* 18.—Hab. ?
grandiflora, *C. Koch, in C. Koch & Fint. Wochenschr.* (1859) 374.—Reg. Caucas.
grata, Salisb. Prod. 357 = Pyrus Malus.
Haleshiehii, Botss. Diagn. Ser. II. ii. 47 = tanacetifolia.
heterophylla, Flueggi. in Ann. Mus. Par. xii. (1808) 428. t. 28 = C. Azarolus.
heterophylla, Stev. Taur. 147 = C. Oxyacantha.
hiemalis, *Lange, in Bot. Tidss.* xiii. (1882–83) 20.— Europ.
hirsuta, Schur, Enum. Pl. Transs. 206 = C. Oxyacantha.
horrida, Medic. Gesch. 84 = C. Crus-galli.
Hostii, Hort. ex Carr. Rev. Hortic. (1873) 470 = Pyrus intermedia.
humulis, Lam. Fl. Fr. iii. 485 = Pyrus Chamaemespilus.
hybrida, Linn. Fauna Suec. ii. App. 557 = Pyrus intermedia.
hypolasia, C. Koch, Dendrol. i. 132 = mexicana.
hyracantha, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. n. 2. p. 248, err. typ. = C. Pyracantha ?
indica, Linn. Sp. Pl. 477 = Raphiolepis indica.
inermis, Linn. Fl. Lapp. 199 = Pyrus intermedia.
Insegnae, *Bertol. Fl. Ital.* vi. 629.—Ital.
integrifolia, Roxb. Fl. Ind. ii. 509 = Cotoneaster microphylla.
intermedia, Fuss, Fl. Transs. 211 = Pyrus intermedia.
kyrtostyla, Fing. ex Schlecht. in Linnæa, iv. (1829) 579 = C. Oxyacantha.
laciniata, Borkh. Handb. Forstb. ii. 1355 = C. Oxyacantha.
laciniata, *Stev. in Bess. Enum. Pl. Volh.* 36.—Tauria.
laciniata, Ucria, in Roem. Arch. i. iii. 69 = tanacetifolia.
laevigata, DC. Prod. ii. 630 = C. Oxyacantha.
laevis, Thunb. Fl. Jap. 204 = Photinia laevis.
lagenaria, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 367 = C. Oxyacantha.
laniocarpa, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 103.—Hispan.
latifolia, Lam. Fl. Fr. iii. 486 = Amelanchier vulgaris.
latifolia, Pers. Syn. ii. 37 = tomentosa.
lavorifolia, Medic. Gesch. 84 = C. Crus-galli.
Lavallei, *Hérincq, ex Lavall. Arbor. Segrez.* 21. t. 7.— Hab. ?
Lennenas, *Warez. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 292.—Hab. ?
leucophloea, Mœnch, Vers. Anal. Baeume, 31 = tomentosa.
linearis, Pers. Syn. ii. 37 = C. Crus-galli.
Lobata, Bosc, Nouv. Cours Agr. ii. 226; et in DC. Prod. ii. 633 = grandiflora.

CRATÆGUS:—
longifolia, DC. Prod. ii. 636, in syn. = Pyrus Aria.
lucida, Ell. Sketch. 549 = acstivalis.
lucida, Mill. Gard. Dict. ed. VIII. n. 6 = C. Crus-galli.
Ludgi, *Carr. in Rev. Hortic.* (1876) 199.—Calif.
lutea, Poir. Encyc. Suppl. iv. 73 = tomentosa.
macracantha, Lodd. in Loud. Arb. Brit. ii. 1819 = coccinea.
macrocarpa, Hegetsch. Fl. Schw. 464 = C. Oxyacantha.
mandshurica, Hort. Bask, ex Maxim. in Bull. Soc. Nat. Mosc. liv. (1879) 1. 20 = chlorosarca.
maroccana, Lindl. Bot. Reg. t. 1855 = sinaica.
maroccana, Pers. Syn. ii. 37 = C. Oxyacantha.
mauva, Linn. f. Suppl. 253 = C. Oxyacantha.
melanocarpa, Bechst. Fl. Taur. Cauc. t. 386 = pentagyna.
melanocarpa, Bory & Chaub. Fl. Pélop. 51 = Heldreichii.
mexicana, *Moç. & Sesse, ex DC. Prod.* ii. 629.—Mexic.
Michauxii, Pers. Syn. ii. 38 = flava.
microcarpa, Lindl. Bot. Reg. t. 1846 = spathulata.
mollis, Scheele, in Linnæa, xxi. (1848) 569 = subvillosa.
monogyna, Jacq. Fl. Austr. iii. 50. t. 292. f. 1 = C. Oxyacantha.
monogyna, Pall. Reise, i. 349 = prænc. ?
multiflora, Hort. ex Lavall. Arbor. Segrez. 57 = heterophylla.
myrtifolia, *Presl, Epim. Bot.* 208.—Peruv.
nigra, Pall. ex Stend. Nom. ed. II. i. 633 = C. Oxyacantha.
nigra, *Waldst. & Kit. Pl. Rar. Hung.* t. 1. 61.—Hungar.
nobilis, S. G. Gmel. Reise, i. 83 = pentagyna.
nudiflora, Nutt. ex Torr. & Gray, Fl. N. Am. i. 468 = acstivalis.
obovatifolia, M. Roem. Syn. Rosifl. 120 = tomentosa.
obtusata, Spach, Hist. Vég. Phan. ii. 104 = Pyrus Aria.
obtusifolia, Pers. Syn. ii. 37 = Osteomeles glabrata.
odorata, Bosc. Cors Nouv. Agr. ii. 321 = tanacetifolia.
odoratissima, Dc. Prod. i. 631 = orientalis.
odoratissima, Hornem. Hort. Hafn. Suppl. 52 = tanacetifolia.
Oliveriana, Bosc, ex DC. Prod. ii. 630 = pentagyna.
opaca, Hook. & Arn. ex Hook. Comp. Bot. Mag. i. (1835) 35 = acstivalis.
orientalis, *Bieb. Fl. Taur. Cauc.* i. 387.—Græcia; As. Min.
ovalifolia, Hornem. Hort. Hafn. Suppl. 52 = C. Crus-galli.
ovalis, *Kit. in Linnæa*, xxxii. (1863) 586 — Europ. Oxyacantha, *Linn. Sp. Pl.* 477.—Europ.; As. temp.
Oxyacantha, Walt. Fl. Carol. 147 = spinifolia.
oxyacanthoides, Thuill. Fl. Par. 245 = C. Oxyacantha.
Pallasii, Griseb. Spicil. Fl. Rumel. i. 89 = pentagyna.
pallida, Salisb. Prod. 357 = Pyrus Aria.
parvifolia, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 169.—Am. bor.
parvifolia, Willd. ex Torr. & Gray, Fl. N. Am. i. 469 = acstivalis ?
pauciflora, Pers. Syn. ii. 37 = C. Pyracantha.
pectinata, *Bosc, ex DC. Prod.* ii. 630.—Reg. Caucas.; Persia.
pentagyna, *Waldst. & Kit. in Willd. Sp. Pl.* ii. 1006. —Europ.
pharmopyrum, Borckh. in Roem. Arch. i. iii. 86 = cordata.
pinnatifida, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 11 = pentagyna.
planatifolia, *Lange, in Bot. Tidsskr.* xiii. (1882–83) 92. —Reg. Caucas.
platyphylla, Lindl. ex Hohen. Enum. Talusch. 190 = pentagyna.
Poiretiana, DC. Prod. ii. 630 = glandulosa.
pomifera, Georgi, It. i. 25 = Pyrus baccata.
pontica, C. Koch, in Verh. Bef. Gartenb. Ver. ii. i. 369 = C. Azarolus.
polyacantha, Jan, ex Nym. Consp. 244 = C. Oxyacantha.
populifolia, Ell. Sketch, i. 553 = coccinea.
populifolia, Walt. Fl. Carol. 147 = cordata.
prunoides, Wendl. in Flora, vi. (1823) 501 = glandulosa.
prunifolia, Bosc, ex DC. Prod. ii. 637 = C. Crus-galli.
prunifolia, Bosc, l. c. = C. Crus-galli.

CRATÆGUS:—
Pseudaria, Spach, Hist. Vég. Phan. ii. 106 = Pyrus Aria.
pubescens, *Stend. Nom.* ed. II. i. 633.—Mexic.
punctata, Jacq. Hort. Vindob. i. 10 = tomentosa.
punicea, Flsch. ex Steud. Nom. ed. II. i. 633 = C. Oxyacantha.
purpurea, Bosc, ex DC. Prod. ii. 638 = sanguinea.
pycnoloba, *Boiss. & Heldr. Diagn.* Ser. II. ii. 46.—Græcia.
Pyracantha, Brandis, For. Fl. 208 = crenulata.
Pyracantha, *Medic. Gesch.* 84.—Europ.
pyracanthifolia, G. Don, in Sweet, Hort. Brit. ed. III. 208 = C. Crus-galli.
pyrifolia, Lam. Encyc. i. 84 = Pyrus angustifolia.
pyrifolia, Moench, Vers. Anal. Baeume, 80 = Pyrus arbutifolia.
pyrifolia, [Soland. in] Ait. Hort. Kew. ed. I. ii. 168 = tomentosa.
quitensis, Benth. Pl. Hartw. 178.—Ecuador.
racemosa, Lam. Encyc. i. 84 = Amelanchier canadensis.
riberaia, Bertol. Misc. Bot. xxii. 14. t. 2 = C. Oxyacantha.
riperia, Salisb. Prod. 357 = Pyrus baccata.
rivularis, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 464.—Am. bor. occ.
rosæformis, Janka, in Oestr. Bot. Zeitschr. xx. (1870) 350 = C. Oxyacantha.
rosea, Hort. ex DC. Prod. ii. 628 = C. Oxyacantha.
rotundifolia, Borckh. in Roem. Arch. i. iii. 87 = glandulosa.
rotundifolia, Lam. Encyc. i. 84 = Amelanchier vulgaris.
rotundifolia, Moench, Vers. Anal. Baeume, 29 = C. Crus-galli.
rubra, Lour. Fl. Cochinch. 290 = Raphiolepis indica.
rubrinervis, *Lange, in Bot. Tidsskr.* xiii. (1882–83) 95.—Hab. ?
ruscinonensis, Gren. & Blanc, in Billotia, i. (1864) 71 = C. Azarolus.
salicifolia, Medic. Bot. Beobacht. 345 = C. Crus-galli.
sanguinea, J. G. Cooper, in Am. Natural. iii. (1870) 452 = Douglasii.
sanguinea, *Pall. Fl. Ross.* i. 25. t. 11.—Rossia; Sibir.
sanguinea, Schrad. Hort. Goett. (1884) = orientalis.
sanguinea, Torr. & Gray, Fl. N. Am. i. 464 = coccinea.
scandica, Wahlenb. Fl. Upsal. 165 = Pyrus intermedia.
Schraderiana, Ledeb. Fl. Ross. ii. 91 = orientalis.
semiglabata, Bluff & Fing. Comp. Fl. Germ. i. 613, 1300 = Pyrus rotundifolia.
serotina, Salisb. Prod. 357 = Pyrus baccata.
serrata, Poir. Encyc. Suppl. i. 292 = Photinia arbutifolia.
serratifolia, Desf. Cat. Hort. Par. ed. III. 408 = Photinia serrulata.
serrulata, Poir. Encyc. Suppl. i. 292 = Photinia arbutifolia.
Shicola, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 238 = Eriobotrya Japonica.
Sieboldi, Steud. Nom. ed. II. i. 634 = Raphiolepis japonica.
sinaica, *Boiss. Diagn.* Ser. II. ii. 48.—Arab.; Persia.
sinensis, Loisel. Herb. Amat. iv. t. 247 = Raphiolepis indica.
songorica, *Regel, Dilect. Sem. Hort. Petrop.* (1876) 36, Iconn.—Turkestan.
sorbifolia, *Desf. Cat. Hort. Par.* ed. III. 408.—Hab.?
sorbifolia, Hort. ex Poir. Encyc. Suppl. iv. 78 = Pyrus aparia.
spathulata, Hort. ex Lavall. Arbor. Segrez. 18 = cuneata.
spathulata, *Michx. Fl. Bor. Am.* i. 288.—Am. bor.
spathulata, Pursh, Fl. Am. Sept. i. 326 = flava.
spectabilis, Carr. in Rev. Hortic. (1872) 471 = C. Oxyacantha.
spectabilis, Salisb. Prod. 357 = Pyrus spectabilis.
spicata, Lam. Encyc. i. 84 = Amelanchier canadensis.
spinosa, Gilib. Fl. Lituan. ii. 321 = C. Oxyacantha.
spinosissima, Lodd. Bot. Cab. t. 1100 = sanguinea.
spinulosa, Steud. Nom. ed. II. i. 634 = Rhaphiolepis indica.
stipulosa, *Steud. l. c.*—Ecuador.
subflava, Lodd. ex Nordm. in Bull. Sc. Acad. Pétersb. ii. (1837) 218 = Pyrus subflava.
subserrata, *Benth. Pl. Hartw.* 10.—Mexic.
subspinosa, *DC. Prod.* ii. 626.—Chili.
sulbellina, *Schrad. ex Torr. & Gray, Fl. N. Am.* i. 466.—As. or.; Am. bor.

CRATÆGUS:—
succulenta, Schrad. ex *Link*, Handb. ii. 78 = sanguinea.
tanacetifolia, *Pers. Syn.* ii. 36.—Cappadoc.
tanacetifolia, Webb, It. Hisp. 48 = lacinista.
terminalis, Gouan, Fl. Monsp. 253 = Pyrus torminalis.
texana, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 454 = subvillosa.
tomentosa, *Linn. Sp. Pl.* 476 (syn. Gronov.).—Am bor.
tomentosa, Michx. Fl. Bor. Am. i. 289 = parvifolia.
torminalis, Linn. Sp. Pl. 476 = Pyrus torminalis.
Tournefortii, Griseb. Spicil. Fl. Rumel. i. 90 = orientalis.
transylvanica, Hort. ex M. Roem. Syn. Rosifl. 107 = C. Oxyacantha.
triflolata, *DC. Prod.* ii. 630.—Am. bor.
trilobe, Polr. Voy. Barb. ii. 176 = C. Azarolus.
trilobata, Labill. Ic. Pl. Syr. Dec. iv. 15 = Pyrus trilobata.
turbinata, Pursh, Fl. Am. Sept. ii. 735 = flava.
umbellata, *Desf. Cat. Hort. Par.* ed. III. 408.—Hab.?
uniflora, Durol, Harbk. Baume. i. 184 = parvifolia.
uniflora, Muenchh. Hausw. v. 147 = tomentosa.
uniistralis, Pers. Syn. ii. 37 = parvifolia.
vernicolor, Rochel, Pl. Rar. Banat. ii. [5] = nigra.
villosa, Thunb. Fl. Jap. 204 = Photinia villosa.
virginica, Lodd. in Loud. Arb. Brit. iii. 1832 = flava.
viridis, Linn. Sp. Pl. 476 = coccinea.
viridis, Medic. Gesch. 85 = C. Crus-galli.
viridis, Walt. Fl. Carol. 147 = flava.
Watsoniana, Steud. Nom. ed. II. i. 634 = C. Crus-galli.
xanthocarpa, Medic. Gesch. 85 = tomentosa.

CRATÆVA, Linn. Syst. ed. I (1735); ed. X. 1044 (1759). *CAPPARIDEÆ*, Benth. & Hook. f. i. 110.
BELOU, Adans. Fam. ii. 408 (1763).
FARQUHARIA, Hilsenb. & Boj. ex Boj. in Ann. Sc. Nat. Sér. II. xx. 1 (1843) 51.
NEVOSMILA, Rafin. Sylva Tellur. 107 (1838).
OTHRYS, Noronha, ex Thou. Nov. Gen. Madag. 18 (1806).
TAPIA, Adans. Fam. ii. 407 (1763).
acuminata, DC. Prod. i. 243 = C. Tapia.
acuminata, Miq. Illustr. Fl. Arch. Ind. 21, in obs. — gynandra.
Adansonii, DC. Prod. i. 243 = religiosa.
apetala, Spreng. Syst. ii. 448 = Niebuhria linearis.
arborea, Schrad. in Goett. Gel. Anz. (1821) 707 = C. Tapia.
avicularis, Burch. ex DC. Prod. i. 244 = Niebuhria triphylla.
Balongas, Koen. in Trans. Linn. Soc. v. (1800) 224 = Feronia Elephantum.
Benthamii, *Eichl. in Mart. Fl. Bras.* xiii. i. 265.—Bras.
Brownii, Korth. ex Miq. Illustr. Fl. Arch. Ind. 21 = membranifolia.
caffra, Burch. ex DC. Prod. i. 243 = Niebuhria triphylla.
capparoides, Andr. Bot. Rep. t. 176 = fragrans.
excelsa, *Boj. Hort. Maurit.* 16.—Madag.
falcata, *DC. Prod.* i. 243.—China.
fragrans, *Wenz, Bot. Mag.* t. 596.—Afr. trop.
Gerardena, Vell. Fl. Flum. 200; v. t. 4 = Gallesia scorododendrum.
Gerardena, Mocq. in DC. Prod. xiii. ii. 8 = prænc.
Gerrardena, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 466.—Madag.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 240 = religiosa.
gynandra, *Linn. Sp. Pl.* ed. II. 636.—Am. trop.
hygrophilæ, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) 11. 292.—Ind. or.
lanta, Dc. Prod. i. 243 = religiosa.
lophosperma, *Kurz, in Journ. Bot.* xii. (1874) 195.—Burma.
macrocarpa, *Kurz, l. c.*—Malaya.
magna, *DC. Prod.* i. 243.—Cochinch.
Marmelos, Linn. Sp. Pl. 444 = Ægle Marmelos.
membranifolia, *Miq. Fl. Ind. Bat. Suppl.* i. 397.—Sumatra.
meschata, Banks, ex Sims, Bot. Mag. xiv. t. 504 = fragrans.
Narvala, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 131 = religiosa.

CRATAEVA:—
oblongifolia, Spreng. Syst. ii. 448 = Niebuhria oblongifolia.
obovata, *Vahl, Symb. Bot.* iii. 61.—Madag.
octandra, Blanco, Fl. Filip. ed. I. 400 = Capparis floribunda.
odora, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 118 = religiosa.
radiatiflora, *DC. Prod.* i. 243.—Ecuador.
religiosa, Ainslie, Mat. Med. Hind. 134 = Aegle Marmelos.
religiosa, *Forst. f. Prod.* 85.—Geront. trop.
Roxburghii, R. Br. in Denh. & Clapp. Trav. App. 224 = religiosa.
sarcocarpa, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 466.—Madag.
Tapia, Burm. f. Fl. Ind. 109 = religiosa.
Tapia, Heyne, ex Wall. Cat. sub n. 6978 = Cadaba trifoliata.
Tapia, *Linn. Sp. Pl.* 444.—Am. austr.
Tapia, Vahl, Symb. Bot. iii. 61 = religiosa.
tapioides, *DC. Prod.* i. 243.—Am. austr.
tumulorum, *Miq. Illustr. Fl. Arch. Ind.* 21. t. 11.—Java.
umbellularis, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 121 = religiosa.
Vallenga, Koen. ex Wight & Arn. Prod. 96 = Feronia Elephantum.

CRATEOLA, Rafin. Fl. Tellur. iv. 84 (1836) = **Anthacanthus**, Nees (Acanthac.).
parviflora, Rafin. l. c. = Hemigraphis elegans!
vincoides, Rafin. l. c. = Anthacanthus vincoides.

CRATERIA, Pers. Syn. i. 485 (1805) = **Casearia**, Jacq. (Samydac.).
capitata, Pers. l. c. = Cas. sylvestris.
fasciculata, Pers. l. c. = Cas. punctata.

CRATERICARPIUM, Spach. in Nouv. Ann. Mus. Par. iv. (1835) 227, 237 = **Cenothera**, Linn. (Onagrar.).
argyrophyllum, Spach, l. c. 238 = O. subulata.

CRATERISPERMUM, Benth. in Hook. Niger Fl. 411 (1849). *RUBIACEAE*, Benth. & Hook. f. ii. 112.
brachymeristum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 161.—Afr. trop.
cerinanthum, *Hiern, l. c.*—Afr. trop.
laurinum, *Benth. in Hook. Niger Fl.* 411.—Afr. trop.
microdon, *Baker, Fl. Maurit.* 145.—Ins. Seychelles.
montanum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 162.—Afr. trop.
Schweinfurthii, *Hiern, l. c.* 161.—Afr. trop.

CRATERITECOMA, Lindl. Veg. Kingd. 677 (1847) = CRATEROTECOMA, Mart. = Lundia, DC. (Bignon.).

CRATERORSTIGMA, Hochst. in Flora, xxiv. (1841) 668. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 954.
auriculatifolium, Benth. & Hook. f. Gen. ii. 954 = pumilum.
plantagineum, *Hochst. in Flora*, xxiv. (1841) 669.—Abyss.
pumilum, *Hochst. l. c.* 670.—Abyss.

CRATEROTECOMA, Mart. ex DC. Prod. ix. 215 (1845) = Lundia, DC. (Bignon.).
parviflora, DC. l. c.—Bras.

CRATEVA, Linn. Syst. ed. I (1735) = Crataeva, Linn. (Capparr.).

CRATOCHWILIA, Neck. Elem. ii. 389 (1790) = Claytia, Boerh. (Euphorb.).

CRATOXYLON, Blume, Bijdr. 148 (1825). *HYPERICINEAE*, Benth. & Hook. f. i. 266.
ANCISTROLOBUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 323. t. 6.
BLUMEA, Jack, in Malay. Misc. ii. (1822) n. VII. 21.
HORSFIELDIA, Blume, Cat. Gew. Buitenz. 15 (1823).
TRIPETERIS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 351. t. 9.
arborescens, Blume, Mus. Bot. Lugd. Bat. ii. 17 = Blumea.
[illegible]

CRATOXYLON:—
Blancoi, *Blume, Mus. Bot. Lugd. Bat.* ii. 17.—Ins. Philipp.
biflorum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 580 = polyanthum.
carneum, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 64 = polyanthum.
celebicum, *Blume, Mus. Bot. Lugd Bat.* ii. 16.—Ins. Celebes.
clandestinum, *Blume, l. c.* 15.—Java.
cochinchinense, Blume, l. c. 17 = polyanthum.
connatum, *Miq. Fl. Ind. Bat.* i. ii. 517.—Malaya.
floribundum, *F. Vill. in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 16.—Ins. Philipp.
formosum, *Benth. & Hook. f. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 258.—Malaya.
glaucum, *Korth. in Flora*, xxxi. (1848) 579.—Borneo.
Harmandii, *Pierre, Fl. For. Cochinch.* t. 58.—Cambod. Hornschuchii, *Blume, Bijdr.* 148.—Java.
lanceolatum, Miq. Fl. Ind. Bat. Suppl. i. 500 = polyanthum.
ligustrinum, Blume, Mus. Bot. Lugd. Bat. ii. 16 = polyanthum.
Maingayi, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 258.—Malaya.
microphyllum, *Miq. Fl. Ind. Bat. Suppl.* i. 500.—Ins. Banca.
myrtifolium, *Blume, Mus. Bot. Lugd. Bat.* ii. 17.—Borneo.
nervifolium, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 238.—Burma.
pentadelphum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 580.—Java.
petiolatum, Blume, Mus. Bot. Lugd. Bat. ii. 17 = polyanthum.
polyanthum, *Korth. in Flora*, xxxi. (1848) 579.—Malaya.
polyneurum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 579.—Sumatra.
pruniflorum, *Kurs, For. Fl. Brit. Burma*, i. 84.—Burma.
pruniflorum, Dyer, in Hook. f. Fl. Brit. Ind. i. 258 = praec.
racemosum, *Blume, Mus. Bot. Lugd. Bat.* ii. 16.—Java.
sumatranum, *Blume, l. c.*—Sumatra.
Wightii, Blume, l. c. 18 = polyanthum.

CRATYLIA, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 131. *LEGUMINOSAE*, Benth. & Hook. f. i. 536.
Desvauxii, *Tul. in Arch. Mus Par.* iv. (1841) 57.—Bras.
floribunda, *Benth. in Ann. Wien. Mus.* ii. (1838) 132.—Bras.
hypargyrea, *Mart. ex Benth. l. c.* 131.—Bras.
mollis, *Mart. ex Benth. l. c.* 132.—Bras.
nitens, Benth. l. c. = floribunda.
nuda, *Tul. in Arch. Mus. Par.* iv. (1841) 68.—Bras.
spectabilis, Tul. l. c. 69 = Camptosema grandiflorum.

CRAWFURDIA, Wall. Tent. Fl. Nep. 68. tt. 47, 48 (1826). *GENTIANEAE*, Benth. & Hook. f. ii. 815.
GOLDENIA, Maxim. in Bull. Acad. Pétersb. iv. (1862) 252.
PTERYGOCALYX, Maxim. Prim. Fl. Amur. 198. t. 9 (1859).
TRIPTEROSPERMUM, Blume, Bijdr. 849 (1825).
affinis, *Wall. Cat.* n. 4370.—Reg. Himal.
angustata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 106.—Reg. Himal.
Blumei, *G. Don, Gen. Syst.* iv. 200.—Java.
campanulacea, *Wall. & Griff. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 106.—Reg. Himal.
fasciculata, Griff. Notul. iv. 92 = japonica.
fasciculata, Wall. Cat. n. 4369 A = affinis.
fascicularis, *Wall. Tent. Fl. Nep.* t. 47.—Reg. Himal.; Ceylon.
japonica, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. 111. (1846) 160.—Reg. Himal.; China; Japon.
Loureiri, *G. Don, Gen. Syst.* iv. 200.—Cochinch.
luteo-viridis, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 448 = japonica.
paniculata, Griseb. Gen. et Sp. Gent. 306 = Pnederia fortida.
puberula, *C. B. Clarke, in Journ. Linn. Soc.* xiv. (1875) 443.—Reg. Himal.
speciosa, *Wall. Cat.* n. 4371.—Reg. Himal.
trimervis, D. Dietr. Syn. Pl. ii. 919 = Blumei.

CREAGHIA, Scortech. in Journ. Bot. xxii. (1884) 369. *RUBIACEAE*.
fagraeoides, *Scortech. l. c.* 370.—Malaya.

CREMANIUM, D. Don, in Mem. Wern. Soc. iv. (1823) 310 = **Miconia**, Ruiz & Pav. (Melastomac.).
ambiguum, Benth. Fl. Hartw. 181 = M. millefiora.
amygdalinum, Griseb. Fl. Brit. W. Ind. 261 = M. integrifolia.
Aschenbornianum, S. Schau. in Linnaea, xx. (1847) 734 = Oxymeris comiodes.
aspergillare, DC. Prod. iii. 198 = M. aspergillaris.
axillare, Macfad. Fl. Jamaic. ii. 92 = M. integrifolia.
Herpestianum, S. Schau. in Linnaea, xx. (1847) 735 = M. Berghesianum.
calophyllum, D. Don, in Mem. Wern. Soc. iv. (1823) 312 = M. myrtifolia.
caracasanum, Triana, in Trans. Linn. Soc. xxviii. (1871) 119 = M. rubiginosa.
chaenopleurodes, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 249 = M. late-crenata.
cinnamomifolium, DC. Prod. iii. 194 = M. Candolleana.
ciathranthum, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 129 = M. clathrantha.
coriaceum, D. Don, in Mem. Wern. Soc. iv. (1823) 312 = M. coelestis.
coriaceum, D. Don, l. c. = M. caerulea.
compressum, Benth. Bot. Voy. Sulph. 96 = M. argentea.
cordifolium, Bunb. in Proc. Linn. Soc. i. (1841) 109.—Bras.
croceum, DC. Prod. iii. 194 = M. croceo.
Elaeodendrum, DC. l. c. 191 = M. Elaeodendrum.
glaberrimum, DC. l. c. 192 = M. millefiora.
glandulosum, DC. l. c. 191 = Tetrazygia bicpida.
granulosum, DC. l. c. 192 = M. granulosa.
hispidissimum, DC. l. c. = M. hispidissima.
inaequidens, DC. l. c. 194 = M. inaequidens.
integrifolium, Macfad. Fl. Jamaic. ii. 93 = M. multispicata.
laterenatum, DC. Prod. iii. 194 = M. laterenata.
latifolium, D. Don, in Mem. Wern. Soc. iv. (1823) 311 = M. latifolia.
laurinum, D. Don, l. c. 312 = M. laurina.
ledifolium, DC. Prod. iii. 193 = M. ledifolia.
ligustroides, DC. l. c. 194 = M. ligustroides.
meaium, D. Don, in Mem. Wern. Soc. iv. (1823) 312 = M. media.
millefiorum, DC. Prod. iii. 193 = M. millefiora.
minutiflorum, DC. l. c. = M. millefiora.
nitidum, D. Don, in Mem. Wern. Soc. iv. (1823) 311 = M. nitida.
nitidum, Griseb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 130 = M. andina.
oligotrichum, DC. Prod. iii. 193 = M. oligotricha.
paludosum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 247 = M. paludosa.
paniculatum, DC. Prod. iv. 799 = M. paniculata.
paniculatum, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 118 = M. minutiflora.
papillosum, DC. Prod. iii. 192 = M. papillosa.
pulverulentum, D. Don, ex Triana, l. c. 126 = M. pulverulenta.
pulverulentum, Triana, (1871) in Trans. Linn. Soc. xxviii. 127 = M. Tiri.
purpurascens, Griseb. ex Triana, l. c. 140 = Mecranium virgatum.
pusilliflorum, DC. Prod. iii. 194 = M. Gardneriana.
pusillum, Cham. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 117 = M. depauperata.
quadrangulare, DC. Prod. iii. 192 = M. quadrangularis.
racemosum, Griseb. Cat. Pl. Cub. 102 = Mecranium integrifolium.
reclinatum, DC. Prod. iii. 192 = M. reclinata.
rigidum, Macfad. Fl. Jamaic. ii. 93 = M. rigida.
Riocarium, Griseb. Fl. Brit. W. Ind. 262 = M. trichotoma.
rotundifolium, D. Don, in Mem. Wern. Soc. iv. (1823) 311 = M. rotundifolia.
rubens, DC. Prod. iii. 191 = M. rubens.
rubens, Griseb. Cat. Pl. Cub. 103 = M. minutiflora.
rubens, Griseb. Fl. Brit. W. Ind. 261 = M. millefiora.
Sellowianum, Cham. ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 206 = M. Sellowiana.
serrulatum, D. Don, in Mem. Wern. Soc. iv. (1823) 312 = M. galeazzoides.
Sieberii, Griseb. Fl. Brit. W. Ind. 262 = M. vulcanica.

CREMANIUM :—
splendens, Macfad. Fl. Jamaic. ii. 94 = Miconia obovalis!
tetrandrum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 186 = M. tetrandra.
theezans, DC. Prod. iii. 194 = M. millefiora.
thyrsiflorum, Triana, in Trans. Linn. Soc. xxviii. (1871) 180 = M. thyrsiflora.
thyrsoideum, D. Don, in Mem. Wern. Soc. iv. (1823) 311 = M. thyrsiflora.
tinctorium, DC. Prod. iii. 193 = M. media.
trinerve, Macfad. Fl. Jamaic. ii. 94 = M. triplinervis.
Trichtatis, Cuveg. in Linnaea, xx. (1847) 111 = M. minutiflora.
vaccinioides, D. Don, in Mem. Wern. Soc. iv. (1823) 312 = M. vaccinioides.
virgatum, Griseb. Cat. Pl. Cub. 102 = Mecranium haemanthum.
virgatum, Griseb. Fl. Brit. W. Ind. 261 = Mecranium virgatum.

CREMANTHODIUM, Benth. in Hook. Ic. Pl. t. 1141, 1142 (1878). *COMPOSITAE*, Benth. & Hook. f. ii. 489.
Decaisnei, *C. B. Clarke, Comp. Ind.* 168.—Reg. Himal.
discoideum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 482.—China.
Hookeri, *C. B. Clarke, Comp. Ind.* 169.—Reg. Himal.
humile, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 481.—China.
linearе, *Maxim. l. c.* 482.—China.
oblongatum, *C. B. Clarke, Comp. Ind.* 168.—Reg. Himal.
palmatum, *Benth. in Hook. Ic. Pl.* t. 1149.—Reg. Himal.
planatifidum, *Benth. l. c.*—Reg. Himal.
plantaginemm, *Maximow. in Bull. Acad. Pétersb.* xxvii. (1881) 481.—China.
ramiforme, *Benth. in Hook. Ic. Pl.* t. 1141.—Reg. Himal. Thomsoni, *C. B. Clarke, Comp. Ind.* 169.—Reg. Himal.

CREMASPORA, Benth. in Hook. Niger Fl. 412 (1849). *RUBIACEAE*, Benth. & Hook. f. ii. 108.
SCHIZOSPERMUM, Bolv. ex Baill. Adansonia, xii. (1879) 295.
africana, *Benth. l. c.*—Afr. trop.
Bocandeana, *Webb, in Hook. New Journ.* ii. (1850) 371. —Ins. Gorgona.
comorensis, *Baill. Adansonia*, xii. (1879) 234.—Madag.
congesta, *Baill. l. c.* 283.—Ins. Comor.
grandis, *Baill. l. c.* 235.—Madag.
heterophylla, F. Didr. in Kjoeb. Vidensk. Meddel. (1854) 186. n. 5 = africana.
maxima, *Baill. Adansonia*, xii. (1879) 236.—Madag. Thomsoni, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 126.— Afr. trop.
tubulosa, *Baill. Adansonia*, xii. (1879) 234.—Madag.

CREMASTOSTEMON, Jacq. Fragm. 68 (1809) = Olinia, Thunb. (Lythrar.).
capensis, Jacq. l. c. = O. capensis.

CREMASTRA, Lindl. Gen. et Sp. Orch. 172 (1833). *ORCHIDEAE*, Benth. & Hook. f. iii. 538.
DACTYLOSTALIX, Reichb. f. in Bot. Zeit. xxxvi. (1878) 74.
HYACINTHORCHIS, Blume, Mus. Bot. Lugd. Bat. i. 48. t. 16 (1849).
mitrata, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 411 = Wallichiana.
Wallichiana, *Lindl. Gen. et Sp. Orch.* 173 ; *et in Wall. Cat.* n. 7849.—Reg. Himal. ; Japon.

CREMASTUS, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 187 = Bignonia, Linn.
angustifolius, Miers, l. c. = B. glutinosa.
Balbisianus, Miers, l. c. = B. Balbisiana.
coriaceus, Miers, l. c. nomen = B. Sellei ?
crassus, Miers, l. c. nomen.—Bras.
glutinosus, Miers, l. c. = B. glutinosa.
grallatorius, Miers, l. c.—Bras.
longipes, Miers, l. c. nomen.—Bras.
loricatus, Miers, l. c. 188, nomen.—Bras.
luridus, Miers, l. c. 187, nomen.—Bras.
sarcosus, Miers, l. c. 188, nomen = Delostoma nervosum ?
planifolius, Miers, l. c. nomen.—Bras.
trichocladus, Miers, l. c. 187, nomen = B. trichoclada.

CREMATOMIA, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. iii. (1869) 301 = Bourreria, P. Br. (Boragin.).
Andricusii, Miers, l. c. 324 = B. Andricusii.
attenuata, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. iii. (1869) 307 = B. domingensis ?
calophylla, Miers, l. c. 306 = B. calophylla.
coriacea, Miers, l. c. 307.—Ind. occ.
cumanensis, Miers, l. c. 302.—Venezuela.
elongata, Miers, l. c. 308.—Jamaic.
fasciculata, Miers, l. c. 313.—Venezuela.
formosa, Miers, l. c. 308 = B. formosa.
grandiflora, Miers, l. c. 308 = B. grandiflora.
guatemalensis, Miers, l. c. = B. grandiflora.
Guildingiana, Miers, l. c. 304.—Ind. occ.
Huanita, Miers, l. c. 313 = B. Huanita.
Yacquiniana, Miers, l. c. 306.—N. Granat.
Kunthiana, Miers, l. c. 311.—Venezuela.
mollisacula, Miers, l. c. 309.—Ind. occ.
revoluta, Miers, l. c. = B. revoluta.
spathulata, Miers, l. c. 310 = B. spathulata.
velutina, Miers, l. c.—Ind. occ.
venosa, Miers, l. c. 305.—Ind. occ.

CREMOBOTRYS, Beer, in Flora, xxxvii. (1854) 848 = Rillbergia, Thunb. (Bromeliac.).
zebrina, Beer, l. c. = B. zebrina.

CREMOCEPHALUM, Miq. Fl. Ind. Bat. ii. 97 (1856) = seq.

CREMOCEPHALUM, Cass. in Dict. Sc. Nat. xxxiv. 390 (1824) = Gynura, Cass. (Compos.).
cernuum, Cass. l. c. = G. cernua.

CREMOCHILUS, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 174 = Stiphocampylus, Pohl (Campanulac.).
meridensis, Turcz. l. c.—Venezuela.

CREMOLOBUS, DC. Syst. iii. 418 (1831). *CRUCIFERAE*, Benth. & Hook. f. i. 90.
sphanopterus, *A. Gray, Bot. U. St. Expl. Exped.* i. 55. —Peru.
chilensis, *DC. Syst.* ii. 419.—Chili.
lineariifolius, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 138.—Chili.
parvifolius, *Weddl. in Ann. Sc. Nat.* Sér. V. i. (1864) 288.—Bolivia.
peruvianus, *DC. Syst.* ii. 419.—Peruv.
pinnatifidus, *Hook. Ic. Pl.* t. 109.—Peruv.
pubescens, *Hook. l. c.* t. 81.—Peruv.
rhomboideus, *Hook. l. c.* t. 82.—Peruv.
sinuatus, *Hook. l. c.* t. 99.—Peruv.
sonchoides, *DC. Syst.* ii. 419.—Peruv.

CREMOPHYLLUM, Scheidw. in Bull. Acad. Brux. ix. 1. (1842) 23 = Dalechampia, Linn. (Euphorb.).
spathulatum, Scheidw. l. c. = D. spathulata.

CREMOPYRUM, Schur, Enum. Pl. Transs. 807 (1866) = Agropyrum, J. Gaertn. (Gramin.).
pectinatum, Schur, l. c. = A. cristatum.

CREMOSPERMA, Benth. Fl. Hartw. 234 (1846) = Bonaria, Linn. (Gesnerac.).
hirtiflorum, Benth. l. c.—Peruv.

CREMOSTACHYS, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 259 = Gelasaria, Zoll. (Euphorb.).
filiformis, Tul. l. c. 260 = G. pedicellata.
fulva, Tul. l. c. 261 = G. fulva.
Lindleyana, Tul. l. c. 262.—Ind. or.

CRENA, Scop. Introd. 220 (1777) = Crenea, Aubl. (Lythrar.).

CRENAEA, Schreb. Gen. i. 317 (1789) = prec.

CRENAMON, Rafin. New Fl. Am. iv. 86 (1836) (Genus dubium).
pectinatum, Rafin. l. c.—Am. bor.

CRENAMUM, Adans. Fam. ii. 113 (1768) = Crepis, Vaill. (Compos.).

CRENEA, Aubl. Pl. Gui. i. 383 t. 909 (1775). *LYTHRARIEAE*.
DOUBCAS, Linn. f. Suppl. 36, 245 (1781).
FAYA, Noch. Elem. ii. 109 (1790).
maritima, *Aubl. l. c.*—Bras.
repens, *G. F. W. Mey. Prim. Fl. Esseq.* 196.—Bras.
serpyllifolia, Kochne, in Engl. Bot. Jahrb. iii. (1882) 220 = repens.

CRENIAS, Spreng. Syst. iv. Cur. Post. 246, 247 (1827) = Miniopsis, Mart. (Podostem.).
scopularum, Spreng. l. c. = M. scopularum.

CRENOSCIADIUM, Boiss. & Heldr. Diagn. Ser. I. x. 80 (1849) = Opopanax, Koch (Umbellif.).
silfolium, Boiss. & Heldr. l. c. 81.—As. Min.

CRENULARIA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 181 = Moriera, Boiss. (Crucif.).
cunomoides, Boiss. Diagn. Ser. II. v. 41.—Cilicia.
glaucescens, Boiss. Fl. Orient. i. 337.—Cilicia.
orbiculata, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 181.—Macedon.
polygaloides, Boiss. Fl. Orient. i. 337.—As. Min.
umbellata, Boiss. l. c. 338 = M. umbellata.

CRESCHITON, Blume, in Flora, xiv. (1831) 506. *MELASTOMACEAE*, Benth. & Hook. f. i. 788.
bibracteata, *Blume, l. c.* 507.—Java.
emarginata, Miq. Fl. Ind. Bat. i. 1. 561 = pudibunda.
pudibunda, *Blume, in Flora*, xiv. (1831) 506.—Java.
superba, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 153 = bibracteata.

CRESCOME, Kunze, in Bot. Zeit. vi. (1848) 133, sphalm. = OREOCOME, Edgew. = Selinum, Linn. (Umbell.).

CREODUS, Lour. Fl. Cochinch. 88 (1790) = Chloranthus, Sw.
odorifer, Lour. l. c. 89 = C. inconspicuus.

CREOLOBUS, Lilja, Fl. Sver. Odl. Vext. 67 (1839) = Montesia, Linn. (Loasac.).
aureus, Lilja, l. c. = M. aurea.

CREPALIA, Steud. Nom. ed. I. 235 (1821) = CRAEPALIA, Schrank = Lolium, Linn. (Gramin.).

CREPIDARIA, Haw. Syn. Pl. Succ. 136 (1812) = Pedilanthus, Neck. (Euphorb.).
carinata, Haw. Suppl. Pl. Succ. 67 = P. carinatus.
cordellata, Haw. Syn. Pl. Succ. 136 = P. cordellatus.
myrtifolia, Haw. l. c. = P. tithymaloides.
padifolia, Haw. l. c. = P. padifolius.

CREPIDISPERMUM, Fries, Epicris. Hierac. 153 (1862) = Hieracium, Linn. (Compos.).

CREPIDIUM, Blume, Bijdr. 887 (1826) = Microstylis, Nutt. (Orchid.).
flavescens, Blume, l. c. 388 = M. flavescens.
Rheedii, Blume, l. c. 387 = M. Rheedii.

CREPIDIUM, Tausch, in Flora, ix. (1826) i. Erg. 80 = Crepis, Vaill. (Compos.).
asperum, Tausch, l. c. = Crepis aspera.
caulescens, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 438 = Crepis glauca.
Dioscoridis, Tausch, in Flora, ix. (1826) i. Erg. 80 = Crepis Dioscoridis.
glaucum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) = Crepis glauca.
runcinatum, Nutt. l. c. = Crepis runcinata.

CREPIDOCARPUS, Klotzsch, ex Boeck. in Linnaea, xxxvi. (1869–70) 414 = Botrycaulon, Linn. (Cyperac.).
cubensis, Klotzsch, ex Boeck. l. c. = S. cubensis.

CREPIDOPTERIS, Benth. in Mart. Fl. Bras. xv. i. 166 (1859), sphalm. = CREPIDOTROPIS, Walp. = Dioclea, H. B. & K. (Legumin.).

CREPIDOSPERMUM, Benth. & Hook. f. Gen. ii. 517, sphalm. = CREPIDISPERMUM, Fries = Hieracium, Linn. (Compos.).

CREPIDOSPERMUM, Hook. f. in Benth. & Hook. f. Gen. i. 826 (1862). *BURSERACEAE*, Benth. & Hook. f. i. 6.
Gogotianum, Triana & Planch. in Ann. Sc. Nat. Sér. V. xiv. (1872) 300 = Spruceanum.
guyanense, *Marchand, ex Triana & Planch. l. c.* in obs.—Guyana.
rhoifolium, *Triana & Planch. l. c.*—N. Granat.
Spruceanum, *Hook. f. in Benth. & Hook. f. Gen.* l. 826.—Peruv.

CREPIDOTROPIS, Walp. in Linnaea, xiv. (1840) 396 = Dioclea, H. B. & K. (Legumin.).
brachiosa, Walp. l. c. = D. brachiosa.

CREPINIA, Reichb. in Mocsal. Handb. ed. II. ii. 1415 (1838) = Pterothrixa, Cass. (Compos.).
Marschalliana, Reichb. l. c. = P. bifida.

CREPIS, [Vaill.] Linn. Gen. ed. I. 240. n. 621 (1737). *COMPOSITÆ*, Benth. & Hook. f. ii. 515.
ARGOSERIS, Stend. Nom. ed. II. i. 80 (1840).
ÆTHEORHIZA, Cass. in Dict. Sc. Nat. xlviii. 425 (1827).
ARIDOSERIS, Cass. l. c. 429 (1827).
ARISOKAMPHUS, DC. Prod. vii. 251 (1838).
ANTHOCHYTRUM, Reichb. lc. Fl. Germ. xix. 89. t. 1482 (1859).
ARACIUM, Neck. Elem. i. 49 (1790).
BARKHAUSIA, Moench, Meth. 587 (1794).
BARKHUSENIA, Hoppe, in Flora, xii. (1829) 512.
BERINIA, Brignol. Fl. Forojul. 30 (1810).
BILLOTIA, Sch. Bip. in Herb. & Winck. Jahrb. Pharmac. iv. (1841) 155, in nota.
BORKHAUSIA, Nutt. Gen. Am. ii. 125 (1818).
BRACHYDERRA, Cass. in Dict. Sc. Nat. xlviii. 429 (1827).
CALLIOPEA, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 309.
CATONIA, Moench, Meth. 535 (1794).
CERAMIOCEPHALUM, Sch. Bip. in Bull. Soc. Bot. Fr. ix. (1862) 284.
CLOSIROSPERMUM, Neck. Elem. i. 54 (1790).
CRENAMUM, Adans. Fam. ii. 112 (1763).
CRPIDIUM, Tausch, in Flora, xi. (1828) i. Erg. 86.
DESOURTIA, Boiss. Diagn. Ser. II. v. 114 (1856).
ENDOPTERA, DC. Prod. vii. 178 (1838).
GATTUMA, Cass. in Dict. Sc. Nat. xviii. 184 (1820)
GERLERA, Andrz. ex Besa. in Flora, xvii. (1834) i. Beibl. 18.
GERACIUM, Reichb. in Mocsal. Handb. ii. 1845 (1838).
HAPALOSTEPHIUM, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 307.
HETERODERIS, Boiss. Fl. Orient. iii. 798 (1875).
HOMALOCLINE, Reichb. Consp. 97 (1828).
HOSTIA, Moench, Meth. Suppl. 221 (1802).
INTYBELLIA, Monn. Essai Hierac. 78 (1829).
INTYBUS, Fries, Novit. Fl. Suec. ed. II. 244 (1842).
LAGOSERIS, Hoffingg. & Link, Fl. Port. ii. 149 (1820); Link, Enum. Hort. Berol. ii. 289 (1822).
LEPICAUNE, Lapeyr. Hist. Abr. Pl. Pyr. 478 (1818).
LEPIDOSERIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 103.
LIMNOCREPIS, Fourr. l. c. 104.
LIMNOSERIS, Peterm. Fl. Lips. 589 (1838).
NEMAUCHENES, Cass. in Bull. Soc. Philom. (1816) 77.
OMALOCLINE, Cass. in Dict. Sc. Nat. xlviii. 431 (1827).
PALEVA, Cass. l. c. xxxix. 398 (1826).
PSAMMOSERIS, Boiss. Diagn. Ser. I. xi. 52 (1849).
PSILOCHENIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 437.
RHYNCHOPAPPUS, Dulac, Fl. Hautes-Pyr. 493 (1867).
SOVERIA, Monn. Essai Hierac. 75 (1829).
SUCCISOCREPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 104.
WIBELIA, Gaertn. Mey. & Scherb. Fl. Wett. iii. 97 (1801).
YOUNGIA, Cass. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 88.

abietina, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 808 (= *Mulgedium abietinum*).—Oriens.
abyssinica, *Sch. Bip. in Flora*, xxii. (1839) 20.—Abyss.
acaulis, *Hook. f. Fl. Brit. Ind.* iii. 396.—Ind. or. ; Burma.
achyrophoroides, *Vatke*, *in Linnæa*, xxxix. (1875) 614.—Abyss.
acuminata, *Boiss. Fl. Orient.* iii. 856.—Syria.
acuminata, *Nuttl. ex Pursh. Am. Phil. Soc.* N. S. vii. (1841) 437.—Am. bor. occ.
adenocaulis, Vis. in Flora, xiii. (1830) 53 = nicæensis.
adenoderis, Sch. Bip. ex Oliver & Hiern, in Fl. Trop. Afr. iii. 430 = Rueppellii.
Adonis, Sprengl. Pugill. i. 54 = chondrilloides.
aetnensis, *Presl. Fl. Sicul.* i. 91.—Sicilia.
agrestis, Bieb. Fl. Taur. Cauc. ii. 256 = setosa.
agrestis, Pries, ex Nym. Consp. 357 = nicæensis.
agrestis, Steller, ex DC. Prod. vii. 164 = biennis.
agrestis, Waldst. & Kit. ex Willd. Sp. Pl. iii. 1602 = virens.
albida, All. Fl. Pedem. i. 219. t. 89 = alpestris.
albida, Bory & Chaub. Fl. Pélop. 54 = incana.
albida, *Vill. Prosp.* 37 ; *Hist. Pl. Dauph.* iii. 139.—Hispan.
alpestris, *Reichb. in Mocsal. Handb.* ed. II. ii. 1403.—Europ.
alpina, *Linn. Sp. Pl.* 806.—Aa. Min. ; Sibir.
altissima, Balb. Cat. Hort. Taur. (1804) 15 = Tolpis virgata.
ambigua, Balb. in Mém. Acad. Turin, xiv. (1805) ii. 69 = Tolpis virgata.
ambigua, A. Gray, Pl. Fendl 114 = Hieracium Fendleri.
ambigua, A. Gray, Pl. Wright. i. 129 = Hieracium Rusbyi.
amplexicaulis, Schur, Enum. Pl. Transs. 374 = blattaroides.
amplexifolia, *Willd. in Willd. & Lange, Prod. Fl. Hisp.* ii. 272.—Hispan.
amygdalina, Lag. Cat. Hort. Madr. (1814) 3 = foetida.
Andersonii, *A. Gray, in Proc. Am. Acad.* vi. (1866) 56.—Calif.
andryaloides, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 25.—Ins. Madera.
angustifolia, Urv. in Mem. Soc. Linn. Par. i. (1822) 357 = tectorum.
Apargia, Pers. Syn. ii. 375 = Chondrilla Peltidium.
aparginoides, Willd. Sp. Pl. iii. 1594 = Chondrilla Prolifera.
arabica, *Boiss. Fl. Orient.* iii. 858.—Arabia.
arenaria, *Pomel, Nouv. Mat. Fl. Atl.* 261, in nota.—Alg.
arenosa, *Scop. Ann.* ii. *Hist Nat.* (1769) 59.—Europ.
armena, *DC. Prod.* vii. 168.—Armen.
aspera, Lam. Encyc. ii. 180 = Picris aspera.
aspera, *Linn. Sp. Pl.* ed. II. 1133.—Syria ; Creta.
aspera, Suter, Fl. Helv. ed. II. ii. 187 = setosa.
athoa, *Boiss. Diagn.* Ser. I. xi. 57.—Macedon.
aurea, *Reichb. in Mocsal. Handb.* ed. II. 1402 ; *Tausch, in Flora*, xi. (1828) i. *Erg.* 78.—Europ.
auriculaefolia, Froel. in DC. Prod. vii. 165 = incarnata.
auriculella, *Sieber, in Spreng. Syst.* iii. 684.—Ins. Creta.
austriaca, Jacq. Enum. Vindob. 140, 270 = blattaroides.
baetica, Lange, in Kjoeb. Vidensk. Meddel. (1877–78) 227 = spathulata.
baetica, Mill. Gard. Dict. ed. VIII. n. 3 = Tolpis barbata.
baicalensis, *Ledeb. in Mém. Acad. Pétersb.* v. (1814) 589.—Sibir.
Balbisiana, F. Schultz, in Flora, xxiii. (1840) 718 = balearica.
balearica, *Costa, Fl. Catal.* 153.—Hispan.
bannatica, Willd. Enum. Hort. Berol. Suppl. 55 = setosa.
barbata, Linn. Sp. Pl. 805 = Tolpis barbata.
barbata, Mill. Gard. Dict. ed. VIII. n. 2 = foetida.
Bauhiniana, Tausch, in Flora, xii. (1829) i. Erg. 88 = virens.
bellidifolia, *Loisel. Fl. Gall.* 537. t. 18.—Corsica.
bellidifolia, Noe, ex Nym. Consp. 360 = cretica.
Bergeri, *Steud. Nom.* ed. II. i. 436.—Graecia.
bicolor, *Reichb. in Mocsal. Handb.* ed. II. ii. 1409, in obs.—Sicil.
biennis, Bieb. Fl. Taur. Cauc. ii. 257 = Marschallii.
biennis, Hook. Fl. Bor. Am. i. 297 = runcinata.
biennis, Lapeyr. Hist. Abr. Fl. Pyr. 84 = taraxacifolia.
biennis, *Linn. Sp. Pl.* 807.—Europ.
biennis, Sebast. & Maur. Fl. Rom. Prod. 272 = lacera.
bithynica, *Boiss. Diagn.* Ser. I. iv. 29.—Aa. Min.
blattarioides, *Vill. Hist. Pl. Dauph.* iii. 138.—Europ.
boetica, Amst. vide baetica.
boliviensis, *Wedd. Chlor. And.* i. 226.—Bolivia.
borealis, Sch. Bip. ex Walp. Rep. ii. 696 = Leontodon borealis.
bractеolata, Nym. Syll. 49 = athoa.
brevifloris, *Delile, ex Steud. Nom.* ed. II. i. 436.—Aegypt.
bullosa, *Tausch, in Flora*, xi. (1828) i. *Erg.* 78.—Reg. Mediterr.
Bungeana, C. A. Mey. ex DC. Prod. vii. 164 = Bungei.
CREPIS :—
Bungei, *Ledeb. ex DC. Prod.* vii. 164.—Sibir.
burejensis, *F. Schmidt, Reisen Amurl.* 52.—Reg. Amur.
bureniana, *Boiss. Fl. Orient.* iii. 852.—Persia.
burseifolia, *Linn. Sp. Pl.* 805.—Sicil.
bursifolia, Pall. ex DC. Prod. vii. 179 = Pterotheca bifida.
caespitosa, *Gren. & Godr. Fl. Fr.* ii. 332.—Corsica.
campestris, Schur, Enum. Pl. Transs. 376 = tectorum.
canariensis, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 161, in textu = Lowei.
Candollei, Sch. Bip. l. c. 457 = virens.
capillaris, *Wallr. Beitr. Fl. Hercyn.* 287.—Europ.
carbonaria, *Sch. Bip. in Flora*, xxii. (1839) 19.—Abyss.
caucasica, *C. A. Mey. in Mém. Acad. Petersb.* Sér. VI. *Sc. Nat.* vii. (1855) 15.—Reg. Caucas.
cernua, Tenore, ex Spreng. Pugill. i. 53 = neglecta.
chondrilloides, Froel. in DC. Prod. vii. 171 = Jacquini.
chondrilloides, *Jacq. Enum. Vindob. App.* 312.—Austria.
chrysantha, *Froel. in DC. Prod. vii.* 165.—Sibir. altaic.
ciliata, *C. Koch, in Linnæa*, xvii. (1843) 277.—Ins. Min.
cinerea, Desf. Tabl. ed. I. 89 = taraxacifolia.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 261, in nota.—Afr. bor.
cochlearifolia, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) 1. 190 (= *Youngia pygmæa*).—Rossia.
Columnæ, *Froel. in DC. Prod.* vii. 167.—Ital. ; Graecia.
comata, *Banks & Soland. ex Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 28.—Ins. Madera.
conyzaefolia, *Dalla Torre, Atl. Alpenfl.* 145.—Europ.
Cooperi, A. Gray, in Proc. Am. Acad. ix. (1874) 214 = virens.
corfacea, *Hornem. Hort. Hafn. Suppl.* ii. 768.—Hab. ?
corniculata, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 54.—As. centr.
coronopifolia, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) 88 = Tolpis coronopifolia.
corymbosa, Tenore, Prod. p. xlvii = neglecta.
cretica, *Boiss. Diagn.* Ser. I xi. 58.—Ins. Creta.
crinita, Soland. ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 24 = Tolpis crinita.
crithmifolia, Link, in Buch, Beschr. Canar. Ins. 147 = Tolpis crithmifolia.
crustica, *Hornem. Hort. Hafn. Suppl.* 90 = Dioscoridis.
crustica, *Schlos. & Vukot. in Rad Jugos. Ahad. Zagreb.* ixlii. (1881) 156.—Croatia.
crocea, *Reichb. in Mocsal. Handb.* ed. II. ii. 1402,—Europ.
crocifolia, *Boiss. & Heldr. Diagn.* Ser. I. vii. 14.—Graecia.
decumbens, *Gren. & Godr. Fl. Fr.* ii. 332.—Gallia.
decumbens, Loac. & Pardo, Ser. inconf. Pl. Aragon. 64 = taraxacifolia.
Dens-leonis, C. Koch, in Linnæa, xxiii. (1850) 689 = sonchifolia.
dentata, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126, nomen.—Malaya.
depressa, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 255.—Reg. Himal.
diffusa, DC. Cat. Hort. Monsp. 98 = virens.
dioritica, *Schott & Kotschy, ex Boiss. Fl. Orient.* iii. 842.—Cilicia.
Dioscoridis, Koch, Syn. Fl. Germ. [ed. II. 1081] = taraxacifolia.
Dioscoridis, *Linn. Sp. Pl.* ed. II. 1133.—Graecia.
Dioscoridis, Pollich, Fl. Hist. Pl. Palat. ii. 309 = tectorum.
Dioscoridis, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 348 = virens.
divaricata, *Boiss. & Heldr. Diagn.* Ser. I. vi. 14.—Graecia.
divaricata, *F. Schultz, in Flora*, xxiii. (1840) 719.—Ins. Madera.
diversifolia, Walp. Rep. vi. 361 = tenuifolia.
djimilensis, *C. Koch, in Linnæa*, xxiii. (1850) 668.—Reg. Caucas.
dubia, *Krock. Fl. Sil.* ii. 11. 224.—German.
dubia, *F. Schultz, in Flora*, xxiii. (1840) 718.—Ins. Madera.

CREPIS :—
achioides, All. Fl. Pedem. i. 222 = Helminthia echioides.
echioides, Ledeb. Ind. Sem. Hort. Dorpat. (1821) 28 = foetida.
albursensis, *Boiss. Diagn.* Ser. I. xi. 56.—Persia.
elegans, Fisch. ex DC. Prod. vii. 161 = balcalensis.
elegans, *Hook. Fl. Bor. Am.* i. 297.—Am. bor. occ.
erucaefolia, *Gren. & Godr. Fl. Fr.* ii. 331.—Gallia.
erucaefolia, Tausch, in Flora, xi. (1830) 1. Erg. 34 = lacera.
fallax, *Boiss. Fl. Orient.* iii. 850.—As Min.
farinosa, *Lam. Fl. Fr.* ii. 132.—Gallia.
fastigata, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 125.—Malaya.
filiformis, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 198 = Tolpis filiformis.
filiformis, *Vis. Fl. Lib. Spec.* 51. t. 21. f. 1.—Afr. bor.
fistulosa, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xxi. ii. (1848) 112 = praemorsa.
flexuosa, *Benth. & Hook. f. Gen.* ii. 515; *C. B. Clarke, Comp. Ind.* 264.—Reg. Himal.
flexuosa, Bernh. ex Stend. Nom. ed. II. i. 496 = Sonchus flexuosus.
floribunda, *Pomel, Nouv. Mat. Fl. Atl.* 261, in nota.—Afr. bor.
fonvielaeana, Froel. in DC. Prod. vii. t. 172 = chondrilloides.
foenciolacea, *Reichb. in Mossl. Handb.* ed. II. ii. 1401. —Europ.
fontana, *Froel. in DC. Prod.* vii. 172.—Armen.
fortens, Link, in Buch, Beschr. Canar. Ins. 147 = foetida.
foetida, *Linn. Sp. Pl.* 807.—Europ.; As. or.
foliosa, Dulac, Fl. Hautes-Pyr. 492 = blattarioides.
Frazzi, Sch. Bip. in Flora, xxv. (1842) 178 = Sieberi.
frutichiana, DC. Prod. vii. 165 = incarnata.
Frutichii, Stend. Nom. ed. II. i. 496 = Jacquini.
fuliginosa, Sibth. & Sm. Fl. Graec. Prod. ii. 138 = neglecta.
fuliginosa, *Webb & Berth. Phyt. Canar.* ii. 455.—Ins. Canar.
fuscipappa, *Benth. & Hook f. Gen.* ii. 514.—Ind. or.
Fussii, Kovats, ex Janka, in Linnaea, xxx. (1859-60) 587 = Hieracium rotundatum.
gaditana, Boiss. Voy. Bot. Espagne, 748 = virens.
glabra, *Boiss. Fl. Orient.* iii. 848.—Reg. Caucas.
glabra, *Koch. Fl. Sil. Supp.* ii. 370.—Europ.
glandulosa, *Bast. in Desv. Journ. Bot.* iii. (1814) 17. —Gallia.
glandulosa, Gmn. Ind. Sem. Hort. Bocead. (1825); Fl. Rar. 329 = foetida.
glauca, Benth. & Hook. f. Gen. ii. 515 = flexuosa.
glauca, *Torr. & Gray, Fl. N. Am.* ii. 488.—Am. bor. occ.
globifera, Hall. f. in Naturw. Anzeig. i. (1818) 91 = Dioscoridis.
glomerata, *Benth. & Hook. f. Gen.* ii. 515.—Reg. Himal.
glomerata, C. B. Clarke, Comp. Ind. 255 = depressa.
Gmelini, Schultes, Oestr. Fl. ii. 419 = biennis.
Gmelini, *Tausch, in in Flora*, (1828) 1. *Erg.* 78.—Sibir.
gracilipes, *Hook f. Fl. Brit. Ind.* iii. 396.—Reg. Himal.
gracilis, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 264 = fuscipappa.
graminifolia, Ledeb. in Mém. Acad. Pétersb. v. (1814) 558 = Lactuca Fischeriana.
grandiflora, Ledeb. Fl. Ross. ii. 827 = djimilensis.
grandiflora, *Tausch, in Flora*, xi. (1828) 1. *Erg.* 80.—Europ.
grandiflora, Willd. Enum. Hort. Berol. Suppl. 56 = albida.
gravcolens, *Schrad. ex Stend. Nom.* ed. II. i. 496.—German.
Hackelii, Lange, in Kjoeb. Vidensk. Meddel. (1877-78) 226 = taraxacifolia.
Haenseleri, F. Schultz, in Flora, xxiii. (1840) 718 = taraxacifolia.
hamata, Vitm. Summa Pl. iv. 417 = setosa.
hastata, *Kit. in Linnaea*, xxii. (1863) 414.—Europ.
heterophylla, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 315 = Lactuca Dubyaea.
heterocarpa, Nym. Syll. 48 = taraxacifolia.
heterogyna, *Froel. in DC. Prod.* vii. 169.—Dalmat.
heterogyrna, Schrad. ex Link, Handb. i. 799; DC. l. c. 179 = Dioscoridis.
heterotricha, *DC. l. c.* 168.—Persia.

CREPIS :—
hiemalis, Biv.ona, Man. i. n. 6. t. 2 = taraxacifolia.
hieracioides, Lam. Fl. Fr. ii. 111 = Picris hieracioides.
hieracioides, *Ledeb. Fl. Alt.* iv. 126.—Sibir. altaic.
hieracioides, Schrank, Baier. Fl. ii. 338 = Picris hieracioides.
hieracioides, F. Schultz, in Flora, xxiii. (1840) 718 = Lowei.
hieracioides, Waldst. & Kit. ex Willd. Sp. Pl. iii. 1601 = succisaefolia.
hieracolymitana, *Boiss. Diagn.* Ser. I. xi. 54.—Syria.
hissara, *Pomel, Nouv. Mat. Fl. Atl.* 261, in nota.—Alb. bor.
hirta, Linn. Sp. Pl. 807 = Leontodon hirtus.
hirta, Pers. Syn. ii. 377 = barbisifolia.
hispida, Waldst. & Kit. Pl. Rar. Hung. i. 42 = setosa.
hispidissima, Bartl. ex Reichb. Fl. Germ. Excurs. 358 = Picris laciniata.
hispidula, Delile, Fl. Egypte, 261. t. 42 = Leontodon hispidulus.
Hookeriana, *Ball, in Journ. Bot.* xi. (1873) 371.—Marocco.
Hookeriana, C. B. Clarke, Comp. Ind. 255 = glomerata.
Hookeriana, *Oliver & Hiern, in Oliver. Fl. Trop. Afr.* iii. 460.—Afr. trop.
Huetii, *Boiss. & Kotschy, in Boiss. Fl. Orient.* iii. 838. —Oriens.
humilis, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) 1. 190 (= *Youngia pygmaea*).—Sibir.; Am. bor.
hybrida, *A. Kern. Sched. Fl. Exsicc.* i. 69.—Tyrol.
hyoseridifolia, *Reichb. in Mossl. Handb.* ed. II. ii. 1403; *Tausch, in Flora*, xi. (1828) 1. *Erg.* 79.—Europ.
hyoseroides, Sibth. & Sm. Fl. Graec. Prod. ii. 139. t. 209 = neglecta.
incana, Lapeyr. Hist. Abr. Pl. Pyr. 483 = Andryala incana.
incana, Ledeb. Fl. Ross. ii. 827 = macropus.
incana, *Sibth. & Sm. Fl. Graec. Prod.* ii. 136; *Fl. Graec.* t. 202.—Graecia.
incarnata, *Tausch, in Flora*, xi. (1828) 1. *Erg.* 79; *Lexi. Syn. Comp.* 137.—Europ.
incarnata, Vis. Stirp. Dalm. Spec. 17. t. 6 = rubra.
incrassata, Banks, ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 24 = Tolpis crinita.
insularis, *Moris & De Not. in Mem. Acc. Tor.* ii. (1840) 141. t. 8. f. 1.—Ins. Capraria.
integra, *Mig. Ann. Mus. Bot. Lugd. Bat.* ii. 190.—Japon.
integrifolia, Dulac, Fl. Hautes-Pyr. 492 = succisaefolia.
integrifolia, *Vest, in Flora*, iii. (1820) 7.—Hab. ?
intermedia, *A. Gray, Syn. Fl. N. Am.* i. 11. 482.—Am. bor. occ.
interrupta, Sibth. & Sm. Fl. Graec. Prod. ii. 137; Fl. Graec. t. 203 = rhoeadifolia.
interrupta, Sieber, ex DC. Prod. vii. 161 = Sieberi.
inthybacea, Brot. Fl. Lusit. i. 231 = taraxacifolia.
Jacquini, *Tausch, in Flora*, xi. (1828) 1. *Erg.* 79.—Europ.
japonica, *Benth. Fl. Hongk.* 194.—As. trop.; Austral.
jubata, *Koch, Taschenbuch*, 231.—Helvet.
juvenalis, *F. Schultz, in Flora*, xxiii. (1840) 719.—Gallia.
Keiskeana, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 528.—Japon.
khorasanica, *Boiss. Fl. Orient.* iii. 835.—Persia.
Kitaibelii, *Prodi. in DC. Prod.* vii. 166.—Croatia.
Kochiana, Boiss. Fl. Orient. iii. 847 = seq.
Kochii, *Tchihat. Asie Min. Bot.* ii. 374.—As. Min.
Kotschyana, *Boiss. Fl. Orient.* iii. 833.—Persia.
Kotschyana, C. B. Clarke, Comp. Ind. 252 = foetida.
Krahlicii, *Pomel, Nouv. Mat. Fl. Atl.* 261, in nota.—Afr. bor.
lacera, *Tenore, Prod.* p. lxxi.—Ital.
Lachenalii, Gochn. Tent. Clchor. 19. t. 8 = tectorum.
laciniata, Schkuhr, ex DC. Prod. vii. 139 = Picris laciniata.
laciniata, *F. Schultz, in Flora*, xxiii. (1840) 718.—Ins. Madera.
lasvigata, *Dum. Fl. Belg. Prod.* 61.—Belgium.
lampsanoides, *Froel. in DC. Prod.* vii. 160.—Mont. Pyren.
lanceolata, Kit. in Linnaea, xxxii. (1863) 414 = virens.

CREPIS :—
lanceolata, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126, nomen.—Malaya.
lappacea, Host, Fl. Austr. ii. 421 = Picris laciniata.
lappacea, Lapeyr. Hist. Abr. Pl. Pyr. 488 = Picris hieracioides.
lappacea, Willd. Sp. Pl. iii. 1599 = Picris paucifloras.
lapsanifolia, Reichb. Fl. Germ. Excurs. 357 = pulchra
latialis, Sebast. Rom. Pl. fasc. ii. tit. 5 = lacera.
latifolia, Balb. ex Pers. Syn. ii. 377 = rigida.
Lechleri, *Sch. Bip. in Lechl. Berb. Am. Austr.* 54.—Chili.
leiocarpa, *Stend. Nom.* ed. II. i. 487.—Graecia.
leucophaea, Gand. ex DC. Prod. vii. 165 = virens.
leontodontoides, *Ali. Auct. Fl. Pedem.* 18.—Europ.
linearifolia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = tectorum.
linguaefolia, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 535.—As. or.
linifolia, Thuill. Fl. Par. 408 = virens.
lodomirensis, Bess. Prim. Fl. Galic. ii. 159 = biennis.
longe-pinnatifida, Chev. Fl. Par. ii. ii. 542 = virens.
longifolia, *Hegetschw. Fl. Schw.* 766.—Helvet.
Lowei, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii 461.—Ins. Canar.
lusitanica, Boiss. ex Sch. Bip. in Webb & Berth. l c 456 = virens.
lyrata, Benth. & Hook. f. Gen. ii. 514 = japonica.
lyrata, *Pomel. in DC. Prod.* vii. 170.—Sibir.
macrocephala, DC. Prod. vii. 172 = albida.
macrophylla, *Desf. Fl. Atlant.* ii. 231.—Afr. bor.
macropus, *Boiss. & Heldr. Diagn.* Ser. I. xi. 57.—As. Min.
macrorhiza, Banks, ex Hook. Bot. Mag. t. 2988 = Tolpis macrorhiza.
marginata, Dulac, Fl. Hautes-Pyr. 492 = lampsanoides.
maritima, Boucher, Fl. Abb. ed. III. 59 = biennis.
Marschallana, Reichb. Fl. Germ. xix. 126, sphalm vide *Crepinia Marschalliana*.
Marschallii, *F. Schultz, in Flora*, xxiii. (1840) 718.—Reg. Casp.
mauritiana, *Froel. in DC. Prod.* vii. 164.—Ins. Maurit.
microcephala, C. B. Clarke, ex Hook. f. Fl. Brit. Ind iii. 415 = Launaea microcephala.
mollis, Aschers. Fl. Brandenb. i. 468 = succisaefolia.
montana, Bernh. Syst. Verz. Erf. 188 = Hieracium sabaudum.
montana, *Reichb. in Mossl. Handb.* ed. II. ii. 1404.—Europ.
montana, Urv. in Mém. Soc. Linn. Par. i. (1822) 357 = Reuterlana.
mucronata, Nym. Syll. 46 = alpestris.
multicaulis, C. Koch, in Linnaea, xxiii. (1850) 690 = Kochiana.
multicaulis, *Ledeb. Fl. Alt.* iv. 126.—Sibir. altaic.
multiflora, *Sibth. & Sm. Fl. Graec. Prod.* ii. 138; *Fl. Graec.* t. 206.—Graecia; As. Min.
Mungieri, Boiss. & Heldr. in Boiss. Diagn. Ser. I. xi 58 = Sieberi.
muralis, Neck. ex Stend. Nom. ed. II. i. 487 = tectorum.
muricata, Gilib. Fl. Lituan. i. 231 = biennis.
muricata, Sibth. & Sm. Fl. Graec. Prod. ii. 138; Fl. Graec. t. 207 = parviflora.
muricata, Vitm. Summa Pl. iv. 417 = hispida.
muricata, Viv. ex Cess. in Bull. Soc. Bot. Fr. xii. (1865) 276 = vesicaria.
muricata, S. G. Gmel. Reise, i. 188 = tectorum.
nana, *Richard, Fraasl. Journ. App.* ed. II. 92.—Sibir.; Am. arct.
neglecta, Bieb. Casp. App. 210 = virens.
neglecta, *Linn. Mant.* i. 107.—Graecia; Oriens.
nemausensis, Bieb. Fl. Tauc. Cauc. iii. 538 = Pterotheca nemausensis.
nemausensis, Gouan, Illustr. 60 = Pterotheca nemausensis.
nemorosa, *Less. in Linnaea*, ix. (1834) 157, nomen.—Reg. Ural.
nemorum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 246 = virens.
Newii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 449.—Afr. trop.
nigricans, *Vis. Fl. Lib. Spec.* Fl. t. 10.—Afr. bor.
Nowae-Zemliae, *Mack. f. Hamilt. New Zeal. Fl.* 164. —N. Zel.
nudis, Lam. Fl. Fr. ii. 114 = Pterotheca nemausensis.

CREPIS :—
nedicensis, *Linn. Sp. Pl.* 805.—Europ.
nedicensis, *Siebor, ex Spreng. Syst.* iii. 634.—Creta.
nudiflora, Vir. Fl. Lib. Spec. 51. t. 13. f. 2 = aspera.
nemoidea, *Pomel, Nouv. Mat. Fl. Atl.* 261, nota.—Afr. bor.
nutans, Geyer, in Hook. Lond. Journ. Bot. vi. (1847) 253 = Calais nutans.
occidentalis, *Nutt. in Journ. Acad. Philad.* vii. (1834) 23.—Am. bor. occ.
olympica, *C. Koch, in Linnaea*, xxiii. (1850) 691.—Bithyn.
oporinoides, *Boiss. Voy. Bot. Espagne*, 868. t. 117.—Hispan.
Oreodis, *Schrenk, Enum. Pl. Nov.* ii. 32.—Soongaria.
Pallasii, *Turc. in Bull. Soc. Nat. Mosc* (1838) 96; xxi. (1848) ii. 110.—As. bor.
pallida, Hort. Par. ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 400 = Tolpis barbata.
paludosa, *Moench, Meth.* 535.—Europ.
paniculata, *Presl, Fl. Sicul.* p. xxxi.—Sicilia.
pannonica, C. Koch, in Linnaea, xxiii. (1850) 689 = rigida.
parviflora, *Desf. Tabl. ed.* 1. 88.—As. Min.; Persia.
parviflora, Moench, Meth. 535 = virens.
patula, Desf. Tabl. ed. 1. 86 = Dioscoridis.
patula, Poir. Voy. Barb. ii. 227 = Lapsana virgata.
pauciflora, *Desf. Tabl.* ed. ii. 108.—Aegypt.
pectinata, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1881) 24.—Ins. Madera.
pectinata, Steph. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) 1. 197 = Crepis tenuifolia.
pergyra, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 837 = macropus.
pinnatifida, *Presl. in DC. Prod.* vii. 167.—As. Min.
pinnatifida, Hort. ex DC. Prod. vii. 93 = Hypochoeris pratensis.
pinnatifida, Willd. Sp. Pl. iii. 1604 = virens.
pleurocarpa, A. Gray, in Proc. Am. Acad. xvii. (1881-83) 231 = intermedia.
Poeppigii, Sch. Bip. in Flora, xxxviii. (1855) 122 = Lechleri.
polymorpha, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126, nomen.—Malaya.
polymorpha, Gilib. Fl. Lituan. i. 231 = tectorum.
polymorpha, Pourr. in Mém. Acad. Toul. iii. (1788) 317 = taraxacifolia.
polymorpha, Wallr. Sched. Crit. 426 = virens.
polytricha, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 96; xxi. (1848) ii. 113.—As. temp.
pontica, *C. A. Mey. in Mém. Acad. Pétersb. Sér.* VI. *Sc. Nat.* vii. (1855) 14.—As. Min.
porrifolia, *D. Don, Prod. Fl. Nep.* 164.—Reg. Himal.
praecox, Balb. Misc. Bot. 37. t. 9 = taraxacifolia.
praemorsa, *Tausch, in Flora*, ix. (1826) 1. *Erg.* 79.—Europ.; As. bor.
primulifolia, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 398 = fuscipappa.
prostrata, *Dum. Fl. Belg. Prod.* 61, in syn.—Belgium.
pterosperma, Hort. ex DC. Prod. vii. 179 = Dioscoridis.
pterotheoides, *Boiss. Fl. Orient.* iii. 850.—Syria.
pulcherrima, Fisch. ex Link, Enum. Hort. Berol. ii. 290 = tenuifolia.
pulchra, Linn. Sp. Pl. 806 = Pterotheca lampsanoides.
pulchra, Soop. ex Steud. Nom. ed. I. 386 = virens.
pulmonariaefolia, *Presl. in DC. Prod.* vii. 169.—Italia.
pungens, *Desf. Tabl.* ed. I. 86, nomen; *et Pers. Syn.* ii. 376.—Hab. ?
purpurea, Bieb. Fl. Taur. Cauc. ii. 255 = Pterotheca bifida.
pygmaea, *Linn. Sp. Pl.* 806.—Europ.
racemifera, *Hook. f. Fl. Brit. Ind.* iii. 397.—Reg. Himal.
racemosa, Tausch, in Flora, xi. (1828) ii. 457 = pana.
radiata, Nym. Comp. 458 = foetida.
radicata, Forsk. Fl. Aegypt. Arab. 145 = Picris radicata.
radicata, Sibth. & Sm. Fl. Graec. Prod. ii. 186; Fl. Graeca, t. 200 = foetida.
ramosissima, *Kit. in Linnaea*, xxiii. (1850) 414.—Banat.
ramosissima, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 258.—Graecia.
raphanifolia, *Willd. Enum. Hort. Berol. Suppl.* 55.—Hab. ?
Raulinii, *Boiss. Diagn. Ser.* I. xi. 56.—Ins. Creta.
recognita, *Gaud. Fl. Helv.* v. 134.—Helvet.
recognita, Hall. f. in Naturw. Anmig. i. (1812) 90 = taraxacifolia.
CREPIS :—
reflexa, *Guss. Fl. Sic. Syn.* ii. 418.—Sicilia.
Reuteriana, *Boiss. & Heldr. Diagn.* Ser. I. xi. 55.—As. Min.
rhaetica, Nym. Consp. 456 = jacquini.
rhagadioloides, Linn. Mant. 108 = Picris Sprengeriana.
rhoeadifolia, Bieb. Fl. Taur. Cauc. ii. 259 = foetida.
rigens, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 127.—Ins. Azores.
rigida, *Waldst. & Kit. Pl. Rar. Hung.* i. 18. t. 19.—Europ. occ.; As. or.
robertioides, *Boiss. Diagn.* Ser. I. xi. 59.—Syria.
rodigeviodes, Sch. Bip. in Linnaea, xix. (1847) 814 = foetida.
Royleï, F. Schultz, in Flora, xxiii. (1840) 718 = Pterotheca bifida.
rubialis, Juss. ex DC. Prod. vii. 154 = taraxacifolia.
rubicaulis, Pers. Syn. ii. 375 = taraxacifolia.
rubra, *Linn. Sp. Pl.* 806.—Ital. ; Graecia.
ruderalis, Boucher, Fl. Abbev. ed. III. 59 = taraxacifolia.
Rueppellii, *Sch. Bip. in Flora*, xxii. (1839) 20.—Abyss.
rumicifolia, *Boiss. & Bal. ex Boiss. Fl. Orient.* iii. 843. —As. Min.
runcinata, *Torr. & Gray, Fl. N. Am.* ii. 487.—Am. bor. occ.
rupestris, Reichb. in Moesl. Handb. ed. II. ii. 1402 = Hieracium rupestre.
Ruprechtii, *Boiss. Fl. Orient* iii 843.—Reg. Caucas
sabauda, Balb. ex DC. Prod. vii. 163 = blennis.
sagittata, *Wender. Ind. Sem. Hort Mark.* (1829); *ex Linnaea*, v. (1830) *Lit.* 54.—Hab. ?
Sahendi, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 141.—Reg. Caucas.
salicifolia, Froel. in DC. Prod. vii. 169 = Bungei.
Sartoriana, Boiss. & Heldr. Diagn. Ser. II. iii. 100 = Columnae.
scabra, Lapeyr. Hist. Abr. Pl. Pyr. 488 = Picris hieracioides.
scabra, Willd. Sp. Pl. iii. 1608 = taraxacifolia.
scanensis, Linn. It. Scan. 170 = biennis.
scariosa, Willd. Sp. Pl. iii. 1598 = vesicaria.
scopulosa, *Kit. in Linnaea*, xxxii. (1863) 413.—Europ.
Schimperi, *Sch. Bip. ex Oliver, Fl. Trop. Afr.* iii. 449. —Afr. trop.
Schreberi, *Froel. in DC. Prod.* vii. 168.—Hab. ?
Schultzii, *Hochst. ex Oliver, Fl. Trop. Afr.* iii. 448.—Abyss.
scorzonerifolia, Rouy, in Bull. Soc. Bot. Fr. xxxi. (1884) 274 = albida.
segetalis, Roth, ex Steud. Nom. ed. II. i. 437 = tectorum.
segetum, *Schkuh. Cat. Pl. Helv.* ed. IV. 14, nomen.—Helv.
semeloides, *Delile, Fl. Egypte*, 262. t. 42. f. 2.—Aegypt.
setosa, *Hall. f. in Roem. Arch.* i. ii. 1.—Europ.; As. Min.
sibirica, Gouan, Illustr. 60 = blattarioides.
sibirica, *Linn. Sp. Pl.* 807.—Europ.; As. Min.; Reg. Himal.
Sibthorpiana, *Boiss. & Heldr. Diagn.* Ser. I. xi. 55.—Ins. Creta.
sicula, *Ucria, in Roem. Arch.* i. i. 69.—Sicilia.
Sieberi, *Boiss. Diagn.* Ser. I. xi. 58.—Graecia.
silenifolia, *Froel. in DC. Prod.* vii. 168.—Hab. ?
silhetensis, *Hook. f. Fl. Brit. Ind.* iii. 397.—Reg. Himal. ; Burma.
simplex, *Froel. in DC. Prod.* vii. 168.—Afr. bor.
sinuata, Lam. Encyc. ii. 179 = Picris sapienloides.
smyrnaea, *DC. Prod.* vii. 170.—As. Min.
sonchifolia, *C. A. Mey. in Schrenk, Enum. Pl. Nov.* ii. 32.—Orient.
spathulata, *Guss. Adnat. Cat. Pl. Boccod.* 5.—Mauritan.
spathulata, Lam. Encyc. ii. 181 = Suffreniana.
Sprengeriana, All. Fl. Pedem. i. 221 = Picris pauciflora.
Sprengeriana, Willd. Sp. Pl. iii. 1598 = Picris Sprengeriana.
stellata, Ball, in Journ. Bot. xi. (1873) 371 = taraxacifolia.
stenoma, *Turcz. ex DC. Prod.* vii. 164.—Dahur.; China.
stenotheca, Fries, ex Walp. Ann. ii. 1009 = Hieracium Crepidifpermeum.
CREPIS :—
Stolickaë, *C. B. Clarke, Comp. Ind.* 397.—Reg. Himal.
striata, Thunb. Prod. Pl. Cap. 189 = Hieracium capense.
stricta, Schultz, Fl. Starg. Suppl. 41 = tectorum.
stricta, Scop. Fl. Carn. ed. II. ii. 99 = neglecta.
succisaefolia, *Tausch, in Flora*, ix. (1826) 1. *Erg.* 79.—Rossia.
succulenta, [*Dryand. in*] Ait. Hort. Kew. ed. I. iii. 126 = Tolpis fruticosa.
Suffreniana, *Steud. Nom.* ed. I. 286; *Lloyd, Fl. Loir. Inf.* 155.—Europ.
sulphurea, *Pourr. ex Willd. & Lange, Prod. Fl. Hisp.* ii. 251.—Hispan.
tanegana, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 198. —Japon.
taraxacifolia, *Thuill. Fl. Par.* 409.—Europ.
taraxacifolia, Willd. Sp. Pl. iii. 1598 = taraxacoides.
taraxacoides, *Desf. Fl. Atlant.* ii. 231.—Afr. bor.
taraxacoides, Pourr. in Mém. Acad. Toul. iii. (1788) 317 = alpina.
taraxacoides, Willd. ex Steud. Nom. ed. II. i. 438 = taraxacifolia.
Taraxacum, Stokes, in With. Bot. Arr. ed. II. ii. 853 = Taraxacum officinale.
taurienensis, Bieb. Fl. Taur. Cauc. iii. 537 = Marschallii.
taurinensis, Willd. Sp. Pl. iii. 1595 = taraxacifolia.
tectorum, Huds. Fl. Angl. ed. I. 301 = virens.
tectorum, *Linn. Sp. Pl.* 807.—Europ.; As. bor.
tectorum, Vill. Hist. Pl. Dauph. iii. 144 = taraxacifolia.
tenerrima, Sch. Bip. ex Oliver, Fl. Trop. Afr. iii. 449 = carbonaria.
tenuifolia, Soland. & Banks, ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 24 = Tolpis pectinata.
tenuifolia, *Willd. Sp. Pl.* iii. 1606.—Reg. Himal. Ass. bor.
tergloucnsis, *A. Kern. Sched. Fl. Exsicc.* 81.—Europ.
thracia, Sprengl. Syst. iii. 634 = foetida.
tinctoria, Dulac, Fl. Hautes-Pyr. 491 = tectorum.
tingitana, *Ball, in Journ. Linn. Soc.* xvi. (1878) 587. —Marocc.; Hisp.
tomentosa, Moench, Meth. 535 = Dioscoridis.
transsilvanica, Schur, Enum. Pl. Transs. 375 = biennis.
triangulata, J. & C. Presl, Delic. Prag. 110 = leontodontoides.
Trinii, *Nym. Syll.* 49.—Hispan.
trichotoma, Moench, Meth. Suppl. 216 = Tolpis conopoilia.
tuberculata, Sch. Bip. ex Nym. Consp. 459 = leontodontoides.
Turczaninowii, C. A. Mey. in Schrenk, Enum. Pl. Nov. ii. 32 = Pallasii.
umbellata, Bernh. Syst. Verz. Erf. 138 = Hieracium umbellatum.
umbellata, Thuill. Fl. Par. 409 = virens.
uniflora, Thuill. l. c. 410 = virens.
varia, Moench, Meth. 534 = tectorum.
variabilis, Krock. Fl. Siles. ii. 11. 326 = virens.
versicolor, Fisch. ex DC. Prod. vii. 151 = Lactuca Fischeriana.
vesicaria, Balb. Elenc. 93 = taraxacifolia.
vesicaria, *Linn. Sp. Pl.* 805.—Hispan.
virens, *Linn. l. c.* ed. II. 1134.—Europ.; Reg. Caucas.
virgata, Desf. in Act. Soc Hist. Nat. Par. i. (1792) 87 = Tolpis virgata.
virgata, Lapeyr. Hist. Abr. Pl. Pyr. 488 = Picris hieracioides.
viscidula, *Froel. in DC. Prod.* vii. 166.—Benat.
willemetioides, *Boiss. Fl. Orient.* iii. 845.—Persia.
youngiformis, C. Koch, in Linnaea, xvii. (1843) 377 = Phocasium leontodontoides.
zacinthia, Nym. Syll. 49 = foetida.

CREPULA, Hill, Veg. Syst. iv. 9 (1762) = Centaurea, Linn. (Compos.).
centaureoides, Hill, Hort. Kew. 58 = Cent. centaurioides.
cornua, Hill, l. c.—Hab. ?

CRESCENTIA, Linn. Syst. ed. I (1735). *BIGNONIACEAE*. Benth. & Hook. f. ii. 1043.
DENDROSICUS, Rafin. Sylva Tellur. 80 (1838).
acuminata, H. B. & K. Nov. Gen. et Sp. iii. 156 = Parmentiera edulis.

CRESCENTIA:—
acuminata, H. B. & K. Nov. Gen. et Sp. iii. 57 = C. Cujete.
aggregata, *Miers, in Trans. Linn. Soc.* xxvi. (1868) 173.—Jamaic.
alata, H. B. & K. Nov. Gen. et Sp. iii. 158 = Parmentiera alata.
angustifolia, Willd. ex Seem. in Trans. Linn. Soc. xxiii. (1862) 90 = C. Cujete.
arborea, Raſn. Sylva Tellur. 81 = C. Cujete.
confertifolia, *Miers, in Trans. Linn. Soc.* xxvi. (1868) 170.—Jamaic.
coriacea, *Miers, l. c.* 177.—Am. trop.
cucurbitina, *Linn. Mant.* 250.—Am. trop.
Cujete, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 138 = cucurbitina.
Cujete, *Linn. Sp. Pl.* 626.—Am. trop.
Cujete, Vell. Fl. Flum. 269; vi. t. 103 = plectantha.
cunefolia, Gardp. in Hook. Journ. Bot. ii. (1840) 422 = C. Cujete.
cuspidata, *Miers, in Trans. Linn. Soc.* xxvi. (1868) 178.—Venezuela.
edulis, Desv. Journ. Bot. iv. (1814) 113 = Parmentiera edulis.
elongata, *Miers, in Trans. Linn. Soc.* xxvi. (1868) 178. —Ind. occ.
fasciculata, *Miers, l. c.* 171.—Ind. occ.
jasminoides, *Lam. Encyc.* i. 559.—Ins. Baham.
latifolia, Mill. Gard. Dict. ed. VIII. n. 2 = cucurbitina.
latifolia, Raſn. Sylva Tellur. 81 = C. Cujete.
lethifera, Tuss. Fl. Antill. iv. 50. t. 17 = cucurbitina.
linearifolia, *Miers, in Trans. Linn. Soc.* xxvi. (1868) 172.—Ind. occ.
macrophylla, Seem. in Hook. Kew Journ. vi. (1854) 278 = Amphitecna macrophylla.
microcarpa, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 294.—Porto Rico.
mucosocarpa, Zaldivar, ex Heller, Reisen in Mexico, 414 = Parmentiera edulis.
nigripes, Linden, Cat. n. 17 (1862) 7 = Amphitecna nigripes.
obovata, Benth. Bot. Voy. Sulph. 130. t. 46 = cucurbitina.
ovata, Burm. f. Fl. Ind. 132 = cucurbitina.
pallustris, Forsyth, ex Seem. in Trans. Linn. Soc. xxiii. (1860) 20 = cucurbitina.
pinnata, Jacq. Coll. iii. 208 (= *Tripinna africana*).—Afr. trop. ?
plectantha, *Miers, in Trans. Linn. Soc.* xxvi. (1868) 170.—Bras.
pumila, Raſn. Sylva Tellur. 81 = C. Cujete.
regalis, *Linden, Fl. Nouv.* (1859) 7.—Am. trop.
spathulata, *Miers, in Trans. Linn. Soc.* xxvi. (1868) 178.—Ecuador.
tuxicaria, Tuss. Fl. Antill. iv. 50. t. 17 = cucurbitina.
trifolia, Blanco, Fl. Filip. ed. I. 489 = Parmentiera alata.

CRESSA, Linn. Dies. Dan. 13 (1747); Amoen. Acad. i. 395 (1749). *CONVOLVULACEÆ*, Benth. & Hook. f. ii. 881.
arabica, *Forsk. Fl. Aegypt. Arab.* 54.—Arabia.
arenaria, Willd. ex Roem. & Schult. Syst. vi. 207 = truxillensis.
australis, R. Br. Prod. 490 = cretica.
cretica, *Linn. Sp. Pl.* 223.—Cosmop. trop.
humifusa, Lam. Fl. Fr. ii. 266 = cretica.
indica, Retz. Obs. iv. 24 = cretica.
intermedia, *T. Anders. in Journ. Linn. Soc. v. Suppl.* i. (1860) 32.—Aden.
latifolia, T. Anders. l. c. 35 = Breweria latifolia.
microphylla, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 63, 128 = cretica.
mucronifolia, Stokes, Bot. Mat. Med. ii. 83 = cretica.
multiflora, *Willd. ex Roem. & Schult. Syst.* vi. 207.—Am. austr.
nudicaulis, *Griseb. in Goett. Abh.* xxiv. (1879) 266.—Reg. Argent.
sericea, *Willd. ex Roem. & Schult. Syst.* vi. 207.—Am. austr.
truxillensis, *H. B. & K. Nov. Gen. et Sp.* iii. 98.—Peruv.
villosa, Hoffmgg. & Link, Fl. Port. i. 373 = cretica.

CRINIPES, Hochst. in Flora, xxxviii. (1855) 279 = Danthonia, DC. (Gramin.).
abyssinicus, Hochst. l. c. 280 = D. tenuiginnis ?

CRINISSA, Reichb. Nom. 95 (1841) = Pyrrhopappus, DC. (Compos.).

CRINITA, Houtt. Handl. vii. 861. t. 40. f. 1 (1777) = Pavetta, Linn. (Rubiac.).
capensis, Houtt. l. c. 862 = P. caffra.

CRINITA, Moench, Meth. 578 (1794) = Aster, Tourn. (Compos.).
linearifolia, Moench, Meth. Suppl. 237 = Chrysocoma Coma-aurea.
punctata, Moench, Meth. 578 = A. acris ?

CRINITARIA, Cass. in Dict. Sc. Nat. xxxvii. 475 (1825) = Aster, Tourn. (Compos.).
bifora, Cass. l. c. 478 = A. acris.
humilis, Hook. Fl. Bor. Am. ii. 24 = A. angustus.
Linosyris, Less. Syn. Comp. 195 = A. Linosyris.
oleæfolia, Cass. in Dict. Sc. Nat. xxxvii. 476 (= *Linosyris villosa*).—Europ. or. ; As. bor.
punctata, Cass. l. c. = A. acris.
villosa, Cass. l. c. (= *Linosyris villosa*).—Europ. or. ; As. bor.
viscidiflora, Hook. Fl. Bor. Am. ii. 24 = Bigelowia viscidiflora.

CRINODENDRON, Molina, Saggio Chili, 179 (1782) = Tricuspidaria, Ruiz & Pav. (Tiliac.).
Hookerianum, C. Gay, Fl. Chil. i. 341 = T. dependens.
Patagua, Molina, Saggio Chili, 179 = T. dependens.

CRINODENDRUM, Juss. Gen. 431 (1789) = præc.

CRINONIA, Banks, ex Tul. in Arch. Mus. Par. viii. (1855) 405 = Hedycarya, Forst. (Monim.).

CRINONIA, Blume, Bijdr. 838 (1825) = Pholidota, Lindl. (Orchid.).
carnea, Blume, l. c. 889 = P. carnea.
globosa, Blume, l. c. = P. globosa.

CRINOPSIS, Herb. Amaryll. 270 (1837) = Crinum, Linn. (Amaryllid.).

CRINUM, Linn. Gen. ed. I. 97 (1737). *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 726.
CRINOPSIS, Herb. Amaryll. 270 (1837).
ERYDOLEA, Salisb. Gen. Pl. Fragm. 115 (1866).
LIRIAMUS, Raſn. Fl. Tellur. iv. 23 (1836).
SCADIANUS, Raſn. Ant. Journ. 164 (1838).
TARNAIS, Salisb. Gen. Pl. Fragm. 115 (1866).
TANGHINOLLI, Adans. Fam. ii. 57 (1763).
abyssinicum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 211.—Abyss.
africanum, Linn. Sp. Pl. 292 = Agapanthus umbellatus.
albiflorum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I Art. iv. 12 = asiaticum ?
album, *Herb. Amaryll.* 272.—Arab.
algense, *Herb. l. c.*—Afr. austr.
amabile, *Donn, Hort. Cantab.* ed. VI. 83.—Sumatra.
amabile, Roxb. ex Stend. Nom. ed. II. i. 438 = latifolium.
americanum, Hort. Prag. ex Steud. l. c. = erubescens.
americanum, *Linn. Sp. Pl.* 292.—Am. austr.
americanum, [DC. in] Red. Lil. t. 332 = asiaticum.
amœnochaneles, *Baker, in Journ. Bot.* xvi. (1878) 195.—Afr. trop.
amœnum, *Roxb. Hort. Beng.* [28] ; *Fl. Ind.* ii. 127.—Ind. or.
angustifolium, Herb. ex Steud. Nom. ed. II. i. 438 = asiaticum.
angustifolium, Linn. f. Suppl. 195 = Cyrtanthus angustifolius.
angustifolium, *R. Br. Prod.* 297.—Austral.
angustum, Steud. Nom. ed. II. i. 438, sphalm. = angustum.
annulatum, Herb. App. 22 = asiaticum.
aquaticum, Burch. ex Spreng. Syst. ii. 55 = campanulatum.
aquaticum, Herb. App. 22 = erubescens.
arenarium, Herb. in Bot. Mag. t. 2355 = angustifolium.
asiaticum, Blanco, Fl. Filip. ed. I. 251 = gracile.
asiaticum, [DC. in] Red. Lil. t. 348 = americanum.
asiaticum, *Linn. Sp. Pl.* 292.—As. trop.
asiaticum, Lour. Fl. Cochinch. 197 = cochinchinense.
asiaticum, Roxb. Hort. Beng. [28] = defixum.
atlenuatum, Willd. ex Schult. f. Syst. vii. 873 = Counselyni.
augustum, *Roxb. Hort. Beng.* [28] ; *Fl. Ind.* ii. 126.—Ins. Mauric.
australasicum, Herb. Amaryll. 259 = angustifolium.
australe, Donn, Hort. Cantab. ed. VI. 83 ; Hornem. Hort. Hafn. 818 = pedunculatum.

CRINUM:—
Balneoli, *Baker, in Gard. Chron.* (1881) ii. 89.—Afr. austr.
Balfouri, *Baker, in Hook. Bot. Mag.* t. 6570.—Ins. Socotra.
bancanum, *Kurz, in Tijdschr. Nederl. Ind.* xxvii. (1864) 321.—Malaya.
barbatum, Linn. ex Herb. Amaryll. 138 = Hippeastrum equestre.
blandum, M. Roem. Syn. Enast. 76 = angustifolium.
brachyandrum, *Herb. in Bot. Mag.* sub t. 2121. p. 8.—Austral.
brachynema, *Herb. in Bot. Reg.* (1842) *Misc.* 86.—Ind. or.
bracteatum, *Willd. Sp. Pl.* ii. 47.—Ins. Seychell.
brevifolium, Roxb. Hort. Beng. [28] ; Fl. Ind. ii. 129 = bracteatum.
brevilimbum, *Carey, ex Herb. in Bot. Mag.* sub t. 2121 p. 7.—Ins. Pacif.
Broussonetii, Herb. l. c. t. 2121 = yuccæflorum.
Broussonetianum, Herb. Amaryll. 260. t. 44. f. 13 = yuccæflorum.
buphanoides, *Welw. ex Baker, in Journ. Bot.* xvi. (1878) 195.—Afr. trop.
caffrum, Herb. in Sweet, Hort. Brit. ed. III. 678 = campanulatum.
campanulatum, *Herb. in Bot. Mag.* sub t. 2121. p. 7. —Afr. austr.
canaliculatum, Carey, ex Steud. Nom. ed. II. i. 439 = capalliflorum.
canaliculatum, Roxb. Hort. Beng. [28] ; Fl. Ind. ii. 132 = pedunculatum.
canaliflorum, *Herb. in Bot. Mag.* sub t. 2121. p. 7.—Ind. or.
Candelabrum, Hort. ex M. Roem. Syn. Enast. 87 = Brunsvigia gigantea.
capense, Herb. Amaryll. 269 = longifolium.
Careyanum, *Herb. in Bot. Mag.* t. 2466.—Ins. Masc.
caribæum, *Baker, in Gard. Chron.* (1881) ii. 40.—Jamaic.
carinifolium, Stokes, Bot. Mat. Med. ii. 215 = asiaticum *caspium*, Pall. Reise, ii. 786 = Allium caspium.
chinense, Lodd. ex Kunth, Enum. Pl. v. 550 = asiaticum cochinchinense, *M. Roem. Syn. Enast. 71.*—Cochinch
Commelinianum, Herb. Amaryll. 254 = Commelyni.
Commelyni, [DC. in] Red. Lil. t. 323 = americanum.
Commelyni, *Jacq. Hort. Schœnb.* ii. 40. t. 209.—Am austr.
concinnum, *Mart. in Schult. f. Syst.* vii. 857.—Bras.
confertum, Herb. in Bot. Mag. t. 2523 = angustifolium.
conicum, M. Roem. Syn. Enast. 87 = americanum.
corantinum, Herb. ex Steud. Nom. ed. II. i. 439 = erubescens.
corantynum, M. Roem. Syn. Enast. 80 = præc.
cordofanum, *Ketschy & Peyr. Pl. Tinn.* 14. t. 21.—Afr. trop.
crassifolium, Herb. App. 23 = variabile.
Cruckshium, D. Dietr. Syn. Pl. ii. 1361 = cruentum.
cruentum, [*Ker-Gawl. in*] *Bot. Reg.* t. 171.—Ind. or.
Cunninghamii, *Baker, in Gard. Chron.* (1881) ii. 72.—Ins. Philipp.
declinatum, Herb. in Bot. Mag. t. 2231 = asiaticum.
defixum, [*Ker-Gawl. in Journ. Sc. & Arts*, iii. (1817) 106.—Ind. or.
densum, Klotzsch, ex Kunth, Enum. Pl. v. 554 = gracilliflorum.
Dieterichii, Schult. Obs. 65 = Commelyni.
distichum, Herb. in Bot. Mag. sub t. 2121. p. 7 = yuccæflorum.
elegans, Carey, ex Herb. l. c. = pratense.
ensifolium, Roxb. Hort. Beng. [28] ; Fl. Ind. ii. 129 = defixum.
erubescens, H. B. & K. Nov. Gen. et Sp. i. 279 = Kunthianum.
erubescens, [*Salisb. in*] *Ait. Hort. Kew.* ed. I. i. 418. —Am. austr.
erythrophyllum, *Carey, ex Herb. in Bot. Mag.* sub t. 2121. p. 7.—Guyana.
exaltatum, Herb. l. c. p. 5 = pedunculatum.
falcatum, Jacq. Hort. Vindob. iii. 24 = Ammocharis falcata.
fimbriatulum, *Baker, in Journ. Bot.* xvi. (1878) 196.—Afr. trop.
firmifolium, *Baker, in Journ. Linn. Soc.* xx. 270.—Madag.
flaccidum, *Herb. in Bot. Mag.* sub t. 2121. p. 7.—Austral.
floridanum, Griseb. Fl. Brit. W. Ind. 588 = erubescens.
floridanum, Steud. Nom. ed. II. i. 439 = sp.

CRINUM :—
floridum, Fraser, ex Herb. in Bot. Mag. t. 2121. p. 8 = asiaticum.
floridum, Salisb. Prod. 298 = Agapanthus umbellatus.
Forbesianum, Herb. Amaryll. 267 = seq.
Forbesii, *Schult. f. Syst.* vii. 864.—Afr. trop.
formosum, *Herb. ex Soub. in Mart. Fl. Bras.* ii. t. 169.—Bras.
giganteum, *Andr. Bot. Rep.* t. 169.—Afr. austr.
giganteum, Blanco, Fl. Filip. ed. II. 175 = asiaticum.
Gordonianum, M. Roem. Syn. Enast. 80 = eubescens.
Gownium, Herb. in Trans. Hort. Soc. iii. (1822) 190 = longifolium !
gracile, *G. F. W. Mey. ex Presl, Rel. Haenk.* ii. 120. —Ins. Luzon.
graciliflorum, *Kunth, Enum. Pl.* v. 565.—Venezuela.
guianense, M. Roem. Syn. Enast. 80 = Kunthianum.
Hayleckii x, Herb. Amaryll. 274.
Herberti, G. Don, ex Loud. Hort. Brit. 120 = Commelyni.
Herbertianum, Schult. f. Syst. vii. 871 = strictum.
Herbertianum, Wall. Pl. As. Rar. ii. 38. t. 145 = zeylanicum.
Hildebrandtii, *Vatke, in Monatsb. Acad. Berl.* (1876) 868.—Ins. Johanna.
Hornemannianum, M. Roem. Syn. Enast. 71 = asiaticum.
humile, *Herb. in Bot. Mag.* t. 2686.—Ind. or.
imbricatum, *Baker, in Gard. Chron.* (1881) ii. 760.—Afr. austr.
imperiale, *Regel, Cat. Pl. Hort. Absakov.* 43.—Hab. ?
inodorum, *Tausch, in Flora*, xix. (1836) ii. 422.—Am. austr.
insigne, Sweet, Hort. Brit. ed. I. 405 = latifolium.
Kerii, M. Roem. Syn. Enast. 93 = lineare.
Kirkii, *Baker, in Hook. Bot. Mag.* t. 6519.—Afr. trop.
Knyffii, *C. Morr. in Belg. Hortic.* ii. (1852) 373. t. 63. —Hab. ?
Kunthianum, *M. Roem. Syn. Enast.* 80.—Nicaragua.
Lancasterum, Willd. ex M. Roem. l. c. 94 = seq.
Lancei, *Herb. ex Sweet, Hort. Brit.* ed. III. 677.—Guiana.
latifolium, Andr. Bot. Rep. t. 478 = zeylanicum.
latifolium, *Linn. Sp. Pl.* 291.—Ind. or.
leucophyllum, *Hook. Bot. Mag.* t. 6788.—Afr. trop.
ligulatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 270.—Madag.
Lindleyanum, *Schult. f. ex Soub. in Mart. Fl. Bras.* iii. t. 159.—Bras. ; Guiana.
Lindleyanum, Herb. Amaryll. 252 = Commelyni.
lineare, *Linn. f. Suppl.* 195.—Afr. austr.
Linnaei, M. Roem. Syn. Enast. 89 = ornatum.
Loddigesianum, Herb. Amaryll. 253, 401 = cruentum.
longiflorum, *Herb. in Bot. Mag.* sub t. 2121.—Jamaic.
longifolium, *Roxb. Hort. Beng.* [38] ; *Fl. Ind.* ii. 130. —Ind. or.
longifolium, Thunb. Prod. Pl. Cap. 59 = capense.
longistylum, Herb. ex Steud. Nom. ed. II. i. 440 = latifolium.
lorifolium, Roxb. Hort. Beng. [23] ; Fl. Ind. ii. 131 = pratense.
Lourieri, M. Roem. Syn. Enast. 85 = zeylanicum !
Macowani, *Baker, in Gard. Chron.* (1878) i. 398.—Afr. austr.
macrocarpum, Carey, ex Kunth, Enum. 1 i. v. 558 = sumatranum.
Makoyanum, Carr. in Rev. Hortic. (1877) 417 = Moorei.
Martinezii, Spreng. Syst. ii. 56 = Lapiedra Martinezii.
mauritianum, *Ledd. Bot. Cab.* t. 650.—Madag.
melocarnum, Herb. ex Reichb. Exot. t. 47 = zeylanicum.
melocarnum, Roxb. Hort. Beng. [38] ; Fl. Ind. ii 140 = latifolium.
Moorei, *Hook. f. Bot. Mag.* t. 6118.—Afr. austr.
multiflorum, *Dauf. Cat. Hort. Par.* ed. III. 47.—Hab. ?
nervosum, L'Hérit. Sert. Angl. 8 = Eurycles zeylvestris.
nervosum, Roxb. Hort. Beng. [38] ; Fl. Ind. ii. 135 = Eurycles amb.
norfolkianum, *A. Cunn. in Hook. Lond. Journ. Bot.* i. (1842) 138.—Ins. Norfolk.
Northianum, *Baker, in Gard. Chron.* (1888) i. 671.—Borneo.
obliquum, Linn. f. Suppl. 195 = Cyrtanthus obliquus.
octoflorum, Herb. ex Steud. Nom. ed. II. i. 440 = eubescens.
odoratissimum, *Tausch, in Flora*, xix. (1836) 422.—Hab. ?

CRINUM :—
ornatum, Herb. Amaryll. 262 = zeylanicum.
Oubechii, *Schult. f. Syst.* vii. 1721.—Hab. ?
pauciflorum, *Baker, in Journ. Bot.* xvi. (1878) 195.—Afr. trop.
Paxtoni, *Herb. Amaryll.* 271. t. 44. f. 24.—Ind. or.
pedunculatum, *R. Br. Prod.* 297.—Austral.
petiolatum, Herb. App. 22 = giganteum.
plicatum, Livingst. ex Hook. Bot. Mag. t. 2908 = asiaticum.
podophyllum, *Hook. Bot. Mag.* t. 6488.—Afr. trop.
pratense, *Herb. Amaryll.* 256.—Ind. or.
procerum, Carey, ex Herb. App. 22 = asiaticum.
purpurascens, *Herb. Amaryll.* 250.—Ins. Fernand. Noronh.
pusillum, *Herb. l. c.* 255. t. 82. f. 3.—Ins. Nicobar.
quitense, Spreng. Syst. ii. 55 = Phaedranassa chloracea.
Redoutéanum, M. Roem. Syn. Enast. 70 = asiaticum.
revolutum, Herb. in Bot. Mag. sub t. 2121. p. 5 = lineare.
revolutum, Lindl. in Trans. Hort. Soc. vi. (1826) 285 = Commelyni.
rigidum, Herb. Amaryll. 248 = sumatranum.
rigaricum, Herb. App. 23 = longifolium.
roseum, *Ecki. ex M. Roem. Syn. Enast.* 84, nomen.—Afr. austr.
rubriculæ, M. Roem. l. c. 70 = pedunculatum.
rubrinervium, Sweet, ex Steud. Nom. ed. II. i. 440 = eubescens.
Ruppellianum, Fresen. in Verh. Beloerd. Gartenb. iii 265.—Arabrum.
Sanderianum, *Baker, in Gard. Chron.* (1884) ii. 102.—Afr. trop.
scabervirum, Herb. in Trans. Hort. Soc. iii. (1822) 196 = scabrum.
scabrum, *Herb. in Bot. Mag.* t. 2180.—Bras.
Schmidti, *Regel, in Act. Hort. Petrop.* vi. (1880) 544. —Afr. austr.
sectitubum, *Teijsm. & Binn. Cat. Hort. Bog.* 877, nomen.—Hab. ?
serrulatum, *Baker, in Gard. Chron.* (1881) i. 786.—Cambodia.
sinicum, Roxb. ex Herb. in Bot. Mag. sub t. 2121. p. 7 = asiaticum.
speciosissimum, Herb. l. c. p. 6 = zeylanicum.
speciosum, Linn. f. Suppl. 195 = Vallota purpurea.
spectabile, Herb. ex Steud. Nom. ed. II. i. 440 = giganteum.
spirale, Andr. Bot. Rep. t. 92 = Carpolyza spiralis.
Spofforthianum, Herb. ex Sweet, Hort. Brit. ed. III. 676 = scabrum.
stenophyllum, *Baker, in Gard. Chron.* (1881) i. 786.—Burma.
Strachеyi, *Baker, l. c.* xvi. (1881) 72.—Reg. Himal.
strictum, *Herb. in Bot. Mag.* t. 2685.—Am. austr.
strictum, *Hornem. Enum. Pl. Hort. Hafn.* (1807) 15 ; *Hort. Hafn.* i. 318.—Hab. ?
subcernuum, *Baker, in Gard. Chron.* (1881) ii. 180.—Afr. trop.
submersum, *Herb. in Bot. Mag.* t. 2468.—Bras.
sumatranum, *Roxb. Hort. Beng.* [38] ; *Fl. Ind.* ii. 131. —Sumatra.
superbum, Roxb. ll. cc. [38] 133 = amabile.
tatinense, [DC.] in Red. Lil. t. 408 = pedunculatum.
tenellum, Jacq. Coll. Suppl. 43 = Carpolyza spiralis.
tenellum, Linn. f. Suppl. 194 = Strumaria filifolia.
Tinneae, Kotschy & Peyr. in Flora, ii. (1858) 508 = seq.
Tinneanum, *Kotschy & Peyr. Pl. Tinn.* 44. t. 21.—Afr. trop.
toxicarium, Roxb. Hort. Beng. [23] ; Fl. Ind. ii. 134 = asiaticum.
umbellatum, Carey, ex Herb. in Bot. Mag. sub t. 2121. p. 7 = amoenum.
undulatifolium, Herb. l. c. p. 8 = scabrum.
undulatum, *Herb. Exot. Fl.* t. 200.—Bras.
uniflorum, *F. Muell. Fragm.* iii. 23.—Austral.
urceolatum, Ruiz & Pav. Fl. Per. iii. 58. t. 287. f. b = Urceolina pendula.
vanilladorum, Walw. ex Baker, in Journ. Bot. xvii. (1878) 196 = giganteum.
variabile, *Herb. Amaryll.* 268. t. 44. f. 28.—Afr. austr.
venosum, *R. Br. Prod.* 297.—Austral.
venustum, *Herb. Carey, ex Steud. Nom.* ed. II. i. 440. —Ind. or.
verecundum, Carey, ex M. Roem. Syn. Enast. 75 = amoenum.
virgineum, *Mart. in Schult. f. Syst.* vii. 865.—Bras.

CRINUM :—
viridifolium, M. Roem. Syn. Enast. 78 = Commelyni.
Wallichianum, M. Roem. Syn. Enast. 85 = zeylanicum.
Weinmanni, M. Roem. l. c. 80 = flaccidum.
Welwitschii, *Baker, in Gard. Chron.* (1884) ii. 40.—Afr. trop.
yuccæflorum, *Jacob. Parad. Lond.* t. 52.—Afr. trop.
yuccoides, Herb. Bot. Mag. sub t. 2121 = yuccæflorum.
zeylanicum, *Linn. Syst.* ed. XII. 286.—Ind. or.

CRIOSANTHES, Rafin. in Am. Monthly Mag. (1818) 268 = **Cypripedium**, Linn. (Orchid.).
borealis, Rafin. l. c. = Cyp. arietinum.
parviflora, Rafin. Fl. Tellur. iv. 46 = prasc. ?

CRISTA-GALLI, Rupp. Fl. Jen. ed. Hall. 240 (1745) = **Rhinanthus**, Linn. (Scroph.).

CRISTARIA, [Heist. Syst. 19 (1748)] Cav. Ic. v. 10 t. 418 (1799). *MALVACEÆ*, Benth. & Hook. f. i. 202.
andicola, *C. Gay, Fl. Chil.* i. 317.—Chili.
argyliæfolia, *Phil. in Linnæa*, xxxiii. (1864-65) 29.—Am austr.
aspera, *C. Gay, Fl. Chil.* i. 320.—Chili.
betonicæfolia, *Cav. Ic.* v. 11, in obs. ; *Pers. Syn.* ii. 248.—Chili.
bipinnata, *Phil. in Linnæa*, xxxiii. (1864-65) 31.—Am. austr.
cocciuea, Pursh, Fl. Am. Sept. ii. 453 = Malvastrum coccineum.
concinna, *Phil. Fl. Atac.* 11.—Chili.
corchoriifolia, *Griseb. in Goett. Abh.* xix. (1874) 91.—Reg. Argent.
cordato-rotundifolia, *C. Gay, Fl. Chil.* i. 328.—Chili.
dissecta, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 153.—Chili.
diversifolia, *Phil. in Anal. Univ. Chil.* (1870) ii. 165.—Chili.
ecristata, *A. Gray, Bot. U. St. Expl. Exped.* i. 165.—Bras.
elegans, *C. Gay, Fl. Chil.* i. 325.—Chili.
ertantha, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 152.—Chili.
foliosa, *Phil. Fl. Atac.* 11.—Chili.
geranifolia, *Presl, Rel. Haenk.* ii. 119.—Chili.
glanchiosa, *Phil. in Linnæa*, xxxiii. (1864-65) 31.—Am. austr.
glaucophylla, *Cav. Ic.* v. 11. t. 418.—Chili.
gracilis, *C. Gay, Fl. Chil.* i. 326.—Chili.
grandiflora, Turcz. in Bull. Soc. Nat. Mosc xxxvi. (1863) i. 564 = eriantha.
Haenkeana, *Presl, Rel. Haenk.* ii. 120.—Chili.
heterophylla, *Hook & Arn. in Hook. Bot. Misc.* iii (1833) 153.—Chili.
heterophylla, Phil. Fl. Atac. 11 = diversifolia.
hirsuta, *Presl, Rel. Haenk.* ii. 119.—Chili.
hispida, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 153 = heterophylla.
insularis, *Phil. in Anal. Univ. Chil.* (1875) 187.—Chili.
integerrima, *Phil. Fl. Atac.* 11.—Chili.
intermedia, *C. Gay. Fl. Chil.* i. 319.—Chili.
lata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 565.—Chili.
loasæfolia, *Phil. in Linnæa*, xxxiii. (1864-65) 28.—Chili.
lobulata, *Phil. Fl. Atac.* 11.—Chili.
Mollinæ, *C. Gay, Fl. Chil.* i. 327.—Chili.
multifida, *Cav. Ic.* v. 13, in obs.—Chili.
multiflora, *C. Gay, Fl. Chil.* i. 321.—Chili.
obtusiloba, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 564 = multifida.
Ovalles, *C. Gay, Fl. Chil.* i. 320.—Chili.
patagonica, *Phil. in Linnæa*, xxxiii. (1864-65) 28.—Patagon.
pinnata, *Phil. in Anal. Univ. Chil.* (1872) 692.—Chili.
pinnatifida, *Hook. & Arn. Bot. Beech. Voy.* 13.—Chili.
Sadæ, *Phil. in Linnæa*, xxxiii. (1864-65) 30.—Chili.
semilifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 197.—Chili.
Spinolæ, *C. Gay, Fl. Chil.* i. 326.—Chili.
trichocaula, *Phil. in Anal. Univ. Chil.* (1872) 692.—Chili.
Urmenetæ, *Phil. l. c.* 681.—Chili.
virgata, *C. Gay, Fl. Chil.* i. 322.—Chili.
viridi-lutcola, *C. Gay, l. c.*—Chili.

CRISTARIA, Sonner. Voy. Ind. iii. 284 (1782) = Combretum, Linn.
coccinea, Sonner. l. c. = Combr. purpureum.

CRISTATHELIA, Nutt. in Journ. Acad. Philad. vii. (1834) 35. t. 11. *CAPPARIDEÆ*, Benth. & Hook. f. i. 105.
CYRBARIUM, Endl. Gen. 891 (1838).
DISPARA, Raffin. Sylva Tellur. 112 (1838).
cross, *Nutt. l. c.* 36. t. 11.—Am. bor.
Jamesii, *Torr. & Gray, Fl. N. Am.* i. 124.—Am. bor.

CRITAMUS, Bess. Enum. Pl. Volh. 93 (1821) = **Falcaria**, Host (Umbell.).
agrestis, Bess. l. c. = F. Rivini.
Falcaria, Reichb. in Moesl. Handb. ed. II. i. 478 = F. Rivini.
heterophyllus, Mert. & Koch, in Roehl. Deutschl. Fl. ii. 441 = Carum Bunius.
saxifragus, Reichb. in Moesl. Handb. ed. II. i. 478 = Carum Bunius.

CRITAMUS, Hoffm. Gen. Umb. ed. II. 182 (1816) = **Aptum**, Tourn. (Umbell.).
daucicus, Hoffm. l. c. 184 = A. lineare.

CRITESION, Rafin. in Journ. Phys. lxxxix. (1819) 103 = **Hordeum**, Linn. (Gramin.).
geniculatum, Rafin. l. c. = H. jubatum.

CRITESIUM, Endl. Gen. 104 (1838) = prec.

CRITHMUM, Linn. Syst. ed. I (1735); Gen. ed. I. 74 (1737). *UMBELLIFERÆ*, Benth. & Hook. f. i. 905.
campestre, Gueld. ex Bieb. Fl. Taur. Cauc. iii. 215 (= *Crenolophium Fischeri*).—Reg. Caucas.
canariense, Cav. in Anal. Clenc. iii. (1801) 35 = maritimum.
latifolium, Linn. f. Suppl. 180 = Astydamia canariensis.
Libanotis, Link, Enum. Hort. Berol. i. 270 = Hippomarathrum Libanotis.
maritimum, *Linn. Sp. Pl.* 246.—Europ.
mediterraneum, Bieb. Fl. Taur. Cauc. iii. 215 (= *Crenolophium Fischeri*).—Reg. Caucas.
pyrenaicum, Forsk. ex Delile, Illustr. Fl. Egyptæ, 10 = Deverra tortuosa.
pyrenaicum, Linn. Sp. Pl. 246 = Seseli Libanotis.
Silaus, Wibel, Prim. Fl. Werth. 197 = Silaus pratensis.

CRITHO, E. Mey. Ind. Sem. Hort. Regiom. (1848) 5 = **Hordeum**, Linn. (Gramin.).
Agricrea, E. Mey. l. c. = H. Agricrea.

CRITHODIUM, Link, in Linnæa, ix. (1834) 132. t. 3 = **Triticum**, Tourn. (Gramin.).
aegilopoides, Link, l. c. = T. monococcum ?
aegypticum, Trin. ex Steud. Nom. ed. II. i. 440 = T. bicorne.

CRITHOPSIS, Jaub. & Spach, Illustr. iv. 80. t. 321 (1850) = **Elymus**, Linn. (Gramin.).
brachystricha, Walp. Ann. iii. 756 = seq.
rhachistricha, Jaub. & Spach, Illustr. iv. 80 = E. Delileanus.

CRITHOPYRUM, Hort. Prag. ex Steud. Syn. Fl. Gram. 344 (1855) = **Agropyron**, J. Gaertn. (Gramin.).
trachycaulum, Hort. Prag. ex Steud. l. c. = A. Richardsonii ?

CRITONIA, P. Br. Hist. Jamaic. 490 (1756); DC. Prod. v. 140 = **Eupatorium**, Tourn. (Compos.).
Dalea, DC. Prod. v. 140 = E. Dalea.
daleoides, DC. l. c. 141 = E. daleoides.
elliptica, Rafin. ex DC. l. c. 178 = E. rotundifolium.
eupatoroides, Gaertn. ex Steud. Nom. ed. II. i. 441 = Kuhnia eupatorioides.
hebebotrya, DC. Prod. v. 141 = E. hebebotrya.
heterosyra, Ernst, in Flora, lvii. (1874) 210.—Venezuela.
imbricata, Grisch. in Mem. Am. Acad. N. S. viii. (1862) 513.—Ind. occ.
Kuhnia, Gaertn. Fruct. ii. 411. t. 174. f. 7 = Kuhnia eupatorioides.
macropoda, DC. Prod. v. 140 = E. macropodum ?
parviflora, DC. l. c. = E. parviflorum.
Pseudodalea, DC. l. c.—Cuba.

CRITONIOPSIS, Sch. Bip. in Pollichia, xx-xxi. (1863) 480 = **Vernonia**, Schreb. (Compos.).
Lindenii, Sch. Bip. l. c. 481 = V. pycnantha.

CROCANTHEMUM, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 370 = **Hallanthemum**, Tourn. (Cistin.).
brasiliensis, Spach, l. c. = H. brasiliense.
canadense, Spach, l. c. = H. canadense.

CROCANTHUS, Klotzsch, ex Klatt, in Peters, Reise Mossamb. Bot. 516 (1868) = **Crocosmia**, Planch. (Irid.).
massambicensis, Klotzsch, ex Klatt, l. c. = Crocosmia aurea.

CROCIDIUM, Hook. Fl. Bor. Am. i. 285. t. 118 (1833). *COMPOSITÆ*, Benth. & Hook. f. iii. 440.
multicaule, *Hook. l. c.*—Am. bor. occ.

CROCIRIS, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 78 (Sert. Pl. Transs.) = **Crocus**, Tourn. (Irid.).
iridiflora, Schur, l. c. = Crocus byzantinus.
speciosa, Schur, l. c. = Crocus byzantinus.

CROCISERIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = **Senecio**, Tourn. (Compos.).
Doronicum, Fourr. l. c. = S. Doronicum.
Gerardi, Fourr. l. c. = S. Gerardi.

CROCKERIA, Greene, ex A. Gray, Syn. Fl. N. Am. i. 11. 445 (1884). *COMPOSITÆ*.
chrysantha, *Greene, ex A. Gray, l. c.*—Calif.

CROCODEILANTHE, Reichb. f. & Warcz. in Bonplandia, ii. (1854) 113 = **Pleurothallis**, R. Br. (Orchid.).
xiphiosa, Reichb. f. & Warcz. l. c. 114 = P. Crocodilanthe.

CROCODILIUM, [Vaill.] Juss. Gen. 173 (1789) = **Centaurea**, Linn. (Compos.).
aureum, Sweet, Hort. Brit. ed. I. 217 = Cent. aurea.
Crocodylium, Hill, Hort. Kew. 68 = Cent. Crocodilium.
cuculiatum, Sweet, Hort. Brit. ed. I. 217 = Cent. Crocodilium.
Galactites, Dum.-Cours. ex Steud. Nom. ed. II. i. 440 = Galactites tomentosa.
muricatum, Sweet, Hort. Brit. ed. I. 217, nomen.—Hispan.
peregrinum, Hill, Hort. Kew. 68 = Cent. peregrina.
salmanticum, Sweet, Hort. Brit. ed. I. 217 = Cent. salmantica.
syriacum, Cass. in Dict. Sc. Nat. xli. 19 = Cent. Crocodilium.
Verutum, Sweet, Hort. Brit. ed. I. 217 = Cent. Verutum.

CROCODILOIDES, Adans. Fam. ii. 127 (1763) = **Berkhaya**, Ehrh. (Compos.).

CROCODYLIUM, Hill, Hort. Kew. 68 (1769) = CROCODILIUM, Vaill. = **Centaurea**, Linn. (Compos.)

CROCOSMA, Klatt, in Peters, Reise Mossamb. Bot. 516 (1868) = seq.

CROCOSMIA, Planch. in Fl. des Serres, Sér. I. vii. (1851–52) 161. *IRIDEÆ*, Benth. & Hook. f. iii. 707.
CROCANTHUS, Klotzsch, ex Klatt, in Peters, Reise Mossamb. Bot. 516 (1868).
aurea, *Planch. l. c.*—Afr. austr. et trop.
crocata, Planch. l. c., in obs. = Tritonia crocata.
densa, Planch. l. c., in obs. = Tritonia densa.
fenestrata, Planch. l. c., in obs. = Tritonia fenestrata.
longiflora, Klatt, in Peters, Reise Mossamb. Bot. 516 = Ixia longiflora.
minutata, Planch. in Fl. des Serres, Sér. I. vii. (1851–52) 161, in obs. = Tritonia minutata.
Rochenaia, Klatt, in Peters, Reise Mossamb. Bot. 516 = Tritonia Rochenaia.
transiflora, Klatt, l. c. = Ixia paniculata, longiflora.

CROCOXYLON, Eckl. & Zeyh. Enum. 138 (1835) = **Monodendrum**, Jacq. (Tiliac.).
excelsum, Eckl. & Zeyh. l. c. = E. crocosum.

CROCUS, [Tourn.] Linn. Syst. ed. I (1735). *IRIDEÆ*, Benth. & Hook. f. iii. 692.
CROCIRIS, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 78 (Sert. Pl. Transs.).
SAFFRAN, Medic. in Act. Theod. Palat. vi. Phys. (1790) 472.
acutiflorus, Seidl, in Opiz, Natural. n. ix. (1825) 128 = vernus.
Adami, J. Gay, in Mém. Bull. Férussac, xxv. (1831) 319 (err. typ. 219) = biflorus.
aerius, *Herb. in Journ. Hort. Soc.* ii. (1847) 283.—Persia.
alatavicus, *Regel & Semenow, in Bull. Soc. Nat. Mosc.* xli. (1868) 1. 434.—Siberia.
albiflorus, Kit. ex Schult. Oestr. Fl. ed. II. i. 101 = vernus.
alopecicus, Baker, in Gard. Chron. (1873) 609 = Gaillardotii.
algarvensis, Baker, l. c. (1877) 1. 45 = nevadensis.
anocyrensis, *Maw, l. c.* (1881) 11. 528.—As. Min.
annulatus, Herb. in Bot. Mag. t. 3868 = chrysanthus.
annulatus, Steph. ex Spreng. & Link, Jahrb. i. 111. 69 = biflorus.
aphyllus, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 223 = nudiflorus.
argenteus, Sabine, in Trans. Hort. Soc. vii. (1830) 481 = biflorus.
asturicus, *Herb. in Bot. Mag.* sub t. 3998.—Hispan.
atlanticus, Pomel, Nouv. Mat. Fl. Alt. 585 = nevadensis.
atticus, Boiss. Diagn. Ser. II. iv. 94 = Sieberi.
atticus, *Orph. in Bull. Congr. Bot. Pétersb.* 1869 (1870) 118.—Graecia.
Aucheri, Boiss. Diagn. Ser. I. xiii. 16 = Suterianus.
aureus, Clarke, Trav. ii. 146 = gargaricus.
aureus, *Sibth. & Sm. Fl. Graec. Prod.* i. 54.—Europ.
autumnalis, Brot. Phytogr. ii. 40. t. 94 = Clusii.
autumnalis, Ker-Gawl. Irid. Gen. 77 = serotinus.
autumnalis, *Mill. Gard. Dict.* ed. VIII. n. 2.—Europ.
autumnalis, Sm. Engl. Bot. t. 848; Bieb. Fl. Taur. Cauc. i. 27 = sativus.
autumnalis, Vis. ex Steud. Nom. ed. II. i. 441 = longiflorus.
Balansae, *J. Gay, ex Maw, in Gard. Chron.* (1879) 1. 394.—Oriens.
balcanensis, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 181 = vallicolens.
banaticus, J. Gay, in Bull. Férussac, xxv. (1831) 320 (err. typ. 290) = byzantinus.
banaticus, *Heuff. in Flora*, xviii. (1835) 235.—Hungar.
biflorus, *Mill. Gard. Dict.* ed. VIII. n. 4.—Europ.; Taur.
Biliotti, *Maw, in Gard. Chron.* (1881) ii. 303.—Pontus.
Boissieri, *Maw, l. c.* 304.—Oriens.
Boryanus, Herb. in Bot. Reg. (1843) Misc. 28 = Boryi.
Boryi, *J. Gay, in Bull. Férussac*, xxv. (1831) 320 (err. typ. 220).—Graecia.
Bulbocodium, Linn. Sp. Pl. 36 = Romulea Bulbocodium.
Bulbocodium, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. i. 28 = reticulatus.
byzantinus, *Ker-Gawl. in Bot. Mag.* sub t. 1111.—Europ.
Cambessedesianus, Herb. in Bot. Reg. (1843) Misc. 33 = seq.
Cambessedesii, *J. Gay, in Bull. Férussac*, xxv. (1831) 320 (err. typ. 220).—Ins. Majoras.
Cambessedesii, Klatt, in Linnaea, xxxiv. (1865–66) 684 = cyprius.
Cambessedianus, Herb. in Bot. Reg. (1843) t. 37. f. 4 = Cambessedesii.
campestris, Herb. l. c. (1843) Misc. 30 = sativus.
cancellatus, *Herb. in Bot. Mag.* sub t. 3864.—As. Min.
candidus, Boiss. Diagn. Ser. I. xiii. 17 = Fleischeri.
candidus, *Clarke, Trav. ii.* 146.—As. Min.
candidus, Schloss. & Vukot. Syll. Fl. Croat. 22 = banaticus.
carpaticus, Herb. f. Fl. Cap. Prod. 2 = Romulea rosea.
carpetanus, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 24. —Hispan.
Cartwrightianus, Herb. in Bot. Reg. (1843) Misc. 32 = sativus.
cashmerianus, Royle, Illustr. Bot. Himal. 99 = sativus.
cassius, *Sieb. & Hesp. ex Nohora. in Bull. Soc. Nat. Mosc.* (1856) 332.—Reg. Casp.

CROCUS :—
chrysanthus, Herb. in Bot. Reg. (1843) Misc. 83 = Soteriana.
chrysanthus, *Herb. in Journ. Hort. Soc.* ii. (1847) 285. —Rumelia.
cilicius, Kotschy, ex Baker, in Journ. Linn. Soc. xvi. (1877) 84 = Schimperi.
circumscissus, Haw. in Trans. Hort. Soc. i. (1812) 137 = biflorus.
Clusianus, Herb. in Bot. Reg. (1845) t. 37. f. 8 = Clusii.
Clusii, *Y. Gay, in Bull. Férussac*, xxv. (1831) 320 (err. typ. 230).—Lusitan.
corsicus, *Vanucc. ex Maw, in Gard. Chron.* (1878) ii. 367.—Ins. Corsica.
crestensis, Nym. Consp. 707 = versicolor.
cretensis, Koern. in Flora, xxxix. (1856) 469 = Veneris.
Crewii, *Hook f. Bot. Mag.* t. 6168.—Archip. Graec.
cristensis, Eugène, in Bull. Soc. Bot. Fr. xv. (1868) 191 = versicolor.
croceus, C. Koch, in Linnaea, xix. (1847) 7 = chrysanthus.
cyprius, *Boiss. & Kotschy, in Unger & Kotschy, Ins. Cyp.* 203.—Ins. Cyprus.
dalmaticus, *Vis. Fl. Daim.* i. 119.—Dalmat.
damascenus, Herb. in Bot. Reg. (1845) Misc. 1 = cancellatus.
Dandordiae, *Maw, in Gard. Chron.* (1881) ii. 781.—Ac. Min.
dianthus, C. Koch, in Linnaea, xxi. (1848) 684 = cancellatus.
discolor, Reuss, Květ. Slov. (1858) 418 = reticulatus.
edulis, Boiss. & Blanch. ex Boiss. Fl. Orient. v. 101 = cancellatus.
etruscus, *Pari. Fl. Ital.* iii. 228.—Italia.
exiguus, Schur, Enum. Pl. Transyl. 652 = banaticus.
fatuus, *Moddic. in Act. Acad. Theod. Palat.* vi. *Phys.* (1790) 472.—Europ.
fimbriatus, Lapeyr. Hist. Abr. Pl. Pyr. 22 = nudiflorus.
flavus, Haw. in Trans. Hort. Soc. i. (1812) 135 = aureus.
Fleischeri, Baker, in Gard. Chron. (1873) 650 = candidus.
Fleischeri, *Y. Gay, in Bull. Férussac*, xi. (1827) 319 (err. typ. 219).—As. Min.
Fleischerianus, Herb. in Journ. Hort. Soc. ii. (1847) 278 = Fleischeri.
floribundus, Haw. in Trans. Hort. Soc. i. (1812) 133 = aureus.
Fontenayi, Reut. ex Nym. Consp. 706 = Tournefortii.
fragrans, Haw. in Trans. Hort. Soc. i. (1812) 136 = versicolor.
Fusianus, J. Gay, ex Nym. Consp. 705 = biflorus.
Gaillardotii, *Boiss. ex Maw, in Gard. Chron.* (1879) i. 294.—Orient.
gargaricus, *Herb. in Bot. Mag.* (1841) sub t. 3866.— Ac. Min.
grandiflorus, Hegetschw. Fl. Schw. 41 = vernus.
graecus, Chapp. in Bull. Soc. Bot. Fr. xx. (1873) 193 = sativus.
granatensis, *Boiss. ex Maw, in Journ. Linn. Soc.* xix. (1882) 368, 373.—Hispan.
graveolens, Boiss. & Reut. in Boiss. Fl. Orient. v. 107 = vitellinus.
hadriaticus, *Herb. in Bot. Reg.* (1845) *Misc.* 82.— Europ. or.
Heuffeliana, Boiss. & Reut. ex Boiss. Fl. Orient. v. 100 = sativus.
hermoneus, *Kotschy, ex Maw, in Gard. Chron.* (1881) ii. 559.—Syria.
Herbertianus, Koern. in Ind. Sem. Hort. Berol. App. (1856) 15 ; et in Walp. Ann. vi. 51 = byzantinus.
Heuffelianus, Herb. in Journ. Hort. Soc. ii. (1847) 278 = banaticus.
Heuffelii, Koern. in Flora, xxxix. (1856) 476 = praec.
hybernus, *Fisv.* in Gribb. Spicil. Fl. Rumel. ii. 374 = sativus.
hyemalis, *Boiss. & Blanche, Diag.* Ser. II. iv. 93.— Syria.
imperati, Colla, Hort. Ripul. App. iv. 19 = suaveolens.
imperati, *Tenore, Mem. Crochi Fl. Nap.* 10. t. 3.— Ital.
Imperatoreus, Herb. in Bot. Mag. sub t. 3998 = Imperati.
incurvatus, *Rchbch., in Corresp. Ver. Naturh. Presb.* ii. (1848) 196.—Europ.
incurvus, Donn, ex Sweet, Hort. Brit. ed. II. i. 441 = Imperati.
indivisus, *Richter, in Corresp. Ver. Naturh. Presb.* ii. (1848) 196.—Europ.
insularis, J. Gay, in Bull. Férussac, xxv. (1831) 321 (err. typ. 231) = minimus.
insularis, Herb. in Bot. Reg. (1843) t. 21 = corsicus.
intermissus, Herb. in Bot. Reg. (1845) Misc. 2 = Gaillardotii.
ionicus, Herb. l. c. 3 = Boryi.
iridiflorus, Heuff. ex Reichb. Ic. Fl. Germ. ix. (1847) 19 = byzantinus.
italicus, Gaud. Fl. Helv. i. 88 = biflorus.
karduchorum, *Kotschy, ex Boiss. Fl. Orient.* v. 99.— Kurdistan.
Kirkii, *Maw, in Gard. Chron.* (1879) i. 284 = candidus.
Korolkowi, *Maw & Regel, in Act. Hort. Petrop.* vi. (1879) 499.—Turkest.
Kotschyanus, Boiss. Fl. Orient. v. 102, in syn. = cancellatus.
Parbinsonii, Steud. Nom. ed. II. i. 441 = biflorus.
parviflorus, *Baker, in Journ. Bot.* xiv. (1876) 266.— Cilicia.
peloponnesiacus, Orph. in Boiss. Diagn. Ser. II. iv. 93 = hadriaticus.
pendulus, Stokes, Bot. Comment. i. 209 = sativus
penicillatus, Steud. ex Baker, in Gard. Chron. (1881) ii. 680 = Fleischeri.
penicillatus, Steud. Nom. ed. II. i. 441 = aureus.
Pestalozzae, Boiss. Diagn. Ser. I. xii. 17 = biflorus.
pholegandrius, Orph. in Bull. Congr. Bot. Petersb. 1869 (1870) 118, 184 = Tournefortii.
pictus, Steud. Nom. ed. II. i. 441 = vernus.
praecox, Haw. in Engl. Bot. Suppl. t. 2645 = biflorus.
pulchellus, *Herb. in Bot. Mag.* sub t. 3862.—Ac. Min.
pulchricolor, Herb. ex Tchihat. Asie Min. Bot. ii. 523 = serius.
pusillus, Tenore, in Mem. Crochi Fl. Nap. 8. t. 1 = biflorus.
pylaeus, J. Gay, ex Tchihat. Asie Min. Bot. ii. 525 = cancellatus.
pyrrenaeus, Herb. in Journ. Hort. Soc. ii. (1847) 255 = nudiflorus.
recurvus, Haw. in Loud. Hort. Brit. Suppl. i. 586 = Imperati.
reflexus, Donn, Hort. Cantab. ed. IV. 11 = Imperati.
Reinwardti, Opiz, Natural. x. (1825) 292 = vernus.
Reinwardti, Reichb. Pl. Crit. x. 14 = versicolor.
reticulatus, *Steven, ex Adam, in Web. & Mohr, Beitr.* i. 45.—Reg. Casc.
revolutus, Haw. in Trans. Hort. Soc. i. (1812) 136 = suaianus.
Rosenianus, C. Koch, in Linnaea, xxi. (1848) 684 = suaianus.
Salzmanni, *Y. Gay, in Bull. Férussac*, xxv. (1831) 320 (err. typ. 220).—Marocc.
Salzmannianus, Herb. in Bot. Mag. sub t. 3868 = praec.
sativus, Bivona, ex Steud. Nom. ed. II. i. 441 = longiflorus.
sativus, *Linn. Sp. Pl.* 36.—Graecia ; As. Min.
sativus, Mill. Gard. Dict. ed. VIII. n. 1 = aureus.
sativus, Tenore, Fl. Med. i. 19 = Thomasii.
Scharojani, *Rupr. in Regel, Gartenfl.* xvii. (1868) 134. —Reg. Caucas.
Schimperi, J. Gay, ex Baker, in Journ. Linn. Soc. xvi. (1877) 84 = cancellatus.
scotinus, Bert. in Nouv. Coll. Opusc. Sc. ii. 150 = longiflorus.
serotinus, *Salisb. Parad. Lond.* t. 30.—Europ. ; As. Min.
setifolius, Stokes, Bot. Mat. Med. i. 104 = sativus.
sibiricus, Barr, in Gard. Chron. (1876) i. 807, 839 = Sieberi.
Sibthorpianus, Herb. in Bot. Reg. (1843) Misc. 83 = Sieberi.
siculus, Tineo, in Guss. Suppl. Fl. Sic. Prod. i. 7 = vernus.
Sieberi, *Y. Gay, in Bull. Férussac*, xxv. (1831) 320 (err typ. 220).—As. Min.
Sieberianus, Herb. in Bot. Mag. sub t. 3866 = praec.
smyrnensis, Poech, Enum. Pl. ins. Cyp. 11, in obs. = Fleischeri.
speciosus, *Bieb. Fl. Taur. Cauc.* i. 37.—Tauria.
speciosus, Griseb. Spicil. Fl. Rumel. ii. 374 = pulchellus.
speciosus, Roche], Pl. Rar. Banat. 5 = byzantinus.
speciosus, Wilson, in Engl. Bot. Suppl. t. 2782 = nudiflorus.
Sprunerі, Boiss. & Heldr. Diagn. Ser. I. vii. 103 = cancellatus.
stellaris, *Haw. in Trans. Hort. Soc.* i. (1812) 136. t. 6. —Europ.
ochroleucus, *Boiss. & Gaill. Diagn.* Ser. II. iv. 93.— Syria.
odorus, Bivona, Strip. Rar. Sicil. Manip. iii. 8 = longiflorus.
odorus, Vis. Strip. Dalm. Spec. 32 = sativus.
officinalis, Hoock. Syn. i. 273 = sativus, vernus.
officinalis, Martyn, Fl. Rustic. ii. t. 56 = sativus.
Olivieri, *Y. Gay, in Bull. Férussac*, xxv. (1831) 319 (err. typ. 219).—As. Min.
Orphanidis, Hook. f. Bot. Mag. t. 5776 = Tournefortii.
Orsinii, Pari. Fl. Ital. iii. 238 = sativus.
Pallasianus, Herb. in Bot. Reg. (1844) t. 5. f. 2 = sativus.
Pallasii, Goldb. in Mém. Soc. Nat. Mosc. v. (1817) 157 = sativus.
Pallasii, Griseb. Spicil. Fl. Rumel. ii. 374 = veluchensis.
Pallasii, Sprun. ex Boiss. Diagn. Ser. I. vii. 103 = cancellatus.
Parbinsonii, Steud. Nom. ed. II. i. 441 = biflorus.
parviflorus, *Baker, in Journ. Bot.* xiv. (1876) 266.— Cilicia.
peloponnesiacus, Orph. in Boiss. Diagn. Ser. II. iv. 93 = hadriaticus.
pendulus, Stokes, Bot. Comment. i. 209 = sativus
penicillatus, Steud. ex Baker, in Gard. Chron. (1881) ii. 680 = Fleischeri.
penicillatus, Steud. Nom. ed. II. i. 441 = aureus.
Pestalozzae, Boiss. Diagn. Ser. I. xii. 17 = biflorus.
pholegandrius, Orph. in Bull. Congr. Bot. Petersb. 1869 (1870) 118, 184 = Tournefortii.
pictus, Steud. Nom. ed. II. i. 441 = vernus.
praecox, Haw. in Engl. Bot. Suppl. t. 2645 = biflorus.
pulchellus, *Herb. in Bot. Mag.* sub t. 3862.—Ac. Min.
pulchricolor, Herb. ex Tchihat. Asie Min. Bot. ii. 523 = serius.
pusillus, Tenore, in Mem. Crochi Fl. Nap. 8. t. 1 = biflorus.
pylaeus, J. Gay, ex Tchihat. Asie Min. Bot. ii. 525 = cancellatus.
pyrrenaeus, Herb. in Journ. Hort. Soc. ii. (1847) 255 = nudiflorus.
recurvus, Haw. in Loud. Hort. Brit. Suppl. i. 586 = Imperati.
reflexus, Donn, Hort. Cantab. ed. IV. 11 = Imperati.
Reinwardti, Opiz, Natural. x. (1825) 292 = vernus.
Reinwardti, Reichb. Pl. Crit. x. 14 = versicolor.
reticulatus, *Steven, ex Adam, in Web. & Mohr, Beitr.* i. 45.—Reg. Casc.
revolutus, Haw. in Trans. Hort. Soc. i. (1812) 136 = suaianus.
Rosenianus, C. Koch, in Linnaea, xxi. (1848) 684 = suaianus.
Salzmanni, *Y. Gay, in Bull. Férussac*, xxv. (1831) 320 (err. typ. 220).—Marocc.
Salzmannianus, Herb. in Bot. Mag. sub t. 3868 = praec.
sativus, Bivona, ex Steud. Nom. ed. II. i. 441 = longiflorus.
sativus, *Linn. Sp. Pl.* 36.—Graecia ; As. Min.
sativus, Mill. Gard. Dict. ed. VIII. n. 1 = aureus.
sativus, Tenore, Fl. Med. i. 19 = Thomasii.
Scharojani, *Rupr. in Regel, Gartenfl.* xvii. (1868) 134. —Reg. Caucas.
Schimperi, J. Gay, ex Baker, in Journ. Linn. Soc. xvi. (1877) 84 = cancellatus.
scotinus, Bert. in Nouv. Coll. Opusc. Sc. ii. 150 = longiflorus.
serotinus, *Salisb. Parad. Lond.* t. 30.—Europ. ; As. Min.
setifolius, Stokes, Bot. Mat. Med. i. 104 = sativus.
sibiricus, Barr, in Gard. Chron. (1876) i. 807, 839 = Sieberi.
Sibthorpianus, Herb. in Bot. Reg. (1843) Misc. 83 = Sieberi.
siculus, Tineo, in Guss. Suppl. Fl. Sic. Prod. i. 7 = vernus.
Sieberi, *Y. Gay, in Bull. Férussac*, xxv. (1831) 320 (err typ. 220).—As. Min.
Sieberianus, Herb. in Bot. Mag. sub t. 3866 = praec.
smyrnensis, Poech, Enum. Pl. ins. Cyp. 11, in obs. = Fleischeri.
speciosus, *Bieb. Fl. Taur. Cauc.* i. 37.—Tauria.
speciosus, Griseb. Spicil. Fl. Rumel. ii. 374 = pulchellus.
speciosus, Roche], Pl. Rar. Banat. 5 = byzantinus.
speciosus, Wilson, in Engl. Bot. Suppl. t. 2782 = nudiflorus.
Sprunerі, Boiss. & Heldr. Diagn. Ser. I. vii. 103 = cancellatus.
stellaris, *Haw. in Trans. Hort. Soc.* i. (1812) 136. t. 6. —Europ.

CROCUS:—
striatus, Link, Enum. Hort. Berol. i. 48 = minimus, biflorus.
suaveolens, *Bertol. Desc. Zaffer. Ital.* n. 2.—Ital.
subapamimus, *Herb. in Journ. Hort. Soc.* ii. (1847) 259.—Europ.
sublimis, Herb. in Bot. Reg. (1845) Misc. 81 = Sieberi.
sulphureus, Griseb. Spicil. Fl. Rumel. ii. 379 = chrysanthus.
sulphureus, *Ker-Gawl. in Bot. Mag.* t. 935.—As. Min.
sulphureus, Klatt, in Linnaea, xxxiv. (1865–66) 676 = Olivieri.
sustanus, Heyne, ex Steud. Nom. ed. II. i. 441 = anlphureus.
susianus, *Ker-Gawl. in Bot. Mag.* t. 652.—As. Min.
Suterianus, *Herb. in Bot. Reg.* (1845) *Misc.* 5.—As. Min.
Swanenburgianus, C. Koch, in Linnaea, xxi. (1848) 633 = vallicola.
syriacus, Boiss. & Gaill. Diagn. Ser. II. iv. 94 = vitellinus.
syriacus, Tchihat. Asie Min. Bot. ii. 522 = Balansae.
Tauri, *Maw, in Gard. Chron.* (1881) ii. 749.—Lycia.
tauricus, Stev. ex Nym. Consp. 706 = speciosus.
thessalus, Boiss. Diagn. Ser. I. xiii. 17 = Sieberi.
Thirbanus, C. Koch, in Linnaea, xxi. (1848) 633 = gargaricus.
Thomasianus, Herb. in Bot. Reg. (1844) t. 3. f. 6 = sativus.
Thomasii, Tenore, in Mem. Crochi Fl. Nap. 12. t. 1 = sativus.
tingitanus, Herb. in Bot. Mag. sub t. 3866 = Salzmanni.
Tomasinianus, *Herb. in Journ. Hort. Soc.* ii. (1847) 273.—Dalmatia.
Tournefortianus, Herb. in Bot. Reg. (1845) t. 37. f. 3 = Tournefortii.
Tournefortii, *Y. Gay, in Bull. Férussac*, xxv. (1831) 290 (err. typ. 230).—Europ. austr.
trifloru, *Burm. f. Fl. Cap Prod.* 2.—Afr. austr.
uniflorus, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 90 = banaticus.
vallicola, Herb. in Bot. Reg. (1845) Misc. 81 = seq.
vallicola, *Herb. l. c. Misc.* 2.—Persia.
variegatus, Hoppe & Horn. Tageb. 127 = reticulatus.
velutchensis, *Herb. in Bot. Reg.* (1845) *Misc.* 80.—Graecia.
velutchensis, Schott, in Bot. Zeit. ix. (1851) 281 = banaticus.
Veneris, *Tappein, ex Poech, Enum. Pl. Ins. Cyp.* 10.—Ins. Cyprus.
vernus, *All. Fl. Pedem.* i. 84.—Europ.
vernus, Curt. Bot. Mag. t. 45 = aureus.
vernus, Desf. Fl. Atlant. i. 32 = nevadensis.
vernus, Gmelinest. ex Ledeb. Fl. Ross. iv. 109 = biflorus.
vernus, Sibth. & Sm. Fl. Graec. Prod. i. 24 = Sieberi.
vernus, Sieber, ex Tenore, Mem. Crochi Fl. Nap. 5. t. 1 = minimus.
vernus, Uechs, Hort. Reg. Panhorm. 46 = longiflorus.
versicolor, Barcelo, Fl. Isl. Balear. 449 = Cambessedesii.
versicolor, *Ker-Gawl. in Bot. Mag.* t. 1110.—Europ.
Visianicus, Herb. in Bot. Reg. (1845) Misc. 85 = sativus
Visianicus, Herb. in Journ. Hort. Soc. ii. (1847) 263 = longiflorus.
vitellinus, *Wahlenb. in Isis*, xxi. (1828) 1000.—Syria.
vittatus, Schloss. & Vukot. Syll. Fl. Croat. 32 = banaticus.
Weldeni, [Hoppe & Foerst. in] Flora, xxiii. (1840) 208 = biflorus.
zonatus, *Y. Gay, in Tchihat. Asie Min. Bot.* ii. 525.—As. Min.

CROCYLLIS, *E. Mey. Zwei Pfl. Docum.* 276 (1848). *RUBIACEAE*, Benth. & Hook. f. ii. 126.
asthosperuoides, *E. Mey. l. c.*—Afr. austr.

CRODISPERMA, Poit. ex Cass. in Dict. Sc. Nat. xlvi. 408 (1827) = Wulffia, Neck. (Compos.).
aspera, Poit. ex Cass. l. c. = W. platyglossa.

CROLOCOS, Rafin. Fl. Tellur. iii. 98 (1836) = Salvia, Tourn. (Labiat.).
cerea, Rafin. l. c. = S. serea.
calicina, Rafin. l. c. = S. calycina.
colorata, Rafin. l. c. = S. serea.
pomifera, Rafin. l. c. = S. pomifera.

CRONYXIUM, Rafin. Fl. Tellur. ii. 28 (1836) = Lloydia, Salisb. (Liliac.).
serotinum, Rafin. l. c. = Ll. alpina.

CROOMIA, Torr. ex Torr. & Gray, Fl. N. Am. i. 663 (1840). *ROXBURGHIACEAE*, Benth. & Hook. f. iii. 747.
TUBERYA, Croom, ex Meisn. Gen. Comm. 340 (1848).
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 138.—Japon.
pauciflora, *Torr. ex Torr. & Gray, Fl. N. Am.* i. 663.—Florida.

CROPTILON, Rafin. Fl. Tellur. ii. 47 (1836) = Aplopappus, Cass. (Compos.).
divaricatum, Rafin. l. c. = A. divaricatus.

CROSPERMA, Rafin. Fl. Tellur. ii. 33 (1836) = Zygadenus, Michx. (Liliac.).
angustifolia, Rafin. l. c. = Z. angustifolius.
lanta, Rafin. l. c. iv. 37 = Z. muscitoxicus.
phalangioides, Rafin. l. c. 33 = Z. muscitoxicus.

CROSSANDRA, Salisb. Parad. Lond. t. 12 (1806). *ACANTHACEAE*, Benth. & Hook. f. ii. 1094.
HARRACHIA, Jacq. f. Eclog. i. 33. t. 21 (1816).
POLYTHRIX, Nees, in DC. Prod. xi. 393 (1846).
axillaris, Nees in Wall. Pl. As. Rar. iii. 93 = undulaefolia.
cercinna, Dalz. & Gibs. Bomb. Fl. Suppl. 71 = undulaefolia.
diffusa, Bech.-Ham. ex Wall. Cat. n. 2388 = Hemigraphis elegans.
fasciculata, Benth. Pl. Hartw. 21 = Stenandrium dulce.
flava, *Hook. Bot. Mag.* t. 4710.—Afr. trop.
Greenstockii, *S. Moore, in Journ. Bot.* xviii. (1880) 87.—Afr. austr.
guineensis, *Nees, in DC. Prod.* xi. 381.—Afr. trop.
Hoenheuene, Nees, l. c. = Aphelandra verticillata.
infundibuliformis, Nees, in Wall. Pl. As. Rar. iii. 98 = undulaefolia.
madagascariensis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 83.—Madag.
nilotica, *Oliver, in Trans. Linn. Soc. xxix.* (1875) 128.—Afr. trop.
nutans, Wight, ex Nees, in DC. Prod. xi. 380 = undulaefolia.
oppositifolia, Wight, ex Nees, in Wall. Pl. As. Rar. iii. 98 = undulaefolia.
pubescens, *Klotzsch, in Peters, Reise Mossamb. Bot.* 214.—Afr. trop.
pubescens, *Klotzsch, l. c.* 213.—Afr. trop.
spicata, *Buch.-Ham. in Wall. Cat.* sub n. 2341.—Ind. or.
undaefolia, *Salisb. Parad. Lond.* t. 12.—Ind. or.

CROSSOCOMA, Hook. Lond. Journ. Bot. vii. (1848) 392 = Crossosoma, Nutt. (Dillen.).

CROSSOLEPIS, Benth. in Enum. Pl. Hueg. 61 (1837) = Angianthus, Wendl. (Compos.).
pusilla, Benth. l. c. = A. pusillus.
pusilla, Hook. fc. Fl. t. 413 = A. tenellus.

CROSSOLEPIS, Less. Syn. Comp. 270 (1832) = Gnephosis, Cass. (Compos.).
brevifolia, A. Gray, in Hook. Kew Journ. iii. (1851) 175 = G. brevifolia.
eriocephala, A. Gray, l. c. 176 = G. eriocephala.
tinifolia, Less. Syn. Comp. 270.—Austral.
tygmaea, A. Gray, in Hook. Kew Journ. iii. (1851) 176 = G. pygmaea.

CROSSONEPHELIS, Baill. Adansonia, xi. (1874) 245. *SAPINDACEAE*.
verticillata, *Baill. l. c.*—Madag.

CROSSOPETALON, Adans. Fam. ii. 224 (1763) = seq.

CROSSOPETALUM, P. Br. Hist. Jamaic. 145 (1756) = Myginda, Linn. (Celastr.).
Cruentanea, Andrz. ex Trautv. in Act. Hort. Petrop. iv. (1884) 20 = G. ciliata.
gentianoides, Roth, Enum. Pl. Phaen. Germ. i. t. 316 = G. ciliata.

CROSSOPHORA, Link, Handb. ii. 498 (1831) = Chrosophore, Neck. (Euphorb.).

CROSSOPHRYS, Klotzsch, in Linnaea, xxiv. (1851) 3 = Cliefkra, Linn. (Ericac.).
brasiliensis, Klotzsch, ex Meisn. in Mart. Fl. Bras. vii. 167 = Cl. brasiliensis.
Chamissonis, Klotzsch, l. c. = Cl. brasiliensis.
guyanensis, Klotzsch, l. c. 170 = Cl. guyanensis.

CROSSOPTERYX, Fenzl, in Nov. Stirp. Dec. Mus. Vind. 45 (1839). *RUBIACEAE*, Benth. & Hook. f. iii. 41.
africana, Baill. Hist. Pl. vii. 430 = Kotschyana.
febrifuga, *Benth. in Hook. Niger Fl.* 361.—Afr. trop.
Kotschyana, *Fenzl, in Nov. Stirp. Dec. Mus. Vind.* 45.—Afr. trop.

CROSSOSOMA, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 150. *DILLENIACEAE*, Benth. & Hook. f. i. 15.
Bigelovii, *S. Wats. in Proc. Am. Acad.* xi. (1876) 122.—Calif.
californicum, *Nutt. in Journ. Acad. Philad. N. S.* i. (1847) 150. t. 22.—Calif.

CROSSOSTEMMA, Planch. ex Benth. in Hook. Niger Fl. 364 (1849). *PASSIFLORAE*, Benth. & Hook. f. i. 813.
laurifolium, *Planch. ex Benth. l. c.* 375.—Afr. trop.

CROSSOSTEPHIUM, Less. in Linnaea, vi. (1831) 220. *COMPOSITAE*, Benth. & Hook. f. ii. 454.
artemisioides, *Less. l. c.*—Ins. Luzon.; China.
turkestanicum, *Regel & Schmalh. in Pl. Nov. Fedsch.* 44.—As. centr.

CROSSOSTIGMA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 339 = Epilobium, Dill. (Onagrac.).
Lindleyi, Spach, l. c. 404 = E. minutum.

CROSSOSTYLIS, Forst. Char. Gen. 87. t. 44 (1776). *RHIZOPHOREAE*, Benth. & Hook. f. i. 681.
TOMOSTYLES, Montr. in Mém. Acad. Lyon, x. (1860) 201.
biflora, *Forst. l. c.* 88. t. 44.—Ins. Soc.
grandiflora, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. xiii. (1870–71) 398.—N. Caled.
Harveyi, *Benth. in Journ. Linn. Soc.* iii. (1859) 77.—Ins. Fiji.
multiflora, *Brongn. & Gris, ex Panch. & Sebert, Not. Bois Nouv. Caléd.* 260.—N. Caled.

CROSSOSTOMA, Spach, Hist. Vég. Phan. ix. 583 (1840) = Sonevola, Linn. (Goodeu.).
lycioides, De Vriese, in Verh. Haarl. x. (1854) 38 = S. spinescens.
oleoides, De Vriese, l. c. 37 = S. spinescens.
spinescens, De Vriese, l. c. 36 = S. spinescens.

CROSSYNE, Salisb. Gen. Pl. Fragm. 116 (1866) = Buphane, Herb. (Amaryll.).
ciliaris, Salisb. l. c. 117 = B. ciliaris.

CROTALARIA, Dill. ex Linn. Gen. ed. I. 218 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 478.
ATOLARIA, Neck. Elem. iii. 31 (1790).
CHRYSOCALYX, Gaill. & Perr. Fl. Senegamb. Tent. 167. t. 49 (1832).
CLAVULA, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 407.
CYSTOLOBIUM, R. Br. in Wall. Cat. in nota sub n. 5482 (1832).
ISCAULON, Rafin. New Fl. Am. ii. 52 (1836).
MARIA-ANTONIA, Parl. Mar. Ant. Nov. Gen. 3 (1854).
PHYLLOCALYX, A. Rich. Tent. Fl. Abyss. i. 166. t. 34 (1847).
QUIROSIA, Blanco, Fl. Filip. ed. II. 393 (1845).
abscondita, *Wight. ex Baker, in Oliver, Fl. Trop. Afr.* iii. 19.—Afr. trop.
abyssinica, *D. Dietr. Syn. Pl.* iv. 982.—Abyss.
acpulcanensis, Humb. & Am. Bot. Beech. Voy. 434 = mayapanensis.
acicularis, *Buch.-Ham. in Wall. Cat.* n. 5206.—As. trop.

CROTALARIA :—
scaminata, *DC. Prod.* ii. 128.—Afr. austr.
scaminata, G. Don. Gen. Syst. ii. 184 = verrucosa.
scutiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 489.—Bras.
scutiflora, Steud. ex A. Rich. Tent. Fl. Abyss. i. 150 = glauca.
segyriaca, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 478.—Aegypt.
sphinx, DC. Prod. ii. 129 = incana.
Spathii, R. Br. ex Baker, in Oliver, Fl. Trop. Afr. ii. 86 = lathyroides.
stata, *Buch.-Ham. ex Roxb. Hort. Beng.* 98 ; *Fl. Ind.* iii. 274.—As. trop.
stata, Wight, ex Steud. Nom. ed. II. i. 442 = rubiginosa.
statipes, *Rafin. New Fl. Am.* ii. 57.— Am. bor.
alba, Linn. Sp. Pl. 716 = Baptisia alba.
sibticaulis, *Franch. in Revoil, Comali*, 26.—Afr. trop.
sibida, *Heyne, ex Roth, Nov. Pl. Sp.* 338.—Ind. or.
sibida, Heyne, ex Wall. Cat. sub n. 5414 = Heynеana.
simoena, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 26.—Afr. trop.
amplexicaulis, Lam. Encyc. ii. 194 = Rafinia perfoliata.
amplexicaulis, Linn. Fl. Afr. Rar. 16 = Rafinia amplexicaulis.
anagyroides, *H. B. & K. Nov. Gen. et Sp.* vi. 404.— Venezuela.
angulata, *Mill. Gard. Dict. ed.* VIII. n. 9.—Honduras.
angulosa, Lam. Encyc. ii. 197 = verrucosa.
angustifolia, Jacq. Hort. Schoenb. ii. 69. t. 219 = Lebeckia sericea.
angustissima, *E. Mey. Comm. Pl. Afr. Austr.* i. 26.— Afr. austr.
anthylloides, D. Don, Prod. Fl. Nep. 241 = calycina.
anthylloides, Lam. Encyc. ii. 195 = sessiliflora.
anthyllopsis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 15.—Afr. trop.
arborescens, Lam. Encyc. ii. 199 = capensis.
arenaria, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 474.— Afr. trop.
arenaria, *Zoll. & Mor. in Nat. an Genesk. Arch. Nederl. Ind.* iii. (1846) 75.—Java.
argentea, Jacq. Hort. Schoenb. ii. 50. t. 220 = Argyrolobium petiolare.
argyraea, *Franch. in Revoil, Comali*, 27.—Afr. trop.
argyraea, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 26.—Afr. trop.
Arnottiana, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 560 = semperflorens.
arvensis, Klotzsch, in Peters, Reise Mossamb. Bot. 58 = podocarpa.
asarifolia, *Rafin. New Fl. Am.* ii. 57.—Am. bor.
Asarina, Berg. Desc. Pl. Cap. 194 = Hallia Asarina.
aspalathoides, *Lam. Encyc.* ii. 202.—Afr. austr.
assamica, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 481.—As. trop.
astragalina, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 155. —Abyss.
atrorubens, *Hochst. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 572.—Afr. trop.
Atusia, Buch.-Ham. ex Steud. Nom. ed. II. i. 442 = sericea.
auriculata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 11.—Malaya.
axillaris, [*Doyaud. in*] *Ait. Hort. Kew.* ed. II. iii. 20. —Afr. trop.
azurea, Eckl. & Zeyh. Enum. 174 = Lotononis azurea.
Balansae, *Micheli, in Mém. Soc. Phys. Genev.* xxviii. (1888) n. VII. 9.—Paraguay.
barbata, *R. Grah. in Wall. Cat.* n. 5394.—Ind. or.
bengalensis, Lam. Encyc. ii. 196 = juncea.
Bernieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 444. —Madag.
Berteriana, DC. Prod. ii. 127 = fulva.
bialata, Schrank, Pl. Rar. Hort. Monac. t. 18 = sagittalis.
bialata, Roxb. Hort. Beng. 54 ; Fl. Ind. iii. 274 = alata.
bifaria, *Linn. f. Suppl.* 323.—Ind. or.
bifaria, Wall. Cat. n. 5399, partim = hirsuta.
bifaria, Herb. Madr. ex Wall. Cat. n. 5381 = nana.
biflora, *Linn. Mant.* ii. 570.—Ind. or. ; Malaya.
bituminosa, Spreng. Syst. iii. 239 = Fagelia bituminosa.
Bojeri, *Sweet, Hort. Brit.* ed. II. 138.—Madag.
brachycarpa, *Benth. in Mart. Fl. Bras.* xv. i. 29.— Bras.
brachystachya, *Benth. in Linnaea*, xxii. (1849) 512.— Bras.

CROTALARIA :—
bracteata, Cham. & Schlecht. in Linnaea, v. (1830) 576 = Schiedeana.
bracteata, *Roxb. Hort. Beng.* 54 ; *Fl. Ind.* iii. 278.— As. trop.
bracteolata, Perr. ex DC. Prod. ii. 128 = Indigofera bracteolata.
bractеosa, Benth. in Ann. Nat. Hist. iii. (1839) 428 = paulina.
brasilе, Schrank, in Syll. Ratisb. ii. (1828) 77 = vitellina.
brevidens, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 585.— Afr. trop.
breviflora, *DC. Prod.* ii. 127.—Bras.
brevipes, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 44 = sessiliflora.
Brownei, Bert. ex DC. Prod. ii. 130 = Saltiana.
buplеurifolia, *Cham. & Schlecht. in Linnaea*, v. (1830) 575.—Mexic.
Burhia, *Buch.-Ham. in Wall. Cat.* n. 5386.—Ind. or.
Burkeana, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 598.—Afr. austr.
Burmanni, *DC. Prod.* ii. 126.—Ind. or.
caerulea, Vell. Fl. Flum. 307 ; vii. t. 107 = Lupinus velutinus vel subsessilis.
caespitosa, Roxb. Hort. Beng. [98] ; Fl. Ind. iii. 269 = linifolia.
calycina, *H. B. & K. Nov. Gen. et Sp.* vi. 405.— Mexic.
calycina, *Schrank, Pl. Rar. Hort. Monac.* t. 12.—Afr. trop.
Campbellii, Arn. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 81 = ramosissima.
candicans, Wight & Arn. Prod. 184 = madurensis.
canescens, Wall. Cat. n. 5415 = ferruginea.
capensis, Baker, in Oliver, Fl. Trop. Afr. ii. 88 = laburnifolia.
capensis, *Jacq. Hort. Vindob.* iii. 26.—Afr. austr.
capitata, *Lindl. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 74.—Reg. Himal.
capitata, Lam. Encyc. ii. 195 = reflexa, Priestleya Candolleana.
carinata, Steud. ex A. Rich. Tent. Fl. Abyss. i. 154 = maxillaris.
Carmioli, *Polakowus. in Linnaea*, xli. (1877) 558.— Costa Rica.
caudata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 18.—Afr. trop.
Cephalotes, Herb. Madr. ex Wall. Cat. n. 5378 = speciosa.
Cephalotes, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 156. —Abyss.
Chincia, Buch.-Ham. ex Wall. Cat. sub n. 5437 = medicaginea.
chinensis, Lam. Encyc. ii. 195 = paniculata.
chinensis, *Linn. Syst. ed.* X. 1165.—China.
chlorantha, Roxb. Hort. Beng. 54 ; Fl. Ind. iii. 268 = hirta.
ciliata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 241 = prostrata.
cistoides, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 18.—Afr. trop.
Claussenii, *Benth. in Mart. Fl. Bras.* xv. i. 28.—Bras.
clavata, Roxb. ex Wight & Arn. Prod. 188 = hirsuta.
clavata, *Wight & Arn. l. c.* 194.—Ind. or.
cleomifolia, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 48.—Afr. trop.
cleomoides, Klotzsch, in Peters, Reise Mossamb. Bot. 59 = lanceolata.
coerulea, *Bedd. in Madr. Journ. Sc.* Ser. II. xix. (1858) 179.—Ind. or.
coerulea, Jacq. Coll. i. 67 = verrucosa.
columnaris, Lam. Encyc. ii. 200 = purpurea.
comosa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 34.—Afr. trop.
cordata, *Welw. ex Baker, l. c.* 27.—Afr. trop.
cordifolia, Linn. Mant. 256 = Hypocalyptus obcordatus.
crassifolia, Buch.-Ham. in Wall. Cat. n. 5416 = ferruginea.
crassipes, *Hook. Ic. Pl.* t. 880.—Austral.
crenata, Wight & Arn. Prod. 181 = seq.
crinita, R. Grah. in Wall. n. 5408 = calycina.
crispata, *F. Muell. ex Benth. Fl. Austral.* ii. 179.— Austral.
ctenoides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 59.—Afr. trop.
cubensis, DC. Prod. ii. 121 = incana.
cuneifolia, *Rafin. New Fl. Am.* ii. 55.—Am. bor.

CROTALARIA :—
cuneifolia, Schrank, in Syll. Ratisb. ii. (1828) 78 = sericea.
cuneiformis, Lam. Encyc. ii. 195 = Rafinia cuneifolia.
Cunninghamii, *R. Br. in Sturt, Exped.* ii. *App.* 71.— Austral.
curtata, *Link, Enum. Hort. Berol.* ii. 229.—Hab. ?
cyanea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 39.—Afr. trop.
cylindrica, A. Rich. Tent. Fl. Abyss. i. 154 = emarginata.
cylindrocarpa, *DC. Prod.* ii. 133.—Afr. trop.
cylindrostachya, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 15.—Afr. trop.
cytisifolia, *Hoffm. ex Heynh. Nom.* ii. 171, nomen.— Peruv.
cytisoides, *Hilsenb. & Bot. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 591.—Madag.
cytisoides, Roxb. Hort. Beng. 54 ; Fl. Ind. iii. 276 = Priotropis cytisoides.
cytisoides, Wight, in Wall. Cat. n. 5494 D = clavata.
decumbens, *Pers. Syn.* ii. 285.—Afr. austr.
decumbens, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 33.—Afr. trop.
deflexa, *Benth. in Miq. Pl. Jungh.* 206.—Java.
densicephala, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 21.—Afr. trop.
depauperata, *Mart. ex Benth. in Mart. Fl. Bras.* xv. i. 30.—Bras.
dichotoma, Boj. ex Sweet, Hort. Brit. ed. II. 128 = Bojeri.
dichotoma, R. Grah. in Edinb. N. Phil. Journ. (Oct.-Dec. 1826) 186 = pumila.
dichotoma, Heyne, ex Roth, Nov. Pl. Sp. 340 = hirsuta
diffusa, Link, Enum. Hort. Berol. ii. 228 = Linkiana.
diffusa, E. Mey. in Linnaea, vii. (1832) 151 = humilis.
diffusa, Roxb. ex Wight & Arn. Prod. 190 = linifolia.
diffusa, Vell. Fl. Flum. 307 ; vii. t. 110 = incana.
digitata, *Hook. Bot. Misc.* ii. (1831) 854.—Ind. or.
digitata, Wight, in Wall. Cat. n. 5430 = Grahamiana.
Dilloniana, *Baker, in Oliver, Fl. Trop. Afr.* ii. 41.— Afr. trop.
diosmaefolia, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 591.—Madag.
dimidiiflora, *Benth. in Mitch. Trop. Austral.* 886.— Austral.
distans, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 582.—Afr. austr.
distichа, Zoll. & Mor. in Nat. en Genesk. Arch. Nederl. Indie, iii. (1846) 75 = sciicularis.
divaricata, *Benth. in Ann. Nat. Hist.* iii. (1839) 429.— Bras.
divaricata, R. Grah. in Wall. Cat. n. 5436 = medicaginea.
divergens, Benth. in Ann. Nat. Hist. iii. (1839) 429 = breviflora.
diversifolia, E. Mey. Comm. Pl. Afr. Austr. i. 27 = Lotononis laxa.
Domboyana, DC. Prod. ii. 132 = anagyroides.
Donkia, *Baker, in Oliver, Fl. Trop. Afr.* ii. 28.—Afr. trop.
Doryenium, A. Rich. ex Baker, in Oliver, Fl. Trop. Afr. ii. 16 = microcarpa.
dubia, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 598.—Ins. Socotr.
dubia, *R. Grah. in Wall. Cat.* n. 5404.—Ind. or.
dumetorum, *Baker, in Oliver, Fl. Trop. Afr.* ii. 22.— Afr. trop.
dumosa, *Franch. in Revoil, Comali*, 25.—Afr. trop.
ebenoides, *Wulp. Rep.* i. 500.—Afr. trop.
Echlonisana, Presl, Bot. Bemerk. 46 = humilis.
Ecklonis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 42.— Afr. austr.
effusa, *E. Mey. Comm. Pl. Afr. Austr.* 25, ex parte.— Afr. austr.
effusa, E. Mey. l. c. ex parte = humilis.
elata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 29. —Afr. trop.
elegans, Bedd. in Madr. Journ. xix. (1858) 178 = pedunculata.
elegans, Hort. ex DC. Prod. ii. 124 = purpurea.
elliptica, *Mart. & Gal. in Bull. Acad. Brux.* x. 11. (1843) 84.—Mexic.
elliptica, Roxb. Hort. Beng. 54 ; Fl. Ind. iii. 279 = uncinella.
elongata, *Thunb. Prod. Pl. Cap.* 135.—Afr. austr.
emarginata, *Soy. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 586.—Afr. trop.

CROTALARIA :—
emarginella, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 109.—Afr. trop.
eminensis, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 581.—Madag.
equicincta, Dals. in Hook. Kew Journ. iii. (1851) 210 = albida.
eremaea, F. Muell. Rep. Greg. Pl. 5 = dissitiflora.
eriantha, Steb. & Zucc. in Abh. Acad. Muench. iv. ii. (1843) 121 = sessiliflora.
eriocarpa, *Benth. Bot. Voy. Sulph.* 80.— Mexic.
eriocaula, *S. Schau. in Linnaea*, xx. (1847) 788.— Mexic.
eriosemoides, *Ficalho et Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 17.—Afr. trop.
errodes, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 19.—Afr. trop.
erythrophleba, *Welw. l. c.* 95.—Afr. trop.
Espeddii, H. B. & K. Nov. Gen. et Sp. vi. 390 = stipularia.
evolvaloides, *Wight, in Wall. Cat.* n. 5410.—Ind. or.
exstipulata, *Mart. ex Schrank, in Syll. Ratisb.* ii. (1832) 78.—Bras.
falcata, Schum. & Thonn. Beskr. Guin. Pl. 325 = gorensis.
falcata, *Vahl, ex DC. Prod.* ii. 133.—Afr. trop.
fascta, *R. Br. in Salt, Abyss. App.* p. lxv, nomen.— Abyss.
fastigiata, E. Mey. in Linnaea, vii. (1832) 152 = Lotononis angustifolia.
fenestrata, Sims, Bot. Mag. t. 1028 = juncea.
ferruginea, *R. Grah. in Wall. Cat.* n. 5896.—As. trop.
ferruginea, Scheele, in Linnaea, xxi. (1848) 571 = Pohliana.
fertilis, *Delile, in Ferr. & Galin. Voy. Abyss.* iii. 122. —Abyss.
filicaulis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 24.—Afr. trop.
filiformis, *Wall. Cat.* n. 5389.—Burma.
filipes, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 475.—Ind. or.
flavicoma, *Benth. in Ann. Nat. Hist.* iii. (1839) 439. —Bras.
flexuosa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 29.—Afr. trop.
flexuosa, Moench, Meth. Suppl. 55 = verrucosa.
floribunda, Ajt. Hort. Kew. ed. II. iii. 10 = Wiborgia sericea.
floribunda, Lodd. Bot. Cab. t. 509 = Wiborgia floribunda.
florida, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 30.—Afr. trop.
foliosa, *Benth. in Ann. Nat. Hist.* iii. (1839) 439.— Bras.
foliosa, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 217 = medicaginea.
Forbesii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 13.—Afr. trop.
formosa, *R. Grah. in Wall. Cat.* n. 5393.—Ind. or.
fruticosa, Herb. Madr. ex Wall. Cat. sub n. 5374 = pulcherrima.
fruticosa, Macfad. Fl. Jamaic. i. 240 = pterocaula.
fruticosa, *Will. Gard. Dict.* ed. VIII. n. 4.—Jamaic.
fulgida, *Baker, in Oliver, Fl. Trop. Afr.* ii. 32.—Afr. trop.
fulva, *Reab. Hort. Beng.* 54; *Fl. Ind.* iii. 266.—As. trop.
furfuracea, *Botss. Diagn.* Ser. I. ii. 6.—Abyss.
Galeotti, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 582.—Mexic.
genistella, Sl. B. & K. Nov. Gen. et Sp. vi. 398 = pterocaula.
genistifolia, Schum. & Thonn. Beskr. Guin. Pl. 325 = glauca.
genistoides, *Lam. Encyc.* ii. 196.—Afr. austr.
genistoides, Willd. Sp. Pl. iii. 987 = Willdenowiana.
glabra, Stend. Nom. ed. II. i. 443 = uncinella.
glabra, Stend. i. c. = glauca.
glabrescens, *Anderss. in Vet. Akad. Handl. Stockh.* 1858 (1865) 246.—Isa. Gelap.
glabrescens, Benth. in Hook. Lond. Journ. Bot. iii. (1843) 563 = obtecta.
glauca, *Willd. Sp. Pl.* iii. 974.—Afr. trop.
glaucifolia, *Baker, in Oliver, Fl. Trop. Afr.* ii. 36.— Afr. trop.
globiflora, *E. Mey. Comm. Pl. Afr. Austr.* i. 24.—Afr. austr.
CROTALARIA :—
globosa, *Wight et Arn. Prod.* 190.—Ind. or.
globulosa, Wight, in Wall. Cat. n. 5413 = globosa.
glycinoa, Lam. Encyc. ii. 200 = Fagelia bituminosa.
goodiaeformis, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 230.—Afr. trop.
goreinsis, *Guill. & Perr. Fl. Senegamb. Tent.* i. 165.— Afr. trop.
gracilis, Pers. ex DC. Prod. ii. 133 = senegalensis.
gracilis, *Wulf. Rep.* i. 591.—Afr. trop.
gracillima, Klotzsch, in Peters, Reise Mossamb. Bot. 55 = hyssopifolia.
Grahamii, Sweet, Hort. Brit. ed. II. 128 = dichotoma.
Grahamiana, *Wight & Arn. Prod.* 194.—Ind. or.
grandiflora, Benth. in Ann. Nat. Hist. iii. (1839) 439 = cuspidata.
grandiflora, Reinw. ex Miq. Fl. Ind. Bat. i. i. 333 = tetragona.
grandis, Hort. Calc. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 80 = fulva.
Grantiana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 43.— Natal.
Grantii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 32.—Afr. trop.
guatemalensis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 3.—Guatem.
guianensis, Aubl. Pl. Gui. ii. 761 = Clitoria guianensis.
Guilleminiana, D. Dietr. Syn. Pl. iv. 932 = atrorubens.
guineensis, D. Dietr. l. c. 933 = gorensis.
hastata, Stend. Nom. ed. I. 339 = verrucosa.
havanensis, *Guss. in Linnaea*, iv. (1829) *Litt.* 86.— Cuba.
Heldeana, A. DC. in Mém. Soc. Phys. Genev. ix. (1841) 97 = Pohliana.
heptaphylla, *Lour. Fl. Cochinch.* 433.—Cochinch.
herbacea, Schweig. ex Schrank, in Syll. Ratisb. ii. (1832) 77 = incana.
hermannioides, Wight & Arn. Prod. 192 = medicaginea.
heterophylla, Linn. f. Suppl. 323 = quinquefolia.
hexaptera, Schrank, in Syll. Ratisb. ii. (1832) ii. 79 = pterocaula.
Heyneana, *R. Grah. in Wall. Cat.* n. 5414.—Ind. or.
Hildebrandtii, *Benth. in Mart. Fl. Bras.* xv. i. 25.—Bras.
Hildebrandtii, Baill. in Bull. Soc. Linn. Par. i. (1885) 445 = orthoclada, *Baker*.
Hildebrandtii, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 230.—Afr. trop.
hirsuta, Roxb. Hort. Beng. 54; Fl. Ind. iii. 270 = mysorensis.
hirsuta, Wall. Cat. n. 5413 A, C = evolvuloides.
hirsuta, *Willd. Sp. Pl.* iii. 978.—Ind. or.
hirta, Lag. Gen. et Sp. Nov. 32 = incana.
hirta, Roth, Nov. Pl. Sp. 339 = bifora.
hirta, *Willd. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 217.—Ind. or.
holopteryx, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 13.—Afr. trop.
holosericea, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 26.—Bras.
Hookeri, Arn. in Ann. Sc. Nat. Ser. II. iii. (1835) 248 = Saltana.
Hookeriana, A. DC. in Mem. Soc. Phys. Genev. ix. (1841) 97 = ovalis.
Hostmanni, Steud. in Flora, xxvi. (1843) 757 = retusa.
humifusa, *R. Grah. in Wall. Cat.* n. 5421.—Ind. or.
humilis, *Guill. & Zeyh. Enum.* 174.—Afr. austr.
hyssopifolia, *Klotzsch, in Peters, Reise Mossamb. Bot.* 55.—Afr. trop.
imbricata, Linn. Fl. Afr. Rar. 16 = Amphithalea densa.
imbricata, Willd. ex Steud. Nom. ed. II. i. 443 = Aspalathus lotoides.
improvisa, Nees, Del. Sem. Hort. Vrat. (1841) 3 = senegalensis.
incana, *Linn. Sp. Pl.* 716.—Am. trop.
incanescens, Linn. f. Suppl. 323 = capensis.
indigofera, Somerat. ex Lam. Encyc. ii. 200 = purpurascens.
inflata, Buch.-Ham. ex Steud. Nom. ed. II. i. 443 = sessiliflora.
intermedia, *Kotschy, in Sitzb. Acad. Wien, Math.-Nat.* I. Abth. i. (1866) 362.—Afr. trop.
involucrata, E. Mey. in Linnaea, vii. (1832) 152 = Lotononis involucrata.
involucrata, Soland. ex Baker, in Oliver, Fl. Trop. Afr. ii. 22 = cuneiolata.
CROTALARIA :—
iodina, Benth. in Lond. Journ. Bot. ii. (1843) 572 = atrorubens.
ivantulensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 26.—Afr. trop.
jamaica, *Spengl. in Hoov. & De Vriese, Tijdschr.* vii. (1840) 308.—Java.
jonces, *Linn. Sp. Pl.* 714.—As. trop.; Austral.
juncea, Willd. Sp. Pl. iii. 976 = sericea.
Kirkii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 26.—Afr. trop.
Kurzii, *Baker, ex Kurz, in Journ. As. Soc. Beng.* xliii. (1873) ii. 229.—Burma.
laburnifolia, *Linn. Sp. Pl.* 715.—As. trop.
laburnoides, Klotzsch, in Peters, Reise Mossamb. Bot. 57 = Saltiana.
lachnocarpa, *Hochst. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 38.—Abyss.
lachnophora, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 151 = lachnocarpa.
lacta, *Mart. ex Benth. in Mart. Fl. Bras.* xv. i. 30.— Bras.
laevigata, *Lam. Encyc.* ii. 198.—Madag.
laevigata, Pursh, Fl. Am. Sept. ii. 469 = Purshii.
lanata, *Brdd. in Madr. Journ. Sc.* Ser. II. xix. (1858) 178.—Ind. or.
lanata, Thunb. Prod. Pl. Cap. 124 = Priestleya villosa.
lanceolata, *E. Mey. Comm. Pl. Afr. Austr.* i. 24.—Afr. austr.
lanceolata, Roxb. Hort. Beng. 54 = Brownei.
lathyroides, *Guill. & Perr. Fl. Senegamb. Tent.* i. 168.— Afr. trop.
latifolia, *Linn. Sp. Pl.* 715.—Jamaic.
latifolia, Roxb. ex Wight & Arn. Prod. 180 = Saltiana.
laxa, *Franch. in Nouvl. Conati*, 24.—Afr. trop.
leiocarpa, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 8.—China.
leiobba, Benth. Ind. Sem. Hort. Gotting. (1837); ex Linnaea, xii. (1838) Litt. 80 = ferruginea.
Lenticula, E. Mey. Comm. Pl. Afr. Austr. i. 26 = Lotononis Lenticula.
Leprieurii, *Guill. & Perr. Fl. Senegamb. Tent.* i. 162.— Afr. trop.
leptoclona, *S. Schau. in Linnaea*, xx. (1847) 787.— Mexic.
leptophylla, Benth. in Ann. Nat. Hist. iii. (1839) 439 = mayparensis.
leptostachya, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 563.—Ind. or.
Leschenaultii, *DC. Prod.* ii. 125.—Ind. or.
Leschenaultii, Macfad. Fl. Jamaic. i. 239 = sericea.
linearis, Herb. Madr. ex Wall. Cat. n. 5370 = calycina.
linearis, *Rafin. New Fl. Am.* ii. 54.—Am. bor.
linearis, Lam. Encyc. ii. 200 = Eriocesna violaceum.
linearis, Thunb. Prod. Pl. Cap. 125 = decumbens.
linifolia, *Linn. f. Suppl.* 322.—As. trop.; Austral.
linifolia, Willd. Sp. Pl. iii. 978 = albida.
Linkiana, *Steud. Nom.* ed. II. i. 443.—Hab.?
littoralis, DC. Prod. ii. 134 = pumila.
littoralis, *H. B. & K. Nov. Gen. et Sp.* vi. 401.— Cuba.
longifolia, Lam. Encyc. ii. 201 = Clitoria guianensis.
longipes, *Rafin. New Fl. Am.* ii. 54.—Am. bor.
longipes, *Wight & Arn. Prod.* 183.—Ind. or.
longirostrata, *Hook. & Arn. Bot. Beech. Voy.* 295.— Mexic.
latifolia, Ait. Hort. Kew. ed. I. iii. 19 = latifolia.
latifolia, Poepp. ex Steud. Nom. ed. II. i. 443 = vitellina.
lotoides, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 575.—Afr. austr.
lotoides, Pers. Syn. i. 284 = latifolia.
lotoides, A. Rich. Tent. Fl. Abyss. i. 157 = microcarpa.
Lotocoela, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 29.—Afr. trop.
lunaria, Burm. f. Fl. Cap. Prod. 21 = Aspalathus lotoides.
lunaria, Linn. Sp. Pl. 716 = Argyrolobium lanceolatum.
lunata, Bedd. in Madr. Journ. Sc. Ser. III. i. (1866) 42 = laxa.
lunulata, *Heyne, in Wall. Cat.* n. 5376.—Ind. or.
lunulata, *Rafin. New Fl. Am.* ii. 55.—Am. bor.
lupulifera, R. Grah. in Wall. Cat. n. 5407 = Leschenaultii.
lupulifolia, Steud. Nom. ed. II. i. 443 = prona.

CROTALARIA:—
lupinoides, *Hochst. ex Benth. in Lond. Journ. Bot.* ii. (1843) 583.—*Afr. trop.*
lupulina, H. B. & K. Nov. Gen. et Sp. vi. 402. t. 590; DC. Prod. ii. 133 = pumila, *Ortog.*
lutescens, *Dala. in Hook. Kew Journ.* ii. (1850) 84.—Ind. or.
luxurians, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 578 = medicaginea.
maclanta, Delile in Caill. Voy. Meroe, ii. (reimp. 85) t. 64 = senegalensis.
macrocalyx, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 579.—Afr. trop.
macrocarpa, *E. Mey. Comm. Pl. Afr. Austr.* i. 24.—Afr. austr.
macrophylla, *Kurz, in Journ. As. Soc. Beng.* xiv. (1876) ii. 266.—Burma.
macrophylla, Welwn. in Syll. Ratisb. ii. (1828) 26 = sericea.
macropoda, Willd. Sp. Pl. iii. 982 = Flemingia congesta.
macropoda, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 157 = ortenesis.
macrostachya, Sond. in Linnæa, xxiii. (1850) 27 = globiflora.
macrostipula, Steud. ex A. Rich. Tent. Fl. Abyss. i. 158 = goreënsis.
macrostyla, D. Don, Prod. Fl. Nep. 242 = Lespedeza macrostyla.
madurensis, *Wight, in Wall. Cat.* n. 5876.—Ind. or.
magnifolia, *Tenore, Ind. Sem. Hort. Nap.* (1829) 16. —Hab.?
malabarica, DC. Prod. ii. 127 = nana.
Marize-Antoniæ, Aschers. ex Baker, in Oliver, Fl. Trop. Afr. ii. 16 = microcarpa.
maritima, *Chapm. Fl. S. U. St Suppl.* 614.—Am. bor
Martiana, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 582 = folicea.
Matthewsiana, *Benth. l. c.* 479.—Fervv.
maxillaris, *Klotzsch, in Peters, Reise Mossamb. Bot.* 58.—Afr. trop.
mayurensis, *H. B. & K. Nov. Gen. et Sp.* vi. 403.—Venezuela.
medicaginea, DC. Prod. ii. 133 = C. Trifollastrum.
medicaginea, *Lam. Encyc.* ii. 201.—As. trop.; Austral.
melanocarpa, Wall. Cat. n. 5883 = linifolia.
mellitoides, Steud. ex A. Rich. Tent. Fl. Abyss. i. 155 = pycnostachya.
Meyeriana, *Steud. Nom.* ed. II. i. 444.—Afr. austr.
micans, *Link, Enum. Hort. Berol.* ii. 228.—Hab.?
micrantha, *Link, Enum. Hort. Berol.* ii. 229.—Zeylan.
micrantha, E. Mey. Comm. Pl. Afr. Austr. i. 27 = Lotonoia microcalyx.
microcarpa, *Hochst. ex Benth. in Lond. Journ. Bot.* ii. (1843) 573.—Nubia.
microphylla, *Vahl, Symb. Bot.* i. 52.—Arab.
Mitchelii, F. Muell. Fragm. iii. 50 = Novae-Hollandiæ.
Mitchellii, *Benth. in Mitch. Journ Trop. Austral.* 122.—Austral.
mollicula, *H. B. & K. Nov. Gen. et Sp.* vi. 408.—Mexico.
mollis, E. Mey. Comm. Pl. Afr. Austr. i. 23 = Meyeriana.
mollis, *Weimm. in Syll. Ratisb.* ii. (1828) 25.—China.
montana, Heyne, ex Roth, Nov. Pl. Sp. 335 = linifolia.
montana, *A. Rich. Tent. Fl. Abyss.* i. 152.—Abyss.
montana, Roxb. Hort. Beng. 54; Fl. Ind. iii. 265 = albida.
mozambicensis, Klotzsch, in Peters, Reise Mossamb. Bot. 60. t. 10 = lanceolata.
mucronata, *Desv. Journ. Bot.* iii. (1814) 76.—Ind. occ.
muldiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 578.—Zeylan.
myuroides, *Roth, Nov. Pl. Sp.* 338.—Ind. or.
nana, *Burm. f. Fl. Ind.* 156. t. 48.—Ind. or.
nepaulensis, DC. Prod. ii. 125 = *nepalensis*.
natalitia, *Meisen. in Hook. Lond. Journ. Bot.* ii. (1843) 67.—Afr. austr.
neglecta, Wight & Arn. Prod. 192 = medicaginea.
nepalensis, Link, Enum. Hort. Berol. ii. 228 = sessiliflora.
nerilifolia, *Wall. Cat.* n. 5862.—Burma.
nervosa, R. Grah. in Wall. Cat. n. 5436 = Pycnospora lachyarachides.
nigricans, *Baker, in Oliver, Fl. Trop. Afr.* ii. 21.—Afr. trop.

CROTALARIA:—
nipaulensis, G. Don, Gen. Syst. ii. 137 = *nepalensis*.
nitens, *H. B. & K. Nov. Gen. et Sp.* vi. 399.—N. Granat.
nitidula, Mart. ex Schrank, in Syll. Ratisb. ii. (1828) 76 = nitens.
Novæ-Hollandiæ, *Wight & Arn. Prod.* 192.—Ind. or. Novae-Hollandiae, *DC. Prod.* ii. 127.—Austral.
nubica, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 581.—Afr. austr.
nummularia, Willd. Sp. Pl. iii. 979 = biflora.
nutans, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 24.—Afr. trop.
obcordata, *Berg. Desc. Pl. Cap.* 195.—Afr. austr.
obtiqua, Buch.-Ham. in Wall. Cat. n. 5888 = ferruginea.
oblanceifolia, Hook. Ic. Pl. sub t. 880 = Novae-Hollandiae.
obovata, *G. Don, Gen. Syst.* ii. 138.—Afr. trop.
obscura, *DC. Prod.* ii. 134.—Afr. austr.
obscura, *R. Grah. in Wall. Cat.* n. 5873.—Ind. or.
occulta, *R. Grah. l. c.* n. 5860.—Reg. Himal.
ochroleuca, *G. Don, Gen. Syst.* ii. 138.—Afr. trop.
Oldhami, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 42 = sessiliflora.
oligostachya, *Baker, in Oliver, Fl. Trop. Afr.* ii. 41.—Afr. trop.
Onobrychis, *A. Rich. Tent. Fl. Abyss.* i. 154.—Abyss
ononoides, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 572.—Afr. trop.
opposita, Linn. f. Suppl. 323 = Rafnia opposita.
ornicularia, E. Mey. in Linnæa, vii. (1832) 151 = Podalyria orbicularis.
orixensis, *Willd. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 217.—Ind. or.; Afr. trop.
orthocladu, *Baker, in Journ. Linn. Soc.* xx. (1883) 124.—Madag.
orthoclada, *Welw ex Baker, in Oliver, Fl. Trop. Afr.* ii. 29.—Afr. trop.
otopteru, *Benth. in Ann. Nat. Hist.* iii. (1839) 429.—Bras.
ovalifolia, Wall. Cat. n. 5411 = rubiginosa.
ovalis, *Pursh, Fl. Am. Sept.* ii. 469.—Georgia.
oxalidifolia, Hort. ex Steud. Nom. ed. II. i. 444 = Lalldi, Steetz oxalidifolia.
oxyptera, E. Mey. Comm. Pl. Afr. Austr. i. 28 = Lotononis oxyptera.
pallida, Blanco, Fl. Filip. ed. I. 570 = sessiliflora.
pallida, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 20.—Afr. trop.
papillosa, Klotzsch, in Peters, Reise Mossamb. Bot. 58 = Saltiana.
paniculata, Pers. Syn. ii. 284 = senegalensis.
paniculata, *Willd. Sp. Pl.* iii. 982.—As. trop.
paramaribensis, Miq. in Ann. & Mag Nat. Hist. xi. (1843) 14 = verrucosa.
parva, R. Grah. in Wall. Cat. n. 5402 = albida.
parviflora, Pursh, Fl. Am. Sept. ii. 469 = Purshii.
parviflora, Roth, Catalect. i. 83 = sagittalis.
parvifolia, Thunb. Prod. Pl. Cap. 124 = Coelidium Thunbergii.
parvula, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 25.—Afr. trop.
patula, R. Grah. in Wall. Cat. n. 5371 = nana.
paulina, *Schrank, Fl. Ratisb. Monac.* t. 88.—Bras.
pedunculariu, Dala. & Gibs. Bomb. Fl. 55 = lutescens.
peduncularis, *R. Grah. in Wall. Cat.* n. 5369.—Ind. or.
pedunculosa, Desv. Journ. Bot. iii. (1814) 76 = laburnifolia.
peguana, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 77.—Burma.
pellita, Bert. ex DC. Prod. ii. 128 = ramosissima.
pendula, Bert. l. c. 130 = laburnifolia.
perfoliata, Linn. Sp. Pl. 714 = Baptisia perfoliata.
perforata, Linn. Fl. Afr. Rar. 15; Amoen. Acad. vi. 92 = Borbonia cillata.
perplexa, E. Mey. Comm. Pl. Afr. Austr. i. 26 = Lotononis perplexa.
Perrottetii, *DC. Prod.* ii. 128.—Afr. trop.
Perrvillei, *Baill. in Bull. Soc. Linn. Par.* i. (1886) 444.—Madag.
petiolaris, *Fresnel. in Rœpll, Comali*, 26.—Afr. trop.
petiolata, *Vog. ex Walp. in Linnæa*, xiii. (1839) 474. —Afr. austr.
Petitiana, *A. Rich. Tent. Fl. Abyss.* i. 159.—Abyss.
physcarpa, Fenzl, ex Baker, in Oliver, Fl. Trop. Afr. ii. 17 = podocarpa.

CROTALARIA:—
pilifera, Klotzsch, in Peters, Reise Mossamb. Bot. 54 = podocarpa.
pilosa, *Mill. Gard. Dict.* ed. VIII. n. 2.—Mexic.
pilosa, *Raffn. New Fl. Am.* ii. 54.—Am. bor.
pilosa, Roxb. ex Mart. in Denkschr. Acad. Muench. vi. (1820) 186 = hirta.
pilosa, Thunb. Prod. Pl. Cap. 125 = obscura.
pilosissima, Miq. Fl. Ind. Bat. i. 1. 327 = ferrugines.
pisicarpa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 16.—Afr. trop.
pisiformis, Guill. & Perr. Fl. Seneg. Tent. i. 162 = Saltiana.
platycalyx, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 160. —Abyss.
platycarpa, Link, Enum. Hort. Berol. ii. 227 = sagittalis.
platyserpala, *Harv. Thes. Cap.* i. 19. t. 29.—Afr. austr.
Plowdeni, *Baker, in Oliver, Fl. Trop. Afr.* ii. 41.—Afr. trop.
podocarpa, *DC. Prod.* ii. 133.—Afr. trop.
Poepplgii, *Presl, Bot. Bemerk.* 128.—Cuba.
Pohliana, *Benth. in Ann. Nat. Hist.* iii. (1839) 428.—Bras.
polycarpa, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 580 = sphaerocarpa.
polycladu, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 17.—Afr. trop.
polygaloides, *Welw. ex Baker, l. c.* 15.—Afr. trop.
polyperma, *Kotschy, Reliq. Kotsch.* (1866) 18. t. 14.—Afr. trop.
porrecta, Wall. Cat. n. 5362 = juncea.
priestleyoides, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 74.—Ind. or.
Pringlei, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 200.—Am. bor. occ.
procumbens, [Moç. & Sesse,] ex DC. Prod. ii. 129 = ovalis.
procumbens, Roxb. Hort. Beng. [98]; Fl. Ind. iii. 278 = medicaginea.
procumbens, C. Sm. in Tuckey, Congo, 352 = Tuckeyana.
procumbens, Wall. Cat. n. 5437 = hemiarioides.
prolifera, E. Mey. in Linnæa, vii. (1832) 152 = Lotononis porrecta.
prolongata, *Baker, in Oliver, Fl. Trop. Afr.* ii. 25.—Afr. trop.
propinqua, *R. Br. in Salt, Abyss. App.* p. lxv. nomen. —Abyss.
prostrata, *Rottl. in Willd. Enum. Hort. Berol.* 747.—As. trop.
prostrata, Wight, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 67 = humifusa.
pubescens, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 35.—Afr. trop.
Pseudoeriosema, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 198.—Afr. trop.
pseudolotoides, D. Don, Prod. Fl. Nep. 242 = Psiotropis cytisoides.
pseudolotoides, Lam. Encyc. ii. 201 = villosa, Rhynchosia pseudolotoides.
pterocaula, *Desv. Journ. Bot.* iii. (1814) 76.—Am. austr.
pteropoda, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 508.—Isl. Socotr.
pubera, *Vahl, Eclog. Am.* ii. 55.—Ind. occ.
pubernia, Hook. f. in Trans. Linn. Soc. xx. (1847) 225 = pumila.
pubescens, Hort. ex Steud. Nom. ed. II. i. 444 = pubera.
pubescens, Moench, Meth. 161 = incana.
pulchella, Andr. Bot. Rep. t. 417 = Lebeckia cytisoides.
pulcherrima, *Roxb. Hort. Beng.* 54; *Fl. Ind.* iii. 267.—Ind. or.
pulchra, Andr. Bot. Rep. t. 601 = pulcherrima.
pulchra, DC. Prod. ii. 126 = fulva.
pumila, Blanco, Fl. Filip. ed. II. 397 = sessiliflora.
pumila, Hochst. & Steud. ex Baker, in Oliver, Fl. Trop. Afr. ii. 17 = microphylla.
pumila, *Ortog. Hort. Matr.* 28.—Cuba.
pumila, *Raffn. New Fl. Am.* ii. 56.—Am. bor.
pumila, Schrank, in Denkschr. Acad. Muench. vi. (1820) 188 = pusilla.
punctata, R. Grah. in Wall. Cat. n. 5401 A, C = albida.
purpurascens, *Lam. Encyc.* ii. 200.—Madag.

CROTALARIA :—
purpurea, *Vent. Yard. Malm.* t. 66.—Afr. austr.
Purshii, *DC. Prod.* ii. 124.—Am. bor.
pusilla, *Heyne, ex Roth, Nov. Pl. Sp.* 335.—Ind. or.
pusilla, Retz. ex Wight & Arn. Prod. 189 = evolvuloides.
pycnostachya, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 554.—Abyss.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* i. 158.—Abyss.
quinata, E. Mey. Comm. Pl. Afr. Austr. i. 27 = Lotononis acutiflora.
quinquefolia, Roxb. ex Roth, Nov. Pl. Sp. 342 = C. Trifoliastrum.
quinquefolia, *Linn. Sp. Pl.* 716.—As. trop.
racemosa, Vell. Fl. Flum. 306; vii. t. 106 = paulina.
ramiflora, Perr. ex DC. Prod. ii. 168 = Tephrosia obcordata.
ramosissima, *Baker, in Oliver, Fl. Trop. Afr.* ii. 36.—Afr. trop.
ramosissima, F. Muell. Fragm. iii. 55 = crispata.
ramosissima, *Roxb. Hort. Beng.* 54; *Fl. Ind.* iii. 288.—Ind. or.
recta, *Stud. ex A. Rich. Tent. Fl. Abyss.* i. 152.—Afr. trop.
reflexa, Thunb. Prod. Pl. Cap. 195 = Priestleya sericea.
Regnellii, Benth. in Linnæa, xxii. (1849) 511 = Pohliana.
remotiflora, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 31 = senegalensis.
remiformis, Lam. Encyc. ii. 194 = Valcos perfoliata.
retusa, *Linn. Sp. Pl.* 715.—Cosmop. trop.
retusifolia, Stokes, Bot. Mat. Med. iii. 316 = retusa.
rhizophylla, R. Grah. in Wall. Cat. n. 5481 = occulta.
rigida, *Heyne, ex Roth, Nov. Pl. Sp.* 343.—Ind. or.
rostrata, Wight & Arn. Prod. 191 = Notonii.
Rothiana, DC. Prod. ii. 127 = bifora.
Rothii, Spreng. Syst. iii. 237 = præc.
rotundifolia, *Petr. Encyc. Suppl.* ii. 402.—Am. bor.
Roxburghiana, DC. Prod. ii. 129 = calycina.
Roxburghii, Spreng. Syst. iii. 239 = præc.
rubiginosa, Buch.-Ham. ex Wall. Cat. n. 5356 = alata.
rubiginosa, Herb. Madr. ex Wall. l. c. n. 5361 = mysorensis.
rubiginosa, Roxb. Hort. Beng. 98; Fl. Ind. iii. 260 = hirsuta.
rubiginosa, Walp. Rep. i. 890 = atrorubens.
rubiginosa, *Willd. Sp. Pl.* iii. 978.—Ind. or.
ruficoma, Franch. Pl. Davidian. i. (1884) 79 = ferruginea.
rufipila, *Benth. in Mart. Fl. Bras.* xv. i. 23.—Bras.
sagittalis, Desv. ex Griseb. Fl. Brit. W. Ind. 178 = stipularia.
sagittalis, *Linn. Sp. Pl.* 714.—Am. bor.
sagittalis, Vell. Fl. Flum. 308; vii. t. 111 = stipularia.
sagittiformis, Wall. Cat. n. 5357 = alata.
Saharæ, *Coss. in Bull. Soc. Bot. Fr.* xi. (1864) 165.—Afr. bor.
salicifolia, *Heyne, in Wall. Cat.* n. 5359 A.—Ind. or.
Saltiana, *Andr. Bot. Rep.* t. 648 (1811).—Geront. trop.
Saltiera, Ait. ex Steud. Nom. ed. I. 239 = præc.
saxatilis, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 219.—Afr. trop.
saxatilis, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indië, iii. (1846) 76 = linifolia.
scabrella, Wight & Arn. Prod. 181 = rubiginosa.
scandens, *Lour. Cochinch.* ii. 435. —Cochinch.
scariosa, *Rafin. New Fl. Am.* ii. 56.—Am. bor.
Schiedeana, *Steud. Nom.* ed. II. i. 445.—Mexic.
Schimperi, A. Rich. Tent. Fl. Abyss. i. 151 = incana.
Schimmacheri, D. Dietr. Syn. Pl. iv. 933 = falcata.
Schweinfurthii, *Deflers, in Bull. Soc. Bot. Fr.* xxxii. (1885) 348.—Afr. trop.
scoparia, Wall. Cat. n. 5418 = albida.
semperflorens, *Vent. Yard. Cels*, t. 17.—As. trop.
senegalensis, *Bocte in DC. Prod.* ii. 128.—Afr. trop.
sericea, Burm. f. Fl. Ind. 156 = Burmanni.
sericea, *Retz. Obs.* v. 26.—Ind. or.
sericea, Willd. Sp. Pl. iii. 979 = juncea.
serpens, E. Mey. in Linnæa, vii. (1832) 158 = Euchlora serpens.
serpilifolia, Lamb. ex DC. Prod. ii. 124 = stipularia.
serpilifolia, Wight, ex Wall. Cat. sub n 5356 = Wightiana.
sessiliflora, *Linn. Sp. Pl.* ed. II. 1004.—China.
sessilifolia, Linn. Mant. ii. 440 = sessiliflora.
setifera, *DC. Prod.* ii. 131.—Mexic.
simplex, A. Rich. Tent. Fl. Abyss. i. 153 = recta.

CROTALARIA :—
sinensis, J. F. Gmel. Syst. 1095 = chinensis.
sobolifera, R. Grah. in Wall. Cat. n. 5490 A = linifolia.
sobolifera, R. Grah. l. c. n. 5490 B = nana.
sparsiflora, *E. Mey. Comm. Pl. Afr. Austr.* i. 26.—Afr. austr.
spartea, *R. Br. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 13.—Afr. trop.
spartioides, *DC. Prod.* ii. 128.—Afr. austr.
spartioides, E. Mey. Comm. Pl. Afr. Austr. i. 23 = virgultalis.
spartioides, Spreng. Syst. iii. 240 = Willdenowiana.
speciosa, *Heyne, ex Roth, Nov. Pl. Sp.* 336.—Ind. or.
spectabilis, Roth, l. c. 341 = sericea.
sphærocarpa, *Perr. ex DC. Prod.* ii. 133.—Afr. tro .
spinosa, *Hochst. in Flora*, xxiv. (1841) 1. *Intall.* 33; *et ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 576.—Abyss.
spinulosa, *Burm. f. Fl. Cap. Prod.* 21.—Afr. austr.
spiendens, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 8 = elliptica.
stenophylla, Eckl. & Zeyh. Enum. 174 = Eckloni.
stenophylla, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 7 = linifolia.
stenoptera, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 12.—Afr. trop.
Steudneri, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xvii. (1868) 651.—Afr. trop.
Stewartii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 32.—Afr. trop.
stipitata, R. Grah. in Wall. Cat. n. 5425 A = C. Trifoliastrum.
stipitata, R. Grah. l. c. n. 5425 B = laevigata.
stipularis, Roxb. Hort. Beng 54; Fl. Ind. iii. 264 = mysorensis.
stipularia, *Desv. Journ. Bot.* iii. (1814) 76.—Guian.; Ind. or.
stipulata, Vell. Fl. Flum. 307; vii. t. 109 = anagyroides.
Stocksii, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 67.—Ind. or.
striata, A. Br. in Flora, xxiv. (1841) 280 = astragalina.
striata, DC. Prod. ii. 131 = Saltiana.
striata, *Schrank, in Syll. Ratisb.* ii. (1828) 76.—Bras.
striata, Schum. & Thonn. Beskr. Guin. Pl. 336 = falcata.
stricta, Roth, Nov. Pl. Sp. 342 = C. Trifoliastrum.
stricta, Roxb. Hort. Beng. 54; Fl. Ind. iii. 265 = strigosa.
strigosa, *Boj. Hort. Maurit.* 88.—Hab. ?
strigulosa, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 584.—Socotr.
Sturtii, *R. Br. in Sturt, Exped. ii. App.* 70.—Austral.
styracifolia, *Desv. Tabl. ed.* II. 214, nomen; *Hornem. Hort. Hafn. Suppl.* 151.—Afr. austr.
subdecurrens, *Mart. ex Benth. in Mart. Fl. Bras.* xv. i. 20.—Bras.
subperfoliata, *Wight, in Wall. Cat.* n. 5377.—Ind. or.
tacta, *Ecklon & Zeyh. Enum. Pl. Sp.* 334.—Ind. or.
tenella, E. Mey. in Linnæa, vii. (1832) 152 = Lotononis tenella.
tenuiflora, *Steud. Nom.* ed. II. i. 445.—Afr. austr.
tomsifolia, Roxb. Hort. Beng. 54; Fl. Ind. iii. 263 = juncea.
townsfolia, Wight, ex Wall. Cat. sub n. 5369 = pedunculata.
tenuirama, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 34.—Afr. trop.
tenuis, *Baker, in Journ. Linn. Soc.* xx. (1883) 124.—Madag.
tenuis, Wall. Cat. n. 5403 = albida.
teplicans, *Hook. & Arn. Bot. Beech. Voy. Suppl.* 414.—Mexic.
tetragona, *Roxb. ex Andr. Bot. Rep.* t. 598; *Fl. Ind.* iii. 263.—As. trop.
tetragona, Wall. Cat. n. 5367 D = obtecta.
tetragonoloba, Roxb. ex Steud. Nom. ed. II. i. 445 = tetragona.
tetraptera, Dietr. ex DC. Prod. ii. 126 = pulcherrima.
thebaica, *DC. Prod.* ii. 128.—Afr. trop.
thebaica, Guill. & Perr. Fl. Seneg. Tent. i. 160 = arenaria.
Thomsoni, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 399.—Afr. trop.
Thunbergiana, Vog. ex Walp. in Linnæa, xiii. (1839) 475 = Lotononis acutiflora.
tigrensis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 36.—Afr. trop.

CROTALARIA :—
tinctoria, Bolv. ex Baill. in Bull. Soc. Linn. Par. i. (1885) 449 = striata.
tomentosa, Perr. ex DC. Prod. ii. 128 = Perrottetii.
tomentosa, Rottl. ex Spreng. Syst. iii. 236 = ramosissima.
tomentosa, *Thunb. Prod. Pl. Cap.* 192.—Afr. austr.
tomentosa, Wight, ex Steud. Nom. ed. II. i. 445 = digitata.
topicansa, Walp. Rep. i. 894, sphalm. = tepicana.
triantha, [Mog. & Sesse,] ex DC. Prod. ii. 135 = pumila.
triantha, Steud. ex Baker, in Oliver, Fl. Trop. Afr. ii. 16 = microcarpa.
trichophora, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 67.—Ind. or.
trichopoda, E. Mey. Comm. Pl. Afr. Austr. i. 154 = Lotononis trichopoda.
trichotoma, *Boj. in Ann. Sc. Nat.* Sér. II. iv. (1835) 265.—Madag.
trifera, Heyne, in Wall. Cat. n. 5387 = hirsuta.
trifera, Linn. Sp. Pl. 715 = Rafnia trifera.
Trifoliastrum, Wall. Cat. n. 5423 G = Notonii.
Trifoliastrum, *Willd. Sp. Pl.* iii. 968.—Ind. or.
trigonaliodes, Webb, in Hook. Niger Fl. 117, in obs. = microphylla.
triphylla, Vell. Fl. Flum. 307; vii. t. 108 = brachystachya.
triquetra, *Dals. in Hook. Kew Journ.* ii. (1850) 84.—Ind. or.
truncata, E. Mey. in Linnæa, vii. (1832) 151 = Lotononis umbellata.
tuberosa, Buch.-Ham. in D. Don, Prod. Fl. Nep. 241 = Eriosema chinense.
Tuckeyana, *Steud. Nom.* ed. II. i. 445.—Afr. austr.
turgida, *Loisel. Herb. Amat.* t. 288.—Hab. ?
Tweediana, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 483.—Reg. Argent.
ukambensis, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 219.—Afr. trop.
umbellata, Wight, in Wall. Cat. n. 5383 = nana.
uncinata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 33.—Afr. trop.
uncinella, *Lam. Encyc.* ii. 200.—China.
undulata, *Knowles & Westc. Fl. Cab.* ii. (1838) 155.—Mexic.
uniflora, *Baker, in Oliver, Fl. Trop. Afr.* ii. 13.—Afr. trop.
uniflora, Koen. ex Roxb. Hort. Beng. [98]; Fl. Ind. iii. 271 = Heylandia latebrosa.
unifoliolata, *Benth. in Ann. Nat. Hist.* iii. (1839) 430.—Bras.
Vachellii, Hook. & Arn. Bot. Beech. Voy. 180 = elliptica.
variegata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 19.—Afr. trop.
vascnicea, *R. Grah. in Wall. Cat.* n. 5427.—Ind. or.
velutina, *Benth. in Ann. Nat. Hist.* iii. (1839) 429.—Bras.
venusta, Wall. Cat. n. 5365 = semiliflora.
verrucosa, *Linn. Sp. Pl.* 715.—Cosmop. trop.
versicolor, *Baker, in Oliver, Fl. Trop. Afr.* ii. 26.—Afr. trop.
versicolor, E. Mey. in Linnæa, vii. (1832) 152 = Lotononis versicolor.
Vespertilio, *Benth. in Ann. Nat. Hist.* iii. (1839) 429.—Bras.
vestita, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 67.—Ind. or.
vestitiata, E. Mey. in Linnæa, vii. (1832) 158 = Lotononis prostrata.
villosa, Linn. Sp. Pl. 715 = Podalyria calyptrata.
villosa, *Thunb. Prod. Pl. Cap.* 192.—Afr. austr.
viminea, Wall. Cat. n. 5397 B = juncea.
viminea, Wall. l. c. n. 5397, pro parte = tecta.
virgata, Benth. in Ann. Nat. Hist. iii. (1839) 429 = paulina.
virgata, Herb. Holm. ex Harv. in Harv. & Sond. Fl. Cap. ii. 30 = Rafnia Thunbergii.
virgata, Mart. in Denkschr. Acad. Muench. vi. (1820) 157 = medicaginea.
virgata, Roxb. ex Wight & Arn. Prod. i. 192 = C. Trifoliastrum.
virgatula, Baker, in Oliver, Fl. Trop. Afr. ii. 30 = virgulata.
virgulata, *Klotzsch, in Peters. Reise Mossamb. Bot.* 86.—Afr. trop.
virgulta, *Spreng. Syst. iv. Cur. Post.* 276.—Afr. austr.

CROTALARIA :—
virgultalis, *Burch. ex DC. Prod.* ii. 126.—Afr. austr.
stacosa, Punb, ex Steud. Nom. ed. II. i. 444, in syn. = paniculata.
vitellina, *Ker-Gawl. in Bot. Reg.* t. 447.—Bras.
Vogelii, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 561.—Afr. trop.
Vogelii, D. Dietr. Syn. Pl. iv. 926 = lintfolia.
volubilis, *Thunb. Prod. Pl. Cap.* 135.—Afr. austr.
Walkeri, Arn. in Nov. Act. Nat. Cur. xviii. i. (1840) 825 = semperflorens.
Wallichiana, Wight & Arn. Prod. 187 = semperflorens.
Waleticshii, *Baker*, *in Oliver*, *Fl. Trop. Afr.* ii. 13.—Afr. trop.
Wightiana, R. Grah. in Wall. Cat. n. 5358 = rubiginosa.
Willdenowiana, *DC. Prod.* ii. 134.—Afr. austr.
xanthoclada, *Boj. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 588.—Madag.
zanzibarica, *Benth. l. c.* 584.—Afr. trop.
Zollingeriana, *Miq. Fl. Ind. Bat.* i. 1. 344.—Java.
Zuccariniana, D. Dietr. Syn. Pl. iv. 925 = striata.

CROTALOPSIS, Michx. ex DC. Prod. ii. 100, in syn. (1825) = **Baptisia**, Vent. (Legum.).

CROTON, Linn. Gen. ed. I. 986 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 293.
ANGELANDRA, Endl. Gen. Suppl. iv. 111. 91 (1850).
AMIERPTA, Rafin. Cat. 13 (1824).
ANISOPHYLLUM, Boiv. ex Baill. Adansonia, i. (1860-61) 153
ARGYRA, Noronha, ex Baill. Adansonia, i. (1860-61) 162.
ARGYRODENDRON, Klotzsch, in Peters, Reise Mossamb. Bot. 100 (1861).
AROTON, Neck. Elem. ii. 336 (1790).
ASYRARA, Klotzsch, in Wiegm. Archiv, vii. (1841) 194.
ASTROGYNE, Benth. Pl. Hartw. 14 (1839).
AUBERTIA, Chapel. ex Baill. Adansonia, i. (1860-61) 162, 166.
BARHAMIA, Klotzsch, in Seem. Bot. Voy. Herald, 104 (1855).
BRACHYSTACHYS, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 47.
BRUNSVIA, Neck. Elem. ii. 337 (1790).
CALYPTRIOPETALUM, Hassk. Retzia, i. 94 (1858).
CASCARILLA, Adans. Fam. ii. 350 (1763).
CINOGASUM, Neck. Elem. ii. 336 (1790).
CLEODORA, Klotzsch, in Wiegm. Archiv, vii. (1841) 196.
CODONOCALYX, Klotzsch, ex Baill. Etud. Gen. Euph. 369 (1858).
COMATOCROTON, Karst. in C. Koch, Wochenschr. ii. (1859) 6.
CROTONANTHUS, Klotzsch, ex Schlecht. in Linnæa, xxvi. (1853–55) 884.
CYCLOSTIGMA, Klotzsch, in Seem. Bot. Voy. Herald, 104 (1853).
DECARINIUM, Rafin. Neogenyt. 1 (1825).
DERPADENIUM, Rafin. l. c. 9 (1825).
ELUTHERIA, Linn. Hort. Cliff. 486 (1737).
ENGELMANNIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 258.
EUTROPIA, Klotzsch, l. c. (1841) 196
FURCARIA, Boiv. ex Baill. Etud. Gen. Euph. 356 (1858).
GISEKLERIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 254.
GYNAMBLOSIS, Torr. in Rep. Marcy Exped. 295 (1858).
HALICUS, Rumpf. ex Rafin. Sylva Tellur. 69 (1838).
HENDECANDRA, Eschsch. in Mem. Acad. Petersb. x. (1826) 287.
HEPTALLON, Rafin. Neogenyt. 1 (1825).
KLOTZSCHIPHYTUM, Baill. Etud. Gen. Euph. 362 (1858).
KUKRAS, Rafin. Sylva Tellur. 69 (1838).
LAMBOGYNE, Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. 416 (1843).
LEONTIA, Reichb. Consp. 198 (1828).
LUCCADENIA, Klotzsch, ex Baill. Adansonia, iv. (1864) 258.
LUNTIA, Neck. Elem. ii. 335 (1790).
MACROCROTON, Klotzsch, in Schomb. Fauna et Fl. Gui. 1186 (1848).

CROTON :—
MEDEA, Klotzsch, in Wiegm. Archiv, vii. (1841) 198.
MEIALISA, Rafin. Sylva Tellur. 68 (1838).
MONGULA, Chapel. ex Baill. Adansonia, i. (1860–61) 147.
MYRIOGONOPHUS, Didr. in Kjoeb. Vidensk. Meddel. (1857) 149.
OCALIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 195.
ODOTALON, Rafin. Sylva Tellur. 66 (1838).
PALAMOSTIGMA, Mart. ex Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 49.
PENTECA, Rafin. Sylva Tellur. 69 (1838).
PILINOPHYTUM, Klotzsch, in Wiegm. Archiv, vii. (1841) 255.
PODOSTACHYS, Klotzsch, l. c. (1841) 198.
SCHRADERA, Willd. in Goett. Journ. Wiss. i. (1797) 1.
SEMILTA, Rafin. Sylva Tellur. 68 (1838).
TIGLIUM, Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. 418 (1843).
TIMANDRA, Klotzsch, in Wiegm. Archiv, vii. (1841) 197.
TRIDESMIS, Lour. Fl. Cochinch. 576 (1790).
TRIPLANDRA, Rafin. Sylva Tellur. 69 (1838).
abaliensis, *Baill. Adansonia*, iv. (1863–64) 337.—Bras.
aberrans, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 232.—Bras.
abnormis, Mart. ex Baill. Adansonia, iv. (1863–64) 354 = timandroides.
abutiloides, *H. B. & K. Nov. Gen. et Sp.* ii. 86.—Peruv.
acerifolius, A. Cunn. ex Hook. in Mitch. Journ. Trop. Austral. 371 = Adriana glabrata.
acervinus, Reinw. ex Blume, Cat. Gew. Buitenz. 104 = Dalechampia bidentata.
acronychioides, *F. Muell. Fragm.* iv. 142.—Austral.
acuminatus, K. Br. in Salt, Abyss. App. p. lxv = macrostachys.
acuminatus, Lam. Encyc. ii. 307 = Mallotus japonicus.
acuminatus, A. Rich. ex Baill. Adansonia, i. (1860–61) 276 = Cephalocroton leucocephalus.
acutus, Thunb. Fl. Jap. 269 = C. Tiglium.
adenanthus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 115.—Bras.
adenocalyx, *Baill. Adansonia*, iv. (1863–64) 344.—Bras.
adenodontus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 267.—Am. austr.
adenopetalus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 90 = Sarothrocrotes.
adenophorus, *Baill. Adansonia*, i. (1860–61) 153.—Madag.
adenophyllus, *Spreng. Syst.* iii. 869.—Jamaica.
adipatus, *H. B. & K. Nov. Gen. et Sp.* ii. 68.—Bras.
adpressus, *Wright & Griseb. in Goett. Nachr.* (1865) 178.—Cuba.
adspersus, *Benth. Pl. Hartw.* 51.—Mexic.
affinis, Geisel. Crot. Monog. 25 = glandulosus.
agoensis, *Baill. Adansonia*, iv. (1863–64) 348.—Bras.
agrarius, *Baill. l. c.* 319.—Bras.
agrophilus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 175.—Bras.
alabamensis, *E. A. Smith, ex Chapman, Fl. S. U. St. Suppl.* 648.—Am. bor.
alagoensis, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 121.—Bras.
albellus, *Muell. Arg. l. c.* 180.—Bras.
albescens, Anders. in Vet. Akad. Handl. Stockh. 1858 (1855) 249 = Scouleri.
albicans, Reichb. f. & Zoll. in Verh. Naturh. Ver. Nederl. Ind. Batav. i. (1856) 21 = Cephalocroton albicans.
albicans, Willd. ex Schlecht. in Linnæa, vi. (1831) 684 = rhamnifolius.
albidus, *Muell. Arg. in DC. Prod.* xv. 11. 645.—Cuba.
Algernoni, Baill. Adansonia, iv. (1863–64) 320 = grandivirsus.
alnifolius, *Lam. Encyc.* ii. 305.—Peruv.
alnoides, *Baill. Adansonia*, iv. (1863–64) 319.—Bras.
althæifolius, *Mill. Gard. Dict.* ed. VIII. n. 9; *Vahl, in Geisel. Cryt. Monog.* 57.—Am. trop.
althæifolius, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 12 (ex. althieoslius).—Malays.
amablis, *Muell. Arg. in Flora*, xlvii. (1864) 587.—Afr. trop.
amazonicus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 102.—Bras.

CROTON :—
ambanivoulensis, *Baill. Adansonia*, i. (1860–61) 165.—Madag.
andinus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 126.—Bolivia.
Andreannus ×, Linden, in André, Illustr. Hortic. xxii. (1876) 56, t. 201.
Andreanus, Linden, ex Rev. Hortic. (1877) 87 = præc.
angolensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 339.—Afr. trop.
angularis, Klotzsch, ex Baill. in Adansonia, iv. (1863–64) 299 = salutaris.
angustifrons, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 223.—Bras.
angustissimus, *Hort. in Gard. Chron.* (1871) 612.—Hab. ?
anisatus, *Baill. Adansonia*, i. (1860–61) 159.—Madag.
anisodontus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 91.—Bras.
antisiphiliticus, *Mart. Reise Bras.* i. 382.—Bras.
apetalum, Blume, Cat. Gew. Buitenz. 104 = Alchornea rugosa.
argenteus, Forsk. Fl. Aegypt. Arab. p. lxxv = Chrozophora obliqua.
argenteus, Linn. Sp. Pl. 1004 = Julocroton argenteus.
argentinus, *Muell. Arg. in Journ. Bot.* xii. (1874) 200.—Reg. Argent.
argothamnoides, Bert. ex Spreng. Syst. iii. 872 = Argithamnia Fendleri.
argutus, Heyne, in Wall. Cat. n. 7768 = Klotzschianus.
argyranthemus, *Micks. Fl. Ber. Am.* ii. 215.—Am. bor.
argyratus, *Blume, Bijdr.* 602.—Ind. or.; Malays.
Argyrodaphne, *Baill. Adansonia*, i. (1860–61) 146.—Madag.
argyroglossus, Baill. l. c. iv. (1863–64) 290 = micans.
argyrophylloides, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 96.—Bras.
argyrophyllus, H. B. & K. Nov. Gen. et Sp. ii. 68 = micans.
arnbemicus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 115.—Austral.
aromaticus, Gaertn. Fruct. ii. 119. t. 107 = candatus.
aromaticus, *Linn. Sp. Pl.* 1005.—Ind. or.
aromaticus, *Mig. Fl. Ind. Bat.* i. 11. 580.—Ins. Molucc.
aromaticus, Wall. Cat. n. 7773 = birmanicus.
asper, Desv. ex Baill. Adansonia, iv. (1863–64) 300 = floribundus.
asper, Kunth, ex Roxb. Hort. Beng. 104; Fl. Ind. iii. 581 = Chrosophora plicata.
asperrimus, *Benth. in Hook. Kew Journ.* vi. (1854) 276.—Bras.
astranatus, Baill. Adansonia, iv. (1863–64) 341 = Klotzschii.
astrogynus, *Baill. l. c.* 310.—Bras.
Astrolites, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 375.—Ins. Thom.
Astrotes, Willd. Sp. Pl. iv. 554 = flavens.
atro-rufus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 194.—Bras.
atrovirens, Wall. Cat. n. 7771 = Mallotus atrovirens.
aucubæfolius, *Andr. Illustr. Hortic.* xix. (1872) 827.—Ins. Pacif.
Augustinianus, Baill. Adansonia, iv. (1863–64) 320 = grandivelum.
auriculatus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 98 = jacobinensis.
avenius, Geisel. Crot. Monog. 11 = luciclus.
axillaris, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 126.—N. Granat.
baccifervs, Wall. Cat. n. 7826 = Mallotus repandus.
bahiensis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 127.—Bras.
Balfourianus, *Muell. Arg. l. c.* 85.—Peruv.
baliospermum, Span. in Linnæa, xv. (1841) 848 = Codiæum variegatum.
balsamens, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 107.—Peruv.
balsamifer, Jacq. Enum. Pl. Carib. 32 = flavens.
barbatus, *H. B. & K. Nov. Gen. et Sp.* ii. 73.—Mexic.
bellulus ×, Linden & André, Illustr. Hortic. xxii. (1875) 104.
Benthamianus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 106.—Bras.
Bentzoe, Linn. Mant. ii. 297 = Terminalia angustifolia.
Benzoe, Murr. Syst. Veg. ed. XIII. 721 = præc.

CROTON :—
Berlandiera, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 141.—Mexic.
Berlandiera, Torr. in Bot. Mex. Bound. 198 = humilis.
Bernlera, *Baill. Adansonia*, i. (1860–61) 152.—Madag.
betaceus, *Baill. l. c.* iv. (1863–64) 341.—Bras.
Betnlester, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 201.—Bras.
betulinus, *Vahl, Symb. Bot.* ii. 98.—Ind. occ.
bicolor, Roxb. Hort. Beng. 69 ; Fl. Ind. iii. 680 = argyrunus.
bidentatus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 140.—Bras.
bifurcatus, *Baill. Adansonia*, i. (1860–61) 164.—Madag.
Billbergianus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 98.—Panama.
birmanicus, *Muell. Arg. l. c.* 112.—Burma.
bispinosus, *Wright, in Sauv. Fl. Cub.* 128.—Cuba.
bixoides, *Vahl, in Geisel. Crot. Monog.* 58.—Am. austr.
Blanchetianus, *Baill. Adansonia*, iv. (1863–64) 301.—Bras.
Bocquilloni, *Baill. l. c.* 161.—Madag.
Boisseri, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 138.—Peruv.
Boivinianus, *Baill. Adansonia*, i. (1860–61) 168.—Madag.
Bojerianus, *Baill. l. c.* 151.—Madag.
bolivianus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 91.—Bolivia.
Bomplandianus, Baill. Adansonia, iv. (1863–64) 339 = C. Persicaria.
Boraginchi, Wall. Cat. n. 7785 = longifolius.
Boutonianus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 80.—Ins. Maurit.
brachiatus, Marc. ex Baill. Etud. Gen. Euph. 368 = serratus.
brachybotryus, *Muell. Arg. in DC. Prod.* xv. 11. 571.—Madag.
bracteatus, *Lam. Encyc.* ii. 308.—Madag.
bractiferus, Roxb. Hort. Beng. 104 ; Fl. Ind. iii. 680 = Codiæum variegatum.
brasiliensis, Mart. ex Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. (1842) 419 = compressus.
brasilianus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 104.—Bras.
Bredemeyeri, *Muell. Arg. in Flora*, lv. (1872) 3.—Venezuela.
brevifolius, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 241 = Scouleri.
brevispicatus, Baill. Adansonia, i. (1860–61) 152 = brachybotryus.
Bridgesii, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 115.—Bolivia.
Brocchianus, Vis. Pl. Aegypt. ac Nub. 39. t. 8. f. 2 = Chrozophora Brocchiana.
Burchellii, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 189.—Bras.
buxifolius, *Muell. Arg. in DC. Prod.* xv. 11. 676.—Bras.
Cajucara, *Benth. in Hook. Kew Journ.* vi. (1854) 276.—Bras.
caldasia, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 93.—Bras.
californicus, *Muell. Arg. in DC. Prod.* xv. 11. 691.—Calif.
calliarpaefolius, *Vahl, in Geisel. Crot. Monog.* 37.—Peruv.
calocephalus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 121.—Bras.
calococcus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 242.—Burma.
calycinus, *Sprang. Neu Entdeck.* iii. 24.—Bras.
campestris, *St. Hil. Fl. Us Bras.* i. 66.—Bras.
canescens, Boj. Hort. Maurit. 221 = mauritianus.
caperasia, Linn. f. Suppl. 422 = Jatropha caperasia.
caperonifolius, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 206.—Bras.
capitatus, *Hook. Fl. Ber. Am.* ii. 214.—Am. bor.
cardiospermus, *Guerin. Fruct.* ii. 190. t. 107.—Zeylan.
carinatus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 126.—Bras.
Caryophyllus, *Benth. in Hook. Kew Journ.* vi. (1854) 274.—Bras.
Casarettoanus, *Muell. Arg. in DC. Prod.* xv. 11. 692.—Bras.
Cascarilla, *Bonpl. in Journ. Linn. Soc.* iv. (1860) 30.—Ins. Bahama.
Cascarilla, Lam. Encyc. ii. 308 = flavens.
Cascarilla, Linn. Sp. Pl. ed. II. 1429 = lineatis.

CROTON :—
cascarilloides, *Geisel. Crot. Monog.* 8.—Ins. S. Doming.
cascarilloides, Raeusch. Nom. ed. III. 260 = Mallotus philippinensis.
casuinoides, *Lam. Encyc.* ii. 211.—Madag.
catanaefolius, H. B. & K. Nov. Gen. et Sp. ii. 70 = Caperonia palustris.
catanaefolius, Linn. Sp. Pl. 1005 = Caperonia castaneaefolia.
catanifolius, Wall. Cat. n. 7760 = Mallotus Wallichianus.
Catarinæ, *Baill. Adansonia*, iv. (1863–64) 349.—Bras.
catiogunus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 263.—Bras.
candatus, *Geisel. Crot. Monog.* 73.—Ind. or.
ceanothifolius, *Baill. Adansonia*, iv. (1863–64) 295.—Bras.
cearensis, *Baill. l. c.* 327.—Bras.
cearensis, Muell. Arg. in Linnæa, xxxiv. (1865–66) 119 = betaceus.
celtidifolius, *Baill. Adansonia*, iv. (1863–64) 331.—Bras.
cerino-dentatus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 186.—Bras.
certusa, *Muell. Arg. in DC. Prod.* xv. 11. 570.—Cuba.
chaetocalyx, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 135.—Bras.
chaetophorus, *Muell. Arg. l. c.* 130.—Bras.
chamaedryfolius, *Griseb. Fl. Brit. W. Ind.* 41.—Guiana.
chamaedryfolius, Lam. Encyc. ii. 214 = Acalypha reptans.
Chamaepitys, Baill. Adansonia, iv. (1863–64) 356 = pycnocephalus.
chapadensis, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 203.—Bras.
Chapelieri, *Baill. Adansonia*, i. (1860–61) 166.—Madag.
Chiamala, Wall. Cat. n. 7775 = Alchornea tiliaefolia.
chilensis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 92.—Chili.
chinensis, Benth. Fl. Hongk. 309 = tomentosus.
chinensis, Geisel. Crot. Monog. 24 = Mallotus Apelta.
chlorocalyx, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 109.—Ind. or.
chlorolencus, *Muell. Arg. l. c.* 106.—Bras.
chrosophoroides, Auct. vide *crosophoroides*.
chrysocladus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 172.—Bras.
Chrysodraphne, *Baill. Adansonia*, i. (1860–61) 147.—Madag.
ciliato-glanduliferus, *Ortega, Hort. Matr.* 51.—Mexic.; Cuba.
cinctus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 285.—Bras.
cinereilus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 137.—Bras.
cirrifolius, *Lam. Encyc.* ii. 210.—Ins. S. Doming.
cladotrichus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 151.—Mexic.
Classenianus, *Baill. Adansonia*, iv. (1863–64) 291.—Bras.
clavuligor, *Muell. Arg. in DC. Prod.* xv. 11. 644.—Cuba.
clusyioides, Forst. f. Prod. 99 = Euphorbia tannensis.
coccorifolius, *Muell. Arg. in DC. Prod.* xv. 11. 676.—Bras.
coccineus, *Noronha, in Verh. Batav. Gen.* v. (1790) ad. I. Art. iv. 11.—Malaya.
coccifolius, Vahl, Symb. Bot. ii. 97 = Mallotus philippinensis.
coccineus, Wall. Cat. n. 7771 = Mallotus atrovirens.
codonocalyx, Baill. Adansonia, ix. (1868–64) 353 = nitrariaefolius.
Coelebogyne, Baill. l. c. 293 = Gandichandi.
Colligaya, *Molina, Sagg. Chil. ed.* ii. 147.—Chili.
Colligaya, Sprang. Syst. iii. 875 = Colligaya odorifera.
collinus, H. B. & K. Nov. Gen. et Sp. ii. 75 = ferrugineus.
collinus, Pauch. ex Baill. Adansonia, ii. (1861–62) 217 = insularis.
collinus, *Phil. Fl. Atac.* 49.—Chili.
comatus, *Vahl. Fl. Hisp.* v. 2. 74.—Bras.
comosus, *Muell. Arg. in DC. Prod.* xv. 11. 667.—Bras.
compressus, *Lam. Encyc.* ii. 208.—Bras.
condylolepsus, H. B. & K. Nov. Gen. et Sp. ii. 80 = rhamnifolius.

CROTON :—
congestus, *Lour. Fl. Cochinch.* ii. 582.—China.
consanguineus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 117.—Ins. Philipp.
conspurcatus, Schlecht. in Linnæa, vii. (1832) 380 = Julocroton conspurcatus.
constrictus, *Baill. Adansonia*, iv. (1863–64) 360.—Bras.
corallatus, Soland. ex Baill. Etud. Gén. Euph. 351 = Ricinocarpus pinifolius.
corchorifolius, Geisel. Crot. Monog. 25 = glandulosus.
Corchorupais, *Baill. Adansonia*, iv. (1863–64) 364.—Bras.
cordatus, Vell. Fl. Flum. x. t. 66 = Julocroton triquetor.
cordisefolius, *Baill. Adansonia*, iv. (1863–64) 334.—Bras.
coriaceus, *H. B. & K. Nov. Gen. et Sp.* ii. 87.—Ecuador.
cornutus, *André, Illustr. Hortic.* xix. (1872) 188.—Ins. Salomon.
corrientesianus, *Baill. Adansonia*, iv. (1863–64) 309.—Am. austr.
Cortesianus, *H. B. & K. Nov. Gen. et Sp.* ii. 83.—Mexic.
coryifolius, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 159 = cubanus.
corylifolius, *Lam. Encyc.* ii. 205.—Jamaic.
corymbulosus, *Rothr. in Wheeler, Rep. Bot.* 242.—Am. bor.
costatus, *H. B. & K. Nov. Gen. et Sp.* ii. 67.—N. Granat.
Cotoneaster, *Muell. Arg. in Flora*, xlvii. (1864) 484.—Madag.
craspedoricbus, *Griseb. in Gœtt. Nachr.* (1865) 173.—Cuba.
crassifolius, Geisel. Crot. Monog. 19 = tomentosus.
crassivenosus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 108 = astrogynus.
cremostachyus, Baill. Adansonia, iv. (1863–64) 321 = grandivelon.
crenatus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 129.—Venezuela.
crosophoroides, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 248 = tomentosus.
cubanus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 94.—Cuba.
Cumingii, *Muell. Arg. l. c.* 101.—Ins. Philipp.
cuneatus, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 49.—Bras.
cuneatus, Miq. in Linnæa, xxii. (1849) 477 = surinamensis.
cyanospermus, Gaertn. Fruct. ii. 120. t. 107 = Phyllanthus cyanospermus.
cynanchifolius, Baill. Adansonia, iv. (1863–64) 329 = celtidifolius.
daphnargyreus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = C. Argyrodaphne.
debilis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 132.—Bras.
decalobus, Muell. Arg. l. c. 80 = Julocroton decalobus.
dedipiens, *Baill. Adansonia*, iv. (1863–64) 310.—Bras.
dentatus, Sm. in Rees, Cyclop. x. n. 40 = populifolius.
denticulatus, Blume, Bijdr. 608 = candatus.
denticulatus, Geisel. Crot. Monog. 72 = Euphorbia lanata.
denticulatus, Wall. Cat. n. 7781 = rhodostachyus.
dentifolius, Baill. Adansonia, iv. (1863–64) 347 = Lundianus.
dentosus, *Griseb. in Gœtt. Abh.* xix. (1874) 97.—Reg. Argent.
dennulatus, Baill. Adansonia, iv. (1863–64) 319 = agrarius.
Deppeanus, Steud. Nom. ed. II. i. 446 = fragilis.
desertorum, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 265.—Bras.
diadenus, Miq. Fl. Ind. Bat. Suppl. 451 = laevifolius.
dichotomus, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 197.—Bras.
dichotomus, Willd. Sp. Pl. iv. 537 = Bernardia dichotoma.
dichrous, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 106.—Bras.
diffusus, Geisel. Crot. Monog. 41 = ovalifolius.
digitatus, Pisch. ex Steud. Nom. ed. I. 240; ed. II. i. 446 = Manihot digitata.
digitatus, Mart. ex G. Don, in trans. Hort. Brit. ed. III. 601 = Manihotifolia.

CROTON :—
digitatus, Hort. Kew. ex Muell. Arg. in DC. Prod. xv. ii. 696 = lacerus.
digitatus, Salzm. ex Schlecht. in Linnæa, xix. (1847) 340 = lobatus.
digitifolius, Baill. Adansonia, iv. (1863–64) 340 = connatus.
dioicus, *Cav. Ic.* i. 4. t. 6.—Mexic.
dioicus, Iserl, ex Geisel. Crot. Monog. 18 = dicolor.
dioicus, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 680.—Malaya.
dioicus, Willd. Sp. Pl. iv. 584 = gracilis.
dicolor, Wright, ex Griseb. in Goett. Nachr. (1865) 173 = origanifolius.
dicolor, *Willd. Sp. Pl.* iv. 582.—Ins. S. Doming.
diepuncticiflorus, Miqu. Fl. Bor. Am. ii. 215 = maritimus.
distans, Wall. Cat. n. 7772 A = Mallotus philippinensis.
divaricatus, Sw. Prod. Veg. Ind. Occ. 100 = glandulosus.
domingensis, A. Rich. Fl. Cub. Fanerog. ii. 213 = sonberrimus.
domingensis, Vahl, in Geisel. Crot. Monog 42 = sidæfolius.
dorstophyllus, Baill. Adansonia, iv. (1863–64) 361 = Julocroton dorstophyllus.
Draco, *Schlecht. in Linnæa*, vi. (1831) 360.—Mexic. draconoides, *Muell. Arg. l. c.* xxxiv. (1865–66) 90.—Ins.
draconoides, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = seq.
Draconopsis, *Muell. Arg. in Journ. Bot.* ii. (1861) 338.—Afr. occ.
draconoloides, *Baill. Adansonia*, iv. (1863–64) 365.—Ins.
drupaceus, Blanco, Fl. Filip. ed. II. 519 = luzoniensis.
drupaceus, Roxb. Hort. Beng. 69; Fl. Ind. iii. 684 = caudatus.
drupaceus, Wall. Cat. n. 7744 = birmanicus!
echinocarpus, Baill. Adansonia, iv. (1863–64) 320 = grandivelum.
echinocarpus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 58.—Ins.
echloides, *Baill. Adansonia*, iv. (1863–64) 324.—Ins. Ehrenbergii, *Schlecht. in Linnæa*, xix. (1847) 248.—Mexic.
Eichleri, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 272.—Bras.
elæagnifolius, Vahl, in Geisel. Crot. Monog. 9 = dioicus.
elæagnoides, *Baill. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 95.—Ins. Socotr.
elæocarpifolius, Wall. Cat. n. 7734 = oblongifolius.
elegans, H. B. & K. Nov. Gen. et Sp. ii. 79 = ferrugineus.
Elliottii, *Chapm. Fl. U.-St.* 407.—Am. bor.
elliptieus, Ell. Sketch. ii. 648 = Elliottii.
elliptieus, *Geisel. Crot. Monog.* 37.—Jave.
elliptieus, Nutt. Gen. Am. ii. 225 = monanthogynus.
elliptieus, A. Rich. Fl. Cub. Fanerog. ii. 213 = pervestitus.
Elsteria, *Benn. in Journ. Linn. Soc.* iv. (1860) 26.—Ins. Baham.
Elsteria, Sw. Prod. Veg. Ind. Occ. 100 = glabellus.
Emetier, Baill. Adansonia, i. (1860–61) 166 = bifurcatus.
emirnensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 252.—Madag.
eremophilus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 154.—Bras.
erianthus, Sm. in Rees, Cyclop. x. n. 21 = laxiflorus.
ericoides, *Baill. Adansonia*, iv. (1863–64) 329.—Bras.
ericoloides, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 228.—Bras.
eriochus, *Muell. Arg. l. c.* 229.—Bras.
eriospermus, Lam. Encyc. ii. 272 = Trigonia crotonoides.
erythradilus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 93.—Peruv.
erythroxyloides, Baill. Adansonia, iv. (1863–64) 327 = chlorolencus.
erythroxyloides, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 111.—Bras.
eucnyphæoides, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 26 = populifolius.
eucnyphæus, *A. Gray, Mem. Bot.* ed. V. 438.—Texas.

CROTON :—
exuberans, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 85.—Bras.
fallax, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 127.—Bras.
farinosus, *Lam. Encyc.* ii. 271.—Madag.
fastuosus, *Baill. Adansonia*, iv. (1863–64) 330.—Bras.
ferrugineliua, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 174.—Bras.
ferrugineus, *H. B. & K. Nov. Gen. et Sp.* ii. 77.—Peruv.
fabellifolius, Ledd. Cat. ex G. Don, in Sweet, Hort. Brit. ed. III. 801 = bibaciifolius.
flavens, *Linn. Pl. Jam. Pugill.* 28; *Syst.* ed. X. 1276.—Am. austr.
flocculosus, Geisel. Crot. Monog. 14 = flavens.
flocculosus, *Kurz, in Journ. As. Soc Beng.* xlii. (1873) 11. 242.—Burma.
floribundus, Lund, ex Didr. in Kjoeb. Videnak. Meddel. (1857) 130 = urticæfolius.
floribundus, *Spreng. Syst.* iii. 873.—Bras.
fœtens, Anders. in Trans. Soc. Arts, xxv. (1807) 302 = heliotidens.
fothergillæfolius, *Baill. Adansonia*, i. (1860–61) 160.—Ins Maurit.
fragilis, *H. B. & K. Nov. Gen. et Sp.* ii. 75.—Am. trop.
fragilis, *Schlecht, in Linnæa*, xxvi. (1855) 688.—N. Granat.
fragrans, *H. B. & K. Nov. Gen. et Sp.* ii. 81.—N. Granat.
Francavillanus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 111.—Cuba.
Franciscanus, *Muell. Arg l. c.* 124.—Mexic.
Fraseri, *Muell. Arg. l. c.* 118.—Peruv.
Friesenaus, *Muell. Arg. in DC. Prod.* xv. ii. 543.—Am. austr. Bras.
Frionis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 176.—Bras.
fruticosus, Mill. Gard. Dict. ed. VIII. n. 6 = lucidus.
fruticulosus, *Torr. Bot. Mex. Bound.* 194.—Texas.
fulvus, *Mart. Reise Bras.* i. 282.—Bras.
fulvus, *A. Rich. Fl. Cub. Fanerog.* ii. 212.—Cuba.
Funckianus, *Muell. Arg. in DC. Prod.* xv. ii. 546.—N. Granat.
furfuraceus, Pers. Syn. ii. 586 = punctatus.
fuscescens, Spreng. Syst. ii. 874 = Julocroton fuscescens.
fuscus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 151.—Bras.
Galeottianus, Baill. Etud. Gen. Euph. 356 = nummulariæfolius.
Garckeanus, *Baill. Adansonia*, iv. (1863–64) 308.—Bras.
Gardneri, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 136.—Bras.
Gardnerianus, *Baill. Adansonia*, iv. (1863–64) 301.—Bras.
Gaudichaudi, *Baill. l. c.* 293.—Bras.
Gibsonianus, *Nimmo, in S. Grah. Cat. Pl. Bomb.* 251.—Ind. or.
Gibsonii, J. Grah. l. c. 192 = Rheedei.
glabellus, Heyne, ex Wall. Cat. n. 8012 = Dimorphocalyx glabellus.
glabellus, *Linn. Syst.* ed. X. 1275.—Am. trop.
glabrescens, *Miq. Fl. Ind. Bat.* i. 11. 562.—Java.
glandulatus, *Veli. Fl. Flum.* x. t. 78.—Bras.
glandulifurus, Vahl, in Geisel. Crot. Monog. 37 = lucidus.
glandulosus, Blanco, Fl. Filip. ed. I. 754 = muricatus.
glandulosus, *Linn. Syst.* ed. X. 1275.—Am. bor. et trop.
glandulosus, Vell. Fl. Flum. x. t. 78 = pungens.
Glaziovii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 94.—Bras.
glechonæfolius, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 126.—Bras.
gleisosus, Sw. Prod. Veg. Ind. Occ. 100 = Mottenia globosa.
glutinosus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 200.—Bras.
glycosmeus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 140 = grewioides.
glycyrrhizoides, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 117.—Bras.
Gnaphalii, *Baill. Adansonia*, iv. (1863–64) 329.—Bras.
gnaphaloides, *Schrad. in Gott. Gel. Anz.* i. (1821) 708.—Bras.

CROTON :—
gnidiaceus, *Baill. Adansonia*, iv. (1863–64) 354.—Bras.
gonocladus, Mart. in Flora, xx. (1837) ii. Beibl. 119 = compressus.
gossypiifolius, *Vahl, Symb. Bot.* ii. 98.—Am. centr.
Goudotii, *Baill. Adansonia*, i. (1860–61) 167. t. 5.—Madag.
goyazensis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 120.—Bras.
gracilescens, *Muell. Arg. l. c.* 110.—Bras.
gracilipes, *Baill. Adansonia*, iv. (1863–64) 333.—Bras.
gracilis, *H. B. & K. Nov. Gen. et Sp.* ii. 69.—Mexic.
grandifolius, Blanco, Fl. Filip. ed. I. 753 = Macaranga moluccana.
grandivelum, *Baill. Adansonia*, iv. (1863 64) 322.—Bras.
grangerioides, *Boj. Hort. Maurit.* 281, nomen; *Baill. Adansonia*, i. (1860–61) 149.—Ins. Maurit.
gratissimus, *Burch. Trav.* ii. 268.—Afr. austr.
grewioides, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 87.—Mexic.
grewioides, *Baill. Adansonia*, iv. (1863–64) 365.—Bras.
Grisebachianus, *Muell. Arg. in Flora*, xlvii. (1864) 484.—Jamaic.
guianensis, *Aubl. Pl. Gui.* ii. 882.—Guian.
Gnilldingii, *Griseb. Fl. Brit. W. Ind.* 38.—Ins. S. Vincent.
gummiferus, A. Cunn. ex Planch. in Hook. Lond. Journ. Bot. iv. (1845) 473 = Bertya gummifera.
guyanensis, Auct. vide guianensis.
Hadrianı, *Baill. Adansonia*, iv. (1863–64) 313.—Bras.
Halecum, Rozb. Hort. Beng. 104; Fl. Ind. iii. 688 = Claoxylon indicum.
Hanceı, *Benth. Fl. Hongk.* 308.—Hongkong
Hasskarlianus, *Muell. Arg. in DC. Prod.* xv. ii. 573.—Ins Madura.
hastatus, Burm. f. Fl. Ind. 205. t. 63. f. 1 = Chrozophora plicata.
hastatus, Linn. Sp. Pl. 1005 = Tragia involucrata.
heeianandrus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 705.—Bras.
helichrysoides, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 123 = seq.
Helichrysum, *Baill. Adansonia*, iv. (1863–64) 358.—Bras.
heliocidens, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 97.—Ins. S. Vincent.
heliotropiifolius, H. B. & K Nov. Gen. et Sp. ii. 88 = rhamnifolius
hemiargyreus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 81.—Bras.
herbaceus, Vell. Fl. Flum. x. t. 69 = glandulosus.
heteroclyx, *Baill. Adansonia*, iv. (1863–64) 324.—Bras.
heterocroton, *Muell. Arg. in DC. Prod.* xv. ii. 621.—Sumatra.
heterochrous, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 121.—Am. centr.
heterodoxus, *Baill. Adansonia*, iv. (1863–64) 300.—Bras.
heteroneurus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 114.—Reg. Argent.
heterophyllus, H. B. & K Nov. Gen. et Sp. ii. 88 = gossypiifolius.
heterolrichus, *Muell. Arg. in DC. Prod.* xv. ii. 620.—Ins. Fiji.
Hialmarsonii, *Griseb. Fl. Brit. W. Ind.* 40.—Ind. occ. hibiscifolius, *Kunth, ex Spreng. Syst.* iii. 876.—N. Granat.
Hieronymi, *Griseb. in Goett. Abh.* xxiv. (1879) 54.—Reg. Argent.
hieronymoides, Muell. Arg. in Linnæa, xxxiv. (1865–66) 98 = Sellowii.
Hilarii, *Baill. Adansonia*, iv. (1863–64) 332.—Bras.
Hillianus, *Veitch, in André, Illustr. Hortic.* xix. (1872) 326.—Ins. Pacif.
hippophaeoides, A. Rich. Fl. Cub. Fanerog. ii. 212 = linearis.
hircinus, Went. Jard. Malm. 80 = populifolius.
hirsutus, Thou. ex Baill. Adansonia, i (1860–61) 167 = Thouarsianus.
hirsutus, Vell. Fl. Flum. x. t. 65 = Julocroton fuscescens.
hirtus, L'Hérit. Stirp. Nov. 17. t. 9 = glandulosus.
hispidus, H. B. & K. Nov. Gen. et Sp. ii. 79 = populifolius.

CROTON :—
Hoffmanni, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 58.—Costa Rica.
Holtonis, *Muell. Arg. l. c.* 115.—N. Granat.
homolepidus, *Muell. Arg. in DC. Prod.* xv. ii. 518.—Cuba.
Hookerianus, Baill. Etud. Gén. Euph. 358 = lucidus.
horminoidenus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = seq.
Hornkium, *Baill. Adansonia*, iv. (1863–64) 315.—Bras.
horridulus, *Baill. Adansonia*, iv. (1863–64) 330.—Bras.
Hostmanni, *Miq. ex Schlecht. in Linnæa*, xxi. (1848) 477.—Guiana.
humilis, *Linn. Syst. ed.* X. 1276 ; *Amœn. Acad.* v. 410, 1759.—Jamaica.
hypoleucus, Dals. in Hook. Kew Journ. Bot. iii. (1851) 193 = reticulatus.
hypoleucus, *Schlecht. in Linnæa*, xix. (1847) 246.—Mexic.
impetiginosus, Poepp. ex Baill. Adansonia, iv. (1863–64) 298 = matourensis.
incanus, Anderss. in Vet. Akad. Handl. Stockh. 1858 (1855) 348 = Sconlerl.
incanus, Blume, ex Baill. Etud. Gén. Euph. 470, obs. = Pretoria incana.
incanus, *H. B. & K. Nov. Gen. et Sp.* ii. 78.—Mexic.
incertus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 177.—Bras.
incisus, *Baill. Adansonia*, i. (1860–61) 159.—Madag.
indivisus, *Vell. Fl. Flum.* x. t. 68.—Bras.
Inophyllum, Forst. f. Prod. 67 = Codiæum Inophyllum.
insularis, *Baill. Adansonia*, ii. (1861–62) 217.—N. Caled.
interodens, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 168.—Bras.
intricata, *Wright, in Sauv. Fl. Cub.* 109.—Cuba.
inversus, Baill. Adansonia, iv. (1863–64) 321 = pianthiensis.
irregularis, *André, Illustr. Hortic.* xix. (1872) 185.—Polynes.
Isabelli, *Baill. Adansonia*, iv. (1863–64) 318.—Bras.
Isertl, *Geisel. Crot. Monog.* 94.—Ins Martin.
Isertl, Griseb. Fl. Brit. W. Ind. 39 = Grisebachianus.
itabirensis, Baill. Adansonia, iv. (1863–64) 296 = cesnotbifolius.
Itacolum, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 246.—Bres.
jacobinensis, *Baill. Adansonia*, iv. (1863–64) 302.—Bras.
jacobinensis, Muell. Arg. in Linnæa, xxxiv. (1865–66) 91 = echioides.
Jacquemanus, *Muell. Arg. in Flora*, lv. (1872) 7.—Ind. occ.
Jamalgota, Buch.-Ham. in Trans. Linn. Soc. xiv. (1824) 258 = C. Tiglium.
Jamesoni, *Muell. Arg. in DC. Prod.* xv. ii. 549.—Ecuador.
japirunia, Muell. Arg. in Linnæa, xxxiv. (1865–66) 129 = laurifolius.
japonicus, Thunb. Fl. Jap. 270 = Mallotus japonicus.
Jardini, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 119.—Ins. Martin.
jatrophe, *Muell. Arg. in DC. Prod.* xv. ii. 667.—Bras.
jatrophæfolius, *Muell. Arg. l. c.* 600.—Java.
Jenynsana, *Gris, ex Baill. Adansonia*, i. (1860–61) 160.—Madag.
Josephinus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 195.—Bras.
Joufas, *Raxb. Hort. Beng.* 104 ; *Fl. Ind.* iii. 685.—Ind. or.
Jolopaidum, *Baill. Adansonia*, iv. (1863–64) 313.—Bras.
juncens, *Baill. l. c.* 329.—Bras.
Kalkmanni, *Muell. Arg. in DC. Prod.* xv. ii. 612.—Bras.
Klotzschi, *Muell. Arg. in Mart. Fl. Bras* xi. ii. 202.—Bras.
Klotzschianus, *Thw. Enum. Pl. Zeyl.* 276.—Ind. or.
Klotzschii, Baill. Adansonia, iv. (1863–64) 346 = glandulosus.
Klotzschii, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 126.—Bras.
Korthalsii, *Muell. Arg. in DC. Prod.* xv. ii. 627.—Borneo.
Kroneanus, *Miq. in Journ. Bot. Néerl.* i. (1861) 97.—China.

CROTON :—
laccifervus, Blanco, Fl. Filip. ed. I. 731 = Macaranga Tanarius.
laccifervus, Linn. Sp. Pl. 1006 = aromaticus.
laccifervus, Wall. Cat. n. 7774 = Mallotus repandus.
lacerus, *Desf. Tabl. ed.* I. 206.—Hab. ?
lachnocarpus, *Benth. in Hook. Kew Journ.* vi. (1854) 5.—Hongkong.
lachnocladus, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* xi. ii. 258.—Bras.
lachnostachys, *Baill. Adansonia*, iv. (1863–64) 363.—Am. austr.
lachnostephanus, Baill. l. c. 357 = pycnocephalus.
lanifolius, Baill. l. c. 321 = campestris.
lanius, Wall. Cat. n. 7788 = Codiocissus lappaceus.
lasvifolius, *Blume, Bijdr.* 608.—Ind. or. ; Malaya.
lasvigatus, *Vahl, Symb. Bot.* ii. 97.—China.
lasvigatus, Wall. Cat. n. 7749 = Actephila excelsa.
lasvigatus, Wall. l. c. 7795 = oblongifolius.
lagoensis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 99.—Bras.
lanatus, Klotzsch, ex Baill. Etud. Gén. Euph. 370 = lanuginosus.
lanatus, *Lam. Encyc.* ii. 211.—Bras.
lanatus, Lour. Fl. Cochinch. ii. 561 = lasianthus.
lancrolatus, Cav. Ic. vi. 38 = Argithamnia tricuspidata.
lancrolatus, Hornem. ex Dietr. in Kjoeb. Vidensk. Meddel. (1857) 148 = Caperonia heteropetala.
Langsdorffii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 168.—Bras.
lanuginosus, *Baill. Adansonia*, iv. (1863–64) 355.—Bras.
lanuginosus, Schum. ex Schweinf. Pl. Nilot. 10 = Chrozophora plicata.
lapanus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 258.—Bras.
Lasegra, *Muell. Arg. l. c.* 184.—Bras.
lasianthus, *Pers. Syn.* ii. 586.—Cochinch.
laurifolius, Muell. Arg. in DC. Prod. xv. ii. 586 = laurinus.
lauriticus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 18.—Malaya.
lauriifolius, *Spach, ex Baill. Adansonia*, iv. (1863–64) 68.—Bras.
laurinus, *Sw. Prod. Veg. Ind. Occ.* 100.—Jamaic.
Lawianus, Nimmo, in J. Grah. Cat. Pl. Bomb. 261 = Trigonostemon Lawianus.
laxiflorus, *Geisel. Crot. Monog.* 44.—Cochinch.
Lechleri, *Muell. Arg. in DC. Prod.* xv. ii. 545.—Peruv.
leiocarpus, Bartl. Ind. Sem. Hort. Gotting. (1855) 3 ; Linnæa, xxviii. (1856) 858 = lobatus.
leiophyllus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 108.—Ins. Philipp.
leprosus, Sprang. ex Griseb. Fl. Brit. W. Ind. 38 = flavens.
leprosus, Willd. Sp. Pl. iv. 558 = Bernardia leprosa.
leptobotrys, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 154.—Bras.
leptophyllus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 108.—Bras.
leptopus, *Muell. Arg. in DC. Prod.* xv. ii. 622.—Ins. Fiji.
leptostachyus, *H. B. & K. Nov. Gen. et Sp.* ii. 79.—N. Granat.
leucadenius, Baill. Adansonia, iv. (1863–64) 338 = fuscus.
leucocephalus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 131 = heterodoxus.
leucophlebius, *Wright, ex Griseb. in Gottt. Nachr.* (1865) 173.—Cuba.
leucophyllus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 126.—Mexic.
Liebmanni, *Muell. Arg. in DC. Prod.* xv. ii. 665.—Mexic.
limbatus, *A. Rich. Fl. Cub. Fenerug.* ii. 212.—Cuba.
Lindheimerianus, *Scheele, in Linnæa*, xxv. (1852) 580.—N. Mexic.
linearifolius, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 162.—Bras.
linearis, *Xacq. Enum. Pl. Carib.* 92.—Ind. occ.
lobatus, Forsk. Fl. Aegypt. Arab. 162 = Jatropha lobata.
lobatus, *Linn. Sp. Pl.* 1005.—Am. trop.
lobatus, Vell. Fl. Flum. x. t. 70 = C. jatrophe.
longifolius, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 170.—Bras.
longifolius, Wall. Cat. n. 7717 = Trigonostemon longifolius.

CROTON :—
longinervius, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 215.—Bras.
Lorentzii, *Muell. Arg. ex Griseb. in Gottt. Abh.* xix. (1874) 96.—Am. trop.
loucombianus, Baill. Adansonia, i. (1860–61) 155 = adenophorus.
lucidus, *Linn. Pl. Jam. Pugill.* 23 ; *Syst.* ed. X. 1275 —Ind. occ.
Lundianus, *Muell. Arg. in DC. Prod.* xv. ii. 662.—Bras.
luridus, *Geisel. Crot. Monog.* 96.—Hab. ?
lutec-brunneus, *Baker, in Journ. Linn. Soc.* xx. (1883) 264.—Madag.
luscinensis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 118.—Ins. Philipp.
Macraei, Hook. f. in Trans. Linn. Soc. xx. (1851) 188 = Sconlerl.
macrobothrys, *Baill. Adansonia*, iv. (1863–64) 329.—Bras.
macrocalyx, Ehrenb. ex Schweinf. Pl. Nilot. 9 = Chrosophora Brocchiana.
macrocalyx, *Mart. ex Baill. Adansonia*, iv. (1863–64) 303.—Bras.
macrocarpus, Reichb. ex Muell. Arg. in DC. Prod. xv. ii. 698 = Excoecaria sebifera.
macrodontus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 128.—Mexic.
macrophyllus, Sw. Prod. Veg. Ind. Occ. 100 = Lasiocroton macrophyllus.
macrostachys, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 251.—Abyss.
maculatus, *Vahl, in Geisel. Crot. Monog.* 30.—Java.
magdalenensis, *Muell. Arg. in Flora*, xlvii. (1864) 484 —N. Granat.
malabaricus, *Bedd. Forest. Man. Bot.* 204.—Ind. or.
malacophyllus, Benth. Pl. Hartw. 248 = leptostachyus.
malacotrichus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 101 = lanuginosus.
Malambo, *Karst. l. c.* xxvii. (1856) 418.—Venezuela
Mandonis, *Muell. Arg. l. c.* xxxiv. (1865–66) 85.—Bolivia.
Manihot, Baill. Adansonia, iv. (1863–64) 341 = lobatus.
marginatus, A. Cunn. ex Benth. Fl. Austral. vi. 78 = Ricinocarpus marginatus.
maritimus, *Walt. Fl. Carol.* 239.—Texas.
Martii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 88.—Bras.
Massangeanum, Linden, ex André, in Illustr. Hortic. xxvi. (1879) 77. t. 347 = Codiæum variegatum.
matourensis, *Aubl. Pl. Gui.* ii. 879. t. 336.—Am. trop
matronalis, Baill. Adansonia, iv. (1863–64) 309 = vestitus.
mauritianus, *Lam. Encyc.* ii. 205.—Ins. Maurit.
mauritianus, *Jacobs. et Willem. in Usteri, Ann. Bot.* xviii. 56.—Maurit.
mauritianus, Thou. ex Baill. Adansonia, i. (1860–61) 150 = tiliæfolius.
medians, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 169.—Bras.
Meduse, *Muell. Arg. in DC. Prod.* xv. ii. 630.—Bras.
megalobotrys, *Muell. Arg. in Flora*, xlvii. (1864) 537. —Afr. trop.
megalocalyx, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 230.—Bras.
megalodendron, *Muell. Arg. in Flora*, lv. (1872) 4.—Venezuela.
megapontius, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 211.—Bras.
Meissneri, *Muell. Arg. in DC. Prod.* xv. ii. 665.—Mexic.
melanocistus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 185.—Bras.
melanosticta, Boiv. ex Baill. Adansonia, i. (1860–61) 268 = Sellingia melanosticta.
membranaceus, *Muell. Arg. in Flora*, xlvii. (1864) 534. —Afr. trop.
menthodorus, *Benth. Pl. Hartw.* 248.—N. Granat.
metallicus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 108.—Ins. Fiji.
mexicanus, *Muell. Arg. l. c.* 113.—Mexic.
micans, *Sw. Fl. Ind. Occ.* ii. 1236.—Jamaic.
microcalyx, Muell. Arg. in Linnæa, xxxiv. (1865–66) 92 = C. Pavlonis.
microcarpus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 148.—Bras.
microphyllus, Lam. Encyc. ii. 212 = Sadaclina microphylla.

CROTON :—
microstachys, *Baill. Adansonia*, iv. (1868–64) 366.—Bras.
migrans, *Caser. Nov. Stirp. Bras. Dec.* 88.—Bras.
Minarum, Muell. Arg. in Linnæa, xxxiv. (1865–66) 188 = cnemothifolius.
misniifolius, Muell. Arg. l. c. = Gaudichaudii.
miradorensis, *Muell. Arg. in DC. Prod.* xv. ii. 627.—Mexic.
Mollii, *Muell. Arg. in Flora*, lv. (1872) 8.—Ind. occ.
mollis, *Benth. in Hook. Kew Journ.* vi. (1854) 375.—Bras.
mollissimus, Geisel. Crot. Monog. 78 = Mallotus ricinoides.
mollissimus, Vahl, ex Geisel. l. c. 56 = lanatus.
moluccanus, Linn. Sp. Pl. 1005 = Mallotus moluccanus.
moluccanus, Willd. Sp. Pl. iv. 551 = Chrozophora plicata.
monanthogynus, *Michx. Fl. Bor. Am.* ii. 215.—Am. bor.
Mongue, *Baill. Adansonia*, i. (1860–61) 158.—Madag.
montanus, Geisel. Crot. Monog. 28 = corylifolius.
montanus, Wall. Cat. n. 7823 = Coelodiscus montanus.
montanus, Willd. Sp. Pl. iv. 545 = Mallotus philippinensis.
montevidensis, *Spreng. Syst.* iii. 873.—Reg. Argent.
monticola, Steud. Nom. ed. II. i. 446 = corylifolius.
Moonii, *Thw. Enum. Pl. Zeyl.* 276.—Zeylan.
morifolius, *Willd. Sp. Pl.* iv. 535.—Am. trop.
moritzbensis, Baill. Adansonia, iv. (1868–64) 332 = rhamnifolius.
Mubango, *Muell. Arg. in Journ. Bot.* ii. (1864) 338.—Afr. trop.
mucronatus, Willd. Sp. Pl. iv. 542 = flavens.
mucronifolius, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 216.—Bras.
multicostatus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 79.—Ins. S. Doming.
multiglandulosus, Reinw. ex Blume, Cat. Gew. Buitenz. 105 = Mallotus moluccanus.
multispicatus, Vell. Fl. Flum. x. t. 75 = urticæfolius.
muricatus, Blanco, Fl. Filip. ed. II. 518 = C. Tiglium.
muricatus, Bo). Hort. Maurit. 281 = lotheriflæfolius.
muricatus, Heyne, in Wall. Cat. n. 7751 = Mallotus muricatus.
muricatus, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 173 = texensis.
muricatus, *Vahl, in Geisel. Crot. Monog.* 47.—Madag.
muricatus, *Zipp. ex Span. in Linnæa*, xv. (1841) 348, nomen.—Ins. Timor.
Muscicapæ, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 199.—Bras.
Mutisianus, *H. B. & K. Nov. Gen. et Sp.* ii. 87.—N. Granat.
myriadenus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 130 = Garckeanus.
myrianthus, *Muell. Arg. l. c.* 188.—Bras.
myrinster, *Baker, in Journ. Bot.* xx. (1882) 268.—Madag.
myrtifolius, *Griseb. in Goett. Nachr.* (1865) 175.—Cuba.
myriodonta, *Muell. Arg. in Journ. Bot.* xii. (1874) 209.—Am. trop.
Myrsinites, *Baill. Adansonia*, iv. (1863–64) 305.—Bras. neo-mexicanus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 141.—N. Mexic.
nepetæfolius, *Baill. Adansonia*, iv. (1863–64) 344.—Bras.
nervosus, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 90 = micans.
nervosus, Rich. ex A. Rich. Fl. Cub. Panerog. ii. 312 = Caperonia castaneæfolia.
nervosus, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 190 = Mallotus rhamnifolius.
Neowiedii, Baill. Adansonia, iv. (1863–64) 320 = grandifolius.
Neowiedii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 100.—Bras.
nigro-viridis, *Thw. Enum. Pl. Zeyl.* 276.—Zeylan.
nitidulus, *Muell. Arg. in Flora*, xlvii. (1864) 587.—Afr. trop.
nitidus, Sw. Prod. Veg. Ind. Occ. 100 = glabellus.
nitidulus, *Baker, in Journ. Linn. Soc. xx.* (1882) 258.—Madag.
nitrariæfolius, *Baill. Adansonia*, iv. (1863–64) 351.—Bras.

CROTON :—
nivosus, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 146 = glabellus.
nivosus, Griseb. Fl. Brit. W. Ind. 40 = helicoides.
niveus, Jacq. *Enum. Pl. Carib.* 32.—Venezuela.
nobilis, *Baill. Adansonia*, iv. (1863–64) 145.—Madag.
Noronhæ, *Baill. l. c.* i. (1860–61) 163.—Madag.
Novi-Friburgi, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 97.—Bras.
nudatus, *Baill. Adansonia*, i. (1860–61) 168.—Madag.
nudus, Willd. ex Schlecht. in Linnæa, vi. (1831) 359 = reflexiflorus.
nummulariæfolius, *A. Rich. Fl. Cub. Fanerog.* ii. 211.—Cuba.
nummularius, *Baill. Adansonia*, iv. (1863–64) 360.—Bras.
nutans, *Forst. f. Prod.* 67 = Carumbium nutans.
obliquifolius, Vis. Pl. Aegypt. ac Nub. 39 = Chrozophora plicata.
obliquus, Vahl, Symb. Bot. i. 78 = Chrozophora obliqua.
oblongifolius, Buch.-Ham. in Wall. Cat. n. 7718 D = C. Joufra.
oblongifolius, Delile, Fl. Egypte, 288 = Chrozophora obliqua.
oblongifolius, *Roxb. Hort. Beng.* 69; *Fl. Ind.* iii. 685.—Ind. or.
oblongifolius, Sieber, ex Spreng. Syst. iii. 850 = Chrozophora tinctoria.
oblongifolius, Wall. Cat. n. 7736 = robustus!
oblongus, *Burm. f. Fl. Ind.* 205 (err. typ. 805).—Malaya!
occidentalis, *Muell. Arg. in Mart. Fl. Bras.* xi ii. 159.—Bras.
odontadenius, *Muell. l. c.* 227.—Bras.
odontopetalus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 79 = C. Myrsinites.
Oerstedianus, *Muell. Arg. l. c.* 105.—Am. centr.
oleoides, Muell. Arg. l. c. 82 = cneorifolius.
oliganthus, *Muell. Arg. l. c.* 106.—Bras.
olivaceus, *Muell. Arg. l. c.* 93.—Peruv.
opponens, *F. Muell. ex Benth. Fl. Austral.* vi. 125.—Austral.
oppositifolius, Geisel. Crot. Monog. 23 = Mallotus oppositifolius.
orbicularis, *Thunb. Fl. Jap.* 28.—Java.
Orbignyanus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 153.—Reg. Argent.
orophilus, Muell. Arg. l. c. 84 = celtidifolius.
organensis, *Baill. Adansonia*, iv. (1863–64) 325.—Bras.
origanifolius, *Lam. Encyc.* ii. 305.—Ind. occ.
orinocensis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 135.—Venezuela.
orthobolus, *Muell. Arg. in Flora*, lv. (1872) 9.—Am. trop.
ovalifolius, *Vahl, in West, Bidr. Ste-Croix*, 307.—Ins. St. Thom.
ovatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 12 = Mallotus tiliæfolius.
oxypetalus, *Muell. Arg. in Journ. Bot.* ii. (1864) 339.—Afr. trop.
oxyphyllus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 101.—Bras.
pallidulus, *Baill. Adansonia*, iv. (1863–64) 296.—Bras.
pallidus, *Muell. Arg. in DC. Prod.* xv. ii. 614.—Ins. S. Doming.
palmensis, Cels, Nouv. Cat. 116, nomen = lacerus.
Palmeri, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 152.—Am. bor. occ.
paludosus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 133.—Bras.
palustris, H. B. & K. Nov. Gen. et Sp. ii. 71 = Caperonia castaneæfolia.
palustris, Linn. Sp. Pl. 1004 = Caperonia palustris.
palustris, Vell. Fl. Flum. x. t. 67 = Vellozianus.
panamensis, *Muell. Arg. in DC. Prod.* xv. ii. 546.—Panama.
panduræformis, *Muell. Arg. l. c.* 671.—Cuba.

CROTON :—
paniculatus, Lam. Encyc. ii. 307 = Mallotus cochinchinensis.
paniculatus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 87.—Bras.
pannosus, *Thunb. Pl. Bras. Dec.* ii. 24.—Bras.
parænius, *Muell. Arg. in Mart Fl. Bras.* xi. ii. 116.—Bras.
pardinus, *Muell. Arg. l. c.* 217.—Bras.
parvifolius, *Muell. Arg. in Flora*, xlvii. (1864) 587.—Bras.
parvulus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 106.—Am. trop.
patulus, Lag. Gen. et Sp. Nov. 21 = Chrosophora verbascifolia.
pauciflorus, *Span. in Linnæa*, xv. (1841) 348, nomen.—Hab.!
pauciscamineus, Muell. Arg. in *Mart. Fl. Bras.* xi. ii. 188.—Bras.
paulinianus, *Muell. Arg. l. c.* 113.—Bras.
paulinus, *Muell. Arg. l. c.* 236.—Bras.
pauperulus, *Muell. Arg. in Flora*, xlvii. (1864) 485.—Chili.
Pavonæ, Bech.-Ham. in Trans. Linn. Soc. xiv. (1825) 259 = C. Tiglium.
Pavonii, Wall. Cat. n. 7722 = præc.
Pavonis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 115.—Peruv.
Payerianus, *Baill. Adansonia*, i. (1860–61) 154.—Madag.
pedicellatus, *H. B. & K. Nov. Gen. et Sp.* ii. 76.—N. Granat.
pedunculatus, *Vell. Fl. Flum.* x. t. 76.—Bras.
pedunculatus, Wall. Cat. n. 7767 = Phyllanthus longipes.
pellitus, *H. B. & K. Nov. Gen. et Sp.* ii. 78.—Peruv.
peltatus, *Thunb. Fl. Japon.* 28.—Java.
peltoideus, Ll. B. & K. Nov. Gen. et Sp. ii. 78 = ferrugineus.
peltophorus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 154.—Bolivia.
pensaccus, Baill. Adansonia, iv. (1863–64) 352 = montevidensis.
pendulus, Hassk. Pl. Jav. Rar. 266 = Paracroton pendulus.
penicillatus, Vent. Choix, 12 = ciliato-glandulosus.
pennimervis, *Scheele, in Linnæa*, xvii. (1843) 343.—Ins. Martin.
pentandrus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 97 = Clausenianus.
perafinis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 166.—Bras.
perdicipes, St. Hil. Pl. Us. Bras. t. 59 = antisiphiliticus.
Perrottetianus, *Baill. Adansonia*, i. (1860–61) 67.—Afr. trop.
Persicaria, *Baill. l. c.* iv. (1863–64) 342.—Bras.
persimilis, Muell. Arg. in Linnæa, xxxiv. (1865–66) 116 = C. Joufra.
pervenitus, *Wright, ex Griseb. in Goett. Nachr.* (1865) 173.—Cuba.
petraeus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 172.—Bras.
phebaloides, A. Cunn. ex Benth. Fl. Austral. vi. 125 = insularis.
phebalioides, *Muell. Arg. in Flora*, xlvii. (1864) 485.—Austral.
philippinensis, Lam. Encyc. ii. 306 = Mallotus philippinensis.
philomoides, Pers. Syn. ii. 585 = C. Astrolites.
phlomoides, Salzm. ex Baill. Adansonia, iv. (1863–64) 369 = Julocroton pycnophyllus.
pinahiensis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 120.—Bras.
pictus, Lodd. Bot. Cab. t. 870 = Codiæum variegatum.
pigmentarius, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 18 = Claoxylon indicum.
pilosus, *Spreng. Pugill.* ii. 87.—Ind. or.
pimeleus, *Baill. Adansonia*, iv. (1863–64) 307.—Bras.
pipocoalyx, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 108.—Bras.
plagiographis, *Muell. Arg. l. c.* 93.—Bras.
planisepalus, *Baker, in Journ. Bot.* xx. (1882) 268.—Madag.
platycladus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 128 = organensis.
plicatus, Vahl, Symb. Bot. i. 78 = Chrosophora plicata.
Pluknetii, Geisel. Crot. Monog. 40 = maritimus.

CROTON :—
Pohlianus, Muell. Arg. in Linnæa, xxxiv. (1865-66) 92 = horridulus.
Polot, *Burm. f. Fl. Ind.* 205 (err. typ. 205).—Malaya.
polyandrus, Kunth. Hort. Berog. 69 ; Fl. Ind. iii. 682 = Ballospermum axillare.
polyandrus, Spreng. Neue Entdeck. ii. 120 = brasiliensis.
polycarpus, *Benth. Pl. Hartw.* 249.—N. Granat.
polycarpus, Wall. Cat. n. 7716 = Chrozophora plicata.
polygynus, *Griseb. Cat. Monog.* 24.—Hab. ?
polygamus, Jacq. Enum. Pl. Carib. 32 = Argithamnia lanciifolia.
polystachyus, Muell. Arg. in Linnæa, xxxiv. (1865-66) 100 = nitrariæfolius.
polystachyus, Spreng. Syst. iii. 868 = matourensis.
Ponaderis, *Baill. in Adansonia*, iv. (1868-64) 311.—Brsa.
populifolius, *Mill. Gard. Dict.* ed. VIII. n. 7 ; *Lam. Encyc.* ii. 208.—Am. trop.
Pottsii, *Muell. Arg. in DC. Prod.* xv. ii. 561.—Mexic.
Pourdiæi, Muell. Arg. vide Purdiæi.
præstervisus, *Muell. Arg. in Mart. Fl. Bras.* xi. 21. 240. —Brsa.
Pringlei, *S. Wats. in Proc. Am. Acad.* xx. (1885) 373. —Am. bor. occ.
procumbens, *Vesq. Enum. Pl. Carib.* 32.—N. Granat.
procumbens, Wright, ex Griseb. in Goett. Nachr. (1865) 174 = cerinus.
prunifolius, Vahl, in Geisel. Crot. Monog. 47 = Lasiocroton prunifolius.
Pseudo-China, Schlecht. in Linnæa, iv. (1830) 84 = niveus.
Pseudo Populus, *Baill. Adansonia*, iv. (1868-64) 335. —Brsa.
pubescens, *Geisel. Crot. Monog.* 14.—Hab. ?
pulchellus, *Baill. Adansonia*, i. (1860-61) 161.—Madag.
pulcher, *Muell. Arg. in DC. Prod.* xv. ii. 644.—Mexic.
pulcherrimus, Willd. ex Schlecht. in Linnæa, vi. (1831) 361 = ciliato-glandulosus.
pulegiodorus, *Baill. Adansonia*, iv. (1868-64) 361.—Brsa.
pulegiodorus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 209.—Brsa.
pulegioides, Muell. Arg. l. c. = pulegiodorus, *Baill.*
punctatus, Anders. in Trans. Soc. Arts, xxv. (1867) 202 = corylifolius.
punctatus, Jacq. Coll. i. 166 = argyranthemus.
punctatus, *Lour. Fl. Cochinch.* 581.—Cochinch.
punctatus, Moon, Cat. Cey. Pl. 65 = Moordi.
punctatus, Retz. Obs. v. 30 = Mallotus philippinensis.
punctatus, Rich. ex Baill. Adansonia, i. (1860-61) 168 = squamigerus.
puncticulatus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 249.—Brsa.
pungens, *Vesq. Ic. Pl. Rar.* iii. 19. t. 622.—Brsa.
Purdiæi, *Muell. Arg. in Flora*, xlvii. (1864) 488.—(Fourdiæi) N. Granat.
pycnadenius, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 216.—Brsa.
pycnanthus, *Benth. Pl. Hartw.* 247.—Peruv.
pycnocarpus, Muell. Arg. in Linnæa, xxxiv. (1865-66) 130 = grandivelum.
pycnocephalus, *Baill. Adansonia*, iv. (1868-64) 356.—Brsa.
pycnophyllus, Salzm. ex Schlecht. in Linnæa, xix. (1847) 348 = Julocroton pycnophyllum.
pycnotrichus, Muell. Arg. l. c. xxxiv. (1865-66) 122 = Vautherianus.
pyrifolius, *Muell. Arg. in Journ. Bot.* ii. (1864) 388.—Afr. trop.
quadripartitus, Labill. Nov. Holl. Pl. ii. 78. t. 225 = Adriana quadripartita.
quadriestosus, *Lam. Encyc.* ii. 208.—Peruv.
quitensis. Sprung. Syst. iii. 872 = alnifolius.
racemosus, *Burm. f. Fl. Ind.* 206 (err. typ. 206).—Java.
ramiflorus, *V. Grah. Cat. Pl. Bomb.* 182.—Ind. or.
reflexifolius, *H. B. & K. Nov. Gen. et Sp.* ii. 68.—Mexic.
refractus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 189.—Brsa.
Regelianus, *Muell. Arg. l. c.* 188.—Brsa.
Regnellianus, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 122.—Brsa.
repandus, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1802) 306 = Mallotus repandus.

CROTON :—
repens, *Schlecht. in Linnæa*, xix. (1847) 237.—Mexic.
reticulatus, *Heyne, in Wall. Cat.* n. 7784.—Ind. or.
reticulatus, *Thunb. Fl. Japan.* 93.—Java.
reticulatus, Willd. Sp. Pl. iv. 545 = Mallotus rhamnifolius.
rhamnifolius, *H. B. & K. Nov. Gen. et Sp.* ii. 75.—Am. trop.
rhamnifolius, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1802) 190 = Mallotus rhamnifolius.
Rheedei, *V. Grah. Cat. Pl. Bomb.* 182.—Ind. or.
rhenanifolius, *Baill. Adansonia*, iv. (1868-64) 313.—Brsa.
rhodocalychus, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 108.—Ind. or.
rhombifolius, Willd. Sp. Pl. iv. 555 = Mallotus repandus.
Richardi, Willd. l. c. 558 = flavens.
ricinocarpus, Linn. Sp. Pl. ed. II. 1427 = Leidesia Sonderiana.
ricinoides, Pers. Syn. ii. 586 = Mallotus ricinoides.
Riedelianus, Muell. Arg. in Linnæa, xxxiv. (1865-66) 104 = buxifolius.
riparius, H. B. & K. Nov. Gen. et Sp. ii. 90 = lobatus.
rivinæfolius, *H. B. & K. l. c.* 80.—N. Granat.
rivularis, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 112.—Afr. austr.
robustus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 242.—Burma.
rosmarinifolius, A. Cunn. ex Benth. Fl. Austral. vi. 78 = Ricinocarpus rosmarinifolius.
rosmarinifolius, A. Cunn. in Field, N. S. Wales, 356 = Bertya rosmarinifolia.
rosmarinifolius, Drumm. ex Muell. Arg. in DC. Prod. xv. ii. 208 = Hippocrepandra ericoides.
rosmarinifolius, *Griseb. in Goett. Nachr.* (1865) 174.—Cuba.
rosmarinifolius, Salisb. Prod. 389 = linearis.
rottleræfolius, *Baill. Adansonia*, iv. (1868-64) 337.—Brsa.
Rottleri, Geisel. Crot. Monog. 54 = Chrozophora plicata.
Rudolphianus, *Muell. Arg. in DC. Prod.* xv. ii. 654.—Brsa.
rufescens, Muell. Arg. in Linnæa, xxxiv. (1865-66) 118 = cordiacifolius.
rufidulus; Baill. Adansonia, iv. (1868-64) 347 = sclerocalyx.
rufo-virgatus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 195.—Brsa.
Ruizianus, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 89.—Peruv.
Sagraeanus, *Muell. Arg. in DC. Prod.* xv. ii. 617.—Cuba.
salicifolius, Geisel. Crot. Monog. 6 = Homonoia riparia.
salinus, *Griseb. in Goett. Abh.* xxiv. (1879) 54.—Reg. Argent.
salutaris, *Casar. Nov. Stirp. Bras. Dec.* 89.—Brsa.
salviæfolius, H. B. & K. Nov. Gen. et Sp. ii. 74 = rhamnifolius.
salvifolius, Mill. Gard. Dict. ed. VIII. n. 10 = humilis.
Sampaioanus, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 94.—Peruv.
sanguifluus, H. B. & K. Nov. Gen. et Sp. ii. 89 = gossypifolius.
Sanguis-draconis, Mart. ex Baill. in Adansonia, iv. (1868-64) 327 = celtidifolius.
sanctalinus, *Baill. Adansonia*, iv. (1868-64) 359.—Am. austr.
sapifolius, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 137.—Brsa.
sarcocarpus, *Baill. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 54.—Ins. Society.
sarcopetalus, *Muell. Arg. in Journ. Bot.* xii. (1874) 201.—Am. trop.
saxatilis, Muell. Arg. in Linnæa, xxxiv. (1865-66) 119 = corchoropsis.
scaber, *Willd. Sp. Pl.* iv. 550.—Peruv.
scaberrimus, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 126.—Cuba.
scabiosus, *Bedd. Fl. Sylv.* ii. t. 268.—Ind. or.
scandens, Sieber, ex Presl, Bot. Bemerk. 109 = Fragia volubilis.
scandens, *Vell. Fl. Flum.* x. t. 72 = Plukenetia tamnoides.
Schlechtendalianus, *Schlecht. in Linnæa*, xix. (1847) 248.—Mexic.
Schimperianus, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 83.—Abyss.

CROTON :—
Schultzeii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 193.—Brsa.
Schultzii, *Benth. Fl. Austral.* vi. 194.—Austral.
sclerocalyx, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 134.—Am. trop.
scoriaceus, Lam. Encyc. ii. 215 = glandulosus.
Scouleri, *Hook. f. in Trans. Linn. Soc.* xx. (1857) 188. —Ins. Galapag.
sebiferus, Linn. Sp. Pl. 1004 = Sapium sebiferum.
Sellowii, *Baill. Adansonia*, iv. (1868-64) 304.—Brsa.
seminudus, *Muell. Arg. in DC. Prod.* xv. ii. 629.—Brsa.
sensitivitus, *Muell. Arg. l. c.* 628.—Brsa.
senegalensis, Lam. Encyc. ii. 212 = Chrozophora senegalensis.
sericeus, Lam. l. c. 210 = matourensis.
sericeus, Schlecht. in Linnæa, v. (1830) 85 = fragilis.
serpyllifolius, *Baill. Adansonia*, iv. (1868-64) 355.—Brsa.
serpyllifolius, Muell. Arg. in DC. Prod. xv. ii. 595 = seq.
serpyloides, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 159 = nummulariæfolius.
serratifolius, *Baill. Adansonia*, iv. (1868-64) 312.—Brsa.
serratus, Hochst. ex Baill. l. c. i. (1860-61) 66 = Caperonia serrata.
serratus, *Muell. Arg. in DC. Prod.* xv. ii. 647.—Brsa.
sessiliflorus, Sw. Prod. Veg. Ind. Occ. 100 = Savia sessiliflora.
setigerus, Hook. Fl. Bor. Am. ii. 141 = Eremocarpus setigerus.
shepherdioides, Schau. in Linnæa, xx. (1847) 729 = hypoleucus.
sidæfolius, *Lam. Encyc.* ii. 204.—Ins. S. Doming.
siderophyllus, *Baill. Adansonia*, iv. (1868-64) 305.—Brsa.
similis, Muell. Arg. in Linnæa, xxxiv. (1865-66) 134 = Landianus.
sincorensis, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* xi. ii. 56.—Brsa.
Sirahi, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 86 = Excœcaria japonica.
Sloanei, Benn. in Journ. Linn. Soc. iv. (1860) 80 = glabellus.
socotranus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 96.—Ins. Socotr.
solanifolius, Geisel. Crot. Monog. 74 = Ballospermum axillare.
Soliman, *Cham. & Schlecht. in Linnæa*, vi. (1831) 361.—Mexic.
Sonderianus, *Muell. Arg. in DC. Prod.* xv. ii. 557.—Brsa.
Sonoræ, *Torr. Bot. Mex. Bound.* 194.—N. Mexic.
sorastensis, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 116.—Bolivia.
sordidus, *Benth. Pl. Hartw.* 247.—N. Granat.
speciosus, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 83.—N. Granat.
sphærocarpus, H. B. & K. Nov. Gen. et Sp. ii. 84. t. 106 = mortifolius.
sphærogynus, *Baill. Adansonia*, iv. (1868-64) 296.—Brsa.
Spica, *Baill. l. c.* 362.—Brsa.
spicatus, Berg. in Phil. Trans. lviii. (1768) 122. t. 7 = incidus.
spicatus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = C. Spica.
spiciflorus, *Thunb. Fl. Japan.* 93.—Java.
spinosus, Forsk. Fl. Aegypt. Arab. 163 = Jatropha spinosa.
spinosus, Linn. Sp. Pl. 1005 = Ricinus communis.
spinosus, *Wright, in Sauv. Fl. Cub.* 197.—Cuba.
spinulis, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 125.—Cuba.
splendidus, Baill. Adansonia, iv. (1868-64) 294 = subgratus.
Spruceanus, *Benth. in Hook. Kew Journ.* vi. (1854) 275.—Brsa.
squamigerus, *Baill. Adansonia*, i. (1860-61) 168.—Madag.
squamulosus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 244.—Brsa.
squarrulosus, Vahl, in Geisel. Crot. Monog. 28 = globellus.
stanleianus, *Muell. Arg. in Linnæa*, xxxiv. (1865-66) 88.—Brsa.

CROTON :—
stenandrus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 84 = velinarsitus.
stenophyllus, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 158.—Cuba.
stenosepalus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 114.—Peruv.
stenotrichus, *Muell. Arg. l. c.* 105.—Bras.
stigmatosus, F. Muell. Fragm. iv. 140 = phebaliotes.
stipulaceus, *H. B. & K. Nov. Gen. et Sp.* ii. 85.—Mexic.
stipulatus, *Veil. Fl. Flum.* x. t. 77.—Bras.
Stoechadis, *Baill. Adansonia*, iv. (1863–64) 314.—Bras.
Storckii, Seem. Viti, 441 = Verreauxii.
strigosus, *Spreng. Neue Entdeck.* iii. 23.—Ins. S. Doming.
styloura, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 128.—Mexic.
suaveolens, *Torr. Bot. Mex. Bound.* 194.—N. Mexic.
suavis, *H. B. & K. Nov. Gen. et Sp.* ii. 76.—N. Granat.
subacutus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 167.—Bras.
subargyrius, *Muell. Arg. l. c.* 174.—Bras.
subcapitatus, Muell. Arg. in Buek, Ind. ad DC. Prod. iv. p. v = humilis.
subcommosus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 284.—Bras.
subcompressus, *Muell. Arg. in DC. Prod.* xv. ii. 557.—Am. austr.
suberosus, *H. B. & K. Nov. Gen. et Sp.* ii. 86.—Mexic.
subferrugineus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 135.—Bras.
subfragilis, *Muell. Arg. l. c.* 111.—Mexic.
subinconus, *Muell. Arg. l. c.* 139.—Guiana.
sublepidotus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 260.—Bras.
subluculdus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 126.—Ins. S. Doming.
subluteus, Lam. Encyc. ii. 210 = guianensis.
sublyratus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 243.—Ins. Andam.
subpannosus, *Griseb. in Goett. Abh.* xix. (1874) 96.—Keg. Argent.
subtomentosus, Linn. Syst. ed. X. 1376 = humilis.
subvillosus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 168.—Bras.
succinber, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 55.—Keg. Argent.
sulcifructus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 94.—Ins. Socotr.
sumatranus, *Mig. Fl. Ind. Bat.* i. ii. 581.—Sumatra.
surinamensis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 82.—Guiana.
sylvaticus, *Hochst. in Flora*, xxviii. (1845) 82.—Afr. austr.
sylvesticus, Schlecht. in Linnæa, xix. (1847) 240 = reflexifolius.
sylvestris, Poepp. ex Griseb. in Goett. Nachr. (1865) 174 = Ricinella sylvestris.
syringæfolius, H. B. & K. Nov. Gen. et Sp. ii. 67 = nivens.
tabacifolius, *Griseb. Crot. Monog.* 26.—Ind. or.
Tamberlikii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 265.—Bras.
tamposensis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 114.—Peruv.
Tartonraira, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 129.—Bras.
tejucensis, *Muell. Arg. l. c.* 176.—Bras.
tenellus, *Muell. Arg. l. c.* 248.—Bras.
tenuis, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 297.—Calif.
tenuissimus, *Baill. Adansonia*, iv. (1863–64) 295.—Bras.
terminalis, *Veil. Fl. Flum.* x. t. 78.—Bras.
tetradenius, *Baill. Adansonia*, iv. (1863–64) 343.—Bras.
Teucridium, *Baill. l. c.* 349.—Bras.
texensis, *Muell. Arg. in DC. Prod.* xv. ii. 692.—Texas.
thebaicolus, *Hort. Sydn. ex Tréfem. & Binn. Cat. Hort. Bog.* 293. nomen.—Hab.?
thermarum, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 151.—Bras.
Thouarsianus, *Baill. Adansonia*, i. (1860–61) 167.—Madag.

CROTON :—
therifer, *H. B. & K. Nov. Gen. et Sp.* ii. 76.—Bras.
Thouinianus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 116 = Klotzschianus.
thymelinus, *Baill. Adansonia*, iv. (1863–64) 352.—Bras.
tiglium, Blume, Bijdr. 602 = laevifolius.
Tiglium, *Linn. Sp. Pl.* 1004.—Ind. or.; Malaya.
tillæfolius, *Lam. Encyc.* ii. 206.—Ins. Maurit.
tiliæfolius, Pers. Syn. ii. 545 = populifolius.
tiliæfolius, Thon. ex Baill. Adansonia, i. (1860–61) 160 = fothergillæfolius.
Timandra, Baill. l. c. iv. (1863–64) 351 = serratus.
timandroides, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 132.—Bras.
tinctorius, Burm. f. Fl. Ind. 304. t. 62 = Chrozophora plicata.
tinctorius, Linn. Sp. Pl. 1004 = Chrozophora tinctoria.
tomentellus, *F. Muell. Fragm.* iv. 141.—Austral.
tomentosus, Hort. Par. ex Link, Enum. Hort. Berol. ii. 406 = dioicus.
tomentosus, Link, l. c. = flavens.
tomentosus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 107 = crassifolius.
tomentosus, Pav. ex Baill. Adansonia, i. (1860–61) 347 = Baillonianus.
Torreyanus, *Muell. Arg. in DC. Prod.* xv. ii. 579.—N. Mexic.
tragifolius, Baill. Adansonia, vi. (1865) 16 = glechomæfolius.
triacros, *F. Muell. Fragm.* vi. 185.—Austral.
triangularis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 128.—Bras.
trichocarpus, Torr. Bot. Mex. Bound. 196 = Cortesianus.
trichocephalus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 102 = hypocrepchalus.
trichophorus, Muell. Arg. l. c. 132 = vestitus.
trichotomus, *Griseb. Crot. Monog.* 50.—Madag.
tricolor, Baill. Adansonia, iv. (1863–64) 291 = argyrophylloides.
tricolor, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 96.—Bras.
tricuspidatus, Lam. Encyc. ii. 212 = Argithamnia tricuspidata.
tridentatus, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* xi. ii. 219.—Bras.
Tridesma, Lcand. ex Baill. Adansonia, iv. (1863–64) 367 = Julocroton fuscescens.
triglandulatus, *Veil. Fl. Flum.* x. t. 79.—Bras.
trigonocarpus, *Griseb. in Goett. Nachr.* (1865) 173.—Cuba.
trilobatus, *Forsk. Fl. Aegypt. Arab.* 163.—Arab.
trilobatus, Willd. Sp. Pl. iv. 566 = lobatus.
triqueter, Lam. Encyc. ii. 215 = Julocroton triqueter.
tristis, Muell. Arg. in Linnæa, xxxiv. (1865–66) 140 = pulegioides.
tuberculatus, Bunge, in Mém. Sav. Etr. Pétersb ii. (1835) 134 = Argithamnia tuberculata.
tucumanensis, *Griseb. in Goett. Abh.* xix. (1874) 95.—Keg. Argent.
Tulasnei, *Baill. Adansonia*, i. (1860–61) 156.—Madag.
umbellatus, Dalz. & Gibs. Bomb. Fl. 231 = Blachia denudata.
umbellatus, Heyne, in Wall. Cat. n. 7768 = Blachia denudata.
umbellatus, Wight, Ic. t. 1874 = Blachia calycina.
umbellatus, Willd. Sp. Pl. iv. 545 = Blachia umbellata.
umbratilis, *H. B. & K. Nov. Gen. et Sp.* ii. 82.—Venezuela.
umbrinus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 86 = heterocalyx.
undulatus, *André. in Illustr. Hortic.* xix. (1872) 265.—Ins. Salomon.
urceolatus, Baill. Adansonia, iv. (1863–64) 299 = salutaris.
urens, Linn. Sp. Pl. 1005 = Tragia involucrata.
urophylla, *Wall. ex Voigt, Hort. Suburb. Calc.* 157, nomen.—Ind. or.
urticæfolius, *Lam. Encyc.* ii. 219.—Bras.
urticoides, A. Cunn. ex Stend. Nom. ed. II. i. 447 = Adriana glabrata.
uruguanus, *Baill. Adansonia*, iv. (1863–64) 335.—Bras.
uruguayensis, *Baill. l. c.* 293.—Uruguay.
vaccinioides, *A. Rich. Fl. Cub. Fanerog.* ii. 211.—Cuba.

CROTON :—
Vahlii, *Griseb. Crot. Monog.* 41.—Hab.?
Vaillanti, *Geisel. l. c.* 10.—Ind. occ.
variegatus, *Blanco, Fl. Filip.* ed. I. 751.—Ins. Philipp
variegatus, Forsk. Fl. Aegypt. Arab. 163 = Jatropha variegata.
variegatus, Linn. Sp. Pl. 1199 = Codiaeum variegatum.
Vauthieri, Muell. Arg. in Linnæa, xxxiv. (1865–66) 93 = macrobothrys.
Vauthieranus, *Baill. Adansonia*, iv. (1863–64) 317.—Bras.
Veitchianus, *André, Illustr. Hortic.* xix. (1872) 185.—Ins. Pacif.
Velame, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 181.—Bras.
vellerifluus, Baill. Adansonia, iv. (1863–64) 353 = nitratæfolius.
Vellozianus, *Muell. Arg. in DC. Prod.* xv. ii. 553.—Bras.
velutinus, *Baill. Adansonia*, iv. (1863–64) 325.—Bras.
venosus, Geisel. Crot. Monog. 42 = Leptonema venosum.
venosus, Spreng. Neue Entdeck. iii. 24 = C. Astroites.
vepretorum, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 172.—Bras.
verbascifolius, Willd. Sp. Pl. iv. 539 = Chrozophora verbascifolia.
verbenæfolius, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 140.—Bras.
Verreauxii, *Baill. Etud. Gén. Euph.* 357.—Austral.
veracolor, *Muell. Arg. in DC. Prod.* xv. ii. 690.—Bras.
vestitus, *Spreng. Syst.* iii. 875.—Bras.
villosus, Forsk. Fl. Aegypt. Arab. 163 = Jatropha villosa.
villosus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 169.—Bras.
villosus, Sibth. & Sm. Fl. Graec. Prod. ii. 249; Fl. Graec. t. 951 = Chrozophora verbascifolia.
vimmalis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 158.—Cuba.
virens, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 142.—N. Mexic.
virgultosus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 104.—Bras.
Virletianus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 138.—Mexic.
viscosus, Labill. Nov. Holl. Pl. ii. 72. t. 222 = Beyeria viscosa.
volubilis, Lianos, in Mem. Acad. Cienc. Madr. iv. (1856) 508 = Mallotus repandus.
vulnerarius, *Baill. Adansonia*, iv. (1863–64) 328.—Bras.
Wagneri, *Muell. Arg. in Flora*, lv. (1872) 5.—Ecuador.
Wallichii, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 118.—Burma.
Warmingii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 105.—Bras.
Welwitschianus, *Muell. Arg. in Journ. Bot.* ii. (1864) 338.—Afr. trop.
Widgrenianus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 104.—Bras.
Wilsonii, *Griseb. Fl. Brit. W. Ind.* 40.—Cuba.
Wrightii, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 160 = Lasiocroton Wrightii.
Wullschlaegelianus, *Muell. Arg. in Flora*, lv. (1872) 10.—Ind. occ.
xalapensis, Hook. f. in Trans. Linn. Soc. xx. (1847) 188 = Scouleri.
xalapensis, *H. B. & K. Nov. Gen. et Sp.* ii. 85.—Mexic.
yparamensis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 166.—Bras.
yunguensis, *Griseb. in Goett. Nachr.* (1865) 171.—Cuba.
zambesicus, *Muell. Arg. in Flora*, xlvii. (1864) 483.—Afr. trop.
zeylanicus, Muell. Arg. in Linnæa, xxxiv. (1865–66) 107 = reticulatus.
Zollingeri, Mlq. Fl. Ind. Bat. i. ii. 581 = argyratus.

CROTONANTHUS, Kloisch, ex Schlecht. in Linnæa, xxvi. (1853–55) 634 = Croton, Linn. (Euphorb.).
podifolius, Klotzsch, l. c. = Croton rhamnifolius.
pradifolius, Muell. Arg. in DC. Prod. xv. ii. 685, ephalon. = præc.

CROTONOGYNE, Muell. Arg. in Flora, xlvii. (1864) 583. *EUPHORBIACEÆ*, Benth. & Hook. f. ii. 396.
Maniana, *Muell. Arg. l. c.*—Afr. trop.

CROTONOPSIS, Michx. Fl. Bor. Am. ii. 185. t. 46 (1803). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 390.
FRIESIA, Spreng. Anleit. ii. 11. 885 (1818).
LEPTEMON, Rafin. in Med. Repos. N. York, v. (1808) 359.
abnormis, Baill. Etud. Gén. Euph. 381 = linearis.
argentea, Pursh, Fl. Am. Sept. i. 206 = linearis.
elliptica, Willd. Sp. Pl. iv. 880 = linearis.
linearis, *Michx. Fl. Bor. Am.* ii. 185. t. 46.—Am. bor.

CROTULARIA, Medic. in Vorles. Churpf. Phys. Gen. ii. (1787) 344 = **Crotalaria**, Linn. (Legum.).

CROTULARIUS, Medic. in Vorles. Churpf. Phys Gen. ii. (1787) 397 = praec.

CROUM, Gled. ' Berl. v. (1749) 127,' sphalm. ex Pfeiff. Nom. i. 1958 = **Urvum**, Tourn. (Legum.).

CROWEIA, Sm. in Trans. Linn. Soc. iv. 222 (1798). *RUTACEAE*, Benth. & Hook. f. i. 298.
angustifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 13.—Austral.
dentata, *R. Sy. ex Benth. Fl. Austral.* i. 380.—Austral.
exalata, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 11.—Austral.
latifolia, Lodd. in G. Don, Gen. Syst. i. 792 = saligna.
saligna, *Andr. Bot. Rep.* t. 79.—Austral.
scabra, R. Grah. in Edinb. Phil. Journ. (1827) 174 = Eriostemon salicifolius.

CROZOPHORA, A. Juss. Tent. Euphorb. 27 (1824) = **Chrosophora**, Neck. (Euphorb.).

CROZOPHYLA, Rafin. Sylva Tellur. 64 (1838) = **Codiaeum**, A. Juss. (Euphorb.).
angustifolia, Rafin. l. c.—Hab. ?
elliptica, Rafin. l. c.—Hab. ?
picta, Rafin. l. c. = Codiaeum variegatum.
variegata, Rafin. l. c. = Codiaeum variegatum.

CRUCIANELLA, Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth. & Hook. f. ii. 150.
LAEMANNIA, S. G. Gmel. ex Trin. in Mém. Acad. Pétersb. vi. (1818) 492. t. 11.
RUBIOLA, Tourn. ex Adans. Fam. ii. 144 (1763).
aegyptiaca, *Linn. Mant.* i. 38.—Aegypt.
americana, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
angustifolia, Koch, ex Nym. Consp. 385 = latifolia.
angustifolia, *Linn. Sp. Pl.* 109.—Europ.; As. Min.
anomala, Balb. ex DC. Prod. iv. 587 = mollugnioides.
aspera, Bieb. Fl. Taur. Cauc. i. 107 = Asperula aspera.
bithynica, Boiss. Diagn. Ser. I. x. 58 = graeca.
bracteata, Boiss. l. c. Ser. I. iii. 29 = Asperula glomerata.
capitata, Labill. Pl. Syr. Dec. i. 12. t. 8 = Asperula glomerata.
chlorantha, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 8 = sec.
chlorostachys, *Fisch. & Mey. l. c.* 25.—As. Min.; Persia.
ciliata, *Lam. Encyc.* ii. 217.—Syria; Arab.
confinis, Jord. ex Nym. Consp. 385 = angustifolia.
Decaisnei, Stend. Nom. ed. II. i. 448 = ciliata.
decumbens, Adam, ex Ledeb. Fl. Ross. ii. 404 = Asperula aspera.
diffusa, Roth, in Usteri, Ann. Bot. x. (1794) 40 = ciliata.
distichа, *Boiss. Diagn.* Ser. I. ii. 35.—As. Min.
exasperata, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 8.—As. Min.; Persia.
fibrosa, Ehrenb. ex Boiss. Fl. Orient. iii. 29 = Asperula glomerata.
filifolia, *Regel & Schmalh. Fl. Turkest.* iii. 18.—Turkestan.
fruticosa, Boiss. Fl. Orient. iii. 20 = macrostachya.
galioides, Boiss. Diagn. Ser. I. iii. 29 = Asperula aspera.
glareica, Trin. in Mém. Acad. Pétersb. vi. (1818) 493 = glauca.
glauca, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 181.—Persia.
glomerata, Bieb. Fl. Taur. Cauc. i. 107 = Asperula glomerata.
graeca, *Boiss. Diagn.* Ser. I. iii. 28.—Graecia.
herbacea, *Forsk. Fl. Aegypt. Arab.* 30.—Aegypt.
hirta, *Pomel, Nouv. Mat. Fl. Atl.* 74.—Afr. bor.

CRUCIANELLA:—
hispida, Mill. Gard. Dict. ed. VIII. n. 4 = Crusea rubra.
hispidula, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 369 = ciliata.
hispidula, Flach. & Mey. Ind. Sem. Hort. Petrop. i. 25 = exasperata.
imbricata, *Boiss. Diagn.* Ser. I. x. 59.—As. Min.
inaignis, *Vatke, in Linnaea*, xxxviii. (1874) 714.—Kurdistan.
Katschyana, Boiss. & Hohen. Diagn. Ser. I. iii. 28 = Asperula Kotschyana.
latifolia, *Linn. Sp. Pl.* 109.—Europ.; As. Min. macrostachys, *Boiss. Diagn.* Ser. I. iii. 27.—As. Min.
maritima, *Linn. Sp. Pl.* 109.—Reg. Mediterr. membranacea, *Boiss. Diagn.* Ser. I. iii. 26 = macrostachya.
mollugineoides, Bieb. Cent. ii. t. 56 = Asperula mollugineoides.
monspeliaca, Linn. Sp. Pl. 109 = latifolia.
monspeliaca, Sibth. & Sm. Fl. Graec. ix. 31. t. 140 = graeca.
monspeliensis, Hill, Veg. Syst. vii. 33 = latifolia.
mucronata, Roth, in Usteri, Ann. Bot. x. (1794) 41 = angustifolia.
orientalis, Willd. ex Schult. Mant. iii. 215 = glauca.
oxyloba, Janka, ex Nym. Consp. 385 = angustifolia.
patula, *Linn. Amoen. Acad.* iii. 401.—Hispan.
persicifolia, Boiss. Diagn. Ser. I. iii. 26 = macrostachya.
pentandra, Dufour, ex Roem. & Schult. Syst. iii. 2, 215 = patula.
pentandra, Willd. ex Ledeb. Fl. Ross. ii. 404 = mollugnioides.
prostrata, Adams, in Weber & Mohr, Beitr. i. 46 = Asperula aspera.
pubescens, Willd. Sp. Pl. i. 602 = Asperula incana.
purpurea, [Honck.] Vers. Gew. Teutschl. 481 = Gallium purpureum.
purpurea, Wulff. ex Steud. Nom. ed. II. i. 448 = Galium Aparine.
rigida, Tausch, in Flora, xviii. (1835) 359 = Asperula rigida.
rupestris, Guss Fl. Sic. Prod. Suppl. 44 = maritima.
sherardioides, Boiss. Diagn. Ser. I. iii. 24 = Asperula sherardioides.
spicata, Lam. Fl. Fr. iii. 372 = angustifolia, latifolia.
stylosa, Trin. in Mém. Acad. Pétersb. vi. (1818) 485 = Asperula ciliata.
suaveolens, *C. A. Mey. Verz. Pfl. Cauc.* 82.—Persia.
subfruticosa, Willd. ex Ledeb. Fl. Ross. ii. 404 = Asperula glomerata.
suffulta, *Woods, Tour. Fl.* 168.—Gallia.
syriaca, Boiss. Diagn. Ser. I. x. 58 = macrostachya.
Tournefortii, Tausch, in Flora, xviii. (1835) 359 = Asperula Tournefortii.
tuberculosa, Cav. Desc. 349 = ciliata.
vulgaris, Gater. ex Steud. Nom. ed. II. i. 448 = latifolia.

CRUCIATA, Gilib. Fl. Lituan. i. 36 (1781) = **Gentiana**, Tourn.
verticillata, Gilib. l. c. = G. Cruciata.

CRUCIATA, Tourn. ex Adans. Fam. ii. 144 (1763) = **Gallum**, Linn. (Rubiac.).
ciliata, Opiz, Seznam, 84 = G. asperuloides.
hirsuta, Tourr. in Ann. Soc. Linn. Lyon, N.S. xvi. (1868) 396 = G. Cruciata.
laevipes, Opiz, Seznam, 84 = G. Cruciata.

CRUCIELLA, Leschen. ex DC. Prod. ii. 74 (1830) = **Xanthosia**, Rudge (Umbellif.).
candida, Leschen. l. c. 75 = X. rotundifolia.

CRUCITA, Linn. Sp. Pl. ed. II. 179 (1762) = **Ixosine**, Linn. (Amarant.).
americana, Lam. Encyc. ii. 218 (Crucita) = I. celosioides?
hispanica, Linn. Sp. Pl. ed. II. 179 = *americana*.
hispanica americana, Roem. & Schult. Syst. iv. 484 = *americana*.

CRUCIUNDULA, Rafin. Fl. Tellur. ii. 100 (1836) = **Pocosana**, Moench (Crucif.).
minima, Rafin. l. c. = N. stylosa.

CRUCKSHANKSIA, Hook. Bot. Misc. ii. 211. t. 90 (1831) = **Balbisia**, Cav. (Geraniac.).
ristifiora, Hook. l. c. = B. verticillata.
peduncularis, Steud. Nom. ed. II. i. 448 = B. verticillata.

CRUCKSHANKSIA, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 361. *RUBIACEAE*, Benth. & Hook f. ii. 97.
ROTHERIA, Meyen, Reise, i. 402 (1834).
Bustillosii, *Phil. in Linnaea*, xxvii. (1856) 696.—Chili.
capitata, *Phil. in Anal. Univ. Chil.* (1872) 731.—Chili.
chrysantha, *Phil. l. c.* 730.—Chili.
densifolia, *Phil. l. c.*—Chili.
glacialis, Poepp. & Endl. Nov. Gen. et Sp. iii. 31. t. 236 = Oreopolus citrinus.
hymenodon, *Hook. f. Arn. in Hook. Bot. Misc.* iii. (1833) 361.—Chili.
macrantha, *Phil. in Linnaea*, xxxiii. (1864–65) 97.—Chili.
montana, *Clos, in C. Gay, Fl. Chil.* iii. 195. t. 33. f. 1.—Chili.
Montitana, *Clos, l. c.*—Chili.
Palma, *Clos, l. c.* 194.—Chili.
pumila, *Clos, l. c.* 196. t. 33. f. 2.—Chili.
tripartita, *Phil. Fl. Atac.* 26.—Chili.

CRUCKSHANKSIA, Miers, Trav. ii. 529 (1826) = **Solenomelus**, Miers (Irid.).
graminea, Miers, l. c. = S. chilensis.

CRUDIA, Schreb. Gen. 282 (1789). *LEGUMINOSAE* Benth. & Hook. f. i. 584.
APALATOA, Aubl. Pl. Gui. 382 (1775) (& t. 147, OPALATOA).
CYCLAS, Schreb. Gen. 282 (1789).
PEYONA, Miq. Fl. Ind. Bat. i. 1. 1081 (1855).
TOUCHIROA, Aubl. Pl. Gui. 384. t. 148 (1775).
WALDSCHMIDTIA, Scop. Introd. 100 (1777).
acuminata, *Benth. Bot. Voy. Sulph.* 89.—Am. centr.
amazonica, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 288.—Bras.
Apalatoa, *Steud. Nom.* ed. II. i. 110.—Guiana.
aromatica, *Willd. Sp. Pl.* ii. 540.—Guiana.
Blanconi, *Kolfr, in Journ. Linn. Soc.* xxi. (1884) 309.—Ins. Philipp.
bractesta, *Benth. in Hook. Journ. Bot.* ii. (1840) 101.—Guinea.
falcata, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1210.—Guiana.
hirtelloides, *Miq. ex Benth. in Mart. Fl. Bras.* xv. ii. 288.—Guiana.
obliqua, *Griseb. Fl. Brit. W. Ind.* 216.—Cuba.
oblonga, *Benth. Bot. Voy. Sulph.* 89.—Ind. occ.
oblonga, Griseb. Fl. Brit. W. Ind. 216 = oblonga.
orientalis, *Hassk. Cat. Fl. Hort. Bog. Alt.* 288.—Java.
Parivoa, *DC. Prod.* ii. 630.—Guiana.
pubescens, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 240.—Bras.
Schreberi, *Y. F. Gmel. Syst.* 674.—Guiana.
senegalensis, *Planch. ex Benth. in Trans. Linn. Soc* xxv. (1866) 314.—Afr. trop.
spicata, Blanco, Fl. Filip. ed. II. 961 = Blancol.
spicata, Willd. Sp. Pl. ii. 539 = Schreberi.
Vouarana, DC. Prod. ii. 630 = Cupania Vouarana.
zeylanica, *Benth. in Trans. Linn. Soc.* xxv. (1866) 314.—Zeylan.

CRUDYA, Batsch, Tab. 98 (1806); DC. Prod. ii. 519 (1825) = praec.

CRUIKSHANKSIA, Benth. & Hook. f. Gen. i. 276 (1862) = **Balbisia**, Cav. (Geran.).

CRUIKSHANKSIA, Reichb. Consp. 212* (1828) = Solenomelus, Miers (Irid.).

CRUMENARIA, Mart. Nov. Gen. et Sp. ii. 68. t. 160 (1826). *RHAMNEAE*, Benth. & Hook. f. i. 396.
choretroides, *Mart. ex Reiss. in Mart. Fl. Bras.* xi. i. 114.—Bras.
decumbens, *Mart. Nov. Gen. et Sp.* ii. 69. t. 160.—Bras.
erecta, *Reiss. in Mart. Fl. Bras.* xi. i. 114.—Bras.
polygaloides, *Reiss. l. c.* 113.—Bras.

CRUMINIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 429 = **Centrosema**, DC. (Legumin.).
giganteum, Desv. l. c. = Cent. Plumieri.

CRUPINA, Dill. ex Linn. Syst. ed. I (1735); Cass. in Dict. Sc. Nat. xii. 67 (1818). *COMPOSITAE*, Benth. & Hook. f. ii. 476.
alpestris, Arvet.—Cf. Bull. Soc. Bot. Fr. xxvi. (1879) Bibl. 89 = vulgaris.
brachypappa, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 84 = vulgaris.
Crupinastrum, *Vis. Fl. Dalm.* ii. 42.—Reg. Mediterr.; Orient.
intermedia, Walp. Ann. i. 447 = Centaurea intermedia, *Mutel*.
Morisii, Boreau, Fl. Centr. Fr. ed. II. 292 = C. Crupinastrum.
pauciflora, *Hoffmgg. & Link, Fl. Port.* ii. 241.—Lusitan.
pauciflora, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 392 = vulgaris.
pseudo-crupina, Walp. Ann. i. 447 = Centaurea pseudo-crupina.
vulgaris, *Cass. Dict. Sc. Nat.* xii. 68.—Reg. Mediterr.; As. occ.
Zuccarinii, Bunge, ex Nym. Consp. 488 = C. Crupinastrum.

CRUPINASTRUM, Schur, in Ber. Bot. Rundsch. (1853) 37; ex oj. Enum. Pl. Transs. 411 = **Serratula**, Linn. (Compos.).
Centaureum, Schur, l. c. = S. coronata.

CRUSEA, Cham. ex DC. Prod. iv. 571 (1830) = **Mitracarpum**, Zucc. (Rubiac.).

CRUSEA, Cham. & Schlecht. in Linnaea, v. (1830) 165. *RUBIACEAE*, Benth. & Hook. f. ii. 163.
acuminata, E. Mey. Zwei Pfl. Docum. 176 = Hedyotis natalensis.
allococca, *A. Gray, in Proc. Am. Acad.* xix. (1883) 78. —Texas.
aspera, *Bartl. in Linnaea*, xiv. (1840) *Litt.* 124.—Mexic.
brachyphylla, *Cham. & Schlecht. in Linnaea*, v. (1830) 165.—Mexic.
calocephala, *DC. Prod.* iv. 567.—Mexic.
coccinea, *DC. l. c.*—Mexic.
glaucescens, E. Mey. ex Harv. in Hook. Lond. Journ. Bot. i. (1842) 22 = Pentanisia variabilis.
hispidula, Hemsl. Biol. Centr. Am. Bot. ii. 57 = aspera.
lanceolata, E. Mey. ex Hochst. in Flora, xxvi. (1843) 70 = Pentanisia variabilis.
longibracteata, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1855) 98.—Nicaragua.
lucida, *Benth. Bot. Voy. Sulph.* 109.—Mexic.
parviflora, *Hook. & Arn. Bot. Beech. Voy.* 430. t. 99.—Mexic.
rubra, *Bartl. in Linnaea*, xiv. (1840) *Litt.* 125.—Mexic.
rubra, Cham. & Schlecht. in Linnaea, v. (1830) 165, excl. syn. = aspera.
subulata, *Hook. & Arn. Bot. Beech. Voy.* 431.—Mexic.
subulata, *A. Gray, in Proc. Am. Acad.* xix. (1883) 78. —Mexic.
subulata, Walp. Rep. ii. 467, sphalm. = subalata.
variabilis, E. Mey. ex Harv. in Hook. Lond. Journ. Bot. i. (1842) 22 = Pentanisia variabilis.
violacea, *Bronga. ex Neumann, in Nov. Hortic.* Sér. II. iv. (1848–46) 268; v. (1846) 61.—Mexic.
Wrightii, *A. Gray, Pl. Wright.* ii. 98.—N. Mexic.

CRUSEA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 304. t. 19. f. 1 = **Chiocca**, DC. (Rubiac.).
glabra, A. Rich. l. c. = Chiocca glabra.
nitida, A. Rich. l. c. t. 19. f. 1 = præc.

CRUZEA, A. Rich. Fl. Cub. Fanerog. ii. 29, in syn. (1853) = CRUSEA, A. Rich. = Chiocca, DC. (Rubiac.).

CRUZETA, Loefl. Iter, 208 (1758) = CRUCITA, Linn. = Trexinus, Linn. (Amarant.).
americana, Moq. in DC. Prod. xiii. ii. 349 = I. celosioides?
hispanica, Loefl. Iter, 208 = *americana*.

CRUZITA, Linn. Syst. ed. XII. 139 (1770) = CRUCITA, Linn. = Trexinus, Linn. (Amarant.).

CRYBE, Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = **Aceratanus**, Linn. (Orchid.).
rosea, Lindl. l. c. = A. rosea.

CRYOSOPHILA, Blume, Rumphia, ii. 58, in nota (1836) = **Copernicia**, Mart. (Palm.).
nana, Blume, l. c. = Cop. nana.

CRYPHAEA, Buch.-Ham. in Brewst. Edinb. Journ. Sc. ii. (1825) 11 = **Chloranthus**, Sw.
erecta, Buch.-Ham. l. c. = Chloranthus officinalis.

CRYPHIA, R. Br. Prod. 508 (1810) = **Prostanthera**, Labill. (Labiat.).
microphylla, R. Br. l. c. = P. microphylla.
serpyllifolia, R. Br. l. c. = P. microphylla.

CRYPHIACANTHUS, Nees, in Del. Sem. Hort. Vrat. (1841); et in Linnaea, xvi. (1842) 298 = **Ruellia**, Linn. (Acanthac.).
acaulis, Nees, in Mart. Fl. Bras. ix. 49.—Bras.
angustifolius, Nees, in DC. Prod. xi. 199 = R. Tweediana.
australis, Nees, l. c. 198 = R. australis.
barbadensis, Nees, Del. Sem. Hort. Vrat. (1841) = R. tuberosa.
clandestinus, Nees in DC. Prod. xi. 156, in syn. = R. tuberosa.
Diptracanthus, Nees, l. c. 190 = R. Diptracanthus.
lactus, Nees, Del. Sem. Hort. Vrat. (1841) = R. lactea.
macrophyllus, Nees, in DC. Prod. xi. 198 = R. lactea.
ovalifolius, Oerst. in Kjoeb Vidensk. Meddel. (1854) 129 = R. ovalifolia.
pumilus, Nees, in DC. Prod. xi. 199 = R. pumila.
tubiflorus, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 129.—Mexic.
udus, Nees, in Mart. Fl. Bras. ix. 49.—Bras.
viscosus, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 128 = R. tuberosa.

CRYPHIANTHA, Eckl. & Zeyh. Enum. 171 (1836) = **Amphithalea**, Eckl. & Zeyh. (Legumin.).
imbricata, Eckl. & Zeyh. l. c. 172 = A. micrantha.

CRYPHIOSPERMUM, Beauv. Fl. Owar. ii. 24. t. 74 (1806) = **Rhyssopteris**, Lour. (Compos.).
abyssinicum, Sch. Bip. in Walp. Rep. vi. 147 = Sigesbeckia abyssinica.
repens, Beauv. Fl. Owar. ii. 24. t. 74 = E. fluctuans.

CRYPSINNA, Fourn. ex Benth. in Journ. Linn. Soc. xix. (1881) 87 = **Epicampes**, Presl (Gramin.).
macroura, Fourn. ex Benth. l. c. = E. macroura.
setifolia, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 506 = E. macroura.
stricta, Fourn. l. c. = E. macroura.

CRYPSIS, Ait. Hort. Kew. i. 48 (1789); iii. 505. *GRAMINEAE*, Benth. & Hook. f. iii. 1139.
ANTITRAGUS, Gaertn. Fruct. ii. 7. t. 80 (1791).
FACHEA, Fourr. ex Steud. Nom. ed. II. i. 449, in syn. (1840).
FALLASIA, Scop. Introd. 72 (1777).
RADDIA, [Pier†] in Ionlos Anthol. iii. (1884) 448.
aculeata, *Ait. Hort. Kew.* i. 48.—Geront. temp. et trop.
acuminata, Trin. in Spreng. Neue Entdeck. ii. 57 = Heleochloa alopecuroides.
aegyptiaca, Trinch. in Flora, xvii. (1834) 120 = Heleochloa alopecuroides.
alopecuroides, Guss. ex Schult. Mant. ii. 569 = Agrostis brachystachys.
alopecurus, Schrad. Fl. Germ. i. 167 = Heleochloa alopecuroides.
alpicola, Hochst. ex Steud. Syn. Pl. Gram. 153 = Heleochloa alpicola.
ambigua, Boiss. & Bal. ex Boiss. Fl. Orient. v. 477 = Heleochloa ambigua.
arenaria, Desf. Fl. Atlant. i. 68 = Phleum arenarium.
Boraeoweri, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) ii. 306.—As. centr.
compacta, *Steud. Syn. Pl. Gram.* 151.—Afr. trop.
cruciannelloides, Boiss. & Bal. ex Boiss. Fl. Orient. v. 477 = Heleochloa cruciannelloides.
dura, Boiss. Diagn. Ser. II. iv. 125 = Heleochloa dura.
fenicularia, Roem. & Schult. Syst. ii. 676 = Heleochloa alopecuroides.
juncea, Steud. Nom. ed. I. 242 = Sporobolus junceus.
macrostachya, Brot. Fl. Lusit. i. 78 = Heleochloa alopecuroides.
macroura, H. B. & K. Nov. Gen. et Sp. i. 140 = Epicampes macroura.

CRYPSIS:—
maritima, Munro, ex Macowan, in Cape Month. Mag. N. S. iii. (1871) Suppl. 7 = Sporobolus virginicus.
Myosurus, *Nees, ex Steud. Syn. Pl. Gram.* 152.—Aegypt.
nigricans, Guss. Fl. Sic. Prod. i. 73 = Agrostis brachystachys.
villaca, Figar. & De Notar. in Mem. Acc. Tor. Ser. II. xiv. (1854) 223 = Heleochloa schoenoides.
phalaroides, Bleb. Fl. Taur. Cauc. i. 45 = Heleochloa alopecuroides.
phleoides, H. B. & K. Nov. Gen. et Sp. i. 140 = Cinna phleoides.
pygmaea, Jaub. & Spach, Illustr. iv. 8 = Heleochloa pygmaea.
schenoides, Beauv. Agrost. 22 = aculeata.
schoenoides, Hochst. ex Steud. Syn. Pl. Gram. 152 = Heleochloa alopecuroides.
schoenoides, Lam. Tabl. Encyc. i. 166 = Heleochloa schoenoides.
setifolia, J. & C. Presl, Reliq. Haenk. i. 345 = Epicampes macroura.
sicula, Jan, Elench. 1 = nigricans.
squarrosa, *Nutt. Gen. Am.* i. 49.—Am. bor.
stricta, H. B. & K. Nov. Gen. et Sp. i. 140 = Epicampes macroura.
tenuissima, *Tausch, in Flora*, xx. (1837) 121.—Sibiria.
vaginiflora, *Opiz, Natural.* viii. (1824) 88.—Aegypt.
virginica, Nutt. Gen. Am. i. 49 = Heleochloa schoenoides.

CRYPSOCALYX, Endl. in R. Br. Verm. Schr. v. 290, in adnot. (1834), sphalm. = CHRYSOCALYX, Guill. & Perr. = **Crotalaria**, Linn. (Legumin.).

CRYPTA, Nutt. in Journ. Acad. Philad. i. (1817) 117. t. 6 = **Elatine**, Linn.
minima, Nutt. l. c. = E. americana.
triandra, A. Braun, ex Walp. Rep. v. 83 = E. triandra.

CRYPTADENIA, Meisn. in Linnaea, xix. (1840) 404. *THYMELAEACEAE*, Benth. & Hook. f. iii. 195.
CALYSERICOS, Eckl. & Zeyh. ex Meisn. in DC. Prod. xv. 573 (1856–57).
brevifiora, *Meisn. in Linnaea*, xix. 406.—Afr. austr.
ciliata, *Meisn. l. c.* 407.—Afr. austr.
filicaulis, *Meisn. l. c.*—Afr. austr.
grandiflora, *Meisn. l. c.* 405.—Afr. austr.
uniflora, *Meisn. l. c.* 406.—Afr. austr.

CRYPTADIA, Lindl. ex Endl. Gen. Pl. Suppl. i. 1381 (1841) = **Gymnarrhena**, Desf. (Compos.).
cuphrantoides, [Lindl. in] Chem. Narrat. Exph. (1858) t. ad p. 441 = G. micrantha.

CRYPTANDRA, Sm. in Trans. Linn. Soc. iv. (1798) 217. *RHAMNEAE*, Benth. & Hook. f. i. 383.
WICHURAEA, Nees, in Lehm. Pl. Preiss. ii. 200 (1847).
albicans, F. Muell. Census, 60 = Trymalium albicans.
alpina, *Hook. f. Fl. Tasm.* i. 75. t. 12.—Tasmania.
amara, *Sm. in Rees, Cycl.* x. n. 2.—Australi.
angustifolia, F. Muell. Census, 60 = Trymalium angustifolium.
anomala, Steud. in Lehm. Pl. Preiss. i. 187 = Trymalium ledifolium.
arbutiflora, *Fenzl, in Enum. Pl. Huegel.* 96.—Austral.
australis, Pers. Syn. i. 350 (sp. delend.).
Behriana, Reiss. in Linnaea, xxix. (1857–58) 298 = tomentosa.
bifida, F. Muell. Census, 61 = Spyridium bifidum.
Billardieri, F. Muell. l. c. 60 = Trymalium Billardieri.
buxifolia, *Fenzl, in Enum. Pl. Huegel.* 98.—Austral.
campanulata, F. Muell. Fragm. iii. 67 = tomentosa.
campanulata, Schlecht. in Linnaea, xx. (1847) 689 = amara.
capitata, Sieber, ex Benth. Fl. Austral. i. 436 = ericifolia.
coactiflolia, F. Muell. Census, 61 = Spyridium coactiflolium.
complicata, F. Muell. l. c. 61 = Spyridium complicatum.
cordata, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 459 = Spyridium cordatum.
coronata, Reiss. in Lehm. Pl. Preiss. ii. 388 = Stenanthemum coronatum.
Daltoni, F. Muell. Census, 60 = Trymalium Daltoni.
divaricata, F. Muell. l. c. 61 = Spyridium divaricatum.
divaricata, Reiss. in Lehm. Pl. Preiss. ii. 388 = tomentosa.
ericifolia, Rudge, in Trans. Linn. Soc. x. (1811) 294. t. 18 = sqq.

CRYPTANDRA :—
ericoides, *Sm. in Rees, Cyclop.* x. (1808) n. 1.—Austral.
eriocephala, Hook. f. Fl. Tasm. i. 72 = Spyridium eriocephalum.
erubescens, F. Muell. ex Reiss. in Linnæa, xxix. (1857–58) 298 = tomentosa.
floribunda, Steud. in Lehm. Pl. Preiss. i. 186 = Trymalium ledifolium.
glabrata, Steud. l. c. 188 = miliaris.
glabriflora, *Benth. Fl. Austral.* i. 441.—Austral.
glaucophylla, Steud. in Lehm. Pl. Preiss. i. 187 = Trymalium ledifolium.
globulosa, F. Muell. Census, 61 = Spyridium globulosum.
Gunnii, Hook. f. Fl. Tasm. i. 75. t. 11 = Spyridium Gunnii.
halmaturina, F. Muell. Census, 61 = Spyridium halmaturinum.
hispidula, *Reiss. in F. Muell. in Linnæa*, xxix. (1857–58) 294.—Austral.
holostyla, Steud. in Lehm. Pl. Preiss. i. 188 = pungens.
humilis, F. Muell. Census, 61 = Stenanthemum humile.
Hookeri, F. Muell. l. c. = Spyridium parvifolium.
intangenda, F. Muell. Fragm. x. 52 = Pomaderris intangenda.
lanuginosa, *F. Muell. l. c.* iii. 65.—Austral.
largiflora, F. Muell. ex Reiss. in Linnæa, xxix. (1857–58) 292 = amara.
laxiphylla, Steud. in Lehm. Pl. Preiss. i. 188 = miliaris.
Lawrencii, Hook. f. Fl. Tasm. i. 72 = Spyridium Lawrencii.
ledifolia, F. Muell. Census, 60 = Trymalium ledifolium.
leucophracta, Schlecht. in Linnæa, xx. (1847) 640 = Stenanthemum leucophracta.
Leucopogon, *Meisn. ex Reiss. in Lehm. Pl. Preiss.* ii. 287.—Austral.
longistaminea, *F. Muell. Fragm.* iii. 64.—Austral.
magnifolia, F. Muell. l. c. 65 = propinqua.
microcephala, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 458 = Spyridium microcephalum.
miliaris, *Reiss. in Lehm. Pl. Preiss.* ii. 288.—Austral.
mollis, Hook. f. Fl. Tasm. i. 73 = Spyridium parvifolium.
mollis, *Nees, ex Reiss. in Lehm. Pl. Preiss.* ii. 289.—Austral.
nervata, Reiss. in Linnæa, xxix. (1857–58) 292 = amara.
nudiflora, *F. Muell. Fragm.* iii. 64.—Austral.
nutans, *Steud. in Lehm. Pl. Preiss.* i. 186.—Austral.
obcordata, Hook. f. Fl. Tasm. i. 71 = Spyridium serpyllaceum.
obovata, Hook. f. l. c. 74 = Spyridium obovatum.
parvifolia, Hook. f. l. c. 73 = Spyridium parvifolium.
parvifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 459.—Austral.
pauciflora, Turcz. l. c. 458 = Spyridium pauciflorum.
phlebophylla, F. Muell. Census, 61 = Spyridium phlebophyllum.
pimelioides, Hook. f. Fl. Tasm. i. 75. t. 12 B = Stenanthemum pimelioides.
pomaderroides, Reiss. in Endl. Nov. Stirp. Dec. 29 = Stenanthemum pomaderroides.
propinqua, *A. Cunn. ex Fenzl, in Enum. Pl. Hueg.* 23.—Austral.
propinqua, Schlecht. in Linnæa, xx. (1847) 638 = tomentosa.
pumila, F. Muell. Census, 61 = Spyridium pumilum.
pungens, *Steud. in Lehm. Pl. Preiss.* i. 187.—Austral.
pyramidalis, Brongn. in Ann. Sc. Nat. Sér. 1. x. (1827) 373 = spinescens.
rotundifolia, F. Muell. Census, 61 = Spyridium rotundifolium.
scoparia, *Reiss. in Lehm. Pl. Preiss.* ii. 285.—Austral. Scortechinii, *F. Muell. in Austral. Chem. & Druggr.* (1884).—Austral.
Sieberi, Fenzl, in Enum. Pl. Hueg. 23 = amara.
spadicea, F. Muell. Census, 61 = Spyridium spadiceum.
spathulata, F. Muell. l. c. = Spyridium spathulatum.
spinescens, *Sieber, ex DC. Prod.* ii. 33.—Austral.
spinosa, A. Cunn. ex G. Don, Gen. Syst. ii. 40 = ericoides!
spyridioides, *F. Muell. Fragm.* iii. 66.—Austral.
suaveis, Lindl. Bot. Reg. (1844) t. 56. Misc. 25 = arbutifiora.
subcoriacea, F. Muell. Census, 61 = Spyridium subcoriaceum.
tomentosa, *Lindl. in Mitch. Three Exped.* ii. 178.—Austral.
tomentosa, Reiss. in Lehm. Pl. Preiss. ii. 286 = patens.
tridentata, Steud. in Lehm. l. c. i. 186 = Spyridium tridentatum.

CRYPTANDRA :—
tubulosa, Fenzl, in Enum. Pl. Hueg. 26 = arbutiflora.
ulicina, Hook. Journ. Bot. i. (1834) 257 = Spyridium ulicinum.
vexilliflora, Hook. l. c. = Spyridium vexilliflorum.
villosa, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 458 = Spyridium villosum.
Waterhousii, F. Muell. Census, 61 = Stenanthemum Waterhousii.
Wayii, F. Muell. l. c. 140 = Trymalium Wayse.
westringioides, F. Muell. l. c. 61 = Spyridium westringiafolium.
Wichurae, F. Muell. l. c. 60 = Trymalium Wichurae.

CRYPTANGIUM, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 168. t. 19 (1842). *CYPERACEAE*, Benth. & Hook. f. iii. 1067.
ACROCARPUS, Nees, in Mart. Fl. Bras. ii. 1. 157. tt. 16, 17 (1842).
CLADOTHECA, Steud. Syn. Pl. Cyp. 178 (1855).
MICROLEPIS, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 157 (1842).
arundinaceum, Boeck. in Flora, lxv. (1882) 351 = Lagenocarpus rigidus.
campestre, *Boeck. in Linnæa*, xxxviii. (1874) 416.—Bras.
comatum, *Boeck. in Flora*, lxv. (1882) 351.—Bras.
densifolium, Boeck. in Linnæa, xxxviii. (1874) 418 = Minarum.
Dracaenula, Boeck. l. c. 415 = Cephalocarpus Dracaenula.
Glaziovii, *Boeck. in Kjoeb. Vidensk. Meddel.* (1879–80) 28.—Bras.
humile, *Boeck. in Linnæa*, xxxviii. (1874) 418.—Bras.
junciforme, *Boeck. l. c.* 415.—Bras.
leptoculmum, *Boeck. l. c.* 420.—Bras.
Minarum, *Boeck. l. c.* 417.—Bras.
pauciflorum, Boeck. in Kjoeb. Vidensk. Meddel. (1879–80) 28 = minarum.
polyphyllum, *Boeck. in Linnæa*, xxxvii. (1874) 414.—Bras.
stellatum, Boeck. l. c. 421 (Lagenocarpi sp.).—Bras.
triquetrum, *Boeck. l. c.* 418.—Bras.
uliginosum, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. 1. 161.—Bras.

CRYPTANTHA, Lehm. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 35 (1836) = **Eritrichium**, Schrad. (Borragin.).
glomerata, Lehm. ex Fisch. & Mey. l. c. = E. cryptanthum.
microcarpa, Fisch. & Mey. l. c. = Krynitzkia clandestina.

CRYPTANTHE, Benth. & Hook. f. Gen. ii. 851 (1876) = prsec.

CRYPTANTHUS, Nutt. ex Moq. in DC. Prod. xiii. ii. 54 (1849) = **Aphanisma**, Nutt. (Chenopod.).
bitoides, Nutt. l. c. = A. bitoides.

CRYPTANTHUS, Otto & Dietr. Allg. Gartenz. iv. (1836) 298. *BROMELIACEAE*, Benth. & Hook. f. iii. 661.
MADVIGIA, Liebm. in Ann. Sc. Nat. Sér. IV. ii. (1854) 378.
PHOLIDOPHYLLUM, Vis. Sem. Hort. Patav. (1847) 4.
acaulis, Beer, Bromel. 75 = undulatus.
Beuckeri, *E. Morr. in Belg. Hortic.* xxx. (1880) 241; xxxi. (1881) 349. t. 17.—Bras.
bivittatus, *Regel, Ind. Sem. Hort. Petrop.* (1864) 15. —Am. austr.
bromelioides, *Otto & Dietr. Allg. Gartenz.* iv. (1836) 298.—Bras.
discolor, *Otto & Dietr. l. c.* 299.—Bras.
diversifolius, Beer, Bromel. 76 = bromelioides.
undulatus, *Otto & Dietr. Allg. Gartenz.* iv. (1836) 299.—Bras.
zonatus, *Beer, Bromel.* 76.—Bras.

CRYPTARRHENA, R. Br. in Bot. Reg. t. 153 (1816). *ORCHIDEAE*, Benth. & Hook. f. iii. 566.
CLINEMMENIA, A. Rich. & Gal. in Compt. Rend. Acad. Sc. Par. xviii. (1844) 513.
ORCHIDOPUNCKIA, A. Rich. in Ann. Sc. Nat. Sér. III. iii. (1845) 24.
Kegolii, *Reichb. f. in Bot. Zeit.* x. (1852) 766.—Guian.
lanata, *R. Br. in Bot. Reg.* t. 153.—Ind. occ.
pallidiflora, *Reichb. f. in Bot. Zeit.* x. (1852) 766.—Mexic.

CRYPTERONIA, Blume, Bijdr. 1151 (1826). *LYTHRARIEAE*, Benth. & Hook. f. i. 783.
HENSLOWIA, Wall. Pl. As. Rar. iii. 13 (1832).
QUILAMUM, Blanco, Fl. Filip. ed. I. 551 (1837).
affinis, Endl. Gen. Suppl. iv. 11. 39 = pubescens.
Cumingii, *Endl. l. c.*—Ins. Philipp.
glabra, *Endl. l. c.*—Reg. Himal.; Malaya.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 574.—Malaya.
Hookeri, A. DC. in DC. Prod. xvi. 11. 679 = pubescens.
leptostachya, *Endl. Gen. Suppl.* iv. 11. 39.—Ins. Philipp.
luzea, *Blume, Mus. Bot. Lugd. Bat.* ii. 138.—Java.
paniculata, *Blume, Bijdr.* 1151.—Java.
paniculata, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) 11. 86 = pubescens.
pubescens, *Endl. Gen. Suppl.* iv. 11. 39.—Burma; Malaya.
Wallichii, *A. DC. in DC. Prod.* xvi. 11. 679.—Burma.

CRYPTINA, Rafin. in Am. Monthly Mag. (1819) 192 = **Illatine**, Lunn.

CRYPTOCALYX, Benth. in Ann. Nat. Hist. Ser. I. ii. (1839) 446 = **Lippia**, Linn. (Verbenac.).
nepetaefolia, Benth. l. c. = L. betulaefolia.

CRYPTOCARIA, C. Gay, Fl. Chil. v. 299 (1849) = **Cryptocarya**, R. Br. (Laurin.).

CRYPTOCARPA, Steud. Nom. ed. II. i. 449, sphalm = seq.

CRYPTOCARPHA, Cass. in Bull. Soc. Philom. (1817) 84 = **Actocarpha**, Juss. (Calycer.).
lanata, Cass. in Dict. Sc. Nat. xii. 86 = A. lanata.
spathulata, Cass. l. c. 85 = A. spathulata.
tribuloides, Cass. l. c. = A. tribuloides.

CRYPTOCARPUS, H. B. & K. Nov. Gen. et Sp. ii. 187. t. 128, 124 (1817). *NYCTAGINEAE*, Benth. & Hook. f. iii. 11.
cordifolius, Moric. Fl. Nouv. Am. 75 = pyriformis.
globosus, *H. B. & K. Nov. Gen. et Sp.* ii. 187. t. 122. —Cuba.
paniculatus, *H. B. & K. ex Schlecht. in Linnæa*, xxvi. (1853–55) 645.—N. Granat.
pyriformis, *H. B. & K. Nov. Gen. et Sp.* ii. 188. t. 124. —Ecuador.
rhomboideus, Moq. in DC. Prod. xiii. ii. 88 = globosus.

CRYPTOCARYA, R. Br. Prod. 402 (1810). *LAURINEAE*, Benth. & Hook. f. iii. 150.
BELLOTA, C. Gay, Fl. Chil. v. 297. t. 59 (1849).
BOLDU, Nees, Syn. Laurin. 177 (1836).
CARYODAPHNE, Blume, ex Nees, Syst. Laurin. 225 (1836).
ENDOCARPA, Rafin. in Sylva Tellur. 135 (1838).
acutifolia, Blume, ex Nees, Syst. Laurin. 214 = costata.
amygdalina, *Nees, in Wall. Pl. As. Rar.* iii. 69.—Reg. Himal.
angustifolia, *E. Mey. Zwei Pfl. Docum.* 176, nomen; *Meisn. in DC. Prod.* xv. i. 74.—Afr. austr.
australis, *Benth. Fl. Austral.* v. 229.—Austral.
Aw-uma, Mart. ex Nees, l. c. 246 = Aydendron pernolle.
Berterreana, C. Gay, Fl. Chil. v. 301 = C. Peumus.
Bidwillii, Meisn. in DC. Prod. xv. i. 74 = glaucescens.
casta, *Blume, Mus. Bot. Lugd. Bat.* i. 285.—Ins. Andam.; Java.
Canellila, *H. B. & K. Nov. Gen. et Sp.* vii. 192.—Am. austr.
caudata, Meisn. in DC. Prod. xv. i. 68 = Griffithiana.
cinnamomifolia, *Benth. Fl. Austral.* v. 228.—Austral.
conolinna, *Hance, in Journ. Bot.* xx. (1882) 79.—China.
costata, *Blume, Bijdr.* 558.—Java.
crassifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 241. —Madag.
crassinervia, *Miq. Fl. Ind. Bat.* i. 1. 924.—Sumatra.
cuneata, Blume, Bijdr. 558 = Tabascia cuneata.
cunneata, Zoll. ex Meisn. in DC. Prod. xv. i. 67 = Phœbe javanica.
Cunninghamii, Meisn. l. c. 73 = glaucescens.
declinata, *Baker, in Journ. Linn. Soc.* xx. (1883) 241. —Madag.
densiflora, *Blume, Bijdr.* 556.—Java.
discolor, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 283 = Litsea timoriana.
diversifolia, *Blume, l. c.* 285.—Sumatra.
Dubia, *H. B. & K. Nov. Gen. et Sp.* ii. 187.—N. Granat.
Durio, Sprang. ex Blume, Syst. Laurin. 209 (= *Magilus duplex phillyreoides*).—Moluc.

CRYPTOCARYA :—
emarginata, *Meisen. in DC. Prod.* xv. i. 76.—Bras.
ferrea, *Blume, Bijdr.* 557.—Malaya.
ferrea. *Kurz, For. Fl. Brit. Burm.* ii. 296 (sp. diversa). —Burma.
ferruginea, Blume, Cat. Gew. Buitenz. 65 = Litsea ferruginea.
ferruginea, A. Cunn. ex Meisen. in DC. Prod. xv. i. 78 = tripilnervia.
floribunda, Miq. ex Meisn. l. c. 40 = Machilus macrantha.
floribunda, Nees, in Wall. Pl. As. Rar. ii. 69 = amygdalina.
floribunda, Wight, Ic. t. 1829 = Wightiana.
glaucescens, *R. Br. Prod.* 402.—Austral.
glaucescens, Hask. Cat. Hort. Bogor. Alt. 89 = tomentosa.
Griffithiana, *Wight, Ic.* t. 1830.—Ind. or.
guianensis, *Meisen. in DC. Prod.* xv. i. 75.—Guiana.
hirsuta, *Schott, in Spreng. Syst. iv. App.* 405.—Bras.
hypoglauca, Meisen. in DC. Prod. xv. i. 78 = glaucescens.
hypospodia, F. Muell. Fragm. v 170 = obovata.
impressa, *Miq. Fl. Ind. Bat.* i. 1. 923.—Malaya.
inflectoria, Miq. l. c. 924 = Griffithiana.
laevigata, *Blume, Bijdr.* 556.—Java.
laevis, Nees, ex Mart. in Flora, xxi. (1838) ii. 64 = Aiouea brasiliensis.
latifolia, *Sond. in Linnaea*, xxiii. (1850) 101.—Afr. austr.
lucida, *Blume, Mus. Bat. Lugd. Bat.* i. 835.—Borneo.
lucidula, *Miq. Fl. Ind. Bat.* i. i. 922.—Japon.
Mackinnoniana, *F. Muell. Fragm.* v. 169.—Austral.
macrophylla, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 14 = Cylicodaphne macrophylla.
mandioccana, *Meisen. in DC. Prod.* xv. i. 75.—Bras.
Meisneri, *F. Muell. Fragm.* v. 170.—Austral.
membranacea, *Thw. Enum. Pl. Zeyl.* 254.—Zeylan.
Mentek, *Blume, ex Nees, Syst. Laurin.* 218.—Java.
micrantha, *Meisn. in DC. Prod.* xv. i. 75.—Bras.
micrococca, Meisn. l. c. 78 = glaucescens.
monticola, Mart. ex Nees, Syst. Laurin. 400 (= *Mespilodaphne phillyreoides*).—Bras.
morstoniana, Meisn. in DC. Prod. xv. i. 74 = glaucescens.
moschata, *Nees & Mart. in Linnaea*, viii. (1833) 37.—Bras.
mucronata, *Spreng. Syst.* ii. 271.—Guiana.
*Mueller*i, Meisn. in DC. Prod. xv. i. 78 = Endiandra pubens.
Murrayi, *F. Muell. Fragm.* v. 170.—Austral.
myristicoides, *Baker, in Journ. Linn. Soc. xx.* (1883) 241.—Madag.
neilgherrensis, Meisn. in DC. Prod. xv. i. 71 = Stocksii.
nitida, *Phil. in Linnaea*, xxxiii. (1864–65) 228.—Chili.
oblique, *Blume, Mus. Bot. Lugd. Bat.* i. 334.—Java.
oblongifolia, *Blume, Bijdr.* 557.—Java.
obovata, *R. Br. Prod.* 403.—Austral.
obscura, *Blume, Mus. Bot. Lugd. Bat.* i. 335.—Java.
obtusifolia, F. Muell. ex Meisn. in DC. Prod. xv. i. 508 (= *Neocinnamomum obtusifolia*).—Austral.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 359.—Sumatra.
patentinervis, *F. Muell. in DC. Prod.* xv. i. 508.—Austral.
pendulifolia, *Dalzell. in Bull. Soc. Linn. Par.* i. (1885) 447. —Madag.
Pennus, *Nees, Syst. Laurin.* 223.—Chili.
pretiosa, Mart. ex Nees, l. c. 287; Reise Bras. iii. 1158 = Cinnam pretiosa.
pyriformis, Nees, Syst. Laurin. 220 (= *Mespilodaphne pyriformis*).—Guiana.
reticulata, *Blume, Mus. Bot. Lugd. Bat.* i. 335.—Java.
retrospiralis, Voigt, Hort. Suburb. Calc. 585 = Cryptocoryne retrospiralis.
Riedelii, *Meisen. in DC. Prod.* xv. i. 75.—Bras.
riparia, Meisn. l. c. 508 = patentinervis.
rubiginosa, *Griff. Notul.* iv. 739.—Burma.
stenantha, Phil. in Linnaea, xxxiii. (1864–65) 228 = C. Pennus.
Stocksii, *Meisen. in DC. Prod.* xv. i. 71.—Ind. or.
Teysmanniana, *Miq. Fl. Ind. Bat. Suppl.* 360.—Sumatra.
tomentosa, *Blume, Mus. Bot. Lugd. Bat.* i. 335.—Java.
triplinervis, *R. Br. Prod.* 402.—Austral.
Wightiana, *Thw. Enum. Pl. Zeyl.* 254.—Ind. or.
Zollingeriana, *Miq. Fl. Ind. Bat.* i. i. 924.—Java.

CRYPTOCENTRUM, Benth. in Journ. Linn. Soc. xviii. (1881) 325. *ORCHIDEAE*, Benth. & Hook. f iii. 557.
Jamesoni, *Benth. in Benth. & Hook. f. Gen.* iii. 557.—Ecuador.

CRYPTOCERAS, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 121 = **Corydalis**, Vent. (Paper.).
bracteatum, C. Muell. in Walp. Ann. vii. 91 = Coryd. bracteata.
giganteum, C. Muell. l. c. = Coryd. gigantea.
longiflorum, C. Muell. l. c. = Coryd. longiflora.
modestum, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 150 = Coryd. rutaefolia.
oppositiflorum, Schott, l. c. iv. (1854) 121 = Coryd. rutaefolia.
pauciflorum, C. Muell. in Walp. Ann. vii. 91 = Coryd. pauciflora.
pulchellum, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 150 = Coryd. rutaefolia.
purpurans, Schott, in Oestr. l. c. = Coryd. rutaefolia.
purpurascens, Boiss. Fl. Orient. i. 128 = praec.
remotum, C. Muell. in Walp. Ann. vii. 91 = Coryd. bulbosa.
rutifolium, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 121 = Coryd. rutaefolia.
verticillare, Schott, l. c. = Coryd. verticillaris.

CRYPTOCHILOS, Spreng. Gen. ii. 676 (1831) = seq.

CRYPTOCHILUS, Wall. Tent. Fl. Nep. 36. t. 36 (1822). *ORCHIDEAE*, Benth & Hook. f. iii. 518.
lutea, *Lindl. in Journ. Linn. Soc.* iii. (1859) 21.—Reg. Himal.
Meirax, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 145 = Eria Meirax.
reticulatus, Reichb. f. in Bot. Zeit. xx. (1862) 214 = Eria reticulata.
sanguinea, *Wall. Tent. Fl. Nep.* 36.—Reg. Himal
Wightii, Reichb. f. in Bot. Zeit. xx. (1862) 214 = Eria Lichenora.

CRYPTOCHLORIS, Benth. in Hook. Ic. Pl. t. 1276 (1882). *GRAMINEAE*, Benth. & Hook. f. iii. 1166.
spathacea, *Benth. l. c.*—Patagon.

CRYPTOCORYNE, Fisch. ex Reichb. Consp. 44 (1828). *AROIDEAE*, Benth. & Hook. f. iii. 968.
MELIOBLASTIS, C. Muell. in Bot. Zeit. iv. (1846) 158, sphalm.
MYRIOBLASTUS, Wall. ex Griff. in Proc. Linn. Soc i. (1845) 264.
alata, Griff. Ic. Pl. iii. tt. 170, 171 = ciliata.
auricsinta, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 302.—Borneo.
Beckettii, *Thw. ex Trim. in Journ. Bot.* xxiii. (1885) 269.—Zeylan.
bulbosa, *Becc. ex Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 302.—Borneo.
caudata, *N. E. Br. in Journ. Linn. Soc.* xvii. (1880) 242.—Borneo.
ciliata, *Fisch. ex Schott, Melet.* 16, nomen : *Blume, Rumphia*, i. 56.—Ind. or.
cognata, *Schott, in Bonplandia*, v. (1857) 232.—Ind. or.
consobrina, *Schott, l. c.*—Ind. or.
cordata, *Griff. Notul.* iii. 138.—Malaya.
Dalzellii, *Schott, in Bonplandia*, v. (1857) 221.—Ind. or.
dicymorrhiza, Zipp. ex Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 122 = ciliata.
egregia, *Schott, l. c.* 128.—Sumatra.
elata, Griff. Notul. iii. 134 = ciliata.
ferruginea, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 302.—Borneo.
Gomezii, *Schott, in Bonplandia*, v. (1857) 221.—Ind. or.
Griffithii, *Schott, Syn. Aroid.* 1.—Malacca.
Hoegelii, *Schott, Aroideae*, 8. t. 12.—Ind. or.
Koenigii, Schott, in Bonplandia, vii. (1859) 81 = Lagenandra Koenigii.
lanceifolia, Schott, l. c. v. (1857) 221 = Lagenandra lanceifolia.
Lingua, *Becc. ex Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 301.—Borneo.
longicauda, *Becc. ex Engl. l. c.* 302.—Borneo.
ovata, Schott, Melet. 16 = Lagenandra toxicaria.

CRYPTOCORYNE :—
pallidinervia, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 301.—Borneo.
pontederiifolia, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 122.—Sumatra.
retrospiralis, *Kunth, Enum. Pl.* iii. 19.—Ind. or.
Roxburghii, Schott, Prod. Aroid. 18 = unilocularis.
spathulata, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 301.—Borneo.
sparsila, *Pisch. ex Wydler, in Linnaea*, v. (1830) 428. —Ind. or.
striolata, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 301.—Borneo.
Thwaitesii, *Schott, in Bonplandia*, v. (1857) 221.—Zeylan.
unilocularis, *Kunth, Enum. Pl.* iii. 18.—Ind. or.
uniloculiara, Schott, in Bonplandia, v. (1857) 223 = Wightii.
Walkeri, *Schott, l. c.* 221.—Zeylan.
Wightii, *Schott, Prod. Aroid.* 17.—Ind. or.

CRYPTODIA, Sch. Bip. in Walp. Rep. ii. 955 (1843), sphalm. = CRYPTADIA, Lindl. = **Gymnarrhena**, Desf. (Compos.).

CRYPTODISCUS, Schrenk, Enum. Pl. Nov. 64 (1841). *UMBELLIFERAE*, Benth. & Hook. f. i. 879.
ammophilus, *Bunge, Lehm. Reliq. Bot.* 139.—Turkestan.
cachroides, *Schrenk, Enum. Pl. Nov.* 65.—Sibir.
persicus, *Boiss. Fl. Orient.* ii. 935.—Persia.
rutaefolius, *Bunge, Lehm. Reliq. Bot.* 138.—Turkestan

CRYPTOGLOCHIN, Heuff. in Flora, xxvii. (1844) 527, 528 = **Carex**, Dill. (Cyperac.).
curvulus, Heuff. l. c. 528 = Carex curvula.

CRYPTOGLOTTIS, Blume, Bijdr. 296. t. 42 (1825) = **Podochilus**, Blume (Orchid.).
serpyllifolius, Blume, l. c. 297.—Java.

CRYPTOGYNE, Cass. in Dict. Sc. Nat. i. 491 (1827) = **Eriocephalus**, Linn. (Compos.).
absinthoides, Cass. l. c. 492.—Afr. austr.

CRYPTOGYNE, Hook. f. in Benth. & Hook. f. Gen ii. 656 (1876). *SAPOTACEAE*, Benth. & Hook. f. ii. 655.
Gerrardiana, *Hook. f. l. c.* 657.—Madag.

CRYPTOLEPIS, R. Br. in Mem. Wern. Soc i. (1809) 69. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 741.
CRYPTOLOBUS, Meisen. ex Steud. Nom. ed. II. i. 450 (1840).
LEPISTOMA, Blume, Fl. Jav. Praef. 7 (1828).
LEPOSMA, Blume, Bijdr. 1049 (1826).
Buchanani, *Kerm. & Schult. Syst.* iv. 409.—Ind. or. *Edithae*, Benth. & Hook. f. Gen. ii. 741 = elegans.
elegans, *Wall. Cat.* n. 1639.—Ind. or.; Malaya.
filiformis, Wall. l. c. n. 4458 = Atherandra acutifolia.
grandiflora, *Wight, Ic.* t. 881.—Ind. or.
javanica, *Blume, Mus. Bot. Lugd. Bat.* i. 146.—Java.
laxiflora, Blume, l. c. 147 = elegans.
longiflora, *Regel, Cat. Pl. Hort. Absahov.* 43.—Hab. ? *pauciflora*, Wight, Ic. t. 498 = elegans.
reticulata, Wall. Cat. n. 1640 = Buchanani.
Wightiana, Wall. l. c. n. 4457 = grandiflora.

CRYPTOLOBUS, Spreng. Anleit. ii. 760 (1818) ; Syst. iii. 154 (1826) = **Amphicarpaea**, Ell. et **Voandzeia**, Thou. (Legumin.).
africanus, Spreng. Syst. iii. 218 = V. subterranea.
americanus, Spreng. l. c. = A. monoica.
amentaceus, Spreng. l. c. 219 = A. monoica.
subterraneus, Spreng. Anleit. ii. 760 = V. subterranea.

CRYPTOLOBUS, Meisn. ex Steud. Nom. ed. II. i. 450, in syn. (1840) = **Cryptolepis**, R. Br. (Asclepiad.).

CRYPTOLOMA, Hanst. in Linnaea, xxix. (1857–58) 506 = **Isoloma**, Decne. (Gesneriac.).
anonymum, Hanst. l. c. 536.—Hab. ?
coloratum, Hanst. l. c.—N. Granat.
cordifolium, Hanst. l. c. 535 = I. cordifolium.
hondense, Klotzsch & Hanst. l. c. 534 = I. hondense.
Hoobarianum, Hanst. l. c. 536 = I. hondense.
pictum, Hanst. l. c. 536 = I. pictum.
pilosum, Hanst. l. c. 538 = I. pilosum.
rhynchocarpum, Hanst. l. c. = I. rhynchocarpum.
strictum, Hanst. l. c. = I. strictum.
trifiorum, Hanst. l. c. 534 = I. trifiorum.

CRYPTOMERIA, D. Don, in Trans. Linn. Soc. xviii. (1841) 166. f. 1. t. 18. *CONIFERÆ*, Benth. & Hook. f. iii. 429.
Fortunei, Otto & Dietr. Allg. Gartens. (1858) 324 = japonica.
japonica, *D. Don, in Trans. Linn. Soc.* xviii. (1841) 167.—Japon.
Lobbiana, Billain, in Otto & Dietr. Allg. Gartenz. (1853) 324 = japonica.
nigricans, Carr. in Rev. Hortic. (1870–71) 119 = japonica.

CRYPTONEMA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) t. 590 = Burmannia, Linn.
malaccensis, Turcz. l. c. 591 = B. coelestis.

CRYPTOPETALON, Cass. in Bull. Soc. Philom. (1817) 19; et in Dict. Sc. Nat. xii. 193 (1818) = Pectis, Linn. (Compos.).
ciliare, Cass. in Dict. Sc. Nat. xii. 193 = P. ciliaris.

CRYPTOPETALUM, DC. Prod. v. 101, in syn. (1836) = prec.

CRYPTOPETALUM, Benth. & Arn. in Hook. Bot. Misc. iii. (1833) 344 = Lepuropetalon, Ell. (Saxifrag.).
pusillum, Hook. & Arn. l. c. 345 = L. spathulatum.

CRYPTOPHORANTHUS, Rodrig. Orch. Nov. ii. 79 (1881) = Pleurothallis, R. Br. (Orchid.).
cryptanthus, Rodrig. l. c. 80.—Bras.
fenestratus, Rodrig. l. c.—Bras.
unctatus, Rodrig. Rev. Engenh. iii. n. 9; et l. c.—Bras.

CRYPTOPHRAGMIA, Benth. & Hook. f. Gen. ii. 1099 (1876) – seq.

CRYPTOPHRAGMIUM, Nees, in Wall. Pl. As. Rar. iii. 76 et 94 (1832) = Gymnostachyum, Nees (Acanth.).
axillare, Nees, in DC. Prod. xi. 96 = G. sanguinolentum.
canescens, Nees, l. c. 95 = G. canescens.
corifolium, Nees, in Wall. Pl. As. Rar. iii. 100 = Andrographis elongata.
elongatum, Nees, in DC. Prod. xi. 95 = Andrographis elongata.
glabrum, Dalz. in Hook. Kew Journ. ii. (1850) 338 = G. glabrum.
latifolium, Dalz. l. c. 137 = G. latifolium.
sanguinolentum, Nees, in DC. Prod. xi. 96 = G. sanguinolentum.
serrulatum, Nees, l. c. 96 = G. canescens.
venustum, Nees, in Wall. Pl. As. Rar. iii. 180 = G. venustum.

CRYPTOPLEURA, Nutt. in Trans. Am. Phil. Soc. Sér. II. vii. (1841) 431 = Troximon, Nutt. (Compos.).
californica, Nutt. l. c. = T. heterophyllum.

CRYPTOPUS, Lindl. Bot. Reg. sub t. 817 (1824). *ORCHIDEÆ*, Benth. & Hook. f. iii. 533.
BICLARDIA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 69.
elatus, *Lindl. Bot. Reg.* sub t. 817.—Madag.

CRYPTOPYRUM, Heynh. Nom. ii. 174 (1846) = Triticum, Linn. (Gramin.).
Richardsonii, Heynh. l. c. = T. violaceum.

CRYPTOSACCUS, Reichb. f. Xenia Orch. i. (1858) 15, in syn. = Leiochilus, Knowles & Westc. (Orchid.).

CRYPTOSANUS, Schaldw. in Otto & Dietr. Allg. Gartenz. xi. (1843) 101 = Leiochilus, Knowles & Westc. (Orchid.).
scriptus, Schaldw. l. c. = L. scriptus.

CRYPTOSEMA, Meisn. in Lehm. Pl. Preiss. ii. 207 (1847) = Jansenia, Kze. Pp. (Legumin.).
fasciculata, Meisn. l. c. = J. formosa.

CRYPTOSEPALUM, Benth. in Benth. & Hook. f. Gen. i. 684 (1867). *LEGUMINOSÆ*, Benth. & Hook. f. i. p.
marcvinense, *Oliver, Fl. Trop. Afr.* ii. 304.—Afr. trop.
microcladum, *Welw. ex Oliver, l. c.* 305.—Afr. trop.
tetraphyllum, *Benth. in Trans. Linn. Soc.* xxv. (1866) 535. t. 45 B.—Afr. trop.

CRYPTOSPERMUM, Stend. Nom. ed. II. i. 460, in syn. (1840), sphalm. = CYRTOSPERMUM, Rafin. = Cryptotænia, DC. (Umbell.).

CRYPTOSPERMUM, Yoong, in Trans. Linn. Soc. iii. (1797) 80 = Operculaëria, Gaertn. (Rubiac.).
Yoongii, Pers. Syn. i. 123 = O. aspera.

CRYPTOSPORA, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 161. *CRUCIFERÆ*, Benth. & Hook. f. i. 101.
falcata, *Kar. & Kir. l. c.*—Soongaria.
Griffithiana, *Fenzl, in Tchihat. Asie Min. Bot.* i. 861.—Persia.
rostrata, Fenzl, l. c. = Pariatoria rostrata.

CRYPTOSTACHYS, Stend. in Flora, xxxiii. (1850) 229 nomen; Syn. Pl. Gram. 181 (1855) = Sporobolus, R. Br. (Gramin.).
vaginata, Stend. ll. cc. = S. vaginæflorus?

CRYPTOSTEGIA, R. Br. in Bot. Reg. t. 485 (1819). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 742.
grandiflora, *R. Br. l. c.*—Afr. trop. (Ind. or. enlt.).
madagascariensis, *Boj. Hort. Maurit.* 213.—Madag.

CRYPTOSTEMMA, R. Br. in Ait. Hort. Kew. ed. II. v. 141 (1813). *COMPOSITÆ*, Benth. & Hook. f. ii. 456.
ALLOLOZONIUM, Knaze, in Linnæa, xvii. (1843) 572.
CYNOTIS, Hoffmgg. Verz. Pfl. Nachtr. ii. (1826) 98.
MICROSTEPHIUM, Less. in Linnæa, vi. (1831) 92.
calendulaceum, *R. Br. in Ait. Hort. Kew.* ed. II. v. 141.—Afr. austr.
Forbesianum, *Harv. in Hook. & Sond. Fl. Cap.* iii. 468.—Afr. austr.
hypochondriacum, R. Br. in Ait. Hort. Kew. ed. II. v. 141 = calendulaceum.
niveum, *Nichols. Dict. Gard.* i. 404.—Afr. austr.
runcinatum, R. Br. in Ait. Hort. Kew. ed. II. v. 141 = calendulaceum.

CRYPTOSTEMON, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 114 = Darwinia, Rudge (Myrtac.).
ericæus, F. Muell. ex Miq. l. c. 115 = D. fascicularis.
fascicularis, F. Muell. ex Miq. l. c. = D. fascicularis.

CRYPTOSTEPHANE, Sch. Bip. in Flora, xxvii. (1844) 769 = Diooma, Cass. (Compos.).
Kraussii, Sch. Bip. l. c. = D. dracanthoides.

CRYPTOSTEPHANUS, Welw. ex Baker, in Journ. Bot. xvi. (1878) 198. t. 197. *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 718.
densiflorus, *Welw. ex Baker, l. c.*—Afr. trop.

CRYPTOSTOMA, D. Dietr. Syn. Pl. i. 561, 625 (1839) = seq.

CRYPTOSTOMUM, Schreb. Gen. 137 (1789) = Moutabea, Aubl. (Polygal.).
coriaceum, Vahl, Eclog. Am. iii. 4 = M. guianensis.
guianense, Raeusch. Nom. ed. III. 81 = M. guianensis.
lanceifolium, Willd. Sp. Pl. i. 1061 = M. guianensis.

CRYPTOSTYLIS, R. Br. Prod. 317 (1810). *ORCHIDEÆ*, Benth. & Hook. f. iii. 506.
ACRONIA, Presl, Rel. Haenk. i. 103 (1827); Symb. Bot. ii. 9 (1858).
ZOSTEROSTYLIS, Blume, Bijdr. 418. t. 22 (1825).
albemifolia, *F. Muell. in Wing. South Sc. Record*, i. (1881) 172.—Ins. Samoa.
Arachnites, *Blume, Orch. Archip. Ind.* 133.—Malaya.
erecta, *R. Br. Prod.* 317.—Austral.
filiformis, *Blume, Orch. Archip. Ind.* 134.—Malaya.
leptochila, *F. Muell. ex Benth. Fl. Austral.* vi. 264.—Austral.
longifolia, *R. Br. Prod.* 317.—Austral.
ovata, *R. Br. l. c.*—Austral.
subulata, Reichb. f. Bot. Syst. Pfl. 15 = longifolia.
Walkeræ, *Blume, Orch. Archip. Ind.* 135.—Zeylan.
zeylanica, *Blume, l. c.*—Zeylan.

CRYPTOTÆNIA, DC. Coll. Mém. v. 49 (1829); et Prod. iv. 116 (1830). *UMBELLIFERÆ*, Benth. & Hook. f. i. 896.
ALACOSPERMUM, Neck. Elem. i. 197 (1790).
CYNOSCIADIUM, Rafin. ex DC. Prod. iv. 255, in syn. (1830).

CRYPTOTÆNIA.—
DERINGA, Adans. Fam. ii. 496 (1763).
MESODISCUS, Rafin. New Fl. Am. iv. 19 (1836).
canadensis, *DC. Coll. Mém.* v. 49; *et Prod.* iv. 119.—Am. bor.
elegans, Webb, ex C. Bolle, Ind. Sem. Hort. Berol. App. (1861) 9 (Asthirici sp. ?).—Ins. Canar.
japonica, Hassk. in Retzia, i. 118 = canadensis.
Thomasii, DC. Prod. iv. 119 = Pimpinella Thomasii.

CRYPTOTHECA, Blume, Bijdr. 1138 (1826) = Amomum, Linn.
spatula, Blume, l. c. 1139 = A. bacciferum.
dichotoma, Blume, l. c. = A. multiflorum.

CRYPTOTONIA, Tausch, in Flora, xvii. (1834) t. 348, sphalm. = Cryptotæenia, DC. (Umbell.).

CRYPTURUS, Link, in Linnæa, xvii. (1843) 887 = Lolium, Linn. (Gramin.).
loliaceus, Link, l. c. = L. leptochloides.

CRYSOPHILA, Benth. & Hook. f. Gen. iii. 998 (1883) = CRYSOPHILA, Blume = Copernicia, Mart. (Palm.).

CRYSTALLOPOLLEN, Steetz, in Peters, Reise Mossamb. Bot. 368 (1862) = Vernonia, Schreb. (Compos.).
angustifolium, Steetz, l. c. 368 = V. Steetziana.
latifolium, Steetz, l. c. 364 = V. Petersii.

CSZERNAEVIA, Endl. Gen. 777 (1839) = CZERNAEVIA, Turcz. = Archangelica, Hoffm. (Umbell.).

CTENANTHUS, Eichl. in Abh. Berl. Acad. (1884) 81. *SCITAMINEÆ*.
compressus, Eichl. l. c. 83 = Laschnathiana.
glabra, *Eichl. l. c.*—Am. trop.
Kunnanziana, *Eichl. l. c.* 84.—Am. trop.
Laschnathiana, *Eichl. l. c.* 83.—Am. trop.
pilosa, *Eichl. l. c.*—Am. trop.
setosa, *Eichl. l. c.* 84.—Am. trop.
Standleri, *Eichl. l. c.*—Am. trop.

CTENIUM, Panz. in Denkschr. Acad. Muench. 1813 (1814) 388. t. 18. *GRAMINEÆ*, Benth. & Hook. f. iii. 1165.
CAMPULOA, Desv. Journ. Bot. i. (1813) 69.
CAMPULOSUS, Desv. in Bull. Soc. Philom. ii. (1810) 189, nomen prius.
MONATHERA, Rafin. in Am. Monthly Mag. (1819) 190.
MONOCERA, Ell. Sketch, i. 126 (1821).
americanum, *Spreng. Syst.* i. 274 = carolinianum.
brachystachyum, *Kunth, Rév. Gram.* ii. 447.—Bras.
canescens, *Benth. in Hook. Niger Fl.* 506.—Afr. trop.
carolinianum, *Panz. in Denkschr. Acad. Muench.* 1813 (1814) 311.—Am. bor.
chapadense, *Doell, in Mart. Fl. Bras.* ii. 111. 78.—Bras.
cirrhosum, *Kunth, Rév. Gram.* ii. 445. t. 136.—Bras.
concinnum, *Nees, Fl. Afr. Austr.* i. 287.—Afr. austr.
digitatum, Spreng. Syst. i. 274 = Chloris digitata.
elegans, *Kunth, Rév. Gram.* i. 98.—Afr. trop.
indicum, Spreng. Syst. i. 274 = Chloris digitata.
nubicum, *In Nat. Ind. Sem. Hort. Gen.* (1857) 28; *et in Ann. Sc. Nat. Sér. III.* xix. (1853) 371.—Afr. trop.
nukuvinse, *Shrad. ex Yard. in Mém. Soc. Sc. Nat. Cherb.* v. (1857) 299, nomen.—Hab.?
planifolium, *Kunth, Enum. Pl.* i. 575.—Mexic.
polystachyum, *Balansa, in Bull. Soc. Bot. Fr.* xxxii. (1885) 264.—Paraguay.
rupestre, *F. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 140.—Ins. Cap. virid.
schollerae, Baker, Fl. Maurit. 463 = Enteropogon Seychellarum.
unyoniense, *Stend. Syn. Pl. Gram.* 362.—Afr. trop.

CTENODAUCUS, Fenzl, Nouv. Mém. Fl. Ad. 146 (1874) = Daucus, Tourn. (Umbellif.).
elegantus, Fenzl, l. c. = D. leucophyllodes.

CTENODON, Beauv. Adansonia, ix. (1870) 294. *LEGUMINOSÆ*.
Wahlbergianum, *Benth. l. c.* 297.—Bras.

CTENOLEPIS, Hook. f. in Benth. & Hook. f. Gen. i. 832 (1867) = **Blastania**, Kotschy & Peyr. (Cucurb.). *cornuformis*, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 630 = B. fimbristipula.
Garcini, Benth. & Hook. f. Gen. i. 832 = B. Garcini.
stipulacea, Benth. & Hook. f. l. c. = B. Garcini.

CTENOLOPHON, Oliver, in Trans. Linn. Soc. xxviii. (1878) 514. *OLACINEAE.*
grandifolius, *Oliver, l. c.* 517.—Malaya.
parvifolius, *Oliver, l. c.* 516.—Malaya.

CTENOMERIA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 29 = **Tragia**, Linn. (Euphorb.).
cordata, Harv. l. c. = T. capensis.
Kraussiana, Hochst. in Flora, xxviii. (1845) 85 = T. capensis.

CTENOPHIS, De Notar. Ind. Sem. Hort. Gen. (1847) 995. *GRAMINEAE*, Benth. & Hook. f. iii. 1171.
pectinella, *De Notar. l. c.*—Afr. trop.

CTENOPSIS, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 19 = **Blastania**, Kotschy & Peyr. (Cucurb.).
cornuformis, Naud. l. c. 13 = B. fimbristipula.
Garcini, Naud. l. c. = B. Garcini.

CTENOSPERMA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 118 = **Cotula**, Tourn. (Compos.).
alpinum, Hook. f. l. c. = Cot. alpina.

CTENOSPERMA, F. Muell. *ex* Pfeiff. Nom. i. 926 (1874) = **Brachycome**, Cass. (Compos.).

CUBA, Soop. Introd. 300 (1777) = **Tachigalia**, Aubl. (Legumin.).

CUBAEA, Schreb. Gen. Pl. 278 (1789) = **Tachigalia**, Aubl. (Legumin.).
bijuga, Spreng. Neue Entdeck. i. 290 = Leptolobium bijugum.
emarginata, Spreng. Syst. ii. 343 = T. emarginata.
paniculata, J. F. Gmel. Syst. 671 = T. paniculata.
speciosa, Spreng. Syst. ii. 345 = T. floribunda.
trigona, J. F. Gmel. Syst. 671 = T. trigona.

CUBEBA, Rafin. Sylva Tellur. 84 (1838); Miq. Comm. Phyt. 33. t. 1 ad 3 (1840) = **Piper**, Linn.
acris, Miq. l. c. 34 = P. acre.
albido-punctata, Miq. l. c. = P. albido-punctatum.
arborescens, Miq. l. c. = Thottea dependens.
arcuata, Miq. l. c. = P. arcuatum.
auriculata, Miq. l. c. = P. ribesioides.
bantamensis, Miq. l. c. 33 = P. bantamensis.
borbonense, Miq. Syst. Pip. 301 = P. borbonense.
canina, Miq. Comm. Phyt. 33. t. 3 = P. caninum.
capensis, Miq. Syst. Pip. 308.—Afr. austr.
celebica, Miq. Comm. Phyt. 33 = P. celebicum.
Chaba, Miq. l. c. 34 = P. Chaba, *Hunt.*
Clusii, Miq. Syst. Pip. 304 = P. Clusii.
coriacea, Miq. Comm. Phyt. 34 = Peperomia coriacea.
costulata, Miq. Syst. Pip. 299 = P. borbonense.
crassipes, Miq. Comm. Phyt. 45 = P. crassipes.
hederacea, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 435 = P. hederaceum.
lanata, Miq. Syst. Pip. 298 = P. caninum.
Lowongi, Miq. Comm. Phyt. 33 = P. Lowong.
majuscula, Miq. l. c. 34 = P. majusculum.
Meismini, Miq. l. c. = P. Malamiri.
minutula, Miq. l. c. 33 = P. minutum.
mollissima, Miq. in Zoll. Syst. Verz. Ind. Archip. 84 = P. mollissimum.
Nessii, Miq. Syst. Pip. 293 = P. pedicellosum.
nigrescens, Miq. Comm. Phyt. 35 = P. nigrescens.
officinalis, Rafin. Sylva Tellur. 84; Miq. Comm. Phyt. 33. t. 1 = P. Cubeba.
phyllostoba, Miq. Fl. Jungh. 294 = P. phyllostachyum.
piperita, Rafin. Sylva Tellur. 165 = P. Cubeba ?
rugulum, Miq. Syst. Pip. 304.—Borneo.
scribae, Miq. Comm. Phyt. 34 = P. Betle.
subcaba, Miq. l. c. 33 = P. subcatum.
sumatranum, Miq. l. c. 33. t. 5. f. A = P. sumatranum.
venosa, Miq. Fl. Jungh. 296 = P. venosum.
Wallichii, Miq. Syst. Pip. 309 = P. ribesioides.

CUBELIUM, Rafin. Cat. 13 (1836) = **Zenkiltrum**, Vent. (Violac.).
connatum, Rafin. l. c. = I. connatum.
parviflorum, Rafin. l. c. = I. glutinosum.

CUBILLA, Blume, Rumphia, iii. 100 (1849). *SAPINDACEAE*, Benth. & Hook. f. i. 400.
Blancoi, *Blume, l. c.* 101.—Ins. Philipp.
Rumphii, *Blume, l. c.*—Ins. Celeben.

CUBOSPERMUM, Lour. Fl. Cochinch. 275 (1790) = **Jussiaea**, Linn. (Onagrar.).
palustre, Lour. l. c. = J. repens.

CUCIFERA, Delile, Fl. Egypte, 145 (1813) = **Hyphaene**, Gaertn. (Palm.).
thebaica, Delile, l. c. = H. Cucifera.

CUCUBALUS, [Tourn.] Linn. Syst. ed. I (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 147.
LYCHNANTHOS, S. G. Gmel. in Nov. Comm. Acad. Petrop. xiv. (1770) 525.
LYCHNANTHUS, C. C. Gmel. Fl. Bad. ii. 249 (1806).
acaulis, Linn. Sp. Pl. 415 = Silene acaulis.
aegyptiacus, Linn. l. c. = Silene aegyptiaca.
alpestris, All. Auct. Fl. Pedem. 29 = Silene Campanula.
alpinus, Lam. Encyc. ii. 200 = Silene Cucubalus.
angustifolius, Mill. Gard. Dict. ed. VIII. n. 3 = Silene Cucubalus.
angustissimus, Nocca & Balb. Fl. Ticin. i. 199. t. 7 = Silene Cucubalus.
antelopum, Vest, in Flora, iv. (1821) 149 = Silene Cucubalus.
astrachanensis, *S. G. Gmel. Reise*, ii. 107 (quid ?).—Reg. Casp.
baccatus, Gueldenst. ex Ledeb. Fl. Ross. i. 333 = seq. baccifer, *Linn. Sp. Pl.* 414.—Europ.; As. bor.
Behen, Linn. l. c. = Silene Cucubalus.
caespitosus, Desf. Tabl. ed. I. 169 = Silene spergulaefolia.
caryophylloides, Poir. Encyc. Suppl. ii. 416 = Silene caryophylloides.
cathaticus, Ficin. Fl. Dresd. i. 318 = Silene italica.
catholicus, Linn. Sp. Pl. 415 = Silene catholica.
chlorafolius, Poir. Encyc. Suppl. ii. 416 = Silene perfoliata.
chloranthus, Willd. Prod. Fl. Berl. 155 = Silene chlorantha.
clowacarpus, Schult. Obs. Bot. 81 = Silene italica.
congestus, Willd. ex Cham. & Schlecht. in Linnaea, i. (1826) 89 = Silene repens.
conicus, Lam. Fl. Fr. iii. 33 = Silene conica.
convoideus, Lam. l. c. 27 = Silene convoidea.
cyclamineus, Sieber, Reise Kreta, ii. 324 = Silene variegata.
Czerri, Stend. Nom. ed. II. i. 451 = Silene Fabaria.
dauricus, Pall. ex Bunge, Verz. Alt. 31 = Silene tenuis.
dichotomus, Lam. Fl. Fr. iii. 32 = Silene Muscipula.
dioicus, Gilib. Fl. Lituan. ii. 167 = Silene Otites.
divaricatus, [Clairv.] Man. Herb. 145 = baccifer.
Douglasii, Eaton, Man. ed. VII; ex ej. ed. VIII. 427 = Silene Douglasii.
dubernus, Mill. Gard. Dict. ed VIII. n. 6 = Silene viscosa.
effusus, Fisch. ex DC. Prod. i. 376 = Silene Otites.
fabarius, Linn. Sp. Pl. 414 = Silene Fabaria.
fabarius, Sieber, Reise Kreta, ii. 324 = Silene variegata.
fabarius, Thore, Chlor. Land. 173 = Silene Thorei.
fasciculatus, Lam. Fl. Fr. iii. 27 = Silene Armeria.
fimbriatus, *Balb. ex Mém. Acad. Turin*, xviii. (1811) 356. t. 6.—Hab. ?
fimbriatus, Bieb. Fl. Taur. Cauc. i. 333 = Silene fimbriata.
flexuosus, Gueldenst. ex Ledeb. Fl. Ross. i. 295 (quid ?). —Rossia.
foecundus, Ficin. Fl. Dresd. i. 313 = Silene italica.
fruticosus, Pall. Reise, ii. 739 = Silene sinaica.
fruticulosus, J. F. Gmel. Syst. 718 = Silene turgida.
glaucus, Spreng. Nachr. i. Bot. Gart. Halle, 19 = Silene glauca.
glutinosus, Retz. Obs. ii. 18 = Silene catholica.
hermaphroditus, Gilib. Fl. Lituan. ii. 167 = Silene Otites.
horizontalis, Moench, Meth. 713 = baccifer.
inapertus, Lam. Fl. Fr. iii. 31 = Silene inaperta.
inflatus, Salisb. Prod. 302 = Silene Cucubalus.
italicus, Linn. Syst. ed. X. 1080 = Silene italica.
lacerus, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 262 = Silene fimbriata.
latifolius, Mill. Gard. Dict. ed. VIII. n. 2 = Silene Cucubalus.
littoralis, Fers. Syn. i. 496 = Silene maritima.
littoralis, Vis. Strip. Dalm. 32 = Silene Cucubalus.

CUCUBALUS—
longicilius, Brot. Fl. Lusit. ii. 180 = Silene nutans.
macrophyllus, Hort. ex Rohrb. Monog. 81 = Silene Fabaria.
marginatus, Schult. Oestr. Fl. ed II. i. 674 = Silene Cucubalus.
maritimus, Lam. Encyc. ii. 220 = Silene maritima.
mollissimus, Bieb. Fl. Taur. Cauc. i. 324 = Silene Marschallii.
mollissimus, Linn. Sp. Pl. ed. II. 593 = Silene mollissima.
mollissimus, Waldst. & Kit. Pl. Rar. Hung. iii. 277 = Silene italica.
montanus, Vest, in Flora, iv. (1821) 149 = Silene Cucubalus.
multifidus, Adam, ex Weber & Mohr, Beitr. i. 57 = Silene fimbriata.
multiflorus, Ehrh. Beitr. vii. 141 = Silene multiflora.
muscosus, Lam. Fl. Fr. iii. 30 = Silene acaulis.
niveus, Nutt. Gen. Am. i. 287 = Silene nivea.
noctiflorus, Mill. Gard. Dict. ed. VIII. n. 8 = Silene noctiflora
nutans, Lam. Fl. Fr. iii. 35 = Silene nutans.
Otites, Linn. Sp. Pl. 415 = Silene Otites.
paniculatus, *Mill. Gard. Dict.* ed. VIII. n. 12 — Hispan.; Ital.
pannonicus, *Wulf. Carol.* 141 = Silene ovata.
polyphyllus, Hort. Gorenk. ed. II. 60, nomen = Silene procumbens.
procumbens, *Y. F. Gmel. Syst.* 712.—Hab. ?
Pumilio, Linn. Mant. i. 73 = Silene Pumilio.
quadrifidus, Linn. Sp. Pl. 415 = Silene quadridentata.
quadrifidus, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 29 = Silene nutans.
reflexus, Linn. Sp. Pl. 416 = Silene reflexa.
roseus, *Desf. Tabl.* ed. I. 159, nomen.—Hab. ?
rotundifolius, Stend. Nom. ed. I. 243 = Silene inflata.
Royeni, Bieb. Casp. 174 = Silene viscosa.
rubens, Roth, Enum. Pl. Phaen. Germ. i. 11. 286 = Silene nutans.
saponariaefolius, Moench, ex Steud. Nom. ed. II. i. 451; Bonata, ex eund. ii. 587 = Silene Cucubalus.
saxatilis, Lam. Fl. Fr. iii. 30 = Silene rupestris.
saxifragus, Lam. l. c. 29 = Silene Saxifraga.
saxifragus, Linn. Mant. i. 71 = Silene dianthoides.
saxifragus, Schang. ex Bunge, Verz. Alt. 32 = Silene turgida.
scabor, Fisch. ex Kegel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) ii. 547 = Silene Cucubalus.
secundus, Gilib. Fl. Lituan. ii. 168 = Silene tatarica.
sibiricus, Linn. Syst. ed X. 1081 = Silene sibirica.
sibiricus, Willd. ex Walp. Rep. ii. 777 = Silene tenuis.
silenoides, Vill. Hist. Pl. Dauph. iii. 614 = Silene italica.
spergulaefolius, Willd. Sp. Pl. ii. 690 = Silene spergulaefolia.
spicatus, Lam. Fl. Fr. iii. 34 = Silene nocturna.
staticaefolius, Poir. Encyc. Suppl. ii. 416 = Silene tenuis.
stellatus, Linn. Sp. Pl. 414 = Silene stellata.
striatus, Willd. ex Walp. Rep. ii. 777 = Silene paucifora.
strictus, *Roth, in Usteri, Ann. Bot.* x (1794) 44.—Hab. ?
strictus, Willd. ex Ledeb. Fl. Ross. i. 306 = Silene paucifora.
sylvestris, Lam. Fl. Fr. iii. 26 = Silene gallica.
tataricus, Linn. Sp. Pl. 415 = Silene tatarica.
variegatus, Lam. Fl. Fr. iii. 28 = Silene gallica.
venosus, Gilib. Fl. Lituan. ii. 165 = Silene Cucubalus.
viridis, Lam. Encyc. ii. 221 = Silene Cucubalus.
viscidus, Krock. Fl. Sil. ii. 1. 40 = seq.
viscosus, Linn. Sp. Pl. 414 = Silene viscosa.
wolgensis, Willd. Enum. Hort. Berol. Suppl. 24 = Silene Otites.

CUCULARIA, Rafin. in Med. Repos. N. York, v. (1808) 252 = **Dicentra**, Bernh. (Papaverac.).
bulbosa, Rafin. l. c. = D. Cucularia.

CUCULINA, Rafin. Fl. Tellur. iv. 49 (1836) = **Catastetum**, Rich. (Orchid.).

CUCULLARIA, Endl. Gen. 859 (1839) = Cucularia, Rafin.

CUCULLARIA, Schreb. Gen. 6 (1789) = Vochysia, Juss.
alpestris, Spreng. Syst. iv. Cur. Post. 9 = V. alpestris.
citrifolia, Schult. Mant. i. 53 = V. citrifolia.
elliptica, Spreng. Syst. iv. Cur. Post. 9 = V. elliptica.
emarginata, Vahl, Enum. i. 5 = V. emarginata.
excelsa, Vahl, l. c. 4 = V. ferruginea.
excelsa, Willd. Sp. Pl. i. 17 = V. guianensis.
ferruginea, Spreng. Syst. iv. Cur. Post. 9 = V. ferruginea.
floribunda, Spreng. l. c. = V. floribunda.
grandis, Spreng. l. c. = V. grandis.
Haenkeana, Spreng. l. c. = V. Haenkeana.
latea, Willd. ex Steud. Nom. ed. II. i. 451 = V. emarginata.
pyramidalis, Spreng. Syst. iv. Cur. Post. 9 = V. pyramidalis.
racemosa, Roem. & Schult. Syst. i. 86 = V. racemosa.
rotundifolia, Spreng. Syst. iv. Cur. Post. 9 = V. rotundifolia.
rufa, Spreng. l. c. = V. rufa.
tetraphylla, G. F. W. Mey. Prim. Fl. Esseq. 13 = V. tetraphylla.
tomentosa, G. F. W. Mey. l. c. 13 = V. ferruginea.
tucanorum, Spreng. Syst. iv. Cur. Post. 9 = V. tucanosum.
Vochy, J. F. Gmel. Syst. 10 = V. guianensis.

CUCULLIFERA, Nees, in Lindl. Introd. Nat. Syst. ed. II. 451 (1836) = Cannamois, Beauv. (Restinc.).
dura, Nees, l. c. = Cannamois simplex.

CUCULLIGERA, Mast. in Journ. Linn. Soc. x. (1869) 238 = prsec.

CUCUMEROIDES, Gaertn. Fruct. ii. 485, t. 180, t. 4 = Trichosanthes, Linn. (Cucurb.).

CUCUMEROPSIS, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 60. *CUCURBITACEÆ*, Benth. & Hook. f. i. 820.
CLADOSICYOS, Hook. f. in Oliver, Fl. Trop. Afr. ii. 584 (1871).
CORYNOSICYOS, F. Muell. Sel. Pl. Indust. Cult. 59 (1876).
edulis, *Cogn. in DC. Monog. Phan.* iii. 518.—Afr. trop.
Mannii, *Naud. in Ann. Sc. Nat.* Sér. V. v. (1866) 30. —Afr. trop.

CUCUMIS, [Tourn.] Linn. Syst. ed. I (1735). *CUCURBITACEÆ*, Benth. & Hook. f. i. 826.
MELO, Adans. Fam. ii. 138 (1763).
RIGOCARPUS, Neck. Elem. i. 228 (1790).
abyssinicus, A. Rich. Tent. Fl. Abyss. i. 294 = C. Figarei.
abyssinicus, Schimp. ex Hook. f. in Oliver, Fl. Trop. Afr. ii. 544 = pustulatus.
acidus, Jacq. Obs. Bot. iv. 14 = C. Melo.
acutanguius, Descourt. Fl. Antill. v. 94, t. 338 = Sechium edule.
acutangulus, Linn. Sp. Pl. 1011 = Luffa acutangula.
acutangulus, Wall. Cat. n. 6736 A = Luffa aegyptiaca.
africanus, Lindl. Bot. Reg. t. 980 = Momordica Charantia.
africanus, *Lindl. f. Suppl.* 423.—Afr. austr.
agrestis, Mill, ex Cogn. in DC. Monog. Phan. iii. 489 = Momordica Balsamina.
agrestis, Reichb. Fl. Germ. Excurs. 294 = Ecballium agreste.
amarissimus, Schrad. Ind. Sem. Hort. Gotting. (1827) = Citrullus vulgaris.
amarus, Stocks, ex *T.* Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 19 = propheturum.
ambigua, Fenzl, ex Hook. f. in Oliver, Fl. Trop. Afr. ii. 543 = dipsaceus, C. Melo.
angolensis, *Hook. f. ex Cogn. in DC. Monog. Phan.* iii. 487.—Afr. trop.
anguinus, T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 19, sphalm. = propheturum.
anguinus, Linn. Sp. Pl. ed. II. 1487 = Trichosanthes Anguina.
angulatus, *Forsk. Fl. Aegypt. Arab.* 168.—Arabia.
Angurie, Forsk. l. c. = propheturum.
Anguria, *Linn. Sp. Pl.* 1011.—Am. trop.

CUCUMIS:—
Anguria, Kodsch. Obs. 56 = Luffa operculata.
angurioides, M. Roem. Syn. Pepon. 79 = C. Anguria.
arabicus, Delile, ex Naud. in Ann. Sc. Nat. Sér. IV. xi. (1859) 14 = propheturum.
arenarius, Schrad. in Linnæa, xii. (1838) 416 = africanus.
arenarius, Schum. & Thonn. Beskr. Guin. Pl. 326 = C. Melo.
aroussicus, Royle, Illustr. Bot. Himal. ii. 220 = C. Melo.
barbatus, *Roth, Catalect.* fasc. ii. 122.—Hab.?
Bardana, Fenzl, ex Naud. in Ann. Sc. Nat. Sér. IV. xi. (1859) 35 = dipsaceus.
bicirrha, Forst. ex Guill. l. c. Sér. II. vii. (1837) 350 = Lagenaria vulgaris.
bipinnatifidus, Wight, ex Naud. l. c. xii. (1859) 99 = Citrullus Colocynthis.
campechianus, H. B. & K. Nov. Gen. et Sp. ii. 123 = C. Melo.
cantalupensis, Haberle, ex M. Roem. Syn. Pepon. 69 = C. Melo.
Cantalupo, Reichb. Fl. Germ. Excurs. 295 = C. Melo.
Chate, Wall. Cat. n. 6727 = C. Melo.
Chate, Hasselq. Iter Palaest. 491; Linn. Syst. ed. X. 1279 = C. Melo.
Chito, Morr. in Ann. Soc. Gand, v. (1849) 341, t. 278 = C. Melo.
chrysocarpa, *Welw. Apont.* 589, n. 62.—Afr. trop.
chrysocomus, Schum. & Thonn. Beskr. Guin. Pl. 327 = C. Melo.
cicatrisatus, Stocks, in Hook. Kew Journ. iv. (1852) 148 = C. Melo.
Citrullus, Ser. in DC. Prod. iii. 301 = Citrullus vulgaris.
cognata, *Fenzl, ex Hook. f. in Oliver, Fl. Trop. Afr.* ii. 543; *Cogn. in DC. Monog. Phan.* iii. 487.—Afr. trop.
Colocynthis, Linn. Sp. Pl. 1011 = Citrullus Colocynthis.
Colocynthis, Thunb. Prod. Pl. Cap. 13 = Citrullus vulgaris.
Conomum, Thunb. Fl. Jap. 324 = C. Melo.
cordifolius, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. ii. 492 = Momordica foetida.
cubensis, Schrad. in Linnæa, xii. (1838) 419 = C. Melo.
deliciosus, Salisb. Prod. 157; Roth, Catalect. fasc. iii. 15 = C. Melo.
dipsaceus, *Ehrenb. ex Spach, Hist. Veg. Phan.* vi. 211. —Arab.; Afr. trop.
dipsaceus, Wender. ex Steud. Nom. ed. II. i. 451 = africanus.
dissectifolius, *Naud. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 23.—Afr. austr.
dissectifolius, Naud. l. c. partim = Citrullus Naudinianus.
dissectus, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 449 = Citrullus vulgaris.
Dudaim, Linn. Sp. Pl. 1011 = C. Melo.
echinatus, Moench, Meth. 654 = C. Anguria.
edulis, Steud. Nom. ed. II. i. 451 = C. Citrullus.
ericarpus, Boiss. & Noë, Diagn. Ser. II. ii. 59 = C. Melo.
erinaceus, Hort. ex Steud. Nom. ed. II. i. 451 = C. Melo.
excisilobus, Salisb. Prod. 157 = sativus.
facifolius, A. Rich. Tent. Fl. Abyss. i. 294, t. 58 = Figarei.
Figarei, *Delile, Cat. ex Naud. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 16.—Afr. trop.
flexuosus, Linn. Sp. Pl. ed. II. 1437 = C. Melo.
foetidissimus, Hamad. Biol. Centr. Am. Bot. i. 483 = Cucurbita foetidissima.
fortidus, Salisb. Prod. 156 = propheturum.
glaber, Walt. Fl. Carol. 240 = Melothria pendula.
grossivalvulus, Hort. ex Steud. Nom. ed. II. i. 451 = myriocarpus.
Gurmia, Wall. Cat. n. 6736 = C. Melo.
Hardwickii, Royle, Illustr. Bot. Himal. 220, t. 47 = sativus.
heptadactylus, *Naud. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 34.—Afr. austr.
Hildebrandtii, Hurk. Madr. ex Wall. Cat. sub n. 6754 = Momordica tuberosa.
hirsutus, Roxb. f. in Oliver, Fl. Trop. Afr. ii. 546, pro parte = Welwitschii.
hirsutus, *Sond. in Harv. & Sond. Fl. Cap.* ii. 497.—Afr. austr.

CUCUMIS:—
Hookeri, *Naud. in Gard. Chron.* (1870) 1508.—Afr. trop.
integrifolius, Roxb. Hort. Beng. 70; Fl. Ind. iii. 734 = Trichosanthes integrifolia.
intermedius, M. Roem. Syn. Pepon. 80 = Momordica Charantia.
jamaicensis, Bert. ex Spreng. Syst. iii. 46 = C. Melo.
jucunda, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 46 = C. Melo.
lacinieus, Eckl. ex Steud. Nom. ed. II. i. 451 = Citrullus vulgaris.
lagenarius, Dum. Fl. Belg. 54 = Lagenaria vulgaris.
lanatus, Thunb. in Journ. Hist. Nat. Par. ii. (1779) 351, t. 27 = Luffa aegyptiaca.
longipes, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 547.—Afr. trop.
lusonicus, Blanco, Fl. Filip. ed. I. 861 = Melothria indica.
macrocarpos, Wender. ex Mart. Reise Bras. i. 536 = Cucurbita Pepo.
maculatus, Willd. Sp. Pl. iv. 614 = C. Melo.
maderaspatanus, Linn. Sp. Pl. 1012 = Melothria maderaspatana.
maderaspatanus, Soland. ex Seem. in Journ. Bot. ii. (1864) 47 = Melothria Gruyana.
madraspatanus, Roxb. Hort. Beng. 70; Fl. Ind. iii. 732 = C. Melo.
magacorpus, G. Don, Gen. Syst. iii. 28 = Luffa aegyptiaca.
Melo, Blanco, Fl. Filip. ed. I. 775 = trigonus.
Melo, *Linn. Sp. Pl.* 1011.—Geront. trop.
melonitrigonus x. *Naud. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 171.
membranifolius, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 545.—Afr. trop.
metuliferus, *E. Mey. ex Schrad. in Linnæa*, xii. (1838) 406, nomen; *et Naud. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 10.—Afr. austr.
missionis, Wall. Cat. n. 6728 = Trichosanthes cucumerina.
Momordica, Roxb. Hort. Beng. 70; Fl. Ind. iii. 730 = C. Melo.
moschatus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 552 = C. Melo.
Muelleri, Naud. in Ann. Sc. Nat. Sér. IV. xi. (1859) 84 = Melothria Muelleri.
muricatus, Willd. Sp. Pl. iv. 613 = sativus.
myriocarpus, *Naud. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 22.—Afr. austr.
Naudinianus, Sond. in Harv. & Sond. Fl. Cap. ii. 496 = dissectifolius, Citrullus Naudinianus.
odoratissimus, Moench, Meth. 654 = C. Melo.
officinarum-malo, Crants, Inst. t. 173 = C. Melo?
operculatus, Roxb. ex Wight & Arn. Prod. 348 = Luffa acutangula.
Panchorianus, Naud. in Ann. Sc. Nat. Sér. IV. xii. (1859) 113, t. 8 = C. Melo.
parviflorus, Salisb. Prod. 157 = C. Anguria.
Pavel, Kostel. in Allg. Med.-Pharm. Fl. ii. 788 = Cocciuea indica.
Pedata, Cranta, Inst. i. 173 = Anguria pedata?
pedatifidus, Schrad. in Linnæa, xii. (1838) 418 = C. Melo.
pedatus, Linn. Syst. ed. X. 1279 = Anguria pedata.
pentandrus, Roxb. ex Wight & Arn. Prod. 348 = Luffa aegyptiaca.
Pepo, Dum. Fl. Belg. 54 = Cucurbita Pepo.
perennis, E. James, Exped. Rock. Mount. ii. 346 = Cucurbita foetidissima.
persicifolius, Norohaa, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 11 = Melothria perpusilla.
persicoides, Seiz. in Verh. Ver. Bef. Gastenb. (1837) 378 = C. Melo.
peruvius, M. Roem. Syn. Pepon. 71 = C. Melo.
pleiocarpus, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 46 = C. Melo.
pictus, Jacq. Hort. Vindob. iii. 17 = C. Melo.
Priscaga, Wender. Ind. Sem. Hort. Marb. (1849); Linnæa, xv. (1841) Lit. 100 = C. Melo.
prophetarum, Jacq. Hort. Vindob. t. 9 = myriocarpus.
propheturum, *Linn. Cent. i.* 32; *Amœn. Acad. iv.* 296. —Arab.; Afr. trop.
prophetarum, Meyer, ex Cogn. in DC. Monog. Phan. iii. 489 = dissectifolius.
prophetarum, Wall. Cat. n. 6735 = trigonus.
Pseudocolocynthis, Royle, Illustr. Bot. Himal. 220, t. 47, f. 2 = trigonus.

CUCUMIS :—
Pseudocolocynthis, Wender. Sel. Sem. Hort. Marb. (1824) ; Linnæa, x. (1836) Litt. 77 = Citrullus Colocynthis.
pubescens, Willd. Sp. Pl. iv. 614 = C. Melo.
purpurea, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 12.—Malaya.
pustulata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 544.—Afr. trop.
pyriformis, Roxb. ex Wight & Arn. Prod. 342 = trigonus.
Rapalitio, Carr. in Rev. Hortic. (1872) 400 = Cucurbita maxima.
reflexus, Zeyh. ex Ser. in DC. Prod. iii. 300 = C. Melo.
Reginæ, Schrad. in Linnæa, xii. (1838) 419 = C. Melo.
Rheedii, Kostel. Allg. Med.-Pharm. Fl. ii. 788 = Coccinia indica.
rigidus, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 497.—Afr. austr.
sagittatus, *Wawra & Peyr. in Sitzb. Acad. Wien. Mat.-Nat.* xxxvii. (1860) 567.—Afr. trop.
sativus, *Linn. Sp. Pl.* 1012.—Ind. or.
Schraderianus, M. Roem. Syn. Pepon. 73 = C. Melo.
sepium, G. F. W. Mey. Prim. Fl. Esseq. 278 = Luffa operculata.
serotinus, Haberle, ex Seis. in Verh. Ver. Bef. Gartenb. (1857) 376 = C. Melo.
setosus, *Cogn. in DC. Monog. Phan.* iii. 491.—Ind. or. Sonderli, *Cogn. l. c.* 489.—Afr. austr.
striatus, A. Rich. Tent. Fl. Abyss. i. 295 = Coccinia adoënsis.
subsericeus, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 545.—Afr. trop.
Tinnæanus, Kotschy & Peyr. Pl. Tinn. 17. t. 8 = metuliferus.
trifoliatus, Linn. Syst. ed. X. 1279 = Anguria trifoliata.
trigonus, Benth. Fl. Austral. iii. 317 = C. Melo.
trigonus, *Roxb. Hort. Beng.* 70 ; *Fl. Ind.* iii. 722.—Ind. or.
trilobatus, Forsk. Fl. Aegypt. Arab. 168 = Figarei.
trilobatus, Linn. Syst. ed. X. 1279 = Anguria trilobata.
tuberosus, Roxb. ex Wight & Arn. Prod. 344 = Momordica tuberosa.
tuberosus, Roxb. ex Wight & Arn. l. c. 348 = Gymnopetalum cochinchinense.
turbinatus, Roxb. Hort. Beng. 70 ; Fl. Ind. iii. 723 = trigonus.
umbilicatus, Salisb. Prod. 157 = C. Melo.
utilissimus, Roxb. Hort. Beng. 70 ; Fl. Ind. iii. 721 = C. Melo.
verrucosus, Herb. Rottl. ex C. B. Clarke, in Fl. Brit. Ind. ii. 623 = Bryonopsis laciniosa.
villosus, Boiss. & Noë, Diagn. Ser. II. ii. 59 = C. Melo.
Wolwitschii, *Cogn. in DC. Monog. Phan.* iii. 490.—Afr. trop.
Zapallito, Carr. in Rev. Hortic. (1878) 298 = Cucurbita maxima.
Zapallo, Stend. Nom. ed. II. i. 452 = Cucurbita Pepo.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* ii. 496.—Afr. austr.

CUCURBITA, [Tourn.] Linn. Syst. ed. I (1735). *CUCURBITACEÆ*, Benth. & Hook. f. i. 838.
Ordovicus, Rafin. Atl. Journ. 140 (1839).
Pepo, [Tourn.] Linn. Syst. ed. I (1735).
Pilgocalyx, Gasparr. in Rendic. Acad. Sc. Nap. vi. (1847) 409.
Sphenantha, Schrad. in Linnæa, xii. (1838) 416.
Tristemon, Scheele, in Linnæa, xxi. (1848) 586.
acutangula, Blume, Bijdr. 982 = Luffa acutangula.
alba, Roxb. ex Wight & Arn. Prod. 344 = Benincasa cerifera.
Angurie, Duchesne, in Lam. Encyc. ii. 158 = Citrullus vulgaris.
aspera, Soland. ex Forst. f. Prod. 92 = Cucumis Melo.
asperata, Gill. in Hook. Bot. Misc iii. (1833) 324 = Cucurbitella asperata.
astrosperma, Schrad. in Linnæa, xii. (1838) 407, nomen = C. Pepo.
australis, Willd. Sp. Pl. iv. 607 = C. Pepo.
baccalaurea, *Ait. ex Steud. Nom.* ed. I. 244.—Hab. ?
cafira, Kotsl. & Zeyh. Enum. 270 = Citrullus vulgaris.
californica, *Torr. ex S. Wats. in Proc. Am. Acad.* xi. (1876) 133.—Am. bor. occ.
Cavanillesiana, Bosch.-Hamb. in Wall. Cat. n. 6716 = Benincasa cerifera.

CUCURBITA:—
campanulata, D. Dietr. Syn. Fl. v. 360 = Luffa umbellata.
cerasiformis, Haberle, ex Mart. Reise Bras. i. 536 = C. Pepo.
cerifera, Pisch. Cat. Hort. Gorenk, ex Savi, in Bibliot. Ital. ix. (1818) 158 = Benincasa cerifera.
cicuerata, *Molina, Sagg. Chil. ed. Germ.* 216.—Chili.
Citrullus, Linn. Sp. Pl. 1010 = Citrullus vulgaris.
Citrullus, M. Roem. Syn. Pepon. 84 = C. Pepo.
cladioïda, Nocca, Fl. Ven. v. 53 = C. Pepo.
coccinea, G. Don, Gen. Syst. iii. 41 = Physedra Barter.
Colocynthis, Link, Handb. i. 648 = Citrullus Colocynthis.
Courgero, Ser. in Mém. Soc. Phys. Genèv. iii. (1825) 11. t. 1 = C. Pepo.
cucullata, Bej. ex Hook. f. in Oliver, Fl. Trop. Afr. ii. 527 = Peponia Vogelii.
cucumifera, Schrad. in Linnæa, xii. (1838) 407, nomen = C. Pepo.
digitata, *A. Gray, Pl. Wright.* ii. 60.—N. Mexic.
dioica, Roxb. ex Wight & Arn. Prod. 347 = Coccinia indica.
elongata, Bean, ex Schrad. in Linnæa, xii. (1838) 407 = C. Pepo.
esculenta, S. F. Gray, Nat. Arr. Brit. Pl. ii. 552 = C. Pepo.
frodicarpa, Hassk. Hort. Bogor. Desc. i. 805 = Sicana odorifera.
exanthematica, Fenzl, ex A. Rich. Tent. Fl. Abyss. i. 296 = Coccinia Moghadd.
Farinæ, Moxz. ex Naud. in Ann. Sc. Nat. Sér. IV. vi. (1856) 17 = maxima.
farinosa, Blume, Bijdr. 981 = Benincasa cerifera.
fastuosa, Sallab. Prod. 157 = C. Pepo.
ficifolia, *Bouché, in Verh. Ver. Gartenb. Berl.* xii. (1837) 205.—As. or.
ficifolia, Wall. Cat. n. 6721 = Trichosanthes himalensis.
foetidissima, *H. B. & K. Nov. Gen. et Sp.* ii. 128.—Mexic.
Galeottii, *Cogn. in DC. Monog. Phan.* iii. 551.—Mexic.
gigantea, Sallab. Prod. 157 = C. Citrullus.
grisea, M. Roem. Syn. Pepon. 86 = C. Pepo.
hieroglyphica, *Rosenth. Syn. Pl. Diaph.* 883.—Hab. ?
hispida, Ser. Fl. Jard. et Cult. ii. 581 = moschata.
hispida, Thunb. Fl. Jap. 322 = Benincasa cerifera.
hybrida, Bertol. ex Naud. in Ann. Sc. Nat. Sér. IV. vi. (1856) 29 = C. Pepo.
idolatrica, Willd. Sp. Pl. iv. 607 = Lagenaria vulgaris.
Lagenaria, Linn. Sp. Pl. 1010 = Lagenaria vulgaris.
leucantha, Duchesne, in Lam. Encyc. ii. 150 = Lagenaria vulgaris.
lignosa, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. austr.
littoralis, Hassk. Cat. Hort. Bogor. Alt. 190 = Benincasa cerifera.
macrocarpa, Gasparr. in Rendic. Acad. Sc. Nap. vi. (1847) 447 = moschata.
maderensis, Molina, Sagg. Chil. 138 = C. Pepo.
mammosa, J. F. Gmel. Syst. 1051 = pracc.
marmiphormis, Haberle, ex M. Roem. Syn. Pepon. 92 = C. Pepo.
maxima, *Duchesne, in Lam. Encyc.* ii. 151.—Aa. trop. (Orb. trop. cult.).
maxima, Wall. Cat. n. 6720 = moschata.
medullaris, *Haberle, ex M. Roem. Syn. Pepon.* 93, nomen.—Hab. ?
melanosternis, *Carr. in Rev. Hortic.* (1880) 187.—Japon.
melanosperma, A. Braun, Cat. Hort. Karls (1854), nomen ; Gasparr. in Rendic. Acad. Sc. Nap. vi. (1847) 448 = Sicana odorifera.
Melanolibon, Linn. Sp. Pl. 1010 = C. Pepo.
Melopepo, Lour. Fl. Cochinch. 598 = moschata.
Melopepo, Wall. Cat. n. 6725 = Trichosanthes pubera.
microcarpa, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 17 = Melothria Muelleri.
moschata, *Duchesne, ex Poir. in Dict. Sc. Nat.* xi. (1818) 294.—Aa. trop.
multifora, Soland. ex Forst. f. Prod. 92, nomen = Luffa aegyptiaca.
oblonga, Link, Handb. i. 648 = C. Pepo.
odorifera, Vell. Fl. Flum. x. t. 99 = Sicana odorifera.
ovifera, Linn. Mant. i. 126 = C. Pepo.

CUCURBITA:—
palmata, *S. Wats. in Proc. Am. Acad.* xi. (1876) 137.—Calif.
Pepo, *Linn. Sp. Pl.* 1010.—Orient.; Afr. trop.
Pepo, Lour. Fl. Cochinch. 598 = Benincasa cerifera.
Pepo, Vell. Fl. Flum. x. t. 100 = Lagenaria vulgaris.
Pepo, Wall. Cat. n. 6722 = moschata.
perennis, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 198 = foetidissima.
perpusilla, Blume, Cat. Gew. Buitenz. 105 = Melothria perpusilla.
pileiformis, M. Roem. Syn. Pepon. 86 = maxima.
pinnatifida, Schrank, in Syll. Ratisb. i. (1824) 190 = Citrullus vulgaris.
polymorpha, Duchesne, in Lam. Encyc. ii. 152 = C. Pepo.
pomiformis, M. Roem. Syn. Pepon. 85 = C. Pepo.
prunosa, Soland. in Forst. f. Prod. 92 = Benincasa cerifera.
purpurea, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 11 (Quid ?).—Java.
pyriformis, Duchesne, ex Poir. in Dict. Sc. Nat. xi. (1818) 287 = C.Pepo.
pyriformis, M. Roem. Syn. Pepon. 89 = Lagenaria vulgaris.
pyxidaria, DC. Fl. Fr. iii. 698 = C. Pepo.
radicans, *Naud. in Ann. Sc. Nat.* Sér. V. vi. (1866) 8.—Mexic.
Rapalitio, Carr. in Rev. Hortic (1872) 399 = maxima.
sagittata, Klein, ex Steud. Nom. ed. II. i. 452 = Melothria heterophylla.
scabra, Blume, Cat. Gew. Buitenz. 105 = Melothria perpusilla.
Schimperiana, Hochst. ex Cogn. in DC. Monog. Phan. iii. 538 = Coccinia Moghadd.
siceraria, Molina, Sagg. Chile, 138 = Lagenaria vulgaris.
simplex, DC. ex Steud. Nom. ed. II. i. 452 = bononiensis.
spathularia, Schrad. in Linnæa, xii. (1838) 406, nomen = moschata.
subcrenata, Willd. Sp. Pl. iv. 606 = C. Pepo.
Succado, Naeg. ex Naud. in Ann. Sc. Nat. Sér. IV. vi. (1856) 29. t. 2 = C. Pepo.
Succedo, Am. in Hook. Journ. Bot. iii. (1841) 278 = pracc. ?
sulcata, Blanco, Fl. Filip. ed. I. 778 = maxima.
texana, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 193 = C. Pepo.
triangulata, Hochst. ex Cogn. in DC. Monog. Phan. iii. 533 = Coccinia Moghadd.
tuberculosa, Schrad. Ind. Sem. Hort. Gotting. (1833) ; Linnæa, x. (1835–36) Litt. 70 = C. Pepo.
turbaniformis, M. Roem. Syn. Pepon. 87 = maxima.
umbellata, Klein, ex Willd. Sp. Pl. iv. 608 ; Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 616 = Luffa umbellata.
umbellata, Wall. Cat. n. 6724 = Gymnopetalum Wightii.
urnigera, Schrad. in Linnæa, x. (1836) Litt. 110 = C. Pepo.
vacua, F. Muell. Fragm. vi. 186 = Benincasa cerifera.
variegata, Steud. Nom. ed. II. i. 452 = C. Pepo.
venosa, Descourt. Fl. Méd. Antill. v. 79. t. 292 = C. Pepo.
verrucosa, Linn. Sp. Pl. 1010 = C. Pepo.
villosa, Blume, Bijdr. 981 = Benincasa cerifera.
vittata, Blume, l. c. 982 = Lagenaria vulgaris.

CUCURBITELLA, Walp. Rep. vi. 50 (1846) (Curcubitella). *CUCURBITACEÆ*, Benth. & Hook. f. i. 834.
Schizostigma, Arn. in Hook. Journ. Bot. iii. (1841) 275.
asperata, *Walp. Rep.* vi. 50.—Reg. Argent.
cucumifolia, *Cogn. in Mart. Fl. Bras.* vi. iv. 70.—Reg. Argent.
Durieui, *Cogn. l. c.* 70. t. 19.—Reg. Argent.
integrifolia, *Cogn. in DC. Monog. Phan.* iii. 783.—Paraguay.

CUDICIA, Buch.-Ham. ex G. Don, Gen. Syst. iv. 80, in syn. (1837).
gymandra, Buch.-Ham. ex Dillw. Rev. Hort. Malab. 41 = Eodyantheres micrantha !
trichotoma, Buch.-Ham. ex G. Don, Gen. Syst. iv. 80 = Pottsia castonensis !

CUDRANIA, Trec. in Ann. Sc. Nat. Sér. III. viii. (1847) 122. t. 3. *URTICACEAE*, Benth. & Hook. f. iii. 374.
acuminata, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 90, 96.—Java.
amboinensis, Miq. Fl. Ind. Bat. i. ii. 200 = javanensis.
frutescens, *Wight, ex Kurz, For. Fl. Brit. Burma*, ii. 434. —Reg. Himal.; Burma.
javanensis, *Trec. in Ann. Sc. Nat.* Sér. III. viii. (1847) 128.—Trop. As.; Afr.; Austral.
obovata, *Trec. l. c.* 126.—Malaya.
pubescens, *Trec. l. c.* 125.—Burma; Java.
rectispina, *Hance, in Journ. Bot.* xiv. (1876) 365 — Hongkong.
Rumphii, Thw. Enum. Pl. Zeyl. 262 = javanensis.
sumatrana, *Miq. Pl. Jungh.* 44.—Sumatra.
triflora, *Hance, in Journ. Bot.* vi. (1868) 49.—China.

CUDRANUS, Miq. Fl. Ind. Bat. i. ii. 200 (1859) = prsec.

CUELLARA, Pers. Syn. i. 488 (1805) = seq.

CUELLARIA, Ruiz & Pav. Prod. 59. t. 10 (1794); Syst. Veg. 103 (1798) = Clethra, Linn. (Ericac.).
ferruginea, Ruiz & Pav. Syst. Veg. 103 = Clethra ferruginea.
obovata, Ruiz & Pav. l. c. 104 = Clethra obovata.
revoluta, Ruiz & Pav. l. c. 103 = Clethra revoluta.

CUEPIA, J. F. Gmel. Syst. 830 (1791) = Couepia, Aubl. (Rosac.).

CUERVEA, Triana, ex Miers, in Trans. Linn. Soc. xxviii. (1872) 870 = Hippocratea, Linn. (Celastrin.).
amplissima, Miers, l. c. 873.—Ind. occ.
grandiceps, Miers, l. c. 871 = H. granatensis.
latifolia, Miers, l. c. 872 = H. integrifolia.
megacarpa, Miers, l. c. = H. megacarpa.
ovalifolia, Miers, l. c. 871 = H. ovalifolia.

CUIETE, Adans. Fam. ii. 307 (1763) = Crescentia, Linn. (Bignoniac.).

CUITLANZINA, Lindl. Orch. Sel. 15 (1836) = seq.

CUITLAUZINA, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 32 (1825) = Odontoglossum, H. B. & K. (Orchid.).
pendula, La Llave & Lex. l. c. 33 = O. citrosmum.

CUITLAUZINIA, Reichb. Nom. 84 (1841) = prsec.

CUJETE, Anet. CUIETE, Adans. = Crescentia, Linn. (Bignoniac.).

CUJUNIA, Alef. in Bonplandia, ix. (1861) 101 = Vicia, Tourn. (Legumin.).
grandiflora, Alef. l. c. = V. grandiflora.

CULCASIA, Beauv. Fl. Owar. i. 2. t. 3 (1805). *AROIDEAE*, Benth. & Hook. f. iii. 980.
Dennamia, Schott, Melet. 19 (1832).
angolensis, *Welw. ex Schott, in Journ. Bot.* iii. (1865) 86.—Afr. trop.
scandens, *Beauv. Fl. Owar.* i. 4. t. 3.—Afr. trop.

CULCITIUM, Humb. & Bonpl. Pl. Aequin. ii. 1. t. 66, 67 (1809). *COMPOSITAE*, Benth. & Hook. f. ii. 444.
Lasiocephalus, Schlecht. in Gen. Nat. Fr. Berl. Mag. viii. (1816) 808.
Oreiogonia, Schlecht. ex Less. Syn. Comp. 332 (1832).
adscendens, *Benth. Pl. Hartw.* 206.—N. Granat.
candidum, D. Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 343 = Senecio Poeppigii.
canescens, *Humb. & Bonpl. Pl. Aequin.* ii. 4. t. 67.—Peruv.
denticulum, D. Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 343 = Senecio Gilliesii.
denticulatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 208.—Peruv.
depressum, D. Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 343 = Senecio depressus.
Gayanum, *Remy, in C. Gay, Pl. Chil.* iv. 180.—Chili.
glaciale, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 278.—Peruv.
CULCITIUM:—
Hankei, *Wedd. Chlor. And.* i. 189.—Peruv.
humile, *DC. Prod.* vi. 236.—Am. austr.
Hyssertia, *Benth. Pl. Hartw.* 204.—N. Granat.
hypoleucum, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 206.—Ecuador.
Hieracium, Spreng. Syst. iii. 481 = Senecio Scrobicaria.
Lechleri, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
ledifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 170.—Ecuador.
lineare, Spreng. Syst. iii. 481 = Bedfordia linearis.
longifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 206.—Ecuador.
magellanicum, *Hombr. & Jacquem. ex Decne. Bot. Voy. Astrol. et Zél.* 45.—Reg. Magellan.
Neaei, *Sch. Bip. ex Wedd. Chlor. And.* i. 140.—Peruv.
nitidum, Spreng. Syst. iii. 481 = Senecio nitidus.
nivale, *H. B. & K. Nov. Gen. et Sp.* iv. 171.—Ecuador.
Pavonii, *Wedd. Chlor. And.* i. 141.—Peruv.
Poeppigii, *DC. Prod.* vi. 324.—Chili.
reflexum, *H. B. & K. Nov. Gen. et Sp.* iv. 171.—Ecuador.
romaniifolium, *Benth. Pl. Hartw.* 207.—Chili.
rufescens, *Humb. & Bonpl. Pl. Aequin.* ii. 1. t. 66.—Ecuador.
salicinum, Spreng. Syst. iii. 481 = Bedfordia salicina.
sarmentosum, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 279.—Peruv.

CULHAMIA, Forsk. Fl. Aegypt. Arab. 96 (1775) = Sterculia, Linn.
hadiensis, J. F. Gmel. Syst. 754 = S. platanifolia !
platanifolia, Stend. Nom. ed. I. 346 = S. platanifolia.
trilobe, Rafin. Sylva Tellur. 78 = S. platanifolia !

CULLAY, Molina, ex Stend. Nom. ed. II. i. 452 (1840) = Quillaja, Molina (Rosac.).

CULLEN, Medic. in Vorles. Charpf. Phys. Ges. ii. (1787) 381. t. Fagonea, Linn. (Legumin.).
corylifolium, Medic. l. c. = F. corylifolia.

CULLENIA, Wight, Ic. tt. 1761, 1762 (1853). *MALVACEAE*, Benth. & Hook. f. i. 212.
excelsa, *Wight, l. c.*—Ind. or.

CULLOMIA, Juss. in Dict. Sc. Nat. xlii. 307 (1826) = Collomia, Nutt. (Polemon.).

CULLUMIA, R. Br. in Ait. Hort. Kew. ed. II. v. 187 (1813). *COMPOSITAE*, Benth. & Hook. f. ii. 460.
adnata, DC. Prod. vi. 499 = setosa.
adpressifolia, Sond. in Linnaea, xxiii. (1850) 69 = Elonia.
binata, *Less. Syn. Comp.* 82.—Afr. austr.
carlinoides, DC. Prod. vi. 499 = cirsioides.
ciliaris, *R. Br. in Ait. Hort. Kew.* ed. II. v. 187.—Afr. austr.
cirsioides, *DC. Prod.* vi. 496.—Afr. austr.
decurrens, *Less. Syn. Comp.* 82.—Afr. austr.
floccosa, *R. Moy. ex DC. Prod.* vi. 499.—Afr. austr.
hispida, *Less. Syn. Comp.* 87.—Afr. austr.
imbricata, DC. Prod. vi. 498 = decurrens.
microcephala, *DC. Prod.* vi. 499.—Afr. austr.
obovata, *E. Mey.* ex DC. l. c. = micrantha.
patula, *Less. Syn. Comp.* 82.—Afr. austr.
pauciflora, DC. Prod. vi. 500 = hispida.
pectinata, *Less. Syn. Comp.* 86.—Afr. austr.
reflexa, Sch. Bip. in Flora, xxvii. (1844) 777 = decurrens.
rigida, *DC. Prod.* vi. 500.—Afr. austr.
setosa, *R. Br. in Ait. Hort. Kew.* ed. II. v. 187.—Afr. austr.
setosa, Sieber, ex DC. Prod. v. 222 = Felicia reflexa.
squarrosa, *R. Br. in Ait. Hort. Kew.* ed. II. v. 186.—Afr. austr.
subinermis, *DC. Prod.* vi. 500.—Afr. austr.
sulcata, Delge, ex Harv. & Sond. Fl. Cap. iii. 461 = hispida.
volcata, *Less. Syn. Comp.* 84.—Afr. austr.

CUMAROUMA, Stend. Nom. ed. II. i. 452 (1840) = COUMAROUMA, Aubl. = Dipteryx, Schreb. (Legumin.).

CUMARUMA, Stend. Nom. ed. I. 345 (1821) = prsec.

CUMARUNA, J. F. Gmel. Syst. 1084 (1791) = prsec.

CUMBATA, Rafin. Sylva Tellur. 160 (1838) = Rubus, Tourn. (Rosac.).
alcefolia, Rafin. l. c. = R. rugosus.
villosa, Rafin. l. c. = R. rugosus.

CUMBEA, Wight & Arn. Prod. 334 (1834) = seq.

CUMBIA, Bech.-Ham. Mysore, iii. 187 (1807) = Careya, Roxb. (Myrtac.).
Canaanea, Bech.-Ham. in Trans. Linn. Soc. xv. (1827) 97 = Careya arborea.

CUMBULA, Stend. Nom. ed. II. i. 453 (1840) = seq.

CUMBALU, Adans. Fam. ii. 199 (1763) = Catalpa, Scop. (Bignon.).

CUMBTEA, Rafin. Sylva Tellur. 106 (1838) = Eugenia, Mich. (Myrtac.).
alba, Rafin. l. c. = Myrcia Cumete.
angustifolia, Rafin. l. c. 107 = E. angustifolia.
divaricata, Rafin. l. c. 106 = Myrcia divaricata.
fragrans, Rafin. l. c. = E. fragrans.
microphylla, Rafin. l. c.—Hab. ?
Mini, Rafin. l. c. = Myrcia Mini.
montana, Rafin. l. c. = E. fragrans.
multiflora, Rafin. l. c. = E. multiflora.
tomentosa, Rafin. l. c. = Myrcia tomentosa.

CUMINGIA, Kunth, Enum. Pl. iv. 631 (1843) = CUMMINGIA, D. Don = Conanthera, Ruiz & Pav. (Haemodor.).

CUMMINGIA, Vidal, Phan. Cuming. Phil. 211 (1886). *MALVACEAE*.
philippinensis, *Vidal, l. c.* 212.—Ins. Philipp.

CUMINIA, Colla, in Mem. Acc. Tor. xxxvii. (1835) 189. t. 47. *LABIATAE*, Benth. & Hook. f. ii. 1184.
brevidens, *Benth. in DC. Prod.* xii. 258.—Ins. Juan Fernand.
eriantha, *Benth. l. c.*—Ins. Juan Fernand.
fernandezia, *Colla, in Mem. Acc. Tor.* xxxvii. (1835) 189. t. 47.—Ins. Juan Fernand.

CUMINIA, J. F. Gmel. Syst. 309 (1791) = Cuminum, Tourn. (Umbell.).
Cyminum, J. F. Gmel. l. c. 464 = Cuminum Cyminum.

CUMINOIDES, Tourn. ex Moench, Meth. 93 (1794) = Lagoecia, Linn. (Umbellif.).
oblignus, Moench, l. c. 94 = L. cuminoides.

CUMINUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. II. 115 (1742). *UMBELLIFERAE*, Benth. & Hook. f. i. 926.
aegyptiacum, Mérat, ex DC. Prod. iv. 201 = C. Cyminum.
Cyminum, *Linn. Sp. Pl.* 264.—Reg. Mediterr.
Cyminum, Wall. Cat. n. 594 = Coriandrum sativum.
hispanicum, Mérat, ex DC. Prod. iv. 201 = C. Cyminum.
minimum, Stend. Nom. ed. II. i. 453, aphalm. = seq.
minutum, Urv. in Mém. Soc. Linn. Par. i. (1822) 288 = Microscias dum tenuifolium.
odorum, Salisb. Prod. 168 = C. Cyminum.

CUMMIN, Hill, Veg. Syst. vi. 7 (1764) = prsec.

CUMMINGIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. I. iii. t. 257 (1828) = Conanthera, Ruiz & Pav. (Haemodor.).
amœna, Rafin. Fl. Tellur. iii. 61 = Con. bifolia.
campanulata, D. Don, in Sweet, Brit. Flow. Gard. Ser. I. iii. t. 257 = Con. campanulata.
parvula, Phil. in Linnaea, xxiv. (1851-52) 74.—Chili.
tenella, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. iv. sub t. 33 = Con. campanulata.
trimaculata, D. Don, l. c. t. 33.—Chili.

CUNCEA, Bech.-Ham. ex D. Don, Prod. Fl. Nep. 136 (1825) = Wendlandia, Linn. (Rubiac.).
bifida, Bech.-Ham. l. c. = W. corymbosa !

CUNILA, Linn. Gen. ed. I. 176 (1767). *LABIATAE*, Benth. & Hook. f. ii. 1183.
MAPVIA, Heist. ex Adans. Fam. ii. 193 (1763).
angustifolia, *Benth. Lab. Gen. et Sp.* 361.—Bras.
bracteolata, A. Dietr. Sp. Pl. i. 854 = Micromeria bracteolata.
Buchanani, Spreng. Neue Entdeck. iii. 88 = Mosla dianthera.
capitata, Linn. f. Suppl. 87 = Ziziphora clinopodioides.
coccinea, Nutt. ex Hook. Exot. Fl. ii. t. 163 = Calamintha coccinea.
fasciculata, *Benth. Lab. Gen. et Sp.* 368.—Bras.
frutiosa, Willd. Sp. Pl. i. 122 = Westringia rosmarinifornis.
galioides, *Benth. Lab. Gen. et Sp.* 363.—Bras.
glabella, Michx. Fl. Bor. Am. i. 13 = Calamintha glabella.
glabella, Torr. Fl. N. U. St. 23 = Calamintha Nuttallii.
glabra, A. Gray, Syn. Fl. N. Am. ii. i. 360, sphalm. = prasc.
hispida, Spreng. Syst. i. 54 = Hedeoma hispida.
incana, *Benth. Lab. Gen. et Sp.* 361.—Bras.
incisa, *Benth. l. c.*—Bras.
leucantha, *Benth. l. c.*—Mexic.
lyrata, Schrank, in Syll. Ratisb. ii. (1828) 57 = Salvia lyrata.
lythrifolia, *Benth. in Bot. Reg.* sub t. 1289.—Mexic.
Mariana, *Linn. Syst.* ed. X. 1259.—Am. bor.
menthoides, *Benth. Lab. Gen. et Sp.* 363.—Bras.
microcephala, *Benth. l. c.* 364.—Bras.
nepalensis, D. Don, Prod. Fl. Nep. 107 = Mosla dianthera.
paniculata, *Benth. ex Steud. Nom.* ed. II. i. 453, nomen.—Am. bor.
piperita, Moç. & Sesse, ex Benth. Lab. Gen. et Sp. 730 = Hedeoma piperita.
polyantha, *Benth. l. c.* 362.—Mexic.
pulegioides, Linn. Sp. Pl. ed. II. 30 = Hedeoma pulegioides.
secunda, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 126.—N. Mexic.
spicata, *Benth. Lab. Gen. et Sp.* 362.—Bras.
stachyoides, *Mart. & Gal. in Bull. Acad. Brux.* xi. ii. (1844) 190.—Mexic.
stricta, *Benth. Lab. Gen. et Sp.* 362.—Bras.
thymoides, Gouan, ex Benth. l. c. 378 = Micromeria filiformis.
thymoides, Linn. Sp. Pl. ed. II. 31 = Thymus Serpyllum.
verticillata, *Moench, Meth. Suppl.* 141.—Hab.?

CUNINA, Clos, in C. Gay, Fl. Chil. iii. 301. t. 34 (1849) = **Nertera**, Banks & Soland. (Rubiac.).
Sanfuentes, Clos, l. c. 303. t. 34 = N. depressa.

CUNNINGHAMIA, R. Br. in Rich. Conif. 149. t. 18 (1826). *CONIFERAE*, Benth. & Hook. f. iii. 486.
BELIS, Salisb. in Trans. Linn. Soc. vii. (1807) 315, nomen prius.
JACULARIA, Rafin. in Loud. Gard. Mag. viii. (1832) 347.
RACOPITYS, Carr. Conif. ed. II. 237 (1867).
RAXOPITYS, [Nelson] Senilis, Pinsonee, 97 (1866).
cupressoides, Zucc. in Sieb. Fl. Jap. ii. 9 = Athrotaxis cupressoides.
lanceolata, Lamb. Pin. ed. I. t. 53 = sinensis.
selaginoides, Zucc. in Sieb. Fl. Jap. ii. 9 = Athrotaxis selaginoides.
sinensis, *R. Br. in Rich. Conif.* 80. t. 18. f. 3.—China.

CUNNINGHAMIA, Schreb. Gen. 789 (1789) = **Malesherbia**, Ruiz. (Rubiac.).
sarmentosa, Willd. Sp. Pl. i. 815 = M. sarmentosa.
verticillata, Willd. l. c. = Danais fragrans!

CUNONIA, Linn. Syst. ed. X. 1025 (1759). *SAXIFRAGEAE*, Benth. & Hook. f. i. 654.
OXYBADICHIA, Burm. Afr. 229. t. 96 (1737).
Bidennea, *Bronguiart & Gris, in Bull. Soc. Bot. Fr.* xix. (1872) 150.—N. Caled.
bullata, *Brongn. & Gris, l. c.*—N. Caled.
capensis, *Linn. Syst.* ed. X. 1025.—Afr. austr.
cauliflora, *Blume, Bijdr.* 206.—Ins. Celebon.
Daubenyella, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 71.—N. Caled.

CUNONIA:—
indica, *Blume, Bijdr.* 867.—Java.
Lemurnandi, *Vieill. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xix. (1872) 147.—N. Caled.
macrophylla, *Brongn. & Gris, l. c.* ix. (1862) 71.—N. Caled.
paichalla, *Brongn. & Gris, l. c.* 73.—N. Caled.
purpurea, *Brongn. & Gris, l. c.*—N. Caled.
simplicifolia, Vieill. ex Brongn. & Gris, l. c. xix. (1872) 150 = Balmense.
Vieillardi, *Brongn. & Gris, l. c.* ix. (1862) 72.—N. Caled.

CUNONIA, Mill. Gard. Dict. ed. VIII (1868) = **Anthoxylum**, Linn. (Irid.).
Anthoxyna, Mill. l. c. = A. Cunonia.

CUNURIA, Baill. Adansonia, iv. 287 (1863–64). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 289.
CLISIOPHYLLUM, Muell. Arg. in Flora, xlvii. (1864) 518.
crassipes, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 510.—Bras.
Spruceana, *Baill. Adansonia*, iv. (1863–64) 288.—Am. austr.

CUNTO, Adans. Fam. ii. 446 (1763) = **Acronychia**, Forst. (Rutac.).

CUPADESSA, Hask. Cat. Hort. Bogor. Alt. 219 (1844) = **Cipadessa**, Blume (Meliac.).

CUPAMENI, Adans. Fam. ii. 356 (1763) = **Acalypha**, Linn. (Euphorb.).

CUPAMENIS, Rafin. Sylva Tellur. 67 (1838) = **Acalypha**, Linn. (Euphorb.).
chamaedrifolia, Rafin. l. c. = A. reptans.
indica, Rafin. l. c. = A. indica.
reptans, Rafin. l. c. = A. reptans.

CUPANIA, Plum. ex Linn. Gen. ed. I. 339 (1737). *SAPINDACEAE*, Benth. & Hook. f. i. 399.
INCTVNOBURA, Blume, Rumphia, iii. 168 (1849).
DIGONOCARPUS, Vell. Fl. Flum. iv. t. 14 (1827).
DIMERIZA, Labill. Sert. Austro-Cal. 51. t. 51 (1824).
DIPLOPETALON, Spreng. Syst. iv. Cur. Post. 146 (1827).
GUIOA, Cav. Ic. iv. 49. t. 373 (1797).
LACCODISCUS, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 496.
MOLINARA, Juss. Gen. 245 (1789).
TRIGONIS, Jacq. Enum. Pl. Carib. 3 (1760).
TRIGONOCARPUS, Vell. Fl. Flom. iv. t. 15 (1827).
TRIPHA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4.
VALENANA, Adbl. Pl. Gui. Suppl. 12. t. 374 (1775).
acuminata, *Miq. Fl. Ind. Bat.* i. ii. 567.—Borneo.
acuta, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 677.—Malaya.
adenophylla, *Planch. ex Hiern, l. c.*—Burma; Malaya.
affinis, *Kletzsch, in Schomb. Faun. et Fl. Gui.* 1160, nomen.—Guian.
Aherzia, Cambess. ex Spach, Hist. Vég. Phan. iii. 60 = Blighia sapida.
alba, Griseb. in Bonplandia, vi. (1858) 4 = cinerea.
Alphandi, F. Muell. Fragm. iv. 158 = Castanospora Alphandi.
altemifolia, *Perr. Syn.* i. 413.—Ins. Borbon.
americana, Gaertn. Fruct. ii. 409 = glabra.
americana, *Linn. Sp. Pl.* 900.—Mexic.
amoenifolia, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 338 = chlorophylla.
anacardioides, *A. Rich. Sert. Astrol.* 33. t. 18.—Amboina.
anodonta, F. Muell. Fragm. ii. 76 = Ratonia anodonta.
apetala, Labill. Sert. Austro-Caled. 73 = Elattostachys apetala.
aphaia, Macfad. Fl. Jamaic. i. 162 = Ratonia apetala.
apetala, Seem. in Bonplandia, ix. (1861) 254 = Capanopsis Storckii.
Aubletii, Miq. Skirp. Surin. Sal. 67 (= *Matayba arborescens*).—Guian.
Bidwilli, *Benth. Fl. Austral.* i. 460.—Austral.
Blumei, Steud. Nom. ed. II. i. 488 = Harpulia cupanioides.
Brackenridgei, A. Gray, Bot. U. St. Expl. Exped. i. 256 (= *Arytera Brackenridgei*).—Ins. Fiji.

CUPANIA:—
brasiliensis, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 563.—Bras.
canescens, Aurs. in Ann. Sc. Nat. Sér. II. ii. (1834) 336 = Pancovia bijuga.
canescens, Pers. Syn. i. 413 = Hemigyrosa canescens.
castaneaefolia, *Mart. in Flora*, xxi. (1838) ii. *Beibl.* 71.—Bras.
Chapelieriana, Cambess. in Mém. Mus. Par. xviii. (1829) 44 (= *Tina madagascariensis*).—Madag.
cinerea, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 38.—Peruv.
clethrodes, Mart. in Flora, xxi. (1838) ii. Beibl. 72 = vernalis.
collina, *Pancb. & Sebert, Not. Bois Nouv. Caléd.* 230.—N. Caled.
Cordieri, F. Muell. Fragm. ix. 92 = Synima Cordieri.
Cunninghamii, Hook. Bot. Mag. t. 4470 = Diploglottis Cunninghamii.
cupanioides, *Camb. in Mém. Mus. Par.* xviii. (1829) 39.—Mauric.
Dasmeliana, F. Muell. Fragm. ix. 96 = Toechima Dasmelianum.
dentata, *Moç. & Sesse, in DC. Prod.* i. 614.—Mexic.
Dimerosa, Steud. Nom. ed. II. i. 453 = glauca.
diphylla, *Vahl, Eclog. Am.* iii. 9.—Guian.
diphyllostegia, F. Muell. Fragm. v. 145 = Diplogiottis Cunninghamii.
diplopetala, *Hassk. Cat. Hort. Bogor. Alt.* 224.—Java.
discolor, Vahl, Eclog. Am. iii. 9 (= *Matayba arborescens*).—Guian.
distachya, Blume, Rumphia, iii. 162 = verrucosa.
edulis, Schum. & Thonn. Beskr. Guin. Pl. 190 = Blighia sapida.
emarginata, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 386.—Bras.
erythrocarpa, F. Muell. Fragm. v. 7 = Toechima erythrocarpa.
erythrorhachis, Miq. Fl. Ind. Bat. Suppl. i. 509 = Lesertiana.
euphoriaefolia, Cambess. in St. Hil. Fl. Bras. Mer. i. 388. t. 80 (= *Matayba guianensis*).—Bras.
exanguiata, F. Muell. Fragm. iv. 156 = Ratonia exanguiata.
excelsa, H. B. & K. Nov. Gen. et Sp. v. 125 = glabra.
falcata, *A. Gray, Bot. U. St. Expl. Exped.* i. 252.—Ins. Fiji.
ferruginea, *Baker, in Oliver, Fl. Trop. Afr.* i. 425.—Afr. trop.
foveolata, F. Muell. Fragm. ix. 95 = Cupaniopsis foveolata.
frigida, Linden, Cat. n. 12 (1857) 13 = dentata.
frondosa, Benth. in Hook. Kew Journ. iii. (1851) 199 = scrobiculata.
frutescens, Mart. in Flora, xxi. (1838) ii. Beibl. 73 = Sapindus frutescens.
fulva, Mart. l. c. 71 = paniculata.
fulva, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 375.—N. Granat.
furfuracea, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 567.—Peruv.
fuscescens, Miq. Fl. Ind. Bat. i. ii. 567 (= *Mischocarpus fuscescens*).—Java.
fuscidula, *Kurz, in Journ. As. Soc. Beng.* xli. ii. (1872) 302.—Burma.
Gelonium, Steud. Nom. ed. II. i. 454 = laevis, et *Tina Gelonium*.
geminata, Poir. Encyc. Suppl. ii. 419 = diphylla.
Glaberrima, Duchass. ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 376 = (*Matayba glaberrima*).—Panama.
glabra, *Sw. Prod. Veg. Ind. Occ.* 61.—Ind. occ.
glabra, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 378 = papillosa.
glabrata, *Kurz, in Journ. As. Soc. Beng.* xli. ii. (1872) 303.—Burma; Malaya.
glauca, *Cambess. in Mém. Mus. Par.* xviii. (1829) 38.—N. Caled.
gracilis, *Pancb. & Sebert, Not. Bois Nouv. Calédon.* 231.—N. Caled.
grandisima, F. Muell. Fragm. iv. 156 = Ratonia grandisima.
Griffithiana, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 188.—Burma.
guatemalensis, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 563.—Guatemala.
guianensis, Miq. Skirp. Surin. Sal. 71 = scrobiculata.
Helferi, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 679.—Burma.

CUPANIA :—
heterophylla, Mart. in Flora, xxi. (1838) 11. Beibl. 69 (= *Matayba heterophylla*).—Bras.
hirsuta, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 585.—Guiana.
hispida, *Radlk. l. c.* 564.—Bras.
inelegans, Spruce, ex Radlk. l. c. 577 (= *Matayba heterophylla*).—Bras.
isomera, *Baker, in Journ. Bot.* xx. (1882) 69.—Madag.
Jacklana, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 678.—Ins. Nicobar.
juglandifolia, A. Rich. Ess. Fl. Cub. 290 = macrophylla.
juglandifolia, *Seem. Fl. Vit.* 46.—N. Caled.
lachnocarpa, F. Muell. Fragm. iv. 107 = Ratonia lachnocarpa.
laevigata, *Mig. Stirp. Surin. Sel.* 72.—Guian.
laevigata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 (= *Matayba arborescens & M. guianensis*).—Guiana.
laevis, *Pers. Syn.* i. 413.—Ins. Macer.
lanuginosa, *Sagot, ex Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 518, nomen; *et Ann. Sc. Nat.* Sér. VI. xii. (1881) 198.—Guiana.
latifolia, *H. B. & K. Nov. Gen. et Sp.* v. 126.—Am. austr.
laxiflora, *Benth. in Hook. Kew Journ.* iii. (1851) 198.—Bras.
lentiscifolia, *Pers. Syn.* i. 413.—Ins. Amic.
leptobotrya, *A. Gray, Bot. U. St. Expl. Exped.* i. 255.—Ins. Fiji.
Lesserliana, *Cambess. in Mém. Mus. Par.* xviii. (1829) 46. t. 3.—Barma ; Malaya.
longifolia, *Benth. in Hook. Kew Journ.* ii. (1850) 211.—Bras.
lucens, F. Muell. Fragm. iii. 44 = Akania Hillii.
Macgillivrayi, *Seem. Fl. Vit.* 46.—N. Caled.
macrophylla, Mart. in Flora, xxi. (1838) 11. Beibl. 74 = Talisia macrophylla.
macrophylla, *A. Rich. Ess. Fl. Cub.* 291.—Cuba.
madagascariensis, G. Don, Gen. Syst. i. 669 (= *Tina Gelonium*).—Madag.
Martyana, F. Muell. Fragm. v. 6 = Sarcopteryx Martyana.
mexicana, Turz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 408 (= *Matayba mexicana*).—Mexic.
micrantha, Mart. in Flora, xxi. (1838) 11. Beibl. 68 (= *Matayba guianensis*).—Bras.
Miquélien, *Blume, Rumphia*, iii. 162.—Java.
Mischocarpus, Steud. Nom. ed. II. i. 454 = Lesserliana.
Mortoniana, F. Muell. Fragm. v. 177 = Rhysotoechia Mortoniana.
multiflora, Mart. in Flora, xxi. (1838) 11. Beibl. 75 = Scyphonychium multiflorum.
multifuga, A. Rich. Ess. Fl. Cub. 287. t. 81 = glabra.
mutabilis, Miq. Fl. Ind. Bat. i. 11. 566 = discadyba.
nervosa, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 27.—Austral.
nitida, DC. Prod. i. 613 = Paullinia triconnia.
oblongifolia, *Mart. in Flora*, xxi. (1838) 11. *Beibl.* 67.—Bras.
oblongifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 587 = Dittelasma Rarak.
obtusa, *Miq. Fl. Ind. Bat.* i. 11. 567.—N. Guinea.
oppositifolia, A. Rich. Ess. Fl. Cub. 292. t. 82 = Ratonia apetala.
O'Shanesiana, F. Muell. Fragm. ix. 96 = Ratonia O'Shanesiana.
pallidula, *Hiern, in Hook. f. Fl. Brit.* i. 676.—Malacca.
Pancheri, Baill. Adansonia, xi. (1874) 246 = Stortheocalyx Pancheri.
paniculata, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 388.—Bras.
papillosa, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 587.—N. Granat.
pentapetala, *Wight & Arn. Prod.* 113.—Ind. or.
pentaphylla, Wight, Ic. t. 402 = pentapetala.
Perrottetii, Cambess. in Mém. Mus. Par. xviii. (1829) 45 (= *Lepidopetalum Perrottetii*).—Ins. Philipp.
platycarpa, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 567.—Bras.
pleurocarpa, *Blume, Rumphia*, iii. 166.—Borneo.
Poiretii, *Kunth, in Ann. Sc. Nat.* Sér. I. i. (1824) 458.—Ins. S. Trinit.
porosa, Miq. Stirp. Surin. Sel. 70 = scrobiculata.
pseudla, *Sagot, in Ann. Sc. Nat.* Sér. VI. xii. (1881) 199.—Guiana.

CUPANIA :—
Psendorhus, *A. Rich. Sert. Astrolab.* 84. t. 14.—Austral.
punctata, Cambess. in St. Hil. Fl. Bras. Mer. i. 389 (= *Matayba punctata*).—Bras.
punctulata, *F. Muell. Fragm.* iii. 19.—Austral.
purgans, Poepp. & Endl. Nov. Gen. et Sp. iii. 88 (= *Matayba purgans*).—Bras.
pyriformis, F. Muell. App. Rep. Intercol. Exhib. 28 = Ratonia pyriformis.
quercifolia, *Kletzsch, in Schomb. Faun. et Fl. Gui.* 1180, nomen.—Guiana.
racemosa, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 568.—Bras.
Ratonia, Cambess. in Mém. Mus. Par. xviii. (1829) 47 = Ratonia domingensis.
regularis, *Blume, Rumphia*, iii. 159.—Borneo.
reticulata, Cambess. in Mém. Mus. Par. xviii. (1829) 41 ; Spliig. in Hoev. & De Vriese, Tijdschr. ix. (1842) 706 = scrobiculata.
retusa, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 1180, nomen.—Guiana.
Roxburghii, Wight, Ic. t. 402 = pentapetala.
rubiginosa, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 566 = Poiretii.
rufescens, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 374.—N. Granat.
rugosa, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 563.—Bras.
rupestris, Cambess. in Mém. Mus. Par. xviii. (1829) 29 = Harpullia capaniodes.
salicifolia, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 445 = Atalaya salicifolia.
sapida, Volgt, Hort. Suburb. Calc. 94 = Blighia sapida.
Saponaria, Pers. Syn. i. 413 = americana, et Sapindus Saponaria.
saponarioides, Sw. Prod. Veg. Ind. Occ. 61 = prsec.
Schomburgkii, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 1180, nomen.—Guiana.
scrobiculata, Hook. & Arn. Bot. Beech. Voy. 281 = dentata.
scrobiculata, H. B. & K. Nov. Gen. et Sp. v. 127 (= *Matayba scrobiculata*).—N. Granat.
scrobiculata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 109.—Guian.
Seemanni, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 375 = sylvatica.
semiglauca, *F. Muell. ex Benth. Fl. Austral.* i. 457.—Austral.
serrata, *F. Muell. Fragm.* iii. 43.—Austral.
setigera, A. Cunn. ex Heynh. Nom. i. 329 = Pseudorhus.
Sieberiyloa, Cambess. in Mém. Mus. Par. xviii. (1829) 36 = Stadmannia oppositifolia.
spectabilis, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 580.—Mexic.
spinosa, *Blanco, Fl. Filip.* ed. I. 184.—Ins. Philipp.
Spruceana, Benth. in Hook. Kew Journ. iii. (1851) 199 (= *Matayba Spruceana*).—Bras.
stipata, F. Muell. *Fragm.* ii. 79 = Ratonia stipitata.
stipitata, *Planch. & Triana, Ann. Sc. Nat. Bois Nouv. Calédon.* 382.—N. Caled.
subalbens, Mart. in Flora, xxi. (1838) 11. Beibl. 73 = Talisia subalbens.
subciseren, *A. Gray, Bot. U. St. Expl. Exped.* i. 256.—Austral.
subrepanda, Mart. in Flora, xxi. (1838) 11. Beibl. 70 = Poiretii.
subsinuata, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 1180, nomen.—Guiana.
subundulata, *Vidal, Phan. Cuming. Philipp.* 6, 105.—Ins. Philipp.
sumatrana, *Miq. Fl. Ind. Bat.* i. 11. 566.—Sumatra.
sylvatica, Cauar. Nov. Stirp. Bras. Dec. 46 (= *Matayba sylvatica*).—Bras.
sylvatica, *Seem. Bot. Voy. Herald*, 92.—Panama.
tenax, A. Cunn. ex Benth. Fl. Austral. i. 462 = Ratonia tenax.

CUPANIA :—
tenuivalvis, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 562.—Bras.
Thouarsiana, Cambess. in Mém. Mus. Par. xviii. (1829) 43 (= *Tina Gelonium*).—Madag.
Tolambitou, *Cambess. l. c.* 43.—Madag.
tomentella, *F. Muell. ex Benth. Fl. Austral.* i. 458.—Austral.
tomentosa, Sw. Prod. Veg. Ind. Occ. 61 = americana.
truncheyana, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 168 = Trichilia spondioides.
trilaka, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 376 = scrobiculata.
triquetra, *A. Rich. Ess. Fl. Cub.* 290.—Cuba.
uruguensis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 159.—Am. austr.
velutina, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 1180, nomen.—Guiana.
vernalea, DC. Prod. i. 613 = laevis.
vernalis, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 387.—Am. austr.
verrucosa, *Blume, Rumphia*, iii. 161.—Java.
vitiensis, Seem. Viti, 434 = Elattostachys vitiensis.
Vous-rana, *Cambess. In Mém. Mus. Par.* xviii (1829) 43.—Guiana.
Wadsworthii, F. Muell. Census, 24 = Cupaniopsis Wadsworthii.
xylocarpa, *A. Cunn. ex F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 27.—Austral.
xanthoxyloides, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 386.—Bras.
Zippeliana, Blume, Rumphia, iii. 160 = Elattostachys Zippeliana.

CUPANIOPSIS, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 498, 584. *SAPINDACEÆ.*
anacardioides, Radlk. l. c. 585 = Cupania anacardioides.
apiocarpa, *Radlk. l. c.* 586.—N. Caled.
exantha, *Radlk. l. c.* 587.—N. Caled.
chytradenia, *Radlk. l. c.*—N. Caled.
crassivalvis, *Radlk. l. c.*—N. Caled.
dictyophora, *Radlk. l. c.* 589.—N. Caled.
foveolata, *Radlk. l. c.* 585.—Austral.
fruticosa, *Radlk. l. c.* 588.—N. Caled.
gnaophloea, *Radlk. l. c.* 589.—N. Caled.
glomerifiora, *Radlk. l. c.* 589.—N. Caled.
inopica, *Radlk. l. c.*—N. Caled.
leptobotrya, Radlk. l. c. 585 = Cupania leptobotrya.
macrocarpa, *Radlk. l. c.* 587.—N. Caled.
myrmoctona, *Radlk. l. c.* 586.—N. Caled.
oedipoda, *Radlk. l. c.* 590.—N. Caled.
petiolaris, *Radlk. l. c.* 586.—N. Caled.
pallocarpa, *Radlk. l.*—N. Caled.
punctulata, Radlk. l. c. 585 = Cupania punctulata.
serrata, Radlk. l. c. = Cupania serrata.
Storckii, *Radlk. l. c.* 587.—Ins. Fiji.
subcuneata, *Radlk. l. c.* 589.—N. Caled.
trigonocarpa, *Radlk. l. c.* 586.—N. Caled.
Wadsworthii, *Radlk. l. c.* 585.—Austral.

CUPARILLA, Rafin. Sylva Tellur. 130 (1838) = **Acacia**, Tourn. (Legumin.).
myrtifolia, Rafin. l. c. = A. myrtifolia.
Sophorina, Rafin. l. c. = A. longifolia.

CUPHAEA, Moench, Meth. 664 (1794) = seq.

CUPHEA, P. Br. Hist. Jamaic. 216 (1756). *LYTHRARIEÆ*, Benth. & Hook. f. i. 778.
Balladonia, Vand. Fasc. Pl. 18 (1771).
Bankesia, Domb. ex DC. Prod. iii. 88 (1828).
Bergenia, Rafin. Sylva Tellur. 109 (1838).
Diptetalon, Rafin. Fl. Tellur. 108 (1838).
Duvernaya, Desp. in DC. Prod. iii. 86, in syn. (1828).
Endecaria, Rafin. Sylva Tellur. 108 (1838).
Maja, Klotzsch, in Schomb. Fauna et Fl. Gui. 1191 (1848).
Melanium, P. Br. Hist. Jamaic. 215 (1756).
Melpona, Rafin. Sylva Tellur. 106 (1838).
Melvilla, Anders. in Journ. Arts & Sc. xxv. (1807) 307, nomen ; Lindl. Bot. Reg. t. 988 (1824).
Parsonsia, P. Br. Hist. Jamaic. 197. t. 21 (1756).
Quirina, Rafin. Sylva Tellur. 108 (1838).
achilleifolia, *St. Hil. Fl. Bras. Mer.* iii. 96.—Bras.
acinos, *St. Hil. l. c.* 101.—Bras.
acuta, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. 1. 373 = C. Spanneocarpa.

CUPHEA :—
acutissima, St. Hil. Fl. Bras. Mer. iii. 105 = antisyphilitica.
aequipetala, *Cav. Ic.* iv. 57. t. 389. f. 2.—Mexic.
aequipetala, Willd. ex Koehne, in Engl. Bot. Jahrb. ii. (1883) 415 = nitidula.
albida, *Raffin. New Fl. Am.* ii. 32.—Am. bor.
anagalloides, *St. Hil. Fl. Bras. Mer.* iii. 122.—Bras.
angustifolia, *Jacq. ex Koehne, in Mart. Fl. Bras.* xiii. 11. 282.—Mexic.
angustifolia, Mart. ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 149 = thymoides.
anisophylla, *Hemsl. Diagn. Pl. Nov.* 51.—Panama.
annulata, *Koehne, in Mart. Fl. Bras.* xiii. 11. 304.—Bras
antisyphilitica, Benth. in Hook. Journ. Bot. ii. (1840) 316 = graciliz.
antisyphilitica, *H. B. & K. Nov. Gen. et Sp.* vi. 202 — Am. austr.
antisyphilitica, Seem. Bot. Voy. Herald, 121 = anisophylla.
Apanxaloa, DC. Prod. iii. 88 = aequipetala.
aperta, *Koehne, in Mart. Fl. Bras.* xiii. 11. 256.—Bras.
apetala, Hort. ex Koehne l. c. 332 = micropetala.
appendiculata, *Benth. Pl. Hartw.* 61.—Mexic.
appendiculata, Seem. Voy. Bot. Herald, 121 = C. Infundibulum.
arenarioides, *St. Hil. Fl. Bras. Mer.* iii. 121.—Bras.
aristata, *Hemsl. Diagn. Pl. Nov.* 51.—Guatemala.
arvensis, Benth. Fl. Hartw. 37 = jorullensis.
ascranens, Moç. & Sesse, ex DC. Prod. iii. 86 = aequipetala.
aspera, *Chapm. Fl. S. U. St.* 185.—Florida.
aspera, Willd. ex Schlecht. & Cham. in Linnaea, v. (1830) 569 = aequipetala.
atrosanguinea, Warsz. ex Koehne, in Mart. Fl. Bras. xiii. 11. 235 = aequipetala.
Baillonis, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 401. —Mexic.
Balsamona, *Cham. & Schlecht. in Linnaea*, ii. (1827) 363.—Bras.
barbigera, Hook. & Arn. Bot. Beech. Voy. 289 = C. Llaves, *Lex.*
bilabralis, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. 11. 256 = aperta.
Boisseriana, *Koehne, in Engl. Bot. Jahrb.* vii. (1885) 42.—Mexic.!
bourieriana, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 314 = fruticosa.
brachiata, *Mart. ex Koehne, in Mart. Fl. Bras.* xiii. 11. 292.—Bras.
bracteata, *Hook. & Arn. Bot. Beech. Voy.* 289.—Mexic.
bracteata, Lag. Gen. et Sp. Nov. 16 = aequipetala.
Brownei, Jacq. Hort. Vindob. ii. 68 = viscosissima.
Bustamants, *Lex. in La Llave & Lex. Nov. Veg. Desc. fasc.* i. 21.—Mexic.
calaminthaefolia, *Schlecht. in Linnaea*, xii. (1838) 274. —Mexic.
calcarata, *Benth. Pl. Hartw.* 7.—Mexic.
calophylla, *Cham. & Schlecht. in Linnaea*, ii. (1827) 361.—Bras.
campestris, *Mart. ex Koehne, in Mart. Fl. Bras.* xiii. 11. 265.—Bras.
campyloceutra, *Grisob. in Gottt. Abh.* xxiv. (1879) 130. —Uruguay.
camaracrarum, *Spruce, ex Koehne, in Mart. Fl. Bras.* xiii. 11. 226.—Venezuela.
cilliata, Koehne, in Engl. Bot. Jahrb. i. (1881) 454 = decandra.
ciliata, Link, ex Koehne, l. c. ii. (1883) 400 = micropetala.
ciliata, Ruiz & Pav. Syst. Veg. 130 = microphylla.
ciliata, Spreng. ex Koehne, in Mart. Fl. Bras. xiii. 11. 226 = Spruceana.
cinnabarina, Planch. in Fl. des Serres, Sér. I. v. (1846) 527 = pinetorum.
circaeoides, *Sw. ex Sims, Bot. Mag.* t. 2201.—Bras.
coccinea, DC. Prod. iii. 85 = cyanea.
coccinea, Mart. ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 405 = palustris.
Commersoniana, *Koehne, in Mart. Fl. Bras.* xiii. 11. 241.—Reg. Argent.
confertiflora, *St. Hil. Fl. Bras. Mer.* iii. 119.—Bras.
conlata, *Ruiz & Pav. Syst. Veg.* 119.—Peruv.
cordifolia, H. B. & K. Nov. Gen. et Sp. vi. 206 = conlata.
cordifolia, Koehne, in Engl. Bot. Jahrb. ii. (1882) 140 = Swartziana.

CUPHEA :—
corniculata, *Koehne, in Mart. Fl. Bras.* xiii. 11. 286.—Mexic.
corymbifera, Nees, in Flora, iv. (1821) 309 = spicata.
costata, *Koehne, in Mart. Fl. Bras.* xiii. 11. 298.—Bras.
cuiabensis, *Mart. ex Koehne, l. c.* 301.—Bras.
cyanaea, *Moç. & Sesse, in DC. Prod.* iii. 88.—Mexic.
dactylophora, *Koehne, in Mart. Fl. Bras.* xiii. 11. 266. —Guiana.
debilis, *Hemsl. Diagn. Pl. Nov.* 51.—Mexic.
decandra, *[Dryand. in] Ait. Hort. Kew.* ed. II. iii. 151. —Jamaic.
densiflora, *Koehne, in Mart. Fl. Bras.* xiii. 11. 247.—Bras.
densiflora, *St. Hil. ex Steud. Nom.* ed. II. i. 454, nomen.—Bras.
denticulata, *H. B. & K. Nov. Gen. et Sp.* vi. 198.— Venezuela.
diosmaefolia, *St. Hil. Fl. Bras. Mer.* iii. 118.—Bras.
dipetala, Koehne, in Engl. Bot. Jahrb. ii. (1882) 422 = verticillata.
disperma, *Koehne, in Mart. Fl. Bras.* xiii. 11. 278.— Bras.
divaricata, Pohl, ex Koehne, l. c. 255 = C. Balsamona.
dodecandra, Hemsl. Diagn. Pl. Nov. 51 = subuliger a.
Donhellavii, Hort. Berol. ex Koehne, in Mart. Fl. Bras. xiii. 237 = nitidula.
elegans, Klotzsch, ex Koehne, in Mart. Fl. Bras. xiii. 11. 301 = C. Melvilla.
elliptica, *Koehne, ex Engl. Bot Jahrb.* ii. (1882) 145. —Mexic.
elongata, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. 11. 270 = Pseudovaccinium.
enervis, Planch. & Linden, in Fl. des Serres, Sér. I. x. (1854–55) 69 = micropetala.
ennecanthera, *Koehne, in Mart. Fl. Bras.* xiii. 11. 280. —Bras.
epilobiifolia, *Koehne, l. c.* 228.—Costa Rica.
erecta, Hoffmgg. Verz. Pfl. 290 = procumbens.
erectifolia, *Koehne, in Mart. Fl. Bras.* xiii. 11. 277.— Bras.
ericoides, *Cham. & Schlecht. in Linnaea*, ii. (1827) 366.—Bras.
euphorbioides, St. Hil. Fl. Bras. Mer. iii. 112 = micrantha.
excorlata, *Mart. ex Koehne, in Mart. Fl. Bras.* xiii. 11. 273.—Bras.
exposa, Schott, ex Koehne, l. c. 256 = aperta.
ferruginea, *Pohl, ex Koehne, l. c.* 274.—Bras.
flava, *Spreng. Nov. Prov.* 14.—Bras.
floribunda, Hook. & Arn. Bot. Beech. Voy. 289 = Hookeriana.
floribunda, Lehm. in Linnaea, vi. (1831) Litt. 11 = aequipetala.
fruticosa, Koehne, in Mart. Fl. Bras. xiii. 11. 244 = spicata.
fruticosa, *Spreng. Neue Entdech.* ii. 156.—Bras.
fruticulosa, Schrad. in Gott. Gel. Anz. i. (1821) 715 = spicata.
fuchsiaefolia, *St. Hil. Fl. Bras. Mer.* iii 106.—Bras.
fulgens, Fenzl, ex Koehne, in Mart. Fl. Bras. xiii. 11. 287 = Hookeriana.
fulgida, Fenzl, ex Koehne, l. c. = Hookeriana.
Galeottii, Hort. Berol. ex Koehne, l. c. = cyanea.
Gardneri, *Koehne, l. c.* 303.—Bras.
glabra, Gill. ex Hook. & Arn. in Hook. Bot. Misc iii. (1833) 314 = Commersoniana.
glandulosa, *Pohl, ex Koehne, in Mart. Fl. Bras.* xiii. 11. 286.—Bras.
glossostoma, *Koehne, l. c.* 234.—Mexic.
glutinosa, *Cham. & Schlecht. in Linnaea*, ii. (1827) 369.—Bras.
gracilifiora, *Koehne, in Mart. Fl. Bras.* xiii. 11. 286.— Mexic.
gracilis, *H. B. & K. Nov. Gen. et Sp.* vi. 199.—Am. austr.
gracilis, Seem. Bot. Voy. Herald, 121 = utriculosa.
grandiflora, *Pohl, ex Koehne, in Mart. Fl. Bras.* xiii. 11. 308.—Bras.
graticioides, Griseb. Cat. Pl. Cub. 105 = mimuloides.
Grisebachiana, *Koehne, in Mart. Fl. Bras.* xiii. 11. 225. —Cuba.
Gúclichii, *Speguzz. in Anal. Soc. Cient. Argent.* xvi. (1883) 99.—Reg. Argent.
heteropetala, *Koehne, in Mart. Fl. Bras.* xiii. 11. 232. —Mexic.
heterophylla, *Benth. Pl. Hartw.* 37.—Mexic.

CUPHEA :—
hirsuta, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 315 = glutinosa.
hirsuta, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. 11. 258 = ingrata.
hirta, DC. Prod. iii. 86 = micrantha.
hirtella, H. B. & K. Nov. Gen et Sp. vi. 197 = decandra.
hispida, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. 11. 297 = sessiliflora.
Hookeriana, *Walp. Rep.* ii. 107.—Mexic.
hypericoides, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. 11. 261 = thymoides.
hyssopifolia, Griseb. in Goett. Abh. xix. (1874) 140 = mesostemon, Griesebachiana.
hyssopifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 199.— Mexic.
hyssopoides, *St. Hil. Fl. Bras. Mer.* iii. 114.—Bras.
hyssopoides, Schrot, ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 148 = ingrata.
ianthina, *Koehne, in Mart. Fl. Bras.* xiii. 11. 238.— Bolivia.
idiostricha, Steud. ex Koehne, l. c. 294 = flava.
ignea, *A. DC. in Fl. des Serres*, Sér. I. v. (1849) 500 C. —Mexic.
imbricata, Moric. ex Koehne, in Mart. Fl. Bras. xiii. 11. 291 = sericeicles.
impatientifolia, *St. Hil. Fl. Bras. Mer.* iii. 118.—Bras.
inaequalifolia, *Koehne, in Mart. Fl. Bras.* xiii. 11. 236. —Bras.
Infundibulum, *Koehne, l. c.* 286.—Costa Rica.
ingrata, *Cham. & Schlecht. in Linnaea*, ii. (1827) 373 —Bras.
intermedia, *Hemsl. Diagn. Pl. Nov.* 52.—Mexic.
ixodes, *Hemsl. l. c.*—Mexic.
jorullensis, Hook. Bot. Mag. t. 5239 = micropetala.
jorullensis, *H. B. & K. Nov. Gen. et Sp.* vi. 208.— Mexic.
Karwinskii, *Koehne, in Mart. Fl. Bras.* xiii. 11. 234.— Mexic.
Klugii, *Rossi, in Belg. Hortic.* xxiv. (1874) 89.— Peruv.
laminuligera, *Koehne, in Mart. Fl. Bras.* xiii. 11. 234. —Mexic.
lanceolata, Baill. Hist. Pl. vi. 481 = procumbens.
lanceolata, *[Dryand. in] Ait. Hort. Kew.* ed. II. iii. 150.—Mexic.
lanceolato-ovata, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. 11. 299 = lutescens.
lancoides, *Koehne, l. c.* 292.—Bras.
leptociada, Cassar. ex Koehne, l. c. 240 = pascuorum.
leptopoda, *Hemsl. Diagn. Pl. Nov.* 52.—Guatemala.
Liebmanni, *Koehne, in Mart. Fl. Bras.* xiii. 11. 231.— Mexic.
ligustrina, Cham. & Schlecht. in Linnaea, ii. (1827) 359 = fruticosa.
linariioides, *Cham. & Schlecht. l. c.* 367.—Bras.
linifolia, *Koehne, in Mart. Fl. Bras.* xiii. 11. 282.— Bras.
Llavea, Hort. ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 410 = aequipetala.
Llaves, *Lex. in La Llave & Lex. Nov. Veg. Desc. fasc.* i. 20.—Mexic.
Llaves, *Lindl. Bot. Reg.* t. 1386.—Guatemala.
Lloveana, Hemsl. Biol. Centr. Am. Bot. i. 422 = C. Llaves, *Lindl.*
lobelioides, *Griseb. Cat. Pl. Cub.* 105.—Cuba.
loboplora, *Koehne, in Mart. Fl. Bras.* xiii. 11. 285.— Mexic.
longiflora, *Koehne, l. c.* 246.—Bras.
longipedunculata, Mart. ex Koehne, l. c. 261 = linarioides.
lophostoma, *Koehne, l. c.* 235.—Mexic.
lourensis, H. B. & K. Nov. Gen. et Sp. vi. 200 = microphylla.
lucida, Langsd. ex Koehne, in Mart. Fl. Bras. xiii. 11. 256 = aperta.
lutescens, *Pohl, ex Koehne, l. c.* 299.—Bras.
lysimachioides, *Cham. & Schlecht. in Linnaea*, ii. (1827) 374.—Bras.
magnoliaemica. Spreng. Syst. iv. Cur. Post. 190 = Commersoniana.
melampyrifolia, *Pohl, ex Koehne, in Mart. Fl. Bras.* xiii. 11. 268.—Bras.
Melanium, R. Br. ex Steud. Nom. ed. I. 945 = viscosissima.
Melvilla, *Lindl. Bot. Reg.* t. 852.—Guiana.
mesostemon, *Koehne, in Mart. Fl. Bras.* xiii. 11. 252.— Bras.

CUPHEA:—
micrantha, *H. B. & K. Nov. Gen. et Sp.* vi. 196.—Venezuela.
microcarpa, Baill. Hist. Pl. vi. 429 = C. Llaves, *Lex.*
micropetala, *H. B. & K. Nov. Gen. et Sp.* vi. 209. t. 551.—Mexic.
microphylla, *H. B. & K. l. c.* 201.—Peruv.
microstyla, *Koehne, in Mart. Fl. Bras.* xiii. ii. 224.—Mexic.
millefoliata, Pohl, ex Koehne, l. c. 258 = ingrata.
mimiloides, *Cham. & Schlecht. in Linnæa*, v. (1830) 570.—Mexic.
minuta, Brongn. in Fl. des Serres, Sér. I. iii. (Janv. 1846) t. 10 = C. Llaves, *Lex.*
montana, Roesl, ex Regel, Gartenfl. viii. (1859) 277 = jorullensis.
multicaulis, *Koehne, in Mart. Fl. Bras.* xiii. ii. 226.—Venezuela.
multiflora, *Lodd. Bot. Cab.* t. 808.—Ins. Trinit.
Neuberti, Hort. ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 397 = C. Llavsi.
nitens, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. ii. 270 = Pseudovaccinium.
nitidula, *H. B. & K. Nov. Gen. et Sp.* vi. 206.—Mexic.
nudicostata, *Hemsl. Diagn. Pl. Nov.* 52.—Mexic.
nudiflora, Willd. ex Stend. Nom. ed. II. i. 454 = spicata.
ocymoides, Decne. in journ. Horticult. iii. t. 8 = aequipetala.
origanifolia, *Cham. & Schlecht. in Linnæa*, ii. (1827) 378.—Bras.
origanifolia, Griseb. in Goetz. Abh. xxiv. (1879) 130 = spicata.
orisabaensis, Peyr. ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 414 = calaminthifolia.
orthodisca, *Koehne, in Mart. Fl. Bras.* xiii. ii. 224.—Mexic.
palustris, Hemsl. Biol. Centr. Am. Bot. i. 445 = procumbens.
palustris, *Koehne, in Mart. Fl. Bras.* xiii. ii. 229.—Mexic.
panamensis, Hemsl. Diagn. Pl. Nov. 52 = utricnlosa.
paradoxa, *Koehne, in Mart. Fl. Bras.* xiii. ii. 280.—N. Granat.; Venezuela.
parviarioides, Koehne, in Engl. Bot. Jahrb. ii. (1882) 170 = petiolata.
Parsonsia, *R. Br. ex Steud. Nom.* ed. I. 245.—Ind. occ.
parvifora, Hook. bxot. Bot. t. 151 = multiflora.
pascuorum, *Mart. ex Koehne, in Mart. Fl. Bras.* xiii. ii. 240.—Bras.
patula, *St. Hil. Fl. Bras. Mer.* iii. 101.—Bras.
pauciflaria, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. ii. 301 = C. Melvilla.
peltidioides, Mart. ex Koehne, l. c. 255 = C. Balsamona.
peltidata, Koehne, in Engl. Bot. Jahrb. ii. (1882) 178 = viscosissima.
petiolata, *Pohl, ex Koehne, in Mart. Fl. Bras.* xiii. ii. 296.—Bras.
pinetorum, *Benth. Pl. Hartw.* 74.—Mexic.
platycentra, Lem. in Fl. des Serres, ii. (1846) t. 180 = ignea.
platycentra, Benth. Pl. Hartw. 7 = C. Bustamanta.
plumbaginea, Mart. in Flora, xxi. (1838) ii. Beibl. 62 = calophylla.
Pohliana, Mart. ex Koehne, in Mart. Fl. Bras. xiii. ii. 272 = C. Spermacoce.
polygalæfolia, Pohl, ex Koehne, l. c. 281 = linarioides.
polymorpha, *St. Hil. Fl. Bras. Mer.* iii. 108.—Bras.
polymorphoides, *Koehne, in Mart. Fl. Bras.* xiii. ii. 269.—Bras.
procumbens, *Cav. Ic.* iv. 55. t. 380.—Mexic.
procumbens, *Ortog. Hort. Matr.* 17.—Cuba.
propinqua, *Hemsl. Diagn. Pl. Nov.* 53.—Mexic.
prunellæfolia, St. Hil. Fl. Bras. Mer. iii. 106 = circacoides.
Pseudomelastomiana, Griseb. Cat. Pl. Cub. 104 = viscosissima.
Pseudoselaine, *Griseb. l. c.* 105.—Cuba.
Pseudovaccinium, *St. Hil. Fl. Bras. Mer.* iii. 102.—Bras.
pteroseperma, *Koehne, in Mart. Fl. Bras.* xiii. ii. 264.—Bras.
pubiflora, Benth. Pl. Hartw. 61 = cyanea.
pulchra, *Moric. Pl. Nouv. Am.* 168. t. 89.—Bras.
pumila, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. ii. 281 = linarioides.
pumila, Schott, ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 162 = retrorsiculispilla.

CUPHEA:—
punctulata, *Koehne, in Mart. Fl. Bras.* xiii. ii. 248.—Bras.
purpurea, Hort. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 412 = procumbens.
pustulata, *Koehne, in Mart. Fl. Bras.* xiii. ii. 224.—Hab. ?
quadrifolia, Pohl, ex Koehne, l. c. 272 = C. Spermacoce.
racemosa, Spreng. Syst. ii. 455 = spicata.
radicans, Macfad. Fl. Jamaic. ii. 87 = C. Parsonsia.
Radula, *Koehne, in Mart. Fl. Bras.* xiii. ii. 287.—Bras.
ramosa, Schott, ex Koehne, l. c. 256 = aperta.
ramosissima, *Pohl, ex Koehne, l. c.* 249.—Bras.
ramulosa, *Mart. ex Koehne, l. c.* 289.—Bras.
rapunculoides, Griseb. Cat. Pl. Cub. 105 = micrantha.
repens, *Koehne, in Mart. Fl. Bras* xiii. ii. 251.—Bras.
reticulata, *Koehne, l. c.* 275.—Bras.
retrorsispilla, *Koehne, l. c.* 278.—Bras.
rigidula, *Benth. in Hook. Journ. Bot.* ii. (1840) 316.—Guiana.
rigidula, Seem. Bot. Voy. Herald, 121 = setosa.
rivularis, *Seem. l. c.*—Panama.
Roezlii, Carr. in Rev. Hortic. (1877) 469 = Hookeriana.
rotundifolia, *Koehne, in Mart. Fl. Bras.* xiii. ii. 224.—Ins. S. Doming.
rubescens, *Koehne, l. c.* 266.—Bras.
ruficapilla, Mart. ex Koehne, l. c. 274 = ferruginea.
salicifolia, *Cham. & Schlecht. in Linnæa*, v. (1830) 566.—Mexic.
scaberrima, *Koehne, in Mart. Fl. Bras.* xiii. ii. 258.—Bolivia.
scabra, Pohl, ex Koehne, l. c. 256 = aperta.
scabrida, H. B. & K. Nov. Gen. et Sp. vi. 208 = aequipetala.
sclerophylla, *Koehne, in Mart. Fl. Bras.* xiii. ii. 268.—Bras.
secundiflora, Moç. & Sesse, ex DC. Prod. iii. 84 = spicata.
serpyllifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 201. t. 550.—N. Granat.
serpyllifolia, Schnech, ex Koehne, in Mart. Fl. Bras. xiii. ii. 281 = thymoides.
serpyllifolia, Sims, in Bot. Mag. t. 2580 = multiflora.
sessilifiora, *St. Hil. Fl. Bras. Mer.* iii. 110.—Bras.
sessilifolia, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 7.—Bras.
setosa, *Koehne, in Mart. Fl. Bras.* xiii. ii. 228.—Mexic.
silenoides, Nees, in Linnæa, x. (1836) Litt. 71 = lanceolata.
Simsii, Sweet, Hort. Brit. ed. I. 151 = multiflora.
speciosa, Hort. ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 396 = Llavea.
spergulioides, *St. Hil. Fl. Bras. Mer.* iii. 121.—Bras.
Spermacoce, *St. Hil. l. c.* 114.—Bras.
spicata, *Cav. Ic.* iv. 56. t. 381.—Am. trop.
spicata, Cham. & Schlecht. in Linnæa, ii. (1827) 358 = origanifolia.
splendens, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. ii. 270 = Pseudovaccinium.
Spruceana, *Koehne, l. c.* 226.—Peruv.
squamuligera, *Koehne, l. c.* 285.—Mexic.
stenopetala, *Koehne, in Engl. Bot. Jahrb.* ii. (1882) 156.—Paraguay.
stricta, Moric. ex Koehne, l. c. 167 = ericoides.
stricta, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. ii. 277 = ericifolia.
strigillosa, Lindl. Bot. Reg. (1846) t. 14 = cyanea.
strigulosa, *H. B. & K. Nov. Gen. et Sp.* vi. 204.—Am. austr.
strigulosa, Lem. in Fl. des Serres, i. (1845) 89. t. 14 = cyanea.
strumosa, Moç. & Sesse, ex DC. Prod. iii. 84 = micropetala.
subciliigera, *Koehne, in Mart. Fl. Bras.* xiii. ii. 281.—Mexic.
Swartziana, *Spreng. Syst.* ii. 455.—Ins. S. Doming.
tenella, Hook. & Arn. Bot. Beech. Voy. 389 = minuloides.
tenuissima, *Koehne, in Mart. Fl. Bras.* xiii. ii. 254.—Bras.
tenuis, *Peyr. in Linnæa*, xxx. (1859–60) 71.—Mexic.
tetrapetala, *Koehne, in Mart. Fl. Bras.* xiii. ii. 222.—Mexic.
tetraphylla, Pohl, ex Koehne, l. c. 273 = C. Spermacoce.

CUPHEA:—
thesioides, Humb. ex Koehne, in Engl. Bot. Jahrb. i. (1881) 454 = denticulata.
thymoides, *Cham. & Schlecht. in Linnæa*, ii. (1827) 368.—Bras.
thymoides, Lorentz, Veg. Nordeste Entrerios, 185 = glutinosa.
tolucana, Peyr. in Linnæa, xxx. (1859–60) 72 = Wrightii.
tricolor, Moç. & Sesse, ex DC. Prod. iii. 84 = jorullensis.
trifoliata, Pohl, ex Koehne, in Mart. Fl. Bras. xiii. ii. 273 = C. Spermacoce.
Trinitatis, DC. Prod. iii. 88 = multiflora.
tuberosa, *Cham. & Schlecht. in Linnæa*, ii. (1827) 372.—Bras.
tubiflora, Hort. ex Lem. in Fl. des Serres, Sér. I. ii. (Dec. 1846) t. 180 = ignea.
Urbaniana, *Koehne, in Engl. Bot. Jahrb.* ii. (1882) 152.—Bras.
utriculosa, *Koehne, in Mart. Fl. Bras.* xiii. ii. 222.—Mexic.
verticillata, *H. B. & K. Nov. Gen. et Sp.* vi. 207. t. 552.—Peruv.
vesicnlosa, *Koehne, in Mart. Fl. Bras.* xiii. ii. 267.—Bras.
villosa, Pohl, ex Koehne, l. c. 299 = intexcens.
violacea, Regel, in Flora, xxxiii. (1850) 354 = aequipetala.
virgata, Cav. Ic. iv. 56. t. 382 = aequipetala.
virgata, DC. Prod. iii. 86 = nitidula.
viscosissima, *Jacq. Hort. Vindob.* ii. 33. t. 177.—Am. bor.
viscosissima, St. Hil. Fl. Bras. Mer. iii. 110 = intexcens.
Warmingh, *Koehne, in Mart. Fl. Bras.* xiii. ii. 209.—Bras.
Weddelliana, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 402.—Bolivia.
Wrightii, *A. Gray, Pl. Wright.* ii. 56.—N. Mexic.
zorulleusis, Spreng. Syst. ii. 456 = jorullensis.
Zimapani, E. Morr. in Belg. Hortic. alv. (1864) 75; Rocsl, ex Regel, in Suppl. ad Ind. Sem. Hort. Petr. (1868) 15 = lanceolata.

CUPHEANTHUS, Seem. Fl. Vit. 76 (1865–73). *MYRTACEÆ*, Benth. & Hook. f. i. 734.
neocaledonicus, *Seem. l. c.*—N. Caled.

CUPHOCARPUS, Decne. & Planch. in Rev. Hortic. (1854) 109. *ARALIACEÆ*, Benth. & Hook. f. i. 937.
aculeatus, *Decne. & Planch. l. c.*—Madag.
inermis, *Baker, in Journ. Linn. Soc.* xxi. (1884) 350.—Madag.

CUPHOEA, Brongn. ex Neumann, in Rev. Hortic. Sér. II. iv. (1845–46) 235 = Cuphea, R. Br. (Lythrar.).

CUPI, Adans. Fam. ii. 158 (1763) = seq.

CUPIA, DC. Prod. iv. 398 (1830) = **Randia**, Linn. (Rubiac.).
auriculata, DC. l. c. 394 = Anomanthodia auriculata.
corymbosa, DC. l. c. (= *Webera corymbosa*).—Hongkong.
corymbosa, Hook. & Arn. Bot. Beech. Voy. 182 (= *Webera attenuata*).—Ind. or.
cymosa, DC. Prod. iv. 394 = Plectronia didyma.
dumetiflora, DC. l. c. = Randia dumetiflora.
macrophylla, DC. l. c. (= *Webera odorata*).—Ind. or.
mollissima, Hook. & Arn. Bot. Beech. Voy. 193 (= *Webera mollissima*).—China.
odorata, DC. Prod. iv. 394 (= *Webera odorata*).—Ind. or.
oppositifolia, DC. l. c. = Randia dumetiflora.
scandens, DC. l. c. = Randia longiflora.
setifera, Korth. Verhand. Akath. ii. ii. (1851) 276 (= *Stylocoryna setigera*).—Sumatra.
thyrsoidea, DC. Prod. iv. 394 = Wendlandia Notoniana.
truncata, DC. l. c. = Triaenodia truncata.

CUPIRANA, Miers, Apocyn. S. Am. 15 (1878) = Clempoa, Aubl. (Apocyn.).
Aublétiana, Miers, l. c. 16 = Geopoxi egestica.
Martiusiana, Miers, l. c. 17.—Am. austr.

CUPRESSINNATA, [Nelson] Senilis, Pinaceæ, 61 (1866) = Taxodium, Rich. (Conif.).
distichæ, [Nelson] Senilis, l. c. = T. distichum.
heterophylla, [Nelson] Senilis, l. c. 62 = T. heterophyllum.
mexicana, [Nelson] Senilis, l. c. 62 = T. mexicanum.
sinensis, [Nelson] Senilis, l. c. = T. distichum.

CUPRESSTELLATA, [Nelson] Senilis, Pinaceæ, 60 (1866) = Pitzroya, Hook. (Conif.).
patagonica, [Nelson] Senilis, l. c. = F. patagonica.

CUPRESSUS, Tourn. ex Linn. Gen. ed. I. 294 (1737).
CONIFERÆ, Benth. & Hook. f. iii. 427.
africana, Mill. Gard. Dict. ed. VIII. n. 6 = Callitris cupressoides.
americana, Catesb. ex Endl. Syn. Conif. 68 = Taxodium distichum.
americana, Trastv. Pl. Imag. Fl. Russ. 12. t. 7 = nootkatensis.
amoena, *C. Koch, in Mgnatsschr. Garten. Pflanz.* i. 110.—China.
Arbor-vitæ, Targ. Tozz. Obs. Bot. Dec. iii–v. 71 = Thuya occidentalis.
arizonica, *Greene, in Bull. Torrey Club.* ix. (1882) 64. —Arizona.
aromatica, Gord. Pinet. 56 = Benthami.
articulata, Forbes, Pinet. Woburn. 191 = Callitris quadrivalvis.
attenuata, Gord. Pinet. 87 = Lawsoniana.
australis, Low, ex Gord Pinet. 72 = sempervirens.
australis, Pers. Syn. ii. 580 = Callitris rhomboides.
bacciformis, Knight, ex Gord. Pinet. Suppl. 88 = Juniperus occidentalis.
bacciformis, Willd. Enum. Hort. Berol. Suppl. 64, nomen = Juniperus phœnicea.
Balfouriana, *Lemoine, ex Gord. Pinet.* ed. II. 79.—Am. bor. occ.
Benthami, *Endl. Syn. Conif.* 59.—Mexic.
Bourgeauii, Hort. ex Gord. Pinet. ed. II. 79 = Goveniana.
*brevirami*s, F. Muell. Sel. Pl. Indust. Cult. 64 = Thuya obtusa.
californica, Carr. Conif. ed. I. 127 = Goveniana.
chilensis, Gill. ex Hook. Lond. Journ. Bot. ii. (1843) 199 = Libocedrus chilensis.
columnaris, Forst. f. Prod. 67 = Araucaria excelsa.
comoides, Spad. Xilolog. i. 189 = sempervirens.
cornuta, Carr. in Rev. Hortic. (1866) 250 = Goveniana.
Coulteri, Forbes, Pinet. Woburn. 190 = Lindleyi.
disticha, Linn. Sp. Pl. 1003 = Taxodium distichum.
Ehrenbergii, Kunze, in Linnæa, xx. (1847) 16 = Benthami.
elongata, Salisb. Prod. 397 = sempervirens.
excelsa, Scott, ex Carr. Conif. 138 = Benthami.
expansa, Targ. Tozz. ex Steud. Nom. ed. II. i. 455 = sempervirens.
fastigiata, DC. Fl. Fr. v. 336 = sempervirens.
flagelliformis, Knight, Syn. Conif. 20 = torulosa.
Fothergilli, Forbes, Pinet. Woburn. 191 = Callitris quadrivalvis.
fragrans, Kellogg, in Proc. Calif. Acad. i. (1856) 103 = Lawsoniana.
funebris, *Endl. Syn. Conif.* 58.—China.
glandulosa, Hook. ex Gord. Pinet. 64 = Macnabiana.
globulifera, Parl. in Ann. Sc. Nat. Sér. IV. xiii. (1860) 377 = sempervirens.
glauca, *Law. Encyc.* ii. 248.—Ind. or.
Goveniana, *Gord. in Journ. Hort. Soc.* iv. (1849) 295. —Calif.
guadalupensis, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 300.—Calif.
Hartwegii, Carr. in Rev. Hortic. (1855) 238 = macrocarpa.
horizontalis, Mill. Gard. Dict. ed. VIII. n. 2 = sempervirens.
humilis, Salisb. Prod. 396 = Callitris arborea.
imbricata, Nutt. ex Gord. Pinet. 307 = Taxodium distichum.
japonica, Linn. f. Suppl. 421 = Cryptomeria japonica.
juniperoides, Linn. Sp. Pl. ed. II. 1422 = Callitris arborea.
Knowltoniana, Regel, Gartenfl. (1857) 246 = Lindleyi.
Knightiana, Perry, ex Gord. Pinet. 61 = Lindleyi.
lævis, Salisb. Prod. 397 = Taxodium distichum.
Lambertiana, Hort. ex Carr. Conif. 136 = macrocarpa.
Lawsoniana, *Andr. Murr. in Edinb. New Phil. Journ.* N. S. i. (Jan.–Apr. 1855) 292. t. 9.—Am. bor. occ.

CUPRESSUS :—
Lindleyi, *Klotzsch, ex Endl. Syn. Conif.* 59.—Mexic.
lugubris, Salisb. Prod. 397 = sempervirens.
lusitanica, Carr. Conif. ed. II. 165 = Benthami.
lusitanica, Lindl. ex Parl. in DC. Prod. xvi. ii. 472 = Lindleyi.
lusitanica, Mill. Gard. Dict. ed. VIII. n. 3 = glauca.
Macnabiana, *Andr. Murr. in Edinb. New Phil. Journ.* N. S. i. (Jan.–Apr. 1855) 293.—Calif.
macrocarpa, *Hartw. in Journ. Hort. Soc.* ii. (1847) 187.—Calif.
majestica, Knight, Conif. 20 = torulosa.
Montezumæ, Humb. & Bonpl. ex Parl. in DC. Prod. xvi. ii. 441 = Taxodium mucronatum.
nepalensis, Loud. Encyc. Trees, 1118 = torulosa.
nootkatensis, *Lamb. Pin.* ed. I. ii. 18.—Am. bor. occ.
nucifera, Hort. ex Carr. Conif. ed. I. 151 = Taxodium heterophyllum.
nutkansus, Torr. in Bot. U. St. Expl. Eped. ii. t. 16 = Lawsoniana.
nuthatensis, Brdly. Bot. Fav. Am. ii. 165 = nootkatensis.
obtusa, C. Koch, Dendrol. ii. ii. 168 = Thuya obtusa.
palustris, Salisb. Prod. 398 = sphæroidalis.
patula, Pers. Syn. ii. 580 = Thuya orientalis.
patula, Spad. Xilolog. i. 193 = sempervirens.
pendens, F. Muell. Sel. Pl. Indust. Cult. 64 = Thuya obtusa.
pendula, [Abel. in] Staunt. Embassy, China, ii. 325 = funebris.
pendula, Griff. Notul. iv. 26 = torulosa.
pendula, L'Hérit. Stirp. Nov. 15. t. 18 = glauca.
pendula, Thunb. Fl. Jap. 265 = Thuya orientalis.
pisifera, C. Koch, Dendrol. ii. ii. 170 = Thuya pisifera.
pyramidalis, Targ. Tozz. Obs. Bot. Dec. iii–v. 73 = sempervirens.
Reyleï, Carr. Conif. ed. I. 128 = sempervirens.
sabinoides, H. B. & K. Nov. Gen. et Sp. ii. 8 = Juniperus tetragona.
Schomburghii, Van Houtte, ex Gord. Pinet. Suppl. 27 = thuriferi.
sempervirens, *Linn. Sp. Pl.* 1002.—Europ. ; Orient.
sinensis, Lee, ex Gord. Pinet. 63 = glauca.
sphærocarpa, Parl. in Ann. Sc. Nat. Sér. IV. xiii. (1860) 378 = sempervirens.
squarrosa, Laws. ex Gord. Pinet. 296 = Thuya pisifera.
stricta, Mill. ex Gord. l. c. 67 = sempervirens.
tetragona, Humb. & Bonpl. ex Carr. Conif. ed. II. 38. = Juniperus tetragona.
Thuia, Targ. Tozz. Osserv. Dec. ii. 52 = Thuya orientalis.
thuioides, Pav. ex Carr. Conif. II. 89 = Libocedrus chilensis.
thujafolia, Knight, ex Gord. Pinet. 69 = sempervirens.
thujaformis, Parker, ex Gord. l. c. Suppl. 26 = sempervirens.
thujaoides, Low, ex Gord. l. c. = sempervirens.
thurifera, *H. B. & K. Nov. Gen. et Sp.* ii. 3.—Mexic.
thurifera, Lindl. Bot. Reg. (1839) Misc. 64 = Lindleyi.
thurifera, Schlecht. in Linnæa, xii. (1838) 493 = Benthami.
Thuya, Targ. Tozz. Obs. Bot. Dec. iii-v. 72 = Thuya orientalis.
thyoides, Linn. Sp. Pl. 1008 = Thuya sphæroidalis.
thyoides, Pav. ex Gord. Pinet. Suppl. 41 = Libocedrus chilensis.
torulosa, *D. Don, Prod. Fl. Nep.* 55.—Reg. Himal.
Tournefortii, Audib. Cat. (1834) ex Carr. Conif. ed. I. 115 = sempervirens.
Tournefortii, Tenore, Ind. Sem. Hort. Neap. (1852).— Cf. Mem. Soc. Ital. Sc. Modena, xxv. ii. (1855) 194 = torulosa.
triquetra, Todd. ex Lond. Hort. Brit. 388 = Callitris quadrivalvis.
Uhdeana, Gord. ex Carr. Conif. ed. I. 129 = thuriferа.
washingtonia, Parl. Bot. Sem. Hort. Florent. (1856) 22.— Cf. Ann. Sc. Nat. Sér. IV. xiii. (1860) 378 = sempervirens.

CUPRESTELLATA, Carr. Conif. ed. II. 115 (1867) = CUPRESSTELLATA, [Nelson] = Pitzroya, Hook. f. (Conif.).

CUPULA, Raf. Princ. Somiol. 28 (1814) = Coupoui, Aubl. (Apocynac.).

CUPULARIA, Godr. & Gren. in Gren. & Godr. Fl. Fr. ii. 160 (1850) = Inula, Linn. (Compos.).
Claussenis, Billot, ex Pomel, Nouv. Mat. Fl. Atl. 36 = Pulicaria Claussenis.

CUPULARIA :—
graveolens, Godr. & Gren. in Gren. & Godr. Fl. Fr. ii. 160 = Inula graveolens.
viscosa, Godr. & Gren. l. c. 161 = I. viscosa.

CUPULISSA, Raf. Fl. Tellur. iii. 57 (1836) = Big nonia, Linn.
grandiflia, Raf. l. c. = B. grandiflora.

CURANGA, Juss. in Ann. Mus. Par. ix. (1807) 319. *SCROPHULARINEÆ*, Benth. & Hook f. ii. 954.
CURANIA, Roem. & Schult. Syst. i. 138 (1817).
DRUPINA, Linn. Pl. Syria. 11 (1776) ; Amoen. Acad. viii. 259 (1785).
PICRIA, Lour. Fl. Cochin. 392 (1790).
SYMPHYLLIUM, Griff. in Madr. Journ. Sc. iv. (1836) 373.
TRISTERIA, Griff. Notul. iv. 113 (1854).
amara, *Juss. l. c.* 320.—Ins. Moluce.
tomentosa, Steud. Nom. ed. II. i. 455 = amara.

CURANIA, Roem. & Schult. Syst. i. 138 (1817) = Curanga, Juss. (Scrophul.).

CURARE, Kunth, ex Humb. Relat. Hist. ii. 349 (1817) = Strychnos, Linn. (Loganiac.).

CURATARI, J. F. Gmel. Syst. 1089 (1791) = Couratari, Aubl. (Myrtac.).

CURATARIA, Spreng. Gen. i. 486 (1830) = prasc.

CURATELLA, Linn. in Locfl. It. Hisp. 260 (1758). *DILLENIACEÆ*, Benth. & Hook. f. i. 12.
PINZONA, Mart. & Zucc. in Abh. Akad. Muench. i. (1823) 871.
alata, *Vent. Choix*, 49.—Guiana.
americana, *Linn. Syst.* ed. X. 1079.—Am. austr.
Çambaiba, St. Hil. Pl. Us. Brés. t. 24 = americana.
glabra, Spruce, ex Benth. in Journ. Linn. Soc. v. (1861) 64 = americana.
Griesbachiana, *Eichl. in Mart. Fl. Bras.* xiii. i. 59.— Bras.

CURCAS, Adans. Fam. ii. 356 (1763) = Jatropha, Linn. (Euphorb.).
Adansonii, Endl. ex Heynh. Nom. ii. 176 = J. Curcas.
cunefolia, Baill. Etud. Gén. Euph. 315 = J. spatulata.
indica, A. Rich. Fl. Cub. Fanerog. ii. 208 = J. Curcas.
pelitata, Baill. Etud. Gén. Euph. 315 = J. hernandiæfolia.
portoricensis, Baill. l. c. 314 = J. hernandiæfolia.
purgans, Medic. Malv. 119 = J. Curcas.

CURCUBITELLA, Walp. Rep. vi. 50 (1846), sphalm. vide Cucurbitella, Walp.

CURCULIGO, Gaertn. Fruct. i. 63. t. 16 (1788) *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 717.
AUROTA, Rafin. Fl. Tellur. iii. 61 (1836).
EMPODIUM, Salisb. Gen. Pl. Fragm. 43 (1866).
FABRICIA, Thunb. in J. C. Fabric. Reise nach Norw. 23 (1779).
FORRESIA, Eckl. Verz. Pfl. Samml. 4 (1827).
MOLINERIA, Colla, Hort. Ripul. App. ii. 331 (1826).
brevifolia, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 258 = orchioides.
ensifolia, K. Br. Prod. 290 = orchioides.
fasciculariа, Spreng. Syst. ii. 60 = Colchicum montanum.
Finlaysoniana, *Wall. Cat.* n. 5162.—Siam.
firma, Kotschy & Peyr. Pl. Tinn. 45. t. 22 B = orchioides.
gallabatensis, *Schweinf. ex Baker, in Trans. Linn. Soc. Ser. II. Bot.* i. (1878) 266.—Afr. trop.
gracilis, *Wall. Cat.* n. 5160.—Reg. Himal.
graminifolia, Nimmo, in J. Grah. Cat. Pl. Bomb. 215 = Hypoxis nurea.
latifolia, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 258. —Burma ; Malaya.
malabarica, Wight, Ic. t. 2042 = orchioides.
orchioides, *Gaertn. Fruct.* i. 63. t. 16.—As. trop.
pauciflora, Zipp. ex Span. in Linnæa, xv. (1841) 477 = orchioides ?
plicata, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 258. —Afr. austr.
recurvata, [*Dryand.*] l. c.—As. trop. ; Austral.
scorzoneræfolia, *Baker, in Journ. Linn. Soc.* xvii. (1878) 194.—Am. trop.
seychellensis, *Boj. Hort. Maurit.* 249, nomen ; *Baker, Fl. Maurit.* 368.—Ins. Masc.

CURCULIGO :—
stans, Labill. Sert. Austr. Caled. 18. t. 24 = orchioides.
sumatrana, Roxb. Hort. Beng. 94; Fl. Ind. ii. 146 = latifolia.
veratrifolia, *Baker*, *in Journ. Linn. Soc.* xvii. (1878) 128.—Ait. austr.
villosa, Wall. Cat. n. 5163 = latifolia, recurvata.

CURCUMA, Linn. Mus. Cliff. 11. n. 4 (1736); Gen. ed. I. 382 (1787). *SCITAMINEAE*, Benth. & Hook. f. iii. 648.
Rendlia, Gieske, Pract. Linn. 199, 209 (1792).
Kua, Medic. in Act. Theod. Palat. vi. Phys. (1790) 394.
Stissera, Gieske, Pract. Linn. 207, 209 (1792).
aeruginosa, *Roxb. in As. Res.* xi. (1810) 353.—Burma.
albiflora, *Thw. Enum. Pl. Zeyl.* 316.—Zeylan.
Amada. *Roxb. in As. Res.* xi. (1810) 341.—Ind. or.
amarissima, *Roxb. Syst.* t. 101.—Ind. or.
americana, Lam. Encyc. ii. 228 = Calathea Allouia.
angustifolia, Dalz. & Gibs. Bomb. Fl. 274 = neilgherrensis.
angustifolia, *Roxb. in As. Res.* xi. (1810) 338. t. 3.—Reg. Himal.
aromatica, *Salisb. Parad. Lond.* t. 96.— Ind. or.
attenuata, *Wall. Cat.* n. 6803.—Burma.
australasica, *Hook. f. Bot. Mag.* t. 5620.—Austral.
caesia, *Roxb. in As. Res.* xi. (1810) 334.—Ind. or.
Careyana, Wall. Cft. n. 6596 = Hitchenia Careyana.
caulina, *J. Grah. Cat. Pl. Bomb.* 210.—Ind. or.
comosa, *Roxb. in As. Res.* xi. (1810) 336.—Burma.
cordata, Wall. Pl. As. Rar. i. 8 = petiolata
cordifolia, Roxb. Hort. Beng. 1, nomen = grac.
decipiens, *Dalz. in Hook. Kew Journ.* ii. (1850) 144.—Ind. or.
elata, *Roxb. Hort. Beng.* 1; *Fl. Ind.* i. 25.—Burma.
erubescens, Wall. Cat. n. 6606 = rubescens.
ferruginea, *Roxb. in As. Res.* xi. (1810) 336.—Ind. or.
glaucophylla, Wall. Cat. n. 6594 = Hitchenia glauca.
grandiflora, *Wall. ex Voigt, Hort. Suburb. Calc.* 565, nomen.—Malaya.
latifolia, *Roxb. Scitam.* t. 108.—Ind. or.
leucorhiza, *Roxb. in As. Res.* xi. (1810) 337.—Ind. or.
longa, *Linn. Sp. Pl.* 3.—An. trop.
longifora, Salisb. in Trans. Hort. Soc. i. (1812) 286 = rubescens.
montana, *Roxb. Pl. Coram.* ii. 28.—Ind. or.
musaces, *Wall. Cat.* n. 6596.—Ind. or.
neilgherrensis, *Wight, Ic.* t. 2006.—Ind. or.
officinalis, Salisb. in Trans. Hort. Soc. i. (1812) 285 = C. Zedoaria.
oligantha, *Trim in Journ. Bot.* xviii. (1885) 245.—Zeylan.
ornata, *Wall. ex Voigt, Hort. Suburb. Calc.* 564, nomen.—Burma.
pallida, *Lour. Fl. Cochinch.* 9.—China.
parviflora, *Wall. Pl. As. Rar.* i. 47. t. 57.—Burma.
petiolata, Hort.—Cf. Gard. Chron. (1871) 6 = seq.?
petiolata, *Roxb. Fl. Ind. ed. Carey*, i. 36.—Burma.
pilosa, *Wall. Cat.* n. 6611.—Burma.
porphyrostalca, *Zipp. ex Span. in Linnaea*, xv. (1841) 479, nomen.—Ins. Timor.
Pseudo-montana, J. Grah. Cat. Pl. Bomb. 210 = montana.
purpurascens, *Blume, Enum. Pl. Jav.* 46.—Jave.
reclinata, *Roxb. in As. Res.* xi. (1810) 342.—Ind. or.
Roscoeana, *Wall. Pl. As. Rar.* i. 8. t. 9.—Burma.
rotunda, Linn. Sp. Pl. 2 = Kaempferia pandurata.
rubescens, *Roxb. in As. Res.* xi. (1810) 336.—Ind. or.
rubricaulis, Link, Enum. Hort. Berol. i. 3 = rubescens.
spaciosa, Hort. Angl. ex Link, i. c. = C. Zedoaria.
strobilifera, *Wall. Cat.* n. 6599.—Burma.
strobilina, Wall. ex J. Grah. Cat. Pl. Bomb. 210 = strobilifera.
sumatrana, *Mig. Fl. Ind. Bat. Suppl.* 615.—Sumatra.
viridiflora, *Roxb. in As. Res.* xi. (1810) 341.—Sumatra.
xanthorrhiza, D. Dietr. Spn. Pl. i. 19 = seq.
xanthorrhiza, *Roxb. Hort. Beng* 1; *Fl. Ind.* i. 25.—Ins. Amboina.
Zedoaria, *Roxb. Scitam.* t. 109.—Reg. Himal.
Zedoaria, Roxb. in As. Res. xi. (1810) 333.—Cf. etiam 336 = aromatica.
Zerumbet, Roxb. l. c. 333 = C. Zedoaria.

CURMELLA, Roem. & Schult. Syst. v. p. xxii (1819). *ASCLEPIADEAE*
Curmella, Rafin. Sylva Tellur. 21 (1838).
Rhaedi, *Roem. & Schult. l. c.* 675.—Malaya.

CURMERIA, Linden & André, in Illustr. Hortic. xx. (1873) 45. t. 191 = Homalomena, Schott (Aroid.).
metallica, Hort.—Cf. Gard. Chron. (1884) i. 711.—Hab.?
picta, Hort.—Cf. l. c. (1874) 92 = seq.?
picturata, Linden & André, in Illustr. Hortic. xx. (1873) 45. t. 191 = H. picturata.
Roezlii, Mast. in Gard. Chron. (1874) ii. 804 = H. Roezlii.
Wallisii, Mast. l. c. (1877) i. 108 = H. Wallisii.

CURNILLA, Rafin. Sylva Tellur. 21 (1838) = Curinda, Roem. & Schult. (Asclep.?).
sarmentosa, Rafin. l. c. = Curinda Rheedi.

CURONDIA, Rafin. Sylva Tellur. 154 (1838) = Christmannia, Desnat. (Inc. sed.).
axillaris, Rafin. l. c. = Christmannia Courondi.

CURRORIA, Planch. ex Benth. in Hook. Niger Fl. 457 (1849). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 741.
decidua, *Planch. ex Benth. l. c.*—Afr. trop.

CURSONIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 429 = Oncosperma, DC. (Compos.).
peruviana, Nutt. l. c. = O. odorata.

CURTIA. Cham. & Schlecht. in Linnaea, i. (1826) 209. t. 4. *GENTIANEAE*, Benth. & Hook. f. ii. 808.
Anonymos, Griseb. Gen. et Sp. Gent. 168 (1839).
Schenkelia, Mart. Nov. Gen. et Sp. ii. 113. tt. 186 ad 188 (1827).
Tinhauserra, Fohl, ex G. Don, Gen. Syst. iv. 201 (1837).
diffusa, *Cham. in Linnaea*, viii. (1833) 14. in obs.—Bras.
gentianoides, *Cham. & Schlecht. l. c.* i. (1826) 210.—Bras.
tenella, *Cham. l. c.* viii. (1833) 13.—Bras.

CURTISIA, Ait. Hort. Kew. ed. I. i. 169 (1789). *CORNACEAE*, Benth. & Hook. f. i. 949.
Doratium, Soland. ex Jaume St. Hil. Expos. ii. 267 (1805).
Junghahnia, J. F. Gmel. Syst. 259 (1791).
Relhamia, J. F. Gmel. l. c. 247 (1791).
fagifolia, Salisb. Prod. 67 = faginea.
faginea, *Ait. Hort. Kew.* ed. I. i. 162.—Afr. austr.

CURTISIA. Schreb. Gen. 199 (1789) = Xanthoxylon, Linn. (Rutac.).
Schreberi, J. F. Gmel. Syst. 498 = Z. Ochroxylum?

CURTOGYNE, Haw. Rev. Pl. Succ. 8 (1821) = Crassula, Dill.
albiflora, Eckl. & Zeyh. Enum. 302 = Cr. albiflora.
Burmanniana, Eckl. & Zeyh. l. c. 304 = Cr. Burmanniana.
dejecta, DC. Prod. iii. 392 = Cr. dejecta.
flava, Eckl. & Zeyh. Enum. 303 = Cr. flava.
undata, Haw. Rev. Pl. Succ. 8 = Cr. dejecta.
undosa, Haw. in Phil. Mag. (1827) 134 = Rochea undosa.
undulata, Haw. Rev. Pl. Succ. 9 = Cr. dejecta.

CURTOISIA, Endl. Gen. Suppl. ii. 60, in syn. (1842) = COURTOISIA, Reichb. = Collomia, Nutt. (Polemon.).

CURTOPOGON, Beauv. Agrost. 89. t. 8. f. 7 (1812) = Aristida, Linn. (Gramin.).
dichotomus, Beauv. l. c. = A. dichotoma.

CUSCUARIA, Schott, in Bonplandia, v. (1857) 45 = Sciadapsus, Schott (Aroid.).
marantifolia, Schott, Gen. Aroid. t. 80 = S. Cuscuaria.
Rumphii, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 120 = S. Cuscuaria.
sparia, Schott, in Bonplandia, ix. (1861) 367 = S. Cuscuaria.

CUSCUTA. [Tourn.] Linn. Syst. ed. I (1735). *CON-VOLVULACEAE*, Benth. & Hook. f. ii. 891.
Anthanema, Rafin. Fl. Tellur. iv. 99 (1836).
Aplostylis, Rafin. l. c. 91 (1836).
Bochingera, F. Schultz, in Jahrb. Pharm. (Mar. 1847).—Cf. Bot. Zeit. vi. (1848) 799.
Cabuzea, Des Moul. Etudes Org. Cusc. 66 (1868).

CUSCUTA:—
Cassytha, Gesner, ex S. F. Gray, Nat. Arr. Brit. Pl ii. 345 (1821).
Cuscutina, Pfeiff. in Bot. Zeit. iv. (1846) 492.
Dastylepis, Rafin. Fl. Tellur. iv. 126 (1836).
Engelmannia, Pfeiff. in Bot. Zeit. iii. (1845) 673.
Epilinella, Pfeiff. l. c. (1845).
Epithymum, Opiz, Seznam, 40 (1852).
Eronema, Rafin. Fl. Tellur. iv. 125 (1836).
Grammica, Lour. Fl. Cochinch. 170 (1790).
Kadula, Rafin. Fl. Tellur. iv. 90 (1836).
Kadurias, Rafin. l. c. 91 (1836).
Lepidanche, Engelm. in Am. Journ. Sc. xliii. (1842) 343. t. 6.
Lepimenes, Rafin. Fl. Tellur. iv. 91 (1836).
Monogynella, Des Moul. Etudes Org. Cusc. 65 (1868).
Nemepis, Rafin. Fl. Tellur. iv. 91 (1836).
Pentaxe, Rafin. l. c. 90 (1836).
Pfeiffera, Buching. in Ann Sc. Nat. Sér. III. v. (1846) 88.
Schrebera, Linn. Sp. Pl. ed. II. 1662 (1763).
Succuta, Des Moul. Etudes Org. Cusc. 74 (1868).
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 78.—Abyss.
acuta, Rafin. Ann. Nat. (1820) 13 = compacta.
acuminata, Nutt. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 498 = californica.
acuminata, *Pomel, Nouv. Mat. Fl. Atl.* 88.—Algeria.
acuta, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 497.—Am. trop.
acutiflora, Kotz, in Parl. Giorn. Bot. Ital. ii. (1847) 286 = C. Epithymum.
acutiloba, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 478.—Am. trop.
acutissima, Buching. ex Engelm. l. c. 490 = hyalina.
africana, Choisy, Cusc. 176 = angulata.
africana, Eckl. & Zeyh. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 475 = nitida.
africana, Krauss, ex Engelm. l. c. 503 = appendiculata.
africana, *Willd. Sp. Pl.* i. 708.—Afr. austr.
aggregata, Roxb. Hort. Beng. 12; Fl. Ind. i. 447 = europaea.
alba, J. & C. Presl, Delic. Prag. 87 = C. Epithymum.
alpina, Hohen. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 479 = kurdica.
alpina, Kotschy, ex Choisy, in DC. Prod. ix. 458 = minor.
americana, Hook. Comp. Bot. Mag. i. (1835) 178 = glomerata.
americana, Hook. Fl. Bor. Am. ii. 77 = obtusiflora.
americana, Jacq. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 484 = corymbosa.
americana, *Linn. Sp. Pl.* 124.—Am. bor.
americana, Poepp. ex Choisy, in DC. Prod. ix. 453 = microstylis.
americana, Thunb. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 479 = chinensis.
americana, Thunb. Prod. Pl. Cap. 32 = africana.
angulata, Edgew. in Trans. Linn. Soc. xx. (1851) 57 = reflexa.
angulata, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 474.—Afr. austr.
aphylla, Rafin. in Am. Monthly Mag. (1818) 40 = glomerata?
aphylla, Steud. Nom. ed. I. 246 = chinensis.
appendiculata, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 503.—Afr. austr.
applanata, *Engelm. l. c.* 479.—Am. bor.
approximata, Bab. in Ann. & Mag. Nat. Hist. xiii. (1844) 253 = planiflora.
arabica, *Fresen. in Mus. Senckenb.* i. (1834) 165.—Orient.
arabica, Wight, Ic. t. 1371 = hyalina.
arvensis, *Beyrich, ex Hook. Fl. Bor. Am.* ii. 77.—Am. bor.
asiatica, Pall. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 465 = planiflora.
astyla, Engelm. in Bot. Zeit. iv. (1846) 278 = monogyna.
aurantiaca, Req. ex Bertol. Fl. Ital. vii. 632 = obtusiflora.
aurea, *Liebm. in Forhandl. Skand. Naturf. Christiania*, (1847) 198.—Hab.?
australis, *R. Br. Prod.* 491.—Austral.
australis, Hook. f. Fl. Tasm. i. 278 = tasmanica.
babylonica, *Aschers. ex Choisy, Cusc.* 174. t. 1. f. 1. —Assyria.
Balbisiana, Bubn. & Kunz. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 499 = planiflora.

CUSCUTA :—
Bidentia, *Berthiot, in Billotia*, t. (1864) 15.—Gallia.
Boissieri, Stocks, in Hook. Kew Journ. iv. (1852) 173 = hyalina.
bonariensis, Hort. ex Engelm. in Bot. Zeit. iv. (1846) 375 = Cronovii.
brachystyla, C. Koch, in Linnæa, xxii. (1849) 747 = europæa.
bracteata, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 509.—Bras.
bractеosa, Gaspar. ex Engelm. l. c. 467 = planiflora.
brevistyla, A. Braun, ex A. Rich. Tent Fl. Abyss. ii. 79 = planiflora.
Burmannii, Choisy, Cusc. 177 = stilida.
californica, *Choisy, Cusc.* 183 ; *Hook. & Arn. Bot. Beech. Voy.* 364.—Calif.
calliopes, Engelm. in Trans. Acad. St. Louis, i. (1859) 462 = C. Epithymum.
callosa, *Pomel, Nouv. Mat. Fl. Atl.* 88.—Algeria.
calycina, Webb & Berth. Phyt. Canar. iii. 87 = planiflora.
campanulata, Nutt. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 482 = americana.
campanulata, Stokes, Bot. Mat. Med. i. 239 = C. Epithymum.
canariensis, Choisy, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 467 = planiflora.
capensis, Choisy, Cusc. 175 = africana.
capillaris, Edgew. in Trans. Linn. Soc. xx. (1851) 86 = europæa.
capillaris, Reichb. Pl. Crit. v. 64 = palæstina.
capillaris, Wall. Cat. n. 1821 = chinensis.
capitata, *Roxb. Hort. Beng.* 13 ; *Fl. Ind.* i. 448.—Reg. Himal.
carinata, R. Br. Prod. 491 = chinensis.
Cassiope, Heldr. & Sart. in Boiss. Diagn. Ser. II. iii. 128 = C. Epithymum.
cassytoides, *Nees, in Linnæa*, xx. (1847) 196, nomen ; *Engelm. in Trans. Acad. St. Louis*, i. (1859) 513.—Afr. austr.
ceanothi, *Behr, in Proc. Calif. Acad.* ed. II. i. (1873) 16.—Calif. Engelm. in Am. Journ. Sc. xliii. (1842) 336. t. 6. f. 1-6 = tenuiflora.
Cesatiana, Bertol. Fl. Ital. vii. 623 = australis.
chilensis, Bert. ex Choisy, in DC. Prod. ix. 456 = racemosa.
chilensis, Hort. Frib. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 508 = Cronovii.
chilensis, *Ker-Gawl. in Bot. Reg.* t. 603.—Chili.
chinensis, *Lam. Encyc.* ii. 229.—China.
chlorocarpa, *Engelm. in A. Gray, Man. Bot. U. St.* ed. I. 350.—Am. bor.
Chrysocoma, Welw. ex Des Moul. Etudes Org. Cusc. 71 = australis.
ciliaris, Hohen. in Boiss. Diagn. Ser. II. iii. 129 = chinensis.
ciliaris, Kotschy, ex Choisy, in DC. Prod. ix. 453 = pedicellata.
cilvicola, Schlecht. in Linnæa, xxii. (1849) 808 = racemosa.
colorans, Maxim. Prim. Fl. Amur. 201 = japonica.
compacta, *Juss. ex Choisy, Cusc.* 185. t. 4. f. 2.—Am. bor.
congesta, Benth. Bot. Voy. Sulph. 138 = americana.
congesta, Beyr, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 502 = inflexa.
corniculata, *Engelm. l. c.* 504.—Am. trop.
corensis, Beyr, ex Hook. Comp. Bot. Mag. i. (1835) 178 = compacta.
Cospii, Engelm. in Am. Journ. Sc. xliii. (1842) 337 = inflexa.
corymbosa, Choisy, Cusc. 180 = racemosa.
corymbosa, Juss. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 478 = foetida.
corymbosa, *Ruiz & Pav. Fl. Per.* i. 69.—Peruv.
crassulata, Engelm. in Hook. Lond. Journ. Bot. ii. (1843) 197 = compacta.
cristata, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 507.—Am. trop.
cupulata, Engelm. in Bot. Zeit. iv. (1846) 378 = planiflora.
cuspidata, *Bingelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 226.—Texas.
cuspidata, *Pomel, Nouv. Mat. Fl. Atl.* 87.—Algeria.
cyanea, *Willd. ex Roem. & Schult. Syst. vi.* 205 = corymbosa.

CUSCUTA :—
decora, *Choisy, ex Engelm. in Trans. Acad. St. Louis*, i. (1859) 501.—Am. trop.
denatiflora, *Hook. f. Fl. N. Zel.* i. 186.—N. Zel.
densiflora, Soyer-Willem. in Mém. Soc. Linn. Par. i. (1823) 98 ; et iv. (1826) 281 = C. Epilinum.
denticulata, *Engelm. in Am. Natural.* ix. (1875) 346.—Am. bor. occ.
desertorum, Matt. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 488 = umbellata.
diaphana, Wender. Fl. Hassiac. 364 = racemosa.
dilatior, Choisy, Cusc. 177 = reflexa.
elegans, *Boiss. & Bal. Diagn.* Ser. II. iii. 129.—Persia.
elegans, Noé, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 468 = planiflora.
epicnidea, Bernh. in Thuer. Gartenz. (1844) n. 4 = europæa.
Epilinum, *Weihe, in Archiv Apoth.* viii. (1824) 54.—Europ., Afr. or.
Epilocanum, Webb, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 466 = planiflora.
Episcochum, Webb & Berth. Phyt. Canar. iii. 86 = planiflora.
Epispartos, Boiss. & Orph. ex Boiss. Fl. Orient. iv. 116 = planiflora.
Epithymiphyta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 194 = C. Epithymum.
Epithymum, Bové, ex Choisy, in DC. Prod. ix. 453 = europæa.
Epithymum, *Murr. Syst.* ed. XIII. 140.—Europ. ; Ax. bor.
Epithymum, Sievers, ex Ledeb. Fl. Alt. i. 293 = pedicellata.
Epithymum, Thuill. Fl. Par. 85 = europæa.
Epithymum, Webb & Berth. Phyt. Canar. iii. 86 = planiflora.
Epitriphyllum, Bernh. in Thuer. Gartenz. (1844) n. 4 = europæa.
europæa, Bové & Engelm. in Trans. Acad. St. Louis, i. (1859) 467 = planiflora.
europæa, *Linn. Sp. Pl.* 124.—Europ. ; Orient.
exalata, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 513.—Am. bor. occ.
exigua, Herb. Par. ex Engelm. l. c. 480 = chinensis.
filiformis, Lam. Fl. Fr. ii. 307 = C. Epithymum, europæa.
fimbricata, Bunge, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 480 = chinensis.
flava, Sievers, ex Ledeb. Fl. Alt. i. 294 = lupuliformis.
floriunda, *H. B. & K. Nov. Gen. et Sp.* iii. 123.—Mexic.
floribunda, *Phil. Fl. Atac.* 37.—Chili.
foetuda, Hook. & Arn. Bot. Beech. Voy. 364 = gracillima.
foetida, *H. B. & K. Nov. Gen. et Sp.* iii. 122.—Ecuador.
fruticosa, Bertol. in Mem. Acad. Sc. Bolog. ii. (1850) 312 = compactum.
funiformis, Willd. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 519 = africana.
fusiformis, Willd. ex Roem. & Schult. Syst. vi. 205 = africana.
gamostyla, Engelm. in Trans. Acad. St. Louis, i. (1859) 513 = exaltata.
gigantea, *Griff. Notul.* i. 243.—Reg. Himal.
globiflora, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 530.—Bolivia.
globularis, Bertol. Fl. Ital. viii. 625 = palæstina.
globularis, Nutt. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 494 = arvensis.
globulosa, Benth. Bot. Voy. Sulph. 138 = americana.
globulosa, *Boiss. & Reut. Diagn.* Ser. II. iii. 126.—Ax. Min. ; Persia.
glomerata, *Choisy, Cusc.* 184. t. 4. f. 1.—Am. bor.
Godronii, Des Moul. Etudes Org. Cusc. 60 = planiflora.
gracillima, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 488.—Am. trop.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 123.—N. Granat.
grandiflora, Wall. Cat. n. 1818 = reflexa.
Grassii, *Delp. in Mem. Acc. Tor.* Ser. II. xxvi. (1871) 132.—Hongkong.
gronovioïna, H. B. & K. Nov. Gen. et Sp. iii. 122 = americana.
Gronovii, *Willd. ex Roem. & Schult. Syst.* vi. 205.—Am. bor.

CUSCUTA :—
Guyoni, Gaspar. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 468 = C. Epithymum.
gymnocarpa, *Engelm. l. c.* 496.—Am. trop.
halophila, Engelm. l. c. 469 = europæa.
halophyta, Fries, Nov. Fl. Suec. Mant. i. 8 = europæa.
hassiaca, Pfeiff. in Bot. Zeit. i. (1843) 705 = racemosa.
hispidula, Engelm. in Am. Journ. Sc. xlv. (1843) 75 = decora.
Hookeri, Sweet, Hort. Brit. ed. I. 290 = reflexa.
hyalina, Boiss. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 469 = europæa.
hyalina, *Roth, Nov. Pl. Sp.* 100.—Ind. or.
hyalina, Wight, Ic. t. 1872 = chinensis.
imbricata, Nutt. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 511 = compacta.
incisa, Cholay, Cusc. 179. t. 2. f. 2 = corymbosa.
incurvata, *Prog. in Mart. Fl. Bras.* viii. 379.—Am. trop.
indecora, Choisy, Cusc. 182. t. 3. f. 3 = decora.
inflexa, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 502.—Am. bor.
inodora, Willd. ex Prog. in Mart. Fl. Bras. vii. 380 = obtusiflora.
intermedia, Choisy, Cusc. 179. t. 2. f. 3 = odorata.
jalapensis, *Schlecht. in Linnæa*, viii. (1833) 515.—Mexic.
japonica, *Choisy, in Zoll. Syst. Vers. Ind. Archip.* 130, 134.—China ; Japon.
Kotschyana, *Boiss. Diagn.* Ser. I. vii. 99.—Persia.
Kotschyi, Des Moul. Etudes Org. Cusc. 56 = C. Epithymum.
kurdica, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 470.—Kurdist.
lasiflora, Benth. Bot. Voy. Sulph. 138 = corymbosa.
Lehmanniana, *Bunge, Lehm. Reliq. Bot.* 220.—Ax. centr.
leiolepis, Miq. in Linnæa, xviii. (1844) 247 = americana.
lentis, *Staff, in Denkschr. Acad. Wien*, i. (1885) 21.—Persia.
leptantha, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 489.—Am. trop.
leucosphæra, Boiss. & Heldr. ex Engelm l. c. 465 = planiflora.
Ligustri, Areschoug, Revis. Cusc. Suec. 17 = europæa.
lupuliformis, *Krock. Fl. Sil.* i. 261. t. 36.—Europ.
macrantha, G. Don, Gen. Syst. iv. 306 = reflexa.
macranthera, Heldr. & Sart. in Boiss. Diagn. Ser. II. iii. 128 = C. Epithymum.
macrocarpa, *G. Don, ex Loud. Hort. Brit. Suppl.* iii. 580.—Sibiria.
macrostyla, Decne. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 464 = abyssinica.
major, Gilib. Fl. Lituan. i. 18 = europæa.
major, Koch & Zis. Cat. Palat. 5 = C. Epilinum.
megalantha, Steud. Nom. ed. II. i. 455 = reflexa.
micrantha, *Choisy, Cusc.* 176. t. 1. f. 8.—Am. trop.
micrantha, Matt. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 506 = parviflora.
micrantha, Tineo, in Guss. Fl. Sic. Syn. ii. 887 = palæstina.
microcephala, D'Escayrac, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 467 = planiflora.
microcephala, *Pomel, Nouv. Mat. Fl. Atl.* 88.—Algeria.
microcephala, Welw. ex Des Moul. Etudes Org. Cusc. 76 = C. Epithymum.
microstyla, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 506.—Am. trop.
Milletti, Hook. & Arn. Bot. Beech. Voy. 201 = obtusiflora.
miniacea, Choisy, in DC. Prod. ix. 458 = seq.
miniata, Mart. Reise Bras. i. 286 = racemosa.
minor, Choisy, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 470 = kurdica.
minor, Gilib. Fl. Lituan. i. 18 = planiflora.
minor, S. F. Gray, Nat. Arr. Brit. Pl. ii. 345 = C. Epithymum.
mitrasformis, *Engelm. ex Hemsl. Diagn. Pl. Nov.* 54.—Mexic.
monogyna, Auct. Fl. Germ. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 516 = lupuliformis.
monogyna, Schmidt, ex Engelm. l. c. 469 = europæa.
monogyna, *Vahl, Symb. Bot.* ii. 32.—Europ. ; Orient.
Mueller, Steud, in Bull. Soc. Bot. Belg. ii. (1863) 336, 337 = C. Epithymum.
neuropetala, Engelm. in Am. Journ. Sc. xlv. (1843) 75 = decora.

CUSCUTA :—
alsidia, *E. Mey. in Drège, Zwei Pfl. Decum.* 176.—Afr. austr.
obtusiflora, H. B. & K. Nov. Gen. et Sp. iii. 122 = australis.
odontolepis, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 486.—Am. trop.
odorata, Poepp. ex Choisy, in DC. Prod. ix. 455 = chilensis.
odorata, *Ruiz & Pav. Fl. Per.* i. 69. t. 105.—Peruv.
oxycarpa, Engelm. in Trans. Acad. St. Louis, i. (1859) 508 = rostrata.
oxypetala, Bolss. Diagn. Ser. II. iii. 130 = hyalina.
palaestina, *Boiss. l. c.* Ser. I. xi. 86.—Graecia; Syria.
paradoxa, Rafin. Am. Nat. (1820) 13 = glomerata.
partita, *Choisy, Cusc.* 188. t. 5. f. 3.—Am. trop.
parviflora, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 508.—Am. trop.
parviflora, Nutt. ex Engelm. l. c. 502 = inflexa.
parviflora, Willd. ex Engelm. l. c. 487 = umbellata.
patens, *Benth. Bot. Voy. Sulph.* 35.—Calif.
pauciflora, *Phil. in Linnaea*, xxxiii. (1864–65) 185.—Chili.
pedicellata, *Ledeb. Fl. Alt.* i. 293. t. 284.—Sibir. altaic.
pedunculosa, Kotschy, ex Choisy, in DC. Prod. ix. 458 = babylonica.
pentagona, Engelm. in Am. Journ. Sc. xliii. (1842) 340 = arvensis.
pentandra, Heyne, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 519 = reflexa.
persica, Decne. ex Engelm. l. c. 470 = Kotschyana.
planiflora, Koch, Syn. Fl. Germ. ed. II. 570 = C. Epithymum.
planiflora, *Tenore, Fl. Nap.* iii. 250.—Hispan.
platyloba, *Prog. in Mart. Fl. Bras.* vii. 381.—Am. trop.
polyantha, Shuttl. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 508 = Gronovii.
Polygonorum, Cesati, in Cat. Sem. Genev. (1849) 22 ; Linnaea, xxiv. (1851) 199 = obtusiflora.
Polygonorum, Engelm. in Am. Journ. Sc. xliii. (1842) 343. t. 6 = chlorocarpa.
popayanensis, H. B. & K. Nov. Gen. et Sp. iii. 123 = corymbosa.
popayanensis, Poepp. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 500 = micrantha.
popayanensis, Poepp. ex Engelm. l. c. 505 = racemosa.
potosina, *Schaffn. ex S. Wats. in Proc. Am. Acad.* xvii. (1882–83) 194.—Mexic.
prismatica, *Pav. ex Choisy, in Cusc.* 182. t. 3. f. 2.—Ecuador.
puberula, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1154, nomen.—Guiana.
pulchella, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 472.—Rossia ; As. bor.
pulcherrima, Schiede, in Linnaea, xi. (1848) 750 = decaea.
pycnantha, Benth. Pl. Hartw. 326 = foetida.
racemosa, *Mart. Reise Bras.* i. 286.—Bras.
reflexa, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 394 = tinctoria.
reflexa, *Roxb. Pl. Corom.* ii. 8. t. 104.—Ind. or.
remotiflora, Bertol. Misc. Bot. x. 29 = compacta.
Ragowitschiana, Trautv. in Bull. Phys.-Math. Acad. Pétersb. xiii. (1855) 378 = australis.
rosea, Jacquem. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 473 = capitata.
rostrata, *Shuttlw. ex Engelm. & A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 295.—Am. bor.
salsa, *Engelm. in S. Wats. Bot. Calif.* i. 536.—Am. bor. occ.
sandwichiana, *Choisy, Cusc.* 184. t. 5. f. 4.—Ins. Sandvic.
Saxatilis, Engelm. in Am. Journ. Sci. xliii. (1842) 339 = Gronovii.
scandens, Brot. Fl. Lusit. 206 = lepuiformis.
schiedeana, Boiss. Diagn. Ser. I. xi. 86 = planiflora.
Schkuhriana, Pfeiff. in Bot. Zeit. iii. (1845) 673 = europaea.
segetum, Rots, in Parl. Giorn. Bot. Ital. ii. Ann. 11. (1847) 286 = europaea.
sicula, Tineo, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 486 = planiflora.
Sideroxea, Liebm. in Forhandl. Skand. Naturf. Christiania (1847), 196 = gracillima.
Solmsi, *Heinsy, in Oestr. Bot. Zeitschr.* xxiv. (1874) 304.—Austria.
spectabilis, Choisy, Cusc. 187. t. 5. f. 1 = americana.

CUSCUTA :—
squamata, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 510.—Am. bor.
stenolepis, *Engelm. l. c.* 508.—Am. trop.
strobilacea, *Liebm. in Forhandl. Skand. Naturf. Christiania* (1847), 194.—Hab. ?
stylosa, Choisy, Cusc. 187. t. 5. f. 2 = corymbosa.
suaveolens, Ser. in Ann. Sc. Phys. Nat. Agric. et Indust. iii. (1840) 519 = racemosa.
subinclusa, *Dur. & Hilg. in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 65.—Am. bor.
subtilis, Chaub. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 489 = gracillima.
subulata, Tineo, in Guss. Fl. Sic. Syn. ii. 888 = C. Epithymum.
subuniflora, *C. Koch, in Linnaea*, xxii. (1849) 747.—As. Min.
sulcata, Roxb. Hort. Beng. 12 ; Fl. Ind. i. 447 = chinensis.
sulcata, Wall. Cat. n. 1320 = hyalina.
surinamensis, Schilling, Desc. Pl. Lepr. 200. t. 2 = americana.
syntyla, Maxim. Prim. Fl. Amur. 200 = japonica.
tasmanica, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 512.—Austral.
tenuiflora, *Engelm. in A. Gray, Man.* ed. I. 350.—Am. bor.
tetrandra, Moench, Meth. 461 = europaea.
tetrasperma, Jan, ex Engelm. in Trans. Acad. St. Louis, i. (1859) 469 = europaea.
timorensis, *Decne ex Engelm. l. c.* 514.—Ins. Timor.
tinctoria, *Mart. ex Engelm. l. c.* 480.—Mexic.
Tinei, Insengs, in Tineo, Pl. Rar. Sicil. 14 = obtusiflora.
trichostyla, *Engelm. in Trans. Acad. St. Louis*, i. (1859) 495.—Am. trop.
trifora, E. Mey. ex Engelm. l. c. 519 = africana.
Trifolii, Bab. in Phytolog. i. (1848) 467 = C. Epithymum.
Triumvirati, *Lange, in Kjoeb. Vidensk. Meddel.* (1851) 98.—Iberia.
tuberosa, J. & C. Presl, Delic. Prag. 215 = europaea.
Ulicis, *Godr. in Mém. Soc. Sc. Nat. Cherb.* xix. (1875)
umbellata, *H. B. & K. Nov. Gen. et Sp.* iii. 121.—Am. austr.
umbrosa, Beyrich, ex Engelm. in A. Gray, Man. ed. I. 351 = reflexa.
umbrosa, Beyrich, ex Hook. Fl. Bor. Am. ii. 78 = Gronovii.
urceolata, Kunze, in Flora, xxix. (1846) 651 = planiflora.
urceolata, Stokes, Bot. Mat. Med. i. 288 = europaea.
verrucosa, Engelm. in Am. Journ. Sc. xliii. (1842) 341. t. 6. f. 25 = arvensis.
verrucosa, Sweet, Brit. Fl. Gard. Ser. I. i. t. 6 = reflexa.
Viciae, F. Sebults, ex Des Moul. Etudes Org. Cusc. 77 = europaea.
vulgaris, Pers. Syn. i. 289 = europaea.
vulgaris, J. & C. Presl, Fl. Cech. 56 = C. Epilinum.
vulgivaga, Engelm. in Am. Journ. Sc. xliii. (1842) 338. t. 6 = arvensis.
xanthochortos, *Mart. ex Engelm. in Trans. Acad. St. Louis*, i. (1859) 486.—Bras.
xanthosiana, Hort. Par. ex Engelm. l. c. 462 = C. Epithymum.

CUSCUTINA, Pfeiff. in Bot. Zeit. iv. (1846) 492 = Cuscuta, Linn. (Convolv.).
cuscutoides, Pfeiff. l. c. = Cuscuta racemosa.

CUSPA, Humb. Relat. Hist. i. 366 (1814) = **Alsodeia**, Thou. (Violar.).

CUSPARIA, D. Dietr. Syn. ii. 1478 (1840) = CASPARIA, Kunth = **Banhinia**, Linn. (Legum.).

CUSPARIA, Humb. Tabl. Géog. ex R. Br. in Flind. Voy. ii. App. iii. 545 (1814). *RUTACEAE* [Benth. & Hook. f. i. 285].
ANGOSTURA, Roem. & Schult. Syn. iv. 188 (1819).
BONPLANDIA, Willd. in *Mém. Acad. Berl.* (1809) 54.
CONCHOCARPUS, Mikan, Delect. fasc. i. t. 2 (1820).
DAUBENTONIA, Vell. Fl. Flum. 97 (1825) i. t. 66 (1827).
DIGLOTTIS, Neas & Mart. in Nov. Act. Nat. Cur. xi. (1825) 151, 176.
LARIOSTEMON, Neas & Mart. l. c. (1825) 129, 171.
OBERTONIA, Vell. Fl. Flum. 16 (1825) i. t. 46 (1827).
ROSENIA, Vell. l. c. 21 (1825) ; i. t. 77 (1827).

CUSPARIA :—
Candolleana, *Engl. in Mart. Fl. Bras.* xii. ii. 116.—Bras.
cuneifolia, *Engl. l. c.* 116.—Bras.
elegans, *Engl. l. c.* 121.—Bras.
febrifuga, *Humb. Tabl. Géogr. ex DC. in Mém. Mus. Par.* ix. (1822) 144.—N. Granat.
Fontanesiana, *Engl. in Mart. Fl. Bras.* xii. ii. 122.—Bras.
Gaudichaudiana, *Engl. l. c.* 116.—Bras.
grandiflora, *Engl. l. c.* 113.—Bras.
heterophylla, *Engl. l. c.* 114.—Bras.
macroanpa, *Engl. l. c.* 121.—Bras.
macrophylla, *Engl. l. c.* 115.—Am. trop.
Martiana, *Engl. l. c.* 119.—Bras.
nodosa, *Engl. l. c.* 122.—Bras.
obovata, *Engl. l. c.* 118.—Bras.
odoratissima, *Engl. l. c.*—Bras.
ovata, *Engl. l. c.* 117.—Bras.
paniculata, *Engl. l. c.* 120.—Bras.
pentagyna, *Engl. l. c.* 122.—Bras.
pentandra, *Engl. l. c.* 120.—Bras.
ramiflora, *Engl. l. c.* 111.—Bras.
silvestris, *Engl. l. c.* 118.—Bras.
toxiaria, *Engl. l. c.* 114.—Bras.
trifoliata, Engl. l. c. 118 = febrifuga.

CUSPIDARIA, DC. in Bibl. Univ. Genèv. xvii. (1838) 192 [Rev. Bign. 9]. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1036.
LOCHMOCYDIA, Mart. ex DC. Prod. ix. 177 (1845).
MANSOA, DC. in Bibl. Univ. Genèv. xvii. (1838) [Rev. Bign. 13].
angustidens, *DC. Prod.* ix. 179.—Bras.
callistegioides, *DC. l. c.* 178.—Bras.
campanulata, *DC. l. c.* 179.—Bras.
crubescens, *Bureau, Monog. Bignon.* 5. t. 2.—Bras.
fasciculata, Sond. in Linnaea, xxii. (1849) 660 = Arrabidaea fasciculata.
lateriflora, *DC. Prod.* ix. 179.—Bras.
multiflora, *DC. l. c.*—Bras.
paniculata, *DC. l. c.* 179.—Bras.
populnea, Mart. ex DC. l. c. 178 = trifoliata.
pterocarpa, DC. l. c. = trifoliata.
puberula, *Mart. Herb. ex DC. l. c.* 178.—Bras.
ramentacea, *Mart. ex DC. l. c.*—Bras.
simplicifolia, *DC. Rev. Bign.* 9.—Bras.
trifoliata, *DC. l. c.*—Bras.

CUSPIDARIA, Link, Handb. ii. 315 (1831) = **Brysimum**, Tourn. (Crucif.).
Bieberstreinii, Rupr. Fl. Cauc. 74 = E. cuspidatum.
cheiranthoides, Link, Handb. ii. 315 = E. cuspidatum.
stenocarpa, Rupr. Fl. Cauc. 75, in textu = E. cuspidatum.

CUSPIDIA, Gaertn. Fruct. ii. 454. t. 171 (1791) = **Berkheya**, Ehrh. (Compos.).
araneosa, Gaertn. l. c. = Didelta cernua.
castrata, Cass. in Dict. Sc. Nat. xii. 252 = Didelta cernua.

CUSPIDOCARPUS, Spenn. in T. Nees, Gen. Fl. Germ. Canop. ii. n. 16 (1848) = **Micromeria**, Benth. (Labiat.).
rupestris, Spenn. l. c. = M. rupestris.

CUSSAMBIUM, [Remph.] Bach.-Ham. in Mem. Wern. Soc. v. ii. (1823) 256 = **Schlechtendera**, Willd. (Sapind.).
glabrum, Bach.-Ham. l. c. = S. trijuga.
pubescens, Bach.-Ham. l. c. 257 = S. trijuga.
spinosum, Bach.-Ham. l. c. 256 = S. trijuga.

CUSSO, Bruce, Trav. v. t. 22, 23 (1790) = **Brayera**, Kunth (Rosac.).

CUSSONIA, Comm. ex Endl. Gen. 1095, in syn. (1840) = **Bilsea**, Cambess. (Hyperic.).

CUSSONIA, Thunb. in Nov. Act. Soc. Sc. Upsa. iii. (1780) 210. t. 12, 13. *ARALIACEAE*, Benth. & Hook. f. i. 944.
QUESNEYA, D. Dietr. Syn. ii. 926 (1840).
SPHAEROSCHEMUM, Seem. in Journ. Bot. iii. (1865) 33. t. 36.
angolensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 82.—Afr. trop.

CUSSONIA :—
arborea, *Hochst. ex A. Rich. Tent Fl. Abyss.* i. 356.—Abyss.
Bartsii, *Seem. in Journ. Bot.* iv. (1866) 299.—Afr. trop.
Hogerii, *Seem. l. c.* 298.—Madag.
calophylla, *Miq. in Ann. Sc. Nat.* Sér. III. i. (1844) 36.—Afr. austr.?
dioica, *Verdik. ex Panch. & Sebert, Not. Bois Nouv. Calédon.* 208.—N. Caled.
fraxinifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 157.—Madag.
Gerardii, *Seem. in Journ. Bot.* iv. (1866) 298.—Afr. austr.
Kirkii, *Seem. l. c.* 299.—Afr. trop.
Kraussii, *Hochst. in Flora*, xxvii. (1844) 431.—Afr. austr.
Lessonii, A. Rich. Fl. N. Zél. i. 285 = Pseudopanax Lessonii.
monophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 156.—Madag.
myriantha, *Baker, l. c.* 157.—Madag.
natalensis, *Sond. in Harv. & Sond. Fl. Cap.* ii. 568.—Afr. austr.
paniculata, *Eckl. & Zeyh. Enum.* 355.—Afr. austr.
paniculata, E. Mey. in Drège, Zwei Pfl. Docum. 176 = umbelliflera.
racemosa, *Baker, in Journ. Linn. Soc.* xx. (1883) 156.—Madag.
spicata, *Thunb. in Nov. Act. Soc. Sc. Ups.* iii. (1780) 213. t. 13.—Afr. austr.
thyrsiflora, *Thunb. l. c.* t. 12.—Afr. austr.
thyrsoidea, Thunb. Nov. Gen. i. 11 = thyrsiflora.
tripteris, *Colla, Hort. Ripul.* 43. t. 26.—Afr. austr.
umbelliflora, *Sond. in Linnaea*, xxiii. (1850) 49.—Afr. austr.
vanteliana, *Baker, in Journ. Linn. Soc.* xx. (1883) 156.—Madag.

CUSSUTHA, Benth. & Hook. f. Gen. ii. 881 (1876) sphalm. = CASSUTHA, Des Moul. = **Cuscuta**, Linn. (Convolv.).

CUSTENIA, Steud. Nom. ed. I. 247 (1821) = seq.

CUSTINIA, Neck. Elem. ii. 223 (1790) = **Salsola**, Linn. (Hippocrat.).

CUTANDIA, Willk. in Bot. Zeit. xviii. (1860) 130. *GRAMINEAE*, Benth. & Hook. f. iii. 1187.
divaricata, *Benth. in Journ. Linn. Soc.* xix. (1881) 118.—Reg. Medit.
incrassata, *Benth. l. c.*—Reg. Medit.
lanceolata, *Benth. l. c.*—Reg. Medit.
maritima, *Benth. l. c.*—Reg. Medit.
memphitica, Benth. l. c. = scleropoides.
phillisaea, *Benth. l. c.*—Reg. Medit.
scleropoides, *Willk. in Bot. Zeit.* xviii. (1860) 130.—Hispan.

CUTARIA, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 66 = **Coutarea**, Aubl. (Rubiac.).

CUTARIA, Jeanne St. Hil. Expos. i. 443 (1805) = præc.

CUTLERA, Rafin. in Am. Monthly Mag. (1818) 176 = seq.

CUTTERA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Gentiana**, Tourn.
Catesbæi, Rafin. in Am. Monthly Mag. (1818) 176 = G. Andrewsii!
ochroleuca, Rafin. in Med. Repos. N. York, v. (1808) 353 = G. ochroleuca.
Saponaria, Rafin. l. c. = G. Saponaria.

CUTTHIA, F. Muell. Fragm. v. 47. t. 40 (1865). *SAXIFRAGEAE*, Benth. & Hook. f. i. 1004.
viburnea, *F. Muell. l. c.*—Austral.

CUTUBEA, Jeanne St. Hil. Expos. i. 332 (1805) = **Contorbea**, Aubl. (Gentian.).

CUVERACEAE, Jones, in As. Res. iv. (1795) 231 = **Cedrela**, P. Br. (Meliac.).

CUVIERA, DC. in Ann. Mus. Par. ix. (1807) 392. t. 22. *RUBIACEAE*, Benth. & Hook. f. ii. 112.
acutiflora, DC. l. c.—Afr. trop.
africana, Spreng. Syst. i. 760 = acutiflora.

CUVIERA :—
asiatica, Spreng. Syst. i. 760 = Stigmanthus cymosus.
longiflora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 157.—Afr. trop.
subuliflora, *Benth. in Hook. Niger Fl.* 407.—Afr. trop.
trilocularis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 157.—Afr. trop.

CUVIERA, Koel. Gram. Gall. et Germ. 328 (1809) = **Hordeum**, Linn. (Gramin.).
europaea, Koel. l. c. = H. sylvaticum.

CYAMOPSIS, DC. Mém. Lég. 280 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 498.
CORDATA, Spreng. Gen. ii. 581 (1831).
psoraloides, *DC. Prod.* ii. 216.—Ind. or.
senegalensis, *Guill. & Perr. Fl. Senegamb. Tent.* i. 171. t. 45.—Afr. trop.

CYAMUS, Sm. Exot. Bot. i. 59 (1804) = **Nelumbium**, Juss. (Nymph.).
flavicomus, Salisb. in Kon. & Sims, Ann. Bot. ii. (1805) 75 = Nelumbium luteum.
luteus, Nutt. Gen. Am. ii. 25 = Nelumbium luteum.
mysticus, Salisb. in Kon. & Sims, Ann. Bot. ii. (1805) 75 = Nelumbium luteum.
Nelumbo, Sm. Exot. Bot. i. (1806) 59. t. 31 = Nelumbium speciosum.
pentapetalus, Pursh, Fl. Am. Sept. ii. 398 = Nelumbium luteum.
reniformis, Pursh, l. c. = Nymphæa odorata.

CYANAEORCHIS, Rodrig. Orch. Nov. i. 112 (1877). *ORCHIDEAE*.
Arundinae, *Rodrig. l. c.* 113.—Bras.

CYANANTHUS, Griff. Notul. iv. 154 (1854) = **Staurantheras**, Benth. (Generac.).
umbrosus, Griff. l. c. = S. umbrosa.

CYANANTHUS, Miers, in Wall. Cat. n. 9007, 9008 (1849) = **Burmannia**, Linn.
pusillus, Miers, l. c. = B. pusilla.

CYANANTHUS, Wall. Cat. n. 1472, 1473 (1830). *CAMPANULACEAE*, Benth. & Hook. f. ii. 567.
barbatus, Edgew. in Trans. Linn. Soc. xx. (1851) 32 = integer.
barbatus, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 9.—China.
Hookeri, *C. B. Clarke, in Hook f. Fl. Brit. Ind.* iii. 485.—Reg. Himal.
incanus, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 20.—Reg. Himal.
inflatus, *Hook. f. & Thoms. l. c.* 21.—Reg. Himal.
integer, *Wall. Cat.* n. 1473.—Reg. Himal.
linifolius, Wall. ex Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 20 = microphyllus.
lobatus, *Wall. Cat.* n. 1478.—Reg. Himal.
microphyllus, *Edgew. in Trans. Linn. Soc.* xx. (1851) 31.—Reg. Himal.
pedunculatus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 484.—Reg. Himal.

CYANASTRUM, Cass. in Dict. Sc. Nat. xlv. 89 (1826) = **Volutarella**, Cass. (Compos.).

CYANEA, Gaudich. in Freyc. Voy. Bot. 457. t. 75 (1826). *CAMPANULACEAE*, Benth. & Hook. f. ii. 546.
KITTELIA, Reichb. Handb. 186 (1837).
MACROCHILUS, Presl, Prod. Lobel. 47 (1836).
arborescens, *H. Mann, in Proc. Am. Acad.* vii. (1867) 138.—Ins. Sandvic.
aspera, A. Gray, i. c. v. (1861) 148 = Rollandia calycina.
Grimesiana, *Gaudich. in Freyc. Voy. Bot.* 458.—Ins. Sandvic.
humille, *Wawra, in Flora*, lxv. (1873) 47.—Ins. Sandvic.
leptostegia, *A. Gray, in Proc. Am. Acad.* v. (1861) 149.—Ins. Sandvic.
lobata, *H. Mann, l. c.* vii. (1867) 138.—Ins. Sandvic.
pilosa, *A. Gray, l. c.* (1861) 149.—Ins. Sandvic.
Rollandia, *A. Gray, l. c.*—Ins. Sandvic.
superba, *A. Gray, l. c.*—Ins. Sandvic.
trichomantha, *A. Gray, l. c.*—Ins. Sandvic.

CYANELLA, Linn. Gen. ed. V. 149 (1754). *HAEMODORACEAE*, Benth. & Hook. f. iii. 679.
PHARETRELLA, Salisb. Gen. Pl. Fragm. 47 (1866).
TRIGELLA, Salisb. l. c. 46 (1866).
alba, Herb. ex Schult. f. Syst. vii. 1700 = Ornithogalum caerulea.
alba, *Linn. f. Suppl.* 201.—Afr. austr.
capensis, *Linn. Gen.* ed. V. 149; *Syst.* ed. X. 985.—Afr. austr.
coerulea, Eckl. Verz. 4 = capensis.
Illiv., Molina, Sagg. Chil. ed. II. 130 = Pasithea caerulea.
lineata, Burch, Trav. ii. 589 = lutea.
lutea, *Linn. f. Suppl.* 201.—Afr. austr.
odoratissima, Lindl. Bot. Reg. t. 1111 = lutea.
orchidiformis, *Jacq. Coll. iv.* 211.—Afr. austr.
pulchella, Salisb. Prod. 249 = capensis.
rosea, Eckl. ex Baker, in Journ. Linn. Soc. xvii. (1879) 498 = lutea.

CYANITIS, Reinw. in Blume, Cat. Gew. Buitenz. 68 (1823) = **Dichrus**, Lour. (Saxifrag.).
chinensis, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 76 = D. febrifuga.
cyboctis, Reinw. in Blume, Cat. Gew. Buitenz. 68; Bijdr. 921 = D. febrifuga.
versicolor, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 76 = D. febrifuga.

CYANOBOTRYS, Zucc. in Abh. Akad. Muench. iv. ii. (1845) 28 = **Muellera**, Linn. (Legumin.).
mexicana, Zucc. l. c. = M. mexicana.

CYANODAPHNE, Blume, Mus. Bot. Lugd. Bat. i. 333 (1851). *LAURINEAE*, Benth. & Hook. f. iii. 151.
cuneata, Blume, l. c. 334 = Dehaasia cuneata.
glabra, *Miq. Fl. Ind. Bat. Suppl.* 360.—Sumatra.
tomentosa, *Blume, Mus. Bot. Lugd. Bat.* i. 334.—Borneo.

CYANOPHYLLUM, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1852) 324 = **Miconia**, Ruiz & Pav. (Melast.).
magnificum, Denis, in Rev. Hortic. (1853) 166 = M. magnifica.
metallicum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852 325 = M. metallica.
vittatum, Linden, ex Kafarin, in Rev. Hortic. (1877) 38 = Clidemia vittata.

CYANOPIS, Blume, Fl. Jav. præf. p. vi (1828); et in DC. Prod. v. 69 (1836) = **Vernonia**, Schreb. (Compos.).
cyrtoloma, DC. in Wight, Contrib. 7 = V. cinerea.
hypoleuca, Sch. Bip. ex Walp. Rep. vi. 98.—Abyss.
Leopoldi, Sch. Bip. ex Walp. l. c. ii. 949 = V. Leopoldi.
madagascariensis, DC. Prod. v. 69 = V. madagascariensis.
moluccensis, DC. l. c. = V. moluccensis.
pubescens, DC. l. c. = V. chinensis.
villosa, DC. l. c. = V. chinensis.

CYANOPIS, Steud. Nom. ed. II. i. 74, in syn. (1840) = CYANOPSIS, Cass. = **Volutarella**, Cass. (Compos.).

CYANOPOGON, Welw. ex C. B. Clarke, in DC. Monog. Phan. iii. 240, 258 (1881) = **Cyanotis**, D. Don, (Commelin.).

CYANOPSIS, Cass. in Bull. Soc. Philom. (1817) 200 = **Volutarella**, Cass. (Compos.).
Chingoyo, Steud. Nom. ed. II. i. 413, in syn. = Pluchea Chingoyo.
decurrens, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 563 = Vernonia cinerea.
microcephala, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 120, nomen.—Malaya.
radicantissima, Cass. in Bull. Soc. Philom. (1817) 200 = Vol. muricata.
Zollingeriana, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 120, nomen = Vernonia cinerea.

CYANOPSIS, Endl. Each. 332 (1841) = CYANOPIS, Blume = **Vernonia**, Schreb. (Compos.).

CYANORCHIS, Thou. ex Stend. Nom. ed. II. i. 457 (1840) = **Pantus**, Lour. (Orchid.) (P. tetragona).

CYANOSERIS, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 43 = **Lactuca**, Tourn. (Compos.).
perennis, Schur; l. c. = L. perennis.

CYANOSPERMUM, Wight & Arn. Prod. 259 = **Rhyncosia**, Lour. (Legum.).
albiflorum, Wight & Arn. l. c. 260 = R. cyanosperma.
angolense, Welw. Apont. 586 = R. calycina ?
calycinum, Hook. f. in Hook. Niger Fl. 312 = R. calycina.
javanicum, Miq. Fl. Ind. Bat. i. 1. 167 = R. rufescens.
tomentosum, Wight & Arn. Prod. 260 = R. cyanosperma.

CYANOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 35. *VERBENACEAE*, Benth. & Hook. f. ii. 1140.
BUNNYA, F. Muell. Fragm. v. 36. t. 39 (1865).
angustifolia, *Turcz. l. c.* 36.—Austral.
Bunnyana, *F. Muell. Fragm.* v. 36.—Austral.
intermedia, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 36.—Austral.
lanceolata, *Turcz. l. c.* 35.—Austral.
Turczaninowii, *F. Muell. Fragm.* vi. 154.—Austral.

CYANOSTREMMA, Benth. ex Hook. & Arn. Bot. Beech. Voy. 415 (1841) = **Calopogonium**, Desv. (Legumin.).
caeruleum, Benth. l. c. = Calop. caeruleum.

CYANOTHAMNUS, Lindl. Swan River App. 18 (1839) = **Boronia**, Sm. (Rutac.).
anethifolius, Bartl. in Lehm. Pl. Preiss. i. 170 = B. ramosa.
ramosus, Lindl. Swan River App. 18 = B. ramosa.
tenuis, Lindl. l. c. = B. tenuis.
tridactylites, Bartl. in Lehm. Pl. Preiss. ii. 227 = B. anemonifolia.

CYANOTIS, D. Don, Prod. Fl. Nep. 45 (1825). *COMMELINACEAE*, Benth. & Hook. f. iii. 851.
BELOSYNAPSIS, Hassk. in Flora, liv. (1871) 259.
CYANOPOGON, Welw. ex C. B. Clarke, in DC. Monog. Phan. iii. 240, 258 (1881).
DALZELLIA, Hassk. in Flora, xlvii. (1866) 593.
ERYTHROTIS, Hook. f. Bot. Mag. t. 6150 (1875).
SIPHOSTOMA, Rafin. Fl. Tellur. ii. 16 (1836).
TOMINGIA, Neck. Elem. iii. 166 (1790).
ZYGOMENES, Salisb. in Trans. Hort. Soc. i. (1812) 271.
abyssinica, Hook. f. in Journ. Linn. Soc. vi. (1862) 21 = Mannia.
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 344 = hirsuta.
adscendens, Dalz. in Hook. Kew Journ. iv. (1852) 343 = tuberosa.
angusta, *C. B. Clarke, in DC. Monog. Phan.* iii. 260.—Afr. occ.
arachnoidea, *C. B. Clarke, l. c.* 250.—Ind. or.
axillaris, *D. Don, Prod. Fl. Nep.* 46.—Ind. or.
barbata, *D. Don, l. c.*—Reg. Himal.
Burmannia, Wight, Ic. t. 2089 = papillonacea.
caespitosa, Kotschy & Peyr. Pl. Tinn. 48 = longifolia.
capitata, *C. B. Clarke, in DC. Monog. Phan.* iii. 243.—Java.
coylanica, *Hassk. Commel. Ind.* 145.—Zeylan.
crescrescens, Hassk. l. c. 114 = tuberosa.
cristata, *D. Don, Prod. Fl. Nep.* 46.—Ind. or.; Malaya.
cucullata, *Kunth, Enum. Pl.* iv. 107.—Ind. or.
decumbens, Wight, Ic. vi. t. 2058 = fasciculata.
dichrotricha, Stocks, in Wight, l. c. 33. t. 2087 = fasciculata.
diurnaphora, Hassk. Commel. Ind. 105 = axillaris, fasciculata.
djurensis, *C. B. Clarke, in DC. Monog. Phan.* iii. 256.—Afr. trop.
erianthe, Hassk. Commel. Ind. 138 = fasciculata.
fasciculata, *Schult. f. Syst.* vii. 1152.—Ind. or.
fasciculata, Wall. Cat. n. 8990 = barbata, arachnoidea.
foecunda, *DC. ex Hassk. Commel. Ind.* 110.—Abyss.
guaniculata, *C. B. Clarke, in DC. Monog. Phan.* iii. 260.—Cochinch.
glaberrima, Hassk. Commel. Ind. 156 = barbata.
Gueinzii, Hassk. l. c. 123 = nodiflora.
hirsuta, Baker, in Trans. Linn. Soc. xxix. (1875) 163 = Commelina aspera.

CYANOTIS :—
hirsuta, *Planch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* viii. 57.—Abyss.
hirtella, Miq. ex C. B. Clarke, in DC. Monog. Phan. iii. 246 = papillonacea.
hispida, Dalz. in Hook. Kew Journ. iii. (1851) 139 = papillonacea.
Huegelii, Hassk. Commel. Ind. 138 = cristata.
imbricata, Kunth, Enum. Pl. iv. 108 = cristata.
incerta, *Hassk. Commel. Ind.* 160.—Asia.
Karliana, *Hassk. l. c.* 146.—Ind. or.
kewensis, *C. B. Clarke, in DC. Monog. Phan.* iii. 243.—Ind. or.
lanata, *Benth. in Hook. Niger Fl.* 542.—Afr. trop.
lanceolata, Wight, Ic. t. 2085 = villosa.
lanceolata, Thw. Enum. Pl. Zeyl. 293 = zeylanica.
Lamiana, Wight, Ic. vi. 33. t. 2086 = fasciculata.
longifolia, *Benth. in Hook. Niger Fl.* 542.—Afr. trop.
longifolia, Wight, Ic. vi. 33. t. 2084 = Wightii.
Mannii, *C. B. Clarke, in DC. Monog. Phan.* iii. 258.—Afr. trop.
nilagirica, Hassk. Commel. Ind. 127 = fasciculata.
nobilis, Hassk. l. c. 148 = barbata.
nodiflora, C. B. Clarke, in Journ. Linn. Soc. xi. (1871) 459 = barbata.
nodiflora, *Kunth, Enum. Pl.* iv. 106.—Afr. austr.
papillonacea, *Schult. f. Syst.* vii. 1151.—Ind. or.
parasitica, *Hochst. ex Schweinf. Beitr. Fl. Aethiop.* 214.—Abyss.
pauciflora, *A. Rich. Tent. Fl. Abyss.* ii. 345.—Abyss.
pilosa, *Schult. f. Syst.* vii. 1154.—Ind. or.
pilosa, Wight, Ic. t. 2088 = arachnoidea.
polyrrhiza, *Hochst. ex Hassk. Commel. Ind.* 150.—Abyss.
racemosa, C. B. Clarke, Commel. et Cyrt. Beng. t. 38 = cristata.
racemosa, Heyne, ex Hassk. Commel. Ind. 125 = villosa.
rosea, Wight, Ic. vi. 33. t. 2086 = fasciculata.
sarmentosa, Wight, l. c. t. 2087 = tuberosa.
sertosa, *Hassk. Commel. Ind.* 144.—As. trop.
Schweinfurthii, Hassk. l. c. 134 = lanata.
speciosa, Hassk. l. c. 108 = nodiflora.
Steckeri, Hassk. l. c. 112 = lanata.
Thomsonii, Hassk. l. c. 133 = tuberosa.
Thwaitesii, Hassk. l. c. 136 = fasciculata.
tuberosa, *Schult. f. Syst.* vii. 1150.—Ind. or.
uniflora, *Hassk. Commel. Ind.* 104.—Malaya.
vaga, *Schult. f. Syst.* vii. 1159.—Java.
vaginata, *Wight, Ic.* vi. 34. t. 2083.—Ind. or.
villosa, *Schult. f. Syst.* vii. 1155.—Ind. or.
vittata, Lindl. in Journ. Hort. Soc. v. (1850) 139 = Zebrina pendula.
vivipara, *Dals. in Hook. Kew Journ.* iii. (1851) 236.—Ind. or.
Wightii, *C. B. Clarke, in DC. Monog. Phan.* iii. 250.—Ind. or.
zebrina, Nees.—Cf. Linnaea, xxiv. (1851) 385 = Zebrina pendula.
zeylanica, C. B. Clarke, in DC. Monog. Phan. iii. 252 = cristata.

CYANOTIS, Miers, in Wall. Cat. n. 9006 (1849) = **Burmannia**, Linn.
nepalensis, Miers, l. c.—Ind. or.

CYANOTRIS, Rafin. in Am. Monthly Mag. (1811) 356; (1815) = Camassia, Lindl. (Liliac.) (1832).
angustifolia, Rafin. l. c. (1819) 192 = Cam. esculenta.
sulcicaules, Rafin. l. c. (1811) 356 = Cam. esculenta.

CYANTHILLIUM, Blume, Bijdr. 889 (1826) = **Vernonia**, Schreb. (Compos.).
malaccense, Blume, l. c. 890 = V. molucconsis.
pubescens, Blume, l. c. = V. chinensis.
villosum, Blume, l. c. 889 = V. chinensis.

CYANUS, [Tourn.] Linn. Syst. ed. I (1735); Juss. Gen. Pl. 174 (1789) = Centaurea, Linn. (Compos.).
arvensis, Moench, Meth. 561 = Cent. Cyanus.
atropurpurea, Besng. Enum. Stirp. Transs. iii. 77 = Cent. atropurpurea.
austriaca, Besng. l. c. 72 = Cent. austriaca.
axillaris, J. & C. Presl, Fl. Cech. 179 = Cent. variegata.
collinus, Guerin. Mey. & Scherb. Fl. Wett. iii. 173 = Cent. Jacea.
coriacea, Besng. Enum. Stirp. Transs. iii. 76 = Cent. Scabiosa.

CYANUS :—
cyanus, Hill, Hort. Kew. 64 = Centaurea Cyanus.
dentatofalcus, Gilib. Fl. Lituan. i. 191 = Cent. Cyanus.
erlophorus, Guerin. Fruct. ii. 385. t. 161 = Cent. eriophora.
fiasculosus, Besng. Enum. Stirp. Transs. iii. 79 = Cent. fiosculosa.
intermedius, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 96 = Cent. Intermedia.
lactatus, Moench, Meth. 561 = Cent. sempervirens.
Jacea, Guerin. Mey. & Scherb. Fl. Wett. iii. 173 = Cent. Jacea.
lacinistus, Gilib. Fl. Lituan. i. 192 = Cent. Scabiosa.
lanceolatus, Moench, Meth. 560 = Cruptna vulgaris.
lugdunensis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 96 = Cent. montana.
lutens, Moench, Meth. 560 = Cent. moschata.
melitensis, Guerin. Fruct. ii. 388 = Cent. melitensis.
mollis, J. & C. Presl, Fl. Cech. 179 = Cent. montana.
montanus, Hill, Hort. Kew. 64 = Cent. montana.
moschatus, Guerin. Fruct. ii. 382 = Cent. moschata.
mucronatus, Moench, Meth. 560 = Volutarella Lippii.
napifolius, Guerin. Fruct. ii. 383 = Cent. napifolia.
niger, Guerin. l. c. 383 = Cent. nigra.
nigrescens, J. & C. Presl, Fl. Cech. 179 = Cent. nigrescens.
orientalis, Besng. Enum. Stirp. Transs. iii. 78 = Cent orientalis.
paniculatus, Gilib. Fl. Lituan. i. 193 = Cent. maculosa.
paniculatus, Moench, Meth. 561 = Cent. paniculata.
phrygius, Guerin. Mey. & Scherb. Fl. Wett. iii. 173 = Cent. phrygia.
plumosus, Gilib. Fl. Lituan. i. 191 = Cent. phrygia.
pullatus, Guerin. Fruct. ii. 383 = Cent. pullata.
pumilus, Besng. Enum. Stirp. Transs. iii. 73 = Cent phrygia.
ragusinus, Guerin. Fruct. ii. 388 = Cent. ragusina.
rosarus, Besng. Enum. Stirp. Transs. iii. 75 = Cent. trinervia.
rotundatus, Moench, Meth. 561 = Cent. ragusina.
Scabiosa, Moench, l. c. = Cent. Scabiosa.
segetum, Hill, Veg. Syst. iv. 29 = Cent. Cyanus.
sibiricus, Guerin. Fruct. ii. 388 = Cent. sibirica.
solstitialis, J. & C. Presl, Fl. Cech. 179 = Cent. solstitialis.
variegatus, Besng. Enum. Stirp. Transs. iii. 74 = Cent variegata.
vulgaris, Delarb. Fl. Auv. ed. II. 208 = Cent. Cyanus.

CYATHANTHERA, Pohl, Pl. Bras. Ic. ii. 130. t. 188 (1831) = **Miconia**, Ruiz & Pav. (Melastom.).
lanceolata, Pohl, l. c. 131 = M. Cyathanthera.

CYATHELLA, Decne. in Ann. Sc. Nat. Sér. II. ii. (1838) 332. t. 12 = **Vincetoxicum**, Rupp. (Asclepiad.).
Bojeriana, Decne. l. c. 333.—Madag.
mucronata, Decne. l. c.—Madag.
repanda, Decne. l. c. t. 12.—Madag.

CYATHIDIUM, Lindl. ex Royle, Illustr. Bot. Himal 251. t. 56 (1839) = **Saussurea**, DC. (Compos.).
taraxacifolium, Lindl. l. c. = S. taraxacifolia.

CYATHOCALYX, Champ. ex Hook. f. & Thoms. Fl. Ind. i. 196 (1855). *ANONACEAE*, Benth. & Hook. f. i. 24.
SOALA, Blanco, Fl. Filip. ed. I. 487 (1837).
Maingayi, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 58.—Malacca.
marginalis, *Schegf. in Ann. Jard. Buitenz.* ii. (1885) 6.—Ins. Banca.
martabanica, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 58.—Burma.
obtusifolius, *Becc. & Schegf. in Ann. Jard. Buitenz.* ii. (1886) 6.—N. Guinea.
pubescens, *Schegf. l. c.* 7.—Sumatra.
sumatranus, *Schegf. in Tijdschr. Nederl. Ind.* xxxii. (1871) 586.—Sumatra.
zeylanicus, *Champ. ex Hook. f. & Thoms. Fl. Ind.* i. 197.—Zeylan.

CYATHOCHAETA, Nees, in Lehm. Pl. Preiss. ii. 86 (1846). *CYPERACEAE*, Benth. & Hook. f. iii. 1061.
TETRALEPIS, *Steud. Syn. Pl. Cyp.* 159 (1855).
avenacea, *Benth. Fl. Australt.* vii. 563.—Austral.
clandestina, *Benth. l. c.*—Austral.
flandica, *Nees, in Lehm. Pl. Preiss.* ii. 86.—Austral.

CYATHOCLINE, Cass. in Ann. Sc. Nat. Sér. I. xvii. (1829) 419. *COMPOSITAE*, Benth. & Hook. f. ii. 361.
flava, C. B. Clarke, Comp. Ind. '87 = lutea.
Lewii, Dalz. & Gibs. Bomb. Fl. 124 = lutea.
Lewii, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 159 = lyrata.
lutea, *Law, ex Wight, l. c.* 158.—Ind. or.
lyrata, *Cass. in Ann. Sc. Nat.* Sér. I. xvii. (1829) 420. —Ind. or.
stricta, DC. Prod. v. 374 = lyrata.

CYATHOCNEMIS, Klotzsch, in Abh. Akad. Berl. (1854) 100 = **Begonia**, Linn.
obliqua, Klotzsch, l. c. 101 = B. obliqua.

CYATHOCOMA, Nees, in Linnaea, ix. (1834) 800 = **Tetraria**, Beauv. (Cyperac.).
Ecklonii, Nees, l. c.; et x. (1836) 195.—Afr. austr.
Neesii, Boeck. in Linnaea, xxxviii. (1874) 313 = Macrochaetum Dregei.
nigrovaginata, Nees, in Linnaea, x. (1836) 196.—Afr. austr.

CYATHODES, Labill. Nov. Holl. Pl. i. 57. t. 81 (1806). *EPACRIDEAE*, Benth. & Hook. f. ii. 612.
ANDROSTOMA, Hook. f. Fl. Antarct. i. 44. t. 30 (1845).
abietina, *R. Br. Prod.* 540.—Austral.
acerosa, *R. Br. l. c.* 539.—Austral.
ascendens, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 268.—Tasman.
Banksii, Gaudich. in Freyc. Voy. Bot. 98 = Tameiameiae.
Baxteri, A. Cunn. in DC. Prod. vii. 741 = Astroloma divaricatum.
Colensoi, *Hook. f. Handb. New Zeal. Fl.* 177.—N. Zel.
dealbata, *R. Br. Prod.* 539.—Tasman.
distichea, Labill. Nov. Holl. Pl. i. 58. t. 82 = Trochocarpa disticha.
divaricata, *Hook. f. Fl. Tasm.* i. 246. t. 74 B.—Tasman.
Douglasii, A. Gray, in Proc. Am. Acad. v. (1861) 324 = imbricata.
empetrifolia, *Hook. f. Fl. N. Zel.* 164.—N. Zel.
glauca, *Labill. Nov. Holl. Pl.* i. 57. t. 81.—Austral.
imbricata, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 10.—Ins. Sandvic.
laurina, R. Br. ex Rudge, in Trans. Linn. Soc. viii. (1807) 298. t. 9 = Trochocarpa laurina.
Macraeana, DC. Prod. vii. 743 = Tameiameiae.
macrantha, Hook. f. Fl. Tasm. i. 245 = stramínea.
Oxycedrus, R. Br. Prod. 540 = acerosa.
parvifolia, *R. Br. l. c.*—Tasman.
Pomaresc, *A. Gray, in Proc. Am. Acad. v.* (1861) 324. —Ins. Sandvic.
pumila, *Hook. f. Handb. New Zeal. Fl.* 785.—N. Zel.
robusta, *Hook. f. l. c.* 177.—N. Zel.
rubicunda, F. Muell. Fragm. iv. 99 = Leucopogon rubicundus.
squamuligera, F. Muell. l. c. 97 = Stenanthera squamuligera.
stramínea, *R. Br. Prod.* 539.—Tasman.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 62.—Ins. Tahiti.
Tameiameiae, *Cham. in Linnaea*, i. (1826) 539.—Ins. Sandvic.

CYATHODISCUS, Hochst. in Flora, xxv. (1842) 240 = **Peddiea**, Harv. (Thymel.).
umbellatus, Hochst. l. c. = P. Dregei.

CYATHOGLOTTIS, Poepp. & Endl. Nov. Gen. et Sp. i. 55. t. 94 (1835) = **Bolbalis**, Ruiz & Pav. (Orchid.).
candida, Poepp. & Endl. l. c. 56 = S. candida.
crocea, Poepp. & Endl. l. c. 55 = S. crocea.
macrantha, Lem. Jard. Fleur. iv. (1854) t. 355 = S. macrophylla ?

CYATHOGYNE, Muell. Arg. in Flora, xlvii. (1864) 396. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 286.
viridis, *Muell. Arg. l. c.*—Afr. trop.

CYATHOPAPPUS, F. Muell. Fragm. ii. 157 (1861) = **Gnaphalia**, Cass. (Compos.).
gnaphaloides, F. Muell. l. c. 158 = G. cyathopappa.

CYATHOPAPPUS, Sch. Bip. in Pollichia, xviii.-xix. (1861) 132 = **Hyptocyaphus**, Sch. Bip. (Compos.).
calcitrapioides, Sch. Bip. l. c. = H. cyathiformis.

CYATHOPHORA, Rafin. Fl. Tellur. iv. 117 (1836) = **Euphorbia**, Linn.
ciliata, Rafin. l. c. = E. heterophylla.
heterophylla, Rafin. l. c. = E. heterophylla.
picta, Rafin. l. c. = E. heterophylla.

CYATHOPSIS, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 66; et in Ann. Sc. Nat. Sér. V. ii. (1864) 152. *EPACRIDEAE*, Benth. & Hook. f. ii. 615.
floribunda, *Brongn. & Gris, ll. cc.* 66, 153.—N. Caled.

CYATHORHACHIS, Nees, ex Steud. Syn. Pl. Gram. 408 (1855) = **Polytoca**, R. Br. (Gramin.).
Wallichiana, Nees, ex Steud. l. c. = P. Wallichiana.

CYATHOSELINUM, Benth. in Benth. & Hook. f. Gen. i. 912 (1867). *UMBELLIFERAE*, Benth. & Hook. f. i. c.
tomentosum, *Benth. & Hook. f. l. c.*—Dalmat.

CYATHOSTELMA, Fourn. in Mart. Fl. Bras. vi. iv. 319 (1888). *ASCLEPIADEAE*.
furcatum, *Fourn. l. c.* 320.—Bras.
latipes, *Fourn. l. c.* 319.—Bras.

CYATHOSTEMMA, Griff. Notul. iv. 707. t. 650 (1854). *ANONACEAE* [Benth. & Hook. f. i. 28].
viridiflora, *Griff. l. c.*—Malaya.

CYATHOSTEMON, Turcz. in Bull. Phys.-Math. Acad. Petersb. x. (1852) 331 = **Baeckea**, Linn. (Myrtac.).
tenuifolius, Turcz. l. c. 332 = B. Drummondii.

CYATHOSTYLES, Schott, ex Meisn. Gen. Comm. 184 (1840) = **Cyphomandra**, Sendtn. (Solanac.).

CYATHULA, Lour. Fl. Cochinch. i. 101 (1790). *AMARANTACEAE*, Benth. & Hook. f. iii. 31.
POLYCNEMUM, Wall. Cat. n. 6338 (1845).
achyranthoides, *Moq. in DC. Prod.* xiii. ii. 326.—Am. trop.
angustifolia, *Moq. l. c.* 328.—Afr. austr.
capitata, *Moq. l. c.* 329.—Reg. Himal.
capitata, Thw. Enum. Pl. Zeyl 249 = ceylanica.
ceylanica, *Hook. f. Fl. Brit. Ind.* iv. 728.—Zeylan.
cylindrica, *Moq. in DC. Prod.* xiii. ii. 328.—Afr. austr.
cylindrica, Schweinf. ex Benth. & Hook. f. Gen. iii. 36 = Pandiaka cylindrica.
geminata, *Moq. in DC. Prod.* xiii. ii. 330.—Afr. trop.
geniculata, *Lour. Fl. Cochinch.* 102.—Geront. trop.
globosa, Moq. in DC. Prod. xiii. ii. 330 = geniculata.
globulifera, *Moq. l. c.* 329.—Afr. austr.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 97.—Afr. austr.
orbiculata, Moq. l. c. 330 = Pupalia orbiculata.
prostrata, Blume, Bijdr. 549 = geniculata.
repens, Moq. in DC. Prod. xiii. ii. 330 = geniculata.
sanguinolenta, *Moq. l. c.* 330.—Ind. or.
Schimperiana, Moq. l. c. 328 = cylindrica.
segnis, Moq. l. c. 327 = tomentosa.
sphaerocephala, *Baker, in Journ. Linn. Soc.* xx. (1883) 238.—Madag.
tomentosa, *Moq. in DC. Prod.* xiii. ii. 327.—Reg. Himal.
triuncinata, *Moq. l. c.*—Madag.

CYBRANTHERA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 87 (1825) = **Limnophila**, R. Br. (Scrophul.).
connata, Buch.-Ham. l. c. = L. hypericifolia.

CYBELE, Falc. in Lindl. Veg. Kingd. 188 C (1847) = **Herminium**, Linn. (Orchid.).

CYBELE, Salisb. in Knight, Prot. 128 (1809) = **Stenocarpus**, R. Br. (Proteac.).
umbelliferа, Salisb. l. c. 124 = S. Forsteri.

CYBELION, Spreng. Syst. iii. 679, 731 (1826) = **Ionopsis**, H. B. & K. (Orchid.).
pallidiflorum, Spreng. l. c. 721 = I. utricularioides.
pulchellum, Spreng. l. c. = I. pulchella.
tenuerum, Steud. Nom. ed. II. i. 468 = I. utricularioides.
teres, Steud. l. c. = I. teres.
testiculatum, Spreng. Syst. iii. 721 = I. testiculata.
Utricularioides, Spreng. l. c. = I. utricularioides.

CYBIANTHUS, Mart. Nov. Gen. et Sp. iii. 87. t. 236 (1829). *MYRSINEAE*, Benth. & Hook. f. ii. 643.
PECKIA, Vell. Fl. Flum. 51; i. t. 184, 185 (1825).
WEIGELTIA, A. DC. in. Trans. Linn. Soc. xvii. (1836) 102.
angustifolius, *A. DC. in Ann. Sc. Nat. Sér.* II. xvi. (1841) 92.—Bras.
Boissieri, *A. DC. l. c.* 93.—Bras.
coriaceus, *Mart. Herb. Fl. Bras.* 359.—Bras.
costaricanus, *Hemsl. Biol. Centr. Am. Bot.* ii. 290.—Costa Rica.
crotonoides, *Schomb. Faun. & Fl. Gui.* 1086, nomen.—Guian.
cuneifolius, *Mart. Nov. Gen. et Sp.* iii. 88.—Bras.
cuspidatus, *Miq. in Mart. Fl. Bras.* x. 297.—Ind. occ.
densiflorus, *Mart. Herb. Fl. Bras.* 360.—Bras.
densiflorus, *Miq. in Mart. Fl. Bras.* x. 300.—Bras.
detergens, *Mart. Herb. Fl. Bras.* 360.—Bras.
fuscus, *Mart. l. c.* 359.—Bras.
glaber, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 94. —Bras.
guyanensis, *Miq. in Mart. Fl. Bras.* x. 298.—Guiana.
Humboldtii, A. DC. in Trans. Linn. Soc. xvii. (1837) 104 = Ardisia acuminata.
Lessertii, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 93.—Bras.
longifolius, *Miq. in Mart. Fl. Bras.* x. 299.—Bras.
luridus, Benth. ex Miq. l. c. 292 = penduliflorus.
macrophyllus, *Miq. l. c.*—Bras.
meridianus, *Miq. l. c.* 297.—Venezuela.
microbotrys, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 94.—Guiana.
multicostatus, *Miq. in Mart. Fl. Bras.* x. 299.—Bras.
multipunctatus, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 94.—Guian.
myranthos, *Miq. in Mart. Fl. Bras.* x. 300.—Guian.
myrtifolius, *Griseb. in Gertt. Abh.* xxiv. (1879) 222.—Reg. Argent.
nitidus, *Miq. in Mart. Fl. Bras.* x. 295.—Bras.
obovatus, *Mart. ex Miq. in Mart. l. c.* 301.—Bras.
parviflorus, C. Muell. in Walp. Ann. v. (1858) 472 — seq.
parvifolius, *Schlecht. in Linnaea*, xxvi. (1853) 672.—N. Granat.
pseudoflorus, *Mart. Nov. Gen. et Sp.* iii. 87.—Bras.
philippinensis, *Hook. f. ex Vidal, Sinops.* p. xxx. t. 61. f. h.—Ins. Philipp.
Priceurii, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 93.—Guian.
Principis, A. DC. in Trans. Linn. Soc. xvii. (1837) 104 = Ardisia acuminata.
subspicatus, *Benth. ex Miq. in Mart. Fl. Bras.* x. 296. —Bras.

CYBIOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 155 = **Ayenia**, Linn. (Stercul.).
abutilifolium, Turcz. l. c. 156 = A. magna.
sidaefolium, Turcz. l. c. = A. mexicana.

CYBISTAX, Mart. in DC. Prod. ix. 198 (1845). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1042.
YANGUA, Spruce, in journ. Linn. Soc. iii. (1859) 197.
antisyphilitica, *Mart. l. c.* 199.—Bras.
macrocarpa, Benth. in Benth. & Hook. f. Gen. ii. 1043 = Godmania macrocarpa.

CYCAS, Linn. Hort. Cliff. 483 (1737). *CYCADACEAE*, Benth. & Hook. f. iii. 444.
TODDA-PANA, Adans. Fam. ii. 25, 611 (1763).
angulata, *R. Br. Prod.* 345.—Austral.
Armstrongii, *Miq. in Arch. Néerl.* iii. (1868) 235.—Austral.
Beddomei, *Dyer, in Trans. Linn. Soc.* Ser. II. ii. (1881) 85.—Ind. or.
caffra, Thunb. in Nov. Act. Soc. Sc. Upsal. ii. (1775) 264 = Encephalartos caffer.
Calmeisiana, *F. Muell. Fragm.* x. 68.—Austral.
celebica, *Miq. Comm. Phyt.* 126.—Ins. Celebes.
circinnalis, *Linn. Sp. Pl.* 1188.—Ins. Moluce.
circinnalis, Roxb. Hort. Beng. 71; Fl. Ind. iii. 744 = Rumphii.
dilatata, Griff. Notul. iv. 1, 15 = pectinata.
glauca, Miq. Comm. Phyt. 127 = Rumphii.
gracilis, *Miq. in Verol. Kon. Akad. Wet.* xv. (1863) 366.—Austral.
hypoleuca, *Presl, Epim. Bot.* 288.—Ins. Philipp.
inermis, *Lour. Fl. Cochinch.* ii. 682.—China.
inermis, Oudem. in Arch. Néerl. ii. (1867) 394 = revoluta.

CYCAS:—
Yenbinianus, Griff. Notul. iv. 9 = pectinata.
Kenneyanus, *F. Muell. in Melb. Chemist* (March 1882).—Austral.
macrocarpa, Griff. Notul. iv. 11 = Rumphii.
madagascariensis, Miq. Comm. Phyt. 127 = circinalis.
media, *R. Br. Prod.* 346.—Austral.
neo-caledonica, *Linden, in Illustr. Hortic.* xxviii. (1881) 82, nomen.—N. Caled.
Normanbyana, *F. Muell. Fragm.* viii. 169.—Austral.
pandanifolia, *Hort.—Cf. Gard. Chron.* (1865) 365.—Hab. ?
papuana, *F. Muell. Papuan Pl. iv.* 76.—N. Guin.
pectinata, *Griff. Notul.* iv. 10.—Reg. Himal. ; Burma.
pluma, *Hort.—Cf. Gard. Chron.* (1878) i. 11.—Hab. ?
revoluta, Bedd. For. Fl. 227 = Beddomei.
revoluta, *Thunb. Fl. Jap.* 233.—Japon.
Riedlei, Gaudich. in Freyc. Voy. Bot. 484 = Macrozamia Fraseri.
Rumphiana, *Regel, Gartenfl.* (1863) 16.—Ins. Philipp. Rumphii, *Miq. in Bull. Sc. Phys. Nat. Néerl.* (1839) 45.—Malaya ; Austral.
Scratchleyana, F. Muell. Papuan Pl. ii. 15 = Rumphii.
Seemanni, *A. Br. in Sitzb. Ges. Naturf. Fr.* (1876) 114.—Ins. Fiji.
siamensis, *Miq. in Bot. Zeit.* xxi. (1863) 334.—Burma ; Siam.
speciosa, D. Don, in Proc. Linn. Soc. i. (1840) 64, 56 = Rumphii.
sphærica, Roxb. Hort. Beng. 71 ; Fl. Ind. iii. 747 = circinalis.
squamosa, Lodd. Cat. ex Loud. Hort. Brit. 408 = circinalis.
squarrosa, Stend. Nom. ed. II. i. 468, sphalm. = præc.
Thouarsii, R. Br. Prod. 347 = circinalis.
undulata, *Desf. ex Gaudich. in Freyc. Voy. Bot.* 484.—Hab. ?
villosa, Herb. Roy. ex A. DC. Prod. xvi. ii. 532 = Encephalartos caffer.
Wallichii, Miq. Mon. Cycad. 32 = Rumphii.

CYCCA, Batsch, Tab. 166 (1802) = CICCA, Linn. = **Phyllanthus**, Linn. (Euphorb.).

CYCLACHAENA, Fresen. Ind. Sem. Hort. Frankf. (1836) 4 ; et in *Linnæa*, xii. (1838) Litt. 78 = **Iva**, Linn. (Compos.).
ambrosiæfolia, Benth. & Hook. *f.* Gen. ii. 358 = I. ambrosiæfolia.
xanthifolia, Fresen. Ind. Sem. Hort. Frankf. (1836) 4 ; ex DC. Prod. vii. 288 = I. xanthifolia.

CYCLADENIA, Benth. Pl. Hartw. 322 (1850). *APOCYNACEÆ*, Benth. & Hook. f. ii. 728.
humilis, *Benth. l. c.* 823.—Calif.
tomentosa, *A. Gray, in Bot. Calif.* i. 474.—Calif.

CYCLAMEN, [Tourn.] Linn. Syst. ed. I (1735) ; Gen. ed. I. 41 (1737). *PRIMULACEÆ*, Benth. & Hook. f. ii. 634.
Cyclaminus, Hall. Stirp. Helv. i. 281 (1768).
æstivum, Reichb. Fl. Germ. Excurs. 407 = europæum.
africanum, *Boiss. & Reut. Pugill. Pl. Nov.* 75.—Afr. bor.
alpestre, Fisch. ex Hoffmgg. Verz. Pfl. 54 = latifolium.
antiochium, *Decne. in Rev. Hortic. Sér. IV.* iv. (1855) 93.—Hab. ?
Atkinsii × T. Moore, Garden Comp. i. 89 ; ex Lem. Jard. Fleur. iii. (1853) t. 297.
autumnale, J. Boos, Schoenbr. Fl. 45 = neapolitanum.
balearicum, Willk. in Oestr. Bot. Zeitschr. xxv. (1875) 111 = repandum.
caucasicum, Willd. ex Boiss. Fl. Orient. iv. 11 = comm.
cilicium, *Boiss. & Heldr. Diagn.* Ser. I. xi. 78.—Cilicia.
Clusii, Lindl. Bot. Reg. t. 1018 = europæum.
cordifolium, Stokes, Bot. Mat. Med. i. 295 = europæum.
coum, *Mill. Gard. Dict.* ed. VIII. n. 8.—Graecia ; Oriens.
creum, Reichb. Fl. Germ. Excurs. 406 = europæum.
cyprium, Ungr & Kotschy, Ins. Cypern, 295 = neapolitanum.
deltoideum, *Tausch, in Flora*, xii. (1829) 667.—Hab. ?
elegans, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 145 = coum.
europæum, *Linn. Sp. Pl.* 145.—Europ. ; Reg. Caucas.
europæum, Pall. ex Bieb. Fl. Taur. Cauc. 140 = coum.
europæum, Savi, Fl. Pis. i. 213 = hederæfolium.
europæum, Sm. Engl. Bot. t. 548 = neapolitanum.
ficariæfolium, Reichb. Fl. Germ. Excurs. 407 = neapolitanum.
floridum, Salisb. Prod. 119 = europæum.
graecum, *Link, in Linnæa*, ix. (1834) 578.—Graecia.
hastatum, *Tausch, in Flora*, xii. (1829) 668.—Hab. ?
hederæfolium, Sieber, ex Steud. Nom. ed. II. i. 458 = latifolium.
hederæfolium, *Ait. Hort. Kew.* ed. I. i. 196.—Europ. austr.
hederæfolium, Sibth. & Sm. Fl. Graec. Prod. i. 128 = latifolium.
hederæfolium, Sims, Bot. Mag. t. 1001 = vernale.
hederæfolium, Willd. Sp. Pl. i. 810 = neapolitanum ?
hyemale, Salisb. Prod. 118 = coum.
ibericum, *Goldie, ex G. Don, in Sweet, Hort. Brit.* ed. III. 560, nomen.—Iberia.
ibericum, Stev. ex Boiss. Fl. Orient. iv. 11 = comm.
immaculatum, [Pieri ?] in Ionios Asthol. v. 192 = hederæfolium.
indicum, *Linn. Sp. Pl.* 145 (cult.?).—Zeylan.
intermedium, *Wender. Ind. Sem. Hort. Marb.* (1825).—Europ. austr.
latifolium, *Sibth. & Sm. Fl. Graec.* ii. 71. t. 185.—Graecia ; As. Min.
linearifolium, *DC. Fl. Fr.* iii. 253.—Gallia.
littorale, *Sadler, ex Reichb. Fl. Germ. Excurs.* 406 ; *Lindl. Bot. Reg.* (1846) t. 56.—Reg. Mediterr.
macrophyllum, *Sieber, in Iris* (1832) 499.—Ins. Helena.
macropus, *Zucc. Del. Sem. Hort. Monac.* (1846) 4.—Oriens.
neapolitanum, *Tenore, Prod. Fl. Nap.* p. lxvi.—Europ.
officinale, Wender. ex Steud. Nom. ed. II. i. 458 = europæum.
orbiculatum, Mill. Gard. Dict. ed. VIII. n. 5 = europæum.
persicum, Mill. l. c. = latifolium.
persicum, Sibth. & Sm. Fl. Graec. Prod. i. 128 = graecum.
Poli, Chiaje, Opusc. 166 = graecum.
purpurascens, Mill. Gard. Dict. ed. VIII. n. 2 = europæum.
pyrolæfolium, Salisb. Prod. 119 = latifolium.
repandum, Sibth. & Sm. Fl. Graec. Prod. i. 128 ; Fl. Graec. t. 186 = hederæfolium.
retroflexum, Moench, Meth. Suppl. 177 = europæum.
romanum, Griseb. Spicil. *Fl.* Rumel. i. 5 = repandum.
subhastatum, Reichb. Fl. Germ. Excurs. 407 = neapolitanum.
utopicum, Hoffmgg. Verz. Pfl. 54 = latifolium.
vernale, C. Koch, in Linnæa, xvii. (1843) 308 = castile.
vernale, *Mill. Gard. Dict.* ed. VIII. n. 4.—Italia.
verum, Reichb. Fl. Germ. Excurs. 407 = hederæfolium.
verum, Sweet, Brit. Flow. Gard. Ser, I. i. t. 9 = coum.

CYCLAMINUS, Hall. Stirp. Helv. i. 281 (1768) = **Cyclamen**, Tourn. (Primul.).
europeus, Scop. Fl. Carn. ed. II. i. 135 = Cyclamen europæum.

CYCLANTHERA, Schrad. Ind. Sem. Hort. Gotting. (1831).—*Cf.* Linnæa, viii. (1833) Litt. 28. *CUCURBITACEÆ*, Benth. & Hook. f. i. 826.
Discanthera, Torr. & Gray, Fl. N. Am. i. 696 (1840).
Achatochilos, Spruce, ex Cogn. in Mém. Cour. Acad. Belg. 8vo. xxviii. (1876) 78 = brachybotrys.
australis, *Cogn. in DC. Monog. Phan.* iii. 857.—Paraguay.
biglandulifera, *Cogn. l. c.* 849.—Mexic.
Bourgeauna, *Naud. in Ann. Sc. Nat.* Sér. V. vi. (1866) 17.—Mexic.
brachybotrys, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1876) 78.—Peruv.
brachystachya, *Cogn. l. c.* 64.—Mexic.
bracteiflorus, *Cogn. l. c.* 74.—Bras.
Burchellii, *Cogn. l. c.* 67.—Bras.
citraceana, *Cogn. in DC. Monog. Phan.* iii. 865.—Venezuela.

CYCLANTHERA:—
cornuta, Hort. Rheed, ex Regel, in Act. Hort. Petrop. iii. (1875) 385 = Calochortus glaucus.
cordifolia, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 76.—N. Granat.
costaricensis, *Cogn. l. c.* 78.—Costa Rica.
digitata, *Ann. in Hook. Journ. Bot.* iii. (1841) 280.—Peruv.
dissecta, *Arn. l. c.*—Am. bor.
dissecta, A. Gray, Pl. Wright. i. 75 ; ii. 61 = Nandiniana.
Eichleri, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 74.—Bras.
elegans, *Cogn. l. c.* 70.—Bras.
eremocarpa, *Cogn. l. c.* 63.—Mexic.
explodens, *Naud. in Ann. Sc. Nat.* Sér. IV. xii. (1859) 160.—N. Granat.
filifera, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 72.—Guatemala.
glazea, *Cogn. l. c.*—Venezuela.
gracillima, *Cogn. l. c.* 71.—Mexic.
Griesebachii, *Cogn. in DC. Monog. Phan.* iii. 864.—Venezuela.
hastata, Cogn. in Mém. Cour. Acad. Belg. 8vo. xxviii. (1878) 64 = tamnoides.
Hystrix, *Arn. in Hook. Journ. Bot.* iii. (1841) 280.—Reg. Argent.
integrifolia, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 65.—Mexic.
Langsd, *Cogn. l. c.* 67.—Mexic.
leptostachya, *Benth. Bot. Voy. Sulph.* 99.—Am. austr.
macropoda, *Cogn. in DC. Monog. Phan.* iii. 858.—Peruv.
Mathewsii, *Arn. in Hook. Journ. Bot.* iii. (1841) 280.—Peruv.
microcarpa, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 75.—Bolivia.
multifoliolata, *Cogn. l. c.* 66.—Mexic.
Nandiniana, *Cogn. in DC. Monog. Phan.* iii. 851.—Texas.
Oerstedii, *Cogn. l. c.* 856.—Costa Rica.
pedata, *Schrad. Ind. Sem. Hort. Gotting.* (1831).—*Cf. Linnæa*, viii. (1833) *Litt.* 28.—Mexic.
quinqueloba, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 64.—Bras.
ribifolia, *Cogn. l. c.* 63.—Mexic.
subincrmis, *Cogn. in Bull. Acad. Belg.* Sér. II. xlv. (1880) 197.—N. Granat.
tamnifolia, *Griseb. in Goett. Abh.* xix. (1847) 147.—Uruguay.
tamnoides, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1878) 64.—Mexic.
tenuifolia, *Cogn. l. c.* 68.—Bras.
tenuisepala, *Cogn. l. c.* 64.—Peruv.
tomentosa, *Cogn. l. c.* 77.—Bolivia.
Triansei, *Cogn. l. c.* 75.—Mexic.

CYCLANTHUS, Poit. in Mém. Mus. Par. ix. (1822) 35. tt. 2, 3. *CYCLANTHACEÆ*, Benth. & Hook. f. iii. 958.
Cyclosanthes, Poepp. in Froriep, Not. xxxi. 312 (1831).
Discanthus, Spruce, in Journ. Linn. Soc. iii. (1859) 196.
bipartitus, *Poit. in Mém. Mus. Par.* ix. (1822) 36. t. 2.—Guiana.
cristatus, *Klotzsch, in 'Linnæa*, xx. (1847) 468.—N. Granat.
discolor, *Hort.—Cf. Gard. Chron.* (1888) i. 510.—Hab. ?
Plumierii, *Poit. in Mém. Mus. Par.* ix. (1822) 37. t. 3.—Ins. Martin.

CYCLAS, Schreb. Gen. 262 (1789) = **Crudia**, Schreb. (Legumin.).
aromatica, J. F. Gmel. Syst. 674 = C. aromatica.
spicata, J. F. Gmel. l. c. = C. Apalatoa.

CYCLEA, Arn. ex Wight, Illustr. i. 23 (1840). *MENISPERMACEÆ*, Benth. & Hook. f. i. 98.
Prepamora, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 30.
Rhaptonema, Miers, l. c. Ser. II. vii. (1851) 41.
Arnottii, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 19 = peltata.
barbata, Miers, l. c. = peltata.
Burmanii, *Miers*, l. c. Ser. II. vii. (1851) 40.—Ind. or. *Burmanni*, Arn. ex Wight, Illustr. i. 23 = peltata.

CYCLEA:—
debiliflora, Miers, Contrib. iii. 243 = peltata.
deltoidea, *Miers, in Hook. Kew Journ.* iii. (1851) 258. —China.
discolor, Miers, in Ann. Nat. Hist. Ser. II. vii. (1851) 41 = peltata.
laniflora, Miers, Contrib. iii. 241 = peltata.
madagascariensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 460.—Madag.
peltata, *Hook. f. & Thoms. Fl. Ind.* i. 201.—Ind. or.; Malaya.
pendulina, *Miers, Contrib.* iii. 248.—Malacca.
peregrina, *Miers, l. c.* 243.—Borneo.
populifolia, Hook. f. & Thoms. Fl. Ind. i. 202 = Lophosyphlum bicristatum.
robusta, *Becc. Malesia.* i. (1877) 157.—Borneo.
versicolor, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 19 = peltata.
villosum, Miers, in Ann. Nat. Hist. Ser. II. vii. (1851) 41 = Pericampylus inconnus.

CYCLIUM, Stend. Nom. ed. II. i. 458 (1840) = Cyclonium, E. Mey. (Scrophul.).

CYCLOBALANOPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 77 = Quercus, Tourn. (Cupul.).
acuta, Oerst. l. c. 78 = Q. acuta.
annulata, Oerst. l. c. = Q. glauca.
argentata, Oerst. l. c. 79 = Q. argentata.
Burgerii, Oerst. l. c. 78 = Q. Buergerii.
Championi, Oerst. l. c. 78 = Q. Championi.
gemelliflora, Oerst. l. c. = Q. gemelliflora.
gilva, Oerst. l. c. 78 = Q. gilva.
glauca, Oerst. l. c. = Q. glauca.
Helferiana, Oerst. l. c. 79 = Q. Helferiana.
Horsfieldii, Oerst. l. c. 78 = Q. semiserrata.
Kamroophii, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. lineata.
laevigata, Oerst. l. c. = Q. laevigata.
lamellosa, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 79 = Q. lamellosa.
lineata, Oerst. l. c. 78 = Q. lineata.
marginata, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. marginata.
Morkusii, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 79 = Q. turbinata.
mespilifolia, Oerst. l. c. = Q. mespilifolia.
muricata, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. sundaica.
myrsinaefolia, Oerst. l. c. = Q. myrsinaefolia.
oidocarpa, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 78 = Q. oidocarpa.
oxydon, Oerst. l. c. 79 = Q. lineata.
paucilamellosa, Oerst. l. c. = Q. lamellosa.
salicina, Oerst. l. c. 78 = Q. salicina.
semiserrata, Oerst. l. c. = Q. semiserrata.
Thomsoniana, Oerst. l. c. 79 = Q. lineata.
velutina, Oerst. l. c. = Q. velutina.

CYCLOBALANUS, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 80 = Quercus, Tourn. (Cupul.).
Beaurittii, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 376 = Q. Beaurittii.
Blancoi, Oerst. l. c. = Q. Blancoi.
Blumeana, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 81 = Q. Blumeana.
celebica, Oerst. l. c. = Q. celebica.
concentrica, Oerst. l. c. = Q. concentrica.
conocarpa, Oerst l. c. = Q. conocarpa.
costata, Oerst. l. c. = Q. costata.
cyrtopoda, Oerst. l. c. 80 = Q. cyrtopoda.
daphnoidea, Oerst. l. c. 81 = Q. daphnoidea.
Diepenhorstii, Oerst. l. c. 80 = Q. Diepenhorstii.
endelotacarpa, Oerst. l. c. 81 = Q. endelotacarpa.
Ewyckii, Oerst. l. c. 80 = Q. Ewyckii.
gracilis, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 376 = Q. gracilis.
Hancei, Oerst. l. c. 375 = Q. Hancei.
Hystrix, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 81 = Castanopsis Hystrix.
indusia, Oerst. l. c. 80 = Q. indusia.
javensis, Oerst. l. c. 81 = Q. javensis.
Lamponga, Oerst. l. c. 81 = Q. Lamponga.
leptogyna, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 378 = Q. leptogyna.
Lionoti, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 81 = Q. Lionoti.
nitida, Oerst. l. c. = Q. nitida.

CYCLOBALANUS:—
Omalokae, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 80 = Quercus Omalokae.
sogyna, Oerst. l. c. 81 = Q. sogyna.
ovalis, Oerst. l. c. = Q. ovalis.
philippensis, Oerst. l. c. = Q. philippensis.
platycarpa, Oerst. l. c. 80 = Q. platycarpus.
Rassa, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 376 = Q. Rassa.
Reinwardtii, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 81 = Q. Reinwardtii.
Tyrmanni, Oerst. l. c. 80 = Q. Korthalsii.

CYCLOBOTHRA, [D. Don, in] Sweet, Brit. Flow. Gard. Ser. I. iii. t. 278 (1828) = Calochortus, Pursh (Liliac.).
alba, Benth. in Trans. Hort. Soc. N. S. i. (1835) 418. t. 14. = Cal. albus.
barbata, [D. Don, in] Sweet, Brit. Flow. Gard. Ser. I. iii. t. 278 = Cal. flavus.
caerulea, Kellogg, in Proc. Calif. Acad. ii. (1863) 4 = Cal. caeruleus.
elegans, Benth. in Bot. Reg. sub t. 1662 = Cal. elegans
elegans, Torr. in Pacif. Rail. Rep. iv. 146 = Cal. Maweanus.
flava, Lindl. Bot. Reg. sub t. 1662 = Cal. flavus.
fusca, Lindl. l. c. = Cal. fuscus.
grandiflora, Mart. & Gal. in Bull. Acad. Brux. ix. 11. (1842) 384 = Cal. Bonplandianus.
Hartwegi, Kunth, Enum. Pl. iv. 231 = Cal. Hartwegii.
lutea, Lindl. Bot. Reg. t. 1663 = Cal. flavus.
monophylla, Lindl. in Journ. Hort. Soc. iv. (1849) 81 = Cal. Benthami.
nitida, Kunth, Enum. Pl. iv. 230 = Cal. nitidus.
nitida, Torr. in Pacif. Rail. Rep. iv. 146 = Cal. Benthami.
pallida, Lindl. Bot. Reg. sub t. 1662 = Cal. flavus.
paniculata, Lindl. l. c. = Cal. albus.
propinqua, Schau. in Linnaea, xix. (1847) 701 = Cal. Bonplandianus.
pulchella, Benth. in Trans. Hort. Soc. N. S. i. (1835) 412 = Cal. pulchellus.
purpurea, Sweet, Brit. Flow. Gard. Ser. II. iv. t. 20 = Cal. Bonplandianus.
uniflora, Kunth, Enum. Pl. iv. 229 = Cal. uniflorus.

CYCLOCAMPE, Steud. Syn. Pl. Cyp. 156 (1855) = Schoenus, Linn. (Cyperac.).
weigeotensis, Steud. l. c. = S. calystachyus.

CYCLOCAMPE, Benth. & Hook. f. Gen. iii. 1068 (1883). *CYPERACEAE.*
arundinacea, Benth. & Hook. f. l. c. = Schoenus arundinaceus.
elongata, Benth. & Hook. f. l. c. = Asterochaeta elongata.
nitens, *Hook. f. Ic. Pl.* sub t. 1484.—Ins. Maurit.

CYCLOCARPA, Miq. Fl. Ind. Bat. iii. 339 (1855) sphalm. = CYCLOCAMPE, Steud. = Schoenus, Linn. (Cyperac.).

CYCLOCARPUS, Jungh. in Hoev. & De Vriese Tijdschr. vii. (1840) 310 = Evodia, Forst. (Rutac.).
japonicus, Jungh. l. c. 311 = E. rutaecarpa?

CYCLOCODON, Griff. Notul. iv. 277 (1854) = Campanumoea, Blume (Campanul.).
asiaticum, Griff. ex Benth. & Hook. f. Gen. ii. 558 = Camp. celebica.
distans, Griff. Notul. iv. 277 = Camp. parviflora.
lancifolium, Kurz, in Flora, liii. (1870) 308 = Camp. celebica.
lancifolium, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 210, partim = Camp. parviflora.
leucocarpum, Miq. Fl. Ind. Bat. ii. 1062 = Camp. celebica.
parviflorum, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 18 = Camp. parviflora.
truncatum, Hook. f. & Thoms. l. c. = Camp. celebica.

CYCLODISCUS, Klotzsch, in Monatsb. Acad. Berl. (1850) 521 = Braganti, Lour. (Aristoloch.).
latifolius, Klotzsch, l. c. = B. tomentosa.
tomentosus, Klotzsch, l. c. = B. tomentosa.

CYCLOGYNE, Benth. in Lindl. Swan River App. 16 (1839) = Swainsona, Salisb. (Legumin.).

CYCLOGYNE:—
canescens, Benth. in Lindl. Swan River App. 16 = S. canescens.
procumbens, F. Muell. in Linnaea, xxv. (1852) 393 = Swainsona procumbens.
swainsonioides, Benth. in Mitch. Journ. Trop. Austral. 397 = S. procumbens.

CYCLOLEPIS, Gill. ex D. Don, in Phil. Mag. xi. (1832) 392. *COMPOSITAE*, Benth. & Hook. f. ii. 491.
denticulata, *Remy, in C. Gay, Fl. Chil.* iii. 292.—Chili. genistoides, *Gill. ex D. Don, in Phil. Mag.* xi. (1832) 392.—Chili.

CYCLOLEPIS, Meq. in Ann. Sc. Nat. Sér. II. i. (1834) 203. t. 8 = Cycloloma, Moq. (Chenopod.).
platyphylla, Moq. l. c. = Cycloloma platyphyllum.

CYCLOLEPSIS, Endl. Gen. Suppl. i. 1385 (1841) = Cyololepis, Gill. (Compos.).

CYCLOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 92. *LEGUMINOSAE*, Benth. & Hook. f. i. 545
Blanchetianum, *Tul. in Arch. Mus. Par.* iv. (1844) 84. —Bras.
brasilicense, *Benth. in Ann. Wien. Mus.* ii. (1838) 92. —Bras.
Clausseni, *Benth. in Hook. Journ. Bot.* ii. (1840) 64 — Bras
Hostmanni, *Brnth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 52.—Guian.

CYCLOLOMA, Moq. Chenop. Enum. 17 (1840). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 80.
Amoreua, Moq. in Delile, Cat. Hort. Monsp. (1844).
Amoreuxia, Moq. in Mém. Soc. Hist. Nat. Monsp. (1826).
Cyclolepis, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 203. t. 9.
Petermannia, Reichb. Nom. 236 (1841).
platyphyllum, *Moq. Chenop. Enum.* 18.—Am. bor.

CYCLOMORIUM, Walp. Rep. ii. 890 (1843) = Desmodium, Desv. (Legumin.).
caracasanum, Walp. l. c. = D. spirale.

CYCLONEMA, Hochst. in Flora, xxv. (1842) 225 = Clerodendron, Linn. (Verben.).
discolor, Klotzsch, in Peters, Reise Mossamb. Bot 260 = Cler. discolor.
hirsutum, Hochst. in Flora, xxv. (1842) 228 = Cler. umbellatum ?
mucronatum, Klotzsch, in Peters, Reise Mossamb. Bot. 262.—Afr. trop
myricoides, Hochst. in Flora, xxv. (1842) 226 = Cler. myricoides.
serratum, Hochst. l. c. 227.—Afr. austr.
spinescens, Oliver, in Journ. Linn. Soc. xv. (1876) 96. —Afr. trop.
spinescens, Klotzsch, in Peters, Reise Mossamb. Bot. 262.—Afr. trop.
sylvaticum, Hochst. in Flora, xxv. (1842) 227.—Afr. austr.
tettense, Klotzsch, in Peters, Reise Mossamb. Bot. 261. —Afr. trop.
triphyllum, Harv. Thes. Cap. i. 17.—Afr. austr.

CYCLOPHYLLUM, Hook. f. in Benth & Hook. f. Gen. ii. pars prima Addend. 585 (1873). *RUBIACEAE*, Benth. & Hook. f. Gen. ii. l. c. et 1229.
Deplanchei, *Hook. f. l. c.* 586, 1229.—N. Caled.

CYCLOPIA, Vent. Dec. Gen. 8 (1808). *LEGUMINOSAE*, Benth. & Hook. f. i. 466.
Ibbetsonia, Sims, in Bot. Mag. t. 1250 (1810).
Boweana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 9.—Afr. austr.
brachypoda, Benth. in Ann. Wien. Mus. ii. (1838) 67 = subternata.
cordifolia, Benth. l. c. = latifolia.
elata, *Smeet, Hort. Brit.* ed. I. 107, nomen.—Afr. austr.
galioides, *DC. Prod.* ii. 101.—Afr. austr.
galioides, E. Mey. ex Benth. in Ann. Wien. Mus. ii. (1838) 67 = genistoides.
genistoides, Sieber, ex Presl, Bot. Bemerk. 42 = galioides.
genistoides, *Vent. Dec. Gen.* 9.—Afr. austr.
grandifolia, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 492 = prnec.

CYCLOPIA :—
grandiflora, A. DC. Not. 8me Pl. Rar. Jard. Genèv. 29 = subtersata.
heterophylla, Eckl. & Zeyh. Enum. 154 = genistoides.
intermedia, E. Mey. Comm. Pl. Afr. Austr. 3 = subtersata.
laricina, E. Mey. l. c. 158 = tenuifolia.
latifolia, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 98.—Afr. austr.
latifolia, Eckl. & Zeyh. Enum. 155 = longifolia !
latifolia, E. Mey. Comm. Pl. Afr. Austr. 3 = subtersata.
lauxiflora, Benth. in Ann. Wien. Mus. ii. (1838) 67 = longifolia !
longifolia, *Vog. in Linnaea*, x. (1836) 595.—Afr. austr.
Meyeriana, Walp. in Linnaea, xiii. (1839) 454 = sessiliflora.
pubescens, *Eckl. & Zeyh. Enum.* 154.—Afr. austr.
sessiliflora, Eckl. & Zeyh. l. c. = subtersata.
sessiliflora, *E. Mey. Comm. Pl. Afr. Austr.* 4.—Afr. austr.
subternata, *Vog. in Linnaea*, x. (1835–36) 595.—Afr. austr.
tenuifolia, *Lehm. Ind. Sem. Hort. Hamb.* (1828) 16 ; *et in Linnaea*, v. (1830) 873.—Afr. austr.
teretifolia, Eckl. & Zeyh. Enum. 154 = genistoides.
ternata, Sweet, Hort. Brit. ed. I. 107 = Aspalathus ternginea.
triphylla, Sweet, l. c. = Argyrolobium sericeum.
Vogelii, Harv. in Harv. & Sond. Fl. Cap. ii. 6 = subternata.

CYCLOPIS, Guill. Archiv. Bot. ii. (1833) 468, sphalm. = **Cyclolepis**, Gill. (Compos.).

CYCLOPOGON, Presl, Rel. Haenk. i. 98. t. 13 (1827) = **Spiranthes**, Rich. (Orchid.).
alho-punctata, Rodrig. Orch. Nov. ii. 281.—Bras.
alpestris, Rodrig. l. c. 283 = S. alpestris !
atroviridis, Rodrig. l. c. 284.—Bras.
cearensis, Rodrig. l. c. 283.—Bras.
ovalifolium, Presl, Rel. Haenk. i. 98. t. 13 = S. Preslii.
procera, Regnell, ex Rodrig. Orch. Nov. ii. 290.—Bras.
trilineata, Rodrig. l. c. 284.—Bras.
variegata, Rodrig. l. c. 282.—Bras.

CYCLOPTERA, Endl. Ench. 118 (1841) = CYCLOPOGON, Presl = **Spiranthes**, Rich. (Orchid.).

CYCLOPTERA, Nutt. ex A. Gray, in Journ. Am. Sc. Ser. II. xv. (1853) 319 = **Abronia**, Juss. (Nyctagin.).
annua, Nutt. l. c. = A. Cyclopters.

CYCLOPTERYGIUM, Hochst. in Flora, xxxi. (1848) 175 = **Schouwia**, DC. (Crucifer.).
brevisiptum, Hochst. l. c. = S. Schimperi.
longisiptum, Hochst. l. c. 176 = S. arabica.

CYCLOPTYCHIS, E. Mey. in Drège, Zwei Pfl. Docum. 176 (1843), nomen ; et Harv. & Sond. Fl. Cap. i. 84 (1859). *CRUCIFERAE*, Benth. & Hook. f. i. 98.
polycaloides, *Sond. l. c.*—Afr. austr.
virgata, *E. Mey. in Drège, Zwei Pfl. Docum.* 176, nomen ; *et Harv. & Sond. l. c.*—Afr. austr.

CYCLOSANTHES, Poepp. in Froriep, Notiz. xxxi. 212 (1831) = **Cyclanthus**, Poit.

CYCLOSIA, Klotzsch, in Otto & Dietr. Allg. Gartenz. vi. (1838) 205 = **Mormodes**, Lindl. (Orchid.).
maculata, Klotzsch, l. c. 206 = M. pardinum.

CYCLOSTEGIA, Benth. in Bot. Reg. sub t. 1289 (1829) = **Blaholstia**, Willd. (Labiat.).
strobilifera, Benth. l. c. = E. strobilifera.

CYCLOSTEMON, Blume, Bijdr. 597 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 279.
DODECASTEMON, Hassk. in Verzl. Kon. Acad. Wetensch. iv. (1855) 141.
LANEASAGUM, Bedd. in Madr. Journ. Sc. Ser. II. xxii. (1861) 71.
PYCNOANDRA, Blume, Mus. Bot. Lugd. Bat. ii. 191 (1856).
SPHRAGIDIA, Thw. in Hook. Kew Journ. vii. (1855) 269. t. 10.
argutus, *Muell. Arg. in DC. Prod.* xv. ii. 485.—Afr. austr.
CYCLOSTEMON :—
coriaceus, *Baill. Etud. Gén. Euph.* 562.—Ins. Pacif. Caminigli, *Baill. l. c.*—Ins. Philipp.
cuspidatus, *Blume, Bijdr.* 599 = Ciconnicria reticulata.
eglandulosus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 240.—Ind. or.
floribundus, *Muell. Arg. in Flora*, xlvii. (1864) 532.—Afr. trop.
glomeratus, *Muell. Arg. l. c.*—Afr. trop.
indicus, *Muell. Arg. in Linnaea*, xxxii. (1863) 81.—Reg. Himal.
leiocarpus, *Kurz, in Journ. Bot. Ind.* iii. (1875) 330.—Ins. Nicobar.
longifolius, *Blume, Bijdr.* 598.—Malaya.
macrophyllus, *Blume, l. c.*—Malaya.
malabaricus, *Bedd. Forest. Man. Bot.* 199.—Ind. or.
macronatus, *Blume, Bijdr.* 598.—Java.
natalensis, *Harv. Thes. Cap.* ii. 64.—Afr. austr.
natalensis, *Muell. Arg. in DC. Prod.* xv. ii. 483.—Afr. austr.
occidentalis, *Muell. Arg. in Flora*, xlvii. (1864) 517.—Afr. trop.
parvifolius, *Muell. Arg. l. c.* 532.—Afr. trop.
priacipum, *Muell. Arg. l. c.* 531.—Afr. trop.
racemosus, Zipp. ex Span. in Linnaea, xv. (1841) 348 = Teysmanni.
rhacodiscus, *Muell. Arg. in DC. Prod.* xv. ii. 484.—Java.
serratus, *Blume, Bijdr.* 599.—Java.
stipularis, *Muell. Arg. in Flora*, xlvii. (1864) 531.—Afr. trop.
subsessilis, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 240.—Burma.
Teysmanni, *Muell. Arg. in DC. Prod.* xv. ii. 481.—Java.
zeylanicus, Baill. Etud. Gén. Euph. 562 = macrophyllus.

CYCLOSTIGMA, Hochst. ex Endl. Gen. Suppl. ii. 56 (1842) ; et in Flora, xxvii. (1844) 528 = **Voacanga**, Thou. (Apocyn.).
natalense, Hochst. l. c. = V. Dregei.

CYCLOSTIGMA, Klotzsch, in Seem. Bot. Voy. Herald, 10 (1853) = Croton, Linn. (Euphorb.).
abutiloides, Klotzsch, l. c. 105 = Croton abutiloides.
denticulatum, Klotzsch, l. c. = Croton panamensis.
Draco, Klotzsch, l. c. = Croton Draco.
hibiscifolium, Klotzsch, l. c. = Croton gossypiifolius.
panamense, Klotzsch, l. c. = Croton panamensis.
xalapense, Klotzsch, l. c. = Croton xalapensis.

CYCLOSTIGMA, Phil. Sert. Mendoc. Alt. 39 (1871) = **Leptoglossis**, Benth. (Solan.).
tenue, Phil. l. c.—Chili.

CYCLOTAXIS, Bion. Diagn. Ser. I. x. 48 (1849) = **Scandix**, Tourn. (Umbellif.).
palaestina, Boiss. l. c. 49 = S. palestina.
turgida, Boiss. & Bal. l. c. Ser. II. v. 108 = S. turgida.

CYCLOTHECA, Moq. in DC. Prod. xiii. ii. 37 (1849) = **Gyrostemon**, Desf. (Phytolac.).
australasica, Moq. l. c. 38 = G. Cyclotheca.

CYCNIA, Griff. Notul. iv. 481 ; Ic. Pl. Asiat. t. 605. f. 5 (1854). *ROSACEAE* ? Benth. & Hook. f. i. 606.
spinosa, *Griff. l. c.*—Burma.

CYCNIA, Lindl. Veg. Kingd. 548 (1847) = **Prinsepia**, Royle (Rosac.).

CYCNIUM, E. Mey. ex Benth. in Hook. Comp. Bot. Mag. i. (1835) 368. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 968.
adonense, *E. Mey. ex Benth. l. c.*—Afr. austr.
brachyoalyx, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xvii. (1868) 681.—Abyssin.
humifusum, *Benth. in Hook. f. Gen. ii.* 969.—Arab.
Krausianum, *Benth. in DC. Prod.* x. 508.—Afr. austr.
longiflorum, Eckl. & Zeyh. ex Oliver, in Trans. Linn. Soc. xxix. (1876) 128 = adonense.
racemosum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 368.—Afr. austr.
Sandersoni, *Harv. Thes. Cap.* i. 31.—Afr. austr.
tubatum, *Benth. in DC. Prod.* x. 508.—Afr. austr.

CYCNOCHES, Lindl. Gen. et Sp. Orch. 154 (1833). *ORCHIDEAE*, Benth. & Hook. f. iii. 552.
aureum, *Lindl. & Paxt. Flow. Gard.* iii. (1852–53) t. 75 & .—Am. centr.
barbatum, Lindl. in Journ. Hort. Soc. iv. (1849) 268 = Polycynchis barbata.
chlorochilon, *Lindl. Sert. Orchid.* (1838) t. 16.—Guiana.
cucullata, Lindl. Bot. Reg. sub t. 1951 = Loddigesii.
Dianae, *Reichb. f. in Bot. Zeit.* x. (1852) 686.—Am. centr.
Egertonianum, *Batem. Orchid. Mex. et Guat.* t. 40.—Guatemala.
Egertonianum, Lindl. Bot. Reg. (1846) t. 46 = Warszewiczii.
Hasgli, *Rodrig. Orch. Nov.* ii. 221.—Bras.
Lehmanni, Nichols. Dict. Gard. I. 419 = Lueddemannia Lehmanni.
Lehmanni, *Reichb. f. Otia Bot. Hamb.* 10.—Ecuador.
Loddigesii, *Lindl. Gen. et Sp. Orch.* 154.—Guiana.
maculatum, *Lindl. Bot. Reg.* (1840) *Misc.* 10.—Mexic.
muscifrons, Lindl. & Paxt. Flow. Gard. iii. (1852–53) 98 = Polycynchis muscifera.
pentadactylon, *Lindl. Bot. Reg.* (1843) *Misc.* 18.—Bras.
Pescatorei, Lindl. & Paxt. Flow. Gard. i. (1850–51) 128 = Lueddemannia Pescatorei.
stelliferum, Lodd. ex Reichb. f. in Walp. Ann. vi. 561 = Warszewiczii.
ventricosum, *Batem. Orchid. Mex. et Guat.* t. 5.—Guatemala.
Warszewiczii, *Reichb. f. in Bot. Zeit.* x. (1852) 784.—Am. centr.

CYCNOGETON, Endl. in Ann. Wien. Mus. ii. (1838) 210 = **Triglochin**, Linn. (Naiad.).
Hasgelii, Endl. Iconogr. t. 78 = T. procerum.
linnaeus, Endl. in Linnaea, xxvii. (1851) 225 = T. procerum.
procerum, Buchen. in Bremen, Abh. i. (1867) 234 = T. procerum.

CYCNOPODIUM, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 52 = **Graffenrieda**, DC. (Melast.).
latifolium, Naud. l. c. = G. latifolia.

CYCNOSERIS, Endl. in Bot. Zeit. i. (1843) 458 = Hypochoeris, Linn. (Compos.).
australis, Endl. l. c. 459 = H. glabra.

CYDENIS, Salisb. Gen. Pl. Fragm. 99 (1866) = **Narcissus**, Tourn. (Amaryll.).
poculiformis, Salisb. l. c. = N. poculiformis.

CYDISTA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 191 = **Bignonia**, Linn.
aequinoctialis, Miers, l. c. = B. aequinoctialis.
amoena, Miers, l. c. nomen.—Ins. S. Doming.
analoga, Miers, l. c. 192, nomen.—Bras.
blanda, Miers, l. c. nomen.—Venezuela.
difficilis, Miers, l. c. = B. difficilis.
diversifolia, Miers, l. c. = B. diversifolia.
eximia, Miers, l. c. nomen.—Uruguay.
incarnata, Miers, l. c. = B. incarnata.
magnifica, Miers, l. c. nomen.—Bras.
picta, Miers, l. c. = B. picta.
praspenea, Miers, l. c. 190.—Bras.
rubicunda, Miers, l. c. 192, nomen.—Bras.
sarmentosa, Miers, l. c. 191 = B. aequinoctialis.
Seemanni, Miers, l. c. 192 = B. difficilis.
spectabilis, Miers, l. c. 191 = B. aequinoctialis.
Vargasiana, Miers, l. c. 192 = Anemopaegma Vargasianum.

CYDONIA, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = **Pyrus**, Tourn. (Rosac.).
Bourgaeuti, Carr. in Rev. Hortic. (1872) 78.—An. Mis. *chinensis*, Trend. Nom. ed. II. i. 459 = P. sinensis, *Poir. communis*, Lobel. in Dalsam. Arb. ed. Nov. iv. 185. t. 86 = P. Cydonia.
europaea, Savi, Alb. Tosc. t. 90 = P. Cydonia.
italica, Spach, Hist. Vég. Phan. ii. 158 = Docynia indica.
japonica, Pers. Syn. ii. 40 = P. japonica.
Lagenaria, Lobel. in Dalsam. Arb. ed. Nov. vi. 225. t. 76 = P. japonica.
lobata, Bosc, Encycl. Agric. vii. 262 = P. Cydonia.

CYDONIA :—
lusitanica, Mill. Gard. Dict. ed. VIII. n. 3 = Pyrus Cydonia.
maliformis, Mill. l. c. n. 2 = P. Cydonia.
Maulei, T. Moore, in Fl. & Pomol. (1875) 49 = P. japonica.
oblonga, Mill. Gard. Dict. ed. VIII. n. 1 = P. Cydonia.
piriformis, Medic. Gesch. Bot. 80 = P. Cydonia.
silvestris, Risso, Hist. Nat. Product. Europ. 296 = P. Cydonia.
sinensis, Thou. in Ann. Mus. Par. xix. (1812) 145. t. 8 = P. sinensis, *Poir.*
speciosa, Sweet, Hort. Suburb. Lond. 113 = P. japonica.
Sumbuchia, Buch.-Ham. in D. Don, Prod. Fl. Nep. 237 = P. Cydonia.
tomentosa, C. Koch, Hort. Dendrol. 180 = P. Cydonia.
Umbuta, M. Roem. Syn. Rosif. 218 = P. Cydonia.
vulgaris, Pers. Syn. fl. 40 = P. Cydonia.

CYLACTIS, Rafin. in Am. Journ. Sc. i. (1819) 377 = **Rubus**, Tourn. (Rosac.).
lyncronotana, Rafin. in Journ. Phys. (Août 1819) t. 97 = R. trifloras.
montana, Rafin. in Am. Journ. Sc. i. (1819) 377 = R. trifloras.

CYLASTIS, Rafin. Sylva Tellur. 160 (1838) = **Rubus**, Tourn. (Rosac.).
arctica, Rafin. l. c. = R. arcticus.
ægopodioides, Rafin. l. c. = R. trifloras.

CYLBANIDA, Noronha, ex Tsl. in Ann. Sc. Nat. Sér. IV. viii. (1857) 135 = **Pittosporum**, Banks.

CYLICADENIA, Lem. Illustr. Hortic. ii. (1855) Misc. 9 = **Odontadenia**, Benth. (Apocyn.).
Hoerisii, Lem. l. c. 10 = O. speciosa !

CYLICHNANTHUS, Dulac, Fl. Hautes-Pyr. 260 (1867) = **Dianthus**, Linn. (Caryophyll.).
barbatus, Dulac, l. c. 261 = D. barbatus.
ciliatus, Dulac, l. c. = D. Seguieri.
deltoides, Dulac, l. c. 262 = D. deltoides.
fimbriatus, Dulac, l. c. = D. fimbriatus.
maculatus, Dulac, l. c. 261 = D. Armeria.
neglectus, Dulac, l. c. 262 = D. neglectus.
plumarius, Dulac, l. c. = D. plumarius.
prolifer, Dulac, l. c. 261 = D. prolifer.
sinibiflorus, Dulac, l. c. = D. Requieni.
virginalis, Dulac, l. c. = D. vaginatus.

CYLICHNIUM, Dulac, Fl. Hautes-Pyr. 68 (1867) = **Gaudinia**, Beauv. (Gramin.).
fragile, Dulac, l. c. = G. fragilis.

CYLICODAPHNE, Nees, in Wall. Pl. As. Rar. ii. 61 (1831); et Laurin. Expos. 18 (1838) = **Litsea**, Lam. (*Tetranthera*, Jacq.) (Laurin.).
accedens, Meisn. in DC. Prod. xv. i. 206.—Borneo.
Bindoniana, F. Muell. Fragm. v. 167 = L. Bindoniana.
chrysopleura, Blume, ex Meisn. in DC. Prod. xv. i. 198 (= *Tetranthera chrysopleura*).—Java.
costata, Blume, Mus. Bot. Lugd. Bat. ii. 12.—Malaya.
cuneata, Blume, Mus. Bot. Lugd. Bat. ii. 12.—Sumatra.
cuspidata, Blume, ex Meisn. in DC. Prod. xv. i. 198 (= *Tetranthera cuspidata*).—Java.
discolor, Meisn. in DC. Prod. xv. i. 204 = L. timoriana.
diversifolia, Blume, Mus. Bot. Lugd. Bat. ii. 12 = L. diversifolia.
Fauvettiana, F. Muell. Fragm. v. 168 = L. reticulata.
ferruginea, Blume, Mus. Bot. Lugd. Bat. ii. 11 = L. ferruginea.
floribunda, Blume, l. c. i. 287.—Ind. or.
fusca, Blume, l. c. ii. 13.—Sumatra.
fuscovenia, Meisn. in DC. Prod. xv. i. 207.—Malaya.
grisea, Blume, Mus. Bot. Lugd. Bat. ii. 13.—Sumatra.
hirta, Blume, l. c.—Java.
Hookeri, Meisn. in DC. Prod. xv. i. 209 = L. khasiana.
inflexoria, Blume, Mus. Bot. Lugd. Bat. ii. 11.—Ind. or.
Laughana, F. Muell. Fragm. v. 169 = L. ferruginea.
longifera, Meisn. in DC. Prod. xv. i. 205.—Ind. or.
lucida, Blume, Mus. Bot. Lugd. Bat. ii. 13 = L. lucida !
macrophylla, Blume, l. c. 14 = Actinodaphne macrophylla.

CYLICODAPHNE :—
magnifica, Miq. Fl. Ind. Bat. i. t. 886.—Sumatra.
malis, Meisn. in DC. Prod. xv. i. 207.—Java.
myristicæfolia, Meisn. l. c. 206.—Malaya.
Nessiana, Miq. Fl. Ind. Bat. i. t. 983.—Malacca.
nitida, Meisn. in DC. Prod. xv. i. 208.—Nepal.
Noronhiana, Blume, Mus. Bot. Lugd. Bat. i. 288 = L. Noronhae.
oblonga, Meisn. in DC. Prod. xv. i. 205 = L. oblonga.
obtusata, Meisn. l. c. 208.—Java.
obtusifolia, Nees, Syst. Laurin. 506 = L. obtusa.
ochracea, Blume, Mus. Bot. Lugd. Bat. ii. 13.—Sumatra.
pallida, Blume, l. c. 11.—Malaya.
polyneura, Meisn. in DC. Prod. xv. i. 205.—Java.
punctata, Blume, Mus. Bot. Lugd. Bat. ii. 13.—Borneo.
rigida, Meisn. in DC. Prod. xv. i. 201 = L. Gardneri.
robusta, Blume, Mus. Bot. Lugd. Bat. ii. 11 = L. robusta.
rubiginosa, Blume, l. c. 13.—Sumatra.
rubra, Blume, l. c. i. 288 = L. rubra.
sebifera, Blume, l. c. = L. sebifera, *Blume*.
Selangi, Miq. Fl. Ind. Bat. Suppl. 361.—Sumatra.
sessiliflora, Meisn. in DC. Prod. xv. i. 206.—Borneo.
sumatrana, Miq. Pl. Jungh. 184.—Sumatra.
tephrophylla, Meisn. in DC. Prod. xv. i. 207.—Java.
Thwaitesii, Meisn. l. c. 208, pro parte = L. ovalifolia.
tuberculata, Blume, Mus. Bot. Lugd. Bat. ii. 12.—Sumatra.
unita, Blume, l. c. i. 287.—Ins. Moluce.
veiutina, Blume, l. c. ii. 13.—Java.
Walkeri, Meisn. in DC. Prod. xv. i. 208 = L. Walkeri, *Wightiana*, Nees, in Wall. Pl. As. Rar. ii. 58 = L. Wightiana.
zeylanica, Meisn. in DC. Prod. xv. i. 201 = L. Gardneri.

CYLIDIUM, Rafin. in Am. Monthly Mag. (1819) 191, nomen (Convolv.).

CYLINDRACHNE, Reichb. Consp. 106 (1828), sphalm. = Cylindrocline, Cass. (Compos.).

CYLINDRIA, Lour. Fl. Cochinch. 69 (1790). (Genus incertæ sedis) Benth. & Hook. f. iii. 165.
rubra, *Lour. l. c.*—China.

CYLINDROCARPA, Regel, in Act. Hort. Petrop. v. (1877) 288. *CAMPANULACEÆ.*
Sewerzowi, *Regel, l. c.*—Turkestan.

CYLINDROCHILUS, Thw. Enum. Pl. Zeyl. 307 (1861) = **Microcoelia**, R. Br. (Orchid.).
pulchellus, Thw. l. c. = S. pulchellus.

CYLINDROCLINE, Cass. in Bull. Soc. Philom. (1817) 11. *COMPOSITÆ*, Benth. & Hook. f. ii. 255.
LEPIDOPOGON, Tausch, in Flora, xii. (1829) i. Erg. 87.
Commersonii, *Cass. l. c.*—Ins. Maurit.

CYLINDROLOBUS, Blume, Præf. Fl. Jav. 6 (1828) = **Eria**, Lindl. (Orchid.).

CYLINDROPUS, Nees, in Linnæa, ix. (1834) 308; et in Edinb. N. Phil. Journ. xv. (Oct. 1834) 366 = **Bolaria**, Berg. (Cyperac.).
juncifolius, Steud. Nom. ed. II. i. 459, sphalm. = seq.
junciformis, Nees, in Linnæa, ix. (1834) 308; et in Edinb. N. Phil. Journ. xv. (Oct. 1834) 366 = S. juncifolius.

CYLINDROSORUS, Benth. in Enum. Pl. Hueg. 62 (1837) = **Angianthus**, Wendl. (Compos.).
flavescens, Benth. l. c. = A. tomentosus.

CYLIPOGON, Rafin. in Journ. Phys. lxxxix. (1819) 97 = **Petalostemum**, Michx. (Legumin.).
capitatum, Rafin. l. c. = Dalea aurea.
virgatum, Rafin. l. c. = P. corymbosum vel Dalea lasiflora.

CYLISTA, Ait. Hort. Kew. ed. I. iii. 512 (1789). *LEGUMINOSÆ*, Benth. & Hook. f. i. 542.
albiflora, Sims, Bot. Mag. t. 1859 = Rhynchosia cyanosperma.
angustifolia, Eckl. & Zeyh. Enum. 258 = Rhynchosia uniflora.
angustifolia, E. Mey. in Linnæa, vii. (1832) 171 = Rhynchosia lescocias.
argentea, Eckl. & Zeyh. Enum. 258 = prsec.
ferruginea, Herb. Madr. ex Wight & Arn. Prod. 258 = Dunbaria ferruginea.
grewioides, Stend. Nom. ed. II. i. 459, sphalm. = *suaveolens*.
lancifolia, Eckl. & Zeyh. Enum. 259 = Rhynchosia Chrysoclas.
placatoria, Blanco, Fl. Filip. ed. I. 589 = Milletia splendens.
polystachya, Spreng. Syst. fv. Cur. Post. 269 = seq.
pycnostachya, DC. Prod. ii. 410 = Rhynchosia calycina.
reticulata, Heyne, in Wall. Cat. n. 5584 = Rhynchosia Heynei.
scariosa, *Roxb. Pl. Corom.* i. 64.—Ind. or.
suaveolens, R. Grah. in Wall. Cat. n. 5587 = Rhynchosia rufescens.
tomentosa, Roxb. Hort. Beng. 55; Fl. Ind. iii. 319 = Rhynchosia cyanosperma.
villosa, *Ait. Hort. Kew.* ed. II. iv. 300.—Afr. austr.

CLIXYLON, Llanos, Fragm. 59 (1851) = **Gymnanthera**, R. Br. (Asclep.).
heterophyllum, Llanos, l. c. = G. pedunculata.

CYLIZOMA, Neck. Elem. iii. 33 (1790) = **Derris**, Lour. (Legumin.).

CYLLENIUM, Schott, Gen. Aroid. t. 9 (1858) = **Biarum**, Schott (Aroid.).
hardwickeanum, Schott, Prod. Aroid. 65 = B. carduchorum.
Sprunerі, Schott, Gen. Aroid. t. 9 = B. Sprunerі.

CYMARIA, Benth. in Bot. Reg. sub t. 1992 (1829) *LABIATÆ*, Benth. & Hook. f. ii. 1222.
ANTHOCOMA, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 569.
acuminata, *Decne. Herb. Timor. Descr.* 71.—Ins. Philipp. et Timor.
dichotoma, *Benth. in Bot. Reg.* sub t. 1992.—Burma.
elongata, *Benth. l. c.*—Burma.
mollis, Miq. Fl. Ind. Bat. ii. 992 = acuminata.

CYMATION, Spreng. Syst. ii. 142 (1825) = **Ornithoglossum**, Salisb. (Liliac.).
laevigatum, Spreng. l. c. = O. glaucum.
undulatum, Spreng. l. c. = O. glaucum.

CYMATOCHLOA, Schlecht. in Bot. Zeit. xii. (1854) 821 = **Paspalum**, Linn. f. (Gramin.).
fluitans, Schlecht. l. c. 822 = P. repens.
repens, Schlecht. l. c. = P. repens.

CYMATOPTERA, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 309 = **Mancorvillea**, R. Br. (Crucif.).
chilensis, Turcz. l. c. = M. linearis.

CYMBA, Dulac, Fl. Hautes-Pyr. 117 (1867) = **Tofieldia**, Huds. (Liliac.).
calyculata, Dulac, l. c. = T. calyculata.

CYMBACHNE, Retz. Obs. vi. 36 (1791) = **Rottboellia**, Linn. f. (Gramin.).
alata, Beauv. Agrost. 159 = K. Cymbachne.
ciliata, Retz. Obs. vi. 36 = R. Cymbachne.

CYMBALARIA, Medic. Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 230 = **Linaria**, Mill. (Scrophul.).
communis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 84 = L. Elatine.
dentata, Schur, Enum. Fl. Transs. 488 = L. sparia.
Elatina, Gaertn. Mey. & Scherb. Fl. Wett. ii. 398 = L. Elatine.
hederacea, S. F. Gray, Nat. Arr. Brit. Pl. ii. 322 = L. Cymbalaria.
muralis, Gaertn. Mey. & Scherb. Fl. Wett. ii. 397 = L. Cymbalaria.
Pseudo-elatine, Schur, Enum. Pl. Transs. 488 = L. Elatine.

CYMBALARIA:—
spuria, Gaertn. Mey. & Scherb. Fl. Wett. ii. 398 = Linaria spuria.
vulgaris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1869) 126 = L. Cymbalaria.

CYMBALLARIA. Stead. Nom. ed. II. i. 459 (1840) = praec.

CYMBANTHES, Salisb. in Trans. Hort. Soc. i. (1812) 329 = **Androcymbium**, Willd. (Liliac.).
foetida, Salisb. l. c. = A. eucomoides.

CYMBARIA, Linn. Gen. ed. II. 282 (1742). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 975.
CYMBOCHASMA, Endl. Gen. 592 (1839).
borysthenica, *Pall. ex Schlecht. in Nov. Act. Hor. Petrop. Berol.* 107. t. 21.—Rossia.
daurica, *Linn. Sp. Pl.* 618.—Dahuria.
mongolica, *Maxim. in Mém. Acad. Pétersb. Sér.* VII. xxix. (1881) n. III. 66.—Mongol.

CYMBICARPOS, Stev. in Mém. Soc. Nat. Mosc. iv. (1832) 266 = **Astragalus**, Tourn. (Legum.).
Christianus, Stev. l. c. = A. Christianus.
tomentosus, Stev. l. c. = A. tomentosus.
trigonus, Stev. l. c. = A. trigonus.

CYMBIDIUM, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 70. *ORCHIDEAE*, Benth. & Hook. f. iii. 536.
APLECTRA, Raf. Cat. 13 (1824).
IRIDORCHIS, Blume, Coll. Orch. Jav. 90. t. 26 (1858).
JENSOA, Raf. Fl. Tellur. iv. 38 (1836).
OLIGASIS, Raf. l. c. ii. 51 (1836).
TRICHORHIZA, Lindl. ex Steud. Nom. ed. II. ii. 702 (1841).
acuiferum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 77 = Eulophia acuiesta.
adenoglossum, Lindl. in Journ. Linn. Soc. vi. (1862) 134 = Eulophia adenoglossa.
affine, Griff. Notul. iii. 336 = Cyperorchis Mastersii ?
alatum, Roxb. Hort. Beng. 63; Fl. Ind. iii. 459 = Thecostele alata.
albuciflorum, F. Muell. Fragm. i. 168 = madidum.
Allagnata, Buch.-Ham. ex Wall. Cat. n. 7327 = Vanda Roxburghii.
aloifolium, Blume, Bijdr. 378. t. 19 = pendulum ?
aloifolium, Heyne, ex Wall. Cat. sub n. 7351 = Diplocentum recurvum.
aloifolium, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 78.—Ind. or.
aloifolium, Wall. Cat. n. 7352 = Finlaysonianum.
altissimum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 74 = Oncidium altissimum.
altum, Willd. Sp. Pl. iv. 106 = Bletia verecunda.
amabile, Roxb. Hort. Beng. 63; Fl. Ind. iii. 457 = Phalaenopsis amabilis.
Andersonii, [Lamb. in] Andr. Bot. Rep. t. 651 = Cyrtopodium Andersonii.
angolense, *Reichb. f. in Flora*, xlviii. (1865) 188.—Afr. trop.
aphyllum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 78 = Dendrobium Plevardi.
appendiculatum, D. Don, Prod. Fl. Nep. 36 = Cremastra Wallichiana.
asamicum, *Linden, in Illustr. Hortic.* xxvii. (1881) 95, nomen.—Assam.
autumnale, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 72 = Eurina saccronota.
bambusifolium, Roxb. Hort. Beng. 63; Fl. Ind. iii. 460 = Arundina bambusifolia.
bicolor, Lindl. Gen. et Sp. Orch. 164 = alolifolium.
bituberculatum, Hook. Exot. Fl. ii. t. 116 = Liparis bituberculata.
boreale, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 76 = Calypso borealis.
Boweri, *F. Muell. in Wing, South S. Record* (Aug. 1888).—Ins. Solomon.
Buchanani, *Reichb. f. in Flora*, lxiv. (1881) 329.—Afr. austr.
Calcolaria, *Willd. Sp. Pl.* iv. 97.—Ind. or.
canaliculatum, *R. Br. Prod.* 331.—Australia.
candidum, *H. B. & K. Nov. Gen. et Sp.* i. 242.—N. Granat.
carnosum, *Griff. Notul.* iii. 339.—Reg. Himal.
chinense, Heynh. Nom. ii. 179 = sinense.
chloranthum, Lindl. Bot. Reg. (1843) Misc. 66.—Austral.

CYMBIDIUM:—
Clypeolum, Willd. Sp. Pl. iv. 99 = Liparis Clypeolum.
coccinum, Sw. in Schrad. Journ. ii. (1799) 214 = Arnithidium coccineum.
cochleare, Lindl. in Journ. Linn. Soc. iii. (1859) 28 = Cyperorchis cochlearis.
Corallorhiza, Sw. in Vet. Acad. Handl. Stockh. (1800) 238 = Corallorhiza innata.
cordatum, Londes, in Mém. Soc. Nat. Mosc. i. (1811) 289 = Listera cordata.
cordigerum, H. B. & K. Nov. Gen. et Sp. i. 341 = Epidendrum atropurpureum.
corniculatum, Spreng. Syst. iii. 722 = Pleurothallis corniculata.
crassifolium, Wall. Cat. n. 7357 = pendulum.
crispatum, *Thunb. Pl. Bras. Dec.* ii. 18.—Bras.
cucullatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 78 = Brassavola cucullata.
cuspidatum, *Blume, Bijdr.* 378.—Java.
cylindricum, Heyne, ex Wall. Cat. sub n. 7317 = Aërides cylindricum.
cyperifolium, *Wall. Cat.* n. 7373.—Reg. Himal.
Dayanum, Reichb. f. in Gard. Chron. (1869) 710 = eburneum.
densiflorum, Griff. Notul. iii. 337 = Cyperorchis elegans.
dependens, Lodd. Bot. Cab. x. 936 = Cirrhaea Loddigesii.
Devonianum, *Paxt. Mag. Bot.* x. (1843) 97.—Reg. Himal.
diurnum, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 78.—Venezuela.
eburneum, *Lindl. Bot. Reg.* (1847) t. 67.—Reg. Himal.
echinocarpon, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 71 = Dichaea echinocarpa.
elegans, Lindl. in Wall. Cat. n. 7354 = Cyperorchis elegans.
emifolium, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 77.—Ind. or. ; Japon.
equitans, Sw. l. c. 72 = Oberonia brevifolia.
erectum, Sw. in Schrad. Journ. ii. (1799) 226 = Cephalanthera pallens.
erectum, Wight, Ic. t. 1753 = aloifolium ?
erythrorhizum, Lindl. in Journ. Linn. Soc. iii. (1859) 30 = aloifolium.
estriatum, Lindl. ex Steud. Nom. ed. II. i. 460 = ensifolium.
falcatum, Sw. in Schrad. Journ. ii. (1799) 226 = Cephalanthera falcata.
Finlaysonianum, *Lindl. in Wall. Cat.* n. 7358.—Malaya.
flabellatum, *Spreng. Syst. iii.* 724.—Madag.
flabelliforme, Griseb. Fl. Brit. W. Ind. 629, sphalm. = Zygopetalum cochleare.
flabelliforme, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 73 = Zygopetalum cochleare.
flavescens, Llanos, Fragm. 96 = Cleisostoma amabile.
floridum, Lindl. Gen. et Sp. Orch. 121 = Bletia florida.
fragrans, Salisb. in Trans. Hort. Soc. i. (1812) 298 = chinense.
furvum, Willd. Sp. Pl. iv. 103 = Vanda furva.
fuscescens, Griff. Ic. Pl. As. iii. p. ii. t. 319 = Tainia fuscescens.
Gibsonii, Lindl. & Paxt. Flow. Gard. iii. (1852-53) 144 = lancifolium.
Gibsonii, *Wall. ex Voigt, Hort. Suburb. Calc.* 627, nomen.—Reg. Himal.
giganteum, Sw. in Schrad. Journ. ii. (1799) 234 (= *Cyrtopera gigantea*).—Afr. austr.
giganteum, *Wall. Cat.* n. 7355.—Reg. Himal.
glandulosum, *H. B. & K. Nov. Gen. et Sp.* i. 340.—Venezuela.
glaucum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 71 = Dichaea glauca.
globosum, Sw. l. c. 73 = Epidendrum globosum.
Goeringii, Reichb. f. in Walp. Ann. iii. 647 = virescens.
gomphocarpum, *Fitzg. in Journ. Bot.* xxi. (1883) 208. —Austral.
graminoides, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 71 = Dichaea graminoides.
grandiflorum, Griff. Notul. iii. 343 = Hookerianum.
grandiflorum, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 76.—Guiana.
guttatum, Willd. Sp. Pl. iv. 106 = Oncidium luridum.
hemistates, Lindl. Gen. et Sp. Orch. 163 = cyperifolium.
Hillii, *F. Muell. in Regel, Gartenfl.* (1879) 138.—Austral.

CYMBIDIUM:—
hirsutum, Willd. Sp. Pl. iv. 94 = Eilsanthus Caravata.
Hookerianum, *Reichb. f. in Gard. Chron.* (1866) 7.—Reg. Himal.
humile, Sm. ex Lindl. Coll. Bot. sub t. 37 = Coelogyne humilis.
Huttoni, *Hook. f. Bot. Mag.* t. 5676.—Java.
hyacinthinum, Sm. Exot. Bot. i. 117 = Bletia hyacinthina.
hyemale, Muhl. ex Willd. Sp. Pl. iv. 107 = Aplectrum hyemale.
imbricatum, Roxb. Hort. Beng. 63; Fl. Ind. iii. 460 = Pholidota imbricata.
iridifolium, *A. Cunn. in Lindl. Bot. Reg.* (1839) *Misc.* 84.—Austral.
iridifolium, Roxb. Hort. Beng. 63; Fl. Ind. iii. 458 = Oberonia iridifolia.
iridifolium, Sw. ex Steud. Nom. ed. II. i. 460 = Oncidium iridifolium.
iridioides, D. Don, Prod. Fl. Nep. 36 (Coelogynes sp. ?).—Nepal.
ixioides, D. Don, l. c. = Spathoglottis ixioides.
javanicum, Blume, Bijdr. 380 = lancifolium.
juncifolium, Willd. Sp. Pl. iv. 102 = Oncidium Cebolleta.
lancifolium, *Hook. Exot. Fl.* t. 51.—As. trop.
latifolium, Spreng. Syst. iii. 725 = Maxillaria latifolia.
latifolium, Sw. in Schrad. Journ. ii. (1799) 225 = Epipactis latifolia.
Leachianum, *Reichb. f. in Gard. Chron.* (1878) ii. 106. —Ins. Formosa.
liliifolium, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 76 = Liparis liliifolia.
limbatum, *Hook. ex Lindl. Gen. et Sp. Orch.* 165.—Ins. Trinitat.
linare, Herb. Heyne, ex Wall. Cat. sub n. 7312 = Saccolabium Wightianum.
linearе, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 72 = Isochilus linearis.
Loreslii, Sw. l. c. 76 = Liparis Loeselii.
longifolium, *D. Don, Prod. Fl. Nep.* 36.—Reg. Himal.
Loniceum, Reichb. f. in Gard. Chron. (1879) i. 332, 405 = giganteum ?
lutеum, Willd. Sp. Pl. iv. 106 = Chloraea crispa.
leucopodioides, *Willd. l. c.* 98.—Ind. or.
macrorrhizum, *Lindl. Gen. et Sp. Orch.* 162.—Reg. Himal.
madidum, *Lindl. Bot. Reg.* (1840) *Misc.* 9.—Austral.
Mannii, Reichb. f. in Flora, lv. (1872) 274 = pendulum.
marginatum, Lindl. Bot. Reg. t. 1530 = Maxillaria gracilis.
Mastersii, Griff. ex Lindl. l. c. (1845) t. 50 = Cyperorchis Mastersii.
Meyenii, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 433 = Arundina Meyenii.
micans, Schau. l. c. = ensifolium.
microrrhizon, Lindl. in Journ. Linn. Soc. iii. (1859) 29 = Cyperorchis Mastersii.
montanum, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 79.—Jamaic.
moschatum, Willd. Sp. Pl. iv. 98 = Dendrobium moschatum.
muricatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 71 = Dichaea muricata.
nervosum, Sw. l. c. 76 = Liparis nervosa.
nitidum, Roxb. Hort. Beng. 63; Fl. Ind. iii. 460 = Coelogyne ocellata.
nitidum, Wall. ex D. Don, Prod. Fl. Nep. 35 = Coelogyne nitida.
nodosum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 78 = Brassavola nodosa.
nutans, Sw. l. c. 77 = Geodorum purpureum.
ochroleucum, Lindl. Gen. et Sp. Orch. 168 = Maxillaria Camaridii.
odontorrhizum, Wall. ex Lindl. Fl. iv. 110 = Corallorhiza odontorrhiza.
ovatum, Willd. l. c. 101 = Dendrobium chlorops.
pallens, Sw. in Schrad. Journ. ii. (1799) 226 = Cephalanthera pallens.
palustre, Sw. l. c. = Epipactis palustris.
Parishii, Reichb. f. in Trans. Linn. Soc. xxx. (1874) 144 = eburneum.
parviflorum, Retzius, ex Lindl. in Journ. Linn. Soc. iii. (1859) 35 = Eria Retzianaidii.
pedicellatum, Sw. in Schrad. Journ. ii. (1799) 224 = Eulophia acuiesta.

CYMBIDIUM:—
pendulum, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 78.—Ind. or.
pictum, R. Br. Prod. 331 = Geodorum pictum.
plantaginifolium, *Willd. Sp. Pl.* iv. 101.—Ind. or.
plicatum, Harv. ex Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 368 = Eulophia scaiesta.
praecox, Sm. ex Lindl. Coll. Bot. sub t. 87 = Cologyne praecox.
praemorsum, Buch.-Ham. ex Wall. Cat. n. 7325 = Acampe multiflora.
praemorsum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 75 = Acampe excavata.
proliferum, Sw. l. c. 71 = Ponera prolifera.
pubescens, *Lindl. Bot. Reg.* (1840) *Misc.* 75.—Malaya.
pulchellum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 75 = Calopogon pulchellus.
purpureum, *Drapiez, Herb. Amat.* v. (1831) 313.—Ind. occ.
purpureum, Stend. Nom. ed. II. i. 559, in syn., sphalm. = Ornithidium vesicum.
pusillum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 74 = Oncidium trifidilabium.
ramosissimum, *Presl, Rel. Haenk.* i. 98.—Am. austr.
reflexum, R. Br. Prod. 331 = Liparis reflexa.
repens, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 71 = Bulbophyllum nutans.
rigidum, Willd. Sp. Pl. iv. 106 = Tetramicra rigida.
rubrum, Sw. in Schrad. Journ. ii. (1799) 226 = Cephalanthera rubra.
Sanderoni, Harv. Gen. S. Afr. Pl. ed. II. 360 = Ansellia gigantea.
sanguinum, Teijsm. & Binn. Cat. Hort. Bog. 51, sphalm. = seq.
sanguinolentum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 318.—Java.
Satyrium, Buch.-Ham. ex Wall. Cat. n. 7334 = Saccolabium baccosum.
scarabaeiforme, Parish, ex Reichb. f. in Gard. Chron. (1865) 942 = Luisia Psyche.
scriptum, Stend. Nom. ed. I. 248 = Eulophia scripta.
scriptum, Sw. in Schrad. Journ. ii. (1799) 218 = Grammatophyllum speciosum.
serrulatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 72 = Epidendrum serrulatum.
sinense, *Willd. Sp. Pl.* iv. 111.—China.
speciosissimum, D. Don, Prod. Fl. Nep. 35 = Cologyne cristata.
speciosum, Reinw. ex Lindl. in Journ. Linn. Soc. iii. (1859) 23 = Arundina densa.
spinosum, Reinw. ex Lindl. l. c. 14 = Dendrobium spinescens.
squamatum, Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 235 = Dipodium squamatum.
stapeliaeflorum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 319.—Java.
stapelioides, Link & Otto, Ic. Pl. Sel. iii. t. 52 = Trygonostalum stapelioides.
stenopetalum, Reinw. ex Lindl. Fol. Orch. Coelog. 14 = Cologyne longifolia.
striatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 77 = Bletia hyacinthina.
strictum, D. Don, Prod. Fl. Nep. 35 = Cologyne elatum.
suave, *R. Br. Prod.* 331.—Austral.
subulatum, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 73.—Jamaic.
syringodorum, Griff. Notul. iii. 338 = eburneum.
tabulare, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 77.—Afr. austr.
tenuifolium, Lindl. Gen. et Sp. Orch. 167 = Luisia tenuifolia.
tenuifolium, Wight, Ic. t. 1689 (non text.) = Luisia teretifolia.
tetrafolium, *Willd. Sp. Pl.* iv. 108.—Ind. or.
teretifolium, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 72 = Epidendrum teretifolium.
teuszianum, Sw. l. c. 75 = Vanda Roxburghii.
teuszioides, Roxb. Hort. Beng. 63; Fl. Ind. iii. 463 = Vanda Roxburghii.
tetragonifolium, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 71 = Platycoblastia tetragonifolia.
tetrapetalum, Sw. l. c. 74 = Oncidium tetrapetalum.
tigrinum, *Parish, ex Hook. Bot. Mag.* t. 5457.—Burmah.
trifidilabium, Spreng. Syst. iii. 731 = Platycoblastia trifidilabia.

CYMBIDIUM:—
trichocarpum, Sw. in Nov. Act. Soc. Upsal. vi. (1799) 71 = Dichaea trichocarpa.
tricolor, Mig. Choix Pl. Buitenz. t. 19 = Finlaysonianum.
trifidum, Sw. in Schrad. Neues Journ. i. (1805) 70 = Bletia verecunda.
triserva, *G. F. W. Mey. Prim. Fl. Esseq.* 258.—Guiana.
tristrum, Sw. in Schrad. Journ. ii. (1799) 214 = Coelia Bauerana.
triquetrum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 74 = Oncidium triquetrum.
triste, Roxb. Hort. Beng. 63; Fl. Ind. iii. 461 = Luisia teretifolia.
tristo, Willd. Sp. Pl. iv. 99 (Luisiae sp.).—Penang; N. Caled.
umbellatum, Spreng. Pugill. ii. 82 = Cirrhopetalum Thouarsii.
undulatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 76 = Oncidium carthagenense.
utriculatum, Bolus, in Journ. Linn. Soc. xx. (1884) 469 (Eulophiae sp.).—Afr. austr.
utriculatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 75 = Govenia utriculata.
variciflorum, Reichb. f. in Bonplandia, iv. (1856) 324 = chloranthum.
variegatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 74 = Oncidium variegatum.
verecundum, Sw. l. c. 75 = Bletia verecunda.
vestitum, *Sw. l. c.* 70.—Jamaica.
vexilliferum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 11.—Mexic.
violaceum, H. B. & K. Nov. Gen. et Sp. i. 341 = Catteya violacea.
virens, Reichb. f. in Walp. Ann. vi. 636, sphalm. = seq.
virescens, *Lindl. Bot. Reg.* (1838) *Misc.* 37.—Japon.
virescens, Willd. Sp. Pl. iv. 106 = Chloraea Piquichen.
viridiflorum, Griff. Itin. Notes, 136 = cyperifolium.
Wallichii, Lindl. Gen. et Sp. Orch. 165 = Finlaysonianum.
xilophyllum, Sw. in Schrad. Journ. ii. (1799) 226 = Cephalanthera ensifolia.
xiphiifolium, *Lindl. in Bot. Reg.* t. 529.—China.

CYMBOCARPA, Miers, in Proc. Linn. Soc. i. (1840) 61; et in Trans. Linn. Soc. xviii. (1841) 543. t. 38. f. 4 = Gymnosiphon, Blume (Burmanniac.).
refracta, Miers, ll. cc. 62, 544 = G. refractum.

CYMBOCARPUM, DC. Prod. iv. 186, in nota (1830); C. A. Mey. Verz. Pfl. Cauc. 182 (1831). *UMBELLIFERAE*. Benth. & Hook. f. i. 907.
TRAGOSMA, C. A. Mey. ex Ledeb. Fl. Ross. ii. 865 (1844).
anethoides, *DC. Prod.* iv. 186, in nota (1830).—Reg. Caucas.; Persia.
erythraeum, *Boiss. Fl. Orient.* ii. 1028.—Persia.
marginatum, *Boiss. l. c.*—Persia.
Wiedemanni, *Boiss. l. c.*—Anatolia.

CYMBOCHASMA, Endl. Gen. 698 (1839) = Cymbaria, Linn. (Scrophul.).

CYMBOCHASMIA, Reichb. Nom. 115 (1841) = praec.

CYMBONOTUS, Cass. in Dict. Sc. Nat. xxxv. 397 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 457.
Lawsonianus, *Gaudich. in Freyc. Voy. Bot.* 462. t. 86.—Austral.
Preissianus, Steetz, in Lehm. Pl. Preiss. i. 486 = Lawsonianus.

CYMBOPETALUM, Benth. in Journ. Linn. Soc. v. (1861) 69. *ANONACEAE*, Benth. & Hook. f. i. 37.
brasilianse, *Benth. l. c.*—Bras.
pendulliflorum, *Baill. Adansonia*, viii. (1868) 268.—Mexic.

CYMBOPHYLLUM, F. Muell. in Hook. Kew Journ. viii. (1856) 302 = Veronica, Tourn. (Scrophul.) (V. densifolia).

CYMBOPOGON, Spreng. Pugill. ii. 14 (1815) = Andropogon, Linn. (Gram.).
arabicus, Nees, ex Stend. Syn. Pl. Gram. 387 = A. laniger.

CYMBOPOGON:—
arundinaceus, Schult. Mant. ii. 457 = Anthistiria gigantea.
citriodorus, Link, Enum. Hort. Berol. i. 107 = A. Schoenanthus.
condensatus, Spreng. Syst. i. 289 = A. condensatus.
distans, Duthie, in [Atkins. Gaz. N. W. Ind.] 640.—Ind. or.
elegans, Spreng. Pugill. ii. 14 = A. cymbarius.
foliosus, Roem. & Schult. Syst. ii. 832 = A. bracteatus.
glandulosus, Spreng. Pugill. ii. 14 = Anthistiria prostrata.
glaucus, Schult. Mant. ii. 459 = A. macrourus.
hirtus, Nees, ex Baker, Fl. Maurit. 446 = Heteropogon hirtus.
Humboldtii, Spreng. Pugill. ii. 15 = A. bractestus.
iwarancusa, Schult. Mant. ii. 458 = A. iwarancusa.
laniger, Duthie, in [Atkins. Gaz. N. W. Ind.] 640.—Ind. or.
Martinianus, Schult. Mant. ii. 459 = A. Schoenanthus.
melancarpus, Spreng. Syst. i. 289 = Heteropogon acuminatus.
prostratus, Sweet, Hort. Brit. ed. II. 561 = Anthistiria prostrata.
reflexus, Roem. & Schult. Syst. ii. 834 = A. bractestus.
Schoenanthus, Spreng. Pugill. ii. 15 = A. Schoenanthus.
strictus, Boj. Hort. Maurit. 375 = A. foveolatus.

CYMBOSEMA, Benth. in Hook. Journ. Bot. ii. (1840) 61. *LEGUMINOSAE*, Benth. & Hook. f. i. 584.
roseum, *Benth. l. c.* 60, 61.—Bras.

CYMBOSERIS, Bolas. Diagn. Ser. I. xi. 50 (1849) = Phascostum, Cass. (Compos.).
palaestina, Boss. l. c. 51.—Syria.

CYMBOSTEMON, Spach, Hist. Vég. Phan. vii. 444 (1839) = Illicium, Linn. (Magnol.).
parviflorus, Spach, l. c. 446 = I. parviflorum.

CYMBURUS, Salisb. Parad. Lond. t. 49, 53 (1806) = Stachytarpheta, Vahl (Verben.).
mutabilis, Salisb. l. c. t. 49 = S. mutabilis.
squamosus, Rafin. Fl. T.llur. ii. 103 = Elytraria caulescens.
urticaefolius, Salisb. Parad. Lond. t. 53 = S. urticaefolia.

CYMBLONEMA, Presl, Epim. Bot. 210 (1849) *RUBIACEAE*, Benth. & Hook. f. i. 735.
memecyloides, *Presl, l. c.*—Ins. Philipp.

CYMICIFUGA, Reichb. in Moesl. Handb. ii. 942, (1828) sphalm. = Cimicifuga, Linn. (Ranunc.).

CYMINON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 65 = Cuminum, Linn. (Umbellif.).
longeinvolucellatum, St. Lag. l. c. = Cum. Cyminum.

CYMINOSMA, Gaertn. Fruct. i. 280. t. 58 (1788) = Acronychia, Forst. (Rutac.).
Ankenda, Gaertn. l. c. = A. laurifolia.
chinensis, G. Don, Gen. Syst. i. 781 = A. laurifolia.
oblongifolia, A. Cunn. ex Hook. Bot. Mag. t. 3322 = A. laevis.
odorata, DC. Prod. i. 722 (= *Jambolifera odorata*).—Cochinch.
pedunculata, Roxb. Hort. Beng. 88; Fl. Ind. ii. 229 = A. laurifolia.
resinosa, DC. Prod. i. 722 = A. laurifolia.

CYMINUM, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 149, in syn. = Microcnidium, Boiss. (Umbell.).
minutum, Boiss. l. c. = M. tenuifolium.

CYMODOCEA, Konig, in Ker. & Sims, Ann. Bot. ii. (1806) 96. t. 7. *NAIADACEAE*, Benth. & Hook. f. iii. 1018.
DIPLANTHERA, Thou. Gen. Nov. Madag. 3 (1809).
HALODULE, Endl. Gen. Suppl. i. 1268 (1841).
PHUCAGROSTIS, Willd. Sp. Pl. iv. 849 (1805).
aequorea, *Konig, in Ker. & Sims. Ann. Bot.* ii. (1806) 96.—Mare Mediterr.
antarctica, *Endl. Gen.* 220.—Austral.
ciliata, *Ehrenb. ex Aschers. in Sitzb. Ges. Naturf. Fr. Berl.* (1867) 3.—Austral.
isoetifolia, *Aschers. l. c.*—Austral.
manatorum, *Aschers. l. c.* (1868) 19.—Ind. occ.
nodosa, Aschers. l. c. (1869) 4 = sequorea.

CYMODOCEA:—
Presnxiana, *Webb & Berth. Phyt. Canar.* iii. 302.—Ins. Canar.
rotundata, *Ascherson & Schweinf. in Sitzb. Ges. Naturf. Fr. Berl.* (1870) 84.—Mare Rubrum.
serrulata, *Aschers. & Magnus, l. c.*—Trop.
Webbiana, *A. Rich. in Ann. Sc. Nat.* Sér. II. xi. (1839) 355.—Ins. Canar.
sotteriifolia, F. Muell. Census, 121 = antarctica.

CYMOPTERIBUS, Bockl. in Proc. Acad. Sc. Philad. 1861 (1862) 455, sphalm. = **Cymopterus**, Raf. (Umbell.).

CYMOPTERUS, Raf. Journ. Phys. lxxxix. (1819) 100. *UMBELLIFERÆ*, Benth. & Hook. f. i. 911.
LEPTOCNEMIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 624 (1840).
ORDOSIS, Raf. in Ser. Bull. Bot. (1830) 217.
PHELLOPTERUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 628 (1840).
PTERYXIA, Nutt. l. c. 634 (1840).
alkiformis, Torr. & Gray, Fl. N. Am. i. 625 = terebinthinus.
alpinus, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 408.—Am. bor.
anisatus, *A. Gray, in Proc. Acad. Philad.* (1863) 68.—Am. bor.
bipinnatus, *S. Wats. in Proc. Am. Acad.* xx. (1885) 368.—Am. bor.
campestris, *Torr. & Gray, Fl. N. Am.* i. 624.—Am bor.
cinerarius, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 535.—Calif.
Fendleri, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) (*Pl. Fendl.* 56).—Texas.
foeniculaceus, Torr. & Gray, Fl. N. Am. i. 624 = terebinthinus.
glaber, *Black, in Hodgson's Res. Hakodate*, 385.—Japon.
glaucus, *Nutt. in Journ. Acad. Philad.* vii. (1834) 28.—Am. bor.
globosus, *S. Wats. in Proc. Am. Acad.* xl. (1876) 141.—Am. bor. occ.
glomeratus, *DC. Prod.* iv. 204.—Am. bor.
littoralis, A. Gray, in Mem. Am. Acad. vi. (1859) 391 = Phellopterus littoralis, *Benth. & Hook. f.*
longipes, *S. Wats. in Bot. King's Exp.* 124.—Am. bor. occ.
macrorhizus, Bockl. in Proc. Acad. Sc. Philad. 1861 (1862) 455 = montanus.
montanus, *Torr. & Gray, Fl. N. Am.* i. 624.—Am. bor. occ.
nevadensis, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 536.—Am. bor. occ.
nivalis, *S. Wats. in Bot. King's Exp.* 123.—Am. bor. occ.
purpureus, *S. Wats. in Am. Natural.* vii. (1873) 300.—Am. bor.
terebinthinus, *Torr. & Gray, Fl. N. Am.* i. 624.—Am. bor. occ.
thapsoides, Torr. & Gray, l. c. 625 = terebinthinus.

CYNANCHICA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 398 = **Asperula**, Linn. (Rubiac.).
tenuifolia, Fourr. l. c. = A. cynanchica.
tinctoria, Fourr. l. c. = A. tinctoria.

CYNANCHUM, Linn. Gen. ed. I. 69 (1737). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 762.
DIPLOGLOSSUM, Meisn. Gen. 269 (1840).
ENDOTROPIA, Endl. Gen. 591 (1838).
EXOSTEMA, Boj. ex Decne. in DC. Prod. viii. 528, 580, in syn. (1844).
PRANCHUM, Neck. Elem. i. 254 (1790).
SYMPHYOGLOSSUM, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 256.
ZIERVOGLIA, Neck. Elem. i. 254 (1790).
abyssinicum, *Decne. in DC. Prod.* viii. 548.—Abyss.
acuminatum, Buch.-Ham. ex Decne. l. c. 622 = Cymnema acuminatum.
acuminatum, *Humb. & Bonpl. ex Schult. Syst.* vi. 111.—Am. austr.
acuminatum, Thunb. Obs. in Cynanch. 5 = Pentatropis microphylla.
acutum, *Linn. Sp. Pl.* 213.—Europ. austr.; Orient; As. bor.
africanum, Hoffmgg. Verz. Pfl. 54 = pilosum.
alatum, *Wight & Arn. Contrib.* 57.—Ind. or.
album, Pers. Syn. i. 274 = Sarcostemma Dombeyanum.

CYNANCHUM:—
altissimum, Jacq. Select. Am. 84 = Gonolobus altissimus.
androsaemifolium, Vahl, ex Decne. in DC. Prod. viii. 500 = Pentopetia androsaemifolia.
angustifolium, Pers. Syn. i. 274 = Vincetoxicum palustre.
angustifolium, Wight & Arn. Contrib. 57 = C. Callialata.
annularia, Heyne, ex Wight, l. c. 55 = Holostemma Rheedianum.
aphyllum, Linn. Syst. ed. XII. iii. App. 235 = Sarcostemma aphyllum.
aphyllum, Vell. Fl. Flum. 122; iii. t. 88 (= *Amplustelma aphyllum*).—Bras.
arboreum, *Forsk. Fl. Aegypt. Arab.* 58.—Arabia.
Argel, Delile, Fl. Egypte, 197. t. 20 = Solenostemma Argel.
Arnottianum, *Wight, Contrib.* 58.—Reg. Himal.
asperum, *Will. Gard. Dict.* ed. VIII. n. 6.—Mexic.
Asterion, Vell. Fl. Flum. iii. t. 75 = Blepharodon Asterion.
astrictum, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 63 = Leptadenia reticulata.
atratum, Bunge, Enum. Pl. Chin. Bor. 45 = Vincetoxicum atratum.
atropurpureum, D. Dietr. Syn. Pl. ii. 906 = Schizoglossum atropurpureum.
auriculatum, Buch.-Ham. ex Wight, Contrib. Bot. Ind. 56 = C. Callialata.
auriculatum, *Royle, ex Wight, l. c.* 58.—Reg. Himal.
barbatum, Linn. f. ex Schult. Syst. vi. 108, in obs. = Vincetoxicum nigrum.
bicolor, Andr. Rep. Bot. t. 562 = Daemia extensa ?
bidens, D. Dietr. Syn. Pl. ii. 906 = Schizoglossum bidens.
bivoltatum, Hook. & Arn. Bot. Beech. Voy. 85 = Oxypetalum Hookeri.
boerhaviifolium, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 294.—Chili.
Bonplandianum, Koem. & Schult. Syst. vi. 98 = Vincetoxicum Kunthii.
Bowmanum, Steud. Nom. ed. II. i. 461 = Glossonema Bowmanum.
bracteatum, Thunb. Obs. in Cynanch. 7 = Tylophora pauciflora.
Brownianum, Schult. Syst. vi. 97 = pedunculatum, *R. Br.*
Brunonis, Gill. ex Hook. Journ. Bot. i. (1834) 299 = Oxypetalum Brunonis.
Bungei, *Decne. in DC. Prod.* viii. 549.—China.
Callialata, *Buch.-Ham. ex Wight, Contrib. Bot. Ind.* 56.—Ind. or.; Burma.
Candollianum, Gill. ex Hook. & Arn. in Hook. Journ. Bot. i. (1834) 291 = Melinia Candolleana.
capense, Linn. f. Suppl. 168 (= *Bumbuvia elliptica*).—Afr. austr.
caperse, Sieber, ex Decne. in DC. Prod. viii. 612 = Tylophora laevigata.
capillare, *Thunb. Obs. in Cynanch.* 8.—Zeylan.; Java.
carnosum, Hort. ex Decne. in DC. Prod. viii. 550 = Hoya carnosa.
carolinense, Jacq. Coll. ii. 288 = Gonolobus carolinensis.
caudatum, Vell. Fl. Flum. 121; iii. t. 77 (= *Orthoria virgata*!).—Bras.
chirenthoides, Domb. ex Decne. in DC. Prod. viii. 587 = Oxypetalum Hookeri.
chinense, *R. Br. in Mem. Wern. Soc.* i. (1809) 44.—China.
ciliatum, Vell. Fl. Flum. 122; iii. t. 84 = Fischeria retusifolia.
cirrhosum, Sw. in Cat. Dorpat. (1810); ex Schult. Syst. vi. 111 (quid ?).
cissoroum, Jacq. Select. Am. 87 = Sarcostemma Brevei.
congestum, Vell. Fl. Flum. 122; iii. t. 82 (= *Orthoria congesta*).—Bras.
convolvuloides, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 294 = Rouliuia convolvuloides.
contiguum, Koch, Syn. Fl. Germ. ed. II. 566 = Vincetoxicum officinale.
cordatum, Vell. Fl. Flum. iii. t. 56 = Rouliuia ferruginea.
cordifolium, *D. Dietr. Syn. Pl.* ii. 905.—Afr. austr.
cordifolium, Retz. Obs. ii. 15 = Daemia extensa.
corymbosum, Pav. ex Decne. in DC. Prod. viii. 601 = Fischeria peruviana.

CYNANCHUM:—
corymbosum, *Wight, Contrib.* 56.—Reg. Himal.; Malaya.
crassifolium, R. Br. in Mem. Wern. Soc. i. (1809) 46 (= *Cynoctonum crassifolium*).—Afr. austr.
crispiflorum, Sw. Prod. Veg. Ind. Occ. 53 = Fischeria scandens.
crispum, Thunb. Prod. Pl. Cap. 46 = Gomphocarpus crispus.
cuspidatum, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 298 = Metastelma cuspidatum.
cuspidatum, Thunb. Obs. in Cynanch. 6 = Gymnema perglarioides.
daharicum, Willd. ex Decne. in DC. Prod. viii. 583 = roseum.
Dalbousiæ, *Wight, Contrib.* 55.—Reg. Himal.
De-Filippii, *Delp. in Mem. Acc. Tor.* Ser. II. xxvi. (1871) 180.—Hongkong.
deltoideum, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 208 = chinense.
deltoideum, *Hook. f. Fl. Brit. Ind.* iv. 24.—Reg. Himal.
denticulatum, Vahl, Eclog. Am. ii. 28 (ordo !).—Guiana.
diffusum, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 298.—Reg. Argent.
Dioscoridis, Thunb. Fl. Ceilan. 5, nomen (quid ?).—Zeyl.
discolor, Sims, in Bot. Mag. t. 1278 = Gonolobus obliquus.
echinatum, Thunb. Obs. in Cynanch. 8 = Daemia extensa.
erectum, Linn. Sp. Pl. 213 = Marsdenia erecta.
erubescens, *R. Br. in Mem. Wern. Soc.* i. (1809) 45.—Austral.
euphorbioides, D. Dietr. Syn. Pl. ii. 906 = Schizoglossum euphorbioides.
excelsum, *Decf. Fl. Atlant.* i. 212.—Afr. bor.
extensum, Jacq. Misc. ii. 358 = Daemia extensa.
filiforme, Jacq. Select. Am. 86 = Sarcostemma Jacquini.
filiformis, *Linn. f. Suppl.* 160.—Afr. austr.
fimbriatum, H. B. & K. Nov. Gen. et Sp. iii. 208. t. 254 = Metaplexis fimbriata.
fissum, *Pomel, Nouv. Mat. Fl. Atl.* 81.—Algeria.
flavens, Thunb. Obs. in Cynanch. 7 = Tylophora paucifiora.
flavum, Thunb. ex Hook. i. Fl. Brit. Ind. iv. 45 = Tylophora asthmatica.
floribundum, *R. Br. in Mem. Wern. Soc.* i. (1809) 45.—Austral.
floridum, *Vell. Fl. Flum.* 123; iii. t. 86.—Bras.
foetidum, H. B. & K. Nov. Gen. et Sp. iii. 202 = Rouliuia Jacquini.
foetidum, L'Hérit. ex Decne. in DC. Prod. viii. 602 = Gonolobus prostratus.
fruticulosum, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 352. t. 12 = Vincetoxicum fruticulosum.
funale, *Poir. Encyc. Suppl.* iii. 427.—Ins. S. Doming.
fuscatum, Link, Enum. Hort. Berol. i. 250 = Vincetoxicum luteum.
fuscum, Schult. Syst. vi. 111 = Vincetoxicum nodium.
gangliosum, Vell. Fl. Flum. 119; iii. t. 73 = Gonolobus gangliosus.
glaucum, *Wall. ex Wight, Contrib.* 55.—As. or.; Reg. Himal.
gracillimum, Wall. l. c. 57 = Adelostemma gracillimum.
grandiflorum, Cav. Ic. i. 14 = Gonolobus grandiflorus.
hamatum, D. Dietr. Syn. Pl. ii. 906 = Schizoglossum hamatum.
hastatum, *Lam. Encyc.* ii. 286.—Orient.
hastatum, Pers. Syn. i. 276 = Leptadenia lanciflolia.
heteromorphum, *Vatke, in Linnaea*, xl. (1876) 215.—Abyss.
heterophyllum, Delile, Cent. Pl. Afr. 47. t. 68 = Leptadenia heterophylla.
Heydei, *Hook. f. Fl. Brit. Ind.* iv. 25.—Tibet. occ.
Heynianum, Schult. Syst. vi. 102 = C. Callialata !
hirsutum, Vell. Fl. Flum. 121; iii. t. 78 (Distance sp.).—Bras.
hirsutum, Pers. Syn. i. 274 = Gonolobus hirsutus.
hirsutum, Vahl, Eclog. Am. iii. 24 = Macrosceptrum obovatum.
hirtum, *Blanco, Fl. Philip. ed. I.* 269.—Ins. Philipp.
hirtum, Linn. Sp. Pl. 213 = Gonolobus obliquus.
Hookerianum, Steud. Nom. ed. II. i. 462 = Metaplexis cuspidatum.

CYNANCHUM:—
Humboldtianum, *Schult. Syst.* vi. 105.—Venezuela.
humile, Falc. ex Hook. f. Fl. Brit. Ind. iv. 26 = Jacquemontianum.
Hystrix, Vell. Fl. Flum. 119; iii. t. 79 = Gonolobus Hystrix.
incongicivum, Griff. Notul. iv. 66 = pauciflorum.
indicum, Burm. f. Fl. Ind. 70 = Tylophora asthmatica.
indicum, Lam. Encyc. ii. 286 = Tylophora laevigata.
inodorum, Lour. Fl. Cochinch. 106 = Gymnema inodorum.
inundatum, Lem. ex Decne. in DC. Prod. viii. 606 = Trychanthera Bergeriana.
Ipecacuanha, Willd. in Jahrb. Pharm. (1795) 169 = Tylophora asthmatica.
Jacquemontianum, *Decne. in Jacquem. Voy. Bot.* 106. t. 113.—Reg. Himal.
japonicum, *R. Morr. & Decne. in Bull. Acad. Brux.* iii. (1836) 172.—Japon.
kraussianum, Wight, Contrib. 58 = C. Vincetoxicum.
laeve, Pers. Syn. i. 274 = Gonolobus laevis.
laevigatum, Vahl, ex Decne. in DC. Prod. viii. 612 = Tylophora laevigata.
lanceolatum, H. B. & K. Nov. Gen et Sp. iii. 203 = Vincetoxicum Kunthii.
lanceolatum, Poir. Encyc. Suppl. ii. 430 = Gymnema subvolubile.
lanciflolium, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 294.—Chili.
lanciflolium, Schum. & Thonn. Beskr. Guin. Pl. 150 = Leptadenia lanciflolium.
latifolium, Schum. & Thonn. l. c. = Leptadenia lanciflolia.
laxum, Bartl. in Koch, Taschenb. 350 = Vincetoxicum officinale.
leucanthum, Jacq. ex J. F. Gmel. Syst. 442 = Sarcostemma Jacquini.
longiflorum, Jacq. Select. Am. 85 (= *Schubertia longiflora*).—Am. austr.
longifolium, Martens, Reise Vened. ii. 570. t. 6 = acutum.
luteolum, Vahl, ex Decne. in DC. Prod. viii. 589 = Melinis Commersoniana.
luteum, Spreng. Syst. i. 892 = glaucum.
lutum, Steud. Nom. ed. 250 = Vincetoxicum luteum.
lycioides, Steud. l. c. ed. II. i. 462; D. Dietr. Syn. Pl. ii. 906 = Tylophora lyciodes.
Macraei, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 295 = Astephanus geminiflorus.
macrorhizon, *Carr. in Rev. Hortic.* (1877) 435.—Japon.
macrophyllum, Pers. Syn. i. 274 = Gonolobus carolinensis.
macrophyllum, *Thunb. Obs. in Cynanch.* 7.—Java.
Magaie, *Buch.-Ham. ex Dillw. Rev. Hort. Malab.* 38.—Ind. or.
maritimum, Jacq. Select. Am. 88. t. 56 = Lachnostoma maritima.
maritimum, Salisb. Prod. 149 = monspeliacum.
mauritianum, Bot. Hort. Maurit. ex Decne. in DC. Prod. viii. 613 = Tylophora laevigata.
mauritianum, Lam. Encyc. ii. 286 = Camptocarpus mauritianus.
medium, R. Br. in Mem. Wern. Soc. i. (1809) 48 = Vincetoxicum nigrum.
melanophleum, *Willd. Enum. Hort. Berol. Suppl.* 13, nomen.—Hab.?
melanthum, Poir. Encyc. Suppl. ii. 430 = Vincetoxicum medium.
micranthum, Thunb. Obs. in Cynanch. 5 = Tylophora asyanica.
microphyllum, H. B. & K. Nov. Gen. et Sp. iii. 204. t. 236 = Vincetoxicum microphyllum.
microphyllum, *Rafin. New Fl. Am.* iv. 59.—Am. bor.
minus, C. Koch, in Linnaea, xix. (1847) 97 = Vincetoxicum fuscatum.
mollis, *Steud. Nom.* ed. II. i. 462; *D. Dietr. Syn. Pl.* ii. 905.—Afr. austr.
monoplectum, Auch. ex Decne. in DC. Prod. viii. 616 = Marsdenia erecta.
monopleurum, Lam. Sp. Pl. 312 = acutum.
monosiphonus, *Spreng. Syst.* i. 851.—Reg. Argent.
mucronatum, Andr. Bot. Rep. t. 516 = Tylophora asthmatica.)
mucronatum, H. B. & K. Nov. Gen. et Sp. iii. 204. t. 236 = Schubertia mucronata.

CYNANCHUM:—
multiflorum, Spreng. Syst. iv. Cur. Post. 110 (= *Schubertia multiflora*).—Bras.
myrtifolium, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 293.—Chili.
myrtifolium, Poepp. ex Decne. in DC. Prod. viii. 580 = Tassadia Poeppigiana.
nanum, Buch.-Ham. ex Wight, Contrib. 59 = Pentabothra nana.
nigricans, Vahl, ex Decne. in DC. Prod. viii. 590 = Rhyssotelma nigricans.
nigrum, Cav. Ic. ii. 45. t. 159 = Gonolobus niger.
nigrum, Pers. Syn. i. 274 = Vincetoxicum nigrum.
nitidum, Vell. Fl. Flum. 120; iii. t. 74 = Blepharodon diffusum?
nivale, Nym. Syll. 108 = Vincetoxicum nivale.
nummularifolium, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 293.—Chili.
obliquum, Jacq. Coll. i. 148 = Gonolobus obliquus.
obtusifolium, Linn. f. Suppl. 169 (= *Cynoctonum crassifolium*).—Afr. austr.
obtusifolium, Rich. ex Pers. Syn. i. 274 = Gonolobus obtusifolius.
odoratissimum, Lour. Fl. Cochinch. 106 = Pergularia odoratissima.
odoratum, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 294 = Morrenia odorata.
oleaefolium, Nect. Voy. Egypte, 20. t. 3 = Solenostemma Argel.
orientale, Willd. ex Spreng. Syst. i. 851 = longifolium?
ovalifolium, *Wight, Contrib.* 57.—Malaya.
ovatum, Thunb. Obs. in Cynanch. 6 = Leptadenia reticulata.
pallidum, Salisb. Prod. 149 = Marsdenia erecta.
parviflorum, Link, ex Fourn. in Mart. Fl. Bras. vi. iv. 210 = Metastelma odoratum.
parviflorum, Sieber, ex Steud. Nom. ed. II. i. 462 = Sarcostemma Brownii.
parviflorum, Sw. Prod. Veg. Ind. Occ. 58 = Metastelma parviflorum.
pauciflorum, *R. Br. in Mem. Wern. Soc.* i. (1809) 45.—Ind. or.
pedunculare, Lam. Encyc. ii. 286 (= Sarcostemmatis sp.).—Guiana.
pedunculatum, *R. Br. in Mem. Wern. Soc.* i. (1809) 45.—Austral.
pedunculatum, *Thunb. Obs. in Cynanch.* 6.—Zeylan.
pendulum, Poir. Encyc. Suppl. ii. 429 (= Leptadeniae sp.).—Afr. trop.
Phylladen, Vell. Fl. Flum. 120; iii. t. 76 = Fischeria adenophylla.
pictum, Vahl, Eclog. Am. ii. 24 = Marsdenia picta.
pilosum, R. Br. in Mem. Wern. Soc. i. (1809) 46 (= *Cynoctonum pilosum*).—Afr. austr.
planiflorum, Jacq. Select. Am. 82 = Gonolobus planiflorus.
planifolium, Linn. Syst. ed. XII. 192 = Gonolobus planiflorus.
procumbens, Medic. ex Steud. Nom. ed. II. i. 462 = Marsdenia erecta.
prostratum, Cav. Ic. i. 5. t. 7 = Gonolobus prostratus.
puberulum, *F. Muell. ex Benth. Fl. Austral.* iv. 388.—Austral.
pubescens, Bunge, Enum. Pl. Chin. Bor. 44 = chinense.
purpurascens, *E. Morr. & Decne. in Bull. Acad. Brux.* iii. (1836) 173.—Japon.
pusillum, Spreng. ex Fourn. in Mart. Fl. Bras. vi. iv. 223 (= *Orthosia virgata*).—Bras.
pycnostachyum, Forsk. Fl. Aegypt. Arab. 58 = Leptadenia Spartum.
racemosum, Jacq. Select. Am. 81 = Roulinia jacquini.
radians, Lam. Encyc. i. 286 = Stelbellia radians.
rectum, Forsk. ex Decne. in DC. Prod. viii. 616 = Marsdenia erecta.
reflexum, Vahl, ex Decne. l. c. 581 = Oxypetalum Banksii.
reticulatum, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 26 = Marsdenia Hamiltonii.
reticulatum, Retz. Obs. ii. 15 = Leptadenia reticulata.
roseum, *R. Br. in Mem. Wern. Soc.* i. (1809) 47.—Sibir.
rostratum, Vahl, Symb. Bot. iii. 45 = Gonolobus rostratus.
rotatum, Vell. Fl. Flum. 121; iii. t. 79 = Exolobus rotatus.
rotundifolium, Thunb. ex Decne. in DC. Prod. viii. 582 (= *Cynoctonum crassifolium*).—Afr. austr.

CYNANCHUM:—
Roylei, *Wight, Contrib.* 58.—Tibet. occ.
scabrum, *Schum. & Thonn. Beskr. Guin. Pl.* 152.—Afr. trop.
Schimperi, Hochst. in Flora, xxiv. (1841) i. Intell. 25 = Marsdenia Schimperi.
scoparium, Nutt. in Ann. Journ. Sc. v. (1822) 291 = Vincetoxicum scoparium.
senegalense, Sieber, ex Decne. in DC. Prod. viii. 621 = Gymnema subvolubile.
serpyllifolium, H. B. & K. Nov. Gen. et Sp. iii. 205 (= *Orthosia serpyllifolia*).—Ecuador.
sibiricum, R. Br. in Mem. Wern. Soc. i. (1809) 48 = Vincetoxicum sibiricum.
sibiricum, Willd. in Ges. Naturf. Fr. Neue Schr. ii. (1799) 124. t. 5. f. 2 = acutum.
speciosum, Nym. Syll. 108 = Vincetoxicum speciosum.
stellatum, Vell. Fl. Flum. 121; iii. t. 80 = Fischeria stellata.
suberosum, H. B. & K. Nov. Gen. et Sp. iii. 205 = Metastelma suberosum.
suberosum, Linn. Sp. Pl. 212 = Gonolobus suberosus.
subulatum, *Vell. Fl. Flum.* 121; iii. t. 85.—Bras.
subvolubile, Schum. & Thonn. Beskr. Guin. Pl. 150 = Gymnema subvolubile.
tamiflolium, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 294 = Roulinia tamiflolia.
tenellum, Blanco, Fl. Filip. ed. I. 204 = Tylophora Perrottetiana.
tenellum, Linn. f. Suppl. 168 (Metastelmatis sp.).—N. Granat.
tenellum, Riedel, ex Fourn. in Mart. Fl. Bras. vi. iv. 250 = Ditasse anomala.
tetragonum, Vell. Fl. Flum. 119; iii. t. 69 = Gonolobus tetragonus.
tingens, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 24 = Marsdenia tinctoria.
tomentosum, Lam. Encyc. ii. 286 = Tylophora asthmatica?
tomentosum, Royle, ex Wight, Contrib. 58 = Roylei.
triforum, Vell. Fl. Flum. 119; iii. t. 70 = Gonolobus retusus.
triste, Griseb. Spicil. Fl. Rumel. ii. 68 = Vincetoxicum speciosum.
umbellatum, Vell. Fl. Flum. 120; iii. t. 78 (Sarcostemmatis sp.).—Bras.
undatum, Andr. Bot. Rep. t. 410 (Tylophorae sp.).—Ind. or.
undulatum, Jacq. Select. Am. 85 = Gonolobus undulatus.
usterion, *Vell. Fl. Flum.* 120; iii. t. 75.—Bras.
versicolor, Bunge, Enum. Pl. Chin. Bor. 44 = Vincetoxicum versicolor.
verticillare, Lam. Encyc. ii. 287 = filiforme.
villosum, Roem. & Schult. Syst. vi. 108 = Lachnostoma Balbisii.
vimivale, *Blanco, Fl. Filip.* ed. I. 203.—Ins. Philipp.
viminale, Linn. Mant. ii. 302 = Sarcostemma viminale.
Vincetoxicum, Pers. Syn. i. 274 = Vincetoxicum officinale.
virens, D. Dietr. Syn. Pl. ii. 906 = Schizoglossum virens.
virens, *Steud. Nom.* ed. II. i. 462; *D. Dietr. Syn. Pl.* ii. 905.—Afr. austr.
virginicum, Banks, ex Schult. Syst. vi. 68 = Gonolobus discolor.
viride, Vell. Fl. Flum. 121; iii. t. 81 = Gonolobus viridiflorus.
viridiflorum, Ker-Gawl. ex Spreng. Syst. i. 852 = Dregea volubilis.
viridiflorum, G. F. W. Mey. Prim. Fl. Esseq. 141 = Gonolobus viridiflorus.
viridiflorum, Sims, Bot. Mag. t. 1929 = Tylophora asthmatica.
vomitorium, Lam. Encyc. ii. 285 = Tylophora asthmatica.
vulgare, Pers. Syn. i. 274 = Vincetoxicum officinale.
Wallichii, *Wight, Contrib.* 56.—Reg. Himal.
Wilfordi, *Hook. f. Fl. Brit. Ind.* iv. 25, in nota.—China; Japon.

CYNAPIUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 640 (1840) = Ligusticum, Linn. (Umbell.).
apiifolium, Nutt. ex Torr. & Gray, l. c. 641 = L. apiifolium.
Bigelowii, Torr. in Facif. Rail. Rep. iv. 94 = Ferula dissoluta.
canadense, Nutt. ex Torr. & Gray, Fl. N. Am. i. 641 = L. apiifolium.

CYNAPIUM, Riv. ex Rupp. Fl. Jen. ed. Hall. 284 (1745) = **Aethusa**, Linn. (Umbell.).

CYNARA, Vaill. ex Linn. Gen. ed. i. 344 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 489.
BOURGAEA, Cos. Notes Crit. 39 (1849).
acaulis, Linn. Sp. Pl. ed. II. 1160 = Centaurea chamaerhapoticum.
acaulis, Sibth. & Sm. Fl. Graec. Prod. ii. 158 = Centaurea pygmaea!
alba, *Boiss. in DC. Prod.* vii. 304.—Hispan.
algarbiensis, *Cass. ex Nym. Consp.* 403.—Hispan.
Cardunculus, Willd. ex Steud. Nom. ed. II. i. 462 = pygmaea.
Cardunculus, *Linn. Sp. Pl.* 827.—Reg. Mediterr.
cornua, Stokes, Bot. Mat. Med. iv. 148 = Cardiua cernua.
colorata, Stokes, l. c. = Cnicus oleraceus.
corsica, Vir. Fl. Lib. Spec. 68 = C. Cardunculus.
diversifolia, Stokes, Bot. Mat. Med. iv. 146 = Cnicus heterophyllus.
eriocephala, Stokes, l. c. 145 = Cnicus eriophorus.
esculenta, Salisb. Prod. 185 = C. Cardunculus.
feros, Tenore, ex Steud. Nom. ed. II. i. 462 = *horrida* ?
glomerata, Thumb. Prod. Pl. Cap. 141 = Platycarpha glomerata.
horrida, Ait. Hort. Kew. ed. I. iii. 145 = C. Cardunculus
hortensis, Mill. Gard. Dict. ed. VIII. n. 2 = C. Cardunculus.
humilis, Juss. Gen. 178 = Centaurea chamaerhapoticum.
humilis, *Linn. Sp. Pl.* 828.—Europ. austr.
humilis, Sibth. & Sm. Fl. Graeca, ix. 25 = Sibthorpiana.
humilis, Vir. Fl. Cors. App. ii. 8 = C. Cardunculus.
Hystrix, *Ball, in Journ. Bot.* xi. (1873) 269.—Marocco.
integrifolia, *Vahl, Symb. Bot.* i. 68.—Hispan.
lamyroides, Webb, Iter Hisp. 84 = alba.
lanceata, Stokes, Bot. Mat. Med. iv. 156 = Cnicus lanceolatus.
palustris, Stokes, l. c. 154 = Cnicus palustris.
pusilla, Labill. Ic. Pl. Syr. Dec. iii. 11 = Centaurea pygmaea.
pygmaea, *Willd. Sp. Pl.* iii. 1692.—Hispan.
repens, Stokes, Bot. Mat. Med. iv. 152 = Cnicus arvensis.
Scolymus, Linn. Sp. Pl. 827 = C. Cardunculus.
Sibthorpiana, *Boiss. & Heldr. Diagn.* Ser. I. x. 94.—Cyprus.
spinosissima, J. & C. Presl, Delic. Prag. 109 = C. Cardunculus.
sylvestris, Lam. Encyc. i. 277 = C. Cardanculus.
syriaca, *Boiss. Diagn.* Ser. I. x. 94.—Syria.
Tournefortii, *Boiss. & Reut. Fl. Nov. Hisp. Diagn.* 18.—Hispan.
tuberosa, Stokes, Bot. Mat. Med. iv. 158 = Cnicus tuberosus.

CYNOCARDAMUM, Webb & Berth. Phyt. Canar. i. 96 (1836) = **Lepidium**, Tourn. (Crucif.).
Mexicanum, Endl. ex Heynh. Nom. ii. 180 = L. Menziesii.
virginicum, Webb & Berth. Phyt. Canar. i. 97 = L. virginicum.

CYNOCRAMBE, Tourn. ex Adans. Fam. ii. 497 (1763); Gaertn. Fruct. i. 362. t. 75 (1788) = **Theligonum**, Linn. (Urtic.).
prostrata, Gaertn. l. c. = T. Cynocrambe.

CYNOCTONUM, J. F. Gmel. Syst. 448 (1791) = **Mitreola**, Linn. (Loganiac.).
petiolatum, J. F. Gmel. l. c. = M. petiolata.
sessilifolium, J. F. Gmel. l. c. = M. sessilifolia.

CYNOCTONUM, E. Mey. Comm. Pl. Afr. Austr. 215 (1837) = **Vincetoxicum**, Moench (Asclep.).
acuminatum, Benth. in Hook. Niger Fl. 458.—Afr. trop.
alatum, Decne. in DC. Prod. viii. 529 = Cynanchum alatum.
angustifolium, Decne. l. c. = Cynanchum Callialata.
bifarium, Phil. in Linnaea, xxxiii. (1864–65) 176.—Chili.
Blumei, Decne. in DC. Prod. viii. 525 = Cynanchum ovalifolium.
boerhaaefolium, Decne. l. c. 580 = Cynanchum boerhaviifolium.
Boivinianum, Decne. l. c.—Madag.
Callialata, Decne. l. c. 528 = Cynanchum Callialata.

CYNOCTONUM:—
copense, E. Mey. Comm. Pl. Afr. Austr. 216 (= *Buxburia elliptica*).—Afr. austr.
coromandel, Teijsm. & Binn. Cat. Hort. Bog. 129, nomen.—Ins. Coron.
chilostoe, Decne. in DC. Prod. viii. 581.—Chili.
corymbosum, Decne. l. c. 528 = Cynanchum corymbosum.
crassifolium, E. Mey. Comm. Pl. Afr. Austr. 216.—Afr. austr.
dimidiatum, Hassk. in Flora, xl. (1857) 98.—Java.
Dombeyanum, Decne. in DC. Prod. viii. 529.—Ecuador.
Dregeanum, Decne. l. c. 531 = *crassifolium*.
erubescens, Decne. l. c. 520 = Cynanchum erubescens.
eurychilos, Decne. l. c. 538.—Madag.
fimbriatum, Decne. l. c. 531 = Melaplexis fimbriata.
floribundum, Decne. l. c. 529 = Cynanchum floribundum.
formosanum, Maxim. in Bull. Acad. Pétersb. xxiii. (1877) 370 (Cynanchi sp.).—Ins. Formosa.
Inculanum, Hance, in Journ. Bot. vi. (1868) 380 (Cynanchi sp.).—Ins. Hai-an.
lycioides, E. Mey. Comm. Pl. Afr. Austr. 217 = Tylophora lycioides.
Meyeri, Decne. in DC. Prod. viii. 581 = V. Meyeri.
microstemma, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 216.—Venezuela.
molle, E. Mey. Comm. Pl. Afr. Austr. 216 = Cynanchum molle.
mucronatum, Decne. in DC. Prod. viii. 531.—Chili.
muricatum, Decne. l. c. 528.—Java.
myrtifolium, Decne. l. c. 531 = Cynanchum myrtifolium.
neurocarum, Phil. in Linnaea, xxxiii. (1864–65) 176.—Chili.
nummularifolium, Decne. in DC. Prod. viii. 581 = Cynanchum nummularifolium.
ovalifolium, Decne. l. c. 529.—Madag.
ovalifolium, Decne. l. c. = Cynanchum ovalifolium.
pachyphyllum, Decne. l. c.—Chili.
patagonicum, Phil. in Linnaea, xxxiii. (1864–65) 175.—Patagon.
pauciflorum, Decne. in DC. Prod. viii. 528 = Cynanchum pauciflorum.
pedunculatum, Decne. l. c. 529 = Cynanchum pedunculatum.
pilosum, E. Mey. Comm. Pl. Afr. Austr. 216.—Afr. austr.
repandum, Decne. in DC. Prod. viii. 580.—Madag.
roseum, Decne. l. c. 532 = V. roseum.
scoparium, Chapm. Fl. S. U. St. 367 = V. scoparium.
undulatum, Decne. in DC. Prod. viii. 531.—Chili.
virens, E. Mey. Comm. Pl. Afr. Austr. 216 = Cynanchum virens.
viride, Phil. Fl. Atac. 35.—Chili.
Wallichii, Decne. in DC. Prod. viii. 528 = Cynanchum Wallichii.
Wilfordi, Maxim. in Bull. Acad. Pétersb. xxiii. (1877) 369 = Cynanchum Wilfordi.
Zollingeri, Miq. Fl. Ind. Bat. ii. 479.—Malaya.

CYNODON, Rich. in Pers. Syn. i. 85 (1805). *GRAMINEAE*, Benth. & Hook. f. iii. 1164.
CAPRIOLA, Adans. Fam. ii. 31 (1763).
DACTYLON, Vill. Hist. Pl. Dauph. ii. 69 (1787).
PERISTA, Koel. Gram. Gall. et Germ. 302 (1802).
altior, F. Muell. Fragm. viii. 118 = tenellus.
arcuatus, *J. & C. Presl, Rel. Haenk.* i. 290.—Ins. Luzon.
ciliaris, *Benth. Fl. Austral.* vii. 610.—Austral.
convergens, F. Muell. Fragm. viii. 118.—Austral.
Dactylon, *Pers. Syn. Pl.* i. 85.—Cosmop.
Dactylon, Steud. Syn. Pl. Gram. 219 = glabratus.
elongatus, *Trin. in Spreng. Neue Entdeck.* ii. 84.—Ind. or.
erectus, *J. & C. Presl, Rel. Haenk.* i. 290.—Mexic.; Peruv.
filiformis, Voigt, Hort. Suburb. Calc. 712 = elongatus.
glabratus, *Steud. Syn. Pl. Gram.* 218.—Arab.; Nubia.
gracilis, Nees, ex Steud. l. c. 218 = Leptochloa uniflora.
hoenpleatus, *Nees, in Linnaea*, vii. (1832) 261.—Afr. austr.
leptochloides, *Steud. Syn. Pl. Gram.* 452.—Ins. Neuilal.
linearis, Willd. Enum. Hort. Berol. 90 = C. Dactylon.

CYNODON:—
maritimus, H. B. & K. Nov. Gen. et Sp. i. 170 = C. Dactylon.
Neesii, Thw. Enum. Pl. Zeyl. 371 = Leptochloa Neesii.
notatus, *Nees, in Linnaea*, vii. (1832) 262.—Afr. austr.
occidentalis, Willd. ex Steud. Nom. ed. II. i. 468 = C. Dactylon.
pascuus, *Nees, Agrost. Bras.* 425.—Bras.
Phragmites, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 308 = Phragmites communis.
pilosissimus, Rasp. Ann. Obs. iii. 8; ex Prlts. Ic. Ind. 384 (quid?).
plicatus, Roem. & Schult. Syst. ii. 412 = Panicum sanguinale.
polystachyus, R. Br. Prod. 187 = Leptochloa polystachya.
portoricensis, Willd. ex Steud. Nom. ed. II. i. 468 = C. Dactylon.
prasocr, Roem. & Schult. Syst. ii. 412 = Panicum sanguinale.
radiatus, *Roth, Nov. Pl. Sp.* 88.—Ind. or.
repens, Dulac, Fl. Hautes-Pyr. 76 = C. Dactylon.
sarmentosus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 100 = C. Dactylon.
stellatus, Willd. Enum. Hort. Berol. 90 = C. Dactyloo.
tenellus, *R. Br. Prod.* 187.—Austral.
tener, *J. & C. Presl, Rel. Haenk.* i. 291.—Ins. Luzon.
tenuis, Trin. in Spreng. Neue Entdeck. ii. 83 = C. Dactylon.
ternatus, *A. Rich. Tent. Fl. Abyss.* ii. 405.—Abyss.
virgatus, *Nees, ex Steud. Syn. Pl. Gram.* 218.—Ind. or.
virgatus, Willd. Enum. Hort. Berol. 90 = Leptochloa chinensis.

CYNOGETON, Kunth, Enum. Pl. iii. 500 (1841) = CYCNOGETON, Endl. = **Triglochin**, Linn. (Nalad.).

CYNOGLOSSUM, [Tourn.] Linn. Syst. ed. I (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 848.
album, Gueldenst. ex Ledeb. Fl. Ross. iii. 166 = officinale.
amplexicaule, Lam. Tabl. Encyc. i. 399 = pictum.
amplexicaule, Michx. Fl. Bor. Am. i. 132 = virginianum.
amplifolium, *Hochst. ex DC. Prod.* x. 149.—Abyss.
anchusoides, Lindl. Bot. Reg. (1842) t. 14 = Paracaryum hellocarpum.
angustifolium, Auch. ex A. DC. Prod. x. 170 = Paracaryum leptophyllum.
angustifolium, Hort. ex Steud. Nom. ed. I. 251 = officinale.
angustifolium, Willd. Sp. Pl. i. 768 = Paracaryum angustifolium.
apenninum, Gouan, Fl. Monsp. 20 = pictum.
apenninum, Linn. Sp. Pl. 134 = Solenanthus apenninus.
apenninum, Roth, ex A. DC. Prod. x. 147 = montanum.
argenteum, Lam. Fl. Fr. ii. 277 = chetrifolium.
arundanum, Cos. Notes Pl. Crit. 41 = chetrifolium.
australe, *R. Br. Prod.* 496.—Austral.
Azoarti, *Phil. in Anal. Univ. Chil.* (1884), reimp. 8.—Chili.
baicalense, Pall. ex Roem. & Schult. Syst. iv. 764 = Craniospermum subvillosum.
Berteri, *Colla, ex DC. Prod.* x. 156.—Chili.
bicolor, Willd. Enum. Hort. Berol. 180 = officinale.
borbonicum, *Bory, Voy.* ii. 362.—Ins. Mascar.
brasiliense, Lehm. Leg. Gen. et Sp. Nov. 10 = Omphalodes amplexicaule.
californicum, A. Gray, Syn. Fl. N. Am. ii. 1. 476, index = Echinospermum californicum!
calycinum, C. A. Mey. Verz. Pfl. Cauc. 100 = Suchtelenia cauthocarpa.
calycinum, Wall. Cat. n. 998 = Wallichii.
canescens, Wall. l. c. n. 918, partim = denticulatum.
canescens, Willd. Enum. Hort. Berol. 180 = orientalium.
cappadocicum, Willd. Sp. Pl. i. 767 = Omphalodes cappadocica.
carolinianum, *Baker, in Journ. Linn. Soc. xx.* (1883) 211.—Madag.
chetrifolium, Lam. Coll. iii. 99 = pictum.
chinkiliense, *Linn. Sp. Pl.* 134.—Reg. Medituer. occ.

CYNOGLOSSUM:—
cheirifolium, Sieber, ex Nym. Consp. 522 = Columnae.
ciliatum, Dougl. ex Lehm. Pugill. ii. 24 = Echinospermum ciliatum.
clandestinum, *Desf. Fl. Atlant.* i. 168. t. 42.—Reg. Mediterr. occ.
clavatum, Viv. Fl. Lib. Spec. 11. t. 5. f. 2 = cheirifolium.
Clusii, Loisel. in Dict. Sc. Nat. xii. 384 = Echinospermum Lappula.
coelestinum, Lindl. Bot. Reg. (1839) t. 36 = Paracaryum caelestinum.
corsulense, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 100 = furcatum.
coeruleum, *Hochst. ex DC. Prod.* x. 148.—Abyss.
Columnae, *Bivona. Stirp. Rar. Sicil. Manip.* ii. [7].—Italia; Graecia; As. Min.
congestum, Poepp. ex DC. Prod. x. 139 = Eritrichium congestum.
creticum, Mill. Gard. Dict. ed. VIII. n. 8 = pictum.
cristatum, Auch. ex DC. Prod. x. 160 = Omphalodes rugulosa.
cristatum, Schreb. in Nov. Act. Nat. Cur. iii. (1767) 476; Lam. Tabl. Encyc. i. 400 = Paracaryum cristatum.
decumbens, *Gueldenst. ex Ledeb. Fl. Ross.* iii. 167.—Russia.
decurrens, *Ruiz & Pav. Fl. Per.* ii. 6.—Chili.
deflexum, Roth, Enum. Pl. Phaen. Germ. i. i. 589 = Echinospermum deflexum.
denticulatum, *A. DC. in DC. Prod.* x. 150.—Ind. or.; Malaya.
diffusum, Roxb. Hort. Beng. [13]; Fl. Ind. i. 4 = Bothriospermum tenellum.
Dioscoridis, Lehb. Fl. Ross. iii. 166 = nebrodense.
Dioscoridis, *Vill. Prosp.* 21; *Hist. Pl. Dauph.* ii. 457. —Reg. Mediterr.
discolor, *Baker, in Journ. Linn. Soc.* xx. (1883) 212. —Madag.
divaricatum, Steph. ex Lehm. Asperif. 161 = officinale.
Drummondii, *Benth. Fl. Austral.* iv. 409.—Austral.
echinatum, Thunb. Prod. Fl. Cap. 84 = Echinospermum cynoglossoides.
Edgeworthii, *A. DC. in DC. Prod.* x. 150.—Reg. Himal.
elongatum, Hornem. Hort Hafn. ii. 656 = Dioscoridis.
emarginatum, Lam. Tabl. Encyc. i. 400 = Paracaryum angustifolium.
Emodi, Schouw, Ind. Sem. Hafn. Coll. (1846) 4; ex Lehm. in Hamb. Gartenz. vi. (1850) 352 = Paracaryum heliocarpum.
enodum, Schouw, in Linnaea, xxiv. (1851) 160 = prae.
enerve, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 259.—Afr. austr.
erectum, *Schweigg. ex Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 29.—Hab.?
fontens, Gilib. Fl. Lituan. i. 22 = officinale.
fulvum, Rudolph, in Schrad. Journ. ii. (1799) 280 = clandestinum.
furcatus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 6.—Ind. or.
germanicum, Jacq. Obs. Bot. ii. 31 = montanum.
glaucinum, Willd. ex Roem. & Schult. Syst. iv. 761 = Solenanthus mollissimus.
glastifolium, Willd. Sp. Pl. i. 764 = Paracaryum glastifolium.
glochidiatum, Benth. in Royle, Illustr. Bot. Himal. 306 = denticulatum.
glochidiatum, *Wall. Cat.* n. 992.—Ind. or.; Burma.
glomeratum, Fraser, Cat. (1818) ex Pursh, Fl. Am. Sept. ii. 729 = Kryaitzkia glomerata.
grande, *Dougl. ex Lehm. Pugill.* ii. 23.—Am. bor. occ.
grandiflorum, Benth. in Royle, Illustr. Bot. Himal. 305 = Lindelofia spectabilis.
Haenkei, Schult. Oestr. Fl. ed. II. i. 863 = montanum.
Hayneï, G. Don, Gen. Syst. iv. 854 = furcatum.
hirsutissimum, *Lehm. Asperif.* 147 (err. typ. 145).—Afr.
hirsutum, Jacq. Hort. Schoenbr. iv. 45. t. 389 = lanceolatum.
hirsutum, *Thunb. Prod. Pl. Cap.* 84.—Afr. austr.
hispidum, Jacq. ex Roem. & Schult. Syst. iv. 78 = micranthum.
hispidum, Pursh, Fl. Am. Sept. ii. 729 = Kryaitzkia glomerata.
hispidum, *Thunb. Prod. Fl. Cap.* 84.—Afr. austr.
holosericeum, *Step. in Mém. Soc. Nat. Mosc.* iii. (1812) 268.—Reg. Caucas.

CYNOGLOSSUM:—
Howardi, A. Gray, Syn. Fl. N. Am. ii. i. 181 = Omphalodes Howardi.
humifusum, *Poepp. ex DC. Prod.* x. 138, in nota.—Chili.
hybridum, Thuill. Fl. Par. 94 = officinale.
intermedium, Fraser, in Mus. Sencksenb. i. (1834) 169 = Paracaryum micranthum.
japonicum, *Thunb. Fl. Jap.* 81.—Japon.
javanicum, Thunb. ex Lehm. Asperif. 118 = Echinospermum javanicum.
laeve, *A. Gray, Syn. Fl. N. Am.* ii. i. 188.—Calif.
laevigatum, Lehm. Asperif. 174 = Rindera pubescens.
laevigatum, Murr. Syst. ed. XIII. 157 = Rindera tetraspis.
lanatum, Auch. ex DC. Prod. x. 169 = Paracaryum Ancheri.
lanatum, Lam. Encyc. ii. 288 (= *Cyphomattus lanata*). —As. Min.; Persia.
lanceolatum, *Forsk. Fl. Aegypt. Arab.* 41.—Arabia.
lanceolatum, *Georgi, Beschr. Russ. Reich.* iii. iv. 754. —Siberia.
lanceolatum, Heyne, ex Wall. Cat. n. 921 = furcatum.
lanceolatum, *Hochst. ex DC. Prod.* x. 148.—Abyss.
lanifolium, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 207.—Afr. trop.
lanuginosum, *Lehm. Asperif.* 148.—Hab.?
Lappula, Scop. Fl. Carn. ed. II. i. 125 = Echinospermum Lappula.
lateriflorum, Aubr. Prog. Morb. x. 25 = Omphalodes littoralis.
lateriflorum, Bert. ex DC. Prod. x. 120 = Pectocarya linearis.
lateriflorum, Hort. ex Steud. Nom. ed. II. i. 464 (*verticillatum, spermum limifolium*).—Hab.?
lateriflorum, Lam. Encyc. ii. 289 = Pectocarya lateriflora.
latifolium, *R. Br. Prod.* 496.—Austral.
laxum, G. Don, Gen. Syst. iv. 856 = Paracaryum glochidiatum.
leptostachyum, *DC. Prod.* x. 152.—Afr. austr.
limense, Willd. Sp. Pl. i. 762 = fecurrens.
linearе, Ruiz & Pav. Fl. Per. ii. 6 = Pectocarya linearis.
lineatum, Russo, ex Ces. Pass. & Gib. Comp. Fl. Ital. 380 = cheirifolium.
linifolium, Bové, ex DC. Prod. x. 159 = Paracaryum micranthum.
linifolium, Linn. Sp. Pl. 134 = Omphalodes linifolia.
lithospermifolium, Lam. Encyc. ii. 240 = Omphalodes myosotoides.
litorale, Spreng. Syst. i. 567 = Omphalodes littoralis.
longiflorum, Benth. in Royle, Illustr. Bot. Himal. 305 = Lindelofia spectabilis.
longiflorum, Lehm. Asperif. 162 = Caccinia Russellii.
Lorcyi, Jord. ex Lange, in Kjoeb. Vidensk. Meddel. (1858) 28 = pictum.
lucidum, Stokes, Bot. Mat. Med. i. 277 = virginicum.
lusitanicum, Fortis, Oaserv. Sopra Cherso ed Osero, 68 = Omphalodes Fortisii.
lusitanicum, Lam. Encyc. ii. 289 = Omphalodes amplexicaulis.
lusitanicum, Mill. Gard. Dict. ed. VIII. n. 6 = Omphalodes linifolia.
macranthum, Regel & Smirnow, in Act. Hort. Petrop. v. (1877) 623 = macrostylum.
macrophyllum, Royle, ex DC. Prod. x. 136 = Paracaryum glochidiatum.
macropterum, Borsscz. ex Trautv. in Act. Hort. Petrop. ix. (1886) 49 = macrostylum?
macrostylum, *Bunge, in Lehm. Reliq. Bot.* (1847) 236. —Turkestan.
magellense, *Tenore, Fl. Nap.* iii. 183. t. 117.—Italia.
marchicum, Roxb. Hort. Beng. [13]; Fl. Ind. i. 457 = Heliocarpum marsilocum.
membranaceum, A. DC. in DC. Prod. x. 150 = Wallichii.
mexicanum, Cham. & Schlecht. in Linnaea, v. (1830) 114 = Echinospermum mexicanum.
micranthum, Dals. & Gibs. Bomb. Fl. 172 = lanceolatum.
micranthum, *Desf. Tabl.* ed. I. 220.—Reg. Himal.; Burma.
microcarpum, Kern. Nov. Pl. Sp. Dec. ii. 13 = Wallichii.
microglochin, *Benth. in Royle, Illustr. Bot. Himal.* 305.—Reg. Himal.
Monnchii, Necca & Roem. ex Steud. Nom. ed. II. i. 464 = Dioscoridis.

CYNOGLOSSUM:—
molle, *Phil. in Linnaea*, xxix. (1857-58) 18.—Chili.
mollissimum, Lehm. Asperif. 166 = Solenanthus mollissimus.
monophlebium, *Baker, in Journ. Linn. Soc.* xx. (1883) 211.—Madag.
montanum, *Linn. Amoen. Acad.* iii. 402; *Lam. Fl. Fr.* ii. 277.—Europ.; Oriens.
Morisoni, DC. Prod. x. 155 = Echinospermum virginicum.
muricatum, Thunb. Prod. Fl. Cap. 84 = Lithospermum flexuosum.
myosotoides, Auch. ex DC. Prod. x. 159 = Paracaryum hirsutum.
myosotoides, Labill. Ic. Pl. Syr. dec. ii. & t. 2 = Paracaryum myosotoides.
myosotoides, Schimp. ex DC. Prod. x. 157 = Omphalodes rugulosa.
nebrodense, *Guss. Fl. Sic. Prod.* i. 216.—Europ. austr.; Afr. bor.
nervosum, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 158.—Reg. Himal.
nitidum, Willd. Enum. Hort. Berol. 181 = Omphalodes amplexicaulis
noblle, *Hook. f. in Gard. Chron.* (1858) 240.—Ins. Chatham.
Nuttallii, Spreng. Syst. i. 565 = Pectocarya penicillata.
occidentale, *A. Gray, in Proc. Am. Acad.* x. (1874) 58. —Calif.
officinale, Brot. Fl. Lusit. i. 295 = clandestinum.
officinale, Desf. Fl. Atlant. i. 158 = pictum.
officinale, Hook. & Arn. Bot. Beech. Voy. i. 152 = grande.
officinale, *Linn. Sp. Pl.* 134.—Europ.; Oriens.
officinale, Willk. in Flora, xxxv. (1852) 217 = Dioscoridis.
Omphalodes, Linn. Syst. ed. X. 914 = seq.
omphalodes, Linn. Sp. Pl. 135 = Omphalodes verna.
ovatifolium, *Grisch. in Goett. Abh.* xxiv. "(1879) 271.—Reg. Argent.
ovatum, Moon, Cat. Pl. Ceyl. 12 = furcatum.
paniculatum, *Hook. & Arn. Bot. Beech. Voy.* 87.—Chili.
papillosum, *Thunb. Fl. Cap.* ii. 352.—Afr. austr.
parvefolium, C. Koch, in Linnaea, xxii. (1849) 645 = nebrodense.
pauciflorum, Bert. Merc. Chil. (Apr. 1829) 510 = Pectocarya pusilla.
pauciflorum, *Ruiz & Pav. Fl. Per.* ii. 6.—Chili.
pellucidum, Lapeyr. Suppl. Hist. Pl. Pyr. 28 = montanum.
penicillatum, Hook. & Arn. Bot. Beech. Voy. 371 = Pectocarya penicillata.
petiolatum, *A. DC. in DC. Prod.* x. 149.—Reg. Himal.
pictum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 179.—Reg. Mediterr.
pilosum, Nutt. Gen. Am. i. 114 = Echinospermum Redowskii.
pilosum, Ruiz & Pav. Fl. Per. ii. 6 = Pectocarya lateriflora.
platyphyllum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 384.—Afr. trop.
prostratum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 100 = Bothriospermum tenellum.
pustulatum, Boiss. Elench. 66 = nebrodense.
racemosum, Roxb. Hort. Beng. [13]; Fl. Ind. i. 456 = lanceolatum.
racemosum, Schreb. in Nov. Act. Nat. Cur. iii. (1767) 478 = Paracaryum angustifolium.
remotum, Moench, Meth. 419 = Dioscoridis.
revolutum, *Ruiz & Pav. Fl. Per.* ii. 6.—Peruv.
Rindera, Lam. f. Suppl. 130 = Rindera tetraspis.
Ritchiei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 157. —Ind. or.
robustum, *Hassk. in Ver. Nat. Ver. Nederl. Ind.* i. (1856) 44.—Malaya.
Rochelii, *A. DC. in DC. Prod.* x. 152.—Ins. Mascar.
Royleï, Wall. Cat. n. 917 = Paracaryum glochidiatum.
ruderale, Salisb. Prod. 116 = officinale.
rugosum, Sesiini, ex Roem. & Schult. Syst. iv. 764 = Moltkia coerulea.
scorpioides, Haenke, in Jacq. Coll. ii. 3 = Omphalodes scorpioides.
sessiliflorum, Poepp. ex DC. Prod. x. 153 = Eritrichium sessiliflorum.
siculum, Guss. Suppl. Fl. Sic. Prod. 52 = pictum.
sphaeroticum, *Boiss. & Heldr. in Boiss. Diagn.* Ser. I. xi. 125.—Ins. Creta.

CYNOGLOSSUM:—
staminosum, Bieb. Fl. Taur. Cauc. iii. 127 = Solenanthus Biebersteinii.
staminosum, Desf. in Ann. Mus. Par. x. (1807) 431 = Solenanthus Tournefortii.
stellulatum, Wall. Cat. n. 924 = glochidiatum.
stylosum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 409 = Solenanthus nigricans.
suaveolens, *R. Br. Prod.* 495.—Austral.
sylvaticum, Haenke, in Jacq. Coll. ii. 77 = montanum.
Tenorii, Guss. ex Nym. Consp. 521 = nebrodense.
tomentosum, *Lehm. Asperif.* 144.—Italia.
Trinanum, *Wedd. Chlor. And.* ii. 90.—N. Granat.
umbellatum, Waldst. & Kit. Pl. Rar. Hung. ii. 158 (= *Mattia umbellata*, Roem. & Schult.).—Hung.
uncinatum, Benth. in Royle, Illustr. Bot. Himal. 305 = Paracaryum glochidiatum.
valentinum, Lag. Gen. et Sp. Nov. 10 = Dioscoridis.
vernale, Salisb. Prod. 116 = Omphalodes verna.
vesiculosum, Wall. Cat. n. 920 = micranthum, Wallichii.
virens, Schreb. Spicil. Fl. Lips. [152] = montanum.
virginianum, *Linn. Sp. Pl.* 134.—Am. bor.
virginicum, J. F. Gmel. Syst. 319 = montanum.
virginicum, Linn. Syst. ed. XII. 146 = virginianum.
virginicum, Willd. Enum. Hort. Berol. 180 = Dioscoridis.
viridiflorum, *Willd. ex Roem. & Schult. Syst.* iv. 757. —Sibir.
vulgare, Gueldenst. ex Ledeb. Fl. Ross. iii. 165 = officinale.
Wallichii, *G. Don, Gen. Syst.* iv. 354.—Reg. Himal.
Xatartii, J. Gay, ex Nym. Consp. 521 = Dioscoridis
zeylanicum, Thunb. ex Lehm. Asperif. 116 = denticulatum.
zeylanicum, Wight, ex Wall. Cat. n. 7018 = Heliotropium paniculatum.

CYNOMETRA, Linn. in Act. Soc. Sc. Upsal. (1741) 78. *LEGUMINOSÆ*, Benth. & Hook. f. i. 586.
CYNOMORA, R. Hedw. Gen. 304 (1806).
IRIPA, Adans. Fam. ii. 508 (1768).
METROCYNIA, Thon. Gen. Nov. Madag. 22 (1809).
Afzelii, *Oliver, Fl. Trop. Afr.* ii. 318.—Afr. trop.
Aguilacha, Spreng. Syst. iii. 237 = Aquilaria Agallocha.
americana, *Vog. in Linnæa*, x. (1836) 602.—Ins. S. Doming.
bauhiniæfolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 99.—Guiana.
bijuga, Span. in Linnæa, xv. (1841) 201 = ramiflora.
cauliflora, *Linn. Sp. Pl.* 383.—Ind. or. ; Malaya.
cauliflora, Wall. Cat. n. 5816 = ramiflora.
Commersoniana, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 876.—Madag.
crassifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 100. —Ind. occ.
cubensis, *A. Rich. Ess. Fl. Cub.* 509.—Cuba.
cuneata, *Tul. in Arch. Mus. Par.* iv. (1844) 179.—Guiana.
falcata, *A. Gray, Bot. U. St. Expl. Exped.* i. 472.—Ins. Fiji.
floribunda, *Benth. in Trans. Linn. Soc. xxv.* (1866) 318.—Afr. trop.
grandiflora, *A. Gray, Bot. U. St. Expl. Exped.* i. 470. t. 69.—Ins. Pacif.
guianensis, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1085, nomen.—Guian.
Hostmanii, *Tul. in Arch. Mus. Par.* iv. (1844) 180.—Guian.
inæquifolia, *A. Gray, in Bot. U. St. Expl. Exped.* i. 473.—Ins. Philipp.
Iripa, *Kostel. Allg. Med.-Pharm. Fl.* iv. 1841.—Ind. or.
laxiflora, *Benth. in Trans. Linn. Soc. xxv.* (1866) 318. —Afr. trop.
madagascariensis, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 875.—Madag.
Mannii, *Oliver, Fl. Trop. Afr.* ii. 317.—Afr. trop.
marginata, *Benth. in Hook. Journ. Bot.* ii. (1840) 100. —Am. trop.
Martiana, Baill. Hist. Pl. ii. 164 = Hymenæa Martiana.
mimosoides, Wall. Cat. n. 5817 = ramiflora.
obtusa, *Benth. in Mart. Fl. Bras.* xv. ii. 248.—Bras.
parvifolia, *Tul. in Arch. Mus. Par.* iv. (1844) 181.—Am. trop.
Parvilliana, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 876.—Madag.
pinnata, *Lour. Fl. Cochinch.* 268.—Cochinch.

CYNOMETRA:—
polyandra, *Roxb. Hort. Beng.* 23; *Fl. Ind.* ii. 372.—Ind. or.
racemosa, *Benth. in Hook. Journ. Bot.* ii. (1840) 100. —Guian.
ramiflora, Miq. Anal. Bot. Ind. 11 = polyandra.
ramiflora, *Linn. Sp. Pl.* 382.—As. et Austral. trop.
Schomburgkiana, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1085, nomen.—Guian.
Spruceana, *Benth. in Mart. Fl. Bras.* xv. ii. 247.—Bras.
tetraphylla, Hook. f. in Hook. Niger Fl. 329 = Cryptosepalum tetraphyllum.
trapezoides, *Bedd. Fl. Sylv.* t. 216.—Ind. or.
Vogelii, *Hook. f. in Hook. Niger Fl.* 328.—Afr. trop.

CYNOMORA, R. Hedw. Gen. 304 (1806) = Cynometra, Linn. (Legumin.).

CYNOMORIUM, Opiz, Fl. Časlav. Cent iii. 255, 261 (1804).—Cf. Lotos, iv. (1854) 43 = Rantunculus, Tourn.
heldreichii, Opiz, ii. cc. = R. arvensis.

CYNOMORIUM, Mich. ex Linn. Syst. ed. I (1735). *BALANOPHOREÆ*, Benth. & Hook. f. iii. 324.
australe, Hook. f. in Trans. Linn. Soc. xxii. (1856) 46, sphalm. = Balanophora fungosa.
Balanophora, Willd. Sp. Pl. iv. 277 = Balanophora fungosa.
capensense, Sw. Prod. Veg. Ind. Occ. 12 = Helosis guianensis.
coccineum, Desconrt. Fl. Med. Antill. ii. 118. t. 96 = Scybalium jamaicense.
coccineum, *Linn. Sp. Pl.* 970.—Reg. Mediterr. ; Orient.
fungosum, Rasnsch. ex Steud. Nom. ed. I. 252 = Balanophora fungosa.
jamaicense, Sw. Prod. Veg. Ind. Occ. 12 = Scybalium jamaicense.
parasiticum, Sw. ex Steud. Nom. ed. I. 252 = Balanophora fungosa.
philippense, *Blanco, Fl. Filip.* ed. i. 665.—Ins. Philipp.
purpureum, Rupr. Sert. Tianseh. 72 = coccineum.
songaricum, *Rupr. l. c.* 73.—Soongaria.

CYNOPHALLA, J. Presl, Rostl. ii. 276 (1825) = Capparis, Tourn.
flexuosa, J. Presl, l. c. = Capp. Cynophallophora.
hastata, J. Presl, l. c. = Capp. Cynophallophora.
linearis, J. Presl, l. c. = Capp. linearis.
lucta, J. Presl, l. c. = Capp. verrucosa.
mollis, J. Presl, l. c. = Capp. mollis.
pascifiora, J. Presl, l. c. = Capp. Cynophallophora.
saligna, J. Presl, l. c. = Capp. Cynophallophora.
zeasita, J. Presl, l. c. = Capp. sessilis.
verrucosa, J. Presl, l. c. = Capp. verrucosa.

CYNOPOA, Ehrh. Phytophyl. n. 72 (1780); Beitr. iv. (1789) 148 = Agropyron, J. Gaertn. (Gramin.) (A. caninum).

CYNOPSOLE, Endl. Gen. 74 (1836) = Balanophora, Forst.
elongata, Endl. l. c. = B. elongata.

CYNOROHIS, Thon. Orch. Iles Afr. in tab. synopt. et t. 13 (1822). *ORCHIDEÆ*, Benth. & Hook. f. iii. 528.
AMPHORCHIS, Thon. ll. cc. tt. 4, 5 (1822).
CYNORCHIS, Thon. ll. cc. tt. 14, 15 (1822).
angustifolia, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 514.—Madag.
Boryana, *Lindl. Gen. et Sp. Orch.* 331.—Ins. Maurit.
brevicornis, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 516. —Madag.
calanthoides, *Kraenzlin, in Bremen Abh.* vii. (1882) 260.—Madag.
fastigiata, *Thou. Orch. Iles Afr.* t. 12.—Madag.
flexuosa, *Lindl. Gen. et Sp. Orch.* 331.—Madag.
gibbosa, Ridl. in Journ. Linn. Soc. xx. (1883) 581 = calanthoides.
grandiflora, *Ridl. l. c.* 332.—Madag.
klpidula, *Ridl. l. c.* xxi. (1885) 517.—Madag.
lilacina, *Ridl. l. c.* 515.—Madag.
purpurascens, *Lindl. Gen. et Sp. Orch.* 331.—Ins. Maurit. et Borbon.
squamosa, *Lindl. l. c.* 332.—Ins. Maurit. et Borbon.
trifphylla, Lindl. l. c. (= *Gymnadenia trifphylla*).—Ins. Maurit.
unifiora, *Lindl. l. c.* 331.—Madag.

CYNORHIZA, Eckl. & Zeyh. Enum. 351 (1837) = Penaeodannum, Tourn. (Umbell.).
alba, Eckl. & Zeyh. l. c. 352 = P. Zeyheri!
monticola, Eckl. & Zeyh. l. c. = P. Zeyheri.
sulcata, Eckl. & Zeyh. l. c. = P. sulcatum, *Sond. typica*, Eckl. & Zeyh. l. c. = P. Cynorhiza.

CYNORRHYNCHIUM, Mitchell, in Act. Phys. Med. Acad. Nat. Cur. viii. (1746) 207, App. = Mimulus, Linn. (Scrophal.).

CYNOSBATA, Reichb. Handb. 290 (1837) = Pelargonium, L'Hérit. (Geraniac.).

CYNOSCIADIUM, DC. Coll. Mém. v. 44, t. 11 (1829); Prod. iv. 140 (1830). *UMBELLIFERÆ*, Benth. & Hook. f. i. 906.
digitatum, *DC. Coll. Mém.* v. 44.—Am. bor.
pinnatum, *DC. l. c.*—Am. bor.

CYNOSORCHIS, Thou. Orch. Iles Afr. in tab. synopt. et tt. 14, 15 (1822) = Cynorchis, Thou. (Orchid.).
Boryana, Reichb. f. Otia Bot. Hamb. 72 = Cynorchis Boryana.
fastigiata, Lindl. Gen. et Sp. Orch. 332, in syn. = Cynorchis fastigiata.
flexuosa, Reichb. f. Otia Bot. Hamb. 72 = Cynorchis flexnosa.
galeata, Reichb. f. in Flora, lxviii. (1885) 536.—Ins. Comor.
purpurascens, Thou. Orch. Iles Afr. t. 15 = Cynorchis purpurascens.
squamosa, Reichb. f. Otia Bot. Hamb. 72 = Cynorchis squamosa.
trifphylla, Thou. Orch. Iles Afr. t. 14 = Cynorchis trifphylla.
unifiora, Reichb. f. Otia Bot. Hamb. 72 = Cynorchis unifiora.

CYNOSURUS, Linn. Gen. ed. I. 18 (1737). *GRAMINEÆ*, Benth. & Hook. f. iii. 1188.
FALCONA, Adans. Fam. ii. 496 (1763).
PHALONA, Dum. Agrost. Belg. 85, 114 (1823).
aegyptiacus, Link, Handb. i. 93, in syn. = acq. *aegyptius*, Linn. Sp. Pl. 72 = Eleusine aegyptiaca.
aristatus, *Bronss. Elench. Hort. Bot. Monsp.* 20, nomen —Hab. ?
aureus, Linn. Sp. Pl. 73 = Lamarckia aurea.
Balansae, *Coss. & Dur. in Bull. Soc. Bot. Fr.* i. (1854) 318.—Afr. bor.
bicolor, *Lond. in Transc. Camb. Phil. Soc.* vi. (1838), reimpr. 8.—Ins. Madera.
cerriuus, Linn. Sp. Pl. 72 = Sesleria caerulea.
callitrichus, *C. & W. Barb. Herbor. Levant*, 165. t. 10. —Syria.
capillaceus, Beauv. Agrost. 159 = Leptochloa tenerrima.
capillaris, Hort. ex Hornem. Hort. Hafn. i. 79 = Leptochloa tenerrima.
capitatus, Wulf. ex Roem. & Schult. Syst. ii. 604 = Sesleria tenella.
coeruleus, Willd. ex Steud. Nom. ed. II. i. 465 = Eleusine aegyptiaca!
Castagnei, Jord. ex Martr. Fl. Tarn, 814 = echinatus!
cerruleus, Auct. vide *caerulus*.
cerruleus, Turra, Nov. Sp. Fl. 11. 14 = Sesleria cylindrica.
coloratus, Lehm. ex Steud. Nom. ed. II. i. 465 = echinatus.
coracanus, Linn. Syst. ed. II. 875 = Eleusine coracana.
Crista-galli, Munby, Fl. Algér. 10 = multibractatus.
cristatus, *Linn. Sp. Pl.* 72.—Europ. ; Reg. Caucas.
cylindricus, Balb. Elench. 86 = Sesleria cylindrica.
distachyon, Rottl. ex Steud. Nom. ed. II. i. 465 = Eleusine aegyptiaca.
distichus, Host, Deutschl. Fl. ed. II. i. t. 49 = Orochloa disticha.
dominguensis, Jacq. Misc. ii. 368 = Leptochloa dominguensis.
durus, Forsk. Fl. Aegypt. Arab. 21 = Eragrostis cynosuroides.
durus, Linn. Sp. Pl. 73 = Sclerochloa dura.
echinatus, *Linn. l. c.*—Europ. ; Orient.
effusus, Link, in Schrad. Journ. ii. (1799) 315 = elegans.
elegans, *Desf. Fl. Atlant.* i. 98.—Reg. Mediterr.
erraceus, Jord. ex Martr. Fl. Tarn, 813 = echinatus!
erucaeformis, [Soland. in] Alt. Hort. Kew. ed. 2. i. 105 = Rostraria cristata.

CYNOSURUS :—
falcatus), Thunb. Prod. Pl. Cap. 28 (= *Campulosus falcatus*).—Afr. enstr.
fertilis, De Less. in Loisel. Fl. Gall. ed. II. i. 68 = echinatus.
filiformis, Vahl, Symb. Bot. ii. 90 = Leptochloa cynosuroides.
flaccifolius, Forsk. Fl. Aegypt. Arab. 21 = Eleusine flocciifolia.
geniculatus, Gueldenst. ex Ledeb. Fl. Ross. iv. 349 (quid ?).—Russia.
gigunteus, Nym. Consp. 998 = echinatus.
glaber, *Beauv. Agrost.* 159.—Hab. ?
gracilis, Cav. Hort. Reg. Matr. i. t. 5. f. 3 ; ex Roem. & Schult. Syst. ii. 806 = Aegopogon cenchroides.
gracilis, Vir. Fl. Lib. Supp. 67 ; Pl. Corr. 8 = elegans.
Hystrix, *Pomel, Nouv. Mat. Et. Ati.* 387.—Algeria.
indicus, Linn. Sp. Pl. 73 = Eleusine indica.
juncifolius, Wulf. ex Mert. & Koch, in Roehl. Deutschl. Fl. ed. III. i. 625 = Sesleria tenuifolia.
lagopoides, Burm. f. Fl. Ind. 29 = Aeluropus pubescens.
Lima, Linn. Sp. Pl. 73 = Wangenheimia disticha.
macrocephalus, Link, Handb. i. 84, sphalm.=seq.
microcephalus, Hoffm. Deutschl. Fl. ed. II. i. i. 49 = Sesleria tenella.
monostachyus, Vahl, Symb. Bot. ii. 20 = Leptochloa monostachya.
multibracteatus, *Roem. & Schult. Syst.* ii. 518.—Afr. bor.
nagicetus, Opiz, Natural. ix. (1825) 151 = cristatus.
obliquatus, Link, in Linnaea, xvii. (1843) 406 = elegans.
ovatus, Hoppe, in Sturm, Deutschl. Fl. Heft vi = Sesleria tenella.
paniceus, Linn. Sp. Pl. 73 = Polypogon monspeliensis.
paniculatus, Thunb. Prod. Pl. Cap. 28 = Poa Uniolae.
paspaloides, Vahl, Symb. Bot. ii. 21 = Chloris petraea.
pectinatus, Lam. Encyc. ii. 188 = Leptochloa pectinata.
penicillatus, Vahl, Symb. Bot. ii. 21 = Chloris penicillata.
phleoides, Desf. Fl. Atlant. i. 82 = Koeleria hispida.
polybracteatus, Gren. & Godr. Fl. Fr. iii. 568 = elegans.
polybracteatus, Poir. Voy. Barb. ii. 97 = cristatus.
polydactylon, Nocca, Cat. Pav. (1813) ; ex Roem. & Schult. Syst. ii. 819 (Quid ?).
retroflexus, Vahl, Symb. Bot. ii. 20 (= *Leptochloa arabica*).—Hab. ?
rupestris, Wulf. ex Stend. Nom. ed. II. i. 465 = Sesleria coerulea.
scoparius, Lam. Encyc. ii. 188 = Chloris radiata.
secundus, Pursh, Fl. Am. Sept. ii. 736 = Bouteloua racemosa.
Seslerie, Schrank, Baier. Fl. i. 350 = Sesleria coerulea.
siculus, Jacq. Obs. Bot. ii. 22 = Desmazeria sicula.
sphaerocephalus, Wulf. in Jacq. Misc. ii. 71 = Sesleria sphaerocephala.
splendens, Tenore, ex Roem. & Schult. Syst. ii. 870 = Apera interrupta.
tenellus, Cav. Hort. Reg. Matr. i. t. 5. f. 2 ; ex Roem. & Schult. Syst. ii. 805 = Aegopogon geminiflorus.
tenurivius, Horwen. Enum. Hort. Hafn. 8 = Leptochloa filiformis.
ternatus, Forsk. Fl. Aegypt. Arab. 21.—Arab.
tristachya, Lam. Encyc. ii. 188 = Eleusine tristachya.
uniflorus, Walt. Fl. Carol. 83 (Quid ?).—Carolina.
Uniolae, Linn. f. Suppl. 110 = Poa Uniolae.
virgatus, Linn. Syst. ed. X. 876 = Leptochloa virgata.

CYNOTIS, Hoffmgg. Verz. Pfl. Nachtr. ii. (1826) 98 = Cryptostemma, R. Br. (Compos.).
hypochondriaca, Hoffmgg. i. c. = Crypt. hypochondriacum.

CYNOTOXICUM, Vell. Fl. Flum. 197 (1825) ; iv. t. 158–159 = Cemmarus ! Linn.
dispermum, Vell. l. c. 198 ; iv. t. 158.—Bras.
monospermum, Vell. l. c. ; iv. t. 157.—Bras.
rufibullatum, Vell. l. c. ; iv. t. 158.—Bras.

CYNOXYLON, Raffin. Alsog. Am. 59 (1836) = Cornus, Tourn.
florida, Raffin. l. c. = Cornus florida.

CYNTHIA, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 308 = Krigia, Schreb. (Compos.).
amplexicaulis, Beck, Bot. N. et Mid. St. 183 = K. amplexicaulis.

CYNTHIA :—
Bocci, DC. Prod. vii. 89 = Krigia Dandelion.
Dandelion, DC. l. c. = K. Dandelion.
Griffithii, Nutt. in Journ. Acad. Philad. vii. (1834). 69 = K. amplexicaulis.
lyrata, Nutt. l. c. 68 = K. Dandelion.
virginica, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 309 = K. amplexicaulis.

CYPELLA, Herb. in Bot. Mag. sub t. 2637 (1826).
IRIDEAE, Benth. & Hook. f. iii. 689.
HERBERTIA, Lindl. in Bot. Reg. sub t. 1031 (1827) ; Herb. *Amaryll.* 179.—Bras.
PHALOCALLIS, Herb. in Bot. Mag. t. 3710 (1839).
POLIA, Tenore, Cat. Ort. Bot. Nap. (1845) 92.
cathartica, Mart. ex Klatt, in Mart. Fl. Bras. iii. i. 526 (= *Lansbergia cathartica*).—Bras.
culciviformis, Sesb. ex Klatt, l. c. 521 = gracilis.
Drummondii, R. Grah. in Edinb. N. Phil. Journ. xx. (1836) 190 = Herbertia Drummondiana.
gigantea, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 864.—Bras.
gracilis, *Baker, in Journ. Linn. Soc.* xvi. (1877) 127.—Bras.
Herbertii, *Herb. Bot. Mag.* sub t. 2637.—Bras.
monticola, Mart. ex Klatt, in Mart. Fl. Bras. iii. i. 526 (= *Lansbergia monticola*).—Bras.
paludosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 866.—Bras.
peruviana, *Baker, in Bot. Mag.* t. 6213.—Peruv.
plumbea, *Lindl. Bot. Reg.* (1838) Misc. 71.—Bras.
purpurea, Mart. ex Klatt, in Mart. Fl. Bras. iii. i. 527 (= *Lansbergia purpurea*).—Bras.
pusilla, *Benth. & Hook. f. Gen.* iii. 690.—Bras.
ramosa, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 864 = Herberti.
violacea, Sesb. ex Klatt, in Mart. Fl. Bras. iii. i. 526 (= *Lansbergia violacea*).—Bras.

CYPELLA, Klatt, in Linnaea, xxxi. (1861–62) 538 = Marica, Ker-Gawl. (Irid.).
brachypus, Baker, in Gard. Chron. (1876) i. 188 = M. brachypus.
caerulea, Sesb. ex Hook. f. Bot. Mag. t. 5612 = M. caerulea.
glauca, Sesb. ex Klatt, in Linnaea, xxxi. (1861–62) 543 = M. glauca.
gracilis, Klatt, l. c. 541 = M. gracilis.
humilis, Klatt, l. c. 540 = M. humilis.
longifolia, Klatt, l. c. 541 = M. longifolia.
lutea, Klatt, in Mart. Fl. Bras. iii. i. 522 = M. lutea.
Northiana, Klatt, in Linnaea, xxxi. (1861–62) 539 = M. Northiana.

CYPELLIUM, Desv. in Hamb. Prod. Ind. Occ. 28 (1825) ! Gen. haud recogn. Benth. & Hook. f. ii. 666.
nervosum, Desv. l. c.—Guiana.

CYPERORCHIS, Blume, Rumphia, iv. 47 (1848) in obs. ; et Mus. Bot. Lugd. Bat. i. 48 (1849). *ORCHIDEAE*, Benth. & Hook. f. iii. 588.
cochleata, *Benth. in Journ. Linn. Soc.* xviii. (1881) 318.—Reg. Himal.
elegans, *Blume, Rumphia*, iv. 47.—Reg. Himal.
Mastersii, *Benth. in Journ. Soc.* xviii. (1881) 318.—Reg. Himal.

CYPERUS, [Mich.] Linn. Syst. ed. I (1735) ; Gen. ed. I. 13 (1737). *CYPERACEAE*, Benth. & Hook. f. iii. 1043.
ANOSPORUM, Nees, in Linnaea, ix. (1834) 289.
ATOMOSTYLIS, Steud. in Flora, xxxiii. (1850) 229 ; Syn. Pl. Cyp. 315 (1855).
BOHEMIA, Linn. Fl. Zeyl. 17 (1747).
BORABORA, Steud. Syn. Pl. Cyp. 71 (1855).
GALILEA, Parl. Fl. Palerm. i. 307 (1845).
HYDROSCHOENUS, Zoll. & Mor. Syst. Verz. Zoll. 95 (1846).
OPETIOLA, Gaertn. Fruct. i. 14. t. 2 (1788).
PAPYRUS, Willd. in Abh. Acad. Berl. (1812–13) 70.
PYRGOSTERNUS, Opiz, Seznam, 90 (1852).
SOROSTACHYS, Steud. in Flora, xxxiii. (1850) 229 ; Syn. Pl. Cyp. 71 (1855).
TORRENTA, Raffin. in journ. Phys. lxxxix. (1819) 105.
TORULINIUM, Desv. in Hamb. Prod. Pl. Ind. Occ. 15 (1825).
TRENTEPOHLIA, Steud. in Bot. Zeit. xvi. (1858) 849.

CYPERUS :—
abnormis, *Steud. Syn. Pl. Cyp.* 48.—Hab. ?
abyssinicus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 474 = Pycreus flavescens.
acaulis, Steud. in Flora, xxv. (1842) 599 = Mariscus cazanensis.
aciculatis, Steud. Syn. Pl. Cyp. 45 (Marisci sp.).—Bras.
aciculatis, With. Bot. Arr. ed. III. ii. 78 = Elescharis aciculatis.
actinostachys, *Welw. ex Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 140.—Afr. trop.
acuminatus, Roxb. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 77 = lucrigatus.
acuminatus, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 435.—Am. bor.
acutangulus, *Boeck. in Linnaea*, xxxv. (1867–68) 551. —Chili.
acutiflorus, Steud. Syn. Pl. Cyp. 33 = protelinolepis.
acutiusculus, *Lag. Gen. et Sp. Nov.* 3.—Mexic.
acutus, R. Br. Prod. 217 (Marisci sp.).—Austral.
adenophorus, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. i. 28.—Bras.
adoensis, Hochst. in Flora, xxiv. (1841) i. Intell. 20 = rigidifolius.
adscendens, *Steud. Syn. Pl. Cyp.* 14.—Hab. ?
aduncus, Nees, ex Boeck. in Linnaea, xxxv. (1867–68) 461 = *Aschenbornianus*.
adustus, J. & C. Presl, Rel. Haenk. i. 167 = melanostachyus.
aegyptiacus, *Clusius. Obs.* 20. t. 3.—Europ.
aequalis, *Vahl, Enum.* ii. 320.—Afr. austr. ; Ins. Madag.
aethiops, Welw. ex Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 129 (Pycrei sp.).—Afr. trop.
affinis, Roem. & Schult. Syst. ii. 213 = comosus.
Afzelii, Boeck. in Linnaea, xxxv. (1867–68) 475 (Pycrei sp.).—Afr. trop.
agrestis, Willd. ex Spreng. & Link, Jahrb. i. 111. 86 = rotundus.
Altoni, *Spreng. Mant.* i. 29.—Hab. ?
alatus, F. Muell. Fragm. viii. 272, in obs. = Kyllinga alba.
albiceps, Ridl. in Journ. Bot. xxii. (1884) 16 = Kyllinga sphaeroceph.
albidus, Lam. Tabl. Encyc. i. 145 = Pycreus sanguinolentus.
albidus, *Retz. Obs.* vi. 21.—China.
albo-marginatus, Mart. & Schrad. ex Nees, in Mart. Fl. Bras. ii. i. c. 9 = Pycreus albo-marginatus.
albostriatus, *Schrad. Anal. Cap. Cyp.* 7.—Afr. austr.
albus, *J. & C. Presl, Rel. Haenk.* i. 175.—Ins. Linon.
alopecuroides, [Koen.] ; Roxb. Hort. Beng. 9 ; Fl. Ind. i. 208 = exaltatus.
alopecuroides, *Rottb. Descr. Nov. Pl.* 88. t. 8 fig. 2 ; *Prog.* 30.—Geront. trop.
alopecuroides, Thunb. Prod. Pl. Cap. 18 = Mariscus riparius.
altivus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 215 (Marisci sp.).—Mexic.
alterniflorus, R. Br. Prod. 216 (Marisci sp.).—Austral.
alterniflorus, Schwein. in Long, Exped. Winnep. ii. 361 = Schweinitzii.
alternifolius, *Linn. Mant.* 28.—Madag.
alternifolius, Steud. Syn. Pl. Cyp. 54 (Marisci sp.).—Austral.
alternifolius, Willd. ex Kunth, Enum. Pl. ii. 80 = longifolius.
altus, Nees, in Wight, Contrib. 84 = exaltatus.
amabilis, *Vahl, Enum.* ii. 318.—Geront. trop.
amauropus, Steud. Syn. Pl. Cyp. 33 (Marisci sp.).—Abyss.
ambiguus, Liebm. Vidensk. Selsk. Skr. V. ii. (1851) 199 = Pycreus propinquus.
ambiguus, Steud. Syn. Pl. Cyp. 18 = Scirpus cernuus.
ambiplexipus, *Steud. l. c.* 30.—Abyss.
ambongensis, Boeck. in Linnaea, xxxvIII. (1874) 261 = obtusiflorus.
amoenulaceus, Hadge, Pl. Gui. 16. t. 19 = Fimbristylis complanata.
amoicola, *Kunth, Enum. Pl.* ii. 82.—Afr. austr.
amoenus, Koen. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 187 = exaltatus.
amoenus, Kunth, Enum. Pl. ii. 86 = longus.
amoyensis, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 249 = procerus.
amphibet, Steud. Syn. Pl. Cyp. 56 = ancepa.
amphibolus, Steud. l. c. 23 = denudatus.
amplissimus, Steud. l. c. 216 = prolixus.

CYPERUS :—
amuricus, *Maxim. Prim. Fl. Amur.* 296.—Reg. Amur.
anabaptistus, *Steud. Syn. Pl. Cyp.* 37.—Ins. Luzon.
ancops, Heyne, ex Steud. Nom. ed. II. i. 466 = *proceruus*.
ancops, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 213.—Cuba.
ancops, Steud. Syn. Pl. Cyp. 23 = Fimbristylis complanata.
Andersoniensis, Boeck. in Engl. Bot. Jahrb. v. (1884) 502 = Mariscus cyperinus.
Andersonii, Boeck. in Linnæa, xxxvi. (1869–70) 388 = Mariscus brachystachys.
andongensis, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 140.—Afr. trop.
Andachæ, A. Rich. Tent. Fl. Abyss. ii. 481 = dichroostachyus.
angolensis, *Boeck. in Flora*, lxiii. (1880) 485.—Afr. trop.
angulatus, Nees, in Wight, Contrib. 73 = Pycreus angulatus.
angulatus, W. Wats. in E. T. Atkinson, Gazetteer, x. (1876) 393 (Pycrei sp.) = *pauper*.
angustatus, Benth. Fl. Austral. vii. 282 (Mariscí sp.) = Luerssenii, *Cuvenii*.
angustatus, R. Br. Prod. 214 (Mariscí sp.).—Austral
angustatus, Thunb. ex Spreng. & Link, Jahrb. i. iii. 84 = leucocaulis.
angustatus, Willd. ex Kunth, Enum. Pl. ii. 88 = difformis.
angustifolius, Buch.-Ham. in Wall. Cat. n. 8876 B = cuspidatus.
angustifolius, A. Dietr. Sp. Pl. ii. 322 (Pycrei sp.) = *rivularis*, Kunth.
angustifolius, *Schum. & Thonn. Beskr. Guin. Pl.* 88.—Afr. trop.
anisos, Roem. & Schult. Mant. ii. 181 (Mariscí sp.) = *Maximæ*.
anisostachyos, Willd. ex Kunth, Enum. Pl. ii. 20 = amabilis.
annuus, Spreng. & Link, Jahrb. i. iii. 88 = Mariscus glomeratus.
annuus, Willd. ex Kunth, Enum. Pl. ii. 338 = Fimbristylis diphylla.
anomalus, Steud. Syn. Pl. Cyp. 37 = Mariscus albescens.
apaturis, *Steud. l. c.* 47.—Hab.?
aphyllus, Boeck. in Linnæa, xxxv. (1867–68) 450 (Pycrei sp.) = *latispicatus*.
aphyllus, Hassk. Cat. Hort. Bogor. Alt. 24 = Kyllinga pungens.
aphyllus, F. Muell. Fragm. viii. 272, in obs. = Kyllinga peruviana.
aphyllus, Vahl, Eclog. Am. ii. 6 = C. Haspan.
apiculatus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 230 (Mariscí sp.) = *Maximæ*.
appendiculatus, Kunth, Enum. Pl. ii. 81 = Mariscus appendiculatus.
aprinus, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 141.—Afr. trop.
aquatilis, R. Br. Prod. 218 = trinervis.
aquatilis, F. Muell. Fragm. viii. 270 = Pycreus polystachyus.
arabicus, Ehrenb. ex Boeck. in Linnæa, xxxvi. (1869–70) 284 = tuberosus.
arcuatus, Boeck. l. c. xxxv. (1867–68) 542 = conglomeratus.
arenarius, Hance, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 178 = stoloniferus.
arenarius, *Retz. Obs.* iv. 9.—As trop.
arenarius, Salzm. ex Steud. Syn. Pl. Cyp. 46 (Mariscí sp.).—Bras.
arenarius, Wight, in Wall. Cat. n. 8374 = aristatus.
arenicola, *Steud. Syn. Pl. Cyp.* 51.—Texas.
areolatus, R. Br. Prod. 216 (Pycrei sp.) = *C. Eragrostis*.
argenteus, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 153.—Afr. trop.
argentinus, Boeck. in Engl. Bot. Jahrb. v. (1884) 502 = Mariscus flavus.
argentinus, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 64 (Pycrei sp.).—Reg. Argent.
argyraeus, *Steud. Syn. Pl. Cyp.* 11.—Afr. trop.
aristatus, Hook. f. & Thoms. partim, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 90 = cuspidatus.
aristatus, *Retth. Descr. Nov. Pl.* 23; *Prog.* 16.—Orbis terrarum.
aristatus, Boeck. in Linnæa, xxxv. (1867–68) 501, partim = Scirpus hamulosus.
Armstrongii, Benth. Fl. Austral. vii. 289 (Mariscí sp.). —Austral.

CYPERUS :—
articulatus, Benth. Fl. Hartw. 194 = nodosus.
articulatus, *Linn. Sp. Pl.* 44.—Cosmop. trop.
articulatus, Schlecht. in Bot. Zeit. vii. (1849) 96, partim = corymbosus.
arundinaceus, Forsk. Fl. Aegypt. Arab. p. cli. n. 45 = conglomeratus.
arundinaceus, Linn. Sp. Pl. 44 = Dulichium spathaceum.
arundinaceus, *Neronha, ex Thou. Prod. Phyt.* 2, in *Milang.* (1811) nomen.—Madag.
Aschersonianus, Boeck. in Linnæa, xxxv. (1867–68) 460 (Pycrei sp.).—Mexic.
asperifolius, Desf. Cat. Hort. Par. ed. III. 387 = textilis.
asperrimus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 218 (Mariscí sp.) = *Manimæ*.
assimilis, Steud. in Flora, xxv. (1842) 584 (Cortoisiæ sp.).—Abyss.
ater, Dalz. & Gibs. Bomb. Fl. 283 = Pycreus capillaris.
ater, Vahl, Enum. ii. 338 (Pycrei sp.) = *C. Eragrostis*.
aterrimus, *Hochst. ex Steud. Syn. Pl. Cyp.* 51.—Abyss.
atrofuscus, Steud. l. c. 2 = Pycreus capillaris.
atroruncatus, Boeck. in Linnæa, xxxvii. (1874) 856 (Pycrei sp.).—Abyss.
atronitens, Hochst. in Flora, xxiv. (1841) i. Intell. 20 (Pycrei sp.).—Afr. trop.
atropurpureus, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 497.—Madag.
atropurpureus, Boj. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1883) 231 (Pycrei sp.) = *Boivini*.
atropurpureus, Pers. Syn. i. 60 = striatus.
atroscapineus, Hemsl. Biol. Centr. Am. Bot. iii. 445 = Mariscus atroscapineus.
atroscapineus, Hochst. ex Steud. Syn. Pl. Cyp. 30 = rigidifolius.
Aucheri, *Jaub. & Spach, Illustr.* ii. t. t. 101.—Persia ; Algeria.
aurantiacus, H. B. & K. Nov. Gen. et Sp. i. 205 = amabilis.
aureolus, *Steud. Syn. Pl. Cyp.* 28.—Guatamala.
aureorufus, Boeck. in Linnæa, xxxvii. (1874) 869 = auricomus.
aureus, Georgi, Beschr. Russ. Reich. Nachtr. 257 = glomeratus.
aureus, H. B. & K. Nov. Gen. et Sp. i. 206 = amabilis.
aureus, J. & C. Presl, Rel. Haenk. i. 168 = squarrosus.
aureus, Tenore, Fl. Nap. iii. 45. t. 101 = esculentus.
auricomus, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 188, pro majore parte = digitatus.
auricomus, *Sieber, ex Spreng. Syst.* i. 230.—Geront. trop.
auriculatus, Nees & Meyen, in Linnæa, ix. (1834) 285 (Mariscí sp.).—Ins. Sandvic.
aurantis, Schrad. Fl. Germ. i. 116 = glomeratus.
autumnalis, Pursh, Fl. Am. Sept. i. 57 = articulatus.
autumnalis, Vahl, Enum. ii. 318 = C. Haspan.
Banaea, Steud. Syn. Pl. Cyp. 216 = reflexus.
Balsaba, *Steud. l. c.* 6.—As. trop.
babakensis, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 152 = prasc.
Bachus, Nees, in Linnæa, ix. (1834) 288 (Pycrei sp.) = *punctiniculatus*.
badius, Boeck. in Linnæa, xxxvi. (1869–70) 276 = Mariscus paniceus.
badius, Desf. Fl. Atlant. i. 45. t. 7 = longus.
badius, Steud. in Flora, xli. (1829) 158 = congestus.
Baeuitzii, *Boeck. in Flora*, lxi. (1878) 140.—Uruguay.
bahiensis, Steud. Syn. Pl. Cyp. 88 = esculentus.
Bakeri, *C. B. Clarke, in Journ. Linn. Soc.* xx. (1883) 290.—Ins. Mascrit.
Baldiniî, Kunth, Enum. Pl. ii. 68 = sphæroletus.
Baldwinii, Boeck. in Linnæa, xxxvi. (1869–70), partim = Mariscus uniformis.
Baldwinii, Chapm. Fl. S. U. St. 511, partim = Mariscus cylindricus.
Baldwinii, Torr. in Ann. Lyc. N. York, iii. (1836) 270 = Mariscus echinatus.
Balfourî, *C. B. Clarke, in Journ. Linn. Soc.* xx. (1883) 289.—Ins. Borbon.
banetious, Kit. ex Nym. Consp. 799 = glaber.

CYPERUS :—
banosanus, *Mig. Fl. Ind. Bat. Suppl.* 899.—Ins. Banca.
barbatus, Poir. Encyc. Suppl. v. 186 = Fimbristylis filamentosa.
Baroni, *C. B. Clarke, in Journ. Linn. Soc.* xx. (1883) 289.—Madag.
Bartoni, Boeck. in Linnæa, xxxv. (1867–68) 460 (Pycrei sp. ?) = pastalatus.
bellus, *Kunth, Enum. Pl.* ii. 58.—Afr. austr.
benguinsis, Spreng. Neue Entdeck. iii. 101 = tegetiformis.
benghalensis, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 151 = C. Babakan.
Bertieri, Kunth, Enum. Pl. ii. 67 = digitatus.
Beyrichii, Steud. Syn. Pl. Cyp. 47 (Mariscí sp.).—Bras.
bicaps, Vahl, Enum. ii. 368 = radiatus.
bicolor, Barton, Comp. Fl. Philad. i. 37 = diandrus.
bicolor, Vahl, Enum. ii. 340 = rotundus.
bidentatus, Poir. Encyc. vii. 350 = alopecuroides.
bidentatus, *Vahl, Enum.* ii. 330.—Afr. trop.
bifidus, *Fourr. Chlor. Hisp.* 174.—Mont. Pyren.
Bigelowii, Steud. Syn. Pl. Cyp. 5 (Pycrei sp.) = *virularis*.
biglumis, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 199 = Mariscus Sieberianus.
bipartitus, Torr. in Ann. Lyc. N. York, iii. (1836) 257 (Pycrei sp.) = *diandrus*, Torr.
Bipontii, *Boeck. in Flora*, xl. (1857) 38.—Mexic.
bispicatus, Steud. Syn. Pl. Cyp. 49 = elatus.
bisumbellatus, Steud. l. c. 46 = prolixus.
bisyncladus, Anders. in Vet. Akad. Handl. Stockh. 1858 (1855) 156 = confertus.
Blanchanus, Desv. ex Boiss. Fl. Orient. v. 371 = glaber.
blandus, *Kunth, Enum. Pl.* ii. 86.—Afr. austr.
blepharoleptos, Steud. Syn. Pl. Cyp. 28 = Scirpus cubensis.
blymoides, *Hochst. in Flora*, xxvii. (1844) 102.—Abyss.
Boehmii, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 498.—Afr. trop.
Boivini, Boeck. in Linnæa, xxxv. (1867–68) 481 (Pycrei sp.).—Madag. ; Ins. Borbon.
borbonicus, Steud. Syn. Pl. Cyp. 31 = articulatus.
Boschianus, *Regel, in Act. Hort. Petrop.* ii. (1873) 316 (cult.).—Hab.!
Bowmanni, F. Muell. ex Benth. Fl. Austral. vii. 287 (Mariscí sp.).—Austral.
brachistus, Poir. Encyc. vii. 350 = compressus.
brachyphyllus, Sp. Link, Jahrb. i. iii. 89 = amabilis.
brachystachys, Andersss. Galapag. Cuenes Veg. 68. t. 18. f. 2 = Mariscus brachystachys.
brachystachys, Nees, ex Boeck. in Linnæa, xxxvi. (1869–70) 210 = digitatus.
brachystachys, Presl, Cyp. et Gram. Sic. 15 = longus.
bractiatus, Hort. Bonn. ex Steud. Syn. Pl. Cyp. 19 = Zollingeri.
bracteiflorus, Steud. l. c. 38 = Mariscus flavus.
bracteolatus, Salzm. ex Steud. l. c. = Mariscus flavus.
bracteolatus, Steud. l. c. 49 (Mariscí sp.) = *ferax*.
bracteosus, *Kunze, ex C. Gay, Fl. Chil.* vi. 166.—Chili.
Brandi, *Vathe, Ind. Sem. Hort. Berol. App.* (1872) 5. —Hab.!
Brauni, Boeck. in Linnæa, xxxv. (1867–68) 559, in cit. = reflexus.
brevicaulis, R. Br. Prod. 214 = trinervis.
brevicaulis, F. Muell. Fragm. viii. 267 = pumilus.
brevifavus, A. Dietr. Sp. Pl. ii. 316 = miserrimus.
brevifolius, Hassk. Cat. Hort. Bogor. Alt. 24 = Kyllinga brevifolia.
brevirachiatus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 214 = Mariscus fusbelliformis.
brevis, *Boeck. in Linnæa*, xxxvi. (1869–70) 241.—Alt. austr.
brevispicatus, Boeck. in Flora, xlii. (1859) 442 (Mariscí sp.).—Hab.!
brizaeocephus, *Steud. Syn. Pl. Cyp.* 5.—Bras.
brizaeus, Forsk. Fl. Am. Sept. i. 51 = Pycreus polystachyus.
brizaeus, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 106.—Ind. occ.
brizaeus, Salzm. ex Steud. Syn. Pl. Cyp. 5 = brizaeocephus.
brizaeus, Vahl, Enum. ii. 336 (Mariscí sp.).—Ind. occ. ; Florida.
bromoides, Lisk, Jahrb. i. iii. 85 = Pycreus angulatus.

CYPERUS :—
Brownii, Roem. & Schult. Syst. ii. 226 = aristatus.
brunneo-ater, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 498. —Madag.
brunneus, Griseb. Fl. Brit. W. Ind. 565 (Mariscí sp.) = *purpurascens*.
brunneus, Sw. Fl. Ind. Occ. i. 116 (Mariscí sp.).—Ind. occ.
Bubani, Nym. Consp. 760 = Pycreus capillaris.
bufonius, Maack, ex Trautv. in Act. Hort. Petrop. ix. (1884) 289 = Juncus bufonius.
bulamensis, *Steud. Syn. Pl. Cyp.* 19.—Afr. trop.
bulbiferus, A. Dietr. Sp. Pl. ii. 324 = bulbosus.
bulbiliosus, Brot. ex Heynh. Nom. ii. 181, nomen = rotundus.
bulbocaulis, Boeck. in Linnæa, xxxvi. (1869–70) 372 = Mariscus bulbocaulis.
bulbo-stoloniferus, Miq. Fl. Ind. Bat. Suppl. 360, 559 = rotundus.
bulbo-stoloniferus, Steud. Syn. Pl. Cyp. 18 = stoloniferus.
bulbosus, Boeck. in Linnæa, xxxvi. (1869–70) 300 = bulbosus, ustatus, hysmoides.
bulbosus, Lag. Gen. et Sp. Nov. 2 = congestus.
bulbosus, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 24 = Martianus.
bulbosus, *Vahl, Enum.* ii. 242.—Afr.; Ind. or.
Burchelli, Schrad. Anal. Fl. Cap. Cyp. 11 = textilis.
caducus, Steud. Syn. Pl. Cyp. 315 = Martianus.
caesius, *Boeck. in Linnæa*, xxxvi. (1869–70) 331.—Venezuela.
caespitosus, *Lianos, Fragm.* 14.—Ins. Philipp.
caespitosus, Poir. Encyc. viii. 250 = Pycreus polystachyus.
caespitosus, St. Hil. Voy. Diam. ii. 396 (Pycreí sp.) = *Hilairenus*.
callei, *Bertol. f. in Mem. Acad. Sc. Bolog.* v. (1854) 464. —Afr. austr. ?
calacaryensis, *Steud. Syn. Pl. Cyp.* 84.—Ins. Luzon.
callidus, *Kern. in Oestr. Bot. Zeitschr.* xiv. (1864) 84. —Europ.
californicus, *S. Wats. Bot. Calif.* ii. 216.—Calif.
calliphorus, *G. F. W. Mey. Prim. Fl. Esseq.* 92.—Guiana.
callistus, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 148.—Afr. trop.
calopterus, Miq. Fl. Ind. Bat. Iii. 282 (Mariscí sp.) = *ferax*.
campestris, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 35 = radiatus.
camphoratus, Lieben. in Vidensk. Selsk. Skr. V. ii. (1841) 216 (Mariscí sp.) = strigosus.
canaliculatus, Retz. Obs. vi. 20 = exaltatus.
canariensis, *Steud. Syn. Pl. Cyp.* 42.—Ins. Canar.
cancellatus, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 131.—Afr. trop.
canescens, Vahl, Enum. ii. 255 = Mariscus albescens.
canus, *Y. & C. Presl, Rel. Haenk.* i. 170.—Mexic.
capensis, Boeck. in Linnæa, xxvi. (1853–70) 378 = Mariscus capensis.
capillaris, [Hochst.] Steud. Syn. Pl. Cyp. 6 (Pycreí sp.) = *stramineus*.
capillaris, Koen. ex Roxb. Fl. Ind. i. 194 = Pycreus capillaris.
capillifolius, A. Rich. Tent. Fl. Abyss. ii. 275 (Pycreí sp.).—Abyss.
capitatus, Burm. f. Fl. Ind. 21 = aristatus.
capitatus, Poir. Encyc. vii. 246 = Mariscus Dregeanus.
capitatus, *Retz. Obs.* iv. 9.—Madag.
capitatus, Retz. l. c. = Fimbristylis barbata.
capitatus, Vand. Fasc. Fl. 5 = aegyptiacus.
capitellatus, Schult. Mant. ii. 101 = Mariscus Dregeanus.
Capito, Steud. Syn. Pl. Cyp. 26 = Mariscus appendiculatus.
carnosonodus, Kunth, Enum. Pl. ii. 86 = Mariscus flabelliformis.
carthusus, Pers. Syn. i. 65 = Fimbristylis monostachya.
caricifolius, Hook. & Arn. Bot. Beech. Voy. 99 = Mariscus albescens.
cariciformis, F. Muell. Fragm. v. 92 = Mariscus cariciformis.
carinus, *D. Don, Prod. Fl. Nep.* 30.—Nepal.
carinatus, Smith, Fl. Austral. vii. 276 = filvus, dactylotes, sesquiorus, subulatus.
carinatus, *R. Br. Prod.* 216.—Austral.
carinatus, Nees, in Lehm. Fl. Preiss. ii. 72 = congestus.

CYPERUS :—
carnosus, Heyne, in Wall. Cat. n. 3855 A = procerus.
castaneus, Bigel. Fl. Bost. ed. III. 19 = *diandrus*, Torr.
castaneus, Hochst. ex Miq. Fl. Ind. Bat. iii. 261 = cuspidatus.
castaneus, *Willd. Sp. Pl.* i. 276.—As. trop.; Austral.
casuarinus, Auct. ex Link, Jahrb. i. iii. 89 = glaber.
cayennensis, Link, l. c. 84 = C. Haspan.
cephalanthus, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 431.—Texas.
cephalophorus, J. & C. Presl, Rel. Haenk. i. 170 (Mariscí sp.) = *ferax*.
cephalostachyus, Steud. Syn. Pl. Cyp. 315 = crassipes.
Cephalotes, Boec, ex Kunth, Enum. Pl. ii. 95 = Mariscus glomeratus.
Cephalotes, *Vahl, Enum.* ii. 311.—As. et Austral. trop.
chalarenthus, *Y. & C. Presl, Rel. Haenk.* i. 177. t. 32.—Am. austr.
Chamissoi, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. 1. 32.—Bras.
chilensis, Boeck. in Linnæa, xxxvi. (1869–70) 323 (Mariscí sp.) = *Conceptionis*.
chlorostachys, *Boeck. in Peters, Reise Mossamb. Bot.* 540.—Afr. trop.
chromatolepis, Steud. Syn. Pl. Cyp. 23 = sphaceltatus.
chrysanthus, Boeck. in Linnæa, xxxv. (1867–68) 476 (Pycreí sp.).—Afr. austr.
chrysoherbos, Steud. Syn. Pl. Cyp. 316, sphalm. = erythrorhizos.
chrysostylus, Link, Hort. Berol. ii. 206 = C. Iria.
chrysostachys, *Boeck. in Flora*, xlii. (1859) 488 [bis].—Am. bor.
ciliatus, *Yungh. in Linnæa*, vi. (1831) 25.—Mexic.
cimicinus, J. & C. Presl, Rel. Haenk. i. 166 (Pycreí sp.).—Am. trop.
cinnamomeus, Retz. Obs. iv. 10 = glomeratus.
clandestinus, Steud. in Flora, xxv. (1842) 594 = Ficinia clandestina.
Cleaveri, Torr. in Ann. Lyc. N. York, iii. (1836) 258 = Pycreus polystachyus.
coercens, Reichb. ex Kunth, Enum. Pl. ii. 20 = amabilis.
cognatus, *Kunth, l. c.* 82.—Afr. austr.
colchicus, C. Koch, in Linnæa, xxi. (1848) 623 = Pycreus capillaris.
collinus, Vahl, Enum. ii. 301 = striatus.
colonatus, Vahl, l. c. 313 = Mariscus coloratus.
colymbinus, Kotschy & Peyr. Pl. Tinn. 49. t. 24.—Afr. trop.
communitatus, Steud. Syn. Pl. Cyp. 10 = pumilus.
comosus, *Poir. Encycl. Suppl.* v. 185.—Am. austr.
comosus, Sibth. & Sm. Fl. Graec. i. 31. t. 44 = rotundus.
comosus, Sieber, ex Steud. Nom. ed. II. i. 467 = longus.
compactus, Lam. Tabl. Encyc. i. 144 = obtusiflorus.
compactus, Retz. Obs. v. 10 = Mariscus microcephalus.
complanatus, Forsk. Fl. Aegypt. Arab. 14 = difformis.
complanatus, Presl, in Oken, Isis, xxi. (1828) 271 = Pycreus capillaris.
complanatus, Willd. Sp. Pl. i. 270 = Fimbristylis complanata.
comporitus, Boeck. in Linnæa, xxxvi. (1869–70) 333 (Mariscí sp.) = *Gunnii*.
compositus, R. Br. Prod. 217 = alopecuroides.
compressus, Jacq. Hort. Vindob. iii. 10. t. 12 = vegetus.
compressus, Krock. Fl. Sil. i. 24 = fuscus.
compressus, *Linn. Sp. Pl.* 46.—Cosmop. trop.
compressus, Muhl. Desc. Gram. 15 (Pycreí sp.).—Am. bor.
compressus, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 271 = surinamensis.
compressus, Thw. Enum. Pl. Zeyl. 342, partim = Zollingeri.
Conceptionis, Steud. Syn. Pl. Cyp. 43 (Mariscí sp.).—Chili.
concinnus, R. Br. Prod. 214.—Austral.
concolor, Steud. Syn. Pl. Cyp. 6 (Pycreí sp.) = *C. Eragrostis*, Vahl.
confertiflorus, Schult. Mant. ii. 121 = radiatus.
confertus, Lam. Tabl. Encyc. i. 146 = Pycreus capillaris.
confusus, *Sw. Prod. Veg. Ind. Occ.* 20.—Am. trop.
congestus, Poir. Encyc. vii. 239 = Mariscus microcephalus.
congestus, *Vahl, Enum.* ii. 260.—Geront. trop.

CYPERUS :—
conglobatus, Link, Jahrb. i. iii. 87 (Mariscí sp.) = *ferax*.
conglomeratus, Cons. Expl. Algér. Bot. iii. 243 = Aucheri.
conglomeratus, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 15 = pungens.
conglomeratus, Hochst. ex Boeck. in Linnæa, xxxv. (1867–68) 522 = proteinolepis.
conglomeratus, *Rottb. Descr. Nov. Pl.* 21; *Prog.* 17. —Geront. trop.
conglomeratus, Thw. Enum. Pl. Zeyl. 343 = pachyrrhizus.
conglomeratus, Vahl, Enum. ii. 334 = Pycreus capillaris.
conglomeratus, Willd. Enum. Hort. Berol. Suppl. 5 = compressus.
comicus, Benth. Fl. Austral. vii. 290, partim = Mariscus albescens.
comicus, Boeck. in Linnæa, xxxviii. (1874) 371 (Mariscí sp.).—Austral.
conjunctus, Steud. Syn. Pl. Cyp. 18 = stoloniferus.
conoideus, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = C. Luzulae.
consanguineus, *Kunth, Enum. Pl.* ii. 42.—Bras.
consocius, *Steud. ex Yard. in Mém. Soc. Sc. Nat. Cherb.* v. (1857) 299, nomen.—Ins. Marquesas.
conspicuus, Steud. Syn. Pl. Cyp. 39 = giganteus.
contractus, Steud. l. c. 18 = cuspidatus.
convallium, *Steud. l. c.* 5.—Bras.
coriaceus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 42 = Martianus rufus.
corifolius, Boeck. in Linnæa, xxxvi. (1869–70) 391 = Mariscus latifolius.
cornutus, Anders. Galapag. Cazues Veg. 53 = Mariscus cornutus.
coromandelinus, Boeck. in Linnæa, xxxv. (1867–68) 480 (Pycreí sp.) = stramineus.
coromandelinus, Spreng. Syst. i. 217 = rubicundus.
coronarius, *Kunth, Enum. Pl.* ii. 44.—Ind. or.
correctus, Steud. Syn. Pl. Cyp. 7 = Pycreus propinquus.
corticalis, *Wright, in Sauv. Fl. Cub.* 171.—Cuba.
corymbosus, Heyne, in Wall. Cat. n. 3852 = tegetum.
corymbosus, *Hook. f. Fl. Tasm. Praef.* 47.—Afr.; Austral.
corymbosus, *Rottb. Descr. Nov. Pl.* 42; *Prog.* 19.—Geront. trop.
corymbosus, Korh. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 51 = Pycreus polystachyus.
corymbosus, Steud. in Flora, xli. (1829) 158 = sphaerospermus.
cosyrensis, Tineo, ex Guss. Fl. Sic. Syn. ii. 779 = laevigatus.
crassipes, *Vahl, Enum.* ii. 299.—Afr. trop.
crinitus, Poir. in Lam. Encyc. vi. 762 (lapsu Scirpus!) = Ficinia filiformis.
crinitus, Spreng. in Flora, xli. (1829) 1. Beibl. 5 (err. typ. pro 'cruentus') = semifirtidus.
crocens, *Vahl, Enum.* ii. 257.—Porto Rico.
cruciformis, Boeck. in Linnæa, xxxv. (1867–68) 572 = vaginatus.
cruentus, Baker, Fl. Maurit. 498 = Pycreus Mundtii.
cruentus, Boeck. in Linnæa, xxxvi. (1869–70) 338 = Mariscus Dregeanus.
cruentus, Retz. Obs. v. 13 (Pycreí sp.) = *C. Eragrostis*, Vahl.
cruentus, Rottb. Descr. Nov. Pl. 21; Prog. 17 = conglomeratus.
cubriformis, Llanos, Fragm. 18 = Mariscus microcephalus.
cuanensis, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 136 (Pycreí sp.).—Afr. trop.
cubanus, Lieben. in Vidensk. Selsk. Skr. V. ii. (1851) 222 (Mariscí sp.) = *ferax*.
cubensis, *Steud. Syn. Pl. Cyp.* 42.—Cuba.
Cumingii, Nees, in Linnæa, ix. (1834) 284 = xanthostachyus.
cuprus, J. & C. Presl, Rel. Haenk. i. 172 = erythrorhizos.
curvatus, Vahl, Enum. ii. 338 = Pycreus capillaris.
curvifolius, *Boeck. in Flora*, lxi. (1878) 188.—Reg. Argent.
curvulus, Boeck. in Linnæa, xxxv. (1867–68) 541 = conglomeratus.
cuspidatus, *H. B. & K. Nov. Gen. et Sp.* i. 204.—Cosmop. trop.
cyclostachyus, Griseb. Fl. Brit. W. Ind. 567 = Mariscus echinatus.

CYPERUS :—
cylindricus, *Boeck. in Flora*, xlii. (1859) 487 [bis].—Ind. or.
cylindricus, Britton, in Bull. Torrey Club, vii. (1880) 48. t. 3. f. 2 = Mariscus cylindricus.
cylindricus, *Chapm. in Coult. Bot. Gaz.* iii. (1877) 18.—Am. bor.
cylindrostachys, Boeck. in Linnæa, xxxvi. (1869–70) 388 = flavus, Haenkei, macer, philippensis, Sieberianus.
cymbaeformis, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 208.—N. Granat.
cymulosus, *Willem. in Uteri, Ann. Bot.* xvii. (1796) 7.—Ins. Mauric.
cyrtolepis, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 436.—Am. bor.
cyrtostachys, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 59 (Pycrei sp.) = *C. Eragrostis*.
dactyliformis, Boeck. in Linnæa, xxxvi. (1869–70) 329 (Marisci sp.).—Afr. austr.
dactylotes, *Benth. Fl. Austral.* vii. 278.—Austral.
damiettensis, A. Dietr. Sp. Pl. ii. 269 = osculentus.
daphanus, Ridl. in Journ. Bot. xxii. (1884) 16 = Mariscus Dregeanus.
debilis, *R. Br. Prod.* 213.—Austral.
decidues, *Boeck. in Flora*, lxii. (1879) 547.—Afr.; Austral.
Deckenii, *Boeck. in Linnæa*, xxxviii. (1874) 361.—Afr. trop.
declinatus, Moench, Meth. 317 = vegetus.
decompositus, F. Muell. Fragm. viii. 267 = Mariscus decompositus.
dehiscens, Kunth, Enum. Pl. ii. 56 = tegetum.
Delavayanus, Webb & Berth. Phyt. Canar. iii. 363 = Meyerianus.
delicatulatus, Steud. Syn. Pl. Cyp. 21 = flavidus.
densiflorus, Link, Jahrb. i. 111. 88 = radiatus.
densiflorus, G. F. W. Mey. Prim. Fl. Esseq. 34 (Marisci sp.) = *ferax*.
densiflorus, Reichb. ex Kunth, Enum. Pl. ii. 65 = giganteus.
densifolius, Kunth, Enum. Pl. ii. 114 = Pycreus densifolius.
densifolius, Steud. Syn. Pl. Cyp. 6 = Pycreus Mundtii.
dentatus, R. Br. in Salt, Abyss. App. p. lxii = conglomeratus.
dentatus, Link, Jahrb. i. iii. 88 (Pycrei sp.) = *Humboldtianus*.
dentatus, *Torr. Fl. U. St.* i. 61.—Am. bor.
denticulatus, Schrad. ex Schult. Mant. ii. 104 = surinamensis.
denudatus, Heyne, in Wall. Cat. n. 3849 = stenostachyus.
denudatus, *Linn. f. Suppl.* 98.—Afr. austr.; Madag.
depauperatus, Steud. Syn. Pl. Cyp. 34 = flabelliformis.
depauperatus, Vahl, Enum. ii. 393 = Eleocharis setacea.
derwellensa, *Steud. in Flora*, xxv. (1842) 556.—Abyss.
Disense, *Steud. Syn. Pl. Cyp.* 98.—Ins. S. Helena.
diandrus, Blvrea, Plant. Insul. 8; ex Parl. Fl. Ital. ii. 28 = fuscus.
diandrus, Torr. Cat. Pl. N. York, 89 (Pycrei sp.).—Am. bor.
diaphanirio, Steud. Syn. Pl. Cyp. 23 = C. Iria.
diaphanus, *Schrad. ex Schult. Mant.* ii. 477.—Ind. or.
dichromaeaeformis, *Kunth, Enum. Pl.* ii. 26.—Brss.
dichrostachyus, *Hochst. in Flora*, xxiv. (1841) 1; *Intell.* 21, nomen.—Abyss.
Dietrichiae, Boeck. in Flora, lviii. (1875) 87 (Marisci sp.).—Austral.
difformis, Blanco, Fl. Filip. ed. I. 32 = procerus.
difformis, *Linn. Cent. Pl.* ii. 6; *Amoen. Acad.* iv. 302.—Gerost. trop.
diffusus, Rosb. Hort. Beng. [61]; Fl. Ind. i. 189 = Schœnus Micheliauus.
diffusus, *Vahl, Enum.* ii. 231.—Reg. trop.
digitatus, Nees, in Wight, Contrib. 38 = radiatus.
digitatus, *Roxb. Hort. Beng.* [81]; *Fl. Ind.* i. 206.—Asia; America.
digitatus, Wall. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 125 = diffusus.
dilatatus, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 86.—Afr. trop.
dilutus, Vahl, Enum. ii. 387 = Mariscus microcephalus.
diluvialis, Schult. Mant. ii. 124 (Pycrei sp.) = *puncticulatus*.
diphyllus, Benth. Fl. Austral. vii. 279 = articulatus, decyriotes.
diphyllus, Retz. Obs. v. 11 = corymbosus.

CYPERUS :—
dispaceus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 219 (Marisci sp.).—Mexic.
distiglumis, *Liebm. l. c.* 218.—Mexic.
distiflorus, Anders. in Vet. Akad. Handl. Stockh. 1858 (1858) 153 = confertus.
distiniflorus, Hort. Vrat. ex Nees, in Linnæa, ix. (1834) 325 ; Torr. in Ann. Lyc. N. York, iii. (1836) 266 (Marisci sp.).—Am. bor.
dimolutus, *Link, Jahrb.* i. iii. 87.—Am. trop.
distachyus, All. Auct. Fl. Pedem. 48. t. 2. f. 5 = laevigatus.
distans, Beyr. ex Kunth, Enum. Pl. ii. 14 = Pycreus polystachyus.
distans, *Linn. f. Suppl.* 108.—Cosmop. trop.
distans, G. F. W. Mey. Prim. Fl. Esseq. 35 (Marisci sp.) = *ferax*.
distans, Purah, Fl. Am. Sept. 53 (Marisci sp.) = *strigosus*.
distichophyllus, Steud. in Flora, xxv. (1842) 582 = Pycreus Mundtii.
distinctus, Steud. Syn. Pl. Cyp. 24 = Drummondii.
diurensis, Boeck. in Flora, lxii. (1879) 556 (Marisci sp.).—Afr. trop.
divaricatus, Lam. Tabl. Encyc. i. 145 = Pycreus capillaris.
divergens, Benth. Pl. Hartw. 28 = scleroides.
divergens, Champ. Fl. S. U. St. 512 (Pycrei sp.) = leucolepis.
divergens, H. B. & K. Nov. Gen. et Sp. i. 206 (Marisci sp.) = *Mariscus*.
dives, Delile, Fl. Egypte, 149 = elopecuroides, exaltatus.
dives, Fenzl, ex Steud. Syn. Pl. Cyp. 36 = radiatus.
divulsus, Ridl. in Journ. Bot. xxii. (1884) 15 (Pycrei sp.).—Madag.
Donianus, A. Dietr. Sp. Pl. ii. 290 = pilosus.
Dregeanus, Kunth, Enum. Pl. ii. 81 = pulcher.
Drummondii, Nees, ex Boeck. in Linnæa, xxxvi. (1869–70) 314 = erythrorhizos.
Drummondii, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 437.—Texas.
dubius, Boeck. in Linnæa, xxxvi. (1869–70) 282 = Mariscus colorstus, Schimperi.
dubius, *Rottb. Descr. Nov. Pl.* 30. t. 4. f. 5.—Gerost. trop.
dubius, Kunth, l'l. Ind. i. 188 = Mariscus Dregeanus.
Duchesingii, Steud. Syn. Pl. Cyp. 45 = sphaeolstus.
durus, Kunth, Enum. Pl. ii. 76 (Marisci sp.).—Afr. austr.
eburneus, Thonn. ex Kunth, Enum. Pl. 46 = margaritaceus.
Ecklonii, Boeck. in Linnæa, xxxvi. (1869–70) 325 = Mariscus humillis.
effusus, Link, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 109 = Atkinsoni.
effusus, Rottb. Descr. et Ic. 22. t. 12. f. 3; Prog. 16 = conglomeratus.
egregius, Kunth, Enum. Pl. ii. 69 = fastigiatus.
Ehrenbergianus, *Boeck. in Linnæa*, xxxvi. (1869–70) 291.—Mexic.
Ehrenbergii, Kunth, Enum. Pl. ii. 89 (Marisci sp.) = *ferax*.
elatior, Boeck. in Linnæa, xxxvi. (1869–70) 327 (Marisci sp.).—Afr. austr.
elatior, *Boeck. in Flora*, lxii. (1879) 553.—Afr. trop.
elatus, Griseb. Fl. Brit. W. Ind. 566 = giganteus, comosus.
elatus, Jacq. Cat. Sem. Hort. Vindob. (1826), ex Linnæa, xi. (1837) Lit. 67 = nutans.
elatus, *Linn. Cent. Pl.* i. 5; *Amoen. Acad.* iv. 301.—Ind. or. ; Afr. trop.
elatus, Nees, in Linnæa, ix. (1834) 325 = surinamus.
elatus, Rottb. Descr. et Ic. 37. t. 10 = distans.
elatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 204 = platyphyllus.
elegans, Kunth, Enum. Pl. ii. 28 = diffusus.
elegans, *Linn. Sp. Pl.* 46.—America.
elegans, Schrad. ex Kunth, Enum. Pl. ii. 88 = Mariscus Meyerianus.
elegans, Schreb. ex Schult. Mant. ii. 122 = virens.
elegans, Wall. Cat. 7O (Pycrei sp.) = flavicomus.
elegantulus, Steud. in Flora, xxv. (1842) 582 (Pycrei sp.) = *strevelitzus*.
elephantinus, *Kunth, Enum. Pl.* ii. 20.—Gerost. trop.
Elliottianus, Schult. Mant. ii. 100 = Pycreus propinquus.
elongatus, Hochst. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 100 = Schimperianus.
elongatus, Lej. ex Nees, in Linnæa, ix. (1834) 325 = longus.
elongatus, Sieber, ex Kunth, Enum. Pl. ii. 59 = rotundus.

CYPERUS :—
elongatus, Steud. Syn. Pl. Cyp. 11 = Pycreus polystachyus.
elytrospidrus, *Steud. l. c.* 27.—Hab. ?
emarginatus, Schrad. Anal. Fl. Cap. Cyp. 5 = longus.
eminens, Klein, ex Kunth, Enum. Pl. ii. 70 = platyphyllus.
enervis, *R. Br. Prod.* 213.—Austral.
Engelmannii, Steud. Syn. Pl. Cyp. 47 (Marisci sp.).—Am. bor.
enodis, Boeck. in Linnæa, xxxvi. (1869–70) 271 = corymbosus.
enzelolius, *Nees & Ehrenb. ex Boeck. in Linnæa*, xxxv. (1867–68) 546.—Arab.
Enzlenii, Pursh, Fl. Am. Sept. i. 53 (Marisci sp.) = *strigosus*.
enterianus, *Boeck. in Flora*, lxi. (1878) 189.—Uruguay.
Eragrostis, Clem. ex Willk. & Lange, Prod. Fl. Hisp. i. 127 = Pycreus Mundtii.
Eragrostis, Kunze, in Flora, xxviii. (1845) 754 (Pycrei sp.) = lanceolatus.
Eragrostis, Lam. Tabl. Encyc. i. 196 = vegetus.
Eragrostis, Rottb. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 48 = Pycreus capillaris.
Eragrostis, Vahl, Enum. ii. 393 (Pycrei sp.).—Gerost. trop.
Eragrostis, Willd. ex Kunth, Enum. Pl. ii. 50 = stoloniferus.
eriophorus, *Steud. Syn. Pl. Cyp.* 50.—China.
erubescens, Link, Enum. Hort. Berol. i. 44 = glaber.
erythraeus, Schrad. ex Schult. Mant. ii. 477 = glaber.
erythrorhizos, *Muhl. Desc. Gram.* 20.—Am. bor.
erythrorhizus, Torr. Fl. U. St. i. 61 (Marisci sp.) = *Michauxianus*.
esculentus, Gouan, Fl. Monsp. 389 = rotundus.
esculentus, *Linn. Sp. Pl.* 45.—Cosmop. trop.
esculentus, Vahl, ex Kunth, Enum. Pl. ii. 60 = longus.
etesius, Schult. Mant. ii. 114 = Mariscus glomeratus.
eumorphus, Steud. Syn. Pl. Cyp. 22 = pelcherrimus.
eurystachys, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 143 (Marisci sp.).—Afr. trop.
evaginatus, *Boeck. in Linnæa*, xxxvi. (1869–70) 351 —Florida.
exaltatus, *Retz. Obs.* v. 11.—As. et Austral. trop.
exaltatus, Soland. ex Kunth, Enum. Pl. ii. 90 (Marisci sp.) = *ferax*.
exaltatus, Strachey, Cat. Kumaon Pl. 74 = nutans.
exctus, *Boeck. in Linnæa*, xxxv. (1867–68) 528.—Afr. trop.
exiguus, Nees, in Linnæa, ix. (1834) 324 = cuspidatus.
exilis, *Willd. ex Kunth, Enum. Pl.* ii. 50.—Ins. Mascar. Madag.
expansus, Boj. Hort. Maurit. 330 = Pycreus albo-marginatus.
expansus, *Poir. Encyc.* viii. 369.—Madag.
falcatus, Ehrenb. ex Boeck. in Linnæa, xxxv. (1867–68) 547 = conglomeratus.
falciculosus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 204 = sclerotus.
familiaris, *Steud. Syn. Pl. Cyp.* 46.—Cuba.
fascicularis, DC. Fl. Fr. iii. 723 = Pycreus capillaris.
fascicularis, Phil. ex Kunth, Enum. Pl. ii. 35 = glaber.
fascicularis, Poir. Voy. Barb. ii. 58 = Pycreus polystachyus.
fasciculatus, Ell. Sketch. i. 68 = Pycreus propinquus.
fastigiatus, Forsk. Fl. Aegypt. Arab. 14 = exaltatus.
fastigiatus, Heyne, ex Wall. Cat. sub n. 3446 = sieuinoides.
fastigiatus, *Roth. Descr. Nov. Pl.* 31; *Prog.* 18.—Afr.; Austral.
fastigiatus, Vahl, Enum. ii. 367 = platyphyllus.
fastigiatus, Willd. ex Kunth, Enum. Pl. ii. 60 = longus, tegetum.
fastigiosus, J. F. Gmel. Syst. 134 = fastigiatus.
fastuosus, Desv. ex Ham. Prod. Pl. Ind. Occ. 12 (Marisci sp.) = *ferax*.
Faudleriauus, *Boeck. in Linnæa*, xxxv. (1867–68) 530. —Mexic.
Fauselianus, *Steud. Syn. Pl. Cyp.* 36.—Afr. trop.
ferax, Benth. Fl. Austral. vii. 286 = subulatus, haematodes.
ferax, Rich. in Ast. Sco. Hist. Nat. Par. i. (1792) 106 (Marisci sp.).—Cosmop. trop.
fernambucensis, Steud. Syn. Pl. Cyp. 42, sphalm. = pernambucensis.
fernambucensis, Colla, in Mem. Acc. Tor. xxxix. (1838) 32. t. 36 = sellowii.
ferox, Vahl, Enum. ii. 387 (Marisci sp.) = *ferax*.

CYPERUS:—

ferruginescens, Boeck. in Linnaea, xxxvi. (1869–70) 896 (Mariscus sp.) = *Michauxianus*.

ferrugineus, Forsk. Fl. Aegypt.-Arab. 14 = fuscus.

ferrugineus, Linn. Sp. Pl. 44 = Dulichium spathaceum.

ferrugineus, Poir. Encyc. vii. 961 = Pycreus polystachyus.

fertilis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 90.—Afr. trop.

festivus, Link, Jahrb. i. iii. 88 = exaltatus.

Fieldingii, Steud. Syn. Fl. Cyp. 11 = flavidus.

filicinus, Nees, ex Steud. Syn. Pl. Cyp. 14 = acuminatus.

filicinus, Vahl, Enum. ii. 282 = Pycreus polystachyus.

filiculmis, *A. Rich. Fl. Cub. Fanerog.* ii. 262.—Cuba.

filiculmis, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 40 = Mariscus filiformis.

filiculmis, Vahl, Enum. ii. 282 = Mariscus glomeratus.

filiformis, Willd. ex Kunth, Enum. Pl. ii. 85 = Mariscus filiformis.

filiformis, Sw. Prod. Veg. Ind. Occ. 20 = Mariscus havanensis.

filipes, *Benth. Fl. Austral.* vii. 271.—Austral.

fimbriatus, Nees, in Wight, Contrib. 86 = pilosus.

firmus, J. & C. Presl, Rel. Haenk. i. 171 = Mariscus albescens.

Fischerianus, *Schimp. ex Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 21, nomen; *et in A. Rich. Tent. Fl. Abyss.* ii. 488.—Abyss.

fissus, *Steud. in Flora*, xxv. (1842) 593.—Abyss.

fistulosus, Ehrenb. ex Boeck. in Linnaea, xxxvi. (1869–70) 275 = articulatus.

flabellaris, Schrad. ex Nees, l. c. vii. (1832) 519 = fastigiatus.

flabelliformis, *Rottb. Descr. Nov. Pl.* 42; *Prog.* 22.—Arabia; Abyss.

flaccidus, *R. Br. Prod.* 213 = trinervis.

flagellatus, Hochst. in Flora, xxiv. (1841) 1. Intell. 21 = flabelliformis.

flavamariscus, Griseb. Fl. Brit. W. Ind. 567 = Mariscus flavus.

flavens, Pall. Tabl. Tanr. 45 = Pycreus flavescens.

flavescens, Linn. Sp. Pl. 46 = Pycreus flavescens.

flavescens, Oliver, in Trans. Linn. Soc. xxix. (1875) 164 (Pycrei sp.) = *Apollii*.

flavescens, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 271 = articulatus.

flavescens, Thw. Enum. Pl. Zeyl. 342 = Pycreus capillaris.

flavescens, Willd. ex Kunth, Enum. Pl. ii. 226 = Fimbristylis dichotoma.

flavicomus, Balb. ex Kunth, Enum. Pl. ii. 92 (Mariscus sp.) = *ferax*.

flavicomus, Michx. Fl. Bor. Am. i. 27 (Mariscus sp.) = *strigosus*.

flavicomus, Pursh, Fl. Am. Sept. i. 58. (Mariscus sp.) = *Michauxianus*.

flavicomus, *Schlecht. in Linnaea*, vi. (1831) 34.—Vera-Cruz.

flavicomus, Torr. Bot. Mex. Bound. 226 = Pycreus albo-marginatus.

flavicomus, Vahl, Enum. ii. 260 (Pycrei sp.).—Am. bor. et austr.

flavicomus, Willd. ex Kunth, Enum. Pl. ii. 14 = Pycreus polystachyus.

flavidus, Bel. Hort. Mauric. 280 = Pycreus polystachyus.

flavidus, *Retz. Obs.* v. 18.—Geront. trop.

flavidus, Vahl, Enum. ii. 256 = Pycreus capillaris.

flavissimus, Schrad. in Goett. Gel. Anz. (1821) iii. 2067 = obtusiflorus.

flavissimus, Steud. in Flora, xli. (1839) 153 = denudatus.

flavus, Boeck. in Linnaea, xxxvi. (1869–70) 864 = Mariscus flavus.

flavus, J. & C. Presl, Rel. Haenk. i. 176 = Pycreus propinquus.

floribibens, Boeck. in Flora, lxii. (1879) 840 = Mariscus strumigninus.

flosculosus, Griseb. Fl. Brit. W. Ind. 566 (Mariscus sp.) = *Vahlii*.

flosculosus, Vahl, Enum. ii. 268. (Mariscus sp.) = *ferax*.

fluitans, Buch.-Ham. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 118 = platycrylis.

fluminalis, Kunth, in Trans. Linn. Soc. Ser. II. ii. (1884) 137. (Pycrei sp.).—Afr. trop.

foliosus, Willd. ex Kunth, Enum. Pl. ii. 11 = Pycreus polystachyus.

fontinalis, Kunth, l. c. ii = Pycreus flavescens.

fontinalis, Kunth, l. c. 86 = marginatus.

CYPERUS:—

formosus, Vahl, Enum. ii. 237 = surinamensis.

Forskohlei, *A. Dietr. Sp. Pl.* ii. 251 = fuscus.

Fortunei, Steud. Syn. Pl. Cyp. 21 = malaccensis.

fossarum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 225 (Mariscus sp.) = *ferax*.

fossarum, *Schiecht. in Linnaea*, xxxi. (1861–62) 316.—Guiana.

fragilis, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 238 (Mariscus sp.) = *ferax*.

fraternus, *Kunth, Enum. Pl.* ii. 42.—Bras.

fraternus, Nees, in Mart. Fl. Bras. ii. i. 19 = reflexus.

Presenti, *Steud. Nom.* ed. II. i. 469.—Abyss.

fscatus, *Boeck. in Flora*, lxv. (1882) 13.—Afr. trop.

fugax, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 196 = Pycreus polystachyus.

fulgurinus, Chapm. Fl. S. Un. St. 511 (Mariscus sp.).—Am. bor.

fulvescens, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 210 = esculentus.

fulvus, *R. Br. Prod.* 215.—Austral.

fulvus, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 126 (Pycrei sp.).—Afr. trop.

fuscescens, *Link, Jahrb.* i. iii. 82.—Am. austr.

fuscus, *Linn. Sp. Pl.* 46.—Europ.; Orient.

galagensis, *C. B. Clarke, in Journ. Linn. Soc.* xx. (1883) 385.—Ins. Galaga.

gangeticus, Roxb. ex Wall. Cat. sub n. 8851 = malaccensis.

Gardneri, Baker, Fl. Maurit. 418 = Bakeri.

Gardneri, *Nees, in Mart. Fl. Bras.* ii. i. 34.—Am. trop.

Gatesii, Torr. in Ann. Lyc. N. York, iii. (1836) 255 = Pycreus polystachyus.

geniculatus, Koen. ex Ainslie, Mat. Med. Hind. 250 = bulbosus.

geminatus, Schrad. ex Schult. Mant. ii. 95 = Fimbristylis bahlensis.

gigantiacus, Griseb. Cat. Pl. Cub. 238 = digitatus.

gigantinus, G. F. W. Mey. Prim. Fl. Esseq. 85 = comosus.

gigantus, *Noronha, ex Thou. Prod. Phyt.* 2, *in Milang.* (1811) nomen.—Madag.

giganteus, *Rottb. ex Kunth, Enum. Pl.* ii. 65.—Jamaica.

gigantus, *Vahl, Enum. Pl.* ii. 264.—Am. trop.

Gilesii, *Benth. Fl. Austral.* vii. 274.—Austral.

Ginge, *Wats. Apont.* 586.—Afr. trop.

glaber, Hoppe, in Sturm, Fl. Deutschl. Heft iii = fuscus.

glaber, *Linn. Mant.* ii. 179.—Europ.; Orient.

glaber, Pourr. ex Willd. & Lange, Prod. Fl. Hisp. i. 138 = fuscus.

glaber, Thou. ex Kunth, Enum. Pl. ii. 12 = Pycreus polystachyus.

glaber, Vill. Hist. Pl. Dauph. ii. 182 = Monti.

glandulosus, Rolfe, in Journ. Bot. xx. (1882) 362 = Mariscus rufus.

glaucinus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 204 = umbellis.

glaucinus, Boeck. in Flora, lviii. (1875) 89 = Mariscus conicus.

glaucus, Steud. Syn. Pl. Cyp. 42. (Mariscl sp.) = *brisaeus*.

glaucus, Willd. ex Kunth, Enum. Pl. ii. 20 = alopecuroides.

Glaziovianus, *Boeck. in Kjoob. Vidensk. Meddel.* (1869) 138; *et in Linnaea*, xxxviii. (1874) 365.—Bras.

globosus, Lam. Encl. Bot. Fl. Pedem. 49 = Pycreus capillaris.

globosus, Balbis, ex Torr. in Ann. Lyc. N. York, iii. (1836) 270 = Mariscus glomeratus.

globosus, Boeck. in Flora, lxi. (1879) 546 = Pycreus polystachyus.

globosus, Forsk. Fl. Aegypt. Arab. 18 = cruentus.

globuliferus, Link, Jahrb. i. iii. 89 = Luzulae.

globuliferus, J. & C. Presl, Rel. Haenk. i. 165 = Andricticlum polycephalum.

globulosus, Aubl. Pl. Gui. i. 47 = Mariscus glomeratus?

glomeratus, Klein, ex Kunth, Enum. Pl. ii. 127 = Courtoisia cyperoides.

glomeratus, *Linn. Cent. Pl.* ii. 5; *Amoen. Acad.* iv. 301.—Europ.; As. temp.

glomeratus, Roxb. ex Ann. in Wight, Contrib. 85 = digitatus.

glomeratus, Walt. Fl. Carol. 70 = virens.

glomeratus, Willd. Sp. Pl. i. 277 (syn. Linn. excl.) = alopecuroides.

glomuliferus, Schrad. ex Schult. Mant. ii. 106 = C. Luzulae.

CYPERUS:—

Goeringii, Steud. Syn. Pl. Cyp. 24 = difformis.

Grabowskianus, Boeck. in Engl. Bot. Jahrb. v. (1884) 503 = Mariscus microcephalus.

gracilescens, Schult. Mant. ii. 100 = esculentus.

gracilinux, *C. B. Clarke, in Journ. Linn. Soc.* xxi. (1884) 102.—Afr. trop.

gracilis, *R. Br. Prod.* 213.—Austral.

gracilis, J. Buch. in Trans. N. Z. Inst. iii. (1871) 210 = vaginatus.

gracilis, Link, Jahrb. i. iii. 84 = esculentus.

gracilis, Muhl. Desc. Gram. 16 = C. Haspan.

gradatus, Forsk. Fl. Aegypt. Arab. 13 = flabelliformis.

Graeffei, *Boeck. in Flora*, lviii. (1875) 84.—Ins. Samos.

gramineus, Roem. & Schult. Syst. ii. 205 = C. Haspan.

gramineus, Steud. Syn. Pl. Cyp. 5 (Pycrei sp.) = *graminicus*.

graminicolus, Steud. l. c. 49 = distans.

graminifolius, Poir. Encyc. vii. 267 = C. Haspan.

grammicus, Kunze, ex Kunth, Enum. Pl. ii. 9 (Pycrei sp.).—Chili.

granadinus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 224. (Mariscl sp.) = *ferax*.

graudiflorus, *Anders. in Vet. Akad. Handl. Stockh.* 1858 (1858) 157.—Ins. Galap.

Grantii, *Boeck. in Flora*, lviii. (1875) 260; *Oliver, in Trans. Linn. Soc.* xxix. (1875) 164.—Afr. trop.

granulatus, Boec. ex Kunth, Enum. Pl. ii. 95 = Mariscus ovularis, glomeratus.

Grayii, Torr. in Ann. Lyc. N. York, iii. (1836) 260 (Mariscl sp.) = *ferax*.

Griffithianus, Boeck. in Linnaea, xxxv. (1867–68) 601 = pilosus.

Griffithii, Steud. Syn. Pl. Cyp. 316 = radians.

grossarius, Koen. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 57 = Pycreus sanguinolentus.

guatemalensis, *Steud. Syn. Pl. Cyp.* 47.—Guatemala.

Gula-Mettii, Schult. Mant. ii. 125 = tegetiformis.

Gunnii, Hook. f. Fl. Tasm. ii. 80. t. 140 (Mariscl sp.).—Austral.

Guzzonii, Casp. in Guss. Fl. Sic. Syn. ii. 779 = Pycreus flavescens.

gymnocaulos, Steud. Syn. Pl. Cyp. 12 = vaginatus.

gymnolepis, Steud. l. c. 2 = *Pycreus nitens*.

gymnos, Schult. Mant. ii. 97 = articulatus.

Gyrnadini, *Steud. Syn. Pl. Cyp.* 28.—Ins. Borbon.

haematodes, Encl. Prod. Fl. Norfolk. 22 (Mariscl sp.).—Ins. D. Howe et Norfolk.

haematostachys, Steud. *Syn. Pl. Cyp.* 41 = reflexus.

Haenkeanus, Kunth, Enum. Pl. ii. 98 = Mariscus microcephalus.

Haenkei, J. & C. Presl, Rel. Haenk. i. 172 (Mariscl sp.) = *ferax*?

Hahnianus, Boeck. in Flora, lxi. (1878) 138 (Pycrei sp.).—Ind. occ.

hakonensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 104. 588.—Japon.

hamatus, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 28 = uncinulatus.

Hamiltonii, Kunth, Enum. Pl. ii. 90 (Mariscl sp.) = *ferax*.

hamulatus, *Kunth, l. c.* 22.—Ind. occ.

hamulosus, Bieb. Fl. Taur. Cauc. i. 35 = Scirpus hamulosus.

Haspan, Benth. Fl. Austral. vii. 270, partim = pulcherrimus.

Haspan, *Linn. Sp. Pl.* 45.—Cosmop. trop.

Haspan, Kunth, Enum. Pl. ii. 96. t. 6 f. 2 = flavidus.

hassanensis, Willd. ex Kunth, Enum. Pl. ii. 84 (Mariscl sp.).—Ind. occ.

hassanicus, H. Mann, in Proc. Am. Acad. vii. (1867) 306 = Mariscus phleoides.

Haworthii, S. F. Gray, Nat. Arr. Brit. Pl. ii. 730 = fuscus.

hebes, Steud. Syn. Pl. Cyp. 315 = pilosus.

Hermannii, Buckl. in Proc. Ac. Sc. Philad. 1862 (1863) 10 = esculentus.

Heldreichianus, Boiss. Diagn. Ser. I. xiii. 39 = longus.

Heliert, *Boeck. in Linnaea*, xxxviii. (1874) 360.—Borneo.

helodes, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 37 = retundus.

helvus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 197 = Pycreus propinquus.

hemisphaericus, Boeck. in Flora, xlii. (1859) 496 ter (err. typ. pro 489 bis) (Mariscl sp.).—Afr. trop.

herbiscovus, Mellise, S. Helena, 848 = rotundus.

herbivagus, Kunth, Enum. Pl. ii. 59 = semitrifidus.

CYPERUS:—
heterocladus, *Baker, in Journ. Linn. Soc.* xx. (1883) 292.—Madag.
heterophyllus, Boeck. in Flora, xlii. (1859) 441 = maritimus.
hexastachyos, Rottb. Descr. Nov. Pl. 38 = rotundus.
hexastachyus, Sw. Fl. Ind. Occ. i. 117 = sphacelatus.
hexastachyus, Nees, in Wight, Contrib. 82 = bulbosus, tuberosus.
Heyneanus, Boeck. in Flora, xlii. (1859) 440 [bis] = pilosus.
Heyneï, Boeck. in Linnaea, xxxv. (1867–68) 600 = procerus.
Hilairenus, Steud. Syn. Pl. Cyp. 99 (Pycreï sp.).—Bras.
Hildebrandtii, Boeck. in Flora, lxiii. (1880) 436 (Mariscï sp.) = *hemisphæricus*.
Hildra, Poir. Encyc. vii. 251 = rotundus.
Hilgendorfianus, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 501.—Japon.
Hillebrandii, Boeck. in Flora, lxiii. (1880) 436 = Mariscus Kunthianus.
hirsutus, Berg. Descr. Pl. Cap. 11 = Fuirena glabra.
hirtus, Thunb. in Hoffm. Phytogr. Bluett. i. (1803) 6 (quid ?).—Afr. austr.
Hochstetteri, *Nees, ex Krauss, in Flora*, xxix. (1846) 755, in nota = Pycreus albo-marginatus.
hoiciflorus, J. & C. Presl, Rel. Haenk. i. 173 = Mariscus albescens.
holophyllus, *Miq. ex De Vriese, Pl. Ind. Bat. Reinw.* 139.—Ins. Celebes.
holoschoenoides, Jan, Cat. Pl. Phasn. 75, ex Schult. Mant. ii. 111 = difformis.
Holoschoenus, R. Br. Prod. 215 (Mariscï sp.).—Austral.
holosericeus, Link, Hort. Berol. i. 217 = Pycreus polystachyus.
homalirius, Steud. Syn. Pl. Cyp. 28 = diffusus.
honestus, Kunth, Enum. Pl. ii. 74 = pilosus.
honoluluensis, *Steud. ex Yard. in Bull. Soc. Linn. Normand.* Sér. II. ix. (1875) 278, 280.—Ins. Sandvic.
Hookeri, Boeck. in Linnaea, xxxvi. (1869–70) 308 = digitatus.
Hookerianus, Arn. ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 280 = Pycreus polystachyus.
Hookerianus, Thw. Enum. Pl. Zeyl. 342 = Cephalotes.
horticolalis, Heyne, ex Wall. Cat. n. 8313 = aguarrosa.
hyalinus, Vahl, Enum. ii. 290 = Pycreus pumilus.
Hydra, H. B. & K. Nov. Gen. et Sp. i. 205 = esculentus.
Hydra, Michx. Fl. Bor. Am. i. 27 = rotundus.
hyemalis, Pursh, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 130 = C. Haspan.
hylæus, *Ridl. in Trans. Linn. Soc. Ser.*-II. ii. (1884) 184.—Afr. trop.
ignoratus, Boeck. in Linnaea, xxxvi. (1869–70) 895 (Mariscï sp.) = *ferax*.
imbecillis, R. Br. Prod. 213 = trinervis.
imbricatus, Llanos, Fragm. 17 = Mariscus albescens.
imbricatus, Retz. Obs. v. 13 = platyphyllus.
imbricatus, Vill. ex Roem. & Schult. Syst. ii. 208 = glaber.
imbricatus, Willd. ex Kunth, Enum. Pl. ii. 20 = alopecuroides.
imerinensis, Boeck. in Engl. Bot. Jahrb. v. (1884) 500 = madagascariensis.
immensus, *C. B. Clarke, in Journ. Linn. Soc.* xx. (1883) 294.—Madag.
impolitus, *Kunth, Enum. Pl.* ii. 78.—Bras.
impubes, Steud. Syn. Pl. Cyp. 45 (Mariscï sp.).—Abyss.
inæquatilis, Boeck, ex Link, Jahrb. i. iii. 84 = C. Haspan.
inæqualis, Link, l. c. 83 (Mariscï sp.) = *Menimae*.
inæqualis, *Willem. in Usteri, Ann. Bot.* xviii. (1796) 5. —Ins. Maurit.
incarnatus, Link, Hort. Berol. i. 206 = congestus.
incompletus, Boeck. in Linnaea, xxxvi. (1869–70) 861 = Mariscus Jacquini.
incompletus, Link, Hort. Berol. i. 219 = Mariscus elatus.
incomtus, *Kunth, Enum. Pl.* ii. 80.—Bras.
inconspicuus, Gennari.—Cf. Linnæa, xxiv. (1851) 900 = rotundus.
inconspicuus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 197 (Pycreï sp.).—Costa Rica.
incrassatus, Berrich, ex Kunth, Enum. Pl. ii. 18 = Pycreus polystachyus.
incurvatus, Roxb. Hort. Beng. [5] ; Fl. Ind. i. 196 = malaccensis.
incurvus, Spreng. ex Kunth, Enum. Pl. ii. 16 = laevigatus.
indecorus, *Kunth, l. c.* ii. 85.—Afr. austr.
indicus, Rich. ex Pers. Syn. i. 65 = Fimbristylis monostachya.
infirmus, *Boeck. in Flora*, xlii. (1859) 438 [bis].—Am. bor.
infirmus, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 8 = Pycreus propinquus.
inflexus, Muhl. Descr. Gram. 16 = aristatus.
infra-apicalis, Nees, ex Aitch. Cat. Punjab Pl. 155 = clausinoides.
infuscatus, Boeck. in Linnaea, xxxvi. (1869–70) 871 = Mariscus longus, fabelliformis.
infuscatus, Kunth, Enum. Pl. ii. 86 = Mariscus longus.
infuscatus, Nees, ex Boeck. in Linnaea, xxxvi. (1869–70) 865 = Mariscus filiformis.
ingratus, *Kunth, Enum. Pl.* ii. 81.—Afr. austr.
inornatus, Boeck. in Flora, lviii. (1875) 86 (Mariscï sp.) = *Gunnii*.
inrigui, Kunth, Enum. Pl. ii. 92 (Mariscï sp.) = *ferax*.
intactus, Vahl, Enum. ii. 339 = Pycreus polystachyus.
interceptus, Steud. Syn. Pl. Cyp. 31 = corymbosus.
intermedius, Gaus. Suppl. Fl. Sic. Prod. 10 = longus.
intermedius, Steud. in Flora, xxv. (1842) 581 (Pycreï sp.) = *lanceolatus*.
intricatus, Kunth, Enum. Pl. ii. 28 = aristatus.
intricatus, *Schrad. ex Schult. Mant.* ii. 98.—Bras.
inundatus, R. Br. Prod. 214 = trinervis.
inundatus, Nees, in Wight, Contrib. 76 = Pycreus pumilus.
inundatus, *Roxb. Hort. Beng.* 6 ; *Fl. Ind.* i. 201.—Ind. or. ; China.
involucratus, Poir. Encyc. vii. 253 = radiatus.
involutus, R. Br. in Salt, Abyss. App. p. lxii = proteinolepis.
Iria, *Linn. Sp. Pl.* 45.—Geront. trop.
iridifolius, A. Dietr. Sp. Pl. ii. 815 = diffusus.
ischnocormus, Steud. Syn. Pl. Cyp. 11 (Pycreï sp.) = *atrovirens*.
ischnos, Schlecht. in Bot. Zeit. vii. (1849) 99 (Mariscï sp.).—Ind. or. ; Am. trop.
ischnostachys, Steud. Syn. Pl. Cyp. 53 = Mariscus cyperinus.
isocladus, Kunth, Enum. Pl. ii. 87 = sequalis.
italicus, *Tod. Ind. Sem. Hort. Panorm.* (1861) 32.—Hab. ?
Jacquemontii, Boeck. in Linnaea, xxxv. (1867–68) 145 (Pycreï sp.).—Ind. occ.
Jacquini, Schrad. in Linnaea, xi. (1837) Litt. 87 = nutans.
jamaicensis, Poir. Encyc. Suppl. v. 166 = Pycreus polystachyus.
japonicus, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 140 (Pycreï sp.).—Japon.
Jardinei, Steud. Syn. Pl. Cyp. 28 = crassipes.
javanicus, Bernh. ex Kunth, Enum. Pl. ii. 21 = castaneus.
jemenicus, Heyne, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 59 = Pycreus anguinholensis.
jemenicus, Retz. Obs. iv. 11 = bulbosus.
jeminus, *Rottb. Descr. Nov. Pl.* 25 ; *Prog.* 24.—Geront. trop.
johnsoniæus, Rudge, Pl. Gui. 17. t. 21 (Mariscï sp.) = *ferax*.
juncens, Link, Jahrb. i. iii. 85 = C. Haspan.
junciformis, Desf. Fl. Atlant. i. 43 = laevigatus.
juncoides, Lam. Tabl. Encycl. i. 147 = C. Haspan.
juncoides, *Rœmer, Ind. Sem. Hort. Petrop.* (1846) 27. —Hab. ?
jurgensenius, Steud. Syn. Pl. Cyp. 3 = Pycreus capillaris.

CYPERUS:—
Junghuhnii, Miq. Fl. Ind. Bat. iii. 260 = Pycreus capillaris.
Kappleri, *Hochst. ex Steud. Syn. Pl. Cyp.* 88.—Guiana.
Kegelianus, Steud. l. c. 29 = Remirea maritima.
Kerstenii, Boeck. in Linnaea, xxxvi. (1869–70) 878 (Mariscï sp.).—Afr. trop.
Kleinianus, Hochst. ex Steud. Syn. Pl. Cyp. 71 = Cuticula cyperoides.
Koenigii, Vahl, Enum. ii. 302 = corymbosus.
Kotschyanus, Fenzl, ex Steud. Syn. Pl. Cyp. 36 = clausinoides.
Kraneri, Franch. & Sav. Enum. Pl. Jap. ii. 104, 539 = Textori.
Kraussii, Boeck. in Linnaea, xxxvi. (1869–70) 879 = Mariscus radiatus, Dregeanus.
Kunthianus, H. Mann, in Proc. Am. Acad. vii. (1867) 208 = Mariscus Kunthianus.
Kunthianus, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 69 = Mariscus Meyenianus.
Kunthii, Parl. Fl. Ital. ii. 41 = longus.
Kurrii, Steud. Syn. Pl. Cyp. 28 = distans.
Kurzii, *C. B. Clarke, in Journ. Linn. Soc.* xxi. (1884) 129.—Ins. Andam.
kyllingioides, Willd. ex Kunth, Enum. Pl. ii. 48 = C. Cephalotes.
kyllingioides, Pursh, Fl. Am. Sept. i. 80 = Mariscus glomeratus.
kyllingioides, Sieber, ex Presl, in Oken, Isis, xxi (1828) 270 = Kyllingia brevifolia.
kyllingioides, Vahl, Enum. ii. 312 = Mariscus Dregeanus.
lacteus, *Steud. Syn. Pl. Cyp.* 37.—Madag.
lacunosus, *Griseb. Cat. Pl. Cub.* 237.—Cuba.
lacustris, *Phil. in Linnaea*, xxxiii. (1864–65) 270.—Chili.
lacustris, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 1 = millifolius.
laetus, *J. & C. Presl, Rel. Haenk.* i. 172.—Chili.
laevigatus, Aceb. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 106 = arenarius.
laevigatus, *Linn. Mant.* ii. 179.—Cosmop. calid.
laevis, *R. Br. Prod.* 213.—Austral.
laevissimus, Steud. Syn. Pl. Cyp. 32 = rotundus.
lagorensis, *Steud. l. c.* 36.—Ins. Philipp.
Lagunetto, Steud. l. c. 5 (Pycreï sp.).—Venezuela.
Lamarchianus, Hochst. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 164 = Fenzelianus.
Lamarchianus, Schult. Mant. ii. 108 = Pycreus capillaris.
lamprocarpus, Boeck. in Linnaea, xxxv. (1867–68) 490 = stoloniferus.
lamprocarpus, *Nees, in Hook. Kew Journ.* vi. (1854) 97.—Sumatra.
lanatus, Ruiz, ex Boeck. in Linnaea, xxxvi. (1869–70) 750 = Scirpus fimbriatus.
lancastriensis, Porter, in A. Gray, Man. Bot. N. U. St ed. V. 555 (Mariscï sp.).—Am. bor.
Lancesis, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 184 = fertilis.
lanceolatus, Poir. Encyc. vii. 245 (Pycreï sp.).—Afr. trop.
lanceolatus, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 271 ; J. & C. Presl, Rel. Haenk. i. 167 = Pycreus capillaris.
lanceus, F. Muell. Fragm. viii. 250 = Pycreus sanguinolentus.
lanceus, Thunb. Prod. Fl. Cap. 18 (Pycreï sp.).—Afr. austr.
lanuginosulus, Ruiz, ex Buchen. Krit. Verz. Juncac. 1 = Luzula gigantea.
lateralis, Forsk. Fl. Aegypt. Arab. 13 = laevigatus.
lateralis, Linn. f. Suppl. 102 = tenellus.
latespiculosus, Steud. in Flora, xli. (1829) 152 = longus.
latestriatus, *Torr. in Bot. Mex. Bound.* 226.—Mexic.
latespiculosus, Boeck. in Flora, xlii. (1859) 438 [bis] (Pycreï sp.).—Ind. or.
latifoliatus, Spreng. Syst. i. 232 = C. Haspan.
latifolius, *Poir. Encyc.* vii. 246.—Afr. austr. ; Ins. Mass.
latifolius, *Poir. l. c.* 270.—Madag.
lauriformis, Sieber, ex Boeck. in Linnaea, xxxv. (1867–68) 575 = sequalis.
laxiusculus, Steud. Syn. Pl. Cyp. 14 = Pycreus elegans.
laxus, R. Br. in Salt, Abyss. App. p. lxi = bulbosus.
laxus, Gmel. Fl. Bad. W. Ind. 561 = diffusus.
laxus, Lam. Tabl. Encyc. i. 146 = elegans.

CYPERUS:—
laxus, Vahl, Enum. ii. 363 = distans.
laxus, *Willem. in Usteri, Ann. Bot.* xvii. (1796) 6.—Ins. Mauric.
Lechleri, Steud. Syn. Pl. Cyp. 37 = xanthostachyus.
Lecontii, *Torr. in Ann. Lyc. N. York*, iii. (1836) 278, in obs.—Am. bor.
Lehmannii, F. Muell. Fragm. viii. 272 = Kyllinga Lehmanni.
leiocaulon, Benth. Fl. Austral. vii. 287 = Mariscus laevis.
lenticularis, Steud. Syn. Pl. Cyp. 48 (Mariscí sp.) = *ferax*.
leoconsis, *Steud. Nom.* ed. II. i. 470.—Afr. trop.
lepidus, Hochst. in Flora, xxv. (1842) 1. Beibl. 134, nomen : Steud. Syn. Pl. Cyp. 14 = amabilis.
lepidus, *F. Muell. ex C. B. Clarke, in Journ. Linn. Soc.* xxi. (1884) 134.—Austral.
leptanthos, *Steud. i. c.* 36.—Bras.
leptocladus, *Kunth, Enum. Pl.* ii. 93.—Afr. austr.
leptophyllus, Hochst. ex Steud. Syn. Pl. Cyp. 38 (Marisci sp.).—Afr. trop.
leptos, Schult. Mant. ii. 105 = C. Haspan.
leptostachys, Nees, in Linnaea, ix. (1834) 285 = flavidus.
leptostachys, Vahl, ex Kunth, Enum. Pl. ii. 93 = distans.
leptostachyus, Griff. Itin. Notes, 321 = rotundus.
Lessonianus, Kunth, Enum. Pl. ii. 39 = trinervis.
lencanthus, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. 1. 18.—Bras.
leucocephalus, Hasak. Pl. Jav. Rar. 87 = Kyllinga monocephala.
leucocephalus, *Rets. Obs.* v. 11.—Ax. et Am. trop.; Austral.
leucocephalus, Wight, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 34 = Cephalotes.
leucolepis, *Boeck. in Linnaea*, xxxv. (1867–68) 610.—Mexic.
leucolepis, Carey, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 61 (Pycrei sp.).—Am. bor.
leucoloma, *Nees, in Linnaea*, ix. (1834) 284 et x. (1836) 138.—Afr. ; Austral.
leucostachys, Boj. ex Baker, in Journ. Linn. Soc. xxi (1883) 460 = Dichromena nodida.
leucostachys, Link, Jahrb. i. iii. 31 = laevigatus.
levis, F. Muell. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 59 = Pycreus sanguinolentus.
Lhotskyanus, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 498.—Austral.
Liebmannii, Steud. Syn. Pl. Cyp. 7 = *inconspicuus*, Liebm.
ligularis, Boeck. in Linnaea, xxxvi. (1869–70) 393, partim = Mariscus albescens.
ligularis, Chapm. Fl. S. U. St. 507, 659 (Marisci sp.) = *brisaeus*.
ligularis, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 196, partim = Mariscus brachystachyus.
ligularis, Hensl. Bot. Voy. Challenger, i. ii. 28 = Mariscus appendiculatus.
ligularis, Hort. Berol. ex Kunth, Enum. Pl. ii. 87 = congestus.
ligularis, Linn. Amoen. Acad. v. 81 = Mariscus rufus.
limosus, *Maxim. Prim. Fl. Amur.* 294.—Reg. Amur.
linctus, Forst. f. Prod. 89 (sphalm. pro 'linearis') = Mariscus Sieberianus.
Linhianus, Schult. Mant. ii. 113 = congestus.
Linnaeanus, Kunth, Enum. Pl. ii. 115 = Pycreus Linnaeanus.
litos, Schult. Mant. ii. 101 = Pycreus polystachyus.
littoralis, R. Br. Prod. 216 = stoloniferus.
litoralis, Heyne, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 46 = Pycreus capillaris.
lividus, *Steud. Syn. Pl. Cyp.* 25.—Guian.
lomatostachys, Nees & Meyen, in Linnaea, ix. (1834) 283 (Marisci sp.) = *ferax*.
longearistatus, *Steud. Syn. Pl. Cyp.* 37.—Mexic.
longiceraria, Boeck. in Linnaea, xxxviii. (1874) 358 (Pycrei sp.).—Reg. Argent.
longifolius, *Poir. Encyc.* vii. 270.—Ins. Mascar. ; Madag.
longifolius, Ruiz, ex Boeck. in Linnaea, xxxvi. (1869–70) 402 (Marisci sp.) = *ferax*.
longiramulosus, Baldwin, ex Boeck. in Linnaea, xxxv. (1867–68) 695 = prolixus.
longus, Baker, Fl. Maurit. 411 = tuberosus.
longus, Boeck. in Journ. Linn. Soc. xviii. (1881) 104 = rotundus.
longus, *Linn. Sp. Pl.* 68.—Europ. ; Orient.

CYPERUS:—
longus, A. Rich. Tent. Fl. Abyss. ii. 484 = Schimperus.
Lorentzianus, Boeck. in Linnaea, xxxviii. (1874) 357 (Pycrei sp.).—Reg. Argent.
Louisianus, Steud. Syn. Pl. Cyp. 10 = Pycreus polystachyus.
luciduliae, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 99 = Zollingeri, Schweinfurthianus.
luciduli, Link, Jahrb. i. iii. 86 = tuberosus.
lucidus, *R. Br. Prod.* 218.—Austral.
lucidus, Kunth, Enum. Pl. ii. 89 (Mariscí sp.) = *ferax*.
lucidus, Rottl. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 48 = Pycreus capillaris.
Luerssenii, *Boeck. in Flora*, lviii. (1875) 88.—Austral.
luteolus, Boeck. i. c. 82 = Pycreus angulatus.
lutescens, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 433.—Am. bor. et austr.
lutens, Boeck. in Linnaea, xxxviii. (1874) 370 (Mariscí sp.).—Madag.
luzonensis, *Y. & C. Presl, Rel. Haenk.* i. 174.—Ins. Luzon.
Luzulae, Hochst. ex Steud. Syn. Pl. Cyp. 25 = vegetus.
Luzulae, *Rottl. ex Willd. Sp. Pl.* i. 276.—Am. trop.
luzuliformis, *Boeck. in Linnaea*, xxxvi. (1869–70) 386.—Afr. austr.
macellus, Kunth, Enum. Pl. ii. 38 = trinervis.
maceri, *C. B. Clarke, in Journ. Linn. Soc.* xxi. (1884) 160.—Burma.
macranthus, Boeck. in Linnaea, xxxv. (1867–68) 462 (Pycrei sp.).—Afr. austr.
macreilema, *Steud. ex Yard. in Mém. Soc. Sc. Nat. Cherb.* iv. (1856) reimp. 10, 36 ; v. (1857) 298.—Ins. Marquesas.
macrocephalus, Boeck. in Linnaea, xxxvi. (1869–70) 380 = Mariscus macrocarpus.
macrophaeus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 231 (Mariscí sp.).—Mexic.
macrophyllus, Boeck. in Linnaea, xxxvi. (1869–70) 376 = Mariscus macrophyllus.
macropus, Boeck. in Flora, lxii. (1879) 550 = Mariscus Dregeanus.
macropus, Miq. Fl. Ind. Bat. Suppl. 599 = radians.
macrorrhizus, Nees, ex Boiss. Fl. Orient. v. 369 = conglomeratus.
macrorrhizus, Nees, in Wight, Contrib. 78 = aegyptiacus.
Macrostachyon, Steud. Syn. Pl. Cyp. 37 = radiatus.
macrostachyus, Lam. Tabl. Encyc. i. 147 = Pycreus pumilus.
macrostachyus, Tenore, ex Boeck. in Linnaea, xxxvi. (1869–70) 324 = rotundus.
maculatus, *Boeck. in Peters, Reise Mossamb. Bot.* 589.—Afr. trop.
madagascariensis, *Roem. & Schult. Syst.* ii. 876.—Afr. trop. ; Madag. ; Austral.
madevastatanus, Willd. Sp. Pl. i. 278 = squarrosus.
malaccensis, *Lam. Tabl. Encyc.* i. 146.—Ax. trop.
manilensis, Boeck. in Engl. Bot. Jahrb. v. (1884) 501 = Mariscus cyperinus.
Manimae, H. B. & K. Nov. Gen. et Sp. i. 209 (Marisci sp.).—Am. trop.
margaritaceus, *Vahl, Enum.* ii. 367.—Afr. trop.
marginatus, *Thunb. Prod. Pl. Cap.* 18.—Afr. austr.
marginellus, Nees, in Wight, Contrib. 83 = pilosus.
martinicides, Nutt. Gen. Am. i. 35 = Mariscus glomeratus.
Mariscus, Nees, in Linnaea, vii. (1832) 515 = albostriatus.
maritimus, Boj. Hort. Maurit. 378 = rotundus.
maritimus, Gouan, ex Spreng. Syst. i. 217 = laevigatus.
maritimus, *Poir. Encyc.* vii. 240.—Madag.
Martianus, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. 1. 23.—Bras.
Martianus, *Schult. Mant.* ii. 108 (sp. ignot.).—Bras.
mauritianus, *Willem. in Usteri, Ann. Bot.* xviii. (1796) 4.—Ins. Mauric.
Maximiliani, Griseb. in Goett. Abh. xxiv. (1879) 311 (Marisci sp.) = *ferax*.
Maximiliani, Schred. ex Nees, in Mart. Fl. Bras. ii. 1. 7 = Pycreus propinquus.
maximus, Roxb. ex Nees, in Wight, Contrib. 85 = platyphyllus.
megapotamicus, Kunth, Enum. Pl. ii. 10 = Pycreus megapotamicus.
melanocephalus, R. Br. in Salt, Abyss. App. p. lxii (Pycrei sp.) = *atronitens*.
melanocephalus, Miq. Fl. Ind. Bat. iii. 287 = Pycreus sanguinolentus.

CYPERUS:—
melanopus, Boeck. in Flora, lxii. (1879) 545 (Pycrei sp.) = *laxevus*.
melanorhizus, Deille, Fl. Aeg. Illustr. 50 = esculentus.
melanostachyus, Anct. ex Walp. Ann. iii. 667 (Pycrei sp.) = *pleurs*.
melanostachyus, H. B. & K. Nov. Gen. et Sp. i. 207 (Pycrei sp.).—Am. austr.
melanostachyus, Palmer, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 64 (Pycrei sp.) = *C. Lagunetto*.
melas, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 127 (Pycrei sp.).—Afr. trop.
melicoides, Poir. Encyc. vii. 278 (Cladii sp.).—Ins. Mascar.
membranaceus, Vahl, Enum. ii. 330 = Pycreus virens.
mexicanus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 227 = digitatus.
Meyenianus, Kunth, Enum. Pl. ii. 88 = Mariscus Meyenianus.
Meyenii, Nees & Arn. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 87 = compressus.
Meyerianus, *Kunth, Enum. Pl.* ii. 49.—Afr. austr.
micans, Kunth, i. c. 12 = Pycreus polystachyus.
Michauxianus, Schult. Mant. ii. 123 (Mariscí sp.).—Am. bor.
Michelianus, Delile, Fl. Aeg. Illustr. 50 = pygmaeus.
micranthus, J. & C. Presl, Rel. Haenk. i. 178 = minutiflorus.
micranthus, Schult. Mant. ii. 121 = dentatus.
microlepis, Steud. Syn. Pl. Cyp. 32 = rotundus.
microcarpus, Boeck. in Bremen Abh. vii. (1880) 37 = flavidus.
microcephalus, R. Br. Prod. 215.—Austral.
microcephalus, Naves, in Blanco, Fl. Philipp. ed. III. App. 804 = Mariscus microcephalus.
microcephalus, Nees, ex Kunth, Enum. Pl. ii. 43 = surinamensis.
microcephalus, Willd. ex Kunth, Enum. Pl. ii. 95 = Mariscus Dregeanus.
microdontus, Torr. in Ann. Lyc. N. York, iii. (1836) 255 = Pycreus polystachyus.
microiris, Steud. Syn. Pl. Cyp. 23 = C. Iria.
microlepis, Baker, Fl. Maurit. 410 = C. Iria.
microlepis, *Boeck. in Flora*, lxii. (1879) 551.—Afr. trop.
microrhizus, *Nees, in Linnaea.* x. (1836) 131.—Afr. austr.
microstachyos, Rottl. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 125 = diffusus.
microstachyus, Vahl, Enum. ii. 315 = amabilis.
millifolius, *Poepp. & Kunth, Enum. Pl.* ii. 93.—Peruv.
minimus, *Forsk. Fl. Aegypt. Arab.* 202.—Mare Marmor.
minimus, Linn. Sp. Pl. 44 = Scirpus capillaris.
minimus, Nutt. Gen. Am. i. 35 = Pycreus polystachyus.
minimus, Roth, ex Nees, in Linnaea, x. (1835–36) 182, cum cit. falsa pro *C. minuto*, Roth.
minutus, *Thunb. Prod. Pl. Cap.* 18.—Afr. ; Austral.
minor, Steud. Syn. Pl. Cyp. 4 = Pycreus polystachyus.
minutiflorus, *Y. & C. Presl, Rel. Haenk.* i. 251.—Ins. Luzon.
minutulus, Vahl, Enum. ii. 314 = Scirpus Michelianus.
minutus, J. F. Gmel. Syst. 130, citatio falsa pro C. minimo.
minutus, Roth, Nov. Pl. Sp. 32 = Scirpus antarcticus.
mitis, Steud. Syn. Pl. Cyp. 316 = tuberosus.
modestulus, Steud. l. c. 16 = Scirpus antarcticus.
moestus, Kunth, Enum. Pl. ii. 81 = diffusus.
mollis, Poir. Encyc. vii. 247 = Mariscus Dregeanus.
monandrus, Roth, Catalecta, fasc. i. 8 = vegetus.
monocephalodes, Roxb. ex Nees, in Wight, Contrib. 92 = Cephalotes.
monocephalus, F. Muell. Fragm. viii. 271 = Kyllinga monocephala.
monocephalus, Roxb. Hort. Beng. [3] ; Fl. Ind. i. 188 = Cephalotes.
monogynus, Boeck. in Linnaea, xxxv. (1867–68) 565 = Cephalotes.
monophyllus, Vahl, Enum. ii. 359 = corymbosus.
monostachyos, Boeck. in Linnaea, xxxvi. (1869–70) 389 (Mariscí sp.) = *Sermentianus*.
monostachyus, Link, in Buch, Beschr. Canar. Ins. 138 = laevigatus.
monostachyus, Linn. Mant. ii. 180 = Fimbristylis monostachya.
monroviensis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 90.—Afr. trop.
Monti, *Linn f. Suppl.* 102.—Europ. ; Orient.

CYPERUS:—
mucronatus, Rottl. Desc. Nov. Pl. 19 = laevigatus.
mucronatus, Steud. in Zoll. Syst. Verz. Ind. Archip. 68 = Pycreus capillaris.
Muehlenbergii, Spreng. Syst. v. 221 = C. Haspan.
Muelleri, *Boeck. in Flora*, xlii. (1859) 484 [bis].—Afr. trop.
Mulm-Pulla, Schult. Mant ii. 99 = pygmaeus.
multibracteatus, Boeck. in Flora, lviii. (1875) 107 (Marisci sp.) = *ferax*.
multiceps, Hook. & Arn. Bot. Beech. Voy. 100 = *ferax*.
multiceps, Link, Hort. Berol. i. 311 = congestus.
multiflorus, Kunth, Enum. Pl. ii. 562 (err. typ.) = minutiflorus.
multiflorus, Steud. Syn. Pl. Cyp. 48, sphalm. = multifolius.
multifolius, *Kunth, Enum. Pl.* ii. 91.—Ecuador.
multimaculatus, *Boeck. in Linnaea*, xxxviii. (1874) 362. —Ind. or.; Burma.
multispicatus, Kunz. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 139 = Kurzii.
multistriatus, Boeck. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 14, lapsus pro 'multispicatus.'
multumbelliferus, Hort. ex Roem. & Schult. Syst. ii. 200 = glomeratus.
Mundtii, Kunth, Enum. Pl. ii. 17 = Pycreus Mundtii.
mundulus, *Kunth, l. c.* 74.—Bras.
musarius, Buch.-Ham. ex Wall. Cat. sub n. 3325 = Scirpus Michelianus.
Mutisii, Andrss. Galapag. Cuenen Veg. 53 = Mariscus brachystachyus.
Mutisii, Griseb. Fl. Brit. W. Ind. 567 = Mariscus Mutisii.
myriostachyus, Tenore, Fl. Nap. iii. 47 = longus.
myrsenias, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 144 (Marisci sp.).—Afr. trop.
mysurensis, Heyne, ex Wall. Cat. sub n. 3346 = cleusinoides.
nanus, Willd. Sp. Pl. i. 272 = Mariscus capillaris.
natalensis, *Hochst. ex Krauss, in Flora*, xxviii. (1845) 755.—Afr. austr.
natans, Buch.-Ham. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 84 = Cephalotes.
Naumannianus, Boeck. in Flora, lxii. (1879) 553 = maculatus.
navicularis, *Steud. Syn. Pl. Cyp.* 93.—Cuba.
Neesii, D. Dietr. Syn. Pl. i. 225 = malaccensis.
Neesii, Kunth, Enum. Pl. ii. 101 = digitatus.
neglectus, Parl. Giorn. Bot. Ital. ii. i. (1846) 181 = rotundus.
nemaculeus, Steud. Syn. Pl. Cyp. 48 = Mariscus filiformis.
nematodes, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 6 = Pycreus megapotamicus.
nemoreus, G. F. W. Mey. Prim. Fl. Esseq. 81 = sphacelatus.
nervosus, Bertol. in Bologn. Opusc. Sc. iii. (1819) 407 = esculentus.
neurotropis, Steud. in Flora, xxv. (1842) 583 = Pycreus sanguinolentus.
niger, *Ruiz & Pav. Fl. Per.* i. 47.—Peruv.
nigricans, Steud. in Flora, xxv. (1842) 584 (Pycrei sp.). —Abyss.
nigricans, With. Arr. Brit. Pl. ed. III. iii. 44 = Schoenus nigricans.
nigro-viridis, Thw. Enum. Pl. Zeyl. 344 = diffusus.
nilagiricus, Hochst. ex Steud. Syn. Pl. Cyp. 2 = Pycreus capillaris.
nilotikus, Forsk. Fl. Aegypt. Arab. 13 = articulatus.
nipponicus, *Franch. & Sav. Enum. Pl. Jap.* ii. 537.—Japon.
nitens, Retz. Obs. v. 13 = aristatus ?
nitens, Rottl. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 104 = rubicundus.
nitens, Vahl, Enum. ii. 331 = Pycreus nitens.
nitidulus, Boeck. in Linnaea, xxxvi. (1869–70) 364 (Marisci sp.) = *ferax*.
nitidus, Boeck. l. c. xxxv. (1867–68) 461 (Pycrei sp.), partim = *lanceus*.
nitidus, Lam. Tabl. Encyc. i. 145 = Pycreus sanguinolentus.
niveus, Herb. Royle, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 100 = Addisonii.
nivens, *Retz. Obs.* v. 12.—Ind. or.
nodosus, *Willd. Enum. Hort. Berol.* 79.—Am. trop.
nodulosus, F. Muell. ex Benth. Fl. Austral. vii. 388 (Marisci sp.) = *Gunnii*.
Notaeus, *Boiss. Fl. Orient.* v. 371.—Anatolia.

CYPERUS:—
Novae-Hanoverae, Boeck. in Engl. Bot. Jahrb. v. (1884) 91 (Marisci sp.) = *ferax*.
Novae-Hollandiae, Boeck. in Linnaea, xxxvi. (1869–70) 344 (Marisci sp.) = *Gunnii*.
nudicaulis, *Poir. Encyc.* vii. 240.—Madag.
nudicuinis, Steber, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 161 = Schimperianus.
nudinsculus, Nees, in Linnaea, x. (1836) 132 = tenellus.
nudus, H. B. & K. Nov. Gen. et Sp. i. 208 = C. Haspan.
nudus, Roxb. Fl. Ind. i. 187 = articulatus.
nudus, Roxb. l. c. 209 = tegetiformis.
nuerensis, *Boeck. in Flora*, lxii. (1879) 555.—Afr. trop.
nutans, Steber, ex Presl, in Oken, Isis, xxi. (1828) 271 = distans.
nutans, *Vahl, Enum.* ii. 363.—Ind. or.
Nuttallii, Beyr. ex Kunth, Enum. Pl. ii. 6 (Pycrei sp.) = *virularis*, Kunth.
Nuttallii, Eddy, in Spreng. Neue Entdeck. i. 240 = Pycreus polystachyus.
Nuttallii, *Torr. in Ann. Lyc. N. York*, iii. (1836) 252. —Am. bor.
obesus, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 211.—Mexic.
obliquus, Nees, in Wight, Contrib. 86 = pilosus.
obscurus, Nees, in Hook. Kew Journ. vi. (1854) 27 = radiatus.
obtrinalis, Steud. Syn. Pl. Cyp. 10 = Pycreus nitens.
obtusiflorus, *Vahl, Enum.* ii. 308.—Afr. austr.; Madag.
obtusus, *Willem. in Usteri, Ann. Bot.* xviii. (1796) 3. —Ins. Mauriti.
occidentalis, Torr. in Ann. Lyc. N. York, iii. (1836) 259 = erythrorhizos.
ochraceus, *Vahl, Enum.* ii. 325.—Ind. occ.
ochreatus, *Boeck. in Linnaea*, xxxvi. (1869–70) 386.— Ind. occ.
ochreoides, Steud. Syn. Pl. Cyp. 84 = Fenzelianus.
ochrocephalus, Steud. in Flora, xxv. (1842) 601 = vegetus.
ochrolucus, Boeck. in Flora, lviii. (1875) 55 = Mariscus albescens ?
octophyllus, Hochst. ex Steud. Syn. Pl. Cyp. 20 = diffusus.
odoratus, Burm. f. Fl. Ind. 20. t 8. f. 2 = exaltatus.
odoratus, Forst. f. Prod. 6 (Marisci sp.) = *ferax*.
odoratus, Kunth, Enum. Pl. ii. 67, partim = giganteus.
odoratus, Linn. Sp. Pl. 46 = malaccensis, Pycreus polystachyus.
odoratus, Osbeck, Voy. China, ed. Forster, i. 361 = rotundus.
Oerstedii, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 294 (Marisci sp.) = *Yahlii*.
officinalis, Nees, ex Gren. & Godr. Fl. Fr. iii. 359 = rotundus.
officinalis, T. Nees, Pl. Medic. Abbild. t. 25. f. B = esculentus.
oleaceus, Roxb. ex Nees, in Wight, Contrib. 81 = bulbosus.
Olfersianus, Kunth, Enum. Pl. ii. 10 = Pycreus propinquus.
olidus, Vahl, Enum. ii. 333 = Pycreus polystachyus.
oligostachyus, H. B. & K. Nov. Gen. et Sp. i. 204 = anabilis.
oliveaceus, Fenzl, ex Steud. Syn. Pl. Cyp. 33 = Fenzelianus.
olivaceus, Vahl, Enum. ii. 333 = Pycreus polystachyus.
olivaris, Targ. Tozz. in Mem. Soc. Ital. Mod. xiii. 11. (1807) 338. t. 13 = rotundus.
omatus, *Steud. Syn. Pl. Cyp.* 90.—Ins. Borbon.
ostachyus, Nees, in Mart. Fl. Bras. ii. i. 39 (Marisci sp.) = *isetus*.
ornatissimus, F. Muell. Fragm. viii. 265 = Pycreus albomarginatus.
ornatus, H. Br. Prod. 217 = procerus.
ornithopodioides, Delile, Fl. Aeg. Illustr. 50 = sanguinolentus.
orochaynus, Pers. Syn. i. 61 = Mariscus havanensis.
orthostachyus, Franch. & Sav. Enum. Pl. Jap. ii. 539 = truncatus.
oryzetorum, Steud. Syn. Pl. Cyp. 37 = exaltatus.
oxystomus, Steud. l. c. 24 = diffusus.
Ottonis, Boeck. in Linnaea, xxxvi. (1869–70) 350 (Marisci sp.) = *brunneus*.
ovatus, Linnæus, Fragm. 15 = distans.
ovuligris, Boeck. in Linnaea, xxxvi. (1871–72) 274, pro majore parte = Sieberianus, umbellatus.
ovularis, *Torr. in Ann. Lyc. N. York*, iii. (1836) 258 = Mariscus ovularis.

CYPERUS:—
owahuensis, Nees, in Meyen, Reise, ii. 119; et in Kunth, Enum. Pl. ii. 80 = Mariscus albescens.
Owanii, Boeck. in Flora, lxi. (1878) 29 (Marisci sp.). —Afr.; Austral.
oxyarcicides, Britton, in Bull. Torrey Bot. Club, xi. (1884) 86 (Marisci sp.) = *macrocephalus*.
oxylepis, *Nees, in Linnaea*, ix. (1834) 265.—Guian.
oxystachys, Nees, ex Boeck. in Linnaea, xxxvi. (1869–70) 380 = longus.
pachythinus, *Nees, l. c.* xxxv. (1867–68) 545.—Ind. or.
pallescens, Boiss. Fl. Orient. v. 375 = Fenzelianus.
pallescens, Desf. Fl. Atlant. i. 45. t. 9 = rotundus.
pallescens, Poir. Encyc. vii. 260 = longus.
pallescens, Sieber, ex Kunth, Enum. Pl. ii. 61 = esculentus.
pallidus, *Kit. ex Kanitz, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 505.—Banat.
pallidus, Link, Jahrb. i. iii. 86 = Mariscus albescens.
pallidus, Nees, in Wight, Contrib. 79 = platystylis.
pallidus, *Semi. Cat. Plant. Epic.* 33. 31 = esculentus.
Pangorei, Retz. Obs. iv. 10 = tuberosus.
Pangorei, Rottb. Desc. Nov. Pl. 31; Prog. 18 = corymbosus.
Pangorei, Roxb. Hort. Beng. 6; Fl. Ind. i. 202 = malaccensis.
Pangorei, Thw. Enum. Pl. Zeyl. 344 = tegetum.
Pangorei, Vahl, ex Kunth, Enum. Pl. ii. 59 = rotundus.
paniceus, Boeck. in Linnaea, xxxvi. (1869–70) 381 = Mariscus paniceus, cyperinus, pictus.
paniciformis, Franch. & Sav. Enum. Pl. Jap. ii. 537 = C. Iria.
panicoides, Lam Tabl. Encyc. i. 146 = C. Iria.
paniculatus, Balb. ex Kunth, Enum. Pl. ii. 63 = sphacelatus.
paniculatus, Blanco, Fl. Filip. ed. I. 32 = Kyagrostis plumosa.
paniculatus, D. Don, Prod. Fl. Nep. 29.—Nepal.
paniculatus, *Hook. & Arn. Bot. Beech. Voy.* 99.—Ins. Sandvic.
paniculatus, Rottb. Desc. Nov. Pl. 40; Prog. 22 = Pycreus polystachyus.
pannonicus, *Jacq. Fl. Austr.* v. 29. t. 6.—Hungar.; Reg. Caucas.
pannonicus, Willd. ex Kunth, Enum. Pl. ii. 18 = laevigatus.
papyraceus, Crantz, Inst. i. 411 = C. Papyrus.
papyroides, Poir. Encyc. vii. 272 = aequalis.
Papyrus, *Linn. Sp. Pl.* 47.—Sicil.; Syria; Afr. trop. paradoxus, *Steud. Syn. Pl. Cyp.* 8.—Ins. Borbon.
parabasis, Mart. & Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 8 = Pycreus flavescens.
paraguayensis, *C. B. Clarke, in Fourn. Linn. Soc.* xxi. (1884) 90.—Paraguay.
Paramatta, Balb. ex Kunth, Enum. Pl. ii. 81 = planifolius.
Paramatta, Mart. Cat. Hort. Erlang. 34 = congestus.
paramattensis, Link, Enum. Hort. Berol. 46 = congestus.
parciflorus, Link, Jahrb. i. iii. 85 = diffusus.
parciflorus, Heyne, ex Wall. Cat. sub n. 3348 = exaltatus.
parciflorus, Muhl. Desc. Gram. 19 = dentatus.
parciflorus, Nees, in Wight, Contrib. 87 = C. Iria.
parciflorus, Vahl, Enum. ii. 352 = Mariscus albescens.
parculus, Steud. Syn. Pl. Cyp. 4 (Pycrei sp.) = *gracilescus*.
parvus, Boeck. in Linnaea, xxxvi. (1869–70) 397 (Marisci sp.) = *Michenianus*.
parvus, Hassk. Cat. Hort. Bogor. Alt. 94 = Kyllinga Pumilio.
patens, Vahl, Enum. ii. 334 = Pycreus nitens.
patuliflorus, Boeck. in Linnaea, xxxv. (1867–68) 478 = Pycreus albomarginatus.
patulus, Bich. Fl. Taur. Cauc. iii. 47 = longus.
patulus, Kit. ex Host, Gram. iii. 49. t. 74 = glaber.
patulus, Willd. ex Link, Enum. Hort. Berol. i. 46 = congestus.
pauciflorus, Steud. Syn. Pl. Cyp. 84 = pilosus.
Pauper, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 476 (Pycrei sp.).—Abyss.; Bay. Hiesal.
Pauper, Pill. in Acad. Univ. Chil. (1872) 388 (Pycrei sp.).—Chili.
pectinatus, Stills. & Bol. ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 384 = malaccensis.
pectinatus, Boeck. Hort. Bonp. [81]; Fl. Ind. 100 = malaccensis.
pectinellus, Retz. ex Boeck. in Linnaea, xxxvi. (1869–70) 360 = Mariscus Jacquemoni.

CYPERUS:—
pectinatus, Vahl, Enum. ii. 396 = radicalis.
pectiniformis, Schult. Mant. ii. 135 = compressus.
pedatus, [Nees in] Meyen, Reise, i. 491; et ex Kunth, Enum. Pl. ii. 85 = C. Haspan.
pedunculosus, *F. Muell. Progn.* viii. 266.—Austral.
pelophilus, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1884) 129 (Pycrei sp.).—Afr. trop.
pennatus, Boeck. in Linnæa, xxxvi. (1869–70) 404 (Mariscí sp.) = *ferax*.
pennatus, Lam. Tabl. Encyc. i. 144 = Mariscus albescens.
pennatus, Steud. in Zoll. Syst. Vers. Ind. Archip. 62 = pilosus.
pentadactylus, Willd. partim, ex Kunth, Enum. Pl. ii. 12 = Pycreus polystachyus.
pentadactylus, Willd. i. c. 50 = exilis.
permutatus, Boeck. in Linnæa, xxxv. (1867–68) 477 = macranthus.
pernambucensis, *Steud. Syn. Pl. Cyp.* 42 (sphalm. *fernambucensis*).—Bras.
perpusillus, *Boeck. in Linnæa*, xxxv. (1867–68) 499.—Mexic.
persicus, Boiss. Diagn. Ser. I. xiii. 39 = arenarius.
pertenuis, Boj. Hort. Maurit. 379 = tuberosus.
pertenuis, Roxb. Hort. Beng. [5]; Fl. Ind. i. 195 = scariosus.
Pervillei, Boeck. in Linnæa, xxxviii. (1874) 359 (Pycrei sp.).—Madag.
Pes-avium, *Bertol. Misc. Bot.* viii. 30. t. 3.—Am. bor.
Peteralanus, *Boeck. in Peters, Reise Mossamb. Bot.* 537.—Afr. trop.
petraeus, Hochst. ex Steud. Syn. Fl. Cyp. 29 = rubicundus.
phaeocephalus, Griseb. in Goett. Abh. xix. (1874) 264 (Mariscí sp.).—Reg. Argent.
philippensis, *y. & C. Presl, Rel. Haenk.* i. 174.—Ins. Luzon.
phleoides, Bleb. ex Georgi, Beschr. Russ. Reich. iii. 674 = glomeratus.
phleoides, H. Mann, in Proc. Am. Acad. vii. (1867) 308 = Mariscus phleoides.
phleoides, Seem. Fl. Vit. 319 (Mariscí sp.) = *ferax*.
phymatodes, Muhl. Desc. Gram. 28 = esculentus.
picrus, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 268 = usitatus.
picus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 200 (Pycrei sp.).—Mexic.
picrus, Boeck. in Linnæa, xxxv. (1867–68) 455, nota err. typ. = præc.
pictus, Steud. Syn. Pl. Cyp. 48 (Mariscí sp.).—Austral.
pictus, Tenore, Fl. Nap. iii. 47 = glaber.
pictus, Wall. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 33 = Mariscus pictus.
pilosus, Baker, Fl. Maurit. 413 = polystachyus.
pilosus, *Vahl, Enum.* ii. 264.—Geront. trop.
pinnatus, A. Rich. Sert. Astrol. p. vii = Mariscus albescens.
pistaloïdes, Steud. Syn. Pl. Cyp. 40 = pilosus.
planiculmis, *Steud. l. c.* 22.—Bras.
planifolius, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 (Mariscí sp.).—Ind. occ.
platyculmis, R. Br. Prod. 214 = C. Haspan.
platyphyllus, *Roem. & Schult. Syst.* ii. 876.—Zeylan.
platystachyus, Griseb. Fl. Brit. W. Ind. 567 = Mariscus flabelliformis.
platystylis, *R. Br. Prod.* 214.—Austral.; Ind. or.
plenus, Heyne, ex Wall. Cat. sub n. 5333 = Pycreus polystachyus.
pleurranthus, Nees, in Wight, Contrib. 78 = laevigatus.
poaeformis, Pursh, Fl. Am. Sept. i. 50 = Pycreus flavescens.
podocarpus, *Boeck. in Flora*, lxii. (1879) 551.—Afr. trop.
poecilos, *Desv. ex Hamb. Prod. Pl. Ind. Occ.* 12.—Ind. occ.
Poeppigii, Kunth, Enum. Pl. ii. 90 = Mariscus castaneus.
Polleni, Steud. Syn. Fl. Cyp. 40 = digitatus.
Polretii, Roem. & Schult. Syst. ii. 228 = alopecuroides.
polycephalus, Lam. Tabl. Encyc. i. 147 = C. Luzulæ.
polycephalus, Link, Enum. Hort. Berol. i. 46 = congestus.
polyphyllus, *Vahl, Enum.* ii. 217.—Afr. trop.
polystachyus, Spreng. ex Kunth, Enum. Pl. ii. 24 = compressus.
polystachyus, Boeck. in Linnæa, xxxv. (1867–68) 478, partim = mirabilis.
polystachyus, Jungh. l. c. vi. (1851) 23 (Pycrei sp.) = *flavus*.
polystachyus, Nees, ex Boeck. l. c. xxxvi. (1869–70) 368 = triformis.

CYPERUS:—
polystachyus, *Rottb. Descr. Nov. Pl.* 39; *Prog.* 21.—Comosp. trop.
polystachyus, Willd. ex Kunth, Enum. Pl. ii. 88 (Mariscí sp.) = strigosus.
ponentinus, Cav. Descr. 39, sphalm. = pannonicus.
pralongatus, Steud. Syn. Pl. Cyp. 46 (Mariscí sp.).—Texas.
praemorsus, *Boeck. in Flora*, xlii. (1859) 486 [bis].—Jamaic.
prasinus, *Kunth, Enum. Pl.* ii. 31.—Afr. austr.
pretensis, *Boeck. in Linnæa*, xxxviii. (1874) 364.—Abyss.
Prescottianus, Hook. & Arn. Bot. Beech. Voy. iii. 100 (Mariscí sp.) = *ferax*.
Presslii, A. Dietr. Sp. Pl. ii. 211 = Androtrichum polycephalum.
Presslii, Parl. Fl. Ital. ii. 40 = longus.
Princeps, Kunth, Enum. Pl. ii. 66 = comosus.
Prionleyi, Britton, in Bull. Torrey Bot. Club, xii. (1885) 7 = Mariscus tetragonus.
prionodes, Steud. Syn. Pl. Cyp. 22 = margiuatus.
prionotropis, *Steud. in Flora*, xxv. (1842) 601.—Chili.
procerulus, Nees, in Wight, Contrib. 82 = tuberosus.
procerus, Bleb. ex Georgi, Beschr. Russ. Reich. iii. iv. 674 = longus.
proceras, *Rottb. Descr. Nov. Pl.* 29.—Ind. or.; Aegypt.
procerus, Roxb. Hort. Beng. 6; Fl. Ind. i. 203, partim = flavus puncticulatus.
procerus, Vahl, ex Kunth, Enum. Pl. ii. 72 = rotundus.
productus, Wright, in Sauv. Fl. Cub. 172 = Mariscus flavus.
prolifer, Lam. Tabl. Encyc. i. 147 = aequalis.
proliferus, Nees, in Linnæa, vii. (1832) 494 = Comosp. montevidensis.
proliferus, Thunb. Prod. Pl. Cap. 18 = Scirpus Holoschoenus.
prolixus, Balb. ex Kunth, Enum. Pl. ii. 67 = digitatus.
prolixus, *H. B. & K. Nov. Gen. et Sp.* i. 206.—N. Granat.
propinquus, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 7 = Pycreus propinquus.
protractus, Steud. Nom. ed. I. 257, sphalm. = fuscus.
proteinolepis, Boeck. in Flora, lxii. (1879) 549 = rotundus.
proteinolepis, Steud. Syn. Pl. Cyp. 15 = conglomeratus.
protractus, Delile, Fl. Aeg. Illustr. 50 = fuscus.
protractus, Link, Enum. Hort. Berol. i. 45 = difformis.
proximus, Steud. Syn. Pl. Cyp. 19 = flabelliformis.
pseudo-bromoides, Boeck. in Linnæa, xxxv. (1867–68) 464 = *latespicatus* et Pycreus sanguinolentus.
pseudo-giganteus, Steud. Syn. Pl. Cyp. 39 = comosus.
pseudoretiguosus, *Steud. l. c.* 46.—Afr. trop.
pseudovegetus, Steud. l. c. 34 = C. Luzulæ.
psilostachys, Steud. l. c. 47 = distans.
pubescens, *y. & C. Presl, Rel. Haenk.* i. 181.—Calif.
pubescens, Steud. Syn. Pl. Cyp. 50 (Fimbristylidis sp. ?).—Ins. Borbon.
pubisetus, Steud. l. c. 16 (Fimbristylidis sp. ?).—Venezuela.
pubisquama, *Steud. l. c.* 20.—Ins. Philipp.
pugioniformis, A. Dietr. Sp. Pl. ii. 230 = pygmæus.
pulchellus, R. Br. Prod. 213 = leucocephalus.
pulcher, D. Don, Prod. Fl. Nep. 38 = Wallichianus.
pulcher, Spreng. ex Kunth, Enum. Pl. ii. 34 = albostriatus.
pulcher, *Thunb. Prod. Pl. Cap.* 18.—Afr. austr.
pulcherrimus, *Willd. ex Kunth, Enum. Pl.* ii. 35.—Ind. or.; Malays.
pulvinulus, Nees & Meyen, in Wight, Contrib. 74 = Pycreus nitens.
Pumilio, Nees, in Linnæa, ix. (1834) 384 = tenellus.
Pumilio, Steud. Syn. Pl. Cyp. 16 = Scirpus cernuus.
pumilus, Linn. Cent. Pl. ii. 6; Amoen. Acad. iv. 302 = Pycreus nitens.
pumilus, Rottb. Descr. Nov. Pl. 29. t. 9. f. 4 = Pycreus sanguinolentus.
pumetetifolius, Steud. Syn. Pl. Cyp. 27 = Mariscus rufus.
punctatus, Lam. Tabl. Encyc. i. 144 = Scirpus pauciflos.
punctatus, Roxb. Hort. Beng. [5]; Fl. Ind. i. 198 = Pycreus nitens.

CYPERUS:—
puncticulatus, Altch. Punjab Pl. 189 = Monti.
puncticulatus, Vahl, Enum. ii. 348 = Pycreus puncticulatus.
punctiviae, Schrad. ex Boeck. in Linnæa, xxxv. (1867–68) 567 = textilis.
pungens, Boeck. l. c. 587 = conglomeratus.
pungens, Hort. Berol. ex Kunth, Enum. Pl. ii. 32 = textilis.
purpurascens, Vahl, Enum. ii. 359 (Mariscí sp.).—Ins. S. Crucis.
Purshii, Roem. & Schult. Syst. ii. 177 = aristata.
pusillus, *Vahl, Enum.* ii. 303.—Ind. or.
pusillus, Wight, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 87, 90 = castaneus.
pustulatus, *Vahl, Enum.* ii. 341.—Afr. trop.
pycnocephalus, Steud. Syn. Pl. Cyp. 5 = Pycreus propinquus.
pycnostachyus, Kunth, Enum. Pl. ii. 77 = Mariscus pycnostachyus.
pygmaeus, Cav. in Anal. Cienc. Nat. iii. (1801) 7 = Scirpus Michelianus.
pygmæus, Lam. Tabl. Encyc. i. 143 = tenellus.
pygmæus, Nutt. in Am. Phil. Trans. N. S. v. (1837) 142 = aristatus.
pygmæus, Retz. Obs. iv. 9 = pusillus.
pygmæus, *Kotth. Descr. Nov. Pl.* 20.—Comosp. trop.
pyrophilus, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 189 = pilosus.
quadriflorus, *Boeck. in Linnæa*, xxxvi. (1869–70) 367.—Abyss.
quinqueflorus, Hochst. ex Steud. Syn. Pl. Cyp. 37 = digitatus.
quitensis, Spreng. Syst. i. 224 = amabilis.
racemosus, Balz. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 142 = elenaënsis.
racemosus, Heyne, ex Boeck. in Flora, lxii. (1879) 555 = exaltatus.
racemosus, Retz. Obs. vi. 20 = elatus.
racemosus, Tenore, Prod. Fl. Nap. p. viii. = longus.
racemosus, Wight, in Wall. Cat. n. 3356 = Mariscus albescens.
radians, *Nees & Meyen, in Linnæa*, ix. (1834) 285, nomen; *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 63.—Burma; China.
radiatus, *Vahl, Enum.* ii. 269.—Ind. or.; Burma.
radicans, Kunth, Enum. Pl. ii. 95, err. typ. = radians.
radiceus, Reichb. ex Kunth, l. c. 26 = simplex.
radiceus, Sibth. & Sm. Fl. Graec. Prod. i. 30 = rotundus.
Ranko, Steud. Syn. Pl. Cyp. 315 = Scirpus perpusillus.
raphistachys, Link, Jahrb. i. iii. 87 (Mariscí sp.) = *ferax*.
rariscinus, Steud. Syn. Pl. Cyp. 16 (Fimbristylidis sp.).—Abyss.
recurvatus, Willd. ex Kunth, Enum. Pl. ii. 22 = cuspidatus.
recurvus, *Vahl, Enum.* ii. 310.—Afr. trop.
recurvus, Willd. ex Kunth, Enum. Pl. ii. 210 (Fimbristylidis sp.).—Afr. trop.
reductus, *Hochst. ex Boeck. in Linnæa*, xxxv. (1867–68) 580.—Abyss.
reflexus, Hort. Par. ex Regel, Ind. Sem. Hort. Petrop. (1866) 70 = juncoides.
reflexus, *Vahl, Enum.* ii. 42.—Reg. Argent.; Chili.
refractus, Engelm. ex Boeck. in Linnæa, xxxvi. (1869–70) 369 (Mariscí sp.) = *lancastriensis*.
Rehmanni, Boiss. Fl. Orient. v. 364 (Pycrei sp.).—As. Min.
Reinschii, *Boeck. in Flora*, lxv. (1882) 13.—Ins. Comor.
repens, Ell. Sketch, i. 68 = esculentus.
reptans, Boeck. in Linnæa, xxxv. (1867–68) 485 = laevigatus.
reptans, Boj. ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 281 = Pycreus Mundtii.
resinosus, Hochst. in Flora, xxv. (1842) i. Beibl. 134, nomen; Steud. Syn. Pl. Cyp. 28 = C. Iria.
retroflexus, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 9 = Mariscus uniflorus.
retrofractus, Engelm. ex Steud. Syn. Pl. Cyp. 47 (Mariscí sp.) = *lancastriensis*.
retrofractus, Torr. Fl. N. York, ii. 344 = Mariscus retrofractus.
retrofractus, Willd. ex Boeck. in Linnæa, xxxvi. (1869–70) 378 = Mariscus ovularis.

retrorsus, Chapm. in Coult. Bot. Gaz. iii. (1878) 175 (Mariscl sp.).—Am. bor.

retusus, Nees, in Linnaea, ix. (1834) 388 = esculentus.

retusus, A. Rich. Tent. Fl. Abyss. ii. 485 = Pycreus albomarginatus.

Retzii, Nees, in Wight, Contrib. 32 = tuberosus.

Retzii, Poir. Encyc. vii. 255 = albidus.

rhaphistachys, Rumph. & Bonpl. ex Kunth, Enum. Pl. ii. 68 (Mariscl sp.) = *ferax*.

rhenanus, Schrad. ex Kunth, l. c. 60 = longus.

rhizophyllus, *Hochst. ex Krauss, in Flora*, xxviii. (1845) 756.—Afr. austr.

Richardi, Steud. Syn. Pl. Cyp. 8 = Kyllinga macrocephala.

rigens, J. & C. Presl, Rel. Haenk. i. 170 (Mariscl sp.).—Peruv.

rigidifolius, *Steud. in Flora*, xxv. (1842) 593.—Abyss.

rigidulus, *Vahl, Enum.* ii. 342.—Ind. or.

rigidus, Vahl, l. c. 309 = maritimus.

riparius, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 88 = C. Haspan.

riparius, Kunth, Enum. Pl. ii 6 (Pycrei sp.).—Am. bor.

riparius, Steud. in Lechl. Berb. Am. Austr. 66 = laevigatus.

robustus, *Kunth, Enum. Pl.* ii. 41.—Bras.

Rostkoviі, Kunth, l. c. 58 = tuberosus.

Rothiі, Boeck. in Flora, lxv. (1882) 13 (Mariscl sp.). —Abyss.

Rothianus, Schult. Mant. ii. 98 = Scirpus antarcticus.

rotundus, Benth. Fl. Austral. vii. 379, partim = bulbosus, stoloniferus, Zollingeri.

rotundus, Benth. Pl. Hartw. 23 = esculentus.

rotundus, Hook. f. in Trans. Linn. Soc. xx. (1847) 177 = lutescens.

rotundus, Kunth, Enum. Pl. ii. 58, partim = bulbosus. rotundus, *Linn. Sp. Pl.* 45.—Cosmop.

rotundus, Miq. Fl. Ind. Bat. Suppl. 260, 600 = Zollingeri.

Roxburghianus, Presl, in Oken, Isis, xxi. (1828) 272 = laevigatus.

Roxburghii, A. Dietr. Sp. Pl. ii. 214 = laevigatus.

Roxburghii, Nees, in Wight, Contrib. 84 = platyphyllus.

Royleі, Arn. in Wight, l. c. 85, in nota = radiatus.

rubescens, Kunth, Enum. Pl. ii. 27, sphalm. = glaber.

rubicundus, *Vahl, Enum.* ii. 308.—Ind. or. ; Afr. trop.

rubicundus, Willd. ex Link, Jahrb. i. iii. 37 = rotundus.

rubiginosus, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 176.—Ins. Galap.

Rudloі, *Boeck. in Flora*, lxv. (1882) 12.—Bras.

rufescens, Schrad. ex Steud. Syn. Pl. Cyp. 27 = Mariscus rufus.

rufescens, Buckl. in Proc. Ac. Sc. Philad. 1862 (1863) 9 = lutescens.

rufescens, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 195.—Texas.

rufidulus, *Steud. Syn. Pl. Cyp.* 316.—Am.

rufinus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 229 (Mariscl sp.) = *Michaxianus*.

rufus, *R. Br. & K. Nov. Gen. et Sp.* i. 210.—Am. austr.

rufus, Nees, ex Kunth, Enum. Pl. ii. 98 = Mariscus microcephalus.

Ruizianus, Boeck. in Linnaea, xxxvi. (1869–70) 860 = Mariscus Jacquinii.

rupestris, Kunth, Enum. Pl. ii. 52.—Afr. austr.

Rusbyi, Britton, in Bull. Torrey Bot. Club, xi. (1884) 39 = Schweinitzii.

sabulicolus, *Poir. in Trans. Linn. Soc.* Ser. II. (1884) 126.—Afr. trop.

sabulosus, Mart. & Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 10 = Pycreus sabulosus.

Salzmanni, Steud. Syn. Pl. Cyp. 15 = cuspidatus.

sanguinalis, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 39 (Mariscl sp.) = *sanguinео-fuscus*.

sanguineo-ater, Boeck. in Engl. Bot. Jahrb. i. (1881) 263 = Mariscus atrocaninginosus.

sanguineo-fuscus, Hook. f. Fl. Tasm. ii. 80. t. 139 (Mariscl sp.).—Austral. ; Tasmania.

sanguineo-rufescens, Nees, in Ann. Nat. Hist. vi. (1841) 48 = lucidus †

sanguineus, Balb. ex Steud. Nom. ed. II. i. 472 (Mariscl sp.) = *purpurascens*.

sanguinolentus, Vahl, Enum. ii. 351 = Pycreus sanguinolentus.

Santenesi, Rottb. Descr. Nov. Fl. 41 ; Prog. 19 = C. Iria.

Santonici, Rottl. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 149 = eleusinoides.

scaber, Boeck. in Flora, lviii. (1875) 107 ; tcl. (1878) 31 = Mariscus scaber.

scaber, Lag. ex Kunth, Enum. Pl. ii. 78 = congestus.

scabrirunus, Nees, in Linnaea, xix. (1847) 697 (Mariscl sp.) = *Maximae*.

scariosus, *R. Br. Prod.* 215.—Ind. or. ; Austral.

schaenoides, Poir. Encyc. Suppl. v. 185 = Fimbristylis squarrulosa.

Schaffneri, *Boeck. in Flora*, lxi. (1878) 20.—Mexic.

Schimperianus, *Steud. Syn. Pl. Cyp.* 84.—Afr. trop.

schoenoides, Griseb. Spicil. Fl. Rumel. ii. 421 = aegyptiacus.

schoenoides, Steud. Syn. Pl. Cyp. 39 = spicigerus.

Schomburgkianus, Nees, in Hook. Journ. Bot. ii. (1840) 398 = leucanthus.

Schottianus, Cat. Sem. Hort. Vindob. (1836) = seq.

Schottii, Hort. Berol. ex D. Dietr. Syn. Pl. i. 220 = prolixus.

Schraderianus, Mart. ex Steud. Syn. Pl. Cyp. 38 = Schweinfurthianus.

Schweinfurthianus, *Boeck. in Flora*, lxii. (1879) 553.—Afr. trop.

Schweinitzii, *Torr. in Ann. Lyc. N. York*, iii. (1836) 276.—Am. bor.

scirpoides, R. Br. in Salt, Abyss. App. p. lxii = Fresenii.

scirpoides, J. & C. Presl, Rel. Haenk. 178 = diffusus.

scirpoides, *Vahl, Enum.* ii. 311.—Afr. trop.

scoparius, Poir. Encyc. vii. 258 = latifolius.

scopiformis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Pycreus polystachyos.

Seemannianus, Boeck. in Linnaea, xxxvi. (1869–70) 890 (Mariscl sp.).—Ins. Fiji.

Sellowianus, Boeck. in Linnaea, xxxv. (1867–68) 446 (Pycrei sp.).—Bras.

Sellowii, Link, Hort. Berol. i. 307 = reflexus.

semiochraceus, Boeck. in Linnaea, xxxvii. (1874) 367 = fastigiatus.

semidivisus, Steud. Syn. Pl. Cyp. 36 = radiatus.

seminudus, Roxb. Hort. Beng. [5] ; Fl. Ind. i. 187 = corymbosus.

seminudus, Zoll. & Moritzi, Syst. Verz. Zoll. 96 = C. Iria.

semiochraceus, *Boeck. in Flora*, lxi. (1878) 29.—Mexic.

semiradicifloras, *Boeck. l. c.* xlii. (1859) 434 [bis].—Hab. ?

semiteres, Heyne, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 49 = Pycreus capillaris.

semitribrachiatus, *Boeck. in Linnaea*, xxxvi. (1869–70) 866.—Ind. occ.

semitrifidus, *Schrad. Anal. Fl. Cap. Cyp.* 6.—Afr. austr.

septentrionalis, Steud. Syn. Pl. Cyp. 46 = Mariscus microcephalus.

serotinus, Rottb. Descr. Nov. Pl. 91 ; Prog. 18 = Monti.

Serra, A. Rich. Tent. Fl. Abyss. ii. 479 = rubicundus.

serrulatus, *Vahl, Enum.* ii. 300.—Madag.

serrulatus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 385.—Am. bor.

serrularinus, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 213.—Nicarag.

seslerioides, *H. B. & K. Nov. Gen. et Sp.* i. 209.—Ind. occ.

setaceus, Retz. Obs. v. 10 = Elecocharis chaetaria.

setaceus, Walter, Fl. Carol. 49 = Pycreus flavescens.

setifolius, *D. Don, Prod. Fl. Nep.* 26.—Nepal.

setifolius, Torr. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 198 (Mariscl sp.) = *Grayii*.

setigerus, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 284.—Am. bor.

sexangularis, Fenzl, ex Steud. Syn. Pl. Cyp. 19 = flabelliformis.

sexangularis, *Nees, in Linnaea*, ix. (1834) 284.—Afr. austr.

sexfloras, *R. Br. Prod.* 215.—Austral.

Siberi, Kunth, Enum. Pl. ii. 96 = Mariscus frivus.

Sieberianus, Link, Hort. Berol. i. 213 = esculentus.

Sieberianus, Spreng. Syst. iv. Cur. Post. 29 = surinamensis.

sillebensis, *Nees, in Wight, Contrib.* 79.—Ind. or. ; Burma.

sillebensis, Thw. Enum. Pl. Zeyl. 343 = pulcherrimus.

simplex, *H. B. & K. Nov. Gen. et Sp.* i. 207.—Am. austr.

sinensis, Debeaux, in Actes Linn. Soc. Bord. xxxi. (1877) t. 2 ; xxxii. (1878) 26 = radiatus.

Smithianus, *Ridl. in Journ. Bot.* xxii. (1884) 15.—Afr. trop.

Smithii, Schrad. Anal. Fl. Cap. Cyp. 12, in adn. = textilis.

solidifolius, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 499. —Madag.

solidus, Kunth, Enum. Pl. ii. 76 (Mariscl sp.) = tabularis, usitatus.

solutus, Steud. Syn. Pl. Cyp. 14 = cuspidatus.

Sonderi, J. A. Schmidt, Beitr. Fl. Cap Verd. Ins. 162 = polystachyus.

soongoricus, *Ker. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 859.—Sibir.

sordidus, J. & C. Presl, Rel. Haenk. i. 171 = virens.

sorostachys, Boeck. in Linnaea, xxxv. (1867–68) 588 = leucocephalus.

sorongoensis, J. & C. Presl, Rel. Haenk. i. 851 = diffusus.

Soyauxii, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 501.—Afr. trop.

spadiceus, Heyne, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 152 = procerus.

spadiceus, Lam. Tabl. Encyc. i. 147 = stoloniferus.

spaniophyllus, Steud. Syn. Pl. Cyp. 21 = malaccensis.

sparteus, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 19 = reflexus.

spathaceus, Linn. Syst. ed. XII. 735 = Dulichium spathaceum.

speciosus, Torr. Bot. Mex. Bound. 226 = californicus.

speciosus, Vahl, Enum. ii. 364 (Mariscl sp.) = *Michauxianus*.

spectabilis, Boeck. in Linnaea, xxxv. (1867–68) 605 (Mariscl sp.) = *Manimae*.

spectabilis, Spreng. Nov. Prov. 15 ; Link, Hort. Berol. i. 218 = conjunctus.

sphaeletatus, Presl, in Oken, Isis, xxi. (1828) 271 = rotundus.

sphaeolatus, *Rottb. Descr. Nov. Pl.* 26 ; *Prog.* 21.—Afr. et Am. trop.

sphaerocephalus, Vahl, Enum. ii. 310 = obtusiflorus.

sphaeroleptis, *Boeck. in Linnaea*, xxxv. (1867–68) 609. —Am. bor.

sphaerosperma, *Schrad. Anal. Fl. Cap. Cyp.* 8.—Afr. austr.

sphaerostachys, Link, Jahrb. i. iii. 39 = C. Luzulae.

sphaeroides, Poir. Encyc. Suppl. v. 187 = C. Sporobolus.

spicato-capitatus, Steud. Syn. Pl. Cyp. 27 = Mariscus rufus.

spicatus, *Y. & C. Presl, Rel. Haenk.* i. 178.—Ins. Luzon.

spicigerus, *Vahl, Enum.* ii. 300.—Bras.

spiculosus, Reichb. Cat. Sem. Hort. Dresd (1828) = congestus.

*spinulifer*eus, Boeck. in Linnaea, xxxv. (1867–68) 465 = Pycreus angulatus.

spinulosus, Roxb. Hort. Beng. 6 ; Fl. Ind. i. 203 = Mariscus microcephalus.

spongioso-coriaceus, Boeck. in Flora, xlii. (1859) 441 = Mariscus albescens.

Sporobolus, *R. Br. Prod.* 215.—Austral.

spretus, Steud. Syn. Pl. Cyp. 21 = denudatus.

squalidus, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 198.—Costa Rica.

squamulatus, Steud. Syn. Pl. Cyp. 49 = distans.

squarrosus, Afzel. ex Kunth, Enum. Pl. ii. 25 = glaber.

squarrosus, Kotschy, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 45 = Pycreus nitens.

squarrosus, *Linn. Cent. Pl.* ii. 6 ; *Amoen. Acad.* iv. 308. —Cosmop. trop.

squarrosus, F. Muell. Fragm. viii. 262 = cuspidatus.

squarrosus, Roxb. Hort. Beng. [31] ; Fl. Ind. i. 190 = pygmaeus.

stachyoborus, C. Koch, in Linnaea, xxi. (1848) 632 = Pycreus retusus.

St-Crucei, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 233 (Mariscl sp.) = *ferax*.

stellatus, Link, Jahrb. i. iii. 44 = C. Haspan.

stellatus, *Novorbac, ex Thou. Prod. Phyt. h, in Mélang.* (1811), nomen.—Madag.

CYPERUS:—
stellatus, Rudge, Pl. Gui. 17. t. 90 = giganteus!
stenolepis, Torr. in Ann. Lyc. N. York, iii. (1836) 268 (Mariscl sp.).—Am. bor.
stenostachyus, *Benth. Fl. Austral.* vii. 280.—Austral.
Steudelianus, Boeck. in Linnaea, xxxvi. (1869–70) 382 = Mariscus Siberianus.
stigmatosus, Steud. Syn. Pl. Cyp. 86 = Mariscus albescens.
stolonifer, Willd. ex Kunth, Enum. Pl. ii. 59 = rotundus.
stoloniferus, Nees, in Wight, Contrib. 81, partim = bulbosus.
stoloniferus, *Rota. Obs.* iv. 10.—Geront. trop.
straminicus, Link, Jahrb. i. iii. 84 = Pycreus flavescens.
straminus, Nees, in Wight, Contrib. 74 (Pycrei sp.).—Ind. or.
striatulus, Vahl, Enum. ii. 311 = obtusiflorus.
striatus, *Rusa & Pav. Fl. Per.* i. 47.—Peruv.
strictus, Lam. Tabl. Encyc. i. 146 = Pycreus capillaris.
strictus, Rottl. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 54 = Pycreus-polystachyus.
strictus, Wight, ex C. B. Clarke, l. c. 46 = Pycreus pumilus.
strigosus, Hook. & Arn. Bot. Beech. Voy. 99 (Mariscl sp.) = *ferax*.
strigosus, Linn. Sp. Pl. 47 (Mariscl sp.).—Am. bor.
strigosus, Schlecht. ex Kunth, Enum. Pl. ii. 59 (Mariscl sp.) = *ferax*.
strigosus, Wight, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 51 = Pycreus polystachyus.
strigosus, Willd. Enum. Hort. Berol. 74 = congestus.
stuppeus, Forst. f. Prod. 89 = Mariscus albescens.
subalatus, Boeck. in Linnaea, xxxvi. (1869–70) 291 = surinamus.
subaiatus, Steud. Syn. Pl. Cyp. 81 = pilosus.
subanceps, *Boeck. in Linnaea*, xxxviii. (1874) 868.—Ind. occ.
subarticulatus, Nees & Meyen, in Linnaea, ix. (1834) 285 = articulatus.
subbulbosus, Roem. & Schult. Syst. iii. 186 = congestus.
subcervinus, *Steud. Syn. Pl. Cyp.* 37.—Reg. Argent.
subfuscus, *Debeaux, in Act. Linn. Soc. Bord.* xxxi. (1877) t. 3; xxxii. (1878) 28.—China.
subnodosus, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 59 = nodosus.
subrotundus, Llanos, Fragm. 14 = diffornis.
submonostachyus, *Steud. & Yard. in Bull. Soc. Linn. Normand.* Sér. II. ix. (1876) 278, 280.—Ins. Marques.!
subulatus, Meyen, Reise, i. 108 = Mariscus Meyenianus.
subulatus, *R. Br. Prod.* 217.—Austral.
sulcinux, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 58 (Pycrei sp.).—Ins. et Afr. trop.
surinamensis, *Rottb. Descr. Nov. Pl.* 35; *Prog.* 20.—Am. trop.
surinamensis, Vahl, ex Kunth, Enum. Pl. ii. 43 = C. Lunulae.
sylvestris, *Ridl. in Trans. Linn. Soc.* Ser. II. *Bot.* ii. (1884) 184.—Afr. trop.
syriacus, Parl. in Mém. Prés. Acad. Sc. Fr. xii. (1854) 498. t. 1 = C. Papyrus.
tabina, Stend. in Lechl. Berb. Am. Austr. 56 (Mariscl sp.).—Peruv.
tabularis, Schrad. Anal. Fl. Cap. Cyp. 10 (Mariscl sp.).—Afr. austr.
tacsonsis, Nees & Meyen, Reise, i. 431; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 64 (Mariscl sp.) = *rigens*.
taltensis, *Steud. Syn. Pl. Cyp.* 37.—Ins. Taiti.
tamyskilus, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1884) 148 (Mariscl sp.).—Afr. trop.
tataricus, Lam. in Linnaea, ix. (1834) 211 (Pycrei sp.).—Sibir. ural.
tegetiformis, Benth. Fl. Austral. vii. 278 = malaccensis.
tegetiformis, *Roxb. Hort. Beng.* 6; *Arn. in Wight, Contrib.* 39.—Ins. trop.
tegetum, C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 190 (exempl. Afric.) = Schimperianus.
tegetum, *Roxb. Hort. Beng.* 6; *Fl. Ind.* i. 208.—Ind. or.; Afr. trop.
temax, *Boeck. in Linnaea*, xxxv. (1867–68) 504.—Afr. austr.
Tenuis, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 278 = Schoenus Tenuis.
tenellus, *Linn. f. Suppl.* 108.—Afr. austr.; Austral.; N. Zel.

CYPERUS:—
tenellus, J. & C. Presl, Rel. Haenk. i. 176 (Pycrei sp.) = rivularis.
tenellus, Torr. in Ann. Lyc. N. York, iii. (1836) 257 = Pycreus polystachyus.
tener, Nees & Ehrenb. ex Boeck. in Linnaea, xxxvi. (1869–70) 301 = bulbosus.
tener, *Vahl, Enum.* ii. 399 (Scirpus!).—Ind. or.
Tenerifae, Nees, in Linnaea, x. (1835–36) 121 = Meyerianus.
Tenerifae, Poir. Encyc. vii. 245 = rubicundus.
tenerrimus, Llebm. in Videnak. Selak. Skr. V. ii. (1851) 209 = humilis.
tenerrimus, *J. & C. Presl, Rel. Haenk.* i. 166.—Mexic.
Tenorianus, Schult. Mant. ii. 544 = esculentus.
Tenorii, Presl, Fl. Sicul. p. xlii = esculentus.
tenuiculmis, Steud. Nom. ed. II. i. 470 = surinamensis.
tenuicomus, Miq. in Linnaea, xix. (1847) 238 = corymbosus.
tenuiculmis, Boeck. in Flora, lxii. (1879) 554 = gracillimus.
tenuicaulis, Boeck. in Linnaea, xxxvi. (1869–70) 286 = Zollingeri.
tenuiculus, Kunth, Enum. Pl. ii. 106 (Pycrei sp.) = rivularis.
tenuiflorus, Ell. Sketch, i. 70 = erythrorhizos.
tenuiflorus, Jacq. Ic. Pl. Rar. ii. & t. 296 = longus.
tenuiflorus, *Rottb. Descr. Nov. Pl.* 30.—Ind. or.; Ins. Mauric.
tenuiflorus, Roxb. Hort. Beng. [5]; Fl. Ind. i. 199 = tuberosus.
tenuifolius, Walp. Ann. i. 898, sphalm. = rotundus.
tenuior, Lapeirn. in A. Gray, Man. Bot. N. U. St. ed. II. 493 (Mariscl sp.) = *Engelmanni*.
tenuis, Muhl. Desc. Gram. 22 = Pycreus polystachyus.
tenuis, Sw. Prod. Veg. Ind. Occ. 20 (Mariscl sp.).—Jamaica.
tenuispica, Steud. Syn. Pl. Cyp. 11 = flavidus.
tenuispicatus, *Boeck. in Linnaea*, xxxv. (1867–68) 525. —Luzulae.
tenuispiculatus, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 500.—Madag.
tephrodes, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 80 = dichromeaeformis.
teretifolius, *A. Rich. Tent. Fl. Abyss.* ii. 477.—Abyss.
teretifructus, Steud. Syn. Pl. Cyp. 5 = Pycreus capillaris.
terminalis, *Marsh, ex Steud. Syn. Pl. Cyp.* 5.—Afr. trop.
terminalis, Michx. ex Steud. Nom. ed. I. 257 = flavescens.
ternus, Link, Jahrb. i. iii. 82 = rubicundus.
tetracarpus, Boeck. in Flora, lvii. (1875) 88 (Mariscl sp.) = *Sumanianus*.
tetragonus, Ell. Sketch, i. 71 (Mariscl sp.).—Am. bor.
tetragonus, Hemsl. Biol. Centr. Am. Bot. iii. 452 = Mariscus Jacquinii.
tetraphyllus, Beauv. ex Kunth, Enum. Pl. ii. 18 = Pycreus polystachyus.
tetraphyllus, *R. Br. Prod.* 214.—Austral.
tetrastachyos, Desf. Fl. Atlant. i. 45 = rotundus.
texensis, Steud. Syn. Pl. Cyp. 9 = Pycreus polystachyus.
textilis, Sieb. Fl. Mauric. 278 = tegetum.
textilis, F. Muell. Fragm. viii. 261 = vaginatus.
textilis, *Thunb. Prod. Pl. Cap.* 18.—Afr. austr.
Thouarsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 141.—Japon.
Teymanni, Boeck. in Flora, lvii. (1875) 259 (Pycrei sp.).—Sumatra.
thermalis, Dum. Fl. Belg. 145 = longus.
Thomsoni, *Boeck. in Linnaea*, xxxvi. (1869–70) 294.—Ind. or.
Thouarsianus, Presl, in Oken, Isis, xxi. (1828) 271 = Mariscus Dregeanus.
Thouarii, Kunth, Enum. Pl. ii. 12 = polystachyus.
Thunbergii, Retzius, ex Kunth, l. c. 72 = radiatus.
Thunbergii, Vahl, Enum. ii. 371 = Mariscus riparius.
thyrsiflorus, Boeck. in Linnaea, xxxvi. (1869–70) 5 = Mariscus flabelliformis, rufus.
thyrsiformis, Jungh. in Linnaea, iv. (1831) 94 = Mariscus Jacquinii.
thyrsiformis, Llebm. in Videnak. Selak. Skr. V. ii. (1851) 216 (Mariscl sp.) = *dissitiflorus*.
tobuscensis, H. B. & K. Nov. Gen. et Sp. i. 204 = diffusus.

CYPERUS:—
tongatabuensis, Boeck. in Flora, lvii. (1875) 85 = Mariscus albescens.
torosus, Sieber, ex Boeck. in Linnaea, xxxv. (1867–68) 84 = Pycreus polystachyus.
torosus, Vahl, Enum. ii. 389 = malaccensis!
torosus, Willd. ex Kunth, Enum. Pl. ii. 60 = longus.
Torreyanus, Schult. Mant. ii. 101 = Pycreus polystachyus.
tortulanus, Vahl, ex Kunth, Enum. Pl. ii. 43 = surinamensis.
tortuosus, Koen. ex Roxb. Hort. Beng. [5]; Fl. Ind. i. 197 = Pycreus capillaris.
trachynotus, *Torr. in Bot. Mex. Bound.* 227.—N. Mexic.
trachyrhachis, Steud. Syn. Pl. Cyp. 3 = Pycreus capillaris.
trachysanthus, *Hook. & Arn. Bot. Beech. Voy.* 99.—Ins. Sandvic.
tremulus, Baker, Fl. Maurit. 409, partim = Pycreus albomarginatus.
tremulus, Poir. Encyc. vii. 264 = Pycreus retusus.
triceps, Nees, in Linnaea, xix. (1847) 697 (Mariscl sp.) = *Meninnae*.
trichodes, *Griseb. Fl. Brit. W. Ind.* 564.—Ind. occ.
trichostachyos, Benth. Fl. Austral. vii. 287 (Mariscl sp.) = *Dietrichiae*.
trifarius, Linn. Mant. ii. 180 (Fimbristylidis sp.) = *trichostachya*.
trigynus, Spreng. Syst. i. 216 = Androtrichum polycephalum.
trinervis, *R. Br. Prod.* 213.—Austral.
Trinitatis, Steud. Syn. Pl. Cyp. 26 = C. Luzulae.
triqueter, *Boeck. in Flora*, lxii. (1879) 548.—Afr. trop.
tripicatus, Boeck. l. c. lxv. (1882) 14 (Mariscl sp.).—Am. austr.
tristachyus, Boeck. in Linnaea, xxxv. (1867–68) 454 (Pycrei sp.).—Am. austr.
tristis, *Kunth, Enum. Pl.* ii. 35.—Afr. austr.
triaulcus, *D. Don, Prod. Fl. Nep.* 39.—Nepal.
truncatulus, Steud. in Flora, xxv. (1842) 581 = Pycreus silens.
truncatus, *C. A. Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* (1888) 108.—Reg. Amur.
truncatus, *A. Rich. Tent. Fl. Abyss.* ii. 487.—Abyss.
tuberiferus, Steud. Nom. ed. II. i. 474 = digitatus.
tuberosus, Boj. Hort. Maurit. 379 = stoloniferus.
tuberosus, Forsk. Fl. Aeg. Arab. Sept. i. 23 = esculentus.
tuberosus, Korth. Desc. Nov. Pl. 28; Prog. 18.—Ind. or.; Afr. bor.!
tuberosus, Roxb. Hort. Beng. [5]; Fl. Ind. i. 199 = rotundus.
tuberosus, Willd. ex Kunth, Enum. Pl. ii. 60, 93 = Mariscus microcephalus.
tunicatus, Roem. & Schult. Syst. Mant. ii. 115 = C. Haspan.
turfosus, Salzm. ex Kunth, Enum. Pl. ii. 7 = Pycreus Mundtii.
turgidulus, *C. B. Clarke, in Journ. Linn. Soc.* xxi. (1884) 180.—Ins. trop.
ubertus, Willd. ex Link, Jahrb. i. iii. 87 = rufus.
uligiuosus, Thou. ex Link, l. c. 83 = laevigatus.
umbellatus, Benth. Fl. Austral. vii. 289, partim = Mariscus albescens.
umbellatus, *Benth. Fl. Hongk.* 386 = Mariscus cyperinus.
umbellatus, Burm. f. Fl. Ind. 21. t. 9. f. 1 = C. Haspan.
umbellatus, Hemsl. Bot. Voy. Challenger, i. 11. 23 = Mariscus appendiculatus.
umbellatus, Oliver, in Trans. Linn. Soc. xxix. (1875) 166, partim = Mariscus macer.
umbellatus, Roxb. Hort. Beng. [31]; Fl. Ind. i. 205 = exaltatus.
umbellatus, Thw. Enum. Pl. Zeyl. 345 = Mariscus pictus, paniceus, cyperinus.
umbraticola, Kunth, Enum. Pl. ii. 13 = Pycreus umbrosus.
umbrosus, Nees, in Linnaea, ix. (1834) 235 = diffusus.
umbrosus, Salzm. ex Steud. Syn. Pl. Cyp. 17 = simplex.
uncinatus, R. Br. Prod. 215 = aristatus.
uncinatus, Desv. ex Ham. Prod. Fl. Ind. Occ. 12 = hamulatus.
uncinatus, *Poir. Encyc.* vii. 247.—Madag.
uncinulatus, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. i. 93.—Bras.
uniceler, *Boeck. in Kjoeb. Videnak. Meddel.* (1879–80) 24.—Am. austr.

CYPERUS:—
uniformis, *Thunb. Fl. Yavan.* 8.—Java.
uniformis, Torr. & Hook. in Ann. Lyc. N. York, iii. (1836) 431 = Mariscus uniformis.
unifolius, Boeck. in Linnaea, xxxvi. (1869–70) 374 = Mariscus havanensis.
unioloides, R. Br. Prod. 216 = Pycreus angulatus.
upolensis, Boeck. in Flora, lviii. (1875) 88 (Mariaci sp.) = *Seemannianus*.
Urvilleanus, Steud. Syn. Pl. Cyp. 16 = Scirpus lanuginus.
ustatus, *Burch. ex Roem. & Schult. Syst. Mant* ii. 477.—Afr. austr.
utriculatus, A. Rich. Ess. Fl. N. Zél. 101. t. 17 (Mariaci sp.).—N. Zel.
vaginatus, *R. Br. Prod.* 213.—Austral.
vaginatus, Link, Jahrb. i. iii. 85 = C. Hespan.
Vahlii, Boeck. in Linnaea, xxxvi. (1869–70) 398 (Mariaci sp.) = *ferax*.
Vahlii, Steud. Syn. Fl. Cyp. 48 (Mariaci sp.).—Ind. occ.
Valdiviae, Steud. l. c. 45 = Mariscus albescens.
variabilis, Salzm. ex Steud. l. c. 46 = esculentus, *ferax*.
variegatus, Boeck. in Linnaea, xxxvi. (1869–70) 387 = Mariscus bulbosaulis, platelleus, Schimperi.
variegatus, Griseb. Fl. Brit. W. Ind. 562 = Pycreus propinquus.
variegatus, H. B. & K. Nov. Gen. et Sp. i. 208 (Pycrei sp.) = *melanostachyus*.
vegetus, Salzm. ex Steud. Syn. Fl. Cyp. 5 = convallium.
vegetus, *Willd. Sp. Pl.* i. 263.—Chili.
ventricosus, R. Br. Prod. 217 = Mariscus albescens.
venustus, Boeck. in Linnaea, xxxvi. (1869–70) 816 = digitatus, auricomus.
venustus, R. Br. Prod. 217 = exaltatus.
venustus, Forst. f. Prod. 89 = Mariscus cyperinus.
venustus, Kunth, Enum. Pl. ii. 68 = digitatus, fastigiatus.
venustus, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 271 = auricomus.
verticulor, Nees, in Wight, Contrib. 78 = aristatus.
verticillatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 206 = radiatus.
vestitus, Hochst. ex Krauss, in Flora, xxviii. (1845) 755 (Mariaci sp.).—Afr. austr.
virens, *Michx. Fl. Bor. Am.* i. 28.—Am. bor.
virens, Reichb. ex Kunth, Enum. Pl. ii. 48 = surinamensis.
virens, Willd. ex Kunth, l. c. 34, 40 = dentatus.
viscosus, Hoffm. Denisch. Fl. ed. II. i. i. 21 = fuscus.
viscosus, Salzm. ex Steud. Syn. Fl. Cyp. 15 = compressus.
virginicus, Jacq. ex Steud. l. c. 47 (Mariaci sp.) = *Michauxianus*.
viridis, Krock. Fl. Sil. i. 64 = fuscus.
viridis, Roxb. ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 173 = rotundus.
viridis, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 270 = glaber.
viridis, Willd. ex Kunth, Enum. Pl. ii. 88 = difformis.
viridulus, Boeck. in Linnaea, xxxv. (1867–68) 485 = laevigatus.
viscosus, Sw. Prod. Veg. Ind. Occ. 20 = elegans.
vitiparus, Steud. Syn. Pl. Cyp. 216 = elegans.
Volckmanni, *Phil. in Linnaea*, xxxiii. (1864–65) 269. —Chili.
vulgaris, Hochst. ex Boeck. in Linnaea, xxxv. (1867–68) 490 = maculatus.
vulgaris, Sieber, ex Kunth, Enum. Pl. ii. 4 = Pycreus capillaris.
vulnerans, Salzm. ex Schlecht. in Bot. Zeit. vii. (1849) 96 = Mariscus radus.
Wallichianus, *Spreng. Syst. iv. Cur. Post.* 28.—Nepal.
Wallichii, Nees, in Wight, Contrib. 83 = malaccensis.
Warmingii, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 139.—Bras.
Watsonianus, *Boeck. in Flora*, xlii. (1859) 485 [bis].—Am. bor.
Wawrai, *Boeck. in Linnaea*, xxxviii. (1874) 368.—Bras.
Webbianus, *Steud. Syn. Pl. Cyp.* 20.—Afr. austr.
Wickgreni, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 499. —Bras.
Wieslii, Steud. Syn. Pl. Cyp. 35 = auricomus.
Wightii, Hance, in Journ. Bot. xvi. (1878) 14 = Zollingeri.
Wightii, Nees, in Wight, Contrib. 78 = rubicundus.

CYPERUS:—
Wolfii, *Wood, in Coult. Bot. Gaz.* i. (1876) 58.—Am. bor.
xanthinus, Presl, in Oken, Isis, xxi. (1828) 271 = Pycreus flavescens.
xanthocomus, Link, Hort. Berol. i. 216 = auricomus.
xanthochlus, Steud. in Flora, xxv. (1842) 595 = cleistinodus.
xanthostachyus, Steud. *l. c.* 600.—Chili.
xanthostachyus, Steud. Syn. Pl. Cyp. 332 (Index) = xanthostachys.
yemenicus, Spreng. Syst. i. 221 = jeminicus, *Roth.* (nec *Rets.*).
Zollingeri, *Steud. Syn. Pl. Cyp.* 17.—As. et Afr. trop.

CYPHAEA, Lem. in Orb. Dict. iv. Corrig. (1849) = Cuphea, P. Br. (Lythrar.).

CYPHANTHERA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 376 = Anthocercis, Labill. (Solanac.).
albicans, Miers, l. c. 379 = A. albicans.
cuneata, Miers, l. c. 378 = A. Endesii.
frondosa, Miers, l. c. 376 = A. Endesii.
microphylla, Miers, l. c. 381 = A. microphylla.
ovalifolia, Miers, l. c. 380 = A. albicans.
scabrella, Miers, l. c. = A. scabrella.
tasmanica, Miers, l. c. 377 = A. tasmanica.
tomentosa, Miers, l. c. 379 = A. albicans.

CYPHIA, Berg. Descr. Pl. Cap. 172 (1767). *CAMPANULACEAE*, Benth. & Hook. f. ii. 554.
CYPHIUM, J. F. Gmel. Syst. 370 (1791).
angustifolia, A. DC. in DC. Prod. vii. 500, sphalm. = digitata.
angustifolia, Eckl. & Zeyh. Enum. 390 = undulata.
angustifolia, Presl, in Eckl. & Zeyh. l. c. 389 = *anomala*, Eckl. & Zeyh. l. c. 390 = undulata.
assimilis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 600.—*Afr. austr.*
atriplicifolia, Presl, in Eckl. & Zeyh. Enum. 389 = crenata.
Botrys, Willd. ex Roem. & Schult. Syst. v. 477 = bulbosa.
bulbosa, *Berg. Descr. Pl. Cap.* 172.—Afr. austr.
campestris, *Eckl. & Zeyh. Enum.* 391.—Afr. austr.
Cardamines, *Willd. Sp. Pl.* i. 953.—Afr. austr.
cinarioides, Presl, in E. Mey. Comm. Pl. Afr. Austr. 293, sphalm. vide linarioides.
corymbosa, *Harv. Thes. Cap.* ii. 39.—Afr. austr.
crenata, *Presl, Prod. Monog. Lobel.* 51.—Afr. austr.
dentariaefolia, *Presl, in E. Mey. Comm. Pl. Afr. Austr.* 295.—Afr. austr.
digitata, *Willd. Sp. Pl.* i. 953.—Afr. austr.
Dregeana, Presl, in E. Mey. Comm. Pl. Afr. Austr. 294 = crenata.
Echloniana, Presl, in Eckl. & Zeyh. Enum. 392 = Zeyheriana.
elata, *Harv. Thes. Cap.* ii. 39.—Afr. austr.
Gerrardi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 601.—Afr. austr.
glandulifera, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 8. —Abyss.
heterophylla, *Presl, in Eckl. & Zeyh. Enum.* 389.—Afr. austr.
incisa, *Willd. Sp. Pl.* i. 953.—Afr. austr.
latipetala, Presl, in Eckl. & Zeyh. Enum. 391 = volubilis.
linarioides, *Presl, l. c.*—Afr. austr.
linarioides, Presl, in E. Mey. Comm. Pl. Afr. Austr. 294, ex parte = campestris.
longipetala, Presl, in Eckl. & Zeyh. Enum. 389 = volubilis.
oblongifolia, *Sond. in Harv. & Sond. Fl. Cap.* iii. 601.—Afr. austr.
pentaschista, *Presl, in E. Mey. Comm. Pl. Afr. Austr.* 296.—Afr. austr.
Phyteuma, *Willd. Sp. Pl.* i. 953.—Afr. austr.
phyteumatifolia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = prace.
pinnata, *Roem. & Schult. Syst.* v. 477.—Ind. occ.
polyadenia, Presl, in E. Mey. Comm. Pl. Afr. Austr. 296 = digitata.
salicifolia, Presl, in Eckl. & Zeyh. Enum. 390 = sylvatica.
servata, Spreng. Neue Entdeck. i. 274 = C. Phyteuma.

CYPHIA:—
simplex, Spreng. ex Presl, Prod. Monog. Lobel. 10 = Lobelia scabra.
Stheno, *Webb, in Hook. Niger Fl.* 148.—Ins. Cap. de Verd.
sylvatica, *Eckl. & Zeyh. in S. Afr. Quart. Journ.* (1830) 378.—Afr. austr.
tomentifolia, A. DC. in DC. Prod. vii. 498 = linarioides.
tomentosa, Presl, in E. Mey. Comm. Pl. Afr. Austr. 295 = digitata.
undulata, *Eckl. & Zeyh. in S. Afr. Quart. Journ.* (1830) 378.—Afr. austr.
volubilis, *Willd. Sp. Pl.* i. 953.—Afr. austr.
Zeyheriana, *Presl, in Eckl. & Zeyh. Enum.* 392.—Afr. austr.

CYPHIUM, J. F. Gmel. Syst. 370 (1791) = prsec. *coperse*, J. F. Gmel. l. c. = Cyphia bulbosa.

CYPHOCALYX, Presl (Bot. Bemerk. 127) (1844), in Abh. Boehm. Ges. Wiss. V. iii. (1846) 557 = Aspalathus, Linn. (Legumin.).
aridus, Presl, l. c. = A. aridus.
major, Presl, l. c. = A. aridus.

CYPHOCARPUS, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 93. *CAMPANULACEAE*, Benth. & Hook. f. ii. 554.
rigescens, *Miers, l. c.*—Chili.

CYPHOKENTIA, Brongn. in Comptes Rend. Acad. Sc. Par. lxxvii. (1873) 397. *PALMAE*, Benth. & Hook. f. iii. 894.
Balansae, Brongn. l. c. 400 = Cyphosperma Balansae.
Billardieri, Brongn. l. c. 401 = Microkentia Billardieri.
bractesta, *Brongn. l. c.* 400.—N. Caled.
Deplanchei, Brongn. l. c. 401 = Microkentia Deplanchei.
eriostachys, Brongn. l. c. = Microkentia eriostachys.
gracilis, Brongn. l. c. = Microkentia gracilis.
Humboldtiana, *Brongn. l. c.* 400.—N. Caled.
macrocarpa, Hort.—Cf. Gard. Chron. (1873) 1. 440 = Kentiopsis macrocarpa.
macrostachya, *Brongn. in Comptes Rend. Acad. Sc. Par.* lxxvii. (1873) 399.—N. Caled.
Pancheri, Brongn. l. c. 400 = Microkentia Pancheri.
robusta, *Brongn. l. c.*—N. Caled.
surculosa, Brongn. l. c. 401 = Microkentia surculosa.
vaginata, *Brongn. l. c.* 402.—N. Caled.
Vieillardi, Brongn. & Hook. f. Gen. iii. 895, sphalm. = Cyphosperma Vieillardi.

CYPHOLOPUS, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 198. *URTICACEAE*, Benth. & Hook. f. iii. 388.
Blumeanus, *Wedd. in Arch. Mus. Par.* ix. (1856–57) 437.—Ins. Moluce.
corruleus, Wedd. in DC. Prod. xvi. 1. 235[12] = Blumeanus.
elliptica, *Wedd. l. c.*—Java.
heterophyllus, *Wedd. l. c.* 235[11].—Ins. Fiji.
latifolius, *Wedd. l. c.*—N. Guinea.
luteocrus, Wedd. l. c. = rufescens.
macrocephalus, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 198.—Ins. Philipp.
melanocarpus, *Miq. Fl. Ind. Bat.* i. ii. 968.—N. Guinea.
melucocensis, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 208 = macrocephalus.
prostratus, *Wedd. in DC. Prod.* xvi. 1. 235[10].—N. Guinea; Ind. bor.
rufescens, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 198.—Java.
vestitus, *Miq. Fl. Ind. Bat.* i. ii. 968.—N. Guinea.

CYPHOMANDRA, Mart. ex Sendtn. in Flora, xxviii. (1845) 102. *SOLANACEAE*, Benth. & Hook. f. ii. 889.
CYATHOSTYLUS, Schott, ex Meisn. Gen. Pl. Comment. 184 (1840).
PALLAVICINIA, De Not. in Flora, xxx. (1847) 667.
PHOMANDRA, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 322.
abutiloides, *Griseb. in Goett. Abh.* xxiv. (1879) 249.—Reg. Argent.
adenobasis, *Sendtn. in Flora*, xxviii. (1845) 176.—Bras.
allophylla, *Hemsl. Biol. Centr. Am. Bot.* ii. 417.—Panama.
betacea, Miers, in Trans. Bot. Soc. Edinburgh, 374 = Hentrogi.

CYPHOMANDRA :—
betacea, *Sendt. in Flora*, xxviii. (1845) 172.—Am. auctr.
brachypoda, *Sendt. l. c.* 171.—Bras.
calycosanensis, *Sendt. ex Walp. Rep.* vi. 579.—Am. trop.
calycina, *Sendt. in Flora*, xxviii. (1845) 167.—Bras.
capsicoides, *Dun. in DC. Prod.* xiii. 1. 896; *Sendt. in Mart. Fl. Bras.* x. 152.—Bras.
carassanae, *Sendt. in Flora*, xxviii. (1845) 168. t. 2. f. 1–8.—Venezuela.
ciliata, *Dun. in DC. Prod.* xiii. 1. 401; *Sendt. ex Walp. Rep.* vi. 578.—Bras.
coriacea, *Dun. l. c.*; *Sendt. ex Walp. l. c.* 579.—Peruv.
cornigera, *Dun. in DC. Prod.* xiii. 1. 401.—Bras.
Cornucopia, *Dun. l. c.* 400.—Bras.
corymbifera, *Sendt. in Flora*, xxviii. (1845) 174.—Bras.
crotonifolia, Sendt. ex Walp. Rep. vi. 579 = Solanum crotonifolium.
cylindrica, *Sendt. in Mart. Fl. Bras.* x. 121.—Bras.
diplocona, *Sendt. in Flora*, xxviii. (1845) 169. t. 3. f. 1–10.—Bras.
divaricata, *Sendt. l. c.* 174. t. 9. f. 1–6.—Bras.
elliptica, *Sendt. in Mart. Fl. Bras.* x. 121.—Bras.
fragrans, Sendt. l. c. 128 = Solanum inaequale.
floribunda, *Dun. in DC. Prod.* xiii. 1. 390.—Bras.
fragrans, *Sendt. in Mart. Fl. Bras.* x. 116.—Reg. Argent.
Fraxinella, *Sendt. in Mart. Fl. Bras.* x. 122.—Bras.
Hartwegi, *Sendt. ex Walp. Rep.* vi. 579.—N. Granat.
inabrensis, *Dun. in DC. Prod.* xiii. 1. 400.—Bras.
laxiflora, *Dun. l. c.* 397.—Bras.
lobata, *Sendt. in Mart. Fl. Bras.* x. 115.—Bras.
nervosa, Sendt. ex Walp. Rep. vi. 579 = Solanum nervum.
obliqua, *Sendt. in Flora*, xxviii. (1845) 172.—Peruv.
Osum-Syringilae, Dun. in DC. Prod. xiii. 1. 394 = Vulcaniana.
oxyphylla, *Dun. l. c.* 396.—Bras.
pendula, *Sendt. in Flora*, xxviii. (1845) 173.—Peruv.
physaloides, *Dun. in DC. Prod.* xiii. 1. 388.—Bras.
piperoides, Dun. l. c. 390.—Hort. Berolinensis.
premnaefolia, *Dun. l. c.* 392.—Bras.
procera, *Wawra, in Oestr. Bot. Zeitschr.* xlii. (1888) 221.—Hab.?
sciadostylis, *Sendt. in Flora*, xxviii. (1845) 170. t. 4.—Bras.
splendens, *Dun. in DC. Prod.* xiii. 1. 395.—Peruv.
syncarpa, *Sendt. in Mart. Fl. Bras.* x. 114.—Bras.
tegorea, Hemsl. Biol. Centr. Am. Bot. ii. 417 = seq.
Tejone, *Sendt. ex Walp. Rep.* vi. 579.—Guiana; Nicaragua.
trachyphylla, Sendt. l. c. = Solanum trachyphyllum.
Velloziana, *Sendt. in Flora*, xxviii. (1845) 175.—Bras.
velutina, *Sendt. in Mart. Fl. Bras.* x. 120.—Bras.
viridiflora, *Sendt. in Flora*, xxviii. (1845) 175.—Costa Rica.

CYPHOMATTIA, Boiss. Fl. Orient. iv. 972 (1879) = Rindera, Pall. (Boragin.).
lanata, Boiss. l. c.—As. Min.; Persia.

CYPHONEMA, Herb. in Bot. Mag. sub t. 3710, 3747 (1839) = Cyrtanthus, Ait. (Amaryll.).
Loddigesiana, Herb. l. c. = Cyrt. helictus.

CYPHOPHOENIX, H. Wendl. ex Benth. & Hook. f. Gen. iii. 938 (1883). *PALMAE*, Benth. & Hook. f. i. s.
CAMPECARPUS, H. Wendl. ex Benth. & Hook. f. l. c. (1883).
elegans, *Benth. & Hook. f. l. c.*—N. Caled.
falcata, *Benth. & Hook. f. l. c.*—N. Caled.

CYPHORIMA, Rafin. in Am. Monthly Mag. (1819) 191, 357; Catal. 18 (1824) = Lithospermum, Linn. (Boragin.).
decumbens, Rafin. in Am. Monthly Mag. (1819) 191 = L. angustifolium.
latifolium, Rafin. l. c. = L. latifolium.
longiflora, Rafin. l. c. = L. angustifolium.
tuberosa, Rafin. Catal. 18 = L. latifolium.

CYPHOSPERMA, H. Wendl. ex Benth. & Hook. f. Gen. iii. 935 (1883). *PALMAE*, Benth. & Hook. f. i. s. Balaense, *H. Wendl. ex Benth. & Hook. f. l. c.* 936, 938.—N. Caled.
Vieillardii, *Benth. & Hook. f. l. c.* 936.—N. Caled.

CYPHOSTIGMA, Benth. in Hook. Ic. Pl. t. 1380 (1883). *SCITAMINEAE*, Benth. & Hook. f. iii. 645.
pulchellum, *Benth. l. c.*—Zeylan.

CYPRIANTHE, Spach, Hist. Vég. Phan. vii. 220 (1839) = Ranunculus, Tourn.
anemonoides, Spach, l. c. 221 = R. asiaticus.
asiatica, Freyn. in Willk. & Lange, Prod. Fl. Hisp. iii. 943 = R. asiaticus.

CYPRIPEDIUM, Linn. Syst. ed. I (1735). *ORCHIDEAE*, Benth. & Hook. f. iii. 634.
ARIETINUM, Beck, Bot. N. and Midl. U. St. 352 (1888).
CALCEOLARIA, Helst. Syst. 5 (1749).
CALCEOLUS, Adans. Fam. ii. 70 (1763).
CORDULA, Rafin. Fl. Tellur. iv. 46 (1836).
CORISANTHES, Steud. Nom. ed. II. i. 474 (1840) in syn.
CRIOSANTHES, Rafin. in Journ. Phys. lxxxix. (1819) 102.
HYPODEMA, Reichb. Nom. 56 (1841).
MESENEPHORA, Rafin. Fl. Tellur. iv. 46 (1836).
SACODON, Rafin. l. c. 45 (1836).
STIMEGAS, Rafin. l. c. (1836).
acaule, Ait. Hort. Kew. ed. I. iii. 308 = humile.
album, Ait. l. c. = spectabile.
Argus, *Reichb. f. in Gard. Chron.* (1878) 608.—Ins. Luzon.
arietinum, *R. Br. in Ait. Hort. Kew.* ed. II. v. 222.—Am. bor.
Atamori, *Morr. in Belg. Hortic.* i. (1851) 171. t. 1. —Japon.
barbatum, *Lindl. Bot. Reg.* (1841) *Misc.* 53.—Java. bifidum, *Rafin. Atl. Journ.* 176.—Am. bor.
Boissierianum, Reichb. f. in Bonplandia, ii. (1854) 116 = Selenipedium Boissierianum.
boreale, Salisb. Prod. 10 = C. Calceolus.
Boxallii, Reichb. f. in Gard. Chron. (1877) i. 367 = villosum.
bulbosum, Blanco, Fl. Filip. ed. I. 627 = Eulophia sumatrana.
bulbosum, Linn. Sp. Pl. 951 = Calypso borealis.
Bullenianum, Reichb. f. in Bot. Zeit. xxiii. (1865) 99 = Hookerae.
Burbidgei, Reichb. f. in Gard. Chron. (1881) ii. 38 = Dayanum.
Calceolus, *Linn. Sp. Pl.* 951.—Europ.
Calceolus, Thunb. Fl. Jap. 30 = macranthum.
Calceolus, Walt. Fl. Carol. 223 = pubescens.
californicum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 389.—Calif.
canadense, Michx. Fl. Bor. Am. ii. 261 = spectabile.
candidum, *Muhl. ex Willd. Sp. Pl.* iv. 146.—Am. bor.
cardinale, Reichb. f. in Gard. Chron. xviii. (1882) 488 = Selenipedium cardinale.
cardiophyllum, *Franch. & Sav. Enum. Pl. Jap.* ii. 39. —Japon.
carinianum, Lindl. & Paxt. Flow. Gard. i. (1850–51) 59 = Selenipedium caricinum.
caudatum, Lindl. Gen. et Sp. Orch. 581 = Selenipedium caudatum.
Chica, *Pfitzer, Veryl. Morph. Orch.* 138.—Hab.?
ciliolare, *Reichb. f. in Gard. Chron.* (1882) ii. 488.—Ins. Philipp.
concolor, *Batem. in Hook. Bot. Mag.* t. 5513.—Burma.
cordigerum, *D. Don, Prod. Fl. Nep.* 37.—Reg. Himal.
cothurnatum, *Vell. Fl. Flum.* ix. t. 63.—Bras.
Crossii, Morr. in Belg. Hortic. xv. (1865) 296 = barbatum.
cruciatum, Dalac, Fl. Hautes-Pyr. 198 = C. Calceolus.
crucifurcae, Zoll. & Mor. in De Vriese, Illustr. Orch. t. 6.—Java.
Curtisii, *Reichb. f. in Gard. Chron.* (1882) ii. 8.—Ins. Sondaic.
debile, *Reichb. f. l. c.* (1878) i. 209, in desc. C. Hincksiani.—Panama.
Dayanum, *Reichb. f. in Bot. Zeit.* xx. (1862) 214.—Borneo.
debile, *Reichb. f. Xenia Orch.* ii. 228.—Japon.
Druriî, *Redd. ex Ait. Hort. Kew.* ed. I. 23.—Ind. or.
epidendriocum, *Vell. Fl. Flum.* ix. t. 64.—Bras.
Fairrieanum, *Lindl. in Gard. Chron.* (1857) 740.—Reg. Himal.
fasciculatum, *Kellogg, ex S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 380.—Am. bor.
ferrugineum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 213 = C. Calceolus.
flavescens, [DC. in] Red. Lil. t. 20 = pubescens.
glanduliferum, *Blume, Rumphia*, iv. 56.—N. Guinea.
grande, Reichb. f. in Gard. Chron. (1881) i. 462 = Selenipedium grande.
guttatum, *Sw. in Vet. Acad. Handl. Stockh.* (1800) 251.—Sibir.
Harrisianum x, Reichb. f. in Gard. Chron. (1869) 168.
Hartwegii, Reichb. f. in Seem. Bot. Voy. Herald, 215 = Selenipedium longifolium.
Hayneldianum, *Reichb. f. Xenia Orch.* ii. 222.—Ins. Philipp.
Hincksianum, Reichb. f. in Gard. Chron. (1878) i. 209 = Selenipedium longifolium.
hirsutissimum, *Lindl. ex Hook. Bot. Mag.* t. 4990.—Java.
hirsutum, Mill. Gard. Dict. ed. VIII. n. 3 = spectabile.
Hookerae, *Reichb. f. in Hook. Bot. Mag.* t. 5862.—Borneo.
Humboldti, *Wawra. ex Reichb. f. in Bot. Zeit.* x. (1852) 691.—Peruv.; Panama.
humile, *Salisb. in Trans. Linn. Soc.* i. (1792) 79. t. 3. f. 4.—Am. bor.
insigne, *Wall. ex Lindl. Coll. Bot.* t. 32.—Reg. Himal.
inpeanum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 10.—Mexic.
japonicum, *Thunb. Fl. Jap.* 30.—Japon.
javanicum, *Reinw. ex Blume, Cat. Gew. Buitenz.* 98, nomen; *in Lindl. & Paxt. Flow. Gard.* i. (1850–51) 88.—Java.
Klotzschianum, Reichb. f. in Linnaea, xxii. (1849) 811 = Selenipedium Klotzschianum.
laevigatum, Batem. in Hook. Bot. Mag. t. 5508 = philippinense.
Lawrenceanum, *Reichb. f. in Gard. Chron.* (1878) ii. 748.—Borneo.
Lexarzae, Scheidw. in Otto & Dietr. Allg. Gartenz. viii. (1839) 265 = inpeanum.
Lindenii, Van Houtte, Fl. des Serres, xviii. (1870) 155 = Selenipedium caudatum.
Lindleyanum, Schomb. ex Lindl. Gen. et Sp. Orch. 58 = Selenipedium Lindleyanum.
lineari-tubulatum, Llanos, Fragm. 99 = Cleisostoma longifolium!
longifolium, Warsc. & Reichb. f. in Bot. Zeit. x. (1852) 690 = Selenipedium longifolium.
Lowei, *Lindl. in Gard. Chron.* (1847) 765.—Borneo.
lutetum, Raf. Med. Fl. (1828) 140. t. 80 = pubescens, parviflorum.
macsanthon, *Sw. in Vet. Acad. Handl. Stockh.* (1800) 251.—Reg. Himal.; Sibir.
Mastersianum, *Reichb. f. in Gard. Chron.* (1879) ii. 102. —Ins. Sondaic.
molle, Lindl. in Beath, Fl. Hartw. 72 = inpeanum.
montanum, *Dougl. ex Lindl. Gen. et Sp. Orch.* 528.—Am. bor.
nigritum, *Reichb. f. in Gard. Chron.* (1882) ii. 102.—Borneo.
niveum, *Reichb. f. in* (1869) 1088.—Burma.
occidentale, *S. Wats. in Proc. Am. Acad.* xi. (1876) 147.—Am. bor. occ.
orientale, Spreng. Syst. iii. 746 = guttatum.
palmifolium, *Lindl. Gen. et Sp. Orch.* 527.—Guian.
pardinum, Reichb. f. in Gard. Chron. (1869) 554 = venustum.
Pariahli, *Reichb. f. in Flora*, lii. (1869) 392.—Burma.
parviflorum, *Salisb. in Trans. Linn. Soc.* i. (1791) 77. t. 2.—Am. bor.
parviflorum, Sims, Bot. Mag. t. 911 = pubescens.
passerinum, *Richards. in Frankl. Narr. 1st Journ. App.* ed. II. 34.—Am. bor.
Pearcei, Hort. ex Reichb. f. in Fl. des Serres, Sér. II. vi. (1866) 78 = Selenipedium Pearcei.
paulistanum, *Rodrig. Orch. Nov.* i. 203.—Bras.
patens x, Reichb. f. in Gard. Chron. (1877) ii. 456.
Petzi, Reichb. f. l. c. (1880) i. 680 = Dayanum.
philippinense, *Reichb. f. in Bonplandia*, x. (1862) 335. —Ins. Philipp.
platytaenium, Reichb. f. in Gard. Chron. (1880) ii. 134 = Stonei.
plectrochilum, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 27.—China.
politum x, Reichb. f. in Gard. Chron. (1880) ii. 525.
pubescens, *Willd. Hort. Berol.* i. 18.—Am. bor.
purpuratum, *Lindl. Bot. Reg.* t. 1991.—Malaya.
purpuratum, Wight, Ic. t. 1760 = barbatum.
Reginae, Walt. Fl. Carol. 223 = spectabile.
reticulatum, Reichb. f. in Gard. Chron. (1882) ii. 520 = Selenipedium Bolsierianum.

CYPRIPEDIUM:—
Roebelenii, Reichb. f. in Gard. Chron. (1888) ii. 684 = philippinense.
Rossii, Regel, Gartenfl. (1873) 97 = Selenipedium longifolium.
Schlimi, Lindm. ex Reichb. f. in Bonplandia, ii. (1854) 278, in syn. = Selenipedium Schlimi.
Schomburghianum, Klotzsch, ex Schomb. Remin. 59 = Selenipedium Klotzschianum.
sinicum, Hance, ex Walp. Ann. iii. 602 = purpuratum.
Socco, *Vell. Fl. Flum.* ix. t. 65.—Bras.
spectabile, *Salisb. in Trans. Linn. Soc.* i. (1791) 78.—Am. bor.
Spicerianum, *Reichb. f. in Gard. Chron.* (1880) i. 40, 363.—Reg. Himal.
splendidum, Scheidw. in Otto & Dietr. Allg. Gartenz. viii. (1839) 366 = irapeanum.
Stonei, *Hook. Bot. Mag.* t. 5349.—Borneo.
superbiens, *Reichb. f. in Bonplandia*, iii. (1855) 297.—Malacca.
Thunbergii, Blume, Orch. Archip. Ind. 169 = macranthon.
tonsum, *Reichb. f. in Gard. Chron.* (1888) ii. 262.—Sumatra.
variegatum, Georgi, Reise, i. 282 = guttatum.
Veitchianum, C. Lem. Illustr. Hortic. xii. (1865) t. 429 = superbiens.
venticosum, Sw. in Vet. Acad. Handl. Stockh. (1800) 251 = macranthon.
venustum, *Wall. ex Sims, Bot. Mag.* t. 2129.—Reg. Himal.
villosum, *Lindl. in Gard. Chron.* (1854) 135.—Burma.
virens, Reichb. f. in Bot. Zeit. xxi. (1863) 198 = javanicum.
vittatum, Vell. Fl. Flum. ix. t. 62 = Selenipedium vittatum.
Warnerianum, Reichb. f. ex Wittm. in Berl. Gart. Zeit. (1888) [575] f. 1-3 = barbatum.
Warscewiczianum, Reichb. f. in Bot. Zeit. x. (1852) 692 = Selenipedium caudatum.

CYPROLEPIS, Stend. in Flora, xxxiii. (1850) 229, nomen = **Kyllingа**, Rottb. (Cyperac.).
denudata, Steud. l. c. = K. vaginata.

CYPRELLA, Turp. in Ann. Mus. Par. vii. (1806) 219. t. 121. *FICOIDEAE*, Benth. & Hook. f. i. 856.
MILLEGRANA, Juss. ex Turp. in Ann. Mus. Par. vii. (1806) 220.
RADIANA, Rafin. Specch. i. 88 (1814), ex DC. Prod. iii. 353.
hamiltana, *Turp. in Ann. Mus. Par.* vii. (1806) 219. t. 121. f. 5.—Ins. S. Doming.

CYPSELOCARPUS, F. Muell. Fragm. viii. 36 (1873). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 52.
halocragoides, *F. Muell. l. c.*—Austral.

CYPSELODONTIA, DC. Prod. vii. 286 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 332.
Eckloniana, *DC. l. c.*—Afr. austr.

CYPSOPHILA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 86 (1800), sphalm. = **Gypsophila**, Linn. (Caryophyll.).

CYRRASIUM, Endl. Gen. 891 (1839) = **Cristatella**, Nutt. (Capparid.).
crassum, Endl. ex Walp. Rep. i. 196 = Crist. crosa.
Yamasii, Endl. ex Walp. l. c. = Crist. Jamesii.

CYRILLA, Garden, ex Linn. Mant. i. 5. (1767). *CYRILLEAE*, Benth. & Hook. f. ii. 1026.
antillana, *Michx. Fl. Bor. Am.* i. 158.—Jamaic.
aquatica, Roxb. Pl. Corom. ii. 47. t. 159 = Limnophila racemosa.
caroliniana, Michx. Fl. Bor. Am. i. 158 = racemiflora.
indica, Koen. ex Wight & Arn. Prod. 364 = Vahlia Oldenlandiae.
oldenlandioides, Koen. ex Wight & Arn. l. c. = Vahlia Oldenlandiae.
paniculata, Nutt. in Am. Journ. Sc. v. (1822) 290 = Ardisia Pickeringia.
racemiflora, *Linn. Mant.* i. 50.—Am. bor.
spinosa, Spreng. Nov. Prod. 15 = Bursaria spinosa.
viscosa, Koen. ex Wight & Arn. Prod. 364 = Vahlia viscosa.

CYRILLA, L'Hérit. Stirpes Nov. 147. t. 71 (1788) = Achimenes, P. Br. (Gesneracc.).
pulchella, L'Hérit. l. c. = A. coccinea.

CYRTA, Lour. Fl. Cochinch. 378 (1790) = **Styrax**, Linn.
agrestis, Buch.-Ham. in Wall. Cat. n. 4402 = S. serrulatum.
agrestis, Lour. Fl. Cochinch. 378 = S. agreste.
dealbata, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iii. (1859) 379 = S. Benzoin ?
Finlaysoniana, Miers, l. c. 378 = S. Finlaysonianum.
japonica, Miers, l. c. 379 = S. japonicum.
luculenta, Miers, l. c. = S. serrulatum.
serrulata, Miers, l. c. 380 = S. serrulatum.
suberifolia, Miers, l. c. 379 = S. suberifolium.
virgata, Miers, l. c. 380 = S. serrulatum.

CYRTACANTHUS, Mart. ex Nees, in Mart. Fl. Bras. ix. 51, in syn. (1847) = **Ruellia**, Linn. (Acanth.).
cordifolius, Mart. ex Nees, l. c. (= *Stephanophysum cordifolium*).—Bras.
corymbosus, Mart. ex Nees, l. c. = R. amoena.

CYRTANDRA, Forst. Char. Gen. 5. t. 3 (1776). *GESNERACEAE*, Benth. & Hook. f. ii. 1012.
GETONIA, Banks & Soland, ex Benn. Pl. Jav. Rar. 122 (1838).
RHYNCHOCARPUS, Reinw. & Blume, Cat. Gew. Buitenz. 84 (1823).
WHITIA, Blume, Cat. Gew. Buitenz. 16 (1823).
acuminata, Kurz, in Journ. Bot. xiii. (1875) 239 = Cyrtandromoea decurrens.
acuminata, Wall. Cat. n. 808 = Cyrtandromoea acuminata.
acutangula, *Seem. in Bonplandia*, ix. (1861) 257, nomen ; *Fl. Vit.* 152.—Ins. Fiji.
Albertisi, *C. B. Clarke, in DC. Monog. Phan.* v. 254. —N. Guinea.
ampla, *C. B. Clarke, l. c.* 259.—Sumatra.
aneitensis, *C. B. Clarke, l. c.* 282.—N. Hebrid.
anisophylla, *C. B. Clarke, l. c.* 249.—Sumatra.
anthropophagorum, *Seem. in Bonplandia*, ix. (1861) 257, nomen ; *Fl. Vit.* 152.—Ins. Fiji.
aplicalata, *C. B. Clarke, in DC. Monog. Phan.* v. 276. —Tahiti.
arborescens, *Blume, Bijdr.* 773.—Java.
arguta, *C. B. Clarke, in DC. Monog. Phan.* v. 229.—Ins. Sandvic.
aristata, Blanco, Fl. Filip. ed. I. 18 = Dopatrium aristatum.
attichioides, *C. B. Clarke, in DC. Monog. Phan.* v. 237.—Ins. Sandvic.
aures, *Yack, in Trans. Linn. Soc.* xiv. (1825) 29.—Sumatra.
auriculata, *C. B. Clarke, in DC. Monog. Phan.* v. 251. —Ins. Philipp.
axillaris, *C. B. Clarke, l. c.* 260.—Borneo.
baccifera, *C. B. Clarke, l. c.* 238.—Ins. Sandvic.
basifixa, *C. B. Clarke, l. c.* 254.—Borneo.
batavicensis, *C. B. Clarke, l. c.* 215.—Java.
Beccarii, *C. B. Clarke, l. c.* 249.—Sumatra.
blonloo, *Yack, in Trans. Linn. Soc.* xiv. (1825) 27.—Sumatra.
Bidwillii, *C. B. Clarke, in DC. Monog. Phan.* v. 272. —Ins. Societ.
bifaria, *Forst. Char. Gen.* 5. t. 3.—Ins. Tahiti.
bifora, Hook. & Arn. Bot. Beech. Voy. 67 = Bidwillii.
Blumeana, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 579 = Jumbula.
bornensis, *C. B. Clarke, in DC. Monog. Phan.* v. 244. —Borneo.
Brighami, *C. B. Clarke, l. c.* 221.—Ins. Sandvic.
Burbidgei, *C. B. Clarke, l. c.* 262.—Borneo.
Butti, *Herne, Fiji*, 260, nomen ; *C. B. Clarke, l. c.* 279.—Ins. Fiji.
calycina, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 229.—N. Guinea.
capitellata, *C. B. Clarke, in DC. Monog. Phan.* v. 259. —Ins. Molucc.
cepaeifolia, Blanco, Fl. Filip. ed. I. 17 = Didymocarpus Blancoi.
cernua, *Yack, in Trans. Linn. Soc.* xiv. (1825) 30.—Malaya.
Carnotiana, G. Don, Gen. Syst. iv. 861 = Carnotiana.
celebica, Blume, Bijdr. 772 = coccinea.
Chippendalei, *Horne, Fiji*, 260, nomen ; *C. B. Clarke, in DC. Monog. Phan.* v. 230.—Ins. Fiji.
chrysea, *C. B. Clarke, l. c.* 213.—Borneo.
ciliata, *Seem. in Bonplandia*, ix. (1861) 257, nomen ; *Fl. Vit.* 152.—Ins. Fiji.
coccinea, *Blume, Bijdr.* 772.—Java.
coleoides, *Seem. in Bonplandia*, ix. (1861) 257, nomen ; *Fl. Vit.* 151.—Ins. Fiji.
compressa, *C. B. Clarke, in DC. Monog. Phan.* v. 279. —Ins. Samoa.
confertiflora, *C. B. Clarke, l. c.* 235.—Ins. Sandvic.
connata, *Nadeaud, Enum. Fl. Tahiti*, 60.—Ins. Tahiti.
cordifolia, De Vriese, Fl. Ind. Bat. Orient. 16 = aurea.
cordifolia, *Gaudich. in Freyc. Voy. Bot.* 446. t. 56.—Ins. Sandvic.
coriacea, *C. B. Clarke, in DC. Monog. Phan.* v. 280.—Ins. Fiji.
Cumingii, *C. B. Clarke, l. c.* 263.—Ins. Philipp.
cuneata, *Blume, Bijdr.* 772.—Java.
cymosa, *Forst. Char. Gen.* 6.—Ins. Tana.
cymosa, Mann, ex C. B. Clarke, in DC. Monog. Phan. v. 222 = laxiflora.
decurrens, *De Vriese, Pl. Ind. Bat. Orient.* 14.—Ins. Amboina.
Denhami, *Seem. Fl. Vit.* 152.—Ins. Fiji.
Desvoeuxii, *Horne, Fiji*, 260, nomen ; *C. B. Clarke, in DC. Monog. Phan.* v. 265.—Ins. Fiji.
dilatata, *C. B. Clarke, l. c.* 238.—Borneo.
discolor, *Hort. Lugd. Bat.*—*Cy. Gard. Chron.* (1865) 865.—Hab. ?
dispar, *DC. Prod.* ix. 282.—Malaya.
distimilis, *C. B. Clarke, in DC. Monog. Phan.* v. 207. —Java.
dolichocarpa, *A. Gray, in Proc. Am. Acad.* vi. (1862) 40.—Ins. Fiji.
elongata, Korth. ex C. B. Clarke, in DC. Monog. Phan. v. 232 = decurrens.
eminens, *C. B. Clarke, l. c.* 208.—Borneo.
Endlicheriana, Walp. in Nov. Act. Nat. Cur. xix Suppl. i. (1843) 359. t. 10 = grandiflora.
eximia, *C. B. Clarke, in DC. Monog. Phan.* v. 210.—Borneo.
falcifolia, *C. B. Clarke, l. c.* 269.—Ins. Samoa.
farinosa, *C. B. Clarke, l. c.* 238.—Borneo.
fenestrata, *C. B. Clarke, l. c.* 233.—Sumatra.
flavescens, *Blume, Bijdr.* 773.—Java.
floribunda, Walp. Ann. iii. 99, sphalm. = Isanthera pernotila.
Forsteri, *C. B. Clarke, in DC. Monog. Phan.* v. 251. —China.
frutescens, *Yack, in Trans. Linn. Soc.* xiv. (1825) 31. —Sumatra.
frutescens, Wall. Cat. n. 807 = decurrens, dispar.
fusca, Herb. ex C. B. Clarke, in DC. Monog. Phan. v 262 = populifolia.
Gaudichaudi, *Gaudich. in Freyc. Voy. Bot.* 447. t. 58.—Ins. Sandvic.
glaberrima, Blanco, Fl. Filip. ed. II. 19 = Bonnaya veronicaefolia.
glabra, Banks, ex Gaertn. f. Fruct. iii. 284. t. 224 = bifora.
glabra, Herb. Korth. ex C. B. Clarke, in DC. Monog. Phan. v. 248 = longepedicellata.
glabra, *Yack, in Trans. Linn. Soc.* xiv. (1825) 28.—Sumatra.
glabrata, *Seland. ex C. B. Clarke, in DC. Monog. Phan.* v. 277.—Ins. Societ.
Gorriei, *Horne, Fiji*, 260, nomen ; *C. B. Clarke, l. c.* 278.—Ins. Fiji.
gracilis, *Hillebrand, ex C. B. Clarke, l. c.* 236.—Ins. Sandvic.
Graeffei, *C. B. Clarke, l. c.* 268.—Ins. Samoa.
grandiflora, *Gaudich. in Freyc. Voy. Bot.* 447.—Ins. Sandvic.
grantia, *Blume, Bijdr.* 772.—Java.
Gray, *C. B. Clarke, in DC. Monog. Phan.* v. 218.—Ins. Sandvic.
hapalantha, *C. B. Clarke, l. c.* 262.—N. Guin.
Harveyi, *Seem. Fl. Vit.* 152.—Ins. Fiji.
Hawaiensis, *C. B. Clarke, in DC. Monog. Phan.* v. 265.—Ins. Sandvic.
heterophylla, *De Vriese, Pl. Ind. Bat. Orient.* 15.—Java.
Hillebrandi, *C. B. Clarke, in DC. Monog. Phan.* v. 226.—Ins. Sandvic.
Hillebrandii, Vatke, in Linnaea, xl. (1876) 228 = Waldaui.
hinata, *Yack, in Trans. Linn. Soc.* xiv. (1825) 27.—Sumatra.
holodasys, *Miq. Fl. Ind. Bat.* ii. 747.—Sumatra.
humulacoides, *Wawra, in Flora*, lx. (1877) 267.—Ins. Sandvic.

CYRTANDRA:—
Hornel, *C. B. Clarke, in DC. Monog. Phan.* v. 281.—Ins. Fiji.
Horsfieldii, Miq. Fl. Ind. Bat. ii. 738 = Pentaphragma begoniaefolium.
humilis, *Blume, Bijdr.* 769.— Malaya.
imminuta, *C. B. Clarke, in DC. Monog. Phan.* v. 254.—Borneo.
impressivenia, *C. B. Clarke, l. c.* 212.—Sumatra.
incisa, *C. B. Clarke, l. c.* 250.—Ins. Philipp.
incompta, Jack, in Trans. Linn. Soc. xiv. (1825) 99.—Sumatra.
induta, *A. Gray, in Proc. Am. Acad.* vi. (1862) 38.—Ins. Societ.
insignis, *C. B. Clarke, in DC. Monog. Phan.* v. 268.—Java.
integrifolia, *C. B. Clarke, l. c.* 248.—Sumatra.
involucrata, *Sonn. in Bonplandia*, ix. (1861) 257, nomen; *Fl. Vit.* 183.—Ins. Fiji.
Kalichii, *Wawra, in Flora*, lv. (1872) 564.—Ins. Sandvic.
kaxaiensis, *Wawra, l. c.* 566.—Ins. Sandvic.
Kealiae, *Wawra, l. c.* 565.—Ins. Sandvic.
labiosa, *A. Gray, in Proc. Am. Acad.* vi. (1862) 40.—Ins. Samoa.
lanuginosa, R. Br. in Wall. Cat. n. 7181 = Isanthera permollis.
latens, *C. B. Clarke, in DC. Monog. Phan.* v. 215.—Borneo.
latifolia, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 238 = cymosa.
laxiflora, *H. Mann, in Proc. Am. Acad.* vii. (1867) 190.—Ins. Sandvic.
Lessoniana, *Gaudich. in Freyc. Voy. Bot.* 447. t. 54.—Ins. Sandvic.
ligulifera, *C. B. Clarke, in DC. Monog. Phan.* v. 252.—N. Guin.
Lobbi, *C. B. Clarke, l. c.* 282.—Ins. Philipp.
longe-petiolata, *De Vriese, Pl. Ind. Bat. Orient.* 12.—Java.
longifolia, *Hillebr. ex C. B. Clarke, in DC. Monog. Phan.* v. 276.—Ins. Sandvic.
longirostris, *De Vriese, Pl. Ind. Bat. Orient.* 12.—Ins Celebes.
lysiosepala, *C. B. Clarke, in DC. Monog. Phan.* v. 235.—Ins. Sandvic.
Macraei, *A. Gray, in Proc. Am. Acad.* v. (1861) 352.—Ins. Sandvic.
macrantha, *C. B. Clarke, in DC. Monog. Phan.* v. 265.—Ins. Sandvic.
macrophylla, Jack, in Trans. Linn. Soc. xiv. (1825) 95. t. 2.—Sumatra.
maculata, Jack, *l. c.* 96.—Sumatra.
malacophylla, *C. B. Clarke, in DC. Monog. Phan.* v. 237.—Ins. Sandvic.
Menziesii, *Hook. & Arn. Bot. Beech. Voy.* 91.—Ins. Sandvic.
microcarpa, *C. B. Clarke, in DC. Monog. Phan.* v. 244.—Borneo.
microcarpa, Herb. Mus. Bot. Lugd. Bat. ex C. B. Clarke, l. c. 195 = Rhyncotechum parviflorum.
microstigma, *C. B. Clarke, l. c.* 284.—Ins. Fiji.
Milnei, *Sonn. in Bonplandia*, ix. (1861) 257, nomen; *Fl. Vit.* 183.—Ins. Fiji.
mirabilis, *C. B. Clarke, in DC. Monog. Phan.* v. 233.—Borneo.
mollis, *De Vriese, Pl. Ind. Bat. Orient.* 16.—Ins. Celebes.
muconota, *Nadeaud, Enum. Pl. Tahiti*, 59.—Ins. Tahiti.
multibracteata, *C. B. Clarke, in DC. Monog. Phan.* v. 258.—Borneo.
Nadeaudi, *C. B. Clarke, l. c.* 264.—Ins. Tahiti.
naviculata, *Zipp. ex C. B. Clarke, l. c.* 240.—Sumatra.
nemorosa, *Blume, Bijdr.* 771.—Java.
nemorosa, De Vriese, Pl. Ind. Bat. Orient. 18 = sulcata.
nigra, *Nadeaud, Enum. Pl. Tahiti*, 60.—Ins. Tahiti.
ulsna, *C. B. Clarke, in DC. Monog. Phan.* v. 278.—Ins. Samoa.
multiflora, *C. B. Clarke, l. c.* 270.—Ins. Samoa.
oblongifolia, *Benth. & Hook. f. Gen.* ii. 1012.—Java.
oenobarba, *H. Mann, in Proc. Am. Acad.* vii. (1867) 189.—Ins. Sandvic.
oligantha, Korth. ex C. B. Clarke, in DC. Monog. Phan. v. 241.—Sumatra.
pallescens, Jack, ex C. B. Clarke, l. c. 281 = pilosa.
paludosa, *Gaudich. in Freyc. Voy. Bot.* 447.—Ins. Sandvic.

CYRTANDRA:—
paradoxa, Hillebr. ex C. B. Clarke, in DC. Monog. Phan. v. 277 = longifolia.
parviflora, *C. B. Clarke, l. c.* 268.—Manilla.
peltata, Jack, in Trans. Linn. Soc. xiv. (1825) 80.—Sumatra.
pendula, *Blume, Bijdr.* 768.—Java.
pendula, Nadeaud, Enum. Pl. Tahiti, 60 = spiculata.
personata, Blanco, Fl. Filip. ed. I. 18; ed. II. 18 = Bonnaya brachiata.
philippinensis, *C. B. Clarke, in DC. Monog. Phan.* v. 260.—Manilla.
phoenicea, *C. B. Clarke, l. c.* 256.—Ins. Ceram.
Pickeringii, *A. Gray, in Proc. Am. Acad.* v. (1861) 350.—Ins. Sandvic.
picta, *Blume, Bijdr.* 769.—Java.
picta, Zoll. Syst. Verz. Ind. Arch. (1854) 55 = repens.
pilosa, *Blume, Bijdr.* 770.—Java.
platyphylla, *A. Gray, in Proc. Am. Acad.* v. (1861) 350.—Ins. Sandvic.
poposantha, *A. Gray, l. c.* vi. (1862) 40.—Ins. Samoa.
polyantha, *C. B. Clarke, in DC. Monog. Phan.* v. 220.—Ins. Sandvic.
populifolia, *Miq. Fl. Ind. Bat.* ii. 741.— Java.
Pritchardii, *Sonn. in Bonplandia*, ix. (1861) 264.—Ins. Fiji.
pubescens, Steud. Nom. ed. II. i. 475 = frutescens.
pulchella, *Rich. ex A. Gray, in Proc. Am. Acad.* vi. (1862) 38.—Ins. Samoa.
radicifiora, *C. B. Clarke, in DC. Monog. Phan.* v. 239.—Borneo.
repens, *De Vriese, Pl. Ind. Bat. Orient.* 14.—Java.
reticosa, *C. B. Clarke, in DC. Monog. Phan.* v. 212.—Java.
rhyncanthera, *C. B. Clarke, l. c.* 233.—Sumatra.
Richii, *A. Gray. in Proc. Am. Acad.* vi. (1862) 39.—Ins. Samoa.
rostrata, *Blume, Bijdr.* 771.—Java.
rubiginosa, Jack, in *Trans. Linn. Soc.* xiv. (1825) 92.—Sumatra.
rubra, *De Vriese, Pl. Ind. Bat. Orient* 16.—Java.
Ruchiana, Meyen, Reise, ii. 126 = grandiflora.
ruseа, *C. B. Clarke, in DC. Monog. Phan.* v. 236.—Borneo.
samoensis, *A. Gray, in Proc. Am. Acad.* vi. (1862) 39.—Ins. Societat.
Sandei, *De Vriese, Pl. Ind. Bat. Orient.* 14.—Java.
sarawakensis, *C. B. Clarke, in DC. Monog. Phan.* v. 209.—Borneo.
scabrella, *C. B. Clarke, l. c.* 277.—Ins. Sandvic.
schizostyla, *C. B. Clarke, l. c.* 204.—Borneo.
serrata, *Blanco, Fl. Filip.* ed. I. 18.—Ins. Philipp.
serrata, *Fauc. in Forbes, Nat. Wand. East. Archip.* 513.—Ins. Timor.
spicata, *De Vriese, Pl. Ind. Bat. Orient.* 13.—Java.
splendens, *C. B. Clarke, in DC. Monog. Phan.* v. 209.—Borneo.
squamulata, Korth. ex C. B. Clarke, l. c. 203 = dispar.
staminea, Vahl, Enum. I. 105 = Agalmyla staminea.
sulcata, *Blume, Bijdr.* 770.—Java.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 60.—Ins. Tahiti.
taitensis, *Rich, ex A. Gray, in Proc. Am. Acad.* vi. (1862) 39.—Ins. Societ.
Tempesti, *Horne, Fiji*, 260, nomen; *C. B. Clarke, in DC. Monog. Phan.* v. 273.—Ins. Fiji.
teres, *C. B. Clarke, l. c.* 207.—Borneo.
Teysmanni, *Miq. Fl. Ind. Bat.* ii. 741.—Sumatra.
trifora, *Gaudich. in Freyc. Voy. Bot.* 447. t. 52.—Ins. Sandvic.
trisepala, *C. B. Clarke, in DC. Monog. Phan.* v. 253.—Borneo.
tristis, *Hillebr. ex C. B. Clarke, l. c.* 227.—Ins. Sandvic.
umbellata, De Vriese, Pl. Ind. Bat. Orient. 18 = cuneata.
urceolata, *C. B. Clarke, in DC. Monog. Phan.* v. 236.—Borneo.
Urvillei, *C. B. Clarke, l. c.* 251.—Ins. Carol.
uriculosa, *C. B. Clarke, l. c.* 230.—Ins. Fiji.
velutina, Herb. Mus. Bot. Lugd. Bat. ex C. B. Clarke, l. c. 260 = aurea.
velutina, *Korth. ex C. B. Clarke, l. c.* 208.—Borneo.
viridescens, *C. B. Clarke, l. c.* 247.—Sumatra.
vitiensis, *Sonn. in Bonplandia*, ix. (1861) 257, nomen; *Fl. Vit.* 182.—Ins. Fiji.
Wrightii, *C. B. Clarke, in DC. Monog. Phan.* v. 237.—Ins. Celebes.

CYRTANDRA:—
Waiolani, *Wawra, in Flora*, lv. (1872) 566.—Ins. Sandvic.
Wawrai, *C. B. Clarke, in DC. Monog. Phan.* v. 228.—Ins. Sandvic.
Zippelii, *C. B. Clarke, l. c.* 289.—Java.
Zollingeri, *C. B. Clarke, l. c.* 268.—Java.

CYRTANDROMOEA, Zoll. Syst. Verz. Ind. Archip. 58 (1854). *GESNERACEAE*, Benth. & Hook. f. ii. 1020.
Bussa, Miq. Fl. Ind. Bat. ii. 732 (1856).
acuminata, *Benth. & Hook. f. Gen.* ii. 1020.—Ins. Penang.
angustifolia, *C. B. Clarke, in DC. Monog. Phan.* v. 186.—Sumatra.
cymulosa, *C. B. Clarke, l. c.*—Burma.
decurrens, *Zoll. Syst. Verz. Ind Archip.* 58.—Java.
dispar, *C. B. Clarke, in DC. Monog. Phan.* v. 187.—Sumatra.
grandiflora, *C. B. Clarke, l. c.* 186.—Malaya.
Miqueliana, *C. B. Clarke, l. c.* 187.—Sumatra.
subintegra, *C. B. Clarke, l. c.*—Sumatra.

CYRTANNUS, Herb. App. 28 (1821), sphalm. = **Cyrtanthus**, Ait. (Amaryll.).

CYRTANTHEMUM, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 130; (Gen. Centr. Am. 56. t. 10. f. 20, 39) = **Besleria**, Linn. (Gesnetrac.).
deflexum, Oerst. ll. cc. = B. deflexa.
hirsutum, Oerst. ll. cc. = B. Cyrtanthemum.

CYRTANTHERA, Nees, in Mart. Fl. Bras. ix. 99. t. 14 (1847) = **Jacobinia**, Moric. (Acanthac.).
aurea, Nees, in DC. Prod. xi. 329 = J. aurea.
aurantiaca, Hook. Bot. Mag. t. 4468 = Beloperone aurantiaca.
catalpaefolia, Nees, l. c. t. 4444 = J. aurea.
Chamissoniana, Nees, in Mart. Fl. Bras. ix. 101.—Bras.
chrysostephana, Hook. f. Bot. Mag. t. 5887 = J. chrysostephana.
citrina, Wawra, in Oestr. Bot. Zeitschr. xxxi. (1881) 281.—Bras.
densiflora, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 147 = J. aurea.
Ghiesbreghtiana, Decne. in Rev. Hortic. Sér. III. ii. (1846) 101.—Mexic.
Libboniana, Lem. Jard. Fleur. iii. (1852–53) t. 311 = J. Libboniana.
longiflora, Nees, in DC. Prod. xi. 329 = J. longiflora.
macrantha, Nees, l. c. 330 = J. macrantha.
magnifica, Nees, in Mart. Fl. Bras. ix. 100.—Bras.
Pohliana, Nees, l. c. 101 = J. Pohliana.
Sellowiana, Nees, l. c. = J. Sellowiana.
umbrosa, Nees, in DC. Prod. xi. 396 = J. aurea.

CYRTANTHERELLA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 148. t. 3. f. 10, 11 = **Jacobinia**, Moric. (Acanthac.).
macrantha, Oerst. l. c. = J. macrantha.

CYRTANTHUS, Schreb. Gen. 122 (1789) = **Posoqueria**, Aubl. (Rubiac.).
longiflorus, Steud. Nom. ed. II. i. 476 = P. longiflora.

CYRTANTHUS, Ait. Hort. Kew. ed. I. i. 414 (1789); iii. 510. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 729.
Cyphonema, Herb. in Bot. Mag. sub t. 3710, et sub t. 3747 (1838).
Eusipho, Salisb. Gen. Pl. Fragm. 189 (1866).
Gastronema, Herb. App. 30 (1821).
Mounella, Salisb. Gen. Pl. Fragm. 189 (1866).
Timmia, J. F. Gmel. Syst. 388 (1791).
alto-luteus, Burch. ex M. Roem. Syn. Ensat. 49 = lutescens.
angustifolius, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 270 = uniflorus.
angustifolius, *Ait. Hort. Kew.* ed. I. i. 414.—Afr. austr.
breviflorus, Harv. Thes. Cap. ii. 25. t. 189 = Anoiganthus breviflorus.
carneus, *Lindl. in Bot. Reg.* t. 1462.—Afr. austr.
coccineus, M. Roem. Syn. Ensat. 80 = angustifolius.
collinus, *Ker-Gawl. in Bot. Reg.* t. 162.—Afr. austr.
cordifolius, Mart. ex Nees, in DC. Prod. xi. 205 sphalm. vide *Cyrtacanthus cordifolius*.

CYRTANTHUS:—
helictus, *Lehm. Delect. Sem. Hort. Hamb.* (1839) 7.—Afr. austr.
intescens, *Herb. Amaryll.* 199. t. 33. f. 14.—Afr. austr.
luteus, *Eckl. & Zeyh. ex M. Roem. Syn. Ensat.* 52, nomen.—Afr. austr.
Mackenii, *Hook. f. in Gard. Chron.* (1869) 641.—Afr. austr.
Macowani, *Baker, in Gard. Chron.* (1875) 98.—Afr. austr.
obliquus, *Ait. Hort. Kew.* ed. I. i. 414.—Afr. austr.
ochroleucus, *Burch. ex Steud. Nom.* ed. II. i. 475.—Afr. austr.
odorus, *Ker-Gawl. in Bot. Reg.* t. 503.—Afr. austr.
pallidus, *Sims, Bot. Mag.* t. 2471.—Afr. austr.
Pumilio, *Schult. f. Syst.* vii. 900.—Afr. austr.
pendens, *Eckl. ex Salm-Dyck, Hort. Dyck.* 87, nomen.—Afr. austr.
purpureus, Herb. in Bot. Mag. sub t. 2113 = Amaryllis purpurea.
sanguineus, *Walp. Ann.* iii. 616.—Afr. austr.
Smithiæ, *Watt, ex Hafw. Gen. S. Afr. Pl.* 388.—Afr. austr.
Smithianus, Herb. in Bot. Mag. sub t. 3779 = præc.
spiralis, *Burch. ex Ker-Gawl. in Bot. Reg.* t. 167.—Afr. austr.
striatus, Herb. in Bot. Mag. t. 284 = angustifolius.
Tuckii, *Baker, in Journ. Bot.* xiv. (1876) 183.—Afr. austr.
uniflorus, *Ker-Gawl. in Bot. Reg.* t. 168.—Afr. austr.
varius, M. Roem. Syn. Ensat. 45 = obliquus.
ventricosus, Willd. Sp. Pl. ii. 49 = angustifolius.
vittatus, *Desf. ex* [*DC. in*] *Red. Lil.* iv. t. 183.—Afr. austr.
Welwitschii, *Hiern, ex Baker, in Journ. Bot.* xvi. (1878) 197.—Afr. trop.

CYRTOCARPA, H. B. & K. Nov. Gen. et Sp. vii. 19. t. 609 (1825). *ANACARDIACEÆ*, [Benth. & Hook. f. i. 493.]
Dastycarya, Liebm. in Kjoeb. Vidensk. Meddel. (1858) 98.
Copallilo, *Schlecht. in Linnæa*, xvi. (1842) 455.—Mexic.
procera, *H. B. & K. Nov. Gen. et Sp.* vii. 20. t. 609.—Mexic.
quinquestila, Blanco, Fl. Filip. ed. II. 269 = Odina speciosa.

CYRTOCERAS, Benn. Pl. Jav. Rar. 90. t. 21 (1838) = **Hoya**, R. Br. (Asclep.).
acuminatum, Regel, Cat. Pl. Hort. Aksakov. 46, nomen.—Hab. ?
coriaceum, Heynh. Nom. ii. 183 = H. multiflora.
elegans, Miq. Fl. Ind. Bat. ii. 515.—Borneo.
floribundum, Maund, Botanist, iv. t. 178 = H. multiflora.
Gibsoni, Regel, Cat. Pl. Hort. Aksakov. 46, nomen.—Hab. ?
laurifolium, Miq. Fl. Ind. Bat. ii. 514.—Java.
Lindleyanum, Miq. l. c. 515 = H. multiflora.
micranthum, Miq. l. c.—Ins. Celebes.
multiflorum, Heynh. Nom. ii. 183 = H. multiflora.
reflexum, Benn. Pl. Jav. Rar. 90 = H. multiflora.
speciosum, Regel, Cat. Pl. Hort. Aksakov. 46, nomen.—Hab. ?
uncinatum, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 408.—Sumatra.

CYRTOCHILOS, Spreng. Syst. iii. 739 (1826) = seq.

CYRTOCHILUM, H. B. & K. Nov. Gen. et Sp. i. 340 (1815) = Oncidium, Sw. (Orchid.).
bictoniense, Batem. Orch. Mex. et Guat. t. 6 = Odontoglossum bictoniense.
citrinum, Hook. Bot. Mag. t. 4454 = Oncid. concolor.
flexum, Lindl. Bot. Reg. (1841) t. 59 = Oncid. graminifolium.
flavescens, Lindl. l. c. t. 1697 = Miltonia flavescens.
flexuosum, H. B. & K. Nov. Gen. et Sp. i. 350 = Oncid. cimiciferum.
graminifolium, Lindl. in Ann. Nat. Hist. iv. (1840) 384 = Oncid. graminifolium.
incurvatum, Reichb. f. in Gard. Chron. (1872) 1085 = Oncid. hastatum.
iridifolium, Lindl. Gen. et Sp. Orch. 211 = Odontoglossum iridifolium.
Jurgensenianum, Lem. Jard. Fleur. iv. (1854) t. 368 = Oncid. hastatum.

CYRTOCHILUM :—
Karwinskii, Lindl. Bot. Reg. sub t. 1992 = Miltonia Karwinskii.
leucochilum, Planch. in Fl. des Serres, Sér. I. v. (1849) t. 522 = Oncid. leucochilum.
maculatum, Lindl. Bot. Reg. (1838) t. 44 = Oncid. maculatum.
mystacinum, Lindl. l. c. Misc. 90 = Odontoglossum mystacinum.
pardinum, Lindl. Gen. et Sp. Orch. 210.—Peruv.
stellatum, Lindl. Sert. Orchid. t. 7 = Miltonia flavescens.
undulatum, H. B. & K. Nov. Gen. et Sp. i. 349. t. 84 = Oncid. undulatum.
volubile, Poepp. & Endl. Nov. Gen. et Sp. i. 35 = Oncid. corymbiferum.
zebrinum, Reichb. f. in Gard. Chron. (1872) 1355 = Oncid. zebrinum.

CYRTOCLADON, Griff. Notul. iii. 147 (1851) = **Komalomena**, Schott (Aroid.).
sanguinolentum, Griff. l. c. (= *Chamæcladon sanguinolentum*).—Ind. or.

CYRTODEIRA, Hanst. in Linnæa, xxvi. (1853) 207 = **Episcia**, Mart. (Gesner.).
chontalensis, Seem. in Gard. Chron. (1867) 655 = E. chontalensis.
coccinea, Hort. Bull.—Cf. Gard. Chron. iii. (1875) 1. 490.—Hab. ?
cuprata, Hanst. in Linnæa, xxvi. (1853) 207. t. 2. f. 20 = E. cuprea.
fulgida, Linden, ex André, Illustr. Hortic. xx. (1873) t. 181 = E. fulgida.
Verschaffelti, Hanst. in Koch, Allgem. Gartenz. xxv. (1857) 237 = E. splendens.

CYRTOGYNE, Reichb. Consp. 156 (1828) = CURTOGYNE, Haw. = **Crassula**, Linn.

CYRTOLEPIS, Less. in Linnæa, vi. (1831) 166 = **Anacyclus**, Linn. (Compos.).
alexandrina, DC. Prod. vi. 17 = A. alexandrinus.
monantha, Less. in Linnæa, vi. (1831) 166 = A. alexandrinus.

CYRTOLOBUM, R. Br. in Wall. Cat. in nota sub n. 5432 (1832) = **Crotalaria**, Linn. (Legumin.).
divaricata, R. Br. l. c. n. 5436 (cf. 5432) = Crot. medicaginea.
elliptica, R. Br. l. c. n. 5438 (cf. 5432) = Crot. elliptica.
medicaginea, R. Br. l. c. n. 5434 (cf. 5432) = Crot. medicaginea.
procumbens, R. Br. l. c. n. 5437 (cf. 5432) = Crot. medicaginea.
spartioides, R. Br. l. c. n. 5435 (cf. 5432) = Crot. Wildenoviana.
Trifoliastrum, R. Br. l. c. n. 5432 = Crot. Trifoliastrum.

CYRTONEMA, Schrad. in Eckl. & Zeyh. Enum. 275 (1836) = **Kedrostis**, Medic. (Cucurbit.).
convolvulacea, Hochst. in Flora, xxv. (1842) i. Hab. 135 = K. foetidissima.
digitatum, Schrad. in Eckl. & Zeyh. Enum. 277 = K. digitata.
dewegens, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 286 = K. foetidissima.
elegans, Fenzl, ex Cogn. in DC. Monog. Phan. iii. 857 = Cosellocarpus Gilgi.
foetidissima, Hochst. ex Hook. f. in Oliver, Fl. Trop. Afr. ii. 564 = K. foetidissima.
hirtellum, Hochst. ex Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1862) 181 = K. hirsella.
latidentatum, Schrad. in Eckl. & Zeyh. Enum. 277 = K. nana.
molle, Kunze, in Linnæa, xx. (1847) 49 = K. mollis.
sphærocarpum, Schrad. in Eckl. & Zeyh. Enum. 276 = K. Zeyheri.
trilobata, Cogn. in DC. Monog. Phan. iii. 836 = præc.
trilobum, Schrad. in Eckl. & Zeyh. Enum. 276 = K. nana.

CYRTONORA, Zipp. ex Miquel, in Bijdr. Nat. Wet. v. (1836) 179, nomen (quid?).
leucochyla, Zipp. l. c.—N. Guinea.

CYRTOPERA, Lindl. in Wall. Cat. n. 7362–64 (1831); et Gen. et Sp. Orch. 189 (1833) = **Eulophia**, R. Br. (Orchid.).
amaonica, Rodrig. Orch. Nov. ii. 180.—Bras.
aurea, Boxall, ex Naves, in Blanco, Fl. Philipp. ed. III. Nov. App. 243, nomen.—Ins. Philipp.
bicarinata, Lindl. in Wall. Cat. n. 7363.—Reg. Himal.
candida, Lindl. in Journ. Linn. Soc. iii. (1859) 31.—Reg. Himal.
Cullenii, Wight, Ic. t. 1754 = *flava*.
densiflora, Reichb. f. Otia Bot. Hamb. 68, in obs. = E. densiflora.
enantiomorpha, Lindl. in Journ. Linn. Soc. iii. (1859) 31 = Cyrtopodium enantiomorphum.
flava, Lindl. in Wall. Cat. n. 7364.—Ind. or.
flavo-purpurea, Reichb. f. Otia Bot. Hamb. 68.—Afr. trop.
foliosa, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 203 = E. foliosa.
fusca, Wight, Ic. t. 1690 = E. nuda.
Gardneri, Thw. Enum. Pl. Zeyl. 302 = E. nuda.
gigantea, Lindl. Gen. et Sp. Orch. 190.—Afr. austr.
Godefroyi, Reichb. f. Otia Bot. Hamb. 85.—Cambodia.
laniflora, Gardn. ex Thw. Enum. Pl. Zeyl. 302 = E. nuda.
longifolia, Reichb. f. in Walp. Ann. vi. 668 = Cyrtopodium Woodfordii.
macrobulbon, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 144.—Burma.
maculata, Boxall, ex Naves, in Blanco, Fl. Philipp ed. III. Nov. App. 242, nomen.—Ins. Philipp.
Mannii, Reichb. f. in Flora, lv. (1872) 274.—Reg. Himal.
mysorensis, Lindl. in Journ. Linn. Soc. iii. (1859) 32 = E. nuda.
nuda, Reichb. f. in Flora, lv. (1872) 274 = E. nuda.
obtusa, Lindl. Gen. et Sp. Orch. 190.—Ind. or.
Olivieriana, Reichb. f. in Flora, lxiv. (1881) 329.—Afr. austr.
pedicellata, Lindl. Gen. et Sp. Orch. 190 = E. scaleata.
plantaginea, Lindl. l. c. 189 = Cyrtopodium plantagineum.
plicata, Lindl. in Wall. Cat. n. 7362 = E. nuda.
rufa, Thw. Enum. Pl. Zeyl. 302 = *sanguinea*.
sanguinea, Lindl. in Journ. Linn. Soc. iii. (1859) 32.—Ind. or.
scabrilingua, Lindl. Gen. et Sp. Orch. 189.—Peruv.
Shuttangæ, Reichb. f. Otia Bot. Hamb. 116.—Afr. trop.
squalida, Reichb. f. in Bonplandia, v. (1857) 38 = E. squalida.
Velloziana, Rodrig. Orch. Nov. ii. 179.—Bras.
Walkeri, Reichb. f. Otia Bot. Hamb. 117.—Afr. trop.
Woodfordii, Lindl. Gen. et Sp. Orch. 189 = Cyrtopodium Woodfordii.
Zollingeri, Reichb. f. in Bonplandia, v. (1857) 38.—Java.

CYRTOPHYLLUM, Reinw. ex Blume, Cat. Gew. Buitenz. 47 (1836); et Bijdr. 1023 (1836) = **Pagraea**, Thunb. (Loganiac.).
fragrans, DC. Prod. ix. 31 = F. fragrans.
lanceolatum, DC. l. c. = F. Wallichiana.
peregrinum, Reinw. ex Blume, Cat. Gew. Buitenz. 47; et Bijdr. 1023 = F. fragrans.
speciosum, Blume, Bijdr. 1023 = F. speciosa.

CYRTOPODIUM, R. Br. in Ait. Hort. Kew. ed. II. v. 216 (1813). *ORCHIDEÆ*, Benth. & Hook. f. iii. 543.
Tylochilus, Nees, in Verh. Gartenb. Ges. Berl. viii. (1832) 194. t. 3.
Andersonii, *R. Br. in Ait. Hort. Kew.* ed. II. v. 216.—Ind. occ.
bicolor, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 472.—Ins. Timor.
Blanchetii, *Reichb. f. in Linnæa*, xxii. (1849) 852.—Bras.
punctatum, *Linden, ex Lindl. Orch. Linden.* 23.—Venezuela.
Reedeanum, *Rodrig. Orch. Nov.* i. 129.—Bras.
sordidum, Lindl. in Journ. Hort. Soc. iv. (1849) 296 = Andersonii.
striatum, *Lindl. Bot. Reg.* (1841) sub t. 8.—Guiana.
stramineum, *Rodrig. in Revist. Exp. iii. 74, ex Orch. Nov.* ii. 129.—Bras.

CYRTOPODIUM :—
elegans, Ham. Prod. Fl. Ind. Occ. 53 = Tetramicra riglida.
Engelii, *Karst. in C. Koch & Fint. Wochenschr.* ii. (1859) 236.—Venezuela.
emiformæ, *Vidal, Phan. Cuming. Philipp.* 76, 80, 150. —Ins. Philipp.
Eugenii, *Reichb. f. & Warm. in Otia Bot. Hamb.* 89. —Bras.
flavescens, *Lindl. ex Steud. Nom.* ed. II. i. 476, nomen. —Mexic.
flavum, Benth. in Journ. Linn. Soc. xvii. (1881) 320 (Eulophiæ sp.).—Ind. or.
flavum, *Link & Otto, ex Reichb. Hort. Bot.* iii. 7.— Bras.
glutiniformo, Raddi, in Mem. Soc. Ital. Mod. xix. ii. (1822) 220 = Woodfordii.
glutinosum, *Meyer, ex Steud. Nom.* ed. II. i. 476, nomen.—Am. austr.
pallidum, *Reichb. f. & Warm. in Otia Bot. Hamb.* 89. —Bras.
palmifrons, *Reichb. f. & Warm. l. c.* 88.—Bras.
parviflorum, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 672.—Guiana.
plantaginea, *Benth. in Journ. Linn. Soc.* xvii. (1881) 320.—Madag.
poecilum, *Reichb. f. & Warm. in Otia Bot. Hamb.* 89. —Bras.
punctatum, *Lindl. Gen. et Sp. Orch.* 186.—Ind. occ.
purpureum, *Reichb. f. & Warm. in Otia Bot. Hamb.* 90.—Bras.
Saintlegerianum, *Reichb. f. in Gard. Chron.* (1885) i. 756.—Paraguay.
spectatissimum, Hort. ex Planch. in Fl. des Serres, xxii. (1877) 179 = punctatum.
squalidum, Vidal, Phan. Cuming. Philipp. 79, 150 = Eulophia squalida.
strictum, *Griseb. Cat. Pl. Cub.* 266.—Cuba.
tigrinum, *Linden, in Illustr. Hortic.* xxviii. (1881) 95, nomen.—Bras.
triste, *Reichb. f. & Warm. in Otia Bot. Hamb.* 90.— Bras.
vernum, *Reichb. f. & Warm. l. c.* 89.—Bras.
verrucosum, *Griseb. Cat. Pl. Cub.* 265.—Cuba.
virescens, *Reichb. f. & Warm. in Otia Bot. Hamb.* 89 —Bras.
Willmorei, Knowles & Westc. Fl. Cab. i. (1837) t. 4 = punctatum.
Woodfordii, *Sims, Bot. Mag.* t. 1814.—Ind. occ.

CYRTOPOGON, Spreng. Syst. i. 266 (1825) = CURTOPOGON, Beauv. = **Aristida**, Linn. (Gramin.).
dichotomus, Spreng l. c. = A. dichotoma.

CYRTORRHYNCHA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 26 (1838) = **Ranunculus**, Tourn.
ranunculina, Nutt. l. c. = R. Nuttallii.

CYRTOSIA, Blume, Bijdr. 396. t. 6 (1825) = **Galeola**, Lour. (Orchid.).
altissima, Blume, l. c.= G. altissima.
javanica, Blume, l. c. = G. javanica.
Lindleyana, Hook. f. & Thoms. in Hook. f. Illustr. Himal. Pl. t. 22 = G. Lindleyana.

CYRTOSIPHONIA, Miq. Fl. Ind. Bat. ii. 401 (1856) = **Rauvolfia**, Linn. (Apocyn.).
amaniensis, Miq. l. c. 402 = R. amsoniæfolia.
maburensis, Teijsm. & Binn. Cat. Hort. Bog. 125.— Java.
reflexa, Miq. Fl. Ind. Bat. ii. 402 = R. reflexa.
spectabilis, Miq. l. c.—Java.
sumatrana, Miq. l. c. 401 = R. sumatrana.

CYRTOSPADIX, C. Koch, Ind. Sem. Hort. Berol. (1853) 13 = **Caladium**, Vent. (Aroid.).
striatipes, C. Koch, l. c. = Cal. striatipes.

CYRTOSPERMA, Griff. Notul. iii. 149 (1851). *AROIDEÆ*, Benth. & Hook. f. iii. 997.
Lasiosperma, Schott, in Bonplandia, v. (1857) 137.
Afzelii, *Engl. in DC. Monog. Phan.* ii. 269.—Afr. trop.
amerlanum, *Engl. in Mart. Fl. Bras.* iii. ii. 117. t. 23. —Bras.
cuspidiflorum, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 284 = Merkusii.
dubium, Schott, l. c. = Merkusii.
edule, Schott, in Bonplandia, ix. (1861) 367 = Merkusii.
intermedium, Schott, in Ann. Mus. Lugd. Bat. i. 284 = Merkusii.
Johnstoni, *N. E. Br. in Gard. Chron.* (1882) ii. 808.— Ins. Salomon.
lasioides, *Griff. Notul.* iii. 150; *Ic. Pl. Asiat.* t. 369. —Malaya.
macrorum, *Becc. ex Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 295.—N. Guin.
Matriefianum, *Hort. ex Gard. Chron.* (1884) i. 711.— Sumatra.
Merkusii, *Schott, in Oestr. Bot. Wochenbl.* (1857) 61. —Sumatra.
senegalense, *Engl. in DC. Monog. Phan.* ii. 270.— Afr. trop.
Sprucеanum, *Engl. in Mart. Fl. Bras.* iii. ii. 118.— Bras.

CYRTOSPERMUM, Benth. in Hook. Kew Journ. iv. (1852) 18 = **Campnosperma**, Thw. (Anacard.).
gummiferum, Benth. l. c. = Camp. gummifera.

CYRTOSPERMUM, Rafin. ex DC Prod. iv. 118 (1830) = **Cryptotaenia**, DC. (Umbellif.).
trifoliatum, Rafin. l. c. 119 = Cryptotaenia canadensis.

CYRTOSTACHYS, Blume, in Bull. Neerl. (1838) 66; ex Rumphia, ii. 101. t. 120 (1848). *PALMÆ*, Benth & Hook. f. iii 892.
ceramica, *H. Wendl. in Kerch. Palm.* 242 —Malaya. Lakka, *Becc. in Ann. Jard. Buitenz.* ii. (1885) 141.— Malaya.
Renda, *Blume, in Bull. Néerl.* (1838) 66; *et Rumphia*, ii. 101.—Sumatra.

CYRTOSTEMMA, Kunze, in Bot. Zeit. i. (1843) 272 = **Clerodendron**, Linn. (Verbenac.).
myricoides, Kunze, l. c. = Clerodendron myricoides.

CYRTOSTEMMA, Spach, Hist. Vég. Phan. x. 321 (1841) = **Scabiosa**, Tourn. (Dipsac.).
atropurpurea, Spach, l. c. 322 = S. atropurpurea.
maritima, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = S. maritima.

CYRTOSTYLIS, R. Br. Prod. 329 (1810). *ORCHIDEÆ*, Benth. & Hook. f. iii. 512.
Huegelii, Endl. in Lehm. Pl. Preiss. ii 6 = reniformis.
oblonga, *Hook. f. Fl. N. Zel.* i. 246.—N. Zel.
reflexa, Spreng. Syst. iii. 714 = Chiloglottis diphylla.
reniformis, *R. Br. Prod.* 322.—Austral.
rotundifolia, *Hook. f. Fl. N. Zel.* i. 246.—N. Zel.

CYRTOTROPIS. Wall. Pl. As. Rar. i. 49. t. 62 (1830) = **Apios**, Moench (Legumin.).
carnea, Wall. l. c. 50. t. 62; Cat. n. 5527 = A. carnea.

CYSSOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) 11. 25 = **Oenanthe**, Tourn. (Umbellif.).
javanicum, Turcz. l. c. 26 = O. stolonifera.

CYSTACANTHUS, T. Anders. in Journ. Linn. Soc. ix. (1867) 457. *ACANTHACEÆ*, Benth. & Hook. f. ii. 1096.
Meninia, Fna, ex Hook. f. Bot. Mag. t. 6048 (1873).
cymosus, *T. Anders. l. c.* 458.—Burma.
insignis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 514.—Burma.
paniculatus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 458.—Burma.
pulcherrimus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 514.—Burma.
tunglda, *Nichols. Dict. Gard.* i. 438.—Cochinch.

CYSTANCHE, Ledeb. Fl. Ross. iii. 314 (1848) = **Cistanche**, Hoffmgg. & Link (Orobanch.).

CYSTANTHE, R. Br. Prod. 555 (1810) = **Richea**, R. Br. (Epacrid.).
acerosa, F. Muell. Fragm. i. 36 = R. acerosa.
Milligani, F. Muell. l. c. = R. Milligani.
procera, F. Muell. l. c. = R. procera.
sprengelioides, R. Br. Prod. 555 = R. sprengelioides.

CYSTIBAX, Heynh. Nom. ii. 184 (1846), sphalm. = **Cybistax**, Mart. (Bignon.).

CYSTICAPNOS, Boerh. ex Linn. Class. Pl. 181 (1747); Gaertn. Fruct. ii. 161. t. 115 (1791) = **Corydalis**, Vent. (Papav.).
africana, Gaertn. l. c. = Coryd. vesicaria.
alexandrina, Link & Otto, ex G. Don, Gen. Syst. i. 141 = Coryd. vesicaria ?
cirrhosa, Moench, Meth. 52 = Coryd. vesicaria.
grandiflora, E. Mey. Zwei Pfl. Docum. 177 = Corydalis Burmanni.

CYSTIDIANTHUS, Hassk. in Hoev. & De Vriese, Tydschr. x. (1843) 125 = **Physostelma**, Wight (Asclep.).
campanulatus, Hassk. l. c. = P. Wallichii.
laurifolius, Blume, Mus. Bot. Lugd. Bat. i. 57.—Java.

CYSTIUM, Stev. in Bull. Soc. Nat. Mosc. iv (1832) 288 = **Astragalus**, Tourn. (Legumin.).
Cicer, Stev. l. c. = A. Cicer.
macrophyza, Stev. l. c. = A. physodes ?
Pallasii, Stev. l. c. = A. Pallasii.
physodes, Stev. l. c. = A. physodes.
Pseudocicer, Opiz, in Lotos, viii. (1858) 65 = A. Cicer.

CYSTOCARPUM, Benth. & Hook. f. Gen. i. 73 (1862) = CISTOCARPIUM, Spach = **Vesicaria**, Lam. (Crucif.).

CYSTOCHILUM, Rodrig. Orch. Novo. i. 197 (1877). *ORCHIDEÆ*.
candidum, *Rodrig. l. c.* 198.—Bras.

CYSTOGYNE, Gasp. Ricerch. Caprif. e Fic. 84. t. 3 (1845) = **Ficus**, Tourn. (Urticac.).
leucocista, Gasp. l. c. = F. leucocista.

CYSTOPUS, Blume, Orch. Archip. Ind. 82 (1858) = **Odontochilus**, Blume (Orchid.).
elongatus, Blume, l. c. 84.—Java.
Hasseltii, Blume, l. c. 86.—Java.
occultus, Blume, l. c. 85.—Java.
pubescens, Blume, l. c. 86.—Java.
spicatus, Blume, l. c. 84.—Borneo.
unifiorus, Blume, l. c. 83.—Java.

CYSTORCHIS, Blume, Orch. Archip. Ind. 87. t. 24, 36 (1858). *ORCHIDEÆ*, Benth. & Hook. f. iii. 599.
glandulosa, *Benth. & Hook. f. Gen.* iii. 598, in nota sub Physuro.—Borneo.
fusca, Benth. & Hook. f. l. c. 599 = Hetaeria fusca.
javanica, *Blume, Orch. Archip. Ind.* 87.—Java.
marginata, *Blume, l. c.* in nota.—Java.
nebularum, *Hance, in Journ. Bot.* xxi. (1883) 282.— China.
obscura, *Blume, Orch. Archip. Ind.* 89.—N. Guin.
variegata, *Blume, l. c.*—Java.

CYSTOSTEMMA, Fourn. in Mart. Fl. Bras. vi. iv. 204 (1885). *ASCLEPIADEÆ*.
umbellatum, *Fourn. l. c.*—Bras.

CYSTOSTEMON, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 89. *BORAGINEÆ*.
socotranum, *Balf. f. l. c.*—Ins. Socotr.

CYTHAREXYLON, Batsch. Tab. 198 (1802) = seq.

CYTHAREXYLUM, Jacq. Enum. Pl. Carib. 26 (1760) = **Citharexylon**, Linn. (Verben.).

CYTHENAEA, Wight & Arn. Prod. 178, in textu (1834) = **Spondias**, Linn. (Anacard.).
dulcis, Wight & Arn. l. c. = S. dulcis.

CYTHEREA, Salisb. in Trans. Hort. Soc. i. (1812) 301 = **Calypso**, Salisb. (Orchid.).
borealis, Salisb. l. c. = Cal. borealis.

CYTHERIS, Lindl. in Wall. Cat. n. 8750 (1830?); et Gen. et Sp. Orch. 128 (1831) = **Nephelaphyllum**, Blume (Orchid.).
cordifolia, Lindl. ll. cc. 8750, 129 = N. cordifolium.
Griffithii, Wight, ic. v. t. 1751–52 = Calanthe ventia.

CYTHISUS, Schrank, Prim. Fl. Salisb. 179 (1792) = **Cytisus**, Linn. (Legum.).

CYTICUS, Link, Handb. ii. 153 (1831), sphalm. = prsec.

CYTINUS, Linn. Gen. ed. VI. 567 (1764). *CYTINACEÆ*, Bentb. & Hook. f. iii. 118.
Bdallophyton, Eichl. in Bot. Zeit. xxx. (1872) 709. t. 8.
Haematolepis, Presl, Epim. Bot. 288 (1849).
Hypocistis, Adans. Fam. ii. 78 (1763).
Hypolepis, Pers. Syn. ii. 598 (1807).
Phelypaea, Thunb. Nov. Gen. Pl. 91 (1784).
Thyrsine, Gleditsch, Syst. Pl. 286 (1764).
americanus, *R. Br. in Trans. Linn. Soc.* xix. (1845) 246.—Mexic.
americanus, Hook. f. in DC. Prod. xvii. 108 = Andrieuxii.
Andrieuxii, *Hemsl. Biol. Centr. Am. Bot.* iii. 41.—Mexic.
canariensis, Webb & Berth. Phyt. Canar. iii. 429 = C. Hypocistis.
Clusii, Nym. Comp. 646 = C. Hypocistis.
dioicus, *Yasz. in Ann. Mus. Par.* xii. (1808) 448.—Af. austr.
Hypocistis, *Linn. Gen.* ed. VI. 566; *Syst. Nat.* ii. 602. —Reg. Mediterr.; Oriena.

CYTISOPSIS, Jaub. & Spach, Illustr. i. 154. t. 84 (1842). *LEGUMINOSÆ*, Benth. & Hook. f. i. 489.
doryenifolia, *Jaub. & Spach, l. c.* 155.—Syria.

CYTISUS, Linn. Syst. ed I (1785). *LEGUMINOSÆ*, Benth. & Hook. f. i. 488.
Aulonix, Rafin. Sylva Tellur. 25 (1838).
Chamaecytisus, Link, Handb. ii. 154 (1831).
Chamaespartium, Adans. Fam. ii. 321 (1763).
Dianzlon, Rafin. Sylva Tellur. 24 (1838).
Lembotropis, Griseb. Spicil. Fl. Rumel. i. 10 (1846).
Lygolis, Rafin. Sylva Tellur. 23 (1838).
Meiemianthera, Rafin. l. c. 25 (1838).
Nubigena, Rafin. l. c. 23 (1838).
Peyronellia, Boiv. ex Webb & Berth. Phyt. Canar. ii. 85, in obs. (1846).
Phyllocytisus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 358.
Sarothamnus, Wimm. Fl. Schles. 278 (1832).
Spartocytisus, Webb & Berth. Phyt. Canar. ii. 49 (1846).
Spartiothamnus, Webb. ex Walp. Ann. i. 221 (1848).
Telinaria, Presl, Bot. Bemerk. 49, 135 (1844).
Teline, Webb & Berth. Phyt. Canar. ii. 84 (1846).
Tubocytisus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 358.
Verzinum, Rafin. Sylva Tellur. 28 (1838).
Virgilia, Moench, Meth. 132 (1794).
absinthioides, *Yenke, in Oestr. Bot. Zeitschr.* xxii. (1872) 176.—Macedon.
acutangulus, *Yaub. & Spach, Illustr.* ii. 70. t. 158.—Galatia.
Adamii x, Polt. in Ann. Soc. Hort. Par. vii. (1880) 501.
aeolicus, *Guss. ex Lindl. Bot. Reg.* t. 1902.—Ins. Lipar.
aethiopicus, Linn. Sp. Pl. 740 = Melolobium cernuum.
affinis, *Boiss. Elench.* 81.—Hispan.
affinis, *Presl, Fl. Sic.* t. p. xix.—Sicil.
africanus, *Loisel. in Duham. Arb. ed. Nov.* v. 154.—Af. bor.
aggregatus, Schur, Enum. Pl. Transe. 149 = prostratus.
albidus, *DC. Cat. Hort. Monsp.* 101. n. 85.—Marocco; Ins. Canar.
albus, Jacquet. Reise Sarm. Karp. i. 49 = austriacus.
albus, *Link, Enum. Hort. Berol.* ii. 241.—Reg. Mediterr. occ.
alpestris, Schur, Enum. Pl. Transe. 148 = supinus.
alpinus, *Lam. Fl. Fr.* ii. 621 = Laburnum vulgare.
alpinus, Mill. Gard. Dict. ed. VIII. n. 2 = Laburnum alpinum.
Alschingeri, *Vis. in Linnaea*, xv. (1841) *Litt.* 101.—Dalmatia.
ambiguus, Schur, Enum. Pl. Transe. 147 = austriacus.
anagyrifolius, Fourr. ex Colm. sec. Willk. & Lange, in Prod. Fl. Hisp. iii. 462 = Adenocarpus hispanicus.
CYTISUS:—
anagyrius, L'Hérit. Stirp. Nov. 184 = Adenocarpus hispanicus.
angulatus, Boiss. ex Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 153 = Genista parviflora.
angustifolius, Moench, Meth. 145 = Laburnum alpinum.
antillanus, *DC. Prod.* ii. 157.—Ins. Carib.
apulus, Tenore & Guss. ex Nym. Syll. 288 = apbinescens.
arabicus, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 78 = Argyrolobium arabicum.
arboreus, *DC. Prod.* ii. 154.—Afr. bor.
arcuatus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1876) 100 = Genista sylvestris.
Ardoini, *Fourn. in Bull. Soc. Bot. Fr.* xiii. (1866) 389. —Reg. Mediterr.
argenteus, Linn. Sp. Pl. 740 = Argyrolobium Linnaeanum.
argenteus, Schur, in Verh. Naturf. Ver. Brunn, xv. 11. (1877) 167 = austriacus.
argyreus, Reichb. Fl. Germ. Excurs. 523 = spinescens.
arvensis, *Vell. Fl. Flum.* 309; vii. t. 117.—Bras.
aspalathoides, Spreng. Syst. iii. 235 = Genista aspalathoides.
atratus, Schur, Enum. Pl. Transe. 147 = nigricans.
atropurpureus, Herb. Pav. ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 91 = Lonchocarpus atropurpureus.
auranticus, *Regel, Cat. Pl. Hort. Ataskov.* 46.—Hab.!
austriacus, *Linn. Sp. Pl.* ed. II. 1042.—Europ.; Reg. Cauc.
austriacus, Medic. Beobacht. (1782) 174 = polycephalus.
baeticus, *Steud. Nom.* ed. II. i. 477.—Reg. Mediterr. occ.
Balansae, *Ball, in Journ. Bot.* xi. (1873) 303.—Orient.
banaticus, Griseb & Schenk, in Wiegm. Archiv, xviii. (1852) i. 292 = austriacus.
Barrelieri, *Ten. in Ann. Sem. Hort. Bot. Panorm.* (1857) 41.—*Cf. Linnaea*, xxix. (1857–58) 748.—Ins. Lipar.
biflorus, Ker-Gawl. in Bot. Reg. t. 808 = elongatus.
biflorus, *L'Hérit. Stirp. Nov.* 184.—Europ.; An bor. *bifloriens*, Host, Fl. Austr. ii. 340 = prostratus.
bouvista, Vell. Fl. Flum. 309; vii. t. 115 (= *Collaea gymnocalycina*).—Bras.
borysthenicus, *Gruner, in Bull. Soc. Nat. Mosc.* xli. (1868) ii. 446, in textu.—Rossia.
brasiliensis, *Vell. Fl. Flum.* 308; vii. t. 114.—Bras.
Cajan, Linn. Sp. Pl. 739 = Cajanus indicus.
calycinus, Bieb. Fl. Taur. Cauc. ii. 166 = Argyrolobium calycinum.
canariensis, *Steud. Nom.* ed. I. 259.—Ins. Canar.
candicans, *Lam. Fl. Fr.* ii. 622; *Boiss. Fl. Orient.* ii. 54.—Reg. Mediterr. occ.
candidus, *Presl, Fl. Sic.* p. xix.—Sicilia.
canescens, *Loisel. in Duham. Arb. ed. nov.* v. 151.—Hab.!
canescens, J. & C. Presl, Delic. Prag. 229 = austriacus.
capensis, Berg. Descr. Pl. Cap. 217 = Rafnia opposita.
capensis, Lam. Encyc. ii. 249 = Lebeckia cytisoides.
capitatus, *Scop. Fl. Carn.* ed. II. ii. 70; *Jacq. Fl. Austr.* v. 38.—Europ. austr.
caramanicus, *Nym. Consp.* 155.—As. Min.
carneus, *Hoffmgg. Prodromus*, ed. X. 275; *ex Linnaea*, xvi. (1842) *Litt.* 276, 276.—Hab.!
cassius, *Boiss. Diagn.* Ser. I. ix. 5.—Syria.
Cayan, Linn. ex Mill. Gard. Dict. ed. VIII. n. 11 = Cajanus indicus.
Chamaecytisus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 104 = austriacus!
chrysobotrys, *Fisch. in Otto & Dietr. Allg. Gartenz.* v. (1837) 122.—Hab.!
chrysotrichus, Boiss. Diagn. Ser. I. ii. 12 = pygmaeus.
ciliatus, Wahlenb. Fl. Carp. 219 = capitatus.
cinctutus, *Ball, in Journ. Linn. Soc.* xvi. (1878) 404.—Marocco.
cinereus, Host, Fl. Austr. ii. 343 = biflorus.
cinereus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 97 = Genista cinerea.
collinus, Schur, in Verh. Naturf. Ver. Brunn, xv. 11. (1877) 169 = biflorus.
communis, Lindem. in Bull. Soc. Nat. Mosc. xl. (1867) i. 404 = biflorus!
CYTISUS:—
complicatus, Raeusch. Nom. ed. III. 212 = Adenocarpus intermedius.
confertus, Schur, Enum. Pl. Transe. 148 = capitatus.
congestus, *Ball, in Journ. Linn. Soc.* xvi. (1878) 404. —Ins. Canar.
cornicalatus, *Gmeldenst. ex Ledeb. Fl. Ross.* i. 521.—Rossia.
coronensis, Schur, Enum. Pl. Transe. 148 = supinus.
creticus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 5.—Ins Creta.
dalmaticus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi (1875) 101 = Genista arlstata?
decaphyllus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 10 (Quid?).—Java.
decumbens, *Spach, in Ann. Sc. Nat.* Sér. III. iii (1845) 156.—Europ.
demissus, *Boiss. Fl. Orient.* ii. 54.—Thessalia.
diffusus, Jacquem. ex Jaub. & Spach. in Ann. Sc. Nat Sér. II. xix. (1843) 48 = Argyrolobium flaccidum.
diffusus, Vis. Fl. Dalmat. iii. 269 = decumbens.
divaricatus, L'Hérit, Stirp. Nov. 184 = Adenocarpus intermedius.
Dorycnium, Fourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 439 = Genista umbellata.
drepanolobus, *Boiss. Diagn.* Ser. I. ix. 6.—Syria.
elongatus, Friveld. ex Nym. Consp. 156 = hirsutus.
elongatus, Waldst. & Kit. Pl. Hung. ii. 200. t. 188 = biflorus.
emeriflorus, Reichb. Fl. Germ. Excurs. 524 = glabrescens.
eriocarpus, *Boiss. Diagn.* Ser. I. ii. 11.—As. Min.
Everestianus, Carr. in Rev. Hortic. (1872) 299; (1873) 390. cum t. = racemosus!
falcatus, DC. Prod. ii. 155 = biflorus.
falcatus, Hort. ex Presl, Bot. Bemerk. 140 = speciosus
falcatus, Waldst. & Kit. Pl. Rar. Hung. iii. 264. t. 238 = hirsutus.
ferox, Hoffmgg. Verz. Pfl. 55 = Genista ferox.
filiformis, Spreng. Syst. iii. 235 = Indigofera sarmentosa.
filipes, *Webb & Berth. Phyt. Canar.* ii. 52. t. 47.—Ins. Canar.
flaccidus, Royle, Illustr. Bot. Himal. 197 = Argyrolobium flaccidum.
foliolosus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 49 = Adenocarpus foliolosus.
foliosus, L'Hérit. Stirp. Nov. 184. t. 96 = Adenocarpus anagyrus.
Fontanesii, *Spach, ex Ball, in Journ. Linn. Soc.* xvi (1878) 405.—Reg. Mediterr. occ.
fragrans, *Lam. Encyc.* ii. 249.—Ins. Teneriff.
fragrans, Weld. in Flora, xv. (1832) 326 = Petteria ramentacea.
frutescens, Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 102 = Genista tinctoria.
gallicus, Kern. ex Willk. & Lange, Prod. Fl. Hisp iii. 464 = supinus.
genistoides, *Regel, Gartenfl.* (1858) 340.—Hab.!
germanicus, Vis. Fl. Dalmat. iii. 268 = Genista germanica.
glaber, Lam. Fl. Fr. ii. 621 = nigricans, sessilifolius.
glaber, Linn. f. Suppl. 325 = biflorus.
glabratus, Link, in Buch, Beschr. Ins. Canar. 156 = scoparius.
glabrescens, *Sartor. Alb. Indig. Ital. Sup.* 382.—Europ. glabrescens, *Schrank, Baier. Fl.* ii. 269.—Bavar.
glomeratus, Boj. Hort. Maurit. 89 = Erioscma parviflorum.
gracilis, Linn. Sp. Pl. 740 = Anthyllis Hermanniae.
grandiflorus, *DC. Prod.* ii. 154.—Lusitan.
graniticus, Rehm. Notis. Veg. Schwarz. Meer. 59 = biflorus!
guineensis, Schum. & Thonn. Beskr. Guin. Fl. 349 = Cajanus indicus.
Hermesii, *Ball, in Journ. Bot.* xi. (1873) 304.—Hispan.
heptaphyllus, Vis. Fl. Dalmat. iii. 268; vii. t. 13 = Lupinus Hilarianus.
heterocaulicus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 100 = Genista germanica.
heterochrous, Webb, ex Willk. & Lange, Prod. Fl. Hisp. iii. 462 = patens.
heterophyllus, *Lagasc. Hist. Adv. Pl. Pyr.* 422.—Mons. Pyren.
Heufelii, Wierzb. ex Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 292 = austriacus.
hirsutissimus, C. Koch, in Linnaea, xix. (1847) 92 = hirsutus.

CYTISUS:—
hirsutus, Cranta, Stirp. Austr. ed. II. fasc. v. 398 = capitatus.
hirsutus, Jacq. Obs. ex Steud. Nom. ed. II. i. 477 = candicans.
hirsutus, *Linn. Sp. Pl.* 789.—Europ. austr.; As. Min.
hirsutus, Pall. Reise, i. 21 = biflorus.
hirsutus, Sibth. & Sm. Fl. Graec. t. 706 = spinescens.
hirtellus, Reichb. ex Nym. Consp. 157 = pygmaeus.
hirtus, Burm. f. Fl. Cap. Prod. 22 = Cneorus aspalathoides.
hispanicus, Lam. Encyc. ii. 248 = Adenocarpus hispanicus.
hispanicus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 100 = Genista hispanica.
hispidus, Willd. Sp. Pl. iii. 1121 = Robinia guineensis.
hoppeanus, Fleisch. in Koch, Syn. Fl. Germ. ed. II. i. 441 = Genista radiata.
hosmariensis, *Ball, in Journ. Linn. Soc.* xvi. (1878) 404.—Marocc.
humifusus, Nym. Consp. 158 = decumbens.
incanus. *Regel, Cat. Pl. Hort. Absakov.* 48.—Hab.?
incarnatus, C. Koch, Dendrol. i. 32 = alburn.
infestus, Guss. Fl. Sic. Prod. ii. 372 = Calycotome infesta.
intermedius, *Salam. ex Presl, Bot. Bemerk.* 51.—Afr. bor.
junceus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 108 = Spartium junceum.
Kitaibelii, Vis. Fl. Dalmat. iii. 209 = procumbens Kunzeanus, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 452.—Hispan.
laburnum, Linn. Sp. Pl. 739 = Laburnum vulgare.
Lamarckii, Tenore, Fl. Nap. v. 94 = hirsutus.
lanceolatus, Herb. Pav. ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 92 = Lonchocarpus lanceolatus.
lanigerus, DC. Prod. ii. 154 = Calycotome villosa.
lanuginosus, Royle, Illustr. Bot. Himal. 197 = Argyrolobium flaccidum.
lasiosemius, *Boiss. in Tchihat. Asie Min. Bot.* i. 12.—As. Min.
leiocarpus, Kern. in Oestr. Bot. Zeitschr. (1868) 90 = biflorus.
leucanthus, Waldst. & Kit. Pl. Rar. Hung. ii. 141. t. 132 = austriacus.
leucotrichus, Schur, in Oestr. Bot. Zeitschr. x. (1860) 179 = hirsutus.
linifolius, *Lam. Fl. Fr.* ii. 624.—Reg. Mediterr. occ
Linkii, *Janka, in Termész. Füzetek*, viii. (1884) 70.—Europ.
lithuanicus, *Gilib. Hist. Pl. Eur.* ii. 275.—Lithuania.
Loeblii, Tausch, in Flora, xxi. (1838) 739 = sessiliflorus.
lotoides, Pourr. in Mém. Acad. Toul. iii. (1788) 318 = supinus.
lotoides, Willd. Sp. Pl. iii. 1137 = Argyrolobium calycinum.
lupinus, Gromow, ex Trautv. in Act. Hort. Petrop. viii. (1883) 196 = supinus?
macrospermus, Bess. ex DC. Prod. ii. 155 = biflorus.
maderensis, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 149.—Madera.
malacitanus, *Boiss. Elench.* 82.—Hispan.
Malyei, Steud. Nom. ed. II. i. 477 = austriacus.
Mannii, Hook. f. in Journ. Linn. Soc. vi. (1862) 8 = Adenocarpus Mannii.
medicaginifolius, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 460 = patens.
microphyllus, Boiss. Diagn. Ser. II. ii. 8 = austriacus, microphyllus, *Lénh, in Buch, Beschr. Ins. Canar.* 156.—Ins. Canar.
Millieri, *Steud. Nom.* ed. II. i. 477.—Lusitan.
mollis, Salisb. Prod. 330 = Cajanus indicus.
mollis, *Willd. Enum. Hort. Berol. Suppl.* 51.—Hab.?
monspessulanus, *Linn. Sp. Pl.* 740.—Europ.
multiflorus, Lindl. Bot. Reg. t. 1191 = hirsutus.
multiflorus, Sweet, Hort. Brit. ed. I. 112 = albus.
myrtifolius, Nym. Consp. 157 = Kitaibelii.
nanus, Sieber, ex Presl, Bot. Bemerk. 51 = candidus.
nanus, Willd. Enum. Hort. Berol. 769 = Argyrolobium calycinum.
nigricans, Linn. Mant. 444 = triflorus.
nigricans, *Linn. Sp. Pl.* 739.—Europ. austr.
nigricans, Pall. Reise, iii. 754 = Calophaca wolgarica.
nubigenus, Link, Enum. Hort. Berol. ii. 240 = fragrans.
obcuneus, Sehm, Enum. Pl. Transs. 147 = austriacus.
obcuneiformis, Schur, l. c. = austriacus.

CYTISUS:—
onosmus, *Tausch. in Flora*, xxi. (1838) 746.—Hab.?
orientalis, *Loisel. in Duham. Arb. ed. Nov.* 156.—As. Min.
ovatus, Vukot. in Rad Jugos. Akad Zagreb. xxxi. (1875) 101 = Genista ovata.
ovalifolius, C. A. Mey. ex Boiss. Fl Orient. ii. 33 = Argyrolobium trigonelloides.
Paivae, Masf. in Anal. Soc. Esp. Hist. Nat. x. (1881) 149 = Genista Paivae.
pallidus, Kit. ex Nym. Consp. 157 = capitatus.
pallidus, *Poir. Encyc. Suppl* ii. 442.—Ins. Canar.
pallidus, Steud. Nom. ed. II. i. 477 = austriacus.
paniculatus, Loisel. in Duham. Arb. ed. Nov. v. 146 = canariensis.
parviflorus, *Boiss. Diagn.* Ser. I. ii 9.—Anatolia.
parviflorus, Hemsl. Pav. ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 91 = Lonchocarpus stropurpureus.
parvifolius, Lam. Fl. Fr. ii. 623 = Adenocarpus intermedius.
patens, *Murr. Syst. Veg.* ed. XIII. 555.—Lusitan.
pauciflorus, Willd. Sp. Pl. iii. 1126 = Argyrolobium calycinum.
pauciflorus, Boiss. Fl. Orient. ii. 1119, sphalm. = praec.
penduliflorus, Stokes, Bot. Mat. Med. iv. 12 = Laburnum vulgare.
pendulinus, Linn. f. Suppl. 328 = patens.
pendulus, Salisb. Prod. 330 = Laburnum vulgare.
pentaphyllus, Salzm. ex Walp Rep. i. 623 = Lotononis lupiniflora.
peregrinus, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 101.—Hab.?
persicus, Burm. f. Fl. Ind. 168. t 51 f. 1 = Crotalaria furfuracea.
philippicanus, Boiss. Fl. Orient. iii. 1119, sphalm. = praec.
pinnatus, Linn. Sp. Pl. 741 = Pongamia glabra.
pinnatus, Pall. Fl Ross. i. 73. t. 47 = Calophaca wolgarica.
plumosus, Spach, ex Nym Consp. 157 = Fontanesii.
polyopethalus, *Tausch, in Flora*, xxi. (1838) t. *Betbl.* 78.—Hab.?
polymorphus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 460 = Adenocarpus intermedius.
polytrichus, *Boiss. Fl. Taur.-Cauc.* iii. 477.—Tauria.
pontiacus, Griseb. Spicil. Fl. Rumel. i. 9 = hirsutus.
pontiacus, Willd. Sp. Pl. iii. 1120 = Adenocarpus intermedius.
Presslii, Nym. Syll. 283 = Calycotome villosa.
procera, *Link, Enum. Hort. Berol.* ii. 241.—Lusitan.
procumbens, Boj. ex Baker, in Journ. Linn. Soc. xx (1883) 134 = Eriosema procumbens.
procumbens, *Sprengl. Syst.* iii. 224.— Europ.
procumbens, *Vell. Bl. Flum.* 309; vii. t. 118.—Bras.
prolifer, Kit. in Linnaea, xxxii. (1863) 611 = pullulans.
prolifera, *Linn. f. Suppl.* 328.—Ins. Teneriff.
propinquus, Schur, Enum. Pl. Transs. 147 = austriacus.
proteiformis, *Scop. Fl. Carn.* ed. II. ii. 70.—Carn.; Tyrol.
proteus, *Zumag. Fl Pedem* ii. 192.—Ital.
Pseudocajan, Jacq. Hort. Vindob. ii. 54 = Cajanus indicus.
Pseudo-nigricans, Schur, Enum. Pl. Transs. 147 = nigricans.
pseudoradiatus Linn. Pl. Rar. Afr. 15; Sp. Pl. 1043 = Indigofera paoraloides.
pterocladus, Boiss. Diagn. Ser. I. ii. 10 = Genista parviflora.
pubescens, Gillib. Fl. Lituan. ii. 81 = supinus.
pubescens, Moench, Meth. Suppl. 43 = candicans, *Lam.*
pulchellus, *Vis. Fl. Dalmat.* iii. 270.—Dalmat.
pullulans, *Kit. in Linnaea*, xxxii. (1863) 611.—Hungar.
pumilus, De Not. ex Nym. Consp. 156 = hirsutus.
pungens, *Sprengl. Syst.* iii. 235.—Ins. Maurit.
purgans, *Spach, in Ann. Sc. Nat.* Sér. III. iii. (1845) 156.—Europ.
purpureus, *Scop. Fl. Carn.* ed. II. ii. 69.— Europ. austr.
pygmaeus, *Willd. Sp. Pl.* iii. 1127.—Macedon.
quinquepetalus, Blanco, Fl. Filip. ed. I. 598 = Desmodium Cephalotes.
racemosus, *Hort.—Cf. Marnoch, Fl. Mag. & Misc. Gard.* ii. (1837) 37. t. 19.—Ins. Teneriff.

CYTISUS:—
radiatus, Koch, Syn. Fl. Germ. ed. I. 167 = Genista radiata.
ramentaceus, Sieber, in Flora, v. (1822) 242 = Petteria ramentacea.
ramosissimus, Poir. Encyc. Suppl. ii. 440 = canariensis.
ramosissimus, Tenore, Syll. 343 = spinescens.
ratisbonensis, Schaeff. Bot. Exp. (1760) tab. in prim. lib.; Koch, Syn. Fl. Germ. ed. II. 171 = biflorus
repens, Wolf. in Lotos, iv. (1854) 176 = hirsutus.
rhombifolius, Parsh, Fl. Am. Sept. ii. 741 = Thermopsis rhombifolia.
Rochelii, Wierzb. ex Griseb. & Schenk, in Wiegm Archiv, xviii. (1852) i. 393 = austriacus.
rostrus, Cambess. in Jacquem. Voy. Bot. 35. t. 40 = Argyrolobium roseum.
rosmarinifolius, *Masf. in Anal. Soc. Esp. Hist. Nat.* v. (1881) 149.—Ins. Canar.
rufulus, *Y. & C. Presl, Delic. Prag.* 36.—Ital.
ruthenicus, Fisch. Ind. Pl. Hort. Petrop. (1824) 25 = biflorus.
sagittalis, Koch, Syn. Fl. Germ. ed. I. 167 = Genista sagittalis.
scariosus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi (1875) 100 = Genista triangularis.
scopulensis, *Kit. in Lunnaea*, xxxii. (1863) 609.—Hungar.
Schimperianus, Steud. ex Baker, in Oliver, Fl. Trop Afr. n. 46 = Argyrolobium Schimperianum.
scoparius, *Link, Enum. Hort. Berol.* ii. 241.—Europ.; Japon.
sericeus, Noronha, in Verh. Batav. Gen. v. (1790) Art. iv. 9 (sericeus) = Cajanus indicus.
sericeus, *Vis Fl. Dalmat.* iii. 269.—Dalmat.
sericeus, Willd Sp. Pl. iii. 1121 = Mundulea sericea.
serotinus, Kit. ex DC. Prod. ii. 156 = biflorus.
sessilifolius, Poir. Encyc. Suppl. ii. 439 = Tephrosia brevipes.
sessilis, Mill. Gard Dict. ed. VIII. n. 4 = sessiliflorus.
sessiliflorus, *Linn. Sp. Pl.* 739.—Europ. austr.
smyrnaeus, *Boiss. Diagn.* Ser. I. ii. 10.—As. Min.
sordidus, Bot. Reg. ex C. Koch, Dendrol. i. 17 = *Adami.*
speciosus, *Presl, Bot. Bemerk.* 140.—Hab.?
spinescens, Presl, Fl. Sic. p. xix (=seq.).—Ital.
spinescens, *Sieber, in Sprengl. Syst.* iii. 225.—Ital.; Graec.
spinosus, Desc. Fl. Med. Ant. i. t. 8 = antillanus.
spinosus, Koch, Syn. Fl. Germ. ed. II. 170 = Calycotome infesta.
spinosus, Lam. Fl. Fr. ii. 625 = Calycotome spinosa.
spinosus, Sieber, ex Steud. Nom. ed. II. i. 477 = spinescens.
strigulosus, Gren. & Godr. Fl. Fr. i. 508, sphalm. = virgultatus.
subleiocarpus, Simk. in Termész. Füzetek, iv. (1885) 44 = leiocarpus.
supinus, Cranz, Stirp. Austr. ed. II. fasc. v. 398; Jacq. Fl. Austr. i. 15. t. 20 = biflorus.
supinus, Griseb. Spicil. Fl. Rumel. i. 7 = pygmaeus.
supinus, *Linn. Sp. Pl.* 740.—Europ.
supinus, Pall. Ind. Taur. ex Steud. Nom. ed. II. i. 478 = hirsutus.
sylvestris, *Vis. Fl. Dalmat.* iii. 268.—Dalmat.
syriacus, *Boiss. Fl. Orient.* ii. 55.—Syria.
telonensis, Loisel. Fl. Gall. 446 = Adenocarpus grandiflorus.
tener, Jacq. Ic. Rar. iii. t. 556 = Genista virgata.
Teneriffae, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 101.—Ins. Teneriff.
tenerrimus, Steud. Nom. ed. II. i. 477, in syn. = praec.
thessalus, Heldr. & Orph. ex Nym. Consp. 157 = austriacus.
Thirkeanus, C. Koch, in Linnaea, xix. (1847) 61 = pygmaeus.
Thommasini, Vis. ex Reichb. Ic. Fl. Germ. t. 2077 = capitatus.
tinctorius, Vis. Fl. Dalmat. iii. 268 = Genista tinctoria tmoleus, *Boiss. Diagn.* Ser. I. ii. 11.—As. Min.
tomentosus, Andr. Bot. Rep. t. 237 = Argyrolobium Andrewsianum.
Tommasinii, Vis. Fl. Dalmat. iii. 265 = capitatus.
Tournefortianus, Loisel. in Duham. Arb. ed. Nov. v. 187 = hirsutus.
tragacanthaeformis, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 421 = Genista horrida.

CYTISUS:—
triangularis, Vis. Fl. Dalmat. iii. 268 = Genista triangularis.
tribracteolatus, *Webb, Otia Hisp.* 8. t. 8.—Hispan.
tridentatus, *Vukot. in Rad Jugos. Akad. Zagreb.* xxxi. (1875) 104.—Reg. Mediterr. occ.
trifloras, *L'Héréit. Stirp.* 184.—Reg. Mediterr. occ.
unibracteatus, Lindem. in Bull. Soc. Nat. Mosc. (1850) ii. 471 = nigricans?
uniflorus, Decne. in Ann. Sc. Nat. Sér. II. iii. (1885) 265 = Argyrolobium uniflorum.
unifoliatus, *Vell. Fl. Flum.* 308; vii. t. 113.—Bras.
wralensis, Hort. ex Ledeb. Fl. Ross. i. 819 = capitatus.
Villarsii, Vis. Fl. Dalmat. iii. 270. t. 58. f. 1 = Genista Villarsii.
villosus, Pourr. in Mém. Acad. Toul. iii. (1788) 317 = triflorus.
villosus, *Ş. & C. Presl. Delic. Prag.* 36.—Austria.
violaceus, Aubl. Fl. Gui. ii. 756 = Eriosema violaceum.
virgatus, Salisb. Prod. 330 = nigricans.
virgatus, Vest, in Syll. Ratisb. i. (1824) 23 = hirsutus.
virgatus, Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 102 = Genista virgata.
virgulatus, *Reichb. Fl. Germ. Excurs.* 524, in obs.—Corsica.
viscosus, *Hill, Hort. Kew.* 288.—Lusitan.
volubilis, Blanco, Fl. Filip. ed. I. 599 = Atylosia mollis.
Weldeni, Vis. in Flora, xiii. (1830) 53 = Pitteria ramentacea.
wolgaricus, Linn. f. Suppl. 327 = Calophaca wolgarica.
Zanoni, Turra, Fl. Ital. Prod. 66 = Argyrolobium Linnæanum.

CYTTARIUM, Peterm. Fl. Lips. Excurs. 608 (1838) (Compos.).
arenarium, Peterm. l. c. = Helichrysum arenarium.
dioicum, Peterm. l. c. = Antennaria dioica.
silvaticum, Peterm. l. c. 609 = Gnaphalium sylvaticum.

CZACKIA, Andrz. Dias. (1818); Spreng. Syst. ii. 90 = **Paradisea**, Mazz. (Liliac.).
Liliastrum, Andrz. l. c. = P. Lillastrum.

CZEKELIA, Schur, in Oestr. Bot. Wochenschr. (1856) 287 = **Musoari**, Tourn. (Liliac.).
transylvanica, Schur, l. c. = M. botryoides.

CZERNAEVIA, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 789 = **Archangelica**, Hoffm. (Umbellif.).
Conioselinum, Turcz. l. c. 740.—Dahuria.
laevigata, Turcz. l. c. = Feucedanum latifolium.

CZERNIAEVIA, Turcz. ex Ledeb. Fl. Ross. iv. 422 (1853) = **Deschampsia**, Beauv. (Gramin.).
meliocides, Turcz. ex Ledeb. l. c.; et in Bull. Soc. Nat. Mosc. xxix. (1856) i. 82 = D. caespitosa.

CZERNIAJEVIA, Turcz. in Bull. Soc. Nat. Mosc. (1838) 93 = CZERNAEVIA, Turcz. = **Archangelica**, Hoffm. (Umbell.).

CZERNYA, Presl, Cyp. et Gram. Sicul. 22 (1818) = **Phragmites**, Trin. (Gramin.).
arundinacea, Presl, l. c. = P. communis?

D

DABEOCIA, C. Koch, Dendrol. ii. 11. 182 (1872) = seq.

DABOECIA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 160. *ERICACEAE*, Benth. & Hook. f. ii. 596.
Borretta, Neck. Elem. i. 212 (1790).
cantabrica, C. Koch, Dendrol. ii. 11. 18 = polifolia.
polifolia, *D. Don, in Edinb. N. Phil. Journ.* xvii. (1834) 160.—Europ.

DACHEL, Adans. Fam. ii. 26 (1763) = **Phoenix**, Linn. (Palm.).

DACRYDIUM, Soland. ex Forst. f. Pl. Escul. 80 (1786). *CONIFERAE*, Benth. & Hook. f. iii. 433.
Lepidothamnus, Phil. in Linnæa, xxx. (1859–60) 730.
araucaroides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 496.—N. Caled.
arthrotaxoides, Carr. Conif. ed. II. 697 = araucaroides.
Balansae, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xvi. (1869) 828.—N. Caled.
Beccarii, *Parl. in DC. Prod.* xvi. 11. 494.—Borneo.
Bidwillii, *Hook. f ex T. Kirk, in Trans. N. Z. Inst.* x. (1878) 388.—N. Zel.
Colensoi, *Hook. fc. Pl.* t. 548.—N. Zel.
cupressiforme, Carr. Conif. 491 = seq.
cupressinum, *Soland. ex Forst. f. Pl. Escul.* 80.—N. Zel.
distichum, D. Don, in Lamb. Pin. ed. II. 190 = Podocarpus taxifolia.
elatum, *Wall. Cat.* n. 6045.—Malaya.
excelsum, D. Don, in Lamb. Pin. ed. II. App. = Podocarpus dacrydioides.
ferrugineum, Van Houtte, ex Gord. Pinet. 290 = Podocarpus dacrydioides.
Fitzgeraldi, *F. Muell. Progrs.* xi. 102.—Austral.
Franklinii, *Hook. f. in Hook. Lond. Journ. Bot.* iv. (1845) 152. t. 6.—Tasmania.
huonense, A. Cunn. ex Carr. Conif. ed. I. 490 = intermedium, *T. Kirk, in Trans. N. Z. Inst.* x. (1878) 386.—N. Zel.
Junghuhnii, Miq. Pl. Jungh. 4 = elatum.
Kirkii, *F. Muell. ex Parl. in DC. Prod.* xvi. 11. 495.—N. Zel.
laxifolium, *Hook. f. in Hook. Lond. Journ. Bot.* iv. (1845) 143.—N. Zel.
lycopodioides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xvi. (1869) 829.—N. Caled.
Mai, A. Cunn. in Ann. Nat. Hist. i. (1838) 213 = Podocarpus spicata.
Mayi, Van Houtte, ex Gord. Pinet. 287 = praec.
Pancheri, Brongn. & Gris, in Bull. Soc. Bot. Fr. xvi. (1869) 830 = Podocarpus pectinata.
plumosum, D. Don, in Lamb. Pin. ed. II. App. 143 = Libocedrus Doniana.
taxifolium, Banks & Soland. ex Lamb. Pin. ed. II. iii. 119. = Podocarpus spicata.
taxoides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 497.—N. Caled.
tenuifolium, Parl. in DC. Prod. xvi. 11. 521 = Podocarpus tenuifolia.
tetragonum, Parl. l. c. 496 = Microcachrys tetragona.
thuioides, Banks & Soland. ex Carr. Conif. ed. I. 479 = Podocarpus dacrydioides.
ustum, Vieill. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. xiii. (1866) 498 = Podocarpus usta.
Vieillardii, Parl. in DC. Prod. xvi. 11. 521 = Podocarpus Vieillardii.
westlandicum, *T. Kirk, in Hook. Ic. Pl.* t. 1218.—N. Zel.

DACRYODES, Vahl, in Dansk. Selsk. Skrift. vi. (1810) 115. *BURSERACEAE*, Benth. & Hook. f. i. 327.
excelsa, *Vahl, l. c.* 116.—Ins. Porto Rico.
hexandra, *Griseb. Fl. Brit. W. Ind.* 174.—Ins. S. Doming.

DACTILIS, Neck. Delic. Gallo-Belg. i. 58 (1768) = **Dactylis**, Linn. (Gramin.).

DACTILON, Vill. Hist. Pl. Dauph. ii. 69 (1787) = **Cynodon**, Rich. (1805) (Gramin.).
officinale, Vill. l. c. = C. Dactylon.
sanguinale, Vill. l. c. = Panicum sanguinale.

DACTIMALA, Rafin. Sylva Tellur. 158 (1838) = **Chrysophyllum**, Linn. (Sapotac.).
angustifolia, Rafin. l. c. = C. angustifolium.
microcarpa, Rafin. l. c. = C. glabrum.
olivaeformis, Rafin. l. c. = C. oliviforme.

DACTIPHYLLON, Rafin. in Am. Monthly Mag. (1818) 268 = **Trifolium**, Tourn. (Legumin.).

DACTIPHYLLUM, Rafin. in Journ. Phys. lxxxix. (1819) 261 = praec.

DACTYLAENA, Schrad. ex Schult. f. Syst. vii. p. ix (1829), in obs. *CAPPARIDRAE*, Benth. & Hook. f. i. 105.
micrantha, *Schrad. ex Schult. f. l. c.*—Bras.

DACTYLAENA:—
microphylla, *Bichl. in Mart. Fl. Bras.* xiii. i. 241.—Bras.
paniculora, *Griseb. in Goett. Abh.* xxiv. (1878) 17.—Am. austr.
Pohliana, *Bichl. in Mart. Fl. Bras.* xiii. i. 242.—Bras.

DACTYLANTHERA, Welw. Apont. 560 (1858), nomen. *GUTTIFERAE*.

DACTYLANTHES, Haw. Syn. Pl. Succ. 182 (1812) = **Euphorbia**, Linn.
anacantha, Haw. l. c. = E. anacantha.
globosa, Haw. in Phil. Mag. (1828) 382 = E. globosa.
hamata, Haw. Syn. Pl. Succ. 183 = E. hamata.
Medusa, Rafin. Fl. Tellur. iv. 112 = E. cervicornis?
petula, Haw. Syn. Pl. Succ. 182 = E. anacantha.
tuberculata, Haw. l. c. 183 = E. tuberculata.

DACTYLANTHUS, Hook. f. in Trans. Linn Soc. xxii. (1859) 425. t. 75. *BALANOPHOREAE*. Benth. & Hook. f. iii. 235.
Taylorii, *Hook. f. l. c.*—N. Zel.

DACTYLIANDRA, Hook. f. in Oliver, Fl. Trop Afr. ii. 557 (1871). *CUCURBITACEAE*.
Welwitschii, *Hook. f. l. c.*—Afr. trop.

DACTYLICAPNOS, Wall. Tent. Fl. Nep. 51. t. 39 (1826) = Dicentra, Bernh. (Papav.).
asplenifolia, Wall. ex Steud. Nom. ed. II. i. 478 = seq.
thalictrifolia, Wall. Tent. Fl. Nep. 51. t. 39 = Dicentra thalictrifolia.

DACTYLIOTA, Blume, Mus. Bot. Lugd. Bat. i. 21 (1849) = **Medinilla**, Gaudich. (Melast.).
bracteata, Blume, l. c. 24 = M. bracteata.
setigera, Blume, l. c. = M. setigera.

DACTYLIS, Linn. Gen. ed. II. 29 (1742). *GRAMINEAE*, Benth. & Hook f. iii. 1193.
Amaxitis, Adans. Fam. ii. 34 (1763).
abbreviata, Bernh. ex Link, Hort. Berol. i. 153 = glomerata.
adscendens, Schrad. ex Schult. Mant. ii. 851 = Lasiochloa adscendens.
alsatica, Bess. in Schult. l. c. 626 = glomerata.
bombycina, *Steud. Syn. Pl. Gram.* 428.—Arab.
brevifolia, Koen.-ex Willd. Sp. Pl. i. 410 = Aeluropus pubescens.
caespitosa, Forst. in Comm. Goett. ix. (1789) 22 = Poa flabellata.
canariensis, Nees, ex Steud. Nom. ed. II. i. 478 = Aeluropus Smithii.
capitata, Linn. f. Suppl. 110 = Lasiochloa laevis.
capitata, *Schult. Mant.* ii. 626.—Sardin.
capitellata, Link, Hort. Berol. i. 153 = glomerata.
caudata, Brot. Fl. Lusit. i. 100 = Koeleria phleoides
ciliaris, Linn. Mant. ii. 185 = Lasiochloa ciliaris.
ciliata, Opiz, ex Nym. Consp. 819 = glomerata.
cristata, Bieb. Fl. Taur. Cauc. i. 67 = Koeleria cristata.
cylindracea, Brot. Fl. Lusit. i. 99 = Koeleria phleoides.
cynosuroides, Linn. Sp. Pl. 71 = Spartina cynosuroides.
cynosuroides, Lodd. ii. 112 = Spartina stricta.
cynosuroides, *Roth, Nov. Pl. Sp.* 74.—Ind. or.
cynosuroides, Walt. Fl. Carol. 77 = Spartina juncea.
dietrichophylla, Brign. Fasc. Pl. Forojul. 16 = Aeluropus laevis.
fasciculata, Lam. Tabl. Encyc. i. 180 = Spartina fasciculata.
fasciculata, C. Sm. ex Link, in Buch, Beschr. Ins. Canar. 139 = Aeluropus Smithii.
geniculata, Burm. f. Fl. Ind. 28 = Spartina geniculata.
glauca, *Roth, Catalecta*, iii. 18.—Hab.?
glaucescens, Willd. Enum. Hort. Berol. 111 = glomerata.
glomerata, *Linn. Sp. Pl.* 71.—Europ.; As. bor.
heterophylla, Steud. Nom. ed. II. i. 478 = Aeluropus laevis.
hirta, Schrad. ex Schult. Mant. ii. 850 = Lasiochloa hirta.
hispanica, Nob. ex Steud. Syn. Pl. Gram. 297 = Nodăna.
hispanica, Roth, Catalecta, i. 8 = glomerata.
hispida, Thunb. Prod. Pl. Cap. 23 = Lasiochloa hispida.
Juncinella, Bory, ex Boiss. Voy. Bot. Espagne, ii. 665 = glomerata.
Kovarscicii, *Opiz, in Lotos*, iii. (1853) 97.—Hab.?
laevis, Thunb. Prod. Pl. Cap. 23 = Lasiochloa laevis.
lagopoides, Linn. Mant. i. 30; Suss. f. Fl. Ind. 28 = Aeluropus laevis.

DACTYLIS:—
Lima, Steud. Syn. Pl. Gram. 446, sphalm. vide *Cynosurus Lima*.
littoralis, Willd. Sp. Pl. i. 408 = Aeluropus laevis.
lobata, Heb. Fl. Taur. Cauc. i. 67 = Koeleria crustata.
longifolia, Schrad. ex Schult. Mant. ii. 551 = Lasiochloa longifolia.
maritima, Cart. Enum. Brit. Gr. 4 = Spartina stricta.
maritima, Schrad. Fl. Germ. i. 318. t. 6. f. i. = Aeluropus laevis.
maritima, Walt. Fl. Carol. 77 = Spartina laevigata.
massoviensis, Steud. Syn. Pl. Gram. 298 = Aeluropus laevis.
memphitica, Spreng. Nachr. i. Bot. Gart. Halle, 20 = Cutandia scleropoides.
mucronata, Steud. Syn. Pl. Gram. 298 = Aeluropus arabica.
Noëana, *Steud. l. c.* 297.—Ins. Oasero.
obtusifolia, Steud. l. c. 298 = Lasiochloa obtusifolia.
Ortmanniana, Opiz, Seznam, 86 = glomerata.
ovata, Steud. Syn. Pl. Gram. 298 = Lasiochloa ovata.
palaestina, *Steud. l. c.* 426.—Palest.
palaeco, Herb. Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 44 = Koeleria crustata.
paspaloides, Willd. Enum. Hort. Berol. 111 = Dinebra arabica.
patens Ait. Hort. Kew. ed. I. 104 = Spartina patens.
pauciceria, *Nees, Fl. Afr. Austr.* 429.—Afr. austr.
pendula, Dum. Obs. Gram. Belg. 146 = glomerata.
phleoides, *Klein, ex Steud. Nom.* ed. II. I. 478.—Ind. or.
pungens, Homem. Enum. Hort. Hafn. (1804) 9; ex Kunth, Enum. Pl. i. 1. 388 = Koeleria phleoides.
pungens, Schreb. Gras. ii. 42 = Echinaria capitata.
repens, Desf. Fl. Atlant. i. 79 = Aeluropus laevis.
repens, Sieber, ex Spreng. Syst. i. 348 = Aeluropus villacus.
rigida, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 60.—Ins. Creta.
scabra, Mann, in Opiz, Natural. vii. (1824) 58 = glomerata
serrata, Thunb. Prod. Pl. Cap. 22 = Lasiochloa serrata.
Smithii, Link, in Buch, Beschr. Canar. Ins. 139 = Aeluropus Smithii.
spicata, Brot. ex Spreng. Syst. i. 332 = Koeleria phleoides.
spicata, Willd. in Ges. Naturf. Fr. N. Schr. iii. (1801) 416 = Elytrophorus articulatus.
stricta, Ait. Hort. Kew. ed. I. i. 104 = Spartina stricta.
strictulosa, Steud. Syn. Pl. Gram. 299 = Lasiochloa strictulosa.
villosa, Tenore, Prod. Fl. Nap. p. lx = glomerata.
villosa, Thunb. Prod. Pl. Cap. 22 = Lasiochloa villosa.

DACTYLIUM, Griff. in Proc. Linn. Soc. ii. (1858) 252 = Erythropogon, Beauv. (Oacta.).
vagum, Griff. l. c. = E. vagum.

DACTYLOCTENIUM, Willd. Enum. Hort. Berol. 1029 (1809) = Eleusine, Gaertn. (Gramin.).
aegyptiacum, Willd. l. c. = E. aegyptiaca.
aristatum, Link, Hort. Berol. i. 89.—Aegypt.
australe, Steud. Syn. Pl. Gram. 299.—Afr. austr.
distachyum, Boj. Hort. Maurit. 270; Trin. ex Steud. Nom. ed. II. i. 479 = E. aegyptiaca.
falcatum, Willd. Enum. Hort. Berol. 1029 (= *Cynosurus falcatus*).—Ind. or.
Figarei, De Not. in Ann. Sc. Nat. Sér. III. ix. (1848) 325 = E. aegyptiaca.
glabrum, Beauv. Agrost. 72, nomen (= *Cynosurus glaber*).—Hab.?
glaucophyllum, Courb. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 188 = E. glaucophylla.
meridionale, Ham. Prod. Pl. Ind. Occ. 6 = E. aegyptiaca.
mucronatum, Willd. Enum. Hort. Berol. 1029 = E. aegyptiaca.
nitidum, Link, ex Steud. Nom. ed. II. i. 479 = Eragrostis nitida.
pilosum, Beauv. Agrost. 72, nomen.—Hab.?
prostratum, Willd. Enum. Hort. Berol. 1029 = E. aegyptiaca.
rubellum, Beauv. Agrost. 72 = E. aegyptiaca.
scindicum, Boiss. Diagn. Ser. II. iv. 121 = E. aristata.
semiglaberatum, Courb. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 126.—Afr. trop.

DACTYLOGRAMMA, Link, Hort. Berol. ii. 446 (1833) = Rhabdochaetopsis, Schrob. (Gramin.).
cimuoides, Link, l. c. = M. glomerata.

DACTYLON, Roem. & Schult. Syst. ii. 411, in syn. (1817) = DACTILON, Vill. = **Cynodon**, Rich. (1805) (Gramin.).

DACTYLOPETALUM, Benth. in Journ. Linn. Soc. iii. (1859) 79. *RHIZOPHOREAE*, Benth. & Hook. f. i. 682.
Barteri, *Hook. f. ex Oliver, Fl. Trop. Afr.* ii. 413.—Afr. trop.
gummiferum, Baill. Adansonia, iii. (1862) 22, 27 = sessiliflorum.
Mannii, *Hook. f. ex Oliver, Fl. Trop. Afr.* ii. 412.—Afr. trop.
sessiliflorum, *Benth. in Journ. Linn. Soc.* iii. (1859) 79.—Afr. trop.

DACTYLOPHYLLUM, Spach, Hist. Vég. Phan. ix. 108 (1840) = Galla, Ruiz & Pav. (Polemon.).

DACTYLOPHYLLUM, Spenn. Fl. Frib. iii. 1084 (1829) = **Potentilla**, Linn. (Rosac.).
Anserina, Spenn. l. c. = P. Anserina.
Ehrharti, Spenn. l. c. = Fragaria collina.
Fragaria, Spenn. l. c. = Fragaria vesca.
Fragariastrum, Spenn. l. c. = P. Fragariastrum.
Sibbaldia, Spenn. l. c. = P. Sibbaldi.

DACTYLORHIZA, Neck. Elem. iii. 129 (1790) = **Orchis**, Tourn.

DACTYLOSTALIX, Reichb. f in Bot. Zeit. xxxvi. (1878) 74 = **Cremastra**, Lindl. (Orchid.).
ringens, Reichb. f. l. c.—Japon.

DACTYLOSTEGIUM, Nees, in Mart. Fl. Bras. ix. 162, t. 81 (1847) = Diclipteru, Juss. (Acanth.).
portoricense, Nees, in DC. Prod. xi. 404, in obs. = Diclipteru portoricensis.
sparsiflorum, Nees, in Mart. Fl. Bras. ix. 164 = Diclipteru sparsiflora.

DACTYLOSTEMON, Klotzsch, in Wiegm. Archiv, vii. (1841) 181 = Actinostemon, Klotzsch (Euphorb.).
angustifolius, Klotzsch, l. c. = *Klotzschii*.
angustifolius, Muell. Arg. in Mart. Fl. Bras. xi. ii. 604 = A. caribaeus!
anisanthus, Griseb. in Goett. Abh. xxiv. (1879) 61.—Reg. Argent.
australis, Muell. Arg. in Mart. Fl. Bras. xi. ii. 608.—Bras.
brasiliensis, Muell. Arg. in Linnaea, xxxii. (1863) 114.—Bras.
communis, Muell. Arg. l. c. 112 = *Klotzschii*.
cuneatus, Klotzsch, ex Muell. Arg. l. c. 114 = A. cuneatus.
desertorum, Muell. Arg. in Mart. Fl. Bras. xi. ii. 608.—Bras.
estrellensis, Muell. Arg. l. c. 609.—Bras.
Gardneri, Muell. Arg. l. c. 602.—Bras.
glabratus, Klotzsch, in Wiegm. Archiv, vii. (1841) 181, nomen = *Klotzschii*.
grandifolius, Klotzsch, l. c. nomen; Muell. Arg. in Linnaea, xxxii. (1863) 111 (= *A. Klotzschianus*).—Guiana.
guianensis, Klotzsch, in Schomb. Faun. et Fl. Gui. 1184, nomen.—Guiana.
Hagenschii, Klotzsch, in Wiegm. Archiv, vii. (1841) 181, nomen = *Klotzschii*.
Klotzschii, Didr. in Kjoeb. Videnak. Meddel. (1857) 127.—Bras.
lagoensis, Muell. Arg. in Mart. Fl. Bras. xi. ii. 602.—Bras.
lasiocarpoides, Muell. Arg. in Linnaea, xxxii. (1863) 114 = A. lasiocarpoides.
lasiocarpus, Klotzsch, in Wiegm. Archiv, vii. (1841) 181, nomen; Muell. Arg. l. c. 111.—Bras.
leptopus, Muell. Arg. in Mart. Fl. Bras. xi. ii. 607.—Bras.
Lundianus, Didr. in Kjoeb. Videnak. Meddel. (1857) 126 = A. lasiocarpoides.
sanctaecrucis, Muell. Arg. in Mart. Fl. Bras. xi. ii. 601.—Bras.
obtusatus, Klotzsch, in Wiegm. Archiv, vii. (1841) 161, nomen = *Klotzschii*.
oligandrus, Muell. Arg. in Linnaea, xxxii. (1863) 115 = A. oligandrus.
Schomburghii, Klotzsch, in Wiegm. Archiv, vii. (1841) 181, nomen; et in Hook. Lond. Journ. Bot. ii. (1843) 45.—Guiana.

DACTYLOSTEMON:—
sparsiflorus, Muell. Arg. in Mart. Fl. Bras. xi. ii. 610.—Bras.
verticillatus, Klotzsch, Ind. Sem. Hort. Berol. App. (1851) 18; et in Linnaea, xxv. (1852) 298 = A. verticillatus.

DACTYLOSTYLES, Scheidw. in Otto & Dietr. Allg. Gartenz. vii. (1839) 405 = **Zygostates**, Lindl. (Orchid.).
fimbrata, Scheidw. l. c. 406 = Z. lunata.

DACTYLUS, Aschers. Fl. Brandenb 810 (1864) = **Cynodon**, Rich. (Gramin.).
officinalis, Aschers. l. c. = C. Dactylon.

DACTYLUS, Burm. f. Fl. Cap. Prod. 3 (1768) = **Pollinia**, Trin. (Gramin.).
spicatus, Burm. f. l. c. = P. villosa.

DACTYLUS, Forsk. Fl. Aegypt. Arab. p. xxxvi (1775) = **Diospyros**, Linn. (Ebenac.).
tropesuntinus, Forsk. l. c. = Diosp. Lotus.

DACTYPHYLLUM, Endl. Gen. 1268, in syn. (1840) = DACTIPHYLLUM, Raf. = **Trifolium**, Tourn. (Legumin.).

DADIA, Vell. Fl. Flum. 347 (1825). *COMPOSITAE*?
Benth. & Hook. f. ii. 228.
lixa, *Vell. l. c.* 348; viii. t. 86.—Bras.

DAEDALACANTHUS, T. Anders. in Thw. Enum. Pl. Zeyl. 229 (1864). *ACANTHACEAE*, Benth. & Hook. f. ii. 1082.
Griffithii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 486.—Reg. Himal.
macrophyllus, *T. Anders. l. c.* 487.—Burma.
microstachyus, *T. Anders l. c.* 486.—Burma.
montanus, *T. Anders. in Thw. Enum. Pl. Zeyl* 229.—Ind. or.
nervosus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 487.—Ind. or
Parishii, *T. Anders l. c.* 489.—Burma.
purpurascens, *T. Anders. l. c.* 488.—Ind. or.
roseus, *T. Anders. l. c.* 487.—Ind. or.
salaccensis, T. Anders. in Journ Agr. Hort. Soc. Ind. N. S. i. (1868) 274 = tetragonus.
scaber, T. Anders. in Journ. Linn. Soc. ix. (1867) 487 = nervosus.
splendens, *T. Anders. l. c.* 486.—Reg. Himal.
strictus, *T. Anders. l. c.* 487.—Reg. Himal.
suffruticosus, *T. Anders. l. c.* 488.—Ins. Andaman.
tetragonus, *T. Anders. l. c.*—Burma.
tubiflorus, *T. Anders. l. c.* 485.—Reg. Himal.

DAEMIA, R. Br. in Mem. Wern. Soc. i. (1809) 50 (Doemia). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 764.
aethiopica, *Decne. in DC. Prod.* viii. 544.—Afr. trop.
angolensis, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 337 = scandens.
atropurpurea, F. Muell. Census, 94 = Pentatropis atropurpurea.
barbata, Klotzsch, in Peters, Reise Mossamb. Bot. 274 = extensa.
bicolor, Sweet, Hort. Brit. ed. II. 361 = extensa?
cordata, *R. Br. in Mem. Wern. Soc.* i. (1809) 50.—As. occ.; Afr. trop.
extensa, *R. Br. l. c.*—Ind. or.; Malaya; Afr. trop.
Forskhâli, Schnit. Syst. vi. 113 = extensa.
garipensis, *E. Mey. Comm. Pl. Afr. Austr.* 220.—Afr. austr.
glabra, *Schult. Syst.* vi. 113.—Arab.
guineensis, G. Don, Gen. Syst. iv. 156 = scandens.
incana, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 336 = cordata.
Kempeana, *F. Muell. Census*, 94.—Austral.
linearis, F. Muell. Census, 94 = Pentatropis linearis.
quinquepartita, F. Muell. l. c. = Pentatropis quinquepartita.
scandens, *G. Don, ex Loud. Hort. Brit.* 94.—Afr. trop.
Schmittiana, *Pomel, Nouv. Mat. Fl. Atl.* 82.—Algeria.
tomentosa, Pomel, l. c. = cordata.

DAEMONOROPS, Blume, ex Schult. f. Syst. vii. 1338 (1830). *PALMAE* [Benth. & Hook. f. iii. 932]. *acerdene*, Blume, Rumphia, ii. p. viii; iii. 13 = D. Draco.
adspersus, *Blume, l. c.* tt. 142, 143 (Calamus in textu). —Java.
angustifolius, *Mart. Hist. Nat. Palm.* iii. 327.—Malaya.
barbatus, *Mart. l. c.* 330.—N. Guinea.
calapparius, *Blume, Rumphia*, iii. 7.—Ins. Amboina.
callicarpus, *Mart. Hist. Nat. Palm.* iii. 326.—Malaya.
cochleatus, *Tejsm. & Binn. Cat. Hort. Bog.* 861; *et ex Miq. in Verh. Akad. Wetensch.* xi. (1868) n. v. 29.—Sumatra †
crinitus, *Blume, Rumphia*, ii. p. viii; iii. 27.—Malaya.
Draco, *Blume, l. c.* iii. 8.—Malaya.
elongatus, *Blume, l. c.* 16.—Borneo.
fasciculatus, Mart. Hist. Nat. Palm. iii. 330 = Calamus Fernandezii.
fusus, *Blume, Rumphia*, iii. 17.—Borneo.
fuscus, *Mart. Hist. Nat. Palm.* iii. 331.—Ins. Philipp. Gaudichaudii, *Mart. l. c.*—Ins. Philipp.
geniculatus, *Mart. l. c.* 329.—Malaya.
grandis, *Mart. l. c.* 327.—Malaya.
Guruba, Mart. l. c. 330 = Calamus Guruba.
heteracanthus, Blume, Rumphia, iii. t. 139 = Calamus heteracanthus.
hirsutus, Blume, l. c. ii. p. viii; iii. 21 = Hystrix.
hygrophilus, *Mart. Hist. Nat. Palm.* iii. 328.—Malaya.
hypoleucus, Kurz, in Journ. As. Soc. Beng. xliii. ii. (1874) 208 = Calamus hypoleucus.
Hystrix, *Mart Hist. Nat. Palm.* iii. 328.—Malaya.
intermedius, *Mart. l. c.* 327.—Malaya.
Jenkinsianus, *Mart. l. c.*—Assam; Burma.
Kirtong, Griff. Palms Brit. E. Ind. 102. t. 216 B = malaccensis.
Korthalsii, *Blume, Rumphia*, iii. 23.—Borneo.
latispinus, Teijsm. & Binn. Cat. Hort. Bog. 74 = Calamus latispinus.
leptopus, *Mart. Hist. Nat. Palm.* iii. 329.—Malaya.
Lewisianus, *Mart. l. c.* 327.—Penang.
longipes, *Mart. l. c.* 329.—Malaya.
longipes, *Miq. Fl. Ind. Bat. Suppl.* 592.—Sumatra.
malaccensis, *Mart. Hist. Nat. Palm.* iii. 327.—Malaya.
manillensis, Mart. l. c. 330 = Calamus manillensis.
marginatus, Blume, Rumphia, iii. 24 = Calamus marginatus.
melanochaetes, *Blume, in Schult. f. Syst.* vii. 1338.—Malaya.
melanolepis, Mart. Hist. Nat. Palm. iii. 331 = Calamus Hugelianus.
mirabilis, *Mart. l. c.* 329.—Malaya.
monticolus, *Mart. l. c.* 328.—Malaya.
niger, *Blume, Rumphia*, iii. 5.—Ins. Amboina.
nutantiflorus, Mart. Hist. Nat. Palm. iii. 326 = Jenkinsianus.
oblongus, Blume, Rumphia, iii. 25. t. 141; Mart. Hist. Nat. Palm. iii. 305 = Calamus oblongus.
ochreatus, Teijsm. & Binn. Cat. Hort. Bog. 74 = Calamus ochreatus.
palembanicus, *Blume, Rumphia*, iii. 20.—Sumatra.
periacanthus, *Miq. Fl. Ind. Bat. Suppl.* 593.—Sumatra.
petiolaris, *Mart. Hist. Nat. Palm.* iii. 326.—Malaya.
platyacanthus, Mart. l. c. 324, 328 = oblongus
Platypetalus, Mart. l. c. 329 = Calamus platypetalus.
ramosissimus, Mart. l. c. 330 = Calamus ramosissimus.
Rheedii, Mart. l. c. 330 = Calamus Rheedei.
ruber, Mart. l. c. 329 = D. Draco.
Rumphii, *Mart. l. c.* 331.—Ins. Amboina.
strictus, Blume, Rumphia, iii. 19 = longipes.
trichroos, *Miq. Fl. Ind. Bat. Suppl.* 592.—Sumatra.
verticillaris, *Mart. Hist. Nat. Palm.* iii. 329.—Malaya.

DAHLIA, Cav. lc. i. 56. t. 30 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 386.
GEORGIA, Spreng. Anleit. ii. ii. 567 (1818).
GEORGINA, Willd. Sp. Pl. iii. 2174 (1803).
arborea, *Regel, Cat. Pl. Hort. Absahor.* 46, nomen.—Am. trop.
acutiflora, Mog. & Sesse, ex DC. Prod. v. 494 = coccinea.
adenocaulon, *G. Don*, in *Sweet, Hort. Brit.* ed. III. 353, nomen.—Mexic.
Barkeriae, Knowles & Westc. Flor. Cab. ii. (1838) 28; iii. (1840) 147. t. 127 = variabilis.
DAHLIA:—
bidentifolia, Sallsb. Parad. Lond. t. 19 = coccinea.
Cervantesii, Lag. ex DC. Prod. v. 494 = coccinea.
coccinea, *Cav. lc.* iii. 2. t. 266.—Mexic.
crocata, Lag. Gen. et Sp. Nov. 30 = variabilis.
crocea, Poir. Encyc. ii. 445 = coccinea.
Decaisneana, Verlot, in Rev. Hortic. (1864) 31 = Merckii.
excelsa, *Benth. in Maund, Botanist*, ii. t. 88.—Mexic.
frustranea, Ait. Hort. Kew. ed. II. v. 88 = coccinea.
glabrata, Lindl. Bot. Reg. (1840) t. 29 = Merkii.
gracilis, *Ortg. in Regel, Gartenfl.* (1876) 97. t. 361.—Mexic.
imperialis, *Roezl, l. c.* (1868) 243. t. 407.—Mexic.
Maximiliana, *Hort.—Cf. Gard. Chron.* (1879) ii. 216, 525; *et ex Hemsl. Biol. Centr. Am. Bot.* ii. 197. —Mexic.
Merckii, *Lehm. Delect. Sem. Hort. Hamb.* (1839), *ex Linnaea*, xiv. (1840) *Litt.* 130.—Mexic.
minor, Vis. in Nuovi Sagg. Acc. Padov. v. (1840) 268 = Merkii.
nana, Andr. Rep. Bot. vis. t. 483 = variabilis.
pannata, Cav. lc. i. 57. t. 30 = variabilis.
platylepis, *A. Brongn. ex Neumann, in Rev Hort. Sér.* II. iv. (1845–46) 305.—Mexic.
pubescens, *A. Brongn. l. c.*—Mexic.
purpurea, Poir. Encyc. Suppl. ii. 444 = variabilis.
pusilla, Zucc. ex DC. Prod. v. 494 = variabilis.
rosea, Cav. lc. iii. 33. t. 265 = variabilis.
Rejoiceana, Knowles & Westc. Flor. Cab. ii. (1840) 186 = variabilis.
sambucifolia, Sallsb. Parad. Lond. t. 16 = variabilis.
scapigera, *Knowles & Westc. Flor. Cab.* iii. (1840) 118. t. 118.—Mexic.
sphondylifolia, Sallsb. in Trans. Hort. Soc. i. (1812) 91 = variabilis.
superflua, Ait. Hort. Kew. ed. II. v. 87 = variabilia.
variabilis, *Desf. Cat. Hort. Par.* ed. III. 182.—Mexic.
Yuarezii, *Hort. ex Gard. Chron.* (1879) ii. 483.—

DAHLIA, Thunb. in Skrivt. Naturh. Selsk. Kiobenh. ii. (1799) 138. t. 4 = Trichoboclus, Pers. (Hamemel.).
crinita, Thunb. l. c. = T. crinitus.

DAHURONIA, Scop. Introd. 219 (1777) = **Moquilea**, Aubl. (Rosac.).

DAIS, Royen, ex Linn. Sp. Pl. ed. II. 556 (1762). *THYMELAEACEAE*, Benth. & Hook. f. iii. 192.
anthyllodes, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 530, 596 = Lasiosiphon anthylloides.
argentea, Eckl. & Zeyh. l. c. 594 = Lasiosiphon Meisnerianus.
canescens, *Bartl. ex Meisn. in Linnaea*, xiv. (1840) 388.—Afr. austr.
coccinea, *Lundii.* in Freyc. Voy. Bot. 443. t. 44 = Pseudais coccinea.
cotinifolia, *Linn. Sp. Pl.* ed. II. 556.—Afr. austr.
cunonia, Linn. ex Decne. in Jacquem. Voy. Bot. 149 = Lasiosiphon cunestus.
disperma, Forst. f. Prod. 33 (= *Drimyospermum Forsteri*). —Ins. Tongataba.
dubiosa, Bleine, Cat. Gew. Bultenz.; Bijdr. 651 = Phaleria Blumei.
dubiosa, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 369 = Phaleria laurifolia.
eriocephala, *Lichtenst. ex Meisn. in Linnaea*, xiv. (1840) 389.—Afr. austr.
glaucescens, *Decne. in Ann. Sc. Nat.* Sér. II. xx. (1843) 51.—Madag.
guidioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 244.—Madag.
laurifolia, Blanco, Fl. Filip. ed. I. 375 = Phaleria Perrottetiana.
laurifolia, Jacq. Coll. i. 146 = cotinifolia.
linifolia, Lam. Encyc. ii. 255 = Lasiosiphon linifolius.
longifolia, *Zipp. ex Span. in Linnaea*, xv. (1841) 335, nomen.—Hab. ?
madagascariensis, Boj. ex Meisn. in DC. Prod. xiv. 529 = glaucescens.
madagascariensis, Lam. Encyc. ii. 254 = Lasiosiphon madagascariensis.
octandra, Linn. Mant. i. 69; Burm. f. Fl. Ind. 104. t. 22. 2 = (*Drimyospermum Burmannii*).—Ind. or.
Ovensii, Eckl. & Zeyh. ex Meisn in DC. Prod. xiv. 595 = Lasiosiphon triphlaevis.

DAIS:—
Passerina, Wendl. ex Meisn. in Linnaea, xiv. (1840) 389 = eriocephala.
pubescens, Lam. Encyc. ii. 255 = Lasiosiphon pubescens.
purpurea, *Gaudich. ex Steud. Nom.* ed. II. i. 479, nomen.—Hab. ?
sericea, Lam. Encyc. ii. 767 = Lasiosiphon anthylloides.
spicata, Endl. Sert. Cabul. 5. t. 3 = Stellera Lessertii.
virgata, Lichtenst. ex Meisn. in DC. Prod. xiv. 530, 561 = Arthrosolon polycephalus.

DAISWA, Rafin. Fl. Telfur. iv. 18 (1836) = **Paris**, Linn. (Liliac.).
polyphylla, Rafin. l. c. = P. polyphylla.

DALBERGARIA, Tussac, Fl. Ant. i. 141. t. 19 (1808) = **Columnea**, Linn. (Gesner.).
phoenicea, Tussac, l. c. = C. sanguinea.
sanguinea, Steud. Nom. ed. II. i. 479 = praec.

DALBERGIA, Linn. f. Suppl. 52 (1781). *LEGU. MINOSAE*, Benth. & Hook. f. i. 544.
AMERIMNON, P. Br. Hist. Jamaic. 288 (1756); DC. Prod. ii. 421 (1825).
ENDESPERMUM, Blume, Cat. Gew. Buitenz. 23 (1823).
LHOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 94.
MISCOLOBIUM, Vog. in Linnaea, xi. (1837) 200.
PODIOPETALUM, Hochst. in Flora, xxiv. (1841) 657.
SALKEN, Adans. Fam. ii. 322 (1763).
SEMBIONOTIS, Schott, in Wien. Zeitschr. iii. (1829) 804.
SOLORI, Adans. Fam. ii. 327 (1763).
TRIPTOLEMEA, Mart. in Flora, xx. (1837) Beibl. 122.
acacicefolia, Dalz. in Hook. New Journ. ii. (1850) 37 = tamaridifolia.
acuminata, *Hassk. in Flora*, xxv. (1842) ii. *Beibl.* 58. —Java.
acuta, *Benth. in Journ. Linn. Soc. lv. Suppl.* (1860) 36; *et in Mart. Fl. Bras.* xv. t. 223.—Bras.
Afzeliana, *G. Don, Gen. Syst.* ii. 276.—Afr. trop.
Afzelii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 284.—Afr. trop.
alata, *Roxb. Hort. Beng.* 53; *Fl. Ind.* iii. 225.—Ind. or.
ambongoensis, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 486.—Madag.
Amerimnum, *Benth. in Journ. Linn. Soc.* lv. *Suppl.* (1860) 36.—Nicaragua.
amorphoides, Spreng. Syst. iv. Cur. Post. 268 = Eysenhardtia amorphoides.
angustifolia, Hassk. in Flora, xxv. (1842) ii. Beibl. 58 = Millettia sericea.
arborea, Heyne, ex Roth, Nov. Pl. Sp. 380 = lanceolaria *arborea*, Willd. Sp. Pl. iii. 901 = Pongamia glabra.
arbutifolia, *Baker, in Oliver, Fl. Trop. Afr.* ii. 282.—Afr. trop.
argentea, Zoll. in Nat. en Geneesk. Arch. Nederl. Indie, iii. (1846) 65 = Derris thyrsiflora.
armata, *E. Mey. Comm. Pl. Afr. Austr.* 152.—Afr. austr.
asamica, *Benth. in Miq. Pl. Yungh.* 256, in adnot.—Assam.
Bakeri, *Weiw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 285.—Afr. trop.
Barclayi, Herb. Kew. ex Baker, in Journ. Linn. Soc xx. (1883) 229 = Mundulea reroluta.
Barclayii, *Telfair, ex Hook. Exot. Fl.* ii. t. 188.—Madag.
Barosi, *Baker, in Journ. Linn. Soc.* xxi. (1884) 337.—Madag.
Bernieri, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 487. —Madag.
Blumei, Hassk. Fl. Jav. Rar. 400 = tamarindifolia.
Bolvini, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 489. —Madag.
brecteolata, *Baker, in Oliver, Fl. Trop. Afr.* ii. 284.—Afr. trop.
brevilinealis, *Vatke, in Oestr. Bot. Zeitschr.* xxvIII. (1878) 266.—Afr. trop.
calycina, *Benth. in Journ. Linn. Soc* lv. *Suppl.* (1860) 36.—Guatemala.
campechiana, *Benth. l. c.* 37.—Mexic.
cana, *R. Grah. in Wall. Cat.* n. 5899.—Burma.
cassioides, Wall. l. c. 5863 = stipulacea.
casdana, *G. Don, Gen. Syst.* ii. 275.—Guiana.

DALBERGIA :—
Championii, *Thw. Enum. Pl. Zeyl.* 94.—Ceylan. ; Malaya.
Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 436.—Madag.
confertiflora, *Benth. in Mig. Pl. Yungh.* 255, in adnot. —Ind. or.
confusa, Blume, ex Miq. Fl. Ind. Bat. i. 1. 129 = Championii.
congesta, *R. Grah. in Wall. Cat.* n. 5872.—Ind. or.
corymbifera, Blume, ex Miq. Fl. Ind. Bat. i. 1. 130 = Cumingiana.
Crowei, DC. Prod. ii. 417 = Derris robusta.
cujabensis, *Benth. in Journ. Linn. Soc. iv. Suppl.* (1860) 36 ; *et in Mart. Fl. Bras.* xv. i. 224.—Bras.
cultrata, *R. Grah. in Wall. Cat.* n. 5861.—Burma.
Cumingiana, *Benth. in Mig. Pl. Yungh.* 255, in adnot. —Ins. Philipp. ; Burma.
Cumingii, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 33 = prasc.
densa, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 217.—Ins. Pacific.
densiconna, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 435.—Madag.
densiflora, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 37 ; *et in Mart. Fl. Bras.* xv. i. 225.—Bras.
Diphaca, Pers. Syn. ii. 276 = Ormocarpum semiodes.
discolor, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 130.— Malaya.
domingensis, Turp. ex Pers. Syn. ii. 276 = Lonchocarpus sericeus.
elliptica, Span. in Linnæa, xv. (1841) 197 = purpurea.
emarginata, Roxb. Hort. Beng. 53 ; Fl. Ind. iii. 224 = latifolia.
emirnensis, *Benth. in Journ. Linn. Soc. iv. Suppl.* (1860) 33.—Madag.
ferruginea, Hochst. ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 42 = sympathetica.
ferruginea, Roxb. Hort. Beng. 98 ; Fl. Ind. iii. 228 = stipulacea.
flexuosa, R. Grah. in Wall. Cat. n. 5875 = reniformis.
foliacea, *Wall. Cat.* n. 5856. A, C, D.—Burma.
foliolosa, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 37 ; *et in Mart. Fl. Bras.* xv. i. 225.—Bras.
frondosa, Roxb. Hort. Beng. 53 ; Fl. Ind. iii. 226 = lanceolaria.
frondosa, Wall. Cat. n. 5855 B = sympathetica.
Gardneriana, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 43 = congesta.
glandulosa, *Benth. l. c.* 35 ; *et in Mart. Fl. Bras.* xv. i. 223.—Bras.
glauca, Wall. Cat. n. 5862 = ovata.
glaucescens, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 37 ; *et in Mart. Fl. Bras.* xv. i. 224.—Bras.
glaucescens, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 149 = Derris elliptica.
glomerata, *Hemsl. Diagn. Pl. Nov.* 8.—Mexic.
glomeriflora, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 70.—Burma.
gracilis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 38 ; *et in Mart. Fl. Bras.* xv. i. 226.—Bras.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 437.—Madag.
Greveana, *Baill. l. c.* 436.—Madag.
guianensis, Spreng. Syst. iv. Cur. Post. 268 = Lonchocarpus sericeus.
Hancei, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 44.—China.
heptaphylla, Poir. Encyc. Suppl. ii. 446 = Lonchocarpus latifolius.
heterophylla, Willd. Sp. Pl. iii. 901 = Derris uliginosa. Hildebrandtii, *Vatke, in Linnæa*, xliii. (1880–82) 101. —Ins. Nossibé.
hircina, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 46.—Reg. Himal.
hircina, Buch.-Ham. in Wall. Cat. n. 5871 A = lanceolaria.
Hookeri, *Boj. Hort. Maurit.* 111, nomen.—Madag.
horrida, R. Grah. in Wall. Cat. n. 5877 = spinosa.
hostilis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 33.—Afr. trop.
hupeana, *Hance, in Journ. Bot.* xx. (1882) 5.—China.
Imaculata, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 42 ; *et in Mart. Fl. Bras.* xv. i. 227.—Bras.
japonica, *G. Don, Gen. Syst.* ii. 276.—Japon.
javanica, Miq. Fl. Ind. Bat. i. 1. 132 = latifolia.
Jenghahnii, *Benth. in Mig. Pl. Yungh.* 254.—Malaya.
Krowee, Roxb. Fl. Ind. iii. 230 = Derris robusta.

DALBERGIA :—
lactea, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 251. —Afr. trop.
lanceolaria, *Linn. f. Suppl.* 316.—Ind. or.
lanceolaria, Moon, Cat. Pl. Ceyl. 51 = Pericopsis Mooniana.
lanceolata, Zipp. ex Span. in Linnæa, xv. (1841) 197 = lanceolaria.
lateriflora, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 37 ; *et in Mart. Fl. Bras.* xv. i. 225.—Bras.
latifolia, *Roxb. Fl. Corom.* ii. 7. t. 113.—Ind. or.
latialiqua, *Desf. Tabl.* ed. I. 198 ; ed. II. 226.—Am. austr.
lineata, R. Grah. in Wall. Cat. n. 5866 A = stipulacea, tamaridifolia.
luzoniensis, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 33.—China.
macrophylla, Spreng. Syst. iv. Cur. Post. 268 = Lonchocarpus sericeus.
macrosperma, *Welw. ex Baker, in Oliver, Fl. Trop Afr.* ii. 236.—Afr. trop.
madagascariensis, *Vatke, in Linnæa*, xliii. (1880–81) 105.—Madag.
malifolia, *Welw. ex Baker, in Oliver, Fl. Trop Afr* ii. 233.—Afr. trop.
marginata, Roxb. Hort. Beng. 53 ; Fl. Ind. iii. 230 = Derris marginata.
melanoxylon, *Guill. & Perr. Fl. Seneg. Tent.* i. 227 t. 53.—Afr. trop.
Meyeriana, Steud. Nom. ed. II. i. 480 = multijuga.
Millettii, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 34.—China.
Miscolobium, *Benth. l. c.* 35 ; *et in Mart. Fl. Bras.* xv. i. 222.—Bras.
Monsalia, Buch.-Ham. ex Wall. Cat. sub n. 5849 = Derris robusta.
Monetaria, Linn. f. Suppl. 317 = Ecastaphyllum Monetaria.
monosperma, *Dals. in Hook. Kew Journ.* ii. (1850) 36.—Ind. or.
Mooniana, Thw. Enum. Pl. Zeyl. 93 = Pericopsis Mooniana.
multiflora, Heyne, ex Wall. Cat. sub n. 5848 = paniculata.
multijuga, R. Grah. in Wall. l. c. n. 5865 = tamaridifolia.
multijuga, *E. Mey. Comm. Pl. Afr. Austr.* 153.—Afr. austr.
myriantha, *Benth. ex A. Gray, Bot. U. St. Expl. Exped.* i. 458.—Bras.
myriantha, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 100 = stipulacea.
nigra, *Allem. ex Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 36 ; *et in Mart. Fl. Bras.* xv. i. 224.—Bras.
nigrescens, *Kurz, in Pegu Rep. App. A.* 48. *et B.* 45. —Burma.
nitidula, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 235.—Afr. trop.
oblongifolia, *G. Don, Gen. Syst.* ii. 276.—Afr. trop.
obovata, *E. Mey. Comm. Pl. Afr. Austr.* 152.—Afr. austr.
ougeinensis, Roxb. Hort. Beng. 53 ; Fl. Ind. iii. 220 (oojeinensis) = Ougeinia dalbergioides.
ovata, *R. Grah. in Wall. Cat.* n. 5854.—Burma.
paniculata, *Roxb. Pl. Corom.* ii. 8. t. 114.—Ind. or.
paniculata, Wall. Cat. n. 5848 J = monosperma.
paniculata, Wall. l. c. E & I = confertiflora.
parviflora, Roxb. Hort. Beng. 98 ; Fl. Ind. iii. 225 = Jangthuhnii.
pendula, Tenore, in Rendic. Soc. Borb. Nap. i. (1842) 409 = D. Sissoo.
pendulifiora, Blume, ex Miq. Fl. Ind. Bat. i. 1. 133 = stipulacea.
pentaphylla, Poir. Encyc. Suppl. ii. 445 = Lonchocarpus latifolius.
Pervillei, *Vatke, in Linnæa*, xliii. (1880–82) 106.— Madag.
phoenicea, Prits. Ic. Ind. 846, sphalm. vide *Dalbergaria phoenicea*.
polyphylla, *Benth. in Miq. Pl. Yungh.* i. 256, in adnot. —Ins. Philipp.
polyphylla, Benth. l. c. partim = Millettii.
polyphylla, Poir. Encyc. Suppl. ii. 446 = Sesbania vesicaras.
prasoca, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 233 = Pterocarpus lucens.
Pseudo-Sissoo, Miq. Fl. Ind. Bat. i. 1. 128 = Championii.

DALBERGIA :—
pubescens, *Hook. f. ex Hook. Niger Fl.* 315.—Afr. trop.
pubinervis, Span. in Linnæa, xv. (1841) 197 = Derris pubinervis.
pulchra, Voigt, Hort. Suburb. Calc. 242 = Millettia pulchra.
punctata, Spreng. Syst. iv. Cur. Post. 268 = Lonchocarpus violaceus, *H. B. & Kr.*
purpuracescens, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 436.—Madag.
purpurea, *Renw. ex Blume, Cat. Gew. Buitens.* 90, nomen : *Hassk. in Flora*, xxv. (1842) 11. *Beibl.* 58 —Java.
purpurea, *Wall. Cat.* n. 5869.—Burma.
radicosa, Zipp. ex Miq. Fl. Ind. Bat. i. 1 141 = Derris uliginosa.
reniformis, *Roxb. Hort. Beng.* 53 ; *Fl. Ind.* iii. 226.— Ind. or. ; Burma.
repens, Span. Ic 1 ; et in Linnæa, xv. (1841) 197 = Derris uliginosa.
retusa, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 436.— Madag.
retusa, *Hemsl. Diagn. Pl. Nov.* 8.—Panama.
Richardi, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 437. —Madag.
rimosa, *Roxb. Hort. Beng.* 58 ; *Fl. Ind.* iii. 233.—Reg. Himal.
riparia, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 31 ; *et in Mart. Fl. Bras.* xv. i. 221.—Bras.
robusta, Miq ex Benth. l. c. 108 = Derris scandens.
robusta, Roxb. Hort. Beng. 58 = Derris robusta.
robusta, Wall. Cat. n. 5849 A = lanceolaria.
rostrata, R. Grah. in Wall. Cat. n. 5867 = stipulacea.
rostrata, *Hassk. in Flora*, xxv. (1842) 11. *Beibl.* 58.— Java. Buch.-Ham.
rotundifolia, *Sond. in Linnæa*, xxiii. (1850) 35.—Afr. austr.
rubiginosa, *Roxb. Pl. Corom.* ii. 9. t. 115.—Ind. or. ; China.
rufa, *G. Don, Gen. Syst.* ii. 276.—Afr. trop.
rufa, R. Grah. in Wall. Cat. n. 5864 = tamarindifolia.
saxatilis, *Hook. f. in Hook Niger Fl.* 314.—Afr. trop
scandens, Roxb. Pl. Corom. ii. 49. t. 192 = Derris scandens.
Schimperiana, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 232 = Lonchocarpus lasiflorus.
sennoides, Blume, ex Miq. Fl. Ind. Bat. i. 1. 134 = lanceolaria.
sericea, *Boj. Hort. Maurit.* 111, nomen.—Madag. ?
sericea, *G. Don, Gen Syst.* ii. 276.—Reg. Himal.
sericea, Spreng. Syst. iii. 198 = Millettia sericea.
sessoides, R. Grah. in Wall Cat. n. 5878 = latifolia.
Sissoo, *Roxb. Hort. Beng.* 58 ; *Fl. Ind.* iii. 228.—Ind. or. ; Afghan.
spinosa, Roxb. ll. cc. [98] ; 238.—Ind. or. ; Afghan.
Spruceana, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 35 ; *et in Mart. Fl. Bras.* xv. i. 223.—Bras.
stenocarpa, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 205.—Reg. Himal.
stipulacea, *Roxb. Hort. Beng.* 58 ; *Fl. Ind.* iii. 233.— Reg. Himal. ; Burma.
stipulata, Wall. Cat. n. 5868 A, B = velutina, reniformis.
Stockii, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 42.—Ind. or.
striata, Boj. Hort. Maurit. 111 = Mundulea suberosa.
suaresensis, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 437.—Madag.
sympathetica, *Nimmo, in J. Grah. Cat. Pl. Bomb.* 55. —Ind. or.
tamarindifolia, *Roxb. Hort. Beng.* 58 ; *Fl. Ind.* iii. 233. —Ind. or. ; Malaya.
Telfairii, Boj. Hort. Maurit. 111 = Mundulea Telfairii.
tephrosioides, Wight & Arn. Prod. 210 = Millettia pulchra.
Thomsoni, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 33.—Reg. Himal.
timoriensis, DC. Prod. ii. 417 = Derris scandens.
tingens, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 438.— Madag.
tingens, Buch.-Ham. in Wall. Cat. n. 5860 = stipulacea.
tomentosa, Vog. in Linnæa, xi. (1837) 197 = variabilis.
torta, R. Grah. in Wall. Cat. n. 5878 = monosperma.
toxicaria, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 438.—Madag.
vacciniifolia, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 268.—Afr. trop.
variabilis, *Vog. in Linnæa*, xi. (1837) 196.—Am. trop.

DALBERGIA:—
velutina, *Benth. in Miq. Pl. Jungh.* 255, in adnot.—Reg. Himal.; Burma.
wessata, *Zipp.* ex Span. in Linnæa, xv. (1841) 197 = Derris scandens.
villosa, *Benth. in Journ. Linn. Soc. lv. Suppl.* (1860) 88; *et in Mart. Fl. Bras.* xv. t. 236.—Bras.
violacea, Hoffmgg. Vers. Pfl. Nachtr. i. 181 = Lonchocarpus violaceus, *H. B. & K.*
volubilis, *Roxb. Fl. Coromn.* ii. 48. t. 191.—Ind. or. *zeylanica*, Roxb. Hort. Beng. 58; Fl. Ind. iii. 228 = lanceolaria.
Zollingeriana, Miq. Fl. Ind. Bat. i. t. 130 = Cumingiana.

DALEA, Gaertn. Fruct. i. 295. t. 61 (1788) = **Mierodon**, Choisy (Selaginch.).
Lippiastrum, Gaertn. l. c. = M. ovatus.

DALEA, Linn. Hort. Cliff. 363. t. 22 (1787); Gen. ed. I. 849 (1737). *LEGUMINOSÆ*, Benth. & Hook. f. i. 492.
ERRAZURIZIA, Phil. in Anal. Univ. Chil. (1872) 689.
JANESIA, Raf. in Atl. Journ. 145 (1832).
PAROSELA, Cav. Descr. 185 (1802).
TRICHOPODIUM, Presl, Bot. Bemerk. 52 (1844).
acutifora, Steud. Nom. ed. II. i. 480 = acutifolia.
acutifolia, *Moç. & Sesse, ex DC. Prod.* ii. 245.—Mexic.
agastachya, Moric. Pl. Nouv. Am. 65. t. 44 = Petalostemum obovatum.
agonocrasis, Steud. Nom. ed. II. i. 480, sphalm. = ayavacensis.
alba, Michx. in Roem. Cat. Hort. Soc. Phys. Turic. (1902) = alopecuroides.
albiflora, *A. Gray, Pl. Wright.* ii. 38.—N. Mexic.
alopecuroides, Blanco, Fl. Filip. ed. II. 289 = nigra.
alopecuroides, *Willd. Sp. Pl.* iii. 1326.—Am. bor.
Alopecurus, *Moç. & Sesse, ex G. Don, Gen. Syst.* ii. 228.—Mexic.
Alopecurus, *Moric. Pl. Nouv. Am.* 10. t. 8.—Mexic.
amœna, *S. Wats. in Am. Natural.* vii. (1873) 800.—Am. bor.
angustifolia, *G. Don, Gen. Syst.* ii. 223.—Mexic.
annua, Thouin, ex Schrank, Pl. Rar. Hort. Monac. t. 9 = Thouini.
arborescens, *Torr. ex A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 316.—Calif.
argentea, *G. Don, Gen. Syst.* ii. 225.—Mexic.
argentea, *Mart. Del. Sem. in Hort. Monac.* (1846); *ex Walp. Ann.* i. 238.—Mexic.
argyræa, *A. Gray, Pl. Wright.* i. 47.—Mexic.
argyrostachys, Hook. & Arn. Bot. Beech. Voy. 385 = lasiostachya.
aristata, Benth. Pl. Hartw. 13 = mucronata.
astragalina, *H. B. & K. Nov. Gen. et Sp.* vi. 484.—Ecuador.
astragalina, Moç. & Sesse, ex DC. Prod. ii. 245 = hypoglottides.
aurea, *Nutt. in Fras. Cat.* (1813), *ex Gen. Am.* ii. 101.—Am. bor.
ayavacensis, *H. B. & K. Nov. Gen. et Sp.* vi. 486.—Mexic.
Berlandieri, *A. Gray, in Proc. Am. Acad.* v. (1861) 177.—Mexic.
bicolor, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 787 = mutabilis.
brachystachya, *A. Gray, Pl. Wright.* ii. 39.—N. Mexic.
californica, *S. Wats. in Proc. Am. Acad.* xi. (1876) 132.—Calif.
calycosa, *A. Gray, Pl. Wright.* ii. 40.—N. Mexic.
canadida, Willd. Sp. Pl. iii. 1337 = Petalostemum candidum.
canescens, *Benth. Bot. Voy. Sulph.* 12.—Calif.
canescens, *Mart. & Gal. in Bull. Acad. Brux.* x. 11. (1843) 48.—Mexic.
carnea, Poir. in Dict. Sc. Nat. xii. 468 = Petalostemum carneum.
chrysothrix, *A. Gray, in Proc. Am. Acad.* v. (1861) 156.—Mexic.
cinerea, *Moric. in Hemsl. Biol. Centr. Am. Bot.* i. 288, nomen.—Mexic.
cinvidoora, Hook. & Arn. ex Hemsl. l. c. 245 = polyphylla.
citriodora, *Willd. Sp. Pl.* iii. 1329 —Mexic.
Clafloriana, Willd. l. c. 1326 = alopecuroides.
comosa, *Schlecht. in Linnæa*, xii. (1838) 299.—Mexic.
compacta, Spreng. Syst. iii. 227 = Petalostemum macrostachyum.

DALEA:—
Coronilla, *G. Don, Gen. Syst.* ii. 226.—Mexic.
corymbosa, Poir. in Dict. Sc. Nat. xii. 462 = Petalostemum corymbosum.
crassifolia, *Hemsl. Biol. Centr. Am. Bot.* i. 288.—Mexic.
crenulata, Hook. & Arn. Bot. Beech. Voy. 385 = nutans.
cylindrica, *Hook. Bot. Misc.* ii. (1831) 213.—Peruv.
decora, S. Schau. in Linnæa, xx. (1847) 743 = pulchella.
diffusa, *Moric. Pl. Nouv. Am.* 8. t. 6.—Mexic.
divaricata, *Benth. Bot. Voy. Sulph.* 12.—Calif.
domingensis, *DC. Prod.* ii. 246.—Ins. S. Doming.
dorycnioides, DC. l. c. 245 = pulchella.
Ehrenbergii, *Schlecht. in Linnæa*, xii. (1838) 290.—Mexic.
elata, *Hook. & Arn. Bot. Beech. Voy.* 416.—Mexic.
elata, *Mart. & Gal. in Bull. Acad. Brux.* x. 11. (1843) 41.—Mexic.
elegans, *Gill. ex Hook. Bot. Misc.* iii. (1833) 183.—Chili.
elegans, Hook. & Arn. Bot. Beech. Voy. 417 = nigra.
Emoryi, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 315.—Mexic.
ennæandra, Fras. Cat. (1813), ex Pursh, Fl. Am. Sept. ii. 741 = laxiflora.
enneaphylla, Bert. ex DC. Prod. ii. 246 = domingensis.
enneaphylla, *Willd. Sp. Pl.* iii. 1338.—N. Granat.
eriophylla, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 340.—Mexic.
ervoides, *Benth. in Hemsl. Diagn. Pl. Nov.* 6.—Mexic.
exilis, *DC. Prod.* ii. 247.—Peruv.
eysenhardtioides, *Hemsl. Diagn. Pl. Nov.* 6.—Mexic.
filiformis, *A. Gray, Pl. Wright.* ii. 39.—N. Mexic.
flava, *Mart. & Gal. in Bull. Acad. Brux.* x. 11. (1843) 40.—Mexic.
flava, Moç. & Sesse, ex DC. Prod. ii. 245 = lutea.
flava, Seem. Bot. Voy. Herald, 280 = simulis.
flavoroses, *Moç. & Sesse, ex DC. Prod.* ii. 246.—Mexic.
formosa, *Torr. in Ann. Lyc. N. York*, ii. (1828) 177.—Am. bor.
Fremontii, *Torr. ex A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 315.—Am. bor. occ.
frutescens, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 175.—Texas.
fruticosa, *G. Don, Gen. Syst.* ii. 226.—Mexic.
gracilis, Kunth, Mim. 126. t. 45 = sericea.
gracilis, Hook. & Arn. Bot. Beech. Voy. 386 = diffusa.
Greggii, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 314.—Mexic.
guatemalensis, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 241, nomen —Guatemala.
Hallii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 625.—Texas.
Hegewischiana, *Steud. Nom.* ed. II. i. 480.—Mexic.
herbacea, Moç. & Sesse, ex G. Don, Gen. Syst. ii. 224 = humilis.
humifuse, *Benth. Pl. Hartw.* 170.—Ecuador.
humilis, *G. Don, Gen. Syst.* ii. 224.—Mexic.
hypoglottoides, *DC. Prod.* ii. 245.—Mexic.
incomposita, *S. Schau. in Linnæa*, xx. (1847) 744.—Mexic.
insignis, *Hemsl. Diagn. Pl. Nov.* 7.—Mexic.
Jamesii, *Torr. & Gray, Fl. N. Am.* i. 295.—Am. bor.
Johnsoni, *S. Wats. in Bot. King's Exp.* 64.—Am. bor. occ.
Kingii, *S. Wats. l. c.* 64. t. 10.—Am. bor. occ.
Kuhnistera, Willd. Sp. Pl. iii. 1337 = Petalostemum corymbosum.
lachnanthe, *S. Schau. in Linnæa*, xx. (1847) 748.—Mexic.
lachnostachys, *A. Gray, Pl. Wright.* i. 46.—Mexic.
laevigata, *A. Gray, l. c.* ii. 38.—N. Mexic.
laevigata, *Moç. & Sesse, ex G. Don, Gen. Syst.* ii. 224.—Mexic.
lagopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 292.
Lagopus, *Willd. Sp. Pl.* iii. 1340.—Mexic.
lanata, *Spreng. Syst.* iii. 227.—Am. bor. occ.
lanuginosa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 307 = lanata.
lasiathera, *A. Gray, Pl. Wright.* i. 48.—Texas.
lasiostachys, *Benth. Pl. Hartw.* 11.—Mexic.
laterijer, Moç. & Sesse, ex DC. Prod. ii. 245 = citriodora.
laxiflora, *Pursh, Fl. Am. Sept.* ii. 741.—Am. bor.
laxiflora, Schlecht. in Linnæa, xii. (1838) 299 = Hegewischiana.
Lemmoni, *Parry, ex A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 206.—Am. bor. occ.

DALEA:—
leporina, Ait. ex Hemsl. Biol. Centr. Am. Bot. i. 287 = alopecuroides.
leptoclados, *Benth. ex Hemsl. l. c.* 242, nomen.—Mexic.
leptostachya, *DC. Prod.* ii. 246.—Am. austr.
leucostachys, *A. Gray, Pl. Fendl.* 12.—Mexic.
leucostoma, *Schlecht. in Linnæa*, xii. (1838) 294—Mexic.
Linnæi, Michx. Fl. Bor. Am. ii. 57. t. 38 = alopecuroides.
longipes, Moç. & Sesse, ex DC. Prod. ii. 245 = sericea.
lutana, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 341.—Mexic.
lutea, *Willd. Sp. Pl.* iii. 1341.—Mexic.
macrostachya, *Moric. Pl. Nouv. Am.* 6. t. 5.—Mexic.
macrotropis, *S. Schau. in Linnæa*, xx. (1847) 742—Mexic.
megacarpa, *S. Wats. in Proc. Am. Acad.* xx. (1885) 359.—N. Mexic.
melantha, *S. Schau. in Linnæa*, xx. (1847) 746.—Mexic.
mollitioides, Moç. & Sesse, ex DC. Prod. ii. 246 = nutans.
microphylla, *H. B. & K. Nov. Gen. et Sp.* vi. 482.—Peruv.
mollis, *Benth. Pl. Hartw.* 206.—Calif.
Moricandi, *D. Dietr. Syn. Pl.* iv. 1015.—Am. austr.
mucronata, *DC. Prod.* ii. 246.—Mexic.
multifoliata, *Phil. f. Cat. Pl. Vasc. Chil.* 55.—Chili.
mutabilis, *Willd. Sp. Pl.* iii. 1339.—Mexic.; Cuba.
Mutisii, *Kunth, Mim.* 161. t. 47.—N. Granat.
nana, *Torr. ex A. Gray, Pl. Fendl.* 31.—Texas.
naviculifolia, *Hemsl. Diagn. Pl. Nov.* 7.—Mexic.
nigra, *Mart. & Gal. in Bull. Acad. Brux.* x. 11. (1843) 43.—Am. austr.
nutans, *Willd. Sp. Pl.* iii. 1339.—Mexic.
obovatifolia, Orteg. Hort. Matr. 82 = mutabilis.
onobrychioides, *Griseb. in Goett. Abh.* xix. (1874) 118.—Reg. Argent.
onobrychioides, Moç. & Sesse, ex DC. Prod. ii. 246 = tomentosa.
Onobrychis, *DC. l. c.* 247.—Peruv.
Orcuttii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 359—Calif.
Ordiae, *A. Gray, l. c.* xvii. (1881–82) 200.—Am. bor. occ.
ornata, Eaton & Wright, N. Am. Bot. 219 = Petalostemum ornatum.
ostafolia, Orteg. Hort. Matr. 80. t. 8 = lutea
Parryi, *Torr. & Gray, ex A. Gray, in Proc. Am. Acad.* vii. (1868) 397.—Am. bor. occ.
parviflora, Pursh, Fl. Am. Sept. ii. 474 = Astragalus gracilis.
parvifolia, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 225.—Ins. Galapag.
pectinata, *Kunth, Mim.* 169. t. 49.—Mexic.
pectinata, Seem. Bot. Voy. Herald, 279 = crassifolia.
pedunculata, Pursh, Fl. Am. Sept. ii. 474 = alopecuroides.
pendulina, Moç. & Sesse, ex DC. Prod. ii. 245 = tuberculata.
penicillata, Moric. Pl. Nouv. Am. 66. t. 45 = laxiflora.
phymatodes, *Willd. Sp. Pl.* iii. 1338.—Venezuela.
platystegia, *S. Schau. in Linnæa*, xx. (1847) 741.—Mexic.
pogonanthera, *A. Gray, Pl. Fendl.* 31.—N. Mexic.
polyosphala, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 244, nomen.—Mexic.
polydenia, *Torr. ex S. Wats. in Bot. King's Exp.* 64. t. 9.—Am. bor. occ.
polygonoides, *A. Gray, Pl. Wright.* ii. 39.—N. Mexic.
polyphylla, *Mart. & Gal. in Bull. Acad. Brux.* x. 11. (1843) 44.—Mexic.
Pringlei, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 201.—Am. bor. occ.
procumbens, *Moç. & Sesse, ex DC. Prod.* ii. 246.—Mexic.
prostrata, *Ortog. Hort. Matr.* 69.—Mexic.
psoraleoides, Moric. Pl. Nouv. Am. 5. t. 4 = verbascoa.
pulchella, *G. Don, Gen. Syst.* ii. 224.—Mexic.
pulchella, *Moric. Pl. Nouv. Am.* 9. t. 7.—Mexic.
pulcherrima, *Moç. & Sesse, ex G. Don, Gen. Syst.* ii. 226.—Mexic.

DALEA :—
purpurea, Vent. Jard. Cels, t. 40 = Petalostemum violaceum.
pyramidalis, *Schlecht. in Linnaea*, xii. (1838) 292.—Mexic.
radicans, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 341.—Mexic.
ramosissima, *Benth. Bot. Voy. Sulph.* 11 t. 10.—Calif.
ramotissima, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 41 = diffusa.
reclinata, *Willd. Sp. Pl.* iii. 1340.—Mexic.
rosea, D. Dietr. Syn Pl. iv. 1017 = Hegewischiana.
rubescens, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 369.—Texas.
scariosa, *S. Wats. l. c.*—N. Mexic.
Schaffneri, *Hemsl. Diagn. Pl. Nov.* 7.—Mexic.
Schottii, *Torr. in Bot. Mex. Bound.* 53.—Am. bor. occ.
scoparia, *A. Gray, Pl. Fendl.* 32.—Texas.
sericea, *Lag. Gen. et Sp. Nov.* 23.—Mexic.
similis, *Hemsl. Diagn. Pl. Nov.* 7.—Mexic.
spinescens, Hemsl. Biol. Centr. Am. Bot. i. 247, sphalm. = sec.
spinosa, A. Gray, in Mem. Am. Acad. N. S. v. (1855) 315 = Asagraea spinosa.
stenophylla, *Griseb. in Goett. Abh.* xix. (1874) 117.—Reg. Argent.
tenuicaulis, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 226.—Ins. Galapag.
tephroloides, *Griseb. Cat. Pl. Cub.* 69.—Cuba.
Thouini, *Schrank, Pl. Rar. Hort. Monac.* t. 9.—Hab.?
thymoides, *Schlecht. & Cham. in Linnaea*, v. (1830) 580.—Mexic.
thyrsiflora, *A. Gray, in Proc. Am. Acad.* v. (1861) 177.—Mexic.
tomentosa, Willd. Sp. Pl. iii. 1241 = verbenacea.
trifoliata, Moric. Pl. Nouv. Am. 3 t. 3 = triphylla.
trifoliata, *Zucc. in Abh. Akad. Muench.* i. (1832) 342; *et in Flora*, xv. (1832) ii. *Beibl.* 69.—Mexic.
triphylla, *Pav. ex Schlecht. in Linnaea*, xii. (1838) 389.—Mexic.
tuberculata, Lag. Gen. et Sp. Nov. 23 = thymoides.
uncifera, *Schlecht. & Cham. in Linnaea*, v. (1830) 580.—Mexic.
unguicularis, Moç. & Sesse, ex DC. Prod. ii. 245 = mutabilis.
velutipes, *Benth. ex Hemsl. Biol. Centr. Am. Bot* i. 248, nomen.—Mexic.
verbenacea, *Schlecht. & Cham. in Linnaea*, v. (1830) 579.—Mexic.
verrucosa, *G. Don, Gen. Syst.* ii. 225.—Mexic.
versicolor, *Zucc. in Abh. Akad. Muench.* i. (1832) 342; *et in Flora*, xv. (1832) ii. *Beibl.* 69.—Mexic.
villosa, Spreng. Syst. iii. 326 = Petalostemum villosum.
violacea, Willd. Sp. Pl. iii. 1337 = Petalostemum violaceum.
virgata, *Lag. Gen. et Sp. Nov.* 23.—Mexic.
vulnerata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 4.—Am. centr.
Wislizeni, *A. Gray, Pl. Fendl.* 32.—N. Mexic.
Wrightii, *A. Gray, Pl. Wright.* i. 49.—N. Mexic.
zimapanica, *S. Schau. in Linnaea*, xx. (1847) 746.—Mexic.
Zuccarinii, Walp. Rep. i. 654 = trifoliata.

DALECHAMPIA, Plum. ex Linn. Gen. ed. I. 375 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 330.
CREMOPHYLLUM, Scheidw. in Bull. Acad. Brux. ix. i. (1842) 28.
RHOPALOSTYLIS, Klotzsch, ex Baill. Adansonia, v. (1864–65) 317.
adscendens, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 640.—Bras.
affinis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 223.—Bras.
alata, *Klotzsch, ex Baill. Adansonia*, v. (1864–65) 309.—Bras.
anisophylla, *Muell. Arg. in Bremen, Abh.* vii. (1880) 29.—Madag.
arctata, *Baill. Adansonia*, v. (1864–65) 314.—Bras.
aristolochiaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 99.—Ecuador.
bahiensis, Muell. Arg. in DC. Prod. xv. ii. 1247 = Ilheotica.
Berneri, *Baill. Adansonia*, i. (1860–61) 277.—Madag.
bidentata, *Blume, Bijdr.* 622.—Java.

DALECHAMPIA:—
bidentata, Thw. Enum. Pl. Zeyl. 270 = indica.
Boiviniana, Baill. Adansonia, i. (1860–61) 277 = tamaifolia.
brasiliensis, Lam. Encyc. ii. 258 = scandens.
brevipes, *Muell. Arg. in Linnaea*, xxxiv. (1865–6) 224.—Bras.
buettnerioides, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1184, nomen.—Guian.
Burchellii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 649.—Bras.
canescens, *H. B. & K. Nov. Gen. et Sp.* ii. 98.—N. Granat.
capensis, Sond. in Linnaea, xxiii. (1850) 106 = natalensis.
capensis, *Spreng. f. Tent. Syst. Veg.* 18.—Afr. austr.
caperonioides, *Baill. Etud. Gén. Euph.* 487, nomen; *Adansonia*, v. (1864–65) 316.—Bras.
cissifolia, Poepp. & Endl. Nov. Gen. iii. 20 = triphylla.
Clausseniana, Baill. Adansonia, v. (1864–65) 310 = triphylla
clematidifolia, *Boy. ex Baill. l. c.* i. (1860–61) 277.—Madag.
colorata, Linn. f. Suppl. 421 = scandens.
colorata, Vell. Fl. Flum. x. t. 58 = brasiliensis.
convolvuloides, *Lam. Encyc.* ii. 256.—Bras.
cordifolia, Domb. ex Lam. l. c. = convolvuloides.
cordofana, Hochst. ex Webb, in Hook. Niger Fl. 178; A. Rich. Tent. Fl. Abyss. ii. 244 = scandens.
crassa, *Klotzsch, ex Baill. Adansonia*, v. (1864–65) 314, nomen; *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 223.—Bras.
coromandelina, Wall. Cat. n. 7798 = indica.
cujabensis, *Mart. ex Baill. Adansonia*, v. (1864–65) 313, nomen; *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 222.—Bras.
denticulata, *Wright, ex Griseb. in Goett. Nachr.* (1865) 181.—Cuba.
digitata, Leandro, ex Baill. Adansonia, v. (1864–65) 309 = pentaphylla.
dioscoreaefolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 20.—Peruv.
ficifolia, *Lam. Encyc.* ii. 258 —Bras.
fimbriata, H. B. & K. Nov. Gen. et Sp. ii. 100 = scandens.
Franciscana, *Baill. Adansonia*, v. (1864–65) 316.—Bras.
Friedrichsthalii, *Muell. Arg. in Flora*, lv. (1872) 45.—Am. centr.
glecomaefolia, *Baill. Adansonia*, v. (1864–65) 314.—Bras.
goyazensis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 643.—Bras.
Grandidiia, *Baill. Adansonia*, v. (1864–65) 312.—Bras.
grandiflora, Mart. ex Baill. l. c. 309 = pentaphylla.
guianensis, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 43.—Guian.
heterophylla, Boj. ex Baill. Adansonia, i. (1860–61) 277 = ternata.
heterophylla, *Vahl, Eclog. Ams.* iii. 44; *Poir. Encyc. Suppl.* ii. 447.—Guian.
hibiscifolia, Spreng. Syst. iii. 86 = scandens.
hibiscoides, H. B. & K. Nov. Gen. et Sp. ii. 102 = scandens.
hispida, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 19.—Am. trop.
Houlletiana, *Baill. Etud. Gen. Euph.* 486. t. 3.—Bras.
humilis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 223.—Bras.
ilheotica, *Wawra, in Oestr. Bot. Zeitschr.* xiii. (1863) 223.—Bras.
indica, *Wight, Ic.* t. 1882.—Ind. or.
ipomoeaefolia, *Benth. in Hook. Niger Fl.* 500.—Afr. trop.
latifolia, Lam. Encyc. ii. 257 = scandens.
Leandri, *Baill. Adansonia*, v. (1864–65) 315.—Bras.
leucophylla, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 219.—Bras.
linearis, *Baill. Adansonia*, v. (1864–65) 316.—Bras.
longipes, *Muell. Arg. in Bremen, Abh.* vii. (1880) 29.—Ins. Nossibé.
magnoliaefolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 219.—Bras.
maximo, Salzm. ex Baill. Adansonia, v. (1864–65) 313 = tiliaefolia.

DALECHAMPIA:—
meridionalis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 648.—Bras.
micrantha, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 19. t. 222.—Bras.
Micromeria, *Baill. Adansonia*, v. (1864–65) 310.—Bras.
microphylla, Muell. Arg. in Linnaea, xxxiv. (1865–66) 225 = glecomaefolia.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 101.—N. Granat.
mollis, Vahl, Eclog. Am. iii. 44 = scandens.
monophylla, Vell. Fl. Flum. x. t. 61 = convolvuloides.
natalensis, *Muell. Arg. in DC. Prod. xv. ii. 1243.*—Afr. austr.
occidentalis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 641.—Bras.
Olfersiana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 220.—Bras.
papposa, Endl. Atakt. 22. t. 20, 21 = scandens
parvifolia, Lam. Encyc. ii. 258 = scandens.
Peckoltiana, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 647.—Bras.
pentaphylla, *Lam. Encyc.* ii. 258.—Bras.
pernambucensis, Baill. Adansonia, v. (1864–65) 311 = scandens.
peruviana, Lam. Encyc. ii. 257 = tiliaefolia.
pruriens, Griseb, Fl. Brit. W. Ind. 51 = tiliaefolia.
Pseudo-clematis, *Baill. Adansonia*, i. (1860–61) 278.—Madag.
pseudotriphylla, *Muell. Arg. in Bremen, Abh.* vii. (1880) 28.—Madag.
psilogyne, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 648.—Bras.
Regnellii, *Muell. Arg. l. c.* 647.—Bras.
Riedeliana, *Muell. Arg. in DC. Prod.* xv. ii. 1286.—Bras.
Rocelians, *Muell. Arg. l. c.* 1283.—Mexic.
rubiformis, Spreng. Syst. iii. 86 = scandens.
rubioides, H. B. & K. Nov. Gen. et Sp. ii. 102 = scandens.
scandens, Kurs, For. Fl. Brit. Burma, 400 (non Linn.).—Burma.
scandens, *Linn. Sp. Pl.* 1054.—Am. trop.
scandens, Vell. Fl. Flum. x. t. 57 = ficifolia.
schizoloba, Muell. Arg. in Linnaea, xxxiv. (1865–66) 221 = arciana.
Sellowiana, *Muell. Arg. l. c.* 220.—Bras.
senegalensis, A. Juss. ex Webb, in Hook. Niger Fl. 174 = scandens.
serrulata, Vell. ex Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 43 = tiliaefolia.
sidaefolia, H. B. & K. Nov. Gen. et Sp. ii. 200 = scandens.
sinuata, *Baill. Adansonia*, i. (1860–61) 278.—Madag.
smilacina, H. B. & K. Nov. Gen. et Sp. ii. 99 = canescens.
spathulata, *Baill. Etud. Gén. Euph.* 487. t. 3. f. 16–30.—Bras.
stenoscapa, *Muell. Arg. in DC. Prod.* xv. ii. 1241.—Bras.
stipulacea, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 221.—Peruv.
subintegra, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 650.—Bras.
subternata, *Muell. Arg. in Bremen, Abh.* vii. (1880) 28.—Madag.
tamaifolia, Muell. Arg. in DC. Prod. xv. ii. 1248 = sec. tamaifolia, *Lam. Encyc.* ii. 256.—Ins. Mascar.; Madag.
tenuiramea, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 222.—Bras.
ternata, *Muell. Arg. in DC. Prod.* xv. ii. 1241.—Zeylan.
tiliaefolia, *Lam. Encyc.* ii. 257.—Bras.
trilobe, Salzm. ex Baill. Adansonia, v. (1864–65) 311 = scandens.
trilobata, Sieber, ex Baker, Fl. Maurit. 217 = tamaifolia.
tripartita, K. Br. in Salt, Abyss. App. p. lxv, nomen = scandens.
triphylla, *Lam. Encyc.* ii. 256.—Bras.
Vahliana, Steud. Nom. ed. II. i. 491 = scandens.
vartiflola, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 221.—Bras.
velutina, *Wight, Ic.* v. t. 1881.—Ind. or.
villosa, Lam. Encyc. ii. 257 = scandens.
volubilis, E. Mey. ex Baill. Etud. Gén. Euph. 487 = capensis.

DALECHAMPIA :—
vulpina, Muell. Arg. in Linnæa, xxxiv. (1865–66) 222 = Leadri.
Weddelliana, *Baill. Adansonia*, v. (1864–65) 315.—Bras.

DALEMBERTIA, Baill. Etud. Gén. Euph. 546. t. 5 (1858). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 339.
Hahniana, *Baill. Adansonia*, xi. (1873) 125.—Mexic.
platanoides, *Baill. l. c.* 124.—Mexic.
populifolia, *Baill. Etud. Gén. Euph.* 546. t. 5.—Mexic.
triangularis, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 218.—Mexic.

DALENIA, Korth. Verh. Nat. Gesch. 248. t. 58 (1842 ?). *MELASTOMACEÆ*, Benth. & Hook. f. i. 758.
Korthalsii, Blume, Mus. Bot. Lugd. Bat. i. 89 = speciosa.
pulchra, Korth. Verh. Nat. Gesch. t. 58 = speciosa.
speciosa, *Korth. l. c.* 244.—Borneo.

DALHOUSIA, Lem. in Orb. Dict. iv. (1849) 593 = seq.

DALHOUSIEA, R. Grah. in Wall. Cat. n. 5339 (1832). *LEGUMINOSÆ*, Benth. & Hook. f. i. 552.
africana, *J. Moore, in Journ. Bot.* xviii. (1880) 3.—Afr. trop.
bracteata, *R. Grah. in Wall. Cat. n.* 5339.—Reg. Himal.
Haematoxylon, Benth. in Ann. Wien. Mus. ii. (1838) 69 = Baphia nitida.
ovalifolia, Benth. l. c. = Baphia polygalacea.
paucisperma, *Griff. Notul.* iv. 445; *Ic. Pl. Asiat.* iv. p. vi. 608.—Assam.
pyrifolia, Benth. in Ann. Wien. Mus. ii. (1838) 69 = Baphia polygalacea.

DALIA, Endl. Gen. 1444 (1841) = DULIA, Adans. = Ledum, Linn. (Ericac.).

DALIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 175 = Dalea, Linn. (Leguminos.).

DALIBARDA, Linn. Sp. Pl. ed. I. 491 (1753) = Rubus, Tourn. (Rosac.).
calycina, DC. Prod. ii. 568 = R. calycinus.
cordata, Steph. in Mém. Soc. Nat. Mosc. i. (1806) 92 = R. Dalibarda.
fragarioides, Michx. Fl. Bor. Am. ii. 300. t. 28 = Waldsteinia fragarioides.
geoides, Pers. Syn. ii. 58 = R. geoides.
latifolia, Blume, Bijdr. 1119 = R. rotundifolius.
lobata, Baldw. in Ell. Sketch, i. 571 = Waldsteinia lobata.
pedata, Steph. in Mém. Soc. Nat. Mosc. i. (1806) 92 = R. pedatus.
pyrifolia, Blume, Bijdr. 1112 = R. pyrifolius.
repens, Linn. Sp. Pl. ed. I. 491 = R. Dalibarda.
ternata, Steph. in Mém. Soc. Nat. Mosc. i. (1806) 129. t. 10 = Waldsteinia sibirica.
violacea, Michx. Fl. Bor. Am. i. 299. t. 27 = R. Dalibarda.

DALLACHYA, F. Muell. Fragm. ix. 140 (1875). *RHAMNEÆ*.
vitiensis, *F. Muell. l. c.*—Austral.

DALRYMPLEA, Roxb. Hort. Beng. 17 (1814); Fl. Corom. iii. 76. t. 279 (1819) = Turpinia, Vent. (Sapindac.).
dumigrensis, Spreng. Syst. i. 771 = T. paniculata.
pomifera, Roxb. Hort. Beng. 17; Fl. Corom. iii. 76. t. 279 = T. pomifera.

DALRYMPLEA, Roxb. Fl. Ind. ed Carey, ii. 421 (1824) = prec.

DALUCUM, Adans. Fam. ii. 393 (1763) = Melico, Linn. (Gramin.).

DALZELLIA, Hassk. in Flora, xlvii. (1865) 593 = Cyanotis, D. Don (Commelin.).
viridiflora, Hassk. l. c. 594 = C. viridiflora.

DALZELLIA, Wight, Ic. v. 84. t. 1919, 1920 (1852) = Terniola, Tul. (Podost.).
foliosa, Wight, l. c. 85 = Ter. foliosa.
Lawii, Wight, l. c. = Ter. Lawii.
pedunculosa, Wight, l. c. = Tristicha pedunculosa.
ramosissima, Wight, l. c. = Ter. ramosissima.
scylanica, Wight, l. c. t. 1919 = Ter. scylanica.

DAMAPANA, Adans. Fam. ii. 393 (1763) = Smithia, Ait. (Legum.).
Mannii, Rafin. Sylva Tellur. 70 = S. sensitiva.

DAMASONIUM, Mill. Gard. Dict. ed. VI (1752). *ALISMACEÆ*, Benth. & Hook. f. iii. 1007.
ACTINOCARPUS, R. Br. Prod. 343 (1810).
Alisma, Mill. Gard. Dict. ed. VIII. n. 1 = stellatum.
angustissimum, Stend. Nom. ed. I. 26 = Echinodorus ranunculoides.
australe, *Salisb. in Trans. Hort. Soc.* i. (1812) 268.—Austral.
Bourgaei, *Coss. Notes Pl. Crit.* 47.—Reg. Mediterr.
californicum, Torr. in Pacif. Rail. Rep. iv. 143. t. 21 = Alisma californicum.
Dalechampii, S. F. Gray, Nat. Arr. Brit. Pl. ii. 217 = stellatum.
flavum, Mill. Gard. Dict. ed. VIII. n. 2 = Limnocharis flava.
maximum, Burm. ex Steud. Nom. ed. II. i. 481 = Limnocharis Plumeri.
minimum, Lange, in Kjoeb. Vidensk. Meddel. (1860) 65 = polyspermum.
minus, Buchen. in Bremen, Abh. ii. (1871) 20 = australe.
polyspermum, *Coss. Notes Pl. Crit.* 47.—Europ.
stellatum, *Thuill. Fl. Par.* ed. II. 186; *Delarb. Fl. Aur.* ed. II. 586.—Europ.
vulgare, Coss. & Germ. Fl. Env. Paris, ii. (1845) 521 = stellatum.

DAMASONIUM, Schreb. Gen. Pl. 242 (1789) = Ottelia, Pers. (Hydroch.).
alismoides, R. Br. Prod. 344, in obs. = O. alismoides.
brasiliense, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 81 = O. brasiliensis.
cynorum, Planch. l. c. 82 = O. ovalifolia.
indicum, Willd. Sp. Pl. ii. 275 = O. alismoides.
javanicum, Blume, Enum. Pl. Jav. i. 80 = O. javanica.
lactucæfolium, Planch. in Ann. Sc. Nat. Sér. III xi. (1849) 81 = O. alismoides.
lanciifolium, Presl, Rel. Haenk. i. 89 = O. alismoides.
ovalifolium, R. Br. Prod. 344 = O. ovalifolia.
seutia, Buch.-Ham. in Wall. Cat. n. 5044 = O. alismoides.
timorense, Zipp. ex Span. in Linnæa, xv. (1841) 477 = O. alismoides.
tranquebaricum, Schult. f. Syst. vii. 1595 = O. alismoides.
ulvæfolium, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 81 = O. ulvæfolia.

DAMATRAS, Reichb. Nom. 93 (1841) = seq.

DAMATRIAS, Reichb. Consp. 111 (1828) = seq.

DAMATRIS, Cass. in Bull. Soc. Philom. (Sept. 1817) 139; et in Dict. Sc. Nat. xli. 471 (1818) = Haplocarpha, Less. (Compos.).
pudica, Cass. in Dict. Sc. Nat. xli. 471.—Afr. austr.

DAMBURNEYA, Rafin. Sylva Tellur. 136 (1838) = Nectandra, Roland. (Laurin.).
maritima, Rafin. l. c. = N. Willdenowiana.

DAMERIA, Endl. Gen. 1444 (1841) = DAUCERIA, Dennst. = Rhabdia, Burm. f. (Myrsin.).

DAMIRONIA, Cass. in Dict. Sc. Nat. lvi. 224 (1828) = Heliyterum, DC. (Compos.).
cernua, Cass. l. c. 225 = H. variegatum.
elegantissima, Cass. l. c. 223 = H. canescens.

DAMMARA, Gaertn. Fruct. ii. 100. t. 103 (1791) = Protium, Burm. f. (Bursera.).
grandiflora, Gaertn. l. c. = P. obtusifolium.

DAMMARA, [Rumph.] Lam. Encyc. ii. 259 (1786), partim = Agathis, Salisb. (Conif.).
alba, Lam. l. c. = A. loranthifolia.
australis, Lamb. Pin. ed. I. ii. 14; ed. II. 78. t. 44 = A. australis.
crassifolia, Knight, Syn. Conif. 45 = A. loranthifolia.
hypoleuca, C. Moore, ex Henk. & Hochst. Nadelhoelz. 217.—N. Caled.
lanceolata, Planch. & Sebert, Not. Bois Nov. Caléd. 169.—N. Caled.

DAMMARA :—
longifolia, Lindl. ex Gord. Pinet. Suppl. 38.—Ins. Fiji.
loranthifolia, Link, Enum. Hort. Berol. ii. 411 = Agathis loranthifolia.
macrophylla, Lindl. in Journ. Hort. Soc. vi. (1851) 271. —Ins. Vanicola.
Moorei, Lem. Jard. Fleur. ii. (1852) Misc. 113 = seq.
Moorii, Lindl. in Journ. Hort. Soc. vi. (1851) 271.—N. Caled.
Motleyi, Parl. in Journ. Bot. i. (1863) 36.—Borneo.
obtusa, Lindl. in Journ. Hort. Soc. vi. (1851) 270.—Ins. N. Hebrid.
orientalis, Lamb. Pin. ed. II. 70 = Agathis loranthifolia.
ovata, C. Moore, ex Gord. Pinet. Suppl. 38.—N Caled.
Peroneti, C. Moore, ex Hook. Kew Journ. iv. (1852) 115. t. 4 = *macrophylla*.
pinnata, Parment. ex Gord. Pinet. Suppl. 42 = Podocarpus latifolius.
robusta, C. Moore, ex F. Muell. in Trans. Pharm. Soc. Vict. ii. (1860) 174.—Austral.
rubriculis, Carr. Comif. ed. II. 618 = A. loranthifolia.
Rumphii, Presl, Epim. Bot. 236 = A. loranthifolia.
silanica, Lam. Encyc. ii. 259 = Diospyros Embliopteris ?
Veitchii, Henk. & Hochst. Nadelhoelz. 216 (quid ?).—Japon.
vitiensis, Seem. in Bonplandia, ix. (1861) 159.—Ins. Fiji.

DAMNACANTHUS, Gaertn. f. Fruct. iii. 18. t. 192 f. 7 (1805). *RUBIACEÆ*, Benth. & Hook. f. ii. 118.
BAUMANNIA, DC. Not. Pl. Rar. Jard. Genèv. vi. 1. t. 1 et 25 (1838).
indicus, *Gaertn. f. l. c.*—Reg. Himal.; Japon.
macrophylla, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 110 = indicus.
major, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 177 = indicus.

DAMPIERA, R. Pr. Prod. 587 (1810). *GOODENOVIEÆ*, Benth. & Hook. f. ii. 540.
LINSCHOTENIA, De Vriese, in Mitch. Trop. Austral. 245 (1849).
adpressa, *A. Cunn. ex DC. Prod.* vii. 508.—Austral.
adpressa, De Vriese, in Verh. Haarl. x. (1854) 100 = lanceolata.
alata, *Lindl. Swan River App.* 27.—Austral.
altissima, *F. Muell. ex Benth. Fl. Austral.* iv. 113.—Austral.
asaurea, De Vriese, in Lehm. Pl. Preiss. i. 400 = linearis.
bicolor, De Vriese, in Nederl. Kruidk. Arch. ii. (1851) 11 = Brownii.
bilocularis, F. Muell. Fragm. ii. 17 = trigona.
Brownii, *F. Muell. l. c.* vi. 29.—Austral.
candicans, *F. Muell. l. c.* x. 86.—Austral.
canescens, De Vriese, in Verh. Haarl. x. (1854) 114. t. 19 = Scævola canescens.
carinata, *Benth. Fl. Austral.* iv. 111.—Austral.
cauloptera, DC. Prod. vii. 504 = alata.
cornuta, Lindl. Swan River App. 27.—Austral.
cuneata, *R. Br. Prod.* 588.—Austral.
Cunninghami, De Vriese, in Nederl. Kruidk. Arch ii. (1851) 11 = lanceolata.
diversifolia, *De Vriese, in Lehm. Pl. Preiss.* i. 402.—Austral.
Drummondi, De Vriese, in Nederl. Kruidk. Arch. ii. (1851) 8 = trilobata.
epiphylloides, De Vriese, in Lehm. Pl. Preiss. i. 402 = alata.
ovata, De Vriese, l. c. 401 = linearis.
ericoephala, *De Vriese, in Verh. Haarl. x.* (1854) 118. t. 21. f. 1–10.—Austral.
eriophora, De Vriese, in Lehm. Pl. Preiss. i. 400 = linearis.
fasciculata, *R. Br. Prod.* 588.—Austral.
fasciculata, DC. Prod. vii. 504 = stricta.
ferruginea, *R. Br. Prod.* 588.—Austral.
ferruginea, Sieber, ex DC. Prod. vii. 504 = stricta.
glabescens, *Benth. Fl. Austral.* iv. 119.—Austral.
glabriflora, *F. Muell. Fragm.* i. 130.—Austral.
hexamastrichea, De Vriese, in Verh. Haarl. x. (1854) 94 = trilobata.

DAMPIERA:—
hederacea, *R. Br. Prod.* 587.—Austral.
incana, *R. Br. l. c.* 588.—Austral.
inundata, De Vriese, in Lehm. Pl. Preiss. i. 404 = Stylidium crassifolium.
junces, *Benth. Fl. Austral.* iv. 115.—Austral.
lanceolata, *A. Cunn. ex DC. Prod.* vii. 503.—Austral.
lanceolata, De Vriese, in Verh. Haarl. x. (1854) 101. t. 17 = adpressa.
laxiflora, De Vriese, l. c. t. 31 = cuneata.
lavandulacea, *Lindl. Swan River App.* 27.—Austral.
leptoclada, *Benth. Fl. Austral.* iv. 116.—Austral.
Lindleyi, De Vriese, in Lehm. Pl. Preiss. i. 402 = alata.
linearis, *R. Br. Prod.* 588.—Austral.
Linschotenii, *F. Muell. Fragm.* vi. 28.—Austral.
loranthifolia, *F. Muell. ex Benth. Fl. Austral.* iv. 116. —Austral.
luteifolia, *F. Muell. Fragm.* x. 11.—Austral.
marifolia, *Benth. Fl. Austral.* iv. 114.—Austral.
melanopogon, Rr. Br. ex De Vriese, in Verh. Haarl. x. (1854) 87 = Brownii.
nervosa, De Vriese, in Nederl. Kruidk. Arch. ii. (1851) 12 = Brownii.
oblongata, R. Br. Prod. 588 = stricta.
oligophylla, *Benth. Fl. Austral.* iv. 115.—Austral.
omissa, De Vriese, in Nederl. Kruidk. Arch. ii. (1851) 10 = Brownii.
ovalifolia, R. Br. Prod. 588 = Brownii.
parvifolia, *R. Br. l. c.* 589.—Austral.
polyraiacea, De Vriese, in Verh. Haarl. x. (1854) 115 = Goodenia phylicoides.
Preissii, *De Vriese, in Lehm. Pl. Preiss.* i. 403.— Austral.
prostrata, *DeVriese, in Lehm. Pl. Preiss.* i. 408 (Quid.!). —Austral.
prostrata, *De Vriese, in Verh. Haarl.* x. (1854) 88 [non prasc.].—Austral.
purpurea, R. Br. Prod. 588 = Brownii.
Reinwardtii, De Vriese, in Verh. Haarl. x. (1854) 97 = Verreuxia Reinwardtii.
repanda, De Vriese, in Lehm. Pl. Preiss. i. 400 = triloba.
repens, DC. Prod. vii. 503 = lavandulacea.
repens, De Vriese, in Verh. Haarl. x. (1854) 114 = Scaevola paludosa.
rosmarinifolia, *Schlecht. in Linnaea*, xx. (1847) 603.— Austral.
rotundifolia, R. Br. Prod. 587 = Brownii.
sacculata, *F. Muell. ex Benth. Fl. Austral.* iv. 111.— Austral.
Scottiana, *F. Muell. Fragm.* xi. 120.—Austral.
sericantha, *F. Muell. ex Benth. Fl. Austral.* iv. 118.— Austral.
spicigera, *Benth. l. c.* 109.—Austral.
stricta, *R. Br. Prod.* 589.—Austral.
stricta, Sieber, ex DC. Prod. vii. 504 = fasciculata.
subspicata, *Benth. Fl. Austral.* iv. 117.—Austral.
suberticillata, De Vriese, in Lehm. Pl. Preiss. i. 403 = fasciculata.
teres, *Lindl. Swan River App.* 27.—Austral.
tomentosa, R. Br. ex DC. Prod. vii. 503 = incana.
triadata, De Vriese, in Lehm. Pl. Preiss. i. 401 = alata.
trigona, *De Vriese, l. c.*—Austral.
triloba, *Lindl. Swan River App.* 27.—Austral.
undulata, R. Br. Prod. 587 = Brownii.
Verreuxii, De Vriese, in Verh. Haarl. x. (1854) 118. t. 20. f. 1–11 = Verreuxia paniculata.
Wellsiana, *F. Muell. Fragm.* x. 13.—Austral.

DANAA, All. Fl. Pedem. ii. 84. t. 68 (1785) = **Physospermum**, Cusson (Umbell.).
aquilegifolia, All. l. c. = P. aquilegifolium.
sylvatica, Lag. Amen. Nat. ii. 97 = P. aquilegifolium.

DANAA, Colla, in Mem. Acc. Torin. xxxvii. (1835) 27. t. 28 = **Senecio**, Tourn. (Compos.).
Vegua, Colla, l. c. 29 = S. discifolius.

DANAË, Medic. Malv. Fam. 79 (1787). *LILIACEAE*, Benth. & Hook. f. iii. 765.
DANAIDIA, Link, Handb. i. 274 (1829).
cestroyna, Webb & Berth. Phyt. Canar. iii. 220. t. 224 = Semele androgyna.
Gayae, Webb & Berth. l. c. = Semele androgyna.
Laurea, *Medic. Malv.* 73.—Gracca ; An. Min. ; Persia.
racemosa, Moench, Meth. 170 = D. Laurea.

DANAIDA, Reichb. Nom. 48 (1841) = prec.

DANAIDIA, Link, Handb. i. 274 (1829) = **Danaë**, Medic. (Liliac.).
racemosa, Link, l. c. = Danaë Laurea.

DANAIS, Comm. ex Vent. Tabl. ii. 584 (1799). *RUBIACEAE*, Benth. & Hook. f. ii. 86.
brevifora, *Baker*, in *Journ. Linn. Soc.* xx. (1883) 163. —Madag.
cernua, *Baker*, in *Journ. Bot.* xx. (1882) 137.—Madag.
Chasseliana, Poir. in Dict. Sc. Nat. xii. 481 = Chasalia capitata.
coronata, Steud. Nom. ed. I. 363 = fragrans.
corymbosa, *Balf. f. in Phil. Trans.* clxvii. (1879) 345. —Ins. Mascar.
denudata, Pers. Syn. i. 198 = fragrans.
fragrans, *Gaertn. f. Fruct.* iii. 88.—Ins. Maurit.; Madag.
Gerrardi, *Baker*, in *Journ. Linn. Soc.* xx. (1883) 160. —Madag.
hispida, *Baker*, *l. c.* 161.— Madag.
laxiflora, DC. Prod. iv. 461 = fragrans.
ligustrifolia, *Baker*, in *Journ. Linn. Soc.* xx. (1883) 162.—Madag.
microcarpa, *Baker*, *l. c.* 168.—Madag.
pauciflora, *Baker*, *l. c.* 163.—Madag.
pubescens, *Baker*, *l. c.* 164.—Madag.
rhamnifolia, *Baker*, *l. c.* —Madag.
rotundifolia, *Poir. Encyc. Suppl.* ii. 450.—Ins. Mascar.
sulcata, *Pers. Syn.* i. 198.—Ins Maurit.
ternata, *Baker*, in *Journ. Linn. Soc.* xx. (1883) 162.— Madag.
verticillata, *Baker*, *l. c.* 164.—Madag.
vestita, *Baker*, *l. c.* xxi. (1885) 408.—Madag.
volubilis, *Baker*, *l. c.* xx. (1883) 161.—Madag.

DANATOPHORUS, Blume, Rumphia, iii. 178 (1847) = **Harpullia**, Roxb. (Sapind.).

DANBYA, Salisb. Gen. Pl. Fragm. 57 (1866) = **Bomarea**, Mirb. (Amaryllid.).
distichiflora, Salisb. l. c. = B. distichiflora.
secundiflora, Salisb. l. c. = B. secundiflola.

DANCERA, Rafin. Sylva Tellur. 92 (1838) = **Clidemia**, D. Don (Melastom.).
hirta, Rafin. l. c. = C. hirta.

DANGERVILLA, Vell. Fl. Flum. 27 (1825) i. t. 66 (1827) = **Cuspuria**, Humb. (Rutac.).
spinosa, Vell. Il. sc. = C. pentagyna.

DANIELIA, Mello, ex Veri. in Rev. Hortic. (1868) 154. *BIGNONIACEAE*.
splendens, *Mello*, *l. c.*—Bras.

DANIELLA, J. J. Benn. in Pharm. Journ. xiv. (1855) 251. *LEGUMINOSAE*. Benth. & Hook. f. i. 580.
oblonga, *Oliver*, *Fl. Trop. Afr.* ii. 300.—Afr. trop.
thurifera, *J. & Benn. in Pharm. Journ.* xiv. (1855) 252.—Afr. trop.

DANTHIA, Steud. Nom. ed. II. i. 483 (1840) = DANTIA, Petit = **Ludwigia**, Linn. (Onagrar.).

DANTHONIA, DC. Fl. Fr. iii. 82 (1805) partim. *GRAMINEAE*, Benth. & Hook. f. iii. 1162.
CHAETOBROMUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836).
CRINIPES, Hochst. in Flora, xxxviii. (1855) 279.
MERATHRETA, Rafin. in Ser. Bull. Bot. i. (1830) 231.
MONACHATHER, Steud. Syn. Pl. Gram. 247 (1855).
PENTAMERIS, Beauv. Agrost. 92. t. 18. f. 3 (1812).
PLINTHANTHESIS, Steud. Syn. Pl. Gram. 14 (1855).
STENOCHAETE, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 421 (1851).
TRIRAPHIS, Nees, Fl. Afr. Austr. 270 (1841).
abyssinica, Hochst. in Flora, xxiv. (1841) i. Intell. 20 = tenuiglumis.
airoides, *Nees*, *Fl. Afr. Austr.* 284.—Afr. austr.
albida, *Hochst. ex Steud. Syn. Pl. Gram.* 244.—Abyss.
Allenii, *Austin*, in *Bull. Torrey Bot. Club*, iii. (1872) 21.—Am. bor.
alpina, J. Buch. in Trans. N. Z. Inst. iv. (1872) 235 = semiannularis.
alpina, Vest, in Flora, iv. (1821) 145 = provincialis.
angulata, *Nees*, *Fl. Afr. Austr.* 313.—Afr. austr.
angustifolia, *Nees*, *l. c.* 302.—Afr. austr.

DANTHONIA:—
Anisopogon, Trin. Sp. Gram. t. 61 = Anisopogon avenaceus.
annosa, *Steud. Syn. Pl. Gram.* 245.—Abyss.
antarctica, *Spreng. Syst.* i. 331; *Hook. f. Fl. N. Zel.* i. 303.—N. Zel.
anthoxanthiformis, *Hochst. in Flora*, xxxviii. (1855) 276.—Abyss.
Archeri, Hook. f. Fl. Tasm. ii. 122. t. 163 = Amphibromus Neesii.
aristidoides, Lehm. ex Nees, Fl. Afr. Austr. 318 = trichotoma.
arundinacea, Steud. Nom. ed. II. i. 482 = Forskälii.
aurcofulva, *E. Desv. in C. Gay, Fl. Chil.* vi. 362.— Chili.
australis, J. Buch. in Trans. N. Z. Inst. iv. (1872) 226 = Raoulii
barbata, *Nees*, *Fl. Afr. Austr.* 301.—Afr. austr.
biparitia, *F. Muell. Fragm.* i. 160.—Austral.
bromoides, *Hook. f. Fl. N. Zel.* i. 303. t. 58.—N. Zel.
Buchananii, J. Buch. Indig. Gr. N. Zeal. 87. t. 35 = Hierochloe redolens.
Buchananii, *Hook. f. Handb. N. Zeal. Fl.* 333.—N. Zel.
Buekeana, *Lehm. ex Nees, Fl. Afr. Austr.* 297.—Afr austr.
cachemyriana, *Jaub. & Spach, Illustr.* iv. 46.—Ind. or.
caespitosa, Gaudich. in Freyc. Voy. Bot. 498 = semiannularis ?
caespitosa, Nees, Lehm Pl. Preiss. ii. 104 = setacea.
californica, *Boland. in Proc. Calif. Acad.* ii (1863) 182. —Calif.
calycina, Reichb. in Moessl. Handb. ed. II. iii. 1880 = provincialis.
calycina, *Rorm. & Schult. Syst.* ii. 691.—Afr. austr.
candida, *Hochst. ex Steud. Syn. Pl. Gram.* 244.— Abyss.
carphoides, *F. Muell. ex Benth. Fl. Austral.* vii. 592. —Austral.
cernua, *Doell*, *in Mart. Fl. Bras.* ii. iii. 101.—Bras.
chilensa, *E. Desv. in C. Gay, Fl. Chil.* vi. 360.—Chili.
cincta, *Nees*, *Fl. Afr. Austr.* 332.—Afr. austr.
Cingula, Steud. Syn. Pl. Gram. 246 = semiannularis ?
circinnata, *Steud. l. c.* 239.—Afr. austr.
cirrhulosa, *Nees*, *Fl. Afr. Austr.* 309.—Afr. austr.
collina, *Phil. in Linnaea*, xxix. (1857–58) 96.—Chili.
collinita, *Nees*, *Fl. Afr. Austr.* 315.—Afr. austr.
colorata, Steud. Syn. Pl. Gram. 241 = tortuosa.
compacta, Boiss. & Heldr. Diagn. Ser. 1. vii. 123 = Avena compacta.
compressa, *Austin*, in *Bull. Torrey, Bot. Club*, iii. (1872) 21.—Am. bor.
coronata, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 70 = lupulina.
crassiuscula, *T. Kirk*, in *Trans. N. Z. Inst.* xvii. (1885) 224.—N. Zel.
crispa, *Nees*, *Fl. Afr. Austr.* 310.—Afr. austr.
Cunninghamii, *Hook. f. Handb. N. Zeal. Fl.* 332.— N. Zel.
curva, *Nees*, *Fl. Afr. Austr.* 328.—Afr. austr.
curvifolia, *Schrad. in Schult. Mant.* ii. 886 —Afr. austr.
cyatophora, *Nees*, *Fl. Afr. Austr.* 296.—Afr. austr.
decora, *Nees*, *l. c.* 332.—Afr. austr.
decumbens, DC. Fl. Fr. iii. 83 = Triodia decumbens.
densiflora, *Nees*, *Fl. Afr. Austr.* 291.—Afr. austr.
denudata, *Nees*, *l. c.* 290.—Afr. austr.
depressa, *Hochst. in Flora*, xxxviii. (1855) 276.—Abyss.
distichа, *Nees*, *Fl. Afr. Austr.* 336.—Afr. austr.
distichophylla, *Lehm. Pugill.* iii. 41.—Afr. austr.
Dregeana, *Steud. Syn. Pl. Gram.* 248.—Afr. austr.
elegans, *Nees*, *Fl. Afr. Austr.* 296.—Afr. austr.
elephantina, *Nees*, *l. c.* 334.—Afr. austr.
elongata, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 20, nomen ; *et ex A. Rich. Tent. Fl. Abyss.* ii. 419.— Abyss.
eriantha, Lindl. in Mitch. Three Exped. ii. 307 = semiannularis.
eriostoma, *Nees*, *Fl. Afr. Austr.* 304.—Afr. austr.
fascicularis, *Steud. Syn. Pl. Gram.* 243.—Afr. austr.
Faxoni, *Austin*, *in Bull. Torrey Bot. Club*, vi. (1875) 190.—Am. bor.
filiformis, *Nees*, *Fl. Afr. Austr.* 292.—Afr. austr.
flaccida, *T. Kirk*, in *Trans. N. Z. Inst.* xvii. (1885) 224 —N. Zel.
flavescens, *Hook. f. Handb N. Zeal. Fl.* 332.—N. Zel.
Forskälii, *Trin. Sp. Gram.* t. 49.—Argypi. ; Arab.
glandulosa, *Schrad. in Schult. Mant.* ii. 885.—Afr. austr.
glauca, *Nees*, *Fl. Afr. Austr.* 327.—Afr. austr.

DANTHONIA :—
glumosa, Benav. Agrost. 92 = sericea.
gracilis, Hook. f. Fl. N. Zel. i. 308. t. 69 = semiannularis.
grandiflora, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 418. —Abyss.
grandiflora, *Phil. in Anal. Univ. Chil.* (1873) 568.—Chili.
Griffithiana, *C. Muell. in Bot. Zeit.* xiv. (1856) 347.—Ind. or.
Gunniana, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 416 = racemosa.
heptamera, *Nees, Fl. Afr. Austr.* 309.—Afr. austr.
hetcropla, *Steud. Syn. Pl. Gram.* 239.—Afr. austr.
hirsuta, *Nees, Fl. Afr. Austr.* 282.—Afr. austr.
hispida, Spreng. Syst. i. 381 = Tristachya leucothrix.
holciformis, *Nees, Fl. Afr. Austr.* 296.—Afr. austr.
intercepta, *Steud. Syn. Pl. Gram.* 248.—Afr. austr.
intermedia, *Vasey, in Bull. Torrey Bot. Club*, x. (1883) 82.—Am. bor.
involucrata, *Schrad. in Schult. Mant.* ii. 388.—Afr. austr.
involuta, *Steud. Syn. Pl. Gram.* 240.—Afr. austr.
juncea, *Trin. in Bull. Sc. Acad Pétersb.* i. (1836) 67. —Afr. austr.
kashmiriana, [*Munro*] *Duthie, in Athins. Gas. N. W. Ind.* 630.—Ind. or.
Koeltlini, Hochst. in Flora, xxiv. (1841) 1. Intell. 20 = D. Streblochaeta.
Kuhlii, *Steud. Syn. Pl. Gram.* 240.—Afr. austr.
lanata, *Schrad. in Schult. Mant.* ii. 386.—Afr. austr.
lappacea, Lindl. in Mitch. Three Exped. i. 313 = Astrebla triticoides.
leonina, Steud. ex Kunth, Enum. Pl. i. 303 = antarctica.
leptophylla, *A. Rich. Tent. Fl. Abyss.* ii. 421.—Abyss.
Lima, *Nees, Fl. Afr. Austr.* 312.—Afr. austr.
Linkii, Kunth, Enum. Pl. i. 315 = semiannularis ?
listida, Trin. Sp. Gram. t. 50 = curvifolia.
longifolia, *R. Br. Prod.* 176.—Austral.
longiglumis, *Nees, Fl. Afr. Austr.* 306.—Afr. austr.
lupulina, *Benav. Agrost.* 152.—Afr. austr.
luzoniensis, *Steud. Syn. Pl. Gram.* 245.—Ins. Luzon.
macrantha, *Schrad. in Schult. Mant.* ii. 385.—Afr. austr.
maxima, Lehm. ex Nees, Fl. Afr. Austr. 337 = Thouarsii.
micrantha, *Nees, Fl. Afr. Austr.* 300.—Afr. austr.
microstoma, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 71 = Thunbergii.
montana, *Doell, in Mart. Fl. Bras.* ii. 111. 101.—Bras.
mutica, *Nees, Fl. Afr. Austr.* 281.—Afr. austr.
nervosa, Hook. f. Fl. Tasm. ii. 121 = Amphibromus Neesii.
neurovillynum, *Steud. Syn. Pl. Gram.* 245.—China.
nuda, *Hook. f. Fl. N. Zel.* ii. 287.—N. Zel.
nutans, *Nees, Fl. Afr. Austr.* 314.—Afr. austr.
obtusifolia, *Nees, in Flora*, xxix. (1846) 190.—Afr. austr.
ovata, *F. Buch. Indig. Grass. N. Zeal.* i. 20.—N. Zel.
pallescens, *Schrad. in Schult. Mant.* ii. 386.—Afr. austr.
pallida, *R. Br. Prod.* 177.—Austral.
pallida, A. Cunn. in Hook. Comp. Bot. Mag. ii. (1836) 370 = Tristemon antarctinum.
papillosa, *Schrad. in Schult. Mant.* ii. 385.—Afr. austr.
papposa, *Nees, Fl. Afr. Austr.* 338.—Afr. austr.
paradoxa, *R. Br. Prod.* 177.—Austral.
patula, *Nees, Fl. Afr. Austr.* 286.—Afr. austr.
pauciflora, *R. Br. Prod.* 177.—Austral.
pauciflora, J. Buch. Indig. Grass. N. Zeal. t. 26 = Triodia exigua.
pectinata, Lindl. in Mitch. Three Exped. ii. 26 = Astrebla pectinata.
penicillata, Benav. Agrost. 92 = racemosa.
penicillata, *F. Muell. Progm.* viii. 135.—Austral.
pentaflora, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 343.—N. Zel.
periantha, Brongn. & Guillem. in Ann. Sc. Nat. Sér. II. xv. (1841) 64, sphalm. *vide orientha.*
picta, *Nees & Meyen, Reise*, i. 311, nomen ; *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 157.—Chili.
pilosa, *R. Br. Prod.* 177.—Austral. ; N. Ze¹
pilosa, Trin. Sp. Gram. t. 51 = setacea.
porosa, *Nees, Fl. Afr. Austr.* 283.—Afr. austr.
propinqua, *Nees, l. c.* 299.—Afr. austr.
provincialis, *DC. Fl. Fr.* iii. 82.—Europ. austr.
pseudo-spicata, *C. Muell. in Bot. Zeit.* xiv. (1856) 348. —Chili.

DANTHONIA :—
pumila, *Nees, Fl. Afr. Austr.* 343.—Afr. austr.
purpurea, *Benav. Agrost.* 160.—Afr. austr.
racemosa, *R. Br. Prod.* 177.—Austral.
radicans, Stend. Syn. Pl. Gram. 248 = Trisetum insulare.
Raoulii, *Steud. l. c.* 246.—N. Zel.
rigida, Raoul, Choix Pl. Nouv. Zél. 12 = Raoulii.
rigida, *Steud. Syn. Pl. Gram.* 243.—Afr. austr.
robusta, *F. Muell. in Trans. Vict. Inst.* i. (1855) 44.—Austral.
rufa, *Nees, Fl. Afr. Austr.* 280.—Afr. austr.
rupestris, *Nees, l. c.* 300.—Afr. austr.
scabra, *Nees, l. c.* 287.—Afr. austr.
scabriflora, *Doell, in Mart. Fl. Bras.* ii. 111. 159.—Bras.
secundiflora, *Y. & C. Presl, Rel. Haenk.* i. 255.—Peruv.
segetalis, *Hochst. in Flora*, xxxviii. (1855) 276.—Abyss.
semiannularis, *R. Br. Prod.* 177.—Austral.
sericantha, *Steud. Syn. Pl. Gram.* 246.—Ecuador.
sericea, *Nutt. Gen. Am.* i. 71.—Am. bor.
sericea, Benav. Agrost. 92 = Avena setacea.
setacea, *R. Br. Prod.* 177.—Austral.
setacea, Hook. f. Fl. Tasm. ii. 121 = semiannularis.
setosa, *Nees, Fl. Afr. Austr.* 325.—Afr. austr.
speciosa, *Lehm. ex Nees, l. c.* 307.—Afr. austr.
spicaeformis, *Benav. Agrost.* 160.—Hab. ?
spicata, *Korn. & Schult. Syst.* ii. 690.—Am. bor.
Streblochaeta, *Steud. Syn. Pl. Gram.* 246.—Abyss.
stricta, *Schrad. in Schult. Mant.* ii. 388.—Afr. austr.
strigosa, Benav. Agrost. 160 = Avena strigosa.
subulata, Hook. f. Fl. Tasm. ii. 121 = setacea.
subulata, *A. Rich. Tent. Fl. Abyss.* ii. 420.—Abyss.
tenella, *Nees, Fl. Afr. Austr.* 324.—Afr. austr.
tenuifolia, *Doell, in Mart. Fl. Bras.* ii. 111. 102.—Bras.
tenuiglumis, *Steud. Syn. Pl. Gram.* 244.—Abyss.
Thomasonii, *F. Buch. Indig. Grass. N. Zeal.* t. 36.—N. Zel.
Thouarii, *Nees, Fl. Afr. Austr.* 337.—Afr. austr.
Thuarii, *Desv. Opusc.* 99.—Afr. austr.
Thunbergii, *Kunth, Rev. Gram.* i. 107.—Afr. austr.
tortuosa, *Trin. Sp. Gram.* t. 68.—Afr. austr.
trichopteryx, *Steud. Syn. Pl. Gram.* 244.—Afr. austr.
trichotoma, *Nees, Fl. Afr. Austr.* 316.—Afr. austr.
trisetoides, *Hochst. ex Steud. Syn. Pl. Gram.* 244.—Abyss.
triticoides, Lindl. in Mitch. Trop. Austral. 365 = Astrebla triticoides.
uberior, *Hochst. in Flora*, xxxviii. (1855) 279.—Abyss.
Unarede, Raoul, Choix Pl. Nouv. Zél. 11. t. 3 = semiannularis.
umbricata, Munro, ex Vasey, Cat. Gr. Un. St. 59, nomen = californica.
varia, Nees, in Lehm. Pl. Preiss. ii. 103 = semiannularis.
villosa, Stend. ex Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1832) 82 = Thunbergii.
violacea, *E. Desv. in C. Gay, Fl. Chil.* vi. 365.—Chili.
virescens, *R. Desv. l. c.* 368.—Chili.
viscidula, *Nees, Fl. Afr. Austr.* 303.—Afr. austr.
Webbiana, *Steud. Syn. Pl. Gram.* 240.—Afr. austr.
Zeyheriana, *Steud. l. c.* 244.—Afr. austr.

DANTIA, Lipp. ex Choisy, in DC. Prod. xiii. ii. 449 (1849) = Escobaxia, Vaill. (Nycteg.).
rubicea, Lipp. l. c. 455 = B. repens.

DANTIA, Petit, ex Linn. Gen. ed. I. 337 (1737) = Ludwigia, Linn. (Onagrac.).
palustris, Des Mont. in Act. Soc. Linn. Bord. xx. (1853) 517 = L. palustris.

DANUBIUNCULUS, Sadler [in Linnev Zeit. 12 Sept. (1845) n. 147] ex Walp. Ann. i. 534 (1848) = Ranunculus, Linn. (Scrophul.).
aconitifolius, Sadler, ll. cc. = L. aquaticus.

DANZLERIA, Bert. ex A. DC. Prod. viii. 224 (1844) = Diospyros, Linn. (Ebenac.).
axillaris, Bert. ex A. DC. l. c. = Diosp. cayennensis.

DAPANIA, Korth. in Nederl. Kruidk. Arch. iii. (1854) 381. *GERANIACEAE*, Benth. & Hook. f. i. 277.
racemosa, *Korth. l. c.*—Sumatra.

DAPHNANDRA, Benth. Fl. Austral. v. 285 (1870). *MONIMIACEAE*, Benth. & Hook. f. iii. 145.
micrantha, *Benth. l. c.*—Austral.
repandula, F. Muell. Fragm. x. 105 = Atherosperma repandulum.

DAPHNE, Tourn. ex Linn. Syst. ed. I (1735). *TH). MELASTACEAE*, Benth. & Hook. f. iii. 190.
ERIOSOLENA, blume, Bijdr. 661 (1825).
GASTRILIA, Rafin. Fl. Tellur. iv. 105 (1836).
LAUREOLA, C. Bauh. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. 147 (1869).
MEZERAUM, C. A. Mey. in Bull. Phys.-Math. Acad Pétersb i. (1843) 358.
MISTRALIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. 147 (1869).
NEMOCTIS, Rafin. Fl. Tellur. iv. 106 (1836).
SANAMUNDA, Neck. Elem. ii. 179 (1790).
SCOPOLIA, Linn. f. Suppl. 60 (1781).
THYMELAEA, Adans. Fam. ii. 285 (1763).
TUMELAIA, Rafin. Fl. Tellur. iv. 105 (1836).
acuminata, Stocks, in Hook. Kew Journ. iv. (1852) 180 : Edgew. & Hohen. Diagn. Ser. I. xii. 103 = oleoides.
albiflora, Wolf, in Oelh. Abbild. Wild. Baeume, ii. t. 33 = D. Mezereum.
alpina, *Linn. Sp. Pl.* 356.—Keg. Mediterr. ; Himal.
alpina, Savi, Due Cent. 98 ; ex Bertol. Fl. Ital. iv. 340 = sericea.
alpina, Sibth. & Sm. Fl. Graec. Prod. i. 261 = oleoides.
altaica, *Pall. Fl. Ross.* i. 58. t. 35.—Sibiria.
americana, Mill. Gard. Dict. ed. VIII. n. 9 (Daphnopsidis sp.).—Ind. occ.
andina, Poepp. & Endl. Nov. Gen. et Sp. ii. 66. t. 191 = Ovidia andina.
angustifolia, C. Koch, in Linnaea, xxii. (1849) 611 = oleoides.
anomala, H. B. & K. Nov. Gen. et Sp. ii. 151 = Ovidia Humboldtii.
Aquilaria, Blanco, Fl. Filip. ed. I. 810 = Wikstroemia ovata.
argentea, Lam. Encyc. iii. 486 = Thymelaea nitida.
argentea, Clarke, Trav. iv. 718 = sericea.
argentea, Sibth. & Sm. Fl. Graec. Prod. i. 258 ; Fl. Graec. t. 356 = Thymelaea Tartonraira.
arsea, *Poir. Encyc. Suppl.* iii. 816.—Oriens.
australis, Cyrill. Pl. Rar. Neap. fasc. ii. t. 16 = sericea.
Bholua, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 68 = cannabina.
Blagwyana, *Freyer, in Flora*, xxi. (1838) 176.—Carniolia.
Blagoyana, Hort. ex Meisn. in DC. Prod. xiv. 585 = sericea.
Bomplandiana, Kunth, Syn. Pl. i. 447 = Daphnopsis Bonplandii.
brasiliensis, Raddi, in Mem. Mod. xviii. Fis. (1820) 391 = Funifera utilis.
buxifolia, Lodch. Fl. Ross. iii. 548 = sericea.
buxifolia, Vahl, Symb. Bot. i. 29 = oleoides.
cachemiriana, Meisn. in DC. Prod. xiv. 585 = oleoides.
cacicola, Berger. Fl. Besses-Pyren. ii. 211 = Thymelaea dioica.
calycina, Lam. Encyc. iii. 488 = Thymelaea nivalis.
calycina, Lapeyr. in Mém. Acad. Toul. i. (1782) 209. t. 15 = Thymelaea calycina.
candicans, Lam. Fl. Fr. iii. 221 = Thymelaea Tartonraira.
candida, Vitm. Summa Pl. ii. 514 = alpina.
canescens, Wall. Cat. n. 1046 = Wikstroemia canescens.
cannabina, Lour. Fl. Cochinch. 336 = Wikstroemia indica.
cannabina, *Wall. in Ax. Res.* xiii. (1830) 385.—Nep. Himal.
canescens, Klotzsch, ex Meisn. in DC. Prod. xiv. 531, 541 = Daphnopsis cannabina.
caucasica, Bieb. Casp. 78 = oleoides.
caucasica, *Pall. Fl. Ross.* i. 58.—Reg. Caucas.
centriflora, H. B. & K. Nov. Gen. et Sp. ii. 150 = Daphnopsis centriflora.
Championi, *Benth. Fl. Hongk.* 296.—China.

DAPHNE:—
chinensis, Spreng. Syst. ii. 237 = odora.
Cneorum, Gueldenst. ex Ledeb. Fl. Ross. iii. 549 = caucasica.
Cneorum, *Linn. Sp. Pl.* 357.—Europ. austr.
Cneorum, Wahlenb. Veg. et Clim. Helv. 72 = striata.
collina, Sm. Spicil. ii. 118 = sericea.
collina, Urv. ex Meisan. in DC. Prod. xiv. 534 = oleoides.
comosa, Adam, in Weber & Mohr, Beitr. i. 53 = glomerata.
cordifolia, Link, Handb. i. 388, sphalm. = *coridifolia*
coriacea, Royle, Illustr. Bot. Himal. 321 = oleoides.
coriacea, Soland. ex Seem. Fl. Vit. 206 = Wikstroemia coriacea.
coridifolia, Lam. Encyc. iii. 437 = Thymelaea coridifolia.
crassifolia, Poir. Encyc. Suppl. iii. 816 = Daphnopsis crassifolia.
cretica, Steud. Nom. ed. II. i. 483 = oleoides.
decandra, *Blume, Bijdr.* 650.—Java.
dioica, Gouan, Illustr. 27. t. 17. *f.* 1 = Thymelaea dioica.
elaeagnoides, Humb. & Bonpl. ex Wiksir. Diss. Daph. ed. II. 40 = Daphnopsis salicifolia.
Elisee, Vis. in Mem. Ist. Venet. vi. (1856) 247 = sericea.
emarginata, Pav. ex Meisan. in DC. Prod. xiv. 522 = Daphnopsis Pavoni.
eriocephala, Royle, Illustr. Bot. Himal. 321 = seq.?
eriocephala, Wall. Cat. n. 1051 = Lasiosiphon eriocephalus.
euphorbioides, Pusch. ex Steud. Nom. ed. II. i. 483 = caucasica
florida, Salisb. Prod. 281 = D. Mezereum.
fortida, Linn. f. Suppl. 228 = Wikstroemia indica.
Fortunei, Benth. in Hook. Kew Journ. v. (1858) 196 = Championi.
Fortuni, Lindl. in Journ. Hort. Soc. i. (1846) 147 = D. Genkwa.
Gardneri, Wall. in Aa. Res. xiii. (1820) 388. t. 9 = Edgeworthia Gardneri.
Genkwa, *Sieb. & Zucc. Fl. Jap.* i. 137. t. 75.—China.
glandulosa, Bertol. Amoen. Ital. 356; Spreng. Syst. ii. 237 = oleoides.
glomerata, *Lam. Encyc.* iii. 438.—Aa. Min.; Reg. Caucas.
gnidioides, *Zaub & Spach, Illustr.* iv. 4. t. 304.—Aa. Min.
gnidioides, Szovits, ex Meisan. in DC. Prod. xiv. 536 = oleoides.
Goldium, *Linn. Sp. Pl.* 357.—Reg. Mediterr.
Hostiana, Lindl. & Paxt. Flow. Gard. i. (1850-51) 170 = D. Mezereum.
humilis, Vahl, ex Willk. & Lange, Prod. Fl. Hisp. i. 299 = Thymelaea thesioides.
hybrida, Lindl. Bot. Reg. t. 1177 = odora.
hypericifolia, Wall. Cat. n. 1048 = Stellera Chamaejasme.
imerina, C. Koch, in Linnaea, xxii. (1849) 613 = glomerata.
inamoena, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 454 = Wikstroemia canescens.
indica, Linn. Sp. Pl. 357 = Wikstroemia indica.
indica, Loisel. Herb. Amat. ii. t. 105 = odora.
indica, Schangin, in Pall. N. Nord. Beitr. vi. 109 = alsatica.
involucrata, *Wall. in Aa. Res.* xiii. (1820) 388.—Ind. or.
japonica, Thunb. in Mus. Nat. Acad. Upsal. xiii. (1792) 106 = odora.
jasminea, Griseb. Spicil. Fl. Rumel. ii. 321 = oleoides.
jasminea, *Sibth. & Sm. Fl. Graec. Prod.* i. 360; *Fl. Graeca*, t. 558.—Graecia.
javanica, Thunb. in Mus. Nat. Acad. Upsal. App. xi. 4 = pendula.
jezoënsis, *Maxim. in Regel, Gartenfl.* (1866) 34.—Japon.
kamtschatica, Maxim. Prim. Fl. Amur. 237 = D. Pseudomezereum.
kiusiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 134.—Japon.
Lagetta, Bonpl. ex Kunth, Syn. Pl. i. 447 = Daphnopsis Bonplandii.
Lagetta, Sw. Prod. Veg. Ind. Occ. 68 = Lagetta lintearia.
lanciifolia, Humb. & Bonpl. ex Wiksir. Diss. Daph. ed. II. 40 = Daphnopsis Humboldtii.
lanuginosa, Lam. Encyc. iii. 436 = Wikstroemia canescens.
laterifiora, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = D. Mezereum.

DAPHNE:—
Laureola, *Linn. Sp. Pl.* 356.—Europ.; Afr. bor.
laurifolia, Willd. ex Kunth, Syn. Pl. i. 446 = Daphnopsis Humboldtii.
Lerchenfeldiana, Schur, Enum. Pl. Transs. 588 = Blagayana ?
linearifolia, *Hart, in Trans. R. Irish Acad.* xxvii. (1866) 442.—Palaestin.
linifolia, Meisan. in DC. Prod. xiv. 528 = Daphnopsis tinifolia.
Liottardi, Vill. Hist. Pl. Dauph. iii. 816 = D. Mezereum.
longifolia, Meisan. in Denkschr. Bot. Ges. Regensb. iii. (1841) 285 = involucrata.
lucida, Loisel. Nouv. Not. Pl. Fr. 17 = oleoides.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 151 = Daphnopsis Humboldtii.
major, Lam. Fl. Fr. iii. 221 = D. Laureola.
Maxii, *Carr. Rev. Hortic.* (1872) 392.—Japon.
mexicana, Spreng. Syst. ii. 236 = Daphnopsis salicifolia.
Meyerum, Gueldenst. ex Ledeb. Fl. Ross. iii. 547 = glomerata.
Mezereum, *Linn. Sp. Pl.* 356.—Europ.; As. bor.
microphylla, Meisan. in DC. Prod. xiv. 533 = jasminea.
monostachya, Willd. Sp. Pl. ii. 420 = Canajera Rheedii
montana, Meisan. in Denkschr. Bot. Ges. Regensb. iii. (1841) 284 = pendula.
mucronata, Royle, Illustr. Bot. Himal. 322. t. 81. f. 2 = oleoides.
multiflora, Thore, ex Reichb. Fl. Germ. Excurs. 165 = D. Laureola.
myrtifolia, Poir. Encyc. Suppl. iii 815 = Thymelaea velutina.
neapolitana, Lodd. Bot. Cab. t. 719 = sericea.
nitida, Vahl, Symb. Bot. iii. 58 = Thymelaea nitida.
obcata, Humb. & Bonpl. in Wiksir. Diss. Daph. ed. II. 40 = Daphnopsis Bonplandii.
occidentalis, Sw. Prod. Veg. Ind. Occ. 63 = Daphnopsis Swartzii.
odora, D. Don, Prod. Fl. Nep. 68 = cannabina.
odora, *Thunb. Fl. Jap.* 159; *Lour. Fl. Cochinch.* 287. —Japon.
odorata, Lam. Fl. Fr. iii. 222 = D. Cneorum.
olearifolia, Lam. Encyc. iii. 440 = sericea.
oleoides, *Schreb. Ic. Descr. Pl. Decad.* i. 13. t. 7; *Linn. Mant.* i. 66.—Europ., As. Min.
oleoides, Tscherniaeff, ex Meisan. in DC. Prod. xiv. 532 = caucasica.
oleoides, Urv. in Mém. Soc. Linn Par. i. (1822) 299 = gnidioides.
oppositifolia, Buch.-Ham. ex Meisan. in DC. Prod. xiv. 547 = Wikstroemia canescens.
orientalis, Willd. ex Steud. Nom. ed. II. i. 483 = Thymelaea villosa.
orthophylla, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 123. t. D. Gnidium.
osyroides, Vahl, ex Steud. Nom. ed. II. i. 483 = Thymelaea Tartonraira.
paniculata, Lam. Fl. Fr. iii. 222 = D. Gnidium.
papyracea, Wall. ex Steud. Nom. ed. II. i. 483 = cannabina.
papyrifera, Buch.-Ham. ex Meisan. in DC. Prod. xiv. 537 = cannabina.
papyrifera, Siebold, in Verh. Bat. Gen. xii. (1830) 24 = Edgeworthia chrysantha.
pauciflora, Span. ex Walp. Ann. i. 589 = Wikstroemia Spanoghbi.
pendula, *Sm. Ic. Ined.* ii. 24. t. 34.—Burma; Malaya.
petraea, *Leybold, in Flora*, xxxvi. (1853) 81.—Tyrol. mer.
Philippi, Gren. & Godr. in Schultz, Arch. Fl. Fr. et All. 282 = D. Laureola.
Pillopillo, C. Gay, Fl. Chil. v. 315 = Ovidia Pillopillo.
polystachya, Willd. Sp. Pl. ii. 420 = Canajera Rheedii.
pontica, *Linn. Sp. Pl.* 357.—As. Min.
prostrata, Salisb. Prod. 281 = D. Cneorum.
Pseudo-mezereum, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858-59) 404.—Japon.
pubescens, Linn. Mant. i. 66 = Thymelaea virgata ?
Rodriguezii, *Texidor, in Rev. Progr. Cienc.* xviii. (1849). —*Cf. Bull. Soc. Bot. Fr.* xvi. (1869) *Rev. Bibl.* 223. —Ins. Minorca.
rosmarinoides, Burm. f. Fl. Cap. Prod. 12 (Passerinae sp.?) —Afr. austr.
rotundifolia, Linn. f. Suppl. 228 = Wikstroemia rotundifolia.
Roumea, *Meisan. in DC. Prod.* xiv. 538.—China.
rupestris, Facch. ex Nym. Suppl. Syll. 67 = petraea.
salicifolia, H. B. & K. Nov. Gen. et Sp. ii. 156 = Daphnopsis salicifolia.

DAPHNE:—
salicifolia, Lam. Encyc. iii. 438 = caucasica.
sempervirens, Salisb. Prod. 281 = D. Laureola.
sericea, D. Don, Prod. Fl. Nep. 69 = Wikstroemia canescens.
sericea, Noë, ex Meisan. in DC. Prod. xiv. 534 = oleoides.
sericea, *Vahl, Symb. Bot.* i. 36.—Ital.; As. Min.
sinensis, Lam. Encyc. iii. 438 = odora.
Sophia, *Kalenicz. in Bull. Soc. Mosc.* xxii. (1849) 1 811. t. 3.—As. occ.
squarrosa, Linn. Sp. Pl. 356 = Gnidia carinata.
striata, *Tratt. in Arch. Gewaechsk.* i. 120. t. 138.—Alp.; Europ. austr.
tangutica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 531.—China.
Tartonraira, Linn. Sp. Pl. 356 = Thymelaea Tartonraira.
tenuiflora, *Span. Ic. n.* 64; *et in Linnaea*, xv. (1841) 335, nomen.—Ins. Timor.
thesioides, Lam. Encyc. iii. 437 = Thymelaea thesioides.
Thesmeinii, Lhotzky, ex Meisan. in Mart. Fl. Bras. v. 1. 67 = Funifera utilis.
Thymelaea, Linn. Sp. Pl. 356 = Thymelaea Sanamunda.
tinifolia, Sw. Prod. Veg. Ind. Occ. 63 = Daphnopsis tinifolia.
tomentosa, Lam. Encyc. iii. 436 = Thymelaea Tartonraira.
triflora, *Lour. Fl. Cochinch.* 286.—China.
trifolia, J. F. Gmel. Syst. 1538, sphalm. = *tinifolia*.
velloziodes, Rodrig. in Bull. Soc. Bot. Fr. xvi. (1869) 238 = Thymelaea velutina
Verlotii, Gren. & Godr, Fl. Fr. iii. 59 = D. Cneorum.
vermiculata, Vahl, Symb Bot. i. 28 = Thymelaea tinctoria.
villosa, Linn. Sp. Pl. ed. II. 510 = Thymelaea villosa.
virgata, Wall. Cat. n. 1047 = Wikstroemia canescens.
viridiflora, Mor. & Zoll. ex Meisan. in DC. Prod. xiv. 542 = Wikstroemia Candolleana.
viridiflora, Wall. Cat. n. 1049 = Wikstroemia indica.
Wallichii, Meisan. in DC. Prod. xiv. 540 = involucrata.

DAPHNEPHYLLUM, Hassk. Cat. Hort. Bogor. Alt. 232 (1844) = **Daphniphyllum**, Blume (Euphorb.).

DAPHNICON, Pohl, in Flora, viii. (1825) 183, nomen = **Hippocratea**, Linn.

DAPHNIDIUM, Nees, in Wall. Pl. As. Rar. ii. 61, 63 (1831); et Laurin. Expos. 19 (1833) = **Lindera**, Thunb. (Laurin.).
acuminatum, Blume, Mus. Bot. Lugd. Bat. i. 351 = L. acuminata.
argenteum, Kurz, in Journ. As. Soc. Beng. xlii (1873) ii. 108.—Burma.
bibracteatum, Nees, Syst. Laurin. 615.—Java.
bifarium, Nees, in Wall. Pl. As. Rar. ii. 63 = L. heterophylla.
caesium, Miq. Pl. Jungh. i. 185 = L. densifolia.
caesium, Nees, Syst. Laurin. 614 = L. caesia.
caesium, Zoll. Verz. 114 = L. acuminata.
caudatum, Nees, ex Wall. Pl. As. Rar. ii. 63.—Ind. or.
Cubeba, Nees, Syst. Laurin. 615 = Litsea Cubeba.
elongatum, Nees, ex Wall. in Pl. As. Rar. ii. 63 = Litsea elongata.
gemmiflorum, Blume, Mus. Bot. Lugd. Bat. i. 352.—Java.
gracile, Nees, in Otto & Dietr. Allg. Gartenz. i. (1833) 91.—Hab.?
lancifolium, Sieb. & Zucc. in Abh. Acad. Muench. iv. iii. (1846) 207 = Litsea lancifolia.
lancifolium, Thw. Enum. Pl. Zeyl. 267.—Zeylan.
macrophyllum, Blume, Mus. Bot. Lugd. Bat. i. 352.—Java.
melastomaceum, Nees, in Wall. Pl. As. Rar. ii. 63 = L. melastomaceus.
Myrrha, Nees, Syst. Laurin. 642 = Litsea trinervia ?
Myrrha, Sieb. & Zucc. in Abh. Acad. Muench. iv. iii. (1846) 207 = L. strychnifolia.
oxyphyllum, Nees, in Wall. Pl. As. Rar. ii. 63.—Ind. or.
pendulifolium, Meisan. in DC. Prod. xv. i. 231.—Hab.?
pulcherrimum, Nees, in Wall. Pl. As. Rar. ii. 63.—Nepal.
sinense, Blume, Mus. Bot. Lugd. Bat. i. 352.—China.
strychnifolium, Sieb. & Zucc. in Abh. Acad. Muench. iv. iii. (1846) 207 = L. strychnifolia.
tenuosum, Meisan. in DC. Prod. xv. i. 231.—Bhotan.

DAPHNIDOSTAPHYLIS, Klotzsch, in Linnæa, xxiv. (1851) 79 = **Arotostaphylos**, Adans. (Ericac.).
acuta, Klotzsch, l. c. 81 = A. Hookeri!
cordifolia, Klotzsch, l. c. 80 = A. tomentosa.
Fendleriana, Klotzsch, l. c. = A. Uva-ursi.
glauca, Klotzsch, l. c. = A. glauca.
Hookeri, Klotzsch, l. c. = A. Hookeri.
pumila, Klotzsch, l. c. 81 = A. pumila.
pungens, Klotzsch, l. c. 80 = A. pungens.
tomentosa, Klotzsch, l. c. 79 = A. tomentosa.

DAPHNIPHYLLOPSIS, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 201 = **Nyssa**, Gronov. (Cornac.).
capitata, Kurz, l. c. = N. sessiliflora.

DAPHNIPHYLLUM, Blume, Bijdr. 1152 (1826).
EUPHORBIACEAE, Benth. & Hook. f. iii. 382.
GOUGHIA, Wight, Ic. v. 22. t. 1878, 1879 (1852).
GYRANDRA, Wall. Cat. n. 8040 (1847 ?).
scutifolium, *Muell. Arg. in DC. Prod.* xvi. 1. 8.—Java.
africanum, *Muell. Arg. in Flora*, xlvii. (1864) 586.—Afr. trop.
bancanum, *Kurz, in Tijdschr. Nederl. Ind.* xxvii. (1864) 188.—Java.
Benthamii, Baill. Etud. Gén. Euph. 565, nomen = himalayense.
Blumeanum, *Baill. l. c.*—Java.
Buergeri, *Muell. Arg. in DC. Prod.* xvi. 1. 8.—Japon.
Calleryanum, Baill. Etud. Gén. Euph. 565 = calycinum.
calycinum, *Benth. Fl. Hongk.* 316.—Hongkong.
Gaudichaudianum, Baill. Etud. Gén. Euph. 565, nomen = calycinum.
glaucescens, *Blume, Bijdr.* 1158.—Ind. or.; Java; Cores.
himalayense, *Muell. Arg. in DC. Prod.* xvi. 1. 4.—Reg. Himal.
humile, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* ii. 488.—Japon.
laurinum, *Baill. Etud. Gén. Euph.* 565. t. 21.—Malaya.
macropodium, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 129.—Japon.
majus, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 76.—Burma.
neilgherrense, Thw. Enum. Pl. Zeyl. 290 = glaucescens.
Pierrei, *Hance, in Journ. Bot.* xiv. (1876) 261.—Cambodia.
Roxburghii, Baill. Etud. Gén. Euph. 565 = glaucescens.
Teysmanni, *Kurz, ex Teysm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 89.—Java.
Zollingeri, *Muell. Arg. in DC. Prod.* xvi. 1. 2.—Java.

DAPHNITIS, Spreng. Syst. i. 370 (1825) = **Botryocrasa**, Willd. (Anacard.).
capensis, Spreng. l. c. 454 = B. laurinum.
madagascarensis, Spreng. l. c.—Madag.

DAPHNOBRYON, Meisn. in DC. Prod. xiv. 586 (1857) = **Drapetes**, Banks (Thymel.).
ericoides, Meisn. l. c.—Borneo.
tasmanicum, Meisn. l. c. = Drapetes tasmanica.

DAPHNOPSIS, Mart. & Zucc. Nov. Gen. et Sp. i. 65 (1824). *THYMELAEACEAE*, Benth. & Hook. f. iii. 191.
COLEOPHORA, Miers, in Ann. Nat. Hist. Ser. II. vii. (1851) 197.
HARGASSERIA, Schlede & Deppe, ex C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 356.
NORDMANNIA, Fisch. & Mey. ex C. A. Mey. l. c. 355.
alpestris, *Benth. & Hook. f. Gen.* iii. 192.—Bras.
angustifolia, *Wright, ex Griseb. Cat. Pl. Cub.* 110.—Cuba.
bogotensis, *Meisn. in DC. Prod.* xiv. 521.—Am. austr.
Bonplandii, *Meisn. l. c.*—Mexic.
brasiliensis, *Mart. & Zucc. Nov. Gen. et Sp.* i. 65.—Bras.
caracasana, *Meisn. in DC. Prod.* xiv. 521.—Venezuela.
caribæa, Griseb. Fl. Brit. Ind. 378 = tinifolia, *Meisn.*
cestrifolia, *Meisn. in DC. Prod.* xiv. 523.—N. Granat.
crassifolia, *Meisn. l. c.* 524.—Ins. S. Doming.
cubensis, Meisn. l. c. 523 = Lasiadenia cubensis.
cuneata, *Radlk. in Sitzb. Acad. Muench.* xiv. (1884) 489.—Cuba.
dioica, Mart. ex Meisn. in Mart. Fl. Bras. v. i. 66 = brasiliensis.
GUADICOS, *Wright, ex Griseb. Cat. Pl. Cub.* 110.—Cuba.

DAPHNOPSIS:—
Humboldtii, *Meisn. in DC. Prod.* xiv. 520.—Am. austr.
Lindeni, *Meisn. l. c.* 523.—Mexic.
Martii, *Meisn. in Mart. Fl. Bras.* v. i. 66. t. 28. f. 2.—Bras.
Pavoni, *Meisn. in DC. Prod.* xiv. 522.—Peruv.
Purdiei, *Meisn. l. c.*—N. Granat.
racemosa, *Griseb. in Goett. Abh.* xxiv. (1879) 134.—Reg. Argent.
salicifolia, *Meisn. in DC. Prod.* xiv. 522.—Mexic.
Swartzii, *Meisn. l. c.*—Ind. occ.
tinifolia, *Griseb. Fl. Brit. W. Ind.* 378.—Jamaic.
tinifolia, *Meisn. in DC. Prod.* xiv. 523.—Ind. occ.
utilis, *Warm. in Kjoeb. Vidensk. Meddel.* (1871) 818.—Bras.

DARBYA, A. Gray, in Am. Journ. Sc. Ser. II. i. (1846) 388 = **Buckleya**, Torr. (Santalac.).
umbellulata, A. Gray, l. c. 396.—Am. bor.

DARGERIA, Decne. in Jacquem. Voy. Bot. 115, t. 121 (1844) = **Leptorhabdos**, Schrenk (Scrophul.).
linifolia, Decne. l. c. 116. t. 121 = L. tinifolia.
pinnatifida, Decne. l. c. = L. Benthamiana.

DARLINGIA, F. Muell. Fragm. v. 152 (1866).
PROTEACEAE, Benth. & Hook. f. iii. 182.
spectatissima, *F. Muell. l. c.*—Austral.

DARLINGTONIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 97 = **Desmanthus**, Willd. (Legum.).
brachyloba, DC. Lég. Mém. xii. t. 66 = Desm. brachylobus.
brevifolia, Rafin. New Fl. Am. i. 42 = Desm. brachylobus.
glandulosa, DC. Lég. Mém. xii. t. 66 = Desm. brachylobus.
illinoënsis, DC. ex Torr. in Ann. Lyc. N. York, ii. (1828) 191 = Desm. brachylobus.
interior, Torr. l. c. = Desm. brachylobus.
virgata, Rafin. New Fl. Am. i. 43 = Neptunia lutea.

DARLINGTONIA, Torr. in Smithson. Contrib. vi. (1854) 4. t. 12. *SARRACENIACEAE*, Benth. & Hook. f. i. 48.
californica, *Torr. l. c.* 5.—Calif.

DARLUCA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 87 = **Faramea**, Aubl. (Rubiac.).
prostrata, Rafin. New Fl. Am. iv. 55 = F. odoratissima.

DARTUS, Lour. Fl. Cochinch. 123 (1790). *SOLANACEAE*? Benth. & Hook. f. ii. 888.
perlarius, *Lour. l. c.* 124.—China; Ins. Moluce.

DARWINIA, Dennst. Schluess. Malab. 31 (1818) = **Litsea**, Lam. (Laurin.).
quinquefiora, Dennst. l. c. = L. zeylanica.

DARWINIA, Rafin. Fl. Ludov. 106 (1817) = **Seebania**, Pers. (Legum.).
exaltata, Rafin. l. c. = S. macrocarpa.

DARWINIA, Rudge, in Trans. Linn. Soc. xi. (1813) 299. t. 22. *MYRTACEAE*, Benth. & Hook. f. i. 697.
CAYPTOSTEMON, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 114.
GENETYLLIS, DC. Prod. iii. 209 (1828).
FRANCISCIA, Endl. Gen. 1226 (1840).
HEDAROMA, Lindl. Swan River App. 7. t. 2 B (1839).
POLYZONE, Endl. in Ann. Wien. Mus. ii. (1838) 191.
SCHUERMANNIA, F. Muell. in Linnæa, xxv. (1852) 386.
axillaris, F. Muell. Census, 51 = Chamaelaucium axillare.
brevifolia, F. Muell. l. c. = Chamaelaucium brevifolium.
brevistyla, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) t. 152 = thymoides.
ciliata, F. Muell. Census, 51 = Chamaelaucium ciliatum.
citriodora, *Benth. in Journ. Linn. Soc.* ix. (1867) 160.—Austral.
dicomoides, *Benth. l. c.*—Austral.
Drummondii, *Turcz. in Bull. Soc. Nat. Mosc.* (1847) t. 155.—Austral.
Endlicheri, F. Muell. Census, 51 = Chamaelaucium virgatum.

DARWINIA:—
fascicularis, *Rudge, in Trans. Linn. Soc.* xi. (1815) 299. t. 22.—Austral.
fimbriata, *Benth. in Journ. Linn. Soc.* ix. (1867) 179.—Austral.
flawescens, A. Cunn. ex Schau. in Nov. Act. Nat. Cur xix. Suppl. ii. (1841) 193 = Homoranthus virgatus.
Forrestii, *F. Muell. Fragm.* xi. 9.—Austral.
gracilis, F. Muell. Census, 51 = Chamaelaucium gracile
helichrysoides, *Benth. in Journ. Linn. Soc.* ix. (1867) 179.—Austral.
heterandra, F. Muell. Census, 51 = Chamaelaucium heterandrum.
Hookeriana, *Benth. in Journ. Linn. Soc.* ix. (1867) 179.—Austral.
intermedia, A. Cunn. ex Schau. in Nov. Act. Nat. Cur xix. Suppl. ii. (1841) 190 = taxifolia.
laxiflora, Schau. l. c. = taxifolia.
macrostegia, *Benth. in Journ. Linn. Soc.* ix. (1867) 179.—Austral.
megalopetala, F. Muell. Census, 51 = Chamaelaucium megalopetalum.
Meisneri, *Benth. in Journ. Linn. Soc.* ix. (1867) 179.—Austral.
micropetala, *Benth. l. c.* 181.—Austral.
Neildiana, *F. Muell. Fragm.* ix. 177.—Austral.
oederoides, *Benth. in Journ. Linn. Soc.* ix. (1867) 179.—Austral.
Oldfieldii, *Benth. l. c.* 180.—Austral.
pauciflora, *Benth. l. c.*—Austral.
pinifolia, *Benth. l. c.* 181.—Austral.
purpurea, *Benth. l. c.* 180.—Austral.
radinophylla, *F. Muell. Fragm.* ix. 175.—Austral.
sanguinea, *Benth. in Journ. Linn. Soc.* ix. (1867) 181.—Austral.
saturejæfolia, Turcz. in Bull. Phys.-Math. Acad Pétersb. x. (1852) 324 = thymoides ?
Schraderi, *Benth. in Journ. Linn. Soc.* ix. (1867) 181.—Austral.
species, *Benth. l. c.* 179.—Austral.
taxifolia, *A. Cunn. in Field, N. S. Wales*, 352.—Austral.
Thomasi, *Benth. in Journ. Linn. Soc.* ix. (1867) 181.—Austral.
thymoides, *Benth. l. c.* 180.—Austral.
Turczaninowii, F. Muell. Census, 51 = Chamaelaucium pauciflorum.
uncinata, F. Muell. l. c. = Chamaelaucium uncinatum.
verticillata, *Benth. in Journ. Linn. Soc.* ix. (1867) 181.—Austral.
vestita, *Benth. l. c.* 180.—Austral.
virescens, *Benth. l. c.* 179.—Austral.
virgata, *F. Muell. Fragm.* ix. 176.—Austral.

DARWYNIA, Reichb. Nom. 176 (1841) = prasc.

DASANTHERA, Rafin. in Am. Monthly Mag. (1818) 267 = **Pentstemon**, Mitch. (Scrophul.).
cuneifolia, Rafin. New Fl. Am. ii. 69 = Herpestis nigrescens.
fruticosa, Rafin. in Am. Monthly Mag. (1818) 267 = F. Menziesii.

DASIANTHERA, Presl, Rel. Haenk. ii. i. 90 (1831) = **Scolopia**, Schreb. (Bixin.).
lusoniensis, Presl, l. c. = S. Dasyanthera.

DASIOGYNA, Rafin. Atl. Journ. 145 (1832) (Quid ?).
glandulosa, Rafin. l. c.—Am. bor.

DASIOLA, Rafin. Neogenyt. 4 (1825) = **Festuca**, Linn (Gramin.).
Biliotes, Rafin. l. c. = F. monandra.

DASIORIMA, Rafin. Fl. Tellur. ii. 45 (1836) = **Solidago**, Vaill. (Compos.).
Biliotî, Rafin. l. c. = S. Elliottii.
imeniophila, Rafin. l. c. = S. sempervirens.
menziesa, Rafin. l. c. = S. sempervirens.

DASISTEMA, Rafin. New Fl. Am. ii. 66 (1836) = DASISTOMA, Rafin.

DASISTEPHA, Rafin. Fl. Tellur. iii. 19 (1836) = **Gentiana**, Tourn.
asclepiadea, Rafin. l. c. = G. asclepiadea.
aspidosa, Rafin. l. c. = G. glauca.

DASISTOMA, Rafin. in Journ. Phys. lxxxix. (1819) 99 = **Gerardia**, Linn. (Scrophul.).
aurea, Rafin. l. c. = *G. flava*.
auriculata, Rafin. New Fl. Am. ii. 67 = G. auriculata.
Drummondii, Benth. in DC. Prod. x. 520 = G. grandiflora.
integrifolia, A. Gray, Man. Bot. N. U. St. 307 = G. laevigata.
lacinistum, Walp. Rep. vi. 649 = Seymeria virgata.
laevigata, Rafin. ex Chapm. Fl. S. U. St. Suppl. 636 = G. laevigata.
macrophylla, Rafin. New Fl. Am. ii. 67 = G. macrophylla.
patula, Chapm. in Coult. Bot. Gaz. iii. (1878) 10 = G. patula.
pectinata, Benth. in DC. Prod. x. 520 = G. pedicularia.
pedicularia, Benth. l. c. 521 = G. pedicularis.
pubescens, Benth. l. c. 520 = G. flava.
quercifolia, Benth. l. c. = G. quercifolia.
tubulosa, Bertol. Mic. Bot. xiii. 17 = Macranthera fuchsioides.

DASSOVIA, Neck. Elem. i. 257 (1790) = **Asclepias**, Linn.

DASTYLEPIS, Rafin. Fl. Tellur. iv. 126 (1836) = **Cuscuta**, Linn. (Convolv.).
Brownei, Rafin. l. c. = C. americana!

DASURUS, Salisb. Gen. Pl. Fragm. 51 (1866) = **Chamaelirium**, Willd. (Liliac.).
luteus, Salisb. l. c. = C. luteum.

DASUS, Lour. Fl. Cochinch. 141. (*INC. SED.*).
DASYS, Lem. in Orb. Dict. iv. 618 (1849).
DASUS, Juss. in Dict. Sc. Nat. xxvii. 201 (1823).
verticillatus, *Lour. l. c.*—Cochinch.

DASYANTHERA, Reichb. Handb. 274 (1837) = DASIANTHERA, Fral = **Scolopia**, Schreb. (Bixin.).

DASYANTHES, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 165 = **Erica**, Linn.
Sparmanni, D. Don, l. c. 166 = E. Sparmanni.

DASYAULUS, Thw. Enum. Pl. Zeyl. 175 (1860) = **Bassia**, Linn. (Sapotac.).
fulvus, Thw. l. c. 176 = B. fulva.
microphyllus, Thw. l. c. 175 = B. microphylla.
Moonii, Thw. l. c. 176 = B. Moonii.
nerifolius, Thw. l. c. 175 = B. nerifolia.

DASYCARPUS, Oerst. in Kjoeb. Vidensk. Meddel (1856) 27 = **Sloanea**, Linn. (Tiliac.).
quadrivalvis, Oerst. l. c. = S. quadrivalvis.

DASYCARYA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 98 = **Cyrtocarpa**, H. B. & K. (Anacard.).
grisea, Liebm. l. c. 99 = C. procera.

DASYCEPHALA, Benth. & Hook. f. Gen. ii. 143 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
alata, *Benth. & Hook. f. l. c.* 144.—Bras.
barbata, *Benth. & Hook. f. l. c.*—Ins. S. Doming.
hispidula, *Benth. & Hook. f. l. c.*—Bras.
indecora, *Hemsl. Biol. Centr.-Am. Bot.* ii. 57.—Mexic.
spicata, *Benth. & Hook. f. Gen.* ii. 144.—Guian.

DASYCOLEUM, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 414. *MELIACEAE*, Benth. & Hook. f. i. 385.
Beccarianum, *Baill. Adansonia*, xi. (1874) 268.—Borneo.
Cuminginnum, *C. DC. in DC. Monog. Phan.* i. 641.—Ins. Philipp.
phillippinum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 415.—Ins. Philipp.
sarawakanum, *C. DC. in DC. Monog. Phan.* i. 641.—Borneo.

DASYLEPIS, Oliver, in Journ. Linn. Soc. ix. (1867) 170. *BIXINEAE*, Benth. & Hook. f. i. 972.
racemosa, *Oliver, l. c.* 171.—Afr. trop.

DASYLIRION, Zucc. in Otto & Dietr. Allg. Gartenz. vi. (1838) 288, 303. *LILIACEAE*, Benth. & Hook. f. iii. 780.
acrotrichum, *Zucc. in Abh. Akad. Munech.* iii. (1843) 238.—Mexic.

DASYLIRION:—
Berlandieri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 240.—Mexic.
Bigelowii, Torr. in Pacif. Rail. Rep. iv. 151 = Nolina Bigelovii.
caespitosum, Scheidw. in C. Koch, Wochenschr. iv. (1861) 326 = Hookeri.
erumpens, Torr. in Bot. Mex. Bound. 216 = Nolina erumpens
flexile, *C. Koch, Ind. Sem. Hort. Berol.* (1867) *App.* i. 5.—Mexic.
glaucophyllum, *Hook. Bot. Mag.* t. 5041.—Mexic.
glaucum, Carr. in Rev. Hortic. (1872) 435 = glaucophyllum.
gracile, Zucc. in Abh. Akad. Muench. iv. ii. (1845) 22 = acrotrichum.
graminifolium, Baker, in Journ. Bot. x. (1872) 297, ex parte = texanum.
graminifolium, S. Wats. in Proc. Am. Acad. xiv. (1879) 240 = acrotrichum.
graminifolium, *Zucc. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 288, nomen; *et in Abh. Akad. Muench.* iii. (1843) 238.—Mexic.
Hartwegianum, Hook. Bot Mag. t. 5099 = Hookeri.
Hartwegianum, Zucc. in Abh. Akad. Muench. iv. ii. (1845) 22 = Nolina Hartwegiana
Hookeri, *Lem. Illustr. Hortic.* vi. (1859) *Misc* 24.—Mexic.
Humboldtii, Kunth, in Abh. Berl. Acad. (1842) 32 = Nolina parviflora.
juncreum, Zucc. in Abh. Akad. Muench, iv. ii. (1845) 19 = Nolina Hartwegiana.
lanciforum, Baker, in Journ. Bot. x. (1872) 299 = serratifolium.
Lindheimerianum, Scheele, in Linnaea, xxv. (1852) 202 = Nolina Lindheimeriana.
longifolium, Zucc. in Abh. Akad. Muench. iii. (1843) 224 = Nolina longifolia.
longissimum, *Lem. Illustr. Hortic.* iii. (1856) *Misc.* 91.—Mexic.
pitrairniaefolium, Karw. ex Zucc. in Otto & Dietr. Allg. Gartenz. vi. (1838) 288 = Hechtia glomerata.
plicatile, *Baker, in Journ. Linn. Soc.* xvii. (1880) 240.—Mexic.
quadrangulatum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 250.—Mexic.
serratifolium, Baker, in Journ. Bot. x. (1872) 298 ex parte = glaucophyllum.
serratifolium, *Zucc. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 288, nomen; *et in Abh. Akad. Muench.* iii. (1843) 225.—Mexic.
tenuifolium, Torr. Bot. Mex. Bound. 215 = Nolina Lindheimeriana.
texanum, *Scheele, in Linnaea*, xxiii. (1850) 140.—Texas.
Wheeleri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 240.—Am. bor. occ.

DASYLOMA, DC. Prod. iv. 140 (1830) = **Osmanthe**, Tourn. (Umbell.).
benghalense, DC. l. c. = O. benghalensis.
corticatum, Miq. Fl. Ind. Bat. i. 1. 742 = O. stolonifera.
glaucum, DC. Prod. iv. 140 = O. benghalensis.
japonicum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 59 = O. stolonifera.
javanicum, Miq. Fl. Ind. Bat. i. 1. 741 = O. stolonifera.
lacinistum, Miq. l. c. = Falcaria laciniata.
latifolium, Lindl. in Royle, Illustr. Bot. Himal. 282 = O. stolonifera.
subspinosum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 59 = O. stolonifera.

DASYMALLA, Endl. Nov. Stirp. Dec. 11 (1839) = **Pityrodia**, R. Br. (Verbenac.).
axillaris, Endl. l. c. = P. racemosa.
terminalis, Endl. l. c. 12 = P. racemosa.

DASYNEMA, Schott, in Spreng. Syst. iv. App. 408 (1827) = **Sloanea**, Linn. (Tiliac.).
almifolium, Walp. Rep. i. 852 = S. sinifolia.
cuneifolium, Walp. l. c. = S. cuneifolia.
hirsutum, Schott, in Spreng. Syst. iv. App. 408 = S. monosperma.
lanciflolium, Benth. in Hook. Journ. Bot iv. (1842) 128 = S. lanciflolia.
obtusum, Splitg. in Hoev. & De Vriese Tijdschr. ix. (1849) 96 = S. parviflora.
obtusum, Walp. Ann. i. 110 = S. obtusa.

DASYNEMA:—
ochrocarpa, Mart. ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 70 = Sloanea ochrocarpa.
pubescens, Poepp. & Endl. Nov. Gen. et Sp. iii. 74 = S. pubescens, *Benth.*
pubescens, Schott, in Spreng. Syst. iv. App. 408 = S. pubescens, *Radlk.*
pubescens, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 224.—Bras.
riparia, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 334 = S. riparia.
salicifolium, Endl. ex Walp. Rep. i. 852 = S. monosperma.

DASYOCHLOA, Willd. ex Steud. Nom. ed. II. i. 484 (1840) = **Triodia**, R. Br. (Gramin.).
avenacea, Willd. l. c. = T. avenacea.
pulchella, Willd. l. c. = T. pulchella.

DASYPHYLLUM, H. B. & K. Nov. Gen. et Sp iv. 17. t. 308 (1820) = Chiquitraga, Juss. (Compos.).
argenteum, H. B. & K. l. c. 18.—Ecuador.

DASYPOGON, R. Br. Prod. 268 (1810); et in Flind. Voy. ii. 608. t. 8 (1814). *JUNCACEAE*, Benth. & Hook. f. iii. 665.
DASYPOGONIA, Reichb. Nom. 46 (1841).
bromelifolius, *R. Br. l. c.*—Austral.
glaber, Labarpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 101 = bromelifolius.
Hookeri, *Drumm. in Hook. Lond. Journ. Bot.* ii. (1843) 169.—Austral.
obliquifolius, Lehm. Pl. Preiss. ii. 52 = bromelifolius.

DASYPOGONIA, Reichb. Nom. 46 (1841) = prec.

DASYRANTHUS, Rafin. ex Steud. Nom. ed. II. i. 484 (1840) = **Gnaphalium**, Linn. (Compos.).

DASYS, Lem. in Orb. Dict. iv. 618 (1849) = **Dasus**, Lour. (Inc. sed.).

DASYSPERMUM, Neck. Elem. i. 295 (1790) (= Farrago Umbelliferarum).

DASYSTACHYS, Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 255. t. 35. *LILIACEAE*, Benth. & Hook. f. iii. 789.
campanulata, *Baker, l. c.* 256.—Afr. trop.
colubrina, *Baker, l. c.*—Afr. trop.
falcata, *Baker, l. c.*—Afr. trop.
pleiostachya, *Baker, l. c.* 255.—Afr. trop.

DASYSTACHYS, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 25 = Chamaedorea, Willd. (Palm.).
Deckeriana, Oerst. l. c. 26 = C. Deckeriana.

DASYSTEMON, DC. Prod. iii. 382 (1828) = **Crassula**, Linn.
calycinum, DC. l. c. = C. calycina.

DASYSTEPHANA, Adans. Fam. ii. 502 (1763); Borckh. in Roem. Archiv, i. i. 25 (1796) = **Gentiana**, Tourn.
adscendens, Borckh. l. c. 26 = G. decumbens.
algida, Borckh. l. c. = G. frigida.
asclepiadea, Borckh. l. c. = G. asclepiadea.
auriculata, Borckh. l. c. = G. auriculata.
glauca, Borckh. l. c. = G. glauca.
punctata, Borckh. l. c. = G. punctata.
purpurea, Steud. Nom. ed. I. 362; ed. II. i. 484, sphalm. = G. punctata.
trifora, Borckh. in Roem. Archiv, i. i. 26 = G. trifora.

DASYSTOMA, Rafin. ex Endl. Gen. 691, in syn. (1839) = DASISTOMA, Rafin. = **Gerardia**, Linn. (Scrophul.).

DATISCA, Linn. Diss. Dass. 21 (1747); Amoen. Acad. i. 415 (1749). *DATISCEAE*, Benth. & Hook. f. i. 844.
TRICERASTES, Presl, Rel. Haenk. ii. 88. t. 64 (1836).
cannabina, *Linn. Sp. Pl.* 1037.—Orient; Reg. Himal.
glabra, Stokes, Bot. Mat. Med. iv. 568 = cannabina.
glomerata, *Baill. Hist. Pl.* ii. 407.—Calif.
hirta, Linn. Sp. Pl. 1037 = Rhus typhina.
nepalensis, D. Don, Prod. Fl. Nep. 303 = cannabina.

DATURA, Linn. Syst. ed. I (1735). *SOLANACEAE*, Benth. & Hook. f. ii. 901.
APEMON, Rafin. Fl. Tellur. ii. 11 (1836).
BRUGMANSIA, Pers. Syn. i. 216 (1805).
CERATOCAULOS, Reichb. Handb. 201 (1837).
STRAMONIUM, Hall. Enum. Stirp. Helv. ii. 489 (1742).
acaulis, Pav. ex Dun. in DC. Prod. xiii. i. 481 = Jaborosa integrifolia.
aegyptiaca, Vent. Pl. Alegrypt. 26 = fastuosa.
alba, F. Muell. Fragm. vi. 144 = Leichhardtii.
alba, Nees, in Trans. Linn. Soc. xvii. (1834) 78 = fastuosa.
albido-lutea, Lem. Jard. Fleur. iv. (1854) Misc. 16 = Dyssochroma albidoflavum.
arborea, *Linn. Sp. Pl.* 179.—Am. austr.
arborea, Mart. in Flora, xxiv. (1841) ii. Beibl. 111 = suaveolens.
Bertolonii, Parl. ex Guss. Fl. Sic. Syn. 207 = ferox.
Bojeri, Delile, Ind. Sem. Hort. Monsp. (1836) 23; et in Ann. Sc. Nat. Sér. II. vii. (1837) 326 = fastuosa.
candida, Pasq. Cat. Ort. Bot. Nap. 36 = arborea.
capensis, Hort. ex Bernh. in Tromms. N. Journ. Pharm xxvi. (1833) 151 = D. Stramonium.
ceratocaula, *Zucc. Hort. Schoenb.* iii. 48.—Am. trop. chlorantha, *Hook. Bot. Mag.* t. 5128.—Hab. ?
cornigera, *Hook. l. c.* t. 4952.—Mexic.
discolor, *Bernh. in Tromms. N. Journ. Pharm.* xxvi. (1833) 149.—*Cf. Linnaea*, viii. (1833) *Litt.* 138.—Mexic.
dubia, *Bianca, ex Tod. Ind. Sem. Hort. Panorm.* (1858) 26.—Hab. ?
dubia, Kich. ex Pers. Syn. i. 216 = fastuosa.
erinacea, *Vell. Fl. Flum.* 70; ii. t. 46.—Bras.
eximia, Lem. Illustr. Hortic. vi. (1859) Misc. 93 = Dyssochroma eximia.
fastuosa, *Linn. Syst. ed.* X. 932.—Geront. trop.
ferox, *Linn. Dist. Dem. Pl. in Amoen. Acad.* iii. 403. —China.
ferox, Nees, in Trans. Linn. Soc. xvii. (1834) 75 = D. Stramonium.
flortbunda, *Feast. Mag. Bot.* ix. (1842) 3 = Am. austr.
fruticosa, Hornem. Hort. Hafn. i. 212 = D. Metel.
Gardneri, Hook. Bot. Mag. sub t. 4252 = suaveolens.
guayaquilensis, H. B. & K. Nov. Gen. et Sp. iii. 8 = D. Metel.
humilis, Desf. Cat. Hort. Par. ed. III. 296 = fastuosa.
Hummata, Bernh. in Tromms. N. Journ. Pharm. xxvi. (1833) 155.—Cf. Linnaea, viii. (1833) Litt. 141 = fastuosa.
hybrida, Tenore, App. iv. Prod. Fl. Neap. 34 = Inermis ?
inermis, *Yacq. Hort. Vindob.* iii. 44. t. 82.—Abyss.
inoxia, Mill. Gard. Dict. ed. VIII. n. 5 = fastuosa.
laevis, Bertol. Fl. Ital. iv. 780 = ferox.
laevis, Linn. f. Suppl. 146 = Inermis.
laevis, Schkuhr, Handb. i. t. 140 = fastuosa.
Leichhardtii, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 20, in obs. nomen.—Austral.
loricata, Sieber, ex Bernh. in Tromms. N. Journ. Pharm. xxvi. (1833) 151 = D. Stramonium.
lurida, Salisb. Prod. 121 = D. Stramonium.
macrocaulis, Roth, Beitr. 159 = ceratocaula.
maxima, Moç. & Sesse, ex Dun. in DC. Prod. xiii. i. 536 = Solandra guttata.
Metel, *Linn. Sp. Pl.* 179.—Am. trop. (Cosmop. valg.).
Metel, Mill. Gard. Dict. ed. VIII. n. 8 = fastuosa.
Metel, Moç. & Sesse, ex Dun. in DC. Prod. xiii. i. 544 = meteloides.
meteloides, *DC. ex Dun. l. c.*—Am. bor. occ.
microcarpa, *Godr. in Mém. Acad. Stanisl.* Sér. IV. v. 1872 (1873) 209.—Hab. ?
muricata, Hort. Berol. ex Godr. l. c. 199 = praecox.
muricata, Link, Enum. Hort. Berol. i. 177 = fastuosa.
nigra, *Hassk. Cat. Hort. Bog. Alt.* 142.—Malaya.
Nilsuammatu, Dun. in DC. Prod. xiii. i. 542 = fastuosa.
parviflora, Salisb. Prod. 121 = D. Stramonium.
praecox, *Godr. in Mém. Acad. Stanisl.* Sér. IV. v. 1872 (1873) 199.—Hab. ?
Pseudostramonium, Sieber, ex Bernh. in Tromms. N. Journ. Pharm. xxvi. (1833) 151 = D. Stramonium.
quercifolia, Godr. l. c. = praecox.
quercifolia, *H. B. & K. Nov. Gen. et Sp.* iii 7.—Mexic.
sanguinea, *Ruiz & Pav. Fl. Per.* ii. 15.—Am. austr.
sarmentosa, Lam. Illustr. ii. 9 = Solandra hirsuta.
scandens, *Vell. Fl. Flum.* 69; ii. t. 46.—Bras.
speciosa, Salisb. Prod. 121 = arborea.

DATURA:—
Stramonium, *Linn. Sp. Pl.* 179. — Cosmop. trop. et temp.
Stramonium, Thunb. Fl. Jap. 91 = fastuosa.
Stramonium, Wall. Cat. Suppl. n. 278 = trapezia.
suaveolens, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 227.—Mexic.
Tatula, Linn. Sp. Pl. ed. II. 256 = D. Stramonium.
Thomasii, Torr. Pacif. Rail. Rep. v. 362 = discolor.
timorensis, Zipp. ex Span. in Linnaea, xv. (1841) 337 = D. Metel.
trapezia, *Nees, in Trans. Linn. Soc.* xvii. (1834) 74.—Ind. or.
Wagmanni, Steud. Nom. ed. II. i. 484 = fastuosa.
Wallichii, Dun. in DC. Prod. xiii. i. 539 = D. Stramonium.
Wrightii, Torr. ex Regel, Gartenfl. (1859) t. 360 = meteloides.

DAUBENTONIA, DC. Mém. Lég. 265 (1825) = **Sesbania**, Pers. (Legumin.).
longifolia, DC. Prod ii. 267 = S. Cavanillesii.
punicea, DC. l. c. = S. punices.
Thurberi, A. Gray, in Mem. Am. Acad. N. S. v. (1855) 313 = S. Thurberi.
Tripetiana Poit. in Ann. Soc. Hortic. Par. xxxi. (1842) 120 = seq.
Tripetii, Poit. l. c. xxvii. (1840) 136.—Reg. Argent.

DAUBENYA, Lindl. Bot Reg. t. 1813 (1835). *LILIACEAE*, Benth. & Hook. f. iii. 806.
aurea, *Lindl. l. c.*—Afr. austr.
coccinea, *Harv. ex Baker, in Journ. Linn. Soc.* xi. (1871) 295.—Afr. austr.
fulva, *Lindl. Bot. Reg.* (1839) t. 53.—Afr. austr.

DAUCALIS, Pomel, Nouv. Mat. Fl. Atl. 145 (1874 = **Caucalis**, Tourn. (Umbell.).
leptophylla, Pomel, l. c. 149 = C. leptophylla.
mauritanica, Pomel, l. c. = C. mauritanica.

DAUCERIA, Dennst Schlüssl. Hort. Malab. 31 (1818) = **Embelia**, Burm. f. (Myrsin.).
acuta, Dennst. l. c. = E. robusta.
obtusa, Dennst. l. c. = E. robusta.

DAUCOSMA, Engelm. & A. Gray, ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 210 = **Discopleura**, DC. (Umbell.).
laciniata, Engelm. & A. Gray, l. c. 211 = Disc. laciniata.

DAUCUS, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 926.
AMMODAUCUS, Coss. & Dur. in Bull. Soc. Bot. Fr. vi. (1859) 893.
BALLIMORA, Rafin. New Fl. Am. iv. 28 (1836).
CTENODAUCUS, Pomel, Nouv. Mat. Fl. Atl. 145 (1874).
DURIAEA, Bolss. & Reut. Diagn. Pl. Nov. Hisp. 14 (1842).
ORLAYA, Hoffm. Gen. Umb. 58 (1814).
PELTACTILA, Rafin. New Fl. Am. iv. 28 (1836).
PLATYDOON, Reichb. Nom. 226 (1841).
PLATYDAUCON, Reichb. l. c. 205 (1841).
PLATYSPERMUM, Hoffm. Gen. Umb. 64 (1814).
STAPLIMUS, Rafin. New Fl. Am. iv. 28 (1836).
TIRACTA, Rafin. l. c. 23 (1836).
abyssinicus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 231.—Abyss.
abyssinicus, C. A. Mey. in Ind. Sem. Hort. Petrop. ix. 56 = D. Carota.
agrestis, Rafin. New Fl. Am. iv. 27; Jord. ex Nym. Consp. 279 = D. Carota.
alatus, *Parr. Voy. Bark.* ii. 135 ; *Encyc. Suppl.* ii. 118. —Algeria.
Allioni, Link, Handb. i. 344 = D. Carota.
anisodorus, Blanco, Fl. Filip. ed. II. 150 = Carum copticum.
Anthriscus, Baill. Hist. Pl. vii. 88 = Caucalis Anthriscus.
arvensis, Pasq. ex Boiss. Fl. Orient. ii. 1075 = setulosus.
Ariedia, Cesats, Inst. ii. 116 = Artedia squamata.
asturiorum, Barnad. ex Colmeiro, Fl. Comp. Madr. 220 = polygamus.

DAUCUS:—
atropatanus, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 252, in textu = pevicus.
aureus, *Desf. Fl. Atlant.* i. 242. t. 61.—Reg. Mediterr.
australis, Guss. Fl. Sic. Prod. Suppl. i. 74 = rupestris.
australis, *Poepp. ex DC. Prod.* iv. 214.—Chili.
bactrianus, *Bunge, Enl. Lehm. Bot.* 136.—Turkestan.
besarabicus, *DC. Prod.* iv. 210.—Ross. austr.
bicolor, Sibth. & Sm. Fl. Graec. iii. 64. t. 270 = Broteri.
Blanchei, *Reut. Ind. Sem. Hort. Genev.* (1867); *Boiss. Fl. Orient.* ii. 1077.—Syria.
Bocconi, *Guss. Fl. Sic. Prod.* i. 329.—Calabria.
brachiatus, *Sieber, ex DC. Prod.* iv. 214.—Austral.
brachiatus, Torr. in Pacif. Rail. Rep. iv. 98 = Cancalis microcarpa.
brachyloba, Boiss. Voy. Bot. Espagne, ii. 258. t. 68 = setifolius.
brevicaulis, Rafin. New Fl. Am. iv. 25 = D. Carota ?
brevifolius, Rafin. l. c. 26 = pusillus.
Broteri, *Tenore, Syll. Fil.* 54.—As. Min. ; Syria.
carnosus, Moench, Meth. 93 = mauritanicus.
Carota, *Linn. Sp. Pl.* 242.—Europ. ; Orient.
Carota, *Tenore, Syll. App.* iv. 10 (sp. dub.).—Italia.
copticus, Lam. Encyc. i. 635 = Carum copticum.
creticus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Graecia ; As. Min.
crinitus, *Desf. Fl. Atlant.* i. 242. t. 62.—Maurit. ; Lusitan.
cuminoides, *Lam. ex Poir. Encyc. Suppl.* ii. 118.—Hab. ?
dentatus, *Bertol. Fl. Ital.* iii. 168.—Gallia ; Hispan.
depressus, Sprenq. Syst. iv. Cur. Post. 110 = Cortia Lindleyi.
divaricatus, Walt. Fl. Carol. 114 = Apium divaricatum.
Durieua, *Lange, in Willk. & Lange, Prod. Fl. Hisp* iii. 23.—Hispan.
esculentus, Salisb. Prod. 102 = D. Carota.
exiguus, Steud. Nom. ed. II. i. 486 = D. Carota.
fortidus, *Rafin. Fl. Ludov.* (sp. dub.).—Louisiana.
foliosus, *Guss. Suppl. Fl. Sic. Prod.* i. 73.—Sicilia.
gaditanus, *Boiss. & Reut. Diagn.* Ser. II. ii. 96.—Hispan.
Gaillardoti, Boiss. Fl. Orient. ii. 1074 = littoralis.
gibbosus, Bertol. Fl. Ital. iii. 160 = foliosus.
Gingidium, Georgi, Beschr. Russ. Reich. Nachtr. 266 = D. Carota.
Gingidium, Linn. Herb. ex DC. Prod. iv. 118 = Ammi visnaga.
Gingidium, *Linn. Sp. Pl.* 242.—Europ. ; Afr. bor.
glaber, Opiz, ex Celak. Prod. Fl. Boehm. iii. 582 = D Carota.
glaberrimus, *Desf. Fl. Atlant.* i. 244.—Afr. bor.
glochidiatus, Fisch, Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 11 = brachiatus.
Golickeanus, *Regel & Schmalh. in Act. Hort. Petrop* v. (1877) 599.—Turkestan.
Gouani, Nym. Syll. 148 = polygamus.
gracilis, *Steinh. in Ann. Sc. Nat.* Sér. II. ix. (1838) 208.—Algeria.
grandiflorus, *Desf. Fl. Atlant.* i. 240. t. 59.—Algeria
grandiflorus, *Scop. Fl. Carn.* ed. II. i. 189.—Europ ; Taur.
guyanensis, Lam. Encyc. i. 634 = D. Gingidium.
guttatus, Sibth. & Sm. Fl. Graec. iii. 63. t. 269 = setulosus.
guttatus, Sieber, ex Nym. Consp. 279 = D. Gingidium.
halophilus, Brot. Phyt. Lusit. t. 168 = D. Gingidium.
heterocarpus, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 60 = muricatus.
heterophyllus, Rafin. New Fl. Am. iv. 26 = D. Carota ?
hirsutus, DC. Prod. iv. 213 = setulosus.
hispanicus, Gouan, Illustr. 9 = D. Gingidium.
hispidifolius, *Clos, in C. Gay, Fl. Chil.* iii. 185.—Chili.
hispidus, Desf. Fl. Atlant. i. 243. t. 63 = D. Gingidium.
hispidus, Gilib. Fl. Lituan. ii. 10 = D. Carota.
hispidus, Mill. Gard. Dict. ed. VIII. n. 4 = polygamus.
insularis, Sibth. & Sm. Fl. Graec. iii. 65. t. 271 = creticus.
jolanis, *Pomel, Nouv. Mat. Fl. Atl.* 146.—Algeria.
Iorulo, Salisb. Prod. 102 = Ammi Visnaga.
Lagoecia, [Weber in] Wigg. Prim. Fl. Holsat. 28 = Caucalis daucoides.

DAUCUS:—
asperitioides, *DC. Prod.* iv. 210.—Mauritan.
latifolia, Baill. Hist. Pl. vii. 193 = Caucalis latifolia.
leptocarpus, *Hochst. in Lorent, Wanderungen*, 337.—Syria; Mesopot.
leptophyllus, Scop. Fl. Carn. ed. II. i. 190 = Caucalis leptophylla.
lewis, Rafin. New Fl. Am. iv. 25 = D. Carota ?
littoralis, *Sibth. & Sm. Fl. Graec.* iii. 65. t. 272.—Syria.
lopadusanus, *Tin. Pl. Ker. Acil.* 85.—Sicilia.
lucidus, Linn. f. Suppl. 179 = D. Gingidium.
marcidus, Timb. in Bull. Soc. Bot. Fr. xiii. (1866) p. clv = D. Carota.
maritimus, Geertn. Fruct. i. 20. t. 4 = pumilus.
maritimus, *Lam. Encyc.* i. 634.—Europ. occ.
maritimus, With. Bot. Arr. ed. III. ii. 290 = D. Carota.
mauritanicus, Linn. Sp. Pl. ed. II. 348; All. Fl. Pedem. ii. 81. t. 61 = D. Carota.
mauritanicus, Salzm. ex DC. Prod. iv. 212 = D. Gingidium.
maximus, *Desf. Fl. Atlant.* i. 241.—Reg. Mediterr.; Oriens.
maydinus, Brot. Phyt. Lusit. i. 82. t. 36 = crinitus.
melananthus, Steud. ex Hochst. in Flora, xxvii. (1844) 19 = Caucalis melanantha.
mexicus, Pers. Syn. i. 305 = Ammi anethifolium.
Michelii, Caruel, Prod. Fl. Tosc. 292 = Broteri.
micranthus, *Pomel, Nouv. Mat. Fl. Atl.* 146.—Algeria.
microphyllus, Presl, ex DC. Prod iv. 213 = pusillus.
microsciadius, Boiss. Diagn. Ser. I. x. 47 = setulosus.
montanus, *Humb. & Bonpl. ex Schult. Syst.* vi. 482.—Venezuela.
montanus, Schmidt, ex Nym. Consp. 279 = D. Carota.
montevidensis, *Link, ex Spreng. Syst. iv. Cur. Post.* 119.—Reg. Aequin.
muricatus, Linn. Herb. ex Guss. Fl. Sic. Syn. i. 336 = Broteri.
muricatus, *Linn. Sp. Pl.* ed. II. 349.—Reg. Mediterr.
nebrodensis, Strobl, in Oestr. Bot. Zeitschr. xxxiv. (1884) 223 = D. Gingidium ?
neglectus, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838). (reimpr. 90).—Ins. Madera.
nitidus, Guep. in Guss. Enum. Pl. Vasc. Inar. 147. t. 5 = D. Gingidium.
nudicaulis, Rafin. New Fl. Am. iv. 25 = D. Carota ?
officinalis, Gueldenst. ex Ledeb. Fl. Ross. ii. 339 = D. Carota.
Paralias, *Pomel, Nouv. Mat. Fl. Atl.* 145.—Algeria.
parviflorus, *Desf. Fl. Atlant.* i. 241. t. 60.—Afr. bor. occ.
parviflorus, [Willk. in] Losc. & Pardo, Sert. Inconf. ed. II. 156 = aureus.
pectinaceus, *DC. Prod.* iv. 212.—Hab. ?
persicus, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 49.—Persia.
platycarpos, Scop. Fl. Carn. ed. II. i. 190 = Caucalis daucoides.
polygamus, *Govan, Illustr.* 9.—Hispan.; Sicil.
polygamus, Jacq. ex Nym. Consp. 279 = D. Carota.
prolifer, J. & C. Presl, Delic. Prag. 180 = Laserpitium hispidum.
pubescens, Koch, Gen. Umbell. 77 = littoralis.
pulcherrimus, *Koch, ex DC. Prod.* iv. 210.—Reg. Cauc.; Persia.
pumilus, *Bail, in Journ. Linn. Soc.* xvi. (1878) 477.—Reg. Mediterr.
pusillus, *Wickst. Fl. Bor. Am.* i. 164.—Am. bor.
rigidus, *Hort. Par. ex DC. Prod.* iv. 215.—Hab. ?
Rouyi, Baill. Hist. Pl. vii. 89, 198 = Caucalis daucoides.
rupestris, *Guss. Fl. Sic. Syn.* i. 335.—Ins. Sicil.
scaber, Nutt. ex Torr. & Gray, Fl. N. Am. i. 636 = pusillus.
scabriusculus, Boiss. & Noë, Diagn. Ser. II. ii. 95 = leptocarpus.
scabrosus, Bert. ex DC. Prod. iv. 211 = setulosus.
scariosus, Rafin. New Fl. Am. iv. 25 = D. Carota ?
sciadophyllus, Rafin. l. c. 24 = D. Carota.
serotinus, *Pomel, Nouv. Mat. Fl. Atl.* 147.—Algeria.
serratus, Moris, in Mem. Acc. Tor. II. ii. (1840) 304 = maritimus.
setifolius, *Desf. Fl. Atlant.* i. 244. t. 68.—Afr. bor.
setulosus, *Guss. ex DC. Prod.* iv. 211.—Ital.; Graecia.
siculus, Tineo, Pugill. Sic. i. 6 = Bocconi.
speciosus, Cusset, in Linnaea, xi. (1837) 623 = setulosus.
strigosus, Rafin. New Fl. Am. iv. 27 = D. Carota ?

DAUCUS:—
sylvestris, Mill. Gard. Dict. ed. VIII. n. 1 = D. Carota.
torilioides, DC. Prod. iv. 214 = montanus.
verticillatus, Hornem. Hort. Hafn. i. 272 = setifolius.
verticillatus, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 477 = crinitus.
Visnaga, Linn. Sp. Pl. 242 = Ammi Visnaga.
vulgaris, Lam. Fl. Fr. iii. 430 = D. Gingidium.
vulgaris, Neck. Delic. Gallo-Belg. i. 139 = D. Carota.

DAUTHONIA, Link. Handb. i. 45 (1829) sphalm. = **Danthonia**, DC. (Gramin.).

DAUVENTONIA, Reichb. in Mossal. Deutsch. Fl. ed. II. p. xlvii (1837), sphalm. = DAUBENTONIA, DC. = **Scobania**, Pers. (Legumin.).

DAVIDIA, Baill. Adansonia, x. (1871) 114. *HAMAMELIDEAE* ?
involucrata, *Baill. l. c.* 115.—Tibet. or.
tibetana, *David, in Nouv. Arch. Mus. Par.* v. (1882) 56.—Tibet. or.

DAVIDSONIA, F. Muell. Fragm. vi. 4 (1867). *SAXIFRAGACEA*.
abnormis, *F. Muell. l. c.*—Austral.
pungens, Hort. ex Gard. Chron. (1876) i. 608 = prunc. ?

DAVIESIA, Poir. Encyc. Suppl. v. 480 (1817) = **Borya**, Labill. (Liliac.).
lucens, Poir. l. c. = B. nitida.

DAVIESIA, Sm. in Trans. Linn. Soc. iv. (1798) 220. *LEGUMINOSAE*, Benth. & Hook. f. i. 469.
abscurens, F. Muell. Fragm. ii. 106 = Latrobea pungens.
acanthoclona, *F. Muell. l. c.* x. 82.—Austral.
acanthoclada, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 262 = Hovea acanthoclada.
aciculata, *Sm. in Ken. & Sims, Ann. Bot.* i. (1805) 506.—Austral.
adnata, F. Muell. Fragm. iii. 105 = striata.
alata, *Sm. in Trans. Linn. Soc.* ix. (1808) 259.—Austral.
alternifolia, *Endl. in Ann. Wien. Mus.* ii. (1838) 199. —Austral.
anceps, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 266.—Austral.
angulata, *Benth. in Lindl. Swan River App.* 57.—Austral.
aphylla, *F. Muell. in Benth. Fl. Austral.* ii. 88. —Austral.
arborea, *F. Muell. & Scort. in Proc. Linn. Soc. N. S. W.* vii. (1883) 221.—Austral.
arthropoda, *F. Muell. Fragm.* viii. 22.—Austral.
Benthamii, Meisn. in Lehm. Pl. Preiss. i. 48 = incrassata.
brachyphylla, Meisn. l. c. 49 = incrassata.
brevifolia, *Lindl. in Mitch. Three Exped. Austral.* ii. 301.—Austral.
buxifolia, *Benth. Fl. Austral.* ii. 75.—Austral.
calystegia, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 264 = crenulata.
cardiophylla, *F. Muell. Fragm.* ii. 105.—Austral.
chordophylla, *Meisn. in Lehm. Pl. Preiss.* i. 48.—Austral.
collateralis, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 75 = genistifolia.
colletioides, *Meisn. in Lehm. Pl. Preiss.* i. 48.—Austral.
colubrinoides, Walp. Rep. i. 870 = colletioides.
concinna, *R. Br. ex Benth. Fl. Austral.* ii. 75.—Austral.
condensata, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 265 = Dillwynia pungens.
cordata, *Sm. in Trans. Linn. Soc.* ix. (1808) 259.—Austral.
corymbosa, *Sm. in Ken. & Sims, Ann. Bot.* i. (1805) 507.—Austral.
crenulata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 265.—Austral.
daphnoides, *Meisn. in Lehm. Pl. Preiss.* i. 54.—Austral.
decurrens, Meisn. l. c. 52 = pectinata.
denudata, Vent. Choix, 6 = Viminaria denudata.
divaricata, *Benth. in Enum. Pl. Hueg.* 31.—Austral.
Drummondii, *Meisn. in Lehm. Pl. Preiss.* i. 53.—Austral.
egena, F. Muell. in Trans. Vict. Inst. i. (1855) 118 = Templetonia egena.

DAVIESIA:—
elongata, *Benth. Fl. Austral.* ii. 74.—Austral.
Epiphyllum, *Meisn. in Bot. Zeit.* xiii. (1855) 27.—Austral.
ericoides, Pers. Syn. i. 454 = Aotus villosa.
euphorbioides, *Benth. Fl. Austral.* ii. 86.—Austral.
filipes, *Benth. in Mitch. Journ. Trop. Austral.* 863. —Austral.
flexuosa, *Benth. in Enum. Pl. Hueg.* 32.—Austral.
genistifolia, *A. Cunn. ex Benth. in Ann. Wien. Mus.* ii. (1838) 75.—Austral.
genistoides, Lodd. Bot. Cab. t. 1552 = ulicina.
glauca, Lodd. l. c. t. 48 = corymbosa.
hakeoides, *Meisn. in Lehm. Pl. Preiss.* i. 47.—Austral.
horrida, *Meisn. l. c.* 54.—Austral.
humifusa, Sieber, ex DC. Prod. ii. 103 = Oxylobium scandens.
incrassata, Meisn. in Lehm. Pl. Preiss. i. 49 = microphylla.
incrassata, *Sm. in Trans. Linn. Soc.* ix. (1808) 255.—Austral.
juncea, Pers. Syn. i. 454 = Viminaria denudata.
junces, *Sm. in Trans. Linn. Soc.* ix. (1808) 260.—Austral.
lanceolata, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 263 = mollis.
latifolia, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iii. 20.—Austral.
latipes, *F. Muell. in Linnaea*, xxv. (1852) 390.—Austral.
laurifolia, Link, Enum. Hort. Berol. i. 408 = corymbosa ?
leptophylla, A. Cunn. ex G. Don, Gen. Syst. ii. 125 = corymbosa.
linearis, Lodd. Bot. Cab. t. 1615 = corymbosa.
longifolia, *Benth. in Lindl. Swan River App.* 61.—Austral.
macrophylla, Endl. Nov. Stirp. Decad. 18 = corymbosa.
microphylla, *Benth. Fl. Austral.* ii. 86.—Austral.
mimosoides, [K. Br. in] Ait. Hort. Kew. ed. II. iii. 20 = corymbosa.
mollis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 263.—Austral.
nematophylla, *F. Muell. ex Benth. Fl. Austral.* ii. 78. —Austral.
nudiflora, *Meisn. in Lehm. Pl. Preiss.* i. 53.—Austral.
obovata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 261.—Austral.
obtusifolia, *F. Muell. Fragm.* ii. 104.—Austral.
oppositifolia, *Endl. in Ann. Wien. Mus.* ii. (1838) 199.—Austral.
ovata, *Benth. Fl. Austral.* ii. 72.—Austral.
pachylima, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 263.—Austral.
pachyphylla, *F. Muell. Fragm.* iv. 15.—Austral.
paniculata, *Benth. in Enum. Pl. Hueg.* 31.—Austral.
parifolia, F. Muell. Fragm. iv. 16 = crenulata.
pectinata, *Lindl. in Mitch. Three Exped. Austral.* ii. 151.—Austral.
pedunculata, *Benth. in Lindl. Swan River App.* 60.—Austral.
phyxodes, A. Cunn. ex G. Don, Gen. Syst. ii. 125 = incrassata.
polyphylla, *Benth. in Lindl. Swan River App.* 58.—Austral.
Preissii, *Meisn. in Lehm. Pl. Preiss.* i. 50.—Austral.
prionodes, Meisn. l. c. 52 = pectinata.
pubigera, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 75 = umbellulata.
pungens, R. Cunn. ex Benth. l. c. = acicularis.
quadrilatera, *Benth. in Lindl. Swan River App.* 59. —Austral.
racemulosa, DC. Prod. ii. 114 = umbellulata.
ramulosa, *Benth. in Lindl. Swan River App.* 56.—Austral.
reclinata, *A. Cunn. ex Benth. Fl. Austral.* ii. 77.—Austral.
reticulata, Sm. in Trans. Linn. Soc. ix. (1808) 256 = Pultenaea reticulata.
reversifolia, *F. Muell. Fragm.* i. 145.—Austral.
rhombifolia, *Meisn. in Lehm. Pl. Preiss.* i. 56.—Austral.
rusciifolia, A. Cunn. ex Hook. Bot. Mag. t. 3196 = ulicina.
spinosissima, *Meisn. in Lehm. Pl. Preiss.* i. 51. —Austral.
squarrosa, *Sm. in Ken. & Sims, Ann. Bot.* i. (1805) 507.—Austral.

DAVIESIA:—
striata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 264.—Austral.
teretifolia, *R. Br. ex Benth. Fl. Austral.* ii. 82.—Austral.
ternata, *Endl. in Ann. Wien. Mus.* ii. (1838) 199.—Austral.
trigonophylla, *Meisen. in Lehm. Pl. Preiss.* ii. 213.—Austral.
ulicifolia, Andr. Bot. Rep. t. 804 = ulicina.
ulicina, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 506. —Austral.
umbellata, Labill. Nov. Holl. Pl. i. 107. t. 137 = ulicina.
umbellata, Sieber, ex DC. Prod. ii. 108 = Oxylobium scandens.
umbellulata, DC. Prod. ii. 114 = ulicina.
umbellulata, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 507.—Austral.
villifera, A. Cunn. ex Benth. in Ann. Wien. Mus ii. (1838) 76 = squarrosa.
virgata, A. Cunn. ex Hook. Bot. Mag. t. 3196 = corymbosa.
Wyattiana, *F. M. Bailey, in Proc. Linn. Soc. N. S. W.* vi. (1881) 145.—Austral.

DAVILLA, Vand. in Roem. Script. 115 (1796). *DILLENIACEÆ*, Benth. & Hook. f. i. 12.
Hibbonia, Vell. Fl. Flum. v. t. 116 (1827).
adenophylla, Mart. ex Eichl. in Mart. Fl. Bras. xiii. i. 98 = flexuosa.
angustifolia, *St. Hil. Fl. Bras. Mer.* i. 19.—Bras.
asperrima, Spligt. in Hoev. & De Vriese, Tijdschr. ix. (1842) 95 = Knuthii.
brasiliana, Billb. ex Beurl. in Vet. Acad. Handl. Stockh. 1854 (1856) 146 = Knuthii.
brasiliana, DC. Syst. i. 405 = rugosa.
castanæfolia, St. Hil. Fl. Bras. Mer. i. 17 = elliptica.
ciliata, A. Rich. Ess. Fl. Cub. 19 = rugosa.
cuspidiuata, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. i. 101.—Bras.
densiflora, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 18.—N. Granat.
elliptica, *St. Hil. Fl. Bras. Mer.* i. 17.—Bras.
flexuosa, *St. Hil. l. c.* 17. t. 2.—Bras.
formosa, Spligt. in Nederl. Kruidk. Arch. i. (1848) 236 = Tetracera surinamensis.
glabrata, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. i. 97. —Bras.
glabrata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 548.—Bras.
Glaziovii, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 178.—Bras.
grandiflora, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 131.—Bras.
grandifolia, *Moric. ex Eichl. in Mart. Fl. Bras.* xiii. i. 99.—Bras.
itaparicensis, Casar. Nov. Stirp. Bras. Dec. 19 = rugosa.
Knuthii, *St. Hil. Pl. Us. Bras.* sub t. 22. p. 6.—Panama.
lacunosa, *Mart. in Flora*, xxi. (1838) ii. *Beibl.* 49.—Bras.
latifolia, *Casar. Nov. Stirp. Bras. Dec.* 19.—Bras.
lucida, Pœpl. Rel. Haenk. ii. 73 = Knuthii.
macrocarpa, *Eichl. in Mart. Fl. Bras.* xiii. i. 96.—Bras.
macroclada, *Moric. ex Eichl. l. c.* 98.—Bras.
macrophylla, St. Hil. Fl. Bras. Mer. i. 18 = multiflora.
Martii, *Eichl. in Mart. Fl. Bras.* xiii. i. 102.—Bras.
multiflora, *St. Hil. Pl. Us. Bras.* sub t. 22. p. 5.—Bras.
multiflora, Seem. Bot. Voy. Herald, 78 = Sagraeana.
ovata, Pœpl. Rel. Haenk. ii. 78 = Knuthii.
pedicellaris, *Benth. in Hook. Kew Journ.* iii. (1851) 161.—Bras.
planibasata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 548.—Bras.
pilosa, Miq. in Linnæa, xix. (1847) 184 = rugosa.
Radula, Mart. in Flora, xxi. (1838) ii. Beibl. 64 = Tetracera Radula.
rugosa, *Poir. Encyc. Suppl.* ii. 457.—Bras.
Sagraeana, *A. Rich. Ess. Fl. Cub.* 17.—Cuba.
Sellowiana, Schlecht. in Linnæa, viii. (1833) 178 = angustifolia.
surinamensis, Miq. l. c. xviii. (1844) 611 = rugosa.
tenuis, *Eichl. in Mart. Fl. Bras.* xiii. i. 107.—Bras.
tintinnabulata, *Schlecht. in Linnæa*, viii. (1833) 178. —Bras.
vaginata, *Eichl. in Mart. Fl. Bras.* xiii. i. 99.—Bras.
villosa, *Eichl. l. c.* 108.—Bras.
wrouisefolia, *Baill. Adansonia*, vi. (1865–66) 272.—Guiana.

DAVVA, DC. Prod. iii. 105 (1828) = **Meriania**, Sw. (Melast.).
adscendens, Griseb. Fl. Brit. W. Ind. 265 = Adelobotrys adscendens.
calophylla, Cham. in Linnæa, ix. (1834) 372 = M. calophylla.
calyptrata, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 136 = M. calyptrata.
ciliata, Griseb. Fl. Brit. W. Ind. 265 = Octopleura ciliata.
ciliata, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 137 = Adelobotrys ciliata.
Clausseni, Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 147 = M. Clausenii.
crassiramis, Naud. l. c. xviii. (1852) 138 = Centronia crassiramis.
excelsa, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 100 = M. Clausenii.
glabra, DC. Prod. iii. 105 = M. glabra.
guianensis, DC. l. c. = Adelobotrys adscendens.
longipes, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 66 = M. longipes.
macrophylla, Benth. Pl. Hartw. 76 = M. nascrophylla.
paniculata, DC. Prod. iii. 105 = M. paniculata.
peruviana, DC. l. c. = Adelobotrys adscendens.
scandens, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 137 = Adelobotrys scandens.
sclerophylla, Naud. l. c. 138 = M. sclerophylla.
sertulata, Rich. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 66 = M. glabra.

DAVVA, Mog. & Sesse, ex DC. Prod. i. 525, 526 (1824) = **Saurauja**, Willd. (Ternstroem.).
serrata, Mog. & Sesse, l. c. 526 = S. serrata.
villosa, Mog. & Sesse, l. c. 525 = S. villosa.

DAVYNA, Monier, ex Mill. Fig. Pl. t. 118 (1756) = **Ayenia**, Linn. (Stercul.).
inermis, Medic. Malv. 123 = A. pusilla.
pusilla, Mill. Gard. Dict. ed. VIII = A. pusilla.

DAVENIA, Michx. ex Jasb. & Spach, in Ann. Sc. Nat. Sér. III. vi. (1846) 137 = **Bleberteinia**, Steph. (Geran.).

DAVENIA, Mill. Gard. Dict. ed. VII (1759) = **Ayenia**, Linn. (Stercul.).

DAZUS, Juss. in Dict. Sc. Nat. xxvii. 201 (1823) = **Dasus**, Lour. (Inc. sed.).

DEASTELLA, Lond. Hort. Brit. 38, in syn. (1830) = **Mimetes**, Salisb. (Protea.).
vaccinifolia, Lond. l. c. = M. vaccinilifolia.

DEBRAEA, Roem. & Schult. Syst. i. 4 (1817) = **Brisma**, Rudge (Vochys.).
floribunda, Roem. & Schult. l. c. 84 = E. floribundum.
violacea, Mart. & Zucc. ex Stend. Nom. ed. II. i. 486 = E. violaceum.

DEBREGEASIA, Gaudich. Bot. Voy. Bonite, t. 90 (1844–66 *). *URTICACEÆ*, Benth. & Hook. f. ii. 390.
Leucocnide, Miq. Fl. Jungh. i. 86 (1851).
Morocarpus, Sieb. & Zucc. in Abh. Math.-Phys. Akad. Muench. iv. iii. (1846) 218; (Fl. Jap. Fam. Nat. ii. 91).
bicolor, Wedd. in DC. Prod. xvi. i. 225* = hypoleuca.
dichotoma, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 463.—Ins. Java.
edulis, *Wedd. l. c.*—Japon.
hypoleuca, *Wedd. l. c.* 463. t. 15.—Ind. or.
longifolia, Wedd. in DC. Prod. xvi. i. 225* = Wallichiana.
longifolia, Wedd. l. c. 225* = velutina.
velutina, *Gaudich. Bot. Voy. Bonite*, t. 90.—Ind. or.
Wallichiana, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 464.—Reg. Himal. ; Burma ; Zeylan.

DECABELONE, Decne. in Ann. Sc. Nat. Sér. V. xiii. (1870–71) 404. t. 3. *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 784.
Barklyi, *Dyer, in Bot. Mag.* t. 6903.—Afr. austr.
elegans, *Decne. in Ann. Sc. Nat.* Sér. V. xiii. (1870–71) 404.—Afr. austr.

DECACERAS, Harv. Thes. Cap. ii. 9. t. 114 (1863) = **Anisotoma**, Fenzl (Asclep.).
Hutoni, Harv. l. c.—Afr. austr.

DECACHAENA, Torr. A. Gray, ex Lindl. Veg. Kingd. 755 (1847) = **Gaylussacia**, H. B. & K. (Vaccin.).

DECACHÆTA, DC. Prod. v. 133 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 388.
Haenkeana, *DC. l. c.*—Mexic.
Seemanni, *Benth. & Hook. f. Gen.* ii. 289.—Mexic.

DECACHAETA, Gardn. in Hook. Lond. Journ. Bot. v (1846) 462 = **Ageratum**, Linn. (Compos.).
conferta, Gardn. l. c. 468 = A. confertum.
longifolia, Gardn. l. c. 462 = A. longifolium.

DECADIA, Lour. Fl. Cochinch. 315 (1790). *INCERT. SED.*—Cf. Benth. & Hook. f. ii. 668.
aluminosa, *Lour. l. c.*—Cochinch.
racemosa, Buch.-Ham. ex Wall. Cat. n. 4430 = Symplocos racemosa.
spicata, Buch.-Ham. l. c. n. 4417 = Symplocos spicata.

DECADON, G. Don, Gen. Syst. ii. 721 (1832) = DECODON, J. F. Gmel. = **Nesæa**, Comm. (Lythrar.).

DECADONTIA, Griff. Notul. iv. 175 (1854) = **Sphenodesma**, Jack (Verbenac.).
caerulescens, Griff. l. c. = S. Griffithiana.

DECAGONOCARPUS, Engl. in Mart. Fl. Bras. xii. ii. 105 (1874). *RUTACEÆ*.
oppositifolius, *Engl. l. c.*—Bras.

DECAISNEA, Brongn. in Duperr. Voy. Coq. Bot. 192. t. 89 (1829) = **Prescottia**, Lindl. (Orchid.).
densiflora, Brongn. l. c. = P. densiflora.

DECAISNEA, Hook. f. & Thoms. in Proc. Linn. Soc. ii. (1854) 250. *BERBERIDEÆ*, Benth. & Hook. f. i. 42.
insignis, *Hook. f. & Thoms. l. c.*—Reg. Himal.

DECAISNEA, Lindl. in Wall. Cat. n. 7388 (1831) = **Tropidia**, Lindl. (Orchid.).
angulosa, Lindl. l. c.—Ind. or.

DECALEPIS, Boeck. in Engl. Bot. Jahrb. v. (1884) 509 = **Tetraria**, Beauv. (Cyperac.).
Dregeana, Boeck. l. c. (= *Cyathocoma nigrovaginata*) —Afr. austr.

DECALEPIS, Wight & Arn. Contrib. 64 (1834) *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 744.
Hamiltonii, *Wight & Arn. l. c.*—Ind. or.

DECALOBA, Rafin. Fl. Tellur. iv. 76 (1836) = **Ipomœa**, Linn. (Convolv.).

DECALOBA, M. Roem. Syn. Pepon. 131, 152 (1846) = **Passiflora**, Linn.
alnifolia, M. Roem. l. c. 138 = P. alnifolia.
Anderssonii, M. Roem. l. c. = P. Anderssonii.
bifora, M. Roem. l. c. 161 = P. innata.
bilobata, M. Roem. l. c. 154 = P. capsularis.
bogotensis, M. Roem. l. c. 153 = P. alnifolia.
bryonioides, M. Roem. l. c. 153 = P. bryonioides.
capsularis, M. Roem. l. c. 154 = P. capsularis.
cirrhiflora, M. Roem. l. c. 164 = P. cirrhiflora.
cuneata, M. Roem. l. c. 158 = P. cuneata.
cynathopora, M. Roem. l. c. 157 = P. auriculata.
dentata, M. Roem. l. c. 156 = P. Naddiana.
elliptica, M. Roem. l. c. 153 = P. elliptica.
exsudans, M. Roem. l. c. 160 = P. bryonioides.
filipes, M. Roem. l. c. 161 = P. filipes.
gemmiflora, M. Roem. l. c. 164 = P. Vespertilio.
hexicycla, M. Roem. l. c. 160 = P. Vespertilio.
holosericea, M. Roem. l. c. 164 = P. holosericea.
Jacquiui, M. Roem. l. c. 156 = P. pulchella.
lorentiensis, M. Roem. l. c. 160 = P. lorentiensis.
lowreuciana, M. Roem. l. c. 156 = P. Naddiana.
lunulifolia, M. Roem. l. c. 159 = P. lunulifolia.
Lschenaultii, M. Roem. l. c. 160 = P. Leschenaultii.
lunata, M. Roem. l. c. 158 = P. lunata, Vent.
lyrasifolia, M. Roem. l. c. 157 = P. obtegens.
mollis, M. Roem. l. c. 162 = P. mollis.
oblongata, M. Roem. l. c. 155 = P. oblongata.
obscura, M. Roem. l. c. 157 = P. foetida.
obtusa, M. Roem. l. c. 156 = P. rugosa?

DECALOBA:—
onychina, M. Roem. Syn. Pepon. 162 = Passiflora amethystina.
organensis, M. Roem. l. c. 160 = P. organensis.
pallida, M. Roem. l. c. 159 = P. mucronata.
pendulifora, M. Roem. l. c. 157 = P. pendulifora.
perfoliata, M. Roem. l. c. = P. perfoliata.
pilifera, M. Roem. l. c. 161 = P. capsularis.
punctata, M. Roem. l. c. 156 = P. punctata.
retusa, M. Roem. l. c. = P. Maximiliana.
Rohrii, M. Roem. l. c. = P. auriculata.
rotundifolia, M. Roem. l. c. 159 = P. rotundifolia.
rubra, M. Roem. l. c. 158 = P. rubra.
rufa, M. Roem. l. c. 159 = P. lunata †
semilunaria, M. Roem. l. c. 154 = P. capsularis.
septenata, M. Roem. l. c. 164 = P. cirrhiflora.
serrifora, M. Roem. l. c. = P. serriflora.
sicyoides, M. Roem. l. c. 163 = P. sicyoides.
silvestris, M. Roem. l. c. 153 = P. silvestris.
Smithii, M. Roem. l. c. 161 = P. capsularis.
subhyfida, Klotzsch, in Schomb. Faun. et Fl. Gui. 1168, nomen.—Guiana.
surinamensis, M. Roem. Syn. Pepon. 158 = P. Vespertilio.
Swartzii, M. Roem. l. c. 160 = P. Swartzii.
tuberosa, M. Roem. l. c. = P. tuberosa.
Vespertilio, M. Roem. l. c. 155 = P. Vespertilio.

DECALOPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 153 = **Chamaelauctium**, Desf. (Myrtac.).
darwinioides, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 226 = C. ciliatum.
juniperinum, Turcz. l. c. 225 = C. megalopetalum.
melaleucum, Turcz. l. c. = C. megalopetalum.
micranthum, Turcz. l. c. 226 = C. gracile.
paucifolorum, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 154 = C. paucifolorum.
rugulosum, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 226 = C. ciliatum.

DECAMERIUM, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1842) 259 = **Gaylussacia**, H. B. & K. (Vaccin.).
dumosum, Nutt. l. c. 260 = G. dumosa.
frondosum, Nutt. l. c. = G. frondosa.
hirtellum, Nutt. l. c. = G. dumosa.
resinosum, Nutt. l. c. = G. baccata.

DECANDOLIA, Bast. Fl. Maine-et-Loire, 28 (1809) = **Agrostis**, Linn. (Gramin.).
alba, Bast. l. c. 29 = A. alba.
effusa, Bast. l. c. 28 = Milium effusum.
pumila, Bast. l. c. = A. vulgaris.
stolonifera, Bast. l. c. 29 = A. stolonifera.
vulgaris, Bast. l. c. 28 = A. vulgaris.

DECANEMA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 338. t. 12. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 765.
Bojerianum, *Decne. l. c.*—Madag.

DECANEURUM, DC. in Guillem. Arch. Bot. ii. (1838) 516 = **Centratherum**, Cass. (Compos.).
amygdalinum, DC. Prod. v. 68 = Vernonia amygdalina.
angustifolium, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 8 = C. Hookeri.
courtallense, Wight, Ic. iii. iv. 6. t. 1081 = C. courtallense.
dinduiglense, DC. in Wight, Contrib. 7 = Vernonia indica.
divergens, DC. l. c. 8 = Vernonia divergens.
ophiopum, DC. l. c. 7 = C. molle.
flumbelliferum, DC. Prod. v. 68.—Ins. Borbon.
frutescens, DC. l. c. 66 = C. frutescens.
glabrum, DC. l. c. 67 = Eupatorium glabrum.
grande, DC. l. c.—Madag.
javanicum, Miq. Fl. Ind. Bat. ii. 20.—Java.
lllachum, Dals. & Gibs. Bomb. Fl. 214.—Ind. or.
microcephalum, Dals. in Hook. Kew Journ. iii. (1851) 231 = Lamprocheantium microcephalum.
molle, Dals. & Gibs. Bomb. Fl. 122 = C. phyllolasenum.
molle, DC. in Deless. Ic. Sel. iv. t. 8 = C. molle.
monanthariaefolium, Klatt, in Ann. Sc. Nat. Sér. V. xvii. (1873) 303.—Ins. Nossibé.
obovatum, DC. Prod. v. 67 = Vernonia scandens.
phyllolasenum, DC. l. c. vii. 264 = C. phyllolasenum.
reticulatum, DC. in Wight, Contrib. 9 = C. reticulatum.
scabridum, DC. l. c. 7 = C. molle.

DECANEURUM :—
scariosum, DC. Prod. vii. 264 = Vernonia scariosa, *Arn.*
senegalense, DC. l. c. v. 68 = Vernonia senegalensis.
silhetense, DC. l. c. 67 = Vernonia bracteata.
silhetense, Wight, Ic. t. 1083 = Vernonia peninsularis.

DECANEURUM, Sch. Bip. Tanacet. 44 (1844) = **Chrysanthemum**, Tourn. (Compos.).
serotinum, Sch. Bip. l. c. 45 = C. serotinum.

DECAPENTA, Rafin. Sylva Tellur. 165 (1838) = **Litsea**, Lam. (Laurin.).
involucrata, Rafin. l. c. = L. sebifera.

DECAPRISMA, Rafin. Fl. Tellur. ii. 78 (1836) = **Canapanula**, Tourn.
cespitosa, Rafin. l. c. = C. caespitosa.

DECAPTHRA, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) ii. 497. *CRUCIFERAE*, Benth & Hook. f. i. 90.
trifida, *Turcz. l. c.* 498.—Chili.

DECARAPHE, Miq. Comm. Phyt. ii. 84 (1838–40) = **Miconia**, Ruiz & Pav. (Melast.).
floribunda, Miq. in Linnaea, xviii. (1844) 280 = M. Miqueliana.
Fockeana, Miq. Comm. Phyt. ii. 84. t. 10 = M. eriopoda.
Hostmanni, Steud. in Flora, xxvii. (1844) 722 = M. platyhedra.

DECARHAPHE, Steud. in Flora, xxvii. (1844) 722 = praec.

DECARINIUM, Rafin. Neogenyt. 1 (1825) = **Croton**, Linn. (Euphorb.).
glandulosum, Rafin. l. c. = C. glandulosum.

DECASCHISTA, Reichb. Nom. 200 (1841) = seq.

DECASCHISTIA, Wight & Arn. Prod. 52 (1834). *MALVACEAE*, Benth. & Hook. f. i. 206.
cramuscula, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 237.—Burma.
crotonifolia, *Wight & Arn. Prod.* 52.—Ind. or.
parviflora, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 68.—Burma.
trilobata, *Wight, Ic.* t. 88.—Ind. or.

DECASPERMUM, Forst. Char. Gen. 78. t. 37 (1776). *MYRTACEAE*, Benth. & Hook. f. i. 716; ii. 103. NELITRIS, Spreng. Syst. ii. 488 (1826).
Blancoi, *Vidal, Phan. Cuming. Philipp.* 172.—Ins. Philipp.
fruticosum, *Forst. Char. Gen.* 74.—Ins. Pacif. †
paniculatum, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 61.—Burma.
rubrum, *Baill. Hist. Pl.* vi. 341.—Ins. Molucc.
sericeum, *Hance, in Journ. Bot.* xv. (1877) 333.—China.

DECASPORA, R. Br. Prod. 548 (1810) = **Trichosarpa**, R. Br. (Epacrid.).
Cliffortii, F. Muell. in Trans. Phil. Soc. Viet. i. (1855) 106.—T. Clarkei.
Cunninghamii, DC. Prod. vii. 758 = T. disticha.
disticha, R. Br. Prod. 548 = T. disticha.
Gunnii, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 270 = T. Gunnii.
involucrata, F. Muell. Fragm. vi. 57 = T. involucrata.
oxycoccoides, A. Cunn. ex DC. Prod. vii. 758 = T. thymifolia.
parviflora, Stschegl. in Bull. Soc. Nat. Mosc. (1859) i. 10 = T. parviflora.
pumila, F. Muell. Fragm. vi. 57 = Pentachondra pumila.
thymifolia, R. Br. Prod. 548 = T. thymifolia.
thymifolia, A. Cunn. ex DC. Prod. vii. 758 = T. disticha.

DECASTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 157 (1861) = **Clusonia**, Linn. (Capparr.).
hirtus, Klotzsch, l. c. = C. hirta.
sanctaharicus, Klotzsch, l. c. = C. strigosa.

DECASTROPHIA, Griff. Notul. iv. 737 (1854); Ic. Pl. Asiat. t. 613 f. 4 = **Erythropsalum**, Blume (Oxalid.).
inconspicua, Griff. l. c. = E. scandens.

DECATELES, Rafin. Sylva Tellur. 35 (1838) = **Hidcroxylon**, Dill. (Sapotac.).
latifolia, Rafin. l. c. = S. decandrum.
lycioides, Rafin. l. c. = Bumelia lycioides.

DECATROPIS, Hook. f. in Benth. & Hook. f. Gen. i. 898 (1869). *RUTACEAE*, Benth. & Hook. f. i. c. Coulteri, *Hook. f. l. c.* 899.—Mexic.

DECHAESIA, F. Muell. Fragm. xi. 71 (1879). *COMPOSITAE*.
hecatocephala, *F. Muell. l. c.* 72.—Austral.

DECEMIUM, Rafin. Fl. Ludov. 34, in obs. (1817) = **Hydrophyllum**, Linn.
hirtum, Rafin. Cat. 18 = H. hispidum †

DECHAMPSIA, Kanth, in Mém. Mus. Par. ii. (1815) 72 = **Deschampsia**, Beauv. (Gram.).

DECKENIA, H. Wendl. ex Seem. in Gard. Chron. (1870) 561 = **Acanthophoenix**, H. Wendl. (Palm.).
nobilis, H. Wendl. l. c.—Ins. Seychelles.

DECKERA, Sch. Bip. in Flora, xvii. (1834) 479 = **Pieris**, Linn. (Compos.).
aculeata, Sch. Bip. l. c. = P. aculeata.
aspleniodes, Sch. Bip. l. c. = P. aspleniodes.
callosa, Pomel, Nouv. Mat. Fl. Atl. 14.—Algeria.
canescens, C. A. Mey. ex Fisch. Mey. & Trautv. Ind. Sem. Hort. Petrop. iv. 43 = P. laevis.
glomerata, Pomel, Nouv. Mat. Fl. Atl. 13.—Algeria.
montana, Pomel, l. c. 14.—Algeria.
nilotica, Sch. Bip. in Flora, xvii. (1834) 479 = P. sulphurea.
racemosa, Pomel, Nouv. Mat. Fl. Atl. 13.—Algeria.
rubiginosa, Pomel, l. c. 270.—Algeria.
scabra, Sch. Bip. in Flora, xvii. (1834) 479 = P. scabra.
sulphurea, Sch. Bip. l. c. = P. sulphurea.

DECKERIA, Karst. in Linnaea, xxviii. (1856) 258 = **Iriartea**, Ruiz & Pav. (Palm.).
Cornets, Karst. l. c. = I. cornuta.
elegans, Linden, Illustr. Hortic. xxviii. (1881) 16.—Hab. ?
Lamarchiana, Karst. in Linnaea, xxviii. (1856) 259 = Dictyocaryum Lamarckianum.
paniculata, Karst. l. c. = I. phaseocarpa.
ventricosa, Karst. l. c. = I. ventricosa.

DECLIEUXIA, H. B. & K. Nov. Gen. et Sp. iii. 352. t. 281 (1818). *RUBIACEAE*, Benth. & Hook. f. ii. 126.
PAVLLOCARPUS, Pohl, ex DC. Prod. iv. 479 (1830).
alba, Zucc. in Schult. Mant. iii. 113 = chiococcoides.
aspalathoides, *Muell. Arg. in Flora*, lix. (1876) 437.—Bras.
brachylobe, *Muell. Arg. l. c.* 436.—Bras.
brasiliensis, Muell. Arg. l. c. 434 = herbacea.
brevicalla, *Muell. Arg. l. c.* 435.—Bras.
cacuminus, *Muell. Arg. l. c.* 436.—Bras.
chiococcoides, Bartl. ex DC. Prod. iv. 479 = mexicana.
chiococcoides, *H. B. & K. Nov. Gen. et Sp.* iii. 354. t. 281.—Venezuela.
clinopodioides, *Muell. Arg. in Flora*, lix. (1876) 434.—Bras.
cornuta, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 106 = Congdonia caerulea.
cordigera, *Mart. & Zucc. in Schult. Mant.* iii. 112.—Bras.
daphnoides, *Mart. & Zucc. l. c.*—Bras.
deltoides, *Muell. Arg. in Flora*, lix. (1876) 434.—Bras.
divergentiflora, *DC. Prod.* iv. 480.—Bras.
follicus, *DC. l. c.*—Bras.
Galeottii, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 231.—Mexic.
gallodes, *DC. Prod.* iv. 479.—Bras.
glabra, DC. l. c. 481 = chiococcoides.
glauca, *Mart. & Schlecht. in Linnaea*, iv. (1829) 8.—Bras.
herbacea, *Cham. & Schlecht. l. c.* 11.—Bras.
imbricata, DC. Prod. iv. 480 = daphnoides.
intermedia, *Muell. Arg. in Flora*, lix. (1876) 435.—Bras.
juniperina, *St. Hil. Voy. Distr. Diam. Bras.* i. 366.—Bras.

DECLIEUXIA :—
latifolia, Hochst. in Flora, xxvi. (1843) 70 = Pentaniaia variabilis.
leiophylla, *Muell. Arg. in Flora*, lix. (1876) 436.—Bras.
lysimachioides, *Zucc. in Schult. Mant.* iii. 112.—Bras.
mariolde, *Mart. & Zucc. l. c.* 111.—Bras.
mexicana, DC. Prod. iv. 479 = chiococcoides.
mollis, *Zucc. in Schult. Mant.* iii. 112.—Bras.
mucronalis, *Mart. ex Cham. & Schlecht. in Linnaea*, iv. (1829) 10.—Bras.
muscosa, *St. Hil. Voy. Distr. Diam. Bras.* i. 365.—Bras.
oenanthioides, *Mart. & Zucc. in Schult. Mant.* iii. 112.—Bras.
origanoides, *Zucc. in Schult. l. c.*—Bras.
orthophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 451.—Bras.
Passerina, *Mart. & Zucc. in Schult. Mant.* iii. 112.—Bras.
polygaloides, *Zucc. in Schult. l. c.* 113.—Bras.
prunicea, DC. Prod. iv. 481 = glauca.
prunelloides, Klotzsch, in Eckl. & Zeyh. Enum. 363 = Pentaniaia variabilis.
psychotrioides, *DC. Prod.* iv. 481.—Am. austr.
pulverulenta, Mart. ex Cham. & Schlecht. in Linnaea, iv. (1829) 9 = chiococcoides.
revoluta, *Muell. Arg. in Flora*, lix. (1876) 436.—Bras.
rhexioides, *Mart. & Zucc. in Schult. Mant.* iii. 112.—Bras.
rubioides, *Zucc. l. c.*—Bras.
sarmentoides, *Mart. & Zucc. l. c.* 111.—Bras.
scabra, Benth. in Linnaea, xxiii. (1850) 465 = diversifolia.
sclerophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 467.—Bras.
Selloana, *Muell. Arg. in Flora*, lix. (1876) 436.—Bras.
spergulifolia, *Mart. & Zucc. in Schult. Mant.* iii. 111.—Bras.
thymbroides, *Mart. & Zucc. l. c.*—Bras.
verticillata, *Muell. Arg. in Flora*, lix. (1876) 436.—Bras.
vincoides, Mart. Reise Bras. i. 397; et ex Zucc. in Schult. Mant. iii. 113 = chiococcoides.

DECODON, J. F. Gmel. Syst. 677 (1791) = **Nesaea**, Comm. (Lythrar.).
aquaticum, J. F. Gmel. l. c. = N. verticillata.
verticillatum, Ell. Sketch. i. 343 = N. verticillata.

DECODONTIA, Haw. Syn. Pl. Succ. 28, in obs. (1812) = **Huernia**, R. Br. (Asclep.).

DECORIMA, Raf. Sylva Tellur. 31 (1838) = **Eisiphus**, Tourn. (Rhamn.).
trinervis, Raf. l. c. = Z. exserta.
umbellata, Raf. l. c. = Karwinskia Humboldtiana.

DECOSTEA, Ruiz & Pav. Prod. 130 (1794) = **Grisellinia**, Forst. (Cornac.).
jodinifolia, Griseb. in Goett. Abh. vi. (1854) 122.—Chili.
racemosa, Phil. in Linnaea, xxviii. (1856) 708.—Chili.
ruscifolia, C. Gay, Fl. Chil. viii. 395. t. 93 b.—Chili.
scandens, Ruiz & Pav. Syst. Veg. 259.—Chili.

DECTIS, Raf. Fl. Tellur. ii. 43 (1836) = **Solidago**, Tourn. (Compos.).
arborescens, Raf. l. c. = Olearia nitida.
decurrens, Raf. l. c. = S. Virgaurea.
Leucadendron, Raf. l. c. = Senecio Leucadendron.
rugosa, Raf. l. c. = S. rugosa.

DECUMARIA, Linn. Sp. Pl. ed. II. App. 1663 (1762). *SAXIFRAGEAE*, Benth. & Hook. f. i. 642.
Forsythia, Walt. Fl. Carol. 124 (1788).
barbara, *Linn. Sp. Pl.* ed. II. 1663.—Am. bor.
Forsythia, Michx. Fl. Bor. Am. i. 282 = barbara.
prostrata, Lodd. ex Lond. Hort. Brit. 188 = barbara.
radicans, Moench, Meth. 107 = barbara.
sarmentosa, Bosc, in Act. Soc. Hist. Nat. Par. i. (1792) 76 = barbara.
scandens, Salisb. Prod. 355 = barbara.

DEDEA, Baill. Adansonia, xii. 286 (1879). *SAXIFRAGEAE*.
major, *Baill. l. c.* 289.—N. Caled.
minor, *Baill. l. c.*—N. Caled.

DEERINGIA, R. Br. Prod. 418 (1810). *AMARANTACEAE*, Benth. & Hook. f. iii. 28.
Cladostachys, D. Don, Prod. Fl. Nep. 76 (1825).
Cyclosperma, Raf. Fl. Tellur. iii. 45 (1836).
altissima, *F. Muell. Fragm.* ii. 92.—Austral.
Amherstiana, Wall. Cat. n. 6889 = Bosia Amherstiana.
baccata, Moq. in DC. Prod. xiii. 2. 236 = celosioides.
celosioides, *R. Br. Prod.* 418.—As. et Austral. trop.
indica, Retz. ex Blume, Bijdr. 582 = celosioides.
polysperma, *Moq. in DC. Prod.* xiii. 2. 236.—Ind. or. tetragyna, *Kunth. Hort. Berol.* 19; *Fl. Ind.* i. 668.—Ins. Moluccas.
virgata, Zipp. ex Span. in Linnaea, xv. (1841) 346 = celosioides.

DEFFORGIA, Lam. Illustr. ii. 71. t. 125 (1793) = **Forgesia**, Comm. (Saxifr.).
borbonica, Lam. l. c. = F. racemosa.

DEGUELIA, Aubl. Pl. Gui. 750. t. 300 (1775) = **Derris**, Lour. (Legumin.).
arborescens, Spreng. Syst. iv. Cur. Post. 269 = Swartzia nitens.
scandens, Aubl. Pl. Gui. ii. 750. t. 300 = Derris guianensis.
uliginosa, Baill. in Bull. Soc. Linn. Par. i. (1885) 442 = Derris uliginosa.

DEHAASIA, Blume, Rumphia, i. 161. tt. 44, 45, 47 (1835). *LAURINEAE*, Benth. & Hook. f. iii. 162.
Haasia, Nees, Syst. Laurin. 372 (1836).
caesia, *Blume, Mus. Bot. Lugd. Bat.* i. 338.—Java.
cuneata, *Blume, Rumphia*, i. 164. t. 46.—Java.
elongata, *Blume, l. c.* 163. t. 47.—Sumatra.
firma, *Blume, Mus. Bot. Lugd. Bat.* i. 338.—Malaya.
media, *Blume, Rumphia*, i. 163. t. 45.—Ins. Amboin.
microcarpa, *Blume, l. c.* 162. t. 44.—Java.
paradoxa, *Blume, Mus. Bot. Lugd. Bat.* i. 338.—Borneo.
pauciflora, *Blume, l. c.*—Malaya.
spathulata, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 113, 112.—Malaya.
squarrosa, *Hassk. Retzia*, 8.—Java.

DEHERAINIA, Decne. in Ann. Sc. Nat. Sér. VI. iii. (1876) 138. *MYRSINEAE*.
smaragdina, *Decne. l. c.* 139.—Mexic.

DELANTRA, Cham. & Schlecht. in Linnaea, i. (1826) 195. *GENTIANEAE*, Benth. & Hook. f. ii. 810.
Callofisma, Mart. Nov. Gen. et Sp. ii. 107. tt. 188, 184 (1827).
diuriaeta, *Garda. ex Benth. & Hook. f. Gen.* ii. 810.—Bras.
erubescens, *Cham. & Schlecht. in Linnaea*, i. (1826) 196.—Bras.
nervia, *Cham. & Schlecht. l. c.* 197.—Bras.
pallescens, Cham. & Schlecht. l. c. 196 = erubescens.

DELDAMIA, Noronha, ex Thou. Hist. Vég. Iles Afr. 61. t. 20 (1807). *PASSIFLOREAE*, Benth. & Hook. f. i. 811.
Thompsonia, R. Br. in Trans. Linn. Soc. xiii. (1821) 221.
alata, *Thou. l. c.*—Madag.
biplinnata, *Thl. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 49.—Madag.
Commersoniana, *DC. Prod.* iii. 327.—Madag.
Noronhiana, DC. l. c. = alata.
setigera, *Thl. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 50.—Madag.
Thompsoniana, *DC. Prod.* iii. 327.—Madag.

DELOSMA, Andrz. ex DC. Syst. ii. 450, 451, in syn. (1831); Spach, Hist. Vég. Phan. vi. 397 (1838) = **Hesperis**, Linn. (Crucif.).
albiflorum, Fuss, Fl. Transs. 57 = H. nivea.
alpinum, Fuss, l. c. = H. alpina.
glabrum, Fuss, l. c. 56 = H. alpina.
inodorum, Fuss, l. c. 57 = H. matronalis.
Kladnii, Fuss, l. c. 56 = H. alpina.
matronale, Andrz. ex DC. Syst. ii. 450 = H. matronalis.
moniliforme, Fuss, Fl. Transs. 57 = H. alpina.
odoratum, Fuss, l. c. = H. matronalis.
runcinatum, Fuss, l. c. 56 = H. matronalis.
sibiricum, Andrz. ex DC. Syst. 451 = H. matronalis.

DELOSMA, —
mucronatum, Andrz. in Bess. Enum. Pl. Volh. 83 = H. matronalis.
triste, Spach, Hist. Vég. Phan. vi. 398 = H. triste.

DELPHANTHUS, Maxim. in Mém. Acad. Pétersb. Sér. VII. x. (1867) n. xvi. p. 2. *SAXIFRAGEAE*.
biflda, *Maxim. l. c.* 3.—Japon.

DELPHOLLIA, Schum. & Thonn. in Danske, Selsk. Afh. iv. (1827) 16; Benkr. Guin. Pl. 249). *SAPINDACEAE*, Benth. & Hook. f. i. 405.
Protea, Cambess. in Mém. Mus. Par. xvlii. (1829) 23. t. 1 C.
borbonica, *Scheff. in Flora*, lii. (1869) 306; *et in Tijdschr. Nederl. Ind.* xxxi. (1870) 17.—Ins. Borbon.
cuneifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 482.—Afr. trop.
grandifolia, *Hook. f. in Hook. Niger Fl.* 249.—Afr. trop.
insignis, *Hook. f. l. c.* 250.—Afr. trop.
laurifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 482.—Afr. trop.
neglecta, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 368.—Madag.
oblongifolia, Radlk. l. c. 369 = Sapindus oblongifolius obovatus, *Radlk. l. c.* 368.—Afr. trop.
Perrillei, *Radlk. l. c.* 369.—Madag.?
pinnata, *Schum. & Thonn. Beskr. Guin. Pl.* 242.—Afr. trop.
xanthocarpa, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 369.—Afr. trop.

DEINOSMOS, Raf. Fl. Tellur. ii. 49 (1836) = **Aster**, Tourn. (Compos.).
sicuius, Raf. l. c. = A. Linosyris.

DELINIA, Mart. & Gal. in Bull. Acad. Brux. xi. ii. (1844) 195. *LABIATAE*, Benth. & Hook. f. ii. 1198.
cocchica, *Mart. & Gal. l. c.*—Mexic.

DELA, Adans. Fam. ii. 499 (1763) = **Seseli**, Linn (Umbellif.).

DELABECHEA, Lindl. in Mitch. Journ. Trop. Austral. 156 (1848) = **Sterculia**, Linn.
rupestris, Lindl. l. c. = S. rupestris.

DELAIRIA, Lem. in Ann. Sc. Nat. Sér. III. i. (1844) 379 = Senecio, Tourn. (Compos.).
odorata, Lem. l. c. 380 = S. mikanioides.

DELARBREA, Vieill. in Bull. Soc. Linn. Norm. ix. (1865) 342. *ARALIACEAE*, Benth. & Hook. f. i. 935.
collina, *Vieill. l. c.*—N. Caled.
Michieana, F. Muell. Fragm. vii. 95 = Porospermum Michieanum.
paradoxa, Vieill. in Bull. Soc. Linn. Normand. ix (1865) 343 = collina.
spectabilis, *Linden & André, in Illustr. Hortic.* xxv. (1878) t. 314.—N. Caled.

DELARIA, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 406 = **Sapphia**, Afzel. (Legumin.).
odorata, Regel, Cat. Pl. Hort. Almakov, 48.—Hab.?
ovalifolia, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 406.—Bras.
pyrifolia, Desv. l. c. = B. polygalaeea.

DELASTREA, A. DC. in DC. Prod. viii. 295 (1844) = Lobranzia, A. DC. (Sapot.).
Bojeri, A. DC. l. c. 296 = L. Bojeri.

DELIA, Dum. Fl. Belg. 110 (1827) = **Spergularia**, J. & C. Presl (Caryophyll.).
segetalis, Dum. l. c. = S. segetalis.

DELILIA, Spreng. Syst. iii. 867 (1826) = **Elvira**, Cass. (Compos.).
Bertorii, Spreng. l. c. 874 = E. Martyni.

DELIMA, Linn. Dis. Dan. 19 (1747); Amoen. Acad. i. 408 (1749) = Tetracera, Linn. (Dillen.).
aspera, Blanco, Fl. Filip. ed. I. 459; ed. II. 293.—Ins. Philipp.
castaneifolia, G. Don, Gen. Syst. i. 71.—Sum.
discophylla, Miq. Stirp. Surin. Sel. 107.—Guian.
fimgulanyfolia, Presl, Rel. Haenk. ii. 78.—Ins. Philipp.
guianensis, Rich. ex DC. Syst. i. 466.—Guiana.

DELIMA :—
hebecarpa, DC. Syst. i. 407 = Tetracera sarmentosa.
intermedia, Blume, Bijdr. 4 = T. sarmentosa.
mexicana, Moç. & Sesse, ex DC. Syst. i. 407 = T. scabrifloru.
nitida, DC. l. c. 408.—Ins. Trinit.
Pirira, DC. l. c. = T. sarmentosa.
sarmentosa, Linn. Gen. ed. V.; Syst. ed. X. 1076 = T. sarmentosa.
Tigarea, Eichl. in Mart. Fl. Bras. xiii. i. 93 = T aspera.
tomentosa, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 816 = Doliocarpus Rolandri.
triquetula, G. Don, Gen. Syst. i. 71 = T. sarmentosa.

DELIMOPSIS, Miq. Fl. Ind. Bat. i. ii. 9 (1859) = **Tetracera**, Linn. (Dillen.).
hirsuta, Miq. l. c. 10.—Java.

DELISSEA, Gaudich. in Freyc. Voy. Bot. 457. t. 76 ad 78 (1826). *CAMPANULACEAE*, Benth. & Hook. f. ii. 546.
acuminata, Gaudich. l. c. t. 76 (Cyanese sp.).—Ins. Sandvic.
ambigua, Presl, Prod. Lobel. 47 = Rollandia ambigua.
angustifolia, Presl, l. c. (Cyanese sp.).—Ins. Sandvic. arbores, *H. Mann, in Proc. Am. Acad.* vii. (1867) 180. —Ins. Sandvic.
arborea, Presl, Prod. Lobel. 47 = Sclerotheca arborea.
asplenifolia, H. Mann, in Proc. Am. Acad. vii. (1867) 183 (Cyanese sp.).—Ins. Sandvic.
calycina, Presl, Prod. Lobel. 47 = Rollandia calycina.
clermontioides, Gaudich. Bot. Voy. Bonite, t. 47 (Clermontiae sp.).—Ins. Sandvic.
coriacea, A. Gray, in Proc. Am. Acad. v. (1861) 147 (Cyanese sp.).—Ins. Sandvic.
Delissetrian, *A. Gray, l. c.*—Ins. Sandvic.
filigera, Wawra, in Flora, lvi. (1873) 31 = Clermontia grandiflora.
fissa, H. Mann, in Proc. Am. Acad. vii. (1867) 182 (Cyanese sp.).—Ins. Sandvic.
hirtella, H. Mann, l. c. 179 (Cyanese sp.).—Ins. Sandvic.
honoluluensis, Wawra, in Flora, lvi. (1873) 11 (Cyanese sp.).—Ins. Sandvic.
Kealiae, Wawra, l. c. 10 = rhytidosperma.
Kunthiana, *Gaudich. Bot. Voy. Bonite*, t. 77.—Ins. Sandvic.
lanceolata, A. Gray, in Proc. Am. Acad. v. (1861) 147 = Rollandia lanceolata.
macrostachys, Presl, Prod. Lobel. 47 = Lobelia macrostachys.
Mannii, W. T. Brigham, ex H. Mann, in Proc. Am. Acad. vii. (1867) 189 (Cyanese sp.).—Ins. Sandvic.
obtusa, A. Gray, in Proc. Am. Acad. v. (1861) 148 (Cyanese sp.).—Ins. Sandvic.
pilosa, H. Mann, in Proc. Am. Acad. vii. (1867) 182 = Cyanea pilosa.
pinnatifida, *Presl, Prod. Lobel.* 47.—Ins. Sandvic.
platyphylla, A. Gray, in Proc. Am. Acad. v. (1861) 148 (Cyanese sp.).—Ins. Sandvic.
racemosa, H. Mann, l. c. vii. (1867) 181 = Rollandia pedunculosa.
recta, Wawra, in Flora, lvi. (1873) 30 (Cyanese sp.). —Ins. Sandvic.
Regina, Wawra, l. c. 9 = Cyanea superba.
rhytidosperma, *H. Mann, in Proc. Am. Acad.* vii. (1867) 180.—Ins. Sandvic.
subcordata, *Gaudich. in Freyc. Voy. Bot.* 457. t. 77. —Ins. Sandvic.
undulata, *Gaudich. l. c.* t. 78.—Ins. Sandvic.
Waihiae, Wawra, in Flora, lvi. (1873) 8 (Clermontiae sp.).—Ins. Sandvic.

DELIVARIA, Miq. Fl. Ind. Bat. ii. 920 (1857) = DILIVARIA, Juss. = **Acanthus**, Tourn.

DELODERIUM, Cass. in Dict. Sc. Nat. xlviii. 430 (1827) = **Leontodon**, Linn. (Compos.).
taraxacifolium, Cass. l. c. = L. creticus.

DELOGNARA, Cogn. in Bull. Soc. Linn. Par. i. (1886) 825. *CUCURBITACEAE*.
Humboldti, *Cogn. l. c.*—Madag.

DELONIX, Rafin. Fl. Tellur. ii. 93 (1836) = **Poinciana**, Tourn. (Leguosa.).
regia, Rafin. l. c. = P. regia.

DELOSTOMA, D. Don, in Edinb. Phil. Journ. ix. (1823) 368. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1043.
CODAZZIA, Karst. & Triana, in Linnaea, xxviii. (1856) 426; Fl. Columb. i. 7. t. 4 (1858).
dentatum, *D. Don, l. c.*—Peruv.
Domheyanum, DC. Prod. ix. 173 = nervosum ?
integrifolium, *D. Don, in Edinb. Phil. Journ.* ix. (1823) 364.—Peruv.
latifolium, Spitg. in Hoev. & De Vriese, Tijdschr. ix. (1842) 11 = Callichlamys riparia.
Lobbii, *Seem. in Bonplandia*, x. (1862) 72.—Peruv.
nervosum, *DC. Prod.* ix. 198.—Peruv.
stenolobium, Steud. Nom. ed. II. i. 467 = Bignonia castaneaefolia.

DELOSTYLIS, Rafin. in Am. Monthly Mag. (1819) 192; Journ. Phys. lxxxix. (1819) 102 = **Trillium**, Linn. (Liliac.).
cernuum, Rafin. l. c. = T. stylosum.
stylosum, Rafin. in Am. Monthly Mag. (1819) 192 = T. stylosum.

DELPECHIA, Montr. in Mém. Acad. Lyon, x. (1860) 221. *INCERT. SED.* Benth. & Hook. f. ii 29.
artensis, *Montr. l. c.* 222.—N. Caled.
floribunda, *Montr. l. c.* 221.—N. Caled.

DELPHIDIUM, Rafin. in Am. Monthly Mag. (1811) 356 = **Delphinium**, Tourn. (Ranunc.).

DELPHINASTRUM, Spach, Hist. Vég. Phan. vii. 336 (1841) = Delphinium, Tourn. (Ranunc.).
elatum, *fissum*, *grandiflorum*, *hybridum*, *ochroleucum*, *punicum*, Spach, Hist. Nat. Vég. vii. 338 ad 345, omnia = Delphinii speciorum homonyma.

DELPHINIUM, Tourn. ex Linn. Syst. ed. I (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 9.
CALLIRIPPA, Heist. Syst. 8 (1748).
CERATOSANTHUS, Schur, Enum. Pl. Transs. 80 (1866).
CONSOLIDA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 711 (1821).
DELPHIDIUM, Rafin. in Am. Monthly Mag. (1811) 356; Med. Fl. ii. 216 (1830).
DELPHINASTRUM, Spach, Hist. Vég. Phan. vii. 336 (1841).
PHLEDINIUM, Spach, l. c. 351 (1839).
PYTIRIUM, Rafin. Princ. Somiol. 29 (1814).
STAPHYSAGRIA, Spach, Hist. Vég. Phan. vii. 347 (1839).
Aconiti, Hohen. in Bull. Soc. Nat. Mosc. (1838) 388 = Hohenackeri.
Aconiti, *Linn. Mant.* i. 77.—Turcia.
addendum, W. R. Mc Nab, in Trans. Bot. Soc. Edinb. ix. (1868) 335 = Ajacis ?
Ajacis, Ledeb. Fl. Ross. i. 58 = orientale.
Ajacis, *Linn. Sp. Pl.* 531.—Europ. austr.
albiflorum, DC. Syst. i. 358 = hybridum.
albocoeruleum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 207.—China.
alpinum, Waldst. & Kit. Pl. Rar. Hung. iii. 273 = elatum.
altissimum, *Wall. Pl. As. Rar.* ii. 25. t. 128.—Reg. Himal.
ambigum, *Linn. Sp. Pl.* ed. II. 749.—Mauritan.
ambiguum, Loisel. in Desv. Journ. Bot. ii. (1809) 341 = Balbisiana.
ambiguum, Mill. Gard. Dict. ed. VIII. n. 8 = Ajacis.
americanum, Mill. l. c. n. 7 = azureum ?
amoenum, Bernh. ex Steud. Nom. ed. II. i. 488 = elatum ?
amoenum, *Stev. ex DC. Syst.* i. 346.—Sibir.
anomalum, Spreng. ex Steud. Nom. ed. II. i. 488 = elatum ?
anthoroideum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 369.—Syria; Cilicia.
atropurpureum, *Pall. Reise Statth. Russ. Reich.* i. 194.—Rossia.
Auckeri, Boiss. in Ann. Sc. Nat. Sér. II. xvi. (1841) 362 = persicum.
anthriscifolium, *Hance, in Journ. Bot.* vi. (1868) 207. —China.
axilliciflorum, G. Don, Gen. Syst. i. 50 = seq.
axilliiflorum, *DC. Syst.* i. 341.—Syria; Cilicia.
azureum, *Wicke. Fl. Bor. Am.* i. 314.—Am. bor.
azureum, Newberry, in Pacif. Rail. Rep. vi. 65 = simplex.
Balansae, *Boiss. & Reut. Diagn.* Ser. II. v. 12.—Algeria.
barbatum, *Bunge, in Arbeit. Nat. Ver. Riga*, i. (1847) 196.—Turkestan.

DELPHINIUM:—
Barlowii x, Lindl. Bot. Reg. t. 1944.
bicolor, *Nutt. in Journ. Acad. Phil.* vii. (1834) 10.—Am. bor. occ.
bicornutum, *Hemsl. Diagn. Pl. Nov.* 17.—Mexic.
bithynicum, Griseb. Spicil. Fl. Rumel. i. 320 = orientale.
Boissierii, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 8 = speciosum.
Bovei, *Decne. in Ann. Sc. Nat.* Sér. II. iv. (1835) 356. —Arab.
brachycentrum, *Ledeb. Fl. Ross.* i. 60.—Kamschatka.
brevicorne, *Vis. Fl. Dalm.* iii. 90.—Dalmat.
Brunonianum, *Royle, Illustr. Bot. Himal.* 56.—Reg Himal.
caeruleum, *Jacquem. Voy. Bot.* 7. t. 6.—Reg. Himal.
californicum, *Torr. & Gray, Fl. N. Am.* i. 31.—Calif
Calleri, *Franchet, in Bull. Soc. Linn. Par.* i. (1882) 229.—China.
camptocarpum, *Fisch. & Mey. in Ledeb. Fl. Ross.* i. 58.—Reg. Casp.
camptocarpum, C. Koch, in Linnaea, xx. (1841) 247 = persicum.
cappadocicum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 367.—As. Min.; Persia.
cardinale, *Hook. Bot. Mag.* t. 4887.—Am. bor. occ.
cardiopetalum, DC. Syst. i. 347 = halteratum.
Carela, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 195 = scabriflolium.
carolinianum, Walt. Fl. Carol. 155 = azureum.
Carumnae, Walp. Rep. ii. 744, sphalm. vide D. *Garumnae*.
casimirianum, *Royle, Illustr. Bot. Himal.* 55. t. 12.—Reg. Himal.
caucasicum, *C. A. Mey. Verz. Pfl. Cauc.* 200.—Reg. Caucas.
Chamissonis, *Prits. ex Walp. Rep.* ii. 744.—Am. bor.
cheilanthum, *Fisch. ex DC. Syst* i. 352.—Sibiria.
chinense, Fisch. ex DC. Prod. i. 53 = grandiflorum.
cilistum, Stev. Cat Hort. Dorpat. (1811) ex Bieb. Fl. Taur. Cauc. iii. 370 = elatum.
cinereum, *Boiss. Diagn.* Ser. I. i. 67.—Lydia.
Cisseanum, Host, Fl. Austr. ii. 67 = elatum.
coccineum, Torr. in Pacif. R. Rep. iv. 42 = cardinale.
confertum, *Guss. Fl. Sic. Prod* ii. 30.—Sicil.
confusum, *Lowe, Man. Fl. Mad.* i. 585.—Ins. Madera
Consolida, Gouan, Hort. Monsp. 258 = pubescens.
Consolida, *Linn. Sp. Pl.* 530.—Europ.; As. bor.
Consolida, Sibth. & Sm. Fl. Graec. Prod. i. 370 = Ajacis.
corymbosum, *Regel, in Act. Hort. Petrop.* vii. (1880) 547.—Turkestan.
crassicaule, *Ledeb. Fl. Ross.* i. 62.—Sibiria.
crassifolium, Schrad. ex Spreng. Gesch. Bot. ii. 201 = speciosum.
cuneatum, Stev. ex DC. Syst. i. 359 = elatum.
cyphopetalum, *Boiss. Diagn.* Ser. I. vi. 7.—Persia.
dahuricum, Stev. ex DC. Prod. i. 53 = cheilanthum.
dasyanthum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 138.—Reg. Altai.
dasycarpum, Stev. ex DC. Syst i. 347 = elatum.
dasycaulon, *Fresen. in Mus. Senckenb.* ii. (1837) 272.—Ind. or.; Abyssin.
dasystachyon, *Boiss. & Bal. Diagn.* Ser. II. vi. 7.—As. Min.
dauricum, Bess. Cat. Hort. Crem. Suppl. iii. (1814) 9 = cheilanthum.
dauricum, Georgi, Beschr. Russ. Reich. iii. iv. 1052 = hybridum.
decorum, A. C. Anders. Cat. Pl. Nev. 117, ex S. Wats. Biblog. Ind. N. Am. Bot. 12 = depauperatum ?
decorum, Benth. Fl. Hartw. 295 = variegatum.
decorum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 33. —Calif.
denudatum, *Wall. Cat.* n. 6719.—Reg. Himal.
depauperatum, *Nutt ex Torr. & Gray, Fl. N. Am.* i. 33.—Am. bor. occ.
deserti, *Boiss. Fl. Orient* i. 58.—Arabia.
dictyocarpum, Steud. Nom. ed. II. i. 488 = dyctiocarpum.
diffusum, Stokes, Bot. Mat. Med. iii. 218 = D. Consolida.
discolor, *Fisch. ex Sweet, Hort. Brit.* ed. II. 9, nomen. —Sibir.
divaricatum, Dulac, Fl. Hautes-Pyr. 223 = D. Consolida.
divaricatum, *Ledeb. in Eichw. Casp. Cauc.* 16. t. 16.—Reg. Caucas. et Casp.

DELPHINIUM:—
dyctiocarpum, *DC. Syst.* i. 360.—Sibiria.
elatum, All. Fl. Pedem. ii. 68 = montanum.
elatum, *Linn. Sp. Pl.* 531.—Europ. ; Reg. Himal.
elatum, Porter, in Haydn, Rep. (1870) 473 = trolliifolium.
elatum, Willd. Enum. Hort. Berol. 573 = laxiflorum.
elegans, *DC. Syst.* i. 356.—Am. bor.?
emarginatum, j. & C. Presl, Delic. Prag. 8 = pentagynum.
exaltatum, *Ait. Hort. Kew.* ed. I. ii. 244.—Am. bor.
exaltatum, Hook. Fl. Bor. Am. i. 25 = scopulorum.
exaltatum, Hook. & Arn. Bot. Beech. Voy. 317 = californicum.
exsertum, DC. Syst. i. 345 = rigidum.
fissum, Waldst. & Kit. Pl. Rar. Hung. i. 58. t. 51 = hybridum.
flammeum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 22.—Am. bor. occ.
flavum, *DC. Syst.* i. 346.—Assyria.
flavum, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 356 = deserti.
flexuosum, Bieb. Fl. Taur. Cauc. ii. 12 = elatum.
flexuosum, *Rafin. in Am. Monthly Mag.* (1811) 356, nomen ; *Rafin. Ann. Nat.* 12.—Am. bor.
foliosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 547.—Calif.
formosum, *Boiss. & Huet, Diagn.* Ser. II. v. 18.—Armenia.
Forskolii, Reichb. Illustr. Sp. Acon. v. t. 5 [63] = ranum.
Garumnae, Lapeyr. Suppl. Hist. Pl. Pyr. 73 = cardiopetalum.
glabellum, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) n. i. 56.—Sibir. batical.
glaciale, *Hook. f. & Thoms. Fl. Ind.* 58.—Reg. Himal.
glandulosum, *Boiss. & Huet, Diagn.* Ser. II. v. 11.—As. Min.
glaucum, *S. Wats. Bot. Calif.* ii. 427.—Calif.
gracile, *DC. Syst.* i. 347.—Hispan.
grandiflorum, Forsk. Fl. Aegypt. Arab. 212 = nanum.
grandiflorum, *Linn. Sp. Pl.* 531.—Sibiria.
gratum, Hoffmgg. Verz. Pfl. Nachtr. i. 192 = elatum.
halteratum, *Sibth. & Sm. Fl. Graec. Prod.* i. 372 ; *Fl. Graec.* t. 507.—Reg. Mediterr.
heliosgenicum, Boiss. in Ann. Sc. Nat. Sér. II. xvi. (1841) 366 = tomentosum.
Henderson, *Goualt, in Rev. Hortic.* Sér. IV. iii. (1854) 441.—Hab.?
hirsutum, Pers. Syn. ii. 82 = hybridum.
hispanicum, Reut. ex Willk. & Lange, Prod. Fl. Hisp. iii. 968 = Lescouii.
hispanicum, *Willk. l. c.* 969.—Hispan.
hispidum, Boiss. Fl. Orient. i. 82 = cappadocicum.
hispidum, Stev. ex DC. Syst. i. 358 = palmatifidum.
Hoffmeisteri, Klotzsch, Bot. Ergeb. Waldem. Reise, 132 = elatum.
Hohenackeri, *Boiss. Fl. Orient.* i. 85.—As. Min.
humulinum, *Basil. Hist. Pl.* i. 92.—China.
hybridum, Linn. in Amoen. Acad. iii. 37 ; Gouan, Illustr. 92 = elatum.
hybridum, *Steph. ex Willd. Sp. Pl.* ii. 1229.—Europ. ; Orient.
incanum, *Clarke, Trav.* ii. 451 = Aconiti.
incanum, *Royle, Illustr. Bot. Himal.* 55.—Reg. Himal.
incisum, Wall. Cat. n. 6717 = elatum.
intermedium, [Soland. in] Ait. Hort. Kew. ed. I. ii. 248 = elatum.
ithaburense, *Boiss. Diagn.* Ser. I. viii. 9.—Palestin.
Jacquemontianum, Cambess. in Jacquem. Voy. Bot. 6. t. 7 = Brunonianum.
junceum, DC. Fl. Fr. v. 641 = peregrinum.
Kotschyi, *T. Moore, Fl. & Pomol.* (1874) 78.—Hab.?
kiovense, Bess. ex Nym. Consp. 20 = elatum.
kurdicum, *Boiss. & Hohen. Diagn.* Ser. I. i. 67.—As. Min.
lanigerum, *Boiss. l. c.* viii. 10.—Persia.
latisepalum, *Hemsl. Diagn. Pl. Nov.* 17.—Mexic.
laxiflorum, *DC. Syst.* i. 360.—Sibiria.
lepidum, *Pisch. & Ave-Lall. Ind. Sem. Hort. Petrop.* ix. 70.—Reg. Caucas.
leptophyllum, *Hemsl. Diagn. Pl. Nov.* 18.—Mexic.
linarioides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 368.—Persia.
longipedunculatum, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 296.—Turkestan.
longipes, Moris, Fl. Sard. i. 59 = peregrinum.

DELPHINIUM:—
Losconii, *Costa, in Anal. Soc. Esp. Hist. Nat.* ii. (1873) 26.—Hispan.
Macsclannsm, *Regel, in Mém. Acad. Pétersb.* Sér. VII. iv. (1861) n. iv. 9.—Reg. Amur.
macrocentron, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 397.—Afr. trop.
macropetalum, DC. Syst. i. 350 = halteratum.
magnificum ×, Paxt. Mag. Bot. xvi. (1849) 258.
maritimum, Cav. ex Balb. Cat. Hort. Taur. (1813) 81 = Requienii.
mauritanicum, *Coss. in Bull. Soc. Bot. Fr.* xxvii. (1880) 68.—Afr. bor.
Maydellianum, *Trauto. in Act. Hort. Petrop.* vi. (1879) 7.—Rossia.
Menziesii, *DC. Syst.* i. 355.—Am. bor. occ.
Menziesii, Durand, in Journ. Acad. Philad. iii. (1855-58) 80 = decorum.
Menziesii, Durand, in Trans. Am. Phil. Soc. xi. (1860) 158 = bicolor.
Menziesii, Torr. & Gray, Fl. N. Am. i. 31 = simplex.
Menziesii, Torr. & Gray, in Pacif. Rail. Rep. ii. 119 = depauperatum.
mesoleucum, *Link, Enum. Hort. Berol.* ii. 80.—Hab. ?
micranthum, *Boiss. & Hohen. Diagn.* Ser. I. i. 68.—Kurdistan.
Middendorffii, *Trautv. in Middend. Reise Sibir.* i. *Bot.* (*Fl. Taimyr.*) 68.—Sibiria.
mixtum, Loisel. Fl. Gén. Fr. 46 = hybridum ?
monophyllum, Gilib. Fl. Lituan. ii. 387 = D. Consolida.
montanum, *DC. Fl. Fr.* v. 641.—Europ. austr.
montanum, DC. Syst. i. 358 = elatum.
moschatum, Munro, ex Hook. f. & Thoms. Fl. Ind. 58 = Brunonianum.
moschatum, Soiler. ex Nym. Consp. 20 = Requienii.
nanum, *DC. Syst.* i. 349.—Aegypt.
Nagellus, Balll. Dict. t. ad fasc. xviii = Aconitum Napellus.
neglectum, *Calla, Hort. Ripul. App.* ii. 346, nomen.—Hab.?
nevadense, *Kunze, in Flora*, xxix. (1846) 647.—Hispan. nudicaule, *Torr. & Gray, Fl. N. Am.* i. 38.—Am. bor. occ.
Nuttallianum, Pritz. ex Walp. Rep. ii. 744 = Menziesii.
obscordatum, *DC. Syst.* i. 350.—Afr. bor.
obscurum, Stev. ex DC. Prod. i. 56 = triste.
occidentale, *S. Wats. ex Coult. Man. Bot. Rocky Mount. Reg.* 11.—Am. bor. occ.
ochroleucum, Stev. ex DC. Syst. i. 353, 548 = hybridum.
officinale, *Wender, in Pharm. Centralbl.* (1836) 812, 813.—Europ. austr.
oliganthum, *Boiss. Fl. Orient.* i. 80.—Syria.
Oliverianum, *DC. Syst.* i. 341.—Assyria.
olopetalum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 364.—Persia.
orientale, *Y. Gay, in Desm. Catal. Dord.* 12.—Europ. ; Oriens.
orientale, Losc. in Trat. Pl. Arag. ii. 10 = Losconii.
ornatum, Bouché, in Bot. Zeit. i. (1848) 26 = orientale.
palmatifidum, *DC. Syst.* i. 358.—Sibiria.
palmatum, *Redde, in Beitr. Kenntn. Russ. Reich.* xxiii. 581.—Sibir. batical.
paniculatum, Host, Fl. Austr. ii. 65 = D. Consolida.
paradoxum, *Bunge, in Arbeit. Nat. Ver. Riga*, i. (1847) 134.—Turkestan.
parviflorum, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 75.—Sibiria.
patens, Benth. Pl. Hartw. 296 = decorum.
patens, Newberry, in Pacif. Rail. Rep. vi. 66 = Menziesii.
pauciflorum, D. Don, Prod. Fl. Nep. 196 = Ajacis.
pauciflorum, Nutt. ex Torr. & Gray, Fl. Bor. Am. i. 38, 661 = Menziesii.
pauciflorum, Reichb. ex Schlecht. in Linnaea, vi. (1831) 589 = Chamissonis.
pauciflorum, Royle, Illustr. Bot. Himal. 55 = denudatum.
pedatisectum, *Hemsl. Diagn. Pl. Nov.* 18.—Mexic.
pesicillatum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 369.—Arabia.
pesicillatum, Hook. f. & Thoms. Fl. Ind. 48 = unchiatum.
pentagynum, Boiss. Voy. Bot. Espagne, 13 = nevadense.
pentagynum, *Lam. Encyc.* ii. 564.—Lusitan. ; Afr. bor.
peregrinum, Delile, Fl. Aeg. Illustr. 65 = nanum.
peregrinum, Lam. Encyc. ii. 564 = halteratum.
peregrinum, *Linn. Sp. Pl.* 531.—Europ. austr. ; Orient.

DELPHINIUM:—
persicum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 363.—As. Min. ; Persia.
phrygium, *Boiss. l. c.* 363.—As. Min.
pictum, Willd. Enum. Hort. Berol. 574 = Requienii.
plenum, *Medic. Bot.* (1788) 87.—Hab. ?
Poltaratskii, *Rupr. Sert. Tianach.* 37.—As. centr.
pseudoperegrinum, *Visch. ex Sweet, Hort. Brit.* ed. II. 9, nomen.—Sibiria.
pubescens, *DC. Fl. Fr.* v. 641.—Gallia.
pubescens, Griseb. Spicil. Fl. Rumel. i. 219 = Ajacis.
pubescens, Henning, in Mém. Soc. Nat. Mosc. vi. (1853) 79, 89 = divaricata.
pubescens, Losc. in Trat. Pl. Arag. ii. 10 = Losconii.
pubiflorum, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 73.—Sibir. batical.
pulchrum ×, Van Houtte, Fl. des Serres, xvii. (1867-68) 157.
puniceum, Pall. Reise, iii. 736 = hybridum.
pusillum, *Labill. Ic. Pl. Syr. Dec.* iv. 3.—Syria.
pygmaeum, Petr. Encyc. Suppl. ii. 468 = pusillum.
Pylaei, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 307.—China.
pyramidale, Royle, Illustr. Bot. Himal. 56 = elatum.
pyrenaicum, Pourr. ex DC. Syst. i. 361 = montanum.
ramosissimum, Stev. ex Ledeb. Fl. Ross. i. 59 = divaricatum.
ranunculifolium, Wall. Cat. n. 6716 = elatum.
Raveyi, *Boiss. Diagn.* Ser. I. i. 66.—As. Min.
rectirsinum, Royle, Illustr. Bot. Himal. 56 = vestitum
Requienii, *DC. Fl. Fr.* v. 642.—Reg. Mediterr.
revolutum, *Desf. Tabl.* ed. II. 51.—Hab. ?
rigidum, *DC. Syst.* i. 344.—Syria.
rugulosum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 361.—Persia ; Turcoman.
sanlculaefolium, *Boiss. Diagn.* Ser. I. vi. 6.—Persia ; Afghanist.
Sarcophyllum, Hook. & Arn. Bot. Beech. Voy. 317 = nudicaule.
Savatieri, *Franch. in Bull. Soc. Linn. Par.* i. (1882) 330.—China.
schaiiforum, *D. Don, in Mem. Wern. Soc.* iii. (1821) 412.—Nepal.
scopsoum, *Greene, in Coult. Bot. Gaz.* vi. (1881) 156. —N. Mexic.
sclerocladum, Boiss. Diagn. Ser. I. viii. 8 = anthoroideum.
scopulorum, *A. Gray, Pl. Wright.* ii. 9.—N. Mexic.
segetum, Lam. Fl. Fr. iii. 595 = D. Consolida.
semibarbatum, *Bien. ex Boiss. Fl. Orient.* i. 92.—Persia.
simplex, *Dougl. in Hook. Fl. Bor. Am.* i. 25.—Am. bor. occ.
simplex, A. Gray, Pl. Wright. ii. 8 = scaureum.
simplex, Salisb. Prod. 375 = Ajacis.
sinense, Pisch. ex Link, Enum. Hort. Berol. ii. 80 = grandiflorum.
sparsiflorum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 307.—China.
speciosum, *Bieb. Fl. Taur. Cauc.* ii. 12.—Reg. Caucas. ; Persia.
speciosum, Jacka, ex Nym. Consp. 20 = elatum.
spurium, Fisch. ex Sweet, Hort. Brit. ed. II. 9 = speciosum.
Staphisagria, *Linn. Sp. Pl.* 531.—Reg. Mediterr.
staphyloides, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = scq.
stenocepalum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 278.—Kamtschatka.
Stockmaiium, *Boiss. Diagn.* Ser. II. i. 12.—Beluchistan.
sulcatum, *Reichb. ex Hoffmgg. Verz. Pfl. Nachtr.* i. 192, nomen.—Hab. ?
sulphureum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 81.—Syria.
sylvaticum, *Pomel, Nouv. Mat. Fl. Atl.* 332.—Algeria.
sylvaticum, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 74 = chellanthum.
Szowitzianum, *Boiss. Fl. Orient.* i. 89.—Reg. Caucas.
tauricum, Pall. ex Bieb. Fl. Taur. Cauc. ii. 12 = hybridum.
tchesnicum, *Boiss. Fl. Orient.* i. 88.—Persia.
tenuissimum, *Sibth. & Sm. Fl. Graec. Prod.* i. 370.—Greece.
thirkeanum, *Boiss. Fl. Orient.* i. 94.—Ad. Min.
tiroliense, *Kern. ex Dalla Torre, Ats. Alpenfl.* 67.—Tirol.

DELPHINIUM :—
tomentosum, *Auch. ex Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 866, partim.—Persia.
tomentosum, Boiss. l. c. pro parte = oliganthum.
tribracteolatum, *DC. Syst.* i. 350.—Afr. bor.
tricolor, *Bernh. ex Steud. Nom.* ed. II. i. 489, nomen. —Orient.
tricorne, *Michx. Fl. Bor. Am.* i. 314.—Am. bor.
tridactylum, Michx. l. c. = exaltatum.
trigonelloides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 866.—Persia.
trigynum, Schott, ex Willk. & Lange, Prod. Fl. Hisp. iii. 970 = peregrinum.
triste, *Ptach. ex DC. Syst.* i. 362.—Davuria ; Sibir.
trollifolium, *A. Gray, in Proc. Am. Acad.* viii. (1872) 275.—Am. bor. occ.
tuberosum, *Auch. ex Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 870.—Persia.
uliginosum, *Curran, in Bull. Calif. Acad.* i. III. (1885) 151.—Calif.
unciinatum, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 24. —Reg. Himal.
urceolatum, Jacq. Coll. i. 158 = exaltatum.
variegatum, *Torr. & Gray, Fl. N. Am.* i. 32.—Am. bor. occ.
velutinum, *Bertol. Excerpt.* 12.—Italia.
velutinum, Schur, Enum. Pl. Transs. 29 = fissum.
venulosum, *Boiss. Fl. Orient.* i. 56.—Cappadocia.
verdunense, Balb. Cat. Hort. Taur. 31 = cardiopetalum.
vericolor, Sallsb. Pred. 375 = D. Consolida.
vestitum, Boiss. Diagn. Ser. II. i. 13 = uncinatum.
vestitum, *Wall. Cat.* n. 4715.—Reg. Himal.
villosum, Stev. ex DC. Prod. i. 65 = elatum.
viminosum, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 374 = azureum.
virescens, Nutt. Gen. Am. ii. 14 = azureum.
virgatum, Jacq. f. ex Spreng. Syst. ii. 617 = grandiflorum.
virgatum, *Poir. Encyc. Suppl.* iii. 458.—Syria.
viscosum, *Hook. f. & Thoms. Fl. Ind.* 52.—Reg. Himal.
vitifolium, *Willd. ex Steud. Nom.* ed. II. i. 489.—Hab. ?
Wallzeri, *Engelm. in Wislis. Tour North Mexico*, 106.—Mexic. bor.

DELTOCARPUS, L'Hérit. ex DC. Syst. ii. 578 (1821) = **Myagrum**, Tourn. (Crucifer.).

DELUCIA, DC. Prod. v. 633 (1836) = **Bidens**, Linn. (Compos.).
ostruthioides, *DC. l. c.* = B. ostruthioides.

DEMATOPHYLLUM, Griseb. in Goett. Abh. xxiv. (1879) 74. *ZYGOPHYLLEAE.*
salsoloides, *Griseb. l. c.*—Reg. Argent.

DEMAXENBIA, Dum. Comm. Bot. 26 (1822). *GRAMINEAE*, Benth. & Hook. f. iii. 1194.
BRIZOPYRUM, Link, Hort. Berol. i. 159 (1827).
castellana, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* i. 113.—Hispan.
loliacea, *Nym. Syll.* 496.—Ins. Balear.
obliters, *Hemsl. Bot. Voy. Challenger*, i. 11. 90.—Ins. S. Helena.
sicula, *Dum. Comm. Bot.* 27.—Europ.

DEMETRIA, Lag. Gen. et Sp. Nov. 30 (1816) = **Grindelia**, Willd. (Compos.).
glutinosa, Lag. l. c. = G. glutinosa.
spathulata, Lag. l. c. = G. inuloides.

DEMIDIUM, DC. Prod. vi. 246 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 500.
filaginoum, *DC. l. c.* 247.—Madag.

DEMIDOFIA, Dennst. Schluss. Hort. Malab. 31 (1818) = **Carallia**, Roxb. (Rhizoph.).
nodosa, Dennst. l. c. = C. integerrima.

DEMIDOFIA, J. F. Gmel. Syst. 458 (1791) = **Dichrandra**, Forst. (Convolv.).
repens, J. F. Gmel. l. c. = Dich. repens.

DEMIDOVIA, Hoffm. Hort. Mosq. (1808) Introd. fol. 3 = **Fuchs**, Linn. (Liliac.).
palophylla, Hoffm. l. c. = P. incompleta.

DEMIDOVIA, Pall. Hort. Demid. 150. t. 1 (1781) = **Tetragonia**, Linn. (Ficoid.).
tetragonioides, Pall. l. c. = T. expansa.

DEMOCRITA, Vell. Fl. Flum. 176 (1825). (*INCERT. SED.*)
resinosa, *Vell. l. c.* 177 ; iv. t. 106.—Bras.

DEMOCRITEA, Doc. Prod. 540 (1830) = **Serissa**, Comm. (Rubiac.).
serissoides, DC. l. c. = S. Democritea.

DENCKEA, Rafin. in Med. Repos. N. York, v. (1808) 353 = **Gentiana**, Tourn.
crinita, Rafin. l. c. = G. crinita.

DENDRAGROSTIS, Nees, in Linnaea, ix. (1834) 467 = **Chusquea**, Kunth (Gramin.).
anelytkra, Nees, l. c. 491 = C. anelythra.
pinifolia, Nees, l. c. 490 = C. pinifolia.
tenella, Nees, l. c. 492 = C. tenella.

DENDRANTHEMA, Des Moul. in Actes Soc. Linn. Bord. xx. (1855) 561 = **Chrysanthemum**, Tourn. (Compos.).
indicum, Des Moul. l. c. 562 = C. indicum.
parthenioides, Des Moul. l. c. 561 = Anthemis parthenioides.
Parthenium, Des Moul. l. c. = C. Parthenium.
sinensis, Des Moul. l. c. 562 = C. sinense.

DENDREMA, Rafin. Sylva Tellur. 14 (1838) = **Corypha**, Linn. (Palm.).

DENDRIUM, Desv. Journ. Bot. i. (1813) 36 = **Lejophyllum**, Hedw. f. (Ericac.).
buxifolium, Desv. l. c. = L. buxifolium.

DENDROBIUM, *Sw.* in Nov. Act. Soc. Sc. Upsal vi. (1799) 82. *ORCHIDEAE*, Benth. & Hook. f. iii. 498.
ACLINIA, Griff. Notul. iii. 320 (1851).
APORUM, Blume, Bijdr. 324 (1825).
BOLBIDIUM, Lindl, Veg. Kingd. 181 (1847).
CALYPSO, Gaudich in Freyc Voy Bot. 422 (1826).
CALCATRIPPA, Heist. Syst. 8 (1748).
CALLISTA, Lour. Fl. Cochinch. 519 (1790).
CERAIA, Lour. l. c. 518 (1790).
CORIANDRIA, Fitzg. Austral. Orch. i. VII. t. 2 (1882).
DESMOTRICHUM, Blume, Bijdr. 320 (1825).
DICHOPUS, Blume, Mus. Bot. Lugd. Bat. ii. 176 (1856).
DITULIMA, Rafin. Fl. Tellur. iv. 41 (1836).
ENDISUS, Rafin. l. c. ii. 51 (1836).
FROSCULA, Rafin. l. c. iv. 44 (1836).
GRASTIDIUM, Blume, Bijdr. 338 (1825).
KERANTHUS, Lour. ex Endl. Gen. 198, in syn. (1836).
LATOURIA, Blume, Rumphia, iv. 41. t. 195, 199 (1850).
MACROSTOMIUM, Blume, Bijdr. 335 (1825).
ONYCHIUM, Blume, l. c. 323 (1825).
ORMOSTEMA, Rafin. Fl. Tellur. iv. 38 (1836).
OXYSTOPHYLLUM, Blume, Bijdr. 335 (1825).
PEDILONIUM, Blume, l. c. 320 (1825).
PIERARDIA, Rafin. Fl. Tellur. iv. 41 (1836).
SARCOPODIUM, Lindl. in Paxt. Flow. Gard. i. (1850) 154.
SARCOSTOMA, Blume, Bijdr. 339 (1825).
SCANDEDERIS, Thou. Orch. Iles Afr. tabl. iii. (1822).
SCHISMOCERAS, Presl, Rel. Haenk. i. 96. t. 18 (1827).
STACHYOBIUM, Reichb. f. in Gard. Chron. (1869) 785.
THELCHITON, Endl. Prod. Fl. Norf. 32 (1833).
THIGUANIA, Rafin. Fl. Tellur. iv. 47 (1836).
TROPILIS, Rafin. l. c. ii. 95 (1836).
acervosum, *Lindl. Bot. Reg.* (1841) *Misc.* 48.—Burma ; Ins. Malaya.
Achillis, Reichb. f. Otia Bot. Hamb. 55 = calcaratum.
aciculare, *Lindl. Bot. Reg.* (1840) *Misc.* 81.—Malaya.
acinaciforme, *Roxb. Hort. Beng.* 68 ; *Fl. Ind.* iii. 487. —Reg. Himal.
Aclinia, Lindl. in Journ. Linn. Soc. iii. (1859) 9 = incurvum.
Aclinia, Reichb. f. in Bonplandia, iv. (1856) 329 = D. Pseudaclinia.

DENDROBIUM :—
acrobaticum, *Reichb. f. in Gard. Chron.* (1871) 802.—Burma.
acuminatissimum, *Lindl. Gen. et Sp. Orch.* 86.—Java.
acuminatum, H. B. & K. Nov. Gen. et Sp. i. 357 = Pleurothallis acuminata.
Adae, *F. M. Bailey, in Proc. Roy. Soc. Queensl.* i (1884) 149.—Austral.
aduncum, *Lindl. Bot. Reg.* (1842) *Misc.* 58 ; (1846) t. 15.—Reg. Himal.
aemulum, *R. Br. Prod.* 333.—Austral.
affine, *Steud. Nom.* ed. II. i. 489 ; *Lindl. Bot. Reg* (1844) *Misc.* 64.—Ins. Timor.
aggregatum, H. B. & K. Nov. Gen. et Sp. i. 358 = Maxillaria aggregata, *Lindl.*
aggregatum, *Roxb. Hort. Beng.* 68 ; *Fl. Ind.* iii. 477.—Burma.
agrostophyllum, *F. Muell. Fragm.* viii. 28.—Austral.
Ainsworthii x, T. Moore, in Gard. Chron. (1874) i. 448.
alatum, Pers Syn. ii. 524 = Maxillaria alata.
albidulum, Thw. ex Trimen, in Journ. Bot. xxiii. (1885) 249 = D. Diodon.
albosanguineum, *Lindl. & Paxt. Flow. Gard.* ii. (1851-52) 98.—Burma.
album, Hook. Exot. Fl. t. 142 = Maxillaria alba.
album, Wight, Ic. t. 1645 = aqueum.
algarum, Reinw. ex Lindl. in Journ. Linn. Soc. iii. (1859) 42, nomen (Taeniophylli sp.).—Java.
alopecurum, Keichb. f. in Walp. Ann. vi. 979 = D. Serra.
aloideum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 48.—Mexic.
alpestre, *Royle, Illustr. Bot. Himal.* 370.—Reg. Himal.
alpestre, Sw. in Nov. Act. Nat. Soc. Sc. Upsal. vi (1799) 84 = Pleurothallis alpestris.
amblyomitis, *Reichb. f. in Gard. Chron.* (1878) i. 832. —N. Guin.
amboinense, *Hook. Bot. Mag.* t. 4987.—Ins. Amboin.
amethystoglossum, *Reichb. f. in Gard. Chron.* (1872) 109.—Ins. Philipp.
amoenum, *Wall. ex Lindl. Gen. et Sp. Orch.* 78.—Reg. Himal.
amplum, *Lindl. in Wall. Cat.* n. 2001.—Reg. Himal.
anceps, Sw. in *Vet. Acad. Handl. Stockh.* (1800) 246. —Ind. or. ; Burma.
Andersonii, J. Scott, in Journ. Agr. Hort. Soc. Ind. N. S. iii. (1872) 117 = dracontis.
angulatum, Lindl. Gen. et Sp. Orch. 76 = homonymum.
angulatum, *Wall. Cat.* n. 2010 —Burma.
angustifolium, *Lindl. Gen. et Sp. Orch.* 76.—Java.
annuligerum, *Reichb. f. in Gard. Chron.* (1871) 675.—‘ Marisa.
anosmum, Lindl. Bot. Reg. (1845) Misc. 32 = superbum.
Antelope, *Reichb. f. in Gard. Chron.* (1888) i. 658.—Ins. Molucc.
antennatum, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 286.—N. Guinea.
Aphrodite, *Reichb. f. in Bot. Zeit.* xx. (1862) 246.—Burma.
aphyllum, Roxb. Hort. Beng. 68, nomen ; et ex Lindl. Gen. et Sp. Orch. 78 = smoenum.
appendiculatum, *Lindl. Gen. et Sp. Orch.* 76.—Java.
aqueum, *Lindl. Bot. Reg.* (1848) *Misc.* 8.—Ind. or.
Arachnites, *Reichb. f. in Gard. Chron.* (1874) ii. 354. —Burma.
Arachnites, Thou. Orch. Iles Afr. t. 88 = Aëranthes Arachnitis.
arachnostachyum, *Reichb. f. in Gard. Chron.* (1877) i. 354.—N. Guinea.
articulatum, *Teijsm. & Binn. Cat. Hort. Bog.* 42, nomen.—Sumatra.
asphale, *Reichb. f. in Gard. Chron.* (1874) ii. 644. —Hab. ?
atractodes, *Ridl. in Journ. Bot.* xxiii. (1885) 128.—Siam.
atropurpureum, *Miq. Fl. Ind. Bat.* iii. 644.—Burma ; Ins. Malaya.
atrosanguineum, E. Morr. & De Vos, Ind. Bibliog. Hort. Belg. 166, sphalm. = albosanguineum.
attenuatum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 17. —Borneo.
aurantiacum, F. Muell. Fragm. vii. 98 = Bulbophyllum aurantiacum.
aureum, Lindl. Gen. et Sp. Orch. 77 = heterocarpum.
auriferum, *Lindl. l. c.* 88.—China.

DENDROBIUM:—
auroroseum, *Reichb. f. in Bonplandia*, v. (1857) 56.—Java.
Halleyi, *F. Muell. Fragm.* viii. 178.—Austral.
Bairdianum, *F. M. Bailey, Suppl. Syn. Queensl. Fl.* (1884), *Ade F. Muell. Census, App.* ii. 3.—Austral.
bambusæfolium, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 149.—Burma.
barbatulum, Batem. in Bot. Mag. t. 5444 = Eytchenarum.
barbatulum, *Lindl. in Wall. Cat.* n. 2013.—Ind. or.
barbatulum, Wight, Ic iii. t. 910 = chlorops.
Barringtoniæ, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 82 = Lycaste Barringtoniæ.
Beckleri, F. Muell. Fragm. v. 95 = Mortii.
Bensoniæ, *Reichb. f. in Bot. Zeit.* xxv. (1867) 280.—Burma.
bicameratum, *Lindl. Bot. Reg.* (1839) *Misc.* 59.—Reg. Himal.
bicaudatum, *Reinw. ex Lindl. in Journ. Linn. Soc.* iii. (1859) 90.—Java.
bicolor, Lindl. Gen. et Sp. Orch. 90 = Eria Lindleyi.
bicolor, Pers. Syn. ii. 524 = Maxillaria bicolor.
bifalce, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 237.—N. Guinea.
bifarium, *Lindl. in Wall. Cat.* n. 2002.—Malaya.
bifforum, A. Rich. Ess. Fl. N. Zél. 167 = Cunninghamii.
bifforum, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 84.—Ins. Societ.
bigibbum, *Lindl. in Paxt. Flow. Gard.* iii. (1852) 25. t. 245.—Austral.
bilobum, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 236.—N. Guinea.
Binnendijkii, *Reichb. f. Xenia Orch.* ii. 74.—Java.
binoculare, *Reichb. f. in Gard. Chron.* (1869) 785.—Ind. or.
Blumii, *Lindl. Gen. et Sp. Orch.* 86.—Java.
Bolbophylli, Griff. It. Notes, 65 = Bulbophyllum Griffithii.
Boothii, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 318.—Java.
bostrychodes, *Reichb. f. in Gard. Chron.* (1880) ii. 748.—Borneo.
Bowmanii, *Benth. Fl. Austral.* vi. 286.—Austral.
Boxallii, Reichb. f. in Gard. Chron. (1874) i. 315 = gratiosissimum.
braccatum, Lindl. Gen. et Sp. Orch. 76 = Eria bractata.
brachycarpum, *A. Rich. Tent. Fl. Abyss.* ii. 282.—Abyss.
brachypetalum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 6.—Java.
brachypus, *Reichb. f. in Linnæa*, xli. (1877) 42.—Ins. Norfolk.
bractesum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 48.—Mexic.
brevifforum, Lindl. in Journ. Linn. Soc. iii. (1859) 14 = bicameratum.
brevifolium, Hort. ex Lindl. l. c. 12 = pulchellum.
brisbanense, Reichb. f. in Walp. Ann. vi. 299 = gracilicaule.
Bryceanum, *Reichb. f. in Gard. Chron.* (1876) ii. 328.—Burma.
Bullenianum, *Reichb. f. in Bot. Zeit.* xx. (1862) 214.—Ins. Philipp.
Bullerianum, Batem. in Bot. Mag. t. 5662 = gratiosissimum.
Burbidgei, *Reichb. f. in Gard. Chron.* (1876) ii. 800.—Ins. Sondlac.
bursigerum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 16.—Ins. Philipp.
calamiformum, Lodd. ex Lindl. Bot. Reg. (1841) Misc. 9 = teretifolium.
calcaratum, Lindl. l. c. (1840) Misc. 89 = cœnostalyx.
calcaratum, *A. Rich. Sert. Astrol.* 18. t. 7.—Ins. Vanikoro.
Calceolaria, *Carey, ex Hook. Exot. Fl.* 184.—Reg. Himal.
calceolum, Roxb. Hort. Beng. 68; Fl. Ind. iii. 488 = Roxburghii.
Caleyi, *A. Cunn. in Hook. Comp. Bot. Mag.* ii. (1836) 277.—Austral.?
calophyllum, *Reichb f. Xenia Orch.* ii. 167.—Malaya.
calopogon, *Reichb. f. l. c.* 23.—Malaya.
Cambridgeanum, *Reichb. f. in Linnæa*, xli. (1877) 89.—N. Caled.
DENDROBIUM:—
cambridgeanum, Paxt. Mag. Bot. vi. (1839) 265 = ochreatum.
canaliculatum, *R. Br. Prod.* 338.—Austral.
candidum, *Wall. ex Lindl. in Bot. Reg.* (1838) *Misc.* 36.—Reg. Himal.
capillipes, *Reichb. f. in Gard. Chron.* (1867) 997.—Burma.
carinatum, *Willd. Sp. Pl.* iv. 138.—Ins. Luzon.
carinianum, *Reichb. f. in Gard. Chron.* (1869) 611.—Reg. Himal.
carnosum, *Presl, Rel. Haenk.* i. 102.—Peruv.
carnosum, Reichb. f. in Walp. Ann. vi. 280 = atropurpureum.
carnosum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1858) 459.—Java.
caryochodes, A. Cunn. in Lindl. Bot. Reg. sub t. 1828 = Calcola casythiodes.
castum, Batem. ex Reichb. f. in Gard. Chron. (1868) 948 = moniliforme.
catenatum, *Lindl. Gen. et Sp. Orch.* 84.—Japon.
catillare, *Reichb. f. in Seem. Fl. Vit.* 304.—Ins. Fiji.
caudatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 315.—Sumatra.
Cerasia, *Lindl. Gen. et Sp. Orch.* 89.—China; Cochinch.
cerealis, *Reichb. f. in Linnæa*, xli. (1877) 40.—Ins. Philipp.
cerinum, *Reichb. f. in Gard. Chron.* (1879) ii. 554.—Malaya.
Chalmersii, *F. Muell. in Wing, South. Sc. Record*, ii. (1885).—N. Guinea.
chlorops, *Lindl. Bot. Reg.* (1844) *Misc.* 44.—Ind. or.
chloropterum, *Reichb. f. & S. Moore, in Journ. Bot.* xvi. (1878) 137.—Hab.?
Christianum, *Reichb. f. in Gard. Chron.* (1882) i. 173.—Siam.
chrysanthum, *Wall. Cat.* n. 2012.—Reg. Himal.
chrysocrepis, *Rev. & Reichb. f. ex Hook. f. Bot. Mag.* t. 6007.—Burma.
chrysotis, Reichb. f. in Gard. Chron. (1870) 1311 = Hookerianum.
chrysotoxum, *Lindl. Bot. Reg.* (1847) sub t. 19.—Burma.
ciliatum, *Parish, ex Hook. Bot. Mag.* t. 5480.—Burma.
ciliatum, Pers. Syn. ii. 523 = Maxillaria ciliata.
cincinnatum, *F. Muell. in Proc. Roy. Soc. Queensl.* i. (1884) 113.—N. Guinea.
cinnabarinum, *Reichb. f. in Gard. Chron.* (1880) ii. 166.—Borneo.
clavatum, Roxb. Hort. Beng. 68; Fl. Ind. iii. 481 = densiflorum.
clavatum, *Wall. Cat.* n. 2004.—Reg. Himal.
Closterium, *Reichb. f. in Linnæa*, xli. (1877) 92.—N. Caled.
Cobbianum, *Reichb. f. in Gard. Chron.* (1881) ii. 780.—Hab.?
Coelogyne, *Reichb. f. l. c.* (1871) 136.—Burma.
coerulescens, Wall. ex Lindl. Sert. Orch. t. 18 = crobile.
comatum, *Lindl. Gen. et Sp. Orch.* 76.—Java.
complanatum, *A. Cunn. in Lindl. Bot. Reg.* (1839) *Misc.* 54.—Australi.
compressum, Lindl. Bot. Reg. (1842) Misc. 63 = lamellatum.
concinnum, *Miq. Fl. Ind. Bat.* iii. 644.—Malaya.
conostalum, *Lindl. Gen. et Sp. Orch.* 89.—Java.
conostalix, *Reichb. f. in Walp. Ann.* vi. 292.—Malaya.
convexum, *Lindl. Gen. et Sp. Orch.* 76.—Java.
corniculatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 83 = Pleurothallis corniculata.
crassinode, Benson & Reichb. f. in Gard. Chron. (1869) 664; et in Flora, lii. (1869) 73 = pendulum.
crassilasiolum, *A. Cunn. in Lindl. Bot. Reg.* (1839) *Misc.* 32.—Austral.
crepidatum, *Griff. Notul.* iii. 319 (sp. dub.).—Ind. or.
crepidatum, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 68.—Ind. or.
crotaceum, *Lindl. Bot. Reg.* (1847) t. 62.—Reg. Himal.
criniferum, *Lindl. l. c.* (1844) *Misc.* 41.—N. Guinea.
crispatum, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 86.—Ins. Societ.
crispum, Dals. in Hook. Kew Journ. iv. (1852) 111 = microbulbon.
cruentum, *Reichb. f. in Gard. Chron.* (1884) i. 604.—Malaya.
DENDROBIUM:—
crumenatum, *Sw. in Schrad. Journ.* ii. (1799) 237.—Ind. or.; China; Malaya.
crystallinum, *Reichb. f. in Gard. Chron.* (1868) 572.—Reg. Himal.; Burma.
cucullatum, R. Br. in Lindl. Bot. Reg. t. 548 = Pierardi.
cucumerinum, *W. McLeay, in Lindl. l. c.* (1842) *Misc.* 58.—Austral.
cultriforme, Thou. Orch. Isles Afr. t. 87 = Polystachya cultrata.
cumulatum, *Lindl. in Gard. Chron.* (1855) 764.—Burma.
Cunninghamii, *Lindl. Bot. Reg.* sub t. 1756.—N. Zel.
Cunninghamii, Steud. Nom. ed. II. i. 490 = Caleyi.
cupreum, Herb. ex Lindl. Bot. Reg. t. 1779 = Calceolaria.
Curtisii, *Reichb. f. in Gard. Chron.* (1881) ii. 102.—Borneo.
cuspidatum, Lindl. in Journ. Linn. Soc. iii. (1859) 4 = Nathanielis.
cuspidatum, *Lindl. in Wall. Cat.* n. 2015.—Burma.
cymbidioides, *Lindl. Gen. et Sp. Orch.* 77.—Java.
dactyliferum, *Reichb. f. in Gard. Chron.* (1884) i. 638.—Hab.?
dactyloides, *Reichb. f. in Journ. Bot.* xv. (1877) 182.—Ins. Samoa.
D'Albertisii, *Reichb. f. in Gard. Chron.* (1878) i. 366.—N. Guinea.
Dalessertianum, Paxt. Mag. Bot. xi. (1844) 145 = pulchellum.
Dalzellii, Hook. Kew Journ. iv. (1852) 292 = Eria Dalzellii.
Dayanum, *Boxall, ex Nawes, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 284, nomen.—Ins. Philipp.
Deasei, *Reichb. f. in Gard. Chron.* (1882) ii. 361.—Ins. Philipp.
densiflorum, D. Don, Prod. Fl. Nep. 34 = Panisea parviflora.
densiflorum, *Wall. Cat.* n. 2000.—Reg. Himal.
denudans, *D. Don, Prod. Fl. Nep.* 34.—Reg. Himal
Deplanchei, *Reichb. f. in Linnæa*, xli. (1877) 90.—N Caled.
Devonianum, *Paxt. Mag. Bot.* vii. (1840) 169.—Reg. Himal.
dicuphum, *F. Muell. Fragm.* viii. 28.—Austral.
Diodon, *Reichb. f. in Linnæa*, xli. (1877) 89.—Zeylan.
discolor, Lindl. Bot. Reg. (1841) t. 52, Misc. 21 = undulatum.
distachyon, *Lindl. in Journ. Linn. Soc.* iii. (1859) 13.—Borneo.
distichum, *Reichb. f. in Linnæa*, xli. (1877) 89.—Ins. Philipp.
dixanthum, *Reichb. f. in Gard. Chron.* (1865) 674.—Burma.
Dominyanum ×, Reichb. f. l. c. (1878) i. 202.
Dracosia, *Reichb. f. in Bot. Zeit.* xx. (1862) 214.—Burma.
eburneum, Reichb. f. ex Batem. in Bot. Mag. t. 5459 = Draconis.
Egertonianum, Lindl. Bot. Reg. (1847) sub t. 36 = amœnum.
elegans, H. B. & K. Nov. Gen. et Sp. i. 253 = Pleurothallis elegans.
elongatum, A. Cunn. in Lindl. Bot. Reg. (1839) Misc. 33 = gracilicaule.
elongatum, *Lindl. Gen. et Sp. Orch.* 77.—Java.
erianthum, *Griff. Notul.* iii. 316.—Reg. Himal.
erosum, *Lindl. Gen. et Sp. Orch.* 86.—Java.
erythropogon, *Reichb. f. in Gard. Chron.* (1885) ii. 196.—Ins. Sondaic.
erythrocaulon, *Reichb. f. l. c.* (1874) ii. 162.—Ins. Philipp.
esuriens, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 15.—Java.
eulophetum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 5.—Burma; Java.
euosmum ×, Reichb. f. in Gard. Chron. (1868) i. 174.
euphlebium, *Reichb. f. ex Lindl. in Journ. Linn. Soc.* iii. (1859) 7.—Java.
excavatum, *Miq. Fl. Ind. Bat.* iii. 644.—Java.
excisum, Lindl. Bot. Reg. (1841) Misc. 77 = bifarium.
exilipinum, *F. Muell.* Fragm. v. 95 = Bulbophyllum exiguum.

DENDROBIUM:—
exsculptum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 216.—Java.
extinctorium, Lindl. Bot. Reg. sub t. 1756 = Eria extinctoria.
Fairfaxii, F. Muell. in Sydney Mail, Sept. 21 (1872) 360; ex Benth. Fl. Austral. vi. 355 = tereticaule.
falcorostre, Fitzg. Austral. Orch. i. v. t. 4 = acq.
Falconeri, *Hook. Bot. Mag.* t. 4944.—Reg. Himal.
falcorostrum, *Fitzg. in Sydney Morning Herald*, Nov. 18 (1876); *Fitzg. Austral. Orch.* i. v. (1879) t. 4. —Austral.
Farmerii, *Pant. Mag. Bot.* xv. (1849) 241.—Reg. Himal.
Fellowsii, *F. Muell. Fragm.* vii. 63.—Austral.
ferox, Hask. Retzia, i. 1 = macrophyllum.
filiforme, Wight, Ic. t. 1845 = Eria Dalzellii.
filiiclonum, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 97.—Ins. Samoa.
fimbriatum, Dalz. in Hook. Kew Journ. iv. (1852) 292 = Eria Dalzellii.
fimbriatum, *Hook. Exot. Fl.* t. 71.—Reg. Himal.
fimbriatum, Lindl. Gen. et Sp. Orch. 76 = plicatile.
Findleyanum, *Parl. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 149.—Burma.
firmum, *Steud. Nom.* ed. II. i. 490.—Java.
Fitzgeraldi, *F. Muell. in Melb. Chemist*, (June 1884). —Austral.
Flabellum, Reichb. f. in Bonplandia, v. (1857) 86 = Macraei.
flavescens, *Lindl. Gen. et Sp. Orch.* 85.—Java.
flavum, *Roxb. Hort. Beng.* 68, nomen.—Ind. or.
flexuosum, Griff. Notul. iii. 317 = longicornu.
floribundum, *D. Don, Prod. Fl. Nep.* 34.—Nepal.
floribundum, *Reichb. f. in Gard. Chron.* (1875) ii. 772. —N. Hebrid.
Foelschei, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 280.—Austral.
foliosum, Jumelle in *Duperr. Voy. Coq. Bot.* 208. t. 41.—Ins. Amboin.
formosum, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 485.— Reg. Himal.
fugax, *Reichb. f. in Gard. Chron.* (1871) 1257 — Ind. or.
funiforme, *Blume, Rumphia*, iv. 40.—N. Guinea.
furcatum, *Renw. ex Lindl. in Journ. Linn. Soc.* iii. (1859) 18.—Ins. Celebes.
fuscatum, Lindl. l. c. 8 = Gibsonii.
fuscescens, *Griff. Notul.* iii. 308.—Reg. Himal.
fuscum, *Fitzg. in Gard. Chron.* (1879) ii. 680.— Austral.
fusiforme, *F. M. Bailey, in Proc. Linn. Soc. N. S. W.* ii (1878) 277.—Austral.
fustiforme, Thou. Orch. Isles Afr. t. 86 = Polystachya fusiformis.
Fytchianum, *Batem in Gard. Chron.* (1864) 100.— Burma.
galeatum, Sw. in Schrad. Neues Journ. i. (1805) 96 = Bulbophyllum galeatum.
gemellum, *Lindl. Gen. et Sp. Orch.* 81.—Java.
geminatum, *Lindl. l. c.* 77.—Java.
Gibsonii, *Paxt. Mag. Bot.* v. (1838) 169.—Reg. Himal.
glaucophyllum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 314.—Java.
glaucum, *Teijsm. & Binn. Cat. Hort. Bog.* 43.—Ins. Amboin.
glossotis, *Reichb. f. Otia Bot. Hamb.* 55.—Ins. Tahiti.
Goldfinchii, *F. Muell. in Wing, South Sc. Record*, (Jan. 1883).—Ins. Solomon.
Goldiei, Reichb. f. in Gard. Chron. (1878) i. 653 = superbiens.
Gordoni, *S. Moore, in Journ. Linn. Soc.* xx. (1883) 373.—Ins. Fiji.
Gouldii, *Reichb. f. in Gard. Chron.* (1867) 901.—Ins. Pacif.
gracile, *Lindl. Gen. et Sp. Orch.* 91.—Java.
gracilicaule, *F. Muell. Fragm.* i. 179.—Ins. Dom. Howe.
graminifolium, *Wight, Ic.* t. 1648.—Ind. or.
graminifolium, Willd. Sp. Pl. iv. 135 = Ottomaria graminifolia.
grandiflorum, H. B. & K. Nov. Gen. et Sp. i. 259. t. 84 = Maxillaria grandiflora.
grandiflorum, *Lindl. Gen. et Sp. Orch.* 77.—Java.
grandiflorum, Reinw. ex Lindl. Fol. Orch. Spec. 4 = Bulbophyllum Reinwardtii.
gratiosissimum, *Reichb. f. in Bot. Zeit.* xxiii. (1865) 90. —Burma.

DENDROBIUM:—
Griffithianum, *Lindl. Bot. Reg.* sub t. 1756.—Burma.
Guibertii, Carr. in Rev. Hortic. (1876) 481 = Griffithianum.
haemoglossum, *Thw. Enum. Pl. Zeyl.* 429.—Ind. or.
Haenkeanum, *Steud. Nom.* ed. II. i. 490.—Ins. Philipp.
Hamburyanum, Reichb. f. in Bonplandia, iv. (1856) 329 = lituiflorum.
Harrisoniae, Hook. Exot. Fl. t. 120 = Maxillaria Harrisoniae.
Harveyanum, *Reichb. f. in Gard. Chron.* (1888) i. 624. —Burma.
Hasseltii, *Lindl. Gen. et Sp. Orch.* 87.—Java.
Hasseltii, *Reichb. f. Xenia Orch.* ii.—Java?
hastatum, Pers. Syn. ii. 524 = Maxillaria hastata.
Hedyarnum, Batem. ex Hook. f. Bot. Mag. t. 5515 = scabrilingue.
Henshallii, Reichb. f. in Bonplandia, iii. (1855) 226 = transparens.
herbaceum, *Lindl. Bot. Reg.* (1840) *Misc.* 69.— Ind. or.
heterocarpum, *Wall. ex Lindl. Gen. et Sp. Orch.* 78. —Reg. Himal.
heteroglossum, *Blume, Rumphia*, iv. 40.—N. Guinea.
heterostigma, *Reichb. f. in Hamb. Gartenz.* xv. (1859) 51.—Java.
Hexadesmia, *Reichb. f. in Gard. Chron.* (1869) 710.— Hab.?
Heyneanum, *Lindl. in Wall. Cat.* n. 1995.—Ind. or.
Hillii, Hook. Bot. Mag. t. 5361 = speciosum.
Hillii, F. Muell. Fragm. i. 88 = Sarcochilus Hillii.
hirsutum, Griff. Notul. iii. 318 = longicornu.
hispidum, *A. Rich. Sert. Astrol.* 13.—Ins. Vanikoro.
homoglossum, *Steud. Nom.* ed. II. i. 490.—Java.
Hookerianum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 8.—Reg. Himal.
Horne, *S. Moore, l. c.* xx. (1883) 378.—Ins. Fiji.
Hughii, *Reichb. f. in Gard. Chron.* (1862) i. 764.— Singapore.
humile, Sm. in Rees, Cyclop. xi. n. 4 = Coelogyne humilis.
humile, Wight, Ic. t. 1643 = microbolbon.
Huttoni, *Reichb. f. in Gard. Chron.* (1869) 686.—Ins. Malay.
hymenanthum, *Reichb. f. in Bonplandia*, iii. (1855) 222.—Ins. Philipp.
Hymenophyllum, *Lindl. Gen. et Sp. Orch.* 86.—Java.
Incrassatum, *Miq. Fl. Ind. Bat.* iii. 581.—Java.
incurvum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 18. —Ind. or.
indivisum, *Miq. Fl. Ind. Bat.* iii. 620.—Malaya.
inflexum, *Teijsm. & Binn. Cat. Hort. Bog.* 43, nomen. —Java.
Infundibulum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 16.—Burma.
insigne, *Reichb. f. ex Miq. Fl. Ind. Bat.* iii. 640.—N. Guinea.
insulare, Steud. Nom. ed. II. ii. 489 = plicatile.
intermedium, *Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind.* v. (1853) 490.—Java.
involutum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 15. —Ins. Societ.
ionopus, *Reichb. f. in Gard. Chron.* (1882) ii. 808.— Burma.
Jamesianum, *Reichb. f. l. c.* (1869) 854.—Ind. or.
Japonicum, Lindl. Gen. et Sp. Orch. 89 = moniliforme.
javanicum, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 46 = Piccoglottis acuminata.
javanicum, Miq. Fl. Ind. Bat. iii. 644 = D. Sarcostoma.
javanicum, *Sw. in Schrad. Neues Journ.* i. 1. (1805) 96.—Java.
Jenkinsii, Lindl. in Journ. Linn. Soc. iii. (1859) 4, lapsu = parciflorum.
Jenkinsii, Wall. in Bot. Reg. (1839) t. 37 = aggregatum.
Jerdonianum, *Wight, Ic.* t. 1644.—Ind. or.
Joconum, *Reichb. f. in Linnaea*, xli. (1877) 91.—N. Caled.
Johannis, *Reichb. f. in Gard. Chron.* (1865) 890.— Austral.
Johnsoniae, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 95.—N. Guinea.
junceum, *Lindl. Bot. Reg.* (1842) *Misc.* 9.—Malaya.
Kefordii, *F. M. Bailey, in Proc. R. Soc. Queensl.* i. (1884) 11.—Austral.

DENDROBIUM:—
Kingianum, *Bidwill, ex Lindl. Bot. Reg.* (1844) *Misc.* 11.—Austral.
Kohlmeyerianum, *Teijsm. & Binn. ex Miq. Choix, Pl Yard. Buitens.* t. 24. f. 1.—Java.
Kuhlii, *Lindl. Gen. et Sp. Orch.* 87.—Java.
labuanum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 6. —Borneo.
lamellatum, *Lindl. Gen. et Sp. Orch.* 89.—Burma; Ins. Malay.
lancoranense, *Reichb. f. in Bonplandia*, v. (1857) 86. —Java.
Lancola, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 83; et Fl. Ind. Occ. iii. 1539 = Pleurothallis Lanceola.
lanceolatum, *Gaudich. in Freyc. Voy. Bot.* 423.—Ins. Moluce.
lancifolium, *A. Rich. Sert. Astrol.* 20. t. 8.—Ins. Moluce.
lasioglossum, *Reichb. f. in Gard. Chron.* (1868) 682. —Burma.
latifolium, H. B. & K. Nov. Gen. et Sp. i. 360 = Maxillaria latifolia.
Lawesium, Lindl. in Journ. Linn. Soc. iii. (1859) 10 = crepidatum.
Lawesii, *F. Muell. in Melb. Chemist* (June 1884). —N. Guinea.
laxum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 84 = Pleurothallis laxa, Stelis alba.
Leechianum x, Reichb. f. in Gard. Chron. (1882) i. 256.
Leonis, *Reichb. f. in Walp. Ann.* vi. 280.—Malaya.
leopardinum, Wall. Tent. Fl. Nep. 39. t. 28 = Bulbophyllum leopardinum.
Lesconii, *Calman, in Trans. N. Z. Inst.* xv. (1883) 326.—N. Zel.
leucochilorum, *Reichb. f. in Gard. Chron.* (1879) i. 202. —Burma.
leucolophotum, *Reichb. f. l. c.* (1882) ii. 552.—Archip. Mal.
Lichenastrum, F. Muell. Fragm. vii. 60 = Bulbophyllum Lichenastrum.
ligulatum, Pers. Syn. ii. 524 = Maxillaria ligulata.
lilacinum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii (1864) 18.—Java.
Linawianum, *Reichb. f. in Walp. Ann.* vi. 284.— China; Japon.
Lindleyanum, Griff. Notul. iii. 309 = nobile.
Lindleyi, Steud. Nom. ed. II. i. 490 = aggregatum.
linearifolium, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 818.—Sumatra.
lineatum, *Teijsm. & Binn. Cat. Hort. Bog.* 43, nomen. —Java.
linguaeforme, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 247.—Austral.; Ins. Pacif.
Lingnelle, *Reichb. f. in Gard. Chron.* (1882) ii. 552.— Ins. Malay.
linguiforme, Sm. Exot. Bot. i. 19. t. 11 = linguaeforme.
lituiflorum, *Lindl. in Gard. Chron.* (1856) 873.—Burma.
lividum, *Teijsm. & Binn. Cat. Hort. Bog.* 43, nomen. —Siam.
lobatum, *Miq. Fl. Ind. Bat.* iii. 631.—Java.
Lobbii, *Lindl. in Journ. Linn. Soc.* iii. (1859) 3.— Borneo.
Lobbii, Teijsm. & Binn. in Tijdschr. Nederl. Ind. v. (1853) 491 = Teysmanni.
longicolle, *Lindl. Bot. Reg.* (1840) *Misc.* 74.—Malaya.
longicornu, *Lindl. in Wall. Cat.* n. 1997.—Reg. Himal.
longifolium, H. B. & K. Nov. Gen. et Sp. i. 360 = Maxillaria longifolia.
longipetalum, Pers. Syn. ii. 523 = Maxillaria longipetala.
Lowii, *Lindl. in Gard. Chron.* (1861) 1046.—Borneo.
Lobbianum, *Reichb. f. l. c.* (1882) i. 400.—Burma.
lucens, *Reichb. f. in Bot. Zeit.* xxii. (1863) 138.— Borneo.
lunatum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 4.— Ins. Philipp.
luteolum, *Batem. in Gard. Chron.* (1864) 269.—Burma.
luscense, *Lindl. Bot. Reg.* (1844) *Misc.* 54.—Ins. Philipp.
lycopodioides, *Lindl. in Journ. Linn. Soc.* iii. (1859) 18.—Borneo.
MacCarthiae, [*Thw. in*] *Bot. Mag.* t. 4886.—Zeylan.
Macfarlanei, *F. Muell. Papuan Fl.* i. 29.—N. Guinea.
Macfarlanei, Reichb. f. in Gard. Chron. (1882) ii. 529 = Johnsoniae.
Macraei, *Lindl. Gen. et Sp. Orch.* 75.—Ind. or.; Ins. Malay.

DENDROBIUM:—

macranthum, De Vriese, ex Lindl. in Journ. Linn. Soc. iii. (1859) 59 = Eria elongata!

macranthum, Mig. Fl. Ind. Bat. iii. 649 = superbum.

macranthum, *A. Rich. Sert. Astrol.* 22.—N. Guinea.

macropus, *Benth. & Hook. f. Gen.* iii. 500.—Ins. Norfolk.

macrostachyum, *Lindl. Gen. et Sp. Orch.* 78.—Ind. or.

maculatum, H. B. & K. Nov. Gen. et Sp. i. 359 = Maxillaria maculata.

manilense, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 430.—Manilla.

marginatum, Batem. ex Hook. Bot. Mag. t. 5454 = xanthophlebium.

marginatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1853) 490.—Java.

marmoratum, *Reichb. f. in Gard. Chron.* iii. (1875) 492.—Burma.

mesochlorum, Lindl. Bot. Reg. (1847) sub t. 36 = amoenum.

metachilium, *Reichb. f. in Bonplandia*, iii. (1855) 222.—Ind. or.

mexicanum, Presl, Symb. Bot. i. 86. t. 25 (= Govenise sp.).—Mexico.

micans v, Retchb. f. in Gard. Chron. (1879) i. 332.

micranthum, Lindl. in Journ. Linn. Soc. iii. (1859) 3 = D. Serra.

microbulbon, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 84 = Stelis micrantha.

microbulbon, *A. Rich. in Ann. Sc. Nat.* Sér. II. xv. (1841) 19. t. 5.—Ind. or.

microchilos, Dals. in Hook. Kew Journ. iii. (1851) 345 = Eria Dalzelli.

micrglaphys, *Reichb. f. in Gard. Chron.* (1868) 1014.—Borneo.

Milligani, F. Muell. Fragm. i. 88. t. 6 = striolatum.

minax, *Reichb. f. Xenia Orch.* ii. 129. t. 145.—Ins. Amboina.

minutissimum, *F. Muell. Fragm.* v. 95.—Austral.

Mirbelianum, *Gaudich. in Freyc. Voy. Bot.* 423.—Ins. Moluc.

miserum, *Reichb. f. in Gard. Chron.* (1869) 388.—Assam.

modestum, *Reichb. f. in Bonplandia*, iii. (1855) 232. —Ins. Philipp.

Mohlianum, *Reichb. f. in Bot. Zeit.* xx. (1862) 214.—Ins. Fiji.

monanthum, *Teijsm. & Binn. Cat. Hort. Bog.* 43, nomen.—Sumatra.

moniliforme, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 85.—Japon.

monophyllum, *F. Muell. Fragm.* i. 189.—Austral.

Mooreanum, *Lindl. in Journ. Hort. Soc.* vi. (1851) 272.—Ins. 'Aniteuse.'

Mooreï, *F. Muell. Fragm.* vii. 26.—Ins. Dom. Howe.

Mortii, *F. Muell. l. c.* i. 214.—Austral.

Mortii, F. Muell. l. c. ii. 93 = Bowmanii.

moschatum, Griff. Notul. iii. 311 = pulchellum.

moschatum, *Sw. in Schrad. Neues Journ.* i. (1806) 94.—Burma.

moschatum, Wall. ex D. Don, Prod. Fl. Nep. 34 = D. Calceolaria.

mucronistum, *Planch. Hort. Donat.* 156.—Ind. or.?

multicaule, Teijsm. & Binn. Cat. Hort. Bog. 43, sphalm. = *musicola*.

multiflorum, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 149 = Nathaniella.

musicicola, Lindl. in Wall. Cat. n. 2017 = Eria musicicola.

mutabile, *Lindl. Gen. et Sp. Orch.* 86.—Java.

Myosurus, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 82 = Oberonia Myosurus.

Nathaniella, *Reichb. f. in Schill. Cat. Orch.* ed. III. (1857) 26.—Burma.

nitidissimum, *Reichb. f. in Journ. Linn. Soc.* xv. (1876) 112.—Ins. Admiralt.

nobile, *Lindl. Gen. et Sp. Orch.* 79.—Reg. Himal.; China.

nodatum, Lindl. in Gard. Chron. (1862) 717 = D. Aphrodite.

nodosum, Dals. in Hook. Kew Journ. iv. (1852) 292 = Macraei.

normale, *Falc. in Proc. Linn. Soc.* i. (1859) 14.—Reg. Himal.

DENDROBIUM:—

nudum, *Lindl. Gen. et Sp. Orch.* 85.—Java.

nummularifolium, R. King, in Gard. Chron. (1879) i. 789 = Bulbophyllum minutissimum.

nutans, *Lindl. Gen. et Sp. Orch.* 90.—Zeylan.

nutans, Presl, Symb. Bot. i. 84. t. 23 = Haenkeanum.

obtusa, Reichb. in Walp. Ann. vi. 1150, ind. lapsu, vide *Dendrocolla obtusa*.

ochreatum, *Lindl. in Wall. Cat.* n. 7410.—Reg. Himal.

ochroleucum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1853) 490.—Java.

odontochilum, *Reichb. f. in Linnaea*, xli. (1877) 91.—N. Caled.

ophioglossoides, Sieber, ex Lindl. Gen. et Sp. Orch. 9 = Pleurothallis floribunda.

ophioglossoides, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 83 = Stelis ophioglossoides.

Ophioglossum, *Reichb. f.* in *Journ. Linn. Soc.* xv. (1876) 113.—Austral.

pachyglossum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 149.—Burma.

palmifolium, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 82 = Maxillaria palmifolia.

Palpebrae, *Lindl. in Journ. Hort. Soc.* v. (1850) 33. —Burma.

panduratum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 19.—Zeylan.

paniculatum, Pers. Syn. ii. 528 = Maxillaria paniculata.

paniculatum, *Sw. in Schrad. Neues Journ.* i. (1805) 97.—Afr. trop.

paradoxum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1858) 516.—Java.

parciflorum, *Reichb. f. ex Lindl. in Journ. Linn. Soc.* iii. (1859) 4.—Reg. Himal.

parcum, *Reichb. f. in Gard. Chron.* (1866) 1042.—Borneo.

pardalinum, Reichb. f. l. c. (1885) ii. 230 = Macraei.

Parishii, *Reichb. f. in Bot. Zeit.* xxi. (1863) 236; *Hook. Bot. Mag.* t. 5488.—Burma.

Parthenium, *Reichb. f. in Gard. Chron.* (1885) ii. 489. —Borneo.

parviflorum, *D. Don, Prod. Fl. Nep.* 34.—Nepal.

pauciflorum, Keinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 185 (= *Mitopetalum plicatum*).—Java.

Paxtoni, Lindl. Bot. Reg. (1839) Misc. 43 = chrysanthum.

Paxtoni, Paxt. Mag. Bot. vi. (1839) 163 = fimbriatum.

peguanum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 19. —Burma.

pendulum, *Roxb. Hort. Beng.* 63; *Fl. Ind.* iii. 484.—Burma.

Perula, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 298. —Assam.

Petri, *Reichb. f. in Gard. Chron.* (1877) i. 107.—Polynesia.

Phalaenopsae, *Fitzg. l. c.* (1880) ii. 38.—Austral.

pictum, Griff. ex Lindl. in Journ. Linn. Soc. iii. (1859) 12 = Devonianum.

pictum, *Lindl. in Gard. Chron.* (1862) 548.—Borneo.

Pierardi, *Roxb. Hort. Beng.* 63; *Fl. Ind.* iii. 482.—Reg. Himal.

planibulbe, *Lindl. Bot. Reg.* (1843) *Misc.* 54.—Manilla.

platyrrachium, *Reichb. f. Otia Bot. Hamb.* 55.—Ins. Fiji.

platypetalum, Pers. Syn. ii. 523 = Maxillaria platypetala.

pleiostachyum, *Reichb. f. in Gard. Chron.* (1882) ii. 590.—Guinea.

plicatum, F. G. Distr. Vollst. Lexik. Gaertn. Nachtr. ii. 651 = Maxillaria paniculata.

plicatile, *Lindl. Bot. Reg.* (1840) *Misc.* 10.—Ins. Philipp.

polyanthum, Wall. Cat. n. 2009 = pendulum!

polycaryum, *Reichb. f. in Gard. Chron.* (1883) ii. 490.—Ins. Sondaic.

polycladium, *Reichb. f. in Linnaea*, xli. (1877) 90.—N. Caled.

polypetalum, Steud. Nom. ed. I. 268, sphalm. = *platypetalum*.

polystachyon, Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 247 = Polystachya luteola.

polystachya, Thou. Orch. Afr. t. 85 = Polystachya mauritiana.

porphyrochilum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 18.—Reg. Himal.

praecultorum, *Reichb. f. in Gard. Chron.* (1877) i. 780. —Ind. or.

praevox, Sm. in Rees, Cyclop. xi. n. 3 = Coelogyne praecox.

DENDROBIUM:—

prasinum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 11. —Ins. Fiji.

primulinum, *Lindl. in Gard. Chron.* (1858) 400.—Reg. Himal.

profusum, *Reichb. f. l. c.* (1884) i. 310.—Ins. Philipp.

proliferum, Pers. Syn. ii. 524 = Maxillaria prolifera.

pruinosum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 214.—Malaya.

Pseudacline, *Lindl. in Journ. Linn. Soc.* iii. (1859) 9. —Ins. Philipp.

pubescens, Hook. Exot. Fl. t. 124 = Eria flava.

pugioniforme, *A. Cunn. in Lindl. Bot. Reg.* (1839) *Misc.* 33.—Austral.

pulchellum, *Lodd. Bot. Cab.* t. 1935 (non Roxb.).—China.

pulchellum, *Roxb. Hort. Beng.* 63; *Fl. Ind.* iii. 486 —Ind. or.

pumilum, *Roxb. ll. cc.* 63, 479.—Burma; Malaya.

pumilum, Sw. in Schrad. Neues Journ. i. (1805) 97 = Bulbophyllum pumilum.

punctatum, Sm. Exot. Bot. i. 21. t. 12 = Dipodium punctatum.

pungentifolium, F. Muell. Fragm. i. 189 = pugioniforme.

purpurascens, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 214.—Java.

purpuraseens, Thw. Enum. Fl. Zeyl. 298 (Coelogynes sp. !).—Zeylan.

purpureum, *Roxb. Hort. Beng.* 63; *Fl. Ind.* iii. 484.— Ins. Moluc.

pusillum, D. Don, Prod. Fl. Nep. 35 = alpestre!

pusillum, H. B. & K. Nov. Gen. et Sp. i. 357 = Pleurothallis pusilla.

pusillum, *Lindl. Gen. et Sp. Orch.* 77.—Java.

pycnostachyum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 19.—Burma.

pygmaeum, A. Cunn. in Hook. Comp. Bot. Mag. ii. (1836) 277 = Caleyi.

pygmaeum, *Lindl. in Wall. Cat.* n. 1999.—Reg. Himal.

pygmaeum, Sm. in Rees, Cyclop. xi. n. 27 = Bulbophyllum pygmaeum.

quadrifidum, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 40 = Pleurothallis quadrifida.

Rabaxi, Lindl. in Journ. Linn. Soc. iii. (1859) 7 = Macraei.

racemiferum, Sw. in Nov. Act. Soc. Sc. Upsal. vi (1799) 83 = Pleurothallis oblongifolia.

radians, *Reichb. f. Xenia Orch.* ii. 130.—Borneo.

ramosissimum, Wight, Ic. t. 1648 = herbaceum.

ramosum, *Lindl. in Wall. Cat.* n. 2003.—Reg. Himal.

ramosum, Pers. Syn. ii. 524 = Maxillaria ramosa.

refractum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 215.—Ins. Malay.

Reinwardti, *Lindl. in Journ. Linn. Soc.* iii. (1859) 17.—Ins. Banda.

reptans, *Franch. & Sav. Enum. Pl. Jap.* ii. 510.— Japon.

reptans, Sw. in Schrad. Neues Journ. i. (1805) 96 = Bulbophyllum nutans.

retusum, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 40 = Pleurothallis retusa.

retusum, Lindl. in Mem. Acad. Clenc. Madr. ii. (1850) 498 = macranthum!

revolutum, *Lindl. Bot. Reg.* (1840) *Misc.* 51.—Burma; Malaya.

rhodocentrum, *Reichb. f. in Gard. Chron.* (1872) 426. —Burma!

rhodopteryginum, *Reichb. f. l. c.* (1875) i. 634.—Burma.

rhodostoma v, Reichb. f. l. c. (1876) i. 795.

rhombeum, Lindl. Bot. Reg. (1843) t. 17 = aureum.

rigens, *Reichb. f. in Walp. Ann.* vi. 280.—Java.

rigescens, Mig. Fl. Ind. Bat. iii. 629 = firmum.

rigidum, *R. Br. Prod.* 333.—Austral.

rigidum, Lindl. Gen. et Sp. Orch. 86 = firmum.

Rimanni, *Reichb. f. in Gard. Chron.* (1882) ii. 650.— Ins. Moluc.

rorulentum, *Teijsm. & Binn. Cat. Hort. Bog.* 44, nomen.—Ins. Amboina.

Rosenbergii, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 217.—Java.

roseum, *Sw. in Schrad. Neues Journ.* i. (1806) 97.— Afr. trop.

Rostkovii, *Lindl. in Journ. Linn. Soc.* iii. (1859) 4. —Ind. or.

Ruckeri, *Lindl. Bot. Reg.* (1843) *Misc.* 25.—Reg. Himal.

DENDROBIUM:—
rugosum, *Lindl. Gen. et Sp. Orch.* 87.—Java.
Rumphiae, Reichb. f. in Walp. Ann. vi. 308 = angustifolium.
Rumphianum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 317.—Ins. Malay.
ruscifolium, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 84 = Pleurothallis ruscifolia.
salaccense, Rchb. ex Reichb. f. in Bot. Zeit. xx. (1862) 214 = Bullenianum.
salaconae, *Lindl. Gen. et Sp. Orch.* 86.—Java.
salliconioides, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 315.—Ins. Malay.
sanguineum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 82 = Bronghtonia sanguinea.
sanguinolentum, *Lindl. Bot. Reg.* (1842) *Misc.* 62. —Penang.
sarcanthum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 20.—Java.
Sarcostoma, *Lindl. Gen. et Sp. Orch.* 80.—Java.
sarcostomum, Teijsm. & Binn. ex Miq. Fl. Ind. Bat. iii. 538 = macrophyllum.
scabriligue, *Lindl. in Journ. Linn. Soc.* iii. (1859) 15.—Burma.
scalpelliforme, *Teijsm. & Binn in Tijdschr. Nederl. Ind.* xxvii. (1864) 17.—Malaya.
scariosum, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 39 = Pleurothallis scariosa.
Schimperianum, *A. Rich. Tent. Fl. Abyss.* ii. 282.—Abyssin.
schizanthum, Lindl. in Paxt. Flow. Gard. i. (1850-51) 184 = striolatum.
Schroederi, Hort. in Flor. Mag. t. 502 = densiflorum.
sclerophyllum, Lindl. t. c. (1844) Misc. 58 = firmum.
Scopa, *Lindl. Bot. Reg.* (1843) *Misc.* 55.—Ins. Philipp.
sculptum, *Reichb. f. in Bot. Zeit.* xxi. (1868) 128.—Borneo.
secundum, *Lindl. in Wall. Cat.* n. 1996; *et Bot. Reg.* t. 1991.—Malaya.
senile, *Parish, ex Reichb. f. in Gard. Chron.* (1865) 434. —Burma.
Serra, *Lindl. in Journ. Linn. Soc.* iii. (1859) 3.—Malaya.
sertularioides, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 88 = Pleurothallis sertularioides.
Shepherdi, F. Muell. Fragm. i. 190 = Bulbophyllum Shepherdi.
signatum, *Reichb. f. in Gard. Chron.* (1884) i. 306.—Siam.
sinuatum, *Lindl. ex Walp. Ann.* vi. 280.—Malaya.
Sindiise, *F. Muell. Fragm.* vi. 94.—Austral.
Spatella, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 298. —Ind. or.
spathaceum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 15. —Reg. Himal.
speciosum, *Sw. Anot. Bot.* i. 17. t. 10.—Austral.
spectabile, *Miq. Fl. Ind. Bat.* iii. 645.—N. Guinea.
spectatissimum, *Reichb. f. in Linnaea*, xli. (1877) 41. —Borneo.
sphegifer[*gl*]*ossum*, Reichb. f. in Bonplandia, ii. (1854) 88 = stupoosum.
spinescens, *Lindl. in Journ. Linn. Soc.* iii. (1859) 14. —Java.
splendidissimum ×, Reichb. f. in Gard. Chron. (1879) i. 396.
squalens, Lindl. in Bot. Reg. t. 782 = Maxillaria squalens.
stenoglossum, *Reichb. f. in Linnaea*, xli. (1877) 88.—N. Caled.
striatum, Griff. Notul. iii. 318 = Bulbophyllum striatum.
Strickladianum, *Reichb. f. in Gard. Chron.* (1877) i. 749.—Japon.
striolatum, *Reichb. f. in Hamb. Gartenz.* xiii. (1857) 313.—Austral.
strongylanthum, *Reichb. f. in Gard. Chron.* (1878) i. 402.—Ind. or.
Stuartii, *F. M. Bailey, in Proc. R. Soc. Queensl.* i. (1884) 12.—Austral.
stupoosum, *Lindl. Bot. Reg.* (1838) *Misc.* 52.—Reg. Himal.
suavissimum, Reichb. f. in Gard. Chron. (1874) i. 406 = chrysotoxum.
subacutum, *Reinw. ex Lindl. in Journ. Linn. Soc.* iii. (1859) 11.—Java.
subarcuatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 316.—Java.

DENDROBIUM:—
subtersa, Lindl. in Journ. Linn. Soc. iii. (1859) 4 = acerosum.
subulatum, *Lindl. Gen. et Sp. Orch.* 91.—Malaya.
sulcatum, *Lindl. Bot. Reg.* (1838) t. 65.—Reg. Himal.
Sumneri, *F. Muell. Fragm.* vi. 94.—Austral.
superbiens, *Reichb. f. in Gard. Chron.* (1876) ii. 516. —Austral.
superbum, *Reichb. f. in Walp. Ann.* vi. 282.—Ins. Philipp.
sylvanum, *Reichb. f. in Linnaea*, xli. (1877) 91.—N. Caled.
Tattonianum, Batem. in Gard. Chron. (1865) 890 = canaliculatum.
taurinum, *Lindl. Bot. Reg.* (1843) t. 28.—Ins. Philipp.
tenellum, *Lindl. Gen. et Sp. Orch.* 91.—Java.
teres, *Lindl. Bot. Reg.* (1840) *Misc.* 51.—Malaya.
teres, Roxb. Hort. Beng. 63; Fl. Ind. iii. 465 = Vanda teres.
teretifolium, *R. Br. Prod.* 333.—Austral.
teretifolium, Lindl. Bot. Reg. (1839) Misc. 32 = striolatum.
terminale, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 109.—Burma.
testiculatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 83 = Ionopsis testiculata.
tetrachromum, *Reichb. f. in Gard. Chron.* (1880) i. 713.—Borneo.
tetraedre, *Lindl. Gen. et Sp. Orch.* 89.—Java.
tetragonum, *A. Cunn. in Lindl. Bot. Reg.* (1839) *Misc.* 33.—Austral.
tetrodon, *Reichb. f. ex Lindl. in Journ. Linn. Soc.* iii. (1859) 10.—Java.
Teysmanni, *Miq. Fl. Ind. Bat.* iii. 640.—Singapore.
thrysiflorum, Reichb. f in Illustr. Hortic. xxii. (1875) 88. t. 207 = densiflorum.
thrysodes, *Reichb. f. in Hamb. Gartenz.* xv. (1859) 51. —Java.
tipuliferum, *Reichb. f. in Gard. Chron.* (1877) i. 72.—Ins. Fiji.
Tmesipteris, *Lindl. in Journ. Linn. Soc.* iii. (1859) 13. —Borneo.
Tokai, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 293 —Ins. Fiji.
tortile, Lour. Cann. in Lindl. Bot. Reg. (1839) Misc. 33 = monophyllum.
tortile, *Lindl. in Gard. Chron.* (1847) 797.—Burma.
transparens, *Wall. Cat.* n. 2008.—Reg. Himal.
Treacherianum, *Reichb. f. ex Hook. f. Bot. Mag.* t. 6591. —Borneo.
triadenium, Lindl. Bot. Reg. (1847) t. 1 = mutabile.
tribuloides, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 83 = Pleurothallis tribuloides.
trichostomum, *Reichb. f. ex Oliver, in Journ. Linn. Soc.* xv. (1876) 30.—N. Guinea.
tricolor, Pers. Syn. ii. 524 = Maxillaria tricolor.
tricuspe, *Lindl. Gen. et Sp. Orch.* 88.—Java.
tridentiferum, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 287.—N. Guinea.
trifiorum, *Lindl. Gen. et Sp. Orch.* 77.—Java.
triptalcides, Roxb. Hort. Beng. 63; Fl. Ind. iii. 478 = Bulbophyllum auricomum.
truncatum, Pers. Syn. ii. 524 = Maxillaria triphylla.
truncatum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 15. —Java.
tuadenum, *Bassll, ex Neves, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 284, nomen.—Ins. Philipp.
umbellatum, *Reichb. f. in Walp. Ann.* vi. 308.—Ins. Moluco.
uncatum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 5.—Java.
undatifiorum, Pers. Syn. ii. 524 = Maxillaria undatiflora.
undulatiflorum, Steud. Nom. ed. I. 368 = Maxillaria undulata.
undulatum, *R. Br. Prod.* 332.—Austral.
undulatum, Pers. Syn. ii. 524 = Maxillaria undulata.
unguiculatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1858) 491.—Java.
uniflorum, Griff. Notul. iii. 305 = revolutum.
uniflorum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 313.—Sumatra.
uniflos, *F. M. Bailey, in Proc. R. Soc. Queensl.* i. (1884) 13.—Austral.
utricularioides, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 83 = Ionopsis utricularioides.
vandiflorum, *Reichb. f. in Gard. Chron.* (1882) ii. 590. —N. Guinea.

DENDROBIUM:—
variegatum, Pers. Syn. ii. 524 = Maxillaria variegata.
Veitchianum, Lindl. Bot. Reg. (1847) t. 25 = macrophyllum.
venustum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 18.—Siam.
veratrifolium, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 286.—N. Guinea.
veratrifolium, Roxb. Hort. Beng. 63; Fl. Ind. iii. 482 = Plaiua veratrifolius.
verruciferum, *Reichb. f. in Linnaea*, xli. (1877) 88.—N. Caled.
vestitum, Wall. Cat. n. 2005 = Eria vestita.
vexabile ×, Reichb. f. in Gard. Chron. (1884) i. 271.—Ind. or.
villosulum, *Wall. Cat.* n. 2006.—Penang.
virgineum, *Reichb. f. in Gard. Chron.* (1884) ii. 520. —Burma.
viridiroseum, *Reichb. f. in Bonplandia*, iii. (1855) 226. —Java.
Wardianum, Warner, Select. Orch. Ser. I. t. 19 = pendulum.
Williamsianum, *Reichb. f. in Gard. Chron.* (1878) i. 652.—N. Guinea.
Williamsoni, *Day & Reichb. f. t. c.* (1869) 78.—Reg. Himal.
xantholeucum, *Reichb. f. Xenia Orch. ii.* 78.—Java.
xanthoplehlum, *Lindl. in Gard. Chron.* (1857) 268. —Burma.
Zollingerianum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 313.—Java.

DENDROBIUM, Ag. Aphor. 138 (1828), sphalm. = **Dendrocalamus**, Sw. (Orchid.).

DENDROCALAMUS, Nees, in Linnaea, ix. (1834) 478. *GRAMINEAE*, Benth. & Hook. f. iii. 1212.
Balcooa, Voigt, Hort. Suburb. Calc. 718 = Bambusa Balcooa.
calostachyus, *Kurs, For. Fl. Brit. Burma*, ii. 562.—Burma.
criticus, *Kurs, l. c.* 559.—Burma.
flagellifer, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 150.—Malacca.
giganteus, *Munro, l. c.*—Burma.
Griffithianus, *Kurs, For. Fl. Brit. Burma*, ii. 562. —Burma.
Hamiltonii, *Nees & Arn. ex Munro, in Trans. Linn. Soc.* xxvi. (1868) 151.—Reg. Himal.
Hookeri, *Munro, l. c.*—Reg. Himal.
latiflorus, *Munro, l. c.* 152. t. 6.—China.
longispathus, *Kurs, For. Fl. Brit. Burma*, ii. 561.—Burma.
membranaceus, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 149.—Burma.
monadelphus, Thw. Enum. Pl. Zeyl. 376 = Oxytenanthera Thwaitesii.
Parishii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 149.—Reg. Himal.
sericeus, *Munro, l. c.* 148.—Ind. or.
strictus, *Nees, in Linnaea*, ix. (1834) 476.—Ind. or.
Tulda, Voigt, Hort. Suburb. Calc. 718 = Bambusa Tulda.

DENDROCHARIS, Miq. in Versl. en Med. Akad. Wet. vi. (1857) 194; et Fl. Ind. Bat. ii. 453 (1859) = **Bodyesanthera**, Hook. (Apocyn.).
inflata, Miq. ll. cc. 194, 453 = Anodendron inflatum.
myrtifolia, Miq. ll. cc. 194, 454.—Malaya.
rubescens, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 403.—Borneo.

DENDROCHILUM, Blume, Bijdr. 398. t. 52 (1825) quoad Sect. I. *ORCHIDEAE*, Benth. & Hook. f. iii. 506 (cf. 496).
abbreviatum, Blume, l. c. 400 = Platyclinis abbreviata.
Arachnites, *Reichb. f. in Gard. Chron.* (1882) i. 256.—Ins. Philipp.
aurantiacum, *Blume, Bijdr.* 398. t. 5.—Java.
aurium, *Reichb. f. in Bonplandia*, iv. (1856) 329.—Java.
brachystemon, *Reichb. f. l. c.* v. (1857) 43 = Platyclinis brachyota.
bracteosum, *Reichb. f. in Walp. Ann.* vi. 241.—Hab.?
Cobblianum, Reichb. f. in Gard. Chron. (1880) ii. 748 = Platyclinis Cobbliana.
cornutum, Blume, Bijdr. 939 = Platyclinis cornuta.

DENDROCHILUM :—
cucumerinum, *Reichb. f. in Gard. Chron.* (1884) ii. 649.—Ins. Philipp.
edentulum, Blume, Bijdr. 399 = Platyclinis edentula.
erosum, *Reichb. f. in Walp. Ann.* vi. 241.—Java.
filiforme, Lindl. Bot. Reg. (1840) Misc. 53 = Platyclinis filiformis.
fuscum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 205.—Ind. or.
glumaceum, André, Illustr. Hortic. xxv. (1878) t. 328 = Platyclinis filiformis.
glumaceum, Lindl. Bot. Reg. (1841) Misc. 28 = Platyclinis glumacea.
junceum, *Reichb. f. Otia Bot. Hamb.* 54.—Ins. Philipp.
latifolium, Lindl. Bot. Reg. (1848) Misc. 56 = Platyclinis latifolia.
longifolium, Reichb. f. in Benplandia, iv. (1856) 329 = Platyclinis longifolia.
magnum, *Reichb. f. in Walp. Ann.* vi. 240.—Hab.?
occultum, *Lindl. Gen. et Sp. Orch.* 34.—Java.
pallidiflavens, *Blume, Bijdr.* 399. t. 52.—Java.
pumilum, *Reichb. f. in Bonplandia*, iii. (1855) 222. —Ins. Philipp.
roseum, Dalz. in Hook. Kew Journ. iv. (1852) 291 = Dendrobium crepidatum.
simile, *Blume, Bijdr.* 400.—Java.
spathaceum, *Reichb. f. in Bonplandia*, v. (1857) 43.—Java.
uncatum, *Reichb. f. l. c.* iii. (1855) 222.—Ins. Philipp. Zollingeri, *Miq. Fl. Ind. Bat.* iii. 626.—Java.

DENDROCNIDE, Miq. Fl. Jungh. 29 (1851) = **Laportea**, Gaudich. (Urtic.).
coerulea, Miq. l. c. 31 = Cyphophapus rufescens.
costata, Miq. l. c. 30 = L. costata.
cremulata, Miq. l. c. 31 = L. crenulata.
pellata, Miq. l. c. 30 = L. peltata.

DENDROCOLLA, Blume, Bijdr. 286. t. 67 (1825) = **Sarcochilus**, R. Br. (Orchid.).
acuminatissima, Blume, l. c. 288 = S. acuminatissimus.
amplexicaulis, Blume, l. c. = S. amplexicaulis.
anceps, Blume, l. c. 292 = S. anceps.
angustifolia, Blume, l. c. 291 = S. angustifolius.
aprinlata, Zoll. ex Reichb. f. in Bonplandia, v. (1857) 39, in obs. (= *Camarotis aprinlata*).—Java!
appendiculata, Blume, l. c. 289 = S. pusillus.
Arachnites, Blume, l. c. 287 = S. Arachnites.
compressa, Blume, l. c. 290 = S. compressus.
emarginata, Blume, l. c. 290 = S emarginatus.
gracilenta, Reichb. f. in Bonplandia, v. (1857) 40 = S. gracilentus.
Hystrix, Blume, Bijdr. 291 = S. Hystrix.
minima, Blume, l. c. 290 = S. minimus.
obtusa, Blume, l. c. 287 = S. obtusus.
pallida, Blume, l. c. 290 = S. pallidus.
pulchella, Thw. Enum. Pl. Zeyl. 480 = S. pulchellus.
purpurascens, Blume, Bijdr. 288 = S. purpurascens.
pusilla, Blume, l. c. 289 = S. pusillus.
rhopalorrhachis, Reichb. f. in Bonplandia, v. (1857) 40 = S. rhopalorrhachis.
serraeformis, Lindl. ex Thw. Enum. Pl. Zeyl. 306 = S. serraeformis.
spuria, Blume, Bijdr. 290 = S. spurius.
subulata, Blume, l. c. 291 = S. subulatus.
teres, Blume, l. c. 289 = S. teres.
Zollingeri, Reichb. f. in Bonplandia, v. (1857) 40 = S. Zollingeri.

DENDRODAPHNE, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 145, 'Genus delendum,' Benth. & Hook. f. iii. 150.
macrophylla, Beurl. l. c.—Panama.

DENDRORIJIUM, Blume, Bijdr. 848. t. 69 (1825) = **Rhta**, Lindl. (Orchid.).
abbreviatum, Blume, l. c. 346 = R. abbreviata.
acuminatum, Blume, l. c. 347 = E. acuminata.
albidotomentosum, Blume, l. c. 345 = E. albidotomentosa.
appendiculatum, Blume, l. c. 352 = E. appendiculata.
bicristatum, Blume, l. c. 346 = E. bicristata.
coriaceum, Blume, l. c. 351 = E. coriacea.
densiflorum, Blume, l. c. 350 = Pinnatia densiflora.
ebulbe, Blume, l. c. 347 = E. ebulbis.
erectum, Blume, l. c. 348 = E. erecta.

DENDRORLIRUM :—
flavescens, Blume, Bijdr. 344 = Erin flavescens.
hyacinthoides, Blume, l. c. 346 = E. hyacinthoides.
lasiflorum, Blume, l. c. 350 = Pinnatia lasiflora.
micranthum, Blume, l. c. 349 = E. micrantha.
multiflorum, Blume, l. c. = E. multiflora.
ornatum, Blume, l. c. 345 = E. ornata.
pusillum, Blume, l. c. 350 = Pinnatia pusilla.
retusum, Blume, l. c. 351 = E. retusa.
robustum, Blume, l. c. 347 = E. robusta.
rugosum, Blume, l. c. 345 = E. stellata.
secundum, Blume, l. c. 350 = E. secunda.
sulcatum, Blume, l. c. 347 = E. sulcata.

ENDROLOBIUM, Benth. in Miq. Pl. Jungh. 216 (1852) = **Desmodium**, Desv. (Legum.).
australe, Benth. l. c. = Desm. Cephalotes.
Cephalotes, Benth. l. c. 216, 218 = Desm. Cephalotes.
Cuminghamum, Benth. l. c. 216.—Ins. Philipp.
umbellatum, Benth. l. c. 218 = Desm. umbellatum.

DENDROMECON, Benth. in Trans. Hort. Soc. Ser. II. i. (1834) 407. *PAPAVERACEAE*, Benth. & Hook. f. i. 54.
Harfordii, *Kellogg, in Proc. Calif. Acad.* v. (1873) 102.—Calif.
rigidum, *Benth. in Trans. Hort. Soc.* Ser. II. i. (1834) 407.—Calif.

DENDROPANAX, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 107. *ARALIACEAE*, Benth. & Hook. f. i. 948.
TEXTURIA, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 12 (1863).
alaris, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 107.—Mexic.
arborcum, *Decne. & Planch. l. c.*—N. Granat.
citrifolium, *Decne. & Planch. l. c.*—Mexic.
cuneatum, Decne. & Planch. l. c. = Gilibertia cuneata.
darienne, Seem. in Journ. Bot. ii. (1864) 300 = Hedera Seemanni.
densiflorum, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 107, nomen = Gilibertia cuneata.
Fendleri, *Seem. in Journ. Bot. ii.* (1864) 301.—Am. austr.
japonicum, *Seem. l. c.*—Japon.
Juergenseni, *Seem. l. c.*—Mexic.
lanceolatum, *Decne. & Planch.* in Rev. Hortic. Sér. IV. iii. (1854) 107 = Gilibertia cuneata.
lanciifolium, Seem. l. c. 302 = laurifolium.
Langsea, *March. in Bull. Acad. Brux.* xlvii. (1879) 12.—Mexic.
laurifolium, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 107, nomen.—Hab.?
monogynum, Seem. in Journ. Bot. iii. (1865) 140 = Gilibertia monogyna.
montanum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 107, nomen.—Hab.?
nutans, *Decne. & Planch. l. c.*—Jamaic.
oblongum, Decne. & Planch. l. c. = Gilibertia cuneata.
obovatum, Decne. & Planch. l. c. = Gilibertia cuneata.
ovatum, Seem. in Journ. Bot. ii. (1864) 301 = Arthrophyllum diversifolium.
parviflorum, *Benth. Fl. Hongk.* 137.—China.
pauciflorum, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 107 = Gilibertia monogyna.
Pavonii, Decne. & Planch. l. c. = Gilibertia umbellata.
pendulum, *Decne. & Planch. l. c.*—N. Granat.
protensum, Benth. Fl. Hongk. 136 = parviflorum.
ramiflorum, Seem. in Journ. Bot. ii. (1864) 312 = Gilibertia cuneata.
Sellowianum, Miq. Ann. Mus. Bot. Lugd. Bat. i. 27 = Gilibertia cuneata.
tomentosum, Seem. in Journ. Bot. ii. (1864) 302 = Didymopanax lucunoides.
trilobum, Seem. l. c. = Gilibertia trilobu.

DENDROPEMON, Blume, in Schult. Syst. f. vii. 1729 (1830) = **Loranthus**, Linn.
anceps, *emarginatum*, *lanciflorum*, *occidentale*, *parviflorum*, *paucanceps*, *uniflorum*, Blume, l. c. Omnes Loranthi species homonymae.

DENDROPHUCON, Endl. Gen. Suppl. v. 19 (1850), aphalm. = prace.

DENDROPHILA, Zipp. ex Blume, Rumphia, iv. 84. t. 190 (1848) = **Tecoma**, Jus. (Bignon.).
trifoliata, Zipp. ex Blume, l. c. 86 = T. Dendrophila.

DENDROPHTHOË, Mart. in Flora, xiii. (1830) i. 109 = Loranthus, Linn.
atropurpurea, Blume, in Schult. f. Syst. vii. 1730 = L. atropurpureus.
bicolor, Mart. in Flora, xiii. (1830) i. 109 = L. longiflorus.
bifora, Blume, in Schult. f. Syst. vii. 1730 = L. Scurrula!
buddlejoides, Blume, l. c. = L. Scurrula.
celastroides, Mart. in Flora, xiii. (1830) i. 110 = L. celastroides.
chrysantha, Blume, in Schult. f. Syst. vii. 1730 = L. chrysanthus.
chrysanthoides, Miq. Fl. Ind. Bat. i. i. 813 = L. chrysanthoides.
cinnamomea, G. Don, Gen. Syst. iii. 420 = L. Scurrula.
coccinea, G. Don, l. c. 419 = L. coccineus.
congener, Mart. in Flora, xiii. (1830) i. 110 = L. pendulus.
cordifolia, Mart. l. c. = L. cordifolius.
Corynitis, Blume, in Schult. f. Syst. vii. 1729 = L. Corynitis.
costulata, Miq. Fl. Ind. Bat. Suppl. 844.—Sumatra.
cuneata, Blume, in Schult. f. Syst. vii. 1730 = L. rhoparcarpus.
curvata, Blume, l. c. = L. curvatus.
cylindrica, Blume, l. c. = L. cylindricus
dasystemon, Miq. Fl. Ind. Bat. i. i. 1097.—Sumatra.
elegans, Mart. in Flora, xiii. (1830) i. 109 = L. elegans.
falcata, Blume, in Schult. f. Syst. vii. 1730 = L. longiflorus.
farinosa, Mart. in Flora, xiii. (1830) i. 110 = L. pentandrus.
fasciculata, Blume, in Schult. f. Syst. vii. 1730 = L. fascicularus.
ferruginea, Blume, l. c. = L. ferrugineus.
Forsteriana, Blume, l. c. = L. Forsterianus.
fulva, Miq. Fl. Ind. Bat. i. i. 815 = L. fulvus.
fusca, Blume, in Schult. f. Syst. vii. 1730 = L. rhopalocarpus.
fuscata, Miq. Fl. Ind. Bat. i. i. 816 = L. fuscatus.
Gaudichaudii, G. Don, Gen. Syst. iii. 419 = L. Gaudichaudii.
glauca, Mart. in Flora, xiii. (1830) i. 109 = L. glaucus.
gracilifora, Mart. l. c. 110 = L. Scurrula.
gracilifolia, Mart. l. c. = prace.
Haenheana, Mart. l. c. = L. Haenkeanus.
Heyneanus, Blume, in Schult. f. Syst. vii. 1730 = L. bractentus.
Heynei, G. Don, Gen. Syst. iii. 420 = L. Scurrula.
ignea, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 357.—Ins. Banca.
incarnata, Blume, in Schult. f. Syst. vii. 1730 = L. incarnatus.
indica, Blume, l. c. = L. longiflorus.
Jungkuhnii, Miq. Fl. Ind. Bat. i. i. 817 = L. Jungkuhnii.
kalabiensis, Miq. l. c. 811 = L. kalabiensis.
Korthigrana, Blume, in Schult. f. Syst. vii. 1730 = L. longiflorus.
Korthalsii, Miq. Fl. Ind. Bat. i. i. 814 = L. Korthalsii.
kramatensis, Miq. l. c.—Borneo.
lepidota, Mart. in Flora, xiii. (1830) i. 110 = L. lepidotus.
lepstantha, G. Don, Gen. Syst. iii. 420 = L. pulverulentus.
leucobotrya, Miq. Fl. Ind. Bat. Suppl. 845.—Ins. Banca.
leucostachys, Miq. l. c. i. i. 819 = L. leucostachys.
ligustrina, Mart. in Flora, xiii. (1830) i. 110 = L. ligustrinus.
linophylla, Steud. Nom. ed. II. i. 491 = L. linophyllus.
longiflora, Blume, in Schult. f. Syst. vii. 1730 = L. longiflorus.
lusonensis, Blume, l. c. = L. lusonensis.
macrocalyx, Miq. Fl. Ind. Bat. i. i. 821 = L. heteranthus.
malifolia, Blume, in Schult. f. Syst. vii. 1730 = L. malifolius.
medinensis, Miq. Fl. Ind. Bat. i. i. 816 = L. medinensis.
obovata, Blume, in Schult. f. Syst. vii. 1730 = L. Scurrula.
obtecta, G. Don, Gen. Syst. iii. 419 = L. Scurrula.
Ovriana, Miq. Fl. Ind. Bat. i. i. 813 = L. faringinus.
ovalifolia, Steud. Nom. ed. II. i. 491 = L. ovalifolius.
pendula, Mart. in Flora, xiii. (1830) i. 110 = L. pendulus.
pentandra, Blume, in Schult. f. Syst. vii. 1730 = Loranthus pentandrus.
pentapetala, G. Don, Gen. Syst. iii. 419 = L. pentapetalus.

DENDROPHTHOË :—
philippensis, Blume, in Schult. f. Syst. vii. 1780 = L. philippensis.
praelonga, Blume, l. c. = L. praelongus.
pulchra, G. Don, Gen. Syst. iii. 419 = L. pulcher.
pulverulenta, Blume, in Schult. f. Syst. vii. 1780 = L. pulverulentus.
racemifera, G. Don, Gen. Syst. iii. 419 = L. coccineus
recurva, G. Don, l. c. 420 = L. recurvus.
repanda, Blume, in Schult. f. Syst. vii. 1780 = L. repandus.
Roxburghiana, Blume, l. c. = L. Scurrula.
rugulosa, Blume, l. c. = L. rugulosus.
Schultesii, Blume, l. c. = L. ferrugineus.
Scurrula, Blume, l. c. = L. Scurrula.
sphenoidea, Blume, l. c. = L. sphenoideus.
tetropetala, Blume, l. c. = L. tetrapetalus.
umbellata, Blume, l. c. = L. loniceroides.
umbellifer, Blume, l. c. = L. umbellifer.
venosa, Mart. in Flora, xiii. (1830) 1. 110 = L. venosus.
verticillata, Scheff. in Ann. Jard. Buitenz. i. (1876) 27. —N. Guinea.
vestita, Blume, in Schult. f. Syst. vii. 1780 = L. vestitus.
Wallichiana, G. Don, Gen. Syst. iii. 419 = L. Wallichianus.

DENDROPHTHORA, Eichl. in Mart. Fl. Bras. v. II. 102. t. 31 (1868). *LORANTHACEAE*, Benth. & Hook. f. iii. 214.
ARCEUTHOBIUM, Griseb. Fl. Brit. W. Ind. 814 (1860).
amoebandra, *Wright, in Sauv. Fl. Cub.* 141.—Cuba
arcuata, *Wright, l. c.* 142.—Cuba.
biserrula, *Eichl. in Mart. Fl. Bras.* v. 11. 104.—Am. trop.
Bonaniae, *Eichl. l. c.* 103.—Cuba.
buxifolia, *Eichl. l. c.* 105.—Ins. S. Doming.
constricta, *Eichl. l. c.* 104.—Cuba.
cubensis, *Eichl. l. c.*—Cuba.
cupressoides, *Eichl. l. c.* 103.—Ind. occ.
cupulata, *Eichl. l. c.* 102.—Ins. S. Doming.
domingensis, *Eichl. l. c.* 108.—Ind. occ.
Epiviscum, *Eichl. l. c.*—Cuba.
glauca, *Eichl. l. c.*—Cuba.
gracilis, *Eichl. l. c.*—Jamaica.
gracilis, *Wright, in Sauv. Fl. Cub.* 141.—Cuba.
grandifolia, *Eichl. in Mart. Fl. Bras.* v. 11. 105.—Cuba.
Griebachii, *Eichl. l. c.* 104.—Cuba.
leptostachya, *Eichl. l. c.*—Ind. occ.
macrostachya, *Eichl. l. c.*—Ins. Martinic.
Mancinellae, *Eichl. l. c.* 103.—Cuba.
myrtilloides, *Eichl. l. c.* 104.—Cuba.
opuntioides, *Eichl. l. c.* 102.—Ind. occ.
Wrightii, *Et Al. l. c.* 108.—Cuba.

DENDROPHYLAX, Reichb. f. in Walp. Ann. vi. 903 (1861). *ORCHIDEAE*, Benth. & Hook. f. iii. 585.
funalis, *Benth. & Hook. f.—Cf. Gen.* iii. 585, 588.—Ind. occ.
hymenanthus, *Reichb. f. l. c.*—Cuba.

DENDROPOGON, Rafin. Neogenyt. 3 (1825) = Tillandsia, Linn. (Bromeliac.).
usneoides, Rafin. l. c. = T. usneoides.

DENDROSERIS, D. Don, in Phil. Mag. xi. (1832) 388. *COMPOSITAE*, Benth. & Hook. f. ii. 504.
REA, Bert. ex Decne. in Guillem. Arch. Bot. i. (1838) 513. t. 9, 10.
Berteriana, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 93.—Ins. J. Fernand.
laevis, *Hemsl. Bot. Voy. Challenger*, i. 111. 99.—Ins. S. Ambrosio.
macrophylla, *D. Don, in Phil. Mag.* xi. (1832) 868.—Ins. J. Fernand.
marginata, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 93.—Ins. J. Fernand.
micrantha, *Hook. & Arn. l. c.*—Ins. J. Fernand.
mollis, *Hook. & Arn. l. c.*—Ins. J. Fernand.
nerilloides, *Hook. & Arn. l. c.*—Ins. J. Fernand.
pinnata, *Hook. & Arn. l. c.*—Ins. J. Fernand.

DENDROSICYOS, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 513. *CUCURBITACEAE*.
Jaubertiana, *Balf. in Bull. Soc. Linn. Par.* i. (1885) 443.—Afr. trop.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 513.—Ins. Socotr.

DENDROSMA, R. Br. ex Cromb. in Journ. Linn. Soc. xvii. (1879) 395, nomen = **Atherosperma**, Labill. † *Mucrosma*.
lucida, R. Br. ex Cromb. l. c. nomen.—Austral.

DENDROSMA, Panch. & Sebert, Not. Bois Nouv. Calédon. 245 (1873). *RUTACEAE*.
Deplanchei, *Panch. & Sebert, l. c.*—N. Caled.

DENDROSPARTUM, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 152 = **Genista**, Linn. (Legumin.).
aetnense, Spach, l. c. = G. aetnensis.
pungens, Spach, ex Walp. Ann. i. 230 = G. purgans.
sardoum, Spach, ex Walp. l. c. = G. aetnensis.

DENDROSTYLIS, Karst. & Triana, in Linnaea, xxviii. (1856) 481. *BIXINEAE*, Benth. & Hook. f. i. 125, 971.
apetiosefolia, *Karst. & Triana, l. c.*—Venezuela.
denticulata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 95.—Guiana.
echinata, *Benth. in Journ. Linn. Soc. v. Suppl.* ii. (1861) 82.—Bras.
grandifolia, *Karst. & Triana, in Linnaea*, xxviii. (1856) 481.—N. Granat.
microphylla, *Karst. & Triana, l. c.*—N. Granat.
pulsecensis, *Karst & Triana, l. c.*—N. Granat.
suaveolens, *Karst. & Triana, l. c.*—N. Granat.

DENDROTROPHE, Miq. Fl. Ind. Bat. i. 1. 779 (1855) = **Henslowia**, Blume (Santalac.).
buxifolia, Miq. l. c. 781 = H. buxifolia.
cassinefolia, Miq. l. c. 780 = H. cassinefolia.
Lobbiana, Miq. l. c. 1096 = H. Lobbiana.
philippinensis, Miq. l. c. = H. varians †
pterinervia, Miq. l. c.—Sumatra.
Reinwardtiana, Miq. l. c. 780 = H. Reinwardtiana.
spicata, Miq. l. c. = H. spicata.
umbellata, Miq. l. c. 779 = H. heterantha †
varians, Miq. l. c. 780 = H. varians.

DENECKIA, Sch. Bip. in Walp. Rep. ii. 952 (1843) = **Denekia**, Thunb. (Compos.).

DENEKIA, Thunb. Prod. Pl. Cap. ii. 158 (1800); Nov. Gen. 177 (1801). *COMPOSITAE*, Benth. & Hook. f. ii. 398.
capensis, *Thunb. Prod. Pl. Cap.* 158.—Afr. austr.
glabrata, *DC. Prod.* v. 462.—Afr. austr.

DENHAMIA, Meisn. Gen. 18 (1837). *CELASTRINEAE*, Benth. & Hook. f. i. 366, 998.
LUCARIUM, A. Rich. Sert. Astrol. 46. t. 46 (1834).
heterophylla, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 29 = obscura.
leucocarpum, Stend. Nom. ed. II. i. 491 = obscura.
obscura, *Meisn. in Walp. Rep.* i. 208.—Austral.
Oleaster, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 29.—Austral.
pittosporoides, *F. Muell. l. c.* 30.—Austral.
xanthosperma, F. Muell. l. c. 28 = obscura.

DENHAMIA, Schott, Melet. 19 (1832) = **Culcasia**, Beauv. (Aroid.).
scandens, Schott, l. c. = C. scandens.

DENIRA, Adans. Fam. ii. 118 (1763) = **Ivа**, Linn. (Compos.).

DENISAEA, Neck. Elem. i. 306 (1790) = **Phryma**, Linn. (Verben.).

DENISONIA, F. Muell. Fragm. i. 124. t. 2 (1859). *VERBENACEAE*, Benth. & Hook. f. ii. 1141.
ternifolia, *F. Muell. l. c.*—Austral.

DENNECKIA, Stend. Nom. ed. II. i. 491 (1840) = **Denekia**, Thunb. (Compos.).

DENS-CANIS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 197 (1745) = **Erythronium**, Linn. (Liliac.).

DENS-LEONIS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 197 (1745) = **Taraxacum**, Linn. (Compos.).

DENTARIA, [Tourn.] Linn. Syst. ed. I (1735) = **Cardamine**, Linn. (Crucif.).
alternifolia, Dalla Torre, Anl. Alpenfl. 62 = *trifolia*.
americana, Bartr. ex DC. Syst. ii. 273 = C. laciniata.
angulata, Torr. in Pacif. Rail. Rep. iv. 65 = C. pauciseta.
bifolia, Stokes, Bot. Mat. Med. iii. 448 = C. diphylla.
bipinnata, C. A. Mey. Verz. Pfl. Cauc. 179.—Reg. Caucas.
bulbifera, Linn. Sp. Pl. 653 = C. bulbifera.
californica, Nutt. ex Torr. & Gray, Fl. N. Am. i. 88 = C. pauciseta.
canescens, Tenore, Ind. Sem. Hort. Neap. (1830) 18, ex Syll. 218 = C. pinnata.
caucasica, Willd. ex DC. Prod. i. 155 = *quinquefolia*.
Clusiana, Reichb. Fl. Germ. Excurs. 671.—Europ.
concatenata, Michx. Fl. Bor. Am. ii. 30 = Thelypodium laciniatum.
dasyloba, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 296 = macrophylla.
digenea ×, Greuil, Nve Beitr. Fl. Schw. 8.—Europ.
digitata, Lam. Encyc. ii. 268 = C. pratensis.
diphylla, Michx. Fl. Bor. Am. ii. 30 = C. diphylla.
dissecta, Leav. in Am. Journ. Sc. vii. (1824) 62 = C. multifida.
enneaphylla, Genera. Elench. n. 604 = *glandulosa*.
enneaphylla, Gueldenst. ex Ledeb. Fl. Ross. i. 130 = *microphylla*.
enneaphylla, Linn. Sp. Pl. 653 = C. enneaphylla.
glabra, Scop. ex Steud. Nom. ed. II. i. 492 = Arabis perfoliata.
glandulosa, Waldst. & Kit. Pl. Rar. Hung. iii. 302. t. 272.—Hungar.
Gmelini, Tausch, in Flora, xix. (1836) 402 = C. macrophylla.
heptaphylla, Vill. Hist. Pl. Dauph. iii. 864 = C. pinnata.
heterophylla, Nutt. Gen. Am. ii. 66 = C. heterophylla.
hirsutula, Andrz. ex DC. Prod. i. 152 = C. macrophylla.
hypanica, Bess. ex DC. Syst. ii. 278 = *quinquefolia*.
integrifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 88 = C. pauciseta.
intermedia, Sond. in Flora, xxxvii. (1855) 129 = C. pentaphylla.
Kitlitzii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.-xxiv. (1880) 73.—Europ.
laciniata, Muhl. Cat. 60; ex Willd. Sp. Pl. iii. 479 = C. laciniata.
leucantha, Tausch, in Flora, xix. (1836) 404.—Hab. † *macrocarpa*, Nutt. ex Torr. & Gray, Fl. N. Am. i. 88. —Am. bor.
macrophylla, Bunge, ex baxim. Prim. Fl. Amur. 45.—Reg. Amur.
maxima, Nutt. Gen. Am. ii. 66 = C. maxima.
microphylla, Willd. Sp. Pl. iii. 479.—Reg. Caucas.
multifida, Muhl. Cat. 63 = C. multifida.
ochroleuca, Gaud. ex DC. Prod. i. 154 = C. macrophylla.
parvifolia, Rafin. Ann. Nat. (1820) 12.—Am. bor.
pentaphyllos, Gueldenst. Reisen, i. 194, 490 = *quinquefolia*.
pentaphyllos, Linn. Sp. Pl. 654 = C. pentaphylla.
pinnata, Lam. Encyc. ii. 268 = C. pinnata.
pinnata, Pall. ex DC. Syst. ii. 277 = *quinquefolia*.
polyphylla, Waldst. & Kit. Pl. Rar. Hung. ii. 174. t. 160.—Hungar.
quinquefolia, Bieb. Fl. Taur. Cauc. ii. 109; Suppl. 486.—Rossia austr.
reperа, Franch. in Bull. Soc. Bot. Fr. xxxii. (1885) 5. —China.
tenella, Pursh, Fl. Am. Sept. ii. 439.—Am. bor. occ.
tenuifolia, Hook. Fl. Bor. Am. i. 46 = prac.
tenuifolia, Ledeb. in Mém. Acad. Pétersb. v. (1815) 547 = C. tuberosa.
trifida, Lam. Illustr. t. 569. f. 3 = C. tenuifolia.
trifolia, Waldst. & Kit. Pl. Rar. Hung. ii. 143. t. 139. —Croatia.
Wallichii, G. Don, Gen. Syst. i. 172 = C. macrophylla.
Willdenowii, Tausch, in Flora, xix. (1836) 403 = C. macrophylla.

DENTELLA, Forst. Char. Gen. 25. t. 13 (1776). *RUBIACEAE*, Benth. & Hook. f. ii. 84.
HEYNEA, Denust. Schlüssel. Hort. Malab. 35 (1818). LITPAVA, Endl. Atakt. 13. t. 13 (1838).

DENTELLA:—
bifida, Buch.-Ham. ex Wall. Cat. n. 6216 = Spiradiclis bifida.
crocea, Roth, Nov. Pl. Sp. 140 = Cephalostigma Schimperi.
geminata, Buch.-Ham. ex Wall. Cat. sub n. 6237 = Polyura geminata.
minima, *Zipp. ex Span. in Linnaea*, xv. (1841) 817, nomen.—Ins. Timor.
perotifolia, Willd. ex DC. Prod. iv. 419 = Cephalostigma Schimperi.
repens, *Forst. Char. Gen.* 26. t. 13.—As. et Austral. trop.
stolonifera, *Monir. in Mém. Acad. Lyon*, x. (1860) 216.—N. Caled.

DENTIDIA, Lour. Fl. Cochinch. 369 (1790) = **Perilla**, Linn. (Labiat.).
nankinensis, Lour. l. c. = P. nankinensis.
purpurascens, Pers. Syn. ii. 135 = P. nankinensis.
purpurea, Poir. Encyc. Suppl. ii. 468 = P. nankinensis.

DEPIERREA, Anon. ex Schlecht. in Linnæa, xvi. (1842) 374 = **Campanula**, Tourn.
campanuloides, Anon. l. c. = C. rotundifolia!

DEPLACHNE, Boiss. Fl. Orient. v. 561 (1881) = **Diplachne**, Beauv. (Gram.).

DEPLANCHEA, Vieill. in Bull. Soc. Linn. Normand. vii. (1863) 96 = **Diplanthera**, Banks & Soland. (Bignon.).
Bulowii, F. Muell. Fragm. v. 92 = Dipl. tetraphylla.
speciosa, Vieill. in Bull. Soc. Linn. Normand. vii. (1863) 96.—N. Caled.

DEPPEA, Cham. & Schlecht. in Linnæa, v. (1830) 167. *RUBIACEÆ*, Benth. & Hook. f. ii. 51.
CHORISTEA, Benth. Pl. Hartw. 63 (1840).
cornifolia, *Benth. Pl. Hartw.* 849.—Mexic.
costa-ricensis, *Polak. in Linnæa*, xli. (1877) 566.—Costa Rica.
erythrorhiza, *Cham. & Schlecht. l. c.* v. (1830) 168.—Mexic.
floribunda, *Hemsl. Diag. Pl. Nov.* 31.—Mexic.
grandiflora, *Schlecht. in Linnæa*, xix. (1847) 748.—Mexic.
hadyotidea, DC. Prod. iv. 618 = erythrorhiza.
obtusiflora, *Benth. Pl. Hartw.* 849.—Mexic.
pubescens, *Hemsl. Diag. Pl. Nov.* 31.—Mexic.
tenuiflora, *Benth. Pl. Hartw.* 849.—Mexic.
umbellata, *Hemsl. Diag. Pl. Nov.* 31.—Mexic.

DEPPIA, Rafin. Fl. Tellur. ii. 51 (1836) = **Lycoasta**, Lindl. (1842) (Orchid.).
mexicana, Rafin. l. c. = L. Deppei.

DEPREA, Rafin. Sylva Tellur. 57 (1838) = **Withania**, Pauq. (Solan.).
crinacantis, Rafin. l. c. = W. ornocensis.
salapensis, Rafin. l. c. = Athenaea xalapensis.

DEPREMESNILIA, F. Muell. Fragm. x. 59 (1876). *LABIATÆ*.
chrysocalyx, *F. Muell. l. c.*—Austral.

DERDERIA, Jaub. & Spach, Illustr. i. 29. t. 67 (1843) = **Jurinea**, Cass. (Compos.).
Auchery, Jaub. & Spach, in Ann. Sc. Nat. Sér. III. xiii. (1849) 363.—Armenia.
berardioides, Jaub. & Spach, Illustr. iii. 134 (= *Aegopogon berardioides*).—Persia.
cheiriifolia, Jaub. & Spach, l. c. ii. 98 = J. Aucheriana.
eriobasis, Jaub. & Spach, l. c. 97 = J. eriobasis.
macrocephala, Jaub. & Spach, l. c. 129. t. 67 = J. macrocephala.

DERINGA, Adans. Fam. ii. 498 (1763) = **Cryptotaenia**, DC. (Umbellif.).

DERINGIA, Stend. Nom. ed. II. i. 492 (1840) = prec.

DERLINIA, Nénad, in Freyc. Voy. Bot. 38 (1826) = **Gratiola**, Rupp. (Scrophul.).

DERMASEA, Haw. Enum. Saxifr. 8 (1821) = **Saxifraga**, Tourn.
elongata, Haw. l. c. 9 = S. virginensis.
nivalis, Haw. l. c. 8 = S. nivalis.
pilosa, Haw. l. c. = S. virginensis.

DERMATOCALYX, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 29. *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 948.
parviflora, *Oerst. l. c.*—Costa Rica.

DERMATOPHYLLUM, Scheele, in Linnæa, xxi. (1848) 455 = **Sophora**, Linn. (Legum.).
Schomburgkii, Sch. Bip. in Schomb. Faun. et Fl. Gui. 1079, nomen.—Guiana.
speciosum, Scheele, in Linnæa, xxi. (1848) 459 = S. secundiflora.

DERMOPHYLLA, Silva Manso, Enum. 30 (1836) = **Cayaponia**, Silva Manso (Cucurb.).
elliptica, Silva Manso, l. c. 32 = C. diffusa.
pendulina, Silva Manso, l. c. 31.—Bras.

DERMOMERA, Reichb. f. Pollr. Orch. Comm. 29 (1852) = **Habenaria**, Willd. (Orch.).
squamata, Reichb. f. l. c. = H. sphylla.

DEROSIPHIA, Rafin. Sylva Tellur. 101 (1838) = **Dissotis**, Benth. (1849) (Melast.).
tuberosa, Rafin. l. c. = Diss. tuberosa.

DEROUETIA, Boiss. & Bal. Diagn. Ser. II. v. 114 (1856) = **Crepis**, Vaill. (Compos.).
Bourgaei, Boiss. Fl. Orient. iii. 883 (= *Lagoseris frigida*).—Lycia.
frigida, Boiss & Bal. Diagn. Ser. II. v. 114 (= *Lagoseris frigida*).—Cappadoc.
robertioides, Boiss. l. c. 115 (= *Lagoseris frigida*).—Cilicia.

DERRIS, Lour. Fl. Cochinch. ii. 483 (1790). *LEGUMINOSÆ*, Benth. & Hook. f. i. 569.
AGANOPE, Miq. Fl. Ind. Bat. i. 1. 151 (1855).
BRACHYPTERUM, Benth. in Ann. Wien. Mus. ii. (1838) 101.
CYLICODMA, Neck. Elem. iii. 88 (1790).
DEGUELIA, Aubl. Pl. Gui. 750. t. 300 (1775).
acuminata, *Benth. in Miq. Fl. Yungh.* 262, in adnot.—Reg. Himal.
acuminata, *Miq. Fl. Ind. Bat.* i. 144.—Java.
affinis, Benth. in Miq. Pl. Jungh. 262, in adnot. = uliginosa!
amœna, *Benth. l. c.*—Burma; Malaya.
Benthamii, Thw. Enum. Pl. Zeyl. 61 = paniculata.
brachypteris, *Baker, in Oliver, Fl. Trop. Afr.* ii. 246.—Afr. trop.
brevipes, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 244.—Ind. or.
canarensis, *Baker, l. c.* 246.—Ind. or.
chinensis, *Benth. in Journ. Linn. Soc. iv. Suppl.* (1860) 104.—China.
Cumingii, *Benth. l. c.*—Ins. Philipp.
cuneifolia, *Benth. in Miq. Pl. Yungh.* 258.—Reg. Himal.; Malaya.
dalbergioides, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 241.—Burma; Malaya.
discolor, *Benth. in Journ. Linn. Soc. iv. Suppl.* (1860) 111.—Reg. Himal.
elegans, *Benth. in Miq. Pl. Yungh.* 252, in adnot.—Burma.
elliptica, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 111.—Burma; Malaya.
ensiata, *Bedd. Ic. Pl. Ind. Or.* i. 42.—Ind. or.
ferruginea, *Benth. in Miq. Pl. Yungh.* 252, in adnot.—Reg. Himal.
floribunda, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 152; *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 105.—Java.
floribunda, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 68 = sinuata.
Fortunensis, Blume, ex Miq. Fl. Ind. Bat. i. 1. 144. t. 8 = uliginosa.
glabrata, *Walp. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 244.—Afr. trop.
guianensis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 106; *et in Mart. Fl. Bras.* xv. 1. 288.—Bras. et Guiana.
Hayneana, *Benth. in Miq. Pl. Yungh.* 252, in adnot.—Ind. or.
javanica, *Miq. Fl. Ind. Bat.* i. 1. 143.—Java.
Korthalsiana, *Blume, ex Miq. l. c.*—Borneo.
laxiflora, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 105.—Formosa.
longifolia, *Benth. l. c.* 107; *et in Mart. Fl. Bras.* xv. 1. 289.—Bras.
lucida, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 243.—Afr. trop.

DERRIS:—
macrolobæ, *Miq. Fl. Ind. Bat. Suppl.* 297.—Sumatra.
macrophylla, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 114.—Java.
Maingayana, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 245.—Singapore.
marginata, *Benth. in Miq. Pl. Yungh.* 252, in adnot.—Reg. Himal.
micropters, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 113.—Reg. Himal.
montana, *Benth. in Miq. Pl. Yungh.* 253.—Java.
multiflora, *Benth. l. c.*—Ins. Philipp.
negrensis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 107; *et in Mart. Fl. Bras.* xv. 1. 288.—Bras.
oblonga, *Benth. l. c.* 112.—Ind. or.
ovalifolia, Benth. in Miq. Pl. Jungh. 252 = oblonga.
paniculata, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 105.—Zeylan.
parviflora, *Benth. l. c.*—Zeylan.
pinnata, Lour. Fl. Cochinch. 483 = tamarindifolia.
platypters, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 245.—Ind. or.
polyanthra, *Miq. Fl. Ind. Bat. Suppl.* 298.—Sumatra.
polyphylla, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 104.—Java.
polystachya, *Benth. l. c.* 114.—Reg. Himal.
prepartbra, C. Muell. in Walp. Ann. vii. 797, sphalm. = pyrrothyrsa.
pubinervis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 109.—Ins. Timor.
pubipetala, *Miq. Fl. Ind. Bat.* i. 1. 145.—Java.
pyrrothyrsa, *Miq. l. c. Suppl.* 297.—Sumatra.
robusta, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 104.—Ind. or.
scandens, *Benth. l. c.* 108.—Austral.
secunda, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 247.—Reg. Himal.
sinuata, *Benth. ex Thw. Enum. Pl. Zeyl.* 93.—Burma; Malaya.
Spanoghena, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 145.—Ins. Timor.
thyrsiflora, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 114.—Reg. Himal.; Malaya.
timorensis, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 138.—Timor; Java; N. Guinea.
trifoliata, Lour. Fl. Cochinch. 488, ed. Willd. ii. 525 = uliginosa.
uliginosa, *Benth. in Miq. Pl. Yungh.* 252.—Geront. trop.
vestita, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 243.—Malaya.
Wightii, *Baker, l. c.* 247.—Ind. or.

DERWENTIA, Rafin. Fl. Tellur. iv. 55 (1836) = Veronica, Tourn. (Scrophul.).
perfoliata, Rafin. l. c. = V. perfoliata.
sanctolina, Rafin. l. c. = V. Derwentia.

DESCANTARIA, Schlecht. in Linnæa, xxvi. (1853) 140 = **Tradescantia**, Linn. (Commelin.).
Balbisii, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 309 = T. elongata.
cumanensis, Schlecht. in Linnæa, xxvi. (1853) 140 = T. cumanensis.
diuretica, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 304 = T. elongata.
grevenkorea, Schlecht. in Linnæa, xxvi. (1853) 141 = T. multiflora.
Schlechtendalii, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 304 = T. elongata.

DESCHAMPSIA, Beauv. Agrost. 91. t. 8. f. 8 (1812). *GRAMINEÆ*, Benth. & Hook. f. iii. 1187.
AIRIDIUM, Steud. Syn. Pl. Gram. 432 (1854).
AVENELLA, Parl. Fl. Ital. i. 244 (1850).
CAMPELLA, Link, Hort. Berol. i. 122 (1827).
CERESIARIA, Tsrcs. ex Ledeb. Fl. Ross. iv. 452 (1853).
LERCHENFELDIA, Schur, Enum. Pl. Transs. 758 (1866).
MONANDRARIA, E. Desv. in C. Gay, Fl. Chil. vi. 341. t. 79 (1853).
PERIBALLIA, Trin. Fund. Agrost. 132 (1820).
PSYGMOPHYLLUM, Fourn. ex Benth. & Hook. f. Gen. iii. 1185 (1883).
PODOSCHOPUS, Dulac, Fl. Hautes-Pyr. 82 (1867).
RETZIOPSEBMA, Steud. Syn. Pl. Gram. 433 (1854).
VAHLODEA, Fries, in Bot. Notiser, (1842) 176.
aenea, Trin. ex Ledeb. Fl. Ross. iv. 451 = caespitosa.

DESCHAMPSIA :—
airæformis, *Benth. & Hook. f. Gen.* iii. 1158.—Chili.
alba, *Roem. & Schult. Syst.* ii. 688.—Groenland.
alpigena, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 326 = cæspitosa.
alpina, *Roem. & Schult. Syst.* ii. 686.—Rossia.
ambigua, *Beauv. Agrost.* 91.—Am. bor.
andina, *Phil. in Anal. Univ. Chil.* (1878) 564.—Chili.
Andræi, Schur, Enum. Pl. Transs. 753 = cæspitosa.
antarctica, E. Desv. in C. Gay, Fl. Chil. vi. 338 = Aira antarctica.
antarctica, *Hook. f. in Phil. Trans.* clxviii. (1879) 18. —Ins. Antarct.
arctica, Regel, in Bull. Soc. Nat. Mosc. xli. (1868) ii. 800 = Bromus crinitus.
argentea, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) reimpr. 7.—Ins. Canar. ; Azor.
arundinacea, Spreng. Nov. Prov. 16 = Danthonia Forskâlii.
atropurpurea, *Scheele, in Flora*, xxvii. (1844) i. 56.— Reg. temp.
australis, *Nees, ex Steud. Syn. Pl. Gram.* 220.—Ins. Hawaii.
bastica, *Willk. & Lange, Prod. Fl. Hisp.* i. 66.— Hispan.
Befteroniana, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 66.—Chili.
Bicheriniana, Koern. & Schult. Syst. Mant. ii. 380 = cæspitosa.
bottnica, Trin. Fund. Agrost. 158 = cæspitosa.
brevifolia, K. Br. in App. Parry, Voy. 101 = cæspitosa.
cæspitosa, *Beauv. Agrost.* 91. t. 18. f. 3.—Cosmop. temp.
calycina, *Ÿ. & C. Presl, Rel. Haenk.* i. 251.—Am. bor.
danthonioides, Munro, in Benth. Pl. Hartw. 342 = calycina.
discolor, Roem. & Schult. Syst. ii. 686 = juncea, flexuosa.
Duriæana, S. F. Gray, Nat. Arr. Brit. Pl. ii. 137 = cæspitosa !
elongata, *Munro, in Benth. Pl. Hartw.* 342.—Am. bor.
flexuosa, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 66. —Reg. bor. temp.
foliosa, *Hack. Cat. Gram. Portugal.* 33.—Ins. Azor.
glomerata, G. Don, ex Loud. Hort. Brit. 28 = cæspitosa !
gracilis, *Vasey, in Coult. Bot. Gaz.* x. (1885) 224.— Calif.
grandiflora, *Nees, ex Steud. Syn. Pl. Gram.* 220.— Patagon.
grandis, Bess. ex Ledeb. Fl. Ross. iv. 422 = bottnica.
holciformis, *Ÿ. & C. Presl, Rel. Haenk.* i. 251.—Calif.
involucrata, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 66.—Hispan.
juncea, *Beauv. Agrost.* 91.—Europ. ; Am. bor.
Kingii, *E. Desv. in C. Gay, Fl. Chil.* vi. 336.—Chili.
koelerioides, *Benth. in Journ. Linn. Soc.* xix. (1881) 96.—Mexic.
koelerioides, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) ii. 399.—As. temp.
laevigata, Roem. & Schult. Syst. ii. 687 = alpina.
laevis, *Schult. Mant.* ii. 380.—Lusitan.
lasiantha, *Phil. in Linnaea*, xxxiii. (1864–65) 290.— Chili.
latifolia, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 413 = cæspitosa.
latifolia, *Phil. in Linnaea*, xxix. (1857–58) 91.—Chili.
latifolia, *Vasey, Cat. Gram. U. St.* 58.—Calif.
laxa, *Phil. in Linnaea*, xxix. (1857–58) 91.—Am. austr.
littoralis, Rent. Cat. Pl. Genèv. ed. II. 326 = cæspitosa.
Mathewsii, *Bull, in Journ. Linn. Soc.* xxii. (1886) 60. —Peruv.
media, Roem. & Schult. Syst. ii. 687 = juncea.
montana, G. Don, ex Loud. Hort. Brit. 28 = Aira montana.
montana, Parl. Fl. Ital. i. 241 = bottnica !
montana, Schur, Enum. Pl. Transs. 753 = cæspitosa.
nitida, J. & C. Presl, Rel. Haenk. i. 251 = Deyeuxia juncifformis.
parviflora, Beauv. Agrost. 91, 160 = cæspitosa.
parvula, *E. Desv. in C. Gay, Fl. Chil.* vi. 339.—Chili.
pulchella, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 66 = Aira lendigera.
pulchra, [*Nees, in*] *Meyen, Reise*, i. 311, 312, nomen.— Chili.
refracta, *Roem. & Schult. Syst.* ii. 687.—Hispan.
setacea, Hack. Cat. Gram. Portugal. 33 = Aira setacea.
stricta, *Hook. f. e. t.* 16.—Lusit.
subdigitata, Parl. Fl. Palerm. i. 101 = juncea.

DESCHAMPSIA :—
Thuillieri, Gren. & Godr. Fl. Fr. iii. 508 = Aira setacea.
uliginosa, *Scheele, in Flora*, xxvii. (1844) 56.—Europ.
Wahlenbergiana, Scheele, l. c. = bottnica !
Wibeliana, Parl. Fl. Palerm. i. 101 = Aira Wibeliana.
Wirbeliana, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 326 = cæspitosa.

DESCLIAEA, [Moç. & Sesse] ex DC. Prod. iv. 488 (1830) = **Symphoriocarpos**, Dill. (Caprifol.).
leucocarpa, [Moç. & Sesse] ex DC. l. c. (= *Margaris nudiflora*).—Mexic.

DESCURAINIA, Webb & Berth. Phyt. Canar. i. 72 (1836) = **Sisymbrium**, Linn. (Crucif.).
Boryi, Des Moul. in Actes Acad. Bord. vii. (1845) 353 = S. pinnatifidum.
Irio, Webb & Berth. Phyt. Canar. i. 73 = S. Irio.
millefolia, Webb & Berth. l. c. = S. Irio !
Peyrusana, Des Moul. in Actes Acad. Bord. vii. (1845) 353 = S. pinnatifidum.
pinnatifida, Webb, ll. Hisp. 75 = Braya pinnatifida.

DESFONTAENA, Vell Fl. Flum. 95 (1825) = **Argithamnia**, Sw. (Euphorb.).
tricocca, Vell. l. c. ; ii. t. 146 = A. tricocca.

DESFONTAINA, Steud. Nom. ed. II. i. 493 (1840) = prec.

DESFONTAINEA, H. B. & K. Nov. Gen. et Sp. vii. 278 (1825) = **Desfontainia**, Ruiz & Pav. (Logan.).

DESFONTAINEA, Reichb. Nom. 195 (1841) = DESFONTAENA, Vell. = **Argithamnia**, Sw. (Euphorb.).

DESFONTAINIA, Hoffmgg. Verz. Pfl. 56 (1824) = **Fontanesia**, Labill. (Oleac.).

DESFONTAINIA, Ruiz & Pav. Prod. 29 t. 5 (1794). *LOGANIACEAE*, Benth. & Hook. f. ii. 794.
LINKIA, Pers. Syn. i. 219 (1805).
acutangula, Dun. in DC. Prod. xiii. 676 = spinosa.
chilensis, *C. Gay, Fl. Chil.* v. 100.—Chili.
fulgens, D. Don, in Trans. Linn. Soc. xvii. (1837) 505 = spinosa.
Hookeri, Dun. in DC. Prod. xiii. 676 = spinosa.
ilicifolia, Phil. in Linnaea, xxix. (1857–58) 25 = spinosa.
parvifolia, *D. Don. in Edinb. Phil. Journ.* (*Jul.-Sept.* 1831) 275.—Peruv.
spinosa, *Ruiz & Pav. Fl. Per.* iii. 47. t. 286.—Peruv.
splendens, Humb. & Bonpl. Pl. Aequin. i. 157 = spinosa.

DESFORGIA, Steud. Nom. ed. II. i. 493 (1840) = DEFFORGIA, Lam. = **Forgesia**, Comm. (Saxifr.).

DESMANTHODIUM, Benth. in Hook. Ic. Pl. t. 1116 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 345.
guatemalense, *Hemsl. Biol. Centr. Am. Bot.* ii. 142.— Guatemal.
ovatum, *Benth. in Hook. Ic. Pl.* sub t. 1116.—Mexic.
perfoliatum, *Benth. l. c.* t. 1116.—Mexic.

DESMANTHUS, Willd. Sp. Pl. iv. 1044, partim (1805). *LEGUMINOSAE*, Benth. & Hook. f. i. 592.
DARLINGTONIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 97.
acinaciformis, Spas. Ic. 60 ; ex Linnaea, xv. (1841) 198 = Neptunia acinaciformis.
acuminatus, *Benth. in Hook. Journ. Bot.* iv. (1842) 357.—Am. bor. occ.
Adenanthera, Wall. Cat. n. 5298 = Neptunia plena.
arborescens, *Boj. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 587.—Madag.
brachylobus, *Benth. in Hook. Journ. Bot.* iv. (1842) 356.—Am. bor.
callistachys, *DC. Prod.* ii. 445.—Hab. ?
casuarinæ, Steud. Nom. ed. II. i. 493 = Entada polystachya.
cinereus, Willd. Sp. Pl. iv. 1048 = Dichrostachys cinerea.
Commersonianus, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 961.—Madag.
comosus, *A. Rich. Ess. Fl. Cub.* 475.—Cuba.

DESMANTHUS :—
depressus, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1046. —Am. bor. et trop.
diffusus, Willd. l. c. = depressus ?
divergens; Willd. l. c. 1048 = Dichrostachys nutans.
falcatus, Scheele, in Linnaea, xxi. (1848) 455 = brachylobus.
incurvus, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 84.—Mexic.
Jamesii, *Torr. & Gray, Fl. N. Am.* i. 402.—Am. bor. occ.
lacustris, Turr. & Gray, l. c. = Neptunia tenuis.
lacustris, Willd. Sp. Pl. iv. 1044 = Neptunia oleracea.
leptolobus, *Torr. & Gray, Fl. N. Am.* i. 402.—Am. bor. occ.
leptophyllus, H. B. & K. Nov. Gen. et Sp. vi. 264 = virgatus.
leptostachys, DC. Lég. Mém. xii. 443 = Dichrostachys nutans.
natans, Willd. Sp. Pl. iv. 1044 = Neptunia oleracea.
nutans, DC. Prod. ii. 446 = Dichrostachys nutans.
obtusus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 371.—Texas.
palustris, *Bertol. f. in Mem. Acc. Sc. Bolog.* iii. (1851) 266.—Afr. trop.
pedunculatus, Buckl. in Proc. Acad. Sc. Philad. 1861, (1862) 454 = velutinus.
Pervillseanus, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 961.—Madag.
plenus, Willd. Sp. Pl. iv. 1045 = Neptunia plena.
polygynius, DC. Prod. ii. 444 = Neptunia plena.
pratarum, Macfad. Fl. Jamaic. i. 211 = depressus.
punctatus, Willd. Sp. Pl. iv. 1047 = Neptunia plena.
pusillus, Zipp. ex Span. in Linnaea, xv. (1841) 198 = Neptunia acinaciformis.
reticulatus, *Benth. in Hook. Journ. Bot.* iv. (1842) 357. —Am. bor. occ.
rhombifolius, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 454 = reticulatus.
salinarum, Steud. Nom. ed. I. 269 = Prosopis julifora.
stolonifer, DC. Prod. ii. 444 = Neptunia oleracea.
strictus, Benth. ex Steud. Nom. ed. II. i. 493 = virgatus.
tenellus, DC. Prod. ii. 445 = depressus.
trichostachyus, DC. Mém. Lég. xii. 444. t. 67 = Dichrostachys nutans.
triquetrus, Willd. Sp. Pl. iv. 1045 = Neptunia triquetra.
tripennus, *Span. Ic.* 62 ; *ex Linnaea*, xv. (1841) 199 (Neptuniæ sp.).—Ins. Timor.
velutinus, *Scheele, in Linnaea*, xxi. (1848) 455.— Mexic.
virgatus, *Willd. Sp. Pl.* iv. 1047.—Am. bor. ; Ind. occ.

DESMAZERIA, Dum. Obs. Gram. Belg. 46 (1823) = **Demarseria**, Dum. (Gramin.).

DESMESIA, Rafin. Fl. Tellur. iii. 68 (1836) (Aroid.). *orixensis*, Rafin. l. c. = Typhonium trilobatum.
venosum, Rafin. l. c. = Sauromatum guttatum.

DESMIA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 153 = **Brica**, Lam.
aequalis, D. Don, l. c. = E. aequalis.
conferta, D. Don, l. c. = E. conferta.
polifolia, D. Don, l. c. = E. polifolia.

DESMIDOCHUS, Reichb. Consp. 130 (1828) = **Boucerosia**, Wight & Arn. (Asclep.).

DESMIDORCHIS, Ehrenb. in Linnaea, iv. (1829) 94 = **Boucerosia**, Wight & Arn. (Asclep.).
acutangula, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 265 = B. acutangula.
crenulata, Decne. l. c. = B. crenulata.
Forskâlii, Decne. l. c. = B. Forskâlii.
pauciflora, Decne. l. c. 266 = B. pauciflora.
retispiciens, Ehrenb. in Abh. Acad. Berl. (1831) 83 = B. Russeliana.
umbellata, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 266 = B. umbellata.

DESMITUS, Rafin. Sylva Tellur. 139 (1838) = **Camellia**, Linn. (Ternstroem.).
reticulata, Rafin. l. c. = C. reticulata.

DESMOCARPUS, Wall. Cat. n. 6978 (1832) = **Cadaba**, Forsk. (Cappar.).
missionis, Wall. l. c. = C. trifoliata.

DESMOCEPHALUM. Hook. f. in Proc. Linn. Soc. i. (1846) 277; et in Trans. Linn. Soc. xx. (1847) 208 = **Elvira,** Cass. (Compos.).
lanigerum, Hook. f. in Trans. Linn. Soc. xx. (1847) 209.—Ins. Galap.

DESMOCEPHALUS, Benth. & Hook. f. Gen. ii. 348 (1873) = praec.

DESMOCHAETA, DC. Cat. Hort. Monsp. 101 (1813) = **Pupalia,** Juss. (Amarant.).
achyranthoides, H. B. & K. Nov. Gen. et Sp. ii. 210 = Cyathula achyranthoides.
alternifolia, DC. Cat. Hort. Monsp. (1813) 103 = Digera arvensis.
atropurpurea, DC. l. c. 102 = P. atropurpurea.
micrantha, DC. Cat. Hort. Monsp. (1813) 102 = Cyathula geniculata.
muricata, DC. l. c. 108 = Digera arvensis.
orbiculata, Wall. Cat. n. 6936 = P. orbiculata.
patula, Roem. & Schult. Syst. v. 550 = P. lappacea.
patula, Wall. Cat. n. 6937 = Cyathula prostrata.
prostrata, DC. Cat. Hort. Monsp. (1813) 102 = Cyathula geniculata.
repens, Roem. & Schult. Syst. v. 552 = Cyathula prostrata.
sanguinolenta, Link, Enum. Hort. Berol. i. 154 = P. lappacea.
sepium, A. Br. in Flora, xxiv. (1841) 288 = Cyathula tomentosa.
sordida, Bunbury, in Proc. Linn. Soc. i. (1841) 109 = Cyathula achyranthoides?
tomentosa, Roem. & Schult. Syst. v. 554 = Cyathula tomentosa.
uncinata, Roem. & Schult. l. c. = Cyathula achyranthoides.
velutina, Wall. Cat. n. 6935 = P. lappacea.
xanthioides, A. Br. in Flora, xxiv. (1841) 285 = P. lappacea.

DESMOCLADUS, Nees, in Lehm. Pl. Preiss. ii. 56 (1846) = **Lonchostyla,** R. Br. (Restiac.).
Brunonianus, Nees, l. c. = L. fasciculata.

DESMODIUM, Desv. Journ. Bot. i. (1813) 122. t. 5 *LEGUMINOSAE,* Benth. & Hook. f. i. 519.
CATENARIA, Benth. in Miq. Pl. Jungh. i. 220 (1852).
CODARIOCALYX, Hassk. in Flora, xxv. (1842) 11. Beibl. 48.
CYCLOMORIUM, Walp. Rep. ii. 890 (1843).
DENDROLOBIUM, Benth. in Miq. Pl. Jungh. i. 215 (1852).
DICRAMA, DC. Mém. Lég. 326 (1825); et Prod. ii. 329.
DOLLINERA, Endl. Gen. 1285 (1841).
EDUSARON, Medic. in Verles. Charpf. Phys. Gen. ii. (1787) 371.
HETZEROLOMA, Desv. ex Retchb. Consp. 152 (1828).
LAGOTIA, C. Muell. in Walp. Ann. iv. 409 (1857), sphalm.
MEIBOMIA, Heist. ex Adans. Fam. ii. 309 (1763).
NICOLSONIA, DC. Prod. ii. 325; Mém. Lég. 311. t. 51 (1825).
OTOTROPIS, Nees, in Del. Sem. Hort. Vratisl. (1838). —Cf. Linnaea, xiii. (1839) Litt. 130.
OXYDIUM, J. J. Bena. Pl. Jav. Rar. 156 (1838).
PERROTTETIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 98.
PHYLLODIUM, Desv. Journ. Bot. i. (1813) 122. t. 5.
PSEUDOLOBUS, Jaume St. Hil. in Nouv. Bull. Soc. Philom. iii. (1813) 193.
PTEROLOMA, DC. Prod. ii. 326, in textu (1825); Benth. in Miq. Pl. Jungh. i. 219 (1852).
SAGOTIA, Walp. in Linnaea, xxiii. (1850) 787.
TETRANEMA, Sweet, Hort. Brit. ed. II. 149 (1830).
TROPITOMA, Rafin. New Fl. Am. ii. 19 (1836).
abyssinicum, DC. Prod. ii. 338 = D. Scalpe.
acanthocladum, *F. Muell. Fragm.* ii. 122.—Austral.
acrocarpum, Hance, in Hook. Lond. Journ. Bot. vii. (1848) 478 = trigonetrum.
acuminatum, *DC. Prod.* ii. 329.—Am. bor.

DESMODIUM:—
adhaesivum, *Schlecht. in Linnaea,* xii. (1838) 514.—Mexic.
adscendens, *DC. Prod.* ii. 329.—Ind. occ.
affine, *Schlecht. in Linnaea,* xii. (1838) 512.—Mexic.
Aitkinianum, Beck, Bot. U. St. 84 = canescens.
alatum, DC. Mém. Lég. 321; Prod. ii. 326 = triquetrum.
Alamani, *DC. Prod.* ii. 330.—Mexic.
albiflorum, *Salzm. ex Benth. & Gerst. in Kjoeb. Vidensk. Meddel.* (1853) 17, nomen; *et Mart. Fl. Bras.* xv. i. 99.—Am. trop.
altissimum, *Sweet, Hort. Brit.* ed. II. 151, nomen.—Ind. or.
ambiguum, *Hemsl. Biol. Centr. Am. Bot.* i. 274.—Mexic.
amoenum, Wall. Cat. n. 5795 = concinnum.
amplexicaule, *Zoll. & Mor. in Nat. en Genees. Arch. Nederl. Indie,* iii. (1846) 77.—Java.
amplifolium, *Hemsl. Biol. Centr. Am. Bot.* i. 274.—Mexic.
ancistrocarpum, DC. Prod. ii. 331 = incanum.
Andersoni, *Seem. Fl. Vit.* 56.—N. Caled.
angulatum, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 101 = floribundum.
angulatum, Wall. Cat. n. 5729 M = Lespedeza eriocarpa.
angulatum, Wall. l. c. n. 5739, ex parte = polycarpum.
angustifolium, *DC. Prod.* ii. 328.—Mexic.
annuum, A. Gray, Pl. Wright. ii. 46 = spirale.
Aparines, DC. Prod. ii. 330 = uncinatum.
Aparines, Hassk. Pl. Jav. Rar. 366 = spirale.
arboreum, Sweet, Hort. Brit. ed. II. 151 = Uraria Lagopus.
arenarium, *H. B. & K. Nov. Gen. et Sp.* vi. 527.—Venezuela.
argenteum, Wall. Cat. n. 5713 = tiliaefolium.
arisonicum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 363.—Arizona.
asperum, *Desv. Journ. Bot.* i. (1813) 122.—Am. trop.
auricomum, *R. Grah. in Wall. Cat.* n. 5704.—Ind. or.
auriculatum, DC. Mém. Lég. 321 = triquetrum.
australe, DC. Prod. ii. 326 = umbellatum.
australe, Hassk. Pl. Jav. Rar. 356 = D. Cephalotes.
axillare, *DC. Prod.* ii. 332.—Ind. occ.
bambusetum, Miq. Fl. Ind. Bat. i. 1. 256 = Gardneri.
barbatum, *Benth. & Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 18.—Am. trop.
barbatum, Wall. Cat. n. 5794 = confertum.
Barclayi, *Benth. Bot. Voy. Sulph.* 88.—N. Granat.
batocaulon, *A. Gray, Pl. Wright.* ii. 47.—Am. bor. occ.
betuloides, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 48.—Ind. or.
biarticulatum, *F. Muell. Fragm.* ii. 121.—As. trop.; Austral.
bicolor, Wall. Cat. n. 5719 = laxiflorum.
bifforum, *DC. Prod.* ii. 326.—Ind. or.
Bigelovii, A. Gray, Pl. Wright. ii. 47 = neo-mexicanum.
Boivinianum, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 481.—Madag.
Bostii, Torr. ex S. Wats. Bibliog. Ind. N. Am. Bot. 218 = Dillenii.
brachypodum, *A. Gray, Bot. U. St. Expl. Exped.* i. 484.—Austral.
brachystachyum, *R. Grah. ex Benth. in Miq. Pl. Jungh.* 220.—Ind. or.
brachypodum, *Schlecht. in Linnaea,* xii. (1838) 521. —Mexic.
brasetanum, *Micheli, in Kjoeb. Vidensk. Meddel.* (1872) 78.—Bras.
brecteum, DC. Prod. ii. 329 = cuspidatum.
brasiliense, Lond. Hort. Brit. 310 = Aeschynomene brasiliana.
brevipes, Vog. in Linnaea, xii. (1838) 100 = cuneatum.
Burgeri, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 45 = polycarpum.
bullenanum, G. Don, Gen. Syst. ii. 294 = triflorum.
canescens, DC. Prod. ii. 328 = heterophyllum.
capense, Eckl. & Zeyh. Enum. 261 = D. Scalpe.
capuniifolium, *DC. Prod.* ii. 331.—Am. trop.
caillifolia, Hemsl. Biol. Centr. Am. Bot. i. 276.—Mexic.
campylocaulon, *F. Muell. ex Benth. Fl. Austral.* ii. 226.—Austral.
campylolobatum, *Hemsl. Biol. Centr. Am. Bot.* i. 276.—Nicaragua.
canadense, *DC. Prod.* ii. 326.—Am. bor.
canescens, *DC. l. c.*—Am. bor.
capitatum, *DC. l. c.* 326.—Malaya.
caripense, *G. Don, Gen. Syst.* ii. 292.—Venezuela.

DESMODIUM:—
cateniformum, Am. in Nov. Act. Nat. Cur. xviii. i. (1836) 321 = laburnifolium.
cavdatum, *DC. Prod.* ii. 327.—Japon.
Cephalotes, *Wall. Cat.* n. 5721.—As. trop.
Chamissonis, Vog. in Linnaea, x. (1835–36) 588 = spurale.
cillare, *DC. Prod.* ii. 329.—Am. bor.
ciliatum, *DC. l. c.* 338.—Afr. austr.
ciliorum, Hook. Comp. Bot. Mag. i. (1835) 28 = cillare?
cinerascens, *A. Gray, Pl. Wright.* ii. 46.—N. Mexic.
cinereum, *DC. Prod.* ii. 330.—Mexic.
cinereum, Poepp. ex Griseb. Fl. Brit. W. Ind. 187 = D. Scorpiurus.
cocruleo-violaceum, DC. Prod. ii. 331 = barbatum.
coeruleum, G. Don, Gen. Syst. ii. 293 = adscendens.
collinum, Loud. Hort. Brit. 310 = gangeticum.
collinum, Wall. Cat. n. 5693 = latifolium.
concinnum, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 101.—Reg. Himal.; Burma.
confertum, *DC. l. c.*—Reg. Himal.
congestum, Benth. in Miq. Pl. Jungh. 222 = Lepto-desmia congesta.
congestum, Wall. Cat. n. 5723 = D. Cephalotes.
cordistipulum, *Hemsl. Biol. Centr. Am. Bot.* i. 277.—Mexic.
cubense, *Griseb. Cat. Pl. Cub.* 78.—Cuba.
cumanense, DC. Prod. ii. 331 = caripense.
cumanense, *Willd. ex Steud. Nom.* ed. II. i. 494.—Venezuela.
cuneatum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 195.—Chili.
cuspidatum, *Hook. Comp. Bot. Mag.* i. (1835) 28.—Am. bor.
dasylobum, *Miq. Fl. Ind. Bat. Suppl.* 305.—Sumatra
dasyphyllum, Miq. Fl. Ind. Bat. i. 1. 253 = Uraria lagopus.
delicatulum, *A. Rich. Tent. Fl. Abyss.* i. 205.—Abyssin.
deltoideum, Walp. Rep. i. 787 = lasiocarpum.
densiflorum, *Hemsl. Biol. Centr. Am. Bot.* i. 277.—Mexic.
denudatum, *DC. Mém. Lég.* 325.—Ins. S. Doming.
dependens, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 248.—Ins. Moluce.; N. Guinea.
dichotomum, DC. Prod. ii. 336 = diffusum.
diffusum, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 100. —Ind. or.; Burma.
diffusum, DC. Prod. ii. 335 = laxiflorum.
Dillenii, *Darl. Fl. Castr.* 414.—Am. bor.
dimorphum, *Wight. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 161.—Afr. trop.
dioicum, *DC. Prod.* ii. 338.—Nepal.
discolor, *Vog. in Linnaea,* viii. (1833) 103.—Bras.
distens, Royle, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 172 = polycarpum.
divergens, *G. Don, Gen. Syst.* ii. 296, nomen.—Ind. or.
diversifolium, *DC. Prod.* ii. 334.—Madag.
diversifolium, Schlecht. in Linnaea, xii. (1838) 513 = incanum.
dolabriforme, *Benth. in Miq. Pl. Jungh.* 226.—Ind. or.
Dregeanum, *Benth. l. c.* 222.—Afr. austr.
dubium, Lindl. Bot. Reg. t. 967 = floribundum.
elatum, *DC. Prod.* ii. 326.—Mexic.
elatum, H. B. & K. Nov. Gen. et Sp. vi. 538 = asperum.
elegans, *Benth. Fl. Hongk.* 85.—China.
elegans, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 100 = tiliaefolium.
elegans, Schlecht. in Linnaea, xii. (1838) 520 = venustum.
elliptifolium, Macfad. Fl. Jamaic. i. 268 = adscendens.
elongatum, Wall. Cat. n. 5715 = laxiflorum.
elongatum, Zoll. in Nat. en Geneesk. Arch. Nederl. Indie, iii. (1846) 56 = Gardneri.
emarginatum, *DC. Prod.* ii. 328.—Ins. Martinic.
erythrocaulon, *DC. l. c.* 331.—Am. austr.
exiguum, A. Gray, Pl. Wright. ii. 46 = neo-mexicanum.
ferrugineum, Wall. Cat. n. 5732 = rufescens.
filiforme, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 227. —Ins. Galapag.
filiforme, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie,* iii. (1846) 77.—Java.
flagellare, *Benth. Fl. Austral.* ii. 228.—Austral.
flexuosum, *Wall. Cat.* n. 5691.—Burma.
floribundum, *G. Don, Gen. Syst.* ii. 297.—Reg. Himal.
floridanum, *Chapm. Fl. S. U. St.* 102.—Florida.

DESMODIUM :—
foliosum, *Hemsl. Biol. Centr. Am. Bot.* i. 278.—Mexic.
formosum, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 29.—China.
fruticulosum, Walp. Rep. i. 787 = mauritianum.
gangeticum, Blanco, Fl. Filip. ed. II. 408 = latifolium.
gangeticum, *DC. Prod.* ii. 327.—As. trop.; Austral.
Gardneri, *Benth. in Mig. Pl. Jungh.* 226.—Ind. or.
Ghiesbreghtii, *Hemsl. Biol. Centr. Am. Bot.* i. 279.—Mexic.
glabellum, *DC. Prod.* ii. 329.—Am. bor.
glabrum, *DC. l. c.* 336.—Honduras.
glaucescens, *Miq. in Linnaea*, xviii. (1844) 569.—Guiana.
gracile, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 182.—Mexic.
Graham, *A. Gray, Pl. Wright.* ii. 48.—N. Mexic.
gramineum, *A. Gray, l. c.* 46.—N. Mexic.
grande, *Kurz, in Journ. Soc. Beng.* xliii. (1874) II. 184.—Burma.
grande, E. Mey. Comm. Pl. Afr. Austr. 124 = paleascens.
grandiflorum, DC. Prod. ii. 338 = acuminatum.
grandifolium, *DC. l. c.*—Ind. or.
granulatum, Walp. Rep. i. 787 = triflorum.
Griffithianum, *Benth. in Miq. Pl. Jungh.* 222.—Reg. Himal.
guianense, *DC. Prod.* ii. 338.—Guiana.
Gunnianum, Benth. ex A. Gray, Bot. U. St. Expl. Exped. i. 435 = varians.
Gunnii, Hook. f. Fl. Tasm. i. 101 = prærc.
gyrans, *DC. Prod.* ii. 326.—As. trop.
gyroides, *DC. Mém. Lég.* 322; Prod. ii. 326.—As. trop.
gyroides, Hask. Pl. Jav. Rar. 362 = polycarpum.
hamosum, Lond. Hort. Brit. 310 = Uraria hamosa.
Hardwickianum, Voigt, Hort. Suburb. Calc. 223 = rhomboideum.
Hartwegianum, *Hemsl. Biol. Centr. Am. Bot.* i. 279.—Mexic.
Helleri, *Peyr. in Linnaea*, xxx. (1859–60) 78—Mexic.
heptaphyllum, *Sweet, Hort. Brit.* ed. II. 151, nomen.—Ind. or.
heterocarpum, DC. Prod. ii. 337 = polycarpum.
heterophyllum, *DC. l. c.* 334.—As. trop.
heterophyllum, *Hook. & Arn. Bot. Beech. Voy.* 417.—Mexic.
heterophyllum, Wall. Cat. n. 5701, C = triflorum.
Hilsenbergii, *Boj. ex Steud. Nom.* ed. II. i. 495, nomen.—Madag.
hippocrepidium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 194 = seq.
Hippocrepis, DC. Prod. ii. 338 = polycarpum.
hirsutum, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 186.—Mexic.
hirtum, *Guill. & Perr. Fl. Seneg. Tent.* 209.—Afr. trop.
Hookerianum, *D. Dietr. Syn. Pl.* iv. 1151.—Mexic.
Horsfieldii, Miq. Fl. Ind. Bat. i. i. 251 = Uraria hamosa.
Humboldtianum, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 481.—Madag.
humifusum, *Beck, Bot. U. St.* 86.—Am. bor.
Illinoense, *A. Gray, in Proc. Am. Acad.* viii. (1870) 389.—Am. bor.
incanum, *DC. Prod.* ii. 332.—Ind. occ.
inflractum, *DC. l. c.* 330.—Mexic.
japonicum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 46 = podocarpum.
javanense, Boj. ex Baker, in Oliver, Fl. Trop. Afr. ii. 166 = paleascens.
jucundum, *Thw. Enum. Pl. Zeyl.* 411.—Zeylan.
karensium, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 238, 239.—Burma.
Kerstenii, *O. Hoffm. in Linnaea*, xliii. (1880–82) 197.—Afr. trop.
Kibalensis, Blume, ex Miq. Fl. Ind. Bat. i. i. 250 = laxiflorum.
labuanifolium, *DC. Prod.* ii. 337.—As. trop.
laburnifolium, Sieber, ex Griseb. Fl. Brit. W. Ind. 187 = cajanifolium.
lactescens, *Boj. Hort. Maurit.* 100.—Ins. Maurit.
lactosum, Steud. Nom. ed. II. i. 495 = prærc.?
lævigatum, *DC. Prod.* ii. 329.—Am. bor.
lamyocarpum, *Hemsl. Biol. Centr. Am. Bot.* i. 280.—Mexic.
lanceolatum, Walp. Rep. i. 787 = gangeticum.
lanceolatum, DC. Prod. ii. 336 = latifolium.

DESMODIUM :—
lateristachyum, Boj. ex Baker, in Oliver, Fl. Trop. Afr. ii. 166 = paleascens.
latifolium, *DC. Prod.* ii. 328.—As. et Afr. trop.
latifolium, Wight, Ic. t. 272 = gangeticum.
laxiflorum, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 100.—As. trop.
laxum, DC. l. c. 102 = podocarpum.
leiocarpum, *G. Don, Gen. Syst.* ii. 294.—Reg. Argent.
leptoclados, *Benth. Hemsl. Diagn. Pl. Nov.* 44.—Mexic.
leptopus, *A. Gray, ex Benth. in Miq. Pl. Jungh.* 226, in obs.—Ins. Philipp.
leptostachyum, Wall. Cat. n. 5697 B = laxiflorum.
Leschenaultii, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 102 = Pseudarthria viscida.
lespedezioides, *Benth. in Miq. Pl. Jungh.* 222.—Madag.
leucanthum, Rafin. New Fl. Am. ii. 19, in obs. (An monstrosa ?).—Am. bor.
lineare, *Hook. Bot. Misc.* iii. (1831) 215.—Peruv.
Lindleyii, Mart. Ausw. Pfl. Muench. t. 17 = incanum.
linearifolium, *DC. Mém. Lég.* 325; *et Prod.* ii. 327.—Hab.?
linearifolium, G. Don, Gen. Syst. ii. 294 = mauritianum.
lineatum, *DC. Prod.* ii. 330.—Am. bor.
lineatum, Span. Ic. 42; ex Linnæo, xv. (1841) 193 = D. Cephalotes.
lupulinum, Schlecht. in Linnæa, xii. (1838) 317 = uncinatum.
lutescens, *DC. Prod.* ii. 326.—China.
macrophyllum, *Desv. Journ. Bot.* i. (1813) 122, nomen.—Hab.?
macropodium, *Hemsl. Diagn. Pl Nov.* 44.—Mexic.
macrostachyum, *Hemsl. l. c.*—Mexic.
maculatum, DC. Prod. ii. 327 = gangeticum.
madrense, *Hemsl. Diagn. Pl. Nov.* 45.—Mexic.
malachophyllum, *DC. Prod.* ii. 338.—Ins. Philipp.
marilandicum, *Boott, ex Darlng. Fl. Cestr.* ed. II. 412.—Am. bor.
marylandicum, DC. Prod. ii. 328 = Dillenii.
mauritianum, *DC. l. c.* 334.—Ins. Maurit.
megaphyllum, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie*, iii. (1846) 77.—Java.
mexicanum, *Sweet, Hort. Brit.* ed. I. 479.—Mexic.
microphyllum, DC. Prod. ii. 337 = parvifolium.
molle, *DC. l. c.* 332.—Am. bor.; Ind. occ.
mollicului, *DC. l. c.* 331.—Am. austr.
monospermum, *Baker, in Journ. Linn. Soc.* xx. (1883) 181.—Madag.
monospermum, Boiv. ex Baill. in Bull. Soc. Linn. Par i. (1884) 481 = Boivinianum.
Muelleri, *Benth. Fl. Austral.* ii. 235.—Austral.
multicaule, *DC. Prod.* ii. 331.—Peruv.
multiflorum, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 101 = floribundum.
natalitium, Sond. in Linnæa, xxiii. (1850) 82 = gangeticum.
Nesel, *DC. Prod.* ii. 338.—Hab.?
nemorosum, *F. Muell. ex Benth. Fl. Austral.* ii. 284.—Austral.
neo-mexicanum, *A. Gray, Pl. Wright.* i. 58.—Mexic.
nervosum, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 28 = polycarpum.
neurocarpum, *Benth. Fl. Australl.* ii. 234.—Austral.
nicaraguense, *Oerst. ex Benth. & Orst. in Kjoeb. Vidensk. Meddel.* (1858) 16.—Nicaragua.
nitidum, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 186.—Mexic.
novo-hollandicum, F. Muell. in Linnæa, xxv. (1852) 394 = Glycine tabacina.
nudiflorum, *DC. Prod.* ii. 330.—Am. bor.
nutans, Wall. Cat. n. 5706 = tiliæfolium.
obcordatum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1878) ii. 230.—Burma.
oblatum, *Baker, ex Kurz, l. c.* 230.—Burma.
oblongifolium, DC. Prod. ii. 332 = axillare ?
oblongum, *Wall. Cat.* n. 5714.—Burma.
obovatum, Wall. l. c. n. 5722 B = capitatum.
obovatum, Vog. in Linnæa, xii. (1838) 106 = adscendens.
obtusum, DC. Prod. ii. 329 = marilandicum.
ochroleucum, *M. A. Curt. ex Canby, in Proc. Acad. Sc. Philad.* (1864) 17.—Am. bor.
Oldhami, *Oliver, in Journ. Linn. Soc.* ix. (1867) 165.—Japon.
orbiculare, *Schlecht. in Linnaea*, xii. (1838) 311.—Mexic.
orbiculatum, *G. Don, Gen. Syst.* ii. 298, nomen.—Hab.?

DESMODIUM :—
orbiculatum, Wall. Cat. n. 5695 = retroflexum.
oriasbanum, *Hemsl. Diagn. Pl. Nov.* 45.—Mexic.
ormocarpoides, *DC. Prod.* ii. 327.—Ind. or.; Malaya.
ospriostreblum, Steud. ex A. Rich. Tent. Fl. Abyss. i. 204 = spirale.
ovalifolium, *Guill. & Perr. Fl. Seneg. Tent.* 208.—Afr. trop.
ovalifolium, Wall. Cat. n. 5730 = polycarpum.
ovalifolium, Walp. Rep. i. 787 = Alysicarpus rugosus.
oxalidifolium, Blume, ex Miq. Fl. Ind. Bat. i. i. 249 = C. Trifoliastrum.
oxalidifolium, G. Don, Gen. Syst. ii. 294 = adscendens.
oxybracteum, DC. Prod. ii. 334 = paleaceum.
oxylobum, C. Muell. in Walp. Ann. vii. 764, sphalm. = dasylobum.
oxyphyllum, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 102—Reg. Himal.
pachyrhiza, *Vog. in Linnaea*, xii. (1838) 97.—Bras.
paleaceum, *Guill. & Perr. Fl. Senog. Tent.* 209.—Afr. trop.
Palmeri, *Hemsl. Diagn. Pl. Nov.* 45.—Mexic.
paniculatum, *DC. Prod.* ii. 329.—Am. bor.
Parkinsoni, *Hemsl. Diagn. Pl. Nov.* 45.—Mexic.
Parryi, *Hemsl. l. c.* 46.—Mexic.
parviflorum, *Baker, in Hook.f. Fl. Brit. Ind.* ii. 172.—Ind. or.
parviflorum, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 185.—Mexic.
parvifolium, Blanco, Fl. Filip. ed. II 408 = triflorum.
parvifolium, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 100.—As. trop.; Austral.
patens, G. Don, Gen. Syst. ii. 298, nomen; Wight, Ic. t. 407 = polycarpum.
pauciflorum, *DC. Prod.* ii. 330.—Am. bor.
pauciovreum, DC in Ann. Sc. Nat. Sér. I. iv. (1825) 101 = concinnum.
pedicellatum, R. Grah. in Wall. Cat. n. 5735 = tortuosum.
pedunculatum, *DC. Prod.* ii. 338.—Am. bor.
penduliflorum, Oudem. Neerl. Plantent. ii. (1866) t. 2 = Lespedeza bicolor.
penduliflorum, Wall. Cat. n. 5727 = concinnum.
pendulum, *F. Muell. Phyt. New Hebr.* 9.—N. Hebrides.
pendulum, Wall. Pl. As. Rar. i. 81. t. 94 = concinnum.
Perrottetii, DC. Mém. Lég. 324; et Prod. ii. 327 = asperum.
peruvianum, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 30.—Peruv.
physocarpos, *Vog. in Linnaea*, xii. (1838) 104.—Bras.
pictum, Walp. Rep. i. 787 = Hedysarum pictum.
plosiuscuium, DC. Prod. ii. 335 = uncinatum.
pilosum, *DC. l. c.* 337.—Japon.
platycarpum, *Benth. in Mart. Fl. Bras.* xv. i. 100.—Bras.
plectocarpum, *Hemsl. Diagn. Pl. Nov.* 46.—Mexic.
plicatum, *Cham. & Schlecht. in Linnaea*, v. (1830) 585.—Mexic.
podocarpum, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 102—Reg. Himal.; China; Japon.
podocarpum, Hook. & Arn. Bot. Beech. Voy. 417. t. 96 = Hookerianum.
podocarpum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 46 = Gardneri.
polycarpum, *DC. Prod.* ii. 334.—As. trop.; Austral.
polycarpum, Wall. Cat. n. 5710 = oxyphyllum.
polycarpum, Wight & Arn. Prod. 227, ex parte = capitatum.
polygonoides, *Welw. ex Baker, in Oliver, Fl. Trop Afr.* ii. 161.—Afr. trop.
polystachyum, *Schlecht. in Linnaea*, xii. (1838) 321.—Mexic.
portoricense, *G. Don, Gen. Syst.* ii. 294.—Porto Rico.
præmorsum, R. Grah. in Wall. Cat. n. 5708 = Lourea obcordata, D. Rottleri.
prehensile, *Schlecht. in Linnaea*, xii. (1838) 315.—Mexic.
Peyoni, DC. Prod. ii. 334 = Wightii.
Pseudo-gyrans, *Miq. Fl. Ind. Bat.* i. i. 244.—Java.
Pseudo-gyroides, Miq. l. c. = gyroides.
Pseudo-triquetrum, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 100 = trigonetrum.
psilocarpum, *A. Gray, Pl. Wright.* ii. 48.—N. Mexic.
psilophyllum, *Schlecht. in Linnaea*, xii. (1838) 310.—Mexic.

DESMODIUM :—
pubens, Young, Fl. Texas, 338; ex S. Wats. Bibliog. Ind. N. Am. Bot. 218 = paucinatum.
pulchellum, *Benth. Fl. Hongk.* 83.—As trop.; Austral.
pulchellum, Blume, ex Miq. Fl. Ind. Bat. i. 1. 242 = polycarpum.
pulcherrimum, Shuttlew. ex Griseb. Fl. Brit. W. Ind. 188 = tortuosum.
punctatum, DC. Prod. ii. 338 = rhynchosia densiflora.
purpureum, Hook. & Arn. Bot. Beech. Voy. 62 = polycarpum.
pycnostachyum, *Benth. in Trans. Linn. Soc.* xxv. (1866) 299.—N. Caled.
quinquangulare, Wight, Ic. t. 298 = diffusum.
quinquangulatum, G. Don, Gen. Syst. ii. 298 = diffusum.
racemiferum, DC. Prod. ii. 331 = adscendens.
racemosum, *DC. l. c.* 337.—Japon.
radiatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 131. —Madag.
radicosa, Macfad. Fl. Jamaic. i. 269 = axillare.
ramosissimum, G. Don, Gen. Syst. ii. 294 = mauritianum.
recurvatum, Wall, Cat. n. 5717 = laxiflorum.
reniforme, *DC. Prod.* ii. 327.—Ind. or.; Malaya.
repandum, DC. l. c. 334 = *D.* Scalpe.
repens, DC. l. c. 338 = lespedeza repens.
reptans, DC. l. c. 333 = axillare.
reticulatum, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 46.—China.
retinens, *Schlecht. in Linnaea*, xii. (1838) 311.—Mexic.
retroflexum, *DC. Prod.* ii. 336.—Reg. Himal.; Burma; China.
retusum, G. Don, Gen. Syst. ii. 297 = polycarpum.
rhombifolium, DC. Prod. ii. 330 = lasvigatum.
rhomboideum, *Sweet, Hort. Brit.* ed. II. 151.—Ind. or.
rhytidophyllum, *F. Muell. ex Benth. Fl. Austral.* ii. 238.—Austral.
rigidum, *DC. Prod.* ii. 330.—Amer. bor.
robustum, E. Mey. Comm. Pl. Afr. Austr. 125 = Pauridarhria Hookeri.
Rottleri, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 174.—Ind. or.; Burma.
Rottleri, G. Don, Gen. Syst. ii. 297 = laxiflorum.
rotundifolium, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 172. —Ind. or.
rotundifolium, *DC. Prod* ii. 330.—Am. bor.
rotundifolium, Mart. ex Benth. n. 5696 = retroflexum.
Roxburghii, Sweet, Hort. Brit. ed. I. 479 = laxiflorum.
Roxburghii, Wall. Cat. n. 5716 = diffusum.
Roxburghii, Walp. Rep. iv. 369 = latifolium.
Roylei, Wight & Arn. Prod. 227 = gyrans.
rubescens, Blume, ex Miq. Fl. Ind. Bat. i. 1. 246 = megaphyllum.
rubigineosum, Benth. in Ann. Nat. Hist. iii. (1839) 484 = asperum.
rubrum, *DC. Prod.* ii. 337.—Cochinch.
rufescens, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 101. —Ind. or.
sagittatum, *DC. Prod.* ii. 337.—Ind. or.
salicifolium, *DC. l. c.* 337.—Ind. or.
salicifolium, Mart. ex Benth. in Mart. Fl. Bras. xv. 1. 106 = Cranocarpus Martii.
Salviæ, *Hemsl. Diagn. Pl. Nov.* 46.—Guatemala.
sambuense, DC. Prod. ii. 335 = floribundum.
sandwicense, E. Mey. Cat. Sem. Hort. Regiom. (1850) 4.—*Cf.* Linnæa, xxiv. (1851) 330 = uncinatum.
scabrellum, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 250 = laxiflorum.
Scalpe, *DC. Prod.* ii. 334.—As. et Afr. trop.
scandens, Blume, ex Miq. Fl. Ind. Bat. i. 1. 246 = megaphyllum.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 205 = *D.* Scalpe.
sclerophyllum, *Benth. in Mart. Fl. Bras.* xv. 1. 102.—Bras.
scoparium, Wall. Cat. n. 5699 = pervifolium.
Scorpiurus, *Desv. Journ. Bot.* i. (1813) 192.—Jamaic.
scutatum, *Hemsl. Biol. Centr. Am. Bot.* i. 287.—Mexic.
securiforme, *Benth. in Miq. Pl. Jungh.* 236.—Ins. Philipp.
semnaerense, *Schweinf. Beitr. Fl. Aethiop.* 8.—Afr. trop.
sequax, *Wall. Pl. As. Rar.* ii. 46.—Reg. Himal.

DESMODIUM :—
sericatum, Presl, Bot. Bemerk. 69 = *D.* Cephalotes.
sericocarpum, *Hemsl. Diagn. Pl. Nov.* 47.—Mexic.
sericophyllum, *Schlecht. in Linnaea*, xii. (1838) 317. —Mexic.
serotinum, *DC. Prod.* ii. 338.—Mexic.
serpens, Wall. Cat. n. 5788 = polycarpum.
serriferum, Wall. l. c. n. 5708 B = oxyphyllum.
sessilifolium, *Torr. & Gray, Fl. N. Am.* i. 368.—Am. bor.
setigerum, E. Mey. Comm. Pl. Afr. Austr. 134 = hirtum.
sikkimense, DC. Prod. ii. 336 = polycarpum.
simplex, *G. Don, Gen. Syst.* ii. 294.—Afr. trop.
Sinclairi, Benth. Bot. Voy. Sulph. 82 = uncinatum.
sinuatum, *Blume, ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 166.—Reg. Himal.; Malaya.
Skinneri, *Benth. in Hemsl. Diagn. Pl. Nov.* 47.—Guatemala.
Sonorense, A. Gray, Pl. Wright. ii. 47 = uncinatum.
spartifolium, G. Don, Gen. Syst. ii. 294 = incanum.
spartioides, DC. Prod. ii. 337 = varians.
spectabile, Miq. in Linnæa, xviii. (1844) 570 = asperum.
spectabile, *Sweet, Hort. Brit.* ed. I. 479.—Ins. Maurit.
spirale, *DC. Prod.* ii. 331.—Cosmop. trop.
spinale, Griseb. Fl. Brit. W. Ind. 188 = neo-mexicanum.
Sprengelii, D. Dietr Syn. Pl. iv. 1154 = spirale.
squarrosum, DC. Prod. ii. 332 = Eriosema squarrosum.
stipulaceum, DC. l. c. 330 = tortuosum.
stipulaceum, Hassk. in Hoev. & De Vriese, Tijdschr. xi. (1844) 75 = pervifolium.
stipulaceum, Wight & Arn. Prod. 229 = trifforum.
stoloniferum, *Steud. Nom.* ed. II. i. 496.—Porto Rico.
strangulatum, Thw. Enum. Pl. Zeyl. 87, partim = Thwaitesii.
strangulatum, Wight & Arn. Prod. 228 = *D.* Scalpe.
striatum, *DC. Prod.* ii. 337.—Japon.
strictum, *DC. l. c.* 328.—Am. bor.
strobilaceum, Benth Pl. Hartw. 11 = Hartwegianum.
strobilaceum, *Schlecht. in Linnaea*, xii. (1838) 316.—Mexic.
subsessilum, *Vog. l. c.* 99.—Bras.
subsessile, *Schlecht. l. c.* 319.—Mexic.
substipulaceum, *Blume, ex Kurz, in Journ. As. Soc Beng.* xlv. (1876) 11. 230.—Burma.
subtile, *Hemsl. Diagn. Pl. Nov.* 47.—Mexic.
sumbuense, G. Don, Gen. Syst. ii. 296 = floribundum.
sulcatum, Wall. Cat. n. 5726 = laxiflorum.
supinum, DC. Prod. ii. 332 = incanum.
sylvaticum, Benth. Pl. Hartw. 116 = spirale.
tenellum, DC. Prod. ii. 338 = spirale.
tenuiflorum, G. Don, Gen. Syst. ii. 294 = mauritianum.
tenue, R. Grah. in Wall. Cat. n. 5731 = polycarpum.
tenuicolum, DC. Prod. ii. 333 = spirale.
tenuifolium, *Torr. & Gray, Fl. N. Am.* i. 363.—Amer. bor.
teres, *Wall. Cat.* n. 5694.—Burma.
terminale, DC. Prod. ii. 327 = molle.
terminale, Guill. & Perr. Fl. Seneg. Tent. 207 = spirale.
Thonningianum, D. Dietr. Syn. Pl. iv. 1147 = Alysicarpus rugosus.
Thunbergii, *DC. Prod.* ii. 337.—Japon.
Thwaitesii, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 169.—Zeylan.
tiliæfolium, *G. Don, Gen. Syst.* ii. 297.—Reg. Himal.; Burma.
timoriense, *DC. Mém. Lég.* 328; *et Prod.* ii. 327.—Ins. Timor.
tomentosum, *DC. Prod.* ii. 337.—Japon.
tortuosum, *DC. l. c.* 333.—Am. bor.; Ind. occ.
tortuosum, Webb, in Hook. Niger Fl. 123 = spirale.
trichocaulon, Hassk. Pl. Jav. Rar. 267 = *D.* Scalpe.
trichocaulon, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 101 = polycarpum.
trichocaulon, *Zoll.* ex Miq. Fl. Ind. Bat. i. 1. 245 = Pseudo-gyrans.
trichostachyum, *Benth. Fl. Austral.* ii. 234.—Austral.
triflorum, *DC. Prod.* ii. 334.—Cosmop. trop.
triflorum, Wall. Cat. n. 5734 *F*, ex parte; Wight & Arn. Prod. 229 = heterophyllum.
Triflolianum, *Miq. Fl. Ind. Bat.* i. 1. 245.—Java.
trigonum, DC. Prod. ii. 332 = uncinatum.
trinerve, R. Grah. in Wall. Cat. n. 5730 = podocarpum.

DESMODIUM :—
trinervium, *Fisch. ex Loud. Hort. Brit.* 310, nomen.—Hab.?
triquetrum, *DC. Prod.* ii. 326.—As. trop.
tuberculatum, G. Don, Gen. Syst. ii. 298 = polycarpum.
tuberosum, G. Don, l. c.; Voigt, Hort. Suburb. Calc. 232 = Pueraria tuberosa.
umbellatum, *DC. Prod.* ii. 335.—As. trop.
uncinatum, *DC. l. c.* 331.—Am. bor. et austr.
varians, *G. Don, Gen. Syst.* ii. 298.—Austral.; N. Caled.
vestitifolium, Steud. Nom. ed. II. i. 496 = incanum.
velutinum, *DC. Prod.* ii. 338.—Am. austr.
venosum, *Vog. in Linnæa*, xii. (1838) 103.—Bras.
venustulum, *Hemsl. Biol. Centr. Am. Bot.* i. 290.—Panama.
venustum, *Steud. Nom.* ed. II. i. 496.—Mexic.
vestitum, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 169.—Burma.
villosum, *DC. Prod.* ii. 338.—Mexic.
villosum, *Hemsl. Biol. Centr. Am. Bot.* i. 291.—Mexic.
violaceum, *G. Don, Gen. Syst.* ii. 298.—Bras.
virgatum, Desv. Journ. Bot. i. (1813) 122 = Lespedeza virgata.
virgatum, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, iii. (1846) 76 = latifolium.
viride, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 29.—China.
viridiflorum, *Beck, Bot. U. St.* 84.—Am. bor.
viridiflorum, DC. Prod. ii. 329 = canescens
viscidum, DC. l. c. 336 = Pseudarthria viscida.
viticiuum, Wall. Cat. n. 5709 = laburnifolium.
Vogelii, Steud. Nom. ed. II. i. 496 = adscendens.
Wade, *DC. Prod.* ii. 331.—Bras.
Walkeri, Arn. in Nov. Act. Nat. Cur. xviii. 1. (1836) 331 = Wightii.
Wallichii, Steud. Nom. ed. II. i. 496 = polycarpum.
Wightii, *R. Grah. in Wall. Cat.* n. 5718.—Ind. or.
Wildenowii, G. Don, Gen. Syst. ii. 296 = diffusum.
Wislizeni, *Engelm. ex A. Gray, Pl. Wright.* i. 53.—Mexic.
Wrightii, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 177.—Texas.
zonatum, Miq. Fl. Ind. Bat. i. 1. 250 = ormocarpoides.

DESMONCHUS, Decf. Cat. Hort. Par. ed. III. 30 (1829) = seq.

DESMONCUS, Mart. Hist. Nat. Palm. ii. 84 (1824). *PALMAE*, Benth. & Hook. f. iii. 949.
acreus, *Drude, in Mart. Fl. Bras.* iii. 11. 307.—Bras.
americanus, *Lodd. ex Loud. Hort. Brit.* 382, nomen. —Ins. S. Vincent.
andicola, *Pasq. Cat. Ort. Bot. Nap.* 36.—Reg. Andin.
ataxiacanthus, *Barb. Rodr. Enum. Palm. Nov.* 35.—Bras.
chinantlensis, *Liebm. in Mart. Hist. Nat. Palm.* iii. 321.—Mexic.
dubius, Lodd. ex Loud. Hort. Brit. 382, nomen = polyacanthos?
grandifolius, *Linden, Illustr. Hortic.* xxviii. (1881) 16. —N. Guinea.
horridus, *Splitg. ex Mart. in Palmet. Orb.* 51.—Guian.
intermedius, *Mart. ex H. Wendl. in Kerch. Palm.* 243. —Hab.?
leptoclonos, *Drude, in Mart. Fl. Bras.* iii. 11. 315.—Bras.
leptospadix, *Mart. Palmet. Orb.* 52.—Bras.
longifolius, *Mart. l. c.*—Peruv.
lophacanthos, *Mart. l. c.* 50.—Bras.
macroscanthos, *Mart. Hist. Nat. Palm.* ii. 86.—Bras.
macroscanthos, Spruce, in Journ. Linn. Soc. xi. (1871) 156 = phenophyllus.
major, *Cruz. ex Griseb. Fl. Brit. W. Ind.* 519.—Ins. Trinit.
melanacanthus, Mart. ex Drude, in Mart. Fl. Bras. iii. 11. 306 = horridus.
mitis, *Mart. Hist. Nat. Palm.* ii. 90.—Bras.
oligacanthus, *Barb. Rodr. Enum. Palm. Nov.* 34.—Bras.
orthacanthos, *Mart. Hist. Nat. Palm.* ii. 87. t. 89.—Bras.
oxyacanthos, *Mart. l. c.* ii. 88.—Ann. trop.

DESMONCUS:—
palustris, Trail, in Journ. Bot. xiv. (1876) 353.—Bras.
panamensis, Linden, Illustr. Hortic. xxvii. (1881) 16.—N. Granat.
phœnophyllus, Drude, in Mart. Fl. Bras. iii. ii. 314.—Bras.
phœnicocarpus, Barb. Rodr. Enum. Palm. Nov. 34 = pyncnacanthos!
polyacanthos, *Mart. Nat. Hist. Palm.* ii. 35. t. 68.—Bras.
polyphyllus, *Pois. ex Desf. Cat. Hort. Par.* ed. III. 80, nomen.—Guiana.
prunifer, *Poepp. in Mart. Hist. Nat. Palm.* ii. 148.—Peruv.
pumilus, *Trail, in Journ. Bot.* xiv. (1876) 353.—Bras.
pycnacanthos, *Mart. Hist. Nat. Palm.* ii. 89.—Bras.
riparius, *Spruce, in Journ. Linn. Soc.* xi. (1871) 156.—Bras.
rudentum, *Mart. Palmet. Orb.* 48.—Bras.
setosus, *Mart. Hist. Nat. Palm.* ii. 89.—Bras.
Wallisi, *Linden, Illustr. Hortic.* xxvii. (1881) 16.—Bras.

DESMONEMA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xx. (1867) 260. *MENISPERMEAE.*
cæfra, *Miers, l. c.*—Afr. austr.

DESMONEMA, Rafin. Atl. Journ. 177 (1838) = **Euphorbia**, Linn.
hirta, Rafin. l. c. 178 = E. pilulifera.

DESMOPHYLA, Rafin. Sylva Tellur. 48 (1838) = **Ehretia**, Linn. (Borag.).
aliena, Rafin. l. c. = E. fasciculata.

DESMOPHYLLUM, Webb & Berth. Phyt. Canar. i. 14 (1836) = **Ruta**, Linn.
pinnatum, Webb & Berth. l. c. = R. pinnata.

DESMOS, Lour. Fl. Cochinch. 353 (1790) = **Unona**, Linn. f. (Anonac.).
chinensis, Lour. l. c. = U. discolor.
cochinchinensis, Lour. l. c. = U. Desmos.

DESMOSCELIS, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 29. *MELASTOMACEAE*, Benth. & Hook. f. i. 740.
calcarata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 84.—Bras.
lychnitides, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 30 = villosa.
stachyoides, Naud. l. c. = villosa.
villosa, *Naud. l. c.*—Bras.

DESMOSCHOENUS, Hook. f. Fl. N. Zel. i. 271 (1853) = Scirpus, Tourn. (Cyperac.).
spiralis, Hook. f. l. c. 272 = S. frondosus.

DESMOSTACHYS, Planch. ex Miers, in Ann. & Mag. Nat. Hist. Ser. II. ix. (1852) 399. *OLACINEAE*, Benth. & Hook. f. i. 350.
acuminata, *Baker, in Journ. Linn. Soc.* xxi. (1884) 383.—Madag.
deltoides, *Baker, l. c.*—Madag.
Planchoniana, *Miers, in Ann. & Mag. Nat. Hist* Ser. II. ix. (1852) 399.—Madag.: Afr. trop.
Renschii, *O. Hoffm. & Hildebr. ex O. Hoffm. Sert. Madag.* 13.—Madag.
tumifolius, *Oliver, Fl. Trop. Afr.* i. 353.—Afr. trop.

DESMOSTEMON, Thw. Enum. Pl. Zeyl. 278 (1861) = **Ostodes**, Blume (Euphorb.).
zeylanicus, Thw. l. c. = O. zeylanicus.

DESMOTRICHUM, Blume, Bijdr. 289. t. 35 (1825) = **Dendrobium**, Sw. (Orchid.).
angulatum, Blume, l. c. 299 = Dendrob. homosymum.
angustifolium, Blume, l. c. 300 = Dendrob. angustifolium.
appendiculatum, Blume, l. c. = Dendrob. appendiculatum.
comatum, Blume, l. c. = Dendrob. comatum.
convexum, Blume, l. c. 301 = Dendrob. convexum.
cymbidioides, Blume, l. c. 302 = Dendrob. cymbidioides.
elongatum, Blume, l. c. = Dendrob. elongatum.
fimbriatum, Blume, l. c. 299 = Dendrob. plicatile.

DESMOTRICHUM:—
geminatum, Blume, Bijdr. 299 = Dendrobium geminatum.
grandiflorum, Blume, l. c. 301 = Dendrob. grandiflorum.
pusillum, Blume, l. c. = Dendrob. pusillum.
triflorum, Blume, l. c. = Dendrob. triflorum.

DESPLATEIA, Bocquill. in Adansonia, vii. (1866–67) 51. *TILIACEAE*, Benth. & Hook. f. i. 235.
subericarpa, *Bocquill. l. c.*—Afr. trop.

DESPRETZIA, Kunth, Rév. Gram. ii. 485. t. 157 (1830) = **Leugitea**, Schreb. (Gramin.).
mexicana, Benth. Pl. Hartw. 82 = Z. Hartwegi.
mexicana, Kunth, Rév. Gram. ii. 485. t. 157 = Z. mexicana.

DESSENIA, Adans. Fam. ii. 285 (1763) = **Gnidia**, Linn. (Thymel.).
daphnefolia, Rafin. Fl. Tellur. iv. 106 = Lasiosiphon daphnefolius!
hirsuta, Rafin. l. c. = Lasiosiphon pubescens!
lanceolata, Rafin. l. c. = G. parviflora.

DESTRUGESIA, Gaudich. Bot. Voy. Bonite, t. 57 (1844–46) = **Capparis**, Linn.
scabrida, Gaudich. l. c. = C. scabrida.

DESTRUGUEZIA, Benth. & Hook. f. Gen. i. 109 (1862) = prec.

DESVAUXIA, R. Br. Prod. 253 (1810) (Devauxia); ed. Nees, 108 (1827) = **Centrolepis**, Labill.
aloyproides, Nees, in Lehm. Pl. Preiss. ii. 71 = C. aloyproides.
aristata, R. Br. Prod. 253 = C. aristata.
Banksii, R. Br. l. c. = C. Banksii.
Billardieri, R. Br. l. c. 252 = C. fasciculata.
brevifolia, Nees, in Lehm. Pl. Preiss. ii. 70 = C. Drummondii.
Drummondiana, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 51 = C. Drummondii.
exserta, R. Br. Prod. 253 = C. exserta.
glabra, F. Muell. in Linnæa, xxviii. (1856) 226 = C. glabra.
longifolia, Gaudich. in Freyc. Voy. Bot. 418 = C. longifolia.
Patersonii, R. Br. Prod. 252 = C. strigosa.
pulvinata, R. Br. l. c. = C. pulvinata.
pusilla, R. Br. l. c. 253 = C. pusilla.
strigosa, R. Br. l. c. 252 = C. strigosa.
tenuior, R. Br. l. c. = C. strigosa.
Urvillei, Steud. Syn. Pl. Cyp. 267 = C. Drummondii.

DETANDRA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiii. (1864) 134; xx. (1867) 17 = **Sychnosepalum**, Eichl. (Menisperm.).
latifolia, Miers, ll. cc. 134; 19.—Bras.
ovata, Miers, ll. cc. = S. microphyllum.

DETARIUM, Juss. Gen. 385 (1789). *LEGUMINOSAE*, Benth. & Hook. f. i. 585.
Beurmanianum, *Schweinf. in Zeitschr. Allg. Erdk.* (1862) 397; *ex C. Muell. in Walp. Rep.* vii. 824.—Afr. trop.
Heudelotianum, Baill. in Adansonia, vi. (1866) 201 = senegalense.
microcarpum, Guill. & Perr. Fl. Senég. Tent. i. 271 = senegalense.
senegalense, *J. F. Gmel. Syst.* 700.—Afr. trop.
zeylanicum, Thw. Enum. Pl. Zeyl. 414 = Crudia zeylanica.

DETHARDINGIA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 80 = **Breweria**, R. Br. (Convolv.).
heterantha, Nees & Mart. l. c.—Bras.

DETHAWIA, Endl. Gen. 775 (1839) = **Sesuli**, Linn. (Umbellif.).
tenuifolia, Endl. l. c. (= *Wallrothia tenuifolia*).—Mont. Fyren.

DETRIDIUM, Nees, Gen. et Sp. Ast. 255 (1833) = **Felicia**, Cass. (Compos.).
Bergianum, Nees, l. c. 256 = Aster Bergianus.

DETRIS, Adans. Fam. ii. 131 (1763) = **Felicia**, Cass. (Compos.).

DEUTZIA, Thunb. Nov. Gen. 19 (1781); Fl. Jap. 10 (1784). *SAXIFRAGEAE*, Benth. & Hook. f. i. 642.
Brunoniana, Wall. Cat. n. 3050 = stamines.
canescens, Hort. ex C. Koch, Dendrol. i. 347 = scabra.
canescens, Siebold, ex Steud. Nom. ed. II. i. 497, sphalm. pro *crenata*.
corymbosa, R. Br. in Wall. Cat. n. 3652 = parviflora.
corymbosa, Lindl. Bot. Reg. (1840) t. 5 = staminea.
crenata, Sieb. & Zucc. Fl. Jap. i. 19 = scabra.
dentata, Hort. ex C. Koch, Dendrol. i. 347 = scabra.
Fortunei, Carr. in Rev. Hortic. (1866) 388 = scabra.
gracilis, *Sieb. & Zucc. Fl. Jap.* i. 22.—Japon.
grandiflora, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1833) 104.—China.
macrantha, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 84.—Reg. Himal.
mexicana, *Hemsl. Diag. Pl. Nov.* 9.—Mexic.
parviflora, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 104.—China.
scabra, Sieb. & Zucc. Fl. Jap. i. 20. t. 7 partim = Siebeldiana.
scabra, *Thunb. Nov. Gen.* 19, 20, 21; *Fl. Jap.* 185.—Japon.
Sieboldi, Koern. in Gartenfl. (1867) 74 = seq.
Sieboldiana, *Maxim., in Mém. Acad. Pétersb.* Sér. VII. x. n. xvi. (1867) 26.—Japon.
staminea, *R. Br. in Wall. Cat.* n. 3655; *et Pl. As. Rar.* ii. 92. t. 191.—Reg. Himal.

DEVAUXIA, Beauv. ex Kunth, Enum. Pl. i. 367, in syn. (1833) = **Glyceria**, Br. (Gramin.).
fluitans, Beauv. ex Kunth, l. c. = G. fluitans.

DEVAUXIA, R. Br. Prod. 258 (1810), sphalm. = DESVAUXIA, R. Br. = **Centrolepis**, Labill.

DEVERRA, DC. Mém. Omb. 45 (1829); Prod. iv. 148 (1830) = **Pituranthos**, Viv. (Umbellif.).
aphylla, DC. Prod. iv. 148.—Afr. austr.
Burchellii, Eckl. & Zeyh. Enum. 347 = *aphylla*.
Burchellii, Sond in Harv. & Sond. Fl. Cap. ii. 549.—Afr. austr.
chlorantha, Coss. & Dur. in Bull. Soc. Bot. Fr. ii. (1855) 248.—Algeria.
juncea, Ball, in Journ. Linn. Soc. xvi. (1878) 468.—Marocco.
Karelkowi, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 589.—Turkestan.
Pituranthos, DC. Prod. iv. 144 = Pituranthos denudatus.
Reboudii, Coss. & Dur. in Bull. Soc. Bot. Fr. ix. (1862) 296.—Algeria.
scoparia, Coss. & Dur. l. c. ii. (1855) 248.—Algeria.
tortuosa, DC. Prod. iv. 148.—Aegypt.
triradiata, Hochst. ex Boiss. Fl. Orient. ii 861.—Arab *virgata*, Coss. & Dur. in Bull. Soc. Bot. Fr. ii. (1855) 248 = *scoparia*.

DEVILLEA, Bert. ex Schult. f. Syst. vii. p. lxvii. 1229 (1830) = **Carraguta**, Lindl. (Bromel.).
speciosa, Bert. l. c. 1229 = C. Berteroniana.

DEVILLEA, Tul. & Wedd in Ann. Sc. Nat. Sér. III. xi. (1849) 107. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 114.
flagelliformis, Tul. & Wedd. l. c.—Bras.

DEWEYA, Torr. & Gray, Fl. N. Am. i. 641 (1840) = **Arrasacia**, Bauer. (Umbellif.).
acaulis, Torr. in Pacif. Rail. Rep. iv. 94 (= *Oreascias dium acaule*).—Am. bor. occ.
arguta, Torr. & Gray, Fl. N. Am. i. 641 = A. arguta (in Addend.).
Hartwegi, A. Gray, in Proc. Am. Acad. vii. (1868) 342.—Am. bor. occ.
Kellogii, A. Gray, l. c. 343.—Calif.
vestita, S. Wats. in Proc. Am. Acad. xvii. (1881–82) 374.—Calif.

DEYEUXIA, Clar. in Beauv. Agrost. 43. t. 9, f. 9, 10 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1152
ACHAETA, Fourn. ex Benth. & Hook. f. Gen. iii 1152 (1883).
BROMIDIUM, Nees & Meyen, in Nov. Act. Acad Nat. Cur. xix. Suppl. i. (1843) 154.
CHAMAECALAMUS, Meyen, Reise, i. 456 (1834).
CINNASTRUM, Fourn. ex Benth. & Hook. f. Gen. iii 1152 (1883).

DEYEUXIA:—
Didymochaeta, Steud. Syn. Pl. Gram. 185 (1854).
Lachnagrostis, Trin. Fund. Agrost. 198 (1819).
Relchela, Steud. Syn. Pl. Gram. 101 (1854).
abietina, Fuss, Fl. Transs. 714 = sylvatica.
acutiflora, *Beauv. Agrost.* 44.—Europ.
arenula, Kunth, Rév. Gram. i. 77 = Forsteri.
aequata, *Benth. Fl. Austral.* vii. 578.—Tasman.
aequivalvis, *Benth. in Journ. Linn. Soc.* xix. (1881) 91.—Am. bor. occ.
agrostides, *Phil. in Linnaea*, xxix. (1857–58) 91.—Chili.
airoides, Roem. & Schult. Syst. ii. 422 = Arundo airoides.
alba, *Y. & C. Presl, Rel. Haenk.* i. 246.—Bolliv.
alestica *Munro, ex Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 845.—Am. bor. occ.
anomala, *Benth. & Hook. f. Gen.* iii. 1158 = plebeia.
anthoxanthoides, *Munro, ex Henderson & Hume, Lahore to Yarkand*, 339.—Turkest.
Anthoxanthum, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 176 (err. typ. 156), 179.—Reg. Andin.
arundinacea, Beauv. Agrost. 160 = Calamagrostis varia.
aurea, *Munro, ex Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 176 (err. typ. 156), 179.—Ecuador.
australis, Benth. & Hook. f. Gen. iii. 1158 = plebeia.
avenacea, Spreng. Syst. i. 254 = Anisopogon avenaceus.
avenoides, *Y. Buch. Man. Grass. N. Zeal.* p. vi.—N. Zel.
avorica, Hochst. ex Seub. Fl. Azor. 19 = Agrostis pallida.
Billardieri, *Kunth, Rév. Gram.* i. 77.—Austral.; N. Zel.
Bolanderi, *Vasey, Descr. Cat. Grass. U. St.* 80.—Calif.
brasiliensis, Spreng. ex Steud. Nom. ed. II. i. 497 (= *Eutriana villosa*).—Bras.
breviaristata, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 179.—Peruv.
brevifolia, *Y. & C. Presl, Rel. Haenk.* i. 248.—Peruv.
breviglumis, *Benth. Fl. Austral.* vii. 584.—Austral.
Breweri, *Vasey, Cat. Gr. U. St.* 80.—Calif.
caespitosa, Hochst. ex Seub. Fl. Azor. 18 = Agrostis pallida.
canadensis, *Munro, ex Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 845.—Am. bor.
capillaris, Clar. fide Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 580 = Poa depauperata.
capitata, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 176 (err. typ. 156), 179.—Bolliv.
Cephalotes, *Wedd. l. c.* 178, 179.—Bolliv.
chalybaea, *Munro, ex Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 846.—Reg. arct.
Chamissonis, *Kunth, Enum. Pl.* i. 242.—Ins. Paschal.
chilensis, *E. Desv. in C. Gay, Fl. Chil.* vi. 322.—Chili.
chrysantha, *Y. & C. Presl, Rel. Haenk.* i. 247.—Peruv.
chrysostachya, *E. Desv. in C. Gay, Fl. Chil.* vi. 323.—Chili.
coarctata, *H. B. & K. Nov. Gen. et Sp.* i. 143.—Ecuador; N. Granat.
compacta, *Munro, ex Duthie, in* [*Athins. Gas. N. W. Ind.*] 238.—Ind. or.
confinis, *Kunth, Rév. Gram.* i. 76.—Am. bor.
crassiglumis, *Vasey, Cat. Gr. U. St.* 80.—Calif.
cryptolophe, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 176 (err. typ. 156), 179.—Bolliv.
curta, *Wedd. l. c.*—Bolliv.
curvula, *Wedd. l. c.* 178, 179.—Peruv.
Cusickii, *Vasey, in Coult. Bot. Gaz.* x. (1885) 224.—Am. bor. occ.
cylindrica, *Benth. Fl. Austral.* vii. 582.—Austral.
dense, *Benth. l. c.*—Austral.
densiflora, *Y. & C. Presl, Rel. Haenk.* i. 247.—Peruv.
deschampsioides, *Vasey, Cat. Gr. U. St.* 80.—Am. bor. occ.
desertícola, *Phil. Fl. Atac.* 55.—Chili.
Drummondiana, *Benth. Fl. Austral.* vii. 580.—Austral.
effusa, *H. B. & K. Nov. Gen. et Sp.* i. 145.—Ecuador.
elata, *Munro, ex Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 846.—Reg. arct.
elata, *Phil. in Linnaea*, xxxiii. (1864–65) 389.—Chili.
elegans, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 179.—Bolliv.
eminens, *Y. & C. Presl, Rel. Haenk.* i. 250.—Peruv.

DEYEUXIA:—
erianthа, *H. B. & K. Nov. Gen. et Sp.* i. 146.—Mexic.
erythrostachya, *E. Desv. in C. Gay, Fl. Chil.* vi. 324.—Chili.
evoluta, *Fourn. in Bull. Soc. Bot. Fr.* xxiv. (1877) 181.—Mexic.
festuciodes, *Wedd. l. c.* xxii. (1875) 178, 179.—Bolliv.
filifolia, *Wedd. l. c.*—Bolliv.
flavescens, Clar. ex Spreng. Syst. i. 254 = planifolia.
Forsteri, *Kunth, Rév. Gram.* i. 77.—Austral.; N. Zel.
frigida, *F. Muell. ex Benth. Fl. Austral.* vii. 588.—Austral.
Funckii, *Steud. ex Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 178.—Reg. Andin.
fuscata, *Y. & C. Presl, Rel. Haenk.* i. 248.—Peruv.
glacialis, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 178, 179.—Bolliv.
gracilis, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 554.—Mexic.
gracilis, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 179.—Bolliv.
groenlandica, *Munro, ex Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 845.—Reg. arct.
Gunniana, *Benth. Fl. Austral.* vii. 584.—Tasman.
Haenkeana, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 179, 180.—Peruv.
Hallerana, *Vasey, Cat. Gr. U. St.* 80.—Am. bor. occ.
Hartmaniana, Munro, ex Hook. f. in Trans. Linn. Soc. xxiii. (1862) 845 = Calamagrostis ambigua.
heteruphylla, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Bolliv.
Hostii, *Desv. ex Andrs. Rys Botan.* ii. (1830) 26.—Kossia.
Howellii, *Vasey, Agric. Grasses U. St.* 70. t. 61.—Am. bor. occ.
imberbis, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Bolliv.
Interrupta, *Y. & C. Presl, Rel. Haenk.* i. 249.—Peruv.
Jamesoni, *Munro, ex Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 176 (err. typ. 156).—Ecuador.
japonica, Spreng. Syst. i. 254 = Arthraxon ciliaris.
juncifolia, *H. B. & K. Nov. Gen. et Sp.* i. 144.—Mexic.
Lagurus, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 176 (err. typ. 156), 180.—Bolliv.
Langsdorffii, *Kunth, Rév. Gram.* i. 77.—Am. bor.
lappacea, *Kunth, l. c.* 76.—Reg. bor. temp.
leiophylla, *Munro, ex Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Bolliv.
leptopoda, *Wedd. l. c.*—Reg. Andin.
Lieblmannіana, *Fourn. in Bull. Soc. Bot. Fr.* xxiv. (1877) 181.—Mexic.
ligulata, *H. B. & K. Nov. Gen. et Sp.* i. 145.—Ecuador.
longearistata, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 176 (err. typ. 156), 180.—Bolliv.
Macounіana, *Vasey, in Coult. Bot. Gaz.* x. (1885) 297.—Am. bor.
Mandoniana, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 179, 180.—Bolliv.
Mann, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 238.—Afr. trop.
megapotamica, Spreng. Syst. iv. Cur. Post. 30 = Anisopogon saccharoides.
mildfordi, Beauv. ex Nym. Consp. 799 = Calamagrostis varia.
minor, *Benth. Fl. Austral.* vii. 582.—Austral.
montana, Beauv. Agrost. 44, 160; Poir. Tabl. Encyc. iii. 538 = Calamagrostis varia.
montana, *Fl. Austral.* vii. 581.—Austral.
montana, Boiss. Fl. Orient. v. 525 = acutiflora.
montevidensis, Spreng. Syst. iv. Cur. Post. 30 = Anisopogon saccharoides.
muhlfordi, Beauv. ex Nym. Consp. 799 = Calamagrostis varia.
minor, *Benth. Fl. Austral.* vii. 582.—Austral.
muticа, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Bolliv.
neglecta, *Kunth, Rév. Gram.* i. 76.—Reg. bor. temp.
nematophylla, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 179, 180.—Bolliv.
nivalis, *Benth. Fl. Austral.* vii. 588.—Austral.
nivalis, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 178 (err. typ. 156), 180.—Bolliv.
nubigenа, *Y. & C. Presl, Rel. Haenk.* i. 250.—Sicua Nutka.
Nuttalliana, *Vasey, Cat. Gr. U. St.* 81.—Am. bor.
obtusata, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Bolliv.
Orbigyana, *Wedd. l. c.* 178, 180.—Bolliv.

DEYEUXIA:—
Orizabae, *Rupr. in Bull. Acad. Brux.* ix. ii. (1842) 238.—Mexic.
ovata, *Y. & C. Presl, Rel. Haenk.* i. 246.—Peruv.
pallens, *Y. & C. Presl, l. c.* 249.—Peruv.; Ecuador.
phalaroides, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Bolliv.
Pickeringii, *Vasey, Cat. Gr. U. St.* 81.—Am. bor.
picta, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Bolliv.
pilosa, *Y. Buch. Man. Grass. N. Zeal.* p. vi.—N. Zel.
planifolia, *H. B. & K. Nov. Gen. et Sp.* i. 145.—Peruv.
plebeia, *Benth. Fl. Austral.* vii. 580.—Austral.
poaeformis, H. B. & K. Nov. Gen. et Sp. i. 146 = Poa subuniflora.
polystachya, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Peruv.
Porteri, *Vasey, Cat. Gr. U. St.* 81.—Am. bor.
Preissi, *Nees, in Linn. Pl. Preiss.* ii. 97.—Austral.
Preslei, Kunth, Enum. Pl. i. 243 = Calamagrostis scabra.
purpurascens, *Kunth, Rév. Gram.* i. 77.—Am. bor. occ.
purpurea, Kunth, l. c. = Calamagrostis purpurea.
quadriseta, *Benth. Fl. Austral.* vii. 581.—Austral.; N. Zel.
rariflora, Fuss, Fl. Transs. 714 = Halleriana?
recta, Cham. & Schlecht. in Linnaea, vi. (1831) 87 = Schiedeana.
recta, *H. B. & K. Nov. Gen. et Sp.* i. 144.—Ecuador; Peruv.
retrofracta, Kunth, Rév. Gram. i. 77 = Forsteri.
rigida, *H. B. & K. Nov. Gen. et Sp.* i. 144.—Ecuador; Peruv.
robusta, *Phil. Fl. Atac.* 54.—Chili.
rubescens, *Vasey, Cat. Gr. U. St.* 81.—Am. bor. occ.
scabra, *Kunth, Rév. Gram.* i. 77; *Benth. Fl. Austral.* vii. 588.—Austral.
scabrescens, *Munro, ex Duthie, in* [*Athins. Gas. N. W. Ind.*] 238.—Ind. or.
scabriuscula, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Peruv.
Schaffneri, *Fourn. l. c.* xxiv. (1877) 181.—Mexic.
Schiedeana, *Rupr. in Bull. Acad. Brux.* ix. ii. (1842) 238.—Mexic.
sciurea, (vide sub Agrostide, p. 65) recte = Dichelachne sciurea.
sedenensis, Beauv. Agrost. 44 = Avena sempervirens.
sesquiflora, *Benth. & Hook. f. Gen.* iii. 1158.—Chili.
sesquiloba, Benth. in Journ. Linn. Soc. xix. (1881) 93, sphalm. =prasc.
setifiora, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 176 (err. typ. 156), 180.—Bolliv.
setifolia, *Hook. f. Fl. N. Zel.* 289. t. 65.—N. Zel.
spathacea, *Clar. ex Steud. Nom.* ed. II. i. 497 = effusa.
spicata, Spreng. Syst. i. 254 = Andropogon condensatus.
spiciger, *Y. & C. Presl, Rel. Haenk.* i. 247.—Peruv.
splendens, Brongn. in Duperr. Voy. Coq. Bot. 23 = Calamagrostis splendens.
Sprucеana, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 178, 180.—Ecuador.
stricta, *H. B. & K. Nov. Gen. et Sp.* i. 146.—N. Granat.
strigosa, *Kunth, Rév. Gram.* i. 77.—Am. bor. occ.
subalpina, Schur, Enum. Pl. Transs. 740 = sylvatica.
subinflata, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 178, 180.—Bolliv.
subspicata, Desv. Opusc. 60, in obs. = Trisetum subspicatum.
Sukstorfii, *Scribn. ex Vasey, Descr. Cat. Grass. U. St.* 81.—Am. bor. occ.
sulcata, Wedd. in Bull. Soc. Bot. Fr. xxii. (1875) 178, 180 = Mandoniana.
sylvatica, *Kunth, Rév. Gram.* i. 77.—Reg. bor. temp.
talosoensis, *H. B. & K. Nov. Gen. et Sp.* i. 143.—Mexic.
*tarifana

DEYEUXIA:—
vicinarum, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 177, 180.—Reg. Andin.
Vidalî, *Phil. in Anal. Univ. Chil.* (1878) 668.—Chili.
violacea, *Wedd. in Bull. Soc. Bot. Fr.* xxii. (1875) 179, 180.—Bolliv.
viridiflavescens, *Kunth, Rh. Gram.* t. 77.—Reg. Argent.
viridis, *Fourn. in Bull. Soc. Bot. Fr.* xxiv. (1877) 181. —Mexic.
viridis, *Phil. in Linnæa*, xxxiii. (1864–65) 388.—Chili.
viridans, *Phil. l. c.* xxix. (1857–58) 90.—Chili.
vulcanica, *Phil. l. c.*—Chili.
Youngii, *Y. Buch. Man. Grass. N. Zeal.* p. vi.—N. Zel.

DIACAECARPIUM, Endl. Gen. 745 (1839) = DIACICARPIUM, Blume = **Marlea**, Roxb. (Cornac.).

DIACANTHA, Lag. Amen. Nat. i. 40 (1811) = **Chuquiraga**, Juss. (Compos.).
ambigua, Cass. in Dict. Sc. Nat. xlii. 129 = C. spinosa.
spinosa, Lag. Amen. Nat. i. 40 = C. spinosa.

DIACANTHA, Less. in Linnæa, v. (1830) 248 = **Barnadesia**, Mutis (Compos.).
arborea, Less. l. c. = B. arborea.

DIACECARPIUM, Hassk. Cat. Hort. Bogor. Alt. 169 (1844) = DIACICARPIUM, Blume = **Marlea**, Roxb. (Cornac.).

DIACHROA, Nutt. ex Steud. Nom. ed. II. i. 497 (1840) = **Glycoeria**, R. Br. (Gramin.).
maritima, Nutt. ex Steud. l. c. = G. maritima.

DIACHYRIUM, Griseb. in Goett. Abh. xix. (1874) 257 = **Sporobolus**, R. Br. (Gramin.).
arundinaceum, Griseb. l. c.—Reg. Argent.

DIACICARPIUM, Blume, Bijdr. 657 (1825) = **Marlea**, Roxb. (Cornac.).
rotundifolia, Hassk. in Bonplandia, vii. (1859) 172 = M. begoniæfolia.
tomentosum, Blume, Bijdr. 657 = M. begoniæfolia.

DIACIDIA, Griseb. in Mart. Fl. Bras. xii. 1. 119 (1858). *MALPIGHIACEÆ*, Benth. & Hook. f. i. 252.
galphimioides, *Griseb. l. c.* 120.—Bras.
vestita, *Benth. & Hook. f. Gen.* i. 253, in nota sub Coleostachyde.—Guiana.

DIACLES, Saliab. Gen. Pl. Fragm. 180 (1866) = **Haemanthus**, Tourn. (Amaryllid.).
ciliaris, Saliab. l. c. = H. albiflos.
pubescens, Saliab. l. c. = H. albiflos.

DIACRIUM, Benth. in Journ. Linn. Soc. xviii. (1881) 312. *ORCHIDEÆ*, Benth. & Hook. f. iii. 526.
bicornutum, *Benth. l. c.*—Guiana.
bidentatum, *Hemsl. Biol. Centr. Am. Bot.* iii. 221.—Mexic.
bigibberosum, *Hemsl. l. c.* 222.—Guatemala.
bilamellatum, *Hemsl. l. c.*—Panama.

DIADENARIA, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 254; et in Abh. Akad. Berl. (1860) 107 = **Poelliantthus**, Neck. (Euphorb.).
articulata, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 108 = P. articulatus.
involucrata, Klotzsch & Garcke, l. c. 107 = P. involucratus.
Pavonii, Klotzsch & Garcke, l. c. 108 = P. Pavonis.
pulcherpa, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 254.—Hab. ?

DIADENIUM, Poepp. & Endl. Nov. Gen. et Sp. i. 41. t. 71 (1835). *ORCHIDEÆ*, Benth. & Hook. f. iii. 567.
CHARNANTHE, Lindl. Bot. Reg. (1838) Misc. 88.
Barkeri, *Benth. & Hook. f. Gen.* iii. 558.—Bras.
micranthum, *Poepp. & Endl. Nov. Gen. et Sp.* i. 41. t. 71.—Peruv.

DIADESMA, Raf. *New Fl. Am.* i. 41 (1836) = **Sida**, Linn. (Malvac.).
carduacea, Raf. l. c. = S. carduacea.
rhombifolia, Raf. l. c. = S. rhombifolia.
stellata, Raf. l. c. = Sphæranthus angustifolia.
virgata, Raf. l. c. = S. virgata.

DIALANTHERA, Raf. Sylva Tellur. 127 (1838) = **Cassia**, Tourn. (Legumin.).
glandulosa, Raf. l. c. = C. glandulosa.

DIALESTA, H. B. & K. Nov. Gen. et Sp. iv. 45. t. 320 (1820) = **Oliganthus**, Cass. (Compos.).
discolor, H. B. & K. l. c. = O. discolor.
stenoloides, Mart. ex DC. Prod. v. 79 = Lychnophora stenoloides.

DIALION, Raf. Sylva Tellur. 89 (1838) = **Heliotropium**, Tourn. (Borragin.).
undulatum, Raf. l. c. = H. undulatum.

DIALISSA, Lindl. in Ann. & Mag. Nat. Hist. Ser. I. xv. (1845) 107 = **Stelis**, Sw. (Orchid.).
pulchella, Lindl. l. c. = S. Dialissa.

DIALIUM, Linn. Mant. i. 3 (1767). *LEGUMINOSÆ*, Benth. & Hook. f. i. 574.
ARGUMA, Aubl. Pl. Gui. i. 16. t. 5 (1775).
CODARIUM, Soland. ex Vahl, Enum. i. 302 (1805).
angolense, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 383.—Afr. trop.
anomalum, *Webb, in Hook. Kew Journ.* ii. (1850) 347. —Ins. Gorgoas.
coromandelianum, Houtt. Handleid. ii. 39 = indum ?
dicolor, Hook. f. in Hook. Niger Fl. 329 = guineense.
divaricatum, *Vahl, Enum.* i. 303.—Guiana; Bras.
guianense, Steud. Nom. ed. II. i. 497 = seq.
guineense, *Willd. in Roem. Arch.* i. 1. 31. t. 6.—Afr. trop.
indum, *Linn. Mant.* i. 24.—Java.
javanicum, Burm. f. Fl. Ind. 12 = indum.
laurinum, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 269.—Malacca.
Maingayi, *Baker, l. c.*—Singapore.
nitidum, Guill. & Perr. Fl. Senegamb. Tent. i. 289 = guineense.
ovoideum, *Thw. Enum. Pl. Zeyl.* 97.—Zeylan.
patens, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 270.—Malacca.
platyspalum, *Baker, l. c.*—Malacca.

DIALLA, Lindl. Veg. Kingd. 390 (1847) = **Diodia**, Griseb. (Malpigh.).

DIALLOBUS, Raf. Sylva Tellur. 128 (1838) = **Cassia**, Tourn. (Legum.).
bicapsularis, Raf. l. c. = C. bicapsularis.
falcatus, Raf. l. c. = C. Tora.
subuniflorus, Raf. l. c. = C. uniflora ?
Tora, Raf. l. c. = C. Tora.
uniflorus, Raf. l. c. = C. uniflora.

DIALLOSPERMA, Raf. Sylva Tellur. 69 (1838) = **Aspalathus**, Linn. (Legumin.).
spinosa, Raf. l. c. = A. spinosa.

DIALLOSTEIRA, Raf. Neogenyt. 2 (1825) = Collinsonia, Linn. (Labiat.).
punctata, Raf. l. c. = C. canadensis.

DIALYCARPA, Mast. in Journ. Linn. Soc. xiv. (1875) 595. *MALVACEÆ*.
Beccarii, *Mast. l. c.* 596.—Borneo.

DIALYPETALUM, Benth. in Benth. & Hook. f. Gen. ii. 555 (1876). *CAMPANULACEÆ*, Benth. & Hook. f. i. c.
floribundum, *Benth. l. c.* 554.—Madag.

DIAMARIPS, Raf. Alsog. Am. 15 (1838) = **Salix**, Tourn.
Crevecœur, Raf. l. c. = S. Crowenna.
rubra, Raf. l. c. = S. rubra.

DIAMONON, Raf. Fl. Tellur. ii. 76 (1836) = **Solanum**, Tourn.
curiacum, Raf. l. c. = S. havanense.

DIAMORPHA, Nutt. Gen. Am. i. 293 (1818). *CRASSULACEÆ*, Benth. & Hook. f. i. 661.
pusilla, *Nutt. l. c.*—Am. bor.

DIANA, Comm. ex Lam. Encyc. ii. 276 (1786) = seq.

DIANELLA, Lam. Encyc. ii. 276 (1786). *LILIACEÆ*, Benth. & Hook. f. iii. 793.
DIANA, Comm. ex Lam. Encyc. ii. 276 (1786).
EXCREMIS, Willd. ex Schult. f. Syst. vii. 354, in adnot. (1829).
RHUACOPHILA, Blume, Enum. Pl. Jav. 18 (1830).
angustifolia, Schult. f. Syst. vii. 353 = caerulea.
Archeri, Hook. f. Fl. Tasm. ii. 58 = tasmanica.
atrata, Hort. Petrop. ex Regel, Gartenfl. (1857) 87 = inconspicua.
austro-caledonica, *Seem. Fl. Vit.* 312.—N. Caled.
banosea, *Mfq. Fl. Ind. Bat. Suppl.* 410.—Ins. Banca.
caerulea, *Sims, Bot. Mag.* t. 805.—Austral.
celebica, *Kunth, Enum. Pl.* v. 55.—Malaya.
cærulea, Willd. Enum. Hort. Berol. 275 = caerulea.
congesta, R. Br. Prod. 280 = caerulea.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 372.—Cuba.
densa, Lindb. in Act. Soc. Sc. Fenn. x. (1871) 132. t. 6 = tasmanica.
divaricata, R. Br. Prod. 280 = revoluta.
dubia, *H. B. & K. Nov. Gen. et Sp.* i. 270. t. 675.—Venezuela.
elegans, Kunth & Bouché, in Ind. Sem. Hort. Berol. (1847) 11 = caerulea ?
elegans, F. Muell. Fragm. vi. 122 = laevis.
ensifolia, (DC. in) Red. Lil. t. 1 = nemorosa.
graminifolia, Kunth & Bouché, Ind. Sem. Hort. Berol. (1847) 11 = caerulea ?
hemicryptæ, Lam. Encyc. ii. 276 = Cordyline hemicryptæ.
Hookeri, Baker, in Journ. Linn. Soc. xiv. (1875) 575 = tasmanica.
humilis, Lodd. ex Steud. Nom. ed. II. i. 498 = nemorosa ?
inconspicua, *Regel, Gartenfl.* (1857) 87.—Hab. ?
intermedia, *Smith. Prod. Fl. Ins. Norfolc.* 28.—N. Zel.; Ins. Norfolk.
javanica, Kunth, Enum. Pl. v. 59 = nemorosa.
laevis, *R. Br. Prod.* 280.—Austral.
laevis, Hook. f. Fl. Tasm. ii. 57 = tasmanica.
longifolia, R. Br. Prod. 280 = laevis.
longifolia, (Lindl. in) Bot. Reg. t. 734 = revoluta.
mauritiana, Blume, Enum. Pl. Jav. i. 12 = nemorosa.
montana, Blume, l. c. = seq.
nemorosa, *Lam. Encyc.* ii. 276. *Yacq. Hort. Schoenb.* t. 94—As. et Austral. trop.
nigra, Colenso, in Trans. N. Z. Inst. xvi. (1884) 389 = intermedia.
obscura, Kunth, Enum. Pl. v. 56 = nemorosa.
odorata, Blume, Enum. Fl. Jav. i. 13 = nemorosa.
paniculata, Hort. Berol. ex Kunth, Ind. Sem. Hort. Berol. (1847) 11 = caerulea.
rara, R. Br. Prod. 280 = laevis.
revoluta, *R. Br. l. c.*—Austral.
sandwicensis, Hook. & Arn. Bot. Beech. Voy. 97 = nemorosa.
strumosa, [Lindl. in] Bot. Reg. t. 751 = laevis.
tasmanica, *Hook. f. Fl. Tasm.* ii. 57. t. 133.—Austral.
triandra, *Afz. Remed. Guin.* 6 (Quid ?).—Afr. trop.

DIANIA, Noronha, ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 142 = Dicoryphe, Thou. (Hamamelid.).
amplifolia, Noronha, ex Tul. l. c. 147 = Dicoryphe Noronhae.

DIANTHELLA, Classon, ex Pomel, Mat. Fl. Atl. 9 (1860) = **Tunica**, Hall. (Caryophyll.).
compressa, Pomel, l. c. = T. compressa.

DIANTHERA, Gronov. Fl. Virgin. 5 (1742); Linn. Gen. ed. II. 13 (1743). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1113.
CHILOGLOSSA, Oerst. in Kjœbh. Vidensk. Meddel. (1854) 180.
DIANANSEA, Raf. Fl. Tellur. ii. 88 (1836).
DIPTANTHERA, Schrank, ex Steud. Nom. ed. II. i. 518 (1840).
LEPTOSTACHYA, Ness, in Wall. Pl. As. Rar. iii. 105 (1832).
LUSTRINIA, Raf. Fl. Tellur. iv. 63 (1836).
MICROSPERMA, Raf. l. c. 64 (1836).
ORTHOTACTUS, Ness, in Mart. Fl. Bras. ix. 181. t. 21 (1847).
PLAGIACANTHUS, Ness, in DC. Prod. xi. 385 (1847).
PORPHYROCOMA, Hook. Bot. Mag. t. 4176 (1845).
RHYTIGLOSSA, Ness, in Lindl. Introd. Nat. Syst. ed. II. 285, 444 (1836).
STETHOMA, Raf. Fl. Tellur. iv. 61 (1836).
acuminata, Benth. & Hook. f. Gen. ii. 1114 = comata.

DIANTHERA:—
acuminata, Ruiz & Pav. Fl. Per. i. 10. t. 16 = Dicliptera acuminata.
alata, Benk. in DC. Prod. Index, iii. 126 = Andrographis alata.
americana, Blanco, Fl. Filip. ed. I. 16 (= *Rostellularia Blancoi*).—Hab ?
americana, Forsk. Fl. Aegypt. Arab. 9 = Barleria flava.
americana, Fraser, ex Nees, in DC. Prod. xi. 340 = ovata.
americana, *Linn. Sp. Pl.* 27.—Am. bor.
Anagallis, *Hemsl. Biol Centr. Am. Bot.* ii. 517.—Mexic.
andrœcamifolia, *Griseb. Fl. Brit. W. Ind.* 455.—Ins S. Doming.
angularis, Salisb. Prod. 102 = Dicliptera sexangularis.
angustifolia, *Benth. & Hook. f. Gen.* ii. 1114.—Bras.
Anselliana, *Benth. & Hook. f. l. c.*—Afr. trop.
appendiculata, Ruiz & Pav. Fl. Per. i. 12. t. 11 = Beloperone appendiculata.
arvensis, Vell. Fl. Flum. 12; i. t. 30 = Justicia arvensis.
bicalyculata, Retz. in Vet. Acad. Handl. (1776) 297 = Peristrophe bicalyculata.
bifida, Moench, Meth. 521 = Peristrophe bicalyculata.
brasiliensis, Vell. Fl. Flum. 11; i. t. 26 = Justicia Arrabidæa.
brevifiora, *Hemsl. Biol. Centr. Am. Bot.* ii. 517.—Mexic.
calycina, *Benth. & Hook. f. Gen.* ii. 1111, 1113, in obs. —Guiana.
campestris, *Benth. & Hook f. l. c.* 1114.—Bras.
Candelariæ, *Hemsl. Biol. Centr. Am. Bot.* ii. 517 — Costa Rica.
candicans, *Benth. & Hook. f. Gen.* ii. 1113.—Mexic.
carolinians, Fraser, ex Nees, in DC. Prod. xi. 340 = ovata.
cayennensis, *Griseb. Cat. Pl. Cub.* 197, in textu.—Guiana.
ciliata, *Benth. & Hook. f. Gen.* ii. 1114.—Costa Rica.
ciliata, Blanco Fl. Filip. ed. II. 19 = Justicia simplex.
ciliata, Medic. Beobacht. (1783) 9 = Schwabea ciliaris.
ciliata, *Ruiz & Pav. Fl. Per.* i. 12 t. 14.—Peruv.
clavata, Forsk. ex Nees, in DC. Prod. xi. 475 = Dicliptera ciliaris.
clavata, Forst. f. Prod. 4 = Dicliptera clavata.
coccinea, Salisb. Prod. 102 = Justicia coccinea.
corrulea, Forst. f. Prod. 4 = Dicliptera pubescens.
collina, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 548.—Reg. Himal.
comata, *Linn. Syst.* ed. X. i. 850.—Mexic.
comata, Vell. Fl. Flum. 12; i. t. 27 = Justicia fumigenalis.
crassifolia, *Chapm. Fl. S. U. St.* 304.—Am. bor.
dasycladus, *Benth. & Hook. f. Gen.* ii. 1113.—Bras.
debilis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 542. —Reg. Himal.
debilis, Forsk. Fl. Aegypt. Arab. 9 = Justicia debilis.
decumbens, Sw. ex Nees, in DC. Prod. xi. 332 = comata.
dichotoma, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 548.—Zeylan.
diotera, *Benth. & Hook. f. Gen.* ii. 1113.—Bras.
elongata, Salisb. Prod. 103 = Thymacanthus bracteolatus.
emiformis, Walt. Fl. Carol. 68 = americana.
enstachiana, *Y. F. Gmel. Syst.* 86.—Ind. occ.
enstachiana, Hook. ex Nees, in DC. Prod. xi. 466 = Blechum trinitense.
flava, Vahl, Symb. Bot. i. 5 = Adhatoda flava.
gemistaefolia, Salisb. Prod. 103 = Justicia orchidioides.
glabra, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 518.—Costa Rica.
glandulosa, *Griseb. Cat. Pl. Cub.* 197.—Cuba.
Griffithii, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 542.—Reg. Himal.
harpochlioides, Griseb. Cat. Pl. Cub. 196 = Jacobinia harpochlioides.
heterophylla, *Benth. & Hook. f. Gen.* ii. 1114.—Bras.
hirsuta, *Y. F. Gmel. Syst.* 85.—Ind. occ.
hirsuta, Ruiz & Pav. Fl. Per. i. 11. t. 13 = Justicia inficiens.
Hookeriana, *Benth. & Hook. f. Gen.* ii. 1113.—Peruv.
humifusa, J. F. Gmel. Syst. 86 = comata.
humilis. Engelm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1847) 280 = ovata.
hyssopifolia, Burm. f. Fl. Ind. 11. t. 5. f. 1 = Peristrophe undulata.

DIANTHERA:—
hyssopifolia, Medic. Beobacht. (1783) 7 = Justicia hyssopifolia.
Inæqualis, *Benth. & Hook. f. Gen.* ii. 1113.—Guatemala.
indica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 548. —Reg. Himal.
japonica, Thunb. Fl. Jap. 21. t. 1 = Dicliptera crinita.
justicioides, Sw. ex Nees, in DC. Prod. xi. 415 = Adhatoda tetranema.
lasia, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877 78) 83. —Bras.
lanciflola, Tafalla, ex Nees, in DC. Prod. xi 357 = Amphiscopia lanciflola.
latifolia, *Benth. & Hook. f. Gen.* ii. 1113.—Mexic.
latifolia, Salisb. Prod. 103 = Adhatoda Vasica.
leptostachya, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 542 = Wallichii.
Lindeniana, *Hemsl. Biol. Centr. Am. Bot.* ii. 518.—Mexic.
lincolata, *Tafalla, ex Nees, in DC. Prod.* xi. 345.—Peruv.
linifolia, *Benth. & Hook. f. Gen.* ii. 1114.—Am. bor.
longifiora, *Benth. & Hook. f. l. c.* 1113.—Bras.
lucida, *Benth. & Hook. f. l. c.*—Ind. occ.
malabarica, Gouan, ex Nees, in DC. Prod. xi. 474 = Dicliptera resupinata.
malabarica, Linn. f. Suppl. 85 = Peristrophe bicalyculata.
martinicensis, J. F. Gmel. Syst. 86 = Dicliptera martinicensis.
mexicana, *Benth. & Hook. f. Gen.* ii. 1114.—Mexic.
mucronata, Math. ex Nees, in DC. Prod. xi. 480 = Dicliptera Hookeriana.
multiflora, Ruiz & Pav. Fl. Per. i. 11. t. 16. f. a = Dicliptera peruviana.
multiflora, Ruiz & Pav. l. c. 10. t. 14. f. b = Dicliptera umbellata.
nitida, J. F. Gmel. Syst. 85 = Thymacanthus nitidus.
nodosa, *Benth. & Hook. f. Gen.* ii. 1113.—Ind. or.
obcordata, Vahl, ex Nees, in DC. Prod. xi. 399 = Thymacanthus tenuiflorus.
oblonga, *Benth. & Hook. f. Gen.* ii. 1114.—N. Granat.
obtusifolia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1868) 525.—Ind. occ.
odora, Forsk. Fl. Aegypt. Arab. 8 = Adhatoda odora.
ovata, *Walt. Fl. Carol.* 68.—Am. bor.
paniculata, Forsk. Fl. Aegypt. Arab. 7 = Peristrophe bicalyculata.
paniculata, Lour. Fl. Cochinch. i. 26 = Rhinacanthus communis.
parviflora, *A. Gray, Syn. Fl. N. Am.* ii. i. 330.—Am. bor.
parviflora, Nordz. ex Nees, in DC. Prod. xi. 471 = Ruspia parviflora.
parvifolia, *Benth. & Hook f. Gen.* ii. 1114.—Texas.
pectoralis, *Y. F. Gmel. Syst.* 86.—Ind. occ.; Mexic.
perploides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1868) 525.—Cuba.
pilosa, Ruiz, ex Nees, in DC. Prod. xi. 405 = Adhatoda pilosa.
punctata, *Vahl, Symb. Bot.* i. 4.—Arabia.
racemosa, *Benth. & Hook f. Gen.* ii. 1114.—Peruv.
repens, Ruiz & Pav. Fl. Per. i. 10. t. 15. f. b = Dicliptera tonsa.
reptans, *Y. F. Gmel. Syst.* 86; *Griseb. Fl. Brit. W. Ind.* 455.—Jamaica.
rigida, *Benth. & Hook. f. Gen.* ii. 1115, in obs.—Bras.
Rugeliana, *Griseb. Cat. Pl. Cub.* 196.—Cuba.
rugosa, Perr. ex Nees, in DC. Prod. xi. 497 = Peristrophe bicalyculata.
Sagraeana, *Griseb. Cat. Pl. Cub.* 196.—Cuba.
sanctitis, Salisb. Prod. 102 = Dicliptera viridis.
secunda, *Griseb. in Goett. Abh.* vii. (1857) 246.—Cuba.
secundiflora, Ruiz & Pav. Fl. Per. i. 11. t. 15. f. a = Justicia secundiflora.
serrata, Vell. ex Nees, in DC. Prod. xi. 440 = Justicia tenuispica.
sessilis, *Y. F. Gmel. Syst.* 85; *Griseb. Fl. Brit. W. Ind.* 456.—Ind. occ.
speciosa, *Benth. & Hook. f. Gen.* ii. 1113.—N. Granat.
spicata, Vell. Fl. Flum. 12; i. t. 28 = Schaueria Schottii.
subserrata, Blanco, Fl. Filip. ed. I. 16 = Justicia Gendarussa.

DIANTHERA:—
sulcata, Griseb. in Goett. Abh. xix. (1874) 224 (Anisacanthi sp.?).—Reg. Argent.
sulcata, Vahl, Symb. Bot. i. 4 = Adhatoda sulcata.
terminalis, *Fawc. in Forbes, Nat. Wand. East. Archip.* 518.—Ins. Timor.
tomentosa, Roxb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 395 = Nelsonia campestris.
trinalca, Forsk. Fl. Aegypt. Arab. 7 = Anisotes trinalcus.
urticifolia, Forsk. l. c. 9 = Dicliptera micranthes.
violacea, *Vahl, Symb. Bot.* i. 6.—Arab.
virgata, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit Ind.* iv. 542.—Reg. Himal.
viridiflora, Benth. & Hook. f. Gen. ii. 1114 = Gutenia laete-virens.
Wallichii, *Benth. & Hook. f. l. c.*—Ins. Philipp.

DIANTHERA, Klotzsch, in Peters, Reise Mossamb. Bot. 160. t. 87 (1861) = Clinonia, Linn. (Capparid.).
abyssinica, Schweinf. Fl. Aethiop. i. 70 = C. didymana.
Burchelliana, Klotzsch, in Peters, Reise Mossamb. Bot. 161 = C. diandra.
grandiflora, Klotzsch, l. c. = C. paradoxa.
lutea, Klotzsch, l. c. 162 = C. lutea.
Petersiana, Oliver, Fl. Trop. Afr. i. 79 = C. diandra.
Peterzii, Klotzsch, in Peters, Reise Mossamb. Bot. 160 = C. diandra.
semitetrandra, Klotzsch, l. c. 162 = C. semitetrandra.

DIANTHOSERIS, Sch. Bip. in Flora, xxv. (1842) 439. *COMPOSITAE*, Benth. & Hook. f. ii. 527.
Schimperi, *Sch. Bip. l. c.* 440.—Abyss.
Rueppellii, *Sch. Bip. l. c.*—Afr. trop.

DIANTHUS, Linn. Syst. ed. I (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 144.
CARYOPHYLLUS, Moench, Meth. 68 (1794).
CYLICHNANTHUS, Dulac, Fl. Hautes-Pyr. 260 (1867).
DIOSANTHUS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 87.
DYANTHUS, P. Br. Hist. Jamaic. 228 (1756), err. typ.
PLEONANTHUS, Ehrh. Beitr. iv. 148 (1789).
PLUMARIA, Opiz, Seznam, 75 (1852).
abyssinicus, R. Br. in Salt, Voy. Abyss. App. p. lxiv = leptolomus.
acicularis, *Fisch. ex Ledeb. Fl. Ross.* i. 284.—An. cent.
acicularis, Schur, Enum. Pl. Transs. 98 = petraeus.
acinifolius, Schur, l. c. 97 = D. Caryophyllus.
acicophyllus, Sieber, ex DC. Prod. i. 368 = juniperinus.
actinopetalus, *Fenzl, Pugill. Pl. Nov. Syr.* 11.—Graecia; An. Min.
acuminatus, Tausch, in Syll. Ratisb. ii. (1828) 242 = monspessulanus.
acuminatus, *F. W. Williams, in Journ. Bot.* xxiii (1885) 347.—Syria.
adulterinus x, Sweet, Hort. Brit. ed. II. 50.
aggericolus, Jord. in Billot, Annot. 48 = sylvestris.
aggregatus, Poir. Encyc Suppl. iv. 124 = barbatus
albens, Eckl. & Zeyh. Enum. 32 = pectinatus.
albens, Jacq. ex Sond. in Harv. & Sond. Fl. Cap. i. 129 = caespitosus.
albens, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 90.—Afr. austr.
albens, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 369 = Zeyheri.
albus, Schkuhr, ex Steud. Nom. ed. II. i. 498 = deltoides.
algetanus, *Graells, ex F. N. Williams, in Journ. Bot.* xxiii. (1885) 347.—Hispan.
alpestris, Balb. in Mém. Acad. Taurin, Phys.-Math. i. (1803-08) 18 = chinensis.
alpestris, Steinh. in Sturm, Deutschl. Fl. Heft 28 = monspessulanus.
alpinus, *Linn. Sp. Pl.* 412.—Europ.; Asia bor.; Am. bor. occ.
alpinus, Sibth. & Sm. Fl. Graec. Prod. i. 298 = haematocalyx.
alpinus, Sturm, ex Steud. Nom. ed. II. i. 498 = glacialis.
alpinus, Vill. Fl. Delph. 41; Hist. Pl. Dauph. iii. 600 = neglectus.
altaicus, Willd. ex Ledeb. Fl. Ross. i. 278 = chinensis.
amanicus, *Pamf. ex F. N. Williams, in Journ. Bot.* xxiii. (1885) 348.—Europ.
amblyonyx, Balbis, Prod. 208 = monspessulanus.
amoenus, *Pomel, Nouv. Mat. Fl. Atl.* 332.—Algeria.

DIANTHUS :—

anatolicus, *Boiss. Diagn.* Ser. I. i. 22.—As. Min.; Tibet. occ.

Andersonii, *F. N. Williams, in Journ. Bot.* xxiii. (1885) 346.—Syria.

angulatus E. Mey, Zwei Pfl. Docum. 178 = caespitosus.

angulatus, *Royle, Illustr. Bot. Himal.* 79.—Reg. Himal. anticarius, *Boiss. & Reut. Pugill. Pl. Nov.* 19.—Hispan.

aragonensis, *Trimb. in Mem. Acad. Toulouse*, Sér. VI. v. (1867) 242.—Hispan.

arborescens, Hoffmgg. Vers. Pfl. 56 = rupicola.

arborescens, *Linn. Sp. Pl.* 418.—Archip. Graec.

arbuscula, *Lindl. Bot. Reg.* t. 1086.—China.

arenarius, All. Fl. Pedem. ii. 77 = D. Caryophyllus.

arenarius, Colm. Fl. Catal. 22 = Requienii.

arenarius, *Linn. Sp. Pl.* 412.—Europ.

arenarius, Pall. Reise, iii. 600 = squarrosus.

arenarius, Thore, Essai Chl. 171 = gallicus.

arenarius, Thuill. Fl. Par. 212 = pungens.

arenarius, Towns. ii. Hung. iii. 408 = plumarius.

aridus, Griseb. ex Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 196 = pubescens.

aristatus, *Boiss. in Tchihat. Asie Min. Bot.* i. 229.—As. Min.

Armeria, *Linn. Sp. Pl.* 410.—Europ. : Reg. Caucas.

Armeriastrum, Wulfen. in Oestr. Bot. Zeitschr. viii. (1858) 318 = D. Armeria.

armerioides, *Rafin. in Desv. Journ. Bot.* iv. (1814) 269.—Am. bor.

arragonensis, Timb. ex Nym. Consp. 105 = Requenii.

arrectus, Dum. Fl. Belg. 106 = D. Caryophyllus.

Arrosti, *Y. & C. Presl, Delic. Prag.* 80.—Sicilia.

asper, Willd. Enum. Hort. Berol. 486 = chinensis.

asperulus, *Boiss. & Huet, Diagn.* Ser. II. i. 51.—As. Min.

atlanticus, *Pomel, Nouv. Mat. Fl. Atl.* 332.—Algeria.

atomarius, *Boiss. Diagn.* Ser. I. viii. 71.—Persia.

atratus, Beauprê, ex Ser. in DC. Prod. i. 256 = polymorphus.

atropurpureus, Gromow, ex Trautv. in Act. Hort. Petrop. viii. (1883) 130 = atrorubens !

atrorubens, *All. Fl. Pedem.* ii. 75.—Europ. austr.

atrpurbens, Baumg. Enum. i. 388 = banaticus.

atrorubens, Bieb. Fl. Taur. Cauc. i. 324 = capitatus.

atrorubens, Friv. ex Nym. Consp. 104 = intermedius.

atrorubens, Gaud. Fl. Helv. iii. 145 = Carthusianorum.

attenuatus, *Sm. in Trans. Linn. Soc.* ii. (1794) 301.—Hispan. ; Maroce.

attenuatus, Xatard, ex Walp. Rep. i. 267 = brachyanthus.

autumnalis, Kit. in Linnæa, xxxii. (1863) 530 = polymorphus.

axillíflorus, Fenzl, Pugill. Pl. Nov. Syr. 10 = multipunctatus.

Balansæ, *Boiss. Fl. Orient.* i. 488.—As. Min.

Balbisii, Ser. in DC. Prod. i. 356 = liburicus.

banaticus, *Heuff. ex Griseb. & Schenk, in Wiegm. Archiv*, xviii. (1852) t. 301.—Banatus.

banaticus, Kit. in Linnæa, xxxii. (1863) 527 = trifasciculatus.

Barrati, Duval-Jouve, in Bull. Soc. Bot. Fr. ii. (1855) 350 = tripunctatus.

barbatus, *Linn. Sp. Pl.* 409.—Europ.

barbatus, Pall. ex Ser. in DC. Prod. i. 355 = D. Pseudarmeria.

bebius, Vis. ex Reichb. Ic. Fl. Germ. vi. 47 = strictus.

benearnensis, H. Loret, in Bull. Soc. Bot. Fr. v. (1858) 827 = attenuatus.

Bertoloni, *Woods, Tour. Fl.* 45.—Italia.

bicolor, Adam, in Web. & Mohr, Beitr. i. 55 = campestris.

bicolor, Bieb. Fl. Taur. Cauc. i. 328 = pallens.

bicolor, Hornem. Hort. Hafn. 407 = leptopetalus.

bifornis, Griseb. Spicil. Fl. Rumel. i. 189 = gracilis.

bifornis, Sibth. & Sm. Fl. Graec. Prod. i. 285 = cinnabarinus.

binatus, Bastl. ex Reichb. Fl. Germ. Excurs. 810 = bifloris !

binatus, Schur, Enum. Pl. Transs. 97 = D. Caryophyllus.

Bingonii, Tenore, Cat. Ort. Bielg. (1805) 18 ; Fl. Nap. i. 252 = rupicola.

biornatus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 11 = banaticus.

bithynicus, *Kotschy, ex Boiss. Fl. Orient.* i. 488.—Armenia.

blandus, Reichb. Fl. Germ. Excurs. 807 = plumarius.

bohemicus, [Mayer] ex Tausch, in Flora, xlii. (1830) 246 = pubescens.

DIANTHUS :—

Bolsseri, *Willk. Ic. et Desc. Pl. Hisp.* i. 22. t. 13.—Hispan.

brachiatus, Sieber, ex Steud. Nom. ed. II. i. 499 (Quid ?).—Hab. ?

brachyanthus, *Boiss. Voy. Bot. Espagne*, ii. 85.—Hispan.

brachyanthus, Gren. & Godr. Fl. Fr. i. 225 = virgineus

brachyanthus, Schur, Enum. Pl. Transs. 96 = microlepis.

brachycalyx, Hnet, ex Nym. Consp. 105 = sylvestris.

bracteatus, Willd. ex Ser. in DC. Prod. i. 360 = Liboschitzianus.

brevicaulis, *Kenal, Pugill. Pl. Nov. Syr.* 10.—Cilicia.

brevifolius, Friv. in Flora, xviii. (1835) 334 = pinifolius.

brevilimbis, *Boiss. ex Trautv. in Act. Hort. Petrop.* ii. (1873) 506.—Armenia.

Broteri, *Boiss. & Reut. Pugill. Pl. Nov.* 22.—Hispan.

Buergeri, *Miq. Ann. Mus. Bot. Lugd Bat.* ii. 77.—Japon.

Burchellii, Ser. in DC. Prod. i. 359 = albens.

burchtormensis, *Ledeb. ex Steud. Nom.* ed. II. i. 499, nomen.—Rossia.

cachemiricus, *Edgew. ex Hook. f. Fl. Brit. Ind.* i. 214.—Reg. Himal.

caesius, [*Sm.*] *Engl. Bot.* (1792) t. 62.—Europ.

caespitosifolius, Plan. Enu. Fl. Gall. 118 = Planellae.

caespitosus, Eckl. & Zeyh. Enum. 81. pro parte = crenatus.

caespitosus, Poir. in Lam. Encyc. iv. 525 = caesius.

caespitosus, *Thunb. Prod. Pl. Cap.* 81.—Afr. austr.

callitzonus, *Schott & Kotschy, in Bot. Zeit.* ix. (1851) 192.—Transsylv.

calocephalus, *Boiss. Diagn.* Ser. I. vi. 28.—As. Min.

Calvert, *Boiss. in Tchihat. Asie Min. Bot.* i. 238.—As. Min.

campestris, *Bieb. Fl. Taur. Cauc.* i. 326.—Reg. Caucas.

campestris, Willd ex Ledeb. Fl. Ross. i. 277 = chinensis.

canescens, C. Koch, in Linnæa, xv. (1841) 710 = orientalis.

capillatus, Meinsh. Bestr. Pfl. Geogr. Sued-Ural-Geb. 6 = capitatus ?

capitatus, *DC. Cat Hort Monsp.* 108.—Rossia austr.

capitatus, Pall. ex Reichb. Fl. Crit. vi. 28 : liburnicus, Carmelitanum, *Reut. ex Boiss. Fl. Orient.* i. 512.—Armenia.

carolinianus, Walt. Fl. Carol. 140 = D. Armeria.

Carthusianorum, All. Fl. Pedem. ii. 74 = chinensis.

Carthusianorum, *Linn. Sp. Pl.* 409.—Europ. ; As. Min.

Carthusianorum, Pall. Ind. Taur. ; Bieb. Casp. App. 209 = capitatus.

Carthusianorum, Sibth. & Sm. Fl. Graec. Prod. i. 288 = intermedius.

caryophylloides, Schult. Obs. Bot. 78 = sylvestris.

Caryophyllus, *Linn. Sp. Pl.* 410.—Europ. ; As. occ.

catalaunicus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 684 = attenuatus.

caucasicus, Sims, Bot. Mag. t. 795 = chinensis.

caucasicus, Bieb. Fl. Taur. Cauc. i. 327 = praec.

Cephalotes, Ser. in DC. Prod. i. 356 = capitatus.

chinensis, *Linn. Sp. Pl.* 411.—Europ. ; As. temp.

chlorophyllus, Schur, Enum. Pl. Transs. 95 = Carthusianorum.

chlorolaucus, Fisch. in Hornem. Hort. Hafn. Suppl. 187 = chinensis.

cibrarius, *Clem. in Mem. Acc. Sc. Torin.* Ser. II. xvi. (1857) 256.—Graecia.

ciliatus, Costa, Fl. Catal. 86 = Costae.

ciliatus, *Guss. Ind. Sem. Hort. Bocc.* (1825) 5 ; *Pl. Rar.* 168.—Italia.

cincinnatus, *Lem. Illustr. Hortic.* xi. (1864) t. 388.—Japon.

cinnabarinus, *Sprun. ex Boiss. Diagn.* Ser. I. vi. 22.—Graecia.

cinnamomeus, Sibth. & Sm. Fl. Graec. Prod. i. 287. t. 400 = pallens.

cintranus, Boiss. & Reut. Pugill. Pl. Nov. 20 = pungens.

clavatus, Desf. ex Steud. Nom. ed. II. i. 499.—Hab. ?

clavatus, Sprang. Neue Entdeck. ii. 169 = Carthusianorum.

cognobilis, Timb. in Bull Soc. Bot. Fr. xi. (1864) 143 = Requienii.

Colensoi, *F. N. Williams, in Journ. Bot.* xxiii. (1885) 344.—Afr. austr.

collinicola, *Sprang. Nachr.* i. *Bot. Gart. Halle*, 20.—Hab. ?

collinus, Waldst. & Kit. Pl. Rar. Hung. i. 51 = chinensis.

collisegrus, Jord. in Billot, Annot. 46 = longicaulis.

DIANTHUS :—

compactus, Kit. in Schult. Oestr. Fl. ed. II. i. 654 = barbatus.

condensatus, Kit. in Linnæa, xxxii. (1863) 532 = monspessulanus.

congestus, Bor. Fl. Centr. Fr. ed. III. ii. 90 = Carthusianorum.

consanguineus, Schur, Enum. Pl. Transs. 93 = crunetus.

consimilis, Jord. in Billot, Annot. 47 = sylvestris.

contortus, Sm. in Rees, Cycl. xi. = fimbriatus.

contractus, Jan, ex Nym. Consp. 105 = Arrosti.

controversus, Gandin, Fl. Helv. iii. 157 = asper, monspessulanus.

cornutus, Hort. ex Steud. Nom. ed. II. i. 499 = clavatus.

coronarius, Lam. Fl. Fr. ii. 536 = D. Caryophyllus.

corsicus, Link, ex Spreng. Syst. ii, 876 = D. Caryophyllus.

corymbosus, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 867 = barbatus !

corymbosus, Poir. Encyc. Suppl. iv. 134 = Poiretianus.

corymbosus, *Sibth. & Sm. Fl. Graec. Prod.* i. 285 ; *Fl. Graec.* t. 395.—Archip. Graec. ; As. Min.

Costae, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 688.—Hispan.

Courtoisii, Reichb. Fl. Germ. Excurs. 806 = superbus.

Courtoisii, Schur, Enum. Pl. Transs. 96 = collinus.

cous, Beiss. Diagn. Ser. I. i. 20 = actinopetalus.

crassipes, *De Roem. in Linnaea*, xxv. (1852) 11.—Hispan.

crenatus, Eckl. & Zeyh. ex Sond. in Harv. & Sond. Fl. Cap. i. 123 = holopetalus.

crenatus, Gilib. Fl. Lituan. ii. 161 = deltoides.

crenatus, [Ker-Gawl.] in Bot. Reg. t. 256 = prostratus.

crenatus, E. Mey. Zwei Pfl. Docum. 172 = scaber.

crenatus, *Thunb. Prod. Pl. Cap.* 81.—Afr. austr.

cretaceus, Adam, in Web. & Mohr, Beitr. i. 56 = leptopetalus.

cibrarius, *Clem. Sert. Or.* 20. t. 2.—As. Min.

crinitus, *Sm. in Trans. Linn. Soc.* ii. (1794) 300.—As. Min. ; Persia ; Ind. bor. occ.

crosticus, *Borbas. ex Yusi, Jahresber.* iv. 1059.—Croatia.

crossopetalus, Fenzl, ex Boiss. Fl. Orient. i. 496 = chinensis.

cruentus, *Fisch. ex Planch. in Fl. des Serres*, Sér. I. v. (1849) t. 488.—Reg. Caucas. ?

cruentus, *Griseb. Spicil. Fl. Rumel.* i. 186.—Graecia.

cyathophorus, Moris, Ind. Sem. Hort. Taurin. (1852) 32 ; et in Ann. Sc. Nat. Sér. III. xix. (1853) 368 = Arrosti.

cyllenreus, Boiss. & Heldr. Diagn. Ser. II. i. 63 = pubescens.

Cyri, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 84.—As. Min. ; Assyria ; Afghanist.

dalmaticus, *Celak. in Oestr. Bot. Zeitschr.* xxxv. (1885) 189.—Dalmatia.

danubialis, Griseb. ex Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) t. 301 = banaticus.

decrescens ×, Borb. in Oestr. Bot. Zeitschr. xxvii. (1877) 378.—Europ.

deltoides, Georgi, Reisen, i. 215 = chinensis.

deltoides, Griseb. Spicil. Fl. Rumel. i. 192 = myrtinervius.

deltoides, *Linn. Sp. Pl.* 411.—Europ. ; Tibet. occ.

deltoides, Patrin, ex Steud. Nom. ed. II. i. 499 = chinensis.

dentosus, Fisch. ex Reichb. Fl. Crit. vi. 32. t. 546 = chinensis.

deserti, Kotschy, in Sitzb. Acad. Wien, Math. Nat. lii. Abth. i. (1866) 262.—Arab.

dichotomus, Pall. Reise Statth. Russ. Reich. ii. 735 = polymorphus !

diffusus, Griseb. Spicil. Fl. Rumel. i. 191 = Friwaldskyanus.

diffusus, Sibth. & Sm. Fl. Graec. Prod. i. 285 ; Fl. Graec. t. 396 = pubescens.

diminutus, Linn. Sp. Pl. ed. II. 587 = prolifer.

discolor, Sims, Bot. Mag. t. 1162 = chinensis.

distinus, Kit. ex Schult. Oestr. Fl. ed. II. i. 655 = polymorphus.

distinus, Reichb. Ic. Fl. Germ. vi. 44 = atrorubens.

divaricatus, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 302.—Archip. Graec.

dubius, Hornem. Hort. Hafn. i. 408 = arenarius, plumarius, suaveolens.

dubius, *Rafin. Caratteri*, 75.—Sicil.

Duffii ×, Hausskn. ex Aschers, in Oestr. Bot. Zeitschr. xxvi. (1876) 259 (an Leitgebii ?).—Europ.

DIANTHUS:—

dumulosus, Boiss. & Huet, Diagn. Ser. II. v. 53 = orientalis.
elatus, *Ledeb. Fl. Alt.* ii. 286.—Sibir. altaic.
elegans, *Bieb. ex Rupr. Fl. Cauc.* 174, in textu.—Reg. Caucas.
elegans, Urv. in Mém. Soc. Linn. Par. i. (1822) 302 = actinopetalus.
ellipticus, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 308.—Japon.
elongatus, *C. A. Mey. Vers. Pfl. Cauc.* 211.—Reg. Caucas.
emarginatus, Ser. in DC. Prod. i. 359 = pallens.
eriostomum, *Fourr. ex Lange, in Kjoeb. Vidensk. Meddel.* (1865) 109, nomen.—Hispan.
erinaceus, *Boiss. Diagn.* Ser. I. i. 21.—Graecia.
erubescens, *Roth, Enum. Pl. Phaen. Germ.* i. ii. 279.—Europ.
erythrocoleus, *Boiss. Fl. Orient.* i. 493.—Armenia.
Falconeri, *Edgew. & Hook f. Fl. Brit. Ind.* i. 214.—Tibet. occ.
fallens, Timb. in Bull. Soc. Bot. Fr. v. (1858) 329 = tener.
fasciculatus, Gilib. Fl. Lituan. ii. 161 = Carthusianorum.
fastigiatus, Pant. in Verh. Ver. Natur- u. Heilk. Presb. ii. 1871–72) 105 = crenatus.
Faurel, *Arvet, in Bull. Soc. Dauph.* (1880) 263.—Gallia.
ferrugineus, *Mill. Gard. Dict. ed.* VIII. n. 9.—Italia.
ferrugineus, Fourr. ex Gren. & Godr. Fl. Fr. i. 232 = Carthusianorum.
filiformis, Lam. Fl. Fr. ii. 537 = Tunica Saxifraga.
fimbriatus, Bieb. Fl. Taur. Cauc. i. 292 = orientalis.
fimbratus, Brot. Fl. Lusit. ii. 177 = Broteri.
fimbriatus, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 292 = crinitus.
fimbriatus, Lam. Fl. Fr. ii. 538 = monspeliacus, superbus.
Fischeri, Spreng. Cat. Sem. Hort. Hal. (1810); Pugill. ii. 62 = chinensis.
flaccidus, Fleber, in Flora, xvii. (1834) 583 = caesius.
floribundus, *Boiss. in Tchihat. Asie Min. Bot.* i. 221. —As. Min. ; Syria.
foliosus, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 184. —China.
fragrans, *Bieb. Fl. Taur. Cauc.* i. 331.—Reg. Caucas.
frigidus, Zucc. in Flora, vii. (1824) 383 = sylvestris.
Friwaldszkyanus, *Boiss. Diagn.* Ser. II. i. 65.—Rumelia.
fruticosus, *Linn. Sp. Pl.* 418.—Archip. Graec.
furcatus, *Balb. in Mém. Acad. Turin, Phys.-Math.* i. (1802–03) 18.—Pedemont.
furcatus, Bourg. ex Nym. Consp. 104 = tener.
furcatus, Hornem. Hort. Hafn. Suppl. 47 = Hornemanni.
gaditanus, *Boiss. Diagn.* Ser. II. i. 67.—Hispan.; Marocco.
gallicus, *Pers. Syn.* i. 495.—Gallia.
Casparrini, *Guss. Fl. Sic. Syn.* i. 479.—Sicil.
golidus, Schott, Nym. & Kotschy, Analect. Bot. 54 = glacialis.
geminatus, Kit. ex Schult. Oestr. Fl. ed. II. i. 656 = Hornemanni.
geminiflorus, Loisel. Fl. Gall. 725 = Seguieri.
giganteus, Urv. in Mém. Soc. Linn. Par. i. (1822) 301 = banaticus.
Girardini, *Lamotte, in Bull. Soc. Bot. Fr.* xxi. (1874) 190.—Gallia.
Gisellae, ×, Borb. in Oestr. Bot. Zeitschr. xxvii. (1877) 378.—Europ.
glacialis, DC. Fl. Fr. vi. 608 = neglectus.
glacialis, *Hoenke, in Vasc. Coll.* ii. 84.—Europ. alp.
glaucophyllus, Boiss. Fl. Orient. i. 514 = capitatus.
glaucophyllus, *Colla, in Mem. Acc. Torino*, xxxv. (1831) 187.—Hab. ?
glaucophyllus, Horoem. ex Ser. in DC. Prod. i. 356 = liburnicus.
glaucophyllus, Wierzb. ex Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) 2. 301 = banaticus.
glaucus, Buch.-Ham. in Wall. Cat. n. 656 = chinensis.
glaucus, Huds. Fl. Angl. ed. II. 185 = caesius.
glaucus, Linn. Sp. Pl. 411 = deltoides.
glaucus, Schousb. ex Ball, in journ. Linn. Soc. xvi. (1877) 355 = gaditanus.
glomeratus, Pall. ex Ledeb. Fl. Ross. i. 276 = polymorphus.
glumaceus, Bory & Chaub. Nouv. Fl. Pélop. 36 = Tunica glumacea.

DIANTHUS:—

glutinosus, *Boiss. & Heldr. Diagn.* Ser. II. i. 61.—Archip. Graec.
Godronianus, Jord. in Billot, Annot. 45 = virgineus.
gracilis, *Sibth. & Sm. Fl. Graec. Prod.* i. 288; *Fl. Graec.* t. 402.—Macedon.
graminifolius, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 11 = Carthusianorum.
graminifolius, *Presl, Fl. Sicul.* 147.—Sicil.
graminifolius, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 144 = tenuifolius.
grandiflorus, *Poir. in Lam. Encyc.* iv. 515.—Hispan.
graniticus, *Jord. Obs. Pl. Crit.* vii. (1849) 13.—Gallia.
gratianopolitanus, Vill. Hist. Pl. Dauph. iii. 598 = caesius.
Grembichi ×, Aschers. in Sitzb. Bot. Ver. Brandenb. xix. (1877) 28.—Europ.
Grisebachii, Boiss. Diagn. Ser. II. i. 62 = viscidus.
Gulise, Janka, in ‘Il. Barl.’ iii. (1874) 422; ex Journ. Soc. xii. (1874) 338 = ferrugineus.
gutsatus, Bieb. Fl. Taur. Cauc. i. 328 = chinensis.
Guyetani, Jord. in Billot, Annot. 46 = sylvestris.
haematocalyx, *Boiss. & Heldr. Diagn.* Ser. II. i. 68.—Graecia.
Hausknechtii, *Boiss. Fl. Orient.* i. 489.—As. Min.
Heldreichi ×, Borb. ex Aschers. in Oestr. Bot. Zeitschr. xxvi. (1876) 256.—Europ.
Hendersonianus ×, Paxt. Fl. Mag. xiv. (1848) 196.
Henteri, Heuff. ex Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 303 = D. Caryophyllus.
heptaneurus, Griseb. & Schenk, i. c. 302 = trifasciculatus.
hirsutus, Lam. Fl. Fr. ii. 538 = D. Armeria.
hirtus, Bieb. Fl. Taur. Cauc. i. 326 = humilis.
hirtus, Lam. Fl. Fr. ii. 538 = D. Armeria.
hirtus, *Vill. Hist. Pl. Dauph.* iii. 593. t. 46.—Gallia; Europ.
hispanicus, Asso, Syn. Stirp. Arag. 53 = pungens.
Hoeltzeri, *Winkler, in Regel, Gartenfl.* (1881) 1.—Turkestan.
holopetalus, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 309.—Afr. austr.
Hornemanni, Salzm. ex Boiss. Diagn. Ser. II. i. 67 = gaditanus.
Hornemanni, *Ser. in DC. Prod.* i. 262.—Reg. Mediterr.
hortensis, Schrad. ex Willd. Enum. Hort. Berol. 469 = plumarius.
humilis, *Willd. ex Ledeb. Fl. Ross.* i. 280.—Rossia austr.
hungaricus, Clem. Sert. Orient. 18 = crinitus.
hungaricus, Griseb. Spicil. Fl. Rumel. i. 198 = petraeus.
hungaricus, Pers. Syn. i. 494 = plumarius.
hybridus, Schmidt, ex Tausch, in Flora, xiii. (1830) 245 = D. Armeria.
hymenolepis, *Boiss. Diagn.* Ser. I. viii. 64.—Mesopotam.
hypanicus, Bess. ex Reichb. Fl. Germ. Excurs. 809 = campestris.
hypochlorus, *Boiss. & Heldr. Diagn.* Ser. I. viii. 67.—As. Min.
hyssopifolius, Linn. Cent. Pl. i. 11 = superbus.
iberica, Stev. ex Ledeb. Fl. Ross. i. 283 = crinitus.
ibericus, Willd. Enum. Hort. Berol. Suppl. 24 = chinensis.
inaequalis, E. Mey. Zwei Pfl. Docum. 178 = crenatus, prostratus.
incertus, Jacquem. ex Edgew. & Hook. f. Fl. Brit. Ind. i. 214 = angularis.
inclusus, Walp. Rep. i. 368 = Saponaria inclusa.
incurvus, Thunb. Prod. Pl. Cap. 81 = albens.
inodorus, Gaertn. Fruct. ii. 297. t. 159. f. 18 = attenuatus.
inodorus, Pall. Reise, ii. 723 = chinensis.
inodorus, Stend. Nom. ed. I. 371 = sylvestris.
insignitus, Timb. in Bull. Soc. Bot. Fr. xi. (1864) 148 = *integer*, Vis. in Flora, xii. (1829) i. Erg. 11 = strictus.
integerrimus, Bunge, in Mém. Acad. Pétersb. Sér. VI. vii. (1856) 583 = Liboschitzianus.
integripetalus, Schur, Enum. Pl. Transs. 98 = petraeus.
intermedius, *Boiss. Fl. Orient.* i. 515.—Rumel.; As. Min.
intermedius, Willd. ex Ledeb. Fl. Ross. i. 276 = polymorphus.
involucratus, Pall. ex Ledeb. l. c. 277 = chinensis.
involucratus, Poir. Encyc. Suppl. iv. 133 = rupicola.
Italicus, *Pers. ex Pasquale, Ind. Sem. Hort. Neap.* (1846) xix.—Italia.
Jacquemontii, *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 214.—Reg. Himal.

DIANTHUS:—

Jacuensis ×, Aschers. in Oestr. Bot. Zeitschr. xxvi. (1876) 257.—Europ.
japonicus, Thunb. Fl. Jap. 188. t. 38 = barbatus.
jemisensis, Less. ex Ledeb. Fl. Ross. i. 278 = chinensis.
judaicus, *Boiss. Diagn.* Ser. I. viii. 66.—Syria.
juniperinus, *Sm. in Trans. Linn. Soc. ii.* (1794) 308.—Ins. Creta.
juratensis, Jord. in Billot, Annot. 47 = sylvestris.
kamiabergensis, *Sond. in Harv. & Sond. Fl. Cap.* i. 124.—Afr. austr.
Karelinianus, Schur, Enum. Pl. Transs. 97 = D. Caryophyllus.
Knappii, Aschers. ex Nym. Consp. 103 = liburnicus.
Kotschyanus, Boiss. & Heldr. Diagn. Ser. I. viii. 68 = anatolicus.
Kremeri, *Boiss. & Reut. Pugill. Pl. Nov.* 21.—Hispan.
Kuschakewiczi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 244.—Sibir. altaic.
lactiflorus, *Fenzl, in Tchihat. Asie Min. Bot.* i. 210. —Cilicia.
lanceolatus, Ster. ex Eichw. Skizze, 172 = bicolor.
lancifolius, Tausch, in Flora, xiv. (1831) 215 = trifasciculatus.
Langeanus, *Willd. in Willd. & Lange, Prod. Fl. Hisp.* iii. 890.—Hispan.
laricifolius, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 7.—Hispan.
latifolius, Willd. Enum. Hort. Berol. 466 = barbatus.
leptolomus, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 42.—Abyss.
leptopetalus, *Willd. Enum. Hort. Berol.* 466.—Macedon.; Reg. Caucas.
leucophaeus, *Sibth. & Sm. Fl. Graec. Prod.* i. 288; *Fl. Graec.* t. 405.—Bithynia.
leucophaeus, Sieber, Reise Kreta, ii. 220 = sphaciotes
Levieri, *Borb. in Oestr. Bot. Zeitschr.* xxvii. (1877) 231.—Ital.
Libanotis, *Labill. Ic. Pl. Syr.* i. 14. t. 5.—Syria.
Liboschitzianus, Hohen. ex Boiss. Fl. Orient. i. 491 = fragrans.
Liboschitzianus, *Ser. ex DC. Prod.* i. 360.—As. Min.
liburnicus, *Bartl. & Wendl. f. Beitr.* ii. 52.—Europ. austr.
ligusticus, Willd. ex Nym. Consp. 103 = liburnicus.
lilacinus, *Boiss. & Heldr. Diagn.* Ser. II. i. 63.—Graecia.
liliodorus, *Panl. ex Nym. Consp.* 105, nomen.—Serb.
linnaeiatus, [Bové] Delile, Ind. Sem. Hort. Monsp (1838) ex Ann. Sc. Nat. Sér. II. vii. (1837) 268 = multipunctatus.
litoralis, Host, Fl. Austr. i. 522 = ciliatus.
longicalyx, *Miq. in Journ. Bot. Néerl.* i. (1861) 127.—China.
longicaulis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = virgineus.
longicaulis, Costa, Fl. Catal. 86 = D. Caryophyllus.
longicaulis, *Tenore, Cat. Hort. Neap.* ed. II. (1819) 77, *ex Fl. Nap.* ii. 379.—Europ. austr.
longiflorus, Poir. in Lam. Encyc. iv. 522 = attenuatus.
longiginnis, *Delile, in Ann. Sc. Nat. Sér.* II. xx. (1843) 80.—Abyss.
Lucae ×, Aschers. in Oestr. Bot. Zeitschr. xxvi. (1876) 259.—Europ.
lusitanus, Brot. Fl. Lusit. ii. 177 = attenuatus.
lydus, *Boiss. Diagn.* Ser. I. i. 20.—As. Min.
macranthus, *Boiss. l. c.* 28.—Persia.
macrolepis, Boiss. l. c. 18 = Cyri.
macrolepis, Fenzl, ex Boiss. l. c. viii. 64 = stenocephalus.
macrosyen, Fenzl, ex Boiss. Fl. Orient. i. 495 = orientalis.
malacitanus, Haensel. ex Boiss. Voy. Bot. Espagne, ii. 85 = Broteri.
marginatus, *Port. Encyc. Suppl.* iv. 181.—Jura.
marsicus, Tenore, Fl. Nap. iv. 61 = monspessulanus.
massmenseus, *Boiss. Diagn.* Ser. II. v. 51.—As. Min.
mauritanicus, *Pomel, Nouv. Mat. Fl. Atl.* 393.—Algeria.
Mayeri, *Presl, Bot. Bemerk.* 18.—Sibiria.
medius ×, Boss. ex Nym. Consp. 108.
membranaceus, Borb. in Oestr. Bot. Zeitschr. xxvi. (1876) 125 = polymorphus.
Mercurii, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 287 = gracilis.
microlepis, *Boiss. Diagn.* Ser. I. i. 23.—Thessalia.

DIANTHUS:—
micropetalus, Eckl. & Zeyh. Enum. 81 = kamisbergensis.
micropetalus, Ser. in DC. Prod. i. 359 = scaber.
Mihi ×, Reichardt, in Verh. Zool. Bot. Ges. Wien, xvii. (1867) 881.—Europ.
minutus, Host, ex Nym. Consp. 105 = multinervis.
moesiacus, *Vis. & Panč. in Mem. Ist. Venet.* xv. (1870) 17.—Europ.
monadelphus, Vent. Jard. Cels, t. 89 = pallens.
monspeliacus, Linn. Syst. ed. X. 1029 = monspessulanus.
monspessulanus, *Linn. Cent.* ii. n. 146 ; *Amœn. Acad.* iv. 313.—Europ. austr.
monspessulanus, Wulf. in Roem. Arch. iii. iii. 360 = erubescens.
montanus, Bieb. Fl. Taur. Cauc. i. 328 = chinensis.
montanus, F. W. Schmidt, in Neu Abh. Boehm. Ges. i. (1791) 80 = Carthusianorum.
Morrisii, *Hance, in Hook. Lond. Journ. Bot.* vii. (1848) 473.—China.
moschatus, J. F. Gmel. Syst. 711 = D. Caryophyllus ?
moschatus, Hort. Par. ex Ser. in DC. Prod. i. 363 = plumarius.
multicaulis, Boiss. & Huet, Diagn. Ser. II. v. 53 = Liboschitzianus.
multicaulis, Pall. ex Ledeb. Fl. Ross. i. 280 = hamilis.
multiceps, *Costa, ex Willk. in Linnæa*, xxx. (1859–60) 88.—Hispan.
multifidus, Gilib. Fl. Lituan. ii. 162 = superbus.
multifidus, Willd. ex Ser. in DC. Prod. i. 364 = fragrans.
multinervis, *Vis. Fl. Dalmat.* iii. 164.—Dalmatia.
multipunctatus, *Ser. in DC. Prod.* i. 362.—Archip. Græc. ; Syria.
multisquamatus, Hochst. ex Boiss. Fl. Orient. i. 497 = sinaicus.
multisquamatus, *F. N. Williams, in Journ. Bot.* xxiii. (1885) 344.—Kurdistan.
muschianus, *Kotschy, ex Boiss. Fl. Orient.* i. 510.—Armenia.
Mussini, Hornem. Hort. Hafn. i. 408 = squarrosus.
mutabilis, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 344. t. 2 = masmenseus.
myrtinervius, *Griseb. Spicil. Rumel.* i. 194.—Macedon.
nanus, Sweet, Hort. Brit. ed. I. 41 = Carthusianorum.
nardiformis, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 196.—Turcia.
nazarenus, Clarke, Trav. ii. 420 = serratifolius.
neglectus, *Loisel. in Desv. Journ. Bot.* ii. (1809) 321. —Europ.
nitidus, *Waldst. & Kit. Pl. Rar. Hung.* ii. 209.—Hungar. ; Macedon.
nodosus, Tausch, in Syll. Ratisb. ii. (1828) 248 = longicaulis.
Noëanus, *Boiss. Diagn.* Ser. II. i. 67.—Rumelia.
Notanus, Boiss. l. c. v. 52 = floribundus.
nudiflorus, Griff. Notul. iv. 466 ; Ic. Pl. Asiat. t. 580 f. 5 = Velezia rigida.
obcordatus, Rent. in Mém. Soc. Phys. Genèv. viii. (1839) 281 = glumaceus.
obtusifolius, *Scheele, in Flora*, xxvi. (1843) 428.—Europ.
ochroleucus, Link, Enum. Hort. Berol. i. 420 = chinensis.
ochroleucus, Pers. Syn. i. 494 = pallens.
ocluatus, *Boiss. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 345. t. 2.—Cilicia.
odoratissimus, Vest, ex Reichb. Fl. Germ. Excurs. 807 = monspessulanus.
odoratus, Vest, ex Steud. Nom. ed. II. i. 500 = plumarius.
omithogalaceus ×, Kern. in Oestr. Bot. Zeitschr. xv. (1865) 309.—Europ.
olympicus, Boiss. Diagn. Ser. I. i. 19 = viscidus.
olympicus, Sibth. ex Boiss. Fl. Orient. i. 467 = leucophæus.
Orsedax, Balb. ex Nym. Consp. 104 = monspessulanus.
orsedeum, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 207 = superbus.
orientalis, *Duval, Hort. Cantab.* ed. IV. 101 ; *Sims, Bot. Mag.* t. 1069.—Orient.
orophilus, Jord. in Billot, Annot. 43 = sylvestris.
pallens, Bieb. Fl. Taur. Cauc. i. 325 = pallidiflorus.
pallens, *Sibth. & Sm. Fl. Græc. Prod.* i. 326 ; *Fl. Græc.* t. 998.—Græcia ; As. Min. ; Persia.
pallidiflorus, *Ser. in DC. Prod.* i. 358.—Reg. Caucas ; Soongaria.
paniculatus, Gueldenst. Reis. i. 68, 110 = pallidiflorus.
papillosus, *Vis. & Panč. in Mem. Ist. Venet.* x. (1861) 494.—Europ.
parnassicus, Boiss. & Heldr. Diagn. Ser. II. i. 64 = viscidus.
parviflorus, Boiss. l. c. Ser. I. i. 21 = anatolicus.
parviflorus, Willd. ex Ledeb. Fl. Koss. i. 279 = pallidiflorus.
patens, Willd. Enum. Hort. Berol. Suppl. 24 = chinensis.
pavonicus, *Tausch, in Flora*, xxii. (1839) 145.—Helvetia.
pectinatus, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 124 = prostratus.
pelviformis, *Heuff. in Flora*, xxxvi. (1853) 625.—Serv.
pendulus, *Boiss. & Blanche, Diagn.* Ser. II. vi. 28.—Syria.
Pennynae ×, Sweet, Hort. Brit. ed. II. 51.
petræus, Bieb. Fl. Taur. Cauc. i. 328 = Liboschitzianus.
petræus, *Waldst. & Kit. Pl. Rar. Hung.* iii. 246.—Hungaria ; Thracia.
Petteri, *Boiss. Diagn.* Ser. II. i. 66.—Dalmatia.
pinifolius, *Sibth. & Sm. Fl. Græc. Prod.* i. 284.—Archip. Græc.
Planellas, *Willk. Ic. et Desc. Pl. Hisp.* i. 79. t. 58.—Hispan.
platyphyllus, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 1. 358.—Japon.
plumarius, Auct. ex Pers. Syn. i. 494 = monspessulanus.
plumarius, Gunn. ex Spreng. Syst. ii. 379 = superbus.
plumosus, *Linn. Sp. Pl.* 411.—Europ. ; As. bor.
plumosus, DC. ex Spreng. Pugill. ii. 64 = monspessulanus.
pogonopetalus, Boiss. & Kotschy, Diagn. Ser. II. vi. 29 = orientalis.
Poiretianus, Ser. in DC. Prod. i. 360 = corymbosus.
polycladus, *Boiss. Diagn.* Ser. I. viii. 55.—Syria.
polylepis, *Bienert, ex Boiss. Fl. Orient.* i. 497.—Persia.
polymorphus, *Bieb. Fl. Taur. Cauc.* i. 324.—Reg. Caucas.
pomeridianus, Bieb. l. c. 329 = leptopetalus.
pomeridianus, *Linn. Sp. Pl.* ed. II. 1673.—Palestin.
Pontederae, A. Kern. Sched. Fl. Exsicc. ii. 67 = D. Carthusianorum.
ponticus, Wahlenb. in Isis, xxi. (1828) 985 = intermedius.
portensis, Libosch. ex Ser. in DC. Prod. i. 363 = plumarius.
præcox, Willd. ex Spreng. Syst. ii. 381 = arenarius.
pratensis, Bieb. Fl. Taur. Cauc. iii. 300 = chinensis.
pratensis, S. F. Gray, Nat. Arr Brit. Pl. ii. 643 = D. Armeria.
pratensis, Lepech. It. iii. 185 = deltoides.
prolifer, Vent. Jard. Cels, Index = pallens.
prolifer. Forsk. Fl. Aegypt. Arab. p. xxv = Aphyllanthes monspeliensis.
prolifer, Linn. Sp. Pl. 410 = Tunica prolifera.
propinquus, Schur, Enum. Pl. Transs. 94 = liburnicus.
prostratus, Eckl. & Zeyh. Enum. 82 = scaber, crenatus.
prostratus, *Jacq. Hort. Schœnbr.* iii. 11. t. 271.—Alp. austr.
prunosus, Boiss. & Orph. Diagn. Ser. II. vi. 28 = hæmatocoalyx.
pruinosus, Janka, Del. Sem. Hort. Vindob. (1858) 4 ; et Linnæa, xxx. (1859–60) 748 = banaticus.
Pseud-Armeria, *Bieb. Fl. Taur. Cauc.* i. 323 ; iii. 297. —Reg. Caucas.
Pseud-Armeria, Wiersb. ex Schur, Enum. Pl. Transs. 91 = D. Armeria.
Pseudo-barbatus, Bess. ex Reichb. Fl. Germ. Excurs. 805 = liburnicus.
Pseudo-carthusianorum, Schur, Enum. Pl. Transs. 98 = petræus.
puberulus, *F. N. Williams, in Journ. Bot.* xxiii. (1885) 344.—Persia.
pubescens, Flach. Cat. Hort. Gorenk. (1612) 59 = leptopetalus.
pubescens, *Sibth. & Sm. Fl. Græc. Prod.* i. 286.—Græcia.
pubescens, Urv. in Mém. Soc. Linn. Par. i. (1822) 303 = glutinosus.
DIANTHUS:—
pulchellus, Fers. Syn. i. 495 = caesius.
pulchellus. Schrad. ex G. Don, Gen. Syst. i. 385 = chinensis.
pulcher, Salisb. Prod. 303 = chinensis.
pulcherrimus, Loisel. in Dict. Sc. Nat. xxxv. 417 = barbatus.
pumilus, Ftvald. ex Boiss. Diagn. Ser. I. i. 22 = microlepis.
pumilus, *Vahl, Symb. Bot.* i. 32.—Arab.
punctatus, *Spreng. Neue Entdeck.* ii. 169.—Hab. ?
pungens, J. Gay, ex Boiss. Voy. Bot. Espagne, ii. 85 = brachyanthus.
pungens, Gren. & Godr. Fl. Fr. i. 284 = furcatus.
pungens, *Linn. Mant.* ii. 240.—Hispan.
pungens, Poir. Encyc. iv. 526 = virgineus.
purpureus, Poir. in Lam. Encyc. iv. 528 = pungens ?
purpureus, *F. N. Williams, in Journ. Bot.* xxiii. (1885) 343.—Syria.
pusillus, *F. W. Schmidt, in Neue Abh. Boehm. Ges.* i (1791) 29.—Europ.
pyrenaicus, *Bernh. ex Steud. Nom.* ed. II. i. 500.—Mont. Pyren.
pyrenaicus, Pourr. in Mém. Acad. Toul iii. (1788) 318 = attenuatus.
quadrangularis, *Haberle, ex Steud. Nom.* ed. II. i. 500. —Hab. ?
quadrilobus, Boiss. in Tchihat. Asie Min. Bot. i. 222 = sulcatus.
Querterii ×, Carr. ex Hérincq, L'Horticulteur Franç. (1866) 43.
racemosus, Vis. in Flora, xii. (1829) 1. Erg. 12 = ciliatus.
ramentaceus, Fenzl, ex Sond. in Harv. & Sond. Fl. Cap. i. 122 = scaber.
ramosissimus, Pall. ex Poir. Encyc. Suppl. iv. 130 = pallidiflorus.
rechcaulis, Ledeb. Fl. Ross. i. 287 = Tunica stricta.
recurvus, Fisch. ex Ledeb. l. c. 284 = squarrosus.
repens, Willd. Sp. Pl. ii. 681 = alpinus.
Requienii, *Godr. in Gren. & Godr. Fl. Fr.* i. 284.—Europ. austr.
Reuterî, Jord. in Billot, Annot. 49 = sylvestris.
revolutus, Tausch, in Flora, xiii. (1830) 245 = superbus.
rigidus, *Bieb. Fl. Taur. Cauc.* i. 325.—Tauria.
robustus, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* i. 492.—Armenia.
rosulatus, Borb. ex Nym. Consp. 103 = liburnicus.
rupestris, Friwald. ex Griseb. Spicil. Fl. Rumel. i. 191 = pubescens.
rupestris, Lam. Fl. Fr. ii. 536 = virgineus.
rupestris, Linn. f. Suppl. 240 = sylvestris.
rupicolus, *Biroena, Sicul. Pl. Cent.* i. 31.—Sicilia.
rupicolus, Jord. ex Nym. Consp. 105 = sylvestris.
rupicolus, Schur, Enum. Pl. Transs. 93 = Carthusianorum.
ruthenicus, Roem. ex Poir. Encyc. Suppl. iv. 131 = chinensis.
sabuletorum, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 26 = Carthusianorum.
sabuli, Kit. in Linnæa, xxxii. (1863) 530 = polymorphus.
sabulosus, Willd. ex Ledeb. Fl. Ross. i. 284 = squarrosus.
saetabensis, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 44.—Hispan.
Samaritani, Heldr. ex Boiss. Fl. Orient. i. 511 = cinnabarinus.
sanguineus, Vis. Stud. Nom. Rar. Hort. Patav. (1845) 1 = atrorubens.
Sartorii, Frenzl, ex Nym. Consp. 107 = velutinus.
saxatilis, Pall. ex Bieb. Fl. Taur. Cauc. i. 329 = pallens.
saxatilis, Pers. Syn. i. 494 = monspessulanus.
saxicola, Jord. Pugill. 29 = sylvestris.
saxifragus, Linn. Sp. Pl. 415 = Tunica saxifraga.
scaber, Chaix, in Vill. Hist. Pl. Dauph. i. 331 = hirtus.
scaber, Eckl. & Zeyh. Enum. 81, partim = crenatus.
scaber, Schleich. ex Suter, Fl. Helv. i. 259 = chinensis.
scaber, *Thunb. Prod. Pl. Cap.* 81.—Afr. austr.
Schenckseri, Jord. Pugill. 29 = longicaulis.
Schenckseri, Reichb. Fl. Germ. Excurs. 811 = sylvestris.
schizopetalus, Wallr. in Linnæa, xiv. (1840) 570 = superbus.

DIANTHUS:—
Schlosseri, *F. N. Williams, in Journ. Bot.* xxiii. (1885) 342.—Europ. austr.
Schweinfurthoi, Cosa. ex Ball, in Journ. Linn. Soc. xvi. (1877) 365 = gaditanus.
Schraderi, Reichb. Hort. Bot. i. 24 = chinensis.
scoparius, *Pavol, ex Boiss. Fl. Orient.* i. 494.—Persia.
Seguieri, Costa, ex Willk. & Lange, Prod. Fl. Hisp. iii. 681 = hirtus.
Seguierii, Vill. Prosp. 48; Chaix, in Vill. Hist. Pl. Dauph. i. 330; iii. 594 = chinensis.
Seidlitzii, *Boiss. Fl. Orient.* i. 506.—Persia.
serotinus, Waldst. Kit. Pl. Rar. Hung. ii. 188. t. 172 = plumarius.
Serpe, *Ficalho & Hiern, in Trans. Linn. Soc. Ser.* II. ii. (1881) 17.—Afr. trop.
serratifolius, *Sibth. & Sm. Fl. Graec. Prod.* i. 287.—Graecia.
serratus, Lapeyr. Hist. Abr. Pl. Pyr. 241 = chinensis.
serrulatus, Boiss. Voy. Bot. Espagne, ii. 84. t. 28 = Broteri.
serrulatus, *Desf. Fl. Atlant.* i. 346.—Afr. bor.
sicalus, *Y. & C. Presl, Delic. Prag.* 59.—Reg. Mediterr.
silenoides, *Poir. in Lam. Encyc.* iv. 514.—Hab.?
sinaicus, *Boiss. Diagn. Ser.* I. i. 23.—Arab.
sinensis, Link, Handb. ii. 287 = chinensis.
speciosus, Reichb. Ic. Fl. Germ. vi. 46 = monspessulanus.
sphaciotieus, *Boiss. & Heldr. Diagn. Ser.* I. viii. 70.—Ins. Creta.
spiculifolius, Schur, Enum. Pl. Transs. 98 = petraeus.
spinosus, Desf. in Ann. Mus. Par. i. (1802) 198. t. 16. f. 1 = Acanthophyllum spinosum.
Sprengelii, G. Don, Gen. Syst. i. 394 = monspessulanus.
sparius ×, Kern. ex Borb. in Rad Jugos. Akad. Zagreb. xxxvi. (1876) 173.—Europ.
squarrosus, *Bieb. Fl. Taur. Cauc.* i. 331.—Tauria.
stenocephalus, *Boiss. Diagn. Ser.* I. i. 19.—Persia.
stenopetalus, *Griseb. Spicil. Fl. Rumel.* i. 187.—Thessal.; Macedon.
sterilis, Stev. ex Boiss. Fl. Orient. i. 505 = humilis.
Sternbergii, Schleich. ex Ser. in DC. Prod. i. 365 = monspessulanus.
Sternbergii, *Sibth. ex [Capelli] Cat. Hort. Taur.* (1931) 24.—Tauria.
stramineus, *Boiss. & Heldr. Diagn. Ser.* I. viii. 70.—As. Min.
striatifolius, Fenzl, Pugill. Pl. Nov. Syr. 10 = multipunctatus.
strictus, *Sibth. & Sm. Fl. Graec. Prod.* i. 288.—Dalmat.; Graecia.
strictus, [Soland. in] Russell, Aleppo, ed. II. ii. 252 = syriacus.
suaveolens, *Spreng. Nov. Prov.* 16.—Hab.?
suavis, Willd. Enum. Hort. Berol. Suppl. 24 = caesius.
subacaulis, Cutanda, Fl. Madrid, 170 = brachyanthus.
subacaulis, Vill. Hist. Pl. Danph. iii. 597 = virgineus.
subbarbatus, *Schur, Enum. Pl. Transs.* 98.—Transylv.
subbiflorus, *Fourr. ex Lange, in Kjoeb. Vidensk. Meddel.* (1865) 109, nomen.—Hispan.
subneglectus, Schur, Enum. Pl. Transs. 95 = Carthusianorum.
subfruticosus, Willd. Enum. Hort. Berol. 466 = ruptícola.
sulcatus, *Boiss. Fl. Orient.* i. 488.—As. Min.
superbiens, Kotschy, ex Boiss. l. c. 492 = robustus.
superbus, Eichw. Casp. Cauc. 31 = monspessulanus.
superbus, *Linn. Amoen. Acad.* iv. 272.—Europ.; As. bor.
supertus, Willk. ex Willk. & Lange, Prod. Fl. Hisp. iii. 684 = Broteri.
supinus, Lam. Fl. Fr. ii. 584 = deltoides.
sylvaticus, *Hoppe, ex Willd. Enum. Hort. Berol.* 467; *Sturm. Deutschl. Fl.* Heft 38.—Europ.
sylvestris, Boiss. Voy. Bot. Espagne, ii. 85 = Boissieri.
sylvestris, *Wulf. in Jacq. Coll.* i. 237.—Europ.
syriacus, *Stend. Nom.* ed. II. i. 501.—Syria.
syriacus, F. N. Williams, in Journ. Bot. xxiii. (1885) 346 = pubescens?
Szowitsianus, *Boiss. Fl. Orient.* i. 503.—Reg. Caucas.
tabrisianus, *Bienert, ex Boiss. l. c.* 496.—Persia; Afghanist.
talyschensis, *Boiss. & Buhse, in Nouv. Mém. Masc.* xii. (1860) 34.—Reg. Caucas.
tanaiticus, *Gueldenst. ex Ledeb. Fl. Ross.* i. 287 (quid?).—Rossia.
tataricus, Fisch. Hort. Gorenk. ed. II. (1812) 59 = chinensis.
Tauscheri, Everam. ex Ledeb. Fl. Ross. i. 284 = acicularis.

DIANTHUS:—
tener, *Balb. in Mém. Acad. Turin, Phys.-Math.* i. (1802–3) 14.—Italia.
tenuiflorus, *Griseb. Spicil. Fl. Rumel.* i. 189.—Thracia; Macedon.
tenuifolius, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 148 = Carthusianorum.
tergestinus, Reichb. Ic. Fl. Germ. vi. t. 266 = virgineus.
thessalus, *Orph. in Bull. Congr. Bot. Pétersb.* 1869 (1870) 132.—Graecia.
tolctanus, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 7.—Hispan.
transylvanicus, Schur, in Verh. Siebenb. Ver. Naturw. v. (1854) 82 = trifasciculatus.
tricolor, Adam, ex Ser. in DC. Prod. i. 360 = pomeridianus.
trifasciculatus, *Kit. ex Schult. Oestr. Fl.* ed. II. i. 654. —Hungar.
tripunctatus, *Sibth. & Sm. Fl. Graec. Prod.* i. 286.—Archip. Graec.
tymphresteus, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. vi. 37.—Graecia.
umbellatus, DC. Cat. Hort. Monsp. 104 = chinensis.
uniflorus, Fork. Fl. Aegypt. Arab. p. cxi = pumilus.
uniflorus, Gilib. Fl. Lituan. ii. 162 = armeria.
vaginatus, Reichb. Ic. Fl. Germ. vi. 44 = strorubens.
vaginatus, Vill. Hist. Pl. Dauph. iii. 594 = Carthusianorum.
valentinus, Willk. Ic. et Descr. Pl. Hisp. i. 14. t. 7 = Broteri.
velutinus, Guss. Pl. Rar. 166 = Tunica velutina.
Verschaffeltii ×, Lem. Ill. Horticr. vi. (1859) t. 220.
versicolor, Fisch. ex Link, Enum. Hort. Berol. i. 430 = chinensis.
villosus, Gilib. Fl. Lituan. ii. 160 = D. Armeria.
virgatus, *Pasq. in Ann. Acc. Asp. Nat.* Ser. III. i. (1861) 28.—Italia.
virgineus, Hall. Taur. 119, ex Bieb. Fl. Taur. Cauc. i. 326 = humilis.
virgineus, *Linn. Sp. Pl.* 412.—Europ. austr.; Afr. bor. occ.
virgineus, Sims, Bot. Mag. t. 1740 = sylvestris.
viridescens, *Vis. Ort. Bot. Padov.* 138.—Dalmatia.
viscidus, *Bory & Chaub. Nouv. Fl. Pélop.* 36.—Graecia.
vivariensis, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 91 = hirtus.
volgensis, Hort. ex Ser. in DC. Prod. i. 361 = deltoides.
Vukotinovičii ×, Borb. in Rad Jugos. Akad. Zagreb. xxxvi. (1876) 172.
vulturius, Guss. & Tenore, Ind. Sem. Hort. Neap. (1827) 3; (1839) 11 = liburnicus.
Waldsteinii, Sternb. in Flora, ix. (1826) 1. Beil. 73 = monspessulanus.
Webbianus, Parol. in Vis. Illustr. Alc. Piant. 20. t. 8. f. 2 = erinaceus.
Willdenowii, Link, Enum. Hort. Berol. i. 430 = chinensis.
Wiemmeri, Wichura, in Jahresb. Schles. Ges. (1854) 75; et Flora, xxxix. (1856) 127 = superbus.
Wolfii ×, Vetter, in Bull. Soc. Murith. xi. (1883) 8, 125 = Courtoisii.
Wulfenii, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 669 = sylvestris.
xylorrhizus, *Boiss. & Heldr. Diagn.* Ser. I. viii. 67.—Graecia.
Zayheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 194.—Afr. austr.
zonatus, *Fenzl, Pugill. Pl. Nov. Syr.* 11.—Graecia; As. Min.

DIAPASIS, Poir. Encyc. Suppl. ii. 468 (1811) = Dianpasis, R. Br. (Goodeu.).

DIAPEDIUM, Koen. Stend. Nom. ed. II. i. 501, 504 (1840) = Diolpetes, Juss. (Acanth.).

DIAPENSIA, Linn. Gen. ed. I. 82 (1737). *DIAPENSIACEÆ*, Benth. & Hook. f. ii. 619.
Rembertia, Adans. Fam. ii. 226 (1763).
americana, Banks, ex Sims, Bot. Mag. sub t. 1108 = Pyxidanthera barbulata.
barbulata, Ell. Sketch, i. 229 = Pyxidanthera barbulata.
cuneifolia, Pursh, Fl. Am. Sept. i. 148 = Pyxidanthera barbulata.
helvetica, Linn. Sp. Pl. 141 = Androsace bryoides.

DIAPENSIA:—
himalaica, *Hook. f. & Thoms. in Hook. Kew Journ.* ix. (1857) 378.—Reg. Himal.
japonica, J. F. Gmel. Syst. 1589, sphalm. = seq.
lapponica, *Linn. Sp. Pl.* 141.—Reg. bor. alp.
obtusifolia, Salisb. Parad. Lond. t. 104 = lapponica.

DIAPENZIA, Dum. Anal. Fam. 28 (1829) = prec.

DIAPERIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 287 = Evax, Gaertn. (Compos.).
multicaulis, Nutt. l. c. = E. multicaulis.
prolifera, Nutt. l. c. = E. prolifera.

DIAPHANE, Salisb. in Trans. Hort. Soc. i. (1812) 304 = Iris, Tourn.
edulis, Salisb. l. c. = Moraea Sisyrinchium.
stylosa, Salisb. l. c. 305 = I. juncea.

DIAPHORA, Lour. Fl. Cochinch. 578 (1790) = Soleria, Berg. (Cyperac.).
cochinchinensis, Lour. l. c. = S. elata.

DIAPHORANTHEMA, Beer, in Flora, xxxvii. (1854) 849; Bronn. 158 (1857) = Tillandsia, Linn. (Bromel.).
bifora, Beer, Bromel. 156 = T. bifora.
capillaris, Beer, l. c. 158 = T. capillaris.
recurvata, Beer, l. c. 156 = T. recurvata.
subulata, Beer, l. c. 155 = T. stricta.
trifora, Beer, l. c. = T. trifora.
umiflora, Beer, l. c. 154 = T. recurvata.
verticolor, Beer, l. c. 155 = T. setacea.
virescens, Beer, l. c. 154 = T. virescens.

DIAPHORANTHUS, Anders. ex Hook. f. Fl. Antarct. 288 (1846) = Pringlea, Anders. (Crucif.).

DIAPHORANTHUS, Meyen, Reise, i. 406 (1834) = Polysachyrus, Lag. (Compos.).
fuscus, Meyen, l. c. = P. sphaerocephalus, niveus.

DIAPHOREA, Pers. Syn. ii. 576 (1807) = DIAPHORA, Lour. = Soleria, Berg. (Cyperac.).

DIAPHYLLUM, Hoffm. Gen. Umbell. 115 (1814) = Bupleurum, Linn. (Umbell.).
aureum, Hoffm. l. c. = B. aureum.
longifolium, Hoffm. l. c. = B. longifolium.
triradiatum, Hoffm. l. c. = B. triradiatum.

DIARINA, Rafin. in Med. Repos. N. York, v. (1808) 352, et Cat. 13 (1824) = Diarrhena, Beauv. (1812) (Gramin.).
festucoides, Rafin. ll. cc. = Diarrhena americana.
sylvatica, Rafin. in Journ. Phys. lxxiix. (1819) 104 = Diarrhena americana.

DIARRHENA, Beauv. Agrost. 142 (1812). *GRAMINEÆ*, Benth. & Hook. f. iii. 1190.
Corycarpus, Zea, ex Spreng. Syst. i. 128 (1825).
Diarina, Rafin. in Med. Repos. N. York, v. (1808) 352; Cat. 13 (1824), nomen prius.
Korycarpus, Zea, in Act. Matr. (1806); ex Lag. Gen. et Sp. Nov. 4 (1816).
Onoa, Trumbull. & Sav. Enum. Pl. Jap. ii. 178 (1879).
Rosmeria, Zea, in Roem. & Schult. Syst. i. 61, 387 (1817).
americana, *Beauv. Agrost.* 142. t. 25.—Am. bor.
japonica, *Franch. & Sav. Enum. Pl. Jap.* ii. 608.—Japon.
setacea, Roem. & Schult. Syst. i. 289 = Festuca setacea.

DIARTHRON, Turcz. in Bull. Soc. Nat. Mosc. (1852) 204; et (1855) ii. 464. t. 11. *THYMELÆACEÆ*, Benth. & Hook. f. iii. 192.
coriaceum, Jaub. & Spach, Illustr. ii. 8. t. 105 = vesiculosum.
linifolium, *Turcz. in Bull. Soc. Nat. Mosc.* (1852) 204. —Mongolia.
vesiculosum, *Endl. Gen. Suppl. iv.* 11. 63.—Persia; Afghan.

DIASCIA, Link & Otto, Ic. Pl. Sel. 7. t. 3 (1820). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 981.
alonsoides, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 17.—Afr. austr.
Barbarae, *Hook. f. Bot. Mag.* t. 5953.—Afr. austr.

DIASCIA:—
Bergiana, Benth. in Hook. Comp. Bot. Mag. ii. (1836) 17 = saccnlata.
Bergiana, *Link & Otto, Ic. Pl. Sel.* 7. t. 2.—Afr. austr.
Burchellii, *Benth. in DC. Prod.* x. 258.—Afr. austr.
capensis, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 16.—Afr. austr.
cuneata, *E. Mey. ex Benth. l. c.* 17.—Afr. austr.
denticulata, *Benth. l. c.* 18.—Afr. austr.
diffusa, *Benth. l. c.* 18.—Afr. austr.
elongata, *Benth. l. c.*—Afr. austr.
heterandra, *Benth. l. c.*—Afr. austr.
hetrandra, *Benth. l. c.*—Afr. austr.
integerrima, *E. Mey. ex Benth. l. c.* 18.—Afr. austr.
macrophylla, *Benth. l. c.* 17.—Afr. austr.
macrophylla, *Spreng. Syst.* ii. 800.—Afr. austr.
montana, Spreng. l. c. = Hemimeris montana.
nemophiloides, *Benth. in DC. Prod.* x. 257.—Afr. austr.
pachyceras, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 16.—Afr. austr.
parviflora, *Benth. l. c.* 17.—Afr. austr.
racemulosa, *Benth. l. c.*—Afr. austr.
rigescens, *E. Mey. ex Benth. l. c.* 18.—Afr. austr.
runcinata, *E. Mey. ex Benth. l. c.* 16.—Afr. austr.
sacculata, *Benth. in DC. Prod.* x. 258.—Afr. austr.
tanyceras, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 17.—Afr. austr.
Thunbergiana, *Spreng. Syst.* ii. 800.—Afr. austr.
unilabiata, *Benth. in DC. Prod.* x. 257.—Afr. austr.

DIASIA, DC. in Bull. Soc. Philom. iii. (1808) 151 = **Molasphaerula**, Ker-Gawl. (Irid.).
graminifolia, DC. l. c.; et in Red. Lil. t. 163 = M. gramines.
intermedia, Heynh. Nom. i. 261 = M. gramines.
iridifolia, DC. in Bull. Soc. Philom. iii. (1808) 151; et in Red. Lil. t. 54 = M. gramines.
iridifolia, Eckl. Top. Verz. 31 = Lapeyrousia Fabricii.
parviflora, Steud. Nom. ed. II. i. 501 = M. parviflora.

DIASPANANTHUS, Miq. in Ann. Mus. Lugd. Bat. ii. 186 (1865) = **Ainsliaea**, DC. (Compos.).
palmatus, Miq. l. c. = A. uniflora.

DIASPASIS, R. Br. Prod. 588 (1810). *GOODENOVIEAE*, Benth. & Hook. f. ii. 540.
filifolia, *R. Br. l. c.* 587.—Austral.

DIASPERUS, Linn. Syst. ed. I. (1735) = **Phyllanthus**, *Linn.* (Euphorb.).

DIASTATEA, Scheidw. in Otto & Dietr. Allg. Gartenz. ix. (1841) 396. *CAMPANULACEAE.*
virgata, *Scheidw. l. c.*—Mexic.

DIASTELLA, Salisb. in Knight, Prot. 61 (1809) = **Leucospermum**, R. Br. (1810) (Proteac.).
bruniformis, Salisb. l. c. 62 = L. ruberum.
ericaefolia, Salisb. l. c. 64 = Mimetes purpurea.
humifusa, Salisb. l. c. 63.—Afr. austr.
myrtifolia, Salisb. l. c. = Mimetes myrtifolia.
parilis, Salisb. l. c. 62 = L. parile.
serpyllifolia, Salisb. l. c.—Afr. austr.
vaccinifolia, Salisb. l. c. 63 = Mimetes vacciniifolia.

DIASTEMA, Benth. Bot. Voy. Sulph. 122 (1844). *GESNERACEAE*, Benth. & Hook. f. ii. 1000.
DIASTEMATION, C. Muell. in Walp. Ann. v. 426 (1858).
DIASTEMELLA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 99.
—Costa Rica.
caracasanum, *Klotzsch & Hanst. in Ann. Sc. Nat. Sér.* IV. vi. (1856) 371.—Venezuela.
comiferum, *Benth. ex Walp. Rep.* vi. 403.—Ecuador.
cristatum, *Hanst. in Linnaea*, xxxiv. (1865–66) 488.—Costa Rica.
discolor, *Benth. ex C. Muell. in Walp. Ann.* v. 425.—Peruv.
divaricatum, *Benth. ex Walp. Rep.* vi. 403.—Peruv.
ericoides, *Benth. ex Walp. l. c.*—Venezuela.
glaziovifolium, Regel, Cat. Hort. Taur. (1868) 84 = Gloxinia glabrata.
gracilis, Regel, in Flora, xxxii. (1849) 180 = Diopyta candida.
inclusum, *Benth. Pl. Hartw.* 235.—Peruv.
longiflorum, *Benth. l. c.* 234.—N. Granat.

DIASTEMA, —
maculatum, *Benth. ex Walp. Rep.* vi. 403.—Peruv.
molle, *Benth. ex Walp. l. c.*—Peruv.
ochroleucum, *Hook. Bot. Mag.* t. 4354.—N. Granat.
pictum, *Benth. ex Walp. Ann.* i. 472.—Guatemala.
pictum, Regel, in Flora, xxxi. (1848) 250 = Isoloma pictum.
quinquevulnerum, *Planch. & Linden, in Fl. des Serres*, Sér. I. viii. (1852–53) 237.—N. Granat.
racemiferum, *Benth. Bot. Voy. Sulph.* 133.—Ecuador.
scabrum, *Benth. ex Walp. Rep.* vi. 403.—Peruv.
tenerrimum, *Benth. ex Walp. l. c.*—Peruv.
urticaefolium, *Benth. ex Walp. l. c.*—Peruv.

DIASTEMANTHE, Steud. Syn. Pl. Gram. 360 (1855) = **Stenotaphrum**, Trin. (Gramin.).
platystachya, Steud. l. c. = S. americanum.

DIASTEMATION, C. Muell. in Walp. Ann. v. 426 (1858) = seq.

DIASTEMELLA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 99 = **Diastema**, Benth. (Gesnerac.).
bracteosa, Oerst. l. c. 100 = Diastema bracteosum.

DIASTEMMA, Lindl. Veg. Kingd. 672 (1847) = **Diastema**, Benth. (Gesnerac.).

DIASTROPHIS, Fisch. & Mey. Ind Sem. Hort. Petrop. ii. 35 (1835) = **Asthionema**, R. Br. (Crucifer.).
cristata, Fisch. & Mey. l. c. 36 = A. Diastrophis.

DIATENOPTERYX, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 284. *SAPINDACEAE.*
sorbifolia, *Radlk. l. c.* 285.—Bras.

DIATOMA, Lour. Fl. Cochinch. 295 (1790) = **Carallia**, Roxb. (Rhizophor.).
brachiata, Lour. l. c. 296 = C. integerrima.

DIATONTA, Walp. Rep. ii. 614 (1843) = DIADONTA, Nutt. = **Coreopsis**, Linn. (Compos.).

DIATOSPERMA, C. Muell. in Walp. Ann. v. 211 (1858) = DIOTOSPERMA, A. Gray = **Ceratogyne**, Turcz. (Compos.).

DIATREMA, Rafin. Fl. Tellur. iv. 72 (1836) = **Ipomoea**, Linn. (Convolv.).
aiba, Rafin. Fl. l. c. = I. alba.
muricata, Rafin. l. c. = I. muricata.
purpurea, Rafin. l. c. = I. purpurea.

DIATROPA, Dum. Fl. Belg. 76 (1827) = **Bupleurum**, Linn. (Umbell.).
rotundifolia, Dum. l. c. = B. rotundifolium.

DIAXULON, Rafin. Sylva Tellur. 24 (1838) = **Cytisus**, Linn. (Legum.).
argentinum, Rafin. l. c. 25 = Argyrolobium Linnæanum.
prolifer, Rafin. l. c. = C. proliferus.

DIAZEUXIS, D. Don, in Trans. Linn. Soc. xvi. (1830) 251 = **Lycoceeris**, Cass. (Compos.).
intermedia, DC. Prod. vii. 22 = L. triplinervia ?
latifolia, D. Don, in Trans. Linn. Soc. xvi.' (1830) 302 = L. latifolia.
Mutisiana, D. Don, l. c. 253, in obs. = L. mexicana.
serrata, D. Don, l. c. 254, 302 = Vernonia arctioides.
trinervis, D. Don, l. c. 256.—Peruv.

DIAZIA, Phil. Fl. Atac. 22. t. 1. E. (1860) = **Calandrinia**, H. B. & K. (Portulac.).
portulacoides, Phil. l. c.—Chili.

DIBERARA, Baill. in Bull. Soc. Linn. Par. i. (1881) 279 = **Nebelia**, Nack. (Brunlac.).

DIBOTHROSPERMUM, Kraf, in Flora, xxix. (1846) 398 = **Malacaria**, Tourn. (Compos.).
agreste, Kraf, l. c. = M. inodora.
pusillum, Kraf, l. c. = M. inodora.

DIBRACHIA, Steud. Nom. ed. II. i. 502 (1840) = DIBRACHYA, Eckl. & Zeyh. = **Pelargonium**, L'Hérit. (Geraniac.).

DIBRACHION, Regel, Ind. Sem. Hort. Petrop. (1865) 51; Gartenfl. xv. (1866) 100. t. 504 = **Homalanthus**, A. Juss. (Euphorb.).
pelitatum, Regel, ll. cc. = H. giganteus.

DIBRACHION, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 138 = **Diplotropis**, Benth. (Legum.).
brasiliense, Tul. l. c. 139 = Dipl. brasiliensis.
guianense, Tul. l. c.—Guiana.
riparium, Spruce, ex Benth. in Mart. Fl. Bras. xv. i. 321 = Dipl. Martiusii.

DIBRACHIUM, Walp. Rep. v. 550 (1845–46) = DIBRACHION, Regel = **Homalanthus**, A. Juss. (Euphorb.).

DIBRACHYA, Eckl. & Zeyh. Enum. 74 (1835) = **Pelargonium**, L'Hérit. (Geraniac.).
cirpratu, Eckl. & Zeyh. l. c. 75 = P. peltatum.
peltata, Eckl. & Zeyh. l. c. 74 = P. peltatum.
scutata, Eckl. & Zeyh. l. c. = P. scutatum.

DICAELOSPERMUM, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 630 (1879). *CUCURBITACEAE.*
Kitchlet, *C. B. Clarke, l. c.*—Ind. or.

DICALIX, Lour. Fl. Cochinch. 663 (1790) = **Symplocos**, Linn. (Styrac.).
acuminatus, Blume, Bijdr. 1119 = S. acuminata.
alsuminosus, Blume, l. c. 1117 (excl. syn.) = S. spicata.
ciliatus, Blume, l. c. 1119 = S. ciliatus, *Miq. cochinchinensis*, Lour. Fl. Cochinch. ii. 663.—Cochinch.
contatus, Blume, Bijdr. 1117 = S. contata.
floridus, Regel & Koern. Ind. Sem. Hort. Petrop. (1858) 45.—Hab. ?
javanicus, Blume, Bijdr. 1117 = S. ferruginea.
odoratissimus, Blume, l. c. 1116 = S. odoratissima.
salaccensis, Blume, l. c. 1118 = S. salaccensis.
sessilifolius, Blume, l. c. = S. subsessilis.
spicatus, Blume, l. c. = S. caryophylloides.
sylvaticus, G. Don, Gen. Syst. ii. 651 = *cochinchinensis. tinctorius*, Blume, Bijdr. 1116 = S. fasciculata.

DICALYMMA, Lem. Illustr. Hortic. ii. (1855) Misc. 87 = **Podachsenium**, Benth. (Compos.).
fragrans, Lem. l. c. = P. paniculatum.

DICALYX, Pour. in Dict. Sc. Nat. xlii. 172 (1819) = DICALIX, Lour. = **Symplocos**, Linn. (Styrac.).

DICARDIOTIS, Rafin. Fl. Tellur. iii. 31 (1836) = **Gentiana**, Tourn. (G. auriculata, *Pall.*).

DICARPAEA, Presl, Symb. Bot. i. 37. t. 26 (1830) = **Linum**, Linn. (Flocod.).
linifolia, Presl, l. c. 38. t. 26 = L. linifolium.

DICARPIDIUM, F. Muell. in Hook. Kew Journ. ix. (1857) 302. *STERCULIACEAE*, Benth. & Hook. f. i. 234.
monoicum, *F. Muell. l. c.*—Austral.

DICARYUM, Willd. ex Roem. & Schult. Syst. iv. 802 (1814). *LOGANIACEAE!*
serrulatum, *Willd. ex Roem. & Schult. l. c.*—N. Granat.
subdentatum, *Willd. ex Roem. & Schult. l. c.*—Ecuador.

DICELLA, Griseb. in Linnaea, xiii. (1839) 250. *MALPIGHIACEAE*, Benth. & Hook. f. i. 362.
bracteosa, *Griseb. l. c.*—Bras.
holosericea, *A. Juss. in Ann. Sc. Nat. Sér. II.* xiii. (1840) 323.—Bras.
lanciflolia, A. Juss. l. c. = bracteosa.
macroptera, *A. Juss. l. c.*—Bras.
ovatifolia, A. Juss. l. c. = bracteosa.

DICELLANDRA, Hook. f. in Benth. & Hook. f. Gen. i. 757 (1867). *MELASTOMACEAE*, Benth. & Hook. f. i. c.
Barteri, *Hook. f. l. c.*—Afr. trop.
setosa, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 459.—Afr. trop.

DICELLOSTYLIS, Benth. in Journ. Linn. Soc. vi. (1863) 123; et Benth. & Hook. f. Gen. i. 207 (1862). *MALVACEAE*, Benth. & Hook. f. i. 207.
axillaris, *Benth. in Journ. Linn. Soc.* vi. (1862) 122.—Ceylon.
jujubifolia, Benth. & Hook. f. Gen. i. 207 = Kydia jujubifolia.

DICENTRA, Bernh. in Linnaea, viii. (1833) 457, 468; xii. (1838) 668. *PAPAVERACEAE*, Benth. & Hook. f. i. 55.
BICUCULLARIA, Juss. ex Steud. Nom. ed. II. i. 202 (1840).
BIKUKULLA, Adans. Fam. ii. (2) 1763, nomen prius.
CAPNORCHIS, Planch. in Fl. des Serres, viii. (1852) 193.
CUCULLARIA, Rafin. in Med. Repos. N. York, v. (1808) 352.
CUCULLARIA, Endl. Gen. 859 (1839).
DACTYLICAPNOS, Wall. Tent. Fl. Nep. 51. t. 39 (1828).
DACTYLISCAPNOS, Spreng. Gen. ii. 568 (1831).
DICLYTRA, Rafin. Cat. 18 (1824).
DICLYTRA, Borck. in Roem. Arch. i. ii. 46 (1797).
DIELYTRA, Dalt. & Schlecht. in Linnaea, i. (1826) 556.
EUCAPNOS, Bernh. in Linnaea, viii. (1833) 468.
LAMPROCAPNOS, Endl. Gen. Suppl. v. 82 (1850).
MACROCAPNOS, Royle, ex Lindl. Introd. Nat. Syst. ed. II. 439 (1836).

canadensis, *Walp. Rep.* i. 118; *Torr. Fl. N. York*, i. 46. —Am. bor.
chrysantha, *Walp. l. c.*—Calif.
Cucullaria, *Bernh. in Linnaea*, viii. (1833) 468.—Am. bor.
eximia, *Torr. Fl. N. York*, i. 46.—Am. bor.
formosa, *Walp. Rep.* i. 118.—Am. bor. occ.
lachenalisflora, *Ledeb. Fl. Ross.* i. 97.—Sibir.; Am. bor. occ.
ochroleuca, *Engelm. in Coult. Bot. Gaz.* vi. (1881) 223.—Calif.
pauciflora, *S. Wats. in Bot. Calif.* ii. 429.—Calif.
pusilla, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1843) 172.—Japon.
Roylei, *Hook. f. & Thoms. Fl. Ind.* i. 273.—Reg. Himal.
saccata, Walp. Rep. i. 118 = formosa.
scandens, *Walp. l. c.*—Reg. Himal.
spectabilis, *Lem. in Fl. des Serres*, Sér. I. iii. (1847) t. 258.—Japon.
tenuifolia, Lodd. Fl. Ross. i. 97 = lachenalisflora.
thalictrifolia, *Hook. f. & Thoms. Fl. Ind.* i. 273.—Reg. Himal.
torulosa, *Hook. f. & Thoms. l. c.* 272.—Reg. Himal.
uniflora, *Kellogg, in Proc. Calif. Acad.* iv. (1871) 141. —Am. bor. occ.

DICENTRANTHERA, T. Anders. in Journ. Linn. Soc. vii. (1864) 82 = **Asystasia**, Blume (Acanth.).
macrophylla, T. Anders. l. c.—Afr. trop.

DICERA, Forst. Char. Gen. 79. t. 40 (1776) = **Blaeocarpus**, Burm. (Tiliac.).
Craspedum, J. F. Gmel. ex DC. Prod. i. 590 = E. tectorius.
dentata, Forst. Char. Gen. 80 = E. dentatus.
serrata, Forst. l. c. = E. dentatus.

DICERA, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 67, 74 (1858) [1856] = **Gironniera**, Gaudich. (Urticac.).
lanceolata, Zipp. l. c. 67 = Celtis paniculata.
rhamnifolia, Zipp. l. c. = G. rhamnifolia.

DICERANDRA, Benth. in Bot. Reg. sub t. 1800 (1830) = **Cerasthaera**, Ell. (Labiat.).
densiflora, Benth. in DC. Prod. xii. 348 = C. densiflora.
linearifolia, Benth. in Bot. Reg. sub t. 1800 = C. linearifolia.
linearis, Benth. Lab. Gen. et Sp. 418 = C. linearifolia.

DICERATELLA, Boiss. Diagn. Ser. I. v. 80 (1844). *CRUCIFERAE*, Benth. & Hook. f. i. 71.
DICERATUM, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 61.
canescens, *Boiss. in Ann. Sc. Nat. Sér.* II. xvii. (1842) 61.—Persia.
floccosum, *Boiss. l. c.*—Persia.
incana, *Bailf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 500.—Ins. Socotr.

DICERATIUM, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 61 = **Diceratella**, Boiss. (Crucif.).
canescens, Boiss. l. c. 62 = Diceratella canescens.
floccosum, Boiss. l. c. 61 = Diceratella floccosum.

DICERATIUM, Lag. Elench. Madr. 20 (1815); Nov. Gen. et Sp. 20 (1816) = **Rotocoras**, R. Br. (Crucif.).
prostratum, Lag. Nov. Gen. et Sp. 20 = N. canariense.

DICERMA, DC. Mém. Lég. 326 (1825); et Prod. ii. 330 = **Desmodium**, Desv. (Legum.).
acanthocladum, F. Muell. Fragm. ii. 122 = Desm. acanthocladum.
biarticulatum, DC. Prod. ii. 339 = Desm. biarticulatum.
elegans, DC. l. c. = Desm. elegans.
pulchellum, DC. l. c. = Desm. pulchellum.
repens, R. Grah. in Wall. Cat. n. 5740 = Desm. heterophyllum.
vestitum, Wall. l. c. n. 5739 = Desm. vestitum.

DICEROCARYUM, Boj. in Ann. Sc. Nat. Sér. II. iv. (1835) 268. t. 10 = **Pretrea**, J. Gay (Pedalin.).
sinuatum, Boj. l. c. 269. t. 10 = P. zanquebarica.

DICEROLEPIS, Blume, Mus. Bot. Lugd. Bat. i. 145. t. 24 (1850) = **Gymnanthera**, R. Br. (Asclep.).
hypoleuca, Miq. Fl. Ind. Bat. ii. 466.—Malaya.
paludosa, Blume, Mus. Bot. Lugd. Bat. i. 146.—Java.
pedunculata, Miq. Fl. Ind. Bat. ii. 466 = G. pedunculata.

DICEROS, Blume, Bijdr. 752 (1826) = **Vandellia**, Linn. (Scrophular.).
caespitosus, Blume, l. c. 753 = V. caespitosa.
*glandulifer*us, Blume, l. c. 752 = V. glandulifera.
latifolius, Blume, l. c. = V. latifolia.
montanus, Blume, l. c. = V. montana.

DICEROS, Lour. Fl. Cochinch. 881 (1790) = **Limnophila**, R. Br. (Scrophular.).
cochinchinensis, Lour. l. c.—Cochinch.
stoloniferus, Blanco, Fl. Filip. ed. II. 349 = L. conferta.

DICEROS, Pers. Syn. ii. 164 (1807) = **Artanema**, D. Don (1834) (Scrophul.).
angustifolius, Benth. in Wall. Cat. n. 3885 = A. angustifolium.
longifolius, Pers. Syn. ii. 164 = A. sesamoides.

DICEROSTYLIS, Blume, Orch. Archip. Ind. i. 116 (1858) = **Hylophila**, Lindl. (Orchid.).
lancolata, Blume, l. c. = H. lancolata.

DICERSOS, Lour. Fl. Cochinch. ii. 861 (1790). sphalm. = DICEROS, Lour. = **Limnophila**, R. Br. (Scrophular.).

DICHAEA, Lindl. Gen. et Sp. Orchid. 206 (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 556.
EPITHECIA, Knowles & Westc. Fl. Cab. ii. (1838) 167. t. 87.
FERNANDEZIA, Ruiz & Pav. Prod. 123 (1794).
brachyphylla, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 228.—Bras.
brachypoda, *Reichb. f. Beitr. Orch. Centr. Am.* 78.—Costa Rica.
bryophila, *Reichb. f. Otia Bot. Hamb.* 90.—Bras.
calycolata, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 4.—Peruv.
coriacea, *Rodrig. Orch. Nov.* ii. 181.—Bras.
echinocarpa, *Lindl. Gen. et Sp. Orch.* 208.—Mexic.
glauca, *Lindl. l. c.* 209.—Jamaic.
gorgonensis, *Reichb. f. in Linnaea*, xli. (1877) 75.—Ins. Gorgona.
graminea, Griseb. Fl. Brit. W. Ind. 625 = seq.
graminoides, *Lindl. Gen. et Sp. Orch.* 209.—Ind. occ.
Hitrio, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 230.—Venezuela.
hystricina, *Reichb. f. in Flora*, xlvii. (1865) 279.—Ind. occ.
Kegelii, *Reichb. f. in Linnaea*, xli. (1877) 190.—Cuba.
Lagotis, *Reichb. f. l. c.* 118.—Venezuela.
latifolia, *Lindl. Gen. et Sp. Orch.* 208.—Ins. S. Vincent.
latifolia, *Rodrig. Orch. Nov.* i. 91.—Bras.
laxa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 4.—Peruv.
Liebmannii, *Reichb. f. in Gard. Chron.* (1879) i. 676, xoxen.—Mexic.
maculata, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 3.—Peruv.

DICHAEA:—
Moritzii, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 228.—Venezuela.
Moseni, *Reichb. f. Otia Bot. Hamb.* 90.—Am. trop.
muricata, *Lindl. Gen. et Sp. Orch.* 209.—Jamaic.
obtrusa, *Lindl. Bot. Reg.* (1839) *Misc.* 54.—Guiana.
Osteadii, *Reichb. f. in Bonplandia*, iii. (1855) 219.—Nicaragua.
panamensis, *Lindl. Gen. et Sp. Orch.* 209.—Panama.
picta, *Reichb. f. in Saund. Refug. Bot.* t. 54.—Ins. Trinit.
pumila, *Reichb. f. Otia Bot. Hamb.* 91.—Bras.
Spitzgerberi, *Reichb. f. in Nederl. Kruidk. Arch.* iv (1859) 227.—Guiana.
squarrosa, *Lindl. in Ann. Nat. Hist.* iv. (1840) 384.—Mexic.
tigrina, *Reichb. f. in Schiller, Cat. Orch.* 27; *ex Regel, Ind. Sem. Hort. Petrop.* (1858) 80.—Am. austr.
trichocarpa, *Lindl. Gen. et Sp. Orch.* 209.—Jamaic.
Trulla, *Reichb. f. Beitr. Orch. Centr. Am.* 104.—Nicaragua.
Weigeltii, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 228.—Guiana.

DICHAELIA, Harv. Gen. S. Afr. Fl. ed. II. 241 (1868) = **Brachystelma**, R. Br. (Asclep.).
Gerrardi, Harv. l. c. = B. Gerrardi.

DICHAETA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 383 = **Baeria**, Fisch. & Mey. (Compos.).
Fremontii, Torr. ex A. Gray, Pl. Fendl. 102 = B. Fremontii.
tenella, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 383 = B. uliginosa.
uliginosa, Nutt. l. c. = B. uliginosa.

DICHAETA, Sch. Bip. in Flora, xxxiii. (1850) 419 = **Schastotellia**, Sch. Bip. (Compos.).

DICHAETANDRA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 81 = **Ernestia**, DC. (Melastom.).
Goudotii, Naud. l. c. 82 = E. Goudotii.

DICHAETANTHERA, Endl. Gen. 1215 (1840). *MELASTOMACEAE*, Benth. & Hook. f. i. 747.
aculeolata, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 61.—Madag.
arborea, *Baker, in Journ. Linn. Soc. xx.* (1883) 147. —Madag.
articulata, *Endl. Gen.* 1215; *et in Walp. Rep.* ii. 187 —Madag.
cordifolia, *Baker, in Journ. Linn. Soc. xx.* (1883) 146. —Madag.
heteromorpha, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 61.—Madag.
madagascariensis, *Triana, l. c.*—Madag.
oblongifolia, *Baker, in Journ. Linn. Soc. xx.* (1883) 147.—Madag.
Rutenbergiana, *Baill. ex Vathe, in Bremen Abh.* ix. (1885) 116.—Madag.

DICHAETARIA, Neck. in Steud. Syn. Pl. Gram. 145 (1855) = **Gymnopogon**, Beauv. (Gramin.).
Wightii, Nees, l. c. = G. rigidus, Thw.

DICHAETOPHORA, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) (Pl. Fendl.) 73 = **Boltonia**, L'Hérit. (Compos.).
campestris, A. Gray, l. c. = B. campestris.

DICHANTHIUM, Willem. in Usteri, Ann. Bot. xviii. (1796) 11 = **Andropogon**, Linn. (Gramin.).
sodosum, Willem. l. c. = A. caricosus.

DICHAPETALUM, Thou. Gen. Nov. Madag. 23 (1806) = **Chailletia**, DC.
Baillonii, Pierre, Fl. For. Cochinch. t. 47.—Cochinch.
Heliforianum, Pierre, l. c. 48 = C. Heliferiana.
Heudelotii, Baill. Hist. Pl. v. 140 = C. Heudelotii.
hispidum, Baill. l. c. 141 = C. hispida.
madagascariensis, Poir. in Dict. Sc. Nat. xlii. 178 = C. Dichapetalum.
pedunculatum, Baill. Hist. Pl. v. 140 = C. pedunculata.
Thouarsianum, Roem. & Schult. Syst. v. 343 = C. Dichapetalum.
toxicarium, Baill. Hist. Pl. v. 139 = C. toxicaria.

DICHELACHNE, Endl. Prod. Fl. Ins. Norf. 20 (1833). *GRAMINEAE*, Benth. & Hook. f. iii. 1163.
comata, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. (1842) 6 = crinita.
crinita, *Hook. f. Fl. N. Zel.* i. 293.—Austral.; N. Zel.
Drummondiana, Steud. Syn. Pl. Gram. 120 = Deyeuxia Drummondiana.
Forsteriana, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. (1842) 6 = crinita.
Hookeriana, Trin. & Rupr. l. c. 8 = crinita.
longiseta, Trin. & Rupr. l. c. 5 = crinita.
montana, Endl. Prod. Fl. Ins. Norf. 24 = sciurea.
procera, Trin. & Rupr. in Mém. Acad. Petersb. Sér. VI. v. Sc. Nat. (1842) 6 = crinita ?
rigida, Trin. & Rupr. l. c. 5 = sciurea ?
sciurea, *Hook. f. Fl. N. Zel.* 294.—Austral.; N. Zel.
setacea, Nees, in Lehm. Pl. Preiss. ii. 98 = Stipa teretifolia.
Sieberiana, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. (1842) 8 = sciurea.
stipoides, Hook. f. Fl. N. Zel. 294. t. 66 = Stipa teretifolia.
vulgaris, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. (1842) 8 = sciurea.

DICHELACTINA, Hance, in Walp. Ann. iii. 375 (1853) = **Phyllanthus**, Linn. (Euphorb.).
nodicaulis, Hance, l. c. 376 = P. Emblica.

DICHELOSTEMMA, Kunth, Enum. Pl. iv. 469 (1843) **Brodiaea**, Sm. (Liliac.).
californica, Wood, in Proc. Acad. Nat. Sc. Philad. 1868) 173 = Stropholirion californicum.
capitata, Benth. ex Wood, l. c. = Milla capitata.
congestum, Kunth, Enum. Pl. iv. 470 = B. congesta.

DICHELOSTYLIS, Endl. Gen. 118, in syn. (1836) = DICHOSTYLIS, Beauv. = **Scirpus**, Linn. (Cyperac.).

DICHERANTHUS, Webb, in Ann. Sc. Nat. Sér. III. v. (1846) 27. t 2. *ILLECEBRACEAE*, Benth. & Hook. f. iii. 18.
plocamoides, *Webb*, *l. c.* 29. t. 2.—Ins. Canar.

DICHILANTHE, Thw. in Hook. Kew Journ. viii. (1856) 269, 276. t. 8. A. *RUBIACEAE*, Benth. & Hook. f. ii. 103.
borneensis, *Baill. Hist. Pl.* vii. 428.—Borneo.
zeylanica, *Thw. in Hook. Kew Journ.* viii. (1856) 270. —Zeylan.

DICHILOS, Spreng. Syst. iv. Cur. Post. 263 (1827) = seq.

DICHILUS, DC. Mém. Lég. 201. t. 35 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 479.
CALYCOTOME, E. Mey. Comm. Pl. Afr. Austr. 118 (1836).
MELINOSPERMUM, Walp. in Linnaea, xiii. (1839) 527.
candicans, E. Mey. in Linnaea, vii. (1832) 154 = Melolobium candicans.
ciliatus, Spreng. f. Syst. Tent. Suppl. 20 = Argyrolobium Andrewsianum.
crassifolius, E. Mey. in Linnaea, vii. (1832) 154 = Argyrolobium crassifolium.
floribundus, Sond. in Linnaea, xxiii. (1850) 28 = Lotononis dichiloides.
gracilis, *Eckl. & Zeyh. Enum.* 188.—Afr. austr.
hypotrichum, *Spreng. Syst.* iv. *Cur. Post.* 278.—Afr. austr.
lanceolatus, E. Mey. in Linnaea, vii. (1832) 154 = Argyrolobium lanceolatum.
lebeckioides, *DC. Mém. Lég.* 202. t. 35.—Afr. austr.
microphyllus, E. Mey. in Linnaea, vii. (1832) 155 = Melolobium microphyllum.
obovatus, E. Mey. l. c. 154 = Argyrolobium obovatum.
patens, E. Mey. Comm. Pl. Afr. Austr. 36 = gracilis.
pusillus, Benth. in Hook. Lond. Journ. Bot. iii. (1841) 554 = gracilis.
sericeus, E. Mey. in Linnaea, vii. (1832) 154 = Argyrolobium sericeum.
sericeus, *Spreng. f. Syst. Tent. Suppl.* 20.—Afr. austr.
spicatus, E. Mey. in Linnaea, vii. (1832) 154 = Melolobium spicatum.
strictus, *E. Mey. Comm. Pl. Afr. Austr.* 36.—Afr. austr.

DICHISMUS, Rafin. in Journ. Phys. lxxxix. (1819) 106 = **Scirpus**, Linn. (Cyperac.).

DICHODON, Bartl. ex Reichb. Nom. 205 (1841) = **Cerastium**, Linn. (Caryophyll.).
anomalum, Reichb. Ic. Fl. Germ. v. 84 = C. *cerastioides*, Reichb. l. c. f. 4915 = C. trigynum.

DICHOESPERMUM, Wight, Ic. vi. 81. t. 2076 (1853) = **Ancistrus**, R. Br. (Commelin.).
aurantiacum, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 207 = A. pauciflorum.
Blumei, Hassk. Commel. Ind. 41 = A. Hamiltonianus.
giganteum, Hassk. l. c. 42 = A. Thomsoni.
juncoides, Wight, Ic. t. 2078 = A. paniculatum.
Koenigii, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 215 = A. Koenigii.
lanceolatum, Wight, Ic. t. 2078 = A. Koenigii.
ochraceum, Hassk. Commel. Ind. 41 = A. ochraceum.
paniculatum, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 215 = A. paniculatum.
repens, C. B. Clarke, Commel. et Cyrt. Beng. t. 28 = A. Hamiltonianum.
repens, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 207 = A. spiratum.
repens, Wight, Ic. t. 2078 = A. ochraceum.
semteres, Hassk. Commel. Ind. 41 = A. paniculatum.

DICHOGLOTTIS, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. (1835) 25 = **Gypsophila**, Linn. (Caryophyll.).
alsinoides, Walp. Ann. i. 90 = G. alsinoides.
austriaca, Schlecht. in Linnaea, xx. (1847) 631 = G. tubulosa.
bellidifolia, Walp. Rep. v. 78 = G. bellidifolia.
floribunda, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 105 = G. floribunda.
frankenioides, Walp. Rep. v. 78 = G. frankenioides.
hispida, Walp. l. c. = G. hispida.
linearifolia, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. (1835) 26 = G. linearifolia.
linifolia, Fisch. & Mey. ex Pritz. Ic. Ind. 862, sphaim. = prsec. ?
muralis, Jaub. & Spach, Illustr. i. 12 = G. muralis.
Scowitzii, Walp. Rep. v. 78 = G. Szowitzii.
spathulatifolia, Fisch. & Mey. ex Ledeb. Fl. Ross. i. 290 = G. spathulatifolia.
trichopoda, Walp. Rep. v. 78 = G. linearifolia.
tubulosa, Jaub. & Spach, Illustr. i. 14 = G. tubulosa.

DICHONDRA, Forst. Char. Gen. 89. t. 20 (1776). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 879.
DEMIDOFIA, J. F. Gmel. Syst. ii. 458 (1791).
STERIPHA, Banks, ex Gaertn. Fruct. ii. 81. t. 94 (1791).
argentea, *Willd. Hort. Berol.* 297. t. 81.—Am. calid.
brevifolia, *Y. Buch. in Trans. N. Z. Inst.* iii. (1871) 208.—N. Zel.
carolinensis, Michx. Fl. Bor. Am. i. 136 = repens.
caroliniana, Willd. ex DC. Prod. ix. 451 = prsec.
cymbalariaefolia, Willd. ex Roem. & Schult. Syst. vi. 209 = Sibthorpia pichinchensis.
macrocalyx, Meisn. in Mart. Fl. Bras. vii. 358 = repens.
parvifolia, *Meism. l. c.* 360.—Bras.
perennis, Pohr. Enzyc. Suppl. ii. 470 = repens.
repanda, Willd. ex Roem. & Schult. Syst. vi. 209 = Sibthorpia pichinchensis.
repens, *Forst. Char. Gen.* 89. t. 20.—Reg. trop. et subtrop.
rotundifolia, Link, Enum. Hort. Berol. i. 249 (= Sibthorpiae sp.).—Peruia.
sericea, Sw. Prod. Veg. Ind. Occ. 54 = repens.
villosa, *Parodi, in Anal. Soc. Cient. Argent.* xiii. (1882) 9; *Contrib.* 32.—Reg. Argent.

DICHONE, Lawa. ex Salisb. in Trans. Hort. Soc. i. (1812) 330 = **Tritomia**, Ker-Gawl. (Irid.).
crispa, Lawa. Cat. 6; ex Salisb. l. c. = T. undulata.

DICHOPETALUM, F. Muell. in Hook. Kew Journ. vii. (1855) 378. t. 11; et in Trans. Phil. Soc. Vict. i. 101 (1855) = **Asorella**, Lam. (Umbellif.).
ranunculaceum, F. Muell. ll. cc. 102; 378 = A. dichopetala.

DICHOPOGON, Kunth, Enum. Pl. iv. 622 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 791.
SIONA, Salisb. Gen. Pl. Fragm. 67 (1866).
humilis, Kunth, l. c. 638 = strictus.
humilis, F. Muell. Fragm. vii. 67 = Arthropodium minus.
leimonophilus, F. Muell. ex Baker, in Journ. Linn. Soc. xv. (1876) 319 = humilis.
setosus, Kunth, Enum. Pl. iv. 623 = strictus.
Sieberianus, *Kunth, l. c.*—Austral.
strictus, *Baker, in Journ. Linn. Soc.* xv. (1876) 319. —Austral.
undulatus, Regel, Gartenfl. ii. (1853) 1. t. 37 = strictus.

DICHOPSIS, Thw. Enum. Pl. Zeyl. 176 (1864). *SAPOTACEAE*, Benth. & Hook. f. ii. 658.
PALAQUIUM, Blanco, Fl. Filip. ed. I. 408 (1837); ed. II. 282 (1845).
argentata, *Benth. & Hook. f. Gen.* ii. 658.—Malaya.
calophylla, Benth. & Hook. f. l. c. = Bassia calophylla.
canaliculata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 541.—Zeylan.
elliptica, *Benth. & Hook. f. Gen.* ii. 658.—Ind. or.
grandis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 540.—Zeylan.
Gutta, *Benth. & Hook. f. Gen.* ii. 658.—Malaya.
Helferi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 542.—Burma.
hexandra, *C. B. Clarke, l. c.* 543.—Malacca.
Hornei, *Hartog, ex Baker, in Journ. Linn. Soc.* xx. (1883) 367.—Ins. Fiji.
Kruetzeana, *Pierre, in Beauvis, Contr. Gutta-Percha*, 60. t. 1.—Cambodia ?
laevifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 541.—Zeylan.
lanceolata, *C. B. Clarke, in Hook. f. l. c.*—Zeylan.
lanceolata, *F. Villar, in Blanco, Fl. Philipp.* ed. III. 124.—Ins. Philipp.
latifolia, *F. Villar, in Blanco, l. c.*—Ins. Philipp.
lusoniensis, *F. Villar, in Blanco, l. c.* 126.—Ins. Philipp.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 543.—Malacca.
obovata, *C. B. Clarke, l. c.* 542.—Malacca.
oliefera, *F. Villar, in Blanco, Fl. Philipp.* ed. III. 125. —Ins. Philipp.
pauciflora, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 541.—Zeylan.
petiolaris, *Thw. Enum. Pl. Zeyl.* 176.—Zeylan.
polyanthe, *Benth. & Hook. f. Gen.* ii. 658.—Burma.
rubens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind*, iii. 542. —Malacca.
rubiginosa, *C. B. Clarke, l. c.* 541.—Zeylan.

DICHOPUS, Blume, Mus. Bot. Lugd. Bat. ii. 176 (1856) = Dendrobium, Sw. (Orchid.).
insignis, Blume, l. c. = D. insigne.

DICHORISANDRA, Mikan, Del. Fl. et Faun. Bras. t. 8 (1820). *COMMELINACEAE*, Benth. & Hook. f. iii. 852.
PETALOXIS, Rafin. Fl. Tellur. ii. 83 (1836).
STICKMANNIA, Neck. Elem. iii. 171 (1790).
affinis, Mart. in Schult. f. Syst. vii. 1185 = Aubletiana.
alba, *Seub. & Warm. in Kjoeb. Vidensk. Meddel.* (1872) 122.—Bras.
albo-marginata, *Linden, in Regel, Gartenfl.* (1868) 34. t. 569.—Bras.
Aubletiana, *Schult. f. Syst.* vii. 1187.—Am. austr.
begoniaefolia, *Kunth, Enum. Pl.* iv. 112.—Bras.
discrepans, Hassk. Commel. Ind. 59 = Gaudichaudiana.
Ehrenbergiana, Klotzsch, ex C. B. Clarke, in DC. Monog. Phan. iii. 209 = Tradescantia holosericea.
foliosa, *Kunth, Enum. Pl.* iv. 112.—Bras.
Gaudichaudiana, *Kunth, l. c.* 113.—Bras.
gracilis, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 14.—Bras.
hirtella, *Mart. in Schult. f. Syst.* vii. 1190.—Bras.
insequalis, *Presl, Rel. Haenk.* ii. 140.—Peruv.
incurva, *Mart. in Schult. f. Syst.* vii. 1184.—Bras.
intermedia, Mart. l. c. 1188 = Aubletiana.
interrupta, Mart. l. c. 1189.—Bras.
latifolia, Kunth, Enum. Pl. iv. 112 = oxypetala.

DICHORISANDRA :—
Leandri, Kunth, Enum. Pl. iv. 112 = villosula.
leucophthalmos, *Hook. Bot. Mag.* t. 4733.—Bras.
longifolia, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 378 = Tradescantia holosericea.
Luschnathiana, Kunth, Enum. Pl. iv. 111 = pubescens.
marginata, Schlecht. in Linnæa, xxvi. (1853–55) 454 = albo-marginata.
mexicana, *Presl, Rel. Haenk.* iii. 140.—Mexic.
micans, *C. B. Clarke, in DC. Monog. Phan.* iii. 281.—Bras.
mollis, Kunth, Enum. Pl. iv. 112 = villosula.
mosaica, *Linden, in Belg. Hortic.* xvii. (1867) 108.—Peruv.
musaica, Koch & Linden, ex C. Koch, Wochenschr. ix. (1866) 346 = mosaica.
ovalifolia, *Presl, Rel. Haenk.* iii. 140.—Panama.
ovata, Mart. in Schult. f. Syst. vii. 1182 = Aublétiana.
ovata, Paxt. Mag. Bot. xv. (1849) t. juxta 5 = Gaudichaudiana.
oxypetala, *Hook. Bot. Mag.* t. 3731.—Bras.
penduliflora, *Kunth, Enum. Pl.* iv. 114.—Bras.
pericrana, *C. B. Clarke, in DC. Monog. Phan.* iii. 281.—Bras.
picta, *Lodd. Bot. Cab.* t. 1667 ; *Hook. Bot. Mag.* t. 4760.—Bras.
procera, Mart. in Schult. f. Syst. vii. 1187 = thyrsiflora.
puberula, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 13. t. 1. f. 2.—Bras.
pubescens, *Mart. in Schult. f. Syst.* vii. 1186.—Bras.
radicalis, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 11.—Bras.
rhizophyra, *Mart. in Schult. f. Syst.* vii. 1191.—Bras.
Saundersii, *Hook. f. Bot. Mag.* t. 6165.—Bras.
scandens, Carth. ex C. B. Clarke, in DC. Monog. Phan. iii. 274 = Aublétiana.
Schomburgkiana, Klotzsch, in Schomb. Fasn. et Fl. Gui. 897 = Aublétiana.
tejucensis, Mart. in Schult. f. Syst. vii. 1186 = villosula.
tenuior, *Mart. l. c.* 1183.—Bras.
thyrsiflora, *Miken, Del. Fl. et Faun. Bras.* 3. t. 3.—Bras.
undata, C. Koch & Linden, ex C. Koch, Wochenschr. ix. (1866) 346 = mosaica.
villosula, *Mart. in Schult. f. Syst.* vii. 1185.—Bras.
Warszewiciana, Planch. Hort. Donat. 80 = Tradescantia Warszewiciana.

DICHOROPETALUM, Fenzl, Pugill. Pl. Nov. Syr. 17 (1843) = Johrenia, DC. (Umbellif.).
alpinum, Fenzl, l. c. = J. alpina.

DICHOSEMA, Benth. in Enum. Pl. Hueg. 85 (1837) = Mirbellia, Sm. (Legumin.).
microphyllum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 353 = M. microphylla.
multicaule, Turcz. l. c. = M. multicaulis.
racemosum, Meisn. in Lehm. Pl. Preiss. i. 78 = Chorizema sericifolium.
spinosa, Benth. Enum. Pl. Hueg. 85 = M. spinosa.
subinermis, Meisn. in Lehm. Pl. Preiss. i. 76 = M. spinosa.

DICHOSMA, DC. ex Loud. Hort. Brit. 85 (1830) = Agathosma, Willd. (Rutac.).
bifida, Loud. l. c. = A. cerefolium.

DICHOSPERMUM, C. Muell. in Walp. Ann. vi. 158 (1861) = DICHOESPERMUM, Wight = Anellema, R. Br. (Commelin.).

DICHOSTACHYS, Kraus, in Flora, xxvii. (1844) 850, sphalm. = Dichrostachys, Wight & Arn. (Legum.).

DICHOSTYLIS, Beauv. in Lestib. Ess. Fam. Cyp. 39 (1819) = Scirpus, Linn. (Cyperac.).
fluitans, Reichb. Fl. Germ. Excurs. i. 75 = S. fluitans.
hamulosa, Nees, in Linnæa, ix. (1834) 289 = S. hamulosus.
Ludwigii, Nees, ex Kunth, Enum. Pl. ii. 189 = S. Ludwigii.
Michelianus, Nees, in Wight, Contrib. 94 = S. Michelianus.
pygmæus, Nees, in Linnæa, ix. (1834) 289 = Cyperus Pygmæus.

DICHOTOMA, Sch. Bip. Pl. Moritz. ex Benth. & Hook. f. Gen. ii. 864 (1878) = Scleria, Berg. (Cyperac.).

DICHOTOMANTHES, Kurz, in Journ. Bot. xi. (1873) 194. t. 138 (? an Rosacea sect. ?).
tristaniacarpa, *Kurz, l. c.* 195.—Yunan.

DICHOTOPHYLLUM, Moench, Meth. 345 (1794) = Cerastophyllum, Linn.
demersum, Moench, l. c. = C. demersum.

DICHROA, Lour. Fl. Cochinch. 301 (1790). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 841.
ADAMIA, Wall. Tent. Fl. Nep. 46. t. 26 (1824).
CYANITIS, Reinw. in Syll. Ratisb. ii. (1828) 10.
CYANITUS, Reinw. in Blume, Cat. Gew. Buitenz. 68 (1823).
Cyanitis, Miq. Fl. Ind. Bat. i. 1. 721 = febrifuga.
febrifuga, *Lour. Fl. Cochinch.* 301.—Reg. Himal.; Malaya; China.
latifolia, Miq. Fl. Ind. Bat. i. 1. 722 = febrifuga.
pubescens, *Miq. l. c.*—Malaya.

DICHROANTHUS, Webb & Berth. Phyt. Canar. i. 65 (1836) = Cheiranthus, Linn. (Crucif.).
cinereus, Webb & Berth. l. c. 67 = C. scoparius.
linifolius, Webb, ex Nym. Consp. 81 = Erysimum linifolium.
mutabilis, Webb & Berth. Phyt. Canar. i. 66 = C. mutabilis.
scoparius, Webb & Berth. l. c. 68 = C. scaber.

DICHROCEPHALA, L'Hérit. ex DC. in Guill. Archiv. Bot. ii. (1883) 517. *COMPOSITÆ*, Benth. & Hook. f. ii. 260.
abyssinica, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) 1. Intell. 26 = latifolia.
Benthami, *C. B. Clarke, Comp. Ind.* 86.—Reg. Himal.
bicolor, Schlecht. in Linnæa, xxv. (1852) 209 = latifolia.
capensis, DC. in Guill. Archiv. Bot. ii. 518 ; Prod. v. 271 = latifolia.
chrysanthemifolia, *DC. ll. cc.* 518, 372.—Ind. or.; Java ; Afric.
erecta, L'Hérit. ex DC. Prod. v. 372 = latifolia.
gracilis, DC. l. c. 371 = latifolia.
grangeæfolia, DC. l. c. 372 = chrysanthemifolia.
Hamiltoni, *Hook. f. Fl. Brit. Ind.* iii. 246.—Ind. or.
latifolia, *DC. in Guill. Archiv. Bot.* ii. 518 ; *et Prod.* v. 372.—As. et Afr. trop. et subtrop.
lyrata, *DC. Prod.* v. 372.—Madag.
macrocephala, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 145 = chrysanthemifolia.
minima, Boj. Hort. Maurit. 179 = seq.
minuta, L'Hérit. ex DC. Prod. vi. 189 = Centipeda orbicularis.
nilagirensis, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 245 = latifolia.
oblonga, Hook. f. in Journ. Linn. Soc. vi. (1862) 12 = chrysanthemifolia.
oblongata, Benth. & Hook. f. Gen. ii. 260 = chrysanthemifolia.
paniculata, Miq. Fl. Ind. Bat. ii. 88 = latifolia.
Schmidii, Wight, Ic. t. 1610 = Centipeda orbicularis.
sonchifolia, DC. in Guill. Archiv. Bot ii. 518 ; et Prod. v. 373 = latifolia.
tanacetoides, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 122 = chrysanthemifolia.

DICHROLEPIS, Welw. Apont. 542, nomen (1858). *ERICACEALEÆ*.
pusilla, *Welw. l. c.*—Afr. trop.

DICHROMA, Carr. in Anal. Clinc. Nat. iii. (1801) 339 ; et Ic. vi. 59. t. 582 (1801) = Ouviria, Comm. (Scrophularin.).
coccineum, Carr. ll. cc. 338, 59 = O. coccinea.

DICHROMA, Pers. Syn. i. 87 (1805) = Dichromena, Michx. (Cyperac.).
corymbosum, Muhl. Desc. Gram. 14 (= *Isolepis stenophylla*).—Am. bor.
ciliatum, Pers. Syn. i. 88 = Dichromena ciliata.
glabra, Pers. l. c. 87 = Dichromena leucocephala.
leucocephalum, Pers. l. c. = Dichromena leucocephala.
rupestre, Pers. l. c. = Dichromena rupens.

DICHROMENA, Michx. Fl. Bor. Am. i. 87 (1803). *CYPERACEÆ*, Benth. & Hook. f. iii. 1047.
DICHROMA, Pers. Syn. i. 87 (1805).
albida, *Steud. Syn. Pl. Cyp.* 136.—Bras.
atrosanguinea, E. Desv. in C. Gay, Fl. Chil. vi. 190 = Scirpus riparius.
blephanophora, Steud. Syn. Pl. Cyp. 136 = ciliata.
blepharophora, J. & C. Presl, Rel. Haenk. i. 251 = paludicola.
caespitosa, Spreng. Syst. i. 202 (= *Isolepis stenophylla*).—Am. bor.
canaliculata, Steud. Syn. Pl. Cyp. 136 = Rynchospora tenuis.
candida, Baker, in Journ. Linn. Soc. xxi. (1885) 450, (Rynchosporæ sp.).—Madag.
candida, Rldi. in Trans. Linn. Soc. Ser. II. ii. (1884) 149 = Rynchospora candida.
capillaris, Kunth, Enum. Pl. ii. 283 = Rynchospora tenuis.
caracasana, *Kunth, l. c.* 281.—Venezuela.
ciliata, J. & C. Presl, Rel. Haenk. i. 197 = pulchella.
ciliata, *Vahl, Enum.* ii. 240.—Am. trop.
compressa, Spreng. Neue Entdeck. i. 288 = Rynchospora rigida ?
consanguinea, Kunth, Enum. Pl. ii. 278 = Rynchospora consanguinea.
contracta, Steud. Syn. Pl. Cyp. 136 (Rynchosporæ sp.).—Bras.
cubensis, Poepp. & Kunth, in Kunth, Enum. Pl. ii. 281 = Rynchospora tenuis.
Drummondiana, Steud. Syn. Pl. Cyp. 135 = leucocephala.
elatior, Kunth, Enum. Pl. ii. 282 = Rynchospora stricta.
emarginata, Steud. Syn. Pl. Cyp. 136 = Rynchospora tenuis.
erostris, Nees, in Linnæa, ix. (1834) 291 = pubera.
filiformis, Kunth, Enum. Pl. ii. 281 = Rynchospora longispicata.
foliosa, Hochst. ex Boeck. in Linnæa, xxxvii. (1873–72) 534 = Rynchospora albiceps.
fragifera, A. Dietr. Sp. Pl. ii. 169 = Calyptrocarya fragifera.
globosa, Roem. & Schult. Syst. ii. 90 = ciliata.
gracilis, Kunth, Enum. Pl. ii. 280 = Rynchospora tenuis.
heterophylla, Boeck. in Flora, xli. (1858) 644 = Rynchospora Chapmanii.
hirsuta, Kunth, Enum. Pl. ii. 281 = Rynchospora hirsuta.
hispida, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 111 = Rynchospora consanguinea.
hispidula, Kunth, Enum. Pl. ii. 279 (Rynchosporæ sp.).—Guiana.
Humboldtiana, Nees, in Mart. Fl. Bras. ii. 1. 111 = pubera.
imbricata, Schrad. ex Steud. Nom. ed. II. i. 503 = pubera.
junciformis, Kunth, Enum. Pl. ii. 279 = Rynchospora junciformis.
latifolia, *Baldw. ex Ell. Sketch*, i. 90.—Am. bor.
leucocephala, *Michx. Fl. Bor. Am.* i. 87.—Am. bor.; Ind. occ.
leucocephala, Willd. ex Steud. Nom. ed. II. i. 503 = ciliata.
leucophylla, Willk. ex Kunth, Enum. Pl. ii. 276 = ciliata.
longifolia, Spreng. Syst. i. 203 = Calyptrocarya longifolia.
micrantha, Kunth, Enum. Pl. ii. 278 = Rynchospora micrantha.
microcephala, Spreng. Syst. i. 202 = pubera.
nardifolia, Kunth, Enum. Pl. ii. 280 = Rynchospora nardifolia.
nervosa, Vahl, Enum. ii. 241 = pubera, ciliata.
nitens, Kunth, Enum. Pl. ii. 282 = Rynchospora nitens.
obtusiflora, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 113 = ciliata.
obtusifolia, Steud. Syn. Pl. Cyp. 136 = ciliata.
Parcensiana, Nees, in Mart. Fl. Bras. ii. 1. 112 = ciliata.
pilosa, Kunth, Enum. Pl. ii. 279 = Rynchospora pilosa.
pilosa, Spreng. Syst. i. 203 = ciliata.
Pohliana, Steud. Syn. Pl. Cyp. 319 = Rynchospora tenuis.
Presliana, A. Dietr. Sp. Pl. ii. 170 = pulchella.
pubera, *Vahl, Enum.* ii. 241 = Am. centr.
pulchella, *Poepp. & Kunth, in Kunth, Enum. Pl.* ii. 277.—Peruv.

DICHROMENA:—
pura, Nees, in Linnaea, ix. (1834) 291 = ellista.
pusilla, Kunth, Enum. Pl. ii. 275 (Rynchosporae sp.).—Ind. occ.
radicans, *Cham. & Schlecht. in Linnaea*, vi. (1831) 28.—Am. trop.
repens, *Vahl, Enum.* ii. 241.—Am. trop.
rigida, Kunth, Enum. Pl. ii. 278 = Rynchospora rigida.
riparia, Steud. Syn. Pl. Cyp. 136 = Rynchospora riparia.
robusta, Kunth, Enum. Pl. ii. 288 = Rynchospora robusta.
rufidula, *Steud. Syn. Pl. Cyp*, 137.—Am. bor.
sabulosa, Kunth, Enum. Pl. ii. 288 = Rynchospora tenuis.
Schiedeana, Kunth, l. c. 282 = Rynchospora eximia.
Sellowiana, Kunth, l. c. (Rynchosporae sp.).—Bras.
setacea, Kunth, l. c. 279 = Rynchospora tenerrima, *Nees*.
setigera, Kunth, l. c. 277 = Rynchospora setigera.
speciosa, Kunth, l. c. 278 = Rynchospora speciosa.
squarrosa, Willd. ex Luk, Jahrb. i. *111*. 77 = Cephalostemon squarrosus.
stolonifera, Steud. Syn. Pl. Cyp. 136 = Rynchospora stoloniféra.
tenella, Steud. Syn. Pl. Cyp. 136 = Rynchospora tenella.
tenera, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 113 = pubera.
tenuifolia, *Spreng. Syst.* i. 203.—Am. austr. trop.
tenuis, Steud. Syn. Pl. Cyp. 135 = radicans.
trachycarpa, *Steud. l. c.* 136.—Hab.?
unicolor, Salzm. ex Steud. l. c. = pubera.
Vahlii, A. Dietr. Sp. Pl. ii. 169 = Scleria communis.
velutina, Kunth, Enum. Pl. ii. 282 = Rynchospora velutina.

DICHRONEMA, Baker, in Journ. Linn. Soc. xxi. (1885) 450 = **Dichromena**, Michx. (Cyperac.).

DICHROPHYLLUM, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 249 = **Euphorbia**, Linn.
bicolor, Klotzsch & Garcke, l. c.; et in Abh. Akad. Berl. (1860) 44 = E. bicolor.
marginatum, Klotzsch & Garcke, ll. cc. = E. marginata.
variegatum, Klotzsch & Garcke, ll. cc. = E. marginata.

DICHROSTACHYS, Wight & Arn. Prod. 271 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 592. CAILLIA, Guill. & Perr. Fl. Senegj. Tent. 239 (1833). Benteriana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 355.—Madag.
brachypus, *Baill. l. c.* 356.—Madag.
cafra, Meisn. ex Krauss, in Flora, xxvii. (1844) 359 = nutans.
callistachys, Hassk. Pl. Jav. Rar. 415 = nutans.
cinerea, Miq. Fl. Ind. Bat. i. 1. 48 = nutans.
cinerea, *Wight & Arn. Prod.* 271.—Geront. trop.
dehiscens, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 511.—Ins. Socotra.
Forbesii, *Benth. in Hook. Journ. Bot.* iv. (1842) 353.—Afr. austr.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 356.—Madag.
Kirkii, *Benth. in Trans. Linn. Soc. xxx.* (1875) 383.—Afr. trop.
macrostachya, *Walp. Rep.* v. 583.—Guiana.
Muelleri, *Benth. Fl. Austral.* vii. 299.—Austral.
nutans, *Benth. in Hook. Journ. Bot.* iv. (1842) 353.—Afr. trop.
platycarpa, *Welw. Apont.* 576.—Afr. trop.
Richardiana, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 355.—Madag.
tenuifolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 353.—Madag.

DICHROTRICHIUM, Benth. & Hook. f. Gen. ii. 1014 (1876) = seq.

DICHROTRICHUM, Reinw. ex De Vriese, Pl. Ind. Bat. Or. 7. t. 1 (1856). *GESNERACEAE*, Benth. & Hook. f. ii. 1014.
THOMSONIOPSIS, [Blume ?] R. Br. in Benn. Pl. Jav. Rar. 124 (1838).
superciliosum, *Benth. & Hook. f. ex C. B. Clarke, in DC. Monog. Phan.* v. 54.—Java.
brevipes, *C. B. Clarke, l. c.*—N. Guinea.

DICHROTRICHUM:—
Chalmersii, *F. Muell. in Melb. Chem. & Drug.* (June 1884).—N. Guinea.
chortesplanum, *C. B. Clarke, in DC. Monog. Phan.* v. 53.—Ins Philipp.
Griffithii, C. B. Clarke, Comm. & Cyrt. Beng. t. 51 = Loxostigma Griffithii.
ternatum, *Reinw. ex De Vriese, Pl. Ind. Bat. Or.* 7. t. 1, 2.—Ins. Ternat.

DICHYNCHOSIA, C. Muell. in Walp. Ann. v. 31 (1858) sphalm. = DIRHYNCHOSIA, Blume = **Spiraeopsis**, Miq. (Saxifr.).

DICKIA, Scop. Introd. 199 (1777) = **Stemodia**, Linn. (Scrophul.).

DICLADANTHERA, F. Muell. Fragm. xii. 23 (1882); Census, 140 (1882). *ACANTHACEAE*. Forrestii, *F. Muell. ll. cc.*—Austral.

DICLEMIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 102 = **Osmaea**, DC. (Melast.).
petiolaris, Naud. l. c. 102 = O. petiolaris.

DICLIDANTHERA, Mart. Nov. Gen. et Sp. ii. 139. t.196, 197 (1827). *STYRACEAE*, Benth. & Hook. f. ii. 671.
PLUCHEA, Vell. Fl. Flum. 157 (1825); iv. t. 20 (1827).
elliptica, *Miers, in Ann. Nat. Hist.* Ser. III. iv. (1859) 186.—Bras.
laurifolia, *Mart. Nov. Gen. et Sp. ii.* 141. t. 196.—Bras.
pendulifloras, *Mart. l. c.* 140. t. 197.—Bras.

DICLIDIUM, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 51 (1842) = **Mariscus**, Gaertn. (Cyperac.).
aciculare, Schrad. ex Nees, in Mart Fl. Bras. ii. t. 55 (= *Cyperus ferax*).—Bras.
auriculatum, Nees, l. c. 51 in obs. (= *Cyperus ferax*).—Bras.
conglobatum, Nees, l. c. in obs. (= *Cyperus ferax*).—Bras.
ferax, Steud. Syn. Pl. Cyp. [Index] 332 (= *Cyperus ferax*) —Bras.
feroz, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 54 (= *Cyperus ferax*).—Bras.
fuscatum, Schrad. ex Nees, l. c. 52 (= *Cyperus ferax*).—Bras.
lentiferum, Schrad. ex Nees, l. c. 53 (= *Cyperus ferax*).—Bras.
lomentaceum, Nees, l. c. 55 (= *Cyperus ferax*).—Bras.
Maximilianei, Schrad. ex Nees, l. c. 52 (= *Cyperus ferax*).—Bras.
odoratum, Schrad. ex Nees, l. c. 54 (= *Cyperus ferax*). —Ind. occ.
Reinwardtii, Schrad. ex Nees, l. c. 51 in obs. (= *Cyperus ferax*).—Bras.
uliginosum, Schrad, ex Nees, l. c. 56 (= *Cyperus ferax*). —Bras.
Vahlii, Schrad. ex Nees, l. c. 53 = Cyperus Vahlii.

DICLIDOCARPUS, A. Gray, Bot. U. St. Expl. Exped. i. 200. t. 14 (1854) = **Trichospermum**, Blume (Tiliac.).
Richii, A. Gray, l. c. = T. Richii.

DICLIDOSTIGMA, Kunze, in Linnaea, xvii. (1843) 576 = **Malachra**, Linn. (Cucurb.).
melothrioides, Kunze, l. c. 577 = M. pervaga.

DICLINOTHRYS, Endl. Gen. Suppl. i. 1257 (1841) = seq.

DICLINOTRYS, Rafin. Neogen. 3 (1825) = **Chamaelirium**, Willd. (Liliac.).
albiflorum, Rafin. l. c. = C. carolinianum.

DICLIPTERA, Juss. in Ann. Mus. Par. ix. (1807) 267. *ACANTHACEAE*, Benth. & Hook. f. ii. 1130. BRACHOSIPHON, Nees, in DC. Prod. xi. 492 (1847). DACTYLOSTEGIUM, Nees, in Mart. Fl. Bras. ix. 102 t. 31 (1847).
DIAPHRAGMA, Kom. ex Steud. Nom. ed. II. i. 501, 504 (1840).

DICLIPTERA:—
GRUNSIA, Neck. Elem. i. 231 (1790).
KUNIRIA, Rafin. Fl. Tellur. iv. 61 (1836).
PANNAYA, Rafin. l. c. (1836).
acuminata, *Juss. in Ann. Mus. Par.* ix. (1807) 268.—Ecuador.
brgolensis, *A. Nees, in Journ. Bot.* xviii. (1880) 362.—Afr. trop.
armata, F. Muell. Fragm. vi. 88 = glabra.
assurgens, *Juss. in Ann. Mus. Par.* ix. (1807) 269.—Am. trop.
bephica, *Nees, in DC. Prod.* xi. 490.—Mexic.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 551.—Ind. or.
bicalyculata, Kostel. Allg. Med.-Pharm. Fl. iii. 934 = Peristrophe bicalyculata.
bivalvis, Blume, Bijdr. 789 = javanica.
bivalvis, *Juss. in Ann. Mus. Par.* ix. (1807) 268.—Ind. or.; Arab.
bivalvis, Nees, in DC. Prod. xi. 475 = xylasica.
blepharostegia, Presl, Bot. Bemerk. 95 = Justicia debilis.
brachiata, Salzm. ex Nees, in DC. Prod. xi. 478 = mucronifolia.
brachiata, *Spreng. Syst.* i. 86.—Am. bor.
bracteata, *Sesm. Fl. Vit.* 184.—Ins. Tahiti.
brasiliensis, Nees & Mart. in Nov. Act. Nat. Cur. xi. i. (1823) 60 (= *Rhytiglossa leucophloea*).—Bras.
brasiliensis, *Spreng. Neue Entdeck.* ii. 128.—Bras.
Buergeriana, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 125; Franch. & Sav. Enum. Pl. Jap. i. 356 = Roxburghiana.
bupleuroides, Nees, in Wall. Pl. As. Rar. iii. 111 = Roxburghiana.
Burmanni, *Nees, l. c.* 112.—Java.
Burmanni, Nees, l. c. partim; T. Anders. in Journ. Linn. Soc. ix. (1868) 520 = D. Leonotis.
canescens, *Nees, in DC. Prod.* xi. 476.—Java.
cayennis, *Nees, in Linnaea*, xv. (1841) 378.—Afr. austr.
cardiocarpa, Nees, in Wall. Pl. As. Rar. iii. 111 = Roxburghiana.
carassana, *Nees, in DC. Prod.* xi. 474.—Venezuela.
chinensis, *Juss. in Ann. Mus. Par.* ix. (1807) 268; *Nees, in DC. Prod.* xi. 477.—As. trop.
chinensis, Roem. & Schult. Syst. Mant. i. 146 = Roxburghiana.
ciliaris, *Juss. in Ann. Mus. Par.* ix. (1807) 268.—Am. trop.
ciliaris, Willd. ex Nees, in DC. Prod. xi. 484 = Schwabea ciliaris.
ciliata, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 384 = Burmanni.
ciliata, Ruiz & Pav. Fl. Per. et Chil. i. 12. t. 14. f. a = Beloperone ciliata.
ciliata, Schau. in ind. Sem. Hort. Vrat. ex Nees, in DC. Prod. xi. 479 = Moritziana.
clavata, *Juss. in Ann. Mus. Par.* ix. (1807) 269.—Ins. Tahiti.
clunopoda, *Nees, in DC. Prod.* xi. 488.—Afr. austr.
coerulea, Blume, Bijdr. 790 = Rungia parviflora.
confinis, *Nees, ex Benth. Bot. Voy. Sulph.* 149.—Ecuador.
congesta, *H. B. & K. Nov. Gen. et Sp.* ii. 337.—Mexic.
crenata, *Miq. Fl. Ind. Bat.* ii. 844.—Samatra.
crinita, Nees, in DC. Prod. xi. 485 = Roxburghiana.
cuneata, *Nees, in Wall. Pl. As. Rar.* iii. 111.—Ind. or.
effusa, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 89.—Ins. Socotra.
erlantha, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 384.—Ins. Timor.
falcata, Poir. ex Steud. Nom. ed. I. 274 = laevigata.
fasciata, Presl, Bot. Bemerk. 95 = Justicia fasciata.
Forsteriana, *Nees, in DC. Prod.* xi. 475.—Hab.?
frondosa, *Juss. in Ann. Mus. Par.* ix. (1807) 269.—Ins. Tahiti.
furcifera, Nees, in Mart. Fl. Bras. ix. 96 = Herpetacanthus melanocholicus.
glabra, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 383.—Ins. Timor.
glandulosa, Schreeb, in Linnaea, xxi. (1848) 765 = brachiata.
Hasskarana, *Nees, in DC. Prod.* xi. 487.—Mexic.
Helei, Riddell, in N. Orl. Med. Journ. (1852); Chapm. Fl. S. U. St. 305 = Gatesia laete-virens.
heterotegia, *Presl, Bot. Bemerk.* 95; *Nees, in DC. Prod.* xi. 478.—Afr. austr.

DICLIPTERA:—
hirtula, Nees, in DC. Prod. xi. 485 = Roxburghiana.
Hookeriana, *Nees, l. c.* 480.—Peruv.
hyalina, *Nees, l. c.* 484.—Ins. Comor.
interrupta, *Blume, Bijdr.* 790.—Java.
javanica, *Nees, in DC. Prod.* xi. 476.—Java.
Vanieri, Webb, in Hook. Niger Fl. 188 = verticillaris.
Kurzii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 551. —Burma.
laevigata, *Juss. in Ann. Mus. Par.* ix. (1807) 268.— Ins. Mascar.
Leonotis, *Dals. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 559.—Ind. or.
lupulina, Presl, Bct. Bemerk. 95 = Adhatoda lupulina.
maculata, *Nees, in DC. Prod.* xi. 485.—Afr.trop.
madagascariensis, *Nees, l. c.* 476.—Madag.
martinicensis, *Juss. in Ann. Mus. Par.* ix. (1807) 268. —Ins. Martinic.
melancholica, Mart. ex Nees, in Mart. Fl. Bras. ix. 96 = Hemigenia melancholica.
micrantha, *Nees, in Wall. Pl. As. Rar.* iii. 112.— Ind. or.
mollis, *Nees, in DC. Prod.* xi. 490.—Mexic.
Morittiana, *S. Schau. in Ind. Sem. Hort. Vrat.* (1848) *ex Nees, l. c.* 479.—Am. trop.
mosambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 230.—Afr. trop.
mucronifolia, *Nees, in Mart. Fl. Bras.* ix. 161.— Bras.
multiflora, *Juss. in Ann. Mus. Par.* ix. (1807) 268.— Ecuador.
multiflora, Nees, in DC. Prod. xi. 487 = Hasenkeana.
ocymoides, *Juss. in Ann. Mus. Par.* ix. (1807) 268. —Am. austr.
ovata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 59.—Ins. Socotra.
ovata, Presl, Bot. Bemerk. 95 (= *Rhytiglossa ovata*). —Afr. austr.
paposana, *Phil. Fl. Atac.* 40.—Chili.
parvibracteata, *Nees, in Wall. Pl. As. Rar.* iii. 111.— Ind. or.
pectinata, Juss. in Ann. Mus. Par. ix. (1807) 269 = Rungia parviflora.
pedunculariа, *Nees, in DC. Prod.* xi. 488.—Mexic.
peruviana, *Juss. in Ann. Mus. Par.* ix. (1807) 268.— Peruv.
pilosa, *H. B. & K. Nov. Gen. et Sp.* ii. 237.—N. Granat.
Pohliana, *Nees, in Mart. Fl. Bras.* ix. 162.—Bras.
portoricensis, *Spreng. in Roem. & Schult. Syst. Mant.* i. 149.—Porto Rico.
procumbens, *Humd. ex Link, Jahrb.* i. 111. 48.—Am. austr.
propinqua, *Nees, in Linnaea*, xv. (1841) 573.—Afr. austr.
pseudoverticillaris, *A. Gray, in Proc. Am. Acad.* xx. (1885) 308.—Arizona.
pubescens, *Juss. in Ann. Mus. Par.* ix. (1807) 269.— Ins. N. Hebrid.
racemifera, F. Muell. Fragm. vi. 89 = spicata.
ramentacea, Spreng. ex Nees, in DC. Prod. xi. 64 = Elytraria frondosa.
recurvata, Spreng. Syst. iv. Cur. Post. 19 (= *Rhytiglossa leucophlœa*).—Bras.
repens, Juss. in Ann. Mus. Par. ix. (1807) 269 = Rungia repens.
repens, Willd. ex Nees, in DC. Prod. xi. 470 = Rungia parviflora.
reptans, *Nees, l. c.* 488.—Burma.
resupinata, *Juss. in Ann. Mus. Par.* ix. (1807) 268.— Mexic.
resupinata, Nutt. ex Nees, in DC. Prod. xi. 480 = brachiata.
retusa, *Juss. in Ann. Mus. Par.* ix. (1807) 268.— Ind. occ.
Rheedii, Kostel. Allg. Med.-Pharm. Fl. iii. 934 = Justicia Gendarussa.
riparia, *Nees, in Wall. Pl. As. Rar.* iii. 112.—Burma. Roxburghiana, *Nees, l. c.* 111.—Ind. or.
Roxburghii, T. Anders. in Journ. Linn. Soc. ix. (1867) 519 = pracc.
rupestris, Nees, in DC. Prod. xi. 486 = Roxburghiana.
samoensis, *Seem. Fl. Vit.* 194.—Ins. Samoa.
scabra, *Nees, in DC. Prod.* xi. 480.—Ecuador.
scorpioides, Juss. in Ann. Mus. Par. ix. (1807) 269 = Tetramerium scorpioides.
scutellata, *Griseb. in Goett. Abh.* xix. (1874) 236.— Reg. Argent.

DICLIPTERA:—
sericea, *Nees, in Mart. Fl. Bras.* ix. 161.—Bras.
serpenginis, *Juss. in Ann. Mus. Par.* ix. (1807) 269. —Jamaic.
sparsiflora, *Nees, in DC. Prod.* xi. 488.—Bras.
speciosa, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 97.—Burma.
speciosa, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 61 = Beloperone involucrata.
spicata, Boj. ex Nees, in DC. Prod. xi. 386 = Adhatoda Bojeriana.
spicata, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 384.—Ins. Timor.
spinosa, Lodd. ex Nees, in DC. Prod. xi. 491 = Barleria lupulina.
spinulosa, Hochst. ex Nees, l. c. 484 = micrantha.
squarrosa, *Nees, in Mart. Fl. Bras.* ix. 161.—Bras.
tetrandra, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 61 = Herpetacanthus Schultzii.
thiaspioides, *Nees, in DC. Prod.* xi. 474.—Am. centr.
timoriensis, Moric. ex Miq. Fl. Ind. Bat. ii. 845 = Burmanni.
tinctoria, Kostel. Allg. Med.-Pharm. Fl. iii. 937 = Periscyphe tinctoria.
tomentosa, *Nees, in DC. Prod.* xi. 481.—Peruv.
Torreyi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 308. —Arizona.
trifurca, *Oerst. in Kjoeb. Vidensk. Meddel.* (1856) 172. —Costa Rica.
trifurcata, Hemsl. Biol. Centr. Am. Bot. ii. 525 = trinervia.
trinervia, Juss. in Ann. Mus. Par. ix. (1807) 269 = Justicia trinervia.
Tweediana, *Nees, in DC. Prod.* xi. 482.—Uruguay.
umbellata, Juss. in Ann. Mus. Par. ix. (1807) 268 = micrantha.
unguiculata, *Nees, ex Benth. Bot. Voy. Sulph.* 149.— Costa Rica.
Vahliana, *Nees, in DC. Prod.* xi. 489.—Am. trop.
velata, *Sonn. Fl. Vit.* 183.—Ins. Tahiti.
verticillaris, Juss. in Ann. Mus. Par. ix. (1807) 268; Roem. & Schult. Syst. i. 170 = Hypoëstes verticillaris.
villosa, Mart. ex Nees, in DC. Prod. xi. 200 = Ruellia villosa.
viridis, *Hassk. in Flora*, xlvii. (1864) 54.—Ins. Philipp.
Welwitschii, *S. Moore, in Journ. Bot.* xviii. (1880) 262.—Afr. trop.
zipotensis, Koem. & Schult. Mant. i. 248 (= *Rhytiglossa leucophlœa*).—Bras.
zeylanica, *Nees, in DC. Prod.* xi. 474.—Ind. or.

DICLIS, Benth. in Hook. Comp. Bot. Mag. ii. (1836) 22. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 963.
ovata, *Benth. l. c.*—Madag.
petiolaris, *Benth. in DC. Prod.* x. 265.—Afr. austr.
reptans, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 22.—Afr. austr.

DICLYTHRA, Raf. Cat. 13 (1824) = DICLYTRA, Borkh. = Dicentra, Benth. (Papav.).

DICLYTRA, Borckh. in Roem. Arch. i. ii. 46 (1797) = Dicentra, Benth. (1858) (Papav.).
bracteata, DC. Syst. ii. 109.—Am. bor. †
canadensis, Borckh. in Roem. Arch. i. ii. 46 = Dicentra Cucullaria.
canadensis, DC. Prod. i. 126 = Dicentra canadensis.
Cucullaria, DC. Syst. ii. 108 = Dicentra Cucullaria.
divaricata, Raf. Cat. 13.—Am. bor.
eximia, Hook. Bot. U. St. 33 = Dicentra canadensis.
eximia, DC. Syst. ii. 109 = Dicentra eximia.
formosa, DC. l. c. = Dicentra formosa.
formosa, Ell. Sketch. ii. 177 = Dicentra eximia.
lachenalieaefora, DC. Syst. ii. 111 = Dicentra lachenalieaefolia.
scandens, D. Don, Prod. Fl. Nep. 198 = Dicentra scandens.
spectabilis, DC. Syst. ii. 110 = Dicentra spectabilis.
tenuifolia, DC. l. c. = Dicentra lachenalieaefolia.
thalictrifolia, D. Don, ex Walp. Rep. i. 118, et Index, 873 = Dicentra thalictrifolia.

DICNECKERIA, Vall. Fl. Flum. 61 (1825) = Euplassa, Salisb. (Proteac.).
ingaifa, Vell. l. c. 61; i. t. 105 (= *Adenostephanus Sellowii*).—Bras.

DICOCCA, Thon. Gen. Nov. Madag. 12, in syn. (1806) = Dicorypha, Thon. (Hamamel.).

DICODON, Ehrb. Beitr. ix. 146 (1789) = Linnaea, Gronov. (Caprifol.).

DICOELIA, Benth. in Hook. Ic. Pl. t. 1289 (1879). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 286.
leucarania, *Benth. l. c.*—Borneo.

DICOLUS, Phil. in Anal. Univ. Chil. (1873) 550; Desc Pl. Nuev. il. 74 = Ephyra D. Don (Hazmodor.)
caerulescens, Phil. ll. cc. 551; 75 = Z. elegans.

DICOMA, Cass. in Bull. Soc. Philom. (1817) 12. *COMPOSITAE*, Benth. & Hook. f. ii. 492.
CRYPTOSTEPHANE, Sch. Bip. in Flora, xxvii. (1844) 782.
MACLEDIUM, Cass. in Dict. Sc. Nat. xxxiv. 39 (1825)
NITELIUM, Cass. l. c. xxxv. 11 (1825).
RHIGIOTHAMNUS, Spach, Hist. Vég. Phan. x. 9 (1841).
SCHAFFNERA, Sch. Bip. in Flora, xxiv. (1841) 1 Intell. 42.
STEIROCOMA, Reich. Nom. 88 (1841).
XEROPAPPUS, Wall. Cat. n. 2980 (1830 ?).
anomala, *Sond. in Linnaea*, xxiii. (1850) 71.—Afr. austr.
argyrophylla, *Oliver, in Hook. Ic. Pl.* t. 1461.—Afr. austr.
Burmanni, *Less. in Linnaea*, v. (1830) 279.—Afr. austr
cana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 841. —Ins. Socotra.
capensis, *Less. in Linnaea*, v. (1830) 277.—Afr. austr.
dicanthoides, *Less. l. c.* 279.—Afr. austr.
karaguensis, *Oliver, in Trans. Linn. Soc.* xxix. (1874) 108.—Afr. trop.
Kirkii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 518.— Afr. austr.
Krausii, Sch. Bip. in Flora, xxvii. (1844) 782 = dicanthoides.
lanuginosa, DC. Prod. vii. 36 = tomentosa.
latifolia, *DC. l. c.* 37.—Afr. austr.
macrocephala, *DC. l. c.* 36.—Afr. austr.
radiata, *Less. in Linnaea*, v. (1830) 278.—Afr. austr.
relhanioides, *Less. l. c.* 279.—Afr. austr.
sessiliflora, *Harv. in Harv. & Sond. Fl. Cap.* iii. 518.—Afr. austr.
speciosa, *DC. Prod.* vii. 37.—Afr. austr.
tomentosa, *Cass. in Bull. Soc. Philom.* (1818) 12.— Ind. or.; Afr. trop.
Zeyheri, *Sond. in Linnaea*, xxiii. (1850) 70.—Afr. austr.

DICONANGIA, Mitchell, in Act. Phyt. Med. Acad. Nat. Cur. viii. (1748) 208, App. = Itea, Linn (Saxifr.).

DICOPHE, Roem. Coll. 205 (1809) = Dicorypha, Thon. (Hamamel.).

DICORIA, Torr. & Gray, in Emory, Rep. 148 (1848). *COMPOSITAE*, Benth. & Hook. f. ii. 358.
Brandigei, *A. Gray, in Proc. Am. Acad.* xi. (1876) 76. —Am. bor. occ.
canescens, *A. Gray, in Emory, Rep.* 148.—Calif.

DICORYNEA, Lindl. Veg. Kingd. 655 (1847) = seq.

DICORYNIA, Benth. in Hook. Journ. Bot. ii. (1840) 82. *LEGUMINOSAE*, Benth. & Hook. f. i. 571.
brevifora, *Benth. in Mart. Fl. Bras.* xv. ii. 83.— Bras.
Baribunda, Spruce, ex Benth. l. c., in obs. = paraënsis ?
paraënsis, *Benth. in Hook. Journ. Bot.* ii. (1840) 82. —Bras.
Spruceana, Benth. in Mart. Fl. Bras. xv. ii. 81 = paraënsis.
sonpensis, Spruce, ex Benth. l. c., in obs. = paraënsis ?

DICORYPHA, R. Hedw. Gen. 81 (1806) = seq.

DICORYPHE, Thon. Gen. Nov. Madag. 12 (1806). *HAMAMELIDEAE*, Benth. & Hook. f. i. 667.
DIANIA, Noronha, ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 149.
DICOCCA, Thon. Gen. Nov. Madag. 12, in syn. (1806).
DICOPHE, Roem. Collect. 205 (1809).

DICORYPHE:—
angustifolia, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 145.—Madag.
buddleoides, *Baker, in Journ. Bot.* xx. (1882) 111.—Madag.
gracilis, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 144.—Madag.
lanceolata, *Tul. l. c.* 146.—Madag.
laurina, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 476.—Madag.
macrophylla, *Baill. l. c.*—Madag.
madagascariensis, Poir. Encyc. Suppl. ii. 475; Illustr. iii. 607 = stipulacea.
Noronhae, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 147.—Madag.
platyphylla, *Tul. l. c.* 146.—Madag.
stipulacea, *Yanme St. Hil. Expos. Fam.* ii. 348.—Madag. *Thouarii*, Roem. & Schult. Syst. iii. 486 = stipulacea.
viticoides, *Baker, in Journ. Linn. Soc.* xx. (1883) 143.—Madag.

DICRAEA, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 100 = seq.

DICRAEIA, Thou. Gen. Nov. Madag. 2 (1806) = **Podostemon**, Michx.
algaeformis, Bedd. in Trans. Linn. Soc. xxv. (1866) 223. t. 24 = P. algaeformis.
algaeoides, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 54 = prasc.
apicata, Tul. in Arch. Mus. Par. vi. (1852) 204 = P. dichotomus.
dichotoma, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 101 = P. dichotomus.
elongata, Tul. l. c. 102 = P. elongatus.
Erythrolichen, Tul. & Wedd. ex Tul. l. c.—Bras.
imbricata, Tul. l. c. 100.—Madag.
longifolia, Wight, Ic. t. 1916. f. 4 = P. dichotomus.
minor, Wedd. in DC. Prod. xvii. 71 = P. minor.
minutiflora, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 100 = P. minutiflora.
pterophylla, Wedd. in DC. Prod. xvii. 71 = P. pterophyllus.
rigida, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 101 = P. dichotomus.
selagnoides, Wedd. in DC. Prod. xvii. 66.—Ind. or.
stylosa, Wight, Ic. t. 1917. f. 2 = P. stylosus.
Wallichi, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 101 = P. Wallichii.
Wightii, Tul. l. c. = P. dichotomus.
Willdenowii, Tul. l. c.—Peruv.; Chili.

DICRAMA, Klatt, in Linnaea, xxxii. (1863) 751, sphalm. = **Dierama**, C. Koch (Irid.).

DICRANACANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136 = **Barleria**, Linn. (Acanthac.)
bispinosa, Oerst. l. c. = B. bispinosa.
buxifolia, Oerst. l. c. = B. buxifolia.
Spina-asylanica, Oerst. l. c. = B. myserensis.
spinicyma, Oerst. l. c. = B. splnicyma.
stimulans, Oerst. l. c. = B. stimulans.

DICRANANTHERA, Presl, Symb. Bot. i. 75. t. 50 (1832) = **Acisanthera**, P. Br. (Melastom.).
hedyotidea, Presl, l. c. 76. t. 50 = A. hedyotidea.
Salzmanni, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 332. t. 13. f. 1 = A. Salzmanni.

DICRANOCARPUS, A. Gray, in Mem. Am. Acad. Ser. II. v. (Pl. Nov. Thurb.) 322 (1855). *COMPOSITAE*, Benth. & Hook. f. ii. 347.
parviflorus, *A. Gray, l. c.*—Mexic.

DICRANOLEPIS, Planch. in Hook. Ic. Pl. t. 798 (1848). *THYMELAEACEAE*, Benth. & Hook. f. iii. 126.
disticha, *Planch. l. c.*—Afr. trop.
Maanii, *Baill. Adansonia*, xi. (1875) 302.—Afr. trop.

DICRANOPETALUM, Presl, in Abh. Boehm. Ges. V. iii. (1845) 464; (Bot. Bemerk. 24) = **Toulicia**, Aubl. (Sapindac.).
polyphyllum, Presl, ii. cc.—Bras.

DICRANOSTACHYS, Trée. in Ann. Sc. Nat. Sér. III. viii. (1847) 58. t. 1 = **Myrianthus**, Beauv. (Urtic.).
serrata, Trée. l. c. = M. serrata.

DICRANOSTIGMA, Hook. f. & Thoms. Fl. Ind. i. 255 (1855) = **Stylophorum**, Nutt. (Papav.).
lactucoides, Hook. f. & Thoms. l. c. = S. lactucoides.

DICRANOSTYLES, Benth. in Hook. Lond. Journ. Bot. v. (1846) 355. *CONVOLVULACEAE*, Benth. & Hook. f. ii. 878.
densa, *Spruce, ex Meisn. in Mart. Fl. Bras.* vii. 328.—Bras.
scandens, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 355.—Guiana.

DICRASTYLIS, Drum. ex Harv. in Hook. Kew Journ. vii. (1855) 56. *VERBENACEAE*, Benth. & Hook. f. ii. 1140.
Beveridgei, *F. Muell. Fragm.* viii. 50.—Austral.
Doranii, *F. Muell. l. c.* 230.—Austral.
fulva, *Drum. ex Harv. in Hook. Kew Journ. Bot.* vii. (1855) 58.—Austral.
Gilesii, *F. Muell. Fragm.* viii. 228.—Austral.
Lewellini, *F. Muell. l. c.* xi. 86.—Austral.
Nicholasii, *F. Muell. l. c.* x. 16.—Austral.
ochrotricha, *F. Muell. l. c.* iv. 161.—Austral.
parviflora, *F. Muell. l. c.* ii. 180.—Austral.
reticulata, *Drum. ex Harv. in Hook. Kew Journ.* vii. (1855) 57.—Austral.
rosmarinifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 326 = parviflora.
Stoechas, *Drum. ex Harv. in Hook. Kew Journ.* vii. (1855) 57.—Austral.

DICRAURUS, Hook. f. in Benth. & Hook. f. Gen iii. 42 (1880). *AMARANTACEAE*, Benth. & Hook. f. iii. 42.
leptoclades, *Hook. f. l. c.* 43.—Texas.

DICROBOTRYON, Endl. Gen. 540 in syn. (1838) = seq.

DICROBOTRYUM, Willd. ex Roem. & Schult. Syst. v. 221 (1819) = **Guettarda**, Linn. (Rubiac.).
divaricatum, Willd. ll. cc. = G. xylosteoides.

DICROPHYLA, Raf. Fl. Tellur. ix. 39 (1836) = **Haemaria**, Lindl. (Orchid.).
elegans, Raf. l. c. = H. discolor.

DICRUS, Reinw. in Blume, Cat. Gew. Buitenz. 47 (1823) = **Voacanga**, Thou. (Apocyn.).
fortidus, Reinw. l. c. = V. foetida.

DICRYPTA, Lindl. Gen. et Sp. Orch. 44, 152 (1830) = **Maxillaria**, Ruiz & Pav. (Orchid.).
Baueri, Lindl. l. c. 44, 152 = M. crassifolia.
bicolor, Benth. ex Lond. Hort. Brit. Suppl. ii. 630 = M. bicolor?
crassifolia, Lindl. ex Loud. l. c. Suppl. iii. 586 = M. crassifolia.
densifolia, Poepp. & Endl. Nov. Gen. et Sp. i. 39. t. 68 = M. densifolia.
discolor, Lodd. ex Lindl. Bot. Reg. (1839) Misc. 91 = M. discolor.
elatior, Reichb. in Linnaea, xvii. (1843) 408 = M. elatior.
iridifolia, Batem. ex Loud. Hort. Brit. Suppl. ii. 630 = M. iridifolia?
irisphyta, Rodrig. Orch. Nov. i. 126.—Bras.
longifolia, Rodrig. l. c. 125.—Bras.
villosa, Rodrig. l. c.—Bras.

DICTAMNUS, Linn. Syst. ed. I (1735). *RUTACEAE*, Benth. & Hook. f. i. 387.
FRAXINELLA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 291 (1745).
albus, *Linn. Sp. Pl.* 383.—Europ.; As. bor.
angustifolius, Sweet, Brit. Flow. Gard. Ser. II. i. t. 93 = albus.
Calodendrum, Poir. Encyc. Suppl. ii. 476; Illustr. ii. 479 = Calodendrum capense.
capensis, Linn. f. Suppl. 232 = Calodendrum capense.
dasycarpus, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 587.—Dahuria.
Fraxinella, Pers. Syn. i. 464 = albus.
Gymnostylis, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 338 = albus.
himalayensis, Royle, Illustr. Bot. Himal. 156 = albus.
hispanicus, Webb, ex Nym. Consp. 142 = albus.
microphyllus, Schur, Enum. Pl. Transs. 141 = albus.

DICTAMNUS:—
obtusiflorus, Koch, Syn. Fl. Germ. ed. II. 180 = albus
odorus, Salisb. Prod. 320 = albus.
sessilis, Wallr. in Linnaea, xiv. (1840) 569 = albus.
solitarius, Stokes, Bot. Mat. Med. ii. 485 = albus.
suffultus, Wallr. in Linnaea, xiv. (1840) 569; et Beitr. Fl. Hercyn. 199 = albus.

DICTAMUS, S. G. Gmel. Fl. Sibir. iv. 177 (1769) sphalm. = prasc.

DICTILIS, Raf. Fl. Tellur. iii. 89 (1836) = **Otostegia**, Benth. (Labiat.).
moluccoides, Raf. l. c. = O. scariosa.

DICTYALOMA, Walp. Rep. i. 519 (1842) = **Dictyoloma**, A. Juss. (Simarub.).

DICTYANDRA, Welw. ex Benth. & Hook. f. Gen. ii. 85 (1873). *RUBIACEAE*, Benth. & Hook. f. i. c.
arborescens, *Welw. ex Benth. & Hook. f. l. c.*—Afr. trop.
involucrata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 86.—Afr. trop.

DICTYANTHES, Raf. in Loud. Gard. Mag. viii. (1832) 247 = **Aristolochia**, Linn.
labiosa, Raf. l. c. = A. cymbiflora.
stapeliaeflora, Reichb. Handb. 178 = Dictyanthus stapeliaeflorus.

DICTYANTHUS, Decne. in DC. Prod viii. 604 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 765.
KYTIDOLOMA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 319.
TYMPANANTHE, Hasak. in Flora, xxx. (1847) 757.
campanulatus, Reichb. Sel. Sem. Hort. Dread. (1850) 4.—Cf. Linnaea, xxiv. (1851) 207 = Pavonii.
parviflorus, *Hemsl. Biol. Centr. Am. Bot.* ii. 329.—Mexic.
Pavonii, *Decne. in DC. Prod.* viii. 605.—Mexic.
reticulatus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 329.—Mexic.
stapeliiflorus, *Reichb. Sel. Sem. Hort. Dresd.* (1850) 4.—Cf. Linnaea, xxiv. (1851) 207.—Mexic.

DICTYOCALYX, Hook. f. in Proc. Linn. Soc. i. (1845) 277; et Trans. Linn. Soc. xx. (1851) 202 = **Cacabus**, Benth. (Solanac.).
Miersii, Hook. f. in Trans. Linn. Soc. xx. (1851) 203 (= *Thingotem Miersii*).—Peruv.; N. Granat.
multivalvis, Hook. f. ex Walp. Rep. vi. 572 = Nicotiana quadrivalvis.
quadrivalvis, Hook. f. ex Walp. l. c. = Nicotiana quadrivalvis.

DICTYOCARPUS, Wight, in Madras Journ. Sc. v. (1837) 309; et in Ann. Sc. Nat. Sér. II. xi. (1839) 169 = **Sida**, Linn. (Malvac.).
truncatus, Wight, ii. cc. = S. Schimperiana.

DICTYOCARYUM, H. Wendl. in Bonplandia, viii. (1860) 106. *PALMAE*, Benth. & Hook. f. iii. 901
fuscum, *H. Wendl. in Bot. Zeit.* xxi. (1863) 131.—N. Granat.
glaucescens, *Linden, Illustr. Hortic.* xxviii. (1881) 81.—N. Granat.
Lamarckianum, *H. Wendl. in Bot. Zeit.* xxi. (1863) 131.—Bras.
Wallisii, *H. Wendl. in Kerch. Palm.* 243; *Linden, Illustr. Hortic.* xxviii. (1881) 81.—N. Granat.

DICTYODAPHNE, Blume, Mus. Bot. Lugd. Bat. i. 270 (1850) = **Endiandra**, R. Br. (Laurin.).
Candolleana, Meisn. in DC. Prod. xv. i. 80 = Dehaasia cuneata.
cuneata, Meisn. l. c. = E. cuneata.
macrophylla, Blume, Mus. Bot. Lugd. Bat. 332.—Malaya.
micrantha, Meisn. in DC. Prod. xv. i. 80.—Java.
rubescens, Blume, Mus. Bot. Lugd. Bat. i. 270 = E. rubescens
sumatrana, Meisn. in DC. Prod. xv. i. 80.—Sumatra
variabilis, Meisn. l. c. = E. rubescens.

DICTYOLOMA, A. Juss. in Mém. Mus. Par. xii. (1825) 499. t. 24. *SIMARUBEAE*, Benth. & Hook. f. i. 312.
BENJAMINA, Vell. Fl. Flum. 93 (1825); ii. t. 139.
DYCTIOLOMA, DC. Prod. ii. 69 (1825).
incanescens, DC. Prod. ii. 69 = Vandellianum.
peruvianum, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 583.—Peruv.
Vandellianum, *A. Juss. in Mém. Mus. Par.* xii. (1825) 499. t. 24.—Bras.

DICTYONEURA, Blume, Rumphia, iii. 168 (1849) = **Cupania**, Linn. (Sapind.).
acuminata, Blume, l. c. = C. acuminata.
obtusa, Blume, l. c. 165 = C. obtusa.

DICTYOPETALUM, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. (1835) 45 [4]; et ex Baill. Dict. ii. 410 (1884) = **Oenothera**, Linn. (Onagrar.).

DICTYOPSIS, Harv. ex Hook. f. in Bot. Mag. t. 5638 (1867) = **Behnia**, Didrich. (Liliac.).
Thunbergi, Harv. l. c. = B. reticulata.

DICTYOSPERMA, Regel, Fl. Nov. Fedsch. 5 (1882). *CRUCIFERAE*.
Olgae, *Regel, l. c.*—Turkestan.

DICTYOSPERMA, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 181. *PALMAE*, Benth. & Hook. f. iii. 890.
album, *H. Wendl. & Drude, in Linnaea*, xxxix. (1875) 181.—Ins. Mascar.
aureum, *H. Wendl. & Drude, in Kerch. Palm.* 248.—Hab.?
furfuraceum, *H. Wendl. & Drude, l. c.*—Hab.?
rubrum, *H. Wendl. & Drude, l. c.*—Hab.?

DICTYOSPERMUM, Wight, Ic. vi. 29. tt. 2069–71 (1853) = **Anellema**, R. Br. (Commel.).
conspicuum, Hassk. Commel. Ind. 22 = A. conspicuum.
montanum, Wight, Ic. vi. t. 2069 = A. montanum.
ovalifolium, Wight, l. c. t. 2070 = A. ovalifolium.
ovatum, Hassk. Commel. Ind. 24 = A. ovatum.
protensum, Wight, Ic. vi. t. 2071 = A. protensum.
Wightii, Hassk. Commel. Ind. 19 = A. ovalifolium.

DICTYOSTEGA, Miers, in Proc. Linn Soc. i. (1840) 61; et Trans. Linn. Soc. xviii. (1841) 538. t. 37. *BURMANNIACEAE*, Benth. & Hook. f. iii. 458.
campanulata, *Karst. in Linnaea*, xxviii. (1856) 422.—N. Granat.
costata, *Miers, in Proc. Linn. Soc.* i. (1840) 61; *Trans. Linn. Soc.* xviii. (1841) 542.—Bras.
longistyla, *Benth. in Hook. Niger Fl.* 528.—Afr. trop.
orobanchioides, *Miers, in Proc. Linn. Soc.* i. (1840) 61; *et in Trans. Linn. Soc.* xviii. (1841) 539.—Bras.
pectinata, *Karst. in Linnaea*, xxviii. (1856) 422.—N. Granat.
Purdieana, *Benth. in Hook. Kew Journ.* vii. (1855) 14.—N. Granat.
Schomburgkii, *Miers, in Proc. Linn. Soc.* i. (1840) 61; *et in Trans. Linn. Soc.* xviii. (1841) 542.—Guian.
umbellata, *Miers, ll. cc.* 61, 541.—Bras.

DICTYOSTEGIA, Benth. & Hook. f. Gen. iii. 458 (1883) = praec.

DICYCLOPHORA, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 88 = **Fyoncocyla**, Lindl. (Umbellif.).
persica, Boiss. l. c. 91.—Persia.

DICYMBE, Spruce, ex Benth. & Hook. f. Gen. i. 504 (1865). *LEGUMINOSAE*, Benth. & Hook. f. i. c.
corymbosa, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 60.—Venezuela.

DICYPELLIUM, Nees, Progr. 14 (1833); Laurin. Expos. 14 (1836). *LAURINEAE*, Benth. & Hook. f. iii. 159.
caryophyllatum, *Nees, Syst. Laurin.* 344.—Am. trop.

DICYRTA, Regel, in Flora, xxxii. (1849) 181. *GESNERACEAE*, Benth. & Hook. f. ii. 1000.
candida, *Hanst. & Klotzsch, in Linnaea*, xxvii. (1854) 714.—Guatemala.

DICYRTA:—
parviflora, Seem. Bot. Voy. Herald, 326. t. 69 = Anetanthus parviflora.
Warszewiczioides, *Regel, in Flora*, xxxii. (1849) 181.—Guatemala.

DIDACTYLE, Lindl. Fol. Orch. Didact. (1852) = **Bulbophyllum**, Thou. (Orchid.).
antennifera, Lindl. Fol. Orch. Didact. 2 = B. geniculatum.
atropurpurea, Rodrig. Orch. Nov. ii. 125 = B. atropurpureum.
bidentata, Rodrig. l. c. 127.—Bras.
candeguelleana, Rodrig. l. c. 126.—Bras.
Clausseni, Lindl. Fol. Orch. Didact. 3 = B. Clausseni.
exaltata, Lindl. l. c. 2 = B. exaltatum.
galeata, Rodrig. Orch. Nov. ii. 129.—Bras.
gladiata, Lindl. Fol. Orch. Didact. 1 = B. meridense.
glutinosa, Rodrig. Orch. Nov. ii. 126 —Bras.
granulosa, Rodrig. l. c. 125 = B. granulosum.
lacinulata, Rodrig. l. c. t. 43.—Bras.
meridensis, Lindl. Fol. Orch. Didact. 3 = B meridense.
nemorosa, Rodrig. Orch. Nov. i. 43.—Bras.
ochracea, Rodrig. l. c. iii. 127.—Bras.
plumosa, Rodrig. l. c. i. 44.—Bras.
quadricolor, Rodrig. l. c. ii. 128.—Bras.
Regnellii, Rodrig. l. c. 128 = B. Regnellii.
tripetala, Lindl. Fol. Orch. Didact. 2 = B. tripetalum.
Weddelii, Lindl. l. c. = B. Weddelii.

DIDACTYLON, Zoll. & Moritzl, Syst. Verz. Zoll. 99 (1845–46) = **Dimeria**, R. Br. (Gramin.).
ramosum, Zoll. & Moritzl, l. c. 100 = Dim. ornithopoda.
simplex, Zoll. & Moritzl, l. c. = Dim. ornithopoda.

DIDASTE, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 472 (1862) = **Acrosanthes**, Eckl. & Zeyh. (Ficoid.).
decandra, E. Mey. ex Harv. & Sond. l. c. = A. angustifolia.
icosandra, E. Mey. ex Harv. & Sond. l. c. 473 = A. teretifolia.
pentandra, E. Mey. ex Harv. & Sond. l. c. 472 = A. humifosa.

DIDELOTIA, Baill. Adansonia, v. (1865) 367. t. 8. *LEGUMINOSAE*, Benth. & Hook. f. i. 1003.
africana, *Baill. l. c.*—Afr. trop.

DIDELTA, L'Hérit. Stirp. Nov. 55. t. 28 (1785). *COMPOSITAE*, Benth. & Hook. f. ii. 461.
BRETEUILLIA, Buchoz, ex DC. Prod. vii. 603, in syn. (1837).
CHORISTZA, Thunb. Prod. Pl. Cap. praef. et 163 (1800); Nov. Gen. 179 (1801).
FAVONIUM, Gaertn. Fruct. ii. 431. t. 174 (1791).
annua, *Less. Syn. Comp.* 62 (Berkheyae sp. !).—Afr. austr.
capensis, Lam. ex DC. Prod. vii. 603 = carnosa.
carnosa, *Ait. Hort. Kew* ed. I. iii 256.—Afr. austr.
cernua, *Less. Syn. Comp.* 61 (Berkheyae sp. !).—Afr. austr.
obtusifolia, Cass. in Dict. Sc. Nat. xxix. 461 = spinosa.
spinosa, *Ait. Hort. Kew.* ed. I. iii. 256.—Afr. austr.
tetragonaefolia, L'Hérit. Stirp. Nov. 56, 178. t. 28 = carnosa.
tomentosa, *Less. Syn. Comp.* 61.—Austral.

DIDEROTA, Comm. ex A. DC. in DC. Prod. viii. 356 (1844) = **Ochrosia**, Juss. (Apocyn.).
amphicarpa, Comm. ex A. DC. l. c. = O. borbonica.

DIDEROTIA, Baill. Adansonia, i. (1860–61) 274 = **Alchornea**, Sw. (Euphorb.).
multispicata, Baill. l. c.—Afr. trop.

DIDESMIS, Lond. Hort. Brit. 266 (1830) = seq.

DIDESMUS, Lesv. Journ. Bot. iii. (1814) 160. t. 25 = **Eupistrum**, Tourn. (Crucifer.).
aegyptiacus, Desv. l. c. = R. aegyptiacum.
aegyptius, DC. Syst. ii. 659 = praec.
bipinnatus, DC. l. c. 659 = R. bipinnatum.
myagroides, Desv. Journ. Bot. iii. (1814) 160 = Erucaria aleppica.
pandurifornis, Stocks, in Proc. Linn. Soc. i. (1848) 267.—Ind. or.
pilosus, Stend. Nom. ed. II. i. 505, sphalm. vide *Didiscus pilosus*.
pinnatus, DC. Syst. ii. 659 = Erucaria aleppica ?

DIDESMUS:—
rostratus, Boiss. Fl. Orient. i. 405.—Palaestin.
tenuifolius, DC. Syst. ii. 659.—Archip. Graec.

DIDIEREA, Baill. in Bull. Soc. Linn. Par. i. (1880) 238. *ANACARDIACEAE*?
madagascariensis, *Baill. l. c.*—Madag.

DIDIMERIA, Lindl. in Mitch. Three Exped. Austral ed. II. ii. 198 (1839) = **Correa**, Sm. (Rutac.).
aemula, Lindl. l. c. = C. aemula.

DIDIPLIS, Rafin. Atl. Journ. (1833) 177. *THRARIEAE*, Benth. & Hook. f. i. 777.
HYPORRICHIA, M. A. Curt. in Torr. & Gray, Fl. N. Am. i. 479 (1840).
PTILINA, Nutt. ex Torr. & Gray, l. c. (1840).
diandra, Wood, Bot. and Flor. 124 = linearis.
linearis, *Rafin. Atl. Journ.* (1833) 177.—Am. bor.

DIDISCUS, DC. in Bot. Mag. t. 2876 (1828) = **Trachymene**, Rudge (Umbellif.).
albiflorus, DC. Prod. iv. 72 = T. incisa.
anisocarpus, F. Muell. in Proc. Roy. Soc. V. D. Land, iii. (1855) 238 = T. australis.
austro-caledonicus. Brongn. & Gris, in Bull. Soc. Bot. Fr. vii. (1860) 372; in Ann. Sc. Nat. Sér. V. iii. (1865) 235 = T. austro-caledonica.
caeruleus, DC. in Hook. Bot. Mag. t. 2875 = T. caerulea.
cyaneus, DC. Mém. Ombell. 28. t. 4 = T. caerulea.
cyanopetalus, F. Muell. Fragm. ix. 46 = T. cyanopetalus
eriocarpus, F. Muell. l. c. = T. eriocarpa.
glandulosus, F. Muell. in Proc. Roy. Soc. V. D. Land. iii. (1855) 238 = T. glandulosa.
glaucifolius, F. Muell. in Linnaea, xxv. (1852) 395 = T. glaucifolia.
grandis, F. Muell. in Proc. Roy. Soc. V. D. Land, iii. (1855) 238 = T. australis.
hemicarpus, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1857) 491 = T. hemicarpa.
humilis, Hook. f. Ic. Pl. t. 804 = T. humilis.
incisus, Hook. Bot Mag. sub t. 2875 = T. incisa.
pilosus, Benth. in Enum. Pl. Hoog. 64 = T. australis.
procumbens, F. Muell. in Proc. Roy. Soc. V. D. Land, iii. (1855) 237 = T. procumbens.
pusillus, F. Muell. Fragm. ix. 47 = T. pilosa.
setulosus, F. Muell. in Proc. Roy. Soc. V. D. Land, iii. (1855) 238 = T. hemicarpa.
villosus, F. Muell. l. c. = T. villosa.

DIDISSANDRA, C. B. Clarke, in DC. Monog. Phan. v. 65 (1888). *CYRTANDREAE*.
elongata, *C. B. Clarke, l. c.* 67.—Sumatra.
frutescens, *C. B. Clarke, l. c.*—Malaya.
lanuginosa, *C. B. Clarke, l. c.* 66.—Reg. Himal.
Miberi, *Franch. in Bull. Soc. Linn. Par.* i. (1885) 450.—China.
ornata, *C. B. Clarke, in DC. Monog. Phan.* v. 69.—Borneo.
rufa, *C. B. Clarke, l. c.* 70.—Borneo.
rufa, *King, in Hook. Ic. Pl.* t. 1437.—Tibet. or.
Sprengelii, *C. B. Clarke, in DC. Monog. Phan.* v. 68.—Sumatra.
trifora, *C. B. Clarke, l. c.* 69.—Borneo.

DIDOTHION, Rafin. Fl. Tellur. iv. 39 (1836) = **Epidendrum**, Linn. (Orchid.).
clavatum, Rafin. l. c. = E. clavatum.

DIDYMAEA, Hook. f. in Benth. & Hook. f. *Gen.* iii. 150 (1879). *RUBIACEAE*, Benth. & Hook. f. i. c.
mexicana, *Hook. f. l. c.*—Mexic.

DIDYMANDRA, Wild. Sp. Pl. iv. 971 (1806) = **Lacistema**, Sw.
dentata, Spreng. Syst. i. 94.—Bras.
purpurea, Willd. Sp. Pl. iv. 971 = L. purpureum.

DIDYMANTHUS, Endl. Nov. Stirp. Dec. 7 (1839). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 60. Moq. *Endl. l. c.* 8.—Austral.

DIDYMANTHUS, Klotzsch, ex Meisn. in DC. Prod. 486 (1857 !) = **Bupleurum**, Salisb. (Prosoc.).
incanus, Klotzsch, l. c. (= *Adenostephanus incanus*).—Bras.
Sellowii, Klotzsch, l. c. (= *Adenostephanus Sellowii*).—Bras.

DIDYMELES, Thou. Hist. Vég. Afr. Austr. 9. t. 1 (1805) Gen. dub. affin.
excelsa, Baill. Hist. Pl. iv. 368 = madagascariensis.
madagascariensis, *Willd. Sp. Pl.* iv. 649.—Madag.

DIDYMERIA, Lindl. ex Ann. Sc. Nat. Sér. II. xv. (1841) 59 = DIDIMERIA, Lindl. = **Correa**, Sm. (Rutac.).

DIDYMOCARPUS, Wall. in Edinb. Phil. Journ. i. (1819) 378. *GESNERACEAE*, Benth. & Hook. f. ii. 1021.
HENCKELIA, Spreng. Anleit. ii. 402 (1817).
LOXOCARPUS, R. Br. in Benn. Pl. Jav. Rar. 120 (1838).
ROTTLERA, Vahl, Enum. i. 88 (1805).
acuminata, *R. Br. in Benn. Pl. Jav. Rar.* 118.—Reg. Himal.
albicaulis, *C. B. Clarke, in DC. Monog. Phan.* v. 78. —Reg. Himal.
alternifolia, *C. B. Clarke, l. c.* 106.—Borneo.
amœna, *C. B. Clarke, l. c.* 87.—Borneo.
Andersonii, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 62. —Reg. Himal.
angovensis, Boj. ex DC. Prod. ix. 271 = Streptocarpus angovensis.
angustifolia, *C. B. Clarke, in DC. Monog. Phan.* v 89. —Borneo.
aristata, *Vill. in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 150.—Ins. Philipp.
aromatica, D. Don, Prod. Fl. Nep. 123 = subalternans.
aromatica, Griff. Ic. Pl. Asiat. t. 440. f. 4 = pulchra.
aromatica, *Wall. Cat.* n. 785.—Reg. Himal.
aromatica, Wall. Pl. As. Rar. t. 141. f. 4–7 = macrophylla.
aurantiaca, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 60. —Reg. Himal.
Auricula, *S. Moore, in Journ. Bot.* xiii. (1875) 229.— China.
bancana, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1873) 438.—Ins. Banca.
barbata, Jack, in Trans. Linn. Soc. xiv. (1825) 88 = Chirita Horsfieldii.
barbinervia, *C. B. Clarke, in DC. Monog. Phan.* v. 85. —Burma.
Beccarii, *C. B. Clarke, l. c.* 89.—Borneo.
bifolia, Jack, ex DC. Prod. ix. 268 = Chirita bifolia.
Blancol, *Hassk. in Flora*, xlvii. (1864) 55.—Ins. Philipp.
bullata, *C. B. Clarke, in DC. Monog. Phan.* v. 92.— Borneo.
cinerea, *D. Don, Prod. Fl. Nep.* 122.—Nepal.
conicapsularis, *C. B. Clarke, in DC. Monog. Phan.* v. 100.—Borneo.
corchorifolia, *Wall. Cat.* n. 792 —Ins. Penang.
cordata, *Wall. l. c.* 781 —Penang.
corniculata, *Jack, in Malay. Misc* i. (1820) v 4.— Sumatra.
crinita, *Jack, l. c.* 1.—Malaya.
cristata, Dalz. in Hook. Kew Journ. iii. (1851) 225 = Chirita hamosa.
cyathiphora, Reichb. f. & Zoll. in Teijsm. & Binn. Cat. Hort. Bog. (1866) 156 = Chirita cærulea.
demissa, *Hance, in Journ. Bot.* xxi. (1883) 166.— China.
detergibilis, *C. B. Clarke, in DC. Monog. Phan.* v. 107.—Malaya.
elegans, *C. B. Clarke, l. c.* 88.—Borneo.
elongata, Jack, in Trans. Linn. Soc. xiv. (1825) 87 = Didissandra elongata.
fasciculariis, C. B. Clarke, in DC. Monog. v. Index, 298 = follicularis.
flocosa, *Thw. Enum. Pl. Zeyl.* 207.—Zeylan.
follicularis, *C. B. Clarke, in DC. Monog. Phan.* v. 98. —Borneo.
frutescens, *Jack, in Malay. Misc.* i. (1820) v. 5.— Malaya.
gracilipes, *C. B. Clarke, in DC. Monog. Phan.* v. 97. —Borneo.
grandiflora, Dietr. ex Steud. Nom. ed. II. i. 506 = Chirita grandiflora.
grandifolia, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. III. 889 = Chirita macrophylla.
Griffithii, Wight, Illustr. ii. 132 = Loxostigma Griffithii.
hamosa, Wall. Cat. n. 788 = Chirita hamosa.
helicteroides, Wall. l. c. 789 = Boea Wallichii.
Hilsenbergii, Boj. ex DC. Prod. ix. 268, 270 = Streptocarpus Hilsenbergii.
Hookeri, C. B. Clarke, Comm. et Cyrt. Beng. t. 54.— Reg. Himal.

DIDYMOCARPUS:—
Humboldtiana, *Gardn. in Calc Journ. Nat. Hist.* vi. (1846) 477.—Zeylan.
incana, *Benth. & Hook. f. ex C. B. Clarke, in DC. Monog. Phan.* v. 93.—Ins. Penang.
Kompsobœa, *C. B. Clarke, l. c.* 92.—Borneo.
Kurzii, C. B. Clarke, Comm. et Cyrt. Beng. t. 66 = Chirita Kurzii.
lanceolata, *C. B. Clarke, in DC. Monog. Phan.* v. 89. —Borneo.
lanuginosa, Maxim. in Bull. Acad. Pétersb. xix. (1874) 535 = Oreocharis Maximowiczii.
lanuginosa, Wall. Cat. n. 791 = Didissandra lanuginosa.
lanuginosa, Wight, ex C. B. Clarke, in DC. Monog. Phan. v. 102 = Rottleriana.
Lawesii, F. Muell. in Wing, South Sc. Record, ii (1882) 229 = Boea Treubii.
leptocalyx, *C. B. Clarke, in DC. Monog. Phan.* v 91. —Borneo.
leucocalyx, *C. B. Clarke, in Hook. f. Fl. Brit. Ind* iv. 348.—Reg. Himal.
longipes, *C. B. Clarke, in DC. Monog. Phan.* v. 86.— Malacca.
longipetiolata, *Gardn. in Calc. Journ. Nat. Hist* vi. (1846) 475.—Zeylan.
lyrata, *Wight, Ic.* iv. t. 1350.—Ind or.
macrophylla, Hook. f. & Thoms. ex C. B. Clarke, in DC. Monog. Phan. v. 74 = Andersoni.
macrophylla, *Wall. ex D. Don, Prod. Fl. Nep.* 122; *et Cat.* n. 1813.—Reg. Himal.
macrophylla, Wall. Cat. n. 784; Royle, Illustr. Bot. Himal. 294. t. 70. f. 1 = pedicellata.
madagascarica, *C. B. Clarke, in DC. Monog. Phan.* v. 108.—Madag.
marginata, *C. B. Clarke, l. c.* 96.—Malacca.
membranacea, Bedd. Ic. Pl. Ind. Or. i. 89. t. 176 = missionis.
Minahassæ, *H. Forbes, in Journ. Linn Soc.* xix. (1882) 298.—Ins. Celebes.
missionis, *Wall. Cat.* n. 6396.—Ind. or.
mollis, *Wall. l. c.* n. 790.—Burma.
monophylla, Wall. ex C. B. Clarke, in DC. Monog. Phan. v. 72 = macrophylla.
Mortoni, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 65.— Reg. Himal.
multiflora, Wall. Cat. n. 793 = Boea multiflora.
nervosa, *C. B. Clarke, in DC. Monog. Phan.* v. 95.— Borneo.
oblonga, *Wall. Pl. As. Rar.* ii. 34. t. 140.—Reg. Himal.
obtusa, Wall. Cat. n. 786 = cinerea.
Oreocharis, Hance, in Ann. Sc. Nat. Sér. V. (1866) v. 230 = Oreocharis Benthami.
ovalifolia, *Wight, Ic.* iv. t. 1351.—Ind. or.
Paraboea, *C. B. Clarke, in DC. Monog. Phan.* v. 106. —Borneo.
paucinervia, *C. B. Clarke, l. c.* 83.—Burma.
pedicellata, *R. Br. in Benn. Pl. Jav. Rar.* 118.— Reg. Himal.
petiolaris, *C. B. Clarke, in DC. Monog. Phan.* v. 109. —Borneo.
platycalyx, *C. B. Clarke, l. c.* 81.—Burma.
platypus, *C. B. Clarke, l. c.* 94.—Malacca.
pœcata, D. Don, Prod. Fl. Nep. 122 = macrophylla.
podocarpa, *C. B. Clarke, in DC. Monog. Phan.* v. 76. —Reg. Himal.
primulæfolia, D. Don, Prod. Fl. Nep. 123 = aromatica, subalternans.
primulæfolia, Gardn. in Calc. Journ Nat. Hist. vi. (1846) 473 = Humboldtiana.
primuloides, Maxim. in Bull. Acad. Pétersb. xix. (1874) 536 = Oreocharis primuloides.
products, *C. B. Clarke, in DC. Monog. Phan.* v. 96.— Sumatra.
pulchra, *C. B. Clarke, l. c.* 79.—Reg. Himal.
punctata, *C. B. Clarke, l. c.* 90.—Borneo.
pundunna, *Wall. Cat.* n. 777.—Reg. Himal.
pygmæa, *C. B. Clarke, in DC. Monog. Phan.* v. 82.— Ind. or.
racemosa, *Jack, in Trans. Linn. Soc.* xiv. (1825) 84. —Sumatra.
repens, *Bedd. Ic. Pl. Ind. Or.* i. 24. t. 120.— Ind. or.
reptans, *Jack, in Malay. Misc.* i. (1820) v. 3.— Malacc.
reticulosa, *C. B. Clarke, in DC. Monog. Phan.* v. 90.— Borneo.

DIDYMOCARPUS:—
Rexii, Bowie, ex Hook. Exot. Fl. iii. t. 227 = Streptocarpus Rexii.
Rottleriana, *Wall. Cat.* n. 778.—Ind. or.
rufescens, *C. B. Clarke, in DC. Monog. Phan.* v. 99.— Borneo.
rufipes, *C. B. Clarke, l. c.* 84.—Burma.
scabrinervia, *C. B. Clarke, l. c.* 91.—Borneo.
Scheffleri, *H. Forbes, in Journ. Linn. Soc.* xix. (1882) 298.—Borneo.
semitorta, *C. B. Clarke, in DC. Monog. Phan.* v. 99.— Malacca.
serrata, *R. Br. in Benn. Pl. Jav. Rar.* 119.—Sumatra.
speciosa, Wall. ex C. B. Clarke, in DC. Monog. Phan. v. 114 = Chirita pumila.
Sprengelii, Korth. ex C. B. Clarke, l. c. 68 = Didissandra Sprengelii.
subalternans, C. B Clarke, Comm. et Cyrt. Beng. t 55 = pulchra.
subalternans, *Wall. Cat.* n. 782 —Reg. Himal.
teres, *C. B Clarke, in DC. Monog. Phan.* v 88.— Borneo
ternata, *Miq. Fl. Ind. Bat Suppl.* 564.—Sumatra.
tomentosa, *Wight, Ic.* iv t. 1349.—Ind. or.
verbenaefiora, *C. B. Clarke, in DC. Monog. Phan.* v. 99. —Borneo.
verticillata, Wall. Cat. n. 783 = oblonga.
vestita, *Baker, in Journ. Linn. Soc.* xxi. (1885) 427.— Madag.
villosa, C. B Clarke, Comm. et Cyrt. Beng. t. 59 = leucocalyx.
villosa, *D. Don, Prod. Fl. Nep* 123.—Reg. Himal.
violoides, *C. B. Clarke, in DC. Monog. Phan.* v. 97.— Borneo.
zeylanica, *R. Br. in Benn. Pl. Jav. Rar.* 119.— Zeylan.

DIDYMOCHAETA, Steud. Syn. Pl. Gram. 185 (1855) = **Deyeuxia**, Clar. (Gramin.).
australis, Steud. l. c. = Deyeuxia plebeia.
chilensis, Steud. l. c. = Deyeuxia chilensis.

DIDYMOCHETON, Blume, Bijdr. 177 (1825) = **Dysoxylum**, Blume (Meliac.).
Gaudichaudianum, A. Juss in Mém. Mus. Par. xix. (1830) 281 = Dys. Gaudichaudianum.
Leschenaultianum, A Juss. l. c.—Java.
littorale, Hassk in Hoev. & De Vriese, Tijdschr. x. (1843) 138.—Java.
nutans, Blume, Bijdr. 177 = Dys. nutans.
pruriens, Zipp. ex Span. in Linnaea, xv. (1841) 183, nomen.—Ins Timor
Richii, A. Gray, Bot. U. St. Expl. Exped. i. 239. t. 20 = Dys. Richii.
urens, Teijsm. & Binn. Cat. Hort. Bog. 391, nomen.— Ins. Batjan.

DIDYMOCHLAMYA, Hook. f. Ic. Pl. t. 1122 (1872). *RUBIACEAE*, Benth. & Hook. f. ii. 67.
Whitei, *Hook. f. l. c.*—N. Granat.

DIDYMOCOCCUS, Blume, Rumphia, iii. 103, in obs. (1847) = **Sapindus**, Linn. (Sapind.).
Donura, Blume, l. c. = S. Donura.
verticillatus, Blume, l. c. = S. Donura.

DIDYMODOXA, E. Mey. ex Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 212, in syn. = **Australina**, Gandich. (Urticac.).
acuminata, Wedd. in Arch. Mus. Par. viii. (1855–56) 549 = A. acuminata.
cuneata, Wedd. in DC. Prod. xvi. i. 236* = A. acuminata.
debilis, E. Mey. ex Wedd. in Ann. Sc. Nat. Sér. IV. i. (1855) 212 = A. capensis.
integrifolia, Wedd. in Arch. Mus. Par. viii. (1855–56) 549 = A Integrifolia.

DIDYMOGYNE, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 207 = **Droguetia**, Gandich. (Urticac.).
abyssinica, Wedd. l. c. = Drog. diffusa.
boehmerioides, Wedd. in DC. Prod. xvi. i. 205 = Boehmeria Didymogyne.

DIDYMOMELES, Spreng. Syst. iii. 899 (1826) = **Didymeles**, Thou. (Inc. sed.).

DIDYMONEMA, Presl, Didym. Nov. Pl. Gen. (1829); et Symb. Bot. i. 5. t. 8 (1830) = **Gahnia**, Forst. (Cyperac.).
filifolia, Presl, ll. cc. = G. tristis.

DIDYMOPANAX, Decne. & Planch. in Rev. Hortic. (1854) 106. *ARALIACEAE*, Benth. & Hook. f. i. 939.
argusticanum, *March. in Mart. Fl. Bras.* xi. 1. 241.—Bras.
argyrophyllum, *Decne. & Planch. in Rev. Hortic.* (1854) 109, nomen.—N. Granat.
Burchellii, *Serm. in Journ. Bot.* vi. (1868) 133. —Bras.
calophyllum, *Decne. & Planch. ex March. in Mart. Fl. Bras.* xi. 1. 240.—Bras.
calvum, *Decne. & Planch. in Rev. Hortic.* (1854) 109. —Bras.
chrysophyllum, Decne. & Planch. l. c. = Morototoni.
Clausenianum, *Decne. & Planch. ex March. in Mart. Fl. Bras.* xi. 1. 239.—Bras.
emarginatum, *Seem. in Illustr. Hortic.* (1854) 153.—Bras.
Gardneri, *Seem. Journ. Bot.* vi. (1868) 132.—Bras.
Houlletii, *Linden, in Gard. Chron.* (1869) 587.—Hab.?
longipetiolatum, *March. in Mart. Fl. Bras.* xi. 1. 234. —Bras.
lucumoides, *Decne. & Planch. in Rev. Hortic.* (1854) 109, nomen.—Bras.
macrocarpum, *Seem. in Journ. Bot.* vi. (1868) 132.—Bras.
marginatum, Decne. & Planch. in Rev. Hortic. (1854) 109, nomen = macrocarpum.
micranthum, *March. in Mart. Fl. Bras.* xi. 1. 239.—Bras.
Morototoni, *Decne. & Planch. in Rev. Hortic.* (1854) 109.—Am. austr.
pachycarpum, *March. in Mart. Fl. Bras.* xi. 1. 236.—Bras.
papyriferum, Koch, Wochenschr. ii. (1859) 69 = Fatsia papyrifera.
parviflorum, Decne. & Planch. in Rev. Hortic. (1854) 109 = vinosum.
Poeppigii, Decne. & Planch. l. c. = Morototoni.
Selloi, *March. in Mart. Fl. Bras.* xi. 1. 236.—Bras.
sericeum, Decne. & Planch. in Rev. Hortic. (1854) 109, nomen = vinosum.
speciosum, Decne. & Planch. l. c. = Morototoni.
splendidum, *Planch. ex Linden, Cat.* n. 10 (1855) 8.—N. Granat.
Spruceanum, *Seem. Journ. Bot.* vi. (1868) 132.—Bras.
tomentosum, Benth. & Hook. f. Gen. ii. 939 = lucumoides.
vinosum, *March. in Mart. Fl. Bras.* xi. 1. 238.—Bras.

DIDYMOPELTA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 669 = **Astragalus**, Tourn. (Legumin.).
turkestanica, Regel & Schmalh. l. c. = A. Schmalhausenii.

DIDYMOPHYSA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1841) 179. *CRUCIFERAE*, Benth. & Hook. f. i. 91.
Aucheri, *Boiss. l. c.*—Persia.
Fedtschenkoana, *Regel, Pl. Nov. Fedsch.* 8.—Turkestan.

DIDYMOPLEXIS, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 363. t. 17. *ORCHIDEAE*, Benth. & Hook. f. iii. 616, 1236.
APETALON, Wight, Ic. v. 22. t. 1756 (1852).
EPIPHANES, Reichb. f. in Seem. Fl. Vit. 295 (1868).
LEUCORCHIS, Blume, Mus. Bot. Lugd. Bat. i. 31 (1849).
micradenia, *Hemsl. in Journ. Linn. Soc.* xx. (1883) 311.—Ins. Fiji.
pallens, *Griff. in Calc. Journ. Nat. Hist.* iv. (1844) t. 17.—Ind. or.

DIDYMOSPERMA, H. Wendl. & Drude, in Kerch. Palm. 243 (1878). *PALMAE*, Benth. & Hook. f. iii. 917.
BLANCOA, Blume, Rumphia, ii. 128 (1836).
caudatum, *H. Wendl. & Drude, l. c.*—Siam.
Horsfieldii, *H. Wendl. & Drude, l. c.*—Java.
nasum, *H. Wendl. & Drude, l. c.*—Ind. or.
porphyrocarpon, *H. Wendl. & Drude, l. c.*—Java.
Reinwardtianum, *H. Wendl. & Drude, l. c.*—Java.
tremulum, *H. Wendl. & Drude, l. c.*—Siam.

DIDYMOTHECA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 278. *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 83.
Drummondii, Moq. in DC. Prod. xiii. 11. 87 = thesioides.
floccosa, F. Muell. Fragm. i. 202; Pl. Vict. i. 198. t. Suppl. 92 = Gyrostemon Cyclotheca.
thesioides, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 279.—Austral.
veroniciformis, F. Muell. in Linnaea, xxv. (1852) 488 = thesioides.

DIDYMOTOCA, E. Mey. Zwei Pfl. Docum. 178 (1848), adopt. diag. = **Australina**, Gaudich. (Urticac.).
debilis, E. Mey. l. c. = A. capensis.

DIDYPLOSANDRA, Wight, Ic. iv. pars iv. 8. t. 1515 (1850) = **Strobilanthes**, Blume (Acanth.) (S. luridus).

DIECTOMIS, Beauv. Agrost. 182. t. 28. f. 5 (1812); H. B. & K. Nov. Gen. et Sp. i. 198. t. 64 (1815) = **Andropogon**, Linn. (Gramin.).
angustata, J. & C. Presl, Kel. Haenk. i. 338 = A. apricus.
fasciculata, Beauv. Agrost. 153. t. 23 = A. leptocomus.
fastigiata, H. B. & K. Nov. Gen. et Sp. i. 198 = A. laniger.
laxa, Nees, in Mart. Fl. Bras. ii. 340 = A. apricus.

DIEFFENBACHIA, Schott, in Wien. Zeitschr. iii. (1829) 830; Meliet. i. 20 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 986.
SIEBUHNIA, Rafin. Fl. Tellur. iii. 66 (1836).
allionii, Regel, in Gartenfl. (1870) t. 672 = humilis.
antiquensis, Linden & André, in Illustr. Hortic. (1875) t. 192 = picta.
Barraquiniana, Versch. & Lem. l. c. xi. (1864) t. 387 = D. Seguine.
Baumannii, Hort. ex Engl. in DC. Monog. Phan. ii. 447 = D. Seguine.
Bauseri, in Gartenfl. (1878) 49 = picta.
Bowmannii, *Carr. in Rev. Hortic.* (1872) 198.—Japon.
brasiliensis, Veitch, Cat. (1875) 12 = picta.
caperata, Schott, Syn. Aroid. 130 = D. Seguine.
consobrina, Schott, l. c. 131 = D. Seguine.
conspurcata, Schott, in Seem. Journ. Bot. ii. (1864) 52 = D. Seguine.
costata, Klotzsch, ex Schott, Syn. Aroid. 128 = macrophylla.
daguensis, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 288. —N. Guinea.
decora, Hort. Versch. ex Engl. in DC. Monog. Phan. ii. 446 = D. Seguine.
Enideri, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 284.—N. Guinea.
gigantea, Versch. in Illustr. Hortic. (1866) t. 470–471 = D. Seguine.
Goldieriana, Schott, in Oestr. Bot. Zeitschr. (1858) 287 = D. Seguine.
grandis, Hort. ex Engl. in DC. Monog. Phan. ii. 447 = D. Seguine.
humilis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 90.—Bras.
imperialis, *Linden & André, in Illustr. Hortic.* (1871) t. 85.—Peruv.
irrorata, Schott, Prod. Aroid. 384 = D. Seguine.
Jennanni, *Veitch, ex Regel, Gartenfl.* (1884) 365.—Hab.?
lancifolia, Linden & André, in Illustr. Hortic. xxi. (1874) t. 167 = picta.
latimaculata, *Linden & André, l. c.* xx. (1873) 8 t. 219.—Bras.
Leopoldi, Hort. Bull, ex Engl. in DC. Monog. Phan. ii. 446 = D. Seguine.
lineata, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1858) 14 = D. Seguine.
lingulata, Schott, Prod. Aroid. 384 = D. Seguine.
liturata, Schott, in Oestr. Bot. Wochenbl. (1853) 68 = D. Seguine.
macrophylla, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 90. —Peruv.
maculata, Sweet, Hort. Brit. ed. III. 632 = D. Seguine.
magnifica, *Linden & Rodigas, in Illustr. Hortic.* xxx. (1883) t. 493.—Venezuela.
membranulifera, *Regel, Ind. Sem. Hort. Petrop.* (1861) 37.—Bras.
mirabilis, Versch. ex Engl. in DC. Monog. Phan. ii. 446 = picta.
neglecta, Schott, in Bonplandia, vii. (1859) 20 = D. Seguine.

DIEFFENBACHIA—
nobilis, Versch. ex Engl. in DC. Monog. Phan. ii. 446 = D. Seguine.
oblique, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 90.—Bras
Oerstedtii, *Schott, in Oestr. Bot. Zeitschr.* (1858) 179 —Costa Rica.
Parlatorii, Linden & André, in Illustr. Hortic. xxiv. (1877) 152. t. 291 = imperialis.
picta, *Schott, in Oestr. Bot. Wochenbl.* (1852) 68.—Bras
Plumieri, Schott, l. c. 69 = D. Seguine.
Poeppigii, Schott, Syn. Aroid. 130 = D. Seguine.
robusta, C. Koch, Ind. Sem. Hort. Berol. (1858) 15 = D. Seguine.
robusta, Schott, Oestr. Bot. Wochenbl. (1854) 65 = macrophylla.
Seguine, *Schott, Meliet.* i. 20.—Am. trop.
Shuttleworthiana, Regel, in Gartenfl. (1878) 313 = picta.
Spruceana, Schott, in Bonplandia, vii. (1859) 20 = humilis.
variegata, Hort. ex Engl. in DC. Monog. Phan. ii. 446 = D. Seguine.
Ventenatiana, Schott, in Bonplandia, vii. (1859) 20 = D. Seguine.
Verschaffeltii, Hort. ex Engl. in DC. Monog. Phan. ii 446 = D. Seguine.
Wallisii, Linden, in Illustr. Hortic. (1870) t. 11 = D Seguine.
Weirii, *Berk. in Journ. Hort. Soc.* N. S. i. (1866) 201.—Bras.
Wendlandii, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 179.—Am. centr.

DIELYTRA, Dam. & Schlecht. in Linnaea, i. (1826) 556.—Cf. Spreng. Syst. iii. 102 (1826) = **Dicentra**, Bernh. (Papav.).
bracteosa, G. Don, Gen. Syst. i. 140 (= *Dielytra bracteosa*).—Am. bor.?
canadensis, G. Don, l. c. = Dicentra canadensis.
chrysantha, Hook. & Arn. Bot. Beech. Voy. 320 = Dicentra chrysantha.
Cucullaria, G. Don, Gen. Syst. i. 140 = Dicentra Cucullaria.
eximia, Don, l. c. = Dicentra eximia.
eximia, Hook. Fl. Bor. Am. i. 35 = Dicentra formosa.
formosa, G. Don, Gen. Syst. i. 140 = Dicentra formosa.
lachenaliaeflora, Cham. & Schlecht. in Linnaea, i. (1826) 556 = Dicentra lachenaliaeflora.
saccata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 67 = Dicentra formosa.
scandens, G. Don, Gen. Syst. i. 140 = Dicentra scandens.
spectabilis, G. Don, l. c. = Dicentra spectabilis.
tenuifolia, Cham. & Schlecht. in Linnaea, i. (1826) 556 = Dicentra lachenaliaeflora.

DIEMENIA, Korth. in Nederl. Kruidk. Arch. iii. (1855) 368 = **Trichocarya**, Miq. (Rosac.).
racemosa, Korth. l. c. = T. racemosa.

DIENA, Reichb. Consp. 69 (1828) = seq.

DIENIA, Lindl. Bot. Reg. sub t. 825 (1824) = **Microstylis**, Nutt. (Orchid.).
calycina, Lindl. Gen. et Sp. Orch. 28 = M. macrostachya.
carinata, Reichb. f. in Bonplandia, iii. (1855) 228 = M. carinata.
congesta, Lindl. Bot. Reg. sub t. 825 = M. congesta.
cordata, Lindl. l. c. (1838) Misc. 78 = M. cordata.
crispata, Lindl. in Ann. & Mag. Nat. Hist. xv. (1845) 338 = M. crispata.
cylindrostachya, Lindl. in Wall. Cat. n. 1934 = M. cylindrostachya.
fusca, Lindl. Gen. et Sp. Orch. 22 = M. congesta.
Gminiai, Lindl. l. c. 28 = M. monophylla.
majoranthemifolia, Reichb. f. in Linnaea, xix. (1847) 809 = M. fastigiata.
monticola, Lindl. in Wall. Cat. n. 1935.—Reg. Himal.
Myurus, Lindl. Gen. et Sp. Orch. 23 = M. Myurus.

DIERAMA, C. Koch, in Ind. Sem. Hort. Berol. App. (1854) 10; et in Walp. Ann. vi. 43 (1861). *IRIDEAE*, Benth. & Hook. f. iii. 703.
DICRAMA, Klatt, in Linnaea, xxxii. (1863) 751.
cupuliflorum, *Klatt, in Ducben, Reisen Ost-Afrika*, iii. 73. t. 3.—Afr. trop.

DIERAMA:—
ensifolium, *C. Koch & Bouché, Ind. Sem. Hort. Berol. App.* (1854) 10.—Afr. austr.
igneum, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 288.—Afr. austr.
pendulum, Baker, in Journ. Linn. Soc. xvi. (1877) 99 = ensifolium.
pulcherrimum, *Baker, l. c.*—Afr. austr.

DIERBACHIA, Spreng. Syst. i. 512, 676 (1825) = **Dunalia**, H. B. & K. (Solanac.).
solanacea, Spreng. l. c. 676 = Dunalia solanacea.

DIERVILLA, Tourn. ex Linn. Syst. ed. I (1735). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 6.
CALYPTROMYRUM, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 46) (1847).
CALYSPHYRUM, Bunge, Enum. Pl. Chin. Bor. 38 (1833).
WAGNERIA, Lem. Illustr. Hortic. iv. (1857) sub t. 115.
WEIGELA, Thunb. in Vet. Acad. Handl. Stockh. (1780) 187. t. 5.
acadiensis, Duham. Arb. i. t. 87 = trifida.
amabilis, Carr. in Rev. Hortic. Sér. IV. ii. (1858) 305 = florida.
canadensis, Willd. Enum. Hort. Berol. i. 222 = trifida.
coraeënsis, *DC. Prod.* iv. 330.—Japon.
floribunda, *Sieb. & Zucc. Fl. Jap.* i. 73. t. 32.—Japon.
florida, *Sieb. & Zucc. l. c.* 75.—China.
grandiflora, Sieb. & Zucc. l. c. 71. t. 31 = coraeensis.
hortensis, *Sieb. & Zucc. l. c.* 70. t. 29.—China; Japon.
humilis, Pers. Syn. i. 214 = trifida.
japonica, *DC. Prod.* iv. 330.—Japon.
longifolia, Rafin. New Fl. Am. iii. 21 = trifida.
Lonicera, Mill. Gard. Dict. ed. VIII = trifida.
lutea, Pursh, Fl. Am. Sept. i. 162 = trifida.
Middendorfiana, *Carr. in Rev. Hort. Sér.* IV. ii. (1858) 306.—Sibir. or.
multiflora, Lem. Illustr. Hortic. v. (1863) t. 383 = floribunda.
parvifolia, Rafin. New Fl. Am. iii. 23 = trifida.
paucifiora, Carr. in Rev. Hortic. Sér. IV. ii. (1858) 310 = florida.
rosea, *Mast. in Journ. Hort. Soc.* N. S. v. (1870) 128. —China.
rosea, Walp. Ann. i. 365 = florida.
sessilifolia, *Buchl. in Am. Journ. Sc.* xlv. (1843) 174. —Am. bor.
splendens, Carr. in Fl. des Serres, viii. (1852–53) 292 = sessilifolia.
Tournefortii, Michx. Fl. Bor. Am. i. 107 = trifida.
trifida, *Mornch. Meth.* 492.—Am. bor.
versicolor, Sieb. & Zucc. Fl. Jap. i. 74. t. 33 = floribunda.

DIERVILLEA, Bartl. Ord. 214 (1830) = **Diervilla**, Tourn.

DIESINGIA, Endl. in Flora, xv. (1832) 118 = **Psophocarpus**, Neck. (Legumin.).
scandens, Endl. l. c. 117 = P. palustris.

DIETERIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 300 = **Aster**, Tourn. (Compos.).
asteroides, Torr. in Emory, Notes Mil. Reconnois. 141 = A. canescens.
canescens, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 300 = A. canescens.
cronopifolia, Nutt. l. c. 302 = A. tanacetifolius.
discoidea, Nutt. l. c. 301 = A. canescens.
gracilis, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 177 = Aplopappus gracilis.
incana, Torr. & Gray, Fl. N. Am. ii. 100 = A. incanus.
puberulenda, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 300 = A. canescens.
rubiginosa, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 177 = Aplopappus rubiginosus.
sessilifolia, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 301 = A. canescens.
spinulosa, Nutt. l. c. 301 = Aplopappus spinulosus.
viscosa, Nutt. l. c. = A. canescens.

DIETES, Salisb. in Trans. Hort. Soc. i. (1812) 307 = **Moraea**, Linn. (Irid.).
bicolor, Sweet, ex G. Don, in Lond. Hort. Brit. Suppl. i. 587 = M. bicolor.

DIETES:—
calenulata, Sweet, Hort. Brit. ed. II. 497 = Moraea calenulata.
compressa, Klatt, in Linnaea, xxxiv. (1865–66) 584 = M. iridioides.
crassifolia, G. Don, in Sweet, Hort. Brit. ed. III. 661 = M. iridioides.
Huttoni, Baker, in Hook. f. Bot. Mag. t. 6174 = M. spathacea.
iridifolia, Salisb. in Trans. Hort. Soc. i. (1812) 307 = M. iridioides.
iridioides, Sweet, Hort. Brit. ed. II. 497 = M. iridioides.
Robinsoniana, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 274 = M. Robinsoniana.

DIETERICA, Ser. in DC. Prod. iv. 8 (1830) = **Caldcluvia**, D. Don (Saxifr.).
paniculata, Ser. l. c. = C. paniculata.

DIETERICHIA, Giseke, Prael. Linn. 228 (1792) = seq.

DIETRICHIA, Giseke, Prael. Linn. 199, 208 (1792) = **Klingtbar**, Adans. (Scrophulin.).
Lampsjanq, Giseke, l. c. 199 = Z. Zerumbet.
Lampuyang, Giseke, l. c. 208 = Z. Zerumbet.
major, Raeusch. Nom. ed. III. 1 = Z. Zerumbet.
minor, Raeusch, l. c. = Z. Zerumbet.
spuria, Giseke, Prael. Linn. 199 = Z. Zerumbet.

DIETRICHIA, Tratt. Arch. Gewaechsk. n. 449 (1812) = **Rochea**, DC. (Crassulac.).
bicolor, Eckl. & Zeyh. Enum. 305 = R. odoratissima.
coccinea, Tratt. Arch. Gewaechsk. n. 449 = R. coccinea.
jasminea, Eckl. & Zeyh. Enum. 305 = R. jasminea.
media, Eckl. & Zeyh. l. c. 304 = R. versicolor.
odoratissima, Tratt. ex DC. Prod. iii. 394 = R. odoratissima.
versicolor, Eckl. & Zeyh. Enum. 304 = R. versicolor.

DIEUDONNAEA, Copn. in Bull. Soc. Bot. Belg. xiv. (1875) 239. *CUCURBITACEAE*.
rhizantha, *Cogn. l. c.*—Peruv.

DIGASTER, Miq. Fl. Ind. Bat. Suppl. 329 (1860) = **Pygeum**, Gaertn. (Rosac.).
sumatranus, Miq. l. c. 330 = P. sumatranum.

DIGERA, Forsk. Fl. Aegypt. Arab. 65 (1775). *AMARANTACEAE*, Benth. & Hook. f. iii. 28.
ECLIOTOPIA, et URTIA, Rafin. Fl. Tellur. iii. 40 (1836).
alternifolia, Aschers. in Schweinf. Beitr. Fl. Aethiop. 180 = arvensis.
arvensis, *Forsk. Fl. Aegypt. Arab.* 65.—As. et Afr. trop.
ciliata, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 285 = arvensis.
Forskaoli, Blume, Bijdr. 542 = arvensis.
fruticosa, Juss. ex Stend. Nom. ed. II. i. 506 = arvensis.
muricata, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 285 = arvensis.

DIGITALIS. [Tourn.] Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 960.
acuta, Moench, Meth. 468 = lutea.
alba, Schrank, Baier. Fl. ii. 170 = purpurea.
ambigua, *Murr. Prod. Stirp. Gotting.* 62.—Europ.; As. occ.
ambigua, Roem. ex Steud. Nom. ed. I. 274 = purpurea.
ambigua, Willd. ex Ledeb. Fl. Ross. iii. 222 = Pentstemon frutescens.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 300.—Algeria.
aurea, Desf. Cat. Hort. Par. ed. III. 396 = D. Fontanesii.
aurea, Hort. ex Benth. in DC. Prod. x. 450 = laevigata.
aurea, Lindl. Digit. Monog. 18. t. 18 = ferruginea.
australis, Tenore, Fl. Med. Univ. 507 = lutea.
brachyantha, Griseb. Spicil. Fl. Rumel. ii. 518 = ferruginea.
canariensis, Linn. Sp. Pl. 868 = Isoplexis canariensis.

DIGITALIS:—
carinata, Boiss. ex Benth. in DC. Prod. x. 450 = orientalis.
carnea, Melg. & Wenig. Syst. Verz. 80 = purpurea.
chinensis, Hook. & Arn. Bot. Beech. Voy. 204, sphalm. vide *sinensis*.
ciliata, *Trautv. in Bull. Acad. Pétersb.* x. (1866) 397. —Reg. Caucas.
cochinchinensis, *Lour. Fl. Cochinch.* 378.—Cochinch.
dasyantha, Pall. ex Benth. in Trans. Linn. Soc. x. (1811) 259 = Pentstemon frutescens.
denticulata x, Tausch, in Flora, xix. (1836) 390.—Europ.
dracocephaloides, Vell. Fl. Flum. 268; vi. t. 101 (Acanthac.).—Bras.
dubia, *Rodrig. in Anal. Soc. Esp. Hist. Nat.* iii. (1874) 48.—Ins. Balear.
epiglottides, Brera, ex Steud. Nom. ed. I. 275 = lanata.
eriostachys, *Bess. ex Reichb. Hort. Bot.* iii. 12. t. 230. —Hab.?
erubescens x, Alt. Hort. Kew. ed. II. iv. 29.
ferruginea, Lam. Encyc. ii. 280 = parviflora.
ferruginea, *Linn. Sp. Pl.* 868.—Europ. austr.
ferruginea, Sibth. & Sm. Fl. Graec. Prod. i. 289 = aurea.
ferruginea, *Wulf. Pl. Rar. Desc.* 75 = laevigata.
flava, Georgi, Reise, ii. 718 = ambigua.
Fontanesii, *Steud. Nom.* ed. II. i. 507.—Hab.?
fucata, Ehrh. Beitr. vii. 151 = purpurascens.
fulva x, Lindl. Digit. Monog. 15. t. 9.—Europ. austr.
fuscescens x, Waldst. & Kit. Pl. Rar. Hung. iii. 304. —Croatia.
glaucinoides, *Carr. in Rev. Hortic.* (1872) 80.—Hab.?
glutinosa, J. Gaertn. in Nov Comm. Petrop. xiv. i. (1770) 544 = Rehmannia chinensis.
grandiflora, Lam. Fl. Fr. ii. 330 = ambigua.
Heidreichii, Jaub. & Spach, Illustr. v. sub t. 409 = orientalis.
integrifolia, Seenus, Reise Istrien, 62 = laevigata.
intermedia, Lapeyr. Hist. Abr. Pl. Pyr. 357 = purpurascens.
intermedia, Pers. Syn. ii. 167 = lutea.
laciniata, *Lindl. Bot. Reg.* t. 1201.—Hispan.
lanigata, C. A. Mey. Verz. Pfl. Cauc. 110 = nervosa.
laevigata, *Waldst. & Kit. Pl. Rar. Hung.* ii. 171. t. 158.—Reg. Danub.; Graecia.
lanata, *Ehrh. Beitr.* vii. 152.—Reg. Danub.; Graecia.
leucocephala, Steud. Nom. ed. II. i. 507, sphalm. = leucophaea.
leucophaea, *Sibth. & Sm. Fl. Graec. Prod* i. 289.—Graecia.
Libertiana, Dum. Fl. Belg. Prod. 86 = purpurea.
Lindleyana x, Tausch, in Flora, xix. (1836) 389.
longebractesta, *Richter, ex Staff, in Denkschr. Acad. Wien*, i. (1885) 91.—Caria.
longiflora, Lej. Rev. Fl. Spa, 126 = purpurascens.
lurida, Salisb. Prod. 101 = Isoplexis canariensis.
lutea, *Linn. Sp. Pl.* 868.—Europ. austr.
lutea, Poll. Hist. Pl. Palat. ii. 199 = ambigua.
lutea, Sibth. & Sm. Fl. Graec. Prod. i. 289 = viridiflora.
lutea-major, Gilib. Fl. Lituan. i. 124 = ambigua.
lutescens x, Lindl. Digit. Monog. 22. t. 21.
lutescens, *Reuy, in Bull. Soc. Bot. Fr.* xxii. (1875) 81. —Gallia.
macrostachya, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 124 = Isoplexis Sceptrum.
magnifiora, Mill. Gard. Dict. ed. VIII. n. 4 = ambigua.
mariana, *Boiss. Voy. Bot. Espagne*, ii. 465. t. 126.—Hispan.
media, Elmig. Hist. Digit. 42 = purpurascens.
media x, Roth, Catalect. Bot. ii. 60.—Europ. austr.
microstha, Roth, Nov. Pl. Sp. 284 = lutea.
Millers, G. Don, Gen. Syst. iv. 506 = ambigua.
minor, Bourg. ex Nym. Consp. 535 = dubia.
minor, *Linn. Mant.* ii. 367.—Hispan.
nervosa, *Steud. & Hochst. ex Benth. in DC. Prod.* x. 450.—Reg. Caucas.; Persia.
newadensis, Kunze, in Flora, xxix. (1846) 673 = purpurea.
nova, Winterl. ex Lindl. Digit. Monog. 19 = lanata.
obscura, *Linn. Sp. Pl.* ed. II. 867.—Hispan.
obtusa, Moench, Meth. 468 = media.
ochroleuca, Jacq. Fl. Austr. i. 36 = ambigua
orientalis, Elmig. Hist. Digit. 45 = lanata.
orientalis, *Lam. Encyc.* ii. 280.—As. Min.

DIGITALIS:—
orientalis, Mill. Gard. Dict. ed. VIII. n. 7 = ambigua.
parviflora, *Jacq. Hort. Vindob.* i. 6. t. 17.—Europ. austr.
parviflora, Lam. Fl. Fr. ii. 333 = lutea.
pilosa, Vell. Fl. Flum. 269 ; vi. t. 102 (Acanthac.).—Bras.
purpurascens, Lej. Fl. Spa, ii. 320 = purpurea.
purpurascens x, Lindl. Digit. Monog. 22 = *Lindleyana*.
purpurascens, *Roth, Catalect. Bot.* ii. 62.—Europ.
purpurea, *Linn. Sp. Pl* 621.—Europ.
purpureo-lutea, G. F. W. Mey. Chloris Hanov. 324 = purpurascens.
regalis, Salisb. Prod. 101 = Isoplexis Sceptrum.
rigida x, Lindl. Digit. Monog. 21. t. 19.
Roxburghii, Roth, Nov. Pl. Sp. 283 = Centranthera humilina ?
Sceptrum, Linn f. Suppl. 282 = Isoplexis Sceptrum.
sibirica, *Lindl. Digit. Monog.* 16. t. 11.—Sibiria
sinensis, Lour. Fl. Cochinch. 378 = Adenosma grandiflorum.
speciosa, Salisb. Prod. 100 = purpurea.
stricta, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 99 = Centranthera hispida.
Thapsi, Bert. ex Nym. Consp. 535 = purpurea.
Thapsi, Cambess ex Nym. l. c. = dubia.
Thapsi, *Linn. Sp. Pl* ed. II. 867.—Hispan.
tomentosa, Link & Hoffmgg. Fl. Port. i. 220. t. 29 = purpurea.
tubiflora x, Lindl Digit. Monog. 22. t. 22.—Europ. austr.
tubulosa x, Tausch, in Flora, xix. (1836) 390.—Europ. ?
Ursenaga, Noé, ex Benth. in DC. Prod. x. 450 = laevigata.
viridiflora, *Lindl. Digit. Monog.* 21. t. 18.—Graecia, Macedon.
Wrisbergii, Roth, Catalect. Bot. i. 71 = lanata.

DIGITARIA, Heist. ex Adans. Fam. ii. 38, 550 (1763) ; Scop. Fl. Carn. ed. II. i. 52 (1772) = **Panicum**, Linn. (Gramin.).
adpressa, Boj. Hort. Maurit. 368 = P. ambigunm.
aegyptiaca, Willd. Enum. Hort. Berol. 93 = P. sanguinale.
affinis, Koem. & Schult. Syst. ii. 470 = P. appressum.
appressa, Pers. Syn. i. 85 = P. fluitans.
aurea, Spreng. Syst. i. 272 = Paspalum pulchrum.
australis, Willd. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. (1835). ii. 201 = P. sanguinale.
barbata, Willd. Enum. Hort. Berol. 91 = P. barbatum.
barbulata, Desv. Opusc. 63.—Ins Borbon.
bicornis, Roem. & Schult. ex Loud. Hort. Brit. 24 = P. bicorne.
biformis, Willd. Enum. Hort. Berol. 92 = P. sanguinale.
borbonica, Desv. Opusc. 63.—Ins. Borbon.
brevifolia, Link, Hort. Berol. i. 225 = P. brevifolium.
chinensis, Hornem. Hort. Hafn. Suppl. 8 = P. chinense ?
chrysoblephara, Fig. & De Not. in Mem. Acc. Tor. Ser. II. xiv. (1853) 356.—Nubia.
ciliaris, Pers. Syn. i. 85 = P. sanguinale.
collina, Salisb. Prod. 19 = Andropogon Ischaemum.
commutata, Schult. Mant. ii. 262 = P. sanguinale.
composita, Willd. Enum. Hort. Berol. 91 = Oplismenus compositus.
concinna, Schrad. ex Steud. Nom. ed. II. i. 507 = P. violascens ?
conjugata, Schult. Mant. ii. 262 = P. conjugatum.
consanguinea, Gaudich. in Freyc. Voy. Bot. 410 = Paspalum dichotomum.
cuspidata, Schult. Mant. ii. 265 = P. cuspidatum.
Dactylon, Scop. Fl. Carn. ed. II. i. 52 = Cynodon Dactylon.
debilis, Willd. Enum. Hort. Berol. 91 = Paspalum debile.
decipiens, Fig. & De Not. in Mem. Acc. Tor. Ser. II. xiv. (1853) 359.—Nubia.
denudata, Link, Hort. Berol. i. 222 = P. violascens.
didactyla, Spreng. Syst. i. 270 = Cynodon tenellus.
didactyla, Willd. Enum. Hort. Berol. 91 = P. didactylum.
digitata, Busse, in Miq. Fl. Jungh. 381.—Java.
distachya, Pers. Syn. i. 85 = P. distachyum.
domingensis, Desv. ex Kunth, Enum. Pl. i. 49 = Paspalum platycaule.

DIGITARIA :—
domingensis, Roem. & Schult. Syst. ii. 475 = Panicum domingense.
elongata, Spreng. Syst. i. 271 = Cynodon elongatus.
elatior, Willd. Enum. Hort. Berol. 91 = P. elatius.
erianthe, Steud in Flora, xii. (1829) 468 = P. ciliare.
eriogona, Link, Hort. Berol. i. 227 = P. sanguinale.
fascicularis, Link, l. c. 223 = P. Schraderi.
fasciculata, Desv. Opusc. 63 = P. fasciculatum ?
filiformis, Delile, Fl. Aeg. Illustr. 51 = P. filiforme.
filiformis, Koel. Desc. Gram. 26 = P. glabrum.
fimbriata, Link, Hort. Berol. i. 226 = P. sanguinale.
foliosa, Lag. Gen. et Sp. Nov. 4 = Paspalum foliosum.
gibbosa, Beauv. Agrost. 160 = P. gibbosum.
glabra, Beauv. l. c. 51 = P. glabrum.
gracilis, Gusa. Enum. Pl. Vasc. Inarim. 375 = P. gracile, *Nym.*
hispida, Spreng. Syst. i. 271 = Arthraxon ciliaris.
hispidula, Willd. Enum. Hort. Berol. 91 = P. Crus-galli.
horizontalis, Willd. l. c. 92 = P. sanguinale.
humifusa, Pers. Syn. i. 85 = P. glabrum.
humifusa, [Pieri ?] in Ionios Anthol. ii. (1834) 464 = P. sanguinale.
inaequalis, Link, Hort. Berol. i. 224 = P. sanguinale.
Ischaemum, Muhl. Desc. Gram. 131 = P. sanguinale.
jamaicensis, Spreng. Syst. i. 272 = Paspalum digitatum.
lanosa, Llanos, Fragm. 28 = Eleusine aegyptiaca.
linearis, Pers. Syn. i. 85 = P. sanguinale.
linearis, Kostal. in Verh. Zool. Bot. Ges. Wien, xxii. (1872) 99 = P. glabrum.
littoralis, Salisb. Prod. 19 = Cynodon Dactylon.
longiflora, Pers. Syn. i. 85 = P. parvulum.
malabarica, Roem. & Schult. Syst. ii. 474 = Centotheca lappacea.
marginata, Link, Enum. Hort. Berol. i. 102 = P. sanguinale.
marithma, Spreng. Syst. i. 272 = Cynodon Dactylon.
miliacea, Willd ex Steud. Nom. ed. II. i. 507 = P. fasciculatum.
mollicoma, Schrad. ex Link, Hort. Berol. i. 229 = P. mollisumum.
monostachya, Willd. ex Steud. Nom. ed. II. i. 508 = P. rottboelioides.
nervosa, Koem. & Schult. Syst. ii. 473 = F. Rottleri
nodosa, Parl. Pl. Nov. 89 = P. Parlatorei.
nuda, Schum in Schum. & Thonn. Beskr. Guin. Pl. 45. —Afr. trop.
obtusifolia, Roem. & Schult. Syst. ii. 989 = P. obtusifolium.
panicea, Willd. ex Steud. Nom. ed. II. i. 508 = P. sanguinale.
papposa, Beauv. Agrost. 51 = P. papposum.
paspaloides, Michx. Fl. Bor. Am. i. 46 = Paspalum vaginatum.
pertensis, Busse, in Miq. Pl. Jungh. 381.—Ins. Banca.
Pes-anseris, Busse, l. c. 380.—Java.
pilosa, Michx. Fl. Bor. Am. i. 46 = P. filiforme.
pilosa, [Pieri ?] in Ionios Anthol. ii. (1834) 464 = P. sanguinale.
pilosa, Willd. Enum. Hort. Berol. 91 = Paspalum scrobiculum.
platiculis, Desv. Opusc. 62.—Hispan.
plebeia, Phil. in Anal. Univ. Chil. (1873) 557.—Chili.
polystachya, Spreng. Syst. i. 272 = Leptochloa polystachya.
praecox, Willd. Enum. Hort. Berol. 91 = P. sanguinale.
procumbens, Hort. ex Steud. Nom. ed. II. i. 508 = P. glabrum.
propinqua, Gaudich. in Freyc. Voy. Bot. 410 = P. timorense.
pruriens, Busse, in Miq. Fl. Jungh. 379 = P. pruriens.
Pseudo-duros, Nees, ex Schlecht. in Linnaea, xxvi. (1853-55) 458 = P. Pseudo-duros.
Pseudo-ischaemum, Busse, in Miq. Fl. Jungh. 382.—Java.
puberula, Link, Hort. Berol. i. 228 = P. puberulum.
pulchella, Griseb. Cat. Pl. Cub. 221.—Cuba.
radiata, Spreng. Syst. i. 272 = Cynodon radiatus.
radicosa, Miq. Fl. Ind. Bat. iii. 437 = P. sanguinale.
reptans, Schum. & Thonn. Beskr. Guin. Pl. 44.—Afr. trop.
repens, Hort. Par. ex Kunth, Enum. Pl. i. 84 = P. domingense.
repens, Willd. ex Kunth, l. c. 51 = P. sanguinale.

DIGITARIA :—
repulsa, Busse, in Miq. Fl. Jungh. 381.—Java.
Rottleri, Koem. & Schult. Syst. ii. 471 = Panicum appressum.
Roxburghii, Spreng. Syst. i. 270 = Paspalum nematodes.
sanguinale, Scop. Fl. Carn. ed. II. i. 53 = P. sanguinale.
sanguinaria, Steud. Nom. ed. I. 376 = seq.
sanguinea, [Weber, in] Wigg. Prim. Fl. Holsat. 6 = P. sanguinale.
serotina, Michx. Fl. Bor. Am. i. 46 = Paspalum serotinum.
setigera, Roth, ex Roem. & Schult. Syst. ii. 474 = P. sanguinale.
setosa, Desv. ex Ham. Prod. Fl. Ind. Occ. 6 = P. Hamiltonii.
sienitica, Hort. Berol. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. (1835) ii. 201 = P. sanguinale.
stolonifera, Schrad. Fl. Germ. i. 165. t. 3. f. 9 = Cynodon Dactylon.
stricta, Gaudich. in Freyc. Voy. Bot. 409 = P. Gaudichaudi.
stricta, Koth, Nov. Pl. Sp. 28 = P. Kunthii.
stricta, Willd. ex Steud. Nom. ed. II. i. 508 = P. sanguinale.
tenuiflora, Beauv. Agrost. 51 = P. brevifolium.
Thunbergii, Beauv. l. c. = Hemarthria compressa.
tristachya, Schult. Mant. ii. 261 = Paspalum distichum.
tristachya, Willd. ex Steud. Nom. ed. II. i. 508 = P. sanguinale.
umbraticola, Doell, in Mart. Fl. Bras. ii. II. 134, in syn. = P. sanguinale.
umbrosa, Link, Hort. Berol. i. 227 = P. sanguinale.
umbrosa, Pers. Syn. i. 85 = P. prostratum.
undulata, Schrank, ex Steud. Nom. ed. II. i. 508 = P. sanguinale.
uniflora, Salzm. ex Steud. l. c. = Paspalum uniflorum.
urvicladoe, Busse, in Miq. Fl. Jungh. 383.—Java.
variabilis, Fig. & De Not. in Mem. Acc. Tor. Ser. II. xiv. (1853) 357.—Nubia.
velutina, Beauv. Agrost. 51 = P. sanguinale.
ventia, Fig. & De Not. in Mem. Acc. Tor. Ser. II. xiv. (1853) 356.—Nubia.
villosa, Hort. Vindob. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. (1835) ii. 198 = P. Schraderi.
villosa, Pers. Syn. i. 85 = P. filiforme.
violascens, Link, Hort. Berol. i. 229 = P. violascens.

DIGLOSSELIS, Rafin. Fl. Tellur. iv. 98 (1836) = **Aristolochia**, Tourn.
cymbifera, Rafin. l. c. = A. cymbifera.
latifolia, Rafin. l. c. = prsec. ?
trinervis, Rafin. l. c. = prsec. ?

DIGLOSSUS, Cass. in Bull. Soc. Philom. (1817) 70 = **Tagetes**, Linn. (Compos.).
variabilis, Cass. in Dict. Sc. Nat. xliii. 241 = T. foeniculacea.

DIGLOTTIS, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 151, 170 = **Cuspuria**, Humb. (Rutac.).
obovata, Nees & Mart. l. c. 170. t. 25 = C. obovata.

DIGLYPHIS, Blume, Pract. Fl. Jav. vi. (1828) p. vi = **Chrysoglossum**, Blume (Orchid.).

DIGLYPHOSA, Blume, Bijdr. 336. t. 60 (1825) = **Chrysoglossum**, Blume (Orchid.).
latifolia, Blume, l. c. = C. latifolium (in Addend.).

DIGLYPHYS, Spach, Hist. Vég. Phan. xii. 176 (1846) = *prsec.*

DIGNATHE, Lindl. in Journ. Hort. Soc. iv. (1849) 268. *ORCHIDEÆ*, Benth. & Hook. f. iii. 540.
pygmæa, *Lindl. l. c.*—Mexic.

DIGOMPHIA, Benth. in Hook. Lond. Journ. Bot. v. (1846) 264. *BIGNONIACEÆ*, Benth. & Hook. f. ii. 1051.
laurifolia, *Benth. l. c.*—Guinea.

DIGOMPHOTIS, Rafin. Fl. Tellur. ii. 97 (1836) = **Habenaria**, Willd. (Orchid.).
cordata, Rafin. l. c. = H. cordata.
undulata, Rafin. l. c. = H. goodyeroides.

DIGONOCARPUS, Vell. Fl. Flum. 158 (1825); iv. t. 14 = **Cupania**, Plum. (Sapind.).
inflatus, Vell. l. c. 154; iv. t. 14 = C. emarginata.

DIGRAPHIS, Trin. Fund. Agrost. 127 (1820) = **Phalaris**, Linn. (Gramin.)
americana, Loud. Hort. Brit. 27 = P. intermedia.
arundinacea, Trin. Fund. Agrost. 127 = P. arundinacea.

DIGYROLOMA, Tercs. in Bull. Soc. Nat. Mosc. xxxv. (1862) ii. 329. *ACANTHACEAE*.—Cf. Benth. & Hook. f. ii. 1071.
Cleghornii, *Turcz. l. c.* 330.—Ind. or.

DILANTHES, Salisb. Gen. Pl. Fragm. 70. (1866) = **Antherium**, Linn. (Liliac.).
revolutus, Salisb. l. c. = A. revolutum.

DILASIA, Rafin. Fl. Tellur. iv. 122 (1836) = **Anellema**, R. Br. (Commel.).
vaginata, Rafin. l. c. 123 = A. vaginatum.

DILATRIS, Berg. Descr. Pl. Cap. 9. t. 3. f. 5 (1767). *HAEMODORACEAE*, Benth. & Hook. f. iii. 674
carolinana, Lam. Tabl. Encyc. i. 127 = Lachnanthes tinctoria.
corymbosa, *Berg. Descr. Pl. Cap.* 9 —Afr. austr.
Heritiera, Pers. Syn. i. 54 = Lachnanthes tinctoria.
hexandra, Lam. Encyc. ii. 282 = Lanaria plumosa.
ixioides, Lam. l. c. = corymbosa.
paniculata, *Linn. f. Suppl.* 103 —Afr. austr.
tinctoria, Pursh, Fl. Am. Sept. i. 30 = Lachnanthes tinctoria.
umbellata, Linn. f. Suppl. 101 = corymbosa.
viscosa, *Linn. f. l. c.*—Afr. austr.

DILEMA, Griff. Notul. iv. 705 (1854) = **Dillenia**, Linn. (D. indica).

DILEPTIUM, Rafin. Fl. Ludov. 85 (1817) = **Lepidium**, Tourn. (Cruciferæ).
diffusum, Rafin. l. c. 85 = L. virginicum.
precox, Rafin. l. c. 86 = L. virginicum.
virginicum, Rafin. in Am. Monthly Mag. (1818) 267 = L. virginicum.

DILEPYRUM, Michx. Fl. Bor Am. i. 40 (1803) = **Muehlenbergia**, Schreb. (Gramin.).
aristosum, Michx. l. c. = M. aristata.
diffusum, Beauv. Agrost. 160 = M. diffusa.
minutiflorum, Michx. Fl. Bor. Am. i. 40 = M. diffusa.
multiflorum, Beauv. Agrost. 160 = M. diffusa.

DILEPYRUM, Rafin. in Med. Repos. N. York, II. v. (1808) 352 = **Oryzopsis**, Michx. (Gramn.).

DILICARIA, T. Anders. in Journ. Linn. Soc. vii. (1864) 36 = seq.

DILIVARIA, Juss. Gen. 103 (1789) = **Asanthus**, Tourn.
abracteata, Pers. Syn. ii. 179 = A. abracteatus.
horrida, Nees, in Linnæa, xv. (1841) 368 = Acanthopsis horrida.
ilicifolia, Juss. Gen. 103 = A. ilicifolius.
longifolia, Poir. Encyc. Suppl. i. 88, in syn. = A. longifolius.
polystachya, Endl. Gen. 703 = A. polystachyus.
scandens, Nees, in DC. Prod. xi. 389 = A. volubilis.
volubilis, Nees, in Wall. Pl. As. Rar. iii. 98 = A. volubilis.

DILKEA, Mast. in Trans. Linn. Soc. xxvii. (1871) 697. *PASSIFLOREAE*.
acuminata, *Mast. l. c.* 698.—Bras.
retusa, *Mast. l. c.*—Bras.
Wallisii, *Mast. in Mart. Fl. Bras.* xiii. t. 622.—Bras.

DILLENIA, Linn. Syst. ed. I (1735). *DILLENIACEAE*. Benth. & Hook. f. i. 18.
COLLETTIA, Salisb. Parad. Lond. sub t. 78 (1807).
DILEMA, Griff. Notul. iv. 705 (1854).
SYALITA, Adans. Fam. ii. 384 (1763).
alata, Banks, ex DC. Syst. i. 494 = Wormia alata.
Andrewsa, *F. Muell. Fragm.* v. 175.—Austral.
angusta, Roxb. Hort. Beng. 43; Fl. Ind. ii. 632 = pentagyna.
DILLENIA:—
aurea, *Sm. Exot. Bot.* ii. 95. t. 92.—Ind. or.
Blanchardii, *Pierre, Fl. For. Cochinch.* t. 14.—Cochinch.
bracteata, *Wight, Ic.* ii. t. 356.—Ind. or.
dentata, Thunb. in Trans. Linn. Soc. i. (1791) 201. t. 20 = Wormia triquetra.
elata, *Pierre, Fl. For. Cochinch.* t. 9.—Cochinch.
elliptica, Thunb. in Trans. Linn. Soc. i. (1791) 200 = indica.
elongata, *Miq. Fl. Ind. Bat.* i. ii. 12.—Sumatra.
eximia, *Miq. l. c. Suppl.* 620.—Sumatra.
floribunda, *Hook. f. & Thoms. Fl. Ind.* i. 71.—Burma.
grandifolia, *Wall. Cat.* n. 946.—Malaya.
grossulariæfolia, Poir. Encyc. Suppl. v. 146 = Hibbertia grossulariæfolia.
Hookeri, *Pierre, Fl. For. Cochinch.* t. 5.—Cambodia
humilis, Dunn, Cat. Hort. Cant. ed. I. 64 = Hibbertia volubilis.
indica, Blanco, Fl. Filip. ed. I. 472 = philippinensis.
indica, *Linn. Sp. Pl.* 535.—As trop.
integra, Moench, Meth. Suppl. t. 76 = Hibbertia volubilis.
integra, Thunb. in Trans. Linn Soc. i. (1791) 199. t. 18 = retusa.
macrophylla, *Reinw. ex Blume, Cat. Gew. Buitenz.* 78, nomen.—Java.
meliosmæfolia, *Hook. f. & Thoms. in Fl. Brit. Ind* i. 36.—Malacca.
ochreata, Teysm. & Binn. Cat. Hort. Bog. 178 = Wormia ochreata.
ornata, Wall. Pl. As. Rar. i. 21. t. 28 = aurea.
ovata, *Wall. Cat.* n. 945.—Ins. Penang.
parviflora, *Griff. Notul.* iv. 704.—Burma.
pentagyna, *Roxb Pl. Corom.* i. 21. t. 26.—Ind. or.
philippinensis, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 307.—Ins. Philipp.
pilosa, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 102 = scabrella.
pilosa, Roxb. Hort. Beng. 43; Fl. Ind. ii. 652 = pentagyna.
procumbens, Labill. Nov. Holl. Pl. ii. 16. t. 156 = Hibbertia procubens.
pulcherrima, *Kurz, in Journ. As. Soc. Beng* xl. (1871) ii. 46.—Burma.
racemosa, Buch.-Ham ex D. Don, Prod. Fl. Nep. 225 = Saurauja nepaulensis.
radicans, *Hort. ex Link, Enum. Hort. Berol.* ii 86.—Hab.?
Reifferscheidia, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 8.—Ins. Philipp.
repanda, Roxb. Fl. Ind. ii. 652 = bracteata.
retusa, *Thunb. in Trans. Linn. Soc.* i. (1791) 200. t. 19. —Ceylan.
scabrella, *Roxb. Hort. Beng.* 43; *Fl. Ind.* ii. 655.—Reg. Himal.
scandens, Willd. Sp. Pl. ii. 1251 = Hibbertia volubilis.
serrata, *Thunb. in Trans. Linn. Soc.* i. (1791) 201.—Malaya.
speciosa, Blanco, Fl. Filip. ed. II. 329 = philippinensis.
speciosa, Curt. Bot. Mag. t. 449 = Hibbertia volubilis.
speciosa, Griff. Notul. iv. 708 = aurea.
speciosa, Thunb. in Trans. Linn. Soc. i. (1791) 200 = indica.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 620.—Sumatra.
tetragyna, Lag. ex DC. Syst. i. 429 = Hibbertia diffusa.
turneræflora, [Ker-Gawl.] Reeves. Fl. Bot. Rep. 27 = Hibbertia volubilis.
volubilis, Vent. Choix, 11 = Hibbertia volubilis.

DILLWINIA, Poir. in Dict. Sc. Nat. xliii. 250 (1819) = **Dillwynia**, Sm. (Legumin.).

DILLWYNIA, Roth, Catalect. fasc. iii. 71 (1806) = **Rothia**, Pers. (Legumin.).
trifoliata, Roth, l. c. = R. trifolistata.

DILLWYNIA, Sm. in Kon. & Sima, Ann. Bot. i. (1805) 510. *LEGUMINOSAE*, Benth. & Hook. f. i. 471.
XEROPETALUM, Reichb. Consp. 154 (1828).
aciculare, Meisn. ex Lehm. Pl. Preiss. i. 62 = cinerascens.
acicularis, Sieber, ex DC. Prod. ii. 109 = prærc. ?
adenophora, Endl. Nov. Stirp. Dec. 80 = floribunda.
brunioides, *Meisn. in Lehm. Pl. Preiss.* i. 62.—Austral.
cinerascens, *R. Br. in Bot. Mag.* t. 2247.—Austral.
cinerascens, DC. Prod. ii. 109 = juniperina.
clavata, Paxt. Mag. Bot. vii. (1840) 117 = floribunda.
DILLWYNIA:—
cuneata, Sieber, ex Benth. Fl. Austral. iii. 136 = Pultenæa cuchifa.
divaricata, *Benth. l. c.* 151.—Austral.
Drummondii, *Van Houtte, in Regel, Gartenfl.* (1863) 305. t. 432.—Austral. ?
elegans, Endl. Nov. Stirp. Dec. 15 = floribunda.
ericifolia, Sims, Bot. Mag. t. 1545 = floribunda.
ericifolia, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 510. —Austral.
ericoides, Sieber, ex Benth. Fl. Austral. iii 148 = ericifolia.
filifolia, Endl. Nov. Stirp. Dec. 15 = ericifolia.
floribunda, *Sm in Kon. & Sims, Ann. Bot.* i. (1805) 510.—Austral.
glaberrima, Sm l. c. = ericifolia.
glycinæfolia, Sm. in Trans. Linn. Soc. ix. (1808) 264 = Chorizema angustifolium
hispida, *Lindl. in Mitch. Three Exped. Austral.* ii. 261.—Austral
hispidula, Sieber, ex Benth. Fl. Austral. iii. 149 = floribunda.
juniperina, *Lodd. Bot. Cab.* t. 401.—Austral.
lanceolata, *Lasiel. ex Drapiez, Herb. Amat. Fl.* vii (1834) t. 478.—Austral.
laxiflora, Benth. in Hueg. Arch. Bot. t. 9 = cinerascens.
macrophylla, Sieber, ex DC. Prod. ii. 109 = ericifolia.
myrtifolia, Sm. in Trans. Linn. Soc. ix. (1808) 269 = Eutaxia myrtifolia.
obovata, Labill. Nov. Holl. Pl. i. 110. t. 140 = Eutaxia myrtifolia.
parviflora, Benth in Ann. Wien. Mus. ii. (1838) 78, in obs. = parvifolia ?
parvifolia, R. Br. in Bot. Mag. t. 1527 = ericifolia.
patula, *F. Muell. Fragm.* vi. 16.—Austral.
peduncularis, Benth. in Ann. Wien. Mus. ii. (1838) 78 = ericifolia.
phylicoides, A. Cunn. in Field, N. S. Wales, 347 = ericifolia.
Preissii, *Benth. Fl. Austral.* ii. 149.—Austral.
pungens, *Mackay, ex Sweet, Hort. Brit.* ed. I. 474.—Austral.
ramosissima, Benth. in Ann. Wien Mus. ii. (1838) 79 = ericifolia.
rudis, Sieber, ex DC. Prod. ii. 109 = floribunda.
scabra, Schlecht. in Linnæa, xx. (1847) 666 = hispida.
sericea, A. Cunn. in Field, N. S. Wales, 347 = floribunda.
seriphioides, Endl. Nov. Stirp. Dec. 17 = ericifolia.
speciosa, Paxt. Mag. Bot. iv. (1838) 99 = ericifolia.
subaphylla, Colla, Hort. Ripul. App. ii. 347 (Dillwyniae sp.).—Hab.?
tenuifolia, Sieber, ex DC. Prod. ii. 109 = ericifolia.
teucrioides, Sieber, ex DC. l. c. 111 = Pultenæa flexilis.

DILOBEIA, Thon. Nov. Gen. Madag. 7 (1806). *PROTEACEAE*, Benth. & Hook. f. iii. 175.
Thouarsi, *Roem. & Schult. Syst.* iii. 476.—Madag.

DILOCHIA, Lindl. Gen. et Sp. Orch. 38 (1830) = **Arundina**, Blume (Orchid.).
pentandra, Reichb. f. in Bonplandia, v. (1857) 54 = A. pentandra.
Wallichii, Lindl. in Wall. Cat. n. 1952 = A. Wallichii.

DILOCHUS, Miq. Fl. Ind. Bat. iii. 669 (1858) = prace.

DILODENDRON, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 355. *SAPINDACEAE*.
bipinnatum, *Radlk. l. c.* 357.—Bras.

DILOMILIS, Rafin. Fl. Tellur. iv. 48 (1836) = **Octadesmia**, Benth. (Orchid.).
serrata, Rafin. l. c. = O. serrafolia.

DILOPHIA, T. Thoms. in Hook. Kew Journ. v. (1853) 19. *CRUCIFERAE*, Benth. & Hook. f. i. 89.
fontana, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 423.—China.
kaschgarica, *Rupr. Sert. Tianach.* 40.—As. centr.
salsa, *T. Thoms. in Hook. Kew Journ.* v. (1853) 20.—Tibet. occ.

DILWYNIA, Pers. Syn. ii. 688 (1807), sphalm. = **Dillwynia**, Sm. (Legumin.).

DIMACRIA, Lindl. in Sweet, Geran. t. 46 (1820) = **Pelargonium**, l'Hérit. (Geraniac.).
affinis, Sweet, Hort. Brit. ed. I. 77 = P. affine.
Andrewsii, Sweet, l. c. = P. Andrewsii.
apiifolia, Sweet, l. c. = P. apioides.
aristata, Sweet, l. c. = P. aristatum.
astragalifolia, Sweet, Geran. t. 103 = P. astragalifolium.
auriculata, Sweet, l. c. 393 = P. longifolium.
barbata, Sweet, Hort. Brit. ed. I. 77 = P. barbatum.
bipartita, Sweet, Geran. t. 142 = P. bipartitum.
carnea, Sweet, Hort. Brit. ed. I. 77 = P. carneum.
coronillaefolia, Sweet, l. c. = P. coronillaefolium.
depressa, Sweet, Geran. t. 390 = P. longifolium.
elegans, Sweet, l. c. t. 302 = P. elegantulum.
fissifolia, Sweet, Hort. Brit. ed. I. 77 = P. fissifolium.
floribunda, Sweet, l. c. = P. floribundum.
foliolosa, Sweet, Geran. t. 108 = P. astragalifolium.
fulgens, Sweet, Hort. Brit. ed. III. 110 = P. homonymum.
heracleifolia, Sweet, l. c. = P. heracleifolium.
imbuta, Sweet, l. c. ed. I. 77 = P. imbutum.
lanceaefolia, Sweet, Geran. t. 387 = P. lanceolatum.
lanceolata, Sweet, Hort. Brit. ed. III. 110 = P. lanceolatum.
linaria, Sweet, l. c. ed. I. 77 = P. lineare.
longiflora, Sweet, l. c. ed. III. 110 = P. longifolium.
longifolia, Sweet, l. c. = P. longifolium.
oxalidifolia, Sweet, l. c. ed. I. 77 = P. oxalidifolium.
punctata, Sweet, Geran. t. 46 = P. vicinifolium.
punctata, Sweet, Hort. Brit. ed. I. 77 = P. punctatum.
radiata, Sweet, l. c. = P. radiatum.
recurvata, Sweet, l. c. = P. recurvatum.
reflexa, Sweet, l. c. = P. reflexum.
revoluta, Sweet, l. c. = P. revolutum.
rumicifolia, Sweet, Geran. t. 318 = P. rumicifolium.
setigera, Sweet, Hort. Brit. ed. III. 110 = P. setigerum.
Smithiana, Sweet, Geran. t. 358 = P. Smithianum.
spathulata, Sweet, Hort. Brit. ed. I. 77 = P. spathulatum.
sulphurea, Sweet, Geran. t. 163 = P. sulphureum.
tenella, Sweet, Hort. Brit. ed. I. 77 = P. tenellum.
undulata, Sweet, l. c. = P. undulatum.
virginea, Sweet, l. c. = P. longifolium.

DIMANISA, Rafin. Fl. Tellur. ii. 83 (1836) = **Dianthera**, Gronov. (Acanthac.).
latifolia, Rafin. l. c. 84 = Dianthera nodosa.
oblongata, Rafin. l. c. = Beloperone plumbaginifolia.

DIMEIANDRA, Rafin. Neogenyt. 2 (1825) = **Amaranthus**, Tourn.

DIMEIANTHUS, Rafin. Fl. Tellur. iii. 41 (1836) = prec.

DIMEIOSTEMON, Rafin. Neogenyt. 4 (1825) = **Andropogon**, Linn. (Gramin.).
macrurus, Rafin. in Ser. Bull. Bot. i. (1830) 221 = A. macrourus.
sessiliflorus, Rafin. l. c. (quid?).
tetrastachys, Rafin. l. c. = A. virginicus.
vaginatus, Rafin. l. c. = A. virginicus.

DIMENOPS, Rafin. Atl. Journ. 144 (1832) = **Krameria**, Loefl. (Polygalac.).
lanceolata, Rafin. l. c. = K. lanceolata.

DIMENOSTEMMA, Steud. Nom. ed. II. i. 509 (1840) = **Dimerostemma**, Cass. (Compos.).

DIMEREZA, Labill. Sert. Austr. Caled. 51. t. 51 (1824) = **Cupania**, Plum. (Sapindac.).
glauca, Labill. l. c. = C. glauca.

DIMERIA, R. Br. Prod. 204 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1126.
DIDACTYLON, Zoll. & Moritzi, Syst. Verz. Zoll. 99 (1846).
HAPLACHNE, J. & C. Presl, Rel. Haenk. i. 284. t. 38 (1830).
PSILOSTACHYS, Steud. Syn. Pl. Gram. 418 (1855).
PTERYGOSTACHYUM, Nees, ex Steud. l. c. 418 (1855).
acinaciformis, *R. Br. Prod.* 204.—Austral.
diandra, Griff. Notul. iii. 71 = ornithopoda.

DIMERIA:—
filiformis, Hochst. ex Miq. in Verh. Nederl. Inst. III. iv. (1851) 35 = ornithopoda.
fuscescens, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 335.—Ind. or.
Hohenackeri, *Hochst. ex Miq. in Verh. Nederl. Inst.* III. iv. (1851) 35.—Ind. or.
laxiuscula, Thw. ex Trimen, in Journ. Bot. xxiii. (1885) 272 = gracilis
ornithopoda, *Trin. Fund. Agrost.* 167.—As. trop.; Austral.
plicosissima, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 336.—Ins. Marian.
psilobasis, F. Muell. Fragm. vii. 104 = ornithopoda.
pusilla, *Thw. Enum. Pl. Zeyl.* 369.—Zeylan.
stipaeformis, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 288 = ornithopoda.
tenera, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 336 = ornithopoda.

DIMERIA, Endl. Gen. 81, in syn. (1836) sphalm. = DIMESIA, Rafin. = **Hierochloë**, S. G. Gmel. (Gramin.).

DIMEROSTEMMA, Cass. in Bull. Soc. Philom. (1817) 11. *COMPOSITAE*, Benth. & Hook. f. ii. 376.
brasiliana, *Cass. l. c.* (1818) 58.—Bras.

DIMESIA, Rafin. in Am. Monthly Mag. (1818) 175 = **Hierochloë**, S. G. Gmel. (Gramin.).
fragrans, Rafin. l. c. = H. borealis.
monticola, Rafin. l. c. = Holcus monticola.

DIMETIA, Meisn. Gen. 160 (1838) = **Hedyotis**, Linn. (Rubiac.).
recurva, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 154 = H. recurva.

DIMETOPIA, DC. Prod. iv. 71 (1830) = **Trachymene**, Rudge (Umbellif.).
anisocarpa, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) 11. 29 = T. australis.
cyanopetala, F. Muell. Fragm. i. 231 = T. cyanopetalea.
eriocarpa, F. Muell. in Trans. Vict. Inst. i. (1855) 127 = T. eriocarpa.
graveolens, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) 11. 29 = T. australis.
hirta, Benth. in Enum. Pl. Hueg. 54 = T. pilosa.
homocarpa, Bunge, in Bot. Zeit. v. (1847) 186 = T. pilosa.
isocarpa, Bartl. Del. Sem. Hort. Gotting. (1846) 3 = T. pilosa.
Preissii, Bunge, in Lehm. Pl. Preiss. i. 284 = T. pilosa.
pusilla, DC. Prod. iv. 71 = T. pilosa.
Walpersii, Bunge, Del. Sem. Hort. Dorp. (1846).—Cf. Linnaea, xxiv. (1851) 156 = T. pilosa.

DIMIA, Spreng. Anleit. ii. i. 485 (1817) = **Dasmia**, R. Br. (Asclep.).

DIMITOPIA, D. Dietr. Syn. ii. 921 (1840) = DIMETOPIA, DC. = **Trachymene**, Rudge (Umbellif.).

DIMOCARPUS, Lour. Fl. Cochinch. 233 (1790), partim = **Nephelium**, Linn. (Sapindac.).
crinitus, Lour. l. c. 234 = N. lappaceum.
informis, Lour. l. c. = Euphoria informis.
Litchi, Lour. l. c. 233 = N. Litchi.
Longan, Lour. l. c. = N. Longana.
Pulasiki, Lour. l. c. Cat. Pl. Cycl. 81 = N. Longana.
undulatus, Wight, ex Wight & Arn. Prod. 113 = N. Longana.
verrucosus, Roxb. ex Wall. Cat. sub n. 8052 = Sapindus Denura.

DIMORPHA, D. Dietr. Syn. ii. 1228 (1840) = **Diamorpha**, Nutt. (Crassulac.).

DIMORPHA, Schreb. Gen. 498 (1791) = **Eperua**, Aubl. (Legumin.).
falcata, Spr. in Rees, Cycl. xi. n. 3 = E. falcata.
grandiflora, Willd. Sp. Pl. iii. 971 = E. grandiflora.
tomentosa, Willd. l. c. = Cradia Parivoa.

DIMORPHANDRA, Schott, in Spreng. Syst. iv. App. 404 (1827). *LEGUMINOSAE*, Benth. & Hook. f. i. 547.
MORA, Schomb. ex Benth. in Trans. Linn. Soc. xvii. (1839) 210. t. 16.

DIMORPHANDRA:—
biretusa, Tul. in Arch. Mus. Par. iv. (1844) 134 = Bras. Gardneriana.
exaltata, *Schott, in Spreng. Syst. iv. App.* 404.—Bras.
excelsa, Baill. Hist. Pl. ii. 167 = D. Mora.
Gardneriana, *Tul. in Arch. Mus. Par.* iv. (1844) 135. —Bras.
guianensis, Baill. Hist. Pl. ii. 149 = D. Mora.
latifolia, *Tul. in Arch. Mus. Par.* iv. (1844) 139.—Bras.
macrostachya, *Benth. in Hook. Journ. Bot.* ii. (1840) 101.—Guiana.
mollis, *Benth. l. c.* 102.—Bras.
Mora, *Benth. & Hook. f. Gen.* i. 588.—Guian.
oleifera, *Triana, ex Hemsl. Biol. Centr. Am. Bot.* i. 345.—Panama.
parviflora, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 251.—Bras.
pennigera, *Tul. in Arch. Mus. Par.* iv. (1844) 137. —Bras.
speciosa, Spruce, ex Benth. in Mart. Fl. Bras. xv. ii. 250 = pennigera.
unijuga, *Tul. in Arch. Mus. Par.* iv. (1844) 135.—Bras.
vernicosa, *Spruce, ex Benth. & Hook. f. Gen.* i. 588; *et Benth. in Mart. Fl. Bras.* xv. ii. 249.—Bras.

DIMORPHANTHES, Cass. in Bull. Soc. Philom. (1818) 30; et in Dict. Sc. Nat. xiii. 254 (1819) = Conyza, Linn. (Compos.).
aegyptiaca, Cass. ii. cc. 30, 256 = C. aegyptiaca.
ambigua, Presl, Fl. Sic. p. xxviii = Erigeron linifolius
angustifolia, Cass. in Dict. Sc. Nat. xxv. 95 = C. fastigiata, Erigeron linifolius.
apurensis, Cass. l. c. 96 = Erigeron apurensis.
bidentata, Cass. l. c. xiii. 256 = C. salicephala.
chinensis, Cass. in Bull. Soc. Philom. (1818) 30 = Erigeron apurensis.
coronopifolia, Cass. in Dict. Sc. Nat. xxv. 96 = C coronopifolia.
crispa, Repr. in Bull. Phys.-Math. Acad. Petersb. xiv. (1856) 285 = Erigeron linifolius.
floribunda, Cass. in Dict. Sc. Nat. xxv. 96 = Erigeron bonariensis.
Gouani, Cass. in Bull. Soc. Philom. (1818) 30; et l. c. xiii. 255 = C. Gouani.
hispida, Cass. l. c. xxv. 96 = C. hispida.
linifolia, Repr. in Bull. Phys.-Math. Acad. Petersb. xiv. (1856) 284 = Erigeron linifolius.
myosotifolia, Cass. in Dict. Sc. Nat. xxv. 96 = C. myosotifolia.
obtusa, Cass. l. c. = C. obtusa.
procera, Cass. in Bull. Soc. Philom. (1821) 175 = C. chilensis.
pulchella, Cass. in Dict. Sc. Nat. xxv. 96 = C. sophiaefolia.
sicula, Cass. in Bull. Soc. Philom. (1818) 30; et l. c. xiii. 255 = Jasonia sicula.
sophiaefolia, Cass. l. c. xxv. 96 = C. sophiaefolia.
stipulacea, Cass. l. c. 94 = seq.
stipulata, Cass. in Bull. Soc. Philom. (1821) 176 = C. incisa.
thesifolia, Cass. in Dict. Sc. Nat. xxv. 96 = C. thesiifolia.

DIMORPHANTHES, Meisn. Gen. Comm. 359 (1843) = seq.

DIMORPHANTHUS, Miq. Comm. Phytogr. 95. t. 19 (1840) = **Aralia**, Tourn.
edulis, Miq. l. c. 96 = A. cordata.
sieboldii, Miq. l. c. 95 = A. spinosa.
mandshuricus, Maxim. Prim. Fl. Amur. 133 = A. mandshurica.

DIMORPHOCALYX, Thw. Enum. Pl. Zeyl. 276 (1861). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 301.
glabellus, *Thw. l. c.*—Zeylan.

DIMORPHOCHLAMYS, Hook. f. in Benth. & Hook. f. Gen. i. 827 (1867). *CUCURBITACEAE*, Benth. & Hook. f. i. c.
Mannii, *Hook. f. l. c.*—Afr. trop.

DIMORPHOCOMA, F. Muell. & Tate, in Trans. Roy. Soc. S. Austral. vi. (1883) 107. *COMPOSITAE*.
minutula, *F. Muell. & Tate, l. c.*—Austral.

DIMORPHOLEPIS, A. Gray, in Hook. Kew Journ. iv. (1852) 227; et in Hook. Ic. Pl. t. 856 = **Heliptorum**, DC. (Compos.).
australis, A. Gray, ll. cc. = H. Dimorpholepis.

DIMORPHOPETALUM, Bert. in 'Merc. Chil. (1829)'. —Cf. Bull. Férussac, xx. (1830) 110; et Am. Journ. Sc. xx. (1831) 256, nomen = **Tetilla**, DC. (Saxifrag.).
Tetilla, Bert. ll. cc. = T. hydrocotylacifolia.

DIMORPHOSTACHYS, Fourn. in Comptes-Rendus Acad. Par. lxxx. (1875) 441 = **Panicum**, Linn. (Gramin.).
adspersa, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 490.—Mexic.
Botterii, Fourn. ex Hemsl. l. c. 14.—Mexic.
Drummondii, Fourn. ex Hemsl. l. c.—Mexic.
Ghiesbreghtii, Fourn. ex Hemsl. l. c.—Mexic.
Langei, Fourn. ex Hemsl. l. c.—Mexic.
monostachya, Fourn. ex Hemsl. l. c.—Mexic.
oajacensis, Fourn. ex Hemsl. l. c.—Mexic.
paspaloides, Fourn. ex Hemsl. l. c.—Mexic.
Schaffneri, Fourn. ex Hemsl. l. c.—Mexic.
variabilis, Fourn. ex Hemsl. l. c.—Mexic.

DIMORPHOTHECA, Vaill. ex Linn. Syst. ed. 1 (1735); Moench, Meth. 580 (1794). *COMPOSITAE*; Benth. & Hook. f. ii. 353.
ACANTHOTHECA, DC. Prod. vi. 73 (1837).
ARNOLDIA, Cass. in Dict. Sc. Nat xxx. 330 (1824).
BLAXIUM, Cass. l. c. 328 (1824).
CASTALIS, Cass. l. c. 332 (1824).
GATTENHOFFIA, Neck. Elem. i. 80 (1790).
LESTIBOSIA, Neck. l. c. 40 (1790).
METERINA, Cass. in Bull. Soc. Philom. (1818) 167.
annua, Less. Syn. Comp. 257 = pluvialis.
aurantiaca, *DC. Prod.* vi. 73.—Afr. austr.
Barbetae, *Harv. in Hook. Bot. Mag.* t. 5387.—Afr. austr.
calendulacea, *Harv. in Harv. & Sond. Fl. Cap.* iii. 423.—Afr. austr.
caulescens, *Harv. l. c.* 420.—Afr. austr.
chrysanthemifolia, *DC. Prod.* vi. 73.—Afr. austr.
chrysanthemoides, Harv. in Harv. & Sond. Fl. Cap. iii. 416, sphalm. = præc.
cuneata, *DC. Prod.* vi. 72.—Afr. austr.
dentata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 423.—Afr. austr.
Dregei, *DC. Prod.* vi. 73.—Afr. austr.
Dregei, Sch. Bip. in Flora, xxvii. (1844) 674 = cuneata.
Ecklonis, *DC. Prod.* vi. 71.—Afr. austr.
fruticosa, *DC. l. c.*—Afr. austr.
graminifolia, DC. l. c. = nudicaulis.
hybrida, DC. l. c. 70 = pluvialis.
incrassata, Moench, Meth. 585 = pluvialis.
integrifolia, *Harv. in Harv. & Sond. Fl. Cap.* iii. 424.—Afr. austr.
Krausii, Sch. Bip. in Flora, xxvii. (1844) 674 = nudicaulis.
leptocarpa, *DC. Prod.* vi. 72.—Afr. austr.
lilacina, *Regel & Herd. Ind. Sem. Hort. Petrop.* (1864) 16.—Afr. austr.
multifida, *DC. Prod.* vi. 72.—Afr. austr.
nudicaulis, *DC. l. c.* 71.—Afr. austr.
perennis, Less. Syn. Comp. 257 = nudicaulis.
pinnata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 423.—Afr. austr.
pluvialis, *Moench, Meth.* 585.—Afr. austr.
polyptera, *DC. Prod.* vi. 72.—Afr. austr.
scabra, DC. l. c. 73 = D. Tragus.
sinuata, *DC. l. c.*—Afr. austr.
Tragus, *DC. l. c.* 71.—Afr. austr.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* iii. 421.—Afr. austr.

DINACRIA, Harv. in Harv. & Sond. Fl. Cap. ii. 380 (1861–62). *CRASSULACEAE*; Benth. & Hook. f. i. 867.
filiformis, *Harv. l. c.* 381.—Afr. austr.

DINAEA, Delile, Fl. Egypt. 96. t. 11. f. 3 (1813) = sq.

DINEBA, Besser, Agrost. 90 (1813) = sq.

DINEBRA, DC. Cat. Hort. Monsp. 104 (1813), partim = Boutelouа, Lagg. (Gramin.).
aristidoides, H. B. & K. Nov. Gen. et Sp. i. 171 = B. aristidoides.
bromoides, H. B. & K. l. c. 172. t. 51 = B. bromoides.
chondrosioides, H. B. & K. l. c. 173. t. 53 = B. ovata.
curtipendula, DC. Cat. Hort. Monsp. 105 = B. racemosa.
repens, H. B. & K. Nov. Gen. et Sp. i. 172. t. 52 = B. bromoides.

DINEBRA, Jacq. Fragm. 77. t. 121. f. 1 (1809) *GRAMINEAE*, Benth. & Hook. f. iii. 1171.
aegyptiaca, Kunth, Enum. Pl. i. 271 = arabica.
americana, Besser, Agrost. 98, 160 = Boutelouа bromoides, juncifolia.
arabica, *Jacq. Fragm.* 77.—Afr. trop.; Ind. or.
brevifolia, *Steud. Syn. Pl. Gram.* 299.—Abyss.
calycina, Wight, ex Ness, in Proc. Linn. Soc. i. (1841) 95 = Tripogon Nessianus.
chloridea, J. & C. Presl, Rel. Haenk. i. 291 (= Eutrematis sp.).—Mexic.
cristata, *J. & C. Presl, l. c. i.* 293.—Mexic.
divaricata, Besser, Agrost. 98, 160 = Cutandia scleropoides.
hirsuta, *J. & C. Presl, Rel. Haenk. i.* 292.—Peruv.
hirta, Nees, ex Steud. Syn. Pl. Gram. 209 = Leptochloa hirta.
juncifolia, Steud. Nom. ed. II. i. 510 = Boutelouа juncifolia.
Lima, Besser, Agrost. 98, 160 = Wangenheimia disticha.
meliconder, Beauv. l. c. 160 = Boutelouа racemosa ?
paspaloides, Beauv. l. c. = præc. ?
retroflexa, Panz. in Denkschr. Acad. Muench. 1813 (1814) 270 = arabica.
secunda, Roem. & Schult. Syst. ii. 711 (= *Eutriana (Ledebouria)*).—Ins. S. Doming.
verticillata, Wight, ex Steud. Syn. Pl. Gram. 209 = Eragrostis Wightiana.

DINEMA, Lindl. Orch. Sel. 16 (1836); et Gen. et Sp. Orch. 111 (1881) = **Epidendrum**, Linn. (Orchid.).
palacearum, Lindl. Bot. Reg. (1840) Misc. 61 = E. auritum.
polybulbon, Lindl. Gen. et Sp. Orch. 111 = E. polybulbon.

DINEMAGONUM, A. Juss. in Arch. Mus. Par. iii. (1843) 585 (Monog. Malpig. 331). *MALPIGHIACEAE*; Benth. & Hook. f. i. 261.
albicaule, *Phil. in Linnaea*, xxxiii. (1864–65) 32.—Chili.
Bridgesianum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 587 (*Monog. Malpig.* 333).—Chili.
Gayanum, *A. Juss. ll. cc.* 585, 331.—Chili.

DINEMANDRA, A. Juss. ex Endl. Gen. 1059 (1840). *MALPIGHIACEAE*, Benth. & Hook. f. i. 261.
ericoides, *A. Juss. in Arch. Mus. Par.* 582. t. 19; (*Monog. Malpig.* 328).—Peruv.
glaberrima, *A. Juss. ll. cc.* 583, 329.—Chili.
ramosissima, *Phil. Fl. Atac.* 12.—Chili.
strigosa, *Phil. l. c.*—Chili.
subaptera, *Phil. l. c.*—Chili.

DINETUS, Buch.-Ham. ex D. Don, in Sweet, Brit. Flow. Gard. t. 127 (1825) = **Porana**, Burm. (Convolv.).
paniculata, Sweet, Hort. Brit. ed. I. 389 = P. paniculata.
racemosa, Sweet, Brit. Flow. Gard. t. 127 = P. racemosa.

DINOCHLOA, Buese, in Miq. Pl. Jungh. 887 (1851–55). *GRAMINEAE*, Benth. & Hook. f. iii. 1214.
andamanica, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) 11. 253.—Ins. Andam.
ciliata, *Kurz, l. c.*—Ins. Philipp.
Maclellandii, *Kurz, l. c.*—Burma.
Tjankorreh, *Buese, in Miq. Pl. Jungh.* 388.—Java.

DINOPHORA, Benth. in Hook. Niger Fl. 555 (1849). *MELASTOMACEAE*, Benth. & Hook. f. i. 745.
spenneroides, *Benth. l. c.*—Afr. trop.

DINOSCHERIS, Griseb. in Goett. Abh. xxiv. (1879) 213. *COMPOSITAE*.
salicifolia, *Griseb. l. c.* 214.—Reg. Argent.

DIOCLEA, H. B. & K. Nov. Gen. et Sp. vi. 437 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 586.
CARPIDOTROPIS, Walp. in Linnaea, xiv. 594 (1840).
HYMENOSPRON, Spreng. Syst. iv. Cur. Post. 283 (1827).
LEPIDAMPHORA, Zoll. ex Miq. Fl. Ind. Bat. i. 1. 217 (1855).
apurensis, *H. B. & K. l. c.*—Venezuela.
argentea, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 420 = Cratylia floribunda.
bicolor, *Benth. in Ann. Wien. Mus.* ii. (1838) 133.—Bras.
Boykinii, *A. Gray, ex S. Wats. Bibl. Ind. N. Am Bot.* 219.—Am. bor.
casuinoides, *Desf. Cat. Hort. Par.* ed. III. 410.—Hab. ?
coriacea, *Benth. in Ann. Wien. Mus.* ii. (1838) 133.—Bras.
Fergusonii, Thw. Enum. Pl. Zeyl. 412 = reflexa.
funalis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 59.—Peruv.
galactioides, *Griseb. Cat. Pl. Cub.* 76.—Cuba.
glabra, Benth. in Ann. Wien. Mus. ii. (1838) 138.—Bras.
glycinoides, *Hort. Belg. ex Pasq. Cat. Orto Bot. Nap.* 86. nomen (cf. casuinoides).—Hab. ?
grandiflora, *Mart. ex Benth. in Ann. Wien. Mus.* ii (1838) 132.—Bras.
guianensis, *Benth. l. c.* 134.—Guiana.
Jacquiniana, *DC. Prod.* ii. 403.—Ins. Martinic.
javanica, Benth. in Miq. Pl. Jungh. 286 = reflexa.
lasiocarpa, *Benth. in Ann. Wien. Mus.* ii. (1838) 133.—Panama.
lasiophylla, *Benth. l. c.* 134.—Bras.
latifolia, *Benth. l. c.* 132.—Bras.
marginata, *Benth. in Mart. Fl. Bras.* xv. i. 166.—Bras.
mollis, *DC. Prod.* ii. 408.—Hab. ?
panamensis, Duchass. & Walp. in Flora, xxxvi. (1853) 229 = guianensis.
panamensis, Seem. Hemsl. Biol. Centr. Am. Bot. i. 302 = reflexa.
panamensis, *Walp. Ann.* vii. 785.—Panama.
pilifera, *Tul. in Arch. Mus. Par.* iv. (1844) 71.—Bras.
purpurea, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 59.—Bras.
reflexa, *Hook. f. in Hook. Niger Fl.* 306.—Afr. et As. trop.
rostrata, *Benth. in Ann. Wien. Mus.* ii. (1838) 133.—Bras.
rubiginosa, Tul. in Arch. Mus. Par. iv. (1844) 72 = rufescens.
rudolphioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 176.—Cuba.
rufescens, *Benth. in Ann. Wien. Mus.* ii. (1838) 133.—Bras.
Schottii, *Benth. l. c.* 134.—Bras.
sericea, *H. B. & K. Nov. Gen. et Sp.* vi. 437.—N. Granat.
violacea, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 132.—Bras.

DIOCLEA, Spreng. Syst. i. 503 (1825) = **Arnebia**, Forsk. (Borag.).
hispidissima, Spreng. l. c. 556 = A. hispidissima.

DIOCTIS, Rafin. Fl. Tellur. iii. 15 (1836) = **Polygonum**, Tourn.
bicorne, Rafin. l. c. = P. pensylvanicum.
equisetifolia, Rafin. l. c. = P. equisetiforme.
maculatum, Rafin. l. c. = P. lapathifolium ?
pensylvanica, Rafin. l. c. = P. pensylvanicum.
vernum, Rafin. l. c. = P. lapathifolium.

DIODELILS, Rafin. Fl. Tellur. iii. 82 (1836) = **Gardoquia**, Ruiz & Pav. (Labiat.).
coccinea, Rafin. l. c. 83 = G. Hookeri.

DIODIA, Gronov. in Linn. Cor. Gen. 11 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 148.
alata, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 11 = Dasycephala alata.
anthospermoides, Cham. & Schlecht. in Linnaea, iii. (1828) 548 = polymorpha.
arenosa, *DC. Prod.* iv. 544.—Bras.
articulata, DC. l. c. = hymenopifolia.
auriculosa, *Bertol. Misc. Bot.* x. 25.—Am. bor.
barbata, DC. Prod. iv. 563 = Dasycephala barbata.

DIODIA:—
berlingera, *Hook. & Arn. Bot. Beech. Voy.* 295.—Mexic.
bogotensis, *Cham. & Schlecht. in Linnaea*, iii. (1828) 347.—N. Granat.
brasiliensis, Spreng. Syst. i. 406 = polymorpha.
brevicaule, Benth. in Hook. Niger Fl. 424 = sarmentosa.
carnosa, *Hochst. in Flora*, xxvii. (1844) 555.—Afr. austr.
conferta, *DC. Prod.* iv. 563.—Bras.
crassifolia, *Benth. Bot. Voy. Sulph.* 108.—Mexic.
cymosa, *Cham. in Linnaea*, ix. (1834) 217.—Bras.
Dasycephala, *Cham. & Schlecht. l. c.* iii. (1828) 848.—Bras.
discolor, *DC. Prod.* iv. 564.—Guiana.
domingensis, *DC. l. c.* 563.—Ins. S. Doming.
elongata, E. Mey. ex Hochst. in Flora, xxvii. (1844) 555 = Spermacoce natalensis.
foliosa, Peyr. in Sitzb. Acad. Wien. Math.-Nat xxxvii. (1860) 579 = Spermacoce natalensis.
Franhii, Stend. & Hochst. ex Presl, Bot. Bemerk. 86 = Hedyotis Franhii.
glabra, Pers. Syn. i. 124 = Spermacoce glabra.
glabra, *Willd. ex H. B. & K. Nov. Gen. et Sp.* iii. 348, in syn.—Peruv.
grandiflora, DC. Prod. iv. 563 = rigida.
hirsuta, Pursh, Fl. Am Sept. i. 106 = virginiana.
hispida, Muhl. ex DC. Prod. iv. 562 = praec.
hispidula, A. Rich. ex DC. l. c. 565 = Dasycephala hispidula.
hysopifolia, *Cham. & Schlecht. in Linnaea*, iii. (1828) 850.—Venezuela.
indecora, DC. Prod. iv. 565 = Dasycephala indecora.
Kirkii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 280.—Afr. trop.
lanigera, *Pohl, ex DC. Prod.* iv. 574.—Bras.
latifolia, DC. l. c. 563 = rigida.
lippioides, *Griseb. Cat. Pl. Cub.* 141.—Cuba.
littoralis, Peyr. in Sitzb. Acad. Wien. Math.-Nat. xxxviii. (1860) 579 = sarmentosa.
maritima, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 75.—Afr. trop.
mitens, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 283.—Porto Rico.
multiflora, *DC. Prod.* iv. 564.—Bras.
muriculata, DC. l. c. = D. Radula.
oblonga, Rafin. Fl. Ludov. 76 = virginiana.
orientalis, Koen. ex Roth. Fl. Ind. ed. Carey, i. 380 = Hydrophylax maritima.
palustris, Cham. & Schlecht. in Linnaea, iii. (1828) 847 = alata.
paradoxa, *Cham. l. c.* ix. (1834) 216.—Bras.
pedunculata, Spreng. Syst. i. 405 = Manettia Pseudodiodia.
pilosa, Schum. & Thonn. Beskr. Guin. Pl. 74 = Spermacoce pilosa.
polymorpha, *Cham. & Schlecht. in Linnaea*, iii. (1828) 844.—Bras.
polysyta, DC. Prod. iv. 563 = rigida.
procumbens, Sw. ex DC. l. c. 562 = prostrata.
prostrata, *Sw. Prod. Veg. Ind. Occ.* 30.—Jamaic.
radicans, Cham. & Schlecht. in Linnaea, iii. (1828) 850 = maritima.
Radula, *Cham. & Schlecht. l. c.* 842.—Bras.
reptans, F. Muell. in Trans. Vict. Inst. i. (1855) 128 = Nertera reptans.
rigida, *Cham. & Schlecht. l. c.* 841.—Am. austr.
rigida, Cham. & Schlecht. l. c. partim = setigera.
romanifolia, *Pohl, ex DC. Prod.* iv. 564.—Bras.
rubricosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 281.—Afr. trop.
rudis, Mig. in Linnaea, xvii. (1844) 300 = rigida.
saponariodes, *Presl, Bot. Bemerk.* 87.—Bras.
sarmentosa, *Sw. Prod. Veg. Ind. Occ.* 30.—Ind. occ.
scabra, *Schum. & Thonn. Beskr. Guin. Pl.* 76.—Afr. trop.
scandens, Benth. in Hook. Journ. Bot. iii. (1841) 286 = sarmentosa.
scandens, *Sw. Prod. Veg. Ind. Occ.* 30.—Ins. S. Doming.
senensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 289. —Afr. trop.
setigera, *DC. Prod.* iv. 563.—Bras.
simplex, *Sw. Prod. Veg. Ind. Occ.* 29.—Jamaic.
spicata, Mlq. Stirp. Surin. Sel. 179 = Dasycephala spicata.
surinamensis, *Mig. l. c.* 180.—Guiana.
teres, *Walt. Fl. Carol.* 87.—Am. bor.

DIODIA:—
tetracocca, Hemsl. Diagn. Pl. Nov. 32 = Crusea allococca.
tetragona, Walt. Fl. Carol. 87 = virginiana.
triccrca, Torr. & Gray, Fl. N. Am. ii. 80 = Crusea allococca.
vaginalis, *Benth. in Hook. Niger Fl.* 424.—Afr. trop.
verticillata, *Vahl, Symb. Bot.* ii. 28.—Ins. S. Crucis.
villosa, *Mog. & Sesse, ex DC. Prod.* iv. 562.—Mexic.
virginiana, *Linn. Sp. Pl.* 104.—Am. bor.
virginica, Linn. Syst. ed. XII. 116; Jacq. Ic. Pl. Rar. t. 29 = praec.

DIODOIS, Pohl, in Flora, vii. (1826) 188, nomen = **Psyllocarpus**, Mart. & Zucc. (Rubiac.).

DIODONTA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. 260 (1841) = **Coreopsis**, Linn. (Compos.).
aristosa, Nutt. l. c. = C. aristosa.
aurea, Nutt. l. c. = C. aurea.
bidentoides, Nutt. l. c. 261 = C. bidentoides.
coronata, Nutt. l. c. 260 = C. coronata.
involucrata, Nutt. l. c. 261 = C. involucrata, *Nutt. leptophylla*, Nutt. l. c. 260 = C. aurea.
mitis, Nutt. l. c. = C. aurea.

DIODONTIUM, F. Muell. in Hook. Kew Journ. ix. (1857) 19 = **Glossogyne**, Cass. (Compos.).
filifolium, F. Muell. l. c. = G. filifolia.

DIODOSPERMA, H. Wendl. in Bot. Zeit. xxxvi. (1878) 118 = **Trithrinax**, Mart. (Palm.).
Burity, H. Wendl. l. c. = T. schizophylla.

DIOLENA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 329. *MELASTOMACEAE*, Benth. & Hook. f. i. 756.
auriculata, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 81.—N. Granat.
agrimonioides, *Triana, l. c.*—N. Granat.
hygrophyla, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 329.—Venezuela.
oblique, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 81.—N. Granat.
pilosiúscs, *Triana, l. c.*—N. Granat.
spicata, *Triana, l. c.*—N. Granat.

DIOMEDEA, Bert. ex Colla, in Mem. Acc. Torino, xxxvii. (1835) 35. t. 31 = **Helianthus**, Linn. (Compos.).
thurifera, Bert. ex Colla, l. c. 37. t. 31 = H. thurifer.

DIOMEDEA, Cass. in Bull. Soc. Philom. (1817) 175 = **Borrichia**, Adans. (Compos.).
argentea, H. B. & K. Nov. Gen. et Sp. iv. 218 = B. argentea.
bidentata, Cass. in Dict. Sc. Nat. xliii. 288 = B. frutescens.
glabrata, H. B. & K. Nov. Gen. et Sp. iv. 218 = B. arborescens.
indentata, Cass. in Dict. Sc. Nat. xlii. 284 = B. arborescens.
linearis, Cass. l. c. 288, in obs. = B. frutescens.

DIOMEDES, Haw. in Phil. Mag. lxii. (1823) 440 = **Maraisseus**, Tourn. (Amaryll.).
Maclayi, Haw. ex Herb. in Journ. Hort. Soc. ii. (1847) 28 = N. Macleayi.
major, Haw. in Phil. Mag. lxii. (1823) 441 = N. Macleayi.
minor, Haw. l. c. 440 = N. Macleayi.
Parkinsoni, Haw. Monog. 5 = N. Macleayi.
Sabini x, Haw. ex Herb. in Journ. Hort. Soc. ii. (1847) 28 = N. Macleayi.

DION, Lindl. Bot. Reg. (1848) Misc. 59 = **Dioon**, Lindl. (Cycad.).

DIONAEA, Ellis, in Nov. Act. Soc. Sc. Upsal. I. (1770) 98. t. 8. *DROSERACEAE*, Benth. & Hook. f. i. 663.
corymbosa, Rafin. Med. Fl. ii. 217 = muscipula.
muscipula, *Ellis, ex Linn. Mant. ii.* 288.—Am. bor.
sensitiva, Salisb. Prod. 221 = muscipula.
sessiliflora, Rafin. Med. Fl. ii. 217 = muscipula.
uniflora, Rafin. Att. Journ. 78 = muscipula.

DIONEA, Rafin. Med. Fl. ii. 217 (1830) = praec.

DIONICHA, Triana, in Trans. Linn. Soc. xxviii. (1871) 61 = seq.

DIONYCHA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 48. *MELASTOMACEAE*, Benth. & Hook. f. i. 745.
Bojerii, *Naud. l. c.*—Madag.

DIONYCHIA, Benth. & Hook. f. Gen. i. 745 (1867) = praec.

DIONYSIA, Fenzl, in Flora, xxvi. (1843) i. 389. *PRIMULACEAE*, Benth. & Hook. f. ii. 632.
GREGORIA, Duby, Bot. Gall. i. 388 (1828); et in DC Prod. viii. 48 (1844).
MACROSYPHONIA, Duby, in Mém. Soc. Phys. Genèv. (1844) 426.
MACROTYBUS, Dulac, Fl. Hautes-Pyr. 425 (1867).
aretioides, *Boiss. Diagn.* Ser. I. vii. 68.—Persia.
Aucheri, Boiss. Fl. Orient. iv. 19 = odora.
bryoides, *Boiss. l. c.* 66.—Persia.
bryoides, Boiss & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 143 = curviflora.
caespitosa, *Boiss. Diagn.* Ser. I. vii. 67.—Persia.
curviflora, *Bunge, in Bull. Acad. Pétersb.* xvi. (1871) 561.—Persia.
diapensiaefolia, *Boiss. Diagn.* Ser I vii. 65.—Persia
drabaefolia, *Bunge, in Bull. Acad. Pétersb.* xvi. (1871) 558.—Persia.
Kotschyi, *Bunge, l. c.* 560.—Persia.
Michauxii, *Boiss. Diagn.* Ser. I. vii. 67.—Persia.
odora, *Fenzl, in Flora*, xxvi. (1843) i. 389.—Assyria
revoluta, *Boiss. Diagn.* Ser. I. vii. 66.—Persia.
rhaptodes, *Bunge, in Bull. Acad. Pétersb.* xvi. (1871) 562.—Persia.
tapetodes, *Bunge, l. c.* 560.—Persia.

DIOON, Lindl. Bot. Reg. (1843) Misc. 59 (Dion). *CYCADACEAE*, Benth. & Hook. f. iii. 445.
PLATYZAMIA, Zucc. in Abh. Akad. Muench. iv (1845) 23. t. 4.
aculeatum, Lem. in Illustr. Hortic. ii. (1855) Misc. 92 = edule.
angustifolium, Miq. in Tijdschr. Wetensch. i. (1848) 37 = edule.
edule, *Lindl. Bot. Reg.* (1843) *Misc.* 59.—Mexic.
imbricatum, Miq. in Tijdschr. Wetensch. i. (1848) 36 = edule.
spinulosum, *Dyer, ex Eichl. in Gart. Zeit.* (1888) 411. —Mexic.
strobilaceum, *Lem. in Illustr. Hortic.* x. (1863) *Misc* 4.—Hab. ?

DIOPOGON, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 46 (1868) = **Sempervivum**, Rupp. (Crassulac.).
Allionii, Jord. & Fourr. l. c. = S. hirtum.
Heuffelii, Jord. & Fourr. l. c. 47 = S. patens.
straminens, Jord. & Fourr. l. c. = S. stramineum.

DIORIMASPERMA, Rafin. Sylva Tellur. 111 (1838) = **Cleome**, Linn. (Capparid.).
violacea, Rafin. l. c. = C. graveolens.

DIORYKTANDRA, Hassk. Retzia, 125 (1855) = **Alsodeia**, Thou. (Violar.).
Roxburghii, Hassk. l. c. = A. Roxburghii.

DIOSANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 87 = **Dianthus**, Linn. (Caryoph.).
arenarium, St. Lag. 194 = D. Amoria.
carthusianicum, St. Lag. l. c. = D. Carthusianorum.
cephalotum, St. Lag. l. c. = D. Cephalotes.
libanotum, St. Lag. l. c. = D. Libanotis.

DIOSCORIDIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 175 = seq.

DIOSCOREA, Plum. ex Linn. Gen. ed. I. 306 (1737). *DIOSCOREACEAE*, Benth. & Hook. f. iii. 745.
BOEDEREA, Miegér. in Bull. Soc. Bot. Fr. xiii. (1866) 374.
BOTRYOSICYOS, Hochst. in Flora, xxvii. (1844) Bes. Beil. 8.
ELEPHANTODON, Salisb. Gen. Pl. Fragm. 12 (1866).
EPIPETRUM, Phil. in Linnaea, xxxiii. (1864–65) 283.
HAMATRIS, Salisb. Gen. Pl. Fragm. 11 (1866).
HELMIA, Kunth, Enum. Pl. v. 414 (1850).
MERIONE, Salisb. Gen. Pl. Fragm. 12 (1866).
POLYNOME, Salisb. l. c. 12 (1866).

DIOSCOREA :—
RICOPHORA, Mill. Gard. Dict. ed. VI. 175. App. (1763).
SIEMONDAEA, Delponte, in Mem. Acc. Torin. Ser. II. xiv. (1854) 394.
STROPHIS, Salisb. Gen. Pl. Fragm. 12 (1866).
URUM, J. F. Gmel. Syst. 838 (1791).
abyssinica, *Hockst. ex Kunth, Enum. Pl.* v. 387.—Abyss.
aculeata, Balb. ex Kunth, l. c. 381 = cayennensis.
aculeata, Buch.-Ham. ex Wall. Cat. n. 5108 = spinosa.
aculeata, *Linn. Sp. Pl.* 1033.—As. trop.
acuminata, *Baker, in Journ. Linn. Soc.* xxi. (1885) 449.—Madag.
acuminata, *Steud. Nom.* ed. II. i. 510.—Chili.
acutangula, Buch.-Ham. ex Wall. Cat. n. 5109 = alata.
acutifolia, *Phil. in Linnaea*, xxix. (1857–58) 64.—Chili.
adenocarpa, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. i. 29.—Bras.
affinis, *Kunth, Enum. Pl.* v. 372.—Guiana
alata, *Linn. Sp. Pl.* 1033.—As. trop.
altissima, Lam. Encyc. iii. 281 = alata l. sativa.
altissima, Sieber, ex Presl, Bot. Bemerk. 116 = multiflora, *Presl.*
altissima, Steud. Nom. ed. II i. 511 = daemona.
amaranthoides, *Presl, Rel. Haenk.* i. 134.—Chili ; Peruv.
amasonum, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. i. 39.—Bras.
amara, Roxb. ex Wight, Ic. t. 811 = daemona.
anguina, *Roxb. Hort. Beng.* 72 ; *Fl. Ind.* iii. 803.—Ind. or.
angulata, *Roxb. ex Steud. Nom.* ed. II. i. 511, nomen (= praec. ?).—Ind. or.
angustifolia, *Lam. Encyc.* iii. 283.—Peruv.
anomala, *Griseb. in Mart. Fl. Bras.* iii. i. 31.—Bras.
arnacana, *Phil. in Anal. Univ. Chil.* (1872) 540.—Chili.
arenaria, *Kunth, Enum. Pl.* v. 344.—Chili.
artifolia, *Presl, Rel. Haenk.* i. 134.—Peruv.
aristolochiaefolia, *Poepp. Fragm. Syn.* 11.—Chili.
articulata, *Steud. Nom.* ed. II. i. 511.—Bras.
aspera, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 794.—Venezuela.
atropurpurea, *Roxb. Hort. Beng.* 72 ; *Fl. Ind.* iii. 800.—Burma ; Malaya.
auriculata, *Poepp. Fragm. Syn.* 13.—Chili.
Batatas, Decne. in Rev. Hortic. Sér. IV. iii. (1854) 248 = divaricata.
Bertheroana, Kunth, Enum. Pl. v. 381 = cayenensis.
Besseriana, *Kunth, l. c.* 345.—Chili.
Bicantacea, Buch.-Ham. ex Wall. Cat. n. 5107 = alata.
Bilbergiana, Kunth, Enum. Pl. v. 354 = macrostachya.
bolyjoncia, Blanco, Fl. Filip. ed. I. 800 = Fuerara phaseoloides.
bonariensis, *Tenore, Ind. Sem. Hort. Neap.* (1838) 3 ; (1839) 11. n. 6.—Reg. Argent.
brachyborya, *Poepp. Fragm. Syn.* 12.—Chili.
brachycarpa, *Schlecht. in Linnaea*, xvii. (1843) 109.—Mexic.
brasiliana, Poir. Encyc. Suppl. iii. 139 = brasiliensis.
brasiliensis, *Willd. Sp. Pl.* iv. 791.—Bras.
Bridgesii, *Griseb. ex Kunth, Enum. Pl.* v. 358.—Chili.
bryoniaefolia, *Poepp. Fragm. Syn.* 13.—Chili.
Buchanani, *Benth. in Hook. Ic. Pl.* t. 1397–98.—Afr. trop.
bulbifera, R. Br. Prod. 294 = sativa.
bulbifera, *Linn. Sp. Pl.* 1033.—As. trop.
bulbifera, Russ. ex Wall. Cat. n. 5106 = alata.
calystegioides, *Kunth, Enum. Pl.* v. 362.—Bras.
Cambessedesiana, Buch.-Ham. ex Wall. Cat. n. 5103 = spinosa.
campestris, *Griseb. in Mart. Fl. Bras.* iii. i. 30.—Bras.
cayennensis, Webb & Berth. Phyt. Canar. iii. 317 = Tamus edulis.
capillaris, *Hemsl. Biol. Centr. Am. Bot.* iii. 354.—Mexic.
carnosanna, *Kunth, Enum. Pl.* v. 354.—Venezuela.
cayennesis, *Lam. Encyc.* iii. 283.—Guiana.
chondrocarpa, *Griseb. in Mart. Fl. Bras.* iii. i. 34.—Bras.
circumvoluta, Hook. Bot. Mag. t. 2935 = Rajania brasiliensis.
cirrhosa, *Lour. Fl. Cochinch.* ii. 625.—Ind. or. ; Cochinch.

DIOSCOREA :—
Cliffortiana, Lam. Encyc. iii. 282 = sativa.
composita, *Hemsl. in Biol. Centr. Am. Bot.* iii. 354.—Mexic.
conferta, Vell. Fl. Flum. x. t. 122 = piperifolia.
convolvulacea, *Cham. & Schlecht. in Linnaea*, vi. (1831) 49.—Mexic.
coriacea, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 794.—Ecuador.
coriacea, Wight, ex Wall. Cat. n. 5104 = oppositifolia.
cornifolia, *Kunth, Enum. Pl.* v. 356.—Java.
cotinifolia, *Kunth, l. c.* 386.—Afr. austr.
crenata, Vell. Fl. Flum. x. t. 127 = sinuata.
cropheana, Buch.-Ham. ex Wall. Cat n. 5105 = glabra.
crinita, *Hook. f. Bot. Mag.* t. 6804.—Natal.
crispata, Roxb. *Hort. Beng.* 72 ; *Fl. Ind.* iii. 802.—Ind. or.
cruenta, Kunth, Enum. Pl. v. 379 = *crenata.*
crumenigera, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. i. 36.—Bras.
cuspidata, Balb. ex Kunth, Enum. Pl. v. 434 = pilosiuscula.
cuspidata, *Humb. & Bonpl. ex Willd Sp. Pl.* iv. 794.—Am. austr.
cuspidata, Klotzsch, ex Kunth, Enum. Pl. v. 429 (= *Helmia consanguinea*).—Guiana
cylindrica, *Burm. Fl. Ind.* 213.—Ind. or.
cymosula, Hemsl. Biol. Centr. Am. Bot. iii. 355 = spinosa.
cynanchifolia, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 156.—Bras.
daemona, *Roxb. Hort. Beng.* 72 ; *Fl. Ind.* iii. 805.—Ind. or. ; Malaya.
Decaisneana, *Carr. in Rev. Hortic.* (1865) 111.—China.
decemangularis, Buch.-Ham. ex Wall. Cat. n 5106 = sativa.
decipiens, *Presl, Rel. Haenk* i. 135.—Peruv.
deflexa, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 157.—Bras.
deltoidea, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 18 = sativa.
deltoidea, *Wall. Cat.* n. 5110.—Ind. or.
densiflora, *Hemsl. in Biol. Centr. Am. Bot.* iii. 356.—Mexic.
Deppei, Schiede, ex Schlecht. in Bot. Zeit. i. (1843) 890 = macrostachya.
Dieckiana, Buch.-Ham. ex Wall. Cat. n. 5109 = alata.
diacantha, *Zipp. ex Span. in Linnaea*, xv. (1841) 479, nomen.—Ins. Timor.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 611.—Sumatra.
digitata, Mill. Gard. Dict. ed. VIII. n. 6 = pentaphylla.
discolor, *Hort. Berol. ex Kunth, Enum. Pl.* v. 384 — Ann. austr.
divaricata, *Blanco, Fl. Filip.* ed. I. 797.—Ins. Philipp.
diversifolia, *Griseb. in Mart. Fl. Bras.* iii. i. 41.—Bras.
dodecaphora, Steud. Nom. ed. II. i. 511 = seq.
dodecaneura, *Vell. Fl. Flum.* x. t. 123.—Bras.
doryphora, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 94.—Ins. Formosa.
eburna, Lour. Fl. Cochinch. 625 = seq.
eburnea, *Lour. l. c. ed. Willd.* ii. 767.—Cochinch.
echinata, Herb. Buch.-Ham. ex Wall. Cat. n. 5103 = spinosa.
effusa, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 161.—Bras.
elegantula, *Kunth, Enum. Pl.* v. 366.—Guiana.
elephantopus, Spreng. Syst. iv. Cur. Post. 148 = Testudinaria elephantipes.
fascicularia, *Roxb. Hort. Beng.* 72 ; *Fl. Ind.* iii. 801.—Ind. or.
fastigeata, *C. Gay, Fl. Chil.* vi. 54.—Chili.
filiformis, *Blume, Enum. Pl. Jav.* i. 22.—Java.
filiformis, Griseb. in Mart. Fl. Bras. iii. i. 44 = Griebachii.
filipendula, Domb. ex Kunth, Enum. Pl. v. 341 = humifusa.
floribunda, *Mart. & Gal. in Bull. Acad. Brux.* ix. 11. (1842) 391.—Mexic.
fodinarum, *Kunth, Enum. Pl.* v. 405.—Bras.
fructa, *Griseb. in Mart. Fl. Bras.* iii. i. 44.—Bras.
furcata, *Griseb. l. c.* 45.—Bras.
Galeottiana, *Kunth, Enum. Pl.* v. 409.—Mexic.
galipaneasis, *Klotzsch, ex Kunth, l. c.* 417.—N. Granat.

DIOSCOREA :—
glabra, *Roxb. Hort. Beng.* 72 ; *Fl. Ind.* iii. 804.—Ind. or.
glabra, Wall. Cat. n. 5105 D, E = oppositifolia.
glabra, Wall. l. c. n. 5105 B = lanrifolia.
glandulosa, *Klotzsch, ex Kunth, Enum. Pl.* v. 352.—Bras.
glauca, Muhl. ex Beck, Bot. U. St. 355 = quaterna.
globosa, *Roxb. Hort. Beng.* 72 ; *Fl. Ind.* iii. 797.—Ind. or.
Goeringiana, *Kunth, Enum. Pl.* v. 402.—Japon.
goyazensis, *Griseb. in Mart. Fl. Bras.* iii. i. 41.—Bras.
gracilis, *Hook. & Arn. Bot. Beech. Voy.* 48.—Chili.
gracillima, *Msg. Ann. Mus. Bot. Lugd. Bat.* iii. 160.—Japon.
grandiflora, Mart. & Gal. in Bull. Acad. Brux. ix. 11. (1842) 392 = Galeottiana.
grandiflora, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. i. 28.—Am. trop.
grandifolia, *Schlecht. in Linnaea*, xvii. (1843) 602.—Mexic.
Griesebachii, *Kunth, Enum. Pl.* v. 353.—Bras.
Haenkeana, *Presl, Rel. Haenk.* i. 135.—Peruv.
hastata, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. austr.
hastata, Presl, Rel. Haenk. i. 133 = Presllii.
hastata, *Vell. Fl. Flum.* x. t. 126 (quid ?).—Bras.
hastifolia, *Nees, in Lehm. Pl. Preiss.* ii. 88.—Austral.
hebantha, Mart. ex Griseb. in Mart. Fl. Bras. iii. i. 39 = dodecaneura.
hederacea, Miers, Trav. ii. 281 = bryoniaefolia.
hederifolia, *Griseb. in Mart. Fl. Bras.* iii. i. 42.—Bras.
helicifolia, *Kunth, Enum. Pl.* v. 348.—Chili.
heptaneura, Vell. Fl. Flum. x. t. 124 = lutea.
heterophylla, Poepp. Fragm. Syn. 11 = arenaria.
heterophylla, Roxb. Hort. Beng. 72 ; Fl. Ind. iii. 804 = sativa.
heteropoda, *Baker, in Journ. Bot.* xx. (1882) 270.—Madag.
hexagona, *Baker, l. c.*—Madag
hexaphylla, *Rafin. New Fl. Am.* ii. 89.—Am. bor.
hirsuta, *Blume, Enum. Pl. Jav.* i. 21.—Ind. or. ; Malaya.
hirsuta, Buch.-Ham. ex Wall. Cat. n. 5103 = spinosa.
hirsuta, *Mart. & Gal. in Bull. Acad. Brux.* ix. 11. (1842) 391.—Mexic.
hirtiflora, *Benth. in Hook. Niger Fl.* 537.—Afr. trop.
hispida, Dennst. Schluess. Hort. Malab. 33 = cylindrica.
humifusa, *Poepp. Fragm. Syn.* 12.—Chili.
humilis, *Bert. ex Colla*, in *Mem. Acc. Tor.* xxxix. (1836) 12. t. 51. f. 2.—Chili.
Hurchucia, Buch.-Ham. ex Wall. Cat. n. 5107 = alata.
intermedia, *Thw. Enum. Pl. Zeyl.* 326.—Zeylan.
japonica, *Thunb. Fl. Jap.* 151.—Japon.
japonica, Herb. Madr. ex Wall. Cat. n. 5107 = alata.
japonica, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1341 = divaricata.
kamoonensis, *Kunth, Enum. Pl.* v. 395.—Reg. Himal.
Kegeliana, *Griseb. in Linnaea*, xxi. (1848) 279.—Guiana.
Kirkiniana, Kunth, Enum. Pl. v. 394 = pentaphylla.
Lacerdaei, *Griseb. in Mart. Fl. Bras.* iii. i. 31.—Bras.
lanata, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 96 = spinosa.
latifolia, *Benth. in Hook. Niger Fl.* 535.—Afr. trop.
laurifolia, *Wall. Cat.* n. 5111.—Ins. Penang.
laxiflora, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. i. 32.—Bras.
laxiflora, Schlecht. in Linnaea, xvii. (1843) 606 = remotiflora.
Leibold*iana*, Kunth, Enum. Pl. v. 355.—Mexic.
leptostachya, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 584.—Bras.
leonensis, *G. Don, in Sweet, Hort. Brit.* ed. III. 690.—Afr. trop.
linearis, *Colla, in Mem. Acc. Tor.* xxxix. (1836) 11. t. 51. f. 1.—Chili.
linearis, *Griseb. Cat. Pl. Cub.* 251.—Cuba.
littoralis, *Phil. in Linnaea*, xxix. (1857–58) 64.—Chili.
longifolia, *Phil. l. c.* xxxiii. (1864–65) 254.—Chili.
longifolia, *Rafin. New Fl. Am.* ii. 89.—Am. bor.

DIOSCOREA :—
lucida, R. Br. Prod. 295 (= Roxburghiae sp.).—Austral.
lunata, Roth, Nov. Pl. Sp. 370 = pentaphylla.
Lnschnthiana, *Kunth, Enum. Pl.* v. 364.—Am. trop.
lutea, *G. F. W. Mey. Prim. Fl. Esseq.* 262.—Guian.
lzsoensis, *Sckau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 444.—Ins. Philipp.
macrocarpa, Mart. & Gal. n. 5100 = dumetorum.
macrophylla, Mart. & Gal. in Bull. Acad. Brux. ix. ii. (1842) 396 = macrostachya.
macrostachya, *Benth. Pl. Hartw.* 78.—Mexic.
macrostachya, Mart. & Gal. in Bull. Acad. Brux. ix. ii (1842) 399 = capillaris
marginata, *Griseb. in Mart. Fl. Bras.* iii. i. 37.—Bras.
Martiana, *Griseb. l. c.* 44.—Bras.
martinicensis, Spreng. Neue Entdeck. iii. 17 = *altissima*.
mayenensis, *Kunth, Enum. Pl.* v. 357.—Bras.
megalantha, *Griseb. in Goett. Abh.* xxiv. (1879) 228.—Reg. Argent.
megalobotrya, *Kunth & R. Schomb. ex Kunth, Enum. Pl.* v. 365.—Guiana.
megapoda, *Rafin. New Fl. Am.* ii. 88.—Am. bor.
meridensis, *Kunth, Enum. Pl.* v. 334.—Am. austr.
micrantha, *Kunth, l. c.* 327.—Bras.
microbotrya, *Griseb. in Goett. Abh.* xxiv. (1879) 222.—Reg. Argent.
microphylla, *Steud. Nom.* ed. II. i. 511.—Chili.
modesta, *Phil. in Linnaea*, xxxiii. (1864–65) 256.—Chili.
mollis, *Kunth, Enum. Pl.* v. 369.—Bras.
mollissima, *Blume, Enum. Pl. Jav.* i. 21.—Malaya.
monadelpha, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 164.—Bras.
montana, Spreng. Syst. iv. Cur. Post. 143 = Testudinaria syrvatica.
multiflora, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. i. 35.—Bras.
multiflora, *Presl, Bot. Bemerk.* 118.—Ins. Martinic.
multiloba, *Kunth, Enum. Pl.* v. 376.—Afr. austr.
multinervis, *Benth. Pl. Hartw.* 52.—Mexic.
myriantha, *Kunth, Enum. Pl.* v. 332.—Ins. Philipp.
nana, *Poepp. Fragm. Syn.* 12.—Chili.
nana, Schlecht. in Linnaea, xviii. (1844) 113 = multinervis.
narcotica, Steud. Nom. ed. II. i. 511 = daemona.
nepalensis, *Sweet, Hort. Brit.* ed. II. 522, nomen.—Reg. Himal.
novemloba, *Steud. ex Phil. f. Cat. Pl. Vasc. Chil.* 286, nomen.—Chili.
nummularia, *Lam. Encyc.* iii. 281.—Ins. Molucc.
obtusifolia, *Hook. & Arn. Bot. Beech. Voy.* 48.—Chili.
Olfersiana, *Klotzsch, ex Griseb. in Mart. Fl. Bras.* iii. i. 38.—Bras.
oligophylla, *Phil. in Linnaea*, xxxiii. (1864–65) 255.—Chili.
opposita, Thunb. Fl. Jap. 151 = oppositifolia.
oppositifolia, *Griseb. in Mart. Fl. Bras.* iii. i. 46.—Bras.
oppositifolia, *Linn. Sp. Pl.* 1033.—Ind. or.
ovata, *Vell. Fl. Flum.* x. t. 117.—Bras.
Ovirala, *Baker, in Journ. Bot.* xx. (1882) 269.—Madag.
pachycarpa, *Kunth, Enum. Pl.* v. 370.—Bras.
pallens, *Schlecht. in Linnaea*, xvii. (1843) 610.—Mexic.
palmata, *Yuss. ex Perr. Syn.* ii. 621.—Am. austr.
paniculata, Michx. Fl. Am. Bor. ii. 239 = villosa.
papillata, *Blanco, Fl. Filip.* ed. I. 801.—Ins. Philipp.
parviflora, *Phil. in Linnaea*, xxxiii. (1864–65) 257.—Chili.
pedicellata, *Phil. in Anal. Univ. Chil.* (1873) 540.—Chili.
peltata, *Perr. Syn.* ii. 621.—Zeylan.
pendula, *Poepp. ex Kunth, Enum. Pl.* v. 367.—Bras.
pentaphylla, *Linn. Sp. Pl.* 1032.—Ax. trop.
pentaphylla, Wall. Cat. n. 5092 E = kamoonensis.
perforata, *Bert. ex Spreng. Syst.* ii. 152.—Ins. Guadalup.
pilosiuscula, *Bert. l. c.*—Ins. S. Doming.
piperifolia, Hort. ex Griseb. in Mart. Fl. Bras. iii. i. 42 = polygonoides.
piperifolia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 795.—Am. austr.

DIOSCOREA :—
piperifolia, Klotzsch, ex Kunth, Enum. Pl. v. 354 = caracasana.
Pirtis, *Nadeaud, Enum. Pl. Tahiti*, 35.—Ins. Tahiti.
Poeppigii, *Kunth, Enum. Pl.* v. 365.—Bras.
Pohli, *Griseb. in Mart. Fl. Bras.* iii. i. 35.—Bras.
polygonoides, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 793.—Am. trop.
polygonoides, Mart. & Gal. in Bull. Acad. Brux. ix. ii. (1842) 398 = pallens.
polystachya, *Turcz. in Bull. Soc. Nat. Mosc.* (1887) n. VII. 158.—China.
prsebensilis, *Benth. in Hook. Niger Fl.* 536.—Afr. trop.
Preslii, *Steud. Nom.* ed. II. i. 511.—Mexic.
propinqua, *Hemsl. Biol. Centr. Am. Bot.* iii. 359.—Mexic.
pruinosa, *Kunth, Enum. Pl.* v. 338.—Am. bor.
pubera, *Blume, Enum. Pl. Jav.* i. 21.—Ins. Molucc.
pubescens, *Poir. Encyc. Suppl.* iii. 127.—Guian.
pulchella, Roxb. Hort. Beng. 73; Fl. Ind. iii. 801 = sativa.
Pumilio, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 162.—Bras.
punctata, R. Br. Prod. 294 = transversa.
purpurea, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 799.—Ind. or.
pusilla, Hook. Ic. Pl. t. 678 = humilis.
pyrenaica, *Bub. & Bordrre, ex Gren. in Bull. Soc. Bot. Fr.* xiii. (1866) 382.—Mont. Pyren.
pyrifolia, *Kunth, Enum. Pl.* v. 354.—Burma; Malacca.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* ii. 316.—Abyssin.
quaternata, *Y. F. Gmel. Syst.* 581.—Am. bor.
quinata, J. F. Gmel. l. c. = villosa.
quinquefolia, *Thunb. Fl. Jap.* 150.—Japon.
quinquelobata, Vell. Fl. Flum. x. t. 126 = articulata.
remotiflora, *Kunth, Enum. Pl.* v. 409.—Mexic.
repanda, *Blume, Enum. Pl. Jav.* i. 22.—Java.
repanda, *Rafin. New Fl. Am.* ii. 89.—Am. bor.
reticulata, *C. Gay, Fl. Chil.* vi. 61.—Chili.
retusa, *Mast. in Gard. Chron.* (1870) 1149.—Afr. austr.
riparia, *Kunth & R. Schomb. ex Kunth, Enum. Pl.* v. 364.—Guiana.
rotundata, Poir. Encyc. Suppl. iii. 139 = cayenensis.
rubella, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 798.—Ind. or.
rubiginosa, *Benth. in Hook. Niger Fl.* 538.—Afr. trop.
rubicaulis, *Kunth, Enum. Pl.* v. 368.—Peruv.
Ruiziana, *Klotzsch, ex Kunth, l. c.* 374.—Peruv.
runcinodes, *Griseb. in Mart. Fl. Bras.* iii. i. 42.—Bras.
rupicola, *Kunth, Enum. Pl.* v. 378.—Afr. austr.
sagittata, *Poir. Encyc. Suppl.* iii. 139.—Guiana.
salicifolia, *Blume, Enum. Pl. Jav.* i. 22.—Java.
samydea, *Griseb. in Mart. Fl. Bras.* iii. i. 33.—Bras.
sapindicoides, *Presl, Rel. Haenk.* i. 153.—Mexic.
sativa, Blanco, Fl. Filip. ed. II. 521 = fasciculata.
sativa, Bowdich, Excurs. Mad. 115 = Tamus edulis.
sativa, Gaertn. Fruct. i. 66 = pruinosa.
sativa, Hort. Par. ex Lam. Encyc. iii. 281 = villosa.
sativa, *Linn. Sp. Pl.* 1033.—Reg. trop.
sativa, Linn. l. c. partim = spinosa.
sativa, Rodsch. Obs. 57 = lutea.
sativa, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 275 = Aneders spicata.
sativa, Willd. ex Kunth, Enum. Pl. v. 336 = quinata.
saxatilis, *Poepp. Fragm. Syn.* 11.—Chili.
scabra, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 794.—Venezuela.
scandens, Kunze, ex Kunth, Enum. Pl. v. 350 = brachybotrya.
Schimperiana, *Hochst. ex Kunth, l. c.* 339.—Abyss.
Schlechtendalii, Kunth, l. c. 411 = multiseta.
Schomburgkiana, *Kunth, ex Schomb. Faun. et Fl. Gui.* 899, nomen.—Guiana.
scorpioides, *Wright, in Sauv. Fl. Cub.* 170.—Cuba.
septemloba, Hort. ex Griseb. in Mart. Fl. Bras. iii. i. 46 = sinuata.
septemloba, *Thunb. Fl. Jap.* 149.—Japon.
septemnervis, *Vell. Fl. Flum.* x. t. 116.—Bras.
Sieberi, Kunth, in Abh. Acad. Berl. (1842) 86 = multiflora, *Presl*.
silvestris, *Vell. Fl. Flum.* x. t. 118.—Bras.
sinuata, *Vell. l. c.* t. 120.—Bras.
sororia, *Kunth, Enum. Pl.* v. 358.—Bras.

DIOSCOREA :—
sparsiflora, *Hemsl. Biol. Centr. Am. Bot.* iii. 360.—Mexic.
spicata, *Roth, Nov. Pl. Sp.* 371.—Ind. or.
spiculata, *Blume, Enum. Pl. Jav.* i. 22.—Malaya.
spiculiflora, *Hemsl. Biol. Centr. Am. Bot.* iii. 361.—Mexic.
spinosa, *Roxb. ex Wall. Cat.* n. 5103.—Ind. or.
spinosa, Wall. l. c. 5103 G, H = anguina.
stenocolpos, *Phil. in Linnaea*, xxxiii. (1864–65) 258.—Chili.
subhastata, Vell. Fl. Flum. x. t. 121 = piperifolia.
Swinhoei, Rolfe, in Journ. Bot. xx. (1882) 359 = dioryphora.
syringaefolia, *Kunth & R. Schomb. in Schomb. Faun et Fl. Gui.* 1119, nomen.—Guiana.
tamifolia, Salisb. Parad. Lond. t. 17 = bulbifera.
tamoides, *Griseb. in Mart. Fl. Bras.* iii. i. 43.—Cuba tenella, *Phil. Fl. Atac.* 51.—Chili.
tenuiflora, *Schlecht. in Linnaea*, xvii. (1843) 608.—Mexic.
tennipes, *Franch. & Sav. Enum. Pl. Jap.* ii. 593.—Japon.
teretiuscula, Hort. ex Griseb. in Mart. Fl. Bras. iii. i. 48 = Kajania brasiliensis.
ternata, *Griseb. in Kjoeb. Vidensk. Meddel.* (1876) 158.—Bras.
tiliaefolia, Kunth, Enum. Pl. v. 401 = spinosa.
tomentosa, *Hohen. ex Kunth, l. c.* 391.—Ind. or.
tomentosa, *Koen. ex Roxb. Hort. Beng.* [105]; *Fl. Ind.* iii. 805.—Ind. or.
tomentosa, *Spreng. Pugill.* ii. 92.—Ind. or.
tomcaria, *Boj. Hort. Maurit.* 252.—Afr. trop.
trachyandra, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 155.—Bras.
trachycarpa, *H. B. & K. Nov. Gen. et Sp.* i. 274.—Guiana.
transversa, *R. Br. Prod.* 295.—Austral.
triandra, Hort. ex Griseb. in Mart. Fl. Bras. iii. i. 48 = sativa.
trichantha, *Baker, in Journ. Linn. Soc.* xx. (1883) 271 —Madag.
trifida, *Linn. f. Suppl.* 427.—Guiana.
trifoliata, *H. B. & K. Nov. Gen. et Sp.* i. 275.—Venezuela.
trifoliata, Mor. ex Kunth, Enum. Pl. v. 417 = galapagensis.
trifoliata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 13 = hirsuta, *Blume*.
trilingnis, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 163.—Bras.
triloba, Karst. ex Kunth, Enum. Pl. v. 418 = trifoliata.
triloba, *Lam. Encyc.* iii. 284.—Guian.
triloba, Willd. Sp. Pl. iv. 791 = trifida.
trinervia, Roxb. ex Wall. Cat. n. 5099 = daemona.
triphylla, Blanco, Fl. Filip. ed. I. 799 = hirsuta.
triphylla, Buch.-Ham. ex Wall. Cat. n. 5099 = daemona.
triphylla, Linn. Sp. Pl. 1032 = pentaphylla.
triphylla, Herb. Russ. ex Wall. Cat. n. 5001 = tomentosa, *Korn*.
triphylla, *Schimp. ex Kunth, Enum. Pl.* v. 436.—Abyss.
tuberosa, Vell. Fl. Flum. x. t. 125 = Kajania brasiliensis.
tubulosa, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 154.—Bras.
Tugui, Blanco, Fl. Filip. ed. I. 800 = fasciculata.
Tunga, Buch.-Ham. ex Wall. Cat. n. 5107 = sativa.
undecimnervis, Vell. Fl. Flum. x. t. 120 = glandulosa.
urophylla, *Hemsl. Biol. Centr. Am. Bot.* iii. 361.—Panama.
varifolia, *Bert. in Mercc. Chil.* n. xiii. p. 613; *ex Colla, in Mem. Acc. Tor.* xxxix. (1836) 13.—Chili.
varifolia, *Kunze, in Linnaea*, xx. (1847) 12.—Bras.
veraicolor, Buch.-Ham. ex Wall. Cat. n. 5106 = sativa.
verticillata, *Lam. Encyc.* iii. 281.—Java.
Vespertilio, *Benth. in Hook. Niger Fl.* 538.—Afr. trop.
villa, *Kunth, Enum. Pl.* v. 400.—Java.
villosa, *Linn. Sp. Pl.* 1033.—Am. bor.
villosa, Willd. ex Kunth, Enum. Pl. v. 418 = trifoliata.
virosa, Wall. Cat. n. 5099 = daemona.
vitata, *Bull, ex Baker, in Bot. Mag.* t. 6408.—Am. trop.
Volckmanni, *Phil. in Linnaea*, xxxiii. (1864–65) 255.—Chili.
vulgaris, Miq. Fl. Ind. Bat. iii. 879 = alata.
Wahlbergi, Decf. Tabl. ed. II. 269 = villosa.
Zollingeriana, *Kunth, Enum. Pl.* v. 334.—Java.

DIOSMA, Linn. Syst. ed. I (1785). *RUTACEAE*, Benth. & Hook. f. i. 289.

abietina, D. Dietr. Syn. Pl. i. 829 = Euchaetis glomerata.

acicularis, Sallsb. Prod. 142 = oppositifolia.

acmaephylla, Eckl. & Zeyh. Enum. 849 = aspalathoides.

acuminata, DC. Prod. i. 715 = Agathosma imbricata.

acuminata, Lodd. Bot. Cab. t. 493 = Adenandra uniflora.

acuta, Lee, ex Bartl. & Wendl. f. Diosm. 182 = Agathosma cuspidata.

alba, Thunb. Prod. Pl. Cap. 84 = Coleonema album.

alpina, D. Dietr. Syn. Pl. i. 823 = Barosma alpina.

ambigua, Bartl. & Wendl. f. Diosm. 42 = oppositifolia.

ambigua, Eckl. & Zeyh. Enum. 108 = aspalathoides.

ambigua, Drège, ex Presl, Bot. Bemerk. 31 = macrosticta.

ambigua, Lodd. Bot. Cab. t. 461 = Agathosma ambigua.

amoena, *Lodd. l. c.* t. 161.—Afr. austr.

angustifolia, Hort. ex Hook. Bot. Mag. sub t. 3340 = Coleonema pulchrum.

apiculata, Spreng. Syst. i. 787 = Agathosma pilifera.

asiatica, Lour. Fl. Cochinch. i. 161 (= Pseudiosma ?). —Cochinch.

aspalathoides, *Lam. Encyc.* ii. 286.—Afr. austr.

barbata, Berg. ex Harv. & Sond. Fl. Cap. i. 480 = Macrostylis villosa.

barbata, DC. Prod. i. 715 = Macrostylis lanceolata.

barbigera, Linn. f. Suppl. 155 = Macrostylis barbigera.

barbigera, Thunb. Prod. Pl. Cap. 48 = Macrostylis lanceolata.

betulina, Thunb. l. c. = Barosma betulina.

bifida, Jacq. Coll. iii. 278. t. 20. f. 1 = Agathosma bifida.

bifurca, Willd. Sp. Pl. i. 1186 = Agathosma bisulca.

biseriata, Spreng. Syst. i. 784 = Adenandra biseriata.

bisulca, Thunb. Prod. Pl. Cap. 84 = Agathosma bisulca.

brachyphylla, D. Dietr. Syn. Pl. i. 828 = Adenandra brachyphylla.

brevifolia, Lam. Encyc. ii. 285 = Agathosma erecta.

brunioides, Link, Enum. Hort. Berol. i. 287 = Agathosma cuspidata.

Bucco, Hort. Berol. ex Regel, Ind. Sem. Hort. Petrop. (1865) 51 = Adenandra umbellata ?

calycina, Steud. in Flora, xiii. (1830) 549 = Coleonema pulchrum.

capensis, Linn. Mant. ii. 243 = Agathosma hispida.

capitata, Linn. f. c. 210; Thunb. Prod. Pl. Cap. 43 = Audouinia capitata.

cerefolia, Vent. Jard. Malm. 93. t. 93 = Agathosma Cerefolium.

ciliata, Lam. Encyc. ii. 287 = Agathosma pubescens.

ciliata, Linn. Sp. Pl. 198 = Agathosma ciliata.

ciliata, Thunb. Prod. Pl. Cap. 48 = Agathosma platypetala.

cistoides, Lam. Encyc. ii. 289 = Adenandra uniflora.

cordata, Mart. Enum. Hort. Erlang. 67 = Agathosma imbricata.

cornicea, DC. Prod. i. 713 = Adenandra coriacea.

corymbosa, Montin, in Phys. Sällsk. Handl. i. (1776) 106 = Agathosma villosa.

crassifolia, Hoffmgg. Vers. Pfl. 57 = succulenta ?

crenata, Linn. Syst. Nat. ed. X. ii. 940 = Barosma crenulata.

crenata, Lodd. Bot. Cab. v. t. 404 = Barosma betulina.

crenulata, Linn. Cent. Pl. ii. 11 = Barosma crenulata.

cupressina, *Linn. Mant.* i. 50.—Afr. austr.

cuspidata, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 94 = Linconia cuspidata.

decussata, Lam. Encyc. ii. 288 = succulenta.

depressa, Eckl. & Zeyh. Enum. 108 = aspalathoides.

deusta, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 95 = Linconia thymifolia.

dichotoma, Berg. Desc. Pl. Cap. 48 = cupressina.

dielsa, Ker-Gawl. in Bot. Reg. t. 502 = Barosma lanceolata.

dubia, Spreng. ex Steud. in Flora, xiii. (1830) 548 = Agathosma guidiolides.

Ecklonianus, *Sond. in Harv. & Sond. Fl. Cap.* i. 375.—Afr. austr.

elata, D. Dietr. Syn. Pl. i. 830 = Euchaetis elata.

ensata, Thunb. Prod. Pl. Cap. 48 = Empleurum serrulatum.

erecta, DC. Prod. i. 715 = Agathosma erecta.

DIOSMA :—

erianthe, Steud. in Flora, xiii. (1830) 550 = Agathosma Schlechtendalii.

ericaefolia, Andr. Bot. Rep. t. 451 = oppositifolia.

ericoides, Lam. Encyc. ii. 285 = aspalathoides.

ericoides, *Linn. Sp. Pl.* 198.—Afr. austr.

ericoides, Sims, Bot. Mag. t. 2382 = oppositifolia.

fervulacea, Hort. Kew. ex Harv. & Sond. Fl. Cap. i. 376 = teretifolia.

floridissima, Spreng. Syst. i. 785 = Barosma foetidissima.

flavescens, *Willd. in Hook. Ic. Pl.* t. 1476.—Afr austr.

flexilis, D. Dietr. Syn. Pl. i. 829 = Euchaetis flexilis.

fragrans, Sims, Bot. Mag. t. 1519 = Adenandra fragrans, *Roem. & Schult.*

glabrata, G. F. W. Mey. ex Bartl. & Wendl. f. Diosm. 32 = oppositifolia.

glandulosa, Hort. ex Bartl. & Wendl. f. l. c. 110 = Barosma ovata.

glandulosa, Thunb. Diss. Diosm. 16 = Agathosma glandulosa.

glomerata, E. Mey. ex Bartl. & Wendl. f. Diosm. 16 = Euchaetis glomerata.

gnidioides, D. Dietr. Syn. Pl. i. 829 = Agathosma gnidioides.

grandiflora, *Burm. f. Fl. Cap. Prod.* 6.—Afr. austr.

graveolens, Licht. ex Roem & Schult. Syst. v. 461 = Barosma pulchella.

hirsuta, Linn. Sp. Pl. 198 = oppositifolia.

hirta, Lam. Encyc. ii. 286 = Agathosma hirta.

hirta, Vent. Jard. Malm. t. 72 = Agathosma Ventenatiana.

hispida, Hort. Angl. ex Bartl. & Wendl. f. Diosm. 159 = Agathosma Cerefolium.

hispida, Hort. ex Bartl. & Wendl. f. Diosm. 168 = Agathosma cuspida.

hispida, Thunb. Prod. Pl. Cap. 42 = Agathosma hispida.

humilis, D. Dietr. Syn. Pl. i. 828 = Adenandra humilis.

hybrida, Hort. Angl. ex Bartl. & Wendl. f. Diosm. 107 = Agathosma pubescens.

hybrida, *Spreng Syst.* i. 787.—Afr austr.

imbricata, Thunb. Prod. Pl. Cap. 43 = Agathosma imbricata.

intermedia, D. Dietr. Syn. Pl. i. 828 = Adenandra marginata.

juniperifolia, Salisb. Prod. 142 = oppositifolia.

juniperina, Moench, Meth. Suppl. 26 = Coleonema album ?

juniperina, Spreng. Syst i. 784 = Coleonema juniperifolium.

larvigata, Spreng. l. c. = Acmadenia laevigata.

lanceolata, Linn. Sp. Pl. 198; Thunb Prod. Pl. Cap. 43 = Barosma lanceolata.

lanceolata, Spreng. Syst. i. 788 = Macrostylis lanceolata.

lancifolia, *F. W. Schmidt, Neue u. Selt. Pfl.* 37.—Hab. ?

laxiophylla, Spreng. Syst. i. 787 = Agathosma villosa.

latifolia, Linn. f. Suppl. 154 = Barosma latifolia.

latifolia, Lodd. Bot. Cab. iii. t. 290 = Barosma crenulata.

laxa, Spreng. Syst. i. 787 = Agathosma villosa.

leiophylla, Spreng. l. c. = Agathosma glabrata.

Lichtensteiniana, D. Dietr. Syn. Pl. i. 828 = Adenandra cuspidata.

linearis, Eckl. & Zeyh. Enum. 108 = aspalathoides.

linearis, Thunb. Prod. Pl. Cap. 43 = Adenandra uniflora.

linifolia, DC. Prod. i. 715 = Agathosma linifolia.

linifolia, Hort. ex Bartl. & Wendl. f. Diosm. 44 = longifolia.

linifolia, Lodd. Bot. Cab. t. 400 = Barosma lanceolata.

longifolia, Eckl. & Zeyh. Enum. 108 = oppositifolia.

longifolia, Wendl. Coll. i. 21. t. 19 = oppositifolia.

lycopodioides, Willd. ex Roem. & Schult. Syst. v. 461 = Agathosma lycopodioides.

macrosticta, *Presl, Bot. Bemerk.* 31.—Afr. austr.

marginata, Linn. f. Suppl. 155 = Adenandra marginata.

Meyeri, D. Dietr. Syn. Pl. i. 828 = Adenandra cuspidata.

Meyeriana, Spreng. Syst. i. 788 = virgata.

Meyeriana, Steud. ex Harv. & Sond. Fl. Cap. i. 378 = Coleonema juniperifolium.

microphylla, Spreng. Syst. i. 787 = Barosma scoparia.

mixta, Hort. Kew. ex Bartl. & Wendl. f. Diosm. 182 = Agathosma cuspidata.

mundtiaefolia, D. Dietr. Syn. Pl. i. 826 = Adenandra mundtiaefolia.

DIOSMA :—

Myrsinites, Lam. Illustr. ii. 82 = Agathosma ciliata.

myrtifolia, Hort. Angl. ex Bartl. & Wendl. f. Diosm. 156 = Agathosma ciliata.

oblonga, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 23 = Barosma oblonga.

obtusa, DC. Prod. i. 714 = Agathosma obtusa.

obtusa, G. F. W. Mey. ex Bartl. & Wendl f. Diosm. 197 = Macrostylis squarrosa.

obtusata, Thunb. Prod. Pl. Cap. 84 = Acmadenia juniperina.

obtusata, Willd. Sp. Pl. i. 1184 = Acmadenia obtusata obtusifolia, *Sond. in Harv. & Sond. Fl. Cap.* i. 377.—Afr. austr.

odorata, DC. Prod. i. 714 = Barosma crenulata.

odoratissima, Montin, in Phys. Sällsk. Handl. i. (1776) 104 = Barosma latifolia.

oppositifolia, Drège, ex Presl, Bot. Bemerk. 31 = Coleonema pulchrum.

oppositifolia, *Linn. Sp. Pl.* 198.—Afr. austr.

oppositifolia, Roem. & Schult. Syst. v. 457 = succulenta.

orbicularis, Hort. ex Bartl. & Wendl. f. Diosm. 110 = Barosma ovata.

orbicularis, Thunb. Prod. Pl. Cap. 84 = Agathosma orbicularis.

ovata, Lodd. ex Bartl. & Wendl. f. Diosm. 80 = Adenandra amoena.

ovata, Thunb. Prod. Pl Cap. 43 = Barosma ovata.

pallida, Sallsb. Prod. 142 = oppositifolia.

parviflora, Willd. ex Roem. & Schult. Syst. v. 462 = Agathosma parviflora.

parvifolia, Harv. & Sond. Fl. Cap. i. 533, Index = Agathosma parviflora.

passerinoides, *Steud. in Flora*, xiii. (1830) 549.—Afr. austr.

patula, Spreng. Syst. i. 786 = Agathosma patula.

pauciflora, D. Dietr. Syn. Pl. i. 829 = Barosma pauciflora.

pectinata, Thunb. Prod. Pl. Cap. 84 = oppositifolia.

perforata, Lam. Illustr. ii. 82 = Agathosma perforata.

pilifera, Steud. in Flora, xiii. (1830) 549 = Agathosma pilifera.

pinifolia, {Finch.} Hort. Gorenk. ed. I. (1808) 94 = oppositifolia.

prolifera, DC. Prod. i. 715; Spreng. Syst. i. 786 = Agathosma prolifera.

puberula, Steud. in Flora, xiii. (1830) 548 = Agathosma guidiolides.

pubescens, Thunb. ex Harv. & Sond. Fl. Cap. i. 482 = Agathosma ericoides.

pubescens, Thunb. Prod. Pl. Cap. 43 = Agathosma villosa.

pulchella, Houtt. Handleid. iii. 326 = Barosma ovata.

pulchella, Linn. Syst. ed. X. 940 = Barosma pulchella.

punctata, Licht. in Roem. & Schult. Syst. v. 461 = Barosma ovata.

pungens, Spreng. Syst. i. 784 = Acmadenia pungens.

purpurea, Desf. Tabl. ed. II. 180 = Agathosma Ventenatiana.

purpurea, Hort. ex Bartl. & Wendl. f. Diosm. 163 = Agathosma villosa.

ramosissima, *Bartl. & Wendl. f. l. c.* 48.—Afr. austr.

recurva, Cham in Linnaea, v. (1830) 51 = ericoides.

reflexa, Lodd. ex DC. Prod. i. 715 = Agathosma reflexa.

reflexa, Soland. ex Harv. & Sond. Fl. Cap. i. 419 = Agathosma imbricata.

rigidula, Willd. Enum. Hort. Berol. Suppl. 12 = succulenta.

rosmarinifolia, Lam. Encyc. ii. 286 = Adenandra marginata.

rotundifolia, D. Dietr. Syn. Pl. i. 828 = Adenandra rotundifolia.

rubra, Berg. Desc. Pl. Cap. 62 = Coleonema album.

rubra, Hort. ex Bartl. & Wendl. f. Diosm. 168 = Agathosma villosa.

rubra, Linn. Sp. Pl. 189 = oppositifolia.

rufescens, Spreng. Syst. i. 787 = Agathosma rubra.

rugosa, Donn, ex Bartl. & Wendl. f. Diosm. 81 = Adenandra umbellata.

rugosa, Hort. Angl. ex Bartl. & Wendl. f. l. c. 162 = Agathosma villosa.

rugosa, Thunb. Prod. Pl. Cap. 43 = Agathosma pubescens.

rugosa, Willd. ex Harv. & Sond. Fl. Cap. i. 425 = Agathosma ciliata.

scabra, Lam. Encyc. ii. 283 = succulenta.

scoparia, D. Dietr. Syn. Pl. i. 829 = Barosma scoparia.

DIOSMA :—
serpyllacea, DC. Prod. i. 715 = Agathosma serpyllacea.
serpyllifolia, Sallab. Prod. 142 = ericoides.
serrata, Blanco, Fl. Filip. ed. I. 168 = Celastrus paniculata.
serratifolia, Curt. Bot. Mag. t. 456 = Barosma serratifolia.
Simsiana, Eckl. & Zeyh. Enum. 108 = oppositifolia.
spartifolia, Stend. in Flora, xiii. (1830) 550 = Barosma scoparia.
speciosa, Sims, Bot. Mag. t. 1271 = Adenandra umbellata.
spicata, DC. Prod. i. 715 = Agathosma spicata.
squamosa, Willd. in Roem. & Schult. Syst. v. 462 = Agathosma squamosa.
squarrosa, Wendl. ex Bartl. & Wendl. f. Diosm. 198 = Macrostylis squarrosa.
stenopetala, Steud. in Flora, xiii. (1830) 549 = Agathosma filipetala.
stenophylla, Spreng. Syst. i. 795 = Barosma angustifolia.
stricta, Willd. ex Roem. & Schult. Syst. v. 462 = Agathosma serpyllacea.
subulata, Wendl. Coll. i. 31. t. 8 = oppositifolia.
succulenta, *Berg. Desc. Pl. Cap.* 68.—Afr. austr.
tenuifolia, Presl, Bot. Bemerk. 31 = Coleonema pulchrum.
tenuifolia, Willd. Enum. Hort. Berol. 258 = oppositifolia.
tenuissima, Willd. Enum. Hort. Berol. Suppl. 12 = Agathosma erecta.
teretifolia, *Hort. Angl. ex Link, Enum. Hort. Berol.* i. 287.—Afr. austr.
ternata, D. Dietr. Syn. Pl. i. 829 = Barosma ternata.
tetragona, Hort. ex Bartl. & Wendl. f. Diosm. 26 = succulenta.
tetragona, Linn. f. Suppl. 155 = Acmadenia tetragona.
Thunbergiana, Spreng. Syst. i. 786 = Agathosma platypetala.
thymifolia, Willd. ex Roem. & Schult. Syst. v. 462 = Agathosma pubescens.
thyoides, Willd. ex Roem. & Schult. l. c. = Agathosma erecta.
thyrsophora, Eckl. & Zeyh. Enum. 107 = ericoides.
tomentosa, Lee, ex Bartl. & Wendl. f. Diosm. 168 = Agathosma pubescens.
uitenhagensis, D. Dietr. Syn. Pl. i. 827 = oppositifolia †
ulicina, Lodd. Cat. (1824) = Acmadenia juniperina.
umbellata, Hort. ex Bartl. & Wendl. f. Diosm. 81 = Adenandra umbellata.
umbellata, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 24 = Agathosma bifida.
undulata, *Thunb. Fl. Cap.* ii. 132.—Afr. austr.
uniceps-lauri, Linn. f. Suppl. 155 = Empleurum serrulatum.
uniflora, Linn. Sp. Pl. 198 = Adenandra uniflora.
ustulata, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 25.—Afr. austr.
Ventenatiana, Spreng. Syst. i. 786 = Agathosma Ventenatiana.
venusta, D. Dietr. Syn. Pl. i. 829 = Barosma venusta.
vestita, DC. Prod. i. 715; Spreng. Syst. i. 787 = Agathosma imbricata.
villosa, Thunb. Prod. Pl. Cap. 48 = Macrostylis villosa.
virgata, Hort. ex Bartl. & Wendl. f. Diosm. 145 = Agathosma imbricata.
virgata, *Lam. Encyc.* ii. 286.—Afr. austr.
virgata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 378 = Coleonema juniperifolium.
virgata, *G. F. W. Mey. ex Bartl. & Wendl. f. Diosm.* 46.—Afr. austr.
virgata, Thunb. Prod. Pl. Cap. 84 = Agathosma virgata.
viscida, D. Dietr. Syn. Pl. i. 828 = Adenandra viscida.
vulgaris, Schlecht. in Linnæa, vi. (1831) 200 = oppositifolia.
Wendlandiana, DC. Prod. i. 715 = Agathosma villosa.

DIOSPYROS, Linn. Gen. ed. I. 148 1737). *EBENACEÆ*, Benth. & Hook. f. ii. 665.
CARGILLIA, R. Br. Prod. 528 (1810), partim.
CAVANILLEA, Desr. in Lam. Encyc. iii. 668 (1789).
DACTYLUS, Forsk. Fl. Aegypt. Arab. p. xxxvi (1775).
DANELSKIA, Bert. ex A. DC. in DC. Prod. viii. 224 (1844).
EMBRYOPTERIS, Gaertn. Fruct. i. 145. t. 29 (1788).

DIOSPYROS :—
GUNISANTHUS, A. DC. in DC. Prod. viii. 219 (1844).
LEUCOXYLUM, Blume, Bijdr. 1189 (1826).
MABOLA, Rafin. Sylva Tellur. 11 (1838).
NOLTIA, Schum. in Danske Selsk. Afh. iii. (1827) 209 (Beskr. Pl. Guin. 189).
PARALEA, Aubl. Pl. Gui. i. 376. t. 231 (1775).
PATONIA, Wight, Illustr. i. 19 (1838), partim.
PERSIMON, Rafin. Sylva Tellur. 104 (1838).
ROSSIDIUM, A. DC. in DC. Prod. viii. 220 (1844).
acapulcensis, H. B. & K. Nov. Gen. et Sp. iii. 254 = Maba acapulcensis.
acuminata, Wall. Cat. n. 4129 (= *Cylicodaphne myristicæfolia*).—Ind. or.
acuta, *Thw. Enum. Pl. Zeyl.* 182.—Zeylan.
affinis, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 169 = Maba.
affinis, *Thw. Enum. Pl. Zeyl.* 179.—Ind. or.
albens, Presl, Rel. Haenk. ii. 62 = Maba albens.
ambigua, Vent. Jard. Malm. t. 17 = Royena ambigua.
amœna, Wall. Cat. n 4139 = lanceolata.
amplexicaulis, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 11. n. 271. f. 139 = leucomelas.
angulata, Poir. Encyc. v. 434 = nodosa.
angustifolia, Auditb. ex Spach, Hist. Vég. Phan. ix. 405 = virginiana.
anonæfolia, *A. DC. in DC. Prod.* viii. 227.—Ins. Mauritius.
apeibacarpos, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 891.—Bras.
apiculata, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 166.—Ins. Penang.
argentea, *Griff. Notul.* iv. 288.—Malacca.
Arnottiana, Miq. ex Thw. Enum. Pl. Zeyl. 423 = Candolleana.
artanthæfolia, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 7.—Peruv.
asiatica, *Dodd. Rep. Forest. Madr.* (1866–67) 20. t. 1.—Ind. or.
asteroculyx, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 198.—Borneo.
attenuata, *Thw. Enum. Pl. Zeyl.* 182.—Zeylan.
aurea, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 408.—Java.
auriculata, Wight, ex Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 188 = foliolosa.
australis, Hiern, l. c. 54, 246 = D. Cargillia.
Bartari, *Hiern, l. c.* 187.—Afr. trop.
batocana, *Hiern, l. c.* 174.—Afr. trop.
Beccarii, *Hiern, l. c.* 204.—Borneo.
Bernieri, *Hiern, l. c.* 268.—Madag.
Bertersi, A. DC. in DC. Prod. viii. 234 = Maba inconstans.
bicolor, Klotzsch, in Peters, Reise Mossamb. Bot. 184 = mespiliformis.
bifloræ, *Blanco, Fl. Filip.* ed. I. 308.—Ins. Philipp.
Blancoi, A. DC. in DC. Prod. viii. 227 = discolor.
Boivini, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 194.—Madag.
borneensis, *Hiern, l. c.* 178.—Borneo.
Boutoniana, A. DC. in DC. Prod. viii. 236 = nodosa.
bracteata, Roxb. Hort. Beng. [98]; Fl. Ind. ii. 539 = montana.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) 71. 72.—Burma.
brasiliensis, Mart. ex Miq. in Mart. Fl. Bras. vii. 5 = D. Ebenaster.
Burchellii, *Hiern, in Journ. Bot.* xii. (1874) 240.—Afr. trop.
burmanica, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 78.—Burma.
buxifolia, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 213 = microphylla.
calophylla, *Hiern, l. c.* 156.—Madag.
calycina, Auditb. ex Spach, Hist. Vég. Phan. ix. 405 = virginiana.
calycina, Bedd. Fl. Sylv. t. 68 = foliolosa.
calycina, Hort. ex C. Koch, Dendrol. ii. 11. 204 = *Lotus*.
canarica, Bedd. Forest. Man. Bot. 145; Ic. Pl. Ind. Or. 28. t. 127 = Candolleana.
Candolleana, Thw. Enum. Pl. Zeyl. 181 = Thwaitesii.
Candolleana, *Wight, Ic.* t. 1321.—Ind. or.
Canonoi, A. DC. in DC. Prod. viii. 237 = multiflora.
capensis, A. DC. l. c. 226 = nodosa.
capitulata, Wight, Ic. t. 1324 = D. Chloroxylon.
capreæfolia, *Mart. ex Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 254.—Bras.

DIOSPYROS :—
Cargillia, *F. Muell. Austr. Veg. Intercol. Exhib. Ess.* (1866–67) 85.—Austral.
caroliniana, Mahl. ex Rafin. Fl. Ludov. 139 = virginiana.
Carthei, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 198.—Ins. Philipp.
caulifloræ, *Blume, Bijdr.* 668.—Java.
cauliflora, Mart. ex Miq. in Mart. Fl. Bras. vii. 7 = Maba cauliflora.
cayennensis, *A. DC. in DC. Prod.* viii. 224.—Amer. trop.
cerasifolia, D. Don, Prod. Fl. Nep. 144 = Eurya symplocina.
chartaceæ, *Wall. Cat.* n. 4135, partim.—Burma.
chinensis, Blume, Cat. Gew. Buitenz. 110 = D. Kaki
Chloroxylon, *Roxb. Pl. Corom.* i. 38. t. 49.—Ind. or. Chrysophyllos, *Poir. Encyc.* v. 438.—Ins. Maurit.
ciliata, *A. DC. in DC. Prod.* viii. 229.—Mexic.
ciliata, Rafin. New Fl. Am. iii. 85 = virginiana.
citrifolia, Wall. ex A. DC. in DC. Prod. viii. 235 = D Embryopteris.
clavigera, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii 555.—Malacca.
coccolobæfolia, *Mart. ex Miq. in Mart. Fl. Bras.* vii 6.—Bras.
Commersonii, Gaertn. f. Fruct. iii. 186. t. 208 = leucomelas.
comorensis, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 220.—Ins. Comoro.
concolor, Moench, Meth. 470 = virginiana.
conduplicata, H. B. & K. Nov. Gen. et Sp. iii. 254 = Maba inconstans.
cordifolia, Roxb. Pl. Corom. i. 38. t. 50 = montana.
cornacea, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 259.—Borneo.
costata, Carr. in Rev. Hortic. (1870) 134 = D. Kaki.
crassiflora, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 260.—Afr. trop.
crumenata, *Thw. Enum. Pl. Zeyl.* 179.—Zeylan.
Cumingi, *A. DC. in DC. Prod.* viii. 237.—Ins. Philipp.
cuneifolia, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 268.—Malac.
cupulosa, F. Muell. Austr. Veg. Intercol. Exhib. Ess (1866–67) 85 = Maba rufa.
cystopus, *Miq. Fl. Ind. Bat. Suppl.* 584.—Sumatra.
dasyphylla, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 71.—Burma.
decandra, Boj. Hort. Maurit. 200 = D. Ebenum.
decandra, *Lour. Fl. Cochinch.* 577.—Cochinch.
decipiens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 568.—Malacca.
Dendo, *Welw. ex Hiern, in Trans. Camb. Phil. Soc* xii. (1873) 198.—Afr. trop.
densiflora, *Wall. Cat.* n. 4140.—Burma.
dictyonea, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 199.—Borneo.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 588.—Sumatra.
digyna, Hort. ex Loud. Encyc. Trees & Shrubs, 627 = virginiana.
digyna, Jacq. Hort. Schoenb. iii. 35. t. 313 = D. Ebenum.
discica, Span. Ic. n. 23; ex Linnæa, xv. (1841) 336 = montana.
discolor, *Willd. Sp. Pl.* iv. 1106.—Ins. Philipp.
distyla, Hort. ex C. Koch, Dendrol. ii. 11. 206 = virginiana.
diversifolia, *Hiern, in Journ. Bot.* xiii. (1875) 353.—Ins. Mascar.
dodecandra, *Lour. Fl. Cochinch.* 398.—Cochinch.
dubia, Wall. Cat. n. 4407 = melanoxylon.
Ebenaster, Retz. Obs. Bot. fasc. v. 31 = D. Ebenum.
Ebenum, Hiern, in Trans. Camb. Phil. Soc. xii. (1876) 206, partim = assimilis.
Ebenum, *Korn. in Phys. Sälsk. Handl.* i. (1776) 176. —Ind. or.; Malaya.
Ebenum, Poir. Encyc. v. 429 = tessellaria.
edulis, Lodd. ex Sweet, Hort. Brit. ed. I. 370 = D. Ebenum.
ehretioides, *Wall. Cat.* n. 4137.—Burma.
elegans, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 571.—Reg. Himal.
emarginata, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 264.—Bras.
Embryopteris, Blanco, Fl. Filip. ed. II. 309 = discolor.

DIOSPYROS :—
Embryopteris, *Pers. Syn.* ii. 624.—Afr. trop.
ebanifera, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 302.—Hong kong.
exsculpta, Bedd. Fl. Sylv. t. 66 = melanoxylon.
exsculpta, Bech.-Ham. in Trans. Linn. Soc. xv. (1826) 110 = tomentosa.
exsculpta, Dalz. & Gibs. Bomb. Fl. 142 = D. Tupru.
fasciculosa, F. Muell. Austr. Veg. Intercol. Exhib. Eas. (1866–67) 35 = Maba fasciculosa.
feminina, Buch.-Ham. ex A. DC. Prod. viii. 288 = Eurya symplocina.
ferruginea, Spitig. ex De Vriese, in Nederl. Kruidk. Arch. i. (1848) 327 = D. Parviea.
fertilis, Lodd. Cat. ex Loud. Arboret. ii. 1197 = virginiana.
flavicans, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 205.—Burma; Malaya.
formince, Buch.-Ham. ex G. Don, Gen. Syst. iv. 39 = Eurya symplocina.
foliolosa, *Wall. Cat.* n. 4143.—Ind. or.
frondosa, Wall. l. c. n. 4125 = Bocages elliptica.
frutescens, *Blume, Bijdr.* 668.—Java.
frutescens, Hask. Pl. Jav. Rar. 407 = Horsfieldii.
fulgines, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 184.—Borneo.
fusco-velutina, *Baker, in Journ. Linn. Soc.* xxi. (1885) 422.—Madag.
Gardneri, *Thw. Enum. Pl. Zeyl.* 181.—Zeylan.
gaultheriacfolia, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 5.—Bras.
geminata, F. Muell. Austr. Veg. Intercol. Exhib. Eas. (1866–67) 35 = Maba geminata.
glabrifolia, Rottl. in Nye Saml. Vidensk. Selsk. Skr. ii. (1783) 340. t. 5 = D. Ebenum.
glauca, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 221 = montana ?
glomerata, *Spruce, in Journ. Linn. Soc.* v. (1861) 7.—Bras.
glutenifera, Wall. Cat. n. 4123 B = D. Embryopteris.
glutinosa, Koen. ex Roxb. Hort. Beng. 40; Fl. Ind. ii. 538 = D. Embryopteris.
Goindu, Dalz. in Hook. Kew Journ. iv. (1852) 111 = montana.
gonoclada, *Baker, in Journ. Linn. Soc.* xxi. (1885) 424.—Madag.
Goudotii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 240.—N. Granat.
gracilifiora, *Hiern, l. c.* 191.—Borneo.
gracilipes, *Hiern, l. c.*—Madag.
grandifolia, *Wall. ex Voigt, Hort. Suburb. Calc.* 345, nomen.—Hab. ?
grata, *Wall. Cat.* n. 2142.—Reg. Himal.
guaiacana, Robin, Voy. Louis, iii. 417 = virginiana.
halesioides, *Griseb. Cat. Pl. Cub.* 168.—Cuba.
haplostylis, *Boiv. ex Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 177.—Madag.
Hasseltii, *Zoll. in Tijdschr. Nederl. Ind.* xiv. (1857) 159.—Java.
hebecarpa, A. Cunn. ex Benth. Fl. Austral. iv. 286 = D. Ebenum.
Hebrastter, Gaertn. Fruct. ii. 478. t. 179 = lencomelas.
Helferi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 560.—Burma.
heterophylla, Wall. Cat. n. 4138 = montana.
Heudeloti, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 215.—Afr. trop.
hexasperma, Hasselt, ex Hask. Pl. Jav. 465. n. 353 = Maba elliptica.
hirsuta, Desf. in Ann. Mus. Par. vi. (1805) 447 = Royena hirsuta.
hirsuta, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 164 = Candolleana, Moonii, Thwaitesii.
hirsuta, *Linn. f. Suppl.* 440.—Zeylan.
hispida, *A. DC. in DC. Prod.* viii. 236.—Bras.
Horsfieldii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 192.—Malaya.
humilis, F. Muell. Austr. Veg. Intercol. Exhib. Eas. (1866–67) 35 = Maba humilis.
incana, Buch.-Ham. ex Wall. Cat. n. 4122 = D. Toposia.
inconstans, Jacq. Enum. Pl. Carib. 34 = Maba inconstans.
inaculpta, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 119 = tomentosa.
insignis, *Thw. Enum. Pl. Zeyl.* 180.—Zeylan.
intermedia, Hort. ex Loud. Encyc. Trees & Shrubs, 627 = virginiana.
japonica, Sieb. & Zucc. in Abh. Acad. Muench. iv. iii. (1846) 186 = D. Lotus.
Kaempferi, *Naud. in Nouv. Arch. Mus. Par.* Sér. II. iii. (1880) 226.—Japon.
Kaki, Blanco, Fl. Filip. ed. I. 302 = discolor.
Kaki, *Linn. f. Suppl.* 439.—Japon.
Kaki, Thunb. Fl. Jap. 157 = Kaempferi.
Kirkii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 199.—Afr. trop.
Korthalsiana, *Hiern, l. c.* 168.—Borneo.
Kuhlii, *Zoll. in Tijdschr. Nederl. Ind.* xiv. (1857) 159. —Java.
Kurzii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 162.—Ins. Andaman. et Nicobar.
laevis, *Boj. ex A. DC. in DC. Prod.* viii. 232.—Madag
lanceaefolia, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 537. —Ind. or.; Burma.
lanceolata, Poir. Encyc. v. 434 = Maba lanceolata.
lanceolata, Mart. ex Miq. n. 4122 = D. Toposia.
lateralis, *Hiern, in Trans. Camb. Phil. Soc.* xii (1873) 167.—Borneo.
laurifolia, A. Rich. Fl. Cub. Fanerog. ii. 86 = D. Ebenum.
laxa, Teijsm. & Binn. in Nederl. Kruidk. Arch. iii. (1855) 406 = truncata.
leucocalyx, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 207.—Madag.
leucomelas, *Poir. Encyc.* v. 432.—Ins. Mascar.
lobata, *Lour. Fl. Cochinch.* 227.—Cochinch.
longifolia, *Spruce, in Journ. Linn. Soc.* v. (1861) 7 = D. Paralea.
Lotus, Linn. Blanco, Fl. Filip. ed. I. 210 = multiflora.
Lotus, *Linn. Sp. Pl.* 1057.—As. temp.
Lotus, Lour. Fl. Cochinch. 226 = Loureiriana.
Loureiriana, *G. Don, Gen. Syst.* iv. 39.—Afr. trop.
lucida, Hort. ex Loud. Gard. Mag. (1841) 394 = virginiana.
lucida, *Wall. Cat.* n. 4127.—Malaya.
lycioides, Desf. in Ann. Mus. Par. vi. (1805) 448 = Royena pallens.
Lycopersicon, *Carr. in Rev. Hortic.* (1878) 470.—Japon.
mabacea, *F. Muell. Austr. Veg. Intercol. Exhib. Eas.* (1866–67) 35.—Australi.
Mabola, Roxb. Hort. Beng. 40; Wall. Cat. n. 4131 A = discolor.
macrocalyx, A. DC. in DC. Prod. viii. 226 = nodosa.
macrocalyx, Klotzsch, in Peters, Reise Mossamb. Bot. 182 = Loureiriana.
macrocarpa, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 242.—N. Caled.
macrocarpa, Korth. ex Hiern, l. c. 168 = Korthalsiana.
macrophylla, *Blume, Bijdr.* 670.—Java.
macrophylla, Wall. Cat. n. 4141 = undulata.
malabarica, Kostel. Allg. Med.-Pharm. Fl. iii. 1099 = D. Embryopteris.
Malacapai, *A. DC. in DC. Prod.* viii. 237.—Ins. Philipp.
Mannii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 205.—Afr. trop.
marttima, *Blume, Bijdr.* 669.—Malaya; Austral.
martabanica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 554.—Burma.
maximowiczii, A. DC. in DC. Prod. viii. 226 = nodosa.
Mazelii, *Thib. & Ketel. ex E. Morr. in Belg. Hortic.* xxiv. (1874) 111.—Japon.
megasperma, *Baker, in Journ. Linn. Soc.* xxi. (1885) 423.—Madag.
melanida, Nenaud, ex A. DC. in DC. Prod. viii. 236 = leucomelas.
melanida, *Poir. Encyc.* v. 431.—Ins. Mascar.
melanida, Sieber, ex Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 201 = discolor.
melanoxylon, Blume, Bijdr. 669 = truncata.
melanoxylon, Hassk. Cat. Pl. Hort. Bogor. Alt. 159 = D. Embryopteris.
melanoxylon, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 169, partim = D. Tupru.
melanoxylon, *Roxb. Pl. Corom.* i. 36. t. 46.—Ind. or.; Zeylan.
melanoxylon, Willd. Sp. Pl. iv. 1109 = D. Ebenum.
membranacea, A. DC. in DC. Prod. viii. 227 = D. Ebenum.
mespiliformis, *Hochst. ex A. DC. in DC. Prod.* viii. 279.—Afr. trop.
mexicana, Schiede, ex Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 238 = texana.
microcarpa, Siebold, in Ann. Soc. Hort. Pays-Bas (1844) 38 = D. Lotus.

DIOSPYROS :—
microcarpa, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 246.—Ins. Timor.
microphylla, *Bedd. Forest. Man. Bot.* 145; *Ic. Pl. Ind. Or.* 27. t. 133.—As. trop.
microthombus, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 187.—Ins. Borbon.
mollis, Wall. ex Steud. Nom. ed. II. i. 514 = chretioides.
montana, Heyne, ex A. DC. in DC. Prod. viii. 259 = D. Tupru.
montana, *Roxb. Pl. Corom.* i. 37. t. 48.—As. trop.
Moonii, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 164 = D. Helferi ?
Moonii, *Thw. Enum. Pl. Zeyl.* 182.—Zeylan.
Morincana, *Hance, in Journ. Bot.* xviii. (1880) 299 — Hongkong.
multiflora, *Blanco, Fl. Filip.* ed. I. 303.—Ins. Philipp.
multiflora, Wall. Cat. n. 4144 = lanceaefolia.
myrmecocarpus, Mart. ex Miq. in Mart. Fl. Bras. vii. 7 = Maba myrmecocarpa.
Nerandii, A. DC. in DC. Prod. viii. 227 = nodosa.
nigra, Blanco, Fl. Filip. ed. II. 211 = D. Ebenum.
nigricans, Dalz. in Hook. Kew Journ. iv. (1852) 110 = assamica.
nigricans, *Wall. Cat.* n. 6351.—Reg. Himal.
nilagirica, *Bedd. Forest. Man. Bot.* 144.—Ind. or.
nodosa, *Poir Encyc.* v. 432.—Ins. Mascar.
oblonga, G. Don, Gen. Syst. iv. 40 = venosa.
oblonga, *Wall. Cat.* n. 4124.—Malaya.
obovata, Jacq. Hort. Schoenb. iii. 34 = tetrasperma.
obovata, Wight, Ic. t. 1226 = lucnandra diplostemon.
obtusifolia, Bert. ex A. DC. in DC. Prod. vii. 234 = Maba inconstans.
obtusifolia, H. B. & K. Nov. Gen. et Sp. iii. 253. t. 247 = D. Ebenum
obtusifolia, *Willd. Sp. Pl.* iv. 1112.—Am. austr.
octandra, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 167 = burmanica.
oleifolia, *Wall. Cat.* n. 4128.—Burma; Malaya.
Olen, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 246.—Ins Fiji.
oligandra, Bedd. Rep. For. Madr. (1867) 68 = Candolleana.
oocarpa, *Thw. Enum. Pl. Zeyl.* 180.—Zeylan.
opaca, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 567. —Zeylan.
oppositifolia, *Thw. Enum. Pl. Zeyl.* 181.—Zeylan.
orizensis, Kiela, ex Willd. Sp. Pl. iv. 407 = ayivatica.
ovicetkis, *Wight, Ic Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 264.—Ind. or.
ovalifolia, *Wight, Ic.* t. 1227.—Ind. or.
ovalis, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 248.—Bras.
pachyphylla, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 566.—Malacca.
Packmanni, *C. B. Clarke, l. c* 564.—Burma.
paniculata, *Dalz. in Hook. Kew Journ.* iv. (1852) 109. —Ind. or.
Paralea, *Steud. Nom.* ed. II. i. 514.—Guian.
parvifolia, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 217.—Madag.
Pearcei, *Hiern, l. c.* 252.—Am. trop.
pellucida, *Hiern, l. c.* 209.—Ins. Philipp.
pendula, *Hasselt, ex Hassk. Pl. Jav.* 465.—Java.
pendulifiora, *Zoll. in Tijdschr. Nederl. Ind.* xiv. (1857) 159.—Java.
pentamera, *Woods & F. Muell. ex F. Muell. Austr. Veg. Intercol. Exhib. Eas.* (1866–67) 35.—Australi.
perfonata, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 243.—Ins. Moluc.
pergamena, *Hiern, l. c.* 234.—Borneo.
Persimon, Wiksr. Jahresb. Bot. 1880 (1884) ed. Berlichm. 92, 96 = virginiana.
peruviana, *Hiern, in Trans. Phil. Camb. Soc.* xii. (1873) 253.—Peruv.
Perrillei, *Hiern, l. c.* 199.—Madag.
philippinensis, *A. DC. in DC. Prod.* viii. 231.—Ins. Philipp.
phyllomega, Steud. Nom. ed. II. i. 514 = macrophylla.
pilosa, *A. DC. l. c.* 210.—Cochinch.
pilosanthera, *Blanco, Fl. Filip.* ed. I. 304.—Ins. Philipp.
planisepala, G. Don, Gen. Syst. iv. 39 = pilosula.
pilosula, *Wall. Cat.* n. 4132.—Ind. or.
platycalyx, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 267.—Ins. Seychelles.
platyphylla, *Hiern, l. c.* 206.—Afr. trop.
plectosepala, *Hiern, l. c.* 201.—Borneo.

DIOSPYROS :—
Poeppigiana, *A. DC. in DC. Prod.* viii. 224.—Bras.
polyalthioides, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 198.—Borneo.
polyandra, *Spruce, in Journ. Linn. Soc.* v. (1861) 7. —Bras.
prunosa, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 208.—Madag.
pruriens, *Dals. in Hook. Kew Journ.* iv. (1852) 110. —Ind. or.
Pseudo-Lotus, *Naud. in Nouv. Arch. Mus. Par.* Sér. II. iii. (1880) 220.—Japon.
psidioides, H. B. & K. Nov. Gen. et Sp. iii. 254 = Maba inconstans.
Pterocalyx, Boj. Hort. Maurit. 200 = melanida.
pterocalycina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 124 = pnæc.
pubescens, Pers. Syn. ii. 625 = Royena hirsuta.
pubescens, Purah, Fl. N. Am. Sept. i. 365 = virginiana.
punctata, Decne. Herb. Tim. Dec. 79 = montana.
punctata, Korth. ex Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 136 = Maba punctata.
pyrrhocarpa, *Miq. Fl. Ind. Bat. Suppl.* 583.— Sumatra.
quæsita, *Thw. Enum. Pl. Zeyl.* 179.—Zeylan.
racemosa, Roxb. Hort. Beng. 40; Fl. Ind. ii. 536 = D. Toposia.
ramiflora, *Roxb. ll. cc* 40; 535.—Ind. or.
reticulata, Sieber, A. DC. in DC. Prod. viii. 236 = leucomelas.
reticulata, Wall. Cat. 159. n. 4120 = D. Ebenum.
reticulata, Willd. Sp. Pl. iv. 1109 = tesellaria.
revoluta, Poir. Encyc. v. 485 = D. Ebenum.
rhodocalyx, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 71.—Siam.
rigida, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 257.—Borneo.
rotundiflora, *Hiern, l. c.* 163.—Borneo.
rotundifolia, *Hiern, l. c.* 181.—Afr. austr.
Roxburghii, Carr. in Rev. Hortic. (1872) 258 = D. Kaki.
sapotoides, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 88.—Burma.
Schi-Tse, Bunge, Enum. Pl. Chin. Bor. 42 = D. Kaki.
senegalensis, Perr. ex A. DC. in DC. Prod. viii. 234 = mespiliformis.
senensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 188.—Afr. trop.
sericea, A. DC. in DC. Prod. viii. 236 = Maba sericea.
sericocarpa, F. Muell. Austr. Veg. Intercol. Exhib. Ess. (1866–67) 85 = Maba rufa.
serrata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 143 = Embya acuminata.
speciosa, Wood, Rep. For. Oudh. (1867–68) 33, nomen = D. Tupru.
sphærocarpa, *Baker, in Journ. Linn. Soc. xxi.* (1885) 493.—Madag.
sphinosa, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 247.—Bras.
Sprucei, *Hiern, l. c.* 210.—N. Granat.
squamosa, *Bej. ex A. DC. in DC. Prod.* viii. 222.— Madag.
squarrosa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 184.—Afr. trop.
stricta, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 201 = elegans.
stricta, Hort. ex Loud. Encyc. Trees & Shrubs, 627 = virginiana.
stricta, *Roxb. Hort. Beng.* 40; *Fl. Ind.* ii. 539.— Ind. or.
subcuta, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 185.—Madag.
suberifolia, *Hiern, l. c.* 189.—Chili.
subrotata, *Hiern, l. c.* 250.—Bras.

DIOSPYROS :—
sumatrana, *Miq. Fl. Sunga.* 208.—Malaya.
sylvatica, *Roxb. Pl. Corom.* i. 56. t. 47.—Ind. or.
sylvatica, Wall. Cat. n. 4117 = montana.
Tectona, Blanco, Fl. Filip. ed. II. 809 = Tectona philippensis.
tesselaria, *Pour. Encyc.* v. 480.—Ins. Mascar.
tetrandra, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 210.—Guiana.
tetrandra, Span. in Linnæa, xv. (1841) 336 = maritima.
tetrasperma, *Sw. Prod. Veg. Ind. Occ.* 62.—Jamaic.
texana, *Scheele, in Linnæa*, xxii. (1849) 145.— Mexic.
Teysmanni, *Miq. Fl. Ind. Bat. Suppl.* 583.—Sumatra.
Thouarsii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 232.—Madag.
Thwaitesii, Bedd. Forest. Man. Bot. 144; Ic. Pl. Ind. Or. 27. t. 135 = hirsuta.
timoriana, Miq. Fl. Ind. Bat. ii. 1045 = D. Ebenum.
tomentosa, Poir. Encyc. v. 486 = chloroxylon.
tomentosa, *Roxb. Hort. Beng.* 40; *Fl. Ind.* ii. 532.— Ind. or.
Toposia, *Buch.-Ham. in Trans. Linn. Soc.* xv. (1827) 115.—Ind. or.
toxicaria, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 175.—Madag.
triculor, *Hiern, l. c.* 183.—Afr. trop.
truncata, *Zoll. & Moritzi, Syst. Verz. Zoll.* 43.—Java.
Tupru, *Buch.-Ham. in Trans. Linn. Soc.* xv. (1827) 111.—Ind. or.
undulata, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 215 = lucida.
undulata, *Wall. Cat.* n. 4186.—Burma; Malaya.
vaccinifolia, Extinguh. in Sitzb. Acad. Wien, xxviii. (1858) 494 = vaccinioides.
vaccinioides, *Lindl. in Hook. Exot. Fl.* ii. t. 139.—China.
vaccinoides, Wall. Cat. n. 4180 = microphylla.
variegata, *Kurz, in Journ. As. Soc. Beng.* xli. (1871) ii. 78.—Burma.
velutina, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 200.—Am. trop.
venosa, *Wall. Cat.* n. 4196.—Ins. Penang.
verrucosa, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 187.—Afr. trop.
Vescol, *Hiern, l. c.* 218.—Madag.
virginiana, *Linn. Sp. Pl.* 1057.—Am. bor.
Waldemarii, Klotzsch, in Bot. Ergeb. Waldem. Reise, 101 = montana.
Weddellii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 253.—Bras.
Wightiana, Wall. Cat. n. 4406 = melanoxylon.
xylopioides, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 8. —Guiana.
Zollingeri, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 222.—Java.

DIOSTEA, Miers, in Trans. Linn. Soc. xxvii. (1870) 103. t. 28 = Bailonia, Bocq. (Verbenac.).
integrifolia, Miers, l. c. 106.—Chili.
infuscata, Miers, l. c. 107 = B. juncea.
juncea, Miers, l. c. 108. t. 28 = B. juncea.
serpens, Miers, l. c. 107.—Chili.
scoparia, Miers, l. c. 104 = Verbena scoparia, *Gill. & Hook.*
stenophylla, Miers, l. c. 105.—Chili.
valdiviana, Miers, l. c. 107 = B. juncea.

DIOTACANTHUS, Benth. in Benth. & Hook. f. Gen. ii. 1100 (1876). *ACANTHACEÆ*, Benth. & Hook. f. l. c.
albiflorus, *Benth. l. c.* 1101.—Ind. or.
grandis, *Benth. l. c.*—Ind. or.

DIOTHONEA, Lindl. in Hook. Journ. Bot. i. (1834) 12. *ORCHIDEÆ*, Benth. & Hook. f. iii. 532.
Gardneana, Lindl. in Ann. & Mag. Nat. Hist. Ser. I. xv. (1845) 197.
Hemisicleia, Lindl. Fol. Orch. (1853).
cottenlaeflora, *Reichb. f. in Linnæa*, xli. (1877) 89.— N. Granat.
gratissima, *Reichb. f. in Bot. Zeit.* x. (1852) 772.— Peruv.
heterothesaes, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 119.—Peruv.
imbricata, *Lindl. Sert. Orch.* 40. n. L.—Mexic.
lindleana, *Lindl. in Hook. Journ. Bot.* i. (1834) 12.— Peruv.

DIOTHONEA :—
opposititolia, Reichb. f. in Linnæa, xxii. (1849) 842 = Hexiea opposititolia.
suavis, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 119.—Peruv.

DIOTIS, Desf. Fl. Atlant. ii. 260 (1798). *COMPOSITÆ*, Benth. & Hook. f. ii. 490.
Nerisa, Spreng. Anleit. ii. ii. 547 (1818).
Otanthus, Hoffngg. & Link, Fl. Port. ii. 264 (1809). candidissima, *Desf. l. c.* 261.—Europ. occ.; Reg. Mediterr.
maritima, Sm. Engl. Fl. iii. 408 = candidissima.

DIOTIS, Schrank. Gen. 688 (1791) = **Eurotia**, Adans (Chenopod.).
atriplicina, Spreng. Syst. iii. 836 = Atriplex pedunculata
atriplicoides, Bieb. Fl. Taur. Cauc. ii. 897 = Atriplex pedunculata.
ceratoides, Willd. Sp. Pl. iv. 868 = Eurotia ceratoides
ferruginea, T. Nees, Gen. Germ. Ic. f. 14–19 = Eurotia ceratoides
lanata, Purah, Fl. Am. Sept. ii. 602 = Eurotia lanata.
revoluta, Nutt. ex Moq. in DC. Prod. xiii. ii. 121 = Eurotia lanata.

DIOTOCARPUS, Hochst. in Flora, xxvi. (1843) 70 = **Pentanisia**, Harv. (Rubiac.).
angustifolius, Hochst. l. c. 71 = P. variabilis.
prunelloides, Hochst. l. c. 70 = P. variabilis.

DIOTOLUS, Tausch, in Flora, xxv. (1842) 284 = **Argyrolobium**, Eckl. & Zeyh. (Legumin.).
Echlonii, Tausch, l. c.—Afr. austr.

DIOTOSPERMA, A. Gray, in Hook. Kew Journ. iv. (1852) 275 = **Ceratogyne**, Turcz. (Compos.).
Drummondii, A. Gray, l. c. 276 = C. oblonoides.

DIOTOSTEMON, Salm-Dyck, in Otto & Dietr. Allg Gartenz. xxii. (1854) = **Cotyledon**, Tourn. (Crassulac.).
Hookeri, Salm-Dyck, l. c.—Mexic.

DIOTOSTEPHUS, Cass. in Dict. Sc. Nat. xlviii. 543 (1827) = **Chrysogonum**, Linn. (Compos.).
repens, Cass. l. c. 544 = C. virginianum.

DIOTOTHECA, Rafin. Fl. Ludov. 74 (1817) = **Sonbiosa**, Tourn. (Dipsac.).
repens, Rafin. Fl. Ludov. 75 (quid ?).—Am. bor.

DIOXIPPE, M. Roem. Syn. Hesper. 33, 45 (1846) = **Glycoosmis**, Corres (Rutac.).
chlorosperma, M. Roem. l. c. 45 = G. chlorosperma.
cyanocarpa, M. Roem. l. c. 46 = G. cyanocarpa.

DIPANAX, Seem. in Journ. Bot. vi. (1868) 180 *ARALIACEÆ*.
Manni, *Seem. l. c.*—Ins. Sandvic.

DIPCADI, Medic. in Act. Acad. Theod. Palat. vi. Phys. 431 (1790). *LILIACEÆ*, Benth. & Hook. f. iii. 699
Polemannia, Berg. ex Schlecht. in Linnæa, i. (1826) 250.
Tricneis, Salisb. Gen. Pl. Fragm. 24 (1866).
Uropetalon, Ker-Gawl. in Bot. Reg. t. 156 (1815).
Zuccagnia, Thunb. in Roem. Archiv, ii. t. 2 (1799).
Zuccagnia, Thumb. Nov. Gen. ix. 127 (1798).
Bakerianum, *Zulue, in Journ. Linn. Soc.* xviii. (1881) 394.—Afr. austr.
Balfourii, *Baker, in Gard. Chron.* (1880) ii. 424.—Ins. Socotra.
ciliare, *Baker, in Journ. Linn. Soc.* xi. (1871) 398.— Afr. austr.
comosum, *Wolw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 347.—Afr. trop.
concanense, *Baker, in Journ. Linn. Soc.* xi. (1871) 399.—Ind. or.
crispum, *Baker, l. c.*—Afr. austr.
erythraeum, *Webb & Benth. Phyt. Canar.* iii. 241.— Arab.; Egypt.
filamentosum, *Medic. in Ustori, Ann. Bot.* ii. (1791) 18.—Afr. austr.
filifolium, *Baker, in Journ. Bot.* xvi. (1878) 232.— Afr. trop.
fulvum, Webb & Benth. Phyt. Canar. iii. 240 = serotinum.

DIPCADI:—
glaucum, *Baker, in Journ. Linn. Soc.* xi. (1871) 401. —Afr. austr.
heterocarpa, *Baker, l. c.* xx. (1858) 274.—Madag.
hyacinthoides, *Baker, l. c.* xl. (1871) 396.—Afr. austr.
hydsuricum, *Baker, l. c.* 397.—Ind. or.
lanceolatum, *Baker, in Journ. Bot.* xvi. (1878) 329.—Afr. trop.
latertitum, *Wolw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 247.—*Ad.* trop.
longifolium, *Baker, in Journ. Linn. Soc.* xi. (1871) 397.—Afr. trop.
minimum, *Webb & Berth. Phyt. Canar.* iii. 241.—Abyss.
montanum, *Baker, in Journ. Linn. Soc.* xi. (1871) 398.—Ind. or.
oxyphem, *Wolw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 246.—Afr. trop.
rigidifolium, *Baker, in Journ. Linn. Soc.* xi. (1871) 399.—Afr. austr.
scrotinum, *Medic. in Act. Acad. Theod. Palat.* vi. *Phys.* (1790) 481.—Europ.; Afr. bor.
setosum, *Baker, in Journ. Linn. Soc.* xi. (1871) 398. —Afr. austr.
taccazeanum, *Baker, l. c.* 400 (taccazeanum).—Abyss.
umbonatum, *Baker, l. c.*—Afr. austr.
unicolor, *Baker, l. c.* 397.—Ind. or.
unifolium, *Baker, l. c.* 399.—Afr. trop.
viride, Moench, Meth. Suppl. 267 = flamentosum.
Welwitschii, *Baker, in Journ. Linn. Soc.* xi. (1871) 400.—Afr. trop.

DIPCADIOIDES, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 493 = **Lachenalia**, Jacq. (Liliac.).
maculata, Medic. l. c. = L. orchidioides.

DIPELTA, Maxim. in Bull. Acad. Pétersb. xxiv. (1878) 60. *CAPRIFOLIACEÆ*.
floribunda, *Maxim. l. c.* 51.—China.

DIPELTA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 578 = **Astragalus**, Tourn. (Legumin.).
turkestanica, Regel & Schmalh. l. c. 579 = A. Dipelta.

DIPERA, Spreng. Syst. iii. 696 (1826) = **Disperis**, Sw. (Orchid.).
capensis, Spreng. l. c. = Disp. capensis.
cucullata, Spreng. l. c. = Disp. cucullata.
secunda, Spreng. l. c. = Disp. secunda.
tenera, Spreng. l. c. = Disp. capensis
villosa, Spreng. l. c. = Disp. villosa.

DIPERIS, Wight, Ic. v. t. 1719 (1852) = prec.

DIPERIUM, Desv. Opusc. 76 (1831) = **Ophiurus**, Gaertn. (Gramin.).
cylindricum, Desv. l. c. = O. perforatus.

DIPETALIA, Rafin. Fl. Tellur. iii. 78 (1836) = **Oligomeris**, Cambess. (1836) (Resedac.).
capensis, Rafin. l. c. = O. capensis.

DIPETALON, Rafin. Sylva Tellur. 108 (1838) = **Cuphea**, P. Br. (Lythrar.).
speciosum, Rafin. l. c. = C. verticillata.

DIPETALUM, Dals. in Hook. Kew Journ. ii. (1850) 38 = **Toddalia**, Juss. (Rutac.).
bilosulare, Dals. l. c. = T. bilosularis.

DIPHACA, Lour. Fl. Cochinch. 458 (1790) = **Ormocarpum**, Beauv. (Legumin.).
Bernieriana, Baill. in Bull. Soc. Linn. Par. i. (1884) 418.—Madag.
cochinchinensis, Lour. Fl. Cochinch. 454 = O. sennoides.
Perrillieana, Baill. in Bull. Soc. Linn. Par. i. (1884) 416.—Madag.

DIPHALANGIUM, Schau. in Linnæa, xix. (1847) 703. *LILIACEÆ*, Benth. & Hook. f. iii. 799.
graminifolium, *Schau. l. c.* 703 (= Milla biflora?).—Mexic.

DIPHEROCARPUS, Llanos, in Mem. Acad. Cienc. Madr. ii. (1838) 501 = **Nephelium**, Linn. (Sapind.).

DIPHOLIS, A. DC. in DC. Prod. viii. 188 (1844). *SAPOTACEÆ*, Benth. & Hook. f. ii. 660.
montana, *Griseb. Fl. Brit. W. Ind.* 401 (1861); *in Mem. Am. Acad. N. S.* viii. (1863) 518.—Ind. occ.
nigra, *Griseb. l. c.* 400.—Ind. occ.
salicifolia, *A. DC. in DC. Prod.* viii. 188.—Ind. occ.

DIPHOREA, Rafin. Neogenyt. 3 (1825) = **Sagittaria**, Linn. (Alismac.).
graminea, Rafin. l. c. = S. graminea.

DIPHRAGMUS, Presl, Bot. Bemerk. 80 (1844); in Abh. Bochm. Ges. V. iii. (1845) 510 = **Spermacoce**, Linn. (Rubiac.).
scaber, Presl, li. cc. 81, 510.—Mexic.

DIPHRYLLUM, Rafin. in Med. Repos. N. York, v. (1808) 357 = **Listera**, R. Br. (Orchid.).
bifolium, Rafin. l. c. = L. bifolia.

DIPHYES, Blume, Bijdr. 310. t. 66 (1825) = **Bulbophyllum**, Thou. (Orchid.).
angustifolia, Blume, l. c. 314 = B. angustifolium.
capitata, Blume, l. c. = B. capitatum.
cernua, Blume, l. c. 318 = B. cernuum.
ciliata, Blume, l. c. 317 = B. ciliatum.
crocea, Blume, l. c. 313 = B. croceum.
flavescens, Blume, l. c. = B. flavescens.
gibbosa, Blume, l. c. 312 = B. gibbosum.
gracilis, Blume, l. c. 319 = B. gracile, *Lindl.*
hirsuta, Blume, l. c. 318 = B. hirsutum.
inæqualis, Blume, l. c. 319 = B. inæquale.
laxiflora, Blume, l. c. 316 = B. laxiflorum.
mucronata, Blume, l. c. 317 = B. mucronatum.
mutabilis, Blume, l. c. 312 = B. mutabile.
obtusa, Blume, l. c. 316 = B. obtusum.
odorata, Blume, l. c. 312 = B. odoratum.
ovalifolia, Blume, l. c. 318 = B. ovalifolium.
pusilla, Blume, l. c. 317 = B. parvulum.
sulcata, Blume, l. c. 314 = B. sulcatum.
tenella, Blume, l. c. 319 = B. tenellum.
tenuifolia, Blume, l. c. 316 = B. tenuifolium.
tortuosa, Blume, l. c. 311 = B. tortuosum.
violacea, Blume, l. c. 316 = B. violaceum.

DIPHYLLEIA, Michx. Fl. Bor. Am. i. 208. t. 19 (1803). *BERBERIDEÆ*, Benth. & Hook. f. i. 44.
Cymosa, *Michx. l. c.*—Am. bor.
Grayi, *F. Schmidt, Reisen Amurl.* 109.—Japon.

DIPHYSA, Jacq. Enum. Pl. Carib. 7 (1760). *LEGUMINOSÆ*, Benth. & Hook. f. i. 500.
carthagenensis, *Jacq. l. c.* 28.—N. Granat.
floribunda, *Peyr. in Linnæa*, xxx. (1859–60) 78.—Am. centr.
humilis, *Oerst. ex Benth. & Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 18.—Mexic.
robinioides, *Benth. ex Benth. & Oerst. l. c.* 11.—Nicaragua.
sennoides, *Benth. ex Benth. & Oerst. l. c.* 12.—Mexic.

DIPHYSTEMA, Neck. Elem. i. 382 (1790) = **Amasonia**, Linn. f. (Verbenac.).

DIPIDAX, Lawa. Cat. 8; ex Saliab. in Trans. Hort. Soc. i. (1812) 330. *LILIACEÆ*, Benth. & Hook. f. iii. 824.
MELANTHIUM, Kunth, Enum. Pl. iv. 155 (1843).
ONIXOTIS, Rafin. Fl. Tellur. ii. 92 (1836).
ciliata, *Baker, in Journ. Linn. Soc.* xvii. (1879) 447. —Afr. austr.
rosea, *Lawa. Cat.* 8; *ex Saliab. in Trans. Hort. Soc.* i. (1812) 330.—Afr. austr.
triquetra, Baker, in Journ. Linn. Soc. xvii. (1879) 447 = rosea.

DIPLACHNE, Beauv. Agrost. 80. t. 16 (1812). *GRAMINEÆ*, Benth. & Hook. f. iii. 1176.
alba, *Hochst. in Flora*, xxv. (1842) 1. *Beibl.* 134, nomen; *et ex Steud. Syn. Pl. Gram.* 248.—Afr. trop.
alopecuroides, Hochst. ex Steud. l. c. 248 = Triodia plumosa.
andropogonoides, Nees, Fl. Afr. Austr. 258 = Avena andropogonoides.

DIPLACHNE:—
arenaria, *Nees, ex Steud. Nom.* ed. II. i. 514; *et ex Hochst. in Flora*, xxxviii. (1855) 427.—Arab.
brevifolia, J. & C. Presl, Rel. Haenk. i. 261 = Festuca Haenkei.
capensis, Nees, Fl. Afr. Austr. 254 (= *Uralepis capensis*).—Afr. austr.
dubia, *Scribner, in Bull. Torrey Bot. Club*, x. (1883) 80.—Mexic.
Elesaine, *Nees, Fl. Afr. Austr.* 254.—Afr. austr.
elongata, Hochst. in Flora, xxv. (1842) 1. Beibl. 134, nomen; et ex Steud. Syn. Pl. Gram. 268 = Eragrostis interrupta.
fascicularis, *Beauv. Agrost.* 80. t. 16. f. 9.—Mexic.
fusca, *Beauv. l. c.* 168—As., Afr. et Austral. trop.
imbricata, *Scribner, in Bull. Torrey Bot. Club*, x. (1883) 80.—Arizona.
indica, *Spreng. Syst.* i. 351.—Ind. or.
livida, *Nees, Fl. Afr. Austr.* 254.—Afr. austr.
lolliformis, *F. Muell. ex Benth. Fl. Austral.* vii. 619.—Austral.
Muelleri, *Benth. l. c.*—Austral.
nana, *Nees, Fl. Afr. Austr.* 259.—Afr. bor. et austr.
nigrescens, *Desv. Opusc* 108.—Am. bor.
parviflora, *Benth. Fl. Austral.* vii. 620.—Austral.
patens, *E. Desv. in C. Gay, Fl. Chil.* vi. 373.—Chili.
patens, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 570.—Mexic.
poæformis, Hochst. in Flora, xxv. (1842) 1. Beibl. 134, nomen = Eragrostis interrupta.
procera, Spreng. Syst. i. 351 = Festuca orgyalis.
rigescens, J. & C. Presl, Rel. Haenk. i. 260 = Festuca rigescens.
rigida, *Benth. ex Vasey, Grass. U. St.* 35.—Texas.
scirpifolia, J. & C. Presl, Rel. Haenk. i. 261 = Festuca scirpifolia.
serotina, *Link, Hort. Berol.* i. 155.—Europ.; As. occ.
simplex, *Doell, in Mart. Fl. Bras.* ii. 111. 97.—Bras.
spicata, *Doell, ex Benth. in Fourn. Linn. Soc.* xix. (1881) 111.—Bras.?
tolucensis, Spreng. Syst. i. 351 = Festuca toluccensis.
sinensis, *Hance, in Journ. Bot.* viii. (1870) 76.—China.
verticillata, *Nees & Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 159.—Chili.
viscida, *Scribner, in Bull. Torrey Bot. Club*, x. (1883) 80.—Am. bor.

DIPLACHNE, R. Br. ex Desf. in Mém. Mus. Par. v. (1819) 272 = **Verticordia**, DC. (Myrtac.).
Baueri, R. Br. ex Desf. l. c. = V. Fontanesii.

DIPLACHYRIUM, Nees, in Flora, xi. (1828) 308 = **Muehlenbergia** ?, Schreb. (Gramin.).
rarum, Nees, l. c.—M. simplex (*vel* Polypogon maritimus ?).

DIPLACRUM, R. Br. Prod. 241 (1810) = **Solaria**, Berg. (Cyperac.).
capitatum, Boeck. in Linnæa, xxxvii. (1874) 436 = S. capitata.
caricinum, R. Br. Prod. 241 = S. caricina.
fygmaeum, Boeck. in Linnæa, xxxvii. (1874) 434 = S. africana.
fygmaeum, Nees, ex Boeck. l. c. = S. pygmæa.
tridentatum, Brongn. in Duperr. Voy. Bot. t. 26 = S. caricina.
seylanicum, Nees, in Wight, Contrib. 119 = S. caricina.

DIPLACTIS, Rafin. Fl. Tellur. ii. 45 (1836) = **Aster**, Tourn. (Compos.).
alpina, Rafin. l. c. = A. alpinus.
blanda, Rafin. l. c. = A. puniceus.
caucasica, Rafin. l. c. = A. caucasicus.
cyanea, Rafin. l. c. = A. laevis.
novanglia, Rafin. l. c. = A. Novae-Angliae.

DIPLACUS, Nutt. in Ann. Nat. Hist. Ser. I. i. (1838) 137 = **Mimulus**, Linn. (Scrophular.).
arachnoideus, Greene, in Bull. Calif. Acad. i. iv. (1885) 210 = M. glutinosus.
auranticus, G. Don, in Sweet, Hort. Brit. ed. III. 816 = M. glutinosus.
cardinalis, Dougl. in Rev. Hortic. Sér. IV. vi. (1857) 137 = M. cardinalis.

DIPLACUS:—
glutinosus, Nutt. in Ann. Nat. Hist. Ser. I. i. (1838) 137 = M. glutinosus.
Godroni, Verach. ex E. Morr. in Belg. Hortic. xiii. (1863) 4 = Mimulus glutinosus!
grandiflorus, Groenl. in Rev. Hortic. Sér. IV. vi. (1857) 408 = M. glutinosus!
latifolius, Nutt. in Ann. Nat. Hist. Ser. I. i. (1838) 138 = M. glutinosus.
leptanthus, Nutt. l. c. = M. glutinosus.
longiflorus, Nutt. l. c. 139 = M. glutinosus.
pumicrus, Nutt. l. c. 137 = M. glutinosus.
rugosus, Benth. in DC. Prod. x. 368 = Berendtia rugosa.
splendidus, Versch. ex E. Morr. in Belg. Hortic. xiii. (1863) 5 = M. glutinosus!
stellatus, Kellogg, in Proc. Calif. Acad. ii. (1863) 18 = M. glutinosus.
Verschaffeltii, Versch. ex E. Morr. in Belg. Hortic. xiii. (1863) 5 = M. glutinosus!

DIPLADENIA, A. DC. in DC. Prod. viii. 491 (1844). *APOCYNACEAE*. Benth. & Hook. f. ii. 726.
MUCRADENIA, Miers, Apocyn. S. Am. 158 (1878).
PRESTONIOPSIS, Muell. Arg. in Bot. Zeit. xviii. (1860) 22.
acuminata, *Hook. Bot. Mag.* t. 4828.—Bras.
alexicaca, A. DC. in DC. Prod. viii. 484 = illustris.
androsaemifolia, A. DC. l. c. = illustris.
amabilis ×, Hort. Backh. ex Flor. & Pomol. (1868) 73.
amoena ×, T. Moore, ex Flor. & Pomol. (1868) 72.
atropurpurea, *A. DC. in DC. Prod.* viii. 486.—Bras.
atropurpurea, Lem. in Fl. des Serres, ii. (1845) t. [29] = atrovioiacea.
atrovioiacea, *A. DC. in DC. Prod.* viii. 484.—Bras.
Billbergii, Beurl. in Ves. Akad. Handl. Stockh. 1854 (1856) 187 = Echites Billbergii.
boliviensis, *Hook. f. Bot. Mag.* t. 5783.—Bolivia.
Brasiliana, Hort. ex Kegel, Gartenfl. (1875) 230, 222 = Odontadenia speciosa!
coccinea, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 132.—Bras.
cordifolia, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1088, nomen.—Guiana.
crassinoda, *A. DC. in DC. Prod.* viii. 486.—Bras.
crassinoda, Lindl. Bot. Reg. (1844) t. 64 = Martiana.
Fendleri, *Muell. Arg. in Linnaea*, xxx. (1859–60) 417. —Venezuela.
flava, *Hook. Bot. Mag.* t. 4709.—N. Granat.
fragrans, *A. DC. in DC. Prod.* viii. 483.—Bras.
Gardneriana, *A. DC. l. c.*—Bras.
gentianoides, *A. DC. l. c.* 486.—Bras.
gentianoides, Muell. Arg. in Mart. Fl. Bras. vi. i. 124 = Echites Pohliana.
Harrisii, Hook. Bot. Mag. t. 4825 = Odontadenia speciosa.
illustris, *A. DC. in DC. Prod.* viii. 483.—Bras.
linariaefolia, A. DC. l. c. 482 = polymorpha.
Lindleyi, *Lem. Illustr. Hortic.* ii. (1855) 13.—Bras.
lineata, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 123.—Bras.
longiloba, *A. DC. in DC. Prod.* viii. 485.—Bras.
Martiana, *A. DC. l. c.*—Bras.
Moricandiana, *A. DC. l. c.* 486.—Bras.
nobilis, *Morr. in Ann. Soc. Gand*, iii. (1847) 331. t. 159; *Lem. in Fl. des Serres*, Sér. I. v. (1849) t. 487.—Bras.
pastorum, A. DC. in DC. Prod. viii. 482 = polymorpha.
peduncularis, A. DC. l. c. = polymorpha.
polymorpha, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 121. —Bras.
pulchella, *A. DC. in DC. Prod.* viii. 485.—Bras.
Riedelii, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 131.—Bras.
Rose-campestris, Lem. in Fl. des Serres, Sér. I. iii. (1847) t. 356 = illustris.
salicifolia, *Regel, Cat. Pl. Hort. Absakov.* 49.—Hab.!
sanctæ, *A. DC. in DC. Prod.* viii. 484.—Bras.
Saponariæ, *A. DC. l. c.* 485.—Bras.
scabra, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 128. —Bras.
Sellowii, *Muell. Arg. l. c.*—Bras.
spigeliaeflora, Muell. Arg. l. c. 122 = longiloba.
splendens, *A. DC. in DC. Prod.* viii. 476.—Bras.

DIPLADENIA:—
stenoloba, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 156.—Peruv.
tenuifolia, *A. DC. in DC. Prod.* viii. 482.—Bras.
urophylla, *Hook. Bot. Mag.* t. 4414.—Bras.
velutina, A. DC. in DC. Prod. viii. 483 = gentianoides.
venulosa, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 126.—Bras.
vincaeflora, *Lem. in Rev. Hortic.* (1853) 183.—Bras.
xanthosiphona, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 133. —Bras.

DIPLANDRA, Bert. in Merc Chil. 612; ex Bull. Férussac, xx. (1830) 110 = **Elodea**, Michx. (Hydroch.). *Potamogeton*, Bert. l. c. = E. chilensis.

DIPLANDRA, Hook. & Arn. Bot. Beech. Voy. 291. t. 80 (1837). *ONAGRARIEAE*, Benth. & Hook. f. i. 792.
lopezoides, *Hook. & Arn. l. c.* 292. t. 80.—Mexic.

DIPLANTHERA, Banks & Soland. ex R. Br. Prod. 449 (1810). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1048.
BULWERIA, F. Muell. Fragm. iv. 147 (1864).
DIPLANCHEA, Vieill. in Bull. Soc. Linn. Normand. vii. (1863) 98.
bancana, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 354; *et ex Hassk. in Flora*, liii. (1870) 220—*Ins. Sunda*.
tetraphylla, *R. Br. Prod.* 449.—Austral.

DIPLANTHERA, Schrank, Pl. Rar. Hort. Monac. t. 62 (1819) = **Justicia**, Linn. (Acanth.).
formosa, Schrank, l. c. = J. furcata.
furcata, Steud. Nom. ed. II. i. 515 = J. furcata.

DIPLANTHERA, Thon. Gen. Nov. Madag. 3 (1806) = **Cynodocea**, Kom. (Nalad.).
indica, Steud. Nom. ed. II. i. 515 = C. isoetaefolia.
madagascariensis, Steud. l. c.—Madag.
tridentata, Steinh. in Ann. Sc. Nat. Sér. II. ix. (1838) 98.—Madag.

DIPLARCHE, Hook. f. & Thoms. in Hook. Kew Journ. vi. (1854) 382. t. 11. *ERICACEAE*, Benth. & Hook. f. ii. 597.
multiflora, *Hook. f. & Thoms. l. c.* 383.—Reg. Himal.
pancifiora, *Hook. f. & Thoms. l. c.*—Reg. Himal.

DIPLARIA, Rafin. ex DC. Prod. vii. 610 (1839) = **Cassandra**, D. Don (Ericac.).

DIPLARINUM, Steud. Nom. ed. II. i. 515 (1840) = seq.

DIPLARINUS, Rafin. in Journ. Phys. lxxxix. (1819) 106 = **Scirpus**, Linn. (Cyperac.).

DIPLARPEA, Triana, in Benth. & Hook. f. Gen. i. 756 (1867). *MELASTOMACEAE*, Benth. & Hook. f. i. c.
paleacea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 80.—N. Granat.

DIPLARRHENA, Labill. Voy. i. 157 (1793) (Diplarrena). *IRIDEAE*, Benth. & Hook. f. iii. 996.
latifolia, *Benth. Fl. Austral.* vi. 400.—Tasmania. Moraea, *Labill. Voy.* i. 157. t. 15.—Austral.

DIPLARRHINUS, Endl. Gen. 119 (1837) = DIPLARINUS, Rafin. = **Scirpus**, Linn. (Cyperac.).

DIPLASANTHUM, Desv. Opusc. 66 (1831) = **Andropogon**, Linn. (Gramin.).
lanosum, Desv. l. c. 67 = A. caricosus.

DIPLASIA, Rich. in Pers. Syn. i. 70 (1805). *CYPERACEAE*, Benth. & Hook. f. iii. 1065.
karstaefolia, *Rich. l. c.*—Guiana.
pycnostachya, *Benth. in Journ. Linn. Soc.* xv. (1877) 513.—Bras.

DIPLAPSIS, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 469 [bis] = **Huanaca**, Cav. (Umbellif.).
cordifolia, Hook. f. Fl. Tasm. i. 157 = H. cordifolia.
hydrocotyloa, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 469 [bis] = H. hydrocotyloa.

DIPLAX, Soland. ex Benn. Pl. Jav. Rar. II (1838) = **Microlsena**, R. Br. (Gramin.).
avenacea, Raoul, Choix Pl. N. Zél. ii. t. 3 = M. avenacea.
polynoda, Hook. f. Fl. N. Zel. 290 = M. polynoda.
tasmanica, Hook. f. Fl. Tasm. ii. 105 = M. tasmanica.

DIPLECOSIA, G. Don, Gen. Syst. iii. 838 (1834) = **Diplycosia**, Blume (Ericac.).

DIPLECTHRUM, Pers. Syn. ii. 508 (1807) = **Satyrium**, Sw. (Orchid.).
amœnum, Thou. Orch. Isles Afr. t. 21, 22 = S. amœnum.
bicallosum, Pers. Syn. ii. 509 = S. bicallosum.
bracteatum, Pers. l. c. = S. bracteatum.
corifolium, Pers. l. c. = S. corifolium.
cucullatum, Pers. l. c. 508 = S. bicorne.
cuculliformum, Salisb. in Trans. Hort. Soc. i. (1812) 267 = pnsc.
erectum, Pers. Syn. ii. 509 = S. corifolium.
foliosum, Pers. l. c. = S. foliosum.
membranaceum, Pers. l. c. = S. membranaceum.
parviflorum, Pers. l. c. = S. parviflorum.
pumilum, Pers. l. c. = S. pumilum.
striatum, Pers. l. c. = S. striatum.

DIPLECTRADEN, Rafin. Fl. Tellur. ii. 90 (1836) = **Habenaria**, Willd. (Orchid.).
leptoceras, Rafin. l. c. = H. leptoceras.

DIPLECTRUM, Endl. Gen. 211 (1837) = **Dipleothrum**, Pers.

DIPLEMIUM, Rafin. Fl. Tellur. ii. 50 (1836) = **Erigeron**, Linn. (Compos.).
caroliniannm, Rafin. l. c. = Soldago tenuifolia.
nervosum, Rafin. l. c. = Chrysopsis graminifolia.
quercifolium, Rafin. l. c. = E. quercifolius.
strigosum, Rafin. l. c. = E. strigosus.

DIPLERISMA, Planch. in Proc. Linn. Soc. i. (1848) 361. et in Tyana. Linn. Soc. xx. (1851) 416. f. 18–20 = **Meliauthus**, Linn. (Sapind.).
comosus, Planch. in Tyana. Linn. Soc. xx. (1851) 416 = M. comosus.
minor, Planch. in Proc. Linn. Soc. i. (1848) 361 = M. minor.

DIPLESTHES, Harv. in Hook. Lond. Journ. Bot. i. (1842) 19 = **Salacia**, Linn. (Celastrin.).
Krausii, Harv. l. c. = S. alternifolia.

DIPLIATHUS, Rafin. Sylva Tellur. 135 (1838) = **Ldtaea**, Lam. (Laurin.).
angulata, Rafin. l. c. = L. glanescens.

DIPLICOSIA, Endl. Gen. 756, in syn. (1839) = **Diplycosia**, Blume (Ericac.).

DIPLIMA, Rafin. Alsog. Am. 13 (1838) = **Salix**, Tourn.
arenaria, Rafin. l. c. = S. repens.
chrysantha, Rafin. l. c. = S. lanata.
lucida, Rafin. l. c. = S. lucida.
stipularis, Rafin. l. c. = S. stipularis.
Stuartiana, Rafin. l. c. = S. Lapponum.
subsessilis, Rafin. l. c. = S. repens!
viminalis, Rafin. l. c. = S. viminalis.
Wrightiana, Rafin. l. c. = S. phyllicifolia.

DIPLISCA, Rafin. Sylva Tellur. 81 (1838) = **Ceanothus**, Linn. (Rhamn.).
ciliatica, Rafin. l. c. = C. reclinatus.

DIPLOCALYMMA, Spreng. Neue Entdeck. iii. 80 (1822). *INC. SED.* Benth. & Hook. f. ii. 868, volabile, *Spreng. l. c.*—Hab.!

DIPLOCALYMNA, Choisy, in DC. Prod. ix. 408 (1845) = pnsc.

DIPLOCALYX, Presl, Bot. Bemerk. 148 (1844) in Abh. Bochm. Ges. V. iii. (1845) 576 = **Mitrasacme**, Cav. (Gesneroc.).
pallidus, Presl, l. c. = M. pallida.
tomentosus, Presl, l. c. 147 = M. tomentella.

DIPLOCALYX, A. Rich. Fl. Cub. Fanerog. ii. 81. t. 54 (1858) = **Schoepfia**, Schreb. (Olacin.).
chrysophylloides, A. Rich. l. c. = S. chrysophylloides.

DIPLOCARDIA, Zipp. ex Blume, Rumphia, iii. 115 (1847) = **Pometia**, Forst. (Sapindac.).
excelsa, Zipp. ex Blume, l. c. (= *Irina Diplocardia*).—Malaya.

DIPLOCEA, Rafin. in Am. Journ. Sc. i. (1819) 252 = **Triplaris**, Beauv. (Cramin.).
barbata, Rafin. l. c. = T. purpurea.

DIPLOCENTRUM, Lindl. Bot. Reg. sub t. 1522 (1832). *ORCHIDEÆ*, Benth. & Hook. f. iii. 582.
congestum, *Wight, Ic.* t. 1682.—Ind. or.
longifolium, Wight, l. c. t. 1681 = recurvum.
recurvum, *Lindl. Bot. Reg.* sub t. 1522.—Ind. or.

DIPLOCERAS, Meisn. Gen. Comm. 340 (1845) = **Parolinia**, Webb (Crucif.).

DIPLOCHAETE, Nees, in Linnæa, ix. (1834) 296 = **Rynchospora**, Vahl (Cyperac.).
guianensis, Nees, l. c. = R. pterocarpa.

DIPLOCHILUS, Lindl. Bot. Reg. sub t. 1499 (1832) = **Diplomeris**, D. Don (Orchid.).
hirsutus, Lindl. l. c. = Diplomeris hirsuta.
longifolius, Lindl. l. c. = Diplomeris pulchella.

DIPLOCHITA, DC. Prod. iii. 176 (1828) = **Miconia**, Ruiz & Pav. (Melastom.).
albicans, Heynh, Nom. i. 877 = M. holosericea.
angustifolia, Benth. Pl. Hartw. 268 = M. pallens.
aurea, DC. Prod. iii. 177 = M. aurea.
bracteata, DC. l. c. 176 = M. holosericea.
bubalina, DC. l. c. 177 = M. bubalina.
caudata, DC. l. c. = M. Donaeana.
florida, DC. l. c. 176 = M. florida.
Fothergilla, DC. l. c. = M. Fothergilla.
leucophæa, DC. l. c. 177 = M. macrophylla.
macrophylla, DC. l. c. = M. macrophylla.
macrotis, Griseb. Fl. Brit. W. Ind. 711.—Ins. Trinitat.
mucronata, DC. Prod. iii. 177 = M. holosericea.
parviflora, Benth. in Hook. Journ. Bot. ii. (1840) 302 = M. verticillata.
parvifora, Griseb. Fl. Brit. W. Ind. 252, partim = M. argentea.
pyramidalis, Heynh. Nom. i. 877 = M. laevigata.
rosea, Macfad. Fl. Jamaic. ii. 74 = M. Fothergilla.
rufipila, Steud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 104 = M. rufipila.
serrulata, DC. Prod. iii. 177 = M. macrophylla.
sessilifolia, Griseb. Fl. Brit. W. Ind. 252, partim = M. ampla, Pinkeneti.
Swartziana, Benth. in Hook. Journ. Bot. ii. (1840) 302 = M. holosericea.
Swartziana, DC. Prod. iii. 177 = M. Fothergilla.
tomentosa, Griseb. Fl. Brit. W. Ind. 252 = M. tomentosa.

DIPLOCHITON, Spreng. Gen. i. 340 (1830) = praec.

DIPLOCHLAENA, Spreng. Gen. i. 369 (1830) = **Diplolaena**, R. Br.

DIPLOCHLAMYS, Muell. Arg. in Flora, xlvii. (1864) 589 = **Mallotus**, Lour. (Euphorb.).
Griffithianus, Muell. Arg. l. c.—Ind. or.

DIPLOCHONIUM, Fenzl, Nov. Stirp. Dec. 57 (1839) = **Trianthema**, Linn. (Ficoid.).
acuminoides, Fenzl, t. 56 = T. hydaspica.
acuminoides, Wawra & Peyr. Sert. Beng. 94 = Sesuvium digynum.

DIPLOCLINIUM, Lindl. Veg. Kingd. 819 (1847) = **Begonia**, Linn.
angustifolium, Miq. Fl. Ind. Bat. i. 1. 687 = B. angustifolia.
apterum, Miq. l. c. 691 = B. apteta, *Blume*.
arcuatum, Miq. l. c. 689 = B. arcuata.
Armattianum, Wight, Ic. t. 1215 = B. cordifolia.
atrichum, Miq. Fl. Ind. Bat. i. 1. 1091 = B. atricha.
bilocularis, Wight, Ic. t. 1214 = B. sinuata, *Wall.*
bonplandianum, Klotzsch, in Abh. Akad. Berl. (1855) 72 = B. insignis.

DIPLOCLINIUM :—
bracteatum, Miq. Fl. Ind. Bat. i. 1. 688 = Begonia bracteata, *Roxb.*
caespitosum, Miq. l. c. 685 = B. caespitosa.
carnosum, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1862) 420 = B. carnosa.
cordifolium, Wight, Ic. t. 1216 = B. cordifolia.
Cumingianum, Miq. Fl. Ind. Bat. i. 1. 691 = B. Cumingiana.
Evansianum, Lindl. Veg. Kingd. ed. I. 818. t. 220 = B. Evansiana.
fasciculatum, Miq. Fl. Ind. Bat. i. 1. 690 = B. fasciculata.
Hasskarlianum, Miq. l. c. 1091 = B. Hasskarliana.
holosericeum, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1862) 421 = B. holosericea.
Horsfieldii, Miq. Fl. Ind. Bat. i. 1. 691 = B. Horsfieldii.
lepidum, Miq. l. c. 689 = B. lepida.
Lindleyanum, Wight, Ic. t. 1217 = B. fallax.
longifolium, Miq. Fl. Ind. Bat. i. 1. 687 = B. longifolia.
orbiculatum, Miq. l. c. 686 = B. orbiculata.
pilosum, Miq. l. c. 688 = B. pilosa.
racemosum, Miq. l. c. 691 = B. racemosa.
repandum, Klotzsch, in Abh. Akad. Berl. (1855) 72 = B. isoptera.
repens, Miq. Fl. Ind. Bat. i. 1. 686 = B. mollis.
Roxburghii, Miq. l. c. 692 = B. Roxburghii.
rubrum, Miq. l. c. 689 = B. robusta.
saxatile, Miq. l. c. 686 = B. saxatilis.
subhirtum, Miq. l. c. 690 = B. subhirta.
tenuifolium, Miq. l. c. 693 = B. tenuifolia.
timorense. Miq. l. c. 692 = B. aptera, *Roxb.*
tuberosum, Miq. l. c. 685 = B. tuberosa.
Wallichianum, Miq. l. c. 690 = B. Wallichiana.

DIPLOCLISIA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 37, 49 = Cocculus, DC. (Menisperm.).
incisyta, Miers, Contrib. iii. 282 = C. macrocarpus.
lepida, Miers, l. c. 284 = C. macrocarpus.
macrocarpa, Miers, in Ann. & Mag. Nat. Hist. Ser. II. viii. (1851) 43 = C. macrocarpus.
pictinervis, Miers, Contrib. iii. 284 = C. macrocarpus.

DIPLOCOEA, Reichb. Consp. 58, in syn. (1828) = DIPLOCEA, Rafin.

DIPLOCOMA, D. Don, in Sweet, Brit. Flow. Gard. t. 346 (1826) = **Heterolepis**, Cass. (Compos.).
villosa, D. Don, l. c. = H. inuloides.

DIPLOCONCHIUM, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 428 = **Agrostophyllum**, Blume (Orchid.).
inocephalum, Schau. l. c.—Ins. Luzon.

DIPLOCOS, Bureau, in DC. Prod. xvii. 215 (1873) = **Taxotrophia**, Blume (Urticac.).
macrophylla, Bureau, l. c. 216.—Ins. Celebes.
scylanica, Bureau, l. c. 215 = T. zeylanica.

DIPLOCRATER, Benth. in Hook. Kew Journ. iii. (1851) 367 = **Cathedra**, Miers (Olacin.).
acuminatus, Benth. l. c. = C. acuminata.

DIPLOCRATER, Hook. f. in Benth. & Hook. f. Gen. ii. 96 (1873). *RUBIACEÆ*, Benth. & Hook. f. l. c.
coriacea, *Hook. f. l. c.*—Afr. trop.
reticulata, *Hook. f. l. c.*—Afr. trop.

DIPLOCYATHA, N. E. Br. in Journ. Linn. Soc. xvii. (1878) 167. *ASCLEPIADEÆ*.
ciliata, *N. E. Br. l. c.*—Afr. austr.

DIPLODISCUS, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 285. *TILIACEÆ*, Benth. & Hook. f. i. 282.
paniculatus, *Turcz. l. c.*—Ins. Philipp.

DIPLODIUM, Sw. in Ges. Nat. Fr. Berl. Mag. iv. (1810) 84. t. 2 = **Pterostylis**, R. Br. (Orchid.).
australe, Sw. l. c. = P. praecox.

DIPLODON, DC. Prod. iii. 94 (a) (1828) = **Diplusodon**, Pohl (Lythrar.).

DIPLODONTA, Karst. in Linnæa, xxviii. (1856) 439 = **Heterotrichum**, DC. (Melastom.).
setosa, Karst. l. c. = H. octonum.

DIPLOGAMA, Opiz, Semam, 38 (1852) = **Silene**, Linn. (Caryophyll.).
Otites, Opiz, l. c. = S. Otites.

DIPLOGASTRA, Welw. ex Reichb. f. in Flora, xlviii. (1865) 188 = **Platylepis**, A. Rich. (Orchid.).
angolensis, Welw. ex Reichb. f. l. c. = P. glandulosa.

DIPLOGENAEA, A. Juss. in Orb. Dict. viii. 75 (1849) = seq.

DIPLOGENEA, Lindl. in Brand, Quart. Journ. (1828) 122 = **Medinilla**, Gaudich. (Melastom.).
viscoides, Lindl. l. c. = M. viscoides.

DIPLOGLOSSIS, Benth. & Hook. f. Gen. ii. 762 (1876), sphalm. = seq.

DIPLOGLOSSUM, Meisn. Gen. Pl. 269 (1840) = **Cynanchum**, Linn. (Asclepiad.).
auriculatum, Meisn. Gen. Comm. 176 = C. auriculatum.
Jacquemontianum, Meisn. l. c. = C. Jacquemontianum.
Roylei, Meisn. l. c. = C. Royleï.

DIPLOGLOTTIS, Hook. f. in Benth. & Hook. f. Gen. i. 895 (1862). *SAPINDACEÆ*, Benth. & Hook. f. i. c.
australis, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 513, 545 = Cunninghamii.
Cunninghamii, *Hook. f. in Benth. & Hook. f. Gen.* i. 895.—Austral.

DIPLOGON, Rafin. in Am. Monthly Mag. (1818) 268 = **Chrysopsis**, Nutt. (Compos.).
ericoides, Rafin. Atl. Journ. 146 = Aster ericaefolius.
graminifolium, Rafin. Cat. 18 = C. graminifolia.
Marianum, Rafin. in Am. Monthly Mag. (1818) 268 = C. Mariana.
stenophyllum, Rafin. ex DC. Prod. v. 327 = C. graminifolia.

DIPLOKENEMA, Pierre, in Arch. Néerl. xix. (1884) 104. *SAPOTACEÆ*.
sebiferu, *Pierre, l. c.* 106.—Borneo.

DIPLOLAENA, R. Br. in Flind. Voy. ii. 546 (1814) *RUTACEÆ*, Benth. & Hook. f. i. 395.
Diplochlaena, Spreng. Gen. i. 369 (1830).
Ventenatum, Leschen. ex Reichb. Consp. 199 (1828).
angustifolia, *Hook. Bot. Mag.* sub t. 4059.—Austral.
Dampieri, *Desf. in Mém. Mus. Par.* iii. (1817) 452.—Austral.
grandiflora, *Desf. l. c.* 451.—Austral.
Hugelii, *Hort. in Hamb. Gartens.* ii. (1846) *Fruitl.* 8.—Austral.
microcephala, *Bartl. in Lehm. Pl. Preiss.* i. 178.—Austral.
salicifolia, Bartl. l. c. = angustifolia.
speciosa, *Brongn. in Nov. Hortic. Sér.* II. iv. (1845–46) 138.—Austral.

DIPLOLEPIS, R. Br. in Mem. Wern. Soc. i. (1809) 41. *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 768.
Sonninia, Poepp. ex Decne. in DC. Prod. viii. 586 = Oxypetalum Hookeri.
Menziesii, *Schult. Syst.* vi. 95.—China.
ovata, *Lindl. in Trans. Hort. Soc.* vi. (1826) 268.—China.
vomitoria, Lindl. l. c. 68 = Tylophora asthmatica.

DIPLOLOBIUM, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 489 = **Swainsona**, Salisb. (Legumin.).
Walcottii, F. Muell. l. c. = S. occidentalis.

DIPLOLOMA, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 185 = **Craniospermum**, Lehm. (Borrag.).
echioides, Schrenk, l. c.—Sibir.

DIPLOLOPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 173. *UMBELLIFERÆ*, Benth. & Hook. f. i. 900.
abyssinicum, Benth. & Hook. f. l. c. = africanum.
africanum, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i. 178.—Afr. trop.
zambesianum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 18. —Afr. trop.

DIPLOMA, Rafin. Fl. Tellur. iii. 19 (1836) = **Gentiana**, Tourn.
altaica, Rafin. l. c. = G. altaica.
angustifolia, Rafin. l. c. = G. angustifolia.
hudsonica, Rafin. l. c. 27 = G. linearis.
pyrenaica, Rafin. l. c. 19 = G. pyrenaica.
tennifolia, Rafin. l. c. 27 = G. linearis ?

DIPLOMERIS, D. Don, Prod. Fl. Nep. 26 (1825). *ORCHIDEÆ*, Benth. & Hook. f. iii. 627.
DIPLOCHILUS, Lindl. Bot. Reg. sub t. 1498 (1833).
PARAGNATHIS, Spreng. Syst. iii. 675 (1826).
hirsuta, *Lindl. Gen. et Sp. Orch.* 331.—Reg. Himal.
pulchella, *D. Don, Prod. Fl. Nep.* 26.—Reg. Himal.

DIPLOMORPHA, Griff. Notul. iv. 479 (1854) = **Agrocia**, Linn. (Euphorb.).
herbacea, Griff. l. c. = A. baccifformis.

DIPLOMORPHA, Meisn. in Denkschr. Regensb. Ges. iii. (1841) 289 = **Wikstroemia**, Endl. (Thymel.).
canescens, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 357 = W. canescens.
Chamædaphne, C. A. Mey. l. c. 358 = W. Chamædaphne.
japonica, Endl. Gen. Suppl. iv. 11. 66 = Stellera japonica.
virgata, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 357 = W. canescens.
viridiflora, C. A. Mey. l. c. 358 = W. viridiflora.

DIPLONEMA, G. Don, Gen. Syst. iv. 43 (1837) = **Bucklea**, Linn. (Ebenac.).
ambigua, G. Don, l. c. = Royena ambigua.
elliptica, G. Don, l. c. = E. polyandra.

DIPLONYX, Rafin. Fl. Ludov. 101 (1817), ex descr. = **Wisteria**, Nutt. (Legumin.).
elegans, Rafin. l. c. = W. speciosa.

DIPLOPAPPUS, Cass. in Dict. Sc. Nat. xlii. 308 (1819) = **Aster**, Tourn. (Compos.).
albus, Lindl. ex Hook. Fl. Bor. Am. ii. 21 = A. ptarmicoides.
alpinus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 404 = A. scopulorum.
amygdalinus, Hook. Fl. Bor. Am. ii. 28 = A. umbellatus.
annuus, Bluff & Fingerh. Comp. Fl. Germ. ii. 368 = Erigeron annuus.
asper, Less. Syn. Comp. 168 = A. hispidus.
asperrimus, DC. Prod. v. 277 = A. trinervius.
aspersus, DC. l. c. = A. asperulus.
australasicus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 171 = Vittadinia australis.
baccharoides, Benth. in Hook. Lond. Journ. Bot. i. (1842) 487 = A. baccharoides.
bellidifolius, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 46 = Aplopappus glutinosus !
bellidioides, Gill. & Don, ex Hook. & Arn. l. c. 50 = Erigeron uniflorus.
canadinus, Gill. & Don, ex Hook. & Arn. l. c. 48 = Leucopsis sericea.
canescens, Hook. Fl. Bor. Am. ii. 21 = Erigeron canadense.
canescens, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 47.—Chili.
caucasicus, DC. Prod. v. 277 = A. caucasicus.
Chamissonis, Less. in Linnæa, vi. (1831) 111 = Aplopappus Chamissonis.
chinensis, Less. Syn. Comp. 165 = Callistephus hortensis.
chrysanthemifolius, Less. in Linnæa, vi. (1831) 116 = Aplopappus chrysanthemifolius.
coquimbensis, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 47.—Chili.
cornifolius, Darlingt. Fl. Cestr. 474 = A. infirmus.
cornifolius, Lindl. ex Torr. & Gray, Fl. N. Am. ii. 183 = A. umbellatus.

DIPLOPAPPUS :—
coronopifolius, Less. in Linnæa, vi. (1831) 113 = Aplopappus coronopifolius.
corymbosus, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 48 = A. tuberosus (in Addend.).
cuneatus, Hook. & Arn. l. c. 46 = Aplopappus scrobiculatus !
delphinifolius, Cass. in Dict. Sc. Nat. xlii. 310, in texte = Erigeron delphinifolius.
diffusus, Less. in Linnæa, vi. (1831) 110, in obs. ; Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 46 = Leucopsis diffusa.
diplostephioides, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 46 = A. diplostephioides.
divaricatus, Hook. Comp. Bot. Mag. i. (1835) 97 = Aplopappus divaricatus.
Douglasii, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 47 = Aplopappus pulchellus.
dubius, Cass. in Bull. Soc. Philom. (1817) 187 = Erigeron annuus.
dysentericus, Bluff & Fingerh. Comp. Fl. Germ. ii. 369 = Pulicaria dysenterica.
elegans, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iii. 250 = A. tricephalus, himalaicus.
elongatus, DC. Prod. v. 276 = A. filifolius.
elongatus, Gill. & Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 49 = A. Gilliesii.
ericoides, Less. in Linnæa, vi. (1831) 117 = Aplopappus ericoides.
ericoides, Torr. & Gray, Fl. N. Am. ii. 182 = A. ericoides, DC.
extenuatus, DC. Prod. v. 276 = A. fruticosus.
filifolius, DC. l. c. = A. filifolius.
filifolius, Hook. Fl. Bor. Am. ii. 21 = Erigeron filifolius.
filifolius, Less. ex Steud. Nom. ed. II. i. 516 = Hysterionica pinifolia.
foliosus, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 45 = Aplopappus foliosus.
foliosus, Hook. & Arn. l. c. 46 = *ilicifolius*.
fruticulosus, Less. in Linnæa, vi. (1831) 118 = A. fruticulosus.
glandulosus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 83 = Olearia adenolasia.
glutinosus, Poepp. ex Less. in Linnæa, vi. (1831) 113 = Aplopappus glutinosus.
gnaphalioides, Cass. in Dict. Sc. Nat. xxv. 96 = Erigeron gnaphalioides.
graminifolius, Less. in Linnæa, v. (1830) 144 = Chrysopsis graminifolia.
grandiflorus, Hook. Fl. Bor. Am. ii. 21 = Erigeron speciosus.
grindelioides, Less. in Linnæa, vi. (1831) 115 = Aplopappus Berterii ?
hispidus, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 48 = Hysterionica jasionoides.
hispidus, Hook. Fl. Bor. Am. ii. 23 = Chrysopsis hispida.
ilicifolius, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 253.—Reg. Argent.
incanus, Lindl. Bot. Reg. t. 1698 = A. canescens.
indgerrimus, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 47.—Chili.
intermedius, Cass. in Dict. Sc. Nat. xlii. 309 = Chrysopsis gossypina ?
inuloides, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 46 = Aplopappus Berterii.
laevigatus, Sond. in Harv. & Sond. Fl. Cap. iii. 35.—Cape.
lanatus, Cass. in Dict. Sc. Nat. xlii. 309 = Chrysopsis gossypina.
lanceolifolius, Cass. l. c. xxv. 96 = A. lavandulaceus.
laxus, Benth. in Hook. Lond. Journ. Bot. i. (1842) 487 = A. trinervius.
leucophyllus, Lindl. in DC. Prod. v. 278 = Corethrogyne filaginifolia.
linariifolius, Lindl. l. c. 277 = A. linariifolius.
linearis, Hook. Fl. Bor. Am. ii. 21 = Erigeron ochroleucus.
luisseanus, Lindl. in DC. Prod. v. 278 = A. ptarmicoides.
macrocephalus, Poepp. ex Less. in Linnæa, vi. (1831) 114 = Aplopappus macrocephalus.
Mariensis, Hook. Comp. Bot. Mag. i. (1835) 97 = Chrysopsis Mariana.
mollisculosus, Lindl. in DC. Prod. v. 277 = A. mollisculus.

DIPLOPAPPUS :—
mucronatus, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 46 = Aplopappus Remyianus.
natalensis, Sch. Bip. ex Walp. Rep. ii. 957.—Afr. austr.
obovatus, Torr. & Gray, Fl. N. Am. ii. 184 = A. rectilatus.
obtusatus, Less. in Linnæa, vi. (1831) 118 = A. obtusatus.
occidentalis, Hook. & Arn. Bot. Beech. Voy. 350 = Erigeron foliosus.
paludosus, Lindl. ex DC. Prod. v. 264 = A. paludosus.
passerinoides, Turcz. in Bull. Soc. Nat. Mosc. xxiv (1851) ii. 83 = Olearia passerinoides.
pinifolius, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 48 = Hysterionica pinifolia.
pinifolius, Less. ex DC. Prod. v. 296, add. = Erigeron Vahlii.
pinnatifidus, Hook. Fl. Bor. Am. ii. 22 = Aplopappus spinulosus, *DC.*
pinnatifidus, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 48.—Chili.
pluriflorus, Steud. Nom. ed. II. i. 516 = A. pluriflorus.
Poeppigianus, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 47 = Aplopappus Poeppigianus.
prunelioides, Poepp. ex Less. in Linnæa, vi. (1831) 111 = Aplopappus prunelioides.
ptarmicoides, Lindl. ex DC. Prod. v. 264 = Aster ptarmicoides.
pubescens, Cass. in Dict. Sc. Nat. xxv. 96 = Erigeron pubescens.
pulicarius, Ledeb. Fl. Ross. ii. 509 = Pulicaria vulgaris.
puncticulus, Less. ex DC. Prod. vii. 280 = Aplopappus pulchellus.
rigidus, Lindl. in DC. l. c. v. 277 = A. linariifolius.
rotundifolius, Less. in Linnæa, vi. (1831) 116 = Olearia dentata.
Royleï, DC. Prod. v. 276 = Erigeron multiradiatus.
scaber, Hook. Fl. Bor. Am. ii. 22 = Heterotheca grandiflora.
scaber, Less. ex Nees, Gen. et Sp. Ast. 188 = A. scaber, *Thunb.*
scrobiculatus, Nees, l. c. 284 = Aplopappus scrobiculatus.
sericeus, Hook. Comp. Bot. Mag. i. (1835) 97 = Chrysopsis graminifolia.
sericeus, Less. in Linnæa, vi. (1831) 110 = Leucopsis sericea.
serrulatus, Harv. in Harv. & Sond. Fl. Cap. iii. 36.—Afr. austr.
setiger, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 47.—Chili.
spinulosus, Hook. & Arn. l. c. 46.—Reg. Argent.
stenophyllus, Hook. & Arn. l. c. 48 = Hysterionica pinifolia.
tortifolius, Less. in Linnæa, vi. (1831) 118 = A. filifolius.
trichophyllus, Hook. Comp. Bot. Mag. i. (1835) 97 = Chrysopsis trichophylla.
trinervius, Volgt, Hort. Suburb. Calc. 408 = A. trinervius.
turbestanicus, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 615 = A. turkestanicus.
umbellatus, Hook. Fl. Bor. Am. ii. 23 = A. umbellatus.
ventosus, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 48 = Leucopsis sericea.
villosus, Cass. in Dict. Sc. Nat. xlii. 309 = A. villosus.
villosus, Hook. Fl. Bor. Am. ii. 22 = Chrysopsis villosa.
villosus, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 48 = Hysterionica montevidensis.
vulgaris, Bluff & Fingerh. Comp. Fl. Germ. ii. 369 = Pulicaria vulgaris.

DIPLOPELEIS, Endl. in Enum. Pl. Hueg. 13 (1837). *SAPINDACEÆ*, Benth. & Hook. f. i. 387.
Huegelii, *Endl. l. c.*—Austral.
Lehmannii, Miq. in Lehm. Pl. Preiss. i. 234 = Huegelii.
pesticlada, *F. Muell. ex Benth. Fl. Austral. i.* 446.—Austral.
Preissii, Miq. in Lehm. Pl. Preiss. i. 233 = Huegelii.
Stuartii, *F. Muell. Fragm.* iii. 13.—Australasia.

DIPLOPENTA, Alef. in Oestr. Bot. Zeitschr. (1862) 10 = Pawonia, Cav. (Malvac.).
leptocarpa, Alef. l. c. 11 = P. typhalaea.
odorata, Alef. l. c. = P. odorata.
pedunculosa, Alef. l. c. = P. sulcicalyx ?

DIPLOPETALON, Spreng. Syst. iv. Cur. Post. 146, 150 (1827) = **Cupania**, Plum. (Sapindac.).
glaucum, Spreng. l. c. 150 = C. glauca.

DIPLOPHRACTUM, Desf. in Mém. Mus. Par. v. (1819) 94. t. 1. *TILIACEAE*, Benth. & Hook. f. i. 232.
auriculatum, *Desf. l. c.*—Java.
salicifolium, Spreng. Syst. iv. Cur. Post. 205 = Mroscemma salicifolia.

DIPLOPHRAGMA, Meissn. Gen. 160 (1838) = **Hedyotis**, Linn. (Rubiac.).
ovalis, Korth. in Nederl. Kruidk. Arch. ii. 11. (1857) 150 = H. ovalis.
tetragularis, Korth. l. c. 149 = H. quadrangularis.

DIPLOPHYLLUM, Lehm. in Ges. Naturf. Fr. Berl. Mag. vii. (1814) 310 = **Veronica**, Tourn. (Scrophul.).
cardiocarpum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 417 = V. biloba.
Crista-galli, Otto, ex Walp. Rep. iii. 335 = V. Crista-galli.
hirsutum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 417 = V. Crista-galli.
veronicaeforme, Lehm. in Ges. Naturf. Fr. Berl. Mag. viii. (1814) 311 = V. Crista-galli.

DIPLOPOGON, R. Br. Prod. 176 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1145.
DIPOGONIA, Beauv. Agrost. 125 (1812).
setaceus, *R. Br. l. c.*—Australia.

DIPLOPRION, Viv. Fl. Lib. Spec. 48. t. 19 (1824) = **Medicago**, Tourn. (Legum.).
medicaginoides, Viv. l. c. = M. coronata.

DIPLOPTERYS, A. Juss. in Deless. Ic. Sel. iii. 20. t. 33 (1837). *MALPIGHIACEAE*, Benth. & Hook. f. i. 260.
Parsiana, *A. Juss. l. c.*—Guiana.

DIPLOPYRAMIS, Welw. Apont. 591 (1858). *POLYGONACEAE*.
aethiopica, *Welw. l. c.*—Afr. trop.

DIPLORHYNCHUS, Welw. ex Ficalho & Hiern, in Trans. Linn. Soc. Ser. II. ii. (1881) 92. *APOCYNACEAE*.
mossambicensis, *Benth. in Hook. Ic. Pl.* t. 1255.—Afr. trop.
pallopus, *Welw. ex Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 28.—Afr. trop.

DIPLOSASTERA, Tausch, Hort. Canal. t. 4 (1832) = **Corsopsis**, Linn. (Compos.).
tinctoria, Tausch, l. c. = C. tinctoria.

DIPLOSCYPHUS, Liebm. in Vidensk. Selsk. Skr. V. ii. (1850) 282 = **Boleria**, Berg. (Cyperac.).
mexicanus, Liebm. l. c. = S. mexicana.

DIPLOSIPHON, Decne. in Jacquem. Voy. Bot. 166. t. 167 (1844) = **Blyxa**, Thou. (Hydrocharid.).
oryzetorum, Decne. l. c. = B. Roxburghii.

DIPLOSPORA, DC. Prod. iv. 477 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 96.
DIPLOSPERMUM, Dals. in Hook. Kew Journ. ii. (1850) 287.
aplocarpa, *Hook. f. Fl. Brit. Ind.* iii. 133.—Ind. or. australis. *Benth. Fl. Australiensis*. iii. 413.—Austral.
confusa, *Hook. f. Fl. Brit. Ind.* iii. 134.—Burma. Dalzellii, *Hook. f. l. c.* 133.—Ind. or.
erythrocarpa, *Hook. f. l. c.* 134.—Ceylon.
Griffithii, *Hook. f. l. c.*—Burma.
ixoroides, *F. Muell. Fragm.* ix. 133.—Austral.
lyrulata, *Miq. Fl. Ind. Bat.* ii. 366.—Java.
Kurzii, *Hook. f. Fl. Brit. Ind.* iii. 135.—Burma.
malaccensis, *Hook. f. l. c.* 134.—Malacca.
pubescens, *Hook. f. l. c.*—Burma.
singularis, *Korth. in Nederl. Kruidk Arch.* ii. 11. (1851) 201.—Ind. or.; Malaya.
singularis, Kurz, For. Fl. Brit. Burma, ii. 50, partim = confusa.
sphaerocarpa, *Hook. f. Fl. Brit. Ind.* iii. 132.—Ind. or.
truncata, Sund. Nom. ed. II. i. 516 = Triscidia truncata.
viridiflora, *DC. Prod.* iv. 477.—China.

DIPLOSTEGIUM, D. Don, in Mem. Wern. Soc. iv. (1823) 296 = **Tibouchina**, Aubl. (Melastom.).
canescens, D. Don, l. c. = T. canescens.

DIPLOSTELMA, Rafin. New Fl. Am. ii. 44 (1836) = **Chastocapsys**, DC. (Compos.).
bellidioides, A. Gray, Pl. Fendl. 78 = C. modesta.
filiformis, Rafin. New Fl. Am. ii. 45 = C. asteroides.
pumila, Rafin. l. c. = C. asteroides.
tadiana, Rafin. l. c. = C. asteroides.

DIPLOSTEMMA, 'Neck.' ex DC. Prod. vii. 306, in nota (1838); lapsu = DIPHYSTEMA, Neck. = **Amasonia**, Linn. f. (Verbenac.).

DIPLOSTEMMA, Steud. & Hochst. ex DC. Prod. vii. 75 (1838) = **Geigeria**, Griesel. (Compos.).
acaulis, Sch. Bip. in Flora, xxv. (1842) 1. Beibl. 134, nomen; et ex Oliver & Hiern, in Fl. Trop. Afr. iii. 368 = G. acaulis.
alatum, DC. Prod. vii. 75 = G. alata.

DIPLOSTEMON, DC. ex Steud. Nom. ed. II. i. 76 (1840); Miq. Fl. Ind. Bat. i. i. 615 (1855) = **Ammania**, Houst. (Lythrar.).
octandrum, Miq. Fl. Ind. Bat. i. i. 615 = A. octandra.

DIPLOSTEPHIUM, H. B. & K. Nov. Gen. et Sp. iv. 96 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 275.
LINOCHILUS, Benth. Pl. Hartw. 197 (1845).
SIMBLOCLINE, DC. Prod. v. 297 (1836).
aculeatum, Nees, Gen. et Sp. Aster. 129 = Olearia ramulosa.
acuminatum, DC. Prod. v. 278 = Aster acuminatus.
affine, *Wedd. Chlor. And.* i. 208.—Peruv.
amygdalinum, Cass. in Dict. Sc. Nat. xxxvii. 486 = Aster umbellatus.
anactinotum, *Wedd. Chlor. And.* i. 201.—N. Granat.
boreale, Spreng. Syst. iii. 544 = Aster reticulatus.
canum, *A. Gray, in Proc. Am. Acad.* xi. (1876) 75.—Calif.
carabayense, *Wedd. Chlor. And.* i. 209.—Peruv.
cornifolium, DC. Prod. v. 278 = Aster infirmus.
Cyparissias, *Wedd. Chlor. And.* i. 203.—N. Granat.
dichotomum, DC. Prod. v. 278 = Aster reticulatus.
eriophyllum, *Wedd. Chlor. And.* i. 206.—N. Granat.
extenuatum, Nees, Gen. et Sp. Aster. 195 = Aster fruticesus.
filifolium, Nees, l. c. 197 = Erigeron filifolius.
floribundum, *Wedd. Chlor. And.* i. 205.—N. Granat.
fruticulosum, Nees, Gen. et Sp. Aster. 194 = Aster fruticosus.
grandiflorum, Spreng. Syst. iii. 544 = Liabum grandiflorum.
Hazenkel, *Wedd. Chlor. And.* i. 208.—Peruv.
hispidum, Nees, ex DC. Prod. v. 278 = Chrysocoma villosa.
ignitarium, Spreng. Syst. iii. 544 = Liabum ignitarium.
lavandulaefolium, H. B. & K. Nov. Gen. et Sp. iv. 97 = Aster lavandulaceus.
Lechleri, *Wedd. Chlor. And.* i. 204.—Peruv.
lepidophyllum, Nees, Gen. et Sp. Aster. 190 = Olearia lepidophylla.
linarifolium, Nees, l. c. 199 = Aster linarifolius.
linifolium, Forbes, Hort. Woburn. 190 = Aster acris.
longipes, Cass. in Dict. Sc. Nat. lvi. 171 = Aster fruticosus.
lyratum, Nees, Gen. et Sp. Aster. 188 = Olearia stelluiata.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 584, nomen.—Bolivia.
melastomoides, Spreng. Syst. iii. 544 = Liabum melastomoides.
microphyllum, Nees, Gen. et Sp. Aster. 191 = Olearia ramulosa.
microphyllum, *Wedd. Chlor. And.* i. 201.—Venezuela.
Meyenii, *Wedd. l. c.*—Peruv.
moranense, Nees, Gen. et Sp. Aster. 198 = Aster moranensis.
nubigenum, Spreng. Syst. iii. 543 = Liabum nubigenum.
obovatum, DC. Prod. v. 278 = Aster retinolatus.
obtusatum, Nees, Gen. et Sp. Aster. 196 = Aster obtusatus.
ochraceum, Nees, l. c. 201 = Tetramolopium ochraceum.
phlogotrichaeum, Nees, l. c. 186 = Olearia stelluiata.
phyllicoides, Weddl. Chlor. And. i. 206 = Tetramolopium phylicoides.
ramulosum, Nees, Gen. et Sp. Aster. 192 = Olearia ramulosa.

DIPLOSTEPHIUM:—
rosmarinifolium, *Wedd. Chlor. And.* i. 202.—N. Granat.
rupestre, *Wedd. l. c.* 206.—N. Granat.
Schultzii, *Wedd. l. c.* 204.—N. Granat.
sessiliflorum, Spreng. Syst. iii. 543 = Liabum sessiliflorum.
sessiliflorum, *Wedd. Chlor. And.* i. 204.—N. Granat.
solidagineum, Spreng. Syst. iii. 544 = Liabum solidagineum.
spinulosum, *Wedd. Chlor. And.* i. 200.—Nog. Andin.
stellulatum, Nees, Gen. et Sp. Aster. 187 = Olearia stellulata.
teretifolium, Nees, l. c. 197 = Erigeron filifolium.
umbellatum, DC. Prod. v. 272 = Aster umbellatus.
verbasciifolium, Spreng. Syst. iii. 544 = Liabum verbasciifolium.
volubilis, Spreng. l. c. 543 = Liabum volubile.

DIPLOSTYLIS, Karst. & Triana, in Linnaea, xxviii. (1856) 483 = **Rochefortia**, Sw. (Boragin.).
fasciculata, Karst. & Triana, l. c.—N. Granat.

DIPLOSTYLIS, Sond. in Linnaea, xxiii. (1850) 112 = **Adenocalina**, Turcz. (Compos.).
angustifolia, Sond. l. c. = A. sessilifolia.
bupleuroides, Sond. l. c. 114 = A. pauciflora.
capfra, Sond. l. c. 115 = A. acuta.
serrata, Sond. l. c. 114 = A. serrata.

DIPLOTAENIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 308 = **Peucedanum**, Tourn. (Umbellif.).
cachrydiifolia, Boiss. l. c. 309.—Persia.

DIPLOTAX, Rafin. Sylva Tellur. 129 (1838) = **Cassia**, Tourn. (Legumin.).
arborescens, Rafin. l. c. = C. glauca ? C. Sophera.

DIPLOTAXIS, DC. Syst. ii. 628 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 84.
acris, Boiss. Fl. Orient. i. 289 = Sisbrei.
apula, *Tenore, Syll.* 226.—Italia.
auriculata, Dur. Expl. Sc. Algér. t. 76 = tenuifolia.
Borrelieri, *DC. Syst.* ii. 634.—Hispan.
biloba, C. Koch, in Linnaea, xx. (1847) 262 = Sisymbrium pannonicum.
brachycarpa, *Godr. in Mém. Acad. Stanislas*, Sér. III. [xvii.] 1859 (1854) 374.—Hab. ?
bracteata, Gren. & Godr. Fl. Fr. i. 81 = Brassica Erucastrum.
brassicaeformis, Koch, ex Boiss. Voy. Bot. Espagne, ii. 84 = Moricandia arvensis.
brassicoides, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 172 = prace.
brassicoides, Roxy, in Bull. Soc. Bot. Fr. xxix. (1882) 40 = Brassica Rouyana.
brevicaulis, Bluff & Fingerh. Comp. Fl. Germ. 158 = viminea.
catholica, *DC. Syst.* ii. 632.—Europ. austr.; Afr. bor.
catholica, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1877) 232 = silicia.
chilensis, *Barn. in C. Gay, Fl. Chil.* i. 132.—Chili.
cineros, *Pomel, Nouv. Mat. Fl. Atl.* 259.—Algeria.
coronopifolia, *Godr. in Mém. Acad. Stanislas*, Sér. III. [xvii.] 1858 (1854) 378.—Hab. ?
crassifolia, DC. Syst. ii. 629 = pendula.
Duveyriereana, *Coss. in Bull. Soc. Bot. Fr.* xi. (1864) 164.—Afr. bor.
Erucastrum, Gren. & Godr. Fl. Fr. i. 81 = Brassica Erucastrum.
ercoides, *DC. Syst.* ii. 631.—Reg. Mediterr. ; As. Min.
erucoides, Schimp. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 35 = Brassica Schimperi.
glabra, Dulac, Fl. Hautes-Pyr. 195 = viminea.
grandiflora, Wall. ex Kurz, Rep. Andam. ed II. 3, sphalm. = Chloroscordum grandiflorum.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 171.—Ind. bor. occ.; Afghan.
Harra, Boiss. Fl. Orient. i. 288 = pendula.
hirsuta, Bertol. Misc. Bot. ii. 14 = pendula.
hispida, DC. Syst. ii. 630 = pendula.
hispidula, *Tenore, Syll.* 225.—Italia.
humilis, Gren. & Godr. Fl. Fr. i. 78 = Brassica humilis.
intermedia, Schur, Enum. Pl. Transs. 60 = tenuifolia.
intricata, *Nym. Syll.* 197.—Hispan.
Lagascana, *DC. Syst.* ii. 629.—Hispan.
Lagascana, Willk. ex Willk. & Lange, Prod. Fl. Hisp. iii. 868 (= *Pendulina Webbiana*).—Hispan.
leucanthemifolia, Jord. Diagn. i. 190, in obs. = Brassica humilis.

DIPLOTAXIS:—
muralis, *DC. Syst.* ii. 684.—Europ.
nana, Boiss. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 3 = pendula.
pachypoda, *Godr. Fl. Fun.* ed. II. 64; *in Mém. Acad. Montp. (Sect. Médic.)* i. (1853) 414.—Hab.?
parvula, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 198.—Soongaria.
pendula, *DC. Syst.* ii. 680.—Reg. Mediterr.; Orient.
platystylis, *Pomel, Nouv. Mat. Fl. Atl.* 224.—Algeria.
platystylos, Willk. in Bot. Zeit. v. (1847) 333 = virgata.
Prolongi, Boiss. Elench. 11; Voy. Bot. Espagne, 33 = viminea.
ramentaristrma, Spreng. Neue Entdeck. iii. 59 = tenuifolia?
repanda, Gren. & Godr. Fl. Fr. i. 79 = Brassica repanda.
saxatilis, *DC. Syst.* ii. 686.—Europ. austr.
saxatilis, Ledeb. Fl. Ross. i. 220 = viminea.
scaposa, *DC. Syst.* ii. 685.—Ins. Lampedusa.
setosa, *DC. l. c.* 699.—Mont. Pyren.
Sieberi, *Presl, Bot. Bemerk.* 10.—Aegypt.; Arab.
silfolia, *Kunze, in Flora*, xxix. (1846) 685.—Hispan.; Marocc.
simplex, *Spreng. Syst.* ii. 914.—Afr. bor.
subcuneata, Jord. Diagn. i. 189 = Brassica humilis.
tenuifolia, *DC. Syst.* ii. 682.—Europ.; Reg. Mediterr.; As. Min.
tenuisiliqua, *Delile, Ind. Sem. Hort. Monsp.* (1847) 7.—Afr. bor.
varicabilis, Rochel, Pl. Banat. Rar. Ind. Spec. = muralis, tenuifolia.
vernalis, *Phil. in Anal. Univ. Chil.* (1872) 670.—Chili.
viminea, *DC. Syst.* ii. 685.—Reg. Mediterr.; As. Min.
virgata, *DC. l. c.* 681.—Hispan.; Afr. bor.

DIPLOTER, Rafin. Sylva Tellur. 165 (1838) = Tetracera, Linn. (Dillen.).
ainifolia, Rafin. l. c. = T. sinifolia.

DIPLOTHECA, Hochst. in Flora, xxix. (1846) 595 = Astragalus, Toem. (Legumin.).
abyssinica, Hochst. l. c. = A. abyssinicus.
atropilosula, Hochst. l. c. 596 = A. venosus.
tigrensis, Hochst. l. c. = A. venosus.
venosa, Hochst. l. c. = A. venosus.

DIPLOTHEMIUM, Mart. Hist. Nat. Palm. ii. 107 (1824). *PALMÆ*, Benth. & Hook. f. iii. 945.
ALLAGOPTERA, Nees, in Nesswied, Reise nach Bras. ii. 335 (1821).
campestre, *Mart. Hist. Nat. Palm.* ii. 109. t. 76. f. 1, 2.—Bras.
caudescens, *Mart. l. c.* iii. t. 70, 77. fig. 1, 2.—Bras.
leucocalyx, *Drude, in Mart. Fl. Bras.* iii. ii. 481.—Bras.
littorale, *Mart. Hist. Nat. Palm.* ii. 110. t. 76. f. 5.—Bras.
maritimum, *Mart. l. c.* 108. t. 76. 77. f. 8.—Bras.
Torrellii, Mart. in Palmet. Orb. 105. t. 15. f. 3 = Jubæa Torellii.

DIPLOTHENIUM, Voigt, in Syll. Ratisb. ii. (1828) 51 = præc.

DIPLOTHRIA, Walp. Rep. ii. 620 (1848) = seq.

DIPLOTHRIX, DC. Prod. v. 611 (1836) = Zinnia Linn. (Compos.).
acerosa, DC. l. c. = Z. acerosa.
juniperifolia, DC. l. c. 612 = Z. juniperifolia.

DIPLOTROPIS, Benth. in Ann. Wien. Mus. ii. (1838) 88. *LEGUMINOSÆ*, Benth. & Hook. f. i. 557.
DIBRACHION, Tul. in Ann. Sc. Nat. Sér. III. xx. (1848) 189.
brachypetalum, *Tul. in Arch. Mus. Par.* iv. (1844) iii. —Guiana.
brasiliensis, *Benth. in Mart. Fl. Bras.* xv. i. 290.—Bras. ferrugines, *Benth. l. c.* 291.—Bras.
grandiflora, *Tul. in Arch. Mus. Par.* iv. (1844) 109.—Bras.
Martini, Benth. in Hook. Journ. Bot. ii. (1841) 72, in obs. sphalm. = seq.
Martiusi, *Benth. in Ann. Wien. Mus.* ii. (1838) 88.—Bras.
nitida, *Benth. in Hook. Journ. Bot.* ii. (1841) 71.—Guiana.

DIPLOUSODON, Meisn. Gen. 88 (1838) = Diplusodon, Pohl (Lythrar.).

DIPLUKION, Rafin. Sylva Tellur. 83 (1838) = Iochroma, Benth. (1845) (Solanac.).
cornifolium, Rafin. l. c. = I. coralloïdis.
lonicum, Rafin. l. c. = I. loxensis.
umbrosum, Rafin. l. c. = I. umbrosa.

DIPLUSION, Rafin. Alsog. Am. 15 (1838) = Salix, Tourn.
cordata, Rafin. l. c. = S. cordata.
lucida, Rafin. l. c. = S. lucida.
rigida, Rafin. l. c. = S. cordata.
undulata, Rafin. l. c. = S. triandra?

DIPLUSODON, Pohl, in Flora, x. (1827) 150. *LYTHRARIEÆ*, Benth. & Hook. f. i. 781.
DIPLODON, DC. Prod. iii. 94 (a) (1828).
DUBYÆA, DC. l. c. in syn. (1828).
FRIEDLANDIA, Cham. & Schlecht. in Linnæa, ii. (1827) 848.
alutaceus, Pohl, in Flora, x. (1827) 152 = lanceolatus.
alutaceus, St. Hil. Fl. Bras. Mer. iii. 155 = helianthemifolius.
amœnus, DC. Prod. iii. 94 (b) = virgatus.
angustifolius, DC. l. c. 94 (a) = hexander.
arborescens, Poepp. & Endl. Nov. Gen. et Sp. ii. 66 = Physocalymma scaberrimum.
buxifolius, *DC. Prod.* iii. 94 (b).—Bras.
Candollei, *Pohl, ex DC. l. c.* 94 (a).—Bras.
Candollei, St. Hil. Fl. Bras. Mer. iii. 146 = hexander, uninervius.
capitatus, *St. Hil. l. c.* 149.—Bras.
decussatus, *Gardn. & Field. Sert. Pl.* t. 1.—Bras.
divaricatus, *Pohl, in Flora*, x. (1827) 150.—Bras.
epilobioides, *DC. Prod.* iii. 94 (a).—Bras.
floribundus, *Pohl, in Flora*, x. (1827) 151.—Bras.
ginnvioides, Griseb. Cat. Pl. Cub. 106 = Ginora Diplusodon.
glaucescens, *DC. Prod.* iii. 94 (c).—Bras.
gracilis, *Koehne, in Mart. Fl. Bras.* xiii. ii. 323.—Bras.
heliantheniifolius, *DC. Prod.* iii. 94 (c).—Bras.
hexander, *DC. l. c.* 94 (a).—Bras.
hirsutus, *DC. l. c.* 94 (c).—Bras.
hirtellus, *G. Don, Gen. Syst.* ii. 719.—Bras.
imbricatus, *Pohl, in Flora*, x. (1827) 150.—Bras.
incanus, *Gardn. & Field. Sert. Pl.* t. 2.—Bras.
kielmeyeroides, *St. Hil. Fl. Bras. Mer.* iii. 151.—Bras.
lanceolatus, *Pohl, in Flora*, x. (1827) 152; *Pl. Bras.* i. 98. t. 51.—Bras.
lanceolatus, St. Hil. Fl. Bras. Mer. iii. 150 = oblongus.
longipes, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 404.—Bras.
lythroides, *DC. Prod.* iii. 94 (e).—Bras.
macrodon, *Koehne, in Mart. Fl. Bras.* xiii. ii. 338.—Bras.
marginatus, *Pohl, in Flora*, x. (1827) 150.—Bras.
microphyllus, *Pohl, l. c.* 151.—Bras.
montanus, Casar. ex Koehne, in Mart. Fl. Bras. xiii. ii. 327 = hirsutus.
Myrsinites, *DC. Prod.* iii. 94 (b).—Bras.
nitidus, *DC. l. c.* 94 (d).—Bras.
oblongus, *Pohl, in Flora*, x. (1827) 152.—Bras.
orbicularis, *Koehne, in Mart. Fl. Bras.* xiii. ii. 329.—Bras.
ovatus, Mart. ex Koehne, l. c. 319 = speciosus.
ovatus, *Pohl, in Flora*, x. (1827) 151.—Bras.
parvifolius, *DC. Prod.* iii. 94 (c).—Bras.
pauciflorus, DC. l. c. = heliantheniifolius.
paberulus, *Koehne, in Mart. Fl. Bras.* xiii. ii. 339.—Bras.
punctatus, *Pohl, in Flora*, x. (1827) 151.—Bras.
quinquelocularis, *Koehne, in Mart. Fl. Bras.* xiii. ii. 316.—Bras.
ramosissimus, *Pohl, in Flora*, x. (1827) 151.—Bras.
resolutifolius, DC. ex Koehne, in Mart. Fl. Bras. xiii. ii. 326 = parvifolius.
rosmarinifolius, *St. Hil. Fl. Bras. Mer.* iii. 147.—Bras.
rotundifolius, *DC. Prod.* iii. 94 (f).—Bras.
scaber, Pohl, in Flora, x. (1827) 152 = lanceolatus.
serpyllifolius, *DC. Prod.* iii. 94 (c).—Bras.
sessilifolius, *Koehne, in Mart. Fl. Bras.* xiii. ii. 333.—Bras.
speciosus, *DC. Prod.* iii. 94 (f).—Bras.
stachyoides, DC. l. c. 94 (e) = villosissimus.

DIPLUSODON:—
strigosus, *Pohl, in Flora*, x. (1827) 151.—Bras.
suberiosus, *Casar. ex Koehne, in Mart. Fl. Bras.* xiii. ii. 329.—Bras.
thymifolius, *DC. Prod.* iii. 94 (b).—Bras.
uninervius, *Koehne, in Mart. Fl. Bras.* xiii. ii. 340.—Bras.
vaccinifolius, DC. Prod. iii. 94 (b) = buxifolius.
villosissimus, *Pohl, in Flora*, x. (1827) 151.—Bras.
villosus, *Pohl, l. c.*—Bras.
virgatus, Gardn. ex Koehne, in Mart. Bras. xiii. ii. 336, 338 = macrodon, incanus.
virgatus, *Pohl, in Flora*, x. (1827) 151.—Bras.

DIPLYCOSIA, Blume, Bijdr. 857 (1826). *ERICACEÆ*, Benth. & Hook. f. ii. 583.
AMPHICALYX, Blume, Fl. Jav. Fragm. p. vii (1828).
acuminata, *Becc. Malesia*, i. (1878) 212.—Borneo.
ambotensis, *Becc. l. c.* 210.—Ins. Amboina.
ciliolata, *Hook. f. in Pl. t.* 894.—Borneo.
consobrina, *Becc. Malesia*. i. (1878) 211.—Borneo.
discolor, *G. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 459.—Reg. Himal.
heterophylla, *Blume, Bijdr.* 858.—Java.
latifolia, *Blume, l. c.*—Java.
macrophylla, *Becc. Malesia*, i. (1878) 212.—Borneo.
microphylla, *Becc. l. c.*—Borneo.
pilosa, *Blume, Bijdr.* 858.—Java.
scabrida, *Becc. Malesia*, i. (1878) 211.—Borneo.
semi-infera, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 459.—Reg. Himal.
Soror, *Becc. Malesia*, i. (1878) 210.—N. Guinea.

DIPODIUM, R. Br. Prod. 330 (1810). *ORCHIDEÆ*, Benth. & Hook. f. iii. 589.
LEOPARDANTHUS, Blume, Rumphia, iv. 47 (1848); et Mus. Bot. Lugd. Bat. i. 47. t. 15 (1849).
WAILESIA, Lindl. in journ. Hort. Soc. iv. (1849) 261.
cruciflorum, *F. Muell. Fragm.* v. 49.—Austral.
fuscum, Buch.-Ham. ex Wall. Cat. n. 7364 (= *Cyrtopera flava*.)—Ind. or.
Hamiltonianum, *F. M. Bailey, in Proc. Linn. Soc N. S. Wales*, vi. (1881) 140.—Austral.
khasyanum, Griff. Notul. iii. 554 (= *Sunipia bicolor*) —Ind. or.
paludosum, *Reichb. f. Xenia Orch.* ii. 15.—Malaya.
pictum, *Reichb. f. l. c.*—Malaya.
picatum, Buch.-Ham. ex Wall. Cat. n. 7369 = Eulophia nuda.
punctatum, *R. Br. Prod.* 331.—Austral.
ramentaceum, Buch.-Ham. ex Wall. Cat. n. 7370 = Eulophia campestris.
Roxiatæ, Buch.-Ham. l. c. n. 7371 = Eulophia nuda.
squamatum, *R. Br. Prod.* 331.—Ins. N. Hebrid.
venosum, F. Muell. Fragm. i. 61 = Eulophia venosa.

DIPOGON, Liebm. Ind. Sem. Hort. Hafn. (1854) 27, ex Ann. Sc. Nat. Sér. IV. ii. (1854) 374 = Dolichos, Linn. (Legumin.).
glycinoides, Liebm. l. c.—Bras.

DIPOGON, Willd. ex Steud. Nom. ed. II. i. 518 (1840) = Chrysopogon, Trin. (Gramin.).
avenaceus, Willd. ex Steud. l. c. = C. avenaceus.

DIPOGONIA, Beauv. Agrost. 125 (1812) = Diplopogon, R. Br. (Gramin.).
setacea, Beauv. l. c. = Diplopogon setaceus.

DIPORIDIUM, Wendl. f. in Bertl. & Wendl. f. Beitr. ii. 24 (1825) = Oehnea, Linn.
arborescens, Wendl. f. l. c. 26 = O. arborea.
atropurpureum, Wendl. l. c. 24 = O. atropurpurea.
dalapoënse, Eckl. & Zeyh. Enum. 118 = O. arborea.
integrifolium, Presl, Bot. Bemerk. 82 = O. arborea?
metallitinum, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 56 = O. atropurpurea.
pulchrum, Walp. Rep. iv. 378 = O. pulchra.
serrulatum, Hochst. in Flora, xxvii. (1844) 304 = O. atropurpurea.

DIPOSIS, DC. Coll. Mém. v. 93. t. 3 (1829); Prod. iv. 61 (1830). *UMBELLIFERÆ*, Benth. & Hook. f. i. 877.
Balbocastanum, *DC. Prod.* iv. 66.—Chili.
canticinclia, *DC. Coll. Mém.* v. 93.—Reg. August.

DIPSACELLA, Opiz, Seznam, 38 (1859) = **Dipsacus**, Linn.
setigera, Opiz, l. c. = Dipsacus pilosus.

DIPSACOZAMIA, Lehm. in Lindl. Veg. Kingd. 225 (1847) = **Ceratozamia**, Brongn. (Cycad.).
mexicana, Liebm. ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 193 = C. mexicana.

DIPSACUS, Linn. Syst. ed. I (1735). *DIPSACEAE*, Benth. & Hook. f. ii. 168.
DIPSACELLA, Opiz, Seznam, 38 (1859).
GALEDRAGON, S. F. Gray, Nat. Arr. Brit. Pl. ii. 475 (1821).
VIRGA, Hill, Veg. Syst. v. 19 (1763).
appendiculatus, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 367.—Abyss.
asper, *Wall. Cat.* n. 4268.—Reg. Himal.
atratus, *Hook. f. & Thomson, ex C. B. Clarke, in Fl. Brit. Ind.* iii. 218.—Reg. Himal.
auriculatus, Royle, Illustr. Bot. Himal. 345 = strictus.
azureus, *Schrenk, Enum. Pl. Nov.* 53.—Sibir.
Bottori, Maly, ex Nym. Consp. 345 = sylvestris.
carminatorius, Salisb. Prod. 55 = fullonum.
comosus, Hoffingg. & Link, Fl. Port. ii. 81 = laciniatus.
connatifolius, Gilib. Fl. Lituan. i. 106 = fullonum.
cuneifolius, *Rafin. New Fl. Am.* ii. 19.—Am. bor.
divaricatus, *Y. & C. Presl, Delic. Prag.* 117.—Sicilia.
elongatus, Salisb. Prod. 55 = pilosus.
ferox, *Loisel. Fl. Gall.* 719. t. 8.—Ins. Corsica.
fullonum, *Linn. Sp. Pl.* 97.—Europ.; Reg. Caucas.
fullonum, S. G. Gmel. Reise, i. 158 = laciniatus.
fullonum, Thore, Ess. Chlor. 36 = sylvestris.
Gmelini, Bieb. Fl. Taur. Cauc. i. 93 = strigosus.
Gmelini, Maxim. Prim. Fl. Amur. 472 = japonicus.
horridus, Opiz, Seznam, 38 = sylvestris.
inermis, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 367.—Reg. Himal.
japonicus, *Miq. in Verzl. en Med. Ahad. Wetensch.* ii. 11. (1868) 83.—China; Japon.
laciniatus, Falk, Beitr. ii. 111 = strigosus.
laciniatus, *Linn. Sp. Pl.* 97.—Europ.; Orient.
lacuiatus, Ucria, Hort. Panorm. 6 = divaricatus.
Leschenaultii, *Coult. in DC. Prod.* iv. 647.—Ind. or.
Leschenaultii, Thw. Enum. Pl. Zeyl. 160 = Walkeri.
longicaulis, Wall. Cat. n. 4256 = inermis.
microcephalus, Martr. Pl. Crit. Tarn, 25 = laciniatus.
minor, Neck. Delic. Gallo-Belg. i. 81 = pilosus.
mitis, D. Don, Prod. Fl. Nep. 161 = inermis.
Morisonii, Bor. Ind. Sem. Hort. Anger. (1842) = sylvestris.
Orsini, Sanguinetti, in Atti Acc. Pont. Lincei, Ser. I. v. (1852) 725. t. 1 = sylvestris.
palustris, Salisb. Prod. 55 = fullonum.
pilosus, *Linn. Sp. Pl.* 97.—Europ.; Reg. Caucas.
pinnatifidus, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 367.—Abyss.
plumosus, C. Koch, in Linnæa, xxiv. (1851) 442 = Scabiosa plumosa.
Pseudo-sylvestris, Schur, Enum. Pl. Transs. 298 = laciniatus.
Roylei, Klotzsch, in Bot. Ergeb. Waldem. Reise, 33 = inermis.
sativus, [Honck.] Vers. Aller Gew. Tentschl. 374; C. C. Gmel. Fl. Bad. i. 314 = fullonum.
Schimperi, A. Br. ex Schweinf. & Ascherson in Beitr. Fl. Aethiop. 287 = pinnatifidus.
setosus, *Hierm. in Oliver, Fl. Trop. Afr.* iii. 250.—Afr. trop.
silvester, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 359 = sylvestris.
sinuatus, Schlecht. ex Roem. & Schult. Syst. iii. 519 = sylvestris.
strictus, *D. Don, Prod. Fl. Nep.* 160.—Reg. Himal.
strigosus, *Willd. ex Roem. & Schult. Syst.* iii. 590.—As. Min.; Persia.
sylvestris, Lam. Fl. Fr. iii. 345 = fullonum.
sylvestris, *Mill. Gard. Dict. ed.* VIII. n. 1.—Europ.; Orient.
vulgaris, C. C. Gmel. Fl. Bad. i. 313 = sylvestris.
Wallichi, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 344.—Zeylan.

DIPSEUDOCHORION, Buchen. in Flora, xlviii. (1865) 345 = **Edmaphyton**, Mig. (Allenac.).
capitellatum, Buchen. l. c. = L. obtusifolium.

DIPTANTHERA, Schrank, ex Steud. Nom. ed. II. i. 518 (1840) = **Dianthera**, Gronov. (Acanthac.).

DIPTERA, Borckh. in Roem. N. Mag. i. (1794) 29 = **Saxifraga**, Linn.
cuscutaeformis, Heynh. Nom. ii. 206 = S. cuscutaeformis.
sarmentosa, Borckh. in Roem. N. Mag. i. (1794) 29 = S. sarmentosa.

DIPTERACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 75 (1832) = **Ruellia**, Linn. (Acanth.).
acaulis, Presl, Bot. Bemerk. 98.—Chili.
affinis, Nees, in Mart. Fl. Bras. ix. 80 = R. speciosa.
angustus, Griseb. in Bonplandia, vi. (1858) 10.—Panama.
australis, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 129 = R. australis.
bahiensis, Nees, in Mart. Fl. Bras. ix. 89 = R. trivialis.
Beyrichianus, Nees, l. c. 83.—Bras.
biflorus, Nees, in Linnæa, xvi. (1842) 294 = Calophanes oblongifolia.
Blanchetianus, Moric. Pl. Nouv. Am. 168.—Bras.
brachysiphon, Nees, in Mart. Fl. Bras. ix. 84 = R. brachysiphon.
bracteatus, Nees, in DC. Prod. xi. 148 = R. bracteata.
brasiliensis, Presl, Bot. Bemerk. 98 = Hygrophila brasiliensis.
brevicaulis, Nees, in Mart. Fl. Bras. ix 85.—Bras.
calvescens, Nees, l. c. 92 = R. solitaria.
calycinus, Champ. in Hook. Kew Journ. v. (1853) 138 = Strobilanthes flaccuiflorus.
calycosus, Nees, in Mart. Fl. Bras. ix. 28.—Bras.
canescens, Nees, in Hook. Lond. Journ. Bot. iv. (1845) 633 = R. geminiflora.
canus, Nees, in Mart. Fl. Bras. ix. 88.—Bras.
capreaefolius, Nees, l. c. 42.—Bras.
Chamaedrys, Nees, l. c. 81.—Bras.
ciliatus, Nees, in Wall. Cat. n. 7151 = Hygrophila phlomoides, Ruellia ciliata.
ciliosus, Nees, in Linnæa, xvi. (1842) 294 = R. ciliosa.
clandestinus, Presl, Bot. Bemerk. 94, 98 = R. tuberosa.
confinis, Nees, in Mart. Fl. Bras. ix. 81.—Bras.
cordifolius, Nees, in Linnæa, xv. (1841) 854 = R. ovata.
cyaneus, Nees, in DC. Prod. xi. 121 = R. cyanea.
dejectus, Nees, in Wall. Pl. As. Rar. iii. 82 = R. prostrata.
diffusus, Nees, in DC. Prod. xi. 124 = R. diffusa.
dissitiflorus, Nees, in Mart. Fl. Bras. ix. 83 = R. dissitiflola.
domingensis, Nees, in DC. Prod. xi. 135 = R. domingensis.
Drummondianus, Nees, l. c. 123 = R. Drummondiana.
Drummondii, Torr. & Gray, ex Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 258 = R. ciliosa.
dubius, Nees, in DC. Prod. xi. 128.—Hab.?
elongatus, Nees, in DC. l. c. 140 = Whitfieldia longifolia.
erectus, Nees, in Wall. Pl. As. Rar. iii. 82 = R. patula.
floribundus, Nees, in DC. Prod. xi. 141 = R. floribunda.
foetidus, Nees, in Benth. Bot. Voy. Sulph. 146 = R. rubicaulis.
furcatus, Nees, in Mart. Fl. Bras. ix. 85.—Bras.
Gardneri, Nees, l. c. 28.—Bras.
geminiflorus, Nees, l. c. 40 = R. geminiflora.
gendarussa, Schwaegr. in Verh. Zool. Bot. Ges. Wien, xviii. (1868) 680.—Nubia.
glanduloso-punctatus, Nees, in Mart. Fl. Bras. ix. 84.—Bras.
grandiflorus, Nees, in DC. Prod. xi. 134 = R. candida.
guttatus, Nees, l. c. 140 = R. guttata.
Haenkei, Nees, l. c. 794 = R. albicaulis.
Helianthus, Nees, in Mart. Fl. Bras. ix. 86.—Bras.
Herbstii, Hook. Bot. Mag. t. 5156 = R. Herbstii.
hirsutus, Vell. Fl. Flum. vi. t. 96 = R. geminiflora.
hispidus, Bertol. Misc. Bot. xvii. 17 = Gratiola pilosa.
Hookerianus, Nees, in DC. Prod. xi. 130 = R. Hookeriana.
humifusus, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 128 = R. humifusa, *Hemsl.*
humilis, Nees, in Mart. Fl. Bras. ix. 29 = R. humilis.
humistratus, Champ. Fl. S. U. St. 303 = Calophanes humistrata.

DIPTERACANTHUS:—
hygrophiloides, Nees, in DC. Prod. xi. 121 = Ruellia hygrophiloides.
hypericoides, Nees, in Mart. Fl. Bras. ix. 39.—Bras.
inaequifolius, Nees, l. c. 48.—Bras.
incomtus, Nees, l. c. 44.—Bras.
interruptus, Presl, Bot. Bemerk. 97.—Bras.
jamaicensis, Nees, in DC. Prod. xi. 140 = Blechum laxiflorum.
lacteus, Presl, Bot. Bemerk. 98 = R. lactea.
lamiifrondis, Nees, in Mart. Fl. Bras. ix. 43.—Bras.
lanato-glandulosus, Nees, in DC. Prod. xi. 130.—Bras.
lanato-viscosus, Nees, l. c. = R. lanato-viscosa.
lanceolatus, Nees, in Wall. Pl. As. Rar. iii. 82 = R. reepens.
leucanthus, Nees, in Benth. Pl. Hartw. 236.—Bras.
Lindheimerianus, Scheele, in Linnæa, xxi. (1848) 764 = R. Drummondiana.
linearis, Champ. Fl. S. U. St. 303 = Calophanes angustus.
linearis, Torr. & Gray, ex Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 258 = Calophanes linearis.
longepetiolatus, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 124 = R. longepetiolata.
longiflorus, Nees, in DC. Prod. xi. 117 = R. longiflora.
longifolius, Stocks, in Hook. Kew Journ. iv. (1852) 177 = R. longifolia.
macranthus, Nees, in Mart. Fl. Bras. ix. 37 = R. macrantha.
macrophyllus, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 125.—Mexic.
matutinus, Presl, Bot. Bemerk. 94 = R. patula.
menthoides, Nees, in Mart. Fl. Bras. ix. 44 = R. menthoides.
microanthus, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 257 = R. strepens.
microcalyx, Nees, in DC. Prod. xi. 116.—Guiana.
microphyllus, Nees, l. c. 143 = Calophanes microphyllus?
Mitchilianus, Nees, l. c. 128 = R. ciliosa.
monanthus, Nees, l. c. 125 = R. monanthos.
multifolius, Nees, in Mart. Fl. Bras. ix. 38 = R. subcaulescens.
Nagchana, Nees, in Wall. Pl. As. Rar. iii. 82 = Calophanes Nagchana.
nanus, Nees, in DC. Prod. xi. 127 = R. geminiflora.
Neesianus, Mart. ex Nees, in Mart. Fl. Bras. ix. 87.—Bras.
Neowedia, Nees, l. c. 30.—Bras.
nitens, Nees, l. c. 41.—Bras.
nocticolorus, Nees, in DC. Prod. xi. 123 = R. ciliosa, nocticlora.
nudiflorus, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 229 = R. tuberosa.
oblongifolius, Champ. Fl. S. U. St. 303 = Calophanes oblongifolius.
ocymoides, Presl, Bot. Bemerk. 98 = R. patula.
pallidus, Nees, in DC. Prod. xi. 126 = R. pallida.
paniculatus, Nees, l. c. 142 = R. paniculata.
parviflorus, Nees, l. c. 128 = Hemigraphis elegans?
patulus, Nees, in Wall. Pl. As. Rar. iii. 82 = R. patula.
Pavonianus, Nees, in DC. Prod. xi. 133 = R. biflora.
pilosus, Nees, l. c. 127 = R. pilosa, *Pav.*
pilosus, Nees, in Linnæa, xv. (1841) 853 = R. pilosa, *Linn.*
Pohlii, Nees, in Mart. Fl. Bras. ix. 49.—Bras.
porrigens, Nees, l. c. 41 = R. porrigens.
procumbens, Nees, in DC. Prod. xi. 138 = R. Schlechtendaliana.
prostratus, Nees, in Wall. Pl. As. Rar. iii. 81 = R. prostrata.
Pumilio, Nees, in DC. Prod. xi. 124 = R. australis.
Purdiaeanus, Nees, l. c. 794 = R. Purdieana.
Puri, Nees, in Mart. Fl. Bras. ix. 88 = R. Puri.
racemosus, Presl, Bot. Bemerk. 98.—Bras.
repens, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 129 = R. repens.
riparius, Champ. Fl. S. U. St. 303 = Calophanes humistratus.
rosmarinus, Nees, in DC. Prod. xi. 139.—Bras.
rubiaeformis, Nees, in Mart. Fl. Bras. ix. 32.—Bras.
rubicaulis, Nees, in DC. Prod. xi. 116 = R. rubicaulis.
scandens, Glendinn. ex Lindl. Bot. Reg. (1847) sub t. 28, 31 = Henfreya scandens.
Schauerianus, Nees, in Linnæa, xvi. (1842) 290 = R. solitaria.
Schlechtendalianus, Nees, in DC. Prod. xi. 138 = R. Schlechtendaliana.
Schottianus, Nees, in Mart. Fl. Bras. ix. 82.—Bras.

DIPTERACANTHUS:—
serpyllifolius, Nees, in DC. Prod. xi. 125.—Madag.
sessilifolius, Nees, in Mart. Fl. Bras. iv. 98.—Bras.
Sibsa, Nees, in Wall. Pl. As. Rar. iii. 81 = R. suffruticosa.
speciosus, Presl, Bot. Bemerk. 98 = R. macrantha.
spectabilis, Hook. Bot. Mag. t. 4494.—Peruv.
squarrosus, Fenzl, Ind. Sem. Hort. Vindob. (1868) 10. —Am. trop.
strepens, Presl, Bot. Bemerk. 97; Nees, in DC. Prod. xi. 121 = R. strepens.
subcapitatus, Nees, l. c. 728 = R. subcapitata.
subdenticulatus, Miq. in Journ. Bot. Néerl. i. (1861) 117.—China.
subringens, Nees, in Mart. Fl. Bras. ix. 31.—Bras.
sudanicus, Schweinf. in Verh. Zool.-Bot. Ges. Wien, xviii. (1868) 679.—Afr. trop.
suffruticosus, Torr. in Bot. Mex. Bound. 122 = R. Parryi.
suffruticosus, Voigt, Hort. Suburb. Calc. 483 = R. suffruticosa.
surinamensis, Miq. in Linnæa, xxii. (1849) 471.—Guiana.
tomentosus, Nees, in Mart. Fl. Bras. ix. 35.—Bras.
tuberosus, Nees, l. c. 42.—Bras.
tubiflorus, Nees, in DC. Prod. xi. 128 = R. tubiflora.
tubiflorus, Seem. Bot. Voy. Herald, 178 = R. densiflora.
turbacensis, Nees, in DC. Prod. xi. 148.—N. Granat.
urophyllus, Nees, in Wall. Pl. As. Rar. iii. 82 = Strobilanthes uropbyllus.
vagans, Benth. ex T. Anders. in Journ. Linn. Soc. ix. (1867) 459 = Calophanes vagans.
ventricosus, Nees, in DC. Prod. xi. 126.—Java.
vincæformis, Nees, in Mart. Fl. Bras. ix. 36 = R. vincoides.
Vindex, Nees, l. c. 42 = R. Vindex.
viscidulus, Nees, in Ind. Sem. Hort. Vrat. (1838) et (1840).—Cf. Linnæa, xvi. (1842) 391 = R. viscidula.
viscidus, Nees, in DC. Prod. xi. 141.—Peruv.
viscosissimus, Nees, in Mart. Fl. Bras. ix. 34.—Bras.
Zeyheri, Sond. in Linnæa, xxiii. (1850) 90 = R. Zeyheri.

DIPTERANTHEMUM, F. Muell. in Wing. South Sc. Record, iii. (1884) 281. *AMARANTACEÆ.* Crosslandi, *F. Muell. l. c.*—Austral.

DIPTERANTHUS, Rodrig. Orch. Nov. ii. 282 (1881). *ORCHIDACEÆ.*
pseudobulbiferus, *Rodrig. l. c.* 283.—Bras.

DIPTERIX, Willd. Sp. Pl. iii. 910 (1801) = **Dipteryx**, Schreb. (Legumin.).

DIPTEROCALYX, Cham. in Linnæa, vii. (1832) 241 = **Lippia**, Linn. (Verben.).
glabrescens, Cham. l. c. 242 = L. hirta.
hirtus, Cham. l. c. 241 = L. hirta.
scaberrimus, Schlecht. in Linnæa, xxvi. (1853–55) 647.—N. Granat.

DIPTEROCARPUS, Gaertn. f. Fruct. iii. 50, t. 187, 188 (1805). *DIPTEROCARPEÆ*, Benth. & Hook. f. i. 191.
Oleoxylon, Wall. Cat. p. 157 n. 953 (1831 ?).
Perigium, Correa, in Ann. Mus. Par. viii. (1806) 397.
acuteangulus, Vesque, in Compt.-Rend. Par. lxxviii. (1874) 628 = appendiculatus.
alatus, A. DC. Prod. xvi. ii. 611 = scaber.
alatus, *Roxb. Hort. Beng.* 49; *Fl. Ind.* ii. 614.—Burma.
angulatus, Dyer, in Journ. Bot. xii. (1874) 104 = geniculatus.
angustifolius, *Wight & Arn. Prod.* 84.—Ind. or.
appendiculatus, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 847.—Ins. Banca.
balsamifer, *Blume, Mus. Bot. Lugd. Bat.* ii. 37.—Borneo.
Bandii, Korth. Verh. Nat. Gesch. Nederl. Bot. 59. t. 5 = pilosus.
Beccarianus, Vesque, in Compt.-Rend. Par. lxxviii. (1874) 627 = globosus
Beccarii, Dyer, in Journ. Bot. xii. (1874) 103 = globosus.
Blancoi, Blume, Mus. Bot. Lugd. Bat. ii. 36 = grandiflorus.
camelliatus, Proc. Linn. Soc. (1873–74) 6, sphalm. = lamellatus.
cordatus, Wall. Cat. n. 956 = tuberculatus.
cornutus, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 296.—Malacca.

DIPTEROCARPUS:—
costatus, Gaertn. f. Fruct. iii. 50. t. 187 = alatus.
costatus, Roxb. Hort. Beng. 43; Fl. Ind. ii. 618 = angustifolius.
crinitus, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 296.—Malaya.
Dryobalanoides, Steud. Nom. ed. II. i. 518 = Dryobalanops aromatica.
elongatus, *Korth. Verh. Nat. Gesch. Nederl. Bot.* 62.—Borneo.
eurhynchus, *Miq. Fl. Ind. Bat. Suppl.* 485.—Borneo.
eurynchioides, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 846.—Ins. Banca.
fagiuus, *Vesque, in Compt.-Rend. Par.* lxxviii. (1874) 625.—Borneo.
fulvus, *Blume, Mus. Bot. Lugd. Bat.* ii. 37.—Ins. Philipp.
geniculatus, *Vesque, in Compt.-Rend. Par.* lxxviii. (1874) 626.—Borneo.
glandulosus, *Thw. Enum. Pl. Zeyl.* 34.—Zeylan.
globosus, *Vesque, in Compt.-Rend. Par.* lxxviii. (1874) 627.—Borneo.
gonopterus, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 376 = alatus.
gracilis, *Blume, Bijdr.* 294.—Java.
grandiflorus, *Blanco, Fl. Filip.* ed. II. 814.—Ins. Philipp.
grandiflorus, Griff. Notul. iv. 515 = Griffithii.
grandiflorus, Wall. Cat. n. 957 = tuberculatus.
grandifolius, *Miq. in Ann. Mus. Lugd. Bat.* i. 214.—Siam.
grandifolius, Teysm. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 214 = tuberculatus.
Griffithii, *Miq. l. c.* 213.—Burma.
Guiso, Blanco, Fl. Filip. ed. II. 813 = Anisoptera Guiso.
Hasseltii, Blume, Fl. Jav. Dipter. 22. t. 6 = trinervis.
hirtus, Vesque, in Compt.-Rend. Par. lxxviii. (1874) 627 = crinitus.
hispidus, *Thw. Enum. Pl. Zeyl.* 33.—Zeylan.
hispidus, *F. Villar*, in Blanco, Fl. Philipp. ed. III. Nov. App. 20 = verrucifluus.
incanus, *Roxb. Hort. Beng.* 42; *Fl. Ind.* ii. 614.—Burma.
indicus, Bedd. in Conserv. For. Rep. (1864–65) 17 = turbinatus.
insignis, *Thw. Enum. Pl. Zeyl.* 34.—Zeylan.
insularis, *Hance, in Journ. Bot.* xiv. (1876) 241.—China.
intricatus, *Dyer, l. c.* xii. (1874) 105.—Borneo.
laevis, Buch.-Ham. in Trans. Wern. Soc. vi. (1829) 399 = turbinatus.
lamellatus, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 159.—Borneo.
lampongus, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 847.—Sumatra.
Lemeslei, *Vesque, in Compt.-Rend. Par.* lxxviii. (1874) 626.—Ins. Palo-Condor.
littoralis, *Blume, Bijdr.* 294.—Java.
Lowii, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 160. —Borneo.
macrocarpus, Vesque, in Compt.-Rend. Par. lxxviii. (1874) 627 = pilosus.
Malaanonan, Blanco, Fl. Filip. ed. II. 812 = Shorea Malaanonan.
Mangahas, Blanco, l. c. 813 = Shorea Mangachapoi.
marginatus, *Korth. Verh. Nat. Gesch. Nederl. Bot.* 64.—Borneo.
Mayapis, Blanco, Fl. Filip. ed. II. 813 = turbinatus.
microcarpus, Zipp. ex Miq. Fl. Ind. Bat. i. ii. 501 = Anisoptera polyandra.
Mottleyanus, Hook. f. in Trans. Linn. Soc. xxiii. (1862) 159 = grandiflorus.
nobilis, Dyer, in Journ. Bot. xii. (1874) 106 = stellatus.
nudus, *Vesque, in Compt.-Rend. Par.* lxxviii. (1874) 626.—Borneo.
obtusifolius, *Blume, Mus. Bot. Lugd. Bat.* ii. 36.—Borneo.
oblongifolius, Thw. Enum. Pl. Zeyl. 33 = hispidus.
oblongus, A. DC. in. DC. Prod. xvi. ii. 608 = hispidus.
obtusifolius, *Teysm. ex Miq. Ann. Mus. Lugd. Bat.* i. 214.—Burma.
Palosapis, *Blanco, Fl. Filip.* ed. II. 813.—Ins. Philipp.
parviflorus, Zipp. in Alg. Konst. en Letter-bode, i. (1829) 297 = Anisoptera polyandra.
pentagonus, *A. DC. in DC. Prod.* xvi. ii. 610.—Borneo.
pentopterus, Dyer, in Journ. Bot. xii. (1874) 106 = macrus.

DIPTEROCARPUS:—
pilosus, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 615.—Burma; Malaya.
plagiatus, Blanco, Fl. Filip. ed. II. 811 = Anisoptera plagata.
polyspermus, *Blanco, l. c.* 812.—Ins. Philipp.
pricnanthus, Dyer, in Journ. Bot. xii. (1874) 104 = fagineus.
pterygocalyx, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 847 = grandiflorus.
quinquenervis, Blume, Mus. Bot. Lugd. Bat. ii. 36 = pentagonus.
retusus, *Blume, Cat. Gew. Buitenz.* 77.—Java.
scaber, *Buch.-Ham. in Mem. Wern. Soc.* vi. (1836) 300. —Ind. or.
scabridus, *Thw. Enum. Pl. Zeyl.* 34.—Zeylan.
Spanoghei, *Blume, Fl. Jav. Dipter.* i. 16. t. 2.—Java.
stellatus, *Vesque, in Compt. Rend. For.* lxxviii. (1874) 626.—Borneo.
stenopterus, *Vesque, l. c.* 626.—Borneo.
Tampacan, *Korth. Verh. Nat. Gesch. Nederl. Bot.* 63 —Borneo.
teres, Steud. Nom. ed II. i. 518 = Dryobalanops aromatica.
thurifer, Blanco, Fl. Filip. ed. II. 810 = Anisoptera thurifera.
trinervis, *Blume, Cat. Gew. Buitenz.* 78.—Java.
turbinatus, *Gaertn. f. Fruct.* iii. 51. t. 186.—Ind. or., Malaya.
tuberculatus, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 614. —Burma.
undulatus, Vesque, in Compt.-Rend. Par. lxxviii. (1874) 626 = Lowii.
validus, *Blume, Mus. Bot. Lugd. Bat.* ii. 36.—Borneo.
vernicifluus, *Blanco, Fl. Filip.* ed. II. 814; *Blume, Mus Bot. Lugd. Bat.* ii. 35.—Ins. Philipp.
vestitus, *Wall. Cat.* n. 954.—Burma.
zeylanicus, *Thw. Enum. Pl. Zeyl.* 33.—Zeylan.

DIPTEROCOMB, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 26 (1835). *COMPOSITÆ*, Benth. & Hook. f. ii. 454.
Jaubertia, Spach, in Jaub. & Spach, Illustr. iii. 181. t. 189 (1849).
pusilla, *Fisch. & Mey. l. c.*—Persia.

DIPTEROSPERMA, Hassk. in Flora, xxv. (1842) ii. Beibl. 28 = Sterculiospermum, Cham. (Bignoniac.).
personatum, Hassk. l. c. = S. caudatum.

DIPTEROSPERMUM, Griff. Notul. iv. 564 (1854) = **Gordonia**, Ellis (Ternstroem.).

DIPTEROTHECA, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) i. Intell. 42; et xxv. (1842) 434 = **Aspilia**, Thou. (Compos.).
Kotschyi, Sch. Bip. ex Hochst. ll. cc. 42, 435 = A. Kotschyana.

DIPTERYGIA, Presl, ex DC. Prod. iv. 80 (1830) = **Asterisculum**, Cham. & Schlecht. (Umbell.).
capitata, Presl, ex DC. l. c. = A. polycephalum.
lasiodiscarpum, Presl, ex DC. l. c. = A. lasidiscarpum.

DIPTERYGIUM, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 66. t. 3. *CRUCIFERÆ*, Benth. & Hook. f. i. 95.
Pteroloma, Hochst. & Steud. ex Steud. Nom. ed. II. ii. 418 (1841).
glaucum, *Decne. l. c.* 67.—Ind. or.; Arab.
scobrum, Decne. ex Boiss. Fl. Orient. i. 417, 958 = glaucum.

DIPTERYX, Schreb. Gen. 495 (1791). *LEGUMINOSÆ*, Benth. & Hook. f. i. 551.
Baryosma, Gaertn. Fruct. ii. 78. t. 93 (1791).
Boldugia, Neck. Elem. iii. 93 (1790).
Coumarouna, Aubl. Pl. Gui. 740. t. 296 (1775), nomen prius.
Rhinhia, Scop. Introd. 301 (1777).
Taralea, Aubl. Pl. Gui. 745. t. 298 (1775).
alata, *Vog. in Linnæa*, xi. (1837) 389.—Bras.
appliculata, Benth. in Hook. Kew Journ. ii. (1850) 234 = oppositifolia.
cornea, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1906, nomen.—Guiana.
crassifolia, *Benth. in Hook. Kew Journ.* ii. (1850) 235. —Guiana.

DIPTERYX:—
emarginata, Spreng. ex Walp. Rep. i. 806 = Pterodon emarginatus.
nodipes, *Tul. in Arch. Mus. Par.* iv. (1844) 100.—Bras.
odorata, *Willd. Sp. Pl.* iii. 910.—Guiana.
oleifera, *Benth. in Hook. Kew Journ.* ii. (1850) 285.—Honduras.
oppositifolia, *Willd. Sp. Pl.* iii. 910.—Guiana.
parviflora, Spruce, ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 136 = oppositifolia.
pteropus, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 127.—Bras.
pterota, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 110 = alata.
reticulata, *Benth. in Hook. Kew Journ.* ii. (1850) 285.—Guiana.
rosea, *Spruce*, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 126.—Bras.
terraphylla, *Benth. l. c.*—Bras.

DIPTYCHANDRA, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 189. *LEGUMINOSAE*, Benth. & Hook. i. 1. 562.
aurantiaca, *Tul. l. c.*—Bras.
cpunctata, *Tul. l. c.*—Bras.
glabra, *Benth. in Mart. Fl. Bras.* xv. ii. 52.—Bras.

DIPTYCHOCARPUS, Trautv. in Bull. Soc. Nat. Mosc. (1850) i. 108. *CRUCIFERAE*, Benth. & Hook. f. i. 67.
ALLOCERATIUM, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 129.
hispidus, *Regel*, *in Act. Hort. Petrop.* v. (1877) 220.—Turkestan.
Olgae, *Regel & Schmalh. l. c.* 231.—Turkestan.
sarawschanicus, *Regel & Schmalh. l. c.*—Turkestan.
strictus, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) i. 108.—Reg. Casp.; Afghan.

DIPTYCHUM, Dulac, Fl. Hautes-Pyr. 81 (1867) = **Scalaria**, Scop. (Gramin.).
cuerulcum, Dulac, l. c. 82 = S. coerulea.

DIPYRENA, Hook. Bot. Misc. i. (1830) 355. *VERBENACEAE*, Benth. & Hook. f. ii. 1146.
WILSONIA, Gill. & Hook. l. c. 372. t. 49 (1830).
dentata, Phil. in Linnaea, xxix. (1857-58) 22 = Baillonia juncea.
glaberrima, *Gill & Hook. in Bot. Misc.* i. (1830) 172.—Chili.
valdiviana, Phil. in Linnaea, xxix. (1857-58) 21 = Baillonia juncea.
Wilsonia, Phil. f. Cat. Pl. Vasc. Chil. 217, sphalm. = glaberrima.

DIRACHMA, Schweinf. ex Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 408. *GERANIACEAE*.
socotrana, *Schweinf. ex Balf. f. l. c.* 404.—Ins. Socotr.

DIRACODES, Blume, Enum. Pl. Jav. i. 55 (1827) = **Amomum**, Linn. (Scitamin.).
javanica, Blume, l. c.—Java.

DIRCA, Linn. Diss. Chen. 17 (1751); Amoen. Acad. iii. 19 (1756). *THYMELAEACEAE*, Benth. & Hook. f. iii. 191.
DOPIA, Adans. Fam. ii. 385 (1763).
occidentalis, *A. Gray*, *in Proc. Am. Acad.* viii. (1873) 631.—Calif.
palustris, *Linn. Sp. Pl.* 858.—Am. bor.

DIRCAEA, Decne. in Rev. Hortic. Sér. III. ii. (1848) 466 = **Geonoma**, Mart.
Blassii, **Regel**, Gartenfl. ii. (1853) 86 = G. Blassii.
bulbosa, Decne. in Rev. Hortic. Sér. III. ii. (1848) 466 = G. bulbosa.
cardinalis, **Regel**, Gartenfl. ii. (1853) 85. t. 41 = G. cardinalis.
Cuspori, Decne in Rev. Hortic. Sér. III. ii. (1848) 466 = G. Cuspori.
fasciculata, Decne. l. c. = G. fasciculis.
Hostiei, Decne. l. c. = G. magnifica.
interrupta, Decne. l. c. = G. interrupta.
lobulata, Lem. jard. Fleur. iii. (1853-55) t. 219 = G. cardinalis.
macrocalix, Lem. Illustr. Hortic. ii. (1855) t. 54 = G. macrocalix.
magnifica, Decne. in Rev. Hortic. Sér. III. ii. (1848) 466 = G. magnifica.
picta, Lem. jard. Fleur. iii. (1852-53) t. 302 = Isoloma pictum.
reflexa, Decne. in Rev. Hortic. Sér. III. ii. (1848) 466 = G. reflexa.
Suttoni, Decne. l. c. = G. Suttoni.
Vauthieri, Decne. l. c. = G. Vauthieri.

DIRHACODES, Lem. in Orb. Dict. v. 73 (1849) = DIRACODES, Blume = **Amomum**, Linn. (Scitam.).

DIRHYNCHOSIA, Blume, Mél. Bot. ex Flora, xli. (1858) 254 = **Spiraeopsis**, Miq. (Saxifrag.).
celebica, Blume, l. c. = S. celebica.

DIRICHLETIA, Klotzsch, in Monatsb. Akad. Berl. (1853) 494. *RUBIACEAE*, Benth. & Hook. f. ii. 56.
glabra, *Klotzsch, l. c.*—Afr. trop.
glaucescens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 51.—Afr. trop.
insignis, Vatke, in Bremen, Abh. ix. (1885) 117 = seq.
Kirondron, Vatke, l. c. = Carphalea Kirondron.
lanceolata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 585.—Ins. Socotra.
obovata, *Balf. f. l. c.*—Ins. Socotra.
Perrelliana, Vatke, in Bremen, Abh. ix. (1885) 117 = Carphalea Perrelliana.
pubescens, *Klotzsch, in Monatsb. Akad. Berl.* (1853) 494.—Afr. trop.
venulosa, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 584.—Ins. Socotra.

DIRTEA, Rafin. Fl. Tellur. iii. 69 (1836) = **Commelina**, Linn.
africana, Rafin. l. c. iv. 122 = C. africana.
japonica, Rafin. l. c. iii. 69 = Aneilema japonicum.

DISA, Berg. Fl. Cap. 348 (1767). *ORCHIDEAE*, Benth. & Hook. f. iii. 630.
GAMARIA, Rafin. Fl. Tellur. iv. 49 (1836).
HERSCHELIA, Lindl. Gen. et Sp. Orch. 362 (1838).
MONADENIA, Lindl. l. c. 356 (1838).
PENTHEA, Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
REPANDRA, Lindl. Orch. Scel. 12 (1826).
SONERILEDIUM, Lindl. Gen. et Sp. Orch. 358 (1838).
aconitoides, *Sond. in Linnaea*, xix. (1847) 91.—Afr. austr.
acuta, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 69.—Afr. austr.
affinis, *N. E. Br. in Gard. Chron.* xxiv. (1885) 402.—Afr. austr.
alpina, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 220.—Afr. trop.
atricapilla, *Bolus, l. c.* xix. (1892) 344.—Afr. austr.
atropurpurea, *Sond. in Linnaea*, xix. (1847) 96.—Afr. austr.
attenuata, *Lindl. Gen. et Sp. Orch.* 351.—Afr. austr.
aurantiaca, *Reichb. f. in Flora*, l. (1867) 98.—Afr. trop.
barbata, *Sw. in Vet. Acad. Handl.* xxi. (1800) 212.—Afr. austr.
Bartoni, *Hort. ex De Puydt, Læ Orchid.* (1880) 275. t. 18.—Afr. austr.
bifida, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 212.—Afr. austr.
Bodkini, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 74.—Afr. austr.
borbonica, *Balf. f. & Moore, in Journ. Bot.* xiv. (1876) 293.—Ins. Borbon.
brachycrea, *Lindl. Gen. et Sp. Orch.* 355.—Afr. austr.
bracteata, Lindl. l. c. 354 = cylindrica.
brachiata, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 211.—Afr. austr.
Buchenaviana, *Kraenzl. in Bremen, Abh.* vii. (1882) 261.—Madag.
capricornis, Reichb. f. in Linnaea, xx. (1847) 692 = gladioliflora.
caulescens, *Lindl. Gen. et Sp. Orch.* 351.—Afr. austr.
Cephalotes, *Reichb. f. Otia Bot. Hamb.* 106.—Afr. austr.
cernua, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 211.—Afr. austr.
Charpentieriana, *Reichb. f. in Linnaea*, xx. (1847) 668.—Afr. austr.
chrysostachya, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 211.—Afr. austr.
coerulea, Reichb. f. Otia Bot. Hamb. 119 = Brownlcea caerulea.
Cooperi, *Reichb. f. in Flora*, lxiv. (1881) 328.—Afr. austr.
cornuta, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 210.—Afr. austr.
crassicornis, *Lindl. Gen. et Sp. Orch.* 348.—Afr. austr.
cylindrica, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 213.—Afr. austr.
Deckenii, *Reichb. f. Otia Bot. Hamb.* 105.—Afr. trop
draconis, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 210.—Afr. austr.
elegans, *Reichb. f. in Flora*, xlviii. (1865) 182.—Afr. austr.
equestris, *Reichb. f. l. c.* 181.—Afr. trop.
excelsa, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 213, excl. syn.—Afr. austr.
excelsa, Thunb. ex N. E. Br. in Journ. Linn. Soc. xx. (1884) 479 = venosa.
extinctoria, *Reichb. f. in Flora*, lxiv. (1881) 328.—Afr. austr.
fascista, *Lindl. Gen. et Sp. Orch.* 350.—Afr. austr.
ferruginea, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 210.—Afr. austr.
filicornis, Thunb Fl. Cap. i. 87 = patens.
flexuosa, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 212.—Afr. austr.
gladioliflora, *Burch. ex Lindl. l. c.* 351.—Afr. austr.
gracilis, Lindl. l. c. 348 = chrysostachya.
graminifolia, *Ker-Gawl. in Brande, Journ.* vi. (1819) 44. t. 1.—Afr. austr.
grandiflora, Linn. f. Suppl. 406 = uniflora.
Harveiana, *Lindl. in Hook. Lond. Journ. Bot.* i. (1842) 15.—Afr. austr.
hemisphaerophora, Reichb. f. Otia Bot. Hamb. 106 = stachyoides.
hians, Spreng. Syst. iii. 698 = lacera.
hircicornis, *Reichb. f. Otia Bot. Hamb.* 105.—Afr. trop.
Huttoni, Reichb. f. l. c. = sanguinea.
incarnata, *Lindl. Gen. et Sp. Orch.* 348.—Madag.
inflexa, *Mundt, ex Lindl. l. c.* 360.—Afr. austr.
lacera, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 212.—Afr. austr.
lacta, *Reichb. f. Otia Bot. Hamb.* 106.—Afr. austr.
leptostachys, Sond. in Linnaea, xix. (1847) 98 = tenuis.
linearis, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 74.—Afr. austr.
longicornis, Thunb. Prod. Pl. Cap. 4 = seq.
longicornu, *Linn. f. Suppl.* 406.—Afr. austr.
longifolia, *Lindl. Gen. et Sp. Orch.* 349.—Afr. austr.
lugens, *Bolus, in Journ. Linn. Soc.* xx. (1884) 483.—Afr. austr.
Macowani, *Reichb. f. Otia Bot. Hamb.* 106.—Afr. austr.
macrantha, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 210.—Afr. austr.
macroceras, Reichb. f. Otia Bot. Hamb. 119 = Brownlea caerulea.
macroglottis, Sond. ex Drège, in Linnaea, xx. (1847) 219, nomen = Charpentieriana.
maculata, Harv. ex Lindl. in Hook. Lond. Journ. Bot. i. (1842) 15 = ocellata.
maculata, *Linn. f. Suppl.* 407.—Afr. austr.
megacrcas, Hook. f. Bot. Mag. t. 6529 = crassicornis.
melaleuca, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 213.—Afr. austr.
minor, *Reichb. f. in Flora*, xlviii. (1865) 182.—Afr. austr.
modesta, Reichb. f. in Linnaea, xx. (1847) 690 = vaginata.
montana, *Sond. in Linnaea*, xix. (1847) 90.—Afr. austr.
multifida, *Lindl. Gen. et Sp. Orch.* 353.—Afr. austr.
natalensis, Lindl. in Hook. Lond. Journ. Bot. i. (1842) 16 = polygonoides.
neglecta, *Sond. in Linnaea*, xix. (1847) 100.—Afr. austr.
nervosa, *Lindl. Gen. et Sp. Orch.* 352.—Afr. austr.
obtusa, *Lindl. l. c.* 355.—Afr. austr.
ocellata, *Bolus, in Journ. Linn. Soc.* xx. (1884) 477.—Afr. austr.

DISA:—
ochrostachya, *Reichb. f. in Flora*, xlviii. (1865) 181.—Afr. trop.
oligantha, *Reichb. f. l. c.* 182.—Afr. austr.
ovalifolia, *Sond. in Linnaea*, xix. (1847) 98.—Afr. austr.
parviflora, Reichb. f. Otia Bot. Hamb. 119 = Brownlea parviflora.
parvilabris, Bolus, in Journ. Linn. Soc. xix. (1882) 344 = oligantha.
patens, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 214.—Afr. austr.
patens, Thunb. Fl. Cap. i. 85 = tenuifolia.
patula, *Sond. in Linnaea*, xix. (1847) 94.—Afr. austr.
physodes, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 211.—Afr. austr.
physodes, Thunb. Fl. Cap. i. 71 = Monadenia inflata.
picta, *Sond. in Linnaea*, xix. (1847) 99.—Afr. austr.
polygonoides, *Lindl. Gen. et Sp. Orch.* 349.—Afr. austr.
porrecta, [Ker-Gawl. in] Journ. Sc. Roy. Institt. v. (1818) t. 1. f. 1 = ferruginea.
porrecta, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 211.—Afr. austr.
prasinata, [Ker-Gawl. in] Bot. Reg. t. 910 = cernua.
propinqua, *Sond. in Linnaea*, xix. (1847) 95.—Afr. austr.
pulchella, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 301. —Abyss.
pulchra, *Sond. in Linnaea*, xix. (1847) 94.—Afr. austr.
purpurascens, *Bolus, in Journ. Linn. Soc.* xx. (1884) 492.—Afr. austr.
pygmæa, *Bolus, l. c.* xxii. (1885) 79.—Afr. austr.
racemosa, *Linn. f. Suppl.* 406.—Afr. austr.
recurvata, Reichb. f. Otia Bot. Hamb. 119 = Brownlea recurvata.
reflexa, Reichb. f. in Flora, xlviii. (1865) 183 = patens.
retinalata, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 73. —Afr. austr.
Richardiana, *Lem. ex Lindl. Gen. et Sp. Orch.* 361. —Afr. austr.
rosea, *Lindl. l. c.* 350.—Afr. austr.
rufescens, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 210.—Afr. austr.
Rutenbergiana, *Kraenzl. in Bremen, Abh.* vii. (1882) 262.—Madag.
sagittalis, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 213.—Afr. austr.
sanguinea, *Sond. in Linnaea*, xix. (1847) 97.—Afr. austr.
schlechtioides, *Sond. l. c.* 92.—Afr. austr.
Scullyi, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 70.—Afr. austr.
scutellifera, *A. Rich. in Ann. Sc. Nat. Sér.* II. xiv. (1840) 373.—Abyss.
secunda, Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 218 = racemosa.
spathulata, *Sw. l. c.*—Afr. austr.
stachyoides, *Reichb. f. in Flora*, lxiv. (1881) 338.—Afr. austr.
stricta, *Sond. in Linnaea*, xix. (1847) 91.—Afr. austr.
tabularis, *Sond. l. c.* 99.—Afr. austr.
Telipogonis, *Reichb. f. l. c.* xx. (1847) 689.—Afr. austr.
tenella, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 212.—Afr. austr.
tenuicornis, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 68.—Afr. austr.
tenuifolia, Sw. in *Vet. Acad. Handl. Stockh.* xxi. (1800) 214.—Afr. austr.
tenuis, *Lindl. Gen. et Sp. Orch.* 354.—Afr. austr.
torta, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 211.—Afr. austr.
trilobe, *Lindl. Gen. et Sp. Orch.* 351.—Afr. austr.
tripartita, *Lindl. l. c.* 353.—Afr. austr.
uncinata, *Bolus, in Journ. Linn. Soc.* xx. (1884) 478. —Afr. austr.
uniflora, *Berg. Desc. Pl. Cap.* 348.—Afr. austr.
vaginata, *Harv. ex Lindl. in Hook. Lond. Journ. Bot.* i. (1842) 15.—Afr. austr.
venosa, *Lindl. Gen. et Sp. Orch.* 351.—Afr. austr.
venosa, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 213.—Afr. austr.
venusta, *Bolus, in Journ. Linn. Soc.* xx. (1884) 482. —Afr. austr.
verdickii, *Reichb. f. in Flora*, xlviii. (1865) 181.—Afr. trop.

DISA :—
Walleri, *Reichb. f. Otia Bot. Hamb.* 105.—Afr. trop.
Welwitschii, *Reichb. f. in Flora*, xlviii. (1865) 181.—Afr. trop.
Zeyheri, Sond. in Linnæa, xix. (1847) 95 = ferruginea.

DISACHOENA, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indië, i. (1844) 488, nomen. *UMBELLIFERÆ* (Ordo?).

DISADENA, Miq. Stirp. Surin. 150 (1850) = **Voyria**, Aubl. (Gentian.).
flavescens, Miq. l. c. = V. flavescens.

DISAKISPERMA, Steud. Syn. Pl. Gram. 287 (1855). *GRAMINEÆ*, Benth. & Hook. f. iii. 1096.
mexicana, *Steud. l. c.*—Mexic.

DISANDRA, Linn. in Murr. Syst. ed. XIII. 290 (1774) = **Sibthorpia**, Linn. (Scrophul.).
Distandra, Link, Enum. Hort. Berol. i. 354 (1821).
africana, Linn. Syst. ed. Reich. ii. 187 = S. africana.
prostrata, Murr. Syst. ed. XIII. 290 = S. peregrina.

DISANTHUS, Maxim. in Bull. Acad. Pétersb. x. (1866) 485. *HAMAMELIDEÆ*, Benth. & Hook. f. i. 1005.
cercidifolia, *Maxim. l. c.* 486.—Japon.

DISARRENUM, Labill. Nov. Holl. Pl. ii. 82. t. 232 (1806) = **Hierochloë**, S. G. Gmel. (Gramin.).
antarcticum, Labill. l. c. 83. t. 232 = H. antarctica.

DISARRHENUM, Beauv. Agrost. 62 (1812) = præc.

DISCANTHERA, Torr. & Gray, Fl. N. Am. 596 (1840) = **Cyclanthera**, Schrad. (Cucurbit.).
dissecta, Torr. & Gray, l. c. 597 = C. dissecta.

DISCANTHUS, Spruce, in Journ. Linn. Soc. iii. (1859) 136 = **Cyclanthus**, Poit.
odoratus, Spruce, l. c. = C. bipartitus.

DISCARIA, Hook. Bot. Misc. i. (1830) 156. t. 64, 65. *RHAMNEÆ*, Benth. & Hook. f. i. 383.
Notophæna, Miers, in Ann. & Mag. Nat. Hist. Ser. III. v. (1860) 267.
Ochetophila, Poepp. ex Reiss. in Endl. Gen. 1099 (1840).
Tetrapasma, G. Don, Gen. Syst. ii. 40 (1832).
americana, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 156.—Reg. Argent.
aphylla, *Meyen, Reise*, i. 488.—Peruv.
articulata, *Miers, Contrib.* i. 305, errat.—Chili.
australis, *Hook. Bot. Misc.* i. 157, in note, t. 45.—Austral.
crenata, *Regel, Ind. Sem. Hort. Petrop.* (1860) 50.—Hab.?
febrifuga, *Mart. Syst. Mat. Med. Veg. Bras.* 37.—Bras.
longipina, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 273.—Reg. Argent.
lycioides, *Miers, l. c.* 272.—Uruguay.
parviflora, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 229.—Galapag.
pauciflora, *Hook. f. l. c.*—Ins. Galapag.
serratifolia, *Benth. & Hook. f. ex* [*Mast. in*] *Gard. Chron.* (1876) ii. 324, in note.—Chili; Patag.
spiculiger, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 274.—Reg. Argent.
Tournaton, *Raoul, Choix Pl. Nouv. Zél.* 29.—N. Zel.

DISCHANTHIUM, Kunth, Enum. Pl. i. 498 (1833) = DICHANTHIUM, Willem. = **Andropogon**, Linn. (Gramin.).

DISCHEMA, Voigt, Hort. Suburb. Calc. 566 (1845) = **Rhabdosia**, Wall. (Sctamin.).
glaucum, Voigt, l. c. = H. glauca.

DISCHIDIA, R. Br. in Mem. Wern. Soc. i. (1809) 82. *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 777.
Collyris, Miq. Fl. Ind. Bat. ii. 518 (1857).
Colyris, Vahl, in Skrift. Nat. Selsk. Kiobenh. vi. (1810) 110.
Conchophyllum, Blume, Bijdr. 1060 (1826).
Leptostemma, Blume, l. c. 1057 (1826).

DISCHIDIA :—
acutifolia, *Maingay, ex Hook. f. Fl. Brit. Ind.* iv. 51. —Malacca.
albida, *Grif. Notul.* iv. 46.—Malacca.
albiflora, Griff. l. c. 47 = D. Collyris 1.
angustifolia, *Miq. Fl. Ind. Bat.* ii. 519.—Java.
bengalensis, *Colebr. in Trans. Linn. Soc.* xii. (1818) 357. t. 18.—Ind. or. ; Malaya.
Brunoniana, Griff. Notul. iv. 44 = hirsuta.
chinensis, *Champ. ex Benth. in Hook. Kew Journ.* v. (1853) 85.—China.
clavata, *Wall. Cat.* n. 4206 ; *Pl. As. Rar.* ii. 86.—Burma
coccinea, *Grif. Notul.* iv. 45.—Malacca.
cochlearia, *Blume, Bijdr.* 1060.—Java.
Collyris, *Wall. Cat.* n. 4207.—Malaya.
complex, *Grif. Notul.* iv. 50.—Malacca.
crassifolia, *Zipp. ex Span. in Linnaea*, xv. (1841) 324, nomen.—Ins. Timor.
cuneifolia, Wall. Cat. n. 4206 = bengalensis.
fasciculata, Decne. in DC. Prod. viii. 633 = hirsuta.
formosana, *Maxim. in Bull. Acad. Pétersb.* xxiii (1877) 385.—Ins. Formosa.
Gaudichaudii, *Decne. in DC. Prod.* viii. 632.—Java, Ins. Timor.
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 51.—Burma.
hirsuta, *Decne. in DC. Prod.* viii. 633.—Burma ; Malaya.
Horsfieldiana, *Miq. Fl. Ind. Bat.* ii. 510.—Java.
imbricata, Steud. Nom. ed. II. i. 519 = D. Collyris.
khasiana, *Hook. f. Fl. Brit. Ind.* iv. 50.—Reg. Himal.
lanceolata, *Decne. in DC. Prod.* viii. 631.—Java.
latifolia, *Decne. l. c.*—Java.
lutescens, *Zipp. ex Span. in Linnaea*, xv. (1841) 324, nomen.—Ins. Timor.
nicobarica, *Dietr. in Kjoeb. Vidensh. Meddel.* (1854) 189.—Ins. Nicobar.
Nummularia, Blume, Bijdr. 1059 = Gandichaudii.
Nummularia, *R. Br. Prod.* 461.—Ind. or. ; Malaya ; Austral.
obovata, *Decne. in DC. Prod.* viii. 633.—Java.
obovata, Griff. Notul. iv. 51 = Griffithii.
orbicularis, Decne. in DC. Prod. viii. 632 = D Nummularia.
ovata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 226.—N. Guinea.
oxyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 141.—Ins. Celebes.
peltata, *Blume, Mus. Bot. Lugd. Bat.* i. 148.—N. Guinea.
picta, *Blume, l. c.*—N. Guinea.
pruinosa, Zipp. ex Blume, Rumphia, iv. 80 = Hoya pruinosa.
puberula, *Decne. in DC. Prod.* viii. 631.—Ins. Mariana.
punctata, *Decne. l. c.*—Java.
Rafflesiana, *Wall. Pl. As. Rar.* ii. 85. t. 149.—As. et Austral. trop.
rhombifolia, *Blume, Bijdr.* 1058.—Java.
Rumphii, *Miq. Fl. Ind. Bat.* ii. 508.—Ins. Amboine.
sagittata, *Decne. in DC. Prod.* viii. 631.—Java.
spatulata, Blume, Bijdr. 1060 = bengalensis.
Spironema, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 324.—Java.
timorensis, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 377.—Ins. Timor.
truncata, *Decne. in DC. Prod.* viii. 632.—Java.
viridescens, Griff. Notul. iv. 49 = Rafflesiana.
Wallichii, *Wight, Contrib.* 48.—Singapore.

DISCHIDIUM, Reichb. Handb. 269 (1837) = **Viola**, Tourn.
bifforum, Opiz, Seznam, 88 = V. bifiora.

DISCHIMIA, Reichb. Consp. 121 (1828) = seq.

DISCHISMA, Choisy, in Mém. Soc. Phys. Genèv. ii. 11. (1823) 93. *SELAGINEÆ*, Benth. & Hook. f. ii. 1123.
arenarium, *E. Mey. Comm. Pl. Afr. Austr.* 251.—Afr. austr.
capitatum, *Choisy, in Mém. Soc. Phys. Genève.* ii. 11. (1823) 94.—Afr. austr.
chenaedryfolium, Walp. Rep. iv. 148 = ciliatum.
ciliatum, *Choisy, in Mém. Soc. Phys. Genève.* ii. 11. (1823) 94.—Afr. austr.
clandestinum, *E. Mey. Comm. Pl. Afr. Austr.* 251.—Afr. austr.
ericoides, *Sweet, Hort. Brit.* ed. II. 414.—Afr. austr.

DISCHISMA :—
flaccum, E. Mey. Comm. Pl. Afr. Austr. 260 = crinoides.
fruticosum, *Rolfe, in Journ. Linn. Soc.* xx. (1883) 351. —Afr. austr.
hispidum, Sweet, Hort. Brit. ed. II. 415; Choisy, in DC. Prod. xii. 7 = ciliatum.
leptostachyum, *E. Mey. Comm. Pl. Afr. Austr.* 261.—Afr. austr.
spicatum, *Choisy, in Mém. Soc. Phys. Genèv.* ii. 11. (1823) 96.—Afr. austr.

DISCHLIS, Phil. Sert. Mendoc. Alt. 51 (1871), sphalm. = **Distichlis**, Rafin. (Gramin.).

DISCHORISTE, D. Dietr. Syn. Fl. iii. 585 (1843) = DYSCHORISTE, Nees = **Calophanes**, D. Don (Acanth.).

DISCIPHANIA, Eichl. in Mart. Fl. Bras. xiii. i. 168 t. 86 (1864). *MENISPERMACEAE*, Benth. & Hook. f. i. 960.
Ernsti, *Eichl. in Yahrb. Bot. Gart. Berl.* ii. (1883) 230.—Venezuela.
lobata, *Eichl. in Mart. Fl. Bras.* xiii. i. 169. t. 86. f. 1. —Bras.

DISCOCACTUS, Pfeiff. in Otto & Dietr. Allg. Gartenz. v. (1837) 241; et in Nov. Act. Nat. Cur. xix. i. (1839) 117. t. 15. *CACTEAE*, Benth. & Hook. f. i. 848.
alticolens, *Lem. ex A. Dietr. in Allg. Gartenz.* xiv. (1846) 209.—Bras.
biformis, Walp. Rep. v. 817, sphalm. = Phyllocactus biformis.
insignis, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 241; *et in Nov. Act. Nat. Cur.* xix. i. (1839) 119.—Ind. occ.
Lehmanni, *Pfeiff. l. c.* xix. i. (1839) 120.—Hab. ?
Linkii, *Pfeiff. l. c.*—Hab. ?
tricornis, *Monv. ex Pfeiff. & Otto, Abbild. Cacteen*, ii. t. 38.—Bras.

DISCOCACTUS, Walp. Rep. v. 877 (1858), sphalm. = DISOCACTUS, Lindl. = **Phyllocactus**, Link (Cact.).

DISCOCAPNOS, Cham. & Schlecht. in Linnaea, i. (1826) 569 = **Fumaria**, Linn. (Papav.).
Mundtii, Cham. & Schlecht. l. c. = F. Mundtii.

DISCOCARPUS, Klotzsch, in Wiegm. Archiv, vii. (1841) 201. t. 9. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 269.
brasiliensis, *Klotzsch, l. c.*—Bras.
caeruleobotrys, *Klotzsch, l. c.*—Guian.
Spruceanus, *Muell. Arg. in Linnaea*, xxxii. (1863) 78. —Bras.

DISCOCARPUS, Lehm. in Vidensk. Selsk. Skr. V. ii. (1851) 808 = **Laportea**, Gaudich. (Urticac.).
mexicana, Liebm. l. c. 809 = L. platycarpa.
nicaraguensis, Liebm. l. c. = L. nicaraguensis.

DISCOLENTA, Rafin. New Fl. Am. iv. 48 (1836); Fl. Tellur. iii. 15 (1836) = **Polygonum**, Tourn.
lapathifolia, Rafin. Fl. Tellur. iii. 15 = P. lapathifolium.
punctata, Rafin. New Fl. Am. iv. 48 = P. punctatum, *Rafin.*
scabra, Rafin. l. c.; Fl. Tellur. iii. 15 = P. lapathifolium ?

DISCOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 105. *LEGUMINOSAE*, Benth. & Hook. f. i. 516.
elongatum, *Benth. in Mart. Fl. Bras.* xv. t. 73.—Bras.
hirtum, *Benth. l. c.*—Bras.
leptophyllum, *Benth. l. c.* t. 72.—Bras.
psoraliaefolium, *Benth. l. c.*—Bras.
pulchellum, *Benth. in Ann. Wien. Mus.* ii. (1838) 106.—Bras.

DISCOMELA, Rafin. Neogenyt. 3 (1825) = **Helianthus**, Linn. (Compos.).
angustifolius, Rafin. l. c. = H. angustifolius.
atrorubens, Rafin. l. c. = H. atrorubens.
sparsiflorus, Rafin. l. c. = H. atrorubens.

DISCOPHORA, Miers, in Ann. & Mag. Nat. Hist. *Ser.* II. x. (1852) 118. *OLACINEAE*, Benth. & Hook. f. i. 393.
KURKERIA, *Mart. Herb. Fl. Bras.* n. 1276 (1840 ?).
guianensis, *Miers, l. c.* 119.—Guiana.

DISCOPHYTUM, Miers, in Lindl. Veg. Kingd. 701 (1847) = **Calycera**, Cav.

DISCOPLEURA, DC. Coll. Mém. v. 88 (1829); et Prod. iv. 106 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 906.
DAUCOSMA, Engelm. & Gray, ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 210.
PTILIMNIUM, Rafin. in Am. Monthly Mag. (1819) 192.
capillaceum, *DC. Coll. Mém.* v. 88. t. 8.—Am. bor.
capillacea, Hook. & Arn. Bot. Beech. Voy. 142 = Aplastrum angustifolium.
costata, Steud. Nom. ed. II. i. 520 = capillacea.
daucifolia, Benth. & Hook. f. Gen. i. 907 = Seseli daucifolium.
juncea, Steud. Nom. ed. II. i. 520 = Nuttallii ?
laciniata, *Benth. & Hook. f. ex S. Wats. Bibliog. Ind. N. Am. Bot.* 430.—Am. bor.
Nuttallii, *DC. Coll. Mém.* v. 89. t. 9.—Am. bor.

DISCOPLIS, Rafin. New Fl. Am. iv. 9 (1836) = Drypetes ! Vahl (Euphorb.).
serrata, Rafin. l. c.—Florida; Cuba.

DISCOPODIUM, Hochst. in Flora, xxvii. (1844) 22. *SOLANACEAE*, Benth. & Hook. f. ii. 893.
penninervium, *Hochst. l. c.*—Abyss.

DISCOPODIUM, Steud. Syn. Pl. Cyp. 150 (1855) = **Trioostularia**, Nees (Cyperac.).
Drummondii, Steud. l. c. = T. compressa, Necsil.

DISCOREA, Miq. Fl. Ind. Bat. iii. 572 (1856) sphalm. = **Dioscorea**, Linn.

DISCOSPERMUM, Dals. in Hook. Kew Journ. ii. (1850) 257 = **Diplospora**, DC. (Rubiac.).
apiocarpum, Dals. l. c. = Diplosp. apiocarpa.
Dalzellii, Thw. Enum. Pl. Zeyl. 138 = Diplosp. Dalzellii.
erythrosporum, Thw. l. c. = Diplosp. erythrospora.
sphaerocarpum, Dalz. in Hook. Kew Journ. ii. (1850) 257 = Diplosp. sphaerocarpa.

DISCOSTIGMA, Hassk. in Flora, xxv. (1842) ii. Beibl. 83 = **Garcinia**, Linn. (Guttif.).
acuminatum, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 364 = G. terpnophylla.
caudatum, A. Gray, Bot. U. St. Expl. Exped. i. 217, in textu = G. rostrata.
corymbosa, Panch. & Sebert, Not. Bois Nouv. Caléd. 234.—N. Caled.
febrile, Miq. Fl. Ind. Bat. Suppl. 496 = G. cornea.
grandifolium, Choisy, in Zoll. Syst. Verz. Ind. Archip. 149, 150 = G. grandifolia.
mergurnse, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 363 = G. merguensis.
rostratum, Hassk. in Flora, xxv. (1842) ii. Beibl. 83 = G. rostrata.
vitiense, A. Gray, Bot. U. St. Expl. Exped. i. 216. t. 16 = G. vitiensis.
zeylanicum, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 364 = G. terpnophylla.

DISCOVIUM, Rafin. in Journ. Phys. lxxxix. (1819) 96 = **Lepidium** ! Tourn. (Crucif.).
gracile, Rafin. l. c. = L. virginicum ?
obtectum, DC. Syst. ii. 700 = L. virginicum ?

DISCURAINIA, Walp. Rep. i. 163 (1842) = DESCURAINIA, Webb = **Sisymbrium**, Linn. (Crucif.).

DISCUREA, Schur, Enum. Pl. Transs. 54 (1866) = **Sisymbrium**, Linn. (Crucif.).
Sophia, Schur, l. c. = S. Sophia.

DISECOCARPUS, Hassk. in Flora, xlix. (1866) 211 = **Commelina**, Linn.
Kotschyi, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 172 = C. Kotschyi.
polygamus, Hassk. Commel. Ind. 9 = C. communis.

DISELMA, Hook. f. Fl. Tasm. i. 353. t. 96 (1860) = **Pittosys**, Hook. f. (Conif.).
Archeri, Hook. f. l. c. = F. Archeri.

DISEMMA, Labill. Sert. Austr. Caled. 78. t. 79 (1824) = **Passiflora**, Linn.
adianthifolia, DC. Prod. iii. 333 = P. glabra.
aurantia, Labill. Sert. Austr. Caled. t. 79 = P. aurantia.
Barclayi, Seem. Fl. Vit. 96 = P. Barclayi.
Baueri, G. Don, Gen. Syst. iii. 56 = P. Baueriana.
Baueriana, Endl. Prod. Fl. Norfolk. 138 = P. Baueriana.
brachystephana, F. Muell. Fragm. i. 56 = P. brachystephana.
caerulescens, Seem. in Bonplandia, x. (1862) 366.—Ins. Tangan.
Calyana, M. Roem. Syn. Pepon. 189 = P. Herbertiana.
coccinea, DC. Prod. iii. 333 = P. Banksii.
filamentosa, Regel & Koern. Ind. Sem. Hort. Petrop. (1858) 43 = P. filamentosa.
floribunda, Fourn. in Rev. Hortic. xl. (1869) 430 = P. sexflora.
Hahnii, Fourn. l. c. = P. Hahnii.
Herbertiana, DC. Prod. iii. 333 = P. Herbertiana.
Horsfieldii, Miq. Fl. Ind. Bat. i. i. 700 = P. Horsfieldii.
Medusaea, Fourn. in Rev. Hortic. xl. (1869) 430 = P. jorullensis.
moluccana, Miq. Fl. Ind. Bat. i. i. 699 = P. moluccana.
Muelleriana, Regel, Ind. Sem. Hort. Petrop. (1866) 101 = P. Muelleriana.
penangiana, Miq. Fl. Ind. Bat. i. i. 700 = Modecca nicobarica ?
Storckii, Seem. Fl. Vit. 96.—Ins. Fiji.
sumatrana, Miq. Fl. Ind. Bat. i. i. 700 = P. sumatrana.
timoriana, Miq. l. c. = P. timoriana.
vitiensis, Seem. Fl. Vit. 96 = P. vitiensis.

DISEPALUM, Hook. f. in Trans. Linn. Soc. xxiii. (1862) 156. t. 20. *ANONACEAE*, Benth. & Hook. f. i. 25.
anomalum, *Hook. f. l. c.*—Borneo.
coronatum, *Becc. in Nouv. Giorn. Bot.* ii. (1870) 155. —Borneo.

DISERIS, Wight, Ic. v. n. 1719, in textu (1853) = **Disperis**, Sw. (Orchid.).

DISERNESTON, Jaub. & Spach, Illustr. i. 78. t. 40 (1848) = Dorema, D. Don (Umbell.).
gummiferum, Jaub. & Spach, l. c. = Dor. Ammoniacum.
hirsutum, Loftus, in Borcs. Ferul. 28 = Dor. Ammoniacum.

DISGREGA, Hassk. in Flora, xlix. (1866) 215 = **Tradescantia**, Linn. (Commelin.).
mexicana, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 205 = T. Disgrega.

DISINSTYLIS, Rafin. Fl. Tellur. iii. 81 (1836) = **Chlora**, Adans. (Gentianac.).
italica, Rafin. l. c. = C. imperfoliata.

DISISOCACTUS, G. K[unze], in Bot. Zeit. iii. (1845) 588; Salm-Dyck, Cact. ed. II. 57 (1850) = DISOCACTUS, Lindl. = **Phyllocactus**, Link (Cactese).
biformis, Salm-Dyck, l. c. = P. biformis.

DISKION, Rafin. Sylva Tellur. 55 (1838) = **Sarracha**, Ruiz & Pav. (Solanac.).

DISMOPHYLA, Rafin. Fl. Tellur. iii. 86 (1836) = **Drosera**, Linn.
binata, Rafin. l. c. = Dros. binata.

DISOCACTUS, Lindl. Bot. Reg. (1845) t. 9 = **Phyllocactus**, Link (Cactese).
biformis, Lindl. l. c. = P. biformis.

DISODEA, Pers. Syn. i. 210 (1805) = **Lygodisodea**, Ruiz & Pav. (Rubiac.).
foetida, Pers. l. c. = L. foetida.

DISODIA, Dum. Anal. Fam. 33 (1829) = prec.

DISOMENE, A. DC. Prod. xvi. ii. 600, in syn. (1868) = DYSEMONE, Soland. = **Gunnera**, Linn. (Halorag.).

DISOON, A. DC. in DC. Prod. xi. 702 (1847) = **Myoporum**, Banks.
Beckeri, F. Muell. Fragm. iv. 48 = M. Beckeri.
corvifolius, F. Muell. l. c. i. 126 = M. salasioides.
floribundus, A. DC. in DC. Prod. xi. 702 = M. floribundum.

DISOXYLON, Reichb. Handb. 318 (1837) = **Dysoxylum**, Blume (Meliac.).

DISOXYLUM, Benth. & Hook. f. Gen. i. 335 (1862) = DYSOXYLUM, Hook. f. & Thoms. = **Amoora**, Roxb. (Meliac.).

DISOXYLUM, A. Juss. in Mém. Mus. Par. xix. (1830) 224 = **Dysoxylum**, Blume (Meliac.).

DISPARA, Rafin. Sylva Tellur. 112 (1838) = **Cristatella**, Nutt. (Capparid.).
filiformis, Rafin. l. c. 113 = C. cross.

DISPARAGO, Gaertn. Fruct. ii. 463. t. 178 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 323.
WIGANDIA, Neck. Elem. i. 95 (1790).
ericoides, *Gaertn. l. c.*—Afr. austr.
gomphrenoides, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 180.—Afr. austr.
Krausii, *Sch. Bip. in Flora*, xxvii. (1844) 693.—Afr. austr.
lasiocarpa, *Cass. in Dict. Sc. Nat.* xxxiv. 42.—Afr. austr.
lasiolia, *DC. Prod.* vi. 257.—Afr. austr.
seriphioides, *DC. l. c.*—Afr. austr.
tortilis, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 181.—Afr. austr.

DISPELTOPHORUS, Lehm. Ind. Sem. Hort. Hamb. (1836) n. 2 = **Menonvillea**, R. Br. (Crucif.).
crassifolius, Lehm. l. c. = M. linearis.

DISPERIS, Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 218. *ORCHIDEAE*, Benth. & Hook. f. iii. 533.
DIPERA, Spreng. Syst. iii. 676 (1826).
DIVOPSIA, Thou. Orch. Iles Afr. t. 1 ad 3 (1822).
DIVOPSIS, Thou. l. c. tabl. 1 (1822).
alata, Labill. Pl. Nov. Holl. ii. 30. t. 210 = Pterostylis praecox.
anthoceros, *Reichb. f. Otia Bot. Hamb.* 103.—Afr. austr.
capensis, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 220.—Afr. austr.
cardiophora, *Harv. Thes. Cap.* ii. 4.—Afr. austr.
Cooperi, *Harv. l. c.* 47.—Afr. austr.
cordata, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 218.—Ins. Maurit.
crassinalis, *Reichb. f. in Linnaea*, xxii. (1849) 860.—Abyss.
cucullata, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 220.—Afr. austr.
Fanninae, *Harv. Thes. Cap.* ii. 46.—Afr. austr.
galerita, *Reichb. f. Otia Bot. Hamb.* 108.—Afr. trop.
graminifolia, Banks, ex Steud. Nom. ed. II. i. 590 = capensis !
Hildebrandtii, *Reichb. f. Otia Bot. Hamb.* 78; *et in Bot. Zeit.* xxxix. (1881) 449.—Ins. Nossibé.
Humblotii, *Reichb. f. in Flora*, lxvii. (1883) 377.—Afr. austr.
Kerstenii, *Reichb. f. Otia Bot. Hamb.* 103.—Afr. trop.
Lindleyana, *Reichb. f. in Flora*, xlviii. (1865) 181.—Afr. trop.
M'Kenii, *Harv. Thes. Cap.* ii. 47, nomen.—Afr. austr.
Macowani, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 77.—Afr. austr.
Meirax, *Reichb. f. Otia Bot. Hamb.* 103.—Afr. trop.
micrantha, *Lindl. Gen. et Sp. Orch.* 370.—Afr. austr.
nanopurpurea, Bolus, in Journ. Linn. Soc. xx. (1884) 486 = purpurata.
neliglerensis, *Wight, Ic.* t. 1719.—Ind. or.
oppositifolia, *Sw. in Rees, Cycl.* xi. n. 6.—Ins. Mascar.
oxyglossa, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 76.—Afr. austr.
paludosa, *Harv. ex Lindl. in Hook. Lond. Journ. Bot.* i. (1842) 14.—Afr. austr.
purpurata, *Reichb. f. in Linnaea*, xli. (1877) 55.—Afr. austr.

DISPERIS:—
Reichenbachiana, *Weiw. ex Reichb. f. in Flora*, xlviii. (1865) 180.—Afr. trop.
secunda, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 220.—Afr. austr.
stenoplectron, *Reichb. f. Otia Bot. Hamb.* 102.—Afr. trop.
tenera, Spreng. ex Bolus, in Journ. Linn. Soc. xix. (1882) 346 = capensis.
tripetaloides, *Lindl. Gen. et Sp. Orch.* 371.—Ins. Borbon.
tripetaloides, Lindl. l. c. partim = zeylanica.
Tysoni, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 79.—Afr. austr.
villosa, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 220.—Afr. austr.
Wealii, *Reichb. f. Otia Bot. Hamb.* 103.—Afr. austr.
Woodii, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 78.—Afr. austr.
zeylanica, *Trim. in Journ. Bot.* xxiii. (1885) 245.—Ind. or.; Zeylan.

DISPERMA, J. F. Gmel. Syst. 885, 892 (1791) = **Mitchella**, Linn. (Rubiac.).
repens, J. F. Gmel. l. c. = M. repens.

DISPLASPIS, Klatt, in Linnaea, xxix. (1858) 711 = DIPLASPIS, Hook. f. = **Huanaca**, Cav. (Umbellif.).

DISPOROCARPA, A. Rich. Tent. Fl. Abyss. i. 307, in syn. (1847) = **Tyllaea**, Linn. (Crassulac.).
pentandra, Aschers. ex Schweinf. & Aschers. in Beitr. Fl. Aethiop. 271 = T. pentandra.
pharmacoides, A. Rich. Tent. Fl. Abyss. i. 307, in syn. = T. pharmacoides.

DISPOROPSIS, Hance, in Journ. Bot. xxi. (1883) 278. *LILIACEAE*.
fusco-picta, *Hance, l. c.*—China.

DISPORUM, Salisb. in Trans. Hort. Soc. i. (1812) 331. *LILIACEAE*, Benth. & Hook. f. iii. 681.
DRAPIEZIA, Blume, Enum. Pl. Jav. i. 8 (1827).
LETHEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2.
PROSARTES, D. Don, in Proc. Linn. Soc. i. (1839) 46; et in Trans. Linn. Soc. xviii. (1841) 531.
calcaratum, *D. Don, in Proc. Linn. Soc.* i. (1839) 45; *et in Trans. Linn. Soc.* xviii. (1841) 516.—Reg. Himal.
ceylanicum, Wight, Ic. t. 2049 = Leschenaultianum.
chinense, D. Don, Prod. Fl. Nep. 50 = pullum.
fuliens, D. Don, l. c. in obs. = pullum.
Hamiltonianum, D. Don, in Proc. Linn. Soc. i. (1839) 46; et in Trans. Linn. Soc. xviii. (1841) 517 = calcaratum.
Hookerii, *Nichols. Dict. Gard.* i. 484.—Calif.
Horsfieldii, D. Don, in Proc. Linn. Soc. i. (1839) 45 = pullum.
lanuginosum, *Nichols. Dict. Gard.* i. 485.—Am. bor. occ.
Leschenaultianum, *D. Don, in Proc. Linn. Soc.* i. (1839) 46; *et in Trans. Linn. Soc.* xviii. (1841) 518.—Ind. or.
Menziesii, *Nichols. Dict. Gard.* i. 485.—Calif.
multiflorum, D. Don, in Trans. Linn. Soc. xviii. (1841) 518 = pullum.
myosorense, Wight, Ic. t. 2049 = Leschenaultianum.
parviflorum, D. Don, Prod. Fl. Nep. 50 = pullum.
Pitcheri, D. Don, l. c. = pullum.
pullum, *Salisb. in Trans. Hort. Soc.* i. (1812) 331.—Ind. or.; Java; China.
Smithii, *D. Don, Prod. Fl. Nep.* 50.—Japon.
smilacinum, *A. Gray, in Perry, Exp. Exped.* ii. 321.—Japon.; Sibir.
trachycarpa, *Benth. & Hook. f. Gen.* iii. 683.—Am. bor. occ.
uniflorum, *Baker, in Journ. Bot.* xii. (1875) 230.—China.
Wallichii, D. Don, in Proc. Linn. Soc. i. (1839) 45; et in Trans. Linn. Soc. xviii. (1841) 516 = calcaratum.

DISQUAMIA, Lem. Jard. Fleur. iii. (1853–55) t. 269 = **Asoëmnes**, Knis & Pav. (Bromel.).

DISSANTHELIUM, Trin. in Linnaea, x. (1836) 305. *GRAMINEAE*, Benth. & Hook. f. iii. 1183.
PHALARIDIUM, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) (Pl. Meyen.) 161.

DISSANTHELIUM:—
STENOCHLOA, Nutt. in Journ. Acad. Philad. Ser. II. i. (1847) 169.
californicum, *Benth. in Hook. Ic. Pl.* t. 1275.—Calif.
sclerochloides, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 572.—Mexic.
supinum, *Trin. in Linnaea*, x. (1836) 305.—Chili.

DISSECOCARPUS, Hassk. Commel. Ind. 9 (1870) = DISECOCARPUS, Hassk. = **Commelina**, Linn.

DISSILIARIA, F. Muell. ex Baill. Adansonia, vii. (1867) 256. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 279.
CHORICERAS, Baill. Adansonia, xi. (1873) 119.
balaghioides, *F. Muell. ex Baill. Adansonia*, vii. (1867) 259.—Austral.
Muelleri, *Baill. l. c.* 259. t. 1.—Austral.
tricornis, *Benth. Fl. Austral.* vi. 91.—Austral.

DISSOCARPUS, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 75 = **Solerolaena**, R. Br. (Chenopod.).
biflorus, *F. Muell. l. c.* = S. biflora.

DISSOCHAETA, Blume, in Flora, xiv. (1831) 493. *MELASTOMACEAE*, Benth. & Hook. f. i. 758.
ancepa, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 70 = Anplectrum divaricatum.
annulata, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 58.—Malaya.
atrostricta, Miq. Fl. Ind. Bat. Suppl. 318 = pallida.
bancana, Miq. l. c. i. 1. 529 = celebica.
Barthei, Hance, ex Benth. Fl. Hongk. 115 = Barthea chinensis.
bibracteata, Baill. Hist. Pl. vii. 25 = Creochiton bibracteata.
biliguiata, Korth. in Verh. Nat. Gesch. Bot. 240 = Anplectrum biliguiatum.
bipulvinata, Korth. l. c. 239 = Marumia bipulvinata.
brachyanthera, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 74 = fusca.
bractsata, *Blume, in Flora*, xiv. (1831) 495.—Malaya.
bracteata, Korth. in Verh. Nat. Gesch. Bot. 237 = Korthalsii.
bracteosa, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 76 = bractsata.
caliothyrsa, *Miq. Fl. Ind. Bat.* i. 1. 528.—Java.
celebica, *Blume, Mus. Bot. Lugd. Bat.* i. 86.—Ins. Malay.
cinnamomea, Blume, l. c. = inappendiculata.
Cumingii, *Naud. in Ann. Sc. Nat. Sér.* III. xv. (1851) 75.—Ins. Philipp.
cyanocarpa, Blume, in Flora, xiv. (1831) 501 = Anplectrum glaucum.
cyanocarpa, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 87 = Anplectrum reformatum.
decipiens, Blume, l. c. 86 = monticola.
Diepenhorstii, Miq. Fl. Ind. Bat. Suppl. 317 = Omphalopus fallax.
divaricata, G. Don, Gen. Syst. ii. 758 = Anplectrum divaricatum.
fallax, Blume, in Flora, xiv. (1831) 493 = Omphalopus fallax.
fallax, Triana, in Trans. Linn. Soc. xxviii. (1871) 65 = gracilis.
fallax, Triana, l. c. = celebica.
fusca, *Blume, in Flora*, xiv. (1831) 497.—Java.
glauca, Blume, l. c. 501 = Anplectrum divaricatum.
gracilis, *Blume, l. c.* 495.—Malaya.
heteromorpha, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 78 = Dichaetanthera heteromorpha.
hirsuta, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 58.—Borneo.
inappendiculata, *Blume, in Flora*, xiv. (1831) 499.—Java.
intermedia, *Blume, l. c.* 498.—Malaya.
intermedia, Herb. Batav. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 54 = inappendiculata.
Korthalsii, *Miq. Fl. Ind. Bat.* i. 1. 530.—Sumatra.
leprosa, *Blume, in Flora*, xiv. (1831) 494.—Java.
liguiata, Blume, Mus. Bot. Lugd. Bat. i. 85 = Anplectrum liguiatum.
microcarpa, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 73 = celebica.
monticola, *Blume, in Flora*, xiv. (1831) 494.—Java.
nemorosa, *G. Don, Gen. Syst.* ii. 763.—Sumatra.
nodosa, Korth. in Verh. Nat. Gesch. Bot. 239 = Anplectrum nodosum.

DISSOCHAETA:—
ovalifolia, Nand. in Ann. Sc. Nat. Sér. III. xv. (1851) 74 = pallida.
palembanica, Miq. Fl. Ind. Bat. Suppl. 817 = Anplectrum divaricatum.
pallida, Blume, Mus. Bot. Lugd. Bat. i. 96. partim = Korthalsii.
pallida, *Blume, in Flora*, xiv. (1831) 500; *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 77.—Malaya.
papericarpa, Nand. l. c. 71 = Anplectrum divaricatum.
punctulata, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxvii. (1871) 53.—Malaya.
reformata, Blume, Mus. Bot. Lugd. Bat. i. 87 = Anplectrum reformatum.
reticulata, Blume, in Flora, xiv. (1831) 499 = Inappendiculata.
rostrata, Korth. Verh. Nat. Gesch. Bot. 239 = Anplectrum Korthalsii.
sagittata, *Blume, in Flora*, xiv. (1831) 500.—Java.
sarcotrica, Bail. Adansonia, xii. (1876-79) 88 (Medinillae sp.).—Madag.
spoliata, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 69. t. 4. f. 1 = Anplectrum glaucum.
superba, Naud. l. c. 77 = pallida.
vacillans, *Blume, in Flora*, xiv. (1831) 495.—Java.
velutina, *Blume, f. c.* 497.—Java.

DISSOLAENA, Lour. Fl. Cochinch. 137 (1790) = **Bauvolfia**, Plum. (Apocyn.).
verticillata, Lour. l. c.—China.

DISSOLENA, G. Don, Gen. Syst. iv. (1837) 99 = praec.

DISSOMERIA, Hook. f. ex Benth. in Hook. Niger Fl. 362 (1849). *SAMYDACEAE*, Benth. & Hook. f. i. 800.
crenata, *Hook. f. l. c.*—Afr. trop.

DISSOPETALUM, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 267 = **Cissampelos**, Linn. (Menisperm.).
mauritianum, Miers, l. c. 286 = C. mauritianum.

DISSORHYNCHIUM, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 434. t. 13 = **Habenaria**, Willd. (Orchid.).
muricatum, Schau. l. c. = H. muricata.

DISSOTHRIX, A. Gray, in Hook. Kew Journ. iii. (1851) 228. t. 5. *COMPOSITAE*, Benth. & Hook. f. ii. 242.
Gardneri, *A. Gray. l. c.*—Mexic.

DISSOTIS, Benth. in Hook. Niger Fl. 346 (1849). *MELASTOMACEAE*, Benth. & Hook. f. i. 746.
ARGYRELLA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 300. t. 6.
DERCHIPHIA, Rafin. Fl. Tellur. 101 (1838), nomen prius.
DUPINETA, Rafin. Sylva Tellur. 101 (1838).
HETEROTIS, Benth. in Hook. Niger Fl. 347 (1849).
LEIOCALYX, Planch. ex Hook. l. c. 350 (1849).
LEPIDANTHEMUM, Klotzsch, in Peters, Mossamb. Reise Bot. 64 (1861).
MELASTOMASTRUM, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 296. t. 5.
ORBECKIASTRUM, Naud. l. c. xiv. (1850) 118.
WEDDELIOPSIS, Planch. ex Benth. in Hook. Niger Fl. 351 (1849).
Afzelii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 449.—Afr. trop.
anisensis, *Hook. f. l. c.*—Afr. trop.
Barteri, *Hook. f. l. c.* 454.—Afr. trop.
cameronii, Hook. f. l. c. 453 = incana.
capitata, *Hook. f. l. c.* 449.—Afr. trop.
cornifolia, *Hook. f. l. c.* 448.—Afr. trop.
debilis, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 58. —Afr. austr.
decandra, *Triana, l. c.*—Afr. trop.
decemnervis, *Triana, l. c.*—Afr. trop.
eximia, *Hook. Gen. S. Afr. Pl.* ed. II. 114.—Afr. austr.
Fischeyanii, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 57.—Siam!
grandiflora, *Benth. in Hook. Niger Fl.* 347.—Afr. trop.
hirsuta, *Hook. f in Oliver, Fl. Trop. Afr.* ii. 450.—Afr. trop.

DISSOTIS:—
incana, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 58. —Afr. austr.
Irvingiana, *Hook. f. Bot. Mag.* t. 5149.—Afr. trop.
laevis, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 451.—Afr. trop.
Melleri, *Hook. f. l. c.*—Afr. trop.
multiflora, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 58.—Afr. occ.
petiolata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 448.—Afr. trop.
phaeotricha, *Harv. Gen. S. Afr. Pl.* ed. II. 114.—Afr. austr.
plumosa, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 452.—Afr. trop.
Princeps, Triana, in Trans. Linn. Soc. xxviii. (1871) 57 = eximia.
prostrata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 452; *Triana, l. c.* 58.—Afr. trop.
radicans, *Hook. f. l. c.* 450.—Afr. trop.
rotundifolia, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 58 = plumosa.
segregata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 448.—Afr. trop.
senegambiensis, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 58.—Afr. trop.
thesiifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 449.—Afr. trop.
tubulosa, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 58.—Afr. trop.
villosa, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 450.—Afr. trop.
villosa, Triana, in Trans. Linn. Soc. xxviii. (1871) 57 = Melastoma villosa.

DISTANDRA, Link, Enum. Hort. Berol. i. 254 (1821) = DISANDRA, Linn. = **Sibthorpia**, Linn. (Scrophul.).

DISTASIS, DC. Prod. v. 279 (1836) = **Chaetopappa**, DC. (Compos.).
concinna, Hook. & Arn. Bot. Beech. Voy. 350 = Erigeron concinnus.
heterophylla, Hemsl. Biol. Centr. Am. Bot. ii. 119 (Erigeron dub.).—Mexic.
modesta, DC. Prod. v. 279 = C. modesta.

DISTEGANTHUS, Lem. in Fl. des Serres, iii. (1847) t. 227. *BROMELIACEAE*, Benth. & Hook. f. iii. 661.
basilateralis, *Lem. l. c.*—Guiana.
scarlatinus, Nichols. Dict. Gard. i. 484 = Distiacanthus scarlatinus.

DISTEGIA, Rafin. New Fl. Am. iii. 20 (1836) = **Lonisera**, Linn. (Caprifol.).
sulans, Rafin. l. c. 21 = L. involucrata.

DISTEGOCARPUS, Sieb. & Zucc. (Fam. Nat. Fl. Jap. ii. 102. t. 3); in Abh. Acad. Muench. iv. iii. (1846) 226 = **Carpinus**, Linn. (Cupulif.).
carpinoides, Sieb. & Zucc. ll. cc. 116, 240 = C. japonica.
Carpinus, Sieb. & Zucc. ll. cc. 103, 227 = C. japonica.
cordatus, A. DC. in DC. Prod. xvi. ii. 128 = C. cordata.
erosa, A. DC. l. c. = C. erosa.
laxiflora, Sieb. & Zucc. Fl. Jap. Fam. Nat. ii. 104 = C. laxiflora.

DISTEIRA, Rafin. Fl. Tellur. iv. 68 (1836) = **Martynia**, Linn. (Pedalin.).
angulosa, Rafin. l. c. = M. diandra.

DISTEMMA, Leon. Hort. Van Houtte, fasc. ii. t. 11 (1847 ?) = DISEMMA, Labill. = **Passiflora**, Linn.
adiantifolium, Lem. Illustr. Hortic. (1867) Misc. 57 = P. glabra.
aurantiacum, Lem. l. c. = P. aurantia.
Bauerianum, Lem. l. c. = P. Bauerana.
carinatum, Lem. Hort. Van Houtte, fasc. ii. t. 11; et in Fl. des Serres (1847) 236, verso.—Mab.!
coccineum, Lem. Illustr. Hortic. (1867) Misc. 57 = P. Banksii.
eglandulosum, Lem. l. c. 56.—Austral.
Hahnii, Mast. in Mart. Fl. Bras. xiii. t. 569 = P. Hahnii.
Herbertianum, Lem. Illustr. Hortic. (1867) Misc. 57 = P. Herbertiana.

DISTEMON, Bonché, in Linnaea, xviii. (1844) 494 = **Canna**, Linn. (Scitamin.).
brasiliensis, Bonché, l. c. 495 = C. brasiliensis.
grandis, Horan. Prod. Scitam. 14 = C. brasiliensis.
Jacquinii, Bonché, in Linnaea, xviii. (1844) 495 = C. Jacquinii.
Kaussii, Bonché, l. c. nomen.—Bras.
Linkii, Bonché, l. c. = C. Linkii.
miniatus, Bonché, l. c. = C. miniata.
Ottonis, Bonché, l. c. 494 = C. Ottonis.
Roscoeanus, Bonché, l. c. 495 = C. denudata.

DISTEMON, Ehrenb. ex Aschers. in Monatsb. Acad. Berl. (1866) 581 = **Antisoharis**, Endl. (Scrophulariin.).
angustifolius, Ehrenb. ex Boiss. Fl. Orient. iv. 423 = A. linearis.
campanulatus, Ehrenb. & Hempr. ex Boiss. l. c. = A. arabica.
glandulosus, Ehrenb. & Hempr. ex Aschers. in Monatsb. Acad. Berl. (1866) 581 = A. glandulosa.

DISTEMON, Wedd. in Arch. Mus. Par. ix. (1856) (Monog. Urt.) 350. t. 20. *URTICACEAE*, Benth. & Hook. f. i. c.
grossum, Wedd. in DC. Prod. xvi. i. 238[sup] = Indicum.
indicum, *Wedd. Monog. Urt.* 351. t. 20.—Ind. or.

DISTEMONANTHUS, Benth. in Benth. & Hook. f. Gen. i. 578 (1865). *LEGUMINOSAE*, Benth. & Hook. f. i. c.
Benthamianus, *Baill. Hist. Pl.* ii. 185.—Afr. trop.
laxus, Oliver, Fl. Trop. Afr. ii. 282 = Benthamianus.

DISTEPHANA, Juss. in Ann. Mus. Par. vi. (1805) 396 = **Passiflora**, Linn.
candida, M. Roem. Syn. Pepon. 199 = P. candida.
citrifolia, M. Roem. l. c. = P. citrifolia.
cuneata, M. Roem. l. c. 200 = P. bicuspidata.
Fackeana, M. Roem. l. c. 199 = P. glandulosa.
glandulosa, Juss. in Ann. Mus. Par. vi. (1805) 396 = P. glandulosa.
pubescens, M. Roem. Syn. Pepon. 199 = P. quadriglandulosa.
quadridentata, M. Roem. l. c. = P. quadriglandulosa.
quadriglandulosa, M. Roem. l. c. = P. quadriglandulosa.
quitensis, M. Roem. l. c. 200 = Tacsonia mixta.
Rohriana, M. Roem. l. c. 199 = P. glandulosa.
spinosa, M. Roem. l. c. = P. spinosa.
stenpyana, M. Roem. l. c. = P. glandulosa.

DISTEPHANIA, Steud. Nom. ed. II. i. 521 (1840) = praec.

DISTEPHIA, Salisb. ex DC. Prod. iii. 335 (1828) = praec.

DISTEPHANUS, Cass. in Bull. Soc. Philom. (1817) 151 = **Vernonia**, Schreb. (Compos.).
capitatus, Boj. ex DC. Prod. v. 74.—Madag.
populifolius, Cass. in Bull. Soc. Philom. (1817) 151 = V. populifolia.
trinervis, Boj. ex DC. Prod. v. 75.—Madag.

DISTEREPTA, Rafin. Sylva Tellur. 126 (1838) = **Cassia**, Tourn. (Legumin.).
pilosa, Rafin. l. c. = C. pilosa.

DISTIACANTHUS, Linden, Cat. n. 28 (1869) 4. *BROMELIACEAE*.
erubescens, *Linden & André.*—*Cf. Illustr. Hortic.* xx. (1873) 171.—N. Granat.
scariatinus, *Linden, Cat.* n. 28 (1869) 4.—Bras.

DISTICHEIA, Rafin. Beitr. iv. 148 (1789) = **Brachypodium**, Beauv. (Gramin.).

DISTICHIA, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 128. *JUNCACEAE*, Benth. & Hook. f. iii. 866.
AGAPANTHA, Steud. in Bot. Zeit. xiv. (1856) 891.
GOUDOTIA, Decne. in Ann. Sc. Nat. Sér. III. iv. (1845) 84. t. 4.
OXYCLOS, Phil. Fl. Atac. 52. t. 6 (1860).
andina, Benth. & Hook. f. Gen. iii. 867 = macrocarpa.
brevifolia, *Phil. ex Benth. & Hook. f. l. c.*—Bolivia.

DISTICHIA:—
clandestina, *Buchen. in Bremen, Abh.* vi. (1879) 370. —Chili.
filamentosa, *Buchen. l. c.* 369.—Bolivia.
macrocarpa, *Weddl. ex Buchen. l. c.* iv. (1874) 128. —Bolivia.
muscoides, *Nees & Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 128.—Peruv.
tolimensis, *Benth. & Hook. f. Gen.* iii. 867.—N. Granat.

DISTICHIS, Thon. ex Lindl. Veg. Kingd. 181 (1847) (Cf. Thou. Orch. Isles Austr. Afr. Tabl. 8me) = **Liparis**, Rich. (Orchid.).

DISTICHLIS, Rafin. in Journ. Phys. lxxxix. (1819) 104. *GRAMINEAE*, Benth. & Hook. f. iii. 1193.
BRIZOPYRUM, J. & C. Presl, Rel. Haenk. 260 (1830).
DISCHLIS, Phil. Sert. Mend. Alt. 51 (1871), sphalm.
ammobia, *Phil. in Anal. Univ. Chil.* (1873) 569.—Chili.
condensata, *Hemsl. Biol. Centr. Am. Bot.* iii. 578.—Mexic.
hirta, *Phil. in Anal. Univ. Chil.* (1873) 570.—Chili.
maritima, *Rafin. in Journ. Phys.* lxxxix. (1819) 104. —Am. bor.
mendocina, *Phil. Sert. Mend. Alt.* 51.—Chili.
nodosa, Rafin. in Journ. Phys. lxxxix. (1819) 104 = Poa Michauxii.
prostrata, *E. Desv. in C. Gay, Fl. Chil.* vi. 398.—Am. trop.; Mexic.
thalassica, *E. Desv. l. c.* 397.—Mexic.; Chili.
Volckmanni, *Phil. in Anal. Univ. Chil.* (1873) 571.—Chili.

DISTICHMUS, Endl. Gen. 119 (1837) = DICHISMUS, Rafin. = **Scirpus**, Linn. (Cyperac.).

DISTICHOCALYX, T. Anders. ex Benth. & Hook. f. Gen. iii. 1080 (1876). *ACANTHACEAE*, Benth. & Hook. f. i. c.
thunbergiaeflora, *T. Anders. l. c.* 1081.—Afr. trop.

DISTICHOSTEMON, F. Muell. in Hook. Kew Journ. ix. (1857) 306. *SAPINDACEAE*, Benth. & Hook. f. i. 410.
hispidulum, Baill. Hist. Pl. v. 411 = phyllopterus.
phyllopterus, *F. Muell. in Hook. Kew Journ.* ix. (1857) 306.—Austral.

DISTICTIS, Mart. ex Meisn. Gen. 300 (1840) ; et ex DC. Prod. ix. 191 (1845). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1027.
MACRODISCUS, Bur. Monog. Big. 46. t. 11 (1864).
arthrerion, *DC. Prod.* ix. 191.—Bras.
lactiflora, *DC. l. c.*—Ins. S. Doming.
Mensoana, *Bur. in Rev. Hortic.* (1868) 154.—Bras.
rigescens, *DC. Prod.* ix. 191.—Ins. S. Doming.

DISTIMAKE, Rafin. Fl. Tellur. iv. 82 (1836) = **Ipomoea**, Linn. (Convolv.).
glaber, Rafin. l. c.=I. glabra.

DISTIMUM, Steud. Nom. ed. II. i. 521 (1840) = seq.

DISTIMUS, Rafin. in Journ. Phys. lxxxix. (1819) 105 = **Pycreus**, Beauv. (Cyperac.).
flavescens, Rafin. l. c. = P. flavescens.
flavicomus, Rafin. l. c. (= *C. flavicomus*, Vahl).—Am. bor

DISTOMAEA, Spenn. Fl. Friburg. i. 245 (1825) = **Listera**, R. Br. (Orchid.).
cordata, Spenn. l. c. 247 = L. cordata.
Nidus-avis, Spenn. l. c. 246 = Neottia Nidus-avis.
ovata, Spenn. l. c. = L. ovata.

DISTOMANTHERA, Turcz. in Bull. Soc. Nat. Mosc. xxxv. (1862) ii. 328. *SAXIFRAGEAE*:—
Cf. Benth. & Hook. f. i. 624.
Dombeyana, *Turcz. l. c.*—Am. austr.

DISTOMISCHUS, Dulac, Fl. Hautes-Pyr. 91 (1867) = **Festuca**, Tourn. (Gramin.).
ciliatus, Dulac, l. c. 92 = F. ciliata.
Myurus, Dulac, l. c. 91 = F. Myurus.
sciuroides, Dulac, l. c. = F. bromoides.
subuniglumis, Dulac, l. c. 92 = F. bromoides.

DISTREPTA, Miers, Trav. Chil. ii. 529 (1826) = **Tecophilaea**, Bert. (Hæmodorac.).
vaginata, Miers, l. c. = T. violaeflora.

DISTREPTUS, Cass. in Bull. Soc. Philom. (1817) 66 ; et Dict. Sc. Nat. xiii. 866 (1819) = **Elephantopus**, Linn. (Compos.).
angustifolius, Cass. in Dict. Sc. Nat. xiii. 868 = E. angustifolius.
crispus, Cass. l. c. lx. 601 = E. spicatus ?
nudiflorus, Cass. l. c. xiii. 868 ; Less. in Linnaea, iv. (1829) 269 = E. spicatus ?
replicatus, Cass. l. c. lx. 601 = E. spicatus.
spicatus, Cass. l. c. xiii. 867 = E. spicatus.
spiralis, Less. in Linnaea, vi. (1831) 690 = E. spicatus ?

DISTYLIS, Gaudich. in Freyc. Voy. Bot. 460. t. 80 (1826) = **Calogyne**, R. Br. (Goodeniac.).
Berardiana, Gaudich. l. c. = C. Berardiana.

DISTYLIUM, Sieb. & Zucc. Fl. Jap. i. 178. t. 94 (1895). *HAMAMELIDEAE*, Benth. & Hook. f. i. 666.
indicum, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 427.—Reg. Himal.
racemosum, *Sieb. & Zucc. Fl. Jap.* i. 179. t. 94.—China ; Japon.

DISYNANTHES, Reichb. Consp. 106 (1828) = seq.

DISYNANTHUS, Rafin. in Am. Monthly Mag. (1818) 268 = **Antennaria**, Gaertn. (Compos.).
plantaginens, Rafin. l. c. = A. plantaginea.

DISYNAPHEIA, Stend. Nom. ed. II. i. 521 (1840) = seq.

DISYNAPHIA, DC. Prod. vii. 267 (1838) = **Eupatorium**, Tourn. (Compos.).
montevidensis, DC. l. c. = E. calyculatum.

DISYNOMA, Rafin. Fl. Tellur. iii. 51 (1836) = **Aethionema**, R. Br. (Crucif.).
cornea, Rafin. l. c. = A. eristatum.

DITASSA, R. Br. in Mem. Wern. Soc. i. (1809) 49. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 767.
alborubra, *Fourn. in Mart. Fl. Bras.* vi. iv. 240.—Bras.
acerosa, *Mart. Nov. Gen. et Sp.* i. 53.—Bras.
adnata, *Fourn. in Mart. Fl. Bras.* vi. iv. 256.—Bras.
angustifolia, *Decne. in DC. Prod.* viii. 575.—Guiana.
anomala, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 99.—Bras.
arietata, *Benth. ex Fourn. in Mart. Fl. Bras.* vi. iv. 247.—Bras.
Benkei, *Roem. & Schult. Syst.* vi. 112.—Bras.
barbata, *Fourn. in Mart. Fl. Bras.* vi. iv. 255.—Bras.
bicolor, *Decne. in DC. Prod.* viii. 575.—Bras.
Blanchetti, *Decne. l. c.*—Bras.
bonariensis, *Decne. l. c.* 577.—Bras.
Burchellii, *Hook. f. Arn. in Hook. Journ. Bot.* i. (1834) 295.—Bras.
campestris, *Griseb. in Gott. Abh.* xxiv. (1879) 228.—Arg. Argent.
capillaris, *Fourn. in Mart. Fl. Bras.* vi. iv. 253.—Bras.
congesta, *Fourn. l. c.* 245.—Bras.
consanguinea, *Decne. in DC. Prod.* viii. 575.—Bras.
crassifolia, *Decne. l. c.* 576.—Bras.
crassinervia, *Decne. l. c.* 577.—Bras.
cucullata, *Fourn. in Mart. Fl. Bras.* vi. iv. 256.—Bras.
decussata, *Mart. Nov. Gen. et Sp.* i. 51. t. 31.—Bras.
dichotoma, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 317.—Guiana.
ericoides, *Decne. in DC. Prod.* viii. 576.—Bras.
eximia, *Decne. l. c.* 574.—Bras.
fallax, *Fourn. in Mart. Fl. Bras.* vi. iv. 248.—Bras.
fasciculata, *Fourn. l. c.* 253.—Bras.
fulva, *Fourn. l. c.* 248.—Bras.
Gardneri, *Decne. in DC. Prod.* viii. 576.—Bras.
glaucescens, *Decne. l. c.*—Guiana.
glaziovii, *Fourn. in Mart. Fl. Bras.* vi. iv. 250.—Bras.
grandiflora, *Fourn. l. c.* 255.—Bras.
Guilleminiana, *Decne. in DC. Prod.* viii. 574.—Bras.
hastata, *Decne. l. c.* 575.—Bras.

DITASSA:—
hemipogonoides, *Fourn. in Mart. Fl. Bras.* vi. iv. 238. —Bras.
Hilariana, *Fourn. l. c.* 254.—Bras.
imbricata, *Fourn. l. c.* 240.—Bras.
Klotzschii, *Fourn. l. c.* 247.—Bras.
laevis, *Mart. Nov. Gen. et Sp.* i. 53.—Bras.
lagoinsis, *Fourn. in Mart. Fl. Bras.* vi. iv. 256.—Bras.
lanceolata, *Decne. in DC. Prod.* viii. 576.—Peruv.; Bras.
linearis, *Mart. Nov. Gen. et Sp.* i. 53.—Bras.
longiloba, *Benth. Fl. Hartw.* 215.—N. Granat.
manicata, *Decne. in DC. Prod.* viii. 575.—Bras.
micromeria, *Decne. l. c.* 576.—Bras.
montana, *Decne. l. c.* 576.—Bras.
mucronata, *Mart. Nov. Gen. et Sp.* i. 52.—Bras.
myrtilloides, *Fenzl, ex Fourn. in Mart. Fl. Bras.* vi. iv. 251.—Bras.
Niruri, *Decne. in DC. Prod.* viii. 576.—Bras.
nitida, *Fourn. in Mart. Fl. Bras.* vi. iv. 241.—Bras.
oboculata, *Mart. Nov. Gen. et Sp.* i. 53.—Bras.
oxypetala, *Decne. in DC. Prod.* viii. 574.—Bras.
oxyphylla, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 260 = lanceolata.
passerinioides, Decne. in DC. Prod. viii. 578 = virgata.
passerinoides, *Mart. Nov. Gen. et Sp.* i. 53.—Bras.
pauciflora, *Decne. in DC. Prod.* viii. 577.—Bras.
Poeppigii, *Fourn. in Mart. Fl. Bras.* vi. iv. 244.—Bras
Pohliana, *Fourn. l. c.*—Bras.
praecincta, *Fourn. l. c.* 253.—Bras.
ramosa, *Fourn. l. c.* 246.—Bras.
reflexa, *Fourn. l. c.* 255.—Bras.
retusa, *Mart. Nov. Gen. et Sp.* i. 53.—Bras.
Riedelii, *Fourn. in Mart. Fl. Bras.* vi. iv. 247.—Bras.
rufescens, *Decne. in DC. Prod.* viii. 574.—Bras.
Salzmanni, *Fourn. in Mart. Fl. Bras.* vi. iv. 252.—Bras.
subtrivialis, *Griseb. in Lechl. Berb. Am. Austr.* 58.—Peruv.
taxifolia, *Decne. in DC. Prod.* viii. 578.—Guiana.
trivialis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 68. t. 277.—Peruv.
umbellata, *Decne. in DC. Prod.* viii. 574.—Bras.
velutina, *Fourn. in Mart. Fl. Bras.* vi. iv. 245.—Bras
venosa, *Decne. ex Fourn. l. c.* 249.—Bras.
virgata, *Fourn. l. c.* 236.—Bras
Warmingii, *Fourn. l. c.* 242.—Bras.

DITAXIA, Endl. Gen. 670 (1839), sphalm. = DITOXIA, Rafin. = **Celsia**, Linn. (Scrophulariac.).

DITAXIS, Vahl, ex A. Juss. Euph. Tent. 37. t. 7 (1824) = **Argithamnia**, Sw. (Euphorb.).
brasiliensis, Baill. Adansonia, iv. (1863–64) 269 = A. brasiliensis.
castaneaefolia, Baill. l. c. 274 = Caperonia castaneaefolia.
chiropetala, Bert. ex Baill. Férussac, xx. (1836) 110 = A. Bertero.
chrysantha, Klotzsch, in Wiegm. Archiv, vii. (1841) 199, nomen = A. Simoniana.
cordata, Baill. Adansonia, iv. (1863–64) 272 = Caperonia cordata.
diocia, H. B. & K. Nov. Gen. et Sp. vii. 170. t. 689 = A. diocia.
fascicularis, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 582 = A. Simoniana.
fusciculata, Vahl, ex A. Juss. Euph. Tent. 110 = A. fascicularis.
glabella, Griseb. in Goett. Abh. vii. (1857) 171 = A. lanciflola.
harmidandra, Griseb. Fl. Brit. W. Ind. 44 = Ricinella hasmidandra.
heteranthe, Zucc. in Abh. Akad. Muench. i. (1829–30) 290 = A. heteranthe.
Hilariana, Baill. Adansonia, iv. (1863–64) 270 = A. Simoniana.
illimanicnsis, Baill. Etud. Gén. Euph. 299 = A. illimanicnsis.
lanciflola, Schlecht. in Linnaea, xxvi. (1853) 636 = A. lanciflola.
linearifolia, Baill. Adansonia, iv. (1863–64) 273 = Caperonia linearifolia.
melochiaefolia, Baill. l. c. 270 = A. Simoniana.
polymorpha, Baill. l. c. 278 = Caperonia heteropetala, heteropetaloides.
Simoniana, Casar. Nov. Stirp. Bras. Dec. x. 87 = A. Simoniana.

DITAXIS:—
trinervia, Baill. Adansonia, iv. (1868–64) 271, sphalm. = seq.
trifoliinervia, Klotzsch, in Wiegm. Arobiv, vii. (1841) 199, nomen = A. montevidensis.

DITELESIA, Rafin. Fl. Tellur. iii. 69 (1836) = Commelina, Linn.
nudiflora, Rafin. l. c. = C. nudiflora.

DITERELA, Rafin. Fl. Tellur. iv. 84 (1836) = Convolvulus, Tourn.
parviflora, Rafin. l. c. = C. parviflorus.

DITHECA, Miq. Fl. Ind. Bat. i. 615 (1855) = Ammannia, Linn. (Lythrar.).
debilis, Miq. l. c. = A. multiflora.
densiflora, Miq. l. c. = A. pentandra.
verticillata, Miq. l. c. = A. lanceolata.

DITHRICHUM, DC. Prod. v. 619 (1836) = DITRICHUM, Cass. = Verbesina, Linn. (Compos.).

DITHYRAEA, Endl. Gen. Suppl. v. 35 (1850) = seq.

DITHYREA, Harv. in Hook. Lond. Journ. Bot. iv. (1845) 77. t. 5 = Biscutella, Linn. (Crucif.).
californica, Harv. l. c. = B. californica.
Wislizenii, Engelm. in Wisliz. Tour. N. Mexico, 95 = B. Wislizenii.

DITHYRIA, Benth. in Hook. Journ. Bot. ii. (1840) 89 = Swartzia, Schreb. (Legumin.).
alterna, Benth. l. c. = S. alterna.
mollis, Benth. l. c. = S. mollis.

DITHYROCARPUS, Kunth, in Ber. Acad. Berl. (1841) 245; et Enum. Pl. iv. 76 (1843) = Floscopa, Lour. (Commelin.).
capensis, Kunth, Enum. Pl. iv. 78 = F. glomerata.
glabratus, Kunth, l. c. 77 = F. glabrata.
glandulosus, Seub in Mart. Fl. Bras. iii. 1. 255 = F. glabrata.
glomeratus, Kunth, Enum. Pl. iv. 78 = F. glomerata.
hirsutus, Kunth, l. c. 77 = F. glabrata.
Meyenianus, Kunth, l. c. 78 = F. scandens.
paniculatus, Kunth, l. c. 79 = F. scandens.
petiolatus, Wight, Ic. t. 2079 = F. scandens.
robustus, Seub. in Mart. Fl. Bras. iii. 1. 255 = F. robusta.
Rothii, Wight, Ic. t. 2079 = F. scandens.
rufus, Kunth, Enum. Pl. iv. 79 = F. scandens.
Schomburghianus, Kunth, l. c. 662 = Ancilema ovato-oblongum.
stemorhachys, Klotzsch, ex C. B. Clarke, in DC. Monog. Phan. iii. 226 = Ancilema ovato-oblongum.
undulatus, Wight, Ic. t. 2079 = F. scandens.

DITHYROSTEGIA, A. Gray, in Hook. Kew Journ. iii. (1851) 100 = Angianthus, Wendl. (Compos.).
amplexicaulis, A. Gray, l. c. = A. amplexicaulis.

DITMARIA, Spreng. Anleit. ii. 11. 704 (1818); et Syst. i. 4 (1825) = Brisma, Rudge (Vochysiac.).
floribunda, Spreng. Syst. i. 18 = E. floribundum.
violacea, Spreng. l. c. iv. Cur. Post. 8 = E. violaceum.

DITOCA, Banks, ex Gaertn. Fruct. ii. 106. t. 126 (1791) = Solenanthus, Linn. (Illecebrac.).
muscosa, Gaertn. l. c. = S. bifforus.

DITOMOSTROPHE, Turcz. in Bull. Soc. Nat. Mosc. (1846) ii. 498 = Guichenotia, J. Gay (Sterculiac.).
angustifolia, Turcz. l. c. 499 = G. semihastata.

DITOXIA, Rafin. Préc. Découv. 39 (1814); et in Desv. Journ. Bot. iv. (1814) 270 = Calsia, Linn. (Scrophul.).
betonicaefolia, Rafin. l. c. 40 = C. betonicaefolia.
cretica, Rafin. ex Desv. l. c. 270 = C. cretica.
lyrata, Rafin. Préc. Découv. 40 = C. cretica.

DITREMEXA, Rafin. Sylva Tellur. 127 (1838) = Cassia, Tourn. (Legumin.).
corallodendron, Rafin. l. c. = C. occidentalis.
fetida, Rafin. l. c. = C. occidentalis.
ligustrina, Rafin. l. c. = C. ligustrina.

DITRICHUM, Cass. in Bull. Soc. Philom. (1817) 33 = Verbesina, Linn. (Compos.).
macrophyllum, Cass. l. c. = V. diversifolia.

DITRICLITA, Rafin. New Fl. Am. i. 63, nomen (1836) = Saxifraga, Linn.

DITRISYNIA, Rafin. Sylva Tellur. 64 (1838) = Stillingia, Garden (Euphorb.).
ligustrina, Rafin. l. c. = S. ligustrina.

DITRITA, Rafin. Sylva Tellur. 115 (1838) = Euphorbia, Linn.
hirta, Rafin. l. c. = E. pilulifera.
obliqua, Rafin. l. c. = E. hypericifolia.
rotundifolia, Rafin. l. c. = E. Chamaesyce.

DITROCHE, E. Mey. ex Moq in DC. Prod. xiii. ii. 19 (1849) = Semonvillea, J. Gay (Ficoid.).
furcata, E. Mey. ex Moq. l. c. 20 = S. fenestrata.

DITRYSINIA, Rafin. Neogenet. 2 (1825) = Sebastiania, Spreng. (Euphorbiac.).
ligustrina, Rafin. l. c. = S. ligustrina.
sylvatica, Rafin. l. c. = S. ligustrina.

DITTA, Grisb. in Mem. Arn. Acad. N. S. viii. (1861) 160 (Pl. Wright. Cub.). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 385.
myricoides, *Griseb. l. c.*—Cuba.

DITTELASMA, Hook. f. in Benth. & Hook. f. Gen. i. 895 (1862). *SAPINDACEAE*, Benth. & Hook. f. l. c.
Rarak, *Hook. f. l. c.* 896.—Malacca.

DITTOCERAS, Benth. & Hook. f. in Hook. Ic. Pl. t. 1422 (1888) *ASCLEPIADEAE*.
Andersoni, *Hook. f. l. c.*—Reg. Himal.

DITTOSTIGMA, Phil. Sert. Mend. Alt. 36 (1871). *SOLANACEAE*, Benth. & Hook. f. ii. 906.
mendocinum, Phil. l. c.—Chili.

DITUILIS, Rafin. Fl. Tellur. iv. 49 (1836) = Liparis, Rich. (Orchid.).
nepalensis, Rafin. l. c. = L. bituberculata.

DITULIMA, Rafin. Fl. Tellur. iv. 41 (1836) = Dendrobium, Sw. (Orchid.).
anceps, Rafin. l. c. = D. anceps.

DIURIS, Sm. in Trans. Linn. Soc. iv. (1798) 222. *ORCHIDEAE*, Benth. & Hook. f. iii. 608.
abbrevinta, *F. Muell. ex Benth. Fl. Austral.* vi. 329.—Austral.
aequalis, F. Muell. ex Benth. l. c. 328 = maculata.
alba, *R. Br. Prod.* 316.—Austral.
aurea, *Sm. Exot. Bot.* i. 15. t. 8.—Austral.
Behrii, Schlecht. in Linnaea, xx. (1849) 572 = pedunculata.
carinata, Lindl. Gen. et Sp. Orch. 510 = setacea.
corymbosa, Lindl. Swan River App. 51 = longifolia.
curvifolia, Lindl. Gen. et Sp. Orch. 507 = maculata.
dendrobioides, *Fitzg. Austral. Orch.* i. VII. (1882).—Austral.
Drummondii, Lindl. Swan River App. 51 = emarginata.
elongata, Sw. in Schrad. N. Journ. i. (1805) 89 = punctata.
emarginata, *R. Br. Prod.* 316.—Australia.
filifolia, Lindl. Swan River App. 51 = setacea.
Fitzna, *Kidl in Forbes, Nat. Wand. East. Archip.* 219.—Ins. Timor.
laertis, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.
lanceolata, Lindl. Gen. et Sp. Orch. 508 = pedunculata.
lanifera, Lindl. Swan River App. 51 = emarginata.
lilacina, F. Muell. in Linnaea, xxvi. (1853) 289 = punctata.
longifolia, *R. Br. Prod.* 316.—Austral.
maculata, *Sm. Exot. Bot.* i. 57. t. 30.—Austral.
Novae-Zelandiae, A. Rich. Fl. Nov. Zel. 163. t. 25 = Orthoceras strictum.
oculata, F. Muell. Fragm. v. 178 = aurea.
oculata, F. Muell. in Linnaea, xxvi. (1853) 241 = sulphurea.
pallens, *Benth. Fl. Austral.* vi. 329.—Austral.
palustris, *Lindl. Gen. et Sp. Orch.* 507.—Austral.
pardina, Lindl. l. c. = maculata.
pauciflora, *R. Br. Prod.* 316.—Austral.
pedunculata, *R. Br. l. c.*—Austral.
porrifolia, Lindl. Swan River App. 51 = longifolia.
punctata, *Sm. Exot. Bot.* i. 13. t. 8.—Austral.

DIURIS:—
secundiflora, *Fitzg. Austral. Orch.* i. IV. (1877).—Austral.
setacea, *R. Br. Prod.* 316.—Austral.
spathulata, Sw. in Schrad. N. Journ. i. (1805) 60 = aurea.
sulphurea, *R. Br. Prod.* 316.—Austral.
tricolor, *Fitzg. in Journ. Bot.* xxiii. (1885) 157. —Austral.

DIUROGLOSSUM, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 157 = Guasuma, Plum. (Sterculiac.).
rufescens, Turcz. l. c. = G. tomentosa.

DIUROSPERMUM, Edgew. in Proc. Linn. Soc. i. (1847) 251 = Utricularia, Linn. (Lentibul.).
album, Edgew. l. c. = U. kumaonensis.

DIZONIUM, Willd. ex Schlecht. in Linnaea, v. (1830) 411, in nota = Geigeria, Griesel. (Compos.).
longifolium, Willd. l. c. = G. africana.

DIZYGANDRA, Meisn. Gen. 294 (1840) = Ruellia, Plum. (Acanthac.).
noctiflora, Shuttlew. ex Nees, in DC. Prod. xi. 128 = R. noctiflora.

DOBERA, Jus. Gen. 425 (1789). *SALVADORACEAE*, Benth. & Hook. f. ii. 680; iii. 162, in obs.
SCHIZOCALYX, Hochst. in Flora, xxvii. (1844) Beil. 1.
TOMEX, Forsk. Fl. Aegypt. Arab. 33 (1776), nomen prius.
coriacea, A. DC. in DC. Prod. xvii. 81 = Roxburghii.
dubia, *Steud. Nom.* ed. II. i. 521.—Africa ?
glabra, A. DC. in DC. Prod. xvii. 81 = Roxburghii.
Roxburghii, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 191.—Ind. or.; Afr. austr.

DOBERA, Rafin. Fl. Tellur. iii. 55 (1836) = Rivina ? Plum. (Phytolac.).

DOBINAEA, Spreng. Syst. iii. 890 (1826) = seq.

DOBINEA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 249 (1825). *SAPINDACEAE*, Benth. & Hook. f. i. 410.
vulgaris, *Buch.-Ham. l. c.*—Reg. Himal.

DOBROWSKIA, A. DC. in DC. Prod. vii. 255 (1839) = seq.

DOBROWSKYA, Presl, Prod. Lobel. 8 (1836) = Lobelia, Linn.
anceps, Presl, l. c. 10 = L. anceps.
aspera, DC. Prod. vii. 356 = L. aspera.
Dregeana, Presl, Prod. Lobel. 9 = L. scabra.
Eckloniana, Presl, l. c. = L. aspera.
laxicaulis, Presl, l. c. 10 = L. scabra.
Massoniana, Presl, l. c. = L. tenella.
polyphylla, Presl, l. c. = L. scabra.
scabra, A. DC. in DC. Prod. vii. 356 = L. scabra.
serratifolia, A. DC. l. c. 356 = L. deblis.
stellarioides, Presl, Prod. Lobel. 10 = L. stellarioides.
stricta, Presl, ex Eckl. & Zeyh. Enum. 398 = L. tenella.
tenella, Sond. in Harv. & Sond. Fl. Cap. iii. 549 = L. tenella.
Thunbergiana, Presl, Prod. Lobel. 9 = L. scabra.
unidentata, A. DC. in DC. Prod. vii. 355 = L. tenella.
Zeyheriana, Presl, Prod. Lobel. 10 = L. scabra.

DOCHAFA, Schott, Syn. Aroid. 24 (1856) = Arisaema, Mart. (Aroid.).
flava, Schott, l. c. = A. flavum.

DOCYNIA, Decne. in Nouv. Arch. Mus. Par. x. (1874) 131. *ROSACEAE*.
Griffithiana, Decne. l. c. t. 14 = indica.
Hookeriana, *Decne. l. c.* t. 15.—Reg. Himal.
indica, *Decne. l. c.* t. 14.—Reg. Himal.; Burma.

DODARTIA, [Tourn.] Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 948.
indica, Linn. Sp. Pl. 633 = Lindenbergia polyantha.
Linaria, Mill. Gard. Dict. ed VIII. n. 2 = Anarrhinum bellidifolium.
orientalis, *Linn. Sp. Pl.* 633.—As. occ. et bor.
virgata, Moench, Meth. 446 = orientalis.

DODECADENIA, Nees, in Wall. Fl. As. Rar. ii. 61 (1831); Laurin. Expos. 19 (1888). *LAURINEAE*, Benth. & Hook. f. iii. 160.
grandiflora, *Nees, in Wall. l. c.* 63.—Reg. Himal.
robusta, Zoll. & Moritzi, Syst. Verz. Zoll. 71 = Litsea sebifera, *Blume*.

DODECADIA, Lour. Fl. Cochinch. 318 (1790). *SAMYDACEAE!* (Gen. dub.) Benth. & Hook. f. i. 1007.
agrestis, *Lour. l. c.* 319.—China.

DODECAS, Linn. f. Suppl. 86, 245 (1781) = **Crenea**, Aubl. (Lythrar.).
maritimus, Griseb. Fl. Brit. W. Ind. 270 = C. maritima.
surinamensis, Linn. f. Suppl. 245 = C. repens.

DODECASPERMA, Rafin. Fl. Tellur. iv. 35 (1836) = **Bomarea**, Mirb. (Amaryll.).
acutifolia, Rafin. l. c. = B. acutifolia.

DODECASPERMUM, Forst. ex Scop. Introd. 219 (1777) (Gen. dub.).

DODECASTEMON, Haask. in Versl. en Med. Kon. Akad. Wetensch. iv. (1855) 141 = **Cyclostemon**, Blume (Euphorbiac.).
Teysmanni, Haask. l. c. = C. Teysmanni.

DODECATHEON, Linn. Dis. Chen. 25 (1751); Amoen. Acad. iii. 16 (1756). *PRIMULACEAE*, Benth. & Hook. f. ii. 634.
MEADIA, Catesb. ex Mill. Gard. Dict. ed. VI. App. 175 (1759).
angustifolium, Rafin. in Am. Monthly Mag. (1818) 40 = D. Meadia.
californicum, A. Murr. Oreg. Circ. 1 = D. Meadia.
cordatum, *crenatum*, *cuneatum*, *elliptic-um*, *flexuosum*, *longifolium*, *obovatum*, *obtusum*, *ovatum*, *parviflorum*, *parvifolium*, *serratum*, *triflorum*, *undatum*, *uniflorum*, Rafin. Act. Journ. 179, 180 = D. Meadia.
dentatum, Hook. Fl. Bor. Am. ii. 119 = D. Meadia.
elliptic-um, Nutt. ex Durand, in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 94 = D. Meadia.
frigidum, Cham. & Schlecht. in Linnaea, i. (1826) 217 = D. Meadia.
integrifolium, Michx. Fl. Bor. Am. i. 123 = D. Meadia.
Jeffreyi, Hort. Fl. des Serres, Sér. II. vi. (1865–67) 99 = D. Meadia.
Meadia, *Linn. Sp. Pl.* ed. I. 144.—Am. bor. et arct.
reflexum, Salisb. Prod. 118 = D. Meadia.

DODONAEA, Linn. Gen. ed. I. 341 (1737). *SAPINDACEAE*, Benth. & Hook. f. i. 410.
EREPLEURISMA, Barth. in Lehm. Pl. Preiss. ii. 228 (1847).
acerosa, Lindl. in Mitch. Journ. Trop. Austral. 273 = filifolia.
adenophora, *Miq. in Linnaea*, xviii. (1844) 95.—Austral.
adenophora, F. Muell. Fragm. i. 98 = concinna.
angustifolia, *Blanco*, *Fl. Filip.* ed. I. 312.—Ins. Philipp.
angustifolia, Linn. f. Suppl. 218 = viscosa.
angustifolia, Thunb. Prod. Pl. Cap. 77 = Thunbergiana.
angustissima, *Hort. ex DC. Prod.* i. 617.—Hab.?
aptera, *Miq. in Lehm. Pl. Preiss.* i. 225.—Austral.
arabica, Hochst. & Steud. in Flora, xxiv. (1841) 2. Intell. 38 = viscosa.
asplenifolia, Rudge, in Trans. Linn. Soc. xi. (1815) 397, t. 30 = viscosa.
attenuata, *A. Cunn. in Field, N. S. Wales*, 358.—Austral.
Baueri, *Endl. in Enum. Pl. Hueg.* 13.—Austral.
bicolata, H. B. & K. Nov. Gen. et Sp. v. 134. t. 448 = viscosa.
boroniaefolia, *G. Don, Gen. Syst.* i. 674.—Austral.
brasiliensis, *Schlecht. in Linnaea*, xviii. (1844) 85 (err. typ. 51).—Bras.
Burmanniana, DC. Prod. i. 616 = viscosa.
burmanniifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 8.—Austral.
cuspa, Eckl. & Zeyh. Enum. 55 = Combretum salicifolium.

DODONAEA:—
Caleyana, G. Don, Gen. Syst. i. 674 = boroniaefolia.
calycina, Cunn. ex A. Gray, Bot. U. St. Expl. Exped. i. 262 = truncatialis.
Candolleana, *Blume, Rumphia*, iii. 190.—Malaya.
ceratocarpa, *Endl. in Enum. Pl. Hueg.* 13.—Austral.
cestroides, Sieber, ex Schlecht. in Linnaea, xvii. (1843) 640 = triquetra.
concinna, *Benth. Fl. Austral.* i. 487.—Austral.
conferta, G. Don, Gen. Syst. i. 674 = viscosa.
conglomerata, Eckl. & Zeyh. Enum. 55 = Combretum salicifolium.
cuneata, *Rudge, in Trans. Linn. Soc.* xi. (1813) 396. t. 19.—Austral.
defexa, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 8 = Baueri.
denticulata, F. Muell. Fragm. i. 96 = ptarmicifolia.
dioica, Roxb. Hort. Beng. [38]; Fl. Ind. ii. 256 = viscosa.
divaricata, *Benth. Fl. Austral.* i. 481.—Austral.
Dominiana, *Blume, Rumphia*, iii. 189.—Peruv.
dubia, Eckl. & Zeyh. Enum. 55 = Combretum salicifolium.
Ehrenbergii, *Schlecht. in Linnaea*, xviii. (1844) 86 (err. typ. 52).—Ins. S. Doming.
elaeagnoides, *Rudolph, in Schrad. N. Journ.* ii. 11. (1807) 292.—Ins. S. Doming.
ericaefolia, *G. Don, Gen. Syst.* i. 674.—Austral.
ericoides, *Miq. in Lehm. Pl. Preiss.* i. 227.—Austral.
erioscypa, *Sm. in Rees, Cycl.* xii. n. 5.—Ins. Sandvic.
filifolia, *Hook. in Mitch. Journ. Trop. Austral.* 241.—Austral.
filiformis, Hort. Angl. ex Link, Enum. Hort. Berol. i. 981 = ericaefolia ?
filicifolia, F. Muell. Fragm. ii. 182 = larreoides.
Forsteri, *Montr. in Mém. Acad. Lyon*, x. (1860) 189.—N. Caled.
heterophylla, Hort. ex G. Don, Gen. Syst. i. 674 = viscosa.
hexandra, *F. Muell. in Trans. Vict. Inst.* i. (1855) 117.—Austral.
hirtella, Miq in Linnaea, xviii. (1844) 94 = boroniaefolia.
hispidula, Endl. Atakt. t. 80 = Distichostemon phyllopterus.
humifusa, *Miq. in Lehm. Pl. Preiss.* i. 226.—Austral.
humilis, *Endl. Atakt.* t. 31.—Austral.
illita, *F. Muell. ex Regel, Ind. Sem. Hort. Petrop.* (1856) 36.—Austral.
inaequifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 408.—Austral.
jamaicensis, DC. Prod. i. 616 = viscosa.
Kingii, G. Don, Gen. Syst. i. 674 = viscosa.
Kobautiana, *Schlecht. in Linnaea*, xiii. (1839) 52.—Afr. trop.
Lamponga, *Miq. Fl. Ind. Bat. Suppl.* i. 511.—Sumatra.
lanceolata, *F. Muell. Fragm.* i. 78.—Austral.
larreaoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 408.—Austral.
latifolia, Salisb. Prod. 276 = viscosa.
lanuriolia, Sieber, ex Lond. Hort. Brit. 154 = seq.?
laurina, Sieber, ex Spreng. Syst. iv. Cur. Post. 153 = triquetra.
leptocarpa, F. Muell. Fragm. i. 219 = inaequifolia.
Lindleyana, F. Muell. Pl. Vict. i. 88 = triangularis.
linearifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 407.—Cuba.
linearis, E. Mey. ex Harv. & Sond. Fl. Cap. i. 249 = Thunbergiana.
lobulata, *F. Muell. in Linnaea*, xxv. (1852) 372.—Austral.
longipes, G. Don, Gen. Syst. i. 674 = triquetra.
lucida, Moench, Meth. 356 = viscosa.
Macgillivrayi, *F. Muell. & Scortechini, in Melb. Chemist*, (Jan. 1882).—Austral.
macrocypa, *F. Muell. Fragm.* iv. 165.—Austral.
madagascariensis, *Radlk. in Bremen, Abh.* viii. (1884) 470.—Madag.
megacarpa, *F. Muell. ex Benth. Fl. Austral.* i. 483.—Austral.
microcarpa, DC. Prod. i. 617 = viscosa.
microcypa, *F. Muell. Ann. Rep. Melb. Bot. Gard.* (1869) 13.—Austral.
mollis, Lindl. in Mitch. Journ. Trop. Austral. 212 = triangularis.
multijuga, *G. Don, Gen. Syst.* i. 674.—Austral.
multijuga, F. Muell. Fragm. i. 218 = larreoides.

DODONAEA:—
Mundiana, Eckl. & Zeyh. Enum. 55 = Thunbergiana
natalensis, Sond. in Linnaea, xxiii. (1850) 28 = viscosa.
nervifolia, A. Cunn. ex A. Gray, Bot. U. St. Expl. Exped. i. 262 = viscosa.
nutans, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 407.—Arab.
oblongifolia, Link, Enum. Hort. Berol. i. 351; et in Bot. Reg. t. 1051 = viscosa.
ovata, Dum.-Cours. Bot. Cult. ed. II. vii. 337 = viscosa.
oxyptera, *F. Muell. in Hook. Kew Journ.* ix. (1857) 197; *Fragm.* i. 74.—Austral.
pallida, Miq. Anal. Bot. Ind. iii. 7 = viscosa.
pauciflorifolia, A. Cunn. ex Steud. Nom. ed. II. 522 = vestita.
peduncularis, *Lindl. in Mitch. Journ. Trop. Austral* 261.—Austral.
pentandra, Griff. Notul. iv. 548 = viscosa.
petiolaris, *F. Muell. Fragm.* iii. 13.—Austral.
physocarpa, *F. Muell. in Hook. Kew Journ.* ix. (1857) 197; *Fragm.* i. 74.—Austral.
pinifolia, *Miq. in Lehm. Pl. Preiss.* i. 227.—Austral.
pinnata, *Sm. in Rees, Cycl.* xii. n. 7.—Austral.
platyptera, *F. Muell. Fragm.* i. 78.—Austral.
polyzyga, *F. Muell. l. c.* 74.—Austral.
Preissiana, Miq. in Lehm. Pl. Preiss. i. 226 = attenuata.
procumbens, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 8.—Austral.
ptarmicaefolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) 11. 185.—Austral.
pterocaulis, Miq. in Lehm. Pl. Preiss. i. 225 = ceratocarpa.
pubescens, Lindl. in Mitch. Journ. Trop. Austral. 242 = peduncularis.
repanda, *Schum. & Thonn. Beskr. Guin. Pl.* 194.—Afr. trop.
salicifolia, DC. Prod. i. 617 = viscosa.
salsolifolia, A. Cunn. in Hook. Journ. Bot. i. (1834) 251 = ericifolia.
scabra, *Lodd. ex Loud. Hort. Brit.* 154, nomen.—Austral.
Schiedeana, Schlecht. in Linnaea, xviii. (1844) 88 (err. typ. 49) = viscosa.
serrulata, *DC. Prod.* i. 617.—Reg. Argent.
sororia, Miq. in Lehm. Pl. Preiss. i. 225 = aptera.
spatulata, Sm. in Rees, Cycl. xii. n. 2 = viscosa.
stenophylla, *F. Muell. Fragm.* i. 72.—Austral.
stenoptera, *F. Muell. l. c.* 98.—Austral.
tenuifolia, Lindl. in Mitch. Journ. Trop. Austral. 248 = adenophora.
Thunbergiana, *Eckl. & Zeyh. Enum.* 54.—Afr. austr.
triangularis, *Lindl. in Mitch. Journ. Trop. Austral.* 219.—Austral.
trifida, *F. Muell. Fragm.* ix. 88.—Austral.
trifoliata, Tenw. Pl. Rar. 12. t. 9 = Pistea trifoliata.
trigona, Lindl. in Mitch. Journ. Trop. Austral. 236 = triangularis.
triquetra, *Forsgk. in Nat. en Genoesk. Arch. Nederl Indie*, ii. (1845) 96.—Java.
triquetra, *Wendl. Beobacht.* 44.—Austral.
truncatialis, *F. Muell. Fragm.* ii. 143.—Austral.
umbellata, G. Don, Gen. Syst. i. 674 = viscosa.
vestita, *Hook. in Mitch. Journ. Trop. Austral.* 265.—Austral.
viscosa, *Jacq. Enum. Pl. Carib.* 19; *Linn. Mant.* 228. —Cosmop. trop. et subtrop.
viscosa, Mart. in Flora, xxii. (1839) 1. Beibl. 57.—Cf. Schlecht. in Linnaea, xviii. (1844) 85 (err. typ. 51) = brasiliensis.
viscosa, Royen, ex Blume, Rumphia, iii. 191 = Candolleana.
viscosa, Sieber, ex Schlecht. in Linnaea, xviii. (1844) 86 (err. typ. 52) = Kobautiana.
Wallichiana, *Blume, Rumphia*, iii. 189.—Java.
Wightiana, Blume, l. c. = viscosa.
Zollingeri, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 587.—Java.

DODONAEA, Plum. ex Adans. Fam. ii. 349 (1763) = Comocladia, P. Br. (Anacard.).
tricuspidata, Pritz. Ic. Ind. 874 = C. tricuspidata.

DOELLIA, Sch. Bip. in Flora, xxv. (1842) 1. Beibl. 124; et in Walp. Rep. ii. 932 (1843) = Mitranea, DC. (Compos.).
Ketschyi, Sch. Bip. l. c. nomen.—Hab.?

DOELLINGERIA, Nees, Gen. et Sp. Aster. 177 (1832) = **Aster**, Linn. (Compos.).
amygdalina, Nees, l. c. 179 = A. umbellatus.
cornifolia, Lindl. in Hook. Comp. Bot. Mag. i. (1835) 98 = A. umbellatus.
cornifolia, Nees, Gen. et Sp. Aster. 181 = A. infirmus.
obovata, Nees, l. c. 182 = A. reticulatus.
ptarmicoides, Nees, l. c. 188 = A. ptarmicoides.
scabra, Nees, l. c. = A. scaber, *Thunb.*
trichocarpa, DC. Prod. v. 265 = A. striatus.
umbellata, Nees, Gen. et Sp. Aster. 178 = A. umbellatus.

DOEMIA, R. Br. in Mem. Wern. Soc. i. (1809) 50, err. typ. = **Daemia**, R. Br. (Asclep.).

DOERRIENA, Borkh. in Rhein. Mag. i. (1793) 528 = **Cerastium**, Dill. (Caryoph.).
erecta, Borkh. l. c. = C. quaternellum.

DOERRIENIA, Dennst. Schlüss. Hort. Malab. 31 (1818) = **Acronychia**, Forst. (Rutac.).
malabarica, Dennst. l. c. = A. laurifolia.

DOERRIERA, Stend. Nom. ed. II. i. 522 (1840) sphalm. = DOERRIENA, Borkh. (Caryoph.).

DOFIA, Adans. Fam. ii. 285 (1763) = **Dirca**, Linn. (Thymel.).

DOGA, Baill. Adansonia, ix. (1869) 205 = **Storckiella**, Seem. (Legumin.).
macrogramma, Panch. ex Baill. Hist. Pl. ii. 132 = S. Pancheri.
Pancheri, Baill. Adansonia, ix. (1869) 205 = S. Pancheri.

DOLIA, Lindl. Bot. Reg. (1844) sub t. 46. *CONVOLVULACEÆ*, Benth. & Hook. f. ii. 880.
ALIBREXIA, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 505.
APLOCARYA, Lindl. Bot. Reg. (1844) sub t. 46.
LALOUTRIA, Gandich. Bot. Voy. Bonite, t. 110 (1844–46 ?).
VELPEAULIA, Gandich. l. c. t. 109 (? 1844–46 ?).
clavata, *Miers*, in *Hook. Lond. Journ. Bot.* iv. (1845) 503.—Chili.
laxa, *Miers*, l. c.—Peruv.
leptophylla, *Miers*, l. c.—Peruv.
revoluta, *Benth. & Hook. f. Gen.* ii. 880.—Peruv.
salsoloides, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
tomentosa, *Benth. & Hook. f. Gen.* ii. 880.—Chili.
verticillata, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
verticillata, Walp. Ann. i. 528, sphalm. = præc.

DOLICHANDRA, Cham. in Linnæa, vii. (1832) 657 = **Macfadyena**, A. DC. (Bignoniac.).
cynanchoides, Cham. l. c. 658 (= *Spathodea Dolichandra*).—Bras.
Kohautiana, Presl, Bot. Bemerk. 92 = Bignonia Ungvis-cati.

DOLICHANDRONE, Fenzl, ex Seem. in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 31. *BIGNONIACEÆ*, Benth. & Hook. f. ii. 1046.
MARKHAMIA, Seem. in Journ. Bot. i. (1863) 226.
MUENTERIA, Seem. l. c. iii. (1865) 329. t. 35, 36.
arcuata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 380. —Ind. or.
Cauda-felina, *Benth. & Hook. f. Gen.* ii. 1046.—China.
crispa, *Seem. in Journ. Bot.* viii. (1870) 381.—Ind. or. falcata, *Seem. l. c.*—Ind. or.
filiformis, Fenzl, ex F. Muell. Fragm. iv. 149 = Spathodea filiformis.
heterophylla, F. Muell. l. c. = Spathodea heterophylla.
Lawii, *Seem. in Journ. Bot.* viii. (1870) 380.—Ind. or.
lutea, *Benth. & Hook. f. Gen.* ii. 1046.—Afr. trop.
Rheedii, *Seem. in Journ. Bot.* viii. (1870) 380.—Ind. or.; Malays.
serrulata, Seem. l. c. 383 = Lawii.
spathacea, Seem. l. c. i. (1863) 226 = crispa, Rheedii.
stipulata, *Benth. & Hook. f. Gen.* ii. 1046.—Burma.
tomentosa, *Benth. & Hook. f. l. c.*—Afr. trop.

DOLICHLASIUM, Lag. Amen. Nat. 33 (1811) = **Trixis**, P. Br. (Compos.).
glandulosiflorum, Lag. ex Hook. & Arn. in Comp. Bot. Mag. i. (1836) 84 = sag.
Lagasca, Gill. ex D. Don, in Phil. Mag. (Apr. 1832) 338.—Chili.

DOLICHODEIRA, Hanst. in Linnæa, xxvi. (1853–55) 205 = **Achimenes**, P. Br. (Gesner.).
tubiflora, Hanst. l. c.—Uruguay.

DOLICHODERIA, Benth. & Hook. f. Gen. ii. 999 (1876) = præc.

DOLICHOGYNE, DC. Prod. vii. 256 (1838) = **Nardophyllum**, Hook. (Compos.).
acaulis, Wedd. ex Benth. & Hook. f. Gen. ii. 259 = Lepidophyllum acaule.
armata, Wedd. Chlor And. i. 181.—Boliv.
Candolii, Remy, in C. Gay, Fl. Chil. iv. 103 = *stachelinoides*.
chlorichoides, Remy, l. c. 104 = N. chlorichoides.
geminiflora, Phil. in Linnæa, xxviii. (1856) 788 = N. geminiflora.
gnaphalioides, DC. Prod. vii. 256 = *stachelinoides*.
lepidophylla, Wedd. Chlor. And. i. 182 = Lepidophyllum lepidophyllum Meyenl.
rigida, Wedd. l. c. = Lepidophyllum rigidum.
rupestris, Wedd. l. c. 183 = Lepidophyllum rupestre.
stachelinoides, DC. Prod. vii. 256.—Reg. Argent.

DOLICHOLASIUM, Spreng. Gen. ii. 623 (1831) = DOLICHLASIUM, Lag. = **Trixis**, P. Br. (Compos.).

DOLICHOLOBIUM, A. Gray, in Proc. Am. Acad. iv. (1860) 308. *RUBIACEÆ*, Benth. & Hook. f. ii. 41.
Knollysii, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 360.—Ins. Fiji.
latifolium, *A. Gray, in Proc. Am. Acad.* iv. (1860) 309.—Ins. Fiji.
longuistmum, *Sprm. in Bonplandia*, iv. (1861) 256; *in Fl. Viti.* 121 t. 25.—Ins. Fiji.
Macgregori, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 360.—Ins. Fiji.
oblongifolium, *A. Gray, in Proc. Am. Acad.* iv. (1860) 309.—Ins. Fiji.

DOLICHOLUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 354 = Rhynchosia, Lour. (1790) (Legumin.).
flavus, Medic. l. c. = seq.?
minimus, Medic. l. c. = R. minima.

DOLICHONEMA, Nees, in Flora, iv. (1821) 308 = **Moldenhawia**, Schrad. (Legumin.).
speciosa, Nees, l. c. = M. floribunda.

DOLICHOS, Linn. Gen. ed. I. 222 (1737). *LEGUMINOSÆ*, Benth. & Hook. f. i. 540.
CHLORYLLIS, E. Mey. Comm. Pl. Afr. Austr. 149 (1885).
DIPOGON, Liebm. in Ann. Sc. Nat. Sér. IV. ii. (1854) 374.
LABLAB, Adans. Fam. ii. 325 (1763).
LABLAVIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 236 (1834).
abyssinicus, Hort. ex Steud. Nom. ed. I. 610 = Phaseolus violaceus.
acinaciformis, *Blanco, Fl. Filip.* ed. I. 578.—Ins. Philipp.
acinaciformis, Jacq. Coll. i. 114 = Canavalia ensiformis.
adenanthus, *Meisner-Sieber*, Forsk. Fl. Aegypt. Arab. 134 = Sesbania aegyptiaca.
Ahipa, *Wedd. in Ann. Sc. Nat.* Sér. IV. vii. (1857) 118.—Peruv.
albus, Lour. Fl. Cochinch. ii. 439 = D. Lablab.
altissimus, Jacq. Enum. Pl. Carib. 27 = Mucuna altissima.
altissimus, Vell. Fl. Flum. 330; vii. t. 154 = Dioclea violacea.
amœnus, Salisb. Prod. 334 = D. Lablab.
andongensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 213.—Afr. trop.
angularis, Willd. Sp. Pl. iii. 1051 = Vigna sinensis.
anguinus, Willd. Enum. Hort. Berol. Suppl. 51 = præc.
anguiosus, DC. Prod. ii. 399 = Phaseolus diversifolius.
angustifolius, *Eckl. & Zeyh. Enum.* 258.—Afr. austr.
angustifolius, Vahl, ex Guill. & Perr. Fl. Senegamb. Tent. i. 230 = Vigna capensis.
angustissimus, E. Mey. Comm. Pl. Afr. Austr. 142 = angustifolius.
arborescens, *G. Don, Gen. Syst.* ii. 358.—Mexic.
arborous, Forsk. Fl. Aegypt. Arab. 134 = Sesbania grandiflora.

DOLICHOS:—
arborous, Roxb. ex Wight & Arn. Prod. 262 = Pongamia glabra.
argenteus, Roxb. ex Wight & Arn. Prod. 249 = ciliatus.
argenteus, *Willd. Sp. Pl.* iii. 1047.—Afr. trop.
ariatatus, *Linn. Sp. Pl.* ed. II. 1021.—Am. austr.
articulatus, Lam. Encyc. ii. 296 = Pachyrhizus angulatus.
arvensis, *Vell. Fl. Flum.* 322; vii. t. 161.—Bras.
axillaris, E. Mey. Comm. Pl. Afr. Austr. i. 144 = biflorus.
barbatus, Wall. Cat. n. 5548 = Atylosia barbata.
benghalensis, Jacq. Hort. Vindob. ii. 57 = D. Lablab.
Benthamii, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 95 = glibosus.
bicolor, Hoffmgg. ex Steud. Nom. ed. II. i. 522 = Dioclea bicolor.
bicontortus, *Dur. in Actes Linn. Soc. Bord.* xxviii. (1869) 58.—Japon.
biflorus, *Linn. Sp. Pl.* 727.—Geront. trop.
blandus, R. Grah. in Wall. Cat. n. 5568 = Atylosia mollis.
bractatus, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 210.—Ind. or.
bracteatus, Wall. Cat. n. 5554 = Rhynchosia bracteata.
brevicaulis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 211.—Afr. trop.
Brunelli, *Zuccagni, in Roem. Collect.* 151.—Hab. ?
bulbosus, Linn. Sp. Pl. ed. II. 1030 = Pachyrhizus angulatus.
callosus, *Blume, Cat. Gew. Buitenz.* 91.—Java.
candicans, *Wall. Cat.* n. 5567.—Burma.
capensis, Linn. Pl. Rar. Afr. 17 = Vigna capensis.
capitatus, *Herb. Madr. ex Wall. Cat.* n. 5549 = Vigna Catjang.
Catjang, Linn. Mant. i. 259 = Vigna Catjang.
Chlorylis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 246.—Afr. austr.
Cienkowskii, Schweinf. & Asch. in Schweinf. Beitr. Fl. Aethiop. 256 = Canavalia ensiformis.
ciliatus, *Klein, in Willd. Sp. Pl.* iii. 1049.—Ind. or.
cochinensis, *Vell. Fl. Flum.* 321; vii. t. 158.—Bras.
comosus, G. F. W. Mey. Prim. Fl. Esseq. 241 = Mucuna comosa.
conspersus, R. Grah. in Wall. Cat. n. 5552 = Dunbaria conspersa.
coriaceus, R. Grah. l. c. n. 5562 = Dioclea reflexa.
crassus, R. Grah. l. c. n. 5553 = Atylosia mollis.
crenatifructus, Steud. ex A. Rich. Tent. Fl. Abyss. i. 228 = D. Lablab.
cultratus, *Forsk. Fl. Aegypt. Arab.* 134.—Arabia.
cultratus, Thunb. in Trans. Linn. Soc. ii. (1794) 340 = D. Lablab.
cultratus, Wight, ex Steud. Nom. ed. II. i. 528 = Canavalia ensiformis.
cuneifolius, Forsk. Fl. Aegypt. Arab. 134 = Crotalaria retusa.
cuneiformis, Steud. Nom. ed. II. i. 528 = præc.
Curtisii, G. Don, Gen. Syst. ii. 357 = D. Lablab.
cuspidatus, R. Grah. in Wall. Cat. n. 5566 = D. Lablab.
cylindricus, Desv. in Ham. Prod. Pl. Ind. Occ. 51 = Vigna capensis.
cylindricus, Mœnch, Meth. Suppl. 48 = Vigna sinensis.
Daltonii, Webb, in Hook. Niger Fl. 125 = biflorus.
dasycarpus, Miq. Fl. Ind. Bat. i. t. 186 = Vigna dolichoides.
debilis, D. Don, ex Hook. Niger Fl. 312 = Rhynchosia debilis.
debilis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 228.—Abyss.
decumbens, *Thunb. Prod. Pl. Cap.* 130.—Afr. austr.
densiflorus, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 214.—Afr. trop.
Didjre, *Forsk. Fl. Aegypt. Arab.* 133.—Arab.
Daltonii, *Dillon, Ind. Sem. Hort. Monsp.* (1847) 7.—Abyss.
dissectus, Lam. Encyc. ii. 300 = Phaseolus acontiifolius.
Dongalata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 214.—Afr. trop.
echinatulus, Blanco, Fl. Filip. ed. II. 401 = Vigna Catjang.
elatus, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 214.—Afr. trop.
elongatus, R. Grah. in Wall. Cat. n. 5548 = Atylosia elongata.

DOLICHOS:—

emarginatus, Jacq. Hort. Schoenb. ii. 50 = Canavalia obtusifolia.

ensiformis, Linn. Sp. Pl. 725 = Canavalia ensiformis.

ensiformis, Thunb. Fl. Jap. 279 = D. Lablab.

erosus, Linn. Sp. Pl. 726 = Pachyrhizus angulatus.

fabaeformis, L'Hérit. Stirp. Nov. 163 = Cyamopsis psoraloides.

Fabæ-indica, *Forsk. Fl. Aegypt. Arab.* 133.—Arab.

Fabæ-nigrita, *Forsk. l. c.* 132.—Arab.

falcatus, *Klein, ex Willd. Sp. Pl.* iii. 1047.—Ind. or.

falciformis, *E. Mey. Comm. Pl. Afr. Austr.* i. 144.—Afr. austr.

farinosus, Hochst. & Steud. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 224 = Rhynchosia minima.

festivus, Wall. Cat. n. 5551 = D. Lablab.

ficifolius, R. Grah. in Wall. l. c. n. 5563 = Pueraria phaseoloides.

filiformis, *Linn. Pl. Jam. Pugill.* 19; *Amoen. Acad.* v. 402.—Jamaic.

Finlaysonianus, R. Grah. in Wall. Cat. n. 5565 = Dunbaria conspersa.

formosus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 223.—Abyssin.

frutescens, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 240.—Reg. Himal.

funarius, *Molina. Sagg. Chile*, 156.—Chili.

galeatus, Gaudich. in Freyc. Voy. Bot. 486. t. 115 = Canavalia galeata.

gangeticus, Roxb. Hort. Beng. 55; Fl. Ind. iii. 310 = Vigna glabra.

geminiflorus, *G. Don, Gen. Syst.* iii. 358.—Mexic.

geminus, *Vell. Fl. Flum.* 321; vii. t. 159.—Bras.

gibbosus, *Thunb. Prod. Pl. Cap.* 130.—Afr. austr.

giganteus, Willd. Sp. Pl. iii. 1041 = Mucuna gigantea.

gladiatus, Jacq. Coll. ii. 276 = Canavalia ensiformis.

glutinosus, Roxb. Hort. Beng. 55; Fl. Ind. iii. 312 = Rhynchosia viscosa.

glycinoides, H. B. & K. Nov. Gen. et Sp. vi. 441 = Vigna glabra.

gracilis, Guill. & Perr. Fl. Seneg. Tent. i. 219 = Vigna gracilis.

grandiflorus, R. Grah. in Wall. Cat. n. 5556 = Pueraria Thomsoni.

Halet, *Wood, Bot. & Florist*, 98; *ex S. Wats. Bibliog. Ind. N. Am. Bot.* 219.—Am. bor.

Hamioo, *Siebold, in Verh. Batav. Gen.* xii. (1830) 55.—Japon.

hastæfolius, *Schmitzl. in Flora*, xxv. (1842) 1. *Beibl.* 146.—Nubia.

hastæformis, *E. Mey. Comm. Pl. Afr Austr.* i. 142.—Afr. austr.

hastatus, *Lour. Fl. Cochinch.* 442.—Afr. or.

Helicopus, *Steud. Nom.* ed. II. i. 528.—Afr. austr.

helvolus, Nutt. Gen. Am. ii. 112 = Phaseolus helvolus.

heterophyllus, *Hornem. Hort. Hafn. Suppl.* 80.—Ins. Canar.

hexandrus, Roxb. Hort. Beng. 55, nomen (quid?).—Ind. or.

hirsutus, Thunb. in Trans. Linn. Soc. ii. (1794) 329 = Pueraria Thunbergiana.

hirtus, Andr. Bot. Rep. t. 446 = Cylista villosa.

hirtus, Hort. ex DC. Prod. ii. 389 = Fagelia bituminosa.

hispidus, Steud. Nom. ed. II. i. 528 = Vigna hispida.

incurvus, Thunb. Fl. Jap. 280 = Canavalia ensiformis.

Jacquini, *DC. Prod.* ii. 397.—Ind. occ.

japonicus, Spreng. Syst. iii. 252 = Wistaria chinensis.

Jungbuhnianus, *Benth. in Miq. Pl. Jungh.* 240.—Java.

Lablab, *Linn. Sp. Pl.* 725.—Reg. trop.

lacteus, Rafin. Fl. Ludov. 108 (quid?).—Am. bor.

lanceolatus, *R. Grah. in Wall. Cat.* n. 5547.—Burma.

leucomelas, *Kunze, Delect. Sem. Hort. Lips.* (1842).—Afr. austr.

lignosus, Jacq. Select. Am. 205 = Jacquini.

ligneosus, Linn. Sp. Pl. 726 = D. Lablab.

linearis, *E. Mey. Comm. Pl. Afr. Austr.* i. 142.—Afr. austr.

linearis, Thunb. Fl. Jap. 280 = Canavalia obtusifolia.

littoralis, Vell. Fl. Flum. 321; vii. t. 160 = Canavalia obtusifolia.

lobatus, Willd. Sp. Pl. iii. 1047 = Vigna lobata.

Lotononis, *Basil. in Bull. Soc. Linn. Par.* i. (1883) 360.—Madag.

DOLICHOS:—

Lubia, *Forsk. Fl. Aegypt. Arab.* 133.—Aegypt.

luteolus, Jacq. Hort. Vindob. i. 39 = Vigna glabra.

luteus, Sw. Prod. Veg. Ind. Occ. 106 = Vigna retusa.

macrodon, R. Grah. in Wall. Cat. n. 5558 = Rhynchosia sericea.

mammosus, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 13 = Pachyrhizus angulatus?

martinicensis, Jacq. ex Medic. Phil. Bot. i. 205 = D. Lablab.

medicagineus, Lam. Encyc. ii. 297 = Rhynchosia minima.

melanophthalmus, DC. Prod. ii. 400 = Vigna sinensis.

Memnonius, Delile, Fl. Egypte, 254 = Rhynchosia Memnonia

mexicanus, Schlecht. in Linnæa, xii. (1838) 329 = Vigna glabra.

miniatus, H. B. & K. Nov. Gen. et Sp. vi. 441 = Canavalia obtusifolia.

minimus, Linn. Sp. Pl. 726 = Rhynchosia minima.

minutus, Roxb. ex Wight & Arn. Prod. 256 = Atylosia scarabæoides.

mitis, *A. Rich. Tent. Fl. Abyss.* i. 224.—Abyssin.

mollis, Jacq. Fragm. 60 = Dioclea mollis.

monachalis, Brot. Fl. Lusit. ii. 125 = Vigna Catjang.

montanus, Lour. Fl. Cochinch. ii. 439 = Pueraria phaseoloides.

monticola, *Mart. ex Benth. in Mart. Fl. Bras.* xv. i. 196.—Bras.

multiflorus, Torr. & Gray, Fl. N. Am. i. 281 = Dioclea Boykinii.

mutabilis, Salisb. Prod. 334 = Canavalia ensiformis.

nervosus, *Schum. & Thonn. Beskr. Guin. Pl.* 342.—*Afr. trop. occ.*

niloticus, Delile, Fl. Egypte, 253. t. 38 = Vigna luteola.

niloticus, Sieber, ex Steud. Nom. ed. II. i. 528 = D. Lablab.

obcordatus, A. Cunn. ex Benth. Fl. Austral. ii. 247 = Hardenbergia retusa.

obcordatus, Roxb. Hort. Beng. 55; Fl. Ind. iii. 308 = Canavalia obtusifolia.

obliquifolius, *Schnitzl. in Flora*, xxv. (1842) 1. *Beibl.* 145.—Nubia.

obovatus, Schum. & Thonn. Beskr. Guin. Pl. 341 = Canavalia obtusifolia.

obtusifolius, *G. Don, Gen. Syst.* iii. 358.—Mexic.

obtusifolius, Jacq. Obs. Bot. iii. 5 = Rhynchosia minima.

obtusifolius, Lam. Encyc. ii. 295 = Canavalia obtusifolia.

olivaceus, Schum. & Thonn. Beskr. Guin. Pl. 340 = Vigna Catjang.

Olivieri, *Schweigf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 556.—Afr. trop.

ovalifolius, Schum. & Thonn. Beskr. Guin. Pl. 341 = Canavalia obtusifolia.

ovatus, R. Grah. in Wall. Cat. n. 5540 = Prophocarpus tetragonolobus.

palmatifidus, *Moç. & Sesse, ex DC. Prod.* ii. 399.—Mexic.

palmatilobus, Sesse, ex DC. l. c. = Pachyrhizus palmatilobus.

palustris, Forsk. ex Steud. in DC. Prod. ii. 394 = Phaseolus aconitifolius.

paraboicus, Nutt. Gen. Am. ii. 112 = Phaseolus helvolus.

Pavonii, *G. Don, Gen. Syst.* iii. 358.—Peruv.

pentaphyllus, E. Mey. Comm. Pl. Afr. Austr. i. 143 = Ilmaria.

phaseoloides, Roxb. Hort. Beng. 55; Fl. Ind. iii. 316 = Pueraria phaseoloides.

pilosus, Klein, ex Willd. Sp. Pl. iii. 1043 = Vigna pilosa.

Plectrotropis, Steud. Nom. ed. II. i. 528 = Vigna capensis.

polystachios, Linn. Sp. Pl. 726 = Phaseolus perennis.

polystachyus, Thunb. Fl. Jap. 281 = Wistaria chinensis.

prostratus, Buch.-Ham. & Keen. ex Roxb. Hort. Beng. [26]; Fl. Ind iii. 316 = ciliata.

prostratus, *Steud. Nom.* ed. II. i. 528.—Afr. austr.

protractus, Steud. l. c. = Vigna triloba.

pruriens, Linn. Syst. ed. X. 1169 = Mucuna pruriens.

Pseudo-sojanus, *Weiw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 215.—Afr. trop.

DOLICHOS:—

psoraloides, Lam. Encyc. ii. 300 = Cyamopsis psoraloides.

pubescens, *Linn. Sp. Pl.* ed. II. 1021.—Am. austr.

pugioniformis, Raeusch. Nom. ed. III. 209 (quid?) —Hab.?

punctatus, Wight & Arn. Prod. 247 = Dunbaria conspersa.

pungens, *Vell. Fl. Flum.* 321; vii. t. 157.—Bras.

purpureus, Linn. Sp. Pl. ed. II. 1021 = D. Lablab.

pyramidalis, Lam. Encyc. ii. 296 = Rhynchosia psaseoloides.

racemosus, Roxb. ex Steud. Nom. ed. II. i. 528 = D. Lablab.

racemosus, *Vell. Fl. Flum.* 322; vii. t. 162.—Bras.

regularis, Linn. Sp. Pl. 726 = Galactia glabella.

repens, *Linn. Pl. Jam. Pugill.* 19; *Amoen. Acad.* v. 402.—Jamaic.

reticulatus, Buch.-Ham. in Wall. Cat. n. 5552 = Atylosia mollis.

reticulatus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 33 = Atylosia reticulata.

reticulatus, *Hochst. in A. Rich. Tent. Fl. Abyss.* i. 225.—Abyssin.

retusus, *E. Mey. Comm. Pl. Afr. Austr.* i. 147.—Afr austr.

rhomboideus, *O. Hoffm. in Linnæa*, xliii. (1880–82) 126.—Afr. trop.

rhynchosioides, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 419.—Peruv.

rhynchosioides, Miq. Fl. Ind. Bat. i. 1. 185 = Dunbaria conspersa.

rhynchosioides, Schlecht. in Linnæa, xxvii. (1854) 581 = lignosus.

roseus, Sw. Prod. Veg. Ind. Occ. 105 = Canavalia obtusifolia.

rotundifolius, Vahl, Symb. Bot. ii. 81 = Canavalia obtusifolia.

ruber, Jacq. Select. Am. 204 = Dioclea Jacquiniana.

rufescens, R. Grah. in Wall. Cat. n. 5544 = Rhynchosia aurea.

rufus, *Vell. Fl. Flum.* 320; vii t. 155.—Bras.

rupestris, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 212.—Afr. trop.

sanguineus, Jacq. Fragm. 45 = D. Lablab.

Savii, *Montr. in Mém. Acad. Lyon*, x. (1860) 197.—N Caled.

scaber, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111 —Guiana.

scabrisusculus, Hassk. in Flora, xxv. (1842) 11. Beibl. 50 = Centrosema virginianum.

scarabæoides, Linn. Sp. Pl. 726 = Atylosia scarabæoides.

sericeus, *E. Mey. Comm. Pl. Afr. Austr.* i. 141.—Afr. austr.

sesquipedalis, Blanco, Fl. Filip. ed. II. 402 = Vigna Catjang.

sesquipedalis, *Linn. Sp. Pl.* ed. II. 1019.—Am. austr.

Sesse, *G. Don, Gen. Syst.* iii. 357.—Mexic.

setæfolius, *Boj. ex Steud. Nom.* ed. II. i. 524.—Ins. Mauric.

sinensis, Forsk. Fl. Aegypt. Arab. 132 = Vigna luteola.

sinensis, Linn. Cent. Pl. ii. 28; Amoen. Acad. iv. 326 = Vigna sinensis.

smilacinus, *E. Mey. Comm. Pl. Afr. Austr.* i. 142.—Afr. austr.

Soja, Linn. Sp. Pl. 727 = Glycine Soja.

Soorootee, Roxb. ex Wight & Arn. Prod. 255 = Mucuna capitata.

sphærospermus, *DC. Prod.* ii. 400.—Jamaic.

spicatus, R. Grah. in Wall. Cat. n. 5557 = Pueraria Thomsoni.

spicatus, R. Grah. l. c. n. 5557 A, B = Pueraria phaseoloides.

spicatus, Koen. ex Roxb. Cat. Hort. Beng. 55 = D. Lablab.

splendens, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 215.—Afr. trop.

spurcatus, G. F. W. Mey. Prim. Fl. Esseq. 242 = Mucuna aspera.

stenocarpus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 224.—Abyssin.

stipulaceus, Lam. Encyc. ii. 300 = Phaseolus trilobus.

stipulosus, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 213.—Afr. trop.

striatus, Steud. Nom. ed. II. i. 524 = Vigna capensis.

subsesquialis, *R. Grah. in Wall. Cat.* n. 5539.—Burma.

subcarnosus, *Wight & Arn. Prod.* 249.—Ind. or.

DOLICHOS :—
subracemosus, *Yacq. Select. Am.* 205.—N. Granat.
suffultus, R. Grah. in Wall. Cat. n. 5564 = Psophocarpus palustris.
tamulicus, *Spreng. Nachv. i. Bot. Gart. Halle*, 21.—Ind. or.
tenuis, *Stend. Nom.* ed. II. i. 524.—Afr. austr.
tetragonolobus, Linn. Syst. ed. X. 1162 = Psophocarpus tetragonoloba.
tetraspermus, Willd. Sp. Pl. iii. 1044 = D. Lablab.
tomentosus, Herb. Madr. ex Wall. Cat. n. 5538 = biflorus.
tomentosus, *Rafin. Fl. Ludov.* 108 (sp. dub.).—Am. bor.
tomentosus, Roth, Nov. Pl. Sp. 345 = Rhynchosia sericea.
tranquebaricus, Jacq. Hort. Vindob. iii. 39. t. 70 = Vigna Catjang.
trilobatus, Linn. Mant. i. 101 = Phaseolus trilobus.
trilobatus, Wall. Cat. n. 5541 = falcatus.
trilobus, Blanco, Fl. Filip. ed. II. 408 = Phaseolus calcaratus.
trilobus, Hontt. Handleid. z. 168. t. 64 = Vigna lobata.
trilobus, Linn. Sp. Pl. 726 = Pachyrhizus angulatus.
trilobus, Lour. Fl. Cochinch. 585 = Pachyrhizus angulatus, Pueraria Thunbergiana.
trilobus, Roth, ex Wall. Cat. n. 5588, partim = Phaseolus aconitifolius.
trilobus, Thunb. Fl. Cap. 590 = Vigna triloba.
truncatus, *Miq. Fl. Ind. Bat.* i. 1. 186.—Java.
tuberosus, Lam. Encyc. ii. 296 = Pachyrhizus tuberosus.
tuberosus, Roxb. ex Wight & Arn. Prod. 249 = falcatus.
tuncifolius, Raeusch. Nom. ed. III. 209, spalm. = cuneifolius.
umbellatus, *Thunb. in Trans. Linn. Soc.* ii. (1794) 339.—Japon.
umbellatus, Willd. ex Steud. Nom. ed. II. i. 524 = glibosus.
uncinatus, Linn. Sp. Pl. ed. II. 1021 = Teramnus uncinatus.
undulatus, *Moq. & Sesse, ex G. Don, Gen. Syst.* ii 857.—Mexic.
unguiculatus, Linn. Sp. Pl. 725 = Vigna Catjang.
unguiculatus, Thunb. Fl. Jap. 279 = umbellatus.
uniflorus, Lam. Encyc. ii. 299 = biflorus.
urens, Linn. Syst. ed. X. 1162; Jacq. Select. Am. t. 182. f. 84 = Mucuna urens.
urens, Roxb. ex Wight & Arn. Prod. 254 = Mucuna monosperma.
varius, *Span. ex Miq. Fl. Ind. Bat.* i. 1. 184.—Java.
vestitus, R. Grah. in Wall. Cat. n. 5545 = Flemingia vestita.
vexillatus, H. B. & K. Nov. Gen. et Sp. vi. 440 = Vigna capensis.
vexillatus, Nutt. Gen. Am. ii. 112 = Phaseolus helvolus.
violaceus, Hort. ex Steud. Nom. ed. II. i. 524 = Phaseolus violaceus.
virgatus, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guian.
viridis, Boch.-Ham. ex Wall. Cat. n. 5559 = Pueraria phaseoloides.
virosus, Roxb. Hort. Beng. 55; Fl. Ind. iii. 301 = Canavalia ensiformis.
Woolawa, Roxb. ex Wight & Arn. Prod. 248 = biflorus.

DOLICHOSIPHON, Phil. in Ann. Univ. Chil. (1873) 534. *SOLANACEÆ.*
Volckmanni, *Phil. l. c.*—Chili.

DOLICHOSTYLIS, Cass. in Dict. Sc. Nat. lvi. 186 (1828) = **Bernadesia**, Mutis (Compos.).
laurifolia, Cass. l. c. 189 (= *Pulicalden laurifolia*).—Peru.

DOLICHOSTYLIS, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 305. *CRUCIFERÆ* [Benth. & Hook. f. i. 996].
LEPTONEMA, Hook. Ic. Pl. t. 699 (1844).
STENONEMA, Hook. ex Benth. & Hook. f. Gen. i. 75 (1862).
Fendleri, *Turcz. l. c.* 306.—Venezuela.
Lindenii, *Turcz. l. c.*—N. Granat.

DOLICHOTHECA, Cass. in Dict. Sc. Nat. li. 475 (1827) = **Corræopsis**, Linn. (Compos.).

DOLICHUS, E. Mey. Comm. Pl. Afr. Austr. i. p. liv. 140 (1835) = **Dolichos**, Linn. (Legumin.).

DOLICOTHECA, Benth. & Hook. f. Gen. ii. 886 (1873) sphalm. = DOLICHOTHECA, Cass. = **Corræopsis**, Linn. (Compos.).

DOLIOCARPUS, Roland. in Vet Akad. Handl. Stockh. (1756) 249. *DILLENIACEÆ*, Benth. & Hook. f. i. 12.
CALINGA, Aubl. Pl. Gui. i. 556. t. 221 (1775).
MAPPIA, Schreb. Gen. Pl. 806 (1791).
OTHLIS, Schott, in Spreng. Syst. iv. App. 407 (1827).
RHIMIUM, Schreb. Gen. Pl. 701 (1791).
RICARDETIA, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 47.
SORAMIA, Aubl. Pl. Gui. i. 552. t. 219 (1775).
TIGAREA, Aubl. l. c. 917. t. 350, 351 (1775).
brevipedicellatus, *Gerche, in Linnæa*, xxii (1849) 47.—Bras.
Calinea, *Y. F. Gmel. Syst* 805.—Guiana; Bras.
Calinea, Griseb. Fl. Brit. W. Ind 8, partim = brevipedicellatus.
calineus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = D. Calinea.
castaneæfolius, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. t. 78.—Bras.
dentosus, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 65.—Bras.
elegans, *Eichl. in Mart. Fl. Bras.* xiii. t. 76. t. 18. f. 1.—Bras.
glomeratus, *Eichl. l. c.* 77. t. 18. f. 2.—Bras.
grandiflorus, *Eichl. l. c.* 78.—Bras.
macrocarpus, *Mart. ex Eichl. l. c.* 77.—Bras.
major, Y. F. Gmel. Syst. 805 = strictus.
nitidi, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 16.—N. Granat.
pischinensis, Mart. ex Eichl. in Mart. Fl. Bras. xiii. i. 76 = dentosus.
pubens, Mart. in Flora, xxi. (1838) ii. Beibl. 49 = Rolandri.
pubescens, Miq. in Linnæa, xix. (1847) 492 = Banara brasiliensis.
Rolandri, *Y. F. Gmel. Syst.* 805.—Guiana; Bras.
scandens, Poir. Encyc. Suppl. ii. 497 = Rolandri.
Schottianus, *Eichl. in Mart. Fl. Bras.* xiii. i. 74.—Bras.
Sellowianus, *Eichl. l. c.* 75.—Bras.
semidentatus, Garcke, in Linnæa, xxii (1849) 48 = Desmondi.
sessiliflorus, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 7.—Bras.
Soramia, *DC. Syst. i.* 406.—Guiana.
soramius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = D. Soramia.
spinuliferus, Miq. in Linnæa, xviii. (1844) 266 = dentosus.
strictus, *Poir. Encyc. Suppl.* ii. 498.—Guiana.
undulatus, *Eichl. in Mart. Fl. Bras.* xiii. i. 78.—Bras.
virgatus, *Sagot, in Ann. Sc. Nat.* Sér. VI. x. (1880) 381.—Guiana.

DOLLINERA, Endl. Gen. 1285 (1841) = **Desmodium**, Desv. (Legumin.).
ruficoma, Endl. ex Walp. Rep. i. 736 = Desm. rufescens.
sambuensis, Endl. l. c. = Desm. floribundum.

DOLLINERIA, Sauter, in Flora, xxxv. (1852) 353 = **Draba**, Dill. (Crucifer.).
ciliata, Sauter, l. c. 354 = D. ciliata.

DOLOMIAEA, DC. in Guill. Archiv. Bot. ii. (1833) 330 = **Jurinea**, Cass. (Compos.).
macrocephala, DC. l. c. = J. macrocephala.

DOLOPHRAGMA, Fenzl, in Ann. Wien. Mus. i. (1836) 63. t. 71 = **Arenaria**, Rupp. (Caryophyll.).
globiflorum, Fenzl, l. c. = A. globiflora.
juniperinum, Fenzl, l. c. = A. juniperina.

DOMA, Lam. Illustr. t. 900 (1823) = **Hyphæne**, Gaertn. (Palm.).

DOMBEIA, Raeusch. Nom. ed. III. 291 (1797) = DOMBEYA, Lam. = **Arenicaria**, Jus. (Comif.).

DOMBEYA, Cav. Dias. 121. t. 58, 61 (1787). *STERCULIACEÆ*, Benth. & Hook. f. i. 221.
ASSONIA, Cav. Dias. 120. t. 49 (1787).
ASTRAPÆA, Lindl. Collect. Bot. t. 14 (1823).
HILSENBERGIA, Boj. in Ann. Sc. Nat. Sér. II. xviii. (1842) 189.
KORNIGIA, Comm. ex Juss. Gen. 275 (1789).
LEEUWENHOECKIA, E. Mey. ex Endl. Nov. Stirp. Dec. 87 (1839).
VAHLIA, Dahl, Obs. Bot. 21 (1787).
WALCUFFA, J. F. Gmel. Syst. 1029 (1791).
XEROPETALUM, Delile, Cent. Pl. Méroé, 34 (1826).
acerifolia, Gaertn. Fruct. ii. 260 = Pterospermum acerifolium.
acutangula, *Cav. Diss.* 123. t. 58. f. 2.—Ins. Mascar.
Ameliæ, *Guill. Archiv. Bot.* i. (1832) 367.—Madag.
angulata, Cav. Diss. 123. t. 59. f. 1 = acutangula.
angulata, Mast. in Gard. Chron. (1867) 74 = Mastersii.
antsanakensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 487.—Madag.
astrapæoides, Boj. Hort. Maurit. 40 = Astiria roseo.
Bernieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 500.—Madag.
biternata, Blanco, Fl. Filip. ed. II. 244 = Helicteres spicata.
Bojeriana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 495.—Madag.
borbonica, *Gaertn Fruct* ii. 259. t. 187.—Ins. Borb.
Breonii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 494.—Madag.
Bruceana, *A. Rich. Tent. Fl. Abyss.* i. 77.—Afr. trop.
Burgessiæ, *Gerr. ex Harv. & Sond. Fl. Cap.* ii. 590.—Afr. austr.
cannabina, *Hilsenb. & Boj. ex Hook. Bot. Mag.* t. 3619.—Madag.
Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 493.—Madag.
cordifolia, DC. Prod. i. 499 = acutangula.
Coria, *Baill. l. c.* 491.—Madag.
cuspidata, *Boj. Hort. Maurit.* 40, nomen.—Madag.
cymosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 589.—Afr. austr.
decanthera, Blanco, Fl. Filip. ed. I. 349 = Helicteres spicata.
decanthera, Cav. Dias. 126 = Trochetia decanthera.
Delilei, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 225 = multiflora?
densiflora, *Planch. ex Harv. & Sond. Fl. Cap.* ii. 589.—Afr. austr.
diversifolia, Spreng. Syst. iv. Cur. Post. 256 = Pterospermum diversifolium.
Dregeana, *Sond. in Linnæa*, xxiii. (1850) 18.—Afr. austr.
elliptica, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 191.—Madag.
erythrocladia, *Boj. l. c.*—Madag.
Erythroxylon, Andr. Bot. Rep. t. 389 = Melhania Melanoxylon.
Erythroxylon, Willd. Sp. Pl. iii. 725 = Melhania Erythroxylon.
ferruginea, *Cav. Diss.* 128. t. 62. f. 2.—Ins. Mascar.
ferruginosa, Poir. Encyc. v. 159, in syn. = præc.
ficulnea, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 488.—Madag.
floribunda, *Baker, in Journ. Linn. Soc.* xxi. (1884) 325.—Madag.
Glechomacifolia, *Baker, l. c.* xx. (1883) 101.—Madag.
globosa, Boj. ex Baker, Fl. Mauri. 28 = populnea.
grandifolia, *Lamk. ex G. Don, Gen. Syst.* i. 536.—Maurit.
Greveana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 492.—Madag.
guazumæfolia, *Baill. l. c.* 495.—Madag.
Hamiltoniana, G. Don, Gen. Syst. i. 536 = Melhania Hamiltoniana.
Hildebrandtii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 493.—Madag.
Hilsenbergii, *Baill. l. c.* 495.—Madag.
hirsuta, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 190.—Madag.
Humblotii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 495.—Madag.
Kirkii, *Mast. in Oliver, Fl. Trop. Afr.* i. 237.—Afr. trop.
Lantziana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 496.—Madag.
laurifolia, *Baill. l. c.* 494.—Madag.
longicuspis, *Baill. l. c.* 488.—Madag.
longifolia, *Baill. l. c.* 496.—Madag.
longipes, *Baill. l. c.* 487.—Madag.
loxocombensis, *Baill. l. c.* 493.—Madag.
lucida, *Baill. l. c.* 496.—Madag.

DOMBEYA:—
macrantha, *Baker, in Journ. Linn. Soc.* xxi. (1884) 225. —Madag.
manaharica, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 491.—Madag.
Mastersii, *Hook. f. Bot. Mag.* t. 6839.—Afr. trop.
Melanoxylon, Roxb. in Beatson, St. Hel. Tracts, 307 = Melhania Melanoxylon.
minor, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 225 = multiflora.
modesta, *Baker, in Journ. Bot.* xx. (1882) 47.—Madag.
mollis, *Hook. Bot. Mag.* t. 4578.—Hab.?
multiflora, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 225.—Afr. trop.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 17.—Afr. austr.
obovalis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 496. —Madag.
ovata, *Cav. Diss.* 127. t. 41. f. 2.—Ins. Borbon.
palmata, Cav. l. c. 122. t. 38. f. 1 = acutangula.
parviflora, *Boiv. ex Baill. in Bull. Soc. Linn. Par.* i. (1885) 492.—Madag.
Pervillei, *Baill. l. c.* 488.—Madag.
phœnicea, Cav. Diss. 129 = Pentapetes phœnicea.
platanifolia, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 190.—Madag.
populnea, *Baill. Hist. Pl.* iv. 67; *Baker, Fl. Maurit.* 35.—Ins. Mascar.
prostrata, Burch. ex DC. Prod. i. 499 = Melhania prostrata.
Pseudo-Populus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 500.—Madag.
punctata, *Cav. Diss.* 125. t. 41. f. 2.—Ins. Mascar.
reflexa, *Drapiez, ex Reichb. Fl. Exot.* vi. 40. t. 846.—Madag.
repanda, *Baker, in Journ. Linn. Soc.* xxi. (1884) 226. —Madag.
reticulata, *Mast. in Oliver, Fl. Trop. Afr.* i. 238.—Afr. trop.
rigida, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 487.—Madag.
rottleroides, *Baill. l. c.* 494.—Madag.
rotundifolia, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 225.—Afr. austr.
rubifolia, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 492. —Madag.
Schimperiana, *A. Rich. Tent. Fl. Abyss.* i. 78.—Afr. trop.
senegalensis, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 225, sub t. 605 = multiflora.
spectabilis, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 191.—Afr. trop.
stipulacea, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 498.—Madag.
tiliacea, *Planch. Fl. des Serres*, Sér. I. vi. (1850–51) 225.—Afr. austr.
tiliæfolia, Cav. Diss. 124. t. 39. f. 2 = acutangula.
tomentosa, *Cav. l. c.* 125. t. 39. f. 3.—Madag.
triumfettæfolia, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 191.—Madag.
umbellata, *Cav. Diss.* 127. t. 41. f. 1.—Ins. Mascar.
Valva, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 494.—Madag.
velutina, Willd. Sp. Pl. iii. 726 = Melhania velutina.
vilosa=filosa, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 191.—Ins. Comor.
Wallichii, *Benth. & Hook. f. Gen.* i. 221.—Madag.?

DOMBEYA, Lam. Encyc. ii. 301. t. 398 (1786) = Araucaria, Juss. (Conif.).
armacea, Raeusch. Nom. ed. III. 291 = A. imbricata.
chilensis, Lam. Encyc. ii. 301 = A. imbricata.
excelsa, Lamb. Pinus, ed. I. 87 = A. excelsa.

DOMBEYA, L'Hérit. Stirp. Nov. 33. t. 17 (1784) = Tourretia, Juss. (Bignoniac.).
lappacea, L'Hérit. l. c. t. 7. lappacea.
lapponica, [. F. Gmel. Syst. 1541, sphalm. = prace.
nodiflora, L'Hérit. ex DC. Prod. ix. 289 = Ecremocarpus scaber.

DOMBEYOA, Phil. Fl. Atac. 25. t. 2 C (1860) = *UMBELLIFERAE*, Benth. & Hook. f. i. 876.
oppositifolia, *Phil. l. c.*—Chili.

DONACIUM, Fries, in Bot. Notiser, (1848) 131 = DONAX, Beauv. = Arundo, Linn. (Gramin.).

DONACODES, Blume, Enum. Pl. Jav. i. 54 (1827) = Amomum, Linn. (Scitamin.).
alliacea, Teijsm. & Binn. Cat. Hort. Bog. 58.—Java.
aperta, Teijsm. & Binn. l. c. 850.—Ins. Banca.
elongata, Teijsm. & Binn. l. c.—Java.
incarnata, Teijsm. & Binn. l. c.—Ins. Amboina.
macrocephala, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 211.—Java.
minor, Teijsm. & Binn. Cat. Hort. Bog. 850.—Java.
paludosa, Blume, Enum. Pl. Jav. i. 54 = Elettaria paludosa.
Pinanga, Blume, l. c. = Alpinia Pinanga.
rosea, Teijsm. & Binn. Cat. Hort. Bog. 58 = Am. roseum.
rubra, Blume, Enum. Pl. Jav. i. 54 = Alpinia rubra.
tomentosa, Blume, l. c. 55 = Alpinia tomentosa.
villosa, Teijsm. & Binn. in Nederl. Kruidk. Arch. iii. (1855) 892 = A. villosum?
Walang, Blume, Enum. Pl. Jav. i. 55 = Alpinia Walang.

DONALDIA, Klotzsch, in Monatsb. Berl. Acad. (Mærz 1854) 127; et in Abh. Akad. Berl. (1854) 78. t. 7 B = Begonia, Linn.
Ottonis, Klotzsch, ll. cc. 127, 79 = B. Ottonis.
ulmifolia, Klotzsch, ll. cc. 127, 78 = B. ulmifolia.

DONATIA, Bert. ex Remy, in C. Gay, Fl. Chil. v. 289 (1849) = Lactarrisea, Remy (Polygon.).

DONATIA, Forst. Char. Gen. 9. t. 5 (1776). *SAXIFRAGEAE*, Benth. & Hook. f. i. 684.
fasciculata, *Forst.* = Rag. Magell. *magellanica*, Lam. Illustr. i. 217. t. 51 = fascicularis. Novæ-Zelandiæ, *Hook. f. Fl. N. Zel.* ii. t. 20.—N. Zel.

DONATIA, Loefl. Iter, 198 (1758) = Aviosmia, Linn. (Verbenac.).

DONATOPHORUS, Zipp. in Bijdr. Natuurk. Wetensch. v. (1830) 181 = Harpullia, Roxb. (Sapindac.).
erythrospermus, Zipp. l. c. = H. thanatophora.

DONAX, Beauv. Agrost. 77 (1812) = Arundo, Tourn. (Gramin.).
acuminatus, Presl, ex Steud. Nom. ed. II. i. 524 = A. Pliniana.
acutiflorus, Beauv. Agrost. 78 = Deyeuxia acutiflora.
Ampelodesmos, Roem. & Schult. Syst. ii. Mant. 835, in syn. = Ampelodesmos tenax.
arundinaceus, Beauv. Agrost. t. 16. f. 4 = A. Donax.
australis, Steud. Nom. ed. II. i. 524 = Phragmites communis.
bengalensis, Beauv. Agrost. 78 = A. bengalensis.
bicolor, Beauv. l. c. = A. bicolor.
bifarius, Trin. in Spreng. Neue Entdeck. ii. 78 = A. bifarius.
borralis, Trin. Fund. Agrost. 156 = Graphephorum arundinaceum.
festuceoides, Beauv. l. c. 78 = Graphephorum arundinaceum.
festucoides, Beauv. l. c. = Ampelodesmos tenax.
Kotschubaeus, Trin. in Repr. in Beitr. Russ. Reich. ii. 65 = Graphephorum pellosachne.
mauritanicus, Beauv. Agrost. 78 = A. Pliniana.
montanus, Beauv. l. c. = Calamagrostis varia.
Plinii, C. Koch, Dendrol. ii. ii. 359 = A. Pliniana.
sativus, Presl, Cyp. et Gram. Sic. 22 = A. Donax.
tenax, Beauv. Agrost. 78 = Ampelodesmos tenax.
Thouarii, Beauv. l. c. t. 16. f. 5 (Thuarii) = A. madagascariensis.
versicolor, Beauv. l. c. = A. Donax.
viridiflosculosus, Roem. & Schult. Syst. ii. 601 = Deyeuxia viridiflavescens.

DONAX, Lour. Fl. Cochinch. 11 (1790) = Clinogyne, Salisb. (Scitamin.).
Arundinastrum, Lour. l. c. = C. dichotoma.

DONCKLAERIA, Hort. ex Lond. Encyc. Pl. Suppl. ii. 1850 (1855) = Centradenia, G. Don (Melastom.).
discerrifolia, Hort. ex Lond. l. c. = C. inæquilateralis.

DONDIA, Adans. Fam. ii. 261 (1763) = Suæda, Forsk. (1775) (Chenopod.).

DONDIA, Spreng. Fl. Umb. Prod. 21 (1813) = Hacquetia, Neck. (Umbell.).
Epipactis, Spreng. l. c. = H. Epipactis.

DONDISIA, DC. Prod. iv. 469 (1830) = Plectronia, Linn. (Rubiac.).
javida, Hassk. Cat. Hort. Bog. Alt. 112 = seq.
foetida, Hassk. Ind. Sem. Hort. Amstelod. (1841) = Vanguieria edulis.
horrida, Korth. in Nederl. Kruidk. Arch. ii. ii. (1851) 238 = P. horrida.
Leschenaultii, DC. Prod. iv. 469 = P. angustiflora.

DONDISIA, Reichb. Consp. 144 (1828) = Hacquetia, Neck. (Umbell.).
Epipactis, Reichb. in Moesal. Handb. ed. II. i. 492 = H. Epipactis.

DONDISIA, Scop. Introd. 316 (1777) = Raphanus, Linn. (Crucif.).

DONIA, R. Br. in Ait. Hort. Kew. ed. II. v. 82 (1813) = Grindelia, Willd. (Compos.).
canariensis, Less. Syn. Comp. 199 = Jasonia lævigata.
ciliata, Nutt. in Journ. Acad. Nat. Sc. Philad. ii. (1821) 118 = Aplopappus ciliatus.
eduis, Lindl. ex Lem. Illustr. Hortic. ii. (1858) Misc. 91.—Mexic.
fruticosa, Desf. Tabl. ed. II. 269 = Grindelia fruticosa.
glutinosa, R. Br. in Ait. Hort. Kew. ed. II. v. 82 = G. glutinosa.
glutinosa, Hook. Fl. Bor. Am. ii. 25 = G. oregana.
inuloides, Hook. l. c. = G. inuloides.
lanceolata, Hook. l. c. = Aplopappus lanceolatus.
squarrosa, Pursh, Fl. Am. Sept. ii. 559 = G. glutinosa.
uniflora, Hook. Fl. Bor. Am. ii. 25 = Aplopappus uniflorus.

DONIA, R. Br. in Ross, Voy. ed. I. App. p. xli (1819) = Oxyris, Hill. (Polygon.).
digyna, R. Br. in Parry, Voy. App. 41 = O. digyna.
sapida, R. Br. in Ross, Voy. App. ed. I. p. xlii = O. digyna.

DONIA, G. & D. Don, in G. Don, Gen. Syst. ii. 467 (1832) = Clianthus, Soland. (Legumin.).
formosa, G. Don, l. c. 468 = C. Dampieri.
punicea, G. Don, l. c. = C. puniceus.
speciosa, G. Don, l. c. = C. Dampieri.

DONIANA, Rafin. in Am. Monthly Mag. (1818) 268 = Grindelia, Willd. (Compos.).

DONIDSIA, G. Don, Gen. Syst. iii. 461 (1834) = DONDISIA, DC. = Plectronia, Linn. (Rubiac.).

DONIOPHYTON, Wedd. Chlor. And. i. 7. t. 4 (1855). *COMPOSITAE*, Benth. & Hook. f. ii. 489. sadicolum, *Wedd. l. c.* 8. t. 4 B.—Chili.

DONKELAARIA, Hort. ex Lem. Illustr. Hortic. ii. (1855) Misc. 78, in nota = Centradenia, G. Don (Melastom.).
discerrifolia, Hort. ex Lem. l. c. = C. inæquilateralis.

DONKELAARIA, Lem. Illustr. Hortic. ii. (1855) Misc. 72 = Guettarda, Linn. (Rubiac.).
dichotoma, Lem. l. c. = G. dichotoma?

DONNINGIA, A. Gray, in Journ. Linn. Soc. xiv. (1875) 29 sphalm. = Downingia, Torr. (Lobeliac.).

DONTOSPERMUM, Sch. Bip. in Walp. Rep. ii. 955 (1843) = Odontospermum, Neck. (Compos.).

DONTOSTEMON, Andrz. ex DC. Prod. i. 190 (1824). *CRUCIFERAE*, Benth. & Hook. f. i. 77.
ANDROSACEA, DC. Prod. i. 190 (1824).
brevipes, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 149 = Malcolmia brevipes.
circinatus, Bunge, l. c. 148 = Malcolmia circinata.
dentatus, *Ledeb. Fl. Ross.* i. 178.—Dahuria.
eglandulosus, Ledeb. l. c. = dentatus.
grandiflorus, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 147 = Malcolmia Bunget.
hispidus, *Stanton. in Bull. Acad. Pétersb.* xviii. (1873) 322.—Manchuria.
integrifolius, *Ledeb. Fl. Ross.* i. 174.—Sibir.

DONTOSTEMON :—
matthioloides, *Franch. Pl. Davidian.* i. (1884) 35.—China.
micranthus, *C.A. Mey. in Ledeb. Fl. Alt.* iii. 190.—China.
oblongifolius, Ledeb. Fl. Ross. i. 175 = dentatus.
pectinatus, *Ledeb. l. c.*—Dahuria; Sibir.
persoonia, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 191.—Sibir.
scorpioides, Bunge, Arb. Naturf. Ver. Riga, i. (1847) 150 = Malcolmia scorpioides.
senilis, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 491.—Mongolia.

DONZELLA, Lem. in Orb. Dict. v. 108 (1849) = seq.

DONZELLIA, Tenore, Ind. Sem. Hort. Neap. (1839) 11. n. 7; et in Ann. Sc. Nat. Sér. II. xiii. (1840) 379. *EUPHORBIACEAE.*
spinosa, *Tenore, ll. cc.* 11. n. 7; 380.—Bras.

DOODIA, Roxb. Hort. Beng. 98 (1814); Fl. Ind. iii. 365 (1832) = Urvaria, Desv. (Legumin.).
alopecuroides, Roxb. ll. cc. 98, 366 = U. Lagopus.
crinita, Roxb. Fl. Ind. iii. 369 = U. crinita.
hamosa, Roxb. l. c. 367 = U. hamosa.
lagopodioides, Roxb. l. c. 366 = U. lagopodioides.
picta, Roxb. l. c. 368 = U. picta.
simplicifolia, Roxb. l. c. 366.—Ind. or.

DOONA, Thw. in Hook. Kew Journ. iii. (1851) t. 12; iv. (1852) 7. *DIPTEROCARPEAE*, Benth. & Hook. f. i. 193.
CARYOLOBIS, Gaertn. Fruct. i. 215. t. 45 (1788).
affinis, *Thw. Enum. Pl. Zeyl.* 35.—Zeylan.
congestiflora, *Thw. l. c.*—Zeylan.
cordifolia, *Thw. l. c.*—Zeylan.
Gardneri, *Thw. l. c.*—Zeylan.
macrophylla, *Thw. l. c.* 403.—Zeylan.
nervosa, *Thw. l. c.* 35.—Zeylan.
oblonga, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 306.—Zeylan.
ovalifolia, *Thw. Enum. Pl. Zeyl.* 402.—Zeylan.
trapezifolia, *Thw. l. c.* 35.—Zeylan.
venulosa, *Thw. l. c.* 403.—Zeylan.
zeylanica, *Thw. in Hook. Kew Journ.* iii. (1851) t. 12; iv. (1852) 7.—Zeylan.

DOORNIA, De Vriese, in Hook. Kew Journ. vi. (1854) 266; et in Tijdsch. Fl. 174 (1855) = Pandanus, Linn. *reflexa*, De Vriese, ll. cc. 267, 174 = P. reflexus.

DOOSERA, Roxb. ex Wight & Arn. Prod. 362 (1834) = Mollugo, Linn. (Ficoid.).
esculenta, Roxb. ex Wight & Arn. l. c. = M. hirta.

DOPATRIUM, Buch.-Ham. ex Benth. Scroph. Ind. 30 (1835). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 958.
aristatum, *Hassk. in Flora*, xlvii. (1864) 56.—Ins. Philipp.
japonicum, *Franch. & Sav. Enum. Pl. Jap.* i. 345.—Japon.
junceum, *Buch.-Ham. ex Benth. Scroph. Ind.* 31.—As. et Austral. trop.
lobelioides, *Benth. l. c.*—Ind. or.
macranthum, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 190.—Afr. trop.
nudicaule, *Benth. Scroph. Ind.* 31.—Ind. or.; Malaya.
senegalense, *Benth. in DC. Prod.* x. 407.—Afr. trop.

DORAENA, Thunb. Nov. Gen. 59 (1783); Gaertn. f. Fruct. iii. 27. t. 184 (1805) = Maesa, Forsk. (Myrsin.).
japonica, Thunb. l. c.; Fl. Jap. 84 = M. Doraena.

DORANTHERA, Steud. Nom. ed. II. i. 525 (1840) = seq.

DORATANTHERA, Benth. ex Endl. Gen. 685 (1839) = Antisharis, Endl. (Scrophularin.).
linearis, Benth. in DC. Prod. x. 347 = A. linearis.
arnegalensis, Walp. Rep. iii. 305 = A. linearis.

DORATIUM, Seland. ex Jaume St. Hil. Expos. ii. 367 (1805) = Xanthoxylum, Linn. (Rutac.).

DORATOLEPIS, Schlecht. in Linnaea, xx. (1847) 593 = Lagenophora, Less. (Compos.).
tetrachaeta, Schlecht. l. c. = L. pulchella.

DORATOMETRA, Klotzsch, in Monatsb. Berl. Acad. (Mærz 1854) 126; et Abh. Akad. Berl. (1854) 63. t. 5 B = Begonia, Linn.
Bowringiana, Seem. Bot. Voy. Herald. 379 = B. lacinata.
Wallichiana, Klotzsch, l. c. = B. Wallichiana.

DORATOPHORA, Lem. in Orb. Dict. v. 119 (1849) = Doryphora, Endl. (Monimiac.).

DORATOXYLON, Thou. sec. Boj. ex Benth. & Hook. f. l. c. (1862). *SAPINDACEAE*, Benth. & Hook. f. l. c.
diversifolia, *Benth. & Hook. f. l. c.*—Maurit.
mauritianum, Thou. ex Baker, Fl. Maurit. 60 = diversifolia.

DORCOCERAS, Bunge, Enum. Pl. Chin. Bor. 54 (1832) = Boea, Comm. (Gesnerac.).
hygrometrica, Bunge, l. c. = B. hygrometrica.

DOREMA, D. Don, in Edinb. Phil. Mag. lx. (1831) 46; et in Trans. Linn. Soc. xvi. (1833) 601. *UMBELLIFERAE*, Benth. & Hook. f. i. 916.
DISERNESTON, Jaub. & Spach, Illustr. i. 78. t. 40 (1848).
Ammoniacum, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 601.—Persia; Afghan.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 329.—Persia.
aureum, *Stocks, in Hook. Kew Journ. Bot.* iv. (1852) 148.—Beluchist.
glabrum, *Fisch. & Mey. ex Ind. Sem. Hort. Petrop.* i. 26.—Armenia.
hirsutum, Loft. in Borcz. Ferul. 28 = D. Ammoniacum.
meifolium, Fenzl, Fugill. Pl. Nov. Syr. 18 = Ferula meifolia.
odoriferum, Loft. ex Borcz. Ferul. 28 (= *Ferulago cordicarpum*?).—Assyria.
robustum, Loft. l. c. = Aucheri.
songaricum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 305.—Songar.

DORIA, Adans. Fam. ii. 124 (1763) = Solidago, Linn. (Compos.).
nemorensis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = Senecio Tournefortii.
paludosa, Fourr. l. c. = Senecio paludosus.
vera, Fourr. l. c. = Senecio Doria.
Virgaurea, Scop. Fl. Carn. ed. II. ii. 176 = Solidago Virgaurea.

DORIA, Thunb. Nov. Gen. xii. 162 (1800) = Othonna, Linn. (Compos.).
abrotanifolia, Harv. in Harv. & Sond. Fl. Cap. iii. 324.—Afr. austr.
acutiloba, DC. Prod. vi. 469 = O. acutiloba.
alata, Thunb. Nov. Gen. 164 = Othonopsis alata.
Arbuscula, DC. Prod. vi. 470 = O. Arbuscula.
bipinnata, Thunb. Nov. Gen. 165 = Senecio pinnatifidus.
campanulata, DC. Prod. vi. 473 = O. campanulata.
carnosa, DC. l. c. 471 = O. pinnatiloba.
Cervaria, Harv. in Harv. & Sond. Fl. Cap. iii. 328.—Afr. austr.
chromochaeta, DC. Prod. vi. 479 = O. chromochaeta.
ciliata, Harv. in Harv. & Sond. Fl. Cap. iii. 322 = Othonopsis ciliata.
clavifolia, DC. Prod. vi. 470 = Othonopsis clavifaea folia.
cneorifolia, DC. l. c. = Othonopsis cneorifolia.
denticulata, Thunb. Nov. Gen. 166.—Afr. austr.
digitata, Jacq. Syst. Syng. 89 = O. digitata.
diversifolia, DC. Prod. vi. 472 = O. diversifolia.
elongata, Thunb. Nov. Gen. 166 = Senecio diversifolius, Harv.
eriocarpa, DC. Prod. vi. 471 = O. eriocarpa.
erosa, Thunb. Nov. Gen. 166 = Cineraria oxyodonta.
glauca, Sch. Bip. in Flora, xxvii. (1844) 696 = Senecio othonnaeiformis.
gymnodiscus, DC. Prod. vi. 472 = O. gymnodiscus.
incisa, Thunb. Nov. Gen. 167 = Senecio incisus.
K-ruasii, Sch. Bip. in Flora, xxvii. (1844) 709 = Othonnopsis Kraussii.
lasiocarpa, DC. Prod. vi. 469 = O. lasiocarpa.
Laureola, DC. l. c. 472 = O. Laureola.

DORIA :—
Lingua, Less. Syn. Comp. 89 = O. Lingua.
linearifolia, DC. Prod. vi. 471 = O. linearifolia.
longipes, Harv. in Harv. & Sond. Fl. Cap. iii. 322.—Afr. austr.
minor, Harv. l. c. 327.—Afr. austr.
nana, DC. Prod. vi. 470 = O. nana.
nivea, Thunb. Nov. Gen. 163 = Senecio niveus.
nodulosa, DC. Prod. vi. 470 = O. nodulosa.
perfoliata, Thunb. Nov. Gen. 164 = O. filicaulis.
pinnatifida, Thunb. l. c. 168 = Senecio incertus.
ramosa, DC. Prod. vi. 472.—Afr. austr.
retrofracta, Less. Syn. Comp. 89 = O. retrofracta.
rigida, Harv. in Harv. & Sond. Fl. Cap. 322 = Senecio spinescens.
semicalva, DC. Prod. vi. 472 = O. semicalva.
serrata, Thunb. Nov. Gen. 165 = Senecio leontodontis.
sonchifolia, Thunb. l. c. 167 = Cineraria sonchifolia.
spinescens, DC. Prod. vi. 470 = O. rhamnoides.
taraxacoides, DC. l. c. = O. taraxacoides.
tortuosa, DC. l. c. 472 = O. tortuosa.
undulata, Thunb. Nov. Gen. 163 = Senecio petiolaris.
undulosa, DC. Prod. vi. 472 = Cineraria sonchifolia.

DORIENA, Endl. Gen. 1148, in syn. (1841) = DOERINGIA, Dennst. = Acronychia, Forst. (Rutac.).

DORITIS, Lindl. Gen. et Sp. Orch. 128 (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 574.
CARTERETIA, A. Rich. Sert. Astrol. 10. t. 4 (1834). bifalcis, *Reichb. f. in Otto, Hamb. Gartenz.* (1860) 116.—N. Guinea.
latifolia, *Benth. & Hook. f. Gen.* iii. 574.—Zeylan.
paniculata, *Benth. & Hook. f. l. c.*—N. Guinea.
pulcherrima, *Lindl. Gen. et Sp. Orch.* 178; *et Wall. Cat.* n. 7848.—Ind. or.
taenialis, *Benth. & Hook. f. Gen.* iii. 574, 577.—Reg. Himal.; Burma.
Wightii, *Benth. & Hook. f. l. c.* 574.—Ind. or.; Burma.

DORNERA, Heuff. ex Schur, Enum. Pl. Transs. 697 (1866) = Carex, Dill. (Cyperac.).
alpina, Heuff. ex Schur, l. c. = C. pyrenaica.

DOROBAEA, Cass. in Dict. Sc. Nat. xlviii. 458 (1827) = Senecio, Toum. (Compos.).
pimpinellaefolia, Cass. l. c. = S. pimpinellaefolius.

DOROBOEA, DC. Prod. vi. 340, in syn. (1837) = praec.

DOROCERAS, Steud. Nom. ed. II. i. 525 (1840) = DORCOCERAS, Bunge = Boea, Comm. (Gesnerac.).

DORONICUM, Tourn. ex Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 440.
ARONICUM, Neck. Elem. i. 27 (1790).
FULLARTONIA, DC. Prod. v. 261 (1836).
GRAMMARTHRON, Cass. in Bull. Soc. Philom. (1817) 32.
acaule, Walt. Fl. Carol. 205 = Arnica radicalis.
altaicum, *Pall. in Act. Acad. Petrop.* (1779) ii. 271. t. 16.—Sibiria.
arabicum, Lam. Encyc. ii. 316 = Gerbera hirsuta.
Arnica, Desf. Tabl. ed. I. 101 = Arnica montana.
Arnottii, DC. in Wight, Contrib. 28 = Senecio polycephalus.
asplenifolium, Lam. Encyc. ii. 315 = Gerbera aspleniifolia, ferruginea.
austriacum, Bieb. Fl. Taur. Cauc. ii. 321 = macrophyllum.
austriacum, *Jacq. Fl. Austr.* ii. 18. t. 180.—Europ.
Bauhini, Saut. ex Reichb. Fl. Germ. Excurs. 324 = hirsutum.
Bellidiastrum, Linn. Sp. Pl. 886 = Aster Bellidiastrum.
Bellidiastrum, Sibth. & Sm. Fl. Graec. Prod. ii. 182 = Bellis sylvestris.
bellidiflorum, Schrank, Reise. Fl. ii. 400 = Stenactis annua.
bellidiflorum, Sallab. Prod. 201 = Aster Bellidiastrum.
Bourgaei, *Sch. Bip. in Bot. Mag.* t. 4994 (Senecionis sp. ?).—Ins. Canar.
cacaliaefolium, *Boiss. & Heldr. Diagn.* Ser. I. xi. 81.—As. Min.
calcaratum, Roxb. Hort. Beng. 61; Fl. Ind. iii. 434 = Vicoa auriculata.
Candolleanum, Wight & Arn. ex Arn. in Nov. Act. Nat. Cur. xvii. (1836) i. 351 = Senecio Hohenackeri.

DORONICUM:—
Candollii, Wight & Arn. ex Wight, Ic. t. 1127 = *Candollesanum*.
carpathicum, Nym. Syll. Suppl. 1 = grandiflorum.
carpetanum, *Boiss. & Reut. ex Willk. & Lange, Prod. Fl. Hisp.* ii. 108.—Hispan.
caucasicum, *Bieb. Fl. Taur. Cauc.* ii. 321.—Europ. austr.; As. Min.
caucasicum, Vis. Fl. Dalmat. ii. 72 = Columnae.
ciliatum, Lam. Encyc. ii. 314 = Hypochoeris grandiflorus.
Clusioni, Poir. Encyc. Suppl. v. 660 = Arnica nudicaulis.
Clusii, *Tausch, in Flora*, xi. (1828) 178.—Mont. Pyren.
Columnae, *Tenore, Fl. Nap. Prod.* 49.—Europ. austr.; As. Min.
cordatum, C. Koch, in Linnaea, xxiv. (1851) 357 = caucasicum.
cordatum, Lam. Fl. Fr. ii. 128 = D. Pardalianches.
cordatum, Sch. Bip. in Oestr. Bot. Wochenbl. iv. (1854) 411 = Columnae.
cordifolium, Stemb. in Denkschr. Acad. Muench. (1813) 147 = Columnae.
cordifolium, Stokes, Bot. Mat. Med. iv. 233 = D. Pardalianches.
corsicum, *Poir. Encyc. Suppl.* ii. 517.—Corsica.
crenatum, Poir. l. c. = Matres crenatae.
croaticum, *Vukot. in Bull. Soc. Bot. Fr.* xxix. (1882) *Bibl.* 6.—Europ. austr.
cruentum, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 236 = Senecio cruentus.
dentatum, *Link, Enum. Hort. Berol.* ii. 337.—Hab.?
echinatum, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 235 = Senecio echinatus.
eriorrhizon, Guss. in Grisch. Spicil. Fl. Rumel. ii. 217 = caucasicum.
Falconeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 333.—Reg. Him. occ.
fulgens, Poir. Encyc. Suppl. v. 660 = Arnica alpina.
Gerardeti, Thw. Enum. Pl. Zeyl. 167 = Senecio Gardneri.
glaciale, *Nym. Syll.* 1.—Hispan.
glutinosum, Willd. Sp. Pl. iii. 2115 = Grindelia glutinosa.
gracile, Schur, Enum. Pl. Transs. 340 = grandiflorum.
grande, Poir. Encyc. Suppl. ii. 517 = Alcíope lanata.
grandiflorum, *Lam. Encyc.* ii. 313.—Europ.
Halleri, Tausch, in Flora, xi. (1828) 180 = grandiflorum.
helveticum, Mill. Gard. Dict. ed. VIII. n. 3 = Senecio Doronicum.
hewerense, Dals. in Dalz. & Gibs. Bomb. Fl. 180 = Senecio hewrensis.
hirsutum, *Lam. Encyc.* ii. 313.—Europ.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 332.—Reg. Himal.
hungaricum, *Reichb. f. Ic. Fl. Germ.* xvi. 84. t. 956.—Europ. austr. or.
incanum, Hill, Veg. Syst. ii. 58 = hirsutum.
incanum, Lam. Encyc. ii. 315 = Tridiocline incana.
incanum, Linn. Sp. Pl. 886 = Senecio Doronicum.
Jacquini, Tausch, in Flora, xi. (1828) 180 = grandiflorum.
japonicum, Lam. Encyc. ii. 314 = Senecio japonicus.
laevifolium, Walt. Fl. Carol. 205 = Erigeron nudicaulis.
lanatum, Poir. Encyc. Suppl. ii. 517 = Alcíope lanata.
Leasingianum, Wight & Arn. ex Arn. in Nov. Act. Nat. Cur. xviii. (1836) t. 350 = Senecio Lessingianus.
linifolium, DC. Prod. vi. 322 = Senecio linifolius.
longifolium, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) 341 = hungaricum.
longifolium, Reichb. Fl. Germ. Excurs. ii. 284 = austriacum.
lucidum, Bernh. ex Pantoč. Adnot. Fl. Herceg. 42 = cordifolium.
macrophyllum, *Fisch. Hort. Gorenk.* ed. II. (1812) 40; *Hornem. Hort. Bot. Hafn. Suppl.* 100.—Reg. Caucas.; Persia.
maloaefolium, Sch. Bip. in Webb & Berth. Phyt. Canar. iii. 230 = Senecio malvaefolius.
maritimum, Lam. Encyc. ii. 314 = Senecio Pseudoarnica.
Matthioli, Tausch, ex Reichb. Fl. Germ. Excurs. ii. 284 = D. Pardalianches.
maximum, *Boiss. & Huet, Diagn.* Ser. II. iii. 81.—Armenia.
mexicanum, Cerr. in Link & Otto, Ic. Pl. Rar. t. 22 = Heterotheca inuloides.
montanum, Lam. Encyc. ii. 313 = Arnica montana.

DORONICUM:—
Nendtveichii, Sadler, in Nentvich, Enum. Pl. Terr. Quin. Eccles. 85. t. 2 = caucasicum.
nudicaule, Michx. Fl. Bor. Am. ii. 121 = Arnica nudicaulis.
oblongifolium, *DC. Prod.* vi. 321.—Reg. Cancas.
oporinum, Poir. Encyc. Suppl. ii. 517 = Olearia operina.
oppositifolium, Lam. Encyc. ii. 313 = Arnica montana.
orientale, Hoffm. in Comm. Soc. Phys. Med. Mosc. t. (1808) 8 = caucasicum.
orientale, Reichb. Fl. Germ. Excurs. i. 256 = Columnae.
Orphanidis, *Boiss. Fl. Orient.* iii. 378.—Macedon.
palmatum, Lam. Encyc. ii. 314 = Senecio Synethesis.
papyraceum, Sch. Bip. in Webb & Berth. Phyt. Canar. iii. 229 = Senecio papyraceus.
Pardalianches, Bourg. ex Rent. & Lange, in Bull. Soc. Bot. Fr. xi. (1864) 46 = carpetanum.
Pardalianches, Lag. ex Willk. & Lange, Prod. Fl. Hisp. ii. 109 = glaciale.
Pardalianches, *Linn. Sp. Pl.* 885.—Europ.
peruvianum, Lam. Encyc. ii. 316 = Werneria rigida.
piloselloides, Lam. l. c. 314 = Gerbera piloselloides.
plantaginifolium, Stokes, Bot Mat. Med. iv. 232 = plantagineum.
plantagineum, *Linn. Sp. Pl.* 885.—Europ.
plantagineum, C. A. Mey. Verz. Pfl. Canc. 80 = oblongifolium.
plantagineum, Poir. Encyc. Suppl. v. 660 = Arnica plantaginea.
plantagineum, Roth, Tent. Fl. Germ. i. 358 = scorpioides.
plantaginifolium, Ledeb. Fl. Ross. ii. 625 = oblongifolium.
procurrens, Dum. Fl. Belg. Prod. 66 = D. Pardalianches.
pyrolaefolium, Lam. Encyc. ii. 315 = Gerbera Burmanni.
ramosum, Walt. Fl. Carol. 205 = Erigeron strigosus.
reticulatum, Bojer, Diagn. Ser. I. iv. 12 = Thiliezi.
retusulatum, Dalz. & Gibs. Bomb. Fl. 180 = Senecio belgaumensis.
reticulatum, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 156; Ic. t. 1151 B = Senecio Grahami.
rotundifolium, Desf. Fl. Atlant. ii. 279. t. 235. f. 1 = Bellium rotundifolium.
Royleï, *DC. Prod.* vi. 321.—Reg. Himal.
rupestre, Wight, Ic. t. 1128 = Senecio Hohenackeri.
scorpioides, C. B. Clarke, Comp. Ind. 169 = Hookeri, Falconeri.
scorpioides, *Lam. Encyc.* ii. 313.—Europ.
scorpioides, Lapeyr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 108 = D. Pardalianches.
scorpioides, Willd. Sp. Pl. iii. 2114 = grandiflorum.
serratum, Poir. Encyc. Suppl. ii. 517 = Gerbera serruginea.
sinuatum, Poir. l. c. = Gerbera serruginea.
spinulosum, Lam. Encyc. ii. 315 = Alcíope tabularis.
stenoglossum, *Maxim. in Bull. Acad. Petersb.* xxvii. (1881) 482.—China.
subalpinum, Schur, in Verh. Siebenb. Ver, Naturw. iv. (1853) 41 = austriacum?
tenuifolium, Wight, Ic. t. 1129 = Senecio tenuifolius.
Thilezii, *Sch. Bip. in Linnaea*, xix. (1847) 35, 314, nomen; *Boiss. Fl. Orient.* iii. 379.—Bithynia.
tomentosum, Wight, Ic. t. 1151 = Senecio Edgeworthii.
torticarpum, Salisb. Prod. 202 = D. Pardalianches.
Tussilaginis, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 231 = Senecio Tussilaginis.
*uginiatum, C. Koch, in Linnaea, xxiv. (1851) 356 = macrophyllum.
Walberi, Arn. in Nov. Act. Nat. Cur. xviii. (1836) t. 350 = Senecio ludens.
Webbii, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 233 (Senecionis sp.).—Ins. Canar.
Wightii, DC. in Wight, Contrib. 23 = Senecio saxatilis.
Wulfenianum, Poir. Encyc. Suppl. v. 660 = Columnae.

DORRIENIA, Endl. Gen. Suppl. ii. 85, in syn. (1842) = DOERRIENIA, Reichb. = Gaudissea, A. St. Hil. (Lamith.).

DORSTENIA, Plum. ex Linn. Gen. ed. I. 336 (1737). *URTICACEAE*. Benth. & Hook. f. iii. 262.
COSARIA, J. F. Gmel. Syst. 71 (1791).
KOSARIA, Forsk. Fl. Aegypt. Arab. 164 (1775).

DORSTENIA:—
SYCMINIUM, Desv. in Mém. Soc. Linn. Par. iv. (1826) 216. t. 12.
Alexiteria, Linn. Syst. ed. X. 899 = D. Houstoni.
appendiculata, *Miq. in Mart. Fl. Bras.* iv. t. 162.—Bras.
argentata, *Hook. f. Bot. Mag.* t. 6795.—Bras.
arifolia, *Lam. Encyc.* ii. 317.—Bras.
asarodes, Gardn. in Hook. Ic. Pl. t. 899 = D. Cayapia.
bahiensis, *Klotzsch, in Linnaea*, xiv. (1840) 268.—Bras.
Barnimiana, *Schweinf. Pl. Quaed. Nilot.* 86. t. 12.—Abyssin.
Bertorii, *Bur. in DC. Prod.* xvii. 272.—Afr. trop.
benguellensis, *Welw. in Trans. Linn. Soc.* xxvii (1869) 71.—Afr. trop.
bicornis, *Schweinf. in Bot. Zeit.* xxix. (1871) 882.—Afr. trop.
Bowmaniana, *Baker, in Saund. Refug. Bot.* v. t. 308.—Bras.
brasiliensis, *Lam. Encyc.* ii. 317.—Bras.
bryoniaefolia, Mart. in Flora, xx. (1837) ii. Beibl. 128, nomen; Miq. in Mart. Fl. Bras. iv. t. 167 = D. Cayapia.
caulescens, Linn. Sp. Pl. 121 = Procris cuinquenervis.
caulescens, Poir. ex Bur. in DC. Prod. xvii. 277 (alius ordinis).—Hab.?
caulescens, Vell. Fl. Flum. 53; t. t. 141 = broeolata.
Cayapia, *Vell. Fl. Flum.* 53; t. t. 137.—Bras.
Ceratosanthes, Lodd. Bot. Cab. t. 1216 = arifolia.
chinensis, Lour. Fl. Cochinch. 90 = Procris chinensis.
Contrajerva, *Linn. Sp. Pl.* 121.—Am. trop.
cordifolia, Mart. ex Miq. in Mart. Fl. Bras. iv. t. 165 = bahiensis.
cordifolia, *Sw. Prod. Veg. Ind. Occ.* 94.—Jamaic.
crassipes, *Griseb. Cat. Pl. Cub.* 58.—Cuba.
crenulata, *C. Wright, ex Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 178.—Cuba.
cuspidata, *Hochst. in Flora*, xxvii. (1844) 108.—Abyssin.
Cyperus, Vell. Fl. Flum. 53; t. t. 140 = arifolia.
digitifolia, Stokes, Bot. Mat. Med. iv. 241 = Procris chinensis.
Drakena, *Linn. Syst.* ed. X. 899.—Mexic.
Drakena, Vell. Fl. Flum. 53; t. t. 139 = arifolia.
elata, *Gardn. in Hook. Ic. Pl.* t. 220.—Bras.
elliptica, *Bur. in DC. Prod.* xvii. 271.—Afr. trop.
erecta, *Vell. Fl. Flum.* 53; t. t. 142.—Bras.
erythrantha, *Griseb. Cat. Pl. Cub.* 58.—Cuba.
exaltata, Herb. Mus. Vindob. ex Miq. in Mart. Fl. Bras. iv. t. 161 = plumeriaefolia.
excentrica, *Moric. Pl. Nouv. Am.* 92. t. 59.—Mexic.
Faria, *Paiva, ex Spreng. in Schrad. Journ.* (1800) ii. 239 (sp. dub.).—Bras.
ficifolia, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 58 = arifolia.
Ficus, Vell. Fl. Flum. 53; t. t. 138 = arifolia.
Fischerii, *Bur. in DC. Prod.* xvii. 277.—Bras.
fluminensis, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 491 = arifolia.
frutescens, Schott, ex Endl. Cat. Hort. Vindob. i. 241 = erecta.
Gigas, *Schweinf. ex Baif. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 93.—Ins. Socotra.
glauca, Pohl, ex Miq. in Mart. Fl. Bras. iv. t. 162 = nervosa.
Griffithiana, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 104.—Burma.
hirta, Desv. in Mém. Soc. Linn. Par. iv. (1826) 218 = erecta.
hispida, Hook. Ic. Pl. sub t. 220 = erecta.
Houstoni, Linn. Sp. Pl. ed. II. 176 = D. Contrajerva.
Hydrocotyle, Mart. ex Miq. in Mart. Fl. Bras. iv. t. 168 = D. Cayapia.
indica, *Wall. Cat.* n. 4689.—Ind. or.
infundibuliformis, Lodd. Bot. Cab. xx. Ind. = tubicina.
ingoidata, *Bur. in DC. Prod.* xvii. 268.—Bras.
Langsdorffiana, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 64 = broeolata.
Lindmana, *Bur. in DC. Prod.* xvii. 269.—Mexic.
longifolia, *Moric. Pl. Nouv. Am.* 90. t. 58.—Mexic.
longipes, Mart. ex Bur. in DC. Prod. xvii. 268 = bahiensis.
lucida, Forst. f. Prod. 11 = Procris cephalicha.
maculata, Lam. Illustr. Hortin. x. (1868) t. 362 = D. Contrajerva.
membranacea, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 59 = arifolia.

DORSTENIA :—
Mannii, *Hook. f. Bot. Mag.* t. 5908.—Afr. trop.
Martiana, Miq. in Mart. Fl. Bras. iv. i. 165 = bahiensis.
mexicana, Benth. Pl. Hartw. 51 = D. Drakena.
minor, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 63 = erecta.
montevidensis, Field. & Gardn. Sert. Pl. sub t. 13 = brasiliensis.
morifolia, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 60 = D. Cayapia.
multiformis, Miq. in Mart. Fl. Bras. iv. i. 165 = arifolia.
nervosa, *Desv. in Mém. Soc. Linn. Par.* iv. (1826) 217. —Bras.
obovata, *Hochst. in Flora*, xxvii. (1844) 103.—Abyss.
ophioglossoides, *Hochst. ex Bur. in DC. Prod.* xvii. 276.—Afr. trop.
opifera, *Fisch. ex Regel, Gartenfl.* (1856) 87.—Bras.
opifera, Mart. Reise Bras. ii. 787 = D. Cayapia.
opponitifolia, Soland. ex Seem. Fl. Vit. 236 = Lecanthus Solandri.
ovalis, Stokes, Bot. Mat. Med. iv. 838 = D. Drakena.
palmata, Pohl, ex Miq. in Mart. Fl. Bras. iv. i. 167 = D. Cayapia.
palmata, Willd. ex Schult. Mant. iii. 317 = D. Contrajerva.
paradoxa, Herb. Berol. ex Bur. in DC. Prod. xvii. 261 = arifolia.
pedunculata, Auct. ex Steud. Nom. ed. II. i. 526 = Procris lucida.
peltata, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 66 = Fischeri.
peltata, *Spreng. Neue Entdeck.* iii. 22.—Ins. S. Doming.
pernambucana, *Arruda, in Koster, Trav. Bras.* 498.— Bras.
petraea, *Wright, ex Griseb. Cat. Pl. Cub.* 58.— Cuba.
picta, *Bur. in DC. Prod.* xvii. 277.—Afr. trop.
placentiodes, Comm. ex Lam. Encyc. ii. 317 = brasiliensis.
plumeriaefolia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 65.—Bras.
Pallurus, *Welw. in Trans. Linn. Soc. xxvii.* (1869) 71. —Afr. trop.
pubescens, Blanco, Fl. Filip. ed. I. 692 = Elatostema obtusum.
pubescens, Forst. f. Prod. 11 = Elatostema sessile.
quadrangularis, Stokes, Bot. Mat. Med. iv. 838 = D. Contrayerva.
quadrata, Desv. in Mém. Soc. Linn. Par. iv. (1826) 218 = arifolia.
radiata, *Lam. Encyc.* iii. 318.—Arabia.
reniformis, Pohl, ex Miq. in Mart. Fl Bras. iv. i. 168 = D. Cayapia.
Riedeliana, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 58 = arifolia.
rotundifolia, *Arruda, in Koster, Trav. Bras.* 498.— Bras.
setosa, *Moric. Pl. Nouv. Am.* 103. t. 64.—Bras.
Sychinsum, Steud. Nom. ed. II. i. 526 = arifolia.
tenera, *Bur. in DC. Prod.* xvii. 271.—Afr. trop.
tentaculata, Fisch. & Mey. Ind. Sem. Hort. Petrop xi. 59 = arifolia.
tenuis, *Bonpl. ex Bur. in DC. Prod.* xvii. 264.—Reg. Argent.
tomentosa, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 61 = brasiliensis.
tropaeolifolia, *Bur. in DC. Prod.* xvii. 276.—Abyss.
tuberosa, *Wright, ex Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 178.—Cuba.
tubicina, *Ruiz & Pav. Fl. Per.* i. 65. t. 102.— Peruv.
turmacefolia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 63.—Bras.
urceolata, *Schott, in Flora*, iv. (1821) 197.—Bras.
vermiciaefolia, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 63 = erecta.
Vitellia, Paiva, ex Spreng. in Schrad. Journ. (1800) ii. 389 = D. Cayapia ?
vitifolia, *Garden. in Field. Sert. Pl.* t. 14 = D. Cayapia.
vivipara, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 70. —Afr. trop.
volubilis, Walp. Ann. i. 786 (alius ordinis).—Java.

DORTMANIA, Neck. Elem. i. 189 (1790) = seq.

DORTMANNA, Linn. Syst. ed. I (1735) = **Lobelia**, Linn. (Campanulac.).
lacustris, G. Don, Gen. Syst. iii. 715 = L. Dortmanna.
paludosa, G. Don, l. c. = L. paludosa.
palustris, G. Don, in Sweet, Hort. Brit. ed. III. 425 = L. paludosa.

DORTMANNIA, Steud. Nom. ed. II. i. 526 (1840) = prasc.

DORVALIA, Comm. ex DC. Prod. iii. 36 (1828) = seq.

DORVALLA, Comm. ex Lam. Encyc. ii. 565 (1786) = **Fuchsia**, Linn. (Onagrar.).
eucharis, Comm. ex Lam. l. c. = F. magellanica.

DORVALLIA, Comm. ex Juss. Gen. 320 (1789) = prasc.

DORYANTHES, Correa, in Trans. Linn. Soc. vi. (1802) 211. t. 25, 24. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 739.
excelsa, *Correa, l. c.* 213.—Austral.
Palmeri, *W. Hill, ex Benth. Fl. Austral.* vi. 452.— Austral.

DORYCHEILE, Reichb. Nom. 56 (1841) = **Cephalanthera**, Rich. (Orchid.).
rubra, Fuss, Fl. Transs. 628 = C. rubra.

DORYCHNIUM, Brongn. Enum. Gen. 128 (1843) = **Doryenium**, Tourn. (Legumin.).

DORYCHNIUM, Royen, ex Moench, Meth. 109 (1794) = **Psoralea**, Linn. (Legumin.).
angustifolium, Moench, l. c. 110 = P. bituminosa.
bracteatum, Moench, Meth Suppl. 37 = P. bracteata.
corylifolium, Moench, Meth. 109 = P. corylifolia.
rotundifolium, Moench, Meth. Suppl. 37 = P. palestina.

DORYCNIUM, Linn. Syst. ed. I (1735); Gen. ed. I. 228 (1737): Vill. Hist. Pl. Dauph. iii. 416 (1789). *LEGUMINOSAE*, Benth. & Hook. f. i. 490.
BONJEANA, Reichb. Fl. Germ. Excurs. 507 (1832).
affine, Boreau, Gnin. Rec. Jard. Grenoble, (1849) 14 = herbaceum.
ambiguum, Poepp. ex Benth. Sem. Hort. Erf. (1837); ex Linnaea, xii. (1838) Litt. 76 (= *Anisolotus Poeppeianus*).—Chili.
anatolicum, *Boiss. & Heldr. Diagn.* Ser. I. ix. 31.— As. Min.
argenteum, Delile, Fl. Egypte, 257. t. 40 = Lotus argenteus.
Broussonetii, *Webb & Berth. Phyt. Canar.* ii. 90.—Afr. bor.
calycinum, *Stocks, in Hook. Kew Journ.* iv. (1852) 145. —Beluchist.
caucasicum, Regel, Ind. Sem. Hort. Petrop. (1856) 49, sphalm. = Doronicum caucasicum.
cinerascens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 22 = suffruticosum.
collinum, Jord. & Fourr. l. c. 23 = suffruticosum.
decumbens, Jord. Obs. Pl. Crit. iii. (1846) 60 = herbaceum.
diffusum, Janka, in Oestr. Bot. Zeitschr. xiii. (1863) 316 = herbaceum.
dispermum, *Ser. in DC. Prod.* ii. 209.—Europ. austr.
dolichocarpum, Clav. in Actes Linn. Soc. Bord. xxxvii. (1884) 234 = suffruticosum.
dumulosum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 22 = suffruticosum.
elongatum, Jord. & Fourr. l. c. 21 = suffruticosum.
eriophthalmum, *Webb & Berth. Phyt. Canar.* ii. 88.— Ins. Canar.
frutescens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 21 = suffruticosum.
fruticosum, Pers. Syn. ii. 354 = suffruticosum.
gracile, Jord. Obs. Pl. Crit. iii. (1846) 70 = herbaceum.
graecum, Ser. in DC. Prod. ii. 208 = latifolium.
Haussknechtii, Boiss. Fl. Orient. ii. 163 = libanoticum.
herbaceum, *Vill. Prosp.* 44; *Hist. Pl. Dauph.* iii. 416. —Europ. austr.; As. Min.
hirsutum, *Ser. in DC. Prod.* ii. 208.—Europ. austr.
hirtum, Poepp. ex Benth. Sem. Hort. Erf. (1837); ex Linnaea, xii. (1838) Litt. 76 = Hosackia subpinnata.

DORYCNIUM :—
humile, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 24 = suffruticosum.
ibericum, Willd. Enum. Hort. Berol. Suppl. 52 = latifolium.
implexum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 21 = suffruticosum.
insulare, Jord. & Fourr. l. c. 28 = suffruticosum.
intermedium, Ledeb. Ind. Sem. Hort. Dorp. (1820) 14 = herbaceum.
Jordani, Loret & Barr. Fl. Montp. 175 = herbaceum.
Jordanianum, Willk. in Willk. & Lange, Prod. Fl. Hisp iii. 336 = prasc.
Kotschyanum, Boiss. Fl. Orient. ii. 163 = libanoticum. Kotschyi, *Boiss. l. c.* 161.—Cilicia.
latifolium, *Willd. Sp. Pl.* iii. 1397.—Graecia; As. Min. libanoticum, *Boiss. Diagn.* Ser. II. ii. 19.—Syria.
lutescens, *Steud. Nom.* ed. II. i. 526.—Abyss.
microcarpum, Ser. in DC. Prod. ii. 209 = Lotus parviflorus.
microphyllum, Sweet, Hort. Brit. ed. II. 137 = Indigofera gracilis.
monspelliense, Bieb. Fl. Taur. Cauc. ii. 221 = pentaphyllum.
monspelliense, Willd. Sp. Pl. iii. 1396 = suffruticosum.
parviflorum, Ser. in DC. Prod. ii. 208 = Lotus parviflorus.
pentaphyllum, Reichb. Fl. Germ. Excurs. 507 = suffruticosum.
pentaphyllum, *Scop. Fl. Carn.* ed. II. ii. 87.—Ross. austr.
procumbens, Lapeyr. Hist. Abr. Pl. Pyr. 441 = Anthyllis Gerardi.
rectum, *Ser. in DC Prod.* ii. 208.—Reg. Mediterr.
sabaudum, Reichb. Fl. Germ Excurs. 507 = herbaceum. *Schimperi*, Jaub. & Spach, Illustr. v. 82. t. 473 = Lotus brachycarpus.
sericeum, Sweet, Hort. Brit. ed. I. 476 = hirsutum.
spectabile, *Webb & Berth. Phyt. Canar.* ii 89.—Ins. Teneriffa.
sphaerocarpum, Clav. in Actes Linn. Soc. Bord. xxxvii. (1884) 524 = suffruticosum.
stenocladium, Jord. & Fourr. Brev. Pl. Nov. fasc ii. 28 = suffruticosum.
strophylium, Schur, Enum. Pl. Transs. 159 = suffruticosum.
subbiflorum, *Ser. in DC. Prod.* ii. 209.—Hispan.
suffruticosum, Griseb. Spicil. Fl. Rumel. i. 41 = herbaceum
suffruticosum, *Vill. Hist. Pl. Dauph.* iii. 417. t. 41.— Europ. austr.
tomentosum, G. Don, in Loud. Hort. Brit. 300 = hirsutum.
torulosum, *Presl, Symb. Bot.* i. 19.—Ins. Teneriffa.
vestitum, Boiss. Fl. Orient. ii. 162 = latifolium.

DORYCINOPSIS, Lem. in Orb Dict. v. 118 (1849) = seq.

DORYCNOPSIS, Boiss. Voy. Bot. Espagne, ii. 163 = **Anthyllis**, Linn. (Legumin.).
Gerardi, Boiss. l. c. 164 = A. Gerardi.

DORYCTANDRA, Hook. f. & Thoms. in Fl. Brit. Ind. i. 186 (1872) = DIORYKTANDRA, Hassk. = **Alsodeia**, Thou. (Violar.).

DORYDIUM, Salisb. Gen. Pl. Fragm. 72 (1866) = **Asphodeline**, Reichb. (Liliac.).
creticum, Saliab. l. c. = A. liburnica.
liburnicum, Saliab. l. c. = A. Taurica.
lutcum, Saliab. l. c. = A. lutea.

DORYPHORA, Endl. Gen. 815 (1837). *MONIMIACEAE*, Benth. & Hook. f. iii. 145.
DORYPTOPHORA, Lem. in Orb. Dict. v. 119 (1849).
LEAROSA, Reichb. Nom. 69 (1841).
austro-caledonica, *Seem. Fl. Vit.* 208.—N. Caled.
Sassafras, *Endl. Ic. Gen. Pl.* t. 10.—Austral.
Vieillardi, *Baill. Adansonia*, ix. (1868–70) 138.—N. Caled.

DORYSTAECHAS, Boiss. & Heldr. ex Benth. in DC. Prod. xii. 261 (1848). *LABIATAE*, Benth. & Hook. f. ii. 1194.
hastata, *Boiss. & Heldr. ex Benth. l. c.*—Lycia.

DORYSTIGMA, Gaudich. Bot. Voy. Bonite, t. 18 (1844–66) [1843] = **Fandanus**, Linn.
madagascariense, Gaudich. l. c. t. 21 = P. madagascariensis.
mauritianum, Gaudich. l. c. t. 18 = P. conglomeratus.

DORYSTIGMA, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 547 = **Jaborosa**, Jus. (Solanac.).
caulescens, Miers, l. c. 548 = J. caulescens.
crispum, Miers, l. c. vii. (1848) 338.—Bolivia.
squarrosum, Miers, l. c. iv. (1845) 548.—Chili.

DORYXYLON, Zoll. in Tijdschr. Nederl. Ind. xiv. (1867) 172 = **Sumbavia**, Baill. (1858) (Euphorb.).
spinosum, Zoll. l. c. = S. rottleroides.

DOSSINIA, C. Morr. in Ann. Soc. Gand, iv. (1848) 171. *ORCHIDEAE*, Benth. & Hook. f. iii. 601.
cristata, Miq. Fl. Ind. Bat. iii. 731 = Hetaeria cristata.
lancelotata, Lindl. in Journ. Linn. Soc. i. (1857) 186 = Hetaeria lancelota.
marmorata, Lindl. l. c. 185 = Anaectochilus brevilabris.
marmorata. *C. Morr. in Ann. Soc. Gand*, iv. (1848) 171.—Java.
Meinertl, *Hort. Makoy, ex E. Morr. in Belg. Hortic.* xxxiv. (1884) 288.—Sumatra.
obliqua, Miq. Fl. Ind. Bat. iii. 731 = Hetaeria obliqua.

DOTHILIS, Rafin. Fl. Tellur. ii. 60 (1836) = **Spiranthes**, Rich. (Orchid.).
grandiflora, Rafin. l. c. = S. grandiflora.

DOTHILOPHIS, Rafin. Fl. Tellur. iv. 89 (1836) = **Epidendrum**, Linn. (Orchid.).
purpurea, Rafin. l. c. = E. Skinneri.

DOUARREA, Montr. in Mém. Acad. Lyon, x. (1860) 223. *INC. SED.* Cf. Benth. & Hook. f. ii. 20.
alba, *Montr. l. c.* 228.—N. Caled.
speciosa, *Montr. l. c.* 222.—N. Caled.

DOUEPEA, Cambess. in Jacquem. Voy. Bot. 18. t. 18 (1837) = **Moricandia**, DC. (Crucif.).
tortuosa, Cambess. l. c. = M. tortuosa.

DOUEPIA, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 172 = prnec.

DOUGLASIA, Lindl. in Brande, Journ. Sc. (Oct. 1827) 385. *PRIMULACEAE*, Benth. & Hook. f. ii. 632.
GREGORIA, Duby, Bot. Gall. i. 333 (1828).
MACROTYRUS, Dulac, Fl. Hautes-Pyr. 425 (1867).
arctica, *Hook. Fl. Bor. Am.* ii. 120.—Am. arct.
dentata, S. Wats. in Proc. Am. Acad. xvii. (1881–82) 374 = nivalis.
laevigata, *A. Gray, l. c.* xvi. (1881) 105.—Am. bor. occ.
montana, *A. Gray, l. c.* vii. (1868) 371.—Am. bor. occ.
nivalis, *Lindl. in Brande, Journ. Sc.* (Oct. 1827) 388.—Am. bor. occ.
Vitaliana, *Benth. & Hook. f. Gen.* ii. 632.—Mont. Pyren.

DOUGLASSIA, Adans. Fam. ii. 200 (1763) = **Clerodendron**, Linn. (Verbenac.).

DOUGLASSIA, Reichb. Consp. 128 (1828) = **Douglasia**, Lindl. (Primul.).

DOUGLASSIA, Schreb. Gen. Pl. 809 (1791) = **Aiouea**, Aubl. (Laurin.).
laurina, Sm. in Rees, Cyclop. xii = A. guianensis.

DOUMA, [Poir.] in Duham. Arb. ed. Nov. iv. 47 (1809) = **Hyphaene**, Gaertn. (Palm.).
thebaica, [Poir.] l. c. 48 = H. thebaica.

DOUPEA, D. Dietr. Syn. Pl. iii. 684, 76 (1843) = DOUEPIA, Cambess. = **Moricandia**, DC. (Crucif.).

DOVEA, Kunth, Enum. Pl. iii. 457 (1841). *RESTIACEAE*, Benth. & Hook. f. iii. 1081.
binata, Steud. Syn. Pl. Cyp. 248 = Hypodiscus binatus.
Boissi, *Mast. in Journ. Linn. Soc.* xxi. (1885) 576.—Afr. austr.
DOVEA (*continued*)—
ebracteata, *Kunth, Enum. Pl.* iii. 458.—Afr. austr.
Hookeriana, *Mast. in Journ. Linn. Soc.* x. (1869) 249.—Afr. austr.
macrocarpa, *Kunth, Enum. Pl.* iii. 458.—Afr. austr.
microcarpa, Kunth, l. c. 459 = ebracteata.
mucronata, *Mast. in Journ. Linn. Soc.* x. (1869) 251.—Afr. austr.
paniculata, *Mast. l. c.* xxi. (1885) 577.—Afr. austr.
racemosa, *Mast. l. c.* 578.—Afr. austr.
thyrsoidea, *Mast. l. c.* x. (1869) 251.—Afr. austr.
tectorum, *Mast. l. c.* 249.—Afr. austr.

DOVYALIS, E. Mey. ex Arn. in Hook. Journ. Bot. iii. (1841) 251. *BIXINEAE*, Benth. & Hook. f. i. 138.
celastroides, *Sond. in Linnaea*, xxiii. (1850) 12.—Afr. austr.
Monteae, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 339.—Madag.
rhamnoides, Burch. ex Harv. & Sond. Fl. Cap. i. 69 = zizyphoides.
rotundifolia, Harv. l. c. 70 = celastroides.
zizyphoides, *E. Mey. ex Arn. in Hook. Journ. Bot.* iii. (1841) 251.—Afr. austr.

DOWEA, Steud. Syn. Pl. Cyp. 248 (1855) = **Dovea**, Kunth (Restiac.).

DOWNINGIA, Torr. in Pacif. Rail. Rep. iv. 116 (Bot. Whipple Exped. 60) (1856). *CAMPANULACEAE*, Benth. & Hook. f. ii. 550.
BOLELIA, Rafin. Atl. Journ. 120 (1832).
CLINTONIA, Dougl. ex Lindl. in Bot. Reg. t. 1241 (1829).
GYNAMPHIA, Rafin. Fl. Tellur. iii. 5 (1836).
elegans, *Torr. in Bot. U. St. Expl. Exped.* ii. 375.—Am. bor. occ.
pulchella, *Torr. in Pacif. Rail. Rep.* iv. 116.—Am. occ.
pusilla, *Torr. in Bot. U. St. Expl. Exped.* ii. 375.—Chili.

DOXANTHA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 188 = **Bignonia**, Linn.
acutistipula, Miers, l. c. 190 = B. acutistipula.
adunca, Miers, l. c. 188 = B. Unguis-cati.
caprolata, Miers, l. c. 190 = B. caprreolata.
chalophora, Miers, l. c. = B. Unguis-cati.
cuspidata, Miers, l. c. nomen.—Reg. Argent.
elongata, Miers, l. c. nomen.—Bras.
exilis, Miers, l. c. nomen.—Reg. Argent.
exalata, Miers, l. c. = B. exalata.
ferruginea, Miers, l. c. = B. ferruginea.
hamigera, Miers, l. c. nomen.—Bras.
inaequa, Miers, l. c. nomen.—Reg. Argent.
lanceolata, Miers, l. c. = B. lanceolata.
lanuginosa, Miers, l. c. = B. lanuginosa.
limbata, Miers, l. c. nomen.—Bras.
longistipulata, Miers, l. c. = B. longistipulata.
mexicana, Miers, l. c. = B. Unguis-cati.
praezigni, Miers, l. c. nomen.—Ins. S. Doming.
radicans, Miers, l. c. = B. Unguis-cati.
scitiflora, Miers, l. c. nomen.—Ins. Antigua.
serratula, Miers, l. c. nomen.—Ins. Guadalup.
speciosa, Miers, l. c. nomen.—Patagon.
tanacetifolia, Miers, l. c. nomen.—Reg. Argent.
tenuicula, Miers, l. c. nomen.—Ins. S. Doming.
Tweediana, Miers, l. c. = B. Tweediana.
torquata, Miers, l. c.—Reg. Argent.
uncata, Miers, l. c. = Spathodea uncata.
unguicalata, Miers, l. c. = B. Unguis-cati.
Unguis, Miers, l. c. = B. Unguis-cati.

DOXANTHES, Rafin. Fl. Tellur. iv. 51 (1836) = **Alpinia**, Linn. (Scitamin.).

DOXEMA, Rafin. Fl. Tellur. iv. 75 (1836) = **Ipomoea**, Linn. (Convolv.).

DOXOMMA, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 98 = **Barringtonia**, Forst. (Myrtac.).
acuminatissima, Miers, l. c. 102 = B. macrostachya.
angustatum, Miers, l. c. 106 = B. augusta.
cochlearifolium, Miers, l. c. 101.—Cochin.
cylindrostachyum, Miers, l. c. 100 = B. macrostachya.
macrostachyum, Miers, l. c. 104 = B. macrostachya.
DOXOMMA (*continued*)—
magnificum, Miers, in Trans. Linn. Soc. . II. i. (1875) 106 = Barringtonia augusta.
neo-caledonicum, Miers, l. c. 108 = B. neo-caledonica.
pendulum, Miers, l. c. 99 = B. macrostachya.
rigidum, Miers, l. c. 104 = Symplocos rigida.
surcostachys, Miers, l. c. 103 = B. macrostachya.
sumatranum, Miers, l. c. 103 = B. sumatrana.
Vriesii, Miers, l. c. 106 = B. Vriesei.

DOXOSMA, Rafin. Fl. Tellur. iv. 9 (1836) = **Epidendrum**, Linn. (Orchid.).
gracilis, Rafin. l. c. = E. gracile.

DOYERREA, Grosourdy, ex Bello, in Anal. Soc. Esp Hist. Nat. x. (1881) 278. *MYRTACEAE*.
emeto-cathartica, *Grosourdy, ex Bello, l. c.*—Porto Rico.

DRABA, Dill. ex Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 74.
AIZODRABA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 285.
DOLLINERIA, Santer, in Flora, xxxv. (1852) 353.
HOLARGMA, Ehrh. Beitr. iv. 148 (1789).
HOLARGIDIUM, Turcz. in Ledeb. Fl. Ross. i. 156 (1842).
MORNCHIA, Roth, Tent. Fl. Germ. i. 278 (1788), partim.
PETROCALLIS, R. Br. in Ait. Hort. Kew. ed. II. iv. 98 (1812).
TORBORTINA, Rafin. Neogenyt. 2 (1825).
acaulis, *Beiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 164.—Cilicia.
Adamsii, Ledeb. Fl. Ross. i. 147 = alpina.
affinis, *Hook. f. Fl. Antarct.* 285.—N. Zealand.
affinis, Host, Fl. Austr. ii. 288 = aizoides.
afinis, *Ledeb. Fl. Ross.* i. 148.—Sibir.
aizoides, Sr. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = aizoides.
aizoides, Cav. Desc. 421 = hispanica.
aizoides, Curt. Bot. Mag. t. 170 = brachystemon.
aizoides, *Linn. Mant.* i. 91.—Europ.
aizoides, Pall. ex Bieb. Fl. Taur. Cauc. ii. 92 = cuspidata.
aizoides, Sibth. & Sm. Fl. Graec. Prod. ii. 4 = olympica.
Aizoon, *Wahlenb. Fl. Carpat.* 198, in nota.—Europ.
aizoonia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = pyrenaica.
alba, Gilib. Fl. Lituan. ii. 47 = Erophila vulgaris.
Alberti, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 327.—Turkestan.
algida, Adams, ex DC. Syst. ii. 337 = alpina.
alpestris, Jord. Diagn. i. 204 = aizoides.
alpicola, Klotzsch, in Bot. Ergeb. Waldem. Reise, 198 = alpina.
alpina, Asso, Syn. Stirp. Arag. 88 = hispanica.
alpina, Baumg. Enum. Stirp. Transs. ii. 281 = Smelowskia alpina.
alpina, Cav. in Roem. Arch. ii. 227 = cuspidata.
alpina, Crantz, Inst. ii. 82 = aizoides.
alpina, *Linn. Sp. Pl.* 642.—Reg. bor. et arct.
altaica, Bunge, Delect. Sem. Hort. Dorp. (1841) 8 = fladnizensis.
alyssoides, Benth. Pl. Hartw. 159 = Hallii.
alyssoides, *H. B. & K. Nov. Gen. et Sp.* v. 79.—Am. austr.
ambigua, *Ledeb. Fl. Ross.* i. 151.—Sibir. altaic.
ambigua, Schur, Enum. Pl. Transs. 66 = pamorosa.
americana, Pers. Syn. ii. 190 = Erophila vulgaris.
andina, Phil. in Linnaea, xxviii. (1856) 669.—Chili.
androsacea, Baumg. Enum. Stirp. Transs. ii. 284 = Kotschyi.
androsacea, Willd. Sp. Pl. iii. 428; Wahlenb. Fl. Lapp. 174. t. 11. f. 5 = fladnizensis.
araliifolia, Hoppeworth, Reisen, ii. 195 = Azalea vochinensis.
Arabis, Pers. Syn. ii. 190 = arabisans.
arabisans, *Michx. Fl. Bor. Am.* ii. 36.—Am. bor.
arabisans, Pursh, Fl. Am. Sept. ii. 434 = ramosissima.
arctoides, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 227.—Peruv.
aureation, *Kunze, Fl. Cauc.* 114, in texts.—Mons Aureat.
aureonana, Phil. in Anal. Univ. Chil. (1872) 673 = Gilliesii.
Arbuscula, *Hook. f. Fl. Antarct.* 287.—Venezuela.

DRABA :—
arctica, J. Vahl, in Fl. Dan. t. 2294 = hirta.
aretioides, *H. B. & K. Nov. Gen. et Sp.* v. 77. t. 485.—Ecuador.
argaea, Kotschy, ex Boiss. Fl. Orient. i. 296 = natolica.
argentea, Lam. Encyc. ii. 329 = Alyssum argenteum.
armata, Schott, Nyman, & Kotschy, Analect. Bot. 49 = longirostra.
armena, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 167.—Armenia.
aspera, *Adams, ex DC. Syst.* iii. 387.—Sibiria.
aspera, Bertol. Amoen. Ital. 384 = aizoides.
asprella, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 125.—Arizona.
astyla, Bunge, ex Schrenk, Reise Nordost. Eur. Russl. ii. 48 = alpina.
athoa, *Boiss. Diagn.* Ser. II. i. 38.—Graecia.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 329.—Algeria.
Aucheri, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 166 = Arabis sulphurea.
aurea, C. Koch, in Linnaea, xix. (1847) 58 = olympica.
aurea, *Vahl, ex Hornem. Plantel.* ed. II. 599 ; *et in Fl. Dan.* t. 1460.—Reg. bor. et arct.
aureola, *S. Wats. Bot. Calif.* ii. 480.—Calif.
australis, *Hook. f. Fl. Antarct.* 285.—Patagonia.
austriaca, Crantz, Stirp. Austr. ed I. fasc. i. 10 = stellata.
batava, *Dum. Fl. Belg. Prod.* 120.—Batav.
Baumgartenі, Schur, Enum. Pl. Transs. 65 = D. Aizoon.
Beckeri, *A. Kern. Sched. Fl. Exsicc.* iii. 76.—Austria.
Bertolonii, Nym. Cónsp. 52 = aizoides.
bifolia, Mühl. Ind. Fl. Lanc. 173 = carolininana !
Boerhaavii, Van Hall, Spec. Bot. 149 = Erophila vulgaris.
Bonplandiana, H. B. & K. Nov. Gen. et Sp. v. 78 = violacea.
borealis, DC. Syst. ii. 342 = incana.
brachycarpa, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 108.—Am. bor.
brachycarpa, Zett. in Bot. Notiser (1854), 151 = stellata.
brachystemon, *DC. Syst.* ii. 334.—Hab. ?
brachystemonis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = prasc.
Brackenridgei, *A. Gray, Bot. U. St. Expl. Exped.* i. 53. —Bolivia.
brunifolia, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 328 = olympica.
bryoides, DC. Syst. ii. 335 = rigida.
caesia, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 108 = stellata.
caespitosa, Hoffm. Hort. Monç. (1808) n. 1225 ; ex Ledeb, Fl. Ross. i. 145 = rigida.
calycina, Boiss. & Bal. Diagn. Ser. II. vi. 14 = calycosa.
calycina, *Desv. Journ. Bot.* iii. (1814) 185.—Peruv.
calycosa, *Boiss. & Bal. ex Boiss. Fl. Orient.* i. 299.—As. Min.
camptocarpa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 11. 299.—Ecuador.
canadensis, Brunet, Pl. Canad. i. 22 ; ex S. Wats. Bibliog. Ind. N. Am. Bot. 55 = arabisans.
cantabrica, Willk. in Flora, xxxiv. (1851) 587 = aizoides.
cappadocica, *Boiss. & Bal. Diagn.* Ser. II. vi. 14.—Cappadoc.
carinthiaca, Hoppe, in Flora, vi. (1823) 487 = sibiquosa.
carnica, Scop. Fl. Carn. ed. II. ii. 6 = Thlaspi montanum.
carnolica, Vitm. Summa, Pl. iv. 13 = Thlaspi montanum.
carolininana, *Walt. Fl. Carol.* 174.—Am. bor.
caucasica, *Spreng. ex Steud. Nom.* ed. I. 264, nomen. —Reg. Caucas.
Chamissonі, G. Don, Gen. Syst. i. 184 = tomentosa.
cheiranthifolia, Lam. Encyc. ii. 328 = Alyssum incanum.
chetranthoides, *Hook. f. Fl. Antarct.* 286.—N. Granat.
chrisfolia, Berg, ex DC. Syst. ii. 321 = Alyssum incanum.
chrisformis, Lam. Fl. Fr. ii. 462 = Alyssum incanum.
chilloensis, *Phil. in Anal. Univ. Chil.* ii. (1862) 877. —Chili.
chrysanthe, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 364.—Am. bor. occ.
ciliaris, Host, Syn. Fl. 354 = fladnizensis.
ciliaris, Kit. ex Pesl, Bot. Bemerk. 9 = Kitaibeliana.

DRABA :—
ciliaris, Linn. Mant. 91 = brachystemon.
ciliaris, Salisb. Prod. 366 ; DC. Fl. Fr. iv. 697 = aizoides.
ciliaris, Schrank, Baier. Fl. ii. 177 = D. Aizoon.
ciliata, *Scop. Fl. Carn.* ed. II. ii. 6.—Austria.
cinerea, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 103.—Sibiria.
clypeata, Lam. Encyc. ii. 328 = Farsetia clypeata.
cochlearioides, Langsd. ex DC. Syst. ii. 365 = Cochlearia officinalis.
cochlearioides, *Turcz. ex Bess. in Flora*, xvii. (1834) 1. *Brixh.* 6.—Sibir. baical.
cognata, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 198 = cuspidata.
colchagensis, *Phil. in Anal. Univ. Chil.* (1879) 673. —Chili.
compacta, Schott, Nyman, & Kotschy, Analect. Bot. 50 = D. Aizoon.
confertifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 11. 301.—Ecuador.
confusa, Ehrh. Beitr. vii. 155 = incana.
confusa, Hartm. Skand. ed. III. 158 = hirta.
contorta, Ehrh. Beitr. vii. 155 = incana.
corrugata, *S. Wats. Bot. Calif.* ii. 480.—Calif.
corsica, Jord. Diagn. i. 205 = Loiseleurii.
corymbosa, R. Br. in Ross, Voy. App. p. cxlii = alpina.
corymbosa, Fries, ex Lange, Cónsp. Fl. Groenl. i. 48 = hirta.
crassifolia, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.–June 1839) 182.—Am. bor.
crenates, *Czern. ex Ledeb. in Bull. Soc. Nat. Mosc.* xxxviii. (1865) i. 177.—Ucrania.
cretica, *Boiss. & Heldr. Diagn.* Ser. I. viii. 27.—Ins. Creta.
cryptantha, *Hook. f. Fl. Antarct.* 284.—Ecuador.
cuneifolia, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 108. —Am. bor.
curtisiliqua, Zett. Pl. Vasc. Pyr. p. xlviii = stellata.
cuspidata, Bertol. Fl. Ital. vi. 467 = aizoides.
cuspidata, *Bieb. Fl. Taur. Cauc.* iii. 434.—As. Min.
cuspidata, DC. Syst. ii. 334 = hispanica.
dahurica, Fisch. ex Turcz. in Enll. Soc. Nat. Mosc. (1838) 87 = hirta.
dasycarpa, *Grenth. Syst. Vers. Erf.* 192.—Europ.
dasycarpa, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 79.—Sibir. altaic.
daurica, *DC. Syst.* ii. 350.—Dauria.
Davilae, Phil. in Anal. Univ. Chil. (1872) 672 = Gilliesii.
davosiana ×, Brügg. in Jahresb. Naturf. Ges. Graub. II. xxiv.–xxiv. (1880) 74.—Europ.
Decleana, *Boiss. & Reut. ex Boiss. Voy. Bot. Espagne*, ii. 718.—Hispan.
dendrifolia, Nutt. in Torr. & Gray, Fl. N. Am. i. 104 = alpina.
dentata, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 192 = ramosissima.
depressa, *Hook. f. Fl. Antarct.* 284.—Ecuador.
dispansioides, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 285.—Bolivia.
dicranoides, Boiss. & Huet, Diagn. Ser. II. v. 80 = rigida.
discoides, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 286.—Bolivia.
diversifolia, Boiss. & Huet, Diagn. Ser. II. v. 80 = olympica.
Dornerі, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 25 = fladnizensis.
Douglasii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 358.—Am. bor. occ.
dorycnifolia, Fries, Summa Veg. Scand. 32 = hirta.
dubia, Suter, Fl. Helv. ii. 46 = tomentosa.
elata, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 150.—Reg. Himal.
elegans, Boiss. in Ann. Sc. Nat. Sér. III. xvii. (1842) 166 = alpina.
ellipsoides, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 153.—Reg. Himal.
elongata, Host, Fl. Austr. ii. 337 = aizoides.
ericaefolia, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 326 = olympica.
eriopoda, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 280.—Dahuria.
eurycarpa, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 590.—Calif.
extensa, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 286. —Ecuador.

DRABA :—
falklandica, *Hook. f. Fl. Antarct.* 288.—Ins. Falkland.
fiscensis, Scheele, in Linnaea, xxi. (1848) 388 = carolininana !
fladnizensis, *Wulf. in Jacq. Misc.* i. 147. t. 17. f. 1.—Reg. bor. et arct.
frigida, Sauter, in Flora, viii. (1825) 72 = tomentosa.
frigida, Schur, Enum. Pl. Transs. 64 = stellata.
fuiniculosa, *Hook. f. Fl. Antarct.* 288. t. 89.—Ins. Falkland.
gelida, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 250 = alpina.
Gigas, Stev. ex Boiss. Fl. Orient. i. 278 = Arabis carduchorum.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 137.—Chili.
glabella, Pursh, Fl. Am. Sept. ii. 434 = arabisans.
glabella, Richards. in Frankl. Narr. 1st Journ. 744 = incana.
glacialis, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 106 = alpina.
glacialis, Hoppe, in Flora, vi. (1823) 424 = aizoides.
globifera, Ledeb. Fl. Ross. i. 146 = olympica.
glomerata, Royle, Illustr. Bot. Himal. 71 = lasiophylla.
Gmelinі, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 107 = repens.
gracilis, R. Grah. in Edinb. N. Phil. Journ. (Apr.–June 1839) 179 = nemorosa.
gracilis, Ledeb. Fl. Ross. i. 152 = hirta.
gracillima, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 153.—Reg. Himal.
grandiflora, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 137, nomen ; *Hook. f. Fl. Antarct.* 286.—Ecuador.
grandiflora, C. A. Mey. in Ledeb. Fl. Alt. iii. 74 = Parrya microcarpa.
grandis, Langsd. ex DC. Syst. ii. 355 = hyperborea.
greggioides, *Griseb. in Goett. Abh.* xxiv. (1879) 16.—Reg. Argent.
Hallii, *Hook. f. Fl. Antarct.* 287.—Ecuador.
Haynaldі, Stur, in Oestr. Bot. Zeitschr. xl. (1861) 186. t. 3 = D. Aizoon.
hederaefolia, *Cass. in Bull. Soc. Bot. Fr.* xxii. (1875) 54, nomen.—Afr. bor.
helvetica, Schleich. Cat. Pl. Helv. ed. III. 18; DC. Syst. ii. 345 = fladnizensis.
Hennsena, Schlecht. in Linnaea, x. (1836) 100 = arabisans.
hesperidiflora, Lam. Encyc. ii. 328 = Aubrietia deltoides.
heterocoma, Fenzl, Pugill. Pl. Nov. Syr. 18 = olympica.
himalayensis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 127 = incana.
hirsuta, Pers. Syn. ii. 190 = repens.
hirsuta, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 64.—Ins. Kurile.
hirta, Gaudin, Fl. Helv. iv. 252 = sibiquosa.
hirta, Gunn. Fl. Norv. ii. 61 = laxa.
hirta, *Linn. Syst.* ed. X. 1197.—Reg. bor. et arct.
hirta, Ocd. in Fl. Dan. t. 143 ; Jacq. Fl. Austr. v. 15. t. 482 = stellata.
hirta, Sibth. & Sm. Fl. Graec. Prod. ii. 5 = Arabis brabaeformis.
hirta, Sm. Fl. Brit. 677 = rupestris.
hirta, Towns. It. Hung. 490, 858 = tomentosa.
hispanica, *Boiss. Elench.* 18.—Hispan. ; Marocc.
hispida, *Willd. Sp. Pl.* iii. 426.—Kossia ; Reg. Cauc.
hispidula, Michx. Fl. Bor. Am. ii. 28 = caroliniana.
Hookeri, *Walp. Ann.* i. 85.—Ecuador.
Hoppeana, Reichb. in Moesel. Handb. ed. II. ii. 1183 = aizoides.
Hoppeana, Rudolphi, ex Reichb. Fl. Germ. Excurs. 666 = sibiquosa.
Hoppii, Trachsel, in Flora, xiv. (1831) 741 = sibiquosa.
Howellii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 354. —Calif.
Huetii, *Boiss. Diagn.* Ser. II. v. 31.—As. Min. ; Persia.
Humboldtii, Desv. Journ. Bot. iii. (1814) 171 = Eutrema Humboldtii.
hyperborea, *Desv. l. c.* 172.—Am. bor. et arct.
Hyrnix, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 148.—Afghan.
imbricata, *C. A. Mey. Verz. Pfl. Cauc.* 182.—Reg. Caucas.
imbricatifolia, *Barn. in C. Gay, Fl. Chil.* i. 158.—Chili.
imervetia, *Rupr. Fl. Cauc.* 201, in textu.—Reg. Caucas.
incana, *Linn. Sp. Pl.* 648.—Reg. bor. et arct. ; Reg Magellan.

DRABA:—

incompta, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 268 = alpina.

inflata, H. C. Wats. ex Syme, Engl. Bot. ed. III. i. 191. t. 134 = Erophila vulgaris.

intermedia, Andrz. ex DC. Syst. ii. 851 = nemorosa.

involuta, Ag. ex Willk. & Lange, Prod. Fl. Hisp. iii. 839 = hispanica.

Jamesonii, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 300.—Ecuador.

japonica, *Maxim. in Bull. Acad. Péterb.* xxii. (1877) 228.—Japon.

Johannis, Host, Fl. Austr. ii. 340 = siliquosa.

Joncolyx, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 299.—Ecuador.

jordilensis, *H. B. & K. Nov. Gen. et Sp.* v. 78.—Mexic.

juvenalis, *Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.)*. i. (1858) 415.—Hab. ?

Khilsiana, *Prasl, Bot. Bemerk.* 9.—Hab. ?

Kotschyi, *Stur, in Oestr. Bot. Zeitschr.* ix. (1859) 38.—Europ.

Krocheri, Schur, Enum. Pl. Transs. 67 = Erophila praecox.

lactea, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 104 = fladnizensis.

lactea, *Neilr. in Oestr. Bot. Zeitschr.* ix. (1859) 92.—Europ.

laevigata, Cham. & Schlecht. in Linnaea, i. (1826) 25 = Extrema Edwardsii.

laevigata, Hoppe, ex Koch, in Flora, vi. (1823) 441 = fladnizensis.

laevipes, DC. Syst. ii. 346 = tomentosa.

lanceolata, *Royle, Illustr. Bot Himal.* 72.—Ind. or. *lapponica*, Schur, Enum. Pl. Transs. 64 = Kotschyi.

lapponica, Willd. ex DC. Syst. ii. 344 = fladnizensis.

latisiliqua, Adams, in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 248 = alpina.

laxicarpa, Rochel, ex DC. Syst. ii. 334 = D. Aizoon.

lasiophylla, *Royle, Illustr. Bot. Himal.* 71.—Reg. Himal. ; Turkestan.

lasiopoda, *Turcz. ex Bess. in Flora*, xvii. (1834) i. *Beibl.* 6.—Sibir. baical.

laxa, *Lindblom, in Lunnaea*, xiii. (1839) 326.—Scandinav.

leiocarpa, Lindbl. in Vet. Akad. Handl. Stockh. 1839 (1841) 66 = hirta.

leiopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = laevipes.

Lemmoni, *S. Wats. Bot. Calif.* ii. 480.—Calif.

leptopetala, T. Fries, in Öfver. Vet. Akad. Stockh. xxvi. (1869) 180 = hirta.

Levieri, *Janka, in Termész. Füzetek*, vii. (1883) 107.—Europ. austr.

Leybold, *Phil. in Linnaea*, xxxiii. (1864–65) 10.—Chili ; Ecuador.

Liljebladii, Wallm. in Liljeb. Svenska Fl. ed. III. 350 = stellata.

linearis, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 167.—Persia ; Tibet occ.

Loiseleurii, *Boiss. Diagn.* Ser. II. i. 34.—Corsica.

Longii, Schwein. ex Torr. & Gray, Fl. N. Am. i. 106 = arabisana.

longirostra, *Schott, Nyman, & Kotschy, Analect. Bot.* 43.—Dalmatia.

lutea, Gilib. Fl. Lituan. ii. 46 = nemorosa.

lutescens, *Cosc. Notes Crit.* 147.—Hispan.

Macleani, *Hook. f. Fl. Antarct.* 238.—Peruv.

macrantha, Vest, ex Rchb. Fl. Germ. Crit. iii. 11 = stellata.

macrocarpa, Adams, in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 248 = alpina.

macrocarpa, Boiss. & Heldr. Diagn. Ser. I. viii. 28 = Erophila macrocarpa.

macroloba, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 301.—Sibir. occ.

macropetala, *Hort. Wirceb. ex Schenk, Ind. Sem. Hort. Wirceb.* (1861).—Hab. ?

macrophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 255.—Sibir.

magellanica, *Lam. Encyc.* ii. 328.—Reg. Magell.

Mandoniana, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 287.—Bolivia.

maritima, Lam. Fl. Fr. ii. 461 = Alyssum maritimum.

Martinsiana, *Y. Gay, in Bibl. Univ. Genève.* N. S. xxviii. (1840) 146.—Spitsbergen.

Mawii, *Hook. f. Bot. Mag.* t. 6196.—Hispan.

micrantha, Nutt. in Torr. & Gray, Fl. N. Am. i. 109 = caroliniana.

DRABA:—

micropetala, Hook. in Parry, 2nd Voy. App. 385 = alpina.

minima, Stend. Nom. ed. II. i. 528 = Erophila minima.

muricaulis, Willd. ex DC. Syst. ii. 357, (err. typ. muniticaulis) = Erophila praecox.

mogollonica, *Greene, in Coult. Bot. Gaz.* vi. (1880) 157.—N. Mexic.

mollis, Scop. Fl. Carn. ed. II. ii. 7 = Arabis vochinensis.

mollissima, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 268.—Reg. Caucas.

mongolica, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 256.—Mongolia.

montana, Berg. ex DC. Syst. ii. 333 = aizoides.

montana, *S. Wats. Bibliog. Ind. N. Am. Bot.* i. 60.—Am. bor.

Moritziana, *Bruegg. in Jahresb. Naturf. Ges. Graub.* ii. xxv. (1882) 49.—Europ.

muralis, Hook. Fl. Bor. Am. i. 56 = nemorosa ?

muralis, *Linn. Sp. Pl.* 643.—Europ. ; As. temp.

muralis, Thunb. Fl. Jap. 259 = nemorosa.

muricella, Wahlenb. Fl. Lapp. 174 = stellata.

muscosa, *Ruiz & Pav. ex Steud. Nom.* ed. II. i. 528.—Peruv.

mutabilis, Desv. Journ. Bot. iii. (1814) 172 = Alyssum mutabile.

myosotidioides, *Hemsl. Diagn. Pl. Nov.* 18.—Mexic.

Nasturtiolum, Scop. Fl. Carn. ed. II. ii. 8 = Smelowskia alpina.

natolica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 168.—As. Min.

nemoralis, Ehrh. Beitr. vii. 153 = nemorosa.

nemorosa, All. Fl. Pedem. i. 244 = muralis.

nemorosa, *Linn. Sp. Pl.* 643.—Reg. bor.

nepalensis, *Wall. Cat.* n. 4785.—Reg. Himal.

nivalis, DC. Fl. Fr. iv. 699 = fladnizensis.

nivalis, Lapeyr. Hist. Abr. Pl. Pyr. 364 = tomentosa.

nivalis, Lilj. in Vet. Acad. Handl. Stockh. (1793) 208 = stellata.

nivea, Sauter, in Flora, xxxv. (1852) 622 = tomentosa.

norvegica, Gunn. Fl. Norv. ii. 106 = hirta.

nubigena, Desv. Journ. Bot. iii. (1814) 171 (= *Eudema nubigena*).—Ecuador.

nummularia, Ehrenb. ex Webb & Berth. Phyt. Canar. i. 93 = Alyssum libycum.

oblongata, R. Br. Voy. Ross. App. p. cxliii, nomen ; DC. Syst. ii. 342 = hirta.

oblongata, Nym. Consp. 52 = aizoides.

obovata, *Benth. Pl. Hartw.* 159.—Ecuador.

occidentalis, Bunge, Verz. Suppl. Fl. Alt. 69 = alpina.

Oederi, Vill.—Cf. Bull. Soc. Bot. Fr. xi. (1864) 152 = siliquosa.

Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 8.—Turkestan.

olympica, Gren. & Godr. Fl. Fr. i. 123 = Loiseleurii.

olympica, *Sibth. ex DC. Syst.* ii. 336.—Graecia ; Orlens.

oligosperma, Hook. Fl. Bor. Am. i. 51 = alpina.

Oreades, *Schrenk, Enum. Pl. Nov.* ii. 56.—Turkestan.

ostetica, Rupr. Fl. Cauc. 112 = mollissima.

oxycarpa, *Boiss. & Heldr. Diagn.* Ser. I. viii. 28.—Syria.

oxycarpa, Sommerf. ex Nym. Consp. 53 = alpina.

Pacheri, Stur, in Oestr. Bot. Wochenbl. v. (1855) 49 = hirta.

pachyrhyncha, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 62.—N. Granat.

parnassica, *Boiss. & Heldr. ex Lindén, ex Triana & Planch. l. c.* 61.—N. Granat.

parnassica, *Boiss. Diagn.* Ser. II. i. 34.—Graecia.

parryoides, Cham. in Linnaea, vi. (1831) 533 = Parrya Ermani.

patagonica, *Phil. l. c.* xxviii. (1856) 669.—Patagon.

pauciflora, R. Br. in Parry, 1st Voy. Suppl. App. p. cclxvi = alpina.

pectica, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 164 = alpina.

petraea, Baumg. Enum. Stirp. Transs. ii. 285 = Hutchinsia petraea.

Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 54.—Peruv.

pilosa, Adams, ex DC. Syst. ii. 336 = alpina.

platycarpa, *Torr. & Gray, Fl. N. Am.* i. 108.—Am. bor. occ.

polygonifolia, Mill. Gard. Dict. ed. VIII. n. 5 = nemorosa.

polytricha, Ledeb. Fl. Ross. i. 146 = alpina.

DRABA:—

pontica, Desf. in Ann. Mus. Par. xi. (1808) 381 = nemorosa.

poposatapetlensis, *Hemsl. Diagn. Pl. Nov.* 18.—Mexic.

porrecta, Trin. ex DC. Syst. ii. 339 = repens.

praecox, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 269 = Erophila praecox.

primuloides, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 250 = alpina.

pulchella, Willd. ex DC. Syst. ii. 354 = alpina.

pulvinata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 298.—Venezuela.

pumila, Miellichh. ex Sauter, in Flora, xxxii. (1849) 666 = tomentosa.

pumila, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = sec.

Pumilio, K. Br. ex DC. Syst. ii. 358 = Capsella procumbens *vel* australis ?

pycnophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 300.—Ecuador.

pygmaea, Turcz. l. c. (1888) 87 = fladnizensis.

pyrenaica, *Linn. Sp. Pl.* 642.—Mont. Pyren.

pyrenaica, Oed. in Fl. Dan. t. 148 = laxa.

radicosa, Royle, Illustr. Bot. Himal. 71 = alpina.

radicata, *Hook. f. Fl. Antarct.* 236.—Ecuador.

ragusana, Vest, ex Reichb. Fl. Crit. iii. 29 = Alyssum moutabile.

ramosa, Gater. Descr. Fl. Montaub. 114 = muralis.

ramosissima, *Desv. Journ. Bot.* iii. (1814) 186.—Am. bor.

repens, *Birb. Fl. Taur. Cauc.* ii. 93.—Reg. Caucas. Sibir.

Reuteri, Boiss & Huet, Diagn. Ser. II. vi. 18 = alpina rhactica, *Brueggr. in Jahresb. Naturf. Ges. Graub.* II xxv. (1882) 49.—Europ.

Rhizobotrya, Steud. Nom. ed. II. i. 528 = Cochlearia brevicaulis.

rigida, Loisel. Fl. Gall. ed. II. ii. 51 = Loiseleurii.

rigida, *Willd. Sp. Pl.* iii. 425.—As. Min.

rivularis, Bernh. ex Reichb. Fl. Germ Excurs 667 = stellata.

rivularis, *Hornem. ex Steud. Nom.* ed. II. i. 528.—Sibir. baical.

Roegneri, Stev. ex Rupr. Fl. Cauc. 291, in textu = inaetica.

Roerneriana, Scheele, in Linnaea, xxi. (1848) 583 = platycarpa *vel* cuneifolia ?

rosea, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 260 = Braya rosea.

rosularis, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 165 = calycosa.

rosulata, Phil. in Linnaea, xxviii. (1856) 669 = Gilliesii.

rubra, Crantz, Crucif. 95. t. 1. f. 5 = pyrenaica.

rudbraiis, Baumg. Enum. Stirp. Transs. ii. 282 = Lepidium Draba.

rupestris, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 91.—Reg. bor. et arct.

rupestris, Willd. ex DC. Syst. ii. 339 = hispida.

sachsiensis, Trautv. in Act. Hort. Petrop. viii. (1883) 92 = uncana.

saxa, Spreng. Syst. ii. 876 = Cochlearia saxa.

samojedorum, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* viii. (1850) 291.—Mont. Ural.

Sauteri, *Hoppe, in Flora*, vi. (1823) 425.—Tyrolia.

saxatilis, Mart. & Koch, in Flora, vi. (1823) 435 = stellata.

saxicola, C. Koch, in Linnaea, xix. (1847) 53 = Arabis drabaeformis.

saxigena, Jord. Diagn. i. 203 = aizoides.

scabra, *C. A. Mey. Vers. Pfl. Cauc.* 182.—Reg. Caucas. scandinavica, *Lindblom, in Linnaea*, xiii. (1839) 322.—Scandinav.

Schiverekia, *Janka, in Termész. Füzetek*, vii. (1883) 109.—Europ. austr.

sclerophylla, Gaudin, Fl. Helv. iv. 255 = fladnizensis.

scopulorum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 286.—Bolivia.

setosa, Royle, Illustr. Bot. Himal. 71 = alpina.

setulosa, C. Koch, in Linnaea, xix. (1847) 53 = Arabis drabaeformis.

setulosa ×, Lereache, in Compt.-Rend. Soc. Hallér. (1852–53) 7.—Europ.

siliquosa, *Bieb, Fl. Taur. Cauc.* ii. 94 ; iii. 432.—As. Min. ; Persia.

siliquosa, Hook. f. Fl. Antarct. 239 = Hookeri.

soratensis, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 287.—Bolivia.

spathulata, Spreng. Syst. ii. 876 = hyperborea.

DRABA:—
spatulata, Lang. ex Reichb. Fl. Germ. Excurs. 665 = Erophila praecox.
spinosa, Lam. Fl. Fr. ii. 461 = Alyssum spinosum.
Spitzelii, *Hoppe, in Sturm, Deutschl. Fl.* Heft lx.—Europ.
Sprengeana, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 286.—Ecuador.
stellaroides, Willd. ex DC. Syst. ii. 339 = repens.
stellata, Cham. in Linnæa, i. (1826) 22 = tomentosa.
stellata, *Jacq. Enum. Vindob.* 113; 256.—Reg. bor. et arct.
stellata, Oed. in Fl. Dan. t. 142 = silliquosa.
stellata, Sieber, ex Steud. Nom. ed. II. i. 528 = Cochlearia brevicaulis.
stellata, With. Arr. Brit. Pl ed. III. 665 = rupestris.
stenocarpa, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 153 = linearis.
stenoloba, Ledeb. Fl. Ross. i. 154 = hirta.
stenopetala, Trauty. in Act. Hort. Petrop. vi. (1879) 11. —As bor.
stenophylla, *Leyb. in Anal. Univ. Chil.* (1859) 678.—Chili.
stolonifera, Barn. in C. Gay, Fl. Chil. i. 155 = Gilliesii.
streptocarpa, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 343.—Am. bor.
stylaris, J. Gay, ex DC. Syst. ii. 348 = incana.
stylosa, Dulac, Fl. Hautes-Pyr. 192 = aizoides.
stylosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 300.—Ecuador.
subamplexicaulis, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 77. —Sibir. altaic.
Subularia, Lam. Tabl. Encyc. iii. 109 = Subularia aquatica.
suffruticosa, *Barn. in C. Gay, Fl. Chil.* i. 157.—Chili.
supranivalis, *Rupr. Fl. Cauc.* 116.—Reg. Caucas.
taurica, *Willd. ex DC. Syst.* ii. 335 = cuspidata.
tenuis, *Barn. in C. Gay, Fl. Chil.* i. 156.—Chili.
Thomasii, Koch, Syn. Fl. Germ. ed. II. 438 = incana.
tibetica, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 152.—Reg. Himal.
toluccana, DC. Syst. ii. 341 = toluccensis.
toluccensis, *H. B. & K. Nov. Gen. et Sp.* v. 78.—Mexic.
tomentosa, *[Clairv.] Man. Herb.* 217; *Wahlb. Fl. Helv.* 123.—Europ.
tomentosa, Schur, Enum. Pl. Transs. 64 = Kotschyi.
Traschelii, *Dalla Torre, Atl. Alpenfl.* 65.—Europ.
transilvanica, Schur, Enum. Pl. Transs. 64 = Kotschyi.
Traunsteineri, Hoppe, in Sturm, Deutschl. Fl. Heft lxv = silliqnosa.
trichella, Fries, Nov. Fl. Suec. Mant. ii. 40 = rupestris
tridentata, DC. Syst. ii. 339 = hispida.
tschukschorum, *Trautv. in Act. Hort. Petrop.* vi. (1879) 13.—Sibir. bor.
turgida, Huet, ex Nym. Consp. 52 = longirostra.
turkestanica, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 7.—Turkestan.
umbellata, Mühl. Cat. 62 = caroliniana.
unalaschkiana, DC. Syst. ii. 350 = incana.
unilateralis, *G. Yones, in Bull. Torrey Bot. Club*, ix. (1882) 134.—Mexic.
uralensis, Willd. ex DC. Syst. ii. 300 = Alyssum podolicum.
velutina, *Boiss. & Huet, Diagn.* Ser. II. vi. 16.—Armenium.
ventosa, *A. Gray, in Am. Natural.* viii. (1874) 212.—Am. bor.
verna, Linn. Sp. Pl. 649 = Erophila vulgaris.
Vesicaria, *Desv. Journ. Bot.* iii. (1814) 186.—Syria.
violacea, *DC. Syst.* ii. 354.—Ecuador.
volcanica, *Kunth. Pl. Hortbn.* 93.—Guatemala.
Wahlenbergii, Hartm. Skand. Fl. ed. I. 249 = fladnizensis.
Wahlenbergii, Schur, Enum. Pl. Transs. 64 = Kotschyi.
Wormskioldii, *Fisch. ex Spreng. Syst.* ii. 874.—Sibiria.
Zahlbruckneri, *Host, Fl. Austr.* ii. 338.—Austria.
Zapateri, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 889.—Hispan.

DRABELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 388 = Draba, Dill. (Crucif.).
muralis, Fourr. l. c. = D. muralis.

DRABOPSIS, C. Koch, in Linnæa, xv. (1841) 253 = Sisymbrium, Tourn. (Crucif.).
verna, C. Koch, l. c. = S. saudum.

DRACAENA, Vand. Diss. (1762); Linn. Mant. i. 68 (1767). *LILIACEÆ*, Benth & Hook. f. iii. 779.
DRACO, Crantz, Duab. Dracon. Arb. 13 (1768).
DRAKAINA, Rafin. Fl. Tellur. iv. 17 (1836)
NEMAMPSIS, Rafin. l. c. (1836).
ORDYLA, Crantz, Duab. Dracon. Arb. 30 (1768).
PLEOMELE, Salisb. Prod. 245 (1796).
TERMINALIS, Medic. Theodors, 88 (1786).
STORERIA, Crantz, Duab. Dracon. Arb. 25 (1768).
acanlis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 252.—Afr. trop.
acuminata, *Thunb. Diss. Dracaena*, 5.—Ins Mascar.
Aitelii, *Baker, in Journ. Bot.* xii. (1874) 167.—Afr. trop.
alho-rosea, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
angustifolia, *Roxb. Hort. Beng.* 24; *Fl. Ind.* ii. 155.—Ind. or.; Malaya.
arborea, *Hort. Angl. ex Link, Enum. Hort. Berol.* i. 341.—Afr. trop.
arborea, C. Koch, Wochenschr. x. (1867) 286 = Kochiana.
atropurpurea, Roxb. Hort. Beng. 24; Fl. Ind. ii. 160 elliptica.
Aubryana, Brongn. ex E. Morr. Belg. Hortic. x. (1880) 348 = thalioides.
aurantiaca, *Wall. Cat.* n. 5144.—Malaya.
aures, *H. Mann, in Proc. Am. Acad*, vii. (1867) 207. —Ins. Sandvic.
Aurora, *Linden & André, in Illustr. Hortic.* xxv. (1878) 26. t. 304.—Ins. Pacif.
australis, Forst. f. Prod. 24 = Cordyline australis.
australis, Hook. Bot. Mag. t. 2885 = Cordyline Baueri.
Bellula, Linden & André, in Illustr. Hortic. xxi. (1874) 41. t. 263 = Cordyline terminalis.
Betschleriana, C. Koch, in Berl. Allg. Gartens. (1858) 284 = concinna.
bicolor, *Hook. Bot. Mag.* t. 5248.—Afr. trop.
Boerhavii, Tenore, in Mem. Acc. Sc. Borb. Nap. iii. 11. (1832) 44. t. 3 = D. Draco.
borealis, Soland. in] Ait. Hort. Kew. ed. I. i. 454 = Clintonia borealis.
Boscii, Hort. ex Steud. Nom. ed. II. i. 528 = Agave geminiflora.
brachyphylla, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 249.—Ins. Andaman.
brasiliensis, Hort ex Schult. f. Syst. vii. 1676 = Cordyline terminalis.
camerooniana, *Baker, in Journ. Bot.* xii. (1874) 166. —Afr. trop.
Candelaria, Hort. ex Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 182 = reflexa.
Cantleyi, *Baker, in Journ. Bot.* xix. (1881) 326.—Singapore.
Cascadæ, *Linden & André, in Illustr. Hortic.* xxi. (1874) 137. t. 181.—N. Hebrid.
cernua, Jacq. Hort. Schoenbr. i. 50. t. 96 = reflexa.
cincta, *Baker, in Journ. Linn. Soc.* xiv. (1875) 580.—Hab.?
cinnabari, *Balf. f. in Trans. Roy. Soc. Edinb.* xxx. (1882) 238.—Ins. Socotra.
cochenchalis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 582 = fruticosa.
compacta, Hort. ex Baker, l. c. 540 = præc.
concinna, *Kunth, Enum. Pl.* v. 8.—Madag.
congesta, Sweet, Hort. Brit. ed. I. 424 = Cordyline congesta.
Cooperi, Hort. ex Regel, Gartenfl. (1871) 148; et Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
cylindrica, *Hook. f. Bot. Mag.* t. 5846.—Afr. trop.
densiflora, *Baker, in Journ. Bot.* xii. (1874) 166.—Afr. trop.
dentata, Pers. Syn. i. 371, in obs. = Lomatophyllum borbonicum.
directicrura, Willd. ex Regel, in Act. Hort. Petrop. i. (1871) 140 = reflexa.
Draco, *Linn. Sp.* ed. XII. 246.—Ins. Canar.
Ehrenbergii, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 538 = Yucca Ehrenbergii.
elliptica, Desf. Cat. Hort. Par. ed. III. 388 = Fontanesiana.
elliptica, *Thunb. Diss. Dracaena*, 6.—Aa. trop.
elliptica, Thunb. l. c., partim = spicata.

DRACAENA:—
ensata, Thunb. Diss. Dracaena, 4 = Dianella ensifolia.
ensifolia, Haw. Syn. Pl. Succ. 67 = Yucca flexilis.
ensifolia, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 222 = Yucca guatemalensis.
ensifolia, Linn. Mant. i. 68 = Dianella ensifolia.
ensifolia, Regel, Gartenfl. (1864) 321. t. 451 = fruticosa.
ensifolia, Wall. Cat. n. 5148 = angustifolia.
ensiformis, Wall. ex Voigt, Hort. Suburb. Calc. 675 = angustifolia.
erecta, Linn. f. Suppl. 204 = Asparagus erectus.
esculenta, Hort. ex Regel, Gartenfl. (1871) 148; et Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
excelsa, Hort. ex Baker, l. c. 532 = fruticosa.
excelsa, Tenore, Ind. Sem. Hort. Neap. (1839) 11. n. 5 = arborea.
ferrea, Linn. Syst ed. XII. 346 = Cordyline terminalis.
filamentosa, Scanag. ex Schult. f. Syst. vii. 729 = Agave geminiflora.
filiformis, Thunb. Diss. Dracaena, 4 = Xerotes filiformis.
Finlaysoni, *Baker, in Journ. Bot. xi.* (1873) 263.—Malaya.
Fintelmanni, Hort. ex Regel, Gartenfl. (1871) 149; et Baker, in Journ. Linn. Soc. xviii. (1880) 222 = Yucca guatemalensis.
flabelliformis, Bory, Voy. i. 270 = Cordyline macrophylla.
Flemingii, Hort ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
flexuosa, Regel, Gartenfl. (1859) 329 = reflexa.
floribunda, *Baker, in Hook. f. Bot. Mag.* t. 6447.—Hab.?
foetida, Salisb. Prod. 254 = Lomatophyllum borboricum.
Fontanesiana, *Schult. f. Syst.* vii. 1676.—Ins. Mascar.
formosa, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
fragrans, *Ker-Gawl. in Bot. Mag.* t. 1081.—Afr. trop.
Fraseri, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
fruticosa, *C. Koch, Wochenschr.* x. (1867) 286.—Hab.?
fruticosa, Regel, in Act. Hort. Petrop. i. (1871) 137 = angustifolia.
Gibsoni, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
gigantea, *Baker, in Journ. Bot.* xii. (1874) 166.—Afr. trop.
gloriosa, Linden, ex E. Morr. in Belg. Hortic. xxiii. (1873) 111 = Cordyline terminalis.
gracilis, Hort. Bull, ex Baker, in Journ. Linn. Soc. xiv. (1875) 580 = cincta.
gracilis, Salisb. Prod. 258 = marginata.
gracilis, *Wall. Cat.* n. 5150.—Malaya.
graminifolia, Linn. Syst. ed. XII. 246 = Liriope spicata.
graminifolia, Wall. Cat. n. 5149 = Finlaysoni.
Griffithii, Regel, Gartenfl. (1871) 146 = Helferiana.
Guilfoylei, Veitch, Cat. ex Regel, Gartenfl. (1871) 149 = Cordyline terminalis.
Helferiana, *Wall. ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 248.—Burma.
hamechrysa, Thunb. Diss. Dracaena, 5 (= Asteliae sp.) —' Ins. Mauriti.'
Heynaana, Wall. Cat. n. 5151 = spicata.
hirsuta, Thunb. Diss. Dracaena, 6 = Palisota thyrsiflora.
Hookeriana, *C. Koch, Wochenschr.* iv. (1861) 394.—Afr. austr.
humilis, *Baker, in Journ. Bot.* xii. (1874) 166.—Afr. trop.
illustris, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
imperialis, Hort. ex Baker, l. c. = Cordyline terminalis.
indivisa, Forst. f. Prod. 24 = Cordyline indivisa.
interrupta, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
interrupta, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 252.—Afr. trop.
interrupta, *Hort. Angl. ex Link, Enum. Hort. Berol.* i. 341 (= præc.?).—Afr. trop.
Jeckiana, *Wall. Cat.* n. 5145.—Malaya.
jamaicæ, Kunth, Enum. Pl. v. 12 = elliptica.
Kirkii, *Baker, in Journ. Bot.* xii. (1874) 164.—Ins. Johanna.
Kuehniana, C. Koch, Wochenschr. iv. (1861) 394 = arborea.
Kochiana, *Regel, Gartenfl.* (1871) 137.—Hab.?

DRACAENA:—

lanceæ, Thunb. Dies. Dracæna, 5 = reflexa.
latifolia, Regel, Gartenfl. (1871) 141 = Hookeriana.
Lennæana, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1876) 588 = Yucca Lennæana.
Lennæana, Regel, Gartenfl. (1871) 149 = Yucca conspicua.
Lennei, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 222 = Yucca guatemalensis.
leonensis, *Ladd. ex Loud. Hort. Brit.* 180, nomen.—Afr. trop.
lineariifolia, Ayres, ex Baker, in Journ. Linn. Soc. xiv. (1875) 581 = reflexa.
linearis, Hort. ex Baker, l. c. 548 = Cordyline terminalis.
lutescens, *A. Versch.—Cf. Belg. Hortic.* xxiii. (1873) 111.—Hab.?
maculata, Roxb. Hort. Beng. 24; Fl. Ind. ii. 157 = elliptica.
maculata, Wall. Cat. n. 5148 A = Porteri.
magnifica, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1876) 540 = Cordyline terminalis.
Mannii, *Baker, in Journ. Bot.* xii. (1874) 164.—Afr. trop.
marginata, Ait. Hort. Kew. ed. I. i. 454 = Lomatophyllum borbonicum.
marginata, Horne, ex Baker, in Journ. Linn. Soc. xiv. (1876) 580 = concinna.
marginata, *Lam. Encyc.* ii. 324.—Madag.
mauritiana, Boj. Hort. Maurit. 348 = Cordyline floribunda.
mauritiana, Lam. Encyc. ii. 325 = Dianella mauritiana.
medeoloides, Linn. f. Suppl. 203 = Asparagus medeoloides.
monostachya, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 253.—Afr. trop.
Moreana, Hort. ex Baker, l. c. 540 = præc.
nemorosa, Stend. Nom. ed. II. i. 529 = Dianella odorata.
nigra, Regel, Gartenfl. (1871) 149 = Fontaneziana.
nilens, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 252.—Afr. trop.
nobilis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1876) 540 = Cordyline terminalis.
obliqua, Thunb. Diss. Dracæna, 6 = Xerotes flexifolia.
obtecta, R. Grah. in Edinb. N. Phil. Journ. (Apr.-June 1837) 175 = Cordyline Banksii.
Ombet, Kotschy & Peyr. Fl. Timn. 47 = D. Draco.
ovata, *Kew-Gawl. in Bot. Mag.* t. 1180.—Afr. trop.
pachyphylla, *Kuru, in Journ. As. Soc. Beng.* xlii. (1872) 11. 249.—Ins. Andamans.
parviflora, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 252.—Afr. trop.
parviflora, Willd. ex Schult. f. Syst. vii. 848 = Nolina parviflora.
Perrottetii, *Baker, in Journ. Bot.* xii. (1874) 165.—Afr. trop.
phrynioides, *Hook. Bot. Mag.* t. 5352.—Afr. trop.
plicata, *Wall. ex Voigt, Hort. Suburb. Calc.* 675, nomen.—Hab.?
Portuana, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
Porteri, *Baker, in Journ. Bot.* xi. (1873) 262.—Penang.
pulchella, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1876) 540 = Cordyline terminalis.
pulcherrima, Hort. ex Baker, l. c. = Cordyline terminalis.
pumila, Hort. ex Link, Enum. Hort. Berol. i. 341 = umbraculifera.
punctata, Hort. Van Houtte, ex Regel, Gartenfl. (1871) 139 = stenophylla.
quilensis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1876) 532 = fruticosa.
Reali, *Linden & André, in Illustr. Hortic.* xx. (1873) 166.—Archip. Ind.
reflexa, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 368 = timorensis.
reflexa, *Lam. Encyc.* ii. 324.—Ins. Mascar.; Madag.
reflexa, F. Muell. Fragm. vi. 130 = angustifolia.
regalis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
reginifera, Sallab. Prod. 258 = D. Draco.
Robinsoniana, *Hort. ex André, in Illustr. Hortic.* xxvi. (1879) 56. t. 342.—Ins. Solomon.
robusta, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1876) 540 = Cordyline terminalis.
Rothiana, *Carr. in Rev. Hortic.* (1877) 68.—Hab.?
rubra, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 12 = Cordyline terminalis?
Rumphii, Regel, Gartenfl. (1873) 140 = Hookeriana.
salicifolia, Regel, l. c. 141 = reflexa.

DRACAENA:—

Saposchnikowi, *Regel, Gartenfl.* (1871) 293. t. 705.—Hab.?
sarmentosa, Boj. ex Baker, in Journ. Linn. Soc. xiv. (1875) 545 = Cordyline floribunda.
schizantha, *Baker, in Journ. Bot.* xv. (1877) 71.—Afr. trop.
septaria, Seem. Fl. Vit. t. 94 = Cordyline terminalis.
siamensis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
Siebôldii, Hort. ex Baker, l. c. = præc.
Siebôldii, Planch. in Fl. des Serres, Sér. I. vi. (1850-51) t. 569 = elliptica.
Smithii, *Baker, in Bot. Mag.* t. 6169.—Afr. trop.
spectabilis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
spicata, *Roxb. Hort. Beng.* 24; *Fl. Ind.* ii. 157.—Ind. or.
splendens, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
stenophylla, *C. Koch, Wochenschr.* iv. (1861) 395.—Afr. trop.
striata, Linn. f. Suppl. 204 = Asparagus striatus.
stricta, Hort. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1847) 11 = Cordyline dracænoides.
stricta, Sims, Bot. Mag. t. 2575 = Cordyline stricta.
sulcata, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
surculosa, *Lindl. in Bot. Reg.* t. 1169.—Afr. trop.
terminalis, Linn. Syst. ed. XII. 246 = Cordyline terminalis.
terminalis, Wight, Ic. t. 2054 = spicata.
ternifora, Roxb. Hort. Beng. 24; Fl. Ind. ii. 159 = spicata.
ternifora, Wall. Cat. n. 5147 A = elliptica.
tessellata, Willd. Enum. Hort. Berol. 374 = marginata.
thalioides, *Hort. Makoy, ex E. Morr. Belg. Hortic.* x. (1860) 348.—Afr. trop.
Thwaitesii, Regel, Gartenfl. (1871) 144 = spicata.
timorensis, *Kunth, in Abh. Acad. Berl.* (1842) 26.—Ins. Timor.
trianæi, Lind. Reised. Guin. 6 = Paliscota thyrsiflora.
Troubetzkoi, *Linden & André, in Illustr. Hortic.* xxi. (1874) 168. t. 187.—Ins. Pacif.
umbraculifera, *Facq. Hort. Schoenbr.* i. 50. t. 95.—Ins. Maurit.
undulata, Linn. f. Suppl. 203 = Asparagus undulatus.
utilis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 540 = Cordyline terminalis.
venenata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 13. (Quid?).—Java.
venusta, *Kerl ex Planch. in Fl. des Serres*, Sér. I. vi. (1850-51) 183.—Hab.?
volubilis, Linn. f. Suppl. 204 = Asparagus scandens, volubilis.
Wallichii, Kunth, in Abh. Acad. Berl. (1842) 26 = spicata.
Warocquei, *Linden & André, Illustr. Hortic.* xxii. (1875) 137. t. 217.—Ins. Solomon.
Weismanni, *T. Moore, Fler. & Pom.* (1878) 202.—Hab.?
xiphophylla, *Baker, in Journ. Linn. Soc.* xxi. (1885) 241.—Madag.
yuccifromis, Vand. Fasc. Pl. 12 = D. Draco.
yuccoides, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1881) 229 = Yucca guatemalensis.
zonlandica, Hort. ex Baker, l. c. xiv. (1873) 544 = Cordyline rubra.

DRACAENOPSIS, Planch. in Fl. des Serres, Sér. I. vi. 110 (1850-51) sub t. 569 = Cordyline, Comm. (Liliac.).
australis, Planch. l. c. = C. australis.
calocoma, H. Wendl. in Bot. Zeit. xvii. (1859) 277 = C. australis.
indivisa, Planch. in Fl. des Serres, Sér. I. vi. (1850-51) 123 = C. indivisa.

DRACO, Crantz, Dusb. Dracon. Arb. 13 (1768) = Dracæna, Vand. (Liliac.), &c.
Cinett, Crantz, l. c. = Dracæna Draco.
Dracunculus, Crantz, l. c. 14 = Pterocarpus Draco.
dragonalis, Crantz, l. c. = Dracæna Draco.
Rhea, Crantz, l. c. 15.—Madag.
Rotenga, Crantz, l. c. 16 = Calamus Rotang.
Sanhilum, Crantz, l. c. 15 = Pterocarpus santalinus?

DRACOCEPHALUM, Hassk. Cat. Hort. Bog. Alt. 292 (1844) = seq.

DRACOCEPHALON, Linn. Syst. ed. I (1735) = Dracocephalum, Linn. (Labiat.).

DRACOCEPHALUM, Linn. Gen. ed. I. 178 (1737).
LABIATAE, Benth. & Hook. f. ii. 1199.
CEPHALOMA, Neck. Elem. i. 314 (1790).
DRACOCEPHALIUM, Hassk. Cat. Hort. Bog. Alt. 292 (1844).
MOLDAVICA, [Tourn.] Linn. Syst. ed. I (1735).
RUYSCHIANA, Linn. l. c. (1735).
RUYSCHIANA, Boeh. ex Mill. Gard. Dict. ed. VII (1759).
acanthoides, Edgew. ex Benth. in DC. Prod. xii. 401 = heterophyllum.
Alberti, *Regel, in Act. Hort. Petrop.* vi. (1879) 362.—Turkestan.
alpinum, Sallsb. Prod. 87 = D. Ruyschiana.
alpinum, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 353.—Sibir. baical.
altaiense, Hort. Vilm. ex Vilm. Fl. Pl. Terre, ed. I 260 = D. Ruyschiana.
altaiense, Laxm. in Nov. Comm. Acad. Petrop. xv (1770) 556. t. 29. f. 3 = grandiflorum.
altaiense, Hiltebr. Dracoc. 50. t. 13 = imberbe.
altaiense, Murr. Syst. ed. XIII. 454 = præc.
angustifolium, Gilib. Fl. Lituan. i. 78 = D. Ruyschiana.
argunense, Fisch. ex Link, Enum. Hort. Berol. ii. 118 = D. Ruyschiana.
aristatum, Bertol. in Nov. Comm. Acad. Bonon. (1844) 325 = Lallemantia iberica.
Aucheri, *Boiss. Diagn.* Ser. I. v. 25.—Persia.
austriacum, *Linn. Sp. Pl.* 595.—Europ.; Kegg. Caucas.
asperum, Boiss. Diagn. Ser. II. iv. 21 = Lallemantia canescens.
balsamicum, Salisb. Prod. 87 = Cedronella triphylla.
bipinnatum, Rupr Sert Tiansch. 65 = Ruprechtii.
botryoides, Boiss. & Buhse, ex Boiss. Fl. Orient. iv. 671 = Ancheri.
botryoides, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 266.—Reg. Caucas.
canariense, Linn. Sp. Pl. 594 = Cedronella triphylla.
canescens, Linn. l. c. 595 = Lallemantia canescens.
chamaedrifolium, *Turcz. ex Benth. in DC. Prod.* xii. 599.—Kamtchat.
chamaedryoides, Balb. Misc. 29 = Sphacele campanulata.
cochinchinense, *Lour. Fl. Cochinch.* 371.—Cochinch.
cordatum, Nutt. Gen. Am. ii. 85 = Cedronella cordata.
crenatifolium, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii (1884) 238.—Turkestan.
cuniloides, *Pall. Reise Statth. Russ. Reich.* ii. 231.—Tauria.
denticulatum, Ait. Hort. Kew. ed. I. ii. 317 = Physostegia virginiana.
discolor, *Bunge, Vers. Suppl. Fl. Alt.* 51.—Sibir. altaic.
diversifolium, *Rupr. Sert. Tiansch.* 66.—Asia centr.
erectum, Koyle, ex Benth. in Hook. Bot. Misc. iii. (1833) 380 = Nepeta erecta.
fastidium, Bunge, in Ledeb. Fl. Alt. ii. 336 = D. Moldavica.
fragile, *Turcz. ex Benth. Lab. Gen. et Sp.* 495.—Dahuria.
fragrans, Sallsb. Prod. 87 = D. Moldavica.
fruticulosum, *Steph. ex Willd. Sp. Pl.* iii. 152.—Sibir.
Germanicum, Benth. in Wall. Pl. As. Rar. ii. 66 = Nepeta Govaniana.
grandiflorum, *Linn. Sp. Pl.* 595.—Sibir.
grandiflorum, Willd. ex Ledeb. Fl. Ross. iii. 355 = imberbe.
heterophyllum, *Benth. Lab. Gen. et Sp.* 738.—Reg. Himal.; Turkest.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 606.—Tibet. or.
hyssopifolium, Mart. ex Steud. Nom. ed. I. 395 = D. Ruyschiana.
ibericum, Bieb. Fl. Taur. Cauc. ii. 54 = Lallemantia iberica.
imberbe, *Bunge, Vers. Suppl. Fl. Alt.* 50.—Sibir. altaic.
inconcum, Sallsb. Prod. 87 = Lallemantia canescens.
inderiense, Lam. ex Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 229 = Lallemantia Royleana.
integrifolium, *Bunge, in Ledeb. Fl. Alt.* ii. 307.—Sibir. altaic.
intermedium, Nutt. in Trans. Am. Phil. Soc. v. (1837) 167 = Physostegia intermedia.
lanchyanthum, *Boiss. Syst. Flavesch.* 65.—As. centr.
lehmannum, *Regel, Pl. Nov. Fedsch.* 66.—As. centr.

DRACOCEPHALUM:—
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 53.—Persia.
lamifolium, Desf. in Ann. Mus. Par. x. (1807) 305 = Lamium striatum.
lancifolium, Moench, Meth. 410 = Physostegia virginiana.
laniflorum, *Rupr. Sert. Tianschi.* 65.—Turkestan.
lignatum, DC. Prod. xii. 397, sphalm. = pinnatum.
mexicanum, H. B. & K. Nov. Gen. et Sp. ii. t. 160 = Cedronella mexicana.
microphyllum, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 364.—Sibir. baical.
Moldavica, *Linn. Sp. Pl.* 595.—Europ.; As. bor.
moldavicum, C. Morr. in Bull. Acad. Belg. iv. (1837) 391 = præc.
multicaule, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 47.—As. Min.; Persia.
nervosum, Rafin. Fl. Ludov. 43 = parviflorum ?
nodulosum, *Rupr. Sert. Tianschi.* 65.—As. centr.
nutans, *Linn. Sp. Pl.* 596.—As. bor.
oblongifolium, *Regel, Pl. Nov. Fedsch.* 67.—Turkestan.
obovatum, Ell. Sketch. ii. 86 = Physostegia virginiana.
ocymifolium, Mill. Gard. Dict. ed. VIII. n. 4 = Lallemantia peltata.
odoratissimum, Poir. Encyc. Suppl. ii. 521 = Dracocephalum taurica.
origanoides, Ledeb. Fl. Alt. ii. 382 = discolor.
origanoides, *Steph. ex Willd. Sp. Pl.* iii. 151.—Sibir.
origanoides, Turcz. ex Benth. in DC. Prod. xii. 398 = pinnatum.
palmatum, *Steph. ex Willd. Sp. Pl.* iii. 151.—Sibir.
parviflorum, *Nutt. in Trans. Am. Phil. Soc.* n. s. v. 136 = Nuttallii.
peltatum, Linn. Sp. Pl. 596 = Lallemantia peltata.
peregrinum, *Linn. Cent. Pl.* ii. 20.—Sibir.
pinnatifidum, Lam. Encyc. ii. 319 = pinnatum.
pinnatum, *Linn. Sp. Pl.* 594.—Sibir.
Royleanum, Benth. in Wall. Pl. As. Rar. i. 65 = Lallemantia Royleana.
rupestre, *Hance, in Journ. Bot.* vii. (1869) 166.—China.
Ruprechti, *Regel, in Act. Hort. Petrop.* vi. (1879) 363.—Tibet. occ.; Turkestan.
Ruyschiana, *Linn. Sp. Pl.* 595.—Europ.; As. bor.
salicifolium, Salisb. Prod. 87 = Lallemantia peltata.
Scheichowii, Turcz. ex Ledeb. Fl. Ross. iii. 388 = palmatum.
scrobiculatum, *Regel, Pl. Nov. Fedsch.* 67.—Turkestan.
setigerum, *Boiss. & Huet, Diagn.* Ser. II. iv. 21.—Armenia.
sibiricum, Linn. Syst. ed. X. 1104 = Nepeta macrantha.
sinense, *S. Moore, in Journ. Linn. Soc.* xvii. (1879) 388.—China.
speciosum, *Benth. in Wall. Pl. As. Rar.* ii. 65.—Reg. Himal.
speciosum, Sweet, Brit. Flow. Gard. t. 98 = Physostegia virginiana.
spicatum, Dulac, Fl. Hautes-Pyr. 399 = D. Ruyschiana.
spinescens, *Wender. Ind. Sem. Hort. Marb.* (1841).—*Cf. Linnæa*, xvi. (1842) Lit. 110.—Hab.?
stamineum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 428.—Reg. Himal.; Sibir.
Stellerianum, *Stend. ex Benth. in DC. Prod.* xii. 402.—Sibiria.
tanguticum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 580.—Mongol.
Tchihatcheffii, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 24 = Salvia Tchihatcheffii.
ternatifolium, Stokes, Bot. Mat. Med. iii. 329 = Cedronella triphylla.
tenuiroides, Bunge, Lab. Pers. 39 = Nepeta lasiocephala.
thymiflorum, *Linn. Sp. Pl.* 596.—Europ.; As. bor.
urticæfolium, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 109.—Japon.
variegatum, Vent. Jard. Cels, t. 44 = Physostegia virginiana.
virginianum, Linn. Sp. Pl. 594 = Physostegia virginiana.

DRACONIA, Heist. Syst. 7 (1748) = **Dracæna** †, Vand. (Liliac.).

DRACONTIUM. Linn. Caroll. Gen. 18 (1787).
AROIDEAE, Benth. & Hook. f. iii. 995.
CHERNEVIRIUM, Schott, in Oester. Bot. Zeitschr. xv. (1866) 73.
ECHIDNIUM, Schott, in Oestr. Bot. Wochenbl. (1857) 62.
EUTERMIA, Rafin. Fl. Tellur. iv. 13 (1836).

DRACONTIUM:—
GODWINIA, Seem. in Journ. Bot. vii. (1869) 313. t. 96, 97.
asperum, *C. Koch, in C. Koch & Fint. Wochenschr.* (1859) 259.—Bras.
bissectum, *Vell. Fl. Flum.* ix. t. 113.—Bras.
camtschaticense, Linn. Sp. Pl. 968 = Lysichiton camtschatcense.
camtschaticum, Spreng. Syst. v. 248 = præc.
Carderi, *Hook. f. Bot. Mag.* t. 6523.—N. Granat.
cordatum, *Auhl. Pl. Gui.* ii. 835.—Guian.
cordatum, Houtt. Handleid. xi. 200. t. 71. f. 2 = Homalomena aromatica.
dubium, *Kunth, Ind. Sem. Hort. Berol.* (1844) 21.—Guiana.
elatum, Mart. in Gard. Chron. (1870) 844 = asperum ?
foecundum, *Hook. f. Bot. Mag.* t. 6808.—Guiana.
foetidum, Linn. Sp. Pl. 967 = Symplocarpus foetidus.
Gigas, *Engl. in DC. Monog. Phan.* ii. 284.—Nicarag.
integerrimum, *Vell. Fl. Flum.* ix. t. 119.—Bras.
hexandrum, Auhl. Pl. Gui. t. 326 = *pentaphyllum*.
lacinialum, Vell. Fl. Flum. ix. t. 120 = Philodendron lacinialum.
lanceæfolium, Jacq. Coll. iv. 118 = Spathiphyllum lanceæfolium.
moluccanum, Zipp. ex Engl. in DC. Monog. Phan. ii. 250 = Epipremnum moluccanum.
pacurinæfolium, Dennst. Schluess. Hort. Malab. 38 = Amorphophallus campanulatus.
pentaphyllum, Aubl. Pl. Gui. ii. 837 = Anthurium pentaphyllum.
pertusum, Forst. f. Prod. 68 n. 381 = Rhaphidophora pertusa.
pertusum, Linn. Sp. Pl. 968 = Monstera pertusa.
polyphyllum, A. Cunn. ex A. DC. in DC. Prod. xvi. 11. 584 = Bowenia spectabilis.
polyphyllum, Dennst. Schluess. Hort. Malab. 38 = Amorphophallus bulbus.
polyphyllum, Forst. f. Pl. Escul. 61. n. 29 = Amorphophallus campanulatus.
polyphyllum, Houtt. Handleid. xi. 199 = Amorphophallus gigauteus.
polyphyllum, *Linn Sp. Pl.* 967.—Am. austr.
Quebitea, J. F. Gmel. Syst. 596 (= *Quebitea guianensis*).—Guiana.
repens, Descourt. Fl. Antill. vii. t. 499 = Anthurium scandens.
sagittæfolium, G. F. W. Mey. Prim. Fl. Esseq. 196 = Urospatha Meyeri.
scandens, Aubl. Pl. Gui. ii. 836 = Anthurium scandens.
spinosum, Linn. Sp. Pl. 967 = Lasia spinosa.
surinamense, Hort. ex Engl. in DC. Monog. Phan. ii. 283 = polyphyllum.
violaceum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. 1. Art. iv. 13 (Quid ?).—Java.
Wallichi, C. Koch, in Hamel. Ind. Arold. 45 = asperum.

DRACONTOCEPHALIUM, Hassk. Cat. Hort. Bog. Alt. 189 (1844) = **Dracocephalum**, Linn. (Labiat.).

DRACONTOMELON, Blume, Mus. Bot. Lugd. Bat. i. 231. t. 49 (1850). *ANACARDIACEAE*, Benth. & Hook. f. i. 427.
COMBUNA, Buill. Adansonia, x. (1872) 239.
costatum, *Blume, l. c.* 233.—Borneo.
Cumingianum, *Baill. in Bull. Soc. Linn. Par.* i. (1877) 152.—Ins. Philipp.
cuspidatum, *Blume, Mus. Bot. Lugd. Bat.* i. 232.—Borneo.
mangiferum, *Blume, l. c.* 231.—Java.
pilosum, *Seem. Fl. Vit.* 92.—Ins. Fiji.
puberulum, *Miq. Fl. Ind. Bat. Suppl.* 524.—Sumatra.
sylvestre, *Blume, Mus. Bot. Lugd. Bat.* i. 231.—Borneo.
vitiense, *Engl. in DC. Monog. Phan.* iv. 253.—Ins. Fiji.

DRACONTOPSIS, Lem. in Orb. Dict. v. 128 (1846) = DRACOPIS, Cass. = **Rudbeckia**, Linn. (Compos.).

DRACOPHYLLUM. Labill. Voy. ii. 210. t. 40 (1798). *EPACRIDEAE*, Benth. & Hook. f. ii. 618.
SPHENOTOMA, R. Fr. Prod. 556 (1810).
acerosum, *Berggr. in Minnesbr. Fiatsg. Sällsk. Lund* (1877) n. VIII. 15.—N. Zel.
affine, Hook. f. Fl. Antarct. 48; Fl. N. Zel. i. 168 = strictum.
amabile, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. ii. (1864) 157.—N. Caled.

DRACOPHYLLUM:—
attenuatum, A. Cunn. ex Walp. Rep. vi. 435 = Urvilleanum.
capitatum, *R. Br. Prod.* 556.—Austral.
cosmeticolor, *Panch. ex Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. ii. (1864) 156.—N. Caled.
Drummondii, *Benth. Fl. Austral.* iv. 368.—Austral.
filifolium, Hook. f. Fl. N. Zel. i. 169 = Urvilleanum.
Fitzgeraldi, *F. Muell. Fragm.* vii. 27.—Austral.
gracile, Brongn. & Gris, in Ann. Sc. Nat. Sér. V. ii. (1864) 156 = gracile.
gracile, *R. Br. Prod.* 556.—Austral.
involucratum, *Brongn. & Gris, l. c.* 157.—N. Caled.
Kirkii, *Berggr. in Journ. Bot.* xviii. (1880) 104.—N. Zel.
latifolium, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 48.—N. Zel.
Lessonianum, A. Rich. Bot. Voy. Astrol. 233 = Urvilleanum.
longifolium, *R. Br. Prod.* 556.—N. Zel.
Lyallii, Hook. f. Fl. N. Zel. i. 169 = longifolium.
Menziesii, *Hook. f. l. c.* 168.—N. Zel.
Milligani, *Hook. fc. Pl.* t. 845.—Tasman.
minimum, *F. Muell. Fragm.* i. 39.—Austral.
muscoides, Hook. f. Handb. N. Zeal. Fl. 188 = minimum.
parviflorum, *F. Muell. ex Benth. Fl. Austral.* iv. 365.—Austral.
Pearsonii, *T. Kirk, in Trans. N. Z. Inst.* xvii. (1885) 223.—N. Zel.
phlogiflorum, *F. Muell. Fragm.* vi. 65.—Austral.
prostratum, *T. Kirk, in Trans. N. Z. Inst.* xiii. (1881) 384.—N. Zel.
ramosum, *Panch. ex Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. ii. (1864) 156.—N. Caled.
recurvum, *Hook. f. Fl. Antarct* 50.—N. Zel.
robustum, Hook. f. l. c. 49 = Urvilleanum.
rosmarinifolium, *R. Br. Prod.* 556.—N. Zel.
scoparium, Hook. f. Fl. Antarct. 46. t. 38 = Urvilleanum.
secundum, *R. Br. Prod.* 556.—Austral.
setifolium, Stchegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 28 = Urvilleanum.
squarrosum, *R. Br. Prod.* 556.—Austral.
squarrosum, *Hook. f. Fl. Antarct.* 48.—N. Zel.
strictum, *Hook. f. l. c.*—N. Zel.
subulatum, *Hook. f. l. c.* 80.—N. Zel.
Thiebautii, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. iii. (1865) 238.—N. Caled.
Traversii, *Hook. f. Handb. N. Zeal. Fl.* 186.—N. Zel.
uniflorum, Bergg. in Minnesbr. Fisiog. Sällsk. Lund (1877) n. viii. t. 4 = Kirkii.
uniflorum, *Hook. f. l. c.* 182.—N. Zel.
Urvilleanum, *A. Rich. Bot. Voy. Astrol.* 221.—N. Zel.
verticillatum, *Labill. Voy.* ii. 211. t. 40.—N. Caled.

DRACOPIS, Cass. in Dict. Sc. Nat. xxxv. 278 (1825) = **Rudbeckia**, Linn. (Compos.).
amplexicaulis, Cass. l. c. = R. amplexicaulis.

DRACUNCULUS, [Rupp.] Ledeb. Fl. Ross. ii. 566, in syn. (1846) = **Arctiolaea**, Linn. (Compos.).
Marschalliana, Ledeb. l. c. (cum cit. falsa) = A. campestris.

DRACUNCULUS, [Tourn.] Adans. Fam. ii. 469 (1763); Schott, Melet. i. 17 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 989.
ANARMODIUM, Schott, in Roeplandia, ix. (1861) 868.
canariensis, *Kunth, Enum. Pl.* iii. 30.—Ins. Canar.
cannæfolius, Rafin. Fl. Tellur. iii. 65 = Spathiphyllum cannæfolium ?
creticus, Schott, Prod. Aroid. 120 = vulgaris.
crinitus, Schott, Melet. i. 17 = Helicodiceros crinitus.
divaricatus, Rafin. Fl. Tellur. iii. 65 = Typhonium divaricatum.
minor, Blume, Rumphia, i. 126 = Helicophyllum crassipes.
muscivorus, Parl. Fl. Ital. ii. 352 = Helicodiceros crinitus.
polyphyllus, Blume, Rumphia, i. 134 = vulgaris.
spadiceus, Rafin. Fl. Tellur. iii. 65 (Quid ?).—Hab.?
trilobatus, Rafin. l. c. = Typhonium divaricatum.
vulgaris, *Schott, Melet.* i. 17.—Europ. austr.

DRAKAEA, Lindl. Swan River App. 55 (1839). *ORCHIDEAE*, Benth. & Hook. f. iii. 511.
ARTHROCHILUS, F. Muell. Fragm. i. 43 (1858).
SPICULARA, Lindl. Swan River App. 56 (1839).

DRAKAEA :—
ciliata, *Reichb. f. Beitr. Syst. Pfl.* 68.—Austral.
elastica, *Lindl. Swan River App.* 56.—Austral.
glyptodon, *Fitzg. in Gard. Chron.* (1893) i. 494.—Austral.
irritabilis, *Reichb. f. Beitr. Syst. Pfl.* 68.—Austral.

DRAKAINA, Rafin. Fl. Tellur. iv. 17 (1836) = **Dracaena**, Vand. (Liliac.).
Draco, Rafin. l. c. = Dracaena Draco.
marginata, Rafin. l. c. = Lomatophyllum borbonicum.

DRAKEA, Endl. Ench. 114 (1841) = **Drakaea**, Lindl. (Orchid.).

DRAKENSTEINIA, DC. Prod. ii. 476 (1825) = seq.

DRAKENSTENIA, Neck. Elem. iii. 38 (1790) = **Boasta-phyllum**, P. Br. (Legumin.).

DRAPARNALDA, St. Lag. in Ann. Soc. Bot. Lyon, viii. 177 (1881) = seq.

DRAPARNAUDIA, Montr. in Mém. Acad. Lyon, x. (1860) 205 = **Xanthostemon**, F. Muell. (Myrtac.).
multiflora, Montr. l. c. 206.—N. Caled.

DRAPERIA, Torr. in Proc. Am. Acad. vii. (1868) 401.
HYDROPHYLLACEAE, Benth. & Hook. f. ii. 830.
nyctyla, *Torr. l. c.*—Calif.

DRAPETES, Banks, ex Lam. in Journ. Hist. Nat. Par. i. (1792) 183. t. 10. *THYMELAEACEAE*, Benth. & Hook. f. iii. 196.
DAPHNORRYON, Meisan. in DC. Prod. xiv. 566 (1857).
KELLERIA, Endl. Gen. Pl. Suppl. iv. 11. 61 (1847).
Dieffenbachii, *Hook. in Lond. Journ. Bot.* ii. (1843) 497.—N. Zel.
ericoides, *Hook. f. in Ic. Pl.* t. 895.—Borneo.
Lyallii, *Hook. f. Fl. N. Zeal.* ii. 836.—N. Zel.
muscoides, Pers. Syn. i. 148 = seq.
muscosus, Hook. f. Fl. N. Zel. 238 = Lyallii.
muscosus, *Lam. in Journ. Hist. Nat. Par.* i. (1792) 189.—Reg. Magell.
tasmanicus, *Hook. f. in Hook. New Journ.* v. (1853) 209. t. 7.—Austral.

DRAPIEZIA, Blume, Enum. Pl. Jav. ii (1827) = **Dissporum**, Salisb. (Liliac.).
multiflora, Blume, l. c. = D. pullum.

DRAYTONIA, A. Gray, Bot. U. St. Expl. Exped. i. 206. t. 15 (1854) = **Saurauja**, Willd. (Ternstroem.).
rubicunda, A. Gray, l. c. = S. rubicunda.

DREBBELIA, Zoll. & Morr. in Nat. en Geneesk. Arch. Nederl. Indie, iii. (1846) 79 = **Spatholobus**, Hassk. (Legumin.).
ferruginea, Zoll. & Morr. l. c. = S. ferrugineus.

DREBBELLIA, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 160. *OLACINEAE*?
subarborescens, *Zoll. l. c.*—Ins. Bali.

DREGEA, Eckl. & Zeyh. Enum. 350 (1837) = **Pouchetdanum**, Tourn. (Umbellif.).
capensis, Eckl. & Zeyh. l. c. 351 = P. capense.
collina, Eckl. & Zeyh. l. c. = P. collinum.
montana, Eckl. & Zeyh. l. c. = P. Ecklonianum.
virgata, Eckl. & Zeyh. l. c. = P. capense.

DREGEA, E. Mey. Comm. Pl. Afr. Austr. 199 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 776.
PTEROPHORA, Harv. Gen. S. Afr. Pl. 293 (1838).
PTERYGOCARPUS, Hochst. in Flora, xxvi. (1843) 78.
WATTAKAKA, Hassk. in Flora, xl. (1857) 99.
arabica, *Decne. in DC. Prod.* viii. 618.—Arab.
floribunda, E. *Mey. Comm. Pl. Afr. Austr.* 199.—Afr. austr.
macrantha, *Klotzsch, in Peters, Reise Mossamb. Bot.* 272.—Afr. trop.
volubilis, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 46.—Ind. or.

DREJERA, Nees, in Mart. Fl. Bras. ix. 112. t. 17 (1847) = Jacobinia, Moric. (Acanthac.).
heterantha, Nees, in DC. Prod. xi. 434.—Bolivia.
Greggii, Torr. in Bot. Mex. Bound. 134 = Anisacanthus Greggii.
juncea, Torr. l. c. = Anisacanthus juncea.

DREJERA :—
parviflora, Beckl. in Proc. Acad. Sc. Philad. 1861 (1862) 462 = Dianthera parviflora.
puberula, Torr. in Bot. Mex. Bound. 128 = Anisacanthus virgularis.
ramosa, Nees, in Mart. Fl. Bras. ix. 112.—Bras.
Thurberi, Torr. in Bot. Mex. Bound. 134 = Anisacanthus Thurberi.
virgata, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 154 = Jacobinia virgata.
Wrightii, Torr. in Bot. Mex. Bound. 129 = Anisacanthus Wrightii.
Willdenowiana, Nees, in DC. Prod. xi. 384 = Jacobinia Mohintii.

DREPACHENIA, Rafin. Neogenyt. 3 (1825) = **Sagittaria**, Rupp. (Alismac.).
lancifolia, Rafin. l. c. = S. lancifolia.

DREPADENIUM, Rafin. Neogenyt. 2 (1825) = **Croton**, Linn. (Euphorb.).
dichotomum, Rafin. Sylva Tellur. 65 = Bernardia dichotoma.
maritimum, Rafin. Neogenyt. 2 = Croton maritimus.

DREPANANDRUM, Neck. Elem. ii. 118 (1790) = **Topobea**, Aubl. (Melastom.).

DREPANANTHUS, Maing. ex Hook. f. Fl. Brit. Ind. i. 56 (1872). *ANONACEAE*.
pruniferus, *Maing. l. c.*—Malacca.
ramuliflorus, *Maing. l. c.*—Malacca.

DREPANIA, Juss. Gen. 169 (1789) = **Tolpis**, Adans. (Compos.).
ambigua, DC. Cat. Hort. Monsp. 105 = T. altissima.
barbata, Desf. Fl. Atlant. ii 232 = T. barbata.
falcata, Roxb. Fl. Calv. 127 = T. barbata.
pallida, Lowe, Nom. ed. II. i. 530 = T. barbata.
umbellata, DC. Cat. Hort. Monsp. 105 = T. umbellata.

DREPANOCARPUS, G. F. W. Mey. Prim. Fl. Esseq. 236 (1818). *LEGUMINOSAE*, Benth. & Hook. f. i. 546.
NEPHROSIS, Rich. ex DC. Prod. ii. 420 (1825).
ONOSERIA, Juss. ex DC. l. c. (1825).
SOMMERFELTIA, Schum. & Thonn. Beskr. Guin. Pl. 331 (1828).
africanus, *O. Don, Gen. Syst.* ii. 378.—Afr. trop.
aristulatus, *Spruce, ex Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 69.—Bras.
Cristo-castroisis, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 96.—Bras.
Cuminghii, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 262 = Dalbergia Cuminghiana.
cynathiformis, *DC. Prod.* ii. 420.—Mexic.
discolor, Benth. in Mart. Fl. Bras. xv. t. 227 = Dalbergia inundata.
dubius, *R. & P. K. Nov. Gen. et Sp.* vi. 390.—Venezuela.
falcatus, *Mig. in Linnaea*, xviii. (1844) 576.—Guiana.
ferox, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 96.—Bras.
floridus, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 118.—Bras.
frondosus, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 96.—Bras.
inundatus, Mart. ex Benth. l. c. = Dalbergia inundata.
lasiolophus, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 507 = Machaerium angustifolium.
lunatus, *G. F. W. Mey. Prim. Fl. Esseq.* 238.—Am. trop.
microphyllus, *E. Mey. in Nov. Act. Nat. Cur.* xii. (1825) 508.—Panama.
microphyllus, *Wawra, in Flora*, xlvii. (1864) 241.—Bras.
monosperma, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 261 = Dalbergia monosperma.
mucronulatus, *Benth. ex Hemsl. Diag. Pl. Nov.* 8.—Mexic.
paniculatus, *G. Don, Gen. Syst.* ii. 378.—Ins. S. Doming.
polyphyllus, Benth. in Ann. Wien. Mus. ii. (1838) 96 = Machaerium acuminatum.
reniformis, Kurz, in Pegu Rep. App. A 49 & B 45 = Dalbergia reniformis.
Schomburgkii, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1906, nomen.—Guiana.
sonquinensis, T. Nees & Ebermaier, Handb. Med. Pharm. Bot. iii. 184 = Pterocarpus erinaceus.
spinosus, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 261 = Dalbergia spinosa.

DREPANOLOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 324 (1840) = **Homokis**, Dougl. (Legumin.).
crassifolius, Nutt. ex Torr. & Gray, l. c. 325 = H. crassifolia.
cytisoides, Nutt. ex Torr. & Gray, l. c. 324 = H. cytisoides.
decumbens, Nutt. ex Torr. & Gray, l. c. = H. decumbens
junceus, Nutt. ex Torr. & Gray, l. c. 325 = H. juncea
lanatus, Nutt. ex Torr. & Gray, l. c. 324 = H. tomentosa.
micranthus, Nutt. ex Torr. & Gray, l. c. = H micrantha.
prostratus, Nutt. ex Torr. & Gray, l. c. 325 = H. prostrata.
rubescens, Nutt. ex Torr. & Gray, l. c. 324 = H. cytisoides.
scoparius, Nutt. ex Torr. & Gray, l. c. 325 = H. glabra
DREPANOPHYLLUM, Wibel, Prim. Fl. Werth. 196 (1799) = **Falcaria**, Host (Umbellif.).
agreste, Hoffm. Umbellif. i. 109 = F. Rivini.
luteum, Eichw. Skizze, 257 = Muretia tanaicensis.
patustre, Hoffm. Umbellif. 110 = Stum latifolium.
riodes, Wibel, Prim. Fl. Werth. 196 = F. Rivini.

DREPANOSPERMUM, Benth. in Benth. & Hook. f. Gen. i. 425 (1862) = **Campnosperma**, Thw. (Anacard.).
gummiferum, Engl. in Mart. Fl. Bras. xii. ii. 393 = C. gummifera.

DREPAPHYLA, Rafin. Sylva Tellur. 120 (1838) = **Acacia**, Willd. (Legumin.).
lanigera, Rafin. l. c. = A. lanigera.
multinervis, Rafin. l. c. = A. lanigera.

DREPILIA, Rafin. New Fl. Am. ii. 52 (1836) = **Thermopsis**, R. Br. (Legumin.).
rhombifolia, Rafin. l. c. 53 = T. rhombifolia.

DRESSSINIA, Korth. Verh. Nat. Gesch. Bot. 251. t 53 (1844). *MELASTOMACEAE*, Benth. & Hook. f. i. 785.
xanthe, *Korth. l. c.*—Borneo.
bulbosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 78.—Malacca.

DRIMIA, Jacq. Ic. Rar. t. 373–377 (1786). *LILIACEAE*, Benth. & Hook. f. iii. 898.
IDOTHEA, Kunth, Enum. Pl. iv. 341 (1843).
IDOTHEARIA, Presl, Bot. Bemerk. 114 (1844).
STREPIDIPHYLA, Rafin. Fl. Tellur. iii. 60 (1836).
acuminata, Lodd. Bot. Cab. t. 1041 = Scilla lanceafolia.
altissima, *Hook. f. Bot. Mag.* t. 5522.—Afr. austr.
altissima, [Ker-Gawl. in] Bot. Mag. t. 1074 = Urginea altissima.
angustifolia, Kunth, Enum. Pl. iv. 340 = Scilla linearifolia.
apertiflora, Baker, in Saund. Refug. Bot. t. 19 = Scilla lorata.
Barteri, *Baker, in Journ. Linn. Soc.* xi. (1871) 423.—Afr. trop.
bifolia, Schweinf. Beitr. Fl. Aethiop. 294 = Scilla Richardiana.
Burchelli, *Baker, in Saund. Refug. Bot. ii. App.* 2.—Afr. austr.
ciliaris, *Jacq. Coll. Suppl.* 41.—Afr. austr.
concolor, *Baker, in Journ. Linn. Soc.* xi. (1871) 422.—Afr. austr.
Cooperi, Baker, in Saund. Refug. Bot. t. 18 = Scilla concolor.
Cowanii, *Ridl. in Journ. Linn. Soc.* xx. (1883) 334.—Madag.
Dregeana, Kunth, Enum. Pl. iv. 340 = Scilla concolor.
Ecklониana, *Schult. f. Syst.* vii. 1710.—Afr. austr.
elata, *Jacq. Coll. Suppl.* 36.—Afr. austr.
exaltata, Eckl. in Linnaea, xx. (1847) 286 = Scilla Ludwigii.
Gasnieri, Schrad. Blumenb. 30 = Scilla revoluta.
hawarthioides, *Baker, in Gard. Chron.* (1875) i. 366.—Afr. austr.
humilis, Berg. ex Eckl. Top. Vers. 9 = Ledebouria.
hyacinthoides, *Baker, in Journ. Bot.* xii. (1874) 6.—Afr. austr.
lanceaefolia, Ker-Gawl. in Bot. Mag. sub t. 1330 = Scilla lanceaefolia.
lanceaefolia, Lodd. Bot. Cab. t. 378 = Scilla revoluta.
lanceolata, Schrad. Blumenb. 30 = Scilla lanceolata.
lanifera, *Baker, in Journ. Linn. Soc.* xi. (1871) 423.—Afr. trop.

DRIMIA :—
lilacina, Schwemf. Beitr. Fl. Aethiop. 294 = Scilla lilacina.
longipedunculata, G. Don, in Loud. Hort. Brit. Suppl. i. 587 = Scilla lanceaefolia.
Ludwigii, Miq. in Bull. Sc. Phys. Néerl. (1839) 89 = Scilla Ludwigii.
media, *Jacq. Coll. Suppl.* 40.—Afr. austr.
ovalifolia, Schrad. Blumenb. 30 = Scilla revoluta.
pendula, Baroh. ex Baker, in Journ. Linn. Soc. xiii. (1873) 254 = Scilla pendula.
petiolata, *C. Koch & Bouche, Ind. Sem. Hort. Berol.* (1851) *App.* 3.—Hab.?
purpurascens, *Jacq. f. Eclog.* t. 30.—Afr. austr.
pusilla, *Jacq. Coll. Suppl.* 42.—Afr. austr.
revoluta, Kunth, Enum. Pl. iv. 341 = Scilla revoluta.
rigidifolia, *Baker, in Journ. Linn. Soc.* xi. (1871) 420. —Afr. austr.
robusta, *Baker, in Saund. Refug. Bot.* t. 190.—Afr. austr.
undulata, Jacq. Coll. Suppl. 41 = Scilla undulata.
villosa, *Lindl. in Bot. Reg.* t. 1846.—Afr. austr.
viridiflora, Eckl. Top. Verz. 2 = Scilla viridiflora.

DRIMIOPSIS, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 73. f. 172. *LILIACEAE*, Benth. & Hook. f. iii. 818.
Barteri, *Baker, in Saund. Refug. Bot.* iii. *App.* 18.—Afr. trop.
botryoides, *Baker, in Journ. Bot.* xii. (1874) 364.—Afr. trop.
Burkei, *Baker, in Saund. Refug. Bot.* iii. *App.* 17.—Afr. austr.
Kirkii, *Baker, in Gard. Chron.* (1874) ii. 644.—Afr. trop.
maculata, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 73.—Afr austr.
minor, *Baker, in Saund. Refug. Bot.* t. 192.—Afr. trop.
perfoliata, *Baker, in Gard. Chron.* (1878) ii. 364. —Afr. trop.

DRIMOPHYLLUM, Nutt. N. Am. Sylv. i. 85 (1842) = Umbellularia, Nutt. (Laurin.).
pauciflorum, Nutt. l. c. = U. californica.

DRIMOPOGON, Rafin. Sylva Tellur. 151 (1838) = Spiraea, Linn. (Rosac.).
Douglasi, Rafin. l. c. = S. Douglasii.
ferruginea, Rafin. l. c. = S. tomentosa.
glomerata, Rafin. l. c. = S. tomentosa.
Menziesii, Rafin. l. c. = S. Douglasii.
parvifolia, Rafin. l. c. = S. tomentosa.
rosea, Rafin. l. c. = S. tomentosa.
tomentosa, Rafin. l. c. = S. tomentosa.

DRIMYCARPUS, Hook. f. in Benth. & Hook f. Gen. i. 424 (1863). *ANACARDIACEAE*, Benth. & Hook. f. i. c.
racemosus, Hook. f. l. c.—Reg. Himal.

DRIMYPHYLLUM, Burch. ex DC. Prod. v. 501 (1836) = Petrobium, R. Br. (Compos.).
hirsutum, Burch. ex DC. l. c. 502 = P. arboreum.

DRIMYS, Forst. Char. Gen. 88. t. 42 (1776). *MAGNOLIACEAE*, Benth. & Hook. f. i. 17.
MAGALLANA, Comm. ex DC. Syst. i. 443 (1818).
MAGELLANIA, Comm. ex Lam. Encyc. ii. 280 (1786).
TASMANNIA, R. Br. in DC. Syst. i. 445 (1818).
WINTERA, *Murr. Syst. ed. XIV.* 567 (1784).
angustifolia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. ii. (1858) 46.—Reas.
aromatica, Descourt. Fl. Méd. Antill. i. 188 = Winteri.
aromatica, *F. Muell. Viet. Pl.* i. 30.—Austral.
axillaris, *Forst. Char. Gen.* t. 42.—N. Zel.
Balansae, *Baill. Adansonia*, x. (1871–72) 335.—N. Caled.
brasiliensis, Miers, in Ann. & Mag. Nat. Hist. Ser. III. ii. (1858) 47 = Winteri.
chilensis, DC. Syst. i. 444 = Winteri.
colorata, *Raoul, Choix Pl. N. Zél.* 24.—N. Zel.
confertiflora, Phil. in Bot. Zeit. xiv. (1856) 641 = Winteri.
crassifolia, *Baill. Adansonia*, viii. (1867–68) 199.—N. Caled.
dipetala, *F. Muell. Pl. Vict.* i. 21.—Austral.
fernandeziana, Steud. in Flora, xxxix. (1856) 408 = Winteri.
granatensis, Linn. f. Suppl. 269 = Winteri.
hatamensis, *Becc. Malesia*, i. (1877) 185.—N. Guinea.
Howeana, *F. Muell. Fragm.* vii. 17.—Ins. D. Howe.
lanceolata, Baill. Hist. Pl. i. 159 = aromatica.
ledifolia, Herb. ex Etchl. in Mart. Fl. Bras. xiii. i. 186 = Winteri.
magnistaminea, Kunth, ex Etchl. in Mart. Fl. Bras. xiii. i. 135 = Winteri.
membranea, *F. Muell Fragm.* v. 175.—Austral.
mexicana, Mocq. & Sesse, ex DC. Syst. i. 444 = Winteri.
montana, Miers, in Ann. & Mag. Nat. Hist. Ser. III. ii. (1858) 44 = Winteri.
paniculata, Steud. in Flora, xxxix. (1856) 408 = Winteri.
Parcheri, *Basil. Adansonia*, x. (1871–72) 336.—N. Caled.
piperita, *Hook. f. Ic. Pl.* t. 896.—Borneo.
punctata, Lam. Encyc. ii. 280 = Winteri.
retorta, Miers, in Ann. & Mag. Nat. Hist. Ser. III. ii. (1858) 45 = Winteri.
uniflora, *Turcz in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 280.—Venezuela.
Winteri, *Forst. Char. Gen.* 84. t. 42.—Am. austr.

DRIMYSPERMUM, Reinw. in Syll. Ratisb. ii. (1828) 15 = Phaleria, Jack (Thymel.).
acuminatum, Seem. Fl. Vit. 209.—Ins Fiji.
ambiguum, Meisn. in DC. Prod. xiv. 605.—Java.
Billardieri, Decne. Bot. Voy. Vénus, 16.—Ins. Amicor.
Blumei, Decne. in Ann. Sc. Nat. Sér. II xix. (1843) 39 = F. Blumei.
Burmanni, Decne. l. c. 40.—Ind. or.
Burnettianum, Seem. Fl. Vit. 208.—Ins. Fiji.
cauliflorum, Thw. Enum Pl. Zeyl. 251 = P. cauliflora.
Clerodendron, F. Muell. Fragm. vii. 1 = P. Clerodendron.
Cumingianum, Decne. Bot. Voy. Vénus, 17 = seq.?
Cumingii, Meisn. in DC. Prod. xiv. 605 = P. Cumingii.
Forsteri, Meisn. l. c.—Ins. Tongatabu.
lanceolatum, A. Gray, in Journ. Bot. iii. (1865) 304. —Ins Fiji.
laurifolium, Decne. in Ann. Sc. Nat. Sér. II. xix. (1843) 39 = P. laurifolia.
laurifolium. Zoll. Verz. ii. 172 = *ambiguum*.
longifolium, Miq. Fl. Ind. Bat. i. i. 885 = *ambiguum*?
macrocarpum, Scheff. in Ann. Jard. Buitenz. (1856) 46. —N. Guin.
macrophyllum, Decne. Bot. Voy. Vénus, 15.—N. Hibern.
montanum, Seem. Fl. Vit 209. t. 54.—Ins. Fiji.
Neumanni, F. Muell. Fragm. v. 26 = P. Neumanni.
Perrottetianum, Decne. in Ann. Sc. Nat. Sér. II. xix. (1843) 40 = P. Perrottetiana.
Phaleria, Meisn. in DC. Prod. xiv. 604 = P. capitata.
pubiflorum, Seem. Fl. Vit. 208.—Ins. Fiji.
revolutum, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 30.—N. Guinea.
subcordatum, Seem. Fl. Vit. 209.—Ins. Fiji.
urens, Reinw. in Syll. Ratisb. ii. (1828) 15.—Ins. Celebes.

DRIPAX, Noronha, ex Thou. Hist. Vég. Afr. ii. 59 (1807) = Alsodeia, Thou. (Violar.).
lanceifolia, Noronha, ex Thou. l. c. = A. angustifolia.

DROBROWSKIA, Brongn. Enum. Gen. 81 (1843) = DOBROWSKYA, Presl = Lobelia, Linn.

DROGOUETIA, Steud. Nom. ed. II. i. 531 (1840) = seq.

DROGUETIA, Gaudich. in Freyc. Voy. Bot. 505 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 394.
DIDYMOCYNE, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 207.
amblgua, *Wedd. in Ann. Sc. Nat. Sér. IV.* i. (1854) 211. —Afr. austr.
cuneata, Buch, Ind. Gen. DC. Prod. iv. 122 = ambigua.
diffusa, *Wedd. in Ann. Sc. Nat. Sér. IV.* i. (1854) 211. —Ind. or.; Afr. trop.
elliptica, Gaudich. in Freyc. Voy. Bot. 505; et Bot. Voy. Bonite, t. 85 = ovata.
leptostachys, Wedd. in DC. Prod. xvi. i. 235* = ovata.
ovata, *Gaudich. in Freyc. Voy. Bot.* 505.—Ins. Mascar.
pauciflora, Wedd. in DC. Prod. xvi. i. 235* = diffusa.
Thouarsiana, *Gaudich. Bot. Voy. Bonite*, t. 87.—Ins. Mascar.

DROGUETIA:—
urticaefolia, Wedd. in DC. Prod. xvi. i. 235* = ambigua.
urticoides, *Wedd. in Ann. Sc. Nat. Sér. IV.* i. (1854) 211.—Ind. or.

DROMOPHYLLA, Lindl. Veg. Kingd. 815 (1847) = DERMOPHYLLA, Manso = Cayaponia, Manso (Cucurb.).

DROSANTHE, Spach, in Ann. Sc. Nat. Sér. II. v (1836) 355 = Hypericum, Tourn.
fimbriata, Spach, Hist. Vég. Phan. v. 378 = H. asyroides.
helianthemoides, Spach, l. c. 379 = H. helianthemoides.
hirtella, Spach, l. c. 377 = H. hirtellum.
hyssopifolia, Spach, l. c. 360 = H. scabrum.
ledifolia, Spach, l. c. 378 = H. nudiflorum.
scabra, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 355; et l. c. 381 = H. scabrum.

DROSANTHUS, R. Br. ex Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 305 = Byblis, Salisb. (Droserac.).

DROSERA, Linn. Syst. ed. I (1735). *DROSERACEAE*, Benth. & Hook. f. i. 662.
ADENOPA, Rafin. Fl. Tellur. iii. 37 (1836)
DIDMOPHYLA, Rafin. l. c. 36 (1836).
EDERA, Neck. Elem. ii. 160 (1790).
FILICINNA, Rafin. Fl. Tellur. iii. 37 (1836).
RORELLA, Hall. ex Ait. Fl. Pedem. ii. 88 (1785).
ROSOLIS, Adans. Fam. ii. 345 (1763).
SONDERA, Lehm. Pugill. viii. 44 (1844).
acaulis, *Thunb. Prod. Fl. Cap.* 57.—Afr. austr.
Adelae, *F. Muell. Progr.* iv. 154.—Austral.
adscendens, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 195 = ascendens.
affinis, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 402.—Afr. trop.
albiflora, Banks, ex Planch. in Ann. Sc. Nat. Sér. III. ix. (1842) 199 = pauciflora.
Aldrovanda, *F. Muell. Fragm.* x. 79.—Austral.
americana, Willd. Enum. Hort. Berol. 340 = longifolia.
anglica, DC. Prod. i. 318 = longifolia.
anglica, *Huds. Fl. Angl.* ed. II. i. 185.—Europ.; Am. bor.
angustifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 7.—Austral.
Arcturi, *Hook. in Journ. Bot.* i. (1834) 247.—Austral.; N. Zel.
ascendens, *St. Hil. Pl. Rem. Brés.* 268.—Bras.
auriculata, *Backh. ex Planch. in Ann. Sc. Nat. Sér. III.* ix. (1848) 196.—Austral.; N. Zel.
Banksii, *R. Br. ex DC. Prod.* i. 319.—Austral.
barbigera, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 287 = Drummondii.
biflora, Willd. ex Roem. & Schult. Syst. vi. 768 = pusilla.
Biliardiera, Tratt. ex Steud. Nom. ed. II. i. 231 = binata.
binata, *Labill. Nov. Holl. Pl.* i. 78. t. 105.—Austral.; N. Zel.
brasiliensis, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. xiv. ii. 396 = graminifolia.
brevifolia, *Pursh, Fl. Am. Sept.* i. 211.—Am. bor.
bulbosa, *Hook. Ic. Pl.* t. 875.—Austral.
Burkeana, *Planch. in Ann. Sc. Nat. Sér. III.* ix. (1848) 192.—Afr. austr.
Burmanni, DC. Prod. i. 318 = Lowreiri.
Burmanni, *Vahl, Symb. Bot.* iii. 50.—As. et Austral. trop.
calycina, *Planch. in Ann. Sc. Nat. Sér. III.* ix. (1848) 209.—Austral.
capensis, *Linn. Sp. Pl.* 282.—Afr. austr.
capillaris, *Poir. Encyc.* vi. 299.—Am. bor.
cistifolia, *Linn. Pl. Rar. Afr.* 9; *Amoen. Acad.* vi. 85. —Afr. austr.
communis, *St. Hil. Fl. Rem. Brés.* 267.—Bras.
cuneifolia, *Thunb. Prod. Fl. Cap.* 57.—Afr. austr.
Cunninghamii, Walp. Rep. i. 229 = binata.
curvipes, *Planch. in Ann. Sc. Nat. Sér. III.* ix. (1848) 196.—Afr. austr.
dentata, Benth. in Hook. Journ. Bot. iv. (1842) 105 = sessilifolia.
dichotoma, Banks & Soland. ex Sm. in Rees, Cycl. xii. = binata.
dichrosepala, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 343.—Austral.
Dietrichiana, *Reichb. f. Beitr. Syst. Pfl.* 78.—Austral.

DROSERA:—
Drummondii, *Lehm. Pl. Preiss.* ii. 235.—Austral.
Drummondii, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 293 = paniculiflora.
erythrorhiza, *Lindl. Swan River App.* 20. n. 91.—Austral.
filiscalix, *Endl. in Enum. Pl. Hueg.* 6.—Austral.
filiformis, *Raf. in Med. Repos. N. York*, v. (1808) 360.—Am. bor.
filipes, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 344.—Austral.
Finlayoniana, Wall. Cat. n. 8753 = indica.
flabellata, *Benth. Fl. Australiæ*. ii. 464.—Austral.
fleetiana, *Walw. ex Oliver, Fl. Trop. Afr.* ii. 408.—Afr. trop.
foliosa, Ell. Sketch. i. 376 = intermedia.
foliosa, Hook. f. ex Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 293 = peltata.
fulva, Planch. l. c. 289 = petiolaris.
gigantea, *Lindl. Swan River App.* 20. n. 86.—Austral.
glanduligera, *Lehm. Pugill.* viii. 37.—Austral.
gracilis, Hook. f. ex Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 297 = peltata.
graminifolia, *St. Hil. Pl. Rem. Brés.* 269. t. 25.—Bras.
grandiflora, *Bartl. in Linnæa*, vii. (1843) 630.—Afr. austr.
Hellanthemum, *Planch. in Ann. Sc. Nat.* Sér. III. ix. (1848) 303.—Afr. austr.
heterophylla, *Lindl. Swan River App.* 20. n. 80.—Austral.
hexagyna, Blanco, Fl. Filip. ed. II. 159 = indica.
hilaris, *Chem. & Schlecht. in Linnæa*, i. (1826) 548.—Afr. austr.
hirtella, *St. Hil. Pl. Rem. Brés.* 262.—Bras.
Huegelii, *Endl. in Enum. Pl. Hueg.* 6.—Austral.
humilis, *Planch. in Ann. Sc. Nat.* Sér. III. ix. (1848) 300.—Austral.
incisa, *A. Rich. Ess. Fl. Cub.* 102.—Cuba.
indica, *Linn. Sp. Pl.* 282.—Geront. trop.
intermedia, R. Cunn. in Ann. Nat. Hist. iv. (1840) 210 = binata.
intermedia, *Hayne, in Schrad. Journ. Bot.* (1800) i. 87; *Drev. & Hayne, Pl. Eur.* iii. 48. t. 75.—Europ.
intermedia, Herb. Royle, ex Hook. f. Fl. Brit. Ind. ii. 435 = peltata.
intermedia, Wight & Arn. Prod. 34 = anglica.
intricata, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 293 = Menziesii.
lciocnema, *Raf. in Fl. Tellur.* iii. 37.—Hab.?
leucoblasta, *Benth. Fl. Australiæ*. ii. 458.—Austral.
linearis, *Goldie, in Edinb. Phil. Journ.* (1832) 325.—Am. bor.
Lobbiana, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 343 = peltata.
longifolia, Drev. & Hayne, Pl. Eur. iii. 42. t. 75 = anglica.
longifolia, *Linn. Sp. Pl.* 282.—Europ.; Am. bor.
longifolia, Sm. Fl. Brit. i. 347 = intermedia.
Lourieii, *Hook. & Arn. Bot. Beech. Voy.* 167.—China.
lunata, Buch.-Ham. ex DC. Prod. i. 319 = peltata.
lusitanica, Linn. Sp. Pl. 282 = Drosophyllum lusitanicum.
macrantha, *Endl. in Enum. Pl. Hueg.* 6.—Austral.
macrophylla, *Lindl. Swan River App.* 20. n. 91.—Austral.
madagascariensis, DC. Prod. i. 318 = ramentacea.
maritima, *St. Hil. Pl. Rem. Brés.* 264. t. 25.—Bras.
Menziesii, *R. Br. ex DC. Prod.* i. 319.—Austral.
microphylla, Pugill. viii. 39 = paleacea.
microphylla, Endl. in Enum. Pl. Hueg. 6 = filicalix.
minor, Schum. & Thonn. Beskr. Guin. Pl. 167 = indica.
minor, Wood, Class-book, 251 = capillaris.
minutiflora, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 296 = paleacea.
montana, *St. Hil. Pl. Rem. Brés.* 260.—Bras.
myriantha, *Planch. in Ann. Sc. Nat.* Sér. III. ix. (1848) 291.—Austral.
Neesii, *Lehm. Pugill.* viii. 42.—Austral.
nitidula, *Planch. in Ann. Sc. Nat.* Sér. III. ix. (1848) 295.—Austral.
obovata, Mert. & Koch, Deutschl. Fl. ii. 502 = longifolia.
paleacea, *DC. Prod.* i. 318.—Austral.
pallida, *Lindl. Swan River App.* 20. n. 81.—Austral.
parvifolia, *St. Hil. Pl. Rem. Brés.* 268.—Bras.
parvula, *Planch. in Ann. Sc. Nat.* Sér. III. ix. (1848) 297.—Austral.
patuliflora, Planch. l. c. 309 = glanduligera!
pauciflora, *Banks, ex DC. Prod.* i. 317.—Afr. austr.

DROSERA:—
pedata, Pers. Syn. i. 387 = binata.
peltata, *Sm. ex Willd. Sp. Pl.* i. 1546.—Austral.
peltata, Wight & Arn. Prod. i. 34 = binata.
pendulifolia, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 301 = ramulosa.
penicillaris, *Benth. Fl. Australiæ*. ii. 457.—Austral.
petiolaris, *R. Br. ex DC. Prod.* i. 318.—Austral.
petiolaris, Sieber, ex Presl, Bot. Bemerk. 14 = peltata.
Planchoni, Hook. f. Fl. Tasm. i. 29 = Menziesii.
planifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 343.—Austral.
platystigma, *Lehm. Pugill.* viii. 37.—Austral.
porrecta, Salisb. l. c. 41 = stoloniféra.
Preissii, *Lehm. l. c.* 45.—Austral.
primulacea, Schlotthaub. in Bonplandia, iv. (1856) 110 = erythrorhiza.
propinqua, R. Cunn. in Ann. Nat. Hist. iv. (1840) 209 = spathulata.
pulchella, *Lehm. Pugill.* viii. 38.—Austral.
purpurascens, Schlotthaub. in Bonplandia, iv. (1856) 111 = stoloniféra.
pusilla, R. Br. ex DC. Prod. i. 317 = pygmæa.
pusilla, *H. B. & K. Nov. Gen. et Sp.* v. 290. t. 490. f. 1.—Venezuela.
pygmæa, *DC. Prod.* i. 317.—Austral.; N. Zel.
pygmæa, Lehm. in Pl. Preiss. i. 250 = paleacea.
ramellosa, *Lehm. Pugill.* viii. 40.—Austral.
ramentacea, *Burch. ex Herv. & Sond. Fl. Cap.* i. 77.—Afr. austr.
Roraimæ, *Klotzsch, in Schomb. Faun. et Fl. Gui. Ined.* 1004.—Guiana.
Roridula, Thunb. Diss. Drosera, 7 = Roridula dentata.
rosulata, Behr. in Linnæa, xx. (1847) 628 = Whittakeri.
rosulata, *Lehm. Pugill.* viii. 36.—Austral.
rotundifolia, *Linn. Sp. Pl.* 282.—Reg. bor.
rotundifolia, Lour. Fl. Cochinch. i. 283 = Lourieii.
scorpioides, *Planch. in Ann. Sc. Nat.* Sér. III. ix. (1848) 288.—Austral.
septentrionalis, Stokes, Bot. Mat. Med. ii. 189 = rotundifolia.
serpens, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 304 = indica.
sessiliflora, G. Don, Gen. Syst. i. 344, sphalm. = sessilifolia.
sessilifolia, Raf. in Atl. Journ. 148 = Dionæa muscipula!
sessilifolia, *St. Hil. Pl. Rem. Brés.* 259. t. 25.—Bras.
spathulata, *Labill. Nov. Holl. Pl.* i. 79. t. 106. f. 1.—Austral.; N. Zel.
speciosa, *Presl, Bot. Bemerk.* 14.—Afr. austr.
spiralis, *St. Hil. Pl. Rem. Brés.* 270.—Bras.
squamosa, *Benth. Fl. Australiæ*. ii. 463.—Austral.
stenopetala, *Hook. f. Fl. N. Zel.* 19. t. 9.—N. Zel.
stoloniféra, *Endl. in Enum. Pl. Hueg.* 5.—Austral.
subintritula, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 292 = macrantha.
subulata, Labill. Pugill. viii. 43 = Neesii.
tenella, *H. B. & K. Nov. Gen. et Sp.* v. 291.—N. Granat.
tenuifolia, Willd. Enum. Hort. Berol. 340 = filiformis.
tomentosa, *St. Hil. Pl. Rem. Brés.* 261.—Bras.
trinervia, Spreng. Neue Entdeck. i. 298 = paniculiflora.
umbellata, *Lour. Fl. Cochinch.* 136.—China.
uniflora, Raf. in Atl. Journ. 148 = Dionæa muscipula.
uniflora, *Willd. Enum. Hort. Berol.* 340.—Reg. Magellan.
verticillata, Linn. Epist. ed. Van Hall, 76 = Roridula dentata.
villosa, *St. Hil. Pl. Rem. Brés.* 267.—Bras.
violacea, *Willd. Enum. Hort. Berol.* 340.—Afr. austr.
Whittakeri, *Planch. in Ann. Sc. Nat.* Sér. III. ix. (1848) 302.—Austral.
sonaria, *Planch. l. c.* 303.—Austral.

DROSOCARPIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 388 = **Hyperiçum**, Tourn.
fimbriatum, Fourr. l. c. = H. Richeri.

DROSODENDRON, M. Roem. Syn. Hesper. 140 (1846) = **Bascikia**, Linn. (Myrtac.).
Rosmarinus, M. Roem. l. c. = B. cochinchinensis.

DROSOPHORUS, R. Br. ex Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 306 = **Byblis**, Salisb. (Droserac.).
cærulea, R. Br. ex Planch. l. c. = B. liniflora.

DROSOPHYLLUM, Link, in Schrad. Neues Journ. i. (1806) ii. 58. *DROSERACEÆ*, Benth. & Hook. f. i. 563.
Spergulus, 'Brot.' ex Steud. Nom. ed. II. ii. 618 (1841).
lusitanicum, *Link, i. c.*—Lusitan.; Marocc.

DROSSERA, Gled. in Mém. Acad. Berl. v. (1749) 126 = **Drosera**, Linn.

DROUGUETIA, Endl. Gen. Suppl. iv. 37 (1847) = **Drogustia**, Gaudich. (Urtic.).

DROZIA, Cass. in Dict. Sc. Nat. xxxiv. 217 (1825) = **Parœnia**, Lag. (Compos.).
dicephala, Cass. l. c. 218 = P. dicephala.
macrocephala, Steud. Nom. ed. II. i. 531 = P. pinnatifida.
monocephala, Cass. in Dict. Sc. Nat. xxxiv. (1825) 218 = P. pinnatifida.

DRUDEA, Griseb. in Goett. Abh. xxiv. (1879) 26. *CARYOPHYLLEÆ*.
lycopodioides, *Griseb. l. c.*—Peruv.

DRUMMONDIA, DC. Prod. iv. 49 (1830) = **Mitella**, Linn. (Saxifrag.).
mitelloides, DC. l. c. 50 = M. pentandra.

DRUMMONDITA, Harv. in Hook. Kew Journ. vii. (1855) 53. *RUTACEÆ*, Benth. & Hook. f. i. 394.
crioclides, *Harv. l. c.*—Austral.

DRUPARIA, [Clairv.] Man. Herb. 159 (1811) = **Prunus**, Tourn. (Rosac.).
Amygdalus, [Clairv.] l. c. 160 = P. Amygdalus.
Avium, [Clairv.] l. c. 159 = P. Avium.
Cerasus, [Clairv.] l. c. = P. Cerasus.
insititicia, [Clairv.] l. c. 160 = P. insititia.
Mahaleb, [Clairv.] l. c. 159 = P. Mahaleb.
Padus, [Clairv.] l. c. = P. Padus.
Prunus, [Clairv.] l. c. = P. domestica.
spinosa, [Clairv.] l. c. = P. spinosa.

DRUPARIA, Manso, Enum. 35 (1836) = **Cayaponia**, Manso (Cucurb.).
racemosa, Manso, l. c.—Bras.

DRUPATRIS, Lour. Fl. Cochinch. 214 (1790). *GEN. DUB.*—*Cl.* Benth. & Hook. f. ii. 668.
cochinchinensis, *Lour. l. c.*—Cochinch.

DRUPIFERA, Rafin. Sylva Tellur. 140 (1838) = **Camellia**, Linn. (Terastroem.).
oleosa, Rafin. l. c. = C. drupifera.

DRUPINA, Linn. Pl. Surin. 11 (1775); Amœn. Acad. viii. 329 (1785) = **Curvanga**, Juss. (Scrophular.).
cristata, Linn. fil. cc. (= *Picria surinamensis*).—Guiana.

DRUSA, DC. in Ann. Mus. Par. x. (1807) 466. t. 28 = **Bowlesia**, Ruiz & Pav. (Umbellif.).
oppositifolia, DC. l. c. = B. oppositifolia.

DRYADAEA, Linn. Syst. ed. I (1735) = **Dryas**, Linn. (Rosac.).

DRYADANTHE, Endl. Gen. 1242 (1840) = **Potentilla**, Linn. (Rosac.).
Bungeana, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 249 = P. tetranda.
pusilla, Walp. Rep. ii. 27 = P. tetranda.

DRYANDRA, R. Br. in Trans. Linn. Soc. x. (1810) 211. t. 8. *PROTEACEÆ*, Benth. & Hook. f. iii. 155.
Josephia, Salisb. in Knight, Prot. 110 (1809).
Hastulata, R. Br. Prod. Suppl. (Prot. Nov.) i. 40 (1830).
Arctotidis, *R. Br. Prod. Nov.* 99.—Austral.
armata, *R. Br. in Trans. Linn. Soc.* x. (1810) 213.—Austral.
Baxteri, *R. Br. Prod. Nov.* 98.—Austral.
bipinnatifida, *R. Br. l. c.* 98.—Austral.
blechnifolia, R. Br. in Trans. Linn. Soc. x. (1810) 215 = quercifolia.

DRYANDRA :—
Brownii, Meisn. in Lehm. Pl. Preiss. i. 595 = nivea.
calophylla, *R. Br. Prot. Nov.* 40.—Austral.
cardiacea, *Lindl. Swan River App.* 33.—Austral.
carlinoides, *Meisn. in Lehm. Pl. Preiss.* ii. 267.—Austral.
cirsioides, *Meisn. in DC. Prod.* xiv. 476.—Austral.
comosa, *Meisn. l. c.* 478.—Austral.
concinna, *R. Br. Prot. Nov.* 38.—Austral.
conferta, *Benth. Fl. Austral.* v. 578.—Austral.
cryptocephala, Meisn. in Lehm. Pl. Preiss. i. 596 = senecifolia.
cuneata, *R. Br. in Trans. Linn. Soc.* x. (1810) 212.—Austral.
Drummondii, Meisn. in Lehm. Pl. Preiss. ii. 267 = calophylla.
elegans, Meisn. in DC. Prod. xiv. 473 = tenuifolia.
falcata, *R. Br. in Trans. Linn. Soc.* x. (1810) 213.—Austral.
fecunda, Lindl. Swan River App. 33 = armata.
ferruginea, Kipp. ex Meisn. in Hook. Kew Journ. vii. (1855) 128 = proteoides.
floribunda, *R. Br. in Trans. Linn. Soc.* x. (1810) 212.—Austral.
foliolata, *R. Br. Prot. Nov.* 38.—Austral.
foliolata, Meisn. in Lehm. Pl. Preiss. ii. 266 = Kippistiana.
formosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 213. t. 3.—Austral.
Fraseri, *R. Br. Prot. Nov.* 39.—Austral.
Hewardiana, *Meisn. in DC. Prod.* xiv. 477.—Austral.
horrida, *Meisn. l. c.* 476.—Austral.
Kippistiana, *Meisn. l. c.* 473.—Austral.
Lindleyana,Meisn. in Lehm. Pl. Preiss. i. 598 = nivea.
longifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 215.—Austral.
mucronulata, *R. Br. l. c.* 213.—Austral.
multisecta, F. Muell. Fragm. v. 185 = obtusa
mutica, Meisn. in DC. Prod. xiv. 475 = foliolata.
nana, *Meisn. in Hook. Kew Journ.* vii. (1855) 121.—Austral.
nervosa, R. Br. in Sweet, Fl. Austral. t. 22 = pteridifolia.
nivea, *R. Br. in Trans. Linn. Soc.* x. (1810) 214.—Austral.
nobilis, *Lindl. Swan River App.* 33.—Austral.
obtusa, *R. Br. in Trans. Linn. Soc.* x. (1810) 214.—Austral.
patens, *Benth. Fl. Austral.* v. 578.—Austral.
plumosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 214.—Austral
polycephala, *Benth. Fl. Austral.* v. 570.—Austral.
praemorsa, *Meisn. in Lehm. Pl. Preiss.* i. 265.—Austral.
Preissii, *Meisn. l. c.* 599.—Austral.
proteoides, *Lindl. Swan River App.* 33.—Austral.
pteridifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 215.—Austral.
pulchella, *Meisn. in DC. Prod.* xiv. 473.—Austral.
quercifolia, *Meisn. l. c.* 467.—Austral.
runcinata, *Meisn. l. c.* 469.—Austral.
sclerophylla, *Meisn. in Hook. Kew Journ.* vii. (1855) 128.—Austral.
senecifolia, *R. Br. Prot. Nov.* 39.—Austral.
Serra, *R. Br. l. c.* 38.—Austral.
serratuloides, *Meisn. in Hook. Kew Journ.* vii. (1855) 128.—Austral.
Shuttleworthiana, *Meisn. l. c.* 192.—Austral.
speciosa, *Meisn. in DC. Prod.* xiv. 478.—Austral.
squarrosa, *R. Br. Prot. Nov.* 38.—Austral.
squarrosa, Meisn. in Lehm. Pl. Preiss. ii. 266 = polycephala.
stenoptera, Meisn. in Hook. Kew Journ. vii. (1855) 122 = nivea.
stuposa, *Lindl. Swan River App.* 33.—Austral.
tenuifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 215.—Austral.
tortifolia, Kipp. ex Meisn. in Hook. Kew Journ. vii. (1855) 121 = Arctotidis.
tridentata, *Meisn. in DC. Prod.* xiv. 479.—Austral.
uncata, A. Cunn. ex Benth. Fl. Austral. v. 561 = tenuifolia.
vestita, *Meisn. in Hook. Kew Journ.* vii. (1855) 121.—Austral.

DRYANDRA, Thumb. Nov. Gen. 60 (1788) = **Aleuritea**, Forst. (Euphorb.).
cordata, Thumb. l. c. = A. triloba.

DRYANDRA :—
oleifera, Lam. Encyc. ii. 329 = Aleurites triloba.
Verniciia, Correa, in Ann. Mus. Par. viii. (1822) 69 = A. triloba.

DRYAS, Linn. Gen. ed. I. 148 (1737). *ROSACEAE*, Benth. & Hook. f. i. 618.
DRYADAZA, Linn. Syst. ed. I (1735).
PTILOTUM, Dulac, Fl. Hautes-Pyr. 213 (1867).
alpina, Salisb. Prod. 383 = octopetala.
anemonoides, Pall. Reise, iii. App. 733 = Geum anemonoides.
chamaedrifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 578 = octopetala.
chamaedrifolia, Richards. in Frankl. Narr. 1st Journ. 740 = Drummondii.
chamaedryoides, Pall. Reise, iii. App. 733 = octopetala.
depressa, Bab. in Trans. Bot. Soc. Edinb. i. (1843) 195 = octopetala.
Drummondii, *Richards. in Hook. Bot. Mag.* t. 2972.—Slbir. or. ; Am. arct.
geoides, Pall. Reise, iii. App. 732 = Coluria geoides.
integrifolia, *Vahl, in Skrivt. Naturh. Selsk. Kjorb.* iv. 11. (1798) 171.—Groenl. ; Am. bor.
octopetala, *Linn. Sp. Pl.* 501.—Reg. bor. temp. et arct.
pentapetala, Linn. l. c. = Geum anemonoides.
pentapetala, Pall. Reise, ii. 569 = Coluria geoides.
pentaphyllaea, Hill, Veg. Syst. xvi. 80 = octopetala.
tenella, Pursh, Fl. Am. Sept. i. 350 = integrifolia.
trifoliata, Pall. ex Ser. in DC. Prod. ii. 555 = Daliabarda fragarioides.

DRYMARIA, Willd. ex Roem. & Schult. Syst. v. p xxxi (1819). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 152.
apetala, *Bartl. in Presl, Rel. Haenk.* ii. 7.—Chili.
arenarioides, *Willd. ex Roem. & Schult. Syst.* v. 406.—Mexic.
ciliaris, Hort.,Berol. ex C. A. Mey. in Ind. Sem. Hort. Petrop. ix. 71 = cordata.
ciliata, Hort. Berol. ex C. A. Mey. l. c. = praec.
cordata, A. Gray, Fl. Fendl 13 = glandulosa.
cordata, *Willd. ex Roem. & Schult. Syst.* v. 406.—Comm. ex Roxb. ctop.
crassifolia, *Benth. Bot. Voy. Sulph.* 16.—Mexic. ; Calif.
cubensis, *Regel. in Otto & Dietr. Allg. Gartens.* viii. (1840) 298.—Cuba.
diandra, Blume, Bijdr. 62 = cordata.
divaricata, *H. B. & K. Nov. Gen. et Sp.* vi. 24.—Peruv.
effusa, *A. Gray, Pl. Wright.* ii. 19.—Mexic.
extensa, Wall. Cat. n. 647 = cordata.
fascinata, *A. Gray, Bot. U. St. Expl. Exped.* i. 125.—Peruv.
Fendleri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 208.—N. Mexic.
filiformis, *Benth. Fl. Austral.* i. 162.—Austral.
frankenioides, H. B. & K. Nov. Gen. et Sp. vi. 21. t. 515 = arenarioides.
glabriuscula, *Bartl. in Presl, Rel. Haenk.* ii. 7.—Peruv.
glandulosa, *Bartl. l. c.* 9.—Mexic.
gracilis, *Cham. & Schlecht. in Linnaea*, v. (1830) 232.—Peruv.
grandiflora, *Bartl. in Presl, Rel. Haenk.* ii. 7.—Chili.
grandiflora, Schlecht. in Linnaea, xii. (1838) 205 = magellanica.
hirsuta, *Bartl. in Presl, Rel. Haenk* ii. 8.—Peruv.
holosteoides, *Benth. Bot. Voy. Sulph.* 16.—Calif. ; Mexic.
laxiflora, *Benth. Pl. Hartw.* 73.—Mexic.
leptoclados, *Hemsl. Diag. Pl. Nov.* 2.—Guatemala.
macrantha, *A. Gray, Bot. U. St. Expl. Exped.* i. 126.—Peruv.
magellanica, *Steud. Nom.* ed. II. i. 582.—Mexic.
moliuginea, Didr. in Linnaea, xxix. (1857–58) 788 = Arenaria moliuginea.
nitida, *Bali, in Journ. Linn. Soc.* xxii. (1885) 31.—Peruv.
nodosa, *Engelm. in A. Gray, Pl. Fendl.* 12.—Mexic.
ortegioides, *Griseb. Cat. Pl. Cub.* 21.—Cuba.
ovata, *Willd. ex Roem. & Schult. Syst.* v. 406.—Ecuador.
Palmeri, *Hemsl. Diag. Pl. Nov.* 22.—Mexic.
palustris, *Cham. & Schlecht. in Linnaea*, v. (1830) 232.—Mexic.

DRYMARIA :—
paposana, *Phil. Fl. Atac.* 10.—Chili.
pauciflora, *Bartl. in Presl, Rel. Haenk.* ii. 8.—Peruv.
polycarpoides, *A. Gray, Pl. Fendl.* 12.—Mexic.
ramosissima, *Schlecht. in Linnaea*, xii. (1838) 206.—Mexic.
retusa, Wight & Arn. Prod. 359 = cordata.
rotundifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 123.—Peruv.
sperguloides, *A. Gray, Pl. Fendl.* 11.—N. Mexic.
stellarioides, *Willd. ex Roem. & Schult. Syst.* v. 416.—Ecuador.
suffruticosa, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 208.—Mexic.
tenella, *A. Gray, Pl. Fendl.* 12.—N. Mexic.
villosa, *Cham. & Schlecht. in Linnaea*, v. (1830) 232.—Mexic.
viscidula, *A. Gray, Bot. U. St. Expl. Exped.* i. 124.—Peruv.
xerophylla, *A. Gray, Pl. Wright.* ii. 18.—N. Mexic.

DRYMEIA, Ehrh. Beitr. iv. 148 (1780) = **Carex**, Linn. (Cyperac.).

DRYMIPHILA, Juss. in Dict. Sc. Nat. iii. Suppl. 53 (1816) = **Drymophila**, R. Br. (Liliac.).

DRYMIS, Juss. Gen. 280 (1789) = **Drimys**, Forst. (Magnol.).

DRYMISPERMUM, Reichb. Nom. 65 (1841) = DRIMYSPERMUM, Reinw. = **Phaleria**, Jack (Thymel.).

DRYMOCALLIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 371 = **Potentilla**, Linn. (Rosac.).
rubricaulis, Fourr. l. c.—Gallia.

DRYMOCODON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 111 = **Campanula**, Tourn.
bononiense, Fourr l. c. = C. bononiensis.
latifolium, Fourr. l. c. = C. latifolia.
rapunculoides, Fourr. l. c. = C. rapunculoides.
Trachelium, Fourr. l. c. = C. Trachelium.

DRYMODA, Lindl. Sert. Orch. t. 8 C (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 506.
picta, *Lindl. l. c.*—Burma.

DRYMONACTES, Steud. Nom. ed. II. i. 532 (1840) sphalm. = seq.

DRYMONAETES, Ehrh. Beitr. iv. 147 (1789) = **Festuca**, Linn. (Gramin.).
giganteus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 187 = F. gigantea.

DRYMONIA, Mart. Nov. Gen. et Sp. iii. 57. t. 224 (1829). *GESNERACEAE*, Benth. & Hook. f. ii. 1007.
alloplectoides, *Hanst. in Linnaea*, xxxiv. (1865–66) 856.—Costa Rica.
bicolor, Mart. ex Lindl. Bot. Reg. xxiv. (1838) t. 4 = serrulata.
calcarata, *Mart. Nov. Gen. et Sp.* iii. 57. t. 224.—Bras.
candida, *Hanst. in Linnaea*, xxxiv. (1865–66) 859.—Peruv.
coerulea, Mart. ex Steud. Nom. ed. II. i. 532 = calcarata.
conchoclyx, *Hanst. in Linnaea*, xxxiv. (1865–66) 860.—Costa Rica.
cristata, *Mig. l. c.* xviii. (1844) 96.—Guiana.
delphinioides, *Seem. Bot. Voy. Herald*, 186.—N. Granat.
grandiflora, G. Don, Gen. Syst. iv. 658 = Besleria grandiflora.
Yacquini, G. Don, l. c. = serrulata.
longifolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 4. t. 204.—Peruv.
marmorata, *Hook. f. Bot. Mag.* t. 6763.—Guian. !
mollis, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 111.—Costa Rica.
mucronulosa, *Hanst. in Linnaea*, xxxiv. (1865–66) 853.—Guatemala.
ochroleuca, Poepp. & Endl. Nov. Gen. et Sp. iii. 5 = Episcia ochroleuca.

DRYMONIA:—
ovata, *Hanst. in Linnaea*, xxxiv. (1865–66) 355.—Costa Rica.
parviflora, *Hanst. l. c.* 352.—Costa Rica.
parvifolia, *Griseb. Fl. Brit. W. Ind.* 463.—Ind. occ.
punctata, Lindl. Bot. Reg. (1842) Misc. 63 = Episcia punctata.
sarmentosula, Lem. in Fl. des Serres, Sér. I. iv (1848) t. 368 = estriata.
serrulata, *Mart. Nov. Gen. et Sp.* iii. 59.—Ind. occ.
speciabilis, *Mart. l. c.*—Panama.
splendens, *Triana, ex Galeotti, in Journ. Hort. Prat. Sér.* II. i. (1857) 97. t. 9.—Reg. Andin.
Turrialvae, *Hanst. in Linnaea*, xxxiv. (1865–66) 359. —Costa Rica.
villosa, Hook. Bot. Mag. t. 4866 = Episcia villosa.
villosa, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 12 (= pracc. †).—Guian. †
Warscewicziana, *Hanst. in Linnaea*, xxxiv. (1865–66) 352.—Costa Rica.

DRYMOPHILA, R. Br. Prod. 292 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 769.
cyanocarpa, *R. Br. Prod.* 292.—Austral.
Moorei, *Baker, in Journ. Linn. Soc.* xiv. (1875) 571. —Austral.
pyrrhocarpa, F. Muell. Fragm. ix. 190 = Moorei.

DRYMOPHLOEUS, Zipp. in Alg. Konst. en Letterbode, i. (1829) 207. *PALMAE*, Benth. & Hook. f. iii. 892.
ambiguus, Becc. Malesia, i. (1877) 42 = Actinopholeus ambiguus (in Addend.).
angustifolius, Mart. Hist. Nat. Palm. iii. 314 = Coleospadix oninensis.
appendiculatus, *Miq. Palm. Arch. Ind Obs. Nov.* 24; *Becc. Malesia*, i. (1877) 45.—Malaya.
bifidus, *Becc. l. c.* 44.—N. Guinea.
ceramensis, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1868) n. v. 5.—Ins. Ceram.
ceramensis, Scheff. Ann. Jard. Buitenz. i. (1876) 121 = oliveaformis.
communis, *Miq. Palm. Arch. Ind. Obs. Nov.* 24 = Coleospadix oninensis.
disaricatus, Benth. & Hook. f. Gen. iii. 893 = Kentiopsis divaricata.
filiferus, Scheff. in Ann. Jard. Buitenz. i. (1876) 178 = Vitiphoenix filifera.
jaculatorius, *Mart. Hist. Nat. Palm.* iii. 314.—N. Guinea.
leprosus, *Becc. in Ann. Jard. Buitenz.* ii. (1885) 119. —N. Guinea.
Normanbyanus, Benth. & Hook. f. Gen. iii. 893 = Normanbya Muelleri.
oliveaformis, *Mart. Hist. Nat. Palm.* iii. 314.—Ins. Amboina.
oliveaformis, Scheff. in Ann. Jard. Buitenz. i. (1876) 157 = bifidus.
paradoxus, Scheff. l. c. 53 = Ptychococcus paradoxus.
propinquus, Becc. Malesia, i. (1877) 43 = Actinopholeus propinquus (in Addend.).
paniculus, *Becc. l. c.* 47 = Pinanga ternatensis.
Rumphianus, *Mart. Hist. Nat. Palm.* iii. 314 = Pinanga ternatensis.
Rumphii, Becc. in Ann. Jard. Buitenz. ii. (1885) 119 = leprosus.
Rumphii, Scheff. l. c. i. (1876) 52 = bifidus.
saxatilis, *Mart. Hist. Nat. Palm.* iii. 314.—Ins. Amboina.
singaporensis, Hook. f. in Kew Report, 1882 (1884) 55 = Ptychoraphis singaporensis.
westiarius, Miq. Palm. Arch. Ind. Obs. 24 = Mischophloeus paniculatus.
Zippelii, Blank. in Hoev. & De Vriese, Tijdschr. ix. (1842) 170 = Caryota mitis.

DRYMOPOGON, Rapp. Fl. Jen. ed. Hall. 124 (1745) = **Spiraea**, Linn. (Rosac.).

DRYMOSPARTUM, Presl, (Bot. Bemerk. 186) (1844) in Abh. Boehm. Ges. V. iii. 567 = **Genista**, Linn. (Legumin.).
aetnense, Presl, l. c. = G. aethnensis.
purgans, Presl, l. c. = G. purgans.
Sardum, Presl, l. c. = G. aethnensis.

DRYMOSPHACE, Opiz, Seznam, 38 (1852) = **Salvia**, Tourn. (Labiat.).
glutinosa, Opiz, l. c. = S. glutinosa.

DRYMYS, Vell. Fl. Flum. 240 (1825) = **Drimys**, Forst. (Magnol.).

DRYOBALANOPS, Gaertn. f. Fruct. iii. 49. t. 186 (1805). *DIPTEROCARPEAE*, Benth. & Hook. f. i. 191.
aromatica, *Gaertn. f. l. c.*—Archip. Ind.
Beccarii, *Dyer, in Journ. Bot.* xii. (1874) 100.—Borneo.
Camphora, Colebr. in As. Res. xii. (1816) 535 = aromatica.
oblongifolia, *Dyer, in Journ. Bot.* xii. (1874) 100.—Borneo.
Scheffleri, *Hance, in Journ. Bot.* xiv. (1876) 307.—Sumatra.
sericea, *Korth. Verh. Nat. Gesch. Bot.* 68.—Borneo.

DRYOPAEIA, Linnaea, ii. (1827) 529 = seq.

DRYOPEIA, Thou. Orch. Iles Afr. t. 1–8 (1822) = **Disperis**, Sw. (Orchid.).
discolor, Thou. l. c. = Disperis cordata.
oppositifolia, Thou. l. c. = Disperis oppositifolia.
tripetaloides, Thou. l. c. = Disperis tripetaloides.

DRYOPETALON, A. Gray, in Smithson. Contrib. v. (Pl. Wright. ii.) 11. t. 14 (1853). *CRUCIFERAE*, Benth. & Hook. f. i. 69.
runcinatum, *A. Gray, l. c.* 12.—N. Mexic.

DRYOPETALUM, Benth. & Hook. f. Gen. i. 69 (1862) = pracc.

DRYOPRIA, Thou. Orch. Iles Afr. Tabl. 1 (1822) = DRYOPEIA, Thou. = **Disperis**, Sw. (Orchid.).

DRYOPSILA, Rafin. Alsog. Am. 25 (1838) = **Quercus**, Tourn. (Cupul.).
aprica, Rafin. l. c. 26 = Q. aprica.
cinerea, Rafin. l. c. 27 = Q. cinerea.
cunesfolia, Rafin. l. c. = Q. cunesfolia, *Rafin.*
laurina, Rafin. l. c. = Q. laurina.
maritima, Rafin. l. c. = Q. nigra.
oligodes, Rafin. l. c. 28 = Q. oligodes.
verrucosa, Rafin. l. c. 26 = Q. verrucosa.
virens, Rafin. l. c. 28 = Q. virens.

DRYPETES, Vahl, Eclog. iii. 49 (1815). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 278.
FREIREODENDRON, Muell. Arg. in DC. Prod. xv. ii. 244 (1866).
LIPARENA, Poit. ex Leman, in Dict. Sc. Nat. xxvii. 6 (1823).
alba, *Poit. in Mém. Mus. Par.* i. (1815) 157. t. 7.—Ins. Testudin.
benguelensis, Spreng. Syst. iii. 902 = Fluggia microcarpa.
crocea, Poit. in Mém. Mus. Par. i. (1815) 159. t. 8 = Xylosma nitidum.
cyathophora, *Muell. Arg. in DC. Prod.* xv. ii. 454.—Bras.
glauca, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 157 = Xylosma nitidum.
lateriflora, *Eclog.* iii. 49.—Porto Rico.
glomerata, *Griseb. in Goett. Abh.* vii. (1857) 172.—Ind. occ.
incurva, *Muell. Arg. in Linnaea*, xxxii. (1863) 82.—Cuba.
mucronata, *Wright, ex Griseb. in Goett. Nachr.* (1865) 165.—Cuba.
sessiliflora, *Allem. Diss. Dry.* (1844).—Bras.
sessiliflora, Balll. Etud. Gén. Euph. Atlas, 45 = Xylosma nitidum.
Spruceana, *Muell. Arg. in DC. Prod.* xv. ii. 454.—Bras. triplinervia, *Muell. Arg. l. c.* 456.—Ind. occ.

DRYPIS, Mich. ex Linn. Syst. ed. I (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 145.
spinosa, *Linn. Sp. Pl.* 413.—Reg. Mediterr.

DRYPSIS, Dach. in Orb. Dict. ix. 424 (1849) = **Dypsis**, Noronha (Palm.).

DRYPTOPETALUM, Am. in Ann. Nat. Hist. i. (1838) 373 = **Gynostroches**, Blume (Rhinoph.).
coriaceum, Arn. l. c. 373 = G. axillaris.

DRYPTOPETALUM:—
membranaceum, Miq. ex Benth. in Journ. Linn. Soc iii. (1859) 78 = Blepharistemma corymbosum.

DUABANGA, Buch.-Ham. in Trans. Linn. Soc. xvii. (1835) 178. *LYTHRARIEAE*, Benth. & Hook. f. i. 788.
grandiflora, Walp. Rep. ii. 114 = sonneratioides.
moluccana, *Blume, Mus. Bot. Lugd. Bat.* i. 199.—Ins. Molucc.
sonneratioides, *Buch.-Ham. in Trans. Linn. Soc.* xvii (1885) 177.—Ind. or.

DUANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 = **Homalanthus**, A. Juss. (Euphorb.)

DUBAEA, Steud. Nom. ed. II. i. 517, in syn. (1840) = DUBYAEA, DC. = **Diplusodon**, Pohl (Lythrar.).

DUBAUTIA, Gaudich. in Freyc. Voy. Bot. 469 t. 84 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 393
lacvigata, *A. Gray, in Proc. Am. Acad.* v. (1861) 135. —Ins. Sandvic.
laxa, *Hook. & Arn. Bot. Beech. Voy.* 87.—Ins. Sandvic
paleata, *A. Gray, in Proc. Am. Acad.* v. (1861) 135 — Ins. Sandvic.
plantaginea, *Gaudich. in Freyc. Voy. Bot.* 468. t. 84.—Ins. Sandvic.

DUBOISIA, R. Br. Prod. 448 (1810). *SOLANACEAE*, Benth & Hook. f. ii. 911.
Hopwoodii, *F. Muell. Fragm.* viii. 282.—Austral.
Leichhardtii, *F. Muell. in Wing, South Sc. Record.* ii. (1882) 292.—Austral.
myoporoides, *R. Br. Prod.* 448.—Austral.

DUBOISIA, Karst. in Otto & Dietr. Allg. Gartenz. xv (1847) 394 = **Pleurothallis**, R. Br. (Orchid.).
Reymondi, Karst. l. c. = P. Raymondi.

DUBOIS-REYMONDIA, Karst. in Bot. Zeit. vi. (1848) 897 = **Pleurothallis**, R. Br. (Orchid.).
lancipetala, Karst. Fl. Columb. i. sub t. 47.—N. Granat.
palpigera, Karst. in Bot. Zeit. vi. (1848) 897, 898.—N Granat.

DUBOSCIA, Bocq. in Adansonia, vii. (1866) 50. *TILIACEAE*, Benth. & Hook. f. i. 265.
macrocarpa, *Bocq. l. c.*—Afr. trop.

DUBOUZETIA, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. viii. (1861) 199. *TILIACEAE*, Benth. & Hook. f. i. 240.
campanulata, *Panch. ex Brongn. & Gris, l. c.*—N Caled.
elegans, *Brongn. & Gris, l. c.* (1863) 476.—N. Caled.
parviflora, *Brongn. & Gris, l. c.*—N. Caled.

DUBREUILIA, Decne. in Nouv. Ann. Mus. Par. iii (1834) 489 = seq.

DUBRUEILIA, Gaudich. in Freyc. Voy. Bot. 495 (1826) = **Pilea**, Lindl. (Urtic.).
callitrichoides, Gaudich. l. c. = P. callitrichoides.
ciliaris, Gaudich. l. c. = P. ciliaris.
crassifolia, Gaudich. l. c. = P. densiflora.
cuneiformis, Gaudich. l. c. = P. cuneiformis.
grandifolia, Gaudich. l. c. = P. grandifolia.
hederacea, Gaudich. l. c. = P. hederacea.
inaequalis, Gaudich. l. c. = P. inaequalis.
lanceolata, Gaudich. l. c. = P. lanceolata.
lucens, Gaudich. l. c. = P. lucens.
melastomoides, Gaudich. l. c. = P. trinervis.
membranacea, Gaudich. l. c. 496 = Urtica membranacea.
microphylla, Gaudich. l. c. 495 = P. microphylla.
multiflora, Gaudich. l. c. = P. multiflora.
orbiculata, Gaudich. l. c. 496 (= *Urtica orbiculata*). —Hab.?
Parietaria, Gaudich. l. c. 495 = P. Parietaria.
pepleides, Gaudich. l. c. = P. pepleides.
pumila, Gaudich. l. c. = P. pumila.
rhombea, Gaudich. l. c. 496 = P. rhombea.
Riedlei, Decne. Herb. Timor. Descr. 161 = P. lucens.
rotundifolia, Gaudich. in Freyc. Voy. Bot. 496 = Pipturus rotundifolia.
rupipendis, Gaudich. l. c. 495 = P. rupipendis.
semidentata, Gaudich. l. c. = P. rupipendis.

DUBRUEILIA:—
serpillifolia, Gaudich. in Freyc. Voy. Bot. 495 = Piles serpillifolia.
sessilifolia, Gaudich. l. c. = P. sessilifolia.
trilobata, Gaudich. l. c. = P. trilobata.
triplinervia, Gaudich. l. c. 496 = P. lucens.
verbascifolia, Gaudich. l. c. 495 = P. verbascifolia.

DUBYAEA, DC. Prod. vii. 94(a), in syn. (1838) = **Diplusodon**, Pohl (Lythrar.).
malastomoides, DC. l. c. 94(a) = Dipl. speciosus.

DUBYAEA, DC. Prod. vii. 247 (1838) = **Lactuca**, Tourn. (Compos.).
crustulata, DC. l. c.—Austral.
hispida, DC. Coll. Mém. ix. t. 19 = L. Dubyaea.
ramosissima, Hance, ex Walp. Ann. ii. 1028 = L. denticulata.
sonchoidea, DC. Prod. vii. 247 = Taraxacum montanum.

DUCHARTREA, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 109. t. 8 = **Pentarhaphia**, Lindl (Gesnerac.)
viridiflora, Decne. l. c. = P. viridiflora.

DUCHASSAINGIA, Walp. in Linnæa, xxiii. (1850) 741 = **Brythrina**, Linn. (Legumin.).
arborescens, Walp. in Flora, xxxvi. (1853) 150 = E. arborescens
caffra, Walp. in Linnæa, xxiii. (1850) 742 = E. caffra.
glauca, Walp. l. c. = E. glauca.
ovalifolia, Walp. l. c. = E. ovalifolia.
remipinata, Walp. in Flora, xxxvi. (1853) 150 = E. respinata.

DUCHEKIA, Kostel. Allg. Med.-Pharm. Fl. i. 213 (1831) = **Pallasia**, Reichb. (Commelin.).
hirsuta, Kostel. l. c. 214 = P. ambigua?

DUCHESNEA, Sm. in Trans. Linn. Soc. x. (1811) 372 = **Fragaria**, Linn. (Rosac.).
chrysantha, Miq. Fl. Ind. Bat. i. 1. 372 = F. indica.
fragaroides, Sm. in Trans. Linn. Soc. x. (1811) 373 = F. indica.
fragiformis, D. Don, Prod. Fl. Nep. 233 = F. indica.
sundaica, Miq. Fl. Ind. Bat. i. 1. 372 = Potentilla Kleiniana.

DUCHESNIA, Cass. in Bull. Soc. Philom. (1817) 153 = **Pulchoria**, Gaertn. (Compos.).
crispa, Cass. in Dict. Sc. Nat. xiii. 546 = P. crispa.

DUCHOLA, Adans. Fam. ii. 357 (1763) = **Omphalea**, Linn. (Euphorb.).

DUCOUDRAEA, Bur. Monog. Bign. 49 (1864) = **Tecoma**, Juss. (Bignon.).
capensis, Bur. l. c. = T. capensis.

DUCROISIA, Endl. Gen. Suppl. v. 10 (1850) = seq.

DUCROSIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 341. *UMBELLIFERAE*, Benth. & Hook. f. i. 923.
anethifolia, *Boiss. l. c.*—Persia.
flabellifolia, *Boiss. l. c.*—Assyria.
Ismelia, *Aschers. in Verh. Bot. Ver. Brand.* xxi. 1879 (1880) 67.—Aegypt.
Olivieri, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 101 = anethifolia.

DUFOUREA, Bory, ex Willd. Sp. Pl. v. 55 (1810) = **Tristicha**, Thou. (Podostem.).
alternifolia, Willd. in Ges. Naturf. Fr. Berl. Mag. vi. (1813) 64; Roem. & Schult. Syst. Veg. i. 50 = T. alternifolia.
Bory, A. Rich. in Dict. Hist. Nat. v. 686 = T. trifaria.
hypnoides, St. Hil. in Mém. Mus. Par. x (1823) 473 = T. hypnoides.
Thouarsii, A. Rich. in Dict. Hist. Nat. v. 686 = T. alternifolia.
trifaria, Bory, ex Willd. Sp. Pl. v. 55 = T. trifaria.

DUFOUREA, Gren. in Act. Soc. Linn. Bord. ix. (1837) 26 = **Azumaria**, Rupp. (Caryophyll.).
crassifolia, Gren. l. c. = A. crassifolia.
purpurascens, Gren. l. c. = A. purpurascens.

DUFOUREA, H. B. & K. Nov. Gen. et Sp. iii. 118. t. 214 (1818) = **Breweria**, R. Br. (Convolvul.).
amazonica, Mart. ex Choisy, in DC. Prod. ix. 487.—Bras.
aurea, Splitg. ex De Vriese, in Nederl. Kruidk. Arch. i. (1848) 150 = *sericea*.
glabra, *H. B. & K. Nov. Gen. et Sp.* iii. 114.—N. Granat.
heterantha, Nees & Mart. in Nov. Act. Nat. Cur. xi. 1. (1823) 70 (= *Aniseia heterantha*).—Bras.
sericea, H. B. & K. Nov. Gen. et Sp iii. 114. t. 214.—N. Granat.

DUFRENOYA, Chatin, in Compt.-Rend. Par. li. (1860) 657 = **Henslowia**, Blume (Santalac.).
heterantha, Chatin, l. c. = H. heterantha.

DUFRESNEA, Meisn. Gen. 173 (1838) = seq.

DUFRESNIA, DC. Coll. Mém. viii. t. 3 (1829); et Prod. iv. 624 (1830) = **Valerianella**, Moench.
leiocarpa, Koch, in Linnæa, xvii. (1843) 34 = V. Dufresnia.
orientalis, DC. Coll. Mém. viii. t. 3 = V. Dufresnia.

DUGAGELIA, Gaudich. in Freyc. Voy. Bot. 513 (1826). 'Delestur,' Benth. & Hook. f. iii. 137.

DUGALDIA, Cass. in Dict. Sc. Nat. lv. 270 (1828) = **Helenium**, Linn. (Compos.).
chrysanthemoides, Cass. l. c. = H. integrifolia.
integrifolia, Cass. l. c. = H. ovalifolia.

DUGESIA, A. Gray, in Proc. Am. Acad. xvii. (1881-82) 215. *COMPOSITAE*
mexicana, *A. Gray, l. c.* 216.—Mexic.

DUGEZIA, Montr. in Mém. Acad. Lyon, x. (1860) 187. *GENUS DUBIUM*, Benth. & Hook. f. i. 979.
artensis, *Montr. l. c.*—N. Caled.

DUGGENA, Vahl, in West, Bidr. Ste-Croix, 269 (1793) = **Gonzales**, Pers. (1805) (Rubiac.).
Richardii, Vahl, l. c. = G. spicata.

DUGLASSIA, Houst. Reliq. 13 (1781) = **Clerodendron**, Linn. (Verbenac.).

DUGORTIA, Scop. Introd. 217 (1777) = **Parinarium**, Juss. (Rosac.).
nucamentosa, J. F. Gmel. Syst. 816.—Guiana.

DUGUETIA, A. St. Hil. Fl. Bras. Mer. i. 35. t. 7 (1825). *ANONACEAE*, Benth. & Hook. f. i. 23.
ABREMOA, Aubl. Pl. Gui. t. 245 (1775), nomen prius.
CARDIOPETALUM, Schlecht. in Linnæa, ix. (1834) 228.
bracteosa, *Mart. Fl. Bras.* xiii. t. 23.—Bras.
Candollei, *Baill. Adansonia*, viii. (1868) 327.—Bras.
coriacea, *Sond. in Linnæa*, xxii. (1849) 557.—Bras.
dicholepidota, *Mart. Fl. Bras.* xiii. t. 22.—Bras
Gardneriana, *Mart. l. c.*—Bras.
inconspicua, *Saget, in Ann. Sc. Nat.* Sér. VI. xi. (1881) 185.—Guiana.
lanceolata, *St. Hil. Fl. Bras. Mer.* i. 35.—Bras.
longisepala, *Benth. in Hook. Kew Journ.* v. (1853) 9.—Bras.
longifolia, *Baill. Adansonia*, viii. (1867-68) 327.—Guiana; Peruv.
Margraviana, *Mart. Fl. Bras.* xiii. t. 25.—Bras.
Moricandiana, *Mart. l. c.* 22.—Bras.
Pohliana, *Mart. l. c.* 24.—Bras.
quitarensis, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 261.—Guiana.
Schlechtendaliana, *Mart. Fl. Bras.* xiii. t. 24.—Bras.
Spixiana, *Mart. l. c.* 23.—Bras.
uniflora, *Mart. l. c.* 24.—Bras.

DUGULDEA, Meisn. Gen. Comm. 122 in syn. (1864) = DUGALDIA, Cass. = **Helenium**, Linn. (Compos.).

DUHALDEA, DC. Prod. v. 366 (1836) = **Inula**, Linn. (Compos.).
chinensis, DC. l. c. = I. Cappa.
eriophora, Steets, in Bonplandia, v. (1857) 308 = I. Cappa.
eupatorioides, Steets, l. c. = I. eupatorioides.
macrophylla, Steets, l. c. = I. macrophylla.

DUHALDIA, Sch.-Bip. in Walp. Rep. ii. 952 (1843) = prec.

DUHAMELA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 82 = seq.

DUHAMELIA, Domb. ex Lam. Encyc. i. 245 (1783) = **Myrsine**, Linn.
Mangillo, Domb. ex Lam. l. c. = M. Mangilila.

DUHAMELIA, Pers. Syn. i. 208 (1805) = **Hamelia**, Jacq (Rubiac.).
chrysantha, Pers. l. c. = H. chrysantha.
glabra, Pers. l. c = Amaioua guianensis.
odorata, Roem. & Schult. Syst. v. 267 = H. suaveolens.
patens, Pers. Syn. i. 208 = H. patens.
sphærocarpa, Pers. l. c. = H. sphærocarpa.
ventricosa, Pers. l. c. = H. ventricosa.

DULACIA. Neck. Elem. ii. 414 (1790) = **Cousinia**, Aubl. (Rosac.).

DULACIA, Vell. Fl. Flum. 32 (1825); i. t. 78 = **Liriosma**, Poepp. (Olacin.).
singularis, Vell. ll. cc. = L. Velloziana.

DULCAMARA, Moench, Meth. 514 (1794) = **Solanum**, Tourn.
flexuosa, Moench, l. c. = S. Dulcamara.
lignosa, Gillib. Fl. Lituan. i. 37 = S. Dulcamara.

DULIA, Adans. Fam. ii. 165 (1763) = **Ledum**, Linn (Ericac.).

DULICHIUM, Pers. Syn. i. 65 (1805). *CYPERACEAE*, Benth. & Hook. f. iii. 1046.
PLEURANTHUS, Rich. ex Pers. Syn. i. 65 (1805).
canadense, Rich ex Pers. Syn. i. 65 = spatheacum. spatheaceum, *Rich. l. c.*—Am. bor.

DULONGIA, H. B. & K. Nov. Gen. et Sp. vii. 76. t. 623 (1825) = **Phyllonoma**, Willd. (Saxifrag.).
acuminata, H. B. & K. l. c. = P. ruscifolia.
integerrima, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 454 = P. ruscifolia.
laticuspis, Turcz. l. c. = P. ruscifolia.

DUMARTROVA, Gaudich. Bot. Voy. Bonite, t. 97 (1844-66?) = **Malaisia**, Blanco (Urtic.).
fagifolia, Gaudich. l. c. = M. tortuosa.

DUMASIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 96. *LEGUMINOSAE*, Benth. & Hook. f. i. 629.
capensis, *Eckl. & Zeyh. Enum.* 245.—Afr. austr.
congesta, R. Grah. in Wall. Cat. n. 5524 = villosa.
cordifolia, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 188.—Ind. or.
glaucescens, *Miq. Fl. Ind. Bat.* i. 1. 227.—Java.
leiocarpa, Benth. in Miq. Pl. Jungh. 231 = villosa.
pubescens, DC. Mém. Lég. 257. t. 45 = villosa.
truncata, *Sieb. & Zucc. in Abh. Akad. Muench*, iv. ii. (1843) 119.—Japon.
villosa, *DC. Mém. Lég.* 257. t. 44.—Ind. or.

DUMERILIA, Lag. ex DC. in Ann. Mus. Par. xix. (1812) 71. t. 6 (15) et 7 (16) = **Jungia**, Linn. f. (Compos.).
auriculata, Steud. Nom. ed. I. 287 = J. auriculata.
axillaris, Lag. ex DC. in Ann. Mus. Par. xix. (1812) 72 = J. axillaris.
crenata, Steud. Nom. ed. I. 287 = J. crenata.
paniculata, Cass. Opusc. i. t. 12 = J. spectabilis.
paniculata, DC. in Ann. Mus. Par. xix. (1812) 72 = J. ferruginea.
paniculata, H. B. & K. Nov. Gen. et Sp. iv. 156 = J. rugosa.
pubescens, Steud. Nom. ed. I. 287 = J. floribunda.
sericea, Steud. l. c. = J. sericea.

DUMERILIA, Less. in Linnæa, v. (1830) 13 = **Perezia**, Lag. (Compos.).
Alamani, DC. Prod. vii. 67 = P. Alamani.
Humboldtii, Less. in Linnæa, v. (1830) 13 = P. Humboldtii.

DUMREICHERA, Hochst. & Steud. in Flora, xxi. (1838) 1. intell. 26 = **Senra**, Cav. (Malvac.).
arabica, Hochst. & Steud. l. c.—S. incana.

DUNALIA, R. Br in Salt, Abyss. Suppl. p. lxiv (1814) = **Torenia**, Linn. (Scrophul.).
scandis, R. Br. l. c.—Abyssin.

DUNALIA, H. B. & K. Nov. Gen. et Sp. iii. 55. t. 194 (1818). *SOLANACEAE*, Benth. & Hook. f. ii. 895.
DIERBACHIA, Spreng. Syst. i. 613. 676 (1825).
acuticalyx, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 338.—Peruv.
asperrima, Dun. in DC. Prod. xiii. i. 527 = Iochroma fuchsioides.
brachycantha, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 339.—Bolivia.
brachystemon, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1861) *App.* 6.—Chili.
cornifolia, Dun. in DC. Prod. xiii. i. 527 = Iochroma cornifolia.
cyanea, *Rowe. ex Dun. l. c.* 484.—Hab. ?
grandiflora, Dun. l. c. 527 = Iochroma grandiflora.
illudens, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 32.—Peruv.
loxensis, Dun. in DC. Prod. xiii. i. 527 = Iochroma loxensis.
lycioides, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 339.—Am. austr.
ramiflora, *Miers, l. c.* 340.—Mexic.
scabicosa, *Miers, l. c.*—Bolivia.
solanacea, *H. B. & K. Nov. Gen. et Sp.* iii. 55. t. 194. —N. Granat.
spathulata, *A. Br. & Ascherson in Ind. Sem. Hort. Berol.* (1861) *App.* 6.—Hab. ?
umbrosa, Dun. in DC. Prod. xiii. i. 528 = Iochroma umbrosa.

DUNALIA, Montr. in Act. Soc. Linn. Bord. xxvi. (1866) 576 = **Amorphophallus**, Blume (Aroid.).
arrhenia, Montr. l. c.—N. Caled.

DUNALIA, Spreng. Pugill. ii. 25 (1815) = **Lucya**, DC. (Rubiac.).
tuberosa, Spreng. l. c. = L. tuberosa.

DUNANTIA, DC. Prod. v. 626 (1836) = **Isocarpha**, R. Br. (Compos.).
Achyranthes, DC. l. c. 627; et in Deless. Ic. Sel. iv. t. 73 = I. oppositifolia.
microcephala, DC. Prod. v. 627.—Patagon.

DUNBARIA, Wight & Arn. Prod. 258 (1834). *LEGUMINOSAE*, Benth & Hook f. i. 541.
acuminatissima, *Miq. Fl. Ind. Bat.* i. i. 180.—Java.
barbata, Benth. in Miq. Fl. Jungh. 243 = Atylosia barbata.
calycina, Miq Fl. Ind. Bat. i. i. 180 = Atylosia barbata.
circinalis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 219.—Burma.
conserpens, *Benth. in Miq. Pl. Jungh.* 241.—As. trop. Cumingiana, *Benth. l. c.* 242.—Ins. Philipp.
debilis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 218.—Reg. Himal.
ferruginea, *Wight & Arn. Prod.* 258.—Ind. or.
fusca, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 186 = Phaseolus fuscus.
Heynei, Span. Ic. n. 9; ex Miq. Fl. Ind. Bat. i. i. 179 = Spanoghel.
Heynei, *Wight & Arn. Prod.* 258.—Ind. or.
Horsfieldii, Miq. Fl. Ind. Bat. i. i. 179 = Atylosia mollis.
latifolia, Wight & Arn. Prod. 258 = ferruginea.
nivea, *Miq. Fl. Ind. Bat.* i. i. 177.—Java.
oblonga, Wight ex Arn. in Nov. Act. Nat. Cur. xviii. (1836) 338 = Heynei.
podocarpa, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 186.—Burma.
polyperma, *Miq. Fl. Ind. Bat.* i. i. 179.—Java.
pulchra, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 218.—Reg. Himal.
punctata, *Benth. in Miq. Pl. Jungh.* 242.—Java.
rubella, *Span. ex Miq. Fl. Ind. Bat.* i. i. 178.—Java. Spanoghel, *Miq. l. c.*—Java.

DUNCANIA, Reichb. Consp. 197 (1828) = **Toddalia**, Juss. (Rutac.).
undulata, Reichb. ex Steud. Nom. ed. II. i. 533 = T. undulata.

DUPARQUETIA, Baill. in Adansonia, vi. (1865–66) 189. t. 4 = **Oligostemon**, Benth. (Gesnerac.).
orchidacea, Baill. l. c. 190 = O. pictus.

DUPATYA, Vell. Fl. Flum. 85 (1825) = **Paepalanthus**, Mart. (Eriocaul.).
aequalis, Vell. l. c. 86; i. t. 85 = P. blepharocnemis.
hirsuta, Vell. l. c.; i. t. 87 = P. Dupatya.
lignlata, Vell. l. c.; i. t. 86 = Eriocaulon Kunthii.

DUPERREYA, Gaudich. in Freyc. Voy. Bot. 452. t. 63 (1826) = **Forana**, Burm. (Convolv.).
sericea, Gaudich. l. c. = P. sericea.

DUPINETA, Rafin. Sylva Tellur. 101 (1838) = **Dissotis**, Benth. (1849) (Melast.).
multiflora, Rafin. l. c. = Dis. multiflora.

DUPINIA, Scop. Introd. 195 (1777) = **Ternstroemia**, Linn. f. (1781).

DUPONTIA, R. Br. in Parry, Voy. App. 290 (1824) = **Graphaphorum**, Desv. (Gramin.).
Fischeri, R. Br. l. c. 191 = G. psilosanthus.
peligrera, Rupr. ex Nym. Consp. 808 = G. psilosanthus.
psilosantha, Rupr. Fl. Samoj. Cisural. (Beitr. Russ. Reich. ii.) t. 6 = G. psilosanthus.
sclerociada, Rupr. l. c. t. 6 = G. psilosanthus ?

DUPRATZIA, Rafin. Fl. Ludov. 54 (1817) = **Phlox**, Linn. (Polemoniac.).
scoparia, Rafin. l. c.—Louisian.

DUPUISIA, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 148. t. 38 (1832) = **Sorindeia**, Thou. (Anacard.).
juglandifolia, A. Rich. l. c. = S. juglandifolia.
longifolia, Hook. f. in Hook. Niger Fl. 387 = S. macrophylla.

DUQUETIA, G. Don, Gen. Syst. i. 97 (1831) = **Duguetia**, A. St. Hil. (Anonac.).

DURANDEA, Delarbr. Fl. Auv. ed. II. 365 (1800) = **Raphanus**, Tourn. (Crucif.).
umbellularis, Delarbr. l. c. = R. Raphanistrum.

DURANDEA, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 594. *LINEAE*, Benth. & Hook. f. i. 245.
serrata, *Planch. l. c.* vii. (1848) 628.—N. Caled.

DURANDOA, Pomel, Nouv. Mat. Fl. Atl. 2 (1860) = **Carthamus**, Tourn. (Compos.).
arborescens, Pomel, l. c. 22 = C. arborescens.
Clausonis, Pomel, l. c. 3 = C. arborescens.

DURANTA, Linn. Gen. ed. I. 867 (1737). *VERBENACEAE*, Benth. & Hook. f. ii. 1150.
ELLISIA, P. Br. Hist. Jamaic. 203 (1756).
HOFFMANNIA, Loefl. It. 194 (1758).
angustifolia, Salisb. Prod. 108 = Plumieri.
argentea, *Ludd. ex G. Don, in Lond. Hort. Brit.* 248. Ind. or.
brachypoda, *Tod. Nuov. Gen. e Sp.* 24.—Hab. ?
buxifolia, Herb. Willd. ex Schau. in DC. Prod. xi. 610 = Citharexylum spinosum.
buxifolia, Poir. ex Schau. in DC. l. c. 616 = triacantha.
dentata, Rich. Herb. ex Pers. Syn. ii. 142 = Plumieri.
Ellisia, Linn. Amœn. Fl. Carib. 36 = Plumieri.
erecta, Linn Sp. Pl. 637 = Plumieri.
Fischeri, Mart. in Flora, xxi. (1838) ii. Beibl. 60 = Cassela integrifolia.
illicifolia, Herb. Willd. ex Schau. in DC. Prod. xi. 609 = Citharexylum ilicifolium.
inermis, Linn. Sp. Pl. 637 = Plumieri.
integrifolia, *Tod. Nuov. Gen. e Sp.* 27.—Hab. ?
latifolia, Salisb. Prod. 108 = Plumieri.
Lorentzii, *Griseb. in Gottt. Abh.* xxiv. (1879) 280.—Reg. Argent.
mancorgoba, *H. B. & K. Nov. Gen. et Sp.* ii. 255.—Mexic.
microphylla, Willd. Enum. Hort. Berol. Suppl. 43 = Plumieri.
Mutisiana, Sm. Introd. Bot. 381 = seq.
Mutisii, *Linn. f. Suppl.* 291.—Am. trop.
obtusifolia, H. B. & K. Nov. Gen. et Sp. ii. 254 = Mutisii.
parietariaefolia, Juss. ex Walp. Rep. iv. 78 = Mutisii.
parviflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 210.—Cuba.
phytolaccaefolia, Juss. Herb. ex Schau. in DC. Prod. xi. 616 = Mutisii.
Plumieri, *Fenz. Select. Am.* 166. t. 176. f. 76.—Am. trop.
racemosa, Mill. Gard. Dict. ed. VIII. n. 2 = Plumieri.
repens, Linn. Sp. Pl. 637 = Plumieri.
spinosa, Mill. Gard. Dict. ed. VIII. n. 3 = Plumieri.
stenostachya, *Tod. Nuov. Gen. e Sp.* 26.—Hab. ?
triacantha, *Juss. in Ann. Mus. Par.* vii. (1806) 77.—Peruv.

DURANTA:—
turbinata, *Tod. Nuov. Gen. e Sp.* 28.—Hab. ?
umbilicata, Miers, Trav. Chil. ii. 580 = Rhaphithamnus cyanocarpus.
vestita, Cham. in Linnæa, vii. (1832) 115 = Plumieri.
xalapensis, H. B. & K. Nov. Gen. et Sp. ii. 254 = Plumieri.

DURETIA, Gaudich. in Freyc. Voy. Bot. 500 (1826) = **Ecehmaria**, Jacq. (Urticac.).
cylindrica, Gaudich. l. c. = B. cylindrica.
palustris, Gaudich. l. c. = B. cylindrica.

DURIA, Scop. Introd. 289 (1777) = **Durio**, Adans. (Malvac.).

DURIEUA, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 14 (1842) = **Daucus**, Tourn. (Umbellif.).
abyssinica, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 51 = Daucus abyssinicus.
graeca, Boiss. l. c. 50 = Daucus creticus.
hispanica, Boiss. & Reut. Diagn. Pl. Nov. 15 = Daucus Durieua.
juncea, Willk. Sert. Fl. Hisp. 57 = Daucus setifolius.

DURIEURA, Mérat, Dis. in Mém. Soc. Acad. Lille, (1829) 432. t. 4 = **Lasuentaea**, Lag. (Scrophul.).
spicata, Mérat, l. c. = L. rotundifolia.

DURIO, Adans. Fam. ii. 399 (1763); Murr. Syst. ed. XIII. 581. *MALVACEAE*, Benth. & Hook. f. i. 213.
carinatus, *Mast. in Journ. Linn. Soc.* xiv. (1875) 500. —Borneo.
ceylanicus, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 1 = Cullenia excelsa.
lanceolatus, *Mast. in Journ. Linn. Soc.* xiv. (1875) 499.—Borneo.
lissocarpus, *Mast. l. c.* 501.—Borneo.
malaccensis, *Planch. ex Mast. in Hook. f. Fl. Brit Ind.* i. 351.—Burma; Malacca.
oblongus, *Mast. in Journ. Linn. Soc.* xiv. (1875) 500 —Borneo.
Oxleyanus, *Griff in Calc. Journ. Nat. Hist.* v. (1845) 115, in note.—Malacca.
stercoraceus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 18 = zibethinus.
zeylanicus, Auct. = *ceylanicus*.
zibethinus, Moon, Cat. Pl. Ceyl. 56 = Cullenia excelsa. zibethinus, *Murr. Syst.* ed. XIII. 581.—Malaya.

DUROIA, Linn. f Suppl. 30, 209 (1781). *RUBIACEAE*, Benth. & Hook. f. ii. 82.
PUSETA, Linn. Diss. Fl. Surin. 16 (1775).
eriophila, *Linn. f. l. c.* 209.—Guiana.
saccifera, *Benth. & Hook. f. Gen.* ii. 82.—Bras.
surinamensis, *Benth. & Hook. f. l. c.*—Guiana.

DUSCHEKIA, Opiz, Seznam, 38 (1852) = **Alnus**, Tourn. (Cupulif.).
ovata, Opiz, l. c. = A. viridis, *DC.*

DUTAILLYEA, Baill. Adansonia, x. (1872) 327. *RUTACEAE*.
trifoliolata, *Baill. l. c.* 328.—N. Caled.

DUTRA, Bernh. ex Steud. Nom. ed. II. i. 484, in syn. (1840) = **Datura**, Linn. (Solanac.).

DUTTONIA, F. Muell. in Hook. Kew Journ. viii. (1856) 74. t. 1 = **Pholidia**, R. Br. (Myopor.).
gibbifolia, F. Muell. l. c. = F. gibbifolia.

DUTTONIA, F. Muell. in Linnæa, xxv. (1852) 409 = **Heliperma**, DC. (Connarac.).
sessiliceps, F. Muell. l. c. 410 = H. Dimorpholepis.

DUVALIA, Bonpl. Descr. Pl. Rar. Malm. 180. t. 53 (1818) = **Hypocalyptus**, Thunb. (Legumin.).
oxalidifolia, Bonpl. l. c. = H. obcordatus.

DUVALIA, Haw. Syn. Pl. Succ. 44 (1812). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 784.
angustiloba, *N. E. Br. in Gard. Chron.* (1888) ii. 280. —Afr. austr.
caespitosa, *Hew. Syn. Pl. Succ.* 45.—Afr. austr.
compacta, *Haw. l. c.* 46.—Afr. austr.
Corderoyi, *N. E. Br. in Bot. Mag.* sub t. 6245.—Hab. ?

DUVALIA:—
deflexa, G. Don, Gen. Syst. iv. 121 = Stapelia deflexa.
elegans, *Haw. Syn. Pl. Succ.* 44.—Afr. austr.
glomerata, *Haw. l. c.* 46; *Suppl.* 14.—Afr. austr.
hirtella, *Sweet, Hort. Brit.* ed. I. 276.—Hab. ?
jacquiniana, *Sweet, l. c.*—Afr. austr.
laevigata, *Haw. Syn. Pl. Succ.* 46; *Suppl.* 14.—Afr. austr.
mastodes, *Sweet, Hort. Brit.* ed. I. 276, nomen; *G. Don, Gen. Syst.* iv. 122.—Afr. austr.
polita, *N. E. Br. in Gard. Chron.* (1878) ii. 180.—Afr. austr.
radiata, *Haw. Syn. Pl. Succ.* 45.—Afr. austr.
reclinata, *Haw. l. c.* 44.—Afr. austr.
replicata, *Sweet, Hort. Brit.* ed. I. 276.—Afr. austr.
tuberculata, Haw. Syn. Pl. Succ. 46; Suppl. 18 = radiata.

DUVALLIA, Haw. Pl. Succ. Suppl. 18 (1819) = prsec.

DUVAUA, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 340 = **Schinus**, Linn. (Anacard.).
crenata, Phil. Fl. Atac. 14 = S. crenatus.
cuneata, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 176 = S. dependens.
dentata, DC. Prod. ii. 74 = S. dependens.
dependens, DC. l. c. = S. dependens.
fasciculata, Griseb. in Goett. Abh. xix. (1874) 116 = S. dependens.
inebrians, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 176 = S. dependens.
latifolia, Gill. ex Lindl. Bot. Reg. t. 1580 = S. latifolius.
longifolia, Lindl. l. c. (1843) t. 59 = S. dependens.
molle, Bert. ex Steud. Nom. ed. II. i. 554 = S. latifolius.
ornata, Phil. f. Cat. Pl. Vasc. Chil. 47, spbalm. = seq.
ovata, Lindl. Bot. Reg. t. 1568 = S. dependens.
pleuropogon, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 467 = Mauria sumplicifolia *et* Lithraea caustica.
polygama, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 340 = S. dependens.
praecox, Griseb. in Goett. Abh. xix. (1874) 116 = S. dependens.
sinuata, Griseb. l. c. xxiv. (1879) 93 = S. sinuatus.
spinescens, Tenore, Cat. Ort. Nap. (1845) 85 = S. dependens.
urticina, Pasq. Cat. Ort. Bot. Nap. 40.—Hab. ?
velutina, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 467 = S. latifolius.

DUVAUCELLIA, Bowd. Mad. 259 (1825). *OLEACEAE*!
tennis, *Bowd. l. c.*—Afr. trop. ?

DUVERNAYA, Desv. ex DC. Prod. iii. 86, in syn. (1828) = **Cuphea**, Jacq. (Lythrar.).

DUVERNOIA, Nees, in DC. Prod. xi. 322 (1847) = seq.

DUVERNOYA, E. Mey. in Drège, Zwei Pfl. Docum. 150 (1843), nomen = **Adhatoda**, Nees (Acanth.).
adhatodoides, E. Mey. l. c.—Afr. austr.

DUVOA, Hook. & Arn. Bot. Beech. Voy. 55 (1832) = DUVAUA, Kunth = **Schinus**, Linn. (Anacard.).

DYANTHUS, P. Br. Hist. Jamaic. 228 (1756) err. typ. = **Dianthus**, Linn. (Caryophyll.).

DYCHOTRIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 85 = **Psychotria**, Linn. (Rubiac.).

DYCKIA, Schult. f. *Syst.* vii. 65, 1194 (1830). *BROMELIACEAE*, Benth. & Hook. f. iii. 667.
ENCHOLIRIUM, Mart. ex Schult. f. Syst. vii. 68, 1233 (1830).
GARRELLIA, Gaudich. Bot. Voy. Bonite, t. 115 (‘1844–46’).
NAVIA, Schult. f. Syst. vii. 65, 1195 (1830).
PRIONOPETALUM, C. Koch, Ind. Sem. Hort. Berol. (1873), App. iv. 3.
altissima, *Lindl. Bot. Reg.* (1841) *Misc.* 84.—Bras.
argentea, *Nichols. Dict. Gard.* i. 497.—Hab. ?
brevifolia, *Hort. ex Baker, in Saunds. Refug. Bot.* t. 286.—Bras.
catharinensis, *C. Koch, Ind. Sem. Hort. Berol.* (1873) *App.* iv. 4.—Bras.
densiflora, *Schult. f. Syst.* vii. 1194.—Bras.
distichiflora, *Schult. f. l. c.*—Bras.
floribunda, *Griseb. in Goett. Abh.* xxiv. (1879) 331.—Reg. Argent.
frigida, *Hook. f. Bot. Mag.* t. 6294.—Bras.
leptostachya, *Baker, in Gard. Chron.* (1884) ii. 198.—Bras.
montevidensis, C. Koch, Ind. Sem. Hort. Berol. (1873) App. iv. 4 = rariflora.
Princeps, Lem. Jard. Fl. ill. (1852–53) t. 224, 225, ex pte = altissima.
rariflora, *Schult. f. Syst.* vii 1195.—Bras.
remotiflora, Otto, in Otto & Dietr. Allg. Gartenz. i. (1833) 129 = rariflora.
sulphurea, *C. Koch, Ind. Sem. Hort. Berol.* (1873) *App.* iv. 3.—Reg. Argent.
tuberosa, Beer, Bromel. 187 = rariflora.

DYCTIOLOMA, DC. Prod. ii. 89 (1825) = **Dictyoloma** A. Juss. (Simarub.).

DYCTIOSPORA, Reinw. ex Korth. in Nederl. Kruidk. Arch. ii. ii. (1851) 157 = **Hedyotis**, Linn. (Rubiac.).
fagraeoides, Reinw. l. c. = H. vaginata.

DYCTISPERMA, Rafin. Sylva Tellur. 159 (1838) = **Rubus**, Tourn. (Rosac.).
apetalus, Rafin. l. c. 160 = R. apetalus.
laciocarpus, Rafin. l. c. = R. laciocarpus.
rigidus, Rafin. l. c. = R. rigidus.
urticifolius, Rafin. l. c. = R. urticaefolius.

DYCTOSPERMA, H. Wendl. in Kerch. Palm. 254 = **Dictyosperma**, H. Wendl. (Palmae).

DYDACTYLON, Zoll. Syst. Verz. Ind. Archip. 55 (1854) = DIDACTYLON, Mor. = **Dimeria**, R. Br. (Gramin.).

DYERA, Hook. f. in Journ. Linn. Soc. xix. (1882) 293. *APOCYNACEAE*.
costulata, *Hook. f. l. c.*—Malaya.
laxiflora, *Hook. f. Fl. Brit. Ind* iii. 644.—Singapore.
Lowii, *Hook. f. in Journ. Linn. Soc.* xix. (1882) 293. —Malaya.

DYLLWINIA, Nees, in R. Br. Verm. Schr. ii. 466 (1826), sphalm. = **Dillwynia**, Sm. (Legum.).

DYMCZEWICZIA, Horan. Prod. Scitam. 26 (1862) = **Zingiber**, Adans. (Scitam.).
capitata, Horan. l. c. = Z. capitatum.
elata, Horan. l. c. = Z. capitatum.
graminea, Horan. l. c. = Z. gramineum.
marginata, Horan. l. c. = Z. marginatum.

DYNAMIDIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 271 = **Potentilla**, Linn. (Rosac.).
agrestum, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 47, 170 = P. verna.
aureum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 271 = P. aurea.
australis, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 47, 170 = P. verna.
Clementi, Fourr. in Ann. Soc. Linn. Lyon N. S. xvi. (1868) 271 = P. subcaulis.
erodorum, Fourr. l. c.—Gallia.
juranum, Fourr. l. c. = P. rubens.
minimum, Fourr. l. c. = P. minima.
montanum, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 47, 170 = P. splendens.
montivagum, Timb. l. c. 47 = P. montivaga.
opacum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 271 = P. opaca.
procumbens, Fourr. l. c. = P. Sibbaldi.
reptans, Fourr. l. c. = P. reptans.
stipulatum, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 171 = P. verna.
subacaule, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 271 = P. subacaulis.
syenitenum, Fourr. l. c. = P. verna.
vernum, Fourr. l. c. = P. verna.
vicarieum, Fourr. l. c. = P. verna.
xerophilum, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 170 = P. verna.

DYNEBA, Lag. Gen. et Sp. Nov. 5 (1816) = **Dinebra**, Jacq. (Gramin.).

DYOSPYROS, Dum. Comm. Bot. 87 (1822) = **Diospyros**, Linn. (Ebenac.).

DYPLECOSIA, G. Don, Gen. Syst. iii. 788 (1834) = **Diplycosia**, Blume (Ericac.).

DYPLOTAXIS, DC. in Mém. Mus. Par. vii. (1821) 248 = **Diplotaxis**, DC. (Crucif.).

DYPONTIA, Dietz. ex Stend. Nom. ed. II. i. 534 (1840) = **Dupontia**, R. Br. (Gram.).

DYPSIS, Noronha, ex Thou. Prod. Phytol. 2; in Mélang. Bot., nomen (1811). *PALMAE*, Benth. & Hook. f. iii. 909.
forficifolia, *Noronha, ex Thou. l. c.*; *Mart. Hist. Nat. Palm.* iii. 180. t. 143. f. 2.—Madag.
hirtula, *Mart. l. c.* 181.—Madag.
nodifera, *Mart. l. c.* 312.—Madag.
pinnatifrons, *Mart. l. c.* 180. t. 158. f. 1.—Madag.

DYPTERYGIA, C. Gay, Fl. Chil. iii. 102 (1847) = DIPTERYGIA, Presl = **Asteriscium**, Cham. & Schlecht. (Umbellif.).

DYSCHORISTE, Nees, in Wall. Pl. As. Rar. iii. 75 (1832) = **Calophanes**, D. Don (Acanthac.).
biloba, Hochst. in Flora, xxviii. (1845) 72 = Asystasia coromandeliana.
cernua, Nees, in Wall. Pl. As. Rar. iii. 81 = C. Nagchana.
depressa, Nees, l. c. = C. Nagchana.
geniculata, Nees, in DC. Prod. xi. 107.—Madag.
literalis, Nees, in Wall. Pl. As. Rar. iii. 81 = C. littoralis.
radicans, Nees, in DC. Prod xi. 106 = C. radicans.

DYSEMONE, Soland. ex Forst. f. in Comm. Gotting. ix. (1789) 45, nomen = **Gunnera**, Linn. (Halorag.).
integrifolia, Banks & Soland. ex Forst. f. l. c. = G. magellanica.
lobata, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 275 = G. lobata.

DYSINANTHUS, Rafin. ex DC. Prod. vi. 270, in syn. (1837) = DISYNANTHUS, Rafin. = **Antennaria**, Gaertn. (Compos.).

DYSMICODON, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1842) 255 = **Specularia**, Heist. (Campan.).
californicum, Nutt. l. c. 256 = S. biflora.
ovatum, Nutt l. c. = S. biflora.
perfoliatum, Nutt. l. c. = S. perfoliata.

DYSODA, Lour. Fl. Cochinch. 145 (1790) = **Serissa**, Comm. (Rubiac.).
fasciculata, Lour. l. c. 146 = S. foetida.
foetida, Salisb. Prod. 60 = S. foetida.

DYSODIA, Cav. in Anal. Cienc. Nat. vi. (1802) 334 (Dysodia). *COMPOSITAE*, Benth. & Hook. f. ii. 406.
BOERREA, Willd. Sp. Pl. iii. 2125 (1803).
CLOMENOCOMA, Cass. in Bull. Soc. Philom. (1816) 199; et in Dict. Sc. Nat. ix. 416 (1817).
COMACLINIUM, Schtdw. & Planch. in Fl. des Serres, viii. (1852). 19. t. 756.
acerosa, DC. Prod. v. 641 = Hymenatherum acerosum.
alternifolia, *Regel & Korm. Ind. Sem. Hort. Petrop.* (1857) 41.—Hab. ?
anthemidifolia, *Benth. Bot. Voy. Sulph.* 29.—Calif.
appendiculata, *Lag. Gen. et Sp. Nov.* 29.—Mexic.
appendiculata, Sch. Bip. ex A. Gray, in Proc. Am. Acad. xix. (1884) 89 = squamosa.
cancellata, *A. Gray, in Proc. Am. Acad.* xix. (1883) 88.—Mexic.
Canantlensis, Lag. Gen. et Sp Nov. 29 = Adenophyllum coccineum.
chrysanthemoides, *Lag. l. c.*—Am. bor.
coccinea, Lag. l. c. = Adenophyllum coccineum.
Cooperi, *A. Gray, in Proc. Am. Acad.* ix. (1874) 201. —Calif.
fastigiata, DC. Prod. v. 640, excl. syn. = chrysanthemoides.
glandulosa, Cav. Desc. 309 = chrysanthemoides.
glandulosa, Less. Syn. Comp. 388 = Adenophyllum Porophyllum.
graveolifera, DC. Prod. v. 640.—Mexic.
incana, DC. l. c. = pubescens.
integrerrima, Lag. Gen. et Sp. Nov. 29 = pubescens.

DYSODIA :—
integrifolia, *A. Gray, in Proc. Am. Acad.* xix. (1883) 97.—Mexic.
montana, *A. Gray, l. c.* 88.—Guatemala.
porophylla, Lag. Gen. et Sp. Nov. 29 = *D. Porophyllum.*
porophylloides, *A. Gray, in Mem. Am. Acad.* N. S. v. (1856) 292.—Calif.
Porophyllum, Cav. in Anal. Cienc. Nat. vi. (1803) = Adenophyllum Porophyllum.
pubescens, *Lag. l. c.*—Mexic.
serratifolia, DC. Prod. v. 641 = Hymenatherum serratifolium.
speciosa, *A. Gray, in Proc. Am. Acad.* v. (1861) 163.—Calif.
squamosa, *A. Gray, l. c.* xix. (1883) 88.—Mexic.
subintegerrima, *Lag. Gen. et Sp. Nov.* 20.—Mexic.
tagetiflora, Lag. l. c. = chrysanthemoides.
tagetoides, Torr. & Gray, Fl. N. Am. ii. 361 = Hymenatherum tagetoides.

DYSODIA, Spreng. Anleit. ii. 11. 564 (1818) = DYSSODIA, Willd. = **Adenophyllum**, Pers. (Compos.).

DYSODIDENDRON, Gardn. in Calc. Journ. Nat Hist. vii. (1847) 1 = **Saprosma**, Blume (Rubiac.).
ceylanica, Gardn. l. c. 2 = S. ceylanicum.
glomeratum, Gardn. l. c. 3 = S. indicum.
Wightii, Gardn. l. c. = S. ceylanicum.

DYSODIUM, Rich. ex Pers. Syn. ii. 489 (1807) = **Melampodium**, Linn. (Compos.).
dinaricatum, Rich. l. c. = M. paludosum.
radiatum, Desf. Cat. Hort. Par. ed. III. 182 = M. paludosum.

DYSOPHYLLA, Blume, Bijdr. 826 (1826). *LABIATAE*, Benth. & Hook. f. ii. 1180.
CHOTEKIA, Opiz & Corda, in Flora, xlii. (1830) i. 33.
EUSTRALIS, Rafin. Fl. Tellur. ii. 95 (1836).
auricularia, *Blume, Bijdr.* 826.—As. trop.
Benthamiana, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 234 = verticillata.
crasicaulis, *Benth. in Wall. Pl. As. Rar.* i. 80.—Reg. Himal.
cruciata, *Benth. l. c.*—Ind. or.
erecta, *Dals. in Hook. Kew Journ.* ii. (1850) 337.—Ind. or.
gracilis, *Dals. l. c.*—Ind. or.
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 641.—Ind. or.
Helferi, *Hook. f. l. c.* 640.—Burma.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 109.—Japon.
lincaris, *Benth. in DC. Prod.* xii. 157.—Reg. Himal.
linearis, Wall. Cat. n. 1540 = quadrifolia.
myosuroides, *Benth. in Wall. Pl. As. Rar.* i. 80.—Ind. or.
myosuroides, Benth. l. c. partim = rugosa.
pentagona, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 641.—Ind. or.
pumila, Benth. in Wall. Pl. As. Rar. i. 80 = crasicaulis.
quadrifolia, *Benth. l. c.*—Ind. or.
ramosissima, Benth. in Wall. Cat. n. 1548 = verticillata.
rugosa, *Hook. f. Fl. Brit. Ind.* iv. 638.—Ind. or.
rupestris, Dalz. in Hook. Kew Journ. iii. (1851) 120 = quadrifolia.
salicifolia, *Dals. ex Hook. f. Fl. Brit. Ind.* iv. 638.—Ind. or.
Sampsoni, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 234.—China.
stellata, *Benth. in Wall. Pl. As. Rar.* i. 80.—Ind. or.
Stocksii, *Hook. f. Fl. Brit. Ind.* iv. 642.—Ind. or.
strigosa, Benth. in Wall. Pl. As. Rar. i. 80 = Pogostemon strigosus.
tetraphylla, Wight, Ic. t. 1444 = cruciata.
tomentosa, *Dals. in Hook. Kew Journ.* ii. (1850) 337.—Ind. or.
velutina, Benth. in Wall. Cat. n. 1538 = quadrifolia.
verticillata, *Benth. in Wall. Pl. As. Rar.* i. 80.—As. et Austral. trop.

DYSOPSIS. Baill. Etud. Gén. Euph. 435 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 264.
MIRABELLIA, Berr. ex Baill. Etud. Gén. Euph. 436 (1858).

DYSOPSIS:—
MOLINA, C. Gay, Fl. Chil. v. 345. t. 61 (1833). Gayana, *Baill. l. c.* 436.—Ins. J. Fernandez.
glechomoides, Muell. Arg. in DC. Prod. xv. ii. 949 = Gayana.

DYSOSMIA, Miq. Fl. Ind. Bat. ii. 325 (1856) = **Saprosma**, Blume (Rubiac.).
dichotoma, Miq. l. c. 326.—Malaya.

DYSOSMIA, M. Roem. Syn. Pepon. 149 (1846) = **Passiflora**, Linn.
acerifolia, M. Roem. l. c. 151 = P. adenopoda.
ciliata, M. Roem. l. c. 149 = P. foetida.
fluminensis, M. Roem. l. c. 150 = P. Vellozii.
foetida, M. Roem. l. c. 149 = P. foetida.
gossypiifolia, M. Roem. l. c. = P. foetida.
hastata, M. Roem. l. c. = P. foetida.
hibiscifolia, M. Roem. l. c. = P. foetida.
nigelliflora, M. Roem. l. c. 150 = P. foetida.
pohyadena, M. Roem. l. c. 151 = P. foetida.

DYSOSMON, Rafin. Fl. Ludov 48 (1817) = **Sesamum**, Linn. (Pedalin.).
amorum, Rafin. l. c. = S. indicum.

DYSOXYLON, Bartl. Ord. 356 (1830) = seq.

DYSOXYLUM, Blume, Bijdr. 172 (1825). *MELIACEAE*, Benth. & Hook. f. i. 332.
CAMANIA, Comm. ex M. Roem. Syn. Hesper. 83, 102 (1846).
DIDYMOCHETON, Blume, Bijdr. 177 (1825).
GONIOCHETON, Blume, l. c. (1825).
GONIOCHETON, G. Don, Gen. Syst. i. 683 (1831).
HARTIGHSEA, A. Juss. in Mém. Mus. Par. xix. (1830) 227, t. 4.
PLAGIOLOM, M. Roem. Syn. Hesper. 83 (1846).
aculeatissimum, Stend. Nom. ed. II. i. 534, aphalm. = seq.
acuminatissimum, Blume, Bijdr. 173 = alliaceum.
acutangulum, *Miq. Fl. Ind. Bat. Suppl.* 508.—Sumatra.
albicans, *Viell. ex C. DC. in DC. Monog. Phan.* i. 522.—N. Caled.
alliaceum, *Blume, Bijdr.* 172.—Java.
alliaceum, Seem. Fl. Vit. 86 = Richii.
amooroides, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 16.—N. Guinea.
arborescens, *Miq. l. c.* 24.—Borneo.
Balansesanum, *C. DC. in DC. Monog. Phan.* i. 508.—N. Caled.
Becccarianum, *C. DC. l. c.* 495.—Borneo.
Beckerianum, *C. DC. l. c.* 509.—Australi.
Beddomei, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 548.—Ind. or.
biyugum, Seem. Fl. Vit. 87 = Lessertianum.
binectariferum, Hiern, in Hook. f. Fl. Brit. Ind. i. 546, ex parte = Amoora filicifornia.
binectariferum, *Hook. f. ex Bedd. in Trans. Linn. Soc.* xxv. (1866) 212.—Ind. or.
binectariflorium, C. DC. in DC. Monog. Phan. i. 502 = Amoora filicifornia.
Blancoi, *Vidal. Cat. Pl. Pr. Man.* 22; *Sinops.* p. xix.—Ins. Philipp.
Blumei, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 25.—Java.
brevipes, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 546.—Malacca.
camalense, *C. DC. in DC. Monog. Phan.* i. 505.—N. Caled.
caulifiorum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 549.—Malacca.
caulostachyum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 12.—N. Guinea.
Championii, Hook. f. & Thoms. ex Thw. Enum. Pl. Zeyl. 61 = Pseudocarapa Championii.
chrysophyllinum, *Viell. ex C. DC. in DC. Monog. Phan.* i. 534.—N. Caled.
costulatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 21.—Sumatra.
costulatum, Miq. l. c. ex parte = Miquelianum.
Cumingianum, *C. DC. in DC. Monog. Phan.* i. 497.—Ins. Philipp.
cuneatum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 549.—Malacca.
cyrtobotryum, *Miq. Fl. Ind. Bat. Suppl.* i. 504.—Sumatra.

DYSOXYLUM:—
dasyphyllum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 19.—Ins. Celebes.
densiflorum, *Miq. l. c.* 9.—Java.
distantinerve, F. Muell. Fragm. ix. 61 = latifolium.
euneuron, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 22.—Ins. Cerum.
excelsum, *Blume, Bijdr.* 176.—Java.
flavescens, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 549.—Malacca.
Forsteri, *C. DC. in DC. Monog. Phan.* i. 507.—Austral foveolatum, *Radlk. in Sitzb. Math.-Phys. Acad. Munech.* ix. (1879) 998.—Ins. Timor.
Fraserianum, *Benth. Fl. Austral.* i. 381.—Austral.
fraternum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 25.—Java.
gaudepense, *C. DC. in DC. Monog. Phan.* i. 524.—N. Caled.
Gaudichaudianum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 15.—Ins. Kawak.
glabrum, *C. DC. in DC. Monog. Phan.* i. 488.—Java.
glomeratum, *Vieill. ex C. DC. l. c.* 521.—N. Caled.
grande, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 547.—Ind. or.
grandiflorum, Wight & Arn. Prod. 128 = Chisocheton grandiflorus.
Griffithii, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 549.—Malacca.
Helmheirae, *C. DC. in DC. Monog. Phan.* i. 488.—Java.
Hamiltonii, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 548.—Reg. Himal.
Klanderi, *F. Muell. Fragm.* v. 176.—Austral.
Kunthianum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 13—N. Guin.
Lampongum, Miq. Fl. Ind. Bat. Suppl. i. 508 = macrothyrsum.
lasiocarpum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 13—N. Guin.
latifolium, *Benth. Fl. Austral.* i. 381.—Austral.
latifolium, F. Muell. Fragm. v. 144 = oppositifolium.
laxiflorum, Blume, Bijdr. 174 = alliaceum.
Lenormandianum, *C. DC. in DC. Monog. Phan.* i. 512.—N. Caled.
Lessertianum, *Benth. Fl. Austral.* i. 382.—Austral.
Lobbii, *C. DC. in DC. Monog. Phan.* i. 484.—Java.
longifolium, Blume, Bijdr. 173 = alliaceum.
macranthum, *C. DC. in DC. Monog. Phan.* i. 525.—N. Caled.
macrocarpum, *Blume, Bijdr.* 175.—Java.
macrocarpum, Thw. Enum. Pl. Zeyl. 60 = binectariferum.
macrophyllum, *Teijsm. & Binn. Cat. Hort. Bog.* 211, nomen.—Java.
macrostachyum, *C. DC. in DC. Monog. Phan.* i. 504.—N. Caled.
macrothyrsum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 20.—Borneo.
Maingayi, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 547.—Malacca.
malabaricum, *Bedd. ex C. DC. in DC. Monog. Phan.* i. 491.—Ind. or.
minutiflorum, *C. DC. l. c.* 487.—N. Caled.
Miquelianum, *C. DC. l. c.* 486.—Malaya.
molle, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 18.—N. Guinea.
mollissimum, *Blume, Bijdr.* 176.—Java.
Mueller, *Benth. Fl. Austral.* i. 381.—Austral.
multijugum, Arn. ex Wight & Arn. Prod. 121 = Chisocheton paniculatus.
Nageliianum, *C. DC. in DC. Monog. Phan.* i. 504.—Java.
Nernsti, *F. Muell. Fragm.* v. 176.—Austral.
nitidum, *C. DC. in DC. Monog. Phan.* i. 532.—N. Caled.
nutans, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 17.—Java.
oppositifolium, *F. Muell. Fragm.* v. 177, 144.—Java.
otophorum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 15.—Java.
pachypodum, *C. DC. in DC. Monog. Phan.* i. 516.—N. Caled.
pallens, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 548.—Reg. Himal.
Pancheri, *C. DC. in DC. Monog. Phan.* i. 514.—N. Caled.
paniculatum, Arn. ex Wight, Ic. t. 146 = Chisocheton paniculatus.

DYSOXYLUM :—
parvifolium, *C. DC. in DC. Monog. Phan.* i. 511.—N. Caled.
Paterson, *F. Muell. Census, Suppl.* i. 9.—Ins. Norfolc.
procerum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 547.—Ind. or.: Burma.
ptychocarpum, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 593.—Austral.
pubescens, *Teijsm. & Binn. Cat. Hort. Bog.* 211, nomen.—Java.
ramiflorum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 10.—Java.
Richii, *C. DC. in DC. Monog. Phan.* i. 511.—Ins Fiji.
roseum, *C. DC. l. c.* 514.—N. Caled.
rufescens, *Vieill. ex Panch. & Sebert, Not. Bois Nouv. Caléd.* 226.—N. Caled.
rufum, *Benth. Fl. Austral.* i. 382.—Austral.
salutare, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 42 = Blanco1.
samoense, *A. Gray, Bot. U. St. Expl. Exped.* i. 241.—Ins. Tutuila.
Schiffneri, *F. Muell. in Melb. Chemist* (1881), 53 ; *et in Trans Bot. Soc. Edinb.* xiv. (1883) 369.—Austral.
schizochitoide, *C. DC. in DC. Monog. Phan.* i. 519.—Malaya.
Schmidii, *C. DC. l. c.* 502.—Austral.
sessile, *Miq. Ann. Mus. Bot. Lugd. Bat* iv. 15.—Ins. Ambolna.
setosum, *Miq. l. c.* 13.—Ins. Timor.
simile, *Blume, Bijdr* 174.—Java.
Spanoghei, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 14.—Ins. Timor.
speciosum, *Miq. l. c.* 11.—Ins. Timor.
spectabile, *Hook. f. Handb. N. Zeal. Fl.* 41.—N. Zel.
spiciflorum, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 35 = Amoora timorensis.
Teysmannii, *C. DC. in DC. Monog. Phan.* i. 510.—Java.
thyrsoideum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 547.—Malacca.
trichostylum, *Miq. Ann Mus. Bot. Lugd. Bat.* iv. 11.—Java.
Turczaninowii, *C. DC. in DC. Monog. Phan.* i. 501.—Ins. Philipp.
unijugum, *C. DC. l. c.* 507.—N. Caled.
Vieillardi, *C. DC. l. c.* 513.—N. Caled.
Vrieseanum, *C. DC. l. c.* 491.—Java.

DYSPHANIA, R. Br. Prod. 411 (1810). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 14.
littoralis, *R. Br. Prod.* 413.—Austral.
littoralis, Moq in DC. Prod. xiii. 11. 86 = myriocephala.
myriocephala, *Benth. Fl. Austral.* v. 165.—Austral.
Plantaginella, *F. Muell. Fragm.* i. 61.—Austral.

DYSSOCHROMA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 250. *SOLANACEAE*, Benth. & Hook. f. ii. 901.
TRIANAEA, Planch. & Linden, in Linden, Prix-Cour. n. 8 (1853), 4.
altidoflavum, *Lem. Illustr. Hortic.* vi. (1859) *Misc.* 40.—Bras.
eximia, *Benth. & Hook. f. Gen.* ii. 904, in nota.—Hab.? longipes, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 253.—Bras.
viridiflora, *Miers, l. c.* 251.—Bras.

DYSSODIA, Cav. in Anal. Clenc. Nat. vi. (1802) 334 = **Dysodia**, Cav. (Compos.).

DYSSODIA, Willd. Enum. Hort. Berol. 900 (1809) = **Adenophyllum**, Pers. (Compos.).
porophylla, DC. Prod. v. 589 = seq.
Porophyllum, Willd. Enum. Hort. Berol. 900 = A. Porophyllum.

DEIRDUSEYCKIA, Rehm. in Oestr. Bot. Zeitschr. xviii. (1868) 374. *NAIADEAE*.
limnobia, *Rehm. l. c.*—Reg. Caucas.

E

EADESIA, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 71 = **Anthocercis**, Labill. (Solanac.).
anthocercidea, F. Muell. l. c. 72 = A. Endesii.

EAPLOSIA, Rafin. New Fl. Am. ii. 51 (1836) = **Baptisia**, Vent. (Legumin.).
longifolia, Rafin. l. c. 52.—Am. bor.
ovata, Rafin. l. c. = B. simplicifolia.

EARINA, Lindl. Bot. Reg. sub t. 1699 (1834). *ORCHIDEAE*. Benth. & Hook. f. iii. 516.
autumnalis, Hook. f. Fl. N. Zel i. 239 = suaveolens.
Deplanchei, *Reichb. f. in Linnaea*, xli. (1877) 96.—N. Caled.
laxior, *Reichb. f. Otia Bot. Hamb.* 54.—Ins Talti.
mucronata, *Lindl. Bot. Reg.* sub t. 1699.—N. Zel.
plana, *Reichb. f. Otia Bot. Hamb.* 54.—Ins. Fiji.
quadriloba, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 325.—N. Zel.
suaveolens, *Lindl. Bot. Reg.* (1843) *Misc.* 61.—N. Zel.
valida, *Reichb. f. in Linnaea*, xli. (1877) 96.—N. Caled.

EARLIA, F. Muell. Fragm. iii. 159 (1863) = **Graptophyllum**, Nees (Acanthac.).
excelsa, F. Muell. l. c. 160 = G. Earlii.

EATONELLA, A. Gray, in Proc. Am. Acad. xix. (1883) 19. *COMPOSITAE*.
Congdoni, *A. Gray, l. c.* 20.—Calif.
nivea, *A. Gray, l. c.* 19.—Am. bor. occ.

EATONIA, Rafin. in Journ. Phys. lxxxix. (1819) 104 *GRAMINEAE*, Benth & Hook. f. iii. 1164.
COLOBANTHUS, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1830) 66.
REDOULEA, Kunth, Rév. Gram. 341 (1830).
obtusata, *A. Gray, in S. Wats. Bot. Calif.* ii. 302.—Am. bor. occ.
purpurascens, *Rafin. in Journ. Phys.* lxxxix. (1819) 104.—Am. bor.

EBELIA, Reichb. Nom. 74 (1841) = **Triodon**, DC. (Rubiac.).

EBELINGIA, Reichb. Consp 199 (1828) = **Harrisonia**, R. Br. (Simarub.).
Brownei, Steud. Nom. ed. II. i. 535 = H. Brownei.

EBENIDIUM, Jaub. & Spach, in Ann Sc. Nat. Sér. II. xix. (1843) 162 = **Ebenus**, Linn. (Legumin.).
Lagopus, Jaub. & Spach, l. c. 163 = Ebenus Lagopus.

EBENOXYLON, Spreng. Gen. i. 40 (1830) = seq.

EBENOXYLUM, Lonr. Fl. Cochinch. ii. 618 (1790) = **Maba**, Forst. (Ebenac.).
serum, Lour. l. c. = M. elliptica.

EBENUS, Linn. Sp. Pl. 764 (1753). *LEGUMINOSAE*, Benth. & Hook. f. i. 512.
ERENDIUM, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 162.
barbigera, *Boiss. Diagn.* Ser. II. ii. 98.—As. Min.
Boissieri, *Heldr.*, *Barbey, in Bull. Soc. Vaud. Sc. Nat.* xxi. (1885) 95.—Orient.
Bourgaei, *Boiss. Fl. Orient.* ii. 554.—Lycia.
capensis, Linn. Mant. 284 = Lebeckia cytisoides.
cilicica, Schott & Kotschy, ex Boiss. Fl. Orient. ii. 556 = Montbretii.
cretica, *Linn. Sp. Pl.* 764.—Ins. Creta.
depressa, *Boiss. & Bal. Diagn.* Ser. II. vi. 60.—Cappadoc.
erinacea, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 161 = stellata.
ferruginea, Jaub. & Spach, l. c. 160 = stellata.
hirsuta, *Jaub. & Spach, Illustr.* iii. 85. t. 251.—As. Min.
horrida, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 159 = stellata.
Lagopus, *Boiss. Fl. Orient.* ii. 558.—Persia.
lagwroides, Boiss. Diagn. Ser. I. ii. 99 = Montbretii.
leucomelas, Comm. ex Poir. Encyc. v. 432 = Diospyros leucomelas.
longipes, *Boiss. & Bal. Diagn.* Ser. II. vi. 60.—Cappadoc.
macrophylla, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. II. xix. (1843) 156.—Cappadoc.
melanida, Comm. ex Poir. Encyc. v. 431 = Diospyros melanida.
moluccs, *Willem. in Ustevi, Ann. Bot.* xviii. (1796) 60.—Ins. Maurit.

EBENUS :—
Montbretii, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. II xix. (1843) 154.—Cappadoc.
pinnata, *Ait. Hort. Kew.* ii. 27.—Barbaria.
pinnata, Sibth. & Sm. Fl. Graec. t. 740 = Sibthorpii.
plumosa, *Boiss. & Bal. Diagn.* Ser. II. vi. 59.—As. Min.
pogonotropis, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xix (1843) 155 = barbigera.
Sibthorpii, *DC. Mém. Leg.* vii. t. 53.—Graecia.
stellata, *Boiss. Diagn.* Ser. I. ii. 100.—Arab ; Persia.
stipulacea, Graeb. Specil. Fl. Rumel. i. 12 = Montbretii.
tessellaria, Comm. ex Poir. Encyc. v. 430 = Diospyros melanida.
tragacanthoides, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 158 = stellata.

EBERLEA, Riddell, ex Nees, in DC. Prod. xi. 85 (1847) = **Hygrophila**, R. Br. (Acanthac.).
glomerata, Riddell, ex Nees, l. c. 86 = H. lacustris.

EBERMAIERA, Nees, in Wall. Pl. As. Rar. iii. 75 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1074.
ERYTHRACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 75 (1832).
STAUROGYNE, Wall. l. c. in 80. t. 186 (1832).
STIFTIA, Pohl, ex Nees, in DC. Prod. xi. 70 (1847).
angustifolia, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 451.—Ind. or.
angozanthus, *Nees, in Mart. Fl. Bras.* ix. 17.—Bras.
argentea, Nees, in DC. Prod. xi. 76 = lanceolata.
aristata, T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 396 = angustifolia.
axillaris, *Nees, in Wall. Pl. As. Rar.* iii. 80.—Penang.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 402.—Burma.
brachiata, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 69.—Bras.
concannula, *Hance, in Journ. Bot.* vi. (1868) 300.—China.
coriacea, *T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 402.—Burma.
debilis, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 452.—Ins. Philipp.
diffusa, Kurz, in Journ. As. Soc. Beng. xl. (1871) 11. 74 = thyrsoidea.
elegans, *Nees, in Mart. Fl. Bras.* ix. 17.—Bras.
elongata, *Nees, in DC. Prod.* xi. 721.—Malaya.
fastigiata, *Nees, in Mart. Fl. Bras.* ix. 19.—Bras.
glauca, T. Anders. in Journ. Linn. Soc. ix. (1867) 450, partim = glutinosa.
glauca, T. Anders. in Thw. Enum. Pl. Zeyl. 224 = zeylanica.
glauca, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 310. —Ind. or.
glutinosa, *Wall. Cat.* n. 9087.—Ind. or.
gracilis, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 453.—Burma.
gracilis. *Wawra, Itin. Princ. Coburgi*, i. 94. t. 10 B.—Bras.
Griffithiana, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 452.—Malacca.
Helferi, *T. Anders. l. c.*—Burma.
Helferi, T. Anders. in Trim. Kew. ex T. Anders. in Hook. f. Fl. Brit. Ind. iv. 400 = E. macrophylla.
hirsuta, *Nees, in Mart. Fl. Bras.* ix. 18.—Bras.
humilis, *Mig. Stirp. Surin. Select.* 138.—Guiana.
humilis, *Nees, in Wall. Pl. As. Rar.* iii. 80.—Burma.
incana, *Hassk. Cat. Hort. Bog. Alt.* 147.—Burma ; Java.
Itatiaiae, *Wawra, Itin. Princ. Coburgi*, i. 93. t. 11.—Bras.
lanceolata, *Hassk. Cat. Hort. Bog. Alt.* 147.—Malaya.
laxiobotrys, *Nees, in DC. Prod.* xi. 732.—Burma.
ligulata, Bedd. Ic. Ind. Or. i. 59 = glauca.
longifolia, *Nees, in DC. Prod.* xi. 76.—Malacca.
Maccellandii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 454.—Burma.
macrophylla, *T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 400.—Burma.
mandioccana, *Nees, in Wall. Pl. As. Rar.* iii. 80.—Bras.
merguensis, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 452.—Burma.

EBERMAIERA :—
minarum. *Nees, in Mart. Fl. Bras.* ix. 17.—Bras.
Nessii. *Vidal, Phan. Cuming. Philipp.* 34.—Ins. Philipp.
nelsonioides, *Miq. Fl. Ind. Bat. Suppl.* 566.—Ins. Banca.
nitida, *Reichb. f. in Gard. Chron.* (1879) i. 812.—Bras.
obtusa, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 468.—Burma.
paniculata, *Wall. ex T. Anders. l. c.*—Ind. or.
parviflora, *T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 402.—Burma.
pedicellata, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 176.—Ind. or
polybotrya, *Nees, in DC. Prod.* xi. 74.—Ind. or.; Burma.
pulchella, Zipp. ex Span. in Linnæa, xv. (1841) 183 = Turræa pinnata.
racemosa, *Miq. Fl. Ind. Bat.* ii. 775.—Malaya.
repens, *Nees, in Mart. Fl. Bras.* ix. 20.—Guiana.
Riedeliana, *Nees, l. c.* 18.—Ilras.
Sanctæ-Catharinæ, *Nees, in DC. Prod.* xi. 72.—Bras.
setigera, T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit Ind. iv. 398 = angustifolia.
setigera, *Nees, in DC. Prod.* xi. 76.—Malaya.
Simonsii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 451.—Reg. Himal.
spathulata, Hassk. Cat. Hort. Bog. Alt. 147 = glauca.
spicifera, *Miq. Fl. Ind. Bat. Suppl.* 566.—Sumatra.
Staurogyne, T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 398 = angustifolia.
Staurogyne, *Nees, in DC. Prod.* xi. 75.—Reg. Himal.
stolonifera, *Nees, in Mart. Fl. Bras.* ix. 19.—Bras.
subpaniculata, Hassk. in Retzia, i. 77 = elongata.
thyrsoides, *Nees, in Wall. Pl. As. Rar.* iii. 80.—Burma.
trichocephala, Miq. Fl. Ind. Bat. ii. 774 = setigera.
Vauthieriana, *Nees, in Mart. Fl. Bras.* ix. 15.—Bras.
velutina, Nees, in DC. Prod. xi. 76 = incana.
venosa, T. Anders. in Journ. Linn. Soc. ix. (1867) 453 = paniculata.
vercoicifolia, *Nees, in Mart. Fl. Bras.* ix. 18.—Bras.
Warmingiana, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 68.—Bras.
zeylanica, T. Anders. in Journ. Linn. Soc. ix. (1867) 451, partim = thyrsoides.
zeylanica, *Nees, in DC. Prod.* xi. 74.—Ind. or.

EBERMAVERA, Meisn. Gen. 298 (1840) = praec.

EBERMEVERA, Endl. Gen. 698 (1839) = praec.

EBERMIERA, Wight, Ic. iv. 17 (1849) = praec.

EBULUM, Garcke, Fl. Deutschl. ed. VII. 184 (1865) = **Sambucus**, Tourn. (Caprifol.).
humile, Garcke. l. c. = S. Ebulus.

EBULUS, Pont. ex Linn. Syst. ed. I. (1735) = **Sambucus**, Tourn. (Caprifol.).

EBURNAX, Rafin. New Fl. Am. i. 42 (1836) = **Mimosa**, Tourn. (Legumin.).
pudica. Rafin. l. c. = M. pudica.
spinalba, Rafin. Sylva Tellur. 119 = Acacia eburnea.

EBUROPETALUM, Becc. in Nuov. Giorn. Bot. iii. (1871) 181. *ANONACEAE.*
borneense, *Becc. l. c.*—Borneo.

ECASTAPHYLLUM, P. Br. Hist. Jamaic. 299 (1756). *LEGUMINOSAE*, Benth. & Hook. f. i. 545.
ACOUROA, Aubl. Pl. Gui. 758. t. 301 (1775).
DRAKENSTENIA, Neck. Elem. iii. 33 (1790).
HECASTOPHYLUM, H. B. & K. Nov. Gen. et Sp. vi. 397 (1824).
PTEROCARPUS, Berg. in Vet. Acad. Handl. Stockh. 116 (1769).
Benthamiorum, Miq. in Linnæa, xviii. (1844) 575 = E. Monetaria.
Berterii, DC. Prod. ii. 421 = E. Monetaria.
Brownei, *Pers. Syn.* ii. 277.—Am. austr.
dubium, *H. B. & K. Nov. Gen. et Sp.* vi. 398.—Venezuela.

ECASTAPHYLLUM :—
foliosum, Benth. in Hook. Journ. Bot. ii. (1840) 64 = glaucum.
glaucum, *Desv. in Ann. Sc. Nat. Sér.* I. ix. (1826) 428.—Porto Rico.
hygrophilum, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 93 = E. Monetaria.
microphyllum, *H. B. & K. Nov. Gen. et Sp.* vi. 398. —Venezuela.
molle, Miq. in Linnæa, xix. (1847) 487 = Brownei.
Monetaria, *Pers. Syn.* ii. 277.—Am. trop.
nitidum, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 93 = E. Monetaria.
Plumieri, Pers. Syn. ii. 277 = E. Monetaria.
pubescens, *DC. Prod.* ii. 421.—Guiana.
Richardi, Pers. Syn. ii. 277 = E. Monetaria.
Sieberi, Reichb. ex Steud. Nom. ed. II. i. 535 = Brownei.
tomentosum, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. i. 239.—Bras.
violaceum, Benth. in Ann. Wien. Mus. ii. (1838) 93 = Geoffrea violacea.

ECBALIUM, DC. Prod. iii. 311, in syn. (1828) = seq.

ECBALLION, Koch, Syn. Fl. Germ. 251 (1835) = seq.

ECBALLIUM, A. Rich. in Dict. Class. Hist. Nat. vi. 19 (1824). *CUCURBITACEAE*, Benth. & Hook. f. i. 826.
ECBALIUM, DC. Prod. iii. 311, in syn. (1828)
agreste, Reichb. Fl. Germ. Excurs. 294 = E. Elaterium.
Elaterium, *A. Rich. in Dict. Class. Hist. Nat.* vi. 19.—Reg. Mediterr.
Lambertianum, M. Roem. Syn. Pepon. 52 = Cucumis trigonus.
officinale, T. Nees, Pl. Off. t. 271 = E. Elaterium.
officinarum, Rich. ex M. Roem. Syn. Pepon. 52 = E. Elaterium.
purgans, Schrad. in Linnæa, xii. (1838) 421 = E. Elaterium.

ECBOLIUM, Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 75. *ACANTHACEAE*, Benth. & Hook. f. ii. 1113.
Linneanum, *Kurz, l. c.*—As. et Afr. trop.
striatum, *Baif. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 89.—Ins. Socotra.

ECBOLIUM, Riv. ex Linn. Syst. ed. I (1735) = **Justicia**, Linn. (Acanth.).

ECCLIMUSA, Mart. ex A. DC. in DC. Prod. viii. 156 (1844) = seq.

ECCLINUSA, Mart. in Flora, xxii. (1839) i. Beibl. 2 (Herb. Fl. Bras. 177). *SAPOTACEAE*, Benth. & Hook. f. ii. 654.
PASSAVERIA, Mart. & Richl. Fl. Bras. vi. 85. tt. 38 et 47. f. 1, 2 (1863).
ramiflora, *Mart. Ill. cc.*—Bras.

ECCOILOPUS, Stend. Syn. Pl. Gram. 128 (1855) = **Eulalia**, Trin. (Gramin.).
andropogonoides, Steud. l. c. 124 = E. cotulifera.

ECCREMANTHUS, Thw. in Hook. Kew Journ. vii. (1855) 279. t. 9 = **Pometia**, Forst. f. (Sapindac.).
eximius, Thw. l. c. = P. tomentosa.

ECCREMIS, Baker, in Journ. Linn. Soc. xv. (1876) 319 aphalm. = **Bixoremis**, Willd. (Liliac.).

ECCREMOCARPUS, Ruiz & Pav. Prod. 90. t. 18 (1794). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1050.
CALAMPELIS, D. Don, in Edinb. N. Phil. Journ. vii. (1829) 89.
longiflorus, *Humb. & Bonpl. Pl. Aequin.* i. 229. t. 65. —Peruv.
rubet, *Regel. Cat. Pl. Hort. Absahov.* 51.—Hab. ?
scaber, *Ruiz & Pav. Syst. Veg.* 167.—Chili.
sepium, Bert. in 'Merc. Chil. 1829'.—Cf. Bull. Férussac, xx. (1829) 111 = scaber.
viridis, *Ruiz & Pav. Prod.* 90. t. 18.—Peruv.

ECDEIOCOLEA, F. Muell. Fragm. viii. 236 (1874). *RESTIACEAE*, Benth. & Hook. f. iii. 1080.
monostachya, *F. Muell. l. c.*—Austral.

ECDYSANTHERA, Hook. & Arn. Bot. Beech. Voy. 195. t. 42 (1836). *APOCYNACEAE*, Benth. & Hook. f. ii. 714.
DENDROCHARIS, Miq. Fl. Ind. Bat. iii. 462 (1856).
barbata, Miq. Fl. Ind. Bat. ii. 451 = Parameria glandulifera.
brachiata, A. DC. in DC. Prod. viii. 448 = micrantha.
densiflora, *Miq. Fl. Ind. Bat.* ii. 452.—Java.
glandulifera, A. DC. in DC. Prod. viii. 448 = Parameria glandulifera.
Griffithii, Wight, Ic. t. 1307 = Parameria glandulifera.
micrantha, *A. DC. in DC. Prod.* viii. 443.—Reg. Himal. ; Burma.
pedunculosa, Miq. Fl. Ind. Bat. Suppl. 857 = Parameria pedunculosa.
rosea, *Hook. f. Arn. Bot. Beech. Voy.* 198.—China.
scandens, Hassk. Cat. Hort. Bog. Alt. 209 = Anodendron infastum.
Schriechii, Heurck & Muell. Arg. in Heurck, Obs. Bot. 191 = Microechites Schriechii.
torosa, Llanos, Fragm. 51 = Parameria philippinensis.

ECHALTIUM, Wight, Ic. ii. 4. t. 472 (1841) = **Melodinus**, Forst. (Apocyn.).
picciduum, Wight, l. c. = M. monogynus.

ECHBANDIA, Orteg. Hort. Matr. 90, 135. t. 18 (1797). *ILLIACEAE*, Benth. & Hook. f. iii. 789.
alnifolia, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) reimpr. 15 = ternifiora.
eleutherandra, *C. Koch, Ind. Sem. Hort. Berol.* (1861) *App.* 6.—Mexic.
gramnea, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) reimpr. 16 = Antherieum leptophyllum.
Haenkeana, Kunth, Enum. Pl. iv. 629 = ternifiora.
leptophylla, Benth. Pl. Hartw. 23 = Antherieum leptophyllum.
linearifolia, Klotzsch, in Otto & Dietr. Allg. Gartenz. viii. (1840) 275 = ternifiora.
scabrella, Wslp. Ann. iii. 1010 = Antherieum scabrellum.
ternifiora, *Ortog. Hort. Matr.* 90, 136.—Mexic.

ECHENAIS, Cass. in Bull. Soc. Philom. (1818) 88 = **Cnicus**, Tourn. (Compos.).
carlinoides, Cass. in Dict. Sc. Nat. xiv. 170 = C. carlinoides.
cirsioides, DC. Prod. vi. 661 = C. argyracanthus.
feros, Edgew. ex C. B. Clarke, Comp. Ind. 118 = C. Wallichii.
nudana, Cass. in Dict. Sc. Nat. xiv. 170 (= *Cirsium scleranthum*).—As. Min. ; Persia.
scleranthe, C. Koch, in Linnæa, xxiv. (1851) 404 (= *Cirsium scleranthum*).—As. Min. ; Persia.
Sieversii, Fisch. & Mey. in Schrenk, Enum. Pl. Nov. 44.—Sibir.

ECHETROSIS, Phil. in Anal. Univ. Chil. (1872) 129 (reimpr. 28). *COMPOSITAE.*
pentasperma, *Phil. ll. cc.*—Reg. Argent.

ECHEVERIA, DC. Prod. iii. 401 (1828) = **Cotyledon**, Linn. (Crassul.).
abyssinica, Otto, in Hamb. Gartenz. xxix. (1873) 6.—Abyss.
acutifolia, Lindl. Bot. Reg. (1843) t. 29 = C. acutifolia.
adunca, Otto, in Hamb. Gartenz. xxix. (1873) 6 = C. aduncа.
agavoides, Lem. Illustr. Hortic. x. (1863) Misc. 78 = C. agavoides.
amœna, L. de Smet, ex E. Morr. in Belg. Hortic. xxv. (1875) 216.—Mexic.
argentea, Hort. ex Lem. Illustr. Hortic. x. (1863) Misc. 78 = C. pulverulenta.
atropurpurea, Hort. ex Baker, in Saund. Refug. Bot. iii. (1870) sub t. 198 = C. atropurpurea.
Bernhardyana, Poerst. ex Lem. Illustr. Hortic. x. (1863) Misc. 78.—Hab. ?
bifida, Schlecht. in Linnæa, xiii. (1839) 411 = C. bifida.
bracteolata, Link, Klotzsch, & Otto, Ic. Pl. ii. 68 = C. bracteolata.
bracteosa, Lindl. & Paxt. Flow. Gard. iii. (1852–53) 60 = C. Pachyphytum.
caespitosa, DC. Prod. iii. 401 = C. caespitosa.
californica, Hort. ex Baker, in Saund. Refug. Bot. i. sub t. 70 = C. californica.

ECHEVERIA :—
calophana, Hort. ex Lem. Illustr. Hortic. x. (1863) Misc. 78 = Cotyledon acutifolia.
campewalata, Kunze, Delect. Sem. Hort. Lips. (1842); et in Linnæa, xvii. (1843) 574 = C. caespitosa ?
canaliculata, Hook. Bot. Mag. t. 4986 = C. canalicula
carnicolor, E. Morr. Belg. Hortic. xxiv. (1874) 158 = C. carnicolor.
clavifolia, Deleuil, ex E. Morr. l. c. 329 = C. Pachyphytum.
coccinea, DC. Prod. iii. 401 = C. coccinea.
Cooperi, Otto, in Hamb. Gartenz. xxix. (1873) 7 = C. Cooperi.
Corderoyi, E. Morr. in Belg. Hortic. xxiv. (1874) 159 = C. Corderoyi.
cymosa, Hort. ex Lem. Illustr. Hortic. x. (1863) Misc. 79 = C. cymosa.
decipiens, E. Morr. in Belg. Hortic. xxiv. (1874) 159 = C. decipiens.
Desmetiana, L. De Smet, ex E. Morr. l. c. = C. Desmetiana.
discolor, L. De Smet, ex E. Morr. l. c.—Mexic.
erecta x, Deleuil, ex E. Morr. l. c. 329 = C. coccinea.
eximia, De Vos, ex E. Morr. l. c. xxv. (1875) 99.—Hab. ?
farinosa, Hort. ex Van Houtte, in Fl. des Serres, xviii. (1873) 1 = C. pulverulenta.
farinosa, Lindl. in Journ. Hort. Soc. iv. (1849) 292 = C. farinosa.
farinulenta, Lem. Illustr. Hortic. xi. (1864) sub t. 392 = C. farinulenta.
ferrea x, Deleuil, ex E. Morr. in Belg. Hortic. xxiv. (1874) 329 = C. Scheerii.
fulgens, Lem. in Hort. Van Houtt. (1845) fasc. i. 8 ; et Jard. Fleur iii. (1853) t. 244 = C. fulgens.
Funki, L. De Smet, ex E. Morr. Belg. Hortic. xxvii. (1877) 169.—Hab. ?
gibbiflora, DC. Prod. iii. 401 = C. gibbiflora.
glauca, Hort. ex Baker, in Saund. Refug. Bot. i. sub t. 61 = C. glauca.
globosa, Hort. ex E. Morr. in Belg. Hortic. xxiv (1874) 161 = C. glauca ?
gracillima, Muehlenpf. ex Otto, in Hamb. Gartenz. xxix. (1873) 7 = C. secunda.
grandiflora, Haw. in Tayl. Phil. Mag. (1828) 268 = C. gibbiflora.
grandifolia, Otto, in Hamb. Gartenz. xxix. (1873) 7 = C. gibbiflora.
grandis, Hort. ex E Morr. in Belg. Hortic. xxiv. (1874) 161 = C. gibbiflora.
grandiserpala x, Deleuil, ex E. Morr. l. c. 329.
Grayii, Mort. l. c. 161 = C. Grayi.
Greeni, L. De Smet, ex Mort. l. c.—Afr. austr.
Hookeri, Lem. Illustr. Hortic. x. (1863) Misc. 80.—Mexic.
imbricata x, Deleuil, ex E. Morr. in Belg. Hortic. xxiv. (1874) 329.
imbricatissima, L. De Smet, in Rev. Hortic. iii. (1877) 147.—Hab. ?
lanceolata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 561 = C. caespitosa.
lanceolata, Torr. in Pacif. Rail. Rep. iv. 89, partim = C. farinosa.
laxa, Lindl. in Journ. Hort. Soc. iv. (1849) 292 = C. caespitosa.
linguæfolia, Lem. Illustr. Hortic. x. (1863) Misc. 81 = C. linguæfolia.
lucida, Steud Nom. ed. II. i. 585 = C. lurida.
lurida, Haw. in Tayl. Phil. Mag. (1831) 418 = C. lurida.
macrophylla, Hort. ex E. Morr. in Belg. Hortic. xxiv. (1874) 163, nomen.—Hab. ?
metallica, Hort. Kew. ex Lem. Illustr. Hortic. x. (1863) Misc. 81 = C. gibbiflora.
mirabilis x, Deleuil, ex E. Morr. in Belg. Hortic. xxv. (1875) 870.
misera, L. De Smet, ex E. Morr. l. c. xxiv. (1874) 162. —Hab. ?
mucronata, Schlecht. in Linnæa, xiii. (1839) 411 = C. mucronata, *Baker*.
mutabilis x, Deleuil, ex E. Morr. in Belg. Hortic. xxiv. (1874) 330.
navicularis, L. De Smet, ex E. Morr. l. c. 163.—Hab. ?
nodulosa, Otto, in Hamb. Gartenz. xxix. (1873) 8 = C. nodulosa.
nuda, Baker, in Gard. Chron. (1888) 280 = C. nuda.
ovata x, Deleuil, ex E. Morr. in Belg. Hortic. xxiv. (1874) 330.
pachyphytoides x, L. De Smet, ex E. Morr. l. c. 163.

ECHEVERIA :—
Pachyphytum, E. Morr. in Belg. Hortic. xxiv. (1874) 157 = Cotyledon Pachyphytum.
paniculata, A. Gray, Pl. Wright. i. 76 = C. Grayii.
paniculata, Moç. & Sesse, ex DC. Prod. iii. 350 = Fouquiera spinosa.
Pascuchii, Baker, in Gard. Chron. (1874) ii. 258 = C. Pascuchii.
peruviana, Meyen, Reise, i. 448 = C. peruviana.
Pfersdorffii, Hort. ex E. Morr. in Belg Hortic. xxiv. (1874) 163.—Hab. ?
prunosa x, E. Morr. l. c.
pubescens, Schlecht. in Linnæa, xiii. (1839) 411 = C. pubescens.
pulverulenta, Nutt. ex Torr. & Gray, Fl. N. Am. i. 560 = C. pulverulenta.
pumila, Van Houtte, Cat. (1846) ; Schlecht. Hort. Hal. 20 = C. pumila.
quitensis, Lindl. in Journ. Hort. Soc. vii. (1852) 268 = C. quitensis.
racemosa, Cham. & Schlecht. in Linnæa, v. (1830) 554 = C. lurida.
retusa, Lindl. in Journ. Hort. Soc. ii. (1847) 306 = C. retusa.
rhombifolia, Otto, in Hamb. Gartenz. xxxix. (1873) 9 = C. rhombifolia.
rosacea, Lind. & André, Illustr. Hortic. xx. (1873) 64. t. 234 = C. secunda.
rosæformis, L. De Smet, ex E. Morr. in Belg. Hortic. xxv. (1875) 100.—Hab. ?
rosea, Lindl. Bot. Reg. (1842) t. 22 = C. rosacea.
sanguinea, Hort. ex E. Morr. in Belg. Hortic. xxiv. (1874) 156 = C. atropurpurea.
scaphylla x, Deleuil, ex E. Morr. l. c. xxii. (1872) 326.
Scheerii, Lindl. Bot. Reg. (1845) t. 27 = C. Sheerii.
Schiedeckeri, L. De Smet, in Rev. Hortic. Belg. iii. (1877) 43.—Hab. ?
secunda, Booth, ex Lindl. Bot. Reg. (1838) Misc. 59 = C. secunda.
securifera x, Deleuil, ex E. Morr. in Belg. Hortic. xxv. (1875) 870.
spathulata, Bull, ex E. Morr. l. c. xxiv. (1874) 176 = Kalanchoë spathulata.
spathulifolia, L. De Smet, ex E. Morr. l. c. 167.—Hab. ?
spicata, Moç. & Sesse, in DC. Prod. iii. 349 = Fouquiria formosa.
spilota, Kunze, ex Walp. Ann. v. 12 = C. secunda.
spiralis x, Deleuil, ex E. Morr. in Belg. Hortic. xxv. (1875) 871.
Sprengelii, E. Morr. l. c. xxiv. (1874) 167 = C. Sprucei.
stellata x, Deleuil, ex E. Morr. l. c. xxv. (1875) 871.
stolonifera, Otto, in Hamb. Gartenz. xxix. (1873) 9 = C. stolonifera.
strictifiora, A. Gray, Pl. Wright. i. 76 = C. strictiflora.
subspicata, E. Morr. in Belg. Hortic. xxiv. (1874) 168 = C. subspicata.
subulifolia, E. Morr. l. c. = C. subulifolia.
teretifolia, DC. Prod. iii. 401 = C. subulifolia.
tortuosa, Hort. ex E. Morr. in Belg. Hortic. xxiv. (1874) 168, nomen.—Hab. ?
undulata x, L. De Smet, ex E. Morr. l. p. = C. undulata.
Van-Celsii, L. De Smet, ex E. Morr. l. c., nomen.—Hab. ?
Verwilrii, Hort. ex E. Morr. l. c., nomen.—Hab. ?
villosa, L. De Smet, ex E. Morr. l. c., nomen.—Hab. ?
yuccoides, Hort. ex E. Morr. l. c. = C. agavoides.

ECHEVERRIA, Lindl. & Paxt. Flow. Gard. iii. (1852–58) 1 = prec.

ECHIDIOCARYA, A. Gray, ex Benth. & Hook. f. Gen. ii. 854 (1876). *BORAGINEÆ*, Benth. & Hook. f. l. c.
arizonica, *A. Gray, in Proc. Am. Acad.* xi. (1876) 89. —Arizona.
californica, A. Gray, l. c. xii. (1877) 164 = Plagiobothrys Cooperi.
ursina, A. Gray, l. c. xix. (1888) 90 = Plagiobothrys ursinus.

ECHIDNIUM, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 62 = **Dracontium**, Linn. (Arold.).
Regelianum, Engl. in DC. Monog. Phan. ii. 286.—Am. trop.

ECHIDNIUM :—
Schomburgkii, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 62.—Guiana.
Spruceanum, Regel, Gartenfl. (1866) 98. t. 503 = *Regelianum*.
Spruceanum, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 350 = Cyrtosperma Spruceanum.

ECHIDNOPSIS, Hook. f. Bot. Mag. t. 5930 (1871). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 781.
cereiformis, *Hook. f. l. c.*—Afr. austr.

ECHINACANTHUS, Nees, in Wall. Pl. As. Rar. iii 75, 90 (1832). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1080.
Andersoni, *C. B Clarke, in Hook. f. Fl. Brit. Ind.* iv. 415.—Reg. Himal.
attenuatus, *Nees, in Wall. Pl. As. Rar.* iii. 90.—Reg. Himal.
Brugmansianus, *Nees, in DC. Prod.* xi. 168.—Ind. or. *calycinus*, Nees, l. c. = Strobilanthes helictus.
longistylus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 416.—Reg. Himal.
Lyallianus, Nees, in DC. Prod. xi. 168 = Nomaphila Lyallinus.
madagascariensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 218.—Madag.
parviflorus, *T. Anders. l. c.* ix. (1867) 459.—Reg. Himal.
Pumilio, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 415.—Reg. Himal.

ECHINACEA, Moench, Meth. 691 (1794). *COMPOSITÆ*. [Benth. & Hook. f. ii. 366.]
BRAUNERIA, Neck. Elem. i. 37 (1790).
HELICHROA, Rafin. Neogenyt. 8 (1825).
LEPACHIS, Rafin. in Am. Monthly Mag. (1819) 195.
angustifolia, *DC. Prod.* v. 554.—Am. bor.
atrorubens, Nutt in Trans. Am. Phil. Soc. N. S. vii. (1841) 354 = Rudbeckia atrorubens.
Dicksoni, *Lindl. Bot. Reg.* (1838) t. 27.—Mexic.
dubia, *Knowles & Westc. Fl. Cab.* iii. (1840) 163.—Mexic.
heterophylla, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. 93 = Isotephana heterophylla.
intermedia, Lindl. ex Paxt. Mag. Bot. xv. (1849) 79 = purpurea.
napifolia, Sweet, Hort. Brit. ed. II. 810 = Rudbeckia napifolia.
pallida, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 354 = angustifolia.
purpurea, *Moench, Meth.* 693.—Am. bor.
sanguinea, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 354 = angustifolia.
serotina, DC. Prod. v. 554 = purpurea.
speciosa, Paxt. Mag. Bot. xv. (1849) 79 = Rudbeckia speciosa.

ECHINAIS, C. Koch, in Linnæa, xxiv. (1851) 408 = ECHENAIS, Cass. = **Onious**, Linn. (Compos.).

ECHINALYSIUM, Trin. Fund. Agrost. 142 (1820) = **Elytrophorus**, Beauv. (Gramin.).
articulatum, Kunth, Enum. Pl. i. 391, sphalm. = seq.
strictum, Trin. Fund. Agrost. 142 = Elyt. articulatus.

ECHINANTHUS, Cerv. in La Naturaleza (1870), 351. *GRAMINEÆ*.

ECHINANTHUS, Neck. Elem. i. 91 (1790) = **Echinops**, Linn. (Compos.).

ECHINARIA, Desf. Fl. Atlant. ii. 385 (1799). *GRAMINEÆ*, Benth. & Hook. f. iii. 1181.
PANICASTRELLA, Moench, Meth. 205 (1794), nomen prius.
capitata, *Desf. l. c.*—Reg. Mediterr.
pumila, Willk. in Bot. Zeit. vi. (1848) 415 = capitata.

ECHINOCACTUS, Link & Otto, in Verh. Preuss. Gartenb. Ver. iii. (1827) 420. t. 18. *CACTEÆ*, Benth. & Hook. f. i. 848.
ASTROPHYTUM, Lemaire, Cact. Gen. Nov. et Sp. 9 (1839).
GYMNOCALYCIUM, Pfeiff. Cat. Hort. Schelh. ex Pfeiff. & Otto, Abbild. Cacteen, ii. t. 19 (1844).
MALACOCARPUS, Salm-Dyck, Cact. Hort. Dyck. ed. II. 24, 141 (1850).

ECHINOCACTUS:—
Acanthion, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 65 = exsculptus.
Acanthion, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 81, 161.—Hab.?
acanthodes, *Lem. Cact. Nov. Gen. et Sp.* 106.—Calif.
acanthostephes, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 102 = Mammillaria acanthostephes.
aciculatus, *Salm-Dyck, Hort. Dyck.* 341.—Bras.
acifer, Hopf. ex Foerst. Handb. Cact. ed. I. 530; Labour. Cact. 329 = spinosus.
acrocanthus, *Stieber, in Bot. Zeit.* v. (1847) 491.—Mexic.
acuatus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 424.—Reg. Argent.
acutangulus, Zucc. ex Pfeiff. Enum. Diagn. Cact. 55 = corynodes.
acutissimus, *Otto & Dietr. in Allg. Gartenz.* iii. (1835) 358.—Chili.
adversispinus, *Muehlenpf. in Otto & Dietr l. c.* xvi. (1846) 10.—Mexic.
agglomeratus, Hort. ex Pfeiff. Enum. Diagn. Cact. 60 = spiralis.
agglomeratus, Karw. ex Pfeiff. l. c. 61 = robustus.
albatus, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 170.—Mexic.
Allardtianus, *A. Dietr. l. c.* xv. (1847) 178.—Mexic.
ancylacanthus, Monv. ex Labour. Monog. Cact. 201 = uncinatus.
anfractuosus, *Mart. ex Pfeiff. Enum. Diagn. Cact.* 63.—Mexic.
arachnoideus, *Scheidw. in Bull. Acad. Brux.* vi. (1839) 90.—Mexic.
araneifer, Lem. ex Labour. Monog. Cact. 248 = Odieri.
aranealarius, *Reichb. in Terschech, Suppl.* 2; *ex Walp. Rep.* ii. 317.—Reg. Argent.
arcuatus, G. Don, Gen. Syst. iii. 168, sphalm. = acuatus.
armatissimus, *Foerst. in Hamb. Gartenz.* xvii. (1861) 162.—Peruv.; N. Granat.
armatus, Salm-Dyck, Hort. Dyck. 341 = Melocactus ferox.
arrectus, Otto, ex Labour. Monog. Cact. 225 = *arrigens*.
arrigens, Link, ex A. Dietr. in Otto & Dietr. Allg. Gartenz. viii. (1840) 161 = criaspatus.
Asterias, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. 11. (1848) 13. t. 3.—Mexic.
aulacogonus, *Lem. Cact. Gen. Nov. et Sp.* 14.—Mexic.
auratus, *Pfeiff. Abbild. Cacteen*, ii. sub t. 14.—Chili.
aureus, Meyen, ex Pfeiff. Enum. Diagn. Cact. 68.—Peruv.
Berteril, *C. Gay, Fl. Chil.* iii. 18.—Chili.
biceras, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xvi. (1846) 370.—Mexic.
bicolor, *Gal. ex Pfeiff. Abbild. Cacteen*, ii. t. 25.—Mexic.
bolivianus, Pfeiff. l. c. sub t. 14 = echinoides.
Bonthilleri, Parm. ex Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 314 = Cereus Schelhasii.
brachiatus, Labour. Monog. Cact. 236 = brachycentrus.
brachycentrus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 91. 160.—Mexic.
brevihamatus, *Engelm. in Proc. Am. Acad.* iii. (1857) 271.—N. Mexic.
brevimammeus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 102 = Mammillaria brevimamma.
Bridgesii, *Pfeiff. Abbild. Cacteen*, ii. t. 14.—Bolivia.
Buckii, *Klein, in Regel, Gartenfl.* (1859) 257. t. 266.—Mexic.
caespitiflus, Pfeiff. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 85 = horridus.
californicus, Hort. ex Foerst. Handb. Cact. ed. II. 472 = viridescens.
californicus, *Mowe. Cat.* (1846); *Cels, ex Labour. Monog. Cact.* 198.—Calif.
campylacanthus, Scheidw. in Otto & Dietr. Allg. Gartenz. viii. (1840) 337 = macrodiscus.
candicans, Hort. ex Pfeiff. Enum. Diagn. Cact. 91 = Cereus candicans.
capricornis, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 274.—Mexic.
castaneoides, *Cels, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 166,—Chili.

ECHINOCACTUS:—
centeterius, *Lehm. ex Pfeiff. Enum. Diagn. Cact.* 65. —Bras.
cephalophorus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 102 = Mammillaria cephalophora.
Ceratites, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 51.—Chili.
Ceratitis, Salm-Dyck, Cact Hort. Dyck. ed. II. 84, 167 = prasc.
cereiformis, *DC. in Mém. Mus. Par.* xvii. (1828) 115. —Mexic.
chlorophthalmus, *Hook. Bot. Mag.* t. 4486.—Mexic.
cinerascens, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 387.—Chili.
cinereus, *Phil. Fl. Atac.* 28.—Chili.
cinnabarinus, Hook. Bot. Mag. t. 4926 (= *Echinopsis cinnabarina*).—Bolivia.
Clava, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 125 = Mammillaria Clava.
coccineus, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 80 (= *Echinopsis rhodacantha*).—Reg. Argent.
columnaris, *Pfeiff. Abbild. Cacteen*, ii. sub t. 14.—Chili.
concinnus, *Hook. Bot. Mag.* t. 4115.—Hab.?
confertus, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 846, nomen.—Hab.?
conglomeratus, *Phil. Fl. Atac.* 28.—Chili.
conoideus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 107 = Mammillaria conoides.
conothelos, *Regel & Klein, Ind. Sem. Hort. Petrop.* (1860) 48.—Mexic.
conquadra, Hort. ex Foerst. Handb. Cact. ed. I. 338 = corynodes.
copiapensis, Pfeiff. Abbild. Cacteen, ii. sub t. 14 = cinerascens.
coptonogonus, *Lem. Cact. Aliq. Nov. Desc.* 23.—Mexic.
corracanthus, *Schridw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 50.—Mexic.
corniferus, Poselg. l. c. xxi. (1858) 102 = Mammillar a cornifera.
cornigerus, *DC. in Mém. Mus. Par.* xvi. (1828) 36. t. 7.—Am. centr.
corrugatus, *Steud. Nom.* ed. II. i. 536.—Chili.
corynodes, *Otto, in Otto & Dietr. Allg. Gartenz.* i. (1833) 364, nomen; *Pfeiff. Enum. Diagn. Cact.* 55. —Reg. Argent.
Coulteri, G. Don, Gen. Syst. iii. 162 = Histrix.
Covanillesii, *Lem. Cact. Aliq. Nov. Desc.* 20.—Bras.
crenatus, Hort. Gallic. ex Pfeiff. Enum. Diagn. Cact. 151 = exsculptus.
crierasus, Lem. ex Labour. Monog. Cact. 178 = latirisetus, *Salm-Dyck.*
crispatus, *DC. in Mém. Mus. Par.* xvii. (1828) 37. t. 8. —Mexic.
Cumingi, *Regel & Klein, Ind. Sem. Hort. Petrop.* (1860) 48.—Chili.
Cumingii, *Hopf. in Otto & Dietr. Allg. Gartenz.* xi. (1843) 222.—Peruv.
Cummingii, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 174.—Hab.?
curvicornis, Miq. in Linnaea, xii. (1838) 5 = spiralis.
curvispinus, *C. Gay, Fl. Chil.* iii. 16.—Chili.
cylindraceus, *Engelm. in Pacif. Rail. Rep.* iv. 32.—Calif.
cylindricus, *Hort. ex Forbes, Hort. Tour Germ.* 152, nomen.—Mexic.
detiliatipinus, *A. Berg, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 131.—Mexic.
Decaisnei, *Steud. Nom.* ed. II. i. 536.—Hab.?
denatus, Stend. l. c. sphalm. = Mammillaria denas?
denudatus, *Link & Otto, lc. Pl. Rar.* 17. t. 9.—Bras.
Deppii, *Link & Otto, ex Steud. Nom.* ed. II. i. 536, nomen.—Mexic.
depressus, *DC. Prod.* iii. 468.—Am. austr.
dichroscanthus, *Mart. ex Pfeiff. Enum. Diagn. Cact.* 62.—Mexic.
dicracanthus, *Hort. ex Forbes, Hort. Tour Germ.* 160, nomen.—Hab.?
Dietrichianus, *Foerst. in Hamb. Gartenz.* xvii. (1861) 160.—Hab.?
Dietrichii, Heynh. Nom. ii. 92 = lanciter.
dolichacanthus, Lem. Cact. Aliq. Nov. Desc. 25 = E. Echidne.
dolichocentrus, Salm-Dyck, Cact. Hort. Dyck. ed. I. 23 = prasc.
Dumeilianus, Cels, ex Salm-Dyck, l. c. ed. II. 30 = Cereus auratus.

ECHINOCACTUS:—
ebenacanthus, *Mowe. ex Labour. Monog. Cact.* 258.—Chili.
Echidne, *DC. Mém. Cact.* 19. t. 11.—Mexic.
echinatus, *Hort. ex Forbes, Hort. Tour Germ.* 152, nomen.—Mexic.
echinoides, *Lem. ex Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 386.—Bolivia.
edulis, *Haage, ex Foerst. Handb. Cact.* ed. I. 346, nomen.—Hab.?
Ehrenbergii, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* vii. (1838) 275.—Mexic.
electracanthus, *Lem. Cact. Aliq. Nov. Desc.* 24.—Mexic.
elegans, Hort. ex Pfeiff. Enum. Diagn. Cact. 86 = Cereus chilensis.
elephantidens, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 102 = Mammillaria elephantidens.
Ellemeetii, *Miq. in Nederl. Kruidk. Arch.* iv. (1859) 337.—Mexic.
elliptica, Lem. Jard. Fleur. iii. (1852–53) 270 = rhodophthalmus.
Emoryi, *Engelm. in Emory, Notes Mil. Reconnois.* 156.—Am. bor. occ.
emiferus, *Lem. Cact. Aliq. Nov. Desc.* 26.—Mexic
equitans, Scheidw. in Bull. Acad. Brux. vi. (1839) 88 = horizonthalonius.
erectus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 126 = Mammillaria erecta.
erinacea, *Lem. Cact. Aliq. Nov. Desc.* 16.—Reg. Argent.
erinaceus, Sweet, Hort. Brit. ed. II. 226 = Cereus erinaceus.
exsculptus, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 65. —Am. austr.
Eyriesii, Turp. Obs. Cact. 58. t. 2 = Cereus Eyriesii.
farinosus, *Cels, ex Foerst. Handb. Cact.* ed. I. 288, nomen.—Hab.?
farnosus, Hort. ex Foerst. l. c. 296 = Cereus Beneckei.
fasciolaris, *Steud. Nom.* ed. II. i. 536.—Peruv.
Fischeri, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 59 = Melocactus ferox.
flavispinus, *Mennsh. in C. Koch & Fint. Wochenschr.* i. (1858) 28.—Mexic.
flavo-virens, Scheidw. in Otto & Dietr. Allg. Gartenz. ix. (1841) 80 = orthacanthus.
flexispinus, Engelm. in Wisliz. Tour N. Mexico, 111 = longihamatus.
flexispinus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 159. —Mexic.
flexuosus, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 347.—Mexic.
fluctuosus, *A. Dietr. l. c.* 154.—Mexic.
Foersteri, *Stieber, in Bot. Zeit.* v. (1847) 491.—Mexic.
foliosus, *Steud. Nom.* ed. II. i. 536.—Chili.
Forbesii, *Lehm. in Terschech, Suppl.* 2; *ex Walp. Rep.* ii. 319.—Hab.?
formosus, Hort. Angl. ex Pfeiff. Enum. Diagn. Cact. 80 (= *Echinopsis formosa*).—Chili.
fossulatus, Scheidw. in Otto & Dietr. Allg. Gartenz. ix. (1841) 49 = hexaedrophorus.
fotevalana, Haage, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 88 = exsculptus.
fuscus, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xvi. (1846) 10.—Chili.
Galeottii, *Schieden. l. c.* ix. (1841) 50.—Mexic.
Gayonus, Hort. Par. ex Lem. Cact. Gen. et Sp. 22 = acutissimus.
germanthus, Hort. ex Otto, in Otto & Dietr. Allg. Gartenz. i. (1833) 365 = Cereus Eyriesii.
Ghiesbrechtii, *Salm-Dyck, l. c.* xviii. (1850) 395.—Mexic.
Ghiesbreghtianus, *Lem. Hort. Univ.* 227. t. 5.—Mexic.
gibbosus, *DC. Prod.* iii. 461.—Mexic.
Gigas, *Pfeiff. ex Foerst. Handb. Cact.* ed. I. 347, nomen.—Hab.?
Gillisii, Hort. ex Forbes, Hort. Tour Germ. 159, nomen = formosus.
gilseu, A. Dietr. in Otto & Dietr. Allg. Gartenz. xiii. (1846) 170 = E. Echidne.
gladiatus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 426.—Mexic.
glanduligerus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 102 = Mammillaria glanduligera.
glaucescens, *DC. in Mém. Mus. Par.* xvii. (1828) 115. —Mexic.
glaucus, Karw. ex Pfeiff. Enum. Diagn. Cact. 57 = recurvus.

ECHINOCACTUS:—
gracilis, Hort. ex Foerst. Handb. Cact. ed. I. 304 = gracillimus.
gracillimus, *Lem. Cact. Gen. Nov. et Sp.* 24.—Hab.?
grandicornis, *Lem. l. c.* 30.—Mexic.
grusoniana, *Jacobi*, *in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 99.—Hab.?
guyannensis, Hort. Gall. ex Pfeiff. Enum. Diagn. Cact. 151 = exsculptus.
Haageanus, *Foerst. Handb. Cact.* ed. II. i. 469.—Peruv.
haematochroanthus, Hemsl. Biol. Centr. Am. Bot. i. 532 = hamatocanthus.
hamatocanthus, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 371.—Mexic.
hamatus, *Hort. ex Forbes*, *Hort. Tour Germ.* 152, nomen.—Reg. Argent.
hamatus, Muehlenpf. in Otto & Dietr. Allg. Gartenz. xvi. (1848) 18 = ectispinus.
hamulosus, *Regel*, *Ind. Sem. Hort. Petrop.* (1856) 34.—Mexic.
Hankeanus, Forst. Handb. Cact. ed. II. i. 471 = humilis, *Auct.*
hastatus, *Hopf. ex Foerst. Handb. Cact.* ed. I. 315, nomen.—Hab.?
Haynii, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 33, 165.—Peruv.
helianthodiscus, Lem. ex Salm-Dyck, l. c. ed. I. 17 = turbiniformis.
helophorus, *Lem. Cact. Gen. Nov. et Sp.* 12.—Mexic.
hemisosus, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 10.—Peruv. vel Boliv.
heteracanthus, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 345.—Mexic.
heteromorphus, Poselg. l. c. xxi. (1858) 126 = Mammillaria heteromorpha.
hexacanthus, *Muehlenpf. l. c.* xiv. (1846) 369.—Mexic.
hexaedrophorus, *Lem. Cact. Gen. Nov. et Sp.* 27.—Mexic.
hexaedrus, *Scheidw. in Bull. Acad. Brux.* vi. (1839) 89.—Mexic.
Heyderi, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 170.—Mexic.
histricacanthus, Hemsl. Biol. Centr. Am. Bot. i. 532 = hystricacanthus.
Histrix, *DC. in Mém. Mus. Par.* xvii. (1828) 115.—Mexic.
holopterus, Miq. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 27 = ornatus.
Hookeri, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 345.—Mexic.
horizontalis, Hort. ex Foerst. Handb. Cact. ed. I. 327 = horizonthalonius.
horizonthalonius, *Lem. Cact. Gen. Nov. et Sp.* 19.—Mexic.
horridus, *C. Gay, Fl. Chil.* iii. 15.—Chili.
horripilus, *Lem. ex Foerst. Handb. Cact.* ed. I. 283; *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 35, 178.—Mexic.
humilis, *Auct. ex Foerst Handb. Cact.* ed. II. i. 471.—Hab.?
humilis, *Phil. Fl. Atac.* 28.—Chili.
Huotii, Cels, Portef. Hort. (1847) 216; *ex Labour. Monog. Cact.* 301 (= *Echinopsis Huotii*).—Chili.
hybocentrus, *Lehm. ex Pfeiff. Enum. Diagn. Cact.* 65.—Bras.
hybogonus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 167.—Chili.
hystricacanthus, *Lem. Cact. Gen. Nov. et Sp.* 21.—Hab.?
hypocrateriformis, *Otto & Dietr. Allg. Gartenz.* vi. (1838) 169.—Reg. Argent.
hystrichacanthus, *Lem. Cact. Gen. Nov. et Sp.* 17.—Mexic.
hystrichocentrus, *A. Berg*, *in Otto & Dietr. Allg. Gartenz.* viii. (1840) 131.—Mexic.
hystrichoides, Monv. ex Labour. Monog. Cact. 215 = hystrichocentrus.
Hystrix, *Haw. in Phil. Mag.* (1830) 116.—Ind. occ.
Hystrix, Monv. ex Labour. Cact. 184 = E. Echidne.
infastus, *Gill. ex Steud. Nom.* ed. II. i. 836, nomen.—Chili.
ingens, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 54.—Mexic.
insculptus, *Scheidw. in Bull. Acad. Brux.* vi. (1839) 89.—Mexic.
interruptus, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 66 = exsculptus.

ECHINOCACTUS:—
interruptus, Scheidw. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 29 = coptonogonus.
intertextus, *Engelm. in Proc. Am. Acad.* iii. (1857) 277.—Mexic.
intertextus, *Phil. in Linnæa*, xxxiii. (1864–65) 81.—Chili.
intortus, *DC. Prod.* iii. 462.—Ind. occ.
intricatus, *Link & Otto*, *in Verh. Preuss. Gartenb. Ver.* iii. (1837) 498.—Reg. Argent.
intricatus, *Salm-Dyck*, *in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 387.—Hab.?
irroratus, *Scheidw. in Bull. Acad. Brux.* vi. (1839) 90.—Mexic.
islayensis, *Foerst. in Hamb. Gartenz.* xvii. (1861) 160.—Peruv.
Jenischianus, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii (1840) 406.—Am. austr.
Johnsoni, *Parry*, *ex Engelm. in Bot. King's Exp.* 117.—Ind. occ.
Jussieui, *Monv. ex Salm-Dyck, Cact. Hort. Dyck*, ed. II. 34, 170.—Chili.
Karwinskii, Zucc. ex Labour. Monog. Cact. 193 = ingens.
Karwinskii, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 50.—Mexic.
Kunzii, *Foerst. Handb. Cact.* ed. I. 298; *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 33, 167.—Chili.
lamellosus, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xv. (1847) 177.—Mexic.
lancifer, *A. Dietr. l. c.* vii. (1839) 154.—Mexic.
lancifer, Reichb. in Terscheck, Suppl. 2; ex Walp. Rep. ii. 920 = electracanthus.
Langsdorfi, *Link & Otto, Ic. Pl. Rar.* 79. t. 40.—Bras.
laticostatus, Engelm. & Bigel. in Pacif. Rail. Rep. iv. 32 = horizonthalonius.
lasiphinus, *Hemsl. Biol. Centr. Am. Bot.* i. 538.—Mexic.
latispinus, Hort. ex Foerst. Handb. Cact. ed. I. 318 = corrugatus.
Lecchii, G. Don, in Sweet, Hort. Brit. ed. III. 288 = Cereus Lecchei.
Lecontei, *Engelm. in Pacif. Rail. Rep.* iv. 29. t. 2. f. 3.—Mexic.
Leeanus, *Hook. Bot. Mag.* t. 4184.—Reg. Argent.
Lemairi, Monv. ex Lem. Cact. Alliq. Nov. Desc. 17 = Mammillaria Lemairei.
leucacanthus, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 66.—Mexic.
leucanthus, Gill. ex Salm-Dyck, Hort. Dyck. 341 = Cereus leucanthus.
leucodictyus, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 34 = gibbosus.
Lindmanni, Ilgelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 246 = texensis.
Lindleyi, *Foerst. in Hamb. Gartenz.* xvii. (1861) 162.—Peruv.
Linkeanus, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 298.—Mexic.
Linkii, *Lehm. ex Pfeiff. Enum. Diagn. Cact.* 48.—Mexic.
longihamatus, *Gal. ex Foerst. Handb. Cact.* ed. I. 321, nomen; *et ex Pfeiff. Abbild. Cacteen*, ii. t. 16.—Mexic.
lophothele, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 305.—Mexic.
loricatus, Poselg. l. c. xxi. (1858) 107 = Mammillaria loricata.
Mackianus, *Hook. Bot. Mag.* t. 3561.—Chili.
Maclaeni, Hemsl. Biol. Centr. Am. Bot. i. 534, sphalm. = Macleni.
macracanthus, *De Vriese, in Hore. & De Vriese, Tijdschr.* vi. (1839) 49.—Mexic.
macracanthus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 26, 143.—Bolivia.
macrocephalus, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 370.—Mexic.
macrodiscus, *Mart. in Nov. Act. Nat. Cur.* xvi. 1. (1833) 341. t. 26.—Mexic.
macromeris, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 102 = Mammillaria macromeria.
macrothele, Poselg. l. c. 125 = Mammillaria macrothele.
Macleni, *Salm-Dyck, ex Foerst. Handb. Cact.* ed. I. 288; *Labour. Monog. Cact.* 363.—Mexic.
Malletianus, *Lem. ex Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 387.—Chili.
mamillosus, Lem. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 19, nomen = mammulosus?

ECHINOCACTUS:—
mammillarioides, Hook. Bot. Mag. t. 3558 = Pfeiffleri, *Zucc.*
mammillifer, *Miq. in Linnæa*, xii. (1838) 8.—Mexic.
mammillosus, Lem. ex Foerst. Handb. Cact. ed. I. 298 = scop.
mammulosus, *Lem. Cact. Aliq. Nov. Desc.* 40.—Bras
marginatus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 386.—Chili.
melanacanthus, Cels, ex Labour. Monog. Cact. 174 = marginatus.
Meimsianus, Wegener, in Otto & Dietr. Allg. Gartenz. xii. (1844) 65 = tricuspidatus.
melocactiformis, *DC. in Mém. Mus. Par.* xvii. (1828) 38. t. 10.—Mexic.
melocactoides, *Lem. Cact. Aliq. Nov. Desc.* 28.—Hab.?
meonacanthus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1837) 428.—Jamaic.
micracanthus, *Fennel, ex Foerst. Handb. Cact.* ed. I. 347, nomen.—Hab.?
minax, Lem. Cact. Aliq. Nov. Desc. 18 = platyceras.
Mirbelii, Lem. l. c. 22 = ornatus.
montevidensis, *G. Don, in Sweet, Hort. Brit.* ed. III. 288, nomen.—Montevideo.
montevidensis, Hort. Dresd. ex Foerst. Handb. Cact. ed. I. 292 = exsculptus.
Monvillii, *Lem. Cact. Aliq. Nov. Desc.* 14.—Paraguay.
Muehlenpfordtii, Fennel, in Otto & Dietr. Allg. Gartenz. xv. (1847) 65 = setispinus.
Muehlenpfordtii, Poselg. l. c. xxi. (1858) 102 = Mammillaria Schereti.
multiflorus, *Hook. Bot. Mag.* t. 4181.—Mexic.
multiplex, Hort. ex Pfeiff. Enum. Diagn. Cact. 70 = Cereus multiplex.
muricatus, Hort. ex Pfeiff. l. c. 49 = tortuosus.
muricatus, *Link & Otto ex Pfeiff. l. c.*—Bras.
mutabilis, *Foerst. in Hamb. Gartenz.* xvii. (1861) 161.—Peruv.
myriostigma, *Salm-Dyck, Cact. Hort. Dyck.* ed. I. 22; ed. II. 24, 155.—Mexic.
napinus, *Phil. in Anal. Univ. Chil.* (1872) 720; *et in Regel, Gartenfl.* (1872) 129.—Chili.
Netrelsianus, *Monv. ex Labour. Monog. Cact.* 248.—Hab.?
Neumannianus, Cels, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 38 = Kunzii.
niger, Hort. ex Salm-Dyck, l. c. 34 = Jussieui.
nobilis, Haw. in Phil Mag. (1830) 113 = gibbosus.
nodosus, *Hemsl. Biol. Centr. Am. Bot.* i. 535.—Mexic.
nummularioides, Steud. Nom. ed. II. i. 536, sphalm. = Pfeiffleri.
obrepandus, Salm-Dyck, in Otto & Dietr. Allg. Gartenz. xiii. (1845) 386 (= *Echinopsis cristata*).—Bolivia.
obvallatus, *DC. in Mém. Mus. Par.* xvii. (1828) 37. t. 9.—Mexic.
obvallatus, Pfeiff. Abbild. Cacteen, ii. t. 22 = lancifer.
occultus, *Phil. Fl. Atac.* 28.—Chili.
ochroleucus, *Jacobi*, *in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 101.—Hab.?
octacanthus, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 10.—Mexic.
octogonus, G. Don, in Sweet, Hort. Brit. ed. III. 288 = Cereus oxygonus.
Odieri, *Lem. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 174.—Chili.
oligacanthus, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 31 = brachycentrus.
oligacanthus, *Mart. ex Pfeiff. Enum. Diagn. Cact.* 58.—Mexic.
Oreptilis, *Haage, ex Foerst. Handb. Cact.* ed. I. 347, nomen.—Hab.?
ornatus, *DC. in Mém. Mus. Par.* xvii. (1828) 114.—Mexic.
orthacanthus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1837) 497.—Mexic.
Ottonianus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 102 = Mammillaria Ottonis.
Ottonis, *Link & Otto, Ic. Pl. Rar.* 81. t. 16.—Mexic.
Ourselianus, Cels, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 34 = multiflorus.
oxyacanthus, *Hort. ex Forbes*, *Hort. Tour Germ.* 152, nomen.—Hab.?
oxygonus, Link & Otto, in Verh. Preuss. Gartenb. Ver. vi. (1839) 419. t. 1 = Cereus oxygonus.
oxypteras, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 57.—Mexic.
pachycentrus, Lehm. ex Pfeiff. l. c 66 = centeterius.

ECHINOCACTUS:—

pachycornis, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 371.—Mexic.

papyracanthus, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1860–68) 198.—Am. bor.

Parryi, *Engelm. in Proc. Am. Acad.* iii. (1857) 276.—Mexic.

parvispinus, *DC. Prod.* iii. 468.—Am. austr.

pectinatus, Scheidw. in Bull. Acad. Brux. v. (1838) 492 = Cereus pectinatus.

pectiniferus, *Lem. Cact. Gen. Nov. et Sp.* 25.—Mexic.

Pepinianus, Cels. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 44 = Cereus Pepinianus.

pentacanthus, Lem. Cact. Allq. Nov. Desc. 27 = anfractuosus.

Pentlandi, Hook. Bot. Mag. t. 4124 (= *Echinopsis Pentlandi*).—Peruv.

Pfeifferi, *Sents, in Otto & Dietr. Allg. Gartenz.* v. (1837) 242.—Mexic.

Pfeifferi, *Zucc. in Pfeiff. Enum. Diagn. Cact.* 58.—Mexic.

phyllacanthoides, Lem. Cact. Nov. Gen. et Sp. 28 = phyllacanthus.

phyllacanthus, *Mart. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 201.—Mexic.

phyllantoides, Lem. ex Labour. Monog. Cact. 211 = phyllacanthus.

pilispinus, Lem. ex Labour. l. c. 186 = pilosus.

pilosus, *Gal. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 148.—Mexic.

piscatisformis, Lehm. ex Foerst. Handb. Cact. ed. II. i. 449 = Discocactus insignis.

Plaschnickii, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1858) 125.—Mammillaria Plaschnickii.

platyacanthus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 428.—Mexic.

platycarpus, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 347, nomen.—Hab.?

platyorpelus, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 9.—Mexic.

platyceros, *Lem. Cact. Allq. Nov. Desc.* 19.—Mexic.

polyacanthus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 422.—Bras.

polyancistrus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 39.—Am. bor. occ.

polycephalus, *Engelm & Bigel. l. c.* 31.—Calif.

polycentrus, Lem. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 32 = orthacanthus.

polyrhaphis, *Pfeiff. ex Salm-Dyck, l. c.* ed. II. 83, 166.—Peruv.

porrectus, *Lem. Cact. Allq. Nov. Desc.* 17.—Mexic.

Poselgerianus, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 346.—Mexic.

Pottsianus, Poselg. l. c. xxii. (1853) 107 = Mammillaria Pottsii.

Pottsii, *Salm-Dyck, l. c.* xviii. (1850) 395.—Mexic.

Pottsii, Schur. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 85 = bicolor.

praegracanthus, *Foerst. in Handb. Gartenz.* xvii. (1861) 160.—Chili.

prulinosus, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 54.—Mexic.

Pseudo-cereus, *Meinsh. in C. Koch & Fint. Wochenschr.* i. (1858) 29.—Mexic.

pubispinus, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1863) 199.—Am. bor.

pulchellus, Mart. in Nov. Act. Nat. Cur. xvi. i. (1832) 342. t. 28. 2 = Cereus pulchellus.

pumilus, *Lem. Cact. Allq. Nov. Desc.* 21.—Hab.?

pulverulentus, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 9.—Bolivia.

pycnacanthus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 102 = Mammillaria pycnacantha.

pycnoxyphus, *Lem. Cact. Gen. Nov. et Sp.* 16.—Hab.?

pyramidalis, Hort. ex Pfeiff. Enum. Diagn. Cact. 86 = Cereus chilensis.

pyramidatus, *Foerst. in Hamb. Gartenz.* xvii. (1861) 162.—Peruv.

quadristatus, *Wegener, in Otto & Dietr. Allg. Gartenz.* xii. (1844) 66.—Mexic.

radiatus, Hort. Belg. ex Pfeiff. Enum. Diagn. Cact. 180 = Mammillaria pycnacantha.

radicans, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 107 = Mammillaria radians.

radicans, Poselg. l. c. = Mammillaria vivipara.

raphidacanthus, Poselg. l. c. 126 = Mammillaria raphidacantha.

ECHINOCACTUS:—

raphidacanthus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 160.—Hab.?

raphidocentrus, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 101.—Hab.?

recurvus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 426; *DC. Prod.* iii. 463.—Mexic.

Reichenbachii, *Terschech, Suppl.* 2; *ex Walp. Rep.* ii. 820.—Mexic.

retusus, *Scheidw. ex Foerst. Handb. Cact.* ed. I. 347, nomen.—Hab.?

rhodacanthus, Salm-Dyck, Hort. Dyck. 841 (= *Echinopsis rhodacantha*).—Reg. Argent.

rhodanthus, *Hort. ex Forbes, Hort. Tour Germ.* 151, nomen.—Hab.?

rhodophthalmus, *Hook. Bot. Mag.* t. 4486.—Mexic.

rincinensis, *Poselg. ex Foerst. Handb. Cact.* ed. II. i. 473.—Mexic.

robustus, Karw. ex Pfeiff. Enum. Diagn. Cact. 60 = spiralis.

robustus, *Otto, in Otto & Dietr. Allg. Gartenz.* i. (1833) 364, nomen; *Pfeiff. Enum. Diagn. Cact.* 61.—Mexic.

roseanus, Hort ex Pfeiff. l. c. 55 = corynodes.

rostratus, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 108.—Chili.

Saglionis, Cels, Portef. Hort. (1847) 180; ex Labour. Monog. Cact. 357 = hybogonus.

salinensis, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 106.—Calif.

Salm-Dyckianus, *Pfeiff. Abbild. Cacteen*, ii. sub t. 14.—Bolivia.

Salm-Dyckianus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 102 = Mammillaria Salm-Dyckiana.

Salmianus, Link & Otto, in Verh. Preuss. Gartenb. Ver. iii. (1827) 428 = Melocactus Salmianus.

Salmii, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 1.—Hab.?

salpingophorus, Pfeiff. ex Labour. Monog. Cact. 302 (= *Echinopsis salpingophora*).—Hab.?

saltillensis, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 101.—Mexic.

Sandillon, *C. Gay, Fl. Chil.* iii. 14.—Chili.

Sandillon, Hort. ex Foerst. Handb. Cact. ed. II. i. 573 = Cereus.

Scheerii, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 155.—Mexic.

Schlechtendalii, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 125 = Mammillaria Schlechtendalii.

sclerothrix, *Lehm. Del. Sem. Hort. Hamb.* (1838). —*Cf. Linnaea*, xiii. (1839) *Litt.* t. 101.—Mexic.

Scopa, *Link & Otto, Ic. Pl. Rar.* 81. t. 41.—Bras.

Sellowianus, Hort. ex Pfeiff. Enum. Diagn. Cact. 55 = corynodes.

Sellowianus, Pfeiff. Enum. Diagn. Cact. 55, sphalm. = Sellowii.

Sellowii, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 425.—Reg. Argent.

sensilliflorus, Hook. Bot. Mag. t. 3569 = Sellowii.

sessiliflorus, *Hort. ex Forbes, Hort. Tour Germ.* 152, nomen = seq.?

sessiliflorus, *Pfeiff. Enum. Diagn. Cact.* 56.—Hab.?

setispinus, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 246.—Texas.

setosus, *Hemsl. Biol. Centr. Am. Bot.* i. 587.—Mexic.

Sickmanni, *Lehm. ex Steud. Nom.* ed. II. i. 587, nomen. —Am. austr.

Sieboldtii, Hort. Berol. ex Foerst. Handb. Cact. ed. I. 347, nomen = praec.?

similis, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 107 = Mammillaria missouriensis.

Simpsoni, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1863) 197.—Am. bor.

sinuatus, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 345.—Texas; Mexic.

Smithii, *Muehlenpf. l. c.* xiv. (1846) 370.—Mexic.

solenacanthus, *Scheidw. l. c.* ix. (1841) 50.—Mexic.

sparsithacanthus, *Mart. ex Foerst. Handb. Cact.* ed. I. 344, nomen.—Mexic.

spectabilis, Hort. ex Pfeiff. Enum. Diagn. Cact. 61 = robustus.

sphaeroideus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 107 = Mammillaria sphaeoleata.

sphaerocephalus, *Muehlenpf. l. c.* xiv. (1846) 370.—Mexic.

Spina-Christi, Zucc. ex Pfeiff. Enum. Diagn. Cact. 59 = Melocactus feros.

ECHINOCACTUS:—

spinosus, *Wegener, in Otto & Dietr. Allg. Gartenz.* xii. (1844) 66.—Mexic.

spinosissimus, *Hort. ex Forbes, Hort. Tour Germ.* 152, nomen.—Hab.?

spiralis, *Karw. ex Pfeiff. Enum. Diagn. Cact.* 60.—Mexic.

Steinesii, Hook. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 27 = pilosus.

stellaria, Hemsl. Biol. Centr. Am. Bot. i. 538 = spiralis.

stellatus, Scheidw. in Otto & Dietr. Allg. Gartenz. viii (1840) 358 = spiralis.

strepitoculon, *Hook. Bot. Mag.* t. 4563.—Bolivia.

strobiliformis, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 107 = Mammillaria strobiliformis, *Scheer subgibbosus*, Haw. in Phil. Mag. (1831) 419 = exsculptus.

subgrandicornis, *Haage, ex Foerst. Handb. Cact.* ed. I. 347, nomen.—Hab.?

submamillosus, *Lem. Cact. Gen. Nov. et Sp.* 20.—Hab.?

subporrectus, Lem. Cact. Allq. Nov. Desc. 25 = leucacanthus.

subuliferus, Hort. ex Pfeiff. Enum. Diagn. Cact. 61 = robustus.

subuliferus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 427. t. 27.—Mexic.

sulcatus, Hort. ex Pfeiff. l. c. 70 = Cereus oxygonus.

sulcatus, *Link & Otto, ex Steud. Nom.* ed. II. i. 587, nomen.—Mexic.

sulcolanatus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 102 = Mammillaria sulco-lanata.

sulphureus, *A. Dietr. l. c.* xiii. (1845) 170.—Mexic.

supertextus, Pfeiff. Abbild. Cacteen, ii. sub t. 14 = Kunzii.

tenuiflorus, *Link, ex Salm-Dyck, Cact. Hort. Dyck.* ed. I. 30, nomen.—Hab.?

tenuispinus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 421.—Bras.

tephracanthus, *Link & Otto, l. c.* 422.—Bras.

Terscheckii, *Knicht. in Terscheck, Suppl.* 3; *ex Walp. Rep.* ii. 815.—Reg. Argent.

tetracanthus, *Lem. Cact. Allq. Nov. Desc.* 15.—Reg. Argent.

tetracentrus, *Lem. Cact. Gen. Nov. et Sp.* 31.—Mexic.

tetraxiphus, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed. I. 30, nomen.—Hab.?

texensis, *Hopf. in Otto & Dietr. Allg. Gartenz.* x. (1842) 297.—Texas.

thriacanthus, Lem. Cact. Gen. Nov. et Sp. 86 = Pfeifferi, *Zucc.*

thrionacanthus, Lem. Cact. Allq. Nov. Desc. 22 = Pfeifferi, *Zucc.*

thelephorus, *Hort. ex Forbes, Hort. Tour Germ.* 152, nomen.—Hab.?

theloideus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 395.—Mexic.

thrincogonos, Lem. Cact. Gen. Nov. et Sp. 22 = acutissimus.

tortuosus, *Link & Otto, Ic. Pl. Rar.* 29. t. 15.—Bras.

tortus, Scheidw. in Bull. Acad. Brux. v. (1838) 493 = ornatus.

Treculianus, *Labour. Monog. Cact.* 202.—Texas.

tribolacanthus, *Monv. Cat. Cact.* (1846); *ex Labour. Monog. Cact.* 221.—Mexic.

tricornis, Monv. ex A. Dietr. in Otto & Dietr. Allg. Gartenz. xiv. (1846) 208 = Discocactus alteolens.

tricuspidatus, Scheidw. l. c. ix. (1841) 51 = phyllacanthus.

tuberculatus, *Link & Otto, in Verh. Preuss. Gartenb. Ver.* iii. (1827) 425.—Mexic.

tuberisulcatus, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 108.—Chili.

tuberosus, Salm-Dyck, ex Foerst. Handb. Cact. ed. I. 287 = leucacanthus.

tubiflorus, Hort. Angl. ex Pfeiff. Enum. Diagn. Cact. 71; Hook. Bot. Mag. t. 3637.—Mexic.

tulensis, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 125.—Mexic.

turbinatus, Hort. ex Pfeiff. in Otto & Dietr. l. c. iii. (1835) 314 = Cereus turbinatus.

turbiniformis, *Pfeiff. l. c.* vi. (1838) 278.—Mexic.

uncinatus, *Engelm. ex Scheer, in Som. Bot. Voy. Herald*, 290; *Engelm. Cact. U. St. et Mex. Bound. Surv.* 20.—Texas.

ECHINOCACTUS:—
uncinatus, *Hopf. ex Forrst. Handb. Cact.* ed. I. 321, nomen; *Gal. in Pfeiff. Abbild. Cactoen*, ii. t. 18. —Mexic.
undulatus, *A. Distr. in Otto & Dietr. Allg. Gartens.* xii. (1844) 187.—Mexic.
wagnipinus, Engelm. in Wisliz. Tour North Mexico, 111 = intertextus, *Engelm.*
Valparaiso, Hort. ex Forrst. Handb. Cact. ed. I. 292 = exsculptus.
Vanderaeyi, Lem. Cact. Aliq. Nov. Desc. 20 = E. Echidne.
Van-Geertii, Hort. ex Forrst. Handb. Cact. ed. I. 378 = Cereus lamprochlorus.
Vargasii, *Regel & Klein, Ind. Sem. Hort. Petrop.* (1860) 48.—Mexic.
verutum, *Hort. Angl. ex Forrst. Handb. Cact.* ed. I. 344, nomen.—Hab. ?
villiferus, *Scheidw. ex Forrst. l. c.* 347, nomen – Hab ? *villosus*, Lem. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 18; ed. II. 88 = polyaspis.
viridescens, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 554. —Calif.
viridiflorus, Pritz. Jc. Ind. ii. 114 = Cereus viridiflorus.
Visnaga, Hook. in Ill. Lond. News (1846); ex Bot. Mag t. 4559 = ingens
wislizenii, Poselg. in Otto & Dietr. Allg. Gartens. xxi. (1858) 107 = Mammillaria wislizenii
Wegeneri, Salm-Dyck, Cact. Hort. Dyck. ed. II. 31, 160 = quadrinatus
Whippiei, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 28. t. 1.—Am. bor. occ.
Williamsii, *Lem. ex Salm-Dyck, in Otto & Dietr. Allg. Gartens.* xiii. (1845) 385.—Hab. ?
Winkleri, Poselg. l. c. xxi. (1853) 102 = Mammillaria Winkleri.
Wippermanni, Muehlenpf. l. c. xiv. (1846) 370 = spinosus.
Wislizeni, *Engelm. in Wisliz Tour North Mexico*, 96. —Am. bor. occ.
xanthacanthus, Miq. in Linnaea, xi. (1837) 155. t. 4 = Melocactus xanthacanthus.
xiphacanthus, Miq. l. c. xii. (1838) 1 = arrigens.

ECHINOCALYX, Benth. in Benth. & Hook. f. Gen. i. 584 (1865) = **Sindora**, Miq. (Legumin.).

ECHINOCARPUS, Blume, Bijdr. 56 (1825). *TILIACEAE*, Benth. & Hook. f. i. 299.
asiaticus, *Benth. in Journ. Linn. Soc* v. *Suppl.* ii. (1861) 72.—Assam.
australis, *Benth. l. c.* 73.—Austral.
dasycarpus, *Benth. l. c.*—Reg. Himal.
Langii, F. Muell. Fragm. v. 28 = Sloanea Langii.
Mackrydei, F. Muell. l. c. vi. 170 = Sloanea Mackrydei.
Murex, *Benth. in Journ. Linn. Soc. v. Suppl.* ii. (1861) 72.—Reg. Himal.
Sigun, *Blume, Bijdr.* 56.—Java.
sinensis, *Hance, in Journ. Bot.* xxii. (1884) 108.—China.
sterculiaceus, *Benth. in Journ. Linn. Soc. v. Suppl.* ii. (1861) 72.—Reg. Himal.
tetragonus, *Teijsm. & Binn. Cat. Hort. Bog.* 184, nomen.—Java.
tomentosus, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 73.—Reg. Himal.
Woolsii, F. Muell. Fragm. vi. 171 = Sloanea Woollsii.

ECHINOCAULON, Spach, Hist. Vég. Phan. x. 521 (1841) = **Polygonum**, Linn.

ECHINOCAULOS, Hask. in Flora, xxv. (1842) ii. Beibl. 20 = **Polygonum**, Linn.
perfoliatus, Hask. l. c. = P. perfoliatum.

ECHINOCEPHALUM, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 294. *COMPOSITAE* (Benth. & Hook. f. ii. 377).
angustifolium, *Gardn. l. c.* 295.—Bras.
discoideum, *Baker, in Mart. Fl. Bras.* vi. iii. 280.—Bras.
lanceolatum, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 294.—Bras.
latifolia, *Gardn. l. c.* 295.—Bras.

ECHINOCEREUS, Engelm. in Wisliz. Tour North Mexico, 91 (1848) = **Cereus**, Haw. (Cact.).
acifer, Lem. Cact. 57, nomen = C. acifer.
adustus, Engelm. in Wisliz. Tour North Mexico, 104 = C. adustus.

ECHINOCEREUS:—
Berlandieri, Hort. ex Palm. in Rev. Horlic. (1865) 92 = C. Berlandieri.
Blankii, Palm. l. c. = C. Blankii.
caespitosus, Engelm. in Wisliz. Tour North Mexico, 110 = C. caespitosus.
candicans, Lem. Cact. 57, nomen = C. candicans.
chiotznis, Console & Lem. in Rev. Hortic. (1864) 173; Illustr. Horlic. xi. (1864) Misc. 15 = C. chiotznis.
chloranthus, Hasge, Preis-Verz. Cact. 22 = C. chloranthus.
cinerascens, Lem. Cact. 56, nomen = C. cinerascens.
cirrhiferus, Engelm. ex Hasge, Preis-Verz. Cact. 22 = C. cirrhiferus.
clavispinus, Hasge, l. c. = C. clavispinus.
coccinnus, Engelm. in Wisliz. Tour North Mexico, 93 = C. phoeniceus.
dasyacanthus, Engelm. l. c. 100 = C. dasyacanthus.
Ehrenbergii, Engelm. ex Hasge, Preis-Verz. Cact. 22 = C. Ehrenbergii
Engelmanni, Lem. Cact. 56, nomen = C. Engelmanni.
enneacanthus, Engelm. in Wisliz. Tour North Mexico, 111 = C. enneacanthus.
Fendleri, Engelm. ex Hasge, Preis-Verz. Cact. 22 = C. Fendleri
flavescens, Engelm. ex Hasge, l. c. = C. flavescens.
gladiatus, Engelm. ex Hasge, l. c. = C. candicans.
gonacanthus, Lem. Cact. 57, nomen = C. gonacanthus.
intricatus, Sencke, ex Hasge, Preis-Verz. Cact. 22 = C. intricatus.
Labouretianus, Lem. Cact. 57, nomen.—Hab ?
lamprochlorus, Engelm. ex Hasge, Preis-Verz. Cact. 22 = C. lamprochlorus.
Lecanus, Lem. Cact. 57, nomen = C. Lecanus.
longisetus, Lem. l. c. nomen = C. longisetus.
multangularis, Engelm. ex Hasge, Preis-Verz. Cact. 23 = C. multangularis.
pectinatus, Engelm. in Wisliz. Tour North Mexico, 110 = C. caespitosus.
peruvianus, A. Linke, in C. Koch & Flnt. Wochenschr. i. (1858) 85.—Peruv.
pentalophus, Lem. Cact. 56, nomen = C. pentalophus.
persistus, Foerst. in Hamb. Gartens. xvii. (1861) 168. —Peruv.
phoeniceus, Lem. Cact. 57, nomen = C. phoeniceus.
polyacanthus, Engelm. in Wisliz. Tour North Mexico, 104 = C. polyacanthus.
Poselgeri, Lem. Cact. 57, nomen.—Hab. ?
Poselgerianus, A. Linke, in Koch, Allg. Gartenz. xxv. (1857) 339.—Mexic.
Princeps, Foerst. in Hamb. Gartenz. xvii. (1861) 168.—Hab. ?
procumbens, Lem. Cact. 56, nomen = C. procumbens.
radians, Engelm. in Wisliz. Tour North Mexico, 104 = C. adustus.
raphidacanthus, Foerst. in Hamb. Gartenz. xvii. (1861) 164.—Hab. ?
Reichenbachianus, Engelm. ex Hasge, Preis-Verz. Cact. 23 = C. caespitosus.
rigidispinus, Engelm. ex Hasge, l. c. = C. rigidispinus, *Muehlenpf.*
Roemeri, Engelm. ex Hasge, l. c. = C. Roemeri, *Engelm.*
rotatus, A. Linke, in C. Koch & Flnt. Wochenschr. i. (1858) 85.—Texas.
ruflospinus, Engelm. in Wisliz. Tour North Mexico, 104 = C. ruflospinus.
Salm-Dyckianus, Scheer, in Seem. Bot. Voy. Herald, 291 = C. Salm-Dyckianus.
scheeri, Lem. Cact. 57, nomen = C. Scheeri.
serpentinus, Lem. l. c. nomen = C. serpentinus.
splendens, Lem. l. c. nomen = C. splendens.
strigosus, Lem. l. c. nomen = C. strigosus.
subinermis, Salm-Dyck, ex Scheer, in Seem. Bot. Voy. Herald, 291 = C. subinermis.
texensis, Jacobi, in Otto & Dietr. Allg. Gartenz. xxiv. (1858) 130.—Texas.
trichacanthus, Engelm. ex Hasge, Preis-Verz Cact. 23 = C. lutescens.
triglochidiatus, Engelm. in Wisliz. Tour North Mexico, 98 = C. triglochidiatus.
viridiflorus, Engelm. l. c. 91 = C. viridiflorus.

ECHINOCHLAENA, Spreng. Gen. i. 65 (1830) = **Echinolaena**, Desv. (Gramin.).

ECHINOCHLOA, Beauv. Agrost. 53. t. 11 (1812) = **Panicum**, Linn. (Gramin.).
colona, Link, Hort. Berol. ii. 209 = P. colonum.
commutata, Schult. Mant. ii. 267 = P. Crus-galli.
composita, Presl, ex Steud. Nom. ed. II. i. 587 = P. Crus-galli.
Crus-corvi, Beauv. Agrost. 53 = P. Crus-galli.
Crus-galli, Beauv. l. c. = P. Crus-galli.
Crus-pavonis, Schult. Mant. ii 269 = P. Crus-galli.
cubensis, Schult. l. c. 596 = P. cubense
divaricata, Anderss. in Peters, Reise Mossamb. Bot. 549.—Afr. trop.
dubia, Roem. & Schult. Syst. ii. 479 = Oplismenus dubius.
echinata, Beauv. Agrost. 53 = P. Crus-galli.
elephantipes, Nees, in Mart. Reise Bras. iii. 1123, nomen = P. elephantipes.
eruciformis, C. Koch, in Linnaea, xxi. (1848) 487 = P. eruciforme.
erythrosperma, Roem. & Schult. Syst. ii. 477 = Setaria italica.
frumentacea, Link, Hort. Berol. i. 204 = P. Crus-galli.
hirta, Schult. Mant. ii. 270 = P. Rothii.
hirtella, Schult. l. c. 269 = Oplismenus hollaceus.
hispida, Schult. l. c. 267 = P. Crus-galli.
Hostii, Link, Hort. Berol ii. 209 = P. Crus-galli.
intermedia, Roem. & Schult. Syst. ii. 477 = Setaria italica.
lanceolata, Beauv. Agrost. 53 = P. lanceolatum.
pulchella, Kunth, Enum. Pl. i. 78 = P. pulchellum.
scabra, Roem. & Schult. Syst. ii. 479 = P. Crus-galli.
setigera, Beauv. Agrost. 53 = P. setigerum.
setigera, Roem. & Schult. Syst. ii 480 = P. prostratum.
setosum, Beauv. Agrost 53 = Setaria setosa.
spectabile, Link, Hort. Berol. ii. 209 = P. spectabile.
squarrosa, Roem. & Schult. Syst. ii. 479 = P. squarrosum.
stagnina, Beauv. Agrost. 53 = P. Crus-galli.
subcordata, Roem. & Schult. Syst. ii. 480 = P. prostratum.
selayensis, Schult. Mant. ii. 269 = P. selayense.
Zenkowskius, Kegel, Ind. Sem Hort. Petrop. (1858); ex Schlecht. in Linnaea, xxvii. (1854) 477 = P. Zenkowskii.
senalis, Parl. Fl. Panorm. i. 119 = P. colonum.

ECHINOCROTON, F. Muell. Fragm. i. 31 (1858) = **Mallotus**, Lour. (Euphorb.).
claoxyloides, F. Muell. l. c. 32 = M. claoxyloides.

ECHINOCYSTIS, Torr. & Gray, Fl. N. Am. i. 542 (1840). *CUCURBITACEAE*, Benth. & Hook. f. i. 835.
ECHINOPEPON, Naud. in Ann. Sc. Nat. Sér. V. iv. (1865) 17.
HEXAMERIA, Torr. & Gray, in Torr. Rep. Pl. N. York, 187 (1889) (Geol. Surv.).
MARAH, Kellogg, in Proc. Calif. Acad. ii. (1863) 18; et i. ed. II. (1873) 87.
MEGARRHIZA, Torr. & Gray, in Pacif. Rail. Rep. vi. 74, nomen (1857); xii. ii. 61 (1860).
australis, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878); *(Diagn. Cucurb.* fasc. ii.) 98 = Cyclanthera australis.
arnesoea, *Griseb. in Goett. Abh.* xxiv. (1879) 135.—Reg. Argent.
Bigelovii, *Cogn. in DC. Monog. Phan.* iii. 804.—Calif.
Coulteri, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878) 88.—N. Mexic.
fabacea, A. Gray, in Proc. Am. Acad. viii. (1873) 384 = oregana.
fabacea, *Naud. in Ann. Sc. Nat. Sér.* IV. xii. (1859) 164.—Calif.
floribunda, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878) 89.—Mexic.
gemella, *Cogn. l. c.* 88.—Mexic.
gilensis, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 189. —N. Mexic.
glutinosa, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878) 93.—Mexic.
guadalupensis, *Cogn. in DC. Monog. Phan.* iii. 819.—Ins. Guadalup.
lanata, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878) 92.—Mexic.
lobata, *Torr. & Gray, Fl. N. Am.* i. 542.—Am. bor. occ.
longispina, *Cogn. l. c.* 91.—Mexic.
macrocarpa, *Greene, in Bull. Calif. Acad* i. iv. (1885) 188.—Calif.

ECHINOCYSTIS:—
Marsh, *Cogn. in DC. Monog. Phan.* iii. 517.—Calif.
millefiora, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878) 88.—Mexic.
minima, *Cogn. in DC. Monog. Phan.* iii. 505.—Calif.
muricata, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878) 88.—Bras.
muricata, Kellogg, in Proc. Calif. Acad. i. 57 = Watsonii.
oregana, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878) 87.—Am. bor. occ.
paniculata, *Cogn. l. c.* 90.—Mexic.
parviflora, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 373.—Am. bor. occ.
pedata, Scheele, in Linnaea, xxi. (1848) 587 = Cyclanthera dissecta.
polycarpa, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 90.—Venezuela.
pubescens, *Cogn. l. c.* 88.—Mexic.
torquata, *Cogn. l. c.*—Mexic.
Watsonii, *Cogn. in DC. Monog. Phan.* iii. 519.—Am. bor. occ.
Wrightii, *Cogn. in Mém. Cour. Acad. Belg.* 8vo, xxviii. (1878) 88.—N. Mexic.

ECHINODENDRUM, A. Rich. Fl. Cub. Fanerog. ii. 18 (1856) = **Scoloxanthus**, Vahl (Rubiac.).
campanulatum, A. Rich. l. c. = S. parviflorus.
parviflorum, A. Rich. l. c. t. 47* = S. parviflorus.

ECHINODISCUS, Benth. in Ann. Wien. Mus. ii. (1838) 94 = **Pterocarpus**, Linn. (Legumin.).
echinatus, Mkq. Fl. Ind. Bat. i. 1. 137 = P. echinatus.
erinaceus, Benth. ex Walp. Rep. i. 728 = P. erinaceus.

ECHINODIUM, Polt. ex Cass. in Dict. Sc. Nat. lix. 235 (1829) = **Acanthospermum**, Schrank (Compos.).

ECHINODORUS, Rich. in Mém. Mus. Par. i. (1815) 365. *ALISMACEAE*, Benth. & Hook. f. iii. 1006.
alpestris, *Micheli, in DC. Monog. Phan.* iii. 47.—Hispan.
brasiensis, *Micheli, l. c.* 59.—Panama.
cordifolius, Griseb. in Goett. Abh. vii. (1857) 257 = rostratus.
ellipticus, *Micheli, in DC. Monog. Phan.* iii. 51.—Am. trop.
floribundus, *Seub. ex Warm. in Kjoeb. Vidensk. Meddel.* (1872) 118.—Bras.
grandiflorus, Micheli, l. c. D. Monog. Phan. iii. 57 = floribundus.
guyanensis, Griseb. Fl. Brit. W. Ind. 505 = Sagittaria guyanensis.
humilis, *Buchen. in Pringsh. Jahrb.* vii. (1869) 28.—Afr. occ.
intermedius, *Griseb. Cat. Pl. Cub.* 218.—Bras.
longipetalus, *Micheli, in DC. Monog. Phan.* iii. 60.—Bras.
macrophyllus, Micheli, l. c. 50 = maricatus.
Martii, *Micheli, l. c.* 49.—Bras.
muricatus, *Griseb. in Bonplandia*, vi. (1858) 11.—Am. trop.
natans, Engelm. in Aschers. Fl. Brand. i. 651 = Elisma natans.
ovalis, *Wright, in Sauv. Fl. Cub.* 153.—Cuba.
paniculatus, *Micheli, in DC. Monog. Phan.* iii. 51.—Am. austr.
parnassifolius, Engelm. in Aschers. Fl. Brand. i. 651 = Alisma parnassifolium.
parvulus, Engelm. in A. Gray, Man. Bot. ed. II. 488 (1856), 492 = tenellus.
pubescens, Senth. ex Warm. in Kjoeb. Vidensk. Meddel. (1872) 118 = ellipticus.
punctatus, *Micheli, in DC. Monog. Phan.* iii. 59.—Bras.
radicans, *Engelm. in A. Gray, Man. Bot.* ed. I. 460; ed. II. 488.—Am. calid.
ranunculoides, *Engelm. in Aschers. Fl. Brand.* i. 651. —Europ. et Afr. bor.
rostratus, *Engelm. in A. Gray, Man. Bot.* ed. I. (1848) 460; ed. II. 488.—Am. bor.
subulatus, Engelm. l. c. = tenellus.
subulatus, *Griseb. Cat. Pl. Cub.* 218.—Guiana.
tenellus, *Buchen. in Bremen, Abh.* ii. (1871) 21.—Am. austr.
virgatus, *Micheli, in DC. Monog. Phan.* iii. 54.—Mexic.

ECHINOGLOSSUM, Reichb. Nom. 54 (1841) = ECHIOGLOSSUM, Blume = **Cleisostoma**, Blume (Orchid.).

ECHINOLAENA, Desv. Journ. Bot. i. (1813) 75. *GRAMINEAE*, Benth. & Hook. f. iii. 1107.
ECHINOCHLAENA, Spreng. Gen. i. 85 (1830).
brachystachya, Kunth, Rév. Gram. i. 54 = Panicum brachystachyum.
hirta, *Desv. Journ. Bot.* i. (1813) 75.—Am. austr.
loliacea, Kunth, Enum. Pl. i. 172 = Ichnanthus candicans.
madagascariensis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 452.—Madag.
naviculoris, Kunth, Rév. Gram. i. 54 = Panicum nemorale.
nemorosa, Kunth l. c. = Ichnanthus nemorosus.
polystachya, H. B. & K. Nov. Gen. et Sp. i. 119 = Panicum uncinatum.
procurrens, Kunth, Rév. Gram. i. 54 = Panicum procurrens.
scabra, H. B. & K. Nov. Gen. et Sp. i. 118. t. 38 = hirta.
Trinii, Zoll. & Moritzi, Syst. Verz. Zoll. 102 = Panicum uncinatum.

ECHINOLEMA, Jacq. f. ex DC. Prod. v. 8 (1836) = **Adiscaphe**, juss. (Calycer.).
armatum, Jacq. f. l. c. = A. spathulata.

ECHINOLITRUM, Steud. Nom. ed. II. i. 537 (1840) = ECHINOLYTRUM, Desv. = **Pimbristylis**, Vahl (Cyperac.).

ECHINOLOBIUM, Desv. Journ. Bot. i. (1813) 128. t. 5 = **Rudyserum**, Linn. (Legumin.).

ECHINOLOMA, Steud. Nom. ed. II. i. 537 (1840), sphalm. = ECHINOLEMA, Jacq. f. = **Adiscaphe**, Juss. (Calycer.).

ECHINOLYSIUM, Benth. in Journ. Linn. Soc. xix. (1881) 114 = ECHINALYSIUM, Trin. = **Elytrophorus**, Beauv. (Gramin.).

ECHINOLYTRUM, Desv. Journ. Bot. i. (1808) 20. t. 1 = **Pimbristylis**, Vahl (Cyperac.).
dipsaceum, Desf. l. c.—Geront. trop.

ECHINOMERIA, Nutt. in Trans. Am. Phil. Soc. vii. (1841) 356 = **Hellanthus**, Linn. (Compos.).
apetala, Nutt. l. c. = H. Radula.

ECHINONYCTANTHUS, Lem. Cact. Nov. Gen. et Sp. 10 (1839) = **Cereus**, Haw. (Cact.).
Decaisneanus, Lem. l. c. 55 = Echinocactus Decalsnei.
Eyriesii, Lem. l. c. 54 = C. Eyriesii.
leucanthus, Lem. l. c. 55 = C. leucanthus.
multiplex, Lem. l. c. = C. multiplex.
nigrispinus, Lem. l. c. (= *Echinopsis nigrispina*).—Hab.?
oxygonus, Lem. l. c. = C. oxygonus.
pictus, Lem. l. c. 54 (= *Echinopsis picta*).—Hab.?
pulchellus, Lem. l. c. 55 = C. pulchellus.
tubiflorus, Lem. l. c. = Echinocactus tubiflorus.
Schelhasii, Lem. l. c. 54 = C. Schelhasii.
turbinatus, Lem. l. c. = C. turbinatus.

ECHINOPANAX, Decne & Planch. in Rev. Hortic. (1854) 105 = **Fatsia**, Decne. & Planch. (Araliac.).
horridus, Decne. & Planch. l. c. = F. horrida.

ECHINOPEPON, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 17 = **Echinopoyatia**, Torr. (Cucurbit.).
glutinosus, Naud. ex Cogn. in DC. Monog. Phan. iii. 583 = Echinoc. glutinosa.
horridus, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 19 = Echinoc. Coulteri.
millefloras, Naud. l. c. 18 = Echinoc. millefiora.
quinquelobatus, Naud. l. c. = Echinoc. torquata.

ECHINOPHORA. Tourn. ex Linn. Syst. ed. II. 20 (1740). *UMBELLIFERAE*, Benth. & Hook. f. i. 861. APIOSCIADIUM, DC. Prod. iv. 224 (1830).
anatolica, *Boiss. & Heldr. Diagn.* Ser. I. x. 56.—As. Min.
cavrifolia, *Boiss. & Bal. l. c.* Ser. II. v. 104.—Cilicia.
caspica, DC. Prod. iv. 225 = Hippomaesthrum caspum.

ECHINOPHORA:—
Exacantha, D. Dietr. Syn. Pl. ii. 985 = Exacantha heterophylla.
flabellifolia, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 92 = Pycnocycla flabellifolia.
platyloba, *DC. Prod. iv.* 225.—Persia.
radiata, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 92.—As. Min.
Sibthorpiana, *Guss. Suppl. Fl. Sic. Prod.* 69.—Graec., Orient.
spinosa, *Linn. Sp. Pl.* 239.—Europ.; Reg. Mediterr
tenuifolia, *Linn l. c.*—Europ. austr.
tenuifolia, Sibth. & Sm. Fl. Graec. t. 266 = Sibthorpiana.
Tournefortii, Jaub. & Spach, Illustr. iii. 50, 241 = anatolica.
trichophylla, *Sm. in Rees, Cycl.* xii. n. 3.—Armen., Persia.

ECHINOPOGON, Beauv. Agrost. 42. t. 9. f. 5 (1812) *GRAMINEAE*, Benth. & Hook. f. iii. 1145.
HYSTERICINA, Steud. Syn. Pl. Gram. 35 (1855).
asper, Trin. Fund. Agrost. 126 = ovatus.
Gunnianus, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 418 = Deyeuxia Gunniana.
ovatus, *Beauv. Agrost.* 42. t. 9.—Austral.
Stoberi, Steud. Syn. Pl. Gram. 188 = ovatus.

ECHINOPS, Linn. Gen. ed. i. 242 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 462.
ECHINANTHUS, Neck. Elem. i. 91 (1790).
ECHINOPUS, Tourn. ex Adans. Fam. ii. 114 (1763)
SPHAEROCEPHALUS, Linn. Syst. ed. I (1735).
Acantholeptus, Jaub. & Spach, Illustr. iii. 99 = Acantholepis orientalis.
acanlis, *Lavy, Vig. Orig.* 11; *ex DC. Prod.* vi. 527.—Hab.?
adenocaulis, Boiss. Diagn. Ser. I. x. 88 = viscosus.
albicaulis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 387.—Soongaria.
albidus, Boiss. & Sprun. Diagn. Ser. I. vi. 97 = sphaerocephalus.
altaicus, Hort. ex DC. Prod. vi. 524 = sphaerocephalus.
amplexicaulis, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 101. t. 47.—Afr. trop.
armatus, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 486.—Assyria.
armatus, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 389 = bannaticus.
Aucheri, *Boiss. Diagn.* Ser. I. vi. 98.—Persia.
australis, Tenore, ex Nym. Comp. 399 = E. Ritro.
bannaticus, *Rochel, ex Schrad. Blumenb.* 48.—Bannat.; Tauria.
bithynicus, Boiss. Diagn. Ser. I. vi. 100 = viscosus.
Blanchianus, *Boiss. Fl. Orient.* iii. 480.—Syria.
Bovei, *Boiss. Diagn.* Ser. I. vi. 99.—Afr. bor.
candidus, Boiss. l. c. x. 86 = Cephalotes.
castaneus, *Munby, Cat. Pl. Algyr.* 17, nomen.—Algeria
Cephalotes, *DC. Prod.* vi. 526.—Persia.
ceratophorus, *Boiss. Diagn.* Ser. I. vi. 100.—Persia.
chamaecephalus, *Hochst. ex A. Rich. Tent. Fl. Abyss* i. 448.—Abyss.
chantavicus, *Trautv. in Bull. Soc. Nat. Mosc.* xxxix. (1866) i. 366.—As. centr.
Chardini, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 124.—Persia.
choetocephalus, *Pomel, Nouv. Mat. Fl. Atl.* 272.—Alger.
chorassanicus, *Bunge, in Bull. Acad. Pétersb.* vi. (1863) 405.—Persia.
coeruleus, *Onerin, ex Trautv. in Act. Hort. Petrop.* viii. (1883) 477, nomen.—Transcauc.
commutatus, *Turczaninow, in Verh. Zool. Bot. Ges. Wien*, viii. (1858) 17.—Europ. austr.
connatus, C. Koch, in Linnaea, xxiv. (1851) 330 = viscosus.
coriarius, C. B. Clarke, Comp. Ind. 213 = cornigerus.
corniger*us*, *DC. Prod.* vi. 526.—Ind. or.
corymbosus, Linn. Sp. Pl. 813 = Cardopatium corymbosum.
creticus, Boiss. & Heldr. Diagn. Ser. I. x. 87 = viscosus.
cynocephalus, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 484.—Persia.
daburicus, *Fisch. Cat. Hort. Gorensk.* ed. II. 37.—Dahur.; Mongol.
dasyanthus, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 47.—As. centr.

ECHINOPS:—
daucifolius, Trevir. Ind. Sem. Hort. Wratislaw. (1819); Ledeb. Fl. Ross. iii. 653 = humilis.
dichrous, *Boiss. & Heldr. in Boiss. Fl. Orient.* iii. 431.—Persia.
echinatus, *Roxb. Hort. Beng.* 62; *Fl. Ind.* iii. 447.—Ind. or.
echinophorus, Boiss. Diagn. Ser. I. x. 87 = spinosus.
eglandulosus, Schur, Enum. Pl. Transs. 426 = exaltatus.
elatus, Bunge, in Bull. Acad. Pétersb. vi. (1868) 407 = horridus.
elegans, Bertol. Fl. Ital. ix. 513 = E. Ritro.
eriophorus, Stev. ex DC. Prod. vi. 652 = humilis.
exaltatus, *Koch, Syn. Fl. Germ.* ed. II. 462 = commutatus.
exaltatus, *Schrad. Hort. Gotting.* ii. 15. t. 9.—Hungar.
fruticosus, Linn. Sp. Pl. 815 = Rolandra argentea.
Gaillardoti, *Boiss. Diagn.* Ser. II. iii. 38.—Syria.
giganteus, Hort. ex DC. Prod. vi. 624 = sphaerocephalus.
giganteus, *A. Rich. Tent. Fl. Abyss.* i. 449 —Abyss.
glabermimus, *DC. in Ann. Sc. Nat.* Sér. II. ii. (1834) 260.—Arab.
glandulosus, E. Weiss, in Verh. Zool. Bot. Ges. Wien, xviii. (1868) 438 = viscosus.
globifer, Janka, in Linnaea, xxx. (1859-60) 581 = exaltatus.
Gmelini, Ledeb. Fl. Alt. iv. 45 = dahuricus.
Gmelini, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 195.—Mongol.
graecus, *Milt. Gard. Dict.* ed. VIII. n. 4.—Graecia.
grandiflorus, Clarke, Trav. ii. 452 = graecus.
Griffithianus, Boiss. Diagn. Ser. II. iii. 39 = echinatus †
Grisebachii, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 221 = dahuricus.
Haussknechtii, *Boiss. Fl. Orient.* iii. 438.—Persia.
hebelepis, *DC. Prod.* vi. 624.—Persia.
Heldreichii, *Boiss. Diagn.* Ser. I. x. 84.—Aa. Min.
heteromorphus, *Bunge, in Bull. Acad. Pétersb.* vi. (1862) 406.—Persia.
hispidus, *Fresen. in Mus. Senckenb.* iii. (1845) 68.—Abyss.
horridus, *Desf. Tabl.* ed. I. 94.—Persia.
horridus, Lank. Enum. Hort. Berol. ii. 367 = sphaerocephalus.
humilis, *Bieb. Fl. Taur. Cauc. Suppl.* 508.—Sibir. altaic.
humilis, Reichb. Fl. Germ. Excurs. 292 = platylepis.
Hussoni, *Boiss. Diagn.* Ser. I. x. 86.—Arabia.
hypoleucus, *Rupr. Sert. Tianschan.* 58.—Turkestan.
illicifolius, *Bunge, in Bull. Acad. Pétersb.* vi. (1863) 408.—Persia.
inermis, *Boiss. & Hausskn. in Boiss. Fl. Orient.* iii. 436.—Persia.
integrifolius, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 446.—Sibiria.
jaxarticus, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 350 (*Lehm. Rel.* 174).—Turkestan.
jenlianus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 135.—Persia.
karatavicus, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 256.—Turkestan.
Kerneri, *Heimerl, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 61.—Persia.
Kotschyi, *Boiss. Diagn.* Ser. I. vi. 99.—Persia.
kurdicus, *Boiss. & Hausskn. ex Boiss. Fl. Orient* iii. 437 (= sec. ?).—Persia.
kurdicus, *Bunge, in Bull. Acad. Pétersb.* vi. (1863) 408.—Persia.
lanatus, Steph. ex DC. Prod. vi. 638 = humilis.
lanuginosus, Lam. Encyc. iii. 317 = graecus.
lanuginosus, Sibth. & Sm. Fl. Graec. t. 926 = microcephalus.
lasiolepis, Boiss. Diagn. Ser. I. x. 89 = Blanchemanus.
lasiolepis, Bunge, in Bull. Acad. Pétersb. vi. (1863) 408 = Cephalotes.
latifolius, Tausch, in Flora, xi. (1828) 486 = dahuricus.
leucographus, *Bunge, in Bull. Acad. Pétersb.* vi. (1863) 406.—Persia.
longifolius, *A. Rich. Tent. Fl. Abyss.* i. 452.—Abyss.
longisetus, *A. Rich. l. c.* 450.—Abyss.
macrochaetus, Boiss. Diagn. Ser. I. x. 85 = viscosus.
macrochaetus, *Fresen. in Mus. Senckenb.* iii. (1845) 69.—Abyss.
macrophyllus, *Boiss. & Hausskn. in Boiss. Fl. Orient.* iii. 436.—Persia.
marscandicus, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 351 (*Rel. Lehm.* 178).—Turkestan.

ECHINOPS:—
maximus, Siev. in Pall. N. Nord. Beitr. vii. 323 = sphaerocephalus.
Sibth. & Sm. Fl. Graec. Prod. ii. 209. —Graecia.
mollis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 84 = commutatus.
multiflorus, Lam. Fl. Fr. ii. 2 = sphaerocephalus.
nanus, *Bunge, in Bull. Acad. Pétersb.* vi. (1863) 411.—Hab.?
Neumayeri, *Vis. Fl. Dalmat.* ii. 95. t. 10. ter.—Europ.
niveus, *Wall. Cat.* n. 2686.—Reg. Himal. occ.
nodiflorus, Lam. Encyc. ii. 337 = Rolandra argentea.
Olivieri, Jaub. & Spach, Illustr. iii. 100 = Acantholepis orientalis.
orientalis, Trautv Diss. Echin. 22 = horridus.
ossicus, *C. Koch, in Linnaea*, xxiv. (1851) 380.—Reg. Caucas.
paniculatus, Jacq. f. Eclog. i. t. 9 = sphaerocephalus.
parviflorus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 124.—Persia.
paucyflorus, Lam. Fl. Fr. ii. 2 = E. Ritro.
persicus, *Stev. ex Fisch. Cat. Hort. Gorenk.* ed. II. (1812) 37.—Reg Caucas.; Persia.
platylepis, *Trautv. Diss. Echin.* 11. f. 1.—Hab.?
polycephalus, Gouan, cx Steud. Nom. ed. II. i. 587 = E. Ritro.
polycrasus, Boiss. Diagn. Ser. I. x. 85 = Blanchemanus.
polygamus, *Bunge, in Bull. Acad. Pétersb.* vi. (1863) 405.—Persia.
pratensis, Turcz. ex Ledeb. Fl. Ross. ii. 653 = dahuricus.
Prescottii, Fisch. & M.y. ex DC. Prod. vi. 625 = glabermimus.
pungens, Trautv. Diss. Echin. 18 = persicus.
Ritro, Georgi, Reise, i. 221 = dahuricus.
Ritro, Gueldenst. ex Ledeb. Fl. Ross. ii. 655 = sphaerocephalus.
Ritro, *Linn. Sp. Pl.* 815.—Reg. Mediterr.; Orient.; As. bor.
Ritro, Schkuhr, Handb. t. 268 = exaltatus.
ritrodes, *Bunge, in Bull. Acad. Pétersb.* vi. (1863) 407 (rytrodes).—Persia.
robustus, *Bunge, l. c.* 405 —Persia.
Rochelianus, *Griseb. Spicil. Fl. Rumel.* ii. 229.—Banat.
ruthenicus, Bess. Enum. Pl. Volh. [77 †]; ex DC. Prod. vi. 623 = dahuricus.
ruthenicus, Rieb. Fl. Taur. Cauc. iii. 597 = E. Ritro.
ruthenicus, Reichb. Ic. Pl. Crit. t. 450 = banaticus, sphaerocephalus.
ruthenicus, Kochel, Pl. Rar. Banat. 77. t. 87 = banaticus.
rytrodes, Bunge, in Bull. Acad. Pétersb. vi. (1863) 407 = ritrodes.
Sartoranus, Boiss. & Heldr. Diagn. Ser. I. vi. 97 = E. Ritro.
serratifolius, Sch. Bip. in Pollichia, xviii-xix. (1861) 186 = longifolius.
siculus, *P. G. Strobl, in Flora*, lxv. (1882) 505.—Sicilia.
sphaerocephalus, *Linn. Sp. Pl.* 814.—Europ.; Orient; As. bor.
sphaerocephalus, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 182 = dahuricus.
sphaerocephalus, Sibth. & Sm. Fl. Graec. t. 923 = viscosus.
spinosissimus, Turra, Farsat. Nov. Gen. 18 = spinosus.
spinosus, Bové, ex DC. Prod. vi. 625 = glabermimus.
spinosus, Herb. Madr. ex Wall. Cat. n. 2985 = echinatus.
spinosus, *Linn. Mant.* i. 119.—Creta; Afr. bor.
spinosus, Urv. in Mém. Soc. Linn. Par. i. (1822) 374 = strictus.
strictus, Eichw. Skizze, 146 = E. Ritro.
strictus, Fisch. ex Sims, Bot. Mag. t. 2457 = exaltatus.
strigosus, Bieb. Casp. App. 211 = sphaerocephalus.
strigosus, *Linn. Sp. Pl.* 815.—Hispan.; Marocc.
subglaber, *Schrenk in Bull. Phys.-Math. Acad. Petersb.* iii. (1845) 110.—Soongaria.
syriacus, Boiss. Diagn. Ser. I. x. 84 = viscosus.
Szowitzii, *Fisch. & Mey. ex DC. Prod.* vi. 625.—Reg. Caucas.
tauricus, Hort. ex DC. l. c. 638 = platylepis.
tauricus, Willd. ex Steud. Nom. ed. II. i. 588 = sphaerocephalus.
taygeteus, Boiss. & Heldr. Diagn. Ser. I. vi. 98 = sphaerocephalus.

ECHINOPS:—
tenuifolius, Fisch. ex Schkuhr, Handb. iii. 181 = E. Ritro.
tenuifolius, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 61 = banaticus.
tenuissimus, Sibth. & Sm. Fl. Graec. x. 19. t. 926 = graecus.
tibeticus, *Bunge, in Bull. Acad. Pétersb* vi. (1863) 404.—Tibet.
Tournefortii, *Ledeb. in Trautv. Diss. Echin.* 21.—Reg. Caucas.
tricholepis, *Schrenk, Enum. Pl. Nov.* 47.—Soongaria.
tunetanus, Vahl, ex Vitm. Summa, v. 175 = spinosus.
Turcaninowii, Trautv. Diss. Echin. 28; Ledeb. Fl. Ross. iii. 657 = Gmelini.
vaginatus, *Boiss. & Hausskn. in Boiss. Fl. Orient.* iii 437.—Syria.
villosissimus, *Bunge, in Bull. Acad. Pétersb.* vi. (1863) 406.—Persia.
villosus, Hort. ex DC. Prod. vi. 624 = sphaerocephalus.
virgatus, Lam. Encyc. ii. 337 = E. Ritro.
viscosus, *DC. Prod.* vi. 625.—Reg. Mediterr.; Orient.
viscosus, Reichb. Fl. Germ. Excurs. 486 = sphaerocephalus.
xanthacanthus, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 46.—Turkestan.

ECHINOPSILON, Moq. in Ann. Sc. Nat. Sér. II. ii. (1834) 127 = Chenoles. Thunb. (Chenop.).
anisacanthoides, F. Muell. in Trans. Phil. Inst. Vict. ix. (1858) 76 = Anisacantha echmopsila.
brachypterus, F. Muell. Fragm. vii. 13 = Kochia brachyptera.
carnosus, Moq. in DC. Prod. xiii. ii. 136 = C. carnosa.
caspicus, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 455.—As. centr.
crassifolius, Moq. in DC. Prod. xiii. ii. 137 = Kochia hirsuta.
dasyphyllus, Moq. l. c. 136 = Kochia dasyphylla.
diffusus, Moq. l. c. 137 = C. diffusa.
dioicanctus, Kar. & Klr. in Bull. Soc. Nat. Mosc. (1841) 736.—Soongaria.
eriophorus, Moq. in Ann. Sc. Nat. Sér. II. ii. (1834) 127 = Kochia latifolia.
eurotioides, F. Muell. Fragm. vii. 13 = C. eurotioides.
hirsutus, Moq. in Ann. Sc. Nat. Sér. II. ii. (1834) 127 = Kochia hirsuta.
hyssopifolius, Moq. in DC. Prod. xiii. ii. 135 = Kochia hyssopifolia.
lanatus, Moq. in Ann. Sc. Nat. Sér. II. ii. (1834) 127 = Kochia hyssopifolia.
muricatus, Moq. l. c. = Kochia muricata.
Reuterianus, Boiss. in Boiss. & Reut. Pugill. Pl. Nov. 105 = Kochia Kesteriana.
sclerolaenoides, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 75 = C. sclaerolenoides.
sedoides, Moq. in Ann. Sc. Nat. Sér. II. ii. (1834) 127 = Kochia sedoides.
sericeus, Moq. Chenop. Enum. 89 = C. diffusa.
stelligerus, F. Muell. Fragm. vii. 13 = Kochia stelligera.

ECHINOPSIS, Linn. Hort. Cliff. 391 (1737) = Echinopus, Linn. (Compos.).

ECHINOPSIS, Zucc. in Abh. Akad. Muench. ii. (1837) 675 = Cereus, Haw. (Cact.).
amoena, A. Dietr. in Otto & Dietr. Allg. Gartenz. xii. (1844) 187 = C. amoena.
amoenissima, Wender. Analect. Kdt. Bemerk. (1852) n. 6.—Hab.?
apiculata, Linke, in C. Koch & Fint. Wochenschr. i. (1858) 85.—Bolivia.
aurata, Salm Dyck, Cact. Hort. Dyck. ed. II. 89, 182 = C. auratus?
Bridgesii, Salm-Dyck, l. c. 181.—Bolivia.
campylacantha, Pfeiff. & Otto, Abbild. Cacteen, i. sub t. 4 = C. leucanthus.
cinnabarina, Labour. Monog. Cact. 288.—Bolivia.
cristata, Salm-Dyck, Cact. Hort. Dyck. ed. II. 88, 178. —Bolivia.
Decaisneana, Walp. Rep. ii. 324 = Echinocactus Decaisnei.
Eyriesii, Pfeiff. & Otto, Abbild. Cacteen, i. sub t. 4 = C. Eyriesii.

ECHINOPSIS :—
Forbesii, Hort. Angl. ex A. Dietr. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 193.—Hab. ?
formosa, Jacobi, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 39.—Chili.
formosissima, Lebour. in Rev. Hortic. Sér. IV. iv. (1855) 26.—Mexic.
gibbosa, Pfeiff. ex Foerst. Handb. Cact. ed. I. 291 = Echinocactus gibbosus.
grandiflora, Linke, in C. Koch, Allg. Gartenz. (1857) 289.—Bras.
Haageanus, Linke, in C. Koch & Fint. Wochenschr. i. (1858) 86 = Echinocactus Haageanus.
Huotii, Labour. Monog. Cact. 301.—Chili.
Jamesianus, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 38 = *Descainiana*.
Kuntii, Schlumb. in Rev. Hortic. Sér. IV. iii. (1854) 348 = *Huotii* !
leucantha, Walp. Rep. ii. 324 = C. leucanthus.
Maximiliana, Heyder, ex A. Dietr. in Otto & Dietr. Allg. Gartenz. xiv. (1846) 250.—Hab. ?
melanacantha, A. Dietr. l. c. 306 = Echinocactus tubiflorus.
Muleyi, Labour. Monog. Cact. 291 = *cristata*.
multiplex, Pfeiff. & Otto, Abbild. Cacteen, i. t. 4 = C. multiplex.
nigrispina, Linke, in C. Koch, Allg. Gartenz. (1857) 289.—Chili.
nigrispina, Walp. Rep. ii. 324.—Hab. !
nodosa, Linke, in C. Koch & Fint. Wochenschr. i. (1858) 85 = Echinocactus nodosus.
octacantha, Muehlenpf. in Otto & Dietr. Allg. Gartenz. xvi. (1848) 19.—Texas.
oxygona, Pfeiff. & Otto, Abbild. Cacteen, i. sub t. 4 = C. oxygonus.
pectinata, Fennel, in Otto & Dietr. Allg. Gartenz. xi. (1843) 282 = C. caespitosus.
Pentlandi, Salm-Dyck, ex A. Dietr. l. c. xiv. (1846) 250.—Peruv.
picta, Walp. Rep. ii. 324.—Hab. ?
pulchella, Zucc. ex Foerst. Handb. Cact. ed. I. 288 = C. pulchellus.
Reichenbachiana, Pfeiff. ex Foerst. l. c. 265 = C. caespitosus.
rhodacantha, Salm-Dyck, Cact. Hort. Dyck. ed. II. 39, 182.—Reg. Argent.
salpingophora, Lem. ex Salm-Dyck, l. c. 181; Labour. Monog. Cact. 302.—Hab. ?
Scheeri, Salm-Dyck, l. c. 179.—Bolivia.
Schelhasii, Pfeiff. & Otto, Abbild. Cacteen, i. sub t. 4 = C. Schelhasii.
Scopa, Carr. in Rev. Hortic. (1875) 374 = Echinocactus Scopa.
setosa, Linke, in C. Koch & Fint. Wochenschr. i. (1858) 86 = Echinocactus setosus.
simplex, Nieb, in C. Koch, Allg. Gartenz. (1857) 288. —Hab. !
Tougardii ×, Hérincq, L'Horticult. Franç. iii. (1853) 198.
tricolor, A. Dietr. in Otto & Dietr. Allg. Gartenz. xvi. (1848) 210.—Hab. ?
tuberculata, Nieb, in C. Koch. Allg. Gartenz. (1857) 287.—Bolivia.
tubiflora, Zucc. ex A. Dietr. in Otto & Dietr. Allg. Gartenz. xiv. (1846) 306 = Echinocactus tubiflorus.
turbinata, Pfeiff. & Otto, Abbild. Cacteen, i. sub t. 4 = C. turbinatus.
valida, Monv. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 181 = *Forbesii*.
Zuccarinii, Pfeiff. & Otto, Abbild. Cacteen, i. sub t. 4 = Echinocactus tubiflorus.

ECHINOPSUS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 109 = **Echinops**, Linn. (Compos.).

ECHINOPTERIS, Lindl. Veg. Kingd. 890 (1847) = seq.

ECHINOPTERYS, A. Juss. in Arch. Mus. Par. iii. (1843) 343 (Monog. 88. t. 9). *MALPIGHIACEAE*, Benth. & Hook. f. i. 264.
ECHINOPTERIS, Lindl. Veg. Kingd. 890 (1847).
Lappula, *A. Juss. ll. cc.*—Mexic.

ECHINOPUS, Tourn. ex Adans. Fam. ii. 114 (1763) = **Echinops**, Linn. (Compos.).
Ritro, Scop. Fl. Carn. ed. II. ii. 121 = Echinops Ritro.
sphaerocephalus, Scop. l. c. = Echinops sphaerocephalus.

ECHINOSCHOENUS, Nees & Meyen, Reise, i. 116 (1834); et in Linnaea, ix. (1834) 297 = **Rynchospora**, Vahl (Cyperac.).
Beyrichii, Nees, in Mart. Fl. Bras. ii. t. 122 = R. Beyrichii.
sparganioides, Lindl. ex Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 108 = R. exaltata.
triceps, Nees & Meyen, Reise, i. 116, 117; et in Linnaea, ix. (1834) 3, 297 = R. exaltata.

ECHINOSPARTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 358 = **Genista**, Linn. (Legumin.).
Barnadesii, Fourr. l. c. = G. Barnadesii.
horridum, Fourr. l. c. = G. horrida.
lugdunense, Fourr. l. c. = G. horrida.

ECHINOSPERMUM, Sw. ex Lehm. Asperif. 118 (1818). *BORAGINEAE*, Benth. & Hook. f. ii. 850.
HETEROCARYUM, A. DC. in DC. Prod. x. 144 (1846).
LAPPULA, [Riv.] Rupp. Fl. Jen. ed. Hall. 12 (1745); Moench. Meth. 416 (1794).
ROCHELIA, Roem. & Schult. Syst. iv. 11 (1819).
affine, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 714. —Sibir. altaic.
alpinum, Schur, Enum. Pl. Transs. 464 = E. Lappula.
anisacanthum, *Turcz. ex Ledeb. Fl. Ross.* iii. 156.—Sibir. baical.
barbatum, *Lehm. Asperif.* 128.—Marocco; Oriens; Sibir.
borbonicum, Lehm. l. c. 119 = Cynoglossum borbonicum.
brachycentrum, Eichw. Casp. Cane. 4 = microcarpum.
brachycentrum, *Ledeb. Fl. Alt.* i. 208.—Sibir. altaic.
brachycephalum, *Claus, in Britz. Opt. Russ. Reich.* viii. 240.—Reg. Casp.
Bungei, *Boiss. Fl. Orient.* iv. 252.—Persia.
caelestinum, Wight, Ic. t. 1304 = Paracaryum caelestinum.
calathicarpum, Stocks, in Hook. Kew Journ. iv. (1852) 175 = Paracaryum calathicarpum.
californicum, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 225.—Calif.
cancellatum, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Medic.) i. (1853) 489 = Rochelia stellulata.
canum, Benth. in Royle, Illustr. Bot. Himal. 306 = Eritrichium strictum.
capense, *DC. Prod.* x. 138.—Afr. austr.
carinatum, Boiss. Diagn. Ser. I. iv. 40 = barbatum.
casanense, Wirz. ex Ledeb. Fl. Ross. iii. 156 = seq.
casanicum, Wirz. Geogr. Pl. Casan. 84; et in Flora, xxv. (1842) 568 = E. Lappula.
caspium, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 34 = semiglabrum.
ciliatum, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 225.—Am. bor. occ.
circinnatum, A. Dietr. ex DC. Prod. x. 164 = Solenanthus circinnatus.
compressum, *Turcz. in Bull. Soc. Nat. Mosc.* xxiii. (1850) 1. 522.—Sibir. baical.
concavum, *F. Muell. Fragm.* ii. 139.—Austral.
condylophorum, *Lehm. Asperif.* 126.—Sibir.
consanguineum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 35.—Sibir. altaic.
cristatum, Bunge, Del. Sem. Dorp. (1839) 8 = Paracaryum cristatum.
cymosum, *Stschegl. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 1. 134.—Soongaria.
cynoglossoides, *Lehm. Asperif.* 131.—Afr. austr.
cynoglossoides, E. Mey. ex A. DC. in DC. Prod. x. 137 = E. Lappula.
deflexum, *Lehm. Asperif.* 120.—Reg. bor. temp.
diffusum, *Lehm. Pugill.* ii. 23.—Am. bor. occ.
diplolomia, *Fisch. & Mey. Enum. Pl. Nov. Schrenk*, 36.—Sibir. altaic.
divaricatum, *Bunge, in Lehm. Rel.* (1847) 284.—As. centr.
enerve, E. Mey. ex DC. Prod. x. 154 = Cynoglossum enerve.
filiforme, Godet, ex DC. Prod. x. 140 = microcarpum.
floribundum, *Lehm. Pugill.* ii. 24.—Am. bor. occ.
Fremontii, Torr. & Gray, in Pacif. Rail. Rep. xii. 46 = Redowskii.
glochidiatum, A. DC. Prod. x. 136 = Paracaryum glochidiatum.
gracile, Lehm. Asperif. 129 = Pectocarya linearis.
Greenei, *A. Gray, in Proc. Am. Acad.* xii. (1877) 168. —Calif.

ECHINOSPERMUM:—
heterocanthum, *Ledeb. Suppl. Ind. Sem. Hort. Dorp.* (1823) 3.—Reg. Caucas.
heterocaryum, *Bunge, in Rel. Lehm.* (1847) 285.—As. centr.
hispidum, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 106. —Am. bor. occ.
intermedium, Ledeb. Fl. Alt. i. 199 = Redowskii.
japonicum, Boiss. Diagn. Ser. I. xi. 128 = E. Lappula.
javanicum, *Lehm. Asperif.* 118.—Java.
Karelini, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 67. —Soongaria; Sibiria.
Kotschyi, Boiss. Diagn. Ser. I. vii. 29 = sinaicum.
laevigatum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 411.—Soongaria.
Lappula, *Lehm. Asperif.* 121.—Europ.; Oriens.
latifolium, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 89. —Abyss.
leiocarpum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 26 = Kryukkia leiocarpa.
macranthum, *Ledeb. Fl. Alt.* i. 205.—Sibir.; Turkestan.
marginatum, *Lehm. Asperif.* 130.—Reg. Caucas. et Ural.
mexicanum, *Hemsl. Biol. Centr. Am. Bot.* ii. 377.—Mexic.
microcarpum, *Ledeb. Fl. Alt.* i. 202.—Persia; Sibiria.
minimum, *Lehm. Asperif.* 126.—Rossia.
myosotiflorum, A. DC. in DC. Prod. x. 141 = Eritrichium strictum.
nervosum, Kellogg, in Proc. Calif. Acad. ii. (1868) 146. t. 42 = diffusum.
oligacanthum, *Boiss. Fl. Orient.* iv. 248.—Persia, Afghan.
oligacanthum, *Ledeb. Fl. Ross.* iii. 161.—Sibir. altaic.
ompholodes, *Schrenk, in Bull. Phys.-Math. Acad. Petersb.* iii. (1845) 211.—Sibiria.
pachypodum, *Hohen. ex A. DC. in DC. Prod.* x. 145.—Mexopot.
paniculatum, E. Mey. ex A. DC. l. c. 149, 143 = Cynoglossum micranthum.
patens, *G. Don, Gen. Syst.* iv. 378.—Am. bor.
patulum, *Lehm. Asperif.* 124.—Reg. Mediterr.; Sibir
patulum, Lehm. in Hook. Fl. Bor. Am. ii. 84 = Redowskii.
pilosum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 469 = Redowskii.
pinetorum, *Greene, ex A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 224.—N. Mexic.
pycnalcum, *Wallich. ex Lehm. in Anal. Soc. Esp. Hist. Nat.* xi. (1882) 117.—Mont. Pyren.
Redowskii, *Lehm. Asperif.* 127.—As. et Am. bor.
rigidum, DC. Prod. x. 141 = microcarpum.
rupestre, *Schrenk, in Bull. Phys.-Math. Acad. Petersb.* ii. (1842) 194.—Sibir. altaic.
scabrosum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 463 = Redowskii.
secundum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 718 = deflexum.
semicinctum, *Stev. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 1. 605, in textu.—Rossia.
semiglabrum, *Ledeb. Fl. Alt.* i. 204.—Sibir. altaic.; Turkestan.
sericeum, Benth. in Royle, Illustr. Bot. Himal. 306 = Eritrichium basifixum.
sessiliflorum, *Boiss. Diagn. Ser.* I. xi. 124.—Persia; Beluchist.
sinaicum, *A. DC. in DC. Prod.* x. 141.—Arab.; Persia.
spathulatum, Benth. in Royle, Illustr. Bot. Himal. 306 = Eritrichium spathulatum.
spinocarpon, Boiss. Fl. Orient. iv. 249 = Vahlianum.
squarrosum, Link, Enum. Hort. Berol. i. 165 = patulum.
squarrosum, Reichb. Fl. Germ. Excurs. 345 = E. Lappula.
strictum, *Ledeb. Fl. Alt.* i. 200.—Sibir. altaic.
strictum, Nees, in Neswied, Reise Nord-Am. ii. 443 = Redowskii.
strigosum, Stend. Nom. ed. II. i. 838 = Cynoglossum borbonicum.
stylosum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 718 = microcarpum.
subdecumbens, Parry, in Proc. Davenp. Acad. i. (1876) 148 = diffusum.
Szovitsianum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 26.—As. Min.; Persia; Soongar.
tenue, *Ledeb. Fl. Alt.* i. 201.—Sibir. altaic.
tenuinum, Schenk, in Linnaea, xxv. (1852) 260 = Redowskii.
thymiphilum, A. DC. in DC. Prod. x. 136 = deflexum.

ECHINOSPERMUM:—
trisrriale, Bunge, Del. Sem. Hort. Dorpat. (1840) 8. n. 13 = barbatum.
trisrriale, Kar. & Kir. ex Ledeb. Fl. Ross. iii. 162 (Quid ?).—Sibir. altaic.
tuberculosum, Ledeb. in Eichw. & Mey. Pl. Casp. 11. t. 6 = Vahlianum.
ursinum, *Greene, ex A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 224.—N. Mexic.
Vahlianum, *Lehm. Asperif.* 182.—Aegypt. ; Orient.
virginicum, *Lehm. l. c.* 117.—Am. bor.
vulgare, Sw. ex Schrad. in Comm. Gotting. iv. (1816–18) 195. *f. 7* = E. Lappula.
zeylanicum, Lehm. Asperif. 116 = Cynoglossum denticulatum.

ECHINOSPHAERA, Sieber, ex Steud. Nom. ed. II. i. 536 (1840) = **Rhinocoarpus**, Desf. (Euphorb.).
rosmarinifolia, Sieber, ex Benth. Fl. Austral. vi. 70 = R. pinifolius.

ECHINOSTACHYS, Brongn. in Planch. Hort. Donat. 25 (1854 ?) = **Aechmea**, Ruiz & Pav. (Bromel.).
Pineliana, Brongn. l. c. = A. Pineliana.
rosea, Beer, Bromel. 149 = A. Pineliana.
Van-Houtteana, Van Houtte, Cat. (1878) = Quesnelia Van-Houttei.

ECHINOSTACHYS, E. Mey. Comm. Pl. Afr. Austr. 248 (1837) = **Pycnostachys**, Hook. (Labiat.).
reticulata, E. Mey. l. c. = P. reticulata.

ECHINUS, Lour. Fl. Cochinch. 633 (1790) = **Mallotus**, Lour. (Euphorb.).
claoxyloides, Baill. Adansonia, vi. (1865–66) 315 = M. claoxyloides.
Dallachyanus, Baill. l. c. 314 = M. Dallachyi.
mollissimus, Baill. l. c. 316 = M. ricinoides.
neraphilus, Baill. l. c. 314 = M. nesophilus.
philippinensis, Baill. l. c. = M. philippinensis.
trisulcus, Lour. Fl. Cochinch. 633 = M. cochinchinensis.

ECHIOCHILON, Desf. Fl. Atlant. i. 166. t. 47 (1800).
BORAGINEAE, Benth. & Hook. f. ii. 852.
fruticosum, *Desf. l. c.*—Afr. bor.
hispidissimum, Tausch, in Flora, xii. (1829) 643 = Arnebia hispidissima.
longiflorum, *Benth. in Hook. Ic. Pl.* t. 1377.—Arab.

ECHIOGLOSSUM, Blume, Bijdr. 864. t. 28 (1825) = Cistacetonia, Blume (Orchid.).
javanicum, Blume, l. c.—Java.
minax, Reichb. f. in Hamb. Gartenz. xvi. (1860) 14.—Java.
muticum, Reichb. f. in Bonplandia, iii. (1855), 225.—Java.
quartum, Reichb. f. in Walp. Ann. vi. 391.—Java.
striatum, Reichb. f. in Gard. Chron. (1879) ii. 390.—Reg. Himal.

ECHIOIDES, Moench, Meth. 416 (1794) = **Myosotis**, Tourn. (Borag.).
annua, Moench, l. c. = M. arvensis.
palustris, Moench, l. c. = M. palustris.
perennis, Moench, l. c. = M. palustris.

ECHIOIDES, [Tourn.] Desf. Fl. Atlant. i. 168 (1800) = **Nonnea**, Medic. (Boragin.).
alba, Poir. in Dict. Sc. Nat. xiv. 218 = N. alba.
alpestris, Poir. l. c. 220 = N. alpestris.
arvensis, Poir. ex Steud. Nom. ed. II. i. 538 = Lycopsis arvensis.
caspica, Poir. in Dict. Sc. Nat. xiv. 219 = N. picta.
ciliata, Poir. l. c. 218 = N. ciliata.
Colmanniana, Poir. l. c. 219 = N. Colmanniana.
elongata, Poir. l. c. = N. elongata.
grandiflora, Desf. Cat. Hort. Par. ed. III. 120 = N. rosea.
lutea, Poir. in Dict. Sc. Nat. xiv. 217 = N. lutea.
nigricans, Desf. Fl. Atlant. i. 168 = N. nigricans.
obtusifolia, Poir. in Dict. Sc. Nat. xiv. 218 = N. obtusifolia.
picta, Poir. l. c. 219 = N. picta.
rosea, Poir. l. c. 220 = N. rosea.
rubra, Desf. Tabl. ed. I. 73 = N. rubra.
ventricosa, Poir. in Dict. Sc. Nat. xiv. 218 = N. alba.
violacea, Desf. Fl. Atlant. i. 164 = N. nigricans.

ECHION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = **Echium**, Tourn. (Boragin.).

ECHIOPSIS, Reichb. Handb. 192 (1837) = **Lobostemon**, Lehm. (Boragin.).

ECHISACHYS, Neck. Elem. iii. 228 (1790) = **Tragus**, Hall. (Gramin.).

ECHITES, P. Br. Hist. Jamaic. 181 (1756). *APOCYNACEAE*, Benth. & Hook. f. ii. 724.
SELUTTA-KAKA, Adans. Fam. ii. 172 (1763).
CHALCOMMA, Miers, Apocyn. S. Am. 110 (1878).
HETEROTHRIX, Muell. Arg. in Mart. Fl. Bras. vi. 188. t. 40 (1860).
MASSICITES, Muell. l. c. 150. t. 46 (1860).
MITOZUS, Miers, Apocyn. S. Am. 218 (1878).
acuminata, Roxb. Hort. Beng. 20 ; Fl. Ind. ii. 15 = Agonosma marginata.
acuminata, *Ruiz & Pav. Fl. Per.* ii. 19. t. 134.—Peruv.
acuminata, Willd. ex DC. Prod. viii. 475 = mucronata.
acutiloba, *A. DC. in DC. Prod.* viii. 451.—Peruv.?
adglutinata, Burm. f. Fl. Ind. 68 (Holarrhenae sp.).—Ind. or.
adglutinata, Jacq. Select. Am. 81. t. 23 = agglutinata.
affinis, Roem. & Schult. Syst. iv. 398 = Ichnocarpus ovatifolius.
agglutinata, *Jacq. Enum. Pl. Carib.* 18.—Ins. S. Doming.
albiflora, Miers, Apocyn. S. Am. 204 = brachyloba.
alexicaca, *Mart. ex Stadelm. in Flora*, xxiv (1841) 1. *Beibl.* 68.—Bras.
almadiandica, *Stadelm. l. c.* 28.—Bras.
amazonica, *Stadelm. l. c.* 50.—Bras.
amplissima, *Zipp. ex Span. in Linnaea*, xv. (1841) 394, nomen (Agonomatis sp.).—Ins. Timor.
andina, *Miers, Apocyn. S. Am.* 204.—Ecuador.
Andrewsii, *Chapm. Fl. S. U. St.* 359.—Ind. occ.
Andrieuxii, Miers, Apocyn. S. Am. 206 = Mandevilla Andrieuxii.
angustata, Vell. Fl. Flum. iii. t. 48 = Macrosiphonia longiflora.
angustifolia, *Benth. in Hook. Journ. Bot.* iii. (1841) 247.—Guiana.
angustifolia, *Poir. Encyc. Suppl.* ii. 537.—Ins. S. Doming.
annularis, Linn. Diss. Pl. Surin. n. 32 ; Amoen. Acad. viii. 254 = seq. ?
annularis, Linn. f. Suppl. 166 = Prestonia annularis.
antennacea, *A. DC. in DC. Prod.* viii. 456.—Peruv.?
antidysenterica, Roxb. ex Flem. in As. Res xi. (1810) 166 = Holarrhena antidysenterica.
apoxya, Voigt, Hort. Suburb. Calc. 522 = Agonosma marginata.
arborea, Vell. Fl. Flum. 114 ; iii. t. 47 = Secondatia arborea.
arborescens, Boj. ex A. DC. in DC. Prod. viii. 488 = Mascarenhasia arborescens.
aspera, *Mart. & Gal. in Bull. Acad. Brux* xi. (1844) 104.—Mexic.
Asperuginis, *Sw. Prod. Veg. Ind. Occ.* 52.—Ind occ.
atropurpurea, Lindl. in Past. Mag. Bot. ix. (1842) 199 = Dipladenia atropurpurea.
atrovioiacea, Stadeln. in Flora, xxiv. (1841) 1. Beibl. 75 = Dipladenia atrovioiacea.
augusta, Vell. Fl. Flum. 114 ; iii. t. 48 = longiflora.
auriculata, *Pohl, ex Stadelm. in Flora*, xxiv. (1841) 1. *Beibl.* 25.—Bras.
bantamensis, *Blume, Bijdr.* 1041.—Java.
barbata, *Desv. Prod. Pl. Ind. Occ.* 30.—Ind. occ.
barbata, D. Dietr. Syn. Pl. i. 651 = Paramena glandulifera.
Benthamii, *A. DC. in DC. Prod.* viii. 467.—Guiana.
Berterii, *A. DC. in DC. l. c.* 447.—Ins. S. Doming.
bicolor, Miq. Stirp. Surin. Sel. 154.—Guiana.
bicornis, *Spruce, ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 114.—Bras.
biflora, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 187 = Billbergii.
biflora, Chapm. in Coult. Bot. Gazette, iii. (1878) 11 = paludosa.
biflora, Ehrenb. ex Schlecht. in Linnaea, xxvi. (1853–55) 686 = Ehrenbergii.
biflora, *Jacq. Enum. Pl. Carib.* 18 ; *Linn. Sp. Pl.* ed. II. 307.—Ind. occ.
bifurcata, Porpp. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 113 = Odontadenia Poeppigii.
bignoniaeflora, Schlecht. in Linnaea, xxvi. (1853–55) 573 = Stemmadenia bignoniaeflora.
Billbergii, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 187.—Panama.

ECHITES:—
bispinosa, Linn. f. Suppl. 167 = Pachypodium bispinosum.
Blanchetii, *A. DC. in DC. Prod.* viii. 448.—Bras.
bogotensis, *H. B. & K. Nov. Gen. et Sp.* iii. 215. t. 243. —N. Granat.
brachiata, Wall. Cat. n. 1668 = Eodyssantha mucrantha.
brachyloba, *Miers, Apocyn. S. Am.* 208.—Peruv.
brachyphon, Torr. in Bot. Mex. Bound. 168 = Macrosiphonia brachysiphon.
brachystachya, Benth in Hook. Journ. Bot. iii. (1841) 248.—Guiana.
bracteata, Griseb. in Goett. Abh. ix. (1874) 208 = Parsonsia bracteata.
bractesta, Heyne, in Wall. Cat. sub n. 1656 = Agonosma cymosa.
bracteata, *H. B. & K. Nov. Gen. et Sp.* iii. 217.—N. Granat.
bractesta, Vell. Fl. Flum. 112 ; ill. t. 41 = Forsteronia bracteata.
brasiliensis, *Thunb. Diss. Echit.* 5.—Bras.
brevipes, *Benth. Pl. Hartw.* 216.—N. Granat.
calycina, Wall. Cat. n. 1668 = Agonosma calycina.
calycosa, *A. Rich. Fl. Cub. Pansrop.* ii. 94.—Ind. occ.
campestris, *Vell. Fl. Flum.* 113 ; ill. t. 43.—Bras.
canescens, *Willd. ex Roem. & Schult. Syst.* iv. 795.—Am. austr.
caryophyllata, Blume, Bijdr. 1041 = Agonosma caryophyllata.
caryophyllata, Roth, Pl. Nov. Sp. 133 = Ichnocarpus ovatifolius.
caryophyllata, Roxb. Hort. Beng. 20 ; Fl. Ind. ii. 11 = Agonosma calycina.
Catesbaei, G. Don, Gen. Syst. iv. 74 = Andrewsii ?
caudata, Blanco, Fl. Filip. ed. I. 106 = Ichnocarpus frutescens.
caudata, Burm. f. Fl. Ind. 68. t. 26 ; Linn. Mant. i. 52 = Strophanthus dichotomus.
chilensis, *A. DC. in DC. Prod.* viii. 468.—Chili.
chlorantha, *Schlecht. in Linnaea*, xxvi. (1853–55) 668 —N. Granat.
chrisophorizas, *Ham. Prod. Pl. Ind. Occ.* 31.—Ind. occ.
ciliata, *Stadelm. in Flora*, xxiv. (1841) 1. *Beibl.* 32.—Bras.
cimicida et *cimscifuga*, Pav. ex A. DC. in DC. Prod. viii 413 = Haplophyton cimicidum.
cinerea, A. Rich. Fl. Cub. Fanerog. ii. 93 = Haplophyton cimicidum.
circinalis, *Sw. Prod. Veg. Ind. Occ.* 52.—Ind. occ.
citrifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 216.—N. Granat.
citrina, *A. DC. in DC. Prod.* viii. 474.—Ind. occ.
clavata, *Roxb. Hort. Beng.* 85 ; *Fl. Ind.* ii. 20.—Ins. Moluce.
collita, *Vell. Fl. Flum.* 112 ; ill. t. 40.—Bras.
coccinea, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 286 = Dipladenia coccinea.
costata, *Stadelm. in Flora*, xxiv. (1841) 1. *Beibl.* 79.—Bras.
concolor, *Ham. Prod. Pl. Ind. Occ.* 31.—Ind. occ.
conferta, Wall. Cat. n. 1655 = Agonosma cymosa.
congesta, *H. B. & K. Nov. Gen. et Sp.* iii. 214.—N Granat.
conoides, *Zipp. ex Span. in Linnaea*, xv. (1841) 824, nomen (Agonomatis sp. ?).—Ins. Timor.
convolvulacea, A. DC. in DC. Prod. viii. 451 = Mandevilla convolvulacea.
cordata, *A. DC. l. c.* 451.—Mexic.
coriacea, *Benth. in Hook. Journ. Bot.* iii. (1841) 249. —Guiana.
coriacea, *Blume, Bijdr.* 1039.—Java.
coriacea, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 6 = Cryptolepis grandiflora.
coriacea, Wall. Cat. n. 1586 = Anodendron paniculatum.
corymbosa, Jacq. Enum. Pl. Carib. 18 = Parsonsia corymbosa.
costata, Forst. f. Prod. 90 = Alstonia costata.
Coulteri, *S. Wats. in Proc. Am. Acad.* xviii. (1883–83) 118.—N. Mexic.
crassifolia, *Spruce, ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 143.—Bras.
crassinoda, Garda. in Hook. Lond. Journ. Bot. i. (1842) 544 = Dipladenia crassinoda.
crassipes, *A. Rich. Fl. Cub. Fanerog.* ii. 91.—Cuba.
cristata, Roth, Nov. Pl. Sp. 135 = Agonosma marginata.

ECHITES :—
cubensis, *Griseb. Cat. Pl. Cub.* 172.—Cuba.
cuiabensis, Muell. Arg. in Mart. Fl. Bras. vi. i. 145 = oxypetalea.
Cururu, *Mart. in Buchn. Rep. Pharm.* (1830) n. 101. —*Cf. Linnaea*, vi. (1831) *Litt.* 30.—Bras.
cuspidata, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 5 = Cryptolepis Buchanani.
cuspidata, Willd. ex Muell. Arg. in Mart. Fl. Bras. vi. i. 152 = trifida.
cuyabensis, *A. DC. in DC. Prod.* viii. 462.—Bras.
cymosa, Roxb. Hort. Beng. [54]; Fl. Ind. ii. 16 = Agnosma cymosa.
Cenaevenulosa, *Stadelm. in Flora*, xxiv. (1841) i. *Beibl.* 47.—Bras.
densiflora, *Blume, Bijdr.* 1040.—Java.
densiflora, Pohl, ex Stadelm. in Flora, xxiv. (1841) i. Beibl. 56 = pycnantha.
denticulata, *Vell. Fl. Flum.* 110; iii. t. 80.—Bras.
dichotoma, Heyne, ex Hook. f. Fl. Brt. Ind. iv. 14 = Tonocarpus Roxburghii.
dichotoma, *H. B. & K. Nov. Gen. et Sp.* iii. 217.—Am. austr.
dichotoma, Roth, Nov. Pl. Sp. 186 = Agnosma caryophyllata.
dichotoma, Roxb. Hort. Beng. 90; Fl. Ind. ii. 19 = Vallaris Heynei.
dichotoma, *Thunb. Diss. Echit.* 6.—Hab.?
didyma, *Vell. Fl. Flum.* 109; iii. t. 27.—Bras.
difformis, Walt. Fl. Carol. 98 = Trachelospermum difforme.
disadena, *Miq. Stirp. Surin. Sel.* 156.—Guiana.
discolor, *Morits, ex Miers, Apocyn. S. Am.* 224.—Venezuela.
dominguensis, *Jacq. Coll.* i. 78; *Sw. Prod. Veg. Ind. Occ.* 52.—Ind. occ.
dubia, *Vell. Fl. Flum.* 114; iii. t. 50.—Bras.
edulis, Thunb. Diss. Echit. 5 = Brachystelma macrorhizum.
Ehrenbergii, *Schlecht. in Linnaea*, xxvi. (1853–55) 666. —Ins. S. Doming.
elastica, *Hort. Angl. ex A. DC. in DC. Prod.* viii. 479, nomen.—Bras.
elegans, Benth. in Hook. Journ. Bot. iii. (1841) 249 = Odontocoma geminata.
elegans, Wall. Cat. n. 1656 = Agnosma cymosa.
elliptica, *Thunb. Diss. Echit.* 4.—N. Granat.
elliptica, Wall. Cat. n. 1666 = Rhynchodia Wallichii.
emarginata, Vell. Fl. Flum. 113; iii. t. 46 = Lasegnea emarginata.
erecta, DC. Prod. viii. 469 = Rhodocalyx rotundifolius.
erecta, *Thunb. Diss. Echit.* 7.—Ind. or.; Java.
erecta, *Vell. Fl. Flum.* 118; iii. t. 45.—Bras.
esculenta, Wall. Cat. n. 1671 = Urceola esculenta.
ferruginea, *A. Rich. Fl. Cub. Fanerog.* ii. 92.—Cuba.
ferruginea, *Thunb. Diss. Echit.* 5 (sp. ignot.)—Java.
floribunda, Sw. Prod. Veg. Ind. Occ. 52 = Parsonsia floribunda.
fluminensis, *A. DC. in DC. Prod.* viii. 452.—Bras.
fragrans, Stadelm. in Flora, xxiv. (1841) i. Beibl. 71 = Diplademia fragrans.
Francisceae, A. DC. in DC. Prod. viii. 452 = violacea.
Franciscea, Lindl. Bot. Reg. (1847) t. 24 = varia.
Fraseri, *Willd. ex Roem. & Schult. Syst.* iv. 796.—Am. bor.
frutescens, Roxb. Hort. Beng. 90; Fl. Ind. ii. 12 = Ichnocarpus frutescens.
funiformis, *Vell. Fl. Flum.* 109; iii. t. 29.—Bras.
galeiformis, *Rudolphi, ex Ledeb. Pl. S. Doming.* 6.—Ins. S. Doming.
ganabarica, Casar. ex Muell. Arg. in Mart. Fl. Bras. vi. i. 144 = funiformis.
geminata, *Hoffmgg. ex Roem. & Schult. Syst.* iv. 795. —Bras.
glandulifera, Wall. Cat. n. 1659 = Parsmeria glandulifera.
glandulosa, Poir. Encyc. Suppl. ii. 687 = laxa.
glandulosa, Ruiz & Pav. Fl. Per. ii. 19 = Prestonia pervviana.
glauca, Roem. & Schult. Syst. iv. 796 = Hancornia speciosa.
glaucescens, Mart. & Gal. in Bull. Acad. Brux. xi. (1844) 356 = Mandevilla oaxacana.
glomerata, *Poir. Encyc. Suppl.* ii. 686.—Ins. S. Doming.
gracilipes, *Stadelm. in Flora*, xxiv. (1841) i. *Beibl.* 22. —Bras.
gracilis, *H. B. & K. Nov. Gen. et Sp.* iii. 219.—Venezuela.

ECHITES :—
grandiflora, Hook. & Arn. in Journ. Bot. i. (1834) 286 = longiflora.
grandiflorea, G. F. W. Mey. Prim. Fl. Esseq. 181 = Odontadenia speciosa.
grandiflora, Roth, Nov. Pl. Sp. 186 = Chonemorpha grandiflora.
grandiflora, Roxb. Hort. Beng. 20; Fl. Ind. ii. 14 = Beaumontia grandiflora.
grandiflora, Stadelm. in Flora (1841) i. Beibl. 49 = Odontadenia sylvestris.
grandis, Wall. Cat. n. 1658 = Chonemorpha macrophylla.
guaranitica, St. Hil. in Mém. Mus. Par. xii. (1825) 324 = E. Velame.
guianensis, *A. DC. in DC. Prod.* viii. 459.—Guian
guianensis, Thonn. in Schem. & Thonn. Beskr. Guin. Pl. 149 = Motandra guineensis.
Haenthii, Boj. ex A. DC. in DC. Prod. viii. 488 = Mascarenhasia Haenthii.
hebecarpa, *Benth. ex Muell. Arg. in Mart. Fl. Bras.* vi. i. 112.—Bras.
heterophylla, J. F. Gmel. Syst. 437 = domingensis.
heterophylla, Miq. in Linnaea, xxv. (1852) 653 = chilensis?
Heynii, Spreng. Syst. i. 632 = Agnosma caryophyllata.
hircus, Roxb. Hort. Beng. 85; Fl. Ind. ii. 18 = Vallaris Pergulana.
hirsuta, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107. —Guiana.
hirsuta, Ruiz & Pav. Fl. Per. ii. 19 = Prestonia hirsuta.
hirsuta, Stadelm. in Flora, xxiv. (1841) i. Beibl. 27 = Mandevilla hispida.
hirsuta, Vell. Fl. Flum. 113; iii. t. 44 = fluminensis.
hirtella, Benth. Pl. Hartw. 67 = Mandevilla oaxacana.
hirtella, H. B. & K. Nov. Gen. et Sp. iii. 213 = Prestonia hirsuta.
hypoleuca, Benth. Pl. Hartw. 23 = Macrosiphonia hypoleuca.
illustris, Vell. Fl. Flum. 114; iii. t. 49 = Diplademia illustris.
infans, *Blume, Bijdr.* 1089.—Java.
insignis, Spreng. Syst. i. 632 = Odontadenia speciosa.
isthmica, Vell. Fl. Flum. 112; iii. t. 39 = Condyloscrpon isthmicum.
jamaicensis, *Griseb. Fl. Brit. W. Ind.* 416.—Ind. occ.
japurensis, *Stadelm. in Flora*, xxiv. (1841) i. Beibl. 19. —Bras.
jasminiflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 357.—Mexic.
javanica, D. Dietr. Syn. Pl. i. 651 (= *Chavannesia javanica*).—Java.
javitensis, Benth. ex Muell. Arg. in Mart. Fl. Bras. vi. i. 143 (= *Amblyanthera Spruceana*).—Bras.
jucunda, *H. B. & K. Nov. Gen. et Sp.* iii. 220.—N. Granat.
Karwinskii, Miers, Apocyn. S. Am. 206 = Mandevilla Karwinskii.
laevigata, Wall. Cat. n. 1669 = Epigynum laevigatum.
laevis, *Thunb. Diss. Echit.* 4.—Zeylan.
lanata, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 359.—Mexic.
lanceolata, Spreng. Syst. i. 634 = Parsonsia lanceolata.
lanuginosa, Mart. & Gal. in Bull. Acad. Brux. xi. (1844) 357 = Macrosiphonia lanuginosa.
lappulacea, *Lam. Encyc.* ii. 341.—Ins. S. Doming.
laxicarpa, A. DC. in DC. Prod. viii. 468 = Mandevilla laxicarpa.
lenticula, Buch.-Ham. ex Wall. Cat. n. 1657 = Chonemorpha macrophylla.
laxa, *Ruiz & Pav. Fl. Per.* ii. 19. t. 184.—Peruv.
lenticula, Zipp. ex Span. in Linnaea, xv. (1841) 824, nomen = Cryptolepis elegans.
leptoloba, *Stadelm. in Flora*, xxiv. (1841) i. *Beibl.* 15. —Bras.
leptophylla, *A. DC. in DC. Prod.* viii. 455.—Bras.
Lindeniana, *Griseb. Cat. Pl. Cub.* 173.—Cuba.

ECHITES :—
linearifolia, *Ham. Prod. Pl. Ind. Occ.* 31.—Ind. occ.
linearifolia, Stadelm. in Flora, xxiv. (1841) i. Beibl 18 = leptophylla.
linearis, Vell. Fl. Flum. 111; iii. t. 36 = Forsteronia linearis.
lisianthiflora, Boj. Hort. Maurit. 211 = Mascarenhasia lisianthiflora.
littores, *H. B. & K. Nov. Gen. et Sp.* iii. 212.—Cuba.
lobata, Afzel. ex A. DC. in DC. Prod. viii. 479 = lorata.
Lockharti, *Steud. Nom.* ed. II. i. 539.—Ins. Trinit
longiflora, *Desf. in Mém. Mus. Par.* v. (1819) 276.—Bras.
longiflora, Miers, Apocyn. S. Am. 194 = umbellata.
lorata, *Afzel. ex Steud. Nom.* ed. II. i. 539.—Afr. trop.
lucida, G. Don, Gen. Syst. iv. 75 = Urceola esculenta.
lucida, *Ruiz & Pav. Fl. Per.* ii. 19.—Peruv.
lucida, Wall. Cat. n. 1670 = Urceola lucida.
lucida, *Willd. ex Roem. & Schult. Syst.* iv. 795.—Venezuela.
luteo, *Vell. Fl. Flum.* 109; iii. t. 25.—Bras.
macrantha, Roem. & Schult. Syst. iv. 795 = Odontadenia speciosa.
macrantha, Spreng. Syst. i. 632 = Chonemorpha macrophylla.
macrocalyx, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 160 —Bras.
macrocarpa, *A. Rich. Fl. Cub. Fanerog.* ii. 94.—Cuba.
macrocarpa, Wall. Cat. n. 1662 = Agnosma marginata.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* iii. 218 —Venezuela.
macrophylla, Roxb. Hort. Beng. 20; Fl. Ind. ii. 13 = Chonemorpha macrophylla.
macrosiphon, Torr. in Bot. Mex. Bound. 158. t. 45 = Macrosiphonia Berlandieri.
macrostoma, *Benth. in Hook. Journ. Bot.* iii. (1841) 248.—Guian.
maculata, *A. DC. in DC. Prod.* viii. 474.—Ins. S. Doming.
madida, *Vell. Fl. Flum.* 112; iii. t. 42.—Bras.
malabarica, Lam. Encyc. ii. 342 = Alstonia scholaris.
Manaosona, A. DC. in DC. Prod. viii. 448 = sulphurea.
manubriata, Wall. Cat. n. 1663 = Anodendron paniculatum.
marambaiensis, G. Don, Gen. Syst. iv. 74 = versicolor
marginata, Roxb. Hort. Beng. [54]; Fl. Ind. ii. 16 = Agnosma marginata.
Martiana, Stadelm. in Flora, xxiv. (1841) i. Beibl. 31 = Diplademia Martiana.
Martii, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 155.—Bras.
Maximiliana, *Stadelm. in Flora*, xxiv. (1841) i. *Beibl.* 48.—Bras.
megapros, Vell. Fl. Flum. 110; iii. t. 33 (Asclepiadeid.?). —Bras.
melaleuca, *Hort. ex Regel, Cat. Pl. Hort. Akakow.* 52. —Hab.?
membranacea, *A. DC. in DC. Prod.* viii. 457.—Peruv.?
mexicana, *Miers, Apocyn. S. Am.* 208.—Mexic.
Meyeriana, Roem. & Schult. ex A. DC. in DC. Prod. viii. 426 = Odontadenia speciosa.
micrantha, Wall. Cat. n. 1667 = Ecdysanthera micrantha.
microcalyx, *A. DC. in DC. Prod.* viii. 456.—Ind. occ.
microphylla, *Stadelm. in Flora*, xxiv. (1841) i. *Beibl.* 35.—Bras.
mollis, Spreng. Syst. i. 634 = Parsonsia lanceolata.
mollis, Willd. in Roem. & Schult. Syst. iv. 795 = macrophylla.
mollissima, *H. B. & K. Nov. Gen. et Sp.* iii. 218.—N. Granat.
monilifera, Wall. Cat. n. 1660 = Parsmeria glandulifera.
montana, *H. B. & K. Nov. Gen. et Sp.* iii. 213.—N. Granat.
macronata, Roem. & Schult. Syst. iv. 796 = trifida.
multinervia, Gomes, ex A. DC. in DC. Prod. viii. 437 = Forsteronia multinervia.
muricata, *A. DC. in DC. l. c.* 474.—Ind. occ.
myrtifolia, Poir. Encyc. Suppl. ii. 686 = Parsonsia myrtifolia.
myrtifolia, Roem. & Schult. Syst. iv. 796 = rosea.

ECHITES:—
Neriandra, Griseb. Fl. Brit. W. Ind. 415 = Andrewsii.
nitida, *Vahl, Eclog. Am.* ii. 19.—Am. austr.
nobilis, Hort. Belg. ex Lem. in Fl. des Serres, Sér. I. v. (1849) sub t. 487 = Diplademia nobilis.
nodosa, *Willd. ex Roem. & Schult. Syst.* iv. 796.—Am. austr.
nummularis, *Burm. f. Fl. Ind.* 69.—Ind. occ.
nutans, Anders. in Trans. Soc. Arts Lond. xxv. (1835!) 208 (= *Haemadictyon ovatum*.).—Ind. occ.
oaxacana, A. DC. in DC. Prod. viii. 451 = Mandevilla oaxacana.
obliqua, Miers, Apocyn. S. Am. 193 = umbellata.
oblongifolia, *Ham. Prod. Pl. Ind. Occ.* 80.—Ind. occ.
obovata, *Nees, ex Steud. Nom.* ed. II. i. 540.—Bras.
obtusa, *Thunb. Diss. Echit.* 6.—Ind. or.
odoratissima, *Grif. Notul.* iv. 76 (sp. dub.).—Burma.
odorifera, *Vell. Fl. Flum.* 109; ill. t. 98.—Bras.
ovalifolia, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 286 = Hookeri.
ovalifolia, *Poir. Encyc. Suppl.* ii. 535.—Ins. S. Doming.
Pala, Buch.-Ham. ex Spreng. Syst. i. 688 = Alstonia scholaris.
pallida, *Miers, Apocyn. S. Am.* 195.—Ecuador.
paludosa, *Vahl, Eclog. Am.* ii. 19; *Ic.* t. 5.—Am. trop.
palustris, Salm, ex A. DC. in DC. Prod. viii. 468 = tomentosa.
pandurata, *A. DC. l. c.* 458.—Mexic.
paniculata, *Poir. Encyc. Suppl.* ii. 536.—Am. austr.
paniculata, Roxb. Hort. Beng. 20; Fl. Ind. ii. 17 = Anodendron paniculatum.
paniculata, Thonn. ex A. DC. in DC. Prod. viii. 428 = Motandra guineensis.
parviflora, *Afzel. ex Steud. Nom.* ed. II. i. 540.—Afr. trop.
parviflora, Roxb. Hort. Beng. 85; Fl. Ind. ii. 20 = Epigynum parviflorum.
pastorum, Mart. ex Stadelm. in Flora, xxiv. (1841) t. Beibl. 59 = Diplademia pastorum.
pauciflora, Rottl. ex Hook. f. Fl. Brit. Ind. iv. 6 = Cryptolepis elegans.
Pavonii, *A. DC. in DC. Prod.* viii. 468.—Peruv.
peduncularis, Stadelm. in Flora, xxiv. (1841) t. Beibl. 54 = Diplademia pedunculatis.
Pellieri, Loud. Encyc. Pl. 1541 = Andrewsii.
peltata, Lockh. ex G. Don, in Loud. Hort. Brit. 67 = Lockhartii.
peltata, *Vell. Fl. Flum.* 110; ill. t. 118.—Bras.
peltigera, *Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 21.—Bras.
petraea, St. Hil. in Mém. Mus. Par. xii. (1825) 322 = Macrosiphonia verticillata.
pilosa, Vell. Fl. Flum. 113; ill. t. 88 (= *Thyrsanthus pilosus*).—Bras.
pinifolia, St. Hil. in Mém. Mus. Par. xii. (1825) 325 = Macrosiphonia verticillata.
plicata, A. DC. in DC. Prod. viii. 454 = peltata.
Pohliana, *Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 78.—Bras.
polyantha, Wall. Cat. n. 1664 = Anodendron paniculatum.
portobellensis, *Benrl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 137.—Panama.
Prieurei, *A. DC. in DC. Prod.* viii. 458.—Guiana.
Primula, Hort. ex Walp. Rep. vi. 742 = Diplademia vincaeflora.
procumbens, Blanco, Fl. Filip. ed. II. 78 = Holarrhena macrocarpa.
pteridifolia, *Mart. ex Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 48.—Bras.
Ptarmica, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 69. t. 278.—Chili.
puberula, Michx. Fl. Bor. Am. i. 120 = Trachelosperma difforme.
pubescens, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 694 = Holarrhena antidysenterica.
pubescens, Hook. & Arn. Bot. Beech. Voy. 84 = chilensis.
pubescens, *Willd. ex Roem. & Schult. Syst.* iv. 796.—Am. austr.
pubiflora, *G. Don, Gen. Syst.* iv. 78.—Bras.
pulchella, Gardn. in Hook. Ic. Pl. t. 470 = Diplademia longiflora.
pulcherrima, *Pohl. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 119.—Bras.

ECHITES:—
puncticulosa, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guiana.
pycnantha, *Steud. Nom.* ed. II. i. 540.—Bras.
quinquangularis, Jacq. Enum. Pl. Carib. 13; Linn. Sp. Pl. ed II. 307 = Prestonia quinquangularis.
religiosa, *Teijsm. & Binn. in Tijdschr. Nederl. Ind* xxvii. (1864) 84.—Siam.
repens, Blanco, Fl. Filip. ed. I. 109 = Holarrhena macrocarpa.
repens, *Yacq. Enum. Pl. Carib.* 13.—Ins. S. Doming.
reticulata, Boj. Hort. Maurit. 211 = Agnosoma marginata.
reticulata, Roth, Nov. Pl. Sp. 184 = Cryptolepis Buchanani.
revoluta, A. DC. in DC. Prod. viii. 457 = circinalis.
rhynchosperma, Wall. Pl. As. Rar. i. 43. t. 49 = Rhynchodia Wallichii.
Richardii, Roem. & Schult. Syst. iv. 891 = hirsuta.
riparia, *H. B. & K. Nov. Gen. et Sp.* iii. 218.—N. Granat.
Rose-constantestris, Endl. in Harting. Parad. Vindob. t. 51 = Diplademia Rosa-campestris.
roseo, *A. DC. in DC. Prod.* viii. 450.—Ind. occ.
rotundifolia, *Mihan, ex A. DC. l. c.* 479.—Bras.
rubricaulis, *Poir. Encyc. Suppl.* ii. 535.—Guiana.
rubro-venosa, *Linden, in R. Morr. Belg. Hortic.* xvii. (1867) 108.—Bras.
rugellosa, *Rich. in Act. Soc. Nat. Hist. Nat. Par.* i. (1792) 107.—Guiana.
rugosa, *Benth. in Hook. Journ. Bot.* iii. (1841) 248.—Guiana.
sagittata, Poir. Encyc. Suppl. ii. 537 = subsagittata.
Sagraei, *A. DC. in DC. Prod.* viii. 450.—Florida.
salicifolia, *Raf. ex Fl. Am.* iv. 59.—Am. bor.
salicifolia, Willd. in Roem. & Schult. Syst. iv. 796 = Allamanda cathartica.
sancta, Stadelm. in Flora, xxiv. (1841) t. Beibl. 59 = Diplademia sancta.
sanguinolente, Tussac, Fl. Antill. 95. t. 11 (= *Haemadictyon ovatum*).—Ind. occ.
scabra, *Mart. ex Roem. & Schult. Syst.* iv. 795.—Bras.
scabra, Lahill. Sert. Austro-Caled. 26. t. 31 = Thenardia scabra.
scandens, Thunb. Diss. Echit. 6.—Java; Zeylan.
schottii, Linn. Mant. 55 = Alstonia scholaris.
Schomburgkii, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1146.—Guiana.
secundiflora, *A. DC. in DC. Prod.* viii. 457.—Mexic.
semidigyna, *Berg, in Verh. Zeeusch. Gen. Wetens.* iii. (1773) 588.—Gulan.
sessilis, Vell. Fl. Flum. 111; ill. t. 85 = Maloutia scabra.
sinensis, Hort. Angl. ex Walp. Rep. vi. 491 (= *Hybanthera cordifolia*).—China.
siphilitica, Auct. = syphilitica.
speciosa, *H. B. & K. Nov. Gen. et Sp.* iii. 219.—Venezuela.
spectabilis, *Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 44.—Bras.
spicata, Jacq. Enum. Pl. Carib. 13 = Forsteronia spicata.
spiciflora, Stadelm. *in Flora*, xxiv. (1841) t. *Beibl.* 58.—Bras.
spinosa, Burm. f. Fl. Ind. 69 = Carissa Carandas.
spiralis, Blanco, Fl. Filip. ed. I. 110 = Parsonsia Rheedii.
splendens, Hook. Bot. Mag. t. 3976 = Diplademia splendens.
splendens, *Span. in Linnaea*, xv. (1841) 324.—Ins. Timor.
Stadelmeyeri, *Mart. ex Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 29.—Bras.
stellaris, *Lindl. in Bot. Reg.* t. 1664.—Bras.
stellulifera, Lem. in Jard. Fleur. i. (1851) t. 67 = viridiflora.
suaveolens, A. DC. in DC. Prod. viii. 452 = Mandevilla suaveolens.
suaveolens, Mart. & Gal. in Bull. Acad. Brux. xi. (1844) 256 = Macrosiphonia hypoleuca.
subsarmosa, *Benth. in Hook. Journ. Bot.* iii. (1841) 247.—Guiana.
subverticta, Andr. Bot. Rep. t. 187 = Andrewsii.
suberecta, Jacq. Enum. Pl. Carib. 13 (= *Haemadictyon suberectum*).—Ind. or.
suberosa, *Vell. Fl. Flum.* 111; ill. t. 84.—Bras.
subsagittata, *Ruiz & Pav. Fl. Per.* ii. 10.—Peruv.
subsessilis, *A. DC. in DC. Prod.* viii. 461.—Peruv.

ECHITES:—
subspicata, *Vahl, Eclog. Am.* ii. 18.—Am. austr.
succulenta, Linn. f. Suppl. 167 = Pachypodium tomentosum.
sulphurea, *Vell. Fl. Flum.* 109; ill. t. 26.—Bras.
surinamensis, *Miq. Stirp. Surin. Sel.* 155.—Guiana.
sylvestris, A. DC. in DC. Prod. viii. 464 = Odontodenia sylvestris.
symphytocarpa, *G. F. W. Mey. Prim. Fl. Esseq.* 132.—Guian.
syphilitica, *Linn. f. Suppl.* 167.—Am. austr.
tenuicaulis, *Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 40.—Bras.
tenuifolia, Mikan, Fl. et Faun. Bras. fasc. 3 = Diplademia tenuifolia.
thyrsoidea, Vell. Fl. Flum. 111; ill. t. 87 = Forsteronia thyrsoides.
tomentosa, *Raf. Fl. Ludov.* 46.—Am. bor.
tomentosa, Roth, Nov. Pl. Sp. 137 = Wrightia tomentosa?
tomentosa, *Vahl, Symb. Bot.* iii. 44.—Gulan.
torosa, Jacq. Enum. Pl. Carib. 13 = torulosa.
torquata, Casar. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 102 = Forsteronia brasiliensis.
torulosa, *Linn. Sp. Pl.* ed. II. 307.—Mexic.
triangularis, Buch.-Ham. ex A. DC. in DC. Prod. viii. 411, 480 = Trachelospermum fragrans.
trichonerma, Zipp. ex Span. in Linnaea, xv. (1841) 324 = Ichnocarpus moluccanus.
trichotoma, *Desf. Cat. Hort. Par.* ed. III. 398.—Hab.?
trifida, *Blanco, Fl. Filip.* ed. I. 109.—Ins. Philipp.
trifida, *Yacq Enum. Pl. Carib.* 13.—Ind. occ.
tropaeifolia, *A. DC. in DC. Prod.* viii. 447.—Bras.
truncata, Lam. Encyc. ii. 340 = Gonolobus caroliniensis.
tuberosa, Haw. ex Steud. Nom. ed. II. i. 540 (= Pachypodii sp.).—Afr. austr.
tubiflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 255.—Mexic.
tubulosa, *Benth. in Hook. Journ. Bot.* iii. (1841) 249.—Guian.
Tweediana, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 370.—Reg. Argent.
umbellata, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 137 = portobellensis.
umbellata, *Yacq. Enum. Pl. Carib.* 13; *Linn. Sp. Pl.* ed. II. 307.—Ind. occ.
Valmazulana, *A. Rich. Fl. Cub. Fanerog.* ii. 93.—Ind. occ.
varia, *Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 17.—Bras.
variegata, Schrad. in Goett. Gel. Anz. i. (1821) 707 = obovata.
Vauthieri, A. DC. in DC. Prod. viii. 457 = coalita.
Velame, *St. Hil. in Bull. Soc. Philom.* (1824) 77.—Bras.
Vellosiana, A. DC. in DC. Prod. viii. 474 = Forsteronia bracteata.
velutina, Mart. ex Stadelm. in Flora, xxiv. (1841) t. Beibl. 72 = Diplademia gentianoides.
velutina, Spreng. Syst. t. 684 = Parsonsia velutina.
venenata, Roxb. Hort. Beng. 90; et ex A. DC. in DC. Prod. viii. 411 = Alstonia venenata.
venenosa, *Mart. ex Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 56.—Bras.
veraguensis, Seem. Bot. Voy. Herald, 168 = Mandevilla veraguensis.
verrucosa, *Willd. ex Roem. & Schult. Syst.* iv. 796.—Bras.
versicolor, *Mart. ex Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 38.—Bras.
verticillis, Buch.-Ham. ex Wall. Cat. n. 1666 = Rhynchodia Wallichii.
vestita, Roem. & Schult. Syst. iv. 894 = Chonemorpha vestita.
violacea, *Vell. Fl. Flum.* 110; ill. t. 81.—Bras.
virescens, *St. Hil. in Bull. Soc. Philom.* (1824) 77.—Bras.
virescens, Stadelm. in Flora, xxiv. (1841) t. Beibl. 63 = Macrosiphonia Martii.
volubilis, *Thunb. Diss. Echit.* 5.—Java.
xanthostoma, *Stadelm. in Flora*, xxiv. (1841) t. *Beibl.* 55.—Bras.
Zuccariniana, *Stadelm. l. c.* 76.—Bras.

ECHITHES, Thunb. Fl. Cap. ii. 150 (1818) = **Echites**, P. Br. (Apocyn.).

ECHIUM, Tourn. ex Linn. Gen. ed. I. 38 (1737).
BORAGINEAE, Benth. & Hook. f. ii. 863.
ARGYREXIAS, Rafin. Fl. Tellur. iv. 86 (1836).
ISOPLESION, Rafin. l. c. ii. (1836).
ISORIUM, Rafin. l. c. ii. 61 (1836).
LAREPHES, Rafin. l. c. iv. 86 (1836).
asuleatum, *Poir. Encyc.* viii. 664.—Ins. Canar.
acutifolium, Willd. ex Lehm. Asperif. 437 = rubrum.
afrocanum, Pers. Syn. i. 168 = Lobostemon fruticosus.
alberezianum, Debeaux, ex Nym. Consp. 515 = italicum.
albicans, *Lag. & Rod. in Anal. Cienc. Nat.* v. (1802) 289.—Hispan.
albicans, Schott, ex Roem. & Schult. Syst. iv. 726 = pomponium.
alopecuroideum, *DC. Prod.* x. 15.—Afr. austr.
altissimum, Jacq. Fl. Austr. v. 85. t. 16 = italicum.
ambiguum, DC. Cat. Hort. Monsp. (1813) 107 = brachyanthum, guanteum.
amoenum, *Pitch. & Mey. Ind. Sem. Hort. Petrop.* iv. 85.—Reg. Casp.
anglicum, Huds. Fl. Angl. ed. I. 70; Mill. Gard. Dict. ed. VIII. n. 1 = vulgare.
angustifolium, *Lam. Tabl. Encyc.* i. 412.—Hispan.; Marocc.
angustifolium, Link, ex Willk. & Lange, Prod. Fl. Hisp. ii. 466 = tuberculatum.
angustifolium, Mill. Gard. Dict. ed. VIII. n. 6 = creticum.
angustifolium, Thunb. Fl. Cap. ii. 248 = Lobostemon Swartii.
arenarium, *Guss. Ind. Hort. Boccad.* (1825); *Pl. Rar.* 88. t. 17.—Reg. Mediterr.
argenteum, Berg. Desc. Pl. Cap. 40; Linn. Mant. 302 = Lobostemon argenteus.
argenteum, Roth, Bot. Abh. 25 = Lobostemon ferocissimus.
armatum, MASS. ex C. Sm. sec DC. Prod. x. 18 = aculeatum.
asperrimum, Lam. Tabl. Encyc. i. 412; Bieb. Fl Taur.-Cauc. i. 185 = italicum.
Auberianum, *Webb & Berth. Phyt. Canar.* iii. 42.—Ins. Canar.
australe, Lam. Tabl. Encyc. i. 413 = creticum.
Benthami, Wall. Cat. n. 931 = Macrotomia Benthami.
Bergianum, E. Mey. ex DC. Prod. x. 6 = Lobostemon fruticosus.
bifrons, *DC. Cat. Hort. Monsp.* (1813) 107.—Ins. Canar.
Boissieri, Steud. Nom. ed. II. i. 540 = pomponium.
bonariense, Poir. Encyc. viii. 674 = violaceum.
brachyanthum, *Hornem. Hort. Hafn.* i. 182.—Ins. Madera.
brachystachyum, Pall. ex Roem. & Schult. Syst. iv. 726 = glomeratum?
callithyrsum, *Webb. ex C. Bolle, Ind. Sem. Hort. Berol.* (1857) *App.* i. 8.—Ins. Canar.
calycinum, *Viv. in Ann. Bot.* i. 11. 164.—Reg. Mediterr.
canaliculatum, Drège, ex DC. Prod. x. 11 = Lobostemon trichotomus.
candicans, DC. Cat. Hort. Monsp. 25 = Marianum.
candicans, Lam. Tabl. Encyc. i. 414 = gigantеum.
candicans, *Linn. f. Suppl.* 131.—Ins. Madera.
canescens, *Presl, ex Steud. Nom.* ed. II. i. 541.—Hab.?
capitatum, Linn. Mant. 42 = Lobostemon capitatus.
caudatum, Thunb. Prod. Fl. Cap. 33 = Lobostemon candatus.
clandestinum, *Pomel, Nouv. Mat. Fl. Atl.* 95.—Alger.
clavatum, Willd. ex Lehm. Asperif. 440 = rubrum.
collinum, Salisb. Prod. 114 = italicum.
Columnae, Ehrenb. ex Boiss. Fl. Orient. iv. 206 = glomeratum.
creticum, Forsk. Fl. Aegypt. Arab. 41 = sericeum.
creticum, Lam. Tabl. Encyc. i. 413 = violaceum.
creticum, *Linn. Sp. Pl.* 139; *DC. Prod.* x. 22.—Reg. Mediterr.
creticum, Pall. ex DC. Prod. x. 20 = rubrum.
creticum, Sibth. & Sm. Fl. Graec. ii. 69. t. 188 = calycinum.
creticum, Sims, Bot. Mag. t. 19, 34 = Simsii.
cynoglossoides, *Desf. Tabl.* ed. II. 86; *Poir. Encyc. Suppl.* v. 487.—Ins. Canar.
cyrenaicum, *Pacho, Voy. Marm.* t. 97.—Afr. bor.

ECHIUM:—
dahuricum, *G. Don, ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 541.—Dahuria.
dalmaticum, Tausch, in Flora, xiv. (1831) 20 = Rauwolfii?
Davazi, *Rouy, in La Natural.* (1892) n. 47; *ex Bolet. Broter.* ii. (1884) 23.—Lusitan.
Decaisnei, *Webb & Berth. Phyt. Canar.* iii. 49.—Ins. Canar.
decipiens, *Pomel, Nouv. Mat. Fl. Atl.* 91.—Alger.
densiflorum, *DC. Cat. Hort. Monsp.* (1813) 108.—Ins. Canar.
densiflorum, *Pomel, Nouv. Mat. Fl. Atl.* 92.—Alger.
diffusum, Gouan, Cat. Hort. Boccad. 23 = arenarium.
diffusum, Jan, ex Nym. Consp. 515 = mantimum.
diffusum, Sibth. & Sm. Fl. Graec. ii. 69. t. 182 = sericeum.
diffusum, Sieber, ex Steud. Nom. ed. II. i. 541 = setosum.
distachyum, Viv. Fl. Libyc. Spec. 8. t. 5. f. 1 = sericeum.
Ecklonianum, *DC. Prod.* x. 14.—Afr. austr.
elegans, Lehm. Asperif. 459 = sericeum.
elegans, Noe, ex Nym. Consp. 514 = vulgare.
elongatum, Lam. Tabl. Encyc. i. 412 = italicum
elongatum, Poir. Encyc. viii. 669 = glomeratum.
eriobotrum, *Pomel, Nouv. Mat. Fl. Atl.* 92.—Alger.
eriostachyum, *DC. Prod.* x. 14.—Afr. austr.
expansum, Boiss. & Haussk. ex Boiss. Fl. Or. Nov dec. ii. 7 = sericeum.
falcatum, Lam. Tabl. Encyc. i. 413 = Lobostemon glaber.
fastuosum, DC. Prod. x. 17 = Marianum.
fastuosum, Herb. Banks, ex Link, in Buch, Beschr. Canar. Inseln. 144 = Ins. Canar.
fastuosum, *Jacq. f. Eclog.* i. 41.—Ins. Canar.
fastuosum, *Salisb. Prod.* 114.—Ins. Tenerifia.
ferocissimum, Andr. Bot. Rep. t. 89 = Lobostemon ferocissimus.
ferox, Pers. Syn. i. 168 = Lobostemon argenteus.
flavum, *Desf. Fl. Atlant.* i. 165.—Hispan.; Afr. bor. occ.
foliosum, *Lehm. Asperif.* 412.—Ins. Canar.
Fontanesii, DC. Prod x. 24 = flavum.
formosum, Pers. Syn. i. 168 = Lobostemon formosus.
frondosum, Salisb. Prod. 114 = Lobostemon fruticosus.
fruticosum, Jacq. Hort. Schoenb. i. 15 = Lobostemon argenteus.
fruticosum, Linn Sp. Pl. 139 = Lobostemon fruticosus.
gaditanum, Boiss. Voy. Bot. Espagne, ii. 422 = tuberculatum.
gigantеum, *Linn. f. Suppl.* 131.—Ins. Tenerifia.
glabrum, Thunb. Prod. Pl. Cap. 33 = Lobostemon glaucophyllus.
glabrum, Vahl, Symb. Bot. iii. 22 = Lobostemon glaber.
glaucophyllum, Jacq. Coll. ii. 325 = Lobostemon glaucophyllus.
glomeratum, Boiss. Voy. Bot. Espagne, 424 = pomponium.
glomeratum, Ledeb. Fl. Ross. iii. 106 = lineariifolium.
glomeratum, *Poir. Encyc.* viii. 670.—Cilicia; Syria.
grandiflorum, Andr. Bot. Rep. (1798) t. 20 = Lobostemon formosus.
grandiflorum, Desf. Fl. Atlant. i. 166 = violaceum.
grandiflorum, Salisb. Prod. 115 = Megacaryon armenum.
hierrense, *Webb, ex C. Bolle, Ind. Sem. Hort. Berol.* (1857) *App.* i. 7.—Ins. Canar.
hispanicum, *Asso, Syn. Arag. Mant.* 162.—Hispan.
hispidum, Burm. f. Fl. Cap. Prod. 4 = Lobostemon capitatus.
hispidum, Sibth. & Sm. Fl. Graec. ii. 68. t. 181 = sericeum.
hispidum, Thunb. Prod. Fl. Cap. 33 = Lobostemon hispidus.
humile, Desf. Fl. Atlant. i. 165 = setosum.
hypertropicum, *Webb, in Hook. Niger Fl.* 155.—Ins. Cap. Verd.
incanum, *Thunb. Prod. Pl. Cap.* 33.—Afr. austr.
indicum, *Bernh. ex Steud. Nom.* ed. II. i. 541.—Hab.?
italicum, Boyé, ex DC. Prod. x. 18 = tuberculatum.
italicum, N. G. Gmel. Itin. i. 119. t. 19 = rubrum.
italicum, Hoffmgg. & Link, ex Steud. Nom. ed. II. i. 541 = lustanicum.
italicum, *Linn. Sp. Pl.* 139.—Europ.; Reg. Mediterr.
italicum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 577 = flavum.
italicum, Sieber, ex DC. Prod. x. 24 = pyramidatum.

ECHIUM:—
laetum, Salisb. Prod. 115 = vulgare.
laevigatum, Lam. Tabl. Encyc. i. 413 = Lobostemon glaucophyllus.
laevigatum, Linn. Syst. ed. X. 916 = Lobostemon laevigatus.
Lagascae, Boiss. Elench. 69 = pomponium.
Lagascae, *Roem. & Schult. Syst.* iv. 27.—Hispan
laxiophyllum, Link, Enum. Hort. Berol. i. 170 = mollc latifolium, *DC. Prod.* x. 14.—Afr. austr.
lineariifolium, *G. Koch, in Linnaea*, xxii. (1849) 629.—Reg. Caucas.
lineariifolium, Moench, Meth. Suppl. 149 = italicum.
lineatum, *Jacq. f. Eclog.* 62. t. 42.—Ins. Canar.
lineolatum, Webb & Berth. Phyt. Canar. iii. 46 = lineatum.
longiflorum, Bertol. Misc. i. 15 = Molikia angustifolia.
longiflorum, Dum.-Cours. Bot. Cult. ed. II. Suppl. vii. 147 = Lobostemon formosus.
longifolium, *Delile, Fl. Egypte*, 184. t. 16. f. 3.—Afr bor.; Arab.
longifolium, Hort. ex DC. Prod. x. 10 = Lobostemon lasiophyllus.
longifolium, Wiest, Herb. Un. Itin. n. 548; ex DC. Prod. x. 19 = sericeum.
longistamineum, Pourr. ex Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 29 = violaceum.
lucidum, Lehm. Sem. Hort. Bot. Hamb. (1827) = Lobostemon lucidus.
lusitanicum, Ail. Fl. Pedem. i. 52 = calycinum.
lusitanicum, DC. Prod. x. 20 = polyanthon.
lusitanicum, Georgi, Beschr. Russ. Reich. Nachtr. 260 = vulgare.
lusitanicum, *Linn. Sp. Pl.* 140.—Hispan.; Lusit.
lusitanicum, Mill. Gard. Dict. ed. VIII. n. 4 = violaceum.
luteum, Lapeyr. Hist. Abr. Pl. Pyr. 91 = italicum.
macranthum, Roem. & Schult. Syst. iv. 20 = violaceum.
macrophyllum, *Lehm. Asperif* 412.—Ins. Canar.
maderense, Hort. ex Steud. Nom. ed. II. i. 541 = candicans
margaritaceum, Lehm. Asperif. 446 = Rauwolfii.
Marianum, *Boiss. Diagn. Ser.* I. xi. 90.—Hispan.
maritimum, Guit. ex Willk. & Lange, Prod. Fl. Hisp ii. 467 = creticum.
maritimum, Tenore, ex Steud. Nom. ed. II. i. 541 = setosum.
maritimum, *Willd. Sp. Pl.* i. 788.—Reg. Mediterr. occ.
megalanthos, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 29 = violaceum.
megalanthum, Steud. Nom. ed. II. i. 541 = praec.
Mertensii, Lehm. Asperif. 458 = albicans.
Mensiesii, Lehm. Pugill. ii. 29 = Amsinckia intermedia
micranthum, Roem. & Schult. Syst. iv. 25 = tenue.
micranthum, Schousb. Vextr. Marokko, 75 = maritimum.
modestum, *Bail, in Journ. Bot.* xi. (1873) 373.—Marocco.
molle, Pet. in Reichb. Fl. Germ. Excurs. 836, in obs. = vulgare.
molle, *Poir. Encyc.* viii. 668.—Ins. Canar.
monstrosum, Schmidt, ex Steud. Nom. ed. II. i. 541 = vulgare.
montanum, *DC. Prod.* x. 15.—Afr. austr.
nervosum, [Dryand. in] Ait. Hort. Kew. ed. II. i. 300 = macrophyllum, candicans.
onosmaefolium, *Webb & Berth. Phyt. Canar.* iii. 46.—Ins. Canar.
onosmoides, *Pomel, Nouv. Mat. Fl. Atl.* 295.—Algeria.
orientale, Linn. Sp. Pl. 139 = Megacaryon armenum.
orientale, C. A. Mey. Verz. Pfl. Casc. 97 = amoenum.
orientale, Steph. Fl. Mosq. n. 124 = violaceum.
ovatum, Poir. Encyc. viii. 666 = calycinum.
pallidum, Salisb. Prod. 114 = candicans.
paniculatum, Drège, ex DC. Prod. x. 8 = Lobostemon paniculaeformis.
paniculatum, Lag. Gen. et Sp. Nov. 10 = Lagascae.
paniculatum, *Thunb. Prod. Pl. Cap.* 33.—Afr. austr.
papillosum, C. Koch, in Linnaea, xvii. (1843) 307 = lineariifolium.
papillosum, Lehm. Asperif. 441 = glomeratum?
papilosum, Thunb. Fl. Cap. ii. 8 = Lobostemon Swartzii.
parviflorum, Moench, Meth. 423; Roth, Catalect. fasc. ii. 14 = calycinum.
parviflorum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = vulgare.
parviflorum, Willd. Enum. Hort. Berol. 187 = tenue.

ECHIUM :—
Pavonianum, *Boiss. Diagn.* Ser. I. xi. 91.—Hispan.
petraeum, Tratt. Thes. Bot. 8 = Moltkia petraea.
pictum, Jan. ex Bolss. Fl. Orient. iv. 207 = Rauwolfia.
Pinianum, *Webb & Berth. Phyt. Canar.* iii. 44.—Ins Canar.
plantagineum, Jacq. Hort. Vindob. i. t. 45 = violaceum.
plantagineum, Kunze, in Flora, xxix. (1846) 761 = creticum.
plantagineum, *Linn. Mant.* ii 202.—Reg. Mediterr.
plantaginifolium, Linn. ex Moris, in Atti Ottav. Riun. Sc. Ital. (1847) 564 = plantagineum.
plantaginicides, Roem. & Schult. Syst. iv. 19 = violaceum.
polycaulon, *Boiss. Diagn.* Ser. I. xi. 92.—Hispan.
pomponum, *Boiss. Voy. Bot. Espagne*, t. 124.—Hispan. ; Afr. bor. occ.
prostratum, Delile, Fl. Egypte, 185. t. 17. f. 1; Poir. Encyc. Suppl. v. 486 = sericeum.
prostratum, Sieber, ex DC. Prod. x. 21 = arenarium.
prostratum, Tenore, Fl. Nap. i. 13 = calycinum.
Pseudo-violaceum, Schur, Enum. Pl. Transs. 470 = violaceum.
pumilum, Lehm. Pugill. i. 16 = sericeum ?
pustulatum, Lehm. Asperif. 448 = tuberculatum.
pustulatum, Sibth. & Sm. Fl. Graec. Prod i. 125 — Rauwolfii ?
pycnanthum, *Pomel, Nouv. Mat. Fl. Atl.* 29.—Algeria.
pyramidale, Lapeyr. Hist. Abr. Pl. Pyr. 90 = italicum.
pyramidatum, DC. Prod. x. 23 = italicum.
pyramidatum, *Dum.-Cours. Bot. Cult.* ed. II. vii. 147. —Afr. austr.
pyrenaicum, Linn. Mant. 234 = italicum.
pyrenaicum, *Mérgen. in Bull. Soc. Bot. Fr.* xii. (1865) 342.—Mont. Pyren.
Rauwolfii, *Delile, Fl. Egypte*, 195. t. 19. f. 3.—Aegypt.
regulariflorum, Ker-Gawl. Recens. 11. n. 42 = Lobostemon formosus.
rosmarinifolium, Schrank, ex Stend. Nom. ed. II. i 541 = Moltkia petraea.
rosmarinifolium, Vahl, Symb. Bot. iii. 22 = Lithospermum rosmarinifolium.
rosmicum, J. F. Gmel. Syst. i. 823 = rubrum.
rosulatum, *Lange, Ind. Sem. Hort. Haun.* (1854) 22. —Hispan.
rubrum, Forsk. Fl. Aegypt. Arab. 41 = sericeum.
rubrum, *Jacq. Fl. Austr.* v. 27.—Europ. austr. ; Reg. Caucas.
rubrum, Moench, Meth. 422 = creticum.
rupestre, Salisb. Prod. 134 = glaucatum.
rusticum, Roem. & Schult. Syst. iv. 28 = rubrum.
sabulicolum, *Pomel, Nouv. Mat. Fl. Atl.* 90.—Algeria.
salmanticum, Lag. Gen. et Sp. Nov. (1816) 10 = lusitanicum.
scabrum, Eckl. & Zeyh. ex DC. Prod. x. 6 = Lobostemon scaber.
scabrum, Thunb. Fl. Cap. ii. 11 = Lobostemon fruticosus, Thunbergianus.
Schöferi, Lange, in Syll. Ratisb. ii. (1828) 218 = vulgare.
sericeum, *Vahl, Symb. Bot.* ii. 35.—Reg. Mediterr. or.
setosum, Bory & Chaub. Fl. Pélop. 13 = sericeum ?
setosum, Delile, Fl. Egypte, 186. t. 17. f. 2 = creticum.
setosum, *Vahl, Symb. Bot.* ii. 35.—Aegypt.
Sibthorpii, Roem. & Schult. Syst. iv. 26 = sericeum.
simplex, *DC. Cat. Hort. Monsp.* (1813) 108.—Ins. Canar.
Stenii, *G. Don, ex A. DC. in DC. Prod.* x. 22.—Europ. austr.
spathulatum, Drège, ex DC. l. c. 10 = Lobostemon obovatus.
spathulatum, Viv. Fl. Libyc. Spec. 8. t. 9. f. 1 = maritimum.
sphaerocephalum, Vahl, Symb. Bot. iii. 22 = Lobostemon sphaerocephalus.
spicatum, Burm. f. Fl. Cap. Prod. 4 = Lobostemon spicatus.
splendens, *Medic. Bot. Beobachk.* (1788) 36.—Hab. ?
splendens, *DC. Prod.* x. 15.—Afr. austr.
Sprengelianum, *DC. l. c.* 16.—Afr. austr.
stenosiphon, *Webb, in Hook. Niger Fl.* 155. t. 15.—Ins. Cap. Verd.
strictissimum, Schur, Enum. Pl. Transs. 471 = italicum.

ECHIUM :—
strictum, *Linn. f. Suppl.* 130.—Ins. Canar.
strictum, Lam. Tabl. Encyc. i. 414 = virescens.
strigosum, Eckl. ex DC. Prod. x. 12 = Lobostemon thymeleaoides.
strigosum, Thunb. Fl. Cap. ii. 8 = Lobostemon strigosus.
Swartzii, Drège, ex DC. Prod. x. 5 = Lobostemon laevigatus.
Swartzii, Lehm. Asperif. 426 = Lobostemon Swartzii.
tenue, *Roth, Catalect.* ii. 18.—Hab. ?
thyrsiflorum, *Mass. ex Lisb, in Buch, Beschr. Canar. Ins.* 144.—Ins. Canar.
thyrsoideum, Vent. Jard. Malm. 98 = rubrum.
tinctorium, Viv. Pl. Aeg. Dec. n. 24 ; ex DC. Prod. x. 25 = Rauwolfii.
trichotomum, Drège, ex DC. Prod x. 10, 26 = Lobostemon cuneeus.
trichotomum, Thunb. Prod. Fl. Cap. 33 = Lobostemon virgatus.
trigonum, Thunb. l. c. = Lobostemon trigonus.
tryguthium, *Pomel, Nouv. Mat. Fl. Atl.* 94.—Algeria.
tuberculatum, Gilib. Fl. Lituan. i. 27 = vulgare
tuberculatum, *Hoffm. & Link, Fl. Port.* i. 183.—Europ. austr ; Afr. bor.
tubiferum, Poir. Encyc. viii. 668 = Lobostemon formosus.
Vahlii, Roem. & Schult. Syst. iv. 14, 715 = Lobostemon glaber.
valentinum, Lag. Gen. et Sp. Nov. (1806) 10 = flavum.
vaudense, Gremli, Beitr. Fl. Schw. 83 = vulgare ?
verecundum, Salisb. Prod. 135 = creticum.
verecundum, Viv. Fl. Aeg. Dec. n. 25 ; ex DC. Prod. x. 25 = setosum.
verrucosum, Thunb. Fl. Cap. ii. 8 = Lobostemon verrucosus.
violaceum, *Linn. Mant.* i. 42.—Europ. austr.
violaceum, Stokes, in With. Bot. Arr. ed. II. i. 201 = vulgare.
virescens, DC. Cat. Hort. Monsp. (1813) 107 = candicans.
virescens, *Vau, Elench.* 8 ; *ex DC. Prod.* x. 17, in nota. —Sicil.
virido-argenteum, *DC. Prod.* x. 18.—Afr. austr.
vulgare, Bourg. ex Reut in Bull. Soc. Bot. Fr. xi. (1864) 46 = polycaulon.
vulgare, Brot. Fl. Lusit. i. 289 = Rauwolfii.
vulgare, *Linn. Sp. Pl.* 139.—Europ.
Wierzbicki, Haberle, ex Reichb Fl. Germ. Excurs. i. 336 = vulgare.
Zeyheri, Eckl. ex Stend. Nom. ed. II. i. 542 = Lobostemon Zeyheri.

ECHTHRONEMA, Herb. in Lindl. Bot. Reg. (1843) Misc. 85 = **Eleyrinchium**, Linn. (Irid.).
californica, Herb. l. c. = S. californicum.
convoluta, Herb. l. c. = S. convolutum.
graminifolia, Herb. l. c. = S. graminifolium.
maculata, Herb. l. c. = S. graminifolium.
pumila, Herb. l. c. = Tapeinia magellanica.
tenuifolia, Herb. l. c. = S. tenuifolium

ECHTRUS, Lour. Fl. Cochinch. 844 (1790) = **Argemone**, Linn. (Papav.).
trivalvis, Stend. Nom. ed. II. i. 543 = A. mexicana.
trivialis, Lour. Fl. Cochinch. 844 = A. mexicana.

ECHYROSPERMUM, Schott, in Isis (1832), 1050. (Legumin.). (Quid ?)

ECKARDIA, Endl. Gen. Suppl. ii. 17, in syn. (1842) = **Paristeria**, Hook. (Orchid.).

ECKARTIA, Reichb. Nom. 58 (1841) = praec.

ECKEBERGIA, Batsch, Tabl. 58 (1802) = **Ekebergia**, Sparrm. (Meliac.).

ECKLONIA, *Stend. in Flora*, vii. (1829) 138. *CYPERACEAE* (Benth. & Hook. f. iii. 1062). TRIANOPTILES, Fenzl, in Endl. Gen. 118 (1836).
australis, *Stend. Nom.* ed. II. i. 542.—Austral.
capensis, *Stend. in Flora*, xii. (1829) 138.—Afr. austr.

ECKLONIA, Schrad. in Comm. Gotting. vii. (1832) 184 = praec.

ECLIPTA, Linn. Mant. ii. 157 (1771). *COMPOSITAE*, Benth. & Hook. f. ii. 361.
MICHELIUM, Forsk. Fl. Aegypt. Arab. 152 (1775).
PALRISTA, Rafin. New Fl. Am. ii. 48 (1836).
adpressa, Moench, Meth. Suppl. 245 = erecta.
alba, Hassk. Fl. Jav. Rar. 528 ; Miq. Fl. Ned. Ind. ii. 65 = erecta.
angustifolia, Presl, Bot. Bemerk. 105 = erecta.
arabica, Stend. Nom. ed. II. i. 542 = erecta.
brachypoda, Michx. Fl. Bor. Am. ii. 130 = erecta.
ciliata, Herb. Madr. ex Wall. Cat. n. 3214 = Carpesium cernuum.
ciliata, *Rafin. Fl. Ludov.* 69.—Am. bor.
dentata, Heyne, ex Wall. Cat. 3211 = erecta.
dichotoma, Rafin. New Fl. Am. ii. 42 = erecta.
dubia, Rafin. l. c. 40 = erecta.
elliptica, *DC. Prod.* v. 491.—Am. austr.
erecta, *Linn. Mant.* ii. 286.—Cosmop. trop.
filicaulis, Schum. & Thonn. Beskr. Guin. Pl. 290 = Spilanthes Acmella.
flexuosa, Rafin. New Fl. Am. ii. 41 = erecta.
heterophylla, Bartl. Ind. Sem. Hort. Gotting. (1857) 8. —Guatemala.
hirsuta, Bartl. in Linnaea, xiii. (1839) Litt. 95 = erecta.
humilis, H. B. & K. Nov. Gen. et Sp. iv. 264. t. 394 = Sabazia humilis.
integrifolia, Spreng. Syst. iii. 602 = Bellia integrifolia.
lanceolata, *DC. Prod.* v. 491.—Bras.
latifolia, Linn. f. Suppl. 376 = Blainvillea rhomboidea.
linearis, Otto, ex Sweet, Hort. Brit. ed. II. 308 = prostrata ?
longifolia, Schrad. in DC. Prod. v. 490 = erecta.
marginata, Stend. Nom. ed. II. i. 542 = erecta.
nutans, Rafin. New Fl. Am. ii. 42 = erecta.
paludicola, *Stend. Nom.* ed. II. i. 542.—Bras.
palustris, DC. Prod. v. 491 = erecta.
palustris, Vell. Fl. Flum. viii. t. 139 = paludicola.
parviflora, Wall. Cat. n. 3213 = erecta.
patula, Schrad ex DC. Prod. v. 491 = erecta.
pedunculosa, Remy, in C. Gay, Fl. Chil. iv. 113 = elliptica.
platyglossa, *F. Muell. Fragm.* ii. 135.—Austral.
procumbens, Michx. Fl. Bor. Am. ii. 130 = erecta.
prostrata, Linn. Mant. ii. 286 = erecta.
pumila, Rafin. New Fl. Am. ii. 42 = erecta.
punctata, Linn. Mant. ii. 286 = erecta.
pusilla, *DC. Prod.* v. 491.—Porto Rico.
quinquenervis, Vell. Fl. Flum. viii. t. 138 (Quid ?).—Bras.
scabriuscula, Wall. Cat. n. 3212 = Wedelia biflora.
sessilis, Sw. Prod. Veg. Occ. 114 = Eubrybia sessilis.
simplex, Rafin. New Fl. Am. ii. 41 = erecta.
spicata, *Spreng. Syst.* iii. 608.—China.
strumosa, Salisb. Prod. 305 = erecta.
sulcata, Rafin. New Fl. Am. ii. 41 = erecta.
thermalis, Bunge, Enum. Pl. Chin. Bor. 89 = erecta.
tinctoria, Rafin. New Fl. Am. ii. 41 = erecta.
trinervis, Vell. Fl. Flum. viii. t. 137 (Quid ?).—Bras.
undulata, Willd. Sp. Pl. iii. 2219 = erecta.
Zippelsana, Blume, Bijdr. 914 = erecta.

ECLOPES, Gaertn. Fruct. ii. 440. t. 169 (1791) = **Helhania**, L'Hérit. (Compos.).
aciculari, Cass. in Dict. Sc. Nat. xiv. 31 = R. pungens.
apiculata, DC. Prod. vii. 289 = R. quinquenervis.
buxifolia, Gaertn. Fruct. ii. 441 = R. speciosa.
centauroides, DC. Prod. vii. 288 = R. centauroides.
cuneata, Less. Syn. Comp. 279 = R. cuneata.
decussata, DC. Prod. vii. 288 = R. cuneata.
grenstaefolia, DC. l. c. = R. genistaefolia.
glutinosa, DC. l. c. = R. affinis.
palensea, Lam. Tabl. Encyc. iii. 268 = R. ericoides.
parallelinervos, Less. Syn. Comp. 280 = R. trinervis.
pumila, Less. l. c. 282 = R. pedunculata, *L'Hérit. punctata*, Cass. in Dict. Sc. Nat. xiv. 32 = R. genistaefolia.
pungens, Cass. l. c. 31 = R. quinquenervis.
schinolepis, DC. Prod. vii. 289 = R. speciosa.
sedifolia, DC. l. c. 286 = R. sedifolia.
speciosa, DC. l. c. 289 = R. speciosa.
stypholeides, DC. l. c. 289 = R. stypholeides.
subpungens, Cass in Dict. Sc. Nat. xiv. 31 = R. pungens.
tenuifolia, Stend. Nom. ed. II. i. 620, in syn. = Helichrysum teretifolium.
trinervis, Less. Syn. Comp. 281 = R. quinquenervis.
viscida, Gaertn. Fruct. ii. 440 = R. genistaefolia.

ECLOTORIPA, Rafin. Fl. Teller. iii. 40 (1836) = **Digera**, Forsk. (Amarant.).
aemula, Rafin. l. c. = sequ ?
fruticosa, Rafin. l. c. = D. arvensis.

ECLYPTA, E. Mey. Comm. Pl. Afr. Austr. 221 (1837) = **Eclipta**, Linn. (Compos.).

ECTADIOPSIS, Benth. in Benth. & Hook. f. Gen. ii. 741 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. l. c.
scutifolia, *Benth. l. c.*—Afr. austr.
brevifolia, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 78.—Ins. Socotr.
nigritana, *Benth. in Hook. Ic. Pl* t. 1187.—Afr. trop.
oblongifolia, *Benth. in Benth. & Hook. f. Gen.* ii. 741. —Afr. austr.
volubilis, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 78.—Ins. Socotr.

ECTADIUM, E. Mey. Comm. Pl. Afr. Austr. ed. II. 231 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 742.
oblongifolium, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 542 (err. typ. 442) = Ex. adiopsis oblongifolium.
virgatum, *E. Mey. Comm. Pl. Afr. Austr.* 188.—Afr. austr.

ECTASIS, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 156 = **Erica**, Linn.
Banksiana, D. Don, l. c. 156 = Erica Banksii.
bruniades, D. Don, l. c. = Erica bruniades
demissa, G. Don, Gen. Syst. iii. 837 = Erica demissa.
filiformis, G. Don, l. c. = Erica filiformis.
follicularis, G. Don, l. c. = Erica Petiveriі.
furfurosa, G. Don, l. c. = Erica monadelpha.
imbricata, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 156 = Erica imbricata.
melastoma, G. Don, Gen. Syst. iii. 837 = Erica Petiveriі.
monadelpha, G. Don, l. c. = Erica Banksii.
penicilliflora, G. Don, l. c. = Erica penicilliflora.
Petiveri, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 156 = Erica Sebana.
picta, G. Don, Gen. Syst. iii. 837 = Erica vestiflua.
placentaeflora, G. Don, l. c. = Erica tiaraeflora.
Plukenetii, D. Don, in Edinb. N. Phil. Journ. xvii, (1834) 156 = Erica Plukenetii.
pudibunda, G. Don, Gen. Syst. iii. 837 = Erica nutans.
Sebana, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 156 = Erica Sebana.
soccifera, G. Don, Gen. Syst. iii. 837 = Erica soccifera.
stellaris, G. Don, l. c. = Erica stellata.
turgida, G. Don, l. c. = Erica turgida.
vellerosiflora, G. Don, l. c. = Erica bruniades.
vestiflua, G. Don, l. c. = Erica vestiflua.

ECTEINANTHUS, T. Anders. in Journ. Linn. Soc. vii. (1864) 45 = **Isoglossa**, Oerst. (Acanthac.).
discoricatus, T. Anders. l. c.—Afr. austr.
Echlonianus, T. Anders. l. c.—Afr. austr.
origanoides, T. Anders. l. c.—Afr. austr.
ovatus, T. Anders. l. c.—Afr. austr.
prolixus, T. Anders. l. c.—Afr. austr.

ECTEMIS, Rafin. Sylva Tellur. 89 (1838) = **Cordia**, Linn. (Boragin.).
lutea, Rafin. l. c. = C. lutea.

ECTINOCLADUS, Benth. in Benth. & Hook. f. Gen. ii. 718 (1876). *APOCYNACEAE*, Benth. & Hook. f. l. c.

ECTOZOMA, Miers, in Ann. Nat. Hist. Ser. II. iv. (1849) 191 = **Juanulloa**, Ruiz & Pav. (Solanac.).
Pavonii, Miers, l. c.—Ecuador.

ECTROSIA, R. Br. Prod. 185 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1188.
agrostoides, *Benth. Fl. Austral.* vii. 634.—Austral.
Gaillveri, *P. Muell. Progrm.* viii. 301.—Austral.
leporina, *R. Br. Prod.* 186.—Austral.
Schultzii, *Benth. Fl. Australiae* vii. 633.—Austral.
spadicea, R. Br. Prod. 186 = leporina.
vaginata, *Trin. ex Steud. Nom.* ed. II. i. 543.—Hab.?

EDDYA, Torr. & Gray, in Pacif. Rail. Rep. ii. 170 (1855) = **Coldenia**, Linn. (Boragin.).
hispidissima, Torr. & Gray, l. c. = C. hispidissima.

EDEMIAS, Rafin. Fl. Tellur. ii. 49 (1836) = **Conyza**, Linn. (Compos.).
egyptiaca, Rafin. l. c. = C. aegyptiaca.
Gouani, Rafin. l. c. = C. Gouani.

EDGARIA, C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 113. *CUCURBITACEAE*.
darjeelingensis, *C. B. Clarke, l. c.* 114. f. 1–9.—Reg. Himal.

EDGEWORTHIA, Falc. in Proc. Linn. Soc. i. (1842) 129 = **Beptonia**, A. DC. (Myrsin.).
buxifolia, Falc. l. c. = B. buxifolia.

EDGEWORTHIA, Meisn. in Denkschr. Bot. Ges. Regensb. iii. (1841) 280. t. 6. *THYMELAEACEAE*, Benth. & Hook. f. iii. 198.
chrysantha, Lindl. in Journ. Hort. Soc. i. (1846) 148 = Gardneri.
Gardneri, *Meisn. in Denkschr. Bot. Ges. Regensb.* iii. (1841) 280. t. 6.—Reg. Himal.; China; Japon.
papyrifera, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 199 = Gardneri.

EDGWORTHIA, Lindl. in Journ. Hort. Soc. i. (1846) 148 = prace.

EDITELES, Rafin. Sylva Tellur. 108 (1838) = **Lythrum**, Linn.
thymifolia, Rafin. l. c. = L. thymifolia.

EDMONDIA, Cass. in Bull. Soc. Philom. (1818) 75; et in Dict. Sc. Nat. xiv. 352 (1819) = **Helichrysum**, Vaill. (Compos.).
bicolor, Cass. in Dict. Sc. Nat. xiv. 254.—Afr. austr
bracteata, Cass. l. c. = H. sesamoides.
spiendens, Cass. l. c. 253 = H. sesamoides.

EDMONDIA, Cogn. in DC. Monog. Phan. iii. 420 (1881). *CUCURBITACEAE*.
spectabilis, *Cogn. l. c.*—Venezuela.

EDMONSTONIA, Seem. Bot. Voy. Herald, 98. t. 18 (1852?) = **Tetrathyriastrum**, Poepp. (Violar.).
pacifica, Seem. l. c. = T. macrophyllum.

ENDOSMIA, Nutt. ex Torr. et Gray, Fl. N. Am. i. 612 (1840) = **Carum**, Linn. (Umbellif.).
Gairdneri, Torr. & Gray, l. c. = C. Gairdneri.
montana, Nutt. ex Torr. & Gray, l. c. = C. Gairdneri.
neurophylla, Maxim. in Bull. Acad. Pétersb. xviii. (1873) 286 = C. neurophyllum.
oregane, Nutt. ex. Torr. & Gray, Fl. N. Am. 612 = C. Gairdneri.
prunella, Nutt. ex Torr. & Gray, l. c. = C. Gairdneri.

EDRAIANTHUS, A. DC. in DC. Prod. vii. 448 (1839) = **Wahlenbergia**, Schrad. (Campanulac.).
carsicinus, Schott, Nyman, & Kotschy, Analect. Bot. 6 = W. tenuifolia.
caudatus, Reichb. Ic. Fl. Germ. xix. 109 = W. dalmatica.
croatscus, Kern. in Oestr. Bot. Zeitschr. xxii. (1872) 290, in textu = W. Kitaibelii.
dalmaticus, A. DC. in DC. Prod. vii. 449 = W. dalmatica.
graminifolius, A. DC. l. c. 448 = W. graminifolia.
Kitaibelii, A. DC. l. c. 449 = W. Kitaibelii.
Observianus, Boiss. Fl. Orient. iii. 886.—Reg. Caucas.
Pumilio, A. DC. in DC. Prod. vii. 449 = W. Pumilio.
serpyllifolius, A. DC. l. c. = Campanula serpyllifolia.
siculus, P. G. Strobl, in Flora, lxvi. (1883) 551 = W. graminifolia ?
tenuifolius, A. DC. in DC. Prod. vii. 449 = W. tenuifolia.
thymifolius, Vis. ex Nym. Consp. 486 = Campanula serpyllifolia.

EDRASTENIA, Rafin. Nem. Fl. Am. iv. 101 (1836) = **Oldenlandia**, Plum. (Rubiac.).
uniflora, Rafin. l. c. = O. glomerata.

EDRISSA, Endl. Suppl. ii. 58 (1842) (phalm. pro Edrisia, Rafin. §) = **Hedyotis**, Linn. (Rubiac.).

EDUSARON, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 371 = **Desmodium**, Desv. (1813) (Legumin.).
canadense, Medic. l. c. = D. canadense.

EDUSARUM, Stend. Nom. ed. II. i. 548 (1840) = prace.

EDWARDIA, Rafin. Specch. i. 158 (1814); ex DC. Prod. ii. 95 (1825) = **Cola**, Schott & Endl. (1832) (Sterculiac.).
turida, Rafin. l. c. = C. acuminata.

EDWARDSIA, Endl. Gen. 618 (1838) = EDWARSIA, Neck. = **Bidens**, Linn. (Compos.).

EDWARDSIA, Salisb. in Trans. Linn. Soc. ix. (1808) 298. t. 26 = **Sophora**, Linn. (Legumin.).
cassioides, Phil. in Bot. Zeit. xxxi. (1873) 741.—Chili.
chilensis, Miers, Trav. ii. 531, nomen; Bot. Reg. t. 1798 = S. macrocarpa.
chrysophylla, Salisb. in Trans. Linn. Soc. ix. (1808) 299 = S. chrysophylla.
denudata, DC. Prod. ii. 97 = S. denudata.
Fernandeziana, Phil. in Bot. Zeit. xiv. (1856) 642 = S. tetraptera.
grandiflora, Salisb. in Trans. Linn. Soc. ix. (1808) 299 = S. tetraptera.
hortensis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat Mosc. xii. (1860) 77 = Keyserlingia Buxbaumii.
kydaspica, Edgew. ex Aitch. in Journ. Linn. Soc viii. (1865) 64.—Ind. or.
intermedia, Tausch, in Flora, xvii. (1834) 489 = S. tetraptera.
Macnabiana, R. Grah. in Edinb. N. Phil. Journ. xxvi. (1838) 195 = S. tetraptera.
macrophylla, Wender. in Linnaea, v. (1830) 202 = S. tetraptera.
madrasspatana, Wight, Ic. t. 1054 = S. glauca, inter rupta.
masafuerana, Phil. in Bot. Zeit. xxxi. (1873) 743 = S. tetraptera.
microphylla, Salisb. in Trans. Linn. Soc. ix. (1808) 299 = S. tetraptera.
minima, Lodd. ex W. Bart. in Loud. Hort. Brit. Suppl ii. 631 = S. tetraptera.
mollis, Royle, Illustr. Bot. Himal. 196 = S. mollis.
myriophylla, Wender. in Linnaea, v. (1830) 202 = S. tetraptera.
nitida, DC. Prod. ii. 97 = S. nitida.
pervifolia, Wight, Ic. sub t. 1054.—China.
persica, Boiss. & Buhse, in Nouv. Mém. Soc. Nat Mosc. xii. (1860) 77 = Keyserlingia Buxbaumii.
Readeana, Phil. in Bot. Zeit. xxxi. (1873) 742 = S. tetraptera.
tetraptera, Poir. Encyc. Suppl. ii. 589, in obs. = S tetraptera.
Toromiro, Phil. in Bot. Zeit. xxxi. (1873) 743. — Chili.

EDWARSIA, Dum. Anal. Fam. 40 (1830) = prace.

EDWARSIA, Neck. Elem. i. 87 (1790) = **Bidens**, Linn. (Compos.).

EGANIA, Remy, in C. Gay, Fl. Chil. iii. 324 (1849) = **Chaetanthera**, Ruiz & Pav. (Compos.).
acerosa, Remy, l. c. 325.—Chili.
apiculata, Remy, l. c. 326.—Chili.
dioica, Remy, l. c. 327. t. 26. f. 1.—Chili.
pallida, Phil. in Linnaea, xxviii. (1856) 712.—Chili.
racemosa, Pritz. Ic. Ind. ii. 114 = *acerosa*.

EGENA, Rafin. Fl. Tellur. ii. 85 (1836) = **Clerodendrum**, Linn. (Verbenac.).
emirnensis, Rafin. l. c. = C. emirnense.

EGERIA, Neraud, in Freyc. Voy. Bot. 28 (1826), nomen. *RUBIACEAE*.

EGERIA, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 79 = **Elodea**, Michx. (Hydrochar.).
chilensis, Planch. ex Benth. & Hook. f. Gen. iii. 450 = Elodea callitrichoides.
densa, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 80 = Elodea densa.
Matthewsii, Planch. ex Benth. & Hook. f. Gen. iii. 450 = Elodea callitrichoides.
Naias, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 80 = Elodea Najas.

EGGERSIA, Hook. f. in Hook. Ic. Pl. t. 1401 (1883). *NYCTAGINEAE*, Benth. & Hook. f. iii. 1217.
buxifolia, *Hook. f. l. c.*—Ind. occ.

EGLETES, Cass. in Bull. Soc. Philom. (1817) 158. *COMPOSITAE*, Benth. & Hook. f. ii. 261.
EVENLIA, Kolelch. Icongr. Exot. iii. 15. t. 242 (1890).

EGLETES:—
Platystephium, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 90.
Xerobius, Cass. in Bull. Soc. Philom. (1817) 153; et in Dict. Sc. Nat. iix. 127 (1829).
arkansana, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 394 = Aphanostephus arkansanus.
bellidiflora, *Lesx. Syn. Comp.* 252.—Am. austr.
californica, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 36 = Erichphilum cespitosum.
domingensis, *Cass. in Dict. Sc. Nat.* xiv. 265.—Ind. occ.
floribunda, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 50.—Brsa.
glabrata, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 103.—Ind. occ.
humifusa, *Lesx. Syn. Comp.* 252.—Ecuador.
humilis, Torr. & Gray, Fl. N. Am. ii. 411 = Aphanostephus humilis.
Liebmanni, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 117.—Mexic.
obovata, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 103.—N. Granat.
ramonissima, A. Gray, Fl. Fendl. 71 = Aphanostephus ramonissimus.
texana, Engelm. ex A. Gray, in Bost. Journ. Nat. Hist. v. (1847) 222 = Aphanostephus arkansanus.
viscosa, *Lesx. Syn. Comp.* 252.—Am. austr.

HIRANDIA, Benth. & Hook. f. Gen. iii. 153 (1880) = EHRHARDIA, Scop. = **Aiouea**, Aubl. (Laurin.).

EHRARTHA, Beauv. Agrost. 60 (1812) = **Ehrharta**, Thunb. (Gramin.).

EHRARTIA, Benth. in Journ. Linn. Soc. xix. (1881) 56 sphalm. = EHRHARTIA, Weber = **Leersia**, Sw. (Gramin.).

EHRENBERGIA, Mart. Nov. Gen. et Sp. ii. 72. t. 163 (1827) = **Tribulus**, Tourn. (Zygophyll.).
tribuloides, Mart. l. c. 78 = T. brasiliensis.

EHRENBERGIA, Spreng. Neue Entdeck. ii. 129 (1821); et Syst. ii. 12 (1825) = **Amaioua**, Aubl. (Rubiac.).
ciliata, Spreng. ii. cc. 129, 188 = A. intermedia.

EHRETIA, Linn. Syst. ed. X. 936 (1759). *BORAGINEÆ*, Benth. & Hook. f. ii. 840.
Carmona, Cav. Ic. v. 22 (1799).
Desmophyla, Rafin. Sylva Tellur. 43 (1838).
Hilsenbergia, Tausch, ex Reichb. Consp. 117 (1828).
Lithothamnus, Zipp. ex Span. in Linnaea, xv. (1841) 384.
Surrisia, Rafin. Sylva Tellur. 43 (1838).
Traxilum, Rafin. l. c. 43 (1838).
abyssinica, *R. Br. in Salt. Abyss. App.* p. lxiv, nomen; *Fresen. in Flora*, xxi. (1838) 606.—Abyss.
acanthophora, DC. Prod. ix. 510 = Rochefortia acanthophora.
acuminata, *R. Br. Prod.* 497.—As et Austral. trop.
affinis, Wall. Cat. n. 900 = laevis.
amoena, *Klotzsch, in Peters, Reise Mossamb. Bot.* 246. —Afr. trop.
Andrieuxii, DC. Prod. ix. 510 = Bourreria Andrieuxii.
articulata, Walp. Rep. iv. 75, sphalm. = Citharexylum reticulatum.
aspera, Willd. Phyt. 4 = laevis.
asperula, *Zoll. & Mor. Syst. Verz. Zoll.* 52.—Java.
Bourreria, Blanco, Fl. Filip. ed. I. 127 = Blancoi.
Bourreria, Chapm. Fl. S. U. St. 329 = Bourreria succulenta.
Blancoi, *DC. Prod.* ix. 511.—Ins. Philipp.
bogotensis, Spreng. Syst. i. 648 = Aegiphila tomentosa?
Bourreria, Desf. in Ann. Mus. Par. i. (1802) 279 = Bourreria recurva.
Bourreria, Linn. Sp. Pl. ed. II. 375 = Bourreria succulenta.
Brandi, *Vatke, in Linnaea*, xliii. (1880–82) 215.—Afr. trop.
buxifolia, *Roxb. Pl. Corom.* i. 42. t. 67.—Ind. or.; Malaya.
calophylla, A. Rich. Fl. Cub. Fanerog. ii. 112 = Bourreria calophylla.
campestris, Salisb. Prod. 112 = tinifolia.
canarensis, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 143 = laevis.
capensis, *Meisen. in Flora*, xxvii. (1844) 330.—Afr. austr.

EHRETIA:—
cassinoides, A. Rich. Fl. Cub. Fanerog. ii. 113 = Bourreria cassinoides.
Catronya, Buch.-Ham. ex Wall. Cat. n. 7012 = laevis.
Championi, Wight & Gardn. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 142 = laevis.
ciliata, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. iii. (1869) 111 = elliptica.
cirrhosa, Lam. Encyc. i. 527 = Maripa scandens.
coromandeliana, Retz. ex DC. Prod. ix. 509 = buxifolia.
corymbosa, *Boj. Hort. Maurit.* 236, nomen; *DC. Prod.* ix. 506.—Ins. Comor.
cuneatensis, *DC. Prod.* ix. 511.—Venezuela.
cuneata, Wight, Ic. t. 1885 = Rhabdia lycioides.
Cutranga, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 142, sphalm. = laevis.
cymosa, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 129.—Afr. trop.
cymosa, Willd. ex Roem. & Schult. Syst. iv. 805 = Tournefortia hirsutissima.
dichotoma, *Blume, Bijdr.* 842.—Java.
dichotoma, Rottl. ex Wall. Cat. n. 904 = laevis.
Dzikzaki, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 234.—China.
diffusa, Vahl, ex DC. Prod. ix. 508 = cymosa.
divaricata, DC. Cat. Hort. Monsp. 108 = Bourreria divaricata.
domingensis, *DC. Prod.* ix. 508.—Ins. S. Doming.
dubia, Jacq. Obs. Bot. i. 19 = Cordia Collococca.
duplicata, Nees, ex Arn. in Linnaea, xi. (1837) 485 = Grabowskia duplicata.
Echioniana, Buek, ex Harv. Thes. Cap. i. 5 = hottentotica.
elliptica, *DC. Prod.* ix. 505.—Texas; Mexic.
exasperata, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. iii. (1867) 112 = elliptica.
exsucca, Bert. ex Griseb. Fl. Brit. W. Ind. 482 = Bourreria domingensis.
exsucca, Linn. Sp. Pl. ed. II. 375 = Bourreria exsucca. fasciculata, *H. B. & K. Nov. Gen. et Sp.* iii. 66.—Venezuela.
ferrea, Willd. Phyt. 4 = Maba buxifolia.
floribunda, Royle, Illustr. Bot. Himal. 306 = laevis.
formosa, *DC. Prod.* ix. 510 = Bourreria formosa.
glabra, Roth, ex Roem. & Schult. Syst. iv. 582; et Nov. Pl. Sp. 127 = Cordia Myxa.
grandiflora, *Poir. Encyc. Suppl.* ii. 8.—Ins. S. Doming.
guatimalensis, DC. Prod. ix. 507 = Bourreria grandiflora.
halimifolia, L'Hérit. Stirp. Nov. 45 = Grabowskia boerhaviaefolia.
halimifolia, Nees, ex Arn. in Linnaea, xi. (1837) 486 = Grabowskia obtusa.
havanensis, Roem. & Schult. Syst. iv. 805 = Bourreria havanensis.
heterophylla, Spreng. Syst. i. 648 = buxifolia.
Heynei, Roem. & Schult. Syst. iv. 582 = laevis.
hottentotica, *Burch. Trav. Afr.* ii. 147.—Afr. austr.
internodia, L'Hérit. Stirp. Nov. 47. t. 24 = petiolaris.
javanica, *Blume, Bijdr.* 842.—Java.
laevis, *Roxb. Pl. Corom.* i. 42. t. 66.—As. et Austral. trop.
laevis, Sieber, ex DC. Prod. ix. 506 = corymbosa.
lanceolata, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 148 = Wightiana.
lanceolata, *Vell. Fl. Flum.* 79; ii. t. 88.—Bras.
latifolia, *Loisel. Herb. Amat.* v. t. 350.—Mexic.
laxiflora, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 395.—Ins. Timor.
laxa, Jacq. Hort. Schoenb. i. 18. t. 41 = petiolaris.
linifolia, J. F. Gmel. Syst. 1542, sphalm. = tinifolia.
longiflora, *Champ. ex Benth. in Hook. Kew Journ.* v. (1853) 58.—China.
longifolia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 110.—Mexic.
lucida, *Span. in Linnaea*, xv. (1841) 384.—Ins. Timor.
lycioides, *Wall. ex G. Don, Gen. Syst.* iv. 891, in syn. —Ind. or.
macrophylla, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 348.—Reg. Himal.
mauritiana, *Willem. in Usteri, Ann. Bot.* xviii. (1796) 18.—Ins. Maurit.
membranifolia, *R. Br. Prod.* 497.—Austral.
microcarpa, *Tenore, Ind. Sem. Hort. Neap.* (1833) 46. —Hab. ?
microphylla, Lam. Tabl. Encyc. i. 495 = buxifolia.

EHRETIA:—
mollis, Wall. Cat. n. 7011 = Mappia ovata.
montevidensis, Spreng. Syst. i. 647 = Citharexylum barbinerve.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 249.—Afr. trop.
obovata, *R. Br. in Salt, Abyss. App.* p. lxiv, nomen; *DC. Prod.* ix. 507.—Ind. bor. occ.; Abyss.
obtusifolia, Hochst. ex DC. l. c. = obovata.
Ovmea, DC. l. c. 511 = acuminata.
ovalifolia, Hasak. Cat. Hort. Bog. Alt. 187 = acuminata.
ovalifolia, *Wight, Ic.* t. 1888.—Ind. or.
ovata, *Sieb. & Zucc. ex Teijsm. & Binn. Cat. Hort. Bog.* 130, nomen.—Japon.
parallela, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 143.—Burma.
petiolaris, *Lam. Encyc.* i. 527.—Ins. Mascar.
philippinensis, A. DC. in DC. Prod. ix. 504 = laevis.
pilicosta, *F. Muell. Progrm.* v. 90.—Austral.
polyantha, *DC. Prod.* ix. 508.—Ins. Philipp.
pubescens, Royle, Illustr. Bot. Himal. 306 = laevis.
punctata, Roth, ex Roem. & Schult. Syst. iv. 582; et Nov. Pl. Sp. 126 = laevis.
pyrifolia, D. Don, Prod. Fl. Nep. 109 = acuminata.
Radula, Poir. Encyc. Suppl. ii. 2 = Bourreria Radula.
resinosa, *Hance, in Journ. Bot.* xviii. (1880) 299. —China.
reticulata, Roem. & Schult. Syst. iv. 805 = Citharexylum reticulatum.
retusa, *Wall. Cat.* n. 903.—Burma.
revoluta, *DC. Prod.* ix. 507 = Bourreria revoluta.
rupestris, Salisb. Prod. 112 = Bourreria succulenta.
saligna, *R. Br. Prod.* 497.—Austral.
scabra, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 12.—Mexic.
scandens, Poir. Encyc. Suppl. iii. 590, in obs. = Maripa scandens.
serrata, Roxb. Hort. Beng. 17; Fl. Ind. i. 596 = acuminata.
setosa, *Roxb. Fl. Ind. ed. Carey*, ii. 349.—Ins. Samoa.
siamensis, *Teijsm. & Binn. Cat. Hort. Bog.* 139, nomen.—Siam.
spicata, Boj. Hort. Maurit. 236 = Maesa indica.
spinifex, Roem. & Schult. Syst. iv. 805 = Bourreria spinifex.
spinosa, *Jacq. Enum. Pl. Carib.* 14; *Linn. Sp. Pl.* ed. II. 375.—Ind. occ.
spinosa, Spreng. ex DC. Prod. ix. 510 = Rochefortia acanthophora.
sulcata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 109.—Ind. occ.
tenuifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 66.—N. Granat.
timorensis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 395 = laevis.
tinifolia, *Linn. Syst.* ed. X. 936.—Ind. occ.
tinifolia, Mart. & Gal. in Bull. Acad. Brux. xi. (1844) 332 = longiflola.
tomentosa, H. B. & K. Nov. Gen. et Sp. iii. 65 = Aegiphila tomentosa.
tomentosa, Lam. Tabl. Encyc. i. 495 = Bourreria tomentosa.
tomentosa, Roth, ex Roem. & Schult. Syst. iv. 582; et Nov. Pl. Sp. 126 = laevis.
triphylla, Hochst. in Flora, xxvii. (1844) 880 (= *Amerina triphylla*).—Afr. austr.
umbellulata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 344 = Ilex Godajam.
uniflora, *Roxb. Hort. Beng.* [84]; *Fl. Ind.* i. 598.—Ins Moluce.
volutina, DC. Prod. ix. 508 = Bourreria tomentosa.
vernalose, Spreng. ex Ragl. in Mart. Fl. Bras. xii. ii. 395 = Lithraea brasiliensis.
viminea, Wall. Cat. n. 906 = Rhabdia lycioides.
violacea, *Kunth, Ind. Sem. Hort. Berol.* (1846) 12.—Afr. austr.
virgata, Blanco, Fl. Filip. ed. I. 127 = acuminata.
virgata, Sw. Prod. Veg. Ind. Occ. 47 = Bourreria virgata.
Wallichiana, *Hook. f. & Thoms. ex Gamble, List Trees Darjeel.* 57; *et Fl. Brit. Ind.* iv. 143.—Reg. Himal.
Wightiana, *Wall. Cat.* n. 7013.—Ind. or.
Zephetiana, Buek, ex Drège, in Linnaea, xx. (1847) 196, nomen; et Harv. Thes. Cap. i. 5 = hottentotica.

EHRHARDIA, Scop. Introd. 107 (1777) = **Aiouea**, Aubl. (Laurin.).

EHRHARDTA, Hedw. f. Gen. 247 (1806) = **Ehrharta**, Thunb. (Gramin.).

EHRHARTA, Thunb. in Vet. Acad. Handl. Stockh. (1779) 216. t. 8. *GRAMINEAE*, Benth. & Hook. f. iii. 1187.

TROCHERA, Rich. in Journ. Phys. xiii. (1779) 295. t. 8.

abyssinica, *Hochst. in Flora*, xxxviii. (1855) 193.—Afr. trop.

acuminata, Spreng. Syst. ii. 114 = Tetrarrhena acuminata.

adscendens, Schrad. in Goett. Gel. Anz. (1821) iii. 2078 = calycina.

aemula, *Schrad. l. c.* 2077.—Afr. austr.

aphylla, *Schrad. l. c.*—Afr. austr.

aristata, Thunb. Prod. Pl. Cap. 66 = longiflora.

auriculata, Steud. in Flora, xii. (1829) 491 = calycina.

avenacea, *Willd. ex Schult. f. Syst.* vii. 1376.—Ins. Borbon.

Banksii, J. F. Gmel. Syst. 549 = longiflora.

barbinodis, *Nees, Fl. Afr. Austr.* 215.—Afr. austr.

brevifolia, *Schrad. in Goett. Gel. Anz.* (1821) iii. 2077. —Afr. austr.; Austral.

bulbosa, *Sm. Pl. Ic. Ined.* sub t. 33.—Afr. austr.

bulbosa, Steud. in Flora, xii. (1829) 491 = semnia.

calycina, Eckl. ex Nees, Fl. Afr. Austr. 208 = melicoides.

calycina, *Sm. Pl. Ic. Ined.* t. 33.—Afr. austr.

capensis, Thunb. in Vet. Acad. Handl. Stockh. (1779) 216. t. 8 = E. Mnemateta.

cartilaginea, Sm. Pl. Ic. Ined. sub t. 33 = E. Mnemateta.

caudata, *Munro, ex A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 490.—Japon.

clandestina, [Weber, in] Wigg. Prim. Fl Holsat. 64 = Leersia oryzoides.

Colensoi, *Hook. f. Fl. N. Zel.* i. 288. t. 65 a.—N. Zel.

contexta, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 111 = Tetrarrhena juncea.

diarrhena, F. Muell. Fragm. vii. 89 = Microlaena tasmanica.

digyna, Thunb. Prod. Pl. Cap. 66 = ramosa.

Diplax, F. Muell. Sel. Pl. Indust. Cult. 74 = Microlaena avenacea.

distichophylla, Labill. Nov. Holl. Pl. i. 90. t. 117 = Tetrarrhena distichophylla.

Drummondiana, Steud. Syn. Pl. Gram. 7 = Tetrarrhena laevis.

dura, *Nees, Fl. Afr. Austr.* 215.—Afr. austr.

Eckloniana, *Schrad. Rev. Alt.* n. 16; *ex Schult. f. Syst.* vii. 1376.—Afr. austr.

erecta, Lam. Encyc. ii. 847 = panicea.

festucacea, *Willd. ex Schult. f. Syst.* vii. 1378.—Madag.

geniculata, Thunb. Prod. Pl. Cap. 199 = calycina.

gigantea, Steud. in Flora, xii. (1829), 491 = villosa.

gigantea, *Thunb. l. c.* 192.—Afr. austr.

hispida, Beauv. ex Steud. Nom. ed. II. i. 544 = Tylaschys leucothrix.

juncea, Spreng. Syst. ii. 114 = Tetrarrhena juncea.

laevis, Spreng. l. c. 115 = Tetrarrhena laevis.

lanifolia, Schrad. in Goett. Gel. Anz. (1821) iii. 2078 = calycina.

longiflora, Eckl. ex Schult. f. Syst. vii. 1376 = Ecklon iana.

longiflora, *Sm. Pl. Ic. Ined.* t. 32.—Afr. austr. (Austral. introd.).

longifolia, *Schrad. in Goett. Gel. Anz.* (1821) iii. 2077.—Afr. austr.

longiseta, Schrad. l. c. 2078 = longiflora.

melicoides, *Thunb. Prod. Pl. Cap.* 192.—Afr. austr.

melicoides, Willd. ex Nees, Fl. Afr. Austr. 209 = calycina.

Microlaena, *Nees, l. c.* 217.—Afr. austr.

Mnemateta, *Linn. f. Suppl.* 209.—Afr. austr.

nudata, Lam. Encyc. ii. 846 = E. Mnemateta.

nutans, Willd. ex Nees, Fl. Afr. Austr. 209 = calycina.

Ottonis, Kunth, Rév. Gram. i. 9 = longiflora.

ovata, *Nees, in Linnaea*, vii. (1832) 286.—Afr. austr.

paniceo, *Sm. Pl. Ic. Ined.* t. 9.—Afr. austr.

paniciformis, *Nees, Fl. Afr. Austr.* 226.—Afr. austr.

paniculata, Poir. Encyc. Suppl. ii. 543 = calycina.

pilosa, Willd. ex Steud. Nom. ed. II. i. 544 = calycina.

punicea, J. F. Gmel. Syst. 1542, sphalm. = panicea.

pusilla, *Nees, Fl. Afr. Austr.* 225.—Afr. austr.

ramosa, Steud. in Flora, xii. (1829) 491 = aphylla.

ramosa, *Thunb. Prod. Pl. Cap.* 192.—Afr. austr.

EHRHARTA:—

ramosa, Willd. ex Nees, Fl. Afr. Austr. 209 = calycina.

rupestris, *Nees, l. c.* 227.—Afr. austr.

setacea, *Nees, l. c.* 228.—Afr. austr.

stipoides, Labill. Nov. Holl. Pl. i. 91. t. 118 = Microlaena stipoides.

stricta, *Nees, Fl. Afr. Austr.* 214.—Afr. austr.

tenacissima, Steud. Syn. Pl. Gram. 7 = Tetrarrhena juncea.

tenella, Spreng. f. Tent. Suppl. Syst. Veg. 11 = Sporobolus tenellus.

Thomsoni, *Petrie, in Trans. N. Z. Inst.* xii. (1880) 356.—N. Zel.

triandra, *Nees, Fl. Afr. Austr.* 221.—Afr. austr.

Trochera, Schrad. in Goett. Gel. Anz. (1821) iii. 2077 = bulbosa.

undulata, Nees, ex Steud. Nom. ed. II. i. 544; et Fl. Afr. Austr. 209 = calycina.

uniflora, *Burch. Trav.* i. 57, nomen.—Afr. austr.

unigiumis, F. Muell. in Trans. Phil. Soc. Vict. i (1855) 111 = Tetrarrhena juncea.

Urvillaeana, Kunth, Rév. Gram. i. 189. t. 6 = longiflora.

varicosa, *Nees, Fl. Afr. Austr.* 200.—Afr. austr.

versicolor, *Schrad. in Goett. Gel. Ans.* (1821) iii. 2078. —Afr. austr.

villosa, *Schult. f. Syst.* vii. 1374.—Afr. austr.

EHRHARTIA, [Weber, in] Wigg. Prim. Fl. Holsat. 63 (1780) = **Leersia**, Sw. (Gramin.).

clandestina, [Weber, in] Wigg. l. c. = L. oryzoides.

EICHHORNIA, Kunth, Enum. Pl. iv. 129 (1843). *PONTEDERIACEAE*, Benth. & Hook. f. iii. 686.

CABANISIA, Klotzsch, ex Schlecht. in Abh. Naturf. Ges. Halle, vi. (1862) 176.

LEPTOSOMUS, Schlecht. l. c. 174 (1862).

PIAROPUS, Kuhn. Fl. Tellur. ii. 81 (1886).

aquatica, Schlecht. in Abh. Naturf. Ges. Halle, vi. (1862) 177 = azurea.

azurea, *Kunth, Enum. Pl.* iv. 129.—Am. trop.

caracasana, Benth. & Hook. f. Gen. iii. 881 = Schultesiana.

cordifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 373.—Cuba.

crassipes, Schlecht. in Abh. Naturf. Ges. Halle, vi. (1862) 177 = speciosa.

crassipes, Solms, in DC. Monog. Phan. iv. 527 = speciosa.

gramminea, Seub. in Mart. Fl. Bras. iii. i. 91 = Heteranthera Seubertiana.

Martiana, *Seub. l. c.*—Am. trop.

Martiusiana, Walp. Ann. i. 848 = praec.

natans, *Solms, in Bremen, Abh.* vii. (1882) 254.—Afr. trop.

paniculata, Solms, in DC. Monog. Phan. iv. 530 = Martiana.

paradoxa, Solms, l. c. 531 = Schultesiana.

pauciflora, *Seub. in Mart. Fl. Bras.* iii. i. 91.—Bras.

Schultesiana, *Seub. l. c.* 95.—Bras.

speciosa, *Kunth, Enum. Pl.* iv. 131.—Am. trop.

subovata, Seub. in Mart. Fl. Bras. iii. i. 91 = Reussia subovata.

tricolor, Seub. l. c. 92 = Martiana.

EICHLERIA, Hartog, in Journ. Bot. xvi. (1878) 72 = Muricia, Hartog (Sapotac.).

albescens, Hartog, l. c. = M. albescens.

discolor, Hartog, l. c. = M. discolor.

EICHLERIA, Progel, in Mart. Fl. Bras. xii. ii. 518 (1877). *OXALIDEAE.*

Blanchetiana, *Progel, l. c.*—Bras.

lucida, *Progel, l. c.*—Bras.

EICHORNIA, A. Rich. Fl. Cub. Fanerog. ii. 373 (1853) = **Hohhornia**, Kunth (Ponted.).

EICHWALDIA, Ledeb. in Eichw. Pl. Casp. 88. t. 34 (1831) = **Beaumuria**, Linn. (Tamaricin.).

asiana, Ledeb. l. c. = R. oxiana.

persica, Boiss. Diagn. Ser. I. iii. 13 = R. silfolia.

EICOSIA, Blume, Fl. Jav. Praef. p. vii (1828), nomen. *ORCHIDEAE.*

EILEMANTHUS, Hochst. in Flora, xxix. (1846) 503 = Indigofera, Linn. (Legumin.).

strobiliferus, Hochst. l. c. = I. strobilifer.

EINADIA, Rafin. Fl. Tellur. iv. 121 (1836) = **Suaeda**, Forsk. (Chenop.).

linifolia, Rafin. l. c. = S. linifolia.

EINOMEIA, Rafin. Med. Fl. i. 62 (1828) = **Aristolochia**, Linn.

Berlandieri, Klotzsch, in Monatsb. Akad. Berl. (1859) 606 = A. micrantha.

bracteata, Rafin. Fl. Tellur. iv. 98 = A. bractesta.

brevipes, Klotzsch, in Monatsb. Akad. Berl. (1859) 606 = A. brevipes.

hastata, Klotzsch, l. c. 625. t. 2 = A. pentandra.

pentandra, Rafin. Med. Fl. i. 62 = A. pentandra.

EIMOMENIA, Meisn. Gen. 884 (1841) = praec.

EINOMERIA, Reichb. Nom. 68 (1841) = praec.

EISENMANNIA, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) i. Intell. 42; et xxv. (1842) i. Beibl. 134 = **Blainvillea**, Cass. (Compos.).

clandestina, Sch. Bip. ex Hochst. ll. cc. = B. rhombifolia.

EIZAGUIRREA, Remy, in C. Gay, Fl. Chil. iii. 401 (1849) = **Leucoria**, Lag. (Compos.).

Candollii, Remy, l. c. 402 = L. floribunda.

cirsioides, Phil. in Linnaea, xxvii. (1856) 719.—Chili.

tonchifolia, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 218.—Chili.

EKEBERGIA, Sparrm. in Vet. Acad. Handl. Stockh. (1779) 282. t. 9. *MELIACEAE*, Benth. & Hook f. i. 335.

ECKEBERGIA, Batsch. Tab. 58 (1802).

benguelensis, *Welw. ex C. DC. in DC. Monog. Phan.* i. 642.—Afr. trop.

capensis, *Sparrm. in Vet. Acad. Handl. Stockh.* (1779) 282. t. 9.—Afr. austr.

convallariaeodora, *Baill. Adansonia*, xi. (1874) 263.—Madag.

discolor, *O. Hoffm. in Linnaea*, xliii. (1880–82) 123 = Afr. trop.

fruticosa, *C. DC. in DC. Monog. Phan.* i. 644.—Afr. trop.

indica, Roxb. Hort. Beng. 33; Fl. Ind. ii. 392 = Cipadessa fruticosa.

integerrima, Wall. Cat. n. 7357 = Cipadessa fruticosa? Meyeri, *Presl, Bot. Bemerk.* 25.—Afr. trop.

Petitiana, A. Rich. Tent. Fl. Abyss. i. 105 = Ruppellana.

Ruppeliana, *A. Rich. l. c.*—Afr. trop.

senegalensis, *Juss. in Mém. Mus. Par.* xix. (1830) 254, 272.—Afr. trop.

Welwitschii, *Hiern, ex C. DC. in DC. Monog. Phan.* i. 643.—Afr. trop.

ELACANTHUS, F. Muell. in Linnaea, xxv. (1852) 410. *COMPOSITAE*, Benth. & Hook. f. ii. 481.

pusillus, *F. Muell. l. c.* 411.—Austral.

ELACHIA, DC. Prod. vii. 266 (1838) = **Chaetanthera**, Ruiz & Pav. (Compos.).

euphrasiaefolia, DC. l. c. = C. multicaulis.

renifolia, Remy, in C. Gay, Fl. Chil. iii. 315.—Chili.

spinulosa, Phil. in Linnaea, xxviii. (1856–57) 115.—Chili.

splendens, Remy, in C. Gay, Fl. Chil. ii. 315.—Chili.

ELACHOCROTON, F. Muell. in Hook. Kew Journ. ix. (1857) 17 = **Sebastiania**, Spreng. (Euphorb.).

asperocaccum, F. Muell. l. c. = S. Chamelaea.

ELACHOPAPPUS, F. Muell. Fragm. iii. 157 (1863) = **Myriocephalus**, Benth. (Compos.).

Rudallii, F. Muell. l. c. = M. Rudallii

ELACHOTHAMNOS, DC. Prod. v. 296 (1836) = **Minuria**, DC. (Compos.).

Cunninghamii, DC. l. c. = M. Cunninghamii.

ELAEAGIA, Wedd. Monog. Cinch. 94 (1849). *RUBIACEAE*, Benth. & Hook. f. ii. 50.

auriculata, *Hemsl. Diagn. Pl. Nov.* 23.—Costa Rica.

Mariae, *Wedd. Monog. Cinch.* 94.—Peruv.

utilis, *Wedd. l. c.*—N. Granat.

ELAEAGNUS, [Tourn.] Linn. Syst. ed. 1 (1735).
ELAEAGNACEÆ, Benth. & Hook. f. iii. 204.
LEPARGYREA, Raﬁn. in Am. Monthly Mag. (1818) 176.
LEPTARGYREIA, Schlecht. in DC. Prod. xiv. 607 (1857).
OCTABILLUM, Lour. Fl. Cochinch. 90 (1790).
acuminata, Hort. Angl. ex Link, Enum. Hort. Berol. i. 111 = latifolia.
Alingaro, *Schlecht. in DC. Prod.* xiv. 615.—Ins. Philipp.
angustifolia, Blanco, Fl. Filip. ed. I. 74 = latifolia.
angustifolia, *Linn. Sp. Pl.* 121.—Europ. ; As. bor.
arborea, Roxb. Fl. Ind. ed. Carey, i. 461 = latifolia.
argentea, Colla, Hort. Ripul. t. 28 = latifolia.
argentea, Moench, Meth. 688 = angustifolia.
argentea, *Pursh, Fl. Am. Sept.* i. 154.—Am. bor.
armata, Buch.-Ham. ex Schlecht. in DC. Prod. xiv. 611 = latifolia.
Breynianstrum, Fisch. ex Schlecht. l. c. = latifolia.
caudata, *Schlecht. in Linnaea*, xxxii. (1863) 201.—Ind. or.
communata, Bernh. in Allg. Thuer. Gartenz. ii. (1843) 137 = argentea.
conferta, Roxb. Hort. Beng. 11 ; Fl. Ind. i. 440 = latifolia.
crispa, Thunb. Fl. Jap. 66 = multiflora.
Cumingii, Schlecht. in DC. Prod. xiv. 613 = latifolia.
cyanea, *Ait. ex Steud. Nom.* i. 544.—Hab. ?
dactyliformis, Hort. ex Schlecht. in DC. Prod. xiv. 615 = angustifolia.
dulcis, *Roxb. Hort. Beng.* 11, nomen.—Ind. or.
edulis, Siebold, ex E. May, in Rev. Hortic. (1876) 18 ; Lavall. Arbor. Segrez 1 = multiflora.
elliptica, Heynе, ex Wall. Cat. n. 3048 = latifolia.
erinaneusis, Fisch. ex Schlecht. in DC. Prod. xiv. 615 = angustifolia ?
fasciculata, *Griff. Notul.* iv. 374.—Assam.
ferruginea, Rich. in Mém. Soc. Hist. Nat. Par. i. (1792) 367, 404 = latifolia.
fusca, *Pepon, ex Low. Hortic. Univ.* iii. 304.—Hab. ?
Gaudichaudiana, *Schlecht. in DC. Prod.* xiv. 612.—Cochinch.
glabra, Hort. ex C. Koch, Dendrol. ii. 1. 389 = argentea.
glabra, *Thunb. Fl. Jap.* 67.—Japon.
gonyanthes, *Benth. in Hook. Kew Journ.* v. (1853) 197.—China.
grandifolia, Bog. ex Schlecht. in DC. Prod. xiv. 612 = latifolia.
Grijsii, *Hance, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 227.—China.
Gussoni, Gasparr. in Ann. Civ. i. (1883) 118 = latifolia.
hortensis, Bleb. Fl. Taur. Cauc. i. 112 = angustifolia.
incana, Lam. Fl. Fr. iii. 476 = angustifolia.
inermis, Mill. Gard. Dict. ed. VIII. n. 2 = angustifolia.
javanica, *Blume, Bijdr.* 638.—Java.
Kologa, Schlecht. in DC. Prod. xiv. 611 = latifolia.
lastevirens, *Lindberg, in Act. Soc. Sc. Fenn.* x. (1871) 125.—Hab. ?
latifolia, *Linn. Sp. Pl.* 121.—As. trop.
latifolia, Lour. Fl. Cochinch. 89 = Loureiri.
longipes, A. Gray, in Mem. Am. Acad. N. S. vi. (1858) 406 = multiflora.
Loureiri, *Champ. in Hook. Kew Journ.* v. (1853) 196.—Hongkong.
macrophylla, *Thunb. Fl. Jap.* 67.—Japon.
macrophylla, Wall. ex DC. Prod. xiv. 612 = latifolia.
Maximowiczii, Wall. Cat. n. 4081 = angustifolia.
multiflora, *Thunb. Fl. Jap.* 66.—Japon.
nepalensis, Salm-Dyck, Hort. Dyck. 99 = latifolia ?
Oldhami, *Maxim. in Bull. Acad. Petersb.* xv. (1871) 377.—Ins. Formosa.
orientalis, Linn. Mant. i. 41 = angustifolia.
pallifolia, Hort. ex C. Koch, Dendrol. ii. 1. 390 = umbellata.
parasyrensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 156 ; *Contrib.* 49.—Reg. Argent.
parvifolia, Wall. Cat. n. 4096, Royle, Illustr. Bot. Himal. 232. t. 51. f. 1 = umbellata.
Perrottetii, *Schlecht. in DC. Prod.* xiv. 613.—Ins. Philipp.
punctata, Schlecht. l. c. 610 = latifolia.
pungens, *Thunb. Fl. Jap.* 68.—Japon.
reflexa, E. Morr. & Decne. in Bull. Acad. Brux. iii. (1836) 171 = umbellata.
rigida, *Blume, Bijdr.* 638.—Java.

ELAEAGNUS :—
rotundifolia, Gagn. in Rev. Hortic. (1870–71) 540 = multiflora ?
salicifolia, *D. Don, ex Loud. Encyc. Trees & Shrubs*, 697. t. 1366.—Hab. ?
Simonii, Carr. in Rev. Hortic. (1869) 100 = latifolia.
songorica, *Fisch. ex Traxvo. in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 87.—Sibir.
spadicea, Savi, in Mem. Soc. Ital. Moden. xxi. (1837) 175 = latifolia.
spinosa, Linn. Cent. ii. 9 ; Amoen. Acad. iv. 305 = angustifolia.
tenuifora, Benth. in Hook. Kew Journ. v. (1853) 197 = glabra.
Thwaitesii, Schlecht. in DC. Prod. xiv. 611 = latifolia.
tiliarnis, Fisch. & Vis. Ort. Bot. Padov. 77 = latifolia.
tomentosa, Moench, Meth. 688 = angustifolia.
trifora, Roxb. Hort. Beng. 11 ; Fl. Ind. i. 439 = latifolia.
umbellata, *Thunb. Fl. Jap.* 66. t. 14.—Japon.
undulata, Hort. ex Ind. Sem. Hort. Genuens (1855) 14, in nota = Olearia Forsteri.
Wallichiana, Schlecht. in DC. Prod. xiv. 612 = latifolia.

ELAEAGRUS, Pall. Fl. Ross. i. 1. 23, 25 (1789) = **Elaeagnus**, Tourn.

ELAEIS, Jacq. Select. Am. 280. t. 172 (1763). *PALMAE*, Benth. & Hook. f. iii. 944.
ALPONSIA, H. B. & K. Nov. Gen. et Sp. i. 306 (1815).
BARCELLA, Drude, in Mart. Fl. Bras. iii. 11. 459 t. 106 (1881).
COROZO, Jacq. ex Glaeke, Prael. 42 (1792).
guineensis, *Jacq. Select. Am* 280.—Afr. trop.
melanococca, *Gaertn. Fruct.* i. 18. t. 6. f. 2.—N Granat.
montana, *Page, ex Steud. Nom.* ed. II. i. 545.—Jamaica.
occidentalis, *Sw. Fl. Ind Occ* i 619.—Ind. occ.
odora, *Trail, in Journ. Bot.* xv. (1877) 81.—Bras.
pernambucana, *Lodd. ex G. Don, in Loud. Hort. Brit* 399.—Bras.
spectabilis, *Lodd ex Sweet, Hort. Brit.* ed. III. 716.—' Ind. or.'

ELAEOCARPUS, Burm. ex Linn. Diss. Dsas. 11 (1747) ; Amoen. Acad. i. 402 (1749). *TILIACEAE*, Benth. & Hook. f. i. 239.
ACRATIUM, DC. Prod. i. 519 (1824).
ACRONODIA, Blume, Bijdr. 123 (1825).
ACROZUS, Spreng. Syst. iv. Cur. Post. 149 (1827).
ADENODUS, Lour. Fl. Cochinch. 294 (1790).
BEYTHEA, Endl. Gen. 1011 (1840).
CRASPEDUM, Lour. Fl. Cochinch. 386 (1790).
DICERA, Forst. Char. Gen. 79. t. 40 (1776).
ERIOSTEMUM, Colla, ex Steud. Nom. ed. II. i. 588 (1840).
GANITRUS, Gaertn. Fruct. i. 271. t. 139 (1791).
LOCHNERIA, Scop. Introd. 271 (1777).
MISIPUS, Raﬁn. Sylva Tellur. 60 (1838).
MONOCERA, Jack, in Malay Misc. i. (1820) v. 42 ; reimp. in Hook. Bot. Misc. ii. (1831) 55.
PERINKA, Raﬁn. Sylva Tellur. 60 (1838).
SKIDANTHERA, Raﬁn. l. c. (1838).
Acuminatus, *Mast. in Hook. f. Fl. Brit. Ind.* i. 408.—Malacca.
acuminatus, *Wall. Cat.* n. 2676.—Ind. or.
adenophyllus, Wall. l. c. n. 6346 = oblongus.
adenopus, *Miq. Fl. Ind. Bat.* i. ii. 209.—Java.
alaternoides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* viii. (1861) 201.—N. Caled.
alnifolius, *Baker, in Journ. Linn. Soc. xx.* (1883) 107.—Madag.
amoenus, *Thw. Enum. Pl. Zeyl.* 83.—Ind. or.
amygdalinus, Wall. Cat. n. 6357 = robustus.
angustifolius, *Blume, Bijdr.* 120.—Java.
angustifolius, Wight, Illustr. 84 = oblongus.
apiculatus, *Mast. in Hook. Fl. Brit. Ind.* i. 407.—Malacca.
argyrodes, *Hance, in Journ. Bot.* xv. (1877) 330.—China.
aristatus, *Roxb. Hort. Beng.* 42 ; *Fl. Ind.* ii. 599.—Ind. or.
aristatus, Wall. Cat. n. 3665 B = robustus.

ELAEOCARPUS :—
armhemicus, *F. Muell. Desc. Not. Papuan Pl.* i. 6.—N. Guin.
atuvirens, *Zoll. & Mor. Syst. Verz. Zoll.* 26.—Java.
Bancroftii, *F. Muell. & Bailey, in Proc. Roy. Soc Queensl.* ii. 11 (1885).—Australi.
Baudouini, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 301.—N. Caled.
bifida, *Hook. & Arn. Bot. Beech. Voy.* 110. t. 24.—Ins Sandvic.
bisectiaris, Roxb. ex Wight & Arn. Prod. 83 = tuberculatus.
bracteatus, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 48.—Burma.
cassinoides, *A. Gray, Bot. U. St. Expl. Exped.* i. 204.—Ins. Fiji.
ceylanicus, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 323.—Zeylan.
Chackrovla, Mast. in Hook. f. Fl. Brit. Ind. i. 408, sphalm. = seq.
Chakrovia, Buch. Ham. in Wall. Cat. n. 2674 (sp. dub.).—Ind. or.
chinensis, *Hook. f. ex Benth. Fl. Hongk.* 48.—China.
cupalliferus, Reta. Obs iv 27 = Vateria acuminata.
coriaceus, *Hook. Ic. Pl.* t. 154.—Ind. or.
Corsonianus, *G. Don, in Loud. Gard. Mag.* xviii. (1842) 370.—Ins. Batagoda
Cumingii, *Turcz. in Bull. Soc. Nat Mosc.* xix. (1846) ii. 491.—Ins. Luzon.
cuneatus, *Wight, Illustr.* i. 83.—Iqd. or.
Cunninghamii, Raoul, Choix Fl. N. Zéland, 25 = dentatus.
cyaneus, *Sims, Bot. Mag* t. 1737.—Austral.
cyanocarpus, *Maingay, ex Mast. in Hook. f. Fl. Brit Ind.* i. 406.—Malacca.
dasyandrus, *Baker, in Journ. Linn Soc* xx. (1883) 108.—Madag.
dentatus, *Vahl, Symb. Bot.* iii. 66.—N. Zel.
Deplanchei, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* x. (1863) 477.—N. Caled.
Dicera, Vahl, Symb. Bot. iii. 67 = dentatus.
dioicus, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) 498.—Japon.
edulis, *Teysm. & Binn. in Tijdschr. Nederl. Ind* xxvii. (1864) 89.—Ins. Amboin.
ellipticus, *Sm. in Rees, Cycl.* xii. n. 3.—Hab. ?
ferrugineus, *Steud. Nom.* ed. II. i. 545 ; *Beddome, Fl Syle.* t. 112.—Ind. or.
Finlaysonianus, Wall. Cat. n. 6851 ¬ E. Varunna.
fissistipula, *Miq. Fl. Ind. Bat.* i. ii. 210 —Java.
floribundus, *Blume, Bijdr.* 120.—As. trop
foveolatus, F. Muell. Fragm. v. 157 = obovatus.
fruticosus, *Roxb. Hort. Beng.* [98] ; *Fl. Ind.* ii. 600.—Ins. Moluce.
Ganitrus, *Roxb. l. c.* 42, 599.—Ind. or. ; Malaya.
geminiflorus, *Brongn. & Gris, in Bull. Soc. Bot. Fr* x. (1863) 477.—N. Caled.
glaber, *Blume, Bijdr.* 122.—Java.
glabrescens, *Mast. in Hook. f. Fl. Brit. Ind.* i. 403.—Malacca.
glandulifer, *Mast. in Hook. f. l. c.* 406.—Zeylan.
glandulosus, Wall. Cat. n. 6859 = oblongus.
Graeffei, *Seem. in Journ. Bot.* ii. (1864) 76.—Ins. Fiji
Grahami, *F. Muell. Fragm.* x. 3.—Austral.
grandiflorus, *Bg. Hort. Maurit.* 45.—Ins. Maurit.
grandiflorus, *Sm. in Rees, Cycl.* xii. n. 5.—Austral.
grandifolius, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 294.—Burma.
grandis, *F. Muell. Fragm.* ii. 81.—Austral.
Griffithii, *A. Gray, in Bot. U. St. Expl. Exped.* i. 203.—Burma : Malaya.
Grijsii, C. Muell. in Walp. Ann. vii. 455, sphalm. = Elaeagnus Grijsii.
grossus, Wall. Cat. n. 2661 = floribundus.
Guillaumii, *Vieill. in Bull. Soc. Linn. Normand.* ix (1865) 336.—N. Caled.
Helferi, *Kurz, ex Mast. in Hook. Fl. Brit. Ind.* i. 402.—Burma.
Henryi, *Hance, in Journ. Bot.* xxiii. (1885) 322.—China.
Hinau, A. Cunn. in Ann. Nat. Hist. iv. (1840) 23 = dentatus.
holopetalus, *F. Muell. Fragm.* ii. 148.—Australi.
Hookerianus, *Raoul, Choix Pl. N. Zéland*, 36.—N. Zel.
hygrophilus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1874) ii. 133.—Burma.
hypadenus, *Miq. Fl. Ind. Bat. Suppl.* i. 406.—Sumatra.
integer, *Wall. Cat.* n. 3668, 3678.—Burma ; Malaya.

ELAEOCARPUS:—
integerrimus, *Lour. Fl. Cochinch.* i. 412.—Cochinch.
integrifolius, Blanco, Fl. Filip. ed. II. 306 = Chailletia gelonioides.
integrifolius, Lam. Encyc. ii. 604 = serratus.
integrifolius, Seber, ex Presl, in Oken, Isis, xxi. (1828) 373 = Ochna integrifolia.
integripetalus, *Miq. Fl. Ind. Bat. Suppl.* i. 407.—Sumatra.
isoticlius, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 31.—Ins. Philipp.
Jackianus, *Wall. Cat.* n. 2679.—Singapore.
japonicus, *Siebold, in Verh. Batav. Gen.* xii. (1830) 63. —Japon.
japonicus, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) 11. 492.—Japon.
lanceaefolius, Wall. Cat. 6858, 9026 = serratus *vel* cuneatus.
lanceaefolius, *Roxb Hort. Beng.* 42; *Fl. Ind.* ii. 598. —As. trop.
lanceolatus, *Blume, Bydr.* 119.—Java.
lanceolatus, Wall. Cat. 2667, partim = lanceaefolius.
laurifolius, *A. Gray, Bot. U. St. Expl. Exped.* i. 208.— —Ins. Fiji.
Lenormandii, *Vieill. in Bull. Soc. Linn. Normand.* ix. (1865) 385.—N. Caled.
leptostachyus, *Wall. Cat.* n. 2672.—Burma; Penang.
littoralis, *Teijsm. & Binn. ex Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 123.—Tenasserim.
Lobbianus, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 236 = floribundus.
longifolius, *Blume, Bydr.* 120.—Java.
longifolius, *Wall. Cat.* n. 6682.—Burma.
lucidus, *Roxb. Hort. Beng.* 42; *Fl. Ind.* ii. 600.—Ind. or.
macrocarpus, *Miq. Fl. Ind. Bat. Suppl.* i. 406.—Sumatra.
macrophyllus, *Blume, Bydr.* 122.—Java.
micranthus, *Teijsm. & Binn. Cat. Hort. Bog.* 390.— Menado.
micranthus, *Vieill. in Bull. Soc. Linn. Normand.* ix. (1865) 392.—N. Caled.
Milnei, *Seem. Fl. Vit.* 28.—Ins Fiji.
moluccanus, *Scheff. in Tydschr. Nederl. Ind.* xxxi. (1870) 345.—Ins. Moluce.
Monocera, *Cav. Ic.* vi. t. 501.—As. trop.
monocerotdes, *Wight, Illustr.* i. 84.—Ind. or.
montanus, *Thw. Enum. Pl. Zeyl.* 32.—Zeylan.
multiflorus, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 31.—Ins. Philipp.
Munroii, *Mast. in Hook. f. Fl. Brit. Ind.* i. 407.—Ind. or.
nitidus, *Jack, in Malay Misc.* i. (1820) n. v. 41.— Malaya.
oblongus, *Gaertn. ex Sm. in Rees, Cycl.* xii. n. 9.— Ind. or.; Malaya.
oblongus, Wall. Cat. n. 2677 = floribundus, robustus.
obovatus, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 328 = coriaceus.
obovatus, *G. Don, Gen. Syst.* i. 559.—Austral.
obtusus, *Blume, Bydr.* 118.—Java.
opposititolius, *Miq. Fl. Ind. Bat.* i. ii. 211.—Ins. Amboin.
ovalifolius, Wall. Cat. n. 2665 = robustus.
ovalis, *Miq. Fl. Ind. Bat. Suppl.* i. 406.—Sumatra.
oviformis, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* x. (1863) 477.—N. Caled.
paucinervatus, *Wall. Cat.* n. 2668.—Malaya.
parvifolius, A. Rich. Sert. Astrol. 67. t. 24 = obovatus.
parviflorus, *Span. in Linnaea*, xv. (1841) 176.—Ins. Timor.
parvifolius, *Wall. Cat.* n. 2662—Malaya.
pauciflorus, *A. Rich. Voy. Astrol.* 67.—N. Zel.
pauciflorus, Walp. Rep. i. 364 = obovatus.
pedunculatus, Labill, Nov. Holl. Pl. ii. 15. t. 155 = Aristotelia pedunculata.
pedunculatus, *Wall. Cat.* n. 2673.—Malaya.
Perrottetiana, DC. Prod. i. 519 = oblongus.
persicifolius, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* (1861) 202.—N. Caled.
petiolatus, *A. Gray, Bot. U. St. Expl. Exped.* i. 208.— Burma.
petiolatus, Wall. ex Steud. Nom. ed. II. i. 545 = integer.
photiniaefolia, *Hook. & Arn. Bot. Beech. Voy.* 350.— Chins.
pilosus, *Roxb. Hort. Beng.* 42.—Ins. Mascal.
Pirincura, Wall. Cat. n. 2664. partim = serratus.
polystachyus, *Wall. l. c.* n. 2671.—Malaya.

ELAEOCARPUS:—
prinodes, *Voigt, Hort. Suburb. Calc.* 128.—Hab.?
pruniifolius, *Wall. Cat.* n. 2669.—Ind. or.
pubescens, Hook. Ic. Pl. t. 155 = subvillosus.
pulchellus, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 300.—N. Caled.
punctatus, *Wall. Cat.* n. 6676.—Singapore.
pyriformis, *A. Gray, Bot. U. St. Expl. Exped.* i. 205.— Ins. Fiji.
quercifolius, *Baker, in Journ. Linn. Soc.* xx. (1883) 108.—Madag.
resinosus, *Blume, Bydr.* 122.—Java.
reticulatus, Sm. in Rees, Cycl. xii. n. 6 = cyaneus.
retusus, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* x. (1863) 477.—N. Caled.
rhodanthus, *Baker, in Journ. Linn. Soc.* xx. (1883) 107.—Madag.
robustus, *Roxb. Hort. Beng.* 42; *Fl. Ind.* ii. 597.—As. trop.
robustus, Wall. Cat. n. 2644 A = integer?
rotundifolius, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* viii. (1861) 202.—N. Caled.
rufovestitus, *Baker, in Journ. Linn. Soc.* xx. (1883) 108.—Madag.
rugosus, *Roxb. Hort. Beng.* 42; *Fl. Ind.* ii. 596.—Ind. or.
rugosus, Wall. Cat. n. 2658 A, B = tuberculatus.
ruminatus, *F. Muell. Fragm.* viii. 1.—Austral.
sericeus, *Baker, in Journ. Linn. Soc.* xx. (1883) 106. —Madag.
sericocarpus, *F. Muell. Fragm.* vi. 171 — Austral.
Seringii, *Montr. in Mém. Acad. Lyon*, x. (1860) 183. —N. Caled.
serratus, Benth. in Hook. Kew Journ. iii. (1851) 264 = lanceaefolius.
serratus, Heyne, in Wall. Cat. n. 2665 = oblongus.
serratus, *Linn. Sp. Pl.* 515.—Ind. or.; Malaya.
serratus, Lam. f. Suppl. 286 = dentatus.
serratus, Retz. Obs. iv. 27 = oblongus ?
serratus, Roxb. ex Wall. Cat. n. 2666 A, B, partim = floribundus.
serrulatus, Benth. in Hook. Kew Journ. iii. (1851) 263 = lanceaefolius.
serrulatus, Roxb. Hort. Beng. 42, nomen = tuberculatus.
sikkimensis, *Mast. in Hook. f. Fl. Brit. Ind.* i. 409.— Reg. Himal.
sikkimensis, Roxb. Encyc. Suppl. ii. 704 = sylvestris.
simplex, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 132.—Burma.
spathulatus, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* viii. (1861) 202.—N. Caled.
speciosus, *Brongn. & Gris, l. c.* 201.—N. Caled.
stipularis, *Blume, Bydr.* 121.—Burma; Malaya.
Storckii, *Seem. in Bonplandia*, x. (1862) 295.—Ins. Fiji.
strictus, Lamb. ex G. Don, Gen. Syst. i. 559 = dentatus.
subdenticulatus, *Miq. Fl. Ind. Bat. Suppl.* 408.— Sumatra.
submonocerus, *Miq. l. c.* 407.—Sumatra.
subpuberus, *Miq. l. c.* 406.—Sumatra.
suberratus, *Baker, in Journ. Linn. Soc.* xx. (1883) 108.—Madag.
subvillosus, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 328.—Zeylan.; Malaya.
sylvestris, Blanco, Fl. Filip. ed. II. 309 = oblongus.
sylvestris, *Poir. Encyc. Suppl.* ii. 704 (silvestris).— Cochinch.
tectorius, *Poir. l. c.*—Cochinch.
tinctorius, Steud. Nom. ed. II. i. 480, sphalm. = praec.
tomentosus, *Blume, Bydr.* 121.—Java.
transultus, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 804.—Soc. Socotra.
tuberculatus, *Roxb. Hort. Beng.* [98]; *Fl. Ind.* ii. 594. —Ind. or.; Malaya.
vaccinifolius, *F. Muell. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 301.—N. Caled.
Varunus, *Buch.-Ham. in Wall. Cat.* n. 2666 G, H.— Ind. or.; Malaya.
venustus, Bedd. Fl. Sylvat. t. 174 = E. Monocera.
verrucosus, Steud. Nom. ed. II. i. 49, sphalm. = Alsarites cordata.
Vieillardii, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* viii. (1861) 202.—N. Caled.
Wallichii, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 133.—Burma.

ELAEOCHARIS, Brongn. Enum. Gen. 12 (1843) = Bleocharis, R. Br. (Cyperac.).

ELAEOCHYTRIS, Fenzl, in Russegg. Reise, i. 957. t. 19 (1848) (Ill. Fl. Syr. 71) = Peucedanum, Tourn. (Umbell.).
cachroides, Fenzl, in Flora, xxvi. (1843) 462 = P. Schischentendalii.
meifolia, Fenzl, in Russeg. Reise, ii. 956. t. 19 = Ferula meifolia.

ELAEOCOCCA, Comm. ex A. Juss. Euph. Tent. 38. t. 11 (1824) = Alsurites, Forst. (Euphorb.).
cordata, A. Juss. l. c.; Blume, Bijdr. 618 = A. cordata
Vernicia, A. Juss. ex Spreng. Syst. iii. 884 = A. cordata
verrucosa, A. Juss. ex Spreng. l. c. = A. cordata.

ELAEOCOCCUS, Spreng. Syst. iii. 884 (1826) = praec.

ELAEODENDRON, Jacq. f. in Nov. Act. Helv. 86 (1787). *CELASTRINEAE*, Benth. & Hook. f. 867.
CROCOXYLON, Eckl. & Zeyh. Enum. 198 (1835).
LAMARCKIA, Hort. ex Endl. Gen. 1088, in syn. (1841).
LAURIDIA, Eckl. & Zeyh. Enum. 194 (1835).
LOUREIRA, Raesach. Nom. ed. III. 67 (1797).
MYSTROXYLON, Eckl. & Zeyh. Enum. 195 (1835).
NERRIJA, Roxb. Fl. Ind. i. 646 (1824).
PARILIA, Dennst. Schlüssel. Hort. Malab. 31 (1818)
PORTENSCHLAGIA, Tratt. Archiv, 250 (1812).
RUBENTIA, Comm. ex Juss. Gen. 378 (1789).
SCHREBERA, Retz. Obs. vi. 35. f. 8 (1791).
SYCOPHYLLUM, Eckl. & Zeyh. Enum. 194 (1835)
aethiopicum, *Oliver, Fl. Trop. Afr.* i. 365.—Afr. trop.
angustifolium, *Presl, Bot. Bemerk.* 84.—Afr. austr.
arboreum, *Pauch. & Sebert, Not. Bois Noue. Calid* 284.—N. Caled.
Argan, Retz. Obs. vi. 26 = Argania Sideroxylon.
arnense, *Montr. in Mém. Acad. Lyon*, x. (1860) 194 — N. Caled.
athranthum, *Presl, Bot. Bemerk.* 84.—Afr. austr.
attenuatum, *A. Rich. Ess. Fl. Cub.* 343.—Ind. occ
australe, *Vent. Yard. Malm.* t. 117.—Austral.
capense, *Eckl. & Zeyh. Enum.* 197.—Afr. austr.
camosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi (1863) i. 602.—N. Zel.
clusiophyllum, *Baill. Adansonia*, xi. (1873-76) 289 — Calif.
croceum, *DC. Prod.* ii. 11.—Afr. austr.
Cunninghami, *Montr. in Mém. Acad. Lyon*, x. (1860) 194.—N. Caled.
curtipendulum, *Endl. Prod. Fl. Ins. Norf.* 141.—Ins. Norfolk.
dioicum, *Griseb. Fl. Brit. W. Ind.* 709.—Ind. occ.
Dregeanum, Presl, Bot. Bemerk. 84 = Pterocelastrus Dregeanus.
ellipticum, *Decne. in Nouv. Ann. Mus. Par.* iii (1834) 478.—Ins. Timor.
Fortunei, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi (1863) i. 602.—China.
glaucum, *Pers. Syn.* i. 341.—As. trop.
griseum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 254.—Madag.
horizontale, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi (1863) i. 603.—Java.
illicifolium, *Tenore, in Atti Terza Ruin. Sc. Ital.* 304. —Hab.?
indicum, Gaertn. Fruct. i. 274. t. 57 = orientale.
indicum, *Poit. in Rev. Hortic. Sér.* II. v. (1846) 155 (= praec.?).—Hab.?
integrifolium, Sweet, Hort. Brit. ed. II. 110 = australe.
japonicum, *Franch. & Sav. Enum. Pl. Jap.* ii. 210.— Japon.
javanicum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 603.—Java.
laurifolium, *Juss. Thes. Cap.* ii. 54.—Afr. austr.
laurinum, Presl, Bot. Bemerk. 84 = Celastrus laurinus.
maculosum, Lindl. in Mitch. Journ. Trop. Austral. 344 = Flindersia maculosa.
melanocarpum, *F. Muell. Fragm.* iii. 92.—Austral.
Meyerianum, Presl, Bot. Bemerk. 84 = Celastrus heterophyllus, *Eckl. & Zeyh.*
microcarpum, Hook. f. in Hook. Lond. Journ. Bot. iii. (1844) 229 = Mellcytus microcarpus.
microcarpum, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 116.—Madag.
nitidissimum, Wall. Cat. n. 6490 = Microtropis ovalifolia.
nitidulum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 268.—Madag.

ELAEODENDRON :—
Novae-Zelandiae, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 602.—N. Zel.
obtusum, Fresl, Bot. Bemerk. 24 = Celastrus oleoides
obganthum, *Baker, in Journ. Linn. Soc.* xx. (1883) 121.—Madag.
orientale, *Spag. Ic. Rar.* t. 48.—Ins. Maurit.; Madag.
oppodon, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 603 = glaucum.
paniculatum, Wight & Arn. Prod. 157 = glaucum
papillosum, *Hochst. in Flora*, xvii. (1844) 305.—Afr. austr.
parvifolium, *Tul. in Ann. Sc. Nat. Sér.* IV. viii. (1857) 109.—Malaya.
pilosum, *Baker, in Journ. Linn. Soc.* xx. (1883) 122—Madag.
punctulatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi (1863) i. 602.—N. Zel.
pyramidale, Salisb. Prod. 143 = orientale.
quadrangulatum, *Reuss. in Mart. Fl. Bras.* xi. i. 33.—Bras.
rotundatum, *DC. Prod* ii. 11.—Ins. S. Thomas.
Roxburghii, Wight & Arn. Prod. 157 = glaucum.
schimoides, Spreng. Syst. i. 780 = Hartogia capensis
sphaerophyllum, *Presl, Bot. Bemerk.* 34; *Bull. Hist Pl.* vi. 27.—Afr. austr.
tomentosum, Willd. ex Roem & Schult. Syst. iii. 345 = Cervantesia tomentosa.
trachycladum, *Baker, in Journ Linn. Soc* xxi (1884) 333.—Madag.
tuberve, *Hort. Madr ex Roem. & Schult. Syst* v. 345.—Ind. or
undulatum, Zipp. ex Span. in Linnaea, xv. (1841) 186—ellipticum.
vaccinioides, *Baker, in Journ. Linn. Soc.* xxi. (1884) 338—Madag.
velutinum, *Harv. Thes. Cap.* ii 55.—Afr. austr.
verrucosum, Kunth, Ind. Sem. Hort Berol. (1845) 12; et in Linnaea, xix. (1847) 393 = croceum.
viride, *Buch.-Ham. ex Wall. Cat.* n. 6826.—Ind or.
xylocarpum, *DC. Prod.* ii. 11.—Ins. S. Thomas.
Zeyheri, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 452.—Afr. austr.

ELAEODENDRUM, Marr. Syst. ed. XIV. 241 (1784) = **Elaeodendron**, Jacq. (Celastr.).

ELAEOGENE, Miq. Fl. Ind. Bat. Suppl. 460 (1860) = *GENUS DUBIUM*. Benth & Hook. f. iii. 256.
sumatrana, *Miq. l. c.*—Sumatra

ELAEOSELINUM. Koch, ex DC. Prod. iv. 215 (1830). *UMBELLIFERAE*, Benth. & Hook f. i. 930.
MARGOTIA, Boiss. Elench 52 (1838).
Asclepium, *Bertol. Fl. Ital.* iii 883.—Europ. austr.
Asclepium, Willk. in Pardo, Ser. Inconf. Arag. 42 = Loscosii.
crinvolucratum, Cos. in Bull. Soc. Bot. Fr xx.(1873) 249; xxi. (1874) 60.—Morocco.
foetidum, *Boiss. Elench.* 91.—Europ. austr.; Afr. bor.
Fontanesii, Boiss. Voy. Bot. Espagne, ii. 363 = foetidum.
humile, *Ball, in Journ. Linn. Soc.* xvi. (1878) 479.—Morocco.
Lagascae, *Boiss. Elench.* 50.—Hispan.
laxum, *Pomel, Nouv. Mat. Fl. Atl.* 151.—Alger.
Loscosii, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* iii. 26.—Hispan.
meoides, *Koch, ex DC. Prod.* iv. 215.—Reg. Medit. occ
millefolium, *Boiss. Elench.* 50.—Hispan.
tenuifolium, Lange, in Willk. & Lange, Prod. Fl Hisp. iii. 26 = Lagascae.
thapsoides, *DC. Prod.* iv. 215.—Mexic.

ELAEOSTICTA, Fenzl, in Flora, xxvi. (1843) 458 = **Conopodium**, Koch (Umbellif.).
glaucescens, Boiss. Diagn. Ser. I. x. 51.—Persia.
millefolium, Fenzl, in Flora, xxvi. (1843) 458.—Syria, Mesopot.
nodosa, Boiss. Diagn. Ser. I. x. 51.—Persia.

ELAIS, Linn. Mant. i. 21 (1767) = **Illaeis**, Jacq. (Palm.).

ELAPHRIUM, Jacq. Enum. Pl. Carib. 3 (1760); Stirp. Am. t. 108, t. 71 (1763) = **Bursera**, Jacq. (1760).
Alatapium, **Schlecht. ex Schlecht. in Linnaea, xvii. (1843) 225 = B. Aloisyion.**

ELAPHRIUM :—
arienae, H. B. & K. Nov. Gen. et Sp. vii. 81 = *capense bicolor*, Schlecht. in Linnaea, xvii. (1843) 625 = B. bicolor.
bipinnatum, Schlecht. l. c. 631 = B. bipinnata.
capense, DC. Prod. i. 724.—Afr. austr.
copalliferum, Moç. & Sesse, ex DC. Prod. i. 724 = B. excelsa.
crenatum, Hensl. Biol. Centr. Am. Bot. i 179, sphalm. = B. cuneata.
cuneatum, Schlecht. in Linnaea, xvii (1843) 629 = B cuneata.
excelsum, H. B. & K. Nov. Gen. et Sp. vii 80 = B. excelsa.
fagaroides, H. B. & K. l. c. 29 = B fagaroides.
glabrifolium, H. B. & K. l. c. 28 = B. penicillata.
glabrum, Jacq Enum. Pl. Carib. 19 = B. glabra.
grandifolium, Schlecht. in Linnaea, xvii. (1843) 249 = B. grandifolia
graveolens, H. B. & K. Nov Gen. et Sp. vii. 81 = B. graveolens.
Hindsianum, Benth. Bot. Voy. Sulph. 11 = B. Hindsiana.
inaequale, DC. Prod i 724 = Clausena inaequalis.
integerrimum, Tul. in Ann. Sc. Nat. Sér. III. vi. (1846) 369 = B. gummifera.
Jacquinianum, H. B. & K. Nov Gen. et Sp. vii 23 = B. tomentosa
jorullense, H. B. & K. l. c. 28 = B. jorullensis
lancifolium, Schlecht. in Linnaea, xvii. (1843) 247 = B. lancifolia.
lanuginosum, H. B. & K. Nov Gen. et Sp. vii. 81 = B lanuginosa.
ovalifolium, Schlecht in Linnaea, xvii. (1843) 248 = B. ovalifolia
penicillatum, Moç. & Sesse, in DC. Prod. i. 724 = B. penicillata
pubescens, Schlecht. in Linnaea, xvi. (1842) 527.—Mexic.
rhoifolium, Benth. Bot. Voy Sulph 10 = B. Hindsiana.
simplicifolium, Schlecht. in Linnaea, xvi. (1842) 532 = B. Schlechtendalii
submoniliforme, March ex Engl. in DC. Monog Phan. iv. 55 = B. submoniliformis.
Tacamahaca, Tul. in Ann. Sc. Nat. Sér. III. vi. (1846) 368 = B. graveolens.
tomentosum, Jacq. Enum. Pl. Carib. 19 = B. tomentosa.
torulosum, Hensl. Biol. Centr. Am. Bot. i. 180.—Mexic.

ELASMATIUM, Dulac, Fl. Hautes-Pyr 121 (1867) = **Goodyera**, R. Br. (Orchid.).
repens, Dulac, l. c. = G. repens.

ELATE, Linn. Mus. Cliff. 12 (1786) = **Phoenix**, Linn. (Palm.).
humilis, Hort. Angl. ex Pasq. Cat. Ort. Bot. Nap. 41, nomen = seq ?
sylvestris, Linn. Sp Pl 1189 = P. sylvestris
versicolor, Salisb. Prod. 264 (= prace. ?).—Ins. Borb.

ELATERIOPSIS, Ernst, in Flora, lvi. (1873) 257 = Cyclanthera, Schrad. (Cucurb.).
caracassana, Ernst, l. c. = C. caracassana.
macropoda, Cogn. Diagn. Cucurb. fasc. ii. 84 = C. macropoda.

ELATERIOSPERMUM, Blume, Bijdr. 620 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 288.
Tapos, *Blume, l. c.* 621.—Java.
Tokbrai, *Blume, l. c.*—Malaya.

ELATERISPERMUM, Reichb. Nom. 194 (1841) = prace.

ELATERIUM, Jacq. Stirp. Am. 241 (1763). *CUCURBITACEAE*, Benth. & Hook. f. i. 835.
MOMORDICA, Neck. Elem. i. 240 (1790).
RYTIDOSTYLIS, Hook. & Arn. Bot. Beech. Voy. 424. t. 97 (1841).
amazonicum, *Mart. ex Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii (1878); (*Diagn. Cucurb.* fasc. ii. 55).—Bras.
bifurcum, A. Dietr. Syn. Pl. v. 372 = Cyclanthera ribifolia.
Bigelovii, S. Wats. in Proc. Am. Acad. xii. (1877) 252 = Echinocystis Bigelovii!

ELATERIUM :—
brachystachium, DC. Prod. iii. 310 = Cyclanthera brachystachya.
carthagenense, *Yacq. Enum. Pl. Carib.* 31; *Linn. Sp. Pl.* ed. II. 1675.—Am. austr.
ciliatum, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii (1878); (*Diagn. Cucurb.* fasc. ii. 54).—Guatemala.
cordatum, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 224.—Ins. Galap.
cordifolium, Moench, Meth. 503 = Echallium Elaterium.
Coulteri, A. Gray, Pl. Wright. ii. 61 = Echinocystis Coulteri.
filiforme, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii (1878); (*Diagn. Cucurb.* fasc. ii. 56).—Mexic.
gemellum, Moç. & Sesse, ex DC. Prod iii. 310 = Echinocystis gemella.
glabrum, *Cogn. in Mém. Cour. Acad. Belg.* Sér. II. xiii. (1880) 199.—N. Granat.
gracile, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii (1878); (*Diagn. Cucurb.* fasc. ii. 51).—Mexic.
hastatum, H. B. & K. Nov. Gen. et Sp. ii. 120 = Cyclanthera tamnoides.
longiflorum, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii (1878); (*Diagn. Cucurb.* fasc. ii. 54)—Panama
minimum, S. Wats. in Proc Am. Acad. xii (1877) 252 = Echinocystis minima
pubescens, Benth. Pl. Hartw. 15 = Echinocystis pubescens.
quadrifidum, *Moç. & Sesse, ex DC. Prod.* iii. 310.—Mexic.
quinquefidum, Hook. & Arn. Bot. Beech. Voy. 292 = Luffa operculata.
quinquelobum, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii (1878); (*Diagn. Cucurb.* fasc. ii. 53).—Venezuela.
ribiforum, Schlecht. in Linnaea, vii. (1832) 388 = Cyclanthera ribiflora.
tamnoides, Willd. Enum. Hort. Berol. 950 = Cyclanthera tamnoides.
torquatum, Moç. & Sesse, ex DC. Prod. iii. 310 = Echinocystis torquata
Trianaei, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii (1878); (*Diagn. Cucurb.* fasc. ii. 57).—N. Granat.
trifoliatum, Linn. Mant. i. 123; Spreng. Syst. iii. 47 = Sicyos angulatus
trilobatum, *Schlecht. in Linnaea*, xxvi. (1853–55) 630.—N. Granat.
Wrightii, A. Gray, Pl. Wright. ii. 61 = Echinocystis Wrightii.

ELATINE, Linn. Gen. ed. I. 118 (1737). *ELATINEAE*, Benth. & Hook. f. i. 162.
BIKOLIA, Bell. in Mém. Acad. Turin, Ser. I. xviii. (1808) 403. t. 11.
CRYPTA, Nutt. in Journ. Acad. Philad. i. (1817) 117. t. 6.
CRYPTINA, Raf in. Am. Monthly Mag. (1819) 192.
ELATINELLA, Opiz, Seznam, 89 (1852).
ELATINOPSIS, Clav. in Act. Linn. Soc. Bord. xxxvii. (1884) p. lxx.
HYDROPIPER, Buxb. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349.
POTAMOPITYS, Buxb. ex Adans. Fam. ii. 256 (1763).
RHIZINUM, Dulac, Fl. Hautes-Pyr 248 (1867).
WILLISELLUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 736 (1821).
alsinoides, Wight, Ic. t. 222 = Bergia sestivosa.
aiata, Roxb. & Schmidt, Fl. Sedin. 179 = Callitriche verna.
Alsinastrum, *Linn. Sp. Pl.* 868.—Europ.; Afr. bor.
ambigua, *Wight, in Hook. Bot. Misc.* ii. (1831) 103.—Ind. or.; Ins. Fiji.
americana, *Arn. in Edinb. Journ. Nat. Geogr. Sc.* i. (1830) 481.—Am. bor.; Austral.; N. Zel.
americana, A. Gray, in Hall, Pl. Tex. 5 = brachy sperma.
americana, A. Gray, in Proc. Acad. Sc. Philad. 1863 (1864) 58 = triandra.
ammannioides, Wight & Arn. Prod. 41 = Bergia ammannioides.
brachysperma, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 361, 368.—Am. bor.
Brochoni, *Clav. in Act. Linn. Soc. Bord.* xxxvii. (1883) p. lxiii.—Gallia.
californica, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 364.—Calif.

ELATINE:—
callitrichoides, Rupr. ex Petrow. in Bull. Soc. Nat. Mosc. xlviii. (1874) ii. 299 = Mandtia.
camphylosperma, *Seub. in Walp. Rep.* i. 284.—Reg. Mediterr.
chilensis, *C. Gay, Fl. Chil.* i. 286.—Chili.
Clintoniana, Peck, Rep. N. York. St. 32, 25 = americana.
conjugata, Lam. Fl. Fr. iii. 11 = major.
Fabri, Gren. in Acad. Sc. Besanç. Séances Publ. (1858) 124 = macropoda.
gratioloides, A. Cunn. in Ann. Nat. Hist. iv. (1840) 26 = americana.
Hardyana, Dum. in Bull. Soc. Bot. Belg. xi. (1872) 267 = E. Hydropiper.
hexandra, *DC. Fl. Fr.* v. 609; *Ser. Ic. Pl. Rar.* i. 14. t. 43. f. 1.—Europ.
hippuroides, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = E. Alsinastrum.
Hydropiper, *Linn. Sp. Pl* 367.—Europ.
Hydropiper, Schkuhr, Handb. t. 109 a = major.
Lindbergii, *Rehrb. in Mart. Fl. Bras.* xiv. ii. 321.—Bras.
lurvoriana, Delile, Fl. Egypte, 216 = Bergia verticillata.
macropoda, *Guss. Fl. Sic. Prod.* i. 475. — Reg. Mediterr.
major, *A. Br. in Syll. Ratisb.* i. (1824) 88.—Europ.
majuscula, Dum. Fl. Belg. 111 = major.
microphylla, *Griseb. in Lechl. Berb. Am. Austr.* 59.—Peruv.
minima, Fisch. & Mey. in Linnæa, x. (1836) 73 = americana.
nodosa, Arn. in Edinb. Journ. Nat. Geogr. Sc. i. (1830) 431 = E. Hydropiper.
orthosperma, Tineb. in Bot. Notiser, i. (1839) 88 = E. Hydropiper.
paludosa, Seub. in Walp. Rep. i. 284 = major, hexandra.
pedicellaris, F. Muell. Fragm. ii. 145 = Bergia pedicellaris.
perennis, F. Muell. l. c. 146 = Bergia perennis.
Schkuhriana, Heyne, in Reichb. Fl. Germ. Exeurs 639 = E. Hydropiper.
Serpillifolium, Gilib. Fl. Lituan. ii. 149 = E. Hydropiper.
serpyllifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 650 = prærc.
sisphosperma, Dum. Fl. Belg. 111 = E. Hydropiper.
spathulata, Gorski, in Eichw. Skizzr, 178 = E. Hydropiper.
texana, Torr. & Gray, Fl. N. Am. i. 678 = Bergia texana.
triandra, *Schkuhr, Handb.* i. 345. t. 119. b. f. 2.—Europ.
trispatala, Sm. Engl. Fl. ii. 248 = hexandra.
verticillaris, F. Muell. Fragm. ii. 148 = Bergia pusilla.
verticillata, Lam. Fl. Fr. iii. 11 = E. Alsinastrum.
verticillata, Wight & Arn. Prod. 41 = Bergia verticillata.

ELATINE, Rupp. Fl. Jen. ed. Hall. 242 (1742) = Linaria, Tourn. (Scrophul.).
cirrhosa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = L. cirrhosa.
Cymbalaria, Moench, Meth. 525 = L. Cymbalaria.
fruticosa, Moench, ex Steud. Nom. ed. II. i. 546 = L. fruticosa.
graeca, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 126 = L. graeca.
hastata, Moench, Meth. 524 = L. Elatine.
lanigera, Moench, ex Steud. Nom. ed. II. i. 546 = L. lanigera.
ovata, Moench, Meth. Suppl. 171 = L. spuria.
pilosa, Moench, ex Steud. Nom. ed. II. i. 546 = L. pilosa.
spuria, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 126 = L. spuria.
tenuis, Moench, Meth. Suppl. 171 = L. cirrhosa.

ELATINELLA, Opiz, Seznam, 39 (1852) = Blatine, Linn.
Hydropiper, Opiz, l. c. = Elatine Hydropiper.
Alsinastrum, Opiz, l. c. = Elatine Alsinastrum.

ELATINOPSIS, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) p. lxx = Blatine, Linn.
Brochoni, Clav. l. c. = Elatine Brochoni.

ELATOSEMA, Franch. & Sav. Enum. Pl. Jap. i. 488 (1875) sphalm. = seq.

ELATOSTEMA, Forst. Char. Gen. 105. t. 53 (1776). *URTICACEAE*, Benth. & Hook. f. iii. 366.
LANGEVELDIA, Gaudich. in Freyc. Voy. Bot. 494 (1826).
acuminatum, *Brongn. Bot. Voy. Coq.* 211.—Ind. or.; Malaya.
affine, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 188. = sessile.
approximatum, Wedd. in DC. Prod. xvi. i. 190 = cuneatum.
Brongniartianum, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 190 = sessile.
bulbiferum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 104.—Burma.
cornutum, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 816.—Reg. Himal.
cuneatum, *Wight, Ic.* vi. t. 2094. f. 3.—Ind. or.; Malaya.
cuspidatum, Wight, l. c. t. 1988 = sessile.
cuspidiforme, Miq. Pl. Jungh. 23 = sasquifolium.
cyrtandræfolium, Miq. l. c. 21 = sessile.
decipiens, *Wedd. in DC. Prod.* xvi. 176.—Reg. Himal.
delicatulum, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 190 = obtusum.
densiflorum, *Franch. & Sav. Enum. Pl. Jap.* i. 439. —Japon.
dasecoma, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 814.—Reg. Himal.
divaricatum, Gaudich. in Freyc. Voy. Bot. 494, in syn. = Pellionia divaricata.
diversifolium, Wedd. in DC. Prod. xvi. ii. 189 = sarculosum.
ellipticum, *Wedd. l. c.* 186.—Reg. Himal.
eurhynchum, *Miq. Fl. Ind. Bat.* i. ii. 244.—Java.
fagtifolium, *Gaudich. in Freyc. Voy. Bot.* 498.—Ins. Mascar.
ficoides, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 806. —Reg. Himal.
filicoides, *Seem. ex Wedd. in DC. Prod.* xvi. i. 188.—Ins. Fidji.
frutescens, Hask. Cat. Hort. Bog. Alt. 79 = Procris frutescens.
gibbosum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 104 (= Pilea sp.).—Ind. or.
glaucescens, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 825.—Ins. Philipp.
Goudotianum, *Wedd. l. c.* 819.—Madag.
gracile, *Hassk. Cat. Hort. Bog. Alt.* 79.—Java.
grande, Wedd. in DC. Prod. xvi. i. 173 = sessile.
hirsutum, *Miq. Fl. Ind. Bat.* i. ii. 243.—Java.
Hookerianum, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 809.—Reg. Himal.
Humbloti, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 883.—Madag.
incisum, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 821. —Madag.
integrifolium, Wedd. in DC. Prod. xvi. i. 179 = sesquifolium.
involucratum, *Franch. & Sav. Enum. Pl. Jap.* i. 439. —Japon.
Japonicum, *Wedd. in Ann. Sc. Nat. Sér.* IV. i. (1854) 189.—Japon.
Junghuhnianum, *Miq. Fl. Jungh.* 18.—Sumatra.
laetum, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 190 = sessile.
laevigatum, Hask. Cat. Bog. Alt. 79 = Procris laevigata.
lanciflorum, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 190.—Java.
lepidotulum, *Miq. Fl. Ind. Bat. Suppl.* 418.—Sumatra.
lineolatum, *Wight, Ic.* t. 1984.—Ind. or.
longifolium, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 189.—Ins. Philipp.
luchuense, Forst. ex Guill. l. c. Sér. II. vii. (1837) 184 = Procris Cephalida.
macrophyllum, *Brongn. Bot. Voy. Coq.* 207. t. 45.—Ins. Ambela.
maculatum, *Gaudich. in Freyc. Voy. Bot.* 498.—Java.
maingayacerinse, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 189.—Madag.
mamillosus, Wedd. l. c. = rostratum.
Mannii, *Wedd. in DC. Prod.* xvi. i. 178.—Afr. trop.

ELATOSTEMA:—
Mariannæ, C. B. Clarke, in Journ. Linn. Soc. xx. (1876) 124 = ficoides.
membranaceum, Hasak. Cat. Hort. Bog. Alt. 79 = acuminatum.
membranifolium, *Kurz, in Journ. As. Soc. Beng.* xlii (1873) ii. 104 = acuminatum.
Miquelianum, Wedd. in Ann. Sc. Nat. Sér. IV. i (1854) 188 = sesquifolium.
molle, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 296.—Reg. Himal.; Malaya.
montanum, Endl. Prod. Fl. Ins. Norf. Bog. 39 = Procris montana.
monticolum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 216.—Afr. trop.
nemorosum, *Seem. Fl. Vit.* 240.—Ins. Fiji.
nigrescens, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 101. —Java.
Novarse, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 149.—Ins. Nicobar.
obovatum, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 190.—Ins. Philipp.
obtusum, *Wedd. l. c.*—Reg. Himal.
oppositifolium, Ins. in Hook. Kew Journ. iii. (1851) 179 = Lecanthus Wightii.
ovatum, Wight, Ic. t. 1985 = Lecanthus Wightii.
Palvaeanum, *Wedd. in DC. Prod.* xvi. i. 176.—Afr. trop.
paludosum, Miq. Pl. Jungh. 19 = macrophyllum.
papillosum, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 837.—Reg. Himal.
parvifolium, Brongn. Bot. Voy. Coq. 210 = Urvilleanum.
parvum, *Blume, ex Miq. in Zoll. Syst. Verz. Ind Archip.* 103.—Java.
pedunculatum, Forst. Char. Gen. 106 = Procris Cephalida.
pedunculosum, Miq. Pl. Jungh. 21 = macrophyllum.
Pellionioides, Gaudich. in Freyc. Voy. Bot. 494, in syn. = Pellionia elatostemoides.
platyphyllum, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 801.—Reg. Himal.
podophyllum, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 189.—Ins. Philipp.
polycephalum, Wedd. l. c. = sessile.
procrilioides, *Wedd. in DC. Prod.* xvi. i. 180.—Reg. Himal.; Burma.
pubescens, Pers. Syn. ii. 557 = sessile.
punctatum, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 189 = sessile.
radicans, Wedd. in Arch. Mus. Par. ix. (1855–56) 832 = Procris radicans.
reticulatum, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 188.—Austral.
rigidum, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 820. —Ins. Philipp.
rostratum, Hask. Cat. Hort. Bog. Alt. 79 = Pellionia elatostemoides.
rostratum, *Miq. in Zoll Syst. Verz. Ind. Archip.* 102.—Java.
rugosum, *A. Cunn. in Ann. Nat. Hist.* i. (1838) 215. t. 9. f. 6–8.—N. Zel.
rupestre, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 804. —Reg. Himal.; Java.
serratum, Forst. ex Wedd. in DC. Prod. xvi. i. 172 = sessile.
sesquifolium, *Hassk. Cat. Hort. Bog. Alt.* 79.—Ind. or.; Malaya.
sessile, *Forst. Char. Gen.* 106.—As. et Afr. trop.
sikkimense, *C. B. Clarke, in Journ. Linn. Soc.* xv (1876) 134.—Reg. Himal.
sinuatum, *Hassk. Cat. Hort. Bog. Alt.* 79.—Java.
spectabile, Miq. Fl. Jungh. 19 = macrophyllum.
stipitatum, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 190.—Austral.
Stracheyanum, *Wedd. l. c.* 188.—Reg. Himal.
strigosum, *Hassk. Cat. Hort. Bog. Alt.* 79.—Java.
subinciscum, *Wedd. in Arch. Mus. Par.* ix. (1855–56) 814.—Reg. Himal.
succosum, Miq. Fl. Jungh. 23 = Procris Cephalida.
sarculosum, *Wight, Ic.* t. 2094. f. 4.—Ind. or.
thibaudiæfolium, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 188 = rostratum.
tomentosum, Wedd. l. c. 189 = sessile.
torrelense, Gandich. in Freyc. Voy. Bot. 498, in syn. = Procris Cephalida.
triserve, Hochst. in Flora, xxviii. (1845) 88 = Procris Cephalida.
ulmifolium, Miq. Fl. Jungh. 21 = sessile.

ELATOSTEMA :—
umbellatum, Blume, Mus. Bot. Lugd. Bat. ii. f. xix = Pellionia umbellata.
umbrosum, A. Gray, ex Wedd. in DC. Prod. xvi. i. 177 = nemorosum.
Urvilleanum, *Brongn. Bot. Voy. Coq.* 210. t. 46. f. A.—Ins. Moluce.
Webbianum, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 190 = cuneatum.
Wightianum, Wedd. l. c. 188 = Procris laevigata
Zollingerianum, Miq. in Zoll. Syst. Verz. Ind. Archip. 101, 105 = rupestre.
Zollingerianum, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 188 = sesquifolium.

ELATOSTEMMA, Endl. Prod. Fl. Ins. Norf. 39 (1833) = pracc.

ELATOSTOMA, Wight, Ic. vi. 35. t. 2091 (1853) = pracc.

ELATTOSTACHYS, Radlk. in Sitzb. Math-Phys. Acad. Muench. ix. (1879) 502, 600. *SAPINDACEAE.*
apetala, *Radlk. l. c.* 602.—N. Caled.
Bidwillii, Radlk. l. c. 601 = Cupania Bidwillii.
duplicato-serrata, *Radlk. l. c.*—Sumatra ?; Celebes ?
falcata, Radlk. l. c. 602 = Cupania falcata.
incisa, *Radlk. l. c.* 601.—N. Caled.
nervosa, Radlk. l. c. 600 = Cupania nervosa.
verrucosa, Radlk. l. c. 601 = Cupania verrucosa.
vitiensis, *Radlk. l. c.* 602.—Ins. Fiji.
xylocarpa, Radlk. l. c. 600 = Cupania xylocarpa.
Zippeliana, *Radlk. l. c.* 601.—N. Guinea.

ELAYUNA, Rafin. Sylva Tellur. 145 (1838) = **Bauhinia**, Linn. (Legumin.).
biloba, Rafin. l. c. = B. reticulata.

ELBUNIS, Rafin. Fl. Tellur. iii. 88 (1836) = **Leucas**, R. Br. (Labiat.).
alba, Rafin. l. c. 89.—Hab. ?

ELCAJA, Forsk. Fl. Aegypt. Arab. 127 (1775) = **Trichilia**, R. Br. (Meliac.).
emetica, Forsk. l. c. = T. emetica.

ELCANA, Blanco, Fl. Filip. ed. II. 584 (1845) = **Cerbera**, Linn. (Apocyn.).
seminalis, Blanco, l. c. = C. Odollam.

ELEAGNUS, Hill, Hort. Kew. 440 (1769) = **Elaeagnus**, Tourn.

ELECTRA, DC. Prod. v. 630 (1836) = **Coreopsis** Linn. (Compos.).
Galeotti, A. Gray, Pl. Wright. i. 110 = C. Galeottii.
mexicana, DC. Prod. v. 630 = C. mexicana.

ELECTRA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 = **Sapindus**, Linn.

ELECTRA, Pans. in Denkschr. Acad. Muench. (1813) 209 = **Schismus**, Beauv. (Gramin.).
brevifolia, Nees, ex Steud. Nom. ed. II. i. 516 = S. brevifolius.
calycina, Pritz. Ind. Ic. 294 = S. brevifolius.
nitida, Pans. ex Steud. Nom. ed. II. i. 516 = S. marginatus.
ovalis, Nees, ex Steud. l. c. = S. ovalis.
scaberrima, Nees, ex Steud. l. c. = S. scaberrimus.

ELECTROSPERMA, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 28 = **Brioscaulon**, Linn.
australasicum, F. Muell. l. c. = Erioc. australasicum.

ELEGIA, Linn. Mant. ii. 163 (1767). *RESTIACEAE*, Benth. & Hook. f. iii. 1092.
CHONDROPETALUM, Rottb. Progr. 12 (1772); Desc. et Ic. Pl. 10. t. 3 (1778).
LAMPROCAULOS, Mast. in DC. Monog. Phan. i. 349 t. 8 (1878).
scaulosta, *Mast. in Journ. Linn. Soc. xxi.* (1885) 580.—Afr. austr.
capensiformis, *Kunth, Enum. Pl.* iii. 474.—Afr. austr.
exilibris, *Labill. ex Steud. Nom. ed.* II. i. 594.—Austral.
bracteolata, Reichb. ex Kunth, Enum. Pl. iii. 466 = Restio erectus.

ELEGIA :—
coleura, *Nees. ex Mast. in A. DC. Monog. Phan.* i. 358.—Afr. austr.
cuspidata, *Mast. in Journ. Linn. Soc. x.* (1869) 240.—Afr. austr.
deusta, *Kunth, Enum. Pl.* iii. 460.—Afr. austr.
Dregeana, Kunth, l. c. 469 = asperifolia.
equisetacea, *Mast. in Journ. Linn. Soc.* xxi. (1885) 588.—Afr. austr.
fastigiata, *Mast. l. c.* 589.—Afr. austr.
fistulosa, *Kunth, Enum. Pl.* iii. 467.—Afr. austr.
glauca, *Mast. in Journ. Linn. Soc.* xxi. (1885) 579.—Afr. austr.
gracilis, Kunth, Enum. Pl. iii. 475 = Lamprecaulos grandis.
juncea, *Linn. Mant.* ii. 297.—Afr. austr.
juncea, Zeyh. ex Kunth, Enum. Pl. iii. 466 = parviflora.
Krausii, Hochst. in Flora, xxviii. (1845) 340 = propinqua.
membranacea, Kunth, Enum. Pl. iii. 474 = juncea.
mucronata, Reichb. ex Kunth, l. c. 475 = Dovea mucronata.
Nessii, Mast. in Journ. Linn. Soc. x. (1869) 246 = Lamprocaulos Nessii.
nuda, *Endl. Gen.* 122.—Afr. austr.
paniculoides, Kunth, Enum. Pl. iii. 470 = Dovea muconata.
paniculata, Pers. Syn. ii 607 = Cannamois virgata.
parviflora, *Kunth, Enum. Pl.* iii. 467.—Afr. austr.
propinqua, *Kunth, l. c.* 473.—Afr. austr.
racemosa, Pers. Syn. ii. 607 = Dovea tectorum.
racemosa, Reichb. ex Kunth, Enum. Pl. iii. 461 = deusta.
rigida, *Mast. in Journ. Linn. Soc* xxi. 1885) 587.—Afr austr.
spathacea, *Mast. l. c.* 588.—Afr. austr.
squamea, *Mast. l. c.* x. (1869) 244.—Afr. austr.
stipularis, *Mast. l. c.* xxi. (1885) 587.—Afr. austr.
thyrsifera, Pers Syn. ii. 607 = propinqua.
vaginulata, *Mast. in Journ. Linn. Soc.* xxi. (1885) 587.—Afr. austr.
Verreauxii, *Kunth, Enum. Pl.* iii. 471.—Afr. austr.
verticillaris, *Kunth, Enum. Pl.* iii. 471.—Afr. austr.

ELEIASTIS, Rafin. Sylva Tellur. 165 (1838) = **Tetracera**, Linn. (Dillen.).
levis, Rafin. l. c. = T. laevis.

ELEIOSINA, Rafin. Sylva Tellur. 150 (1838) = **Spiraea**, Linn. (Rosac.).
altaica, Rafin. l. c. 151 = S. laevigata.
bracteata, Rafin. l. c. = S. laevigata.
cuneifolia, Rafin. l. c. = S. laevigata.
levigata, Rafin, l. c. = S. laevigata.
obovata, Rafin. l. c. = S. salicifolia.
trilobo, Rafin. l. c. = S. trilobata.
virgata, Rafin. l. c. = S. virgata.

ELEIOTIS, DC. Mém. Lég. vii. 348 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 538.
monophylla, DC. l. c. 350 = E. Sororia.
Rottleri, Wight & Arn. Prod. 281 = Desmodium Rottleri.
Sororia, *DC. Mém. Lég.* vii. 350.—Ind. or.

ELELIS, Rafin. Fl. Tellur. iii. 92 (1836) = **Salvia**, Tourn. (Labiat.).
austriaca, Rafin. l. c. 93 = S. austriaca.

ELEMANTHUS, [Schlecht. in] Bot. Zeit. v. (1847) 150 = EILEMANTHUS, Hochst. = **Indigofera**, Linn. (Legumin.).

ELEMI, Adans. Fam. ii. 342 (1763) = **Amyris**, Linn. (Rutac.).

ELEMIFERA, Linn. Hort. Cliff. 486 (1737) = pracc.

ELEOCARIS, Sanguin. in Atti Acc. Pont. Lincei, Ser. I. v. (1852) 498, 516 = seq.

ELEOCHARIS, R. Br. Prod. 224 (1810). *CYPERACEAE*, Benth. & Hook. f. iii. 1047.
BULBOSTYLIS, Stev. in Mém. Soc. Nat. Mosc. v. (1813) 355.
CHAETOCYPERUS, Nees, in Linnaea, ix. (1834) 289.

ELEOCHARIS :—
CLAVULA, Dum. Fl. Belg. 143 (1827), pro parte.
ELEOGENUS, Nees, in Linnaea, ix. (1834) 294.
LIMNOCHARIS, Kunth, Enum. Pl. ii. 188 (1837).
LIMNOCHLOA, Beauv. ex Lestib. Essai Cyp. 41 (1819).
SKVIPIDIUM, Nees, in Linnaea, ix. (1834) 293.
acicularis, Boj. Hort. Maurit. 284 = E. Chaetaria.
acicularis, *R. Br. Prod.* 224.—Reg. bor.; Am. austr.
acuminata, *Nees, in Linnaea*, ix. (1834) 294.—Am. bor.
acuta, *R. Br. Prod.* 224.—Austral.
acutangula, Schult. Mant. ii. 93 = fistulosa.
acutisquamata, Boeckl. in Proc. Acad. Sc. Philad. 1862 (1863) 10 = palustris.
affinis, C. A. Mey. ex Claus, in Beitr. Pfl. Russ. Reich viii. 261 = palustris.
affinis, *Steud. Syn. Pl. Cyp.* 76.—China; Java.
albibracteata, *Nees & Meyen, in Linnaea*, ix. (1834) 294, nomen; *Meyen, Reise*, i. 454, nomen.—Peruv.
albida, *Torr. in Ann. Lyc. N. York*, iii. (1836) 304.—Am. bor.; Ind. occ.
albivaginata, Boeck. in Linnaea, xxxvi. (1869–70) 436 = ochreata.
ambigua, T. Kirk & Buchanan, in Trans. N. Z. Inst. vi. (1873) 225 = acuta.
amphibia, *Durv. in Act. Soc. Linn. Bord.* xxi. (1856) 287. t. 2.—Gallia.
anceps, *Ridl. in Trans. Linn. Soc. Ser.* II. ii. (1884) 148.—Afr. trop.
apposiciculata, Phil. in Linnaea, xxix. (1857–58) 85 = palustris.
arcuata, Kunze, ex Kunth, Enum. Pl. ii. 147 = maculosa.
arenaria, *Benth. in Hook. Kew Journ.* ii. (1850) 244, in textu.—Bras.
arenicola, Torr. ex Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 237 = montana.
argyrolepis, Kier. in Mém. Sav. Etr. Péterb. vii. (1851) 518 = palustris.
articulata, Kunth, Enum. Pl. ii. 157 = interstincta.
ascasmensis, *Phil. Fl. Atac.* 58.—Chili.
atricha, *Benth. Fl. Austral.* vii. 295 (sp. diverse).—Austral.
atricha, *R. Br. Prod.* 225.—Austral.
atropurpurea, *Kunth, Enum. Pl.* ii. 151.—Reg. temp. et trop.
atropurpurea, J. & C. Presl, Rel. Haenk. i. 196 = capitata.
Baeothryon, Nees, in Linnaea, ix. (1834) 294 = Scirpus pauciflorus.
Baeothryon, *Schult. Mant.* ii. 92.—Am. bor.
Balansaiana, *Boeck. in Flora*, lxii. (1879) 159.—Am. trop.
Baldwinii, *Chapm. Fl. S. U. St.* 519.—Am. bor.
Baroni, Baker, in Journ. Linn Soc. xx. (1883) 297 = limosa.
Bartolinus, De Not. Ind. Sem. Hort. Genuens. (1846) 24; et in Ann. Sc. Nat. Sér. III. xi. (1849) 255 = carniolica.
Bartolominea, Stend. Syn. Pl. Cyp. 80, sphalm. = praec. bicolor, *Chapm. Fl. S. U. St.* 517.—Florida.
biocreata, Nees, in Linnaea, ix. (1834) 294 = ochreata.
bisetiata, Steud. Syn. Pl. Cyp. 82 = sphacelata.
bivaginata, *Steud. l. c.* 77.—Paraguay.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 392.—Calif.
bonariensis, *Nees, in Hook. Journ. Bot.* ii. (1840) 298. —Am. austr.
brizantha, *Steud. Syn. Pl. Cyp.* 79.—Guatemala.
caduca, *Schult. Mant.* ii. 88.—Aegypt.; Syria; Sardinia.
caespitosa, Link, Hort. Berol. i. 288 = Scirpus caespitosus.
caespitosissima, Baker, in Journ. Linn. Soc. xxi. (1885) 450 = subvivipara.
calva, Torr. Fl. N. York, ii. 346, in obs. = palustris.
calyptrata, Steud. Syn. Pl. Cyp. 81 = sulcata.
cancellata, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 170.—Mexic.
capillacea, *Kunth, Enum. Pl.* ii. 139.—Bras.
capitata, *R. Br. Prod.* 223.—Reg. trop.
carniolica, *Koch, Syn. Fl. Germ.* ed. II. 853.—Europ. austr.
cellulosa, *Torr. in Ann. Lyc. N. York*, iii. (1836) 298. —Am. bor.
Chaetaria, *Roem. & Schult. Syst.* ii. 154.—Reg. trop.
chlorocarpa, Boeck. in Flora, lxi. (1878) 34 = atricha.

ELEOCHARIS :—
chrysocarpa, *Boeck. in Kjoeb. Vidensnh. Meddel.* (1869) 134.—Bras.
cognata, Steud. Syn. Pl. Cyp. 81 = interstincta.
compacta, *R. Br. Prod.* 224.—Austral.
complanata, *Boeck. in Flora*, lxii. (1879) 562.—Afr. trop.
compressa, Sullivant, in Am. Journ. Sc. Ser. I. xlii. (1842) 50 = acuminata.
confervoides, *Steud. Syn. Pl. Cyp.* 82.—Madag.
congesta, *D. Don, Prod. Fl. Nep.* 41.—Nepal.
conica, J. & C. Presl, Delic. Prag. 149 = palustris.
consanguinea, Kunth, Enum. Pl. ii. 148 = nodulosa.
constricta, Schult. Mant. ii. 87 = geniculata.
coronata, *Steud. Syn. Pl. Cyp.* 77.—Chili.
costata, J. & C. Presl, Fl. Cech. 11 = acicularis.
costulata, *Nees & Meyen, in Linnaea*, ix. (1834) 294, nomen.—Chili
crassa, C. A. Mey. ex Boeck. in Flora, xli. (1858) 413, in obs.=palustris
crassiculis, Steud. Syn. Pl Cyp. 81 = geniculata.
Cunninghamii, *Boeck in Flora*, xli. (1858) 412.—N. Zel.
cylindrica, *Buchl. in Proc. Acad. Sc. Philad.* 1862 (1862) 10.—Texas.
cylindrostachys, *Boeck. in Flora*, lviii. (1875) 108.—Austral.
cyperina, Roem. & Schult. Syst. ii. 155 = Mariscus Jacquinii.
debilis, *Kunth, Enum. Pl.* ii. 143.—Bras.
dense, Benth. Pl. Hartw. 27 = geniculata.
densesquamata, Steud. Syn. Pl. Cyp. 83 = geniculata.
depauperata, Kunth, Enum. Pl. ii. 140 = E. Chastaria.
diandra, *Wright, in Bull. Torrey Bot. Club*, x. (1883) 101.—Am. bor.
Dietrichiana, *Boeck. in Flora*, lviii. (1875) 107.—Austral.
dispar, *E. Hill, in Coult Bot. Gaz. vi.* (1882) 3.—Am. bor.
Doenbeyana, *Kunth, Enum. Pl.* ii. 145.—Peruv.; Mexic.
Dregeana, Steud. Syn. Pl. Cyp. 78 = palustris.
dubia, Phil. in Linnaea, xxix. (1857–58) 77 = pachycarpa.
elata, *Boeck. in Kjoeb. Vidensk. Meddel.* (1871) 151 — Bras
elegans, Roem. & Schult. Syst. ii. 150 = geniculata.
Ellisotti, A. Dietr. Sp. Pl. ii. 88 = interstincta.
elliptica, Kunth, Enum. Pl. ii. 146 = tenuis, montana.
elongata, *Chapm. Fl. S. U. St.* 515.—Florida.
emarginata, Klotzsch, ex Boeck. in Linnaea, xxxvi. (1869–70) 443.—Bras.
Engelmanni, Steud. Syn. Pl. Cyp. 79 = ovata.
equisetina, *Y. & C. Presl, Rel. Haenk.* i. 195.—Ins. Luzon.
equisetoides, Torr. in Ann. Lyc. N. York, iii. (1836) 296 = interstincta.
erratica, Steud. Syn. Pl. Cyp. 79 = atropurpurea.
erythrochlamys, Miq. Fl. Ind. Bat. iii. 500 = tetraquetra.
erythropoda, Steud. Syn. Pl. Cyp. 76 = glaucescens.
exigua, *Roem. & Schult. Syst.* ii. 154.—N. Granat.
filiculmis, Kunth, Enum. Pl. ii. 144 = sulcata.
filiculmis, Schur, Enum. Pl. Transs. 690 = palustris.
filiformis, Kunth, Enum. Pl. ii. 146 = tenuis.
fistulosa, *Schult. Mant.* ii. 89.—Geront. trop.
fistulosa, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 269 = variegata.
fluitans, Hook. Brit. Fl. ed. I. 24 = Scirpus fluitans.
fusco-sanguines, *Boeck. in Linnaea*, xxxvi. (1869–70) 495.—Chili.
geniculata, *R. Br. Prod.* 224.—Am. bor. et austr.
glauca, *Boeck. in Kjoeb. Vidensk. Meddel.* (1871) 150.—Bras.
glaucescens, *Schult. Mant.* ii. 89.—Am. bor.
Glazioviana, *Boeck. in Kjoeb. Vidensk. Meddel.* (1871) 150.—Bras.
gracilis, R. Br. Prod. 224.—Austral.
gracilis, Hook. f. Fl. Tasm. ii. 35 = acuta, Cunninghamii.
gracilis, Nees, in Linnaea, ix. (1834) 294 = maculosa.
gracillima, *Boeck. in Linnaea*, xxxvi. (1869–70) 459 = maculosa.
gracillima, Hook. f. Handb. N. Zeal. Fl. 745 = Cunninghamii.
Graefeana, Boeck. in Flora, lviii. (1875) 108 = variegata.
grandis, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 134 (= Fimbristylidis sp.).—Bras.
harmatolepis, Steud. Syn. Pl. Cyp. 79 = palustris.

ELEOCHARIS :—
Hildebrandtii, *Boeck. in Flora*, lxi. (1878) 34.—Afr. trop.
homotyma, *Steud. Syn. Pl. Cyp.* 79.—Guiana.
Hookeri, *Boeck. in Linnaea*, xxxvi. (1869–70) 430.—N. Zel.
incurvata, Schult. Mant. ii. 92 = Scirpus articulatus.
intermedia, *Schult. l. c.* 91.—Am. bor.
interstincta, *R. Br. Prod.* 224.—Am. trop.
japonica, *Boeck. in Linnaea*, xxxvi. (1869–70) 492.—Japon.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 142 = affinis.
Jelskiana, *Boeck. in Linnaea*, xxxviii. (1874) 376.—Guiana.
juncoides, Schult. Mant. ii. 90 = Scirpus debilis.
lastvirens, Steud. Syn. Pl. Cyp. 79 = atropurpurea.
Lechleri, *Boeck. in Linnaea*, xxxvi. (1869–70) 432.—Chili.
Lehmanni, Kier. in Mém. Sav. Etr. Pétersb. vii. (1851) 517 = palustris.
leptocaulis, Steud. Syn. Pl. Cyp. 77 = nana.
leptophylla, Schult. Mant. ii. 88 = Scirpus pungens.
Leveschii, Shuttl. in Flora, xx. (1837) 241 = atropurpurea.
leucocarpa, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 132.—Bras
Liebmanniana, *Boeck. in Linnaea*, xxxvi. (1869–70) 439.—Mexic.
limosa, Drège, in Linnaea, xx. (1847) 246 = palustris.
limosa, A. Rich. Tent. Fl. Abyss. ii. 496 = marginulata.
limosa, *Schult. Mant.* ii. 87.—Afr. austr; Madag.
llocarpa, *Phil. in Anal. Univ. Chil.* (1873) 553.—Chili.
litoralis, *Phil. in Linnaea*, xxix. (1857–58) 77.—Chili.
macra, Kunth, Enum. Pl. ii. 142 = debilis.
macrantha, *Boeck. in Linnaea*, xxxvi. (1869–70) 453 — Syria.
macrorhiza, *Boeck. in Flora*, xli. (1858) 413.—Ins. Faïal.
macrostylis, Spreng. Neue Entdeck. iii. 11 = Ryuchospora pterocarpa.
maculosa, *R. Br. Prod.* 224.—Am. austr.; Austral.
marginulata, *Hochst. ex Steud. Syn. Pl. Cyp.* 78.—Abyss.
media, Schult. Mant. ii. 91 = fistulosa.
melanocarpa, Fahl. in Linnaea, xxix. (1857–58) 85 = ochresta.
melanocarpa, *Torr. in Ann. Lyc. N. York*, iii. (1836) 311.—Am. bor.
melanocephala, *E. Desv. in C. Gay, Fl. Chil.* vi. 175.—Chili.
melanosperma, *Steud. Syn. Pl. Cyp.* 78.—Bras.
mendocina, *Phil. in Anal. Univ. Chil.* (1873) 553.—Chili.
mexicana, Peyr. in Linnaea, xxx. (1859–60) 314 = palustris?
microcarpa, *Torr. in Ann. Lyc. N. York*, iii. (1836) 312.—Am. bor. et centr.
microformis, Buchl. in Proc. Acad. Sc. Philad. 1862 (1863) 10 = interstincta.
minima, *Kunth, Enum. Pl.* ii. 139.—Bras.
minor, [Pleri?] in Ionica, Anthol. ii. (1834) 460 = palustris?
mitracarpa, Steud. Syn. Pl. Cyp. 77 = palustris.
minuta, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 508.—Madag.
monandra, Hochst. in Flora, xxv. (1842) 1. Beibl. 134, nomen; Steud. Syn. Pl. Cyp. 78 = atropurpurea.
montana, *Roem. & Schult. Syst.* ii. 158.—N. Granat.
montevidensis, *Kunth, Enum. Pl.* ii. 144.—Reg. Argent.
mucronulata, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 46 = acuta.
Muehlenbergiana, *Steud. Nom.* ed. II. i. 847.—Am. bor.
multicaulis, *Sm. Engl. Fl.* i. 64.—Europ.
multiflora, Chapm. Fl. S. U. St. 517 = atropurpurea.
mutata, *R. Br. Prod.* 224.—Am. austr.
mutata, Schlecht. in Bot. Zeit. vii. (1849) 134 = nodulosa.
nana, *Kunth, Enum. Pl.* ii. 140.—Bras.; Mexic.
Nassauniana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 92.—Afr. trop.
nebrodensis, Parl. Fl. Ital. ii. 62 = palustris.
nigrescens, *Kunth, Enum. Pl.* ii. 137.—Hab.?

ELEOCHARIS :—
nodulosa, *Schult. Mant.* ii. 87.—Bras.
obsoleta, Steud. Syn. Pl. Cyp. 81 = interstincta.
obtusa, Schult. Mant. ii. 89 = ovata.
obtusangula, *Kunth, Enum. Pl.* ii. 137.—Hab.?
obtusetrigona, Steud. Syn. Pl. Cyp. 80 = minuta.
ochracea, *Nees, in Linnaea*, ix. (1834) 294, nomen.—Am. trop.
ochrostachys, Boeck. in Linnaea, xxxvi. (1869–70) 452 = affinis.
ochrostachys, *Steud. Syn. Pl. Cyp.* 80.—Java.
olivacea, *Torr. in Ann. Lyc. N. York*, iii. (1836) 300.—Am. bor.
ovata, *R. Br. Prod.* 224.—Reg. temp. et trop.
oxycoryna, Dur. in Bull. Soc. Bot. Fr. ii. (1855) 609 = amphibia.
pachycarpa, *E. Desv. in C. Gay, Fl. Chil.* vi. 174.—Chili.
pallida, *Shuttl. ex Steud. Syn. Pl. Cyp.* 76.—Am. bor.
palustris, *R. Br. Prod.* 224.—Europ.; Am. bor.
palustris, F. Muell. Fragm. viii. 240 = acuta.
pauciflora, Link, Hort. Berol. i. 284 = Scirpus pauciflorus.
pauciflora, *S. Wats. in Bot. Calif.* ii. 221.—Am. bor. occ.
pellucida, *Y. & C. Presl, Rel. Haenk.* i. 196.—Ins. Luzon.
pianikiensis, Steud. Syn Pl. Cyp 77 = sulcata.
pilcata, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 417 = palustris.
planiculmis, Steud. Syn. Pl. Cyp. 80 = fistulosa.
planifolia, Nees, in Linnaea, ix. (1834) 294 = Scirpus planifolius.
plantaginea, *R. Br. Prod.* 224.—Ind. or.; Madag.
plantaginea, F. Muell. Fragm. viii. 238 = sphacelata
polycaula, Wender. in Marb. Schrift. i. (1822) 182 = palustris.
polymorpha, Nees, in Linnaea, ix. (1834) 294 = ochresta, etc.
prolifera, *Torr. in Ann. Lyc. N. York*, iii. (1836) 316.—Am. bor.
punctata, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 132.—Bras.
punctata, Hochst. ex Steud. Syn Pl. Cyp. 75 = sulcata.
purpurascens, Boeck. in Linnaea, xxxvi. (1869–70) 455 = congesta.
pusilla, *R. Br. Prod.* 225.—Austral.
pygmæa, Torr. in Ann. Lyc N. York, iii. (1836) 313 = Scirpus nanus.
quadrangulata, *R. Br. Prod.* 224.—Am. bor.
Rabenii, *Boeck. in Kjoeb. Vidensk. Meddel.* (1871) 149.—Bras.
radicans, Kunth, Enum. Pl. ii. 142 = costulata.
reclinata, Kunth, l. c. 143 = intermedia.
recurvata, Nees, in Linnaea, ix. (1834) 294 = E. Chastaria.
riparia, Nees, ex Spreng. Syst. iv. Cur. Post. 27 capitata.
riparia, Phil. in Linnaea, xxxiii. (1864–65) 270 = acicularis.
Robbinsii, *Oakes, in Am. Gard. Magaz.* vii. 178; *et A. Gray, Man.* ed. I. 524.—Am. bor.
rostellata, *Torr. Fl. N. York*, ii. 347.—Am. bor.
Rothiana, *Boeck. in Flora*, xliii. (1860) 3.—Bras.
Salzmanniana, *Steud. Syn. Pl. Cyp.* 78.—Bras.
scariosa, Steud. l. c. 80 = mutata.
Schottiana, Steud. Syn. Pl. Cyp. 79 = maculosa.
Schweinfurthiana, Boeck. in Flora, lxii. (1879) 562 = microcarpa.
Sellowiana, *Kunth, Enum. Pl.* ii. 149.—Bras.
septata, Miq. in Linnaea, xvii. (1843) 58 = interstincta
setacea, R. Br. Prod. 224, in obs.=E. Chastaria.
setacea, R. Br. l. c. 225 = capitata.
Sieberi, Kunth, Enum. Pl. ii. 138 = variegata.
simplex, A. Dietr. Sp. Pl. ii. 78 = tortilis.
simplex, Kunth, Enum. Pl. ii. 143 = albida.
singularis, Steud. Syn. Pl. Cyp. 81 = geniculata.
Soveria, Kunth, Enum. Pl. ii. 148 = lineata.
sphacelata, *R. Br. Prod.* 224.—Austral.
spiralis, *R. Br. l. c.* 226.—Ind. or.
stenolepis, *Boeck. in Linnaea*, xxxvi. (1869–70) 424.—Mexic.
striata, Hochst. ex Steud. Syn. Pl. Cyp. 76 = marginulata.
subsetacea, *E. Desv. in C. Gay, Fl. Chil.* vi. 173.—Chili.
stylosa, Steud. Syn. Pl. Cyp. 76 = sulcata.

ELEOCHARIS:—
subarticulata, *Boeck. in Linnaea*, xxxvi. (1869–70) 455. —Bras.
subnodosa, *Miq. in Linnaea*, xix. (1847) 925.—Guiana.
subnodulosa, *Steud. Syn. Pl. Cyp.* 81.—Guatemala.
subpolifera, Steud. l. c. 80 = affinis.
subsphacelata, Steud. l. c. 87 = sphacelata.
subtilis, *Boeck. in Linnaea*, xxxvi. (1869–70) 426.—Bras.
subulata, *Boeck. in Flora*, xli. (1858) 412.—Ind. or.
subvivipara, *Boeck. in Linnaea*, xxxvi. (1869–70) 424. —Ind. or.
sulcata, *Nees, in Linnaea*, ix. (1834) 294.—N. Mexic.
sulcicaulis, Reichb. ex Kunth, Enum. Pl. ii. 150 = ochreata.
tenuis, *Schult. Mant.* iii. 89.—Am. bor. et austr.
tenuissima, *Boeck. in Linnaea*, xxxvi. (1869–70) 419. —Bras.
tetraquetra, *Nees, in Wight, Contrib.* 113.—Nepal.
texana, *Britton, in Bull. Torrey Bot. Club*, xi. (1884) 87.—Texas.
Thomsoni, Boeck. in Linnæa, xxxvi. (1869–70) 451 = afflata.
Torreyana, Boeck. l. c. 440 = microcarpa.
tortilis, *Schult. Mant.* ii. 92.—Am. bor.
triangulata, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 368 = Scirpus mucronatus.
trichoides, Kunth, Enum. Pl. ii. 141 = acicularis.
tricostata, *Torr. in Ann Lyc. N. York*, iii. (1836) 310. —Am. bor.
trifora, Boeck. in Flora, lxiii. (1880) 437 = E. Chaetaria.
truncata, Schlecht. in Bot. Zeit. vii. (1849) 118 = Dombeyana.
tuberculosa, *R. Br. Prod.* 224.—Am. bor.
tuberosa, *Schult. Mant.* ii. 86.—Ind. or.
tumida, Schult. l. c. = plantaginea.
uniglumis, Schult. l. c. 88 = palustris.
valdiviana, *Phil. in Linnaea*, xxix. (1857–58) 77.—Chili.
valida, Boeck. in Flora, lxv. (1882) 350 = geniculata.
variegata, *Presl, in Oken, Isis*, xxi. (1828) 369; *Kunth, Enum. Pl.* ii. 158.—Madag.
vivipara, *Link, Hort. Berol.* i. 283.—Am. bor.
Watsoni, Bab. in Ann. & Mag. Nat. Hist. Ser. II. x. (1852) 20 = palustris.
Wickurai, Boeck. in Linnæa, xxxvi. (1869–70) 448 = tetraquetra.
Widgrenii, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 503. —Bras.
Zanardinii, Parl. Fl. Ital. ii. 67 = atropurpurea.

ELOGENUS, Nees, in Linnæa, ix. (1834) 294 = **Mesoeharis**, R. Br. (Cyperac.).
atropurpurea, Nees, in Wight, Contrib. 113 = Eleocharis atropurpurea.
capitatus, Nees, in Wight, l. c. 112 = Eleocharis capitata.
laxiovirens, Nees, in Mart. Fl. Bras. ii. i. 108 = Eleocharis atropurpurea.
nodulosus, Nees, l. c. 104 = Eleocharis nodulosa.
ochreatus, Nees, in Linnæa, ix. (1834) 294 = Eleocharis ochreata.
ovatus, Nees, l. c. = Eleocharis ovata.
Schottianus, Nees, in Mart. Fl. Bras. ii. i. 102 = Eleocharis maculosa.
Sellovianus, Nees, l. c. 108 = Eleocharis Sellowiana.

ELEOGITON, Link, Hort. Berol. i. (1827) 238 = Scirpus, Linn. (Cyperac.).
atropurpureus, A. Dietr. Sp. Pl. ii. 97 = Eleocharis atropurpurea.
caricoides, A. Dietr. l. c. 99 = Hemichlaena angustifolia.
cernuus, A. Dietr. l. c. 96 = S. cernuus.
curvulus, Nees, in Wight, Contrib. 110 = S. fluitans.
digitatus, Nees, in Linnæa, ix. (1834) 291; x. (1836) 168 = S. flaccifolius.
disciulus, Nees, l. c. ix. (1834) 291 = S. flaccifolius.
Donatii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 178 = S. lacustris.
elongatulus, A. Dietr. Sp. Pl. ii. 96 = S. pumilus.
exiguus, A. Dietr. l. c. 98 = Eleocharis ochularis.
fluviatilis, Nees, in Linnæa, x. (1836) 165 = S. Ludwigii.
filiformis, A. Dietr. Sp. Pl. ii. 96 = Eleocharis tenuis.
fluitans, Link, Hort. Berol. i. 236 = S. fluitans.

ELEOGITON :—
lacustris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 173 = S. lacustris.
littoralis, Fourr. l. c. = S. littoralis.
longifolius, Nees, in Linnæa, ix. (1834) 291; x. (1836) 164 = S. flaccifolius.
monostachya, A. Dietr. Sp. Pl. ii. 97 = Fimbristylis schoenoides.
parvula, Link, Hort. Berol. i. 235 = S. nanus.
planifolius, A. Dietr. Sp. Pl. ii. 98 = S. planifolius.
radicans, A. Dietr. l. c. 97 = Eleocharis costulata.
Rothii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 178 = S. pungens.
rubicunda, Nees, in Linnæa, x. (1836) 164 = S. Ludwigii.
scabra, A. Dietr. Sp. Pl. ii. 99 = Fimbristylis acuminata.
scariosa, A. Dietr. l. c. 98 = Ficinia scariosa.
spadiceus, A. Dietr. l. c. 99 = S. spadiceus.
striata, Nees, in Linnæa, ix. (1834) 219; x. (1836) 165 = S. capillifolius.
Tabernaemontani, Fourr. in Ann. Soc. Linn Lyon, N. S. xvii. (1869) 178 = S. lacustris.
tenuissima, Nees, in Linnæa, x. (1836) 168 = S. tenuissimus.
triqueter, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 178 = S. triqueter.
verruculosa, A. Dietr. Sp. Pl. ii. 98 = S. verruculosus.

ELEPHANTINA, Bertol. Fl. Ital. vi. 279 (1844) = Rhynchoospora, Griseb. (Scrophul.).
proboscidea, Bertol. l. c. 280 = R. Elephas.

ELEPHANTODON, Salisb. Gen. Pl. Fragm. 12 (1866) = Dombeya, Linn.
eburnea, Salisb. l. c. = D. eburnea.

ELEPHANTOPSIS, A. Dietr. Syn. Pl. iv. 1372 (1847) = ELEPHANTOSIS, Less. = seq.

ELEPHANTOPUS, Linn. Gen. ed. I. 249 (1787). *COMPOSITAE*, Benth. & Hook. f. ii. 327.
DISTREPTUS, Cass. in Bull. Soc. Philom. (1817) 66; et in Dict. Sc. Nat. xiii. 366 (1819).
ELEPHANTOSIS, Less. in Linnæa, iv. (1829) 322.
MATAMORIA, La Llave & Lex. Nov. Veg. Desc. i. 8 (1824).
PERUDELEPHANTOPUS, Rohr, in Skrift. Nat. Selsk. Kiøbenh. iii. (1792) 214.
SPIROCHAETA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 166.
angustifolius, *Sw. Prod. Veg. Ind. Occ.* 115.—Am. trop.
biflorus, *Sch. Bip. in Linnaea*, xx. (1847) 519; *D. Dietr. Syn. Pl.* iv. 1372.—Bras.
carolinensis, G. F. W. Mey. Prim. Fl. Esseq. 256, 318 = scaber.
carolinianus, *Willd. Sp. Pl.* iii 2390.—Am. bor.
cernuus, Vell. Fl. Flum. viii. t. 143 = scaber.
cuneifolius, Vell. ex DC. Prod. v. 56, sphalm. = prace.
crispus, D. Dietr. Syn. Pl. iv. 1372 = spicatus.
cuneifolius, *Fourn. in Bull. Soc. Bot. Fr.* xxx. (1883) 186.—Reg. Argent.
dubius, Blanco, Fl. Filip. ed. II. 442 = spicatus.
elatus, Bertol. Misc. Bot. xi. 21 = tomentosus.
elongatus, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 426.—Bras.
flexuosus, Rafin. Fl. Ludov. 78 = carolinianus.
hirtiflorus, *DC. Prod.* v. 56.—Bras.
listrioides, Steud. Nom. ed. II. i. 549 = angustifolius.
Mertii, R. Grah. in Edinb. N. Phil. Journ. (Jan.–March 1830) 378 = scaber.
micropappus, *Less. in Linnaea*, vi. (1831) 689.—Bras.
mollis, H. B. & K. Nov. Gen. et Sp. iv. 26 = scaber.
nudatus, A. Gray, in Proc. Am. Acad. xv. (1880) 47 = radicalis ?
radicalis, Ell. Sketch. ii. 481 = tomentosus.
radicalis, *Poir. Encyc. Suppl.* ii. 543.—Am. bor.
rudiflorus, Willd. Sp. Pl. iii. 2390; Spreng. Syst. iv. 454 = angustifolius.
palustris, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 287.—Bras.
paniculatus, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 173 = hirtiflorus.
pratensis, *Wright, in Sauv. Fl. Cub.* 75.—Cuba.
quadriflorus, D. Dietr. Syn. Pl. iv. 1372 = angustifolius.
racemosus, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 427.—Bras.
Riedelii, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 175.—Bras.

ELEPHANTOPUS:—
riparius, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 425.
scaber, *Linn. Sp. Pl.* 814.—Cosmop. trop.
senegalensis, *Oliver & Hiern, in Oliver, Fl. Trop.* iii. 299.—Afr. trop.
setiferus, *R. Grah. in Edinb. N. Phil. Journ.* (Jan. March 1831) 378.—Ind. occ.
serratus, Blanco, Fl. Filip. ed. I. 635 = mollis.
sinuatus, *Zoll. & Mor. Syst. Verz. Zoll.* 99.—Java.
sordidus, Salisb. Prod. 212 = scaber.
spicatus, *Aubl. Pl. Gui.* ii. 808.—Guian.
strigosus, Willd. ex DC. Prod. v. 87 = spicatus.
tomentosus, *Linn. Sp. Pl.* 814.—Am. bor.
vaginatus, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 426.—Bras.
violaceus, *Sch. Bip. in Linnaea*, xx. (1847) 517.—Am. bor.
virgatus, *Desv. in Ham. Prod. Pl. Ind. Occ.* 52.—Guian.

ELEPHANTORRHIZA, Benth. in Hook. Journ. Bot. iv. (1842) 344. *LEGUMINOSAE*, Benth. & Hook. f. i. 590.
Burchellii, *Benth. l. c.*—Afr. austr.
Burkii, *Benth. l. c.* v. (1846) 21.—Afr. austr.
Petersiana, *Bolle, in Petersi, Reise Mossamb. Bot.* 9.—Afr. trop.

ELEPHANTOSIS, Less. in Linnæa, iv. (1829) 322 = **Elephantopus**, Linn (Compos.).
angustifolia, DC. Prod. v. 87 = Elephantopus angustifolius.
biflora, Less. in Linnæa, iv. (1829) 322 = Elephantopus biflorus.
listrioides, Flsch. & Mey. Ind. Sem. Hort. Petrop. i. 27 = Elephantopus angustifolius.
quadrifolia, Less. in Linnæa, iv. (1829) 323 = Elephantopus angustifolius.

ELEPHANTUSIA, Willd. Sp. Pl. iv. 1156 = **Phytelephas**, Ruiz & Pav. (Palm.).
macrocarpa, Willd. l. c. = P. macrocarpa.
microcarpa, Willd. l. c. 1157 = P. microcarpa.

ELEPHAS, Tourn. ex Adans. Fam. ii. 211 (1763); Guss. Fl. Sic. Prod. ii. 155 (1828) = **Rhynohoocrys**, Griseb. (Scrophul.).
Columnae, Guss. l. c. = R. Elephas.
incana, G. Don, Gen Syst. iv. 619 = R. orientalis.
orientalis, Guss. Fl. Sic. Prod. ii. 156, in obs. = R. orientalis.
recta, G. Don, Gen Syst. iv. 619 = R. Elephas.

ELETTARIA, Maton, in Trans. Linn. Soc. x. (1811) 250. *SCITAMINEAE*, Benth. & Hook. f. iii. 646.
CARDAMOMUM, Salisb. in Trans. Hort. Soc. i. (1812) 282.
alba, *Blume, Enum. Pl. Jav.* 58.—Java.
anthodiodes, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 892.—Java.
atropurpurea, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv (1862) 326.—Sumatra.
cannaecarpa, Wight. Ic. t. 2007 = Amomum aculeatum.
Cardamomum, *Maton, in Trans. Linn. Soc.* x. (1811) 254.—Ind. or.
coccinea, Blume, Enum. Pl. Jav. 59 = Amomum coccineum.
costata, Volgt, Hort. Saburb. Calc. 568 = Amomum costatum
Diepenhorstii, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 326.—Sumatra.
floribunda, Blume, Enum. Pl. Zeyl. 319 = Amomum floribundum.
fusiosa, Blume, Enum. Pl. Jav. 59 = Amomum fusiosa.
globerrima, Zoll. & Mor. Syst. Verz. Zoll. 84 = Amomum globerrimum.
hemisphaerica, Blume, Enum. Pl. Jav. 51 = Alpinia hemisphaerica.
involucrata, Thw. Enum. Pl. Zeyl. 319 = Amomum involucratum.
linguiformis, *Schult. Mant.* i. 35.—Ind. or.
macrocephala, *Miq. Fl. Ind. Bat.* iii. 608.—Java.
major, Sm. in Rees, Cyclop. xxxix = E. Cardamomum.
media, Link, ex Horan. Prod. Scitam. 31 = costata.
minor, *Blume, Enum. Pl. Jav.* 58.—Java.
minuta, *Blume, l. c.* 53.—Java.
mollis, Blume, l. c. = Alpinia mollis.
musacea, *Horan. Prod. Scitam.* 31.—Ins. Amboina.

ELETTARIA :—
nemoralis, Thw. Enum. Pl. Zeyl. 319 = Amomum nemorale.
pallida, Blume, Enum. Pl. Jav. 52.—Java.
paludosa, *Mig. Fl. Ind. Bat.* iii. 602.—Java.
Pininga, Miq. l. c. = Alpinia Pininga.
punicea, Schult. Mant. i. 24.—Sumatra.
rosea, Teijsm. & Binn. ex Benth. & Hook. f. Gen. iii. 644 = Amomum roseum.
rubra, Miq. Fl. Ind. Bat. iii. 602 = Alpinia rubra.
rufescens, Thw. Enum. Pl. Zeyl. 480 = Amomum rufescens.
Scottiana. *F. Muell. Fragm.* viii. 24.—Austral.
solaris, *Blume, Enum. Pl. Jav.* 52.—Java.
speciosa, *Blume*, l. c. 51.—Java.
tomentosa, Miq. Fl. Ind. Bat. iii. 603 = Alpinia tomentosa.
villosa, Miq. l. c. = Amomum villosum.
Walang, Miq. l. c. = Alpinia Walang.

ELEUSINE, Gaertn. Fruct. i. 7. t. 1 (1788). *GRAMINEÆ*, Benth. & Hook. f. iii. 1172.
ACRACHNE, Wight & Arn. in Lindl. Introd. Nat. Syst ed. II. 381 (1836).
DACTYLOCTENIUM, Willd. Enum. Hort. Berol. 1029 (1809).
aegyptiaca, *Desf. Fl. Atlant.* i. 85.—Cosmop. trop. et subtrop.
arabica, Hochst. ex Steud. Syn. Pl. Gram. 211 = flagellifera.
aristata, *Ehrenb. ex Boiss. Fl. Orient.* v. 557, in syn.—Baluchist.
barbata, *Desv. Opusc.* 104.—Bras.
barcinonensis, *Costa, Ind. Sem Hort. Barcin.* (1859); *et Willk. in Linnaea*, xxx. (1860) 125.—Hispan.
brevifolia, Hochst. ex Steud. Syn. Pl. Gram. 211 = Eragrostis coelachyrum.
caespitosa, *A. Rich. Tent Fl. Abyss.* ii. 412.—Abyss.
calycina, Roxb. Hort. Beng. 8; Fl. Ind. i. 346 = Leptochloa calycina.
cerealis, *Ehrenb. ex Sweet, Hort. Brit.* ed. II. 571.—Aegypt.
cerealis, Salisb. Prod. 19 = coracana.
chinensis, F. Muell. Census, 134 = Leptochloa chinensis.
ciliata, *Rafin. in Desv. Journ. Bot.* iv. (1814) 278 = aegyptiaca.
coracana, *Gaertn. Fruct.* i. 8. t. 1.—Am. austr.; Ind. or.; Aegypt.
cruciata, Lam. Illustr. t. 208 = aegyptiaca.
Dagussa, Schimp. in Regel, Gartenfl. (1872) 205 = E. Tocussa.
Digitaria, Willd. ex Steud. Nom. ed. II. 549 = Leptochloa Digitaria.
digitata, Spreng. Syst. iv. Cur. Post. 86 = Leptochloa subdigitata.
distachya, Trin. ex Steud. Nom. ed. II. i. 549 = Indica.
distans, *Link, Enum. Hort. Berol.* i. 61.—Hab.?
distans, Moench, Meth. 210 = indica.
domingensis, Pers. Syn. i. 87 = Leptochloa domingensis.
domingensis, Sieber, ex Schult. Mant. ii. 323 = indica.
dura, Lam. Illustr. i. 208 = Poa dura.
elongata, Willd. ex Steud. Nom. ed. II. i. 549 = Leptochloa filiformis.
erythrosperma, Jan, ex Steud. l. c. = oligostachya.
falcata, Eckl. ex Steud. l. c. = Michrochloa caffra.
falcata, Spreng. Syst. i. 349 = Harpechloa capensis.
filiformis, Pers. Syn. i. 87 = Leptochloa filiformis.
flagellifera, *Nees, in Linnaea*, xvi. (1842) 220.—Aa. occ. et Afr. trop.
flocciflola, *Spreng. Syst.* i. 350.—Arabia.
geminata, *Lange, Ind. Sem. Hort. Haun.* (1861) 28.—Hispan.
glabra, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 58.—Afr. trop.
glaucophylla, *Munro, ex Benth. in Journ. Linn. Soc.* xix. (1881) 107.—Afr. trop.
gunantha, *Schrank, in Syll. Ratisb.* i. (1824) 191.—Bras.
Gussini, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 563 = indica.
gracilis, Salisb. Prod. 19 = indica.
gracilis, *Spreng. Syst.* i. 350 = Leptochloa gracilis.
inaequalis, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 563 = indica.
indica, *Gaertn. Fruct.* i. 8.—Cosmop. trop. et subtrop.

ELEUSINE:—
indica, Steud. Syn. Pl. Gram. 211 = japonica.
japonica, *Steud. l. c.*—Japon.
juncea, *Boj Hort. Maurit.* 270.—Madag.
leucosperma, Bernh. ex Steud. Nom. ed. II. i. 549 = oligostachya.
Limo, Lam. Illustr. i. 208 = Cynosurus Limo.
macrosperma, Stokes, Bot. Mat. Med. i. 150 = Cynosurus indicus.
macrostachya, *Benth. in Journ. Linn. Soc.* xix. (1881) 107.—Madag.
marginata, Lindl. in Mitch. Three Exped. i. 319 = indica.
monostachya, Spreng. Syst. i. 349 = Leptochloa monostachya.
mucronata, Michx. Fl. Bor. Am. i. 65 = Leptochloa mucronata.
mucronata, Stokes, Bot. Mat. Med. i. 150 = aegyptiaca.
multiflora. *Hochst. in Flora*, xxiv. (1841) t. *Intell.* 20 nomen; *et in A. Rich. Tent. Fl. Abyss.* ii. 412.—Aegypt.
oligostachya, *Link, Hort. Berol.* i. 60.—Bras.
ovalis, *Ehrenb. ex Sweet, Hort. Brit.* ed. II. 571.—Aegypt.
pectinata, Moench, Meth. Suppl. 68 = aegyptiaca.
penicillata, Spreng. Syst. i. 350 = Chloris penicillata.
polydactyla, *Steud. Syn. Pl. Gram.* 211.—Afr. trop.
polystachya, F. Muell. Fragm. i. 216 = Leptochloa subdigitata.
procera, Spreng. ex Steud. Nom. ed. II. i. 549 = Uniola racemiflora.
prostrata, Spreng. Syst. i. 350 = aegyptiaca.
racemosa, Heyne, in Roth, Nov. Sp. Pl. 80 = verticillata.
radulans, R. Br. Prod. 186 = aegyptiaca.
rigida, *Spreng. Syst. iv. Cur. Post.* 86.—Hab.?
rigidiflola, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 566 = indica.
scabra, Fourn. ex Hemsl. l. c. = indica.
sparsa, Mudd. Desc. Gram. Uber. 135 = Leptochloa filiformis.
sphaerosperma, Stokes, Bot. Mat. Med. i. 149 = coracana.
stolonifera, *R. Br. in Salt, Abyss. App.* 63 = Abyssin.
stricta, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 345.—Ind. or.
stricta, Willd. ex Steud. Nom. ed. II. i. 549 = Leptochloa chinensis.
tenerrima, Hornem. Hort. Hafn. i. 79 = Leptochloa chinensis.
Tocussa, *Fresen. in Mus. Senckenb.* ii. (1837) 141.—Hab.?
tristachya, *Schrank, in Syll. Ratisb.* i. (1824) 191.—Hab.?
tristachya, *Kunth, Rév. Gram.* i. 92.—Reg. Argent.
tristachya, Lam. Illustr. i. 208 = indica.
unioloides, Willd. ex Steud. Nom. ed. II. i. 549 = Leptochloa virgata.
verticillata, Roxb. Hort. Beng. 8; Fl. Ind. i. 346.—Geront. trop.
virgata, Pers. Syn. i. 87 = Leptochloa virgata.

ELEUTHERANTHERA, Poit. ex Bosc, in Nouv. Dict. Hist. Nat. ed. I. vii. 498 (1803). *COMPOSITAE*, Benth. & Hook. f. ii. 371.
CHALARIUM, Poit. ex DC. Prod. v. 546, in syn. (1836).
FINGALIA, Schrank, in Syll. Ratisb. i. (1824) 67.
KEGELIA, Sch. Bip. in Linnaea, xxi. (1848) 245.
OCHERA, Cass. in Bull. Soc. Philom. (1818) 32; et in Dict. Sc. Nat. xxxv. 445 (1825).
ovata, *Poit. ex Steud. Nom.* ed. II. i. 549.—Am. trop.
pinnata, Sch. Bip. in Bot. Zeit. xxiv. (1866) 329 = ovata.
ruderalis, Sch. Bip. l. c. = ovata.

ELEUTHERINE, Herb. in Bot. Reg. (1843) t. 57. *IRIDEAE*, Benth. & Hook. f. iii. 695.
GALATEA, Salisb. in Trans. Hort. Soc. i. (1812) 330, nomen prius.
anomala, *Herb. in Bot. Reg.* (1843) t. 57.—Am. trop.
plicata, *Herb. l. c.* sub t. 57.—Ind. occ.

ELEUTHEROCARPUM, Schlecht. in Lechl. Berb. Am. Austr. 59 (1857) = Cotocomolaes, Lindl. (Rosac.).
escalloniaefolium, Schlecht. l. c.—Am. austr.

ELEUTHEROCOCCUS, Maxim. Prim. Fl. Amur. 123 (1859). *ARALIACEAE*, Benth. & Hook. f. i. 941.
senticosus, *Maxim. l. c.*—Manchur.

ELEUTHEROPETALUM, H. Wendl. ex Oerst. in Vidensk. Meddel. 1858 (1859) 6 = **Chamaedorea**, Willd. (Palm.).
Ernesti-Augusti, Oerst. l. c. 7 = C. Ernesti-Augusti.
Sartorii, Liebm. ex Oerst. Amérique Centrale, 18. t. 3 f. 1–6 = C. Sartorii.

ELEUTHEROSPERMUM, C. Koch, in Linnaea, xvi. (1842) 365. *UMBELLIFERAE*, Benth. & Hook. f. i. 872.
cicutarium, Boiss. Fl. Orient. ii. 924 = grandifolium
grandifolium, *C. Koch, in Linnaea*, xvii. (1843) 81.—Armenia.
laxicum, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 924.—Aa. Min.

ELEUTHEROSTEMON, Klotzsch, in Linnaea, xii. (1838) 219 = **Philippia**, Klotzsch (Ericac.).
galioides, Klotzsch, l. c. = P. galioides.
multiglandulosum, Klotzsch, l. c. 220 = P. viscosa

ELEUTHRANTHERON, Steud. Nom. ed. II. i. 549 (1840) = **Elantheranthera**, Poit. (Compos.).

ELEUTHRANTHES, F. Muell. Fragm. iv. 92 (1864). *RUBIACEAE*, Benth. & Hook. f. ii. 142.
opercularina, *F. Muell. l. c.*—Austral.

ELIAEA, Cambess. in Ann. Sc. Nat. Sér. I. xx. (1830) 400. t. 13. *HYPERICINEAE*, Benth. & Hook. f. i. 166.
CUSSONIA, Endl. Gen. 1035, in syn. (1840).
LANIGEROSTEMMA, Chapelier, ex Endl. l. c. (1840).
articulata, *Cambess. in Ann. Sc. Nat. Sér.* I. xx. (1830) 401.—Madag.

ELICHRYSUM, Tourn. ex Linn. Gen. ed. I. 250 (1737) = **Helichrysum**, Vaill. (Compos.).

ELICTOTRICHON, Bess. ex Andrz. Rys Botan. i (1823) 9 = **Avena**, Linn. (Gramin.).
sempervirens, Bess. l. c. = A. Besseri.

ELIDE, Medic. Phil. Bot. ii. 71 (1791) = **Asparagus**, Tourn. (Liliac.).

ELIDURANDIA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 450 = **Fugosia**, Juss. (Malvac.).
texana, Buckl. l. c. = F. Drummondii.

ELIEA, G. Don, Gen. Syst. i. 613 = **Eliaea**, Cambess. (Hypericin.).

ELIGIA, Dum. Anal. Fam. 62 (1829) = **Elegia**, Linn (Restiac.).

ELIMUS, Nocca, in Usteri, Ann. Bot. v. (1793) 44 = **Elymus**, Linn. (Gramin.).

ELIOKARMOS, Rafin. Fl. Tellur. ii. 94 (1836) = Ornithogalum, Tourn. (Liliac.).
asarum, Rafin. l. c. = O. arabicum.
caudatum, Rafin. l. c. = O. caudatum.
coarctatum, Rafin. l. c. = O. coarctatum.
maculatus, Rafin. l. c. iii. 59 = O. maculatum.
thyroides, Rafin. l. c. ii. 24 = O. thyroides.

ELIONURUS, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 941 (1806) (Elyonurus). *GRAMINEAE*, Benth. & Hook. f. iii. 1199.
CALLICHLOEA, Spreng. ex Steud. Nom. ed. II. i. 257 (1840).
argenteus, *Nees, Fl. Afr. Austr.* 96.—Abyss.; Afr. austr.
bilingula, *Hoch. in Mart. Fl. Bras.* ii. iii. 308.—Bras.
candidus, *Hack. l. c.* 306.—Am. austr.
ciliaris, H. B. & K. Nov. Gen. et Sp. i. 193. t. 63 = tripsacoides.
ciliaris, Nees, Agrost. Bras. 356 = candidus.
citreus, *Munro, ex Benth. Fl. Austral.* vi 310.—Austral.
dubius, Nees, Agrost. Bras. 356 = Andropogon tener.
elegans, *Kunth, Rév. Gram.* i. 161.—Afr. trop.

ELIONURUS :—
Griesbachii, J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 154 = E. Royleanus.
hirsutus, *Munro, ex Benth. in Journ. Linn. Soc.* xix. (1881) 68.—Persia.
latiflorus, *Nees, ex Steud. Syn. Pl. Gram.* 364.—Bras. ; Parag.
Nuttallii, Vasey, Cat. Gr. U. St. 25 = tripsacoides.
pertusus, Nees, ex Steud. Syn. Pl. Gram. 364 = Andropogon pertusus.
platypus, *Hack.*—*Cf. Bolet. Soc. Broter.* iii. (1885) 135.—Afr. trop.
prostratus, Kunth, Enum. Pl. i. 480, sphalm. = rostratus.
ostratus, *Nees, Agrost. Bras.* 357.—Bras.
Royleanus, *Nees, ex A. Rich. Tent. Fl. Abyss.* ii. 474.—Ind. or. ; Arab. ; Afr. trop.
thymolodorus, *Nees, Fl. Afr. Austr.* 95.—Afr. austr.
tripsacoides, *H. B. & K. Nov. Gen. et Sp.* i. 192. t. 62.—Am. bor. et trop.

ELIOPIA, Rafin. Sylva Tellur. 90 (1838) = **Heliotropium**, Tourn. (Borag.).
riparia, Rafin. l. c.—Hab. ?
serrata, Rafin. l. c. = H. indicum.

ELIOTTIA, Stend. Nom. ed. I. 326 (1821) = **Elliottia**, Muhl. (Ericac.).

ELISANTHE, Reichb. Nom. 206 (1841) = **Silene**, Linn. (Caryophyll.).
Bridgesii, Rupr. Fl. Cauc. 200 = S. Bridgesii.
Chamissowii, Rupr. l. c. 201 = S. verecunda.
Drummondii, Rupr. l. c. 200 = S. Scouleri.
Hindsii, Rupr. l. c. 200.—Calif.
noctiflora, Rupr. Fl. Ingric. i. 161 = S. noctiflora.
Scouleri, Rupr. Fl. Cauc. 200 = S. Scouleri.
viscosa, Rupr. l. c. = S. viscosa.
Zawadskii, Fuss, Fl. Transs. 106 = S. Zawadskii.

ELISARRHENA, Benth. & Hook. f. Gen. i. 961 (1867), sphalm. = **Elisarrhena**, Miers.

ELISENA, Herb. Amaryll. 201 (1837). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 784.
LIRIOPE, Herb. App. 41 (1821).
LIRIOPSIS, Reichb. Consp. 22 (1828).
longipetala, *Lindl. Bot. Reg.* (1838) *Misc.* 45.—Peruv.
marginata, M. Roem. Syn. Enast. 63 = Nerine marginata.
ringens, *Herb. Amaryll.* 201.—Peruv.
sublima, *Herb. in Bot. Mag.* t. 3873.—Peruv.

ELISMA, Buchen. in Prisgsh. Jahrb. vii. (1869) 25. *ALISMACEAE*, Benth. & Hook. f. iii. 1005.
natans, *Buchen. l. c.*—Europ.

ELISSARRHENA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiii. 194 (1864) ; xx. (1867) 169 = **Anomospermum**, Miers (Menisperm.).
longipes, Miers, ll. cc. = A. grandifolium.

ELISABETHIA, Schomb. ex Benth in Hook. Journ. Bot. ii. (1840) 92. *LEGUMINOSAE*, Benth. & Hook. f. i. 577.
coccinea, *Schomb. ex Benth. l. c.*—Guiana.
macrocalyx, *Benth. in Mart. Fl. Bras.* xv. ii. 214.—Bras.
Princeps, *Schomb. ex Benth. in Hook. Journ. Bot.* ii. (1840) 92.—Guiana.

ELIZALDIA, Willk. Strand- u. Steppengeb. Iber. 128 (1859) = **Wonnea**, Moench (Boragin.).
nouvoides, Willk. l. c. 129 = N. multicolor.

ELKAJA, M. Roem. Syn. Hesper. 116 (1846) = ELCAJA, Fersk. = **Trichilia**, Jum. (Rutac.).

ELKANIA, Schlecht. ex Wedd. in DC. Prod. xvi. i. 227, in syn. (1869) = **Fouquiera**, Gaudich. (Urticac.).
multinervis, Schlecht. ex Wedd. l. c. = P. cymosa.

ELLBANTEWS, Presl, Rel. Haenk. i. 97 (1827). *ORCHIDEAE*, Benth. & Hook. f. iii. 522.
EVALWA, Poepp. & Endl. Nov. Gen. et Sp. i. 32. t. 56 (1838).
cochleariformis, *Reichb. f. in Walp. Ann.* vi. 479.—Peruv.
erypophyllostachya, *Reichb. f. l. c.*—Venezuela.

ELLBANTHUS:—
aurantiacus, *Reichb. f. in Walp. Ann.* vi. 482.—Bras.
aureus, *Reichb. f. l. c.* 484.—Peruv.
Bonplandi, *Reichb. f. l. c.* 488.—Peruv.
bractescens, *Reichb. f. l. c.* 475.—Mexic.
brasiliensis, *Reichb. f. l. c.*—Bras.
capitatus, *Reichb. f. l. c.*—Mexic.
Carawata, *Reichb. f. Otia Bot. Hamb.* 92, in obs.—Guian.
casapanis, *Reichb. f. in Walp. Ann.* vi. 475.—Peruv.
cephalophorus, *Reichb. f. l. c.* 476.—Peruv.
colommaris, *Reichb. f. l. c.* 483.—N. Granat.
cosifer, *Reichb. f. l. c.* 474.—Peruv.
corilifolius, *Reichb. f. l. c.* 478.—N. Granat.
crinipes, *Reichb. f. Otia Bot. Hamb.* 92.—Bras.
cyanrocephalus, *Reichb. f. in Walp. Ann.* vi. 476.—Hab. ?
discolor, *Reichb. f. l. c.* 480.—Peruv.
ensatus, *Reichb. f. l. c.* 482.—Ecuador.
flavescens, *Reichb. f. l. c.* 479.—Mexic.
furfuraceus, *Reichb. f. l. c.* 480.—Mexic.
gracilis, *Reichb. f. l. c.* 481.—Venezuela.
haematozanthus, *Reichb. f. l. c.* 485.—N. Granat.
Hallii, *Reichb. f. l. c.* 482.—Peruv.
hymenophorus, *Reichb. f. l. c.* 480.—Mexic.
kermesinus, *Reichb. f. l. c.* 478.—Venemela.
lanciflorus, *Presl, Rel. Haenk.* 97.—Ecuador.
lepidus, *Reichb. f. Otia Bot. Hamb.* 92, in obs.—Hab. ?
Lindenii, *Reichb. f. in Walp. Ann.* vi. 477.—Venezuela.
linifolius, *Presl, Rel. Haenk.* 97.—Peruv.
lupulinus, *Reichb. f. in Walp. Ann.* vi. 483.—Mexic.
maculatus, *Reichb. f. l. c.* 482.—Ecuador.
Myrosmatis, *Reichb. f. l. c.* 473.—Peruv.
oliganthus, *Reichb. f. l. c.* 481.—Peruv.
purpureus, *Reichb. f. l. c.* 478.—Am. trop.
rhodolepis, *Reichb. f. l. c.* 482.—Peruv.
robustus, *Reichb. f. l. c.* 474.—Peruv.
Ruizii, *Reichb. f. in Linnaea*, xxii. (1849) 842.—Peruv.
strobiliifer, *Reichb. f. in Walp. Ann* vi. 483.—Peruv.
Wagneri, *Reichb. f. l. c.* 474.—Venezuela.
xanthrocoma, *Reichb. f. in Bot. Mag* t. 6016.—Peruv.

ELLEBORUS, Vill. Hist. Pl. Dauph. iii (1789) 716 = **Helleborus**, Tourn. (Ranunc.).

ELLERTONIA, Wight, Ic. iv. 2. t. 1295 (1848). *APOCYNACEAE*, Benth. & Hook. f. ii. 705.
KANETTA, Kostel. Allg. Med. Pharm. Fl. iii. 1062 (1834), nomen prius.
madagascariensis, Radlk. in Bremen, Abh. viii. (1883) 402 = Alyxia erythrocarpa.
Rheedii, *Wight, Ic.* iv. t. 1295.—Ind. or.

ELLIMIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 125 (1838) = **Oligomeris**, Cambess (Compos.).
ruderalis, Nutt. l. c. = O. subulata.

ELLIOTIA, Spach, Hist. Vég. Phan. ix. 443 (1840) = seq.

ELLIOTTIA, Muhl. ex Nutt. Gen. Am. Add. (1817). *ERICACEAE*, Benth. & Hook. f. ii. 598.
TRIPETALRIA, Sieb. & Zucc. in Abh. Akad. Muench. iii. ii. (1843) 721. t. 3. f. 2.
Muhlea, Stend. Nom. ed. I. 226 = racemosa.
racemosa, *Muhl. ex Ell. Sketch.* i. 448.—Am. bor.

ELLIPANTHUS, Hook. f. in Benth. & Hook. f. Gen. i. 484 (1862). *CONNARACEAE*, Benth. & Hook. f. i. c.
calophyllus, *Kurz, in Journ. As. Soc. Beng.* xli. (1882) ii. 205.—Malaya.
Griffithii, *Hook. f. Fl. Brit. Ind.* ii. 56.—Malaya.
Helferi, *Hook. f. l. c.* 55.—Burma ; Malaya.
monophyllus, *Hook. f. in Benth. & Hook. f. Gen.* i. 484.—Burma.
Thwaitesii, Hook. f. Fl. Brit. Ind. ii. 55 = unifoliolatus.
tomentosus, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 205 = monophyllus.
unifoliolatus, *Hook. f. in Benth. & Hook. f. Gen.* i. 484.—Zeylan.

ELLIPHIA, Hook. f. & Thoms. Fl. Ind. i. 104 (1855). *ANONACEAE*, Benth. & Hook. f. i. 33.
cinerascens, *Miq. Ann. Mus. Lugd. Bat.* ii. 10.—Sumatra.
coriacea, *Scheff. in Ann. Jard. Buitenz.* ii. (1883) 4.—Borneo.

ELLIPSIA:—
cuneifolia, *Hook. f. & Thoms. Fl. Ind.* i. 104.—Malacca.
ferruginea, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 52.—Burma.
gilva, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 9.—Borneo.
glabra, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 52.—Malacca.
lalia, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 10.—Sumatra.
nervosa, *Hook. f. & Thoms. Fl. Brit. Ind.* i. 52.—Malacca.

ELLISIA, P. Br. Hist. Jamaic. 263 (1756) = **Duranta**, Linn. (Verbenac.).
acuta, Linn. Amoen. Acad. v. 400 = D. Plumieri.

ELLISIA, Linn. Sp. Pl. ed. II. 1663 (1763). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 827.
EUCRYPTA, Nutt. in Journ. Acad. Philad. N. S. i. (1847) (Pl. Gamb.) 158.
MACROCALYX, Trew, in Act. Nat. Cur. ii. (1761) 382.
NYCTELEA, Scop. Introd. 775 (1777).
ambigua, Nutt Gen. Am. i. 118 = E. Nyctelea.
chrysanthemifolia, *Benth. in Trans. Linn. Soc.* xvii. (1837) 274.—Calif.
membranacea, *Benth. l. c.*—Calif.
microcalyx, Nutt. in Trans. Am. Phil. Soc. v. (1837) 191 = Nemophila microcalyx.
Nyctelea, *Linn. Sp. Pl.* ed. II. 1662.—Am. bor.
ranunculacea, Nutt. in Trans. Am. Phil. Soc. v. (1837) 191 = Nemophila microcalyx.
Torreyi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 302.—Am. bor. occ.

ELLISIOPHYLLUM, Maxim. in Bull. Acad. Pétersb. xvi. (1871) 228. *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 828.
reptans, *Maxim. l. c.*—Japon.

ELLOBIUM, Lilja, in Linnaea, xv. (1841) 262 = **Fuchsia**, Linn. (Onagrar.).
fulgens, Lilja, l. c. = F. fulgens.

ELLOBUM, Bisme, Bijdr. 746 (1826) = **Vandellia**, Linn. (Scrophul.).
montanum, Blume, l. c. 747 = V. Ellobum.

ELMICERA, Reichb. Consp. 128 (1828) = **Pentstemon**, Mitch. (Scrophul.).
barbata, Reichb. ex Steud. Nom. ed. II. i. 550 = P. barbatus.

ELODEA, Jack, in Malay. Misc. ii. (1822) n. VII. 21 = **Oratoxylon**, Blume (Hypericin.).
chinensis, Hance, in Hook. Lond. Journ. Bot. vii. (1848) 472 = C. polyanthum.
egyptica, Jack, in Malay. Misc. ii. (1822) n. VII. 25, in obs. = Hypericum aegyptiacum.
formosa, Jack, l. c. n. 24 = C. formosum.
pulchella, Hort. Chels. ex Lond. Hort. Brit. Suppl. i. 587 = C. polyanthum.
sumatrana, Jack, in Malay. Misc. ii. (1822) n. VII. 22 = C. sumatranum.

ELODEA, Juss. Gen. 255, in obs. (1789) ; Vent. Tabl. iii. 144 (1798) = **Hypericum**, Tourn.
axillaris, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 170 = H. paludosum.
campanulata, Pursh, Fl. Am. Sept. ii. 379 = H. virginicum.
crassifolia, Blume, Mus. Bot. Lugd. Bat. ii. 15.—Japon.
Drummondii, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 167 = H. virginicum.
emarginata, Steud. Nom. ed. II. i. 550 = H. virginicum.
floribunda, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 169 = H. paludosum.
Fraseri, Spach, l. c. 168 = H. virginicum.
japonica, Blume, Mus. Bot. Lugd. Bat. ii. 15 = H. japonicum ?
palustris, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 171 = H. Elodes.
pauciflora, Spach, l. c. 169 = H. virginicum.
petiolata, Pursh, Fl. Am. Sept. ii. 379 = H. paludosum.
Ruseggeri, Walp. Rep. i. 891 (= *Triadenia Ruseggeri*).—Syria.
tubulosa, Pursh, Fl. Am. Sept. ii. 379 = H. paludosum.
virginica, Nutt. ex Torr. & Gray, Fl. N. Am. i. 167 = H. virginicum.

ELODEA. Michx. Fl. Bor. Am. i. 20 (1803). *HYDROCHARIDEÆ*, Benth. & Hook. f. iii. 450.
ANACHARIS, Planch. in Ann. & Mag. Nat. Hist. Ser. II. i. (1848) 25.
APALANTHE, Planch. l. c. 27 (1848).
DIPLANDRA, Bert. in Mercv. Chil. 612; ex Bull. Férussac, xx. (1830) 110.
EGERIA, Planch. in Ann. Sc. Nat. Sér. III. xi (1849) 79.
PHILOTRIA, Rafin. in Am. Monthly Mag. (1818) 175.
UDORA, Nutt. Gen. Am. ii. 242 (1818).
callitrichoides, *Casp. in Monatsb. Berl. Acad.* (1857) 47.—Am. austr.
canadensis, *Michx. Fl. Bor. Am.* i. 20.—Am. bor.
chilensis, *Casp. in Monatsb. Berl. Acad.* (1857) 47.—Chili.
densa, *Casp. l. c.* 49.—Reg. Argent.
grandensis, *Irmsch. & Bongl. Pl. Arquin.* ii. 50.—N. Granat.
guianensis, *Rich. in Mém. Institt. Pr.* ii. (1811) 4, 75. —Guiana.
latifolia, Casp. in Monatsb. Berl. Acad. (1857) 46 = canadensis.
Nalas, *Casp. l. c.* 49.—Bras.
orinocensis, *Rich. in Mém. Institt. Pr.* ii. (1811) 75.—Venezuela.
Planchonii, *Casp. in Monatsb. Berl. Acad.* (1857) 47.—Am. bor.
Schweinitzii, Casp. l. c. 46 = canadensis.

ELODES, Adans. Fam. ii. 444 (1763); Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 171 = **Hypericum**, Tourn.
aegyptiaca, Payer, Organog. 8. t. 1 = H. aegyptiacum.
palustris, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 172 = H. Elodes.
petiolata, Rafin. Fl. Tellur. iii. 79 = H. paludosum.
tubulosa, Rafin. l. c. = H. paludosum.

ELOPIUM, Schott, in Oestr. Bot. Zeitschr. (1855) 84 = **Philodendron**, Schott (Aroid.).
surinamense, Schott, l. c. 85 = P. surinamense.

ELPHEGEA, Cass. in Bull. Soc. Philom. (1818) 31; et in Dict. Sc. Nat. xiv. 361 (1819) = **Palædia**, Jacq. (Compos.).
crenata, Cass. in Dict. Sc. Nat. xiv. 362 = P. balsamica.
dentata, Cass. l. c. 363 = P. dentata.
hirta, Cass. l. c. 364 = Conyza lithospermifolia.
lanceolata, Cass. l. c. 363 = P. linearifolia.
latifolia, Cass. l. c. 362 = P. balsamica.
minor, Cass. l. c. = P. integerrima.
quinquenervia, Cass. l. c. 363 = P. integerrima.

ELPHEGEA, Less. Syn. Comp. 182 (1832) = **Felicia**, Cass. (Compos.).
Borgiana, Less. l. c. = Aster Bergianus.
ciliata, Less. in Linnæa, vi. (1831) 126 = Mairia tastifolia.
microcephala, Less. Syn. Comp. 184 = Mairia microcephala.
reflexa, Less l. c. = F. reflexa.

ELSBOLZIA, Reichb. Consp. 116 (1828) sphalm. = **Blaholtsia**, Willd. (Labiat.).

ELSCHOTZIA, Brongn. Enum. Gen. 68 (1843) = prsec.

ELSHOLTIA, Koch, Syn. Fl. Germ. ed. II. 631 (1837) = prsec.

ELSHOLTZIA, Willd. in Roem. & Usteri, Mag. iv. xi. (1790) 3. *LABIATÆ*, Benth. & Hook. f. ii. 1181.
APHANOCHILUS, Benth. in Bot. Reg. sub t. 1932 (1839).
CYCLOSTEGIA, Benth. l. c. (1839).
barbinervis, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 104 = Comanthosphace barbinervis.
Beddomei, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 648.—Burma.
blanda, *Benth. Lab. Gen. et Sp.* 169.—As. trop.
cristata, *Willd. in Roem. & Usteri, Mag.* iv. xi. (1790) 5.—Europ.; As. bor.
densa, *Benth. Lab. Gen. et Sp.* 714.—Reg. Himal.
elata, *Zoll. & Mor. in Nat. en Genees. Arch. Nebrl. Indie*, ii. (1845) 5.—Java.
eriantha, *Benth. in DC. Prod.* viii. 161.—Java.

ELSHOLTIA:—
eriostachya, *Benth. Lab. Gen. et Sp.* 168.—Reg. Himal.
eriostachya, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 645 = densa.
flava, *Benth. Lab. Gen. et Sp.* 161.—Reg. Himal.
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 644.—Reg. Himal.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 108 = eriostachya.
incisa, *Benth. Lab. Gen. et Sp.* 162.—Reg. Himal.; Burma.
integrifolia, *Benth. l. c.* 714.—China.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 103 = Comanthosphace japonica.
javanica, Blume, Bijdr. 825 = Plectranthus javanicus.
leptostachya, *Benth. Lab. Gen. et Sp.* 715.—Reg. Himal.
mollissima, *Benth. l. c.* 168.—Java.
scimoides, Pers. Syn. ii. 114 = Geniosporum prostratum.
oppositifolia, Poir. in Dict. Sc. Nat. xiv. 366 = Colebrookea oppositifolia.
paniculata, Willd. Sp. Pl. iii. 59 = Pogostemon paniculatus.
Patrini, Garcke, Fl. Deutschl. ed. IV. 357 = cristata.
pilosa, *Benth. Lab. Gen. et Sp.* 163.—Reg. Himal.
polystachya, *Benth. l. c.* 161.—Reg. Himal.
pubescens, *Benth. l. c.* 169.—Java.
pusilla, Benth. l. c. 714 = eriostachya.
Schimperi, *Hochst. ex Vatke, in Linnæa*, xxxvii. (1871–73) 325, nomen.—Abyss.
Staunton, *Benth. l. c.* 161.—China.
stellipila, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 103 = Comanthosphace stellipila.
strobilifera, *Benth. Lab. Gen. et Sp.* 163.—Reg. Himal.
subdentata, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 103 = Comanthosphace sublanceolata.
Thompsoni, *Hook. f. Fl. Brit. Ind.* iv. 649.—Ind. or.
villosa, *Bonth. Hort. Beng.* [95]; *Fl. Ind.* iii. 4 = Pogostemon villosus.

ELSHOLZIA, Moench, Meth. 389 (1794) = prsec.

ELSHOTZIA, Roxb. Fl. Ind. iii. 4 (1832) = prsec.

ELSNERIA, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) (Pl. Meyen.) 346. t. 8 = **Bowlesia**, Ruiz & Pav. (Umbellif.).
crataegifolia, Walp. l. c. = B. multiradiata.

ELSOTA, Adans. Fam. ii. 358 (1763) = **Securidaca**, Linn. (Polygal.).

ELSSHOLZIA, Garcke, Fl. Deutschl. ed. VI. 307 (1863) = **Blaholtsia**, Willd. (Labiat.).

ELTROPLECTRIS, Rafin. Fl. Tellur. ii. 51 (1836) = **Pelexia**, Poit. (Orchid.).
acaenoides, Rafin. l. c. = P. setacea.

ELUTERIA, Steud. Nom. ed. I. 296 (1821) = ELUTHERIA, Linn. = **Croton**, Linn. (Euphorb.).

ELUTHERIA, P. Br. Hist. Jamaic. 369 (1756) = **Cascarilla**, Juss. (Meliac.).

ELUTHERIA, Linn. Hort. Cliff. 486, 487 (1737) = Croton, Linn. (Euphorb.).

ELUPHERIA, M. Roem. Syn. Hesper. 122 (1846). *MELIACEÆ*, Benth. & Hook. f. i. 389.
SCHMARDAEA, Karst. Fl. Columb. i. 187. t. 93 (1861?).
microphylla, *M. Roem. l. c.*; *C. DC. in DC. Monog. Phan.* i. 735.—Peruv.

ELVASIA, DC. in Ann. Mus. Par. xvii. (1811) 422. t. 20. *OCHNACEÆ*, Benth. & Hook. f. i. 318.
HOSTMANNIA, Planch. in Hook. Ic. Pl. t. 709 (1848).
calophylla, *DC. in Ann. Mus. Par.* xvii. (1811) 422. t. 20.—Bras.
casuarinoides, *Engl. in Mart. Fl. Bras.* xii. 354.—Am. trop.
Hostmannia, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 648.—Guiana.
quinquestylosa, *Sprunc, ex Engl. in Mart. Fl. Bras.* xii. 355.—Am. trop.

ELVINA, Steud. Nom. ed. II. i. 550 (1840) = suq.

ELVIRA, Cass. in Dict. Sc. Nat. xxx. 67 (1824). *COMPOSITÆ*, Benth. & Hook. f. ii. 343.
DELILIA, Spreng. Syst. iii. 337 (1826).
DESMOCEPHALIUM, Hook. f. in Proc. Linn. Soc. i. (1845) 277; et Trans. Linn. Soc. xx. 306 (1847).
MERATIA, Cass. in Dict. Sc. Nat. xxx. 65 (1824).
MICROCOCCIA, Hook. f. in Trans. Linn. Soc. xx. (1847) 309.
biflora, DC. Prod. v. 508 = Martynl.
Martyni, *Cass. in Dict. Sc. Nat.* xxx. 68.—Mexic.

ELWENDIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 140 = **Ctenum**, Linn. (Umbell.).
cornifera, Boiss. l. c. = C. Elwendia.

ELWERTIA, Rafin. Sylva Tellur. 103 (1838) = **Clusia**, Plum. (Guttifer.).
retusa, Rafin. l. c. = C. roseo.

ELYMUS, Linn. Hort. Upsal. 22 (1748); Amœn. Acad. iii. (1787) 20. *GRAMINEÆ*, Benth. & Hook. f. iii. 1206.
CRITHOPSIS, Jaub. & Spach, Illustr. iv. 80. t. 321 (1850).
LEYMUS, Hochst. in Flora, xxxi. (1848) 118, in adnot.
ORTHOSTACHYS, Ehrh. Beitr. iv. 146 (1789).
POLYANTHERIX, Nees, in Ann. Nat. Hist. Ser. I. i. (1838) 384.
SITANION, Rafin. in Journ. Phys. lxxxix. (1819) 103.
SITOSPELOS, Adans. Fam. ii. 36 (1763).
aegyptiacus, Spreng. Syst. i. 338 = Delileanus.
agropyroides, *Y. & C. Presl, Rel. Haenk.* i. 265.—Chili.
Alberti, *Regel, in Act. Hort. Petrop.* vii. (1880) 581.—Am. centr.
alaicus, Spreng. f. Tent. Suppl. 5 = junceus.
andinus, *Trin. in Linnæa*, x. (1836) 304.—Peruv.
angulatus, *Y. & C. Presl, Rel. Haenh.* i. 264.—Peruv.
angustatus, Steud. Nom. ed. II. i. 550, sphalm. = seq.
angustus, *Trin. ex Ledeb. Fl. Alt.* i. 119.—Sibir. antarcticus, *Hook. f. Fl. Antarct.* 388.—Reg. Magellan
arenarius, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) ii 285.—Reg. Casp.
arenarius, Georgi, Beschr. Russ. Reich. Nachtr. 357 = dasytachys.
arenarius, S. G. Gmel. Reise, iv. 44; ex Ledeb. Fl Ross. iv. 332. = giganteus.
arenarius, *Linn. Sp. Pl.* 83.—Europ.; Am. bor. occ.
arenarius, Mert. in Linnæa, v. (1830) 61 = mollis?
arenarius, Pall. Reise, i. 243 = sabulosus.
asper, *Nees, ex Steud. Syn. Pl. Gram.* 349.—Chili.
brachystachys, Walp. Ann. iii. 786 = Delileanus.
caducus, *Boiss. Fl. Orient.* v. 691.—Afghan.
canadensis, *Linn. Sp. Pl.* 83.—Am. bor.
caninus, Linn. Syst. Suc. 39 = Agropyron caninum.
caninus, Roth, Tent. Fl. Germ. i. 51 = Agropyron repens.
cappadocicus, *Boiss. & Bal. in Bull. Soc. Bot. Fr.* iv. (1857) 806.—Cappadoc.
Caput-Medusæ, Forsk. Fl. Aegypt. Arab. 25 = Cenchrus catharticus.
Caput-Medusæ, *Linn. Sp. Pl.* 84.—Europ. austr., Afr. bor.; Orient.
carolinianus, *Walt. Fl. Carol.* 82.—Am. bor.
chaetotus, *Phil. in Linnæa*, xxix. (1857–58) 104.—Chili.
cladostachys, *Turcz. in Bull. Soc. Nat. Mosc.* xxix. (1856) i. 64.—Sibir. or.
compositus, Steud. Syn. Pl. Gram. 348 = europæus.
condensatus, *Y. & C. Presl, Rel. Haenh.* i. 265.—Am. bor. occ.
cordalensis, *Phil. in Linnæa*, xxxiii. (1864–65) 308.—Chili.
crinitus, Schreb. Beschr. Gräs. 15. t. 24 = E. Caput-Medusæ.
dahuricus, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 105, nomen; et xxiii. i. (1857) 62.—Tauria.
dasytachys, *Trin. ex Ledeb. Fl. Alt.* i. 120.—Sibir. altaic.
Delileanus, *Schult. Mant.* ii. 494.—Syria; Aegypt.
desertorum, *Kar. & Kir. in Bull. Sc. Nat. Mosc.* (1841) 567.—Sibir. altaic.
dictaeus, *Dem. Opusc.* 97.—Hab.?
diven, *Y. & C. Presl, Rel. Haenh.* i. 265.—Staus Nootica.

ELYMUS:—
dumetorum, Hoffm. Deutschl. Fl. ed. II. i. t. 64 = Agropyron repens.
durus, Hort. ex Steud. Nom. ed. II. i. 550, 551 = virginicus.
europaeus, *Linn. Mant.* i. 35.—Europ.; Reg. Casc.
europaeus, Willd. Fl. Berol. Prod. 61 = arenarius.
excelsus, *Turcz. in Bull. Soc. Nat. Mosc.* xxix. (1856) i. 62.—As. bor.
flexuosus, *Tausch, in Flora*, xx. (1837) 120.—Hab. ?
fragilis, Boiss. Diagn. Ser. I. vii. 128 = Hordeum fragile.
Gayanus, *E. Desv. in C. Gay, Fl. Chil.* vi. 467.—Chili.
geniculatus, Vahl, *Symb. Bot.* iii. 10.—Kosa.; Sibir.
glaucifolius, Muhl. in Willd. Enum. Hort. Berol. 181 = canadensis.
glaucus, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 99 = sibiricus.
glaucus, *Regel, in Act. Hort. Petrop.* vii. (1880) 585.—Turkestan.
gracilis, *Phil. in Linnaea*, xxxiii. (1864–65) 801.—Chili.
hirsutus, *y. & C. Presl, Rel. Haenk.* i. 264.—Sinus Nootka.
hirsutus, Schreb. ex Roem. & Schult. Syst. ii 776 = villosus.
hordeiformis, Desf. Tabl. ed. I. 15, nomen; et Cat. Hort. Par. ed. III. 387 = striatus.
hyalanthus, *Rupr. Sert. Tianschi.* 36.—As. centr.
Hystrix, Linn. Sp. Pl. 560 = Asperella Hystrix.
intermedius, Bieb. Fl. Taur. Cauc. i. 82 = E. Caput-Medusae.
interruptus, *Buchl. in Proc. Acad. Soc. Philad.* 1862 (1863) 99.—Am. bor.
jemisensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxix. (1856) i. 64.—Sibir.
jubatus, Link, Hort. Berol. i. 19 = Hordeum jubatum.
juncus, *Fisch. in Mém. Soc. Nat. Mosc.* i. (1811) 25. t. 4.—Rosa.; As. bor.
Karelini, *Turcz. in Bull. Soc. Nat. Mosc.* xxix. (1856) i. 64.—Turkestan.
lanuginosus, *Trin. ex Ledeb. Fl. Alt.* i. 121.—Sibir. altaic.
latiglumis, *Phil. in Linnaea*, xxxiii. (1864–65) 802.—Chili.
Lechleri, Steud. Syn. Pl. Gram. 480 = Hordeum jubatum.
leucostachys, Everam. ex Griseb. in Ledeb. Fl. Ross. iv. 338 = dasystachys.
litoralis, Turcz. ex Griseb. l. c. = dasystachys.
Macounii, *Vasey, Gr. U. St.* 46, nomen.—Am. bor.
macrostachys, Spreng. in Schrad. Journ. (1800) ii. 196 = gigantens.
mexicanus, Cav. Desc. 314 = sabulosus.
mollis, R. Br. ex Richards. in Frankl. Narr. 1st Journ. 723 = villosus.
mollis, *Trin. in Spreng. Neue Entdeck.* ii. 72.—As. bor. or.; Am. bor. occ.
monococcus, *Linn. Syst. ed.* X. 879.—Hab. ?
montanus, Pall. ex Georgi, Beschreib. Russ. Reich. Nachtr. 258 = E. Caput-Medusae.
multicaulis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 868.—Sibir. altaic.
muticus, *Phil. in Linnaea*, xxxiii. (1864–65) 800.—Chili.
neogenes, *Steud. Syn. Pl. Gram.* 349.—' Terra-nova.'
nutans, *Griseb. in Gott. Nachr.* (1868) 72.—Reg. Himal.
obtusus, Kunth, Enum. Pl. i. 556 = ovatus.
Orcuttianus, *Vasey, in Coult. Bot. Gaz.* x. (1885) 258.—Calif.
ovatus, *Trin. ex Ledeb. Fl. Alt.* i. 121.—Sibir. altaic.
paposanus, *Phil. Fl. Atac.* 58.—Chili.
parciflorus, *Law. Tabl. Encyc.* i. 207.—Hab. ?
philadelphicus, Linn. Cent. Pl. I. 6; Amoen. Acad. iv. 296 = canadensis.
pilifer, [Soland. in] Russell, Aleppo ed. II. ii. 244 = Agropyron orientale.
plantagineus, Link, Hort. Berol. i. 18 = E. Caput-Medusae.
praetervisus, *Steud. Syn. Pl. Gram.* 348.—*Reg.* Ural.
pratensis, *Phil. in Linnaea*, xxxiii. (1864–65) 801.—Chili.
procipuus, *Prunen. Ind. Sem. Hort. Franc.* (1826); *ex Steud. Syn. Pl. Gram.* 349.—Am. bor.

ELYMUS:—
Pseudoagrogyrum, Trin. ex Turcz. in Bull. Soc. Nat. Mosc. (1858) 106; (1856) i. 63 = Triticum Pseudo-agropyrum.
Pseudohystrix, Schult. Mant. ii. 427 = Asperella Hystrix.
racemosus, Lam. Tabl. Encyc. i. 207 = sabulosus.
racemosus, Poir. Encyc. Suppl. ii. 546 = sibiricus.
ramosus, Desf. Tabl. ed. I. 15, nomen; Cat. Hort. Par. ed. III. 387 = sibiricus.
rhachitrichus, Hochst. ex Jaub. & Spach, Illustr. iv. 80 = Delileanus.
rigescens, *Trin. in Linnaea*, x. (1836) 304.—Peruv.
sabulosus, *Rieb. Pl. Taur. Cauc.* i. 81.—Reg. Caucas.
salsuginosus, Turcz. ex Ledeb. Fl. Ross. iv. 333 = dasystachys.
Saunderesii, *Vasey, in Bull. Torrey Bot. Club*, xi. (1884) 136.—Am. bor. occ.
secaliformis, Trin. ex Steud. Nom. ed. II. i. 551 = Hordeum fragile, *Boiss.*
secalinus, Claus, in Goebel, Reise, ii. 319; et ex Ledeb. Fl. Ross. iv. 330 = praec ?
sibiricus, *Linn. Sp. Pl.* 83.—As. bor.; Am. bor.
Sitanion, *Schult. Mant.* ii. 426.—Am. bor.
striatus, *Besa. Enum. Pl. Volkyn.* 50 (quid ?).—Volhynia.
striatus, *Willd. Sp. Pl.* i. 470.—Am. bor.
subpaniculatus, *Steud. Syn. Pl. Gram.* 350.—Austral.
subulatus, Forsk. Fl. Aegypt. Arab. 26 = Delileanus ?
tener, Linn. f. Suppl. 114 = sibiricus.
triticoides, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 99 = condensatus.
vaginatus, *Phil. in Linnaea*, xxxiii. (1864–65) 800.—Chili.
Veldiviae, *Steud. Syn. Pl. Gram.* 349.—Chili.
villosus, *Muhl. ex Willd. Enum. Hort. Berol.* 131.—Am. bor.
virginicus, *Linn. Sp. Pl.* 84.—Am. bor.

ELYMUS, Mitchell, in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) 210, App. = Eisanis, Linn. (Gramin.).

ELYNA, Schrad. Fl. Germ. i. 155 (1806) = **Kobresia**, Willd. (Cyperac.).
Bellardi, Willd. in Vet. Akad. Handl. Stockh. (1814) 107, et (1818) 154 = K. caricina.
Bellardi, C. Koch, in Linnaea, xxi. (1848) 616 = K. scirpina.
capillifolia, Decne. in Jacquem. Voy. Bot. 178. t. 174 = K. capillifolia.
capillifolia, Henderson, Yarkand, 339 = Carex stenophylla.
caricina, Mertens & Koch, Deutschl. Fl. i. 458 = K. caricina.
filifolia, Turcz. in Bull. Soc. Nat. Mosc. xxviii. (1855) i. 353 = K. filifolia.
filiformis, Steud. Syn. Pl. Cyp. 245.—Am. bor.
humilis, C. A. Mey. ex Trautv. in Act. Hort. Petrop. i. (1871) 21 = K. schoenoides.
hokanica, Regel, in Act. Hort. Petrop. vii. (1880) 563 (Kobresiae sp.).—Turkestan.
laxa, Kunth, Enum. Pl. ii. 534 = K. laxa.
schoenoides, C. A. Mey. in Ledeb. Fl. Alt. iv. 235 = K. schoenoides.
sibirica, Turcz. in Bull. Soc. Nat. Mosc. (1838) 103; Ledeb. Fl. Ross. iv. 262 = K. schoenoides.
spicata, Boiss. Fl. Orient. v. 394 = K. capillifolia.
spicata, Schrad. Fl. Germ. i. 155 = K. scirpina.
stenocarpa, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 536 = K. Royleana.
stricta, Hoppe, ex Steud. Nom. ed. II. i. 551 = K. scirpina.

ELYNANTHUS, Beauv. ex Lestib. Essai Fam. Cyp. 23 (1819). *GENUS DUBIUM* (Cyperac.).—Cf. Benth. & Hook. f. iii. 1064.

ELYNANTHUS, Nees, in Linnaea, vii. (1832) 530; ix. (1835) 296 = **Tetraria**, Beauv. (Cyperac.).
anceps, Nees, in Linnaea, ix. (1834) 298, nomen.—Afr. austr.
aphyllus, F. Muell. ex Boeck. in Linnaea, xxxviii. (1874) 311 = Schoenus aphyllus.
arenarius, Nees, in Linnaea, ix. (1834) 298 = T. compar.
aristatus, Boeck. l. c. xxxviii. (1874) 260 (Tetrariae sp.).—Afr. austr.
auritus, Nees, in Linnaea, ix. (1834) 296 (Tetrariae sp.) = *cylindricus*.

ELYNANTHUS:—
australis, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 48 = Cladium elynanthoides.
bifidus, Nees, l. c. = Schoenus bifidus.
borbonicus, Steud. Syn. Pl. Cyp. 155 (Carphae sp.).—Ins. Borbon.
capillaceus, Benth. Fl. Austral. vii. 377 (Cladii sp.).—Austral.
capitatus, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 48 = Schoenus capitatus.
compar, Boeck. in Linnaea, xxxviii. (1874) 256, partim = *cylindricus*.
compar, Nees, l. c. vii. (1832) 530 = T. compar.
cuspidatus, Nees, l. c. ix. (1834) 298; x. (1836) 188 (Tetrariae sp.).—Afr. austr.
Dregeanus, Boeck. l. c. xxxviii. (1874) 262 (Tetrariae sp.).—Afr. austr.
ensifolius, Boeck. l. c. 264 = Cladium ensifolium.
gracilis, Nees, l. c. x. (1835–36) 189 = *cuspidatus*.
grandiflorus, Nees, in Lehm. Pl. Preiss. ii. 78 = Schoenus grandiflorus.
Kraussii, Hochst. in Flora, xxviii. (1845) 761 = Macrochaetium Dregei.
ligulatus, Boeck. in Linnaea, xxxviii. (1874) 262 (Tetrariae sp.).—Afr. austr.
loreus, Nees, l. c. ix. (1834) 298 (Tetrariae sp.) = *cuspidatus*.
microstachyus, Boeck. l. c. xxxviii. (1874) 261 = *cuspidatus*.
obtusifolius, Nees, in Lehm. Pl. Preiss. ii. 78 = Schoenus obtusifolius.
octandrus, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 48 (alieni gen.).—Austral.
pictus, Boeck. in Linnaea, xxxviii. (1874) 254 (Tetrariae sp.).—Afr. austr.
revolutus, Nees, in Lehm Pl. Preiss. ii. 78 = *octandrus*.
sculptus, Nees, l. c. 79 = Schoenus sculptus.
spathaceus, Nees, in Linnaea, x. (1835–36) 186 = Schoenus nigricans
sylvaticus, Nees, l. c. ix. (1834) 298; x. (1835–36) 188 (Tetrariae sp.).—Afr. austr.
ustulatus, Nees, l. c. x. (1835–36) 198 (Tetrariae sp.).—Afr. austr.
viscosus, Nees, l. c. 186 = T. compar.

ELYONORUS, Bartl. Ord. 31 (1830) = seq.

ELYONURUS, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 941 (1806) = **Elionurus**, Kunth (Gramin.).
tripsacoides, Humb. & Bonpl. l. c. = Elionurus tripsacoides.

ELYTHRANTHE, Reichb. Nom. 78 (1841) = **Loranthus**, Linn.

ELYTHRARIA, D. Dietr. Syn. i. 80, 64 (1839) = **Elytraria**, Michx. (Acanth.).

ELYTHROPHORUS, Dum. Anal. Fam. 68 (1829) = **Elytrophorus**, Beauv (Gramin.).

ELYTHROSPERMUM, Steud. Nom. ed. II. i. 552 (1840) = ELYTROSPERMUM, C. A. Mey. = **Scirpus**, Linn. (Cyperac.).

ELYTHROSTAMNA, Boj. Hort. Maurit. 230 (1837) = **Ipomoea**, Linn. (Convolv.).
convolvulacea, Boj. l. c. = I. pileata.

ELYTRANTHE, Blume, in Schult. f. Syst. vii. 1730 (1830) = **Loranthus**, Linn.
albida, Blume, l. c. = L. albidus.
ampullacea, G. Don, Gen. Syst. iii. 425 = L. ampullaceus.
Arnottiana, Miq. Fl. Ind. Bat. I. i. 882 = L. Arnottianus.
avenis, G. Don, Gen. Syst. iii. 426 = L. avenis.
carinatula, G. Don, l. c. = L. ampullaceus.
cochinchinensis, G. Don, l. c. = L. cochinchinensis.
cupulata, G. Don, l. c. 427 = L. cupulatus.
farinosa, G. Don, l. c. = L. pentandrus.
formosa, G. Don, l. c. 426 = L. formosus.
gemmiflora, G. Don, l. c. 427 = L. formosus.
globosa, Blume, in Schult. f. Syst. vii. 1730 = L. globosus.
involucrata, G. Don, Gen. Syst. iii. 427 = L. involucratus.
lonicroides, G. Don, l. c. 427 = L. lonicioides.

ELYTRANTHE :—
obmunita, G. Don, Gen. Syst. iii. 428 = Loranthus obmunitus.
sinuata, G. Don, l. c. 426 = L. globosus.
pallens, G. Don, l. c. = L. ampullaceus.
patula, G. Don, l. c. = L. patulus.
retusa, G. Don, l. c. 425 = L. retusus.
rigida, G. Don, l. c. 427 = L. pentandrus.
sphaerocarpa, G. Don, l. c. 426 = L. globosus.
subglobosa, G. Don, l. c. = L. globosus.
subumbellata, G. Don, l. c. 427 = L. subumbellatus.
tetragona, G. Don, l. c. = L. tetragonus.
umbellata, G. Don, l. c. = L. Scurrula.
viridiflora, G. Don, l. c. 426 = L. globosus.

ELYTRARIA, Michx. Fl. Bor. Am. i. 8 (1803); Vahl, Enum. i. 106 (1805). *ACANTHACEAE*, Benth. & Hook. f. ii. 1073.
amara, Blanco, Fl. Filip. ed. II. 3 = crenata.
asperifolia, Nees, in DC. Prod. xi. 65 = tridentata.
bromoides, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 116.—Mexic.
caroliniensis, Pers. Syn. i. 23 = virgata.
caulescens, Ledeb. Ind. Sem. Hort. Dorp. (1820) = tridentata.
crenata, *Vahl, Enum.* i. 106.—Reg. trop.
cupressina, Nees, in DC. Prod. xi. 65 = virgata.
fasciculata, H. B. & K. Nov. Gen. et Sp. ii. 235 = tridentata.
frondosa, H. B. & K. l. c. 116, 234 = tridentata.
herbacea, Michx. ex Nees, in DC. Prod. xi. 63 = virgata.
imbricata, Pers. Syn. i. 23 = tridentata.
indica, Pers. l. c. = crenata.
lyrata, Vahl, Enum. i. 106 = crenata.
marginata, Vahl, Enum. i. 108 = crenata.
microstachya, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 114 = tridentata.
pachystachya, Oerst. l. c. 116. t. 3 = tridentata.
ramosa, H. B. & K. Nov. Gen. et Sp. ii. 235 = tridentata.
scorpioides, *Roem. & Schult. Syst. Mant.* i. 128.—Ins. St. Martha.
tridentata, *Vahl. Enum.* i. 107.—Reg. trop.
Vahliana, Michx. Fl. Bor. Am. i. 9. t. 1 = virgata.
virgata, *Michx. l. c.*—Am. trop.
virgata, Nees, in DC. Prod. xi. 63 = crenata.

ELYTRIGIA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190 = **Agropyron**, J. Gaertn. (Gramin.).
repens, Desv. l. c. = A. repens.

ELYTRIGIUM, Benth. in Journ. Linn. Soc. xix. 130 (1881) = prsec.

ELYTROPAPPUS, Cass. in Bull. Soc. Philom. (1816) 199. *COMPOSITAE*, Benth. & Hook. f. ii. 338.
ACHYROCOMA, Schrank, in Denkschr. Akad. Munenh. viii. (1821-22) 147, 170.
CYATHOPAPPUS, Sch. Bip. in Pollichia, xviii-xix. (1861) 153.
adpressus, *Hemsl. in Hara. & Sond. Fl. Cap.* iii. 274.—Afr. austr.
ambiguus, *DC. Prod.* vi. 256.—Afr. austr.
canescens, *DC. l. c.*—Afr. austr.
cinerascens, DC. l. c. 281 = glandulosus.
cyathiformis, *DC. l. c.* i. 287.—Afr. austr.
glandulosus, *Less. Syn. Comp.* 348.—Afr. austr.
Rhinocerotis, *Less. l. c.* 344.—Afr. austr.
spinellosus, *Cass. in Bull. Soc. Philom.* (1816) 199.—Afr. austr.
spinulosus, Less. Syn. Comp. 343 = spinellosus.

ELYTROPHORUM, Folr. in Dict. Sc. Nat. xiv. 377 (1819) = seq.

ELYTROPHORUS, Beauv. Agrost. 67. t. 14. f. 3 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1183.
ECHINALYSIUM, Trin. Fund. Agrost. 142 (1820).
articulatus, *Beauv. l. c.*—Ind. or.

ELYTROPUS, Muell. Arg. in Bot. Zeit. xviii. (1860) 21. t. 1. *APOCYNACEAE*, Benth. & Hook. f. ii. 726.
chilensis, *Muell. Arg. l. c.* 22; *et in Linnaea*, xxx. (1859-60) 440.—Reg. Argent. Chili.

ELYTROPUS:—
heterophyllus, Miers, Apocyn. S. Am. 116 = Echites chilensis.
Pturmica, Miers, l. c. 115 = Echites Pturmica.
subsecurus, Miers, l. c. 114 = Echites chilensis.
spectabilis, Miers, l. c. 116 = Echites spectabilis.

ELYTROSPERMUM, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1830) 300. t. 3 = Betlpsus, Linn. (Cyperac.).
californicum, C. A. Mey. l. c. = S. riparius.

ELYTROSTAMNA, Cholsy, in DC. Prod. ix. 265 (1845) = ELYTHROSTAMNA, Boj. = Ipomoea, Linn. (Convolv.).

EMBAMMA, Griff. Notul. iv. 694 (1854) = Pteristanthes, Blume (Ampelid.).
caudigerum, Griff. l. c. 695 = P. cissoides.
heteranthum, Griff. l. c. 694 = P. heteranthe.

EMBELIA, Burm. f. Fl. Ind. 62. t. 63 (1768). *MYRSINEAE*, Benth. & Hook. f. ii. 644.
CHORIPETALUM, A. DC. in Trans. Linn. Soc. xvii. (1834) 181.
DAMERIA, Endl. Gen. 1444 (1841).
DAUCKIA, Dennst. Schlüssl. Hort. Malab. 31 (1818).
GHEZAEMBILLA, Adans. Fam. ii. 449 (1763).
PATTARA, Adans. l. c. 447 (1763).
PLOTIA, Neck. Elem. ii. 58 (1790).
RIBESIOIDES, Linn. Fl. Zeyl. 190 (1747).
SAMARA, Linn. Mant. 144, excl. syn. 199 (1771).
abyssinica, *Baker, in Oliver, Fl. Trop. Afr.* iii. 497.—Afr. trop.
adnata, *Bedd. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 514.—Ind. or.
amentacea, *C. B. Clarke, l. c.*—Malacca.
angustifolia, *A. DC. in DC. Prod.* viii. 84.—Ins. Borbon.
arbores, *A. DC. l. c.* 85.—Madag.
Bassei, A. DC. in Trans. Linn. Soc. xvii. (1834) 181 = robusta.
borneensis, *Scheff. Comm. Myrsin.* 48.—Borneo.
bracteosa, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 366.—Malaya.
Burmanni, Retz. Obs. iv. 28 = E. Ribes et Salvadora persica.
caesecana, *Zoch, ex Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 299.—Malaya.
clusiaefolia, *Miq. Fl. Sumgh.* 187.—Java.
concinna, *Baker, in Journ. Linn. Soc.* xx. (1883) 199.—Madag.
coriacea, *Wall. Cat.* n. 3214; *A. DC. in Trans. Linn. Soc.* xvii. (1834) 185.—Malaya.
cyrtobotrya, *Miq. Fl. Ind. Bat.* ii. 1013.—Java.
dasythyrsa, *Miq. l. c. Suppl.* 573.—Ins. Banca.
dentata, Buch.-Ham. ex Wall. Cat. sub n. 2304 = E. Ribes.
esculenta, D. Don, Prod. Fl. Nep. 147 = floribunda.
ferruginea, Wall. Cat. n. 2310 = robusta.
floribunda, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 291.—Reg. Himal.; Malaya.
Gamblei, *Kurz, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 516.—Reg. Himal.
garcinioides, Miq. Fl. Jungh. 187 = E. Ribes.
Gardneriana, *Wight, Ic.* t. 1308.—Ind. or.
glandulifera, Wight, l. c. t. 1307 = E. Ribes.
Grossularia, Retz. Obs. iv. 24 = Salvadora persica.
guineensis, *Baker, in Oliver, Fl. Trop. Afr.* iii. 496.—Afr. trop.
indica, J. F. Gmel. Syst. 320 = E. Ribes.
javanica, *A. DC. in DC. Prod.* viii. 84.—Java.
Jussiaei, *A. DC. l. c.*—Madag.
Kraussii, *Herv. Thes. Cap.* ii. 17. t. 197.—Afr. austr.
Lampasi, *Scheff. Comm. Myrsin.* 37.—Malaya.
lucida, *Wall. Cat.* n. 3215.—Singapore.
macrophylla, Blume, ex Scheff. Comm. Myrsin. 36 = coriacea.
madagascariensis, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 81.—Madag.
micrantha, *A. DC. l. c.*—Ins. Maurit.
microcalyx, *Kurz, in Journ. Bot.* xiii. (1875) 236.—Ins. Nicobar.
Myrsillea, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 67.—Burma; Malaya.
Nagushia, *D. Don, Prod. Fl. Nep.* 147.—Reg. Himal.
nervosa, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 81.—Cochinch.
nilotica, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 106. t. 73.—Afr. trop.
nummularisfolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 198.—Madag.
ostans, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 290.—Reg. Himal.; Burma.
oblongata, Miq. Fl. Ind. Bat. Suppl. 573 = coriacea.
ovata, *Scheff. Comm. Myrsin.* 41.—Sumatra.
parvifora, *Wall. Cat.* n. 2307.—Reg. Himal.; Malaya peninsula.
pergamacea, *A. DC. in DC. Prod.* viii. 85.—Java.
philippinensis, *A. DC. l. c.* 83.—Ins. Philipp.
picta, A. DC. in Trans. Linn. Soc. xvii. (1834) 180 = robusta.
racemosa, *Hassk. ex Scheff. Comm. Myrsin.* 36.—Java.
ramosa, *Wall. Cat.* n. 6522 (Urticeae ?).—Ind. or.
reticulata, Wall. l. c. 6521 = robusta.
Ribes, *Burm. f. Fl. Ind.* 62. t. 23.—As. trop.
robusta, *Roxb. Hort. Beng.* 16; *Fl. Ind.* i. 587.—Ind. or.
sarmentosa, *Baker, in Journ. Linn. Soc.* xx. (1883) 198.—Madag.
Schimperi, *Vatke, in Linnaea*, xl. (1876) 206.—Abyss.
sessilifora, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 66.—Burma.
singalangensis, *Scheff. Comm. Myrsin.* 37.—Sumatra.
sumatrana, Miq. Fl. Jungh. 185 = E. Ribes.
Tajeran-Cottam, A. DC. in Trans. Linn. Soc. xvii. (1834) 181 = robusta.
umbellulata, Wall. Cat. n. 2312 = Myrsine capitellata.
wrophylla, Wall. l. c. n. 3209 = Parastemon urophyllum
westiia, Kurz, For. Fl. Brit. Burma, ii. 108 = nutans.
vestita, *Roxb. Fl. Ind. ed. Carey*, ii. 288.—Reg. Himal.
villosa, *Baker, in Journ. Linn. Soc.* xx. (1883) 199.—Madag.
villosa, A. DC. in Trans. Linn. Soc. xvii. (1834) 180 = robusta ?
villosa, Wall. Cat. n. 2313, ex parte = Lampani.
villosa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 289 = robusta.
viridiflora, *Scheff. Comm. Myrsin.* 48.—Ind. or.; Malaya.

EMBELICA. Boj. Hort. Maurit. 279 (1837) = seq.

EMBLICA, Gaertn. Fruct. ii. 122. t. 108 (1791) = Phyllanthus, Linn. (Euphorb.).
annua, Rafin. Sylva Tellur. 91 = Agynsia baccifermis.
arborea, Rafin. l. c. = P. Emblica.
grandis, Gaertn. Fruct. ii. 123 = Agynsia baccifermis.
officinalis, Gaertn. l. c. 122. t. 108. f. 2 = P. Emblica.
Pulasia, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 507 = Cleistanthus collinus.
pilosa, Spreng. Syst. iii. 20 = P. pilosus.
piscfornia, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 506.—Ind. or.
racemosa, Spreng. Syst. iii. 20 = Agynsia baccifermis.

EMBLINGIA, F. Muell. Fragm. ii. 2. t. 11 (1860). *CAPPARIDEAE*, [Benth. & Hook. f. i. 868].
calcicolifora, *F. Muell. l. c.* ii. 3.—Austral.

EMBOTHRIUM, Forst. Char. Gen. 15. t. 8, partim. (1776). *PROTEACEAE*, Benth. & Hook. f. iii. 153.
CATAS, Dondi ex Lam. Encyc. ii. 354 (1785).
ORREOCALLIS, R. Br. in Trans. Linn. Soc. x. (1810) 196.
alnifolium, Poepp. ex Meisn. in DC. Prod. xiv. 450 = Lomatia obliqua.
angustifolium, Willd. Sp. Pl. i. 589, in syn. = Grevillea linearis.
bipinnatum, Willd. ex Roem. & Schult. Syst. i. 583 = Lomatia ferruginea.
buxifolium, Sm. Icones, Bot. N. Holl. 29. t. 10 = Grevillea buxifolia.
Chapparo, Humb. in Magaz Ausein.—Cf. R. Br. in Trans. Linn. Soc. x. (1810) 221 = Roupala complicata.
cinereum, Thb. ex Meisn. in DC. Prod. xiv. 385, 444 = Grevillea setosa.
coccineum, *Forst. Char. Gen.* 16. t. 8.—Reg. Magell.
cordifolium, Willd. ex Roem. & Schult. Syst. i. 583 = Roupala cordifolia.
crithmifolium, Sieud. Nass. ed. II. i. 282 = Lomatia silaifolia.

EMBOTHRIUM:—
curvatum, Willd. ex Roem. & Schult. Syst. i. 538 = Roupala complicata.
cytisoides, Cav. Ic. iv. 60 = Grevillea linearis, sericea.
dentatum, Ruiz & Pav. Fl. Per. i. 62. t. 94 = Lomatia dentata.
emarginatum, Ruiz & Pav. l. c. t. 95 = grandiflorum.
ferrugineum, Cav. Ic. iv. 59 = Lomatia ferruginea.
fraxinifolium, Cav. l. c. 60 = Grevillea buxifolia.
Gilliesii, *Meisn. in DC. Prod.* xiv. 444.—Reg. Argent.
grandiflorum, Bonpl. ex Meisn. l. c. 446 (= *Oreocallis Ruizii*).—Peruv.
grandiflorum, *Lam. Encyc.* ii. 354.—Ecuador; Peruv.
herbacrum, Cav. Ic. iv. 58 = Lomatia silaifolia.
hirsutum, Lam. Encyc. ii. 355 = Lomatia obliqua.
ilicifolium, Poir. Encyc. Suppl. ii. 551 = Lomatia illicifolia.
lanceolatum, *Ruiz & Pav. Fl. Per.* i. 62. t. 96.—Chili.
linare, Andr. Bot. Rep. t. 272 = Grevillea linearis.
linearifolium, Cav. Ic. iv. 59 = Grevillea linearis.
longifolium, Poir. Encyc. Suppl. ii. 551 = Lomatia longifolia.
mayus, Willd. Sp. Pl. i. 539, in syn. = Grevillea sp.?
minus, Willd. l. c. = Grevillea punicea.
monospermum, Ruiz & Pav. Fl. Per. i. 63 = Roupala peruviana.
macronatum, Willd. ex Roem. & Schult. Syst. i. 538 = grandiflorum.
myricoides, Gaertn. f. Fruct. iii. 215. t. 218 = Lomatia longifolia.
myrtifolium, Poepp. ex Meisn. in DC. Prod. xiv. 425, 444 = Roupala myrtoidea.
obliquum, Ruiz & Pav. Fl. Per. i. 63. t. 97 = Roupala diversifolia.
pinnatum, Ruiz & Pav. l. c. = Roupala diversifolia.
Pinal, Domb. ex Meisn. in DC. Prod. xiv. 449, 444 = Lomatia dentata.
polystachyum, Willd. ex Roem. & Schult. Syst. i. 538 (= *Andrieptatium polystachyum*).—N. Granat.
rubricaule, Giord. Cas. (1837).—Cf. Meisn. in DC. Prod. xiv. 451 = Stenocarpus saligna.
rursale, Domb. ex Meisn. in DC. Prod. xiv. 450, 444 = Lomatia obliqua.
salicifolium, Veni. Jard. Cels, 8 = Hakea saligna.
salignum, Andr. Rep. Bot. t. 215 = Hakea saligna.
sericeum, Sm. Spec. Bot. N. Holl. 25. t. 9 = Grevillea sericea.
silaifolium, Sm. l. c. 28. t. 8 = Lomatia silaifolia.
spathulatum, Cav. Ic. iv. 60 = Telopea speciosissima.
speciosissimum, Sm. Spec. Bot. N. Holl. i. 19. t. 7 = Telopea speciosissima.
speciosum, Salisb. Parad. Lond. t. 111 = Telopea speciosissima.
strobilinum, Labill. Nov. Holl. Pl. ii. 116. t. 265 = Knightia strobilina.
sylvaticum, Poepp. ex Meisn. in DC. Prod. xiv. 449 = Lomatia dentata.
tinctorium, Labill. Nov. Holl. Pl. i. 81. t. 48 = Lomatia tinctoria.
truncatum, Labill. l. c. 22. t. 44 = Telopea truncata.
umbellatum, Lam. f. Suppl. 128 = seq.
umbelliferum, Forst. Char. 16 = Stenocarpus Forsteri.
Wickhami, *Hill & F. Muell. Fragm.* viii. 164.—Austral.

EMBOTRIUM, Ruiz & Pav. Prod. 15 (1794) = prec.

EMBRYOGONIA, Blume, Mus. Bot. Lugd. Bat. ii. 192, 1852 [1856]= Combretum, Linn.
acuminata, Blume, l. c. 193 = C. acuminatum.
arborea, Teijsm. & Binn. in journ. Bot. Néerl. i. (1861) 366 = Terminalia Chebula.
latifolia, Blume, Mus. Bot. Lugd. Bat. ii. 193 = C. latifolium.
lucida, Blume, l. c. = C. trifoliatum.

EMBRYOPTERIS, Gaertn. Fruct. i. 165. t. 39 (1786) = Diospyros, Linn. (Ebenac.).
discolor, G. Don, Gen. Syst. iv. 41 = D. discolor.
glutinifera, G. Don, l. c. = D. Embryopteris.
glutinifera, Roxb. Pl. Corom. i. 49 = D. Embryopteris.
glutinifolia, Hort. ex Link, Enum. Hort. Berol. ii. 433 = pruna, ?
Kaki, G. Don, Gen. Syst. iv. 41 = D. Kaki.
lanceolata, G. Don, l. c. = D. Toposia.
Lourieboiana, G. Don, l. c. = D. Sideroxylon.
peregrina, Gaertn. Fruct. i. 165 = D. Embryopteris.
racemosa, G. Don, Gen. Syst. iv. 41 = D. Toposia.

EMELIA, Wight, Ic. iii. 11 (1846), sphalm. = Emilia, Cass. (Compos.).

EMELISTA, Rafin. Sylva Tellur. 127 (1838) = Cassia, Linn. (Legumin.).
obtusifolia, Rafin. l. c. = C. Tora.

EMBORHIZA, Pohl, in Flora, viii. (1825) 183, nomen.
RUBIACEAE, Benth. & Hook. f. ii. 146.
ENDLICHERA, Presl, Symb. Bot. i. 78. t. 49 (1832). brasiliensis, *Walp. Rep.* ii. 409.—Am. trop.
Pohliana, *Presl, Bot. Bemerk.* 86.—Bras.

EMERICIA, Roem. & Schult. Syst. iv. 88, 401 (1819) = Vallaris, Burm. f. (Apocyn.).
divaricata, Roem. & Schult. l. c. 401 = Strophanthes divaricatus.
Perrgularia, Roem. & Schult. l. c. = V. Pergulana.
sinensis, Roem. & Schult. l. c. 402 = V. sinensis.

EMERUS, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = Coronilla, Linn. (Legumin.).
aculeatus, Hornem. Hort. Hafn. 696 = Sesbania aculeata.
alpestris, Scheele, in Flora, xxvi. (1843) 448 = C. Emerus.
Caesalpina, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 368 = C. Emerus.
fruticosa, Hornem. Hort. Hafn. 695 = C. Emerus.
grandiflorus, Hornem. l. c. = Sesbania grandiflora.
herbaceus, Mill. Gard. Dict. ed. VIII. n. 3 = Sesbania occidentalis.
horizeus, Desv. in Mém. Soc. Linn. Par. iv. (1826) 299 = C. Emerus.
major, Mill. l. c. n. 1; Moench, Meth. 157 = C. Emerus.
minor, Mill. l. c. n. 2 = C. Emerus.
pubescens, Schum. & Thonn. Beskr. Guin. Pl. 354 = Sesbania pubescens.
Sesban, Hornem. Hort. Hafn. 696 = Sesbania aegyptiaca.

EMEX, Neck. Elem. Bot. ii. 214 (1790). *POLYGONACEAE*, Benth. & Hook. f. iii. 101.
CENTOPODIUM, Burch. Trav. i. 340 (1824).
NEO, Stend. Nom. ed. I. 553 (1821).
PODOCENTRUM, Burch. ex Meisn. in Linnaea, xiv. (1840) 489.
VIXO, Medic. Phil. Bot. i. 178 (1789).
acuta, Rafin. Fl. Tellur. iii. 46 = Rumex acutus.
australis, *Steinh. in Ann. Sc. Nat.* Sér. II. ix. (1838) 198. t. 7.—Austral.
Centropodium, Meisn. in Linnaea, xiv. (1840) 490 = australis.
spinosa, *Campd.* Fl. t. 1. f. 1.—Europ.; Afr. bor.

EMILIA, Cass. in Bull. Soc. Philom. (1817) 68; et in Dict. Sc. Nat. xiv. 405 (1819). *COMPOSITAE*, Benth. & Hook. f. ii. 445.
adscendens, *DC. Prod.* vi. 308.—Madag.
amplexicaulis, *Baker, in Journ. Linn. Soc.* xx. (1883) 190.—Madag.
angulata, *DC. Prod.* vi. 308.—Arab.
angustifolia, DC. l. c. = praemorsa.
casapitosa, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 100.—Afr. trop.
citrina, *DC. Prod.* vi. 308.—Madag.
coccinea, Sweet, Hort. Brit. ed. III. 332 = sagittata.
flacida, *Mig. Fl. Ind. Bat.* ii. 102.—Sumatra.
flammea, *Cass. in Dict. Sc. Nat.* xiv. 406.—Ind. or.; Ins. Philipp.
gracilis, *DC. Prod.* vi. 308.—Afr. trop.
graminea, *DC. l. c.*—Madag.
humifusa, *DC. l. c.* 302.—Afr. trop.
mucronata, Wall. Cat. n. 316 = sonchifolia.
prenanthoides, *DC. Prod.* vi. 308.—Reg. Himal.
pteranthoides, Thw. Enum. Pl. Zeyl. 167 = zeylanica, Walkeri.
pumila, *DC. Prod.* vi. 302.—Afr. trop.
purpurea, Cass. in Dict. Sc. Nat. xxiv. 292 = sonchifolia.
rigidula, DC. Prod. vi. 302 = sonchifolia.
sagittata, DC. l. c. = flammea.
scabra, DC. l. c. 308 = sonchifolia.
sinica, Miq. in Journ. Bot. Néerl. i. (1861) 105 = sonchifolia.
sonchifolia, Benth. in Hook. Niger Fl. 429 = flammea.
sonchifolia, *DC. Prod.* vi. 302.—As. et Afr. trop.
Walkeri, *Hook. f. Fl. Brit. Ind.* iii. 387.—Zeylan.
zeylanica, *C. B. Clarke, Comp. Ind.* 178.—Zeylan.

EMINIUM, Schott, Syn. Aroid. 16 (1856) = Heliocophyllum, Schott (Aroid.).
Ledebouri, Schott, l. c. 17 = H. crassifolium.

EMMENANTHE, Benth. in Trans. Linn. Soc. xvii. (1836) 281. *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 830.
MILTITZIA, A. DC. in DC. Prod. ix. 296 (1845).
glaberrima, *Torr. ex S. Wats. in Bot. King's Exp.* 257.—Am. bor. occ.
glandulifera, *Torr. ex S. Wats. l. c.*—Am. bor. occ.
lutea, *A. Gray, in Proc. Am. Acad.* x. (1875) 329.—Am. bor. occ.
parviflora, *A. Gray, in Pacif. Rail. Rep.* vi. 84. t. 15.—Am. bor. occ.
pendulifora, *Benth. in Trans. Linn. Soc.* xvii. (1835) 281.—Calif.
pusilla, *A. Gray, in Proc. Am. Acad.* xi. (1876) 87.—Am. bor. occ.

EMMENANTHUS, Hook. & Arn. Bot. Beech. Voy. 217 (1836) = Ixonanthes, Jack (Linear).
chinensis, Hook. & Arn. l. c. = I. chinensis.

EMMENOSPERMA, F. Muell. Fragm. iii. 62 (1863). *RHAMNEAE*, Benth. & Hook. f. i. 999. alphitonoides, *F. Muell. l. c.* 63.—Austral.
Cunninghamii, *Benth. Fl. Austral.* i. 415.—Austral.
Panchericanum, *Baill. Adansonia*, xi. (1874) 289.—N. Caled.

EMMENOSPERMUM, C. B. Clarke, ex Hook. f. Fl. Brit. Ind. iv. 249, 304 (1883) = Phthetirospermum, Bonge (Scrophul.).
Parrasia, C. B. Clarke, ex Hook. f. l. c. 304 = P. Parishii.

EMMENOSPERMUM, F. Muell. Census, 60 (1882) = Emmenosperma, F. Muell. (Rhamneae).

EMMEORHIZA, Endl. Gen. 865 (1838) = Emeorhiza, Pohl (Rubiac.).

EMMOTIUM, Meisan. Comm. 112 (1838) = seq.

EMMOTUM, Desv. in Ham. Prod. Pl. Ind. Occ. 29 (1825). *OLACINEAE*, Benth. & Hook. f. i. 352.
PODOPETALUM, Benth. in Trans. Linn. Soc. xvii. (1841) 684. t. 42.
SCHNIZLEINIA, Mart. ex Engl. in Mart. Fl. Bras. xii. 11. 46 (1872).
acuminatum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. x. (1852) 178.—Bras.
affine, Miers, l. c. 180 = acuminatum.
apogon, *Griseb. in Gottt. Abh.* xxiv. (1879) 149.—Reg. Argent.
fagifolium, *Desv. in Ham. Prod. Pl. Ind. Occ.* 29.—Guiana.
glabrum, *Benth. ex Miers, in Ann. & Mag. Nat. Hist.* Ser. III. iv. (1859) 366.—Bras.
nitens, *Miers, l. c.* Ser. II. x. (1852) 180.—Bras.
orbiculatum, *Miers, l. c.* 178.—Bras.

EMORYA, Torr. in Bot. Mex. Bound. 121. t. 26 (1858). *LOGANIACEAE*, Benth. & Hook. f. ii. 794.
suaveolens, *Torr. l. c.*—Am. bor. occ.

EMPEDOCLEA, A. St. Hil. Fl. Bras. Mer. i. 19. t. 5 (1825). *DILLENIACEAE*, Benth. & Hook. f. i. 11.
alnifolia, *A. St. Hil. l. c.* 20.—Bras.

EMPEDOCLIA, Rafin. Caratt. Gen. Sic. 78 (1810) = Sideritis, Tourn. (Labiat.).
montana, Rafin. l. c. 79 = S. sicula.

EMPETRUM, [Tourn.] Linn. Syst. ed. I (1735). *EMPETRACEAE*, Benth. & Hook. f. iii. 414.
aciculare, Bertol. Misc. Bot. xiii. 20 = Ceratiola ericoides.
album, Linn. Sp. Pl. 1022 = Corema album.
andinum, Phil. ex A. DC. in DC. Prod. xvi. i. 96 = nigrum.
Camradii, Torr. in Ann. Lyc. N. York, iv. (1846) 88 = Corema Conradii.
crassifolium, Rafin. New Fl. Am. iii. 51 = nigrum.

EMPETRUM:—
nodium, Carmich. in Trans. Linn. Soc. xii. (1818) 506 = nigrum.
nigrum, *Linn. Sp. Pl.* 1022.—Europ.; As. bor.
pinnatum, Lam. Encyc. i. 567 = Margyricarpus setosus.
procumbens, Gilib. Exercit. ii. 398 = nigrum.
purpureum, Rafin. New Fl. Am. iii. 50 = nigrum.
rubrum, Vahl, ex Willd. Sp. Pl. iv. 718 = nigrum.
scoticum, Hook. ex Stend. Nom. ed. II. i. 553 = nigrum.
serrulatum, Pritz. Ic. Ind. i. 895, sphalm. = Empleurum serrulatum.

EMPHYSOPUS, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 118 = Lagenophora, Cass. (Compos.).
Gunnii, Hook. f. l. c. 114 = L. Emphysopus.

EMPLEOTOCLADUS, Torr. in Smithson. Contrib. vi. (1854) (Pl. Frémont.) 10. t. 5. *ROSACEAE*, Benth. & Hook. f. i. 614.
fasciculatus, *Torr. l. c.*—Am. bor. occ.

EMPLEURIDIUM, Sond. & Harv. ex Harv. Thes. Cap. i. 49 (1859). *RUTACEAE*, Benth. & Hook. f. i. 291.
juniperinum, *Sond. & Harv. l. c.*—Afr. austr.

EMPLEUROSMA, Bartl. in Lehm. Pl. Preiss. ii. 228 (1847) = Dodonæa, Linn. (Sapind.).
virgata, Bartl. l. c. = D. pinifolia.

EMPLEURUM, [Soland. in] Ait. Hort. Kew. ed. I. iii. 340 (1789). *RUTACEAE*, Benth. & Hook. f. i. 291.
Aitoni, *J. F. Gmel. Syst.* 268 = serrulatum.
crenatum, Eckl. & Zeyh. Enum. 117 = serrulatum.
serrulatum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. iii. 340. —Afr. austr.

EMPODIUM, Salisb. Gen. Pl. Fragm. 43 (1866) = Curculigo, Gaertn. (Amaryll.).
plicatum, Salisb. l. c. 44 = C. plicata.

EMPOGONA, Hook. f. in Hook. Ic. Pl. t. 1091 (1871). *RUBIACEAE*, Benth. & Hook. f. iii. 94.
Kirkii, *Hook. f. l. c.*—Afr. trop.

EMPUSA, Lindl. Bot. Reg. sub t. 825 (1824) = Liparis, Rich. (Orchid.).
pardoxa, Lindl. l. c. = L. odorata.

EMPUSARIA, Reichb. Consp. 69 (1828) = prec.

EMULARIA, Rafin. Fl. Tellur. iv. 81 (1836) = Justicia, Linn. (Acanth.).

EMULINA, Rafin. Fl. Tellur. iv. 81 (1836) = Convolvulus, Tourn.
parviflora, Rafin. l. c. = C. parviflorus.

EMURTIA, Rafin. Sylva Tellur. 106 (1838) = Eugenia, Mich. (Myrtac.).
emarginata, Rafin. l. c. = E. emarginata.
guayaquillensis, Rafin. l. c. = E. guayaquillensis.
micrantha, Rafin. l. c. = E. Maximiliana.
punicifolia, Rafin. l. c. = E. punicæfolia.

ENAIMON, Rafin. Sylva Tellur. 9 (1838) = Olea, Tourn.
undulata, Rafin. l. c. = O. laurifolia.

ENALCIDA, Cass. in Bull. Soc. Philom. (1819) 81 = Thymenæa, Linn. (Compos.).
farnicultifolia, Cass. in Dict. Sc. Nat. xiv. 448 = T. coronopifolia.
pilifera, Cass. in Bull. Soc. Philom. (1819) 81 = T. coronopifolia.

ENANTIA, Falc. in Hook. Journ. Bot. iv. (1842) 75 = Sabia, Colebr.

ENANTIA, Oliver, in Journ. Linn. Soc. ix. (1867) 174. *ANONACEAE*, Benth. & Hook. f. i. 998.
chlorantha, *Oliver, l. c.* 175.—Afr. trop.

ENANTIOSPARTON, C. Koch, Dendrol. i. 34 (1869) = Genista, Linn. (Legumin.).
radiatum, C. Koch, l. c. = G. radiata.

ENANTIOTRICHUM, E. Mey. ex DC. Prod. vi. 448 (1837) = Euryops, Cass. (Compos.).
Athanasiae, E. Mey. ex DC. l. c. 444 = Euryops speciosissimus.
Serra, E. Mey. ex DC. l. c. = Euryops Serra.

ENARGEA, Banks, ex Gaertn. Fruct. i. 283 (1788) = Lasiarrhiza, Ruiz & Pav. (1802) (Liliac.).
marginata, Gaertn. l. c. t. 59 = L. marginata.

ENARTEA, Stend. Nom. ed. II. i. 553 (1840), sphalm. = prasc.

ENARTHROCARPUS, Labill. Ic. Pl. Syr. Dec. v. 4. t. 2 (1812). *CRUCIFERAE*, Benth. & Hook. f. i. 99.
HUASONIA, Boiss. Diagn. Ser. I. viii. 46 (1849).
aneops, *Godr. in Mém. Acad. Montp. (Sect. Médic.)* i. (1858) 413.—Hab.?
arcuatus, *Labill. Ic. Pl. Syr. Dec.* v. 4.—Reg. Mediterr.
arcuatus, Loisel. Nouv. Not. 29 = clavatus.
Blaisei, *Castagne, Suppl. Cat. Pl. Marseille*, 11.—Gallia.
clavatus, *Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.)* i. (1858) 412.—Afr. bor.
lyratus, *DC. Syst.* ii. 661.—Graecia; Aegypt.
pterocarpus, *DC. l. c.*—Aegypt.
pterocarpus, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1858) 413 = aneops.
recurvus, *Pomel, Nouv. Mat. Fl. Alt.* 357.—Algeria.
strangulatus, *Boiss. Diagn.* Ser. I. viii. 44.—Aegypt.; Arab.
trabalis, *Pomel, Nouv. Mat. Fl. Alt.* 358.—Algeria.

ENARTOCARPUS, Poir. in Dict. Sc. Nat. xiv. 444 (1819) = Enarthrocarpus, Labill. (Crucif.).

ENCELIA, Adans. Fam. ii. 128 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 378.
ARMANIA, Bert. ex DC. Prod. v. 576 (1836).
BARATTIA, A. Gray & Engelm. in Proc. Am. Acad. i. (1846) 48.
EUCALIA, Raeusch. Nom. ed. III. 251, 385 (1797).
EUCNIDA, Lam. in Verh. Dict. v. 300 (1549).
GERAEA, Torr. & Gray, in Proc. Am. Acad. i. (1847) 48.
PALLASIA, L'Hérit. Dias. (1784) ex Ait. Hort. Kew. iii. 198 (1789).
SIMSIA, Pers. Syn. ii. 478 (1807).
albescens, *A. Gray, in Proc. Am. Acad.* viii. (1873) 356.—Mexic.
alterniflolia, Raeusch. Nom. ed. III. 251 = canescens.
amplexicaulis, *Hemsl. Biol. Centr. Am. Bot.* ii. 183.—Mexic.
argophylla, A. Gray, in Proc. Am. Acad. viii. (1873) 357 = Helianthella argophylla.
californica, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 287.—Calif.
calva, *A. Gray, in Proc. Am. Acad.* viii. (1873) 358.—Texas.
canescens, *Cav. Ic.* i. 46. t. 61.—Peruv.
conspersa, Benth. Bot. Voy. Sulph. 26 = halimifolia?
conspersa, A. Gray, Bot. Mex. Bound. 88 = californica.
cordata, *Hemsl. Biol. Centr. Am. Bot.* ii. 183.—Mexic.
dentata, Poir. Encyc. Suppl. v. 665 = Verbesina dentariocephala.
dentariocephala, *A. Gray, in Proc. Am. Acad.* viii. (1873) 357.—Am. bor. occ.
exaristata, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 183.—Am. bor. occ.
farinosa, *A. Gray, ex Torr. in Emory, Notes Mil. Reconnais.* 148.—Calif.
fendleri, *Hemsl. Biol. Centr. Am. Bot.* ii. 183.—Mexic.
frutescens, *A. Gray, in Proc. Am. Acad.* viii. (1873) 357.—Am. bor. occ.
Gheiseghetii, *A. Gray, l. c.* 358.—Mexic.
grandiflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 184.—Nicaragua.
halimifolia, *Cav. Ic.* iii. 6. t. 210.—Mexic.
heterophylla, *Hemsl. Biol. Centr. Am. Bot.* ii. 184.—Mexic.
hispida, *Anderss. in Galeap. Veget. Ins.* ii. 186.—Ins. Galapag.
hispida, *Hemsl. Biol. Centr. Am. Bot.* ii. 184.—Mexic.
lagascaeformis, *A. Gray, ex Hemsl. l. c.*—Mexic.

ENCELIA:—
linensis, Stend. Nom. ed. II. i. 490, in syn. = canescens.
mexicana, *Mart. ex DC. Prod.* v. 578.—Mexic.
microcephala, A. Gray, in Proc. Am. Acad. viii. (1873) 357 = Helianthella microcephala.
microphylla, *A. Gray, l. c.* xv. (1880) 87.—Mexic.
nivea, Benth. Bot. Voy. Sulph. 27 = Viguiera nivea.
nudicaulis, A. Gray, in Proc. Am. Acad. viii. (1872) 656 = Helianthella nudicaulis.
obionifolia, *DC. Prod.* v. 567.—Chili.
parvifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 206.—Peruv.
polycephala, *Hemsl. Biol. Centr. Am. Bot.* ii. 184.—Nicaragua.
sanguinea, *Hemsl. l. c.* 185.—Mexic.
scaposa, *A. Gray, in Proc. Am. Acad.* viii. (1873) 657 —N. Mexic.
sericea, *Hemsl. Biol. Centr. Am. Bot.* ii. 185.—Guatemala.
stenophylla, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 41.—Am. bor. occ.
suberistata, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot* ii. 185.—Mexic.
tomentosa, *Walp. in Linnaea*, xiv. (1840) 804.—Peruv.
viscida, *A. Gray, in Proc. Am. Acad.* xi. (1876) 78.—Calif.

ENCENTRUS, Presl, Bot. Bemerk. 38 (1844); in Abh. Boehm. Ges. V. iii. (1845) 483 = Gymnosporia, Benth. & Hook. f. (Celastrin.).
linearis, Presl, ll. cc. (= *Celastrus linearis*).—Afr. austr.

ENCEPHALARTOS, Lehm. Pugill. vi. 3. t. 1 ad 5 (1834). *CYCADACEAE*, Benth. & Hook. f. iii. 445.
ARTHROZAMIA, Reichb. Consp. 40 (1828).
acanthus, *Mart. in Gard. Chron.* (1878) 11. 810.—Afr. austr.
Altensteinii, *Lehm. Pugill.* vi. 11. t. 4 et 5.—Afr. austr.
Barteri, *Carruth. ex Miq. in Arch. Neerl.* iii. (1868) 248.—Afr. trop.
brachyphyllus, Lehm. & De Vriese, in Hoev. & De Vriese, Tijdschr. iv. (1837–38) 414 = caffer.
caffer, Hook. Bot. Mag. t. 4908 = longifolius.
caffer, Lehm. Pugill. vi. 14 = lanuginosus.
caffer, *Miq. Monogr. Cycad.* 58.—Afr. austr.
cafforum, Hort. Slon, ex Yates, in Proc. Linn. Soc. ii. (1849) 19 = caffer.
cycadifolius, *Lehm. Pugill.* vi. 12.—Afr. austr.
Cycadis, Sweet, Hort. Brit. ed. III. 636 = caffer.
Denisonii, F. Muell. in Journ. Pharm. Soc. Vict. ii. 90 = Macrozamia Peroviskiana.
Douglasii, F. Muell. Census, 110 = Macrozamia Douglasii.
Dyeri, F. Muell. in Melb. Chemist (June 1885) (= Macrozamiae sp.).—Austral.
elongatus, *Miq. in De Hoev. & De Vriese, Tijdschr.* iv. (1837–38) 419.—Afr. austr.
ferox, *Bertol. f. in Mem. Acc. Sc. Bolog.* iii. (1851) 264.—Afr. trop.
Frazeri, Miq. in Verol. en Med. Akad. Wetensch. xv. (1863) 268 = Macrozamia Frazeri.
Friderici-Guilielmi, Lehm. Pugill. vi. 8 = cycadifolius.
Ghellinckii, *Lem. Illustr. Hortic.* (1868) 79. t. 567.—Afr. austr.
Hildebrandtii, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1874) 8.—Afr. trop.
horridus, *Lehm. Pugill.* vi. 14.—Afr. trop.
lanuginosus, Hort. ex Miq. in Verh. Kon. Acad. Wetensch. II. iii. (1860) 55 = Macrozamia Paulli-Guilielmi.
lanuginosus, *Lehm. Pugill.* vi. 14.—Afr. austr.
latifolius, Sweet, Hort. Brit. ed. III. 636 = Zamia latifolia.
latifolius, *Lehm. in Hoev. & De Vriese, Tijdschr.* iv. (1837–38) 424.—Hab.?
Lehmanni, *Lehm. Pugill.* vi. 14.—Afr. austr.
longifolius, *Lehm. l. c.*—Afr. austr.
Macdonnellii, F. Muell. ex Miq. in Verol. en Med. Kon. Akad. Wet. xv. (1863) 276 = Macrozamia Frazeri.
Mackenzii, Hort. ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1860) 55 = Macrozamia Paulli-Guilielmi.

ENCEPHALARTOS:—
Marumii, De Vriese, in Hoev. & De Vriese, Tijdschr. v. (1838–39) 187 = Altenensteinii.
mauritianus, Miq. Monog. 48 = Lehmanni.
Miquelii, F. Muell. Fragm. iii. 38 = Macrozamia Miquelii.
Moorei, F. Muell. l. c. xi. 125 (Macrozamiae sp.).—Austral.
nanus, Lehm. in Hoev. & De Vriese, Tijdschr. iv. (1837–38) 421 = horridus.
Oldfieldii, Miq. in Verh. en Med. Kon. Akad. Wetensch. xv. (1863) 370 = Macrozamia Fraseri.
Paulo-Guilielmi, F. Muell. in Trans. Pharm. Soc. Vict. ii. 91 = Macrozamia Paul-Guilielmi.
Poggei, *Aschers. in Verh. Bot. Ver. Brand.* xx. (1878) p. xxxv.—Afr. trop.
Preissii, F. Muell. in Journ. Pharm. Soc. Vict. ii. 90 = Macrozamia Fraseri.
pruniferus, *Sweet, Hort. Brit.* ed. III. 626.—Hab.?
pumilus, Sweet, l. c. = Zamia pumila.
pungens, *Lehm. Pugill.* 24.—Afr. austr.
repandus, Sweet, Hort. Brit. ed. III. 626 = Zamia repanda.
spinulosus, Lehm. in Hoev. & De Vriese, Tijdschr. iv. (1837–38) 420 = Lehmanni.
spiralis, Lehm. Pugill. vi. 13 = Macrozamia spiralis.
tridentatus, *Lehm. l. c.*—Afr. austr.
Vanhallii, De Vriese, in Hoev. & De Vriese, Tijdschr. iv. (1837–38) 422 = horridus.
Verschaffelti, *Regel, in Act. Hort. Petrop.* iii. (1875) 111; *Gartenfl.* (1875) 32. t. 932.—Hab.?
villosus, *Lem. Illustr. Hortic.* (1868) *Misc.* 69. t. 557. — Afr. austr.

ENCEPHALLARTES, Endl. Gen. Suppl. ii. Add. 108 (1842) = prec.

ENCHELVA, Lem. in Orb. Dict. v. 300 (1849) = **Enoella**, Adans. (Compos.).

ENCHIDION, Muell. Arg. in DC. Prod. xv. ii. 1256 (1866) = seq.

ENCHIDIUM, Jack, in Malay. Misc. ii. (1822) n. vii. 89 : reimpr. in Hook. Comp. Bot. Mag. i. (1836) 257 = **Trigostemon**, Blume (Euphorb.).
verticillatum, Jack, l. c. = T. indicus.

ENCHOLIRION, Benth. & Hook. f. Gen. iii. 667 (1883) = seq.

ENCHOLIRIUM, Mart. in Schult. f. Syst. vii. 68, 1238 (1830) = **Dyckia**, Schult. f. (Bromel.).
Augustae, R. Schomb. in Verh. Preuss. Gartenb. Ver. (1846) 18. t. 2.—Cf. Bot. Zeit. iv. (1846) 454 = Caragata Augustae.
corallinum, Linden, Illustr. Hortic. xviii. (1871) t. 70 = Tillandsia corallina.
Garreli, Beer, Bromel. 37 = D. altissima.
Jonghii, Libon, in Koch, Allg. Gartenz. (1857) 22 = T. Jonghii.
Saundersii, Hort. in Rev. Hortic. (1872) 300; André, in Illustr. Hortic. xx. (1873) t. 159 = Tillandsia Saundersii.
spectabile, Mart. ex Schult. f. Syst. vii. 11. 1233 (Dyckiae sp.).—Bras.
Vaghii, Carr. in Rev. Hortic. (1878) 140.—Hab.?

ENCHYDRA, F. Muell. Fragm. iii. 139 (1863) sphalm. = **Enhydra**, Lour. (Compos.).

ENCHYLAENA, R. Br. Prod. 407 (1810). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 61.
Eucnylaena, Spreng. Gen. i. 928 (1830).
aegyptiaca, Spreng. Syst. i. 938 = Suaeda baccata.
Forsythenia, Spreng. l. c. = Suaeda baccifera.
marginata, *Benth. Fl. Austral.* v. 182.—Austral.
micrantha, *Benth. l. c.* 181.—Austral.
microphylla, *Moq. in DC. Prod.* xiii. ii. 198.—Austral.
pavonicae, R. Br. Prod. 408 = tomentosa.
physophora, Spreng. Syst. i. 938 = Suaeda physophora.
pubescens, Moq. in DC. Prod. xiii. ii. 198 = tomentosa.
tomentosa, *R. Br. Prod.* 408.—Austral.
villosa, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 76.—Austral.

ENCHYLIS, Ehrh. Beitr. iv. 147 (1789) = **Sedum**, Tourn. (Crassul.).

ENCHYSIA, Presl, Prod. Monog. Lobel. 40 (1836) = **Laurentia**, Neck. (Campanulac.).
Baueri, Presl, l. c. = Isotoma fluviatilis?
dentata, A. DC. in DC. Prod. vii. 409 = L. repens.
erecta, A. DC. l. c. = L. repens.
erinodes, Presl, Prod. Monog. Lobel. 40, excl. syn. = L. repens.
Gaudichaudii, Presl, l. c. 41 = Isotoma fluviatilis?
Lessonii, Presl, l. c. 40 = Isotoma fluviatilis?
repens, Presl, l. c. = L. repens.
scapigera, Presl, l. c. 41 = Isotoma scapigera.
secunda, Sond. in Harv. & Sond. Fl. Cap. iii. 551 = L. repens.

ENCILIA, Reichb. Nom. 166 (1841) = **Brcilia**, A. Juss. (Phytolac.).

ENCKEA, Kunth, in Linnaea, xiii. (1839) 590 = **Piper**, Linn.
Amalago, Griseb. Cat. Pl. Cub. 64 = P. cubense.
Amalago, Griseb. Fl Brit. W. Ind. 169, partim = P. unguiculatum.
articulata, Griseb. Cat. Pl. Cub. 65 = P. tenue.
blattarum, Kunth, in Linnaea, xiii. (1839) 596 = P. blattarum.
californica, Kunth, l. c. 601 = P. californicum.
californica, Miq. Syst. Pip. 368 = P. triquetrum.
crenathifolia, Kunth, in Linnaea, xiii. (1839) 597 = P. crenathifolium.
decrescens, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 440 = P. decrescens.
dolcei, Kunth, in Linnaea, xiii. (1839) 595 = P. Bertcroanum.
dubia, Kunth, l. c. 606 = P. Poiteanum.
glaucescens, Kunth, l. c. 593 = P. unguiculatum.
heptandra, Miq. Syst. Pip. 350 = P. heptandrum.
Kunthii, Miq. l. c. 368 = P. Kunthii.
laevigata, Kunth, in Linnaea, xiii. (1839) 602 = P. laevigatum.
laevigata, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 440 = P. tenue.
lata, Kunth, in Linnaea, xiii. (1839) 606 = P. latum.
Lessertiana, Miq. Syst. Pip. 349 = P. tenue.
Lindenii, Miq. l. c. 358 = P. Lindenii.
Martiana, Miq. in Linnaea, xx. (1847) 132 = P. Enckea.
miconiaefolia, Benth. Bot. Voy. Sulph. 167 = P. miconiaefolium.
orthostachya, Kunth, in Linnaea, xiii. (1839) 601 = P. orthostachyum.
plantaginea, Kunth, l. c. 599 = P. medium.
platyphylla, Benth. Bot. Voy. Sulph. 167 = P. platyphyllum.
reticulata, Miq. Syst. Pip. 365 = P. reticulatum.
Schlechtendalii, Miq. l. c. 362 = P. Schlechtendalianum.
Sieberi, Griseb. Fl. Brit. W. Ind. 170 = P. medium.
Sieberi, Miq. Syst. Pip. 358 = P. Sieberi.
smilacifolia, Kunth, in Linnaea, xiii. (1839) 605 = P. smilacifolium.
stipulacea, Kunth, l. c. 600 = P. stipulaceum.
subpanduraeformis, Miq. in Linnaea, xxvi. (1853) 217 = P. panduratum.
Swartzii, Griseb. Fl. Brit. W. Ind. 169 = P. medium.
tenuis, Kunth, in Linnaea, xiii. (1839) 603 = P. tenue.
unguiculata, Kunth, l. c. 592 = P. unguiculatum.
verrucifolia, Miq. Syst. Pip. 355 = P. verrucifolium.
vernicea, Miq. l. c. 359 = P. orthostachyum.

ENCKIANTHUS, Desf. Cat. Hort. Par. ed. III. 187 (1829) = **Enkianthus**, Lour. (Ericac.).

ENCKLEIA, Pfeiff. Nom. i. 1199 (1874) = ENKLEIA, Griff. = **Linostoma**, Wall. (Thymel.).

ENCLIANDRA, Zucc. in Abh. Akad. Muench. ii. (1837) 385 = **Fuchsia**, Linn. (Onagrar.).
parviflora, Zucc. l. c. 387 = F. Encliandra.
thymifolia, *Lilja, in Linnaea*, xv. (1841) 262 = F. thymifolia.

ENCOPA, Griseb. Cat. Pl. Cub. 184 (1866). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 987.
tenuifolia, *Griseb. l. c.*—Cuba.

ENCOPEA, Presl, Bot. Bemerk. 88 (1844); in Abh. Boehm. Ges. V. iii. (1845) 513 = **Faramea**, Aubl. (Rubiac.).
umbellifera, Presl, l. c. = F. umbellifera.

ENCOPIA, Benth. & Hook. f. Gen. ii. 126 = prec.

ENCUREA, Walp. Rep. i. 414 (1842) = ENOUREA, Aubl. = **Paullinia**, Linn. (Sapindac.).

ENCYANTHUS, Spreng. Syst. ii. 267 (1825) = **Enkianthus**, Lour. (Ericac.).

ENCYCLA, Benth. in DC. Prod. xiv. 10, in syn. (1856) sphalm. = EUCYDA, Nutt. = **Briogonium**, Michx. (Polygon.).

ENCYCLIA, Hook. Bot. Mag. t. 3881, 3013 (1838) = **Epidendrum**, Linn. (Orchid.).
amazonica, A. Brongn. ex Neum. in Rev. Hortic. Sér II. iv. (1845–46) 137.—Bras.
chloroleucum, Neum. l. c. 138 = Epid. chloroleucum.
Duboisiana, Neum. l. c. 137 = Epid. Duboisianum.
Hermentiana, A. Brongn. ex Neum. l. c. 138.—Venezuela.
macrochyla, Neum. l. c. 137 = Epid. atropurpureum.
mexicana, Hort. Berol. ex Otto & Dietr. Allg. Gartenz. v. (1837) 38, nonem.—Mexic.
patens, Hook. Bot. Mag. t. 3018 = Epid. odoratissimum.
phoenicea, Neum. in Rev. Hortic. Sér. II. iv. (1845–46) 137 = Epid. phoeniceum.
viridiflora, Hook. Bot. Mag. t. 2881.—Bras.

ENCYCLIA, Poepp. & Endl Nov. Gen. et Sp. ii. 10 (1838) = **Polystachya**, Hook. (Orchid.).
macrostachya, Poepp. & Endl. l. c. 11 = Epidendrum armeniacum.
nana, Poepp. & Endl. l. c. 10.—Peruv.
polystachya, Poepp. & Endl. l. c.—Peruv.

ENCYCLIUM, Neum. in Rev. Hortic. Sér. II. iv. (1845–46) 137 = **Epidendrum**, Linn. (Orchid.).

ENDALLEX, Rafin. in Ser. Bull Bot. i. (1830) 220 = **Phalaris**, Linn. (Gramin.).
arundinacea, Rafin. l. c. = P. arundinacea.

ENDAMMIA, Rafin. New Fl. Am. iii. 49 (1836) = Coremа. D. Don (Empetr.).
ericoides, Rafin. l. c. = C. Conradi.

ENDECARIA, Rafin. Sylva Tellur. 102 (1838) = **Cuphea**, P. Br. (Lythrar.).
coccinea, Rafin. l. c. = C. Llavea.

ENDEISA, Rafin. Fl. Tellur. ii. 51 (1836) = **Dendrobium**, Sw. (Orchid.).
flava, Rafin. l. c. = D. densiflorum.

ENDEMA, Pritz. Ic. Ind. i. 398 (1855) sphalm. = EUDEMA, Humb. & Bonpl. = **Brayа**, Sternb. & Hoppe (Crucif.).

ENDERA, Regel, Gartenfl. (1872) 226. t. 732 = **Tacsarum**, Brongn. (Aroid.).
conophalloides, Regel, l. c. = T. cylindricum.

ENDESPERMUM, Blume, Cat. Gew. Buitenz. 92 (1823) = **Dalbergia**, Linn. (Legumin.).
diversifolium, Blume, ex Miq. Fl. Ind. Bat. i. 1. 128 = D. Sissoo.
penduliflorum, Zipp. ex Miq. l. c. 128 = D. stipulacea.
retusum, Blume, ex Miq. l. c. 131 = D. tamarindifolia.
scandens, Blume, Cat. Gew. Buitenz. 92 = D. Championi ?
seplentrionum, Champ. ex Thw. Enum. Pl. Zeyl. 94 = D. Championi.

ENDIANDRA, R. Br. Prod. 409 (1810). *LAURINEAE*, Benth. & Hook. f. iii. 154.
Dictyodaphne, Blume, Mus. Bot. Lugd. Bat. i. 270 (1850).
Silvara, Meisn. in DC. Prod. xv. i. 84 (1864).
Silvia, Allem. Diss. Rio Janeiro (1845).
Candolleana, Meisn. ex Kurz, For. Fl. Brit. Burma, ii. 295 = Dehaasia cuneata.
cuneata, *Miq. Fl. Ind. Bat.* i. 1. 919.—Sumatra.
discolor, *Benth. Fl. Austral.* v. 201.—Austral.
firma, *Nees, in Wall. Pl. As. Rar.* ii. 61.—Ind. or. glauca, *R. Br. Prod.* 403.—Austral.
hypotephra, *F. Muell. Fragm.* v. 166.—Austral.
jamaicensis, Spreng. Syst. i. 176 = Acrodiclidium jamaicense.
Muelleri, *Meisn. in DC. Prod.* xv. i. 209.—Austral.
pubens, *Meisn. l. c.*—Austral.

ENDIANDRA, —
rubescens, *Blume, ex Miq. Pl. Yungh.* 176.—Java.
Sieberi, *Nees, Syst. Laurin.* 194.—Austral.
sumatrana, *Miq. Fl. Ind. Bat.* i. 1. 919.—Sumatra.
viscos, *F. Muell. Fragm.* ii. 90.—Austral.
viridis, J. Presl, Rostl. ii. 68 = Acrodiclidium jamaicense.

ENDIPLUS, Raf. in Am. Monthly Mag. (1811) 856; (1812) 191 = **Phacelia**, Juss. (Hydrophyll.).
bifidus, Raf. l. c. (1811) 856.—Am. bor.
phacelioides, Raf. in Journ. Phys. lxxxix. (1819) 99 = P. Endiplus.

ENDIUSA, Alef. in Oestr. Bot. Zeit. ix. (1859) 359 = **Vicia**, Tourn. (Legumin.).
hirsuta, Alef. l. c. 360 = V. hirsuta.
Loiseleurii, Alef. l. c. (= *Ervum Loiseleurii*).—Europ.

ENDLICHERA, Presl, Symb. Bot. i. 78. t. 49 (1832) = **Emeorhiza**, Pohl (Rubiac.).
brasiliensis, Presl, l. c. = Emeorhiza brasiliensis.

ENDLICHERIA, Nees, in Linnaea, viii. (1833) 87 = **Aydendron**, Nees (Laurin.).
anomala, Nees, ex Meisn. in Mart. Fl. Bras. v. ii. 233 (= *Goeppertia anomala*).—Bras.
hirsuta, Nees, in Linnaea, viii. (1833) 88 = Cryptocarya hirsuta.
sericea, Nees, l. c. (= *Goeppertia sericea*).—Ind. occ.

ENDOCARPA, Raf., Sylva Tellur. 125 (1838) = Cryptocarya, R. Br. (Laurin.).
corymbosa, Raf. l. c. = C. dubia.

ENDOCELLION, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xxxvii. (1865) i. 375, in nota. *COMPOSITARUM* genus dubium.
boreale, *Turcz. l. c.*—Rossia.

ENDOCODON, Raf., Fl. Tellur. iv. 49 (1836) = **Calathea**, G. F. W. Mey. (Scitamin.).
zebrina, Raf. l. c. = C. zebrina.

ENDOCOMA, Raf., Fl. Tellur. ii. 57 (1836) = **Bottionea**, Colla (Liliac.).
parviflora, Raf. l. c. = B. thysanotoides.
peruviana, Raf. l. c. = B. thysanotoides.

ENDODACA, [Schlecht. in] Linnaea, ix. (1834) Litt. 98 = seq.

ENDODECA, Raf., Med. Fl. i. 62 (1828) = **Aristolochia**, Linn.
Bartonii, Klotzsch, in Monatsb. Akad. Berl. (1859) 600 = A. Serpentaria.
dodecandra, Raf., Med. Fl. i. 62 = A. Serpentaria?
hastata, Raf., Fl. Tellur. iv. 99 = A. Serpentaria?
polyrrhiza, Klotzsch, in Monatsb. Akad. Berl. (1859) 601 = A. Serpentaria.
sempervirens, Raf., Fl. Tellur. iv. 99 = A. sempervirens.
Serpentaria, Raf., Med. Fl. i. 62 = A. Serpentaria.

ENDODESMIA, Benth. in Benth. & Hook. f. Gen. i. 166 (1862). *HYPERICINEAE*, Benth. & Hook. f. i. 166.
calophylloides, *Benth. l. c.*—Afr. trop.

ENDODIA, Raf., Neogenyt. 4 (1825) = **Leersia**, Sw. (Gramin.).
lenticularis, Raf., l. c. = L. lenticularis.

ENDOCLES, Salisb. Gen. Pl. Fragm. 51 (1866) = **Stigmatanus**, Michx. (Liliac.).
lustrus, Salisb. l. c. = Z. muscatetorios.

ENDOGONA, Raf., Fl. Tellur. ii. 37 (1836) = **Anthericum**, Linn. (Liliac.).
Adrantheres, Raf., l. c. 38 = Dianella ensifolia.
ramosa, Raf., l. c. = A. ramosum.

ENDOGONIA, Lindl. Veg. Kingd. 656 (1847) = **Trigonotis**, Stev. (Boragin.).

ENDOISILLA, Raf., Sylva Tellur. 114 (1838) = **Euphorbia**, Linn.
Myrsinites, Raf., l. c. = E. Myrsinites.

ENDOLASIA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 579 = **Manettia**? Mutis (Rubiac.).
Funchii, Turcz. l. c. 580.—N. Granat.

ENDOLEPIS, Torr. ex A. Gray, in Pacif. Rail. Rep. xii. ii. (Bot. Stev. Exped.) 47 (1860) = **Atriplex**, Linn. (Chenop.).
Suckleyi, Torr. l. c. = A. Endolepis.

ENDOLEUCA, Cass. in Bull. Soc. Philom. (1819) 47; et Dict. Sc. Nat. xiv. 474 = **Metalasia**, R. Br. (Compos.).
pulchella, Cass. in Dict. Sc. Nat. xiv. 474 = M. Cephalotes.
sphaerocephala, Cass. l. c. 475 = M. Cephalotes.

ENDOLOMA, Rafin. Sylva Tellur. 79 (1838) = **Bignonia**, Linn.
purpurea, Rafin. l. c. 80 = B. purpurea.

ENDOMELAS, Rafin. Fl. Tellur. iv. 67 (1836) = **Thunbergia**, Linn. f. (Acanth.).
alata, Rafin. l. c. = T. alata.

ENDONEMA, A. Juss. in Ann. Sc. Nat. Sér. III. vi. (1846) 26. t. 8. *PENAEACEAE*, Benth. & Hook. f. iii. 202.
GLISCHROCOLLA, A. DC. in DC. Prod. xiv. 490 (1856). retsioides, *Sond. in Linnaea*, xxiii. (1850) 103.—Afr. austr.
Thunbergii, *A. Juss. in Ann. Sc. Nat. Sér. III.* vi. (1846) 19, 26, t. 3.—Afr. austr.

ENDOPAPPUS, Sch. Bip. in Bonplandia, viii. (1860) 369 = **Chrysanthemum**, Tourn. (Compos.).
macrocarpus, Sch. Bip. l. c. (= *Pyrethrum macrocarpum*).—Alger.

ENDOPLECTRIS, Rafin. Fl. Tellur. iii. 56 (1836) = **Epimedium**, Tourn. (Berberid.).
tricolor, Rafin. l. c. 57 = Epim. macranthum.

ENDOPOGON, Nees, in Wall. Pl. As. Rar. iii. 76 (1832) = **Strobilanthes**, Blume (Acanth.).
Amomum, Nees, in Wall. l. c. 99 = S. consanguineus.
argutus, Nees, in DC. Prod. xi. 104 = S. viscosus.
Beddomianus, Wight, Ic. iv. t. 1499 = S. foliosus
consanguineus, Nees, in Wall. Pl. As. Rar. iii. 99 = S. exarcolatus.
cuspidatus, Benth. in Linnaea, xxiv. (1851) 646 = S. cuspidatus.
decurrens, Nees, in DC. Prod. xi. 105 = S. adnatus.
digitalis, Nees, l. c. 104 = S. viscosus.
foliosus, Wight, Ic. iv. t. 1501 = S. foliosus.
Gardnerianus, Nees, in DC. Prod. xi. 723 = S. Gardnerianus.
hybridescens, Nees, in Wall. Pl. As. Rar. iii. 99 = S. consanguineus.
integrifolius, Dalz. in Hook. Kew Journ. ii. (1850) 343 = S. pilosiflatus.
khasyanus, Nees, in DC. Prod. xi. 104 = S. adnatus.
macrostegius, Nees, l. c. = S. macrostegius.
rhamniflolius, Wight, Ic. iv. t. 1511 = S. rhamniflolius.
Strobilanthes, Wight, l. c. t. 1500 = S. Zenkerianus.
urticolor, Wight, l. c. t. 1497 = S. cuspidatus.
viscosus, Am. ex Nees, in Hook. Comp. Bot. Mag. ii. (1836) 313 = S. viscosus.
vitellinus, Nees, in DC. Prod. xi. 723 = Phlogacanthus asperulus.

ENDOPOGON, Rafin. Fl. Tellur. ii. 54 (1836). *RUBIACEAE* (Quid?).
pumilum, *Rafin. l. c.* 55.—Am. bor.

ENDOPTERA, DC. Prod. vii. 178 (1838) = **Crepis**, Linn. (Compos.).
aspera, DC. l. c. = C. aspera.
dichotoma, Boiss. & Bal. Diagn. Ser. II. iii. 98 = C. multiflora.
Dioscoridis, DC. Prod. vii. 179 = C. Dioscoridis.
intermedia, Boiss. & Bal. Diagn. Ser. II. iii. 99.—Syria.

ENDORIMA, Rafin. in Am. Monthly Mag. (1819) 195 = **Baldwinia**, Nutt. (Compos.).
modesta, Rafin. Fl. Tellur. ii. 91 = B. uniflora?

ENDOSIPHON, T. Anders. ex Benth. & Hook. f. Gen. ii. 1085 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
pumilioides, *T. Anders. l. c.*—Afr. trop.

ENDOSPERMUM, Benth. Fl. Hongk. 204 (1861). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 322.
CAPELLENIA, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1866) 289.
borneense, *Benth. ex Muell. Arg. in Flora*, xlvii. (1864) 469.—Borneo.
chinense, *Benth. Fl. Hongk.* 204.—China; Malaya.
formicarum, *Becc. Malesia*, ii. (1884) 44.—N. Guinea.
malaccense, *Benth. ex Muell. Arg. in Flora*, xlvii (1864) 469.—Malaya.
moluccanum, *Becc. Malesia*, ii. (1884) 88.—Ins. Amboina.

ENDOSPERMUM, Endl. Gen. 1304 (1841) = ENDOSPERMUM, Blume = **Dalbergia**, Linn. (Legumin.)

ENDOSTEIRA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 575. *TILIACEAE*? Genus dubium, Benth. & Hook. f. i. 205.
oppositifolia, *Turcz. l. c.*—Ins. S. Vincent.

ENDOSTEPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 227 = **Galipea**, Aubl. (Rutac.).
jasminoides, Turcz. l. c. = G. multiflora.

ENDOTHECA, Rafin. Fl. Tellur. iv. 98 (1836) = ENDODECA, Rafin. = **Aristolochia**, Linn.

ENDOTIS, Rafin. Fl. Tellur. ii. 21 (1836) = **Allium**, Tourn. (Liliac.).
pyrenaica, Rafin. l. c. = A. suaveolens.

ENDOTRICHE, Steud. Nom. ed. I. 298 (1821) = **Gentiana**, Tourn.
campestris, Steud. l. c. = G. campestris.

ENDOTROPIS, Endl. Gen. 591 (1838) = **Cynanchum**, Linn. (Asclep.).
auriculata, Decne. in DC. Prod. viii. 546 = C. auriculatum.
caudata, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 128 = C. Wilfordi.
Jacquemontiana, Decne. in DC. Prod. viii. 547 = C. Jacquemontianum.
Meyeri, Decne. l. c. 546 = C. virens.
Roylei, Decne. l. c. 547 = C. Royleï.

ENDOTROPIS, Rafin. Sylva Tellur. 81 (1838) = **Rhamnus**, Linn.
oleifera, Rafin. l. c. = R. californica.

ENDRACHIUM, Juss. Gen. 133 (1789) = **Humbertia**, Comm. (Convolv.).
madagascariense, J. F. Gmel. Syst. 339 = H. madagascariensis.

ENDRESSIA, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 228 = **Lignatiouam**, Linn. (Umbellif.).
pyrenaica, J. Gay, l. c. 224 = L. pyrenaicum.

ENDUSA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. viii. (1851) 173 (nomen tantum). *OLACINEAE*? Benth. & Hook. f. i. 345.

ENDUSIA, Benth. & Hook. f. Gen. i. 535 (1865) = ENDIUSA, Alef. = **Vicia**, Tourn. (Legumin.).

ENDYMION, Dum. Fl. Belg. 140 (1827) = **Scilla**, Linn. (Liliac.).
amoenulus, Nym. Consp. 731 = S. sibirica.
campanulatus, Parl. Fl. Ital. ii. 478 = S. hispanica.
cedretorum, Pomel, Nouv. Mat. Fl. Atl. 364.—Algeria.
cernuus, Dum. Fl. Belg. 140 = S. festalis.
non-scriptus, Garcke, Fl. Deutschl. ed. I. 322 = S. festalis.
nutans, Dum. Fl. Belg. 140 = S. festalis.
patulus, Dum. l. c. = S. hispanica.

ENEMION, Rafin. in Journ. Phys. xci. (1820) 70 = **Leucocrinum**, Linn. (Anemonisac.).
biternatum, Rafin. l. c. = L. biternatum.
Raddeanum, Regel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) ii. 91 = L. Raddeanum.

ENEMIUM, Steud. Nom. ed. II. i. 554 (1840) = prsec.

ENEMOSYNE, Lehm. Pl. Preiss. ii. 236 (1847), sphalm. = **Bremosyne**, Endl. (Saxifrag.).

ENEODON, Rafin. Fl. Tellur. iii. 86 (1836) = **Leucas**, R. Br. (Labiat.), *urticifolia*, Rafin. 1. c. = L. urticæfolia.

ENGELHARDTIA, Leschen. ex Blume, Bijdr. 598 (1825); Fl. Jav. fasc. viii. 5 (1828). *JUGLANDEÆ*, Benth. & Hook. f. iii. 399.
Oremounnsea, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 52.
Pterilema, Reinw. in Syll. Ratisb. ii. (1828) 18.
acerifiora, *Blume, Fl. Jav.* t. 2, 5.—Reg. Himal.; Malaya.
chrysolepis, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 227 = Wallichiana.
Colebrookiana, *Lindl. in Wall. Pl. As. Rar.* iii. 4. t. 208.—Reg. Himal.; China.
Oreommnea, *C. DC. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 36.—Am. centr.
palembanica, Miq. Fl. Ind. Bat. Suppl. 846 = serrata.
parviflora, *C. DC. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 35.—Ins. Philipp.
philippinensis, C. DC. l. c. = spicata.
polystachya, *Roxlb. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 385.—Ind. or.
rigida, *Blume, Bijdr.* 598.—Java.
Roxburghiana, Lindl. in Wall. Cat. n. 4940 = spicata.
sclanica, Blume, Fl. Jav. Jugl. 8 = Shorea selanica.
serrata, *Blume, Fl. Jav.* t. 2, 5.—Malaya.
spicata, *Blume, Bijdr.* 598.—Reg. Himal.; Malaya.
villosa, Kurz, For. Fl. Brit. Burma, ii. 491 = Colebrookiana.
Wallichiana, *Lindl. in Wall. Cat.* n. 4942.—Malaya.

ENGELIA, Karst. ex Nees, in DC. Prod. xi. 721 (1847) = **Mendoncia**, Vell. (Acanthac.).
tovernis, Klotsch & Karst. ex Nees, l. c.—N. Granat.
villosa, Klotsch & Karst. ex Nees, l. c.—N. Granat.

ENGELMANNIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 253 = **Croton**, Linn. (Euphorb.).
Nuttalliana, Klotzsch, l. c. = C. monanthogynus.

ENGELMANNIA, Pfeiff. in Bot. Zeit. iii. (1845) 678 = **Cuscuta**, Linn. (Convolv.).
nigrana, Pfeiff. l. c. 674 = C. racemosa.
racemosa, Pfeiff. l. c. iv. (1846) 21 = C. racemosa.

ENGELMANNIA, Torr. & Gray, Fl. N. Am. ii. 383 (1841). *COMPOSITÆ*, Benth. & Hook. f. ii. 361.
Angelandra, Endl. Gen. Suppl. iii. 69 (1843).
Englemannia, A. Gray, ex Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 343.
pinnatifida, *Torr. & Gray, l. c.*—Am. bor. occ.

ENGLEMANNIA, A. Gray, ex. Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 343 = prsec.

ENHALUS, Rich. in Mém. Inst. Par. ii. (1811) (Math.-Phys.), 64. *HYDROCHARIDEÆ*, Benth. & Hook. f. iii. 454.
acoroides, Rich. ex Chatin, Anat. Pl. Aquat. 15. t. 6 = Koenigii.
Koenigii, *Rich. in Mém. Inst. Par.* ii. (1811) (*Math.-Phys.*), 78.—Malaya; Austral.
marinus, Griff. Notul. iii. 178 = Koenigii.

ENHYDRA, Lour. Fl. Cochinch. 510 (1790) (Hydrx). *COMPOSITÆ*, Benth. & Hook. f. ii. 360.
Cryptogrammum, Beauv. Fl. Owar. ii. 24. t. 74 (1805).
Hingcha, Roxb. Hort. Beng. 62 (1814); DC. Prod. v. 636 (1836).
Meyera, Schreb. Gen. 570 (1791).
Pfeillichera, Blume, ex DC. Prod. v. 636, in syn. (1836).
Soberya, Rafn & Pav. Prod. 109. t. 23 (1794).
Soseya, Pers. Syn. ii. 478 (1807).
Tetractis, Endl. Gen. 419, in syn. (1836).
Tetragotis, Raf. in Blume, Bijdr. 989 (1826).
Waelbergeria, Schum. & Thonn. Beskr. Guin. Pl. 302 (1827).
Beccarii, Asch. Sitzb. Ges. Nat. Fr. Berl. (1867) 5.—Borneo.
brevifolia, Rich. ex Chatin, Anat. Pl. Aquat. 15. t. 6.—Koenigii.
crispata, Blume, Bijdr. 989.—Java.
Dioscoreæ, Miq. Fl. Ind. Bat. Suppl. 846 = Koenigii.
Koenigii, Spreng. Syst. i. 605 (1825).
Lepinema, Rafin. Fl. Tellur. iii. 96 (1836).
Meyeriana, Zoll. & Mor. Verz. Pfl. Lomb. 116 (1846).
Rheedii, Wight, Ic. t. 1956.—Ind. or.
Roxburghii, Wight, Ic. t. 1957.—Ind. or.
Salisburia, Steud. Nom. ed. II. i. 554 (1840) = prsec.
Soberya, Rafin. Fl. Tellur. iii. 96 (1836).
Soseya, Rafin. Fl. Tellur. iii. 96 (1836).
Tetragotis, Rafin. Fl. Tellur. iii. 96 (1836).
Waelbergeria, Rafin. Fl. Tellur. iii. 96 (1836).
zeylanica, Wight, Ic. t. 1958.—Ceylon.

ENHYDRA:—
affinis, *Didricks. Ind. Sem. Hort. Haun.* (1849) 14.—Peruv.
Angeallia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 409.—Am. austr.
canaliculata, Cass. in Dict. Sc. Nat. xiv. 554 = fluctuans.
fluctuans, *Lour. Fl. Cochinch.* 511; *ed. Willd.* 626.—Cochinch.
Helonchæ, DC. Prod. v. 637 = paludosa.
integrifolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 410.—Bras.
longifolia, DC. Prod. v. 637 = paludosa.
maritima, *DC. l. c.*—Peruv.
maritima, Lam. in Linnæa, vi. (1831) 159 = sessilis.
oblonga, *DC. Prod.* v. 637.—Peruv.
paludosa, *DC. l. c.*—Ind. or.; Malaya; Austral.
rivularis, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 410.—Bras.
sessilis, *DC. Prod.* v. 637.—Ind. occ.
Sobreyra, *DC. l. c.*—Peruv.
subcaulescens, DC. Prod. l. c. = sessilis.
Woolfii, F. Muell. Fragm. iii. 139 = paludosa.

ENHYDRIA, Kanitz, in Mart. Fl. Bras. xii. ii. 380 (1882) = **Enydria**, Vell. = **Myriophyllum**, Ponted. (Haloragˇ.).

ENICOSANTHUM, Becc. in Nuov. Giorn. Bot. iii. (1871) 183. *ANONACEÆ*.
paradoxum, *Becc. l. c.* 184.—Borneo.

ENICOSTEMA, Blume, Bijdr. 848 (1826). *GENTIANEÆ*, Benth. & Hook. f. ii. 807.
Adenema, G. Don, Gen. Syst. iv. 201 (1837).
Henicostemma, Endl. Gen. 605 (1838).
Hiffinia, Spreng. Syst. i. 605 (1825).
Lepinema, Rafin. Fl. Tellur. iii. 96 (1836).
Slevogtia, Reich. Consp. 138 (1828).
littorale, *Blume, l. c.*—Reg. trop.

ENICOSTEMMA, Steud. Nom. ed. II. i. 555 (1840) = prsec.

ENIPEA, Rafin. Fl. Tellur. iii. 92 (1836) = **Salvia**, Tourn. (Labiat.).
formosa, Rafin. l. c. = S. leonuroides.

ENKEA, Walp. Ann. i. 603 (1846-49) = ENCKEA, Kunth = **Piper**, Linn.

ENNIANTHUS, Lour. Fl. Cochinch. 276 (1790). *ERICACEÆ*, Benth. & Hook. f. ii. 588.
Meisteria, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 137. t. 1.
Melidora, Noronha, ex Salisb. in Trans. Hort. Soc. ii. (1817) 156 partim.
Tritomodon, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 584.
biflorus, Lour. Fl. Cochinch. 277 = Rhododendron indicum ?
campanulatus, *Nichols. Dict. Gard.* i. 810.—Japon.
himalaicus, *Hook. f. & Thomt. in Hook. Kew Journ.* vii. (1855) 195.—Reg. Himal.
japonicus, *Hook. Bot. Mag.* t. 5823.—Japon.
quinqueflorus, *Lour. Fl. Cochinch.* 277.—China.
reticulatus, Lindl. in Bot. Reg. t. 885 = quinqueflorus.
uniflorus, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 489.—China.

ENKLEIA, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 284 = **Idnostoma**, Wall. (Thymel.).
malaccensis, Griff. l. c. 285 = I. scandens.
malayana, Griff. Notul. iv. 668 = prsec.

ENKYANTHUS, DC. Prod. viii. 783 (1839) = **Enkianthus**, Lour. (Ericac.).

ENKYLIA, Griff. Pl. Cantor. 26 (1887 ?); and in Journ. As. Soc. Beng. xxiii. (1854) 645 = **Gynostemma**, Blume (Cucurb.).
digyna, Griff. ll. cc. 27, 645 = G. pedatum.
trigyna, Griff. ll. cc. 27, 646 = G. pedatum.

ENKYLISTA, Benth. & Hook. f. Gen. ii. 86 (1873), sphalm. = **Eukylista**, Benth. = **Calycophyllum**, DC. (Rubiac.).

ENNEAPOGON, Desv. ex Beauv. Agrost. 81 (1812); et Journ. Bot. i. (1813) 70 = **Pappophorum**, Schreb. (Gramin.).
Desvauxii, Beauv. Agrost. 82, 161 = P. nigricans.
gracilis, Desv. Journ. Bot. i. (1813) 70 = P. nigricans.
mollis, Lehm. Pugill. iii. 40 = P. molle.
nigricans, Desv. Journ. Bot. i. (1813) 70 = P. nigricans.
pallidus, Desv. l. c. = P. nigricans.
persicus, Bolss. Diagn. Ser. I. v. 71 = P. persicum.
phleoides, Roem. & Schult. Syst. ii. 616 = P. phleoides.
purpureascens, Desv. Journ. Bot. i. (1813) 70 = P. nigricans.
scaber, Lehm. Pugill. iii. 41 = P. scabrum.

ENNEPTA, Rafin. Sylva Tellur. 52 (1838) = **Ilex**, Tourn.
atomaria, Rafin. l. c. = I. lucida.
coriacea, Rafin. l. c. = I. lucida.
myricoides, Rafin. l. c. = I. glabra.

ENODIUM, Gaudin, Agrost. Helv. i. 145 (1811) = **Molinia**, Schrank (Gramin.).
atrovirens, Dum. Obs. Gram. Belg. 108 = M. cærulea.
cæruleum, Gaudin, Agrost. Helv. i. 145 = M. cærulea.
litorale, Kunth, Enum. Pl. i. 350 = M. cærulea.
tylesticum, Link, Enum. Hort. Berol. i. 40 = M. cærulea.

ENOMEIA, Spach, Hist. Vég. Phan. x. 360 (1841) = EINOMEIA, Rafin. = **Aristolochia**, Linn.

ENOSANTHES, A. Cunn. ex Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 191 = **Homoranthus**, A. Cunn. (Myrtac.).
flavescens, A. Cunn. ex Schau. l. c. 192 = H. virgatus.
virgata, A. Cunn. ex Schau. l. c. 193 = H. virgatus.

ENOTHREA, Rafin. Fl. Tellur. iv. 48 (1836) = **Octomeria**, R. Br. (Orchid.).
graminifolia, Rafin. l. c. 44 = O. graminifolia.

ENOUREA, Aubl. Pl. Gui. i. 587 (1775) = **Paullinia**, Linn. (Sapindac.).
caprodata, Aubl. l. c. = P. connarifolia.
fagínea, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 379 = P. fagínea.

ENRILA, Blanco, Fl. Filip. ed. I. 709 (1837) = **Ventilago**, Gaertn. (Rhamn.).
dichotoma, Blanco, l. c. = V. maderaspatana.

ENSETE, Bruce, Trav. v. App. 86; ex Horan. Prod. Monog. Scitam. 40 (1862) = **Musa**, Linn. (Scitam.).
edule, Bruce, ex Horan. l. c. = M. Ensete.

ENSKIDE, Rafin. Fl. Tellur. iv. 109 (1836) = **Utricularia**, Linn. (Lentib.).
barbata, Rafin. l. c. = U. barbata.
chrysantha, Rafin. l. c. = U. chrysantha.
flava, Rafin. l. c. = U. chrysantha.

ENSLENIA, Nutt. Gen. Am. i. 164 (1818). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 757.
Ampelamus, Rafin. in Am. Monthly Mag. (1819) 192.
Enslinia, Reichb. Consp. 181 (1828).
Nematalis, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 264.
albida, *Nutt. Gen. Am.* i. 165.—Am. bor.
cinerea, Spreng. Syst. i. 850 = albida.
jamaicensis, Griseb. Fl. Brit. W. Ind. 418 = Roulinia jamaicensis.
lingulata, *Benth. Pl. Hartw.* 290.—Mexic.
racemosa, Griseb. Fl. Brit. W. Ind. 418, 785 = Roulinia jacquinii.
volubilis, *Karst. Fl. Columb.* ii. 117. t. 162.—N. Granat.

ENSLENIA, Rafin. Fl. Ludov. 85 (1817) = **Nelensia**, Poir. = **Pedicularis** ?, Tourn. (Scrophul.).
bibracteata, Rafin. l. c. (*N. bibracteata*).—Am. bor.

ENSOLENANTHE, Schott, in Bonplandia, ix. (1861) 268 = **Alocasia**, Schott (Aroid.).

ENSTOMA, A. Juss. in Orb. Dict. vi. 78 (1849) sphalm. = **Rustoma**, Salisb. (Gentian.).

ENTADA, Adans. Fam. ii. 318 (1763). *LEGUMINOSÆ*, Benth. & Hook. f. i. 589.
ADENOPODIA, Presl, Epim. Bot. 206 (1850).
GUGALORIUM, P. Br. Hist. Jamaic. 362 (1756).
PERIMA, Rafin. Sylva Tellur. 118 (1838).
STREPSILOBUS, Rafin. l. c. 117 (1838).
abyssinica, *Steud. in A. Rich. Tent. Fl. Abyss.* i. 234. —Abyss.
acaciæfolia, *Benth. in Trans. Linn. Soc.* xxx. (1875) 566.—Am. trop.
Adenanthera, DC. Mém. Lég. 422 = scandens.
africana, *Guill. & Perr. Fl. Seneg. Tent.* 233.—Afr. trop.
arborea, *Grif. Notul.* iv. 452.—Burma.
chiliantha, DC. Mem. Lég. 422; Prod. ii. 425 = polystachya.
cirrhosa, Rafin. Sylva Tellur. 119 = polystachya.
Duparquetiana, *Baill. Adansonia*, vi. (1866–66) 210. —Afr. trop.
durissima, Baill. l. c. 208 = Prosopis oblonga.
Gandu, Hoffngg. Preiss. Verz. Pfl. viii. 27 = scandens.
Gigalobium, DC. Mém. Lég. 421; Prod. ii. 424 = scandens.
ingæfolia, Presl, Epim. Bot. 205 = Piptadenia inæqualis.
Kirkii, *Oliver, Fl. Trop. Afr.* ii. 327, in nota.—Afr. trop.
monostachya, DC. Mém. Lég. 422. t. 61; Prod. ii. 425 = scandens.
myriadenia, Benth. in Hook. Journ. Bot. ii. (1840) 133 = Mimosa myriadenia.
natalensis, *Benth. l. c.* iv. (1842) 353.—Afr. austr.
Parrana, Spreng. Syst. ii. 325 = scandens.
Plumerii, Spreng. Syst iv. Car. Post. 164 = polystachya.
polyphylla, *Benth. in Hook. Journ. Bot.* ii. (1840) 133 —Guiana.
polystachya, *DC. Mém. Lég.* 422, 434. t. 61, 62; *Prod.* ii. 425.—Am. trop.
Pursætha, DC. ll. cc. 421; 425 = scandens.
Rheedii, Spreng. Syst. iii. 325 = scandens.
Rumphii, Scheff. in Tijdschr. Nederl. Ind. xxxii. (1871) 413 = scandens.
scandens, *Benth. in Hook. Journ. Bot.* iv. (1842) 332. —Am. trop.
sudanica, *Schweinf. Rel. Kotsch.* (1868) 8. t. 8.—Afr. trop.
sufruticosa, *Vatke, in Linnæa*, xliii. (1880–82) 108.— Madag.
Wahlbergia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 277. —Afr. austr.
Werhæna, Presl, Epim. Bot. 206 = Pentaclethra filamentosa.

ENTAGONUM, Polr. in Dict. Sc. Nat. xxix. 528 (1823) = ENTOGANUM, Banks = **Melloope**, Forst. (Rutac.).

ENTASIKON, Rafin. Nov. Fl. Am. iv. 98 (1836) = **Tropocarpus**, Nutt. (Umbell.).
tenuifolium, Rafin. l. c. 29 = T. Æthusa.
tuberosum, Rafin. l. c. = prasc.?

ENTATICUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 198 (1821) = **Habenaria**, Willd. (Orchid.).
albidus, S. F. Gray, l. c. 208 = Fl. albida.
viridis, S. F. Gray, l. c. 206 = H. viridis.

ENTELEA, R. Br. in Bot. Mag. t. 2480 (1824). *TILIACEÆ*, Benth. & Hook. f. i. 204.
arborescens, *R. Br. l. c.*—N. Zel.
australis, Walp. Rep. i. 353 = prasc.
palmata, *Lindl. Bot. Reg.* (1838) *Misc.* 66.—Hab.?
pubescens, Sweet, Hort. Brit. ed. III. 35, sphalm. = prasc.
sloanæoides, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 574 = arborescens.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 69.—Ins. Tahiti.

ENTEROLOBIUM, Mart. in Flora, xx. (1837) ii. Beibl. 117. *LEGUMINOSÆ*, Benth. & Hook. f. i. 596.
Blanchetii, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 234 = Pithecolobium Blanchetii.
cyclocarpum, *Griseb. Fl. Brit. W. Ind.* 236.—Jamaic.
ellipticum, *Benth. in Hook. Lond. Journ.* iii. (1844) 234.—Bras.
glaucocarpum, Mart. in Flora, xx. (1837) ii. Beibl. 117. = E. Timbouva.

ENTEROLOBIUM:—
Gomboril, Steud. Nom. ed. II. i. 555, sphalm. = E. Timbouva.
Mongollo, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 117.— Bras.
polycephala, *Griseb. in Goett. Abh.* xxiv. (1879) 123.— Reg. Argent.
Schomburgkii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 599.—Am. trop.
Tamboril, Mart. in Flora, xx. (1837) ii. Beibl. 117 = E. Timbouva.
Timbouva, *Mart. l. c.* 128.—Bras.

ENTEROPOGON, Nees, in Lindl. Introd. Nat. Syst. ed. II. 448 (1836). *GRAMINEÆ*, Benth. & Hook. f. iii. 1165.
leptophylla, *Benth. in Journ. Linn. Soc.* xix. (1881) 101.—Madag.
macrostachya, *Munro, ex Benth. l. c.*—Abyss.
mellocides, *Nees, in Lindl. Introd. Nat. Syst.* ed. II. 448, incl. syn.
Seychellarum, *Benth. in Journ. Linn. Soc.* xix. (1881) 101.—Ins. Maurit.

ENTEROSPERMUM, Hiern, in Oliver, Fl. Trop. Afr. iii. 92 (1877). *RUBIACEÆ*.
littorale, *Hiern, l. c.* 93.—Afr. trop.

ENTOGANUM, Banks, ex Gaertn. Fruct. i. 331. t. 68 (1788) = **Melloope**, Forst. (Rutac.).
laevigatum, Gaertn. l. c. = M. ternata.

ENTOSIPHON, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 44 = Parinarium, Juss. (Rosac.).
indicus, Bedd. l. c. 45 = P. indicum.

ENTRECASTEAUXIA, Montr. in Mém. Acad. Lyon, x. (1860) 244. *INCERT. SED.* Benth. & Hook. f. ii. 1124.
elliptica, *Montr. l. c.*—N. Caled.

ENULA, Neck. Elem. i. 4 (1790) = **Inula**, Linn. (Compos.).

ENUREA, J. F. Gmel. Syst. 748 (1791) = ENOUREA, Aubl. = **Paullinia**, Linn. (Sapindac.).

ENVDRA, Lour. Fl. Cochinch. 510 (1790) = **Enhydra**, Lour. (Compos.).

ENVDRIA, Vell. Fl. Flum. 56 (1825); i. t. 150 = **Myriophyllum**, Ponted. (Haloragˆ.).
aquatica, Vell. l. c. 57; i. t. 150 = M. brasiliense.

ENYMONOSPERMUM, Spreng. ex DC. Prod. iv. 244, in syn. (1830) = **Pleurospermum**, Hoffm. (Umbellif.).

EOMECON, Hance, in Journ. Bot. xxii. (1884) 346. *PAPAVERACEÆ*.
chionanthe, *Hance, l. c.*—China.

EOPEPON, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 31 = **Trichosanthes**, Linn. (Cucurb.).
anantherum, Naud. in Huber, Cat. (1870) Suppl. 6 = T. Kirilowii.
viticifolius, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 32 = T. Kirilowii.

EOPHYLON, A. Gray, in Journ. Linn. Soc. xi. (1869) 22 = Cotyzianthera, Blume (Gentian.).
Lobbii, A. Gray, l. c. 23.—Java.
tenellum, A. Gray, l. c.—Ins. Pacif.

EOPHYTON, Benth. & Hook. f. Gen. ii. 908 (1876) = prasc.

EPACRIS, Forst. Char. Gen. 19 (1776); Cav. Ic. iv. 35. t. 344, 345 (1797). *EPACRIDEÆ*, Benth. & Hook. f. ii. 615.
acuminata, *Benth. Fl. Austral.* iv. 240.—Austral.
alpina, *Hook. f. Fl. N. Zel.* i. 106.—N. Zel.
apiculata, *A. Cunn. in Field, N. S. Wales*, 340.— Austral.
attenuata, Ledd. Bot. Cab. t. 98 = Lysinema pungens.
aurieulata, Benth. in Enum. Pl. Hong. 76 = microphylla.

EPACRIS:—
autumnalis x, Paxt. Mag. Bot. xi. (1844) 195.
bicolor x, Paxt. l. c. xv. (1849) 193.
Calvertiana, *F. Muell. Fragm.* viii. 52.—Austral.
campanulata, Ledd. Bot. Cab. t. 1925 = impressa.
capitata, Polr. Encyc. Suppl. ii. 556 = Dracophyllum capitatum.
carinata, A. Cunn. ex DC. Prod. vii. 762 = rigida.
cerasifera, R. Grah. in Edinb. N. Phil. Journ. (July 1832) 168 = impressa.
corinthioides, Labill. Nov. Holl. Pl. i. 48. t. 59 = Prionotes corinthioides.
ciliata, Polr. Encyc. Suppl. ii. 555 = Lysinema ciliatum.
coccinea, *Paxt. Mag. Bot.* vi (1839) 123.—Hab.?
conspicua, Polr. Encyc. Suppl. ii. 556 = Lysinema conspicuum.
coriacea, *A. Cunn. ex DC. Prod.* vii. 763.—Austral.
corymbiflora, Hook. f. Fl. Tasm. i. 261 = myrtifolia.
crassifolia, *R. Br. Prod.* 551.—Austral.
delicata x, Paxt. Mag. Bot. xv. (1849) 194.—Austral
Cunninghami, Hort. ex Paxt. Mag. Bot. xi. (1844) 196 = *autumnalis*.
densiflora, *Hort. Angl. ex Lem. Illustr. Hortic* i (1854) t. 19.—Hab.?
diosmæfolia, *Ledd. Bot. Cab.* t. 1834.—Austral.
dubia, Lindl. Bot. Reg. (1846) t. 38 = heteronema?
exserta, *R. Br. Prod.* 551.—Tasman.
fasciculata, Forst. f. Prod. 13 = Leucopogon fasciculatus.
fimbriata, F. Muell. Fragm. viv. 125 = Lysinema fimbriatum.
Franklinii, Hook. f. Fl. Tasman. i. 261. t. 79 = mucronulata.
frondosa, Gaertn. Fruct. ii. 77 = Dracophyllum longifolium.
gracilis, Poir. Encyc. Suppl. ii. 556 = Dracophyllum gracile.
grandiflora, Hort. ex Paxt. Mag. Bot. i. (1841) 52 = seq.?
grandiflora, Willd. Sp. Pl. i. 884 = longiflora.
Gunnii, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 272 = microphylla.
heteronema, *Labill. Nov. Holl. Pl.* i. 42. t. 56.— Austral.
hirtella, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 271 = Archeria hirtella.
Hookeri, *Regel. Gartenfl.* (1856) 281.—Austral.
hyacinthiflora, *W. G. Smith, in Gard. Chron.* (1876) ii. 548. t. 105. f. 80, nomen.—Hab.?
impressa, *Labill. Nov. Holl. Pl.* i. 48. t. 58.—Austral.
juniperina, Forst. Char. Gen. 20 = Cyathodes abietina.
Kinghornii x, Hort. ex Lem. Jard. Fleur. ii. (1852) t. 175. n. 3.
lanuginosa, *Labill. Nov. Holl. Pl.* i. 42. t. 57.— Austral.
lasianthe, Polr. Encyc. Suppl. ii. 555 = Lysinema lasianthum.
latifolia, Banks & Soland. ex A. Cunn. in Ann. Nat Hist. ii. (1839) 48 = Dracophyllum latifolium.
longiflora, *Cav. Ic.* iv. 35. t. 344.—Austral.
longifolia, Banks & Soland. ex Hook. f. Fl. Antarct. 48 = Dracophyllum latifolium.
longifolia, Forst. Char. Gen. 20 = Dracophyllum longifolium.
microphylla, F. Muell. Fragm. vi. 72 = Archeria serpyllifolia.
microphylla, *R. Br. Prod.* 550.—Austral.
microphylla, Hook. f. in Hook. Lond. Journ. Bot vi. (1847) 272 = pectophylla.
minuta, Lindl. Bot. Reg. (1845) t. 5 = longiflora.
mucronulata, *R. Br. Prod.* 552.—Tasman.
mucronulata, Hook. f. Fl. Tasman. i. 258 = acuminata.
Muelleri, Sond. in Linnæa, xxvi. (1853–55) 252 = rigida.
multiflora, *Lem. Illustr. Hortic.* vii. (1860) t. 266.— Austral.
myrtifolia, *Labill. Nov. Holl. Pl.* i. 41. t. 55.— Tasman.
nivalis, Lodd. Bot. Cab. t. 1921. R. Grah. in Edinb. N. Phil. Journ. (July 1832) 168 = impressa.
nivea, DC. Prod. vii. 762 = impressa.
obtusifolia, *Sm. Exot. Bot.* i. 17. t. 40.—Austral.
odorata, Sm. ex Nouv. & Schult. Syst. iv. 362 = Lysinema pungens.
ommaneyfora, A. Cunn. in Field, N. S. Wales, 340 = purpurascens.

EPACRIS:—
paludosa, *R. Br. Prod.* 551.—Austral.
pauciflora, *A. Rich. Exp. Fl. N. Zel.* 213. t. 29.—N. Zel.
pedicellata, DC. Prod. vii. 761 = microphylla.
pentapetala, Poir. Encyc. Suppl. ii. 555 = Lysinema ciliatum.
petrophila, *Hook. f. Fl. Tasman.* i. 261.—Austral.
pulchella, *Cav. Ic.* iv. 26. t. 345.—Austral.
pulchella, Sims, Bot. Mag. t. 1170 = microphylla.
pumila, Forst. Char. Gen. 20 = Pentachondra pumila.
pungens, Cav. Ic. iv. 26. t. 346 = Lysinema pungens.
pungens, Sims, Bot. Mag. t. 644 = purpurascens.
purpurascens, *R. Br. Prod.* 550.—Austral.
purpurascens, Sieber, ex D.C. Prod. vii. 760 = pulchella.
racemosa, *Hook. f. Fl. N. Zel.* i. 166.—N. Zel.
reclinata, *A. Cunn. ex Benth. Fl. Austral.* iv. 234.—Austral.
refulgens, Hort. ex Paxt. Mag. Bot. xi. (1844) 195 = autumnale.
rigida, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 64.—Austral.
riparia, R. Br. Prod. 550 = pulchella.
riparia, Sieber, ex DC. Prod. vii. 765 = Lysinema pungens.
rivularis, Sieber, ex Spreng. Syst. iv. Cur. Post. 64 = microphylla.
robusta, *Benth. Fl. Austral.* iv. 237.—Austral.
rosea, Lodd. Bot. Cab. t. 883 = Lysinema pungens.
rosmarinifolia, Forst. f. Prod. 13 = Dracophyllum rosmarinifolium.
rubra, Benth. Fl. Austral. iv. 242, in syn. = purpurascens.
rubra, Spreng. Syst. i. 639 = Cosmelia rubra.
ruscifolia, R. Br. Prod. 550 = impressa.
ruscifolia, Sieber, ex Benth. Fl. Austral. iv. 234 = reclinata.
secunda, Poir. Encyc Suppl. ii. 556 = Dracophyllum secundum.
serpyllifolia, *R. Br. Prod.* 551.—Austral.
Sieberiana, *Hook. f. Handb. N. Zeal. Fl.* 179.—N. Zel.
sparsa, *R. Br. Prod.* 551.—Austral.
spuria, Cav. Ic. iv. 37. t. 347 = Leucopogon ericoides.
squarrosa, *Hook f. Fl. Tasman.* i. 259.—Tasman.
squarrosa, Poir. Encyc. Suppl. ii. 556 = Dracophyllum squarrosum. *R. Br.*
taxotonioniu x, Paxt. Fl. Mag. xiv. (1846) t. 195.
tomentosa, Lindl. in Mitch. Three Exped. ii. 177 = impressa.
vaccinioides, Lhotzky, ex Sond. in Lehm. Pl. Preiss. i. 235 = Pentachondra pumila.
variabilis, Lodd. Bot. Cab. t. 1816 = impressa.
villosa, Cav. Ic. iv. 37. t. 347. f. 2 = Leucopogon juniperinus.
Vilmoreana, *Planch. Hort. Donat.* 45.—Austral.
virgata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 273 = exserta.

EPACTIUM, Willd. ex Schult. Mant. iii. 15 (1827). *INC. SED.*
arenarium, *Willd. ex Schult. l. c.*—Peruv.

EPALLAGE, DC. Prod. vi. 8 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 369.
Helicta, Less. Syn. Comp. 221 (1832).
alternifolia, Steud. Nom. ed. II. i. 103 = dentata.
asemonifolia, *DC. Prod.* vi. 4.—Madag.
dentata, *DC. l. c.*—Madag.
humilis, *Baker, in Journ. Linn. Soc.* xx. (1883) 189.—Madag.
minima, *Baker, l. c.*—Madag.
rupestris, *DC. Prod.* vi. 4.—Madag.
salmifolia, DC. l. c. 3 = Sphaerophyllum madagascariense.

EPALTES, Cass. in Bull. Soc. Philom. (1816) 189; et in Dict. Sc. Nat. xv. 8 (1819). *COMPOSITAE*, Benth. & Hook. f. ii. 398.
Ethulopsis, F. Muell. Fragm. ii. 154 (1861).
Gymbanthes, Steetz, in Peters, Reise Mossamb. Bot. 457 (1864).
Litogyne, Harv Thes. Cap. ii. 55. t. 155 (1863).
Pachytheca, Steetz, in Peters, Reise Mossamb. Bot. 462 (1864).
alata, *Steetz, in Peters, Reise Mossamb. Bot.* 463.—Afr. trop.
australis, DC. Prod. v. 462 = Cunninghamii.

EPALTES:—
australis, *Less. in Linnaea*, v. (1831) 148.—Austral.
brasiliensis, *DC. Prod.* v. 461.—Bras.
Cunninghamii, *Benth. Fl. Austral.* iii. 530.—Austral.
divaricata, *Cass. in Bull. Soc. Philom.* (1818) 189.—Ind. or.
gariepina, *Steetz, in Peters, Reise Mossamb. Bot.* 451.—Afr. trop.
Harrisii, *F. Muell. Fragm.* xi. 101.—Austral.
hirsuta, *Less. Syn. Comp.* 206.—Ind. or.
itinerariiolia, DC. Prod. v. 461 = divaricata.
littoralis, *Less. in Linnaea*, vi. (1831) 151.—Ind. or.
mexicana, *Less. l. c.* v. (1830) 147.—Mexic.
pleiocephala, *F. Muell. Fragm.* x. 100.—Austral.
pygmaea, *DC. Prod.* v. 461.—Ind. or.
Tatei, *F. Muell. in Trans. Roy. Soc. S. Austral.* vi. (1883) 81.—Austral.
umbelliformis, *Steetz, in Peters, Reise Mossamb. Bot.* 452.—Afr. trop.

EPALTHES, Walp. Rep. ii. 600 (1843) · prasc.

EPATTIS, Rafin. New Fl. Am. ii. 17 (1836). *COMPOSITAE.*
oblique, *Rafin. l. c.*—Oregon.

EPERUA, Aubl. Pl. Gui. i. 369. t. 142 (1775).
LEGUMINOSAE, Benth. & Hook. f. i. 580.
Adleria, Neck. Elem. ii. 542 (1790).
Dimorpha, Schreb. Gen. 493 (1791).
Panzera, Willd. Sp. Pl. ii. 540 (1799).
Parivoa, Aubl. Pl. Gui. 756. t. 303 (1775).
Rotmannia, Neck. Elem. ii. 450 (1790).
bijuga, *Mart. ex Benth. in Mart. Fl. Bras.* xv. ii. 226.
decandra, Blanco, Fl. Filip. ed. I. 368 = Afzelia bijuga.
falcata, *Aubl. Pl. Gui. i.* 369. t. 142.—Guian.
falcata, Blanco, Fl. Filip. ed. I. 369 = Afzelia rhomboidea.
grandiflora, *Benth. & Hook. f. Gen.* i. 580; *Baill. Hist. Pl.* ii. 110.—Guian.
leucantha, *Benth. in Mart. Fl. Bras.* xv. ii. 225.—Bras.
purpurea, *Benth. l. c.* 226.—Bras.
rhomboidea, Blanco, Fl. Filip. ed. II. 260 = Afzelia rhomboidea.
rubiginosa, *Miq. Stirp. Surin. Sel.* 12.—Guia

EPETETIORHIZA, Steud. Nom. ed. II. ii. 328 (1841) = seq.

EPETORHIZA, Steud. Nom ed. II. i. 556 (1840) = **Physalis**, Linn. (Solanac.).

EPHAIOLA, Rafin. Sylva Tellur. 124 (1838) = **Aemistus**, Schott (Solanac.).
odorata, Rafin. l. c. = A. Plumieri.

EPHEBOPOGON, Nees & Meyen, ex Steud. Nom. ed. II. i. 556 (1840). *GRAMINEAE.*
gratus, *Nees & Meyen, ex Steud. l. c.*—China.

EPHEDRA, Tourn. ex Linn. Gen. ed. I. 319 (1737). *GNETACEAE*, Benth. & Hook. f. iii. 418.
Chaetocladus, Senilis, [Nelson] Pin. 161 (1866).
‘ alata, *Decne. in Ann. Sc. Nat. Sér.* I. ii. (1834) 299.—Afr. bor. ; Arab.
alata, Schimp. ex Carr. Conif. ed. I. 655 = E. Alte.
Alte, Brand. For. Fl. 501. t. 69 = foliata.
‘ Alte, *C. A. Mey. Monog. Ephedr.* 78. t. 8. f. 4.—Syria ; Arab.
altissima, Bové, in Ann. Sc. Nat. Sér. II. i. (1834) 162 = alata.
altissima, Buch, Beschr. Phys. Ins. Canar. 159, 168 ; Tommas. in Flora, xviii. (1835) ii. Beibl. 56 = fragilis.
altissima, Delile, Desc. Aegypt. 110 = E. Alte.
‘ altissima, *Desf. Fl. Atlant.* ii. 371. t. 253.—Afr. bor.
‘ americana, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 860.—Am. austr.
andina, Poepp. & Endl. Syn. Conif. 255 = americana.
antisiphylitica, S. Wats. Bot. King's Exp. 338 = nevadensis.
‘ antisyphilitica, *Berland. ex C. A. Mey. Monog. Ephedr.* 101.—Am. bor. ; Mexic.
aphylla, Forsk. Fl. Aegypt. Arab. 170 = E. Alte.
arborea, Lag. ex Bertol. Fl. Ital. x. 394 = distachya.

EPHEDRA:—
arborea, F. Muell. Syst. Arr. Pl. Carpentaria, 14 (Calycopeplí sp.).—Austral.
asperagoides, *Griff. Itin. Notes*, 340 = foliata.
‘ aspera, *Engelm. ex S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 157.—Am. bor. occ.
boryoides, Fisch. in C. A. Mey. Monog. Ephedr. 99. t. 5. f. 12 = distachya.
bractoata, Miers, Trav. Chili, ii. 531 = americana.
‘ californica, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 200.—Calif.
‘ chilensis, *Presl, Bot. Bemerk.* 109.—Chili.
campylopoda, DC. Prod. xvi. ii. 356 = fragilis.
chilensis, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xi. (1863) 253 = americana.
ciliata, Aitch. in Journ. Linn. Soc. xix. (1882) 167 = foliata.
ciliata, Fisch. & Mey. in C. A. Mey. Monog. Ephedr. 100 = E. Alte.
Clusii, Dufour, in Bull. Soc. Bot. Fr. vii. (1860) 445 = distachya, nebrodensis.
‘ dahurica, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) i. 421.—Dahuria.
dissoluta, Webb & Berth. Phyt. Canar. iii. 275 = fragilis.
distachya, Brot. Fl. Lusit. ii. 6 = fragilis.
distachya, Forsk. Fl. Aegypt. Arab. 77 = E. Alte.
distachya, *Gaud. Fl. Helv.* vi. 304 = helvetica.
‘ distachya, *Linn. Sp. Pl.* 1040.—Europ. ; As. bor.
distachya, Vill. Hist. Pl. Dauph. iii. 816 = nebrodensis.
dubia, Regel, in Act. Hort. Petrop. vi. (1879) 482 = distachya.
‘ dumosa, *Miers, Contrib. Bot.* ii. 168. t. 77 A.—Chili.
equisetiformis, Webb & Berth. Phyt. Canar. iii. 275 = nebrodensis.
‘ equisetina, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 499.—Sibiria ; China.
‘ foemina, *Forsk. Fl. Aegypt. Arab.* 219.—Arab.
‘ foliata, *Boiss. & Kotschy, ex Boiss. Diagn.* Ser. I. vii. 101.—Arab. ; Persia.
‘ fragilis, *Desf. Fl. Atlant.* ii. 372.—Reg. Mediterr ; Arab.
fragilis, Fresen. in Mus. Senckenb. i (1834) [1838] 73 ; Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 240 = E. Alte.
fragilis, Lowe, in Journ. Linn. Soc. v. (1860) 44 = altissima ?
fragilis, Moris, Stirp. Sard. Elench. ii. 8 = nebrodensis.
‘ frustillata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III xi. (1863) 262.—Patagon.
Gerardiana, Wall. Cat. n. 6048 = distachya.
‘ glubaltica, *Boiss. Fl. Orient.* v. 714, obs.—Gibraltar.
‘ glauca, *Regel, in Act. Hort. Petrop.* vi. (1879) 484.—Turkestan.
graeca, C. A. Mey. Monog. Ephedr. 93. t. 5 = nebrodensis.
‘ helvetica, *C. A. Mey. l. c.* 87. t. 8. f. 10.—Europ.
humilis, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 251 = americana.
‘ intermedia, *Schrenk & C. A. Mey. Monog. Ephedr.* 88.—As. centr. ; Himal. ; China.
kokanica, Regel, in Act. Hort. Petrop. vi. (1879) 479 = foliata.
‘ lomatolepis, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1844) 210.—Soongaria.
macrocephala, Bertol. Misc. Bot. xxiii. 17 = distachya.
major, Host, Fl. Austr. ii. 71 = nebrodensis.
major, Puller, Ins. Fl. Dalmat. 90 = fragilis.
meritima, St. Lager, Cat. Fl. Rhone = distachya.
media, C. A. Mey. Monog. Ephedr. 80 = distachya.
minor, Host, Fl. Austr. ii. 671 = distachya.
‘ monosperma, *S. G. Gmel. ex Willd. Sp. Pl.* iv. 859.—As. bor. ; China.
monostachya, Gmel. Pl. Rar. Icon. 375, nomen = nebrodensis.
monostachya, Linn. Sp. Pl. 1040 = distachya.
monostachya, Tourn. in Bull. Soc. Nat. Mosc. xxvii. (1854) i. 422 = monosperma.
monticola, Miers, Contrib. Bot. ii. 166. t. 76 A = americana.
‘ nebrodensis, *Tineo, in Guss. Fl. Sic. Syn.* ii. 688.—Reg Mediterr. ; As. bor.
‘ nevadensis, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 208.—Am. bor. occ.
occidentalis, Torr. ex Parl. in DC. Prod. xvi. 354 = antisyphilitica.
‘ ochreata, *Miers, Contrib. Bot.* ii. 169. t. 77 B.—Reg. Argent.

EPHEDRA :—
* pachyclada, *Boiss. Fl. Orient.* v 713.—Persia; Afghan.
pedunculata, Boiss. l. c. 717 = foliata.
pedunculata, Engelm. ex S. Wats. in Proc. Am. Acad. xviii. (1883) 157 = antisyphilitica.
peruviana, Bert. ex Carr. Conif. ed. I. 549 = americana.
podostylax, Boiss. Fl. Orient. v. 715 = distachya.
polygonoides, Pall. Fl. Ross. ii. 87 = distachya.
polygonoides, Siev. in Pall. *N. Nord. Beitr.* viii. 174 = monosperma.
polyphylla, Boiss. & Hausskn. ex Boiss. Fl. Orient. v. 716 = foliata.
procera, C. A. Mey. Ind. Sem. Hort. Petrop. x. (1844) 45 = nebrodensis.
rigida, St. Lager, Cat. Pl. Rhone, 687 = helvetica.
rupestris, Benth. Pl. Hartw. 258 = americana.
scandens, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xi. (1863) 361 = Tweedliana.
scoparia, Lange, in Kjoeb. Vidensk. Meddel. (1862) 33 (Fugill. Pl. Hisp. 88) = nebrodensis.
stenosperma, Schrenk & C. A. Mey. Monog. Ephedr. 77. t. 3. f. 5 = distachya.
* strobilacea, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 499 (*Lehm. Reliq. Bot.* 252).—Persia; Ass. centr.
subtristachya, C. A. Mey. Monog. Ephedr. 80 = distachya.
* Torreyana, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 296.—Am. bor. occ.
* triandra, *Tul. in Ann. Sc. Nat.* Sér. IV. x. (1858) 125. —Reg. Argent.
trifaria, Parl. in DC. Prod. xvi. 11. 359, sphalm. = seq.
* trifurca, *Torr. in Bot. Emory Rep.* (1848) 152.—Am. bor. occ.
trifurca, Parry, Bot. Obs. S. Utah, 351 = Torreyana.
* Tweediana, *C. A. Mey. Monog. Ephedr.* 96. t. 5. f. 9.—Reg. Argent.
Villarsii, Gren. & Godr. Fl. Fr. iii. 161 = nebrodensis.
vulgaris, Rich. Conif. 26 = distachya.
vulgaris, Willk. in Flora, xxxv. (1852) 319 = nebrodensis.

EPHEMERUM, Retchb. Fl. Germ. Excurs. 409 (1831) = Lysimachia, Linn. (Primulac.).
nemorum, Retchb. l. c. = L. nemorum.
Nummularia, Schur, Enum. Pl. Transs. 549 = L. Nummularia.

EPHEMERUM, Tourn. ex Moench, Meth. 387 (1794) = Tradescantia, Linn. (Commelin.).
bicolor, Moench, Meth. Suppl. 78 = Rhoeo discolor.
congestum, Moench, Meth. 388 = Tradescantia virginica.
cristatum, Moench, l. c. = Cyanotis cristata.
racemosum, Moench, l. c. = Tinantia fugax.

EPHIALIS, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. i. (1838) 461 = Vitex, Linn. (Verbenac.).
pentaphylla, Banks & Soland. ex A. Cunn. l. c. = V. littoralis, *A. Cunn.*
simplicifolia, Soland. ex Seem. in Journ. Bot. iii. (1865) 256 = Gmelina macrophylla.

EPHILIS, Banks & Soland. ex Seem. Fl. Vit. 189 (1866) = prsec.

EPHILIS, Schreb. Gen. 253 (1791) = **Ratonia**, DC. (Sapindac.).
frutinea, Willd. Sp. Pl. ii. 238.—Am. austr.
guyanensis, Pers. Syn. i. 413 = *frutinea*.
Patriciana, Spreng. Syst. ii. 253.—Guian.
Vouarana, Spreng. l. c.—Guian.

EPHIPPIANDRA, Decne. in Ann. Sc. Nat. Sér. IV. ix. (1858) 378. t. 7. *MONIMIACEÆ*, Benth. & Hook. f. iii. 141.
myrioides, *Decne. l. c.*—Madag.

EPHIPPIANTHUS, Retchb. f. in F. Schmidt, Reisen Amurl. 180. t. 5 (1868) = **Lipuris**, Rich. (Orchid.).
sachalinensis, Retchb. f. l. c.—Insc. Sachall.
Schmidtii, Retchb. f. in Flora, lii. (1866) 33 = *sachalinensis*.

EPHIPPIORHYNCHIUM, Nees, in Mart. Fl. Bras. ii. 1. 184 (1842) = **Rynchospora**, Vahl (Cyperac.).
amazonicum, Nees, ex Boeckl. in Linnæa, xxxvii. (1873–78) 626 = R. amazonica.
longirostre, Nees, in Mart. Fl. Bras. ii. 1. 136 = R. amazonica.

EPHIPPIORHYNCHIUM :—
polycephalum, Nees, in Mart. Fl. Bras. ii. 1. 184 = R. cyperoides.
stenocarpum, Nees, l. c. 136 = R. stenocarpa.
tenuirostre, Nees, l. c. 135 = R. cyperoides.
triflorum, Nees, l. c. 136 = R. triflora.
tripicatum, Nees, l. c. 136 = R. triflora.

EPHIPPIUM, Blume, Bijdr. 808. t. 65 (1825) = Cirrhopetalum, Lindl. (Orchid.).
capitatum, Blume, l. c. 809 = C. capitatum.
ciliatum, Blume, l. c. = C. Blumii.
cornutum, Blume, l. c. 810 = C. cornutum.
elongatum, Blume, l. c. 809 = C. elongatum.
grandiflorum, Blume, Rumphia, iv. 43 = C. grandiflorum !
lepidum, Blume, Bijdr. 810.—Java.
nigricans, Tetjam. & Binn. Cat. Hort. Bog. 50.—Java.
reclinatum, Tetjam. & Binn. l. c.—Java.
uniflorum, Blume, Bijdr. 809.—Java.

EPHYNES, Rafin. Sylva Tellur. 101 (1838) = **Mono chaetum**, Nand. (Melastom.).
Bonplandii, Rafin. l. c. = M. Bonplandii.

EPIADENA, Rafin. Fl. Tellur. iii. 94 (1836) = **Salvia**, Tourn. (Labiat.).
bicolor, Rafin. l. c. = S. bicolor.

EPIANDRA, Benth. & Hook. f. Gen. iii. 1066 (1883) sphalm. = seq.

EPIANDRIA, Presl, in Oken, Isis, xxi. (1828) 270 = Gaulia, Forst. (Cyperac.).
tenuiflolia, Presl, l. c. = G. tristis.

EPIBATERIUM, Forst. Char. Gen. 107. t. 54 (1776) = Cocculus, DC. (1818) (Menisperm.).
pendulum, Forst. f. l. c. 108 = C. Epibaterium.
scandens, Hausch. Nom. ed. III. 275 = C. Epibaterium.
tomentosum, Pers. Syn. ii. 561 = C. Chondodendrum.

EPIBATHERIUM, Raeusch. Nom. ed. III. 275 (1797) = prsec.

EPIBIASTRUM, Scop. Introd. 258 (1777) = prsec.

EPIBLEMA, R. Br. Prod. 315 (1810). *ORCHIDEÆ*, Benth. & Hook. f. iii. 607.
grandiflorum, *R. Br. l. c.*—Austral.

EPICAMPES, J. & C. Presl, Rel. Haenk. i. 325. t. 39 (1830). *GRAMINEÆ*, Benth. & Hook. f. iii. 1148.
Cryptsinna, Foum. ex Benth. in Journ. Linn. Soc. xix. (1881) 87.
berlandieri, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 548.—Mexic.
Bourgæi, *Fourn. ex Hemsl. l. c.*—Mexic.
Buchingeri, *Fourn. ex Hemsl. l. c.*—Mexic.
coerulea, *Griseb. in Goett. Abh.* xix. (1874) 256.—Reg. Argent.
distichophylla, *Vasey, Cat. Grass. U. St.* 45.—Am. bor.
expansa, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 548.—Mexic.
gigantea, *Fourn. ex Hemsl. l. c.*—Mexic.
gracilis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1846) 271.—Mexic.
lanata, *J. & C. Presl, Rel. Haenk.* i. 326.—Mexic.
laxiuscula, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 549.—Mexic.
macroura, *Benth. in Journ. Linn. Soc.* xix. (1881) 87. —Mexic.
pubescens, *J. & C. Presl, Rel. Haenk.* i. 326.—Mexic.
rigens, *Benth. in Journ. Linn. Soc.* xix. (1881) 88.—Mexic.
robusta, Foum. ex Hemsl. Biol. Centr. Am. Bot. iii. 550 = gracilis.
stricta, *J. & C. Presl, Rel. Haenk.* i. 325. t. 39.—Mexic.

EPICARPURA, Hassk. Cat. Hort. Bog. Alt. 74 (1844) = seq.

EPICARPURUS, Blume, Bijdr. 488 (1825) = **Strobilus**, Lour. (Urtic.).
asper, Steud. Nom. ed. II. i. 556 = S. asper!
Gaudichaudii, Steud. l. c. = S. asper!
tomentosus, Zipp. ex Span. in Linnæa, xv. (1841) 344.—Ins. Timor.

EPICARPURUS :—
microphyllus, Raoul, Choix Pl. N. Zél. 14.—N. Zel.
orientalis, Blume, Bijdr. 488 = S. asper.
spinosus, Steud. Nom. ed. II. i. 556; Wight, Ic. vi. t. 1962 = Phyllochlamys spinosa.
timorensis, Decne. Herb. Timor. 171 = Phyllochlamys spinosa.
zeylanica, Thw. in Hook. Kew Journ. iv. (1852) 1. Taxotrophis zeylanica.

EPICHARIS, Blume, Bijdr. 166 (1825). *MELIACEÆ*, Benth. & Hook. f. i. 388.
altissima, Blume, l. c. 167 = Dysoxylum densiflorum.
Baillonii, *Pierre, in Bull. Soc. Linn. Par.* i. (1881) 992 —Cambod.
Balansæana, Baill. Adansonia, xi. (1874) 257 = Dysoxylum Balansæanum.
caulifiora, Blume, Bijdr. 166 = Dysoxylum ramiflorum.
densiflora, Blume, l. c. 167 = Dysoxylum densiflorum.
exarillata, Arn. ex Wight & Arn. Prod. 120 = Dysoxylum binectariferum.
exarillata, Nimmo, in J. Grah. Cat. Pl. Bomb. t. 31 = Amoora Lawi.
Fraserana, C. DC. in DC. Monog. Phan. i. 506 = Dysoxylum Fraserianum.
Gotadhora, M. Roem. Syn. Hesper. 108 = Dysoxylum binectariferum.
juglans, *Hance, in Journ. Bot.* xv. (1877) 330.—China.
Kunthiana, A. Juss. in Mém. Mus. Par. xix. (1830) 229 = Dysoxylum Kunthianum.
Leschenaina, C. DC. in Bull. Soc. Bot. Fr. xxii. (1875) 231 = Dysoxylum Lessertianum.
Loureiri, *Pierre, in Bull. Soc. Linn. Par.* i. (1881) 291. —Cambodia.
macrocarpa, *Miq. Fl. Ind. Bat. Suppl.* 506.—Sumatra
minutiflora, Baill. Adansonia, xi. (1874) 258 = Dysoxylum minutiflorum.
molis, Wall. ex Voigt, Hort. Suburb. Calc. 138 = Dysoxylum Hamiltonii.
pachypoda, Baill. Adansonia, xi. (1874) 259 = Dysoxylum pachypodum.
Pancheri, Baill. l. c. 258 = Dysoxylum Pancheri.
rosea, Baill. l. c. 260 = Dysoxylum roseum.
sericea, Blume, Bijdr. 166 = Dysoxylum ramiflorum.
setosa, Span. in Linnæa, xv. (1841) 129 = Dysoxylum setosum.
speciosa, A. Juss. in Mém. Mus. Par. xix. (1830) 230 = Dysoxylum speciosum.

EPICHROCANTHA, Eckl. & Zeyh. ex Meisn. in DC Prod. xiv. 580 (1857) = **Gnidia**, Linn. (Thymel.).
biflora, Eckl. & Zeyh. ex Meisn. l. c. 588 = G. decurrens.
debilis, Eckl. & Zeyh. ex Meisn. l. c. = G. parvifora.
juniperifolia, Eckl. & Zeyh. ex Meisn. l. c. 587, 588 = G. juniperifolia, subulata.
ovata, Eckl. & Zeyh. ex Meisn. l. c. 586 = G. coriacea
pungens, Eckl. & Zeyh. ex Meisn. l. c. 587 = G. stylpheloides.
simplex, Eckl. & Zeyh. ex Meisn. l. c. 587, 588 = G. juniperifolia, subulata.
villosa, Eckl. & Zeyh. ex Meisn. l. c. 587 = G. Sonderiana.

EPICHYSIANTHUS, Voigt, Hort. Suburb. Calc. 533 (1845) = Chonemacrapha, G. Don (Apocyn.).
macrophyllus, Voigt, l. c. = C. macrophylla.

EPICILINASTRUM, Boj. ex DC. Prod. vii. 1. 16 (1838) = **Garbura**, Gronov. (Compos.).
Candollei, Boj. ex DC. l. c. 19 = G. Candollei.
hirsutum, Boj. ex DC. l. c. = G. Bojeri.

EPICOILA, Rafin. Sylva Tellur. 196 (1838) = **Loranthus**, Linn.
grandiflora, Rafin. l. c. = L. grandiflorus, *Ruiz & Pav.*

EPICOSTORUS, Rafin. Atl. Journ. 144 (1832) = **Hellia**, D. Don (Rosac.).
canadensis, Rafin. l. c. = N. Torreyi.

EPICRANTHES, Blume, Bijdr. 806. t. 9 (1825) = **Bulbophyllum**, Thou. (Orchid.).
barbata, Retchb. f. in Walp. Ann. vi. 225 = Monomeria barbata.
javanica, Blume, Bijdr. 806.—Java.

EPICRIANTHES, Blume, Fl. Jav. Pracf. p. vii (1828) = EPICRANTHES, Blume = Bulbophyllum, Thou. (Orchid.).

EPIDENDRON, Linn. Syst. ed. I (1735) = sec.

EPIDENDRUM, Linn. Gen. ed. I. 273 (1737). *ORCHIDEAE*, Benth. & Hook. f. iii. 528.
AMPHIGLOTTIS, Salisb. in Trans. Hort. Soc. i. (1812) 294.
ARACHNELLUM, Hoffmgg. Verz. Orch. 21; ex Linnaea, xvi. (1842) Litt. 299.
ARISTYLIS, Rafin. Neogenyt. 4 (1825).
AULIZA, Salisb. in Trans. Hort. Soc. i. (1812) 294.
BARKERIA, Knowles & Westc. Fl. Cab. iii. 7. t. 49 (1838).
CAULARTHRON, Rafin. Fl. Tellur. ii. 40 (1836).
COILOSTYLIS, Rafin. l. c. iv. 87 (1836).
DIDOTHION, Rafin. l. c. 39 (1836).
DIMERA, Lindl. Orch. Scel. 16 (1896); et Gen. et Sp. Orch. 111 (1881).
DOTHILOPHIS, Rafin. Fl. Tellur. iv. 39 (1836).
DOXOSMA, Rafin. l. c. 9 (1836).
ENCYCLIA, Hook. Bot. Mag. t. 2831, 2912 (1828).
ENOPHYA, Rafin. Fl. Tellur. ii. 62 (1836).
ETDISANTHEMA, Neck. Elem. iii. 133 (1790).
LARNANDRA, Rafin. Neogenyt. 4 (1825).
NAMOSIRA, Lindl. Bot. Reg. t. 1541 (1832).
NYCTOSMA, Rafin. Fl. Tellur. ii. 9 (1836).
ORESTEDELLA, Reichb. f. in Bot. Zeit. x. (1852) 932.
PHADROSANTHUS, Neck. Elem. iii. 133 (1790).
PHYSINGA, Lindl. Bot. Reg. (1838) t. 38.
PROSTHECHEA, Knowles & Westc. Fl. Cab. ii. 111 (1838).
PSEUDEPIDENDRUM, Reichb. f. in Bot Zeit. x. (1852) 733.
PSYCHILIS, Rafin. Fl. Tellur. iv. 40 (1836).
SULPITIA, Rafin. l. c. 37 (1836).
TRITELANDRA, Rafin. l. c. ii. 85 (1836).

acicalare, *Batem. ex Lindl. Bot. Reg.* (1841) *Misc.* 46.—Mexic.
acinaciforme, Pav. ex Lindl. Gen. et Sp. Orch. 198 = Oncidium pusciflorum.
acinatoides, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14 (Quid !).—Java.
Acklandiae, Reichb. f. in Walp. Ann. vi. 312 = Cattleya Acklandiae.
aculeatum, Poir. Encyc. Suppl. i. 376 = Eulophia aculeata.
acuminatum, *Ruiz & Pav. Syst. Veg.* 248.—Peruv.
acutissimum, *Lindl. Fol. Orch. Epid.* 46.—N. Granat.
adenocarpon, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 24.—Mexic.
adenoscalum, *La Llave, l. c.* 22.—Mexic.
adenoglossum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 36.—Peruv.
advenum, *Reichb. f. in Gard. Chron.* (1872) 1194.—Brsa.
aemulum, Lindl. Bot. Reg. t. 1896 = fragrans.
Aérides, Raeusch. Nom. ed. III. 265 = Aérides odoratum.
aéridiflolium, Reichb. f. in Walp. Ann. vi. 1169 = sec.
aéridiforme, Booth, in Lindl. Bot. Reg. (1848) Misc. 9 = Harrisonise.
aeruginosum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69 = *E. Flav-aérie.*
affine, Focke, in Bot. Zeit. xl. (1858) 841 = oncidioides.
affine, *Reichb. f. in Bonplandia*, iv. (1856) 297.—Guatemala.
affine, A. Rich. Fl. Cub. Fanerog. ii. 237 = fucatum.
agathosmioides, *Reichb. f. in Linnaea*, xxii. (1849) 841.—Venezuela.
agrarium, *Lindl. in Hook. Journ. Bot.* iii. (1841) 84.—Peruv.
aggregatum, Roxb. ex Steud. Nom. ed. II. i. 556 = Dendrobium aggregatum.
alatum, *Batem. Orch. Mexic.* t. 18.—Guatemala.
alatum, Lindl. Bot. Reg. (1847) t. 53 = ambiguum.
alatum, Lindl. in Benth. Pl. Hartw. 92 = trachyphilum.
albidum, *Vell. Fl. Flum.* ix. t. 56.—Brsa.
albo-marginatum, *Reichb. f. in Linnaea*, xli. (1877) 113.—Ecuador.
albo-purpureum, *Rodrig. Orch. Nov.* i. 51.—Brsa.
Allardii, *Regel, in Act. Hort. Petrop.* v. (1877) 998.—Mexic.
Allemagnianum, Rodrig. Orch. Nov. ii. 139 = Allemanii.

EPIDENDRUM:—
Allemanii, *Rodrig. l. c.* i. 54.—Brsa.
alnoides, Curt. Bot. Mag. t. 867 = Cymbidium pendulum.
aloifolium, Batem. Orch. Mex. & Guat. t. 25 = Parkinsonianum.
aloifolium, Linn. Sp. Pl. 953 = Cymbidium aloifolium.
alpestre, Sw. Prod. Veg. Ind. Occ. 135 = Pleurothallis alpestris.
alpicolum, *Reichb. f. in Bonplandia*, ii. (1854) 110.—N. Granat.
alternans, Lindl. in Hook. Journ. Bot. iii. (1841) 88 = fimbriatum.
altissimum, *Batem. in Bot. Reg.* (1838) *Misc.* 38.—Ins. Baham.
altissimum, Jacq. Enum. Pl. Carib. 30 = Oncidium altissimum.
altum, Poir. Encyc. Suppl. i. 374 = Bletia verecunda.
amabile, *Linden & Reichb. f. in Bonplandia*, iii. (1855) 219.—Mexic.
amabile, Linn. Sp. Pl. 953 = Phalaenopsis amabilis.
ambiguum, *Lindl. Fol. Orch. Epid.* 18.—Guatemala.
Amblostoma, Reichb. f. in Bonplandia, ii. (1854) 89 = Amblostoma cornua.
amethystinum, *Reichb. f. in Gard. Chron.* (1867) 876.—Am. trop.
amethystoglossum, Reichb. f. in Walp. Ann. vi. 319 = Cattleya guttata.
amictum, *Linden & Reichb. f. in Bonplandia*, iii. (1855) 219.—Hab. ?
ammophilum, *Rodrig. Orch. Nov.* ii. 149.—Brsa.
amplobulbum, A. Rich. Fl. Cub. Fanerog. ii. 240 = aloeps.
amplexicaule, *Lindl. Fol. Orch. Epid.* 51.—Peruv.
anceps, *Jacq. Select. Am.* 284. t. 188.—Am trop.
anceps, Sw. Prod. Veg. Ind. Occ. 121 = bifarium.
angustifolium, *Sw. l. c.* 138.—Jamaic.
angustistamum, *Lindl. Fol. Orch. Epid.* 84.—N. Granat.
anisatum, *La Llave & Lex Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 27.—Mexic.
ansiferum, *Reichb. f. in Bonplandia*, ii. (1854) 111.—Venezuela.
antennifernm, Humb. & Bonpl. Pl. Aequin. i. 98. t. 28 = Trichocerus antennifer.
antenniferum, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 16.—Mexic.
anthoceros, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 281.—N. Granat.
anthropophorum, *Reichb. f. l. c.* iv. (1856) 215.—Hab. ?
aphyllum, Poir. Encyc. Suppl. i. 371 = Dendrobium Pierardi.
apiculatum, *Vell. Fl. Flum.* ix. t. 25 (Quid !).—Brsa.
apiculatum, *Reichb. f. in Linnaea*, xli. (1877) 81.—Hab. ?
aquaticum, *Lindl. in Ann. & Mag. Nat. Hist.* xii. (1843) 393.—Brsa.
arachnoglossum, *Reichb. f. ex André, in Rev. Hortic.* (1882) 554.—N. Granat.
arachnoideum, *Rodrig. Orch. Nov.* i. 60.—Brsa.
arbusculum, *Lindl. in Benth. Pl. Hartw.* 93.—Mexic.
Aristotelia, Raeusch. Nom. ed. III. 265 = Spiranthes australis.
armeniacum, *Lindl. Bot. Reg.* t. 1867.—Brsa.; Peruv.
Arminii, *Reichb. f. in Bonplandia*, iii. (1855) 67 — Venezuela.
aromaticum, *Batem. Orch. Mex.* t. 10.—Guatemala.
articulatum, Burm. f. Fl. Ind. 189 (Quid !)—Ind. or.
articulatum, Klotzsch, in Otto & Dietr. Allg. Gartenz. vi. (1838) 397 = tessellatum.
asperum, *Lindl. in Hook. Journ. Bot.* i. (1834) 6.—Am. trop.
atropurpureum, Hensel. in Gard. Chron. (1868) 11. 42, spharicum = bifolium.
atropurpureum, *Willd. Sp. Pl.* iv. 115.—Mexic.
attenuatum, Lindl. Fol. Orch. Epid. 69 = Hookeriannm.
attenuatum, *Lindl. l. c.* 41.—Venezuela.
aurantiacum, *Batem. in Lindl. Bot. Reg.* (1838) *Misc.* 8.—Guatemala.
aureum, Lindl. Fol. Orch. Epid. 4 = aurantiacum.
Auriculs, *Vell. Fl. Flum.* ix. t. 31.—Brsa.
auritum, *Lindl. Bot. Reg.* (1848) *Misc.* 4.—Am. cont.
auropurpureum, Lindl. Gen. et Sp. Orch. 99 = bifolium.
autumnale, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 239 = Earina mucronata.
autumnale, Forst. f. Prod. 60 = Earina mucronata.

EPIDENDRUM:—
aviculæ, *Lindl. in Hook. Journ. Bot.* iii. (1841) 85.—Brsa.
Bacnlus, *Reichb. f. in Bonplandia*, iv. (1856) 214.—Hab. ?
bahamense, *Griseb. Fl. Brit. W. Ind.* 614.—Ins. Baham.
bahiense, *Reichb. f. in Hamb. Gartenz. xv.* (1859) 53.—Brsa.
Barbae, *Reichb. f. Beitr. Orch. Centr. Am.* 88.—Costa Rica.
Barkeriola, *Reichb. f. in Gard. Chron.* (1884) ii. 816—Hab. ?
Barringtoniae, Sm. Ic. Pict. 15 = Lycaste Barringtoniae.
basilare, Klotzsch, in Otto & Dietr. Allg. Gartenz. xi. (1843) 193 = Stamfordianum.
bellense, *Reichb. f. in Linnaea*, xli. (1877) 78.—Gnian.
Bettianum, *Rodrig. Orch. Nov.* ii. 147.—Brsa.
bicameratum, *Reichb. f. in Gard. Chron.* (1871) 1194.—Mexic.
bicolor, Reichb. f. in Walp. Ann. vi. 311 = Cattleya bicolor.
bicornutum, Hook. Bot. Mag. t. 3332 = Discrium bicornutum.
bidentatum, Koen. in Retz. Obs. vi. 54 = Limodorum bidentatum.
bidentatum, Lindl. Gen. et Sp. Orch. 98 = Discrium bidentatum.
bifarium, *Sw. in Schrad. Journ.* ii. (1799) 212.—Jamaic.
bifidum, *Aubl. Pl. Gut.* ii. 824.—Am. austr.
bifidum, *Lindl. ex Reichb. f. in Hamb. Gartenz.* xv. (1859) 59.—Ins. S. Doming.
bifornm, Forst. f. Prod. 60 = Dendrobium bifforum.
bifforum, *Rodrig. Orch. Nov.* i. 52.—Brsa.
bifforum, *Ruiz & Pav. Syst. Veg.* 247.—Peruv.
bifurcatum, Lindl. in Lodd. Cat. ed. II. n. 566, ex Bot. Reg. (1844) Misc. 15 = cuculiatum.
bigibberosum, Reichb. f. in Walp. Ann. vi. 346 = Discrium bigibberosum.
bilamnellatum, Reichb. f. l. c. 345 = Discrium bilamellatum.
bipaplare, *Reichb. f. in Flora*, xlviii. (1865) 277.—Ind. occ.
Bisetra, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 112 (err. typ. 105).—Guian.
bisetum, Lindl. Bot. Reg. (1841) Misc. 68 = polyanthum.
bispiculum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14 (Quid !).—Java.
bivalve, *Lindl. Fol. Orch. Epid.* 47.—Venezuela.
blepharistes, *Barker, ex Lindl. in Bot. Reg.* (1844) *Misc.* 20.—Venezuela.
blepharoclinium, *Reichb. f. in Linnaea*, xli. (1877) 87.—Ecuador.
bletioides, *Griseb. Fl. Brit. W. Ind.* 616.—Jamaic.
Boothianum, *Lindl. Bot. Reg.* (1838) *Misc.* 5.—Cuba.
bracciferum, *Reichb. f. in Linnaea*, xli. (1877) 114.—Ecuador.
brachiatum, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 20.—Mexic.
brachyphillum, *Lindl. Orch. Linden.* 9.—Venezuela.
brachyrachidum, *Lindl. Fol. Orch. Epid.* 60.—Am. austr.
brachyglossum, *Lindl. Bot. Reg.* (1844) *Misc.* 33.—Peruv.; N. Granat.
brachyphyllum, *Lindl. Fol. Orch. Epid.* 78.—Bolivia.
brachystachyum, Thou. Fl. Ins Austr. Afr. t. 84 = Acenia brachystachya.
brachystem, *Rodrig. Orch. Nov.* i. 59.—Brsa.
bractium, *Vell. Fl. Flum.* ix. t. 16.—Brsa.
bracteolatum, Presl, Rel. Haenk. i. 100 = radiatum.
bracteosum, Lindl. Bot. Reg. (1840) Misc. 58 = sciculare.
Bradfordii, *Griseb. Fl. W. Ind.* 613.—Ind. occ.
Brassavolae, *Reichb. f. in Bot. Zeit.* x. (1852) 729.—Guatemala.
brevirostrum, *Lindl. Fol. Orch. Epid.* 46.—Ecuador.
bronghtonicolas, *Griseb. Cat. Pl. Cub.* 261.—Cuba.
balbosum, *Vell. Fl. Flum.* ix. t. 11.—Brsa.
caespitosum, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 1. t. 101.—Peruv.
caespitosum, *Rodrig. Orch. Nov.* i. 60.—Brsa.
calamarium, *Lindl. Bot. Reg.* (1838) *Misc.* 86.—Brsa.
calanthum, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 111.—Peruv.
calcaratum, *Vell. Fl. Flum.* ix. t. 3.—Brsa.
calcolare, Bech.-Ham. ex D. Don, Prod. Fl. Nep. 32 = Saccolabium calceolare.

EPIDENDRUM :—
Calcolaria, Auct. = sec.
Calcolariae, Koen. in Retz. Obs. vi. 45 = Cymbidium Calcolaria.
caldenae, *Rodrig. Orch. Nov.* ii. 148.—Bras.
calligarium, *Reichb. f. in Gard. Chron.* (1869) 1110.—Mexic.
Calista, Raeusch. Nom. ed. III. 265 (= *Calista amabilis*).—Coehinch.
callilorum, Lem. Jard. Fleur. iv. (1854) t. 414, Misc. 66 = raniferum.
calocheilum, Bot. Mag. t. 3898 = slatum.
campestre, *Lindl. Bot. Reg.* (1844) *Misc.* 17.—Bras.
campylostalyx, *Reichb. f. in Bot. Zeit.* x. (1852) 780.—Am. centr.
canaliculatum, Poir. Encyc. Suppl. i. 875 = Tetramicra rigida.
canaliculatum, *Vell. Fl. Flum.* ix. t. 10.—Bras.
candidum, *Rodrig. Orch. Nov.* ii. 150.—Bras.
candidum, *Vell. Fl. Flum.* ix. t. 19.—Bras.
Candollei, *Lindl. Bot. Reg.* (1839) *Misc.* 55.—Mexic.
caninum, Burm. f. Fl. Ind. 189 (Quid ?).—Ind. or.
caprese, Linn. f. Suppl. 407 = Myrtacidium filicorae.
caracasanum, *Regel, Gartenfl.* (1859) 324.—Venezuela.
cardioglossum, *Reichb. f. in Linnaea*, xxii. (1849) 841.—Venezuela.
carinatum, *Lindl. Fol. Orch. Epid.* 61.—Mexic.
carinatum, Linn. Sp. Pl. 953 = Dendrobium carinatum.
carinatum, *Vahl, in West, Bidr. Ste-Croix*, 303.—Ins. S. Cruz.
carneum, *Lindl. Orch. Linden.* 8.—Venezuela.
carnosum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 87.—Bras.
carolinianum, Lam. Encyc. i. 182 = nocturnum.
carpophorum, *Rodrig. Orch. Nov.* ii. 148.—Bras.
carthagenense, Jacq. Enum. Pl. Carib. 30 = Oncidium carthagineae.
cartilagiuiflorum, *Reichb. f. Xenia Orch.* iii. 24.—Bolivia.
catenatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14 (= Grammatophylli sp. ?).—Java.
Catillus, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 113.—N. Granat.
caudatum, Linn. Syst. ed. X. 1246 = Brassia caudata.
cauliflorum, *Lindl. Bot. Reg.* (1838) *Misc.* 47.—Bras.
cauliflorum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14 (Quid ?).—Java.
cearense, *Rodrig. Orch. Nov.* ii. 141.—Bras.
Cebolletta, Jacq. Enum. Pl. Carib. 30 = Oncidium Ceboletta.
centradenia, Reichb. f. in Walp. Ann. vi. 1163 = centropetalum.
centropetalum, Reichb. f. in Bonplandia, iii. (1855) 68 = viridibrunneum.
centropetalum, *Reichb. f. in Bot. Zeit.* x. (1852) 782.—Venezuel.
cepiformé, Hook. Bot. Mag. t. 3765 = Candollei.
Cerasia, Raeusch. Nom. ed. III. 265 = Dendrobium Cerasia.
cerasiiles, *Lindl. Bot. Reg.* (1844) *Misc.* 91.—N. Granat.
cerreum, *H. B. & K. Nov. Gen. et Sp.* i. 353.—N. Granat.
cespitosum, *Lam. Encyc.* i. 187.—Ins. Borbon.
chaenoides, *Reichb. f. in Bonplandia*, ii. (1854) 20.—Venezuela.
chioneum, *Lindl. Bot. Reg.* (1845) *Misc.* 73.—N. Granat.
chlorinense, *Reichb. f. in Bot. Zeit.* x. (1852) 780.—Am. centr.
Chloë, *Reichb. f. in Bonplandia*. iv. (1856) 227.—Guatemala.
chlorochilum, Lindl. Bot. Reg. (1838) Misc. 25 = chlorolcucum.
chlorochilum, *Rodrig. Orch. Nov.* i. 55.—Bras.
chlorinum, Rodrig. l. c. ii. 139 = chlorochilum.
chlorolcucum, *Hook. Bot. Mag.* t. 3567.—Guian.
chlorops, *Reichb. f. in Gard. Chron.* (1880) ii. 584.—Mexic.
chondylobolbon, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 20.—Mexic.
Christi, *Reichb. f. in Linnaea*, xli. (1877) 112.—Ecuador.
Christyanum, *Reichb. f. in Gard. Chron.* (1884) ii. 86.—Bolivia.

EPIDENDRUM :—
chrysostomum, *Reichb. f. in Otto & Dietr. Allg. Gartens.* xxiv. (1856) 98.—Peruv.
cicendarium, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14 (Quid ?).—Java.
ciliare, *Linn. Syst.* ed. X. 1246; *Jacq. Enum. Pl. Carib.* 29.—Am. trop.
cingillum, *Reichb. f. in Gard. Chron.* (1889) i. 380.—Hab. ?
cinnabarinum, *Salom. ex Lindl. Gen. et Sp. Orch.* 106.—Bras.
cinnamomeum, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 19 = pterocarpum.
citrinum, Reichb. f. in Walp. Ann. vi. 817 = Cattleya citrina.
citrinum, *Rodrig. Orch. Nov.* i. 53.—Bras.
clavatum, Koen. in Retz. Obs. vi. 50 = Saccolabium clavatum.
clavatum, *Lindl. Bot. Reg.* t. 1870.—Am. trop.
clavicaulatum, Sw. Prod. Veg. Ind. Occ. 120 = Vanilla clavicalata.
Clowesii, *Batem. ex Lindl. Bot. Reg.* (1844) *Misc.* 16.—Guatemala.
Clypeolum, Forst. f. Prod. 61 = Liparis Clypeolum.
cnemidophorum, *Lindl. Fol. Orch. Epid.* 58.—Guatemala.
coccineum, Jacq. Enum. Pl. Carib. 29 = Ornithidium coccineum.
coccineum, *Reichb. f. in Bonplandia*, iii. (1855) 68.—Venezuela.
cochleariifolium, Sw. Prod. Veg. Ind. Occ. 126 = Lepanthes cochlearifolia.
cochlatum, Cart. Bot. Mag. t. 159 = fragrans.
cochlatum, *Linn. Sp. Pl.* ed. II. 1351.—Am centr. ; Ind. occ.
cochlidium, *Lindl. in Ann. Nat. Hist.* iv. (1840) 382.—Peruv.
collare, *Lindl. Bot. Reg.* (1848) *Misc.* 60.—Guatemala.
colorans, Klotzsch, in Otto & Dietr. Allg. Gartenz xix. (1851) 250 = polyanthum.
complanatum, Koen. in Retz. Obs. vi. 50 (= Sarcochilli sp.).—Zeylan.
complicatum, Griseb. Cat. Pl. Cub. 262 = scalpellilgerum.
compositum, *Vell. Fl. Flum.* ix. t. 39.—Bras.
compressum, *Griseb. Fl. Brit. W. Ind.* 617.—Ins. Trinit.
conchaechilum, *Rodrig. Orch. Nov.* i. 58.—Bras.
concinnum, Steud. Nom. ed. I. 299, sphalm. = Lepanthes concinna.
concolor, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 25.—Mexic.
concretum, *Jacq. Enum. Pl. Carib.* 30 ; *Select. Am.* 228.—Am. austr.
conivens, *Vell. Fl. Flum.* ix. t. 44.—Bras.
conopseum, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 219.—Am. bor.
conspicuum, *Lem. Illustr. Hortic.* xvi. (1869) t. 592.—Bras.
Cooperianum, *Batem. in Hook. Bot. Mag.* t. 5654.—Bras.
coordinatum, *Reichb. f. in Linnaea*, xli. (1877) 7.—N. Granat.
corallorhizon, Poir. Encyc. Suppl. i. 877 = Corallorhiza innata.
cordatum, *Ruiz & Pav. Syst. Veg.* 244.—Peruv.
cordatum, Vell. Fl. Flum. ix. t. 38 = cordifolium.
cordifolium, *Steud. Nom.* ed. II. i. 557.—Bras.
coriaceum, Parker, ex Hook. Bot. Mag. t. 5595 = coriaceum.
coriaceum, Poir. Encyc. Suppl. i. 886 = Aërides coriaceum.
corifolium. *Lindl. in Journ. Hort. Soc.* vi. (1851) 218.—Am. centr.
coriophorum, *Reichb. f. in Walp. Ann.* vi. 847.—Ecuador.
cornigerum, Sw. Prod. Veg. Ind. Occ. 123 = Pleurothallis emarginata.
cornutum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 86.—Peruv.
cornutum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14 (Quid ?).—Java.
coronatum, *Ruiz & Pav. Syst. Veg.* 243.—Peruv.
corymbosum, *Lindl. Fol. Orch. Epid.* 61.—Bras.
corymbosum, Ruiz & Pav. l. c. 246 = umbelliforum.
costaricense, *Reichb. f. in Bot. Zeit.* x. (1852) 927.—Costa Rica.
costatum, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 21.—Mexic.

EPIDENDRUM :—
Coxianum, *Reichb. f. in Gard. Chron.* (1877) ii. 388.—Am. trop.
cranifolium, Endl. Gen. et Sp. Orch. 107 = ellipticum.
crassilabium, Poepp. & Endl. Nov. Gen. et Sp. ii. t. 102 = variegatum.
crenatum, *Vell. Fl. Flum.* ix. t. 6.—Bras.
crinlferum, *Reichb. f. in Gard. Chron.* (1871) 1291.—Costa Rica.
crispatum, Forst. f. Prod. 60 = Dendrobium crispatum.
crispatum, Knowles & Weste. Fl. Cab. ii. (1838) 79 = adenocarpum.
cristum, Lam. Encyc. i. 186 = Oncidium carthagineae
cristatum, *Ruiz & Pav. Syst. Veg.* 243.—Peruv.
crithmifolium, *Voigt, in Syll. Ratisb.* ii. (1828) 52.—Hab. ?
croceum, *Ruiz & Pav. Syst. Veg.* 246.—Peruv.
cubense, *Lindl. Bot. Reg.* (1843) *Misc.* 17.—Cuba.
cucullatum, *Lindl. l. c.* (1838) *Misc.* 33.—Bras.
cucullatum, Linn. Sp. Pl. ed. II. 1350 = brassavola cucullata.
cuspidatum, Lodd. Bot. Cab. t. 10 = ciliare.
cyclotellum, *Reichb. f. in Gard. Chron.* (1880) i. 72.—Mexic.
cycnostachys, Benth. in Journ. Linn. Soc. xviii. (1881) 314 = Stamfordianum.
cycnostalix, Reichb. f. in Bot. Zeit. x. (1852) 781 = Stamfordianum.
cylindraceum, *Lindl. Bot. Reg.* (1844) *Misc.* 86.—N Granat.
cylindrostachys, *Reichb f. & Warsc. in Bonplandia*, ii. (1854) 110.—N. Granat.
decipiens, *Lindl. Fol. Orch. Epid.* 70.—N. Granat, Guiana.
delicatissimum, *Reichb. f. in Linnaea*, xli. (1877) 36—Am. centr.
Daudrobii, *Reichb. f. l. c.* xvii. (1849) 841.—Venezuela.
dendrobioides, *Thunb. Pl. Bras. Dec.* ii. 17.—Bras.
densiflorum, *Hook. Bot. Mag.* t. 3791.—Mexic ; Venezuela.
denticalatum, *Reichb. f. Orch. Nov.* ii. 148.—Bras.
dichotomum, Lindl. Bot. Reg. (1838) Misc. 79 = corymbosum.
dichotomum, Paxt. ex Steud. Nom. ed. II. i. 557 = Paxtonianum.
dichotomum, *Presl, Rel. Haenk.* i. 101.—Peruv.
dichromum, *Lindl. Bot. Reg.* (1848) *Misc.* 78.—Bras.
difforme, *Jacq. Enum. Pl. Carib.* 29 ; *Select. Am.* 231 t. 186.—Am. trop.
difturum, Sw. Prod. Veg. Ind. Occ. 121 = Seraphyta multiflora.
dilotum, *Lindl. Bot. Reg.* (1848) *Misc.* 65.—Guatemala
dipterum, Sieber, ex Rïdl. in Journ. Linn. Soc. xxi. (1886) 498 = Cryptopus elatus.
dipus, Lindl. Bot. Reg. (1845) t. 4 = nutans.
discoidale, *Lindl. Fol. Orch. Epid.* 54.—Ins. S. Doming
discolor, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 22 = nocturnum.
distantiflorum, *A. Rich. & Gal. l. c.* 19. t. 7.—Mexic
distichum, Forst. f. Prod. 92, nomen (Quid ?).—Ins Societ.
distichum, Lam. Encyc i. 189 = Oberonia brevifolia.
diurnum, Poir. Encyc. Suppl. i. 875 = Cymbidium diurnum.
dolosum, Reichb. f. Xenia Orch. ii. 224 = Cattleya dolosa.
domesticum, Linn. Sp. Pl. 953 = Vanilla aromatica.
Dubois-anum, Bronge. ex A. Rich. Fl. Cub. Fanerog. ii. 239 = atropurpureum.
Dummanitu, Buch.-Ham. ex Lindl. Gen. et Sp. Orch. 90 = Dendrobium densiflorum.
Durum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 87.—Bras.
eburneum, *Reichb. f. in Gard. Chron.* (1867) 404.—Panama.
echinocarpum, Sw. Prod. Veg. Ind. Occ. 124 = Dichaea echinocarpa.
elatius, Reichb. f. in Walp. Ann. vi. 819 = Cattleya guttata.
elegans, *Reichb. f. l. c.* 874.—Mexic.
elegans, *Vell. Fl. Flum.* ix. t. 13.—Bras.
ellipticum, *R. Grah. in Hook. Exot. Fl.* iii. t. 207 ; *et in Edinb. N. Phil. Journ.* i. (1826) 171.—Bras.
elongatum, Jacq. Coll. iii. 269 ; *Ic. Pl. Rar.* iii. 17. t. 605 = secundum.
emarginatum, *Ruiz & Pav. Syst. Veg.* 247.—Peruv.
Endresii, *Reichb. f. in Gard. Chron.* (1883) i. 423.—Costa Rica.

EPIDENDRUM :—
ensatum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 22.—Mexic.
ensicaulon, A. Rich. & Gal. l. c. 19 = venosum.
ensifolium, Linn. Sp. Pl. 954 = Cymbidium ensifolium.
ensiforme, *Vell. Fl. Flum.* ix. t. 2.—Bras.
equitans, Forst. f. Prod. 60 = Oberonia glandulosa.
equitans, *Lindl. Bot. Reg.* (1838) *Misc.* 44.—Mexic.
equitans, *Ruiz & Pav. Syst. Veg.* 249.—Peruv.
erubescens, *Lindl. in Hook. Journ. Bot.* iii. (1841) 87. —Mexic.
esculentum, Hort. Kew. ex Lindl. Fol. Orch. Epid. 15 = aciculare.
exostum, *Reichb. f. in Linnæa*, xli. (1877) 79.—Bras.
erectum, *Hook. f. Bot. Mag.* t. 5902.—N. Granat.
Evelynæ, *Reichb. f. Xenia Orch.* iii. 23.—Bolivia.
exasperatum, *Reichb. f. Beitr. Orch. Centr. Am.* 87.— Costa Rica.
excisum, *Lindl. Bot. Reg.* (1844) *Misc.* 86.—N. Granat
expansum, *Reichb. f. in Linnæa*, xli. (1877) 115.—N. Granat.
falcatum, *Lindl. in Ann. Nat. Hist.* iv. (1840) 382.— Mexic.
fallax, *Lindl. Orch. Linden.* 9.—N. Granat.
falsiloquum, *Reichb. f. in Gard. Chron.* (1885) i 565. —Hab. ?
Fascicula, Forst. f. Prod. 60 = Vanilla Fasciola.
fastigiatum, *Lindl. Fol. Orch. Epid.* 56.—N. Granat.
fastigiatum, Lindl. l. c. 71 = quitensium.
favorita, *Reichb. f. in Gard. Chron.* (1874) ii. 98.— Mexic.
ferrugineum, *Ruiz & Pav. Syst. Veg.* 245.—Peruv.
filicaule, *Lindl. Gen. et Sp. Orch.* 101.—Bras.
filiforme, Sw. Prod. Veg. Ind. Occ. 128 = Acianthus fliformis.
Fimbria, *Reichb. f. in Bonplandia*, ii. (1854) 282.— Mexic
fimbriatum, *H. B. & K. Nov. Gen. et Sp.* i. 351.—N. Granat.
fimbriatum, *Vell. Fl. Flum.* ix. t. 4.—Bras.
firmum, *Reichb. f. Beitr. Orch. Centr. Am.* 87.—Costa Rica.
flabellatum, *Lindl. Fol. Orch. Epid.* 6.—Mexic.
flabelliforme, Sw. Prod. Veg. Ind. Occ. 128 = Zygopetalum cochleare.
Flabellum-Veneris, Koen. in Retz. Obs. vi. 57 (Quid ?). —Ind. or.
flavidum, Lindl. Gen. Linden. 8 = leucochilum.
flavo-virens, *Regel, Ind. Sem. Hort. Petrop.* (1855) 19 ; *et in Linnæa*, xxviii. (1856) 371.—Bras.
flavo-virens, *Reichb. f. Beitr. Orch. Centr. Am.* 85.— Guatemala.
flavum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 88.— Bras.
flexuosum, G. F. W. Mey. Prim. Fl. Esseq. 260 = Imatophyllum.
floribundum, *H. B. & K. Nov. Gen. et Sp.* i. 353. t. 96.—Am. austr.
floridum, *Vell. Fl. Flum.* ix. t. 22.—Bras.
florilinguum, *Rodrig. Orch. Nov.* i. 57.—Bras.
Flos-aëris, Koen. in Retz. Obs. vi. 64 = Aërides lasiopetala.
Flos-aëris, Linn Sp. Pl. 952 = Arachnanthe moschifera.
Forbesii, Reichb. f. in Walp. Ann. vi. 316 = Cattleya Forbesii.
formosum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 201 = alatum.
fractiflexum, *Rodrig. Orch. Nov.* ii. 151.—Bras.
fragrans, *Sw. Prod. Veg. Ind. Occ.* 123.—Am. trop.
Friderici-Guilielmi, *Warsc. ex Reichb. f. in Bonplandia*, ii. (1854) 110.—Peruv.
frigidum, *Linden, ex Lindl. Bot. Reg.* (1845) *Misc.* 76. —Venezuela.
Fruticicaseum, Norouha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 15 (Quid ?).—Java.
Frutex, *Reichb. f. Xenia Orch.* i. 96. t. 97.—Peruv.
fruticosum, *Pœp. ex Lindl. Gen. et Sp. Orch.* 101.— Mexic.
fuscatum, *Lindl. Bot. Reg.* (1838) *Misc.* 15.—Cuba.
Fuchsii, Kegel, Ind. Sem. Hort. Turic. (1850) ; ex Schweiz. Zeitschr. Gartenb. ix. (1851) 302 = Skinneri.
fulgens, A. Braun. in Dupurr. Voy. Coq. Bot. 196. t. 45 = Schomburgkii.
Fusula, Sw. Prod. Veg. Ind. Occ. 126 = Dendrophylax funalis.
Funckianum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 21.—Mexic.

EPIDENDRUM :—
funiformum, Morr. in Ann. Soc. Gand, iv. (1848) 211. t. 196 = polyanthum.
Funkii, *Reichb. f. in Linnæa*, xxii. (1849) 889.—N. Granat.
furvum, Linn. Sp. Pl. ed. II. 1346 = Vanda furva.
fuscatum, Sm. Spicil. 24. t. 23 = anceps.
fusiforme, *Reichb. f. in Walp. Ann.* vi. 292.—Hab. ?
galeatum, *Vell. Fl. Flum.* ix. t. 3.—Bras.
Galeola, Raeusch. Nom. ed. III. 265 = Cranichis nudifolia.
Galeottianum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 21.—Mexic.
gallopavinum, *Reichb. f. in Bonplandia*, iii. (1855) 219.—Bras.
gastropodium, *Reichb. f. in Walp. Ann.* vi. 405.—N. Granat.
geminiflorum, *H. B. & K. Nov. Gen. et Sp.* i. 354.— N. Granat.
geniculatum, *Rodrig. Orch Nov.* ii. 146.—Bras.
Gheiselhuijtianum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 19.—Mexic.
gigantæum, Hort. ex Planch. Hort. Donat. 164 = gravidum !
gigantæum, Poir. Encyc. Suppl. i. 376 (= *Cyrtopera gigantea*).—Afr. austr.
Gigas, *Vell. Fl. Flum.* ix. t. 20.—Bras.
gilvum, *Vell. l. c.* t. 23.—Bras.
Girondianum, *Reichb. f. in Bonplandia*, iv. (1856) 327.—Peruv.
gladiatum, Lindl. Gen. et Sp. Orch. 106 = anisatum.
glanciopetalum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14. (Quid ?).—Java.
glaucum, *Lindl. Bot. Reg.* (1840) *Misc.* 29.—Mexic.
glaucum, Sw. Prod. Veg. Ind. Occ. 134 = Dichæa glauca.
globosum, *Jacq. Enum. Pl. Carib.* 29 ; *Select. Am.* 292. t. 183.—Am. trop.
glossaspis, *Reichb. f. in Linnæa*, xli. (1877) 87.— Ecuador.
glossoceras, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 110.—Peru.
glumaceum, *Lindl. Bot. Reg.* (1839) *Misc.* 88.—Bras.
glumibracteum, *Reichb. f. in Hamb. Gartenz.* xix. (1881) 1.—Costa Rica.
glutinosum, Schedw. in Otto & Dietr. Allg. Gartenz. xi. (1843) 110 = odoratissimum.
Goncalvi, *Reichb. f. Orch. Nov.* ii. 152.—Bras.
gracile, *Lindl. Bot. Reg.* t. 1763.—Ins. Baham.
gracilicaule, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 111.—Peruv.
gracillimum, *Reichb. f. & Warsc. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 314.—Peruv.
Grahami, Hook. Bot. Mag. t. 3885 = phœnicsenm.
graminifolium, *Lindl. Gen. et Sp. Orch.* 108.—Peruv.
graminifolium, Linn. Sp. Pl. ed. II. 1353 = Octomeria graminifolia.
graminifolium, Pav. ex Lindl. Gen. et Sp. Orch. 114 = Porta lanceolata.
graminoides, Sw. Prod. Veg. Ind. Occ. 125 = Dichæa graminoides.
grammatoglossum, *Reichb. f. in Linnæa*, xxii. (1849) 887 = Venezuela.
grandiflorum, Humb. & Bonpl. Pl. Aequin. i. 94. t. 27 = Stanhopea Bucephalus.
grandiflorum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 86.—Peruv.
grandiflorum, Poir. Encyc. Suppl. 377 (= *Cleistes Guianensis*).—Guiana.
granuliferum, Lindl. in Hook. Journ. Bot. iii. (1841) 88 = cucullioides.
graveolens, Reichb. f. in Walp. Ann. vi. 319 = Cattleya granulosa.
gratiosum, *Reichb. f. in Bonplandia*, iv. (1856) 215. —Am. trop.
gravidum, *Lindl. in Journ. Hort. Soc.* iv. (1849) 114. —Mexic.
guatemalense, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 350 = cucullioides.
Guillemianum, Lindl. ex Planch. Hort. Donat. 164 = gravidum !
guttatum, Linn. Sp. Pl. 953 = Ocidium luridum.
guttatum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 20.—Mexic.
haematochilum, *Reichb. f. in Flora*, xliii. (1865) 277. —Ind. occ.
Hamkeanum, *Presl, Rel. Haenk.* i. 100.—Peruv.
Hanburii, *Lindl. Bot. Reg.* (1844) *Misc.* 46—Mexic.

EPIDENDRUM :—
Harrisoniæ, *Hook. Bot. Mag.* t. 3209.—Bras.
Harrisonianum, Reichb. f. in Walp. Ann. vi. 317 = Cattleya Loddigesii.
Hartwegii, *Lindl. in Benth. Pl. Hartw.* 150.—Peruv.
hastatum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 82.— N. Granat.
havanense, *Ledd. ex W. Bast. in Loud. Hort. Brit. Suppl.* iii. 544, nomen.—Cuba.
Helleri, Fenzl, ex Hemsl. in Gard. Chron. (1882) ii. 304 = gracile.
Hemiscieria, Reichb. f. in Walp. Ann. vi. 333 (= *Hemiscieria nutans*).—Am. trop.
Hensch.nii, *Rodrig. Orch. Nov.* ii. 147.—Bras.
herbaceum, *Ledd. ex W. Bast. in Loud. Hort. Brit Suppl.* iii. 544, nomen.—Bras.
heterodoxum, *Reichb. f. in Bonplandia*, ii. (1854) 20. —Venezuela.
hexadactylum, *Rodrig. Orch. Nov.* i. 56.—Bras.
hexandrum, Koen. in Retz. Obs. vi. 45 (Appendicular sp.).—Ind. or.
Hilchmm, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 32 = Rynchostylis retusa.
hircinum, A. Rich. Fl. Cub. Fanerog. ii. 236 = fucatum.
hirsutum, Poir. Encyc. Suppl. i. 367 = Elleanthus Caravata.
hololeucum, *Rodrig. Orch Nov.* ii. 145.—Bras.
Hookerianum, *Reichb. f. in Linnæa*, xli. (1877) 87.— N. Granat.
Huegelianum, Reichb. f. in Walp. Ann. vi. 31 = Cattleya Skinneri.
Humboldtii, Reichb. f. in Linnæa, xxii. (1849) 886 = Schomburgkia Humboldtii.
humile, Sm. Exot. Bot. ii. 75 = Coelogyne humilis.
humile, *Vell. Fl. Flum.* ix. t. 24.—Bras.
hyemale, Poir. Encyc. Suppl. i. 375 = Aplectrum hyemale.
hymenodes, *Lindl. Fol. Orch. Epid.* 58.—N. Granat.
ibaguense, Lindl. Gen. et Sp. Orch. 107 = ybaguense.
imatophyllum, *Lindl. l. c.* 106.—Guian.
imbricatum, *Lam. Encyc.* i. 169.—Guian.
imbricatum, *Lindl. Gen. et Sp. Orch.* 110.—Bras.
incisum, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 112.—Peruv.
incisum, *Vell. Fl. Flum.* ix. t. 18.—Bras.
incompitum, *Reichb. f. in Linnæa*, xxv. (1852) 735.— Am. centr.
incumbens, Lindl. Bot. Reg. (1840) Misc. 45 = aromaticum.
indicum, Poir. Encyc. Suppl. i. 384 = Rynchostylis retusa.
indivisum, Bradf. ex Griseb. Fl. Brit. W. Ind. 614 (Diarii sp.).—Ins. Trinit.
indusiatum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxii. (1854) 177 = lacertinum.
infaustum, *Reichb. f. in Hamb. Gartenz.* xix. (1863) 18. —Bras.
inflexum, *Vell. Fl. Flum.* ix. t. 20.—Bras.
insectiferum, *Lindl. Fol. Orch. Epid.* 57.—Bolivia.
insidiosum, *Reichb. f. in Linnæa*, xli. (1877) 80.— Hab. ?
intermedium, Reichb. f. in Walp. Ann. vi. 318 = Cattleya intermedia.
inversum, *Lindl. Bot. Reg.* (1839) *Misc.* 85.—Bras.
ionocentrum, *Reichb. f. in Gard. Chron.* (1888). ii. 8. —Am. trop.
ionophelebium, *Reichb. f. Beitr. Orch. Centr. Am.* 108. —Costa Rica.
ionosmum, *Lindl. Bot. Reg.* (1838) *Misc.* 49.—Guiana.
Isabelia, Reichb. f. Walp. Ann. vi. 318 = Cattleya Isabella.
isochilum, *Reichb. f. in Bonplandia*, iv. (1856) 326.— Ind. occ.
jajense, *Reichb. f. l. c.* (1854) 20.—Venezuela.
jamaicense, *Lindl. Fol. Orch. Epid.* 82.—Jamaica.
Jamiesonis, *Reichb. f. in Bonplandia*, iv. (1856) 327. —Ecuador.
jejunum, *Reichb. f. Otia Bot. Hamb.* 12.—Ecuador.
Jenischianum, *Reichb. f. in Fl. des Serres*, Sér. I. ix. (1853–54) 98 ; *et in Bonplandia*, ii. (1854) 89.— Bras.
juncifolium, Linn. Sp. Pl. ed. II. 1351 = Oncidium Cebolleta.
Karstenii, *Reichb. f. in Linnæa*, xli. (1877) 37.— N. Granat.
Karwinskii, *Reichb. f. in Bonplandia*, iv. (1856) 327. —Mexic.
kermesinum, *Lindl. Fol. Orch. Epid.* 31.—Venezuela.

EPIDENDRUM:—
Klotzscheanum, *Reichb. f. in Linnaea*, xxii. (1849) 886. —Venezuela.
Kraenzlinii, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 116.—Porto Rico.
Krugii, *Bello, l. c.* 117.—Guiana.
labiatum, Reichb. f. in Walp. Ann. vi. 313 = Cattleya labiata.
labiatum, Sw. Prod. Veg. Ind. Occ. 124 = Leiochilus cochlearis.
labiosum, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 113 (err. typ. 105).—Cuba.
lacerinum, *Lindl. Bot. Reg.* (1841) *Misc.* 58.—Guatemala.
lacerum, *Lindl. l. c.* (1836) *Misc.* 17.—Cuba.
lactiflorum, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 32 = falcatum.
lacustre, *Lindl. Fol. Orch. Epid.* 50.—Venezuela.
laeve, Lindl. Bot. Reg. (1844) Misc. 24 = paniculatum.
lagotis, *Reichb. f. in Bonplandia*, iii. (1855) 68.—Venezuela.
Lambda, *Linden, ex Reichb. f. l. c.* ii. (1854) 281.—N. Granat.
lamellatum, Westc. ex Lindl. Bot. Reg. (1843) Misc. 46 = stenopetalum.
lamproculon, *Reichb. f. in Bot. Zeit.* x. (1852) 782.—Mexic.
Lancoia, Sw. Prod. Veg. Ind. Occ 123 = Pleurothallis Lancoia.
lancolatum, *Bradf. ex Griseb. Fl. Brit. W. Ind.* 616. —Ins. Trinit.
lancifolium, *Pav. ex Lindl. Gen. et Sp. Orch.* 98.—Mexic
lanipes, *Lindl. Fol. Orch. Epid.* 91.—Peruv.
Lansbergii, Regel, Ind. Sem. Hort. Petrop. (1855) 19 —polyanthum.
Lansbergii, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 316.—Venezuela.
lasiopetalum, Poir. Encyc. Suppl. i. 384 = Aëridos lasiopetala.
latifolium, Reichb. f in Walp. Ann. vi. 848, sphalm. = falcatum.
latilabrum, *Lindl. Bot. Reg.* (1841) *Misc.* 77.—Am. trop.
laxum, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 2.—Peruv.
laxum, Sw. Prod. Veg. Ind. Occ. 125 = Pleurothallis laxa.
Lechleri, *Reichb. f. in Linnaea*, xli. (1877) 58.—Peruv.
ledifolium, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 21.—Mexic.
Lehmanni *Reichb. f. Otia Bot. Hamb.* 11.—Ecuador.
leiobolbon, Hook. Journ. Bot. iii. (1841) 308. t. 10 = varicosum.
Leochilus, *Reichb. f. in Flora*, xlviii. (1865) 277.—Ind. occ.
leoparvinum, *Reichb. f. in Linnaea*, xli. (1877) 112. —Ecuador!
leucochilum, Lindl. Orch. Linden. 8 = lacustre.
leucochilum, *Link, Klotzsch & Otto, in Otto & Dietr. Allg. Gartenz.* xi. (1843) 145.—N. Granat.
leucomelanum, *Reichb. f. in Linnaea*, xli. (1877) 58. —Mexic.
ligneosum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 28.—Mexic.
ligulatum, La Llave, l. c. 26.—Mexic.
ligulatum, Vell. Fl. Flum. ix. t. 15 = Oncidium pumilum.
liliacinum, *Linden, in Illustr. Hortic.* xxviii. (1881) 96, nomen.—N. Granat.
Liliastrum, Salisb. ex Lindl. Gen. et Sp. Orch. 177 = Sobralia Liliastrum.
lilliflorum, Koen. in Retz. Obs. vi. 61 (Quid?).—Ind. or.
Lima, *Lindl. Fol. Orch. Epid.* 67.—Peruv.
limbatum, *Lindl. Bot. Reg.* (1843) *Misc.* 69.—Guatemala.
Lindbergii, *Reichb. f. Otia Bot. Hamb.* 91.—Bras.
Lindenianum, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 20 = auritum.
Lindenii, *Lindl. Bot. Reg.* (1845) *Misc.* 48.—Venezuela.
Lindenii, Lindl. in Ann. & Nat. Hist. xii. (1843) 397 = fallax.
Lindleyanum, *Reichb. f. in Walp. Ann.* vi. 875.—Costa Rica.
lineare, Blanco, Fl. Filip. ed. I. 444 = Cleisostoma tenuemum.
lineare, Jacq. Enum. Pl. Carib. 29 = Isochilus linearis.
linearis, *Ruiz & Pav. Syst. Veg.* 249.—Peruv.
linearifolium, Hook. Bot. Mag. t. 4572 = acicnlare.

EPIDENDRUM:—
linearilingue, *Reichb. f. in Linnaea*, xli. (1877) 114.—N. Granat.
lineatum, Salisb. Prod. 10 = fragrans.
lineatum, *Vell. Fl. Flum.* ix. t. 36.—Bras.
Linkianum, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* vi. (1838) 209.—Mexic.
lividum, Lindl. Bot. Reg. (1838) Misc. 51 = tessellatum.
Llavei, Steud. Nom. ed. II. i. 558 = ligulatum.
Loddigesii, Reichb. f. in Walp. Ann. vi. 316 = Cattleya Loddigesii.
longibulbe, *Ledd. ex W. Bant. in Loud. Hort. Brit. Suppl.* iii. 544, nomen.—Guiana.
longicolle, *Lindl. Bot. Reg.* (1838) *Misc.* 84.—Guiana.
longifolium, *Rodrig. Orch. Nov.* ii. 132.—Bras.
longiflorum, *H. B. & K. Nov. Gen. et Sp.* i. 354.—N. Granat.
longiflorum, Koen. in Retz. Obs. vi. 55 (Quid?).—Ind. or.
longihastatum, *Rodrig. Orch. Nov.* i. 59.—Bras.
longipes, *Reichb. f. Otia Bot. Hamb.* 10.—Ecuador.
longipetalum, Lindl. & Paxt. Flow. Gard. i. (1850–51) 149. t. 30 = ciliatum.
longipetalum, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 29. t. 17 = antennifernm.
longispathum, *Rodrig. Orch. Nov.* i. 56.—Bras.
longovarium, *Rodrig. l. c.* 57.—Bras.
Lunaeanum, A. Rich. ex Lindl. Fol. Orch. Epid. 23 = Isochilus.
luteolum, Reichb. f. in Walp. Ann. vi. 313 = Cattleya luteola.
luteo-virens, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 19.—Mexic.
lutetum, Poir. Encyc. Suppl. i. 374 = Chloraea crispa.
lycopodioides, Koen. in Retz. Obs. vi. 55 = Cymbidium lycopodioides.
macropodium, *Reichb. f. in Bonplandia*, iv. (1856) 327. —Peruv.
macrostachya, Thun. Fl. Iles Austr. Afr. t. 88 = Aëonia macrostachya.
macrostachyum, *Lindl. Bot. Reg.* (1845) *Misc.* 70.—N. Granat.
macrothyrsoides, *Reichb. f. in Linnaea*, xli. (1877) 113. —N. Granat.
maculatum, *Anbl. Pl. Gui.* ii. 825.—Guiana.
maculatum, *Ruiz & Pav. Syst. Veg.* 248.—Peruv.
Magnolion, Mnhl. Cat. 81 = conopseum.
maicum, *Lindl. Bot. Reg.* (1844) *Misc* 83.—Peruv.
marginatum, Link, Klotzsch, & Otto, Ic. Pl. Rar. 90. t. 86 = radiatum.
marginatum, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 113 (err. typ. 105).—Guiana.
marmoratum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 31.—Mexic.
Martianum, *Lindl. in Ann. Nat. Hist.* iv. (1840) 382. —Bras.
Matheusii, *Reichb. f. in Linnaea*, xli. (1877) 81.—Peruv.
matutinum, Poir. Encyc. Suppl. i. 384 = Aëridos matutinum.
matutinum, *Reichb. f. in Linnaea*, xxii. (1849) 840.—Venezuela.
maximum, Reichb. f. in Walp. Ann. vi. 316 = Cattleya maxima.
maysifolium, Lindl. Fol. Orch. Epid. 92, sphalm. = maculosum.
megagastrium, *Lindl. l. c.* 49.—Peruv.
megalanthum, *Rodrig. Orch. Nov.* i. 51.—Bras.
megalopsathum, *Reichb. f. in Linnaea*, xli. (1877) 114. —N. Granat.
melanoculon, *Reichb. f. in Walp. Ann.* vi. 875.—Mexic.
mellosmum, *Reichb. f. in Gard. Chron.* (1869) 980.—Mexic.
mesocircon, *Lindl. Fol. Orch. Epid.* 51.—Bolivia.
michuacanum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 26.—Mexic.

EPIDENDRUM:—
micranthum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 88.—Peruv.
micranthum, Sw. Prod. Veg. Ind. Occ. 125 = Stelis micrantha.
microbulbon, *Hook. Ic. Pl.* t. 847.—Mexic.
microcharis, *Reichb. f. in Gard. Chron.* (1870) 1246.—Guatemala.
microdendron, *Reichb. f. Beitr. Orch. Centr. Am.* 84.—Costa Rica.
micron, *Vell. Fl. Flum.* ix. t. 41.—Bras.
microphyllum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 85.—Guiana.
microptus, *Reichb. f. in Hamb. Gartenz.* xix. (1863) 18. —Hab.!
microtos, *Reichb. f. in Bonplandia*, iv. (1856) 326.—Peruv.
Miersii, *Lindl. Fol. Orch. Epid.* 59.—Bras.
minimum, *Anbl. Pl. Gui.* ii. 825.—Guiana.
minutum, Anbl. l. c. 824 = Polystachya luteola.
miserrimum, *Reichb. f. in Bonplandia*, iii. (1855) 230. —Ind. occ.
misorum, Lindl. Bot. Reg. (1841) Misc. 87 = Hormidium misorum.
modestum, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 110.—Peruv.
molle, *Reichb. f. in Linnaea*, xli. (1877) 84.—Bras.
monanthum, Steud. Nom. ed. II. i. 558 = Hormidium uniflorum.
monile, Thunb. Fl. Jap. 30 = Dendrobium catenatum.
moniliforme, Linn. Sp. Pl. 954 = Dendrobium moniliforme.
monophyllum, Hook. Exot. Fl. ii. t. 109 = Pleurothallis emarginata.
monophyllum, *Vell. Fl. Flum.* ix. t. 14.—Bras.
Monroeanum, *Ledd. ex W. Bant. in Loud. Hort. Brit Suppl.* iii. 544, nomen.—Guatemala.
montanum, Sw. Prod. Veg. Ind. Occ. 121 = Octadesmia serratifolia.
Moritzii, *Reichb. f. in Linnaea*, xxii. (1849) 837.—Bras.
moschatum, [Buch.-Ham. in] Symes, Embassy, ed. II. iii. 315 = Dendrobium Calceolaria.
Mossei, *Reichb. f. in Gard. Chron.* (1880) ii. 390.—Hab.!
mucronatum, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 239 = Earina mucronata.
multibulbe, Norcsha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 15 (Quid?).—Java.
muricatum, Poir. Encyc. Suppl. i. 368 = Dichaea muricata.
muricatum, *Reichb. f. in Bonplandia*, vi. (1858) 237.—Peruv.
muscifrons, Lindl. in Hook. Journ. Bot. i. (1834) 6 = sacerpe.
myodes, *Reichb. f. Beitr. Orch. Centr. Am.* 86.—Costa Rica.
Myosurus, Forst. f. Prod. 80 = Oberonia Myosurus.
myrianthum, *Lindl. Fol. Orch. Epid.* 59.—Guatemala.
naevosum, *Lindl. l. c.* 4.—Mexic.
nanum, *Reichb. f. in Hamb. Gartenz.* xix. (1868) 281. —Cuba.
nanum, *Reichb. f. in Bonplandia*, ii. (1854) 20.—Hab.!
necatocaulon, *A. Rich. Fl. Cub. Fanerog.* ii. 398.—Cuba.
nemorale, *Lindl. in Hook. Journ. Bot.* iii. (1841) 82. —Mexic.
nervosum, *Lem. Encyc.* i. 186.—Ins. Amboina.
nervosum, Thunb. in Trans. Linn. Soc. ii. (1794) 327 = Liparis nervosa.
nitens, *Reichb. f. Beitr. Orch. Centr. Am.* 89.—Guatemala.
nocturnum, *Jacq. Enum. Carib.* 29; *Select. Am.* 235. t. 128.—Am. trop.
nodosum, Linn. Sp. Pl. 953 = Brassavola nodosa.
nonchinense, *Reichb. f. in Walp. Ann.* vi. 854.—Costa Rica.
novo-granatense, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 111.—N. Granat.
nubium, *Reichb. f. Beitr. Orch. Centr. Am.* 81.—Guatemala.
nudum, *Kunth, in Nov. Gen. et Sp.* i. 352.—Ind. or.
nudum, Ruiz & Pav. Syst. Veg. 245 = Epidendrum.
nutans, *Sw. Prod. Veg. Ind. Occ.* 121.—Jamaica.
oblongatum, A. Rich. Fl. Cub. Fanerog. ii. 399 = phoeniceum.
obtusifolium, *Willd. Sp. Pl.* iv. 113.—Ind. occ.

EPIDENDRUM:—
obtusum, *A. DC.* 9ᵐᵉ *Not. Gard. Genève*. 17.—Mexic.
ochraceum, *Lindl. Bot. Reg.* (1838) t. 36. *Misc.* 14.—Mexic.
ochranthum, A. Rich. Fl. Cub. Fanerog. ii. 387 = virens.
ochriodes, *Lindl. Fol. Orch. Epid.* 59.—N. Granat.
ochrochlorum, *Rodrig. Orch. Nov.* ii. 140.—Bras.
octandrum, *Vell. Fl. Flum.* ix. t. 21.—Bras.
odontorhizum, Poir. Encyc. Suppl. i. 377 = Corallorrhiza odontorhiza.
odontospathum, *Rchb. f. Xenia Orch.* iii. 23.—Bolivia.
odoratissimum, *Lindl. Bot. Reg.* t. 1415.—Bras.
odoratum, Poir. Encyc. Suppl. i. 385 = Aërides odoratum.
Oerstedii, *Reichb. f. in Bot. Zeit.* x. (1852) 937.—Costa Rica.
ollare, *Vell. Fl. Flum.* ix. t. 5.—Bras.
oncidioides, *Lindl. Bot. Reg.* t. 1638.—Guian.
ophioglossoides, Jacq. Enum. Pl. Carib. 29 = Stelis ophioglossoides.
Ophrydis, *Koen. in Retz. Obs.* vi. 46.—Ind. or.
Ophrydis, *Rœusch. Nom.* ed. III. 263, nomen.—Jamaica.
oppositifolium, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 21 = Hexisea oppositifolia.
orchidium, Koen. in Retz. Obs. vi. 48 (= *Ceratochilus orchideus*).—Ind. or.
orchidiflorum, *Salzm. ex Lindl. Gen. et Sp. Orch.* 108.—Bras.
oreonastes, *Reichb. f. Xenia Orch.* iii. 22.—Bolivia.
orgyale, *Lindl. in Ann. & Mag. Nat. Hist.* xv. (1845) 256.—N. Granat.
ornatum, Lem. in Fl. des Serres, Sér. I. iv. (1848) 384 b = floribundum.
osmanthum, *Rodrig. Orch. Nov.* ii. 134.—Bras.
Ottonis, *Reichb. f. in Hamb. Gartenz.* xiv. (1858) 213.—Venezuela.
ovale, Swr. Prod. Veg. Ind. Occ. 125 = Lepanthes concinna.
ovalifolium, *Lindl. Gen. et Sp. Orch.* 104.—Mexic.
ovatum, Linn. Sp. Pl. 953 = Dendrobium chlorops.
ovulum, *Lindl. Bot. Reg.* (1848) *Misc.* 55.—Mexic.
oxypetalum, *Lindl. Orch. Linden.* 8.—Cuba.
pachyanthum, *Lindl. Bot. Reg.* (1838) *Misc.* 31.—Guian.
pachyspalum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 374 = variegatum.
palaceum, Relchb. f. Beitr. Orch. Centr. Am. 80 = auritum.
pallens, *Reichb. f. l. c.* 82.—Costa Rica.
pallidiflorum, *Hook. Bot. Mag.* t. 2980.—Ins. S. Doming.
pallidum, *Court. in Mag. Hort.* i. (1833) 256.—Hab.?
palmarum, Salzm. ex Lindl. Gen. et Sp. Orch. 486 = Vanilla palmarum.
palmifolium, Swr. Prod. Veg. Ind. Occ. 123 = Maxillaria palmifolia.
palpigerum, *Reichb. f. in Gard. Chron.* (1879) ii. 40.—Mexic.
pamplonense, *Reichb. f. in Linnæa*, xxii. (1849) 837.—N. Granat.
panchryum, *Reichb. f.* & *Warsc. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 313.—Peruv.
paniculatum, Retzius. f. in Walp. Ann. vi. 1166 = Ionopsis paniculata.
paniculatum, *Ruiz & Pav. Syst. Veg.* 248.—Am. austr.
paniculicum, *Rodrig. Orch. Nov.* i. 58.—Bras.
panthera, *Reichb. f. in Bonplandia*, iv. (1856) 326.—Mexic.
Papilio, *Vell. Fl. Flum.* ix. t. 28.—Bras.
papilionaceum, Lam. Encyc. i. 179 (Quid?).—Japon.
papilionaceum, Vahl, in West, Bidr. Ste-Croix, 303 = bifidum.
papilionum, Batem. in Lindl. Bot. Reg. (1838) Misc. 7 = adenocaryum.
parahybanense, *Rodrig. Orch. Nov.* ii. 141.—Bras.
parallelum, *Lindl. Fol. Orch. Epid.* 24.—N. Granat.
parasiticum, *Rodrig. Orch. Nov.* ii. 139.—Bras.
parenthicum, *Reichb. f. in Bot. Zeit.* x. (1852) 782.—Guatemala.
Parkinsonianum, *Hook. Bot. Mag.* t. 3778.—Mexic.
parviflorum, *Ruiz & Pav. Syst. Veg.* 245.—Peruv.
parvilabre, *Lindl. in South. Pl. Hartw.* 180.—Peruv.
pasculi, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 23.—Mexic.
pastoris, Link & Otto, Ic. Pl. Rar. t. 12 = Linklauum.
patens, *Sw. Fl. Ind. Occ.* iii. 1496.—Ind. occ.; Guatemala.
pauciflorum, *Rodrig. Orch. Nov.* i. 49.—Bras.
pauper, *Vell. Fl. Flum.* ix. t. 18.—Bras.
Pavonianum, *Reichb. f. in Bonplandia*, iv. (1856) 215.—Hab.?
Paxtonianum, *Strud. Nom.* ed. II. i. 558.—Bras.
paysenue, *Reichb. f. in Bonplandia*, iii. (1855) 220.—Peruv.
pedicellatum, Poir. Encyc. Suppl. i. 376 = Eulophia scutellata.
pendulum, Roxb. Pl. Corom. i. 55. t. 44 = Cymbidium pendulum.
pentadactylum, *Reichb. f. in Bonplandia*, ii. (1854) 89.—Peruv.
pentotis, *Reichb. f. in Linnæa*, xli. (1877) 81.—Bras.
Peperomiæ, *Reichb. f. in Bonplandia*, ii. (1854) 20 = Venezuela.
pergameneum, *Reichb. f. Beitr. Orch. Centr. Am.* 86.—Costa Rica.
Philippii, *Reichb. f. in Linnæa*, xxii. (1849) 840.—Reg. Andin.
phœnicerum, *Lindl. Bot. Reg.* (1841) *Misc.* 57.—Cuba.
phyllocharis, *Reichb. f. Xenia Orch.* iii. 11. t. 208.—Costa Rica.
physodes, *Reichb. f. in Bot. Zeit.* x. (1852) 781.—Mexic.
physodes, *Reichb. f. in Gard. Chron.* (1873) 289.—Costa Rica.
pictum, *Lindl. Bot. Reg.* (1838) *Misc.* 32.—Guian.
pilestum, *Reichb. f. in Bonplandia*, iii. (1855) 68.—Venezuela.
pilliferum, *Reichb. f. in Linnæa*, xli. (1877) 83.—Panama.
piperinum, Lindl. in Ann. & Mag. Nat. Hist. xv. (1845) 256.—N. Granat.
Pipio, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 98.—Bras.
pium, *Reichb. f.* & *Warm. in Otia Bot. Hamb.* 92.—Bras.
plantaginifolium, Koen. in Retz. Obs. vi. 60 = Cymbidium plantaginifolium.
platyglossum, *Reichb. f. in Linnæa*, xli. (1877) 6.—N. Granat.
platyotis, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 318.—Venezuela.
platystigma, *Reichb. f. Beitr. Orch. Centr. Am.* 88.—Costa Rica.
plicatum, *Lindl. Bot. Reg.* (1847) t. 35.—Cuba.
polyanthemum, Reichb. f. in Walp. Ann. vi. 1166, sphalm. = fucatum.
polyanthum, Hort. Gall. ex Lindl. Fol. Orch. Epid. 13 = fucatum.
polyanthum, *Lindl. Gen. et Sp. Orch.* 106.—Mexic.
polybulbon, *Sw. Prod. Veg. Ind. Occ.* 124.—Mexic.; Ind. occ.
polyclonium, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 332.—Cuba.
polyphyllum, *Vell. Fl. Flum.* ix. t. 17.—Bras.
polystachyum, Thos. Fl. Iles Austr. Afr. t. 82 = Aeonia polystachya.
polystachyum, *H. B.* & *K. Nov. Gen. et Sp.* i. 352.—Hab.?
polystachyum, Pav. ex Lindl. Gen. et Sp. Orch. 106 = polyanthum.
Porpax, *Reichb. f. in Bonplandia*, iii. (1855) 220.—Nicaragua.
Porpax, *Reichb. f. in Flora*, xlviii. (1865),378.—Ind. occ.
porphyreum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 56.—N. Granat.
porphyroglossum, Reichb. f. Xenia Orch. ii. 33 = Cattleya porphyroglossa.
porphyroxiphium, *Reichb. f. in Linnæa*, xli. (1877) 80.—Hab.?
praecox, Sm. Exot. Bot. ii. 73 = Coelogyne praecox.
praemorsum, Retz. Pl. Corom. i. 24. t. 43 (Saccolabii sp.).—Ind. or.
praetervisum, *Reichb. f. in Linnæa*, xli. (1877) 82.—Venezuela.
pratense, *Reichb. f. Beitr. Orch. Centr. Am.* 84.—Guatemala.
primuliflorum, Batem. ex Lindl. & Paxt. Flow. Gard. i. (1850–51) 151 = rufum.
prismatocarpum, *Reichb. f. in Bot. Zeit.* x. (1852) 799.—Am. centr.
proliferum, Swr. Prod. Veg. Ind. Occ. 124 = Ponera prolifera.
prolifer, *Rodrig. Orch. Nov.* i. 61.—Bras.
propinquum, *A. Rich.* & *Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 21.—Mexic.
prolinosum, *A. Rich. & Gal. l. c.* 20.—Mexic.
Pseudepidendrum, *Reichb. f. Xenia Orch.* i. 160. t. 53.—Panama.
pterocarpum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 83.—Mexic.
pugionatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14. (Quid?).—Java.
pulchellum, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 21 = Hormidium pulchellum.
pulchellum, Swr. Prod. Veg. Ind. Occ. 125 = Lepanthes pulchella.
pulcherrimum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxii. (1854) 283 = cochlidium.
punctatum, Linn. Syst. ed. X. 1246 = Cyrtopodium punctatum.
punctatum, Reichb. f. in Walp. Ann. vi. 1166, sphalm. = tripunctatum.
punctiferum, *Reichb. f. in Gard. Chron.* (1881) ii. 88.—Bras.
punctulatum, *Reichb. f. l. c.* (1885) ii. 70.—Mexic.
puniceum, Reichb. f. in Walp. Ann. vi. 923 = Alamania punicea.
purpurachylum, *Rodrig. Orch. Nov.* i. 49.—Bras.
purpurascens, Focke, in Tijdschr. Nat. Wetensch. iv. (1851) 64 = clavatum.
purpureum, *Rodrig. Orch. Nov.* i. 59.—Bras.
purum, *Lindl. Bot. Reg.* (1844) *Misc.* 75.—Venezuela; N. Granat.
pusillum, Koen. in Retz. Obs. vi. 49 (Sarcochili sp.).—Ind. or.
pusillum, Linn. Sp. Pl. ed. II. 1352 = Oncidium iridifolium.
pygmæum, Hook. Journ. Bot. i. (1834) 49 = Hormidium uniflorum.
pyriforme, *Lindl. Bot. Reg.* (1847) sub t. 10.—Cuba.
quadrangulare, *Reichb. f.* in Otto & Dietr. Allg. Gartenz. xviii. (1850) 402 = variegatum.
quadratum, *Lindl. Fol. Orch. Epid.* 33.—Am. trop.
quitensium, *Reichb. f. in Walp. Ann.* vi. 992.—Peruv.
racemiflorum, Sw. Prod. Veg. Ind. Occ. 125 = Pleurothallis obliquifolia.
racemosum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 15. (Quid?).—Java.
radiatum, Hoffmgg. Verz. Orch.—Cf. Bot. Zeit. i. (1848) 333 = difforme?
radiatum, *Lindl. Bot. Reg.* (1841) *Misc.* 58.—Mexic.
radicans, *Pav. ex Lindl. Gen. et Sp. Orch.* 104.—Mexic.
ramonense, *Reichb. f. Beitr. Orch. Centr. Am.* 81.—Costa Rica.
ramosum, *Jacq. Enum. Pl. Carib.* 29; *Select. Am.* 221. t. 152.—Am. trop.
rasiferum, *Lindl. Gen. et Sp. Orch.* 109.—Mexic.
raphidophorum, *Lindl. Fol. Orch. Epid.* 52.—Venezuela.
recurvatum, *Lindl. Bot. Reg.* (1845) *Misc.* 77.—Venezuela.
refractum, *Lindl. in Ann. & Mag. Nat. Hist.* xii. (1843) 298.—Venezuela.
Renanthera, Rœusch. Nom. ed. III. 265 = Renanthera coccinea.
replicatum, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 167.—N. Granat.
resectum, *Reichb. f. in Linnæa*, xli. (1877) 82.—Panama.
resupinatum, Forst. f. Prod. 61 = Microstylis Rheedii.
retusum, Linn. Sp. Pl. 953 = Rhynchostylis retusa.
revolutum, *Rodrig. Orch. Nov.* i. 55.—Bras.
rhaphidophorum, Reichb. f. in Walp. Ann. vi. 1167, sphalm. = raphidophorum.
rhizomaniacum, *Reichb. f. Otia Bot. Hamb.* 12.—Ecuador.
rhizophorum, Batem. in Lindl. Bot. Reg. (1838) Misc. 8 = radicans.
rhynchophorum, *A. Rich.* & *Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 20.—Mexic.
rigidum, *Jacq. Enum. Pl. Carib.* 29; *Select. Am.* 222. t. 154.—Am. trop.
rigidum, Lodd. Bot. Cab. t. 1600 = ramosum.
rivulare, *Lindl. in Ann.* & *Mag. Nat. Hist.* Ser. III. i. (1858) 330.—Cuba.

EPIDENDRUM:—
rostrigerum, *Reichb. f. in Linnaea*, xli. (1877) 58.—Peruv.
rubroviolaceum, Lindl. Bot. Reg. (1843) Misc. 9 = densiflorum.
rubrum, *Lem. Encyc.* i. 178.—Ins. S. Doming.
Ruckeanum, *Reichb. f. in Hamb. Gartens.* xxl. (1865) 336.—Hab.?
rufum, *Lindl. Bot. Reg.* (1845) *Misc.* 33.—Ins. Baham.
Ruizianum, *Steud. Nom.* ed. II. i. 558.—Peruv.
rupestre, *Lindl in Hook. Journ. Bot.* iii. (1841) 84.—Peruv.
ruscifolium, Jacq. Enum. Pl. Carib. 29 = Pleurothallis ruscifolia.
saaronicum, Koen. in Retz. Obs. vi. 58 = Aërides matutinum.
Sagraeanum, A. Rich. Fl. Cub. Fanerog. ii. 235 = fuscatum.
sanguinarum, Sw. Prod. Veg. Ind. Occ. 124 = Brougtonia sanguinea.
sarcochilum, *Lindl. & Reichb. f. in Bonplandia*, ii. (1854) 281.—N. Granat.
sarcodes, *Lindl. Fol. Orch. Epid.* 45.—N. Granat.
sarcophyllum, Focke, in Tijdschr. Nat. Wetensch. iv. (1851) 67 = microphyllum.
sarcostalix, *Reichb. f. & Warsc. in Bonplandia*, ii (1854) 111.—Peruv.
satyrioides, Sw. Prod. Veg. Ind. Occ. 123 = Ionopsis testiculata.
Saururus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14. (Quid?).—Java.
saxatile, *Lindl. in Hook. Journ. Bot.* iii. (1841) 84.—Bras.
scabrum, *Ruiz & Pav. Syst. Veg* 248.—Peruv.
scalpelligerum, *Reichb. f. in Flora*, xlviii. (1865) 278.—Ind. occ.
Sceptrum, *Lindl. Orch. Linden.* 8.—Venezuela.
Schilleriannum, Reichb. f. in Walp. Ann. vi. 318 = Cattleya Schilleriana.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 333 = Eulophia Schimperiana.
Schlimii, Lindl. Fol. Orch. Epid. 74 = sociorense.
Schlimii, *Reichb. f. in Linnaea*, xxii. (1849) 838.—Venezuela.
Schomburgkii, *Lindl. Bot. Reg.* (1838) *Misc.* 15.—Guian.
Schreineri, *Rodrig. Orch. Nov.* ii. 142.—Bras.
sclerocladium, *Lindl. ex Reichb. f. in Bonplandia*, iv. (1856) 336.—Peruv.
scopulorum, *Reichb. f. Xenia Orch.* iii. 38.—Bolivia.
scriptum, Linn. Sp. Pl. ed. II. 1351 = Grammatophyllum speciosum.
scriptum, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 22.—Mexic.
sculptum, *Reichb. f. in Bonplandia*, ii. (1854) 89.—Panama.
Scutella, *Lindl. Bot. Reg.* (1844) *Misc.* 83.—Peruv.
secundum, *Vacq. Enum. Pl. Carib.* 29; *Select. Am.* 284. t. 187.—Am. trop.
secundum, Sw. Bot. 325 = anceps.
selligerum, *Batem. ex Lindl. in Bot. Reg.* (1838) *Misc.* 46.—Mexic.; Guatemala.
Sellowii, *Reichb. f. in Linnaea*, xxii. (1849) 839.—Bras.
seriatum, *Lindl. Fol. Orch. Epid.* 59.—Mexic.
serpens, *Lindl. in Benth. Pl. Hartw.* 149.—Peruv.
Serronianum, *Rodrig. Orch. Nov.* i. 50.—Bras.
serrulatum, Sw. *Prod. Veg. Ind. Occ.* 121.—Ind. occ.
serrulatioides, Sw. l. c. 122 = Pleurothallis serrulatioides.
sessile, Koen. in Retz. Obs. vi. 60 = Bulbophyllum clandestinum.
sessile, *Sw. Prod. Veg. Ind. Occ.* 122.—Ind. occ.
setiferum, *Lindl. in Ann. Nat. Hist.* iv. (1840) 388.—Bras.
simplex, *Spreng. Syst.* iii. 736.—Ins. Mascar.
sinense, Andr. Bot. Rep. iii. t. 216 = Cymbidium sinense.
Sintenisii, *Reichb. f. in Bot. Deutsch. Bot. Ges.* iii. (1885) 277.—Porto Rico.
sinuosum, *Lindl. Fol. Orch. Epid.* 46.—N. Granat.
sisyrinchifolium, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 19.—Mexic.
Skinneri, *Batem. in Bot. Reg.* t. 1851.—Guatemala.
smaragdinum, *Lindl. Bot. Reg.* (1838) *Misc.* 32.—Guian.; Bras.
soccorrense, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 90, in note.—N. Granat.

EPIDENDRUM:—
Sophronitis, *Lindl. & Reichb. f. in Gard. Chron.* (1857) 853.—Ecuador.
Soratae, *Reichb. f. Xenia Orch.* iii. 24.—Bolivia.
Spatella, *Reichb. f. in Bonplandia*, iv. (1856) 214.—Hab.?
spathaceum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 85.—Peruv.
spathulatum, Linn. Sp. Pl. 953 = Vanda spathulata.
spectabile, Focke, in Bot. Zeit. xi. (1853) 229 = cuniculatum.
spectabile, *Reichb. f. in Walp. Ann. vi.* 375.—Mexic.; Guatemala.
spectatissimum, *Reichb. f. in Bonplandia*, iii. (1855) 68.—Venezuela.
spherostachyum, *Reichb. f. in Linnaea*, xli. (1877) 115.—Ecuador.
spiculatum, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 180.—Ins. Galapag.
spinescens, *Lindl. Fol. Orch. Epid.* 72.—Bras.
spondylanthes, *Reichb. f. in Bot. Zeit.* x. (1852) 781.—Costa Rica.
Spruceanum, Lindl. Fol. Orch. Epid. 80 = nocturnum.
squalidum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 24.—Mexic.
squamatum, Poir. Encyc. Suppl. i. 376 = Dipodium squamatum.
squamatum, *Rodrig. Orch. Nov.* ii. 134.—Bras.
Stamfordianum, *Batem. Orch. Mex.* t. 11.—Mexic.
Stangeanum, *Reichb. f. in Gard. Chron.* (1881) i. 462.—Panama.
stellatum, *Lindl. Fol. Orch. Epid.* 21.—N. Granat.
stenopetalum, *Hook. Bot. Mag.* t. 3410.—Jamaic.
stenopetalum, *Knowles & Westc. Fl. Cab.* ii. (1838) 178.—Mexic.
sterile, Lam. Encyc. i. 189 (Bulbophyll sp.).—Ind. or.
stramineum, *Lindl. Fol. Orch. Epid.* 38.—N. Granat
stratiotes, Thunb. in Trans. Linn. Soc. ii. (1796) 327 = Bletia hyacinthina.
strobiliferum, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 363.—Ind. occ.
Strophinx, *Reichb. f. in Linnaea*, xli. (1877) 78.—Guatemala.
subadnatum, *Reichb. f. l. c.* 6.—N. Granat.
subaequale, *Eggers, Fl. Ste-Croix*, 113.—Ins. S. Thom.
subequilinum, Reichb. f. in Walp. Ann. vi. 1167, nom. nud.—Mexic.
subequilinum, *Lindl. Bot. Reg.* (1846) sub t. 64.—Mexic.
subpurpureum, *Reichb. f. in Bonplandia*, ii. (1854) 21.—Venezuela.
subulatifolium, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 21.—Mexic.
subulatum, Koen. in Retz. Obs. vi. 51 = Aërides cylindricum?
subulatum, Sw. Prod. Veg. Ind. Occ. 123 (= Brassavolae sp. ?).—Ind. occ.
subumbellatum, Hoffmgg. in Linnaea, xvi. (1858-55) 232 = difforme.
sulphuroleucum, *Rodrig. Orch. Nov.* i. 56.—Bras.
superbum, Reichb. f. Xenia Orch. ii. 26 = Cattleya violacea (in Addend.).
syringaeiflorum, Warsc. ex Reichb. f. in Bonplandia, ii. (1854) 110 = laeve.
syringothyrsis, *Reichb. f. ex Hook. f. Bot. Mag.* t. 6145.—Bolivia.
tabulare, Poir. Encyc. Suppl. i. 376 = Cymbidium tabulare.
tampense, *Lindl. Bot. Reg.* (1847) sub t. 35.—Florida.
tenax, *Reichb. f. in Bonplandia*, ii. (1854) 20.—Venezuela.
tenellum, Lindl. Fol. Orch. Epid. 44 = Schlimii.
tenue, *Lindl. in Hook. Journ. Bot.* iii. (1841) 86.—Bras.
tenuiflorum, Hort. ex Lindl. Fol. Orch. Epid. 86 = difforme.
tenuifolium, Linn. Sp. Pl. 953 = Cymbidium tenuifolium.
teres, *Reichb. f. in Bonplandia*, iii. (1855) 230.—Mexic.
teres, Thunb. Fl. Jap. 30 = Luisia teres.
tentaculiferum, *Sw. Prod. Veg. Ind. Occ.* 122.—Ind. occ.
tenuatum, *Vell. Fl. Flum.* ix. t. 48.—Bras.
terrestre, Linn. Syst. ed. X. 1348 = tuberosum.
tessellatum, *Batem. in Lindl. Bot. Reg.* (1838) *Misc.* 7.—Guatemala.

EPIDENDRUM:—
tessellatum, Roxb. Pl. Corom. i. 34 = Vanda Roxburghii.
tesselatum, Steud. Nom. ed. II. i. 558 = Vanda Roxburghii.
tessulatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. (Quid?).—Java.
testaefolium, Sw. Prod. Veg. Ind. Occ. 122 = Pleurothallis testaefolia.
tetraceros, *Reichb. f. in Bot. Zeit.* x. (1852) 782.—Ancentr.
tetragonum, Thou. Fl. Ins Austr. Afr. t. 84 = Phajus tetragonus.
tetrapetalum, Jacq. Enum. Pl. Carib. 30 = Oncidium tetrapetalum.
tetrapetalum, *Vell. Fl. Flum.* ix. t. 29.—Bras.
Thrixspermum, Raeusch Nom. ed. III. 265 (= *Thrixspermum centipeda*).—Cochinch.
thrombodes, *Reichb. f. in Linnaea*, xli. (1877) 79.—Peruv.
tibicinis, Batem. in Lindl. Bot. Reg. (1838) Misc. 8 = Schomburgkia tibicinis.
tigrinum, *Linden, ex Lindl. Orch. Linden.* 9.—Venezuela.
tipaticolum, *Lindl. Fol. Orch. Epid.* 32.—Venezuela
tolimense, *Lindl. Bot. Reg.* (1845) *Misc.* 66.—N. Granat.
tomentosum, Koen. in Retz. Obs. vi. 53 (= Erias sp.) —Ind. or.
torquatum, *Lindl. in Benth. Pl. Hartw.* 149.—N Granat.
tovarense, *Reichb. f. in Linnaea*, xxii. (1849) 838.—Venezuela; N. Granat.
trachycarpum, *Lindl. in Benth. Bot. Voy. Sulph.* 172.—Mexic.
trachychilum, *Lindl. Fol. Orch. Epid.* 19.—Mexic.
triangulatum, Griseb. Cat. Pl. Cub. 262 = trianguliferum.
trianguliferum, *Reichb. f. in Flora*, xlviii. (1865) 277.—Ind. occ.
tribuloides, Sw. Prod. Veg. Ind. Occ. 123 = Pleurothallis tribuloides.
trichocarpum, Sw. l. c. 124 = Dichaea trichocarpa.
tricrure, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 113.—Peruv.
tridactylum, Lindl. Bot. Reg. (1838) Misc. 46 = Amblostoma cernuum.
tridens, *Lodd. ex W. Baxt. in Loud. Hort. Brit Suppl.* iii. 304.—Guian.
trident, Poepp. & Endl. Nov. Gen. et Sp. ii. 2. t. 103 nocturnum.
trideutatum, Sw. Prod. Veg. Ind. Occ. 125 = Lepanthes tridentata.
triflorum, *Ruiz & Pav. Syst. Veg.* 248.—Peruv.
trigonifolium, Sw. Prod. Veg. Ind. Occ. 122 = Stelis ophioglossoides.
trilabiatum, *Vell. Fl. Flum.* ix. t. 42.—Bras.
Trinitatis, *Lindl. Bot. Reg.* (1840) *Misc.* 60.—Ins. Trinit.
tripartitum, *Vell. Fl. Flum.* ix. t. 7.—Bras.
tripterum, Lindl. in Hook. Journ. Bot. iii. (1841) 83 = Linkianum.
tripterum, Sm. Ic. Pict. 14. t. 14 = Coelia Bauerana.
tripunctatum, *Lindl. Bot. Reg.* (1841) *Misc.* 66.—Mexic.
tripunctatum, Lindl. Fol. Orch. Epid. 41 = punctiferum
tripunctum, Sw. Prod. Veg. Ind. Occ. 122 = Oncidium triquetrum.
tristi, Forst. f. Prod. 80 (Luisiae sp.).—Pensang; N. Caled.
triste, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 20 = ochraceum.
Trulla, *Reichb. f. in Bonplandia*, iv. (1856) 214.—Hab.?
tuberosum, *Forst. f. Prod.* 59.—N. Caled.
tuberosum, Linn. Sp. Pl. ed. II. 1352 (sp. delend.).—Ind. or.; Bras.
turbinatum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 41.—Mexic.
Turialvae, *Reichb. in Gard. Chron.* (1871) 1078.—Costa Rica.
umbellatum, Buch.-Ham. ex Lindl. Gen. et Sp. Orch. 221 = Saccolabium dayanum.
umbellatum, Forst. f. Prod. 60 = Cirrhopetalum Thouarsii.
umbellatum, Sw. Prod. Veg. Ind. Occ. 121 = difforme.
umbellatum, Vell. Fl. Flum. ix. t. 30 = Cheiropterocephalus corymbiferus.

EPIDENDRUM:—
umbelliferum, J. F. Gmel. Syst. 65 = difforme.
undulatum, Sw. Prod. Veg. Ind. Occ. 123 = Oncidium carthagenense.
uniflorum, Lindl. Bot. Reg. (1839) Misc. 15 = Hormidium uniflorum.
uniflorum, *Kodrig. Orch. Nov.* i. 61.—Bras.
uniflorum, *Vell. Fl. Flum.* ix. t. 34.—Bras.
urnuoides, D. Don, Prod. Fl. Nep. 37 = Sarcochilus urnoides.
utricularioides, Sw. Prod. Veg. Ind. Occ. 122 = Ionopsis utricularioides.
utriculatum, Poir. Encyc. Suppl. i. 375 = Govenia utriculata.
vaccinoides, *Reichb. f. ex Linden, in Illustr. Hortic.* xxix. (1882) 51, nomen.—Venezuela.
vandiflorum, *Lindl. in Journ. Hort. Soc.* iv. (1849) 369.—Mexic.
Vanilla, Linn. Sp. Pl. 952 = Vanilla angustifolia, aromatica.
vanillum, St. Lag in Ann. Soc. Bot. Lyon, vii. (1880) 125 = præc.
Vanneriarum, *Reichb. f. in Gard. Chron.* (1885) ii. 676.—Hab. ?
varicosum, *Batem. in Bot. Reg.* (1838) *Misc.* 30.—Am. centr.
variegatum, *Hook. Bot. Mag.* t. 3151.—Bras.
variegatum, Koen. in Retz. Obs. vi. 44 (Quid ?).—Ind. or.
varsegatum, Sw. Prod. Veg. Ind. Occ. 122 = Oncidium variegatum.
venosum, *Lindl. Gen. et Sp. Orch.* 99.—Mexic.
Ventilabrum, *Vell. Fl. Flum.* ix. t. 32.—Bras.
ventricosum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 86.—Peruv.
vernicium, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 110.—Peruv.
verrucosum, Lindl. Bot. Reg. (1844) t. 51 = nemorale.
verrucosum, *Sw. Fl. Ind. Occ.* iii. 1497.—Jamaica
vesicatum, *Lindl. Bot. Reg.* (1838) *Misc.* 50.—Bras.
Vespa, *Vell. Fl. Flum.* ix. t. 37.—Bras.
vestitum, Sw. Prod. Veg. Ind. Occ. 124 = Ornithidium vestitum.
Vieji, *Reichb. f. in Bonplandia*, iii. (1855) 220.—Nicaragua.
vincentinum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 88.—Ins. S. Vincent.
violaceum, *Spog. Select. Am.* 230.—Am. austr.
violaceum, Lodd Bot. Cab. t. 887 = Cattleya Loddigesii.
violaceum, Reichb. f. in Walp. Ann. vi. 218 = Cattleya violacea (in Addend.).
violodora, Gal. ex Lindl. Fol. Orch. Epid. 10 = selligeum.
virens, Hoffmgg. in Linnæa, xvi. (1842) Litt. 238 = latilabrum.
virens, *Lindl. & Past. Flow. Gard.* i. (1850–51) 152.—Guatemala.
virescens, Lodd. Bot. Cab. t. 1867 = ancepa.
virescens, Poir. Encyc. Suppl. i. 374 = Chloraea Piquichen.
virgatum, *Lindl. in Hook. Journ. Bot.* iii. (1841) 88.—Mexic.
virgatum, *Vell. Fl. Flum.* ix. t. 40.—Bras.
viride, *Ruiz & Pav. Syst. Veg.* 244.—Peruv.
viridibrunneum, *Reichb. f. in Walp. Ann.* vi. 868.—N. Granat.
viridiflorum, *Lindl. Bot. Reg.* (1842) *Misc.* 80.—Bras.
viride-purpureum, Hook. Bot. Mag. t. 3666 = anceps.
viscidum, Lindl. Bot. Reg. (1840) Misc. 81 = ciliare.
vitellinum, *Lindl. Gen. et Sp. Orch.* 97.—Mexic.
viviparum, *Lindl. Bot. Reg.* (1841) *Misc.* 10.—Guian.
volubile, *Ruiz & Pav. Syst. Veg.* 247.—Peruv.
voluta, Thou. Fl. Ile Austr. Afr. t. 81 = Aeonia Auberti.
volutum, *Lindl. & Past. Flow. Gard.* ii. (1851–52) 151.—Am. centr.
vomiforme, *Sw. Prod. Veg. Ind. Occ.* 124.—Jamaica.
Wagneri, Klotzsch in Otto & Dietr. Allg. Gartens. xix. (1851) 250 = virens.
Walkeriauum, Reichb. f. in Walp. Ann. vi. 416 = Cattleya Walkeriana.
Wallisii, *Reichb. f. in Gard. Chron.* (1876) ii. 96.—N. Granat.
Warneckianum, *Reichb. f. in Bot. Zeit.* x. (1852) 742.—Costa Rica.
Weddelii, *Lindl. Fol. Orch. Epid.* 67.—Bras.
Widgrenii, *Lindl. l. c.* 89.—Bras.
Wilsonianum, Reichb. f. in Gard. Chron. (1877) ii. 72 = Cattleya Wilsoniana.
Wrightii, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. (1858) 331.—Cuba.
xanthinum, *Lindl. Bot. Reg.* (1844) *Misc.* 18.—Bras.
xantholeucum, *Reichb. f. in Linnæa*, xxii. (1849) 839. —Venezuela.
xipheres, *Reichb. f. in Fl. des Serres*, Sér. I. ix. (1853–54) 98.—Peruv.
xylostachyum, *Lindl. Bot. Reg.* (1845) *Misc.* 70.—N. Granat.
xytriophorum, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 112.—Peruv.
ybaguense, *H. B & K. Nov. Gen. et Sp.* i 352.—N. Granat.

EPIFAGUS, Nutt. Gen. Am. ii. 60 (1818). *OROBANCHACEAE*, Benth. & Hook. f. ii. 986.
LEPTAMNIUM, Raf. in Am. Monthly Mag. (1818) 367.
MYLANCHE, Wallr. Orob. 75 (1825).
americanus, *Nutt. l. c.*—Am. bor
virginianus, Bart. Com. Fl. Philad. ii. 50 = americanus.

EPIGAEA, Linn. Diss Chen. 27 (1751); in Amoen Acad. iii. 17 (1756). *ERICACEAE*, Benth. & Hook. f. ii. 585.
MEMARCYLUM, Mitch. in Act. Phys. Med. Acad. Nat. Cur. vii. (1748) 218, App.
PARAFYROLA, Miq. in Ann. Mus. Lugd. Bat iii. 191 (1867).
asiatica, *Maxim. in Bull. Acad. Petersb.* xi. (1867) 431.—Japon.
cordifolia, Sw. *Prod. Veg. Ind. Occ.* 73.—Am. austr.
repens, *Linn. Sp. Pl.* 395.—Am. bor.
rubescens, Stend. Nom. ed. II. i. 559 = repens.

EPIGENIA, Vell. Fl. Flum. 188 (1825); iv. t. 137, 138 = Styrax, Tourn.
crenata, Vell. ll. cc. 184; iv. t. 138 = Symplocos lanceolata.
integerrima, Vell. ll. cc. 183; iv. t. 137 = Styrax glabratum.

EPIGOGIUM, Koch, Syn Fl. Germ. ed. I. 698, sphalm. (1837) = **Epipogum**, S. G. Gmel. (Orchid.).

EPIGYNANTHUS, Blume, in Flora, viii. (1825) 679 = **Hydrilla**, Rich. (Hydrocharid.).
Blumei, Hasak. Cat. Hort. Bog. Alt. 53 = H. ovalifolia.

EPIGYNIUM, Klotzsch, in Linnæa, xxiv. (1852) 49 = **Acotinum**, Linn.
Acosta, Klotzsch, l. c. 58 = V. Acosta.
acuminatum, Klotzsch, l. c. 52 = Corallobotrys acuminata.
affine, Klotzsch, l. c. 51 = V. Donianum.
coriaceum, Klotzsch, l. c. 52 = V. coriaceum.
cuneifolium, Klotzsch, l. c. = V. cuneifolium.
Donianum, Klotzsch, l. c. 51 = V. Donianum.
Dunalianum, Klotzsch, l. c. = V. Dunalianum.
elipticum, Klotzsch, l. c. 52 = V. ellipticum.
floribundum, Klotzsch, l. c. = V. floribundum.
Griffithianum, Klotzsch, l. c. 51 = V. Griffithianum.
laurifolium, Klotzsch, l. c. 52 = V. laurifolium.
Leschenaultii, Klotzsch, in Linnæa, xxiv. (1852) 50 = V. Leschenaultii.
leucobotrys, Nutt. ex Hook. Bot. Mag. t. 5108 = V. serratum.
lucidum, Klotzsch, l. c. 52 = V. lucidum.
malacense, Klotzsch, l. c. 50 = V. malacoense.
myrtoidermum, Klotzsch, l. c. 52 = V. myrtoides ?
neilgherrense, Klotzsch, l. c. 50 = V. neilgherrense.
obovatum, Klotzsch, l. c. 51 = Agapetes obovata.
rotundifolium, Walp. Ann. ii. (1852) 1095 = V. Leschenaultii.
serratum, Klotzsch, in Linnæa, xxiv. (1852) 50 = V. serratum.
Sprengelii, Klotzsch, l. c. 51 = V. Donianum.
variegatifolium, Klotzsch, l. c. 52 = V. variegatifolium.
venosum, Klotzsch, l. c. 50 = V. venosum.

EPIGYNUM, Wight, Ic. t. 1308 (1848). *APOCYNACEAE*, Benth. & Hook. f. ii. 717.

EPIGYNUM:—
LEGOUIXIA, Muell. Arg. in Heurck, Pl. Nov. 145 (1871).
Griffithianum, *Wight, Ic.* t. 1308.—Burma.
laevigatum, *Hook. f. Fl. Brit. Ind.* iii. 666.—Reg. Himal.
Maingayi, *Hook. f. Fl. Brit. Ind.* iii. 666.—Malacca.
parviflorum, *Hook. f. l. c.*—Ind. or.

EPILASIA, Benth. & Hook. f. Gen. ii. 582 (1873) *COMPOSITAE*, Benth. & Hook. f. ii. 583.
Acrolasia, C. B. Clarke, Comp. Ind. 279 = Bungei.
ammophila, C. B. Clarke, l. c. = Bungei.
Bungei, *C. B. Clarke, l. c.*—As. temp.
cenopleura, C. B. Clarke, l. c. = Bungei.
hemilasia, C. B. Clarke, l. c. = Bungei.
intermedia, C. B. Clarke, l. c. = Bungei.

EPILATORIA, Comm. ex Steud Nom. ed. II. i. 559 (1840) = ELPHEGEA, Cass. acc. Steud. = **Faradia** ?, Jacq (Compos.).

EPILEPIS, Benth. Fl. Hartw. 17 (1839) = **Coreopsis**, Linn. (Compos.).
rudis, Benth. l. c. = C. rudis.

EPILEPSIS, Lindl. Veg Kingd. 711 (1847) = EPILEPIS, Benth. = **Coreopsis**, Linn. (Compos.).

EPILINELLA, Pfeiff. in Bot. Zeit. iii. (1845) 673 = **Cuscuta**, Linn. (Convolvul.).
cuscutoides, Pfeiff. l. c. = C. Epilinum.

EPILITHES, Blume, Bijdr. 784 (1826) = **Serpicula**, Linn. (Halorag.).
coccinea, Blume, l. c. (= *Laurembergia coccinea*).—Java.

EPILOBIUM, Dill. ex Linn. Syst. ed. I (1736). *ONAGRARIEAE*, Benth. & Hook. f. i. 787.
CHAMAENERION, [Tourn.] Adans. Fam. ii. 85 (1763).
CROSSOSTIGMA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 174.
abortivum, Hausek. Monog. Epilob. 178 = lanceolatum.
acidulum, Borb. in Oesterr. Bot. Zeitschr. xxix. (1879) 182 = obscurum.
acinifolium, Reichb. ex Schur, Enum Pl. Transs. 218 = collinum.
acutum, Dalac, Fl. Hautes-Pyr. 331 = Duriaei.
adenocaulon, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 119.—Am. bor.
Adi, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 341, nomen.—Rossia.
adnatum, Griseb. in Bot. Zeit. x. (1852) 851 = tetragonum.
affine, Hong. Obs. Vég. Sitcha, 17 = glandulosum.
affine, Maxim. in Ind. Sem. Hort. Petrop. (1869) 16 = japonicum.
aggregatum, Celak. in Sitzb. Boehm. Ges. Wiss. (1873) = montanum.
alatum, Hegetschw. Fl. Schw. 357 = alsinifolium.
alsidiforum, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 341, nomen.—Rossia.
algidum, *Bieb. Fl. Taur. Cauc.* i. 296.—Reg. Caucas.
alpestre, Eichw. Skizze, 161 = algidum.
alpestre, Hoppe, ex Murith, Guide Val. 66 = trigonum.
alpestre, Jacq. Obs. Vindob. 35 = alpinum ?
alpestre, Schmidt, Fl. Bohem. iv. 81 = alsinifolium.
alpicolum x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 63.—Europ.
alpinum, Griseb. in Bot. Zeit. x. (1852) 853 = Hornemanni.
alpinum, Hads. Fl. Angl. ed. I. 162 = alsinifolium.
alpinum, *Linn. Sp. Pl.* 348.—Europ.; As. bor. et Arct. Reg.
alsinefolium, Wirtg. ex Haussk. Monog. Epilob. 187 = palustre.
alsinifolium, Bivt, Norges Fl. iii. (1826) 1120 = Hornemanni.
alsinifolium, *Vill. Prosp.* 45; *Hist. Pl. Dauph.* iii. 511. —Europ.
alsinoides, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 93. —N. Zel.
alternans, Link, ex Haussk. Monog. Epilob. 92 = lanceolatum.
ambigens, *Haussk. Monog. Epilob.* 178, nomen.—Hab. ?

EPILOBIUM:—

ambiguum, Fries, Summa Veg. 177 = obscurum.
americanum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 113.—Am. bor.
amoenum, *Rafin. in Med. Repos. N. York*, II. v. (1808) 354, nomen; *Am. Monthly Mag.* (1818) 266.—Am. bor.
amphibolum, *Haussk. Monog. Epilob.* 174, nomen.—Hab. ?
amplectens, *Benth. in Wall. Cat.* n. 6880.—Reg. Himal.
amplexicaule, Lam. Fl. Fr. iii. 479 = hirsutum.
amurense, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 85.—Reg. Amur.
anagallidifolium, Lam. Encyc. ii. 376 = alpinum.
anatolicum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 89.—As. Min.
anceps, Fries, Nov. Fl. Suec. ii. 90 = alsinifolium.
anceps, Lamotte, Prod. i. 389 = lanceolatum.
anditicum, Gill. ex Hook. Bot. Misc. iii. (1888) 309 = pedicellare.
andicolum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 118.—Am. austr.
angustifolium, Lam. Fl. Fr. iii. 482 = Dodonaei.
angustifolium, *Linn. Sp. Pl.* 347.—Europ.; As. bor.; Am. bor.
angustissimum, Weber, Pl. Mimus Cogn. Dec. 8 = Dodonaei.
Antonianum, Anct. ex Pers. Syn. i, 409 = angustifolium.
approximatum ×, Hausskn. Monog. Epilob. 177, nomen.—Europ. ?
aquaticum, Thuill. Fl. Par. ed. II. 191 = hirsutum.
Arnottianum, Gill. ex Hook. Bot. Misc. iii. 309 = pedunculare.
Aschersonianum ×, Hausskn. Monog. Epilob. 178, nomen.—Europ.
atriplicifolium, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 82.—N. Zel.
attenuatum, Schur, Enum. Pl. Transs. 209 = parviflorum.
australe, *Poepp. & Hausskn. in Haussk. Monog. Epilob.* 269.—Am. austr.
Balansae, Boiss. Fl. Orient. ii. 750 = origanifolium.
Baueri, Endl. in Enum. Pl. Hong. 44 = junceum.
Beckhausii, *Haussk. Monog. Epilob.* 213.—Europ.
Behringianum, *Haussk. l. c.* 277.—As. bor.
biforum, Schur, ex Hausskn. l. c. 110 = Lamyi.
Billardierianum, *Ser. in DC. Prod.* iii. 41.—Austral.
Blytianum ×, Hausskn. Monog. Epilob. 177.—Europ.
Boissieri ×, Hausskn. l. c.—Europ. ?
Bolezi, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 90.—Madag.
Bollianum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II xxiii.-xxiv. (1858) 57.—Europ.
Bongardi, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 87.—Alaska.
Bonplandianum, *H. B. & K. Nov. Gen. et Sp.* vi. 95.—Reg. Andin.
Borbasianum ×, Hausskn. Monog. Epilob. 177.—Europ.
Borderianum ×, Hausskn. l. c.—Europ.
boreale, *Haussk. l. c.* 279.—Am. bor. occ.
brachiatum, Celak. in Sitzb. Bohm. Ges. Wiss. (1879) = obscurum.
brachycarpum, Leight. in Ann. Nat. Hist. viii. (1841) 401 = angustifolium.
brachycarpum, Presl, Rel. Haenk. ii. 80 = repens.
Brandzai, Floriano, in Magyar. Növényt. Lapok, ix. (1885) 137 = alsinifolium.
brasiliense, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 119.—Bras.
brevifolium, *D. Don, Prod. Fl. Nep.* 232.—Reg. Himal.
brevipes, *Hook. f. Fl. N. Zel.* ii. 236.—N. Zel.
brevipilum ×, Hausskn. Monog. Epilob. 177.—Europ.
brevistylum, Barb. in Bot. Calif. i. 220 = glandulosum.
caesium, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 91.—Am. austr.
caespitosum, *Haussk. Monog. Epilob.* 261.—N. Zel.
californicum, *Haussk. l. c.* 260.—Calif.
calycinum, *Haussk. l. c.* 196.—Japon.
canescens, *Endl. in Enum. Pl. Hong.* 134.—Austral.
canescens, White, in Ann. Nat. Hist. i. (1838) 206; et in Proc. Bot. Soc. Lond. i. (1838) 208 = Dodonaei.
capense, *Buching. ex Krauss, in Flora*, xxvii. (1844) 625.—Afr. austr.
carpinifolium, Willk. Sert. Fl. Hisp. 80 = montanum.
Celakovskyanum ×, Haussk. Monog. Epilob. 177.—Europ.

EPILOBIUM:—

cephalostigma, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 91.—Japon.
cernuum, Pall. ex Haussk. Monog. Epilob. 158 = alpinum.
cerulescens, *Rafin. in Med. Repos. N. York*, II. v. (1808) 354, nomen.—Am. bor.
chilense, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 118.—Chili.
chlorandrum, *Haussk. l. c.* 149.—N. Zel.
chlorasefolium, *Haussk. l. c.*—N. Zel.
chordorrhizum, Fries, Summa Veg. 177 = obscurum.
ciliatum, *Rafin. in Med. Repos. N. York*, II. v. (1808) 361.—Am. bor.
cinereum, *A. Rich. Bot. Voy. Astrol.* i. 880.—N. Zel.
Clarkeanum, *Haussk. Monog. Epilob.* 220.—Reg. Himal.
cochinchinense, Wall. Cat. n. 6884 = hispidum.
collinum, *C. C. Gmel. Fl. Bad.* iv. 865.—Europ.
coloratum, *Muhl. ex Willd. Enum. Hort. Berol.* 411; *Spreng. Fl. Hal. Mant.* 39.—Am. bor.
commutatum ×, Haussk. Monog. Epilob. 178.—Europ.
conferruminatum, *Hook. f. Ic. Pl.* t. 685.—Tasman.
confertum, A. Cunn. in Ann. Nat. Hist. iii. (1839) 84 = junceum.
confine ×, Haussk. Monog. Epilob. 177.—Europ.
confusum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 151.—Armenia; Reg. Casp.
conversum ×, Haussk. Monog. Epilob. 178.—Europ.
consimile, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 58.—Reg. Cancas.
consociatum ×, Haussk. Monog. Epilob. 178.—Europ.
conspersum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879).—Ind. or.
cordatum, Bivona, Andr. Plant. Ined. 9 = parviflorum.
cordifolium, D. Don; ex Haussk. Monog. Epilob. 166 = alsinifolium.
cordifolium, *A. Rich. Tent. Fl. Abyss.* i. 274. t. 50.—Abyss.
corymbosum, Roth, in Skr. Dansk. Selsk. x. (1770) 440. t. 3. f. 28 = latifolium.
crassicaule, Gremli, Beitr. Fl. Schw. 78 = montanum.
crassifolium, Lehm. Ind. Sem. Hort. Hamb. (1824) 8, 30 = Dodonaei.
crassum, *Hook. f. Fl. N. Zel.* ii. 238.—N. Zel.
Cunninghamii, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 118.—Austral.
cupreum, *Lange, Ind. Sem. Hort. Haun. App.* (1872) 2.—Hab ?
cylindricum, *D. Don, Prod. Fl. Nep.* 232 = roseum.
dacicum, Borb. in Erdk. ix. (1879) XVI. 5 = obscurum.
dasycarpum, Fries, Mant. ii. 21 = alpinum.
davuricum, *Pisch. in Hornem. Hort. Hafn. Suppl.* 44; *et in Spreng. Nov. Prov.* 17.—Dahur.
decipiens, Haussk. in Oestr. Bot. Zeitschr. xxix. (1879) 57 = tetragonum.
decumbens, Dum. Fl. Belg. Prod. 89 = obscurum.
decurrens, Hort. Hal. (1813); ex Kelchb. Ic. Fl. Germ. ii. 89 = tetragonum.
densifolium, *Haussk. Monog. Epilob.* 246.—Chili.
denticulatum, Kunze, ex Haussk. l. c. 276 = pedicellare.
dentosum, Rafin. Précr. Découv. 42; et in Desv. Journ. Bot. ii. (1814) 271 = lineare.
denticulatum, C. Gay, Fl. Chil. ii. 347 = paberulum.
denticulatum, Ruiz & Pav. Fl. Per. iii. 76 = junceum, pedicellare.
denticulatum, Wender. Ind. Sem. Hort. Marb. (1824) 9 = Dodonaei.
differme, Gillb. Fl. Lituan. ii. 190 = angustifolium.
divaricatum, Rafin. in Med. Repos. N. York, II. v. (1808) 354; Précr. Découv. 42 = coloratum.
diversifolium, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 151.—Tasmania.
Dodonaei, *Vill. Prosp.* 45; *Hist. Pl. Dauph.* iii. 507.—Europ.
doriphyllum, *Haussk. Monog. Epilob.* 257.—Mexic.
dovrense ×, Hausskn. l. c. 177.—Europ. hor.
draconcellense, Ser. ex Gand. Fl. Helv. iii. 9 = Dodonaei.
Dregeanum, E. Mey. ex Haussk. Monog. Epilob. 109 = tetragonum.
dubium, Borb. in Erdk. ix. (1879) XVI. 6 = hirsutum?
dubium, Curt. Cat. n. 507; ex With. Arr. Brit. Pl. ed. II. i. 390 = tetragonum?
Duftii ×, Haussk. Monog. Epilob. 178.
Durieui, *F. Gay, in Ann. Sc. Nat.* Ser. II. vi. (1836) 288 (ex. typ. 128).—Mosel. Pyren.

EPILOBIUM:—

Duthiei, *Haussk. Monog. Epilob.* 205.—Reg. Himal.
ecostatum, Chaix, ex Haussk. l. c. 87 = collinum.
elatum, Munro, ex Haussk. l. c. 44 = angustifolium.
emirnifolium, Kunze, ex Haussk. l. c. 156 = alpinum.
erosum, *Haussk. Monog. Epilob.* 268.—Tasmania.
eximum, Spachy ex Haussk. l. c. 290 = junceum.
erronenum ×, Haussk. l. c. 178.—Europ.
erubescens, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 150.—N. Zel.
euphrasioides, Sieber, ex Haussk. Monog. Epilob. 290 = junceum.
Fauchieri, Hausm. ex Nym. Consp. 845 = alsinifolium.
fallacinum ×, Haussk. Monog. Epilob. 177.
fastigiatum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 489 = glandulosum.
Fendleri, *Haussk. Monog. Epilob.* 261.—N. Mexic.
fennicum, Laest. ex Haussk. l. c. 120 = obscurum.
finitimum ×, Haussk. Monog. Epilob. 177.—Europ.
fissipetalum, Stend. ex A. Rich. Tent. Fl. Abyss. i. 278 = stereophyllum.
flaccidum, Brot. Fl. Lusit. ii. 18 = Toumefortii.
flaccidum, Soland. ex Haussk. Monog. Epilob. 300 = rotundifolium.
flavescens, *E. Mey. ex Horv. & Sond. Fl. Cap.* ii. 507 —Afr. austr.
Fleischeri, Hochst. in Flora, ix. (1826) 28 = Dodonaei.
flexuosum, Nutt. ex Haussk. Monog. Epilob. 249 = minutum.
foliosum, Hochst. in Flora, xxiv. (1841) t. Intell. 30 = hirsutum.
foliosum, Hoff. & Heynh. Fl. Dresd. 297 = parviflorum.
foliosum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 490 = minutum.
fontanum, Chaix, ex Haussk. Monog. Epilob. 166 = alsinifolium.
Franciscanum, *Barbey, in Bot. Calif.* i. 220.—Am bor. occ.
Freynii, Celak. in Arch. Naturw. Landesd. Boehm. iv (1881) 881 = tetragonum.
frigidum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 51.—As. Min.; Reg. Caucas.
frigidum, Retz. Fl. Scand. Prod. ed. II. 89 = latifolium.
fruticosum, Lour. Fl. Cochinch. 236 = Jussiaea suffruticosa.
gelidum, Vahl, ex Haussk. Monog. Epilob. 82 = Dodonaei.
gemmascens, Lange, in Kjoeb. Vidensk. Meddel. (1865) 187 = obscurum.
gemmascens, C. A. Mey. Verz. Pfl. Cauc. 178 = roseum.
gemmiflorum, Bor. in Bull. Soc. Agricn. Maine-et-Loire (1855); ex Mem. Compt. 247 = alsinifolium.
gemmiflorum, Willk. & Lange, Prod. Fl. Hisp. iii. 184 = obscurum.
Gerardianum, Wall. Cat. n. 6896 = latifolium.
Gesneri, Vill. Prosp. 45 = angustifolium.
glabellum, R. Br. ex Haussk. Monog. Epilob. 294 = Billiardierianum.
glabellum, *Forst. f. Prod.* 37.—N. Zel.
glaberrimum, *Barbey, in Bot. Calif.* i. 220.—Am. bor. occ.
glabrum, Gillib. Fl. Lituan. ii. 186 = montanum.
glanduligerum ×, K. Knaf, ex Celak. in Arch. Naturw. Landesd. Boehm. iv. (1881) 882 = Europ.
glandulosum, *Lehm. in Hook. Fl. Bor. Am.* i. 206.—Am. bor. occ.
glaucinum, Haussk. in Oestr. Bot. Zeitschr. xxix. (1879) 51 = urolepe.
glaucum, Phil. in Linnaea, xxxiii. (1864–65) 70 = pedicellare.
gorgoneum, *Lowe, ex Haussk. Monog. Epilob.* 58.—Ins. Madera.
gracile, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxv. (1882) 70 = angustifolium.
grandiflorum, Jacquem. Voy. Bot. 57 = latifolium.
grandiflorum, [Weber, in] Wigg. Prim. Fl. Holsat. 80 = hirsutum.
Griffithianum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 151.—Afghan.
Gunckelianum, *Haussk. l. c.* 149.—Tasmania.
Haenkeanum, *Haussk. l. c.* 146.—Am. austr.
Hallori, Retz. Fl. Scand. Prod. ed. II. 89 = latifolium.
haloxanthium, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 84.—N. Zel.
Hausmannianum ×, Borb. in Oestr. Bot. Zeitschr. xxix. (1879) 128.—Europ.
Hegetschweilerum ×, Haussk. Monog. Epilob. 177.—Europ.

EPILOBIUM :—
Hectori, *Hausskn. Monogr. Epilob.* 298.—N. Zel.
Heerianum ×, Brueggr. in Jahresb. Naturf. Ges. Graub. II. xxiii.-xxiv. (1880) 56.—Europ.
heterocaule, Borb. in Ertek. ix. (1879) xvi. 15 = montanum.
heterophyllum, Hegetschw. Fl. Schw. 859; Beitr. Crit. Anfs. 337 = nutans.
heterophyllum, Nutt. ex Hausskn. Monog. Epilob. 275 = tetragonum.
himalayense, *Hausskn. l. c.* 218.—Reg. Himal.
himalense, Royle, Illustr. Bot. Himal. i. 211 = hirsutum.
hirsutum, Huds. Fl. Angl. ed. I. 161 = parviflorum.
hirsutum, *Linn. Sp. Pl.* 347.—Europ.; As. bor.
hirtigerum, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 33.—N. Zel.
hispidulum, *Steud. Nom.* ed. II. i. 559.—Chili.
hispidum, *Weinm. in Bull. Soc. Nat. Mosc.* xxiii. (1850) 1. 545.—Kossia.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 585.—Reg. Himal.; Japon.
Hornemanni, Bong. Veg. Sitka, in Mém. Acad. Péterb. vi. (1838) 186 = roseum.
Hornemanni, Dietr. Fl. Univers. ii. sect. 69. t. 579 = lactiflorum.
Hornemanni, Reichb. Ic. Fl. Germ. ii. 78 = alpinum.
Hornemanni, Schur, Enum. Pl. Transs. 212 = nutans.
Hugueninii ×, Brueggr. in Jahresb. Naturf. Ges. Graub. II. xxiii.-xxiv. (1880) 66.—Europ.
humile, Wild. ex Stev. in DC. Prod. iii. 40 = latifolium.
Hutchi ×, Borb ex Hausskn. Monog. Epilob. 177.—Europ.
hybridum, Schur, Enum. Pl. Transs. 209 = hirsutum.
hybridum, *Tausch. Hort. Canal. fasc.* 1. n. 7; *in Flora*, vii. (1824) 88.—Bohem.
incanum, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 33.—N. Zel.
incanum, Vers. Syn. i. 410 = hirsutum.
indicum, *Hausskn. Monogr. Epilob.* 199.—Reg. Himal.
insulare, *Hausskn. l. c.* 260.—N. Zel.
intermedium, Hegetschw. Fl. Schw. 856; Beitr. Crit Anfs. 341 = montanum.
intermedium, Métat, Fl. Par. 247 = hirsutum.
intermedium, Wormsk. ex Ser. in DC. Prod. iii. 40 = angustifolium.
intersitum ×, Hausskn. Monog. Epilob. 177.
japonicum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 55.—Japon.
jonanthum, *Hausskn. Monogr. Epilob.* 231.—Afr. austr.
jucundum, *A. Gray, in Proc. Am. Acad.* xii. (1877) 57.—Calif.
junceum, *Forst. f. Prod.* 90.—N. Zel.; Am. austr.
kamtschaticum, Lepech. in Nov. Act. Acad. Petrop. xi. (1798) 879. t. 6 = latifolium.
Kerneri, Borb. in Oestr. Bot. Zeitschr. xxvi. (1876) 17 = palustre.
khasianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 585.—Reg. Himal.
Knafii, Celak. in Arch. Naturw. Landesd. Boehm. iii. (1875) 551 = parviflorum.
Kreuseri, Uechtr. in Bot. Zeit. xxiv. (1874) 240 = palustre.
Krulleanum, *Hausskn. Monogr. Epilob.* 305.—N. Zel.
lactiflorum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 89.—Suecia.
lactum, Wall. Cat. n. 6329 = hirsutum.
Lamottianum ×, Hausskn. Monog. Epilob. 178.—Europ.
Lamyi, *Schultz, in Flora*, xxvii. (1844) 806.—Europ.; Orient.
lanceolatum, *Sebast. & Mauri, Prod. Fl. Rom.* 138. t. 1. f. 2.—Europ.
Langeanum ×, Hausskn. Monog. Epilob. 178.—Europ.
lapponicum, Baser, ex Hausskn. l. c. 34 = collinum.
lapponicum, Steud. Nom. ed. II. i. 559 = palustre.
Lavrangianum, F. Schultz, Arch. Fl. 272 = montanum.
Leschianum ×, Hausskn. Monog. Epilob. 177.—Europ.
latifolium, *Linn. Sp. Pl.* 347.—Reg. bor. et sept.
latifolium, Mattusch. Fl. Sil. i. 363 = angustifolium.
latifolium, F. W. Schmidt, Fl. Bohem. iv. 75. n. 372 = angustifolium.
lanum, Roxb., Illustr. Bot. Himal. 211 = tetragonum.
Lechleri, *Phil. & Hausskn. in Monogr. Epilob.* 270.—Am. austr.

EPILOBIUM :—
lelophyllum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 53.—Ind. or.
leiospermum, *Hausskn. Monogr. Epilob.* 206.—Tibet.
leiotylon, Petersn. Deutschl. Fl. (1849) 180 = angustifolium.
leptocarpum, *Hausskn. Monogr. Epilob.* 258.—Am. bor. occ.
leptophyllum, *Raffin. in Med. Repos. N. York*, ii. v. (1808) 258; *Prod. Monog.* 41.—Am. bor.
ligulatum, Baker, in Phytol. N. S. ii. (1857) 90, 404 = palustre.
limosum, Schur, Enum. Pl. Transs 213 = montanum.
Lindblomianum ×, Hausskn. Monog. Epilob. 178.—Europ.
linearе, Fries, Nov. Ment. iii. 185 = davuricum.
linearе, Mühl. Cat. 39 = palustre.
linearifolium, Beurl. ex Hausskn. Monog. Epilob. 148 = davuricum.
linifolium, *Bieb. Tabl. Casp.* 58.—Reg. Caucas.
linnaeoides, *Hook. f. Fl. Antarct.* 10. t. 6.—Tasman. et N. Zel.
lividum, *Haussk. Monogr. Epilob.* 201.—Ind. or.
Lobelii, Vill. Fl. Delph. 36 = Dodonaei.
longipes, Hort Kew. ex Hausskn. Monog. Epilob. 308 = pedunculare.
Lorentzianum, Hausskn l. c. 263 = Watsoni.
luteum, *Pursh, Fl. Am. Sept.* i. 259.—Am. bor. occ.
macranthum, Hook Ic. Pl. t. 297 = pallidiflorum.
macrocarpum, Steph. in Ann. Nat. Hist. vili. (1842) 170 = angustifolium.
macropus, *Hook. Ic. Pl.* t. 812.—N. Zel.
maderense, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 90.—Ins. Madera.
magellanicum, *Phil. & Haussk. in Haussk. Monog. Epilob.* 271.—Reg. Magellan.
Mallei, Wm. F. Schultz, Arch. Fl. 115 = Duriaei.
matrense, Borb. in Oestr. Bot. Zeitschr. xxix. (1879) 182 = obscurum.
Maximowiczii, Hausskn. l. c. 57 = glandulosum.
melanodictyon, *Hook. Ic. Pl.* t. 818.—N. Zel.
menthoides, Bolss. & Heldr. Diagn. Ser. II. ii. 53 = parviflorum.
merxmullerianum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 148.—Venezuela.
metallicum, Barbey, ex Hausskn. Monog. Epilob. 304 = glabellum.
mexicanum, Moç. & Sesse, ex DC. Prod. iii. 41 = Bonplandianum.
mexicanum, *Schlecht. in Linnaea*, xii. (1838) 266.—Mexic.
micranthum, Nutt. ex Hausskn. Monog. Epilob. 247 = paniculatum.
microanthum, Pall. ex Hausskn. l. c. 109 = tetragonum.
microphyllum, *A. Rich. Bot. Voy. Astrol.* i. 325.—N. Zel.
microphyllum, Wall. ex Hausskn. Monog. Epilob. 218 = nepalense.
minutiflorum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 55.—Syria.
minutum, *Lindl. in Hook. Fl. Bor. Am.* i. 207.—Am. bor. occ.
mirtum, Simk. in Termész. Füzetek, i. (1877) 104; et Oests. Bot. Zeitschr. xxvii. (1877) 158 = tetragonum.
modestum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 55.—Ind. or.
molle, Lam. Fl. Fr. iii. 479 = parviflorum.
molle, Torr. Fl. U. St. i. 393 = strictum.
mollissimum, Schur, Enum. Pl. Transs. 209 = parviflorum.
montanoformе, K. Knaf, ex Celak. in Arch. Naturw. Landesd. Boehm. iv. (1881) 382 = montanum.
montanum, Bolss. Fl. Orient. ii. 747 = anatolicum.
montanum, Bolss. Voy. Bot. Espagne, 212 = lanceolatum.
montanum, Hacq. It. Dac. Carp. i. 178 = angustifolium.
montanum, *Linn. Sp. Pl.* 348.—Europ.; As. bor.
montanum, Pall. ex Hausskn. Monog. Epilob. 127 = roseum.
Mundtii, *Haussk. l. c.* 235.—Afr. austr.
muscharlile, Bolss. & Rest. Diagn. Ser. II. ii. 54 = lanceolatum.
mutabile, Wender. ex Hausskn. Monog. Epilob. 88 = hypericifolium.
nataleuse, *Hausskn. l. c.* 235.—Afr. austr.
nebrodense ×, Strobl, ex Hausskn. l. c. 178.—Sicil.

EPILOBIUM :—
neogradiense, Borb. in Oestr. Bot. Zeitschr. xxix. (1879) 183 = lanceolatum.
nepalense, *Haussk. l. c.* xxix. (1879) 53.—Reg. Himal.
neriophyllum, *Haussk. Monog. Epilob.* 236.—Afr. austr.
nerterioides, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 32.—Austral.; N. Zel.
nervosum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 88 = roseum.
neutrum, Pall. ex Hausskn. Monog. Epilob. 71 = parviflorum.
nitidum, Host, Fl. Austr. i. 469 = collinum.
nitidum, Sauter, Fl. Salzb. 245 = alsinifolium.
nivale, *Meyen, Reise*, i. 815.—Chili.
norvegicum ×, Hausskn. Monog. Epilob. 178.—Norveg.
Nova-Zelandiae, *Hausskn. l. c.* 256.—N. Zel.
novo-mexicanum, *Hausskn. l. c.* 260.—N. Mexic.
novum, Winterl. Ind. Hort. Pest. 44. f. 17 = tetragonum.
nubigenum, *Phil. in Linnaea*, xxxiii. (1864–65) 71.—Chili.
nudum, Schur. in Fl. Dan. t. 1815 = roseum.
nummularifolium, *R. Cunn. ex A. Cunn. in Ann. Nat. Hist.* iii. (1839) 31.—N. Zel.
nutans, G. Don, Gen. Syst. ii. 681 = hypericifolium.
nutans, Hornem. in Fl. Dan. t. 1387 = lactiflorum.
nutans, Kern. in Oestr. Bot. Zeitschr. xix. (1869) 301 = palustre.
nutans, Lej. Rév. Fl. Spa, 76 = collinum.
nutans, *F. W. Schmidt, Fl. Boem.* iv. 82. n. 380.—Europ.
nutans, Sommerf. Fl. Lapp. Suppl. 17 = davuricum.
obcordatum, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 582.—Calif.
obscurum, Boiss. Fl. Orient. ii. 749 = confusum.
obscurum, Eckl. & Zeyh. Enum. 272 = capense.
obscurum, Host, Mant. ex Bieb. Fl. Taur. Cauc. iii. 282 = Tournefortii.
obscurum, Lowe, Man. Fl. Madeira, 274 = Lamyi.
obscurum, Reichb. Ic. Fl. Crit. ii. 89 = virgatum.
obscurum, F. W. Schmidt, Fl. Boem. iv. (1794) 81 = tetragonum.
obscurum, *Schreb. Spicil. Fl. Lips.* 147.—Europ.
obtruncatum, *Raffin. in Med. Repos. N. York*, ii. v. (1808) 254, nomen.—Am. bor.
oliganthum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 245.—Madag.
oligostachyum, Michx. Fl. Bor. Am. i. 228 = palustre.
oligodontum, *Haussk. in Oestr. Bot. Zeitschr.* xxix. (1879) 58.—Japon.
opacum, *Lehm. Pugill.* ii. 14.—Am. bor. occ.
opacum, Peterm. Deutschl. Fl. (1849) 180 = parviflorum.
oregonense, *Hausskn. Monog. Epilob.* 276.—Am. bor. occ.
organifolium, Anders. in Bot. Notiser (1844), 154 = Hornemanni.
organifolium, Bolss. Fl. Orient. ii. 750 = anatolicum.
organifolium, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 586 = leiophyllum.
organifolium, Kirschl. Prod. 88 = Duriaei.
organifolium, Lam. Encyc. ii. 376 = alsinifolium.
Otites, Hoffingg. ex Steud. Nom. ed. II. i. 560 = lanceolatum.
ovatum, Phil. ex Hausskn. Monog. Epilob. 272 = chilense.
Osanesii, *F. Schultz, Arch. Fl.* 278 = montanum.
palustinum, F. Schultz, *l. c.* 278 = Lamyi.
pallidiflorum, *Soland. ex A. Cunn. in Ann. Nat. Hist.* iii. (1839) 34.—N. Zel.
pallidum, Tausch, ex Hausskn. Monog. Epilob. 82 = montanum !
palustre, *Linn. Sp. Pl.* 348.—Reg. bor. temp.
palustre, Uerin, Hort. Panorm. 109 = tetragonum.
palustre, Wild. Sp. Pl. ii. 815 = parviflorum.
paniculatum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 490.—Am. bor.
pannosum, *Haussk. Monogr. Epilob.* 209.—Ind. or.
parviflorum, *Schreb. Spicil. Fl. Lips.* 148.—Europ.; As. occ.
parviflorum, Dulac, Fl. Hautes-Pyr. 382 = alsinifolium.
parviflorum, Schrank, in Denkschr. Bot. Ges. Regensb. i. 11. (1818) 18 = latifolium.
pedicellare, *Presl, Rel. Haenk.* ii. 30.—Chili.
pedunculare, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 31.—N. Zel.
pedunculatum, *Phil. in Anal. Univ. Chil.* (1872) 718.—Chili.

EPILOBIUM:—
peradenatum ×, Borb. in Oestr. Bot. Zeitschr. xxix. (1879) 368.—Europ.
perpusillum, *Hausskn. Monogr. Epilob.* 301.—Tasman.
perramosum, Schur, Enum. Pl. Transs. 213 = montanum.
persicifolium, Vill. Hist. Pl. Dauph. i. 328 = angustifolium.
persicinum, Reichb. Fl. Germ. Excurs. 685 = montanum.
peruvianum, *Hausskn. Monogr. Epilob.* 353.—Peruv.
phyllonemos, Knaf, ex Borb. in Ertek. ix. (1879) xvi. 21 = obscurum.
pilosum, Kit. ex Hausskn. Monogr. Epilob. 138 = roseum.
polyclonum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 150.—N. Zel.
ponticum, *Hausskn. Monogr. Epilob.* 203.—As. Min.
prionophyllum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 58.—Armenia.
propinquum, Hausskn Monog. Epilob. 218 = roseum.
pruinosum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 90.—Am. bor.
Pseudo-linearе, *Hausskn. l. c* 120—Calif.
Pseudo-obscurum, *Hausskn. l. c.* 53.—Ind. or.
Pseudo-scaposum, *Hausskn. l. c.* 89.—Unalask.
Pseudo-trigonum, Borb. l. c. xxvii. (1877) 188 = montanum.
Pseudo-virgatum, F. Schultz, ex Hausskn. Monog. Epilob. 109 = Lamyi.
ptarmicaefolium, F. Schultz, in Pollichia, xiii. (1855) 27 = tetragonum.
pubens, *A. Rich. Bot. Voy. Astrol.* i. 329. t. 86.—N. Zel.
puberulum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1832) 309.—Ins. Chiloe.
pubescens, Le. & Court. Comp. ii. 52 = montanum.
pubescens, Roth, Tent. Fl. Germ. i. 167 = parviflorum.
pumilum, Steud. Nom. ed. i. 301 = alpinum.
purpurascens, Gilli. Fl. Lituan. ii. 189 = obscurum.
purpuratum, *Hook f. Handb. N. Zeal. Fl.* 77.—N. Zel.
purpureum, Fries, Fl. Hal. 65 = palustre.
pusillum, Gill. ex Hook. Bot. Misc. iii. (1832) 309 = alpinum.
pycnostachyum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 150.—N. Zel.
pyrenaicum ×, Hausskn. Monog. Epilob. 177.—Europ.
pyrricholophum, *Franch. & Sav. Enum. Pl. Jap.* ii. 270.—Japon.
radicosa, Hausskn. Monog. Epilob. 81 = montanum.
ramiflorum, Hegetschw. Fl. Schw. 359; Beitr. Crit. Aufs. 542 = palustre.
ramosissimum, Gren. & Godr. Fl. Fr. i. 579, in syn. = obscurum.
ramosum, Huds. Fl. Angl. ed. I. 141 = hirsutum.
Reichenbachii, Bnegg. in Jahresb. Naturf. Ges. Graub. II. xxii.-xxiii. (1880) 67.—Europ.
repandum, Spach, ex Hausskn. Monog. Epilob. 290 = junceum.
repandum, *Steud. Nom.* ed. II. i. 560.—Chili.
repens, Hill, Brit. Herbal, 148 = alpinum.
repens. *Schlecht. in Linnaea*, xii. (1838) 267.—Mexic.
reticulatum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 583.—Reg. Himal.
rhenitiflorum, Fisch. ex Hausskn. Monog. Epilob. 194 = latifolium.
rhynchocarpum, Boiss. Diagn. Ser. II. ii. 53 = palustre.
rhynchopetmum, *Hausskn. Monog. Epilob.* 211.—Reg. Himal.
rigidum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 51.—Am. bor.
riphaeum, Tausch, ex Hausskn. Monog. Epilob. 166 = alsinifolium.
riparium, Hegetschw. Fl. Schw. 355 = parviflorum.
riparium, Kirschl. Fl. Als. 266 = hirsutum.
riparium, Nolte, Fl. Holsat. 41 = obscurum.
riparium, Wahlenb. Fl. Upsal. 196 = palustre.
rivulicolum ×, Hausskn. Monog. Epilob. 177.—Europ.
roseum. DC. Fl. Fr. iv. 432 = trigonum.
roseum, Fries, ex Steud. Nom. ed. II. i. 560 = montanum.
roseum, Kit. Addit. 274; ex Hausskn. Monog. Epilob. 166 = alsinifolium.
roseum, *Schrad. Spicil. Fl. Lips.* 147.—Europ.; As. occ.
roseum, Turcz. in Ledeb. Fl. Ross. ii. 116 = Hornemanni.
rosmarinifolium, Haenke, in Jacq. Coll. ii. 50 = Dodonaei.
rosmarinifolium, Pursh, Fl. Am. Sept. i. 359 = linearе.

EPILOBIUM:—
rotundifolium, *Forst. f. Prod.* 27.—N. Zel.
Roylеanum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 55.—Reg. Himal.
rubricaule, Soland. ex Hausskn. Monog. Epilob. 294 = Billardierianum.
rubrum, Lucé, Topog. Nachr. Oesel, 117 = angustifolium.
salicifolium, Facch. in Zeitschr. Ferdinand. Tirol, III. iv. (1854) 43 = alsinifolium.
salicifolium, Stokes, Bot. Mat. Med. ii. 351 = angustifolium.
saligнum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 90.—Madag.
Salisianum ×, Bnegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 67.—Europ.
Salomannii, Boiss. & Reut. ex Nym. Consp. 247 = Tournefortii.
sarmentaceum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 149.—Tasman.
sarmentosum, Celak. Prod. iv. 883 = palustre.
saxifragifolium, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 119.—Am. bor. occ.
scaturiginum, Wimm. ex Uechtr. in Bot. Zeit. xxiv. (1874) 341 = palustre.
Schimperianum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 273.—Abyss.
Schmidianum, Rouk. Fl. Sedin. 173 = palustre.
Schmidtii, Flcrik. ex Hausskn. Monog. Epilob. 166 = alsinifolium.
Schulzeanum ×, Hausskn. Monog. Epilob. 177.
scrotinatum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 868.—Europ.
scrobiculorum ×, Borb. l. c. 185.—Europ.
sericeum, Benth. in Wall. Cat. n. 6829 = hirsutum.
sericeum, *Schum. ex Hausskn. Monog. Epilob.* 64.—Ind. or.
serratum, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 583 = hirsutum.
serrulatum, Pourr. ex Willk. in Willk. & Lange, Prod. Fl. Hi p. iii. 187 = tetragonum.
serrulatum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 52.—Kamtsch.
sikkimense, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 52.—Reg. Himal.
silvaticum, Anct. = *sylvaticum*.
simplex, Tratt. Archiv, i. 37 = palustre.
smyrnaeum, Boiss. Diagn. Ser II. ii. 52 = roseum.
spectabilis, Dum. Fl. Belg. Prod. 89 = lanceolatum.
speciosum, Decne. in Jacquem. Voy. Bot. iv. 57 = latifolium.
spicatum, Lam. Fl. Fr. iii. 482; Encyc. ii. 573 = angustifolium.
spumatum, Nutt. Fl. Am. Sept. i. 250 = linearе.
Stefanescui, Florianu, in Magyar, Novént. Lapok, ix. (1886) 137 = alsinifolium.
Steigeri ×, Bnegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 65.—Europ.
stenophyllum, Borb. in Ertek. ix. (1879) xvi. 11 = parviflorum.
stereophyllum, *Fresen. in Mus. Senckenb.* ii. (1837) 163.—Abyss.
Stewart, Boiss. Diagn. Ser. II. ii. 55 = Dodonaei.
Strachеyanum, *Hausskn. Monog. Epilob.* 214.—Reg. Himal.
strictum, *Mukl. Cat.* 39; *et ex Sprengl. Syst.* ii. 238.—Am. bor.
subalpigнum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 59.—Scoogar.
subcordatum, *Hausskn. l. c.* 56.—China.
subhirsutum, *Gennar.—Cf. Linnaea*, xxiv. (1851) 201. —Italia.
subtectum, Benrl. in Bot. Notizer (1851), 185 = alpinum.
suffruticosum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 488.—Am. bor. occ.
sylvaticum, Bor. Fl. Cent. Fr. ed. III. ii. 239 = montanum.
sylvestre, Dierb. in Geiger, Mag. (Oct. 1826); ex Mert. & Koch, Deutschl. Fl. iii. 16 = montanum.
Sinicum ×, Hausskn. Monog. Epilob. 178.—Syria.
tanguticum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 56.—China.
tasmanicum, *Hausskn. Monog. Epilob.* 296.—Tasman.
Tauschii, Knaf, ex Hausskn. l. c. 78 = parviflorum.
tenellum, Rafin. Prec. Découv. 41 = palustre.
tenuipes, Hook. f. Fl. Tasmac. i. 116 = confertifolium.
terratum, *Schur, Enum. Pl. Transs.* 208 (sp. dub.).—Transylv.

EPILOBIUM:—
tetragoniforme, Simk. Term. Rajzi. (1877) 105 = parviflorum.
tetragonum, *Aitch. in Journ. Linn. Soc.* xix. (1881) 183 = palustre.
tetragonum, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 586 = Wallichianum.
tetragonum, Cass. Fl. Sic. Syn. i. 444 = Tournefortii.
tetragonum, *Linn. Sp. Pl.* 348.—Europ.
tetragonum, Lour. Fl. Cochinch. i. 281 = Jussiaea suffruticosa.
tetragonum, Pollich, Hist. Pl. Palat. i. 377 = obscurum.
tetrapetalum, Pall. ex Steud. Nom. ed. II. i. 560 = latifolium.
thuringiacum ×, Hausskn. Monog. Epilob. 177.—Europ.
thymifolium, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 33.—N. Zel.
tibetanum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 59.—Tibet.
tingitanum, Salzm. ex Ball, in Journ. Linn. Soc. xvi (1878) 459 = tetragonum.
tomentosum, Gilib. Fl. Lituan. ii. 189 = palustre.
tomentosum, Vent. Jard. Cels, t. 90 = hirsutum.
Tournefortii, Freyn, Fl. Sued. Istr. 97 = tetragonum.
Tournefortii, *Michal. in Bull. Soc. Bot. Fr.* ii. (1855) 731.—Oriens.
Trevirnianum, Ausserd. ex A. Kern. Sched. ii. 41 = palustre.
trichonerurum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix (1879) 54.—Ind. or.
trichophyllum, *Hausskn. l. c.* 53.—Reg. Himal.
trigonum, Boiss. Fl. Orient. ii. 749 = algidum.
trigonum, Hort. ex Steud. Nom. ed. II. i. 560 = montanum.
trigonum, Ledeb. Fl. Ross. ii. 111 = prionophyllum.
trigonum, *Schrank, Baier. Fl.* i. 644.—Europ.
udicolum ×, Hausskn. Monog. Epilob. 178.—Europ.
Uechtritzianum ×, Pax, in Deutsche Bot. Monatsschr. i. (1883) 139.—Europ.
uliginosum, Fries, ex Hausskn. Monog. Epilob. 198 = roseum.
umbrosum, Dum. Fl. Belg. Prod. 89 = parviflorum.
uralense, Hort. Petrop. ex Hausskn. Monog. Epilob. 72 = parviflorum.
uralense, *Rupr. in E. Hoffm. Ural. Bor.* ii. *Suppl.* B, 32.—Reg. Himal.
valdivianum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix (1879) 138.—Chili.
variabile, Lucé, Topog. Nachr. Oesel, 187 = angustifolium.
vernutum, Dougl. ex Steud. Nom. ed. II. i. 560 = latifolium.
verticillatum, Schur, Enum. Pl. Transs. 213 = montanum.
verticillatum, Tenore, Fl. Neap. Prod. 23; Fl. Neap. i. 194 = angustifolium.
vestitum, Benth. in Wall. Cat. n. 6827 = parviflorum
villosum, Curt. Fl. Lond. fasc. ii. t. 22 = parviflorum.
villosum, Thunb. Prod. Pl. Cap. 78 = hirsutum.
virgatum, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 33. —N. Zel.
virgatum, Koch, Syn. Fl. Germ. 866 = tetragonum.
virgatum, Lam. Encyc. ii. 575 = obscurum.
virgatum, Lamy, ex Koch, Syn. Fl. Germ. ed. II. 1093 = Lamyi.
vogesiacum, Hausskn. Monog. Epilob. 178 = palustre.
Wallichianum, *Hausskn. in Oestr. Bot. Zeitschr.* xxix. (1879) 54.—Reg. Himal.
Watsoni, *Barbey, in Bot. Calif.* i. 219.—Am. bor. occ.
Watsianum, *Hausskn. Monog. Epilob.* 304.—Reg. Himal.
weissenburgense, F. Schultz, in Pollichia, xviii-xix (1861) 166 = tetragonum.
Wimmeri, F. Schultz, l. c. (1859) 109 = obscurum.
Winkleri, Kern. in Oestr. Bot. Zeitschr. xxvi. (1876) 113 = alsinifolium.

EPIMEDIUM, [Tourn.] Linn. Syst. ed. I (1735).
BERBERIDEAE, Benth. & Hook. f. i. 44.
ENDOPLECTRIS, Raf. Fl. Tellur. iii. 66 (1836).
alpinum, *Linn. Sp. Pl.* 717.—Europ.
alpinum, Sibth. & Sm. Fl. Graec. t. 150 = pubigerum.
concinnum, *Vahl, in Regel, Gartenfl.* (1872) 268.—Japon.
diphyllum, Lodd. Bot. Cab. t. 1858 = Asteractylus diphyllus.
elatum, *Morr. & Decne. in Ann. Sc. Nat.* Bot. II. ii. (1834) 256.—Reg. Himal.

EPIMEDIUM:—
grandiflorum, Morr. in Belg. Hortic. ii. (1884) 141. t. 35. f. A = macranthum ?
hexandrum, Hook. Fl. Bor. Am. i. 30 = Vancouveria hexandra.
Hydaspidis, Falc. in Proc. Linn. Soc. i. (1889) 18 = elatum.
Ibarisa, Siebold, ex Regel, Ind. Sem. Hort. Petrop. (1866) 89 = macranthum ?
japonicum, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 71 = Acerantherus diphyllus.
longifolium, Decne. in Rev. Hortic. Sér. IV. iii. (1854) 241 = macranthum !
macranthum, *Morr. & Decne. in Ann. Sc. Nat* Sér II. ii. (1884) 352.—Japon.
Musschianum, *Morr. & Decne. l. c.* 358.—Japon
Pernideianum, *Cosa. in Bull. Soc. Bot. Fr.* ix. (1862) 167.—Algeria.
pinnatum, *Fisch. in DC. Syst.* ii. 29.—Persia.
pterocrus, *Morr. in Ann. Soc. Gand*, i. (1845) 145 t. 14.—Reg. Causas. !
pubescens, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 309.—Chin.
pubigerum, *Morr. & Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 355.—Byzant. ; Reg. Cauc.
rubrum, *E. Morr. in Belg. Hortic.* iii. (1853) 33. t 6. —Japon.
sagittatum, *Baker, in Gard. Chron.* (1880) i. 683.—Japon.
sinense, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 71 = sagittatum.
sorsicolor, E. Morr. in Belg. Hortic. iv. (1854) 54. t. (3) f. 6 = macranthum.
violaceum, Morr. & Decne. in Ann. Sc. Nat. Sér. II ii. (1834) 354 = macranthum.

EPIMENIDION, Rafin. Fl. Tellur. ii. 13 (1836) = **Stellia**, Linn. (Liliac.).
hyacinthoides, Rafin. l. c. = S. hyacinthoides.
undulatum, Rafin. l. c. = Urginea undulata.

EPIONIX, Rafin. Fl. Teller. ii. 31 (1836) = **Baccametra**, Salisb. (Liliac.).
fama, Rafin. l. c. = B. colonellaris.
rubra, Rafin. l. c. = B columellaris.

EPIPACTIS, Adans. Fam. ii. 70 (1763); R. Br. in Alt. Hort. Kew. v. 201 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 519.
abortiva, All. Fl. Pedem. ii. 151 = Limodorum abortivum.
alba, Crantz, Stirp. Austr. ed. II. fasc. vi. 460 = Cephalanthera ensifolia.
alpina, Schrank, Prim. Fl. Salisb. 90 = Herminium alpinum.
americana, Lindl. in Ann. Nat. Hist. iv. (1840) 385 = Royleana.
amœna, Buch.-Ham. ex Wall. Cat n. 7408 = consimilis.
Arachnites, F. W. Schmidt, in Mayer, Phys. Aufs. i. (1791) 249 = Ophrys Arachnites.
atherasis, Lej. Fl. Spa, ii. 196 = microphylla.
atropurpurea, Rafin. Caratteri, 87. n. 208 = latifolia.
atrorubens, Rostk. ex Spreng. Syst. iii. 707 = microphylla.
atrorubens, *Schult. Oestr. Fl.* ed. II. i. 58 ; *Reichb. Fl. Germ. Excurs.* 138.—Europ.; As. occ.
bahiniifolia, *Rosb. Hort. Beng* 63, nomen.—Ind. or.
bicarinata, Broth.-Ham. ex Wall. Cat. n. 7385 (= *Cymbopora bicarinata*).—Reg. Himal.
camtschatica, Sw. in Vet. Acad. Nya Handl. Stockh. (1800) 225 = Neottia camtschatica.
carinata, Roxb. Hort. Beng. 63 ; Fl. Ind. iii. 454 = Pogonia carinata.
caudata, Poir. Encyc. Suppl. ii. 571 = Acianthus caudatus.
condensata, Boiss. Diagn. Ser. II. iv. 91 = latifolia.
consimilis, D. Don, Prod. Fl. Nep. 28 = latifolia.
consimilis, *Wall. Cat.* n. 7403.—Persia ; Reg. Himal.
convallarioides, Sw. in Vet. Acad. Nya Handl. Stockh. (1800) 223 = Listera convallarioides.
Corallorhiza, Crantz, Stirp. Austr. ed. II. fasc. vi. 464 = Corallorhiza innata.
cordata, All. Fl. Pedem. ii. 152 = Listera cordata.
cucullata, Labill. Nov. Holl. Pl. ii. 61. t. 211. f. 2 = Eriochilus autumnalis.
Dalhousiæ, Wight, Ic. t. 1782 = latifolia.
dimorica, Steinh. Bot. Mat. Med. iv. 261 = Spiranthes dimorica.

EPIPACTIS:—
ensifolia, F. W. Schmidt, in Mayer, Phys. Aufs. i. (1791) 251 = Cephalanthera ensifolia.
Epipogium, All. Auct. Fl. Pedem. 22 = Epipogum aphyllum.
Epipogum, Crantz, Stirp. Austr. ed. II. fasc. vi. 477 = prasc.
erecta, Sw. in Schrad. Neues Journ. i. (1805) 65 = Cephalanthera pallens.
exserta, Poir. Encyc. Suppl. ii. 571 = Acianthus exsertus.
falcata, Sw. in Schrad. Neues Journ. i. (1805) 66 = Cephalanthera falcata.
formicata, Poir. Encyc. Suppl. ii. 571 = Acianthus formicatus.
gigantea, Dougl. ex Hook. Fl. Bor. Am. ii. 202. t. 202 = Royleana.
grandiflora, Roxb. Hort. Beng. 63 ; Fl. Ind. iii. 456 = Spathoglottis pubescens.
grandiflora, Sm. Engl. Bot. t. 271 ; Gaud Fl. Helv. v. 406 = Cephalanthera pallens.
grandifolia, All. Fl. Pedem. ii. 152 = Cephalanthera ensifolia.
Helleborine, Crantz, Stirp. Austr. ed. II. fasc. vi. 467 = prasc.
herbacea, Lindl. in Royle, Illustr. Bot. Himal. 368 = latifolia.
intermedia, Schur, Enum. Pl. Transs. 649 = microphylla.
intrusa, Lindl. in Journ. Linn. Soc. i. (1857) 175 = latifolia.
Jullana, Roxb. Hort. Beng. 63 ; Fl. Ind. iii. 455 = Pogonia Juliana.
kamtschatica, Lindl. Gen. et Spec. Orch. 458 = Neottia camtschatica.
lancifolia, F. W. Schmidt, in Mayer, Phys. Aufs. i. (1791) 253 = Cephalanthera pallens.
latifolia, *All. Fl. Pedem.* ii. 151 ; *Sw. in Vet. Akad. Nya Handl.* Stockh. (1800) 222.—Europ. ; As. bor.
latifolia, Less. in Linnæa, ix. (1834) 156 = strorubens.
Lessonii, Virv. in Mém. Soc. Linn. Par. iv. (1826) 605 = Pogonia tetraphylla.
longifolia, All. Fl. Pedem. ii. 152 = palustris.
longifolia, Blume, Orch. Archip. Ind. 185 = Thunbergii.
lutea, Steud. Nom. ed. II. i. 561 = Chloraea Gaudichaudii
macropodia, Peterm. Fl. Bienitz, 21 = strorubens.
macrostachya, Lindl. in Wall. Cat. n. 7404 = latifolia.
media, Fries, Mant. ii. 54 = strorubens.
microphylla, Sieber, ex Nym. Consp. 688 = strorubens, microphylla. *Sw. in Vet. Akad. Nya Handl. Stockh.* (1800) 222.—As. Min. ; Reg. Caucas.
monophylla, F. W. Schmidt, in Mayer, Phys. Aufs. i. (1791) 245 = Microstylis monophylla.
Monorchis, F. W. Schmidt, l. c. 246 = Herminium Monorchis.
myodes, F. W. Schmidt, l. c. 248 = Ophrys muscifera.
Nidus-avis, Crantz, Stirp. Austr. ed. II. fasc. vi. 475 ; Sw. in Vet. Acad. Nya Handl. Stockh. (1800) 222 = Neottia Nidus-avis.
ochroleuca, Baumg. Enum. Stirp. Transs. iii. 174 = Cephalanthera pallens.
ovalifolia, Stokes, Bot. Mat. Med. iv. 292 = Listera ovata.
ovalis, Bab. in Engl. Bot. Suppl. t. 2884 = latifolia.
ovata, Crantz, Stirp. Austr. ed. II. fasc. vi. 473 = Listera ovata.
pallens, Sw. in Schrad. Neues Journ. i. (1805) 65 = Cephalanthera pallens.
pallida, Sw. in Vet. Acad. Nya Handl. Stockh. (1800) 229 = Cephalanthera pallens.
palustosa, F. W. Schmidt, in Mayer, Phys. Aufs. i. (1791) 245 = Malaxis paludosa.
palustris, *Crantz, Stirp. Austr.* ed. II. fasc. vi. 462.—Au. bor. ; Am. bor.
papillosa, *Franch. & Sav. Enum. Pl. Jap.* ii. 519.—Japon.
Phyllanthes, G. E. Sm. in Gard. Chron. (1852) 660 = latifolia.
Phyllanthus, Lindl. in Journ. Linn. Soc. i. (1857) 175 = prasc.
plicata, Roxb. Hort. Beng. 63 ; Fl. Ind. iii. 454 = Pogonia plicata.
porrifolia, Sw. in Vet. Acad. Nya Handl. Stockh. (1800) 228 = Microtis porrifolia.
pubescens, Pursh, Fl. Am. Sept. ii. 591 = Posthlevia glandulosa.
purpurata, Sm. Engl. Fl. iv. 41 = latifolia.

EPIPACTIS:—
purpurea, Crantz, Stirp. Austr. ed. II. fasc. vi. 457 = Cephalanthera rubra.
pycnostachys, *C. Koch, in Linnaea*, xxii. (1849) 289.—Bithynia.
refracta, Labill. Nov. Holl. Pl. ii. 60. t. 211. f. 1 = Chiloglottis diphylla.
repens, Crantz, Stirp. Austr. ed. II. fasc. vi. 473 = Goodyera repens.
Royleana, *Lindl. in Royle, Illustr. Bot. Himal.* 368.—Reg. Himal. ; Am. bor. occ.
rubiginosa, Crantz, Stirp. Austr. ed. II. fasc. vi. 467. 468 = strorubens.
rubra, F. W. Schmidt, in Mayer, Phys. Aufs. i. (1791) 253 = Cephalanthera rubra.
Salassia, Pers. Syn. ii. 518 = Liparis purpurascens.
salina, Schur, Enum. Pl. Transs. 650 = palustris.
scariosa, Buch.-Ham. in Wall. Cat. n. 3739 = Pachystoma senile.
semifolia, Peterm. in Flora, xxvii. (1844) 370 = latifolia.
spiralis, Crantz, Stirp. Austr. ed. II. fasc. vi. 478 = Spiranthes autumnalis.
Thunbergii, *A. Gray, in Perry, Exp. Jap.* ii. 319.—Japon.
trifervis, *Roxb. Hort. Beng.* [102] ; *Fl. Ind.* iii. 455. —Ins. Moluce.
uliginosa, Vest, in Flora, viii. (1825) 493 = latifolia.
varians, Crantz, Stirp. Austr. ed. II. fasc. vi. 468, 471 = latifolia.
veratrifolia, Boiss. Diagn. Ser. I. xiii. 11 = consimilis.
violacea, Bor. Fl. Centr. fr. ed. III. 651 = latifolia.
viridans, Crantz, Stirp. Austr. ed. II. fasc. vi. 467, 476 = latifolia.
viridiflora, Reichb. in Moessl. Handb. ed. II. ii. 1575 = strorubens.
viridiflora, Rupr. in Beitr. Pfl. Russ. Reich. iv. 88 = latifolia.
xiphophylla, Sw. in Schrad. Neues Journ. i. (1805) 65 = Cephalanthera ensifolia.

EPIPACTIS, Hall. Enum. Stirp. Helv. i. 277 (1742) = **Goodyera**, R. Br. (Orchid.).

EPIPACTUM, Rlfg. in Marb. Schrift. ii. (1881) 125 = **Epipactis**, Adans. (Orchid.).

EPIPETRUM, Phil. in Linnæa, xxxiii. (1864–65) 253 = **Dioscorea**, Linn.
humile, Phil. l. c. = D. humilis.

EPIPHANES, Blume, Bijdr. 421. t. 4 (1825) = **Gastrodia**, R. Br. (Orchid.).
javanica, Blume, l. c. = G. javanica.

EPIPHANES, Reichb. f. in Seem. Fl. Vit. 295 (1868) = **Didymoplexis**, Griff. (Orchid.).
micradenia, Reichb. l. c. = D. micradenia.
pachystomoides, F. Muell. Fragm. viii. 174 = Pogonia pachystomoides.
pallens, Reichb. f. in Seem. Fl. Vit. 296 = D. pallens.

EPIPHEGUS, Spreng. Neue Entdeck. i. 264 (1820) = **Epifagus**, Nutt. (Orobanch.).

EPIPHEJUS, Walp. Rep. iii. 457 (1845) = prasc.

EPIPHORA, Lindl. in Hook. Comp. Bot. Mag. ii. 201 (1836) = **Polystachya**, Hook. (Orchid.).
pubescens, Lindl. l. c. = P. pubescens.

EPIPHYLLUM, Haw. Syn. Pl. Succ. 197 (1812) partim ; Pfeiff. Enum. Diagn. Cact. 127 (1837) *CACTEAE*. [Benth. & Hook. f. i. 849. Schumannberera, Lem. in Rev. Hortic. Sér. IV. vii. (1858) 258.
Ackermanni, Haw. in Phil. Mag. (Aug. 1829) 109 = Phyllocactus Ackermanni.
Aitoni, Hort. ex Steud. Nom. ed. II. i. 561 = Cereus speciosus.
alatum, Haw. Suppl. Pl. Succ. 84 ; et in Phil. Mag. (Aug. 1829) 109 = Rhipsalis pachyptera, Swartziana.
Altensteinii, *Pfeiff. Enum. Diagn. Cact.* 128 = Bma.
anguliger, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1880 = Phyllocactus anguliger.
biformis, G. Don, l. c. 1878 = Phyllocactus biformis.
Bridgesii, Lem. Illustr. Hortic. viii. (1861) Misc. 5 = truncatum.

EPIPHYLLUM:—
caulorhizum, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1380 = Phyllocactus caulorhizus.
ciliare, Hort. ex Pfeiff. Enum. Diagn. Cact. 130 = Rhipsalis ramulosa.
cilistum, Hort. ex Pfeiff. l. c. = Rhipsalis ramulosa.
crenulatum, Hort. ex Pfeiff. l. c. = Rhipsalis rhombea.
crispatum, Haw. in Phil. Mag. (Feb. 1830) 109 = Rhipsalis crispata.
elegans, Cels, ex Foerst. Handb. Cact. ed. I. 446 = truncatum?
Guesneyri, *Houllet, in Rev. Hortic.* (1876) 230.—Hab.?
Hitchini, Hort. ex Steud. Nom. ed. II. i. 561 = Cereus splendidus.
Hookeri, Haw. in Phil. Mag. (Aug. 1829) 108 = Cereus Hookeri.
hybridum, Hort. ex Pfeiff. Enum. Diagn. Cact. 121 = Cereus speciosissimus.
Jenkinsonii, G. Don, Gen. Syst. iii. 170 = Phyllocactus phyllanthoides.
latifrons, Zucc. ex Pfeiff. Enum. Diagn. Cact. 126 = Phyllocactus latifrons.
oxypetalum, Haw. in Phil. Mag. (Aug. 1829) 109 = Phyllocactus latifrons.
phyllanthoides, Sweet, Hort. Brit. ed. I. 172 = Phyllocactus phyllanthoides.
Phyllanthus, Haw. Syn. Pl. Succ. 197 = Phyllocactus Phyllanthus.
platycarpum, Zucc. ex Pfeiff. Enum. Diagn. Cact. 131 = Rhipsalis platycarpum.
ramulosum, Hort. ex Pfeiff. l. c. 130 = Rhipsalis ramulosa.
rhombeum, Hort. ex Pfeiff. l. c. = Rhipsalis rhombea.
Ruckeri, Paxt. Mag. Bot. xii. (1846) 46 = truncatum.
Ruckerianum, Hort. ex Lem. Illustr. Hortic. viii. (1861) Misc. 5 = truncatum.
Russellianum, *Hook. Bot. Mag.* t. 3717.—Bras.
Smithianum, Marnock, Floricult. Mag. viii. t. 13 = truncatum.
speciosum, Haw. Suppl. Pl. Succ. 84 = Phyllocactus phyllanthoides.
splendens, Hort. ex Ann. Fl. et Pom. (1839) 848 (ex Sched. DC.) = seq.
splendidum, Paxt. Mag. Bot. i. (1841) 49 = Cereus splendidus.
truncatum, *Haw. Suppl. Pl. Succ.* 85.—Bras.
Vandesii v. G. Don, Gen. Syst. iii. 170.
violaceum, Cels, ex Foerst. Handb. Cact. ed. I. 446 = truncatum?

EPIPHYSTIS, Trin. Fund. Agrost. 97 (1820) = **Mariscus**, Gaertn. (Cyperac.).
ophiuroides, Trin. l. c. = M. havanensis.

EPIPOGION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = seq.

EPIPOGUM, Ehrh. Beitr. iv. (1789) 149 = seq.

EPIPOGON, Ledeb. Fl. Ross. iv. 77 (1853) = seq.

EPIPOGUM, S. G. Gmel. Fl. Sibir. i. 11. t. 2. f. 2 (1747). *ORCHIDEÆ*, Benth. & Hook. f. iii. 617.
CERATOSIA, Lindl. Gen. et Sp. Orch. 388 (1840).
GALERA, Blume, Bijdr. 416. t. 2 (1826).
PODANTHERA, Wight, Ic. v. 22. t. 1769 (1851).
aphyllum, *Sw. Summ. Veg. Scand.* (1814) 85; *Sw. Bot.* t. 519.—Europ.; As. bor.
Gmelini, Rich. in Mém. Mus. Par. iv. (1818) 58 = aphyllum.
Guilyyiti, F. Muell. Fragm. viii. 30 = sutana.
nutans, *Reichb. f. in Bonplandia*, v. (1857) 56.—As. et Afr. trop.; Austral.
roseum, Lindl. in Journ. Linn. Soc. i. (1857) 177 = sutana.

EPIPREMNUM, Schott, in Bonplandia, v. (1857) 45. *AROIDEÆ*, Benth. & Hook. f. iii. 998.
RAPHIDOPHORA, Hassk. Cat. Hort. Bog. Alt. 58, (1844) partim.
amplissimum, *Engl. in Engl. Bot. Jahrb.* i. (1881) 128.—N. Guinea.
asperatum, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 270.—Ins. Moluce.
Beccarii, Engl. l. c. 229 = Rhaphidophora Beccari.
elegans, *Engl. l. c.*—N. Guinea.

EPIPREMNUM:—
giganteum, *Schott, in Bonplandia*, v. (1857) 45.—Ind. or.
magnificum, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 270.—Ins. Celebes.
medium, *Engl. in A. DC. Monog. Phan.* ii. 250.—Java.
mirabile, *Schott, Gen. Aroid.* t. 79.—Java.
moluccanum, *Schott, in Mig. Ann. Mus. Lugd. Bat.* i. 190.—Ins. Amboina.
nobile, *Engl. in A. DC. Monog. Phan.* ii. 250.—Ins. Celebes.
Zippelianum, Engl. in Bot. Jahrb. i. (1881) 182 = asperatum.

EPIPRINUS, Griff. Notul. iv. 487 (1854). *EUPHORBIACEÆ*. Benth. & Hook. f. ii. 325.
malayanus, *Grigf. l. c.*—Malacca.

EPIRIXANTHES, Blume, Cat. Gew. Buitenz. 25 (1823) Salomonia, Lour. (Polygal.).
cylindrica, Blume, l. c. 82 = S. sphylla.
elongata, Blume, l. c.—Java.
linearis, Blume, l. c.—Java.
tenella, Hook. f. in Trans. Linn. Soc. xxiii. (1862) 158 = S. sphylla.

EPIRHIZANTHES, Benth. & Hook. f. Gen. i. 186 (1862) sphalm. = prasc.

EPIRHIZANTHUS, Endl. Gen. 738 (1839) = prasc.

EPIRIZANTHE, Blume, Fl. Jav. Praef. p. vii (1828) = prasc.

EPISANTHERA, Hochst. ex Nees, in DC. Prod. xi. 308 in syn. (1847) = **Paulo-Wilhelmia**, Hochst. (Acanthac.).

EPISCIA, Mart. Nov. Gen. et Sp. iii. 89 (1829). *GESNERACEÆ*, Benth. & Hook. f. ii. 1006.
ALSOBIA, Hanst. in Linnæa, xxvi. (1853) 207.
CREMNOPOLENIA, Benth. in Hook. Lond. Journ. v. (1846) 369.
CYRTODEIRA, Hanst. in Linnæa, xxvi. (1853) 207.
NAUTILOCALYX, Linden, Cat. ex Hanst. in Linnæa, xxvi. (1853) 207.
PARADRYMONIA, Hanst. in Linnæa, xxvi. (1853) 207.
PETRODEIRA, Hanst. in Linnæa, xxvi. (1853) 207.
SKIOPHILA, Hanst. l. c. (1853).
TRICHODRYMONIA, Gerst. in Vidensk. Selsk. Skrift. V. v. (1861) 113 (Gen. Centr. Am. 88).
bicolor, *Hook. Bot. Mag.* t. 4390.—N. Granat.
bracteascens, *Hanst. in Linnæa*, xxxiv. (1865–66) 351. —Am. trop.
chontalensis, *Hook. Bot. Mag.* t. 5925.—Nicaragua.
ciliosa, *Hanst. in Mart. Fl. Bras.* viii. 408.—Am. trop.
congesta, *Hanst. in Linnæa*, xxxiv. (1865–66) 347.—Mexic.
cupreata, *Hanst. l. c.* 340.—Nicaragua.
decumbens, *Mart. Nov. Gen. et Sp.* iii. 89. t. 216.—Bras.
elongata, *Ann. Fl. et Pom.* (1838) 188 (ex Sched DC.). —Hab.?
erythropus, *Hook. f. Bot. Mag.* t. 6219.—N. Granat.
fulgida, *Hook. f. l. c.* t. 6186.—N. Granat.
glabra, *Hanst. in Linnæa*, xxxiv. (1865–66) 349.—Am. trop.
guadalupensis, *Hanst. l. c.* 345.—Ind. occ.
hirsuta, *Hanst. l. c.* 351.—Am. trop.
hirtiflora, *Sprace, ex Hanst. in Mart. Fl. Bras.* viii. 402.—Bras.
hispida, *DC. Prod.* vii. 547.—Am. austr.
Hookeri, *Hanst. in Linnæa*, xxxiv. (1865–66) 350.—Am. trop.
lanuginosa, *Hanst. l. c.* 346.—Costa Rica.
lilacina, *Hanst. l. c.* 342.—Costa Rica.
longifolia, *Hanst. l. c.* 347.—Am. trop.
Luciani, *Linden & Fourn. in Illustr. Hortic.* xxiii. (1876) 43. t. 236.—N. Granat.
meltitrifolia, *Mart. Nov. Gen. et Sp.* iii. 42.—Ind. occ.; Bras.
minutoides, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 365.—Guiana.
ochrolenca, *Hanst. in Mart. Fl. Bras.* viii. 408.—Afr. trop.
picta, *Hanst. l. c.*—Am. trop.
pulchella, *Mart. Nov. Gen. et Sp.* iii. 89.—Ins. Teinit.
punctata, *Hanst. in Linnæa*, xxxiv. (1865–66) 343.—Am. trop.

EPISCIA:—
reptans, *Mart. Nov. Gen. et Sp.* iii. 89. t. 217.—Bras.
rosea, Mart. & Gal. in Bull. Acad. Bruz. ix. (1842) reimp. 8 = Tetranema mexicana.
splendens, *Hanst. in Linnæa*, xxxiv. (1865–66) 341.—N. Granat.
subacaulis, *Griseb. Fl. Brit. W. Ind.* 462.—Ind. occ.
tessellata, *Linden, ex Lem. Illustr. Hortic.* xvi. (1869) sub t. 607.—Peruv.
villosa, *Hanst. in Linnæa*, xxxiv. (1865–66) 348.—Guian.

EPISCOPIA, Moritz, ex Klotzsch, in Linnæa, xxiv. (1851) 42 in syn. = **Themistoclesia**, Klotzsch (Vaccin.).
pendula, Moritz, ex Klotzsch, l. c. = T. pendula.

EPISIPHIS, Rafin. Fl. Tellur. iii. 78 (1836) = **Hypericum**, Linn.
parvifolia, Rafin. l. c. = H. sagyptiacum.

EPISTEIRA, Rafin. Sylva Tellur. 90 (1838) = **Phyllanthus**, Linn. (Euphorb.).
coccinea, Rafin. l. c. = P. coccineus.

EPISTEMUM, Walp. in Linnæa, xiii. (1839) 473 = **Amphithales**, Eckl. & Zeyh. (Legumin.).
ferrugineum, Walp. l. c. = A. cuneifolia.

EPISTEPHIUM, Kunth, Syn. Pl. Aequin. i. 240 (1822). *ORCHIDEÆ*, Benth. & Hook. f. iii. 591.
amplexicaule, *Poepp. & Endl. Nov. Gen. et Sp.* i. 52. t. 91.—Peruv.
elatum, *H. B. & K. Nov. Gen. et Sp.* vi. t. 532.—N. Granat.
Frederici-Augusti, *Reichb. f. in Bonplandia*, ii. (1854) 97.—N. Granat.
laxiflorum, *Rodrig. Orch. Nov.* i. 173.—Bras.
monanthum, *Poepp. & Endl. Nov. Gen. et Sp.* 53. t. 93.—Peruv.
parviflorum, *Lindl. Gen. et Sp. Orch.* 433.—Ind. occ.
Portellianum, *Rodrig. Orch. Nov.* i. 173.—Bras.
racemosum, *Rodrig. l. c.*—Bras.
sclerophyllum, *Lindl. Gen. et Sp. Orch.* 433.—Bras.
sessiliflorum, *Lindl. Orchid. Linden.* 36.—N. Granat.
smilacifolium, *Reichb. f. in Linnæa*, xli. (1877) 65.—N. Caled.
speciosum, *Rodrig. in Rev. de Engenh.* 74.—Bras.
Williamsii, *Hook. f. in Bot. Mag.* t. 5465.—Bras.

EPISTYLIUM, Sw. Fl. Ind. Occ. ii. 1095. t. 23 (1800) = **Phyllanthus**, Linn. (Euphorbiac.).
axillare, Sw. l. c. 1097 = P. axillaris.
cauliflorum, Sw. l. c. 1099 = P. caulifloras.
cordifolium, Baill. Etud. Gén. Euph. 648 = P. Baillonianus.
floribundum, Thw. Enum. Pl. Zeyl. 258 = P. cinereus, affinis.
glaucescens, Baill. Etud. Gén. Euph. 648, nomen = P. pulcher.
latifolium, Thw. Enum. Pl. Zeyl. 258 = P. Baillonianus.
leptoclados, Hance, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 229 = P. leptoclados.
montanum, Thw. Enum. Pl. Zeyl. 258 = P. oreophilus.
phyllanthoides, Baill. Etud. Gén. Euph. 648, nomen = P. pulcher.
polyphyllum, Thw. Enum. Pl. Zeyl. 258 = P. anabaptistifolius.
pulchrum, Baill. Etud. Gén. Euph. 648 = P. pulcher.
Roxburghii, Baill. l. c. = P. Roxburghii.
septemlocum, Baill. l. c. = P. anabaptistatus.

EPITHECIA, Knowles & Westc. Fl. Cab. ii. (1838) 167. t. 17 = **Dichaea**, Lindl. (Orchid.).
glauca, Knowles & Westc. l. c. = D. glauca.

EPITHECIUM, Benth. & Hook. f. Gen. 899, 1289 (1883) = prasc.

EPITHEMA, Blume, Bijdr. 737 (1826). *GESNERACEÆ*, Benth. & Hook. f. ii. 1017.
ARKINIA, R. Br. in Wall. Pl. As. Rar. iii. 65. t. 268 (1832).
CARROCALYMMA, Zipp. in Alg. Konst.- en Letterbode, i. (1826) 197.
Benthami, *C. B. Clarke, in DC. Monog. Phan.* v. 159. —Ins. Philipp.

EPITHEMA:—
Brunonis, *Decne. in Nouv. Ann. Mus. Nat. Par.* iii. (1834) 404.—Ins. Timor.
carnosum, *Benth. Scroph. Ind.* 57.—Ind. or.
ceylanicum, Wight, Illustr. 182. t. 159 = carnosum.
diffusum, *Span. in Linnaea*, xv. (1841) 881.—Java.
Horsfieldii, *R. Br. in Wall. Pl. As. Rar.* iii. 66.—Java.
Roxburghii, *DC. Prod.* ix. 279.—Ins. Moluce.
sexstile, *Blume, Bijdr.* 787.—Java.
tenue, *C. B. Clarke, in DC. Monog. Phan.* v. 181.—Afr. trop.
triandrum, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 150.—Ins. Philipp.
violaceum, Span. in Linnaea, xv. (1841) 880 = Brunonis.
zeylanicum, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 492 = carnosum.

EPITHINIA, Jack, in Malay. Misc. i. (1820) n. v. 12 = **Scyphiphora**, Gaertn. (Rubiac.).
malayana, Jack, l. c. = S. hydrophyllacea.

EPITHYMUM, Opiz, Seznam, 40 (1852) = **Cuscuta**, Tourn. (Convolvulac.).
cuscutoides, Opiz, l. c. = C. Epithymum.

EPITRACHYS, C. Koch, in Linnaea, xxiv. (1851) 396 = **Onioua**, Linn. (Compos.).
adunca, C. Koch, l. c. 402 (= *Cirsium fallax*).—Reg. Caucas.
arachnoidea, C. Koch, l. c. 400 = C. arachnoideus.
caucasica, C. Koch, l. c. = C. horridus.
ciliata, C. Koch, l. c. 397 = C. ciliatus.
erythrolepis, C. Koch, l. c. 398 (= *Cirsium erythrolepis*).—Reg. Caucas.
Kormelii, C. Koch, l. c. = C. lappaceus.
lanceolata, C. Koch, l. c. 400 = C. lanceolatus.
lanifera, C. Koch, l. c. = C. laniferus.
lappacea, C. Koch, l. c. 399, in obs. = C. lappaceus.
leucocephala, C. Koch, l. c. 401 = C. leucocephalus.
longifolia, C. Koch, l. c.—Armenia.
macrobotrys, C. Koch, l. c. 399.—Armenia.
microcephala, C. Koch, l. c. 402 = Carbenia benedicta ?
munita, C. Koch, l. c. 398 = C. muniticus.
numoralis, C. Koch, l. c. 400 = C. lanceolatus.
ossetica, C. Koch, l. c. 398 = C. fimbriatus.
paniculata, C. Koch, l. c. 399 = C. lappaceus.
pontica, C. Koch, l. c. 398, in obs. (= *Cirsium polycephalum*).—Pontus.
propontica, C. Koch, l. c. in obs. = C. eriophorus.
rigida, C. Koch, l. c. 400 = C. arachnoideus.
serrulata, C. Koch, l. c. 397 = C. serrulatus.
Szovitsii, C. Koch, l. c. (= *Cirsium Szovitsii*).—Reg. Caucas.
tenuiloba, C. Koch, l. c. 400.—Armenia.
tricholoma, C. Koch, l. c. 398 (= *Cirsium tricholoma*).—Reg. Caucas.
vulgaris, C. Koch, l. c. 402, in obs. = C. italicus.

EPITRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 74 = **Angianthus**, Wendl. (Compos.).
cuspidatus, Turcz. l. c. 75 = A. demissus.

EPLATEIA, Rafin. Sylva Tellur. 87 (1838) = **Withania**, Pauq. (Solanac.).
arborescens, Rafin. l. c. = W. arborescens.

EPLEIENDA, Rafin. Sylva Tellur. 107 (1838) = **Eugenia**, Mich. (Myrtac.).
sinemariensis, Rafin. l. c. = E. sinemariensis.

EPURGA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = **Euphorbia**, Linn.
Lathyris, Fourr. l. c. = Euphorbia Lathyris.

ERACLISSA, Forsk. Fl. Aegypt. Arab. 208 (1775) = **Andrachne**, Linn. (Euphorb.).
hexagyna, Forsk. l. c. = A. telephioides.

ERACLYSSA, Soop. Introd. 289 (1777) = prec.

ERAGROSTIS, Host, in Gram. iv. 14 (1809); Beauv. Agrost. 70. t. 14. f. 11 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1186.
Coelachyrum, Nees, in Linnaea, xvi. (1842) 221.
Eragrostiella, Rafin. Neogenyt. 4 (1825).
Eragrostis, *Steud. Nom.* ed. II. i. 632 (1840).

ERAGROSTIS:—
Harpachne, Hochst.—Cf. Flora, xxiv. (1841), Intell. 20, nomen; et ex A. Rich. Tent. Fl. Abyss. ii. 431 (1850).
Macroblepharus, Phil. in Linnaea, xxix. (1857–58) 100.
Megastachya, Beauv. Agrost. 74, 167 (1812).
abortiva, *Steud. Syn. Pl. Gram.* 279.—Austral.
abessinica, Link, Hort. Berol. i. 192 = Poa abyssinica.
abyssinica, Schrad. in Linnaea, xii. (1838) 450 = prec.
aciculata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 406.—Bras.
acuminata, *Doell, in Mart. Fl. Bras.* ii. iii. 153.—Bras.
aculiflora, *Nees, Agrost. Bras.* 501.—Am. trop.
aegyptiaca, *Delile, Fl. Egypt.* 157. t. 4. f. 2.—Afr. bor.
affinis, Nees & Meyen, ex Steud. Syn. Pl. Gram. 265 = aciculata.
affinis, Steud. Nom. ed. II. i. 569 = Vahlii.
airoides, *Nees, Agrost. Bras.* 509.—Bras.
alba, *N. & C. Presl, Rel. Haenk.* i. 279.—Calif.
albescens, *Steud. Syn. Pl. Gram.* 269.—Afr. trop.
amabilis, *Wight & Arn. in Hook. & Arn. Bot. Beech. Voy.* 251.—Ind. or.; Malaya.
amboinensis, *Trin. ex Steud. Nom.* ed. II. i. 652.—Ins. Amboina.
amoena, *N. & C. Presl, Rel. Haenk.* i. 275.—Mexic.
apiculata, *Doell, in Mart. Fl. Bras.* ii. iii. 146.—Bras.
arabica, Jaub. & Spach, Illustr. iv. 81 = ciliaris.
articulata, *Nees, Agrost. Bras.* 502.—Bras.
aspera, *Nees, Fl. Afr. Austr.* 408.—Abyss.; Afr. austr.; Ind. or.
atropurpurea, Hochst. ex Steud. Syn. Pl. Gram. 267 = tenuifolia.
atrovirens, Lange, in Kjoeb. Vidensk. Meddel. (1860) 47 = papposa.
atrovirens, *Nees, Fl. Afr. Austr.* 400.—Afr. austr.
aturensis, *Trin. ex Steud. Nom.* ed. II. i. 562.—Venezuela.
aulacosperma, Steud. l. c. = papposa.
aurea, Steud. Syn. Pl. Gram. 267 = plumosa.
bagdadensis, Boiss. Fl. Orient. v. 581 = pilosa.
bahiensis, *Roem. & Schult. Syst.* ii. Mant. 318.—Reg. trop.
barbata, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 70. —Bonariensis.
benguelensis, *Peyr. in Sitzb. Acad. Wien, Math.-Nat.* xxxvii. (1860) 583.—Afr. trop.
beniensis, *Steud. Nom.* ed. II. i. 569.—Afr. trop.
Bergiana, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 70.—Afr. austr.
Berteroiana, Steud. Nom. ed. II. ii. 361, in syn., sphalm. Berterroiana.
bicolor, Boiss. Fl. Orient. v. 581 = pilosa.
bicolor, *Nees, Fl. Afr. Austr.* 407.—Afr. austr.
bifaria, *Steud. Syn. Pl. Gram.* 264.—Ind. or.
bifaria, Wight, ex Steud. Nom. ed. II. i. 562 = coromandelina.
biformis, *Benth. in Hook. Niger Fl.* 568.—Afr. trop.
Boivini, *Steud. Syn. Pl. Gram.* 426.—Madag.
Boryana, *Steud. Nom.* ed. II. i. 569.—Ins. S. Doming.; Borbonia.
brasiliana, Nees, Agrost. Bras. 210 = elegans.
brasiliensis, Nees, l. c. 497 = bahiensis, interrupta, *Beauv.*, Nees.
brevifolia, *Benth. in Hook. Ic.* sub t. 1368 (p. 51).—Ind. or.
brizantha, *Nees, Fl. Afr. Austr.* 411.—Afr. austr.
bromoides, *Coste, Supplem. Fl. Catal.* 81.—Hispan.
brizoides, *Nees, in Linnaea*, vii. (1832) 226.—Afr. austr.
brizoides, Roem. & Schult. Syst. ii. Mant. 319 = Nettalliana.
bromoides, *Steud. Syn. Pl. Gram.* 276.—Peruv.
Brownеana, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 305 = acq.
Brownei, *Nees, ex Steud. Nom.* ed. II. i. 562; *et Nees, Fl. Afr. Austr.* 396, in syn.—Geront. trop. et subtrop.
bulbilifera, Steud. Syn. Pl. Gram. 267 = Brownei.
Caelachyrum, Boiss. Fl. Orient. v. 584 = E. Coelachyrum.
Calotheca, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 414 = Briza Lamarckiana.
Cambessedиana, *Steud. Syn. Pl. Gram.* 269.—Afr. trop.
campestris, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 70.—Am. bor.

ERAGROSTIS:—
capensis, Trin. in Mém. Acad. Pétersb. Sér. VI. i (1831) 400 = briziodes.
capillaris, *Nees, Agrost. Bras.* 505.—Am. bor.; Ind. occ.
capillaris, *Steud. ex Doell, in Mart. Fl. Bras.* ii. iii. 120.—Bras.
capillifolia, *Nees, Fl. Afr. Austr.* 403.—Afr. austr.
candata, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 573.—Mexic.
caudata, Nees, Fl. Afr. Austr. 410, adn. = plumosa.
cernua, Steud. Syn. Pl. Gram. 266 = Brownei.
chaetophylla, *Steud. Syn. Pl. Gram.* 279.—Austral.
chalcantha, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i (1831) 401.—Afr. austr.
Chapelieri, Nees, Fl. Afr. Austr. 392 = elegans.
chiliensis, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 464 = virescens.
chloromelas, *Steud. Syn. Pl. Gram.* 271.—Afr. austr.
ciliaris, *Kunth, Enum. Pl.* i. 227. n. 9.—Am. trop.
ciliaris, *Link, Hort. Berol.* i. 192.—Reg. trop.; Afr. austr.
ciliata, *Nees, Agrost. Bras.* 513, obs. I.—Ind. or.
cinnaeformis, *Schrad. in Linnaea*, xii. (1838) 452.—Hab.?
Coelachyrum, *Benth. in Hook. Ic. Pl.* t. 1368.—Aegypt.; Arab.
cognata, Steud. Syn. Pl. Gram. 273 = pectinacea.
collina, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 413.—Rossia.
compacta, Steud. l. c. 275 = Vahlii.
compta, *Link, Hort. Berol.* i. 193.—Reg. Argent.
concinna, *Steud. Syn. Pl. Gram.* 279.—Austral.
condensata, *Steud. l. c.* 278.—Mexic.
conferta, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 409.—Am. trop.
confinis, Nees, ex Steud. Syn. Pl. Gram. 265 = amabilis ?
congesta, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 178.—Afr. trop.
conricta, [*Nees, in*] *Meyen, Reise*, i. 484, nomen.—Peruv.
coromandelina, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 415.—Ind. or.
cretacea, Nees, in Ann. Nat. Hist. i. (1838) 284 = coromandelina.
Cumingii, Steud. Syn. Pl. Gram. 266 = interrupta, *Beauv.*
curtipedicellata, *Buchi. in Proc Acad. Sc. Philad.* 1862 (1863) 97.—Am. bor.
curvula, *Nees, Fl. Afr. Austr.* 397.—Afr. austr.
cuspidata, Link, Hort. Berol. ii. 298 = inconstans.
cylindrica, Nees, ex Hook. & Arn. Bot. Beech. Voy. 251 = geniculata.
cylindrica, Steud. Syn. Pl. Gram. 267 = ciliata *vel* plumosa.
cylindriflora, *Hochst. in Flora*, xxxviii. (1855) 324.—Abyss.
cynosuroides, *Beauv. Agrost.* 71.—Ind. or.
cyperoides, *Beauv, l. c.* 70.—Afr. austr.
decidua, *Hochst. in Flora*, xxxviii. (1855) 324.—Abyss.
decipiens, Steud. Syn. Pl. Gram. 279 = Leptochloa chinensis.
delicatula, *Trin. in Bull. Sc. Acad. Pétersb.* Sér. VI. i. (1838) 70.—Bras.
depauperata, *Anderss. in Peters, Reise Mossamb. Bot.* 559.—Afr. trop.
descrticola, *Phil. Fl. Atac.* 55.—Chili.
despiciens, Roem. & Schult. Syst. Mant. ii. 318 = plumosa.
diandra, Kth. ex Boiss. Fl. Orient. v. 584, in syn. = interrupta, *Beauv.*
diandra, *Steud. Syn. Pl. Gram.* 279.—Austral.
diarrhena, Steud. l. c. 266 = interrupta, *Beauv.*
diffusa, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 97 = Parishii.
dinodeorum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 418.—Ins. Galapag.
diplachnoides, Steud. Syn. Pl. Gram. 268 = interrupta, *Doell.*
domingensis, *Steud. l. c.* 278.—Ins. S. Doming.
elegans, *Nees, Agrost. Bras.* 510.—Bras.
elegantula, Nees, ex Miq. Anal. Bot. Ind. ii. 27 = Brownei.
elongata, *Yacy. Relag. Gram.* t. 3.—Ind. or.
elytroblephara, *Steud. Syn. Pl. Gram.* 280.—Ins. Societ.?

ERAGROSTIS :—
equitans, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 413.—Ins. Sandvic.
eriopoda, *Benth. Fl. Austral.* vii. 648.—Australi.
erythrogyna, Nees, ex Steud. Syn. Pl. Gram. 273 = Frankii.
Euchron, *Steud. l. c.* 367.—Japon.; Java.
excelsa, *Griseb. Cat. Pl. Cub.* 237.—Mexic.
exinta, *Steud. Syn. Pl. Gram.* 279.—Austral.; N. Zel.
expansa, *Link, Hort. Berol.* i. 190.—Reg. Argent.
falcata, *Gaudich. in Freyc. Voy. Bot.* 508. t. 25.—Austral.
fasciculata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 403.—Afr. trop.
Fendleriana, *Steud. Syn. Pl. Gram.* 278.—Mexic.
ferruginea, *Beauv. Agrost.* 71.—Japon.
filiformis, *Link, Hort. Berol.* i. 191.—Am. bor.
filiformis, Nees, in Linnæa, vii. (1832) 330 = Thunbergiana.
firma, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 70.—Bras.
flavescens, J. & C. Presl, Rel. Haenk. i. 278 = poæoides.
flexuosa, Steud. Syn. Pl. Gram. 266 = minor!
floribunda, Roem. & Schult. Syst. ii. Mant. 819 = Vahlii.
Frankii, Steud. Syn. Pl. Gram. 273.—Am. bor.
gangetica, Steud. l. c. 266 = major!
geniculata, Dur. in Expl. Sc. Algér. t. 40. f. 4 = papposa!
geniculata, *Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl.* i. (1843) 203.—As. trop.
Geyeri, *Steud. Syn. Pl. Gram.* 272.—Am. bor.
glaucina, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 403.—Ins. S. Doming.
glabrata, *Nees, in Linnæa*, vii. (1832) 332.—Afr. austr.
glaucoon, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 406 = articulata.
glumacea, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 578.—Mexic.
glutinosa, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 397.—Jamaica.
gracilis, *Schrad. in Linnæa*, xii. (1838) 451.—Chili.
guineensis, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 404, in nota = fascicularis!
gummiflua, *Nees, Fl. Afr. Austr.* 803.—Afr. austr.
Haenkei, *J. & C. Presl, Rel. Haenk.* i. 279.—Mexic.
hapalantha, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 409, partim = interrupta, *Beauv.*, Haenkeii, *Steud. Syn. Pl. Gram.* 265.—Java.
hirsuta, *Nees, Agrost. Bras.* 508.—Am. bor.
hirta, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 573.—Mexic.
Hochstetteri, *Steud. Syn. Pl. Gram.* 268.—Nubia.
homomalla, *Nees, Fl. Afr. Austr.* 406.—Afr. austr.
Hornemanniana, *Nees*, 398.—Afr. trop.
imbecilla, *Benth. Fl. Austral.* vii. 643.—Austral.
imbecillis, Steud. Syn. Pl. Gram. 279 = Leptochloa chinensis.
incrassata, Nees, Agrost. Bras. 495 = Vahlii.
indica, Link, Hort. Berol. i. 192, in syn. = Poa abyssinica.
indica, Steud. Syn. Pl. Gram. 264 = pilosa.
inflexa, Steud. Nom. ed. II. i. 553.—N. Granat.
interrupta, *Beauv. Agrost.* 71.—Afr., As, et Austral. trop.
interrupta, *Doell, in Mart. Fl. Bras.* ii. iii. 157 (sp. diversa).—Bras.
Jacquini, *Schrad. in Linnæa*, xii. (1838) 450.—Hab. ?
japonica, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 405 = plumosa.
Jardinei, *Steud. Syn. Pl. Gram.* 269.—Afr. trop. ?
Kunthii, Link, Hort. Berol. ii. 394, in syn. = interrupta, *Beauv.*
lacunaria, *F. Muell. First Gen. Report* (1858) 20.—Austral.
Lamarckii, *Steud. Syn. Pl. Gram.* 269.—Afr. trop.
laniflora, *Benth. Fl. Austral.* vii. 648.—Austral.
Lappula, *Nees, Fl. Afr. Austr.* 413.—Afr. austr.
lata, *Baker, Fl. Maurit.* 456.—Ins. Mascar.
laxiflora, *Schrad. in Linnæa*, xii. (1838) 451.—Afr. austr.
laxiloloides, *Guss. Fl. Sic. Syn.* i. 108; *Steud. Syn. Pl. Gram.* 263.—Sicil.

ERAGROSTIS :—
Lehmanniana, *Nees, Fl. Afr. Austr.* 402.—Afr. austr.
lepida, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 454 = plumosa.
Leprieurii, *Steud. Syn. Pl. Gram.* 269.—Afr. trop.
leptostachya, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 406.—Bras.
leptocarpa, *Benth. Fl. Austral.* vii. 644.—Austral.
leptostachya, *Steud. Syn. Pl. Gram.* 279.—Austral.
leucosticta, *Nees, ex Doell, in Mart. Fl. Bras.* ii. iii. 144.—Bras.
limbata, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 578.—Mexic.
Lindeniana, *Steud. Syn. Pl. Gram.* 278.—N. Granat.
linearis, *Benth. in Hook. Niger Fl.* 567.—Afr. trop.
linearis, *Trin. ex Steud. Nom.* ed. II. i. 553.—China.
Linkii, *Steud. l. c.* 273.—Am. bor.
lobata, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 396 = cylindrica.
longeradiata, *Steud. Syn. Pl. Gram.* 272.—Am. bor.
longifolia, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 32, nomen; *et in A. Rich. Tent. Fl. Abyss.* ii. 455.—Abyss.
lugens, *Nees, Agrost. Bras.* 507.—Am. trop.
lugens, Rupr. in Bull. Acad. Brux. ix. (1842) relmp. 9 = capillaris.
luzula, *J. & C. Presl, Rel. Haenk.* 276.—Peruv.
luxoniensis, Steud. Syn. Pl. Gram. 266 = Brownei.
macilenta, *Steud. l. c.* 268.—Abyss.
madagascariensis, *Steud. l. c.* 272.—Madag.
major, *Host, Gram.* iv. t. 34.—Reg. trop. et subtrop.
mangalorica, Hochst. ex Miq. Anal. Bot. Ind. ii. 26 = pilosa.
Mauritii, *Steud. Syn. Pl. Gram.* 272.—Ins. Mascar.
maxima, *Baker, in Journ. Linn. Soc.* xxi. (1885) 454.—Madag.
maxima, *Fourn. ex Hemsl. Biol. Am. Centr. Bot.* iii. 574.—Mexic.
mayparensis, *Steud. Syn. Pl. Gram.* 276; *Doell, in Mart. Fl. Bras.* ii. iii. 158.—Am. austr.
megalosperma, *F. Muell. ex Benth. Fl. Austral.* vii. 644.—Austral.
megapotamica, Roem. & Schult. Syst. ii. Mant. 617 = purpurascens.
megastachya, Link, Hort. Berol. i. 187 = major.
Meratiana, *Steud. Syn. Pl. Gram.* 276.—Bras.
mexicana, *Link, Hort. Berol.* i. 190.—Mexic.
microstachya, Link, l. c. ii. 394 = virgata.
milleflora, *Steud. Syn. Pl. Gram.* 265.—Ins. Timor.
Millettii, *Hook. & Arn. Bot. Beech. Voy.* 383.—China.
minor, *Host, Fl. Austr.* i. 135.—Reg. trop. et temp.
minutiflora, *J. & C. Presl, Rel. Haenk.* i. 274.—Ind. or.
Montufari, *Steud. Syn. Pl. Gram.* 275.—Ecuador.
mosambicensis, Steud. l. c. 264 = pilosa, interrupta, *Doell.*
mucronata, *Roem. & Schult. Syst.* ii. 877.—Ind. or.
multicaulis, Steud. Syn. Pl. Gram. 266 = pilosa.
multiflora, Aschers. ex Boiss. Fl. Orient. v. 580 = virgata.
multiflora, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 401 = amabilis!
nanegumensis, Hochst. ex Steud. Syn. Pl. Gram. 264 = pilosa.
nanegumensis, Nees, ex Schrad. in Linnæa, xii. (1838) 453 = interrupta, *Beauv.*
nardoides, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 415 = coromandelica.
Neesii, *Trin. l. c.* 405.—Bras.
nervosa, *Hochst. in Flora*, xxxviii. (1855) 327.—Abyss.
Newrikii, *Hance, in Journ. Bot.* xviii. (1880) 302.—China.
nigra, *Nees, ex Steud. Nom.* ed. II. i. 553; *et Syn. Pl. Gram.* 267.—Afr. trop.; Ind. or.; Austral.
nigricans, *Steud. Nom. l. c.*—Ecuador.
nindensis, *Planchu & Hieron, in Trans. Linn. Soc. Ser.* II. ii. (1881) 93.—Afr. trop.
nitida, *Chapm. Fl. S. U. St.* 564.—Florida.
nitida, *Link, Hort. Berol.* i. 195.—Aegypt.
nodosa, Boiss. Fl. Orient. v. 585 = interrupta, *Beauv.*
nodosa, Nees, ex Steud. Nom. ed. II. i. 553 = Brownei.
Nortuliana, *Steud. Nom.* ed. II. i. 553.—Amer.

ERAGROSTIS :—
obtusa, *Munro, ex Planchu & Hieron, in Trans. Linn. Soc. Ser.* II. ii. (1881) 92.—Afr. trop.
Olmedi, *Steud. Nom.* ed. II. i. 554.—Peruv.
orientalis, Trin. in Mém. Acad. Pétersb. Sér. VI. iv. ii. (1838) Suppl. 72 = elongata.
orina, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 458 = Brownei.
oxuriensis, Steud. Syn. Pl. Gram. 269 = beninensis.
oxylepis, *Torr. in Rep. Bot. Whipple Exped.* 156.—Mexic.
Palmeri, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 83 183.—Mexic.
panamensis, J. & C. Presl, Rel. Haenk. 277 = mayparensis.
paniciformis, *Steud. Syn. Pl. Gram.* 268.—Abyss.
panicoides, *Steud. l. c.* 278 (sp. dub.).—Mexic.
paniculata, Steud. l. c. = saperr.
papposa, *Steud. Nom.* ed. II. i. 554.—Reg. Mediterr.; Orient.
parviflora, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 411 = pilosa.
parviglumis, Hochst. ex Steud. Syn. Pl. Gram. 267 = nigra!
parvula, Steud. l. c. = Parishii.
pastoënsis, *Trin. in Bull. Sc. Acad. Pétersb.* (1836) 70.—N. Granat.
patens, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 178.—Afr. trop.
patula, *Steud. Nom.* ed. II. i. 554.—Ecuador.
pauciflora, *Trin. fils Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 575.—Am. bor.; Mexic.
pectinacea, *Nees, Fl. Afr. Austr.* 408 in syn.—Mexic
pellucida, Steud. Syn. Pl. Gram. 279 = pilosa.
pensylvanica, *Scheele, in Flora*, xxvii. (1844) 56.—Am. bor.
perennis, *Doell, in Mart. Fl. Bras.* ii. iii. 144.—Bras.
peruviana, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 396.—Am. austr.
Petersii, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 70.—Ins. Tahiti.
Pierrii, *Benth. in Hook. Ic. Pl.* t. 1370.—Baluchist.
pilifera, *Schenk, in Linnæa*, xxii. (1849) 344.—Texas.
pilosa, *Beauv. Agrost.* 71.—Reg. calid.
pilosiuscula, Link, Hort. Berol. i. 189 = articulata.
plana, *Nees, Fl. Afr. Austr.* 390.—Afr. austr.
planiculmis, *Nees, l. c.* 391.—Afr. austr.
plumosa, *Link, Hort. Berol.* i. 192.—As. et Afr. trop.
poæformis, Link, l. c. 188 = minor.
poæoides, Beauv. Agrost. 71. t. 14 = minor.
pogonia, Steud. Syn. Pl. Gram. 267 = Brownei.
polymorpha, Roem. & Schult. Syst. ii. 876 = bahiensis.
polymorpha, Steud. Syn. Pl. Gram. 265 = major.
polymorpha, Trin. ex Steud. Nom. ed. i. 554, 562 = amabilis.
polytricha, *Nees, Agrost. Bras.* 507.—Bras.
porosa, *Nees, Fl. Afr. Austr.* 401.—Afr. austr.
procera, Steud. Syn. Pl. Gram. 266 = Brownei!
procumbens, *Nees, Fl. Afr. Austr.* 396.—Afr. austr.
prolifera, *Steud. Syn. Pl. Gram.* 177.—Ind. occ.
propinqua, *Steud. l. c.* 271.—Ins. Mauriti.
psammodes, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 400.—Bras.
puberula, *Steud. Syn. Pl. Gram.* 263.—Abyss.
pubescens, Steud. l. c. 279 = Brownei.
pulchella, Parl. ex Webb, in Hook. Niger Fl. 188 = pilosa.
punctata, *Link, ex Steud. Nom.* ed. II. i. 561.—Ind. or.
purpurascens, *Roem. & Schult. Syst.* ii. *Mant.* 817.—Bras.
Purshii, *Hort. ex Schrad. in Linnæa*, xii. (1838) 451.—Am. bor.
quitensis, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 70.—Am. austr.
racemosa, *Steud. Syn. Pl. Gram.* 271.—Afr. austr.
rachitricha, Hochst. ex Miq. Anal. Bot. Ind. ii. 25 = tenuila.
reptans, *Nees, Agrost. Bras.* 514.—Reg. trop.
retinervulosa, Steud. Syn. Pl. Gram. 268 = pilosa.
rhachitricha, Hochst. ex Miq. Anal. Bot. Ind. ii. 25 = tremula.
riparia, Nees, Agrost. Bras. 512, pdn. = ciliata.
riparia, Steud. Syn. Pl. Gram. 265 = plumosa.
Rothii, Link, Hort. Berol. ii. 394 = interrupta, *Beauv.*
rubens, Hochst. ex Miq. Anal. Bot. Ind. ii. 24 = elongata.
rubens, Steud. Syn. Pl. Gram. 265 = amabilis.

ERAGROSTIS:—
rubiginosa, *Trin. in Mém. Acad. Pétersb* Sér. VI. i. (1831) 401.—Afr. trop.
rufescens, Roem. & Schult. Syst. Mant. ii. 819 = inconstans.
rupestris, Steud. Syn. Pl. Gram. 365 = ciliata.
Salzmanni, Steud. l. c. 277 = prolifera.
sarmentosa, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 398.—Afr. austr.
saxatilis, *Hemsl. Bot. Voy. Challenger*, i. 11. 90.—Ins. S. Helena.
scabra, *Phil. Fl. Atac.* 55.—Chili.
scaligera, Steud. Syn. Pl. Gram. 277 = Vahlii.
Schimperi, *Benth. in Hook. Ic. Pl.* t. 1371.—Abyss.
Schnittii, *Benth. Fl. Austral.* vii. 646.—Austral.
sclerantha, *Nees, Fl. Afr. Austr.* 388.—Afr. austr.
sclerophylla, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 412.—Bras.
secunda, Nees, ex Steud. Syn. Pl. Gram. 264 = coromandelina.
secundiflora, *J. & C. Presl, Rel. Haenk.* i. 276 (sp. dub.).—Mexic.
seminuda, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 406.—Bras.
senegalensis, *Nees, Sem. Hort. Vratisl.* (1834).—*Cf. Linnæa*, xii. (1838) 452.—Afr. trop.
sessiliflora, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 97 = Diplachne rigida.
setifolia, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 419 = chaetophylla.
simpliciflora, *Steud. Syn. Pl. Gram.* 278 (sp. dub.).—Mexic.
solida, Nees, Agrost. Bras. 501 = rufescens.
spartinoides, *Steud. Syn. Pl. Gram.* 265.—Ins. Phillip.
speciosa, Steud. l. c. 279 = Brownei.
spectabilis, *Steud. Nom.* ed. II. i. 564.—Am. bor.
spiralostachya, Cosn. & Dur. ex Lange, in Kjoeb. Vidensk. Meddel. (1860) 47 = papposa.
spinosa, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 415.—Afr. austr.
squamata, *Steud. Syn. Pl. Gram.* 274.—Bras.
stenoclada, *J. & C. Presl, Rel. Haenk.* i. 278.—Peruv.
stenophylla, Hochst. ex Miq. Anal. Bot. Ind. ii. 27 = Brownei.
stenostachya, *Steud. Syn. Pl. Gram.* 279.—Austral.
striata, Nees, Fl. Afr. Austr. 385 = Bergiana.
stricta, Steud. Syn. Pl. Gram. 264 = Brownei.
strigosa, *Anderss. in Peters, Reise Mossamb. Bot.* 559. —Afr. trop.
suaveolens, Becker, ex Claus, in Beitr. Pflanz. Russ Reich. viii. 266 = collina.
subulata, *Nees, Fl. Afr. Austr.* 399.—Afr. austr.
sudana, *Griseb. Cat. Pl. Cub.* 237.—Cuba.
superba, *Payr. in Sitzb. Acad. Wien, Math.-Nat.* xxxviii. (1860) 584.—Afr. trop.
Taffanga, *Steud. Syn. Pl. Gram.* 268.—Abyss.
tenax, *Steud. Nom.* ed. II. i. 564.—Ecuador.
tenella, Benth. Fl. Hongk. 481 = interrupta, *Beauv.*
tenella, Nees, Fl. Afr. Austr. 405 = pilosa.
tenella, Roem. & Schult. Syst. ii. 676 = plumosa.
tenellula, Steud. Syn. Pl. Gram. 279 = plumosa.
tenuiflora, *Rupr. ex Steud. l. c.* 268.—Afr. trop.
tenuifolia, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 20, nomen; *et in A. Rich. Tent. Fl. Abyss.* i. 425.—Abyss.; Ind. or.
tenuis, *A. Gray, Man. Bot. N. U. St.* ed. V. 632.—Am. bor.
tenuis, Steud. Syn. Pl. Gram. 273 = pilosa.
tenuissima, Schrad. ex Nees, Fl. Afr. Austr. 410 = interrupta, *Beauv.*
tephrosanthos, *Roem. & Schult. Syst. Mant.* ii. 816.—Ins. Martin.
tephrosanthos, Steud. Syn. Pl. Gram. 278 = limbata, Parishii.
Thunbergiana, *Steud. Nom.* ed. II. i. 564.—Afr. austr.
thyrsiflora, *Willd. & Lange, Prod. Fl. Hisp.* i. 83.—Hispan.
tremula, *Hochst. in Flora*, xxv. (1842) 1. *Beibl.* 134, nomen; *et ex Steud. Syn. Pl. Gram.* 269.—Gerront. trop.
trichophora, Coss. & Dur. in Bull. Soc. Bot. Fr. ii. (1855) 313 = papposa?
trichophylla, *Benth. Fl. Austral.* vii. 644.—Austral.
Trinii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 414 = Teicidia capensis.
trilicea, *Steud. Nom.* ed. II. i. 564.—Sicil.

ERAGROSTIS:—
unineria, *Steud. Syn. Pl. Gram.* 278 (sp. dub.).—Mexic.
Unioloa, *Nees, Agrost Bras.* 494.—Bras.
unioloides, *Nees, ex Steud. Syn. Pl. Gram.* 264.—Ins. Banca.
Uniola, Steud. l. c. 278 = pectinacea.
Urvillei, Steud. l. c. 279 = Brownei?
Vahlii, *Nees, Agrost. Bras.* 499.—Bras.
variabilis, Gaudich. in Freyc. Voy. Bot. 408 = wahoowensis.
velutina, *Schrad. in Linnæa*, xii. (1838) 451.—Am. bor.
Verae-Crucis, Rupr. in Bull. Acad. Brux. ix. (1842) reimp. 9 = oxylepis.
verticillata, Beauv. Agrost. 71; Roem. & Schult. Syst. ii. 678 = pilosa.
verticillata, Cosn. ex Lange, in Kjoeb. Vidensk. Meddel. (1860) 47 = papposa.
villosa, Steud. Syn. Pl. Gram. 276 = velutina.
villosa, Trin. Fund. Agrost. 137 = ciliaris.
virescens, *J. & C. Presl, Rel. Haenk.* i. 276.—Chili.
Virletii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 476.—Mexic.
viscosa, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i (1831) 397.—Ax. trop.
vulgaris, Presl, ex Steud. Nom. ed. II. i. 561 = major?
wahoowensis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 412.—Ins. Sandvic.
Wightiana, *Benth. in Hook. Ic. Pl.* t. 1881.—Ind. or.
Wildenowiana, Nees, ex Hook. & Arn. Bot. Beech. Voy. 252; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 205 = minor.
Willdenowii, Nees, ex Steud. Nom. ed. II. i. 565 = pyrasa.
zeylanica, Manro, ex Hance, in Journ. Linn. Soc. xiii. (1873) 187 = bahiensis.
zeylanica, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 204 = elongata.

ERANGELIA, Renealm. ex Linn. Sp. Pl. ed. I. 288 (1753) = Gaiianthus, Linn. (Araryllid.).

ERANTHEMUM, Linn. Diss. Dass. 1 (1747); Amoen. Acad. i. 384 (1749). *ACANTHACEAE*, Benth. & Hook. f. ii. 1097.
FIGARETTA, Adans. Fam. ii. 223 (1763).
SALZINGANTHA, Hort. ex Lem. in Fl. des Serres, Sér. I. iii. (1847) 240.
UPUDALIA, Rafin. Fl. Tellur. iv. 66 (1836).
acanthoides, *Spreng. Syst. i.* 88.—Ins. S. Doming.
acanthophorum, Roem. & Schult. Syst. i. Mant. 154 = Haplanthus verticellaris.
aciculare, R. Br. ex Roem. & Schult. Syst. i. 177 = Anthacanthus jamaicensis.
acuminatiseimum, *Miq. Fl. Ind. Bat.* iii. 835.—Java.
affine, Spreng. Syst. i. 89 = hypocrateriforme.
alatum, *Nees, in DC. Prod.* xi. 460.—Mexic.
albiflorum, *Hook. Bot. Mag.* t. 4225.—Bras.
album, *Nees, in DC. Prod.* xi. 455.—Burma; Penang.
ambiguum, Roem. & Schult. i. Mant. 154 = Anthacanthus acicularis.
amoenum, Boch.-Ham. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 420 = Daedalacanthus suffruticosus.
anderedi, *Mast. in Gard. Chron.* (1869) 134, 136.—Malaya.
angustatum, Linn. Mant. 171 = Agathelpis angustifolia.
angustifolium, Murr. Syst. ed. XIII. 55 = Agathelpis angustifolia.
armatum, R. Br. ex Roem. & Schult. Syst. i. 176 = Anthacanthus armatus.
asperrum, *Hook. f. Bot. Mag.* t. 5711.—Ins. Salomon. atropurpureum, *Hort.—Cf. Gard. Chron.* (1875) i. 619. —Polynes.
barlerioides, Roxb. Hort. Beng. 3; Fl. Ind. i. 113 = Daedalacanthus suffruticosus.
bicolor, *Schrank, Pl. Rar. Hort. Monac.* t. 8.—Ins. Luzon.
bifarium, *Nees, in DC. Prod.* xi. 458.—Burma.
Blumei, Hassk. Cat. Hort. Bog. Alt. 151 = crenulatum.
Blumei, Teijsm. in Miq. Fl. Ind. Bat. ii. 886 = Andersoni.
bornense, *Hook. f. Bot. Mag.* t. 6701.—Borneo.
capense, Linn. Sp. Pl. 9 = Daedalacanthus nostanus.
Carruthersii, *Seem. Fl. Vit.* 185.—Ins. Fiji.
cassabarinum, *Wall. Pl. As. Rar.* i. 20. t. 21.—Burma.

ERANTHEMUM:—
coccineum, Lem. in Fl. des Serres, Sér. I. iii. (1847) 240 = Thyrsacanthus Lemairianus.
Cooperi, *Hook. Bot. Mag.* t. 5467.—N. Caled.
corcovadense, *Regel, Petrop. Ind. Sem. Hort.* (1865) 47.—Bras.
cordatum, *Nees, in Benth. Bot. Voy. Sulph.* 147.—Panama.
crenulatum, *Lindl. in Bot. Reg.* t. 879.—Reg. Himal.
crenulatum, Nees, in Wall. Pl. As. Rar. iii. 107 = malacoense.
crenulatum, Wall. Cat. n. 2491 = malabaricum.
crenulatum, Wall. i. c. D, E = album.
crenulatum, Wall. i. c. F = pulchellum.
cristatum, *Humb. & Bonpl. ex Spreng. & Link, Jahrb.* i. 111. 46.—Am. austr.
cubense, Spreng. ex Nees, in DC. Prod. xi. 736 = Barleriola solanifolia.
cuspidatum, *Nees, in Benth. Pl. Hartw.* 147.—Mexic.
decurrens, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 43, nomen; *et in DC. Prod.* xi. 458.—Abyss.
detruncatum, *Nees & Mart. in Nov. Act. Acad. Nat Cur.* xi. (1823) 64.—Bras.
diantherum, Blume, Bijdr. 793 = crenulatum.
diantherum, *Roxb. Hort. Beng.* 3; *Fl. Ind.* i. 112.—Ins. Molucc.
diversifolium, *Miq. Fl. Ind. Bat.* iii. 837.—Java.
Echolium, T. Anders. in Thw. Enum. Pl. Zeyl. 235 = Echolium Linnaeanum.
Edgeworthianum, Nees, in DC. Prod. xi. 44 = Daedalacanthus nervosus.
elatum, Kurz, in Journ. Bot. x. (1872) 46 = pulchellum.
elegans, *R. Br. ex Roem. & Schult. Syst.* i. 174.—Afr. trop.
elegans, Mast. in Gard. Chron. (1868) 1234 = Andersoni.
enarginatum, *Desf. ex Spreng. & Link, Jahrb.* i. III. 47.—Ins. S. Thomas.
erectum, *Hort. Bogor. ex T. Anders. in Journ. Agri-Hort. Soc. Ind.* N. S. i. (1868) 386.—Hab. ?
exsequatum, *Nees, in Mart. Fl. Bras.* iv. 158.—Bras.
fasciculatum, Blume, Bijdr. 793 = bicolor.
fasciculatum, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 166.—Am. centr.
fastigiatum, R. Br. ex Roem. & Schult. Syst. i. 174; Spreng. Syst. i. 89 = Daedalacanthus nostanus.
filiforme, Willd. ex Nees, in DC. Prod. xi. 451 = cuspidatum.
flavum, Hort. ex Steud. Nom. ed. II. i. 565 = Ruellia flava.
flavum, Willd. Enum. Hort. Berol. Suppl. 2 = Barleria flava.
foecundum, *Lindl. in Bot. Reg.* t. 1494.—Bras.
gracilliforum, *Nees, in Wall. Pl. As. Rar.* iii. 107.—Penang.
grandiflorum, Zipp. ex Span. in Linnaea, xv. (1841) 328 = bicolor.
Haenkeanum, *Nees, in DC. Prod.* xi. 457.—Ins. Luzon.
heterophyllum, *Nees, l. c.*—Bras.
hispidulum, *Nees, l. c.* 456.—Mexic.
hispidum, *Nees, l. c.* 456.—Afr. trop.
hypocrateriforme, *R. Br. ex Roem. & Schult. Syst.* i. 175.—Afr. trop.
hyssopifolium, Herb. Linn. ex T. Anders. in Journ. Linn. Soc. vii. (1864) 118 (non Acanthac.).—Hab. ?
ignanum, *Linden, in Fl. des Serres*, vii. (1847–48) 19. —Peruv.
indicum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 497.—Reg. Himal.
insolarum, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 349.—Ins. Fiji.
lanceum, *Nees, in DC. Prod.* xi. 451.—N. Granat.
lapathifolium, *Nees, l. c.* 454.—Ind. or.
latifolium, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) 11. 79 = pulchellum.
laxiflorum, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 349.—Ins. Fiji.
Leptanthus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 500.—Reg. Himal.
leptocalathyrus, *Nees, in Mart. Fl. Bras.* ix. 157.—Bras.
leuconeuron, Hort. ex Regel, Gartenfl. (1856) 291. t. 174 = seq.
leuconeurum, *Hort. Petrop. ex Schlecht. in Linnaea*, xxvii. (1854) 500; *Fisch. ex Regel, Ind. Sem. Hort. Petrop.* (1858) 48.—Bras.
longifolium, *Regel, Ind. Sem. Hort. Petrop.* (1865) 47. —Am. austr.

ERANTHEMUM:—
longifolium, *Sesm. Fl. Vit.* 185.—N. Caled.
lycioides, *Desf. ex Sprong. & Link, Fabrb.* i. 111. 47.—Ind. occ.
macrophyllum, *Nees, in Mart. Fl. Bras.* ix. 157.—Bras.
macrophyllum, Wall. Cat. n. 7179 = Dasdalacanthus macrophyllus.
macrostachyum, Mart. ex Nees, in DC. Prod. xi. 385 = Stenandrium serpens.
malabaricum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 497.—Ind. or.
malaccense, *C. B. Clarke, l. c.* 498.—Malacca.
marmoratum, *Hort. ex Regel, Ind. Sem. Hort. Petrop.* (1865) 48.—Am. austr.
micranthum, Hort. Berol. ex Nees, in DC. Prod. xi. 456 = variabile.
microphyllum, R. Br. ex Roem. & Schult. Syst. i. 176 = Asthacanthus microphyllus.
modestum, *Nees, ex Mart. in Nov. Act. Nat. Cur.* xi. i. (1832) 68.—Bras.
montanum, *Hook. Bot. Mag.* t. 4031.—Ind. or.
montanum, Roxb. Hort. Beng. (80); Fl. Ind. i. 100 = Dasdalacanthus montanus.
montanum, Russ. ex Nees, in DC. Prod. xi. 449 = Wightianum.
montanum, Wall. Cat. n. 2492 = Dasdalacanthus purpurascens.
Moorei, *Hort. ex Gard. Chron.* (1875) i. 619.—Polynes.
nervosum, R. Br. ex Roem. & Schult. Syst. i. 174 = Dasdalacanthus nervosus.
nervosum, Dalz. & Gibs. Bomb. Fl. 195 = Dasdalacanthus purpurascens.
nigritianum, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 51.—Afr. trop.
nigrum, *Linden*, in *Illustr. Hortic.* xxvii. (1880) t. 404.—Ins. Salomon.
obovatum, E. Mey. ex Nees, in DC. Prod. vii. 462 = Calophanes Persoonii.
palustiforum, Hook. Bot. Mag. t. 5957 = malaccense, cinnabarinum.
palustiforum, *Nees, in Wall. Pl. As. Rar.* iii. 108.—Reg. Himal.
palustiforum, Steud. Nom. ed. II. i. 5 = praec.
Parishii, T. Anders. in Journ. Linn. Soc. ix. (1867) 489 = Dasdalacanthus Farishii.
Parishii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 449.—Burma.
parviflorum, Berg. Desc. Pl. Cap. 2 = Agathelpis parviflila.
parviflorum, *R. Br. ex Sprong. Syst.* i. 88.—Mexic.
parviflorum, Willd. Sp. Pl. i. 52 = Justicia salviaeflora.
parvifolium, Link, Enum. Hort. Berol. i. 28 = Asthacanthus aciculatis.
parvifolium, Linn. Mant. 171 = Agathelpis parvifolia.
pelagicum, *Seem. Fl. Vit.* 185.—Ins. N. Hebrid.
praecox, *Benth. Pl. Hartweg.* 291.—Mexic.
pubescens, *Roth, Nov. Pl. Sp.* 1.—Ind. or.
pulchellum, Andr. Bot. Rep. t. 88 = Dasdalacanthus nervosus.
pulchellum, Hort. in Gastenaag. (1810) 176. t. 17; ex Nees, in DC. Prod. xi. 456 = bicolor.
pulchellum, Roxb. Hort. Beng. 8; Fl. Ind. i. 111 = Dasdalacanthus purpurascens.
punctatum, *Nees, in DC. Prod.* xi. 455.—Ind. or.
purpurascens, Wight, ex Nees, in Wall. Pl. As. Rar. iii. 106 = Dasdalacanthus purpurascens.
racemosum, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 113.—Ins. Moluc.
repandum, *R. Br. ex Roem. & Schult. Syst.* i. 175.—Ins. Tanna.
reticulatum, *Hort. ex Gard. Chron.* (1875) i. 619.—Polynes.
Riedelianum, *Nees, in Mart. Fl. Bras.* ix. 156.—Bras.
roseum, R. Br. Prod. i. 477 = Dasdalacanthus roseus.
roseum, *Linden & Pourn. in Illustr. Hortic.* xxiii. (1876) t. 285.—Bras.
salaccense, *Blume, Bijdr.* 792.—Java.
salsoloides, Linn. f. Suppl. 92 = Campylanthus salsoloides.
sanguinolentum, *Hort. ex Veitch, in Fl. des Serres*, Sér. II. v. (1849) 187. t. 1488.—Madag.
scabrum, Wall. ex T. Anders. in Journ. Linn. Soc. ix. (1867) 487 = Dasdalacanthus nervosus.

ERANTHEMUM:—
Schomburgkii, Hort. in Illustr. Hortic. xxvi. (1879) t. 949 = reticulatum.
semperflorens, *Roth, Nov. Pl. Sp.* 2.—Ind. occ.
senense, *Klotzsch, in Peters, Reise Mossamb. Bot.* 219.—Afr. trop.
serpyllifolium, Willd. ex Roem. & Schult. Mant. i. 145 = Andrographis serpyllifolia.
sessile, *Nees, in Mart. Fl. Bras.* ix. 156.—Bras.
smaillinum, *Regel & Hord. Ind. Sem. Hort. Petrop.* (1861) 39.—Bras.
sinuatum, R. Br. ex Roem. & Schult. Syst. i. 176 = Asthacanthus sinuatus, repandus.
spinosum, R. Br. l. c. 176 = Asthacanthus spinosus.
spinosum, Lour. Fl. Cochinch. i. 19 = Hapianthus verticillaris.
strictum, Colebr. ex Roxb. Fl. Ind. ed. Carey, i. 114 = Dasdalacanthus strictus.
succisifolium, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii 149 = cinnabarinum.
suffruticosum, Roxb. Hort. Beng. 8; Fl. Ind. i. 111 = Dasdalacanthus suffruticosus.
Teijsmanni, *T. Anders. in Journ. Agri-Hort. Soc. Ind. N. S.* i. (1855) 325.—Malaya.
tenellum, *Benth. Fl. Austral.* iv. 555.—Austral.
tetragonum, *A. Dietr. ex Nees, in DC. Prod.* xi. 454.—Ind. occ.
tetragonum, Wall. Cat. n. 7180 = Dasdalacanthus tetragonus.
tricolor, *Nichols. Dict. Gard.* i. 518.—Polynes.
tuberculatum, *Hook. f. Bot. Mag.* t. 5406.—Hab. ?
variabile, *R. Br. Prod.* 477.—Austral.
varians, Blllb. in Linnæa Samfund. i. 1822 (1823) 11 = Dasdalacanthus nervosus.
verbenaceum, *Nees, ex Mart. in Nov. Act. Nat. Cur.* xi. i. (1825) 65.—Bras.
versicolor, *Hort.—Cf. Gard. Chron.* (1875) i. 619.—Hab. ?
verticillatum, Spreng. Syst. i. 88 = Hapianthus verticillare.
viscidum, R. Br. ex Roem. & Schult. Syst. i. 176 = Asthacanthus vincoides.
viscidum, *Blume, Bijdr.* 793.—Java.
Wightianum, Wall. Cat. n. 2495 = Dasdalacanthus montanus.
Zollingerianum, *Nees, in DC. Prod.* xi. 455.—Ins. Philipp.

ERANTHIS, Salisb. in Trans. Linn. Soc. viii. (1807) 303. *RANUNCULACEÆ*, Benth. & Hook. f. i. 7. HELLEBOROIDES, Adans. Fam. ii. 458 (1763). KOELLIA, Biria, Dis. Renonc. 21 (1811).
cilicica, *Schott & Kotschy, in Oesty. Bot. Wochenbl.* (1854) 118.—Cilicia.
hyemalis, *Salisb. in Trans. Linn. Soc.* viii. (1807) 304. —Europ.
Keiskei, *Franch. & Sav. Enum. Pl. Jap.* ii. 269.—Japon.
longistipitata, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) i. 244.—As. bor.
pinnatifida, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 238.—Japon.
sibirica, *DC. Syst.* i. 315.—Sibir.
stellata, *Maxim. Prim. Fl. Amur.* 22.—Mandshur.
uncinata, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 55.—Sibir.

ERANTHUS, G. Don, Gen. Syst. i. 45 (1831) = prasc.

ERASMA, R. Br. in Abel, Narr. Journ. Chin. 374 (1818), nomen (Brunlac.).

ERASMIA, Miq. in Inst. Veral. en Med. 1842 (1843) 81 = Peperomia, Ruiz & Pav. (Piperac.).
floribunda, Miq. l. c. = P. Erasmia.

ERATO, DC. Prod. v. 317 (1836) = Idaburea, Adans. (Compos.).
polymnioides, DC. l. c. 318.—Peruv.

ERATOBOTRYS, Fenzl, ex Endl. Gen. Suppl. ii. 18 (1842) = Stellia, Linn. (Liliac.).
bifolia, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 326 = S. Richardiana.
illaciua, Fenzl, ex Kunth, Enum. Pl. iv. 379 = S. Blacina.

ERBLICHIA, Scem. Bot. Herald, 130. t. 27 (1853) = Piriqueta, Aubl. (Turnerac.).
madagascariensis, O. Hoffm. Sert. Madag. (1881) 19 = P. madagascariensis.
odorata, Seem. Bot. Voy. Herald, 130. t. 27 = P. odorata.

ERCILLA, Endl. Gen. 977 (1840) = seq.

ERCILLA, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 11. t. 5. *PHYTOLACCACEÆ*, Benth. & Hook. f. iii. 84.
APODOSTACHYS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 577.
BRIDGESIA, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 168. t. 102.
ERCILIA, Endl. Gen. 977 (1840).
SURIANA, Domb. ex Meisn. Gen. Comm. 235 (1841).
spicata, *Moq. in DC. Prod.* xiii. 11. 35.—Chili.
volubilis, *A. Juss. in Ann. Sc. Nat.* Sér. I. xxv. (1832) 11. t. 5. f. 1.—Peruv.

EREBENNUS, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 14 = Hibiscus, Linn. (Malvac.).
canarensus, Alef. l. c. = H. tetraphyllus.

EREBINTHUS, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) 210, App. = Tephrosia, Pers. (Legumin.).

ERECHTHITES, Less. Syn. Comp. 395 (1832) = seq.

ERECHTITES, Raña. Fl. Ludov. 65 (1817). *COMPOSITÆ*, Benth. & Hook. f. ii. 448.
NECCIA, Cass. in Bull. Soc. Philom. (1820) 90.
PTILERIA, Rafin. in Am. Monthly Mag. (1818) 268.
ambigua, DC. Prod. vi. 295 = praecalta.
asperifolia, Nand. in Linnæa, xxv. (1852) 524 = arguta.
arguta, *DC. Prod.* vi. 296.—Austral.; N. Zel.
Atkinsoniae, *F. Muell. Fragm.* v. 88.—Austral.
Bathurstiana, DC. Prod. vi. 297 = arguta.
cacalioides, Less. Syn. Comp. 395 = praealta.
candicans, Hook. f. in Hook. Lond. Journ. Bot. vi (1847) 199 = Senecio quercifolius.
carduifolia, DC. Prod. v. 294 = praealta.
elongata, A. Gray, Syn. Fl. N. Am. i. ii 396 = praealta.
glabrescens, DC. Prod. vi. 295 = quadridentata.
glabrescens, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 550.—N. Zel.
glandulosa, DC. Prod. vi. 295 = quadridentata.
glomerata, DC. l. c. 297 = arguta ?
glossanthe, Nand. in Linnæa, xxv. (1852) 524 = Senecio brachyglossus.
Gunnii, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 122 = quadridentata.
hieracifolia, Rafin. ex DC. Prod. vi. 294 = praealta.
hispidula, *DC. l. c.* 296.—Austral.
ignobilis, *Baker, in Mart. Fl. Bras.* vi. iii. 299.—Bras.
incana, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii 55 = quadridentata.
lacerata, F. Muell. in Linnæa, xxv. (1852) 617 = arguta.
minima, *DC. Prod.* vi. 427, in syn.—Austral.
mixta, *DC. l. c.* 297.—Austral.
Muelleri, Lange, ind. Sem. Hort. Haun. (1861) 22 = praecalta.
argumentosa, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 490 = valerianaefolia.
paniculata, *Benth. Pl. Hartw.* 209.—N. Granat.
pleridioides, Nand. & Mudl. in Linnæa, xxv. (1852) 523 = mixta.
pleridioides, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 200 = prenanthoides.
praealta, *Rafin. Fl. Ludov.* 65; *Less. in Linnæa*, vi. (1831) 411.—Am. bor. et austr.
prenanthoides, *DC. Prod.* vi. 296.—Austral.
pumila, *Arnott, in Trans. N. Z. Inst.* xiii. (1881) 338. —N. Zel.
pumila, DC. Prod. vi. 297 = arguta ?
pyrrophylla, *Sch. Bip. ex Miq. Pl. Ind. Bat.* ii. 96.—Java.
quadridentata, *DC. Prod.* vi. 295.—Austral.

ERECHTITES :—
Richardiana, DC. Prod. vi. 297 = hispidula.
runcinata, *DC. l. c.* 295.—Mexic.
scabrella, *Hook. f. Handb. N. Zeal. Fl.* 157.—N. Zel.
sonchoides, DC. Prod. vi. 296 = prenanthoides.
suicata, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 419 = prenella.
tenuiflora, DC. Prod. vi. 296 = quadridentata.
valerianæfolia, *DC. l. c.* 295.—Bras.

EREMÆA, Lindl. Swan River App. 11 (1839). *MYRTACEÆ*, Benth. & Hook. f. i. 707.
scutifolia, *F. Muell. Fragm.* ii. 30.—Austral.
beaufortioides, *Benth. Fl. Austral.* iii. 103.—Austral.
chractesta, *F. Muell. Fragm.* ii. 29.—Austral.
cricifolia, Lindl. Swan River App. 11 = pilosa.
fimbriata, *Lindl. l. c.*—Austral.
pilosa, *Lindl. l. c.*—Austral.
violacea, *F. Muell. Fragm.* xi. 10.—Austral.

EREMANTHE, Spach, Hist. Vég. Phan. v. 421 (1836); et in Ann. Sc. Nat. Sér. II. v. (1836) 368 = Hypericum, Linn.
calycina, Spach, ll. cc. 423; 368 = H. calycinum.
patula, C. Koch, Hort. Dendrol. 65 = H. patulum.

EREMANTHUS, Less. in Linnæa, iv. (1829) 317. *COMPOSITÆ*, Benth. & Hook. f. ii. 385.
CHRESTA, Vell. Fl. Flum. viii. t. 150, 151 (1827), nomen prius.
LAGOPTALUM, Pohl, ex Baker, in Mart. Fl. Bras. vi. 11. 168 (1873).
PRESTELIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 386 (1873).
PYCNOCEPHALUM, DC. Prod. v. 88 (1836).
SPHÆROPHORA, Sch. Bip. in Pollichia, xx-xxi. (1863) 402.
STACHYANTHUS, DC. Prod. v. 84 (1836).
angustifolius, *Baker, in Mart. Fl. Bras.* vi. 11. 170.—Bras.
bicolor, *Baker, l. c.* 185.—Bras.
cinctus, *Baker, l. c.* 163.—Bras.
crotonoides, Sch. Bip. in Pollichia, xx-xxi. (1863) 396 = Vernonia crotonoides.
Elæagnus, *Sch. Bip. l. c.* xviii-xix. (1861) 166.—Bras.
eriopus, *Baker, in Mart. Fl. Bras.* vi. 11. 169.—Bras.
exsuccus, *Baker, l. c.* 166.—Bras.
glomerulatus, *Less. in Linnæa*, iv. (1829) 317.—Bras.
goyazensis, *Sch. Bip. in Pollichia*, xviii-xix. (1861) 165.—Bras.
incanus, *Less. in Linnæa*, vi. (1831) 662.—Bras.
Martii, *Baker, in Mart. Fl. Bras.* vi. 11. 167.—Bras.
mollis, *Sch. Bip. in Pollichia*, xviii-xix. (1861) 166.—Bras.
pallidiortus, Sch. Bip. l. c. 165 = glomerulatus.
pandurifolius, *Baker, in Mart. Fl. Bras.* vi. 11. 169.—Bras.
pannosus, *Baker, l. c.* 164.—Bras.
plantaginifolius, *Baker, l. c.* 166.—Bras.
pycnocephalus, *Baker, l. c.* 166.—Bras.
scapigerus, *Baker, l. c.* 168.—Bras.
speciosus, *Baker, l. c.* 169.—Bras.
sphærocephalus, *Baker, l. c.* 167.—Bras.
stellatus, Sch. Bip. in Pollichia, xviii-xix. (1861) 164 = glomerulatus.
verbascifolius, Sch. Bip. l. c. xx-xxi. (1863) 397 = Vernonia crotonoides.
Weddellii, Sch. Bip. l. c. xviii-xix. (1861) 165 = goyazensis.

EREMIA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 156. *ERICACEÆ*, Benth. & Hook. f. ii. 592.
HEXASTEMON, Klotzsch, in Linnæa, xii. (1838) 230.
Barklingiana, *Klotzsch, in Linnæa*, xii. (1838) 218.—Afr. austr.
brevifolia, *Benth. in DC. Prod.* vii. 700.—Afr. austr.
lanata, *Benth. l. c.*—Afr. austr.
parviflora, *Klotzsch, in Linnæa*, xii. (1838) 498.—Afr. austr.
recurvata, *Klotzsch, l. c.*—Afr. austr.
totta, *D. Don, in Edinb. N. Phil. Journ.* xvii. (1834) 156.—Afr. austr.
tuberosularia, *Benth. in DC. Prod.* vii. 700.—Afr. austr.

EREMIASTRUM, A. Gray, in Mem. Am. Acad. Sér. II. v. (1855) 320 (Pl. Nov. Thurb.). *COMPOSITÆ*, Benth. & Hook. f. ii. 370.
bellidiforme, *A. Gray, l. c.* 321.—Calif.

EREMITES, Benth. in Journ. Linn. Soc. xix. (1881) 52 = eeq.

EREMITIS, Doell, in Mart. Fl. Bras. ii. 11. 338. t. 48 (1877) = **Pariana**, Aubl. (Gramin.).
monothalamia, Doell, l. c. = P. parviflora.

EREMOBIUM, Boiss. Fl. Orient. i. 156 (1867) = **Malcolmia**, R. Br. (Crucif.).
linoare, Boiss. l. c. 157 = M. segyptiaca.

EREMOCALLIS, Salisb. ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 598 (1821) = **Erica**, Tourn.
cinerea, S. F. Gray, l. c. = E. cinerea.
glomerata, S. F. Gray, l. c. = E. Tetralix.

EREMOCARPUS, Benth. Bot. Voy. Sulph. 53. t. 26 (1844). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 297.
setigerus, *Benth. l. c.*—Calif.

EREMOCARPUS, Lindl. Veg. Kingd. 778 (1847), sphalm. = EREMODAUCUS, Benge = **Trachydium**, Lindl. (Umbellif.).

EREMOCHARIS, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 78 (1849) = **Citanthus**, Soland. (Legumin.).

EREMOCHARIS, Phil. Fl. Atac. 25. t. 2 B (1861) = **Asteriscotum**, Cham. (Umbell.).
flexuosa, Phil. f. Cat. Pl. Vasc. Chil. 103, sphalm. = seq.
fruticosa, Phil. Fl. Atac. 25.—Chili.

EREMOCHLOA, Busse, in Miq. Pl. Jungh. 357 (1851-54) = **Panicum** (Wall. Gramin.).
Horsferi, Busse, l. c.—Sumatra.

EREMOCHLOË, S. Wats. in Rot. King's Exped. 282. t. 40 (1871). *GRAMINEÆ*, Benth. & Hook. f. iii 1177.
Bigelovii, *S. Wats. l. c.* 282.—N. Mexic.
Kingii, *S. Wats. l. c.* t. 40.—Calif.
Thurberi, S. Wats. l. c. f. 1 = Bigelovii.

EREMODAUCUS, Bunge, Del. Sem. Hort. Dorp. (1843). —Cf. Linnæa, xvii. (1844) 151 = **Trachydium**, Lindl. (Umbellif.).
Lehmanni, Bunge, tl. cc. 6; et 151.—Turkestan.

EREMODENDRON, DC. ex Meisn. Gen. Comm. 201 = **Bremophila**, R. Br. (Myoporin.).
Cunninghamii, DC. ex Meisn. Gen. Comm. 201 = Bremophila oppositifolia.

EREMOGONE, Fenzl, Verbreit. Alsin. 18. tab. ad p. 18 (1888) = **Arenaria**, Linn. (Caryophyll.).
aizoifolia, Fenzl, ex Steud. Nom. ed. II. i. 566 = A. lychnides.
capillaris, Fenzl, Verbreit. Alsin. 87 = A. capillaris.
Cephalotes, Fenzl, l. c. t. ad p. 46 = A. Cephalotes.
cucubaloides, Hohen. in Bull. Soc. Nat. Mosc. (1838) 401 = A. cucubaloides.
dawurica, Fenzl, Verbreit. Alsin. 87 = A. juncea.
formosa, Fenzl, l. c. = A. capillaris.
Gmelinii, Fenzl, l. c. t. ad p. 18 = Gypsophila Gmelinii.
graminea, C. A. Mey. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 7 = A. graminea.
graminifolia, Fenzl, Verbreit. Alsin. 87 = A. graminifolia, *Schrad.*
gypsophiloides, Fenzl, l. c. t. ad p. 57 = A. gypsophiloides.
Holostea, Rupr. Fl. Caucas. 217 = A. Holostea.
juncea, Fenzl, Verbreit. Alsin. 87 = A. juncea.
longifolia, Fenzl, l. c. = A. saxatilis.
lychnoides, Rupr. Fl. Caucas. 219 = A. lychnides.
micrantha, Bohnt. Enum. Pl. Transs. 913, Index = A. graminifolia, *Schrad.*
nardifolia, Fenzl, Verbreit. Alsin. 87 = A. capillaris.
ottitesiana, Fenzl, l. c. = A. ottitesides.
procera, Reichb. fo. Fl. Germ. v. 88 = A. graminifolia, *Schrad.*
pungens, Fenzl, Verbreit. Alsin. t. ad pp. 57 et 59 = A. pungens.
rigida, Fenzl, l. c. t. ad p. 46 = A. rigida.
stenophylla, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 7 = A. graminifolia, *Schrad.*
subulata, Fenzl, Verbreit. Alsin. 87 = A. subulata.

EREMOLÆNA, Baill. in Bull. Soc. Linn. Par. i. (1884) 413. *CHLÆNACEÆ*.
Humbloniana, *Baill. l. c.*—Madag.

EREMOLEPIS, Griseb. in Goett. Abh. vi. (1854) 134. *LORANTHACEÆ*, Benth. & Hook. f. iii. 215.
IXIDIUM, Eichl. in Mart. Fl. Bras. v. ii. 130. t. 31 (1868).
punctulata, *Griseb. l. c.*—Chili.
verrucosa, *Griseb. l. c.*—Chili.
Wrightii, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 102.—Cuba.

EREMOPANAX, Baill. Adansonia, xii. (1878) 158. *ARALIACEÆ*.
angustata, *Baill. l. c.* 159.—N. Caled.
Balansæ, *Baill. l. c.* 160.—N. Caled.
otopyrena, *Baill. l. c.* 158.—N. Caled.
Vieillardi, *Baill. l. c.* 161.—N. Caled.

EREMOPHILA, R. Br. Prod. 518 (1810). *MYOPORINEÆ*, Benth. & Hook. f. ii. 1195.
EREMODENDRON, DC. ex Meisn. Gen. 392 (1840).
STENOCHILUS, R. Br. Prod. 517 (1810).
adenotricha, F. Muell. ex Benth. Fl. Austral. v. 15 = Pholidia adenotricha.
albicans, F. Muell. in Proc. Roy. Soc. Tasm. ii. (1859) 297 = Brownii.
alternifolia, *R. Br. Prod.* 518.—Austral.
arborescens, A. Cunn. ex A. DC. in DC. Prod. xi. 718 = oppositifolia.
Beckeri, F. Muell. Fragm. i. 156 = Myoporum Beckeri.
Behriana, F. Muell. in Proc. Roy. Soc. Tasm. iii. (1859) 296 = Pholidia Behriana.
Berryi, *F. Muell. Fragm.* viii. 228.—Austral.
bignoniiflora, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 294.—Austral.
Bowmani, *F. Muell. Fragm.* ii. 189.—Austral.
brevifolia, F. Muell. Census, 104 = Pholidia brevifolia.
Brownii, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 297.—Austral.
Christophori, *F. Muell. Fragm.* ix. 120.—Austral.
Clarkei, *Oldfield & F. Muell. l. c.* i. 208.—Austral.
crassifolia, F. Muell. in Proc. Roy. Soc. Tasm. iii. (1859) 297 = Pholidia crassifolia.
Cunninghamii, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 84 = oppositifolia.
Dalyana, F. Muell. Fragm. v. 23 = Pholidia Dalyana.
Delisseriæ, F. Muell. l. c. 108. t. 41 = Pholidia Delisseri.
Dempsteri, *F. Muell. l. c.* v. 60.—Austral.
densifolia, F. Muell. l. c. ii. 160 = Pholidia densifolia.
denticulata, *F. Muell. l. c.* i. 125.—Austral.
divaricata, F. Muell. in Trans. Roy. Soc. Tasm. iii. (1859) 298 = Pholidia divaricata.
Drummondi, *F. Muell. Fragm.* vi. 147.—Austral.
Duttoni, *F. Muell. Rep. Babb. Exped.* 16.—Austral.
Elderi, *F. Muell. Fragm.* viii. 228.—Austral.
eriocalyx, *F. Muell. l. c.* i. 226.—Austral.
exilifolia, *F. Muell. l. c.* x. 88.—Austral.
Forrestii, *F. Muell. l. c.* vii. 49.—Austral.
Fraseri, *F. Muell. l. c.* xi. 51.—Austral.
Freelingii, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 295.—Austral.
glabrifolia, F. Muell. Rep. Babb. Exped. 18 = Pholidia glabrifolia.
Gibsoni, *F. Muell. Fragm.* viii. 227.—Austral.
Gilesii, *F. Muell. l. c.* 49.—Austral.
Goodwinii, *F. Muell. Rep. Babb. Exped.* 17.—Austral.
gracilifolia, *F. Muell. Fragm.* i. 208.—Austral.
Grahamii, F. Muell. in Proc. Roy. Soc. Tasm. iii. (1859) 297 = Brownii.
Hughesii, *F. Muell. Fragm.* viii. 228.—Austral.
imbricata, F. Muell. Census, 104 = Pholidia imbricata.
incana, F. Muell. in Proc. Roy. Soc. Tasm. iii. (1859) 297 = Brownii.
Laanii, *F. Muell. in Melb. Chemist* (Jan. 1886).—Austral.
latifolia, *F. Muell. in Linnæa*, xxv. (1852) 423.—Austral.
Latrobei, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 294.—Austral.
leucophylla, *Benth. Fl. Austral.* v. 18.—Austral.
longifolia, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 295.—Austral.
Macdonnelli, *F. Muell. Rep. Babb. Exp.* 18.—Austral.

EREMOPHILA:—
Mackinlayi, *F. Muell. Fragm.* iv. 80.—Austral.
maculata, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 397.—Austral.
Maitlandi, *F. Muell. ex Benth. Fl. Austral.* v. 19.—Austral.
microtheca, F. Muell. ex Benth. l. c. 14 = Phyllodia microtheca.
Mitchelli, *Benth. in Mitch. Trop. Austral.* 31.—Austral.
myoporoides, F. Muell. Fragm. v. 28 = Myoporum laxiflorum.
Oldfieldii, *F. Muell. l. c.* t. 208.—Austral.
oppositifolia, *R. Br. Prod.* 518.—Austral.
Palalayi, *F. Muell. Rep. Babb. Exped.* 17.—Austral.
Pantoni, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 261.—Austral.
platycalyx, *F. Muell. Fragm.* v. 109.—Austral.
polyclada, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 294.—Austral.
racemosa, F. Muell. l. c. 297 = maculata.
resinosa, F. Muell l. c. 296 = Phyllodia resinosa.
rotundifolia, *F. Muell. Fragm.* i. 207.—Austral.
sanctifiae, F. Muell. in Proc. Roy. Soc. Tasm. iii. (1859) 295 = Phyllodia santalina.
scoparia, F. Muell. l. c. 296 = Phyllodia scoparia.
strongylophylla, *F. Muell. Fragm.* x. 87.—Austral.
Sturtii, *R. Br. in Sturt, Exped. Centr. Austral.* ii. *App.* 85.—Austral.
subfloccosa, *Benth. Fl. Austral.* v. 28.—Austral.
tuberculata, F. Muell. in Proc. Roy. Soc. Tasm. iii. (1859) 294 = Latrobei.
Turtoni, *F. Muell. Fragm.* x. 87.—Austral.
viscida, *Endl. in Ann. Wien. Mus.* ii. (1838) 208.—Austral.
viscosa, Stend. Nom. ed. II i. 566, sphalm. = præc.
Weldii, *F. Muell. Fragm.* vii. 109.—Austral.
Willisii, *F. Muell. l. c.* iii. 21. t. 20.—Austral.
Woollsiana, F. Muell. l. c. i. 195. t. 7 = Phyllodia Woollsiana.
Youngii, *F. Muell. l. c.* x. 16.—Austral.

EREMOPYRUM, Jaub. & Spach, Illustr. iv. 26 (1850–53) = **Agropyron**, J. Gaertn. (Gramin.).
cristatum, Willk. & Lange, Prod. Fl. Hisp. i. 108 = A. cristatum.
orientale, Jaub. & Spach, Illustr. iv. 26 = A. orientale.
squarrosum, Jaub. & Spach, l. c. 28 = A. petalum.

EREMOPYXIS, Baill. Adansonia, ii. (1861–62) 238 = **Thryptomene**, Endl. (Myrtac.).
camphorata, Baill. l. c. 239 = T. saxicola.

EREMOSPARTON, Fisch. & Mey. Enum. Pl. Schrenk, 75 (1841). *LEGUMINOSÆ*, Benth & Hook. f. i. 504.
aphyllum, *Fisch. & Mey. l. c.* 76.—Reg. Casp.
tarkestanum, *Franch. in Ann. Sc. Nat. Sér.* VI. xv. (1883) 352.—Turkestan.

EREMOSPATHA, Mann & H. Wendl. in Kerch. Palm. 244 (1878). *PALMÆ*, Benth. & Hook. f. iii. 926.
cuspidata, *Mann & H. Wendl. l. c.*—Afr. trop.
Hookeri, *Mann & H. Wendl. l. c.*—Afr. trop.
macrocarpa, *Mann & H. Wendl. l. c.*—Afr. trop.

EREMOSPORUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 355; et Hist. Vég. Phan. v. 375 (1836) = **Hypericum**, Tourn.
Olivierii, Spach, ll. oc. = H. Olivieri.

EREMOSTACHYS, Bunge, in Ledeb. Fl. Alt. ii. 414 (1830). *LABIATÆ*, Benth. & Hook f. ii. 1215.
CLUERIA, Rafin. Fl. Tellur. iii. 37 (1836).
acanthocalyx, *Boiss. Diagn.* Ser. II. iv. 49.—Beluchist.
adenantha, *Jaub. & Spach, Illustr.* iv. 18. t. 412.—Persia.
affinis, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1844) 211.—Soongaria.
azalensis, *Bunge, Lehm. Rel. Bot.* 266.—Turkestan.
Bungei, *Regel, in Act. Hort. Petrop.* vi. (1879) 380.—Persia.
discolor, *Bunge, Lab. Persic.* 79.—Persia.
diversifolia, *Regel, in Act. Hort. Petrop.* vi. (1879) 380.—Kokand.
fulgans, *Bunge, Lab. Persic.* 80.—Persia.

EREMOSTACHYS:—
glabra, *Boiss. ex Benth. in DC. Prod.* xii. 548.—Persia.
gymnocalyx, *Schrenk, Enum. Pl. Nov.* 84.—Turkestan.
hyoscyamoides, *Botss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 161.—Persia.
iberica, Vis. Sem. Kar. Patav. (1846) 64; et ex Jaub. & Spach, Illustr. v. 13 = laciniata.
lilienis, *Regel, in Act. Hort. Petrop.* vi. (1879) 375.—Turkestan.
Kaufmanniana, *Regel, l. c.* 379.—Turkestan.
Krauseana, *Regel, l. c.* 378.—Turkestan.
labiosa, *Bunge, Lab. Persic.* 79.—Persia.
laciniata, Ahch. Cat. Pl. Panjab et Scinde, 118 = acanthocalyx.
lacinata, *Bunge, in Ledeb. Fl. Alt.* ii. 416.—Aa. Min.; Syria.
laevigata, *Bunge, Lab. Persic.* 81.—Persia.
Lehmanniana, *Bunge, l. c.*—Turkestan.
leucasfolia, *Benth. in DC. Prod.* xii. 547.—Afghan.
macrochlita, Jaub. & Spach, Illustr. v. 18 = laciniata.
macrophylla, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 64.—Aa. occ.
molinioides, *Bunge, in Ledeb. Fl. Alt.* ii. 415.—Sibir.
napuligera, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 237.—Turkest.
Neumani, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 50.—Persia.
Olgæ, *Regel, Pl. Nov. Fedsch.* 70.—Turkestan.
paniculata, *Regel, in Act. Hort. Petrop.* vi. (1879) 381.—Turkestan.
phlomoides, *Bunge, in Ledeb. Fl. Alt.* ii. 414.—Reg. Cancas.; Sibir.
pulvinata, *Jaub. & Spach, Illustr.* v. 18.—Persia.
pyramidalis, Jaub. & Spach, l. c. t. 462 = macrophylla.
rotata, *Schrenk, ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* ix. *Suppl.* 3, 1.—Soongaria.
sanguinea, *Jaub. & Spach, Illustr.* v. 18.—Hab.?
Sewerzovii, *Herd. in Bull. Soc. Nat. Mosc.* xli. (1868) 1. 892.—Turkestan.
speciosa, *Rupr. Sert. Tianach.* 98.—Aa. centr.
Stochsii, Boiss. Diagn. Ser. II. iv. 48 = leucasfolia.
superba, Bunge, Lehm. Rel. Bot. 265 = Lehmanniana.
superba, *Regel, ex Benth. in Hook. Bot. Misc.* iii. (1833) 381.—Reg. Himal.
thyrsiflora, *Benth. in DC. Prod.* xii. 548.—Afghan.
Tournefortii, *Jaub. & Spach, Illust.* iv. 12. t. 412.—Aa. Min.
tranoxana, *Bunge, Lehm. Rel. Bot.* 265.—Turkestan.
tuberosa, *Bunge, in Ledeb. Fl. Alt.* ii. 416.—Reg. Cancas.; Sibir.
Vicaryi, *Benth. in Aitch. Cat. Pl. Panjab et Scinde*, 119.—Ind. bor. occ.

EREMOSYCE, Stend. Nom. ed. II. i. 566 (1840) = seq.

EREMOSYNE, Endl. in Enum. Pl. Hueg. 58 (1837). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 636.
pectinata, *Endl. l. c.*—Austral.

EREMURUS, Bieb. Cent. Pl. Ross. i. 61 (1816). *LILIACEÆ*, Benth. & Hook. f. iii. 787.
AMMOLIRION, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1842) 515.
HENNINGIA, Kar. & Kir. l. c. 516 (1842).
SELONIA, Regel, l. c. xii. (1861) 1. 457.
Aitchisoni, *Baker, in Journ. Linn. Soc.* xvii. (1880) 105.—Afghan.
Alberti, *Regel, in Act. Hort. Petrop.* viii. (1884) 868.—Aa. centr.
alboviridens, *Baker, in Journ. Bot.* xvii. (1879) 17.—Orient.
altaicus, *Stev. in Bull. Soc. Nat. Mosc.* iv. (1832) 355.—Soongar.; Sibir.
angustifolius, *Baker, in Journ. Linn. Soc.* xv. (1876) 329.—Persia; Afghanist.
anisopterus, *Regel, Gartenfl.* (1878) 260.—Oriens.
Ancheriensis, *Boiss. Diagn.* Ser. I. viii. 130.—Persia; Turkest.
aurantiacus, Baker, in Journ. Linn. Soc. xv. (1876) 368 = Bungei.
bachtiaricus, *Boiss. Fl. Orient.* v. 828.—Persia.
bucharicus, *Regel, in Act. Hort. Petrop.* viii. (1884) 870.—Bokhara.
Bungei, *Baker, in Journ. Bot.* xvii. (1879) 17.—Persia.
cappadocicus, *Y. Gay, ex Baker, in Journ. Linn. Soc.* xv. (1876) 351.—Aa. Min.

EREMURUS:—
Capusi, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 260.—Turkestan.
caucasicus, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 351 = spectabilis.
Griffithii, *Baker, in Journ. Linn. Soc. xv.* (1876) 353 = Afghan.
himalaicus, *Baker, l. c.*—Reg. Himal.
inderiensis, *Regel, Gartenfl.* (1878) 259.—Sibir.; Turkestan.
Kaufmanni, *Regel, l. c.* 260.—Turkestan.
Karolkowi, *Regel, in Act. Hort. Petrop.* iii. (1875) 116.—Turkestan.
libanoticus, Boiss. Diagn. Ser. II. iv. 97 = spectabilis.
luteus, *Baker, in Journ. Bot.* xvii. (1879) 18.—Persia.
Olgæ, *Regel, Gartenfl.* (1878) 260.—Turkestan.
pauciflorus, Baker, in Journ. Bot. xvii. (1879) 18 = persicus.
persicus, *Boiss. Diagn.* Ser. I. vii. 191.—Persia, Afghan.
robustus, *Regel, Gartenfl.* (1878) 257.—Turkestan.
sibiricus, Weinm. Hort. Paulowsk. 167; ex Schult. f. Syst. vii. 483 = spectabilis.
spectabilis, *Bieb. Pl. Ross.* 2. t. 61.—Aa. Min.; Persia.
spectabilis, Ledeb. Fl. Alt. ii. 25 = altaicus.
stenophyllus, *Baker, in Journ. Linn. Soc.* xv. (1876) 351.—Persia.
Stocksii, *Baker, l. c.* 353.—Beluchistan.
Szawtowi, *Regel, in Act. Hort. Petrop.* viii. (1884) 872.—Aa. centr.
tauricus, *Stev. in Bull. Soc. Nat. Mosc.* iv. (1832) 353 —Tauria; Persia; Soongar.
tauricus, Weinm. Hort. Paulowsk. 167; ex Schult. f. Syst. vii. 483 = spectabilis.
turkestanicus, *Regel, Gartenfl.* (1878) 260.—Turkestan.
volkhonskii, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 217 = persicus.

ERESDA, Spach, Hist. Vég. Phan. vii. 101 (1839) = **Reseda**, Linn.
alba, Spach, l. c. 102 = R. alba.

ERESIMUS, Rafin. Sylva Tellur. 60 (1838) = **Cephalanthus**, Linn. (Rubiac.).
stellatus, Rafin. l. c. 61 = C. stellatus.

ERETIA, Stokes, Bot. Mat. Med. i. (1812) 421 = **Ehretia**, Linn. (Boragin.).

ERHARTA, Jus. Gen. 93 (1789) = **Ehrharta**, Thunb (Gramin.).

ERHETIA, Hill, Hort. Kew. 440 (1769) = **Ehretia** Linn. (Borag.).

ERIA, Lindl. Bot. Reg. t. 904 (1826). *ORCHIDEÆ*, Benth. & Hook. f. iii. 509.
AGGREIANTHUS, Wight, Ic. v. 18. t. 1787 (1852).
ALVISIA, Lindl. Fol. Orch. (1859).
BRYORIUM, Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
CALLOSTYLIS, Blume, Bijdr. 340. t. 74 (1825).
CREATIUM, Blume, l. c. 341. t. 68 (1825).
CONCHIDIUM, Griff. Notul. iii. 321. t. 310 (1851).
CYLINDROLOBUS, Blume, Prael. Fl. Jav. 6 (1858).
DENDROLIRIUM, Blume, Bijdr. 343. t. 69 (1825).
ERIOCANTHA, Rafin. in Lond. Gard. Mag. viii. (1832) 247.
EXERIA, Rafin. Fl. Tellur. iv. 49 (1836).
LICHENORA, Wight, Ic. v. t. 1788 (1852).
MYCARANTHES, Blume, Bijdr. 359. t. 87 (1825).
OCTOMERIA, D. Don, Prod. Fl. Nep. 31 (1825).
PORPAX, Lindl. Bot. Reg. (1845) Misc. 68.
PINALIA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 31, in syn. (1825); Lindl. Orch. Sccl. 31 (1826).
TRICHOSIA, Blume, Bijdr. Tabellen, 11 (1835).
TRICHOTOSIA, Blume, Bijdr. 343 (1825).
TYLOSTYLIS, Blume, Prael. Fl. Jav. 6 (1858).
XIPHOSIUM, Griff. Itin. Not. 78 (1845).
abbreviata, *Lindl. Gen. et Sp. Orch.* 86.—Java.
acervata, *Lindl. in Journ. Hort. Soc.* vi. (1851) 87.—Reg. Himal.
aeriadenchyma, Benth. in Journ. Linn. Soc. xviii. (1881) 393, sphalm. = aleiadenchyma.
acuminata, *Lindl. Gen. et Sp. Orch.* 86.—Java.
acuminata, *Lindl. Bot. Reg.* (1843) Misc. 86.—Ind. or.

ERIA :—

acutissima, *Reichb. f. in Gard. Chron.* (1876) i. 567.—Hab. ?

aeridostachya, *Reichb. f. ex Lindl. in Journ. Linn. Soc.* iii. (1859) 48; *et in Seem. Fl. Vit.* 301.—Malaya; Ins. Fiji.

affinis, Griff. Notul. iii. 297 = bracteosa.

alba, *Lindl. Gen. et Sp. Orch.* 67.—Reg. Himal.

albidotomentosa, *Lindl. l. c.* 66.—Java.

Ambroxia, Hance, in Journ. Bot. xxi. (1883) 232 (Bulbophyllum sp.).—China.

amica, *Reichb. f. Xenia Orch.* ii. 103.—Reg. Himal.

angulata, *Reichb. f. in Flora*, lv. (1872) 272.—Ind. or. *Anta*, Reichb. f. in Walp. Ann. vi. 270 = Tainia latifolia.

annulata, *Blume, Mus. Bot. Lugd. Bat.* ii. 184.—Java.

annulata, Lindl. in Journ. Linn. Soc. iii. (1858) 57 = rufinula.

aporoides, *Lindl. in Journ. Linn. Soc.* iii. (1859) 60. —Ins. Philipp.

appendiculata, *Lindl. Gen. et Sp. Orch.* 69.—Java.

armeniaca, Lindl. Bot. Reg. (1841) Misc. 88. t. 42 = ornata.

articulata, *Lindl. in Journ. Linn. Soc.* iii. (1859) 47. —Zeylan.

bambusifolia, *Lindl. l. c.* 61.—Reg. Himal.

barbarosa, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 420.—Hab. ?

barbata, *Reichb. f. in Walp. Ann.* vi. 270.—Reg Himal.

Barringtoniana, *Reichb. f. in Gard. Chron.* (1872) 666. —Borneo.

bicolor, *Lindl. Gen. et Sp. Orch.* 65.—Zeylan.

bicolor, Lindl. in Journ. Linn. Soc. iii. (1859) 58 = Lindleyi.

bicornis, Reichb. f. in Walp. Ann. vi. 269 = Tainia bicornis.

bicristata, *Lindl. Gen. et Sp. Orch.* 67.—Java.

bifalcis, *Lindl. in Journ. Linn. Soc.* iii. (1859) 56.— Borneo.

biflora, *Griff. Notul.* iii. 302.—Reg. Himal.

biflora, *Lindl. in Journ. Linn. Soc.* iii. (1859) 56.— Malacca.

bigibba, *Reichb. f. in Gard. Chron.* (1884) ii. 680.— Borneo.

bipunctata, *Lindl. Bot. Reg.* (1841) *Misc.* 83.—Reg. Himal.

braccata, Dalz. & Gibs. Bomb. Fl. 269 = reticosa.

braccata, *Lindl. in Journ. Linn. Soc.* iii. (1859) 46.— Zeylan.

brachystachya, *Reichb. f. in Bonplandia*, iii. (1855) 228. —Ins. Philipp.

bracteosa, *Lindl. Bot. Reg.* (1841) *Misc.* 18.— Burma; Malaya.

canaliculata, *Blume, Mus. Bot. Lugd. Bat.* ii. 184.— Borneo.

capitellata, *Lindl. in Journ. Linn. Soc.* iii. (1858) 57. —Java.

carinata, *Gibson, in Calc. Journ. Nat. Hist.* v. (1845) 365.—Reg. Himal.

carinata, Reichb. f. in Seem. Fl. Vit. 301 = Thelasis carinata.

Carolettae, *Hance, in Journ. Bot.* xv. (1877) 337.— Cambodia!

cauligera, Reichb. f. Otia Bot. Hamb. 55 = Phreatia cauligera.

ciliata, *Mig. Fl. Ind. Bat.* iii. 659.—Java.

clavilophylax, Reichb. f. in Bonplandia, v. (1857) 54 (Phreatiae sp.).—Ins. Carol.

clavicaulis, *Wall. ex Lindl. Bot. Reg.* (1840) *Misc.* 90.—Reg. Himal.

clavicaulis, *Wall. ex Lindl. in Journ. Linn. Soc.* iii. (1858) 59 (*non Bot. Reg.*).—Reg. Himal.

cochlearis, *Lindl. Bot. Reg.* (1844) *Misc.* 14.—Ins. Philipp.

compressa, *Blume, Mus. Bot. Lugd. Bat.* ii. 183; *Reichb. f. in Bonplandia*, v. (1857) 55.—Java.

concolor, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 168.—Burma.

convallarioides, *Lindl. Gen. et Sp. Orch.* 70.—Reg. Himal.

coriacea, *Reichb. f. in Bonplandia*, v. (1857) 54.—Java.

Cornuti, *Reichb. f. in Gard. Chron.* (1878) ii. 106. —Ins. Formosa.

coronaria, Reichb. f. in Walp. Ann. vi. 271 = Trichosma suavis.

Cumingi, *Reichb. f. in Gard. Chron.* (1880) ii. 685.— Borneo.

ERIA :—

cylindripoda, Griff. Notul. iii. 299 = Trichosma suavis.

Dalzelli, *Lindl. in Journ. Linn. Soc.* iii. (1859) 47.— Ind. or.

dasyphylla, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 147.—Reg. Himal.; Burma.

dasypus, *Reichb. f. in Bot. Zeit.* xxii. (1864) 415.— Burma.

Dayana, *Reichb. f. in Gard. Chron.* (1877) ii. 102.— Ind.

dendrophylax, Reichb. f. in Bonplandia, v. (1857) 54 (Phreatiae sp.).—Ins. Carol.

dentiflora, [*Lodd. Cat. Orch.* 6]; *Lindl. ex Loud. Hort. Brit. Suppl.* ii. 633.—Ind. or.

Dillwynia, Hook. Bot. Mag. t. 4163 = bracteosca.

discolor, *Lindl. in Journ. Linn. Soc.* iii. (1859) 51.— Reg. Himal.

ebulbis, *Lindl. Gen. et Sp Orch.* 69.—Java.

eburnea, *Reichb. f. in Hamb. Gartenz.* xv. (1859) 56. —Ind. or.

elegans, Reichb. f. in Seem. Fl. Vit. 301 = Phreatia elegans.

elongata, *Blume, Mus. Bot. Lugd. Bat.* ii. 183.— Java.

elongata, *Lindl. in Journ. Linn. Soc.* iii. (1859) 49.— Burma.

Elwesii, Reichb. f. in Gard. Chron. (1888) i. 402 = E. *Meirax*.

Ephemera, Reichb. f. in Walp. Ann vi. 272 = Lindleyi.

crecta, *Lindl. Gen. et Sp. Orch.* 68.—Java.

eriopetala, *Par & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 148.—Burma.

excavata, *Lindl in Wall. Cat.* n. 1974.—Reg Himal.

extinctoria, *Oliver, in Hook. Bot Mag* t. 6910.— Burma.

ferox, *Blume, Mus. Bot. Lugd. Bat.* ii. 184.—Malaya.

ferruginea, *Lindl. Bot Reg.* (1839) t. 35.—Reg. Himal.

ferruginea, *Tejsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 312.—Java.

filiformis, Reichb. f. in Walp. Ann. vi. 268 = Dalzelli.

Fitzalani, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 252.—Austral

flava, Griff. Notul. iii. 301 = elongata.

flava, *Lindl. in Wall. Cat.* n. 1973.—Reg. Himal.

flavescens, *Lindl. Gen. et Sp. Orch.* 66.—Java.

floribunda, *Lindl. in Wall. Cat.* n. 7408.—Malaya.

fragrans, *Reichb. f. in Bot. Zeit.* xxii. (1864) 415.— Burma.

fusca, *Blume, Mus. Bot. Lugd. Bat.* ii. 183.—Java.

gladiata, *Reichb. f. in Seem. Fl. Vit.* 300.—N. Zel.

graminifolia, *Lindl. in Journ. Linn. Soc.* iii. (1859) 54.—Reg. Himal.

Griffithii, *Reichb. f. Xenia Orch.* ii. 102, 168.— Burma.

hemilaea, *Reichb. f. in Hamb. Gartenz.* xix. (1863) 11.—Java.

hirta, *Blume, Mus. Bot. Lugd. Bat.* ii. 184.—Ins. Celebes.

hologlossa, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 148 (= Tainiae sp.).—Burma.

hyacinthoides, *Lindl. Gen. et Sp. Orch.* 66.—Java.

ignea, *Reichb. f. in Gard. Chron.* (1881) i. 782.— Borneo.

javensis, *Zoll. & Mor. Syst. Verz. Zoll.* 86.—Java.

Jerdoniana, Reichb. f. in Walp. Ann. vi. 267 = E. Lichenora.

khasiana, Lindl. in Journ. Linn. Soc. iii. (1859) 59 = Thelasis.

Kingii, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 71.—Ins. Salomon.

lamorganensis, *Reichb. f. in Bonplandia*, v. (1857) 55. —Java.

lanata, Griff. Notul. iii. 301 = flava.

lanicaps, Reichb. f. in Hamb. Gartenz. xix. (1863) 10 = flava.

latifolia, Blume, Mus. Bot. Lugd. Bat. ii. 183 = sumatrana.

latifolia, *Reichb. f. in Bonplandia*, v. (1857) 55.— Java.

laxiflora, Miq. Fl. Ind. Bat. iii. 665 = Phreatia laxiflora.

lelophylla, *Lindl. in Journ. Linn. Soc.* iii. (1859) 57. —Borneo.

leucostachya, Lindl. in Journ. Hort. Soc. iii. (1848) p. xv = floribunda.

leucovarekha, Hort. ex Reichb. f. in Gard. Chron. (1880) ii. 388 = Tainia latifolia.

ERIA :—

Lichenora, *Lindl. in Journ. Linn. Soc.* iii. (1859) 46. —Ind. or.

limenophylax, *Reichb. f. in Bonplandia*, v. (1857) 54. —Java.

limenophylax, Reichb. f. Xenia Orch. ii. 97. t. 130 partim = Phreatia limenophylax.

Lindleyana, Griff. Notul. iii. 300 = obesa.

Lindleyi, *Thw. Enum. Pl. Zeyl.* 299.—Zeylan.

linearis, *Lindl. in Journ. Linn. Soc.* iii. (1859) 53.— Java.

lineoligera, *Reichb. f. in Gard. Chron.* (1885) ii. 262. —Siam.

littoralis, *Tejsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 312.—Malaya.

lobata, *Reichb f. in Bonplandia*, v. (1857) 55.—Java.

longicaulis, *Tejsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 313.—Malaya.

longilabris, *Lindl. Bot. Reg.* (1841) *Misc.* 38.—Ins. Philipp.

Matthewsi, *Reichb f. Otia Bot. Hamb* 55.—Ins. Tahiti.

Meirax, *N. E. Br. in Gard. Chron.* (1880) ii. 608.— Burma.

mergnensis, *Lindl. in Journ. Linn. Soc.* iii. (1859) 52.—Burma.

micrantha, *Lindl. in Journ. Linn. Soc.* iii. (1859) 47 = Dalzelli.

microchilos, Lindl. in Journ. Linn. Soc. iii. (1859) 47 = Dalzelli.

Microdiris, Reichb. f. in Seem. Fl Vit. 301 = Phreatia Microdiris.

microphylla, *Blume, Mus. Bot. Lugd. Bat.* ii. 184.— Java.

minutiflora, Reichb f. in Seem. Fl. Vit. 301 = Phreatia minutiflora.

monostachya, *Lindl. in Journ. Linn. Soc.* iii. (1859) 55.—Java.

muconata, *Lindl. Bot. Reg.* (1842) *Misc.* 25.—Ins. Philipp.

multiflora, *Lindl. Gen. et Sp. Orch.* 68.—Java.

muscicola, *Lindl. in Journ. Linn. Soc.* iii. (1859) 47. —Reg. Himal.; Zeylan.

Myosurus, Reichb. f. in Bonplandia, v. (1857) 54 = Phreatia Myosurus.

Myosurus, Reichb. f. in Seem. Fl. Vit. 300 = Oberonia Myosurus.

myristicaeformis, *Hook. Bot. Mag.* t. 5415.—Burma.

myosorensis, *Lindl. in Journ. Linn. Soc.* iii. (1859) 54. —Ind. or

nana, *A. Rich. in Ann. Sc. Nat.* Sér. II. xi. (1842) 19. —Ind. or.

nutans, *Lindl. Bot. Reg.* (1840) *Misc.* 88.—Malaya.

obesa, *Lindl. in Wall. Cat.* n. 1976.—Burma.

oblique, *Lindl. in Journ. Linn. Soc.* iii. (1859) 55; *Reichb. f. in Walp. Ann.* vi. 271.—Singapore.

obliterata, *Reichb. f. in Bonplandia*, v. (1857) 55.— Java.

odoratissima, *Tejsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 17.—Sumatra.

oreophylax, Reichb. f. Otia Bot. Hamb. 55 = Phreatia oreophylax.

ornata, *Lindl. Gen. et Sp. Orch.* 66.—Java.

ornata, *Lindl. in Journ. Linn. Soc.* iii. (1859) 48 (*non Gen. et Sp. Orch.*).—Reg. Himal.

ovata, *Lindl. Bot Reg.* (1844) sub t. 29.—Ins. Philipp.

pachysachya, *Lindl. in Journ. Linn. Soc.* iii. (1859) 60.—Java.

palcata, Reichb. f. in Linnaea, xli. (1877) 87 = Phreatia palcata.

paniculata, *Lindl. in Wall. Pl. As. Rar.* i. 32. t. 36.— Reg. Himal.

paniculata, Reichb. f. ex Lindl. in Journ. Linn. Soc. iii. (1859) 58 = monostachya.

pannea, *Lindl. Bot. Reg.* (1842) *Misc.* 64.—Reg. Himal.; Malaya.

Parishii, *Lindl. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 147.—Burma.

pauciflora, *Blume, Mus. Bot. Lugd. Bat.* ii. 183.— Java.

pauciflora, *Wight, Ic.* v. t. 4. t. 1636.—Ind. or.

pauciflola, *Reichb. f. in Walp. Ann.* vi. 270.—Java.

perpusilla, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 148 (Phreatiae sp.).—Burma.

planicaulis, Wall. ex Lindl. Bot. Reg. (1840) Misc. 8 = Agrostophyllum khasianum.

Pleurothallis, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 147.—Burma.

plenaroides, Reichb. f. in Bonplandia, v. (1857) 54 (Phreatiae sp.).—Ins. Samoa.

ERIA:—
polystachya, *A. Rich. in Ann. Sc. Nat.* Sér. II. xv. (1841) 20. t. 9.—Ind. or.
polystachya, Wight, Ic. v. t. 4. t. 1684 (sphalm. nomen textu) = pubescens.
polyura, *Lindl. Bot. Reg.* (1841) *Misc.* 55.—Ins. Philipp.
profusa, *Lindl. l. c.* (1842) *Misc.* 2.—Zeylan ?
prorepens, Reichb. f. Otia Bot. Hamb. 54 = Phreatia prorepens.
pubescens, Lindl. in Bot. Reg. sub t. 904 = flava.
pubescens, *Wight, Ic.* v. t. t. 1634 (polystachya).—Ind. or.
pulchella, Griff. Notul. iii. 297 = Griffithii.
pulchella, *Lindl. in Wall. Cat* n. 7407.—Malaya.
pulvinata, *Lindl. in Journ. Linn. Soc.* iii. (1859) 56. —Burma.
pumila, *Lindl. in Wall. Cat.* n. 1972.—Reg. Himal.; Burma.
pusilla, *Lindl. in Journ. Linn. Soc.* iii. (1859) 48.—Reg. Himal.
pusilla, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 312.—Java.
Reinwardtii, *Lindl. in Journ. Linn. Soc.* iii. (1859) 53.—Java.
reticosa, *Wight, Ic.* v. t. 4. t. 1687.—Ind. or.
reticulata, *Benth. & Hook. f. Gen.* iii. 509.—Ind. or.
retrofiexa. *Lindl. in Journ. Linn. Soc.* iii. (1859) 50. —Ins. Philipp.
retusa, *Reichb. f. in Bonplandia*, v. (1857) 54.—Java.
rhodopteres, *Reichb. f. in Gard. Chron.* (1882) ii. 586. —Hab.?
Richardiana, *Reichb. f. in Seem. Fl. Vit.* 300.—N. Zel.
rigida, *Blume, Mus. Bot. Lugd. Bat.* ii. 183.—Borneo.
rigida, Reichb. f. in Bonplandia, v. (1857) 55 = pulchella.
Rimanni, *Reichb. f. in Gard. Chron.* (1885) ii. 712.—Burma.
ringens, *Reichb. f. in Bonplandia*, ii. (1855) 222.—Burma.
robusta, *Lindl. Gen. et Sp. Orch.* 89.—Java.
rosea, *Lindl. in Bot. Reg.* t. 978.—China.
rosea, Wall. Cat. n. 7408—carinata.
rostriflora, *Reichb. f. in Seem. Fl. Vit.* 301.—Ins. Fiji.
rubiginosa, *Blume, Mus. Bot. Lugd. Bat.* ii. 184.—Borneo.
rufinula, *Reichb. f. in Hamb. Gartenz.* xix. (1868) 13. —Reg. Himal.
rugosa? Lindl. Gen. et Sp. Orch. 66 = stellata.
scabrilinguis, *Lindl. in Journ. Linn. Soc.* iii. (1859) 51.—Reg. Himal.
sclerophylla, *Lindl. l. c.* 56.—Java.
secunda, *Reichb. f. in Bonplandia*, v. (1857) 54.—Java.
secundiflora, Griff. Notul. iii. 302 = stricta.
sicaria, *Lindl. in Journ. Linn. Soc.* iii. (1859) 50.—Burma.
sinica, *Lindl. l. c.* 48.—China.
sonkaria, *Reichb. f. in Bonplandia*, v. (1857) 55.—Java.
spaciosa, Reichb. f. in Walp. Ann. vi. 269 = Tainia speciosa.
sphaerocarpa, *Reichb. f. in Seem. Fl. Vit.* 301.—Ins. Fiji.
sphaerochila, Lindl. in Journ. Linn. Soc. iii. (1859) 54 = excavata.
stachyurus, Reichb. f. in Seem. Fl. Vit. 301 = Phreatia Mycosurus.
stellata, *Lindl. in Bot. Reg.* t. 904.—Ins. Philip. stenostachya, *Reichb. f. in Seem. Fl. Vit.* 301.—Ins. Fiji.
stricta, *Lindl. Coll. Bot.* t. 41 B.—Reg. Himal.
suavis, Lindl. in Journ. Linn. Soc. iii. (1859) 52 = Trichosma suavis.
suluata, *Lindl. Gen. et Sp. Orch.* 89.—Java.
sumatrana, *Miq. Fl. Ind. Bat.* iii. 558.—Sumatra.
sumbawensis, *Reichb. f. in Bonplandia*, v. (1857) 55. —Java.
tahitensis, Reichb. f. in Linnaea, xli. (1877) 87; et in Seem. Fl. Vit. 301 (1868) = Phreatia tahitensis.
tentaculata, Griff. Notul. iii. 298 = pannea.
Thwaitesii, *Trimen, Cat. Pl. Cyl.* 88.—Zeylan.
tomentella, Reichb. f. in Walp. Ann. vi. 271 = javensis.
tricolor, *Thw. Enum. Pl. Zeyl.* 429.—Zeylan.
trilophota, *Lindl. in Journ. Linn. Soc.* iii. (1859) 54. —Borneo.
truncata, *Lindl. l. c.* 58.—Burma.

ERIA :—
uniflora, Dalz. in Hook. Kew Journ. iv. (1852) 111 = bracteata.
ustulata, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 147.—Borneo.
vaginata, *Benth. in Journ. Linn. Soc.* xviii. (1881) 503.—Java.
valida, *Lindl. l. c.* iii. (1859) 60.—Java.
velifera, Wight, ex Lindl. l. c. 46 = bracteata ?
velutina, *Lodd. ex Lindl. Bot. Reg.* (1840) *Misc.* 86. —Singapore.
velutina, Thw. Enum. Pl. Zeyl. 299 = Thwaitesii.
vestita, *Lindl. Bot. Reg.* (1844) 79.—Malaya.
Vieillardi, *Reichb. f. in Linnaea*, xli. (1877) 86.—N. Caled.
vittata, *Lindl. in Journ. Linn. Soc.* iii. (1859) 51.—Reg. Himal.
Vrieseana, *Reichb. f. in C. Koch, Allg. Gartenz.* (1858) 379.—Malaya.
vulpina, *Reichb. f. in Bonplandia*, iii. (1855) 222.—Java.
Wightii, Reichb. f. in Walp. Ann. vi. 270 = Tainia bicornis.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 55.—Java.

ERIACHAENIUM, Sch. Bip. in Flora, xxxviii. (1855) 190. *COMPOSITAE*, Benth. & Hook. f. ii. 456. magellanicum, *Sch. Bip. l. c.* 191.—Reg. Magell.

ERIACHNE, R. Br. Prod. 168 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1185.
ACINERIA, Beauv. Agrost. 72 (1812).
MEGALACHNE, Thw. Enum. Pl. Zeyl. 378 (1864).
agrostoides, *F. Muell. Fragm.* vii. 33.—Austral.
ampla, *Nees, Fl. Afr. Austr.* 377.—Afr. austr.
aristidea, *F. Muell. Fragm.* v. 205.—Austral.
Armitti, *F. Muell. ex Benth. Fl. Austral.* vii. 627 (Armittii).—Austral.
assimilis, *Steud. Syn. Pl. Gram.* 286.—Afr. austr.
aures, *Nees, Fl. Afr. Austr.* 376.—Afr. austr.
avenacea, *R. Br. Prod.* 184.—Austral.
brevifolia, R. Br. l. c. = mucronata.
capensis, *Steud. in Flora*, xii. (1829) 470.—Afr. austr.
capillaris, *R. Br. Prod.* 184.—Austral.
chinensis, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 228 = pallescens.
ciliata, *R. Br. Prod.* 184.—Austral.
Eckloni, *Nees, Fl. Afr. Austr.* 373.—Afr. austr.
festucacea, *F. Muell. Fragm.* v. 205.—Austral.
glauca, *R. Br. Prod.* 184.—Austral.
gracilis, *Brongn. in Duperr. Voy. Bot.* 25.—Ins. Molucc.
Hookeri, Munro, in Journ. Linn. Soc. vi. (1862) 42 = pallescens.
maloumensis, *Steud. Syn. Pl. Gram.* 286.—Reg. Magell.
melicacea, *F. Muell. Fragm.* v. 205.—Austral.
microphylla, *Nees, Fl. Afr. Austr.* 377.—Afr. austr.
montana, Steud. Nom. ed. II. i. 567, sphalm. = Panicum subigenum.
mucronata, *R. Br. Prod.* 184.—Austral.
obtusa, *R. Br. l. c.* 184.—Austral.
ovata, *Nees, in Hook. Lond. Journ. Bot.* ii. (1843) 416.—Austral.
pallescens, *R. Br. Prod.* 184.—Austral.
pallida, *F. Muell. ex Benth. Fl. Austral.* vii. 631.—Austral.
pallida, *Nees, Fl. Afr. Austr.* 375.—Afr. austr.
Preissiana, Nees, in Lehm. Pl. Preiss. ii. 102 = ovata.
rara, R. Br. Prod. 183.—Austral.
sclerachnoides, *F. Muell. Fragm.* viii. 233.—Austral.
Schultziana, F. Muell. l. c. 187 = stipacea.
setacea, *F. Muell. ex Benth. Fl. Austral.* vii. 629.—Austral.
squarrosa, *R. Br. Prod.* 183.—Austral.
Stendelii, *Nees, Fl. Afr. Austr.* 378.—Afr. austr.
stipacea, *F. Muell. Fragm.* v. 206.—Austral.
stricta, *Steud. Syn. Pl. Gram.* 287.—Hab. ?
tristis, *Nees, ex Steud. l. c.*—Afr. austr.
tuberculata, *Nees, Fl. Afr. Austr.* 374.—Afr. austr.

ERIACHNE, Phil. Sert. Mend. Alt. 49 (1871) = Pentoum. Lin. (Graminis.).
rigida, Phil. l. c. 50 = P. Phaeotoma.

ERIADENIA, Miers, Apocyn. S. Am. 117 (1878). *APOCYNEAE*.
obovata, *Miers, l. c.*—Am. austr.

ERIANDROSTACHYS, Baill. Adansonia, xi. (1874) 289. *SAPINDACEAE*.
Chapelieri, *Baill. l. c.*—Madag.

ERIANTHERA, Benth. in Hook. Bot. Misc. iii. (1833) 360 = Laxmium, Tourn. (Labiat.).
rhomboidea, Benth. l. c. = L. rhomboideum.

ERIATHERA, Nees, in Wall. Pl. As. Rar. iii. 77 (1832) = Andrographis, Wall. (Acanthac.).
lobelioides, Nees, l. c. 115 = A. echioides.
Neesiana, Wight & Arn. ex Nees, in DC. Prod. xi. 514 = A. serpyllifolia.
serpyllifolia, Nees, in Wall. Pl. As. Rar. iii. 115 = A. serpyllifolia.

ERIANTHUS, Michx. Fl. Bor. Am. i. 54 (1803). *GRAMINEAE*, Benth. & Hook. f. iii. 1126.
RIPIDIUM, Trin. Fund. Agrost. 169 (1820).
alopecuroides, Ell. Sketch, 88 = saccharoides.
angustifolius, Nees, Agrost. Bras. ii. 316 = saccharoides.
articulatus, F. Muell. Fragm. viii. 118 = Pollinia articulata.
asper, *Nees, Agrost. Bras.* ii. 315.—Bras.; Paraguay.
aureus, Beauv. ex Roem. & Schult. Syst. ii. 824 = Pollinia aurea
aureus, Nees, ex Steud. Syn. Pl. Gram. 412 = Pollinia Cumingii.
brasilianus, Anderss. in Ofvers. Vet. Akad. Stockh. (1855) 162 = asper.
brevibarbis, Michx. Fl. Bor. Am. i. 55 = saccharoides.
capensis, *Nees, Fl. Afr. Austr.* 93.—Afr. austr.
contortus, Baldw. ex Ell. Sketch, i. 140 = saccharoides.
Cumingii, F. Muell. Fragm. viii. 118 = Pollinia Cumingii.
cupidatus, Anderss. in Ofvers. Vet. Akad. Stockh. (1855) 162 = asper.
Ecklonii, Nees, Fl. Afr. Austr. 93 = capensis.
fastigiatus, *Nees, ex Steud. Syn. Pl. Gram.* 409.—Ind. or.
filiculus, *Nees, ex Steud. l. c.*—Ind. or.
Fischerianus, Rupr. ex Doell, in Mart. Fl. Bras. ii. iii. 258 = saccharoides.
floridulus, Anderss. in Ofvers. Vet. Akad. Stockh. (1855) 162 = maximus.
floridus, Schult. Mant. iii. 568 = (*Miscanthus japonicus*).—Chira; Japon.; Malaya; Polynes.
fulvus, Kunth, Rév. Gram. i. 160 = Pollinia Cumingii.
fulvus, *Nees, ex Steud. Syn. Pl. Gram.* 409.—Reg. Himal.
giganteus, Muhl. Desc. Gram. Uber. 92 = saccharoides.
glaucus, Hassk. in Flora, xxv. (1842) ii. Beibl. 8 (Pollinise sp.).—Java.
hamatachyus, Hochst. ex Miq. in Verh. Nederl. Inst. III. iv. (1851) (Anal. Bot. Ind. ii.) 85 = Pollinia argentea.
Hostii, *Griseb. Spicil. Fl. Rumel.* ii. 548.—Europ. austr. ; Orient.
irritans, Kunth, Rév. Gram. i. 160 = Pollinia irritans.
Jacquemontii, *Munro, ex Duthie*, [*in Athins. Gas. N. W. Ind.*] 686.—Ind. or.
jamaicensis, Anderss. in Ofvers. Vet. Akad. Stockh. (1855) 163 = Ravennae.
japonicus, Beauv. Agrost. 14 (= *Miscanthus sinensis*). —China.
longestatus, *Anderss. in Ofvers. Vet. Akad. Stockh.* (1855) 163.—Ind. or.
longistatus, Anderss. ex Benth. in Journ. Linn. Soc. xix. (1881) 67 = pteus.
Mackinlayi, F. Muell. Census, 131 = Pollinia Mackinlayi.
maximus, *Brongn. in Duperr. Voy. Coq. Bot.* 97.—Ins. Polynes.
mollis, *Griseb. in Goett. Nachr.* (1866) 93.—Reg. Himal.
Monnieri, *Carr. in Rev. Hortic.* (1872) 25.—Olympus.
nepalensis, *Steud. Nom.* ed. II. i. 567.—Reg. Himal.
nudipes, Griseb. in Goett. Nachr. (1866) 92 (Enlalise sp.).—Ind. or.
orientalis, C. Koch, in Linnaea, xix. (1847) 5 = Hostii.
pungens, Hochst. ex Krauss, in Flora, xxiv. (1846) 117 = capensis.
purpurascens, Anderss. in Ofvers. Vet. Akad. Stockh. (1855) 161 = Ravennae.

ERIANTHUS:—
Ravenne, *Beauv. Agrost.* 14.—Reg. Mediterr. ; Orient ; Ind. or.
repens, Beauv. l. c. (Tricholaenae sp.).—Afr. trop.
Roxburghii, F. Muell. Fragm. viii. 117 = Pollinia argentea.
rufiflus, Griseb. in Goett. Nachr. (1868) 98 = filvus.
rufus, Nees, ex Steud. Syn. Pl. Gram. 400 = Pollinia argentea.
saccharoides, *Michx. Fl. Bor. Am.* i. 55.—Am. bor. et austr.
saccharoides, Willd. Enum. Hort. Berol. 46 = Andropogon saccharoides.
Sorghum, *Nees, Fl. Afr. Austr.* 92.—Afr. austr.
speciosus, Debeaux, in Act. Linn. Soc. Bord. xxxii. (1878) 88 (Polliniae sp.).—China.
strictus, *Baldw. in Ell. Sketch.* i. 89.—Am. bor.
strictus, Bluff. & Fingerh. Consp. Fl. Germ. ed. II. i. 105 = Hostii.
tunctorius, Siebold, in Verh. Batav Gen. xii. (1830) 384 (Mischanthi sp.).—Japon.
tristachyus, Hook. & Arn. Bot. Beech. Voy. 241 (Polliniae sp.).—China.
velutinus, *Munro, ex Duthie, in [Atkins. Gaz. N. W. Ind.]* 686.—Ind. or.
vernicolor, *Nees, ex Steud. Syn. Pl. Gram.* 409.—Reg. Himal.
villosus, F. Muell. Fragm viii. 118 = Pollinia Mackinlayi.
villosus, Steud. Nom. ed. II. i. 567 = Pollinia villosa.

ERIAXIS, Reichb. f. in Linnaea, xli. (1877) 63 = **Galeola**, Lour. (Orchid.).
rigida, Reichb. f. l. c. = G. rigida.

ERICA, [Tourn.] Linn. Syst. ed. I (1735). *ERICACEAE*, Benth. & Hook. f. ii 590.
ARSACE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 119.
CALLISTA, CERAMIA, CHONA, DASVANTHES, DRSMIA, ECTASIS, ERIODESMIA, EURYLEPIS, EURYLOMA, EURYSTEGIA, LAMPROTIS, LOPHANDRA, OCTOPERA, PACHYSA, SYRINGODRA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 153–156.
EREMOCALLIS, GYPOCALLIS, Salisb. ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 308 (1821).
LOPHARINA, Neck, Elem. i. 215 (1790).
MICROTREMA, Klotzsch, in Linnaea, xii. (1838) 499.
abietina, Andr. ex G. Don, Gen. Syst. iii. 823 = concinna.
abietina, Berg. Desc. Pl. Cap. 105 = mammosa.
abietina, *Linn. Sp. Pl.* 355.—Afr. austr.
abrotanoides, *Burm. f. Fl. Cap. Prod.* 11.—Afr. austr.
absinthioides, Spreng Syst. ii. 196 = scoparia.
absinthoides, Linn. Mant. i. 66 = hispidula.
abietinoides, E. Mey. ex Klotzsch, in Linnaea, xii. (1838) 218 = Philippia Leeana.
accommodata, *Klotzsch, ex Benth. in DC. Prod.* vii. 630.—Afr. austr.
acrophya, Fenzl, in Flora, xxi. (1838) 604 = arborea.
Actaea, Link, Enum. Hort. Berol. i. 371 = peltata.
Actea, [Forbes] Hort. Eric. Woburn. 1 = imbricata.
acuminata, Andr. Heathery, iii. t. 101 = Massoni.
acuta, *Andr. l. c.* i. t. 1.—Afr. austr.
acutangula, Lodd. Bot. Cab. t. 1868 = nigrita.
aculiflora, *Tausch, in Flora*, xxii. (1839) ii. 636.—Hab. ?
acutifolia, Bartl. in Linnaea, vii. (1832) 651 = purpurea.
adsequata, *Tausch, in Flora*, xxii. (1839) ii. 634.—Hab. ?
adenophora, *Spreng. Syst.* iii. 188.—Afr. austr.
adenostoma, *Kunze, in Linnaea, xx.* (1847) 38.—Afr. austr.
adpressa, Klotzsch, l. c. xii. (1838) 509 = Massoni.
adunca, *Benth. in DC. Prod.* vii. 618.—Afr. austr.
aequalis, *Benth. l. c.* 615.—Afr. austr.
affinis, *Benth. l. c.* 656.—Afr. austr.
aggregata, Roxb. ex Salisb. in Trans. Linn. Soc. vi. (1802) 266 = cumuliflora.
aggregata, *Wendl. Eric.* ii. fasc. xiii. 11.—Afr. austr.
Aitoni, Willd. Sp. Pl. ii. 898 = E. Aitonia.
Aitonia, *Masson, in Bot. Mag.* t. 429.—Afr. austr.
Aitoniana, Lodd. Bot. Cab. t. 144 = E. Aitonia.
albens, *Linn. Mant.* ii. 231.—Afr. austr.
albens, Thunb. Prod. Pl. Cap. 70 = Scyphogyne inconspicua ?
albescens, *Klotzsch, ex Benth. in DC. Prod.* vii. 679.—Afr. austr.
albida, Thunb. Fl. Cap. ed. Schult. 347 = albens.

ERICA:—
albiflora, Klotzsch, ex Benth. in DC. Prod. vii. 674 = mollis.
Alopecias, Tausch, in Flora, xx. (1837) 498 = corifolia.
alopecuroides, Lodd. Bot. Cab. t. 874 = nudiflora.
alopecuroides, Wendl. Eric. fasc. xx. ii. 119. t. 45 = tintervallaria.
Alopecurus, *Harv. Thes. Cap.* i. 31.—Afr. austr.
alsinifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 266 = gilva.
amabilis, Salisb. l. c. 385 = inflata.
ambigua, Wendl. Eric. ii. fasc. xvi. 61. t. 24 = Massoni.
amoena, Salisb. in Trans. Linn. Soc. vi. (1802) 329 = incarnata.
amoena, Wendl. Bot. Beobacht. 48 = plumosa.
ampullacea, *Curt. Bot. Mag.* t. 908.—Afr. austr.
ampullacea, Tratt. ex Tausch, in Flora, xvii. (1834) ii. 594 = Andrew-iana.
ampulliformis, Sallab. in Trans. Linn. Soc. vi. (1802) 331 = ampullacea.
Andrewsiana, *Tausch, in Flora*, xvii. (1834) ii. 594.—Hab. ?
Andrewsii, Klotzsch, in Linnaea, x. (1836) 314 = Massoni.
andromedaeflora, *Andr. Col. Heaths*, iii. t. 146.—Afr. austr.
angustata, *Bartl. in Linnaea*, vii. (1832) 651.—Afr. austr.
angustiscilla, Bartl. l. c. 638 = Zeyheri.
anthina, Spreng. Syst. ii. 196 = mauchata.
anthura, Link, in Flora, xxviii. (1845) 190 = vagans.
aperta, Spreng. Syst. ii. 200 = borbonisaefolia.
Aphanes, *Spreng. l. c.* 196.—Afr. austr.
appressa, *Spreng. l. c.* iv. *Cur. Post.* 146.—Afr. austr.
approximata, Schlecht. ex Spreng l. c. ii. 196 = hispidula.
aprica, Klotzsch, ex Benth. in DC. Prod. vii. 660 = cardiuifolia.
arachnoides, Klotzsch, ex Benth. l. c. 691 = hispidula.
aragonensis, Willk. in Linnaea, xxv. (1852) 46 = australis.
arborea, Brot. [Fl. Lusit. ii. 221] = lusitanica.
arborea, *Linn. Sp. Pl.* 353.—Reg. Mediterr. et Caucas.
arborea, Thunb. Diss. Eric. 40 = strigosa.
Arbuscula, Lodd. Bot. Cab. t. 848 = lateralis.
arbutifiora, Wendl. Eric. iii. fasc. xxvi. 10 = trifiora.
Archeria, *Andr. Heathery*, i. t. 8.—Afr. austr.
Archeriana, Lodd. Bot. Cab. t. 1465 = praec.
arctica, Waitz, Beschr. Gatt. Art. Heid. 189 = Phyllodoce taxifolia.
ardens, *Andr. Heathery*, ii. t. 51.—Afr. austr.
argentea, *Klotzsch, ex Benth. in DC. Prod.* vii. 686.—Afr. austr.
argyrea, Andr. Col. Heaths, iv. t. 218 = purpurea.
argutifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 364 = E. Lee.
aristata, *Andr. Col. Heaths*, iii. 147.—Afr. austr.
aristata, *[Forbes] Hort. Eric. Woburn.* 2.—Afr. austr.
aristifolia, *Niven, ex Benth. in DC. Prod.* vii. 676.—Afr. austr.
armata, *Klotzsch, ex Benth. l. c.* 672.—Afr. austr.
armata, Spreng. Syst. ii. 184 = Sparmanni.
artemisioides, E. Mey. ex Klotzsch, in Linnaea, xii. (1838) 218 = Salsola artemisioides.
articularis, Curt. Bot. Mag. t. 428 = corifolia.
articularis, *Linn. Mant.* i. 85.—Afr. austr.
articularis, Thunb. Diss. Eric. 37 = pulchella.
articularis, Thunb. Prod. Pl. Cap. 71 (= *Syndesmanthus articularis*).—Afr. austr.
asiatica, Waitz, Beschr. Gatt. Art. Heid. 187 = Casiope ericoides.
aspera, Andr. Heathery, iii. t. 104 = Sparmanni.
asperifolia, Sallab. in Trans. Linn. Soc. vi. (1802) 324 = setacea.
assurgens, Link, Enum. Hort. Berol. i. 372 = persoluta
atrovirida, Klotzsch, ex Benth. in DC. Prod. vii. 666 = decipiens.
aurea, *Andr. Col. Heaths*, ii. 76.—Afr. austr.
auricularis, Sallab. in Trans. Linn. Soc. vi. (1802) 327 = tronolaris.
australis, *Linn. Ment.* ii. 231.—Reg. Mediterr. occ.
axillaris, Sallab. in Trans. Linn. Soc. vi. (1802) 325 = strigosa.
axillaris, Soland. ex Benth. in DC. Prod. vii. 692 = coarctata.
axillaris, Thunb. Diss. Eric. 16 = Salaxis axillaris.
axilliflora, *Bartl. in Linnaea*, vii. (1832) 640.—Afr. austr.
azalaeaflora, Steud. Nom. ed. II. i. 618. in syn. = seq.

ERICA:—
azalaeifolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 334.—Afr. austr.
azorica, *Hochst. ex Seub. Fl. Azor.* 40.—Ins. Azor.
baccaeflorumis, Salisb. in Trans. Linn. Soc. vi. (1802) 352 = baccans.
baccans, *Linn. Mant.* ii. 232.—Afr. austr.
baculiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 346 = Sebana.
Baudinnia, *Andr. Col. Heaths*, iv. 220 = lyrbana.
Banksia, *Andr. Col. Heaths*, i. 5.—Afr. austr.
Banksiana, Pax. Mag. Bot. vii. (1840) 248 = praec.
Banksii, Willd. Sp. Pl. ii. 895 = praec.
barbata, *Andr. Heathery*, i. t. 4.—Afr. austr.
barbata, Dredge, ex Benth. in DC. Prod. vii. 671 = perlata.
barbata, Hort. Prag ex Tausch, in Flora, xvii. (1834) ii. 616 = leucopelta.
barbigera, *Klotzsch, ex Benth. in DC. Prod.* vii. 616.—Afr. austr.
barbigera, Sallab. in Trans. Linn. Soc. vi. (1802) 341 = Blaeria barbigera.
Bartlingiana, Klotzsch, ex Benth. in DC. Prod. vii. 656.—Afr. austr.
Bartlingiana, *Klotzsch, in Linnaea*, xii. (1838) 521.—Afr. austr.
Bauera, Andr. Col. Heaths, iv. 221 = Bowieana.
Beaumontia, *Andr. l. c.* 222.—Afr. austr.
Beaumontiana, Kollisson, ex Lodd. Bot. Cab. t. 1686 = praec.
Beken, E. Mey. ex Klotzsch, in Linnaea, xii. (1838) 519 = Savilea praecox.
bella, Spreng. Syst. ii. 197 = scriphifolia.
Bergiana, *Linn. Mant.* ii. 235.—Afr. austr.
Bergiana, Wendl. Eric. i. fasc. ii. 29 = florida.
bibax, Salisb. in Trans. Linn. Soc. vi. (1802) 358 = flammea.
bibracteata, Klotzsch, in Linnaea, xii. (1838) 516 = Massoni.
bisalycolata, *Moench, Meth. Suppl.* 18.—Afr. austr. ?
bicolor, Andr. Heathery, ii. t. 54 = dichrus.
bicolor, *Thunb. Diss. Eric.* 36.—Afr. austr.
biflora, Link, Enum. Hort. Berol. i. 367 = tianthifolia.
Blaeria, Thunb. Prod. Pl. Cap. 72 = Blaeria ericoides.
blanda, *Andr. Heathery*, iii. t. 107.—Afr. austr.
blanda, Sallab. in Trans. Linn. Soc. vi. (1802) 331 = cornuta.
Blandfordia, *Andr. Col. Heaths*, iii. 153.—Afr. austr.
Blandfordiana, Ait. Hort. Kew. ed. II. ii. 392 = praec.
blenna, Sallab. in Trans. Linn. Soc. vi. (1802) 379 = vernix.
Bonplandia, Lodd. Bot. Cab. t. 345 = seq.
Bonplandiana, Sims, Bot. Mag. t. 2126 = E. Muscari.
borbonisaefolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 286.—Afr. austr.
borbonica, Willd. ex Klotzsch, in Linnaea, xii. (1838) 220 = Philippia galioides.
botuliformis, Sallab. in Trans. Linn. Soc. vi. (1802) 369 = E. Tetralix.
Bouchrana, Kegel, in Gartenfl. (1852) 78.—Hab. ?
Bowerii, Donovan, in Nat. Repos. iii. (1824) t. 83, sphalm. = Bowieana.
Bowia, [Forbes], Hort. Eric. Woburn. 4 = Bowieana.
Bowieana, *Lodd. Bot. Cab.* t. 843.—Afr. austr.
brachialis, *Salisb. in Trans. Linn. Soc.* vi. (1802) 367.—Afr. austr.
brachycerata, *Benth. in DC. Prod.* vii. 686.—Afr. austr.
brachycrosa, *Tausch, in Flora*, xxii. (1839) ii. 634.—Hab. ?
bracata, Moench, Meth. 50 = carnea.
bracteata, *Thunb. Diss. Eric.* 18.—Afr. austr.
bracteolaris, *Lam. Encyc.* i. 481.—Afr. austr.
brevifolia, *Soland. ex Salisb. in Trans. Linn. Soc.* vi. (1802) 338.—Afr. austr.
brevipes, Bartl. in Linnaea, vii. (1832) 643 = coarctata.
Broadleyana, *Andr. Col. Heaths*, iii. 154.—Afr. austr.
Brockenthalii, Spreng. Neue Entdeck. i. 271 = Bruckenthalia spiculifolia.
brunioides, *Linn. Sp. Pl.* 354.—Afr. austr.
brunicefolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 341.—Afr. austr.
brunnea-alba, Bartl. in Linnaea, vii. (1832) 632 = imbricata.
bryantha, Thunb. Diss. Eric. 15 = Bryanthus Gmelini.
buccinaeformis, *Salisb. Prod.* 294.—Afr. austr.
buccinifiora, Sims, Bot. Mag. t. 2465 = Massoni.
bullaria, Steud. Nom. ed. II. i. 569, sphalm. = seq.
bullularia, Salisb. Prod. 296 = ramentacea.

ERICA :—
Burchellii, *Benth. in DC. Prod.* vii. 682.—Afr. austr.
byssina, Salisb. in Trans. Linn. Soc. vi. (1802) 333 = capitata.
caduca, *Thunb. Prod. Pl. Cap.* 71.—Afr. austr.
caduciflora, Salisb. in Trans. Linn. Soc. vi. (1802) 370 = palchella.
casta, Salisb. l. c. 349 = imbricata.
caffra, Andr. Col. Heaths, t. 7 = persoluta.
caffra, Linn. Dias. 9. n. 22 ; Amoen. Acad. viii. 50, 55 = arborea.
caffra, Linn. Sp. Pl. 353 = urceolaris.
calamiformis, Salisb. in Trans. Linn. Soc. vi. (1802) 361 = Massoni.
calathiflora, Salisb. l. c. 328 = bicolor.
caledonica, *Spreng. f. Tent. Suppl. Syst.* 18.—Afr. austr.
callosa, Wendl. Eric. ii. fasc. XVIII. 37. t. 33 = brevifolia.
calostoma, Lodd. Bot. Cab. t. 1759 = Massoni.
calycanthoides, Klotzsch, in Linnaea, xii. (1838) 539 = patula.
calyciflora, *Tausch, in Flora*, xvii. (1834) ii. 617.—Afr. austr.
calycina, Andr. Col. Heaths, t. 8 = corifolia, bracteata.
calycina, *Linn. Sp. Pl.* ed. II. 507.—Afr. austr.
calycina, Thunb. Diss. Eric. 47 = vespertina.
calycinades, [Forbes] Hort. Eric. Woburn. 4 = glomerata.
calyculata, Wendl. Eric. i. fasc. IV. 5 = penicilliflora.
campanularis, Salisb. in Trans. Linn. Soc. vi. (1802) 310 = campanulata.
campanulata, *Andr. Col. Heaths*, t. 9.—Afr. austr.
campanulata, F. W. Schmidt, Neue u. Selt Pfl. 47 = pyramidalis.
campylophylla, Spreng. Syst. ii. 190 = florida.
canaliculata, *Andr. Col. Heaths*, iii. 157.—Afr. austr.
candida, Bartl. in Linnaea, vii. (1832) 628 = E. Loea.
candida, *Spreng. Pugill.* i. 30.—Hab. ?
canescens, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 407 = villosa.
canescens, Wendl. Eric. ii. fasc. XXIII. 171. t. 65 = pusilla.
cantharoxiformis, Lodd. Bot. Cab. t. 1961 = reflexa.
capax, Salisb. in Trans. Linn. Soc. vi. (1802) 381 = obbata.
capillaris, Bartl. in Linnaea, vii. (1832) 647.—Afr. austr.
capillaris, Drège, ex Benth. in DC. Prod. vii. 680 = tenuis.
capitata, *Linn. Sp. Pl.* 355.—Afr. austr.
capitata, Thunb. Diss. Eric. 17 = brunioides.
carbonina, Salisb. in Trans. Linn. Soc. vi. (1802) 333 = brunioides.
cardulifolia, *Salisb. l. c.* 330.—Afr. austr.
carinata, *Klotzsch, ex Benth. in DC. Prod.* vii. 680.—Afr. austr.
carinata, Lodd. Bot. Cab. t. 1071 = Massoni.
carinula, Steud. Nom. ed. II. i. 570, sphalm. = inflata.
carnea, *Linn. Sp. Pl.* ed. II. 504.—Europ. austr.
carneola, G. Don, Gen. Syst. iii. 798 = inflata.
carminia, Lodd. Bot. Cab. t. 926 = inflata.
caroliniana, Hort. ex Benth. in DC. Prod. vii. 685 = varia.
catervaeflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 372.—Afr. austr.
catervaeflora, Pers. Syn. i. 431, sphalm. = praec.
Celsiana, Lodd. Bot. Cab. t. 177 = Thunbergii.
Cephalotes, *Thunb. Diss. Eric.* 21.—Afr. austr.
Cephalotes, Willd. ex Steud. Nom. ed. II. i. 570 = splendens.
cerinthoides, Berg. Desc. Pl. Cap. 104 = elongata.
cerinthoides, *Linn. Sp. Pl.* ed. II. 505.—Afr. austr.
cernua, Andr. Col. Heaths, t. 13 = verscundu.
cernua, *Linn. f. Suppl.* 322.—Afr. austr.
cernua-lanceolata, Wendl. Eric. i. fasc. VIII. 13 = praec.
cervicillora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 362.—Afr. austr.
Chamaetetralix, Tausch, in Flora, xvii. (1834) ii. 616 = strigosa.
Chamiacmia, *Klotzsch, ex Benth. in DC. Prod.* vii. 685.—Afr. austr.
chlamydiflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 338.—Afr. austr.
chloroloma, *Lindl. Bot. Reg.* (1836) t. 17.—Afr. austr.
chlorosepala, *Benth. in DC. Prod.* vii. 649.—Afr. austr.

ERICA :—
chrysantha, *Klotzsch, ex Benth. in DC. Prod.* vii. 630.—Afr. austr.
ciliaris, Huds. Fl. Angl. ed. I. 144 = Calluna vulgaris.
ciliaris, *Linn. Sp. Pl.* 354.—Europ. austr. occ.
ciliciflora, Salisb. in Trans. Linn. Soc. vi. (1802) 339 = Cristebachia velleriflora.
cinerascens, *Willd. Enum. Hort. Berol. Suppl.* 21.—Hab. ?
cinerea, *Linn. Sp. Pl.* 353.—Europ.
cistifolia, *Hort. Angl. ex Link, Enum. Hort. Berol.* i. 369.—Afr. austr.
clavaeflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 365.—Afr. austr.
clavata, Andr. Col. Heaths, iii. 159 = viridiflora.
Cliffordiana, Lodd. Bot. Cab. t. 34 = tenuiflora.
Cleoniana, Hort. ex Benth. in DC. Prod. vii. 643 = pinguis.
coarctata, *Wendl. Eric.* ii. fasc. XIX. 99. t. 37.—Afr. austr.
coccinea, *Berg. Desc. Pl. Cap.* 92.—Afr. austr.
coccinea, Linn. ex Benth. in DC. Prod. vii. 634 = tubiflora.
coccinea, Linn. Sp. Pl. 355 = Sebana.
coccinea, *Thunb. Diss. Eric.* 23.—Afr. austr.
cochlearifformis, Wendl. ex Klotzsch, in Linnaea, xii. (1838) 542 = caledonica.
codonodes, Lindl. Bot Reg. t. 1698 = lusitanica.
corrulea, Willd. Sp. Pl. ii. 393 = Bryanthus taxifolius.
Colletor, Spreng. Syst. ii. 192 = hispidula.
colorans, *Andr. Col. Heaths*, iv. 228.—Afr. austr.
comosa, *Linn. Mant.* ii. 254.—Afr. austr.
compacta, Bartl. in Linnaea, vii. (1832) 644 = catervaeflora.
complanata, Nois. ex Spreng. Syst. ii. 196 = crassifolia.
Comptoniana, Andr. Col. Heaths, iv. 224 = curvifolia.
concava, Lodd. Bot. Cab. t. 134 = bicolor.
concinna, Drège, ex Benth. in DC. Prod. vii. 629 = cruenta.
concinna, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 23.—Afr. austr.
condensata, *Benth. in DC. Prod.* vii. 681.—Afr. austr.
conferta, *Andr. Heathery*, ii. t. 59.—Afr. austr.
confertiflora, Steud. Nom. ed. II. i. 570, sphalm. = seq.
confertifolia, Wendl. Eric. ii. fasc. XX. 131. t. 46 = labrilis.
congesta, Lodd. Bot. Cab. t. 1743 = stylaris.
congesta, Wendl. Eric. ii. fasc. XVII. 75. t. 29 = persoluta.
conica, *Lodd. Bot. Cab.* t. 1179.—Afr. austr.
conifera, *Tausch, in Flora*, xxii. (1839) ii. 632.—Afr. austr.
coniflora, Klotzsch, ex Benth. in DC. Prod. vii. 655 = pumila.
conniveus, Klotzsch, ex Benth. l. c. 670 = filiformis.
conspicua, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 22.—Afr. austr.
Constantia, *Nois. ex Benth. in DC. Prod.* vii. 672.—Hab. ?
copiosa, *Wendl. Eric.* iii. fasc. XXV. 3. t. 1.—Hab. ?
coralloides, Hort. ex Tausch, in Flora, xvii. (1834) ii. 600 = mammosa.
corallina, *Andr. Col. Heaths*, iii. 160.—Afr. austr.
corifolia, Berg. Desc. Pl. Cap. 108 ; Thunb. Diss. Eric. 46 = obtecta.
corifolia, Linn. ex Benth. in DC. Prod. vii. 652 = bracteata.
corifolia, *Linn. Sp. Pl.* 355.—Afr. austr.
cornuta, *Roxb. ex Salisb. in Trans. Linn. Soc.* vi. (1802) 331, in syn.—Afr. austr.
coronata, Andr. Col. Heaths, t. 15 = fascicularis.
coronifera, *Benth. in DC. Prod.* vii. 683.—Afr. austr.
corsica, DC. Fl. Fr. iii. 677 = stricta.
corsica, Lichtst. ex Benth. in DC. Prod. vii. 679 = quadriceps.
Corydalis, *Salisb. in Trans. Linn. Soc.* vi. (1802) 334.
corymbosa, Bartl. in Linnaea, vii. (1832) 648 = catervaeflora.
corymbosa, *Tausch, in Flora*, xvii. (1834) ii. 618.—Afr. austr.
costaeflora, Salisb. in Trans. Linn. Soc. vi. (1802) 363 = E. Loea.
costata, Andr. Col. Heaths, t. 16 = versicolor.
cothurnalis, Salisb. Prod. 307 = Sebana.
Coventryna, Andr. Col. Heaths, iv. t. 226 = pavetteiflora.
Coventryana, Lodd. Bot. Cab. t. 432 = praec.
crassifolia, *Andr. Col. Heaths*, iv. 227.—Hab. ?

ERICA :—
crassifolia, Klotzsch, ex Benth. in DC. Prod. vii. 669 = latifolia.
crassinepala, *Benth. l. c.* 619.—Afr. austr.
crenata, E. Mey. ex Benth. l. c. 659 = mucosa.
cretica, Tausch, in Flora, xvii. (1834) ii. 604 = verticillata.
crinifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 367 = cerinthoides.
crinita, *Lodd. Bot. Cab.* t. 1432.—Afr. austr.
crinaeflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 332 —Afr. austr.
crocata, Spreng. Syst. ii. 201 = acuta.
cruciformis, Andr. Col. Heaths, iv. 228 = E. Muscari.
cruenta, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 16.—Afr. austr.
cubica, *Linn. Mant.* ii. 253.—Afr. austr.
cubica, Thunb. Diss. Eric. 31 = scriphifolia.
cubitalis, *Linn. Amoen. Acad.* viii. 51 = viscaria ?
cucullata, Tausch, in Flora, xvii. (1834) 615 = nigrita.
cuicilaeflora, Salisb. in Trans. Linn. Soc. vi. (1802) 357 = Massoni.
cumulata, Klotzsch, ex Benth. in DC. Prod. vii. 690 = variabilis.
cumuliflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 336.—Afr. austr.
cuneiformis, Benth. in DC. Prod. vii. 648 = E. Muscari
cupressifolia, Wendl. ex Klotzsch, in Linnaea, xii. (1838) 218 = Philippea Chamiasonis.
cupressiformis, Salisb. in Trans. Linn. Soc. vi. (1802) 354 = discolor.
cupressina, [Forbes] Hort. Eric. Woburn. 7 = turgera.
curviflora, *Linn. Sp. Pl.* 354, partim ; *Thunb. Diss. Eric.* 24.—Afr. austr.
curviflora, Linn. Sp. Pl. 354, partim = buccinaeformis.
curvifolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 380 —Afr. austr.
curvirostris, *Salisb. l. c.* 375.—Afr. austr.
cuspidata, Klotzsch, ex Benth. in DC. Prod. vii. 615 = pubescens.
cuspidigera, Salisb. in Trans. Linn. Soc. vi. (1802) 358 = ignescens.
cystiformis, Salisb. l. c. 378 = laevis.
cylindrica, Andr. Heathery, ii. t. 60 = Massoni.
cylindrica, Thunb. Diss. Eric. 24 = tenuiflora.
cylindrica, *Wendl. Eric.* i. fasc. XI. 7.—Hab. ?
cylindrifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 356 = serratifolia.
cymosa, *E. Mey. ex Benth. in DC. Prod.* vii. 670.—Afr. austr.
cyrillaeflora, Salisb. in Trans. Linn. Soc. vi. (1802) 357 = palustris.
Dabci, Crantz, Inst. ii. 350 = Daboecia polifolia.
Daboecia, Georgi, Beschr. Russ. Reich. III. iv. 936 = Bryanthus taxifolius.
Daboecia, Linn. Sp. Pl. ed. II. 509 = Daboecia polifolia.
daphniflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 334.—Afr. austr.
daphnoides, Lodd. Bot. Cab. t. 154 = praestans.
decipiens, *Spreng. f. Tent. Suppl. Syst.* 18.—Afr. austr.
decipiens, St. Amans, Fl. Agen. 159 = vagans
declinata, Lodd. Bot. Cab. t. 1662 = curvirostris.
decoriceus, *Willd. Enum. Hort. Berol. Suppl.* 21, nomen.—Hab. ?
decora, *Andr. Col. Heaths*, iii. 162.—Afr. austr.
decora, Salisb. in Trans. Linn. Soc. vi. (1802) 354 = E. Uhria.
decumata, Steud. Nom. ed. I. 304 = curvirostris.
decurrens, Klotzsch, ex Benth. in DC. Prod. vii. 632 = oblongiflora.
defexa, [*Forbes*] *Hort. Eric. Woburn.* 8.—Hab. ?
delecta, *Tausch, in Flora*, xxii. (1839) ii. 633.—Hab. ?
delicious, *Wendl. f. ex Benth. in DC. Prod.* vii. 666 = Afr. austr.
demissa, Hort. ex [Forbes] Hort. Eric. Woburn. 8 = soccifora.
demissa, *Klotzsch, ex Benth. in DC. Prod.* vii. 666.—Afr. austr.
densa, Andr. Col. Heaths, iii. 163 = ventricosa.
densiflora, Bartl. in Linnaea, vii. (1832) 623 = imbricata.
densiflora, Drège, ex Benth. in DC. Prod. vii. 629 = discolor.
densifolia, Willd. Sp. Pl. ii. 380 = E. Uhria.

ERICA :—
dentata. *Lam. Encyc.* i. 486.—Afr. austr.
dentata, Thunb. Diss. Eric. 28 = denticulata.
dentata, Wendl. Eric. ii. fasc. XIX. 101. t. 85 = venusta.
denticularis, Salisb. in Trans. Linn. Soc. vi. (1802) 334 = seq.
denticulata, *Linn. Mant.* ii. 229.—Afr. austr.
depressa, Andr. Heathery, i. t. 17 = nana.
depressa, *Linn. Mant.* ii. 230.—Afr. austr.
desmantha, *Bentth. in DC. Prod.* vii. 620.—Afr. austr.
dianthera, Mœnch, Meth. 50 = multiflora.
dianthiflora, *Tausch, in Flora*, xvii. (1834) ii. 594.—Hab.?
dianthiolita, *Salisb. in Trans. Linn. Soc.* vi. (1802) 333.—Afr. austr.
diaphana, Spreng. Syst. ii. 178 = transparens.
dichromata, Lodd. Bot. Cab. t. 1818 = dichrus.
dichrus, *Spreng. Syst.* ii. 178.—Afr. austr.
Dichrocarpi, Lodd. Bot. Cab. t. 1798 = pallens.
dicranifolia, *Tausch, in Flora*, xxii. (1839) ii. 634.—Hab.?
didyma, Stokes, in With. Bot. Arr. ed. II. i. 400 = vagans.
dilatata, *Wendl. f. ex Benth. in DC. Prod.* vii. 656.—Hab.?
dilecta, Hort. ex Klotzsch, in Linnæa, xii. (1838) 521 = fastigiata.
diosmæfolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 330.—Afr. austr.
diotæflora, *Salisb. l. c.* 342.—Afr. austr.
discolor, *Andr. Col. Heaths*, i. 20.—Afr. austr.
distans, *Benth. in DC. Prod.* vii. 646.—Afr. austr.
distans, Spreng. f. Tent. Suppl. Syst. 13 = thymifolia.
distorta, *Bartl. in Linnæa*, vii. (1832) 644.—Afr. austr.
divaricata, [Forbes] Hort. Eric. Woburn. 8 = tenuis.
divaricata, Lodd. ex Steud. Nom. ed. I. 805 = calycina.
divaricata, Wendl. Eric. i. fasc. VII. 5 = imbricata.
divergens, Wendl. ex Benth. in DC. Prod. vii. 670 = filiformis
doliiformis, Salisb. in Trans. Linn. Soc. vi. (1802) 368 = blanda.
Dregeana, Klotzsch, in Linnæa, xii. (1838) 501 = chlorioloma.
droseræfolia, *Tausch, in Flora*, xvii. (1834) ii. 600.—Hab.?
drosoroides, Lam. Encyc. i. 480 ; Andr. Col. Heaths, i. 21 = glutinosa.
ducalis, *Klotzsch, in Linnæa*, x. (1836) 347.—Afr. austr.
dumosa, *Andr. Col. Heaths*, iv. 220.—Afr. austr.
dumosa, Salisb. Prod. 296 = Blæsria purpurea.
dura, Hort. ex G. Don, Gen. Syst. iii. 791 = hirta.
dyscntha, *Benth. in DC. Prod.* vii. 674.—Afr. austr.
ebracteata, Hort. ex Benth. l. c. 679 = persoluta.
echiiflora, Andr. Col. Heaths, iii. 164 = coccinea.
Edelinia, *Bompl. Desc. Pl. Rar. Malm.* 48. t. 16.—Afr. austr.
effusa ×, Nichols. Dict. Gard. 592.
Ekloniana, *Tausch, in Flora*, xxii. (1839) ii. 625.—Hab.?
elata, Andr. Col. Heaths, ii. 87 = conspicua.
elata, Hoffmgg. & Link, Fl. Port. i. 411 = arborea.
elegans, *Andr. Heathery*, iii. t. 111.—Afr. austr.
elongata, *Lodd. Bot. Cab.* t. 788.—Afr. austr.
elongata, Roxb. ex G Don, Gen. Syst. iii. 793 = intervallaria.
emarginata, Andr. Col. Heaths, iii. 166 = nigrita.
embolifera, Salisb. in Trans. Linn. Soc. vi. (1802) 340 = Simochellus depressus.
emboliriifolia, Salisb. l. c. 379 = Nivenii.
empetrifolia, *Linn. Sp. Pl.* ed. II. 507.—Ads. austr.
empetrina, Linn. Syst. ed. X. 1008 = præc.?
empetroides, *Andr. Heathery*, i. t. 19.—Afr. austr.
ephemera, *Tausch, in Flora*, xxii. (1839) ii. 636.—Hab.?
epipteta, *Wildd. Enum. Hort. Berol. Suppl.* 21, nomen.—Hab.?
epistemia, Lodd. Bot. Cab. t. 1186 = saccifera.
equestris, Klotzsch, in Linnæa, x. (1836) 350 = arinella.
ericinifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 343 = Blæsria purpurea.
erectiuscula, Wendl. ex Klotzsch, in Linnæa, xii. (1838) 536 = brachiata.

ERICA :—
erianthe, Klotzsch, ex Benth. in DC. Prod. vii. 617 = brunialoides.
eriocephala, Andr. Heathery, ii. t. 61 = villosa.
eriocephala, Lam. Encyc. i. 489 (= *Syndesmanthus articulatus*).—Afr. austr.
eriopus, *Benth. in DC. Prod.* vii. 650.—Afr. austr.
eros, Lodd. Bot. Cab. t. 138 = Massoni.
erubescens, *Andr. Heathery*, iii. t. 118.—Afr. austr.
Esenbeckii, [Dryand. in] Hort. Kew. ed. II. ii. 268 = E. Uhria.
excelsa, *Tausch, in Flora*, xvii. (1834) ii. 596.—Hab.?
exigua, *Salisb. in Trans. Linn. Soc.* vi. (1802) 373.—Afr. austr.
exilis, Salisb. l. c. 340 (= *Syndesmanthus scaber*).—Afr. austr.
eximia, Lodd. Bot. Cab. t. 1105 = aristella.
expansa, Hort. ex Benth. in DC. Prod. vii. 686 = patens.
exposita, Lodd. Bot. Cab. t. 1521 = pellucida.
expromita, Spreng. Syst. ii. 195 = Simochellus depressus.
exquisita, *Carr. in Rev. Hortic.* (1880) 362.—Hab.?
exserens, Klotzsch, ex Benth. in DC. Prod. vii. 679 = persoluta.
exserta, [Forbes] Hort. Eric. Woburn. 9 = peltata.
exserta, Hort. ex Benth. in DC. Prod. vii. 689 = lasciva.
exsudans, Anct. vide *exudans*.
exsurgens, *Andr. Col. Heaths*, i. t. 22.—Afr. austr.
exudans, Andr. l. c. iv. 238 = pellucida.
fabrilis, *Salisb. in Trans. Linn. Soc.* vi. (1802) 338.—Afr. austr.
Fairesa, *Carr. in Rev. Hortic.* (1880) 467.—Hab.?
fallax, Salisb. in Trans. Linn. Soc. vi. (1802) 371 = parilis.
fascicularis, *Linn. f. Suppl.* 219.—Afr. austr.
fasciculata, Thunb. Prod. Pl. Cap. 71 = Simochellus glabellus.
fasciformis, Salisb. in Trans. Linn. Soc. vi. (1802) 332 = Blæsria purpurea.
fastigiata, Andr. Heathery, ii. t. 62 = E. Walkeria.
fastigiata, *Linn. Mant.* i. 66.—Afr. austr.
fastuosa, Salisb. in Trans. Linn. Soc. vi. (1802) 359 = curviflora.
fausta, *Salisb. l. c.* 326.—Afr. austr.
favosa, Salisb. Prod. 298 = spicata.
Ferguseni, *Gentili. & Carr. in Rev. Hortic.* (1882) 467.—Hab.?
ferox, Salisb. in Trans. Linn. Soc. vi. (1802) 324 = Ecremis Bartlingiana.
ferrea, Berg. Desc. Pl. Cap. 112 = mucosa.
ferruginea, Andr. Col. Heaths, iii. 168 = squarrosa.
ferta, Salisb. in Trans. Linn. Soc. vi. (1802) 371 = parilis.
Fibula, Link, Enum. Hort. Berol. i. 368 = lateralis.
filamentosa, *Andr. Heathery*, i. t. 22.—Afr. austr.
filifolia, *Regel, Ind. Sem. Hort. Petrop.* (1856) 29.—Hab.?
filiformis, Bartl. in Linnæa, vii. (1832) 636 = teretiuscula.
filiformis, *Salisb. in Trans. Linn. Soc.* vi. (1802) 345.—Afr. austr.
filipendula, *Benth. in DC. Prod.* vii. 668.—Afr. austr.
fimbriata, *Andr. Heathery*, ii. t. 63.—Afr. austr.
finitima, *Lodd. ex G. Don, in Loud. Hort. Brit.* 158.—Afr. austr.
fistulæflora, Salisb. in Trans. Linn. Soc. vi. (1802) 333 = tenuiflora.
flacca, *E. Mey. ex Benth. in DC. Prod.* vii. 670.—Afr. austr.
flaccida, Link, Enum. Hort. Berol. i. 367 = lanata.
flagellaria, Link, l. c. 365 = teretiuscula.
flagelliformis, Andr. Col. Heaths, iv. 234 = teretiuscula.
flammea, *Andr. Heathery*, i. t. 23.—Afr. austr.
flava, Andr. l. c. ii. t. 64 = parilis.
flava, Waltz, Beschr. Gatt. Art. Heid. 198 = Fetiverii.
flavicans, *Klotzsch, ex Benth. in DC. Prod.* vii. 651.—Afr. austr.
flavicoma, *Bartl. in Linnæa*, vii. (1832) 639.—Afr. austr.
flavida, Klotzsch, ex Benth. in DC. Prod. vii. 670 = filiformis.
flexicaulis, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 395.—Hab.?
flexilis, Salisb. Prod. 296 = racemosa.

ERICA :—
flexuosa, Andr. Col. Heaths, i. 23 = imbricata.
floccifera, *Benth. in DC. Prod.* vii. 660.—Afr. austr.
floccifloros, *Tausch, in Flora*, xxii. (1839) ii. 629.—Hab.?
floccosa, *Bartl. in Linnæa*, vii. (1832) 640.—Afr. austr.
floccosa, Salisb. in Trans. Linn. Soc. vi. (1802) 360 = conspicua.
floribunda, *Lodd. Bot. Cab.* t. 176.—Afr. austr.
floribunda, Wendl. Eric. ii. fasc. XIV. 19 = nudiflora.
florida, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 428 = anthina.
florida, Lodd. Bot. Cab. t. 284 = moschata.
florida, *Thunb. Diss. Eric.* 40.—Afr. austr.
floscullosa, Salisb. in Trans. Linn. Soc. vi. (1802) 340 (= *Syndesmanthus paucyfolius*).—Afr. austr.
foliacea, *Andr. Col. Heaths*, iv. 226.—Afr. austr.
foliacea, Klotzsch, ex Benth. in DC. Prod. vii. 691 = hispidula.
folliculoris, Salisb. in Trans. Linn. Soc. vi. (1802) 348 = Fetiverii.
Forbesiana, *Klotzsch, in Linnæa*, x. (1836) 349.—Afr. austr.
formosa, Andr. Heathery, iii. t. 114 = purpurea.
formosa, *Thunb. Diss. Eric.* 49. t. 3.—Afr. austr.
Forskåhlii, Vitm. Summa Pl. ii. 414 = vagans.
fragilis, Salisb. in Trans. Linn. Soc. vi. (1802) 346 = E. Banksia.
fragrans, *Andr. Col. Heaths*, ii. 95.—Hab.?
fragrans, Salisb. in Trans. Linn. Soc. vi. (1802) 363 = E. Muscari.
frondosa, Salisb. Prod. 296 = coccinea.
fucata, *Klotzsch, ex Benth. in DC. Prod.* vii. 667.—Afr. austr.
fucata, Thunb. Diss. Eric. 15 = scoparia.
fugax, Salisb. in Trans Linn. Soc. vi. (1802) 351 = trifora.
fulgens, Klotzsch. in Linnæa, xii. (1838) 507 = foliacea.
fulgida, [*Forbes*] *Hort. Eric. Woburn.* 11.—Afr. austr.
furcæflora, Salisb. Prod. 294 = pulchella.
furfuracea, Salisb. in Trans. Linn. Soc. vi. (1802) 348 = monadelpha.
fuscata, F. G. Dietr. ex Steud. Nom. ed. I. 805, 806 = lanuginosa.
fuxiformis, Salisb. Prod. 297 = Plukenetii.
galiiflora, Bartl. in Linnæa, vii. (1832) 648 = floribunda.
galiiflora, Salisb. Prod. 296 = comosa.
galioides, Klotzsch, ex Benth. in DC. Prod. vii. 691 = hispidula.
galioides, Poir. Encyc. Suppl. i. 728 = Philippia galioides.
geisela, Andr. Heathery, i. t. 24 = gilva.
gemmifera, Lodd. Bot. Cab. t. 457 = Massoni.
genistæfolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 337.—Afr. austr
gibbosa, *Klotzsch, ex Benth. in DC. Prod.* vii. 678.—Afr. austr.
gigantea, *Klotzsch, ex Benth. l. c.* 656.—Afr. austr.
Gilii, *Benth. l. c.* 684.—Afr. austr.
gilva, *Wendl. Bot. Beobacht.* 48.—Afr. austr.
glabella, Thunb. Prod. Pl. Cap. 73 = Simochellus glabellus.
glabra, Gilib. Fl. Lituan. i. 3 = Calluna vulgaris.
glabra, Link, Enum. Hort. Berol. i. 362 = ventricosa.
glabra, Thunb. Prod. Pl. Cap. 69 = Simochellus glaber.
glandulifera, *Klotzsch, in Linnæa*, x. (1836) 353.—Afr. austr.
glandulosa, Andr. Col. Heaths, ii. 97 = flexicaulis.
glandulosa, Thunb. Diss. Eric. 25 = pellucida.
glandulosa, Wendl. Eric. ii. fasc. XIII. 5 = coccinea.
glauca, *Andr. Col. Heaths*, i. 24.—Afr. austr.
glaucocerus, Bartl. in Linnæa, vii. (1832) 651 = E. Lees.
glaucifolia, Klotzsch, ex Benth. in DC. Prod. vii. 620 = imbricata.
globosa, *Andr. Heathery*, iii. t. 116.—Afr. austr.
globosa, *Burm. f. Pl. Cap. Prod.* 11.—Afr. austr.
globosa, *Willd. Sp. Pl.* ii. 406.—Afr austr.
globulifera, Klotzsch, ex Benth. in DC. Prod. vii. 667 = demissa.
glomerata, *Andr. Col. Heaths*, iv. 237.—Afr. austr.
glomerata, [Forbes] Hort. Eric. Woburn. 11 = plumosa.
glomerata, Salisb. Prod. 293 = E. Tetralix.
glomiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 330 = reflexa.
glumæflora, *Klotzsch, ex Benth. in DC. Prod.* vii. 689.—Afr. austr.

ERICA :—
glutinosa, Andr. Col. Heaths, i. 25 = E. Lees, onosmæflora.
glutinosa, *Berg. Desc. Pl. Cap.* 98.—Afr. austr.
glutinosa, Roxb. ex G. Don, Gen. Syst. iii. 795 = verniciflua.
gnaphalisflora, Salisb. in Trans. Linn. Soc. vi. (1802) 357 = gnaphalodes.
gnaphaloides, Berg. Desc. Pl. Cap. 119 = calycina.
gnaphaloides, *Thunb. Diss. Eric.* 45.—Afr. austr.
gnaphaloides, *Linn. Sp. Pl.* 356 (sp. dub.).—Afr. austr.
gnaphaloides, E. Mey. ex Klotzsch, in Linnæa, xii. (1838) 541 = Meyeriana.
gnidiacifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 336 = vespertina.
Gordonia, Forbes, Hort. Woburn. 81 = nitida.
gorteriæfolia, Salisb. in Trans. Linn. Soc. vi. (1802) 381 = retorta.
gracilis, *Salisb. l. c.* 375.—Afr. austr.
gracilis, *Wendl. Bot. Beobacht.* 47.—Afr. austr.
grandiflora, *Linn. f. Suppl.* 223.—Afr. austr.
grandinosa, Lodd. Bot. Cab. t. 627; Andr. Col. Heaths, iv. 238 = formosa.
granulata, Linn. Mant. ii. 224 = ramentacea.
grisea, Klotzsch, ex Benth. in DC. Prod. vii. 674 = hirtiflora.
grossa, Salisb. Prod. 292 = halicacaba.
guttæflora, Salisb. in Trans. Linn. Soc. vi. (1802) 374 = lateralis.
haematostoma, *Wendl. Eric.* iii. fasc. xxvi. 18.—Afr. austr.
hallicacaba, *Linn. Pl. Rar. Afr.* 11; *Mant.* ii. 274.—Afr. austr.
halocalycina, Buek, in DC. Prod. Index, iii. 166 = variabilis.
Hartnellii, Hort. ex Klotzsch, in Linnæa, xii. (1838) 510 = crinita.
hebecalyx, *Benth. in DC. Prod.* vii. 630.—Afr. austr.
helianthemifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 326 = mariifolia.
hemisphærica, Klotzsch, ex Benth. in DC. Prod. vii. 678 = veltiaris.
herbacea, Georgi, Beschr. Russ. Reich. III. iv. 985 = Calluna vulgaris.
herbacea, Linn. Sp. Pl. 352 = carnea.
Hesseana, *Wendl. ex Klotzsch, in Linnæa*, ix. (1834) 636.—Afr. austr.
Hibbertia, *Andr. Heathery*, iii t. 118.—Afr. austr.
hibernica, Syme, Eng. Bot. ed. III. vi. 42 = mediterranea.
hirsuta, *Klotzsch, ex Benth. in DC. Prod.* vii. 669.—Afr. austr.
hirsuta, Lodd. Bot. Cab. t. 754 = persicina.
hirsuta, Salisb. in Trans. Linn. Soc. vi. (1802) 339 = Grisebachia hispida.
hirsuta, Thunb. Prod. Pl. Cap. 72 (= *Acrastemon hirsutus*).—Afr. austr.
hirta, Andr. Col. Heaths, ii. 101 = speciosa.
hirta, *Thunb. Diss. Eric.* 36. t. 2.—Afr. austr.
hirta, Wendl. Eric. iii. fasc. xxvi. 16 = dilatata.
hirtiflora, *Curt. Bot. Mag.* t. 481.—Afr. austr.
hirtifolia, *Hornem. Hort. Hafn.* i. 270.—Afr. austr.
hispida, Andr. Heathery, ii. t. 69 = racemosa.
hispida, *Burm. f. Fl. Cap. Prod.* 11.—Afr. austr.
hispida, Thunb. Diss. Eric. 19 = hispidula.
hispidula, *Linn. Sp. Pl.* ed. II. 1672.—Afr. austr.
halocalycina, Klotzsch, ex Benth. in DC. Prod. vii. 690 = variabilis.
holosericea, Salisb. in Trans. Linn. Soc. vi. (1802) 352 = andromedæflora.
horizontalis, Audr. Heathery, ii. t. 70 = cumuliflora.
hottentæflora, Salisb. in Trans. Linn. Soc. vi. (1802) 331 = cubica.
Humeana, Lodd. Bot. Cab. t. 389 = fastigiata.
humifusa, *Hibb. ex Salisb. in Trans. Linn. Soc.* vi. (1802) 333.—Afr. austr.
humilis, *Benth. in DC. Prod.* vii. 615.—Afr. austr.
humilis, Neck. Delic. Gallo-Belg. i. 182 = cinerea.
humilis, Salisb. in Trans. Linn. Soc. vi. (1802) 329 = depressa.
hyacinthoides, Andr. Col. Heaths, iii. 174 = praestans.
hybrida, Hort. ex Benth. in DC. Prod. vii. 688 = Mussoni.
hypocrateriformis, Tausch, in Flora, xx. (1837) 402 = pavettæflora.
hyssopifolia, Salisb. in Trans. Linn. Soc. iv. (1802) 337 = tricteata.
hystricifloræ, Salisb. l. c. 367 = Sparmanni.
ignescens, *Andr. Heathery*, i. t. 27.—Afr. austr.

ERICA :—
ignorata, Klotzsch, ex Benth. in DC. Prod. vii. 676 = verecunda.
imbecilla, Sweet, Hort. Brit. ed. I. 268 = gracilis, *Salisb.*
imbellis, Salisb. Prod. 298 = lutea.
imbricata, *Linn. Sp. Pl.* ed. II. 508.—Afr. austr.
imperialis, Andr. Col. Heaths, iv. 239 = fascicularis.
inaperta, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 429 = cerviciflora.
incana, Wendl. Eric. ii. fasc. xviii. 89 = pubescens.
incarnata, Andr. Col. Heaths, i. 28 = lateralis.
incarnata, *Thunb. Diss. Eric.* 50.—Afr. austr.
incarnata, Klotzsch, in Linnæa, xii. (1838) 521 = daphniflora.
Incluse, *Wendl. f. ex Benth. in DC. Prod.* vii. 654.—Afr. austr.
inclyta, Sweet and Salisb. in Trans. Linn. Soc. vi. (1802) 330 = tubercularis.
incurva, *Klotzsch, ex Benth. in DC. Prod.* vii. 690.—Afr. austr.
inconspicua, *Bartl. in Linnæa*, vii. (1832) 642.—Afr. austr.
inconspicua, *Thunb. Prod. Pl. Cap.* 71.—Afr. austr.
incurvata, Andr. Col. Heaths, iii. 178 = floridula.
incurva, *Thunb. Prod. Pl. Cap.* 188.—Afr. austr.
incurva, Wendl. Bot. Beobacht. 47 = stellata.
indica, Walts, Beschr. Gatt. Art. Heid. 165 = Menziesia trifora.
indigesta, Klotzsch, in Linnæa, xii. (1838) 523 = daphniflora.
inermis, Klotzsch, l. c. 505 = exigua.
inflata, Hort. ex Benth. in DC. Prod. vii. 644 = jasminiflora.
inflata, *Thunb. Diss. Eric.* 41.—Afr. austr.
inflexa, Pers. Syn. i. 428 = incurva.
infundibuliformis, Andr. Col. Heaths, iv. 240 = pavettæflora.
infundibuliformis, Bartl. in Linnæa, vii. (1832) 638 = Lawsoniana.
Ingrami ×, Hort. ex Morr. Belg. Hortic. vii. (1857) 232. t. 6. f. 1.
innocua, *Hoffmgg. Vers. Pfl. Nachtr.* ii. 111.—Afr. austr.
insulsa, [*Forbes*] *Hort. Eric. Woburn.* 18.—Hab.?
intermedia, *Klotzsch, ex Benth. in DC. Prod.* vii. 621.—Afr. austr.
intertexta, Lodd. Bot. Cab. t. 1034 = empetrifolia.
intervallaris, *Salisb. in Trans. Linn. Soc.* vi. (1802) 375.—Afr. austr.
involucrata, *Klotzsch, ex Benth. in DC. Prod.* vii. 617.—Afr. austr.
involucrata, *Benth. l. c.* 619.—Afr. austr.
Irbyana, *Andr. Col. Heaths*, iii. 176.—Afr. austr.
irregularis, *Benth. in DC. Prod.* vii. 680.—Afr. austr.
isanthera, *Benth. l. c.* 660.—Afr. austr.
Jacksonii ×, Paxt. Mag. Bot. viii. (1841) 149.
jasminiflora, *Andr. Col. Heaths*, t. 29. t. 26.—Afr. austr.
jasminiflora, *Salisb. Prod.* 293.—Afr. austr.
jasminoides, *Carr. in Rev. Hortic.* (1852) 362.—Hab.?
jubata, Lodd. ex Sprengel, St. 198 = melanthera.
Juliana, Lodd. Bot. Cab. t. 799 = E. Walkeria.
juncea, Salisb. in Linnæa, vii. (1832) 648 = regerminans.
juniperifolia, *Salisb. Prod.* 296.—Afr. austr.
Kennedyana, Klotzsch, in Linnæa, xii. (1838) 506 = lactiflora.
Krausiana, *Klotzsch, ex Walp. Rep.* ii. 738.—Afr. austr.
labialis, Salisb. in Trans. Linn. Soc. vi. (1802) 340 = Sympieza capitellata.
Lachnaea, *Andr. Heathery*, iii. t. 190.—Afr. austr.
lachnaeoides, *G. Don, Gen. Syst.* iii. 795.—Afr. austr.
lachneæfolia, Salisb. in Trans. Linn. Soc. vi. (1802) 385 = E. Lachnaea.
lactea, Lees, ex Klotzsch, in Linnæa, xii. (1838) 506 = lactiflora.
lactiflora, *Lodd. Bot. Cab.* t. 990.—Hab.?
lacunæflora, Salisb. Prod. 297 = Bergiana.
laeta, *Bartl. in Linnæa*, vii. (1832) 648.—Afr. austr.
lacvigata, *Bartl. l. c.* 636.—Afr. austr.
laevis, *Andr. Col. Heaths*, iii. 182.—Afr. austr.
laevis, *Sprengl. Pugill.* i. 20.—Hab.?
lagenæformis, Salisb. in Trans. Linn. Soc. vi. (1802) 362 = jasminiflora.
Lambertia, Andr. Col. Heaths, ii. 104 = physodes.
lamellaris, Salisb. in Trans. Linn. Soc. vi. (1802) 327 = urceolaris.
lamprotes, *Klotzsch, in Linnæa*, xii. (1838) 536.—Afr. austr.
lanata, *Andr. Heathery*, iii. t. 121.—Afr. austr.

ERICA :—
lanata, Wendl. Bot. Beobacht. 45 = conspicua.
lanceolaris, Stend. Nom. ed. 2. 806, sphalm. = hirta.
lanceolata, *Pers. Syn.* i. 424.—Hab.?
laniflora, F. W. Schmidt, Neue u. Selt. Pfl. 44 = ignescens.
lanuginosa, Wendl. Eric. i. fasc. ii. 28 = conspicua.
lanuginosa, *Andr. Heathery*, iii. t. 122.—Afr. austr.
larices, *Burm. f. Fl. Cap. Prod.* 11.—Afr. austr.
laricifolia, Lam. Encyc. i. 487 = imbricata.
laricina, Berg. Desc. Pl. Cap. 94 = nigrita.
laricina, *Spreng. f. Tent. Suppl. Syst.* 19.—Afr. austr.
lasciva, *Salisb. in Trans. Linn. Soc.* vi. (1802) 349.—Afr. austr.
lasiandra, Klotzsch, ex Benth. in DC. Prod. vii. 677 = rubens.
lasiocephala, Klotzsch, ex Benth. l. c. 617 = bruniades.
lasiophylla, Spreng. Syst. ii. 195 = strigosa.
lateralis, *Willd. Sp. Pl.* ii. 880.—Afr. austr.
latifolia, *Andr. Heathery*, ii. t. 72.—Afr. austr.
lavandulæfolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 332.—Afr. austr.
Lawsoni, Sims, Bot. Mag. t. 1720 = E. Lawsonia.
Lawsonia, *Andr. Col. Heaths*, iv. 242.—Afr. austr.
laxa, *Andr. Heathery*, ii. t. 73.—Afr. austr.
laxa, Lam. Encyc. i. 480 = mammosa.
laxa, *Thunb. Prod. Pl. Cap.* 189.—Afr. austr.
laxiflora, *Buek, in DC. Prod. Index*, iii. 167.—Afr austr.
Lees, *Andr. ex Willd. Sp. Pl.* ii. 400, *et Col. Heaths*, i. 31.—Afr. austr.
Leeana, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 376 = praec.
Lehmanni, *Klotzsch, ex Benth. in DC. Prod.* vii 618 —Afr. austr.
leiophylla, *Benth. l. c.* 684.—Afr. austr.
lenitformis, Salisb. Prod. 294 = umbellata.
lepidota, *Rach, in Linnæa*, xxvi. (1855) [1855] 786 — Afr. austr.
leptocarpha, Spreng. f. Tent. Suppl. Syst. 18 = Lawsonia.
leptocephala, Klotzsch, ex Benth. in DC. Prod. vii. 620 = imbricata.
leptoclada, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 94.—Afr. austr.
leptophylla, Klotzsch, ex Benth. in DC. Prod. vii 620 = tiarseflora.
leptopus, *Benth. l. c.* 680.—Afr. austr.
leucantha, G. Don, Gen. Syst. iii 792 = regerminans.
leucantha, *Link, Enum. Hort. Berol.* i. 371.—Afr. austr.
leucanthera, Andr. Heathery, i. t. 28 = praeo.
leucanthera, *Linn. f. Suppl.* 223.—Afr. austr.
leucodesmia, *Benth. in DC. Prod.* vii. 681.—Afr. austr
leucopelta, *Tausch, in Flora*, xvii. (1834) ii. 616.—Hab.?
leucophylla, Klotzsch, in Linnæa, xii. (1838) 507 = modesta.
leucosepala, Klotzsch, ex Benth. in DC. Prod. vii. 689 = glumæflora.
leucostoma, *Tausch, in Flora*, xvii. (1834) ii. 596 — Hab.!
Lichtensteinii, Klotzsch, in Linnæa, xii. (1838) 504 = regerminans.
lilacina, Klotzsch, ex Benth. in DC. Prod. vii. 657 = gnaphalodes.
linaria, *Benth. l. c.* 622.—Afr. austr.
linifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 386 = tenuifolia.
Linnæa, Andr. Heathery, ii. t. 75 = persicina.
Linnæanna, Alt. Hort. Kew. ed. II. ii. 375 = persicina.
linnaeoides, Andr. Col. Heaths, iii. 107 = persicina.
litoralis, *Regel, Cat. Pl. Hort. Akushow.* 55, nomen.—Hab.!
littuiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 356 = perspicua.
longiaristata, *Benth. in DC. Prod.* vii. 668.—Afr. austr.
longiflora, Klotzsch, ex Benth. l. c. 680 = tenuis.
longiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 359 = conspicua.
longifolia, *Donn, Hort. Cantab.* ed. I. 43; *ex Willd. Sp. Pl.* ii. 398.—Afr. austr.
longipedicellata, Hoffmgg. Verz. Pfl. 59 = longipedunculata.
longipedunculata, Lodd. Bot. Cab. t. 198 = dumosa.
longipedunculata, *Wender. ex Steud. Nom.* ed. II. i. 576.—Hab.?
longipes, Bartl. in Linnæa, vii. (1832) 649 = coarctata.
longipes, *Klotzsch, ex Benth. in DC. Prod.* vii. 684.—Afr. austr.

ERICA :—
lucida, *Salisb. in Trans. Linn. Soc.* vi. (1802) 337 ; *Andr. Col. Heaths*, ii. 109.—Afr. austr.
lugubris, Salisb. Prod. 295 = carnea.
lusitanica, *Rudolph, in Schrad. Journ.* ii. (1799) 286. —Europ. occ.
lutea, *Berg. Desc. Pl. Cap.* 115 ; *Linn. Mant.* ii. 254. —Afr. austr.
luteo-alba, [Forbes] Hort. Eric. Woburn. 14 = leucantha.
lutescens, Hort. ex C. Koch, Dendrol. ii. 11. 138 = Calluna vulgaris.
lychnides, *Wendl. ex Steud. Nom.* ed. I. 807, nomen.—Hab. ?
Lycopodiastrum, *Lam. Tabl. Encyc.* ii. 428.—Hab. ?
lycopodistifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 361 = Massoni.
lycopodioides, *Hornem. Hort. Hafn.* i. 374.—Afr. austr.
lycopodioides, *Lodd. Cat. ex Benth. in DC. Prod.* vii. 619.—Hab. ?
lycopodioides, Waitz, Beschr. Gatt. Art. Heid. 156 = Cassiope lycopodioides.
lyngrea, Salisb. in Trans. Linn. Soc. vi. (1802) 336 = nigrita.
lysimachiaeflora, Salisb. l c. 339 = melanthera.
Mackaiana, Bab. in Mackay, Fl. Hibern. 181 = E. Tetralix.
Machaii, Hook. in Comp. Bot. Mag. i. (1835) 158 = prasc.
Macnabiana ×, Paxt. Mag. Bot. vii. (1840) 135.
macrolonia, *Benth. in DC. Prod.* vii. 661.—Afr. austr
macrophylla, *Klotzsch, ex Benth. l. c.* 669.—Afr. austr.
macrostoma, Klotzsch, ex Benth. l c. 688 = floribunda.
maculosa, *Tausch, in Flora*, xxii. (1839) ii. 629.—Hab. ?
magnifica, Andr. Col. Heaths, iv. 244 = Massoni
mallotiana, Salisb. in Trans. Linn. Soc. vi. (1802) 370 = empetrifolia.
mammosa, *Linn. Mant.* ii. 284.—Afr. austr.
mammosa, Thunb. Diss. Eric. 42 = blanda.
manipuliflora, Salisb. in Trans. Linn. Soc. vi. (1802) 344 = verticillata.
margaritacea, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 20. —Afr. austr.
marginata, Bartl. in Linnæa, vii. (1832) 687 = caledonica.
marginata, Benth. in DC. Prod. vii. 687, sphalm. = nigrita.
marifolia, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 15.—Afr. austr.
maricoides, Hort. ex Benth. in DC. Prod. vii. 678 = nidularia.
Martinesii, Lag. ex Benth. l. c 665 = E. Tetralix.
Massoni, *Linn. f. Suppl.* 221.—Afr. austr.
Mauxenia, Andr. Col. Heaths, i. 36 = prasc.
mauritanica, Linn. Syst. ed. X. 1002 = pelviformis.
media, *Klotzsch, ex Benth. in DC. Prod.* vii. 682.—Afr. austr.
mediiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 331 = Thunbergii.
mediterranea, *Linn. Mant.* ii. 229.—Europ. austr.
mediterranea, Reichb. ex Nym. Coasp. 489 = carnea.
mediterranea, Sieber, ex Tausch, in Flora, xvii. (1834) ii. 604 = verticillata.
melanthera, *Linn. Mant.* ii. 232.—Afr. austr.
melanthera, Thunb. Prod. Pl. Cap. 89 = cristaeflora.
melastoma, Andr. Col. Heaths, i. 37 = Petiverii.
mellifera, Hort. Angl. ex Link, Enum. Hort. Berol. i. 370 = turgida.
melliflua, Salisb. Prod. 293 = cruenta.
mera, Klotzsch, in Linnæa, ix. (1834) 644 = fulgida.
Mertensiana, *Wendl. ex Klotzsch, l. c.* 659.—Hab. ?
metulaeflora, Curt. Bot. Mag. t. 812 = Massoni.
metulaeflora, Klotzsch, in Linnæa, xii. (1838) 502 = Meuroni.
Meuroni, *Benth. in DC. Prod.* vii. 636.—Afr. austr.
Meyeriana, *Klotzsch, in Linnæa*, xii. (1838) 541.—Afr. austr.
microcalyx, *Regel, Ind. Sem. Hort. Petrop.* (1856) 29. —Hab. ?
microstoma, Berg. Desc. Pl. Cap. 97 = rudiflora.
millefiora, Berg. l. c. 96 = penicularia.
minuta, Klotzsch, ex Benth. in DC. Prod. vii. 691 = hispidula.
minutiflora, Andr. Col. Heaths, iv. 245 = coarctata.
minutissima, *Klotzsch, ex Benth. in DC. Prod.* vii. 691. —Afr. austr.
mira, *Klotzsch, ex Benth. l. c.* 680.—Afr. austr.

ERICA :—
mirabilis, Andr. Col. Heaths, iv. 246 = praestans.
mitraeformis, Salisb. in Trans. Linn. Soc. vi. (1802) 372 = hirtiflora.
modesta, [Forbes] Hort. Eric. Woburn. 15 = mollis.
modesta, *Salisb. in Trans. Linn. Soc.* vi. (1802) 352.—Afr. austr.
mollearis, Pers. Syn. i. 422, sphalm. = empetrifolia.
mollis, *Andr. Col. Heaths*, iv. 247.—Afr. austr.
mollissima, *Lodd. ex G. Don, in Loud. Hort. Brit.* 150.—Hab. ?
monadelpha, Curt. Bot. Mag. t. 1370 = E. Bankeia.
monaliphia, *Andr. Col. Heaths*, i. 38.—Afr. austr.
monilliformis, Salisb. Prod. 293 = baccans.
Monsoniana, *Linn. f. Suppl.* 223.—Afr. austr.
montana, [Forbes] Hort. Eric. Woburn. 15 = fabrilis.
monticola, Klotzsch, ex Benth. in DC. Prod. vii. 684 = lavandulaefolia.
Mooreana ×, Lem. Jard. Fleur. iii. (1852) t. 259.
moschata, *Andr. Col. Heaths*, iv. 246.—Afr. austr.
moschata, Lodd. Bot. Cab. t. 614 = E. Muscari.
mucosa, *Linn. Mant.* ii 232.—Afr. austr.
mucronides, Lodd. ex [Forbes,] Hort. Eric. Woburn. 15 ; Bot. Cab. t. 1202 = mucosa.
mucronata, *Andr. Col. Heaths*, iii. 186.—Afr. austr.
multicaulis, Salisb. in Trans. Linn. Soc. vi. (1802) 369 = stricta.
multiflora, Bivona, Sic. Pl. Cent. i. 33 = peduncularis.
multiflora, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 353 = verticillata.
multiflora, Huds. Fl. Angl. ed. II. 166 = vagans.
multiflora, *Linn. Sp Pl.* 355 —Reg. Mediterr.
multiglandulosa, Willd. ex Klotzsch, in Linnæa, xii. (1838) 230 = Phillipsia viscosa.
multumbellata, Benth. in DC. Prod. vii. 658, sphalm. = ramentacea.
multumbellifera, Berg. Desc Pl. Cap. 110 = ramentacea.
multumbellifera, Hort. ex Benth. in DC. Prod. vii. 679 = pelviformis.
multumbelliflora, *Tausch, in Flora*, xxii. (1839) ii. 628. —Hab. ?
multumbracteata, *Tausch, in Flora*, xxii. (1839) ii. 630.—Hab. ?
munda, Salisb. in Trans. Linn. Soc. vi. (1802) 337 = nigrita.
Mundtiana, Klotzsch, ex Benth. in DC. Prod. vii. 684 = lutea.
mundula, Andr. Col. Heaths, iv. 249 = fastigiata.
muricata, Wendl. f. ex Benth. in DC. Prod. vii. 685 = varia.
murina, Klotzsch, ex Benth. l. c. 679 = distorta.
Muscari, *Andr. Col. Heaths*, i. 40.—Afr. austr.
muscicapa, *Tausch, in Flora*, xvii. (1834) ii. 595.—Hab. ?
mutabilis, Andr. Col Heaths, iii. 187 = cruenta.
mutabilis, Salisb. in Trans. Linn. Soc. vi. (1802) 369 = cinerea.
mutica, *Tausch, in Flora*, xvii. (1834) ii. 597.—Hab. ?
myrantha, Bartl. in Linnæa, vii. (1832) 688 = imbricata.
narmadicola, Nym. in Bot. Notiser (1851), 133 = verticillata.
nana, Hort. ex C. Koch, Dendrol. ii. 11. 138 = Calluna vulgaris.
nana, *Salisb. in Trans. Linn. Soc.* vi. (1802) 355.—Afr. austr.
neglecta, G. Don, Gen. Syst. iii. 792 = gracilis, *Wendl. Willd.* x., Hort. ex Paxt. Mag. Bot. ix. (1842) 195.
nemorosa, Klotzsch, ex Benth. in DC. Prod. vii. 688 = floribunda.
nidicularia, G. Don, Gen. Syst. iii. 799, sphalm. = nidularia.
nidiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 333 = E. Muscari.
nidularia, *Lodd. Bot. Cab.* t. 764.—Hab. ?
nigricans, Lodd. ex G. Don, in Loud. Hort. Brit. 151 = seq.
nigrita, *Linn. Mant.* i. 15.—Afr. austr.
nitens, Bartl. in Linnæa, vii. (1832) 652 = incida.
nitens, [*Forbes*] *Hort. Eric. Woburn.* 16.—Hab. ?
nitens, Lee, ex Stend. Nom. ed. II. i. 576 = Bergiana.
nitida, *Andr. Heathery*, iii. t. 181.—Afr. austr.
nitidula, Hort. ex Benth. in DC. Prod. vii. 687 = nigrita.
nivalis, *Andr. Col. Heaths*, iv. 250.—Afr. austr.
nives, [*Forbes*] *Hort. Eric. Woburn.* 16.—Afr. austr.
Nivenia, *Andr. Heathery*, ii. t. 76.—Afr. austr.

ERICA :—
nodiflora, *Klotzsch, in Linnæa*, xii. (1838) 539.—Afr. austr.
nodiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 340 = Grisebachia plumosa.
nolaeflora, Salisb. l. c. 371 = plumosa.
notabilis, Wendl. ex Spreng. Syst. ii. 184 = Eremia tuberculoris.
nudicaulis, Berg. Desc. Pl. Cap. 118 = pillaliflora.
nudiflora, Hort. ex Benth. in DC. Prod. vii. 678 = nidularia.
nudiflora, *Linn. Mant.* ii. 229.—Afr. austr.
nutans, G. Don, Gen. Syst. iii. 796 = rubens.
nutans, *Wendl. Eric. t. fasc.* iii. 5.—Hab. ?
obbata, *Andr. Heathery*, t. t. 32.—Afr. austr.
obcordata, [Forbes] Hort. Eric. Woburn. 17 = bracteata.
obera, Salisb. Prod. 294 = margaritacea.
obesa, *Tausch, in Flora*, xxii. (1839) ii. 628.—Hab. ?
obliqua, *Ait. Exot. Pl.* t. 3 ; *Thunb. Diss. Eric.* 44.—Afr. austr.
oblonga, [Forbes] Hort. Eric. Woburn. 17 = Massoni.
oblongiflora, *Benth. in DC. Prod.* vii. 632.—Afr. austr.
oblongifolia, Steud. Nom. ed. II. i. 577, sphalm. = prasc.
obpyramidalis, Salisb. Prod. 298 = pyramidalis.
obtecta, *Tausch, in Flora*, xxii. (1839) ii. 638.—Afr. austr.
obtusa, Lodd. Bot. Cab. t. 1027 = brevifolia.
obtusata, *Klotzsch, ex Benth. in DC. Prod.* vii. 615.—Afr. austr.
oboallaris, Salisb. in Trans. Linn. Soc. vi. (1802) 336 = bracteata.
ochroleuca, Wendl. f. ex Benth. in DC. Prod. vii. 687 = nigrita.
octogona, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 147.—Hab. ?
octophylla, Thunb. Diss. Eric. 44 = fascicularis.
odorata, *Andr. Col. Heaths*, iii. 191.—Afr. austr.
odorata, Spreng. Neue Entdeck. i. 271 = laevis.
oellula, Andr. Col. Heaths, iv. 251 = cruenta.
olympiaca, Sibth. ex Salisb. in Trans. Linn. Soc. vi. (1802) 324 = Bruckenthalia spiculifolia.
onosmaeflora, *Salisb. l. c.* 363.—Afr. austr.
ophthylla, *Benth. in DC. Prod.* vii. 672.—Afr. austr.
oppositifolia, Andr. Col. Heaths, iii. 192 = lutea.
opulenta, *Wendl. ex Klotzsch, in Linnæa*, xii. (1838) 499.—Afr. austr.
orbicularis, Lodd. Bot. Cab. t. 153 = Blaeria ericoides.
ornata, Klotzsch, ex Benth. in DC. Prod. vii. 620 = imbricata.
ostrina, Lodd. Bot. Cab. t. 1218 = coccinea.
ovata, Lodd. l. c. t. 417 = hirtiflora.
ovata, Wendl. ex Steud. Nom. ed. II. i. 577 = versicolor.
ovina, *Klotzsch, ex Benth. in DC. Prod.* vii. 674.—Afr. austr.
oxycoccoides, *Salisb. in Trans. Linn. Soc.* vi. (1802) 324.—Afr. austr.
Pabsti, *Regel, Gartenfl.* (1858) 50.—Hab. ?
pachycephala, Klotzsch, ex Benth. in DC. Prod. vii. 689 = lasciva.
pachyphylla, Spreng. Syst. ii. 199 = brevifolia.
padibunda, Pers. Syn. i. 431, sphalm. = autana.
palaeacea, Klotzsch, ex Benth. in DC. Prod. vii. 620 = imbricata.
palaeacea, Salisb. in Trans. Linn. Soc. vi. (1802) 341 (= *Syndesmanthe arbiculatus*).—Afr. austr.
pallens, *Andr. Col. Heaths*, iii. 194.—Hab. ?
pallens, *Spreng. Pugill.* i. 30.—Hab. ?
pallescens, Klotzsch, in Linnæa, xii. (1838) 541 = Meyeriana.
pallida, Lodd. Bot. Cab. t. 1355 = Massoni.
pallida, Salisb. in Trans. Linn. Soc. vi. (1802) 326 = pubescens.
pallida, Wendl. Eric. iii. fasc. xxv. 11. t. 5 = pallidiflora.
pallidiflora, Klotzsch, ex Benth. in DC. Prod. vii. 679 = persoluta.
pallide-purpurea, Linn. Sp. Pl. 354 = vagans.
pallidiflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 351. —Afr. austr.
paludosa, Salisb. Prod. 293 = concinna.
palustris, *Andr. Heathery*, ii. t. 77.—Afr. austr.
paniculata, *Linn. Sp. Pl.* ed. II. 508.—Afr. austr.
paniculata, Lodd. Bot. Cab. t. 1194 = gnaphaloides.
paniculata, Thunb. Prod. Pl. Cap. 72 = Simocheilus obovatus.

ERICA :—

paniculata, Wendl. ex Stend. Nom. ed. II. i. 577 = persoluta.
pannosa, Salisb. in Trans. Linn. Soc. vi. (1802) 289 = barbata.
parilis, *Salisb. l. c.* 371.—Afr. austr.
Parmentieri, Lodd. Bot. Cab. t. 197 = praestans.
parviflora, *Linn. Sp Pl.* ed. II. 506.—Afr. austr.
Passerinae, *Linn. f. Suppl.* 231; *Thunb. Diss. Eric.* 18.—Afr. austr.
passerinaefolia, Salisb. in Trans. Linn. Soc. vi. (1802) 282 = E. Passerinae.
patens, *Andr. Col. Heaths*, ii. 115.—Hab. ?
Paternoma, Thunb. Prod. Pl. Cap. 189; Andr. Col. Heaths. i. 43 = abietina.
paterzoinoides, [Forbes] Hort. Eric. Woburn. 18 = Massoni.
patula, *Klotzsch, in Linnaea*, xii. (1838) 538.—Afr. austr.
patula, K. Mey. ex Benth. in DC. Prod. vii. 680 = leptopsa.
pavettaeflora, *Salisb. Prod.* 297.—Afr. austr.
Paxtoni, *Gentilh. & Carr. in Rev. Hortic.* (1882) 407.—Hab. ?
pectinata, Bartl. in Linnaea, vii. (1832) 647 = Eremia totta.
pectinata, *Klotzsch, in Linnaea*, xii. (1838) 509.—Afr austr.
pectiniifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 367 = erubescens.
pedicellata, *Klotzsch, in Linnaea*, xii. (1838) 525.—Afr. austr.
peduncularis, J. & C. Presl. Delic. Prag. 89 = multiflora.
peduncularis, *Salisb. in Trans. Linn. Soc.* vi. (1802) 339.—Afr. austr.
pedunculata, Andr. Col. Heaths, iv 252 = pilulifera.
pedunculata, Wendl. Eric. ii. fasc. XXIII. 173. t. 66 = lateralis.
pellucida, *Andr. Col. Heaths*, iii. 197.—Afr. austr.
pellucida, Salisb. in Trans. Linn. Soc. vi. (1802) 384, ex parte = venusta.
pellucidoides, [Forbes] Hort. Eric. Woburn. 18 = pellucida.
peltata, *Andr. Col. Heaths*, iv. 254.—Afr. austr.
peltiformis, *Salisb. Prod.* 298.—Afr. austr.
pendula, Lodd. Bot. Cab. t. 902 = rubens.
pendula, Wendl. Bot. Beobacht. 48 = stricta.
penicillata, *Andr. Col. Heaths*, ii. 116.—Afr. austr.
penicillata, Lodd. Bot. Cab. t. 1918 = Plukenetii.
penicilliflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 348.—Afr. austr.
penicilliformis, Salisb. Prod. 297 = praec. ?
pentaphylla, Linn. Sp. Pl. ed. II. 506 = urceolaris.
peripiocaefolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 387.—Afr. austr.
perlata, [*Forbes*] *Hort. Eric. Woburn.* 18.—Afr. austr.
perlata, Lee, ex Benth. in DC. Prod. vii. 672 = globosa.
persoluta, Curt. Bot. Mag. t. 842 = peliviformis.
persoluta, *Linn. Mant.* ii. 280.—Afr. austr.
perspicua, Andr. Col. Heaths, iv. 255 = perspicua, *Wendl.* ?
perspicua, [Forbes] Hort. Eric. Woburn. 18 = Massoni.
petaphora, *Wendl. Bot. Beobacht.* 45.—Afr. austr.
peroviscoides, [Forbes] Hort. Eric. Woburn. 18 = Massoni.
petiolaris, *Lam. Encyc.* i. 487.—Afr. austr.
petiolaris, Salisb. in Trans. Linn. Soc. vi. (1802) 334 = petiolata.
petiolata, *Thunb. Diss. Eric.* 15.—Afr. austr.
Petraea, Linn. Mant. ii. 280 = Sebana.
Petiveriana, Andr. Col. Heaths, i. 44 = seq.
Petiverii, *Willd. Sp. Pl.* ii. 894.—Afr. austr.
petraea, *Benth. in DC. Prod.* vii. 668.—Afr. austr.
Petrea, Lodd. Bot. Cab. t. 965 = rivalis.
pharetraeformis, Salisb. Prod. 295 = exsurgens.
phylicaefolia, Salisb. in Trans. Linn. Soc. vi. (1802) 364 = purpurea.
phylicoides, Willd. Sp. Pl. ii. 861 = pulchella.
physantha, *Benth. in DC. Prod.* vii. 619.—Afr. austr.
physodes, *Linn. Syst. ed.* X. 1002.—Afr. austr.
physophylla, *Benth. in DC. Prod.* vii. 682.—Afr. austr.
pocta, [Forbes] Hort. Eric. Woburn. 19 = vestiflua.
pilifera, Klotzsch, in Linnaea, xii. (1838) 505 = trichophora.
pilifera, *Thunb. Prod. Pl. Cap.* 186.—Afr. austr.
pilosa, Lodd. Bot. Cab. t. 606 = villosa.
pilulaeformis, Benth. in DC. Prod. vii. 680, sphalm. = pilulifera.

ERICA :—

pilularis, Berg. Desc. Pl. Cap. 111 = ramentacea.
pilularis, Lodd. Bot. Cab. t. 1568 = mucosa.
pilulifera, Andr. Heathery, t. 278 = mucosa.
pilulifera, *Linn. Sp. Pl.* 355.—Afr. austr.
pilulifera, Wendl. Eric. ii. fasc XIX. 107. t. 41 = strigosa.
piluliformis, Salisb. in Trans. Linn. Soc. vi. (1802) 378 = pilulifera.
pinea, Andr. Col. Heaths, t. 46 = longifolia.
pinea, *Thunb. Diss. Eric.* 28.—Afr. austr.
pinea, Wendl. Eric. i. fasc. I. 11 = E. Lcea.
pinguis, *Klotzsch, in Linnaea*, x. (1836) 351.—Afr. austr.
pinifolia, Andr. Col. Heaths, ii. 119 = longifolia.
pinifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 362 = vestita.
piniformis, Wendl. ex Klotzsch, in Linnaea, ix. (1834) 526 = longifolia.
pistillaris, Salisb. in Trans. Linn. Soc. vi. (1802) 368 = australis.
pityophylla, Synst. ii. 161 = longifolia.
placentaeflora, Salisb. in Trans. Linn. Soc. vi. (1802) 346 = tiaraeflora.
plantifolia, *Linn. Sp. Pl.* ed. II. 508.—Afr. austr.
plantifolia, Wendl. Eric. ii. fasc. XVI. 59 = thymifolia.
Plukenetii, *Linn. Sp. Pl.* 356.—Afr. austr.
Plukenetiana, Ait. Exot. Pl. t. 9 = præc.
Plukenetii, Berg. Desc. Pl. Cap. 91 = Sebana.
plumigera, Bartl. in Linnaea, vii. (1832) 636 = chlamydiflora.
plumosa, *Andr. Heathery*, i. t. 36.—Afr. austr.
plumosa, Thunb. Prod. Pl. Cap. 73 = Griesbachia ciliaris.
plumosa, Wendl. Eric. i. fasc. XII. 5 = serratifolia.
podophylla, *Benth. in DC. Prod.* vii. 681.—Afr. austr.
pogonanthera, *Bartl. in Linnaea*, vii. (1832) 634.—Afr. austr.
Pohlmanni, Lodd. Bot. Cab. t. 1852 = E. Lachnaea.
polifolia, *Salisb. ex Benth. in DC. Prod.* vii. 615.—Afr. austr.
polyantha, *Klotzsch, ex Benth. l. c.* 688.—Afr. austr.
polycodon, *Benth. l. c.*—Afr. austr.
polygonaeflora, *Lodd. Bot. Cab.* t. 1805.—Afr. austr.
polygalaeflora, *Klotzsch, in Linnaea*, xii. (1838) 537.—Afr. austr.
polytricha, Sweet, Hort. Brit. ed. II. 338 = speciosa.
polytrichifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 329 = lusitanica.
pomifera, Hort. ex Benth. in DC. Prod. vii. 654 = andromedaeflora.
porrigens, Bartl. in Linnaea, vii. (1832) 638 = imbricata.
praecox, *Klotzsch, in Linnaea*, xii. (1838) 517.—Afr. austr.
praecox, Lodd. Bot. Cab. t. 1418 = strigosa.
praegnans, Andr. Col. Heaths, iii. 203 = ventricosa.
presnitera, *Tausch, in Flora*, xvii. (1834) ii. 596.—Hab. ?
prenestas, *Andr. Col. Heaths*, iv. 257.—Afr. austr.
prenestis, Bartl. in Linnaea, vii. (1832) 645 = publigera.
primuloides, Andr. Col. Heaths, iii. t. 203 = fastigiata.
Princeps, Andr. l. c. ii. 121 = Massoni.
procera, Salisb. in Trans. Linn. Soc. vi. (1802) 338 = arborea.
procera, Wendl. Bot. Beobacht. 45 = ignescens.
procumbens, *Lodd. Bot. Cab.* t. 1993.—Hab. ?
prolifera, Salisb. in Trans. Linn. Soc. vi. (1802) 376 = persoluta.
propendens, *Andr. Col. Heaths*, ii. 122.—Afr. austr.
prostrata, Hort. ex C. Koch, Dendrol. ii. 1. 138 = Calluna vulgaris.
protrudens, *Link, Enum. Hort. Berol.* i. 372.—Afr. austr.
protrusa, Salisb. Prod. 293 = australis.
Pseudoventilata, Benth. in Mssnd, Botanist, iii. t. 104 = Massoni ?
puberula, Bartl. in Linnaea, vii. (1832) 644 = hirtiflora.
puberula, *Klotzsch, l. c.* ix. (1834) 529.—Afr. austr.
pubescens, Andr. Col. Heaths, i. t. 48 = hirtiflora.
pubescens, Curt. Bot. Mag. t. 460 = parviflora.
pubescens, *Linn. Sp. Pl.* ed. II. 506.—Afr. austr.
pubescens, Sieber, ex Benth. in DC. Prod. vii. 674 = mollis.
pubigera, *Salisb. in Trans. Linn. Soc.* vi. (1802) 372.—Afr. austr.
pudibunda, Salisb. l. c. 345 = nutans.
pugioniflolia, Salisb. l. c. 327 = tetragona.

ERICA :—

pulchella, *Houtt. Handleid.* iv. 504; *Andr. Col. Heaths*, i. 49.—Afr. austr.
pulchella, Thunb. Diss. Eric. 22 = E. Lcea.
pulchella, Willd. ex Stend. Nom. ed. II. i. 578 = Onosmaefiora.
pulcherrima, G. Don, in Sweet, Hort. Brit. ed. III 434 = jasminiflora.
pulchra, Salisb. in Trans. Linn. Soc. vi. (1802) 384 = E. Walkera.
pulverulenta, [*Forbes*] *Hort. Eric. Woburn.* 20—Afr austr.
pulviniformis, Salisb. Prod. 295 = coccinea.
pumilla, *Andr. Col. Heaths*, iv. 258.—Afr. austr.
punctata, Bartl. in Linnaea, vii. (1832) 646 = corolata
pura, Andr. ex Benth. in DC. Prod. vii. 626 = E. Lcea
pura, Lodd. Bot. Cab. t. 73 = pubescens.
puriatis, Hoss, ex [Forbes] Hort. Eric. Woburn. 20 = pubescens.
purpurascens, Lam. Encyc. i. 488 = multiflora.
purpurascens, Linn. Sp. Pl. ed. II. 508 = mediterranea.
purpurea, *Andr. Col. Heaths*, i. 50.—Afr. austr.
purpurea, Thunb. Prod. Pl. Cap. 71 = Blaeria pur purea.
pusilla, *Salisb. in Trans. Linn. Soc.* vi. (1802) 374.—Afr. austr.
pusilla, Thunb. Prod. Pl. Cap. 70 = Blaeria pusilla.
pycnantha, *Benth. in DC. Prod.* vii. 654.—Afr. austr
pygmaea, Andr. Col. Heaths, iv. 259 = sicaefolia.
pyramidalis, *Ait. Hort. Kew.* ed. I. iii. 491.—Afr. austr
pyramidalis, Salisb. in Trans. Linn. Soc. vi. (1802) 349 = imbricata.
pyramidiformis, *Wendl. Eric.* ii. fasc. XXIV. 181 — Hab. ?
pyrolaeflora, Salisb. in Trans. Linn. Soc. vi. (1802) 351 = trifiora.
pyxidiflora, Salisb. l. c. 371 = empetroides.
quadriaeflora, *Salisb. l. c.* 375.—Afr. austr.
quadrangularis, Andr. Col. Heaths, iv. 260 = Massoni.
quadrangularis, Salisb. Prod. 297 = denticulata.
quadrata, Lodd. Bot. Cab. t. 1948 = formosa.
quadriflora, Andr. Heathery, t. t. 43 = Bergiana.
quadriflora, Klotzsch, in Linnaea, ix (1834) 565 = dichrus.
quadriflora, *Willd. Sp. Pl.* ii. 879.—Afr. austr.
quadrifolia, Prits. Ic. Ind. i. 421, sphalm. vide *quadriflora*, Andr.
quadrifossa, Salisb. Prod. 293 = mammosa.
quinquangularis, Berg. Desc. Pl. Cap. 112 = imbricata
racemifera, Andr. Col. Heaths, iii. 204 = regerminans.
racemosa, *Thunb. Diss. Eric.* 31.—Afr. austr.
Rachii, *Regel, Ind. Sem. Hort. Petrop.* (1856) 30.—Hab. ?
radiata, Andr. Col. Heaths, i. 52 = Massoni.
radiiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 360 = fasciculata.
ramentacea, *Linn. Mant.* ii. 282.—Afr. austr.
ramosissima, Kunth, ex Klotzsch, in Linnaea, xii. (1838) 520 = Philippis gallicides.
ramosissima, Roxb. ex Stend. Nom. ed. I. 809 = salsa.
ramosissima, Wendl. Eric. ii. fasc. XVIII. 93. t. 36 = mollis.
ramulosa, Bartl. in Linnaea, vii. (1832) 633 = imbricata.
ramulosa, Viv. Ann. Bot. i. 17. 189 = stricta.
Rastellum, Sprang. Syst. ii. 196 = penicilliflora.
recurvata, *Andr. Col. Heaths*, iv. 262.—Afr. austr.
reflexa, *Link, Enum. Hort. Berol.* i. 371.—Afr. austr.
refulgens, Andr. Col. Heaths, iv. 264 = versicolor.
regerminans, Andr. l. c. 265 = peliviformis.
regerminans, Hort. ex Benth. in DC. Prod. vii. 679 = persoluta.
regerminans, *Linn. Mant.* ii. 282.—Afr. austr.
regia, *Bartl. in Linnaea*, vii. (1832) 630.—Afr. austr.
Regiana, Hort. ex C. Koch, Dendrol. ii. 1. 138 = Calluna vulgaris.
resinosa, Sims, Bot. Mag. t. 1139 = E. Vernix.
retorta, *Linn. f. Suppl.* 230.—Afr. austr.
retroflexa, Wendl. Eric. i. fasc. VIII. 7 = pulchella.
retusa, *Tausch, in Flora*, xvii. (1834) ii. 598.—Hab. ?
revoluta, Bartl. in Linnaea, vii. (1832) 631 = Plukenetii.
rigens, Benth. in DC. Prod. vii. 687, sphalm. = laxa.
rigescens, Bartl. in Linnaea, vii. (1832) 635 = laxa.
rigida, Lodd. Bot. Cab. t. 1906 = Massoni.
rigidifolia, Wendl. Eric. ii. fasc. XX. 125. t. 46 = incida.
rigidissima, Wendl. ex Klotzsch, in Linnaea, ix. (1834) 547 = purpurea.
Rinzli, *Regel, Ind. Sem. Hort. Petrop.* (1857) 46.—Afr. austr. ?

ERICA :—
Kollissonia, [*Forbes*], *Hort. Eric. Woburn.* 21.—Afr. austr.
rosea, Andr. Heathery, ii. t. 83 = vestita.
rosea, Lichtenst. ex Klotzsch, in Linnæa, xii. (1838) 504 = regerminans.
rostellis, [Forbes], Hort. Eric. Woburn. 21 = penicilliflora.
rostrata, Bartl. in Linnæa, vii. (1832) 632 = penicilliflora.
Roxburghii, *Benth. in DC. Prod.* vii. 682.—Afr. austr.
rubella, Ker-Gawl. in Bot. Mag. t. 2165 = glomerata.
rubens, *Andr. Heathery*, i. t. 43.—Afr. austr.
rubens, Thunb. Diss. Eric. 49 = peduncularia.
rubercalyx, Andr. Col. Heaths, iv. 266 = Massoni ?
rubicunda, Klotzsch, in Linnæa, xii. (1838) 540 = teretiascula.
rubida, Lodd. Bot. Cab. t. 1166 = Massoni.
rubrocalyx, Gentlib. & Carr. in Rev. Hortic. (1852) 306, sphalm. = Massoni.
rubrosepala, Sweet, Hort. Brit. ed. II. 389 = Massoni.
rufescens, *Klotzsch, in Linnæa*, x. (1836) 882.—Afr. austr.
rugosa, Andr. Col. Heaths, iv. 267 = Massoni.
rupestris, Andr. t. c. ii. 128 = depressa.
rupestris, Salisb. in Trans. Linn. Soc. vi. (1802) 253 = halicacaba.
ruptcola, *Klotzsch, in Linnæa*, xii. (1838) 504.—Afr. austr.
Russeliana, *Andr. Col. Heaths*, iv. 269.—Hab. ?
Russelliana, *Lodd. Bot. Cab.* t. 1018.—Hab. ?
sacciflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 255. —Afr. austr.
sagittata, *Klotzsch, ex Benth. in DC. Prod.* vii. 681.—Afr. austr.
sagittifolia, Stokes, Bot. Mat. Med. ii. 370 = Calluna vulgaris.
Sainsburyana, Andr. Col. Heaths, iv. 270 = infusa.
salax, *Salisb. in Trans. Linn. Soc.* vi. (1802) 256.—Afr. austr.
Salisburia, Andr. Col. Heaths, iv. 271 = purpurea.
sanguinea, *Lodd. Bot. Cab.* t. 86.—Afr. austr.
sanguinolenta, Lodd. Bot. Cab. t. 468 = sicaefolia.
saturejæfolia, *Tausch, in Flora*, xxii. (1839) ii. 686.—Hab. ?
Savilea, *Andr. Heathery*, t. 238.—Afr. austr.
Savileana, Sweet, Hort. Brit. ed. I. 261 = præc.
Savilleia, Andr. Col. Heaths, iii. t. 205 = præc.
Savillia, Lodd. Bot. Cab. t. 95 = præc.
saxatilis, Salisb. Prod. 295 = canea.
scabra, F. W. Schmidt, Neue u. Selt. Pfl. 42 = hispida.
scabra, Thunb. Prod. Pl. Cap. 72 (= *Syndesmanthus scaber*).—Afr. austr.
scabriuscula, Drège, ex Benth. in DC. Prod. vii. 678 = strigosa.
scabriuscula, *Link, Enum. Hort. Berol.* i. 372.—Hab. ?
scabriuscula, *Lodd. Bot. Cab.* t. 517.—Afr. austr.
scariosa, Berg. Desc. Pl. Cap. 102 = spumosa.
scariosa, Lodd. Bot. Cab. t. 477 = acuta.
scariosa, Thunb. Prod. Pl. Cap. 188 = capillaris.
sceptriformis, Salisb. in Trans. Linn. Soc. vi. (1802) 256 = spicata.
Schelliana, Lodd. Bot. Cab. t. 538 = plumosa.
scolastoma, Klotzsch, in Linnæa, x. (1836) 822 = E. Muscari.
scoparia, C. Koch, in Linnæa, xxiii. (1850) 624 = arborea.
scoparia, *Linn. Sp. Pl.* 353.—Reg. Mediterr. occ.
scoparia, Thunb. Diss. Eric. 48 = arborea.
Sebana, *Donn, Hort. Cantab.* 48 ; *ex Willd. Sp. Pl.* ii. 395.—Afr. austr.
secundiflora, Tausch, in Flora, xvii. (1834) ii. 617 = decora.
selaginifolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 253.—Afr. austr.
semimicrota, Drège, ex Benth. in DC. Prod. vii. 680 = depressa.
senilis, *Klotzsch, ex Benth. l. c.* 637.—Afr. austr.
serpens, Salisb. Prod. 293 = physodes.
serpiniflolia, *Salisb. l. c.* 257.—Afr. austr.
serpyllifolia, Andr. Heathery, vi. 389 ; Lodd. Bot. Cab. t. 746 = hispidula.
serrata, Thunb. Prod. Pl. Cap. 69 = hispidula.
serratifolia, *Andr. Col. Heaths*, i. 58.—Hab. ?
serrulata, [Forbes], Hort. Eric. Woburn. 23 = diosmæfolia.

ERICA :—
sertiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 242 = rudiflora.
sessiliflora, Andr. Heathery, ii. t. 86 = clavæflora.
sessiliflora, Linn. f. Suppl. 223 = spicata.
sessiliflora, Wendl. Eric. ii. fasc. XXII. 149 t. 56 = genistæfolia.
setacea, *Andr. Col. Heaths*, i. 59.—Afr. austr.
setacea, Drège, ex Benth. in DC. Prod. vii. 679 = persoluta.
setifera, *Klotzsch, ex Benth. l. c.* 616.—Afr. austr.
setosa, *Bartl. in Linnæa*, vii. (1832) 646.—Afr. austr.
setula, Klotzsch, ex Benth. in DC. Prod. vii. 689 = glumæflora.
setulosa, *Benth. l. c.* 682.—Afr. austr.
sexfaria, *Ait. Hort. Pl.* t. 11.—Afr. austr.
Shuttleworthii, Hort. Berol. ex Klotzsch, in Linnæa, xii. (1838) 498 = Eremia parviflora.
Shannonea, *Andr. Col. Heaths*, iv. 273 ; *Heathery*, t. 239.—Afr. austr.
Shannoneana, Spreng. Syst. ii. 185 = præc.
Shannoni, Lodd. Bot. Cab. t. 168 = præc.
Shannonius, Sweet, Hort. Brit. ed. I. 261 = præc.
Skannoniana, Benth. in DC. Prod. vii. 644 = præc.
sicæfolia, *Salisb. in Trans. Linn. Soc.* vi. (1802) 226. —Afr. austr.
sicula, Guss. Adnot. Cat. Pl. Boccad. 6 = Pentapera sicula.
Sieberiana, Klotzsch, ex Benth. in DC. Prod. vii. 673 = caterviæflora.
simplicifolia, Willd. Sp. Pl. ii. 402 = cervifloræ et buccinæformis.
simplicifolia, Steud. Nom. ed. I. 809 = præc.
sitiens, *Klotzsch, in Linnæa*, xii. (1838) 505.—Afr. austr.
Smithiana, Lodd. Bot. Cab. t. 1614 = regerminans.
socciflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 247. —Afr. austr.
sodalis, Klotzsch, ex Benth. in DC. Prod. vii. 690 = paniculata.
Soldeck, *Andr. Heathery*, ii. t. 89.—Afr. austr.
Solandriana, Benth. in DC. Prod. vii. 658 = præc.
solandroides, Andr. Col. Heaths, iv. 274 = stellata.
sordida, Andr. t. c. i. 60 = spicata.
sordida, Drège, ex Benth. in DC. Prod. vii. 684 = tubiflora.
Sparmanni, *Linn. f. in Vet. Acad. Handl. Stockh.* (1772) 24. t. 2.—Afr. austr.
sparsa, Klotzsch, ex Benth. in DC. Prod. vii. 620 = imbricata.
spuria, Lodd. Bot. Cab. t. 1467 = floribunda.
speciosa, *Andr. Col. Heaths*, ii. 138.—Afr. austr.
speciosa, Schaeev. lc. Pl. Rar. 3 = mammosa.
speciosissima, Klotzsch, in Linnæa, ix. (1834) 645 = fulgida.
spectabilis, *Klotzsch, ex Benth. in DC. Prod.* vii. 659. —Afr. austr.
spectabilis, Walts, Beschr. Gatt. Art. Heid. 220 = grandiflora.
sphærantha, Link, ex Stend. Nom. ed. II. i. 580 = tenera.
sphæranthа, *Spreng. Syst. iv. Cur. Post.* 146.—Afr. austr.
sphærocephala, *Wendl. ex Benth. in DC. Prod.* vii. 686.—Hab. ?
sphenanthera, *Tausch, in Flora*, xxii. (1839) ii. 626.—Hab. ?
spicata, *Thunb. Diss. Eric.* 42. t. 4.—Afr. austr.
spiculifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 224 = Bruckenthalia spiculifolia.
spiraeæfolia, Salisb. l. c. 250 = leucanthera.
spinula, *Lodd. ex G. Don, in Loud. Hort. Brit.* 153, nomen.—Hab. ?
spiraea, Hoffmgg. Vers. Pfl. Nachtr. ii. 28 = odorata.
spiralifolia, Salisb. Prod. 298 = abietina.
splendens, Andr. Col. Heaths, iv. 275 = tomida.
splendens, Wendl. Eric. i. fasc. VIII. 5 = conspicua.
splendida, *Mackay, ex G. Don, in Loud. Hort. Brit.* 146, nomen.—Hab. ?
Sprengelii, Endl. in Harting. Parad. Vindob. t. 67. f. 2 = aristata.
Sprengelii, Sweet, ex Stend. Nom. ed. II. i. 580 = lactiflora.
spumosa, *Linn. Pl. Rar. Afr.* 11 ; *Mant.* ii. 275.—Afr. austr.
spumosa, Thunb. Diss. Eric 17 = sexfaria.
spumosa, Wendl. Eric. ii. fasc. XVI. 51. t. 19 = tricops.
spuria, Andr. Col. Heaths, i. 62 = Massoni.

ERICA :—
squamæflora, Salisb. in Trans. Linn. Soc. vi. (1802) 249 = imbricata.
squamosa, *Andr. Heathery*, ii. t. 91.—Afr. austr.
squarrosa, *Salisb. in Trans. Linn. Soc.* vi. (1802) 260. —Afr. austr.
stagnalis, Salisb. l. c. 259 = sulphurea.
staminea, *Andr. Col. Heaths*, iii. 208.—Afr. austr.
Steinbergiana, *Wendl. f. ex Klotzsch, in Linnæa*, xii. (1838) 531.—Afr. austr.
stellaris, Nois. ex G. Don, Gen. Syst. iii. 827 = seq.
stellata, *Lodd. Bot. Cab.* t. 893.—Afr. austr.
Stelleriana, Willd. Sp. Pl. ii. 387 = Cassiope Stelleriana.
stellifera, Andr. Col. Heaths, iv. 276 = Massoni.
stenantha, *Klotzsch, ex Benth. in DC. Prod.* vii. 685.—Afr. austr.
stenantha, Sweet, Hort. Brit. ed. II. 840 = tenuiflora
stenoma, *Tausch, in Flora*, xxii. (1839) ii. 633.—Hab. ?
stenophylla, *Benth. in DC. Prod.* vii. 679.—Afr. austr.
stricta, *Donn, Hort. Cantab.* 45 ; *ex Willd. Sp. Pl.* ii. 396.—Europ. austr.
strictiflora, Steud. Nom. ed. II. i. 580, sphalm. = seq.
strictifolia, Klotzsch, in Linnæa, ix. (1834) 646 = grandiflora.
strigiflora, Steud. Nom. ed. II. i. 580, sphalm. = seq.
strigilifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 267 = elongata, erubescens.
strigosa, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 17.—Afr. austr.
strigosa, Wendl. Eric. i. fasc. II. 25 = persoluta.
struthiolaefora, Lodd. ex G. Don, in Loud. Hort. Brit. 148 = seq. ?
struthiolaefolia, Bartl. in Linnæa, vii. (1832) 636 = teretiuscula.
strutiaefolia, Steud. Nom. ed. II. i. 580, sphalm. = præc.
stylaris, *Spreng. Syst.* ii. 198.—Afr. austr.
stylosa, Rudolph, in Schrad. Journ. (1800) ii. 229 = arborea.
stylosa, Salisb. in Trans. Linn. Soc. vi. (1802) 250 = imbricata.
suaveolens, *Andr. Col. Heaths*, iv. 277.—Afr. austr.
suaveolens, Lodd. Bot. Cab. t. 24 = latifolia.
suavis, Bartl. in Linnæa, vii. (1832) 641 = cornuta.
subdivaricata, Berg. Desc. Pl. Cap. 114 = persoluta.
subterminalis, *Klotzsch, ex Benth. in DC. Prod.* vii. 670.—Afr. austr.
subulata, *Wendl. Eric.* ii. fasc. XX. 123. t. 47.—Afr. austr.
suffulta, *Wendl. ex Benth. in DC. Prod.* vii. 619.—Afr. austr.
sulcata, *Benth. l. c.* 682.—Afr. austr.
sulcata, *Hornem. Hort. Hafn.* i. 371.—Afr. austr.
sulphurea, *Andr. Col. Heaths*, iv. 278.—Afr. austr.
superba, *Hoffmgg. Vers. Pfl. Nachtr.* iii. 35.—Hab. ?
superba, Sweet, Hort. Brit. ed. II. 340 = purpurea ?
Swainsonia, Andr. Col. Heaths, iii. 209 = Massoni.
Syndriana, *Hort. ex Gentlib. & Carr. in Rev. Hortic.* (1852) 306.—Hab. ?
tardiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 273 = parviflora.
taxifolia, [*Dryand. in*] *Ait. Hort. Pl.* t. 19 ; *Wendl. Bot. Beobacht.* 44.—Afr. austr.
tegulaefolia, Salisb. in Trans. Linn. Soc. vi. (1802) 251 = spumosa.
Templea, Lee, ex Andr. Col. Heaths, iv. 280 = Massoni ?
tenella, *Andr. Heathery*, ii. t. 94.—Hab. ?
tenera, Klotzsch, ex Benth. in DC. Prod. vii. 677 = gracilis.
tenera, *Steud. Nom.* ed. II. i. 580.—Hab. ?
tenuicaulis, *Klotzsch, ex Benth. in DC. Prod.* vii. 669. —Afr. austr.
tenuiflora, *Andr. Heathery*, iii. t. 146.—Afr. austr.
tenuifolia, Hort. ex Salisb. in Trans. Linn. Soc. vi. (1802) 231 = campanulata.
tenuifolia, *Linn. Syst.* ed. X. 1092.—Afr. austr.
tenuis, *Mœnch, Meth. Suppl.* 17.—Hab. ?
tenuis, *Salisb. in Trans. Linn. Soc.* vi. (1802) 229.—Afr. austr.
tenuissima, Wendl. Eric. i. fasc. VI. 9 = gracilis.
teretiscula, *Wendl. l. c.* ii. fasc. XXI. 141.—Afr. austr.
terminalis, Klotzsch, in Linnæa, xii. (1838) 509 = curvifolia.

ERICA :—
terminalis, *Salisb. Prod.* 296.—Ins. Corsicus.
tetragona, *Linn f. Suppl.* 228 ; *Thunb. Diss. Eric.* 14. —Afr. austr.
Tetralix, *Linn. Sp. Pl.* 353.—Europ
tetraloba, Roxb. ex Salisb. in Trans. Linn. Soc. vi. (1802) 337 = genistaefolia.
tetrahecolica, *Benth. in DC. Prod.* vii. 683.—Afr. austr.
teucrifolia, *Spreng. Pugill* i. 31.—Afr. austr.
thelitrifora, Lodd. Bot. Cab. t. 1294 = staminea.
thimifolia, *Wendl. Bot. Beobacht.* 48.—Hab. ?
Thomsonii x , Lem. Jard. Fleur, iii. (1852) t. 259.
Thunbergii, *Linn. f. Suppl.* 220.—Afr. austr.
thymifolia, *Andr. Col. Heaths*, ii 186.—Afr. austr.
thymoides, Klotzsch, ex Benth. in DC. Prod. vii. 670 = flacca.
thyrsoidea, Tausch, in Flora, xviii. (1834) ii. 619 = curviostris.
tiaraeflora, *Andr. Col. Heaths*, iii. 213.—Afr. austr.
togata, Sims, Bot. Mag. t. 1636 = borbonaefolia.
togatoides, Forbes, Hort. Woburn. 89 = bracteata.
tomentosa, *Salisb. in Trans. Linn. Soc.* vi. (1802) 327.—Afr. austr.
torosa, *Mœnch, Meth. Suppl.* 18.—Hab. ?
tortuosa, *Ludd. ex G. Don, in Loud. Hort. Brit.* 158, nomen.—Hab. ?
totta, Barl. in Linnaea, vii. (1832) 647 = Eremia Bartlingiana.
totta, Thunb. Diss. Eric. 18 = Eremia totta.
tragulifera, Salisb. in Trans. Linn. Soc. vi. (1802) 374 = nitida.
translucens, Andr. Heathery, t. 295 = Massoni ?
translucens, *Wendl. ex Spreng. Syst.* ii. 186, in syn ; *Tausch, in Flora*, xvii. (1834) ii. 598.—Afr. austr. ?
transparens, *Andr. Col. Heaths*, iv. 283.—Afr. austr.
transparens, Berg. Desc. Pl. Cap. 108 = comosa.
transparens, Thunb. Prod. Pl. Cap. 71 = elongata.
transpicuaca, Willd. ex Klotzsch, in Linnaea, xii. (1838) 217 = Bruckenthalia spiculifolia.
trassula, Klotzsch. l. c. 515, vide *troesula*.
tricepa, *Link, Enum. Hort. Berol.* i. 371.—Afr. austr.
trichophora, *Benth. in DC. Prod.* vii. 673.—Afr. austr.
trichophylla, *Benth. l. c.* 679.—Afr. austr.
trichroma, *Benth. l. c.* 642.—Afr. austr.
tricolor, Niven, ex Benth. in Trans. = trichroma.
tricolor, Spreng. Syst. ii. 193 = cumulifiora.
trifaria, Klotzsch, ex Benth. in DC. Prod. vii. 620 = lubricata.
trifiora, *Linn. Sp. Pl.* 354.—Afr. austr.
triphylla, *Link, Enum. Hort. Berol.* i. 360.—Afr. austr.
tristis, *Bartl. in Linnaea*, vii. (1832) 643.—Afr. austr.
tristis, Klotzsch, l. c. xii. (1838) 519 = Savilea.
triumphans, Lodd. Bot. Cab. t. 267 = andromedaeflora.
trivialis, Klotzsch, ex Benth. in DC. Prod. vii. 666 = decipiens.
trossula, Lodd. Bot. Cab. t. 668 = F. Muscari.
tuberculeris, Salisb. in Trans. Linn. Soc. vi. (1802) 330 = Eremia tuberculaeria.
tubiflora, Linn. Sp. Pl. ed. II. 505 = ignescens.
tubiflora, Thunb. Diss. Eric. 25 = bucciniaeformis.
tubiflora, *Willd. Sp. Pl. ii.* 408.—Afr. austr.
tubiuscula, Bedf. Hort. Eric. Woburn. 26 ; Lodd. Bot. Cab. t. 1157 = parviflora.
tubulosa, Sm. ex Steud. Nom. ed. I. 310 = tubiflora.
tubulosa, Wendl. Eric. ii. fasc. xvi. 57, t. 22 = Massoni.
tumida, *Kew-Gemal. in Bot. Reg.* t. 65.—Afr. austr.
tunicata, Bartl. in Linnaea, vii. (1832) 636 = cortifolia.
turbinata, *Andr. Heathery*, vi. 297.—Afr. austr.
turbiniflora, *Salisb. in Trans. Linn. Soc.* vi. (1802) 377.—Afr. austr.
turgida, Link, Enum. Hort. Berol. i. 365 = taxifolia.
turgida, *Salisb. in Trans. Linn. Soc.* vi. (1802) 345.—Afr. austr.
turmalis, *Salisb. l. c.* 342.—Afr. austr.
turrigera, *Salisb. l. c.* 377.—Hab. ?
Uhria, *Andr. Col. Heaths*, ii. 141.—Afr. austr.
umbellata, Aspo, Syn. Arag. 49 = australis.
umbellata, Barti. in Linnaea, vii. (1832) 647 = deliciosa.
umbellata, *Linn. Sp. Pl.* 352.—Reg. Mediterr. occ.
umbellifera, Loisel. in Desv. Journ. Bot. ii. (1809) 287 = multiflora.
umbellifera, Wendl. Eric. iii. fasc. xxv. 13. t. 6 = tenella.

ERICA :—
umbelliflora, *Klotzsch, ex Benth. in DC. Prod.* vii. 659.—Afr. austr.
umbelliformis, C. Koch, Dendrol. ii. ii. 277. Ind. sphalm. = multiflora.
uncifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 369 = regerminans.
undulata, Andr. Col. Heaths, iv. t. 288 ; Lodd. Bot. Cab. t. 1792 = Massoni ?
unibracteata, *Klotzsch, ex Benth. in DC. Prod.* vii. 692.—Afr. austr.
unica, Spreng. Syst. ii. 188 = pilulifera.
unicolor, Wendl. Eric. iii. fasc. xxv. 7. t. 3 = virescens.
uniflora, *Burm. f. Fl. Cap. Prod.* 11.—Afr. austr.
unilateralis, *Klotzsch, ex Benth. in DC. Prod.* vii. 667. —Afr. austr.
urceolaris, *Berg. Desc. Pl. Cap.* 107.—Afr. austr.
urceolaris, Salisb. in Trans. Linn. Soc. vi. (1802) 326 = hirta.
urceolaris, Sibth. & Sm. ex Steud. Nom. ed. II. i. 581 = vagans.
urnina, *Lav. ex Sweet, Hort. Brit.* ed. I. 268, nomen.—Hab. ?
vaga, Benth. in DC. Prod. vii. 667 = verticillata.
vaga, Salisb. Prod. 294 = vagans.
vagans, Desf. Fl. Atlant. i. 299 ; DC. Fl. Fr. iii. 679 = multiflora.
vagans, Koch, Syn. Fl. Germ. ed. II. 548 = verticillata.
vagans, *Linn. Mant.* ii. 230.—Reg. Mediterr.
Vanheurckii, *Muell. Arg. in Heurck, Obs. Bot.* 36.—Afr. austr.
varia, *Lodd. Bot. Cab.* t. 1325.—Afr. austr.
variabilis, *Klotzsch, ex Benth. in DC. Prod.* vii. 690.—Afr. austr.
varicifolia, Salisb. Prod. 298 = Monsoniana.
velitaria, *Salisb. in Trans. Linn. Soc.* vi. (1802) 367. —Afr. austr.
vellerifiora, Salisb. l. c. 333 = bruniades.
velutina, Bartl. in Linnaea, vii. (1832) 645 = tomentosa.
venosa, *Gentils. & Carr. in Rev. Hortic.* (1882) 406.—Hab. ?
ventricosa, *Thunb. Diss. Eric.* 27.—Afr. austr.
ventrosa, *Sweet, Hort. Brit.* ed. II. 341.—Afr. austr. ?
venusta, *[Forbes] Hort. Eric. Woburn.* 27 ; *Benth. in DC. Prod.* vii. 647.—Afr. austr.
venusta, Hort. ex Benth. in DC. Prod. vii. 647 = denticulata.
venusta, Klotzsch, in Linnaea, ix. (1834) 695 = elongata.
venusta, Salisb. Prod. 297 = ventricosa.
verecunda, Lodd. Bot. Cab. t. 1827 = Meuroni.
verecunda, *Salisb. in Trans. Linn. Soc.* vi. (1802) 379. —Afr. austr.
vernalis, Lodd. Bot. Cab. t. 1608 = pyramidalis.
verticillus, *Salisb. in Trans. Linn. Soc.* vi. (1802) 335. —Afr. austr.
Vernix, *Andr. Col. Heaths*, iii. t. 214.—Afr. austr.
Vernoni, *Gentils. & Carr. in Rev. Hortic.* (1882) 408. —Hab. ?
versicolor, *Andr. Col. Heaths*, i. 67.—Afr. austr.
verticillaris, Salisb. in Trans. Linn. Soc. vi. (1802) 360 = capitata.
verticillata, Andr. Col. Heaths, i. 66 = mammosa.
verticillata, Berg. Desc. Pl. Cap. 99 = concinna.
verticillata, *Forsk. Fl. Aegypt. Arab.* 210.—Reg. Mediterr. or.
vesicaria, Soland. ex Salisb. in Trans. Linn. Soc. vi. (1802) 330 = glomiflora.
vesicularis, *Salisb. l. c* 335.—Hab. ?
vespertina, *Linn. f. Suppl.* 221.—Afr. austr.
vestiflua, *Salisb. in Trans. Linn. Soc.* vi. (1802) 346.—Afr. austr.
vestita, Trége, ex Benth. in DC. Prod. vii. 625 = longifolia.
vestita, *Thunb. Diss. Eric.* 22.—Afr. austr.
vestitoides, *Regel, Ind. Sem. Hort. Petrop.* (1856) 30. —Hab. ?
villosa, *Andr. Col. Heaths*, iii. 216.—Afr. austr.
villosa, Wendl. Eric. ii. fasc. xvi. 53. t. 21 = bruniades.
villosiuscula, Lodd. Bot. Cab. t. 1844 = pusilla.
viminalis, Salisb. Prod. 298 = albens.
violacea, Klotzsch, ex Benth. in DC. Prod. vii. 620 = lubricata.
virescens, *Thunb. Diss. Eric.* 27 = pelviformis.
virgata, *Berg. Desc. Pl. Cap.* 97.—Afr. austr.

ERICA :—
virgata, Thunb. Diss. Eric. 18 = bispidula.
virgata, *Wendl. ex Spreng. Syst.* ii. 197.—Afr. austr.
virginalis, *Klotzsch, ex Benth. in DC. Prod.* vii. 658.—Afr. austr.
virgularis, Salisb. in Trans. Linn. Soc. vi. (1802) 324 = bispidula.
virgulata, Wendl. Eric. ii. fasc. xxi. 181. t. 49 = scoparia.
viridescens, Lodd. Bot. Cab. t. 238 = virescens.
viridiflora, *Andr. Col. Heaths*, iv. 287.—Afr. austr.
viridipurpurea, Gouan, Hort. Reg. Monsp. 195 = cinerea.
viridipurpurea, Linn. Sp. Pl. 353 = pelviformis.
viridis, Andr. Col. Heaths, ii. 140 = E. Lesa.
viscaria, Alt. Exot. Pl. t. 1 = decora.
viscaria, Bory, ex Benth. in DC. Prod. vii. 696 = Philippia viscosa.
viscaria, *Linn. Mant.* ii. 231.—Afr. austr.
viscida, Salisb. in Trans. Linn. Soc. vi. (1802) 373 = viscaria.
viscida, Wendl. Eric. ii. fasc. xiii. 7 = onosmaeflora.
viscosa, Steud. Nom. ed. I. 311, sphalm. = prsec.
volutaeflora, Salisb. in Trans. Linn. Soc. vi. (1802) 335 = nigrita.
vulgaris, Linn. Sp. Pl. 352 = Calluna vulgaris.
Walkeria, *Andr. Col. Heaths*, i. 72.—Afr. austr.
Walkeriana, Sweet, Hort. Brit. ed. II. 341 = prsec.
Watsoni x, Benth. in DC. Prod. vii. 665.—Anglia.
Wendlandiana, *Klotzsch, in Linnaea*, ix. (1834) 652. —Afr. austr.
westphalingia, Hort. ex Benth. in DC. Prod. vii. 689 = Massoni ?
Willmorei x, Knowles & Westc. Fl. Cab. ii. (1858) 115.
xeranthemifolia, Salisb. in Trans. Linn. Soc. vi. (1802) 339 = Eremia lanata.
Zeyheri, Bartl. in Linnaea, vii. (1832) 635 = affinis.
Zeyheri, Spreng. f. Tent. Suppl. Syst. 12.—Afr. austr.

ERICALA, Renealm. ex S. F. Gray, Nat. Arr. Brit Pl. ii. 336 (1821) = Gentiana, Tourn.
acuta, G. Don, Gen. Syst. iv. 190 = G. Amarella.
aestiva, G. Don, l. c. = G. verna.
alpina, G. Don, l. c. 189 = G. acaulis.
altaica, G. Don, l. c. = G. altaica.
angulosa, G. Don, l. c. 191 = G. verna.
angustifolia, G. Don, l. c. 190 = G. angustifolia.
aquatica, G. Don, l. c. 192 = G. aquatica.
argentea, G. Don, l. c. = G. argentea.
bavarica, S. F. Gray, Nat. Arr. Brit. Pl. ii. 336 = G. bavarica.
canaliculata, Steud. Nom. ed. II. i. 582 = Jaeschkea latiepala.
capitata, D. Don, ex G. Don, Gen. Syst. iv. 198 = G. capitata.
carinata, D. Don, ex G. Don, l. c. 189 = G. carinata.
carpathica, G. Don, l. c. 190 = G. nivalis.
coronata, D. Don, ex G. Don, l. c. 198 = G. coronata.
cruciata, G. Don, l. c. 191 = G. cruciata.
decemfida, D. Don, in Trans. Linn. Soc. xvii. (1837) 514 = G. decemfida.
depressa, D. Don, ex G. Don, Gen. Syst. iv. 189 = G. depressa.
Douglasiana, G. Don, l. c. = G. Douglasiana.
flava, G. Don, l. c. = G. Amarella.
humilis, G. Don, l. c. 198 = G. humilis.
imbricata, G. Don, l. c. 190 = G. imbricata.
linearis, G. Don, l. c. 190 = G. linearis.
Lourseri, G. Don, l. c. 192 = G. indica.
marginata, D. Don, ex G. Don, l. c. 192 = G. carinata.
nivalis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 336 = G nivalis.
nutans, G. Don, Gen. Syst. iv. 198 = G. prostrata.
orbiculata, G. Don, l. c. 193 = G. quadrifaria.
pedicellata, D. Don, in Trans. Linn. Soc. xvii. (1837) 514 = G. quadrifaria.
procumbens, G. Don, Gen. Syst. iv. 192 = G. quadrifaria.
prostrata, G. Don, l. c. 190 = G. prostrata.
pumila, G. Don, l. c. 189 = G. pumila.
pyrenaica, G. Don, l. c. = G. pyrenaica.
quadrifaria, G. Don, l. c. 190 = G. quadrifaria.
Ruploana, G. Don, l. c. 191 = G. decemfida.
salifolia, G. Don, l. c. 189 = G. saxifraga.
thalassea, G. Don, l. c. 191 = G. spathacea.
squarrosa, G. Don, l. c = G. squarrosa.
Thunbergii, G. Don, l. c. 198 = G. Thunbergii.
tubiflora, G. Don, l. c. 189 = G. tubiflora.

ERICALA :—
utriculosa, G. Don, Gen. Syst. iv. 191 = Gentiana utriculosa.
venusta, G. Don, l. c. 189 = G. venusta.
verna, S. F. Gray, Nat. Arr. Br. Pl. ii. 335 = G. verna.

ERICAMERIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 318 = **Aplopappus**, Cass. (Compos.).
diffusum, Benth. Bot. Voy. Sulph. 23 = Bigelovia diffusa.
erecta, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 226 = A. Bloomeri.
microphylla, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 319 = A. ericoides.
nana, Nutt. l. c. = A. nanus.
resinosa, Nutt. l. c. = A. nanus.

ERICAULON, Lour. Fl. Cochinch. i. 60 (1790) sphalm. = **Eriocaulon**, Linn.

ERICILLA, Steud. Nom. ed. II. i. 582 (1840) = **Brotlla**, Juss. (Phytolac.).

ERICINELLA, Klotzsch, in Linnæa, xii. (1838) 222.
ERICACEAE, Benth. & Hook. f. ii. 591.
gracilis, *Benth. in DC. Prod.* vii. 697.—Afr. austr.
Mannii, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 16.—Afr. trop.
multiflora, *Klotzsch, in Linnæa*, xii. (1838) 228.—Afr. austr.
passerinoides, *Bolus, in Journ. Linn. Soc.* xviii. (1881) 393.—Afr. austr.

ERICOILA, Renealm. ex Borckh. in Roem. Arch. i. i. 23 (1796) = **Gentiana**, Tourn.
aestiva, Bercht. & Presl, Rostl. i. Gentian. 15 = G. verna.
altaica, Borckh. in Roem. Arch. i. i. 27 = G. altaica.
anguiosa, Bercht. & Presl, Rostl. i. Gentian. 15 = G. verna.
aquatica, Borckh. in Roem. Arch. i. i. 27 = C. aquatica.
aurea, Bercht. & Presl, Rostl. i. Gentian. 14 = G. aurea.
bavarica, Borckh. in Roem. Arch. i. i. 27 = G. bavarica.
Chirayita, Bercht. & Presl, Rostl. i. Gentian. 14 = Swertia Chirata.
cruciata, Borckh. in Roem. Arch. i. i. 27 = G. cruciata.
humilis, Bercht. & Presl, Rostl. i. Gentian. 16 = G. humilis.
imbricata, Bercht. & Presl, l. c. 15 = G. imbricata.
Tirashii, Borckh. in Roem. Arch. i. i. 27 = G. verna.
macrophylla, Bercht. & Presl, Rostl. i. Gentian. 14 = G. macrophylla.
nivalis, Borckh. in Roem. Arch. i. i. 27 = G. nivalis.
prostrata, Borckh. l. c. = G. prostrata.
pumila, Borckh. l. c. = G. pumila.
pyrenaica, Borckh. l. c. = G. pyrenaica.
quinquefiora, Bercht. & Presl, Rostl. i. Gentian. 14 = G. quinquefiora.
umbellata, Bercht. & Presl, l. c. = G. aurea.
utriculosa, Borckh. in Roem. Arch. i. i. 27 = G. utriculosa.
verna, Borckh. l. c. = G. verna.
verticillata, Borckh. l. c. = Enicostema littorale.

ERICOMA, Vasey, in U. St. Dep. Agr. Rep. (1881–82) 247 = ERIOCOMA, Nutt. = **Orysopsis**, Michx. (Gramin.).

ERICOMYRTUS, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 154 = **Baeckea**, Linn. (Myrtac.).
Drummondii, Turcz. l. c. 155 = B. pulchella.

ERIGENIA, Nutt. Gen. Am. i. 187 (1818). *UMBELLIFERAE*, Benth. & Hook. f. i. 886.
ERIGONIA, A. Juss. in Orb. Dict. ix. 95 (1849).
bulbosa, *Nutt. l. c.*—Am. bor.

ERIGERON, Linn. Hort. Cliff. 407 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 279.
ASTRADELPHUS, Remy, in Ann. Sc. Nat. Sér. III. xii. (1849) 186.
CARDOTUS, Rafin. Fl. Tellur. ii. 50 (1836).
CONYZELLA, Regel, in Mém. Acad. Pétersb. Sér. VII. xiv. (1869) n. 4 (Sert. Tianch. 51).

ERIGERON :—
DIPLEMIUM, Rafin. Fl. Tellur. ii. 50 (1836).
GUSMANIA, Remy, in C. Gay, Fl. Chil. iv. 12 (1849).
HETEROCHAETA, DC. Prod. v. 282 (1836).
LEPTILON, Rafin. in Am. Monthly Mag. (1818) 268.
LEPTOSTELMA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. i. 35 (1830).
PANTOL, Adans. Fam. ii. 124 (1763).
PAPPOCHROMA, Rafin. Fl. Tellur. ii. 48 (1836).
PHALACROLOMA, Cass. in Dict. Sc. Nat. xxxix. 404 (1826).
POLYACTIDIUM, DC. Prod. v. 281 (1836).
POLYACTIS, Less. Syn. Comp. 188 (1832).
STENACTIS, Cass. in Dict. Sc. Nat. xxxvii. 462 (1825); l. 488 (1827).
TANAXION, Rafin. Fl. Tellur. ii. 51 (1836).
TERRANEA, Colla, in Mem. Acc. Torin. xxxviii. (1835) 11. t. 28.
TESSENIA, Bubani, in Nouv. Giorn. Bot. It. v. (1873) 318.
TRIMORPHA, Cass. in Dict. Sc. Nat. iii. Suppl. 65 (1816).
WOODVILLEA, DC. Prod. v. 318 (1836).
abyssinicus, Sch. Bip. ex Schweinf. & Aschers. in Beitr. Fl. Aethiop 284 = Conyza abyssinica.
acris, C. B. Clarke, Comp. *Ind.* 52 = alpinus.
acris, *Linn. Sp. Pl.* 863.—Reg. bor. temp.
adscendens, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 83.—Am. bor.
aegyptiacus, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 198 = asteroides.
aegyptius, *Linn. Mant.* i. 112.—Afr. bor.
affinis, *DC. Prod.* v. 289.—Mexic.
alatus, D. Don, Prod. Fl. Nep. 171 = Laggera alata.
albidus, A. Gray, in Proc. Am. Acad. v. (1862) 319 = bonariensis.
alpestris, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 123 = maximus.
alpestris, Hoppe, ex Nym. Consp. 388 = alpinus.
alpinus, Lam. Fl. Fr. ii. 140 = acris.
alpinus, Ledeb. Fl. Alt. iv. 90 = glabratus.
alpinus, Less. in Linnæa, vi. (1831) 128 = pulchellus.
alpinus, *Linn. Sp. Pl.* 864.—Reg. bor. et arct.
alpinus, Purah, Fl. Am. Sept. ii. 532 = uniflorus.
ambiguus, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859).—Austral.
ambiguus, Nutt. Gen. Am. ii. 147 = strigosus.
ambiguus, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 208 = linifolius.
amorphoglossus, *Boiss. Diagn.* Ser. I. vi. 80.—Persia.
amphibolus, Ledeb. Fl. Ross. iii. 686 = pulchellus.
amplexicaulis, *Hort. ex Poir. Encyc. Suppl.* v. 464; *Desf. Cat. Hort. Par.* ed. III. 401.—Am. bor.
amplexicaulis, Torr. ex Torr. & Gray, Fl. N. Am. ii. 171 = philadelphicus.
Anderssonii, A. Gray, in Proc. Am. Acad. vi. (1864–65) 540 = Aster Andersoni.
andicola, *DC. Prod.* v. 287.—Chili.
andinus, *Phil. in Linnæa*, xxviii. (1856) 722.—Chili.
andryaloides, *Benth. ex C. B. Clarke, Comp. Ind.* 52. —Reg. Himal.
angulosus, Gaudin, Fl. Helv. v. 265 = acris.
angustatus, Fries, ex Nym. Consp. 389 = acris.
angustatus, Greene, in Bull. Calif. Acad. i. n. 111. (1885) 88 = inornatus.
angustissimus, *Lindl. ex DC. Prod.* v. 286.—Ind. or.
annuus, *Pers. Syn.* ii. 431.—Am. bor.
anomalus, DC. Prod. v. 298 = Brachyactis menthodora.
aperuinianus, Georgi, Reise, i. 250; ex Ledeb. Fl. Ross. ii. 490 = uniflorus.
apiculatus, *Benth. Pl. Hartw.* 204.—N. Granat.
apurensis, *Griseb. Fl. Brit. W. Ind.* 265.—Ind. occ.
argentatus, *A. Gray, in Proc. Am. Acad.* viii. (1872) 649.—Am. bor. occ.
arisonicus, *A. Gray, l. c.* xix. (1883) 2.—Arizona.
Armenus, Boiss. & Huet, ex Boiss. Fl. Orient. iii. 164 = pulchellus.
armerifolius, *Turcz. ex DC. Prod.* v. 291.—Sibir.; Am. bor. occ.
artemisiaefolius, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 81.—Bolivia.
asper, Nutt. Gen. Am. ii. 147 = glabellus.
aspergillatus, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 91.—Am. bor. occ.

ERIGERON :—
asperulus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 13, nomen.—Chili?
asteroides, Andrz. ex Bess. Enum. Pl. Volh. 33 = acris.
asteroides, Hoppe, ex Nym. Consp. 388 = alpinus.
asteroides, *Hort. Angl. ex Link, Enum. Hort. Berol.* ii. 324.—Hab.?
asteroides, *Roxb. Hort. Beng.* 61; *Fl. Ind.* iii. 432.—Ind. or.
asteroides, Wall. Cat. n. 2975 B = Blumea amplectens.
atticus, Vill. Hist. Pl. Dauph. iii. 237 = Villarsii.
Aucheri, *Boiss. Fl. Orient.* iii. 170.—Syria; Persia.
auranticus, *Regel, Gartenfl.* (1879) 289; *et in Act. Hort. Petrop.* vi. (1879) 305.—Turkestan.
australis, Hornem. ex Spreng. Syst. iii. 517 = strigosus.
australis, *Phil. in Anal. Univ. Chil.* (1873) 488.—Chili.
bahamensis, *Scop. Del. Insub.* iii. 14.—Ins. Baham.
bellidiastroides, *Grisch. Cat. Pl. Cub.* 149.—Cuba.
Bellidiastrum, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 307.—Am. bor.
bellidifolius, *Muhl. ex Willd. Sp. Pl.* iii. 1958.—Am. bor.
bellidioides, *Benth. & Hook. f. Gen.* ii 281.—Reg. Himal.
bellidioides, Spenn. Fl. Friburg. ii. 586 = annuus.
bellioides, *DC. Prod.* v. 288.—Porto Rico.
Berterianus, *DC. l. c.* 286.—Chili.
Beyrichii, Hort. Berol. ex Torr. & Gray, Fl. N. Am. ii. 176 = strigosus.
biflorus, *Phil. in Anal. Univ. Chil.* (1873) 488.—Chili.
Bigelovii, *A. Gray, in Torr. Bot. Mex. Bound.* 78.—Am. bor. occ.
Blanchetii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 111. 36 = Conyza Blanchetii.
Bloomeri, *A. Gray, in Proc. Am. Acad.* vi. (1865) 540. —Calif.
bonariensis, Hort. ex Link, Enum. Hort. Berol. ii. 323 = linifolius.
bonariensis, *Linn. Sp. Pl.* 863.—Am. austr.
Bovei, *Boiss. Fl. Orient.* iii. 168.—Aegypt.; Arab.
Brandegei, *A. Gray, Syn. Fl. N. Am.* i. ii. 210.—Am. bor. occ.
brasoensis, Buckl. in Proc. Acad. Sc. Philad. 1861 (1863) 456 = tenuis.
Brewer, *A. Gray, in Proc. Am. Acad.* vi. (1866) 541. —Calif.
Bungei, *Boiss. Fl. Orient.* iii. 164.—Afghan.
caespitosus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 307.—Am. bor. occ.
camphoratus, Linn. Sp. Pl. 864 = Pluchea camphorata.
camporum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. iii. 34 = Aster camporum.
canadensis, Brot. Fl. Lusit. i. 359 = rufescens.
canadensis, *Linn. Sp. Pl.* 863.—Amphig.
canadensis, Tenore, Syll. Fl. Neap. 428 = linifolius.
Candollei, F. Muell. ex Benth. Fl. Austral. iii. 499 = Minuria denticulata, integerrima.
canescens, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 50 = stenophyllus.
canescens, Parry, in Jones, Exped. n. 139; ex A. Gray, Syn. Fl. N. Am. i. ii. 213 = ochroleucus.
canescens, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 534, nomen.—Bolivia.
canescens, Torr. & Gray, Fl. N. Am. ii. 179 = caespitosus.
canescens, *Willd. Sp. Pl.* iii. 1953.—Hab.?
canus, *A. Gray, Pl. Fendl.* 87.—N. Mexic.
capensis, Houtt. Handleid. x. 629 = Vernonia pinifolia.
cardaminifolius, Wedd. Chlor. And. i. 198, in obs. = Conyza cardaminifolia.
carolinensis, Cranta. Inst. i. 311 = seq.
caroliniauus, Linn. Sp. Pl. 863 = Solidago tenuifolia.
carpaticus, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 336 = Villarsii.
caucasicus, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 63 = pulchellus.
chilensis, D. Don, ex G. Don, in Loud. Hort. Brit. 343 = Conyza chilensis.
chinensis, Jacq. Hort. Schoenb. iii. 30. t. 303 = apurensis.
chinensis, Siebold, ex DC. Prod. v. 449 = Laggera aurita.
chionophilus, *Boiss. Fl. Orient.* iii. 168.—Persia.
chionophilus, *Wedd. Chlor. And.* i. 191.—N. Granat.

ERIGERON:—
Chrysocoma, Walp. Rep. iv. 396, sphalm. = Chrysocoma Coma-aurea.
Chrysopsidis, *A. Gray, Syn. Fl. N. Am.* i. ii. 210.—Am. bor. occ.
ciliaris, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 170.—Japon.
ciliaris, *Phil. in Linnaea*, xxviii. (1856) 722.—Chili.
ciliatus, Ledeb. Fl. Alt. iv. 92 = Aster angustus.
ciliatus, Boiss. ex Tchihat. Asie Min. Bot. ii. 235 = uniflorus.
ciliatus, Boiss. Fl. Orient. iii. 165, in syn. = seq.
cinerascens, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Chili.
cinereus, A. Gray, Pl. Fendl. 68 = divergens.
cinereus, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 90.—Patagon.
clematideus, *Sch. Bip. ex Walp. Rep.* ii. 968 = Nidorella mespilifolia.
cochinchinensis, *Lour. Fl Cochinch.* 500, in obs.—Cochinch.
cochinchinensis, Spreng. ex DC. Prod. v. 446 = Blumea aromatica.
collinus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 536, nomen.—Bolivia.
Coma-aurea, Sch. Bip. in Flora, xxvii. (1844) 670 = Chrysocoma Coma-aurea.
compositus, *Pursh, Fl. Am. Sept.* ii. 535.—Am. bor.; Reg. arct.
conchophyllum, Walp. Rep. ii. 583, sphalm. = armeriifolius.
concinnus, *Torr. & Gray, Fl. N. Am.* ii. 174.—Am. bor. occ.
contortus, Desf. Tabl. ed. I. 102, nomen; Pers. Syn. ii. 432 = linifolius.
conyzoides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 106.—Austral.
coronopifolius, Clem. ex Willk. & Lange, Prod. Fl. Hisp. ii. 24 = linifolius.
coronopifolius, *Schrank, in Denkschr. Bot. Ges Regensb.* ii. (1822) 69.—Afr. austr.
corymbosus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 308.—Am. bor. occ.
corymbosus, Wallr. Beitr. Fl. Hercyn. 272; et in Linnæa, xiv. (1840) 642 = acris.
Coulteri, *Porter & Coulter, Fl. Colorad.* 61.—Am. bor. occ.
Cuxi, *Phil. in Linnaea*, xxxiii. (1864–65) 134.—Chili.
crassicaulis, *Sch. Bip. l. c.* xxxiv. (1865–66) 584, nomen.—Bolivia.
crispus, Pourr. in Mém. Acad. Toul. iii. (1788) 318 = linifolius.
crocifolius, Wedd. Chlor. And. i. 191 = Aster crocifolius.
cuneifolius, DC. Prod. v. 288 = Aster jamaicensis.
cuspidatus, Spreng. ex DC. l. c. 444 = Blumea oxyodonta.
cymosus, *Walt. Fl. Carol.* 206.—Carolina.
Darrellianus, *Hemsl. in Journ. Bot.* xxi. (1883) 104.—Ins. Bermuda.
decumbens, Benth. Pl. Hartw. 316 = foliosus.
decumbens, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 309.—Am. bor. occ.
decurrens, DC. Prod. v. 298 = Pterigeron decurrens.
decurrens, Vahl, Symb. Bot. i. 72 = Conyza arabica.
delphinifolius, A. Gray, Pl. Wright. ii. 77 = neomexicanus.
delphinifolius, *Willd. Enum. Hort. Berol.* 873 — Mexic.
dentatus, Burm. f. Fl. Cap. Prod. 26 = Conyza scabrida?
denticulatus, *Burm. f. Fl. Ind.* 180.—Ind. or.
dianthifolius, *Griseb. in Goett. Abh.* xxiv. (1879) 174.—Keg. Argent.
diaphanus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 594, nomen.—Bolivia.
diffusus, Pers. Syn. ii. 431 = Leucopsis diffusa.
diplopappoides, *Schau. in Linnaea*, xix. (1847) 732.—Mexic.
discoideus, Kellogg, in Proc. Calif. Acad. v. (1873) 55 = Conyza Coulteri.
divaricatus, *Michx. Fl. Bor. Am.* ii. 128.—Am. bor.
diversicostus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 311 = divergens.
divergens, Hook. in Lond. Journ. Bot. vi (1847) 342 = flagellaris.
divergens, *Torr. & Gray, Fl. N. Am.* ii. 175.—Am. bor. occ. et trop.

ERIGERON:—
diversifolius, Cav. in Anal. Clenc. Nat. iv. (1801) 88 = Conyza Gouani.
diversifolius, Rich. ex Reichb. Fl. Germ. Excurs. 389 = annuus.
Dougiasii, Torr. & Gray, Fl. N. Am. ii. 177 = foliosus.
drobachensis, O. Muell. in Fl. Dan. t. 874 = acris.
dryophyllus, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 210.—Mexic.
dubius, Spreng. Syst. iii. 519 = Hysterionica montevidensis.
Eatoni, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 91.—Am. bor. occ.
schioides, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop 146 = Conyza tigrensis.
elbursensis, *Boiss. Diagn.* Ser. I. xi. 1.—Persia.
Elliati, *Hook. fc. Fl.* t. 147.—Reg. Himal.
elongatus, Ledeb. Fl. Alt. iv. 91 = acris.
elongatus, Mœnch, Meth. Suppl. 247 = Villarsii.
engadinensis ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 105.
eriocephalus, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 613.—Turkestan.
eriocephalus, J. Vahl, in Fl. Dan. t. 2299 = uniflorus.
eriophyllus, *A. Gray, Pl. Wright.* ii. 77.—Mexic.
Ervendbergii, *A. Gray, in Proc. Am. Acad.* viii. (1878) 550.—Mexic.
expansus, Poepp. ex Spreng. Syst. iii. 518 = Aster inconspicuus.
exstipulatus, Schum. & Thonn. Beskr. Guin. Pl. 367 = Conyza dentata.
fastidus, Burm. f. Fl. Cap. Prod. 26 = Nidorella foliosa.
falcatus, D. Don, Prod. Fl. Nep. 172 = Laggera flava.
fasciculatus, Colla, in Mem. Acc. Tor. xxxvii. (1885) 37.—Apiculosus.
fastigiatus, *Hornem. Hort. Hafn.* ii. 971.—Hab. ?
ferruginens, *Wedd. Chlor. And.* i. 195.—Bolivia.
filifolius, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 308.—Am. bor. occ.
filiformis, Spreng. Syst. iii. 520 = Hysterionica pinifolia.
flagellaris, Boiss. & Hausk. ex Boiss. Fl. Orient. iii. 172 = Lachnophyllum Noëanum.
flagellaris, *A. Gray, Pl. Fendl.* 68.—N. Mexic.
floribundus, Sch. Bip. in Bull. Soc. Bot. Fr. xii. (1865) 81; et in Linnæa, xxxiv. (1865–66) 584 = bonariensis.
florifer, Hook. Fl. Bor. Am. ii. 20 = Townsendia floriferi.
foetidus, Linn. Sp. Pl. ed. II. 1213 = Nidorella foliosa.
foliosus, Don, ex Steud. Nom. ed. II. i. 588, sphalm. = Conyza falcata.
foliosus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 309.—Am. bor. occ.
fragrans, *Spreng. Syst.* iii. 518.—Hab. ?
frigidus, Boiss. & Buhse, ex Boiss. Fl. Orient. iii. 164 = Leucheilus.
frigidus, *Boiss. in DC. Prod.* vii. 274.—Hispan.
frigidus, *Wedd. Chlor. And.* i. 231.—Bolivia.
fruticosus, *Phil. in Linnaea*, xxxiii. (1864–65) 134.—Chili.
fruticosus, *DC. Prod.* v. 283.—Ins. J. Fernand.
Funkii, Sch. Bip. ex Nym. Consp. 388 = alpinus.
Gaudichaudii, *DC. Prod.* vii. 274.—Peruv.
Gayanus, *Remy, in C. Gay, Fl. Chil.* iv. 26.—Chili.
glabellus, *Nutt. Gen. Am.* ii. 147.—Am. bor.
glaber, Hoppe, ex Nym. Consp. 388 = alpinus.
glaberrimus, Schrank, ex Nym. l. c. 389 = acris.
glabratus, Endr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 23 = alpinus.
glabratus, Hook. Fl. Bor. Am. ii. 18 = acris.
glabratus, *Hoppe & Hornsch, ex Bluff & Pingerh. Comp. Fl. Germ.* ii. 364.—Helvet.
glabrifolius, *DC. Prod.* vii. 287.—Chili.
glacialis, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxvii. (1865) i. 389 = uniflorus.
glandulosus, Poir. Encyc. viii. 487 = Chrysopsis graminifolia.
glandulosus, *Porter & Coulter, Fl. Colorad.* 60.—Am. bor. occ.
glareosus ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 105.
glaucus, *Ker-Gawl. in Bot. Reg.* t. 10.—Am. bor. occ.
glomeratus, Salisb. Prod. 195 = Conyza Gouani.
glomeratus, Spreng. ex DC. Prod. v. 297 = Blumea Wightiana.

ERIGERON:—
glutinosus, Desf. Cat. Hort. Par. ed. III. 174 = Felicia glutinosa.
glutinosus, Linn. Sp. Pl. ed. II. 1212 = Jasonia glutinosa.
guaphalioides, *H. B. & K. Nov. Gen. et Sp.* iv. 88. t. 331.—Mexic.
Gouani, Jacq. Hort. Vindob. iii. 48. t. 79 = Conyza aegyptiaca.
Gouani, Linn. Mant. ii. 469 = Conyza Gouani.
graminifolius, Salisb. Reitr. ii. 240 = armeriifolius.
gramineus, Linn. Sp. Pl. 864 (= *Erigeron graminifolium*).—Sibir.
graminifolius, Pall. ex Herd. in Bull. Soc. Nat. Mosc. xxxvii. (1865) i. 388 (= *Arctogeron gramineum*).—Sibir.
graminifolius, *Phil. in Linnaea*, xxviii. (1856) 723.—Chili.
graminorus, Russ. ex DC. Prod. v. 444 = Blumea oxyodonta.
grandiflorus, *Hook. Fl. Bor. Am.* ii. 18. t. 128.—Am. bor. occ.
grandiflorus, Hoppe, ex DC. Prod. v. 291 = alpinus.
grandiflorus, Nutt. in Journ. Acad. Philad. vii. (1834) 81 = macranthus.
grandiflorus, Willd. ex Spreng. Syst. iii. 519 = pulchellus.
Grantii, *Oliver & Hiern, in Trans. Linn. Soc.* xxix. (1874) 93.—Afr. trop.
gravesiens, Linn. Cent. i. n. 80; Amoen. Acad. iv. 290 = Inula graveolens.
Gunnii, Hook. f. Fl. Tasman. i. 188 = pappochroma.
Hegetschweileri, *Bruegg. in Berlepsch, Schweiz*, 109.—Helvetia.
helvetiens ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 105.—Europ.
heterophyllus, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 11.—Mexic.
heterophyllus, Muhl. ex Willd. Sp. Pl. iii. 1956 = annuus.
hieraciifolius, D. Don, Prod. Fl. Nep. 172 = Blumea hieraciifolia.
hieraciifolius, Poir. Encyc. viii. 491 = Podocoma hieraciifolia.
hieracioides, *Wedd. Chlor. And.* i. 194.—Peruv.
hirsutus, Hoppe, ex DC. Prod. v. 291 = alpinus.
hirsutus, *Lour. Fl. Cochinch.* 500.—China.
hirsutus, Pursh, Fl. Am. Sept. ii. 742 = pumilus.
hirsutus, Wall. Cat. n. 3976 = asteroides.
hirtellus, *DC. Prod.* v. 290.—Chili.
hirtus, Thunb. Prod. Pl. Cap. 153 = Conyza incisa.
hispidus, *Baker, in Mart. Fl. Bras.* vi. iii. 29.—Bras.
hispidus, DC. in Wight, Contrib. 9 = asteroides.
hispidus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 310 = glaucus.
Hochstetteri, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 147 = Conyza Hochstetteri.
Howellii, *A. Gray, Syn. Fl. N. Am.* i. ii. 209.—Am. bor. occ.
Huelsenii, *Vatke, in Oestr. Bot. Zeitschr.* xxi. (1871) 346.—Europ.
humilis, *R. Grah. in Edinb. N. Phil. Journ.* (Oct.–Dec. 1828) 175.—Am. sept.
humilis, Spreng. ex DC. Prod. v. 287 = monticulus.
hyemalis, *Rafin. Fl. Ludov.* 65.—Am. bor.
hyoseroides, *Griseb. Cat. Pl. Cub.* 149.—Cuba.
hypoleucus, Sch. Bip. ex Schweinf. & Aschers. in Beitr. Fl. Aethiop. 384 = Conyza hypoleucus.
hypopolius, *Phil. in Anal. Univ. Chil.* (1878) 487.—Chili.
hymopifolius, *Michx. Fl. Bor. Am.* ii. 128.—Am. bor.
hyssopifolius, Sch. Bip. in Flora, xxvii. (1844) 670 = Nidorella foliosa.
incanus, Vahl, Symb. Bot. i. 73 = Conyza incana.
incisus, Sieber, ex DC. Prod. v. 281 = trilobus ?
incisus, Lamk. Prod. Fl. Cap. 256 = Conyza incisa.
incomptus, *A. Gray, Syn. Fl. N. Am.* i. ii. 218.—Calif.
indicus, *Burm. f. Fl. Ind.* 180.—Java.
inomatus, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 88.—Am. bor. occ.
integrifolius, Bertol. Misc. Bot. vi. t. 9 = uniflorus.
integrifolius, Bigel. Fl. Bost. ed. II. 302 = strigosus.
intermedius, Reichb. ex Nym. Consp. 388 = alpinus.
intermedius, Schleich. ex Nym. l. c. 389 = acris.
intermedius, Trachs. ex Dalla Torre, Art. Alpenfl. 135 = *Trachselii*.

ERIGERON:—
inuloides, Poir. Encyc. Suppl. v. 464 = Pulicaria inuloides.
involucratus, *Link, ex Sweet, Hort. Brit.* ed. II. 294, nomen.—Bras.
ivaefolius, Sch. Bip. in Flora, xxvii. (1844) 670 = Conyza ivaefolia.
jamaicensis, Linn. Syst. ed. X. 1213 = Aster jamaicensis.
japonicus, Thunb. Fl. Jap. 312 = Conyza japonica.
javanicus, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 120, nomen; *Miq. Fl. Ind. Bat.* ii. 30.—Malaya.
Jurzitzei, Spreng. Syst. iii. 519 = Conyza melastomoides.
kamtschaticus, DC. Prod. v. 290 = acris.
Karvinskianus, *DC. l. c.* 285.—Mexic.
khorsanicus, *Boiss. Fl. Orient.* iii. 170.—Persia.
Kotschyi, Boiss. Diagn. Ser. I. vi. 82 = Aucheri.
Kraussii, Sch. Bip. ex Walp. Rep. ii. 958 = Nidorella auriculata.
laevigatus, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 113 (err. typ. 105).—Guian.
lanatus, *Hook. Fl. Bor. Am.* ii. 17. t. 121.—Am. bor. occ.
lanceolatus, *Wedd. Chlor. And.* i. 198.—Bolivia.
Landbecki, *Phil. in Linnaea*, xxxiii. (1864–65) 188.—Chili.
lanciflius, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 308.—Ins. Galapag.
latisquamatus, Maxim. Prim. Fl. Amur. 478 = Aster angustus.
latuquamus, *Boiss. Fl. Orient.* iii. 167.—Persia.
laxiflorus, *Baker, in Mart. Fl. Bras.* vi. iii. 81.—Bras.
Lechleri, *Sch. Bip. in Flora*, xxxviii. (1855) 114.—Chili.
Lehmanni, *Boiss. Fl. Orient.* iii. 171.—Turkestan.
Lehmanni, *Spreng. Syst.* iii. 517.—Aegypt.
leiomerus, *A. Gray, Syn. Fl. N. Am* i. ii. 211.—Am. bor. occ.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 2.—Arizona.
lepidotus, Less. in Linnaea, vi. (1831) 502 (= Tetramolophi sp.).—Ins. Sandvic.
leptopetalus, *Phil. in Linnaea*, xxxiii. (1864–65) 136.—Chili.
leptorhizon, *DC. Prod.* v. 288.—Am. austr.
Leschenaultii, DC. l. c. 292 = alpinus.
leucanthemifolius, *Schau. in Linnaea*, xix. (1847) 728.—Mexic.
leucanthus, D. Don, Prod. Fl. Nep. 171 = Blumea leucanthema.
leucophyllus, *Boiss. Fl. Orient.* iii. 171.—Turkestan.
liatroides, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 172 = Pteirgeron liatroides.
limnophilus, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 81.—Bolivia.
linearifolius, Cav. in Anal. Cienc. Nat. iv. (1801) 89 = linifolius.
linearis, Spreng. ex DC. Prod. v. 274 = Vicoa auriculata.
linifolius, Bert. ex DC. Prod. v. 254 = Aster subulatus.
linifolius, *Willd. Sp. Pl.* iii. 1955.—Reg. calid.
Linosyris, [Clairv.] Man. Herb. 248 = Aster Linosyris.
lonchophyllus, Hook. Fl. Bor. Am. ii. 18 = armeriifolius.
longifolius, Baill. Hist. Pl. viii. 86 = chilensis?
longifolius, Brouss. ex Webb & Berth. Phyt. Canar. ii. 208, in obs. = Conyza Gouani.
longifolius, Desf. Tabl. ed. I. 102, nomen; Cat. Hort. Par. ed. III. 400 = Jasonia longifolia.
longipes, *DC. Prod.* v. 288.—Mexic.
lyratus, *Kuhn. Ati. Veurm.* 147.—Am. bor.
macranthus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 310.—Am. bor. occ.
macrophyllus, Herbich, Stirp. Rar. Bucov. 57 = glabratus.
macromischus, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 147, nomen.—Abyss.
Mandonii, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 81, nomen.—Bolivia.
mariticus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 310 = glaucus.
maximus, *Otto, ex DC. Prod.* v. 284.—Mexic.
mexifolius, Sch. Bip. ex Walp. Rep. ii. 958 = Nidorella mexifolia.
microcephalus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 584, nomen.—Bolivia.

ERIGERON:—
microspermus, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 120, nomen.—Java.
minuriloides, *Benth. Fl. Austral.* iii. 495.—Australi.
miser, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 379.—Calif.
mixtus x, Arvet.—Cf. Bull. Soc. Bot. Fr. xxvi. (1879) Bibl. 89.—Europ.
modestus, *A. Gray, Pl. Fendl.* 68.—Texas.
mollis, D. Don, Prod. Fl. Nep. 172 = Blumea Wightiana.
monocephalus, Schur, Enum. Pl. Transs. 309 = uniflorus.
Monorchis, *Griseb. in Goett. Abh.* xxiv. (1879) 175.—Bras.
montevidensis, *Baker, in Mart. Fl. Bras.* vi. iii. 80.—Uruguay.
montevidensis, Spreng. Syst. iii. 519 = Hysterionica montevidensis.
monticolus, *Wall. Cat.* n. 2977.—Reg. Himal.
mucronatus, *DC. Prod.* v. 285.—Mexic.
Muelleri, Lund, ex Nym. Consp. 389 = acris.
Muiriî, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 210.—Alaska.
multiradiatus, Wall. Cat. n. 3286 = alpinus.
multiflorus, Hook. & Arn. Bot. Beech. Voy. 87 = Aster exilis.
multiradiatus, *Benth. & Hook. f. Gen.* ii. 281.—Reg. Himal.
muralis, Lapeyr. Suppl. Hist. Pl. Pyr. 133 = acris.
Myosotus, *Pers. Syn.* ii. 431.—Reg. Magell.
Myosotis, *Remy, in C. Gay, Fl. Chil.* iv. 22.—Chili.
myosotoides, Juss. ex Pers. Syn. ii. 431 = E. Myosotis, *Pers.*
nanus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 308 = radicatus.
nanus, Sch. Bip. ex Schweinf. & Aschers. in Beitr. Fl. Aethiop. 284 = Conyza nana.
nanus, Schur, Enum. Pl. Transs. 309 = alpinus.
natalensis, Sch. Bip. ex Walp. Rep. ii. 958 = Nidorella mespilifolia.
neglectus, Kern. ex Nym. Consp. 388 = alpinus.
neo-mexicanus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 2.—N. Mexic.
nervosus, Willd. Sp. Pl. iii. 1953 = Chrysopsis graminifolia.
nevadensis, *A. Gray, in Proc. Am. Acad.* viii. (1878) 649.—Am. bor. occ.
nevadensis, *Wedd. Chlor. And.* i. 194.—Venezuela.
nigromontanus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 114 = Aucheri.
nivalis, Boiss. Diagn. Ser. I. vi. 82 = chionophilus.
nivalis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 311.—Am. bor.
niveus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 584, nomen.—Bolivia.
Novæ-Zelandiæ, *Y. Buch. in Trans. N. Z. Inst.* xvii. (1885) 287.—N. Zel.
nudatus, *A. Gray, in Proc. Am. Acad.* xx. (1885) 297.—Am. bor. occ.
nudicaulis, *Michx. Fl. Bor. Am.* ii. 124 (err. typ. 224).—Am. bor.
nudiflorus, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 456 = divergens.
nudiscapus, Sch. Bip. in Linnaea, xxii. (1849) 571 = Aster camporum.
obliquus, Linn. Mant. ii. 578 = Blumea amplectens.
obovatus, *Boiss. Fl. Orient.* iii. 167.—Kurdistan.
obovatus, Spreng. ex DC. Prod. v. 289 = Laggera flava.
occidentalis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 311.—Am. bor. occ.
ochroleucus, Eaton, in Bot. King's Exped. 152 = Eatoni.
ochroleucus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 309.—Am. bor. occ.
Olgæ, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 44.—As. centr.
olympicus, Schott & Kotschy, in Oestr. Bot. Wochenbl. vii. (1857) 380 = alpinus.
oolepis, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 586, nomen.—Bolivia.
Ovsczen, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. *Suppl.* 17.—Soongaria.
oreganus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 2.—Am. bor. occ.
orientalis, Boiss. Diagn. Ser. II. iii. 7 = acris.
orizabensis, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 91.—Mexic.
othonnæfolius, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 30.—Chili.

ERIGERON:—
Palmeri, *A. Gray, in Proc. Am. Acad.* xv. (1880) 32.—Mexic.
palustris, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 128 = maximus.
paniculatus, Lam. Fl. Fr. ii. 141 = canadensis.
paniculatus, Meig. & Wenig. Syst. Verz. 101 = acris.
paposanus, *Phil. Fl. Atac.* 29.—Chili.
pappochroma, *Labill. Nov. Holl. Pl.* ii. 47. t. 193.—Austral.
paradoxus x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 105.—Europ.
Parishii, *A. Gray, Syn. Fl. N. Am.* i. ii. 212.—Calif.
pauciflorus, Hook. & Arn. Bot. Beech. Voy. 87 = Vittadenia Chamissonis.
paucifolius, Less. ex Baker, in Mart. Fl. Bras. vi. iii. 84 = Conyza notobellidiastrum.
paxensis, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 80, nomen.—Bolivia.
pectinatus, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 147 = Conyza pectinata.
pedatus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 308 = compositus.
pellitus, Wedd. Chlor. And. i. 190 = Aster pellitus.
perramosus, *Schur, in Oestr. Bot. Zeitschr.* xix. (1869) 207.—Europ.
persicifolius, Benth. in Hook. Niger Fl. 430 = Conyza persicæfolia.
persicus, *Boiss. Diagn.* Ser. I. vi. 81.—Persia.
peucephyllus, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 89.—Am. bor. occ.
philadelphicus, Bart. Veg. Mat. Med. t. 20 = strigosus.
philadelphicus, *Linn. Sp. Pl.* 863.—Am. bor.
philadelphicus, Lour. Fl. Cochinch. 500 = cochinchinensis.
philadelphicus, Michx. Fl. Bor. Am. ii. 123 = quercifolius.
philadelphicus, Willd. Sp. Pl. iii. 1957 = acris.
Philippii, *Sch. Bip. ex Wedd. Chlor. And.* i. 192.—Chili.
phlogotrichus, Spreng. Syst. iii. 520 = pappochroma.
pilosus, *Pers. Syn.* ii. 431.—Hab.?
pilosus, Wall. Fl. Carol. 206 = Chrysopsis gossypina.
pinifolius, Poir. Encyc. viii. 490 = Hysterionica pinifolia.
pinnatifidus, D. Don, Prod. Fl. Nep. 172 = Conyza stricta.
pinnatifidus, Thunb. Prod. Fl. Cap. 158 = Conyza pinnatifida.
pinnatus, Linn. f. Suppl. 368 = Conyza pinnatifolia.
pinnatus, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 178.—Peruv.
Pisonis, *Spreng., Pl. Jul.* 160.—Bras.
podolicus, Bess. Enum. Pl. Volh. 76 = acris.
Poeppigii, *DC. Prod.* v. 287.—Chili.
policephalus, *A. Gray, Syn. Fl. N. Am.* i. ii. 210.—Am. bor. occ.
politus, *Fries, Summa Veg. Scand.* 3, 184.—Europ. bor.
polymorphus, Soop. Fl. Carn. ed. II. ii. 162 = alpinus.
polyphyllus, *Phil. in Linnaea*, xxxiii. (1864–65) 135.—Chili.
Prantlii, *Dalla Torre, Atl. Alpenfl.* 124.—Europ.
primulæfolius, Juss. ex Cass. in Dict. Sc. Nat. xlii. 61 = Podocoma primulæfolia.
Pringlei, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 210.—Arizona.
pseudo-acris, Schur, Enum. Pl. Transs. 308 = acris.
pubescens, *H. B. & K. Nov. Gen. et Sp.* iv. 88.—Mexic.
pulchellus, *DC. Prod.* v. 287.—Oriens; An. bor.
pulchellus, Hook. Fl. Bor. Am. ii. 19, partim = glabellus.
pulchellus, Michx. Fl. Bor. Am. ii. 124 = bellidifolius.
pulchellus, Turcz. in Bull. Soc. Nat. Mosc. 1840 = uniflorus.
pulcher, *Phil. in Anal. Univ. Chil.* (1873) 487.—Chili.
pulvinatus, *Wedd. Chlor. And.* i. 194.—Bolivia.
pumilus, Hook. Lond. Journ. Bot. vi. (1847) 242 partim = ochroleucus.
pumilus, *Nutt. Gen. Am.* ii. 147.—Am. bor. occ.
purpureus, Ait. Hort. Kew. ed. I. iii. 186 = philadelphicus.
pusillus, Lehm. Ind. Sem. Hort. Hamb. (1828) 8; Pugill. i. 18 = Lehmanni, *Spreng.*
pusillus, Nutt. Gen. Am. ii. 148 = canadensis.
pycnotrichus, Schott & Kotschy, ex Tchihat. Asie Min. Bot. ii. 284 = uniflorus.
pyrifolius, Benth. Fl. Hongk. 176 = Microglossa volubilis.
pyrrhopappus, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop 147 = Conyza pyrrhopappa.
quercifolius, DC. Prod. v. 285 = tenuis.

ERIGERON :—
quercifolius, *Lam. Tabl. Encyc.* iii. 258.—Am. bor.
quercifolius, Pursh, Fl. Am. Sept. ii. 538 = philadelphicus.
racemosus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 413 = armerifolius.
radiatus, *Hook. Fl. Bor. Am.* ii. 17. t. 128.—Am. bor.
radiatus, Spreng. ex DC. Prod. v. 299 = multiradiatus
ramosus, Rafin. Fl. Ludov. 66 = divaricatus !
Regnellii, Sch. Bip. in Linnæa, xxii. (1849) 571 = Aster Regnellii.
Remyanus, *Wedd. Chlor. And.* i. 195.—Chili.
repens, *A. Gray, Syn. Fl. N. Am.* i. ii. 217.—Texas.
repens, Wedd. Chlor. And. i. 191 = Aster repens.
resinosum, Spreng. Syst. iii. 519 = Hysterionica montevidensis.
retrofexus, *Poir. Encyc. Suppl.* v. 464.—Am. bor.
rhastioides v., Benegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 105.—Europ.
rigidus, *Pers. Mant.* iii. 107.—Norveg.
rivularis, Spreng. ex DC. Prod. v. 288 = bellioides.
rivularis, *Sw. in Prod. Veg. Ind. Occ.* 113.—Ind. occ.
rosulatus, *Wedd. Chlor. And.* i. 198.—Bolivia.
Rottleri, *Spreng. Syst.* iii. 518.—Ind. or.
Royleï, DC. Prod. v. 292 = alpinus!
rubricaulis, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 536, nomen.—Bolivia.
ruderalis, Salisb. Prod. 195 = canadensis.
rufescens, *Link, ex Steud. Nom.* ed. I. 312.—Lusitan.
runcinatus, Bert. ex DC. Prod. v. 288 = rivularis.
rupestris, Hoppe, ex DC. l. c. 291 = alpinus.
rupestris, *Schleich. Cat. Pl. Helv.* ed. IV. 18.—Europ.
rupicolus, *Phil. in Bot. Zeit.* xiv. (1856) 644.—Ins. J. Fernand.
rutilus, Poir. Encyc. Suppl. v. 464 = Conyza calocephala, sericea.
saginoides, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 98.—Am. bor. occ.
scaber, Pers. Syn. ii. 431 = Conyza meiastomoides.
scaber, Thunb. Prod. Fl. Cap. 168 = Pulicaria capensis.
scaber, Willd. ex Spreng Syst. iii. 517 = acris.
scabrivimus, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 80 = maximus.
scandens, *Thunb. Fl. Jap.* 318.—Japon.
scandinavicus, Scheele, ex Nym. Consp. 389 = politus.
scaposus, *DC. Prod.* v. 287.—Mexic.
scaposus, Torr. & Gray, Fl. N. Am. ii. 170 = repens.
scariosus, Moench, Meth. Suppl. 247 = Conyza Gouani.
Schiedeanus, *Less. in Linnæa*, v. (1830) 145.—Mexic.
Schimperi, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 147 = Conyza stricta.
Schuechtii, Sch. Bip. in Linnæa, xxii. (1849) 571 = maximus.
Schultzii, *Zoll. Syst. Vers. Ind. Archip.* 120, 127.—Java.
scorzonerifolius, *Remy, in C. Gay, Fl. Chil.* iv. 26.—Chili.
semiamplexicaulis, *Meyen, Reise*, i. 311.—Chili.
semiharbatus, DC. Prod. v. 292 = alpinus.
Senecio, Auct. Am. ex Steud. Nom. ed. I. 312 = canadensis.
Senecio, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 218 = Senecio vulgaris.
senecioides, *Wedd. Chlor. And.* i. 198.—Peruv.
sericeus, *Retz. Obs.* v. 38.—Java.
serotinus, Wibthe, ex Reichb. Fl. Germ. Excurs. 289 = acris.
serpentarius, Banks, ex Steud. Nom. ed. I. 312 = bellidifolius.
serratus, Forsk. Fl. Aegypt. Arab. 148 = aegyptiacus.
sessiliflia, *R. Mudl. Fragm.* xi. 100.—Austral.
sicanus, Linn. Sp. Pl. 864 = Pulicaria sicula.
solidaginoides, *Schlecht. in Linnæa*, xxv. (1852) 218. —Canada.
sordidus, Gill. ex Hook. & Arn. Comp. Bot. Mag. ii. (1836) 254 = bonariensis.
spathulatus, *Schum. & Thonn. Beskr. Guin. Pl.* 355. —Afr. trop.
spathulatus, *Vahl, in West, Bidr. Ste Croix*, 303.— Ins. Antigua.
spathulatus, *Wedd. Chlor. And.* i. 197.—Ecuador.
speciosus, *DC. Prod.* v. 294.—Am. bor. occ.
spiciformis, *Griseb. in Gœtt. Abh.* xix. (1874) 171.— Reg. Argent.
spiculosus, *Hook. & Arn. Bot. Beech. Voy.* i. 32.— Chili.

ERIGERON :—
spinosus, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 148, nomen = Conyza spinosa.
spinulosus, Phil. in Linnæa, xxviii. (1856–58) 18 = spiculosus.
Sprengelii, *Sch. Bip. ex Walp. Rep.* ii. 958.—Afr. austr.
squarrosus, [Clairv.] Man. Herb. 245 = Inula squarrosa.
squarrosus, Lindl. Bot. Reg. (1841) Misc. 44 = glaucus !
squarrosus, Walt. Fl. Carol. 206 = Chrysopsis gossypina.
stenophyllus, Eaton, in Bot. King's Exp. 152. t. 17 = Aster arenarioides.
stenophyllus, *A. Gray, in Pacif. Rail. Rep.* iv. 98.— Am. bor. occ.
stenophyllus, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 49.—Chili.
stenophyllus, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 176 = foliosus.
Steudelii, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 148 = Conyza Steudelii.
stipulatus, Schum. & Thonn. Beskr. Guin. Pl. 355 = Laggera auriis.
Stocksianus, *Boiss. Diagn.* Ser. II. iii. 8.—Afghan.
strictissimus, Wender. Ind. Sem. Hort. Mark. (1825); ex Literatur. Flora, lii. (1838) 237 = Conyza chilensis.
strictus, DC. Prod. v. 289 = canadensis.
strictus, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 49.—Chili.
strygosus, Bigel. Fl. Bost. ed. II. 302 = annuus.
strigosus, *Muhl. ex Willd. Sp. Pl.* iii. 1956.—Am. bor.
subandinus, *Phil. in Linnæa*, xxviii. (1856) 723.— Chili.
subdecurrens, A. Gray, in Pacif. Rail. Rep. iv. 97 = Conyza Coulteri.
subdecurrens, *Sch. Bip. ex A. Gray, Syn. Fl. N. Am.* i. ii. 230.—Arizona.
subvestitus, Rozb. ex DC. in Wight, Contrib. 9 = asteroides.
subspicatus, *Benth. in Oerst. Kjoeb. Vidensk. Meddel.* (1852) 92.—Costa Rica.
sulcatus, DC. Prod. v. 284 = maximus.
sulcatus, Meyen, Reise, i. 311 = Conyza chilensis.
sulcatus, Nees, in Pr. Neuwied, Reise Nord. Am. ii. 439 = pumilus.
Sulivani, *Hook. f. Fl. Antarct.* 306.—Chili.
sumatrensis, *Retz. Obs.* v. 38.—Sumatra.
supplex, *A. Gray, in Proc. Am. Acad.* vii. (1866) 358. —Calif.
sylvaticus, *Phil. in Linnæa*, xxviii. (1856) 723.— Chili.
tasmanicus, Hook. f. Fl. Tasm. i. 158. t. 46 = pappochromus.
tenellus, *DC. Prod.* v. 288.—Mexic.
tenellus, Flœh. ex Herd. in Bull. Soc. Nat. Mosc. xxvii. (1866) i. 398 = armerifolius.
tener, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 91.— Calif.
tenuifolius, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 207.—Ins. Galap.
tenuifolius, Sch. Bip. in Flora, xxvii. (1844) 670 = Chrysocoma tenuifolia.
tenuis, *Torr. & Gray, Fl. N. Am.* ii. 176.—Am. bor. chrincicoles, *Griseb. Cat. Pl. Cub.* 148.—Cuba.
Thunbergii, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 895.—Japon.
Trachselii v., Dalla Torre, Atl. Alpenfl. 125.—Europ.
transsilvanicus, Schur, Enum. Pl. Transs. 307 = linifolius.
trichodepis, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 536, nomen.—Bolivia.
tricuneatus, Linn. f. Suppl. 368 = Baccharis conferta.
trifoliolatus, DC. Prod. v. 290 = Vittadinia trifoveata.
trifidus, Hook. Fl. Bor. Am. ii. 17 = compositus.
trifidus, *Schlecht. in Linnæa*, xxvii. (1854) 518.—N. Granat.
trifoveatus, Gill. & Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 49 = Vittadinia trifoveata.
trilobus, *Boiss. Fl. Orient.* iii. 168.—Arabia.
trilobus, Sond. in Hamb. Gartenz. xii. (1856) 78 = Vittadinia australis.
triplinervis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 111. 33 = Conyza tripliner via.

ERIGERON :—
tripolioides, *Phil. in Linnæa*, xxxiii. (1864–65) 185.— Chili.
trinervis, D. Don, Prod. Fl. Nep. 171 = Conyza stricta.
tuberosus, Linn. Sp. Pl. 864 = Jasonia tuberosa.
Turczaninowii, Wedd. Chlor. And. i. 198 = pinnatus.
Tweediei, Benth. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 50 = Leucocyta Tweediei.
uliginosus, *Benth. Pl. Hartw.* 204.—N. Granat.
uliginosus, Turcz. ex Ledeb. Fl. Ross. ii. 489 = armerifolius.
umbellatus, Gilib. Fl. Lituan. i. 215 (err. typ. 125) = acris.
undulatus, Moench, Meth. 598 = bonariensis.
uniflorus, *Linn. Sp. Pl.* 864.—Reg. bor. et arct.
uniflorus, Sm. Engl. Fl. iii. 423 = alpinus.
unifolius, err. typ. p. 607, col. 1, sub *Conyza ambigua*, lege 'linifolius.'
uralensis, Less. in Linnæa, ix. (1834) 154 = acris.
ursinus, *Eaton, in Bot. King's Exp.* 148.—Am. bor. occ.
utahensis, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 89.—Am. bor. occ.
Vahlii, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 108.—Ins. Falkland.
variegatus, Sch. Bip. ex Schweinf. & Aschers. in Beitr. Fl. Aethiop. 285 = Conyza variegata.
varius, *Webb, in Hook. Niger Fl.* 134.—Ins. Cap. Verd.
velutipes, *Hook. & Arn. Bot. Beech. Voy.* 134.—Mexic.
vernus, Torr. & Gray, Fl. N. Am. ii. 176 = nudicaulis.
Villarsii, *Bell. Ann. Fl. Pedem.* 38. t. 7.—Europ.
Villarsii, Willd. Sp. Pl. iii. 1958 = acris.
villosus, Steud. Nom. ed. I. 312 = Aster villosus.
virens, Ruiz. ex DC. Prod. v. 439 = Blumea virens.
virgatus, *Gueldenst. ex Ledeb. Fl. Ross.* ii. 491 (quid ?).—Ross. austr.
viscosus, Desf. Tabl. ed. I. 109 = Psiadia glutinosa.
viscosus, Linn. Sp. Pl. 863 = Inula viscosa.
Vittadinia, F. Muell. Fragm. v. 87 = Vittadinia scabra.
vulgaris, Scheele, ex Nym. Consp. 389 = acris.
Wightianus, Wall. Cat. n. 3264 = Pulicaria Wightiana.
Wightii, DC. Prod. v. 286 = alpinus.
serdanus, Boiss. Diagn. Ser. I. vi. 82 = amorphoglossus.
Zollingeri, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 120, nomen; *Miq. Fl. Ind. Bat.* ii. 31.—Java.

ERIGERUM, Linn. Gen. ed. I. 254 (1737) = **Erigeron**, Linn. (Compos.).

ERIGONE, Salisb. Gen. Pl. Fragm. 115 (1866) = **Crinum**, Linn. (Amaryllid.).
Govenica, Salisb. l. c. 116 = C. longifolium !
longifora, Salisb. l. c. = C. longiflorum.
longifolia, Salisb. l. c. = C. longifolium.
revolute, Salisb. l. c. = C. lineare.
riparia, Salisb. l. c. = C. longifolium.
viridiflora, Salisb. l. c. = C. Commelyni.

ERIGONIA, A. Juss. in Orb. Dict. ix. 95 (1849) = **Erigenia**, Nutt. (Umbellif.).

ERIMATALIA, Roem. & Schult. Syst. v. 27 (1819) = **Eryedbe**, Roxb. (Convolv.).
Rheedii, Roem. & Schult. l. c. 331 = Eryc. paniculata.

ERINACEA, [Tourn.] Adans. Fam. ii. 321 (1763). *LEGUMINOSÆ*, Benth. & Hook. f. i. 462. pungens, *Boiss. Voy. Bot. Espagne*, 145.—Hispan.

ERINGIUM, Neck. Delic. Gallo-Belg. i. 138 (1768) = **Eryngium**, Tourn. (Umbell.).

ERINIA, Noulet, Fl. Sous-Pyr. 407 (1837) = **Campanula**, Tourn.
Campanula, Noulet, l. c. = C. Erinus.

ERINNA, Phil. in Linnæa, xxxiii. (1864–65) 266. *LILIACEÆ*. Benth. & Hook. f. iii. 865.
gilliesioides, *Phil. l. c.*—Chili.

ERINOCARPUS, Nimmo, ex J. Grah. Cat. Bomb. Pl. 21 (1839). *TILIACEÆ*, Benth. & Hook. f. i. 284.
Keimoni, Hassk. Retzia, i. 127 = Nimmonii.
Nimmoanus, Mast. in Hook. f. Fl. Brit. Ind. i. 294 = seq.
Nimmonii, *J. Grah. Cat. Bomb. Pl.* 21.—Ind. or.

ERINOSMA, Herb. Amaryll. 330 (1837) = **Leucojum**, Linn. (Amaryllid.).
carpathica, Herb. in Bot. Mag. t. 1998 = L. vernum.
vernæ, Herb. Amaryll. 331 = L. vernum.

ERINUS, Linn. Hort. Cliff. 320 (1737). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 961.
AGERATUM, Tourn. ex Adans Fam. ii. 210 (1763).
asthiopicus, *Thunb. Prod. Pl. Cap.* 102.—Afr. austr.
africanus, Linn. Sp. Pl. 630 = Zaluzianskya villosa.
alpinus, *Linn. l. c.*—Europ.
americanus, Mill. Gard. Dict. ed. VIII. n. 3 = Buchnera americana.
bilabiatus, Roxb. Hort. Beng. 47; Fl. Ind. iii. 92 = Adenosma capitatum.
campanulata, Nym. Consp. 481 = Campanula Erinus.
capensis, Linn. Mant. ii. 252 = Zaluzianskya capensis.
fragrans, Ait. Hort. Kew. ed. I. ii. 357 = Zaluzianskya lychnidea.
fruticescens, Mill. Gard. Dict. ed. VIII. n. 4 = Capraria asxifrageæfolia.
glabratus, Nym. Consp. 544 = alpinus.
gracilis, *Lehm. in Linnæa*, x. (1836) *Litt.* 76.—Afr. austr.
hispanicus, Pers. Syn. ii. 147 = alpinus.
incisus, Thunb. Prod. Pl. Cap. 103.—Afr. austr.
lanistatus, Linn. Sp. Pl. 630 = Verbena erinoides.
lanceolatus, Kitt. Taschenb. ed. II. 385 = alpinus.
lychnideus, Lam. Encyc. Suppl. ii. 336 = Zaluzianskya lychnidea.
lychnidoides, Linn. f. Suppl. 287 = Zaluzianskya capensis.
lychnideus, Thunb. Prod. Pl. Cap. 102 = Lyperia fragrans.
maritimus, Linn. f. Suppl. 287 = Zaluzianskya maritima.
patens, *Thunb. Prod. Pl. Cap.* 102.—Afr. austr.
peruvianus, Linn. Sp. Pl. 630 = Verbena chamædrifolia.
procumbens, Mill. Gard. Dict. ed. VIII. n. 6 = Herpestis chamædryoides.
selaginoides, Thunb. Prod. Pl. Cap. 102 = Zaluzianskya selaginoides.
simplex, *Thunb. l. c.*—Afr. austr.
tomentosus, *Mill. Gard. Dict.* ed. VIII. n. 2.—Vera Cruz.
tomentosus, *Thunb. Prod. Pl. Cap.* 103.—Afr. austr.
tristis, *Linn. f. Suppl.* 287.—Afr. austr.
verticillatus, Mill. Gard. Dict. ed. VIII. n. 5 = Stemodia parviflora.
villosus, Thunb. Prod. Pl. Cap. 102 = Zaluzianskya villosa.
viscosus, Salisb. Prod. 94 = Sphenandra viscosa.

ERIOBOTRYA, Lindl. in Trans. Linn. Soc. xiii. (1822) 102. *ROSACEÆ* [Benth. & Hook. f. i. 637].
angustissima, *Hook. f. Fl. Brit. Ind.* ii. 373.—Reg. Himal.
bengalensis, *Hook. f. l. c.* 371.—Reg. Himal.; Malaya.
chinensis, G. Don, Gen. Syst. ii. 608 = Crataegus chinensis.
cordata, Lindl. in Trans. Linn. Soc. xiii. (1822) 102 = Osteomeles latifolia.
cuneata, D. Dietr. Syn. Pl. iii. 163 (= *Hesperomeles cuneata*).—Peruv.
dubia, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 145.—Reg. Himal.
elliptica, *Lindl. in Trans. Linn. Soc.* xiii. (1822) 102. —Reg. Himal.
fragrans, *Champ. ex Benth. in Hook. Kew Journ.* ix. (1857) 80.—Hongkong.
glabrata, Steud. Nom. ed. II. i. 585 = Osteomeles glabrata.
heterophylla, *Lindl. in Trans. Linn. Soc.* xiii. (1822) 102.—Peruv.
Hookeriana, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 146.—Reg. Himal.
integrifolia, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 304 = Photinia Notoniana.
japonica, *Lindl. in Trans. Linn. Soc.* xiii. (1822) 102. —Japon.; China.
latifolia, *Hook. f. Fl. Brit. Ind.* ii. 370.—Burma.
longifolia, *Hook. f. l. c.*—Reg. Himal.
macrocarpa, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 306.—Burma.
oblonga, D. Dietr. Syn. Pl. iii. 163 (= *Hesperomeles oblonga*).—N. Granat.
obtusifolia, DC. Prod. ii. 632 = Osteomeles glabrata.
petiolata, *Hook. f. Fl. Brit. Ind.* ii. 370.—Reg. Himal.

ERIOCACHRYS, DC. Prod. iv. 241, in syn. (1830) = **Magydaris**, Koch (Umbellif.).

ERIOCALIA, Sm. Exot. Bot. 37. t. 78, 79 (1806) = **Actinotus**, Labill. (Umbellif.).
leucocephala, Heynh. Nom. ii. 239 = A. leucocephala.
major, Sm. Exot. Bot. ii. 37 = A. Helianthi.
minor, Sm. l. c. 39 = A. minor.

ERIOCALYX, Endl. Gen. 1264, in syn. (1840) = ERIOCYLAX, Neck. = **Aspalathus**, Linn. (Legumin.).

ERIOCARPÆA, Bertol. in Nov. Comm. Acad. Bonon. vi. (1843) 284; (Misc. Bot. ii. 20) = **Onobryohis**, Gaertn. (Legumin.).
pinnata, Bertol. ll. cc. = O. Gaillardoti.

ERIOCARPHA, Cass. in Dict. Sc. Nat. lix. 236 (1829) = **Montanoa**, La Llave (Compos.).

ERIOCARPHA, Lag. ex DC. Prod. vi. 38 (1837) = **Lasiospermum**, Lag. (Compos.).
pedunculatis, Lag. ex DC. l. c. = L. pedunculare.

ERIOCARPUM, Nutt. in Trans. Am. Phil. Soc. N. S vii. (1841) 320 = **Aplopappus**, Cass. (Compos.).
grindelioides, Nutt. l. c. = A. Nuttalii.

ERIOCAULON, Linn. Gen. ed. II. 55 (1742). *ERIOCAULEÆ*, Benth. & Hook. f. iii. 1020.
CRAETODISCUS, Steud. Syn. Pl. Cyp. 261 (1855).
ELECTROSPERMA, F. Muell. in Trans. Phil. Soc. Vict. i. (1854) 23.
LASIOLEPIS, Boeck. in Flora, lvi. (1873) 90.
LEUCOCEPHALA, Roxb. Fl. Ind. iii. 612 (1832).
NASMYTHIA, Huds. Fl. Angl. ed. I. 414 (1762).
RANDALIA, Beauv. ex Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 47. t. 5.
SPHAEROCHLOA, Beauv. ex Desv. l. c. (1828).
SYMPHACHNE, Beauv. ex Desv. l. c. (1828).
abyssinicum, *Hochst. in Flora*, xxvii. (1845) 341.—Abyss.
Achiton, Koern. in Linnæa, xxvii. (1854) 630 = nigricans.
aciphyllum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 636 = Paepalanthus aciphyllus.
aciphyllum, Reichb. ex Koern. in Mart. Fl. Bras. iii. i. 356 = Paepalanthus subtilis.
acrotrichum, Steud. Syn. Pl. Cyp. 277 = Paepalanthus Schrader.
affine, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 624 = Paepalanthus affinis.
africanum, *Hochst. in Flora*, xxviii. (1845) 340.—Afr. austr.
Afzelianum, *Wikstr. ex Koern. in Linnæa*, xxvii. (1854) 630.—Afr. trop.
albidum, Steud. Syn. Pl. Cyp. ii. 278 = Paepalanthus albidus.
alpestre, *Hook. f. & Thoms. ex Koern. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 163.—Ind. or.
amoenum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 637 = Paepalanthus amœnus.
amplexicaule, Rottb. in Act. Lit. Univ. Hafn. I. (1778) 272 = Tonina fluviatilis.
ancepe, Walt. Fl. Carol. 88 = Lachnocaulon Michauxii.
angustifolium, *Koern. in Mart. Fl. Bras.* iii. i. 495.—Bras.
anthemiflorum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 636 = Paepalanthus anthemidiflorus.
aquaticum, Segot, ex Koern. in Mart. Fl. Bras. iii. i. 498 = melanocephalum.
aquatile, *Koern. in Linnæa*, xxvii. (1854) 600, nomen; *et in Mart. Fl. Bras.* iii. i. 495.—Bras.
arctiotiflorum, Steud. Syn. Pl. Cyp. 282 = Paepalanthus anthemidiflorus.
arenarium, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 442 = Paepalanthus arenarius.
arenarium, Salzm. ex Koern. in Mart. Fl. Bras. iii. i. 356 = Paepalanthus subtilis.
argenteum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 631 = Paepalanthus argenteus.
argentinum, Mart. in Wall. Pl. As. Rar. iii. 28 = quinquangulare.
argyraeum, Steud. Syn. Pl. Cyp. 271 = quinquangulare.
stratum, *Koern. in Linnæa*, xxvii. (1854) 610.—Zeylan.
atrocephalum, Steud. Syn. Pl. Cyp. 276 = Paepalanthus tortilis.

ERIOCAULON:—
australasicum, *Koern. in Linnæa*, xxvii. (1854) 616. —Austral.
australe, *R. Br. Prod.* 254.—Austral.
bahiense, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 629 = Paepalanthus bahiensis.
Benthami, *Kunth, Enum. Pl.* iii. 545.—Mexic.
Benthami, *Seem. Bot. Voy. Herald*, 221.—Panama.
Benthamii, Schlecht. in Linnæa, xviii. (1844) 434 = Ekenbergianum.
bifidum, Schrad. ex Schult. Mant. ii. 468 = Paepalanthus Schraderi.
bifurcatum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 105.—Afr. trop.
bisumbellatum, Steud. Syn. Pl. Cyp. 275 = Paepalanthus Steudellanus.
blepharophoron, Bong. in Mém. Acad. Pétersb. Ser VI. i. (1831) 626. t. 16 = Paepalanthus blepharophoron.
Blumei, *Koern. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 240.—Java.
Bongardii, St. Hil. Voy. Distr Diam. ii. 444 = Paepalanthus Hilairei.
Bomplandianum, Steud. Syn. Pl. Cyp. 275 = Paepalanthus Humboldtii.
borbonicum, Willd. ex Kunth, Enum. Pl. iii. 560 = striatum.
brachypeplon, *Koern. in Linnæa*, xxvii. (1854) 665.—N. Irlandia †
brachyphyllum, Willd. ex Kunth, Enum. Pl. iii. 534 = Paepalanthus tenuis.
brachypus, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 622, 646 · Paepalanthus brachypus.
brachypus, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 96.—Peruv.
bracteotum, Steud. Syn. Pl. Cyp. 272 = robustum.
brasiliense, Mart. in Flora, xxiv. (1841) ii. Beibl. 36 = Paepalanthus brasiliensis.
brevifolium, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1116, nomen; *et Koern. in Mart. Fl. Bras.* iii. i. 496.—Guian.
brevifolium, Mart. ex Koern. in Mart. Fl. Bras. iii. i. 485 = Sellowianum.
breviscapum, *Koern. in Linnæa*, xxvii. (1854) 676.—Ind. or.
brisoides, Steud. Syn. Pl. Cyp. 281 = Paepalanthus gracilis.
Brownianum, *Mart. in Wall. Pl. As. Rar.* iii. 25.—Ind. or.
bryoides, Ried. ex Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 624 = Paepalanthus bryoides.
Buergerianum, *Koern. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 163.—Japon.
caesium, *Griseb. Fl. Brit. W. Ind.* 526.—Ind. occ.
caespitosum, Poepp. ex Koern. in Mart. Fl. Bras. iii. i. 361 = Paepalanthus Schraderi.
caespitosum, Wikstr. in Vet. Acad. Handl. Stockh. (1820) 78. t. 4 = Paepalanthus caespitosus.
calocephalum, Bong. in Mém. Acad Pétersb. Sér. VI. i. (1831) 625 = Paepalanthus curvifolius.
canescens, Bong. l. c. 623. t. 50 = Paepalanthus canescens.
cantonense, *Hook. & Arn. Bot. Beech. Voy.* 219.—China.
capillare, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 625. t. 48 = Paepalanthus capillaris.
caricifolium, Steud. Syn. Pl. Cyp. 281 = Paepalanthus laricifolius.
caulescens, Hook. f. & Thoms. ex Thw. Enum. Pl. Zeyl. 341 = striatum.
caulescens, Poir. Encyc Suppl. iii 162 = Paepalanthus caulescens.
caulescens, Willd. ex Koern. in Mart. Fl. Bras. iii. i. 363 = Paepalanthus dendroides.
centauroides, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 635. t. 45 = Paepalanthus centauroides.
Cephalotes, Poir. Encyc. Suppl. iii. 161, in obs. = gnaphalodes.
ceylanicum, *Koern. in Linnæa*, xxvii. (1854) 667.—Zeylan.
chrysanthemiflorum, Schnizl. Ic. t. 46. f. 3 = Paepalanthus anthemidiflorus.
chrysanthus, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 623 = Paepalanthus chrysanthus.
ciliatum, Bong. l. c. 632 = Paepalanthus ciliatus.
ciliiflorum, F. Muell. Fragm. i. 95 = cinereum.
cinereum, *R. Br. Prod.* 254.—Austral.
cinereum, Buch.-Ham. ex Mart. in Wall Pl. As. Rar. iii. 29 = Hamiltonianum.

ERIOCAULON :—

circinnatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 632 = Paepalanthus circinnatus.
cladocaulon, Steud. Syn. Pl. Cyp. 276 = Paepalanthus Gardnerianus.
Clausseniauum, Steud. l. c. 279 = Paepalanthus rufulus.
compactum, Steud. l. c. 274 = Paepalanthus compactus.
compressum, *Lam. Encyc.* iii. 276.—Am. bor.
concretum, *F. Muell. Fragm.* i. 92.—Austral.
congestum, H. B. & K. Nov. Gen. et Sp. i. 252 = Paepalanthus fasciculatus.
consanguineum, Kunth, Enum. Pl. iii. 566 = Wallichianum.
corymbosum, Boug. in Mem. Acad. Pétersb. Sér. VI. i. 1831) 629 = Paepalanthus corymbosus.
crassifolium, Boug. l. c. 634 = Paepalanthus crassifolius.
crassiscapum, *Boug. l. c.* 628.—Bras.
cristatum, Mart. ex Koern. in Linnæa, xxvii. (1854) 607 = micrurum.
cristatum, *Mart. in Wall. Pl. As. Rar.* iii. 28.—Ind. or.
curvifolium, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 627 = Paepalanthus curvifolius.
cuspidatum, *Dals. in Hook. Kew Journ.* iii. (1851) 281.—Ind. or.
cuyabense, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 634. t. 10 = Philodice cuyabensis.
Dalzellii, *Koern. in Linnæa*, xxvii. (1854) 605.—Ind. or.
decangulare, Hope, in Phil. Trans. lix. (1770) 248. t. 12 = septangulare.
decangulare, *Linn. Sp. Pl.* 87.—Am. bor.
decangulare, Wall. Fl. Carol. 33 = compressum.
decangulare, Willd. ex Kunth, Enum. Pl. iii. 544 = Humboldtii.
decemangulare, H. B. & K. Nov. Gen. et Sp. i. 254 = Humboldtii.
decumbens, Steud. syn. Pl. Cyp. 276 = Paepalanthus Bongardi.
dendroides, H. B. & K. Nov. Gen. et Sp. i. 252 = Paepalanthus dendroides.
depressum. *R. Br. Prod.* 255.—Austral.
desutum, *R. Br. l. c.*—Austral.
dictyophyllum, *Koern. in Linnæa*, xxvii. (1854) 600, nomen ; *et in Mart. Fl. Bras.* iii. i. 486.—Bras.
distichophyllum, Mart. in Flora, xxiv. (1841) ii. Beibl. 36 = Paepalanthus distichophyllus.
divaricatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 631, 641 = Paepalanthus divaricatus.
Dregei, *Hochst. in Flora*, xxviii. (1845) 341.—Afr. austr.
echinospermum, *Wright, in Sauv. Fl. Cub.* 162.—Cuba.
echinulatum, *Mart. in Wall. Pl. As. Rar.* iii. 29.—Ind. or.
Ehrenbergianum, *Klotzsch, ex Koern. in Mart. Fl. Bras.* iii. i. 491.—Mexlc.
elatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 631. t. 63 = Paepalanthus elatus.
Electrospermum, F. Muell. Census, 123 = australasicum.
elegans, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 635 = Paepalanthus elegans.
elichrysoides, *Boug. l. c.* 631.—Bras.
elichrysoides, Kunth, Enum. Pl. iii. 546 = Kunthii.
elongatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 630. t. 61 = Paepalanthus helichrysoides.
enatifolium, H. B. & K. Nov. Gen. et Sp. i. 254. t. 70 = Paepalanthus enatifolius.
erythropodum, Miq. ex Koern. in Linnæa, xxvii. (1854) 643 = quinquangulare.
eurypeplon, *Koern. l. c.* 605.—Ind. or.
exiguum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 637. t. 59 = Paepalanthus exiguus.
falcatum, *Boug. l. c.* 630.—Bras.
falcatum, Mart. ex Koern. in Mart. Fl. Bras. iii. i. 432 = Paepalanthus flavescens.
flaviculare, Weigelt, ex Kunth, Enum. Pl. iii. 506 = Paepalanthus congestus.
fasciculatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 634 = Paepalanthus Schraderi.
fasciculatum, Lam. Encyc. iii. 276 = Paepalanthus Lamarckii.
fasciculatum, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 271 = Paepalanthus congestus.
fastigiatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 624. t. 60 = Paepalanthus fastigiatus.

ERIOCAULON :—

fenestratum, *Bot. ex Koern. in Linnæa*, xxvii. (1854) 671.—Madag.
filiforme, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 634. t. 58 = Paepalanthus nitens.
filiformo, *Rafin. Atl. Journ.* 121.—Am. bor.
Fischerianum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 637. t. 29 = Paepalanthus Fischerianus.
fistulosum, *R. Br. Prod.* 255.—Austral.
flaccidum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 636 = Paepalanthus flaccidus.
flagellare, Gulll. in Deless. Ic. Sel. iii. 60. t. 96 = Paepalanthus flagellaris.
flavescens, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 628 = Paepalanthus flavescens.
flavidulum, Michx. Fl. Bor. Am. ii. 166 = Paepalanthus flavidulus.
flavidum, A. Gray, Man. Bot. N. U. St. ed. II. 489 = Paepalanthus flavidulus.
fluitans, Baker, in Journ. Linn. Soc. xx. (1883) 277 = melanocephalum.
fluviatile, *Trim. in Journ. Bot.* xxiii. (1885) 270.—Zeylan.
Freycynii, Thunb. Pl. Bras. Dec. i. 7. t. 1. f. 2 = Paepalanthus paludosus.
fuliginosum, *Griseb, Cat. Pl. Cub.* 226.—Cuba.
Gardneri, Steud. Syn. Pl. Cyp. 274 = Paepalanthus polyanthus.
geniculatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 625 = Paepalanthus geniculatus.
geratense, Steud. Syn. Pl. Cyp. 276 = Paepalanthus caulescens.
gibbosum, *Koern. in Linnæa*, xxvii. (1854) 600, nomen ; *et in Mart. Fl. Bras.* iii. i. 489.—Bras.
giganteum, Afzel. ex Koern. in Linnæa, xxvii. (1854) 573 = Mesanthemum radicans.
glabrum, Steud. Syn. Pl. Cyp. 281 = Paepalanthus gracilis.
glaucosum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 625 = Paepalanthus glaucosus.
glaucum, *Griff. Notul.* 118.—Burma.
gnaphaloides, Ell. Sketch, ii. 565 = decangulare.
gnaphaloides, Michx. Fl. Bor. Am. ii. 165 = compressum.
gomphrenoides, *Kunth, Enum. Pl.* iii. 548.—Bras.
gracile, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 634. t. 46 = Paepalanthus gracilis.
gracile, *Mart. in Wall. Pl. As. Rar.* iii. 29.—Ind. or.
gramineum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 627 = Paepalanthus gramineus.
gregale, *Koern. in Linnæa*, xxvii. (1854) 606.—Ind. or.
grisæum, *Koern. in Mart. Fl. Bras.* iii. i. 479.—Bras.
guianense, Steud. Syn. Pl. Cyp. 278 = Mesanthemum radicans.
guianense, D. Dietr. Syn. Pl. v. 263 = Humboldtii.
guyanense, *Koern. l. c.* 478.—Guian.
Hamiltonianum, *Mart. in Wall. Pl. As. Rar.* iii. 29.—Ind. or.
helichrysoides, Steud. Syn. Pl. Cyp. 283 = Paepalanthus helichrysoides.
heteranthum, Benth. Fl. Hongk. 382 = cinereum.
heterochiton, *Koern. in Miq. Ann Mus. Bot. Lugd. Bat.* iii. 240.—Madag.
heterogynum, F. Muell. Fragm. i. 98 = depressum.
heterolepis, *Steud. Syn. Pl. Cyp.* 271.—Ind. or.
heterostachyum, *Bong. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 626. t. 39.—Bras.
hirsutum, Wikstr. in Vet. Acad. Handl. Stockh. (1820) 79. t. 4 = Paepalanthus hirsutus.
hirtellum, Steud. Syn. Pl. Cyp. 280 = Paepalanthus gracilis.
homomallum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 626 = Paepalanthus homomallus.
Hostmanni, Steud. Syn. Pl. Cyp. 281 = Paepalanthus simplex.
Humboldtii, *Kunth, Enum. Pl.* iii. 544.—Guian.
incanum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 633. t. 61 = Paepalanthus incanus.
infirmum, Steud. Syn. Pl. Cyp. 271 = gracile.
intermedium, Koern. in Linnæa, xxvii. (1854) 601 = setaceum.
isidiifolium, Steud. Syn. Pl. Cyp. 278 = Paepalanthus monticolus.
itkyphyllum, Mart. in Flora, xxiv. (1841) ii. Beibl. 35 = Paepalanthus ithyphyllus.
japonicum, *Koern. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 169.—Japon.

ERIOCAULON :—

junciformis, Steud. Syn. Pl. Cyp. 283 = Paepalanthus flaccidus.
Koernickianum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 101.—Texas.
Kunthianum, Mart. ex Koern. in Mart. Fl. Bras. iii. i. 365 = Paepalanthus caespititius.
Kunthii, *Koern. in Linnæa*, xxvii. (1854) 599, nomen ; *et in Mart. Fl. Bras.* iii. i. 482.—Bras.
Lamarckii, Steud. Syn. Pl. Cyp. 276 = Paepalanthus Lamarckii.
lanceolatum, *Miq. ex Steud. l. c.* 271.—Ind. or.
Langsdorffii, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 632. t. 51 = Paepalanthus Langsdorffii.
lanuginosum, Boug. l. c. 627 = Paepalanthus lanuginosus.
latifolium, Boug. l. c. 631. t. 64 = Paepalanthus latifolius.
laxifolium, *Koern. in Linnæa*, xxvii. (1854) 600, nomen ; *et in Mart. Fl. Bras.* iii. i. 494.—Bras.
leptophyllum, *Kunth, Enum. Pl.* iii. 549.—Bras.
leucocephalum, *Steud. Syn. Pl. Cyp.* 272.—Ins. Borbon.
leucomelas, Steud. Nom. ed. II. i. 585 = melaleucum.
Lhotzkyi, Steud. Syn. Pl. Cyp. 270 = Smithii.
lingulatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 626 = Paepalanthus lingulatus.
linearifolium, *Koern. in Linnæa*, xxvii. (1854) 601, nomen ; *et in Mart. Fl. Bras.* iii. i. 496.—Bras.
lividum, *F. Muell. Fragm.* i. 92.—Austral.
longifolium, *Nees, ex Kunth, Enum. Pl.* iii. 567.—As. trop. ; Madag.
longipes, Mart. in Flora, xxiv. (1841) ii. Beibl. 35 = Paepalanthus amoenus.
lusulaefolium, *Mart. in Wall. Pl. As. Rar.* iii. 28.—Ind. or.
macrobolax, *Koern. in Linnæa*, xxvii. (1854) 599, nomen ; *et in Mart. Fl. Bras.* iii. i. 484.—Bras.
macrocephalum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 630 = Paepalanthus macrocephalus.
macrorhizon, Boug. l. c. 631 = Paepalanthus macrorhizus.
Martianum, Wall. Cat. n. 7379 = quinquangulare.
Maximiliani, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 620 = Paepalanthus Hilairei.
Maximiliani, Mart. in Herb. Fl. Bras. n. 897 = Paepalanthus nitens.
Maximiliani, Schrad. in Schult. Mant. ii. 470 = Paepalanthus ramosus.
melaleucum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 629 = Paepalanthus melaleucus.
melanocephalum, *Mart. in Wall. Pl. As. Rar.* iii. 29.—Ind. or.
melanocephalum, *Kunth, Enum. Pl.* iii. 549.—Bras.
Meteniumum, C. Muell. in Walp. Ann. vi. 1171, sphalm. = lanceolatum.
microcephalum, Cham. & Schlecht. in Linnæa, vi. (1831) 43 = Schlechtendahlii.
microcephalum, Hook. & Arn. Bot. Beech. Voy. 311 = Benthami, Seem.
microcephalum, *H. B. & K. Nov. Gen. et Sp.* i. 253.—Ecuador.
microphyllum, Gulll. in Deless. Ic. Sel. iii. 58. t. 96 = Paepalanthus microphyllus.
minimum, *Lam. Encyc.* iii. 275.—Zeylan.
minutulum, Steud. Syn. Pl. Cyp. 270 = Paepalanthus pusillus.
Miquelianum, *Koern. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 169.—Japon.
miserum, *Koern. in Linnæa*, xxvii. (1854) 607.—Ind. or.
modestum, *Kunth, Enum. Pl.* iii. 547.—Bras.
molle, Mart. ex Koern. in Mart. Fl. Bras. iii. i. 487 = crassiscapum.
molle, Steud. Syn. Pl. Cyp. 282 = Paepalanthus plumosus.
monocaspum, *F. Muell. Fragm.* i. 94.—Austral.
monticolum, Steud. Syn. Pl. Cyp. 278 = Paepalanthus monticolus.
myrtulum, Steud. l. c. 281 = Paepalanthus chrysanthus.
mucronatum, Boug. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 636. t. 10 = Paepalanthus mucronatus.
myocephalum, Mart. in Flora, xxiv. (1841) ii. Beibl. 60 = Paepalanthus myocephalus.
nanum, *R. Br. Prod.* 254.—Austral.
nardifolium, Steud. Syn. Pl. Cyp. 281 = Paepalanthus Fischerianus.
Nessianum, *Koern. in Linnæa*, xxvii. (1854) 622.—Zeylan.

ERIOCAULON :—
nepalense, *Bong. in Mém. Acad. Pétersb.* Sér. VI. i. (1841) 610.—Nepal.
nepalense, Kunth, Enum. Pl. iii. 554 = viride.
nigrescens, D. Dietr. Syn. Pl. v. 265 = melaleucum.
nigricans, *R. Br. Prod.* 254.—Austral.
nigrovirescum, St. Hil. Voy. Distr. Diam. i. 392 = Paepalanthus tortilis.
nilagiricum, *Steud. Syn. Pl. Cyp.* 271.—Ind. or.
nitens, Bong, in Mém. Acad. Pétersb. Sér. VI. i. (1831) 633 = Paepalanthus nitens.
nitidum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 636 = Paepalanthus nitidus.
nitidum, Buch.-Ham. ex Wall. Cat. n. 6073 = sexangulare.
niveum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 635. t. 37 = Paepalanthus niveus.
niveum, Hoffmgg. ex Koern. in Mart. Fl. Bras. iii. i. 304 = Philodice Hoffmannseggii.
nubigenum, Steud. Syn. Pl. Cyp. 281 = Paepalanthus nubigenus.
obtusifolium, Steud. l. c. 277 = Paepalanthus obtusifolius.
odoratum, *Dals. in Hook. Kew Journ.* iii. (1851) 280. —Ind. or.
officinale, *Koern. in Mart. Fl. Bras.* iii. i. 480, in nota. —China.
oryzetorum, *Mart. in Wall. Pl. As. Rar.* iii. 28.—Ind. or.
pallidum, *R. Br. Prod.* 254.—Austral.
paludosum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 625 = Paepalanthus plantagineus.
palustre, *Salzm. ex Steud. Syn. Pl. Cyp.* 280.—Am. trop.
papillosum, *Koern. in Mart. Fl. Bras.* iii. i. 489.—Bras. paraguayense, *Koern. l. c.* 497.—Paraguay.
pardinum, D. Dietr. Syn. Pl. v. 263 = elichrysoides.
parvum, *Koern. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 168.—Japon.
pedunculatum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 630 = Paepalanthus falcatus
pellucidum, Michx. Fl. Bor. Am. ii. 166 = septangulare.
perpusillum, Steud. Syn. Pl. Cyp. 277 = Paepalanthus tortilis.
pilyferum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 631 = Paepalanthus pilifer.
pilosum, H. B. & K. Nov. Gen. et Sp. i. 251 = Paepalanthus pilosus.
planifolium, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 629 = Paepalanthus monticolus.
Plantaginella, Crants, Inst. i. 360 = Paepalanthus triangularis.
plantaginoides, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 625 = Paepalanthus zosteraefolius.
plantaginoides, Desv. in Ham. Prod. Pl. Ind. Occ. 16 = Paepalanthus plantaginoides.
plumosum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 632. t. 62 = Paepalanthus plumosus.
polyanthum, Bong. l. c. 632 = Paepalanthus polyanthus.
polyanthum, Mart. in Flora, xxiv. (1841) ii. Beibl. 35 = Paepalanthus Claussenianus.
polyphyllum, Salzm. ex Steud. Syn. Pl. Cyp. 277 = Paepalanthus tortilis.
polytrichoides, Steud. l. c. 275 = Paepalanthus polytrichoides.
Prescottianum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 635. t. 36 = Mesanthemum Prescottianum.
proliferum, Bong. l. c. 632. t. 28 = Paepalanthus prolifer.
proliferum, Hochst. ex Steud. Syn. Pl. Cyp. 277 = Paepalanthus viviparus.
pruinum, Steud. Syn. Pl. Cyp. 280 = modestum.
pubescens, *Lam. Encyc.* iii. 276.—Madag.
pubigerum, *Bong. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 628. t. 42.—Bras.
pulchellum, *Koern. in Linnaea*, xxvii. (1854) 632.—Afr. trop.
pumilum, *Afzel. ex Koern. l. c.* 621.—Afr. trop.
pumilum, Rafin. Atl. journ. 121 = septangulare ?
pusillum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 634. t. 30 = Paepalanthus pusillus.
pusillum, *R. Br. Prod.* 254.—Austral.
pusillum, Poepp. ex Koern. in Mart. Fl. Bras. iii. i. 356 = Paepalanthus tortilis.
pusillum, Willd. ex Kunth, Enum. Pl. iii. 546 = microcephalum.
Pygmaeum, Dals. in Hook. Kew Journ. iii. (1851) 281 = E. Xeranthemum ?
pygmaeum, *Koern. in Mart. Fl. Bras.* iii. i. 477.—Bras.

ERIOCAULON :—
pygmaeum, Mart. in Flora, xxiv. (1841) ii. Beibl. 60 = Paepalanthus Schraderi.
quadrangulare, *Lour. Fl. Cochinch.* 60.—Cochinch.
quinquangulare, Boj. Hort. Maurit. 361 = Paepalanthus repens.
quinquangulare, Heyne, ex Mart. in Wall. Pl. As. Rar. iii. 29 = melaleucum.
quinquangulare, *Linn. Sp. Pl.* 87.—Ind. or. ; Austral.
quinquangulare, Mart. ex Koern. in Linnaea, xxvii. (1854) 645 = trilobum.
radians, *Benth. in Hook. Niger Fl.* 547.—Afr. trop.
ramosum, Wikstr. in Vet. Acad. Handl. Stockh. (1820) 76. t. 3 = Paepalanthus ramosus.
Ravenelii, *Chapm. Fl. S. U. St.* 508.—Carolina.
repens, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 623, 648. t. 7 = Paepalanthus bongardi.
repens, Lam. Encyc. iii. 275. t. 50. f. 2 = Paepalanthus repens.
Riedelianum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 630. t. 34 = Paepalanthus Vauthierianus.
rigidulum, Steud. Syn. Pl. Cyp. 278 = Paepalanthus rigidulus.
rigidum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 621 = Paepalanthus rigidus.
rigidum, Mart. in Flora, xxiv. (1841) ii. Beibl. 35 = Paepalanthus falcifolius. Steud. Syn. Pl. Cyp. 270.—Ind. or.
rivulare, Dals. in Hook. Kew Journ. iii. (1851) 280 = Dalzellii.
rivulare, *G. Don, ex Benth. in Hook, Niger Fl.* 547. —Afr. trop.
robustum, *Steud. Syn. Pl. Cyp.* 271.—Ind. or.
rosulatum, *Koern. in Linnaea*, xxvii. (1854) 600, nomen; *et in Mart. Fl. Bras.* iii. i. 487.—Bras.
Rouxianum, *Steud. Syn. Pl. Cyp.* 270.—Ind. or.
rufulum, St. Hil. Voy. Distr. Diam. i. 391 = Paepalanthus rufulus.
rupestre, Steud. Syn. Pl. Cyp. 278 = Paepalanthus rupestris.
saxatile, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 626. t. 30 = Paepalanthus saxatilis.
Scabiosa, Crants, Inst. i. 360 = quinquangulare.
scariosum, *R. Br. Prod.* 255.—Austral
Schiedeanum, *Koern. in Linnaea*, xxvii. (1854) 600, nomen; *et in Mart. Fl. Bras.* iii. i. 492.—Mexic.
Schukxii, *Benth. Fl. Austral.* vii. 195.—Austral.
scirpoides, *Griseb. Cat. Pl. Cub.* 226.—Cuba.
Sieberianus, Benth. Pl. Hartw. 260 = Paepalanthus sclaginoides.
Sellowianum, *Kunth, Enum. Pl.* iii. 545.—Bras.
separatum, Steud. Syn. Pl. Cyp. 280 = Paepalanthus hirsutus.
septangulare, *With. Bot. Arr.* ed. III. ii. 184.—Reg. Brit. ; Am. bor.
sericeum, Mart. in Wall. Pl. As. Rar. iii. 29 = Wightianum.
serotinum, Walt. Fl. Carol. 88 = decangulare.
sessiliflorum, Mart. in Flora, xxiv. (1841) ii. Beibl. 60 = Paepalanthus sessiliflorus.
setaceum, Heyne, ex Wall. Cat. n. 6073 = sexangulare.
setaceum, *Linn. Sp. Pl.* 87.—Ax. trop. ; Austral.
setaceum, *Lour. Fl. Cochinch.* 60.—Cochinch.
sexangulare, *Linn. Sp. Pl.* 87.—Ind. or. ; Madag.
Sieboldianum, Sieb. & Zucc. ex Steud. Syn. Pl. Cyp. 273 (Sieboldtianum) = sexangulare.
sigmoideum, *Wright, in Sauv. Fl. Cub.* 168.—Cuba.
similimum, Steud. Syn. Pl. Cyp. 277 = Paepalanthus caniescens.
simplex, Steud. l. c. 280 = Paepalanthus simplex.
sinicum, *Miq. in Journ. Bot. Néerl.* i. (1861) 87.—China.
Smithii, *R. Br. Prod.* 254.—Austral.
Soldanella, Royle, Illustr. Bot. Himal. 409. t. 97. f. 1 = trilobum.
Sonderianum, *Koern. in Linnaea*, xxvii. (1854) 669.—Afr. austr.
spadiceum, Lam. Encyc. iii. 277 = Scirpus spadiceus.
sparganioides, *Bong. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 627. t. 58.—Bras.
spathaceum, *Rafin. Atl. Journ.* 121 (Quid ?).—Am. bor.
speciosum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 636. t. 57 = Paepalanthus speciosus.
spectabile, *F. Muell. Fragm.* i. 96.—Austral.
sphaeralbum, Willd. ex Kunth, Enum. Pl. ii. 206 (= (sperospermum).—Cuba.
sphagnoides, Poepp. ex Koern. in Mart. Fl. Bras. iii. i. 302 = Torresia fluviatilis.

ERIOCAULON :—
spirale, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 634. t. 45 = Paepalanthus spiralis.
Spixianum, Steud. Nom. ed. II. i. 586 = Paepalanthus Spixianus.
splendens, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 633. t. 66 = Paepalanthus caulescens.
Spruceanum, *Koern. in Mart. Fl. Bras.* iii. i. 488.—Bras.
Statices, Crants, Inst. i. 360 = decangulare.
stellare, Guill. in Deless. Ic. Sel. iii. 20 = Paepalanthus plumosus.
stellulatum, *Koern. in Linnaea*, xxvii. (1854) 620.—Ind. or.
stramineum, *Koern. in Mart. Fl. Bras.* iii. i. 478.—Bras.
striatum, *Lam. Encyc.* iii. 275.—Ins. Borbon.
subalatum, Boj. ex Koern. in Linnaea, xxvii. (1854) 645 = trilobum.
subtile, Steud. Syn. Pl. Cyp. 281 = Paepalanthus subtilis.
subuncinatum, Steud. l. c. 277 = Paepalanthus caulescens.
supinum, Herb. Vind. ex Koern. in Mart. Fl. Bras. iii. i. 352 = Paepalanthus supinus.
surinamense, Steud. Syn. Pl. Cyp. 275 = Paepalanthus caulescens.
tenue, Buch.-Ham. ex Wall. Cat. n. 6073 = sexangulare.
tenue, H. B. & K. Nov. Gen. et Sp. i. 253 = Paepalanthus tenuis.
tenue, Poepp. ex Koern. in Mart. Fl. Bras. iii. i. 359 = Paepalanthus polytrichoides.
tenuifolium, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1116, nomen; *et Koern. in Mart. Fl. Bras.* iii. i. 496.—Guiana.
texense, *Koern. in Linnaea*, xxvii. (1854) 594.—Am. bor.
Thunbergii, *Wikstr. ex Koern. in Linnaea*, xxvii. (1854) 677.—Afr. trop.
Thwaitesii, *Koern. l. c.* 627.—Zeylan.
tortile, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 624. t. 49 = Paepalanthus tortilis.
tortile, Steud. Syn. Pl. Cyp. 275 = Paepalanthus flaccidus.
tortuosum, *F. Muell. Fragm.* i. 91.—Austral.
triangulare, Linn. Sp. Pl. ed. II. 128 = Paepalanthus triangularis.
trichophyllum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 636. t. 54 = Paepalanthus trichophyllus.
trichosepalum, *Wright, in Sauv. Fl. Cub.* 161.—Cuba.
trilobum, *Buch-Ham. ex Wall. Cat.* n. 6072 D.—Ind. or.
Trinianum, Mart. in Flora, xxiv. (1841) ii. Beibl. 58 = Paepalanthus rufulus.
truncatum, *Buch-Ham. ex Mart. in Wall. Pl. As. Rar.* iii. 29.—Ind. or.
tuberum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 629 = Paepalanthus tuberosus.
umbellatum, Bong. l. c. 633 = Paepalanthus helminthorrhizus.
umbellatum, H. B. & K. Nov. Gen. et Sp. i. 252 = Paepalanthus Humboldtii.
umbellatum, Lam. Encyc. iii. 277 = Paepalanthus umbellatus.
umbrosum, Salzm. ex Koern. in Mart. Fl. Bras. iii. i. 357 = Paepalanthus myocephalus.
uncinatum, Steud. Syn. Pl. Cyp. 281 = Paepalanthus uncinatus.
usuuriense, *Koern. ex Regel, Tent. Fl. Ussur.* 157.—Manchur.
vaginatum, *Koern. in Linnaea*, xxvii. (1854) 599, nomen; *et in Mart. Fl. Bras.* iii. i. 483.—Bras.
Vanheurckii, *Muell. Arg. in Heurck, Obs. Bot.* 98.—Ind. or.
Vauthierianus, Guill. in Deless. Ic. Sel. iii. 57 = Paepalanthus Vauthierianus.
vermiculoides, Steud. Syn. Pl. Cyp. 280 = Paepalanthus vermonioides.
verticillatum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 638 = Paepalanthus verticillatus.
villosum, Michx. Fl. Bor. Am. ii. 196 = Lachnocaulon Michauxii.
villosum, Salzm. ex Steud. Syn. Pl. Cyp. 277 = Paepalanthus Schraderi.
villosum, Willd. ex Koern. in Linnaea, xxvii. (1854) 596 = decangulare.
viride, *Koern. l. c.* 637.—Ind. or.
viviparum, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 632. t. 28 = Paepalanthus viviparus.
Wahlbergii, Wikstr. ex Koern. in Mart. Fl. Bras. iii. i. 480 = Paepalanthus Wahlbergii.
Wallichianum, *Mart. in Wall. Pl. As. Rar.* iii. 26. t. 249.—Ind. or.

ERIOCAULON:—
Wallichianum, Thw. Enum. Pl. Zeyl. 341 = sexangulare.
Wightianum, *Mart. in Wall. Pl. As. Rar.* iii. 98.—Ind. or.
xeranthemoides, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 685 = Paepalanthus xeranthemoides.
xeranthemoides, *Henrck & Muell. Arg. in Henrck, Obs. Bot.* 108.—Afr. trop.
Xeranthemum, *Mart. in Wall. Pl. As. Rar.* iii. 99.—Ind. or.
yuncæfolium, Mertens, ex Mart. in Flora, xxiv. (1841) ii. Beibl 80 = Paepalanthus tortilis.
Zollingerianum, *Koern. in Linnæa*, xxvii. (1854) 682.—Java.
zosterifolius, Steud. Syn. Pl. Cyp. 277 = Paepalanthus zosteræfolius

ERIOCEPHALUS, Linn. Gen. ed. I. 265 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 416.
BRACHYGYNE, MICROGYNE, SPHONOGYNE, STENOGYNE, Cass. in Dict. Sc. Nat. i. 498 (1827).
CRYPTOGYNE, Cass. l. c. 491 (1827).
MONOCHLAENA, Cass. l. c. 496 (1827).
affinis, DC. Prod. vi. 147 = racemosus.
africanus, Burm. f. Prod. Fl. Cap. 25 = sericeus.
africanus, *Linn. Sp. Pl.* 926.—Afr. austr.
africanus, Sieber, ex DC. Prod. vi. 147 = umbellulatus.
aspalathoides, *DC. Prod.* vi. 146.—Afr. austr.
capitellatus, *DC. l. c.* 146.—Afr. austr.
corymbosus, Moench, Meth. 590 = africanus.
decussatus, Burch. Trav. i. 272 = glaber.
eximius, *DC. Prod.* vi. 147.—Afr. austr.
frutescens, R. Br. in Ait. Hort. Kew. ed. II. v. 180 = africanus.
glaber, *Thunb. Prod. Pl. Cap.* 168.—Afr. austr.
laricinus, DC. Prod. vi. 148 = glaber.
microcephalus, *DC. l. c.*—Afr. austr.
microphyllus, DC. l. c. = glaber.
paniculatus, Cass. in Dict. Sc. Nat. i. 496 = umbellulatus.
pectinifolius, Linn. Syst. ed. XII. 579 = Hippia frutescens.
pedicellaris, DC. Prod. vi. 146 = punctulatus.
pteronioides, *DC. l. c.*—Afr. austr.
pteronoides, Sch. Bip. in Flora, xxvii. (1844) 676 = punctulatus.
pteraphilioides, *DC. Prod.* vi. 146.—Afr. austr.
pubescens, *DC. l. c.* 148.—Afr. austr.
punctulatus, *DC. l. c.* 146.—Afr. austr.
purpureus, Burch. ex G. Don, in Loud. Hort. Brit. 305 = glaber.
racemosus, Gaertn. Fruct. ii. t. 168 = umbellulatus.
racemosus, *Linn. Pl. Rar. Afr.* 25.—Afr. austr.
racemosus, Zeyh. ex Steud. Nom. ed. II. i. 586 = Elytropappus glandulosus.
scariosus, *DC. Prod.* vi. 147.—Afr. austr.
septifer, Cass. in Dict. Sc. Nat. i. 494 = africanus.
septulifer, *DC. Prod.* vi. 125.—Afr. austr.
sericeus, *Gaudich. ex DC. l. c.*—Afr. austr.
sessiliflorus, Sond. in Harv. & Sond. Fl. Cap. iii. 204 = glaber.
simplicifolius, Salisb. Prod. 211 = racemosus.
spicatus, Burm. ex DC. Prod. vi. 147 = racemosus.
spinescens, *Burch. Trav.* i. 272.—Afr. austr.
tenuifolius, *DC. Prod.* vi. 146.—Afr. austr.
tuberculosus, *DC. l. c.* 146.—Afr. austr.
umbellatus, Auct. = seq.
umbellulatus, *Cass. in Dict. Sc. Nat.* i. 493.—Afr. austr.
varii folius, Salisb. Prod. 211 = africanus.

ERIOCHAENIUM, C. Muell. in Walp. Ann. v. 849 (1858) = **Eriachaenium**, Sch. Bip. (Compos.).

ERIOCHAETA, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xiv. (1854) 874 = **Pennisetum**, Rich. (Gramin.).
densiflora, Fig. & De Not. l. c. 876.—Nubia.
reversa, Fig. & De Not. l. c. 878.—Nubia.
secundiflora, Fig. & De Not. l. c. 875.—Nubia.

ERIOCHAETA, Torr. ex Steud. Nom. ed. II. i. 586 (1840) = **Rynchospora**, Vahl (Cyperac.).

ERIOCHILOS, Spreng. Syst. iii. 714 (1826) = seq.

ERIOCHILUM, Ritg. in Marb. Schrift. ii. (1831) 121 = seq.

ERIOCHILUS, R. Br. Prod. 328 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 611.
autumnalis, *R. Br. l. c.*—Austral.

ERIOCHILUS:—
cucullatus, Reichb. f. Beitr. Syst. Pfl. 27 = autumnalis.
dilatatus, *Lindl. Swan River App.* 53.—Austral.
fimbriatus, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 192.—Austral.
latifolius, *Lindl. Swan River App.* 53.—Austral.
Lindleyi, Endl. in Lehm. Pl. Preiss. ii. 10 = scaber.
multiflorus, *Lindl. Swan River App.* 53.—Austral.
scaber, *Lindl. l. c.*—Austral.
tenuis, *Lindl. l. c.*—Austral.

ERIOCHITON, F. Muell. 2nd Gen. Report, 15 (1854), nomen = **Chenolea**, Thunb. (Chenopod.).
sclerolaenoides, F. Muell. l. c. = C. sclerolaenoides.

ERIOCHLAENA, Spreng. Syst. iii. 128 (1826) = **Eriolaena**, DC. (Sterculiac.).

ERIOCHLAMYS, Sond. et F. Muell. in Linnaea, xxv. (1852) 488. *COMPOSITAE*, Benth. & Hook. f. ii. 317.
Behrii, *Sond. & F. Muell. l. c.*—Austral.
Knappii, *F. Muell. in Melb. Chemist* (May 1888).—Austral.

ERIOCHLOA, H. B. & K. Nov. Gen. et Sp. i. 94. t. 30, 31 (1815). *GRAMINEAE*, Benth. & Hook. f. iii. 1089.
AGLYCIA, Willd. ex Steud. Nom. ed. II. i. 37 (1840).
ALYCIA, Willd. ex Steud. l. c. 66 (1840).
HEXOPUS, Trin. Fund. Agrost. 100 (1820).
MONACHNE, Beauv. Agrost. 49. t. 10 (1812), nomen prius.
ORDIOPHNE, Link, Hort. Berol. i. 51 (1827).
acuminata, Kunth, Enum. Pl. i. 72 = polystachya.
annulata, *Kunth, Rév. Gram.* i. 30.—Reg. trop.
brasiliensis, Spreng. Syst. i. 249 = Paspalum dissitiflorum.
distachya, *H. B. & K. Nov. Gen. et Sp.* i. 95.—Bras.
globosa, Kunth, Rév. Gram. i. 30 = Milium globosum.
grandiflora, *Benth. in Journ. Linn. Soc.* xix. (1881) 89.—Am. trop.
grandiflora, Vasey, Gr. U. St. 11 = Lemmoni.
Kunthii, *G. F. W. Mey. Prim. Fl. Esseq.* 47.—Am. austr.
Lemmoni, *Vasey & Scribner, in Coult. Bot. Gaz.* ix. (1883) 185.—Arizona.
mollis, Kunth, Rév. Gram. i. 30 = punctata.
polystachya, *H. B. & K. Nov. Gen. et Sp.* i. 95.—Mexic.
pulchella, Kunth, Rév. Gram. i. 30 = Panicum pulchellum.
punctata, *Ham. Prod. Pl. Ind. Occ.* 5.—As. trop.; Austral.
punctata, Hochst. ex Steud. Syn. Pl. Gram. 101 = polystachya.
purpurascens, Hochst. in Flora, xxiv. (1841) i. Intell. 19 = Panicum gossypinum, *A. Rich.*
sericea, *Munro, ex Vasey, Gr. U. St.* 11, nomen.—Texas.
succincta, *Kunth, Enum. Pl.* i. 73.—Rossia.
sundaica, *Miq. Fl. Ind. Bat. Suppl.* 606.—Ins. Banca.
trichopus, *Hochst. ex Benth. in Journ. Linn. Soc.* xix. (1881) 89.—Afr. trop.
vestita, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 97.—Ins. Socotra.
villosa, *Kunth, Rév. Gram.* i. 28. t. 18.—Aa. or.

ERIOCHRYSIS, Beauv. Agrost. 8. t. 4. f. 11 (1812) = **Saccharum**, Linn. (Gramin.).
attenuata, Nees, ex Steud. Syn. Pl. Gram. 411 = S. fuscum.
copaynala, Beauv. Agrost. 8. t. 4. f. 11 = S. cayennense.
fusca, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 315 = S. fuscum.
longifolia, Munro, ex Benth. in Journ. Linn. Soc. xix. (1881) 66.—Ind. or.
Narenga, Nees, ex Steud. Syn. Pl. Gram. 411 = S. Narenga.
pallida, Munro, in Harv. Gen. S. Afr. Pl. ed. II. 440.—Afr. austr.
pulchra, Beauv. Agrost. 162 = S. cayennense.
porphyrocoma, Hance, in Journ. Bot. xiv. (1876) 294.—China.

ERIOCHYLUS, Steud. Nom. ed. II. i. 586 (1840) = **Brachylus**, R. Br. (Orchid.).

ERIOCLADIUM, Lindl. Swan River App. 24 (1839) = **Anglanthus**, Wendl. (Compos.).
pyramidatum, Lindl. l. c. = A. Cunninghamii.

ERIOCLEPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 95 = ERIOLEPIS, Cass. = **Onicus**, Linn. (Compos.).

ERIOCLINE, Cass. in Dict. Sc. Nat. xv. 191 (1819) = **Osteospermum**, Linn. (Compos.).
obovata, Cass. l. c.; et xxx. 383 = O. spinosum.

ERIOCNEMA, Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 144. *MELASTOMACEAE*, Benth. & Hook. f. i. 736.
acaulis, Triana, in Trans. Linn. Soc. xxviii. (1871) 23 t. 1. f. 2 = Hilariana.
aenea, Naud. in Rev. Hortic. Sér. III. ii. (1848) 381 = Bertolonia marmorata.
fulva, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 144.—Bras.
Hilariana, *Naud. l. c.* 145.—Bras.
marmorata, Naud. in Rev. Hortic. Sér. III. ii. (1848) 381 = Bertolonia marmorata.

ERIOCOCCUS, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 143 = **Phyllanthus**, Linn. (Euphorb.).
caudatus, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 178.—Java.
glaucescens, Zoll. l. c. = P. pulcher.
gracilis, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 143 = P. gracilipes.

ERIOCOELUM, Hook. f. in Benth. & Hook. f. Gen. i. 400 (1862). *SAPINDACEAE*, Benth. & Hook. f. i. c.
paniculatum, *Baker, in Oliver, Fl. Trop. Afr.* i. 428.—Afr. trop.
racemosum, *Baker, l. c.* 427.—Afr. trop.

ERIOCOMA, H. B. & K. Nov. Gen. et Sp. iv. 267. t. 396 (1820) = **Montanoa**, La Llave (Compos.).
arborescens, Alam. ex DC. Prod. v. 566 = M. arborescens.
floribunda, H. B. & K. Nov. Gen. et Sp. iv. 267. t. 396 = M. floribunda.
fragrans, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 44 = M. tomentosa.
frutescens, Alam. ex DC. Prod. v. 565 = M. frutescens.
grandiflora, Alam. ex DC. l. c. = M. grandiflora.
heterophylla, Schrad. Sem. Hort. Gotting. (1838) 8 = M. tomentosa.

ERIOCOMA, Natt. Gen. Am. i. 40 (1818) = **Oryngopsis**, Michx. (Gramin.).
cuspidata, Natt. l. c. = O. cuspidata.
mexicana, Auct. ex Steud. Nom. ed. II. i. 586.—Am. bor.
Webberi, Thurb. in S. Wats. Bot. Calif. ii. 283 = O Webberi.

ERIOCORYNA, Steud. Nom. ed. II. i. 586 (1840) = seq

ERIOCORYNE, Wall. ex DC. Prod. vi. 541 (1837) = **Saussurea**, DC. (Compos.).
nidularia, Wall. ex DC. l. c. = S. gossypiphora.

ERIOCYCLA, Lindl. in Royle, Illustr. Bot. Himal. 282 (1885) = **Pituranthos**, Viv. (Umbellif.).
nuda, Lindl. l. c. = P. nuda.

ERIOCYCLAX, Neck. Elem. iii. 25 (1790) = **Aspalathus**, Linn. (Legumin.).

ERIODAPHUS, Spach, Hist. Vég. Phan. vi. 286 (1838) = ERIUDAPHUS, Nees, = **Scoliopia**, Schreb. (Bixin.).

ERIODENDRON, DC. Prod. i. 479 (1824). *MALVACEAE*, Benth. & Hook. f. i. 210.
CAMPYLANTHERA, Schott & Endl. Melet. 25 (1832).
CEIBA, Medic. Malv. 15 (1787); Gaertn. Fruct. ii. 244. t. 128 (1791).
ERIONE, Schott & Endl. Melet. 24 (1832).
GOSSAMPINUS, Buch.-Ham. in Trans. Linn. Soc. xv. (1826) 156.
anceolifolium, *DC. Prod.* i. 479.—Mexic.
anfractuosum, *DC. l. c.*—As. et Afr. trop.
caribaeum, G. Don, in Loud. Hort. Brit. 292 = anfractuosum.

ERIODENDRON:—
guineense, G. Don, in Loud. Hort. Brit. 292 = anfractuosum.
jasminiodorum, *St. Hil. Fl. Bras. Mer.* i. 265.—Bras.
leianthemum, *DC. Prod.* i. 479.—Bras.
occidentale, *G. Don, Gen. Syst.* i. 518.—Mexic.
orientale, Kostel. Allg. Med.-Pharm. Fl. v. 1875 = anfractuosum.
pentandrum, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 113 = anfractuosum.
pubiflorum, *St. Hil. Fl. Bras. Mer.* i. 266.—Bras.
rhaseoanthum, *Decne. ex Kunth*, *in Journ. Soc. Hortic. Fr.* Sér. II. iv. (1870) 90.—Bras.†
Riviéri, *Decne. in Fl. des Serres*, Sér. II. xii. (1877) 167.—Bras.
Samauma, *Mart. Nov. Gen. et Sp.* i. 91.—Bras.
trischistandrum, *A. Gray, Bot. U. St. Expl. Exped.* i. 192.—Peruv.†

ERIODESMIA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 156 = **Erica**, Tourn.
capitata, D. Don, l. c. 158 = Erica capitata.

ERIODICTYON, Benth. Bot. Voy. Sulph. 35 (1844).
HYDROPHYLLACEÆ, Benth. & Hook. f. ii. 881.
ERIODYCTON, Benth. in DC. Prod. x. 133 (1846).
angustifolium, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 181.—Am. bor. occ.
californicum, Decne. Bot. Voy. Vénus, 22 = crassifolium.
crassifolium, *Benth. Bot. Voy. Sulph.* 35.—Calif.
glutinosum, *Benth. l. c.* 36.—Calif.
Lobbii, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 202.—Calif.
sessilifolium, *Greene, l. c.* 201.—Calif.
tomentosum, *Benth. Bot. Voy. Sulph.* 36.—Calif.

ERIODRYS, Rafin. Alsog. Am. 30 (1838) = **Quercus**, Tourn. (Cupulif.).
lanata, Rafin. l. c. 30 = Q. lanuginosa†

ERIODYCTION, Benth. in DC. Prod. x. 133 (1846) = **Eriodictyon**, Benth. (Hydrophyll.).

ERIOGENIA, Steud. Nom. ed. II. i. 587 (1840) = ERIOGYNIA, Hook. = **Spiræa**, Linn. (Rosac.).

ERIOGLOSSUM, Blume, Bijdr. 229 (1829). *SAPINDACEÆ*, Benth. & Hook. f. i. 396.
MOULINSIA, Cambess. in Mém. Mus. Par. xviii. (1829) 37. t. 3.
PANCOVIA, Willd. Sp. Pl. ii. 385 (1799).
UITENIA, Noronha, in Verh. Bat. Gen. v. (1790) ed. I. Ari. iv. 4.
alliaceum, *Zipp. ex Span. in Linnæa*, xv. (1841) 181.—Ins. Timor.
caulifloram, *Guill. & Perr. Fl. Senieg. Tent.* i. 118. t. 26.—Afr. trop.
edule, *Blume, Bijdr.* 229.—As. trop.
membranifolium, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 193.—Malaya.
rubiginosum, Brand. For. Fl. 106 = edule.

ERIOGONUM, Michx. Fl. Bor. Am. i. 246. t. 24 (1803). *POLYGONACEÆ*, Benth. & Hook. f. iii. 92.
EPINOSA, Lag. Gen. et Sp. Nov. 14 (1816).
EUCYCLA, Nutt. in Journ. Acad. Philad. N. S. i. 166 (1847).
STENOGONUM, Nutt. l. c. 170 (1847).
Abertzianum, *Torr. in Bot. Emory Rep.* 150.—N. Mexic.
acaule, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 160.—Am. bor. occ.
acetosoides, Torr. ex Benth. in DC. Prod. xiv. 16 = gracile.
æstivum, Benth. l. c. 13 = undum.
alatum, *Torr. in Stkpr. Rep.* 168. t. 8.—Am. bor. occ.
album, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 164 = dichotomum.
alpinum, *Bongard. in Coult. Bot. Gaz.* vii. (1882) 6.—Am. bor.
ambiguum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 160 = cæspitosum.
androsaceum, Benth. in DC. Prod. xiv. 9 = cæspitosum.
angulosum, *Benth. in Trans. Linn. Soc.* xvii. (1837) 406.—Am. bor. occ.
angustifolium, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 164 = hersleioides.
annuum, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 164. —Am. bor. occ.
apiculatum, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 378.—Am. bor.
arachnoideum, Esch. in Mém. Acad. Pétersb. (1821–22) 287 = latifolium.
arachnoideum, Hook. & Arn. Bot. Beech. Voy. 158 = undum.
arborescens, *Greene, in Bull. Calif. Acad.* i. i. (1884) 11.—Am. bor. occ.
atrorubens, *Engelm. Pl. Wisliz.* 24.—N. Mexic.
aureum, Nutt. ex Benth. in DC. Prod. xiv. 8 = flavum.
auriculatum, Benth. in Trans. Linn. Soc. xvii. (1837) 413 = undum.
Baileyi, *S. Wats. in Proc. Am. Acad.* x. (1875) 346.—Am. bor. occ.
brachypodium, *Torr. & Gray, l. c.* viii. (1870) 180.—Calif.
breviscale, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 163.—Am. bor. occ.
cæspitosum, *Nutt. in Journ. Acad. Philad.* vii. (1834) 50. t. 8. f. 2.—Am. bor. occ.
californicum, *A. Murr. Oreg. Circ.* 1.—Am. bor. occ.
campanulatum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 163 = breviscale.
cernuum, *Nutt. l. c.* 162.—Am. bor. occ.
chrysocomum, *A. Gray, in Proc. Am. Acad.* xi. (1876) 101.—Am. bor. occ.
ciliatum, *Torr. ex Benth. in DC. Prod.* xiv. 20.—N. Mexic.
cinereum, *Benth. Bot. Voy. Sulph.* 45.—Calif.
compositum, *Dougl. ex Benth. in Bot. Reg.* t. 1774 (1836).—Am. bor. occ.
convertifolium, Benth. in DC. Prod. xiv. 17 = microthecum.
cordatum, Torr. ex Benth. l. c. 21 = inflatum.
corymbosum, *Benth. l. c.* 17.—Am. bor. occ.
crassifolium, Benth. in Trans. Linn. Soc. xvii. (1837) 408 = flavum.
cymosum, Benth. in DC. Prod. xiv. 19 = sinuum.
dasyanthemum, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 177.—Calif.
decumbens, Benth. in Trans. Linn. Soc. xvii. (1837) 413 = sinuum.
deflexum, *Torr. in Ives, Rep. Colorado River, Bot.* 24. —Am. bor. occ.
dellatulum, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 379.—Am. bor.
denudatum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 161.—Calif.
dichotomum, *Dougl. ex Benth. in Trans. Linn. Soc.* xvii. (1837) 415.—Am. bor. occ.
dioicum, *Rafin. New. Fl. Am.* iv. 53.—Am. bor.
divaricatum, *Hook. Kew Journ.* v. (1853) 265.—Am. bor. occ.
Douglasii, *Benth. in DC. Prod.* xiv. 9.—Am. bor. occ.
effusum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 164 = microthecum.
elatum, *Dougl. ex Benth. in Trans. Linn. Soc.* xvii. (1837) 413.—Am. bor. occ.
elliptieum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 161 = umbellatum.
elongatum, *Benth. Bot. Voy. Sulph.* 45.—Calif.
elongatum, Nutt. ex Benth. in DC. Prod. xiv. 10 = ovalifolium.
elongatum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 168 = virgatum.
eriophyllum, A. Gray, in Proc. Am. Acad. viii. (1870) 170 = fasciculatum.
fasciculatum, *Benth. in Trans. Linn. Soc.* xvii. (1837) 411.—Am. bor. occ.
flavum, *Nutt. in Fras. Cat.*; *Benth. in Trans. Linn. Soc.* xvii. (1837) 408.—Am. bor. occ.
foliosum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 371. —Calif.
Fremonti, Torr. ex Benth. in DC. Prod. xiv. 18 = breviscale.
giriculatum, Dur. & Hilg. in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1856) 48 = Hoermanni.
gracilentatum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 163 = sphærocephalum.
glandulosum, *Nutt. ex Benth. in DC. Prod.* xiv. 21.—Am. bor. occ.
gnaphalodes, Benth. in Hook. Kew Journ. v. (1853) 268 = multiceps.
Gordonii, *Benth. in DC. Prod.* xiv. 20.—Am. bor. occ.
gossypinum, *Curran, in Bull. Calif. Acad.* i. iv. (1885) 274.—Calif.
gracile, *Benth. Bot. Voy. Sulph.* 46.—Calif.
gracillimum, *S. Wats. in Bot. Calif.* ii. 480.—Calif.
Greenei, A. Gray, in Proc. Am. Acad. xii. (1877) 83 = dichotosum.
Greggii, *Torr. & Gray, l. c.* viii. (1870) 187.—Mexic.
gypophyllum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 163 = hersleioides.
Havardi, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 194.—Am. bor.
Heermanni, *Dur. & Hilg. in Pacif. Rail. Rep.* v. 14. t. 17.—Am. bor. occ.
helianthemifolium, Benth. in DC. Prod. xiv. 15 = polyciadon, Wrightii.
hemisteum, Torr. ex Torr. & Gray, in Proc. Am. Acad. viii. (1870) 184 = hieraciifolium.
hersleioides, *Nutt. in Journ. Acad. Philad.* vii. (1834) 49.—Am. bor. occ.
hieraciifolium, *Benth. in DC. Prod* xiv. 6.—N. Mexic.
hirtiflorum, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xii. (1877) 259.—Calif.
Hookeri, *S. Wats. l. c.* xiv. (1879) 295.—Am. bor. occ.
incanum, *Torr. & Gray, l. c.* viii. (1870) 161.—Calif.
infatum, *Torr. in Frémont, Rep.* ii. 317.—Am. bor. occ.
insigne, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 295. —Am. bor. occ.
intricatum, *Benth. Bot. Voy. Sulph.* 46. t. 22.—Calif.
Jamesii, *Benth. in DC. Prod.* xiv. 7.—Am. bor. occ.
Kelloggii, *A. Gray, in Proc. Am. Acad.* viii. (1870) 295.—Calif.
Kennedyi, *Porter, ex S. Wats. l. c.* xii. (1877) 263.—Am. bor. occ.
Kingii, *Torr. & Gray, l. c.* viii. (1870) 165.—Am. bor. occ.
lachnogynum, *Torr. ex Benth. in DC. Prod.* xiv. 8.—Am. bor. occ.
lateriflorum, *Rafin. New Fl. Am.* iv. 53.—Am. bor.
latifolium, *Sm. in Rees, Cycl.* xiii. n. 8.—Calif.
laxiflorum, Nutt. ex Benth. in DC. Prod. xiv. 18 = microthecum.
Lemmoni, *S. Wats. in Proc. Am. Acad.* xii. (1877) 266.—Am. bor. occ.
leptocladon, Torr. & Gray, in Pacif. Rail. Rep. ii. 129 = microthecum.
leucocladon, Benth. Pl. Hartw. 338 = gracile.
Lindheimerianum, Scheele, in Linnæa, xxii. (1849) 149 = sinuum.
Lobbii, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 162.—Calif.
lonchophyllum, *A. Gray, l. c.* 173.—Am. bor.
longifolium, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 164.—Florida.
marifolium, *A. Gray, in Proc. Am. Acad.* viii. (1870) 163.—Calif.
microcladon, Nutt. in Journ. Acad. Philad. i. (1847) 164 = breviscale.
microthecum, *Nutt. l. c.* 162.—Am. bor. occ.
mohavense, *S. Wats. in Proc. Am. Acad.* xii. (1877) 266.—Calif.
molestum, *S. Wats. l. c.* xvii. (1881–82) 379.—Am. bor.
multiceps, *Nees, in Pr. Neuwied, Reise Nord-Amer.* ii. 446.—Am. bor. occ.
multiflorum, *Benth. in Trans. Linn. Soc.* xvii. (1837) 412.—Texas.
niveum, *Dougl. ex Benth. l. c.* 414.—Am. bor. occ.
nudum, *Dougl. ex Benth. l. c.* 413.—Calif.
nutans, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 181.—Am. bor. occ.
Nuttallii, Gamb. ex Nutt. in Journ. Acad. Philad. N. S. i. (1847) 166 = ovalifolium.
oblongifolium, Benth. in Trans. Linn. Soc. xvii. (1837) 412 = latifolium.
obtusum, Benth. in DC. Prod. xiv. 15 = racemosum.
ochrocephalum, *S. Wats. in Bot. Calif.* ii. 480.—Calif.

ERIOGONUM :—
Orcuttianum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 371.—Calif.
ovalifolium, Torr. ex Benth. in DC. Prod. xiv. 15 = racemosum.
ovalifolium, Benth. in Trans. Linn. Soc. xvii. (1837) 409 = Douglasii.
ovalifolium, *Nutt. in Journ. Acad. Philad.* vii. (1834) 50. t. 8. f. 1.—Am. bor. occ.
Palmeri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 267. —Am. bor. occ.
panduratum, *S. Wats. in Bot. Calif.* ii. 480.—Calif.
Parishii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 379.—Am. bor.
Parryi, *A. Gray, l. c.* x. (1874) 77.—Utah.
parviflorum, Nutt. Gen. Am. i. 261 = pauciflorum.
parvifolium, *Sm. in Rees, Cycl.* xiii. n. 2.—Calif.
pauciflorum, *Pursh, Fl. Am. Sept.* ii. 735.—Am. bor. occ.
pharnaceoides, *Torr. in Stigr. Rep.* 167. t. 11.—N. Mexic.
platyphyllum, Torr. ex Benth. in DC. Prod. xiv. 20 = tenellum.
Plumatella, *Dur. & Hilg. in Pacif. Rail. Rep.* v. 14. t. 16.—Am. bor. occ.
polifolium, Benth. in DC. Prod. xiv. 12 = fasciculatum.
polyanthum, Benth. l. c. = umbellatum.
polycoyx, Nutt. ex Torr. & Gray, in Proc. Am. Acad. viii. (1870) 165, nota = ovalifolium.
polycladon, *Benth. in DC. Prod.* xiv. 16.—N. Mexic.
Pratenianum, *Dur. in Journ. Acad. Philad.* Ser. II. iii. (1855) 100.—Am. bor.
proliferum, Torr. & Gray, in Proc. Am. Acad. viii. (1870) 164 = ovalifolium.
puberulum, *S. Wats. l. c.* xiv. (1879) 295.—Am. bor. occ.
purpureum, Nutt. ex Benth. in DC. Prod. xiv. 10 = ovalifolium.
pusillum, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 154.—Am. bor. occ.
pyrolæfolium, *Hook. ex A. Murr. Oreg. Circ.* 2; *et in Kew Journ.* v. (1853) 295. t. 10.—Calif.
racemosum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 161.—Am. bor. occ.
reniforme, *Torr. ex Benth. in DC. Prod.* xiv. 21.—Am. bor. occ.
robustum, *Greene, in Bull. Calif. Acad.* i. iii. (1885) 126.—Am. bor. occ.
rosense, Dur. & Hilg. in Pacif. Rail. Rep. v. 14 = virgatum.
rosmarinifolium, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 164 = fasciculatum.
rotundifolium, *Benth. in DC. Prod.* xiv. 21.—N. Mexic.
salsuginosum, *Hook. Kew Journ.* v. (1853) 264.—Am. bor. occ.
sanctæ, *S. Wats. in Proc. Am. Acad.* xii. (1877) 267. —Calif.
scalare, *S. Wats. l. c.* 261.—Calif.
sericeum, Pursh, Fl. Am. Sept. i. 277 = flavum.
sericeum, Torr. & Gray, in Proc. Am. Acad. viii. (1870) 155 = jamesii.
shastense, *A. Murr. Oreg. Circ.* 2.—Calif.
Shockleyi, *S. Wats. in Proc. Am. Acad.* xvii. (1882–83) 194.—Am. bor. occ.
Simpsonii, Benth. in DC. Prod. xiv. 18 = microrhecum, annuum.
spathulatum, *A. Gray, in Proc. Am. Acad.* x. (1874) 76.—Am. bor.
spergulinum, *A. Gray, l. c.* vii. (1868) 380.—Calif.
sphærocephalum, *Dougl. ex Benth. in Trans. Linn. Soc.* xvii. (1837) 407.—Am. bor. occ.
staticifolium, Hook. in Kew Journ. v. (1853) t. 10 = pyrolæfolium.
stellatum, Benth. in Trans. Linn. Soc. xvii. (1837) 409 = umbellatum.
strictum, *Benth. l. c.* 414.—Am. bor. occ.
subreniforme, *S. Wats. in Proc. Am. Acad.* xii. (1877) 260.—Am. bor. occ.
suffruticosum, *S. Wats. l. c.* xx. (1885) 370.—Texas.
sulcatum, *S. Wats. l. c.* xiv. (1879) 296.—Am. bor.
tenellum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 165.—Am. bor. occ.
tenellum, Torr. in Ann. Lyc. Nat. Hist. N. York, ii. (1826) 241 = ovalifolium.
tenuerame, Schoele, in Linnæa, xxii. (1849) 150 = longifolium.

ERIOGONUM :—
Thomasii, *Torr. in Pacif. Rail. Rep.* iv. 264.—Am. bor. occ.
Thompsonæ, *S. Wats. in Am. Natural.* vii. (1873) 302.—Am. bor.
Thurberi, *Torr. Bot. Mex. Bound.* 176.—Am. bor. occ.
thymoides, *Benth. in DC. Prod.* xiv. 9.—Am. bor. occ.
Tolmieanum, Hook. Fl. Bor. Am. ii. 134 = umbellatum.
tomentosum, *Micha. Fl. Bor. Am.* i. 246. t. 24.—Am. bor. occ.
Torreyanum, *A. Gray, in Proc. Am. Acad.* viii. (1870) 155.—Calif.
trachygonum, Torr. ex Benth. in DC. Prod. xiv. 15 = Wrightii.
trichopes, *Torr. in Emory, Notes Mil. Reconnois.* 150. —Am. bor. occ.
trichopodum, Benth. in DC. Prod. xiv. 20 = trichopes.
trisne, *S. Wats. in Proc. Am. Acad.* x. (1875) 347.—Am. bor. occ.
truncatum, *A. Gray, in Proc. Am. Acad.* viii. (1870) 154.—Calif.
umbellatum, Benth. in Trans. Linn. Soc. xvii. (1837) 410 = heracleoides.
umbellatum, *Torr. in Ann. Lyc. Nat. Hist. N. York*, ii. (1826) 241.—Am. bor. occ.
undulatum, *Benth. in DC. Prod.* xiv. 7.—Mexic.
ursinum, *S. Wats. in Proc. Am. Acad.* x. (1875) 347.—Calif.
vagans, *S. Wats. l. c.* xx. (1885) 370.—Am. bor.
verticillatum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 165 = virgatum.
villiflorum, *A. Gray, in Proc. Am. Acad.* viii. (1872) 630.—Am. bor. occ.
vimineum, *Dougl. ex Benth. in Trans. Linn. Soc.* xvii. (1837) 416.—Am. bor. occ.
virgatum, *Benth. in DC. Prod.* xiv. 16.—Calif.
Watsonii, *Torr. & Gray, in Proc. Am. Acad.* viii. (1873) 138.—Calif.
Wrightii, *Torr. ex Benth. in DC. Prod.* xiv. 15.—Am. bor. occ.

ERIOGYNIA, Hook. Fl. Bor. Am. i. 255. t. 88 (1838) = Spiræa, Linn. (Rosac.).
pectinata, Hook. l. c. = S. pectinata.

ERIOLÆNA, DC. in Mém. Mus. Par. x. (1823) 102. t. 5. *STERCULIACEÆ*, Benth. & Hook. f. i. 220.
ERIOCHLÆNA, Spreng. Syst. iii. 128 (1826).
GUMELIA, Buch.-Ham. ex Wall. Cat. sub n. 1178 (1832).
JACKIA, Spreng. Syst. iii. 85 (1826).
MICROCHLÆNA, Wight & Arn. Prod. 70 (1834).
MICROLÆNA, Wall. Cat. n. 1178 (1832).
SCHILLERA, Reichb. Consp. 204 (1828).
WALLICHIA, DC. in Mém. Mus. Par. x. (1823) 104. t. 6.
Candolleï, *Wall. Pl. As. Rar.* i. 57.—Ind. or.
Hookerianæ, *Wight & Arn. Prod.* 70.—Ind. or.
lepidocalycis, *Wight, Ic.* t. 895.—Ind. or.
Roxburghii, Spreng. Syst. iii. 128 = Pterospermum semisagittatum.
spectabilis, *Planch. ex Hook. f. Fl. Brit. Ind.* i. 372.—Reg. Himal.
Stocksii, *Hook. f. & Thoms. ex Mast. in Fl. Brit. Ind.* i. 370.—Ind. or.
Wallichii, *DC. in Mém. Mus. Par.* x. (1823) 102.—Ind. or.

ERIOLEPIS, Cass. in Dict. Sc. Nat. xli. 331 (1825) = Cnidoscolus, Linn. (Compos.).
achinata, Cass. l. c. i. 470 = C. echinatus.
feros, Cass. l. c. = C. ferox.
italica, Cass. l. c. = C. Italicus.
lanceolata, Cass. l. c. xli. 331 = C. lanceolatus.
lanciflora, Cass. l. c. i. 470 = C. lanciflorus.
lanigera, Cass. l. c. = C. cynaroides.

ERIOLITHIS, Geærtn. Fruct. ii. 277 (1791). *ROSACEÆ*.
mirabilis, *Geærtn. l. c.*—Perav.

ERIOLOBUS, M. Roem. Syn. Rosif. 104, 216 (1847) = Pyrus, Tourn. (Rosac.).
trilobatus, M. Roem. l. c. 216 = P. trilobata.

ERIOLYTRUM, Desv. ex Kunth, Rév. Gram. i. 219 (1829) = Panicum, Linn. (Gramin.).
junceum, Desv. ex Kunth, l. c. = P. racemosum.

ERIONE, Schott & Endl. Melet. 34 (1832) = Ericdendron, DC. (Malvac.).
jasminiodora, Schott & Endl. l. c. = Eriodendron jasminiodorum.

ERIONIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. ? Art. IV. 2, nomen (Gen. ignot.).

ERIOPAPPUS, Arn. in Lindl. Introd. Nat. Syst. ed. II. 448 (1836) = Layia, Hook. (Compos.).
glandulosus, Arn. l. c. = L. glandulosa.

ERIOPAPPUS, Hort. ex Lond. Encyc. Pl. 688 (1829) = Eupatorium, Tourn. (Compos.).
paniculatus, Hort. ex Loud. l. c. = Eupatorium paniculatum.

ERIOPE, Humb. & Bonpl. ex Benth. Lab. Gen. et Sp. 142 (1888). *LABIATÆ*, Benth. & Hook. f. ii. 1178.
alpestris, *Mart. ex Benth. l. c.* 145.—Bras.
complicata, *Mart. ex Benth. l. c.* 144.—Bras.
crassifolia, *Mart. ex Benth. l. c.* 143.—Bras.
crassipes, *Benth. l. c.* 144.—Bras.
filifolia, *Benth. in DC. Prod.* xii. 141.—Bras.
fœtida, *St. Hil. ex Benth. Lab. Gen. et Sp.* 145.—Bras.
hypenioides, *Mart. ex Benth. l. c.* 143.—Bras.
macrostachya, *Mart. ex Benth. l. c.* 145.—Bras.
micrantha, *Benth. in DC. Prod.* xii. 141.—Bras.
monticola, *Mart. ex Benth. Lab. Gen. et Sp.* 143.—Bras.
nudiflora, *Humb. & Bonpl. ex Benth. l. c.* 143.—Am. trop.
obtusata, *Benth. l. c.* 143.—Bras.
pallens, *Benth. in DC. Prod.* xii. 143.—Bras.
parviflora, Steud. Nom. ed. II. i. 587 = seq.
parvifolia, *Mart. ex Benth. Lab. Gen. et Sp.* 144.—Bras.
polyphylla, *Mart. ex Benth. l. c.* 143.—Bras.
stricta, *Benth. in DC. Prod.* xii. 142.—Bras.
tenuicaules, *St. Hil. ex Benth. Lab. Gen. et Sp.* 144.—Bras.

ERIOPETALUM, Wight, Contrib. 35 (1834) = Brachystelma, R. Br. (Asclepiad.).
attenuatum, Wight, l. c. = B. attenuatum.
lasvigatum, Wight, l. c. = B. lasvigatum.
parviflorum, Wight, l. c. = B. parviflorum.

ERIOPHA, Hill, Hort. Kew. 69 (1769) = Centaurea, Linn. (Compos.).
Eriopha, Hill, l. c. = C. eriophora.

ERIOPHILA, Reichb. in Mossel. Handb. ii. 1127 (1828) = Erophila, DC. (Crucif.).

ERIOPHORUM, Linn. Syst. ed. I (1735). *CYPERACEÆ*, Benth. & Hook. f. iii. 1052 (partim).
LINAGROSTIS, Adans. Fam. ii. 41 (1763).
TRICHOPHORUM, Pers. Syn. i. 69 (1805).
alpicolum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 78 = angustifolium.
alpinum, *Linn. Sp. Pl.* 53.—Europ.; Am. bor.
alpinum, Vill. Hist. Pl. Dauph. ii. 164 = Scheuchzeri.
angustifolium, *Roth, N. Beitr.* i. 94.—Reg. bor. temp. et frig.
arundinaceum, Neos, in Wight, Contrib. 110 = comosum.
brachyanthrum, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 96) = vaginatum.
caespitosum, Host, Gram. Austr. i. 30. t. 39 = vaginatum.
callithrix, Lange, in Fl. Dan. Suppl. x. 19 = vaginatum.
callithrix, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 208. t. 2.—Ins. S. Laurent.
capitatum, Host, Gram. Austr. i. 30. t. 38 = Scheuchzeri.
Chamissonis, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 204. t. 2.—Sibir. alt.
comosum, *Neos, in Wight, Contrib.* 110; (*Wall. Cat.* n. 5448).—Ind. or.
comosum, Neos, l. c. partim = microstachyum.
cyperinum, Linn. Sp. Pl. ed. II. 77 = Scirpus Eriophorum.
dubium, Hegetschw. Fl. Schw. 62 = angustifolium.
dubium, *Steud. Syn. Pl. Cyp.* 146.—Moth.?
filamentosum, Reichb. in Mossel. Bot. Jahrb. v. (1834) 206 (Magnolias sp.).—Madagas.

ERIOPHORUM :—
gracile, *Koch, in Roth, Catal.* ii. (1799) *Addend.* 259. —Reg. bor. temp.
gracile, Sm. Engl. Bot. t. 2409 = angustifolium.
Hoppii, F. Nylander, in Act. Soc. Sc. Fenn. iii. (1849) 12. t. 2 = gracile.
Hudsonianum, Michx. Fl. Bor. Am. i. 24 = alpinum.
humile, Turcz. in Bull. Soc. Nat. Mosc. (1838) 108 = Scheuchzeri.
intermedium, Bast. in Desv. Journ. Bot. iii. (1814) 19 = angustifolium.
intermedium, Cham. ex C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 304 = Chamissonis.
latifolium, *Hoppe*, *Taschenb.* (1800) 103.—Reg. bor. temp. et frig.
leucocephalum, Boeck. in Flora, xli. (1858) 419 = Scheuchzeri.
Lockharti, *Steud. Nom.* ed. II. i. 587.—Ins. Trinit.
medium, Anderss. in Bot. Notiser (1857), 62 = Scheuchzeri.
microstachyum, *Boeck. in Linnaea*, xxxvlii. (1874) 399. —Ind. or.
minus, Dalla Torre, Atl. Alpenfl. 216 = angustifolium.
monostachion, Grosshk. in Keyserl. & Dersch. Beschr. Prov. Kurl. 116 = angustifolium.
montevidense, Link, Hort. Berol. i. 881 = Androtrichum polyocephalum.
pallidum, Neos. in Linnaea, ix. (1834) 293 = gracile.
polystachion, *Linn. Sp. Pl.* 52.—Reg. bor. temp. et frig.
polystachion, Sm. Engl. Fl. i. 67 = angustifolium.
pubescens, Sm. l. c. 68 = latifolium.
rufescens, Anderss. in Bot. Notiser (1857), 79 = Scheuchzeri.
russolum, Fries, Novit. Mant. iii. 67 = Scheuchzeri.
Scheuchzeri, *Hoppe*, *Taschenb.* (1800) 104. *App.* t. 7.—Reg. bor. temp. et frig.
sphærocephalum, Welfen, Fl. Norica, 62 = Scheuchzeri.
strictum, R. Br. in Richards. App. Frankl. Journ. 3 = latifolium.
tenellum, Nutt. Gen. Am. 2. Addit. = gracile.
triquetrum, Hoppe, Taschenb. (1800) 106 = gracile.
vaginatum, Curt. Fl. Lond. fasc. ii. t. 11 = Scheuchzeri.
vaginatum, *Linn. Sp. Pl.* 52.—Reg. bor. temp. et frig.
Vaillantii, Poll. & Turp. Fl. Paris, t. 52 = angustifolium.
virginicum, *Linn. Sp. Pl.* 52.—Am. bor.
vulgare, Pers. Syn. i. 70 = latifolium.
Zeyheri, Steud. Nom. ed. II. i. 588 (sphalm. = *Eriosema*) = Rhychosia Zeyheri.

ERIOPHORUS, Vaill. ex DC. Fl. Fr. iv. 86 (1805) = **Andryala**, Linn. (Compos.).

ERIOPEXYLLUM, Lag. Gen. et Sp. Nov. 26 (1816). *COMPOSITAE*, (Benth. & Hook. f. ii. 402.)
ambiguum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 26.—Calif.
caespitosum, *Dougl. ex Lindl. Bot. Reg.* t. 1167.—Am. bor. occ.
confertiflorum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 26.—Calif.
gracile, *A. Gray, l. c.* 26.—Am. bor. occ.
lanatum, Forbes, Hort. Woburn. 183 = caespitosum.
lanosum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 25. —Am. bor. occ.
multicaule, *A. Gray, l. c.* 24.—Am. bor. occ.
Nevinii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 459.—Calif.
nubigenum, *Greene, ex A. Gray, in Proc. Am. Acad.* xix. (1883) 26.—Calif.
opposititolium, Heynh. Nom. i. 310 = Bahia oppositifolia.
Pringlei, *A. Gray, in Proc. Am. Acad.* xix. (1883) 25. —Am. bor. occ.
stachadifolium, *Lag. Gen. et Sp. Nov.* 26.—Calif.
trollifolium, Lag. l. c. = Bahia trollifolia.
Wallacei, *A. Gray, in Proc. Am. Acad.* xix. (1883) 25. —Am. bor. occ.
Watsoni, *A. Gray, l. c.* 26.—Am. bor. occ.

ERIOPEXYTON, Benth. in Wall. Pl. As. Rar. i. 68 (1830). *LABIATAE*, Benth. & Hook. f. ii. 1215.
Wallichianum, *Benth. l. c.*—Reg. Himal.

ERIOPODIUM, Hochst. in Flora, xxiv. (1846) 115, in obs. = Anthospermum, Linn. (Gramin.).
Krauseli, Steud. l. c. = A. canescens.

ERIOPSIS, Lindl. Bot. Reg. (1847) t. 18. *ORCHIDEAE*, Benth. & Hook. f. iii. 546.
Pseuderiopsis, Reichb. f. in Linnaea, xxii. (1849) 852.
biloba, *Lindl. Bot. Reg.* (1847) t. 18.—Peruv.
rhytidobulbon, Lem. Jard. Fleur. ii. (1852) Misc. 77 = sec.
rutidobulbon, *Hook. Bot. Mag.* t. 4437.—Peruv.
Sceptrum, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 98.—Peruv.
Schomburgkii, *Reichb. f. l. c.* iii. (1855) 67.—Peruv.

ERIORHAPHE, Miq. Pl. Jungh. 280 (1851–55) = Fentapetes, Linn. (Sterculiac.).
punicea, Miq. l. c. = P. phoenicea.

ERIOSEMA, DC. [Prod. ii. 388 (1825)]; Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 421. (sphalm. Eurioema). *LEGUMINOSAE*, Benth. & Hook. f. i. 543.
Muxiria, Welw. Apont. 573 (1858).
Pyrnotrichia, Wight & Arn. Prod. 288 (1834).
Afzelii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 225.—Afr. trop.
ambiguum, Meisn. ex Kranss, in Flora, xxvii. (1844) 357 = Rhynchosia Ecklonel.
argenteum, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 423 = Argyrolobium sericeum.
barbatum, Desv. l. c. = crinitum.
Benthamianum, *Mart. ex Benth. in Linnaea*, xxii. (1849) 521.—Bras.
Bojeri, *Benth. ex Baker, in Journ. Linn. Soc.* xx. (1883) 139.—Madag.
Bojeriana, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 888. —Madag.
Burkei, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 260.—Afr. austr.
cajanoides, *Hook. f. ex Hook. Niger Fl.* 314.—Afr. trop.
campestre, *Benth. in Mart. Fl. Bras.* xv. i. 212.—Bras.
capitatum, E. Mey. Comm. Pl. Afr. Austr. i. 130 = Psoralea tomentosa.
chinense, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 81.—China.
chrysopotum, E. Mey. ex Steud. Nom. ed. II. i. 588 = Rhynchosia chrysopotis.
congestum, *Benth. in Mart. Fl. Bras.* xv. i. 214.—Bras.
consanguineum, Klotzsch, in Peters, Reise Mossamb. Bot. 32 = parviflorum.
cordatum, *E. Mey. Comm. Pl. Afr. Austr.* i. 128.—Afr. austr.
cordifolium, *Benth. in Linnaea*, xxii. (1849) 520.—Bras.
cordifolium, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 237.—Abyss.
crinitum, *Benth. in Hook. Journ. Bot.* ii. (1840) 62.—Am. trop.
crinitum, *G. Don, Gen. Syst.* ii. 348.—Mexic.
defoliatum, *Benth. in Linnaea*, xxii. (1849) 524.—Bras.
diffusum, *G. Don, Gen. Syst.* ii. 347.—Am. trop.
Dregei, Meisn. ex Kranss, in Flora, xxvii. (1844) 357 = parviflorum.
Dregei, E. Mey. Comm. Pl. Afr. Austr. 129 = squarrosum.
ellipticum, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 232.—Afr. trop.
elongatum, Baill. Adansonia, vi. (1866–66) 227 = glomeratum.
flavifforum, *Miq. in Linnaea*, xviii. (1844) 578.—Guian.
flemingioides, *Baker, in Oliver, Fl. Trop. Afr.* ii. 229. —Afr. trop.
floribundum, *Benth. in Linnaea*, xxii. (1849) 524.—Bras.
floribundum, Klotzsch, in Peters, Reise Mossamb. Bot. 33 = cajanoides.
glabrum, *Mart. ex Benth. in Linnaea*, xxii. (1849) 523.—Bras.
glomeratum, *Hook. f. in Hook. Niger Fl.* 313.—Afr. trop.
gracile, Klotzsch, in Peters, Reise Mossamb. Bot. 34 = glomeratum.
grandiflorum, *G. Don, Gen. Syst.* ii. 347; *Seem. Bot. Voy. Herald*, 345.—Mexic.

ERIOSEMA :—
griseum, *Baker, in Oliver, Fl. Trop. Afr.* ii. 228.—Afr. trop.
Guienzii, *Sond. in Linnaea*, xxiii. (1850) 34.—Afr. austr.
heterophyllum, *Benth. l. c.* 520.—Afr. trop.
incanum, Klotzsch, in Peters, Reise Mossamb. Bot. 35 = cajanoides.
insignis, *O. Hoffm. in Linnaea*, xliii. (1880–82) 128.—Afr. trop.
Krausianum, *Meisn. in Hook. Lond. Journ. Bot.* ii. (1843) 91.—Afr. austr.
lagoënse, *Michel, in Kjoeb. Vidensk. Meddel.* (1875) 93.—Bras.
lanceolatum, *Benth. in Hook. Journ. Bot.* ii. (1840) 62.—Am. trop.
lancifolium, Benth. ex Hemsl. Biol. Centr. Am. Bot. i. 313 = simplicifolium.
longepedunculatum, *A. Rich. Tent. Fl. Abyss.* i. 236.—Abyss.
longiforum, Benth. in Linnaea, xxii. (1849) 523 = Riedelii.
longifolium, *Benth. l. c.* 519.—Bras.
macrophyllum, Klotzsch, in Peters, Reise Mossamb. Bot. 34 = cajanoides.
Muxiria, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 229.—Afr. trop.
oblongum, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 259. —Afr. austr.
obovatum, *Benth. in Mart. Fl. Bras.* xv. i. 224.—Bras.
parviflorum, *E. Mey. Comm. Pl. Afr. Austr.* 130.—Afr. austr.
pauciflorum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 31.—Afr. trop.
platycarpon, *Michels, in Mém. Soc. Phys. Genèv.* xxvii. (1883) n. vii. 34.—Paraguay!
podocatchyum, Hook. f. ex Hook. Niger Fl. 314 = parviflorum.
polystachyum, *Baker, in Oliver, Fl. Trop. Afr.* ii. 225. —Afr. trop.
polystachyum, E. Mey. Comm. Pl. Afr. Austr. 130 = cajanoides.
populiflorum, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 259.—Afr. austr.
procumbens, *Benth. ex Baker, in Journ. Linn. Soc.* xx. (1883) 134.—Madag.
prorepens, *Benth. in Linnaea*, xxii. (1849) 524.—Bras.
psoraleoides, *G. Don, Gen. Syst.* ii. 348.—Madag.
psoraloides, Baill. in Bull. Soc. Linn. Par. i. (1888) 837 = prasc.
puberulum, Eckl. & Zeyh. Enum. 256 = Rhynchosia pubescens. *Steud.*
pulchellum, *G. Don, Gen. Syst.* ii. 348; *Benth. in Hook. Journ. Bot.* ii. (1840) 62.—Mexic.
pycnanthum, *Benth. in Mart. Fl. Bras.* xv. i. 212.—Bras.
pygmaeum, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 225.—Afr. trop.
radicosum, *A. Rich. Tent. Fl. Abyss.* i. 238.—Abyss.
reticulatum, *Benth. in Linnaea*, xxii. (1849) 521.—Bras.
reticulatum, E. Mey. Comm. Pl. Afr. Austr. 129 = Rhynchosia Ecklonel.
rhynchosioides, *Baker, in Journ. Linn. Soc.* xv. (1876) 94.—Afr. trop.
Riedelii, *Benth. in Linnaea*, xxii. (1849) 523.—Bras.
rigidum, *Benth. l. c.* 522.—Bras.
robustum, *Baker, in Oliver, Fl. Trop. Afr.* ii. 229.—Afr. trop.
rotundifolium, *Michels, in Mém. Soc. Phys. Genève.* xxvii. (1883) n. vii. 35.—Paraguay.
rufum, Baill. Adansonia, vi. (1866) 226 = glomeratum.
rufum, *G. Don, Gen. Syst.* ii. 347; *Benth. in Hook. Journ. Bot.* ii. (1840) 62.—Am. trop.
saligonum, *E. Mey. Comm. Pl. Afr. Austr.* 129.—Afr. austr.
sericeum, *Baker, in Oliver, Fl. Trop. Afr.* ii. 226.—Afr. trop.
sericeum, Eckl. & Zeyh. Enum. 256 = Rhynchosia Orthodanum.
sessiliforum, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 422.—Peruv.
simplicifolium, *G. Don, Gen. Syst.* ii. 346.—Panama.
speciosum, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 230.—Afr. trop.

ERIOSEMA :—
spicatum, Hook. f. in Hook. Niger Fl. 513 = parviflorum.
squarrosum, *Walp. in Linnaea*, xiii. (1839) 536.—Afr. austr.
stipulare, *Benth. l. c.* xxii. (1849) 519.—Bras.
stictum, *Benth. l. c.* 520.—Bras.
trinerve, *E. Mey. Comm. Pl. Afr. Austr.* 130.—Afr. austr.
tuberosum, *A. Rich. Tent. Fl. Abyss.* i. 227.—Abyss.
venulosum, *Benth. in Mart. Fl. Bras.* xv. i. 211.—Bras.
violaceum, *G. Don, Gen. Syst.* iii. 847; *Benth. in Hook. Journ. Bot.* ii. (1840) 92.—Am. trop.
volubile, *Michell, in Mém. Soc. Phys. Genèv.* xxviii. (1888) n. VII. 86.—Paraguay.
Zeyheri, E. Mey. Comm. Pl. Afr. Austr. 129 = squarrosum.

ERIOSERMUM, Thunb. Fl. Cap. ii. 294 (1818), sphalm. = Briospermum, Jacq. (Liliac.).

ERIOSOLENA, Blume, Bijdr. 651 (1825) = Daphne, Linn. (Thymel.).
affinis, Zoll. Syst. Verz. Ind. Archip. 116 (Rubiaces). —Malaya.
montana, Blume, Bijdr. 651 = D. pendula.
viridiflora, Zoll. & Mor. in Nat. Geneesk. Arch. Neêrl. Indie, i. (1844) 615 = Wikstroemia Candolleana.
Wallichii, Meisn. in DC. Prod. xiv. II. 540 = D. involucrata.

ERIOSOLENIA, Benth. & Hook. f. Gen. iii. 190 (1880) = prec.

ERIOSPERMUM, Jacq. Coll. v. Suppl. 72 (1796). *LILIACEAE*. Benth. & Hook. f. iii. 760.
PHYLLOGLOTTIS, Salisb. Gen. Pl. Fragm. 15 (1866).
TRAUMAZA, Salisb. l. c. (1866).
abyssinicum, *Baker, in Journ. Linn. Soc.* xv. (1876) 363.—Afr. trop.
albucoides, *Baker, in Gard. Chron.* (1876) i. 716.—Afr. austr.
andongense, *Welw. ex Baker, in Trans. Linn. Soc. Ser.* II. i. (1878) 261.—Afr. trop.
Bellendeni, *Sweet, Hort. Brit.* ed. I. 420.—Afr. austr.
Boviesanum, *Baker, in Journ. Linn. Soc.* xv. (1876) 367.—Afr. austr.
brevipes, *Baker, l. c.* 363.—Afr. austr.
calcaratum, *Baker, in Gard. Chron.* (1876) i. 716.—Afr. austr.
capense, Thunb. Fl. Cap. ii. 294 = latifolium.
Cooperi, *Baker, in Journ. Linn. Soc.* xv. (1876) 365. —Afr. austr.
corymbosum, *Baker, l. c.* 366.—Afr. austr.
Dregeanum, Presl, Bot. Bemerk. 118 = parvifolium.
flexuosum, *Welw. in Baker, in Trans. Linn Soc.* Ser. II. i. (1878) 261.—Afr. trop.
follicliforme, *Andr. Bot. Rep.* t. 521.—Afr. austr.
folioliferum, Ker-Gawl. in Bot. Reg. t. 795 = proliferum.
Kirkii, *Baker, in Journ. Linn. Soc.* xv. (1876) 367.—Afr. trop.
lanceaefolium, *Jacq. Coll.* v. *Suppl.* 72.—Afr. austr.
lanuginosum, *Jacq. Hort. Schoenb.* iii. 7. t. 265.—Afr. austr.
latifolium, *Jacq. Coll.* v. *Suppl.* 83.—Afr. austr.
latifolium, Ker-Gawl. in Bot. Mag. t. 1862 = Bellendeni.
Mackenii, *Baker, in Journ. Linn. Soc.* xv. (1876) 366. —Afr. austr.
ophioglossoides, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 262.—Afr. austr.
ornithogaloides, *Baker, in Journ. Linn. Soc.* xv. (1876) 366.—Afr. austr.
palmiceum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 261.—Afr. trop.
paradoxum, *Ker-Gawl. in Bot. Mag.* sub t. 1862.—Afr. austr.
parviflorum, *Jacq. Coll.* v. *Suppl.* 74.—Afr. austr.
proliferum, *Baker, in Journ. Linn. Soc.* xv. (1876) 367.—Afr. austr.
pubescens, *Jacq. Hort. Schoenb.* iii. 8. t. 264.—Afr. austr.
spinulo, *Berg. ex Schnitt. f. Syst.* vii. 1696.—Afr. austr.
stenophyllum, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 261.—Afr. trop.

ERIOSPHAERA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 221 (1817) = Leucospermum, Leg. (Compos.).
multiflda, F. G. Dietr. l. c. = L. pedunculare.

ERIOSPHAERA, Less. Syn. Comp. 270 (1832). *COMPOSITAE*. Benth. & Hook. f. ii. 251.
apiculata, DC. Prod. vi. 166 = Helichrysum marifolium.
catipes, DC. l. c. 167 = Helichrysum cotipes.
coriacea, DC. l. c. = Helichrysum rotundatum.
dubia, DC. l. c. = Leontonyx squarrosa ?
Oculus-cati, DC. l. c. 166 = Helichrysum marifolium.
Oculus-cati, *Less. Syn. Comp.* 270.—Afr. austr.
rotundifolia, DC. Prod. vi. 166 = Helichrysum rotundatum.
umbellata, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 79 = Helichrysum umbellatum.

ERIOSPORA, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 502 (1850). *CYPERACEAE*, Benth. & Hook. f. iii. 1070.
CATAGYNA, Besuv. in Lestib. Essai Fam. Cyp. 26 (1819), nomen prius.
abyssinica, *Hochst. ex A. Rich. l. c.*—Abyss.
pilosa, *Benth. in Hook. Ic. Pl.* xiv. 30. t. 1432. —Afr. trop.

ERIOSTAX, Rafin. Fl. Tellur. iv. 25 (1836) = Aechmea, Ruiz & Pav. (Bromel.).
glauca, Rafin. l. c. = A. bromeliaefolia.

ERIOSTEMON, Less. Syn. Comp. 12 (1832) = Sausurea, DC. (Compos.).
taraxacifolium, Less. l. c. = S. Eriostemon.

ERIOSTEMON, Sm. in Trans. Linn. Soc. iv. (1798) 221. *RUTACEAE*, Benth. & Hook. f. i. 292.
alpinus, F. Muell. Fragm. ii. 108 = Phebalium squamulosum.
ambiens, *F. Muell. l. c.* vi. 166.—Austral.
amblycarpus, F. Muell. l. c. i. 102 = Phebalium amblycarpum.
amplifolius, *F. Muell. in Melb. Chemist* (Dec. 1884). —Austral.
anceps, *F. Muell. Fragm.* v. 109.—Austral.
ancops, Spreng. Syst. ii. 322 = Phebalium argenteum.
australasicus, Sm. ex Steud. Nom. ed. II. i. 586 = salicifolius.
Baileyi, Prits. Ic. Ind. ii. 119, sphalm. = Phebalium Ralstoni.
Banksii, *A. Cunn. ex Endl. in Enum. Pl. Hueg.* 15.—Austral.
Beckleri, *F. Muell. Fragm.* ix. 109.—Austral.
Benthamii, *F. Muell. l. c.* 108.—Austral.
bilobus, F. Muell. l. iv. 103 = Phebalium rude.
brevifolius, A. Cunn. ex Endl. in Enum. Pl. Hueg. 16 = diffusus.
buxifolius, *F. Muell. Fragm.* vii. 36.—Austral.
buxifolius, *Sm. in Rees, Cycl.* xiii. n. 2.—Australia.
calycinus, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 14 = nodiflorus.
capparideus, Sm. Syn. i. 465 = Adenandra uniflora.
capitatus, F. Muell. Fragm. i. 106 = Microcybe pauciflora.
corymbiflorus, F. Muell. l. c. 105 = Asterolasia corrifolia, Muelleri.
corymbosus, *Labill. Sert. Austro-Caled.* t. 54.—N. Cal.
Cosdi, *F. Muell. in Melb. Chemist* (Dec. 1884).—Austral.
Cunei, F. Muell. Vict. Pl. i. 119 = Crowea exalata, saligna.
Cunninghamii, *F. Muell. Fragm.* ix. 107.—Austral.
cuspidatus, A. Cunn. in Field, N. S. Wales, 321 = myoporoides.
decussatus, Colla, Hort. Ripul. 52. t. 30 = Eleaocarpus dentatus.
diffornis, *A. Cunn. ex Endl. in Enum. Pl. Hueg.* 15. —Austral.
Drummondii, F. Muell. Fragm. i. 106 = Asterolasia phebalioides.
obovatissimus, Endl. in Enum. Pl. Hueg. 15 = salicifolius.
ophurus, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 14 = splendens.

ERIOSTEMON :—
elaeagnifolius, Baill. Hist. Pl. iv. 388 = Phebalium squamulosum.
elatior, F. Muell. Fragm. i. 181 = Phebalium Billardieri.
ericifolius, *A. Cunn. ex Benth. Fl. Austral.* i. 885.—Austral.
filifolius, *F. Muell. Fragm.* ix. 108.—Austral.
Gelesnowii, F. Muell. l. c. i. 107 = Geleznowia verrucosa.
glaucescens, *Ait. ex Sweet, Hort. Brit.* ed. III. 766 (Quid ?).—Austral.
gracilis, *R. Grah. in Edinb. N. Phil. Journ.* xvi. (1834) 175.—Austral.
grandiflorus, F. Muell. Fragm. i. 105 = Asterolasia grandiflora.
halmaturorum, F. Muell. in Linnaea, xxv. (1852) 376 = linearis.
Hillebrandii, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 10 = Phebalium bilobum.
hispidulus, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 164. —Austral.
Hookeri, F. Muell. Fragm. i. 104 = Asterolasia squamuligera.
intermedius, Hook. Bot. Mag. t. 4489 = myoporoides.
Kaulack, Monit. in Mém. Acad. Lyon, x. (1860) 191.—N. Caled.
lamprophyllus, *F. Muell. in Journ. Pharm. Soc. Vict.* ii. (1858) 43 = Phebalium lamprophyllum.
lanceolatus, Gaertn. f. Fruct. iii. 154. t. 210 = salicifolius.
lanicifolius, F. Muell. in Trans. Vict. Inst. i. (1855) 32 = myoporoides.
Leichhardti, *F. Muell. Fragm.* v. 5.—Austral.
lepidotus, Spreng. Syst. ii. 322 = Phebalium squamulosum.
linearifolius, DC. Prod. i. 730 = Geijera parviflora.
linearis, *A. Cunn. ex Endl. in Enum. Pl. Hueg.* 16. —Austral.
marginatus, Sm. in Rees, Cycl. xiii. n. 5 = Adenandra marginata.
Maxwelli, *F. Muell. Fragm.* ix. 106.—Austral.
microphyllus, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1854) 99.—Austral.
mollis, F. Muell. Census, 11 = Asterolasia mollis.
montanus, F. Muell. Pl. Vict. i. 129 = Phebalium montanum.
Mortoni, *F. Muell. Fragm.* ix. 108.—Austral.
myoporoides, *DC. Prod.* i. 730.—Austral.
nerifolium, Sieber, ex Spreng. Syst. iv. Cur. Post 164 = myoporoides.
nodiflorus, *Lindl. Swan River App.* 17.—Austral.
Nottii, *F. Muell. Fragm.* vi. 32.—Australia.
obcordatus, A. Cunn. ex Hook. Journ. Bot. i. (1834) 254 = obovalis.
oblongifolium, Steud. Nom. ed. II. i. 586 = Acronychia laevis.
obovalis, *A. Cunn. in Field, N. S. Wales*, 321.—Austral.
Oldfieldii, F. Muell. Fragm. i. 3 = Phebalium Oldfieldii.
ovatifolius, F. Muell. l. c. 108 = Phebalium ovatifolium.
oxothamnoides, F. Muell. l. c. = Phebalium oxothamnoides.
pallidus, F. Muell. Census, 11 = Asterolasia pallida.
parvatamus, Sm. in Rees, Cycl. xiii. n. 6 = Boronia ledifolia.
parvifolius, *R. Br. ex Benth. Fl. Austral.* i. 885.—Austral.
phyllicifolius, F. Muell. Fragm. i. 105 = Phebalium phyllicifolium.
phyllicoides, F. Muell. l. c. 107 = Phebalium diosmeum.
pleurandroides, F. Muell. l. c. 106 = Asterolasia pleurandroides.
pulchellum N. Lem. Jard. Fleur. iv. (1854) t. 396.
pungens, Lindl. in Mitch. Three Exped. ii. (1839) 156 = Phebalium pungens.
racemosum, Endl. in Enum. Pl. Hueg. 15 = spicatum.
Ralstoni, F. Muell. Fragm. ii. 101. t. 14 = Phebalium Ralstoni.
rhomboeus, Lindl. in Mitch. Journ. Trop. Austral. 326 = diffornis.
rotundifolius, A. Cunn. ex Endl. in Enum. Pl. Hueg. 15 = Phebalium rotundifolium.
salicifolius, Sm. in Rees, Cycl. xiii. n. 1.—Austral.
saligneus, Endl. Prod. Fl. iv. 307 = Crowea saligna.

ERIOSTEMON :—
salsolifolius, Sm. in Rees, Cycl. xiii. n. 8 = Philotheca australis.
Sandfordii, F. Muell. Fragm. i. 107 = Gelesnowia calycina.
scaber, *A. DC. Nat.* 10^mt *Yard. Bot. Genèv.* 8.—Hab.!
scaber, *Pant. Mag. Bot.* xiii. (1847) 137.—Austral.
sedifornis, F. Muell. Fragm. i. 102 = Phebalium glandulosum.
serrulatus, F. Muell. l. c. 4 = Phebalium bilobum.
Smithianus, W. Hill, ex Benth. Fl. Austral. i. 385 = difformis.
spicatus, *A. Rich. Sert. Astrol.* ii. 76. t. 27.—Austral.
squameus, Lehill. Nov. Holl. Pl. i. 111. t. 141 = Phebalium Billardierii.
trachyphyllus, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1856) 99.—Austral.
trianeria, *Hook. Journ. Bot.* i. (1884) 254.—Tasman.
trymalioides, F. Muell. Fragm. i. 106; Vict. Pl. i. 134 = Asterolasia trymalioides.
tuberculosus, F. Muell. Pl. Vict. i. 130 = Phebalium tuberculosum.
Turczaninowii, F. Muell. l. c. 120 = Correa angustifolia.
umbellatum, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) *ii.* 15 = Phebalium dentatum.
uniflorus, Sm. in Rees, Cycl. xiii. n. 4 = Adenandra uniflora.
verrucosus, A. Rich. Sert. Astrolab. ii. 74. t. 26 = obovalis.
virgatus, *A. Cunn. ex Hook. f. in Hook. Journ. Bot.* ii. (1840) 417.—Austral.

ERIOSTEMUM, Colla, ex Steud. Nom. ed. II. i. 588 (1840) = **Blaxocarpus**, Burm. (Tiliac.).
dentatum, Colla, ex Steud. l. c. = Blaxocarpus cyaneus.

ERIOSTEMUM, Poir. in Dict. Sc. Nat. xv. (1819) 196 = **Eriostemon**, Sm. (Rutac.).

ERIOSTEMUM, Steud. Nom. ed. II. i. 588 (1840) = seq.

ERIOSTOMUM, Hoffngg. & Link, Fl. Port. i. 105. t. 7 (1809) = **Stachys**, Tourn. (Labiat.).
alpinum, Hoffngg. & Link, ex Steud. Nom. ed. I. 806, in syn. = S. alpinum.
creticum, Hoffngg. & Link, Fl. Port. i. 105, in obs. = S. cretica.
germanicum, Hoffngg. & Link, l. c. = S. germanica.
lanatum, Hoffngg. & Link, l. c. = S. lanata.
lusitanicum, Hoffngg. & Link, l. c. = S. germanica.
polystachyum, Presl, Fl. Sicul. 87 = S. germanica.
salviaefolium, Presl, l. c. = S. italica.

ERIOSYCIS, Phil. in Anal. Univ. Chil. (1872) 721.
CACTACEÆ.
Sandilion, *Phil. l. c.*—Chili.

ERIOSYNAPHE, DC. Prod. iv. 175 (1830) = **Ferula**, Tourn. (Umbellif.).
Kotschyana, Fenzl, in Tchihat. Asie Min. Bot. i. 483 = Johrenia selinoides.
longifolia, DC. Prod. iv. 175 = F. longifolia.
tortuosa, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 35 = Johrenia fœnicea.

ERIOTHECA, Schott & Endl. Melet. 35 (1832) = **Bombax**, Linn. (Malvac.).
glabrescens, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xvii. (1842) 211 = B. glabrescens.
parviflora, Schott & Endl. Melet. 35 = B. parviflora.
pubescens, Schott & Endl. l. c. = B. pubescens.
tomentosa, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xvii. (1842) 211 = B. tomentosa.

ERIOTHERIX, Cass. in Bull. Soc. Philom. (1817) 32 (Eriotrix). *COMPOSITÆ*, Benth. & Hook. f. ii. 443.
juniperifolia, *Cass. in Dict. Sc. Nat.* xv. 260.—Ins. Barbou.
lycopodioides, DC. Prod. vi. 299 = juniperifolia.

ERIOTHYMUS, J. A. Schmidt, in Mart. Fl. Bras. viii. 171. t. 83 (1866) = **Kettikia**, Benth. (Labiat.).
rubicunda, J. A. Schmidt, l. c.—Bras.

ERIOTRICHIUM, Lem. in Orb. Dict. v. 406 (1849) = **Eritrichium**, Schrad. (Boragin.).

ERIOXANTHA, Rafin. in Lond. Gard. Mag. viii. (1832) 247 = **Eria**, Lindl. (Orchid.).

ERIPHIA, P. Br. Hist. Jamaic. 270 (1756) = **Bealeria**, Linn. (Acanth.).
jamaicensis, Roem. & Schult. Syst. iii. 337 = B. lutea.
lutea, Rafin. Sylva Tellur. 77 = B. lutea.
pallida, Rafin. l. c. = præc.

ERIPHILEMA, Herb. in Lindl. Bot. Reg. (1843) Misc. 85 = **Staynchium**, Linn. (Irid.).
grandiflora, Herb. l. c. = S. grandiflorum.

ERIPHLEMA, Baker, in Journ. Linn. Soc. xvi. (1877) 115 = ERIPHILEMA, Herb. = **Staynchium**, Linn. (Irid.).

ERISIMUM, Neck. Delic. Gallo-Belg. i. 280 (1768) = **Erysimum**, Linn. (Crucif.).

ERISMA, Rudge, Pl. Gui. Rar. i. 7. t. 1 (1805). *VOCHYSIACEÆ*, Benth. & Hook. f. i. 976.
DENDRÆA, Roem. & Schult. Syst. i. 4 (1817).
DITMARIA, Spreng. Anleit. ii. 11. 704 (1818); et Syst. i. 4 (1825).
calcaratum, *Warm. in Mart. Fl. Bras.* xiii. ii. 111.—Am. trop.
floribundum, *Rudge, Pl. Gui. Rar.* i. 7, 8. t. 1.—Guiana.
Japura, *Spruce, ex Warm. in Mart. Fl. Bras.* xiii. ii. 106.—Bras.
lasiflorum, *Warm. l. c.*—Am. trop.
multiflorum, *Link, ex Warm. l. c.* 114.—Bras.
micranthum, *Spruce, ex Warm. l. c.* 113.—Am. trop.
nitidum, *DC. Prod.* iii. 80.—Guian.
nitrum, *Link, ex Warm. in Mart. Fl. Bras.* xiii. ii. 114.—Bras.
uncinatum, *Warm. l. c.* 110.—Bras.
violaceum, *Mart. Nov. Gen. et Sp.* i. 187.—Bras.

ERISMANTHUS, Wall. Cat. n. 8011 (1846). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 325.
obliquus, *Wall. l. c.*—Ins. Penang.

ERITHALIA, Bunge, ex Steud. Nom. ed. II. i. 671, in syn. (1840) = **Gentiana**, Tourn.

ERITHALIS, P. Br. Hist. Jamaic. 165 (1756). *RUBIACEÆ*, Benth. & Hook. f. ii. 105.
HERRERA, Adans. Fam. ii. 158 (1763).
angustifolia, *DC. Prod.* iv. 465.—Cuba.
ellipitica, Rafin. Sylva Tellur. 123 = fruticosa.
fruticosa, *Linn. Syst. ed.* X. 930.—Ind. occ.
inodora, Jacq. Select. Am. 73, 178. f. 23 = fruticosa.
parviflora, *Griseb. Cat. Pl. Cub.* 134.—Cuba.
pentagonia, *DC. Prod.* iv. 465.—Cuba.
procumbens, Rafin. Sylva Tellur. 123 = fruticosa!
rotundata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 507.—Ind. occ.
sumiflora, Cuerun. f. Trect. iii. 93. t. 196 (= *Polyphragmon minus*).—Hab.!
vaccinifolia, *Benth. & Hook. f. Gen.* ii. 105.—Cuba.

ERITHALIS, Forst. f. Prod. 17 (1786) = **Timonius**, Rumph. (Rubiac.).
cymosa, Forst. f. ex DC. Prod. iv. 465 = T. Forsteri.
obovata, Forst. f. Prod. 88, sphalm. = seq.
polygama, Forst. f. l. c. f. 17 = T. Forsteri.
polygama, Willd. Sp. Pl. i. 997, partim = T. Rumphii.
Timon, Spreng. Pugill. i. 18 = T. Rumphii.

ERITHEIS, S. F. Gray, Nat. Arrr. Brit. Pl. ii. 464 (1821) = Intula, Linn. (Compos.).
maritima, S. F. Gray, l. c. = I. crithmoides.

ERITHRÆA, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 80 = **Erythræa**, DC. (Gentian.).

ERITRICHIUM, Schrad. in Comm. Gotting. iv. (1830) 186, in obs. (Eritrichum) (Comm. Asperif. 16). *BORAGINEÆ*, Benth. & Hook. f. ii. 850.

ERITRICHIUM :—
ANOPLOCARYUM, Ledeb. Fl. Ross. iii. 154 (1847).
CRYPTANTHA, Lehm. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 35 (1836).
MEGASTOMA, Coss. & Dur. ex Benth. & Hook. f. Gen. ii. 851 (1876).
albiflorum, *Griseb. in Goett. Abh.* vi. (1854) 131.—Reg. Magell.
alyssoides, A. DC. in DC. Prod. x. 131 = Krynitzkia alyssoides.
angustifolium, A. Gray, in Proc. Am. Acad. v. (1862) 165 = Krynitzkia micromeris.
angustifolium, Torr. in Pacif. Rail. Rep. v. 363 = Krynitzkia angustifolia.
angustifolium, S. Wats. Bot. King's Exped. 241 = Krynitzkia ambigua.
apricum, *Phil. in Linnæa*, xxxiii. (1864–65) 190.—Chili.
arctioides, DC. Prod. x. 125 = nanum.
asperum, *Phil. in Linnæa*, xxix. (1857–58) 16.—Chili.
australasicum, *DC. Prod.* x. 134.—Austral.
barbigerum, A. Gray, Syn. Fl. N. Am. ii. i. 194 = Krynitzkia barbigera.
basiflxum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 165.—Ind. or.
brevipes, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 446 = Trigonotis brevipes.
californicum, DC. Prod. x. 130 = Krynitzkia californica.
calycinum, *Phil. Fl. Atac.* 88.—Chili.
canescens, A. Gray, in Proc. Am. Acad. x. (1874) 87 = Plagiobothrys canescens.
capitulliforum, *Clos, in C. Gay, Fl. Chil.* iv. 467.—Chili.
chaetocalyx, *Phil. Fl. Atac.* 89.—Chili.
Chamissonis, DC. Prod. x. 125 = nanum.
Choristianum, DC. l. c. 130 = Krynitzkia Chorisiana.
chrysanthum, *Phil. in Linnæa*, xxxiii. (1864–65) 191.—Chili.
ciliatum, DC. Prod. x. 127 = pectinatum.
circumscissum, A. Gray, in Proc. Am. Acad. x. (1874) 88 = Krynitzkia circumscissa.
clandestinum, A. DC. in DC. Prod. x. 129 = Krynitzkia clandestina.
collinum, *Phil. in Linnæa*, xxix. (1857–58) 17.—Chili.
congestum, *DC. Prod* x. 129.—Chili.
cornutifolium, Kellogg, in Proc. Calif. Acad. ii. (1863) 163 = Krynitzkia Chorisiana.
Cooperi, A. Gray, in Proc. Am. Acad. xix. (1883) 89 = Krynitzkia Cooperi.
crassisepalum, Torr. & Gray, in Pacif. Rail. Rep. ii. 171 = Krynitzkia crassisepala.
cryptanthum, *A. DC. in DC. Prod.* x. 129.—Chili.
cynoglossoides, *Phil. in Linnæa*, xxix. (1857–58) 16.—Chili.
Czeknanowskii, *Trautv. in Act. Hort. Petrop.* v. (1877) 86.—Sita.
diffusum, *Phil. in Linnæa*, xxxiii. (1864–65) 191.—Chili.
dimorphum, *Phil. l. c.* xxix. (1857–58) 16.—Chili.
elongatum, A. DC. in DC. Prod. x. 124 = Mertensia elongata.
falcatum, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 64.—Reg. Argent.
Fetiowi, *Regel, in Act. Hort. Petrop.* vi. (1879) 840.—Turkestan.
filiforme, *Phil. Fl. Atac.* 89.—Chili.
floribundum, Torr. in Bot. Mex. Bound. 140 = Krynitzkia floribunda.
fruticosum, *Phil. in Linnæa*, xxix. (1857–58) 15.—Chili.
fruticulosum, Klotzsch, in Bot. Ergeb. Waldem. Reise, 96 = strictum.
fulvescentescens, A. Gray, in Proc. Am. Acad. x. (1874) 91 = Krynitzkia fulvocanescens.
fulvescentescens, S. Watz. l. c. xviii. (1883) 121 = Krynitzkia Jamesii.
fulvum, DC. Prod. x. 132 = Plagiobothrys rufescens.
fulvum, A. Gray, in Proc. Am. Acad. x. (1874) 87 = Plagiobothrys acrochilvus.
Gayanum, *Wedd. Chlor. And.* ii. 58.—Peruv.
Gilliesi, *Phil. in Anal. Univ. Chil.* (1872) 517.—Chili.
glaucosum, *Phil. in Linnæa*, xxxiii. (1864–65) 189.—Chili.
globuliferum, *Clos, in C. Gay, Fl. Chil.* iv. 464.—Chili.

ERITRICHIUM:—
glomeratum, DC. Prod. x. 181 = Krynitzkia glomerata.
glomeratum, A. Gray, in Am. Journ. Sc. Ser. II. xxxiv. (1862) 255 = Krynitzkia virgata.
gnaphaloides, *A. DC. in DC. Prod.* x. 181.—Chili.
Gullisleni, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 408 = radicans.
Hacquetii, Koch, Syn. Fl. Germ. ed. I. 507 = nanum.
heliotropioides, Torr. in Bot. Mex. Bound. 140 = Krynitzkia heliotropioides.
hispidum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 402 = Krynitzkia ramosa.
hispidum, *Phil. Fl. Atac.* 88.—Chili.
holopterum, A. Gray, in Proc. Am. Acad. xii. (1877) 81 = Krynitzkia holoptera.
humile, *DC. Prod.* x. 183.—Peruv.
incanum, A. DC. in DC. l. c. 127 = pectinatum.
intermedium, A. Gray, in Proc. Am. Acad. xvii. (1881–82) 225 = Krynitzkia intermedia.
involucratum, *Phil. in Anal. Univ. Chil.* (1878) 517. —Chili.
Jacquemontii, Decne. in Jacquem. Voy. Bot. 122. t. 127 = strictum.
Jamesii, Torr. in Marcy, Rep. 294 = Krynitzkia Jamesii.
japonicum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 96 = Trigonotis peduncularis.
jenisseense, *Turcz. ex DC. Prod.* x. 126.—Sibir. alt.
Kingii, S. Wats. in Bot. King's Exped. 243. t. 28 = Plagiobothrys Kingii.
latifolium, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 407.—Soongaria.
latifolium, Rupr. Beitr. Pfl. Russl. ii. 50 = villosum.
leiocarpum, S. Wats. in Bot. King's Exped. 244 = Krynitzkia Torreyana.
leucophaeum, A. DC. in DC. Prod. x. 129 = Krynitzkia leucophaea.
limense, *A. DC. l. c.* 183.—Peruv.
lineare, DC. l. c. 181 = Krynitzkia linearis.
linifolium, Wedd. Chlor. And. ii. 89 = Krynitzkia linifolia.
longifolium, Decne. in Jacquem. Voy. Bot. 124. t. 129 = strictum.
longisetum, *Phil. in Linnaea*, xxxiii. (1864–65) 189.— Chili.
Maackii, *Maxim. Prim. Fl. Amur.* 208.—Mandshuria.
Mandonii, *Bail*, in *Journ. Linn. Soc.* xxii. (1885) 51. —Bolivia.
micranthum, *Phil. Fl. Atac.* 88.—Chili.
micranthum, Torr. in Bot. Mex. Bound. 141 = Krynitzkia micrantha.
microcarpum, DC. Prod. x. 123 = Trigonotis microcarpa.
micromeris, A. Gray, in Proc. Am. Acad. xix. (1883) 90 = Krynitzkia micromeris.
minutiflorum, *Phil. in Linnaea*, xxxiii. (1864–65) 190. —Chili.
mollis, A. Gray, in Proc. Am. Acad. xix. (1883) 89 = Krynitzkia mollis.
multicaule, DC. Prod. x. 128 = Trigonotis multicaulis.
multicaule, Torr. in Bot. Mex. Bound. 140 = Krynitzkia Jamesii.
Munroi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 168.—Ind. or.
muricatum, *A. DC. in DC. Prod.* x. 182.—Chili.
muriculatum, A. DC. l. c. 132 = Krynitzkia muricolata.
muriculatum, Torr. in Bot. U. St. Expl. Exped. ii. 416. t. 18 = Krynitzkia ambigua.
myosotideum, Maxim. Prim. Fl. Amur. 208 = Trigonotis myosotidea.
nanum, *Schrad. in Comm. Getting.* iv. (1820) 186, in obs.; *Asperifol.* 16.—Reg. bor. et frigid.
nemorosum, DC. Prod. x. 128 = Mertensia racemosa.
nothofilosum, A. Gray, in Proc. Am. Acad. xvii. (1881–82) 227 = Plagiobothrys nothofilvus.
obovatum, *DC. Prod.* x. 128 = Krynitzkia obovata.
ovalifolium, DC. l. c. 123 = Trigonotis ovalifolia.
oxycarpum, A. Gray, in Proc. Am. Acad. x. (1874) 58 = Krynitzkia oxycarpa.
oxygonum, A. Gray, l. c. xix. (1883) 89 = Krynitzkia oxygona.
pachnophilum, *Wedd. Chlor. And.* ii. 87.—Reg. Andin.
parviflorum, *Phil. Fl. Atac.* 89.—Chili.
patens, Decne. in Jacquem. Voy. Bot. 125 = strictum.
pauciflorum, DC. Prod. x. 127 = rupestre.
pectinatum, *DC. Prod.* x. 127.—Sibir. altaic.
pedunculare, DC. l. c. 123 = Trigonotis pedunculoris.

ERITRICHIUM:—
persicum, *Boiss.* Diagn. Ser. I. xi. 129.—Orient.
phaeocalicis, *Clos, in C. Gay, Fl. Chil.* iv. 468.—Chili.
plebeium, A. DC. in DC. Prod. x. 128 = Krynitzkia plebeia.
plebeium, Torr. in Pacif. Rail. Rep. iv. 124 = Krynitzkia Scouleri.
pratense, *Phil. in Linnaea*, xxxiii. (1864–65) 193.—Chili.
primuloides, Decne. in Jacquem. Voy. Bot. 124. t. 128 = Mertensia primuloides.
procumbens, DC. Prod. x. 133 = Plagiobothrys procumbens.
pterocaryum, Torr. in Bot. Mex. Bound. 142 = Krynitzkia pterocarya.
pusillum, Torr. & Gray, in Pacif. Rail. Rep. ii. 171 = Krynitzkia pusilla.
pustulosum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 164.—Reg. Himal.
pygmaeum, *C. B. Clarke, l. c.* 165.—Reg. Himal.
pygmaeum, Wedd. Chlor. And. ii. 89 = Krynitzkia linifolia.
racemosum, S. Wats. ex A. Gray, in Proc. Am. Acad. xvii. (1881–82) 226 = Krynitzkia ramosissima.
radicans, DC. Prod. x. 126 = Trigonotis radicans.
ramosum, A. DC. in DC. l. c. 129 = Krynitzkia ramosa.
rotundifolium, DC. l. c. 132 = Trigonotis rotundifolia.
rupestre, *Bunge, Verz. Suppl. Fl. Alt.* 14.—Sibir. altaic.
Scouleri, A. DC. in DC. Prod. x. 130 = Krynitzkia Scouleri.
secundiflorum, DC. l. c. 124 = Mertensia echioides.
sericeum, Aitch. in Journ. Linn. Soc. xix. (1882) 178 = strictum.
sericeum, *DC. Prod.* x. 126.—Sibir. baical.
sessiliflorum, *DC. l. c.* 133.—Am. austr.
setosissimum, A. Gray, in Proc. Am. Acad. xii. (1877) 80 = Krynitzkia setosissima.
spathulatum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 164.—Ind. or.
strictum, *Decne. in Jacquem. Voy. Bot.* 125.—Ind. or.
strictum, *Phil. Fl. Atac.* 89.—Chili.
subamoenum, *Phil. l. c.*—Chili.
tenellum, A. Gray, in Proc. Am. Acad. x. (1874) 57 = Plagiobothrys tenellus.
tenuicaule, *Phil. in Linnaea*, xxix. (1857–58) 16.— Chili.
tenuifolium, Schlecht. ex A. Gray, in Proc. Am. Acad. xx. (1885) 267 = Krynitzkia tenuifolia.
torgilonense, A. Kern. Sched. Fl. Exsicc. i. 52 = seq.
torgiloviense, *Endl. ex Putterl. in T. Nees, Gen. Pl. Fl. Germ.* n. 59*. *fasc.* xvi. f. 27–32 = nanum.
texanum, A. DC. in DC. Prod. x. 130 = Krynitzkia texana.
tibeticum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 165.—Tibet.
tinctorium, DC. Prod. x. 128 = Plagiobothrys tinctorius.
Torreyi, A. Gray, in Proc. Am. Acad. x. (1874) 58 = Plagiobothrys Torreyi.
turkestanicum, *French. in Ann. Sc. Nat. Sér.* VI. xviii. (1883) 216.—Turkestan.
uliginosum, *Phil. in Anal. Univ. Chil.* (1878) 519.— Chili.
verrucosum, *Phil. in Linnaea*, xxix. (1857–58) 17.— Chili.
villosum, *Bunge, Verz. Suppl. Fl. Alt.* 14.—Rossia arct.
villosum, DC. Prod. x. 126 = hastilobum.
virgatum, Porter, in Hayden, Geol. Rep. (1870) 479 = Krynitzkia virgata.
Volckmanni, *Phil. in Anal. Univ. Chil.* (1861) i. 54; *et Linnaea*, xxxiii. (1864–65) 188.—Chili.
Walpersii, *Wedd. Chlor. And.* ii. 90.—Peruv.

ERITRONIUM, Soop. Fl. Carn. ed. I. 385 (1760), ophalm. = Erythronium, Linn. (Liliac.).

ERIUDAPHUS, Nees, in Eckl. & Zeyh. Enum. 271 (1836) = Sociopta, Schreb. (Bixin.).
Classianus, Balli. Adansonia, i. (1860–61) 151.— Hab.?
Ecklonii, Nees, in Eckl. & Zeyh. Enum. 271 (= *Phaberos Ecklonii*).— Afr. austr.
Mundtii, Nees, l. c. 272 (= *Phaberos Mundtii*).—Afr. austr.
serratus, Harv. Gen. S. Afr. Pl. 417.—Afr. austr.
Zeyheri, Nees, in Eckl. & Zeyh. Enum. 272 (= *Phaberos Zeyheri*).—Afr. austr.

ERLANGEA, Sch. Bip. in Flora, xxxvi. (1853) 84. *COMPOSITAE*, Benth. & Hook. f. ii. 225.
JARDINIA, Sch. Bip. in Flora, xxxvi. (1853) 84, in syn.
plumosa, *Sch. Bip. l. c.*—Afr. trop.

ERMANIA, Cham. in Linnaea, vi. (1831) 533 = Parrya, R. Br. (Crucif.).
parryoides, Cham. l. c.= P. Ermani.

ERMANNIA, Endl. Gen. 865, in syn. (1839) = prec.

ERNDELIA, Neck. Elem. ii. 325 (1790) = Passiflora, Linn.
reflexiflora, Rafin. Fl. Tellur. iv. 104 = P. reflexiflora.

ERNDLIA, Gleske, Pracl. Linn. 199, 209 (1792) = Curcurma, Linn. (Scitamin.).
Zerumbet, Gleske, ll. cc. = C. Zedoaria.

ERNESTIA, DC. Prod. tii. 121 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 740.
DICHAETANDRA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 91.
cordifolia, *Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 36. t. 2. f. 2 b.—Am. austr.
Goudotii, *Triana, l. c.*—N. Granat.
quadriseta, *Berg, ex Triana, l. c.* 35. t. 2. f. 2 a.— Peruv.
tenella, *DC. Prod.* iii. 121.—N. Granat.

ERNODEA, Sw. Prod. Veg. Ind. Occ. 29 (1788). *RUBIACEAE*, Benth. & Hook. f. ii. 143.
KNOXIA, P. Br. Hist. Jamaic. 140 (1754).
calabrica, Link, Enum. Hort. Berol. i. 135 = Putoria calabrica.
litoralis, *Sw. Prod. Veg. Ind. Occ.* 29.—Ind. occ.
maritima, Sibth. & Sm. Fl. Graec. Prod. i. 96 = Putoria calabrica.
nepalensis, Spreng. Syst. iv. Cur. Post. 40 = Knoxia corymbosa.
pedunculata, Poir. Encyc. Suppl. ii. 581 = Isidorea amoena.
pungens, Lam. Tabl. Encyc. i. 276 = Isidorea amoena.

ERNSTINGIA, Scop. Introd. 322 (1777) = Ratonia, DC. (Sapindac.).

EROBATHOS, Spach, Hist. Vég. Phan. vii. 301 (1839) = Rigella, Turcz. (Ranunculac.).
coarctatum, Spach, l. c. 304 = N. coarctata.
damascenum, Spach, l. c. 303 = N. damascena.

EROBATOS, Reichb. Handb. 277 (1837) = prec.

EROCHLOA, Stend. Syn. Pl. Gram. 373 (1855) = seq.

EROCHLOË, Rafin. Neogenyt. 4 (1838) = Zizagrostis, Beauv. (Gramin.).
amabilis, Rafin. in Ser. Bull. Bot. i. 221 = E. amabilis.
spectabilis, Rafin. l. c. = E. amabilis.

ERODENDRON, Medsen. in DC. Prod. xiv. 290, in syn. (1856) = seq.

ERODENDRUM, Salisb. Parad. Lond. t. 67 (1807) = Protea, Linn.
acutifolium, Knight, Prot. 49 = P. roseoa.
amplexicaule, Salisb. Parad. Lond. t. 67 = P. amplexicaulis.
angustum, Knight, Prot. 89 = P. grandiflora.
bombycinum, Knight, Prot. 47 = P. bombycina.
Schnlt. Syst. iii. Mant. 268 = P. lanuginosa.
conspicuum, Knight, Prot. 48 = P. turbinifolora.
coronarium, Knight, l. c. 41.—Afr. austr.
cynaroides, Knight, l. c. 43 = P. cynaroides.
echinatum, Knight, l. c. 41 = P. laxifolia.
fastidium, Knight, l. c. 40 = P. teneax.
formosum, Salisb. Parad. Lond. t. 76 = P. formosa.
glaucophyllum, Knight, Prot. 47 = P. glaucophylla.
grandiflorum, Knight, l. c. 43 = P. grandiflora.
hallmiifolium, Knight, l. c. 49.—Afr. austr.
holosericeum, Knight, l. c. 40 = P. speciosa.
incanophilum, Knight, l. c. 37 = P. incanopha.
lignifolium, Knight, l. c. 36 = P. longifolia.
linnaeifolium, Knight, l. c. 47 = P. nanuna.
longifolium, Knight, l. c. 46 = P. teretifolia.

ERODENDRUM :—
longiperum, Knight, Prot. 85 = Protea longifolia.
lorifolium, Knight, l. c. 41.—Afr. austr.
magnificum, Knight, l. c. 27 = P. speciosa.
mellifluum, Knight, l. c. 34 = P. mellifera.
mucronifolium, Knight, l. c. 48 = P. mucronifolia.
neriifolium, Knight, l. c. 40 = P. Lepidocarpon.
obtusum, Knight, l. c. 38 = P. speciosa.
ovmulum, Knight, l. c.—Afr. austr.
paucniflorum, Knight, l. c. 46 = P. canaliculata.
pinifolium, Knight, l. c. 45.—Afr. austr.
pulchellum, Knight, l. c. 36 = P. pulchella.
radulæfolium, Knight, l. c. 44 = P. Radula ?
restiniifolium, Knight, l. c. 45.—Afr. austr.
scolopendrifolium, Knight, l. c. 43 = P. Scolopendrium ?
scolymifiorum, Knight, l. c. 48 = P. Scolymus.
serpentifolium, Knight, l. c. 44 = P. repens.
speciosum, Knight, l. c. 39 = P. speciosa.
staticifolium, Knight, l. c. 45 = P. staticefolia ?
tenax, Salisb. Parad. t. 70 = P. tenax.
turbiniflorum, Salisb. l. c. t. 108 = P. turbiniflora.
umbonale, Knight, Prot. 35 = P. longifolia.
umbonale, Salisb. ex Steud. Nom. ed. II. i. 589 = P. longiflora.

ERODION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = **Erodium**, L'Hérit. (Geran.).
botrydium, St. Lag. l. c. = Erodium Botrys.
malachioides, St. Lag. l. c.; et vii. (1881) 183 = Erodium malachioides.

ERODIOPHYLLUM, F. Muell. Fragm. ix. 119 (1876). *COMPOSITAE*, Benth. & Hook. f. ii. 1223.
Elderi, *F. Muell. l. c.* 120.—Austral.

ERODIUM, L'Hérit. Geran. t. 1 ad 6 (1787). *GERANIACEAE*, Benth. & Hook. f. i. 272.
ERODION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135.
MYRRHINA, Rupr. in Mém. Acad. Pétersb. Sér. VII. xv. n. 11. (1869) Fl. Sec. 277.
SCOLOPACIUM, Eckl. & Zeyh. Enum. 89 (1835).
absinthoides, Sibth. & Sm. Fl. Graec. t. 652 = chrysanthum.
absinthoides, *Willd. Sp. Pl.* iii. 627.—Macedon.; As. Min.
aegyptiacum, *Boiss. Diagn.* Ser. II. i. 111.—Aegypt.; Syria.
affine, Fenzl, ex Steud. Nom. ed. II. i. 589 = Friedrichsthalianum.
affine, Tenore, Ind. Sem. Hort. Neap. (1830) 13; ex Fl. Neap. iv. 95 = laciniatum.
alsbidum, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1887) 111 = cicutarium.
alsiefolium, Moris, in Mem. Acc. Torin. xxxviii. (1835) 29 = alnifolium.
alexandrinum, *Delile, Fl. Aeg. Illustr.* 63.—Aegypt.
alleticklcum, Steud. ex A. Rich. Tent. Fl. Abyss. i. 116 = cicutarium.
alnifolium, *Guss. Fl. Sic. Prod.* ii. 307.—Sicili.
alpinum, Hort. Par. [Desf. Tabl. ed. II. 166 ?] = romanum.
alpinum, *L'Hérit. Geran.* t. 3.—Graecia; Ital.
alpinum, Sibth. & Sm. Fl. Graec. t. 659 = hirtum.
alsiefolium, *Delile, Ind. Sem. Hort. Monsp.* (1847) 7; *et in Ann. Sc. Nat. Sér.* III. x. (1848) 329.—Hab.?
althaeoides, Jord. Pugill. Pl. Nov. (1852) 41 = malacoides.
amanum, *Boiss. & Kotschy, in Boiss. Fl. Orient.* i. 887.—Syria.
amblyrum, *Pomel, Nouv. Mat. Fl. Atl.* 338.—Alger.
angulatum, Pomel, l. c. 341 = malacoides.
anthemifolium, *Bieb. Fl. Taur. Cauc.* ii. 131.—Reg. Caucas.
anthemifolium, Tenore, Syll. 330 = alpinum.
apenninum, Tenore, Syll. App. iv. 35 = alpinum.
arabicum, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 336 = alnifolium.
arboreum, *Willd. Sp. Pl.* iii. 638.—Afr. bor.
arduanum, *Willd. l. c.* 637.—Afr. austr.
arenarium, Jord. Pugill. Pl. Nov. (1852) 44 = cicutarium.
arenarium, Pomel, Nouv. Mat. Fl. Atl. 340 = laciniatum.

ERODIUM:—
arenarium, Pourr. ex Lange, in Kjoeb. Vidensk. Meddel. (1846) 192 = Jacquinianum.
asplenioides, Boiss. Voy. Bot. Espagne, 128 = Boissieri.
asplenioides, *Willd. Sp. Pl.* iii. 635.—Afr. bor.
astragaloides, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 130.—Hispan.
atlanticum, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 56; ex. (1878) 343.—Marocc.
atomarium, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1848) 423 = cicutarium.
australe, Salzm. ex Nym. Consp. 189 = moschatum.
Barrandieanum, Loret, in Loret & Barrand. Fl. Monsp. i. 123, in obs. = romanum.
bipinnatum, Willd. Sp. Pl. iii. 628 = cicutarium.
Bocconi, Steud. Nom. ed. II. i. 589 = chamaedryoides.
Bocconi, Viv. Prod. Fl. Cors. App. 5 = maritimum.
Boissieri, *Coss. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 364.—Hispan.
Boraeanum, Jord. Pugill. 47 = cicutarium.
Botrys, *Bertol. Amoen. Ital.* 35.—Reg. Mediterr.
Botrys, Hohen. in Bull. Soc. Nat. Mosc. (1838) 895 = oxythinchum.
Botrys, C. A. Mey. Verz. Pfl. Cauc. 209 = laciniatum.
Bourgaei, Boiss. Fl. Orient. i. 892 = tmolenm.
Bovei, Delile, Ind. Sem. Hort. Monsp. (1838) [6]; ex Linnaea, xiii (1839) Litt. 104 = alnifolium.
Bowei, Steud. Nom. ed. II. i. 589 = praec.
brevicaule, Bertol. in Nov. Comm. Acad. Bonon. vi. (1844) 229. t. 9. f. 2 = ciconium.
bryoniaefolium, *Boiss. Diagn.* Ser. I. i. 61.—Aegypt.; Persia.
cadmeum, Jaub. & Spach, Illustr. ii. 107. t. 189 = laciniatum.
Campsianum, Coss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 535 = astragaloides.
Cariati, Jord. Pugill. Fl. Nov. (1852) 49 = romanum.
carneum, Jord. l. c. 47 = cicutarium.
carvifolium, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 9.—Hispan.
cascalhotium, Sweet. Geran. t. 6 = romanum.
Cavanillesii, *Willd. Pl. Haloph.* 111.—Hispan.
cedronum, *Schott, in Oestr. Bot. Wochenbl.* iv. (1854) 301.—Cilicia.
Charophyllum, Steud. Nom. ed. I. 314; Coss. Notes Crit. ii. 32 = cicutarium.
chamaedryoides, *L'Hérit. Geran.* t. 6.—Ins. Balear.; Corsica.
cheilanthifolium, Boiss. Elench. 27 = trichomanefolium.
chium, Sibth. & Sm. Fl. Graec. t. 657 = laciniatum.
chium, *Willd. Phyt.* 10.—Reg. Mediterr.
chrysanthum, *L'Herit. ex DC. Prod.* i. 645.—Graecia.
chumbalense, Munro, ex Wight, Ic. t. 1074 = Monsonia senegalensis.
cicutarium, Janka, ex Nym. Consp. 189 = E. Botrys.
ciconium, *Willd. Sp. Pl.* iii. 629.—Reg. Mediterr.; Orient.
cicutarifolium, Salisb. Prod. 311 = cicutarium.
cicutarium, *L'Hérit. ex Ait. Hort. Kew.* ed. I. ii. 414. —Reg. Mediterr.; Aa. med.
cinerarium, Moris, in Mem. Acc. Torin. xxxviii. (1835) 29 = laciniatum.
cinereum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 115.—As. Min.
corsicum, Steud. Nom. ed. II. i. 589, sphalm. = *Enodium corsicum*.
commixtum, Jord. ex Schultz. Arch. Fl. Fr. et Allem. 156 = cicutarium.
communitatum, Todaro, Ind. Sem. Hort. Panorm. (1868) 35 = malachoides.
conicum, *Lenan, in DC. Fl. Fr.* iv. 842.—Corsic.; Sardin.
crassifolium, *Ait. Hort. Kew.* ii. 414.—Afr. bor.; Cyprus.
crassifolium, DC. Prod. i. 646 = hirtum.
crassum, Steud. Nom. ed. I. 314 = crassifolium.
crenatum, Pomel, Nouv. Mat. Fl. Atl. 341 = alnifolium ?
crispum, Lapeyr. Hist. Abr. Pl. Pyr. 394 = petraeum.
cuneatum, Viv. Prod. Fl. Cors. App. 5 = chium.
Cygnorum, *Nees, in Lehm. Pl. Preiss.* i. 162.—Austral.
daucoides, *Boiss. Elench.* 28.—Hispan.
decipiens, Delile, ex Tenore, Ind. Sem. Hort. Neap. (1838) 18; et Fl. Neap iv. 95 = laciniatum.
dentatum, Dum. in Bull. Soc. Bot. Belg. iv. (1865) 345 = moschatum.
dryadifolium, Salisb. Prod. 311 = chamaedryoides.
elongatum, Todaro, ex Parl. Fl. Ital. v. 246 = alnifolium.

ERODIUM:—
fallax, Jord. Cat. Gr. Réc. Gren. (1849) 14 = romanum.
filicinum, *Pomel, Nouv. Mat. Fl. Atl.* 339.—Alger.
foetidum, L'Hérit. ex DC. Prod. iv. 645 = petraeum.
Friedrichsthalianum, *Steud. Nom.* ed. II. i. 589.—Graecia.
fumarioides, Stev. in Mém. Soc. Nat. Mosc. iv. (1812) 49 = absinthoides.
Gaillardoti, *Boiss. Diagn.* Ser. II. vi. 41.—Syria.
Gasparrinii, Guss. Fl. Sic. Prod. ii. 801 = E. Botrys.
gruinum, Desf. Tabl. ed. II. 166 = hymenodes.
gruinum, *Murphy, Fl. Algér.* 71.—Algeria.
gruoides, *St. Hil. Fl. Bras. Merid.* i. 80.—Am. austr.
glabellum, *Delile, ex Steud. Nom.* ed. I. 314.—Aegypt
glaucum, Pomel, Nouv. Mat. Fl. Atl. 342 = glaucophyllum ?
glandulosum, Dum. in Bull. Soc. Bot. Belg. iv. (1865) 345 = sabulicola.
glandulosum, Willd. Sp. Pl. iii. 628 = macradenum.
glaucophyllum, *Ait. Hort. Kew.* ed. I. ii. 416.—Afr. bor.; Arab.
glaucophyllum, Bové, ex Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 285 = Monsonia nivea.
glauco-virens, *Lojac. in Nuovist. Sicil.* i. (1882) 105. —Sicil.
glutinosum, Dulac, Fl. Hautes-Pyr. 238 = malachoides.
glutinosum, Dum. in Bull. Soc. Bot. Belg. iv. (1865) 345 = cicutarium.
graveolens, Lapeyr. Hist. Abr. Pl. Pyr. 390 = macradenum.
gruinum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 415.—Europ. austr.; Orient.
gruinum, Tenore, Prod. Fl. Nap. 39 = laciniatum.
Guicciardii, *Heldr. in Boiss. Diagn.* Ser. II. vi. 40.—Graecia.
Gussoni, *Tenore, Prod. Fl. Nap.* 39.—Ital.
guttatum, Ledeb. Fl. Ross. i. 478 = oxythinchum.
guttatum, *Willd. Sp. Pl.* iii. 636.—Reg. Mediterr. occ. hellotropioides, *Willd. Sp. Pl.* iii. 638.—Hab.?
himalayanum, Royle, Illustr. Bot. Himal. 150 = cicutarium?
hirsutum, Schur, in Verh. Naturf. Ver. Brucnn, xv. ii. (1877) 184 = cicutarium.
hirtum, Jacq. Eclog. i. 82. t. 58 = Jacquinianum.
hirtum, Willd. ex Steud. Nom. ed. II. i. 589 = morainense.
hirtum, *Willd. Sp. Pl.* iii. 632.—Afr. bor.; Arab.; Syria.
hispidum, Presl, Fl. Sic 208 = laciniatum.
Hoefliauum, *C. A. Mey. in Mém. Acad. Pétersb. Sér.* VI. Sc. Nat. vii. (1855) 3.—Sibiria.
Hohenackeri, Ledeb. Fl. Ross. i. 478 = oxythinchum.
Hussoni, *Boiss. Diagn.* Ser. I. viii. 119.—Aegypt.; Arabia.
hymenodes, *L'Hérit. Geran.* t. 4.—Alger.
incarnatum, *L'Hérit. l. c.* t. 5.—Afr. austr.
involucratum, Kunze, in Flora, xxix. (1846) 740 = laciniatum.
Jacquinianum, *Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop. ix. Suppl.* 11.—Europ. austr.; Persia.
Kotschyanum, Boiss. Fl. Orient. i. 889 = cedronum.
Kotschyanum, Koechel, ex Boiss. l. c. 896 = bryoniaefolium.
lacinlatum, *Willd. Sp. Pl.* iii. 633.—Reg. Mediterr.; Persia.
Lebelii, *Jord. Pugill. Pl. Nov.* (1852) 48.—Gallia.
leucanthum, *Boiss. Diagn.* Ser. I. i. 60.—As. Min.
leucophyllum, Presl, Bot. Bemerk. 26 = laciniatum.
litoreum, Badaro, in Moretti, Bot. Ital. 26 = chium.
littoreum, *Léman, ex DC. Fl. Fr.* iv. 848.—Reg. Mediterr. occ.
longicaule, Todaro, Ind. Sem. Hort. Panorm. (1868) 35 = alnifolium.
lucidum, Lapeyr. Hist. Abr. Pl. Pyr. 394 = petraeum.
macradenum, *L'Hérit. Geran.* t. 1.—Mont. Pyren.
macrophyllum, *Hook. & Arn. Bot. Beech. Voy.* 357.—Am. bor. occ.
maculatum, Salzm. ex Presl, Bot. Bemerk. 26 = cicutarium.
malachoides, Bové, ex Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 286 = alnifolium.
malachoides, *Willd. Phyt.* 10.—Reg. Mediterr.; Orient.
malacitanum, Amo, Fl. Iber. vi. 68 = Jacquinianum.
malopoides, Loisel. Fl. Gall. ed. II. ii. 88 = cicutarium.
malopoides, Willd. Sp. Pl. iii. 640 = alnifolium.
malvaceum, Jord. ex Loret & Barrand. Fl. Monsp. 122 = malachoides.
Manescavi, *Coss. in Ann. Sc Nat.* Sér. III. vii. (1847) 206. t. 11.—Mont. Pyren.

ERODIUM :—
Marmocci, *Peri. Fl. Ital.* v. 208.—Italia.
maritimum, Eckl. & Zeyh. Enum. 59 = Pelargonium chamaedrifolium.
maritimum, *Sm. Fl. Brit.* ii. 732.—Europ.
mauritanicum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 209.—Afr. bor.
medicense, *Battand. in Bull. Soc. Bot. Fr.* xxx. (1883) 264.—Alger.
melanostigma, Mart. ex Delile, Ind. Sem. Hort. Monsp. (1838); ex Linnæa, xiii. (1839) Litt. 104 = cicutarium.
micropetalum, *Boiss. & Hausskn. in Boiss. Fl. Orient.* i. 889.—As. Min.
microphyllum, *Pomel, Nouv. Mat. Fl. Atl.* 339.—Alger.
millefolium, *Willd. ex H. B. & K. Nov. Gen. et Sp.* v. 238.—Ecuador.
minutiflorum, Godr. in Mém. Soc. Sc. Nat. Cherb. xix. (1875) 173 = cicutarium.
monsoniodes, Steud. ex Boiss. Fl. Orient. i. 892 = grulnum.
montanum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iii. (1856) 737.—Algeria.
moranense, *Willd. ex H. B. & K. Nov. Gen. et Sp.* v. 238.—Mexic.
moschatum, *L'Hérit. ex Ait. Hort. Kew.* ed. I. ii. 414. —Reg. Mediterr.; Orient.
multicaule, Link, Enum. Hort. Berol. ii. 184 = serotinum. Munbyanum, *Boiss. ex Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 283.—Alger.
murcicum, Willd. Sp. Pl. iii. 636 = chium.
narbonnense, Delile, Ind. Sem. Hort. Monsp. (1836); ex Ann. Sc. Nat. Sér. II. vii. (1837) 286 = chium.
Neitzschii, Janka, in Oestr. Bot. Zeitschr. xvii. (1867) 101 = E. Botrys.
nervulosum, *L'Hérit. ex DC. Prod.* i. 643.—Sicil.
neuradaefolium, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1858) 425 = aggregatum.
nitagiricum, Schlecht. ex Hook. f. Fl. Brit. Ind. i. 485 = Pelargonium ossularioides.
niveum, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 285 = Monsonia nivea.
numidicum, Salzm. ex Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1858) 421 = scandicinum.
obliquifolium, Tenore, Ind. Sem. Hort. Nasp. (1830) 13; ex Fl. Neap. iv. 95 = alnifolium.
obtusifolium, Karel. ex Ledeb. Fl. Ross. i. 477 = oxyrhinchum.
olympicum, Clem. Sertul. Or. 17 = absinthoides.
oxyrhinchum, *Bieb. Fl. Taur. Cauc.* ii. 133.—Orient.; Turkestan.
pachyrhizum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ix. (1862) 432.—Alger.
pallidiflorum, Jord. Cat. Gr. Réc. Gren. (1849) 2 = cicutarium.
parviflorum, Jord. l. c. (1849) 15 = cicutarium.
pauciflorum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 419.—Chili.
pelargoniiflorum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 118.—Orient.
pentadenum, Dalac, Fl. Hautes-Pyr. 237 = cicutarium.
peristeroides, *Turcz. in Bull. Soc. Nat. Mosc.* (1863) i. 592.—N. Zel.
petraeum, Sibth. & Sm. Fl. Græc. t. 651 = absinthoides.
petraeum, Tenore, ex Nym. Consp. 141 = alpinum.
petraeum, *Willd. Sp. Pl.* iii. 625.—Reg. Mediterr. occ.
Petrosilenum, L'Hérit. ex DC. Prod. i. 647 = cicutarium.
pilosum, *Menyh. Kalocsa Vidh. Novény.* 53 (= sec. !). —Hungar.
pilosum, Steud. Nom. ed. I. 814 = cicutarium.
pilosum, Walp. Rep. i. 446 = laciniatum.
pimpinellaefolium, Sibth. Fl. Oxon. 211 = cicutarium.
pimpinellifolium, Willd. Sp. Pl. iii. 630 = præc.
praecox, Willd. Sp. Pl. iii. 631 = cicutarium.
praetermissum, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 181 = cicutarium.
primulaceum, *Welw. ex Lange, Ind. Sem. Hort. Bot. Haun.* (1865) 24.—*Cf. Linnæa*, xxviii. (1856) 859. —Hispan.
procumbens, Jord. Pugill. Pl. Nov. (1852) 51 = romanum.
pulchellum, Karel. ex Ledeb. Fl. Ross. i. 476 = laciniatum.
pulchrum, Salisb. Prod. 311 = incarnatum.
pulverulentum, Willd. Sp. Pl. iii. 632 = laciniatum.
punctatum, Steud. Nom. ed. II. i. 679, in syn. sphalm. = Pelargonium punctatum.

ERODIUM :—
pyramidatum, Presl, Bot. Bemerk 26 = laciniatum.
radicatum, Lapeyr. Fl. Pyr. t. t. 1 = macradenum.
reflexum, *Delile, Fl. Aeg. Illustr.* 68.—Aegypt.; Arab.
Reichardi, DC. Prod. i. 649 = chamaedryoides.
ribifolium, Jacq. Coll. Suppl. 121 = malachoides.
Robertianum, C. A. Mey. ex Nym. Consp. 140 = moschatum.
romanum, *Willd. Sp. Pl.* iii. 630.—Europ. austr.; Syria.
rupestre, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 538 = supracanum.
rupicolum, *Boiss. Voy. Bot. Espagne*, ii. 724.—Hispan.
ruthenicum, Bieb. Fl. Kar. Cent. 1. t. 48 = serotinum.
sabuliculum, Jord. ex Nym. Consp. 140 = cicutarium.
sabulicolum, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* iii. 537.—Hispan.
Salzmanni, Boiss. & Reut. ex Nym. l. c. 140 = primulaceum.
Salzmanni, Delile, Ind. Sem. Hort. Monsp. (1838) 6.— Cf. Linnæa, xiii. (1839) Litt. 104 = cicutarium.
scandicinum, *Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.)* i. (1858) 421.—Hispan.
sebaceum, *Delile, Ind. Sem. Hort. Monsp.* (1838) 6.— *Cf. Linnæa*, xiii. (1839) *Litt.* 105.—Galloprov.
Semenovii, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) i. 553.—As. centr.
Sempervivum, Roxb. in Beatson, St. Helena Tracts, 108 = Pelargonium Cotyledonis.
serotinum, *Stev. in Mém. Acad. Pétersb.* iii. (1809–10) 297. t. 15. f. 2.—Reg. Caucas.
Sibthorpianum, Boiss. Diagn. Ser. I. i. 60 = absinthoides.
solutinum, Todaro, Ind. Sem. Hort. Panorm. (1868) 26 = laciniatum.
speciosum, Salisb. Prod. 312 = Monsonia speciosa.
staphylinum, Bertol. Fl. Ital. vii. 135 = Jacquinianum.
stellatum, Delile, Ind. Sem. Hort. Monsp. (1838) 6.— Cf. Linnæa, xiii. (1839) Litt. 104 = primulaceum.
Stephanianum, *Willd. Sp. Pl.* iii. 625.—Reg. Himal.; As. bor.
Steveni, *Bieb. Fl. Taur. Cauc.* ii. 132.—Reg. Caucas.
stipaceum, *Eugenio & Hook. f. in Fl. Brit. Ind.* i. 485. —Reg. Himal.
stipulaceum, Duf. ex Nym. Consp. 139 = malachoides.
strigosum, Karel. ex Ledeb. Fl. Ross. i. 475 = laciniatum.
stylatum, *Lman, ex DC. Prod* i. 646.—Hab. ?
subalbidum, Jord. Pugill. Pl. Nov. (1852) 43 = cicutarium.
subtrilobum, Jord. l. c. 42 = malachoides.
supracanum, *L'Hérit. Geran.* t. 8.—Hispan.
tataricum, *Willd. Sp. Pl.* iii. 625.—Sibir.
tenuisectum, Godr. & Gren. in Gren. & Godr. Fl. Fr. i. 311 = Jacquinianum.
tenuisectum, Lange, in Kjoeb. Vidensk. Meddel. (1865) 189 = cicutarium.
texanum, *A. Gray, Gen. Illustr.* ii. 180.—Am. bor. occ.
tibetanum, *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 484. —Reg. Himal.
tmolum, *Reut. ex Boiss. Fl. Orient.* i. 892.—As. Min.
Tolassianum, Jord. Pugill. Pl. Nov. (1852) 50 = romanum.
tordilioides, Viv. Fl. Lib. Spec. 89 = grulnum.
tordylioides, *L'Hérit. ex DC. Prod.* i. 647.—Alger.
Touchonum, *Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.)* i. (1858) 425.—Hab. ?
trichomanefolium, Iiossa. Voy. Bot. Espagne, 120. t. 27 = pro.
trichomanefolium, *L'Hérit. ex DC. Prod.* i. 645.— Syria.
trifolium, Cav. Descr. 150 = Geranium trifolium.
trilobatum, Jacq. Coll. iv. 200 = hymenodes.
trilobatum, Jord. ex Nym. Consp. 139 = malachoides.
triviale, Jord. Pugill. Pl. Nov. (1852) 43 = cicutarium.
tuberosum, Boiss. Diagn. Ser. I. viii. 118 = hirtum.
tunetanum, *G. Don, Gen. Syst.* i. 723.—Afr. bor.
undulatum, Salisb. Prod. 311 = Monsonia emarginata.
valentinum, Boiss. & Reut. ex Willk. & Lange, Prod. Fl. Hisp. iii. 538 = petraeum.
verbenaefolium, *Delile, Ind. Sem. Hort. Monsp.* (1847) 7; *et in Ann. Sc. Nat.* Sér. III. x. (1848) 229.— Hab. ?
violaefolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 592.—Ind. or.
viscosum, Salzm. ex Delile, in Linnæa, xiii. (1839) 104 = cicutarium.
viscosum, *Steud. Nom.* ed. I. 815.—Hab. ?

ERONEMA, Rafin. Fl. Tellur. iv. 126 (1836) = Cuscuta, Tourn. (Convolv.).
Robinsoni, Rafin. l. c.—Hab. ?

EROPHACA, Boiss. Voy. Bot. Espagne, ii. 176 (1840) = **Astragalus**, Tourn. (Legumin.).
bætica, Boiss. l. c. 177 = A. lusitanicus.

EROPHILA, DC. Syst. ii. 356 (1821). *CRUCIFERÆ*, Benth. & Hook. f. i. 75.
GANSBLUM, Adans. Fam. ii. 430 (1763).
affinis, *ambigens*, *andegavensis*, *Bardini*, *brevifolia*, *brevipila*, *breviscapa*, *cabillonensis*, *campestris*, *chlorotica*, *cinerea*, *clavisiformis*, *confinis*, *cuneifolia*, *curtipes*, *decipiens*, *dentata*, *fallacina*, *furcipila*, *lepida*, *leptophylla*, *lucida*, *lugdunensis*, *medioxima*, *micrantha*, *muricola*, *oblongata*, *obovata*, *occidentalis*, *Onasmi*, *patula*, *procerula*, *pulocarpa*, *pyrenaica*, *Revelieri*, *rigidula*, *rubella*, *rubricaulis*, *ruriensis*, *serrata*, *sparsipila*, *spathulæfolia*, *subintegra*, *subulata*, *subtilis*, *tenuis*, *vestita*, *virescens*, *wiesbadensis*, Jord. Diagn. 207–244, omnes = vulgaris.
americana, DC. Syst. ii. 356 = vulgaris.
Boerhaavii, Dum. Fl. Belg. 120 = vulgaris.
brachycarpa, Jord. Pugill. 9 = vulgaris.
Draba, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 952 = vulgaris.
glabrescens, Jord Pugill. 10 = vulgaris.
hirta, Jord. ex Nym. Consp. 54 = vulgaris.
hirtella, Jord. Pugill. 10 = vulgaris.
inflata, H. C. Wats. ex Hook. f. Bot. Soc. Edinb. xii. 35 = vulgaris.
Krockeri, Andrz. ex Nym. Consp. 54 = vulgaris.
macrocarpa, *Boiss. Fl. Orient.* i. 804.—As. Min.
majuscula, Jord. Pugill. 11 = vulgaris.
minima, *C. A. Mey. Verz. Pfl. Cauc.* 184.—Reg. Casp., Syria.
minutissima, DC. Syst. ii. 357 = præcox.
muscosa, DC. Syst. ii. 358 = Draba muscosa.
præcox, *DC. l. c.* 357.—Europ.; Orient.
propinqua, Jord. ex Pomel, in Bull. Soc. Bot. Fr. xviii. (1871) 220 = vulgaris.
setulosa, *Boiss. & Blanche, Diagn.* Ser. II. v. 81.— Syria.
siliquosa, Boiss. & Heldr. Diagn. Ser. I. viii. 98, in syn. = macrocarpa.
spathulata, Opiz, Seznam, 41 = vulgaris.
stenocarpa, Jord. Pugill. 11 = vulgaris.
verna, E. Mey. ex Garcke, Fl. Deutschl. ed. IV. 85 = vulgaris.
vulgaris, *DC. Syst.* ii. 356.—Europ.; Orient.

EROSMA, Booth, Cat. 118; ex Endl. Suppl. iv. ii. 24, in syn. (1847) = **Picus**, Tourn. (Urtic.).

EROSMIA, A. Juss. in Orb. Dict. xi. 254 (1840), sphalm. = **Eucosmia**, H. B. & K. (Rubiac.).

EROTEUM, Sw. Prod. Veg. Ind. Occ. 85 (1788) = **Freziera**, Sw. (1799) (Ternstroem.).
theaceides, Sw. l. c. = Cleyera theoides.
undulatum, Sw. l. c. = F. undulata.

ERPETINA, Nand. in Ann. Sc. Nat. Sér. III. xv. (1851) 299 = **Madinilla**, Gaudich. (Melastom.).
radicans, Naud. l. c. = M. Erpetina.

ERPETION, DC. ex Sweet, Brit. Flow. Gard. ii. t. 170 (1826) = **Viola**, Tourn.
Cymbalorian, DC. ex Sweet, l. c. = V. hederacea.
hederacea, G. Don, Gen. Syst. i. 836 = V. hederacea.
petiolaris, G. Don, l. c. = V. hederacea.
reniformis, Sweet, Brit. Flow. Gard. t. 170 = V. hederacea.
spathulata, G. Don, Gen. Syst. i. 835 = V. hederacea.

ERRAZURIZIA, Phil. in Anal. Univ. Chil. (1872) 689 = **Dalea**, Linn. (Legumin.).
glandulifera, Phil. l. c. = D. multifoliolata.

ERTELA, Adans. Fam. ii. 368 (1763) = **Monniera**, Linn. (Rutac.).

ERTELIA, Steud. Nom. ed. II. i. 590 (1840) = præc.

ERUCA, Tourn. ex *Adans. Fam.* ii. 419 (1763). *CRUCIFERÆ*, Benth. & Hook. f. i. 84.
EUSEBIUM, Link, Enum. Hort. Berol. ii. 174 (1822).
VELLERUCA, Pomel, Mat. Fl. Atl. 12 (1860).

ERUCA :—
alba, Noulet, Fl. Sous-Pyr. 33 = Brassica alba.
aquatica, Moench, ex Steud. Nom. ed. 1. 215, 788 = Nasturtium aquaticum.
arvensis, Noulet, Fl. Sous-Pyr. 54 = Moricandia arvensis.
aspera, Mill. Gard. Dict. ed. VIII. n. 4 = Brassica Erucastrum.
Barbarea, Lam. Fl. Fr. ii. 497 = Barbarea vulgaris.
bellidis-folia, Mill. Gard. Dict. ed. VIII. n. 2 = Diplotaxis viminea.
brevirostris, *Pomel, Nouv. Mat. Fl. Atl.* 226.—Alger.
cappadocica, *Reut. Cat. Hort. Geneo.* (1857).—As. Min.
decumbens, Moench, Meth. 257 = Diplotaxis muralis.
deserti, *Pomel, Nouv. Mat. Fl. Atl.* 226.—Alger.
elongata, Baumg. Enum. Stirp. Transs. ii. 264 = Brassica elongata.
erecta, Lag. Gen. et Sp. Nov. 20 = Brassica Tournefortii.
Erucastrum, Gaertn. Mey. & Scherb. Fl. Wett. ii. 458 = Brassica Erucastrum.
fortida, Moench, Meth. 256 = sativa.
glabrescens, Jord. Diagn. i. 193 = sativa.
grandyflora, Cav. Desc. 426 = sativa.
hispida, Cav. l. c. 425 = Brassica valentina.
hispida, DC. Syst. ii. 638 = sativa.
lanceolata, *Pomel, Nouv. Mat. Fl. Atl.* 267.—Alger.
lativalvis, *Boiss. Fl. Orient.* i. 396.—Persia.
longirostris, *Uechtr. in Oestr. Bot. Zeitschr.* (1874) 138.—Hispan.
longistyla, *Pomel, Nouv. Mat. Fl. Atl.* 268.—Alger.
muralis, Bess. Enum. Pl. Volh. 27 = Diplotaxis muralis.
muralis, Gaertn. Mey. & Scherb. Fl. Wett. ii. 459 = Diplotaxis tenuifolia.
orthoscpala, *Lange, Ind. Sem. Hort. Haun.* (1857) 28.—*Cf. Linnæa*, xxix. (1857-58) 736.—Hispan.
perennis, Mill. Gard. Dict. ed. VIII. n. 8 = Diplotaxis tenuifolia.
pervista, Jord. Diagn. i. 193 = sativa.
pinnatifida, *Pomel, Nouv. Mat. Fl. Atl.* 267.—Alger.
Ruchella, Stend. Nom. ed. II. i. 690, sphalm. = seq.
Ruchetta, Spach, Hist. Vég. Phan. vi. 355 = sativa.
sativa, *Mill. Gard. Dict.* ed. VIII. n. 1.—Reg. Mediterr.; As. occ.
sctulea, *Boiss. Diagn.* Ser. II. v. 26.—Algeria.
stenocarpa, Boiss. & Kent. Pugill. Pl. Nov. 8 = sativa.
sylvestris, Lam. Fl. Fr. ii. 497 = Brassica Erucastrum.
tanacetifolia, Mill. Gard. Dict. ed. VIII. n. 5 (tenacetifolia.) = Sisymbrium tanacetifolium.
tenuifolia, Moench, Meth. 257 = Diplotaxis tenuifolia.
vesicaria, *Cav. Desc.* 426.—Hispan.
viminea, Mill. Gard. Dict. ed. VIII. n. 6 (viminia) = Diplotaxis viminea.

ERUCAGO, Tourn. ex Adans. Fam. ii. 423 (1763) = **Bunias**, Linn. (Crucif.).
arvensis, Fourn. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 386 = B. Erucago.
aspera, Hornem. Hort. Hafn. ii. 692 = B. Erucago.
brachyptera, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 386 = B. Erucago.
campestris, Desv. Journ. Bot. iii. (1814) 168 = B. Erucago.
dentata, Moench, Meth. 278 = B. Erucago.
glandulosa, Roehl. Deutschl. Fl. ed. II. 856 ; ex Koch, Syn. Fl. Germ. ed. I. 77 = B. Erucago.
macroptera, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 386 = B. Erucago.
mexicana, Jaume St. Hil. Expos. i. 510 = B. Erucago.
monspeliensis, Cav. Desc. 418 = B. Erucago.
runcinata, Hornem. Hort. Hafn. ii. 692 = B. Erucago.

ERUCARIA, Cerv. in La Naturaleza (1870), 347. *GRAMINEAE.*
glabra, Cerv. l. c. 350 = Boutelous racemosa.
glandulosa, *Cerv. l. c.* 347.—Mexic.
hirsuta, *Cerv. l. c.* 348.—Mexic.
longifolia, *Cerv. l. c.* 350.—Mexic.
luteiscens, *Cerv. l. c.* 349.—Mexic.
mexicana, *Cerv. l. c.*—Mexic.
tetrastachia, *Cerv. l. c.* (arr. syn. tretastachia).—Mexic.
villosa, *Cerv. l. c.* 348.—Mexic.

ERUCARIA, Gaertn. Fruct. ii. 398. t. 163. f. 9 (1791). *CRUCIFERAE*, Benth. & Hook. f. i. 100.
PACHILA, Rafin. in Loud. Gard. Mag. viii. (1832) 246.
REBOUDIA, Coss. & Dur. in Bull. Soc. Bot. Fr. ii. (1856) 704.
Aegiceras, J. Gay, ex Boiss. Fl. Orient. i. 367 (= *Hussonia arvensis*).—Arab.
aleppica, Bieb. Fl. Taur. Cauc. iii. 451 = Brassica adpressa.
alepyca, *Gaertn. Fruct.* ii. 398.—Graecia ; Arab. ; Orient.
crassifolia, *Delile, Fl. Egypte*, 244 —Aegypt.
eucarioides, *C. Muell. in Walp. Ann.* vii. 178.—Alger.
grandiflora, *Boiss. Diagn.* Ser. I. vi. 18.—Persia.
hypogaea, Viv. Fl. Cors. Sp Nov. 11 = Morisia hypogaea.
hyrcanica, DC. Syst. ii. 676 = Brassica adpressa.
latifolia, DC. l. c. 676 = aleppica.
linearifolia, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 390.—Persia.
microcarpa, *Boiss. Diagn.* Ser. I. viii. 47.—Aegypt.; Arab.
Olivieri, *Spreng. Syst.* ii. 915.—Mesopotam.
persica, Gorter, ex Ledeb. Fl. Ross. i. 219 = Brassica adpressa.
persica, Hort. ex W. Baxt in Loud. Hort. Brit. Suppl. ii. 638 = Brassica incana.
tenuifolia, *DC. Syst.* ii. 675.—Hispan.
uncata, Boiss. Diagn. Ser. I. viii. 47, in syn. (= *Hussonia uncata*).—Arab.

ERUCASTRUM, Presl, Fl. Sic i. 92 (1826) = **Brassica**, Tourn. (Crucif.).
arabicum, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 35 = B. Schimperi.
boeticum, Lange, in Kjoeb. Videnak. Meddel. (1865) 70 = B. boetica.
canariense, Webb & Berth. Phyt. Canar. i. 81.—Ins. Canar.
cheiranthiflorum, Link, Handb. ii. 317 = B. Cheiranthus.
Cheiranthus, Link, l. c. = B. Cheiranthus.
Cossonianum, Reut. Cat. Hort. Genev. (1854) = B. Cossoniana.
Decandollii, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 947 = Moricandia arvensis.
elongatum, Reichb. Fl. Germ. Excurs. 694 = B. elongata.
fruticulosum, Presl, Fl. Sic. i. 92 = B. fruticulosa.
Gmelini, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 947 = B. Erucastrum !
heterophyllum, Nym. Syll. 197 = B. adpressa.
incanum, Koch, Syn. Fl. Germ. ed. I. 56 = B. adpressa.
inodorum, Reichb. Fl. Germ. Excurs. 698 = B. Erucastrum.
intermedium, Jord. Diagn. i. 191 = B. Erucastrum.
Lamarkii, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 947 = B. Erucastrum.
laxicaule virens, Boiss. Fl. Orient. i. 389.—Cilicia.
leucanthum, Coss. & Dur. in Bull. Soc. Fr. ii. (1855) 307.—Alger.
longirostre, Nym. Consp 48 = B. longirostris.
longirostris, Uechtr. in Oestr. Bot. Zeitschr. v. (1855) 308 = B. longirostre.
montanum, Hegewschw. Fl. Schw. 656 = B. Cheiranthos.
monspeliense, Reichb. Fl. Germ. Excurs. 698 = B. Erucastrum.
oleraceum, Presl, Fl. Sic. i. 93 = B. laevigata !
palustre, Vis.—Cf. Linnaea, xxviii. (1856) 305 = B. elongata !
Pollichii, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 946 = B. Erucastrum.
Pseudotanaceti, Lange, in Kjoeb. Videnak. Meddel. (1865) 79.—Hispan.
pubescens, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 861.—Hispan.
Richerii, Link, Handb. ii. 317 = B. Richerii.
sabularia, Link, l. c. = B. sabularia.
subhirsutifolium, Nym. Consp. 48.—Hispan.
Tournefortii, Link, Handb. ii. 317 = B. Tournefortii.
varium, Dur. Expl. Sc. Algér. t. 75 = B. varia.
virgatum, Presl, Fl. Sic. i. 94 = B. boetica.
Zanonii, Ball, in Bull. Soc. Bot. Fr. vii. (1860) 252.—Italia.

ERVILIA, Link, Enum. Hort. Berol. ii. 240 (1822) = **Vicia**, Tourn. (Legumin.).
canarea, Alef. in Bonplandia, ix. (1861) 126 = V. cassubica.
cassubica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = V. cassubica.
hirsuta, Opiz, Seznam, 41 = V. hirsuta.
monanthos, Opiz, l. c. = V. monanthos.
Orobus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = V. cassubica.
pinformis, Schur, l. c. = V. pisiformis.
sativa, Link, Enum. Hort. Berol. ii. 240 = V. Ervilia.
sylvatica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = V. sylvatica.
tetrasperma, Opiz, Seznam, 41 = V. gemella.
vulgaris, Godr. Fl. Lorr. i. 173 = V. hirsuta.

ERVUM, Tourn. ex Linn. Gen. ed. I. 217 (1737) = **Vicia**, Tourn. (Legumin.).
agrigentinum, Guss. ex Ser. in DC. Prod. ii. 367 = V. lutcosa.
alpestre, Trautv. in Act. Hort. Petrop. iii. (1875) 50 = V. alpestris
amoenum, Trautv. l. c. 53 = V. amoena.
aristatum, Rafin. Précis Découv. 38 = V. gracilis, *Loisel.*
aurantiacum, Trautv. in Act. Hort. Petrop. iii. (1875) 56 = V. aurantia.
australe, Vell Fl. Flum. 316 ; vii. t. 141 (Lathyri sp.). —Bras.
Biebersteinii, Guss. Fl. Sic. Prod. ii. 445 = V. pubescens.
Boissieri, Hort. ex Boiss. Fl. Orient. ii. 598 = *orientale.*
calcaratum, Trautv. in Hort. Act. Petrop. iii. (1875) 45 = V. calcarata.
camelorum, Spreng. in Schrad. Journ. (1800) ii. 197 = Lens esculenta.
caprealatum, Lowe, Man. Fl. Mad. i. 199 = V. capreolata.
capricornium, Vell. Fl. Flum. 315 ; vii. t. 140 (Vicise sp.).—Bras.
cassubicum. Peterm. Fl. Lips. 553 = V. cassubica.
chilense, Fisch. & Mey. ex Kegel, Ind. Sem. Hort. Petrop. (1856) 40.—Chili.
cirrhosum, Vell. Fl. Flum. 316 ; vii. t. 142 (Lathyri sp.).—Bras.
cochinchinense, Pers. Syn. ii. 398.—Cochinchina.
corsicum, Nym. Syll. 210.—V. corsica.
costatum, Trautv. in Act. Hort. Petrop. iii. (1875) 52 = V. costata.
Cracca, Trautv. l. c. 49 = V. Cracca.
cyanesum, Boiss. & Hohen. Diagn. Ser. I. ix. 115.—Persia.
dalmaticum, Presl, in Weitenw. Beitr. ii. (1837) 21.—Dalmat.
diphyllum, Bess. Cat. Jard. Volh. Suppl. (1812) 15, ex Reich. Cat. Hort. Taur. (1815) App. i. 11.—Volhynia.
dispermum, Roxb. ex Willd. Enum. Hort. Berol. 766 = Lens esculenta.
emarginatum, Engelm. ex Alef. in Oestr. Bot. Zeitschr. ix. (1859) 368 = V. micrantha.
erectum, Walt. Fl. Carol. 187 = Galactia sessiliflora.
Ervilia, Linn. Sp. Pl. 738 = V. Ervilia.
filiforme, Kozb. in Wall. Cat. n. 5955 = V. hirsuta.
glabrum, Gilib. Fl. Lituan. ii. 107 = V. gemella.
gracile, DC. Cat. Hort. Monsp. 109 = V. gracilis, *Loisel.*
gracile, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 156 = V. pubescens.
himalayense, Ax. Bc. & Bouché, in Ind. Sem. Hort. Berol. (1855) App. 3.—Ind. or.
hirsutum, Linn. Sp. Pl. 738 = V. hirsuta.
hirsutum, Lour. Fl. Cochinch. 481 = *cochinchinense.*
Hohenackeri, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 35 = Lens Lenticula.
Kotschyanum, Boiss. Diagn. Ser. I. vi. 48 = V. Monbreti.
kurdistanum, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 364.—Kurdistan.
Lens, Linn. Sp. Pl. 738 = Lens esculenta.
Lens, Wall. Cat. n. 6954 C = V. hirsuta.
Lenticula, Schreb. ex Sturm, Deutschl. Fl. Heft 32 = Lens Lenticula.
lenticuloseum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = praec.
lentoides, Tenore, Prod. Fl. Nap. 70 = Lens nigricans.

ERVUM :—
Loiseleurii, Bieb. Fl. Taur. Cauc. iii. 475 = Vicia pubescens.
Loiseleurii, C. A. Mey. Verz. Pfl. Cauc. 148 = V. Meyeri.
longifolium, Tenore, Prod. Fl. Nap. 42 = V. gracilis, *Loisel.*
lunatum, Boiss. Diagn. Ser. II. v. 93 = V. lunata.
macrocarpum, Presl, Fl. Sic. p. xxiii.—Sicilia.
mastichinum, Hort. ex Steud. Nom. ed. II. i. 591 = Phaseolus ferrugineus.
mediterraneum, Vell. Fl. Flum. 315; vii. t. 186 (Viciae sp.).—Bras.
megalotropis, Trautv. in Act. Hort. Petrop. iii. (1875) 51 = V. megalotropis.
Meyeri, Trautv. i. c. 46 = V. Meyeri.
microanthum, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 364 = V. Lenaeana.
monanthos, Schrank, Baier. Fl. ii. 264 = *monanthos*, Linn.
monanthos, Georgi, Beschr. Russ. Reich. III. iv. 1171 = V. gracilis.
monanthos, Linn. Sp. Pl. 738 = V. monanthos.
multiflorum, Pursh, Fl. Am. Sept. ii. 739 = Astragalus multiflorus.
nigricans, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 364 = V. nigricans.
nigricans, Bieb. Fl. Taur. Cauc. ii. 164 = Lens nigricans.
nigrum, Hort. ex Heynh. Nom. i. 312, nomen = prsec. !
ochroleucum, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 365 = Lathyrus ochroleucus.
orientale, Boiss. Diagn. Ser. I. ix. 115.—Persia.
Orobus, Kitt. Taschenb. ed. II. 1163 = V. Orobus.
parviflorum, Bertol. Amoen. Ital. 88 = V. disperrimum.
poucijugum, Trautv. in Act. Hort. Petrop. iii. (1875) 76.—Transcaucasus.
pendulum, Roth, Nov. Pl. Sp. 351.—Hab. !
pictum, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 365 = V. biennis.
pilosissimum, Schur, Enum. Pl. Transs. 172 = Lens esculenta.
pilosum, Engelm. ex Alef. in Oestr. Bot. Zeitschr. ix. (1859) 363 = V. ludovicianum.
pisiforme, Peterm. Fl. Lips. 559 = V. pisiformis.
plicatum, Moench, Meth. 147 = V. krvilia.
Pseudo-Orobus, Trautv. in Act. Hort. Petrop. iii. (1875) 53 = V. Pseudo-Orobus.
pubescens, DC. Cat. Hort. Monsp. 109 = V. pubescens.
punctatum, Hort. ex Steud. Nom. ed. II. i. 591 = V. krvilia.
Salzii, J. Gay, ex Salis-Marsch. in Flora, xvii. (1834) II. Beibl. 61 = V. pubescens.
sardoum, Moris, ex Spreng. Syst. iv. Cur. Post. 846 = V. hirsuta.
silvaticum, Peterm. Fl. Lips. 552 = V. sylvatica.
solonicum, Linn. Cent. Pl. ii. 28; Amoen. Acad. iv. 296 = V. lathyroides.
solonicum, Thuill. Fl. Par. ed. II. 371 = V. gemella.
solonicum, Wulf. ex Schrank, Fl. Rar. Hort. Monac. t. 48 = Lens Lenticula.
stipulaceum, Bast. in Desv. Journ. Bot. iii. (1814) 18 = V. monanthos.
subglobosum, Relchb. Sel. Sem. Hort. Dresd. (1839); ex Linnaea, xiv. (1840) Litt. 192.—Hab. !
sylvaticum, Anct. plur. vide *silvaticum*.
sylvaticum, Fisch. Cat. Hort. Gorenk. (1808) 116 = Lens nigricans.
Tenorii, Steud. Nom. ed. II. i. 591, sphalm. vide *Terronii*.
tenuifolium, Lag. Gen. et Sp. Nov. 23 = V. gracilis, *Loisel.*
tenuifolium, Trautv. in Act. Hort. Petrop. iii. (1875) 50 = V. tenuifolia.
tenuissimum, Bieb. Casp. 185 = V. gemella.
tenuissimum, Pers. Syn. ii. 309 = V. gracilis, *Loisel.*
Terronii, Tenore, Fl. Neap. Prod. App. v. 23 = V. hirsuta.
tetragonum, Hoffm. Cat. Kourak. 47 = V. gemella.
tetraphyllum, Vell. Fl. Flum. 315; vii. t. 189 (Lathyri sp.).—Bras.
tetrospermum, Bieb. Fl. Taur. Cauc. ii. 164 = V. pubescens.
tetrospermum, Linn. Sp. Pl. 738 = V. gemella.
trideatatum, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 363 = V. ludovicianum.
trijugum, Alef. in Boopiandia, ix. (1861) 212 = V. trijuga.

ERVUM :—
uniflorum, Tenore, Fl. Nap. v. 121. t. 179. f. 2 = Lens Lenticula.
unijugum, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 365 = Vicia unijuga.
varium, Brot. Fl. Lusit. ii. 152 = V. gracilis, *Loisel.*
venosum, Trautv. in Act. Hort. Petrop. iii. (1875) 55 = V. venosa.
venulosum, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 364 = V. venulosa.
vicioeforme, Spreng. Syst. iii. 270 = seq.
vicioides, Desf. Fl. Atlant. ii. 168. t. 198 = V. cerviformis.
villosum, Pomel, Nouv. Mat. Fl. Atl. 194.—Alger.
villosum, Trautv. in Act. Hort. Petrop. iii. (1875) 47 = V. villosa.
volubile, Wall. Fl. Carol. 187 = Galactia glabella.

ERXLEBENIA, Opiz, Seznam, 41 (1852) = **Pyrola**, Tourn. (Ericac.).
rosea, Opiz, l. c. = P. minor.

ERXLEBIA, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 494 = Commelina, Linn.
fusiformis, Medic. l. c. = C. tuberosa.

ERYBATHOS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 326 = EROBATHOS, Spach = **Nigella**, Moench (Ranunc.).

ERYCIBE, Roxb. Pl. Corom. ii. 21. t. 159 (1798).
CONVOLVULACEAE, Benth. & Hook. f. ii. 868.
CATONIA, Vahl, in Skrivt. Nat. Hist. Selsk. Kiobenh. vi. (1810) 98.
ERIMATALIA, Roem. & Schult. Syst. v. 27 (1819).
cambodiensis, Miq. Fl. Ind. Bat. Suppl. 576 = Prineol.
citrinifiora, Griff. Notul. iv. 284 = glomerata.
coriacea, *Wall. Cat.* n. 1837.—Reg. Himal.
expansa, *Wall. l. c.* n. 1831.—Burma.
ferruginea, Griff. Notul. iv. 283 = expansa.
fragrans, Wall. Cat. n. 1836 = coriacea.
glaucescens, *Wall. l. c.* n. 1834.—Ind. or.; China.
glomerata, *Blume, Bijdr.* 1047.—Java.
glomerata, Wall. Cat. n. 1838 (= prsec. !).—Burma.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 193.—Burma.
laevigata, *Wall. Cat.* n. 1833.—Reg. Himal.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 193.—Malacca.
malaccensis, *C. B. Clarke, l. c.*—Burma; Malacc.
obtusifolia, *Benth. Fl. Hongk.* 236.—China.
paniculata, *Roxb. Pl. Corom.* ii. 21. t. 159.—As. et Austral. trop.
Princei, *Wall. Cat.* n. 1835.—Malaya.
Rheedii, Blume, Bijdr. 1047 = paniculata.
subpicata, *Wall. Cat.* n. 1832.—Reg. Himal. ; Burma.
tomentosa, *Blume, Bijdr.* 1048.—Java.
Wightiana, J. Grah. Cat. Pl. Bomb. 137 = paniculata.

ERYCINA, Lindl. Fol. Orch. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 565.
echinata, *Lindl. l. c.*—Mexic.

ERYNGIUM, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 878.
KLONION, Rafin. New Fl. Am. iv. 25 (1836).
LESSONIA, Bert. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 352, in syn.; et in Deless. Ic. Sel. iii. 48. t. 79 (1837).
STEELANTHUS, Rafin. Atl. Journ. 149 (1832).
agavifolium, *Griseb. in Gottl. Abh.* xix. (1874) 155.—Argent.
albovillosum, *Forsh. Fl. Aegypt. Arab.* p. xiii, nomen. —Ins. Melita.
alsinifolium, *Mart. ex Urban, in Mart. Fl. Bras.* xi. i. 309.—Bras.
alpinum, Lapeyr. Hist. Abr. Pl. Pyr. 137 = Bourgati.
alpinum, *Linn. Sp. Pl.* 233.—Europ.
americanum, Walt. Fl. Carol. 102 = prostratum !
amethystinum, Linn. Fl. Fr. iii. 401 = Bourgati.
amethystinum, *Linn. Sp. Pl.* 233.—Europ.
amethystinum, Salem. ex DC. Prod. i. 83 = dilatatum.
Andesonii, *Lag. ex Loud. Hort. Brit.* 101, nomen.— Hab. !
angustifolium, DC. Prod. iv. 95 = prostratum.
anomalum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 350.—Chili.
antihystericum, Rottl. in Act. Lit. Univ. Hafn. i. (1776) 238 = foetidum.
aquaticum, Hook. & Arn. Bot. Beech. Voy. 142 = petiolatum.

ERYNGIUM :—
aquaticum, Linn. Sp. Pl. 233 = yuccaefolium.
aquaticum, Michx. Fl. Bor. Am. i. 168 = prsealtum.
aquaticum, Vell. Fl. Flum. 126; iii. t. 99 = floribundum.
aquifolium, *Cav. in Anal. Cienc. Nat.* iii. (1801) 32.—Hispan.
arenalicum, *Baldw. in Ell. Sketch*, i. 344.—Florida.
articulatum, Hook. Lond. Journ. Bot. vi. (1847) 232 = petiolatum.
arvense, *Phil. in Linnaea*, xxx. (1859–60) 190.—Chili.
arvense, Vell. Fl. Flum. 125; iii. t. 88 = elegans.
asperifolium, Delar. Eryng. 36. t. 11 = giganteum.
australe, Wulf. & West, ex Schult. Syst. vi. 884 = amethystinum.
axilliflorum, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i. 172.—Mexic.
Baldwii, *Spreng. Syst.* i. 870.—Am. bor.
Barrelieri, *Boiss. in Ann. Sc. Nat. Sér.* III. i. (1844) 125.—Reg. Mediterr. or.
Beecheyanum, *Hook. & Arn. Bot. Beech. Voy.* 294.—Mexic.
bellidifolium, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 28 = nudicaule.
Billardieri, *Delar. Eryng.* 25. t. 2.—As. occ.
Billardieri, Heldr. ex Boiss. Diagn. Ser. I. x. 20 = Heldreichii.
bithynicum, *Boiss. in Ann. Sc. Nat. Sér.* III. i. (1844) 128.—As. Min.
Bocconi, Lam. Encyc. iv. 754 = tricuspidatum.
Bonplandi, *Delar. Eryng.* 52. t. 22.—Mexic.
Bonplandianum, H. B. & K. Nov. Gen. et Sp. v. 81 = prsec.
Bourgati, *Gouan, Illustr.* 7. t. 3.—Hispan. ; Marocc.
Bourgati, Vahl, Symb. Bot. ii. 46 = dilatatum.
Bovei, *Boiss. in Ann. Sc. Nat. Sér.* III. i. (1844) 124 —Alger.
bracteosum, *Griseb. in Gottl. Abh.* xxiv. (1879) 146.— Reg. Argent.
bromeliaefolium, *Delar. Eryng.* 60. t. 28.—Mexic.
Bungei, *Boiss. Fl. Orient.* ii. 824.—Persv.
buplenrifolium, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 353.—Ins. Juan Fernandez.
caeruleocreas, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 679 = Billardieri.
caeruleum, *Bieb. Fl. Taur. Cauc.* i. 200.—As. occ.
caeruleum, Gilib. Fl. Lituan. ii. 3 = planum.
caeruleum, Link, Enum. Hort. Berol. i. 261 = amethystinum.
campestre, *Linn. Sp. Pl.* 233.—Europ.
canaliculatum, *Cham. & Schlecht. in Linnaea*, i. (1826) 238 (err. typ. 338).—Brasa.
Cardesii, *Clus, in C. Gay, Fl. Chil.* iii. 118.—Chili.
Carlinae, *Delar. Eryng.* 58. t. 28.—Mexic.
carlinoides, *Boiss. Fl. Orient.* ii. 825.—Afghan.
carthamoides, *Sw. in Rees, Cycl.* xiii.—Mauritan.
cancasicum, *Fisch. ex Steud. Nom.* ed. I. 315, nomen. —Caucas.
caucasicum, Trautv. in Act. Hort. Petrop. i. (1871) 23 = caeruleum.
Cervantesii, Chapm. Fl. S U. St. 140 = Baldwii.
Cervantesii, *Delar. Eryng.* 47. t. 18. f. 1.—Mexic.
Chamissonis, *Urban, in Mart. Fl. Bras.* xi. i. 315 —Bras.
ciliatum, *Cham. & Schlecht. in Linnaea*, i. (1826) 349 (err. typ. 349).—Bras.
ciliatum, Vell. Fl. Flum. 126; iii. t. 100 = canaliculatum.
concavum, *Delar. Eryng.* 30. t. 7.—Mexic.
coquimbanum, *Phil. ex Urban, in Mart. Fl. Bras.* xi. i. 306, in obs.—Chili.
corniculatum, *Lagasc.* ex Urv. 756.—Hispan.
coronatum, Donn, Hort. Cantab. ed. IV. 56 = prsec.
coronatum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 350.—Reg. Argent.
corymbosum, Torr. & Gray, Fl. N. Am. i. 604 = Hookeri.
coscopolifolium, *Decne. in Bull. Soc. Bot. Fr.* xx. (1873) 27.—Reg. Argent.
crantzioides, *Griseb. in Gottl. Abh.* vi. (1854) 232. —Chili.
creticum, Jan, ex Guss. Fl. Sic. Prod. i. Add. 7 = triquetrum.
creticum, *Linn. Encyc.* iv. 754.—Reg. Mediterr. or.
crinitum, Webb, Iter Hisp. 46 = glaciale.
crinitum, Lange, in Kjoch. Vidensk. Meddel. (1865) 82 = aquifolium.
crinitum, Presl, Delic. Prag. 186 = dilatatum.

ERYNGIUM:—
cyanreum, Sibth. & Sm. Fl. Graec. Prod. i. 175 = orcticum.
cylindraceum, Willd. ex Spreng. Syst. i. 872 = Phyteuma.
cymosum, *Delar. Eryng.* 62. t. 21.—Mexic.
Decaisneanum, *Urban*, in *Mart. Fl. Bras.* xi. t. 816.—Bras.
Deppeanum, *Cham. & Schlecht. in Linnaea*, v. (1830) 207.—Mexic.
depressum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 351.—Chili.
dichotomum, *Desf. Fl. Atlant.* i. 226. t. 55.—Reg. Mediterr.
diffusum, *Torr. in Ann. Lyc. N. York*, ii. (1828) 206.—Am. bor.
dilatatum, Gras. Fl. Sic. Syn. i. 302 = amethystinum.
dilatatum, *Lam. Encyc.* iv. 755.—Reg. Mediterr. occ.
discolor, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 198.—Arizona.
divaricatum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 350.—Reg. Argent.
dubium, *Tratt. Archiv*, t. 186.—Hab.?
Duriaeanum, *Y. Gay*, in *Ann. Sc. Nat. Sér.* III. ix. (1848) 155.—Hispan.
Duriaei, J. Gay, l. c. 156 = praec.
ebracteatum, *Lam. Encyc.* iv. 759.—Bras.
elatum, *Decne. in Bull. Soc. Bot. Fr.* xx. (1873) 23.—Bras.
echinatum, *Urban, in Mart. Fl. Bras.* xi. i. 806.—Bras.
elegans, *Cham. & Schlecht. in Linnaea*, i. (1826) 348 (err. typ. 248).—Bras.
eriophorum, *Cham. & Schlecht. l. c.* 342 (err. typ. 242).—Bras.
expansum, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 236.—Austral.
falcatum, *Delar. Eryng.* 40. t. 18.—Syria.
filiforme, Shuttl. ex A. Gray, Pl. Wright. i. 78 = Baldwini.
flaccidum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 350.—Reg. Argent.
floribundum, *Cham. & Schlecht. in Linnaea*, i. (1826) 346 (err. typ. 246).—Bras.
fluminense, *Urban, in Mart. Fl. Bras.* xi. i. 826.—Bras.
foetidum, Forsk. Fl. Aegypt. Arab. p. xxiii = dilatatum.
foetidum, *Linn. Sp. Pl.* 382, partim.—Ind. occ.; Florida.
fuscatum, Linn. Sp. Pl. 382, partim = yuccaefolium.
foetidum, Walt. Fl. Carol. 113 = virginianum!
foliosum, *Scheele, in Linnaea*, xvii. (1843) 389.—Bras.
fraxineum, Moris, in Mem. Acc. Torin. xxxviii. (1835) 45 = bupleuroides.
galioides, *Lam. Encyc.* iv. 757.—Lusitan.
Ghiesbreghtii, *Decne. in Bull. Soc. Bot. Fr.* xx. (1873) 21.—Mexic.
giganteum, *Bieb. Fl. Taur. Cauc.* i. 201.—Armen. glaciale, *Boiss. Elench.* 44.—Hispan.
glaucum, Hoffm. Hort. Monç. (1808) n. 1309 = giganteum.
Glaziovianum, *Urban, in Mart. Fl. Bras.* xi. t. 825.—Bras.
glomeratum, Auch. ex Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 338 = Cardopatium orientale.
glomeratum, *Lam. Encyc.* iv. 755.—Ax. Min.; Syria.
gracile, Baldw. in Ell. Sketch. i. 345 = Baldwini.
gracile, *Delar. Eryng.* 54. t. 24.—Mexic.
gramineum, *Delar. l. c.* 60. t. 27.—Mexic.
gramineum, *Hornem. Enum. Pl. Hort. Hafn.* (1807) 16.—Hab.?
Gynnocephalum, Steud. Nom. ed. II. i. 593 = canaliculatum.
Hamiltoni, *Presl, ex DC. Prod.* iv. 84.—Mexic.
Harkesnelli, *Curran, in Bull. Calif. Acad.* i. n. III. (1885) 189.—Calif.
Haldreichii, *Boiss. Diagn.* Ser. I. x. 20.—Syria.
hemisphaericum, *Urban, in Mart. Fl. Bras.* xi. i. 824.—Bras.
heterophyllum, *Engelm. in Wisliz. Tour North Mexico*, 106.—Mexic.
Hookeri, *Walp. Rep.* ii. 369.—Am. bor. occ.
hurumbeides, *DC. Prod.* iv. 84.—Mexic.
Humboldtianum, H. B. & K. Nov. Gen. et Sp. v. 27 = paniculatum.
Humboldtii, Delar. Eryng. 61. t. 29 = paniculatum.
humifusum, *Clos, in C. Gay, Fl. Chil.* iii. 112.—Chili.
humile, *Cav. in Anal. Hist. Nat.* ii. (1800) 115.—Peruv.
ilicifolium, Brot. Fl. Lusit. i. 419 = Duriaeanum.
Ilicifolium, *Lam. Encyc.* iv. 757.—Reg. Mediterr. occ.
integrifolium, Walt. Fl. Carol. 112 = prostratum!
intermedium, Weinm. in Bull. Soc. Nat. Mosc. (1837) n. vii. 89 = planum.
junceum, *Cham. & Schlecht. in Linnaea*, i. (1826) 341 (err. typ. 241).—Bras.
juncifolium, Mart. ex Urban, in Mart. Fl. Bras. xi. i. 833 = juncosum.
Koehneanum, *Urban, l. c.* 823.—Bras.
Kotschyi, Boiss. Diagn. Ser. II. v. 97 = Billardieri.
lacuum, *Pohl, ex Urban, in Mart. Fl. Bras.* xi. i. 816.—Bras.
laeve, *Brousс. ex Schult. Syst.* vi. 887, nomen.—Hab.?
Lassauxii, *Decne. in Bull. Soc. Bot. Fr.* xx. (1873) 22.—Reg. Argent.
lateriflorum, Lam. Encyc. iv. 756 = Agriophyllum laterale.
latifolium, Gilib. Fl. Lituan. ii. 8 = planum.
latifolium, Hoffmgg. & Link, ex Willk. & Lange, Prod. Fl. Hisp. iii. 13 = campestre!
Leavenworthii, *Torr. & Gray, Fl. N. Am.* i. 604.—Mexic.
leucacanthum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = E. Spinalbe.
linearе, Pohl, ex Urban, in Mart. Fl. Bras. xi. i. 835 = juncosum.
linearifolium, Vell. Fl. Flum. 126; iii. t. 102 = canaliculatum.
Lingua-Tucani, Mart. Reise, i. 291 = E. Pristis.
longifolium, *Cav. in Anal. Hist. Nat.* ii. (1800) 183.—Mexic.
longiramosum, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i. 171.—Mexic.
lundelioides, *Cham. & Schlecht. in Linnaea*, i. (1826) 340 (err. typ. 240).—Bras.
macrocalyx, *Schrenk, Enum. Pl. Nov.* 60.—Soongaria.
marginatum, *Pohl, ex Urban, in Mart. Fl. Bras.* xi. i. 831.—Bras.
maritimum, *Linn. Sp. Pl.* 233.—Europ.; Ax. Min.
maritimum, *Pomel, Nouv. Mat. Fl. Atl.* 138.—Alger.
Mettenii, *Wood, Cl. Book*, 279.—Am. bor.
microcephalum, *Weber, ex Steud. Nom.* ed. II. i. 592, nomen.—Dalmatia.
microcephalum, *Willd. ex Spreng. Syst.* i. 871.—Am. trop.
monocephalum, *Cav. in Anal. Hist. Nat.* ii. (1800) 116.—Mexic.
multifidum, Sibth. & Sm. Fl. Graec. Prod. i. 175 = auriculatum.
nasturtiifolium, *Juss. ex Delar. Eryng.* 46. t. 17.—Am. bor.
nigrantothemum, Boiss. & Bahse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 95 = Billardieri.
Noëanum, *Boiss. Diagn.* Ser. II. ii. 72.—Persia.
nudicaule, *Law. Encyc.* iv. 759.—Reg. Argent.
nudiflorum, Willd. ex Spreng. Syst. i. 869 = abractеatum.
nudum, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 334.—Chili.
odoratum, Hort. Par. ex DC. Prod. iv. 92 = corniculatum.
odoratum, Lam. Encyc. iv. 756 = pusillum.
oligodon, *Griseb. in Goett. Abh.* xxiv. (1879) 146.—Reg. Argent.
Oliverianum, *Delar. Eryng.* 37. t. 12 (*Pl. hybrid.* !).—Orient.
orientale, Mill. Gard. Dict. ed. VIII. n. 6 = amethystinum!
ovalifolium, Michx. Fl. Bor. Am. i. 168 = virgatum.
ovinum, A. Cunn. in Field, N. S. Wales, 338 = rostratum.
palliscens, Mill. Gard. Dict. ed. VIII. n. 5 = amethystinum.
paniculatum, *Paul. & Vis. in Mem. Ist. Veget.* xv. (1870) 20.—Serbia.
Palmito, *Boiss. & Heldr. Diagn.* Ser. I. x. 21.—Graecia.
palustre, Hort. Madr. ex DC. Prod. iv. 92 = corniculatum.
pandanifolium, *Cham. & Schlecht. in Linnaea*, i. (1826) 336 (err. typ. 236).—Bras.
paniculatum, *Cav. & Dombey, ex Delar. Eryng.* 59. t. 26.—Chili.
parviflorum, Sibth. & Sm. Fl. Graec. Prod. i. 175 = glomeratum.
pauciflorum, Hoffmgg. & Link, Fl. Port. ii. 379 = galioides.
pectinatum, *Presl. ex DC. Prod.* iv. 86.—Mexic.
pentaphyllum, *Link, ex Steud. Nom.* ed. I. 316, nomen.—Hab.?
pentechinum, Sieber, ex DC. Prod. iv. 89 = glomeratum.
petiolatum, *Hook. Fl. Bor. Am.* i. 259.—Am. bor. occ.
Phyteumae, *Delar. Eryng.* 51. t. 21.—Mexic.
phyteumatos, Benth. Fl. Hartw. 86 = praec.
pinnatifidum, Bunge, in Lehm. Pl. Preiss. i. 293 = rostratum.
planifolium, Pall. Reise, iii. 316 = planum.
planiginosum, *F. Muell. in Proc. Roy. Soc. Tasm.* iii. (1859) 235.—Austral.
planum, Lapeyr. Hist. Abr. Fl. Pyr. 187 = Bourgati.
planum, Lindl. in Royle, Illustr. Bot. Himal. 223 = caeruleum.
planum, *Linn. Sp. Pl.* 233.—Europ.; Ax. bor.
platyphyllum, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 24.—E. Sem.
Plukenetii, Ell. Sketch. i. 332 = virginianum.
Pohlianum, *Urban, in Mart. Fl. Bras.* xi. i. 836.—Bras.
polyphlorum, *Clos, in C. Gay, Fl. Chil.* iii. 113.—Chili.
poteriolides, *Griseb. in Goett. Abh.* xix. (1874) 155.—Reg. Argent.
praealtum, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 210.—Am. bor. occ.
pratense, *Phil. in Linnaea*, xxx. (1859–60) 189.—Chili.
Prietia, *Cham. & Schlecht. l. c.* 1. (1826) 337 (err. typ. 237).—Bras.
prostratum, James, ex S. Wats. Bibliog. Ind. N. Am. Bot. 421 = diffusum.
prostratum, *Nutt. ex DC. Prod.* iv. 92.—Am. bor.
prostratum, Wood, Cl. Book, 278 = Baldwini.
proteaeflorum, *Delar. Eryng.* 62. t. 30.—Mexic.
Pseudojunceum, *Clos, in C. Gay, Fl. Chil.* iii. 114.—Chili.
pulchellum, *Phil. Fl. Atac.* 25.—Chili.
pulchellum, Wedd. Chlor. And. ii. 201 = humile.
pumilum, Gilib. Chor. Grodn. 30 = planum.
purpuratum, Sm. in Rees, Cycl. xiii = dilatatum.
pusillum, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 125 = viviparum.
pusillum, Delar. Eryng. 44. t. 16 = galioides.
pusillum, Desf. Fl. Atlant. i. 223 = Barrelieri.
pusillum, Gilib. Fl. Lituan. ii. 8 = planum.
pusillum, *Linn. Fl. Suec.* ed. II. 238.—Europ. austr.; Afr. bor. pyramidale, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 890.—Persia.
radiatum, Willd. ex Spreng. Syst. i. 871 = Carlinae.
radiciflorum, *Tratt. Archiv.* t. 764; *ex DC. Prod.* iv. 93.—N. Granat.
ranunculoides, *Benth. Pl. Hartw.* 86.—Mexic.
ranunculoides, Willd. ex Spreng. Syst. i. 871 = humile.
Ravenelii, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 209.—Mexic.
reptens, Willd. ex Spreng. Syst. i. 870 = Baldwini.
rigidum, *Law. Encyc.* iv. 752.—Europ.
rostratum, *Cav. in Anal. Hist. Nat.* ii. (1800) 118.—Austral.; Chili.
Sanguisorba, *Cham. & Schlecht. in Linnaea*, i. (1826) 339 (err. typ. 239).—Bras.
sarcophyllum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 352.—Ins. Juan Fernand.
saxposum, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i. 172.—Mexic.
scariosum, Delar. Eryng. 38. t. 5 = glomeratum.
Schiedeanum, *Cham. & Schlecht. in Linnaea*, v. (1830) 206.—Mexic.
scirpinum, *Cham. l. c.* viii. (1833) 324.—Bras.
scrbicum, *Panč. in Verh. Zool.-Bot. Ges. Wien*, vi. (1856) 590.—Serbia.
Serra, *Cham. & Schlecht. in Linnaea*, i. (1826) 346 (err. typ. 246).—Bras.
serratum, *Cav. in Anal. Hist. Nat.* ii. (1800) 189.—Mexic.
serratum, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 208 = E. Serra.
sparganoides, *Clos, in C. Gay, Fl. Chil.* iii. 113.—Chili.

ERYNGIUM:—
Spina-alba, Auct. = sec.
Spinalba, *Vill. Prosp.* 26; *Hist. Pl. Dauph.* ii. 660.—Europ.
stellatum, *Mutis, ex Delar. Eryng.* 55. t. 25.—Am. austr.
stenophyllum, *Urban, in Mart. Fl. Bras.* xi. t. 880.—Bras.
suaveolens, Broess. ex Desf. Tabl. ed. I. 122 = corniculatum.
subscaule, *Cav. in Anal. Hist. Nat.* ii. (1800) 116.—Mexic.
subulatum, Vell. Fl. Flum. 126; ill. t. 101 = paniculatum.
syriacum, Lam. Encyc. iv. 750 = creticum.
tenue, Hook. & Arn. Bot. Beech. Voy. 393 = Wrightii.
tenue, *Lam. Encyc.* iv. 755.—Reg. Mediterr. occ.
ternatum, *Poir. Encyc. Suppl.* iv. 235.—Ins. Creta.
tetracephalum, Bunge, in Lehm. Pl. Preiss. i. 293 = rostratum.
thoracifolium, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 122.—As. Min.
thyrsoideum, *Boiss. l. c.* 121.—Persia; Armen.
Tournefortianum, Sternb. in Syll. Ratisb. i. (1824) 32 = ternatum.
tricuspidatum, Domb. ex DC. Prod. iv. 62 = Asteriscum chilense.
tricuspidatum, *Linn. Amoen. Acad.* iii. 405.—Reg. Mediterr.
tricuspidatum, Pand. in Verh. Zool.-Bot. Ges. Wien, vi. (1856) 220 = palmatum.
trifidum, Linn. Diss. Demons. in Amoen. Acad. iii. 406 = campestre.
tripartitum, *Desf. Cat. Hort. Par.* ed. III. 405.—Hab.? triquetrum, *Vahl, Symb. Bot.* ii. 46.—Afr. bor.; Sicil.
tuberosum, *Rafin. Fl. Ludov.* 60.—Am. bor.
Tuscanu, Vell. Fl. Flum. 126; ill. t. 103 = E. Pristis.
uncinatum, Cham. & Schlecht. in Linnaea, i. (1826) 347 (err. typ. 247) = elegans.
uniflorum, *Clos, in C. Gay, Fl. Chil.* iii. 115.—Chili.
variflolium, *Cost. in Bull. Soc. Bot. Fr.* xxii. (1875) 59.—Marocc.
vesiculosum, *Labill. Nov. Holl. Pl.* i. 78. t. 98.—Austral.; N. Zel.
virens, Link, in Linnaea, ix. (1834) 570 = campestre.
virgatum, *Lam. Encyc.* iv. 757.—Am. bor.
virginianum, Ell. Sketch. i. 340 = praealtum.
virginianum, *Lam. Encyc.* iv. 759.—Am. bor.
viviparum, *Y. Gay, in Ann. Sc. Nat.* Sér. III. ix. (1848) 171.—Gall. occ.
vulgare, Lam. Fl. Fr. iii. 401 = campestre.
Wrightii, *A. Gray, Pl. Wright.* i. 76.—Texas.
yuccaefolium, *Michx. Fl. Bor. Am.* i. 104.—Am. bor.
Zanonii, Lam. Encyc. iv. 754 = tricuspidatum.

ERVOCYCLA, Pritz. in. Ind. 421 (1855), err. typ. = ERIOCYCLA, Lindl. = Pituranthos, Viv. (Umbell.).

ERYSIBE, G. Don, Gen. Syst. iv. 274, 292 (1837) aphalm. = Erycibe, Roxb. (Convolv.).

ERYSIMASTRUM, Rupr. Fl. Cauc. 76 (1869) = Erysimum, Tourn. (Crucif.).
ancepe, Rupr. Fl. Cauc. 86 = Erysimum ancepe.
andinum, Rupr. l. c. 83, in textu = Erysimum leptophyllum.
asperum, Rupr. l. c. 76 = Erysimum asperum.
borcale, Trautv. in Act. Hort. Petrop. viii. (1883) 105 = Erysimum boreale.
brachycarpum, Trautv. l. c. = Erysimum brachycarpum.
Chorinthium, Trautv. l. c. = Erysimum cheiranthoides.
cretaceum, Trautv. l. c. 106 = Erysimum cretaceum.
gelidum, Trautv. l. c. = Erysimum gelidum.
ibericum, Rupr. Fl. Cauc. 77 = Erysimum ibericum.
macrocarpum, Trautv. in Act. Hort. Petrop. viii. (1883) 106 = Erysimum ibericum.
Meyerianum, Rupr. Fl. Cauc. 82 = Erysimum Meyerianum.
substrigosum, Rupr. l. c. 80 = Erysimum leptophyllum.

ERYSIMUM, (Tourn.) Linn. Syst. ed. I (1735); Gen. ed. I. 198 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 79.
CHEIRINIA, Link, Enum. Hort. Berol. ii. 170 (1822).
CUSPIDARIA, Link, Handb. ii. 315 (1831).
ERYSIMASTRUM, Rupr. Fl. Cauc. 76 (1869).

ERYSIMUM:—
STROPHADES, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 82.
acaules, Jord. Diagn. i. 176 = ochroleucum.
aciphyllum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 80.—As. Min.
aducumbens, Kotschy, ex Boiss. Fl. Orient. i. 202 = alpestre.
adpressum, Mann, ex Opiz, Natural. xi. (1826) 422 = hieracifolium.
adscendens, Nym. Syll. Suppl. 84 = *ascendens*.
alliaceum, Salisb. Prod. 370 = Sisymbrium Allaria.
Allioria, Lian. Sp. Pl. 660 = Sisymbrium Allaria.
alpestre, Jord. Diagn. i. 171 = lanceolatum.
alpestre, *Kotschy, ex Boiss. Fl. Orient.* i. 202.—As. Min.
alpinum, Fries, Summ. Veg. Scand. 229 = hieracifolium.
alpinum, Pers. Syn. ii. 200 = lanceolatum.
alpinum, Roth, Tent. Fl. Germ. i. 382 = Arabis brassicaeformis.
altaicum, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 158.—Reg. Caucas.; Sibiria.
altissimum, Lej. Fl. Spa, ii. 70 = hieracifolium.
alyssoides, *Franch. in Nouv. Arch. Mus. Par.* Sér. II. v. (1884) 189.—As. or.
amplexicaule, *A. Rich. Tent. Fl. Abyss.* i. 20.—Abyss.
anceps, *Stev. ex Ledeb. Fl. Ross.* i. 187.—Reg. Caucas.
Andrzejowicianum, Bess. Enum. Pl. Volh. 97 = canescens.
angustatum, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 166 = hieracifolium.
angustifolium, Baumg. Enum. Stirp. Transs. ii. 262 = pannonicum.
angustifolium, *Ehrh. Beitr.* vii. 155.—Hungar.
angustifolium, *Rafin. Atl. Journ.* 153.—Am. bor.
arcuatum, Opiz, ex J. & C. Presl, Fl. Cech. 138 = Barbarea arcuata.
arboreum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 95 = asperum.
armeniacum, Boiss. Fl. Orient. i. 202 = gelidum.
armeniacum, J. Gay, Erysim. Nov. 4 = ibericum.
arvense, Thore, Chlor. Land. 295 = Brassica Cheiranthos.
ascrifens, Jord. Diagn. i. 180 = ochroleucum.
asperulum, *Boiss. & Heldr. Diagn.* Ser. II. vi. 11.—Graecia.
asperum, *DC. Syst.* ii. 505.—Am. bor.
atticum, Heldr. & Sart. in Boiss. Diagn. Ser. II. i. 25 = Boryanum.
Auctori, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 78.—Persia.
Auctorianum, J. Gay, Erysim. Nov. 9 = Gayanum.
auranticum, *Leyb. in Flora*, xxxviii. (1855) 588.—Tyrolis.
aureum, *Bieb. Fl. Taur. Cauc.* ii. 117.—Reg. Caucas.
aurigeranum, Jeanb. & Timb. in Bull. Soc. Sc. Phys. & Nat. Toulouse, iii. (1875-76) 555 = cheiranthoides.
aurosicum, Jord. Diagn. i. 169 = montosicolum.
australe, *Y. Gay, Erysim. Nov.* 6.—Hispan.; Marocc.
australe, Heldr. ex Boiss. Fl. Orient. i. 198 = Raulini.
austriacum, Roth, Tent. Fl. Germ. i. 382 = Conringia austriaca.
autasticum, *Veri. Cat. Yard. Grenob.* (1857) 94.—Gallia.
Barbarea, Bieb. Fl. Taur. Cauc. ii. 116 = Barbarea arcuata.
Barbarea, Forsk. Fl. Aegypt. Arab. p. lxix = Nasturtium palustre.
Barbarea, Linn. Sp. Pl. 660 = Barbarea vulgaris.
Barbarea, Pall. Tabl. Taur. 54 = Barbarea arcuata.
Baumgartenianum, Schur, in Verh. Siebenb. Ver. Naturw. v. (1858) 166 = crepidifolium.
biculosum, *DC. Syst.* ii. 509.—Helvetia.
bicorne, [Soland. in] Ait. Hort. Kew. ed. I. ii 394 = Notoceras canariense.
bieraciifolium, F. Muell. in Linnaea, xxv. (1852) 367 = Blennodia lasiocarpa.
Boccani, Pers. Syn. ii. 200 = canescens, ochroleucum.
Boissieri, Walp. Rep. i. 164 = Sisymbrium parviflorum.
Bonannianum, Presl, Fl. Sic. i. 76 = lanceolatum.
borcale, *C. A. Mey. ex Rupr. Fl. Cauc.* 76.—Sibir.
Boryanum, *Boiss. Diagn.* Ser. I. i. 71.—Graecia.
Bourgeauianum, Coss. ex Nym. Consp. 41 = australe.
brachycarpon, *Spreng. Syst.* ii. 908.—Reg. Caucas.
brachycarpum, *Boiss. Fl. Orient.* i. 198.—Armenia; Persia.
Brassica, Crantz, Inst. ii. 101 = Conringia orientalis.

ERYSIMUM:—
brayoides, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 57 = Braya alpina.
brevicaule, Jord. Diagn. i. 173 = lanceolatum.
brevipes, F. Muell. in Linnaea, xxv. (1852) 367 = Blennodia brevipes.
brevirostre, *Tausch, in Flora*, xix. (1836) 408.—Hab.?
caespitosum, *DC. Syst.* ii. 497.—Persia.
calycinum, *Griseb. Spicil. Fl. Rumel.* i. 260.—Macedon.
campestre, Scop. Fl. Carn. ed. II. ii. 37 = Conringia orientalis.
canescens, Bory & Chaub. Fl. Pélop. 46 = graecum.
canescens, DC. Syst. ii. 501 = australe.
canescens, Moench, Meth. Suppl. 86 = cinereum.
canescens, *Roth, Catalecta*, i. 76.—Europ. austr., Reg. Caucas.
capsellinum, *F. Muell. Nativo Pl. Vict.* i. 35.—Austral.
cardaminoides, F. Muell. Fragm. xi. 59 = Sisymbrium cardaminoides.
carium, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 77.—As. Min.
carnicolum, Dollin. in Flora, x. (1827) 254; Host, Fl. Austr. ii. 376 = pannonicum.
carnuicolum, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 143 = crepidifolium.
caucasicum, *Trautv. in Bull. Soc. Nat. Mosc.* xii. (1868) 1, 480.—Reg. Caucas.
cheiranthifolium, Gilib. Fl. Lituan. ii. 58 = cheiranthoides.
cheiranthoides, Crantz, Stirp. Austr. fasc. i. 29 = lanceolatum.
cheiranthoides, *Linn. Sp. Pl.* 661.—Europ.; Am. bor.
Cheiranthus, Baumg. Enum. Stirp. Transs. ii. 261 = pannonicum.
Cheiranthus, J. & C. Presl, Fl. Cech. 138 = crepidifolium.
Cheiranthus, Roth, Tent. Fl. Germ. i. 282 = lanceolatum.
Cheiri, Crantz, Class. Cruc. f. 116 = Cheiranthus Cheiri.
cheiriformum, Bolss. Diagn. II. v. 24 = E. Pseudocheiri.
cheiriflorum, Wallr. Sched. Crit. 367 = pannonicum.
chlidopodum, *Tausch, in Flora*, xix. (1836) 408.—Hab.?
chrysanthemum, Pand. ex Nym Consp. 41 = pannonicum.
cinerascens, Jord. Diagn. i. 168 = canescens.
cinereum, *Moench, Meth.* 265.—Europ.
clausum, Walp. Rep. i. 169 = Conringia clavata.
collinum, *Andrz. in DC. Prod.* i. 198.—Reg. Caucas.
collispermum, Jord. Diagn. i. 165 = canescens.
Competianum, *Cerm. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 304.—Soongar.
confertum, Jord. Diag. i. 151 = virgatum.
confine, Jord. l. c. 169 = canescens.
constrictum, Jord. l. c. 177 = ochroleucum.
cordifolium, Pall. Tabl. Taur. 54; Stokes, Bot. Mat. Med. iii. 458 = Sisymbrium Allaria.
cornutum, Pall. Reise, iii. t. Mm = Malcolmia torulosa
cornutum, Pers. Syn. ii. 200 = Syrenia restiliflora.
crassicaule, *Boiss. Fl. Orient.* i. 199.—Persia.
crassipes, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 27.—As. Min.; Pers.
crassistylum, *Presl, Fl. Sic.* i. 77.—Sicilia.
crepidifolium, Griseb. Spicil. Fl. Rumel. ii. 258 = amygdalum.
crepidifolium, *Reichb. fil. Fl. Germ.* i. 8.—Europ.
creticum, *Rupr. Fl. Cauc.* 76.—Reg. Caucas.
creticum, *Boiss. & Heldr. Diagn.* Ser. II. i. 36.—Ins. Creta.
Cunninghamii, F. Muell. Census, 5 = Blennodia Cunninghamii.
cuspidatum, Jord. Diagn. i. 164 = canescens.
curvipes, F. Muell. in Linnaea, xxv. (1852) 368 = Blennodia curvipes.
cuspidatum, *DC. Syst.* ii. 498.—Graec.; As. Min.; Persia.
Custeauanum, Schur, Enum. Pl. Transs. 57 = crepidifolium?
Decumbens, Schleich. ex Steud. Nom. ed. II. i. 546 = ochroleucum.
deflexum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 166.—Reg. Himal.
delphinense, Jord. Diagn. i. 185 = virgatum.
dentatifolium, Jord. l. c. 194 = virgatum.
dentatocalycinum, *J. & C. Presl, Fl. Cech.* 139.—Bohem.
diffusum, Ehrh. Beitr. vii. 157 = canescens.

ERYSIMUM:—
divaricatum, Czern. ex Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 304.—Szongaria.
divaricatum, Waldst. Beitr. Fl. Hercyn. 228.—Europ.
dubium, DC. Syst. ii. 504.—Hab.?
Durieui, *Boiss. Diagn.* Ser. II. i. 26.—Hispan.
durum, J. & C. Presl, Delic. Prag. 226 = virgatum.
elatum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 95 = asperum.
elatum, *Pomel, Nouv. Mat. Fl. Atl.* 369.—Europ.
elatum, Schum. Enum. Pl. Transs. 58 = Marschallianum.
elbrusense, *Boiss. Fl. Orient.* i. 302.—Persia.
erocastifolium, *Rupr. Fl. Cauc.* 86.—Reg. Caucas.
Erucastrum, Scop. Fl. Carn. ed. II. ii. 27 = Brassica Erucastrum.
etnense, Jord. Diagn. i. 161 = cheiranthoides.
exaltatum, Andrz. in Bess. Enum. Pl. Volh. 73, 83 = hieracifolium.
exasperatum, Gmel. ex DC. Syst. ii. 497 = caespitosum.
filifolium, F. Muell. in Linnaea, xxv. (1852) 368 = Blennodia filifolia.
firmum, Sweet, Hort. Brit. ed. I. 95 = virgatum.
Fischerianum, Presc. ex Ledeb. Fl. Ross. i. 188 = altaicum.
foliosum, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 166 = hieracifolium.
fruticulosum, J. Gay, Exysim. Nov. 5 = caespitosum.
funiculosum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 166.—Reg. Himal.
Fustianum, Schur, Enum. Pl. Transs. 58 = Marschallianum.
Gayanum, *Boiss. Fl. Orient.* i. 196.—Persia.
gelidum, *Bunge, ex Tchihat. Asc Min. Bot.* i. 386.—Armenia.
glaberrimum, Hook. & Arn. Bot. Beech. Voy. 322 = Sisymbrium linifolium?
glabrum, *Presl, Fl. Sic.* i. 76.—Sicilia.
glareosum, Jord. Diagn. i. 178 = ochroleucum.
glastifolium, Crantz, Class. Crucif. 117 = Arabis perfoliata.
glaucum, Moench, Meth. 255 = Coringia orientalis.
goniocaulon, *Boiss. Diagn.* Ser. I. viii. 28.—As. Min.; Syria.
gracile, DC. Syst. ii. 504 = versicolor.
gracile, J. Gay, Exysim. Nov. 9 = hieracifolium.
graecum, *Boiss. & Heldr. Diagn.* Ser. II. i. 27.—Graecia.
gramineum, *Pomel, Nouv. Mat. Fl. Atl.* 371.—Algeria.
grandiflorum, Bieb. Fl. Taur. Cauc. ii. 117 = leptostylum.
grandiflorum, Czern. ex J. Gay, Exysim. Nov. 3 = ucranicum.
grandiflorum, Desf. Fl. Atlant. ii. 85 = longifolium.
grandiflorum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 96 = Cheiranthus asper.
grandiflorum, Wallp. Ann. ii. 49 = Coringia grandiflora.
Griffithianum, *Boiss. Diagn.* Ser. II. i. 28.—Persia; Afghanist.
Griffithii, Hook. f. & T. Anders. in Fl. Brit. Ind. i. 158 = Griffithianum.
helveticum, Cat. Fl. Madrit. 128 = australe.
helveticum, DC. Fl. Fr. iv. 858 = ochroleucum.
helveticum, Griseb. Spicil. Fl. Rumel. i. 269 = olympicum?
Hesperis, Scop. Fl. Carn. ed. II. ii. 28 = lanceolatum.
hieracifolium, Jacq. Fl. Austr. i. 72 = pannonicum.
hieracifolium, *Linn. Cent. Pl.* i. 18.—Europ.
hieracifolium, Pall. Tabl. Taur. 54 = cuspidatum.
hieracifolium, Schleich. ex DC. Syst. ii. 496 = virgatum.
hieracifolium, Urv. in Mém. Soc. Linn. Par. i. (1822) 387 = smyrnaeum.
hirsutum, Lbsk. Enum. Hort. Berol. ii. 268 = Sisymbrium runcinatum.
Hookeri, *Boiss. Fl. Orient.* i. 208.—Afghanist.
humile, Pers. Syn. ii. 200 = lanceolatum.
hybridum, *Gremer, Dec.* 6.—Holsat.
Ibericum, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 43 = Smyrnaeum.
Ibericum, *DC. Syst.* ii. 498.—Reg. Caucas.
incanum, *Kunze, in Flora*, xxix. (1846) 752.—Hispan.
integrifolium, *Tenore, ex Steud. Nom.* ed. II. i. 394.—Hab.?

ERYSIMUM:—
jugicolum, Jord. Diagn. i. 178 = pumilum.
junceum, Willd. Sp. Pl. iii. 518 = Sisymbrium junceum.
Kotschyanum, *F. Gay, Exysim. Nov.* 5.—As. Min.
Kunzeanum, Boiss. & Reut. Diagn. Ser. II. i. 27 = incanum.
lactescens, Boiss. l. c. Ser. I. i. 71 = rupestre.
lancolatum, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 116.—Europ.
lanceolatum, Cham. & Schlecht. in Linnaea, i. (1826) 23 = Redowskii.
lanceolatum, Colin. Apunt. Fl. Cost. 28 = australe.
lanceolatum, Kitchw.,Casp. Cauc. 81 = hieraciifolium.
lanceolatum, Hook. Fl. Bor. Am. i. 64 = parviflorum.
lanceolatum, Pursh. Fl. Am. Sept. ii. 69 = asperum.
lanceolatum, Reichb. ex Nym. Consp. 41 = pannonicum.
lanceolatum, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 808 = ochroleucum.
laxiflorum, *F. Gay, Exysim. Nov.* 4.—Persia.
leiostachyum, *Rupr. Fl. Cauc.* 75.—Reg. Caucas.
leptocarpum, *F. Gay, Exysim. Nov.* 4.—Armenia.
leptopetalum, C. Koch, in Linnaea, xix. (1847) 52 = cuspidatum.
leptophyllum, *Andrz. in DC. Prod.* i. 198.—Reg. Caucas.
leptostachyum, Hohen. ex Boiss. Fl. Orient. i. 208 = leptophyllum.
leptostylum, *DC. Syst* ii. 494.—Reg. Caucas.
leucophaeum, Jord. Diagn. i. 167 = canescens.
linariaefolium, Tausch, in Flora, xiv. (1831) 212 = lanceolatum.
linearе, Boiss. Diagn. Ser. I. viii. 24 = oleaefolium.
linearifolium, Moench, Meth. Suppl. 85 = ochroleucum.
linifolium, *F. Gay, Exysim. Nov.* 3.—Hispan.
longifforum, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 166 = hieraciifolium.
longifolium, *DC. Syst.* ii. 504.—Afr. bor.
longisiliquum, *Tausch, in Flora*, xix. (1836) 408.—Hab.!
longisiliquosum, *Willd. Enum. Hort. Berol.* 680.—Hab. auct.
longisiliquum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 166.—Reg. Himal.
longisiliquum, Schleich. Cat. Pl. Helv. ed. III. 14 = virgatum.
Lucae, F. Muell. Fragm. xi. 59 = Sisymbrium Lucae.
lucidum, Salisb. Prod. 270 = Barbarea vulgaris.
lyratum, Gilib. Fl. Lituan. iii. 80 = Barbarea vulgaris.
lyrifolium, Stokes, Bot. Mat. Med. iii. 459 = Barbarea vulgaris.
maclaurum, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 80.—China.
maclovianum, C. Gay, ex Spreng. Syst. iv. Cur. Post. 243 = Arabis maclioviana.
macranthum, *F. Gay, Exysim. Nov.* 3.—Mexic.
macranthum, *Tausch, in Flora*, xix. (1836) 409.—Austria.
macrocarpum, Rupr. Fl. Cauc. 75, nomen = Ibericum.
macrostigma, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 61.—Persia.
Marschallianum, *Andrz. in Bieb. Fl. Taur. Cauc.* iii. 447.—Podol.; Sibir.
Meyerianum, *Rupr. Fl. Cauc.* 75, nomen.—Reg. Caucas.
microcarpum, Tausch, ex Willk. & Lange, Prod. Fl. Hisp. iii. 808 = ochroleucum.
minus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Hab.?
montanum, Crantz, Class. Crucif. 116 = hieraciifolium.
montosicolum, *Jord. in Billot, Ann. Fl. Fr. All.* 128.—Gallia.
murale, Cat. Fl. Madrit. 128 = australe.
murale, DC. ex Nym. Consp. 41 = lanceolatum.
murale, *Desf. Tabl.* ed. I. 129.—Europ.
murale, Lam. Fl. Fr. ii. 514 = Cheiranthus Cheiri.
mutabile, *Boiss. & Heldr. Diagn.* Ser. I. viii. 26.—Graecia.
myriophyllum, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 102.—Hispan.
nanum, *Boiss. & Hohen. Diagn.* Ser. I. viii. 23.—Persia.
Nasturtium, F. Muell. in Linnaea, xxv. (1852) 368 = Blennodia nasturtioides.
nemorale, *Pomel, Nouv. Mat. Fl. Atl.* 370.—Algeria.
nervosum, *Pomel, l. c.*—Algeria.
nevadense, Reut. Cat. Jard. Genèv. (1855) 4.—Cf Linnaea, xxviii. (1856) 855 = australe.

ERYSIMUM:—
obtusangulum, [Clairv.] Man. Herb. 219 = Sisymbrium obtusangulum.
ochroleucum, *DC. Fl. Fr.* iv. 858.—Hispan.
ochroleucum, Zawad. ex Niesal. in Verh. Naturf. Ver. Brunn, vii. i. (1870) 59 = crepidifolium.
odoratum, Baumg. Enum. Stirp. Transs. ii. 262 = crepidifolium.
odoratum, Ehrh. Beitr. vii. 157 = pannonicum.
officinale, Linn. Sp. Pl. 660 = Sisymbrium officinale.
oleaefolium, *F. Gay, Exysim. Nov.* 6.—Persia.
olympicum, *Boiss. Fl. Orient.* i. 191.—Thessalia.
orsites, Jord. Diagn. i. 178 = pumilum.
orientale, Mill. Gard. Dict. ed. VIII. n. 4; [R. Br. in] Ait. Hort. Kew. ed. II. iv. 117 = Coringia orientalis.
pachycarpum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 167.—Reg. Himal.
pallens, Boiss. Voy. Bot. Espagne, 716 = australe.
pallens, Pers. Syn. ii. 200; Koch, Syn. Fl. Germ. ed. I. 59 = ochroleucum.
pallens, Wallr. Sched. Crit. 883 = crepidifolium.
pallescens, *Herbich, Stirp. Rar. Bucov.* 49.—Europ.
pallidum, *Boiss. Fl. Orient.* i. 207.—Lycia.
pannonicum, *Crantz, Stirp. Austr.* ed. I. fasc. i. 30.—Europ.
Parnassi, Nym. Consp. 40 = Boryanum.
parviflorum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 95.—Am. bor.
parviflorum, Pers. Syn. ii. 199 = cheiranthoides.
parvulum, Jord. Diagn. i. 173 = pumilum.
pasgalense, *Boiss. Fl. Orient.* i. 196.—Persia.
patens, J. Gay, Exysim. Nov. 9 = hieraciifolium.
patsiliquosum, Wender. Del. Sem. Hort. Marb. (1838) 4.—Cf. Linnaea, xxvi. (1853) 751 = Coringia planisiliqua.
patsiliquosum, Wender. l. c. (1856) 3.—Cf. Linnaea, xxix. (1857–58) 797 = praec.
patulum, *Iljenkam. Illustr. Hayn. Suppl.* 73.—Hab.? pectinatum, *Bory & Chaub. Fl. Pélop.* 45. t. 26.—Graecia.
perfoliatum, Crantz, Stirp. Austr. ed. I. fasc. i. 27 = Coringia orientalis.
Perofskianum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 56.—Reg. Caucas.; Afghanist.
persepolitanum, *Boiss. Diagn.* Ser. I. vi.—Persia.
persicum, Boiss. in Ann. Sc. Nat. Ser. II. xvii. (1842) 70 = subulatum.
petrophilum, Jord. Diagn. i. 166 = canescens.
pinnatum, Walt. Fl. Carol. 174 = Sisymbrium canescens.
planisiliquum, Steud. Nom. ed. II. i. 394; Ledeb. Fl. Ross. i. 192 = Coringia planisiliqua.
platypkyllum, Schur, Enum. Pl. Transs. 56 = canescens.
polyceratium, Link, Enum. Hort. Berol. ii. 168 = Sisymbrium polyceratium.
polyceratum, Pall. Reise, iii. 741 = Malcolmia africana, torulosa.
praecox, Sm. Fl. Brit. iii. 707 = Barbarea praecox.
Pseudoodoratum, *Boiss. Fl. Orient.* i. 201.—Armenia.
Pseudo-odoratum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 84 = hieraciifolium!
pulchellum, J. Gay, Exysim. 10 = rupestre.
pulvinatum, J. Gay, l. c. 5 = Aucheri.
pumilum, *DC. Syst.* ii. 510.—Helvetia.
pumilum, Gand. Fl. Helv. iv. 355 = lanceolatum.
pumilum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 95.—Am. bor.
pumilum, Schur, in Oestr. Bot. Zeitschr. viii. (1858) 22 = crepidifolium.
purpureum, *Auch. ex Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 53.—Armenia; Syria.
pusillum, *Bory & Chaub. in Bory, Morée*, t. 23.—Graecia.
pusillum, *Gill. in Hook. Bot. Misc.* iii. (1833) 140.—Chili.
pycnophyllum, J. Gay, Exysim. Nov. 3 = thyrsoideum.
pygmaeum, J. Gay, l. c. 6 = Cheiranthus pygmaeus.
pyrenaicum, Jord. Diagn. i. 176 = ochroleucum.
pyrenaicum, Vill. Prosp. 39 = Sisymbrium austriacum.
quadrangulum, Desf. Tabl. ed. I. 129 = Syrenia sessiliflora.
quadricorne, Steph. ex Willd. Sp. Pl. iii. 514 = Tetracme quadricornis.
radicale, Pall. ex DC. Syst. ii. 511 = Leptaleum filifolium.
ramosissimum, Crantz, Stirp. Austr. ed. I. fasc. i. 31; ed. II. 29 = repandum.

ERYSIMUM :—
varifolium, C. Koch, in Linnaea, xix. (1847) 52 = cuspidatum.
Rasiliei, *Bolss. Fl. Orient.* i. 192.—Ins. Creta.
Radowskii, *Weinm. Cat. Hort. Dorp.* (1810) 65.—Sibir.
repandum, *Linn. Amoen. Acad.* iii. 416.—Europ.; Afr. bor.; Aa. occ.
repandum, Losc. & Pard. Ser. Inc. ed. II. 24 = incanum.
retrofractum, Torr. in Bot. U. St. Expl. Exped. ii. 230 = Sisymbrium reflexum.
rhaeticum, DC. Syst. ii. 503 = ochroleucum.
rhaticum, Kunze, ex Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 806 = australe.
Richardsii, F. Muell. Fragm. x. 106 = Sisymbrium Richardsii.
rigens, Jord. Diagn. i. 168 = canescens.
rigidum, DC. Syst. ii. 505 = repandum.
robustum, D. Don, Prod. Fl. Nep. 202 = hieraciifolium.
rubellum, *Reis. in Hoffm. Phyt. Blatt.* i. (1803) 42. —Europ.
runcinatum, Gilib. Fl. Lituan. ii. 58 = Sisymbrium officinale.
runcinatum, Link, Handb. ii. 313 = Sisymbrium runcinatum.
rupestre, Bory & Chaub. Fl. Pélop. t. 25 = Borysmum.
rupestre, *DC. Syst.* ii. 494.—Aa. Min.
rupicola, Schott & Kotschy, in Oestr. Bot. Wochenbl. (1855) 145 = Kotschyanum.
ruscinonense, Jord. Diagn. i. 162 = canescens.
sabulosum, *Fisch. ex Sweet, Hort. Brit.* ed. III. 37.—Sibir.
scabrosum, Banks, ex DC. Syst. ii. 603 = Brassica strigosa.
scabrum, *DC. l. c.* 505.—Syria.
Schaererianum, *Wallr. Sched. Crit.* 363, in obs.—Helvetia.
Schultesii, Schrad. Cat. Hort. Gotting. (1831).—*Cf.* Linnaea, viii. (1833) Lit. 28 = Syrenia sessiliflora.
segetiaceum, Jord. Diagn. i. 168 = cheiranthoides.
sessiliflorum, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 116 = Syrenia sessiliflora.
siculum, Heldr. ex Nym. Consp. 41 = lanceolatum.
sinuatum, *Spreng. Neue Entdeck.* iii. 81.—Sicilia.
sisymbrioides, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 150.—Aa. Min.; Persia; Turkest.
smyrnaeum, *Boiss. & Bal. Diagn.* Ser. II. v. 38.—Aa. Min.
stenophyllum, Boiss. ex Boiss. Fl. Orient. i. 204 = sclpyrium.
stigmatosum, *Franch. Pl. Davidian.* i. (1884) 88.—China.
Stockiisianum, *Boiss. Fl. Orient.* i. 199.—Belochist.
stenchadifolium, *Tausch, in Flora*, xix. (1836) 409.—Sicilia.
strictissimum, C. C. Gmel. Fl. Bad. iii. 77, in syn. = hieraciifolium.
strictum, DC. Prod. i. 197 = pannonicum.
strictum, Gaertn. Mey. & Scherb. Fl. Wett. ii. 451 = hieraciifolium.
striatum, J. Gay, ex Com. in Bull. Soc. Bot. Fr. iv. (1857) 484 = incanum.
strigosum, *DC. Syst.* ii. 506.—Sibir.
strophades, *Boiss. Diagn.* Ser. I. viii. 24.—Mesopotam.
subhastatum, Horsem. Enum. Pl. Hort. Hafn. (1807) 87 = Sisymbrium orientale.
substrigosum, *Rupr. Fl. Cauc.* 75, nomen.—Reg. Caucas.
subulatum, *J. Gay, Erysim. Nov.* 8.—Persia.
suecicum, Jord. Diagn. i. 159 = virgatum.
suffruticosum, Spreng. Nov. Prov. 17 = murale.
sulphureum, Schrank, Baier. Fl. ii. 308 = Sterigma sulphureum.
supinum, Link, Enum. Hort. Berol. ii. 168 = Sisymbrium supinum.
syriacum, *Bieb. Fl. Taur. Cauc.* iii. 441.—Ucrania.
sylvestre, Desf. Tabl. ed. II. 186 = lanceolatum.
sylvestre, Scop. Fl. Carn. ed. II. ii. 28 = canescens.
Szowitsianum, *Boiss. Fl. Orient.* i. 196.—Lac. Ourmah.
tanacetifolium, [Cleirv.] Man. Herb. 219 = Sisymbrium tanacetifolium.
tenellum, *DC. Syst.* ii. 500.—Cappadoc.
tenuifolium, [Cleirv.] Man. Herb. 220 = Diplotaxis tenuifolia.
tenuifolium, Stokes, Bot. Mat. Med. iii. 460 = Barbarea praecox.

ERYSIMUM :—
tenuissimum, *J. Gay, Erysim. Nov.* 7.—Armenia.
Thalianum, Kitt. Taschenb. ed. II. 399 = Sisymbrium Thalianum.
Thomsoni, *Hook. f. in Journ. Linn. Soc.* v. (1861) 165. —Reg. Himal.
thracicum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 79.—Aa. Min.
Tilingi, *J. Gay, Erysim. Nov.* 10.—Am. bor.
transsilvanicum, Schur, Enum. Pl. Transs. 57 = ochroleucum.
tuberosum, Domb. ex DC. Syst. ii. 584 = Cardamine tuberosa.
turritum, Lam. Fl. Fr. ii. 514 = lanceolatum, cheiranthoides.
uncanilcatum, *Boiss. & Huet, Diagn.* Ser. II. v. 28.—Armenia; Syria.
Urvillaeanum, J. Gay, ex Nym. Consp. 41 = smyrnaeum.
verrucosum, Mill. Gard. Dict. ed. VIII. n. 3 = Barbarea praecox.
verrucosum, *Boiss. & Gaill. Diagn.* Ser. II. vi. 12.—Syria.
versicolor, *Andrz. ex DC. Prod.* i. 198.—Reg. Caucas.; Turkest.
violaceum, D. Don, Prod. Fl. Nep. 202 = Cardamine violacea.
virgatum, DC. Prod. i. 197 = hieraciifolium.
virgatum, Lej. Fl. Spa, ii. 71 = murale.
virgatum, *Roth, Catalect.* fasc. i. 75.—Europ.
virgatum, Schleich. ex DC. lc. Gall. Rar. t. 36 = longisiliquosum.
vulgare, Rupr. Fl. Cauc. 91 = Sisymbrium officinale.
Wheleri, *Kohr. in Wheeler, Rep. Bot. U. St. Geogr. Surv.* vi. 64.—Am. bor.
Witmanni, Zawadsk. Enum. 194 = crepidifolium.

ERYSSIMUM, Opiz, Natural, xi. (1826) 422, sphalm. = Bryssimum, Tourn. (Crucif.).

ERYTHALIA, Delarb. Fl. Auv. ed. II. 30 (1800), sphalm. = ERYTHALLIA, Renealm. = Gentiana, Tourn.

ERYTHAEA, Pfeiff. Nom. i. 1256 (1874), sphalm. = Erythraea, Borck. (Gentian.).

ERYTHRA, S. Watts. Bot. Calif. ii. 211 (1880). *PALMAE*, Benth. & Hook. f. iii. 937.
armata, *Watts, l. c.* 212.—Calif.
edulis, *S. Wats. l. c.*—Calif.

ERYTHEREMIA, Endl. Gen. Suppl. ii. 50 (1842) = ERYTHREMIA, Nutt. = Lygodesmia, D. Don (Compos.).

ERYTHRACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 75 (1832) = Ebermaiera, Nees (Acanth.).
acutangulus, Presl, Bot. Bemerk. 97.—Bras.
calycinus, Presl, l. c.—Bras.!
coccineus, Presl, l. c. = Ruellia coccinea.
elongatus, Dala. & Gibs. Bomb. Fl. 184 = Eberm. elongata.
elongatus, Nees, in DC. Prod. xi. 78 = Eber. elongata.
elongatus, Presl, Bot. Bemerk. 97.—Bras.
Griffithianus, Nees, in DC. Prod. xi. 78 = Eberm. Griffithiana.
obtusus, Nees, l. c. = Eberm. obtusa.
obtusus, Wight, ic. t. 1291 = Eberm. Griffithiana.
pedunculosus, Presl, Bot. Bemerk. 97.—Bras.!
pilosus, Presl, l. c. = Ruellia pilosa.
Pohlianus, Presl, l. c.—Bras.!
racemosus, Nees, in Wall. Pl. As. Rar. iii. 80 = Eberm. racemosa.
rubet, Presl, Bot. Bemerk. 97 = Ruellia rubra.

ERYTHRAEA, Renealm. ex Borck. in Roem. Arch. i. t. (1796) 28; Rich. in Pers. Syn. i. 283 (1805). *GENTIANACEAE*, Benth. & Hook. f. ii. 809.
CENTAUREILLA, Delarb. Fl. Auv. ed. II. 30 (1800).
CENTAURIUM, Rupp. Fl. Jen. ed. Hall. 29 (1745).
CENTAURIUM, Gilib. Fl. Lituan. i. 85 (1781); Moench, Meth. 449 (1794).
CYRANDRA, Griseb. in DC. Prod. ix. 44 (1845).
HIPPOCENTAURIA, J. A. Schultes, Oestr. Fl. ed. II. i. 866 (1814).
SCHENKIA, Griseb. in Bonplandia, i. (1853) 236.
XANTHAEA, Reichb. Nom. 123 (1841).

ERYTHRAEA :—
acutiflora, *Schott, in Oken, Isis* (1818), 821.—Am. sept.
aggregata, *Sweet, Brit. Flow. Gard.* Ser. I. ii. t. 127 (= sec. ?).—Hab.?
aggregata, Thouin, ex Mert. & Koch, Deutschl. Fl. ii. 324 = linariifolia.
albiflora, *Kit. in Linnaea*, xxxii. (1863) 481.—Hungar.
alopecuroides, *G. Don, Gen. Syst.* iv. 305.—Hab.?
altaica, Griseb. in DC. Prod. ix. 57 = Meyeri.
anatolica, C. Koch, in Linnaea, xix. (1847) 27 = ramosissima.
angustifolia, Wallr. Sched. Crit. 504 = linariifolia
arenaria, Presl, Delic. Prag. 88 = ramosissima.
australis, *R. Br. Prod.* 451.—Austral.
babylonica, *Griseb. in DC. Prod.* xi. 60.—Persia; Ind. or.
Barrelieri, Dnf. in Bull. Soc. Bot. Fr. vi. (1860) 851 = major.
Beyrichii, *Torr. in Marcy, Rep.* 291. t. 18.—Am. bor. occ.
Boissieri, Willk. in Linnaea, xxv. (1852) 48 = grandiflora.
Cachanlahuen, Roem. & Schult. Syst. iv. 167 = chilensis.
caespitosa, Hoffmgg. & Link, Fl. Port. i. 332. t. 66 = conferta.
calycosa, Beckl. in Proc. Acad. Sc. Philad. 1862 (1863) 7 = chironioides.
Candollii, *Des Moul. in Act. Linn. Soc. Bord.* xx (1855) 597.—Europ.
capitata, Willd. ex Roem. & Schult. Syst. iv. 168 = E. Centaurium.
caspica, Fisch. in Griseb. Gen. et Sp. Gent. 145 = ramosissima.
cretaeroides, Voigt, Hort. Suburb. Calc. 520 = Roxburghii.
Centaurium, Borck. in Roem. Arch. i. t. 30 = Muehlenbergii.
Centaurium, *Pers. Syn.* i. 283.—Europ.
centauroides, Hort. Gotting. ex Griseb. Gen. et Sp. Gent. 141 = ramosissima.
chilensis, Benth. Pl. Hartw. 87 = quitensis.
chilensis, *Pers. Syn.* i. 283.—Chili.
chironioides, *A. Gray, in Bot. Calif.* i. 479.—Mexic. *chironioides*, Torr. in Bot. Mex. Bound. 156 = venusta.
chloodes, Gren. & Godr. Fl. Fr. ii. 684 = conferta.
chloraefolia, Lehm. Pl. Preiss. ii. 289 = Schnea ovata.
cochinchinensis, *Spreng. Syst.* i. 580.—Cochinch.
compar, *R. Br. in Salt, Abyss. App.* 64.—Abyss.
compressa, Hayne, in Kunth, Fl. Berol. 65 = linariifolia.
conferta, *Pers. Syn.* i. 283.—Hispan.
corymbosa, Dalac, Fl. Hautes-Pyr. 458 = E. Centaurium.
corymbosa, *Willd. ex Steud. Nom.* ed. II. i. 595.—Barbaria.
diffusa, Woods, in Hook. Comp. Bot. Mag. ii. (1836) 274 = Massonii.
divaricata, Schaffn. ex Schlecht. in Bot. Zeit. xiii. (1855) 920 = quitensis.
Douglasii, *A. Gray, in Bot. Calif.* i. 480.—Am. bor. occ.
elata, Nutt. ex A. Gray, l. c. = Douglasii.
Elodes, *Roem. & Schult. Syst.* iv. 178.—Mont. Pyren.
emarginata, Waldst. & Kit. Pl. Pan. Hung. iii. 305. t. 275 = ramosissima.
exaltata, Benth. & Hook. f. Gen. ii. 811, in nota, sub Chironia.—Douglasii.
fastigiata, *Wallr. Beitr. Fl. Hercyn.* 186.—Europ.
floribunda, *Benth. Pl. Hartw.* 233.—Calif.
Gerardi, Bauing. Enum. Stirp. Transs. i. 171 = ramosissima.
germanica, Hoffmgg. & Link, Fl. Port. i. 331 = E. Centaurium.
glomerata, *Wittr. in Bot. Notiser* (1854), 115.—Scand.
gracilliflora, *Pomel, Nouv. Mat. Fl. Atl.* 12.—Algeria.
gracilis, *Schleich. Cat. Pl. Helv.* ed. IV. 18.—Helvet.
grandiflora, *Biasoa, Ser. Cent.* 8.—Lusitan.
gypsicola, *Boiss. & Reut. ex Boiss. Diagn.* Ser. I. v. 18.—Hispan.
helodes, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = E. Elodes.
inaperta, Schleicht. Fl. Berol. i. 121 = ramosissima.
intermedia, Pollini, Viagg. 16 = ramosissima.
jurulana, H. B. & K. Nov. Gen. et Sp. iii. 177 = Schultesia stenophylla.
latifolia, Sm. Engl. Fl. i. 328 = E. Centaurium.
linariifolia, Wittr. in Bot. Notiser (1854), 112 = linariifolia.

ERYTHRAEA:—
linariifolia, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 428 = babylonica.
linariifolia, Griseb. Spicil. Fl. Rumel. ii. 62 = E. Centaurium.
linariifolia, *Pers. Syn.* i. 283.—Europ.
linariifolia, Willd. ex Willd. & Lange, Prod. Fl. Hisp. ii. 663 = major.
linearifolia, Boiss. Fl. Orient. iv. 68 = E. Centaurium.
littoralis, Fries, Nov. Fl. Suec. 80 = linariifolia.
littoralis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 338; Sm. Engl. Fl. i. 320 = conferta†
longiflora, Nutt. ex S. Wats. Bot. King's Exped. 277 = Nuttalii.
lutea, Roem. & Schult. Syst. iv. 171 = maritima.
luteola, Pers. Syn. i. 283 = Cicendia pusilla.
macrantha, *Hook. & Arn. Bot. Beech. Voy.* 438.—Mexic.
madrensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 346.—Mexic.
major, Boiss. Voy. Bot. Espagne, 412 = grandiflora.
major, *Hoffmgg. & Link, Fl. Port.* i. 349. t. 65.—Lusitan.
maritima, *Pers. Syn.* i. 283.—Reg. Mediterr.
maritima, Stend. Nom. ed. II. i. 596 = Massoni.
Massoni, *Sweet, Hort. Brit.* ed. II. 363; *H. C. Wats. in Hook. Lond. Journ. Bot.* iii. (1844) 595.—Ins. Azor.
mexicana, Griseb. ex Hook. & Arn. Bot. Beech. Voy. 303 = macrantha.
Meyeri, *Bunge, in Ledeb. Fl. Alt.* i. 220.—As. centr.; Ind. or.
microcalyx, *Boiss. & Reut. Diagn.* Ser. II. iii. 121.—Lusitan.
microphylla, Benth. & Hook. f. Gen. ii. 811, in nota sub Cicendia = Sebaea khasiana.
Muehlenbergii, *Griseb. Gen. et Sp. Gent.* 146.—Am. bor. occ.
Muehlenbergii, Griseb. l. c., partim = ramosissima.
Mussoni, Willk. & Lange, Prod. Fl. Hisp. ii. 664, in syn. sphalm. = Massoni.
nana, Hegetschw. Fl. Schw. 202 = ramosissima.
nudicaulis, *Engelm. in Proc. Am. Acad.* xvii. (1882) 223.—Arizona.
numulariifolia, Stend. Nom. ed. II. i. 596 = Massoni.
Nuttallii, *S. Wats. in Bot. King's Exped.* 276. t. 29, partim.—Am. bor. occ.
Nuttallii, S. Wats. l. c. partim = Douglasii.
occidentalis, Roem. & Schult. Syst. iv. 171 = maritima.
pallida, Willd. ex Griseb. Gen. et Sp. Gent. 138 = chilensis.
paludosa, Schrad. ex Griseb. l. c. 144 = linariifolia.
palustris, Schrad. ex Griseb. l. c. 148 = linariifolia.
paposana, *Phil. Fl. Atac.* 35.—Chili.
pauciflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 373.—Mexic
Pickeringii, Oakes, in Hovey, Mag. ex A. Gray, Syn. Fl. ii. 1. 113 = spicata.
Plumieri, H. B. & K. ex Griseb. Gen. et Sp. Gent. 118 = Eustoma silenifolium.
portensis, Hoffmgg. & Link, Fl. Port. i. 351 = Massoni.
pulchella, Coss. ex Willk. & Lange, Prod. Fl. Hisp. ii. 661 = ramosissima.
pulchella, A. Gray, ex Griseb. in DC. Prod. ix. 60 = Muehlenbergii.
pulchella, Hornem. in Fl. Dan. t. 1637 = ramosissima.
pumila, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 246 = ramosissima.
pyrenaica, Rich. ex Pers. Syn. i. 283 = ramosissima.
quitensis, *H. B. & K. Nov. Gen. et Sp.* iii. 178.—Ecuador.
ramosissima, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 428 = Roxburghii, babylonica.
ramosissima, Griseb. in DC. Prod. ix. 57 = Meyeri.
ramosissima, *Pers. Syn.* i. 283.—Europ.; As. temp.
rhodensis, Boiss. & Reut. Diagn. Ser. II. vi. 121 = E. Centaurium.
Roxburghii, *G. Don, Gen. Syst.* iv. 306.—Ind. or.
sabaeoides, *A. Gray, in Proc. Am. Acad.* vi. (1866) 41.—Ins. Sandvic.
sexguelinea, Mabille, Rech. Pl. Corse, fasc. ii. 45 = major.
sicifolia, Chaub. ex Post, in Bull. Soc. Bot. Fr. vii. (1860) 608 = Massoni.
setacea, Benth. Bot. Voy. Sulph. 128.—Mexic.
spathifolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 50 = ramosissima.
spatiosa, *A. Gray, in Bot. Calif.* i. 479.—Mexic.

ERYTHRAEA:—
spicata, *Pers. Syn.* i. 283.—Reg. Mediterr.; Orient.
stricta, *Schlecht. in Bot. Zeit.* xiii. (1855) 918.—Mexic.
suffruticosa, Stend. ex Griseb. in DC. Prod. ix. 58 = grandiflora.
syivatica, *Kanitz, in Kit. Add. Fl. Hungar.* 127. —Hungar.
szegsardensis, *Menyh. Kalocsa, Vidik-Növény.* 117, nomen.—Hungar.
tenella, Nutt. ex A. Gray, in Bot. Calif. i. 480 = Douglasii.
tenuiflora, Hoffmgg. & Link, Fl. Port. i. 354. t. 67 = ramosissima.
tenuifolia, Griseb. Gen. et Sp. Gent. 143 = linariifolia.
tenuifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 373.—Mexic.
tetramera, *Schlecht. in Bot. Zeit.* xiii. (1855) 919.—Peruv.
texensis, *Griseb. Gen. et Sp. Gent.* 139.—Texas.
trichantha, Durand, in Pacif. Rail. Rep. v. t. 9 = venusta.
trichanthera, *Griseb. Gen. et Sp. Gent.* 146.—Calif.
tripkylla, W. L. E. Schmidt, Diss. t. 1 = tenuifolia.
uliginosa, W. L. E. Schmidt, l. c. 27 = E. Elodes.
uliginosa, Waldst. & Kit. Pl. Rar. Hung. iii. t. 258 = linariifolia.
umiflora, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 255 = Zygostigma uniflorum.
variabilis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 50 = ramosissima.
venusta, *A. Gray, in Bot. Calif.* i. 479.—Calif.
virescens, *Willd. ex Stend. Nom.* ed. II. i. 596.—Creta.
viridiana, *C. Bolle, in Bonplandia*, ix. (1861) 2.—Ins. Can. Verd.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 337 = E. Centaurium.

ERYTHRANTHE, Spach, Hist. Vég. Phan. ix. 312 (1840) = Mimulus, Linn. (Scrophulariin.).
cardinalis, Spach, l. c. 313 = M. cardinalis.

ERYTHRANTHUS, Oerst. ex Hanst. in Linnaea, xxvi. (1855) 215, nomen; et in Vidensk. Selsk.-Skr. V. v. (1861) 120 (Gen. Centr. Am. 46. t. 8) = Allopleurus, Mart. (Gesner.).
coriaceus, Oerst. ll. cc. 215, 120 = A. coriaceus.

ERYTHREMIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 445 = Lygodesmia, D. Don (Compos.).
aphylla, Nutt. l. c. 446 = L. aphylla.
grandiflora, Nutt. l. c. 445 = L. grandiflora.

ERYTHRINA, Linn. Gen. ed. I. 216 (1787). *LEGUMINOSAE*, Benth. & Hook. f. i. 581.
CHIROCALYX, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 97.
CORALLODENDRON, Tourn. ex Rupp. Fl. Jen. ed. Hall. 257 (1745).
DUCHASSAINGIA, Walp. in Linnaea, xxiii. (1850) 741.
HYPAPHORUS, Hassk. Hort. Bogor. Desc. 197 (1858).
MACROCYMBIUM, Walp. in Flora, xxxvi. (1853) 149.
MICROPTERYX, Walp. in Linnaea, xxiii. (1850) 739.
MOURICOU, Adans. Fam. ii. 326 (1763).
STENOTROPIS, Hassk. Retzia, i. 188 (1855).
XYPHANTHUS, Rafin. Fl. Ludov. 108 (1817).
abyssinica, *Lam. Encyc.* ii. 392.—Abyss.
acanthocarpa, *E. Mey. Comm. Pl. Afr. Austr.* 151.—Afr. austr.
acanthophora, Harv. & Sond. Fl. Cap. ii. 609, sphalm. = prec.
aculeatissima, *Desf. Tabl.* ed. I. 191, nomen.—Hab.? *alba*, Roxb. ex Wight & Arn. Prod. 261 = suberosa.
Amasias, *Spruce, in Journ. Linn. Soc.* iii. (1859) 202. —Peruv.
americana, Mill. Gard. Dict. ed. VIII. n. 5 = carnea.
arborescens, *Roxb. Hort. Beng.* 58; *Fl. Ind.* iii. 256.—Reg. Himal.
argentea, Blume, ex Miq. Fl. Ind. Bat. i. 1. 207 = ovalifolia.
bidentata, Lindl. Bot. Reg. (1847) t. 9.
bidentata, Nisell. in Hook. Kew Journ. ix. (1856) 21 = Vespertilio.
blastoua, *Grif. Notul.* iv. 441.—Burma.
brasiensis, *Presl, Symb. Bot.* i. 70.—Am. austr.
brevibracteata, *Mog. & Saese, ex DC. Prod.* ii. 413.—Mexic.
Brucei, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 688.—Afr. trop.
caffra, Blanco, Fl. Filip. ed. II. 394 = ovalifolia.
caffra, Ker-Gawl. in Bot. Reg. t. 786 = Humeana.

ERYTHRINA:—
caffra, *Thunb. Prod. Pl. Cap.* 121.—Afr. austr.
carnea, Blanco, Fl. Filip. ed. I. 564 = indica.
carnea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 8.—Am. austr.
compacta, Bull, Cat. (1871) 4; Rev. Hortic. (1882) 248 = E. Crista-galli†
corallifera, Salisb. Prod. 335 = E. Corallodendron.
Corallodendron, Herb. Madr. ex Wall. Cat. n. 5959 = suberosa.
Corallodendron, Lam. Encyc. ii. 390 = indica.
Corallodendron, *Linn. Sp. Pl.* 706.—Am. bor.; Ind. occ.
Corallodendron, Lour. Fl. Cochinch. ii. 497 = Loureiri.
corallioides, *Mog. & Sesse, ex DC. Prod.* ii. 413.—Mexic.
cordyloia, Jacq. ex DC. l. c. 414 = Rudolphia volubilis.
Crista-galli, *Linn. Mant.* i. 99.—Bras.
cubensis, *Wright, in Sauv. Fl. Cub.* 26.—Cuba.
cuneata, R. Grah. in Wall. Cat. n. 5967 = indica.
diversiora, *Mog. & Sesse, ex DC. Prod.* ii. 414.—Mexic.
eqneandra, *DC. Cat. Hort. Monsp.* 109.—Hab.?
erythrostachia, *Morr. in Ann. Soc. Gand*, v. (1849) 457.—Hab.?
excelsa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 183.—Afr. trop.
falcata, *Benth. in Mart. Fl. Bras.* xv. i. 173.—Bras.
fusciculata, Benth. in Linnaea, xxii. (1849) 518 = E. Crista-galli.
fusa, *Presl, Symb. Bot.* i. 69.—Hab.?
fulgens, *Sweet, Hort. Brit.* ed. I. 130, nomen.—Ind. or.
fusca, *Lour. Fl. Cochinch.* 627.—Cochinch.
glauca, *Willd. in Ges Naturf. Freunde Neue Schr.* iii. (1801) 428.—Venezuela.
Greeffrei, Tineo, Cat. Fl. Hort. Panorm. (1827) 278 = speciosa.
guineensis, *G. Don, Gen. Syst.* ii. 371 = senegalensis.
Hamiltoniana, Stend. Nom. ed. II. i. 596 = suberosa.
hastifolia, Bertol. f. in Mem. Acc. Sc. Bolog. ii. (1850) 568 = Humeana.
hederaefolia, Spreng. Syst. iii. 244 = herbacea.
herbacea, *Linn. Sp. Pl.* 706.—Am. bor.; Ind. occ.
holosericea, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 69.—Burma.
horrida, *Mog. & Sesse, ex DC. Prod.* ii. 413.—Mexic.
huillensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 183.—Afr. trop.
Humeana, *Spreng. Syst.* iii. 243.—Afr. austr.
Humei, E. Mey. Comm. Pl. Afr. Austr. 150 = prasc.
humilis, Salisb. Prod. 335 = herbacea.
incana, *Wild. Enum. Hort. Berol. Suppl.* 50, nomen. —Am. austr.
indica, *Lam. Encyc.* ii. 391.—As. trop.; Austral.
indica, Zoll. in Nat. en Geneesk. Arch. Neêrl. Ind. iii. (1846) 61 = lobulata.
inermis, *Abbt. Pl. Gui.* ii. 760.—Guian.
inermis, Mill. Gard. Dict. ed VIII. n. 6 = mitis.
insignis, *Tod. Nuovi Gen. e Sp.* 66.—Hab.?
lapsiala, Lam. Encyc. ii. 393 = Camptosema erythrinoides.
lacta, *Dekuh. Cat. Pl. Hort. Camald.* ed. II. 20.—Hab.?
latifolia, Schum. & Thonn. Beskr. Guin. Pl. 238 = senegalensis.
latissima, E. Mey. Comm. Pl. Afr. Austr. 151 = tomentosa.
laurifolia, Jacq. Obs. Bot. iii. 1 = E. Crista-galli.
leptorrhiza, *Mog. & Sesse, ex DC. Prod.* ii. 413.—Mexic.
lithosperma, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 209.—Burma: Malaya.
Livingstoniana, *Baker, in Oliver, Fl. Trop. Afr.* ii. 182.—Afr. trop.
lobulata, *Miq. Fl. Ind. Bat.* i. 1. 208.—Java.
longipes, *Mog. & Sesse, ex DC. Prod.* ii. 413.—Mexic.
Loureiri, *G. Don, Gen. Syst.* ii. 372.—Cochinch.
macrophylla, *DC. Prod.* ii. 411.—Hab.?
marmorata, *Veitch, ex Planch. in Fl. des Serres*, xxiii. (1880) 21.—Hab.?
maxima, Roxb. ex Wight & Arn. Prod. 261 = suberosa.
mediterranea, *Vell. Fl. Flum.* 308; vii. t. 103.—Bras.
microptera, *Poepp. ex Walp. in Linnaea*, xxiii. (1850) 740.—Peruv.
mitis, *Jacq. Hort. Schoenb.* ii. 47. t. 216.—Venezuela.
monosperma, *Gaudich. in Freyc. Voy. Bot.* 486. t. 114. —Ins. Sandvic.
monosperma, Lam. Encyc. ii. 391 = Butea frondosa.
Moodi, *Tod. Hort. Bot. Panorm.* ii. 7. t. 96.—Ind. or.?
Mulungu, *Mart. ex Benth. in Mart. Fl. Bras.* xv. i. 173.—Bras.

ERYTHRINA:—
Nakaseta, Buch.-Ham.ex Wall. Cat. n. 5966 = suberosa.
nervosa, *DC. Prod.* ii. 413.—Bras.
orientalis, Merr. in Comm. Gotting. viii. (1787) 35. t. 1 = indica.
ovalifolia, *Roxb. Hort. Beng.* 53; *Fl. Ind.* iii. 254.—Ind. or.; Malaya.
ovalis, *Wall. ex Steud. Nom.* ed. II. i. 596.—Ind. or.
pandurneformis, Blume, ex Miq. Fl. Ind. Bat. i. 1. 909 = lobulata.
Parcellii, *Hort. Bull. in Gard. Chron.* (1874) ii. 393. f. 89.—Ins. Pacif.
patens, *Moç. & Sesse, ex DC. Prod.* ii. 414.—Mexic.
poiteyana, Fenzl, in Flora, xxvii. (1844) 312 = tomentosa.
picta, Linn. Sp. Pl. ed. II. 998 = indica?
picta, Wall. Cat. n. 5960 = ovalifolia.
placidoides, Sweet, Hort. Brit. ed. I. 120; DC. Prod. ii. 414 = acq.?
Placipula, Linn. Sp. Pl. 707 = Placidia Erythrina.
plantaginea, Linn. Syst. ed. X. 1158 = Rudolphia pelleta.
polanthes, *Brot. in Trans. Linn. Soc.* xiv. (1824) 342. —Am. austr.
polysperma, Pritz. Ic. Ind. i. 436, sphalm. = monosperma.
portoricensis, Desf. Tabl. ed. I. 191 = Rudolphia volubilis.
Princeps, *A. Distr. in Otto & Dietr. Allg. Gartens.* ii. (1834) 305.—Mexic.
pulcherrima, *Tod. Nuovi Gen. e Sp.* 70.—Hab.?
Raja, Mclean, in Hook. Lond. Journ. Bot. ii. (1843) 96 = Humeana.
reniformis, Buch.-Ham. in Wall. Cat. n. 5968 = suberosa.
resupinata, Mor. Syst. Vers. Zoll. 4 = lobulata.
resupinata, *Roxb. Hort. Beng.* 53; *Fl. Ind.* iii. 257.—Ind. or.
reticulata, *Presl, Symb. Bot.* ii. 22. t. 68.—Mexic.
rosea, *A. Distr. in Otto & Dietr. Allg. Gartens.* ii. (1834) 258.—Mexic.
rubicunda, *Zipp. ex Span.* 75.—Hab.?
rubrinervia, *H. B. & K. Nov. Gen. et Sp.* vi. 434.—N. Granat.
Schottii, Harv. Thes. Cap. i. 39 = tomentosa.
secundiflora, Benth. in *Trans. Linn. Soc.* xix. (1824) 346. t. 12.—Bras.
secundiflora, Hamk. Pl. Jav. Rar. 378 = lithosperma.
senegalensis, *DC. Prod.* ii. 413.—Afr. trop.
senegalensis, A. Rich. Tent. Fl. Abyss. i. 213 = Brucei.
setosa, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 194.—Mexic.
spathacea, *DC. Prod.* ii. 412.—Ins. S. Doming.
spathacea, Wall. Cat. n. 5965 = indica.
speciosa, *Andr, Bot. Rep.* t. 448.—Ind. occ.
speciosa, Tod. Ind. Sem. Hort. Panorm. (1860) 11 = E. Crista-galli.
spinosa, Mill. Gard. Dict. ed. VIII. n. 3 = E. Corallodendron.
spinosa, *Voigt, in Syll. Ratisb.* ii. (1828) 55.—Hab.?
stricta, *Roxb. Hort. Beng.* 53; *Fl. Ind.* iii. 251.—Ind. or.
suberifera, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 132.—Afr. trop.
suberosa, *Roxb. Hort. Beng.* 53; *Fl. Ind.* iii. 253.—Ind. or.
sublobata, Roxb. ll. cc. [96], 254 = suberosa.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 304.—Malaya.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 80.—Ins. Tahiti.
tomentosa, *R. Br. in Salt, Abyss. App.* p. 65, nomen; *A. Rich. Tent. Fl. Abyss.* i. 213.—Abyss.
tomentosa, Benth.-Ham. in Wall. Cat. n. 5964 = suberosa.
triacantha, *Willd. Enum. Hort. Berol. Suppl.* 50, nomen.—Hab.?
umbrosa, *H. B. & K. Nov. Gen. et Sp.* vi. 434.—Am. austr.
variegata, Linn. Dia. Herb. Amb. in Amoen. Acad. iv. 132 = indica.
velutina, Jacq. Hort. Schoenb. iv. t. 406 = arenarden.
velutina, *Willd. in Ges. Naturf. Freunde Neue Schr.* iii. (1801) 426.—Venezuela.
versa, *Vell. Fl. Flum.* 304; vii. t. 102.—Bras.
versicolor, *Boj. Hort. Maurit.* 110, nomen.—Madag.
Vespertilio, *Benth. in Mitch. Journ. Trop. Austral.* 218.—Austral.

ERYTHRINA:—
viarum, *Tod. Nuovi Gen. e Sp.* 62.—Hab.?
Vogelii, Hook. f. in Hook. Niger Fl. 307 = senegalensis.
Zeyheri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 286.—Afr. austr.

ERYTHROCARPUS, Blume, Bijdr. 604 (1822) = **Gelonium**, Roxb (Euphorb.).
glomeruliatus, Blume, l. c. 605 = G. bifarium.
spicatus, Blume, l. c. = G. spicatum.

ERYTHROCARPUS, M. Roem. Syn. Pepon. 264 (1846) = **Modecca**, Lam. (Passiflor.).
populifolius, M. Roem. l. c. = M. populifolia.

ERYTHROCEPHALUM, Benth. in Benth. & Hook. f. Gen. ii. 488 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
longifolium, *Benth. ex Oliver, in Trans. Soc. Linn.* xxix. (1873) 102.—Afr. trop.
minus, *Oliver, l. c.* 103.—Afr. trop.
nutans, *Benth. ex Oliver, l. c.* 102.—Afr. trop.
zambesianum, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 441.—Afr. trop.

ERYTHROCHAETE, Sieb. & Zucc. in Abh. Acad. Muench. iv. iii. (1846) 188 (Fam. Nat. Fl. Jap. ii. 64) = **Senecio**, Tourn. (Compos.).
denisate, A. Gray, in Mem. Am. Acad. N. S. vi. (1858-59) 395 (sp. dub.).—Japon.
palmatifida, Sieb. & Zucc. in Abh. Acad. Muench. iv. iii. (1846) 188 = S. japonicus.

ERYTHROCHILUS, Reinw. ex Flora, viii. (1825) i. 108 = **Claoxylon**, A. Juss. (Euphorbiac.).
indicus, Reinw. l. c. 615 = C. indicum.
longifolius, Blume, Bijdr. 616 = C. longifolium.
minor, Blume, l. c. = C. indicum.
mollis, Blume, l. c. 615 = C. indicum.
multiflorus, Zipp. ex Span. in Linnaea, xv. (1841) 849. —Ins. Timor.

ERYTHROCHITON, Griff. in Proc. Linn. Soc. i. (1846) 283 = **Ternstroemia**, Linn.
Wallichianum, Griff. l. c. = T. penanglana.

ERYTHROCHITON, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 151, 165. *RUTACEAE*, Benth. & Hook. f. i. 284.
PENTAMORPHA, Scheidw. in Bull. Acad. Brux. ix. i. (1842) 20.
TOXOSIPHON, Baill. Adansonia, x. (1872) 311.
brasiliensis, *Nees & Mart. l. c.* 170. t. 25.—Bras.
hypophyllanthus, *Planch. & Linden, in Ann. Sc. Nat.* Sér. III. xix. (1853) 75.—N. Granat.
Lindeni, *Planch. & Linden, Hort. Lind.* 35. t. 13; *Hemsl. Biol. Centr. Am. Bot.* i. 166.—Mexic.
macrophyllum, Makoy, ex Hook. Bot. Mag. t. 4948 = Cusparia microphylla.

ERYTHROCHYLUS, Reinw. ex Blume, Cat. Gew. Buitenz. 110, nomen (1823) = ERYTHROCHILUS, Reinw. = Claoxylon, A. Juss. (Euphorbiac.) (1844).

ERYTHROCOCCA, Benth. in Hook. Niger Fl. 506 (1849). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 306.
scalenta, *Benth. l. c.*—Afr. trop.

ERYTHRODANUM, Thou. Fl. Tristan d'Acuga. 41 (1811) = **Wurtzia**, Banks & Soland. (Rubiac.).
alsinaeflorum, Thou. l. c. 42 = N. depressa.
majus, Thou. l. c. t. 11 = N. depressa.

ERYTHRODES, Blume, Bijdr. 410. t. 72 (1825) = **Physurus**, Rich. (Orchid.).
latifolia, Blume, l. c. 411 = P. latifolius.

ERYTHROGYNE, Vis. in Cesp. Ricensch. Caprif. e Fic. 86 (1845) = **Ficus**, Tourn. (Urticac.).
luteacens, Vis. l. c. = F. diversifolius.

ERYTHROLAENA, Sweet, Brit. Flow. Gard. Ser. I. t. 184 (1825) = **Cnicus**, Linn. (Compos.).
conspicua, Sweet, l. c. = C. conspicuus.

ERYTHRONIUM, Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 819.
DENS-CANIS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 197 (1745).
MITHRYDATIUM, Adans. Fam. ii. 48 (1763).
albidum, *Nutt. Gen. Am.* i. 223.—Am. bor.
alticaum, Bess. ex Baker, in Journ. Linn. Soc. xiv. (1874) 297 = E. Dens-canis.
americanum, *Ker-Gawl. in Sims, Bot. Mag.* t. 1118.—Am. bor.
angustatum, Rafin. in Med. Repos. N. York, II. v. (1808) 354 = E. Dens-canis.
aquatile, Salisb. in Trans. Hort. Soc. i. (1812) 381 = americanum.
aureum, Hort. Angl. ex Loisel. Herb. Amat. i. sub t. 51; Link, Enum. Hort. Berol. i. 328 = americanum.
bifolium, Sweet, Hort. Brit. ed. II. 597 = E. Dens canis.
bracteatum, Bigel. in Beck, Bot. 365 = albidum.
bulbosum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = E. Dens-canis.
caninum, Dulac, Fl. Hautes-Pyr. 113 = E. Dens canis.
carolinianum, Walt. Fl. Carol. 122 = americanum.
Dens-canis, *Linn. Sp. Pl.* 305.—Europ.; As. bor. *Dens-canis*, Michx. Fl. Bor. Am. i. 193 = americanum.
flavescens, Loisel. Herb. Amat. i. 51 = americanum.
flavum, Sm. in Rees, Cyclop. xiii. n. 3 = americanum.
gigantteum, Lindl. Bot. Reg. t. 1786 = grandiflorum.
grandiflorum, Benth. Pl. Hartw. 339 = Hartwegi.
grandiflorum, *Pursh, Fl. Am. Sept.* i. 231.—Am. bor occ.
Hartwegi, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 261.—Am. bor. occ.
Indicum, Kotll. ex Spreng. Syst. ii. 97 = Scilla indica.
japonicum, Polt. in Rev. Hort. Sér. IV. iii. (1854) 284 = grandiflorum.
lanceolatum, Pursh, Fl. Am. Sept. i. 230 = americanum.
latifolium, Schur, Enum. Pl. Transs. 663 = E. Dens-canis.
longifolium, Mill. Gard. Dict. ed. VIII. n. 2 = E. Denscanis.
maculatum, DC. Fl. Fr. iii. 197 = E. Dens-canis.
maculosum, Lam. Fl. Fr. iii. 286 = E. Dens-canis.
maximum, Dougl. ex Baker, in Journ. Linn. Soc. xiv. (1874) 298 = grandiflorum.
Nuttallianum, Schult. f. Syst. vii. 1681 = americanum.
obtusiflorum, Opiz, Seznam, 42 = E. Dens-canis?
ovatifolium, Polr. Encyc. viii. 660 = E. Dens-canis.
propullans, *A. Gray, in Am. Natural.* v. (1871) 300. t. 7A.—Am. bor.
purpurascens, *S. Wats. in Proc. Am. Acad.* xii. (1877) 277.—Calif.
revolutum, Baker, in Gard. Chron. (1876) i. 138 = purpurascens.
revolutum, Sm. in Rees, Cyclop. xiii. n. 2 = grandiflorum.
speciosum, Nutt. ex Baker, in Journ. Linn. Soc. xiv. (1874) 298 = grandiflorum.
vernale, Salisb. Prod. 238 = E. Dens-canis.

ERYTHROPALLA, Hamk. Cat. Hort. Bog. Alt. 191 (1844) = Erythropalum, Blume (Olacin.).

ERYTHROPALUM, Blume, Bijdr. 921 (1826). *OLACINEAE*, Benth. & Hook. f. i. 347.
BALIOSPATHUM, Blanco, Fl. Filip. ed. I. 187 (1837).
DACTYLIUM, Griff. in Proc. Linn. Soc. ii. (1853) 252.
DECASTROPHIA, Griff. Notul. iv. 737 (1854): Ic. Pl. Asiat. t. 618, 614.
ERYTHROPALUM, Blume, Fl. Jav. praef. p. vii (1828).
MACKAYA, Ann. in Jard. Mag. Zool. & Bot. ii. (1838) 250.
MODECCOPSIS, Griff. in Proc. Linn. Soc. i. (1845) 171.
populifolium, *Mast. in Hook. f. Fl. Brit. Ind.* i. 678. —Ind. or.
scandens, *Blume, Bijdr.* 922.—Reg. Himal.; Malaya. vagans, *Mast. in Hook. f. Fl. Brit. Ind.* i. 678.—Reg. Himal.; Malaya.

ERYTHROPHILA, Arn. in Hook. Journ. Bot. iii. (1841) 356, sphalm. = Erythrophysa, E. Mey. (Sapindac.)

ERYTHROPHLAEUM, Reichb. Consp. 157 (1828) = seq.

ERYTHROPHLEUM, Afzel. ex R. Br. in Tuckey, Congo, 480 (1818). *LEGUMINOSAE*, Benth. & Hook. f. i. 583.
FILLAEA, Guill. & Perr. Fl. Seneg. Tent. i. 242. t. 55 (1833).
LABOUCHERIA, F. Muell. in Journ. Linn. Soc. iii. (1859) 126.
MAVIA, Bertol. f. in Mem. Acc. Bolog. ii. (1850) 870. t. 20.
chirostachya, Baill. Hist. Pl. ii. 150 = Laboucheri.
Commingo, *Bull. Adansonia*, x. (1871) 105.—Madag.
Fordii, *Oliver, in Hook. Ic. Pl.* t. 1409.—China.
guineense, *G. Don, Gen. Syst.* ii. 434.—Afr. trop.
judiciale, Procter, in Am. Journ. Pharm. xviii. (1852) 195 = guineense.
Labouchetii, *F. Muell. ex Benth. Fl. Austral.* ii. 297. —Austral.
ordale, Bolle, in Peters, Reise Mossamb. Bot. i. 10 = guineense.

ERYTHROPHYSA, E. Mey. Zwei Pfl. Docum. 183 (1848); et ex Harv. & Sond. Fl. Cap. i. 237 (1860). *SAPINDACEAE*, Benth. & Hook. f. i. 397.
ascalina, *Bail. Adansonia*, xi. (1878-76) 289.—Madag.
undulata, *E. Mey. Zwei Pfl. Docum.* 188; *et ex Harv. & Sond. Fl. Cap.* i. 238.—Afr. austr.

ERYTHROPOGON, DC. Prod. vi. 254 (1837) = **Metalasia**, R. Br. (Compos.).
imbricatum, DC. l. c. 255 = M. uniflora.
umbellatum, DC. l. c. = M. umbellata.

ERYTHROPSIS, Lindl. in Brand. Journ. Sc. (Sept. 1827) 119, in obs. = **Sterculia**, Linn.
Roxburghiana, Schott & Endl. Melet. 33 = S. alata.

ERYTHRORCHIS, Rhume, Rumphia, i. 200. t. 70 (1835) = **Galeola**, Lour. (Orchid.).
altissima, Blume, l. c. = G. altissima.
aphylla, F. Muell. Fragm. ii. 167 = G. casuthoides.
foliata, F. Muell. l. c. = G. foliata.
Kuhlii, Reichb. f. Xenia Orch. ii. 78 = G. Kuhlii.

ERYTHRORHIZA, Michx. Fl. Bor. Am. ii. 84. t. 36 (1803) = **Galax**, Linn. (Diapens.).
rotundifolia, Michx. l. c. 85 = G. aphylla.

ERYTHROSPERMUM, Lam. Tabl. Encyc. ii. 407. t. 374 (1791). *BIXINEAE*, Benth. & Hook. f. i. 127.
PECTINEA, Gaertn. Fruct. ii. 126. t. 111. f. 2 (1791).
amplexicaule, DC. Prod. i. 258 = verticillatum.
amplifolium, Thou. Vég. Iles Austr. Afr. t. 21. f. 2 = verticillatum.
coronarium, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 338.—Madag.
cuneifolium, *Boj. Hort. Maurit.* 19.—Ins. Maurit.
elliptieum, Poir. Encyc. Suppl. ii. 585 = verticillatum.
erythroxyloides, Boj. Hort. Maurit. 19 = verticillatum.
lanceolatum, *Boj. l. c.*—Ins. Maurit.
laxiflorum, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 258 = verticillatum.
leucocarpum, Clos, l. c. = Denhamia obscura.
longifolium, Boj. Hort. Maurit. 19 = verticillatum.
macrophyllum, Poir. Encyc. Suppl. ii. 585 = verticillatum.
mauririanum, Baker, Fl. Maurit. 11 = verticillatum.
mauritianum, Thou. Vég. Iles Austr. Afr. 67 = verticillatum.
paniculatum, Poir. Encyc. Suppl. ii. 585 = verticillatum.
parviflorum, Thou. Vég. Iles Austr. Afr. t. 21. f. 1 = verticillatum.
phytolaccoides, *Gardn. in Calc. Journ. Nat. Hist.* vii. (1847) 451.—Zeylan.
polyandrum, *Oliver, in Hook. Ic. Pl.* t. 1333.—Ins. Ramos.
pubescens, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 258 = verticillatum.
pyrifolium, Lam. Tabl. Encyc. ii. 407 = verticillatum.
verticillatum, *Lam. l. c.*—Ins. Mascar.

ERYTHROSTAPHYLE, Hasse, in Journ. Bot. xi. (1873) 246 = **Xodes**, Blume (Oleacin.).
salicifolia, Staxon, l. c. = I. virgiana.

ERYTHROSTEMON, Klotzsch, in Link, Klotzsch, & Otto, Ic. Pl. i. 97. t. 39 (1841) = **Caesalpinia**, Linn. (Legumin.).
Gilliesii, Klotzsch, l. c. 98 = C. Gilliesii.

ERYTHROSTICTUS, Schlecht. in Linnaea, i. (1826) 90 = **Androsaemum**, Willd. (Liliac.).
europaeus, Lange, in Kjoeb. Vidensk. Meddel. (1860) 69 = A. punctatum.
graminaeus, Schlecht. in Linnaea, i. (1826) 90 = A. punctatum.
palaestinus, Boiss. ex Baker, in Journ. Linn. Soc. xvii (1879) 445 = A. palaestinum.
punctatus, Schlecht. in Linnaea, i. (1826) 90 = A. punctatum.

ERYTHROSTIGMA, Hassk. in Flora, xxv. (1842) ii. Beibl. 45 = Connarus, Linn.
diversifolia, Hassk. l. c.—Malaya.
elliptieum, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 174.—Malaya.
macrophyllum, Zoll. l. c.—Malaya.
obliquum, Zoll. l. c.—Malaya.
villosum, Zoll. l. c. 175.—Malaya.

ERYTHROTIS, Hook. f. Bot. Mag. t. 6150 (1875) = Cynanchum, D. Don (Connelin.).
Beddomei, Hook. f. l. c. = C. kewensis.

ERYTHROXYLON, Linn. Syst. ed. X. 1085 (1759) = seq.

ERYTHROXYLUM, P. Br. Hist. Jamaic. 278 (1756). *LINEAE*, Benth. & Hook. f. i. 344
ROZLANA, Comm. ex DC. Prod. i. 575 (1824).
SETHIA, H. B. & K. Nov. Gen. et Sp. v. 175 (1821).
STRUDELIA, Spreng. Neue Entdeck iii. 59 (1822).
acaminatum, *Ruiz & Pav. Fl. Per.* iv. t. 399.—Peruv.
acuminatum, Walp. Rep. i. 407 = lucidum, *Moon.*
acutifolium, *Standl. ex Peyr. in Mart. Fl. Bras.* xii. i. 168.—Bras.
affine, A. Rich. Ess. Fl. Cub. 258 = obovatum.
affine, A. St. Hil. Fl. Bras. Mer. ii. 97 = Kunthianum, *A. St. Hil.*
alsterniifolium, *A. Rich. Ess. Fl. Cub.* 256.—Cuba.
amazonicum, *Peyr. in Mart. Fl. Bras.* xii. i. 167 — Bras.
ambiguum, *Peyr. in Mart. Fl. Bras.* xii. i. 142.—Bras.
americanum, *Fisch. ex Steud. Nom. ed.* I. 319, nomen. — Hab. ?
amplum, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 372.—Bras.
angustifolium, *Mart. in Flora*, xxi. (1838) ii. *Beibl.* 84. — Bras.
areolatum, Billh. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 186, in nota = Bumelia ferox.
areolatum, *Linn. Syst.* ed. X. 1085.—Ind. occ.
areolatum, Poepp. ex Griseb. Fl. Brit. W. Ind. 118 = obtusum.
areolatum, Vell. Fl. Flum. 194; iv. t. 175 = suberosum.
aristiguerum, *Peyr. in Mart. Fl. Bras.* xii. i. 157.—Bras.
australe, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 22.—Austral.
bahiense, *Peyr. in Mart. Fl. Bras.* xii. i. 160.—Bras.
bentiacum, *Mart. in Abh. Akad. Munech.* iii. ii. (1840) 390.—Am. austr.
Bongardianum, *Meyer, ex Peyr. in Mart. Fl. Bras.* xii. i. 145.—Bras.
brevipedicellum, St. Leg. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = brevipes.
brevipes, *DC. Prod.* i. 573.—Ind. occ.
burmanicum, *Griff. Notul. iv.* 668.—Burma; Malaya.
buxifolium, *Lam. Encyc.* ii. 394.—Madag.
Buxus, *Peyr. in Mart. Fl. Bras.* xii. i. 137.—Bras.
cagfrum, Sond. in Linnaea, xxiii. (1850) 22 = emarginatum.
campestre, *A. St. Hil. Fl. Bras. Mer.* ii. 97.—Bras.
carthagenense, Jacq. Enum. Pl. Carib. 21 = areolatum.
casuarinoides, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 340.— N. Granat.
catarinense, *Spruce, ex Peyr. in Mart. Fl. Bras.* xii. i. 149.—Bras.
cinclanthum, *Mart. in Abh. Akad. Munech.* iii. ii. (1840) 357.—Bras.
citrifolium, *A. St. Hil. Fl. Bras. Mer.* ii. 94.—Bras.

ERYTHROXYLUM:—
Coca, *Lam. Encyc.* ii. 393.—Peruv.
coelophlebium, *Mart. in Abh. Akad. Munech.* iii. ii. (1840) 396.—Bras.
columbinum, *Mart. l. c.* 379.—Bras.
compressum, *Peyr. in Mart. Fl. Bras.* xii. i. 163.— Bras.
concinnatum, Pritz. Ic. Ind. i. 486, err. typ. = cinnatum.
cornutum, Steud. ex Peyr. in Mart. Fl. Bras. xii. i. 174 = mucronatum.
cotinifolium, A. St. Hil. Fl. Bras. Mer. ii. 98 = campestre.
cumanense, *H. B. & K. Nov. Gen. et Sp.* v. 177.— Venezuela.
cuneatum, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 135 = burmanicum.
cuspidifolium, *Mart. in Abh. Akad. Munech.* iii. ii. (1840) 359.—Bras.
Daphnites, *Mart. l. c.* 363.—Bras.
decadnum, A. St. Hil. Fl. Bras. Mer. ii. 95 = nitidum.
discolor, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 184.—Madag.
distortum, *Mart. in Abh. Akad. Munech.* iii. ii. (1840) 346.—Bras.
divaricatum, *Peyr. in Mart. Fl. Bras.* xii. i. 146.— Bras.
ectinocalyx, *Mart. in Abh. Akad. Munech.* iii. ii. (1840) 355.—Bras.
ellipticum, *R. Br. ex Benth. Fl. Austral.* i. 384.— Austral.
ellipticum, *Peyr. in Mart. Fl. Bras.* xii. i. 145.— Bras.
emarginatum, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 224.—Afr. trop
eugenifolium, Mart. ex Peyr. in Mart. Fl. Bras. xii. i. 150 = Kunthianum, *A. St. Hil.*
exaltatum, *Benj. ex Peyr. l. c.* 151.—Bras.
ferrugineum, *Cav. Diss.* 404.—Madag.
ficifolium, *Peyr. in Mart. Fl. Bras.* xii. i. 169.— Bras.
fimbriatum, *Peyr. l. c.* 162.—Bras.
firmum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 327. —Madag.
flaccidum, *Salzm. ex Peyr. in Mart. Fl. Bras.* xii. i. 141.—Bras.
floribundum, *Mart. in Abh. Akad. Munech.* iii. ii. (1840) 398.—Bras.
floribundum, Seem. Bot. Voy. Herald, 90 = macrophyllum.
frangulaefolium, *A. St. Hil. Fl. Bras. Mer.* ii. 99.— Bras.
Gaudichaudii, *Peyr. in Mart. Fl. Bras.* xii. i. 169.— Bras.
Gerrardi, *Baker, in Journ. Linn. Soc.* xx. (1883) 109. —Madag.
glaucocalyx, *Peyr. in Mart. Fl. Bras.* xii. i. 154.— Bras.
gomphioides, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 341.—N. Granat.
gracilipes, *Peyr. in Mart. Fl. Bras.* xii. i. 159.—Bras.
grandifolium, *Peyr. l. c.* 161.—Bras.
Griesbachii, *Peyr. l. c.*—Bras.
havanense, H. B. & K. Nov. Gen. et Sp. v. 177 = obtusum.
havanense, Jacq. Enum. Pl. Carib. 21 = ovatum.
havanense, A. Rich. Ess. Fl. Cub. 257 = squamatum.
hondense, *H. B. & K. Nov. Gen. et Sp.* v. 176.—N. Granat.
hypericifolium, *Lam. Encyc.* ii. 394.—Ins. Mascar.
indicum, Bedd. Fl. Sylvat. t. 81 = monogynum.
intermedium, *Peyr. in Mart. Fl. Bras.* xii. i. 151.— Bras.
josinickiea, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 184.—Madag.
Kapplerianum, *Peyr. in Mart. Fl. Bras.* xii. i. 159.— Bras.
Kunthianum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 294.—Ind. or.; Burma.
Kunthianum, *A. St. Hil. Fl. Bras. Mer.* ii. 96.— Bras.
lanceolatum, *Walp. Rep.* i. 407.—Ind. or.
lanceum, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 185.—Ins. Comoro.
lanciifolium, *Peyr. in Mart. Fl. Bras.* xii. i. 144.— Bras.
laurifolium, *Lam. Encyc.* ii. 394.—Ins. Mascar.

ERYTHROXYLUM :—
laxinum, *Planch. & Triana, in Ann. Sc. Nat. Sér.* IV. xviii. (1862) 341.—N. Granat.
ligustrinum, *DC. Prod.* i. 574.—Guiana.
lineolatum, *DC. l. c.* 575.—Guiana.
longifolium, Lam. Encyc. ii. 334 = laurifolium.
lucidum, *H. B. & K. Nov. Gen. et Sp.* v. 179.—N. Granat.
lucidum, *Maun, Cat. Pl. Ceyl.* 36.—Zeylan.
macrocalyx, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 384.—Bras.
macrochaetum, *Miq. in Linnaea*, xxii. (1849) 798.—Bras.
macronemium, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 402.—Peruv.
macrophyllum, *Cav. Diss.* 401. t. 237.—Guiana.
macrophyllum, Mart. in Abh. Akad. Muench. iii. 11. (1840) 369 = Martii.
magnifolium, *A. St. Hil. Fl. Bras. Mer.* ii. 93; *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 400.—Bras.
Mamacrux, *Mart. l. c.* 385.—Peruv.
Mangifilia, Ucopp. ex Peyr. in Mart. Fl. Bras. xii. 1. 165 = citrifolium.
Manail, *Oliver, Fl. Trop Afr.* i. 274.—Afr. trop.
Martii, *Peyr. in Mart. Fl. Bras.* xii. 1. 160.—Bras.
mauritianum, Rottl. ex Wall. Cat. n. 6851 = laurifolium.
mexicanum, *H. B. & K. Nov. Gen. et Sp.* v. 178.—Mexic.
micranthum, *Bong. ex Peyr. in Mart. Fl. Bras.* xii. 1. 164.—Bras.
microphyllum, *A. St. Hil. Fl. Bras. Mer.* ii. 100.—Bras.
Mikanii, *Peyr. in Mart. Fl. Bras.* xii. 1. 153.—Bras.
minutifolium, *Griseb. Cat. Pl. Cub.* 41.—Cuba.
monogynum, *Roxb. Pl. Corom.* i. 61. t. 88.—Ind. or.; China.
mucronatum, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 372.—Guiana.
Myrsinites, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 345.—Bras.
myrtoides, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 185.—Madag.
nanum, A. St. Hil. Fl. Bras. Mer. ii. 97 = nitidum.
nitidum, *Baker, in Journ. Linn. Soc.* xx. (1883) 110. —Madag.
nitidum, *Spreng. Syst.* ii. 390.—Bras.
Nummularia, *Peyr. in Mart. Fl. Bras.* xii. 1. 188.—Bras.
obovatum, *Macfad. Fl. Jamaic.* i. 143.—Jamaic.
obtusifolium, *Hook. f. Fl. Brit. Ind.* i. 415.—Zeylan.
obtusum, *DC. Prod.* i. 574.—Cuba.
ochranthum, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 371.—Bras.
orinocoense, *H. B. & K. Nov. Gen. et Sp.* v. 178.—Venezuela.
ovalifolium, *Peyr. in Mart. Fl. Bras.* xii. 1. 135.—Bras.
ovatum, *Cav. Diss.* 404. t. 238.—Ind. occ.
panamense, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 551.—Panama.
paradisæ, *Peyr. in Mart. Fl. Bras.* xii. 1. 164.—Bras.
parvistipulatum, *Peyr. l. c.* 136.—Bras.
passerinum, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 386.—Bras.
Pelleterianum, *A. St. Hil. Fl. Bras. Mer.* ii. 100. t. 102. —Bras.
peruvianum, *Mitchel. & Pascal. ex Steud. Nom.* ed. II. i. 596.—Peruv.
petiolatum, *Peyr. in Mart. Fl. Bras.* xii. 1. 158.—Bras.
pictum, *E. Mey. in Drège, Zwei Pfl. Docum.* 183, nomen; *Harv. & Sond. Fl. Cap.* i. 284.—Afr. austr.
platyclados, *Boj. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 186.—Madag.
polyneuridos, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 360.—Bras.
popayanense, *H. B. & K. Nov. Gen. et Sp.* v. 179.—N. Granat.
pulchrum, *A. St. Hil. Fl. Bras. Mer.* ii. 94.—Bras.
pyrifolium, *Baker, in Journ. Linn. Soc.* xx. (1883) 109.—Madag.
retusum, Bauer, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 284 = burmanicum.
revolutum, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 351.—Bras.
rigidulum, *DC. Prod.* i. 575.—N. Granat.

ERYTHROXYLUM :—
Roraimæ, Klotzsch, in Schomb. Faun. et Fl. Gui. 1096 = rufum.
rufum, *Cav. Diss.* 404. t. 239.—Ind. occ.
Ruizii, *Peyr. in Mart. Fl. Bras.* xii. 1. 159.—Bras.
Schomburgkii, *Peyr. l. c.* 148.—Bras.
sempervirens, Salzm. ex Peyr. l. c 154 = passerinum.
sideroxyloides, Lam. Encyc. ii. 293 = laurifolium.
sideroxyloides, Retz. Fl. Ind. ii. 449 = monogynum ?
spinescens, *A. Rich. Ess. Fl. Cub.* 255.—Cuba.
Spruceanum, *Peyr. in Mart. Fl. Bras.* xii. 1. 160.—Bras.
squamatum, *Sw. Prod. Veg. Ind. Occ.* 75; *Vahl, Symb. Bot.* iii. 60. t. 63.—Guiana.
strobilaceum, *Peyr. in Mart. Fl. Bras.* xii. 1. 140.—Bras.
subcordatum, *Bert. ex DC. Prod.* i. 575.—Ins. S. Doming.
subcrosum, *A. St. Hil. Pl. Us. Bras.* t. 69.—Bras.
subglaucescens, Mart. ex Peyr. in Mart. Fl. Bras. xii. 1. 146 = virgultosum.
subracemosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 390.—Bras.
subrotundum, *St. Hil. Fl. Bras. Mer.* ii. 99.—Bras.
sumatranum, Miq. Fl. Ind. Bat. Suppl. 512 = burmanicum.
surinamense, *Miq. in Hoev. & De Vriese, Tijdschr.* x. (1843) 86.—Guiana.
tetrastachyum, *Peyr. in Mart. Fl. Bras.* xii. 1. 170.—Bras.
tortuosum, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 399.—Bras.
utile, *Sald. in Ann. Sc. Nat.* Sér. V. xix. (1874) 212. —Bras.
vaccinifolium, *Mart. in Abh. Akad. Muench.* iii. 11. (1840) 387.—Bras.
virgultosum, *Mart. l. c.* 389.—Bras.
Warmingii, *Peyr. in Mart. Fl. Bras.* xii. 1. 140.—Bras.

ERYTROCHILUS, Blume, Bijdr. 614 (1825) = ERYTHROCHILUS, Reinw. = Claoxylon, A. Juss. (Euphorb.).

ERYTROCHITON, [Schlecht. in] Bot. Zeit. iv. (1846) 656 = ERYTHROCHITON, Griff. = Ternstroemia, Linn.

ERYTRONIUM, Scop. Fl. Carn. ed. I. 235 (1760) = Erythronium, Linn. (Liliac.).

ESCALLONIA, Mutis, ex Linn. f. Suppl. 21 (1781). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 644.
STERRHOLYON, Ruiz & Pav. Prod. 88. t. 6 (1794).
(Species 8, vel. Fl. Flum. i. t. 78, 74 (1827).
affinis, Rupr. ex A. Gray, Bot. U. St. Expl. Exped. i. 664 = revoluta.
alpina, *Poepp. & Endl. Nov. Gen. et Sp.* i. 8. t. 13.—Chili.
angustifolia, Phil. in Linnæa, xxxiii. (1864–65) 85 = virgata.
angustifolia, *Presl, Rel. Haenk.* ii. 47.—Peruv.
araucana, *Phil. in Linnæa*, xxvii. (1856) 690.—Chili.
arguta, *Presl, Rel. Haenk.* ii. 48. t. 58.—Chili.
bellidifolia, *Phil. in Anal. Univ. Chil.* (1872) 734.—Chili.
berberifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 296.—Ecuador.
Berteriana, DC. Prod. iv. 665 = pulverulenta.
bicolor, Mart. ex Engl. in Linnæa, xxxvi. (1869–70) 556 = chlorophylla.
bifida, Link & Otto, ex Engl. in Mart. Fl. Bras. xiv. 11. 196 = montevidensis.
Callcottiæ, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 343.—Jun. J. Fernand.
candida, *Lem. Jard. Fleur.* iv. (1854) t. 408. *Misc.* 57. —Bras.
carassana, A. St. Hil. in Guill. Arch. Bot. ii. (1838) 237 = chlorophylla.
coracasana, H. B. & K. Nov. Gen. et Sp. iii. 297 = floribunda.
cormoides, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 344 = sco.
carmelitana, *Meyen, Reise*, i. 313.—Chili.
chlorophylla, *Cham. & Schlecht. in Linnæa*, i. (1826) 542.—Bras.
chonotica, *Phil. in Linnæa*, xxvii. (1856) 690.—Chili.
citidula, Rosm. & Schult. Syst. v. 239 = Miersenyrtus microphylla.

ESCALLONIA :—
Clusseniî, *Miq. in Linnæa*, xix. (1847) 484.—Bras.
commutata, Regel, Gartenfl. (1857) 159 = rubra ?
coccinea, *Phil. in Anal. Univ. Chil.* (1861) t. 61, *et in Linnæa*, xxxiii. (1864–65) 86.—Chili.
cocquimbensis, *Remy, in C. Gay, Fl. Chil.* iii. 61.—Chili.
corymbosa, *Pers. Syn.* i. 284.—Peruv.
crenulata, Roem. & Schult. Syst. v. 239 = Bacckea crenulata.
cuneifolia, Roem. & Schult. l. c. 328 = discolor ?
cupularis, Gill. ex Hook. & Arn. in Bot. Misc. iii. (1833) 343 = illinita.
dense, *Planch. & Linden, Cat.* n. lx. (1854) 3.—N. Granat.
discolor, *Vent. Choix*, 54. t. 54.—N. Granat.
dumetorum, *Phil. in Linnæa*, xxxiii. (1864–65) 86.—Chili.
duplicato-serrata, Remy, in C. Gay, Fl. Chil. iii 58 = macrantha, *Hook. & Arn.*
farinacea, *A. St. Hil. in Guill. Arch. Bot.* ii. (1838) 225.—Bras.
farinosa, Steud. Nom. ed. II. i. 598, sphalm. = pracc.
fernandeziana, Phil. in Bot. Zeit. xiv. (1856) 644 = Callcottiæ.
frescens, *Presl, Rel. Haenk.* ii. 48.—Chili; Peruv.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iii. 297.—N. Granat.
floribunda, Retchb. Hort. Bot. t. 202 = montevidensis.
florida, *Poepp. & Endl. Nov. Gen. et Sp.* i. 8. t. 14.—Chili.
Foæi, *Phil. in Linnæa*, xxvii. (1856) 692.—Chili.
glabrata, *Phil. l. c.* xxxiii. (1864–65) 87.—Chili.
glandulosa, Lodd. Bot. Cab. t. 1291 = rubra.
glandulosa, Sweet, Brit. Flow. Gard. Ser. II. iv. t. 81 = Grahamiana.
Grahamiana, *Gill. ex Hook. & Arn. in Bot. Misc.* iii (1833) 343.—Chili.
grandiflora, Hort. Berol. ex Engl. in Mart. Fl. Bras xiv. 11. 184 = Claussenii.
hirsuta, *Presl, Bot. Bemerk.* 73.—Chili.
hirta, Willd. ex Roem. & Schult. Syst. iii. 329 = discolor.
humilis, A. St. Hil. in Guill. Arch. Bot. ii. (1838) 225 = Claussenii ?
illinita, *Presl, Rel. Haenk.* ii. 49. t. 59.—Chili.
leucantha, *Remy, in C. Gay, Fl. Chil.* iii. 58.—Chili.
littoralis, *Phil. in Linnæa*, xxx. (1859–60) 189.—Chili.
longifolia, *Phil. in Anal. Univ. Chil.* (1872) 734.—Chili.
macrantha, *Hook. & Arn. in Hook. Bot. Misc.* iii (1833) 341.—Ins. Chiloe.
macrantha, *Wedd. Chlor. And.* ii. 312.—Peruv.
megapotamica, Spreng. Syst. iv. Cur. Post. 94 = Sellowiana.
Meriteriana, Rupr. ex A. Gray, Bot. U. St. Expl Exped. i. 664 = illinita.
microcarpa, Gill. ex Hook. & Arn. in Bot. Misc. iii. (1833) 343 = arguta.
millegrana, *Griseb. in Goett. Abh.* xxiv. (1879) 141.—Reg. Argent.
mollis, Phil. in Linnæa, xxvii. (1856) 692 = rubra.
montana, *Phil. l. c.* xxxiii. (1864–65) 87.—Chili.
montevidensis, *DC. Prod.* iv. 4.—Reg. Argent.
multiflora, Presl, Rel. Haenk. ii. 48 = resinosa.
myrtilloides, *Linn. f. Suppl.* 156.—N. Granat.
myrtoides, Bert. ex DC. Prod. iv. 665; Gill. ex Hook. & Arn. in Bot. Misc. iii. (1833) 343 = arguta.
obtusissima, *A. St. Hil. Fl. Bras. Mer.* iii. 91.—Bras.
organensis, *Gardn. in Hook. Ic. Pl.* t. 614.—Bras.
paniculata, *Phil. in Linnæa*, xxviii. (1856) 691.—Chili.
paniculata, Roem. & Schult. Syst. v. 236 = floribunda.
pendula, *Pers. Syn.* i. 285.—Peruv.; N. Granat.
Philippiana, Engl. in Linnæa, xxxvi. (1869–70) 571; Mast. in Gard. Chron. (1873) 947 = virgata.
Poeppigiana, DC. Prod. iv. 3 = rubra.
pollicida, *Hook. Ic. Pl.* t. 114.—Peruv.
pterocladon, *Hook. Bot. Mag.* t. 4827.—Patagon.
pulverulenta, *Pers. Syn.* i. 285.—Chili.
punctata, DC. Prod. iv. 3 = macrantha, *Hook. & Arn.*
resinosa, Gill. ex Hook. & Arn. in Bot. Misc. iii. (1833) 343 = arguta.
resinifluxa, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 343 = pulverulenta.
resinosa, *Pers. Syn.* i. 285.—Peruv.

ESCALLONIA:—
revoluta, Bert. ex Steud. Nom. ed. II. i. 598 = pulverulenta.
revoluta, *Pers. Syn.* i. 285.—Chili.
rosea, *Griseb. in Goett. Abh. vi.* (1856) 121.—Chili.
rosea, Phil. in Linnæa, xxviii. (1856) 692 (= præc. ?).—Chili.
rubra, Bert. ex Steud. Nom. ed. II. i. 598 = illinita.
rubra, *Pers. Syn.* ii. 235.—Chili.
rubricaulis, Colla, in Mem. Acc. Torin. xxxvii. (1834) 79 = Callcottia.
rupestris, *Phil. in Linnæa*, xxviii. (1856) 693.—Chili.
sanguinea, Hort. Veitch. ex Mast. in Gard. Chron. (1872) 947 = macrantha.
Sellowiana, *DC. Prod.* iv. 4.—Bras.
serrata, *Sw. Ic. Ined.* 2. t. 81.—Patagon.
spathulata, Hort. Monac. ex DC. Prod. iv. 4 = montevidensis.
spinulosa, Kunth, ex Engl. in Mart. Fl. Bras. xiv. ii. 140 = pulverulenta.
spiræoides, A. St. Hil. Fl. Bras. Mer. iii. 88 = Sellowiana.
stricta, Kunze, in Linnæa, xx. (1847) 44 = illinita.
stricta, Remy, ex C. Gay, Fl. Chil. iii. 53 = virgata.
thyrsoides, *Bert. in Mercr. Chil.* xiii. 616; *ex Bull. Férussac*, xv. (1830) 112.—Chili.
tortuosa, *H. B. & K. Nov. Gen. et Sp.* iii. 295.—Ecuador.
Tubar, Mutis, ex Humb. Tabl. Nat. i. 113 = discolor.
vaccinoides, *A. St. Hil. Fl. Bras. Mer.* iii. 87.—Bras.
virgata, *Pers. Syn.* i. 284.—Am. austr.
viscosa, *Forbes, Hort. Woburn.* 231, nomen. (*Link. & Otto, ex Steud. Nom.* ed. II. i. 598).—Chili.

ESCHENBACHIA, Moench, Meth. 573 (1794) = Conyza, Linn. (Compos.).
ambigua, Moris, Fl. Sard. ii. 372 = Erigeron linifolius.
globosa, Moench, Meth. 573 = C. aegyptiaca.

ESCHERIA, Regel, in Flora, xxxii. (1849) 179, in syn. = Gloxinia, L'Hérit. (Gesnerac.).
gloxiniæfolia, Regel, l. c. = G. maculata.

ESCHHOLTZIA, Reichb. Consp. 187 (1828) = seq.

ESCHHOLZIA, Cham. in Nees, Hor. Phys. Berol. t. 15 (1830) = seq.

ESCHOLTZIA, Dum. Anal. Fam. 52 (1829) = seq.

ESCHSCHOLTZIA, Bernh. in Linnæa, viii. (1833) 464 = seq.

ESCHSCHOLTZIA, Cham. in Nees, Hor. Phys Berol. 73 (1830). *PAPAVERACEÆ*, Benth. & Hook. f. i. 54.
CHEVREUX, Lindl. Bot. Reg. t. 1677 (1835) et t. 1948 (1839).
ONONOIA, Rafin. Fl. Tellur. ii. 92 (1836).
Aculeata, *Greene, in Bull. Calif. Acad.* i. n. iii. (1885) 89.—Calif.
crespitosa, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 408 = californica.
californica, *Cham. in Nees, Hor. Phys. Berol.* 74.—Calif.
compacta, Steud. Nom. ed. II. i. 598 = californica.
crocea, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 407 = californica.
Douglasii, Benth. Pl. Hartw. 296 = californica.
elegans, *Greene, in Bull. Calif. Acad.* i. n. iv. (1885) 121.—Calif.
fumariæfolia, Heynh. Nom. i. 316 = Hunnemannia fumariæfolia.
glyptosperma, *Greene, in Bull. Calif. Acad.* i. n. iii. (1885) 70.—Calif.
hypecoides, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 408 = californica.
maritima, *Greene, in Bull. Calif. Acad.* i. n. iii. (1885) 89.—N. Mexic.
minutiflora, *S. Wats. in Proc. Am. Acad.* xi. (1876) 123.—Am. bor. occ.
Parishii, *Greene, in Bull. Calif. Acad.* i. n. iv. (1885) 138.—Calif.
peninsularis, *Greene, l. c.* n. iii. (1885) 68.—Calif.
rhombipetala, *Greene, l. c.* 71.—Calif.
rosea, Hort.—Cf. Gard. Chron. (1868) 1336 = californica.
tenuifolia, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 408 = californica.

ESCHSHOLZIA, DC. Prod. iii. 844 = præc.

ESCHWEILERA, Mart. in DC. Prod. iii. 293 (1828) = Lecythis, Linn. (Myrtac.).
acuminata, Miers, in Trans. Linn. Soc. xxx. (1874) 272 = L. ovata.
acuminatissima, Miers, l. c. 272 = L. acuminatissima.
albiflora, Miers, l. c. 268 = L. albiflora.
amara, Mart. ex Berg, in Mart. Fl. Bras. xiv. i. 493 = L. turbinata.
angustifolia, Mart. in Flora, xx. (1837) ii. Beibl. 89 = L. angustifolia.
Blanchetiana, Miers, in Trans. Linn. Soc. xxx. (1874) 256 = L. Blanchetiana.
bracteosa, Miers, l. c. 274 = L. bracteosa.
compressa, Miers, l. c. 248 = L. angustifolia.
cordata, Miers, l. c. 270 = L. cordata.
coriacea, Mart. ex Berg, in Mart. Fl. Bras. xiv. i. 491 = L. coriacea.
corrugata, Miers, in Trans. Linn. Soc. xxx. (1874) 258 = L. corrugata.
elegans, Miers, l. c. 268 = L. elegans.
fendleriana, Miers, l. c. 250.—Venezuela.
flaccifolia, Miers, l. c. 264.—Guiana.
gracilis, Miers, l. c. 273 = L. ovata.
grandiflora, Miers, l. c. 287, sphalm. = seq.
grandifolia, Mart. ex DC. Prod. iii. 293 = L. grandifolia.
guanabarica, Miers, in Trans. Linn. Soc. xxx. (1874) 251.—Bras.
Idatimonoides, Mart. ex Berg, in Mart. Fl. Bras. xiv. i. 498 = L. ovata.
idatimonoides, Miers, in Trans. Linn. Soc. xxx. (1874) 254 = L. idatimonoides.
laevifolia, Miers, l. c. 266 = L. laevifolia.
laurifolia, Miers, l. c. 272 = L. ovata.
longipes, Miers, l. c. 258 = L. longipes.
luridа, Miers, l. c. 262.—Bras.
Luschnathii, Miers, l. c. 258 = L. Luschnathii.
lutea, Miers, l. c. 254 = L. lutea.
macrophylla, Miers, l. c. 263 = L. macrophylla.
micrantha, Miers, l. c. 260 = L. micrantha.
Mortoniana, Miers, l. c. 271.—Venezuela.
nana, Miers, l. c. 261 = L. nana.
obovata, Miers, l. c. 267 = L. obovata.
odora, Miers, l. c. 278 = L. odora.
ovata, Mart. ex Miers, l. c. 257 = L. ovata.
pachyopala, Mart. ex Berg, in Mart. Fl. Bras. xiv. i. 494 = L. pachyopala.
pallida, Miers, in Trans. Linn. Soc. xxx. (1874) 267.—Guian.
paniculata, Miers, l. c. 265 = L. paniculata.
parviflora, Miers, l. c. 260 = L. parviflora. *Aubl.*
parvifolia, Mart. ex DC. Prod. iii. 293 = L. parvifolia.
pilosa, Miers, in Trans. Linn. Soc. xxx. (1874) 274 = L. pilosa.
platycarpa, Miers, l. c. 268 = L. platycarpa.
Riedeliana, Miers, l. c. 269 = L. Riedeliana.
rigida, Miers, l. c. 249.—Venezuela.
Sagotiana, Miers, l. c. 262 = L. parviflora, *Sagot.*
serrulata, Miers, l. c. 250.—Bras.
Sieberiana, Miers, l. c. 241 = L. Sieberiana.
simplex, Miers, l. c. 264.—Guian.
subglandulosa, Miers, l. c. 266 = L. subglandulosa.
tenax, Miers, l. c. 270 = L. tenax.
tenuifolia, Miers, l. c. 268 = L. tenuifolia.
trochiformis, Miers, l. c. 269.—Bras.
verruculosa, Miers, l. c. 255 = L. verruculosa.

ESCLERONA, Rafin. Sylva Tellur. 120 (1838) = Acacia, Willd. (Legumin.).
montana, Rafin. l. c. = A. xylocarpa.

ESCOBEDIA, Ruiz & Pav. Prod. Fl. Per. 91. t. 18 (1794). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 985.
SILVIA, Vell. Fl. Flum. 55 (1825).
laevis, *Cham. & Schlecht. in Linnæa*, v. (1830) 108.—Mexic.
linearis, Schlecht. l. c. viii. (1833) 246 = laevis.
scabrifolia, *Ruiz & Pav. Syst. Veg.* 159.—Am. trop.
Silvia, Steud. Nom. ed. II. i. 598 = scabrifolia.
stricta, Zucc. in Flora, xv. (1832) ii. Beibl. 100 = laevis.

ESCULUS, Linn. Gen. ed. i. 109 (1737) = Aesculus, Linn. (Sapindac.).

ESDRA, Salisb. Gen. Pl. Fragm. 60 (1866) = Trillium, Linn. (Liliac.).
sessilis, Salisb. l. c. = T. sessile.

ESEMBECKIA, Barb.-Rodr. in Rev. Engenh. v. (28 Jul. 1882) = Esenbeckia, H. B. & K. (Rutac.).

ESENBECKIA, Blume, Bijdr. 118 (1825) = Neesia, Blume (Malvac.).
altissima, Blume, l. c. 119 = N. altissima.

ESENBECKIA, H. B. & K. Nov. Gen. et Sp. vii. 246. t. 655 (1825). *RUTACEÆ*, Benth. & Hook. f. i. 299.
COLYTHRUM, Schott, Rutac. 13. t. 7 (1834).
KUALA, Karst. & Triana, in Linnæa, xxviii. (1856) 428.
METRODOREA, A. St. Hil. Fl. Bras. Mer. ii. t. 16 (1825).
POLEMBRYUM, A. Juss. in Mém. Mus. Par. xii. (1825) 519 (Mém. Rut. 126).
alata, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 306.—N. Granat.
attenuata, *Griseb. Fl. Brit. W. Ind.* 136.—Ins. Trinit.
Berlandieri, *Baill. Adansonia*, x. (1871) 151.—Mexic.
castanocarpa, Griseb. Fl. Brit. W. Ind. 135 = pilocarpoides.
choisyoides, *Schlecht. in Linnæa*, xxvii. (1854) 529.—Hab.?
cornuta, *Engl. in Mart. Fl. Bras.* xii ii. 146.—Peruv fasciculata, *Barb.-Rodr. in Rev. Engenh.* v. (28 Jul. 1882).—Bras.
febrifuga, *A Juss. in Mém. Mus. Par.* xii. (1825) 486, in obs.—Bras.
grandiflora, *Mart. Nov. Gen. et Sp.* iii. 85.—Bras.
intermedia, *Mart. Syst. Mat. Med. Bras.* 39.—Bras.; Chili.
latifolia, Mart. Nov. Gen. et Sp iii. 84 = pumila.
leiocarpa, *Engl. in Mart. Fl. Bras.* xii. ii. 145.—Bras.; Chili.
leucophylla, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 440 = pumila.
maurozoides, Mart. Nov. Gen. et Sp. iii. 82 = pilocarpoides.
mollis, *Miq. in Linnæa*, xxiii. (1849) 706.—Bras.
pentaphylla, *Griseb. Fl. Brit W. Ind.* 135.—Jamaica.
pilocarpoides, *H. B. & K. Nov. Gen. et Sp.* vii. 248.—Am. trop.
pumila, *Pohl, Pl. Bras.* ii. 44.—Bras.
Riedeliana, *Engl. in Mart. Fl. Bras.* xii. ii. 142.—Bras.
Warmeciwii, *Engl. l. c.* 148.—Peruv.

ESERA, Neck. Elem. ii. 160 (1790) = Drosera, Linn.

ESMARCHIA, Reichb. Fl. Germ. Excurs. 793 (1832) = Cerastium, Dill. (Caryophyl.).
cerastoides, Reichb. l. c. = C. tetrandrum.

ESMERALDA, Reichb. f. Xenia Orch. ii. 38 (1874) = Arachnanthe, Blume (Orchid.).
Catacarti, Reichb. f. l. c. 39 = A. Catacarti.
Sanderiana, Reichb. f. in Gard. Chron. (1882) i. 588 = Vanda Sanderiana.

ESMERALDIA, Fourn. in Ann. Sc. Nat. Sér. VI. xiv. (1882) 367. *ASCLEPIADEÆ*.
stricta, *Fourn. l. c.* 368.—Venezuela.

ESOPON, Rafin. Fl. Ludov. 149 (1817) = Pronanthes, Linn. (Compos.).
glaucum, Rafin. l. c. = *P. glauca* (nomen delendum).—Am. bor.

ESPADANA, A. Rich. Fl. Cub. Fanerog. ii. 147 (1858). *VERBENACEÆ*, Benth. & Hook. f. ii. 1148.
ARMENIASTRUM, Lem. Jard. Fleur. iv. (1854) Misc. 76.
amœna, *A. Rich. l. c.* 148.—Cuba.
apiculata, *Miers, in Trans. Linn. Soc.* xxvii. (1870) 194.—Cuba.

ESPEJOA, DC. Prod. v. 660 (1836) = Jaumea, Pers. (Compos.).
mexicana, DC. l. c. = J. mexicana.

ESPELETIA, Mutis, ex Humb. & Bompl. Pl. Aequin. ii. 10. t. 70-72 (1800). *COMPOSITÆ*, Benth. & Hook. f. ii. 347.
LIBANOTHAMNUS, Ernst, in Vazgesia (1870), 186.
argentea *Humb. & Bompl. l. c.* 14. t. 71.—N. Granat.

ESPELETIA :—
banksiaefolia, *Sch. Bip. ex Wedd. Chlor. And.* i. 67.—Venezuela.
corymbosa, *Humb. & Bonpl. Pl. Aequin.* ii. 16.—N. Granat.
corymbosa, Sch. Bip. ex Wedd. Chlor. And. i. 63 = Schultzii.
Funckii, *Sch. Bip. ex Wedd. l. c.* 64.—N. Granat.
grandiflora, *Humb. & Bonpl. Pl. Aequin.* ii. 11. t. 70.—N. Granat.
Hartwegiana, Sch. Bip. ex Wedd. Chlor. And. i. 62 = grandiflora.
Lindenii, *Sch. Bip. ex Wedd. l. c.* 67.—Venezuela.
Moritziana, *Sch. Bip. ex Wedd. l. c.* 65.—Venezuela.
nerifolia, *Sch. Bip. ex Wedd. l. c.* 67.—Venezuela.
nivea, Moritz. ex Wedd. i. c. 65 = argentea.
oppositifolia, Sch. Bip. ex Wedd. l. c. 62 = grandiflora, Moritziana.
platylepis, Sch. Bip. ex Wedd. l. c. 64 = corymbosa.
rigida, Humb. & Bonpl. Pl. Aequin. t. 73 = corymbosa. Schultzii, *Wedd Chlor. And.* i. 63.—Venezuela.
spicata, *Sch. Bip. ex Wedd. l. c.* 65.—Venezuela.
Weddelii, *Sch. Bip. ex Wedd. l. c.* 66.—Venezuela.

ESPELETIA, Nutt. in Journ. Acad. Philad. vii. (1834) 87. t. 4 = **Balsamorhiza**, Hook. (Compos.).
ampiexicaulis, Nutt. l. c. 88 = Wyethia amplexicaulis.
helianthoides, Nutt. l. c. 88 = B. sagittata.
sagittata, Nutt. l. c. 89 = B. sagittata.

ESPELETIOPSIS, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 414 (1873) = **Helenium**, Linn. (Compos.).

ESPERA, Willd. in Ges. Naturf. Freunde Neue Schr. iii. (1801) 449 = **Berria**, Roxb. (Tiliac.).
cordifolia, Willd. l. c. 450 = B. Ammomilla.

ESPINOSA, Lag. Gen. et Sp. Nov. 14 (1816) = **Eriogonum**, Michx. (Polygon.).
verticillata, Lag. l. c. = Eriog. tomentosum.

ESSENHARDTIA, Sweet, Hort. Brit. ed. III. 704 (1839) = **Hysenhardtia**, H. B. & K. (Legumin.).

ESTERHAZIA, Bartl. Ord. 186 (1830) = seq.

ESTERHAZYA, Mikan. Del. Faun. et Fl. Bras. t. 5 (1820). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 972.
LAGIOSTEMON, Schott, ex Endl. Gen. 690, in syn. (1839).
alpestris, Spix, ex Mart. Reise Bras. i. 397 = Gerardia brachyphylla.
campestris, Spix, ex Mart. l. c. = splendida.
macrodonta, *Cham. & Schlecht. in Linnaea*, viii. (1833) 93.—Bras.
montana, Spix, ex Mart. Reise Bras. i. 397 = splendida.
nervosa, *Benth. in DC. Prod.* x. 514.—Bras.
splendida, *Mikan. Del. Faun. et Fl. Bras.* t. 5.—Bras.

ESULA, Rupp. Fl. Jen. ed. Hall. 271 (1745); Haw. Syn. Pl. Succ. 168 (1812) = **Euphorbia**, Linn.
alopica, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 149 = E. aleppica.
amygdaloides, Haw. Syn. Pl. Succ. 164 = E. amygdaloides.
angustifolia, Haw. l. c. 166 = E. Esula.
arenica, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. Esula.
biumbellata, Haw. Syn. Pl. Succ. 156 = E. biumbellata.
Characias, Haw. l. c. 163 = E. Characias.
cupressina, S. F. Gray, Nat. Arr. Brit. Pl. ii. 389 = E. Cyparissias.
Cyparissias, Haw. Syn. Pl. Succ. 155 = E. Cyparissias.
Dalechampii, Haw. l. c. = E. Esula.
dendroides, Haw. l. c. 158 = E. dendroides.
diffusa, Haw. l. c. 158 = E. exigua.
exigua, Haw. l. c. = E. exigua.
falcata, Haw. l. c. 159 = E. falcata.
Gerardiana, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. Gerardiana.
juncoides, Haw. Syn. Pl. Succ. 157 = E. aleppica.
minima, Haw. l. c. 158 = E. Peplus.
multicornymboa, Haw. l. c. 156 = E. multicornymboa.
nicaeensis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. nicaeensis.
obscura, Fourr. l. c. = E. falcata.
Parolias, Fourr. l. c. = E. Parolias.
peploides, Fourr. l. c. = E. peploides.
ESULA :—
Peplus, Haw. Syn. Pl. Succ. 158 = Euphorbia Peplus.
pinea, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. pinea.
Pithyusa, Fourr. l. c. = E. Pithyusa.
portlandica, Haw. Syn. Pl. Succ. 154 = E. portlandica.
provincialis, Haw. l. c. 167 = E. Terracina.
Pseudo-cyparissias, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. Esula.
retusa, Fourr. l. c. 149 = E. exigua ?
riparia, Fourr. l. c. 150 = E. Esula.
rotundifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 387 = E. Peplus.
salicifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. androsaemifolia.
segetalis, Haw. Syn. Pl. Succ. 156 = E. segetalis.
sulcata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. sulcata.
sylvatica, Haw. Syn. Pl. Succ. 153 = E. sylvatica.
taurinensis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. taurinensis.
tenuifolia, Fourr. l. c. = E. tenuifolia.
Terracina, Fourr. l. c. = E. Terracina.
tricuspidata, Fourr. l. c. 149 = E. exigua.
valentina, Haw. Syn. Pl. Succ. 157 = E. Terracina.
virgata, Haw. l. c. 155 = E. virgata.
vulgaris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = E. Esula.

ETABALLIA, Benth. in Hook. Journ. Bot. ii. (1840) 99 = **Inocarpus**, Forst. (Legumin.).
guianensis, Benth. l. c.—Guian.
macrophylla, Benth. l. c. = I. edulis.

ETAERIA, Blume, Bijdr. 409. t. 14 (1825) = **Hetaeria**, Blume (Orchid.).

ETHERISCIUS, Desv. in Ham. Prod. Pl. Ind. Occ. 28 (1825). *RUBIACEAE* ?
parasiticus, *Desv. l. c.*—Guiana.

ETERISCUS, Steud. Nom. ed. II. i. 599 (1840) = prace.

ETHANIUM, Salisb. in Trans. Hort. Soc. i. (1812) 281 = **Renealmia**, Linn. f. (Scitamin.).
racemosum, Salisb. l. c. = R. racemosa.

ETHEIRANTHUS, Kostel. Ind. Hort. Prag. 56 (1844) = **Muscari**, Tourn. (Liliac.).
constrictus, Kostel. l. c. = M. constrictum.
Yacquini, Kostel. l. c. = M. racemosum ?
maritimus, Kostel. l. c. = M. maritimum.
Millori, Kostel. l. c. = M. comosum ?

ETHEOSANTHES, Rafin. Neogenyt. 3 (1825) = **Tradescantia**, Linn. (Commelin.).
ciliata, Rafin. l. c. = T. ciliata.

ETHESIA, Rafin. Fl. Tellur. iii. 60 (1836) = **Ornithogalum**, Tourn. (Liliac.).
fragrans, Rafin. l. c. = O. praecinum.

ETHESIA, Rafin. Fl. Tellur. iv. 63 (1836) = **Jacobinia**, Mart. (Acanthac.).
carnea, Rafin. l. c. 64 = J. Pohliana.

ETHIONEMA, Brongn. Enum. Gén. 92 (1843) = **Aethionema**, R. Br. (Crucif.).

ETHULIA, Linn. Sp. Pl. ed. II. 1171 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 234.
KAMISIA. Forsk. Fl. Aegypt. Arab. 158 (1775).
LEUCHIA, Scop. Introd. 198 (1777).
PTERADIA, Adans. Fam. ii. 109 (1763).
ageratoides, Spreng. Syst. iii. 458 = Alomia ageratoides.
alata, Sond. in Linnaea, xxiii. (1850) 60.—Afr. austr.
angustifolia, Boj. ex DC. Prod. v. 13 = conyzoides.
aquatica, Roxb. ex DC. l. c. 116 = Adenostemma viscosum.
auriculata, Thunb. Prod. Pl. Cap. 141 = Dichrocephala latifolia.
bechioides, *Klatt, in Ann. Sc. Nat. Sér.* V. xvii. (1873) 361.—Ins. Comor.
Bidentis, Linn. Mant. i. 110 = Flaveria Contrayerba.
brasiliensis, *Link, Enum. Hort. Berol.* ii. 210.—Bras.
conyzoides, Boj. ex DC. Prod. v. 69 = Vernonia madagascariensis.
corymbosa, Herb. Madr. ex Wall. Cat. n. 3089 = Laggera alata.
ETHULIA :—
conyzoides, *Linn. Sp. Pl.* ed. II. 1171.—As. et Afr. trop.
Cunninghamii, Hook. in Mitch. Journ. Trop. Austral. 63 = Epaltes Cunninghamii.
divaricata, Linn. Mant. i. 110 = Epaltes divaricata.
divaricata, Wall. Cat. n. 3046 = Brachyscome saxatilis.
divergens, Rusa. ex Wall. l. c. 3089 = Laggera alata.
gariepina, DC. Prod. v. 15 = Epaltes gariepina.
glomerata, Spreng. ex DC. Prod. v. 372 = Dichrocephala latifolia.
integrifolia, D. Don, Prod. Fl. Nep. 182 = Dichrocephala latifolia.
involucrata, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 3 = Otospermum involucratum.
Krausii, *Sch. Bip. ex Walp. Rep.* ii. 946.—Afr. austr.
megacephala, Sch. Bip. ex Miq. Fl. Ind. Bat. ii. 8 = conyzoides.
meyogalitana, Sch. Bip. ex De Vriese, Pl. Ind. Bat. Reinw. 128 = conyzoides.
nepalensis, Spreng. ex DC. Prod. vi. 141 = Cotula hemisphaerica.
paniculata, Houtt. Handleid. x. 551 = Hippia integrifolia.
paniculata, Schkuhr, Handb. iii. 67. t. 285 = Dichrocephala latifolia.
parviflora, *Klatt, in Linnaea*, xxxvii. (1871–73) 507.—Madag.
ramosa, Roxb. Hort. Beng. 61; Fl. Ind. iii. 413 = conyzoides.
repanda, Nees, ex DC. Prod. v. 663 = Cephalaphora glauca.
Rueppellii, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 372 = Gutenbergia Rueppellii.
sinapifolia, Steud. Nom. ed II. i. 599 = Dichrocephala latifolia.
spergunophora, Linn. Sp. Pl. ed. II. 1171 = Spergunophorus Vaillantii.
Strachiana, Sw. Prod. Veg. Ind. Occ. 111 = Spergunophorus Vaillantii.
tomentosa, *Linn. Mant.* i. 110.—China.
uniflora, Walt. Fl. Carol. 195 = Sclerolepis verticillata.

ETHULIOPSIS, F. Muell. Fragm. ii. 154 (1861) = Epaltes, Cass. (Compos.).
discoa, F. Muell. l. c. 155 = Epaltes Cunninghamii.

ETHUSA, Linn. Syst. ed. I (1735); Gen. ed. I. 78 (1737) = **Aethusa**, Linn. (Umbellif.).

ETLINGERA, Giseke, Prael. Linn. 199, 209. (1792) = Amomum, Linn. (Scitamin.).
litteralis, Giseke, l. c. = A. littorale.

ETORLOBA, Rafin. Sylva Tellur. 79 (1838) = Jaoranda, Juss. (Bignoniac.).
caerulea, Rafin. l. c. = J. caroliniana.

ETTLINGERA, Raesach. Nom. ed. III. 1 (1797) = ETLINGERA, Giseke.

ETUBILA, Rafin. Sylva Tellur. 125 (1838) = **Loranthus**, Linn.
brasiliensis, Rafin. l. c. = L. brasiliensis.
dichotoma, Rafin. l. c. = L. dichotomus.
ferruginea, Rafin. l. c. 159 = L. ferrugineus.
longiflora, Rafin. l. c. 135 = L. longiflorus.
maculata, Rafin. l. c. 159 = L. maculatus.

ETUSA, Roy. ex Steud. Nom. ed. II. i. 599 (1840) = **Aethusa**, Linn. (Umbellif.).

EUACER, Opiz, Seznam, 42 (1852) = **Acer**, Tourn. (Sapind.).
acutifolium, Opiz, l. c. = A. platanoides ?
affine, *austriacum*, *campestre*, *arboreum*, *erythrocarpon*, *hebbithornum*, *leiocarpon*, *macrocarpon*, *microcarpon*, *molle*, *obtusilobum*, *orthocarpon*, *pallens*, *palmatifolium*, *polycarpon*, *pubigerum*, *rubescens*, *stenolepis*, *subglaucophyllum*, Opiz, Seznam, 42 = A. campestre.
barbatum, Opiz, l. c. = A. barbinerve.
lanuginosum, Opiz, l. c. = A. obtusatum.
platanoides, Opiz, l. c. = A. platanoides.
rubrotinctum, Opiz, in Lotos (1856) 68 = A. campestre ?
scharfianum, Opiz, l. c. = A. campestre ?
tataricum, Opiz, l. c. = A. tataricum.

EUADENIA, Oliver, in Benth. & Hook. f. Gen. i. 969 (1867). *CAPPARIDEAE*, Benth. & Hook. f. l. c. eminens, *Hook. f. Bot. Mag.* t. 6578.—Afr. trop. Kirkii, *Oliver, Pl. Trop. Afr.* i. 91.—Afr. trop. trifoliata, *Benth. & Hook. f. Gen.* i. 969.—Afr. trop. *trifoliolata*, Oliver, *Pl. Trop. Afr.* i. 91 = trifoliata.

EUALCIDA, Hemsl. Biol. Centr. Am. Bot. ii. 222 (1881) sphalm. = ENALCIDA, Cass. = **Tagetes**, Linn. (Compos.).

EUARTHROCARPUS, Endl. Ench. 451 (1841) sphalm. = **Enarthrocarpus**, Labill. (Crucif.).

EUARTHRONIA, Nutt. ex A. Gray, in Proc. Am. Acad. iv. (1860) 49 in syn. = **Coprosma**, Forst. (Rubiac.). *foliosa*, Nutt. ex A. Gray, l. c. = C. foliosa.

EUBASIS, Salisb. Prod. 68 (1796) = **Aucuba**, Thunb. (Cornac.). *dichotoma*, Salisb. l. c. = A. japonica.

EUBOTRYS, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1841) 369 = **Leucothoë**, D. Don (Ericac.). *bracteata*, Nutt. l. c.—Florida. *racemosa*, Nutt. l. c. = L. racemosa.

EUBRACHION, Hook. f. Fl. Antarct. 291 (1846). *LORANTHACEAE*, Benth. & Hook. f. iii. 216. Arnottii, *Hook. f. i. c.*—Uruguay. brasiliense, *Eichl. in Mart. Fl. Bras.* v. ii. 133.—Bras.

EUCALIA, Raeusch. Nom. ed. III. 385, 251 (1797) = **Encelia** † Adans. (Compos.).

EUCALLIAS, Raña. Fl. Tellur. iv. 25 (1836) = **Billbergia**, Thunb. (Bromeliac.). *versicolor*, Raña. l. c. = B. sebrina.

EUCALYPTON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = **Eucalyptus**, L'Hérit. (Myrtac.).

EUCALYPTUS, L'Hérit. Sert. Angl. 18 (1788) *MYRTACEAE*, Benth. & Hook. f. i. 707.
AROMADENDRON, Andr. ex Steud. Nom. ed. II. i. 184 (1840).
AROMADENDRUM, Andr. ex Steud. l. c. 599 in syn. (1840).
EUDESMIA, R. Br. in Flind. Voy. ii. 599. t. 8 (1814).
SYMPHYOMYRTUS, Schau. in Lehm. Pl. Preiss. i. 126 (1844).
Abergiana, *F. Muell. Fragm.* xi. 41.—Austral.
acervula, Hook. f. Fl. Tasm. i. 135 = Stuartiana.
acervula, Miq. in Nederl. Kruidk. Arch. iv. (1859) 137 = macrorhyncha.
acervula, Sieber, ex DC. Prod. iii. 217 = piperita.
acmenioides, *Schau. in Walp. Rep.* ii. 924.—Austral.
acuminata, Hook. in Mitch. Journ. Trop. Austral. 390 = rostrata.
acutangula, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 338.—Austral.
alba, *Reinw. ex Blume, Bijdr.* 1201.—Ins. Timor. ; Austral.
albens, *Miq. in Nederl. Kruidk. Arch.* iv. (1859) 138. —Austral.
albicans, *F. Muell. Fragm.* vii. 42.—Austral.
alpina, *Lindl. in Mitch. Three Exped.* ii. 175.— Austral.
amblyon, DC. Prod. iii. 219 = amygdalina.
amygdalina, *Labill. Nov. Holl. Pl.* ii. 14. t. 154.— Austral.
amygdalina, Schau. in Lehm. Pl. Preiss. i. 130 = loxophleba.
androsamoides, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 113.— Austral.
angulosa, Schau. in Walp. Rep. ii. 926 = incrassata.
angustifolia, *Deaf. ex Link, Enum. Hort. Berol.* ii. 80. —Austral.
angustifolia, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 337.—Austral.
angustissima, *F. Muell. Fragm.* iv. 25.—Austral.
annulata, *Benth. Fl. Austral.* iii. 224.—Austral.
aspera, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 93. —Austral.
aureusiace, F. Muell. l. c. 91 = miniata.
Baileyana, *F. Muell. Fragm.* xi. 37.—Austral.
Banksiana, Miq. in Nederl. Kruidk. Arch. iv. (1859) 137 = Stuartiana.

EUCALYPTUS:—
Baueriana, Schau. in Walp. Rep. ii. 924 = polyanthemos.
Bauteri, R. Br. ex Benth. Fl. Austral. iii. 207 = santalifolia.
Behriana, *F. Muell. in Trans. Vict. Inst.* i. (1855) 84. —Austral.
bicolor, *A. Cunn. ex Hook. in Mitch. Journ. Trop. Austral.* 390.—Austral.
bigalerita, F. Muell. in Journ. Linn. Soc. iii. (1859) 96 = platyphylla.
bistrigata, *Sm. in Trans. Linn. Soc.* iii. (1797) 286.— Austral.
Bowmani, *F. Muell. ex Benth. Fl. Austral.* iii. 219.— Austral.
brachyandra, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 97.—Austral.
brachypoda, Benth. Fl. Austral. iii. 228, partim = microthecа.
brachyphylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 21.—Austral.
brevifolia, F. Muell. in Journ. Linn. Soc. iii. (1859) 84 = microthecа.
bupreutium, *F. Muell. Fragm.* iii. 57.—Austral.
caesia, *Benth. Fl. Austral.* iii. 227.—Austral.
Cayuputea, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 126 = odorata.
calophylla, *R. Br. in Journ. Geog. Soc.* i. 1831 (1832) 20, nomen; *Lindl. Bot. Reg.* (1841) *Misc.* 72.— Austral.
calycogonа, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 338.—Austral.
camaldulensis, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 20.—Austral.
capitellata, *Sm. Bot. N. Holl.* 42.—Austral.
celastroides, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 338.—Austral.
cinerea, *F. Muell. ex Benth. Fl. Austral.* iii. 229.— Austral.
citriodora, *Hook. in Mitch. Journ. Trop. Austral.* 395. —Austral.
cladocalyx, F. Muell in Linnaea, xxv. (1852) 388 = corynocalyx.
clavigera, *A. Cunn. ex Schau. in Walp. Rep.* ii. 926. —Austral.
Cloeziana, *F. Muell. Fragm.* xi. 44.—Austral.
cneorifolia, *DC. Prod.* iii. 220.—Austral.
cocciferа, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 477.—Tasman.
colossea, F. Muell Fragm. vii. 42 = diversicolor.
comitifolia, *Schau. in Lehm. Pl. Preiss.* i. 129.— Austral.
confertiflora, Kipp. ex F. Muell. in Journ. Linn. Soc. iii. (1859) 96 = ferruginea.
connata, *Dum.-Cours. Bot. Cult.* ed. II. vii. 280.— Austral.
conoidea, *Benth. Fl. Austral.* iii. 227.—Austral.
Cooperiana, *F. Muell. Fragm.* xi. 88.—Austral.
cordata, *Labill. Nov. Holl. Pl.* ii. 13. t. 152.—Tasmania.
cordata, Lodd Bot. Cab. t. 838 = pulverulenta.
coriacea, A. Cunn. ex Schau. in Walp. Rep. ii. 925 = gunnifolia.
cornuta, *Labill. Voy.* i. 408. t. 90 ; *Nov. Holl. Pl.* ii. 121.—Austral.
corynocalyx, *Sm. Bot. N. Holl.* 43 ; *et in Trans. Linn. Soc.* iii. (1797) 287.—Austral.
corynocalyx, *F. Muell. Fragm.* ii. 48.—Austral.
cosmophylla, *F. Muell. in Trans. Vict. Inst.* (1855) 83.—Austral.
costata, F. Muell. l. c. 88 = incrassata.
crebra, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 87.— Austral.
Cunninghami, Sweet, Hort. Brit. ed. II. 209 = stricta.
curvula, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 195.— Austral.
cuspidata, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 21 = incrassata.
daphnoides, Miq. in Nederl. Kruidk. Arch. iv. (1859) 138 = occidens.
dealbata, *A. Cunn. ex Schau. in Walp. Rep.* ii. 924.— Austral.
Decmeniana, *Blume, Mus. Bot. Lugd. Bat.* i. 83.— Ins. Timor.
deglupta, *Endl. in Enum. Pl. Hueg.* 49.—Austral.
descrta, *F. Muell. Fragm.* iii. 130.—Austral.
dealupta, *Blume, Mus. Bot. Lugd. Bat.* i. 82.—Ins. Celebes ; Timor.

EUCALYPTUS:—
dichromophloia, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 89.—Austral.
discolor, *Desf. Tabl.* ed. II. 198, nomen ; *Cat. Hort. Par.* ed. III. 408.—Austral.
diversicolor, *F. Muell. Fragm.* iii 131.—Austral.
diversifolia, Bonpl. Desc. Pl. Jard. Malm. 85. t. 18 = viminalis.
dives, *Schau. in Walp. Rep.* ii. 926.—Austral.
dorsiexylon, *F. Muell. Fragm.* ii. 55.—Austral.
drepanophylla, *F. Muell. ex Benth. Fl. Austral.* iii. 223.—Austral.
Drummondii, *Benth. l. c.* 227.—Austral.
dumosa, *A. Cunn. ex Schau. in Walp. Rep.* ii. 925.— Austral.
elaeophora, F. Muell. Fragm. iv. 82 = goniocalyx.
elata, Dehnh. Cat. Pl. Hort. Camald. 26 = viminalis.
elongata, *Link, Enum. Hort. Berol.* ii. 80.—Austral.
enervifolia, Walp. Rep. ii. 164, sphalm. = cneorifolia.
erythrocalyx, F. Muell. Fragm. ii. 32 = pyrniformis.
erythrocorys, *F. Muell. l. c.* 33.—Austral.
erythronema, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 337.—Austral.
endesmoides, *F. Muell. Fragm.* ii. 35.—Austral.
eugenioides, *Sieber, ex Spreng. Syst. iv. Cur. Post.* 195. —Austral.
eximia, *Schau. in Walp. Rep.* ii. 926.—Austral.
exserta, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 85. —Austral.
fabrorum, Schlecht. in Linnaea, xx (1847) 687 = viminalis.
falcata, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i. 168.—Austral.
falcifolia, Miq. in Nederl. Kruidk. Arch. iv. (1858) 126 = oblique.
fasciculosa, F. Muell. in Trans. Vict. Inst. i. (1855) 84 = paniculata.
ferruginea, *Schau. in Walp. Rep.* ii. 926.—Austral.
fibrosa, F. Muell. in Journ. Linn. Soc. iii. (1859) 87 = siderophloia.
fissilis, *F. Muell. Fragm.* ii. 55.—Austral.
flexilis, *Reppl, Gartenfl.* (1856) 264.—Australia.
floribunda, Hueg. in Enum. Pl. Hueg. 42 = marginata.
floribunda, F. Muell. in Journ. Linn. Soc. iii. (1859) 96 = ferruginea.
foecunda, *Schau. in Lehm. Pl. Preiss.* i. 130.—Austral.
Foelschcana, *F. Muell. in Melbourne Chem. & Drugg.* (Nov. 1882).—Austral.
fruticetorum, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1858) 131 = gracilis.
gamophylla, *F. Muell. Fragm.* xi. 40.—Austral.
gigantea, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 20.— Austral.
gigantea, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 479 = oblique.
glandulosa, *Dorf. Cat. Hort. Par.* ed. III. 408.—Austral.
glauca, *DC. Prod.* iii. 221.—Austral.
glaucophylla, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 113.— Austral.
globularis, Hort. ex DC. Prod. iii. 219 = amygdalina.
globulosus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = sec.
Globulus, *Labill. Voy.* i. 158. t. 18 ; *Nov. Holl. Pl.* ii. 121.—Australia.
gunnifolia, G. Don, Gen. Syst. ii. 920, sphalm. = cneorifolia.
gomphocephala, *DC. Prod.* iii. 220.—Austral.
gonisantha, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i. 168.—Austral.
goniocalyx, *F. Muell. Fragm.* ii. 48.—Austral.
gracilis, *F. Muell. in Trans. Vict. Inst.* i. (1855) 85. —Austral.
gracilis, Sieber, ex Benth. Fl. Austral. iii. 222 = crebra.
grandifolia, *R. Br. ex Benth. l. c.* 230.—Austral.
granulosa, Sieber, ex Benth. l. c. 240 = viminalis.
grossa, *F. Muell. ex Benth. l. c.* 228.—Austral.
Gunnii, *Hook. f. in Hook. Lond. Journ. Bot.* iii. (1844) 499.—Australia.
Gunnii, Miq. in Nederl. Kruidk. Arch. iv. (1858) 126 = viminalis.
Gunnii, F. Muell. Fragm. ii. 62 = Stuartiana.
haemastoma, Miq. in Nederl. Kruidk. Arch. iv. (1859) 130 = bicolor.
haemastoma, *Sm. in Trans. Linn. Soc.* iii. (1797) 286. —Austral.
hemilampra, F. Muell. in Journ. Linn. Soc. iii. (1859) 85 = rainiflora.
hemiphloia, *F. Muell. Fragm.* ii. 62.—Australia.

EUCALYPTUS :—
heterophylla, Miq. in Nederl. Kruidk. Arch. iv. (1859) 141 = obliqua.
hirsuta, Link, Enum. Hort. Berol. ii. 31 = Angophora cordifolia.
Hookeri, F. Muell. in Journ. Linn. Soc. iii. (1859) 90 = tesselaris.
Howittiana, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 171.—Austral.
hypericifolia, R. Br. ex Benth. Fl. Austral. iii. 208 = amygdalina.
hypericifolia, *Dum.-Cours. Bot. Cult.* ed. II. vii. 279. —Austral.
hypoleuca, Schau. in Lehm. Pl. Preiss. i. 181 = marginata.
incrassata, *Labill. Nov. Holl. Pl.* ii. 12. t. 150.—Austral.
incrassata, Sieber, in DC. Prod. iii. 217 = pilularis.
inamprocarpa, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 129 = oblonga
largiflorens, F. Muell. in Trans. Vict. Inst. i. (1855) 84 = bicolor.
latifolia, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 95. —Austral.
Lehmanni, *Preiss, ex Schau. in Lehm. Pl. Preiss.* i. 127.—Austral.
leptophleba, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 86.—Austral.
leptophylla, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 123 = uncinata.
leptopoda, *Benth. Fl. Austral.* iii. 238.—Austral.
Leucadendron, *Rrinw. ex De Vriese, Pl. Ind. Bat. Rrinw.* 68.—Ins. Timor.
leucoxylon, *F. Muell. in Trans. Vict. Inst.* i. (1855) 33.—Austral.
ligustrina, DC. Prod. iii. 219 = amygdalina.
ligustrina, Miq. in Nederl. Kruidk. Arch. iv. (1859) 134 = Gunnii.
Lindleyana, DC. Prod. iii. 219 = amygdalina.
linearis, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 20.— Austral.
longifolia, Lindl. in Bot. Reg. t. 947 = amygdalina.
longifolia, *Link, Enum. Hort. Berol.* ii. 29.—Austral.
longirostra, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 125 = rostrata.
lozophleba, *Benth. Fl. Austral.* iii. 252.—Austral.
Luehmanniana, *F. Muell. Fragm.* xi. 38.—Austral.
macrandra, *F. Muell. ex Benth. Fl. Austral.* iii. 235.— Austral.
macrocalyx, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 339 = pyriformis.
macrocarpa, *Hook. Ic. Pl.* t. 405–407.—Austral.
macrorrhyncha, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) 11. 20 = Lehmanni.
macrorhyncha, *F. Muell. First Gen. Rep.* 12; *Benth. Fl. Austral.* iii. 207.—Austral.
maculata, *Hook. Ic. Pl.* t. 619.—Austral.
Mahagoni, F. Muell. Fragm. ii. 41 = marginata.
mannifera, A. Cunn. ex Benth. Fl. Austral. iii. 240 = viminalis.
mannifera, *Mudie, in Trans. Med. Bot. Soc.* 1832–33 (1834), 24.—Austral.
marginata, *Sm. in Trans. Linn. Soc.* vi. (1802) 302.— Austral.
media, *Link, Enum. Hort. Berol.* ii. 30.—Austral.
megacarpa, *F. Muell. Fragm.* ii. 70.—Austral.
melanophloia, *F. Muell. in Journ. Linn. Soc.* iii. (1858) 93.—Austral.
melliodora, *Lindl. in Mitch. Journ. Trop. Austral.* 238.—Austral.
metasideros, F. Muell. in Journ. Linn. Soc. iii. (1859) 86 = peltata.
micrandra, *A. Cunn. ex Schau. in Walp. Rep.* ii. 924. —Austral.
microantha, DC. Prod. iii. 217 = haemastoma.
microcarpa, *F. Muell. ex Benth. Fl. Austral.* iii. 216. —Austral.
microcorys, *F. Muell. Fragm.* ii. 50.—Austral.
microphylla, A. Cunn. in Field, N. S. Wales, 350 = stellulata, sticta.
microphylla, *Willd. Enum. Hort. Berol.* 515.—Austral.
microtheca, *F. Muell. in Journ. Linn. Soc.* iii. (1858) 97.—Austral.
minista, *A. Cunn. ex Schau. in Walp. Rep.* ii. 925.— Austral.
melucana, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 498.— Ins. Moluce.
mucronata, *Link, Enum. Hort. Berol.* ii. 30.—Austral.

EUCALYPTUS :—
Muelleri, Miq. in Nederl. Kruidk. Arch. iv. (1859) 130 = incrassata.
multiflora, Poir. Encyc. Suppl. ii. 594 = resinifera !
multiflora, *Rich. ex A. Grey, Bot. U. St. Expl. Exp.* 554.—Ins. Philipp.
myrtifolia, *Link, Enum. Hort. Berol.* ii. 30.—Austral.
nervosa, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 129 = oblique.
nitida, Hook. f. Fl. Tasm. i. 127. t. 29 = amygdalina.
nutans, *F. Muell. Fragm.* iii. 152 = platypus.
obcordata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 337.—Austral.
obliqua, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 434 = Decaneana.
obliqua, *L'Hérit. Sert. Angl.* 18. t. 20.—Austral.
oblonga, *DC. Prod.* iii. 217.—Austral.
obtusiflora, *DC. l. c.* 220.—Austral.
obtusifolia, G. Don, Gen. Syst. ii. 820, sphalm. = obtusiflora.
occidentalis, *Endl. in Enum. Pl. Hueg.* 49.—Austral.
ochrophloia, *F. Muell. Fragm.* xi. 96.—Austral.
odontocarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 98.—Austral.
odorata, *Behr, in Linnæa*, xx. (1847) 657.—Austral. Oldfieldii, *F. Muell. Fragm.* ii. 37.—Austral.
oleosa, *F. Muell. l. c.* 56.—Austral.
oleosa, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 127 = uncinata.
oligantha, *Schau. in Walp. Rep.* ii. 926.—Austral.
oppositifolia, Desf. Tabl. ed. 1. 232 = purpurescens.
orbifolia, *F. Muell. Fragm.* v. 60.—Austral.
ornata, Sieber, ex Benth. Fl. Austral. iii. 208 = pilularis.
ovata, Labill. Nov. Holl. Pl. ii. 13. t. 153 = brachypoda !
pachyloma, *Benth. Fl. Austral.* iii. 237.—Austral.
pachyphylla, *A. Cunn. ex F. Muell. in Journ. Linn. Soc.* iii. (1859) 98. nomen, in nota.—Austral.
pachyphylla, *F. Muell. l. c.*—Austral.
pachypoda, *F. Muell. Fragm.* vii. 41.—Austral.
pallens, *DC. Prod.* iii. 219.—Austral.
pallidifolia, *F. Muell. Fragm.* iii. 131.—Austral.
paniculata, *Sm. in Trans. Linn. Soc.* iii. (1797) 287. —Austral.
papuana, *F. Muell. Papuan Pl.* i. 8.—N. Guinea.
parvifolora, F. Muell. in Journ. Linn. Soc. iii. (1859) 90 = bicolor.
patellaris, *F. Muell. l. c.* 84.—Austral.
patens, *Benth. Fl. Austral.* iii. 247.—Austral.
patentiflora, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 125 = melliodora.
patentiflora, F. Muell. Fragm. ii. 64, in syn. = viminalis.
pauciflora, *Sieber, ex Spreng. Syst. iv. Cur. Post.* 198. —Austral.
pellita, *F. Muell. Fragm.* iv. 159.—Austral.
peltata, *Benth. Fl. Austral.* iii. 254.—Austral.
pendula, Page, ex Stersl. Nom. ed. II. f. 600 = bicolor.
pericillata, Hort. ex DC. Prod. iii. 218 = piperita.
perfoliata, *R. Br. ex Benth. Fl. Austral.* iii. 258.— Austral.
perfoliata, *Desf. Cat. Hort. Par.* ed. III. 408.—Austral.
persicifolia, DC. Prod. iii. 217 = pilularis, sideraphloia.
persicifolia, Lodd. Bot. Cab. t. 501 = viminalis.
persicifolia, Miq. in Nederl. Kruidk. Arch. iv. (1859) 137 = Stuartiana.
phlebophylla, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 140 = pauciflora.
phoenicea, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 91.—Austral.
pilularis, DC. Prod. iii. 218 = viminalis.
pilularis, *Sm. in Trans. Linn. Soc.* iii. (1797) 284.— Austral.
piperata, Stokes, Bot. Mat. Med. iii. 69 = piperita.
piperita, Hort. ex DC. Prod. iii. 221 = viminalis.
piperita, *Sm. in Trans. Linn. Soc.* iii. (1797) 286.— Austral.
piperita, Sm. in White, Journ. Voy. N. S. Wales, 236 = capitellata.
Planchoniana, *F. Muell. Fragm.* xi. 48.—Austral.
platyphylla, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 98.—Austral.
platypoda, Cav. Ic. iv. 32. t. 341 = botryoides.
platypus, *Hook. Ic. Pl.* t. 949.—Austral.
pleurocarpa, Schau. in Lehm. Pl. Preiss. i. 182 = tetragona.
pluriloculoris, F. Muell. Fragm. ii. 70 = Preissiana.

EUCALYPTUS :—
polyanthemos, *Schau. in Walp. Rep.* ii. 924.— Austral.
polycarpa, F. Muell. in Journ. Linn. Soc. iii. (1859) 88 = terminalis.
polyclada, F. Muell. l. c. 98 = clavigera.
populifolia, *Desf. Cat. Hort. Par.* ed. III. 408.— Hab. !
populifolia, Hook. Ic. Pl. t. 879 = populnea.
populifolia, Hook. in Mitch. Journ. Trop. Austral. 204 = platypoda ?
populnea, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 93.—Austral.
porosa, Miq. in Nederl. Kruidk. Arch. iv. (1859) 131 = odorata.
Preissiana, *Schau. in Lehm. Pl. Preiss.* i. 181.— Austral.
procera, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 20.— Austral.
prunosa, *Schau. in Walp. Rep.* ii. 926.—Austral.
pruinosa, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) 11. 23 = pyriformis.
ptychocarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 90.—Austral.
pulchella, *Desf. Cat. Hort. Par.* ed. III. 408.— Austral.
pulverulenta, Link, Enum. Hort. Berol. ii. 31 = glauca.
pulverulenta, *Sims, Bot. Mag.* t. 2087.—Austral.
pulvigera, A. Cunn. in Field, N. S. Wales, 350 = pulverulenta.
punctata, *DC. Prod.* iii. 217.—Austral.
purpurascens, *Link, Enum. Hort. Berol.* ii. 31.— Austral.
pyriformis, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) 11. 22.—Austral.
pyrophora, *Benth. Fl. Austral.* iii. 257.—Austral.
racemosa, *Cav. Ic.* iv. 24.—Austral.
radiata, Sieber, ex DC. Prod. iii. 218 = amygdalina.
Rameliana, *F. Muell. Fragm.* x. 84.—Austral.
Raveretiana, *F. Muell. l. c.* 99.—Austral.
redunca, *Schau. in Lehm. Pl. Preiss.* i. 127.—Austral
regnans, *F. Muell. in Rep. Acclim. Soc. Vict.* (1870) 20.—Austral.
resinifera, *Sm. in White, Journ. Voy. N. S. Wales*, 231. —Austral.
reticulata, *Link, Enum. Hort. Berol.* ii. 29.—Austral.
rigida, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 114.—Austral.
rigida, Sieber, ex Benth. Fl. Austral. iii. 205 = obtusiflora.
Risdoni, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 477.—Tasmania.
robusta, *Sm. in Bot. N. Holl.* 39; *et in Trans. Linn. Soc.* iii. (1797) 283.—Austral.
rostrata, Cav. Ic. iv. 23 = robusta.
rostrata, *Schlecht. in Linnæa*, xx. (1847) 655.— Austral.
rubricaulis, *Desf. Cat. Hort. Par.* ed. III. 408.— Austral.
rudis, *Endl. in Enum. Pl. Hueg.* 49.—Austral.
salicifolia, *Cav. Ic.* iv. 24.—Austral.
saligna, *Sm. in Trans. Linn. Soc.* iii. (1797) 285.— Austral.
salmonophloia, *F. Muell. Fragm.* xi. 11.—Austral.
salubris, *F. Muell. l. c.* 54.—Austral.
santaliflora, Miq. in Nederl. Kruidk. Arch. iv. (1859) 138 = dumosa.
santalifolia, *F. Muell. in Trans. Vict. Inst.* i. (1855) 35.—Austral.
santaliflora, F. Muell. l. c. partim = cneorifolia.
Saxesa, *Blume, Mus. Bot. Lugd. Bat.* i. 84.—Ins. Moluce.
scabra, Dum.-Cours. Bot. Cult. ed. II. vii. 280 = pilularis.
semicorticata, F. Muell. in Journ. Linn. Soc. iii. (1859) 81 = pilularis.
septentrionalis, *F. Muell. Eucalypt. Decad.* viii. (1883).— Austral.
serica, *Schau. in Walp. Rep.* ii. 926.—Austral.
sideraphloia, *Benth. Fl. Austral.* iii. 230.—Austral.
sideroxylon, A. Cunn. in Mitch. Journ. Trop. Austral. 239 = leucoxylon.
Sieberiana, F. Muell. Eucalypt. Decad. ii. (1879) = virgata.
signata, F. Muell. in Journ. Linn. Soc. iii. (1859) 85 = haemastoma.
smithii, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 123 = oleosa.

EUCALYPTUS:—
spathulata, *Hook. Ic. Pl.* t. 611.—Austral.
spectabilis, F. Muell. Fragm. v. 45 = pellita.
spicheriocarpa, Hook. Bot. Mag. t. 4086 = calophylla.
spodophylla, F. Muell. Fragm. ii. 71 = pruinosa.
Staigeriana, *F. Muell. ex Bailey, Syn. Queensl. Fl.* 176.—Austral.
stellulata, *Sieber, ex DC. Prod.* iii. 217.—Austral.
stenophylla, *Link, Enum. Hort. Berol.* ii. 30.—Austral.
stricta, *Sieber, ex Spreng. Syst. iv. Cur. Post.*, 195.—Austral.
Stuartiana, *F. Muell. ex Miq. in Nederl. Kruidk. Arch.* iv. (1859) 181.—Austral.
subemarginipetala, Miq. l. c. 188 = paucifora.
subulata, A. Cunn ex Schau. in Walp. Rep. ii. 924 = terreticornis.
sylvicultrix, F. Mnell. ex Benth. Fl. Austral. iii. 201 = paucifora.
tectifica, F. Muell. in Journ. Linn. Soc. iii. (1859) 92 = alba.
tenuiramis, Miq. in Nederl. Kruidk. Arch. iv. (1859) 128 = amygdalina.
terticornis, *Sm. Bot. N. Holl.* 41; *et. in Trans. Linn. Soc.* iii. (1797) 284.—Austral.
terminalis, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 89.—Austral.
terminalis, Sieber, ex Benth. Fl. Austral. iii. 211 = paniculata.
teucelliaria, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 88.—Austral.
tetragona, *F. Muell. Fragm.* iv. 51.—Austral.
tetraptera, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 22.—Austral.
tetrodonta, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 97.—Austral.
Todtiana, *F. Muell. in Wing. South Sc. Record*, ii. (1882) 171.—Austral.
Toreliana, *F. Muell. Fragm.* x. 106.—Austral.
trachyphloia, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 90.—Austral.
transversa, Link, Enum. Hort. Berol. ii. 30 = capitellata.
tuberculata, *Parm ex DC. Prod.* iii. 221.—Austral.
turbinata, Behr & F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 137 = oleosa.
umbellata, *Dum.-Cours. Bot. Cult.* ed. II. vii. 279.—Austral.
uncinata, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 23.—Austral.
undulata, F. Muell. in Journ. Linn. Soc. iii. (1859) 95 = ferruginea.
urnigera, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 477.—Tasmania.
variegata, F. Muell. in Journ. Linn. Soc. iii. (1859) 88 maculata.
vernicosa, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 478.—Tasmania.
versicolor, *Blume, Mus. Bot. Lugd. Bat.* i. 84.—Ins. Moluc.
viminalis, Hook. in Mitch. Journ. Trp. Austral. 157 = tesselaris.
viminalis, *Labill. Nov. Holl. Pl.* ii. 12. t. 151.—Austral.
virgata, *Sieber, ex DC. Prod.* iii. 217.—Austral.
Watsoniana, *F. Muell. Fragm.* x. 98.—Austral.
Woollsii, F. Muell. l. c. ii. 50 = longifolia.
xanthonema, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 168 = reducta.
Youngiana, *F. Muell. Fragm.* x. 5.—Austral.

EUCAPNIA, Rafin. Fl. Tellur. iii. 74 (1836) = Nicotiana, Linn. (Solanac.).
repanda, Rafin. l. c. = N. repanda.

EUCAPNOS, Bernh. in Linnaea, viii. (1833) 468 = Dicentra, Bernh. (Papav.).
eximius, Bernh. l. c. = D. eximia.
formosus, Bernh. l. c. = D. formosa.
spectabilis, Sieb. & Zucc. in Abh. Akad. Muench. iii. (1846) 721 = D. spectabilis.

EUCARPHA, Spach, Hist. Veg. Phan. x. 408 (1841) = Knightia, R. Br. (Proteac.).

EUCARYA, Mitch. Three Exped. Austral. ii. 100 (1838) = Fusanus, R. Br. (Santalac.).
Merrapenea, Mitch. l. c. = F. pentandra ?

EUCENTRUS, Endl. Gen. Suppl. v. 82 (1850) = ENCENTRUS, Presl = Celastrus, Linn.

EUCEPHALUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 396 = Aster, Tourn (Compos.).
albus, Nutt. l. c. 399 = A. prairicoides.
elegans, Nutt. l. c. 398 = A. elegans.
ericoides, Nutt. l. c. 399 = A. ericoides.
glaucus, Nutt. l. c. = A. glaucus.

EUCERARIA, Mart. Nov. Gen et Sp. iii. 90. t. 238 (1829). *SAMYDACEAE*, Benth. & Hook. f. i. 797.
nitida, *Mart. l. c.*—Bras.

EUCHARIS, Bartl. & Wendl. f. Diosm. 15. t. A (1824). *RUTACEAE*, Benth. & Hook. f. i. 289.
abietina, Eckl. & Zeyh. Enum. 104 = glomerata.
dubia, Sond. in Harv. & Sond. Fl. Cap. i. 373 = Macrostylis dubia.
elata, *Eckl. & Zeyh. Enum.* 104.—Afr. austr.
flexilis, *Eckl. & Zeyh. l. c.* 105.—Afr. austr.
glomerata, *Bartl. & Wendl. f. Diosm.* 16.—Afr. austr.
glomerata, Eckl. & Zeyh Enum. 104 = linearis.
lacvigata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 488.—Afr. austr.
linearis, *Sond. in Harv. & Sond. Fl. Cap.* i. 372.—Afr. austr.

EUCHARIDIUM, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 36 (1835). *ONAGRARIEAE*, Benth., & Hook. f. i. 790.
Brewer, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 532.—Calif.
concinnum, *Fisch. & Mey. Ind. Sem Hort. Petrop.* ii. 36.—Calif.
grandiflorum, *Fisch. & Mey. l. c.* vii. 48.—Calif.

EUCHARIS, Planch. & Linden, Cat. n. viii. (1853) 3; et in Fl. des Serres, Sér. I. viii. (1852–53) 107. *AMARYLLIDEAE*, Benth. & Hook f. iii. 721.
MATHIRUA, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 387.
amazonica, Linden, in Illustr. Hortic. xxviii. (1881) 30 = grandiflora.
candida, *Planch. & Linden, Cat.* n. viii. (1853) 3; *et in Fl. des Serres*, Sér. I. viii. (1852–53) 107.—N. Granat.
galanthoides, *Planch. & Linden, Cat.* xvii. i. (1862) 4.—Bras.
grandiflora, *Planch. & Linden, in Fl. des Serres*, Sér. I. ix. (1853–54) 255.—N. Granat.
Hartwegiana, Nichols. Dict. Gard. i. 536 = Calipharuia Hartwegiana.
Masterii, *Baker, in Bot. Mag.* t. 6831.—Hab.?
paradoxa, T. Moore, in Gard. Chron. (1876) i. 242 = subedentata.
Sanderi, *Baker, in Bot. Mag.* t. 6676.—N. Granat.
subedentata, *Benth. & Hook. f. Gen.* iii. 731; in nota sub Calipharuia (subedentula).—N. Granat.

EUCHILOPSIS, F. Muell. in Melbourne Chemist, June (1882); ex Census, 87. *LEGUMINOSAE.*
linearis, *F. Muell. l. c.*—Austral.

EUCHIDIUM, Endl. Gen. Suppl. v. 92 (1850) = ENCHIDIUM, Jack = Trigonostemon, Blume (Euphorb.).

EUCHILOS, Spreng. Gen. i. 866 (1830) = seq.

EUCHILUS, R. Br. in Ait. Hort. Kew. ed. II. iii. 17 (1811) = Pultenaea, Sm. (Legumin.).
calycinus, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 276 = P. calycina.
crinitpodus, F. Muell. Fragm. i. 145 = P. rotundifolia.
cuspidatus, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 68 = P. ternata.
linearis, Benth. in Enum. Pl. Hueg. 104 = Sphaerolobium Erichilus.
obcordatus, R. Br. in Ait. Hort. Kew. ed. II. iii. 17 = P. obcordata.
purpureus, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 276 = P. conferta ?
rotundifolius, Turcz. l. c. 277 = P. rotundifolia.
spinulosus, Turcz. l. c. 275 = P. spinulosa.

EUCHITON, Cass. in Dict. Sc. Nat. lvi. 214 (1828) = Gnaphalium, Linn. (Compos.).
collinus, Cass. l. c. 216 = G. collinum.
decipiens, Cass. l. c. 217 = G. decipiens.
Forsteri, Cass. l. c. 216 = G. japonicum.
pulchellus, Cass. l. c. 215 = G. japonicum.

EUCHLAENA, Schrad. Ind. Sem. Hort. Gotting. (1839).—Cf. Linnaea, viii. (1833) Litt. 25. *GRAMINEAE*, Benth. & Hook f. iii. 1114.
KRANA, Brign. Ind. Sem. Hort. Mutin. (1849).—Cf. Ann. Sc. Nat. Sér. III. xii. (1849) 365.
Bourgaei, *Fourn. in Bull. Soc. Bot. Belg.* xv. (1876) 468.—Mexic.
Gouaneiniii, Fourn. l. c. = mexicana.
luxurians, Durieu & Aschers. in Bull. Soc. Linn. Par. i. (1877) 107 = mexicana.
mexicana, *Schrad. Ind. Sem. Hort. Gotting.* (1832).—*Cf. Linnaea*, viii. (1833) *Litt.* 25.—Mexic.

EUCHLORA, Eckl. & Zeyh. Enum. 171 (1836). *LEGUMINOSAE*, Benth. & Hook. f. i. 476.
MICROTROPIS, E. Mey. Comm. Pl. Afr. Austr. 65 (1835).
serpens, *Eckl. & Zeyh. l. c.*—Afr. austr.

EUCHLORIS, D. Don, in Mem. Wern. Soc. v. (1826) 548 = Helichrysum, Vaill. (Compos.).
nudifolia, D. Don. l. c. = H. nudifolium.

EUCHRESTA, Benn. Pl. Jav. Rar. 148. t. 31 (1838). *LEGUMINOSAE*, Benth. & Hook. f. i. 550.
Horsfieldii, *Benn. l. c.*—Java.
japonica, *Benth. ex Oliver, in Fourn. Linn. Soc.* ix. (1867) 164.—Japon.

EUCHROMA, Nutt. Gen. Am. ii. 54 (1818) = Castilleja, Linn. (Scrophular.).
angustifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 46 = C. parviflora.
Bradburii, Nutt. l. c. 47 = G parviflora.
coccinea, Nutt. Gen. Am. ii. 55 = C. coccinea.
grandiflora, Nutt. l. c. = C. sessiliflora.
integrifolia, Nutt. ex Benth. in DC. Prod. x. 582 = C. miniata.
pallescens, Nutt. ex A. Gray, Syn. Fl. N. Am. ii. i. 299 = Orthocarpus pallescens.
purpurea, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 180 = C. purpurea.

EUCHYLAENA, Spreng. Gen. i. 228 (1830) = Euchylaena, R. Br. (Chenop.).

EUCHYLIA, Dulac. Fl. Hautes-Pyr. 468 (1867) = Loniosra, Linn. (Caprifol.).
alpigena, Dulac, l. c. = L. alpigena.
caerulea, Dulac, l. c. = L. caerulea.
nigra, Dulac, l. c. = L. nigra.
pyrenaica, Dulac, l. c. = L. pyrenaica.
verticillata, Dulac, l. c. 464 = L. Periclymenum.
villosa, Dulac, l. c. = L. Xylosteum.

EUCHYLUS, Foir. in Dict. Sc. Nat. xv. (1819) 216 = EUCHILUS, R. Br. = Pultenaea, Sm. (Legumin.).

EUCLADUS, Nutt. ex Hook. Ic. Pl. t. 650 (1844) = Schiedea, Cham. & Schlecht. (Caryophyll.).
suffruticosus, Nutt. ex Hook. l. c. = S. Nuttallii.

EUCLEA, Murr. Syst. Veg. ed. XIII. 747 (1774). *EBENACEAE*, Benth. & Hook. f. ii. 664.
BRACHYCHILA, Harv. in Linnaea, xx. (1847) 192.
DIPLONEMA, G. Don, Gen. Syst. iv. 42 (1837).
KELLAUA, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 95.
RYMIA, Endl. Gen. 748 (1839).
acutifolia, *E. Mey. in Drège, Zwei Pfl. Docum.* 184.—Afr. austr.
angustifolia, Benth. in Hook. Niger Fl. 443 = E. Pseudebenus.
Balfourii, *Hiern, ex Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 406.—Ins. Socotra.
bilocularis, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 102.—Afr. trop.
coriacea, *A. DC. in DC. Prod.* viii. 216.—Afr. austr.
daphnoides, *Hiern, in Trans. Camb. Phil. Soc.* xii. t. (1873) 103.—Afr. austr.

EUCLEA:—
deserforum, Eckl. & Zeyh. in Linnæa, xx. (1847) 192 = lanceolata.
divinorum, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 99.—Afr. austr.
Dregeana, A. DC. in DC. Prod. viii. 216 = polyandra.
elliptica, A. DC. l. c. = polyandra.
ferruginea, Bernh. in Flora, xxvii. (1844) 825 = polyandra.
fructuosa, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 101.—Afr. trop.
herbacea, Lour. Fl. Cochinch. ii. 629 (Euphorbiacea).—China.
humilis, Eckl. & Zeyh. in Linnæa, xx. (1847) 192 = undulata.
Kellau, *Hochst. in Flora*, xxvi. (1843) 88.—Abyss.
Krausziana, Bernh. in Flora, xxvii. (1844) 824 = tomentosa.
lances, *Thunb. Prod. Pl. Cap.* 95.—Afr. austr.
lanceolata, *E. Mey. in Dregε, Zwei Pfl. Docum.* 184.—Afr. austr.
laurina, *Hiern, ex Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 406.—Ins. Socotr.
linearis, *Zeyh. in Linnæa*, xx. (1847) 192.—Afr. austr.
macrophylla, E. Mey. in Drège, Zwei Pfl. Docum. 184 = natalensis.
multiflora, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 100.—Afr. austr.
myrtina, Burch. Trav. Afr. ii. 538 = undulata.
natalensis, *A. DC. in DC. Prod.* viii. 218.—Afr. austr.
ochracea, E. Mey. in Drège, Zwei Pfl. Docum. 184 = lanceolata.
ovata, *Burch. Trav. Afr.* i. 337.—Afr. austr.
pilosa, Lour. Fl. Cochinch. ii. 629 = Diospyros pilosa.
polyandra, *E. Mey. in Drège, Zwei Pfl. Docum.* 184.—Afr. austr.
Pseudobenus, *E. Mey. l. c.*—Afr. austr.
pubescens, Eckl. & Zeyh. in Linnæa, xx. (1847) 192 = polyandra.
racemosa, *Murr. Syst. Veg.* ed. XIII. 747.—Afr. austr.
rigida, E. Mey. in Drège, Zwei Pfl. Docum. 184 = lanceolata.
rufescens, E. Mey. l. c. = ovata.
tomentosa, *E. Mey. l. c.*—Afr. austr.
undulata, *Thunb. Nov. Gen. Pl.* v. (1784) 86.—Afr. austr.

EUCLIANDRA, Steud. Nom. ed. II. i. 600 (1840) = ENCLIANDRA, Zucc. = **Fuchsia**, Linn. (Onagr.).

EUCLIDIUM, R. Br. in Ait. Hort. Kew. ed. II. iv. 74 (1813). *CRUCIFERÆ*, Benth. & Hook. f. i. 97.
ORNITHORYNCHIUM, Roehl. Deutschl. Fl. ed. II. 886 (1819–13).
SORIA, Adans. Fam. ii. 421 (1763).
aegyptiacum, Andrz. ex DC. Syst. ii. 488 = Ochthodium aegyptiacum.
syriacum, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 74.—Reg. Casp. ; Afghan.
tataricum, *DC. Syst.* ii. 492.—Europ. or. ; Orient.

EUCNEMIA, Reichb. Nom. 88 (1841) = seq.

EUCNEMIS, Lindl. Gen. et Sp. Orch. 161 (1838) = **Govenia**, Lindl. (Orchid.).
brevilabris, Lindl. l. c. = G. brevilabris.

EUCNIDE, Zucc. Del. Sem. Monac.—*Cf.* Linnæa, xvii. (1844) 508; et in Abh. Akad. Muench. iv. 1. (1845) 1. = **Mentzelia**, Linn. (Loasac.).
bartonioides, Zucc. ll. cc. = M. grosoviæfolia.
floribunda, S. Wats. in Proc. Am. Acad. xvii. (1881–82) 356.—Calif.
lobata, A. Gray, Pl. Lindh. 192 = M. lobata.
urens, Parry, in Am. Natural. ix. (1875) 144 = M. urens.

EUCODONIA, Hanst. in Linnæa, xxvi. (1853) 201 = **Achimenes**, P. Br. (Gesnerac.).
Ehrenbergii, Hanst. l. c. 201 = A. longa.
illaticaulis, Van Houtte, Fl. des Serres, xvii. (1867–68) 79. t. 1737–38.—Hab. ?

EUCODONOPSIS, Van Houtte, in Fl. des Serres, xvi. (1865) t. 1606 = **Achimenes**, P. Br (Gesnerac.).
saxgnioides x, Van Houtte, l. c.

EUCOLUM, Salisb. Prod. 98 (1796) = **Gloxinia**, L'Hérit. (Gesner.).
crassifolium, Salisb. l. c. = G. maculata.

EUCOMEA, Soland. ex Salisb. Prod. 218 (1796) = **Eucomis**, L'Hérit. (Liliac.).
clata, Salisb. l. c. = Eucomis punctata.
humilis, Salisb. l. c. = Eucomis nana.
regalis, Salisb. l. c. = Eucomis regia.

EUCOMIS, L'Hérit. Sert. Angl. 17. t. 18 (1788). *LILIACEÆ*, Benth. & Hook. f. iii. 813.
BASILARA, Juss. ex Lam. Encyc. i. 362 (1783) ; et Gen. 59 (1789).
EUCOMEA, Soland. ex Salisb. Prod. 218 (1796).
amaryllidifolia, *Baker, in Gard. Chron.* (1878) ii. 492.—Afr. austr.
bicolor, *Baker, l. c.*—Afr. austr.
bifolia, Jacq. Coll. iv. 215 = Whitheadia latifolia.
clavata, Baker, in Saund. Refug. Bot. t. 298 = regia.
nana, *L'Hérit. Sert. Angl.* 17.—Afr. austr.
macrophylla, Hort. ex Baker, in Journ. Linn. Soc. xiii. (1873) 325 = regia.
punctata, *L'Hérit. Sert. Angl.* 18.—Afr. austr.
purpureo-caulis, Andr. Bot. Rep. t. 359 = nana.
regia, *Ait. Hort. Kew.* ed. I. i. 438.—Afr. austr.
regia, L'Hérit. Sert. Angl. 17 = undulata.
striata, Donn, Hort. Cantab. ed. II. 40 = punctata.
undulata, *Ait. Hort. Kew.* ed. I. i. 438.—Afr. austr.

EUCOSIA, Blume, Bijdr. 415. t. 18 (1825). *ORCHIDEÆ*, Benth. & Hook. f. iii. 605.
carnea, *Blume, i. c.*—Java.

EUCRINUM, Nutt. ex Lindl. Veg. Kingd. 204 (1847), nomen. *LILIACEÆ*.

EUCRIPHIA, Pers. Syn. ii. 92 (1807) = **Eucryphia**, Cav.

EUCROSIA, Ker-Gawl. in Bot. Reg. t. 207 (1817). *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 730.
bicolor, *Ker-Gawl. l. c.*—Am. austr.

EUCRYPHIA, Cav. Ic. iv. 48. t. 372 (1797). *ROSACEÆ*, Benth. & Hook. f. i. 615.
CARPODONTOS, Labill. Voy. ii. 18. t. 18 (1798).
PELLINIA, Molina, Sagg. Chile, ed. II. 160 (1810).
Billardieri, *Spach, Hist. Vég. Phan.* v. 345.—Austral.
cordifolia, *Cav. Ic.* iv. 49. t. 372.—Chili.
Milliganii, Hook. f. Fl. Tasm. i. 54 = Billardieri.
Moorei, *F. Muell. Fragm.* iv. 2.—Austral.
pinnatifolia, *C. Gay, Fl. Chil.* i. 353. t. 8.—Chili.

EUCRYPTA, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 166 = **Hitchcockia**, Linn. (Hydrophyll.).
chrysanthemifolia, Greene, in Bull. Calif. Acad. i. iv. (1885) 200 = Ellisia chrysanthemifolia.
foliosa, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 159 = Ellisia chrysanthemifolia.
paniculata, Nutt. l. c. = Ellisia chrysanthemifolia.

EUCYCLA, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 166 = **Eriogonum**, Michx. (Polygonac.).
ovalifolia, Nutt. l. c. = Eriogonum ovalifolium.
purpurea, Nutt. l. c. = Eriogonum ovalifolium.

EUDEMA, Humb. & Bonpl. Pl. Aequin. ii. 138. t. 125 (1808) = **Erato**, Sternb. & Hoppe (Cruciferæ).
grandiflora, Planch. in Hook. Lond. Journ. Bot. iii. (1844) 630. t. 22.—Am. austr.
nubigena, Humb. & Bonpl. Pl. Aequin. ii. 136. t. 124.—Ecuador.
rupestris, Humb. & Bonpl. l. c. 138. t. 128 = Extrema Humboldtii.
thlaspiformis, Phil. in Anal. Univ. Chil. (1872) 673.—Chili.

EUDESMIA, R. Br. in Flind. Voy. ii. 599. t. 8 (1814) = **Eucalyptus**, L'Hérit. (Myrtac.).
erythrocorys, F. Muell. Fragm. ii. 83 = Eucalyptus erythrocorys.
eucalyptoides, F. Muell. l. c. 85 = Eucalyptus eudesmoides.
tetragona, R. Br. in Flind. Voy. ii. 600 = Eucalyptus tetragona.

EUDESMIS, Rafin. Fl. Tellur. iii. 80 (1836) = **Colchicum**, Linn. (Liliac.).
fascicularis, Rafin. l. c. = C. montanum.

EUDIANTHE, Reichb. Nom. 206 (1841) = **Lychnis**, Linn. (Caryophyll.).
Coeli-rosea, Reichb. Ic. Fl. Germ. vi. 55. t. 208 = L. Coeli-rosea.
corsica, Reichb. ex Willk. Ic. et Desc. Pl. Hisp. i. 29 = L. corsica.
laeta, Reichb. ex Willk. l. c. = Silene laeta.
oculata, A. Braun. Del. Sem. Hort. Friburg. (1849) 3 = L. Coeli-rosa.

EUDIPETALA, Rafin. Fl. Tellur. iii. 68 (1836) = **Commelina**, Linn.
deficiens, Rafin. l. c. = C. virginica.

EUDISTEMON, Rafin. Med. Fl. ii. 219 (1830) = **Senebiera**, DC. (Cracif.).

EUDODECA, Steud. Nom. ed. II. i. 601 (1840) = ENDODECA, Rafin. = **Aristolochia**, Linn.

EUDOLON, Salisb. Gen. Pl. Fragm. 127 (1866) = **Strumaria**, Jacq. (Amaryllid.).
undulata, Salisb. l. c. = S. undulata.

EUDORUS, Cass. in Bull. Soc. Philom. (1816) 185 ; et in Dict. Sc. Nat. xv. 525 (1819) = **Senecio**, Linn (Compos.).
senecioides, Cass. ll. cc. = S. Eudorus.

EUDOXIA, D. Don, ex G. Don, Gen. Syst. iv. 202 (1838) = **Gentiana**, Tourn.
formosissima, D. Don, ex G. Don, l. c.—Peruv.
pendula, D. Don, ex G. Don, l. c.—Peruv.

EUFORBIA, Tenore, Fl. Nap. i. 258 (1811–15) = **Euphorbia**, Linn.

EUFRAGIA, Griseb. Spicil. Fl. Rumel. ii. 13 (1844) = **Bartsia**, Linn. (Scrophul.).
atrovioiacea, Dur. ex Gass. Enum. Pl. Inar. 243 = B. latifolia.
latifolia, Griseb. Spicil. Fl. Rumel. ii. 14 = B. latifolia.
viscosa, Benth. in DC. Prod. x. 548 = B. viscosa.
Virtanii, Cos. in Bull. Soc. Bot. Fr. xii. (1865) 279, 282 = B. latifolia.

EUGENISSOA, *Griff. in* Calc. Journ. Nat. Hist. v. (1845) 301. *PALMÆ*, Benth. & Hook. f. iii. 934.
minor, *Becc. in Nuov. Giorn. Bot.* iii. (1871) 18.—Borneo.
insignis, *Becc. l. c.* 22.—Borneo.
triste, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 301.—Malaya.
utilis, *Becc. in Nuov. Giorn. Bot.* iii. (1871) 26.—Borneo.

EUGENIA, Mich. ex Linn. Syst. ed. I (1735). *MYRTACEÆ*, Benth. & Hook. f. i. 718.
ACMENA, DC. in Dict. Class. Hist. Nat. xi. 401 (1826).
AMYRIS, Rafin. Syiva Tellur. 106 (1838).
CALOPHYLLOIDES, Smeathm. ex DC. Prod. iii. 272 (1828).
CALYCISANTHUS, Blume, Bijdr. 1089 (1826).
CARYOPHYLLUS, Linn. Syst. ed. I (1735).
CATINGA, Aubl. Pl. Gui. 511. t. 203 (1775).
CEROCARPUS, Hamk. in Flora, xxv. (1842) ii. Beibl. 96 ; Cat. Hort. Reg. Alt. 209 (1844).
CLAVIMYRTUS, Blume, Mus. Bot. Lugd. Bat. i. 115. t. 49 (1849).
CLINITOCALYX, Blume, l. c. 84. t. 36 (1849).
CUMETRA, Rafin. Syiva Tellur. 106 (1838).
EMURTIA, Rafin. l. c. (1838).
EPLESIUDA, Rafin. l. c. 107 (1838).
GELPEKIA, Blume, Mus. Bot. Lugd. Bat. i. 86. t. 35 (1849).
GREGGIA, Geertn. Fruct. i. 162. t. 33 (1788).
GUAPURIUM, Juss. Gen. 384 (1789).
HEXACHLAMYS, Berg, in Linnæa, xxvii. (1854) 345.
JAMBOLIFERA, Houtt. Handleid. ii. 279 (1774).
JAMBOS, Adans. Fam. ii. 88 (1763).
JOSSINIA, Comm. ex DC. Prod. iii. 267 (ex typ. 267) (1828).
LOMASTELMA, Rafin. Syiva Tellur. 107 (1838).
MACROSTYLUS, Miq. Fl. Ind. Bat. i. 1. 469 (1855).

EUGENIA :—
MALIDRA, Rafin. Syiva Tellur. 107 (1838).
MICROJAMBOSA, Blume, Mus. Bot. Lugd. Bat. i. 117 (1849).
MYRCIARIA, Berg, in Linnæa, xxvii. (1854) 930.
OLYNTHIA, Lindl. Collect. 19, in obs. (1825).
OPA, Lour. Fl. Cochinch. 308 (1790).
OPANEA, Rafin. Syiva Tellur. 106 (1838).
PHYLLOCALYX, Berg, in Linnæa, xxvii. (1854) 606.
PLINIA, Plum. ex Linn. Syst. ed. I (1735).
RUGENIA, Neck. Elem. ii. 78 (1790).
SIPHONEUGENIA, Berg, in Linnæa, xxvii. (1854) 844.
STENOCALYX, Berg, l. c. 809 (1854).
STRONGYLOCALYX, Blume, Mus. Bot. Lugd. Bat. i. 88, t. 64 (1849).
SUARDA, Nocca, ex Steud. Nom. ed. II. ii. 651 (1841).
SYLLYNIUM, Meyen & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 384.
SYZYGIUM, Gaertn. Fruct. i. 166. t. 38 (1788).

acapulcensis, *Steud. Nom.* ed. II. i. 601.—Mexic.
Acca, Berg, in Linnæa, xxvii. (1854) 793, sphalm., est nomen generis.
acetosæa, *Pav. Encyc. Suppl.* iii. 125.—Guian.
Acka, *DC. Prod.* iii. 277.—Peruv.
acris, *Wight & Arn. Prod.* 331.—Ind. or.
acrophylla, *Berg, in Linnæa*, xxvii. 1. 575.—Bras.
acuminata, Link, Enum. Hort. Berol. ii. 28 = Myrtus serices.
acuminata, *Roxb. Hort. Beng.* [37]; *Fl. Ind.* ii. 492. —Ins. Molucc.
acuminatissima, *Berg, in Mart. Fl. Bras.* xiv. i. 315.—Bras.
acuminatissima, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 67; For. Fl. Burma, i. 487 = E. Thwaa.
acuminatissima, Miq. in Linnæa, xix. (1847) 440 (= *Blepharocalyx acuminatissima*).—Bras.
acutangula, Linn. Sp. Pl. 471 = Barringtonia acutangula.
acutata, *Miq. in Linnæa*, xxii. (1849) 535.—Bras.
acutilobis, *DC. Prod.* iii. 281.—Jamaic.
adamantium, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 342. —Bras.
adenantha, *Berg, in Mart. Fl. Bras.* xiv. i. 378.—Bras.
adenocalyx, *DC. Prod.* iii. 271.—Guian.
adenocarpa, *Berg, in Mart. Fl. Bras.* xiv. i. 369.—Bras.
aegicroides, *Korth. ex Miq. Anal. Bot. Ind.* i. 22.—Malaya.
aeruginea, *DC. Prod.* iii. 282.—Am. trop.
affinis, *DC. l. c.* 279.—Ind. occ.
affinis, Gill. in Hook. Bot. Misc. iii. (1833) 321 = Hookeri.
alaternifolia, *Benth. ex Berg, in Linnæa*, xxvii. (1854) 249.—N. Granat.
alba, Roxb. Hort. Beng. [37]; Fl. Ind. ii. 493 = javanica.
albanensis, *Sond. in Hors. & Sond. Fl. Cap.* ii. 522.—Afr. austr.
albicans, A. Rich. ex Berg, in Linnæa, xxx. (1859–60) 683 = Calyptranthes cuprea.
albida, *Steud. & Bongé Pl. Aequin.* ii. 107. t. 113.—N. Granat.
albiflora, *Duthie, ex Kurz, For. Fl. Burma*, i. 491.—Burma.
albo-tomentosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 349.—Bras.
alpigena, *DC. Prod.* iii. 265.—Bras.
alpina, *Willd. Sp. Pl.* ii. 961.—Jamaic.
alternifolia, Benth. Fl. Hartw. 175 = alaternifolia.
alternifolia, *Wight, Ic.* t. 587; *Illustr.* ii. 16.—Ind. or.
amazonica, *Berg, in Mart. Fl. Bras.* xiv. i. 322.—Bras.
amblyoptera, *Berg, l. c.* 341.—Bras.
amboinensis, Hort. ex DC. Prod. iii. 294 = Metrosideros vera.
amboinensis, *Note. ex Link, Enum. Hort. Berol.* ii. 26.—Ins. Amboin.
americana, Hort. Makoy, ex E. Morr. in Belg. Hortic. x. (1860) t. ad p. 25 = mexicana?
amplexicaulis, *A. Gray, Bot. U. St. Expl. Exped.* i. 524. t. 52.—Ins. Fiji.
amoena, *Thw. Enum. Pl. Zeyl.* 114.—Zeylan.

EUGENIA :—
amplexicaulis, *Roxb. Hort. Beng.* [37]; *Fl. Ind.* ii. 488. —Ind. or.; Burma.
amplexicaulis, Vell. Fl. Flum. 310; v. t. 44 (= *Gomidesia amplexicaulis*).—Bras.
anastomosana, *DC. Prod.* iii. 269.—Guian.
ancepa, *Berg, in Mart. Fl. Bras.* xiv. i. 360.—Bras.
andina, *Berg, in Linnæa*, xxvii. (1854) 274.—Peruv.
andrearomoides, Bedd. Fl. Sylv. Anal. Gen. p. cvii = cordifolia.
androsaemoides, *DC. Prod.* iii. 284.—Ind. or.
androsaemoides, Herb. Wight, ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 491 = operculata.
angophoroides, *F. Muell. Fragm.* v. 33.—Austral
angustifolia, Blume, in Flora, vii. (1824) 291 = Blumei.
angustifolia, *Lam. Encyc.* iii. 208.—Ins. S. Doming.
angustifolia, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 490 = ventenata.
angustifolia, Spreng. Nov. Prov. 18 = Sprengelii.
angustistima, *Berg, in Mart. Fl. Bras.* xiv. i. 369.—Bras.
anisosepala, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 481. —Malacca.
aphithosa, *Vieill. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 469.—N. Caled.
apiculata, *DC. Prod.* iii. 276.—Chili.
apiocarpa, *Berg, in Mart. Fl. Bras.* xiv. i. 384.—Bras.
aprica, *Berg, l. c.* 218.—Reg. Argent.
aquea, *Burm. f. Fl. Ind.* 114.—Burma; Zeylan.
araujoana, *Berg, in Mart. Fl. Bras.* xiv. i. 319.—Bras.
Arayan, *Seem. Bot. Voy. Herald*, 125.—N. Granat.
arbutifolia, *Berg, in Mart. Fl. Bras.* xiv. i. 306.—Bras.
arctostaphylodes, *Berg, l. c.* 317.—Bras.
arenaria, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 349.—Bras.
areolata, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 490.—Reg. Himal.
argentea, *Bedd. Fl. Sylv. Anal. Gen.* p. cix.—Ind. or.
Arivos, *Aubl. Pl. Gui.* i. 510.—Guian.
armenia, *Seem. Bot. Voy. Herald*, 125.—N. Granat.
Armstrongi, *Benth. Fl. Austral.* iii. 286.—Austral.
Arnottiana, *Wight, Ic.* t. 999; *Illustr.* ii. 17.—Ind. or.
aromatica, Baill. Hist. Pl. vi. 311, 345 = caryophyllata.
aromatica, *Berg, in Linnæa*, xxvii. (1854) 268.—Venezuela.
Arrabidae, *Berg, l. c.* 302.—Bras.
arvensis, Vell. Fl. Flum. 309; v. t. 37 = oxyphylla.
asperifolia, *Berg, in Linnæa*, xxx. (1859–60) 674.—Ind. occ.
astringens, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 361. —Bras.
atropunctata, *Steud. in Flora*, xxvi. (1843) 762.—Guian.
aurata, *Berg, in Mart. Fl. Bras.* xiv. i. 273.—Bras.
australis, Colla, in Mem. Acc. Torin. xxxi. (1827) 132. t. 8 = Serteriana.
australis, Wendl. ex Link, Enum. Hort. Berol. ii. 28 = myrtifolia, *Sims*.
australis-dulcis, *Seem. Fl. Vit.* 76.—N. Caled.
axillaris, Vell. Fl. Flum. 309; v. t. 41 = supraaxillaris.
axillaris, Willd. Sp. Pl. ii. 960 = monticola.
azurensis, *Berg, in Mart. Fl. Bras.* xiv. i. 294.—Bras.
badia, *Berg, l. c.* 286.—Bras.
bagensis, *Berg, l. c.* 231.—Reg. Argent.
balansis, *DC. Prod.* iii. 271.—Bras.
Balbisiana, *Berg, in Linnæa*, xxvii. (1854) 268.—Ind. occ.
Balfourii, *Baker, Fl. Maurit.* 116.—Ins. Mascar.
balsamea, *Wight, Illustr.* ii. 16.—Reg. Himal.; Burma.
balsamifera, *Vacq. Fragm.* 40. t. 45. f. 2.—Jamaic.
Barneoudii, Berg, in Linnæa, xxvii. (1854) 263 = Hookeri.
barrosensis, *Berg, in Mart. Fl. Bras.* xiv. i. 313.—Bras.
barosensis, Bal. ex Berg, l. c. xxix. (1857–58) 238 = lanceolata.
barosensis, Griseb. in Gostt. Abh. vii. (1857) 214 = proxima.

EUGENIA :—
barosensis, *Vacq. Coll.* iii. 188; *Ic. Rar.* t. 486.—Ins. S. Doming.
batamarensis, *Berg, l. c.* 262.—Reg. Argent.
Bauangucca, Blanco, Fl. Filip. ed. I. 418 = laeta.
Beddomei, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 476.—Ind. or.
bella, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 362.—Bras.
bella, Phil. in Linnæa, xxviii. (1856) 641 = pulchra.
Benthamiana, Wight, ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 484 = Arnottiana.
Benthamii, *Berg, in Linnæa*, xxvii. (1854) 164.—Guian.
Bergiana, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 182.—Cuba.
berlyensis, *Berg, in Linnæa*, xxvii. (1854) 468.—Guian.
Berteriana, *Spreng. Syst* ii. 487.—N. Granat.
bicolor, *Berg, in Mart. Fl. Bras.* xiv. i. 293.—Bras.
biforis, Celeb. in Wall. Cat. n. 5606 = Wallichiana.
bifaria, *Wall. Pl. As. Rar.* ii. 47. t. 161.—Reg. Himal.
biflora, *DC. Prod.* iii. 276.—Jamaic.
bimarginata, *Berg, in Mart. Fl. Bras.* xiv. i. 291.—Bras.
bimarginata, DC. Prod. iii. 271 = dicrosea.
Blanchetiana, *Berg, in Mart. Fl. Bras.* xiv. i. 278.—Bras.
Blumei, *Steud. Nom.* ed. II. i. 601.—Java.
Boseri, *Baker, Fl. Maurit.* 115.—Ins. Mascar.
bolivianeta, *Berg, in Mart. Fl. Bras.* xiv. i. 321.—Am. trop.
Bonplandiana, *Berg, in Linnæa*, xxvii. (1854) 238.—N. Granat.
bonnensana, *Miq. Anal. Bot. Ind.* i. 24.—Borneo.
brachiata, *Roxb. Hort. Beng.* [37]; *Fl. Ind.* ii. 488.—Ins. Amboin.
brachybotrya, *DC. Prod.* iii. 277.—Guian.
brachycnema, *F. Muell. Fragm.* iv. 59.—Austral.
brachypoda, *DC. Prod.* iii. 274.—Guian.
brachystachya, *Berg, in Linnæa*, xxvii. (1854) 301.—Ins. Guadelup.
Brackenridgei, *A. Gray, Bot. U. St. Expl. Exped.* i. 521. t. 61.—Ins. Fiji.
bracteata, Raeusch. ex DC. Prod. iii. 264 = Myrcia bracteata.
bracteata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Myrcia laxiculata.
bractesta, *Rozb. Hort. Beng.* [37]; *Fl. Ind.* ii. 490.—Ind. or.
bracteata, Vell. Fl. Flum. 309; v. t. 40 = involucrata.
bracteolaris, Lam. ex DC. Prod. iii. 267 = brasiliensis.
bracteolata, *Wight, Ic.* t. 581; *Illustr.* ii. 15.—Burma; Malaya.
brasiensa, *DC. Prod.* iii. 276.—Bras.
brasiliana, *Aubl. Pl. Gui.* i. 511.—Guian.
brasiliensis, *Lam. Encyc.* iii. 208.—Bras.
brevifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 521.—Ins. Samoa.
brevipes, *A. Rich. Exs. Fl. Cub.* 584; *et ex Berg, in Linnæa*, xxx. (1859–60) 676.—Ind. occ.
Bridgesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 322.—Chili.
Bullockii, *Hence, in Journ. Bot.* xvi. (1878) 297.—China.
buxifolia, Lam. Encyc. iii. 209 = coriacea.
buxifolia, *Phil. in Linnæa*, xxviii. (1856) 640.—Chili.
buxifolia, Spreng. ex DC. Prod. iii. 275 = guadalupensis.
buxifolia, *Willd. Sp. Pl.* ii. 960.—Am. bor.; Ind. occ.
cabanisiana, *Berg, in Linnæa*, xxvii. (1854) 235.—Florida.
cachoeirensis, *Berg, in Mart. Fl. Bras.* xiv. i. 265.—Bras.
calcaeensis, *Bedd. Forest. Man. Bot.* 110.—Ind. or.
calliphylla, Reinw. ex De Vriese, Fl. Ind. Bat. Reinw. 69 = aquea.
calophylloides, *Wight, Ic.* t. 1000.—Ind. or.
Calophyllodes, *DC. Prod.* iii. 272.—Afr. trop.
calycina, *Benth. in Hook. Niger Fl.* 558.—Afr. trop.
calycina, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 352. —Bras.
calycolpoides, *Griseb. Fl. Brit. W. Ind.* 288.—Ind. occ.
calycoreticoides, *Berg, in Linnæa*, xxix. (1857–58) 286.—Mexic.
calycosemæ, *Berg, in Mart. Fl. Bras.* xiv. i. 276.—Bras.

EUGENIA:—
calyculata, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 271.—Porto Rico.
calycistrata, Roxb. ex Wight & Arn. Prod. 329 = E. Jambolana.
camarioca, *Wright, in Sauv. Fl. Cub.* 42.—Cuba.
Cambessedeana, *Berg, in Mart. Fl. Bras.* xiv. i. 230.—Reg. Argent.
campestris, *DC. Prod.* iii. 274.—Bras.
campestris, *Vell. Fl. Flum.* 206; v. t. 35.—Bras.
canna, *DC. Prod.* iii. 284.—Bras.
Candolleana, *DC. l. c.* 281.—Bras.
caseloneuia, *Berg, in Mart. Fl. Bras.* xiv. i. 232.—Reg. Argent.
canescens, *Berg, in Linnaea*, xxix. (1857–58) 229.—Ind. occ.
capensis, *Harv. Gen. S. Afr. Pl.* 416; *et in Fl. Cap.* ii. 292.—Afr. austr.
capitellata, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 335 = Memecylon Arnottianum.
capitullera, *Berg, in Mart. Fl. Bras.* xiv. i. 268.—Bras.
capparidifolia, *DC. Prod.* iii. 281.—Bras.
Capuli, *Schlecht. in Linnaea*, v. (1830) 580.—Mexic.
carassoides, *F. Muell. Fragm.* iii. 130.—Austral.
carragnensis, *Berg, in Linnaea*, xxvii. (1854) 240.—Costa Rica.
carthagenensis, *Jacq. Enum. Pl. Carib.* 28.—N. Granat.
caryophylla, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = E. Pseudo-caryophyllus.
caryophyllæa, *Wight, Ic.* t. 540; *Illustr.* ii. 15.—Ind. or. ; Malaya.
caryophylla, *Thunb. Diss.* 1.—Ins. Moluce.
caryophyllifolia, Lam. Encyc. iii. 198 = E. Jambolana.
caryophyllioides, Benth. in Hook. Niger Fl. 359 = calophylloides.
Casarettoana, *Berg, in Mart. Fl. Bras.* xiv. i. 230.—Bras.
casearioides, *DC. Prod.* iii. 275.—Venezuela.
casapensis, *Berg, in Linnaea*, xxvii. (1854) 296.—Peruv.
cassinoides, *Berg, in Mart. Fl. Bras.* xiv. i. 266.—Bras.
cassinoides, *Lam. Encyc.* iii. 205.—Madag.
Catharinæ, *Berg, in Mart. Fl. Bras.* xiv. i. 259.—Bras.
Catinga, *Baill. Hist. Pl.* vi. 344.—Guian.
catinguifera, *Griseb. Cat. Pl. Cub.* 89.—Cuba.
catuiliora, *Berg, in Mart. Fl. Bras.* xiv. i. 268.—Bras.
cauliflora, DC. Prod. iii. 273 = edulis.
cauliflora, Miq. in Linnæa, xxii. (1849) 587 (= *Myrciaria truncifloræ*).—Bras.
crassiflora, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 283 = Kurzii.
cerasiflora, *Miq. in Linnæa*, xxii. (1849) 793.—Bras.
cerasiformis, *DC. Prod.* iii. 274.—Java.
cerasoides, Roxb. Hort. Beng. [95]; Fl. Ind. ii. 488 = operculata.
Chekam, DC. Prod. iii. 278 = seq.
Chequen, *Molina, Sagg. Chile*, ed. II. 148.—Chili.
chinensis, *Regel, Cat. Pl. Hort. Absahov.* 57.—China.
chiquitensis, *Berg, in Mart. Fl. Bras.* xiv. i. 219.—Bras.
chloranthæ, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 487.—Malaya.
chlorocarpa, *Berg, in Mart. Fl. Bras.* xiv. i. 220.—Bras.
chlorophylla, *Berg, l. c.* 258.—Bras.
chrysantha, *Berg, l. c.* 272.—Bras.
chrysobalanoides, *DC. Prod.* iii. 285.—Ind. occ.
chrysocarpa, *Berg, in Linnaea*, xxvii. (1854) 268.—Chili.
chrysophylloides, *DC. Prod.* iii. 276.—Guian.
Chrysophyllum, *Pelr. Encyc. Suppl.* iii. 129.—Guian.
chrysteinsis, Berg, in Mart. Fl. Bras. xiv. i. t. 4. f. 75 = cristaisnsis.
ciarensis, *Berg, l. c.* 235.—Bras.
ciliolata, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 344.—Bras.
cincta, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 183.—Cuba.
cinerascens, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1846) 109.—Bras.
cinerea, Kurz, in Pegu For. Rep. Append. B. 50 = brachiata.
cinnamomea, *Vidal, Phan. Cuming. Philipp.* 178.—Ins. Philipp.
cisplatensis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 342.—Bras.

EUGENIA:—
citrifolia, *Poir. Encyc. Suppl.* iii. 129.—Guian.
clavifiora, *Roxb. Hort. Beng.* [97]; *Fl. Ind.* ii. 488.—As. trop.
clinocarpa, *DC. Prod.* iii. 287.—Bras.
clusiæfolia, *A. Grey, Bot. U. St. Expl. Exped.* i. 526.—Ins. Samoa.
clusioides, *Bronga. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 180.—N. Caled.
cœsiocarpa, *Berg, in Mart. Fl. Bras.* xiv. i. 220.—Bras.
coarensis, *DC. Prod.* iii. 267.—Bras.
coccifera, *Berg, in Mart. Fl. Bras.* xiv. i. 278.—Bras.
codyensis, *Munro, ex Wight, Illustr.* ii. 18.—Ind. or.
coffeæfolia, *DC. Prod.* iii. 272.—Guian.
colchaguiensis, *Phil. in Linnæa*, xxxiii. (1864–65) 72.—Chili.
collpensis, *Berg, in Linnæa*, xxix. (1857–58) 243.—Mexic.
collina, *DC. Prod.* iii. 281.—Bras.
colorata, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 492.—Malacca.
columbiensis, *Berg, in Linnæa*, xxvii. (1854) 269.—Venezuela.
Comete, Aubl. Pl. Guin. t. 196 = Commeta Myrcia.
commutata, *Berg, in Linnæa*, xxvii. (1854) 174.—Jamaic.
comosa, Wall. Cat. n. 3566 D = oblata.
comosa, *Cambess., Spring, ex Mart. in Flora*, xx. (1837) *Beibl.* ii. 81.—Bras.
complanata, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 353.—Bras.
complicata, *Berg, in Mart. Fl. Bras.* xiv. i. 218.—Bras.
compressa, *DC. Prod.* iii. 276.—Peruv.
compta, *Griseb. ex Berg, in Linnæa*, xxx. (1859–60) 677.—Ind. occ.
conconina, *Phil. in Linnæa*, xxviii. (1856) 640.—Chili.
concinna, Thw. Enum. Pl. Zeyl. 416 = Thwaitesii.
condensata, *Baker, in Journ. Bot.* xx. (1882) 112.—Madag.
confertifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 528. t. 61.—Ins. Fiji.
confusa, *DC. Prod.* iii. 279.—Ind. occ.
conglomerata, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 497.—Malaya.
contracta, Polr. Encyc. Suppl. iii. 125 = caryophyllæa.
contracta, Wall. Cat. n. 3602 = pellucida.
cordata, *DC. Prod.* iii. 272.—Ind. occ.
cordata, *Lam. in Oliver, Fl. Trop. Afr.* ii. 458.—Afr. trop.
cordifolia, *Wight, Ic.* t. 544; *Illustr.* ii. 16.—Zeylan.
coriacea, *F. G. Dietr. Vollst. Lexih. Gærtn. Nachtr.* iii. 268.—Ins. Maurit.
cornifiora, *F. Muell. Fragm.* v. 28.—Austral.
coronata, *Schum. & Thonn. Beskr. Guin. Pl.* 230.—Afr. trop.
coronata, *Vahl, ex DC. Prod.* iii. 271.—Afr. trop.
corralina, *Phil. in Linnæa*, xxxiii. (1864–65) 72.—Chili.
Correæ, *Berg, in Mart. Fl. Bras.* xiv. i. 277.—Bras.
corrensella, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 319 (= *Myrceugenia corrensella*).—Chili.
corticosa, Lour. Fl. Cochinch. 306 = E. Jambolana.
corymbosa, *Berg, in Linnæa*, xxxi. (1861–62) 256.—Peruv.
corymbosa, Lam. Encyc. iii. 199 = caryophyllæa.
corymbosa, Roxb. Hort. Beng. [97]; Fl. Ind. ii. 497 = densiflora.
corymbosa, Wall. Cat. n. 3566 F = lineata.
corymbantha, *F. Muell. Fragm.* ix. 145.—Austral.
corynacarpa, *A. Gray, Bot. U. St. Expl. Exped.* i. 526.—Ins. Fiji.
costaricensis, *Berg, in Linnæa*, xxvii. (1854) 218.—Costa Rica.
costata, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 272.—Porto Rico.
costata, *Berg, in Mart. Fl. Bras.* xiv. i. 277.—Bras.
costata, *Bronga. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 178.—N. Caled.
costata, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 359.—Bras.
cotinifolia, *Jacq. Obs. Bot.* iii. 3; *Lam. Encyc.* ii. 304.—Ins. Macar
Cometa, DC. Prod. iii. 245 = seq.
Cometa, Aubl. Pl. Gui. i. 497 = Myrcia Cometa.
crassifolia, *DC. Prod.* iii. 266.—Bras.
crassifolia, *Mog. in Linnæa*, xix. (1847) 439.—Bras.

EUGENIA:—
crassifolia, *Vieill. ex Bronga. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 469.—N. Caled.
crenata, *Berg, in Linnæa*, xxvii. (1854) 226.—Jamaic
crenata, Vell. Fl. Flum. 209; v. t. 39 = Velloziana.
crenulata, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 490.—Malacca.
crenulata, *Wild. Sp. Pl.* ii. 961.—Ins. S. Doming.
cristæensis, *Berg, in Mart. Fl. Bras.* xiv. i. 228.—Bras.
cristata, *Wright, in Sauv. Fl. Cub.* 41.—Cuba.
Cruckshanksii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 291.—Chili.
cryptophlebia, *F. Muell. Fragm.* ix. 144.—Austral.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 178.—Ins. Philipp.
Cumingii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 319.—Ins. Chiloe.
cuneata, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 495.—Reg. Himal.
cuneata, Wall. Cat. n. 3625 = E. Josinia.
cuneifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 144.—Madag.
cuspidata, *Berg, in Mart. Fl. Bras.* xiv. i. 286.—Bras.
cuspidata, Wall. Cat. n. 3637 = Decaspermum paniculatum.
cuspidifolia, *DC. Prod.* iii. 279.—Bras.
cyclophylla, *Baker, in Journ. Bot.* xx. (1882) 111.—Madag.
cyclophylla, *Berg, in Mart. Fl. Bras.* xiv. i. 287.—Bras.
cyclophylla, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 494.—Zeylan.
cydoniæfolia, *Berg, in Mart. Fl. Bras.* xiv. i. 229.—Bras.
cylindrica, *Wight, Ic.* t. 527; *Illustr.* ii. 14.—Zeylan.
cymatodes, *Berg, in Mart. Fl. Bras.* xiv. i. 278.—Bras.
cymosa, *Lam. Encyc.* iii. 199.—Ins. Maurit.
cymosa, Roxb. Hort. Beng. [97]; Fl. Ind. ii. 492 = grandis.
cymosa, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = mascarensis.
cyphophloea, *Griseb. Cat. Pl. Cub.* 87.—Cuba.
Dallachyana, *F. Muell. ex Benth. Fl. Austral.* iii. 287.—Austral.
Darwinii, *Hook. f. Fl. Antarct.* 277.—Chili.
dasyantha, *Berg, in Mart. Fl. Bras.* xiv. i. 272.—N. Granat.
decora, Salisb. Prod. 353 = E. Jambos.
decora, *Vell. Enum. Pl. Zeyl.* 115.—Zeylan.
decora, Wall. Cat. n. 3606 = javanica.
decorticosa, *Berg, in Mart. Fl. Bras.* xiv. i. 240.—Bras.
decumbens, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 339.—Bras.
deflexa, Polr. Encyc. Suppl. iii. 124 = Myrcia deflexa.
delicatula, *DC. Prod.* iii. 278.—Bras.
densiflora, *DC. l. c.* 287.—Malaya.
densiflora, Wild. ex Berg, in Linnæa, xxvii. (1854) 247 = follosa.
depauperata, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 346.—Bras.
deserti, *Cambess. l. c.* 343.—Bras.
Desvauxiana, *Berg, in Linnæa*, xxvii. (1854) 196.—Ind. occ.
diantha, *Berg, in Mart. Fl. Bras.* xiv. i. 239.—Bras.
dichotoma, *DC. Prod.* iii. 278.—Am. bor.; Ind. occ.
dichroma, *Berg, in Mart. Fl. Bras.* xiv. i. 290.—Bras.
dicrana, *Berg, in Linnæa*, xxvii. (1854) 259.—Florida.
dicrocea, *Berg, in Mart. Fl. Bras.* xiv. i. 291.—Bras.
dictyophleba, *Berg, l. c.* 236.—Bras.
dimorpha, *Berg, l. c.* 264.—Reg. Argent.
diospyrifolia, *Wall. Cat.* n. 3617.—Reg. Himal.
dipoda, DC. Prod. iii. 262 = paniculosa.
discolor, *DC. l. c.* 277.—Reg. Andin.
discolorans, *Wright, in Sauv. Fl. Cub.* 41.—Cuba.
disperma, *Vell. Fl. Flum.* 209; v. t. 98.—Bras.
distans, *Berg, in Mart. Fl. Bras.* xiv. i. 262.—Bras.
distichæ, *DC. Prod.* iii. 274.—Jamaic.
distonæ, *Berg, in Linnæa*, xxvii. (1854) 185.—Bras.
dittocarpa, *Berg, in Mart. Fl. Bras.* xiv. i. 262.—Bras.
divaricata, *Benth. in Hook. Journ. Bot.* ii. (1840) 319.—Guian.
divaricata, *Berg, in Linnæa*, xxvii. (1854) 257.—Chili.

EUGENIA:—
diversiorata, Lam. Encyc. iii. 209 = Myrcia diversiorata.
diversicolor, Willd. ex Berg, in Linnæa, xxvii. (1854) 199 = barnensis.
diversiflora, *Berg, l. c.* xxx. (1859–60) 675.—Bras.
diversifolia, *Bruguv. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 180.—N. Caled.
Djonat, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 116. —Ins. Philipp.
dodonæifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 364.—Bras.
Dombeyana, *DC. Prod.* iii. 276.—Peruv.
domestica, Baill. Hist. Pl. vi. 245 = malaccensis.
domingensis, *Berg, in Linnæa*, xxvii. (1854) 296.— Ind. occ.
Doniana, Berg, in Mart. Fl. Bras. xiv. i. 643, Index = Myrcia Doniana?
Duartiana, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 348. —Bras.
Duchassaingiana, *Berg, in Linnæa*, xxvii. (1854) 225. —Ind. occ.
dulcis, *Berg, in Mart. Fl. Bras.* xiv. i. 256.—Bras.
dumetorum, *DC. Prod.* iii. 284.—Cochinch.
dumosa, *DC. l. c.* 276.—Am. calid.
Dupontii, *Baker, Fl. Maurit.* 116.—Ins Mascar.
dysantha, *Benth. Pl. Hartw.* 178.—N. Granat.
dysenterica, *DC. Prod.* iii. 268.—Bras.
edulis, *Benth. & Hook. f. ex Griseb. in Goett. Abh.* xxiv. (1879) 126.—Reg. Argent.
edulis, *Vell. Fl. Flum.* 208; v. t. 84.—Bras.
effusa, *A. Gray, Bot. U. St. Exspl. Exped.* i. 524.—Ins. Pacific.
egensis, *DC. Prod.* iii. 281.—Bras.
Ehrenbergiana, *Berg, in Linnæa*, xxvii. (1854) 153.— Ind. occ.
elegans, *Berg, in Mart. Fl. Bras.* xiv. i. 232.—Reg. Argent.
elliptica, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 323.—Uruguay.
elliptica, Lam. Encyc. iii. 206 = ostinifolia.
elliptica, *Phil. in Anal. Univ. Chil.* (1856) ii. 315.— Chili.
elliptica, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = venosa.
elliptica, Sm. in Trans. Linn. Soc. iii. (1797) 281 = Smithii.
elliptica, Wall. Cat. n. 3587 = valdevenosa.
emarginata, *DC. Prod.* iii. 280.—Venezuela.
emarginata, *Vahl, in West, Bidr. Ste-Croix*, 290.— Ind. occ.
enumenaia, *Baker, in Journ. Linn. Soc.* xx. (1883) 145.—Madag.
erythrocarpa, *DC. Prod.* iii. 270.—N. Granat.
erythroxyloides, *Mart. ex Berg, in Mart. Fl. Bras.* xiv. i. 246.—Bras.
Eschholtziana, *Berg, l. c.* 582.—Bras.
esculenta, *Berg, in Linnæa*, xxvii. (1854) 278.—Ins. S. Doming.
estrellensis, *Berg, in Mart. Fl. Bras.* xiv. i. 250.— Bras.
eucalyptoides, *F. Muell. Fragm.* iv. 55.—Austral.
eucosma, *Berg, in Mart. Fl. Bras.* xiv. i. 233.—Bras.
eurychilla, *Berg, in Linnæa*, xxvii. (1854) 213.— Guian.
exalata, *Rich. ex Berg, l. c.* xxx. (1859–60) 687.— Ind. occ.
excelsa, *Berg, in Mart. Fl. Bras.* xiv. i. 277.—Bras.
excoriata, *Berg, l. c.* 213.—Bras.
exechuas, *Berg, l. c.* 268.—Bras.
expallens, *Berg, l. c.* 260.—Bras.
expansa, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 491.— Penang.
expansa, *Spring, ex Mart. in Flora*, xx. (1837) *Beibl.* ii. 84.—Bras.
exsucca, *DC. Prod.* iii. 278.—Peruv.
fallax, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Myrcia fallax.
faramoides, *A. Rich. Ess. Fl. Cub.* 588.—Cuba.
fascicularis, Wall. Cat. n. 3923 = brasiensis.
fascicularifera, *Berg, in Linnæa*, xxvii. (1854) 238.— Guian.
Fehri, *Berg, in Mart. Fl. Bras.* xiv. i. 238.—Bras.
Feliebertii, *DC. Prod.* iii. 266.—Bras.
Fendleana, *Berg, in Mart. Fl. Bras.* xiv. i. 238.—Bras.
ferruginea, Bass. in C. Gay, Fl. Chil. ii. 392 = Myrtus ferruginea.
Ferreiresana, *Berg, in Mart. Fl. Bras.* xiv. i. 265.— Bras.

EUGENIA:—
ferruginea, *Hook. f. Arn. in Hook. Bot. Misc.* iii. (1833) 319.—Chili.
ferruginea, Poir. Encyc. Suppl. iii. 194 = Myrcia ferruginea.
ferruginea, *A. Rich. Ess. Fl. Cub.* 591.—Cuba.
ferruginea, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = Psidium ferrugineum.
ferruginea, Wight, lc. t. 554 = E. Thwaria.
fetida, Poir. Encyc. Suppl. iii. 139 = foetida.
ficifolia, Wall. ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 476 = grandis.
Fieldingii, *Berg, in Linnæa*, xxix. (1857–58) 242.— Mexic.
filiformis, *Macfad. Fl. Jamaic.* ii. 116.—Jamaic.
filiformis, *Wall. ex Duthie, in Hook. f. Fl. Brit. Ind.* ii. 478.—Malaya.
fimbriata, *DC. Prod.* iii. 278.—Peruv.
firma, DC. Prod. iii. 282, partim = capparidifolia.
firma, Wall. Cat. n. 3603 = grandis.
flamingensis, *Berg, in Mart. Fl. Bras.* xiv. i. 276.— Bras.
flava, *Berg, l. c.* 278.—Bras.
flavescutata, *DC. Prod.* iii. 272.—Bras.
flavo-nigra, *Rich. ex Berg, in Linnæa*, xxx. (1859–60) 691.—Ind. occ.
flavo-virens, *Berg, l. c.* xxvii. (1854) 184.—Ind. occ.
flocciferæ, *Thw. Enum. Pl. Zeyl.* 115.—Zeylan.
floccosa, *Bedd. Fl. Sylv.* i. t. 200.—Ind. or.
floribunda, Spreng. ex DC. Prod. iii. 279 = confusa.
floribunda, *West. Bidr. Ste-Croix*, 216; *ex Willd. Sp. Pl.* ii. 960.—Ins. S. Crucis.
florida, *DC. Prod.* iii. 288.—Bras.
fluminensis, *Berg, in Mart. Fl. Bras.* xiv. i. 265.— Bras.
foetida, *Pers. Syn.* ii. 29.—Guian.
foliosa, *DC. Prod.* iii. 277.—Am. trop.
formosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 354.— Bras.
formosa, *Wall. Pl. As. Rar.* ii. 6. t. 108.—Ind. or.; Ind. occ.
forsteri, *F. Muell. Fragm.* v. 13 = grandis.
foveolata, *Berg, in Mart. Fl. Bras.* xiv. i. 261.—Bras.
fragrans, *Willd. Sp. Pl.* ii. 964.—Ind. occ.
Francavillana, *Berg, in Linnæa*, xxx. (1859–60) 686. —Bras.
franciscensis, *Berg, in Mart. Fl. Bras.* xiv. i. 238.— Bras.
Freireana, *Berg, l. c.* 305.—Bras.
Friedrichsthalii, *Hemsl. Biol. Centr. Am. Bot.* i. 410. —Guatemala.
frondosa, *Wall. Cat.* sub n. 3890.—Reg. Himal.
frutescens, Wall. Cat. n. 3560 G = E. Jambolana.
fruticosa, *Roxb. Hort. Beng.* [87]; *Fl. Ind.* ii. 487.— Reg. Himal.; Burma.
fruticulosa, *DC. Prod.* iii. 270.—Bras.
fulgipes, *Berg, in Mart. Fl. Bras.* xiv. i. 233.—Bras.
fulva, *Thw. Enum. Pl. Zeyl.* 115.—Zeylan.
fulvescens, *Mart. ex DC. Prod.* iii. 283.—Bras.
fulvipes, *Zagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 189.—Guian.
Funciana, *Berg, in Linnæa*, xxix. (1857–58) 238.— Am. austr.
fusca, *Berg, in Mart. Fl. Bras.* xiv. i. 290.—Bras.
fusiformis, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 479.— Malaya.
Gacognei, *Montr. in Mém. Acad. Lyon*, x. (1860) 208. —N. Caled.
Gardneri, *Bedd. For. Man. Bot.* 108.—Ind. or.
Gardneriana, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 85 = Cuminglana.
Gardneriana, *Berg, in Mart. Fl. Bras.* xiv. i. 216.— Bras.
Gaudichaudiana, *Berg, l. c.* 238.—Bras.
Gayana, *Barn. in C. Gay, Fl. Chil.* ii. 390.—Chili.
geminiflora, *Berg, in Mart. Fl. Bras.* xiv. i. 228.— Bras.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 320.—Chili.
glabrata, *DC. Prod.* iii. 276.—Ins. S. Doming.
glandulifera, Roxb. Hort. Beng. [87]; Fl. Ind. ii. 496 = zeylanica.
glandulosa, *Blanco, Fl. Filip.* ed. I. 417.—Ins. Philipp.
glandulosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 383.—Bras.

EUGENIA:—
glaucea, *Berg, in Mart. Fl. Bras.* xiv. i. 279.—Bras.
glaucescens, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 368.—Bras.
glomerata, *Lam. Encyc.* iii. 199.—Ins. Mascar.
glomerata, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = Calyptranthes Spyglium.
glomerata, *Spring, ex Mart. in Flora*, xx. (1837) *Beibl.* ii. 82.—Bras.
Gomeziana, *Berg, in Mart. Fl. Bras.* xiv. i. 254.— Bras.
gracilenta, *Hance, in Journ. Bot.* xxiii. (1885) 7.— China.
gracilipes, *A. Gray, Bot. U. St. Expl. Exped.* i. 513.— Ins. Fiji.
gracilis, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 46 = Moonlana.
gracilis, *Berg, in Mart. Fl. Bras.* xiv. i. 222.—Bras.
grandiflora, *Berg, in Linnæa*, xxix. (1857–58) 236.— Venezuela.
grandifolia, *Berg, in Mart. Fl. Bras.* xiv. i. 224.— Bras.
grandis, *Wight, Illustr.* ii. 17.—As. et Austral. trop.
granulata, *Berg, in Linnæa*, xxvii. (1854) 200.— Cuba.
grata, *Wight, Illustr.* ii. 15.—Ind. or.; Burma.
Gray, Sem. *Mus. Vit.* 498; *Fl. Vit.* 79.—Ins. Fiji.
Greggii, *Poir. Encyc. Suppl.* iii. 196; *DC. Prod.* iii. 280.—Ind. occ.
Griffithii, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 481.— Malacca.
Grijsii, *Hance, in Journ. Bot.* ix. (1871) 5.—China.
Guabiju, *Berg, in Mart. Fl. Bras.* xiv. i. 251.—Reg. Argent.
guadalupensis, *DC. Prod.* iii. 275.—Ind. occ.
Guapurium, *DC. l. c.* 278.—Peruv.
Guayabillo, *A. Rich. Ess. Fl. Cub.* 590.—Cuba.
guayequinia, *DC. Prod.* iii. 275.—Ecuador.
Guayavillo, *Benth. Pl. Hartw.* 174.—N. Granat.
Gudilla, *Barn. in C. Gay, Fl. Chil.* ii. 396.—Chili.
Guillauti, *Sond. in Harv. & Sond. Fl. Cap.* ii. 528.— Afr. austr.
guianensis, *Aubl. Pl. Gui.* i. 506.—Guian.
Hasskelliana, *Trim. in Journ. Bot.* xxiii. (1885) 207.— Zeylan.
Hallii, *Berg, in Linnæa*, xxvii. (1854) 250.— Ecuador.
Heckelii, *Pancher & Sebert, Not. Bois Nouv. Calédon.* 259.—N. Caled.
Helferi, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 480.— Burma; Malaya.
hemilampra, *F. Muell. Fragm.* ix. 45.—Austral.
hemispherica, *Wight, Illustr.* ii. 14.—Ind. or.
Henryi, *Hance, in Journ. Bot.* xxiii. (1885) 7.—China.
herbacea, *Berg, in Mart. Fl. Bras.* xiv. i. 270.—Bras.
Heritieriana, Steud. Nom. ed. II. i. 603 = L'Heritierana.
heterocilia, *Tuss. Fl. Antill.* i. 98.—Ind. occ.
heteropbylla, *A. Rich. Ess. Fl. Cub.* 586.—Cuba.
Heyneana, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 500.— Ind. or.
Hilariana, *DC. Prod.* iii. 269.—Bras.
hirsuta, *Ruiz & Pav. Fl. Per.* iv. t. 417.—Peruv.
hirta, *Berg, in Mart. Fl. Bras.* xiv. i. 274.—Bras.
Hodgkinsoniæ, *F. Muell. Fragm.* ix. 145.—Austral.
Hoffmannseggii, *Berg, in Mart. Fl. Bras.* xiv. i. 315 —Bras.
Hozuel, *Sonn. Fl. Vit.* 76.—N. Caled.
Hookeri, *Steud. Nom.* ed. II. i. 608.—Chili.
horizontalis, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 179.—N. Caled.
humifusa, Phil. in Linnæa, xxviii. (1856) 689 (= *Myrteola humifusa*).—Chili.
humilis, *Phil. l. c.* 689.—Chili.
humilis, Vell. Fl. Flum. 210; v. t. 43 = hypericifolia.
hyemalis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 360. —Bras.
hypericifolia, *Gærtn. in Hook. Lond. Journ. Bot.* ii. (1843) 384.—Bras.
hypoleuca, Thw. ex Bedd. For. Man. Bot. 112 = ostinifolia.
hypogæatica, F. Muell. Fragm. v. 15 = cariocoides.
ibitipocensis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 365.—Bras.
illawarrie, Berg, in Mart. Fl. Bras. xiv. i. 233 = maschima.
imbricata, *Berg, l. c.* 586.—Bras.
impunctata, *Berg, l. c.* 289.—Bras.

EUGENIA:—
- inaequilobe, *DC. Prod.* iii. 282.—Guiana.
- inaequicosta, *Benth. in Hook. Journ. Bot.* ii. (1840) 231.—Guiana.
- inocarpa, *DC. Prod.* iii. 264.—Bras.
- inophylla, *Roxb. Hort. Beng.* [37]; *Fl. Ind.* ii. 496.—Ins. Moluce.
- inophyloides, *A. Gray, Bot. U. St. Expl. Exped.* i. 231. —Ins. Samoa.
- insignis, *Thw. Enum. Pl. Zeyl.* 416.—Ceylan.
- insipida, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 360.— Bras.
- *insipida*, Mart. in Flora, xx. (1837) ii. Beibl. 85 = prasina.
- insularis, *Berg, in Linnaea*, xxix. (1857-58) 288.— Ind. occ.
- intermedia, *Berg, l. c.* xxvii. (1854) 258.—Ins. S. Doming.
- inundata, *DC. Prod.* iii. 280.—Bras.
- involucrata, *DC. l. c.* 264.—Bras.
- iraxnensis, *Hemsl. Biol. Centr. Am. Bot.* i. 410.— Costa Rica.
- irirensis, *Berg, in Mart. Fl. Bras.* xiv. 1. 586.—Bras.
- ischnoscelos, *Berg, l. c.* 309.—Bras.
- isocalmensis, *Berg, l. c.* 258.—Bras.
- ithypensis, *Berg, in Mart. Fl. Bras.* xiv. 1. 580.— Bras.
- itajurensis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 340. —Bras.
- itapemirimensis, *Cambess. l. c.* 346.—Bras.
- jamaicensis, *Berg, in Linnaea*, xxvii. (1854) 237.— Jamaic.
- jambolides, *Wender. ex Steud. Nom.* ed. II. i. 603.— Hab. ?
- Jambolana, *Lam. Encyc.* iii. 198.—As. et Austral. trop.
- *Jambolana*, Willd. ex Berg, in Linnaea, xxvii. (1854) 98 = Myrcia fallax.
- *jambolifera*, Roxb. ex Wight & Arn. Prod. 329 = E. Jambolana.
- *Jambos*, Roxb. Hort. Beng. [37], nomen = E. Jambos.
- *Jambos*, Blanco, Fl. Filip. ed. I. 416 = malaccensis.
- Jambos, *Linn. Sp. Pl.* 470.—As. trop.
- *jambos*, Crantz, Inst. ii. 201 = E. Jambos.
- jambosoides, *Wright, ex Griseb. Cat. Pl. Cub.* 89.— Cuba.
- janeirensis, *Berg, in Mart. Fl. Bras.* xiv. 1. 578.— Bras.
- javanica, *Lam. Encyc.* iii. 200.—Malaya.
- *javanica*, Lam. l. c. partim = squea.
- jequitinhonhensis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 352.—Bras.
- Jostinia, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 500.— Ind. or.
- jucunda, *F. Muell. ex Benth. Fl. Austral.* iii. 251.— Austral.
- Jungkhuhniana, *Miq. Fl. Ind. Bat.* i. 1. 441.—Java.
- kalahiensis, *Miq. Anal. Bot. Ind.* i. 28.—Borneo.
- Karsteniana, *Berg, in Linnaea*, xxvii. (1854) 277.— Venezuela.
- Karwinskyana, *Berg, l. c.* xxix. (1857-58) 244.— Mexic.
- khasiana, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 491.— Reg. Himal.
- Klotzschiana, *Berg, in Mart. Fl. Bras.* xiv. 1. 255.— Bras.
- Kochiana, *DC. Prod.* iii. 268.—Bras.
- Korthalsiana, *Miq. Anal. Bot. Ind.* i. 28.—Borneo.
- Kunthiana, *DC. Prod.* iii. 264.—Bras.
- Kurzii, *Duthie, ex Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 68.—Reg. Himal.; Burma.
- laeta, *Buch.-Ham. in Mem. Wern. Soc.* v. (1832) 338. —Ind. or.
- laevicaulis, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 492. —Malacca.
- *laevigata*, DC. Prod. iii. 283 (= *Eugeniopsis laevigata*). —Bras.
- laevigata, *Miq. Anal. Bot. Ind.* i. 25.—Borneo.
- laevis, *Berg, in Linnaea*, xxvii. (1854) 177.—Ins. S. Doming.
- Lambertiana, *DC. Prod.* iii. 270.—Ind. occ.
- lancea, *Poir. Encyc. Suppl.* iii. 123.—Ins. S. Thomas.
- *lanceaefolia*, Roxb. Hort. Beng [37]; Fl. Ind. ii. 494 = Wallichii.
- lanceolaris, *Rumph. Il.* cc.—Ind. or.
- lanceolata, *Berg, in Mart. Fl. Bras.* xiv. 1. 276.— Bras.
- *lanceolata*, Lam. Encyc. iii. 200 = Wightiana.

EUGENIA:—
- lancifolia, *Miq. Anal. Bot. Ind.* i. 17.—Borneo.
- Langsdorffii, *Berg, in Mart. Fl. Bras.* xiv. 1. 568.— Bras.
- Larsotteana, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 350.—Bras.
- laterifiora, *Willd. Sp. Pl.* ii. 749.—Ins. S. Crucis.
- *laterifolia*, Pers. ex DC. Prod. iii. 278 = ecostifiora.
- latifolia, *Aubl. Pl. Gui.* i. 502. t. 199.—Guiana.
- *latifolia*, Mart. ex Berg, in Mart. Fl. Bras. xiv. 1. 294 = platyphylla.
- *latifolia*, Spreng. ex DC. Prod. iii. 266 = barueusis.
- laurifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 357. —Bras.
- *laurifolia*, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 489 = bifaria.
- *laurifolia*, Spring, ex Mart. in Flora, xx. (1837) ii. Beibl. 83 = Springiana.
- *laurina*, Willd. Sp. Pl. ii. 967 = Symplocos spicata.
- laxa, *DC. Prod.* iii. 268.—Bras.
- *laxiflora*, Poir. Encyc. Suppl. iii. 123 = Myrcia splendens.
- Leandraeana, *Berg, in Mart. Fl. Bras.* xiv. 1. 311.— Bras.
- lepidocarpa, *Wall. Cat.* n. 3618, partim.—Burma; Malaya.
- lepidota, *Berg, in Linnaea*, xxvii. (1854) 226.—Costa Rica.
- leptanthe, *Benth. in Hook. Journ. Bot.* ii. (1840) 231. —Guian.
- leptantha, *Wight, Ic.* t. 528; *Illustr.* ii. 15.—Malaya; Austral.
- leptoclada, *Berg, in Mart. Fl. Bras.* xiv. 1. 248.— Bras.
- leptocarpa, *Roxb. Hort. Beng.* [92]; *Fl. Ind.* ii. 495. —Ins. Romoa.'
- *leptospermoides*, Barn. in C. Gay, Fl. Chil. ii. 386 = parvifolia.
- leptospermoides, *DC. Prod.* iii. 266.—Chili.
- leptostemon, *Miq. Fl. Ind. Bat.* i. 1. 442.— Borneo.
- Leucadendron, *Berg, in Linnaea*, xxvii. (1854) 202.— Costa Rica.
- leucanthera, *Berg, in Mart. Fl. Bras.* xiv. 1. 277.— Bras.
- leucocalyon, *Miq. Anal. Bot. Ind.* i. 26.—Borneo.
- L'Heritieriana, *DC. Prod.* iii. 272.—Ind. occ.
- Lhotskyana, *Berg, in Mart. Fl. Bras.* xiv. 1. 305.— Bras.
- *ligustrina*, Cambess. in St. Hil. Fl. Bras. Mer. ii. 364, partim = myrtifolia, *Cambess.*
- ligustrina, *Miq. in Linnaea*, xvii. (1849) 529.—Bras.
- ligustrina, *Willd. Sp. Pl.* ii. 962.—Ind. occ.
- limbata, *DC. Prod.* iii. 278.—Peruv.
- limbosa, *Berg, in Linnaea*, xxvii. (1854) 294.— Peruv.
- Lindbergiana, *Berg, l. c.* xxx. (1859-60) 685.—Bras.
- Lindeniana, *Berg, l. c.* xxix. (1857-58) 240.—Bras.
- Lindleyana, *DC. Prod.* iii. 278.—Am. austr.
- linearifolia, *Berg, in Mart. Fl. Bras.* xiv. 1. 280.— Bras.
- linearis, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 486. —Ind. or.
- linearis, *Rich. ex Berg, in Linnaea*, xxx. (1859-60) 678.—Ind. occ.
- linearis, *DC. Prod.* iii. 278.—Ind. occ.
- lineata, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 487.— Malaya.
- Lingua, *Berg, in Mart. Fl. Bras.* xiv. 1. 266.—Bras.
- linguaeformis, *Berg, l. c.* 268.—Bras.
- lissophylla, *Bedd. For. Man. Bot.* 108.—Ind. or.
- litoralis, *Pauch. ex Brongn. et Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 178.—N. Caled.
- livida, *Berg, in Mart. Fl. Bras.* xiv. 1. 584.—Bras.
- loasaleuricoides, *Baker, in Journ. Linn. Soc.* xxi. (1884) 341.—Madag.
- *lonchophylla*, Voigt, Hort. Suburb. Calc. 49 = lanceolaria.
- longifolia, *DC. Prod.* iii. 269.—Bras.
- longipes, *Berg, in Linnaea*, xxvii. (1854) 150.— Florida.
- lucida, *Banks, ex Gaertn. Fruct.* i. 167.—Hab. ?
- lucida, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 356.— Bras.
- *lucida*, Lam. Encyc. iii. 202 = cotinifolia.
- *lucidula*, Miq. Fl. Ind. Bot. i. 1. 444 = polyantha.
- luschnoeth, *Bert. ex DC. Prod.* iii. 260.—Ins. S. Doming.

EUGENIA:—
- lugens, *Berg, in Linnaea*, xxvii. (1854) 299.—Peruv.
- *Luma*, Berg, l. c. 251 = apiculata.
- *Lumilla*, Phil. in Bot. Zeit. xiv. (1856) 648 = Myrtus ferruginea.
- Luschnathiana, *Klotzsch, ex Berg, in Mart. Fl. Bras.* xiv. 1. 338.—Bras.
- lutescens, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 341. —Bras.
- mabaeoides, *Wight, Illustr.* ii. 18.—Zeylan.
- macahensis, *Berg, in Mart. Fl. Bras.* xiv. 1. 589.— Bras.
- macarensis, *Berg, in Linnaea*, xxvii. (1854) 288.— Venezuela.
- macrantha, *Berg, in Mart. Fl. Bras.* xiv. 1. 301.— Bras.
- macrocalyx, *Mart. ex Berg, l. c.* 332.—Bras.
- *macrocarpa*, Cham. & Schlecht. in Linnaea, xiii. (1839) 417 = mexicana.
- macrocarpa, *Roxb. Hort. Beng.* [37]; *Fl. Ind.* ii. 497. —Ind. or.; Malaya.
- macrocarpa, *Schlecht. in Linnaea*, v. (1830) 560.— Mexic.
- macrocalada, *Berg, in Mart. Fl. Bras.* xiv. 1. 248.— Bras.
- macrophylla, *Berg, l. c.* 268.—Bras.
- *macrophylla*, Lam. Encyc. iii. 196 = malaccensis.
- macrorhyncha, *Miq. Anal. Bot. Ind.* i. 21.—Borneo.
- macrosepala, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 501.—Ind. or.
- macrosperma, *DC. Prod.* iii. 277.—Bras.
- macrostemon, *Berg, in Linnaea*, xxvii. (1854) 188.— Cuba.
- maculata, *Berg, in Mart. Fl. Bras.* xiv. 1. 297.— Bras.
- magnifica, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 178.—N. Caled.
- magnifica, *Spring, ex Mart. in Flora*, xx. (1837) ii. *Beibl.* 66.—Bras.
- Maingayi, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 484. —Malacca.
- Maire, *A. Cunn. in Ann. Nat. Hist.* iii. (1839) 114 = N. Zel.
- malabarica, *Bedd. Fl. Sylv.* i. t. 199.—Ind. or.
- *malaccensis*, Blanco, Fl. Filip. ed. I. 415 = E. Jambos.
- malaccensis, *Linn. Sp. Pl.* 470.—As. trop.
- *malaccensis*, Lour. Fl. Cochinch. 306 = squea.
- *malaccensis*, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 96 = javanica.
- *maleolens*, Pers. Syn. ii. 29 = foetida.
- malpighioides, *DC. Prod.* iii. 276.—Bras.
- *Manangaqui*, Blanco, Fl. Filip. ed. II. 290 = javanica.
- mandioccana, *Berg, in Mart. Fl. Bras.* xiv. 1. 585.— Bras.
- mangifolia, *Wall. Cat.* n. 3619.—Reg. Himal.
- Manzani, *Berg, in Mart. Fl. Bras.* xiv. 1. 323.—Bras.
- margaritaceata, *G. Don, Gen. Syst.* ii. 854.—Bras.
- Marchiana, *Griseb. Fl. Brit. W. Ind.* 238.—Ind. occ.
- marginata, *Pers. Syn.* ii. 17.—Ins. S. Doming.
- mariquitensis, *Berg, in Linnaea*, xxvii. (1854) 267.— N. Granat.
- maritima, *Benn. in C. Gay, Fl. Chil.* ii. 390.—Chili.
- maritima, *DC. Prod.* iii. 271.—Bras.
- *maritima*, DC. l. c. 283 = scapulicosta.
- marowynensis, *Miq. Stirp. Surin. Sel.* 39.—Guian.
- Martini, *Berg, in Linnaea*, xxvii. (1854) 230.—Ind. occ.
- Martiusiana, *DC. Prod.* iii. 269.—Bras.
- mascarensis, *Proel, in Obss. Isis*, xxi. (1828) 274.— Ins. Mauric.
- maschalantha, *Berg, in Mart. Fl. Bras.* xiv. 1. 278.— Reg. Argent.
- matanzasis, *Berg, in Linnaea*, xxvii. (1854) 285.— Cuba.
- Mato, *Griseb. in Gottt. Abh.* xxiv. (1879) 128.—Reg. Argent.
- Maximiliana, *Berg, in Mart. Fl. Bras.* xiv. 1. 250.— Bras.
- Maximiliana, *DC. Prod.* iii. 270.—Bras.
- mayensis, *Berg, in Mart. Fl. Bras.* xiv. 1. 312.— Bras.
- megalopetala, *Griseb. Cat. Pl. Cub.* 89.—Cuba.
- Meissneana, *Sagot, in Ann. Sc. Nat. Bot.* VI. xi. (1880) 194.—Guian.
- membranacea, *Berg, in Mart. Fl. Bras.* xiv. 1. 589.— Bras.
- *memoryoides*, *Benth. in Hook. Niger Fl.* 389.—Afr. trop.

EUGENIA :—
mespiloides, *Lem. Encyc.* ii. 908.—Ins. Maurit.
mexicana, *Steud. Nom.* ed. II. i. 608.—Mexic.
Meyeriana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 588.—Bras.
Michelii, Lam. Encyc. iii. 908 = uniflora.
micrantha, *Bedd. For. Man. Bot.* 108.—Zeylan.
micrantha, *Bertol. Fl. Guatimal.* 93.—Guatemala.
micrantha, *DC. Prod.* iii. 283.—N. Granat.
microcarya, *Miq. Anal. Bot. Ind.* i. 37.—Borneo.
microcalyx, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 493.—Malacca.
microcarpa, *Berg*, in *Mart. Fl. Bras.* xiv. i. 574.—Bras.
microcarpos, Lam. Encyc. iii. 201 = Myrcia Mini, splendens.
microcepala, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 108.—Bras.
microphylla, Abel, Narr. Journ. China, 364 = pyxophylla !
microphylla, *Bedd. For. Man. Bot.* 110.—Ind. or.
microphylla, *A. Rich. Ess. Fl. Cub.* 584.—Cuba.
microphylla, *Rich. ex Berg*, in *Linnaea*, xxx. (1859–60) 678.—Ins. S. Doming.
micropoda, *Baker*, in *Journ. Linn. Soc.* xx. (1883) 148.—Madag.
micropora, *DC. Prod.* iii. 269.—Bras.
Mieridana, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 108.—Bras.
Mikaniana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 297.—Bras.
Mikaniana, DC. Prod. iii. 283 = Myrcia Mikaniana.
mikanioides, *Berg*, in *Mart. Fl. Bras.* xiv. i. 298.—Bras.
Mini, Aubl. Pl. Gui. i. 498 = Myrcia Mini.
Mini, Mart. ex DC. Prod. iii. 283 = Myrcia Pseudo-Mini.
Mini, Willd. ex Berg, in Mart. Fl. Bras. xiv. i. 315 = Hoffmanneggii.
minima, *Blume*, *Cat. Gew. Buitenz.* 75.—Java.
minutiflora, *Hance*, in *Journ. Bot.* ix. (1871) 5.—China.
modesta, *DC. Prod.* iii. 278.—Bras.
modesta, Phil. in Linnaea, xxvii. (1856) 628 = proba.
Mollitana, *Berg*, in *Linnaea*, xxvii. (1854) 295.—Am. trop.
molliconia, *Mart. ex Berg*, *l. c.* 313.—Bras.
mollis, Willd. ex Berg, i. c. 121 = Myrcia Humboldtiana.
monosperma, Vell. Fl. Flum. 209; v. t. 49 = compactiflora.
montana, Aubl. Pl. Gui. i. 495 = fragrans.
montana, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 365.—Bras.
montana, *Wight*, *Ic.* iii. t. 1060.—Ind. or.
monticola, *DC. Prod.* iii. 276.—Am. bor.; Ind. occ.
Moonlana, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 535.—Bras.
Moonlana, *Wight*, *Illustr.* ii. 13.—Zeylan.
moonlodes, *Berg*, in *Mart. Fl. Bras.* xiv. i. 262.—Bras.
Moorei, F. Muell. Fragm. v. 33 = E. jambolana.
moraviana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 304.—Bras.
Moritziana, *Karst. Ausw. Gew. Venez.* 18. t. 6.—Venezuela.
mosquitensis, *Berg*, in *Linnaea*, xxxi. (1861–62) 257.—Am. centr.
mucronata, *Berg*, *l. c.* xxvii. (1854) 172.—Ind. occ.
mucronata, *Phil. in Anal. Univ. Chil.* (1868) ii. 227.—Chili.
Mueller, *Miq. Anal. Bot. Ind.* i. 23.—Borneo.
mugiensis, *Berg*, in *Mart. Fl. Bras.* xiv. i. 271.—Bras.
multiflora, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 361.—Bras.
multiflora, *Bedd. & Arn. in Hook. Bot. Misc.* iii. (1833) 225.—Chili.
multiflora, Lam. Encyc. iii. 202 = Myrcia multiflora.
multiflora, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 110.—Guian.
Munroii, Miq. Anal. Bot. Ind. i. 18 = seq.
Munroii, *Wight*, *Illustr.* ii. 14.—Ind. or.
muricata, *DC. Prod.* iii. 283.—Bras.
myrtifolia, *Berg*, in *Mart. Fl. Bras.* xiv. i. 311.—Bras.
myrobistida, *Hance*, in *Journ. Bot.* xxiii. (1885) 6.—China.
myrobistida, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 358.—Bras.
myrtophylla, *Casaretti*, *Dec.* 77.—Bras.

EUGENIA :—
myristicifolia, *Sagot*, in *Ann. Sc. Nat. Sér. VI.* xx. (1885) 194.—Guian.
myrobalana, *DC. Prod.* iii. 277.—Bras.
myrtifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 367.—Bras.
myrtifolia, *Rosb. Hort. Beng.* [87]; *Fl. Ind.* ii. 490.—Ind. or.; Malaya.
myrtifolia, Sallab. Prod. 258 = uniflora.
myrtifolia, *Roxb. Fl. Ind. Mag.* v. 2980.—Austral.
myrtillifolia, *DC. Prod.* iii. 268.—Bras.
myrtoides, *Bronga. & Gris*, in *Bull. Soc. Bot. Fr.* xii. (1865) 180.—N. Caled.
myrtoides, *G. Don*, *Gen. Syst.* ii. 854.—Afr. trop.
myrtoides, Poir. Encyc. Suppl. iii. 125 = buxifolia, *Willd.*
naua, *Berg*, in *Mart. Fl. Bras.* xiv. i. 244.—Bras.
nataliia, *Sond. in Harv. & Sond. Fl. Cap.* ii. 522.—Afr. austr.
Neesiana, *Wight*, *Illustr.* ii. 15.—Zeylan.
neglecta, *Berg*, in *Mart. Fl. Bras.* xiv. i. 320.—Bras.
nemoralis, *DC. Prod.* iii. 267.—Bras.
nervosa, Bedd. For. Man. Bot. 106 = operculata.
nervosa, *Lour. Fl. Cochinch.* i. 308.—Cochinch.
neurocalyx, *A. Gray*, *Bot. U. St. Expl. Exped.* i. 512.—Ins. Pacific.
Nhanica, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 362.—Bras.
nigra, *DC. Prod.* iii. 268.—Bras.
nigrescens, Poir. Encyc. Suppl. iii. 123 = cymosa.
nitens, *DC. Prod.* iii. 285.—Ins. Maurit.
nitida, *Benth. in Hook. Journ. Bot.* ii. (1840) 322.—Guian.
nitida, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 349.—Bras.
nitida, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 496.—Malacca.
nitida, Vell. Fl. Flum. 208; v. t. 35 (= *Gomidesia Chamissoana*).—Bras.
nodiflora, Aubl. Pl. Gui. ii. 11. 140 = aquea !
nummularia, *Baker*, *Fl. Maurit.* 118.—Ins. Maurit.
nummularia, *Wright*, *ex Grisb. Cat. Pl. Cub.* 86.—Cuba.
nutans, *Berg*, in *Mart. Fl. Bras.* xiv. i. 299.—Bras.
oaxacana, *Berg*, in *Linnaea*, xxx. (1859–60) 683.—Mexic.
obcordata, *Raoul*, in *Ann. Sc. Nat.* Sér. III. ii. (1844) 193.—N. Zel.
oblata, *Roxb. Hort. Beng.* [87]; *Fl. Ind.* ii. 493.—Ind. or.; Malaya.
oblongum, *Berg*, in *Mart. Fl. Bras.* xiv. i. 302.—Bras.
oblongifolia, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 491.—Malacca.
obovata, *Berg*, in *Mart. Fl. Bras.* xiv. i. 289.—Bras.
obovata, *Poir. Encyc. Suppl.* iii. 124.—Ins. Maurit.
obovata, Wall. ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 491, sphalm. pro *Syzygium obovatum*) (Wall. Cat. 3552) = operculata.
obscura, *DC. Prod.* iii. 285.—Bras.
obtusa, *DC. l. c.* iii. 266.—Peruv.
obtusata, *Willd. ex Berg*, in *Linnaea*, xxvii. (1854) 182.—Jamaic.
obtusifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 365.—Bras.
obtusifolia, Roxb. Hort. Beng. [37]; *Fl. Ind.* ii. 485.—Jambolana.
obtusissima, Steud. Nom. ed. II. i. 608 = javanica.
obversa, *Berg*, in *Mart. Fl. Bras.* xiv. i. 295.—Bras.
obversa, *Miq. Anal. Bot. Ind.* i. 18.—Borneo.
ocsneasis, *Linden*, *ex Regel*, *Cat. Hort. Absehow.* 57.—N. Granat.
occlana, *Kurz*, in *Journ. As. Soc. Beng.* xlv. (1876) ii. 130.—Malaya.
ochra, *Berg*, in *Linnaea*, xxvii. (1854) 216.—Guian.
octopetala, Bech.-Ham. ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 493 = E. Thwaea.
odorata, *Berg*, in *Linnaea*, xxvii. (1854) 171.—Ins. S. Doming.
odorata, Wight, Illustr. ii. 16 = brachiata.
oeideocarpa, *Berg*, in *Mart. Fl. Bras.* xiv. i. 270.—Bras.
Oerstediana, *Berg*, in *Linnaea*, xxvii. (1854) 295.—Costa Rica.
oleaefolia, *DC. Prod.* iii. 270.—Am. trop.
oleina, Wight, Illustr. ii. 15 = myrtifolia.

EUGENIA :—
oleoides, *Planch. & Linden*, *Cat.* ix. (1854); *ex Bot. Exc.* xii. (1854) 264.—Hab. ?
oleosa, *F. Muell. Fragm.* v. 18.—Austral.
oligantha, *Bedd. For. Man. Bot.* 108.—Ind. or.
oligoneura, *Berg*, in *Mart. Fl. Bras.* xiv. i. 321.—Bras.
oligophylla, *Rich. ex Berg*, in *Linnaea*, xxx. (1859–60) 685.—Guian.
oliveana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 308.—Bras.
olivifolia, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 495.—Zeylan.
opaca, *Berg*, in *Mart. Fl. Bras.* xiv. i. 278.—Reg. Argent.
operculata, *Roxb. Hort. Beng.* [87]; *Fl. Ind.* ii. 486.—As trop.
orbicularis, *Berg*, in *Linnaea*, xxx. (1859–60) 678.—Ind. occ.
orbiculata, Lam. Encyc. iii. 202 = cotinifolia.
Orbignyana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 314.—Bras.
orcinoma, *Berg*, in *Linnaea*, xxvii. (1854) 158.—Costa Rica.
orignoides, *Berg*, *l. c.* xxix. (1857–58) 229.—Mexic.
orthostemon, *Berg*, *l. c.* xxvii. (1854) 179.—N. Granat.
Ottonis, *Berg*, *l. c.* 298.—Venezuela.
ovalifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 350.—Bras.
ovalis, *Berg*, in *Linnaea*, xxvii. (1854) 156.—Ecuador.
ovata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1888) 319.—Chili.
ovigera, *Bronga. & Gris*, in *Bull. Soc. Bot. Fr.* xii. (1865) 179.—N. Caled.
owariensis, *Beauv. Fl. Owar.* ii. 20. t. 70.—Afr. trop.
oxyphylla, *Berg*, in *Mart. Fl. Bras.* xiv. i. 251.—Bras.
paehnantha, *Berg*, *l. c.* 585.—Bras
pachyphylla, *Kurz*, in *Journ As. Soc. Beng.* xlii. (1873) ii. 282.—Burma; Malaya.
pacifica, *Benth. Bot. Voy. Sulph.* 98.—Ins. Cocos.
pallens, *DC. Prod.* iii. 284.—Amer.
paltera, Poir. Encyc. Suppl. iii. 122 = Calyptranthes Chytraculia.
pallida, *Berg*, in *Mart. Fl. Bras.* xiv. i. 281.—Reg. Argent.
palmiflora, *Panch. ex Brongn. & Gris*, in *Bull. Soc. Bot. Fr.* xii. (1865) 178.—N. Caled.
pamatensis, *Miq. Anal. Bot. Ind.* i. 22.—Borneo.
Pancheri, *Brongn. & Gris*, in *Bull. Soc. Bot. Fr.* xii. (1865) 180.—N. Caled.
Paniala, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 489 = operculata.
paniculaeflora, Steud. in Flora, xxvi. (1843) 762 = Myrcia Kegeliana.
paniculata, *Banks*, *ex Gaertn. Fruct.* i. 167; *Lam. Encyc.* iii. 199.—Ins. Borbon.
paniculata, *Bello*, in *Anal. Soc. Esp. Hist. Nat.* x. (1881) 271.—Porto Rico.
paniculata, Cambess. in St. Hil. Fl. Bras. Mer. ii. 338 (= *Eugeniopsis paniculata*).—Bras.
paniculata, Hilsenb. & Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = Myrcia ferruginea.
paniculata, Jacq. Coll. ii. 108 = fragrans.
pantagensis, *Berg*, in *Mart. Fl. Bras.* xiv. i. 246.—Bras.
papillosa, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 495.—Malacca.
paracatuana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 286.—Bras.
parafluta, *Berg*, *l. c.* 301.—Bras.
paranahybensis, *Berg*, *l. c.* 287.—Bras.
paraleuis, *Berg*, *l. c.* 288.—Bras.
Parkeri, *Baker*, in *Journ. Linn. Soc.* xx. (1883) 144.—Madag.
Parkeriana, *DC. Prod.* iii. 271.—Ins. Trinit.
parviflora, *Lam. Encyc.* iii. 200.—Java.
parviflia, *DC. Prod.* iii. 266.—Peruv.
patagonica, *Phil. in Linnaea*, xxxiii. (1864–65) 72.—Patagon.
patens, *Poir. Encyc. Suppl.* iii. 124.—Guian.
Patriti, *Vahl*, *Eclog. Am.* ii. 36.—Guian.
patula, *DC. Prod.* iii. 284.—Bras.
pauciflora, *DC. l. c.* 280.—Bras.
pauciflora, Wight, Illustr. ii. 14 = laeta.
pellucida, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 485.—Burma.

EUGENIA:—
pellucida, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 85 = cinnamomea.
penangiana, *Duthie*, in *Hook. f. Fl. Brit Ind.* ii. 486. —Ins. Penang.
pendens, *Duthie*, *l. c.* 475.—Malacca.
pendula, *DC. Prod.* iii. 284.—Java.
perforata, *Berg*, in *Linnæa*, xxvii. (1854) 236.—Bras.
periplocifolia, Jacq. Coll. ii. 108 = Myrcia splendens.
pernitifolia, *Berg*, in *Mart. Fl. Bras.* xiv. i. 226.—Bras.
phæa, Berg, l. c. 308 = disperma.
Phillippii, *Berg*, in *Linnæa*, xxvii. (1854) 145.—Bras.
phillyreoides, Willd. ex Berg, in Mart. Fl. Bras. xiv. i. 288 = vaga.
phillyreifolia, *Baker*, in *Journ. Linn. Soc.* xx. (1883) 145.—Madag.
phillyreoides, *Trim. in Journ. Bot.* xxiii. (1885) 207.—Zeylan.
phyllireæfolia, *A. Rich. Ess. Fl. Cub.* 585.—Cuba.
pia, *DC Prod.* iii. 263.—Bras.
piauhiensis, *Berg*, in *Mart. Fl. Bras.* xiv. i. 285.—Bras.
pilosela, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 357.—Bras.
Pimenta, DC. Prod. iii. 285 = Pimenta officinalis.
pinifolia, *Mart. ex Berg*, in *Mart. Fl. Bras.* xiv. i. 275.—Bras.
Piresiana, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 341.—Bras.
pisiformis, *Cambess. l. c.* 356.—Bras.
Pisonis, *Berg*, in *Mart. Fl. Bras.* xiv. i. 228.—Bras.
platacaefolia, *DC. Prod.* iii. 270.—Bras.
Pitra, *Berg*, in *Linnæa*, xxvii. (1854) 264.—Chili.
planipes, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 253.—Ins. Chiloe.
planiramea, *Berg*, in *Mart. Fl. Bras.* xiv. i. 234.—Bras.
platycalda, *Berg*, *l. c.* 257.—Bras.
platyphylla, *Berg*, *l. c.* 294.—Bras.
platysema, *Berg*, *l. c.* 276.—Bras.
pleurantha, *Berg*, *l. c.* 280.—Bras.
plicato-costata, *Berg*, *l. c.* 275.—Bras.
plicatula, *Wright*, in *Sauv. Fl. Cub.* 40.—Cuba.
pluriflora, *DC. Prod.* iii. 270.—Bras.
pluriflora, Mart. in Flora, xx. (1837) ii. Beibl. 88 = rotundifolia.
Poeppigiana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 282.—Bras.
Pohliana, *DC. Prod.* iii. 264.—Bras.
Poiretii, *DC. l. c.* 274.—Ins. S. Doming.
Poiteaui, *Berg*, in *Linnæa*, xxvii. (1854) 181.—Guian.
polita, Wall. Cat. sub n. 3606 = Wallichii.
polyadema, *Berg*, in *Linnæa*, xxxi. (1861–62) 258.—Peruv.
polyantha, *Phil. in Linnæa*, xxviii. (1856) 689.—Chili.
polyantha, Wall. Cat. sub n. 3608 = grandis.
polyantha, *Wight*, *Illustr.* ii. 17.—Burma; Java.
polycarpa, *Berg*, in *Mart. Fl. Bras.* xiv. i. 231.—Reg. Argent.
polyocephala, *Miq. Anal. Bot. Ind.* i. 19.—Borneo.
polygama, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 491 = Decaspermum paniculatum.
polypetala, *Wall. Cat.* n. 3616.—Reg. Himal.
polyphylla, *Berg*, in *Mart. Fl. Bras.* xiv. i. 240.—Bras.
polystachya, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 110.—Guian.
pomatensis, *Miq. Anal. Bot. Ind.* i. 22. t. 4.—Borneo.
populifolia, *Baker*, *Fl. Maurit.* 118.—Ins. Maurit.
porphyrocalda, *Berg*, in *Linnæa*, xxvii. (1854) 266.—Chili; Peruv.
portoricensis, *DC. Prod.* iii. 266.—Porto Rico.
praecox, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 488 = Wallichii.
prasina, *Berg*, in *Mart. Fl. Bras.* xiv. i. 235.—Bras.
Prianri, *Berg*, in *Linnæa*, xxx. (1859–60) 681.—Guian.
Priauri, *Sagot*, in *Ann. Sc. Nat.* Sér. VI. xx. (1885) 188.—Guian.
probe, *Berg*, in *Linnæa*, xxix. (1857–58) 235.—Chili.
procera, *Poir. Encyc. Suppl.* ii. 129.—Am. bor.; Ind. occ.
products, *DC. Prod.* iii. 266.—Guian.
prominens, *Berg*, in *Mart. Fl. Bras.* xiv. i. 285.—Bras.
prosoneura, *Berg*, in *Linnæa*, xxxi. (1861–62) 255.—Bras.
protracta, *Berg*, *l. c.* 264.—Bras.
protracta, Steud. in Flora, xxvi. (1843) 762 = Salzmanni.
pruniformis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 340.—Bras.
Pseudocaryophyllus, *DC. Prod.* iii. 282.—Bras.
Pseudomalaccensis, Hort. Par. ex Lindm. in Fl. des Serres, Sér. I. v. (1849) 429 = malaccensis.
Pseudo-Mini, Steud. Nom. ed. II. i. 604 = Myrcia Pseudo-Mini.
Pseudopedilum, *Jacq. Enum. Pl. Carib* 23.—Ins. Martinic.
pseudoflora, *Berg*, in *Mart. Fl. Bras.* xiv. i. 233.—Bras.
psidioides, *DC. Prod.* iii. 268.—Guian.
psychotrioides, *Mart. ex Berg*, in *Mart. Fl. Bras.* xiv. i. 281.—Bras.
pterocarpa, *Baill. Hist. Pl.* vi. 338.—N. Caled.
pterocaulis, Miq. Anal. Bot. Ind. i. 18 = Memecylon appendiculatum.
pubescens, *DC. Prod.* iii. 283.—Venezuela.
pulchella, *Rosb. Hort. Beng.* [87]; *Fl. Ind.* ii. 496.—Ins. Molucc.
pulchra, *Berg*, in *Linnæa*, xxix. (1857–58) 294.—Chili.
pumila, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 109.—Bras.
punctata, Vahl, in West, Bidr. Ste-Croix, 289 = Myrcia punctata.
puugens, *Berg*, in *Mart. Fl. Bras.* xiv. i. 224.—Bras.
punicaefolia, *DC. Prod.* iii. 267.—Venezuela.
purpurascens, Baill. Hist. Pl. vi. 345 = malaccensis.
purpurea, Roxb. Hort. Beng. [87]; Fl. Ind. ii. 483 = malaccensis.
pustulata, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 495.—Malacca.
pycnacantha, *Benth. Pl. Hartw.* 174.—N. Granat.
pyramidalis, *Berg*, in *Mart. Fl. Bras.* xiv. i. 236.—Bras.
pyriflora, *Berg*, *l. c.* 299.—Bras.
pyrifolia, *Desv. in Ham. Prod. Pl. Ind. Occ.* 44.—Guian.
pyrifolia, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 487.—Sylhet.
pyrifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 336.—Bras.
pyrrocalda, *Berg*, in *Linnæa*, xxvii. (1854) 165.—Bras.
pyxophylla, *Hance*, in *Journ. Bot.* ix. (1871) 6.—China.
quadrangularis, *Duch. ex Griseb. in Goett. Abh.* vii. (1857) 214.—Guadeloup.
quadranguiata, *A. Gray, Bot. U. St. Expl. Exped.* i. 511.—Ins. Fiji.
quaternifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 363.—Bras.
quitarensis, *Benth. in Hook. Journ. Bot.* ii. (1840) 232.—Guian.
racemifera, *Berg*, in *Mart. Fl. Bras.* xiv. i. 232.—Bras.
racemifera, *Sagot*, in *Ann. Sc. Nat.* Sér. VI. xx. (1885) 196.—Guian.
racemiflora, *Berg*, in *Linnæa*, xxvii. (1854) 262.—Peruv.
racemosa, *DC. Prod.* iii. 281.—Bras.
racemosa, Linn. Sp. Pl. 471 = Barringtonia racemosa.
racemosa, *Berg*, in *Mart. Fl. Bras.* xiv. i. 304.—Bras.
ramiflora, *Desv. in Ham. Prod. Pl. Ind. Occ.* 43.—Guian.
ramiflora, Miq. in Linnæa, xxii. (1849) 536 = sphenophylla.
ramosissima, *Wall. Cat.* n. 3604.—Reg. Himal.
rampana, *Baker*, *Fl. Maurit.* 116.—Ins. Maurit.
Ranan, *Berm. in C. Gay, Fl. Chil.* ii. 388.—Chili.
ratiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 221.—Ins. Fiji.
reenvista, *Berg*, in *Mart. Fl. Bras.* xiv. i. 319.—Bras.
Regellana, *Berg*, *l. c.* 278.—Bras.
Regnelliana, *Berg*, *l. c.* 245.—Bras.
Reinwardtiana, *DC. Prod.* iii. 267.—Ins. Molucc.
repanda, *Berg*, in *Mart. Fl. Bras.* xiv. i. 264.—Bras.
reticulata, *Berg*, in *Linnæa*, xxvii. (1854) 270.—Bras.
EUGENIA:—
reticulata, Wight, Ic. t. 541; Illustr. ii. 16 = mangifolia.
retinadenia, *Wright*, in *Sauv. Fl. Cub.* 48.—Cuba.
retivenia, *Wright*, *l. c.* 44.—Cuba.
revoluta, *Berg*, in *Linnæa*, xxix. (1857–58) 224.—Cuba.
revoluta, *Wight*, *Ic.* t. 534; *Illustr.* ii. 17.—Ind. or.
revoluta, Willd. ex Berg, in Linnæa, xxvii. (1854) 184 = tuberculata.
rhododendrifolia, *Miq. Anal. Bot. Ind.* i. 19.—Borneo.
rhodomelea, Comm. ex DC. Prod. iii. 259 = cymosa.
rhomboearpa, *Berg*, in *Mart. Fl. Bras.* xiv. i. 229.—Bras.
rhopaloides, *DC. Prod.* iii. 278.—Peruv.
Ribeireana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 307.—Reg. Argent.
Richardiana, *Berg*, in *Linnæa*, xxx. (1859–60) 694.—Ins. Antigua.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 510.—Ins. Pacific.
Riedeliana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 261.—Bras.
rigida, *DC. Prod.* iii. 278.—Bras.
rigidifolia, *A. Rich. Ess. Fl. Cub.* 585; *et ex Berg*, in *Linnæa*, xxx. (1859–60) 679.—Cuba.
rimosa, *Wright*, in *Sauv. Fl. Cub.* 40.—Cuba.
riparia, A. Cunn. ex Hook. Bot. Mag. sub t. 4133 = Backhousia myrtifolia.
riparia, *DC. Prod.* iii. 288.—Bras.
rivularis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 337.—Bras.
rivularis, *Serm. Viti*, 486, nomen; *Fl. Vit.* 80.—Ins. Fiji.
rivularum, *Thw. Enum. Pl. Zeyl.* 115.—Zeylan.
robusta, Berg, in Linnæa, xxvii. (1854) 296 = caparidifolia.
romana, *Berg*, in *Mart. Fl. Bras.* xiv. i. 264.—Bras.
rorsimana, *Berg*, in *Linnæa*, xxvii. (1854) 219.—Guian.
roseo, *DC. Prod.* iii. 264.—Bras.
rosmarinifolia, *Poir. Encyc. Suppl.* iii. 129.—Ind. occ.
rostrata, Bedd. ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 493 = cymosa.
rostrata, *Berg*, in *Mart. Fl. Bras.* xiv. i. 282.—Bras.
Rottleriana, *Wight & Arn. Prod.* 331.—Ind. or.
rotundifolia, *Casaretti*, *Dec. Stirp. Nov. Bras.* (iv.) 40.—Bras.
rotundifolia, *Wight*, *Illustr.* ii. 17.—Zeylan.
Roxburghiana, Wall. Cat. sub n. 3608 = javanica.
Roxburghii, DC. Prod. iii. 271 = bracteata.
rubens, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 496.—Ind. or.; Burma.
rubescens, *A. Gray, Bot. U. St. Expl. Exped.* i. 525.—Ins. Pacific.
rubicunda, *Wight*, *Ic.* t. 538; *Illustr.* ii. 15.—Ind. or
rubiginosa, Cambess. in St. Hil. Fl. Bras. Mer. ii. 338 (= *Engenioptis rubiginosa*).—Bras.
rubra, Retzw. ex De Vriese, Pl. Ind. Bat. Reinw. 76 (= *Nelitris rubra*).—Malaya.
rubricaulis, *Duthie*, in *Hook. f. Fl. Brit. Ind.* ii. 487.—Java; Malaya.
rubrocincta, *Berg*, in *Mart. Fl. Bras.* xiv. i. 291.—Bras.
rufa, Barn. in C. Gay, Fl. Chil. ii. 387 = ferruginea.
rufescens, *DC. Prod.* iii. 279.—Bras.
rufo-fulva, *Thw. Enum. Pl. Zeyl.* 116.—Zeylan.
rugosa, Ruiz & Pav. ex DC. Prod. iii. 280 = polystachya.
Ruiziana, *Berg*, in *Linnæa*, xxvii. (1854) 167.—Guian.
rupestris, *Berg*, in *Mart. Fl. Bras.* xiv. i. 276.—Bras.
rusciifolia, *Poir. Encyc. Suppl.* iii. 128.—Am. merid.
rutidocarpa, *Ruiz & Pav. Fl. Per.* iv. t. 415.—Peruv.
sabulosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 355.—Bras.
Sagraea, *Berg*, in *Linnæa*, xxx. (1859–60) 690.—Bras.
salicifolia, *Berg*, in *Mart. Fl. Bras.* xiv. i. 312.—Bras.
salicifolia, Bach.-Ham. ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 473 = polypetala.
salicifolia, *DC. Prod.* iii. 278.—N. Granat.
salicifolia, *Lamk. in Oliver, Fl. Trop. Afr.* ii. 457.—Afr. trop.
salicifolia, *Wight*, *Illustr.* ii. 16 = Hæneana.
Salzmanni, *Benth. in Hook. Journ. Bot.* ii. (1840) 219.—Bras.

EUGENIA:—
aomarangensis, Berg, in Mart. Fl. Bras. xiv. i. 646, Index = jevenica.
sancta, *DC. Prod.* iii. 267.—Bras.
sandwicensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 519. —Isa. Sandvic.
savaiensis, *A. Gray, l. c.* 520.—Isa. Navigat.
scabrida, Wall. Cat. n. 3564 D = grata.
scandens, *Baker, Fl. Maurit.* 118.—Isa. Maurit.
scaphephylla, *Wright, in Sauv. Fl. Cub.* 41.—Cuba.
Schaueriana, *Miq. Stirp. Surin. Sel.* 41.—Guian.
Schiedeana, *Schlecht. in Linnaea*, xiii. (1839) 418.— Mexic.
Schlechtendaliana, *Berg, in Mart. Fl. Bras.* xiv. i. 321. —Bras.
Schomburgkii, *Benth. in Hook. Journ. Bot.* ii. (1840) 321.—Guian.
Schottiana, *Berg, in Mart. Fl. Bras.* xiv. i. 286.— Bras.
Schueckiana, *Berg, l. c.* 258.—Bras.
sclerophylla, *Bedd. For. Man. Bot.* 108.—Zeylan.
scoparia, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 489.— Singapore.
seechellarum, *Baker, in Proc. Roy. Irish Acad.* Ser. II. *Sc.* ii. (1875-77) 160.—Isa. Seychelles.
Seikirkii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 318.—Isa. J. Fernand.
Selloi, *Hort. ex Berg, in Mart. Fl. Bras* xiv. i. 327.— Bras.
Sellowiana, *DC. Prod.* iii. 262.—Bras.
sericea, *Berg, in Mart. Fl. Bras.* xiv. i. 280.—Bras.
sericiflora, *Benth. Bot. Voy. Sulph.* 98.—N Granat.
sessiliflora, *Vahl, in West, Bidr. Ste- Croix*, 290.— Isa. S. Doming.
sessiliflola, *DC. Prod.* iii. 263.—Bras.
Shepherdi, F. Muell. Fragm. ix. 148 = Myrtus Shepherdi.
Sieberiana, *DC. Prod.* iii. 277.—Ins. Trinit.
sinemariensis, *Aubl. Pl. Gui.* i. 501. t. 198.—Guian.
singapattiana, *Bedd. Ic. Pl. Ind. Or.* i. 85.—Ind. or.
sisophylla, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 486.— Penang.
smaragdina, *Berg, in Linnaea*, xxvii. (1854) 218.— Guadelup.
Smithii, *Poir. Encyc. Suppl.* iii. 126.—Austral.
Smithii, Spreng. ex Berg, in Linnaea, xxvii (1854) 191 = buxifolia, *Willd.*, monticola.
solimoensis, *Berg, in Mart. Fl. Bras.* xiv. i. 293.— Bras.
Sonderiana, *Berg, l. c.* 270.—Bras.
sparsiflora, *DC. Prod.* iii. 263.—Bras.
spathophylla, *Berg, in Mart. Fl. Bras.* xiv. i. 241.— Bras.
spathulata, Bedd. For. Man. Bot. 108 = clivifolia.
spathulata, *Berg, in Mart. Fl. Bras.* xiv. i. 249.— Bras.
speciosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 351.— Bras.
spectabilis, *Phil. in Linnaea*, xxviii. (1856) 689.— Chili.
sphaerocarpa, *DC. Prod.* iii. 279.—Bras.
sphenoides, *Berg, in Mart. Fl. Bras.* xiv. i. 259.— Bras.
sphenophylla, *Berg, l. c.* 288.—Bras.
spicata, Buch.-Ham. ex Wall. Cat. n. 3621 = bracteata.
spicata, Lam. Encyc. iii. 201 = sylanica.
spicifiora, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 62.—Bras.
splendens, *Berg, in Mart. Fl. Bras.* xiv. i. 226.—Bras.
Sprengelii, *DC. Prod.* iii. 276.—Bras.
Springiana, *Berg, in Mart. Fl. Bras.* xiv. i. 290.—Bras.
Spruceana, *Berg, l. c.* 257.—Bras.
squamata, Willd. ex Berg, in Linnaea, xxvii. (1854) 17 = Marlieria umbraticola.
stenophylla, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 322.—Ins. Chiloe.
Stephani, *Berg, in Mart. Fl. Bras.* xiv. i. 242.—Bras.
stictopetala, *Mart. ex DC. Prod.* iii. 270.—Bras.
stigmatosa, *DC. l. c.* 266.—Bras.
stipularis, *Miq. Fl. Ind. Bat.* i. 1. 441.—Ins. Amboin.
Stockii, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 498.— Ind. or.
strelitzensis, *Berg, in Mart. Fl. Bras.* xiv. i. 256.—Bras.
stricta, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 179.—N. Caledon.
strigipes, *Berg, in Mart. Fl. Bras.* xiv. i. 268.—Bras.
strigosa, Sieb. ex Berg, in Linnaea, xxx. (1859-60) 699 = Calyptranthes cuprea.

EUGENIA:—
suaveolens, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 339. —Bras.
subalterna, *Benth. in Hook. Journ. Bot.* ii. (1840) 320. —Guian.
subamplexicaulis, *DC. Prod.* iii. 277.—Bras.
subavenia, *Berg, in Mart. Fl. Bras.* xiv. i. 271.— Bras.
subavenis, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 489.— Bras.
subcordata, *Berg, in Mart. Fl. Bras.* xiv. i. 300.— Bras.
subcordata, *Wight & Arn. Prod.* 881.—Ind. or.
subcorymbosa, *Berg, in Mart. Fl. Bras.* xiv. i. 295.— Bras.
subdecussata, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 491. —Malaya.
suberosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 364. —Bras.
subobliqua, *Benth. in Hook. Journ. Bot.* ii. (1840) 322.—Guian.
suborbicularis, *Benth. Fl. Austral.* ii. 385.—Austral.
subterminalis, *DC. Prod.* iii. 263.—Bras.
subvericillata, *Berg, in Linnaea*, xxix. (1857-58) 284.—Am. centr.
suffruticosa, *Berg, in Mart. Fl. Bras.* xiv. i. 242.— Bras.
sulcata, *Spring, ex Mart. in Flora*, xx. (1837) ii. *Beibl.* 85.—Bras.
supraaxillaris, *Spring, l. c.* 83.—Bras.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 748; *et* xxvii. (1854) 182.—Guian.
sylvatica, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 337. —Bras.
sylvatica, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 352 (= *Eugenioptis? sylvatica*).—Bras.
sylvestris, *Moon, ex Wight, Ic.* t. 582; *Illustr.* ii. 15.— Zeylan.
sylvicola, *Wright, in Sauv. Fl. Cub.* 41.—Cuba.
Tabasco, G. Don, Gen. Syst. ii. 866 = Pimenta officinalis.
tanaensis, *Baker, in Journ. Bot.* xx. (1882) 111.— Madag.
tapacumensis, *Berg, in Linnaea*, xxvii. (1854) 222.— Guian.
teffensis, *Berg, in Mart. Fl. Bras.* xiv. i. 255.—Bras.
Temu, *Hook. & Arn. Bot. Beech. Voy.* 56.—Chili
tenella, *DC. Prod.* iii. 272.—Bras.
tenella, *Miq. in Linnaea*, xxii. (1849) 582.—Bras.
tenuifolia, *Berg, in Mart. Fl. Bras.* xiv. i. 249.—Bras.; Venezuela.
tenuiramea, Miq. Stirp. Surin. Sel. 39 = egensis.
tenuis, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 500.—Ind. or.
tephrodes, *Hance, in Journ. Bot.* xxiii. (1885) 7.— China.
ternatifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 336.—Bras.
ternifolia, *Berg, in Linnaea*, xxvii. (1854) 246.— Venezuela.
ternifolia, Roxb. Hort. Beng. [37]; Fl. Ind. ii. 489 = formosa.
terpnophylla, *Thw. Enum. Pl. Zeyl.* 114.—Zeylan.
tetraedra, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 476.— Malaya.
tetragona, *Wight, Illustr.* ii. 16.—Reg. Himal.; Burma.
tetrasperma, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 271.—Porto Rico.
tetrastichia, *Poepp. ex Berg, in Mart. Fl. Bras.* xiv. i. 234.—Bras.
Teyu-ibe, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 116; *Contrib.* 196.—Reg. Argent.
thamnoides, *Berg, in Linnaea*, xxvii. (1854) 179.— Chili.
thomasiana, *Berg, l. c.* 183.—Ins. S. Thomas.
Thumm, *Roxb. Hort. Beng.* [92]; *Fl. Ind.* ii 495.— Burma.
Thwaitesii, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 506. —Zeylan.
Tierneyana, *F. Muell. Fragm.* iv. 14.—Austral.
tinguyensis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 358. —Bras.
tinifolia, Lam. Encyc. iii. 209 = mespiloides.
toeniana, *Berg, in Mart. Fl. Bras.* xiv. i. 269.—Bras.
toddalioides, *Wight, Ic.* t. 542; *Illustr.* ii. 16.—Reg. Himal.; Burma.
tomentosa, Aubl. Pl. Gui. i. 504 = Myrcia tomentosa.
tomentosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 353. —Bras.

EUGENIA:—
trichoscypa, *Phil. in Linnaea*, xxviii. (1856) 668.— Chili.
triflora, *Ham. Prod. Fl. Ind. Occ.* 44.—N. Granat.
triflora, Willd. ex Berg, in Mart. Fl. Bras. xiv. i. 462 = Campomanesia triflora.
trinervia, DC. Prod. iii. 279 = Rhodamnia trinervia.
trinervia, Vahl, Eclog. Am. ii. 36 = Vahlii.
Trinitatis, *DC. Prod.* iii. 280.—Ins. Trinit.
triphylla, *Berg, in Mart. Fl. Bras.* xiv. i. 243.—Bras.
triplinervia, Berg, in Linnaea, xxvii. (1854) 190 = buxifolia *Willd.*, monticola.
triquetra, *Berg, l. c.* 141.—N. Granat.
tristis, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 233.—Burma; Ind. or.
truncata, *Berg, in Linnaea*, xxvii. (1854) 157.—Costa Rica.
truncifiora, *G. Don, Gen. Syst.* ii. 867.—Mexic.
tuberculata, *DC. Prod.* iii. 273.—Cuba.
tumida, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 487.— Malacca.
turbinata, *Berg, in Mart. Fl. Bras.* xiv. i. 218.—Bras.
tutulienis, *A. Gray, Bot. U. St. Expl. Exped.* i. 529.— Isa. Navigat.
Tweediei, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 328.—Uruguay.
ubensis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 351.— Bras.
Ugni, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 318 = Myrtus Ugni.
umbelleria, Berg, in Mart. Fl. Bras. xiv. i. 292 = dicrosna.
umbellata, DC. Prod. iii. 273 = dicrosna.
umbellata, Spreng. Neue Entdeck. ii. 169 = erythrocarpa.
umbelliflora, *Berg, in Mart. Fl. Bras.* xiv. i. 290.— Bras.
umbrosa, Bedd. For. Man. Bot. 108 = subavenis.
umbrosa, *Berg, in Mart. Fl. Bras.* xiv. i. 282.—Bras.
undulata, *Aubl. Pl. Gui.* i. 508. t. 202.—Guiana.
unedyfolia, Spreng. ex Berg, in Linnaea, xxvii. (1854) 193 = buxifolia, *Willd.*, monticola.
unifiora, *Berg, in Mart. Fl. Bras.* xiv. i. 215.—Bras.
unifiora, *Linn. Sp. Pl.* 470.—Am. trop.
uruguayensis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 362.—Bras.
Uvalha, *Cambess. l. c.* 367.—Bras.
vacciniifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 148.—Madag.
vagae, Berg, in Mart. Fl. Bras. xiv. i. 238 = paniculifolia.
Vahlii, *DC. Prod.* iii. 279.—Guian.
valdevenosa, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 489 —Malaya.
Valenzuelana, *A. Rich. Ess. Fl. Cub.* i. 393.—Cuba.
variabilis, *Baill. Hist. Pl.* vi. 341.—Bras.
varians, *Miq. Anal. Bot. Ind.* i. 21.—Malaya.
Vauthieriana, *Berg, in Mart. Fl. Bras.* xiv. i. 217.— Bras.
Velloziana, *Berg, in Linnaea*, xxvii. (1854) 202.— Bras.
Vellozii, *Berg, in Mart. Fl. Bras.* xiv. i. 255.—Bras.
velutina, *Berg, l. c.* 317.—Bras.
venezuelensis, *Berg, in Linnaea*, xxvii. (1854) 188.— Venezuel.
venosa, *Lam. Encyc.* iii. 200.—Isa. Mascar.
Ventenatii, *Benth. Fl. Austral.* iii. 383.—Austral.
venulosa, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 490.— Malaya.
venusta, *Roxb. Hort. Beng.* [87]; *Fl. Ind.* ii. 491.— Burma.
verecunda, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 496. —Malaya.
verniciosa, *Berg, in Mart. Fl. Bras.* xiv. i. 274.—Bras.
verrucosa, *A. Rich. Ess. Fl. Cub.* 389.—Cuba.
verrucosula, Steud. Nom. ed. II. i. 605 = seq.
verruculosa, *DC. Prod.* iii. 284.—Bras.
verticillaris, *Berg, in Mart. Fl. Bras.* xiv. i. 281.—Bras.
vetula, *DC. Prod.* iii. 271.—Bras.
vicosensis, *Berg, in Mart. Fl. Bras.* xiv. i. 279.— Bras.
Vieillardi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 180.—N. Caled.
villosa, Poir. Encyc. Suppl. iii. 124 = Monimia rotundifolia.
viminalis, *Berg, in Mart. Fl. Bras.* xiv. i. 287.—Bras.
vincaefolia, *Berg, l. c.* 270.—Bras.
violacea, *Lam. Encyc.* iii. 200.—Isa. Mascar.
violacea, *Berg, in Mart. Fl. Bras.* xiv. i. 281.— Bras.

EUGENIA :—
virgata, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 102.—Bras.
virgata, *Macfad. Fl. Jamaic.* ii. 121.—Jamaic.
virgultosa, *DC. Prod.* iii. 280.—Ind. occ.
virkilifora, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 844. —Bras.
viridis, *Berg, in Linnæa*, xxvii. (1854) 209.—Bras.
vismeaefolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 270.—Guian.
vismioides, *DC. Prod.* iii. 267.—Ins. Molucc.
Vitis-idæa, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 122 = Myrtus pedunculata.
Vitis-idæa, *Wright, in Sauv. Fl. Cub.* 49.—Cuba.
vulgaris, Baill. Hist. Pl. vi. 346 = E. Jambos.
Wallenii, *Macfad. Fl. Jamaic.* ii. 118.—Jamaic.
Wallichii, *Wight, Ic.* t. 536; *Illustr.* ii. 17.—Reg. Himal.; Burma.
Warszewiczii, *Hemsl. Biol. Centr. Am. Bot.* i. 411.— Panama.
Widgrenii, *Sond. ex Berg, in Mart. Fl. Bras.* xiv. i. 327.—Bras.
Wightiana, *Wight, Ic.* t. 599; *Illustr.* ii. 15.—Ind. or. *Wightii*, Bedd. for. Man. Bot. 109 = lasta.
Willdenowii, DC. Prod. iii. 265 = unifora.
Willdenowii, Wight, Illustr. ii. 13 = bracteata.
Willconii, *F. Muell. Fragm.* v. 19.—Austral.
Wrightii, *Baker, in Proc. Irish Acad.* Ser. II. *Sc.* ii. (1875–77) 160.—Ins. Seychell.
wyandensis, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 47.—Ind. or.
xalapensis, *DC. Prod.* iii. 277.—Mexic.
xanthocarpa, Mart. Syst. Mat. Med. Bras. 31, 62 = Campomanesia xanthocarpa.
xanthocarpa, *Thw. Enum. Pl. Zeyl.* 416.—Zeylan.
xanthoclora, *Berg, in Mart. Fl. Bras.* xiv. i. 316.— Bras.
xanthoxyloides, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 866.—Bras.
xylopifolia, *DC. Prod.* iii. 179.—Guiana.
xylophyllum, *Berg, in Linnæa*, xxvii. (1854) 196.— Cuba.
Ybaviya, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 69; *Contrib.* 119.—Reg. Argent.
ypanemensis, *Berg, in Mart. Fl. Bras.* xiv. i. 316.— Bras.
yrirensis, Berg, l. c. 647, Index = irirensis.
yumuryensis, *Berg, in Linnæa*, xxvii. (1854) 234.— Cuba.
Zeyheri, *Harv. Gen. S. Afr. Pl.* 416; *et in Fl. Cap.* ii. 593.—Afr. austr.
zeylanica, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 490 = bracteata.
zeylanica, *Wight, Ic.* i. 78; *Illustr.* ii. 15.—Ind. or.; Malaya.
zeylanica, Willd. Sp. Pl. ii. 968 = unifora.
Zuccarinii, *Berg, in Mart. Fl. Bras.* xiv. i. 257.—Bras.

EUGENIOIDES, Linn. Fl. Zeyl. 192 (1747) = Symplocos, Linn. (Styrac.).

EUGENIOPSIS, Berg, in Linnæa, xxvii. (1854) 80 = Marlieria, Cambess. (Myrtac.).
acuminatissima, Berg, in Mart. Fl. Bras. xiv. i. 144. —Bras.
affinis, Berg, l. c. 149.—Bras.
angustifolia, Berg, l. c. 148.—Bras.
cannæfolia, Berg, l. c. 148.—Bras.
Clausseniana, Berg, l. c. 148.—Bras.
Gardneriana, Berg, l. c.—Bras.
Gaudichaudiana, Berg, l. c. 147.—Bras.
grandifolia, Berg, l. c. 149.—Bras.†
laurigrate, Berg, in Linnæa, xxvii. (1854) 81.—Bras.
Luschnathiana, Berg, in Mart. Fl. Bras. xiv. i. 147.— Bras.
ovata, Berg, l. c. 148.—Bras.
paniculata, Berg, in Linnæa, xxvii. (1854) 82.—Bras.
polygama, Berg, in Mart. Fl. Bras. xiv. i. 147.— Bras.
Richardiana, Berg, in Linnæa, xxx. (1859–60) 665.— Ind. occ.
Riedeliana, Berg, in Mart. Fl. Bras. xiv. i. 561.— Bras.
rubiginosa, Berg, in Linnæa, xxvii. (1854) 82.—Bras.
Schottiana, Berg, in Mart. Fl. Bras. xiv. i. 148.— Bras.
sylvatica, Berg, in Linnæa, xxvii. (1854) 81.—Bras.
Treucheriana, Berg, l. c. xxxi. (1861–62) 249.—Bras.

EUGONE, Salisb. Prod. 388 = Gloriosa, Linn. (Liliac.).
superba, Salisb. l. c. = G. superba.

EUILUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 149 = Astragalus, Tourn. (Legumin.).
roduncus, Stev. l. c. = A. roduncus.

EUKLASTAXON, Steud. in Flora, xxxiii. (1850) 228, nomen: Syn. Fl. Germ. 413 (1855) = Andropogon, Linn. (Gramin.).
tenuifolius, Steud. l. c. = A. leucostachys.

EUKRANIA, Rafin. Alsog. Amer. 59 (1838) = Cornus, Tourn.
canadensis, Rafin. l. c. = C. canadensis.
cyanocarpa, Rafin. l. c. 60 = C. cyanocarpa.
mascula, Rafin. l. c. 59 = C. Mas.
suecica, Rafin. l. c. = C. suecica.

EUKYLISTA, Benth. in Hook. Kew Journ. v. (1853) 330 = Calycophyllum, DC. (Rubiac.).
Spruceana, Benth. l. c. 331.—Peruv.

EULALIA, Kunth, Rév. Gram. i. 160. t. 93 (1829) = Pollinia, Trin. (Gramin.).
argentea, Brongn. in Duperr. Voy. Coq. Bot. 92 = P. argentea.
aurea, Kunth, Rév. Gram. i. 160 = P. aurea.
caprasis, Hochst. in Flora, xxix. (1846) 117 = P. nuda.
concinna, Nees, ex Steud. Syn. Pl. Gram. 412 = P. articulata.
glabrata, Brongn. in Duperr. Voy. Coq. Bot. 93 = P. glabrata.
mexicana, Steud. Syn. Pl. Gram. 413 = Andropogon saccharatus?
villosa, Nees, Fl. Afr. Austr. 91 = P. villosa.

EULALIA, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 332, partim = Miscanthus, Anderss. (Gramin.).
cotulifera, Munro, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 189 = M. cotulifer.
densa, Munro, in Seem. Bot. Voy. Herald, 424 = M. japonicus.
japonica, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 332 = M. sinensis.
nepalensis, Trin. l. c.—Reg. Himal.

EULEUCUM, Rafin. Fl. Tellur. iii. 56 (1836) = Coreusa, D. Don (Empetrac.).
album, Rafin. l. c. =C. album.

EULOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 514 (1838–40). *ONAGRARIEÆ*, Benth. & Hook. f. i. 789.
californicus, *Nutt. l. c.* 515.—Am. bor. occ.

EULOPHIA, R. Br. [ex Lindl.] in Bot. Reg. t. 666 (1823). *ORCHIDEÆ*, Benth. & Hook. f. iii. 535.
CYRTOPERA, Lindl. in Wall. Cat. n. 7362–64 (1831!); et Gen. et Sp. Orch. 189 (1835).
ORTHOCHILUS, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 264 t. 82 (1851).
abyssinica, *Reichb. f. in Linnæa*, xxii. (1849) 866.— Abyss.
acaliscta, *Spreng. Syst.* iii. 730.—Afr. austr.
adenoglossa, *Reichb. f. Otia Bot. Hamb.* 66.—Afr. trop.
albiflora, Edgew. ex Lindl. in Journ. Linn. Soc. iii. (1859) 30 = barbacata.
alismatophylla, *Reichb. f. in Flora*, lxviii. (1885) 543. —Afr. austr.
aloifolia, *Welw. ex Reichb. f. l. c.* i. (1867) 104.—Afr. trop.
andamanensis, *Reichb. f. l. c.* iv. (1872) 276.—Burma.
articulata, *Lindl. Gen. et Sp. Orch.* 181.—Afr. trop.
Arundinæ, *Reichb. f. in Linnæa*, xxii. (1849) 854.— Bras.
atrovirens, *Lindl. Gen. et Sp. Orch.* 188.—Ins. Maurit.
baglisensis, *Reichb. f. Otia Bot. Hamb.* 66.—Afr. trop.
barbata, *Spreng. Syst.* iii. 730.—Afr. austr.

EULOPHIA:—
beravensis, *Reichb. f. in Bot. Zeit.* xxxix. (1881) 649; *Otia Bot. Hamb.* 74.—Madag.
bicolor, Blume, Orch. Archip. Ind. 181 = Cyrtopodium bicolor.
bicolor, Dals. in Hook. Kew Journ. iii. (1851) 843 = nuda.
bicolor, *Reichb. f.* & *Sond. ex Reichb. f. in Flora*, xlviii. (1865) 186.—Afr. austr.
brachyrrhiza, Lindl. in Journ. Linn. Soc. iii. (1859) 24 = barbacata.
bracteosa, *Lindl. in Wall. Cat.* n. 7366.—Reg. Himal.; Burma.
cafra, *Reichb. f. in Flora*, xlviii. (1866) 186.—Afr. austr.
callichroma, *Reichb. f. Otia Bot. Hamb.* 116.—Afr. austr.
campestris, *Wall. Cat.* n. 7367.—Ind. or.
capensis, Bolus, in Journ. Linn. Soc. xix. (1882) 336 = aculeata.
carinata, Lindl. Gen. et Sp. Orch. 183 = virens.
carunculifera, *Reichb. f. in Flora*, lxiv. (1881) 329.— Afr. austr.
celebica, *Blume, Orch. Archip. Ind.* 182.—Ins. Celebes.
ciliata, Reichb. f. in Walp. Ann. vi. 647 = gracilis.
clavicornis, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 209 = hians.
cochlearis, *Lindl. l. c.*—Afr. austr.
cochlearis, Steud. Nom. ed. II. i. 605 = Zygopetalum cochlearе.
comosa, *Sond. in Linnæa*, xix. (1847) 72.—Afr. austr.
concolor, *Lindl. Gen. et Sp. Orch.* 181.—Ins. Borbon.
Cooperi, *Reichb. f. in Flora*, lxiv. (1881) 330.—Afr. austr.
cordylophylla, *Reichb. f. l. c.* lxviii. (1885) 541.—Ins. Comoro.
crinita, G. Don, in Loud. Hort. Brit. Suppl. i. 588 = Zygopetalum crinitum.
cristata, *Lindl. Gen. et Sp. Orch.* 186.—Afr. trop.
cucullata, Lindl. l. c. = Polystachya grandiflora.
Cullenii, *Blume, Orch. Archip. Ind.* 182, in obs.— Ind. or.; Chinа.
decipiens, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 155 = gramines.
densiflora, *Lindl. in Journ. Linn. Soc.* iii. (1859) 25. —Reg. Himal.
dilatata, Lindl. in Ann. & Mag. Nat. Hist. x. (1842) 184 = Bletia secunda.
dipodiiformis, *Reichb. f. in Bonplandia*, v. (1857) 38. —Malaya.
Dregeana, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 208.—Afr. austr.
elongata, *Blume, Orch. Archip. Ind.* 182.—Sumatra.
emarginata, *Blume, l. c.* 180.—Sumatra.
emarginata, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 209 = hians.
ensata, *Lindl. in Bot. Reg.* t. 1147.—Afr. austr.
euglossa, *Reichb. f. in Bot. Mag.* t. 5561.—Afr. trop.
exaltata, *Reichb. f. in Bonplandia*, v. (1857) 38.— Java.
explanata, *Lindl. Gen. et Sp. Orch.* 180.—Reg. Himal.
filiculalis, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 184.—Mexic.
Fitzalani, *F. Muell. Fragm.* viii. 30; *ex Benth. Fl. Austral.* vi. 260.—Austral.
foliosa, *Bolus, in Journ. Linn. Soc.* xix. (1882) 337.— Afr. austr.
fusca, Blume, Orch. Archip. Ind. 182, in obs. = nuda.
galbana, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 469. —Madag.
gamoleia, *Lindl. in Bot. Reg.* t. 742.—Afr. trop.
graminea, *Lindl. in Wall. Cat.* n. 7372.—Ind. or.; Malaya.
grandiflora, Lindl. Gen. et Sp. Orch. 181 = tenuissima.
guineensis, (*Lindl. in*) *Bot. Reg.* t. 696.—Afr. trop.
hastata, Lindl. in Journ. Linn. Soc. iii. (1859) 25 (Taiwan sp.).—Reg. Himal.
hellebrina, Hook. f. Bot. Mag. t. 5875 = Habenaria hellebrina.
hemilæca, Lindl. in Journ. Linn. Soc. iii. (1859) 25 = campestris.
herbacea, *Lindl. Gen. et Sp. Orch.* 183.—Ind. or.
hians, *Spreng. Syst.* iii. 730.—Afr. austr.
incompleta, Coll. Motel. iii. 249 = gracilis.

EULOPHIA:—
tamelata, *Lindl. Gen. et Sp. Orch.* 184.—Afr. austr.
teretiglossa, *Reichb. f. in Flora*, lxiv. (1881) 229.—Afr. austr.
Teschenaultii, *Blume, Orch. Archip. Ind.* 182.—Java.
iliacea, Lindl. Gen. et Sp. Orch. 185 = Govenia iliacea.
limochiloides, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 208.—Afr. austr.
lonchophylla, *Reichb. f. in Flora*, lxvii. (1885) 542.—Ins. Comoro.
longicollis, *Lindl. in Journ. Linn. Soc.* vi. (1862) 132.—Afr. trop.
longicornis, Spreng. Syst. iii. 720 = Mystacidium filicorne.
lurida, *Lindl. Gen. et Sp. Orch.* 182.—Afr. trop
lutea, *Blume, Orch. Archip. Ind.* 181.—Java.
lutea, *Lindl. in Journ. Linn. Soc.* vi. (1862) 132.—Afr. trop.
Machaiana, Lindl. Bot. Reg. t. 1438 = Zygopetalum Mackaii.
macrochlas, *Blume, Orch. Archip. Ind.* 183.—Java.
macrostachya, *Lindl. Gen. et Sp. Orch.* 183.—Ind. or.
maculata, *Reichb. f. in Walp. Ann.* vi. 647.—Bras.
madagascariensis, *Kraenzl. in Bremen, Abh.* vii. (1882) 256.—Madag.
maxillaris, G. Don, in Loud. Hort. Brit. Suppl. i. 588 = Zygopetalum maxillare.
megistophylla, *Reichb. f. in Gard. Chron.* (1885) i. 787; *et in Flora*, lxviii. (1885) 879.—Ins. Comoro.
Melesgris, *Reichb. f. in Linnaea*, xx. (1847) 683.—Afr. austr.
micrantha, *Lindl. Gen et Sp. Orch.* 184.—Afr. austr.
Milnei, *Reichb. f. Otia Bot. Hamb.* 116.—Afr. trop.
monilie, *Reichb. f. in Flora*, l. (1867) 105.—Afr. trop.
monophylla, *S. Moore, in Baker, Fl. Maurit.* 360.—Ins. Mascar.
monticola, *Blume, Orch. Archip Ind.* 182.—Java.
Murrayana, Steud. Nom. ed. II. i. 605 = Zygopetalum Murrayanum.
natalensis, *Reichb. f. in Flora*, xlviii. (1865) 186.—Afr. austr.
nuda, *Lindl. in Wall. Cat.* n. 7371.—Ind. or.
nutans, *Sond. in Linnaea*, xix. (1847) 73.—Afr. austr.
ochreata, *Lindl. in Journ. Linn. Soc.* iii. (1859) 24.—Ind. or.
odontoglossa, Reichb. f. in Linnaea, xix. (1847) 373 = sclista.
ovalis, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 302 = barbata.
parvilabris, *Lindl. l. c.* 201.—Afr. austr.
pedicellata, Spreng. Syst. iii. 720 = aculeata.
Peterzii, *Reichb. f. in Flora*, xlviii. (1865) 186.—Afr. trop.
pileata, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 468.—Madag.
platypetala, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 209.—Afr. austr.
plicata, Bolus, in Journ. Linn. Soc. xix. (1882) 336, excl. syn. pro parte = aculeata.
pratensis, *Lindl. in Journ. Linn. Soc.* iii. (1859) 25.—Ind. or.
promensis, Lindl. in Wall. Cat. n. 7365 (Tainise sp.).—Burma.
proteranra, *Reichb. f. in Flora*, l. (1867) 104.—Afr. trop.
pulchra, *Lindl. Gen. et Sp. Orch.* 183.—Ins. Borbon.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* ii. 334.—Abyss.
ramentacea, Lindl. in Wall. Cat. n. 7370 = campestris.
ramentacea, Wight, Ic. t. 1866 = pratensis.
ramosa, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 470.—Madag.
reticulata, *Ridl. l. c.*—Madag.
rostrata, Steud. Nom. ed. II. i. 605 = Zygopetalum rostratum.
rupestris, Lindl. in Wall. Cat. n. 7368 = campestris.
rupestris, *Reichb. f. in Linnaea*, xx. (1847) 683.—Afr. austr.
rupicola, Reichb. f. in Benjkudia, v. (1857) 38 = rupestris.
Rumohrigiana, *Kraenzl. in Bremen, Abh.* vii. (1882) 256.—Madag.
Saundersiana, *Reichb. f. in Bot. Zeit.* xxiv. (1866) 378.—Ash. Insp. ?

EULOPHIA:—
Schimperiana, *A. Rich. Tent. Fl. Abyss.* ii. 333.—Abyss.
sclerophylla, *Reichb. f. in Flora*, lxviii. (1885) 542.—Afr. austr.
scripta, *Lindl. Gen. et Sp. Orch.* 181.—Ins. Mascar.
sinensis, *Rolfe, in Journ. Bot. Néerl.* i. (1861) 91.—China.
sphaerocarpa, *Sond. in Linnaea*, xix. (1847) 74.—Afr. austr.
squalida, *Lindl. Bot. Reg.* (1841) *Misc.* 77.—Malaya.
stachyodes, *Reichb. f. Otia Bot. Hamb.* 66.—Afr. trop.
stenochila, Steud. Nom. ed. II. i. 605 = Zygopetalum crinitum.
stenopetala, *Lindl. in Journ. Linn. Soc.* iii. (1859) 26.—Reg. Himal.
streptopetala, *Lindl. Bot. Reg.* t. 1002.—Afr. austr.
stricta, Lindl. Gen. et Sp. Orch. 185 = Bletia stricta.
tenella, *Reichb. f. in Linnaea*, xx. (1847) 681.—Afr. austr.
tristis, *Spreng. Syst.* iii. 720.—Afr. austr.
vaginata, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 467.—Madag.
venosa, *Reichb. f. ex Benth. Fl. Austral.* vi. 300; *et Xenia Orch.* ii. 198.—Austral.
vernicosa, *Reichb. f. Otia Bot. Hamb.* 116.—Afr. trop.
violacea, *Reichb. f. in Linnaea*, xx. (1847) 683.—Afr. austr.
virens, *Spreng. Syst.* iii. 720.—Ind. or.
virens, Stocks, ex Lindl. in Journ. Linn. Soc. iii. (1859) 25 = pratensis.
viridiflora, Steud. Nom. ed. II. i. 605 = virens
viriis, *Lindl. in Journ. Linn. Soc.* vi. (1862) 132.—Afr trop.
Zeyheriana, *Sond. in Linnaea*, xix. (1847) 73.—Afr. austr.

EULOPHIUS, R. Br. in Bot. Reg. sub t. 573 (1821) = Eulophia, R. Br. (Orchid.).
carinatus, R. Br. l. c. = Eulophia virens.
virens, R. Br. l. c. = Eulophia virens.

EULOPHUS, Nutt. ex DC. Coll. Mém. v. 69 (1829); et Prod. iv. 248 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 886.
PEUCEDANINA, Reichb. Handb. 219 (1837).
americanus, *Nutt. ll. cc.*—Am. bor.
peucedanoides, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* i. 56.—Mexic.
texanus, *Benth. & Hook. f. Gen.* i. 885.—Am. bor.

EULOPHUS, Nutt. in Journ. Acad. Philad. vii. (1834) 27 = Peucedanum, Tourn. (Umbell.).
ambiguus, Nutt. l. c. = P. ambiguum.
triternatus, Nutt. l. c. = P. triternatum.

EULYCHNIA, Phil. Fl. Atac. 28. t. 11 A (1861). *CACTEAE*, Benth. & Hook. f. i. 1008.
acida, *Phil. in Linnaea*, xxxiii. (1864–65) 80.—Chili.
breviflora, *Phil. Fl. Atac.* 34.—Chili.
castanea, *Phil. in Linnaea*, xxxiii. (1864–65) 80.—Chili.

EUMACHIA, DC. Prod. iv. 178 (1830) = Ixora, Linn. (Rubiac.).
coccinea, DC. l. c. 479 = I. carnea.

EUMECANTHUS, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 348 = Euphorbia, Linn.
arenarius, Klotzsch & Garcke, l. c. 249 = Euph. arenaria.
arienaia, Klotzsch & Garcke, l. c. = Euph. arienaia.
Benthamianus, Klotzsch & Garcke, l. c. = Euph. arienaia.
triphyllus, Klotzsch & Garcke, l. c. = Euph. scabrella.

EUMOLPE, Decne. ex Jacquem & Hérincq, Man. Gén. Pl. ii. 678 (1849 ?) = Glorinia, L'Hérit. (Gesnerac.).
fimbriata, Decne. l. c. = G. glabrata.

EUMORPHA, Eckl. & Zeyh. Enum. 77 (1836) = Pelargonium, L'Hérit. (Geraniac.).
catarractae, Eckl. & Zeyh. l. c. 78 = P. patulum.
elegans, Eckl. & Zeyh. l. c. 77 = P. elegans.
grandiflorum, Eckl. & Zeyh. l. c. 76 = P. grandiflorum.
laevigata, Eckl. & Zeyh. l. c. = P. laevigatum.
marmorata, Eckl. & Zeyh. l. c. = P. patulum.
nobilis, Eckl. & Zeyh. l. c. 77 = P. nobile.

EUMORPHIA:—
tabularis, Eckl. & Zeyh. Enum. 77 = Pelargonium tabulare.
truncilobe, Eckl. & Zeyh. l. c. 78 = P. patulum.
variegata, Eckl. & Zeyh. l. c. = P. patulum.

EUMORPHIA, DC. Prod. vi. 2 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 415.
Dregeana, *DC. l. c.* 8.—Afr. austr.

EUNANUS, Benth. in DC. Prod. x. 374 (1846) = Mimulus, Linn. (Scrophul.).
angustatus, Greene, in Bull. Calif. Acad. i. 111. (1885) 99.—Calif.
bicolor, A. Gray, in Proc. Am. Acad. vii. (1868) 381.—Calif.
Bigelovii, A. Gray, in Pacif. Rail. Rep. iv. 121 = M. Bigelovii.
Bolanderi, Greene, in Bull. Calif. Acad. i. 111. (1885) 105 = M. Bolanderi.
brevipes, Greene, l. c. = M. brevipes.
Breweri, Greene, l. c. 101 = M. rubellus.
Coulteri, Harv. & A. Gray, ex Benth. Pl. Hartw. 329 = M. tricolor.
Douglasii, Benth. in DC. Prod. x. 374 = M. atropurpureus.'
Douglasii, Benth. Pl. Hartw. 329, partim = *Kelloggii*.
Douglasii, A. Gray, in Proc. Am. Acad. xi. (1876) 95, partim = M. Douglasii.
Fremontii, Benth. in DC. Prod. x. 374 = M. Fremontii.
Fremontii, S. Wats. Bot. King's Exped. 226 = M. nanus, *Hook. & Arn.*
Kelloggii, Curran, ex Greene, in Bull. Calif. Acad. i. 111. (1885) 100.—Calif.
latifolius, Greene, l. c. 99 = M. latifolius.
Layneae, Greene, l. c. 104 = M. leptalcas.
leptalcus, Greene, l. c. 101 = M. leptalcus.
mephiticus, Greene, l. c. 103 = M. mephiticus.
mohavensis, Greene, l. c. 106 = M. mohavensis.
Parryi, Greene, l. c. 104 = M. Parryi.
pictus, Greene, l. c. 106.—Calif.
Rattani, Greene, l. c. 105 = M. Rattani.
Tolmiei, Benth. in DC. Prod. x. 374 = M. nanus, *Hook. & Arn.*
Torreyi, Greene, in Bull. Calif. Acad. i. 111. (1885) 104 = M. Torreyi.
tricolor, Greene, l. c. 99 = M. tricolor.

EUNOMIA, DC. Syst. ii. 555 (1821) = Asthlonema, R. Br. (Crucif.).
Bourgaei, Boiss. Fl. Orient. i. 344 = A. Bourgaei.
chlorocephalia, DC. Syst. ii. 556 = A. chlorocephalum.
cochlearina, F. Muell. in Linnaea, xxv. (1852) 369 = Thlaspi cochlearinum.
cordata, DC. Syst. ii. 557 = A. cordatum.
cordata, Pinard, ex Boiss. Diagn. Ser. I. v. 68 = A. Moricandianum.
iberidea, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 191 = A. iberideum.
Menziesii, Janb. & Spach, Illustr. i. 101 = A. iberideum.
Moricandiana, Boiss. Diagn. Ser. I. v. 88 = A. Moricandianum.
oppositifolia, DC. Syst. ii. 556 = A. oppositifolium.
orbicularis, Griseb. Spicil. Fl. Rumel. i. 289 (= *Crenularia orbicularis*).—Macedon.
rotundifolia, C. A. Mey. Verz. Pfl. Casc. 194 = A. rotundifolium.
rubescens, Schost & Kotschy, ex Tchihat. Asie Min. Bot. i. 339 = A. rubescens.
subulata, Boiss. & Heldr. Diagn. Ser. I. viii. 42 = A. tabulatum.

EUNOXIS, Rafin. New Fl. Am. iv. 84 (1836) = Lactuca, i Tourn. (Compos.).
amabilis, Rafin. l. c.—Am. bor.

EUODIA, Forst. Char. Gen. 13 (1776) = Evodia, Forst. (Rutac.).

EUODIA, Gaertn. Fruct. ii. t. 108 (1791) = EVODIA, Gaertn. = Ravensara, Sonner. (Laurin.).

EUONYMUS, Linn. Hort. Cliff. 89 (1737). *CELASTRINEAE*, Benth. & Hook. f. i. 390.
EUONYMUS, Tourn. ex Linn. Gen. ed. I. 99 (1737).
MELANOCARYA, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 452.
VYENOMUS, Prosl, in Abh. Boehm. Ges. V. Folge, iii. (1845) 602.

EUONYMUS:—
acuminatus, *Benth. Pl. Hartw.* 59.—Mexic.
acuminatus, *Rafin. New Fl. Am.* iii. 59.—Am. bor.; Sibir.
acutangulus, Wight, Illustr. i. 178 = angulatus.
alatus, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. 1856 (1857) 185, in obs. = Thunbergianus.
alternifolius, Moench, Meth. 71 = americanus.
americanus, *Linn. Sp. Pl.* 197.—Am. bor.
angulatus, *Wight, Ic.* t. 1053.—Ind. or.
angustifolius, Gillib. Fl. Lituan. ii. 128; Vill. Hist. Pl. Dauph. ii. 540 = europaeus.
angustifolius, Pursh, Fl. Am. Sept. i. 168 = americanus.
atropurpureus, Hook. Fl. Bor. Am. i. 119 = occidentalis.
atropurpureus, *Jacq. Hort. Vindob.* ii. 55.—Am. bor.
atropurpureus, Roxb. Hort. Beng. 18; Fl. Ind. i. 627 = Hamiltonianus.
attenuatus, *Wall. Cat.* n. 4298.—Reg. Himal.
australianus, *F. Muell. Fragm.* iv. 116.—Austral.
bancanus, *Miq. Fl. Ind. Bat. Suppl.* 513.—Ins. Banca.
bullatus, *Wall. Cat.* n. 4299.—Reg. Himal.
Bungeanus, *Maxim. Prim. Fl. Amur.* 470.—Reg. Amur.
calocarpus, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 209.—Burma.
capillaceus, Wall. Cat. n. 7370 = Microtropis bivalvis.
carolinensis, Marsh. Arbust. Am. 43 = atropurpureus.
chinensis, *Lindl. in Trans. Hort. Soc.* vi. (1826) 74.—China.
chinensis, Lour. Fl. Cochinch. 156 (alien. gen.).—China.
cinereus, *Laws. in Hook. f. Fl. Brit. Ind.* i. 611.—Reg. Himal.
Colpoon, Lam. Encyc. ii. 574 = Colpoon compressum.
Colpoon, Linn. Mant. ii. 210 = Cassine capensis.
crenulatus, *Wall. Cat.* n. 4297.—Ind. or.
cuneatus, *Rafin. New Fl. Am.* iii. 59.—Am. bor.
cuneifolius, *Wright, ex A. Gray, in Mem. Am. Acad.* N. S. viii. (1861) 171.—Cuba.
dichotomus, *Heyne, ex Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 410.—Ind. or.
echinatus, *Wall. l. c.*—Reg. Himal.
effusus, *Vis. in Act. Ist. Ven. Sc.* Ser. III. iv. (1858–59) 141.—Hab.?
europaeus, *Linn. Sp. Pl.* 197.—Europ.; As. Min.
fibrilliferus, *Planch. & Miq. in Bull. Soc. Nat. Moscu.* (1859) 335.—Reg. Caucas.
filiformis, Wall. Cat. n. 4295 = Microtropis bivalvis.
fimbriatus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 408.—Reg. Himal.
floribundus, Sterv. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 122 = europaeus.
Forbesii, Hance, in Journ. Bot. xvii. (1880) 259 = Bungeanus.
friculus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 409.—Reg. Himal.
garcinifolius, Roxb. Hort. Beng. 18; Fl. Ind. i. 628 = Microtropis discolor.
gibber, *Hance, in Journ. Bot.* xx. (1882) 77.—China.
gibbiflora, Maack, ex Trautv. in Act. Hort. Petrop. viii. (1883) 197, aphalm. = Lamiera chrysantha.
glaber, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 628.—Reg. Himal.
glaucus, Teroc. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 599 = angulatus.
Goughii, Wight, Ic. t. 215 = indicus.
gracilis, Sieber. Cat. Pl. Jap. et Chin. (1836) 98 = radicans.
grandiflorus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 404.—Reg. Himal.
grandis, *Pall. Reise Statth. Russ. Reich.* i. 317.—Astrakh.
Griffithii, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 78.—Burma.
gyrosus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 408 = Elaeodendron glaucum.
Hamiltonianus, *Wall. l. c.* 408.—Reg. Himal.
hederaceus, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 883.—Hongkong.
heterophyllus, *Rafin. New Fl. Am.* iii. 59.—Am. bor.
Horsfieldii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 598.—Java.
indicus, *Heyne, ex Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 409.—Ind. or.
inermis, Forsk. Fl. Aegypt. Arab. 204 = Ochna Wightiana.

EUONYMUS:—
japonicus, *Linn. f. Suppl.* 154; *Thunb. Fl. Jap.* 100. —China; Japon.
japonicus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 405 = pendulus.
javanicus, *Blume, Bijdr.* 1146.—Burma; Malaya.
lacerus, Buch.-Ham. in D. Don, Prod. Fl. Nep. 191 = grandiflorus.
laetus, Wall. Cat. n. 4294 = Microtropis bivalvis.
latifolius, S. G. Gmel. Reise, i. 130; Gillib. Fl. Lituan. ii. 129 = europaeus.
latifolius, A. Gray, in Mem. Am. Acad. N. S. vi. (1857) 384 = oxyphyllus.
latifolius, Marsh. Arbust. Am. 44 = atropurpureus.
latifolius, *Mill. Gard. Dict.* ed. VIII. n. 3.—Europ.; As. Min.
lanciflorus, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 86 = oxyphyllus.
laxiflorus, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 883.—China.
leiophloeus, Sterv. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 122 = latifolius.
lencocarpos, *Boj. Hort. Maurit.* 68, nomen.—Maurit.
longifolius, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 882.—Hongkong.
longifolius, *Medic. Beobacht.* (1782) 311.—Hab.?
lucidus, D. Don, Prod. Fl. Nep. 191 = pendulus.
Manckii, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1857) 356.—Reg. Amur.
macropterus, *Rupr. l. c.* 359.—Reg. Amur.
Maximowiczianus, Siebold, in Verh. Batav. Gen. xii. (1830) 49 = Sieboldianus.
Maximowiczianus, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 188, nomen.—Podol.
medius, *Kit. in Linnaea*, xxxii. (1863) 641.—Hungar.
melanostictus, *Franch. & Sav. Enum. Pl. Jap.* ii. 312. —Japon.
mexicanus, *Benth. Pl. Hartw.* 86.—Mexic.
microcarpus, Bunge, Enum. Pl. Chin. Bor. 14 = Bungeanus.
microcarpus, D. Don, Prod. Fl. Nep. 191 = fimbriatus.
multiflorus, Opiz, ex Bercht. Oek.-Techn. Fl. Boehm. ii. 153 = europaeus.
muricatus, Rafin. New Fl. Am. iii. 59 = americanus.
myrtifolius, *Boj. Hort. Maurit.* 68, nomen.—Borbon.
nanus, *Bieb. Fl. Taur. Casuc.* iii. 160.—Reg. Caucas.; As. temp.
nipponicus, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 447.—Japon.
nitidus, Benth. in Hook. Lond. Journ. Bot. i. (1842) 488 = chinensis.
obovatus, Nutt. *Gen.* Am. i. 155 = americanus.
occidentalis, *Nutt. ex Torr. in Pacif. Rail. Rep.* iv. 74. —Am. bor. occ.
ovatus, Wall. Cat. n. 4300 = fimbriatus.
oxyphyllus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 86.—Japon.
paniculatus, *Wight, ex Laws. in Hook. f. Fl. Brit. Ind.* i. 609.—Ind. or.
parviflorus, *Hemsl. Diagn. Pl. Nov.* 6.—Nicaragua.
pauciflorus, *Maxim. Prim. Fl. Amur.* 74.—Reg. Amur.
pendulus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 406, in textu.—Reg. Himal.
pinnatus, Mill. Gard. Dict. ed. VIII. n. 4 = Trichilia hirta.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 451.—China.
pterocladus, Hochst. ex Laws. in Hook. f. Fl. Brit. Ind. i. 610 = angulatus.
pubescens, Sterv. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 121, in textu = europaeus.
punctatus, Wall. Cat. n. 4286 = nitidus.
quadrigonus, Guuldenst. ex Ledeb. Fl. Ross. i. 497 = europaeus.
recurvus, *Sieb. Fl. Ind. Bat. Suppl.* 513.—Sumatra.
revolutus, *Wight, Illustr.* i. 178.—Zeylan.
rigidus, Walp. Rep. i. 833, aphalm. = Gymnosporia Wallichiana.
rosmarinifolius, *Vis. in Act. Ist. Ven. Sc.* Ser. III. iv. (1858–59) 141.—Hab.?
sachalinensis, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 446.—As. or.
sarmentosus, G. Don, in Lond. Hort. Brit. 94; et Gen. Syst. ii. 5 = americanus.
scandens, R. Grah. in Edinb. N. Phil. Journ. (Jan.–Mar. 1827) 386 = echinatus.

EUONYMUS:—
scandens, Hort. ex G. Don, in Lond. Hort. Brit. 84 = americanus.
schensianus, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 444.—China.
Schottii, *Ettingsh. ex Vis. in Act. Ist. Ven. Sc.* Ser. III. iv. (1858–59) 140.—Hab.?
sclerocarpus, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 209 = Glyptopetalum sclerocarpum.
Semenovii, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) i. 557.—As. centr.
semperviens, Marsh. Arbust. Am. 44 = americanus.
sempervirens, *Rupr. ex Boiss. Fl. Orient.* ii. 10.—Reg. Caucas.
serratifolius, *Bedd. For. Man. Bot.* p. lxiv.—Ind or.
serrulatus, Wall. Cat. n. 4296 = glaber.
Sieboldianus, *Blume, Bijdr.* 1147.—Japon.
sphaerocarpus, Hassk. Cat. Hort. Bog. Alt. 229 = javanicus.
subtriflorus, Blume, Bijdr. 1147 = Thunbergianus.
sumatranus, Miq. Fl. Ind. Bat. i. ii. 589 = javanicus.
thesefolius, *Wall. Cat.* n. 4293.—Reg. Himal.
Thunbergianus, *Blume, Bijdr.* 1147.—As. or.
Thwaitesii, *Laws. in Hook. f. Fl. Brit. Ind.* i. 608.—Zeylan.
timorensis, Zipp. ex Span. in Linnaea, xv. (1841) 186 = javanicus.
Tinus, Buch.-Ham. in Wall. Cat. n. 4292 = Elaeodendron glaucum.
tingens, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 406.—Reg. Himal.
Tobira, Thunb. Fl. Jap. 99 = Pittosporum Tobira.
tristis, Salisb. Prod. 142 = atropurpureus.
usuriensis, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 449.—Mandschuria.
vaguus, Wall. Cat. n. 4284, partim = echinatus.
vagans, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 412 = Reg. Himal.
velutinus, *Finch. & Hey. in Bull. Soc. Nat. Mosc.* (1886) 337.—Reg. Caucas.
verrucosus, *Scop. Fl. Carn.* ed. II. i. 166.—Europ. or.; As. Min.
Vidalii, *Franch. & Sav. Enum. Pl. Jap.* ii. 312.—Japon.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 1 = europaeus.
Walkerii, *Wight, Illustr.* i. 178.—Zeylan.
zeylanicus, Moon, Cat. Pl. Ceyl. 17 = Walkerii.

EUOPIS, Bartl. Ord. 148 (1830) = EVOPIS, Cass. = **Berkheya**, Ehrh. (Compos.).

EUOSMA, Andr. Bot. Rep. t. 620 (1811) = **Logania**, K. Br.
albiflora, Andr. l. c. = L. floribunda.

EUOSMA, Willd. ex Schult. Mant. iii. 6, 138 (1827) = sec.

EUOSMIA, Humb. & Bonpl. Pl. Aequin. ii. 165 (1812) (Evosmia). *RUBIACEAE*, Benth. & Hook f. ii. 71.
aggregata, Spreng. Syst. i. 417 (non generis).—Peruv.
canipensis, *Humb. & Bonpl. Pl. Aequin.* ii. 165.—Venezuela.
corymbosa, Benth. in Hook. Journ. Bot. iii. (1841) 218 = Bothriospora corymbosa.
obovata, Spreng. Syst. i. 417 (= *Higginsia obovata*).—Peruv.
verticillata, Spreng. l. c. (= *Higginsia verticillata*).—Peruv.

EUOSMUS, Nutt. Gen. Am. i. 258 (1818) = **Lindera**, Thunb. (Laurin.).
australis, Nutt. l. c. 259 = L. Benzoin.
alticola, Nutt. l. c. = Sassafras alticolum.
Benzoin, Nutt. l. c. = L. Benzoin.
Diospyros, Nutt. l. c. = L. melissaefolia.
gymnicola, Nutt. l. c. = L. gymnicola.
Sassafras, Nutt. l. c. = Sassafras officinale.

EUOTHONAEA, Reichb. f. in Bot. Zeit. x. (1852) 773 = **Maxillaria**, Lindl. (Orchid.).
imbricata, Reichb. f. l. c. = M. imbricata.
oppositifolia, Reichb. f. l. c. = M. oppositifolia.

EUPARAEA, Steud. Nom. ed. II. i. 606 (1840) = sec.

EUPAREA, Banks & Soland. ex Gaertn. Fruct. i. 380, t. 80 (1788) = **Anagallis**, Tourn. (Primul.)
amoena, Gaertn. Fruct. i. 380. t. 80 = A. alternifolia.
bractealis, Bando, in Ann. Sc. Nat. Sér. II. xx. (1843) 345 = A. tenella.
chilensis, Bando, l. c. = A. alternifolia.
linarina, Bando, l. c. = A. alternifolia.
parvula, Bando, l. c. = A. alternifolia.

EUPATORIUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 247 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 245.
AGERATOPSIS, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 246 (1873).
BATSCHIA, Moench, Meth. 567 (1794).
BOLBOSTYLIS, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 467.
CAMPULOCLINIUM, DC. Prod. v. 136 (1836).
CHROMOLAENA, DC. l. c. 138 (1836).
CHRONE, Dulac, Fl. Hautes-Pyr. 512 (1867).
CONOCLINIUM, DC. Prod. v. 125 (1836).
CRITONIA, DC. l. c. 140 (1836).
DISYNAPHIA, DC. l. c. vii. 267 (1838).
KRIGIOPPUS, Hort. ex Lond. Encyc. Fl. 688 (1829).
GYPTIS, Cass. in Bull. Soc. Philom. (1818) 139; et in Dict. Sc. Nat. xvi. 10 (1820); xx. 177 (1821).
HABERLEA, Pohl, ex Baker, in Mart. Fl. Bras. vi. 11. 341 (1876).
HEBECLINIUM, DC. Prod. v. 126 (1836).
HETEROLAENA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 245 (1873).
KYRSTENIA, Neck. Elem. i. 81 (1790).
OOCLIDIUM, DC. Prod. v. 128 (1836).
OSMIA, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 251.
PRAXELIS, Cass. in Dict. Sc. Nat. xliii. 261 (1826).
TRAGANTHES, Wallr. Sched. Crit. i. 456, in nota (1822).
WIKSTROMIA, Spreng. Syst. iii. 484 (1826).
acuminatum, D. Don, Prod. Fl. Nep. 170 = Vernonia divergens!
acuminatum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1885) 241 = serrulatum.
acuminatum, *H. B. & K. Nov. Gen. et Sp.* iv. 107.—N. Granat.
acuminatum, Spreng. ex DC. Prod. v. 83 = Vernonia saliguа.
acuminatum, Wall. ex DC. l. c. 82 = Vernonia arborea!
adenantium, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 477.—Bras.
adenachnium, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 75.—Mexic.
adenanthum, *DC. Prod.* v. 164.—Bras.
adenanthum, Gardn. ex Baker, in Mart. Fl. Bras. vi. 11. 214 = Trichogonia Gardneri.
adenochaetium, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 91.—Mexic.
adenolepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. 11. 393.—Bras.
adenophorum, Hort. Berol. ex Kunth, Ind. Sem. Hort. Berol. (1847) 12 = trapezoideum.
adenophorum, Spreng. Syst. iii. 420 = glandulosum.
adenospermum, *Sch. Bip. in Seem. Bot. Voy. Herald*, 299.—Mexic.
affine, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1885) 240 = conyzoides.
affine, *H. B. & K. Nov. Gen. et Sp.* iv. 126.—Peruv.
affine, Malt. ex DC. Prod. vii. 66 = Perezia hebeclada.
affine, *Wender. Ind. Sem. Hort. Mark.* (1841); *et in Linnaea*, xvi. (1842) *Litt.* 111.—Hab.?
africanum, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 301.—Afr. trop.
ageratifolium, *DC. Prod.* v. 178.—Am. bor.; Ind. occ.
ageratoides, Bert. ex DC. Prod. v. 178 = ageratifolium.
ageratoides, *Linn. f. Suppl.* 358.—Am. bor.
alienatum, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 445 = monardifolium.
alkinense, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 92.—Mexic.
album, *Hemsl. l. c.*—Mexic.
album, *Linn. Mant.* i. 111.—Am. bor.
album, Thunb. Fl. Jap. 306 = cannabinum.
alpestre, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 118 = serratum.
altervisum, DC. Prod. v. 144 = laevigatum.

EUPATORIUM:—
alternifolium, Ard. Animad. ii. 40 = Kuhnia eupatorioides.
alternifolium, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. 11. 338.—Bras.
altissimum, *Linn. Sp. Pl.* 837.—Am. bor.
altissimum, Linn. Syst. ed. XII. 587; et ex Murr. ed. XIII. 619 = ageratoides.
amarum, Vahl, Symb. Bot. iii. 98 = Mikania amara.
ambiguum, Hook. Comp. Bot. Mag. i. (1885) 96 = serotinum, semiserratum.
amblyolaenum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 11. 332 = bupleurifolium.
amoenum, Pursh, Fl. Am. Sept. ii. 514 = purpureum.
amphidyctium, *DC. Prod.* v. 168.—Bras.
amplifolium, *A. Gray, in Proc. Am. Acad.* xv. (1880) 28.—Mexic.
amplum, *Benth. Pl. Hartw.* 200.—N. Granat.
amygdalinum, *Lam. Encyc.* ii. 408.—Am. trop.
anamensens, Steud. Nom. ed. II. i. 606, sphalm. = aromatisans.
anethifolium, *DC. Prod.* v. 182.—Bras.
Angosturae, *Polak. in Linnaea*, xli. (1877) 575.—Costa Rica.
angulicaule, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. 11. 287.—Bras.
angustifolium, *Spreng. Syst.* iii. 415.—N. Granat.
angustissimum, *Spreng. ex Baker, in Mart. Fl. Bras.* vi. 11. 325.—Bras.
apiculatum, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 476.—Bras.
appendiculatum, Less. ex Baker, in Mart. Fl. Bras. vi. 11. 365 = hecantanthum.
arallaefolium, *Less. in Linnaea*, vi. (1831) 408.—Mexic.
arboreum, *H. B. & K. Nov. Gen. et Sp.* iv. 131.—Ecuador.
arborescens, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 129 = Vernonia cinerea.
arbutifolium, *Benth. Pl. Hartw.* 199.—N. Granat
Arechtitae, *Baker, in Journ. Bot.* xvi. (1878) 78.—Reg. Argent.
arcolare, *DC. Prod.* v. 169.—Mexic.
arcolatum, *Perr. ex DC. l. c.* 12 = Ethulia conyzoides.
argentinum, Wall. Cat. n. 3166 = cannabinum.
argutum, *H. B. & K. Nov. Gen. et Sp.* iv. 121.—Mexic.
Armani, Balb. Pl. Kar. Hort. Taur. (1810) 27. t. 6 = Clibadium rotundifolium.
Arnottianum, *Griseb. in Goett. Abh.* xix. (1874) 167.—Reg. Argent.
Arnottii, *Baker, in Mart. Fl. Bras.* vi. 11. 323.—Bras.
aromaticum, *Linn. Sp. Pl.* 839.—Am. bor.
aromaticum, Sieber, ex Steud. Nom. ed. II. i. 606 = Brickellia diffusa.
aromatisans, *DC. Prod.* v. 150.—Ind. occ.
arrayanum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 489.—Bras.
artemisiifolium, *Griseb. in Goett. Abh.* xxiv. (1879) 171.—Reg. Argent.
ascendens, Mart. ex Baker, in Mart. Fl. Bras. vi. 11. 344 = decumbens.
ascendens, *Sch. Bip. ex Baker, l. c.* 296.—Bras.
Aschenbornianum, *Schau. in Linnaea*, xix. (1847) 720.—Mexic.
asclepiadeum, DC. Prod. v. 148 = bupleurifolium.
asperifolium, Sch. Bip. in Linnaea, xxx. (1859–60) 182 = monardifolium.
aspermium, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. 11. 292.—Bras.
asperulaeosum, *Baker, l. c.* 342.—Bras.
asperum, Roxb. Hort. Beng. 61; Fl. Ind. iii. 415 = Vernonia Roxburghii.
atriplicifolium, Hort. ex Lam. Encyc. ii. 407 = repandum.
atriplicifolium, *Vahl, in West, Bidr. Ste-Croix*, 302; *Symb. Bot.* iii. 96.—Ind. occ.
auriculatum, Hort. ex Lam. Encyc. ii. 411 = Senecio deltoideus.
auriculatum, *Vahl, Symb. Bot.* iii. 95.—Bras.
australe, Bartl. Ind. Sem. Hort. Gotting. (1837); et in Linnaea, xii. (1838) Litt. 81 = Bartlingianum.
australe, Thunb. Fl. Bras. Den. ii. 26 = laevigatum.
axillare, [Mœnch & Sesse,] ex DC. Prod. v. 138 = Brickellia californica.
axilliflorum, *Griseb. in Goett. Abh.* xix. (1874) 169.—Reg. Argent.
Ayapana, Vent. Jard. Malm. t. 8 = triplinerve.
ayapanoides, *Grisel. Cat. Pl. Cub.* 146.—Cuba.

EUPATORIUM:—
azangaroënse, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
azureum, *DC. Prod.* v. 168.—Mexic.
bacchariifolium, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 118.—Bras.
baccharoides, *H. B. & K. Nov. Gen. et Sp.* iv. 132.—N. Granat.
Baciennum, *DC. Prod.* v. 157.—Bras.
Ballii, *Oliver, in Hook. Ic. Pl.* t. 1463.—Peruv.
ballotaefolium, *H. B. & K. Nov. Gen. et Sp.* iv. 121.—Bras.; N. Granat.
Barclayanum, Benth. Bot. Voy. Sulph. 119 = amygdalinum
Bartlingianum, *Steud. Nom.* ed. II. i. 606.—Chili.
bartsiaefolium, *DC. Prod.* v. 147.—Bras.
bartsioides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 11. 295 = prasc.
bellidifolium, *Benth. Pl. Hartw.* 48.—Mexic.
Benthamii, *Klatt, in Leopoldina*, xx. (1884) 90.—Mexic.
Berlandieri, DC. Prod. v. 167 = ageratifolium.
Berlandieri, A. Gray, in Torr. Bot. Mex. Bound. 76 = occidentale.
Bertersanum, Colla, in Mem. Acc. Torin. xxxiii. (1829) 130 = guadalupense.
Bertholdii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 299.—Mexic.
betonicaefolium, *Mill. Gard. Dict.* ed. VIII. n. 9.—Mexic.
betonicaeforme, *Baker, in Mart. Fl. Bras.* vi. 11. 362.—Bras.
betonicum, *Hemsl. Biol Centr. Am. Bot.* ii. 93.—Am. bor. occ.
bicolor, Lam. ex DC. Prod. vi. 326 = Gynoxis buxifolia.
Bigelovii, *A. Gray, in Torr. Bot. Mex. Bound.* 75.—Mexic.
Billbergianum, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 184.—Panama.
birmanicum, DC. Prod. v. 179 = cannabinum.
Blanchetii, *Sch. Bip. in Flora*, xxxiii. (1850) 30.—Bras.
blepharilepis, *Sch. Bip. in Seem. Bot. Voy. Herald*, 300.—Mexic.
bogotense, *DC. Prod.* v. 181.—N. Granat.
brachiatum, Sw. ex Wiksr. in Vet. Acad. Handl. Stockh. (1827) 71 = odoratum.
bracteatum, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 472.—Bras.
bracteatum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1885) 239 = tozziaefolium.
brasilianum, Spreng. Syst. iii. 417 = Brickellia pinifolia.
Braunii, *Polak. in Linnaea*, xli. (1877) 576.—Costa Rica.
brevipes, *DC. Prod.* v. 168.—Mexic.
brevipetiolatum, *Klatt, in Leopoldina*, xx. (1884) 90.—Mexic.
brevipetiolatum, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. 11. 336, in obs.—Bras.
brevisetum, *DC. Prod.* v. 169.—Mexic.
Brickellia, DC. l. c. 182 = Brickellia cordifolia.
Brunetii, *A. Gray, Syn. Fl. N. Am.* i. 11. 96.—Am. bor. occ.
brunneolum, *Baker, in Mart. Fl. Bras.* vi. 11. 338.—Bras.
buddleaefolium, *Benth. Pl. Hartw.* 185.—Peruv.
bunifolium, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1885) 240 = pinnatifidum.
bupleurifolium, *DC. Prod.* v. 148.—Bras.
Burchellii, *Baker, in Mart. Fl. Bras.* vi. 11. 356.—Bras.
Bustamanta, *DC. Prod.* v. 168.—Mexic.
cacalioides, *H. B. & K. Nov. Gen. et Sp.* iv. 130.—Ecuador.
caduciseta, *DC. Prod.* v. 166.—N. Granat.
calaminthaefolium, *H. B. & K. Nov. Gen. et Sp.* iv. 129. t. 343.—Mexic.
callilepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. 11. 336.—Bras.
calocephalum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 388 = ivaefolium.
calycinum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 11. 336 = acq.
calyculatum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1885) 242.—Bras.
campestre, *DC. Prod.* v. 168.—Bras.
candicans, Wall. Cat. n. 3999 = Lescomeris spectabilis.

EUPATORIUM:—
Candollesanum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1836) 348.—Bras.
Candollesanum, Hook. & Arn. l. c. partim = betonicæformæ.
canescens, Orteg. Hort. Matr. Dec. 84 = Kuhnia rosmarinifolia.
canescens, Vahl, in West, Bidr. Ste-Croix, 302; Symb. Bot. iii. 95 = sinuatum.
cannabifolium, Salisb. Prod. 187 = seq.
cannabinum, *Linn. Sp. Pl.* 838.—Europ.; As. bor.
cannabis-folium, Gilib. Fl. Lituan. i. 272 = præc.
capense, Spreng. f. Tent. Suppl. 22 = Vernonia mespilifolia.
capillare, *Baker, in Mart. Fl. Bras.* vi. ii. 341.—Bras.
capillare, Desv. in Ham. Prod. Pl. Ind. Occ. 51 = Brickellia diffusa.
capilliper, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 79 = guadalupense.
carolinianum, *Hort. Vind. ex Steu. in Bull. Soc. Nat. Mosc.* xxix. (1856) ii. 372.—Hab.?
cazinæfolium, Bertol. Misc. Bot. v. 17. t. 6. f. a = cuneifolium.
caucasicum, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) ii. 371.—Reg. Caucas.
ceanothifolium, Muhl. ex Willd. Sp. Pl. iii. 1755 = aromaticum.
ceanothifolium, Spreng. Syst. iii. 418 = triste.
celebicum, Blume, Bijdr. 908 = Vernonia arborea.
celtidifolium, *Lam. Encyc.* ii. 406.—Ind. occ.; N. Granat.
cerasifolium, *Baker, in Mart. Fl. Bras.* vi. ii. 308 — Bras.
ceratophyllum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1838) 340.—Bras.; Reg. Argent.
chamæsdrifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 113. —Peruv.
charopadifolium, Poir. Encyc. Suppl. ii. 606 = Mikania scandens.
Chilco, *H. B. & K. Nov. Gen. et Sp.* iv. 135.—Peruv.
chilense, *Benth. in Merc. Chil.* (1829); *ex Am. Journ. Sc. sc.* (1831) 260.—Chil.
chilense, Molina, Sagg. Chile, 142, 354 = Flaveria Contrayerba.
chinense, *Linn. Sp. Pl.* 837.—China; Japon.
chlorolepis, *Baker, in Mart. Fl. Bras.* vi. ii. 357.—Bras.
Christianum, *Baker, l. c.* 298.—Bras.
cicutricaulon, Sch. Bip. ex Baker, l. c. 326 = angustissimum.
ciliatum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 340 = conyzoides.
ciliatum, *Less. in Linnæa*, vi. (1831) 404.—Mexic.
cinerascens, Wall. Cat. n. 3066 = Microglossa albescens.
cinereo-viride, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 294.—Bras.
cinereum, *Baker, in Journ. Bot.* xx. (1882) 226.—Bras.
cinereum, *Griseb. Fl. Brit. Ind.* 359.—Ind. occ.
cinereum, Linn. f. Suppl. 354 = Pteronia incana.
cinereum, *Raffin. New Fl. Am.* iv. 80.—Am. bor.
cinereum, Wall. ex Steud. Nom. ed. II. i. 606 = Microglossa albescens.
Clausenii, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 443 = stachyophyllum.
clavulatum, *Griseb. in Goett. Abh.* xix. (1874) 168.—Reg. Argent.
clematidum, *Griseb. l. c.* xxiv. (1879) 172.—Reg. Argent.
clematidoides, Less. ex Baker, in Mart. Fl. Bras. vi. ii. 256 = Mikania phaseolodes.
clematidoides, Sch. Bip. in Pollichia, xxii.–xxiv. (1866) 258 = Reevesii.
Clematitis, DC. Prod. v. 144 = conyzoides.
coahuilense, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 205.—Mexic.
coelestinum, *Linn. Sp. Pl.* 838.—Am. bor.; Ind. occ.
cognatum, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 13.—Mexic.
colliæum, *DC. Prod.* v. 164.—Mexic.
coloratum, Spreng. ex DC. Prod. v. 143 = Microglossa albescens.
coloratum, Willd. Sp. Pl. iii. 1768 = Vernonia senegalensis.
compactum, *Benth. Bot. Voy. Sulph.* 112.—Peruv.
compactifolium, Walt. Fl. Carol. 199 = corymbopilifolium.

EUPATORIUM:—
compressum, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 478 = Symphyopappus polystachyus.
conyzoidifolium, DC. Prod. v. 148 = serratum.
concinnum, *DC. l. c.* 156.—Bras.
concinnum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1836) 339 = ivæfolium.
condensatum, Reichb. ex Less. in Linnæa, iv. (1839) 337 = Oliganthes condensata.
confertifolium, *Klatt, in Abh. Naturf. Ges. Halle.* xv. (1882) 194.—N. Granat.
confertum, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 116 = serratum.
congestum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1836) 339 = rozzioliifolium.
conglobatum, *DC. Prod.* v. 165.—Bras.
connatum, Michx. Fl. Bor. Am. ii. 99 = perfoliatum.
conocliniifolium, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 478 = Trichogonia podocarpa.
consanguineum, *DC. Prod.* v. 166.—Bras.
conspicuum, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 13.—Mexic.
conyzoides, Mill. Gard. Dict. ed. VIII. n. 14 = Vernonia arborescens.
conyzoides, *Vahl, Symb. Bot.* iii. 96.—Am. bor. et trop.
cordatum, Barm. f. Fl. Ind. 178 = Mikania scandens.
cordatum, Walt. Fl. Carol. 199 = aromaticum?
cordifolium, *Sw. Prod. Veg. Ind. Occ.* 111.—Jamaica.
cordiforme, Poir. Encyc. Suppl. ii. 600 = aromaticum.
coriaceum, *Scheele, in Linnæa*, xviii. (1844) 457.—Bras.
coriaceum, *Spreng. Syst.* iii. 410.—Ins. S. Doming.
coriaceum, Vahl, Symb. Bot. iii. 94 (= *Decaneurum ferdinandifilium*).—Ins. Borbon.
cornifolium, Willd. ex Less. in Linnæa, iv. (1829) 337 (= *Odontoloma acuminatum*).—Venezuela.
coronopifolium, *Willd. Sp. Pl.* iii. 1750.—Am. bor.; Ind. occ.
corsicum, Req. ex Loisel. in Mém. Soc. Linn. Par. vi. (1827) 428 = cannabinum.
corymbiferum, *Griseb. Fl. Brit. W. Ind.* 361.—Jamaica.
corymbosum, *Asbl. Pl. Gui.* ii. 799.—Guian.
cotinifolium, *Willd. Phyt.* 11. t. 7. f. 1.—Ins. Martinic.
crassifolium, Raffin. Fl. Lndov. 62 (sp. falsa).
crassifolium, Shuttl. ex A. Gray, Syn. Fl. N. Am. i. ii. 97 = mikanioides.
crenatum, DC. Prod. v. 191 = Mikania stipulacea.
crenatum, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 441 = squalidum.
crenatum, Gomez, in Mem. Ac. Lisboa, iii. (1812) Mem. Corresp. 28 = Mikania cordifolia.
crenulatum, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 474 = Trichogonia hirtiflora.
crithmifolium, *Griseb. in Goett. Abh.* xix. (1874) 169.—Reg. Argent.
crithmifolium, Less. ex Baker, in Mart. Fl. Bras. vi. ii. 262 = Mikania verticillata.
critonioides, *Pœpp. et Endl. in Nov. Act. Phys.-Med. Acad. Leopold.* xiii. (1827) 1.—Venezuela.
cryptanthum, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 302.—Bras.
cubense, DC. Prod. v. 173 = villosum?
cubense, Pers. Syn. ii. 402 = Kuhnia rosmarinifolia.
cunninghamii, DC. Prod. v. 149 = Symphyopappus cuneatus.
cuneatum, Engelm. in Torr. & Gray, Fl. N. Am. ii. 88 = perfoliatum.
cuneifolium, DC. Prod. v. 177 = semiserratum.
cuneifolium, *Willd. Sp. Pl.* iii. 1753.—Am. bor.
cuspidatum, Pav. ex A. Gray, in Proc. Am. Acad. xvii. (1881–82) 206 = Barrettia Pavonii.
cuspidatum, Spreng. ex DC. Prod. v. 82 = Vernonia acuminata.
cuspidatum, Willd. ex Less. in Linnæa, iv. (1829) 315 = Oliganthes discolor.
cydoniæfolium, Willd. ex Less. l. c. 269 = Vernonia dichocarpa.
cylindrocephalum, *Sch. Bip. in Flora*, xxxiii. (1850) 30, nomen; *Baker, in Mart. Fl. Bras.* vi. ii. 263.—Bras.
cynanchifolium, *DC. Prod.* v. 172.—Cuba.
Dalea, H. B. & K. Nov. Gen. et Sp. iv. 106 (= *Critonia Faradæalea*).—Cuba.
Dalea, *Lingt. Syst.* ed. X. 1304.—Jamaica; Cuba.
delaeides, *Hemsl. Biol. Centr. Am. Bot.* ii. 98.—Mexic.

EUPATORIUM:—
decumbens, *DC. Prod.* v. 154.—Peruv.
decipiens, *Baker, in Mart. Fl. Bras.* vi. ii. 347.—Bras.
decipiens, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 340 = Ophryosporus triangularis.
decumbens, *Baker, in Mart. Fl. Bras.* vi. ii. 344.—Bras.
decussatum, Sieber, ex Steud. Nom. ed. II. i. 607 = cordifolium.
deltoideum, Raffin. ex DC. Prod. v. 136 = coelestinum.
deltoidesm, *Zucc. Hort. Schœnb.* iii. 63. t. 308.—Mexic.
deltoideum, Poepp. ex Spreng. Syst. iii. 415 = Pœppigii.
deltoideum, Steud. Nom. ed. II. i. 607, sphalm. = coelestinum.
dendroides, *Spreng. Syst.* iii. 415.—Am. austr.
denticulatum, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 281 = multiflosculosum.
dentatum, *Benth. Pl. Hartw.* 306.—N. Granat.
dentatum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 443.—Bras.
denticulatum, Vahl, Symb. Bot. iii. 98 = Mikania scandens.
dichotomum, Sch. Bip. ex Miq. Stirp. Surin. Sel. 184 = squalidum.
dicline, Endlicher, in Trans. Linn. Soc. xx. (1851) 63 = cannabinum.
diffusum, Vahl, Symb. Bot. iii. 94 = Brickellia diffusa.
dimorpholepis, *Baker, in Mart. Fl. Bras.* vi. ii. 331.—Bras.
discolor, *DC. Prod.* v. 161.—Bras.
discolor, Spreng. Syst. iii. 412 = Vernonia discolor.
disjunctum, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 117 = pyrifolium.
dissectum, *Benth. Bot. Voy. Sulph.* 113.—Mexic.
dissectum, A. Gray, in Proc. Am. Acad. xviii. (1883) 100 = Greggii.
dissolvens, *Baker, in Mart. Fl. Bras.* vi. ii. 308.—Bras.
divaricatum, Berg. Desc. Pl. Cap. 229; Thunb. Prod. Pl. Cap. 142 = Pteronia divaricata.
divaricatum, Sch. Bip. in Linnæa, xxii. (1849) 871 = guadalupense.
divergens, Less. l. c. v. (1880) 188 = conyzoides.
divergens, Roxb. Hort. Beng. 61; Fl. Ind. iii. 414 = Vernonia divergens.
diversifolium, Schrad. Ind. Sem. Hort. Gotting. (1829) = pycnocephalum.
dodoneæfolium, DC. Prod. v. 161 = amygdalinum.
dodonæifolium, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1838) 44.—Reg. Argent.
Dombeyanum, *DC. Prod.* v. 167.—Am. austr.
domingense, *Spreng. Syst.* iii. 412.—Ins. S. Doming.
Donianum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 348 = macrocephalum, leptolepis.
dryadeum, DC. Prod. vii. 269 = macrophyllum.
dubium, Pohr. Encyc. Suppl. ii. 606 = purpureum.
dumosum, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 81; *et in Linnæa*, xxxiv. (1865–66) 585.—Bolivia.
duodecimflorum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 318 = corgale.
dyctiophyllum, *DC. Prod.* v. 158.—Bras.
Ehrenbergii, *Hemsl. Biol. Centr. Am. Bot.* ii. 94.—Mexic.
elæagnifolium, Hort. Calc. ex DC. Prod. v. 32 = Vernonia elæagnifolia.
elatum, Salisb. Prod. 187 = altissimum.
elatum, *Steetz, in Seem. Bot. Voy. Herald*, 146.—Panama.
elegans, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 467 = amygdalinum.
elegans, *H. B. & K. Nov. Gen. et Sp.* iv. 133.—Ecuador.
elegans, Steud. Nom. ed. II. i. 607 = Liatris elegans.
elliptícum, *DC. Prod.* v. 154.—Ins. Penang.
elliptícum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 340.—Reg. Argent.
elongatum, Hook. f. Am. l. c. 241 = oblongifolium.
Engelmannianum, Raegel.—Cf. Torr. & Gray, Fl. N. Am. ii. 302 = aromaticum.
ensifolium, *Griseb. in Goett. Abh.* xxiv. (1879) 170.—Reg. Argent.
epilobioides, *H. B. & K. Nov. Gen. et Sp.* iv. 126.—N. Granat.
erioclados, *DC. Prod.* v. 150.—Bras.
erigeroides, DC. l. c. 171 = corymbiflorum.

EUPATORIUM:—
- erectifolium, *DC. Prod.* v. 156.—Bras.
- erythrolepis, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 583, nomen.—Boliv.
- Espinosarum, *A. Gray, in Proc. Am. Acad.* xv. (1880) 96.—Mexic.
- *exacutidens*, G. Don, ex Baker, in Mart. Fl. Bras. vi. III. 154 = Ichthyothere Cunabi.
- exerto-venosum, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 234.—Peruv.
- *exiguum*, Klotzsch, ex Steets, in Seem. Bot. Voy. Herald, 145 = ireinoides.
- extenuatum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 440.—Bras.
- *extrorsum*, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 290 = hvezolinum.
- *falcatum*, Michx. Fl. Bor. Am. ii. 99 = purpureum.
- fasciculare, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 54.—Peruv.
- fastigiatum, *H. B. & K. Nov. Gen. et Sp.* iv. 125.—N. Granat.
- Fendleri, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 205.—N. Mexic.
- *ferruginascens*, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 348 = hebeciadum.
- ferrugineum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 443.—Bras.
- *ferrugineum*, Lahlll. Nov. Holl. Pl. ii. 88. t. 180 = Helichrysum ferrugineum.
- *Finlaysonianum*, Wall. Cat. n. 7138 = cannabinum.
- *flaccidum*, Spreng. ex DC. Prod. v. 174 = Brickellia diffusa.
- *flexuosum*, Lam. Encyc. ii. 411 = Faujasia flexuosa.
- *flexuosum*, Kozb. Hort. Beng. 61; Fl. Ind. iii. 415 = odoratum †
- *floribundum*, H. B. & K. Nov. Gen. et Sp. iv. 118 = conyzoides.
- *floridanum*, Rafin. ex Torr. & Gray, Fl. N. Am. ii. 86 = altissimum.
- fomicicaulescum, *Willd. Sp. Pl.* iii. 1760.—Am. bor.; Ind. occ.
- *fomicicaulescens*, Walt. Fl. Carol. 199 = fomicicaulescum.
- *foliolosum*, DC. Prod. v. 176 = Ophryosporus triangularis.
- *foliolosum*, Wall. ex DC. l. c. 451 = Pluchea indica.
- *Fortunei*, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 170 = japonicum.
- *Frasieri*, Pohr. Encyc. Suppl. ii. 600 = ageratoides.
- fraternum, *DC. Prod.* v. 163.—Am. austr.
- *Freyreyst*, Thunb. Fl. Bras. Dec. ii. 25 = Ophryosporus Freyreissii.
- *fruticosum*, Desf. Tabl. ed. I. 98 = Kuhnia fruticosa.
- fruticosum, *Linn. Syst.* ed. X. 1204.—Hab.†
- fruticosum, *Mill. Gard. Dict.* ed. VIII. n. 6.—Mexic.
- fuligineum, *H. B. & K. Nov. Gen. et Sp.* iv. 110.—Am. austr.
- *fulvum*, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1836) 341 = Mikania fulva.
- *fulvum*, Raln. New Fl. Am. iv. 80 (sp. falsa).
- furcatum, *Lam. Encyc.* ii. 407.—Ind. occ.
- *fuscorubrum*, Wall. Fl. Carol. 199 = purpureum.
- *gaiapathifolium*, Gard. in Hook. Lond. Journ. Bot. vi. (1847) 446 = supestre.
- Gaudichaudianum, *DC. Prod.* v. 148.—Bras.
- Gayanum, *Wedd. Chlor. Andi.* i. 210.—Peruv.
- glaberrimum, *DC. Prod.* v. 144.—Mexic.
- *glabratum*, Hort. Berol. (1839); ex Kunth, Ind. Sem. Hort. Berol. (1847) 13 = modestum.
- glabratum, *H. B. & K. Nov. Gen. et Sp.* iv. 127.—Mexic.
- *glabriusculum*, DC. Prod. v. 161 = Vasthieranum.
- glabrum, *Heyne, ex Wall. Cat.* n. 2323.—Ind. or.
- *glandulosum*, Hort. Berol. ex Kunth, Ind. Sem. Hort. Berol. (1847) 13 = cognatum.
- glandulosum, *H. B. & K. Nov. Gen. et Sp.* iv. 122.—Jamaic.; Mexic.
- *glandulosum*, Michx. Fl. Bor. Am. ii. 98 = album.
- *glandulosum*, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 314 = amygdalinum.
- *glastifolium*, Mertel. Misc. Bot. v. 16. f. 4 = Trilisa odoratissima.
- *glaucescens*, Kll. Sketch. ii. 297 = cuneifolium.
- glaucum, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 99.—Mexic.
- Glaziovii, *Baker, in Mart. Fl. Bras.* vi. II. 357.—Bras.
- *gnaphalioides*, Mlq. ex DC. Prod. v. 180 = Brickellia verbesinoides.

EUPATORIUM:—
- *gnidiomorphilum*, *Less. in Linnaea*, vi. (1831) 105.—Chili.
- *Gnkini*, F. Schmidt, ex Trautv. in Act. Hort. Petrop. viii. (1883) 432 = japonicum †
- *glomeratum*, DC. Prod. v. 154 = pallescens.
- glumaceum, *DC. l. c.* 181.—Cuba.
- glutinosum, *Lam. Encyc.* ii. 408.—Ecuador.
- gnidioides, *DC. Prod.* v. 150.—Bras.
- gnaphalioides, *DC. l. c.* 150.—Mexic.
- *Gouani*, Hort. ex Spreng. Nov. Prov. 32 = Salmea Espatoria.
- *gracile*, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 348 = glechonophyllum.
- gracile, *H. B. & K. Nov. Gen. et Sp.* iv. 124.—N. Granat.
- gracilifiorum, *DC. Prod.* v. 145.—Mexic.
- grande, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 347.—Bras.
- grandiceps, *Wright, in Sauv. Fl. Cub.* 76.—Cuba.
- grandidentatum, *DC. Prod.* v. 167.—Mexic.
- *grandiflorum*, Hook. Fl. Bor. Am. ii. 26 = Brickellia grandiflora.
- grandiflorum, *Regel, in Gartenfl.* i. (1852) 102. t. 12. —Guatemala.
- Greggii, *A. Gray, Syn. Fl. N. Am.* i. II. 102.—Am. bor.; Mexic.
- *grisco-viride*, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 353 = adamantium.
- gnadalupense, *Sprang. Syst.* iii. 414.—Ins. Guadalup.
- guapalense, *Klatt, in Leopoldina*, xx. (1884) 90.—Mexic.
- guatemalense, *Regel, Ind. Sem. Hort. Turic.* (1850) 4 *Adnot.*; *ex Walp. Ann.* ii. 816.—Guatemala.
- gymnocephalum, *Wedd. Chlor. And.* i. 216.—N. Granat.
- Haageanum, *Regel & Korm. Ind. Sem. Hort. Petrop* (1857) 40.—Hab.†
- Hsenkeanum, *DC. Prod.* v. 158.—Mexic.
- halimifolium, *L.* v. 1207.—Am. austr.
- Hartwegi, *Benth. Pl. Hartw.* 19.—Mexic.
- *hastatum*, Linn. Syst. ed. X. 1204 = Mikania hastata.
- hastile, *Sch. Bip. in Linnaea*, xiv. (1847) 719.—Mexic.
- havanense, *H. B. & K. Nov. Gen. et Sp.* iv. 128.—Cuba.
- hebebotrys, *Hemsl. Biol. Centr. Am. Bot.* ii. 95.—Mexic.
- hebeciadum, *DC. Prod.* v. 164.—Bras.
- hecatanthum, *Baker, in Mart. Fl. Bras.* vi. II. 365.—Bras.
- *hecatanthum*, Sch. Bip. in Linnaea, xxx. (1859–60) 182 = macrocephalum.
- hederaefolium, *A. Gray, Pl. Fendl.* 65.—Mexic.
- helianthifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 127.—Peruv.
- hemisphaericum, *DC. Prod.* v. 158.—Bras.
- heptanthum, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
- heterociinium, *Griseb. Fl. Brit. W. Ind.* 358.—Jamaic.
- *heterolasenum*, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 326 = serratum.
- *heterophyllum*, DC. Prod. v. 180 = cannabinum.
- heterophyllum, *A. Rich. Fl. Cub. Fanerog.* ii. 39.—Cuba.
- *hexanthum*, DC. Prod. v. 148 = bartsiaefolium.
- hirsutissimum, *Baker, in Mart. Fl. Bras.* vi. II. 311.—Bras.
- hirsutum, *Baker, l. c.* 359.—Bras.
- hirsutum, *DC. Prod.* v. 170.—Mexic.
- *hirsutum*, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 339 = bartsiaefolium.
- *hirtum*, Less. ex Baker, in Mart. Fl. Bras. vi. II. 360 = Mikania hirsutissima.
- hispidum, *Pers. Syn.* ii. 402.—Afr. trop.
- *Hookerianum*, Griseb. in Goett. Abh. xix. (1874) 166 = conyzoides.
- hormisoides, *Baker, in Mart. Fl. Bras.* vi. II. 300.—Bras.
- Horsfieldii, *Mlq. Fl. Ind. Bat.* ii. 27.—Java.
- *Houstonianum*, Linn. Sp. Pl. 836 = Mikania Houstonis.
- *Houstonis*, Hort. ex DC. Prod. v. 193 = Salmea Espatoria.
- *Houstonis*, Linn. Syst. ed. X. 1204 = Mikania Houstonis.
- *Houstonis*, Sw. Obs. Bot. 300 = Mikania Swartziana.
- hyperbolifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 118.—N. Granat.
- hypoleucum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 519.—Ind. occ.
- hyssopifolium, *Linn. Sp. Pl.* 836.—Am. bor.
- hyssopifolium, *A. Gray, in Proc. Am. Acad.* xv. (1880) 28.—Mexic.
- ianthinum, *Hemsl. Biol. Centr. Am. Bot.* ii. 96.—Mexic.
- impetiolare, *Griseb. Fl. Brit. W. Ind.* 357.—Ins Dominic.
- *inamoenum*, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 313 = Trichogonia villosa.
- *incanum*, Sch. Bip. ex Baker, l. c. 327 = halimifolium.
- incarnatum, *Wall. Fl. Carol.* 200.—Am. bor.; Mexic
- incasicum, *Wedd. Chlor. And.* i. 217.—Boliv.
- incisum, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 112 (err. typ. 105).—Guiana.
- incomptum, *DC. Prod* v. 178.—Mexic.
- inconspicuum, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 535.—Boliv.
- Indigofera, *Parodi, in Anal. Soc. Cient. Argent.* v. (1877) 38.—Paraguay.
- innumerosum, *DC. Prod.* v. 164.—Am. austr.
- *integerrimum*, Spreng. ex Baker, in Mart. Fl. Bras. vi. II. 324 = velutinum.
- integrifolium, *Dest. ex Spreng. Syst.* iii. 410.—Ins. Guadalup.
- intermedium, *DC. Prod.* v. 146.—Bras.
- inulaefolium, *H. B. & K. Nov. Gen. et Sp.* iv. 109.—Am. trop.
- *involucratum*, Sch. Bip. in Linnaea, xxii. (1849) 572 = corlaceum.
- ireinoides, *H. B. & K. Nov. Gen. et Sp.* iv. 106. t. 340.—Am trop.
- itacolumense, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 310.—Bras.
- ivaefolium, *Linn. Syst.* ed X. 1205.—Am. bor.; Ind. occ.
- ixiocladon, *Benth. ex Cornt. in Kjoeb. Vidensk. Meddel.* (1862) 77.—Am. centr.
- *ixodes*, Benth. in Hook. Journ. Bot. ii. (1840) 41 = amygdalinum.
- Jamesonii, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 169.—Peruv.
- japonicum, *Thunb. Fl. Jap.* 308.—Japon.
- *javanicum*, Blume, Bijdr. 908 = Vernonia arborea.
- Karwinskianum, *DC. Prod.* v. 163 = Mexic.
- *Kirilowii*, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 158 = Lindleyanum.
- kleinioides, *H. B. & K. Nov. Gen. et Sp.* iv. 120.—N. Granat.; Bras.; Peruv.
- *Kuhnia*, Crants, Inst. i. 264 = Kuhnia eupatorioides.
- *lactescens*, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 340 = stevlaefolium.
- laeve, *DC. Prod.* v. 169.—Bras.
- laevigatum, *Lam. Encyc.* ii. 408.—Am. trop.
- *laevigatum*, Torr. Cat. Pl. N. York, 92 = purpureum
- Lallavei, *Batll. Hist. Pl.* viii. 296.—Mexic.
- *Lamberti anum*, Wall. Cat. n. 3287 = cannabinum.
- lamiifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 113.—Ecuador.
- *lamiifolium*, Link, Enum. Hort. Berol. ii. 306 = sonchsonchifolium.
- *lanatum*, Spreng. ex DC. Prod. v. 169 = Isnia Cappa.
- *lanceolatum*, Muhl. ex Willd. Sp. Pl. iii. 1752 = laeviifolium.
- lanigerum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1836) 342.—Bras.
- lantanifolium, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 511.—Ind. occ.
- lasiocaulon, *Hook. & Arn. Bot. Beech. Voy.* 297.—Mexic.
- lasiophthalmum, *Griseb. in Goett. Abh.* xix. (1874) 167.—Reg. Argent.
- *latifolium*, Less. ex Baker, in Mart. Fl. Bras. vi. II. 297 = Mikania Argyreiae.
- latipes, *Benth. Pl. Hartw.* 200.—N. Granat.
- lavandulaefolium, *DC. Prod.* v. 154.—Peruv.
- *laxum*, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 476 = guadalupense.
- leiophyllum, *Less. in Linnaea*, vi. (1831) 402.—Mexic.
- leptocephalum, *DC. Prod.* v. 148.—Peruv.
- leptolepis, *Baker, in Mart. Fl. Bras.* vi. II. 359.—Bras.
- *leptophyllum*, DC. Prod. v. 176 = fomicicaulescum.
- *leptopodum*, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 478 = Brickellia diffusa.
- *leucanthum*, Klotzsch, ex Walp. Rep. vi. 133 = Conyza modesta, *Kunth*.
- leucocephalum, *Benth. Pl. Hartw.* 86.—Mexic.

EUPATORIUM:—
leucolepis, *Torr. & Gray, Fl. N. Am.* ii. 84.—Am. bor.
leucophyllum, *H. B. & K. Nov. Gen. et Sp.* iv. 116.—Peruv.
leucotrichium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 299 = verbenaceum.
listrideum, *DC. Prod.* v. 142.—Bras.
lithanthicum, *Sch. Bip. in Journ. Bot.* i. (1863) 234.—Cuba.
Liebmanni, *Hemsl. Biol. Centr. Am. Bot.* ii. 96.—Mexic.
Liebmannii, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 76.—Mexic.
ligulaefolium, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 342 = guidioides.
ligustrinum, DC. Prod. v. 181 = micranthum.
Lindenianum, *A. Rich. Fl. Cub. Fanerog.* ii. 42.—Cuba.
Lindleyomerianum, Scheele, in Linnæa, xxi. (1848) 599 = ageratifolium.
Lindleyanum, *DC. Prod.* v. 180.—China.
Lindleyanum, F. Muell. Fragm. v. 93 = cannabinum.
linearifolium, Michx. Fl. Bor. Am. ii. 97, partim = lucolepis.
linearifolium, Walt. Fl. Carol. 199 = hyssopifolium.
lineatum, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 336.—Bras.
linifolium, DC. Prod. v. 149 = bupleurifolium.
longepetiolatum, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 535, nomen.—Bolivia.
longicaule, *DC. Prod.* v. 178 (sp. dub.).—Nepal.
longicaule, Wall. Cat. n. 3169 = Vernonia saligna.
longipes, *A. Gray, in Proc. Am. Acad.* xv. (1880) 26.—Mexic.
longipetiolatum, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 317 = Trichogonia salviæfolia.
lonicerioides, H. B. & K. Nov. Gen. et Sp. iv. 116 = amygdalinum.
lucidum, *Ortog. Hort. Matr. Dec.* 85.—Mexic.
ludovicianum, *DC. Prod.* v. 138.—Am. bor.
Lundianum, *DC. l. c.* 170.—Bras.
lupulinum, *Baker, in Mart. Fl. Bras.* vi. ii. 301.—Bras.
lutescens, Rafin. in Med. Repos. N. York, v. (1808) 361 (sp. falsa).
lusoniense, Llanos, Fragm. 88 = tripliner've.
macranthum, *Sw. Fl. Ind. Occ.* iii. 1815.—Ind. occ.
macrocephalum, *Less. in Linnæa*, v. (1830) 126.—Bras.
macrolon, DC. Prod. v. 145 = laevigatum.
macrophyllum, *Linn. sp. Pl.* ed. ii. 1175.—Am. trop.
macropodum, *Baker, in Mart. Fl. Bras.* vi. ii. 279.—Guiana.
maculatum, Linn. Cent. Pl. i. 27; Amoen. Acad. iv. 288 = purpureum.
Mairetianum, *DC. Prod.* v. 167.—Mexic.
malvaefolium, *DC. l. c.* 160.—Mexic.
Mandonii, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 533, nomen.—Boliv.
marginatum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 54.—Peruv.
Marrubium, Walt. Fl. Carol. 199 = rotundifolium.
Martii, Mart. in Flora, xxi. (1838) ii. Beibl. 88 = suq.
Martiusii, DC. Prod. vii. 262 = aqualichum.
Maximiliani, Schrad. ex DC. l. c. v. 143 = conyzoides.
megacephalum, *Mart. ex Baker, in Mart. Fl. Bras.* vi. ii. 354.—Bras.
megalophyllum, *Klatt, in Leopoldina*, xx. (1884) 90.—Mexic.
megaphyllum, *Baker, in Mart. Fl. Bras.* vi. ii. 322.—Bras.
megapotamicum, *Spreng. Syst.* iii. 420.—Bras.
mehaudenium, *Hance, in Journ. Bot.* xxiii. (1885) 295.—China.
melanolepis, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 97.—Mexic.
melissæfolium, *D. Dietr. Syn. Pl.* iv. 1894.—Mexic.
melissæfolium, Lam. Encyc. ii. 411 (= *Nothites melissæfolia*).—Peruv.
melissoides, Willd. Sp. Pl. iii. 1754 = aromaticum.
Mendezii, *DC. Prod.* v. 160.—Mexic.
menthæfolium, *Poepp. ex Spreng. Syst.* iii. 412.—Cuba.
menthæfolium, Wall. ex C. B. Clarke, Comp. Ind. 18 = Vernonia chinensis.
microcharactium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 319 = stevisefolium.
microanthum, J. F. Gmel. Syst. 1196 = Mikania amara.
microanthum, Lag. Gen. et Sp. Pl. Nov. 25 = Ageratum corymbosum.
microanthum, *Less. in Linnæa*, v. (1830) 128.—Mexic.

EUPATORIUM:—
microcephalum, *Regel, Ind. Sem. Hort. Petrop.* (1860) 25.—N. Granat.
microphyllum, *Linn. f. Suppl.* 355.—N. Granat.
microrhombeum, Cass. in Dict. Sc. Nat. xxv. 482 = guadalupense.
millanioides, *Chapm. Fl. S. U. St.* 195.—Am. bor.
Millieti, *Steud. Nom.* ed. II. i. 606.—Mexic.
modestum, *Kunth, Ind. Sem. Hort. Berol.* (1847) 18.—Mexic.
molle, Bert. ex DC. Prod. v. 173 = peberulum.
molle, *H. B. & K. Nov. Gen. et Sp.* iv. 109.—N. Granat.
molle, Sw. Prod. Fl. Ind. Occ. 111 = macrophyllum.
mollissimum, *Baker, in Mart. Fl. Bras.* vi. ii. 331.—Bras.
monanthum, *Sch. Bip. in Seem. Bot. Voy. Herald*, 299.—Mexic.
monardaefolium, *Walp. in Linnæa*, xiv. (1840) 506.—Bras.
montanum, *Sw. Prod. Veg. Ind. Occ.* 111.—Jamaic.
montserratense, *Spreng. Syst.* iii. 417.—Reg. Argent.
mortifolium, *Mill. Gard. Dict.* ed. VIII. n. 10.—Mexic.
Morisi, *Viz. Ort. Bot. Padov.* (1842) 80.—Hab.?
mucronatum, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 440 = conyzoides.
Mueller, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 90.—Mexic.
multicrenulatum, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 335.—Bras.
multiflorum, DC. Prod. v. 182 = anethiifolium.
multiflorum, Less. ex Baker, in Mart. Fl. Bras. vi. ii. 281 = Mikania buddleiæfolia, nodulosa.
multiflorum, Mart. ex Baker, l. c. 361 = organense.
multiloconlosum, *DC. Prod.* v. 141.—Bras.
multinerve, Benth. Pl. Hartw. 76 = pycnocephalum.
multiserratum, *Sch. Bip. in Seem. Bot. Voy. Herald*, 301.—Mexic.
myginda efolium, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 101.—Mexic.
myrsinifolium, *Spreng. Syst. Ic. Pl. Rar.* iii. 15. t. 582; *Coll.* ii. 341.—Am. austr.
myriadenium, *Schau. in Linnæa*, xix. (1847) 731.—Mexic.
myrianthum, Less. ex Baker, in Mart. Fl. Bras. vi. ii. 280 = Mikania myriocephala.
myriocephalum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 443.—Bras.
myrtilloides, *DC. Prod.* v. 165.—Bras.
myzorense, Heyne, ex DC. l. c. 451 = Pluchea tomentosa.
Neasanum, *DC. l. c.* v. 160.—Mexic.
nemorense, Sch. Bip. in Linnæa, xxxiv. (1865–66) 535, nomen; Baker, in Mart. Fl. Bras. vi. ii. 304 = Vanillerianum.
nepalense, Spreng. Syst. iii. 412 = Vernonia divergens?
nepetaefolium, *H. B. & K. Nov. Gen. et Sp.* iv. 112.—Mexic.
nepetaefolium, *Weure, in Oestr. Bot. Zeitschr.* xiii. (1863) 219.—Hab.?
nepetoides, Lindl. ex Baker, in Mart. Fl. Bras. vi. ii. 260 = bellotaefolium.
nervosum, Sieber, ex DC. Prod. v. 209 = Baccharis nervosa.
nervosum, *Sw. Prod. Veg. Ind. Occ.* 111.—Ind. occ.
neurolepis, Torr. ex Torr. & Gray, Fl. N. Am. ii. 81 = ivaefolium.
neurophyllum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 321 = dendroides.
nigrescens, Hook. & Arn. Bot. Beech. Voy. 397 = collinum.
nigrum, *Bull. Hist. Pl.* viii. 81.—Hab.?
nitidum, *Baker, in Mart. Fl. Bras.* vi. ii. 351.—Bras.
nitidum, DC. Prod. v. 150 = Kaunia nitida.
niveum, *H. B. & K. Nov. Gen. et Sp.* iv. 115. t. 342.—N. Granat.
nodiflorum, Wall. Cat. n. 3166 = connabinum.
nubigenum, *Benth. Pl. Hartw.* 55.—Guatemala.
nudiflorum, *A. Rich. Fl. Cub. Fanerog.* ii. 41.—Cuba.
nudum, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 475 = Ageratum melissæfolium.
Nummularia, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 341.—Bras.
nutans, H. B. & K. Nov. Gen. et Sp. iv. 108 = Brickellia secundiflora.
oaxacanum, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 234.—Mexic.

EUPATORIUM:—
oblongifolium, *Baker, in Mart. Fl. Bras.* vi. ii. 333.—Bras.
obovatum, Rafin. in Med. Repos. N. York, v. (1808) 359 = rotundifolium?
obovatum, Willd. ex Less. in Linnæa, iv. (1829) 271 = Vernonia scabra.
obscurum, DC. Prod. v. 142 = ivaefolium.
obtusifolium, Willd. Sp. Pl. iii. 1768 = Vernonia punctata.
obtusissimum, *DC. Prod.* v. 178.—Ins. S. Doming.
occidentale, *Hook. Fl. Bor. Am.* i. 306.—Am. bor. occ.
odoratum, *Linn. Syst.* ed. X. 1205. partim.—Am. trop.
odoratum, Linn. l. c., partim = conyzoides.
odoratum, Walt. Fl. Carol. 200 = ageratoides?
Oerstedianum, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1855) 74.—Am. centr.
officinale, Balli. Dict. ii. 748 = Mikania officinalis.
oliganthes, Less. in Linnæa, v. (1830) 137 = Brickellia oliganthes.
oligocephalum, *DC. Prod.* v. 166.—Mexic.
oligolepis, *Hemsl. Biol. Centr. Am. Bot.* ii. 98.—Mexic.
omphalaefolium, *Kunth & Bouché, Ind. Sem. Hort Berol.* (1844) *Admt.* n. 13; *ex Walp. Rep.* v. 112.—Mexic.
opiferum, Balli. Dict. ii. 748 = Mikania cordifolia.
oppositifolium, *Hoffmgg. Vers. Pfl. Nachtr.* iii. 26.—Am. bor.
orbiculatum, *DC. Prod.* v. 172.—Bras.
Orbignyanum, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 234.—Reg. Argent.
oregonum, Nutt. in Trans. Am. Phil. Soc. N. S. vii (1841) 386 = occidentale.
organense, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 117.—Bras.
orgyale, *DC. Prod.* v. 174.—Bras.
origanoides, *H. B. & K. Nov. Gen. et Sp.* iv. 114.—N. Granat.
origanoides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 257.—Peruv.
Orinabae, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 98.—Mexic.
orphalaefolium, Regel, Gartenfl. (1853) 4, sphalm. = omphalaefolium.
Ossaanum, *DC. Prod.* v. 144.—Ind. occ.
Ossaanum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 287 = laevigatum.
ovalifolium, *Hook. & Arn. Bot. Beech. Voy.* 297.—Mexic.
ovatum, Bigel. Fl. Bost. ed. II. 296 = rotundifolium.
oxychlaenum, DC. Prod. v. 169 = amygdalinum.
oxyhedrum, *Baker, in Mart. Fl. Bras.* vi. ii. 259, in obs.—Bras.
oxylepis, *DC. Prod.* v. 146.—Bras.
oxylepis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 339 = Warmingii.
Pabstii, *Regel, in Ann. Sc. Nat.* Sér. IV. xii. (1859) 376.—Hab.!
palaefolium, *Klatt, in Leopoldina*, xx. (1884) 76.—Mexic.
pallescens, *DC. Prod.* v. 154.—Bras.
pallidum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 341 = stevisefolium, pallescens.
palmare, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 294.—Bras.
palustre, *Baker, l. c.* 363.—Bras.
pandurifolium, *Baker, l. c.* 310.—Bras.
paniculatum, Lindl. ex DC. Prod. v. 23 = Vernonia Donianæ.
paniculatum, Mill. Gard. Dict. ed. VIII. n. 15 = E. Verse-crucis.
paniculatum, Schrad. Ind. Sem. Hort. Gotting. (1832) 3; ex Linnæa, viii. (1833) Litt. 36 = guadalupense.
papantlense, *Less. in Linnæa*, vi. (1831) 498.—Mexic.
paradoxum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 340 = Ophryosporus paradoxus.
paranense, *Hook. & Arn. l. c.* 341.—Bras. Argent.
parietaefolium, *Schenle, in Linnæa*, xvii. (1844) 468.—Mexic.
Parryi, *A. Gray, in Torr. Bot. Mex. Bound.* 75.—Mexic.
parviflorum, Ashl. Pl. Gal. ii. 797 = Mikania scandens.
parviflorum, Ell. Sketch. ii. 296 = micranthum.
parvilloenm, *Sw. Prod. Veg. Ind. Occ.* 111.—Jamaic.
patulifolium, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 234.—Patagon.

EUPATORIUM:—
patens, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 348.—Reg. Argent.
patens, *Phil. Sert. Mens. Alt.* 21.—Chili.
paucidentatum, *Sch. Bip. in Linnaea*, xxiv. (1865–66) 535, nomen ; *et Baker, in Mart. Fl. Bras.* vi. ii. 297.—Boliv.
pauciflorum, *H. B. & K. Nov. Gen. et Sp.* iv. 120.—N. Granat. †
paucifloroum, *Raßn. New Fl. Am.* iv. 80.—Am. bor.
paucifloum, Wall. ex DC. Prod. v. 15 = Vernonia teres.
paulinum, *DC. l. c.* 158.—Bras.
paulinum, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 282 = squalidum.
pauperstum, *H. B. & K. Nov. Gen. et Sp.* iv. 121.—N. Granat.†
pauperculum, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 205.—Arizona.
pavonianifolium, Sch. Bip. in Flora, xxxiii. (1850) 80 = Symphyopappus decussatus.
pascuarense, *H. B. & K. Nov. Gen. et Sp.* iv. 123.—Mexic.
pectinatum, *Raßn. Atl. Journ.* 17.—Am. bor.
pectinatum, Wall. Cat. n. 8172 = Vernonia pectinata.
pedale, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 295.—Bras.
peduncolosum, A. Gray, Pl. Wright. i. 86 = longipes.
pedunculosum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 240.—Reg. Argent.
pellucidum, *H. B. & K. Nov. Gen. et Sp.* iv. 108.—N. Granat.
pendulum, Schrad. Ind. Sem. Hort. Gotting. (1830) ; ex Linnæa, vi. (1831) Litt. 72 = brickellia pendula.
pentanthum, Sch. Bip. in Linnæa, xxii. (1849) 572 = dendroides.
Pentlandianum, *DC. Prod.* v. 157.—Am. austr.
perfoliatum, *Linn. Sp. Pl.* 838.—Am. bor.
perforatum, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 289.—Bras.
perornatum, Klatt, in Leopoldina, xx. (1884) 90 = Liebmanni, *Hemsl.*
Perrinianum, *Spreng. Syst.* iii. 412.—Am. austr.
persicifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 120.—Ecuador.
Petasites, *Griseb. Insul. Brit. Gotting.* (1877) 8.—Am. centr.
petiolare [*Mey. & Sesse,*] *ex DC. Prod.* v. 166.—Mexic.
phlogifolium, *DC. l. c.* 147.—Bras.
piauhyense, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 473 = dendroides.
pichinchense, *H. B. & K. Nov. Gen. et Sp.* iv. 122.—Ecuador.
pictum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 443.—Bras.
pilosum, Walt. Fl. Carol. 199 = teucrifolium †
pinnatifidum, DC. Prod. v. 149 = pinnatisectum.
pinnatifidum, *Ell. Sketch.* i. ii. 295.—Am. bor.
pinnatifidum, Buek, Ind. ad DC. Prod. ii. p. v = pinnatisectum.
pinnatipartitum, *Sch. Bip. in Linnæa*, xxii. (1849) 572, nomen ; *Baker, in Mart. Fl. Bras.* vi. ii. 338.—Bras.
pinnatisectum, *Steud. Nom.* ed. II. i. 608.—Am. austr.
pitcheiæfolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 346 = pyrifolium.
piptopappum, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 82 ; *et in Linnæa*, xxxiv. (1865–66) 535, nomen.—Bolivia.
piquerioides, DC. Prod. v. 175 = Ophryosporus piquerioides.
plantaginifolia, *Baker, in Mart. Fl. Bras.* vi. ii. 355.—Bras.
plectranthifolium, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1859) 76.—Costa Rica.
plenibracteatum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 511.—Cuba.
podachænoides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 356 = purpurascens.
Poeppigii, *Sprengel. Syst.* iii. 418.—Cuba.
polycnemiæfolium, Less. ex Baker, in Mart. Fl. Bras. vi. ii. 359 = Mikania salliæfolia.
polycnemium, Spreng. Syst. iii. 482 = Vernonia polycnemia.
polycnemium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 279 = subsquarrosum.
polycnemium, Wall. Cat. n. 3171 = Vernonia divergens.
polyhystrum, *DC. Prod.* v. 174.—Mexic.
polyrhabdium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 324 = bracteatum.

EUPATORIUM:—
polystachyum, DC. Prod. v. 149 = Symphyopappus polystachyus.
polystachyum, Willd. ex Less. in Linnæa, iv. (1829) 238 = Vernonia mollis.
pomaderrifolium, *Benth. Pl. Hartw.* 199.—N. Granat.
ponticum, Pall. ex Georgi, Beschr. Russ. Reich. viii. 306 = cannabinum.
popocatapetlense, *Schlecht. ex Hemsl. Biol. Centr. Am. Bot.* ii. 99, nomen.—Mexic.
populifolium, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 242 = hecatanthum.
populifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 111.—Mexic.
populifolium, Mart. in Flora, xx. (1837) ii. Beibl. 105 = sacrophyllum.
porphyranthum, *A. Gray, in Proc. Am. Acad.* xv. (1880) 27.—Mexic.
porphyrolepis, *Baker, in Mart. Fl. Bras.* vi. ii. 280.—Bras.
poterioides, Sch. Bip. in Linnæa, xxii. (1849) 57 = sphærocephalum.
pruuifolium, *Griseb. in Goett. Abh.* xix. (1874) 167.—Reg. Argent.
propinquum, DC. Prod. v. 146 = laevigatum.
prunellæfolium, *H. B. & K. Nov. Gen. et Sp.* iv. 123.—Mexic.
prunifolium, *H. B. & K. l. c* 132.—Peruv.
Pseudochilca, *Benth. Pl. Hartw.* 198.—N. Granat.
Pseudoperfoliatum, *Sch. Bip. in Leopoldina*, xx. (1884) 76.—Mexic.
psiadiæfolium, DC. Prod. v. 146 = laevigatum.
puberulum, *DC. l. c.* 172.—Ins. S. Doming.
pubescens, Muhl. ex Willd. Sp. Pl. iii 1755 = rotundifolium.
pubescens, Wall. ex DC. Prod. vi. 281 = Carpesium cernuum.
pulchellum, *H. B. & K. Nov. Gen. et Sp.* iv. 119. t 345.—Mexic.
pulchrum, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 444 = stachyophyllum.
pumilum, *Wender. ex Steud. Nom.* ed. II. i. 608, nomen.—Hab. ?
punctatum, *Lam. Encyc.* ii. 408.—Ind. occ.
punctatum, Mill. Gard. Dict. ed. VIII. n. 11 = Milleri.
punctatum, Mkq. in Linnæa, xvii. (1843) 66 = laevigatum.
puncticulosum, Willd. Enum. Hort. Berol. 853 = purpureum.
punctulatum, *DC. Prod.* v. 147.—Bras.
punduanum, Wall. Cat. n. 3170 = cannabinum.
pungens, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 474 = oxylepis.
purpurascens, *Sch. Bip. in Linnæa*, xxx. (1859–60) 183, nomen ; *Baker, in Mart. Fl. Bras.* vi. ii. 356.—Bras.
purpureum, *Linn. Sp. Pl.* 838.—Am. bor.
purpureum, Lour. Fl. Cochinch. 457 = quaternum.
pyanocephalum, *Less. in Linnæa*, vi. (1831) 404.—Am. bor. ; Mexic.
pyramidale, D. Don, Prod. Fl. Nep. 170 = Vernonia Roxburghii.
pyramidale, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1885) 234.—Bolivia.
pyrifolium, *DC. Prod.* v. 158.—Bras.
quadrangulare, *DC. l. c.* 150.—Mexic.
quaternum, *DC. l. c.* 158.—Cochinch.
quinquenervioides, Hemsl. Biol. Centr. Am. Bot. ii. 90, sphalm. = Fleischmannia rhodostylis.
quinquenectum, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1859) 79.— Nicaragua.
quinquepilosum, Less. ex Baker, in Mart. Fl. Bras. vi. ii. 240 = Mikania laevis.
racemosum, Bertol. Misc. Bot. vi. 26. t. 1 = coronopifolium.
racemosum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 339 = bracteatum.
ramosissimum, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 444 = squalidum.
ramosum, *Mill. Gard. Dict.* ed. VIII. n. 13.—Am. bor.
Rauwenti, *Wall. Cat.* n. 3166.—China.
reflexum, Moench, Meth. Suppl. 234 = Blumea chinensis.
Regnellii, Sch. Bip. in Linnæa, xxx. (1859–60) 182 = Ophryosporus Regnellii.
remotifolium, DC. Prod. v. 165 = E. Vitalbæ.
Remyanum, *Phil. Fl. Atac.* 29.—Chili.

EUPATORIUM:—
repandum, *Willd. Sp. Pl.* iii. 1767.—Ind. occ.
reticulatum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 54.—Peruv.
reticulatum, *Torr. ex DC. Prod.* v. 176.—Am. bor.
reticulatum, *Dew. in Ham. Prod. Pl. Ind. Occ.* 51.—Jamaic.
reticulatum, Hook. & Arn. Bot. Beech. Voy. 29 = E. Salvia.
reticulatum, *A. Rich. Fl. Cub. Fanerog.* ii. 89.—Cuba
retifolium, Sch. Bip. in Flora, xxxiii. (1850) 80, nomen = Mikania sessilifolia.
retrofractum, Thunb. Prod. Pl. Cap. 149 = Iphiona polygalifolia.
revolutum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 444.—Bras.
revolutum, Pohl, ex Baker, in Mart. Fl. Bras. vi. ii. 314 = amygdalinum.
rhinanthaceum, *DC. Prod.* v. 146.—Bras.
rhodanthum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 314 = amygdalinum.
rhodochlamydeum, *A. Gray, in Proc. Am. Acad.* xv. (1880) 26.—Mexic.
rhomboideum, *H. B. & K. Nov. Gen. et Sp.* iv. 127.—Mexic.
Riedelianum, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 478.—Bras.
Riedelii, *Baker, in Mart. Fl. Bras.* vi. ii. 355.—Bras.
rigidulum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 167.—Japon.
rigidum, Benth. Pl. Hartw. 88 = Brickellia Hartwegi.
rigidum, *Sw. Prod. Veg. Ind. Occ.* 111.—Jamaic.
rigidum, Wall. Cat. n. 3925 = Vernonia teres.
riparium, *Regel, Gartenfl.* xv. (1866) 324. t. 525.—Mexic.
Rosalesia, DC. Prod. v. 163 = Brickellia Rosalesia.
roseum, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 106.—Bras.
rosmarinifolium, Labill. Nov. Holl. Pl. ii. 38 = Helichrysum rosmarinifolium.
rosmarinifolium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 319 = stevisefolium.
Rothrockii, *A. Gray, Syn. Fl. N. Am.* i. ii. 102.—Arizona.
rotundifolium, *Linn. Sp. Pl.* 837.—Am. bor.
rubricaule, H. B. & K. Nov. Gen. et Sp. iv. 124 = guadalupense.
rufescens, *Lund, ex DC Prod.* v. 168.—Bras.
rufidulum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 243.—Bras.
rufum, Spreng. ex DC. Prod. v. 83 = Vernonia extensa.
rugosum, *H. B. & K. Nov. Gen. et Sp.* iv. 114.—Ecuador.
rugosum, Schrad. ex DC. Prod. v. 174 = Schraderi.
rugosum, Spreng. Neue Entdeck. iii. 39 = Sprengelianum.
rupestre, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 474.—Bras.
rupestre, Kafin. Atl. Journ. 17 = altissimum.
Sabeanum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 456 = conyzoides.
sagittatum, *A. Gray, Pl. Wright.* i. 88.—Am. bor. occ.
salicinum, *Lam. Encyc.* ii. 409.—Ecuador.
Salvia, *Colla, in Mem. Acc. Tor.* xxxviii. (1835) 8.—Chili.
salviæfolium, *H. B. & K. Nov. Gen. et Sp.* iv. 131.—Ecuador.
salviæfolium, Sims, Bot. Mag. t. 2019 = perfoliatum.
salvifolium, *Balb. ex Spreng. Neue Entdeck.* iii. 40.—Hab.?
salvifolium, Bert. ex Spreng. Syst. iii. 412 = pubescens.
Salsmennianum, DC. Prod. v. 159 = amygdalinum.
saturneiæfolium, Lam. Encyc. ii. 411 = Stevia saturneiæfolia.
scaberrimum, *Walp. in Linnæa*, xiv. (1840) 505.—Bras.
scabridum, Ell. Sketch, ii. 299 = rotundifolium.
scabrum, *Linn. f. Suppl.* 354.—N. Granat.
scandens, Hook. & Arn. Comp. Bot. Mag. i. (1835) 243 = Mikania lagoënsis.
scandens, Link, Enum. Hort. Berol. ii. 307 = Senecio diclitoides.
scandens, Linn. Sp. Pl. 836 = Mikania scandens.
scandens, Thunb. Prod. Pl. Cap. 149 = Mikania capensis.
scariosum, Wallr. ex DC. Prod. v. 138 = Brickellia pendula.

EUPATORIUM:—
Schaffneri, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 100, nomen.—Mexic.
schiedeanoides, Sch. Bip. ex A. Gray, in Torr. Bot. Mex. Bound. 76 = seq.
Schiedeanum, Schrad. Ind. Sem. Hort. Gotting. (1832) 3 = pycnocephalum.
Schimperianum, *Schenk, in Linnæa*, xviii. (1844) 458. Arabia.
schizanthum, *Griseb. Fl. Brit. W. Ind.* 361.—Jamaic.
Schlechtendalii, DC. Prod. vii. 269 = sordescens.
Schomburgkii, *Benth. in Hook. Journ. Bot.* ii. (1840) 41.—Guiana.
Schraderi, *DC. Prod.* v. 174.—Am. austr.
Schultzii, *Sch. Bip. in Zeitschr. Gartenb. Darmst.* (1857) 6.—Mexic.
scopuliorum, *Wedd. Chlor. And.* i. 216.—Bolivia.
scoroloniolides, *A. Gray, in Proc. Am. Acad.* xv. (1880) 97.—Mexic.
secundiflorum, Berl. ex DC. Prod. v. 48 = Vernonia arborescens.
secundiflorum, Lag. Gen. et Sp. Nov. 25 = Brickellia secundiflora.
Selloi, *Baker, in Mart. Fl. Bras.* vi. 11. 361.—Bras.
semialatum, *Benth. Pl. Hartw.* 76.—Mexic.
semiserratum, *DC. Prod.* v. 177.—Am. bor.
semistriatum, *Baker, in Mart. Fl. Bras.* vi. 11. 318.—Bras.
serocinoideum, *Baker, l. c.* 335.—Bras.
septuplinervium, *Klatt, in Leopoldina*, xx. (1884) 90.—Mexic.
sertorum, *H. B. & K. Nov. Gen. et Sp.* iv. 110.—N. Granat.
serotinum, *Micks. Fl. Bor. Am.* ii. 100.—Am. bor.; Mexic.
serratifolium, *DC. Prod.* v. 181.—N. Granat.
serratuloides, *H. B. & K. Nov. Gen. et Sp.* iv. 117.—Ecuador.
serratum, *Spreng. Syst.* iii. 415.—Bras.
serrulatum, *DC. Prod.* v. 147.—Bras.
sessilifolium, *Linn. Sp. Pl.* 837.—Am. bor.
setiferum, Pav. ex A. Gray, in Proc. Am. Acad. xvii. (1881–82) 206 = Barrocetra Pavonii.
Sideritidis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1859) 77.—Costa Rica.
Siebertianum, DC. Prod. v. 181 = lresinoides.
silenifolium, Mart. ex Baker, in Mart. Fl. Bras. vi. 11. 215 = Trichogonia campestris.
silphiifolium, Mart. in Flora, xx. (1837) 11. Beibl. 105 = pallescens.
Simonsii, C. B. Clarke, Comp. Ind. 39 = cannabinum.
Sinclairii, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1859) 79 = guadalupense.
sinense, J. F. Gmel. Syst. 1198 = chinense.
sinuatum, *Lam. Encyc.* ii. 407.—Ind. occ.
sinuatum, Lour. in Cochinch. 487 = hispidum.
smilacinum, *H. B. & K. Nov. Gen. et Sp.* iv. 111.—N. Granat.
Solierii, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 498 = cannabinum.
solidaginifolium, *A. Gray, Pl. Wright.* i. 87.—Am. bor. occ.
solidaginoides, *H. B. & K. Nov. Gen. et Sp.* iv. 126.—Ecuador.
Sondrei, Sch. Bip. in Linnæa, xxii. (1849) 571 = hepleriifolium.
Sonoræ, A. Gray, Pl. Wright. ii. 74 = pycnocephalum.
sophiæfolium, *Linn. Sp. Pl.* 1175.—Ind. occ.
sophioides, *DC. Prod.* v. 170.—Ins. S. Doming.
Sorstae, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 585, nomen.—Bolivia.
sordescens, *DC. Prod.* v. 167.—Bras.
sordidum, *Less. in Linnæa*, vi. (1831) 408.—Mexic.
spathulatum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 242 = halimifolium.
speciosum, Vent. Jard. Cels, 79 = Liatris elegans.
spectabile, Spreng. Syst. iii. 410 = Leucomeris spectabilis.
sphaerocephalum, *Sch. Bip. in Linnæa*, xxx. (1859–60) 182, nomen; *Baker, in Mart. Fl. Bras.* vi. 11. 317.—Bras.
spicatum, *Lam. Encyc.* ii. 409.—Reg. Argent.
spinosifolium, *A. Gray, in Torr. Bot. Mex. Bound.* 78, note.—Mexic.
spissumfolium, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 101, nomen.—Mexic.
spinulosum, *Klatzsch, ex Polak. in Linnæa*, xli. (1877) 678.—Costa Rica.

EUPATORIUM:—
Sprengelianum, *DC. Prod.* v. 184.—Ind. occ.
squalidum, *DC. l. c.* 142.—Bras.
squamosum, D. Don, Prod. Fl. Nep. 170 = Roxburgii.
squarrosum, Cav. Ic. i. 68 = Brickellia Cavanillesii.
squarrulosum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 239 = latridiflorum.
stachyophyllum, *Spreng. Syst.* iii. 490.—Bras.
stenolepis, *Steetz, in Seem. Bot. Voy. Herald*, 148.—N. Granat.
Sternbergianum, *DC. Prod.* v. 167.—Peruv.
sterlaefolium, *DC. l. c.* 158.—Bras.
stigmatosum, Bertol. Misc. Bot. v. 15. t. 5 = album.
stigmatosum, Meyen & Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 357 = corymbosum.
stillingiaefolium, *DC. Prod.* v. 160.—Mexic.
stipulaceum, Vahl, Symb. Bot. iii. 94 = Mikania stipulacea.
stoechadifolium, *Linn. f. Suppl.* 355.— N. Granat.
stoechadosmum, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 222.—China.
stramineum, *DC. Prod.* v. 150.—Peruv.
strobilanthum, Mart. ex Baker, in Mart. Fl. Bras. vi. 11. 300 = horminoides.
suaveolens, Chapm. in Coult. Bot. Gaz. iii. (1878) 5 = stramoniifolium.
suaveolens, *H. B. & K. Nov. Gen. et Sp.* iv. 109.—Am. austr.
suaveolens, Wall. Cat. n. 3290 = cannabinum.
subalternifolium, *DC. Prod.* v. 152.—Bras.
subcordatum, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1859) 77.—Am. centr.
subdentatum, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 478 = adamantium.
subhastatum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 239 = hartiaefolium.
subinclusum, *Klatt, in Leopoldina*, xx. (1884) 75.—Mexic.
subobtusum, DC. Prod. v. 161 = amygdalinum.
subpannivervium, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 89.—Mexic.
subpinnatum, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 242 = erodiifolium.
subserratum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 440.—Bras.
subsessilifolium, Miq. in Journ. Bot. Néerl. i. (1861) 99 = Lindleyanum.
subtriplinerve, Sch. Bip. in Linnæa, xxx. (1859–60) 189 = rufescens.
subtruncatum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 448.—Bras.
subuliginum, A. Gray, Pl. Wright. i. 86 = Barrocetra subuliginea.
subvelutinum, DC. Prod. vii. 268 = squalidum.
subverticillatum, Less. ex Baker, in Mart. Fl. Bras. vi. 11. 295 = Mikania subverticillata.
sulcatum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 242 = Mikania pentstemonoides.
surinamense, Sch. Bip. ex Miq. Stirp. Surin. Sel. 185 = sessilifolium.
syriacum, Jacq. Ic. Pl. Rar. i. 17. t. 170; Misc. ii. 349 = cannabinum.
syriacum, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) ii. 278.—Syria.
syringaefolium, *Tuscz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 169.—Peruv.
tanacetifolium, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1836) 242 = erodiifolium.
tectum, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 117.—Bras.
tenuifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 107.—Venezuel.
tepicanum, *Hemsl. Biol. Centr. Am. Bot.* ii. 101.—Mexic.
ternifolium, Ell. Sketch, ii. 306 = purpureum.
tetragonum, *Schrad. Ind. Sem. Hort. Gotting.* (1830); *ex Linnæa*, vi. (1831) *Litt.* 78.—Mexic.
tetranthum, *Griseb. Fl. Brit. W. Ind.* 360.—Jamaic.
tetranthum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 11. 337 = dentatum.
texcatlicum, *Willd. Sp. Pl.* iii. 1758.—Am. bor.
texcatlifolium, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 239 = hartiaefolium.
theaefolium, *Benth. Pl. Hartw.* 199.—N. Granat.
theopentaefolium, *DC. Prod.* v. 164.—Mexic.
thyrsoideum, *Mey. ex DC. l. c.* 150.—Mexic.
tinifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 122.—N. Granat.

EUPATORIUM:—
tomentellum, *Schrad. Ind. Sem. Hort. Gotting.* (1838) 8. t. 3.—Mexic.
tomentosum, Lam. Encyc. ii. 410 = Senecio paniculatus.
tomentosum, Steud. Nom. ed. II. i. 606, sphalm. = tomentellum.
Torreyanum, Short, ex Torr. & Gray, Fl. N. Am. ii. 84 = hyssopifolium.
tortifolium, Chapm. in Coult. Bot. Gaz. iii. (1878) 5 = hyssopifolium.
toxolepis, *DC. Prod.* v. 146.—Bras.
trapezoideum, *Kunth, Ind. Sem. Hort. Berol.* (1847) 13.—Mexic.
tremulum, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 241 = dendroides.
triangulare, Moç. ex DC. Prod. v. 166 = deltoideum.
triangulare, Poir. Encyc. Suppl. ii. 604 = atriplicifolium.
triangulatum, *Alam. ex DC. Prod.* v. 172.—Mexic.
tricephalotes, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. 11. 329.—Bras.
trichobasis, *Baker, l. c.* 364.—Bras.
trichophorum, DC. Prod. v. 147 = hartiaefolium.
trichosanthum, *A. Rich. Fl. Cub. Fanerog.* ii. 41.—Cuba.
trichotomum, Sch. Bip. in Linnæa, xxxiv. (1865–66) 585, nomen; Baker, in Mart. Fl. Bras. vi. 11. 306 = Vauthierianum.
trifidum, *Vahl, Symb. Bot.* iii. 94.—Ind. occ.
triflorum, Aubl. Pl. Gui. ii. 795. t. 314 = Piptocarpa triflora.
trifoliatum, Hort. Dorp. ex Step. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 271 = cannabinum.
trifoliatum, Linn. Sp. Pl. 837 = purpureum.
trifoliatum, Lucé, Topog. Nachr. Cesel, 270 = cannabinum.
trigonocarpum, *Griseb. Fl. Brit. W. Ind.* 359.—Ins. Dominic.
trigonum, *Garan. in Hook. Lond. Journ. Bot.* vi. (1847) 446.—Bras.
trinervium, *Sch. Bip. in Seem. Bot. Voy. Herald*, 300.—Mexic.
triplinerve, *Vahl, Symb. Bot.* iii. 97.—Am. trop.
triasctum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 242.—Chili.
triste, *DC. Prod.* v. 166.—Jamaic.
trizoides, *Mart. ex Baker, in Mart. Fl. Bras.* vi. 11. 311.—Bras.
truncatum, Ell. Sketch. ii. 298 = sessilifolium.
truncatum, Muhl. ex Willd. Sp. Pl. iii. 1751 = perfoliatum.
tubiflorum, *Benth. Pl. Hartw.* 76.—Mexic.
Tuerckheimii, *Klatt, in Leopoldina*, xx. (1884) 96.—Mexic.
tulanum, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 324.—Mexic.
turbinatum, *A. Gray, in Proc. Am. Acad.* xv. (1880) 96.—Mexic.
Tweedianum, Hook. & Arn. in Hook. Comp. Bot Mag. i. (1835) 242 = serrisefolium.
umbellulatum, Spreng. Syst. iv. App. 409 = Piptocarpha umbellulata.
umbrosum, *Benth. Pl. Hartw.* 198.—Peruv.
urticæfolium, *Linn. f. Suppl.* 354.—N. Granat.
urticæfolium, Banks, ex Griseb. Fl. Brit. W. Ind. 362 = guadalupense.
vaccinioides, *Benth. Pl. Hartw.* 200.—N. Granat.
vagans, Wall. Cat. sub n. 3040 = Vernonia scandens.
vagans, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 477 = Gaudichaudianum.
vallincola, *DC. Prod.* v. 168.—Peruv.
vanilloides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 11. 346 = pyrifolium.
Vargasianum, *DC. Prod.* v. 158.—Venezuel.
vatifolium, *Benth. Ind. Sem. Hort. Gotting.* (1840); *ex Linnæa*, xv. (1841) *Litt.* 93.—Am. bor.
Vauthierianum, *DC. Prod.* v. 159.—Bras.
velutinum, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 475.—Bras.
venosum, Mart. ex Baker, in Mart. Fl. Bras. vi. 11. 281 = squalidum.
Verae-crucis, *Steud. Nom.* ed. II. i. 606.—Mexic.
verbenaceum, *DC. Prod.* v. 146.—Bras.
verbenæfolium, Micks. Fl. Bor. Am. ii. 99 = cuneifolium.
vernonia, *Vahle & Kurtz. Ind. Sem. Hort. Havn.* (1871) *App.* 7; *ex Cavanill.* (1872) 38.—Mexic.

EUPATORIUM:—
vernoniopsis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 284.—Bras.
vernoniæfolium, H. B. & K. Nov. Gen. et Sp. iv. 112. t. 341 = Brickellia veroniæfolia.
verticilare, Wall. Cat. n. 3167 = Vernonia divergea.
verticillatum, Lam. Encyc. ii. 405 = purpureum.
vestitum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 55.—Peruv.
vestitum, Spreng. Syst. iii. 416 = Salmea hirsuta.
viburnoides, *DC. Prod.* v. 171.—Mexic.
villosum, Bert. ex DC. Prod. v. 173 = obtusissimum.
villosum, *Sw. Prod. Veg. Ind. Occ.* 111.—Am. bor.; Ind. occ.
vincæfolium, Lam. Encyc. ii. 410 = Mikania amara.
vindex, *DC. Prod.* v. 160.—Bras.
violaceum, Bert. ex Spreng. Syst. iii. 414 = guadalupense.
violaceum, Raffin. Fl. Ludov. 69 (sp. falsa).
virgatum, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 241 = pinnatifidum.
virgatum, Schrad. Ind. Sem. Hort. Gotting. (1829); ex DC. Prod. v. 159 = pycnocephalum.
viridiflorum, *Baker, in Mart. Fl. Bras.* vi. ii. 309.—Bras.
viscidum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 241.—Reg. Argent.
viscosum, *H. B. & K. Nov. Gen. et Sp.* iv. 129.—Ecuador.
viscosum, Wall. Cat. n. 3527 = cannabinum.
Vitalbae, *DC. Prod.* v. 168.—Bras.
vitifolium, *Klatt, in Leopoldina*, xx. (1884) 90.—Mexic.
Volckmanni, *Phil. in Anal. Univ. Chil.* (1861) 51.—Chil.
volkameriæfolium, Wall. Cat. sub n. 3001 = Vernonia volkameriæfolia.
volubile, Vahl, Symb. Bot. iii. 93 = Mikania scandens.
vulcanicum, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 78.—Am. centr.
Wallichii, DC. Prod. v. 179 = japonicum.
Warmingii, *Baker, in Mart. Fl. Bras.* vi. ii. 289.—Bras.
Warmingii, Sch. Bip. ex Baker, l. c. 340 = dictyophyllum.
Wawanenianum, Regel & Koern. Ind. Sem. Hort. Petrop. (1857) 41 = tricranthum.
Wrightii, *A. Gray, Pl. Wright.* i. 87.—N. Mexic.
xalapense, *H. B. & K. Nov. Gen. et Sp.* iv. 128.—Mexic.
xerolepis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 322 = patens, *Hook. & Arn.*
xylophyloides, DC. Prod. v. 149 = dendroides.
xylorhizum, *Sch. Bip. in Linnæa*, xxx. (1859–60) 189, nomen; *Baker, in Mart. Fl. Bras.* vi. ii. 292.—Bras.
zeylanicum, Linn. Sp. Pl. 837 = Vernonia zeylanica.

EUPETALUM, Lindl. Introd. Nat. Syst. ed. II. 440 (1836) = Begonia, Linn.
Candichaudii, Klotzsch, in Monatsb. Berl. Acad. (1854) 191 = B. Candichaudii.
geranifolium, Klotzsch, l. c. = B. geranifolia.
Kunthianum, Klotzsch, l. c. = B. Candichaudii.
Lindleyanum, Gaudich. Bot. Voy. Bonite, t. 50 = B. Candichaudii.
petaloides, Lindl. Introd. Nat. Syst. ed. II. 440 = B. petaloides.
tuberosum, Klotzsch, in Monatsb. Berl. Acad. (1854) 191 = B. geranifolia.

EUPHOCARPUS, Andens. ex Steud. Nom. ed. II. i. 622, 600 (1840) = Correa, Sm. (Rutac.).

EUPHORBE, Blume, ex Meisn. in DC. Prod. xv. i. 58, 59, in syn. (1864) = Alsodaphne, Nees (Lauric.).
cauwensis, Blume, ex Meisn. l. c. 59 (= *Nothaphoebe cauwensis*).—Sumatra.
chartacea, Blume, ex Meisn. l. c. 58 (= *Nothaphoebe chartacea*).—Java.
cuneata, Blume, ex Meisn. l. c. (= *Nothaphoebe cuneata*).—Malaya.
falcata, Blume, ex Meisn. l. c. 59 (= *Nothaphoebe falcata*).—Sumatra.
macrophylla, Blume, ex Meisn. l. c. (= *Nothaphoebe macrophylla*).—Malaya.
rufescens, Blume, ex Meisn. l. c. (= *Nothaphoebe rufescens*).—Malaya.
umbellifloræ, Blume, ex Meisn. l. c. 58 (= *Nothaphoebe umbellifloræ*).—Malaya.

EUPHORA, Griff. Notul. iv. 547 (1854) = Aglaia, Lour. (Meliac.).
exstipulata, C. DC. in DC. Monog. Phan. i. 616 = A. minutiflora.
exstipulata, Griff. Notul. iv. 547 = A. minutiflora.

EUPHORBION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = Euphorbia, Linn.
anacampserum, St. Lag. l. c. 126 = Euphorbia Anacamperos.
apium, St. Lag. l. c. = Euphorbia Apios.
dendroideum, St. Lag. l. c. = Euphorbia dendroides.
exulum, St. Lag. l. c. = Euphorbia Esula.
heliocopium, St. Lag. l. c. = Euphorbia Heliosocpia.
lathyrum, St. Lag. l. c. = Euphorbia Lathyris.
myrsinitum, St. Lag. l. c. = Euphorbia Myrsinites.
peraltum, St. Lag. l. c. = Euphorbia Peralias.
peplum, St. Lag. l. c. 125 = Euphorbia Peplus.

EUPHORBIA, Linn. Gen. ed. I. 162 (1737). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 258.
ADENOPETALUM, ARTHROTHAMNUS, DICHROPHYLLUM, EUMECANTHUS, EUPHORBIASTRUM, LEPTOPUS, STERIGMANTHE, TITHYMALOPSIS, TRICHEROSTIGMA, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 248–252.
ADENORIMA, AGALOMA, AKLEMA, ALLOBIA, CTATHUPHORA, DESMOREMA, KANOPICON, KESASELMA, LEPADENA, LOPHOBIS, MURTEKIAS, NIHOMENES, PYTHIUS, TIRUCALLA, TORFOSIDIS, TUMALIS, VALLARIA, XAMESIKE, Rafin. Fl. Tellur. iv. 119–117 (1836).
ALECTOROCTONUM, Schlecht. in Linnaea, xix. (1847) 259.
AMBOPHILLUM, Haw. Syn. Pl. Succ. 159 (1812).
ANTHACANTHA, Lem. Illustr. Hortic. v. (1858) Misc. 64.
APLARIMA, Rafin. New Fl. Am. iv. 99 (1836).
ATHYMALUS, Neck. Elem. ii. 353 (1790).
CHAMAESYCE, S. F. Gray, Nat. Arr. Brit. Pl. ii. 260 (1821).
CHAMAESYCE, S. F. Gray, l. c. 259 (1821).
CHYLOCALA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150.
DACTYLANTHUS, Haw. Syn. Pl. Succ. 182 (1812).
DITRITRA, ENDOISILA, PECCANA, Rafin. Sylva Tellur. 114–115 (1838).
EPURGA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150.
ESULA, Rupp. Fl. Jen ed. Hall. 271 (1745); Haw. Syn. Pl. Succ. 158 (1812).
GELARGEROIS, Haw. l. c. 148 (1812).
KERASELMA, Neck. Elem. ii. 358 (1790).
LACANTHIS, Rafin. Fl. Tellur. ii. 94 (1836).
LYCIOPSIS, Schweinf. Beitr. Fl. Aethiop. 37 (1867).
MEDUSEA, Haw. Syn. Pl. Succ. 153 (1812).
PETALANDRA, F. Muell. ex Boiss. in DC. Prod. xv. ii. 27 in syn. (1862).
PETALOMA, Rafin. ex Boiss. l. c. 63 (1862).
PLEURADENIA, Rafin. Atl. Journ. 182 (1838).
POINSETTIA, R. Grah. in Edinb. N. Phil. Journ. xx. (1836) 419.
TITHYMALUS, Tourn. ex Hall. Enum. Stirp. Helv. i. 189 (1742).
TREISIA, Haw. Syn. Pl. Succ. 131 (1812).
TRIADENUM, Rafin. New Fl. Am. iv. 99 (1836).
abortiva, *Forsk. Fl. Aegypt. Arab.* p. cxii, nomen.—Arabia.
abyssinica, *J. F. Gmel. Syst.* 759; *A. Rich. Tent. Fl. Abyss.* ii. 329.—Abyss.
acalyphoides, *Hochst. ex Boiss. in DC. Prod.* xv. ii. 98.—Nubia.
acanthothamnos, *Heldr. ex Sart. in Boiss. Diagn.* Ser. II. iv. 86.—Graecia.
acmitis, Roxb. Hort. Beng. [86]; Fl. Ind. ii. 472 = fusiformis.
acocensis, *Boiss. in DC. Prod.* xv. ii. 55.—Bolliv.
achanocarpa, Spreng. Syst. iii. 797, sphalm. = akenocarpa.
aculeata, *Forsk. Fl. Aegypt. Arab.* 94.—Arab.
aculeata, E. Mey. in Drège, Zwei Pfl. Docum. 184 = sclerophylla.
acuminatoria, Lam. Encyc. ii. 496 = falcata.
acuta, *Engelm. in Terr. Bot. Mex. Bound.* 139.—Texas.
admochlora, *E. Mey. & Decne. in Bull. Acad. Brux.* iii. (1836) 174.—Japon.!
adnophylla, *Hort. ex Steud. Nom.* ed. II. i. 609, nomen.—Hab.!

EUPHORBIA:—
adenopoda, *Baill. Adansonia*, i. (1860–61) 141.—Madag.
adenoptera, *Bertol. Misc. Bot.* iii. 20. t. 23.—Am. trop.
adiantholdes, *Lam. Encyc.* ii. 496.—Peruv.
Adriana, *St. Hil. ex Klotzsch & Garcke, in Abh. Akad. Berl.* (1860) 78.—Bras.
aegyptiaca, *Boiss. Cent. Euph.* 13.—Aegypt.
affinis, Boiss. l. c. 19 = Balbisii.
affinis, DC. Fl. Fr. vi. 363 = E. Terracina.
agowensis, *Hochst. ex Boiss. in DC. Prod.* xv. ii. 70.—Abyss.
agraria, *Bieb. Fl. Taur. Casuc.* i. 375.—Europ. austr.; Reg. Caucas.
agrorum, Willd. ex Ledeb. Fl. Ross. iii. 579 = agraria.
akenocarpa, *Guss. Adnot. Cat. Pl. Boccad.* (1821) 7.—Reg. Mediterr. occ.
akenocarpa, Roux & Blaise, ex Boiss. in DC. Prod. xv. ii. 118 = cyhlrensis.
alata, *Hook. Ic. Pl.* t. 700.—Jamaic.
alatavica, *Boiss. Cent. Euph.* 33.—Soongaria.
albomarginata, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 174.—N. Mexic.
aleppica, *Linn. Sp. Pl.* 458.—Europ. austr.; Orient.
alexandrina, Delile, Fl. Egypte, 284. t. 30 = E. Terracina.
algeriensis, *Boiss. Diagn.* Ser. II. iv. 85.—Alger.
alpigena, Kern. in Oestr. Bot. Zeitschr. xvi. (1866) 357 = dulcis.
alpina, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 156.—Sibir.
alsinaefolia, *Baill. Adansonia*, vi. (1866) 288.—Austral.
alsinaefolia, *Boiss. Cent. Euph.* 16.—Bras.
alsinoides, *Miq. in Linnaea*, xxvi. (1853) 229.—Ind. or.
altaica, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 190.—Sibir.
altissima, *Boiss. Diagn.* Ser. I. v. 59.—As. Min.; Syria.
amblyrus, Waldst. & Kit. Ic. Pl. Hung. ii. 144. t. 135 = carniolica.
ammanioides, *H. B. & K. Nov. Gen. et Sp.* ii. 85.—Panama; Mexic.
ammaniotrichia, *Boiss. Cent. Euph.* 10.—Mexic.
amoena, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1164, nomen.—Guian.
ampla, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 20.—Afr. trop.
amplexicaulis, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 158.—Ins. Galapag.
amplexicaulis, Ledeb. Fl. Ross. iii. 567 = condylocarpa.
amygdaloides, Lam. Encyc. ii. 439 = nicaeensis.
amygdaloides, *Linn. Sp. Pl.* 463.—Eu.-op.; Orient.
anacampseroides, Lam. Encyc. ii. 430 = Pedilanthus padifolius.
Anacampseros, Bal. ex Boiss. in DC. Prod. xv. ii. 174 = denticulata.
Anacampseros, *Boiss. Diagn.* Ser. I. v. 55.—As. Min.
anacantha, *Ait. Hort. Kew.* ed. I. ii. 136.—Afr. austr.
ancepa, Benth. Bot. Voy. Sulph. 169 = adenoptera.
andina, *Phil. in Linnaea*, xxiv. (1857) 41.—Chili.
andrachnoides, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 197 = subcordata.
andromonoica, Pruel, Delic. Prag. 67 = Preslii.
androsaemifolia, [*Schousb.*] *Willd. Enum. Hort. Berol. Suppl.* 27.—Lusitan.
androsaemifolia, Stend. Nom. ed. II. i. 610 = lucida.
androsaemoides, Dennst. Schlüss. Hort. Malab. 36 = hypericifolia.
angularis, Klotzsch, in Monatsb. Akad. Berl. (1859) 250, nomen; et in Peters, Reise Mossamb. Bot. 92 = abyssinica.
anguiata, *Yacq. Coll.* ii. 309; *Ic. Pl. Rar.* iii. t. 481. —Europ. med.
angusta, *Engelm. in Torr. Bot. Mex. Bound.* 139.—Texas.
angustifolia, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 63.—Nepal.
angustifolia, Lockh. ex Boiss. in DC. Prod. xv. ii. 73 = heterophylla.
angustifolia, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 49.—Reg. Argent.
angustifolia, Sweet, Hort. Brit. ed. I. 357 = E. Esula.
anomala, Salzm. ex Boiss. in DC. Prod. xv. ii. 59 = lanulana.
antiquorum, Forsk. Fl. Aegypt. Arab. 93 = E. Cactus.

EUPHORBIA :—
antiquorum, *Linn. Sp. Pl.* 450.—Ind. or.
antiquorum, E. Mey. in Drège, Zwei Pfl. Docum. 184 = cerviornis.
antiquorum, Wall. Cat. n. 7688 = E. Nivalia.
antisiphylitica, *Zucc. in Abh. Akad. Muench.* i. (1829–30) 209.—Mexic.
anychioides, *Boiss. Cent. Euph.* 19.—Mexic.
aphylla, *Brouss. in Willd. Enum. Hort. Berol.* 501.—Teneriffe.
apiculata, *Anderss. in Vet. Akad. Handl. Stockh.* 1858 (1855) 234.—Ins. Galapag.
Apios, *Linn. Sp. Pl.* 457.—Graecia; As. Min.
apocynoides, *Klotzsch, in Seem. Bot. Voy. Herald.* 99.—Panama.
arabica, *Hochst. & Steud. ex Boiss. in DC. Prod.* xv. ii. 38.—Arabia; Abyss.
aragonensis, Losc. & Pard. Ser. Inconf. Arag. 96 = nevadensis.
arariea, Jord. in Billot, Annot. 27 = E. Esula.
arborea, Boiss. in DC. Prod. xv. ii. 108, sphalm. = Tuckeyana.
arborescens, Hort. Angl. ex Salm-Dyck, Hort. Dyck. 104 = grandidens.
arborescens, Hort. Par. ex Boiss. in DC. Prod. xv. ii. 82 = macroglypha.
arborescens, E. Mey. in Drège, Zwei Pfl. Docum. 184 = Synadenium arborescens.
arborescens, Roxb. Hort. Beng. [36]; Fl. Ind. ii. 468 = antiquorum ?
arborescens, C. Sm. in Tuckey, Congo, 351 = Tuckeyana.
Arbuscula, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 98.—Ins. Socotra.
arceuthoides, *Boiss. Cent. Euph.* 20.—Afr. austr.
arenaria, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 260 = zygophylloides.
armata, *H. B. & K. Nov. Gen. et Sp.* ii. 57.—N. Granat.
armata, Nutt. in Trans. Am. Phil. Soc. v. (1837) 171 = petaloides.
armata, Willd. ex Ledeb. Fl. Ross. iii. 570 = falcata.
argusa, *[Soland. in] Russ. Aleppo, ed.* II. ii. 253.—Graecia; Syria.
arienais, *H. B. & K. Nov. Gen. et Sp.* ii. 57.—Mexic.
arillata, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1018 = granulata.
arizonica, *Engelm. in Torr. Bot. Mex. Bound.* 186.—N. Mexic.
arbanensa, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 261 = dictyosperma.
armata, Thunb. Prod. Pl. Cap. 86 = E. Hystrix.
armena, Boiss. in DC. Prod. xv. ii. 1268 = petrophila.
armeniaca, Boiss. Fl. Orient. iv. 1118 = petrophila.
Armstrongiana, *Boiss. in DC. Prod.* xv. ii. 47.—Austral.
Artaudiana, DC. Fl. Fr. vi. 360 = pinea.
artiensis, *Anderss. in Vet. Akad. Handl. Stockh.* 1858 (1855) 236.—Ins. Galapag.
articulata, Aubl. Pl. Gui. i. 480 = linearis.
articulata, Desnst. Schlecht. Hort. Malab. 20 = E. Atoto.
arvalis, *Boiss. & Heldr. Diagn.* Ser. I. xii. 116.—As. Min.; Persia.
arvensis, Schleich. ex Steud. Nom. ed. I. 328 = falcata.
ascendens, Willd. ex Boiss. in DC. Prod. xv. ii. 81 = orbiculata.
aspera, *Bieb. Fl. Taur. Cauc.* i. 377.—Reg. Caucas.
asperifolia, Engelm. ex Boiss. in DC. Prod. xv. ii. 74 = strigosa.
astrachanica, C. A. Mey. ex Claus, in Beitr. Pfl. Russ. Reich. viii. 254 = leptocaula.
astroites, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* x. 44.—Mexic.
astyla, *Engelm. ex Boiss. in DC. Prod.* xv. ii. 40.—Mexic.
atlantica, *Coss. ex Boiss. l. c.* 130.—Algeria.
atlantica, Pers. Syn. ii. 18 = bupleuroides.
Atoto, *Forst. f. Prod.* 86.—As. et Austral. trop.; Ins. Pacif.
Atoto, Guill. in Ann Sc. Nat. Sér. II. vii. (1837) 137 = falcata.
atropurpurea, *Brouss. Elench. Pl. Hort. Bot. Monsp.* (1806) 24, nomen; *Willd. Enum. Hort. Berol.* 501.—Teneriffe.
atrorubra, Engelm. ex Boiss. in DC. Prod. xv. ii. 101 = E. Ipecacuanha.

EUPHORBIA :—
atrosanguinea, Poepp. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 70 = portulacoides.
Aubryana, *Baill. Adansonia*, ii. (1861–62) 213.—N. Caled.
Auchori, *Boiss. Diagn.* Ser. I. vii. 94.—Persia.
aulacosperma, *Boiss. l. c.* xii. 117.—Syria.
auricularia, Boiss. Cent. Euph. 17 = rosea ?
australis, *Boiss. l. c.* 15.—Austral.
avenia, Thibaut, ex Pers. Syn. ii. 15 = imbricata.
axorica, Hochst. in Seub. Fl. Axor. 27 = pinea.
axorica, Walw. ex Boiss. in DC. Prod. xv. ii. 144 = medicaginea.
baetica, vide boetica.
bahiensis, *Boiss. in DC. Prod.* xv. ii. 24.—Bras.
Baillonii, Boiss. ex Baill. Adansonia, i. (1860–61) 142 = Bosseri.
Balbisii, *Boiss. Cent. Euph.* 11.—Ins. Baham.
balearica, Poir. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 73 = imbricata.
balsamifera, *Ait. Hort. Kew.* ed. I. ii. 137.—Ins. Canar.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 463.—Ins. Banca.
barbellata, *Engelm. in Torr. Bot. Mex. Bound.* 190.—Mexic.
Barrelieri, *Savi, Bot. Etrusc.* i. 145.—Ital.
Basilicia, Tenore, Prod. Fl. Nap. p. xxix = Barrelieri.
Baueri, *Engelm. ex Boiss. in DC. Prod.* xv. ii. 27.—Austral.
Beaumlerana, *Hook. f. & Coss. in Bull. Soc. Bot. Fr.* xxi. (1874) 184.—Marocc.
begoniaefolia, *Lehm. ex Steud. Nom.* ed. II. i. 610, nomen.—Hab. ?
bejarensis, DC. in Atti Reun. Second. Sc. Ital. Tor. (1844) 707 = lamygiasta.
belgradica, *Forsk. Fl. Aegypt. Arab.* 211.—Byzant.
Berteriana, *Balb. ex Spreng. Syst.* iii. 794.—Ind. occ.
Bertheloti, *Bolle, ex Boiss. in DC. Prod.* xv. ii. 109.—Ins. Canar.
Berythea, *Boiss. & Blanch. Diagn.* Ser. II. iv. 82.—Syria.
Besseri, *Boiss. in DC. Prod.* xv. ii. 49.—Chili.
bialata, Link, Enum. Hort. Berol. ii. 13 = pterococca.
bicephala, Bertol. Misc. Bot. iii. 20 = Berteriana.
bicolor, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 233.—Texas.
bifida, *Hook. & Arn. Bot. Beech. Voy.* 218.—China.
bifida, Thw. Enum. Pl. Zeyl. 269 = E. Atoto.
biformis, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 151.—Mexic.
bifurcata, *Engelm. in Torr. Bot. Mex. Bound.* 190.—N. Mexic.
biglandulosa, Boiss. in DC. Prod. xv. ii. 165 = micaceiada.
biglandulosa, *Desf. in Ann. Mus. Par.* xii. (1808) 114.—Graecia; As. Min.
biglandulosa, Willd. Enum. Hort. Berol. Suppl. 27 = pendula ?
bilobata, *Engelm. in Torr. Bot. Mex. Bound.* 190.—N. Mexic.
binervia, Miq. Fl. Ind. Bat. i. II. 420 = congenera.
biumbellata, *Poir. Voy. Barb.* ii. 174.—Afr. bor.
biumbellata, Us. Itin. ex Boiss. in DC. Prod. xv. ii. 149 = Cupani.
Bivonae, *Steud. Nom.* ed. II. i. 610.—Afr. bor.; Sicil.
Blanchetii, Miq. in Linnaea, xxii. (1849) 797 = Madag.
blepharophylla, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 188.—Soongaria.
boerhaviifolia, *Boiss. in DC. Prod.* xv. ii. 54.—Am. centr.
boetica, *Boiss. Cent. Euph.* 36.—Hispan.
Boissieri, *Baill. Adansonia*, i. (1860–61) 142.—Madag.
Boivini, *Boiss. in DC. Prod.* xv. ii. 77.—Madag.
Bojeri, *Hook. Bot. Mag.* t. 5527.—Madag.
bombensis, Jacq. Enum. Pl. Carib. 23 = ammenioides.
bongensis, *Kotschy & Peyr. Pl. Timor.* 40. t. 19.—Afr. trop.
bonifaciensis, Requien, ex Gren. & Godr. Fl. Fr. iii. 56 = E. Pithyusa.
Bonplandii, Swnt, Hort. Brit. ed. II. 456 = tortida.
borbonica, *Boiss. in DC. Prod.* xv. ii. 156.—Ins. Borbon.
bothlospermea, *Boiss. & Kotschy, in Boiss. Cent. Euph.* 36.—Armenia.

EUPHORBIA :—
botryoides, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 14 (Quid ?).—Java.
Bottae, *Boiss. in DC. Prod.* xv. ii. 95.—Arabia.
bounophila, *Boiss. l. c.* 154.—Persia.
Bourgaeana, *J. Gay, ex Boiss. l. c.* 108.—Ins. Teneriff.
brachiata, Jan, Elench. 7 = palustris.
brachiata, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 74.—Afr. austr.
brachyora, *Engelm. in Torr. Bot. Mex. Bound.* 192.—N. Mexic.
bracteata, Jacq. Hort. Schoenb. iii. 14. t. 276 = Pedilanthus bracteatus.
bracteolaris, Boiss. Cent. Euph. 8 = hypericifolia.
brasiliensis, *Lam. Encyc.* ii. 428.—Am. trop.
Brousi, Ann. Fl. & Pom. (1833) 189 = Bojeri.
Bridgesii, Berr. ex Boiss. in DC. Prod. xv. ii. 103 = portulacoides.
Brittingeri, Opiz, ex Reichb. Fl. Germ. Excurs. 757 = verrucosa.
Broteri, Daveau, in Bol. Soc. Broter. iii. (1885) 33 = nicaeensis.
Brousseauii, Willd. ex Link, in Buch, Beschr. Canar. Ins. 158 = obtusifolia.
Brownii, *Baill. Adansonia*, vi. (1866) 290.—Australi.
bubalina, *Boiss. Cent. Euph.* 26.—Afr. austr.
buchtormensis, *C. A. Mey. ex Ledeb. Fl. Alt.* iv. 189.—Sibir.
Bubani, *Boiss. in DC. Prod.* xv. ii. 167.—Persia.
Bungei, *Boiss. l. c.* 118.—Persia.
bupleurifolia, *Jacq. Hort. Schoenb.* i. 55. t. 106.—Afr. austr.
bupleurifolia, E. Mey. in Drège, Zwei Pfl. Docum. 184 = oxyleuja.
bupleuroides, *Desf. Fl. Atlant.* i. 337. t. 103.—Afr. bor.
bupleuroides, Willd. ex Boiss. in DC. Prod. xv. ii. 159 = leptocaula.
bupleuroides, Willd. ex Ledeb. Fl. Ross. iii. 571 = tenuifolia.
Burchelli, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 680.—Bras.
Burmanni, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 75.—Afr. austr.
Burmanniana, J. Gay, in Webb & Berth. Phyt. Canar. iii. 239 = thymifolia.
buxifolia, *Lam. Encyc.* ii. 421.—Ind. occ.
Cactus, *Ehrenb. ex Boiss. in DC. Prod.* xv. ii. 82.—Arabia.
cacorum, *Mart. ex Boiss. l. c.* 51.—Bras.
caelalensis, *Boiss. Diagn.* Ser. II. iv. 81.—Beluchist.
caerulescens, Haw. in Phil. Mag. (1826) 276 = virosa.
caesaraugustana, *Lacat. ex Willk. & Lange, Prod. Fl.* Hisp. i. 505, nomen.—Hispan.
caesia, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 743 = E. Esula.
caespitosa, vide cespitosa.
caespitosa, Tenore, Fl. Nap. iv. t. 261. t. 143 = pinea.
Cafogala, Ehrh. in Hannov. Mag. 14tes Stück (1783) 213; Beitr. ii. 109 = Geraldina.
calcarea, Coss. & Dur. ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 227 = pinea.
calendulaefolia, Delile, Fl. Egypte, 233. t. 80. f. 1 = nigvia.
californica, *Benth. Bot. Voy. Sulph.* 49. t. 228.—Calif.
californica, Boiss. Cent. Euph. 6 = Deppeana.
callisedens, Engelm. ex Boiss. in DC. Prod. xv. ii. 87 = hispida.
callitrichioides, H. B. & K. Nov. Gen. et Sp. ii. 58 = prostrata.
callitrichoides, Schan. in Linnaea, xx. (1847) 737 = adenoptera.
calycoclada, *H. B. & K. Nov. Gen. et Sp.* ii. 59.—Mexic.
calyptrata, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 473, nomen; *Boiss. in DC. Prod.* xv. ii. 113.—Algeria.
campestris, *Chem. & Schlecht. in Linnaea*, v. (1830) 84.—Mexic.
canadiensis, Lam. Encyc. ii. 417 = E. Cliusa.
canaliculata, Ledd. Bot. Cab. t. 737 = Pedilanthus tithymaloides.
canaliculata, Pers. Syn. ii. 13 = graminea.
canariensis, Forsk. Fl. Aegypt. Arab. p. xcii = polycnemifolia ?

EUPHORBIA:—
canariensis, *Linn. Sp. Pl.* 450.—Ins. Canar.
canariensis, Thunb. Fl. Cap. ed. Schult. 408 = tetragona.
canariensis, Trémaut, Voy. Soudan Orient. Atlas, tt. 13, 14 = E. Candelabrum.
Candelabrum, *Trémaut, ex Kotschy, in Mittheil. Geogr. Gesells. Wien*, i. (1857) 199.—Afr. trop.
canescens, Linn. Sp. Pl. ed. II. 652 = E. Chamaesyce.
Cannii, *Parl. Fl. Ital.* ix. 466.—Alp. marit.
capitata, Buch.-Ham. ex Wall. Cat. n. 7708 A = crinita.
capitata, Lam. Encyc. ii. 422 = pilulifera.
capitellata, *Engelm. in Torr. Bot. Mex. Bound.* 188.—Mexic.
capitulata, *Reichb. Fl. Germ. Excurs.* 878.—Thessal.; Graecia.
Caput-Medusae, *Linn. Sp. Pl.* 452.—Afr. austr.
caracassana, *Boiss. in DC. Prod.* xv. ii. 60.—Venezuela.
cardiophylla, *Boiss. & Heldr. Diagn.* Ser. I. xii. 107.—Lycia.
Careyi, *F. Muell. Fragm.* xi. 64.—Austral.
carinata, Doon, Hort. Cantab. ed. VI. 181 = Pedilanthus tithymaloides.
carinthiaca, Traunf. ex Boiss. in DC. Prod. xv. ii. 116 = pilosa.
carniolica, Brot. Fl. Lusit. ii. 317 = boetica.
carniolica, DC. Fl. Fr. iii. 542 = flaviccma.
carniolica, *Jacq. Fl. Austr.* v. 34. t. 14.—Tyrol.
carnitolica, Lapeyr. Hist. Abr. Pl. Pyr. 372 = hybernia.
carpatica, Ehrenb. ex Boiss. in DC. Prod. xv. ii. 98 = cuneata.
casabonensis, Royle, Illustr. Bot. Himal. 329. t. 82 = pilosa!
Cassia, *Boiss. Diagn.* Ser. I. xii. 108.—Syria.
cassioides, Presl, Bot. Bemerk. 109 = hypericifolia.
casuarhoides, *Boiss. Cent. Euph.* 30.—Cuba.
Cattimandoo, W. Ell. in Wight, Ic. t. 1993 = trigona.
caudata, Boiss. & Haussk. in DC. Prod. xv. ii. 1267 = falcata.
caudiculosa, *Boiss. in DC. Prod.* xv. ii. 154.—Syria.
cebrina, Hochst. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 86 = cerebrina.
cechica, Opiz, Seznam, 62 = pilosa.
celastroides, *Boiss. in DC. Prod.* xv. ii. 11.—Ins. Sandvic.
centunculoides, *H. B. & K. Nov. Gen. et Sp.* ii. 51.—Cuba.
ceratocarpa, *Tenore, Fl. Nap.* i. 268. t. 48.—Sicil.
cerebrina, *Hochst. ex Boiss. in DC. Prod.* xv. ii. 146.—Abyss.
cereiformis, *Linn. Sp. Pl.* 451.—Afr. austr.
cerinthifolia, Fisch. ex Boiss. in DC. Prod. xv. ii. 123 = alpina.
cernua, *Cos. & Dur. ex Boiss. l. c.* 188.—Algeria.
cervicornis, *Boiss. Cent. Euph.* 27.—Afr. austr.
cespitosa, *Lam. Encyc.* ii. 431.—Reg. Argent.
cestrifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 59.—Peruv.
Chaixiana, Timb. in Mém. Acad. Toul. Sér. IV. vi. (1856) 153 = amygdaloides.
Chamaebuxa, *Bernard. ex Gren. & Godr. Prosp. Fl. Fr.* 3.—Mont. Pyren.
Chamaepeplus, *Boiss. & Gaill. Diagn.* Ser. II. iv. 88.—Palaest.
Chamaerrhodos, *Boiss. Cent. Euph.* 3.—Bras.
Chamaesaxi, Benth. Fl. Austral. vi. 49, sphalm. = *E. Chamaesyce*, Bail.
Chamaesula, *Boiss. Cent. Euph.* 38.—N. Mexic.
Chamaesyce, Baill. Adansonia, vi. (1866) 285 = Drummondii.
Chamaesyce, *Linn. Sp. Pl.* 455.—Reg. Mediterr.
Chamaesyce, C. A. Mey. in Ledeb. Fl. Alt. iv. 196 = hamiltona.
Chamaesyce, Roxb. Hort. Beng. [36]; Fl. Ind. ii. 478 = microphylla !
Chamissonis, *Boiss. in DC. Prod.* xv. ii. 14.—Ins. Fiji.
Characias, Host, Fl. Austr. ii. 508 = Wulfenii.
Characias, *Linn. Sp. Pl.* 468.—Reg. Mediterr. occ.
Characias, Sibth. & Sm. Fl. Graec. Prod. i. 332 = Sibthorpii.
chelonophora, *Boiss. & Heldr. Diagn.* Ser. I. xii. 112.—Persia.
chersonesa, *Pluck. & Mey. in Bull. Soc. Nat. Mosc.* (1856) 171, nomen; *Boiss. in DC. Prod.* xv. ii. 70.—Transcauc.

EUPHORBIA:—
chenopodifolia, *Boiss. in DC. Prod.* xv. ii. 1264.—Bolivia.
Chesneyi, *Boiss. l. c.* 152.—Assyria.
chilensis, Echeg. Symb. 852. n. 79 = Schickendantii.
chilensis, C. Gay, Fl. Chil. v. 385 = portulacoides.
chrysophylla, *Klotzsch, ex Boiss. in DC. Prod.* xv. ii. 104.—Bras.
ciliata, *Sprang. Fl. Hal. Mant.* 43.—Hab.!
ciliata, Boiss. Diagn. Ser. II. iv. 86 = denticulata.
cinerascens, Engelm. in Torr. Bot. Mex. Bound. 186 = melanadenia.
clandestina, *Jacq. Hort. Schoenb.* iv. 43. t. 484.—Afr. austr.
Clava, *Jacq. Ic. Pl. Rar.* i. 9. t. 85; *Coll.* i. 104.—Afr. austr.
clavarioides, *Boiss. Cent. Euph.* 25.—Afr. austr.
clavata, Salisb Prod. 389 = E. Clava.
Clementei, *Boiss. Elench.* 92.—Hispan.
Clementei, Bourg. ex Boiss. in DC. Prod. xv. ii. 131 = rupicola.
Cleopatra, *Baill. Adansonia*, ii. (1861–62) 213.—N. Caled.
clusiaefolia, *Hook. & Arn. Bot. Beech. Voy.* 95.—Ins. Sandvic.
coccinea, *Roth, Nov. Pl. Sp.* 228.—Ind. or.
coccinea, Willd. ex Boiss. in DC. Prod. xv. ii. 71 = pulcherrima.
Coderiana, DC. Fl. Fr. vi. 365 = platyphylla.
corcorum, Muell. Arg. in Mart. Fl. Bras. xi. ii. 675 = cacorum.
coriadenia, vide caeladenia.
corniculacea, vide caerulescens.
cornuta, Boiss. in DC. Prod. xv. ii. 120 = pilosa.
collctioides, *Benth. Bot. Sulph. Voy.* 168.—Mexic.
collina, *Phil. in Linnaea*, xxix. (1857–58) 41.—Chili.
collina, Willd. ex Ledeb. Fl. Ross. iii. 578 = nicaeensis.
colorata, *Engelm. in Torr. Bot. Mex. Bound.* 190.—Mexic.
Commelini, DC. Cat. Hort. Monsp. 110 = E. Caput-Medusae.
commutata, *Engelm. ex A. Gray, Man. Bot. N. U. St.* ed. II. 389.—Am. bor.
comosa, *Vell. Fl. Flum.* 202; v. t. 15.—Bras.
compressa, *Boiss. in DC. Prod.* xv. ii. 41.—Venezuela.
Conceptionis, Rupr, ex Boiss. in DC. Prod. xv. ii. 102 = portulacoides.
condylocarpa, *Bieb. l. c.* i. 377.—Armenia.
confusa, Blume, ex Boiss. in DC. Prod. xv. ii. 30 = hirsutissima.
congemera, *Blume, Bijdr.* 634.—Malaya.
congesta, Willd. Enum. Hort. Berol. Suppl. 28 = pinea.
consimilis, Steph. ex Boiss. in DC. Prod. xv. ii. 123 = pachyrrhiza.
coniosperma, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 196.—Persia.
connata, Boiss. in DC. Prod. xv. ii. 112.—Persia.
consanguinea, Engelm. ex Boiss. in DC. Prod. xv. ii. 48 = serpyllifolia.
consanguinea, Klotzsch, in Bot. Ergeb. Waldem. Reise, 114 = pilosa.
consanguinea, *Schrenk, Enum. Pl. Nov.* i. 88.—Sibir.
convolvuloides, *Hochst. ex Boiss. in DC. Prod.* xv. ii. 46.—Afr. trop.
copiapina, *Phil. in Linnaea*, xxix. (1857) 42.—Chili.
corallioides, *Linn. Sp. Pl.* 460.—Italia.
corallioides, Sprun. ex Nym. Consp. 649 = oblongata.
corallioides, Thunb. ex Boiss. in DC. Prod. xv. ii. 168 = Stokoldina.
cordata, *Meyen, Reise*, ii. 150.—Ins. Sandvic.
cordata, Schrank, Baier. Fl. i. 747 = dulcis.
cordelilata, Haw. Misc. 185 = Pedilanthus cordatus.
cordifolia, *Ell. Sketch*, ii. 656.—Am. bor.
cordifolia, C. A. Mey. ex Boiss. Cent. Euph. 40 = rumicifolia.
corticosa, C. Koch, in Linnaea, xxi. (1848) 780 = magalantha.
corticosifolia, Boiss. in DC. Prod. xv. ii. 15 = Hookeri.
cortifolia, *Lam. Encyc.* ii. 431.—Afr. austr.
cornigera, Boiss. in DC. Prod. xv. ii. 122 = pilosa.

EUPHORBIA:—
cornuta, *Pers. Syn.* ii. 17.—Aegypt.; Arabia.
corollata, *Linn. Sp. Pl.* 459.—Am. bor.
coronata, *Thunb. Prod. Pl. Cap.* 56.—Afr. austr.
correntina, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 49.—Reg. Argent.
corrigioloides, *Boiss. Cent. Euph.* 11.—Ind. or.
corsica, *Requien, in Ann. Sc. Nat.* Sér. I. v. (1825) 884.—Ins. Corsic.
cosmosperma, Reichb. Ic. Fl. Germ. v. & pinea.
Cossoniana, *Boiss. in DC. Prod.* xv. ii. 135.—Alger.
costata, Schur, in Verh. Siebenb. Ver. Naturwiss. iii. (1852) 132; iv. (1853) 55 = pilosa.
cotinifolia, H. B. & K. Nov. Gen. et Sp. ii. 46 = carcassana.
cotinifolia, *Linn. Sp. Pl.* 458.—Ind. occ.
cotinoides, *Mig. in Linnaea*, xxi. (1848) 478; *Stirp. Surin. Sel.* 96.—Guian.
cotyliformis, Steud. Nom. ed. I. 324, sphalm. = seq.
cotyliophora, *Sprang. in Schrad. Journ.* (1800) ii. 197.—Hab.
craspedia, *Boiss. Diagn.* Ser. I. vii. 95.—Syria.
crenulata, Engelm. in Torr. Bot. Mex. Bound. 192 = leptocera.
cretica, Mill. Gard. Dict. ed. VIII. n. 28 = E. Characias.
crispata, Hornem. Hort. Hafn. Suppl. 58 = pubescens.
crispata, Lem. Illustr. Hortic. iv. (1857) Misc. 71 = Lemaireana.
cristata, D. Dietr. Syn. Pl. v. 321 = denticulata.
cristata, *Heyne, ex Roth, Nov. Pl. Sp.* 226.—Ind. or.
crotonoides, *Boiss. in DC. Prod.* xv. ii. 98.—Nubia.
cruenta, Steud. Nom. ed. II. i. 611 = seq!
cruentata, *R. Grah. in Edinb. N. Phil. Journ.* (Oct. 1832) 361.—Am bor.
cubensis, *Boiss. in DC. Prod.* xv. ii. 1265.—Cuba.
cucumerina, *Willd. Sp. Pl.* ii. 886.—Afr. austr.
Cumbrae, *Boiss. Cent. Euph.* 16.—Mexic.
cuneata, *Vahl, Symb. Bot.* ii. 88.—Arabia.
cunefolia, *Guss. Pl. Rar.* 190. t. 38.—Reg. Mediterr.
cunnifolia, Roxb. Hort. Beng. [91]; Fl. Ind. ii. 471 = prolifera.
Cupani, *Guss. Fl. Sic. Prod.* i. 548.—Ins. Sicil. et Sardin.
Cupani, Schultz, ex Gren. & Godr. Fl. Fr. iii. 86 = E. Pithyusa.
cuphosperma, *Boiss. in DC. Prod.* xv. ii. 78.—N. Mexic.
cupularis, Boiss. Cent. Euph. 33 = Synadenium arborescens.
curtifolia, Chaub. Nouv. Fl. Pélop. 80 = E. Myrsinites.
Curtisii, *Engelm. in Chapm. Fl. S. U. St.* 401.—Am. bor.
cuspidata, Bernh. ex Krauss, in Flora, xxviii. (1845) 86 = striata.
cuspidata, Bertol. Misc. Bot. ii. 9; iii. 17 = hypericifolia.
cyanogala, *Wright, ex Griseb. Cat. Pl. Cub.* 21.—Cuba.
cyathophora, Murr. in Comm. Gotting. vii. (1786) 81 = heterophylla.
cybirensis, *Boiss. Diagn.* Ser. I. vii. 89.—Reg. Mediterr. occ.
cymosa, Polr. Encyc. Suppl. ii. 611 = nudiflora.
Cyparissias, *Linn. Sp. Pl.* 461.—Europ.
cypria, *Boiss. in DC. Prod.* xv. ii. 116.—Ins. Cyprus.
Dallachyana, Baill. Adansonia, vi. (1866) 285 = Drummondii.
dalmatica, Vis. Fl. Dalmat. iii. 228 = graeca.
damascena, Boiss. Diagn. Ser. I. xii. 112 = tinctoria.
daphnoides, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 21.—Ins. Masc.
Darlingtonii, *A. Gray, Man. Bot. N. U. St.* ed. I. 404.—Am. bor.
decaryana, *Cos. in Bull. Soc. Bot. Fr.* xxii. (1875) 67, nomen.—Marocc.
decipiens, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 197.—Persia.
decumbens, Willd. Enum. Hort. Berol. Suppl. 27 = hypericifolia.
decussata, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 74.—Afr. austr.
decussata, Salisb. Prod. 389 = E. Lathyris.
deflexa, *Sibth. & Sm. Fl. Graec. Prod.* i. 328.—Graecia.
delicatula, *Boiss. Cent. Euph.* 19.—Mexic.
Delorti, Timb. ex Nym. Consp. 655 = E. Terracina.
deltoides, *Engelm. ex Chapm. Fl. S. U. St. Suppl.* 647.—Am. bor.

EUPHORBIA:—
dendroides, *Linn. Sp. Pl.* 462.—Reg. Mediterr.
dens, Schott & Kotschy, ex Tchihat. Asie Min. Bot. i. 187.—Schottians.
dens, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1846) 308.—Persia.
densiflora, *Klotzsch, in Peters, Reise Mossamb. Bot.* 94, in obs.—Mexic.
dentatifolia, *C. Koch, in Linnaea*, xxi. 1848 [1849] 722. —Reg. Caucas.
dentata, *Michx. Fl. Bor.-Am.* ii. 211.—Am. bor.
denticulata, *Lam. Encyc.* ii. 481.—As. Min.: Syria.
denudata, Bertol. Fl. Ital. v. 91 = corallioides.
depauperata, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 22, nomen; *A. Rich. Tent. Fl. Abyss.* ii. 241.—Abyss.
Deppeana, *Boiss. Cent. Euph.* 6.—'Calif.'
deprimæ, C. Gay, Fl. Chil. v. 386 = Engelmanni.
depressa, Phil. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 26 = ovalifolia.
depressa, Torr. ex Spreng. Syst. iii. 794 = maculata.
Desegliesi, Bor. ex Boiss. in DC. Prod. xv. ii. 128 = angulata.
desertícola, F. Muell. in Linnaea, xxv. (1852) 440 = cremophila.
desertorum, *Wemm. in Bull. Soc. Nat. Mosc.* (1837) n. vii. 78.—Rossia.
dichotoma, *Forsk. Fl. Aegypt. Arab.* 98.—Egypt.
dichotoma, Roxb. Hort. Beng. [91]; Fl. Ind. ii. 471 = elegans?
dictyosperma, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 37.—Am. bor.
diffusa, Defour, in Bull. Soc. Bot. Fr. vii. (1860) 445 = rupicola.
diffusa, *Hook. f. in Trans. Linn. Soc. xx.* (1851) 184. —Ins. Galapag.
diffusa, Jacq. Misc. ii. 311 = exigua.
dilatata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 240. —Abyss.
dilatata, E. Mey. in Drège, Zwei Pfl. Docum. 184 = Meyeri.
dilatata, Torr. & Gray, in Pacif. Rail. Rep. ii. 175 = alata.
dioeca, H. B. & K. *Nov.* Gen. et Sp. ii. 58 = adenoptera.
dioica, *Hieron. Ic. Pl. Argent.* 46.—Reg. Argent.
dioscoreoides, *Boiss. Cent. Euph.* 29.—Mexic.
discoidalis, *Chapm. Fl. S. U. St.* 401.—Florida.
discolor, Bertol. Misc. Bot. x. 87 = corollata.
discolor, *Boiss. in DC. Prod.* xv. ii. 54.—Am. centr.
discolor, Klotzsch, ex Boiss. in DC. l. c. 104 = elodes.
discolor, Ledeb. Fl. Ross. iii. 577 = E. Esula.
discolor, Shuttlew. ex Boiss. in DC. Prod. xv. ii. 119 = Darlingtonii.
distícha, Kogelm. ex Boiss. l. c. 82 = corrigioloides.
distincta, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 65 = pilosa.
distincta, Stschegl. in Bull. Soc. Nat. Mosc. xxvii. (1854) i. 196 = subcordata.
distinguenda, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 122 = pilosa.
divaricata, A. Cunn. ex Benth. Fl. Austral. vi. 50 = Wheeleri.
diversicoma, Jacq. Ic. Pl. Rar. i. 9. t. 87 = dendroides.
diversgens, Klotzsch, in Bot. Ergeb. Waldem. Reise, 114 = Rothiana.
diversifolia, Hochst. ex Boiss. in DC. Prod. xv. ii. 146 = pilosa.
diversifolia, Pohl. Encyc. Suppl. ii. 618; Willd. Enum. Hort. Berol. Suppl. 37 = E. Characias.
diversifolia, *Schrank*, ex Steud. Nom. ed. I. 324 = buplenroides.
diversifolia, Willd. ex Boiss. in DC. Prod. xv. ii. 71 = pulcherrima.
djumlensis, *Boiss. Fl. Orient.* iv. 1104.—As. Min.
domingensis, Spreng. ex Boiss. in DC. Prod. xv. ii. 84 = lasciocaula.
dracunculoides, *Lam. Encyc.* ii. 493.—As. et Afr. trop.
drastica, Stevens, in Pall. N. Nord. Beitr. vii. 286 = alpina.
Dregeana, *E. Mey. in Drège, Zwei Pfl. Docum.* 184; nomen; *Boiss. in DC. Prod.* xv. ii. 95.—Afr. austr.
Drummondii, *Boiss. Cent. Euph.* 14.—Australi.
dryadum, *Thunb. in Schum. & Thonn. Beskr. Guin. Pl.* 280.—Afr. trop.
dubia, Dierb. Fl. Heidelb. ii. 137 = stricta.
dulcis, Asso, Syn. Stirp. Arag. 59 = lucidifolia.
dulcis, Bertol. Fl. Ital. v. 90 = verrucosa.
dulcis, *Linn. Sp. Pl.* 467.—Europ.

EUPHORBIA:—
dulcis, Ruegel, ex Boiss. in DC. Prod. xv. ii. 190 = ruscinonensis.
dulcis, Sibth. & Sm. Fl. Græc. v. 50. t. 464 = thamnoides.
dumetorum, Coss. & Dur. ex Boiss. in DC. Prod. xv. ii. 181 = tripolensis.
dumosa, Boiss. Diagn. Ser. II. xii. 119 = thamnoides.
dumosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 188.—Afr. austr.
dumosa, H. Rich. Fl. Cub. Taxorog. ii. 196 = trichotoma.
Duvalii, *Lecoq & Lamotte, Cat. Pl. Vasc. Fr.* 327.—Gallia.
echinata, Salm-Dyck, Hort. Dyck. 342 = cereiformis.
echinocarpa, *Presl, Rel. Haenk.* i. (1835) 102; *ex Bot. Zeit.* i. (1843) 561, nomen.—Hab.?
echinocarpa, Sieber, ex Boiss. in DC. Prod. xv. ii. 119 = cyparissias.
Echinus, *Hook. f. & Coss. in Bull. Soc. Bot. Fr.* xxi. (1874) 164.—Marocc.
Edgeworthii, *Boiss. in DC. Prod.* xv. ii. 120.—Reg. Himal.
edulis, *Lour. Fl. Cochinch.* 298.—Cochinch.
effusa, Ehrenb. ex Boiss. in DC. Prod. xv. ii. 71 = squamosa.
Ehrenbergii, Sweet, Hort. Brit. ed. II. 455 = E. Terracina.
Eichleri, *Muell Arg. in Journ. Bot.* xii. (1874) 332.—Reg. Argent.
elegans, Heync, in Wall. Cat. n. 7718 = longistyla.
elegans, *Spreng. Syst.* iii. 794.—Ind. or.
elliptica, Boiss. Cent. Euph. 18 = saccharata.
elliptica, Lam. Encyc. ii. 495 = geniculata.
elliptica, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.
elodes, *Boiss. Cent. Euph.* 23.—Texas.
elongata, Poir. Encyc. Suppl. ii. 621 = buplenroides.
emarginata, Ait. Hort. Kew. ed. I. ii. 145 = hyberna.
emarginata, *Boiss. in DC. Prod.* xv. ii. 93.—Bras.
emarginata, Lam. Encyc. ii. 496 = myrtilliifolia.
emirnensis, *Baker, in Journ. Linn. Soc. xx.* (1883) 251. —Madag.
Engelmanni, Boiss. Cent. Euph. 15 = ovalifolia.
ennaegona, Haw. Misc. 184 = cereiformis.
enopia, *Boiss. Cent. Euph.* 27.—Afr. austr.
enstifolia, *Baker, in Journ. Linn. Soc. xx.* (1883) 251. —Madag.
ephedroides, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 76.—Afr. austr.
epicyparissias, Kraus, ex Boiss. l. c. 168 = involucrata.
epicyparissias, E. Mey. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 38 = Wahlbergii.
epicyparissias, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 168.—Afr. austr.
epiphylloides, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) 11. 247.—Ins. Andam.
epithymoides, All. Fl. Pedem. i. 324 = verrucosa.
epithymoides, Bab. Fl. Bath, 44 = pilosa.
epithymoides, Brot. Fl. Lusit. ii. 311 = lusitanica.
epithymoides, *Linn. Sp. Pl.* ed. II. 656; *Jacq. Fl. Austr.* iv. 38. t. 344.—Europ.
eremophila, *A. Cunn. ex Hook. in Mitch. Journ. Trop. Austral.* 348.—Austral.
eriantha, *Benth. Bot. Voy. Sulph.* 31.—Calif.
ericostorum, *Zumag. Fl. Pod.* ii. 168.—Italia.
ericoides, *Lam. Encyc.* ii. 480.—Afr. austr.
erinacea, *Boiss. & Kotschy, Diagn.* Ser. II. iv. 87.—Syria.
erioearpa, Bertol. in Nov. Comm. Acad. Sc. Bonon. iii. (1839) 174 = E. Characias.
eriocalda, Boiss. & Heldr. Diagn. Ser. I. vii. 88 = oblongata.
eriocalda, Sart. ex Nym. Consp. 649 = pubescens.
eriophora, *Boiss. Diagn.* Ser. I. v. 51.—As. Min.; Pers.
eriophylla, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 764 = subcordata.
erosa, Willd. Enum. Hort. Berol. Suppl. 37 = cereiformis.
erubescens, Boiss. Diagn. Ser. I. vii. 90 = macrostegia.
erubescens, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 110.—Afr. austr.
erythradenia, *Boiss. Diagn.* Ser. I. vii. 92.—Persia.
erythranthe, F. Muell. Fragm. ii. 152 = australis.
erythrina, *Link, Enum. Hort. Berol.* ii. 12.—Afr. austr.
erythrocarpa, *Klotzsch, in Schomb. Reun. & Pl. Brit. Gui.* 848, nomen.—Guianas.
erythrodada, *Boiss. in DC. Prod.* xv. ii. 98.—Ind. or.
erythradenia, *Boiss. & Heldr. in Boiss. Diagn.* Ser. I. xii. 134.—As. Min.

EUPHORBIA:—
erythrophylla, Bertol. Fl. Guat. 19. t. 6 = pulcherrima.
erythrorhiza, *Boiss. in DC. Prod.* xv. ii. 108.—Bras.
erythrocyclades, *Baker, in Journ. Linn. Soc.* xx. (1883) 253.—Madag.
Esula, Bieb. Fl. Taur. Cauc. i. 376 = virgata.
Esula, Forsk. Fl. Aegypt. Arab. 94 = Schimperiana.
Esula, Kotschy, ex Boiss. in DC. Prod. xv. ii. 160 = virgata.
Esula, *Linn. Sp. Pl.* 461.—Europ.
Esula, Pollich, Hist. Pl. Palat. ii. 18 = Gerardiana.
Esula, Tenore, Fl. Nap. i. 373 = E. Cyparissias.
esuloeformis, Schau. in Linnæa, xx. (1847) 739 = camposria.
esuloides, Tenore, Syll. Fl. Neap. 388 = E. Cyparissias.
exclusa, *S. Wats. in Proc. Am. Acad.* xviii. (1883–83) 150.—Mexic.
exigua, *Linn. Sp. Pl.* 461.—Europ.
expansa, Janka, in Linnæa, xxx. (1859–60) 600 = carniolica.
exstipulata, *Engelm. in Torr. Bot. Mex. Bound.* 189. —Texas.
falcata, *Linn. Sp. Pl.* 456.—Reg. Mediterr.; As. calid.
fasciculata, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.
Fendleri, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 175.—N. Mexic.
Ferdinandi, Baill. Adansonia, vi. (1866) 234 = Drummondii.
fidjiana, *Boiss. in DC. Prod.* xv. ii. 110.—Ins. Fiji.
filicina, *Porteuscl. Enum. Pl. Daim.* 15. t. 8.—Dalmat.
filipes, *Benth. Fl. Austral.* vi. 51.—Austral.
fimbriata, Heyne, ex Roth, Nov. Pl. Sp. 237 = crenata.
fimbriata, Hort. Par. ex Boiss. in DC. Prod. xv. ii. 81 = Lamarckii.
fimbriata, Scop. Delic. Insub. iii. 8 = Scopolina.
fimbriata, Wall. ex Boiss. in DC. Prod. xv. ii. 126 = khasyana.
fimbrilligera, Mart. in Linnæa, xix. (1847) 407 = ocymoides.
firma, Ledeb. Fl. Ross. iii. 563 = Gerardiana.
Fischeriana, Steud. Nom. ed. II. i. 611 = Pallasii.
flabellaris, *Anders. ex Boiss. in DC. Prod.* xv. ii. 17.—Ins. Galapag.
flagelliformis, *Engelm. in Hayden, Bull. Geol. Geogr. Surv.* ii. (1876) 245.—Am. bor. occ.
flaviconia, *DC. Fl. Fr.* vi. 364.—Hispan.
flavo-purpurea, Willd. in Cenir. Bot. Zeitschr. xxv. (1876) 118 = platyphylla.
Fleurotii, Jord. in Billot, Annot. 97 = E. Esula.
flexicaulis, Sonder, in Linnæa, xxii. (1849) 153 = serpens.
flexuosa, H. B. & K. Nov. Gen. et Sp. ii. 55 = buxifolia.
floribunda, *Engelm. ex Boiss. in DC. Prod.* xv. ii. 89. —Mexic.
floridæ, *Engelm. in Torr. Bot. Mex. Bound.* 189.—Mexic.
floridana, Chapm. Fl. S. U. St. 401 = sphærosperma.
foetida, Hoppe, ex Koch, Syn. Fl. Germ. ed. II. 734 = stricta.
foetida, Schult. ex Reichb. Fl. Germ. Excurs. 756 = platyphylloa.
foliata, Buch.-Ham. ex Dillw. Rev. Hort. Malab. 52 = thymifolia.
folioloses, *Boiss. in DC. Prod.* xv. ii. 94.—Am. trop.
fontanesii, Steud. Nom. ed. II. i. 611 = dentata.
Forskalei, Bourg. ex Boiss. in DC. Prod. xv. ii. 47 = prostrata.
Forskålii, J. Gay, in Webb & Berth. Phyt. Canar. iii. 240 = granulata, ægyptiaca.
fourcadei, *Boiss. & Gaill. Diagn.* Ser. II. iv. 87 = anisacanthera.
fragifera, *Jan, ex Link, Enum. Hort. Berol.* ii. 16.—Italia.
fragifera, Schur, Enum. Fl. Transs. 592 = epithymoides.
fragilis, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 241 = granulata.
Franconna, *Boiss. Cent. Euph.* 22.—Hispan.
frankenifolia, H. B. & K. Nov. Gen. et Sp. ii. 62 = graminifolia.
Fraseri, *Boiss. in DC. Prod.* xv. ii. 56.—Kentuc.
Friedrichsthalii, *Boiss. l. c.* 61.—Guatemala.
Fructus-Pini, Mitt. Gard. Diet. ed. VIII. n. 28 = E. Caput-Medusæ.
fruticosa, Bivona, Sic. Pl. Cent. i. 85 = Biverona.
fruticosa, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1219 = cuneata.
fruticosa, *Forsk. Fl. Aegypt. Arab.* 94.—Arab.

EUPHORBIA:—
fruticulosa, Engelm. ex Boiss. in DC. Prod. xv. ii. 88. —Mexic.
fulgens, Karw. ex Klotzsch, in Otto & Dietr. Allg. Gartenz. ii. (1834) 96.—Mexic.
furcillata, H. B. & K. Nov. Gen. et Sp. ii. 60.—Mexic.
fusiformis, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 68.—Ind. or.
gaditana, Coss. Notes Crit. 46.—Hispan.
Gaillardoti, Boiss. & Blanche, Diagn. Ser. II. iv. 84. —Mesopotam.
galilæa, Boiss. Diagn. Ser. I. xii. 116 = falcata.
galioides, Boiss. Cent. Euph. 14 = microphylla.
Garberi, Engelm. ex Chapm. Fl. S. U. St. Suppl. 646.—Florida.
garipina, Boiss. Cent. Euph. 28.—Afr. austr.
Garkeana, Boiss. in DC. Prod. xv. ii. 88.—Ins. S. Doming.
Gasparrinii, Boiss. l. c. 125.—Sicilia.
Gaudichaudii, Boiss. Cent. Euph. 7.—Ins. Marian.
Gayi, Salis, in Flora, xvii. (1834) ii. *Beibl.* 6.—Ins. Contic.
gemella, Lag. Gen. et Sp. Nov. 17 = pilulifera.
geminispina, How. ex Steud. Nom. ed. II. i. 612. —Mexic.
geniculata, Ortag. Hort. Matr. Dec. 16.—Am. trop.
geniculoides, Berg. Desc. Pl. Cap. 146; *Linn. Mant.* ii. 254.—Afr. austr.
Gerardiana, *Jacq. Fl. Austr.* v. 17. t. 486.—Europ.
Germaini, *Phil. in Linnæa*, xxix. (1857–58) 40. —Chili.
Geyeri, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 260.—Am. bor.
glaberrima, Klotzsch. in Seem. Bot. Voy. Herald, 276. —Mexic.
glaberrima, C. Koch, in Linnæa, xxi. (1848) 736 = tirophylla.
glabrata, Sw. Prod. Veg. Ind. Occ. 76 = buxifolia.
glabriflora, Vis. in Mem. Ist. Venet. xii. (1864) 477 = pubescens!
glaricos, *Pall. ex Bieb. Fl. Taur. Cauc.* i. 378.—Europ.; Orient.
glauca, Ehrenb. ex Boiss. in DC. Prod. xv. ii. 140 = dracunculoides.
glauca, *Forst. f. Prod.* 36.—N. Zel.
glauca, Roxb. Fl Ind. ii. 478 = Rothiana.
glaucescens, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 138 = Gerardiana.
glaucophylla, *Petr. Encyc. Suppl.* ii. 618.—Afr. trop.
globulosa, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 498.—Algeria.
globosa, *Sims, Bot. Mag.* t. 2934.—Afr. austr.
globulifera, H. B. & K. Nov. Gen. et Sp. ii. 56 = pilulifera.
globulosa, Boiss. in DC. Prod. xv. ii. 189 = globosa.
glomerata, *Bieb. ex Steud. Nom.* ed. I. 325, nomen.—Reg. Caucas.
glyptosperma, *Engelm. in Torr. Bot. Mex. Bound.* 186. —Mexic.
Gmelini, Steud. Nom. ed. II. i. 612 = virgata.
Goeringii, Steud. ex Boiss. in DC. Prod. xv. ii. 30 = humifusa.
Goliana, *Lem. Encyc.* ii. 438.—Ins. Borbon.
Gollmariana, *Klotzsch, ex Boiss. in DC. Prod.* xv. ii. 58.—Venezuela.
Gondoti, *Boiss. l. c.* 106.—N. Granat.
goyazensis, *Boiss. Cent. Euph.* 28.—Bras.
gracillima, Baill. Adansonia, i. (1860–61) 140 = pyrifolia.
gracilis, Bess. Ind. Hort. Crem. (1816) 56 = E. Esula.
gracilis, Ell. Sketch, ii. 657 = E. lpecacuanha.
gracilis, Loisel. Fl. Gall. ed. I. 738 = tenuifolia.
gracilis, Pav. ex Hoq. in DC. Prod. xiii. ii. 369 = Amazanthes crosolatus.
graeca, *Boiss. & Sprun. Diagn.* Ser. I. v. 58.—Græc.; An Mis.
graminea, *Jacq. Select. Am.* 151.—Am. centr.
graminea, Koch. ex Boiss. in DC. Prod. xv. ii. 156 = Rothiana.
graminea, Schlecht. & Cham. in Linnæa, v. (1830) 88 = scabrella.
graminifolia, Michx. Fl. Bor. Am. ii. 210 = granulata.
graminifolia, Vill. Fl. Delphin. 47 = tenuifolia.
graminifolia, How. in *Phil. Mag.* (1832) 88.—Afr. austr.
graminifolia, Haw. Syn. Pl. Succ. 126 = dregeana.
granulata, Lam. Illustr. Botriz. iv. (1837) Misc. 101 = stipulacea.
Grantii, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 144. —Afr. trop.
granulata, *Forsk. Fl. Aegypt. Arab.* 94.—Afr. et As. trop.
Greggii, *Engelm. ex Boiss. in DC. Prod.* xv. ii. 147.—Mexic.
griffithii, Boissier, ex Boiss. l. c. 41.—Mexic.
Guachanca, *Ataxa, Voy.* ii. 497.—Peruv.
Gueinzii, *Boiss. in DC. Prod.* xv. ii. 71.—Afr. austr.
Guilfoilei, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–9) 406 = Sieboldiana.
guineensis, *Brot. Cat. Hort. Ajuda, ex Bot. Zeit.* i. (1843) 561, nomen.—Afr. trop.
gummifera, *Boiss. Cent. Euph.* 26.—Afr. austr.
Guyoniana, *Boiss. & Reut. Pugill. Pl. Nov.* 109.—Algeria.
gymnoclada, *Boiss. Cent. Euph.* 20.—Bras.
gymnoclada, Engelm. in Proc. Am. Acad. v. (1861) 171 = Xanti.
haematantha, *Boiss. in DC. Prod.* xv. ii. 63.—Ecuador.
haematodes, *Boiss. l. c.* 179.—Hab.?
halophila, Miq. Anal. Bot. Ind. iii. 16 = E. Atoto.
hamata, *Sweet, Hort. Brit.* ed. III. 597.—Afr. austr.
Hartwegiana, *Boiss. in DC. Prod.* xv. ii. 31.—Mexic.
Hausknechtii, Boiss. l. c. 126? = E. helioscopia.
hawaiensis, Wild. ex Boiss. l. c. 73 = heterophylla.
Hawortkii, Sweet, Hort. Brit. ed. I. 356 = E. Clava.
Hawortkii, Sweet, l. c. 357 = spinosa.
hebecarpa, *Boiss. Diagn.* Ser. I. vii. 90.—Persia.
Heldreichii, *Orph. ex Boiss. l. c.* Ser. II. iv. 90.—Græcia.
helicotheke, Lem. Illustr. Hortic. iv. (1857) Misc. 100 = E. Characias.
helioscopia, Haussk. ex DC. Prod. xv. ii. 126? = Hausknechtii.
heliscopioides, Losc. & Pard. Ser. Inconf. Arag. 93 = heloscopia.
heptagona, *Linn. Sp. Pl.* 450.—Afr. austr.
Herminieri, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 63. —Afr. trop.
hermlariæfolia, *Willd. Sp. Pl.* ii. 902.—As. Min.
hermitoides, Nutt. in Trans. Am. Phil. Soc. v. (1837) 171 = serpens.
Herronii, Riddell, Syn. Fl. W. St. 82 = dentata.
heterodena, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Moscou, xii. (1860) 198 = erythradenia.
heteradena, Jamb. & Spach, Illustr. ii. 42. t. 131 = megalantha.
heteranthe, Nutt. in Trans. Am. Phil. Soc. v. (1837) 171 = hexagona.
heterodoxa, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 701. —Bras.
heterophylla, Desf. Fl. Atlant. i. 385. t. 102 = E. Terracina.
heterophylla, *Linn. Sp. Pl.* 458.—Am. bor.
hexagona, *Nutt. ex Spreng. Syst.* iii. 791.—Texas.
Heyneana, Boiss. in DC. Prod. xv. ii. 35, partim = microphylla?
Heynlana, Spreng. Syst. iii. 791 = microphylla.
hiberna, Lesch. ît. i. 291 = dulcis.
hiberna, Welw. ex Nym. Consp. 649 = Welwitschii.
hibernica, Spreng. Syst. iii. 800 = hyberna.
hieroglyphica, *Coss. & Dur. ex Boiss. in DC. Prod.* xv. ii. 144.—Algeria.
hierosolymitana, Boiss. Diagn. Ser. I. xii. 110 = thamnoides.
himalayensis, Boiss. in DC. Prod. xv. ii. 118 = sec.?
himalayensis, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 118.—Reg. Himal.
Hindsiana, *Benth. Bot. Voy. Sulph.* 51. t. 24.—Calif.
hirsuta, Kit. ex Boiss. in DC. Prod. xv. ii. 116 = pilosa.
hirsuta, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 66 = E. Esula.
hirta, Linn. Sp. Pl. 454 = pilulifera.
hirtella, *Boiss. Cent. Euph.* 7.—Bras.
hirtula, *Engelm. ex S. Wats. in Bot. Calif.* ii. 74.—Calif.
hispida, *Boiss. Cent. Euph.* 8.—Reg. Himal.
Hoffmanniana, *Boiss. in DC. Prod.* xv. ii. 99.—Costa Rica.
Hohenackeri, Hochst. ex Boiss. l. c. 167 = Gerardiana.
Hohenackeri, Orph. ex. Boiss. Diagn. Ser. II. iv. 89 = Orphanidis.
homophylla, Lange, ex Boiss. in DC. Prod. xv. ii. 167 = Gerardiana.
Hookeri, *Steud. Nom.* ed. II. i. 612.—Ins. Sandvic.
horrida, *Boiss. Cent. Euph.* 27.—Afr. austr.
hortensis, Engelm. ex Boiss. l. c. 15 = Engelmanni.
huschanahana, *Ruiz, ex Klotzsch & Garcke, in Abh. Akad. Berl.* (1860) 71.—Peruv.; Boliv.
huanchahana, Boiss. in DC. Prod. xv. ii. 108 = pnæc.
Humboldtii, Willd. Enum. Hort. Berol. 608 = graminea.
humifusa, Hort. ex Boiss. in DC. Prod. xv. ii. 42 = Engelmanni.
humifusa, *Willd. Enum. Hort. Berol. Suppl.* 27.—Sibir.
humilis, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 185.—Sibir.
humilis, Royle, Illustr. Bot. Himal. 259. t. 62. f. 2 = fusiformis.
humistrata, *Engelm. ex A. Gray, Man. Bot. N. U. St.* ed. II. 386.—Am. bor.
hyberna, *Linn. Sp. Pl.* 462.—Europ.
hyberna, Viv. Fl. Corn. 7 = insularis.
Hydnorae, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 95.—Afr. austr.
hypericifolia, Aust. Flor. ex Boiss. l. c. 22 = Preslii.
hypericifolia, Hochst. ex Boiss. l. c. 22 = indica.
hypericifolia, *Linn. Sp. Pl.* 454.—Amphig. trop.
hypericifolia, Phil. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 34 = thyrsiperma.
hyssopifolia, Linn. Syst. ed. X. 1049 = brasiliensis.
Hystrix, *Jacq. Hort. Schoenb.* iii. 48. t. 307.—Afr. austr.
Iberica, *Boiss. Cent. Euph.* 28.—Reg. Caucas.
illirica, Lam. Encyc. ii. 436 = pilosa.
imbricata, *Vahl, Symb. Bot.* ii. 54.—Lusitania.
imperfolata, Vis. Fl. Dalmat. iii. 527 = Silicina.
inæquilatera, Engelm. in Torr. Bot. Mex. Bound. 187 = serpyllifolia.
inæquilatera, Sond. in Linnæa, xxiii. (1850) 105 = sanguinea.
incana, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 197 = salicifolia.
incisa, *Engelm. in Torr, Rep. Colorado River Bot.* 27. —Arizona.
inclinata, Hort. ex Boiss. in DC. Prod. xv. ii. 42 = Engelmanni.
incompta, Cesati, in Bibl. Ital. xci. (1838) 348 = dulcis.
incongrua, Bull, in Journ. Bot. xiii. (1875) 205.—Marocc.
indersiana, Less. ex Ksr. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 448 = pygmæa.
indica, Lam. Encyc. ii. 433 = hypericifolia.
indica, Wall. Cat. n. 7710 = roxburghiana.
indica, Willd. Gard. Dict. ed. VIII. n. 13.—Hab.?
inermis, Pand. ex Boiss. in DC. Prod. xv. ii. 126? = fragifera!
ingens, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. l. c.* 87.—Afr. austr.
insulana, *Vell. Fl. Flum.* 202; v. t. 14.—Bras.
insularis, *Boiss. Cent. Euph.* 25.—Corsica, Sicil.
intermedia, Brth. Fl. Normand. ed. III. 271 = E. Esula.
intermedia, Fisch. & Mey. ex Boiss. in DC. Prod. xv. ii. 163 = iberica.
intermedia, Hochst. ex Boiss. l. c. 35 = sanguinea.
inundata, *Torr. ex Chapm. Fl. S. U. St.* 402.—Florida.
invoiucrata, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 168.—Afr. austr.
involucrata, *Wall. Cat.* n. 7696.—Reg. Himal.
Ipecacuanha, *Linn. Sp. Pl.* 455.—Am. bor.
ischidiifolia, *Lem. Encyc.* ii. 430.—Hispan.
ixaia, Pers. Syn. ii. 15 = pnæc.
japonica, Boiss. Diagn. Ser. I. vii. 91 = megalantha.
Italica, *Lam. Encyc.* ii. 434.—Italia.
italica, Salzm. ex Boiss. Voy. Bot. Espagne, ii. 570 = medicaginea.
italica, Tineo, Syn. Hort. Panorm. (1802) 18 = E. Terracina.
tirophylla, *Boiss. Cent. Euph.* 29.—Reg. Caucas.
Jacquemontii, *Boiss. in DC. Prod.* xv. ii. 113.—Reg. Himal.
Jacquini, *Fenzl, ex Boiss. l. c.* 106.—Hab.?
jacquiniæflora, Hook. Bot. Mag. t. 3678 = fulgens.
Jamesoni, *Boiss. Cent. Euph.* 11.—Ecuador.
japonica, *Siebold, ex Boiss. in DC. Prod.* xv. ii. 1266, ex parte.—Japon.
japonica, Zoll. ex Boiss. l. c. 127 = helioscopia.
jappigica, Tenore, Fl. Nap. iv. 266 = nicæensis.
javanica, *Pungh. in Nat. en Genees. Arch. Nederl. Ind.* ii. (1845) 44.—Java.
Jolkini, *Boiss. Cent. Euph.* 23.—Japon.

EUPHORBIA:—
juncea, Ait. Hort. Kew. ed. I. ii. 142 = aleppica.
juncoides, Stend. Nom. ed. II. i. 612 = prace.
juniperifolia, Kth. ex Boiss. in DC. Prod. xv. II. 164, 1267 = pauciflora.
kabirensis, Rasanch. Nom. ed. III. 140 = cornuta.
Kaleniczenkii, Cserm. ex Trautv. in Act. Hort. Petrop. ix. (1884) 159, nomen = E. Cyparissias.
kanaiensis, *Boiss. in DC. Prod.* xv. II. 1265.—N. Caled.
kanaorica, *Boiss. l. c.* 154.—Reg. Himal.
karwinskyi, *Boiss. Cent. Euph.* 6.—Mexic.
kaschgarica, *Regel in Act. Hort. Petrop.* vi. (1879) 401.—Turkestan.
Kerneri, *Huter, ex A. Kern. Sched. Fl. Exsicc.* ii. 48. —Europ. austr.
khasyana, *Boiss. in DC. Prod.* xv. II. 120.—Reg. Himal.
Klotzschiana, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Sel.) 95 = brasiliensis.
Kotschyana, *Fenzl, Pugill. Pl. Nov. Syr.* 7.—As. Min.
Krausiana, Bernh. ex Krauss, in Flora, xxviii. (1845) 87 = crubescens.
Kunzei, Lehm. ex Ind. Sem. Hort. Hamb. (1830) 4; Linnæa, v. (1830) 375 = Lagascæ.
kurdica, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* iv. 1191.—Persia.
lacera, *Boiss. Euph. Cent.* 22.—Mexic.
lactea, *Ham. Syn. Pl. Succ.* 137.—Ind. or.
lacten, *Roxb. Hort. Beng.* [86]; *Fl. Ind.* ii. 468.—Ins. Moluce.
lactiflua, *Phil. Fl. Atac.* 48.—Chili.
laeta, Ait. Hort. Kew. ed. I. ii. 141 = dendroides.
laeta, Heyne, ex Roth, Nov. Pl. Sp. 230 = Rothiana.
lævigata, *Lam. Encyc.* ii. 421.—Bras.
lævigata, Vahl, Symb. Bot. ii. 84 = E. Atoto.
lævis, vide levis.
Lagascæ, *Spreng. Neue Entdeck.* ii. 115.—Sardinia.
Lagascæ, Spun. ex Nym. Consp. 654 = graeca.
Lamarckii, Sweet, Hort. Suburb. Lond. 107 = obtusifolia.
lanata, *Sieber, ex Spreng. Syst.* iii. 792.—Syria; Persia.
lancolaria, Herb. Heyne, ex Wall. Cat. n. 7891 = Rothiana.
lanceolata, Spreng. Fl. Hal. Mant. 41 = dracunculoides.
lanciflola, *Schlecht. in Linnæa*, vii. (1832) 148.—Mexic.
lanifera, *Haw. ex Steud. Nom.* ed. II. i. 612, nomen.—Mexic.
lanuginosa, Lam. Encyc. ii. 436 = corallioides.
lanuginosa, Thunb. Fl. Par. ed. II. 238 = platyphylla.
lanica, *Boiss. Cent. Euph.* 24.—Persia.
lastocarpa, *Klotzsch, in Nov. Act. Nat. Cur.* xix. Suppl. i. (1843) 414.—Ind. occ.; Mexic.
lasiocarpa, C. Koch, in Linnæa, xxi. (1848) 721 = eriophora.
lasiocaula, Boiss. in DC. Prod. xv. II. 1266 = pekinensis.
lata, *Engelm. in Torr. Bot. Mex. Bound.* 188.—Texas.
Latasi, *H. B. & K. Nov. Gen. et Sp.* ii. 58.—N. Granat.
laterifiora, Janb. & Spach, Illustr. ii. 41 = osyrides.
laterifiora, *Schum. & Thonn. Beskr. Guin. Pl.* 262.—Afr. trop.
Lathyris, Georgi, Beschr. Russ. Reich. III. iv. 998 = Pallasii.
Lathyris, *Linn. Sp. Pl.* 457 (Lathyrus).—Europ. austr.
latifolia, *Gueldenst. ex Ledeb. Fl. Ross.* iii. 579 (quid?). —Rossia.
latifolia, [Hoben.] Boiss. Cent. Euph. 39 = iberica.
latifolia, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 168.—Sibir.
latifolia, Salzm. ex Boiss. Voy. Bot. Espagne, ii. 870 = saxicaginea.
laurifolia, *Lam. Encyc.* ii. 418.—Peruv.
Ledebourii, *Boiss. Cent. Euph.* 36.—Taurin.
leiosperma, Boiss. l. c. 5 = obliqua.
leiosperma, Sibth. & Sm. Fl. Græc. v. 51. t. 465 = E. Terracina.
Lemaireana, *Boiss. in DC. Prod.* xv. II. 81.—Afr. trop.
lepidota, Schan. in Linnæa, xx. (1847) 738 = graminea.
leptocaula, *Boiss. in DC. Prod.* xv. II. 159.—Rossia.

EUPHORBIA:—
leptocera, *Engelm. ex Torr. in Pacif. Rail. Rep.* iv. 135.—Am. bor. occ.
leptoclada, *Baill. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 410.—Ins. Socotr.
leptophylla, Vill. Hist. Pl. Dauph. iii. 825 = tenuifolia.
leucanthis, *Boiss. in DC. Prod.* xv. II. 43.—Mexic.
leucoloma, Kaſin. Herb. (1838) 29; Atl. Journ. 177 = marginata.
leuconeura, *Boiss. in DC. Prod.* xv. II. 78.—Madag.
leucophylla, *Benth. Bot. Voy. Sulph.* 50.—Calif.
leucotricha, Boiss. Voy. Bot. Espagne, ii. 585. t. 160 = pubescens.
levis, Polr. Encyc. Suppl. ii. 612 = E. Atoto.
ligularia, Roxb. Hort. Beng. [86]; Fl. Ind. ii. 465 = nerilifolia.
ligularia, Chaub. in Ann. Sc. Obs. iv. t. 6 = amygdaloides.
ligustrina, *Boiss. Cent. Euph.* 21.—Hispan.
Linaria, Link, in Buch, Beschr. Canar. Ins. 158 = E Terracina.
linariæfolia, Desf. Tabl. ed. I. 204 = E. Terracina.
linariæfolia, Lam. Encyc. ii. 437 = Gerardiana.
linariæfolia, Polr. Encyc. Suppl. ii. 611 = spinosa.
Lindeniana, A. Rich. Fl. Cub. Fanerog. ii. 197 = adenopters.
Lindheimeriana, Engelm. ex Boiss. in DC. Prod. xv. II. 142 = longicuria.
linearifolia, *Roth, Nov. Pl. Sp.* 224.—Ind. or.
linearifolia, Willd. Sp. Pl. ii. 893 = spinosa.
linearis, Heyne, ex Roth, Nov. Pl. Sp. 225 = linearifolia.
linearis, Heyne, in Wall. Cat. n. 7712 = linearifolia.
linearis, *Lam. Encyc.* iii. 93.—Ind. occ.
linearis, Schrank, in Denkschr. Akad. Muench. vi. 1816–17 (1820) 131 = cornuta.
linguiata, Pieuf. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 192 = epithymoides.
linifolia, *Burm. f. Fl. Cap. Prod.* 14.—Afr. austr.
linifolia, Herb. Russ. ex Wall. Cat. n. 7691 = linifolia.
linifolia, Jacq. Coll. i. 135 = spinosa.
linifolia, Tenore, Prod. Fl. Nap. p. xxix = pinea.
linifolia, Vahl, Sym. Bot. ii. 58 = restiacea.
linifolia, Wall. ex Boiss. in DC. Prod. xv. II. 157 = prolifera.
literata, Jacq. Coll. ii. 340 = platyphyllos.
litoralis, H. B. & K. Nov. Gen. et Sp. ii. 54 = buxifolia.
litoralis, Noé, ex Nym. Consp. 654 = pinea.
littorea, Miq. ex Boiss. in DC. Prod. xv. II. 102 = corallioides.
livida, *C. A. Mey. in Delge, Zwei Pfl. Docum.* 184 nomen; *Boiss. in DC. Prod.* xv. II. 14.—Afr. austr.
Lockhartii, Stend. Nom. ed. II. i. 613 = heterophylla.
longetuberculata, Boiss. l. c. 1264 = seq.
longetuberculosa, *Hochst. ex Boiss. l. c.* 85.—Abyss.
longibracteata, DC. Fl. Fr. v. 356 = segetalis.
longicruria, *Schoele, in Linnæa*, xxii. (1849) 152.—Texas.
longifolia, Baill. ex Boiss. in DC. Prod. xv. II. 78 = Boissieri.
longifolia, *D. Don, Prod. Fl. Nep.* 163.—Reg. Himal.
longifolia, Gueldenst. Reise, i. 192 = leptocaula.
longifolia, Lam. Encyc. ii. 417 = multiflora.
longitradiata, *Labrey. Suppl. Hist. Pl. Pyr.* 65.—Mont. Pyren.
longiuscula, *Boiss. Cent. Euph.* 9.—Ind. or.
lophogona, *Lam. Encyc.* ii. 417.—Madag.
Lorentii, Hochst. in Lorent, Wanderungen, 344 = tinctoria.
Lorentzii, *Msall. Aeg. in Journ. Bot.* xii. (1874) 231. —Reg. Argent.
Lowyei, Jard. in Billot, Annot. 26 = E. Esula.
lucida, Lam. Encyc. ii. 416 = E. Hystrix.
lucida, Auct. Gall. ex Boiss. in DC. Prod. xv. II. 162 = sadrosemifolia.
lucida, *Waldst. & Kit. Pl. Rar. Hung.* i. 54. t. 54.—Mexic.
lucorum, *Rupr. in Maxim. Prim. Fl. Amur.* 389.—Reg. Amur.
ludoviciana, Raſin. Fl. Ludov. 111 = corollifolia.
lunulata, *Bunge, Enum. Pl. Chin. Bor.* 69.—China.
lupulina, *Boiss. Cent. Euph.* 28.—N. Granat.
lurida, *Engelm., in Ives, Rep. Colorado River Bas.* 96. —Calif.
lustanica, *Steud. Nom.* ed. II. i. 613.—Lusitan.

EUPHORBIA:—
lutea, Alam. ex Boiss. in DC. Prod. xv. II. 73 = pulcherrima.
luteola, *Cass. & Dur. ex Boiss. l. c.* 150.—Algeria.
lutescens, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 194.—Sibir.
lycia, *Boiss. in DC. Prod.* xv. II. 172.—Lycia.
lycioides, *Boiss. Cent. Euph.* 39.—Rossia.
Maackii, *Meinshaus. Nachr. Wilni-Geb.* 104; *ex Trautv. Act. Hort. Petrop.* ix. (1884) 160.—Sibir.
Macgillivrayi, *Boiss. in DC. Prod.* xv. II. 96.—Anstral.
macrocarpa, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 197.—Persia.
macrocerus, *Pirsch, & Mey. Ind. Sem. Hort. Petrop.* iv. 36.—Reg. Caucas.
macrocladæ, Boiss. Diagn. Ser. I. v. 54 = tinctoria.
macroglypha, *Lem. Illustr. Hortic.* iv. (1857) *Misc* 73.—Madag.
macropus, *Boiss. in DC. Prod.* xv. II. 53.—Mexic.
macrorhiza, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 191.—Sibir.
macrorhiza, Pall. ex Ledeb. Fl. Ross. iii. 565 = Pallasii.
macrostegia, *Boiss. in DC. Prod.* xv. II. 171.—Persia.
maculata, Andrews, in Vet. Acad. Handl. Stockh. 1858 (1855) 283 = fabellaris.
maculata, Aubl. Pl. Gui. i. 479 = thymifolia.
maculata, Lechl. ex Boiss. in DC. Prod. xv. II. 42 = Meyenlana.
maculata, *Linn. Sp. Pl.* 455.—Am. bor.
madagascariensis, Comm. ex Lam. Encyc. ii. 417 = lophogona.
Maddeni, *Boiss. in DC. Prod.* xv. II. 141.—Reg. Himal.
Magdalenæ, *Benth. Bot. Voy. Sulph.* 50.—Calif.
magnidens, Haw. ex Salm-Dyck, Hort. Dyck. 105 = grandidens.
magnimamma, *Haw. ex Steud. Nom.* ed. II. i. 613.—Mexic.
malacophylla, Clarke, Trav. ii. 354 = lanata.
mallicata, *Boiss. in DC. Prod.* xv. II. 152.—Persia.
mamillana, Willd. ex Boiss. in DC. Prod. xv. II. 28 = ammanioides.
Marschalliana, *Boiss. Diagn.* Ser. I. vii. 94.—Ins. Min.; Persia.
Marschalliana, Kotschy, ex Boiss. in DC. Prod. xv. II. 173 = E. Myrsinites.
Martiusiana, Steud. Nom. ed. II. i. 613 = Lagascæ.
Mauryæna, Phil. in Bot. Zeit. xiv. (1856) 647 = Wahlenbergia tuberosa.
massiliensis, DC. Fl. Fr. vi. 357 = E. Chamæsyce.
mastrichensis, *Boiss. Cent. Euph.* 35.—Hispan.
mauritanica, Lam. Encyc. ii. 418 = obtusifolia.
mauritanica, *Linn. Sp. Pl.* 459.—Afr. austr.
mauritanica, Webb, Florul. Lanzerot. 31; ex Webb & Berth. Pl. Canar. iii. 230 = E. Regis-Jubæ.
medicaginea, *Boiss. Elench.* 83.—Reg. Mediterr. occ.
Meduse, *Thumb. Prod. Pl. Cap.* 96.—Afr. austr.
megalalistion, *Bail, in Journ. Bot.* xiii. (1875) 205.—Marocc.
megalantha, *Boiss. Diagn.* Ser. I. vii. 95.—Persia.
mekaniensis, Kit. in Roehel, Pl. Banat. Rar. 96, nomen; et Linnæa, xxxii. (1863) 556 = epithymoides.
melanadenia, *Torr. in Pacif. Rail. Rep.* iv. 135.—Calif.
melanocarpa, *Boiss. in DC. Prod.* xv. II. 43.—Ecuador.
melanostieta, *E. Mey. in Delge, Zwei Pfl. Docum.* 134, nomen; *et ex Boiss. in DC. Prod.* xv. II. 98.—Afr. austr.
melapetala, *Gasparr. Ind. Sem. Hort. Boccod.* (1839) 9; *ex Guss. Suppl. Fl. Sic. Prod.* 168.—Sicil.
mellifera, Pers. Fl. Ind. iv. 471 = spinosa.
mellifera, *Ait. Hort. Kew.* ed. I. ii. 468.—Ins. Madera.
mellifera, Semb. Fl. Amer. 37 = stygiana.

EUPHORBIA:—
meloformis, *Ait. Hort. Kew.* ed. I. ii. 135.—Afr. austr.
melonoformis, Lem. Jard. Fleur. iv. (1854) Misc. 65 = pmec.
Mendesii, *Boiss. Cent. Euph.* 16.—Mexic.
mercurialina, *Michx. Fl. Bor. Am.* ii. 212.—Am. bor.
Mercurialis, Pers. Syn. ii. 13 = pmec.
mesembrianthemifolia, Jacq. Enum. Pl. Carib. 22; Select. Am. 150 = buxifolia.
Meyenana, *Klotzsch*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 414.—Bolivia.
Meyeri, *Boiss. Cent. Euph.* 35.—Afr. austr.
Meyeri, Steud. Nom. ed. II. i. 618 = rupestris.
micans, Scheele, in Linnæa, xvii. (1843) 843 = amygdaloides.
micracantha, *Boiss. Cent. Euph.* 25.—Afr. austr.
micractina, *Boiss. in DC. Prod.* xv. ii. 137.—Reg. Himal.
microdenia, *Boiss. l. c.* 77.—Austria.
micrantha, Steph. ex Willd. Sp. Pl. ii. 905 = stricta.
microcephala, *Boiss.* in *DC. Prod.* xv. ii. 1262 — Bolivia.
micromera, *Boiss. l. c.* 44.—N. Mexic.
microphylla, *Heyne, ex Roth, Nov. Pl. Sp.* 229.—Ind. or.
microphylla, Lam. Encyc. ii. 423 = thymifolia.
microscadia, Boiss. Diagn. Ser. I. vii. 89 = striatella microsphæra, *Boiss. l. c.* 87.—Mexicopet.
Milii, Desmoul. ex Boiss. in DC. Prod. xv. ii. 79 = splendens.
minima, Hort. ex Link, Enum. Hort. Berol. ii. 12 = pepioides.
minuta, Losc. & Pard. Ser. Inconf. Arag. 96 = paucifora.
minutiflola, *Boiss. in DC. Prod.* xv. ii. 1263.—Austral.
minutula, *Boiss. l. c.*—Cuba.
mitera, *Benth. Bot. Voy. Sulph.* 51.—Calif.
missourica, *Rafin. Atl. Journ.* 146.—Am bor.
Mitchelliana, *Boiss. in DC. Prod.* xv. ii. 25.—Austral.
modesta, Boiss. Cent. Euph. 34 = E. Terracina.
Momchiana, *Stransl. Nom. ed.* II. i. 618.—Hab.?
mohammmerensis, Boiss. in DC. Prod. xv. ii. 118 = microphaera.
mollis, C. C. Gmel. Fl. Bad. ii. 330 = pilosa.
monantha, *Wright, ex Boiss. in DC. Prod.* xv. ii. 55.—Cuba.
montana, *Engelm. in Torr. Bot. Mex. Bound.* 192.—N. Mexic.
Montetri, *Hook. Bot. Mag* t. 5534.—Afr. trop.
montevidensis, Casar. ex Boiss. in DC. Prod. xv. ii. 43 = ovalifolia.
montevidensis, Klotzsch, ex Boiss. l. c. 105 = papilosa.
monticola, Boiss. Diagn. Ser. I. vii. 93 = bourgæana.
monticola, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 242.—Abyss.
Morisoniana, Klotzsch, in Seem. Bot. Voy. Herald. 100 = heterophylla, barbellata.
mosana, Lej. Fl. Spa, i. 318 = E. Esula.
mosambicana, *Boiss. in DC. Prod.* xv. ii. 36.—Afr. trop.
mucronata, *Clarke, Trav.* ii. 260.—As. Min.
mucronata; Lam. Encyc. ii. 427 = falcata.
mucronata, Lapeyr. Hist. Abr. Pl. Pyr. 271 = E. Pithyusa.
Mueller, *Boiss. in DC. Prod.* xv. ii. 27.—Australi.
multicaulis, *Engelm. in Torr. Bot. Mex. Bound.* 191. —Mexic.
multicaulis, Thuill. Fl. Par. ed. II. 238 = nicæensis.
multicorymbosa, *Sweet, Hort. Brit.* ed. I. 357.—Hab.?
multiflora, Willd. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 31 = adenopetes.
multiformis, *Gaudich. in Freyc. Voy. Bot.* 100, nomen; *Hook. & Arn. Bot. Beech. Voy.* 95.—Ins. Sandvic.
multiformis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 65 = pilosa.
multiseta, *Benth. Pl. Hartw.* 51.—Mexic.
muricata, Bieb. Fl. Taur. Cauc. i. 378 = aspera.
muricata, Sm. Engl. Fl. iv. 65, in texts = alexocarpa.
muricata, Thunb. Prod. Pl. Cap. 36 = brachiata.
Myrsinites, Brot. Fl. Lusit. ii. 317 = nicæensis.
Myrsinites, *Linn. Sp. Pl.* 463.—Europ. austr. or.
Myrsinites, Pall. Ind. Taur. ex Ledeb. Fl. Ross. iii. 579 = Marschalliana.
myrtifolia, Hook. & Arn. Bot. Beech. Voy. 95 = Ekebergii.

EUPHORBIA:—
myrtifolia, Lam. Encyc. ii. 419 = Pedilanthus tithymaloides.
myrtifolia, Linn. Sp. Pl. ed. II. 653 = myrsilifolia.
myrtifolia, E. Mey. in Drège, Zwei Pfl. Docum. 184 = sclerophylla.
myrtilifolia, *Linn. Syst.* ed. X. 1048.—Jamaic.
myrtilloides, Schlecht. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 85 = buxifolia.
Myrtoides, *Boiss. in DC. Prod.* xv. ii. 15.—Austral.
Naglezi, *Boiss. l. c.* 26.—Java.
nana, Klotzsch, ex Boiss. Cent. Euph. 8 = E. Chamaerrhodos.
nana, Royle, Illustr. Bot. Himal. 329, t. 82 = fusiformis.
natalensis, *Bernh. ex Krauss, in Flora*, xxviii. (1845) 86.—Afr. austr.
neapolitana, Tenore, Fl. Nap. ii. 266 = E. Terracina.
nemoralis, Darling. Fl Cestr. ed. II. 518 = Darlingtonn nemoralis, *Kit. in Rochel, Pl. Banat. Rar.* 26, nomen. —Europ.
nemoralis, Salisb. Prod. 390 = amygdaloides.
neo-caledonica, *Boiss. in DC. Prod.* xv. ii. 1263.—N. Caled.
nepalensis, Boiss. l. c. 157 = prolifera.
nerrifolia, Roxb. Hort. Beng. 86; Fl. Ind. ii. 467 = E. Nivulia.
nerifolia, *Linn. Sp. Pl.* 451.—Ind. or.; Malaya.
nervosa, Kit. in Linnæa, xxxii. (1863) 559 = rugosa.
nevadensis, *Boiss. & Reut. Pugill. Pl. Nov.* 110.—Hispan.
nicæensis, *All. Fl. Pedem.* i. 285. t. 69.—Reg. Mediterr
nicæensis, Coss. ex Willk. & Lange, Prod. Fl Hisp. iii. 801 = matritensis.
nicæensis, St. Am. Fl. Agen. 192 = E. Esula.
nicæensis, Sebast. & Maur. Fl. Rom. Prod. 164 = E. Terracina.
nilagirica, Miq. Anal. Bot. Ind. iii. 17 = sanguinea.
nitens, Trevir. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 149 = agraria.
Nivulia, *Buch.-Ham. in Trans. Linn. Soc.* xiv. (1825) 236.—Afr. or.
nodiflora, Steud. Nom. ed. II. i. 618 = pilulifera.
Notana, Boiss. in DC. Prod. xv. ii. 166 = tinctoria.
norfolkiana, *Boiss. l. c.* 110.—Ins. Norfolk.
notadarnia, Boiss. & Hohen. Diagn. Ser. I. xii. 111 = orientalis
notata, Engelm. ex Boiss. DC. Prod. xv. ii. 43 = serpillifolia.
notopters, *Boiss. in l. c.* 26.—Ind. or.
nuda, Pers. Syn. ii. 13, in syn. = graminea.
nudicaulis, *Chapm. Fl. S. U. St.* 402.—Florida.
nudicaulis, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 116.—Malaya.
nudiflora, *Jacq. Coll.* iii. 180.—Jamaic.
nudiflora, Lam. Encyc. ii. 426 = graminea.
Nummularia, *Hook f. in Trans. Linn. Soc.* xx. (1851) 183.—Ins. Galapag.
nummulariæfolia, Willd. Enum. Hort. Berol. 608 = imbricata.
nutans, Lag. Gen. et Sp. Nov. 17 = Preslii.
obconica, *Boy. Hort. Maurit.* 287.—Afr. occ.
obcordata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 93.—Ins. Socotr.
oblanceolata, *Balf. f. l. c.*—Ins. Socotr.
oblonga, *Endl. Prod. Fl. Norf.* 35.—Ins. Norfolk.
obliquata, Forsk. Fl. Aegypt. Arab. 93 = E. Terracina.
obliquifolia, Kunze, ex Boiss. Cent. Euph. 15 = ovalifolia.
obtetorata, Jacq. Enum. Pl. Carib. 22 = pilulifera.
oblongata, *Griseb. Spicil. Fl. Rumel.* i. 136.—Bytant.
oblongata, C. Koch, in Linnæa, xix. (1847) 17 = stricta.
oblongifolia, C. Koch, l. c. xxi. (1848) 726 = amygdaloides.
obovalifolia, *A. Rich. Tent. Fl. Abyss.* ii. 239.—Abyss.
obovata, *Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 241.—Arab.
obscura, Lang, ex Reichb. Fl. Germ. Excurs. 762 = virgata.
obscura, Loisel. in Desv. Journ. Bot. ii. (1809) 332 = falcata.
obtusata, *Pursh, Fl. Am. Sept.* ii. 606.—Am. bor.
obtusifolia, Lam. Encyc. ii. 430 = E. Terracina.
obtusifolia, *Poir. Encyc. Suppl.* ii. 600.—Ins. Teneriff.
obtusifolia, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 66.—Transylv.
occulta, Klotzsch, in Seem. Bot. Voy. Herald, 277 = antisyphilitica.

EUPHORBIA:—
ocellata, *Durand & Hilgard, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 46; *et Pacif. Rail. Rep.* v. 15.—Am. bor.
octogona, *Hort. ex Steud. Nom.* ed. II. i. 613, nomen. —Hab.?
ocymoides, *Linn. Sp. Pl.* 453.—Mexic.
ocymoides, Hook. & Arn. Bot. Beech. Voy. 310 = adenopters.
odontadenia, *Boiss. in DC. Prod.* xv. ii. 148.— N. Mexic.
odontophylla, Boiss. Diagn. Ser. II. iv. 82 = Stocksiana.
odontophylla, Willd. Enum. Hort. Berol. Suppl. 28 = cereiformis.
Oerstediana, *Boiss. in DC. Prod.* xv. ii. 59.—Am. centr.
officinalis, Forsk. Fl. Aegypt. Arab. 94 = officinarum?
officinalis, Jackson, Acc. Maroc. ed. III. 184 = resinifera.
officinarum, Boiss. in DC. Prod. xv. ii. 84 = Beaumieriana.
officinarum, Hochst. ex Boiss. l. c. 96 = Schimperi.
officinarum, *Linn. Sp. Pl.* 451.—Afr. bor.
obtistica, Steud. & Hochst. ex Boiss. in DC. Prod. xv. ii. 142 = communtata.
oleaefolia, Gouan, Herbor. 29 = nicæensis.
oleaefolia, Noé, ex Nym. Consp. 659 = lucida.
oleracea, Pers. Syn. ii. 14 = E. Peplus.
oliganiha, Boiss. in DC. Prod. xv. ii. 19 = pycnostegia.
Oncei, *Fronch. & Sav. Enum. Pl. Jap.* ii. 186.— Japon.
opaca, Lang, ex Reichb. Fl. Germ. Excurs. 762 = virgata
ophthalmica, Pers. Syn. ii. 18 = pilulifera.
oraria, F. Muell. ex Boiss. in DC. Prod. xv. ii. 12 = E. Atoto.
orbiculata, *H. B. & K. Nov. Gen. et Sp.* ii. 52.— N. Granat.
orbicalata, Miq. Fl. Ind. Bat. i. ii. 421 = microphylla.
orcophila, Miq. Anal. Bot. Ind. iii. 17 = Kothiana.
orientalis, Bertol. Fl. Ital. v. 75 = carpatorum.
orientalis, *Linn. Sp. Pl.* 460.—Armen.; Persia.
origanoides, Bert. ex Boiss. in DC. Prod. xv. ii. 21 = Berteriana.
origanoides, *Linn. Sp. Pl.* 453.—Ins. Adscenac.
Orizabae, *Boiss. in DC. Prod.* xv. ii. 147.—Mexic.
ornithopus, *Jacq. Fragm. Bot.* 78. t. 130.—Afr. austr.
Orphanidis, *Boiss. Diagn.* Ser. II. iv. 89.—Græcia.
osyrides, *Boiss. l. c.* Ser. I. vii. 87.—Persia.
ovalifolia, *Engelm. ex Klotzsch & Garcke, in Abh. Akad. Berl.* (1860) 26, in syn.—Chili.
ovalifolia, Kostel. Allg. Med.-Pharm. Fl. v. 1724 = hypericifolia.
ovata, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 167.—Afr. austr.
oxycoccoides, *Boiss. Cent. Euph.* 14.—Madag.
oxyodonta, *Boiss. in DC. Prod.* xv. ii. 1267.—Syria; Mesopot.
oxyphylla, *Boiss. l. c.* 1268.—Hispan.
oxystegia, *Boiss. Cent. Euph.* 21.—Afr. austr.
pachyrhiza, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 745.—Soongaria.
Pallasii, *Turcz. l. c.* xxvii. (1854) i. 358.—Dahuria.
pallens, Dulw. Rev. Hist. Malab. 84 = E. Atoto.
pallida, Willd. Sp. Pl. ii. 923 = salicifolia, lucida.
Palmeri, *Engelm. ex S. Wats. in Bot. Calif.* ii. 75.— Calif.
palustris, *Linn. Sp. Pl.* 462.—Europ.
panacea, Webb & Berth. Phyt. Canar. iii. 247 = E. Terracina.
Pancheri, *Baill. Adansonia*, ii. (1861-62) 211.—N. Caled.
paniculata, Benth. Cat. Pl. Pyrén. 78 = sero-comans.
paniculata, Desf. Fl. Atlant. i. 336 = algeriensis.
paniculata, Ell. Sketch, ii. 660 = corollata.
paniculata, Loisel. Fl. Gall. ed. II. i. 345 = pilosa.
paniculata, Tenore, Prod. Fl. Nap. p. lxvii = platyphylla.
pannonica, Host, Fl. Austr. ii. 566 = glareosa.
papillaris, Jan, Elench. 7 = Bivonæ.
papilligera, Boiss. Cent. Euph. 6 = hypericifolia.
papillosa, *St. Hil. Pl. Ut. Bras.* t. 18.—Reg. Argent.
papillosa, Pousolz, Cat. Pl. Gard. 18 = Duvalii.
paradoxa, Schur, Enum. Pl. Transs. 596 = E. Esula.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 50.—Paraguay.

EUPHORBIA:—
Parsiana, *Linn. Sp. Pl.* 458.—Europ. occ.; Reg. Mediterr.
pavonsquarrata, Blanco, Fl. Filip. ed. II. 386 = serrulata.
parasitica, Pav. ex Boiss. in DC. Prod. xv. ii. 5 = Pedilanthus ramosissimus.
Parryi, *Engelm. in Am. Natural.* ix. (1875) 350.—Am. bor. occ.
parviflora, Lam. Encyc. ii. 494 = roses.
parvifolia, Linn. Syst. ed. X. 1047 = hypericifolia.
parvifolia, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 34.—Afr. austr.
parvimamma, *Boiss. l. c.* 86.—Afr. austr.
parvula, *Delile, Fl. Egypte*, 285. t. 30. f. 4.—Aegypt.
parvula, C. Koch, in Linnaea, xxi. (1848) 761 = arvalis.
patagonica, *Hieron. in Bol. Acad. Nac. Cordova*. iii. (1879) 387.—Patagon.
patens, *Kit. in Linnaea*, xxxii. (1863) 558.—Europ.
patula, Mill. Gard. Dict. ed. VIII. n. 13 = succaulis.
pauciflora, *Dufour, in Bull. Soc. Bot. Fr.* vii. (1860) 442.—Hispan.
pauciflora, *Hill, Veg. Syst.* x. 87 (Quid?).—Ind. or.
pauciflora, *Nutt. ex Seem. Fl. Vit.* 216.—Ins. Sandvic.
paucifolia, Klotzsch, in Lehm. Pl. Preiss. i. 174 = Calycopeplus ephedroides.
pedicellata, *Engelm. in Torr. Bot. Mex. Bound.* 186.—Mexic.
peduncularis, Klotzsch, in Seem. Bot. Voy. Herald, 277 = strigosa.
pedunculosa, A. Rich. Fl. Cub. Faserog. ii. 198 = graminea.
pegmoides, *Boiss. Cent. Euph.* 21.—Mexic.
peltinervis, *Rupr. in Maximow. Prim. Fl. Amur.* 389; *Boiss. l. c.* 81.—China.
peltata, *Roxb. Hort. Beng.* [86]; *Fl. Ind.* ii. 474.—Ind. or.
peltigera, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 91.—Afr. austr.
pendula, *Link, Enum. Hort. Berol.* ii. 10; *Boiss. l. c.* 78.—Hab.?
pentadactyla, *Griseb. in Gentt. Abh.* xxiv. (1879) 63.—Reg. Argent.
pentagona, Blanco, Fl. Filip. ed. I. 418 = neriifolia.
pentagona, *Haw. in Phil. Mag.* (1827) 187.—Afr. austr.
pentagona, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14 = neriifolia.
pentagona, Royle, Illustr. Bot. Himal. 339 = Royleana.
Pentlandi, *Boiss. in DC. Prod.* xv. ii. 69.—Bolivia.
peperomioides, *Boiss. Cent. Euph.* 17.—Bras.
Peplidion, *Engelm. in Torr. Bot. Mex. Bound.* 191.—Texas.
Peplis, *Linn. Sp. Pl.* 455.—Europ. occ.; Reg. Mediterr.
peploides, *Gouan, Fl. Monsp.* 174.—Europ.; Afr. bor.
peploides, Griseb. Spicil. Fl. Rumel. i. 188 = E. Peplus.
peploides, *E. Mey. in* Drège, Zwei Pfl. Docum. 184 = phylloclada.
peploides, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 178 = longicornis.
Peplus, *Linn. Sp. Pl.* 456.—Europ.; An. bor.
pereskiaefolia, *Haullet, ex Baill. Adansonia*, i. (1860–61) 105.—Afr. trop.
perforata, Gmel. Fl. Sic. Prod. i. 540 = prostrata.
Perrottetii, Jaub. & Spach, Illustr. v. 72. t. 464 = cuneata.
personifoliana, Boiss. Diagn. Ser. I. vii. 93 = striatella.
persica, C. A. Mey. ex Boiss. in DC. Prod. xv. ii. 111 = magalanthe.
persica, Stev. ex Boiss. l. c. 160 = virgata.
Pervilleana, *Baill. Adansonia*, i. (1860–61) 141.—Hab.?
Pestalozzae, *Boiss. Diagn.* Ser. I. xii. 114.—Lycia.
petaloides, *Engelm. in Torr. Bot. Mex. Bound.* 185.—Am. bor.
petiolaris, *Sims, Bot. Mag.* t. 583.—Ins. S. Thomas.
petiolata, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 258.—Syria.
Petitiana, A. Rich. Tent. Fl. Abyss. ii. 241 = cerobrina.
petrophila, *C. A. Mey. in Mém. Sc. Pétersb. Sér.* VI. *Sc. Nat.* vii. (1855) 9.—Tauria.
Phillippiana, *Boiss. in DC. Prod.* xv. ii. 117.—Chili.
phillippina, J. Gay, ex Boiss. l. c. 47 = thymifolia.
phosphorea, *Mart. Reise Bras.* ii. 612, 756.—Bras.
phyllanthoides, Boiss. Cent. Euph. 4 = oblongata.
phylloclada, *Boiss. in DC. Prod.* xv. ii. 66.—Afr. austr.

EUPHORBIA:—
phymatoclada, *Boiss. Cent. Euph.* 24.—Afr. austr.
phymatosperma, *Boiss. & Gaill. Diagn.* Ser. II. iv. 88.—Syria.
physalifolia, *Boiss. Cent. Euph.* 19.—Mexic.
physoclada, *Boiss. l. c.* 24.—Madag.
picta, Jacq. Coll. iii. 173 = graminea.
pilosa, Brot. Fl. Lusit. ii. 315 = pubescens.
pilosa, Chaub. Nouv. Fl. Pélop. 81 = oblongata.
pilosa, D. Don, ex Bab. in Engl. Bot. Suppl. t. 2767.—palmaris.
pilosa, *Linn. Sp. Pl.* 460.—Europ.; As. bor.
pilosa, Pursh, Fl. Am. Sept. ii. 607 = Darlingtonii.
pilosa, Vill. Hist. Pl. Dauph. iii. 823 = flavicoma.
pilosula, *Engelm. ex Boiss. in DC. Prod.* xv. ii. 89.—Mexic.
pilulifera, Linn. ex Boiss. in DC. Prod. xv. ii. 20 = parviflora.
pilulifera, *Linn. Sp. Pl.* 454.—Amphig. trop.
pinea, *Linn. Syst. ed.* XII. 883.—Reg. Mediterr.
pinea, Tenôr. Agenn. 65, ex Willk. & Lange, Prod. Fl. Hisp. iii. 500 = portlanzlica.
pinifolia, DC. Fl. Fr. ii. 861 = E. Esula.
pinifolia, Willd. Enum. Hort. Berol. Suppl. 28 = sieppica.
piscatoria, *Ait. Hort. Kew.* ed. I. ii. 137.—Ins. Mader.
piscatoria, Link, in Buch, Beschr. Canar. Ins. 158 = E. Esula.
pistaciifolia, *Boiss. in DC. Prod.* xv. ii. 93.—Afr. austr.
Pithyusa, *Linn. Sp. Pl.* 458.—Reg. Mediterr.
platyphylla, Pritz. in Linnaea, xxxvi. 466, nomen = platyphyllos.
platyleiga, Decne. in Jacquem. Voy. Bot. 156. t. 157 = elegans.
platyphyllos, *Linn. Sp. Pl.* 460.—Europ.; Afr. bor.
platyphyllos, *Engelm. ex S. Wats. in Bot. Calif.* ii. 482.—Calif.
plebeia, *Boiss. Diagn.* Ser. I. vii. 98.—Persia austr.
plumerioides, *Trijtm. ex Hassk. Hort. Bog* i. 20.—Java.
podadenia, *Boiss. in DC. Prod.* xv. ii. 40.—Mexic.
Poeppigii, *Boiss. l. c.* 56.—Peruv.
Poinettiana, Brist, ex R. Grah. in Edinb. N. Phil. Journ. (1836) 414 = pulcherrima.
polyacantha, *Boiss. Cent. Euph.* 25.—Abyssin.
polycarpa, *Benth. Bot. Voy. Sulph.* 50.—Calif.
polycarpa, Boiss. & Hohen. Diagn. Ser. I. xii. 112 = decipiens.
polychroma, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 309 = epithymoides.
polycnemoides, *Boiss. Cent. Euph.* 10.—Texas.
polycnemoides, *Hochst. ex Boiss. in DC. Prod.* xv. ii. 46.—Abyss.
polygaloides, *Boiss. & Reut. in Boiss. Cent. Euph.* 84.—Hispan.
polygona, *How. Misc.* 134.—Afr. austr.
polygona, Lodd. Bot. Cab. t. 1244 = cereiformis.
polygonifolia, Forsk. Fl. Aegypt. Arab. p. xxiii = E. Peplis.
polygonifolia, Hook. Fl. Bor. Am. ii. 140 = glyptosperma.
polygonifolia, Jacq. Coll. v. t. 18. f. 3 = maculata.
polygonifolia, Link, in Buch, Beschr. Canar. Ins. 158 = mauritanica.
polygonifolia, *Linn. Sp. Pl.* 455.—Am. bor.
polygonifolia, Michx. ex Boiss. in DC. Prod. xv. ii. 101 = E. Ipecacuanha.
polygonifolia, Michx. Fl. Bor. Am. ii. 211 = Curtisii.
polygonisperma, Uren. & Godr. Fl. Fr. iii. 75 = humifusa.
pomifera, Thunb. Prod. Pl. Cap. 86 = meloformis.
portlandica, *Linn. Sp. Pl.* 458.—Europ. occ.
portlandifolia, Salis, in Flora, xvii. (1834) ii. Beibl. 5 = pinea.
portlandica, Sibth. & Sm. Fl. Graec. Prod. i. 297 = E. Terracina.
portulacoides, Linn. Herb. ex Boiss. in DC. Prod. xv. ii. 109 = E. Ipecacuanha, heliscopia.
portulacoides, *Linn. Sp. Pl.* 456, partim; *Spreng. Syst.* iii. 792.—Chili.
Postii, *Boiss. Fl. Orient.* iv. 1089.—Syria.
potentillaefolia, *Boiss. Cent. Euph.* 8.—Bras.
praecox, Fisch. Cat. Hort. Gorenk. (1812) 68 = leptocaula.
pseudoasia, *Greenway, ex Trimen. in Ast. Hort. Petrop.* ix. (1884) 168, nomen.—Rom.
Pseudofalcata, *Fl. Stc. Prod.* i. 539.—Ann. bor.
Pseudofalcata, Spenn. ex Nym. Consp. 664 = graeca.
Pseudofalcata, Baill. Adansonia, i. (1860–61) 61 = convolvuloides.

EUPHORBIA:—
pseudofalcata, *Baker, in Journ. Linn. Soc.* xviii. (1881) 276.—Madag.
procera, Bieb. Fl. Taur. Cauc. i. 378 = pilosa.
procumbens, DC. Cat. Hort. Monsp. 111 = pilulifera.
procumbens, Hort. ex Boiss. in DC. Prod. xv. ii. 43 = ovalifolia.
procumbens, Meerburg, Fl. Rar. t. 55 = stellata.
procumbens, *Mill. Gard. Dict.* ed. VIII. n. 19.—Afr. austr.
prolifera, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 62.—Reg. Himal.
prolifera, Ehrenb. ex Boiss. in DC. Prod. xv. ii. 16 = obovata.
propinqua, *R. Br. in Salt, Abyss. App.* p. lxv, nomen.—Abyss.
prostrata, *Ait. Hort. Kew.* ed. I. ii. 139.—Am. trop.
prostrata, Burchell, ex Hensl. Bot. Challeng. i. ii. 82 = E. Chamaesyce.
prostrata, J. Grah. Cat. Pl. Bomb. 179 = thymifolia.
prostrifolia, *Boiss. in DC. Prod.* xv. ii. 92.—Afr. austr.
provincialis, Noé, ex Nym. Consp. 663 = glareosa.
provincialis, Willd. Sp. Pl. ii. 914 = E. Terracina.
prunifolia, Jacq. Hort. Schoenb. iii. 18 = geniculata.
Pseudochamaesyce, Fisch. & Mey. Ind. Sem. Hort. Petrop. ix. 78 = humifusa.
Pseudocyparissias, Jord. in Billot, Annot. 26 = E. Esula.
Pseudo-Esula, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 66 = E. Cyparissias.
Pseudo-lucida, Schur, l. c. iii. (1852) 124 = lucida.
pterococca, *Boiss. Fl. Orient.* iii. 812.—Reg. Mediterr.
pubentissima, *Michx. Fl. Bor. Am.* ii. 212.—Am. bor.
pubera, Blume, Bijdr. 588 = indica.
pubescens, *Vahl, Symb. Bot.* ii. 86.—Reg. Mediterr.
pubigera, Friwald. ex Boiss. in DC. Prod. xv. ii. 126 = E. Apios.
pugniformis, Boiss. l. c. 92 = procumbens.
pulchella, Brouss. in DC. Cat. Hort. Monsp. 111 = imbricata.
pulchella, Hort. ex Poir. Encyc. Suppl. ii. 615 = nummulariaefolia.
pulchella, H. B. & K. Nov. Gen. et Sp. ii. 56 = brasiliensis.
pulchella, *Lap. & Rodr. in Anal. Clenc. Nat.* v. (1802) 71.—Hab.?
pulcherrima, *Willd. ex Klotzsch, in Otto & Dietr. Allg. Gartenz.* ii. (1834) 97 ... Mexic.
pulverulenta, Kit. ex Reichb. Fl. Germ. Excurs. 762 = glareosa.
pumila, Pall. Ind. Taur. ex Ledeb. Fl. Ross. iii. 573 = nicaeensis.
pumila, Sibth. & Sm. Fl. Graec. Prod. i. 324 = hemlariaefolia.
punctata, *Delile, Fl. Egypte*, 324. t. 30. f. 3.—Afr. bor.
punctata, *Krock. Fl. Siles.* ii. i. 97.—Europ.
punctulata, *Anderss. in Vet. Akad. Handl. Stockh.* 1858 (1855) 285.—Ins. Galapag.
pungens, Lam. Encyc. ii. 481 = spinosa.
pungens, E. Mey. in Drège, Zwei Pfl. Docum. 184 = striata.
pungens, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 268.—Cilicia.
punicea, *Sw. Prod. Veg. Ind. Occ.* 76.—Jamaic.
purpurascens, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 690 = E. Terracina.
purpurascens, *Schousb. in Hornem. Hort. Hafn.* 508.—Marocc.
purpurascens, Schum. & Thonn. Beskr. Guin. Pl. 258 = indica.
purpurata, Thunll. Fl. Par. ed. II. 325 = dulcis.
pusilla, *Lag. Gen. et Sp. Nov.* 17.—Hispan.
pycnantha, *Engelm. in Torr. Bot. Mex. Bound.* 188.—Mexic.
pycnophylla, C. Koch, in Linnaea, xix. (1847) 17 = hemlariaefolia.
pycnotegia, *Boiss. Cent. Euph.* 9.—Ind. or.
pygmaea, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 171, nomen; *Boiss. in DC. Prod.* xv. ii. 99.—Persia, Scamandria.
pygmaea, Ledeb. Fl. Ross. iii. 569 = Ledebourii.
pygmaea, Phil. in Linnaea, xxix. (1857–58) 41 = Philippiana.
pyrenaica, Jord. Obs. Fl. Crit. iii. (1846) 237 = E. Chamaebuxus.
pyrifolia, *Lam. Encyc.* ii. 479.—Ins. Mauric.
Pythiusa, Friwald. ex Nym. Consp. 664 = dulcis.
quinoa, *Boiss. in DC. Prod.* xv. ii. 42.—Ecuador.

EUPHORBIA :—
racemosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 75.—Afr. austr.
racemosa, Tausch, ex Reichb. Fl. Germ. Excurs. 701 = E. Esula.
radians, *Benth. Pl. Hartw.* 8.—Mexic.
radiata, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 90.—Afr. austr.
radiata, Thunb. Prod. Pl. Cap. 86 = stellata.
radicans, Moric. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 24 = serpens.
radioloides, *Boiss. in DC. Prod.* xv. ii. 45.—Mexic.
ragusana, Keichb. Fl. Germ Excurs. 873 = pinea.
ramosissima, Boiss. in DC. Prod. xv. ii. 14 = Sparmanni.
ramosissima, *Hook. & Arn. in Bot. Beech. Voy.* 69.—Ins. Pacif.
ramosissima, Loisel. in Mém. Soc Linn. Par. vi (1827) 415 = E. Terracina.
rapulum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 448.—Soongaria.
Rattani, *S. Wats. in Proc. Am. Acad.* xx. (1885) 372. —Calif.
recta, Klotzsch, in Seem. Bot. Voy. Herald, 277 = calletioides.
recurva, *Hook. f. in Trans. Linn. Soc.* xx (1847) 182. —Ins. Galapag.
reflexa, *Spreng. Syst.* iii. 802.—Hab.?
reflexta, Lowe, in Trans. Camb. Phil. Soc., vi. (1838) reimp. 11 = Preslii.
Regis-Jubae, *Webb & Berth. Phyt. Canar.* iii. 250.— Ins. Teneriff.
Reichenbachiana, Willk. in Flora, xxxv. (1852) 310 = pilosa.
Reinwardtiana, Steud. Nom. ed II. i. 614 = serrulata.
Remyi, *A. Gray, ex Boiss. in DC. Prod.* xv. ii. 1262.— Ins. Sandvic.
reniformis, *Blume, Bijdr.* 684.—Java.
repanda, *Sweet, Hort. Brit.* ed. I. 356.—Ind. or.
repens, *C. Koch, in Linnaea*, xxi. (1848) 728.— Oriens.
repetita, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 240.— Abyss.
resinifera, *Berg, in Berg & C. F. Schmidt, Offic. Gew.* t. 84 d verso.—Marocc.
resinosa, *Benth. Bot. Voy. Sulph.* 162.—Mexic.
retusa, Bieb. Fl. Taur. Cauc. i. 371 = exigua.
retusa, Cav. Ic. i. 21 = sulcata.
retusa, Forsk. Fl. Aegypt. Arab. 93 = cornuta.
Resteriana, *Boiss. Diagn.* Ser. I. xii. 115.—Syria.
revoluta, *Engelm. in Torr. Bot. Mex. Bound.* 186 — Texas.
rhadodes, *Boiss. Cent. Euph.* 30.—Bras.
rhipsaloides, Lem. Illustr. Hortic. xv. (1857) Misc. 72 = E. Tirucalli.
rhombea, Willd. ex Link, in Buch, Beschr. Canar. Ins. 158 = E. Terracina.
rhombifolia, *Boiss. Cent. Euph.* 19.—Afr. austr.
rhytidosperma, *Boiss. & Bal. Diagn.* Ser. II. iv. 84.— Cilicia.
rhytisperma, *Engelm. ex Klotzsch & Garcke, in Abh. Akad. Berl.* (1860) 34, in syn.—Chili.
Richardiana, Baill. Adansonia, i. (1860-61) 105 = abyssinica.
Riedeliana, *Boiss. Cent. Euph.* 30.—Bras.
rigens, Sweet, Hort. Brit. ed. II. 455 = spinosa.
rigida, Bieb. Fl. Taur. Cauc. i. 375 = biglandulosa.
rigida, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 414 = corsica.
rigidula, Steud. Nom. ed. II. i. 614 = spinosa.
rimarum, *Cass. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 261.—Marocc.
riparia, Jord. in Billot, Annot. 21 = E. Esula.
Rochebrunl, *Franch. & Sav. Enum. Pl. Jap.* ii. 485.— Japon.
Rosmeriana, *Scheele, in Linnaea*, xxii. (1849) 151.— Texas.
Rosseri, Orph. ex Boiss. in DC. Prod. xv. ii. 173 = Heldreichii.
rosea, *Retz. Obs.* iv. 26.—Ind. or.
rosea, Roth, in Ges. Naturf. Fr. Neue Schr. iv. (1803) 185 = glanconcnsis.
rosea, Roxb. in Beatson, St. Hel. Tracts, 308 = E. Chinensoyon.
Rothiana, Miq. Fl. Ind. Bat. i. ii. 619 = javanica.
Rothiana, *Sprang. Syst.* iii. 796.—Ind. or.

EUPHORBIA :—
Rothiana, Wight, Ic. t. 1864 = oreophila.
rotundata, Hochst. in Lorent, Wanderungen, 344 = denticulata.
rotundifolia, *Hook. & Arn. Bot. Beech. Voy.* 44.— Chili.
rotundifolia, Loisel. in Desv. Journ. Bot. ii. (1809) 331 = peploides.
rotundifolia, Phil. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 26 = ovalifolia.
Royleana, *Boiss. in DC. Prod.* xv. ii. 83.—Reg. Himal
rubescens, Link, in Buch, Beschr. Canar. Ins. 158 = E. Peplis.
rubicunda, Blume, Cat. Gew. Buitenz. 73 = thymifolia.
rubra, Cav. Ic. i. 21. t. 34. f. 2 = falcata.
rubra, DC. Fl. Fr. vi. 359 = exigua.
rubra, *Pall. Tabl. Taur.* 51.—*Cf. Ledeb. Fl. Ross.* iii. 580 (sp. dub.).—Tauria.
rubricalis, Kegel, Ind. Sem. Hort. Petrop. (1856) 37 = rupicola.
ruderalis, *Dum. Fl. Belg.* 87 (Quid?).—Europ.
ruderalis, Scheele, in Linnaea, xvii. (1843) 343 = arvalis.
Ruegeliana, Shuttlew. ex Boiss. in DC. Prod. xv. ii. 67 = Curtisii.
rugosa, *Kit. in Rochel, Pl. Rar. Banat.* 26, nomen.— Banat.
Rumiana, *Boiss. in DC. Prod.* xv. ii. 31.—Peruv.
ramiciformis, *Boiss Cent. Euph.* 39.—As. Min.
rupestris, Rudolphi ex Boiss. in DC. Prod. xv. ii. 105 = Barrelieri.
rupestris, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 190.— Sibir.
rupicola, *Boiss. Elench.* 81.—Reg. Mediterr. occ.
rupicola, Scheele, in Linnaea, xxii. (1849) 153 = Fendleri.
ruscinonensis, *Boiss. Cent. Euph.* 33.—Mont. Pyren.
sabulicola, *Boiss. l. c.* 6.—Bras
saccharata, *Boiss. l. c.* 18.—Mexic.
Sagraeana, A. Rich. ex Boiss. in DC. Prod. xv. ii. 105 = trichotoma.
salicetorum, Jord. Pugill. Pl. Nov. 138 = androsaemifolia.
salicifolia, DC. Fl. Fr. vi. 363 = E. Esula.
salicifolia, *Host, Syn. Fl. Austr.* 267.—Europ.
salina, Willd. ex Boiss. in DC. Prod. xv. ii. 15 = buxifolia.
Sampsoni, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 240 = peknensis.
sanguinea, *Hochst. & Steud. ex Boiss. in DC. Prod.* xv. ii. 35.—Afr trop.
sanguinea, Hort. Berol. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 39 = haematoides.
Sarati, *Ardoino, Fl. Alp. Marit.* 335.—Europ. austr.
sarawschanica, *Regel, Pl. Nov. Fedsch.* 78.—Turkestan.
sarcodes, *Boiss. Cent. Euph.* 29.—Bras
sarpiana, *Becker, in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 18; *Boiss. in DC. Prod.* xv. ii. 159.— Rossia.
satureioides, Lem. Encyc. ii. 624 = pulcra.
Sauliana, *Bor. ex Boiss. in DC. Prod.* xv. ii. 1266. —Lydia.
saxatilis, Bieb. Fl. Taur. Cauc. iii. 919 = petrophila.
saxatilis, *Georgi, Beschr. Russ. Reich.* III. iv. 995 (sp. dub.).—Rossia.
saxatilis, *Jacq. Fl. Austr.* iv. 28. t. 345.—Austria.
saxatilis, Loisel. Fl. Gall. ed. II. i. 342 = Gerardiana.
scandens, *H. B. & K. Nov. Gen. et Sp.* ii. 58.— Mexic.
scabrella, *Boiss. in DC. Prod.* xv. ii. 55.—Mexic.
scabrifolia, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 247.—Burma.
scopiformis, Buek, in DC. Prod. Ind. iv. 148, sphalm. = scopiformis.
Schickendantzii, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 17.—Reg. Argent.
Schimperi, *Presl, Bot. Bemerk.* 109.—Arabia.
Schimperiana, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 242.—Abyss.
schinoedenia, Boiss. & Hohen. ex Boiss. Diagn. Ser. I. xii. 118 = cheiradenia.
schizocoras, Boiss. l. c. v. 58 = tinctoria.
schizoleuka, *F. Muell. ex Boiss. in DC. Prod.* xv. ii. 20.—Australi.

EUPHORBIA :—
schizoloba, Engelm. in Proc. Am. Acad. v. (1862) 173 = ineisa.
Schlechtendalii, *Boiss. Cent. Euph.* 18.—Mexic.
Schottiana, *Boiss. Diagn.* Ser. II. iv. 85.—Cilicia.
Schultzii, *Benth. Fl. Austral.* vi. 47.—Austral.
Schweinfurthii, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 410.—Ins. Socotr.
sciadophila, *Boiss. in DC. Prod.* xv. ii. 57.—Bras.
sclerophylla, *Boiss. Cent. Euph.* 37.—Afr. austr.
scolopendrina, Donn, Cat. Hort. Cantab. ed. III. 88 = uncinata.
scopiformis, *Boiss. in DC. Prod.* xv. ii. 1269.—Afr. austr.
Scopoliana, *Steud. Nom.* ed. II. i. 615.—Cilicia.
scordifolia, *Jacq. Coll.* v. 113.—Afr. trop.
Scotanum, *Schlecht. in Linnaea*, xix. (1847) 250.— Mexic.
Seemanni, Klotzsch, in Seem. Bot. Voy. Herald, 276 = campestris.
segetalis, J. Grah. Cat. Pl. Bomb. 179 = Rothiana?
segetalis, *Linn. Sp. Pl.* 458.—Europ.
segetalis, Pall. Reise, i. 164 = E. Esula.
segetalis, Pall. Tabl. Taur. 51 = agraria.
segetalis, Raul. Crète, 549 = E. Terracina.
segovienais, *Boiss. in DC. Prod.* xv. ii. 58.—Mexic.
Seguieri, Nym. Conap. 655 = nicaeensis.
Seguieriana, Neck. in Act. Akad. Theod. Pal. ii (1770) 498 = Gerardiana.
Seguierii, All. Fl. Pedem. i. 288 = prsec.
Sellos, Boss. Cent. Euph. 30 = clodes.
Selloi, *Klotzsch, ex Boiss. in DC. Prod.* xv. ii. 50.— Bras.
semiperforlata, *Vis. Fl. Cors Sp. Non.* 7.—Sardin.
sens-comans, *Bubani, in Nuov. Ann. Sc. Nat. Bolog.* ix. (1848) 95.—Mont. Pyren.
serotina, Host, Fl. Austr. ii. 562 = nicaeensis.
serpens, Baill. Adansonia, ii. (1861-62) 311 = neocaledonica.
serpens, Bub ex Boiss. Cent. Euph. 12 = Balbisii.
serpens, *H. B. & K. Nov. Gen. et Sp.* ii. 52.—Amer. bor.
serpyrrhiza, Boiss. in DC. Prod. xv. ii. 75, sphalm. = scopiformis.
serpyllifolia, Baib ex Boiss. l. c. 49 = adenopters.
serpyllifolia, *Boj. Hort. Maurit.* 288.—Ins. Galega.
serpyllifolia, *Pers. Syn.* ii. 14.—Amb. bor.
serrata, S. G. Gmel. It. ii. 257 = pilosa ?
serrata, *Linn. Sp. Pl.* 459.—Lusitania.
serrula, *Engelm. in Torr. Bot. Mex. Bound.* 188.— N. Mexic.
serrulata, *Renew. ex Blume, Bijdr.* 685.—Malaya; Chuna.
serrulata, Thuill. Fl. Par. ed. II. 287 = stricta.
serrulata, Vell. Fl. Flum. 208; v. t. 17 = brasiliensis.
sessiliflora, Boiss. & Sprun. Diagn. Ser. I. v. 52 = oblongata.
sessiliflora, E. Mey. in Drège, Zwei Pfl. Docum. 184 = gummifera.
sessiliflora, *Roxb. Hort. Beng.* [38]; *Fl. Ind.* ii. 471.— Burma.
sessilifolia, *Klotzsch, ex Boiss. in DC. Prod.* xv. ii. 64. —Bras.
setiformis, Poir. Voy. Barb. ii. 178 = E. Terracina.
setigera, E. Mey. in Drège, Zwei Pfl. Docum. 184 = sanguinea.
setiloba, *Engelm. ex Torr. in Pacif. Rail. Rep.* v. 364. —Calif.
setosa, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 672.— Bras.
sexangularis, Henckel, Adumbr. 7 = glauca.
sharkottensis, Baill. Adansonia, vi. (1866) 287 = myrtoides.
sibirica, Fisch. ex Boiss. in DC. Prod. xv. ii. 123 = alpina.
Sibthorpii, *Boiss. Cent. Euph.* 39.—Graecia.
Sieboldiana, *Morr. & Decne. in Bull. Acad. Brux.* iii. (1836) 174.—Japon.
silkimensis, *Boiss. in DC. Prod.* xv. ii. 113.—Reg. Himal.
silenifolia, Sweet, Hort. Brit. ed. II. 454 = elliptica.
simplex, C. Koch, in Linnaea, xxi. (1848) 730 = segetalis.
sinuata, Hochst. ex Boiss. in DC. Prod. xv. ii. 142 = E. Chamaepeplus.
Sinclairiana, *Benth. Bot. Sulph. Voy.* 163.—Ins. Gorgon.
socialis, Zoll. in Nat. en Genesk. Arch. Nederl. Indie, i. (1844) 484 = javanica.

EUPHORBIA:—
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 93.—Ins. Socotr.
solisfera, Vis. ex Boiss. in DC. Prod. xv. ii. 138 = capitulata.
soltsaqua, Reichb. Fl. Germ. Excurs. 756 = dulcis.
somagarica, *Boiss. Centr. Euph.* 32.—Sibir.
sordida, Klotzsch, ex Boiss. in DC. Prod. xv. ii. 21 = Berteriana.
sororia, *Schrenk*, in *Bull. Phys.-Math. Acad. Péterab.* iii. (1845) 308.—Persia.
Sparrmanni, *Boiss. Cent. Euph.* 5.—Austral.
spartioides, *Hornem. ex Schrank*, in *Denkschr. Bot. Ges. Regensb.* ii. (1822) 48.—Hab. ?
spartioides, Jacq. Hort. Schoenb. iv. 44 = Cupani.
spathulaefolia, *G. Don*, in *Sweet, Hort. Brit* ed. III. 599.—Ind. or.
spathulata, *Lam. Encyc.* ii. 428.—Bras.
sphaerococca, Salam. ex Cos. Notes Crit. 45 = akenocarpa.
sphaerorhiza, *Benth. Pl. Hartw.* 8.—Mexic.
sphaerosperma, *Shuttlew. ex Boiss. in DC. Prod.* xv. ii. 102.—Florida.
spicata, *E. Mey. in Drège, Zwei Pfl. Docum.* 184, nomen; *Boiss. in DC. Prod.* xv. ii. 97.—Afr. austr.
spinosa, *Linn. Sp. Pl.* 457.—Europ. austr.
spinosa, Sibth. & Sm. Fl. Graec. v. 49. t. 463 = acanthothamnos.
spinosa, Vis. Fl. Lib. Spec. 36 = Bivonae.
spiralis, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 410.—Ins. Socotr.
splendens, *Boj. ex Hook. in Bot Mag.* t. 2902.—Madag.
spongiosa, Ledeb. ex Schrank, in Syll. Ratisb. i. (1824) 214 = E. Lathyris.
Spruceana, *Boiss. in DC. Prod.* xv. ii. 53.—Peruv.
squamigera, Loisel. Fl. Gall. ed. II. i. 345 = rupicola.
squamosa, Willd. Sp. Pl. ii. 918 = nereis.
squarrosa, *Haw. in Phil. Mag.* (1826) 276.—Afr. austr.
stapelioides, *Boiss. Cent. Euph.* 26.—Afr. austr.
stellaespina, *Haw. in Phil. Mag.* (1826) 275.—Afr. austr.
stellata, *Willd. Sp. Pl.* ii. 886.—Afr. austr.
stellulata, Salsm. ex Loisel. in Mém. Soc. Linn. Par. vi. (1827) 415 = cuneifolia.
stellulata, Salzm. in Flora, iv. (1821) 110 = pterococca.
stenophylla, *Boiss. in DC. Prod.* xv. ii. 104.—Bras.
stenophylla, Schur, Enum. Pl. Transs. 595 = E. Cyparissias.
stictospora, *Engelm. in Torr. Bot. Mex. Bound.* 187. —N. Mexic.
stipulacea, Engelm. ex Boiss. in DC. Prod. xv. ii. 30 = alboncarpiata.
Stocksiana, *Boiss. Centr. Euph.* 34.—Beluchist.
Strachеyi, *Boiss in DC. Prod.* xv. ii. 114.—Reg. Himal.
striata, Boiss. Diagn. Ser. I. vii. 91 = striatella.
striata, Eckl. & Zeyh. ex Boiss. in DC. Prod. xv. ii. 169 = E. Erythrina.
striata, *Thunb. Prod. Pl. Cap.* 86.—Afr. austr.
striatella, *Boiss. in DC. Prod.* xv. ii. 152.—Persia.
stricta, *Linn. Syst.* ed. X. 1048.—Europ. med.
stricta, Tenore, Ad. Fl. Neap. Syll. App. iv. 18 = platyphyllos.
strigosa, *Hook. & Arn. Bot. Beech. Voy.* 310.—Mexic.
strobilifera, Dels. in Hook. Kew Journ. iii. (1851) 229 = elegans.
stygiana, *H. C. Wats. in Hook. Lond. Journ. Bot.* iii. (1844) 605.—Ins. Azor.
subamplexicaulis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 744.—Soongaria.
subcilidata, Pers. Syn. ii. 18 = platyphylla.
subcordata, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 184.—Soongaria.
subcordata, Schur, Enum. Pl. Transs. 594 = pilosa.
subhastata, Vis. & Pant. in Mem. Ist. Venet. x. (1861) 444 = agraria.
subserrata, Engelm. ex Boiss. in DC. Prod. xv. ii. 43 = serrulifolia.
subtuberculata, *C. A. Mey. ex Boiss. l. c.* 118.—Persia.
subumbellata, Stend. ex Boiss. l. c. 108 = Coplapina.
suffruticosa, *Forsk. Fl. Aegypt. Arab.* p. xxvi.—Ins. Rhodos.
suffruticulosa, Lecoq & Lamotte, Cat. Pl. Vasc. Fr. 237 = flavicoma.
sulcata, *De Lens, ex Loisel. Fl. Gall.* ed. II. i. 339.—Algar.
sulcata, Lam. ex Boiss. in DC. Prod. xv. ii. 176 = peroniaefolia.
sumbawensis, *Boiss. l. c.* 36.—Ins. Sumbawa.
supina, Rafin. ex Boiss. l. c. 47 = maculata.

EUPHORBIA:—
sylvatica, Linn. Sp. Pl. 468; Jacq. Fl. Austr. iv. 39. t. 378 = amygdaloides.
Synadenia, Baill. Adansonia, iii. (1862-63) 142 = Synadenium arborescens.
syriaca, Spreng. Syst. iii. 792 = lanata.
syrmatesia, *Kit. ex Boiss. in DC. Prod.* xv. ii. 170, nomen.—Banat.
syspirensis, C. Koch, in Linnaea, xxi. (1848) 727 = tinctoria.
syrtyla, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) ii. 1218.—Arab.
Szovitsii, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 27. —As. Min. à Pers.
taitensis, *Boiss. Cent. Euph.* 5.—Ins. Taiti.
talyschensis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 196 = aspera.
tamanduana, *Boiss. Cent. Euph.* 4.—Bras.
tanaicensis, Guss. Fl. Sic. Prod. i. 547 = Cupani.
tannensis, Hort. ex Boiss. in DC. Prod. xv. ii. 138 = platyphylla.
tannensis, *Spreng. Fl. Hal. Mant* 42.—Ins. Tanna.
tatarica, Jacquem. ex Boiss. in DC. Prod. xv. ii. 114 = falcata.
taurica, Bess. ex Link, Enum. Hort. Berol. ii. 14 = segetalis.
taurinensis, *All. Fl. Pedem.* i. 287.—Europ. austr.
taxifolia, *Lam. f. Fl. Cap. Prod.* 14.—Afr. austr.
teheranica, *Boiss. Cent. Euph.* 36.—Persia.
telephioides, *Chapm. Fl. S. U. St.* 402.—Florida.
tenella, H. & K. Nov. Gen. et Sp. ii. 53 = prostrata.
tenera, *S. Wats. in Proc. Am. Acad.* xviii. (1882-83) 150.—Mexic.
tenuifolia, Bieb. Fl. Taur Cauc. i. 379 = leptocaula.
tenuifolia, *Lam. Encyc.* ii. 428.—Gallia.
tenuis, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 62.—Nepal.
Terracina, Griseb. Spicil. Fl. Rumel. i. 139 = graeca.
Terracina, Lag. Elench. Hort. Matrit. (1816) 7 = Lagascae.
Terracina, *Linn. Sp. Pl.* ed. II. 654.—Reg. Mediterr.
Terracina, Reichb. Fl. Germ. Excurs. 759 = taurinensis.
tetracreas, Lange, in Kjoeb. Vidensk. Meddel. (1865) 127 = portlandica.
Tetracreas, Browits, ex Boiss. in DC. Prod. xv. ii. 166 = tinctoria.
tetragona, *Baker, in Saund. Refug. Bot.* t. 80.—Afr. austr.
tetragona, *Haw. in Phil. Mag.* (1826) 276.—Afr. austr.
tetragona, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 238 = polycantha.
tetragona, *Engelm. in Torr. Bot. Mex. Bound.* 191.— Texas.
tetrapteros, *Baker, in Journ. Linn. Soc.* xxi. (1885) 440.—Madag.
tettensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 94; *Boiss. in DC. Prod.* xv. ii. 49.—Afr. trop.
texana, *Boiss Cent. Euph.* 30.—Texas.
thamnoides, *Boiss.* l. c. 38.—As. Min.; Syria.
thelesperma, Hochst. ex Boiss. in DC. Prod. xv. ii. 98 = acalyphoides.
Thi, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 660.—Afr. trop.
thinophila, *Phil. in Anal. Univ. Chil.* (1873) 587.— Chili.
Thomasiana, *Boiss. in DC. Prod.* xv. ii. 113.—Reg. Himal.
Thuarsiana, *Baill. Adansonia*, i. (1860-61) 139.— Madag.
thymifolia, Forsk. Fl. Aegypt. Arab. 94 = scordifolia.
thymifolia, *Linn. Sp. Pl.* 454; *Burm. f. Fl. Ind.* 112. —Amphig. trop.
thymifolia, Pursh, Fl. Am. Sept. ii. 606 = maculata.
thymifolia, Thunb. Fl. Jap. 196 = humifusa.
thymifolia, Wall. Cat. n. 7710 D = rosea.
thymifolia, Wall. l. c. E = granulata.
thymelaea, Griseb. Spicil. Fl. Rumel. i. 143 = agraria.
thymoides, *Boiss. in DC. Prod.* xv. ii. 164.—Reg. Himal.
tinctoria, *Boiss. l. c.* 114.—Reg. Himal.
tigridis, Boiss. l. c. 84 = turcomanica.
tinctoria, *Boiss. & Huet, ex Boiss. in DC. Prod.* xv. ii. 166.—Persia.
Tirucalli, Forsk. Fl. Aegypt. Arab. p. cxii = Schimperi.
Tirucalli, *Linn. Sp. Pl.* 452.—Afr. or.; Ind. or.
Tirucalli, Thunb. Fl. Cap. 406 = Burmanni.
tithymaloides, Linn. Sp. Pl. 466 = Pedilanthus tithymaloides.

EUPHORBIA:—
tomentella, *Engelm. ex Boiss. in DC. Prod.* xv. ii. 32. —Mexic.
tomentella, *Zipp. ex Span. in Linnaea*, xv. (1841) 350, nomen.—Ins. Timor.
tomentosa, Pers. Syn. ii. 18 = scordifolia.
Tommasiniana, Bertol. Fl. Ital. v. 78 = virgata.
torrida, *DC. in Atti Reun. Second. Sc. Ital. Tor.* (1841) 158.—Mexic.
tortilis, *Retzl. ex Wight, Ic.* t. 898.—Ind. or.
tovarensis, *Boiss. Cent. Euph.* 18.—Venezuela.
toxicaria, Afzel. ex Steud. Nom. ed. II. i. 615 = drupifera.
toxicaria, *Neuwette, ex Link, Enum. Hort. Berol.* ii. 10, nomen.—Hab. ?
trachyphylla, A. Rich Fl. Cub. Fanerog. ii. 198 = heterophylla.
trachysperma, *Engelm. in Torr. Bot. Mex. Bound.* 189.—Mexic.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 124 = agraria.
transtagana, *Boiss. Diagn.* Ser. II. iv. 88.—Lusitan
trapezoidalis, *Boiss. in DC. Prod.* xv. ii. 1268.—Afr. trop.
trapezoidalis, Viv. Fl. Lib. Spec. 25. t. 4. f. 1 = E. Terracina.
triacantha, *Ehrenb. ex Boiss. in DC. Prod.* xv. ii. 85. —Abyss.
tricaulcata, *Forsk. Fl. Aegypt. Arab.* 94.—Arab.
triangularis, *Desf. Cat. Hort. Par.* ed III. 389.—Hab ? *tribuloides*, Lam. Encyc. ii. 415 = canariensis.
trichogona, Bertol. Misc. Bot. iii. 20. t. 6. f. 3 = prostrata.
trichophylla, *Baker*, in *Journ Linn. Soc.* xx. (1883) 250.—Madag.
trichotoma, *H. B. & K. Nov. Gen. et Sp.* ii. 60.— Florida; Cuba.
tricuspidata, Lapeyr. Hist. Abr. Pl. Pyr. 271 = exigua
tridentata, Lam. Encyc. ii. 416 = anacantha.
triflora, *Schott, Nyman, & Kotschy, Analect. Bot.* 68 —Dalmatia.
trigona, *Haw. Syn. Pl. Succ.* 127.—Ind. or ; Malaya.
trigona, Mill. Gard. Dict. ed. VIII. n. 8 = antiquorum.
trigona, *Roxb. Hort. Beng.* [36]; *Fl. Ind.* ii. 468.— Ins. Moluce.
trigonocarpa, Fisch. Ind. Sem. Hort. Petrop. vii. 48 = pilosa.
trinervia, *Boiss. Elench.* 32 = boetica.
trinervia, *Schum. & Thonn. Beskr. Pl. Guin.* 253.— Afr. trop.
trinervis, Bertol. Fl. Ital. v. 37 = Preslii.
triphylla, Willd. ex Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 249 = scabrella.
tristis, Bess. Ind. Hort. Crem. (1811) Suppl. iv. 27 = E. Esula.
Triumfetti, Bertol. Fl. Ital. v. 35 = E. Esula.
tuberculata, *Jacq. Hort. Schoenb.* ii. 43. t. 208.—Afr. austr.
tuberosa, *Linn. Sp. Pl.* 456.—Afr. austr.
Tuckeyana, *Steud. Nom.* ed. II. i. 615.—Ins. Cap. Verd.
turcomanica, *Boiss. Cent. Euph.* 13.—Turcoman.
Turczaninowii, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 447.—Turkest.
turkestanica, *Franch. in Ann. Sc. Nat. Sér.* VI. xviii. (1884) 248.—Turkestan.
turkestanica, *Regel, Pl. Nov. Fedsch.* 78.—Turkest.
Turpini, *Boiss. Cent. Euph.* 12.—Ins. N. Doming.
ucranica, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 168, nomen.—Ucrania.
uliginosa, Lange, in Kjoeb. Videnak. Meddel. (1865) 126 = flavicoma.
uliginosa, *Welw. ex Boiss. in DC. Prod.* xv. ii. 127.— Lusitan.
umbellulata, *Engelm. x Boiss. l. c.* 40.—Mexic.
umbrosa, *Bert. ex Spreng. Syst.* iii. 791.—Ins. S. Doming.
uncinata, *DC. Pl. Grass.* t. 151.—Afr. austr.
undulata, *Bieb. Fl. Taur. Cauc.* i. 371.—Reg. Casp.; Sibir.
undulata, Schweigg. Enum. Fl. Hort. Regiom. 84 = salicifolia.
undulata, Willd. Enum. Hort. Berol. Suppl. 28 = pilosa.
uniflora, Dals. & Gibs. Bomb. Fl. 237 = microphylla.
uniflora, *G. Don, in Lond. Hort. Brit. Suppl.* i. 385. —Am. austr.

EUPHORBIA :—
uniflora, *Rafin. in Med. Repos. N. York*, v. (1808) 360. —Am. bor.
uniflora, *Resb. Hort. Beng.* [30]; *Fl. Ind.* ii. 473.—Ind. or.
uralensis, Fisch. ex Link, Enum. Hort. Berol. ii. 14 = virgata.
urceolophora, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 50.—Reg. Argent.
Vaccaria, Ball, Adansonia, vi. (1866) 286 = australis.
Vachellii, Hook. & Arn. Bot. Beech. Voy. 213 = serrulata.
vaginulata, *Griseb. Fl. Brit. W. Ind.* 52.—Ins Isham.
Vahlii, Willd. ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 87 = linearia.
valerifolia, Orteg. Hort. Matr. 127; Willd. Enum. Hort. Berol. 804 = E. Terracina.
Valerianæ, Pers. Syn. ii 18 = seq.
valerianæfolia, Lam. Encyc. ii. 485 = cyparissa.
varinbilis, *Ceæti, in Bibl. Ital.* xcl. (1838) 348.—Reg. Mediterr.
varians, Haw. Syn. Pl. Succ. 130 = E. Nivula.
variegata, Heyne, ex Roth, Nov. Pl. Sp. 225 = elegans.
variegata, Sims, Bot. Mag t. 1747 = marginata.
Vauthieriana, *Boiss. Cent. Euph.* 12 — Bras.
velleriflora, *Boiss. in DC. Prod.* xv. ii. 40.— Mexic.
velligera, *Schau in Linnaea*, xx. (1847) 728.— Mexic.
Velutschi, Boiss. & Rent. Pugill. Pl. Nov. 108 = Welwitschii.
venefica, *Trémaux, ex Boiss. in DC. Prod.* xv. ii 178. —Afr. trop.
venenata, Schlecht. in Linnæa, xix. (1847) 250 = E. Scotanum.
veneta, Tenore, Fl. Med. Univ. 894 = K. Characias.
veneta, Willd. Enum. Hort. Berol. 807 = Wulfenii.
verna, Salsm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 660 = E. Terracina.
verrucosa, Bertol. Fl. Ital. v. 45 = akenocarpa
verrucosa, Desf. Fl. Atlant. i. 303 = atlantica.
verrucosa, Georgi, Beschr. Russ. Reich. III. iv. 994 = dulcis.
verrucosa, Guss. Fl. Sic. Syn. i. 342 = Gasparrinii.
verrucosa, *Lam. Encyc.* ii. 481.—Europ.
verrucosa, Linn. Mant ii. 293 = cyparissa.
verrucosa, Linn. Sp. Pl. 459 = pubescens.
verrucosa, Pall. Tabl. Taur. E. 5 pilosa.
versicolor, *Greene, in Coult. Bot. Gaz.* vi. (1881) 184. —Am. bor.
verticillata, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 81 = Pallasii.
verticillata, Orph. ex Boiss. Diagn. Ser. II. iv. 90 = Heldreichii.
verticillata, Poir. Encyc. Suppl. ii 611 = petiolaris.
verticillata, Vell. Fl. Flum. 202; v. t. 16 = pilulifera.
vestita, *Boiss. Cent. Euph.* 7.—Mexic.
Vieillardi, *Baill. Adansonia*, ii. (1861-62) 212.— N. Caled.
villifera, *Schrebr, in Linnæa*, xxii. (1849) 153.— Texas.
villosa, Friwald. ex Boiss. in DC. Prod. xv. ii. 125 = epithymoides.
villosa, Waldst. & Kit. ex Willd. Sp. Pl. ii. 909 = pilosa.
villosissima, Klotzsch, ex Boiss. in DC. Prod. xv. ii. 51 = Sellol.
vimmalis, Burm. f. Fl. Cap. Prod. 14 = Burmanni.
viminalis, Lnn. Sp. Pl. 452 = Sarcostemma viminale.
vminalis, Mill. Gard. Dict. ed. VIII. n. 15 = E. Tirucalli.
viminea, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 184. —Ins. Galapag.
virgata, Desf. Tabl. ed. I. 204 = obtusifolia.
virgata, Not, ex Nym. Consp. 652 = lucida.
virgata, *Waldst. & Kit. Pl. Rar. Hung.* ii. 176. t. 162. —Europ.
viridiflora, Waldst. & Kit. l. c. ii. 309 = dulcis.
viridis, *Boiss, ex Klotzsch & Garcke, in Abh. Akad. Berl.* (1860) 40, in syn.—Persv.
virosa, *Willd. Sp. Pl. ii.* 883.—Afr. austr.
viscidula, *Boiss. Cent. Euph.* 5.—Bras.
viscillina, Lour. & Pard. Ser. Inconf. Aneg. 98 = isotidifolia.
Wahlbergia, *Boiss. in DC. Prod.* xv. ii. 169.—Afr. austr.
Wallichiana, Boiss. l. c. 30 = microphylla.

EUPHORBIA :—
Welwitschii, *Boiss. & Reut. Pugill. Pl. Nov.* 105 (Welwitschii); *Boiss. in DC. Prod.* xv. ii. 130.— Lusit.
Wheeleri, *Baill. Adansonia*, vi. (1866) 286.— Austral.
Wightiana, Boiss Cent. Euph. 37 = Rothiana.
Wittmanni, *Boiss. l. c* 31.—Reg. Caucas.
Wrightii, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 174.— Texas.
Wulfenii, Heldr. ex Nym. Consp. 651 = Sibthorpii.
Wulfenii, *Hoppe, in Flora*, xii (1829) 159.—Dalmatia.
xalapensis, *H. B. & K. Nov. Gen. et Sp.* ii. 61.— Mexic.
Xanti, *Engelm. ex Boiss. in DC. Prod.* xv. ii. 62.— Calif.
xylosylloides, *Bronyn. ex Lem. in Illustr. Hortic.* iv. (1857) *Misc* 72.—Madag.
yemenica, *Boiss Cent. Euph* 37.—Arab.
zambesiana, *Benth in Hook. Ic. Pl. t.* 1305.—Afr. trop.
zieriolides, *Boiss. in DC. Prod.* xv. ii. 58.—Mexic.
zonosperma, *Muell Arg. in Mart. Fl. Bras.* xi. ii. 606.—Bras.
zornioides, *Boiss. in DC. Prod.* xv. ii. 19.—Ind. or.
zygophylloides, *Boiss. Cent Euph.* 10.—Texas.

EUPHORBIASTRUM, Klotzsch & Garcke, in Monatsb Akad. Berl. (1859) 252 = **Euphorbia**, Linn.
Hoffmannianum, Klotzsch & Garcke, l. c. = Euphorbia Hoffmanniana.

EUPHORIA, Comm. ex Juss. Gen. 248 (1789), partm. *SAPINDACEÆ*, Benth. & Hook. f. i. 406.
annularis, Blanco, Fl. Filip. ed. I. 285 = Ratonia montana.
atterodata, Planch. ex Hiern, in Hook. f. Fl. Brit. Ind. i. 684 = Sapindus attenuatus.
chrysea, Korth. ex Miq. Fl. Ind. Bat. i. ii. 554 = Nephelium chryseum.
cinerea, *Kadik in Sitzb. Phys.-Math. Acad Muench.* viii. (1878) 299.—Ins. Philipp.
crinita, Poir. Encyc. Suppl. iii. 478 = Nephelium lappaceum.
Cubili, Blanco, Fl. Filip. ed. I. 287 = Cubilia Rumphii.
Danura, Wall. Cat. n. 8051 = Sapindus Danura.
didyma, Blanco, Fl. Filip. ed. I. 288 = Nephelium glabrum.
elongata, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1878) 96.—Borneo.
fuscata, Benth. & Hook. f. Gen. i. 406 = Glenniea zeylanica.
Gardneri, Benth. & Hook. l. c ; Bedd. Ic. Pl. Ind. Or. t. 87 = Nephelium Gardneri.
glabra, Blume, Bijdr. 283 = Nephelium glabrum.
informis, *Poir. Encyc. Suppl.* iii. 478.—Cochinch.
Leichhardtii, *Benth. Fl. Austral.* i. 468.—Austral.
Litchi, Blanco, Fl Filip. ed. I. 285 = Nephelium Longana.
Litchi, Juss. Gen. 248 ; Desf. Tabl. ed. I. 135 = Nephelium Litchi.
Litchi, Wall. Cat. n. 8048 G, H = Nephelium rubescens.
Longana, Blume, Bijdr. 283 = Nephelium mutabile.
Longana, Lam. Encyc. iii. 574 = Nephelium Longana.
malaiensis, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1878) 95.—Malacca.
Nephelium, DC. Prod. i. 612 = Nephelium lappaceum.
Nephelium, Wall. Cat. n. 8052 B, partim = Mallotus subpeltatus.
Noronhiana, Blume, Bijdr. 284 = Xerospermum Noronhianum.
pinnata. Poir. Encyc. Suppl. iii. 479 = Schmidelia pinnata.
Pometia, Steud. Nom. ed. II. i. 616 = præc.
punicea, Lam. Encyc. iii. 573 = Nephelium litchi.
pupillum, Steud. Nom. ed. II. ii. 192, in syn. = Nephelium Longana.
rubra, Royle, Illustr. Bot. Himal. 138 = Sapindus attenuatus.
sinensis, J. F. Gmel. Syst. 611 = Nephelium Litchi, Longana.
stellulata, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 308.—Ins. Philipp.
undulata, Herb. Heyne, ex Wall. Cat. n. 8049 = Nephelium Longana.
verrucolosa, Salisb. Prod. 280 = Nephelium Longana.

EUPHORBIA :—
verticillata, Lindl. Bot. Reg. t. 1059 = Sapindus Danura.
xerocarpa, Blume, Bijdr. 284 = Xerospermum Noronhianum.

EUPHORIOPSIS, Radlk. in Sitzb. Math.-Phys Acad Muench viii. (1878) 301, 302. *SAPINDACEÆ*. longifolia, *Radlk. ll. cc.*—Ind. or.

EUPHORNA, Steud. Nom. ed. II. i. 616 (1840) = **Euphrona**, Vell. (Quid ?).

EUPHRASIA, Linn. Syst. ed. I (1735). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 976.
agrestis, Arrond. Cat. Pl. Morb. 76 = officinalis.
alpina, *R. Br. Prod.* 436.—Austral.
alpina, Lam. Encyc. ii. 490 = officinalis.
alsa, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 107 = antarctica.
anclicola, *Benth. in DC. Prod.* x. 554.—Chill.
antarctica, *Benth. l. c* 555.—Austral.; Reg. Magell.
arctica, Lange, in Fl. Dan. t. 2910 = officinalis.
arguta, *R. Br. Prod.* 436.—Austral.
argula, A. Kern Sched. Fl Exsice. i. 40 = officinalis.
aromatica, *Noronha, in Verh Batav. Gen.* v. (1790) ed. I. Art. iv. 15 (Quid ?).—Java.
aspera, Host. Fl. Austr. i. 385 = Bartsia aspera.
aspera, Willd. ex Spreng. Syst. ii. 776 = Siriga aspera.
asperrima, Link, ex Spreng. l. c. 777 = Bartsia aspera ?
azorica, H. C. Wats. in Hook. Lond. Journ. Bot. iii. (1844) 598 = grandiflora.
Biancae, Guss. Pl. Suc Syn. ii. 841 = Bocconi.
Bocconi, *Guss. Adnot. Cat. Pl. Boccad.* (1821) 8.— Sicilia.
Bromnii, F. Muell. Fragm. v. 88 = speciosa, collina, alpina, striata.
caerulea, Tausch, in Flora, xvii. (1834) 445 = officinalis
campestris, Jord. Pugill. 131 = officinalis.
capensis, *Burm f. Fl. Cap. Prod.* 16.—Afr. austr.
carniolica, A. Kern. Sched. Fl. Exsice. ii. 117 = officinalis.
cebennensis, Martin, in Billot, Annot. Fl. Fr. Allem. 147 = officinalis.
chrysantha, *Phil. in Anal. Univ. Chil.* i. (1861) 66.— Chili.
caerulea, Auct. vide *caerulea*.
collina, *R. Br. Prod.* 436.—Austral.
condensata, Jord. Pugill. 135 = officinalis.
Cortx, Crantz, Stirp. Austr. ed. II. fasc. iv. 298 = Bartsia lutea.
coromandelsina, Rottl. ex Spreng. Pugill. i. 43 = Sopubia delphinifolia.
corsica, Lousel. Fl. Gall. ii. 867 = Bartsia Odontites.
cuneata, *Forst. f. Prod.* 43.—N. Zel.
cuprea, Jord. Pugill. 136 = officinalis.
cuspidata, *Hook. f. Fl. Tasman.* i. 298.—Tasman.
cuspidata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = officinalis
cuspidatissima, St. Lag. l. c. = officinalis.
cyanaica, *Pacho, Voy. Marm* t. 98.—Cyren.
danica, T. Ghem. Geogr. Beschr. Isl. 144.—Cf. Bab. in Journ. Linn. Soc. xi. (1870) 320 = officinalis.
depauperata, Benth. in Wall. Cat. n. 8686 = officinalis.
diemenica, Spreng. Syst. ii. 777 = alpina.
disperma, Hook. f. in Ic. Pl. t. 1333 = longiflora, *T. Kirk.*
divergens, Jord. ex F. Schultz, Arch. Fl. Fr. et Allem. 191 = Bartsia Odontites.
eriectorum, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 494 = officinalis.
exigua, Reut. in Compt. Rend. Soc. Hallér. (1854–56) 123 = officinalis.
Flifolia, Link, ex Steud. Nom. ed. II. i. 616 = officinalis.
flava, *Poir. Voy. Barb.* ii. 191.—Afr. bor.
frutescens, Sieber, ex Benth. in DC. Prod. x. 551.= Bocconi.
glandulosa, Benth. in DC. l. c. 555 = Phtheirospermum glandulosum.
glutinosa, *Bieb. Fl. Taur. Cauc.* ii. 70.—Tauris.
gracilis, Fries, Mant. iii. 82 = officinalis.
grandiflora, *Hochst. in Seub. Fl. Anor.* 89. t. 8.—Ins. Azores.
helmithospermum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 15 (Quid ?).—Java.

EUPHRASIA:—
hirsuta, *T. Gliem. Geogr. Reachr. Island*; *ex Bab. in Journ. Linn. Soc. xi.* (1870) 920.—Island.
hirtella, Jord. ex Rent. in Compt. Rend. Soc. Hallér. (1854–56) 120 = officinalis.
hispida, Willd. ex Spreng. Syst. ii. 776 = Bartsia melampyroides.
Hollians, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 20 = Bartsia scordifolia.
imbricata, Pers. Syn. ii. 149 = officinalis.
imbricata, *Phil. in Anal. Univ. Chil.* (1865) ii. 331.—Chili.
Jaubertiana, Bor. in Ann. Sc. Nat. Sér. II. vi. (1836) 264 = Bartsia Odontites.
kalifolia, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 617 = seq.
kalifornica, Pourr. ex Willd. Enum. Pl. Hort. Berol. 635, 636 = Bartsia purpurea.
Kochii, F. Schultz, Arch. f.t. 315 = Bartsia Odontites.
lactuca, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 14 (Quid?).—Java.
laevis, Gater. Fl. Montaub. 111 = Bartsia lutea.
lanceolata, Gaud. Fl. Helv. iv. 116 = Bartsia lutea.
Lapeyrousii, Soyer-Willem. Obs. Pl. Fr. 106 = officinalis.
latifolia, Linn. Sp. Pl. 604 = Bartsia latifolia.
latifolia, Pursh, Fl. Am. Sept. ii. 430 = officinalis.
latifolia, Schur, Enum. Pl. Transs. 510 = Bartsia Odontites.
lepontica ×, Brugg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 101.—Europ.
limifolia, Lam. Fl. Fr. ii. 349 = Bartsia viscosa.
limifolia, Linn. Sp. Pl. 604 = Bartsia lutea.
limifolia, Schousb. ex Bail, in Journ. Linn. Soc. xvi. (1878) 602 = Bartsia purpurea.
litoralis, *Fries, Mant.* iii. 196.—Europ.
longiflora, *T. Kirk, in Trans. N. Z. Inst.* x. (1878) 446.—N. Zel.
longiflora, *Vahl, Symb. Bot.* iii. 78.—Hispan.
lutea, Dubois, Fl. Orl. 341 = Bartsia Odontites.
lutea, Linn. Sp. Pl. 604 = Bartsia lutea.
maculata, Gilib. f. Lituan. i. 125 = officinalis.
maialis, Jord. Pugill. 134 = officinalis.
melastha, *Clos, in C. Gay, Fl. Chil.* v. 147.—Chili.
melampyroides, H. B. & K. Nov. Gen. et Sp. ii. 333 = Bartsia melampyroides.
micrantha, *Brenn. in Moddel. Soc. Faun. et Fl. Fenn.* v. (1880) 77.—Finland.
micrantha, Reichb. Fl. Germ. Excurs. 858 = officinalis.
minima, Jacq. ex DC. Fl. Fr. iii. 473 = officinalis.
minor, Link, Handb. i. 514 = officinalis.
mollis, Ledeb. ex Benth. in DC. Prod. x. 558 = officinalis.
montana, Jord. Pugill. 132 = officinalis.
multicaulis, Benth. in DC. Prod. x. 558 = collina.
Munroi, *Hook. f. Handb. New Zeal. Fl.* 220.—N. Zel.
muscosa, *Phil. in Anal. Univ. Chil.* (1873) 528.—Chili.
mutica, H. B. & K. Nov. Gen. et Sp. ii. 333 = Bartsia mutica.
nemorosa, Pers. Syn. ii. 149 = officinalis.
nitidula, Rent. in Compt. Rend. Soc. Hallér. (1854–56) 123 = officinalis.
nivalis, Beck, in Verh. Zool. Bot. Ges. Wien, (1883) 235 = officinalis.
Odontites, Linn. Sp. Pl. 604 = Bartsia Odontites.
officinalis, *Linn. f. c.*—Reg. bor. temp. et arct.
paludosa, R. Br. Prod. 436 = collina.
parviflora, Fries, Summa Veg. Scand. 196 = officinalis.
pectinata, Tenore, Fl. Nap. i. Prod. 26 = officinalis.
perpusilla, *Phil. in Linnaea*, xxix. (1857–58) 28.—Chili.
picta, *Phil. l. c.* xxiii. (1864–65) 213.—Chili.
picta, Wimm. Fl. Schles. ed. III. (1857) 407 = officinalis.
pinifolia, *Poir. Voy. Barb.* ii. 193.—Afr. bor.
praecox, Lam. Fl. Fr. ii. 351 = Bartsia latifolia.
pratensis, Reichb. Fl. Germ. Excurs. 859 = officinalis.
pratensis, Sebast. & Mauri, Fl. Rom. Prod. 199 = Bartsia viscosa.
pubervula, Jord. Pugill. 133 = officinalis.
pubescens, *Benth. in DC. Prod.* x. 555.—Peruv.
pulchella, A. Kern. Sched. Fl. Exsicc. i. 48 = officinalis.
pumila, A. Kern. l. c. 48 = officinalis.

EUPHRASIA:—
purpurea, Desf. Fl. Atlant. ii. 36 = Bartsia purpurea.
pygmaea, C. Koch, in Linnaea. xxii. (1849) 686 = officinalis.
ramosa, Pers. Syn. ii. 149 = officinalis.
ramosissima, Rent. in Compt.-Rend. Soc. Hallér. (1854–56) 119 = tricuspidata.
repens, *Hook. f. Fl. N. Zel.* i. 199.—N. Zel.
repens, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 14 (Quid?).—Java.
revoluta, *Hook. f. Fl. N. Zel.* i. 199.—N. Zel.
rigidifolia, *Bivona, ex Spreng. Syst.* ii. 775.—Sicilia.
rigidifolia, Klotzsch, ex Benth. in DC. Prod. x. 550 = *Odontites Auchers*).—Graecia; Oriens.
rigidifolia, Link, ex Nym. Consp. 550 (= *Odontites Linkii*).—Graecia.
rigidula, Jord. Pugill. 134 = officinalis.
rosmarinifolia, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 15 (Quid?).—Java.
Rothoviana, Hayne, Arzneigew. ix. t. 7 = officinalis.
rubra, Hoppe, Bot. Taschenb. (1791) 215; J. F. Gmel. Syst. 928 = Bartsia Odontites.
salisburgensis, Funk, ex Hoppe, l. c. (1794) 190 = officinalis.
santolinaefolia, H. B. & K. Nov. Gen. et Sp. ii. 333. t. 166 = Bartsia santolinaefolia.
saxatilis, Schur, Enum. Pl. Transs. 510 = officinalis.
scabra, *R. Br. Prod.* 437.—Austral.
serotina, Lam. Fl. Fr. ii. 350 = Bartsia Odontites.
simplex, D. Don, Prod. Fl. Nep. 95 = officinalis.
Soyeri, Timb. in Mém. Acad. Toul. Ser. V. vi. (1862) 36. t. 6 = officinalis.
speciosa, *R. Br. Prod.* 437.—Austral.
speciosa, A. Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 115 = officinalis.
spectabilis, *Phil. in Anal. Univ. Chil.* i. (1861) 66.—Chili.
spiralis, Pourr. ex Poir. Encyc. Suppl. ii. 596 = tenuifolia.
squarrosa, Salzm. ex Reichb. Fl. Germ. Excurs. 860 = Bartsia purpurea.
striata, *R. Br. Prod.* 436.—Austral.
stricta, Host, Fl. Austr. ii. 185 = officinalis.
stricta, H. B. & K. Nov. Gen. et Sp. Pl. ii. 333 = Bartsia stricta.
stricta, Schleich. Cat. Pl. Helv. ed. IV. 16 = officinalis?
subalpina, Schur, Enum. Pl. Transs. 510 = officinalis.
subexserta, *Benth. in DC. Prod.* x. 555.—Chili.
tatarica, Fisch. ex Spreng. Syst. ii. 777 = officinalis.
tenuifolia, *Pers. Syn.* ii. 150.—Lusitan.
tetragona, R. Br. Prod. 436 = collina.
tetraquetra, Arrond. in Bull. Soc. Polym. (1862) 96; ex Cat. Pl. Morb. 76 = officinalis.
tricuspidata, *Linn. Sp. Pl.* 604.—Tyrol.
trifida, *Poepp. ex Benth. in DC. Prod.* x. 554.—Chili.
Trixago, Vis. Fl. Dalmat. ii. 175 = Bartsia Trixago.
Ucchtritziana, Jung. & Engl. ex Jung. in Oestr. Bot. Zeitschr. xvii. (1867) 141 = officinalis.
uliginosa, Ducom. ex Rent. in Compt.-Rend. Soc. Hallér. (1854–56) 121 = officinalis.
verecunda, Salisb. Prod. 95 = Bartsia Odontites.
verna, Bell. in Mém. Acad. Turin, v. 1790–91 (1793) 289. t. 5 = Bartsia Odontites.
versicolor, A. Kern. Sched. Fl. Exsicc. i. 47 = officinalis.
violacea, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 14 (Quid?).—Java.
virgata, Link, ex Steud. Nom. ed. II. i. 616 = tenuifolia.
viscidula, Dum. Fl. Belg. 82 = seq.?
viscosa, Linn. Mant. 86 = Bartsia purpurea.
viscosa, Fall. Ind. Taur. ex Bleb. Fl. Taur. Cauc. ii. 70 = glutinosa.
viscosa, Sieber, ex Steud. Nom. ed. II. i. 616 = Bartsia viscosa.
vulgaris, Gmelodenst. ex Ledeb. Fl. Ross. iii. 362 = officinalis.
Willkommii, Freyn, in Flora, lxi. (1884) 681 = officinalis.

EUPHROBOSCIS, Wight, Ic. v. t. 1782 (1852), sphalm. = EUPROBOSCIS, Griff. = **Thelasis**, Blume (Orchid.).

EUPHRONIA, Vell. Fl. Flum. 177; iv. t. 107 (1952) (Quid?).
ternata, *Vell. Il. cc.*—Bras.

EUPHRONIA, Mart. Nov. Gen. et Sp. i. 121. t. 73 (1824). *ROSACEAE*, Benth. & Hook. f. i. 616. hirtelloides, *Mart. l. c.*—Bras.

EUPHROSINE, Allem. in Nov. Act. Nat. Cur. iv. (1770) 93 (Quid?).

EUPHROSINE, Endl. Gen. Suppl. i. 1381 (1841) = seq.

EUPHROSINIA, Reichb. Nom. 100 (1841) = seq.

EUPHROSYNE, DC. Prod. v. 380 (1836) = **Iva**, Linn. (Compos.).
ambrosiaefolia, A. Gray, Pl. Wright. i. 102 = Ambrosiaefolia.
parthenifolia, DC. Prod. v. 530.—Mexic.
xanthifolia, A. Gray, Pl. Wright. ii. 85 = I. xanthifolia.

EUPHYLEIA, Rafin. Fl. Tellur. iv. 16 (1836) = **Cordylina**, Roym. (Liliac.).
odorata, Rafin. l. c. 17 = C. Baueri?

EUPLASSA, Salisb. in Knight, Prot. 101 (1809). *PROTEACEAE*, Benth. & Hook. f. iii. 177.
ADENOSTEPHANUS, Klotzsch, in Linnaea, xv. (1841) 61.
DIDYNECKERIA, Vell. Fl. Flum. 41 (1825).
DIDYMANTHUS, Klotzsch, ex Meisn. in DC. Prod. xiv. 486 (1857 ?).
meridionalis, *Salisb. l. c.*—Guiana.

EUPLOCA, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 180 = **Heliotropium**, Tourn. (Borag.).
convolvulacea, Nutt. l. c. 190 = H. convolvulaceum.
grandiflora, Torr. in Emory, Milit. Reconnois. 146 = H. convolvulaceum.

EUPODIA, Rafin. Fl. Tellur. iii. 29 (1836) = **Chironia**, Linn. (Gentian.).
purpurea, Rafin. l. c. = C. peduncularis.

EUPOGON, Desv. Opusc. 67, in obs. (1831) = **Andropogon**, Linn. (Gramin.).

EUPOMATIA, R. Br. App. Flind. Voy. ii. 597. t. 2 (1814). *ANONACEAE*, Benth. & Hook. f. i. 29. Bennettii, *F. Muell. Fragm.* i. 45.—Austral.
laurina, *R. Br. App. Flind. Voy.* ii. 597. t. 2.—Austral.
laurina, Hook. in Bot. Mag. t. 4848 = Bennettii.

EUPREPIA, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 268 = **Astragalus**, Tourn. (Legumin.).

EUPROBOSCIS, Griff. in Calc. Journ. Nat. Hist. v. (1845) 371. t. 26 = **Thelasis**, Blume (Orchid.).
Griffithii, Reichb. f. in Bonplandia, v. (1857) 37 = T. carinata.
pygmaea, Griff. in Calc. Journ. Nat. Hist. v. (1845) 373. t. 26 = T. pygmaea.

EUPTELEA, Sieb. & Zucc. Fl. Jap. i. 138. t. 72 (1835). *MAGNOLIACEAE*, Benth. & Hook. f. i. 954.
Davidiana, *Baill. Adansonia*, xi. (1873–76) 305—Tibet.
pleiosperma, *Hook. f. & Thoms. in Journ. Linn. Soc* vii. (1864) 248.—Reg. Himal.
polyandra, *Sieb. & Zucc. Fl. Jap.* i. 134.—Japon.

EUPTERON, Miq. Fl. Ind. Bat. i. 1. 762 (1855) = **Polyosma**, Forst. (Araliac.).
acuminatum, Miq. l. c.=P. acuminata.
nodosum, Miq. l. c.=P. nodosa.

EUPTILIA, Rafin. Fl. Tellur. iv. 95 (1836) = **Scabiosa**, Tourn. (Dipsac.).
cretica, Rafin. l. c. = S. papposa.

EUPYRENA, Wight & Arn. Prod. 422 (1834) = **Timonius**, Rumph. (Rubiac.).
glabra, *Wight & Arn. l. c.* 422 = T. Jambosella.
hexasperma, Wight & Arn. ex Walp. Rep. ii. 486 = T. Rumphii?

EURANTHEMUM, Nees, ex Steud. Nom. ed. II. i. 616 (1840), sphalm. = **Brunthemum**, Linn. (Acanth.).

EURAPHIS, Trin. ex Lindl. Veg. Kingd. 115 (1847) = **Pappophorum**, Schreb. (Gramin.).

EUREIANDRA, Hook. f. in Benth. & Hook. f. Gen. i. 826 (1867). *CUCURBITACEAE*, Benth. & Hook. f. l. c.
Balfourii, *Cogn. & Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 404.—Ins. Socotr.
formosa, *Hook. f. in Benth. & Hook. f. Gen.* i. 825.—Afr. trop.
Schweinfurthii, *Cogn. in DC. Monog. Phan.* iii. 416.—Afr. trop.

EURHAPHIS, Trin. ex Steud Nom. ed. II. i. 616 (1840) = **Pappophorum**, Schreb. (Gramin.).

EURHOTIA, Neck. Elem. i. 207 (1790) = **Cephalis**, Sw. (Rubiac.).

EURIOSMA, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 421, sphalm. = **Eriosema**, DC. (Legumin.).

EURIPLES, Rafin. Fl. Tellur. iii. 94 (1836) = **Salvia**, Tourn. (Labiat.).
rugosa, Rafin. l. c. = S. virgata.

EUROSCHINUS, Hook. f. in Benth. & Hook. f. Gen. i. 422 (1862). *ANACARDIACEAE*, Benth. & Hook. f. l. c.
elegans, *Engl. in Bot. Jahrb.* i. (1881) 424.—N. Caled.
falcatus, *Hook. f. in Benth. & Hook. f. Gen.* i. 422.—Austral.
obtusifolius, *Engl. in Bot. Jahrb.* i. (1881). 424.—N. Caled.
verrucosus, *Engl. l. c.*—N. Caled.
Vieillardi, *Engl. l. c.*—N. Caled.

EUROSTORHIZA, G. Don, ex Steud. Nom. ed. II. i. 617 (1840) = **Physalis**, Linn. (Solanac.).

EUROSTORHIZA, Steud. Syn. Pl. Cyp. 265 (1855) = **Caustis**, R. Br. (Cyperac.).
Urvillei, Steud. l. c. = C. pentandra.

EUROSTORRHIZA, Benth. & Hook. f. Gen. iii. 1067 (1883) - prace.

EUROTIA, Adans. Fam. ii. 260 (1763). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 55.
CERATOSPERMUM, Pers. Syn. ii. 551 (1807).
DIOTIS, Schreb. Gen 638 (1791).
GUELDENSTAEDTIA, Neck. Elem. ii. 204 (1790).
KRANIKOFA, Rafin. Princ. Somiol. 30 (1814).
KRASCHENINNIKOVIA, Gueldenst. in Nov. Comm. Petrop. xvi. (1772) 548. t. 17.
capensis, E. Mey. ex Moq. in DC. Prod. xiii. ii. 307 = Sericocoma capensis.
ceratoides, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 239.—Europ.; As. bor.; Am. bor. occ.
Eversmanniana, *Stschegl. ex Transe. in Act. Hort. Petrop.* ix. (1884) 127, nomen.—Sibir.
ferruginea, Boiss. Fl. Orient. iv. 918, in syn. = ceratoides.
glabra, E. Mey. ex Mey. in DC. Prod. xiii. ii. 308 = Sericocoma pungens.
lanata, C. Koch, in Linnaea, xxii. (1849) 184 = ceratoides.
lanata, *Moq. Chenop. Enum.* 81.—Mexic.

EURYA, Thunb. Nov. Gen. 67 (1783). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 183.
GERRIA, Blume, Bijdr. 194 (1825).
acuminata, *DC. Mém. Ternstr.* 26; *Prod.* i. 525.—Reg. Himal.; Malaya.
acuminata, Wall. Cat. n. 1464, partim = symplocina.
aceps, *Blume, Mus. Bot. Lugd. Bat.* ii. 111.—Java.
angustifolia, Miq. ex Dyer, in Hook. f. Fl. Brit. Ind. i. 292 = Camellia theifera.
angustifolia, Wall. Cat. n. 1465 = acuminata.
angustifolia, *Wulf. Rep.* i. 389.—Java.
attenuata, Blume, Mus. Bot. Lugd. Bat. ii. 105 = Pyrenaria attenuata.
bifaria, Wall. Cat. n. 8791 = acuminata.
Blumeana, *Korth. Verh. Nat. Gesch. Bot.* 119.—Java.
causerosa, Blume, Mus. Bot. Lugd. Bat. ii. 106 = Freziera causerosa.
celebica, Relouv. ex Korth. Verh. Nat. Gesch. Bot. 118 = obovata.

EURYA:—
ceylanica, *Wight, Illustr.* i. 98.—Zeylan.
chinensis, *R. Br. in Abel, Journ.* 379.—China.
chinensis, Hook. f. & Thoms. ex Dyer, in Fl. Brit. Ind. i. 285 = acuminata.
chrysophylla, Blume, Mus. Bot. Lugd. Bat. ii. 105 = Freziera sericea.
clandestina, *Blume, l. c.* 121.—Java.
conocarpa, Korth. Verh. Nat. Gesch. Bot. 115 = japonica.
confinis, *Blume, Mus. Bot. Lugd. Bat.* ii. 110.—Java.
crupa, Siebold, ex Blume, l. c. 106 = japonica.
dioica, Blume, l. c. 105 = Elaeodendron dioscum.
elliptica, Gardn. in Calc. Journ Nat. Hist. vii. (1847) 443 = japonica.
euprista, Korth. Verh. Nat. Gesch. Bot. 113 = acuminata.
fasciculata, Buch.-Ham. in Wall. Cat. n. 1468 = acuminata.
fasciculata, Wall. Cat. n. 8662 = japonica.
glabra, Blume, Mus. Bot. Lugd. Bat. ii. 109 = obovata.
grandis, *Choisy, in Zoll. Syst. Vers. Ind. Archip.* 147 —Java.
Hasseltii, *Blume, Mus. Bot. Lugd. Bat.* ii. 112.—Java.
hirsuta, *Miq. Fl. Ind. Bat. Suppl.* 477.—Sumatra.
hortensis, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1848) 163 = japonica.
Humboldtiana, Blume, Mus. Bot. Lugd. Bat. ii. 105 = Freziera reticulata.
integrifolia, Blume, l. c. = Cleyera integrifolia.
Jacquemartii, *Carr. in Rev. Hortic* (1869) 369 — Japon.
japonica, *Thunb. Nov. Gen.* 67.—Japon.
lasiopetala, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 446 = Adinandra lasiopetala.
latifolia, Hort. ex C. Koch, Dendrol. i. 490 = japonica.
littoralis, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1848) 163 = chinensis.
lucida, Wall. Cat. n. 1462 = acuminata.
Macartneyi, *Champ. in Proc. Linn. Soc.* ii. (1850) 99. —Hongkong.
membranacea, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 444 = acuminata.
microphylla, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1848) 163 = japonica.
montana, Sieb. & Zucc. l. c. = japonica.
multiflora, DC. Mém. Ternstr. 25; Prod. i. 525 = acuminata.
multiflora, Siebold, ex Blume, Mus. Bot. Lugd. Bat ii. 106 = japonica.
myrtifolia, *Blume, l. c.* 113.—Java.
nervosa, Blume, l. c. 105 = Freziera nervosa.
nitida, Korth. Verh. Nat. Gesch. Bot. 115. t. 7 = japonica.
obovata, *Korth. l. c.* 118.—Java.
parvifolia, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 445 = chinensis.
phyllanthoides, Blume, Mus. Bot. Lugd. Bat. ii. 110 = japonica.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* 211.—Ins. Navigat.
pumila, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 14 = japonica.
pusilla, Siebold, ex Blume, Mus. Bot. Lugd. Bat. ii. 107 = japonica.
reticulata, *Korth. Verh. Nat. Gesch. Bot.* 118.—Sumatra.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* 212.—Ins. Samoa.
rigida, *Champ. in Mem. Soc. Phys. Genèv.* xiv. (1855) 126.—Java.
rostrata, *Blume, Mus. Bot. Lugd. Bat.* ii. 119.—Sumatra.
Roxburghii, Wall. Cat. n. 1465 = japonica.
salicifolia, Blume, Mus. Bot. Lugd. Bat. ii. 118 = acuminata.
salicifolia, *Choisy, in Zoll. Syst. Vers. Ind. Archip.* 148, 146.—Java.
sandwichensis, *A. Gray, Bot. U. St. Expl. Exped.* 209. —Ins. Sandvic.
sericea, Blume, Mus. Bot. Lugd. Bat. ii. 105 = Freziera sericea.
serrata, Walp. Rep. i. 389 = acuminata.
symplocina, *Blume, Mus. Bot. Lugd. Bat.* ii. 114.—Reg. Himal.; Java.

EURYA:—
systyla, Miq. ex Dyer, in Hook. f. Fl. Brit. Ind. i. 84 = japonica.
ternatana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 105.—Ins. Ternat.
theoides, Blume, Mus. Bot. Lugd. Bat. ii. 105 = Cleyera theoides.
timorensis, *Zipp. ex Span. in Linnaea*, xv. (1841) 177, nomen.—Ins. Timor.
trichocarpa, *Korth. Verh. Nat. Gesch. Bot.* 114.—Ins. Amboin.
trichogyna, Blume, Mus. Bot. Lugd. Bat. ii. 114 = prace.
tristyla, Wight & Arn. Prod. 86 = japonica.
undulata, Blume, Mus. Bot. Lugd. Bat. ii. 105 = Freziera undulata.
uniflora, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1848) 163 = japonica.
virens, *Blume, Mus. Bot. Lugd. Bat.* ii. 112.—Sumatra.
vitiensis, A. Gray, Bot. U. St. Expl. Exped. 209 = japonica.
Wallichiana, Planch. ex Dyer, in Hook. f. Fl. Brit. Ind. i. 286 = symplocina.
Wallichiana, Steud. Nom. ed. II. i. 617 = acuminata.
Wightiana, Wall. Cat. n. 8662 = japonica.
zeylanica, Walp. Rep. i. 389 = ceylanica.
Zollingeri, *Choisy, in Zoll. Syst. Vers. Ind. Archip.* 148, 147.—Java.

EURYALE, Salisb. in Kon. & Sims, Ann. Bot. ii. (1806) 73. *NYMPHAEACEAE*, Benth. & Hook. f. i. 47.
ANNESLEA, Roxb. ex Anda. Bot. Rep. t. 618 (1804); Roxb. Fl. Ind ii. 578 (1832).
amazonica, Poepp. in Froriep, Notizen, xxxv. (1832) 181 = Victoria regia.
brasiliana, Steud. Nom. ed. II. i. 617 = Victoria regia?
Cruziana, Orb. ex Walp. Ann. i. 22 = Victoria regia.
ferox, *Salisb. in Kon. & Sims, Ann. Bot.* ii. 74.—Ind. or.; China.
indica, Planch. in Ann. Sc. Nat. Sér. III. xix. (1853) 29 = ferox.

EURVALES, Steud. Nom. ed. II. i. 440, in syn. (1840) sphalm. = **Euryales**, Salisb. (Amaryll.).

EURYANDRA, Forst. Char. Gen. 81 (1776) = **Tetracera**, Linn. (Dillen.).
scandens, Forst. l. c. = T. Euryandra.

EURVANDRA, Hook. f. in Oliver, Fl. Trop. Afr. ii 583 (1871) = **Eureiandra**, Hook. f. (Cucurb.).
Sumbul, Kauffm. l. c. = F. Sumbul.

EURYANTHE, Cham. & Schlecht. in Linnaea, v. (1830) 224 = **Amoreuxia**, Moç. & Sesse (Bixin.).
Schiedeana, Cham. & Schlecht. l. c. 226 = A. palmatifida.

EURYBIA, Cass. in Dict. Sc. Nat. xvi. 46 (1820) = **Olearia**, Moench (Compos.).
aculeata, DC. Prod. v. 370 = O. ramulosa.
adenophora, F. Muell. Fragm. i. 111 = O. adenophora.
affinis, Steetz, in Lehm. Pl. Preiss. i. 421 = O. helecophila.
albida, Hook. f. Fl. N. Zel. i. 118 = O. albida.
alpicola, F. Muell. in Proc. Roy. Soc. Tasm. iii. (1859) 229 = O. alpicola.
alpina, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 106 = O. personioides.
alpina, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 84 = O. nitida.
argophylla, Cass. in Dict. Sc. Nat. xxxvii. 467 = O. argophylla.
artemisioides, Sond. & F. Muell. in Linnaea, xxv. (1852) 456 = O. tubuliflora.
aspera, Steetz, in Lehm. Pl. Preiss. i. 420 = O. strigosa.
asteroidea, F. Muell. Fragm. i. 111 = O. asteroidea.
avicenniaefolia, Hook. f. Fl. N. Zel. i. 120 = O. avicenniaefolia.
axillaris, DC. Prod. v. 366 = O. axillaris.
Beckleri, F. Muell. Veg. Chath. Isl. 21 = O. cydoniaefolia.

EURYBIA :—
brachyglossa, DC. Prod. v. 265 = Olearia axillaris.
brachyphylla, F. Muell. ex Sond. in Linnæa, xxv. (1852) 465 = O. lepidophylla ?
candidissima, Steetz, in Lehm. Pl. Preiss. i. 418 = O. axillaris.
capitellata, DC. Prod. v. 266 = O. axillaris.
cardiophylla, F. Muell. in Linnæa, xxv. (1852) 398 = O. pannosa.
chrysophylla, DC. Prod. v. 266 = O. chrysophylla.
chrysotricha, Tenore, ex Lindl. Bot. Reg. (1841) Misc. 19.—Austral.
ciliata, Benth. in Enum. Pl. Hueg. 58 = O. ciliata.
Colensoi, F. Muell. Veg. Chath. Isl. 22 = O. Colensoi.
collina, F. Muell. ex Sond. in Linnæa, xxv. (1852) 465 = O. ramulosa.
commixta, Nees, Gen. et Sp. Aster. 142 = Aster Herveyi.
conocephala, F. Muell. in Trans. Vict. Inst. i. (1855) 36 = O. conocephala.
corymbosa, Cass. in Dict. Sc. Nat. xxvii. 487 = Aster corymbosa.
cunefolia, Walp. in Linnæa, xiv. (1840) 816 = O. pimeleoides.
Cunninghamii, Hook. f. Fl. N. Zel. i. 117. t. 30 = O. Cunninghamii.
cydoniæfolia, DC. Prod. v. 267 = O. cydoniæfolia.
Dampieri, DC. l. c. 266 = O. axillaris.
decurrens, DC. l. c. 269 = O. decurrens.
dentata, Hook. f. Fl. N. Zel. 118 = O. ilicifolia.
elæophila, DC. Prod. v. 269 = O. heleophila.
epilepia, DC. l. c. 270 = O. ramulosa.
ericoides, Hook f. Fl. Tasm. ii. 368 = O. Hookeri.
ericoides, F. Muell. ex Sond. in Linnæa, xxv. (1852) 464 = O. floribunda.
ericoides, Steetz, in Lehm. Pl. Preiss. i. 423 = O. ramulosa.
erubescens, DC. Prod. v. 267 = O. myrsinoides.
Ferresii, F. Muell. Fragm. iii. 18. t. 18 = O. Ferresii.
floribunda, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 109 = O. floribunda.
Forsteri, Hook. f. Fl. N. Zel. 119 = O. Forsteri.
fulgida, Steud. Nom. ed. II. i. 616, sphalm. = seq.
fulvida, Cass. in Dict. Sc. Nat. xvi. 48 = O. stellulata.
furfuracea, DC. Prod. v. 267 = O. furfuracea.
glandulosa, DC. l. c. 269 = O. glandulosa.
glomerata, Benth. ex Nees, Gen. et Sp. Aster. 139 = Aster macrophyllus.
glutescens, F. Muell. ex Sond. in Linnæa, xxv. (1852) 462 = O. decurrens.
glutescens, Sond. l. c. = O. glutinosa.
glutinosa, Lindl. Bot. Reg. (1839) Misc. 88 = O. glutinosa.
gracilis, Benth. in Enum. Pl. Hueg. 58.—Austral.
Gunniana, DC. Prod. v. 268 = O. stellulata.
Hookeri, F. Muell. ex Sond. in Linnæa, xxv. (1852) 463 = O. Hookeri.
hygrophila, DC. Prod. v. 269 = O. hygrophila.
illita, F. Muell. Fragm. i. 16 = O. elliptica.
imbricata, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 61 = O. imbricata.
iodochroa, F. Muell. Fragm. ii. 110 = O. iodochroa.
Iwasseni, Cass. in Dict. Sc. Nat. xxxvii. 487 = Aster Herveyi.
lamprophylla, F. Muell. ex Sond. in Linnæa, xxv. (1852) 459 = O. myrsinoides.
ledifolia, DC. Prod. v. 269 = O. ledifolia.
Lehmanniana, Steetz, in Lehm. Pl. Preiss. i. 422 = O. heleophila.
lepidophylla, DC. Prod. v. 270 = O. lepidophylla.
leptophylla, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 171.— Austral.
linearifolia, DC. Prod. v. 266 = O. axillaris.
linifolia, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 109 = O. glutinosa.
lirata, DC. Prod. v. 267 = O. stellulata.
Lyallii, Hook. f. Fl. Antarct. Suppl. 848 = O. Lyallii
macrophylla, Cass. in Dict. Sc. Nat. xxvii. 487 = Aster macrophyllus.
megalophylla, F. Muell. in Proc. Roy. Soc. Tasm. iii. (1859) 238 = O. megalophylla.
microphylla, DC. Prod. v. 270 = O. ramulosa.
Muelleri, Sond. in Linnæa, xxv. (1852) 459 = O. Muelleri.
muricata, Steetz, in Lehm. Pl. Preiss. i. 423 = O. muricata.
myrsinoides, Nees, Gen. et Sp. Aster. 146 = O. myrsinoides.

EURYBIA :—
nitida, Hook. f. Fl. N. Zel. i. 117 = Olearia nitida.
nummularifolia, Hook. f. l. c. 116 = O. nummularifolia.
obcordata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 106 = O. obcordata.
oligantha, DC. Prod. v. 266 = O. axillaris.
operina, F. Muell. Veg. Chath. Isl. 22 = O. operina.
opposita, F. Muell. Fragm. iii. 55 = O. chrysophylla.
paniculata, Steetz, in Lehm. Pl. Preiss. i. 422 = O. heleophila.
pannosa, F. Muell. Pl. Vict. t. 32 = O. pannosa.
pauciclenata, Steetz, in Lehm. Pl. Preiss. i. 420 = O. pauciclenata.
persoonioides, DC. Prod. v. 267 = O. persoonioides.
picridifolia, F. Muell. in Linnæa, xxv. (1852) 397 = O. picridifolia.
pimeleoides, DC. Prod. v. 268 = O. pimeleoides.
pinifolia, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 108 = O. pinifolia.
propinqua, DC. Prod. v. 270 = O. ramulosa.
quercifolia, Cass. in Dict. Sc. Nat. xvi. 47 = O. stellulata.
ramosissima, DC. Prod. v. 270 = O. ramosissima.
ramulosa, DC. l. c. = O. ramulosa.
rosmarinifolia, DC. l. c. 268 = O. rosmarinifolia.
rudis, Benth. in Enum. Pl. Hueg. 58 = O. rudis.
rugosa, F. Muell. in Journ. Linn. Soc. v. (1861) 22 = O. stellulata.
scabra, Benth. in Enum Pl. Hueg. 58 = O. rudis.
Schreberi, Nees, Gen. et Sp. Aster. 138 = Aster macrophyllus
semidentata, F. Muell. Veg. Chath. Isl. 21 = O. semidentata.
seriocarpa, F. Muell. ex Sond. in Linnæa, xxv. (1852) 459 = O. pimeleoides.
Solandri, Hook. f. Fl. N. Zel. 119 = O. Solandri.
stregua, Steetz, in Lehm. Pl. Preiss. i. 419 = O. strigosa.
Stuartii, F. Muell. Fragm. i. 202 = O. Stuartii.
subrepanda, DC. Prod. v. 268 = O. stellulata.
subspicata, Hook. in Mitch. Journ. Trop. Austral. 298 = O. subepicata.
tenuifolia, DC. Prod. v. 269 = O. tenuifolia.
teretifolia, Sond. in Linnæa, xxv. (1852) 464 = O. teretifolia.
Traversii, F. Muell. Veg. Chath. Isl 19.—Ins. Chath.
tubuliflora, Sond. & F. Muell. in Linnæa, xxv. (1852) 455 = O. tubuliflora.
virgata, Hook. f. Fl. N. Zel. 110 = O. virgata.
viscidula, F. Muell. Fragm. i. 50 = O. viscidula.
viscosa, Cass. in Dict. Sc. Nat. xxxvii. 487 = O. viscosa.
xerophila, F. Muell. Fragm. i. 51 = O. xerophila.

EURYBIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 464 (1821) = **Aster**, Tourn. (Compos.).
maritima, S. F. Gray, l. c. = A. Tripolium.

EURYBIOPSIS, DC. Prod. v. 260 (1836) = **Wittadinia**, A. Rich. (Compos.).
australis, Hook. f. Fl. N. Zel. 125 = V. australis.
gracilis, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 110 = V. australis.
Hookeri, F. Muell. in Linnæa, xxv. (1852) 453 = V. australis.
intricata, F. Muell. l. c. 396 = Minuria Cunninghamii.
macrorrhiza, DC. Prod. v. 260 = V. macrorhiza.
scabrida, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 110 = V. australis.

EURYCHAENIA, Griseb. Fl. Brit. W. Ind. 258 (1860); et in Goett. Abh. ix. (1861) 46 = **Micoania**, Ruiz & Pav. (Melastom.).
ferruginea, Griseb. Fl. Brit. W. Ind. 259, 788 = M. ferruginea.
fulva, Griseb. l. c. 259 = M. dolichorrhyncha.
punctulata, Griseb. l. c.; et in Goett. Abh. ix. (1861) 46 = M. fulva.

EURYCHANES, Nees, in Mart. Fl. Bras. ix. 52. t. 3 (1847) = **Eualitis**, Plum. (Acanthac.).
verbasciformis, Nees, l. c. 53.—Bras.

EURYCHITON, Nimmo, in J. Grah. Cat. Pl. Bomb. Addenda (1839) = **Statice**, Linn. (Plumbag.).
adenota, Nimmo, l. c. = S. axillaris ?

EURYCLES, Salisb. in Trans. Hort. Soc. i. (1812) 337. *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 734.
PROSPHYS, Herb. App. 42 (1821).
STEMONIX, Rafin. Atl. Journ. 165 (1838).
alata, Sweet, Hort. Brit. ed. I. 406 = sylvestris.
amboinensis, [Lindl. in] Loud. Encyc. Pl. 345 = sylvestris
australasica, [Lindl. in] Loud. l. c. = sylvestris.
australis, Schult. f. Syst. vii. 911 = sylvestris.
coronata, Sweet, Hort. Brit. ed. I. 406 = sylvestris.
Cunninghamii, *Ait. ex Loud. Hort. Brit. Suppl.* i. 588. —Austral.
javanica, M. Roem. Syn. Enast. 157 = sylvestris.
nervosa, G. Don, ex Loud. Hort. Brit. 117 = nuda.
nuda, *Sweet, Hort. Brit.* ed. I. 406.—Ind. or.
rotundifolia, M. Roem. Syn. Enast. 156 = sylvestris.
sylvestris, *Salisb. in Trans. Hort. Soc.* i. (1812) 337 —Malaya, Austral.

EURYCOMA, Jack, in Malay. Misc. ii. (1822) VII. 44. *SIMARUBEÆ*, Benth. & Hook. f. i. 312.
apiculata, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 522.—Penang.
longifolia, *Jack, in Malay. Misc.* ii. (1822) VII. 45.— Burma ; Malaya.
mergurnsis, Planch. in Hook. Lond. Journ. Bot. v (1846) 584 = longifolia.
tanayana, Wall. Cat. n. 8528 = longifolia.

EURYGANIA, Klotzsch, in Linnæa, xxiv. (1851) 26. *VACCINIACEÆ*, Benth. & Hook. f. ii. 568.
angustifolia, *Klotzsch, l. c.*—Peruv.
antisiensifolia, *Klotzsch, l. c.*—N. Granat.
bifora, *Benth. & Hook. f. Gen.* iii. 568.—Peruv.
mellifora, *Benth. & Hook. f. Gen.* ii. 568.—Peruv.
multiflora, Klotzsch, in Linnæa, xxiv. (1851) 27 = seq.
ovata, *Hook. f. Bot. Mag.* t. 6893.—Peruv.
parvifolia, *Benth. & Hook. f. Gen.* ii 568.—N. Granat.
phillyreæfolia, *Benth. & Hook. f. l. c.*—Peruv.
polyantha, *Benth. & Hook. f. l. c.*—Peruv.

EURYLEPIS, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 164 = **Erica**, Tourn.
albens, D. Don, l. c. = Erica albens.
azaleæflora, Steud. Nom. ed. II. i. 618 = seq.
azaleæfolia, G. Don, Gen. Syst. iii. 808 = Erica azaleæfolia.
bracteata, G. Don, l. c. = Erica bracteata.
crossata, G. Don, l. c. = Erica acuta.
diosmæfolia, G. Don, l. c. 809 = Erica diosmæfolia.
fragrans, G. Don, l. c. 808 = Erica fragrans.
glabra, G. Don, l. c. = Erica ventricosa.
halicacaba, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 164 = Erica halicacaba.
Massoni, D. Don, l. c. = Erica Massoni.
Monsoniæ, G. Don, Gen. Syst. iii. 809 = Erica Monsoniana.
mucronata, G. Don, l. c. 808 = Erica mucronata.
obcordata, G. Don, l. c. = Erica bracteata.
pachyphylla, G. Don, l. c. 809 = Erica brevifolia.
pallida, Steud. Nom. ed. II. i. 615 = seq.
palliflora, G. Don, l. c. = Erica palliflora.
petiolata, G. Don, Gen. Syst. iii. 808 = Erica petiolata.
pyrolæflora, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 164 = Erica trifora.
rigidifolia, G. Don, Gen. Syst. iii. 809 = Erica lucida.
sexfaria, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 164 = Erica sexfaria.
tetragona, G. Don, Gen. Syst. iii. 808 = Erica tetragona.
Thunbergii, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 164 = Erica Thunbergii.
trifiora, D. Don, l. c. = Erica trifora.

EURYLOBIUM, Hochst. in Flora, xxv. (1842) 228. *VERBENACEÆ*, Benth. & Hook. f. ii. 1128.
serrulatum, *Hochst. l. c.*—Afr. austr.

EURYLOMA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 165 = **Erica**, Tourn.
Alteni, D. Don, l. c. = Erica Alticola.
ampullacea, G. Don, Gen. Syst. iii. 816 = Erica ampullacea.
ampullisvoides, G. Don, l. c. 817 = Erica Forbesiana.
aristata, G. Don, l. c. = Erica aristata.
aristella, G. Don, l. c. = Erica aristella.
capax, G. Don, l. c. = Erica obbata.
cerinita, G. Don, l. c. = Erica cerinita.

ERYLOMA :—
curvifolia, G. Don, Gen. Syst. iii. 816 = Erica curvifolia.
Hartwelli, G. Don, l. c. = Erica crinita.
imperialis, G. Don, l. c. = Erica fascicularis.
Irbyana, G. Don, l. c. 817 = Erica Irbyana.
jasminiflora, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 156 = Erica jasminiflora.
mirabilis, G. Don, Gen. Syst. iii. 816 = Erica praestans.
notabilis, G. Don, l. c. = Erepia tubercularis.
obbata, G. Don, l. c. 817 = Erica obbata.
oblonga, G. Don, l. c. = Erica Massoni.
Princeps, G. Don, l. c. = Erica Massoni.
recurvata, G. Don, l. c. = Erica recurvata.
retorta, G. Don, l. c. = Erica retorta.
Shannonensa, G. Don, l. c. 816 = Erica Shannonea.
squarrosa, G. Don, l. c. 817 = Erica squarrosa.
tricolor, G. Don, l. c. 816 = Erica aristata.

ERYLOMA, Rafin. Fl. Tellur. iv. 75 (1836) = **Ipomoea**, Linn. (Convolv.).
grandiflora, Rafin. l. c. = I. Bona-nox.
latiflora, Rafin. l. c. = I. Bona-nox.
leucantha, Rafin. l. c. = I. leucantha.

ERYNEMA, Endl Gen. Suppl ii. 93 (1842) = **Hermannia**, Linn. (Sterculac.).

ERYNOMA, Stend. Nom. ed. II. i. 618 (1840) = seq.

ERYNOME, DC. Prod. iv. 475 (1830) = **Coprosma**, Forst. (Rubiac.).
cantkioides, DC. l. c. = C. Billardieri.

ERYOMVKTUS, Schau. in Linnaea, xvii. (1843) 239 = **Baeckea**, Linn. (Myrtac.).
alpina, Schau. l. c. = B. diffusa.
diffusa, Schau. l. c. = B. diffusa.
parviflora, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 149 = B. diffusa.
Stuartiana, F. Muell. ex Miq. l. c. = B. diffusa.
thymifolia, Schau. in Linnaea, xvii. (1843) 239 = B. diffusa.

ERYOPS, Cass. in Bull. Soc. Philom (1818) 140. *COMPOSITAE*, Benth. & Hook. f. ii. 452.
ENANTIOTRICHUM, E. Mey. ex DC. Prod. vi. 443 (1837).
HERTIA, Neck. Elem. i. 8 (1790).
abrotanifolius, *DC. Prod.* vi. 443.—Afr. austr.
algoensis, *DC. l. c.* 443.—Afr. austr.
arabicus, *Steud. ex Yaub. & Spach, Illustr.* iv. 88. t. 355.—Arab.
asparagoides, *Less. Syn. Comp.* 394.—Afr. austr.
Athanasiae, DC. Prod. vi. 444 = abrotanifolius.
Athanasiae, *Less. Syn. Comp.* 394.—Afr. austr.
calvescens, *DC. Prod.* vi. 445.—Afr. austr.
Candollei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 412.—Afr. austr.
carnosus, Cass. in Dict. Sc. Nat. xvi. 52 = tenuisatinus.
comosus, Cass. l. c. 53 = abrotanifolius.
densifolius, Sond. in Harv. & Sond. Fl. Cap. iii. 416 = sulcatus.
diversifolius, *Harv. in Harv. & Sond. l. c.* 415.—Afr. austr.
Dregeanus, *Sch. Bip. in Flora*, xxviii. (1845) 51.—Afr. austr.
empetrifolius, *DC. Prod.* vi. 446.—Afr. austr.
flabelliformis, Cass. in Dict. Sc. Nat. vi. 31 = virgineus.
imbricatus, DC. Prod. vi. 446 = Candollei.
imbricatus, *Less.* Syn. Comp. 394 = laterifforus.
intermedius, DC. Prod. vi. 444 = abrotanifolius.
laterifforus, *Less. Syn. Comp.* 394.—Afr. austr.
linearis, *Harv. Thes. Cap.* ii. 34.—Afr. austr.
linifolius, *DC. Prod.* vi. 446.—Afr. austr.
longipes, *DC. l. c.* 445.—Afr. austr.
multifidus, *DC. l. c.* 444.—Afr. austr.
oligoglossus, *DC. l. c.* 446.—Afr. austr.
pectinatus, *Cass. in Dict. Sc. Nat.* xvi. 51.—Afr. austr.
pinifolia, *A. Rich. Tent. Fl. Abyss.* i. 445. t. 80.—Afr. trop.
punctatus, *DC. Prod.* vi. 445.—Afr. austr.
racemosus, DC. l. c. 446 = oligoglossus.
Serra, *DC. l. c.* 444.—Afr. austr.
sootranus, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 841.—Ins. Socotra.

ERYOPS:—
spathaceus, *DC. Prod.* vi. 446.—Afr. austr.
speciosissimus, DC. l. c. 444 = Athanasiae.
strictus, DC. l. c. 447 = tenuisatinus.
subcarnosus, *DC. l. c.* 445.—Afr. austr.
sulcatus, *Less. Syn. Comp.* 394; *Harv. in Harv. & Sond. Fl. Cap.* iii. 416.—Afr. austr.
subsessilis, *Sch. Bip. in Flora*, xxviii. (1845) 52.—Afr. austr.
tenuisatinus, *Less. Syn. Comp.* 394.—Afr. austr.
trifidus, *Less. l. c.*—Afr. austr.
trifurcatus, *Cass. in Dict. Sc. Nat.* xvi 52.—Afr. austr.
trilobus, *Harv. in Harv. & Sond. Fl Cap* iii. 411—Afr. austr.
triqueter, Less. ex Harv. & Sond. l. c. 414 = oligoglossus.
virgineus, *Less. Syn. Comp.* 394; *DC. Prod.* vi. 445—Afr. austr.

ERYOSMA, Walp. Rep. i. 785 (1842) = **Eriosema**, DC. (Legumin.).

ERYPTERA, Nutt ex Torr. & Gray, Fl N. Am. i. 629 (1840) = **Feuceodanum**, Linn. (Umbellif.).
lucida, Nutt ex Torr. & Gray, l. c. = F. Emyptera.

ERYSPERMUM, Salisb. Parad. Lond. t. 75 (1807) = **Leucadendron**, K. Br. (1810) (Proteac.).
aemulum, Knight, Prot. 21 = L. aemulum.
aetherum, Knight, l. c. 19.—Afr austr.
argenteum, Knight, l. c. 22 = L. argenteum.
concolor, Knight, l. c. 24 = L. concolor.
cuspidatifolium, Knight, l. c. 20 = L. strictum.
decorum, Knight, l. c. 24 = L. decorum.
densum, Knight, l. c. 23.—Afr. austr.
frondosum, Knight, l. c. 19 = L. adscendens.
grandiflorum, Salisb Parad. Lond. t. 105 = L. grandiflorum.
humiflusum, Knight, Prot. 19 = L. adscendens.
microcephalum, Knight, l. c. 18.—Afr. austr.
multiflorum, Knight, l. c. 20.—Afr. austr.
proserum, Knight, l. c.—Afr. austr.
salicifolium, Salisb. Parad. Lond. t. 75 = L. strictum.
saligmum, Knight, Prot. 22 = L. saligmum.
sericeum, Knight, l. c. 21 = L. sericeum.
spissifolium, Knight, l. c. 23.—Hab.?
ulginosum, Knight, l. c. 21 = L. ulginosum.
volutaefolium, Knight, l. c. 23.—Afr. austr.

ERYSTEGIA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 154 = **Erica**, Tourn.
andromedaeflora, D. Don, l. c. = Erica andromedaeflora.
Corydalis, G. Don, Gen. Syst. iii. 810 = Erica Corydalis.
elegans, G. Don, l. c. 809 = Erica glauca.
emarginata, G. Don, l. c. = Erica nigrita.
glauca, D. Don, in Edinb N. Phil. Journ xvii. (1834) 154 = Erica glauca.
lachneaeflora, G. Don, Gen. Syst. iii. 809 = Erica Lachnaea.
lanuginosa, G. Don, l. c. = Erica lanuginosa.
marginata, Benth in DC. Prod. vii. 687, sphalm. = Erica nigrita.
pannosa, G. Don, l. c. 810 = Erica barbata.
pomifera, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 154 = Erica andromedaeflora.
regnalisfolia, G. Don, Gen. Syst. iii. 809 = Erica squamosa.
triceps, G. Don, l. c. = Erica triceps.
trumphans, G. Don, l. c. = Erica andromedaeflora.

ERYSTYLES, Wawra, in Oestr. Bot. Zeitschr. xiii. (1863) 228. *SCITAMINEAE*.
Cotyledon, *Wawra, l. c.*—Bras.

ERYSTYLUS, Bouché, in Linnaea, xviii. (1844) 485 = **Canna**, Linn. (Scitam.).
flaccidus, Bouché, l. c. = C. flaccida.
Reevesii, Bouché, l. c. = C. Reevesii.

ERYTAENIA, Torr. & Gray, Fl. N. Am. i. 638 (1838–40). *UMBELLIFERAE*, Benth. & Hook. f. i. 907.
macrophylla, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 455 = texana.
texana, *Torr. & Gray, Fl. N. Am.* i. 638.—Am. bor. occ.

ERYTALIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 119 = ERYTHALIA, Renealm. = **Gentiana**, Tourn.

ERYTENIA, Buckl. in Proc. Acad. Soc. Philad. 1861 (1862) 455 = **Eurytaenia**, Torr. & Gray (Umbellif.).

ERYTHALIA, D. Don, in Trans. Linn. Soc. xvii. (1837) 515 = ERYTHALIA, Renealm. = **Gentiana**, Tourn.

EUSARCOPS, Rafin. Fl. Tellur. iv. 11 (1836) = **Hippeastrum**, Herb. (Amaryllid.).
reticulata, Rafin. l. c. = H. reticulatum.

EUSCAPHIS, Sieb. & Zucc. Fl. Jap. i. 124. t. 67 (1835). *SAPINDACEAE*, Benth. & Hook. f. i. 412.
HENOKIA, Rafin. Alsog. Am. 47 (1838).
simplicifolia, *Sieb. & Zucc. l. c.* 189, in obs.—Japon. staphyleoides, *Sieb. & Zucc. l. c.* 124.—Japon.

EUSIDEROXYLON, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 292. *LAURINEAE*, Benth. & Hook f. ii. 455.
BIHANIA, Meisn. in DC. Prod. xv. i. 96 (1864).
SALGADA, Blanco, Fl Filip. ed. II. 221 (1845)
Zwageri, *Teijsm. & Binn. l. c.*—Borneo.

EUSIPHO, Salisb. Gen. Pl. Fragm. 139 (1866) = **Cyrtanthus**, Ait. (Amaryllid.).
angustifolia, Salisb l. c. = C. angustifolius.
odora, Salisb. l. c. = C. odorus.

EUSMIA, Humb. & Bonpl. Pl. Aequin. ii. t. 184 (1808) = **Eucosmia**, H. B. & K. (Rubiac.).

EUSOLENANTHE, Benth. & Hook f. Gen. iii. 975 (1883), sphalm. = ENSOLENANTHE, Schott = **Alocasia**, Neck. (Aroid.).

EUSTACHIA, Rafin. Cat. 14 (1824) = seq.

EUSTACHYA, Rafin. in Am. Monthly Mag. (1819) 190 = **Veronica**, Tourn. (Scrophul.).
alba, Rafin. Cat. 14 = V. virginica.
cornuta, Rafin. ex DC. Prod. x. 464 = V. virginica.
japonica, Rafin. ex Steud. Nom. ed. II. i. 618 = V. virginica.
oppositifolia, Rafin. New Fl. Am. ii. 21.—Am. bor.
purpurea, Rafin. in Am. Monthly Mag. (1819) 190 = V. virginica.

EUSTACHYS, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = **Chloris**, Sw. (Gramin.).
dactichophylla, Nees, in Mart. Fl. Bras. ii. 418 = C. dactichophylla
floridana, Chapm. Fl. S. U. St. 557 = C. floridana.
glauca, Chapm. l. c. = C. glauca.
petraea, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = C. petraea.
polystachya, Kunth, Rev. Gram. i. 88 = C. inermis.
submutica, Roem. & Schult. Syst. ii. 614 = C. submutica.

EUSTACHYS, Salisb. Gen. Pl. Fragm. 88 (1866) = **Ornithogalum**, Tourn. (Liliac.).
latifolia, Salisb. l. c. = O. latifolium.
pyramidale, Salisb. l. c. = O. pyramidale.

EUSTATHES, Spreng. Syst. ii. 319 (1825) = **Bystathes**, Lour. (Sapindac.).

EUSTAXIA, Rafin. Fl. Tellur. 56 (1836) = **Veronica**, Tourn. (Scrophul.).
alba, Rafin. l. c. = V. virginica.
japonica, Rafin. l. c. = V. virginica?
oppositifolia, Rafin. l. c. = V. spicata.
purpurea, Rafin. l. c. = V. virginica.
sibirica, Rafin. l. c. = V. virginica.
villosa, Rafin. l. c. = V. longifolia.

EUSTEGIA, R. Br. in Mem. Wern. Soc. i. (1809) 51. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 765.
filiformis. *Roem. & Schult. Syst.* vi. 120.—Afr. austr.
hastata, *R. Br. in Mem. Wern. Soc.* i. (1809) 51.—Afr. austr.
Asotata, Sieber, ex Decne. in DC. Prod. viii. 511 = Microloma sagittatum.
humilis, *E. Mey. Comm. Pl. Afr. Austr.* 221.—Afr. austr.
Lonchitis, *E. Mey. l. c.*—Afr. austr.

EUSTEGIA, Rafin. Sylva Tellur. 95 (1838)† (Melastom.).
Mutisi, Rafin. l. c.= Centronia Mutisii.
pulverulenta, Rafin. l. c. = Pogonanthera pulverulenta.
tomentosa, Rafin. l. c. = Miconia tomentosa.

EUSTEPHIA, Cav. Ic. iii. 20 t. 238 (1794). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 732.
brevicuba, D. Dietr. Syn. Pl. ii. 1175 = Hippeastrum bicolor.
coccinea, *Cav. Ic.* iii. 20.—Am. austr.
cornuta, D. Dietr. Syn. Pl. ii. 1175 = Hippeastrum bicolor.
cyrtanthoides, D. Dietr. l. c. = Hippeastrum bicolor.
Herbertiana, D. Dietr. l. c.=Hippeastrum Herbertianum.
ignea, D. Dietr. l. c. = Hippeastrum bicolor.
Macleanica, Herb. in Bot. Mag. sub t. 3865, nomen = Hippeastrum advenum.

EUSTERALIS, Rafin. Fl. Tellur. ii. 95 (1836) = **Dysophylla**, Blume (Labiat.).
pumila, Rafin. l. c. = D. verticillata.

EUSTIGMA, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 313. *HAMAMELIDEAE*, Benth. & Hook. f. i. 668.
oblongifolium, *Gardn. & Champ. l. c.*—China.

EUSTOMA, Salisb. Parad. Lond. t. 34 (1806). *GENTIANEAE*, Benth. & Hook. f. ii. 812.
ARENBERGIA, Mart. & Gal. in Bull. Acad. Brux. xi. 1. (1844) 378.
ENSTOMA, A. Juss. in Orb. Dict. vi. 78 (1849).
URANANTHUS, Benth. Pl. Hartw. 46 (1840).
chironioides, Griseb. in DC. Prod. ix. 51 = silenifolium.
exaltatum, Salisb. Parad. Lond. sub t. 34 = silenifolium.
gracile, Engelm. in Fl. Calif. i. 621 = Russellianum.
lactewm, Liebm. Ind. Sem. Hort. Hamb. (1847), ex Sond. in E. Otto, Allg. Gartenz. iv. (1848) 369 = silenifolium.
maculatum, Benth. Pl. Hartw. 292 = Sabbatia maculata.
Russellianum, *G. Don, in Sweet, Hort. Brit.* ed. III. 478.—Texas ; Mexic.
silenifolium, *Salisb. Parad. Lond.* t. 34.—Am. bor.

EUSTREPHIA, D. Dietr. Syn. Pl. ii. 1175 (1840) = **Eustephia**, Cav. (Amaryll.).

EUSTREPHUS, R. Br. Prod. 281 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 768.
SPIRANTHERA, Rafin. Fl. Tellur. iv. 31 (1836).
angustifolius, R. Br. l. c. = latifolius.
Brownii, F. Muell. Fragm. vii. 78 = latifolius.
celebicus, D. Dietr. Syn. Pl. ii. 1117 = Dianella celebica.
javanicus, D. Dietr. l. c. = Dianella nemorosa.
latifolius, *R. Br. Prod.* 281.—Austral.
leucanthus, Hassk. Pl. Jav. Rar. 115 = latifolius.
timorensis, *Kidl. in Forbes, Nat. Wand. East. Archip.* 290.—Ins. Timor.
Watsonianus, Miq. in Lianaea, xviii. (1844) 84 = latifolius.

EUSTYLIS, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1847) 286 (Pl. Lindh. 27, 28) = **Nemastylis**, Nutt. (Irid.).
purpurea, Engelm. & Gray, l. c. = N. purpurea.

EUSTYLIS, Hook. f. Fl. N. Zel. t. 19 (ANISOTOME in textu, p. 90) = **Angelica**, Kiv. (Umbell.).
geniculata, Hook. f. l. c. = A. geniculata.

EUSTYLUS, Baker, in Journ. Linn. Soc. xvi. (1877) 102 = EUSTYLIS, Engelm. & Gray = **Nemastylis**, Nutt. (Irid.).

EUSYNAXIS, Griff. Notul. iv. 660. t. 608 (1854) = **Pyrenaria**, Blume (Ternstroem.).
barringtoniifolia, Griff. l. c. 660, 561 = P. barringtoniaefolia.

EUSYNETRA, Rafin. Fl. Tellur. ii. 57 (1836) = **Columnea**, Plum. (Gesneriac.).
bicolor, Rafin. l. c. = C. bicolor (in Addend.).

EUTACTA, Link, in Linnaea, xv. (1841) 548 = **Araucaria**, Juss. (Conif.).
Cookii, Carr. Conif. ed. II. 612 = A. Cookii.
Cunninghamii, Link, in Linnaea, xv. (1841) 545 = A. Cunninghamii.
excelsa, Link, l. c. 544 = A. excelsa.
humilis, Carr. Conif. ed. II. 616 = A. Cookii.
minor, Carr. l. c. 614 = A. Cookii.
Muelleri, Carr. in Rev. Hortic. (1866) 392 = A. Muelleri.
Panchenii, Carr. Conif. ed. II. 615 = A. Cookii.
Rulei, Carr. in Rev. Hortic. (1866) 279 = A. Rulei.
subulata, Carr. Conif. ed. II. 614 = A. Cookii.

EUTASSA, Salisb. in Trans. Linn. Soc. viii. (1807) 316 = **Araucaria**, Juss. (Conif.).
Cunninghamii, G. Don, in Sweet, Hort. Brit. ed. III. 623 = A. Cunninghami.
heterophylla, Salisb. in Trans. Linn. Soc. viii. (1807) 316 = A. excelsa.

EUTAXIA, R. Br. in Ait. Hort. Kew. ed. II. iii. 16 (1811). *LEGUMINOSAE*, Benth. & Hook. f. i. 471.
SCLEROTHAMNUS, R. Br. in Ait. Hort. Kew. ed. II. iii. 16 (1811).
Baxteri, Knowl. & Westc. Flor. Cab. i. (1837) 89 (in Eutaxia).—Austral.
cuneata, *Meisn. in Lehm. Pl. Preiss.* i. 65.—Austral.
densifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 1. 271.—Austral.
diffusa, F. Muell. Fragm. i. 7 = empetrifolia.
dillwynioides, *Meisn. in Lehm. Pl. Preiss.* i. 63.—Austral.
divaricata, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 270 = Dillwynia divaricata.
empetrifolia, *Schlecht. in Linnaea*, xx. (1847) 667.—Austral.
ericoides, Meisn. l. c. 64 = virgata.
leptophylla, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 268 = empetrifolia.
myrtifolia, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 16.—Austral.
obovata, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 271 = parvifolia.
parvifolia, *Benth. in Enum. Pl. Hueg.* 34.—Austral.
patula, F. Muell. in D. N. F. Dietr. Fl. Univ. N. S. t. 17 = Dillwynia patula.
punctata, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 272 = Mastrolobium reticulatum.
pungens, Sweet, Fl. Australt. t. 28 = Dillwynia pungens.
reticulata, Meisn. in Lehm. Pl. Preiss. i. 65 = Gastrolobium reticulatum.
sparsiflora, *F. Muell. in Hook. Kew Journ.* viii. (1856) 42 = Dillwynia patula.
Strangeana, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 270 = Pultenaea prostrata ?
uncinata, Turcz. l. c. 269 = Dillwynia patula.
virgata, *Benth. in Enum. Pl. Hueg.* 34.—Australi.

EUTELIA, R. Br. ex DC. Prod. iii. 76, in syn. (1828) = **Ammannia**, Linn. (Lythrar.).
ammannioides, R. Br. ex DC. l. c. = A. pentandra.

EUTELINE, Rafin. Sylva Tellur. 24 (1838) = **Genista**, Linn. (Legumin.).
decumbens, Rafin. l. c. = Cytisus decumbens.
germanica, Rafin. l. c. = G. germanica.
sagittalis, Rafin. l. c. = G. sagittalis.

EUTERELA, Rafin. Fl. Tellur. iv. 12 (1836) = **Dracocephalum**, Linn. (Arold.).
nigricans, Rafin. l. c. = D. polyphyllum.

EUTERPE, Gaertn. Fruct. i. 24 (1788). *PALMAE*, Benth. & Hook. f. iii. 896.
aculeata, Spreng. Syst. ii. 140 = Martinezia Aiphanes.
acuminata, *H. Wendl. in Kerch. Palm.* 244.—Hab. ?
andicola, *Brongn. ex Mart. Palmet. Orb.* 8. t. 2. f. 2 —Bolivia.
antioquensis, Linden, Illustr. Hortic. xxviii. (1881) 31 = nomen.
brevirugata, *Mart. Hist. Nat. Palm.* iii. 309.—Ind. occ.
Caatinga, Spruce, in Journ. Linn Soc. xi. (1869) 137 = E. Catinga.
caribaea, Spreng. Syst. ii. 140 = Oreodoxa oleracea.
Catinga, *Wallace, Palm-Trees, Amaz.* 27. t. viii.—Bras
decurrens, *H. Wendl. in Kerch. Palm.* 244, nomen.—Hab. ?
disticha, *H. Wendl. ex Linden, Cat.* n. 23 (1868).—N. Granat.
edulis, *Mart. Hist. Nat. Palm.* iii. 33. t. 32.—Am trop.
elegans, *Linden, Illustr. Hortic.* xxviii. (1881) 31.—N. Granat.
emiformis, *Mart. Hist. Nat. Palm.* ii. 32. t. 31.—Peruv.
flamentosa, Blume, ex Kunth, Enum. Pl. iii. 185 = Oncosperma filamentosa.
globosa, Gaertn. Fruct. i. 24 = edulis.
gracilis, *Linden, Cat.* n. 17 (1862).—Bras.
Haenkeana, *Brongn. ex Mart. Palmet. Orb.* 9. t. 2. f. 3.—Bolivia.
Karsteniana, *Engel, in Linnaea*, xxxiii. (1864–65) 670. —N. Granat.
longepetiolata, *Oerst. in Kjoeb. Vidensh. Meddel.* (1858) 31.—Costa Rica.
longevaginata, *Mart. Palmet. Orb.* 11. t. 12. f. 1.—Bolivia.
longibracteata, *Barb.-Rodr. Enum. Palm. Nov.* 17.—Bras.
macrospadix, *Oerst. in Kjoeb. Vidensh. Meddel.* (1858) 31.—Nicaragua.
Manselle, *Griseb. & H. Wendl. ex Griseb. in Mem Acad. Am.* N. S. viii. (1862) 530.—Ind. occ.
Manicki, Lodd. ex H. Wendl. in Kerch. Palm. 240 (Guild ?).—Hab. ?
Mensiesii, Pritz. Ic. Ind. 450, sphalm. = *Euterea Mensiesii*.
mollissima, Barb.-Rodrig. Enum. Palm. Nov. 16 = E. Catinga.
mollissima, Spruce, in Journ. Linn. Soc. xi. (1869) 139 = precatoria.
montana, *R. Grah. in Bot. Mag.* t. 3874 (? Prestoeae sp.).—Ins. Grenat.
oleracea, *Engel, in Linnaea*, xxxiii. (1864–65) 671.—N. Granat.
oleracea, *Mart. Hist. Nat. Palm.* ii. 29. tt. 29, 30.—Bras.
pisifera, *Gaertn. Fruct.* i. 24.—Hab. ?
Prapa, *Mart. Gen. Palm.* (1824) 24 ; *Spreng. Syst.* ii. 142.—Venezuela.
precatoria, *Mart. Palmet. Orb.* 10. t. 8. f. 2.—Bolivia.
purpurea, *Engel, in Linnaea*, xxxiii. (1864–65) 669.—N. Granat.
punctata, *Linden, Illustr. Hortic.* xxviii. (1881) 32. —Bras.
vinifera, Mart. Hist. Nat. Palm. i. t. s. ii. f. 18, 19 = Gaussia vinifera.
Zamora, Hort. Linden, ex H. Wendl. in Kerch. Palm. 245, nomen (Guild ?).—Hab. ?

EUTERPAS, A. Gray, in Proc. Am. Acad. xv. (1880) 89. *COMPOSITAE*.
Palmeri, *A. Gray, l. c.* 40.—Mexic.

EUTHALES, R. Br. Prod. 279 (1810) = **Vellela**, Sm. (Goodenov.).
filiformis, De Vriese, in Lehm. Pl. Preiss. i. 414 = V. trinervis.
macrophylla, Lindl. Bot. Reg. (1840) Misc. 54 = V. macrophylla.
pilosella, De Vriese, in Lehm. Pl. Preiss. i. 414 = V. trinervis ?
trinervis, R. Br. Prod. 579 = V. trinervis.

EUTHALES, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 307 (1817) = **Tovomita**, Aubl. (Guttif.).
fructipendula, Steud. Nom. ed. II. i. 619 = T. fructipendula.
peruviana, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 307 = T. fructipendula.

EUTHALIA, Rupr. Fl. Cauc. 230 (1869) = **Arenaria**, Rupp. (Caryophyll.).
rotundifolia, Rupr. l. c. = A. rotundifolia.
serpyllifolia, Rupr. l. c. = A. serpyllifolia.

EUTHALIS, Banks & Soland. ex Hook. f. Fl. Antarct. 264 (1847) = **Maytenus**, Molina (Celastrin.).
lucida, Banks & Soland. ex Hook. f. l. c. = M. magellanicus.

EUTHAMIA, Ell. Sketch, ii. 301, in syn. (1824) = **Solidago**, Vaill. (Compos.).
graminifolia, Ell. l. c. = S. lanceolata.
occidentalis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 326 = S. occidentalis.
tenuifolia, Ell. Sketch. ii. 302, in syn. = S. tenuifolia.

EUTREMIS, Jack, in Malay. Misc. i. (1820) v. 15.
OCHNACEÆ, Benth. & Hook. f. i. 310.
elegantissima, Wall. in Roxb. Fl Ind. ed. Carey, ii. 305 = Gomphia sumatrana!
leucocarpa, *Jack, in Malay. Misc.* i. (1820) v. 16.—Malaya.
minor, *Jack, l. c.* 18.—Malaya.
obtusifolia, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 163.—Borneo.
pulcherrima, Wall. Cat. n. 2518 = Gomphia sumatrana.
robusta, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 163.—Borneo.

EUTHODON, Griff. Notul. iv. 84, t. 458 (1854) = **Pottsia**, Hook. & Arn. (Apocyn.).
paniculata, Griff. l. c. = P. cantonensis.

EUTHRIXIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 267 = **Chaetanthera**, Ruiz & Pav. (Compos.).
affinis, D. Don, in Phil. Mag. (1829) 391; Gunl. Arch. Bot. ii. (1838) 467 = C. linearis.
salsoloides, D. Don, in Trans. Linn. Soc. xvi. (1830) 259 = C. linifolia.

EUTHYRA, Salisb. Gen. Pl. Fragm. 61 (1866) = **Paris**, Linn. (Liliac.).

EUTHYSTACHYS, A. DC. in DC. Prod. xii. 606 (1848). *VERBENACEÆ*, Benth. & Hook. f. ii. 1138.
abbreviata, *A. DC. l. c.*—Afr. austr.

EUTMON, Rafin. Atl. Journ. 177 (1832) = **Talinum**! Adans. (Portulac.).
napiforme, Rafin. l. c. = T. teretifolium!

EUTOCA, R. Br. in App. Frankl. Voy. 764. t. 27 (1823) = **Phacelia**, Juss. (Hydrophyll.).
acaulis, Mart. & Gal. in Bull. Acad. Brux. xii. ii. (1845) 276.—Mexic.
albiflora, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 156 = P. viscida.
Andreuxii, A. DC. in DC. Prod. ix. 294 = P. pimpinelloides.
arctioides, Hook. & Arn. Bot. Beech. Voy. 374 = Cosmibuena arctioides.
brachyloba, Benth. in Trans. Linn. Soc. xvii. (1837) 277 = P. brachyloba.
burrifolia, Spreng. Syst. i. 569.—Mexic.
congesta, Dougl. ex Lehm. Pugill. ii. 18.—Am. bor. occ.
Cumingii, Benth. in Trans. Linn. Soc. xvii. (1837) 276.—Chili.
divaricata, Benth. l. c. 278 = P. divaricata.
Douglasii, Benth. l. c. 276 = P. Douglasii.
echioides, Dougl. ex A. DC. in DC. Prod. ix. 294 = P. Menziesii.
Franklinii, R. Br. App. Frankl. Voy. t. 27; et in Trans. Linn. Soc. xvii. (1837) 277 = P. Franklinii.
frigida, Phil. Fl. Atac. 37.—Chili.
glabra, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 254.—Am. bor.
glandulosa, Nutt. in Hook. Kew Journ. iii. (1851) 293 = P. glandulosa, *Nutt.*
glomerata, Dougl. ex A. DC. in DC. Prod. ix. 294 = P. Menziesii.
gracilis, Mart. & Gal. in Bull. Acad. Brux. xii. ii. (1845) 276.—Mexic.
EUTOCA :—
grandiflora, Benth. in Trans. Linn. Soc. xvii. (1837) 278 = P. grandiflora.
hetrophylla, Torr. in Stansbury, Exped. Utah, 398 = P. Menziesii.
hirsuta, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 256 = P. parviflora !
loasaefolia, Benth. in Trans. Linn. Soc. xvii. (1837) 277 = P. loasaefolia.
lutea, Hook. & Arn. Bot. Beech. Voy. Suppl. 374 = Emmenanthe lutea.
Menziesii, R. Br. in App. Frankl. Voy. 764 = P. Menziesii.
mexicana, Benth. in Trans. Linn. Soc. xvii. (1837) 277 = P. pimpinelloides.
mexicana, Hort. ex C. A. Mey. in Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 74 = P. divaricata.
multiflora, Dougl. ex Lehm. Pugill. ii. 19 = P. Menziesii.
Origerssana, Heer, in Regel, Gartenfl. (1861) 309.—Mexic.
parviflora, R. Br. App. Frankl. Voy. ii. 51 = P. parviflora.
patuliflora, Engelm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1845) 258 = P. patuliflora.
pedunculosa, Phil. in Anal. Univ. Chil. (1884) reimpr. 7.—Chili.
phaceloides, Benth. in Trans. Linn. Soc. xvii. (1837) 279 = P. circinatiformis.
pimpinelloides, Spreng. Syst. i. 569 = P. pimpinelloides.
pinnatifida, Phil. in Anal. Univ. Chil. (1870) ii. 189.—Chili.
pusilla, Lehm. Pugill. ii. 18 = P. sericea.
pusilla, Phil. in Linnaea, xxxiii. (1864–65) 186.—Chili.
sericea, R. Grah. in Edinb. N. Phil. Journ. (Apr.-June 1830) 372 = P. sericea.
speciosa, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 158 = P. grandiflora.
stricti flora, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 258 = P. stricti flora.
viscida, Benth. in Bot. Reg. t. 1808 = P. viscida.
vuscosa, Hook. Bot Mag. t. 3572 = præc.
Wrangeliana, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 37 = P. divaricata.

EUTRALIA, Rafin. Fl. Tellur. iii. 46 (1836) = **Rumex**, Tourn. (Polygon.).
luxurians, Rafin. l. c. = R. luxurians.

EUTREMA, R. Br. in Parry, 1st Voy. App. 267. t. A (1824). *CRUCIFERÆ*, Benth. & Hook. f. i. 78.
alpestre, *Ledeb. Fl. Ross.* i. 198.—Sibir.
arenicola, Richards. ex Hook Fl. Bor. Am. i. 67 = Parrya arenicola.
Boneandii, Spreng. Syst. ii. 880 = *Eudema nubigena*.
cordifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 283.—Sibir. bascal.
Edwardsii, *R. Br. in Parry, 1st Voy. App.* 9. t. A.—Am. arct.
hederaefolia, *Franch. & Sav. Enum. Pl. Jap.* ii. 288.—Japon.
himalaicum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 164.—Reg. Himal.
Humboldtii, *Spreng. Syst.* ii. 880.—Ecuador.
integrifolium, *Bunge, Del. Sem. Hort. Dorp.* (1839) 8.—Hab.!
intermedium, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 283 = Edwardsii.
labradoricum, *Turcz. l. c.* xxvii. (1854) ii. 305.—Am. arct.
parviflorum, Turcz. l. c. xv. (1842) 283 = Edwardsii.
pilliferum, Turcz. l. c. 283 = Parrya microcarpa.
primulaefolium, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 164.—Reg. Himal.
Roxeri, Spreng. Syst. ii. 880 = Cochlearia officinalis.
septigerum, *Bunge, Verz. Suppl. Fl. Alt.* 78.—Sibir.
Wsanbi, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 283.—Japon.

EUTRIANA, Trin. Fund. Agrost. 161 (1820) = **Boutelous**, Lag. (Gramin.).
abyssinica, K. Br. ex Fresen. in Mus. Senckenb. ii. (1837) 142.—Abyss.
EUTRIANA :—
affinis, Hook. f. in Trans. Linn. Soc. xx. (1851) 174 = Bouteloua racemosa.
Antillarum, Steud. Syn. Pl. Gram. 217.—Ind. occ.
aristidoides, Trin. Gram. Unifl. 242; Kunth, Rév. Gram. i. 96 = B. aristidoides.
barbata, Kunth, l. c. 96 = B. barbata.
bromoides, Trin. Fund. Agrost. 161; et in Gram. Unifl. 241 = B. bromoides.
chloroides, Kunth. Enum. Pl. i. 280.—Mexic.
cristata, Kunth, l. c. 281.—Mexic.
cristata, Trin. Gram. Unifl. 241 = B. cristata.
curtipendula, Trin. Fund. Agrost. 161 = B. racemosa.
Enteropogon, Steud. Syn. Pl. Gram. 216 = Enteropogon mellicoides.
gracilis, Hook. f. in Trans. Linn. Soc. xx. (1851) 175.—Ins. Galapag.
gracilis, Trin. Gram. Unifl. 240 = B. oligostachya.
Heterostega, Trin. l. c. 242 = B. bromoides.
hirsuta, Kunth, Enum. Pl. i. 280.—Peruv.
hirta, Trin. Gram. Unifl. 240 = B. hirsuta.
humilis, Trin. l. c 239 = B. prostrata.
juncea, Trin. l. c. 238 = B. juncifolia.
juncifolia, Kunth, Rév. Gram. i. 95 = B. juncifolia.
Karwinskiana, Trin. ex Steud. Nom. ed. II. i. 620 (Quod !).—Mexic.
Lagascæ, Kunth, Rév. Gram. i. 95 = B. juncifolia.
Ledebourii, Trin. Gram. Unifl. 238.—Louisiana; Ins. S. Doming.
multiseta, Nees, Agrost. Bras. 413 = R. multiseta.
oligostachya, Kunth, Rév. Gram. i. 96 = B. oligostachya.
Pentarrhaphis, Trin. Gram Unifl. 239.—Mexic.
pilosa, Hook. f. in Trans. Linn. Soc. xx. (1851) 175.—Ins. Galapag.
Polydon, Trin. Gram. Unifl. 242 = B. distichia.
repens, Trin. l. c. 241; Kunth, Rév. Gram. i. 95 = B. bromoides.
tenuis, Trin. l. c. 240 = B. tenuis.
Trianæa, Trin. l. c. 239 = B. Trinthera.
villosa, Steud. Nom. ed. II. i. 620.—Bras.

EUTROPIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 196 = **Croton**, Linn. (Euphorbiac..
brasiliensis, Klotzsch, l. c. = C. brasiliensis.
obovata, Klotzsch, l. c. = C. brasiliensis.

EUTROPIS, Falc. in Proc. Linn. Soc. i. (1839) 15 = **Pentatropis**, R. Br. (P. microphylla) (Asclepiad.).

EUXENIA, Cham. in Nees, Hor. Phys. Berol. 75. t. 16 (1820) = **Podanthus**, Lag. (Compos.).
grata, Cham. l. c. = P. ovatifolius.
Mitiqui, DC. Prod. v. 501 = P. Mitiqui.
radiata, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 7 (= *Gymnopus euxenioides*).—Bras.

EUXOLUS, Rafin. Fl. Tellur. iii. 42 (1836) = **Amaranthus**, Linn.
Blitum, Gren. Fl. Ch. Jurass. 652 = A. Blitum.
caudatus, Moq. in DC. Prod. xiii. ii. 274 = A. viridis.
crassipes, Hieron. in Bol. Acad. Nac. Córdova, iv. (1881) 18 = A. polygonoides.
crispus, Less. in Bull. Soc. Bot. Fr. vi. (1859) 656.—Am. bor.
deflexus, Rafin. Fl. Tellur. iii. 42 = A. deflexus.
emarginatus, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1851) 18 = A. viridis.
enervis, F. Muell. Fragm. i. 140 = A. enervis.
interruptus, Moq. in DC. Prod. xiii. ii. 275 = A. interruptus.
lineatus, Moq. l. c. 276 = A. interruptus.
lividus, Moq. l. c. 273 = A. Blitum.
macrocarpus, F. Muell. Census, 29 = A. macrocarpus.
Mitchelii, F. Muell. in Giles, Geogr. Trav. Centr. Austr. 214 = A. Mitchelli.
muricatus, Moq. in DC. Prod. xiii. ii. 276 = A. muricatus.
oleraceus, Moq. l. c. 278 = A. Blitum.
paraguayensis, Parodi, in Anal. Soc. Cient. Argent. v. (1878) 276; Contrib. 31.—Paraguay.
polygamus, Moq. in DC. Prod. xiii. ii. 272 = A. polygamus.
polygonoides, Thw. Enum. Pl. Zeyl. 248 = A. polygamus.
polystachyus, Mlq. Fl. Ind. Bat. i. i. 1036 = A. polystachyus.

EUXOLUS—
pumilus, Chapm. Fl. S. U. St. 381 = Amarantus pumilus.
rhombeus, Moq. in DC. Prod. xiii. ii. 275 = A. interruptus.
scleranthoides, Anders. in Vet. Akad. Handl. Stockh. 1858 (1855) 163.—Ins Galapag.
undulatus, Moq. in DC. Prod. xiii. ii. 272 = A. interruptus.
viride, Moq. l. c. 272 = A. viridis.

EUZOMODENDRON, Coss. Not. Pl. Espagne, 144 (1852); et in Webb, Otia Hisp. 47. t. 43 (1858). *CRUCIFERAE*, Benth. & Hook. f. i. 85. Bourgeanum, *Coss. Il. cc.*—Hispan.

EUZOMUM, Link, Enum. Hort. Berol. ii. 175 (err. typ. 174) (1822) = **Bruca**, Tourn. (Crucif.).
acre, Webb, Fragm. Fl. Aethiop. 17 = Diplotaxis Sieberi.
crucoides, Spach, Hist. Vég. Phan. vi. 382 = Diplotaxis crucoides.
hespéridiforme, Spach, ex Steud. Nom. ed. II. i. 620 = Diplotaxis Sieberi.
hispidum, Link, Handb. ii. 319 = Eruca sativa.
sativum, Link, Enum. Hort. Berol. ii. 175 (err. typ. 174) = Eruca sativa.
vesicarium, Link, Handb. ii. 319 = Eruca vesicaria.

EVACIDIUM, Pomel, Nouv. Mat. Fl. Atl. 287 (1875). *COMPOSITAE*.
atlanticum, *Pomel, l. c.* 288.—Afr. bor.

EVACOPSIS, Pomel, Nouv. Mat. Fl. Atl. 41 (1874) = **Evax**, Gaertn. (Compos.).
discolor, Pomel, l. c. 43 = Evax discolor.
exigua, Pomel, l. c. 41 = Evax exigua.
Heldreichii, Pomel, l. c. 42 = Evax Heldreichii.
montana, Pomel, l. c.—Afr. bor.
polycephala, Pomel, l. c. 43.—Afr. bor.

EVAIEZOA, Rafin. Fl. Tellur. ii. 70 (1836) = **Saxifraga**, Tourn.
axoides, Rafin. l. c. = S. axoides.
aygiifolia, Rafin. l. c. = S. aygiifolia.
androsacea, Rafin. l. c. 71 = S. androsacea.
bifera, Rafin. l. c. 70 = S. bifera.
bulbifera, Rafin. l. c. = S. bulbifera.
Burseriana, Rafin. l. c. 71 = S. Burseriana.
callosa, Rafin. l. c. = S. longifolia.
cespitosa, Rafin. l. c. 70 = S. cespitosa.
cuneata, Rafin. l. c. = S. cuneata.
diapensoides, Rafin. l. c. 71 = S. diapensoides.
erosa, Rafin. l. c. = S. erosa.
Eskoltzi, Rafin. l. c. = S. serpyllifolia.
exarata, Rafin. l. c. = S. exarata.
exilis, Rafin. l. c. = S. exilis.
geranioides, Rafin. l. c. 70 = S. geranioides.
granulata, Rafin. l. c. = S. granulata.
hieraciifolia, Rafin. l. c. 71 = S. hieraciifolia.
hirta, Rafin. l. c. 70 = S. hypnoides.
ladanifera, Rafin. l. c. = S. pedatifida.
Laurentiana, Rafin. l. c. 71 = S. rivularis.
mutata, Rafin. l. c. = S. mutata.
mutans, Rafin. l. c. = Romanoffia unalaschcensis, *opposititolia*, Rafin. l. c. 70 = S. oppositifolia.
palmata, Rafin. l. c. = S. geranioides.
pedatifida, Rafin. l. c. = S. pedatifida.
pensylvanica, Rafin. l. c. 71 = S. pensylvanica.
punctata, Rafin. l. c. 70 = S. punctata.
pungens, Rafin. l. c. 71 = S. Vandellii.
retusa, Rafin. l. c. 70 = S. retusa.
rivularis, Rafin. l. c. = S. rivularis.
serpyllifolia, Rafin. l. c. 71 = S. serpyllifolia.
sileniflora, Rafin. l. c. = S. sileniflora.
trichodes, Rafin. l. c. = S. sedoides.
trifurcata, Rafin. l. c. 70 = S. trifurcata.
venosa, Rafin. l. c. 71 = S. venosa.

EVALLARIA, Neck. Elem. iii. 189 (1790) = **Polygonatum**, Adans. (Liliac.).
polygonata, Neck. l. c. = P. officinale.
verticillata, Neck. l. c. = P. verticillatum.

EVALTHE, Rafin. Fl. Tellur. iii. 77 (1836) = **Chiromia**, Linn. (Gentian.).
jasminoides, Rafin. l. c. = C. jasminoides.

EVANDRA, R. Br. Prod. 239 (1810) *CYPERACEAE*, Benth. & Hook. f. iii. 1067.
aristata, *R. Br. l. c.*—Austral.
pascifiora, *R. Br. l. c.*—Austral.

EVANESCA, Rafin. Sylva Tellur. 105 (1838) = **Myrtus**, Tourn.
crassifolia, Rafin. l. c. = M. dioica.

EVANSIA, Salisb. in Trans. Hort. Soc. i. (1812) 303 = **Iris**, Tourn.
chinensis, Salisb. l. c. = I. japonica.
cristata, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 373 = I. cristata.
dichotoma, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 302 = I. dichotoma.
fimbriata, Decne. l. c. = I. japonica.
gracilipes, Klatt, in Abh. Naturf. Ges. Halle, xii (1882) 373 = I. gracilipes.
japonica, Klatt, l. c. = I. japonica.
lacustres, Klatt, l. c. = I. lacustris.
nepalensis, Klatt, l. c. = I. nepalense.
speculatrix, Klatt, l. c. = I. speculatrix.
syrtica, Klatt, l. c. = I. Sisyrinchium.
tectorum, Klatt, l. c. = I. tectorum.
vespertina, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 302. —Mongol.

EVAX, Gaertn. Fruct. ii. 393. t. 165 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 296.
CALYMMANDRA, Torr. & Gray, Fl. N. Am. ii. 262 (1841).
DIAPERIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii (1841) 387.
EVACOPSIS, Pomel, Nouv. Mat. Fl. Atl. 41 (1874).
FILAGINOPSIS, Torr. & Gray, Fl. N. Am. ii. 262 (1841).
FILAGO, Loefl. in Linn. Gen. ed. V. 397 (1754).
HESPERBVAX, A. Gray, in Proc. Am. Acad. vii. (1868) 356, in obs.
anatolica, *Boiss. & Heldr. Diagn.* Ser. I. xi. 2.—As. Min.
argentea, *Pomel, Nouv. Mat. Fl. Atl.* 41.—Alger.
asteriscifora, *Pers. Syn.* ii. 422.—Reg. Mediterr.
candida, *A. Gray, Syn. Fl. N. Am.* i. ii. 330.—Texas.
carpetana, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 69.—Hispan.
caulescens, *A. Gray, in Pacif. Rail. Rep.* iv. 101.—Calif.
contracta, *Boiss. Diagn.* Ser. I. xi. 3.—Syria; Persia.
Crocidion, *Pomel, Nouv. Mat. Fl. Atl.* 40.—Alger.
desertorum, *Pomel, l. c.*—Alger.
discolor, *DC. Prod.* v. 459.—Sicilia.
ericoides, Schrank, in Denkschr. Akad. Muench. viii. 1821–22 (1824) 172 = Helichrysum teretifolium.
eriosphaera, *Boiss. & Heldr. Diagn.* Ser. I. xi. 3.—As. Min.
exigua, *DC. Prod.* v. 458.—Archip. Graec. ; Persia.
filaginoides, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 279.—Songaria.
Funkii, Sch. Bip. ex Nym. Consp. 395 = asteriscifera.
Heldreichii, *Parl. in Giorn. Tosc. Sc. Med. Fis. e Nat.* i. (1840) 138.—Reg. Mediterr. occ.
indica, Roxb. Hort. Ben. ex Wall. Cat. n. 2956 = Gnaphallum pulvinatum.
involucrata, Schrank, in Denkschr. Akad. Muench. viii. 1821–22 (1824) 171 = Petalacte coronata.
laxiscapa, Lange, ex Willk. & Lange, Prod. Fl. Hisp. ii. 65 = carpetana.
marocica, DC. Prod. v. 459 = Filago marocica.
microcephalos, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 65.—Hispan.
multicaulis, *DC. Prod.* v. 459.—Mexic.
nevadensis, *Boiss. Fl. Orient.* iii. 245, in nota.—Hispan.
palaestina, Boiss. Diagn. Ser. I. xi. 2 = anatolica.
perpusilla, *Boiss. & Heldr. l. c.* Ser. II. iv. 18.—Graecia.
pruiflora, *Nutt. ex DC. Prod.* v. 459.—Am. bor. occ.
pygmaea, Brot. Fl. Lusit. i. 363 = umbellata.
rotundata, *Moris, in Atti Terza Ruin. Sc. Ital.* 481.—Corsic. ; Sardin.
spathulata, Pers. Syn. ii. 422 = Gnaphalium bracteatum.
tenuifolia, *Guss. Fl. Sic. Syn.* ii. 460.—Sicil.
umbellata, *Gaertn. Fruct.* ii. 393.—Reg. Mediterr.
verna, Rafin. Act. Journ. 176 = multicaulis.

EVEA, Aubl. Pl. Gui. i. 100. t. 39 (1775) = **Cephalis**, Sw. (Rubiac.).
brasiliensis, Cham. & Schlecht. in Linnaea, ix. (1834) 237 = Salemannia nitida.
guianensis, Aubl. Pl. Gui. i. 100 = C. Evea.
palacea, A. Rich. in Mém. Soc. Hist. Nat. Par. v (1834) 198.—Guian.

EVELTRIA, Rafin. Fl. Tellur. iv. 80 (1836) = **Orthosanthus**, Sweet (Irid.).
multiflora, Rafin. l. c. = O. multiflorus.

EVELEVNA, Steud. Nom. ed. II. i. 620 (1840) = seq.

EVELYNA, Poepp. & Endl. Nov. Gen. et Sp. i. 32 t. 56 (1835) = **Illeanthus**, Presl (Orchid.).
amethystina, Reichb f. & Warsz. in Bonplandia, ii (1854) 113 = Ell. amethystinus.
argophyllostachys, Reichb. f. l. c. 21 = Ell. argophyllostachys.
aurantiaca, Lindl. in Benth. Pl. Hartw. 149 = Ell aurantiacus.
aurea, Poepp. & Endl. Nov. Gen. et Sp. 33. t. 57 = Ell. aureus.
Bonplandii, Reichb. f. in Bot. Zeit. x. (1852) 711 = Ell. Bonplandi.
bractescens, Lindl. Orch. Linden. 10 = Ell. bractescens.
brasiliensis, Lindl. in Hook. Lond. Journ. Bot. ii (1843) 661 = Ell. brasiliensis.
capitata, Poepp. & Endl. Nov. Gen. et Sp. i. 32. t. 56 = Ell. capitatus.
Caravata, Lindl. in Fl. des Serres, xiv. (1861) 85. t. 1497 = Ell. Caravata
casapensis, Reichb. f. in Bot. Zeit. (1852) 709 = Ell casapensas.
cephalophora, Reichb. f. l. c. = Ell. cephalophorus.
columnaris, Lindl. Orch. Linden. 11. = Ell. columnaris
confera, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 118 = Ell. confer.
corrifolia, Reichb. f. ex Linden, Illustr. Hortic xxix. (1882) 51 = Ell. corrifolius.
crinipes, Reichb. f. Otia Bot. Hamb. 92, in obs. = Ell crinipes.
cynarocephala, Reichb. f. in Bonplandia, iv. (1856) 216 = Ell. cynarocephala.
discolor, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 113 = Ell. discolor.
ensata, Lindl. Orch. Linden. 11. = Ell. ensatus.
flavescens, Lindl l. c. = Ell. flavescens.
furfuracea, Lindl. l. c. 12 = Ell. furfuraceus.
gracilis, Reichb. f. in Linnaea, xxii. (1849) 842 = Ell. gracilis.
graminifolia, Poepp. & Endl. ex Linden, Illustr. Hortic. xxix. (1882) 15.—Peruv.
haematoxantha, Reichb. f. ex Linden, l. c. = Ell. haematoxanthus.
Hallii, Reichb. f. in Bot. Zeit. x. (1852) 711 = Ell. Hallii.
Hooheriana, Rodrig. Orch. Nov. ii. 166.—Bras.
hymenophora, Reichb. f. in Bot. Zeit. x. (1852) 710 = Ell. hymenophorus.
hermesina, Lindl. Orch. Linden. 11 = Ell. hermesinus.
lepida, Reichb. f. in Hamb. Gartenz. xv. (1859) 56 = Ell. lepidus.
Lindenii, Reichb. f. ex Linden, Illustr. Hortic. xxix (1882) 15 = Ell. Lindenii.
lupulina, Lindl. Orch. Linden. 11 = Ell. lupulinus.
maculata, Lindl. in Benth. Pl. Hartw. 149 = Ell. maculatus.
Myrosmatis, Reichb. f. in Bot. Zeit. x. (1852) 708 = Ell. Myrosmatis.
oeconomica, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 113.—N. Granat.
oligantha, Poepp. & Endl. Nov. Gen. et Sp. i. 33. t. 57 = Ell. oliganthus.
purpurea, Reichb. f. in Bot. Zeit. x. (1852) 710 = Ell. purpureus.
rhodolepis, Reichb. f. l. c. 711 = Ell. rhodolepis.
robusta, Reichb. f. l. c. 708 = Ell. robustus
Rutaii, Reichb. f. in Linnaea, xxii. (1849) 842 = Ell. Rutaii.
simplex, Reichb. f. ex Linden, Illustr. Hortic. xxix. (1882) 15.—N. Granat.
strobiliflora, Poepp. & Endl. Nov. Gen. et Sp. 33 = Ell. strobiliflor.
Wagneri, Reichb. f. in Bonplandia, ii. (1854) 21 = Ell. Wagneri.

EVELYNA, Raf. Sylva Tellur. 164 (1838) = **Lindera**, Thanb. (Laurin.).

EVERION, Raf. Sylva Tellur. 124 (1838) = **Alternanthera**, Forsk. (Amarant.).
interrupta, Raf. l. c. = A. procumbens.

EVERSMANNIA, Bunge, in Goebel, Reise, ii. 267. t. 6 (1838). *LEGUMINOSAE*, Benth. & Hook. f. i. 510.
astragaloides, *Regel & Schmalh. Pl. Nov. Fedsch.* 18. —Turkestan.
hedysaroides, *Bunge, in Goebel, Reise* ii. 267. t. 6.— Reg. Casp.

EVIA, Comm. ex Juss. Gen. 373 (1789) = **Spondias**, Linn. (Anacard.).
acida, Blume, Mus. Bot. Lugd. Bat. i. 234 = S. dulcis.
amara, Comm. ex Blume, l. c. = S. mangifera.
dulcis, Comm. ex Blume, l. c. 233 = S. dulcis.

EVODEA, Kunth, Syn. Pl. iii. 327 (1824) = seq.

EVODIA, Forst. Char. Gen. 18. t. 7 (1776) (Euodia). *RUTACEAE*, Benth. & Hook. f. i. 396.
ATITARA, Mazugr. ex Juss. in Dict. Sc. Nat. iii. 277 (1816).
BOYMIA, A. Juss. in Mém. Mus. Par. xii. (1825) 507 (Mem. Rut. 134).
LEPTA, Lour. Fl. Cochinch. 92 (1790).
MEGABOTRYA, Hance, in Walp. Ann ii. 259 (1851).
PHILAGONIA, Blume, Cat. Gew. Buitenz. 20 (1823); Bijdr. 250 (1825).
TETRADIUM, Lour. Fl. Cochinch. 91 (1790).
accedens, *Blume, Bijdr.* 246.—Java.
acronychioides, *F. Muell. Fragm.* iv. 117.—Austral.
alata, *F. Muell. l. c.* vii. 142.—Austral
alba, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 166. —Borneo.
aromatica, Blume, Bijdr. 246 = glabra.
Aubertin, *Benth. & Hook. f. Gen.* i. 396.—Ins. Borbon.
Baudouini, *Baill. Adansonia*, x. (1871–72) 326.—N. Caled.
Bintoco, Blanco, Fl. Filip. ed. II. 80 = latifolia.
Bonwickii, *F. Muell. Fragm.* v. 56.—Austral.
bracteata, *Nadeaud, Enum. Pl. Tahiti*, 76.—Ins. Tahiti.
Brombyana, F. Muell. Fragm. v. 4 = Brombya platynema.
celastracea, *Baker, in Journ. Linn. Soc.* xx. (1883) 117.—Madag.
Chapelieri, *Baill. Adansonia*, x. (1871–72) 326.— Madag.
contermina, F. Muell. Fragm. vii. 144 = Melicope contermina.
Cunninghamii, F. Muell. l. c. iii. 2 = Medicosma Cunninghamii.
densiflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 117.—Madag.
divaricata, Poepp. ex Engl. in Mart. Fl. Bras. xii. 11. 279 = Protium divaricatum.
drupacea, *Labill. Sert. Austr. Caled.* 73. t. 75.—N. Caled.
elaeodendroides, *Baker, Fl. Maurit.* 39.—Ins. Mascar.
Elleryana, *F. Muell. Fragm.* v. 4.—Austral.
erythrococca, F. Muell. l. c. i. 26 = Melicope erythrococca.
euneura, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 243.— Sumatra.
Farquhariana, *F. Muell. Fragm.* ix. 101.—Austral.
febrifuga, St. Hil. in Bull. Soc. Philom. (1822) 129 = Esenbeckia febrifuga.
fraxinifolia, *Hook. f. Fl. Brit. Ind.* i. 490.—Reg. Himal., Malaya.
glabra, *Blume, Bijdr.* 245.—Java.
glauca, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 23 = melliodella.
gracilis, Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 43 = triphylla.
haplophylla, *F. Muell. Fragm.* v. 179.—Austral.
hortensis, *Forst. Char. Gen.* 18. t. 7.—Ins. N. Hebrid.
incerta, Blume, Bijdr. 245 = Acronychia Minahassae.
Lamarckiana, Benth. Fl. Hongk. 59 = triphylla.
laxiuscula, *Baill. Adansonia*, xi. (1874) 179.—N. Caled.
latifolia, *DC. Prod.* i. 724.—Malaya.
litterosella, *Endl. Prod. Fl. Norf.* 36. n. 146.—Ins. Norfolk.

EVODIA:—
longifolia, *A. Rich. Bot. Voy. Astrol.* ii. 61.—N. Guinea.
lucida, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 244.— Sumatra.
macrophylla, Blume, Bijdr. 246 = accedens.
macrophylla, Wall. Cat. n. 1230 = Roxburghiana.
madagascariensis, *Baker, in Journ. Bot.* xx. (1882) 48. —Madag.
Maruwahong, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 244 = Roxburghiana.
meliaefolia, *Benth. Fl. Hongk.* 58.—China.
micrococca, *F. Muell. Fragm.* i. 144.—Austral.
Minahassae, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 255 = Acronychia Minahassae.
neurococca, F. Muell. Fragm. i. 28 = Melicope neurococca.
obtusifolia, *DC. Prod.* i. 724.—Ins. Mascar.
octandra, F. Muell. Fragm. ii. 102 = Melicope australasica.
pedicuiloss, *Hook. f. Fl. Brit. Ind.* i. 489.— Singapor.
pentacocca, F. Muell. Fragm. iii. 41 = Bosistoa sapindiformis.
polybotrya, *C. Moore & F. Muell. in F. Muell. l. c.* vii. 143.—Austral.
pomaderrioila, *Baill. Adansonia*, xi. (1875) 306.— N. Caled.
ramiflora, *A. Gray, in Mem. Am. Acad.* N. S. vi (1858–59) 383.—Japon.
roxburghiana, *Hook. f. Fl. Brit. Ind* i. 488.—Malaya. Roxburghiana, *Benth. Fl. Hongk.* 59.—As. trop.
rutaecarpa, *Hook. f. & Thoms. in Fl. Brit Ind.* i. 490 —Reg. Himal.; Chuna; Japon.
saxicocca, *Baill. Adansonia*, xi. (1875) 301.—N. Caled.
speciosa, *Reichb. f. & Zoll. ex Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 255.—Java.
suaveolens, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 11.—N. Guinea.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 77 —Ins. Tahiti.
triphylla, Bedd. Fl. Sylv. Anal. Gen. xli. t. 6. f. 2 = Roxburghiana.
triphylla, *DC. Prod.* i. 724.—Burma; Malaya.
Vieillardi, *Baill. Adansonia*, xi. (1874) 179.—N. Caled.
viticina, *Wall. Cat.* n. 1219.—Ind. or.
vitiflora, *F. Muell. Fragm.* viii. 144.— Australd.
xanthoxyloides, *F. Muell. l. c.* iv. 155.—Austral.

EVODIA, Gaertn. Fruct. ii. 100 [t. 108] (1791) = **Ravensara**, Sonner. (Laurin.).
aromatica, Pers. Syn. ii. 1 = R. aromatica.
Ravensarae, Gaertn. Fruct. ii. 101 = R. aromatica.

EVODIANTHUS, Oerst in Kjoeb. Vidensk. Meddel. 194 (1857) = **Carludovica**, Ruiz & Pav. (Cyclanth.)
angustifolius, Oerst. l. c. 195 = C. Oerstedii.

EVOISTA, Raf. Sylva Tellur. 56 (1838) = **Lycium**, Linn. (Solan.).
carolinianum, Raf. l. c. = L. carolinianum.
spinosa, Raf. l. c. = L. tetrandrum.

EVOLVOLUS, Sw. Prod. Veg. Ind. Occ. 55 (1788) = seq.

EVOLVULUS, Linn. Sp. Pl. ed. II. 391 (1763). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 875.
CAMDESSIA, Scop. Introd. 190 (1777).
CLADOSTYLES, Humb. & Bonpl. Pl. Aequin. i. 202. t. 57 (1808).
LEUCONALLA, Phil. Sert. Mendoc. Alt. 81 (1871).
MERIANA, Vell. Fl. Flum. 198 (1825).
PLESILIA, Raf. New Fl. Am. iv. 55 (1836); Fl. Tellur. iv. 88 (1836).
acapulcensis, *Willd. ex Roem. & Schult. Syst.* vi. 199. —Mexic.
agrestis, *Schweinf. Beitr. Fl. Aethiop.* 92.—Afr. trop.
albiflorus, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 259.—Mexic.
albiflorus, *Schlecht. in Linnaea*, xxvi. (1853–55) 654. —N. Granat.
alopecuroides, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 96; *et ex Meisn. in Mart. Fl. Bras.* vii. 834.—Am. trop.
alsinoides, Garda. in Hook. Lond. Journ. Bot. i (1842) 585 = pusillus.

EVOLVULUS:—
alsinoides, *Linn. Sp. Pl.* ed. II. 392.—Amphig. trop.
alsinoides, Wall. Cat. n. 1317, partim = breweria evolvuloides.
alsinoides, Willd. Sp. Pl. i. 1517, partim = mucronatus
anagalloides, *Meisen. in Mart. Fl. Bras.* vii. 848.— Am. trop.
angustifolius, Roxb. Hort. Beng. (84); Fl. Ind. ii. 107 = alsinoides.
angustissimus, H. B. & K. Nov. Gen. et Sp. iii. 116 = incanus.
anomalus, *Meisen. in Mart. Fl. Bras.* vii. 858.—Am trop.
arucanus, *Phil. in Anal. Univ. Chil.* (1878) 513.— Chili.
Arbuscula, *Poir. Encyc. Suppl.* iii. 459.—Ins. S. Doming.
argentreus, R. Br. Prod. 489 = alsinoides.
argenteus, *Pursh, Fl. Am. Sept.* i. 187.—Am. bor. occ.
argyreus, *Choisy, Conv. Rar.* 153.— Ecuador.
arizonicus, *A. Gray, Syn. Fl. N. Am.* ii. i. 216.—N. Mexic.
aurigenius, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 100 —Bras.
azureus, Vahl, ex Schum & Thonn. Beskr. Guin. Pl 166 = villosus.
barbatus, *Meisen. in Mart. Fl Bras.* vii. 851.—Am. trop.
brevipedicellatus, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1153, nomen.—Guiana.
canescens, *Meisen. in Mart. Fl. Bras.* vii. 850.— Bras.
canus, Mart. ex Meisn l. c. 837 = helichrysoides.
canus, Spreng. ex Choisy, Conv. Rar. 151 = Arbuscula.
capensis, *E. Mey. ex Choisy, in DC. Prod.* ix. 444.— Afr. austr.
capitatus, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 80.—Bras.
caprolatus, Mart. ex Choisy, in DC. Prod. ix. 445 = alsinoides.
cardiophyllus, *Schlecht. in Linnaea*, xxvi. (1853–55) 653.—N. Granat.
Chamaepitys, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 98.— Am. trop.
chinensis, Choisy, Conv. Rar. 155 = alsinoides.
chrysotrichus, *Meisen. in Mart. Fl. Bras.* vii. 851.— Am. trop.
columbianus, *Meisen. l. c.* 847.—Am. trop.
Commersonii, Steud. Nom. ed. II. i. 630 = incanus.
cordatus, Choisy, in DC. Prod. ix. 447 = anagalloides.
cordatus, *Morr. Pl. Nouv. Am.* 137. t. 82.—Bras.
crassoides, Mart. in Flora, xxiv. (1841) ii. Beibl. 100 = incanus.
cuneifolius, *Raf. Fl. Tellur.* iv. 82.—Hab. ?
cuspidatus, H. B. & K. Nov. Gen. et Sp. iii. 116 = incanus.
daphnoides, *Morr. Pl. Nouv. Am.* 59. t. 40.—Bras.
debilis, H. B. & K. Nov. Gen. et Sp. iii. 115 = alsinoides.
decumbens, N. Br. Prod. 489 = alsinoides.
desertorum, Mart. ex Choisy, in DC. Prod. ix. 442 = Martii.
dichondroides, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 117.—Afr. trop.
diffusus, Chapm. Fl. S. U. St. 345 = alsinoides.
diosmoides, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 97. —Am. trop.
discolor, Benth. Pl. Hartw. 6 = holosericeus.
distichophyllus, Mart. in Flora, xxiv. (1841) ii. Beibl. 101 = holosericeus.
dominguensis, Spreng. ex Choisy, Conv. Rar. 150 = alsinoides.
echioides, *Moric. Pl. Nouv. Am.* 55. t. 37.—Bras.
elegans, *Moric. l. c.* t. 36.—Bras.
emarginatus, Herm. f. Fl Ind. 77 = Ipomoea reniformis.
ericaefolius, Mart. ex Schrank, Pl. Rar. Hort. Monac. t. 94 = gypsophiloides.
ericoides, Mart. in Flora, xxiv. (1841) ii. Beibl. 97 = glomeratus.
exilis, *Meisn. in Mart. Fl. Bras.* vii. 842.—Am. trop.
falcatus, *Griseb. in Gottt. Abh.* xix. (1874) 239.—Reg. Argent.
ferrugineus, Wall. Cat. n. 1316 = Convolvulus pluricaulis.
filiformis, Willd. ex Steud. Nom. ed. II. i. 630 = alsinoides.
filipes, *Mart. in Flora*, xxiv. (1844) ii. *Beibl.* 100.— Am. trop.
frankenioides, *Moric. Pl. Nouv. Am.* 49. t. 33.—Bras.

EVOLVULUS:—
fagacissimus Hochst. in Flora, xxiv. (1841) i. Intell. 35 = alsinoides.
fuscus, *Meisn. in Mart. Fl. Bras.* vii. 330.—Am. trop.
gangeticus, *Linn. Sp. Pl.* ed. II. 391.—Ind. or.
glaber, Spreng. Syst. i. 861 = mucronatus.
glabriusculus, Choisy, Conv. Rar. 156 = mucronatus.
Giechoma, *Welw. Apont.* 589.—Afr. trop.
glomeratus, Choisy, Conv. Rar. 148 = Martii.
glomeratus, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. i. (1823) 81.—Bras.
graphaloides, *Moric. Pl. Nouv. Am.* 61. t. 41.—Bras.
gracillimus, *Mig. Fl. Ind. Bat.* ii. 629.—Sumatra.
grandiflorus, *Parodi, Contrib.* 30.—Paraguay.
guianensis, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1153, numen.—Guiana.
gypsophiloides, *Moric. Pl. Nouv. Am.* 52. t. 35.—Bras.
hederaceus, Burm. f. Fl. Ind. 77 = Convolvulus flavus.
heliantheroides, *Meisn. in Mart. Fl. Bras.* vii. 345.—Am. trop.
helichrysoides, *Moric. Pl. Nouv. Am.* 184. t. 80.—Am. trop.
heterophyllus, Labill. Sert. Austr. Caled. 24. t. 29 = alsinoides.
hirsutulus, Herb. Brit. Mus. ex Choisy, Conv. Rar. 154 = alsinoides.
hirsutus, Lam. Encyc. iii. 538 = alsinoides.
holosericeus, *H. B. & K. Nov. Gen. et Sp.* iii. 116.—N. Granat.
holosericeus, Torr. Bot. Mex. Bound. 150 = arizonicus, incanus.
imbricatus, Mart. ex Choisy, Conv. Rar. 150 = rufus.
incanus, H. B. & K. Nov. Gen. et Sp. iii. 117 = argyreus.
incanus, *Pers. Syn.* i. 288.—Am. trop.
jacobinus, *Moric. Pl. Nouv. Am.* 135. t. 81.—Am. trop.
javanicus, Blume, Bijdr. 724 = alsinoides.
kramerioides, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 98.—Am. trop.
lactus, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 228.—Arizona.
lagopodioides, *Meisn. in Mart. Fl. Bras.* vii. 333.—Bras.
lagopus, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 96.—Bras.
lanceaefolius, Span. in Linnæa, xv. (1841) 341 = alsinoides.
lanceolatus, Poir. Encyc. Suppl. iii. 459 = villosus.
latifolius, *Ker-Gawl. in Bot. Reg.* t. 401.—Bras.
linarioides, *Meisn. in Mart. Fl. Bras.* vii. 348.—Bras.
linifolius, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 353 = filipes.
linifolius, Linn. Sp. Pl. ed. II. 392 = alsinoides.
limoides, Moric. Pl. Nouv. Am. 139. t. 83 = alsinoides.
lithospermoides, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 99.—Bras.
longifolius, *Choisy, Conv. Rar.* 159.—Am. austr.
macroblepharis, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 101.—Am. austr.
macrophyllus, Willd. ex Roem. & Schult. Syst. vi. 199 = Convolvulus floribundus.
madagascaricus, *Vatke, in Linnæa*, xliii. (1880-82) 521.—Madag.
Martii, *Meisn. in Mart. Fl. Bras.* vii. 337.—Am. trop.
Maximiliani, *Mart. ex Choisy, in DC. Prod.* ix. 444.—Am. trop.
microphyllus, Mart. & Gal. in Bull. Acad. Brux. xii. ii. (1845) 257 = alsinoides.
modestus, *Choisy, in DC. Prod.* ix. 448.—Bras.
modestus, Hance, in Walp. Ann. iii. 116 = alsinoides.
mucronatus, *Sw. ex Willd. Sp. Plant. Enum. Hort. Berol. Stockh.* (1837) 61.—Am. trop.
Mushlenbergii, Spreng. Pugill. i. 27 (non ordinis).—Am. bor.
nanus, *Meisn. in Mart. Fl. Bras.* vii. 346.—Am. trop.
nataloensis, *Sond. in Linnæa*, xxiii. (1850) 80.—Afr. austr.
niveus, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 97; *et ex Meisn. in Mart. Fl. Bras.* vii. 332.—Am. trop.
nummularius, Linn. Sp. Pl. ed. II. 391 = alsinoides.
nummularius, Nutt. Gen. Am. i. 174 = mucronatus?
Nuttallianus, Roem. & Schult. Syst. vi. 198 = argenteus.
ovalifolius, Torr. in Bot. Mex. Bound. 150 = Beweria ovalifolia.

EVOLVULUS:—
paniculatus, *Sprang. Syst.* i. 862.—N. Granat.
passerinoides, *Meisn. in Mart. Fl. Bras.* vii. 355.—Bras.
phylicoides, Mart. in Flora, xxiv. (1841) ii. Beibl. 98 = glossnoides.
phyllicoides, Schrad. in Gostt. Gel. Anz. i. (1821) 707 = glomeratus.
phyllanthoides, *Moric. Pl. Nouv. Am.* 52. t. 34.—Bras.
pilosissimus, Mart. & Gal. in Bull. Acad. Brux. xii. ii. (1845) 257 = alsinoides.
pilosus, Nutt. Gen. Am. i. 174 = argenteus.
pilosus, Roxb. Hort. Beng. 23; Fl. Ind. ii. 106 = Convolvulus pluricaulis.
pimulus, Span. in Linnæa, xv. (1841) 341, sphalm. = pumilus.
Pohlii, *Meisn. in Mart. Fl. Bras.* vii. 344.—Am. trop.
procumbens, *Montr. in Mém. Acad. Lyon*, x. (1860) 238.—N. Caled.
Pseudo-incanus, *Span. in Linnæa*, xv. (1841) 341, nomen.—Ins. Timor.
pterocaulon, *Moric. Pl. Nouv. Am.* 140. t. 84.—Bras.
pterygophyllus, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 96.—Bras.
pudicus, Hance, in Walp. Ann. iii. 115 = alsinoides.
pulchellus, *Meisn. in Mart. Fl. Bras.* vii. 341.—Am. trop.
pumilus, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 348, nomen.—Ins. Timor.
purpuro-coeruleus, *Hook. Bot. Mag.* t. 4202.—Jamaic.
pusillus, *Choisy, Conv. Rarv.* 155.—Bras.
ramiflorus, Boj. ex Choisy, in DC. Prod. xi. 447 = alsinoides.
reniformis, Salzm. ex Choisy, Conv. Rar. 150 = alsinoides.
Riedelii, *Meisn. in Mart. Fl. Bras.* vii. 351.—Bras.
rufus, *St. Hil. Voy. Brés.* i. 377, 138.—Bras.
Rutenbergianus, *Vatke, in Bremen, Abh.* ix. (1885) 198.—Madag.
saxifragus, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 99.—Bras.
scoparioides, *Mart. l. c.* 337.—Bras.
sericeus, Benth. in Hook. Lond. Journ. Bot. v. (1846) 355 = holosericeus.
sericeus, Lesnd. ex Choisy, Conv. Rar. 147 = latifolius.
sericeus, Sw. Prod. Veg. Ind. Occ. 55 = incanus.
sericeus, Wall. Cat. n. 1315 = alsinoides.
serpyllioides, *Meisn. in Mart. Fl. Bras.* vii. 345.—Am. trop.
serpyllioides, *Wright, in Sauv. Fl. Cub.* 108.—Cuba.
Sherardi, Choisy, Conv. Rar. 160 = Convolvulus mucrathus.
simplex, *Anderss. Galap. Oarnes Veget.* 211.—Ins. Galapag.
sinicus, Mig. in Journ. Bot. Néerl. i. (1861) 112 = alsinoides.
speciosus, *Moric. Pl. Nouv. Am.* 50. t. 34.—Bras.
stricticaulis, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 354.—Guian.
tenuis, *Mart. ex Choisy, Conv. Rar.* 156.—Bras.
thymiflorus, *Choisy, l. c.* 147.—Bras.
tridentatus, Linn. Sp. Pl. ed. II. 393 = Ipomœa tridentata.
veronicaefolius, H. B. & K. Nov. Gen. et Sp. iii. 117. t. 215 = alsinoides.
villosus, R. Br. Prod. 489 = alsinoides.
villosus, *Ruiz & Pav. Fl. Per.* iii. 30. t. 253. f. b.—Am. austr.
virgatus, Choisy, in DC. Prod. ix. 446 = Convolvulus chondrilloides.
virgatus, Willd. ex Sprang. Syst. i. 862 = incanus.

EVONIMUS, Neck. in Act. Acad. Theod. Palat. ii. (1770) 490 = Euonymus, Linn. (Celastr.).

EVONYMODAPHNE, Nees, Syst. Lauria. 263 (1836) = Ocotea, Aubl. (Lauria.).
armeniaca, Hort. Berol. ex Meisn. in DC. Prod. xv. i. 189 (= *Oreodaphne leptobotra*).—Peruv.
armeniaca, Nees, Syst. Laurin. 264 (= *Oreodaphne Evonymodaphne*).—Peruv.

EVONYMOIDES, Isnard, ex Medic. Phil. Bot. i. 178 (1789) = Celastrus, Linn.
buxifolia, Moench, Meth. 70 = C. buxifolius.
excelsa, Soland. ex DC. Prod. i. 617 = Alectryon excelsum.
scandens, Medic. Phil. Bot. i. 178 = C. scandens.

EVONYMUS, Tourn. ex Linn. Gen. ed. I. 29 (1737) = Euonymus, Linn. (Celastr.).

EVONYXIS, Rafin. Fl. Tellur. ii. 29; iv. 27 (1836) = Melanthium, Linn. (Liliac.).
glauca, Rafin. l. c. = Zygadenus Fremonti.
hybrida, Rafin. l. c. = M. virginicum.
monoica, Rafin. l. c. = M. virginicum.
undulata, Rafin. l. c. (quid?).—Hab.?
virginica, Rafin. l. c. = M. virginicum.

EVOPIS, Cass. in Bull. Soc. Philom. (1818) 32; et in Dict. Sc. Nat. xvi. 65 (1820) = Berkheya, Ehrh. (Compos.).
heterophylla, Cass. in Dict. Sc. Nat. xvi. 66 = B. cynaroides.

EVOSMA, Steud. Nom. ed. I. 330 (1821) = EUOSMA, Andr. = Logania, R. Br.

EVOSMIA, Kunth, Syn. iii. 66 (1824) = Buoemia, Humb. & Bonpl. (Kubiac.).

EVOSMUS, Rafin. Sylva Tellur. 138 (1838) = EUOSMUS, Basil. = Lindera, Thunb. (Lauria.).

EVOTROCHIS, Rafin. Fl. Tellur. ii. 76 (1836) = Primula, Linn.
involucrata, Rafin. l. c. = P. verticillata.
odorata, Rafin. l. c. = P. verticillata.

EVRARDIA, Adans. Fam. ii. 342 (1763) = Pistacia! Linn. (Anacard.).

EWALDIA, Klotzsch, in Monatsb. Berl. Acad. (Mærz, 1854) 123; et in Abh. Akad. Berl. (1854) 53. t. 3 C = Begonia, Linn.
ferruginea, Klotzsch, ll. cc. = B. multinervia.
lobata, Klotzsch, ll. cc. 123, 54 = B. multinervia.

EWYCKIA, Blume, in Flora, xv. (1832) 524 = Pteroxandra, Jack (Melastom.).
capitellata, Walp. Rep. v. 724 = P. capitellata.
coerulea, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 261 = P. cæruleascens.
cordata, Korth. Verh. Nat. Gesch. Bot. 255 = P. cordata.
cyanea, Blume, in Flora, xiv. (1831) 525 = P. cæruleascens.
echinata, Walp. Rep. v. 724 = Kibessia pubescens.
galeata, Korth. Verh. Nat. Gesch. Bot. 254. t. 67 (= *Rectomitra galeata*).—Borneo.
Jackiana, Walp. Rep. v. 724 = P. cæruleascens.
Korthaisiana, Miq. Fl. Ind. Bat. Suppl. 321 = P. Korthalsiana.
latifolia, Blume, Mus. Bot. Lugd. Bat. i. 6 = P. paniculata.
medinilliformis, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 261 = P. capitellata.
paniculata, Miq. Fl. Ind. Bat. Suppl. 321 = P. cæruleascens.
tuberculata, Korth. Verh. Nat. Gesch. Bot. 255 = Kibessia tuberculata.

EXACON, Adans. Fam. ii. 509 (1763) = seq.

EXACUM, Linn. Dias. Dass. 6 (1747); Amoen. Acad. i. 116 (1749). *GENTIANEÆ*, Benth. & Hook. f. ii. 803.
CHONDROPIS, Rafin. Fl. Tellur. iii. 29 (1836).
FLOYERIA, Neck. Elem. i. 386 (1790).
PARACLESIA, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 13.
affine, *Balf. f. in Regel, Gartenfl.* (1883) 84. t. 1108.—Ins. Socotra.
alatum, Heyne, in Roth, Nov. Pl. Sp. 86 = Canscora perfoliata.
albens, Blanco, Fl. Filip. ed. I. 58 = tetragonum.
albens, Linn. f. Suppl. 123 = Sebæa albens.
amplexifolium, D. Dietr. Syn. Pl. i. 440 = nervosum.
anamallayanum, *Bedd. Ic. Pl. Ind. Or.* i. 23.—Ind. or.
apkyllum, Willd. Sp. Pl. i. 838 = Voyria aphylla.
arenarium, Schult. ex Griseb. in DC. Prod. ix. 47 = Centaurium racemosum.
atropurpureum, *Bedd. Ic. Pl. Ind. Or.* i. 24.—Ind. or.
aureum, Linn. f. Suppl. 123 = Sebæa aurea.
auillare, *Thw. Enum. Pl. Zeyl.* 202.—Ind. or.
Beddomei, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 427 = atropurpureum.

EXACUM:—
bellum, Hance, in Hook. Lond. Journ. Bot. vii. (1848) 472 = tetragonum.
bicolor, *Roxb. Hort. Beng.* 88; *Fl. Ind.* i. 397.—Ind. or.
brachiatum, Spix, in Mart. Nov. Gen. et Sp. ii. 397 = Curtia diffusa.
bulbilliferum, *Baker, in Journ. Linn. Soc.* xx. (1883) 209.—Madag.
caeruleum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 50.—Ins. Socotra.
Candollei, Bast. Fl. Maine-et-Loire, Suppl. 22 = Cicendia pusilla.
carinatum, Roxb. Hort. Beng. [10]; Fl. Ind. i. 399 = petiolare.
chilense, Bert. in Merc. Chil. (1829); ex Bull. Férussac, xx. (1830) 111 = Microcala quadrangularis.
chironioides, Griseb. Gen. et Sp. Gent. 109 = tetragonum.
Cleistostoma, Buch.-Ham. ex Wall. Cat. n. 4356 = tetragonum.
concinnum, Miq. Anal. Bot. Ind. iii. 10; et ex C. B. Clarke, in Hook. f. Fl Brit. Ind. iv. 100 = Hoppea fastigiata.
cordatum, Linn. f. Suppl. 124 = Belmonia cordata.
courtallense, *Arn. in Ann Sc. Nat.* Scr. II. xi. (1839) 172.—Ind. or.
cubense, Poepp. ex Griseb. Gen. et Sp. Gent. 126 = Schultesia stenophylla.
dichotomum, Herb. Madr. ex Wall. Cat. n. 4359 = pedunculatum.
diffusum, Willd. Sp. Pl. i. 637 = Canscora diffusa.
erectum, Koth, Nov. Pl. Sp. 83 = Canscora diffusa.
exaltatum, Spreng. Syst. iv. Cur. Post. 41 = pallescens.
fastigiatum, Arn. ex Griseb. Gen. et Sp. Gent. 158 = Hoppea fastigiata.
filiforme, Hook. & Arn. Bot. Beech. Voy. 84 = Microcala quadrangularis.
filiformе, Willd. Sp. Pl. i. 638 = Microcala filiformis.
foliosum, Griseb. in DC. Prod. ix. 45 = Walkeri, Wightianum.
glaucum, Bast. ex Roem. & Schult. Syst. Mant. iii. 97 = Cicendia pusilla.
gracilipes, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 80.—Ins. Socotr.
grandiflorum, Gaertn. Fruct. ii. 158. t. 114. f. 5 = Sebaea capitata.
grandiflorum, Wall. Cat. n. 4358 = bicolor.
guianense, Aubl. Pl. Gui. i. 68 = Schultesia stenophylla.
Hamiltonii, G. Don, Gen. Syst. iv. 212 = tetragonum.
heteroclitum, Willd. Sp. Pl. i. 639 = Canscora sessiliflora.
Horsfieldianum, Miq, Fl. Ind. Bat. ii. 556 = tetragonum.
hyssopifolium, Willd. Sp. Pl. i. 640 = Enicostema littorale.
inflatum, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 285 = Microcala quadrangularis.
Lawii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 98. —Ind. or.
macranthemum, Miq. Anal. Bot Ind. iii. 10 = sessile.
macranthum, Arn. in Ann. Sc. Nat. Ser. II. xi. (1839) 176 = zeylanicum.
Meisnianum, Hohen. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 98 = bicolor.
minus, Roem. & Schult. Syst. iii. Mant. 98 = Coutoubea spicata.
nervosum, *Spreng. Syst. iv. Cur. Post.* 41, 338.—Bras.
ovale, Griseb. Gen. et Sp. Gent. 111 = Walkeri.
ovatum, Labill. Nov. Holl. Pl. i. 58. t. 52 = Sebaea ovata.
pallescens, *Spreng. Syst.* iv. *Cur. Post.* 41, 338.—Bras.
patens, Willd. ex Griseb. Gen. et Sp. Gent. 139 = Erythraea quitensis.
pedunculare, Arn. in Ann. Sc. Nat. Sér. II. xi. (1839) 176 = seq.
pedunculatum, *Linn. Sp. Pl.* 112.—Ind. or.
perfoliatum, D. Dietr. Syn. Pl. i. 440 = pallescens.
Perocttetii, *Griseb. in DC. Prod.* ix. 46.—Ind. or.
petiolare, *Griseb. l. c.* 46.—Ind. or.
pinnathum, *Wall. Cat.* n. 4360.—Burma.
pulchellum, Purah, Fl. Am. Sept. i. 100 = Cicendia pulchella.
pumilum, *Griseb. in DC. Prod.* ix. 46.—Ind. or.
punctatum, *Linn. f. Suppl.* 124.—Ind. or.
purpureum, Lam. Encyc. ii. 647 = Schultesia stenophylla.

EXACUM:—
pusillum, DC. Fl. Fr. iii. 668 = Cicendia pusilla.
quadrangulare, Willd. Sp. Pl. i. 636 = Microcala quadrangularis.
quinquenervium, *Griseb. Gen. et Sp. Gent.* 112.—Madag.
quitense, Spreng. Syst. i. 425 = Erythraea quitensis.
racemosum, Roem. & Schult. Syst. iii. Mant. 99 = Coutoubea ramosa.
ramosum, Vahl, Symb. Bot. iii. 17 = Coutoubea racemosa.
rivulare, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 175 = pedunculatum.
rosum, Royle, Illustr. Bot. Himal. 276 = tetragonum.
rosulatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 210. —Madag.
sessile, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 427 = Linné.
sessile, *Linn. Sp. Pl.* 112.—Ind. or.
spathulatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 210.—Madag.
spicatum, Vahl, Symb. Bot. iii. 17 = Coutoubea spicata.
stellatum, Buch.-Ham. ex Wall. Cat. n. 4359 = pedunculatum.
strictum, Willd. ex Griseb. Gen. et Sp. Gent. 132 = Coutoubea spicata.
stylosum, Wall. Cat. n. 4355 = tetragonum.
sulcatum, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 425 = tetragonum.
sulcatum, Roxb. Hort. Beng. [10]; Fl. Ind. i. 400 = pedunculatum.
sulcatum, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 96 = bicolor.
tenellum, Herb. Madr. ex Wall. Cat. n. 4362 = Canscora diffusa.
tenuifolium, *Aubl. Pl. Gui.* i. 70.—Guian.
teres, *Wall. Cat.* n. 4354.—Reg. Himal.
ternifolium, Roem. & Schult. Syst. iii. 159 = Coutoubea spicata.
tetragonum, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 398. —Ind. or. ; Burma ; China.
tetrapterum, Meyen, ex Griseb. in DC. Prod. ix. 561 = tetragonum.
transmoricum, Bedd. Ic. Ind. Or. i. 24 = courtallense.
tricolor, Zoll. & Mor. Syst. Verz. Zoll. 56 = sulcatum.
vaginale, Labill. Nov. Holl. Pl. i. 37. t. 51 = Logania latifolia.
verticillatum, Willd. Sp. Pl. i. 640 = Enicostema littorale.
violaceum, Lam. Encyc. ii. 647 (= *Apophragma tenuifolium*).—Guiana.
viscosum, Sm. Ic. Pict. 18 = Ixanthus viscosus.
Walkeri, *Arn. ex Griseb. Gen. et Sp. Gent.* 110.—Zeylan.
Wightianum, *Arn. in Ann. Sc. Nat.* Sér. II. xi. (1839) 176.—Ind. or.
zeylanicum, *Roxb. Hort. Beng.* 88; *Fl. Ind.* i. 398.—Zeylan.
zeylanicum, Wall. Cat. n. 4357 A = Wightianum.

EXADENUS, Griseb. Obs. Gentian. 36 (1836) = **Halenia**, Borckh. (Gentian.).
alatus, Mart. & Gal. in Bull. Acad. Brux. xi. i (1844) 372 = H. alata.
brevicornis, Griseb. Gen. et Sp. Gent. 328 = H. brevicornis.
parviflorus, Griseb. l. c. 322 = H. parviflora.
parvifolius, Walp. Rep. vi. 506, sphalm. = seq.
paucifolius, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 372 = H. paucifolia.

EXAGROSTIS, Steud. Nom. ed. II. i. 622 (1840) sphalm. = **Eragrostis**, Host (Gramin.).

EXALLOSIS, Rafin. Fl. Tellur. iv. 88 (1836) = **Convolvulus**, Tourn.
biflora, Rafin. l. c. = C. biflorus.

EXARRHENA, R. Br. Prod. 495 (1810) = **Myosotis**, Linn. (Boragin.).
Lyallii, Hook. f. Handb. N. Zeal. Fl. 105 = M. Lyallii.
macrantha, Hook. f. l. c.—N. Zel.
petiolata, Hook. f. l. c. = M. petiolata.
saxosa, Hook. f. l. c. 196 = M. saxosa.
scaveolens, R. Br. Prod. 495 = M. scaveolens.

EXCOECARIA, Linn. Syst. ed. X. 1288 (1759). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 337.
COMMIA, Lour. Fl. Cochinch. 605 (1790).
SCLEROCROTON, Hochst. in Flora, xxvii, (1845) 85.
SPIROSTACHYS, Sond. in Linnæa, xxiii. (1850) 106.
TARNIOSAPIUM, Muell. Arg. in DC. Prod. xv. ii. 1200 (1866).
abyssinica, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 217.—Abyss.
acerifolia, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1857) 129.—Reg. Himal.
acerosa, Muell. Arg. in DC. Prod. xv. ii. 1207 = Sapium aucuparium.
affinis, Endl. Prod. Fl. Norf. 38 = E. Agallocha.
affinis, Griff. Notul. iv. 486 = Sapium baccatum.
africana, *Muell. Arg. in Linnæa*, xxxii. (1863) 128.—Afr. austr.
Agallocha, *Linn. Syst.* ed. X. 1288.—As. trop ; Malaya ; Ins. Pacif.
albicans, Griseb. in Goett. Nachr. (1866) 179 = Sebastiania Griesebachiana.
arguta, *Muell. Arg. in Mart. Fl. Bras.* xi. ii 614.—Bras.
baccata, Muell. Arg. in DC. Prod. xv. ii. 1211 = Sapium baccatum.
bantamensis, *Muell. Arg. in Linnæa*, xxxii. (1863) 124.—Ins. Bantam.
bicalcarata, *Muell. Arg. l. c.* 125.—Bras.
bicolor, *Hassk. Retzia*, i. 158.—Malaya.
biglandulosa, Muell. Arg. in DC. Prod. xv. ii. 1204 = Sapium aucuparium.
cochinchinensis, *Lour. Fl. Cochinch.* ii. 612.—Cochinch.
cochinchinensis, Muell. Arg. in DC Prod. xv. ii. 1215 = crenulata.
Coligunaya, Baill. Hist. Pl. v. 184 = Adenopeltis Coligunaya.
concolor, *Spreng. Syst.* iii. 24.—Bras.
crenulata, *Wight, Ic.* t. 1865.—Ind. or.
cubensis, *Muell. Arg. in Linnæa*, xxxii. (1863) 122.—Cuba.
cuneata, Muell. Arg. in DC. Prod. xv. ii. 1208 = Sapium cuneatum.
Dallachyana, *Benth. Fl. Austral.* vi. 158.—Austral.
diandra, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 615.—Bras
discolor, Muell. Arg. in DC. Prod. xv. ii. 1210 = Sapium discolor.
discolor, *Spreng. Syst.* iii. 24.—Bras.
diversifolia, *Muell. Arg. in DC. Prod.* xv. ii. 1211.—Sumatra.
eglandulata, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 628. —Bras.
eglandulosa, Muell. Arg. in DC. Prod. xv. ii. 1209 = Bonania adenodon.
emarginata, *Muell. Arg. in DC. Prod.* xv. ii. 1212.—Cuba.
brachyandra, Muell. Arg. l. c. 1224 = Actinostemon concolor.
brachypoda, Griseb. in Goett. Nachr. (1865) 178 = Actinostemon caribæus.
brasiliensis, Spreng. Neue Entdeck. ii. 117 (= *Dactylostemon brasiliensis*).—Bras.
Brolgenii, *Muell. Arg. in Linnæa*, xxxii. (1863) 124. —Bolivia.
Carnettia, Willd. Sp. Pl. iv. 864 = E. Agallocha.
canjoerensis, *Dennst. Schluessl. Hort. Malab.* 31.—Ind. or
caribæa, Griseb. Fl. Brit. W. Ind. 51 = Actinostemon caribæus.
Chamelaea, Baill. Adansonia, vi. (1865–66) 328 = Sebastiania Chamelaea.
erythrosperma, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 161 = Bonania erythrosperma.
farinosa, Griseb. in Goett. Abh. vii. (1857) 169 = Gymnanthes hypoleuca.
glandulosa, *Sw. Fl. Ind. Occ.* ii. 1194.—Am. trop.
glauca, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 84.—Reg. Argent.
Goudotiana, *Muell. Arg. in Linnæa*, xxxii. (1863) 122 —Madag.
guianensis, Baill. Hist. Pl. v. 188 = Maprounea guianensis.
guineensis, *Muell. Arg. in Linnæa*, xxxii. (1863) 123. —Afr. trop.
haematosperma, Muell. Arg. in DC. Prod. xv. ii. 1209 = Sapium haematospermum.
herbertiaefolia, *Jacq. in Lem. Hort. Univ.* (1844) 289. —Austral.

EXOECARIA :—
himalayensis, Muell. Arg. in Linnæa, xxxii. (1863) 122 = conrifolia.
hippophaefolia, *Griseb. in Goett. Abh.* xxiv. (1879) 61. —Reg. Argent.
Hookstatriviana, Muell. Arg. in Linnæa, xxxii. (1863) 123 = reticulata.
holophylla, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 247.—Burma.
ilicifolia, *Spreng. Neue Entdeck.* ii. 117.—Bras.
indica, Muell. Arg in Linnæa, xxxii. (1863) 123 = Sapium indicum.
insignis, Muell. Arg. in DC. Prod. xv. ii. 1212 = Sapium insigne.
integerrima, Muell. Arg. l. c. 948 = reticulata.
integrifolia, *Roxb Hort. Beng.* [105]; *Fl. Ind.* iii. 757. —Ins. Moluicc.
japonica, *Muell. Arg. in Linnæa*, xxxii. (1863) 123. —Japon.
Klotzschii, Beill. Hist. Pl. v. 185 (= *Dactylostemon Klotzschii*).—Bras.
laevis, Blanco, Fl. Filip. ed. I. 788 = Homalanthus populifolius.
lanceolaris, *Muell. Arg. in DC. Prod.* xv. ii. 1221.— Ins. Penang.
Lastellei, *Muell. Arg. l. c.* 1218.—Madag.
Laurocerasus, Muell. Arg. l. c. 1202 = Sapium Laurocerasus.
leucogyna, Muell. Arg. l. c. 1208 = Sapium leucogynum.
leucosperma, Muell. Arg. l. c. 1209 = Sapium leucospermum.
Laureiranus, *Muell. Arg. l. c.* 1217.—Cochinch.
lucida, *Sw. Fl. Ind. Occ.* ii. 1121.—Jamaic.
macrocarpa, Muell. Arg. in DC. Prod. xv. ii. 1207 = Sapium macrocarpum.
madagascariensis, *Muell. Arg. l. c.* 1219 —Madag.
Manniana, *Muell. Arg. in Flora*, xlvii. (1864) 453.— Afr. trop.
marginata, *Griseb. in Goett. Abh.* xix. (1874) 97.— Reg. Argent.
marginata, Kunze, ex Beill, Etud. Gén. Euph. 582 = Adenopeltis Colligusya.
marginata, Muell. Arg. in DC. Prod. xv. ii. 1208 = Sapium marginatum.
Martii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 614.—Bras.
melanostieta, *Muell. Arg. in Linnæa*, xxxii. (1863) 122.—Madag.
myricifolia, Griseb. in Goett. Nachr. (1865) 178 = Bosaula myricifolia.
oblongifolia, *Muell. Arg. in Journ. Bot.* ii. (1864) 337.—Afr. trop.
obovata, Muell. Arg. in DC. Prod. xv. ii. 1208 = Sapium obovatum.
obtusifolia, *Muell. Arg. in Linnæa*, xxxii. (1863) 125. —Penv.
obtusiloba, Muell. Arg. in DC Prod. xv. ii. 1208 = Sapium obtusilobum.
occidentalis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 615. —Bras.
oppositifolia, *Griff. in Calc. Journ. Nat. Hist.* iv. (1844) 886.—Reg. Himal.
oppositifolia, Muell. Arg. in DC. Prod. xv. ii. 1219 = crenulata.
ovalis, Endl. Prod. Fl. Norfolk. 38 = E. Agallocha.
ovatifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 14 = Cerbera Odollam.
pallens, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 161 = Gymnanthes pallens.
pallida, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 628.— Bras.
parvifolia, *Muell. Arg. in Flora*, xlvii. (1864) 453.— Austral.
polyandra, Griseb. in Goett. Nachr. (1865) 180 = Gymnanthes polyandra.
potamophila, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 637.—Bras.
quadrangularis, *Muell. Arg. in Linnæa*, xxxii. (1863) 124.—Malaya.
reticnervis, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 146.—Ins. Nicobar.
reticulata, *Muell. Arg. in DC. Prod.* xv. ii. 1213.— Afr. exter.
riparia, Schlecht. in Linnæa, vii. (1832) 386 = Gymnanthes riparia.
Sagraei, *Muell. Arg. l. c.* xxxii. (1863) 121.—Cuba.
salpingodentata, Muell. Arg. in DC. Prod. xv. ii. 1209 = Sapium salpingodentatum.
sebifera, Muell. Arg. l. c. 1210 = Sapium sebiferum.

EXOECARIA :—
serrata, [R. Br. in] Ait. Hort. Kew. ed. II. v. 418 = Adenopeltis Colligusya.
serrulata, *Miq. in Linnæa*, xix. (1847) 446.—Bras.
sicca, Blanco, Fl. Filip. ed. I. 787 = Homolanthus populifolius.
stylaris, Muell. Arg. in DC. Prod. xv. ii. 1204 = Sapium stylare.
suberosa. Muell. Arg. l. c. 1202 = Sapium suberosum.
subsessilis, *Muell. Arg. l. c.* 1203.—Bras.
subulata, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 637.— Bras.
tenax, Griseb. in Goett. Nachr. (1865) 179 = Gymnanthes pallens.
Thouarsiana, *Muell. Arg. in DC. Prod.* xv. ii. 1218.— Madag.
tijuccana, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 616. —Bras.
tinifolia, *Sw. Fl. Ind. Occ.* ii. 1119.—Jamaic.
triatia, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 615.— Bras.
venulosa, Wright, ex Griseb. in Goett. Nachr. (1865) 159 = Sebastiana Grisebachiana.
virgata, Zoll. & Mor. ex Miq. Fl. Ind. Bat. i. ii. 416 (Sapii sp.).—Java.
Warmingii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 626. —Bras.

EXOERMIS, Willd. ex Schult. f. Syst. vii. 354 in nota (1830). *LILIACEAE*, Benth. & Hook. f. iii. 794.
coarctata, Baker, in Journ. Linn. Soc. xv. (1876) 320 = ramosa.
ramosa, *Willd. ex Schult. f. Syst.* vii. 354 in syn.— Am. trop.

EXERIA, Rafin. Fl. Tellur. iv. 49 (1836) = **Eria**, Lindl. (Orchid.).

EXITELES, Miers, in Journ. Linn. Soc. xvii. (1879) 336 = acq.

EXITELIA, Blume, Fl. Jav. Praef. p. vii (1838) = **Farinarium**, Juss. (Rosac.).
corymbosa, Blume, l. c. = P. Maranthes.
multiflora, Walp. Rep. v. 115 = P. Griffithianum.

EXOACANTHA, Labill. Ic. Pl. Syr. i. 10. t. 2 (1791). *UMBELLIFERAE*, Benth. & Hook. f. i. 927.
heterophylla, *Labill. l. c.*—Syria.

EXOCARPUS, Labill. Voy. i. 355. t. 14 (1798) (Exocarpo); Pers. Syn. ii. 561 (1807). *SANTALACEAE*. Benth. & Hook. f. iii. 230.
CANOPUS, Presl, Epim. Bot. 248 (1849).
SARCOCALYX, Zipp. ex Mackl. in Bijdr. Wet. v. 179 (1850).
XYNOPHYLLA, Montr. in Mém. Acad. Lyon, x. (1860) 250.
aphylla, *R. Br. Prod.* 357.—Austral.
Bidwillii, *Hook. f. Fl. N. Zel.* 223. t. 52.—N. Zel.
Casuarinae, *Baill. Adansonia*, iii. (1863-63) 109.— Ins. Sandvic.
ceramicus, *A. DC. in DC. Prod.* xiv. 691.—Malaya.
ceramicus, Naves, in Blanco, Fl. Philipp. ed. III. t. 317 = Moehlenbeckia platyclada.
cupressiformis, Hook. & Arn. Bot. Beech. Voy. 95 = Gaudichaudii.
cupressiformis, *Labill. Voy.* i. 155. t. 14.—Austral.
dacrydioides, Schlecht. in Linnæa, xx. (1847) 580 = aphylla.
Gaudichaudii, *A. DC. in DC. Prod.* xiv. 690.—Ins. Sandvic.
glandulosus, Miq. in Lehm. Pl. Preiss. i. 619 = sparteus.
homaeocladus, *C. Moore & F. Muell. in F. Muell. Progra.* viii. 9.—Austral.
humifusus, *R. Br. Prod.* 357.—Ins. Tasman.
humifusus, Hook. f. Fl. Tasm. i. 326 = nanus.
lactiflus, *R. Br. Prod.* 356.—Austral.
leptomerioides, F. Muell. Miq. in Nederl. Kruidk. Arch. iv. (1859) 108 = aphyllus.
linnonensis, Presl, Epim. Bot. 248 = latifolius.
minutulus, Zipp. in Bull. Férussac, xviii. (18) 92 = latifolius.
nanus, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 231.—Ins. Tasman.
odoratus, *A. DC. in DC. Prod.* xiv. 689.—Austral.
ovatus, Blume, Mus. Bot. Lugd. Bat. i. 191 = latifolius.

EXOCARPUS:—
pendulus, F. Muell. in Trans. Vict. Inst. i. (1855) 42 = sparteus
phyllanthoides, *Endl. Prod. Fl. Norfolk.* 46.—Ins. Norfolk.
sandvicensis, *Baill. Adansonia*, iii. (1862-63) 109.— Ins. Sandvic.
sparteus, *R. Br. Prod.* 356.—Austral.
spicatus, A. DC. in DC. Prod. xiv. 689 = sparteus.
strictus, *R. Br. Prod.* 357.—Austral.
strictus, Sieber, ex A. DC. in DC. Prod. xiv. 689 (Casuarinae sp.).—Austral.
xylophylloides, *Baker, in Journ. Linn. Soc.* xx. (1883) 249.—Madag.

EXOCARYA, Benth. in Hook. Ic. Pl. t. 1206 (1877). *CYPERACEAE*, Benth. & Hook. f. iii. 1056.
sclerioides, *Benth. l. c.*—Austral.

EXOCHAENIUM, Griseb. in DC. Prod. ix. 55 (1845) = **Belmontia**, E. Mey. (Gentian.).
debile, Welw. in Trans. Linn. Soc. xxvii. (1869) 48.— Afr. trop.
grande, Griseb. in DC. Prod. ix. 55 = B. grandis.
primuloflorum, Welw. in Trans. Linn. Soc. xxvii. (1869) 47.—Afr. trop.

EXOCHORDA, Lindl. in Gard. Chron. (1858) 925. *ROSACEAE*, Benth. & Hook. f. i. 613.
Alberti, *Regel, in Act. Hort. Petrop.* viii. (1884) 696. —As. centr.
Davidiana, Baill. Adansonia, ix. (1868-70) 149 = Nuttallia Davidiana.
grandiflora, *Lindl. in Gard. Chron.* (1858) 925.— China.
Korolkowi, *Lavall. Arbor. Segrez.* 39.—Soongar.?
serratifolia, *S. Moore, in Hook. Ic. Pl.* t. 1255.— China.

EXOCROA, Rafin. Fl. Tellur. iv. 80 (1836) = **Convolvulus**, Linn.
aegyptiaca, Rafin. l. c. = C. althaeoides.

EXODECONUS, Rafin. Sylva Tellur. 57 (1838) = **Physalis**, Linn. (Solan.).
prostratus, Rafin. l. c. = P. prostrata.

EXODICLIS, Rafin. Sylva Tellur. 97 (1838) = **Acisanthera**, P. Br. (Melast.).
angustifolia, Rafin. l. c. 98 = A. brevifolia.
latifolia, Rafin. l. c. = A. brevifolia.

EXOGONIUM, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 443 (Conv. Or. 61) = **Ipomoea**, Linn. (Convolv.).
arenarium, Choisy, Conv. Rar. 129. t. 1 = I. arenaria.
bracteatum, Choisy, in Mém. Soc. Phys. Genèv. vi (1833) 448 = I. bracteata.
curviflorum, Moç. & Sesse, ex Choisy, in DC. Prod. ix. 336, 347 = I. coccinea.
dumosum, Benth. Pl. Hartw. 46 = I. Purga.
eriospermum, Choisy, Conv. Rar. 130 = I. Desrousseauxii.
filiforme, Choisy, l. c. 129 = I. filiformis.
Olivae, Bâroma, Viaje Cav. Cacahuam. 29 = I. bracteata.
pedatum, Choisy, Conv. Rar. 130 = I. pedata.
Purga, Lindl. Bot. Reg. (1847) t. 49 = I. Purga.
racemosum, Choisy, Conv. Rar. 128 = I. repanda.
repandum, Choisy, l. c. = I. racemosa.
spicatum, Choisy, l. c. = I. bracteata.
umbellatum, Moç. & Sesse, ex Choisy, in DC. Prod. ix. 336, 347 = I. peduncularis.

EXOLEPTA, Rafin. in Am. Monthly Mag. (1819) 198 = **Cassandra**, D. Don (Ericac.).
calyculata, Rafin. l. c. = C. calyculata.

EXOLOBUS, Fourn. in Mart. Fl. Bras. vi. iv. 218 (1868). *ASCLEPIADEAE*.
grandiflorus, *Fourn. l. c.* 219.—Bras.
patens, *Fourn. l. c.* 218.—Bras.
rotatus, *Fourn. l. c.* 220.—Bras.
Selloanus, *Fourn. l. c.* 219.—Bras.
stenolobus, *Fourn. l. c.*—Bras.

EXOMIS, Fenzl, ex Moq. Chenop. Enum. 49 (1840). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 58.
albicans, Moq. in DC. Prod. xiii. 11. 89 = atriplicioides. atriplicioides, *Moq. Chenop. Enum.* 49.—Afr. austr.
axyrioides, *Fenzl, ex Moq. l. c.*—Afr. austr.

EXOPHYA, Rafin. Fl. Tellur. ii. 63 (1836) = **Epidendrum**, Linn. (Orchid.).
fuscata, Rafin. l. c. = E. patens.

EXORRHIZA, Becc. in Ann. Jard Buitenz. ii. (1885) 198. *PALMAE*.
Wendlandiana, *Becc. l. c.*—Ins. Fiji.

EXOSTEGIA, Boj. ex Decne in DC. Prod. viii 528, 530, in syn. (1844) = **Cynanchum**, Linn. (Asclep.).
filicoulis, Boj. ex Decne. l. c. 530 (= *Cynoctonum repandum*).—Madag.
laevigata, Boj. ex Decne. l. c. 528 (= *Cynoctonum curvychiton*).—Madag.

EXOSTEMMA, Rich. ex Humb. & Bonpl. Pl. Aequin. i. 131. t. 38 (1808) (Exostema). *RUBIACEAE*, Benth. & Hook. f. ii. 42.
angustifolium, *Roem. & Schult. Syst.* v. 19.—Ins. S. Doming.
australe, *St. Hil. Pl. Us. Bras.* t. 1. 3.—Bras.
bicolor, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 32.—Peruv.
brachycarpum, *Roem. & Schult. Syst.* v. 19.—Jamaic.
canescens, *Bartl. ex DC. Prod.* iv. 359.—Mexic.
capitatum, *Spreng. Neue Entdeck.* ii. 143.—Am. austr.
caribaeum, *Roem. & Schult. Syst.* v. 18.—Ind occ.
coriaceum, *Roem. & Schult. l. c.* 20.—Ins. S. Doming.
corymbiferum, *Roem. & Schult. l. c.*—Ins. Pacif.
corymbosum, *Spreng. Syst.* i. 706.—Peruv.
Coulteri, *Hook. f. ex Hemsl. Diag. Pl. Nov.* 32.—Mexic.
cuspidatum, *St. Hil. Pl. Us. Bras.* i. t. 3.—Bras.
dissimiliforum, *Roem. & Schult. Syst.* v. 17.—Am. austr.
ellipticum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 504.—Ind. occ.
floribundum, *Roem & Schult. Syst.* v. 19.—Ind. occ.
formosum, *Cham. & Schlecht. in Linnaea*, iv. (1829) 179.—Bras.
grandiflorum, *Hort. ex Regel, Cat. Pl. Hort. Asakaw.* 58.—Hab. ?
linearum, *Roem. & Schult. Syst.* v. 18.—Ins. S. Doming.
longicuspe, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 48.—Costa Rica.
longiflorum, *Roem. & Schult. Syst.* v. 18.—Ins. S. Doming.
Macrocnemia, G. Don, Gen. Syst. iii. 482 = Remija Macrocnemia.
maynense, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 31. t. 237.—Bras.
mexicana, *A. Gray, in Proc. Am. Acad.* v. (1861) 180.—Mexic.
myrtifolium, *Griseb. Cat. Pl. Cub.* 125.—Cuba.
nerilifolium, *A. Rich. Pl. Cub. Fanerog.* ii. 7.—Cuba.
occidentale, Benth. Bot. Voy. Sulph. 104 = Rustia occidentalis.
parviflorum, *Rich. ex Humb. & Bonpl. Pl. Aequin.* i. 132.—Ind. occ.
peruvianum, *Humb. & Bonpl. l. c.* 132.—Peruv.
philippicum, Blanco, Fl. Filip. ed. II. 118 = Hymenodictyon excelsum.
phillyreum, *Roem. & Schult. Syst.* v. 20.—Ins. Philipp.
purpureum, *Griseb. Cat. Pl. Cub.* 125.—Cuba.
rotundatum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 504.—Ind. occ.
salicifolium, *Griseb. Cat. Pl. Cub.* 125.—Ind. occ.
Sonnazzae, *Mart. Reise Bras.* ii. 789.—Bras.
triflorum, *G. Don, Gen. Syst.* iii. 481.—Jamaic.
Valenzuelae, A. Rich. Fl. Cub. Fanerog. t. 48 = parviflorum.
Vavassorii, *Griseb. Fl. Brit. W. Ind.* 393.—Ins. S. Doming.

EXOSTYLIS, Schott. ex Spreng. Syst. iv. Cur. Post. 406 (1827). *LEGUMINOSAE*, Benth. & Hook. f. i. 566.
glabra, *Pap. in Linnaea*, xi. (1837) 266.—Bras.
venosa, *Schott, ex Spreng. Syst.* iv. *Cur. Post.* 406.—Bras.

EXOSTYLIS, G. Don, Gen. Syst. ii. 454 (1832) = praec.

EXOTANTHERA, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 342. *VIOLARIEAE*?—Cf. Benth. & Hook. f. i. 970.
racemosa, *Turcz. l. c.*—Madag.

EXOTHAMNUS, D Don, ex Hook. Comp. Bot. Mag. ii. (1836) 46 = **Abtsilofia**, Tourn. (Compos.).
ilicifolius, D. Don, l. c. (= *Diplopappus ilicifolius*).—Reg. Argent.

EXOTHEA, Macfad. Fl. Jamaic. i 232 (1837) = **Bypelate**, P. Br. (Sapindac.).
oblongifolia, Macfad. l. c. = H. paniculata.
paniculata, Walp. Rep. v. 366, in syn. = H. paniculata.

EXOTHECA, Anderss. in Nov. Act. Upsal. Ser. III ii. (1856) 253 = **Anthisistria**, Linn. (Gramin.).
abyssinica, Anderss. l. c. = A. abyssinica.

EXOTHOSTEMON, G Don, Gen. Syst. iv. 62 (1836) = **Prestonia**, R. Br. (Apocyn.).
bracteatum, G. Don, Gen. Syst. iv. 82 = Echites bracteata.
contortum, Miers, Apocyn. S. Am. 241 = P. contorta.
gracile, G. Don, Gen. Syst. iv. 82 = Echites gracilis.
javitense, G. Don, l. c. = Echites javitensis.
macrophyllum, G. Don, l. c. = Echites macrophylla.
mollissimum, G. Don, l. c. = Echites mollissima.
paludorum, G. Don, l. c. 83 = Echites paludosa.
sericeum, Miers, Apocyn. S. Am. 241 = P. sericea.
speciosum, G. Don, Gen. Syst. iv. 82 = Echites speciosa.

EXPHALOSCHOENUS, Nees, in Hook. Journ. Bot. ii. (1840) 398, sphalm = CEPHALOSCHOENUS, Nees = **Hynchospora**, Vahl (Cyperac.).

EXTELMA, Dum. Anal. Fam. 26 (1829) = **Oxystelma** ? R. Br. (Asclep.).

EXYDRA, Endl. Fl. Posoyn. 119 (1830) = **Glycoria**, R. Br. (Gramin.).
aquatica, Endl. l. c. = G. aquatica.

EYDISANTHEMA, Neck. Elem. iii. 133 (1790) = **Epidendrum**, Linn. (Orchid.).

EYDOUXIA, Gaudich. Bot. Voy. Bonite, t. 18 (1844-52) = **Pandanus**, Linn.
Delessertii, Gaudich. l. c. t. 17. ff. 7, 8 = P. odoratissimus.
macrocarpa, Gaudich. c. t. 18. ff. 1-6 = P. Eydouxia.

EVREA, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 331 = **Turpinia**, Vent. (Sapindac.).
vernalis, Champ. ex Benth. l. c. = T. arguta.

EVREA, F. Muell. in Linnaea, xxv. (1852) 402 = **Pluchea**, Cass. (Compos.).
rubelliflora, F. Muell. l. c. = P. Eyreas.

EVRYTHALIA, Reneslm. ex Borckh. in Roem. Arch. i. 1. 28 (1796) = **Gentiana**, Tourn.
Amarella, Borckh. l. c. = G. Amarella.
angustifolia, D. Don, in Sweet, Hort. Brit. ed. III. 471 = G. angustifolia.
auriculata, G. Don, Gen. Syst. iv. 187 = G. auriculata.
centaurium, Borckh. in Roem. Arch. i. 1. 33 = G. campestris.
canaliculata, D. Don, ex Griseb. in DC. Prod. ix. 96 = G. Moocroftiana.
carinata, D. Don, in Trans. Linn. Soc. xvii. (1837) 516 = G. carinata.
carinthiaca, Berchtd. & Presl, Rostl. i. Gent. 18 = Pleurogyne carinthiaca.
caucasica, Berchtd. & Presl, l. c. = G. caucasica.
conferta, G. Don, in Sweet, Hort. Brit. ed. III. 471 = G. conferta.
coronata, D. Don, in Trans. Linn. Soc. xvii. (1837) 515 = G. coronata.
dichotoma, Berchtd. & Presl, Rostl. i. Gent. 18 = G. dichotoma.
floribunda, G. Don, Gen. Syst. iv. 187 = Swertia Chirata.
Gekleri, G. Don, in Sweet, Hort. Brit. ed. III. 471 = G. decumbens.

EVRYTHALIA :—
Gebleria, Stend. Nom. ed. II. i. 616 = praec.
germanica, Schrank, Pl. Rar. Hort. Monac. t. 278 = Gentiana germanica.
glacialis, Berchtd. & Presl, Rostl. i. Gent. 18 = G. tenella.
lancifolia, Berchtd. & Presl, l. c. = G. Amarella.
nana, Borckh. in Roem. Arch. i. 1. 28 = G. tenella.
obtusifolia, Berchtd. & Presl, Rostl. i. Gent. 18 = G. obtusifolia.
pedunculata, D. Don, in Trans. Linn. Soc. xvii. (1837) 516 = G. tenella.
pratensis, Berchtd. & Presl, Rostl. i. Gent. 18 = G. Amarella.
pulchella, S. F. Gray, Nat. Arr. Brit. Pl. ii. 336 = G. Amarella.
rotata, Berchtd. & Presl, Rostl. i. Gent. 19 = Pleurogyne carinthiaca.
septemfida, Borckh. in Roem. Arch. i. 1. 28 = G. septemfida.
sulcata, Berchtd. & Presl, Rostl. i. Gent. 19 = Pleurogyne carinthiaca.
tenella, Buck, DC. Prod. Index, iii. 182 = G. tenella.
uliginosa, Berchtd. & Presl, Rostl. i. Gent. 17 = G. Amarella.
umbellata, D. Don, ex G. Don, Gen. Syst. iv. 187 = G. aurea.
uniflora, Berchtd. & Presl, Rostl. i. Gent. 17 = G. germanica.

EVSELIA, Neck. Elem. i. 199 (1790) = **Gallum**, Linn. (Rubiac.).

EVSELIA, Reichb. Iconog. Exot. iii. 15. t. 242 (1830) = **Egletes**, Cass. (Compos.).
bellidiflora, Reichb. l. c. = Egletes bellidiflora.

EYSENHARDTIA, H. B. & K. Nov. Gen. et Sp. vi. 489. t. 592 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 492.
VARENNEA, DC. Prod. ii. 522 (1825).
VIBORQUIA, Orteg. Hort. Matr. Dec. 66. t. 9 (1798), nomen prius.
adenostylis, *Baill. Adansonia*, ix. (1870) 239.—Guatemala.
amorphoides, *H. B. & K. Nov. Gen. et Sp.* vi. 489.—Mexic.
Drummondii, Torr. & Gray, Fl. N. Am. i. 690 = amorphoides.
orthocarpa, S. Wats. in Proc. Am. Acad. xvii. (1881-82) 331 = amorphoides.
spinosa, *Engelm. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 174.—Texas.
texana, Scheele, in Linnaea, xxi. (1848) 462 = amorphoides.

EYSTATHES, Lour. Fl. Cochinch. 284 (1790). *INC. SED.*—Cf. Benth. & Hook. f. i. 892.
sylvestris, *Lour. l. c.* 285.—Cochinch.

F

FABA, [Tourn.] Linn. Syst. ed. I (1735) = **Vicia**, Linn. (Leguin.).
angustifolia, Bernh. Syst. Verz. Erf. 250 = V. sativa.
bengalensis, Roxb. Fl. Ind. ed. Carey, ii. 435, in obs. = Strychnos Ignatii ?
Bona, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 350 = V. narbonensis.
equina, Medic. l. c. = V. narbonensis.
major, Desf. Tabl. ed. I. 196 = V. Faba.
minor, Roxb. Fl. Ind. iii. 323, in obs. = V. Faba.
narbonensis, Schur, in Verh. Naturf. Ver. Brunnn. xv. ii. (1877) 193 = V. narbonensis.
sativa, Bernh. Syst. Verz. Erf. 250 = V. Faba.
sepium, Bernh. l. c. = V. sepium.
serratifolia, Fras. Fl. Transc. 179 = V. narbonensis.
vulgaris, Moench, Meth. 150 = V. Faba.

FABAGO, Tourn. ex Adans. Fam. ii. 507 (1763) = **Zygophyllum**, Linn.
alata, Moench, Meth. Suppl. 90 = Z. Fabago.
major, Sweet, Brit. Flow. Gard. vi. t. 226 = Z. Fabago.
portulacaefolius, Medic. in Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 213 = Z. simplex.

FABERA, Sch. Bip. in Nov. Act. Nat. Cur. xxi. (1845) 129 = **Hypochoeris** Linn. (Compos.).
cretensis, Sch. Bip. l. c. 136 = H. cretensis.
hispida, Sch. Bip. l. c. 131 – H. cretensis.

FABIANA, Ruiz & Pav. Prod. 22. t. 24 (1794). *SOLANACEÆ*, Benth. & Hook. f. ii. 905.
bifora, *Remy, in C. Gay, Fl. Chil.* v. 41.—Chili.
bryoides, *Phil. Fl. Atac.* 40.—Chili.
cordillola, *Dun. in DC. Prod.* xiii. 580.—Chili.
densa, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 297.—N. Granat.
denudata, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 168.—Chili.
ericoides, *Dun. in DC. Prod.* xiii. 590.—Chili.
grandiflora, *Dun. l. c.* 591.—Chili.
heterophylla, *Dun. l. c.* 593.—Bras.
Hieronymi, *Niederl. in Roca, Exped. Rio-Negro*, ii. 258.—Patagon.
imbricata, *Ruiz & Pav. Fl. Per.* ii. 19. t. 122.—Pervv.
lanuginosa, *Hook. & Arn. Bot. Beech. Voy.* 35.—Chili.
Peckii, *Niederl. in Roca, Exped. Rio-Negro*, ii. 258.—Patagon.
petuniolden, *Grisb. in Goett. Abh.* xxiv. (1879) 242.—Reg. Argent.
Sellowiana, *Dun. in DC. Prod.* xiii. 592.—Bras.
squamuligera, *Dun. l. c.* 591.—Chili.
thymifolia, St. Hil. in Mem. Mus. Par. xii. (1825) 317 = Petunia thymifolia.
viscosa, *Hook. & Arn. Bot. Beech. Voy.* 36.—Chili.

FABRENIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 = **Anacardium** ? Linn.

FABRIA, E. Mey. Zwei Pfl. Docum. 185, nomen (1848) = **Ruellia**, Linn. (Acanth.).
cordyfolia, Nees, in DC. Prod. xi. 114 = R. ovata.
pilosa, Nees, l. c. = R. pilosa, *Linn.*
rigida, E. Mey. Zwei Pfl. Docum. 185 = R. pilosa, *Linn.*

FABRICIA, Adans. Fam. ii. 168 (1763) = **Lavandula**, Tourn. (Labiat.).

FABRICIA, Gaertn. Fruct. i. 175. t. 35. f. 4 (1788) = **Leptospermum**, Forst. (Myrtac.).
angustifolia, *Otto, ex DC. Prod.* iii. 226 = Agonis marginata.
bracteata, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 476.—Ins. Molucc.
coriacea, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 147 = L. laevigatum.
incana, Dum.-Cours. Bot. Cult. ed. II. v. 387 = L. pubescens.
laevigata, Gaertn. Fruct. i. 175. t. 35 = L. laevigatum.
minor, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 387 = L. pubescens.
myrtifolia, Hort. Angl. ex Dum. Cours. Bot. Cult. ed. II. vii. 278 = L. pubescens.
myrtifolia, Hort. ex Heynh. Nom. ii. 70 = Agonis marginata.
myrtifolia, Sieber, ex Benth. Fl. Austral. iii. 103 = L. laevigatum.
pubescens, Hort. ex Schau. in Linnæa, xv. (1841) 422 = L. pubescens.
sericea, Sweet, Hort. Brit. ed. I. 165 = L. pubescens.
stricta, Lodd. Bot. Cab. t. 1219 = Agonis marginata.

FABRICIA, Scop. Introd. 307 (1777) = **Alyssocarpus**, Neck. (Legum.).

FABRICIA, Thunb. in J. C. Fabric. Reise nach Norw. 23 (1779) = **Curouligo**, Gaertn. (Amaryllid.).
alba, Thunb. l. c. 26 = Hypoxis alba.
amœna, Salisb. Prod. 244 = Veltheimia viridifolia.
glauca, Salisb. l. c. 245 = Veltheimia glauca.
minuta, Thunb. in J. C. Fabric. Reise nach Norw. 25 = Pauridia hypoxidioides.
plicata, Thunb. l. c. 29 = C. plicata.
serrata, Thunb. l. c. = Hypoxis serrata.
stellata, Thunb. l. c. 27 = Hypoxis stellata.
villosa, Thunb. l. c. 31 = Hypoxis villosa.

FABRITIA, Medic. in Staatsw. Vorlts. Churpf. Phys.-Oek. Ges. i. (1791) 230 = FABRICIA, Adans. = **Lavandula**, Tourn. (Labiat.).

FACCHINIA, Reichb. Ic. Fl. Germ. xv. 29 (1841) = **Arenaria**, Rupp. (Caryophyll.).
lanceolata, Reichb. l. c. = A. lanceolata.

FACELIS, Cass. in Bull. Soc. Philom. (1819) 94; et in Dict. Sc. Nat. xvi. 104 (1820). *COMPOSITÆ*, Benth. & Hook. f. ii. 304.
LEPTALEA, D. Don, ex Hook. & Arn. in Comp. Bot. Mag. i. (1835) 102.
PTEROPOGON, Fenzl, in Flora, xxii. (1839) ii. 727.
apiculata, *Cass. in Dict. Sc. Nat.* xvi. 104.—Am. austr.
plumosa. *Sch. Bip. in Linnæa*, xxiv. (1865–66) 632, nomen.—Boliv.
retusa, *Sch. Bip. l. c.*, nomen.—Boliv.

FADGENIA, Lindl. Veg. Kingd. 295 (1847), sphalm. = FADVENIA, Endl. = **Garrya**, Dougl. (Cornac.).

FADOGIA, Schweinf. Rel. Kotsch. 47. t. 32 (1868). *RUBIACEÆ*, Benth. & Hook. f. ii. 111.
PACHYSTIGMA, Hochst. in Flora, xxv. (1842) 284.
agrestis, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 154.—Afr. trop.
ancylantha, *Schweinf. l. c.* 155.—Afr. trop.
Cienkowskii, *Schweinf. Rel. Kotsch.* 47.—Afr. trop.
fuchsioides, *Schweinf. ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 85. t. 50.—Afr. trop.
glaberrima, *Welw. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 155.—Afr. trop.
lactiflora, *Welw. ex Hiern, l. c.* 156.—Afr. trop.
leucophloea, *Hiern, l. c.* 153.—Afr. trop.
stenophylla, *Welw. ex Hiern, l. c.* 155.—Afr. trop.
Zeyheri, *Sond. in Linnæa*, xxiii. (1850) 56.—Afr. austr.

FADVENIA, Endl. Gen. Suppl. ii. 30 (1842) = **Garrya**, Dougl. Cornac.).
Hookeri, Endl. l. c. Suppl. iv. 36 = G. Fadyena.
laurifolia, Endl. l. c. = G. laurifolia.
macrophylla, Endl. l. c. = G. macrophylla.
oblonga, Endl. l. c. = G. oblonga.
ovata, Endl. l. c. = G. ovata.

FAENICULUM, Hill, Hort. Kew. 103 (1769) = **Foeniculum**, Tourn. (Umbell.).

FAETIDIA, Comm. ex Juss. Gen. 325 (1789) = **Foetidia**, J. F. Gmel. (Myrtac.).

FAGARA, Linn. Syst. ed. X. 897 (1759) = **Xanthoxylum**, Linn. (Rutac.).
acuminata, Sw. Prod. Veg. Ind. Occ. 38 = Z. acuminatum.
affinis, Schult. Mant. iii. 224 = Z. affine.
alata, Eckl. & Zeyh. Enum. 118 = Z. alatum, *Steud. Ariscorina*, Kuxb. Hort. Beng. 11.—Ind. or.
arnata, Thunb. Prod. Pl. Cap. 28 = Z. Thunbergii.
australis, Cunn. ex Steud. Nom. ed. II. i. 622 = Z. australe.
Avicennæ, Lam. Encyc. ii. 445 = Z. Avicennæ.
Budrunga, Roxb. Hort. Beng. 11; Fl. Ind. i. 417 = Z. Budrunga.
capensis, Thunb. Diss. Med. Afric. 6 (= *Elaphrium capense*).—Afr. austr.
decandra, Blanco, Fl. Filip. ed. I. 66 = Buchanania florida.
dubia, Willd. ex Schult. Mant. iii. 228 = Fraxinus Schiedeana.
dumosa, Griseb. in Mem. Am. Acad. N. S. vii. (1861) 170 = Z. dumosum.
Elaphrium, Willd. Sp. Pl. i. 668 = Canarium mauritianum.
emarginata, Sw. Prod. Veg. Ind. Occ. 38 = Z. emarginatum.
Enodia, Linn. f. Suppl. 195 = Evodia hortensis.
frazinifolia, Lam. Illustr. i. 334 = Z. Clava-Herculis.
glabra, Blume, Cat. Gew. Buitenz. 40 = Evodia glabra.
gracilis, Griseb. Cat. Pl. Cub. 48 = Z. gracile.
guianensis, Lam. Encyc. ii. 446 = Z. hermaphroditum.
heterophylla, Lam. l. c. 448 = Z. heterophyllum.
horrida, Thunb. in Trans. Linn. Soc. ii. (1794) 329 = Z. horridum (*sed* Gleditsia japonica).
inermis, Willd. ex Schult. Mant. iii. 227 = Z. Caribæum.
lentiscifolia, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 165 = Z. Pterota.

FAGARA :—
Lunu-Ankenda, Gaertn. Fruct. i. 334 = Evodia Roxburghiana.
martinicensis, Lam. Illustr. i. 334 = Zanthoxylum martinicense.
microphylla, Desf. Tabl. ed. I. 300 = Z. tragodes.
monophylla, Lam. Illustr. i. 334 = Z. ochroxylum.
nitida, Roxb. Hort. Beng. 11; Fl. Ind. i. 419 = Z. nitidum.
obtusifolia, Schult. Mant. iii. 222 = Evodia obtusifolia.
octandra, Blanco, Fl. Filip. ed. I. 64 = Melicope ternata.
octandra, Linn. Mant. i. 40 = Bursera tomentosa.
pentandra, Aubl. Pl. Gui. i. 78 = Z. hermaphroditum.
peruviana, Willd. ex Schult. Mant. iii. 228 = Z. affine
phylloptera, Griseb. Cat. Pl. Cub. 48 = Z. phylloptherum.
pimpinelloides, Lam. Illustr. i. 334 = Z. pimpinelloides.
piperita, Blanco, Fl. Filip. ed. I. 64 = Z. oxyphyllum.
piperita, Linn. Syst. ed. X. 897 = Z. pimpinelloides.
piperita, Lour. Fl. Cochinch. i. 101 = Z. nitidum.
Pterota, Blanco, Fl. Filip. ed. I. 66 = Z. Aviennæ.
Pterota, *Linn. Syst.* ed. X. 897 = Z. Pterota.
Rhetsa, Roxb. Hort. Beng. 11; Fl. Ind. i. 417 = Z. Rhetsa.
schinifolia, Salisb. Prod. 67 = Z. tragodes.
spinifex, Jacq. Fragm. 10 = Z. spinifex.
spinosa, Sw. Prod. Veg. Ind. Occ. 38 = Z. spinosum.
stenoptera, Griseb. Cat. Pl. Cub. 48 = Z. stenopterum.
tragodes, Jacq. Enum. Pl. Caribæ. 19 = Z. tragodes.
trifoliata, Prod. Veg. Ind. Occ. 38 = Z. ternatum.
triphylla, Lam. Encyc. ii. 447 = Evodia triphylla.
triphylla, Lour. Fl. Cochinch. 82.—Cochinch.
triphylla, Roxb. Hort. Beng. 11; Fl. Ind. i. 416 = Evodia Roxburghiana ?
xanthoxyloides, Lam. Encyc. ii. 446 = Z. senegalense.
zeylanica, J. F. Gmel. Syst. 258 = Evodia Roxburghiana.

FAGARASTRUM, G. Don, Gen. Syst. ii. 67 (1832) = **Clausena**, Burm. f. (Rutac.).
anisatum, G. Don, Gen. Syst. ii. 87 = C. anisata.
capense, G. Don, l. c. = Zanthoxylum capense.
inæquale, G. Don, l. c. = C. inæqualis.
natalense, Walp. Rep. iv. 413, Index = Casetis natalensis.
Thunbergii, G. Don, Gen. Syst. ii. 87 = Zanthoxylum Thunbergii.

FAGELLIA, Neck. Elem. iii. 41 (1790). *LEGUMINOSÆ*, Benth. & Hook. f. i. 541.
luminosa, *DC. Prod.* ii. 475.—Afr. austr.
flexuosa, Meisen. in Hook. Lond. Journ. Bot. ii. (1843) 93 = luminosa.
pubescens, Eckl. & Zeyh. Enum. 257 = Rhynchosia viscidula.
resinosa, Hochst. in Flora, xxiv. (1841) 1. Intell. 32 = Rhynchosia resinosa.
viscida, Eckl. & Zeyh. Enum. 257 = Rhynchosia viscidula.

FAGELIA, Schwencke, in Verh. Bat. Gen. Roterd. i. (1774) 274. t. 13 = **Calosolaria**, Linn. (Scrophul.).
flavicans, J. F. Gmel. Syst. 40 = C. pinnata.

FAGOIDES, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. ii. (1839) 210 = **Ahouai**, A. Cunn. (Caprifol.).
trilobe, Banks & Soland. ex A. Cunn. l. c. = A. Banksii.

FAGONIA, Tourn. ex Linn. Gen. ed. I (1735). *ZYGOPHYLLEÆ*, Benth. & Hook. f. i. 267.
acerosa, Botss. Diagn. Ser. I. viii. 124 = cretica.
arabica, Linn. Sp. Pl. 386 = cretica.
armata, R. Br. in Salt, Abyss. App. 64 = cretica.
aspera, *C. Gay, Fl. Chil.* i. 470.—Chili.
Aucheri, Botss. Diagn. Ser. I. i. 97 = cretica.
Bruguieri, DC. Prod. i. 704 = cretica.
californica, Benth. Bot. Voy. Sulph. 10 = cretica.
chilensis, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 165 = cretica.
cisoides, Delile, ex T. Anders. in Journ. Linn. Soc. v. Suppl. 1. (1860) 11 = cretica.
cretica, *Linn. Sp. Pl.* 384.—Europ. austr. ; Orient ; Afr. bor. ; Am. bor. et austr.
deflexa, Moench, Meth. Suppl. 82 = cretica.

FAGONIA :—
diversifolia, Boiss. Diagn. Ser. II. i. 113 = cretica.
echinella, Boiss. l. c. Ser. I. viii. 123 = cretica.
Ehrenbergii, Schweinf. in Verh. Zool.-Bot. Ges. Wien, xviii. (1868) 639 = cretica.
elongata, Salisb. Prod. 319 = cretica.
cretica, Mill. Gard. Dict. ed. VIII. n. 1 = cretica.
Forskhlii, Pfund, in Flora, vii. (1874) 414 = cretica.
fruticosa, Coss. ex T. Anders. in journ. Linn. Soc. v. Suppl. i. (1860) 13 = cretica.
getula, Pomel, Nouv. Mat. Fl. Atl. 888 = cretica.
glutinosa, Delile, Fl. Egypte, 230. t. 28. f. 2 = cretica.
grandiflora, Boiss. Diagn. Ser. I. viii. 121 = cretica.
hispanica, Linn. Sp. Pl. 386 = cretica.
indica, Burm. f. Fl. Ind. 109 = cretica.
kahirina, Boiss. Diagn. Ser. I. viii. 122 = cretica.
latifolia, Delile, Fl. Egypte, 230. t. 28. f. 3 = cretica.
Mayana, Schlecht. in Bot. Zeit. ix. (1851) 844 = cretica.
microphylla, Pomel, Nouv. Mat. Fl. Atl. 888 = cretica.
mollis, Delile, Fl. Egypte, 230. t. 28. f. 3 = cretica.
montana, Miq. Fl. Ind. Bat. i. ii. 596 = Azima tetracantha.
myriacantha, Boiss. Diagn. Ser. I. viii. 123 = cretica.
myoornasis, Heyne, ex Roth, Nov. Sp. Pl. 215 = cretica.
Olivieri, DC. Prod. i. 704 = cretica.
orientalis, *Prsl. Delic. Prag.* 38.—Oriens.
parviflora, Boiss. Diagn. Ser. I. viii. 124 = cretica.
persica, DC. Prod. i. 704 = cretica.
scabra, Forsk. Fl. Aegypt.-Arab. 88 = cretica.
Schimperi, Prsl, Bot. Bemerk. 30 = cretica.
sinaica, Boiss. Diagn. Ser. I. i. 61 = cretica.
subinermis, Boiss. l. c. 60 = cretica.
tenuifolia, Stend. & Hochst. ex Boiss. Fl. Orient. i. 909 = cretica.
thebaica, Boiss. Diagn. Ser. I. viii. 122 = cretica.
virens, Coss. ex T. Anders. in journ. Linn. Soc. v. Suppl. i. (1860) 12 = cretica.
viscida. Prsl, Bot. Bemerk. 30 = cretica.
viscosa, Hochst. ex Boiss. Fl. Orient. i. 904 = cretica.

FAGOPYRUM, Tourn. ex Hall. Enum. Stirp. Helv. i. 172 (1742). *POLYGONACEÆ*, Benth. & Hook. f. iii. 99.
FAGOTRITICUM, Linn. in Vet. Acad. Handl. (1744) 117. t. 4.
KUNOKALE, Rafin. Fl. Tellur. iii. 12 (1836).
PHEGOPYRUM, Peterm. Fl. Bienitz, 92 (1841).
carinatum, Moench, Meth. 290 = Polygonum Convolvulus.
cereale, Rafin. Fl. Tellur. iii. 10 = esculentum.
chinense, Rafin. l. c. = cymosum ?
crassifolium, *Rafin. l. c.*—Hab. ?
cymosum, *Meisn. in Wall. Pl. As. Rar.* iii. 63.—Reg. Himal.; China.
dentatum, Moench, Meth. 290 = tataricum.
Doyendi, Hort. ex Fenzl, in Linnæa, xxiv. (1851) 282 = esculentum.
dumetorum, Schreb. Spicil. Fl. Lips. 42 = Polygonum dumetorum.
emarginatum, Moench, Meth. Suppl. 100 = esculentum.
esculentum, *Moench, Meth.* 290.—Europ.; As. bor.
membranaceum, Moench, l. c. = Polygonum dumetorum.
perfoliatum, Rafin. Fl. Tellur. iii. 10 = Polygonum perfoliatum !
rotundatum, Bab. in Trans. Linn. Soc. xvii. (1841) 137 = tataricum.
sagittatum, Gilib. Exercit. ii. 485 = esculentum.
sarracenicum, Dum. Fl. Belg. Prod. 12 = esculentum.
subdentatum, Gilib. Exercit. ii. 486 = esculentum !
suffruticosum, *F. Schmidt, Reis. Amuri.* 170.—Ins. Sachalin.
tataricum, *Gaertn. Fruct.* ii. 182. t. 119. f. 6.—Europ.; As. bor.
triangulare, Meisn. ex Wall. Pl. As. Rar. iii. 63 = cymosum.
vulgare, T. Ness, Gen. Pl. Monochl. 58 = esculentum.

FAGOTRITICUM, Linn. in Vet. Acad. Handl. Stockh. (1744) 117. t. 4 = **Fagopyrum**, Tourn. (Polygonac.).

FAGRÆA, Thunb. in Vet. Acad. Handl. Stockh. (1782) 236 ; et Nov. Gen. 24 (1783). *LOGANIACEÆ*, Benth. & Hook. f. ii. 704.

FAGRÆRA :—
CYRTOPHYLLUM, Reinw. ex Blume, Cat. Gew. Buitenz. 47 (1823) ; et Bijdr. 1092 (1826).
KENTIA, Stend. Nom. ed. II. i. 845 (1840).
KUMLIA, Reinw. in Blume, Cat. Gew. Buitenz. 51 (1823).
PICROPHLOEUS, Blume, Bijdr. 1019 (1826).
UTANIA, G. Don, Gen. Syst. iv. 668 (1837).
amboinensis, *Blume, Mus. Bot. Lugd. Bat.* i. 166.—Ins. Amboin.
appendiculata, Blume, l. c. 169 = morindæfolia.
auricularia, Benth. in journ. Linn. Soc. i. (1857) 96 = scq.
auriculata, *Yach, in Malay. Misc.* ii. (1822) n. VII. 82. —Burma : Malaya.
Berteriana, *A. Gray, ex Benth. in Journ. Linn. Soc.* i. (1857) 98.—Ins. Pacific.
Blumei, G. Don, Gen. Syst. iv. 69 = obovata.
bornensis, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 31.—Borneo.
camosa, *Yach, in Malay. Misc.* ii. (1822) n. VII. 81.—Burma ; Malaya.
cellanica, *Thunb. in Vet. Akad. Handl. Stockh.* (1782) 182 ; *Nov. Gen.* 85.—Zeylan.
celebica, Blume, Mus. Bot. Lugd. Bat. i. 167 = obovata.
coarctata, Blume, Bijdr. 1021 = racemosa.
congesta, *Blume, Mus. Bot. Lugd. Bat.* i. 168.—Sumatra.
cordifolia, *Blume, Rumphia*, ii. 88.—Ins. Molucc.
coromandelina, Wight, Ic. iv. t. 1316 = obovata.
cvassifolia, Blume, Rumphia, ii. 31. t. 78. f. 1 = obovata.
crassipes, *Benth. in Journ. Linn. Soc.* i. (1857) 99.—Borneo.
crenulata, *Maingay, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 38.—Malacca.
cuspidata, Blume, Mus. Bot. Lugd. Bat. i. 170 = morindæfolia.
dubia, Wall. Cat. n. 4456 = Ternstroemia penangiana.
elliptica, *Roxb. Hort. Beng.* 84 ; *Fl. Ind.* i. 462.—Ins. Molucc.
euneura, *Scheff. in Flora*, lii. (1869) 308 ; *et in Tijdschr. Nederl. Ind.* xxxi. (1870) 20.—Ins. Molucc.
fastigiata, *Blume, Rumphia*, ii. 89, in obs. t. 76.—Ins. Sunda.
Forsteni, *Blume, Mus. Bot. Lugd. Bat.* i. 166.—Ins. Celebes.
fragrans, *Roxb. Hort. Beng.* 84 ; *Fl. Ind.* i. 461.—Burma ; Malaya.
fuscescens, *Blume, Mus. Bot. Lugd. Bat.* i. 168.—Sumatra.
Gaudneri, Rhw. Enum. Pl. Zeyl. 200 = obovata.
globosa, Wall. Cat. n. 1598 = obovata.
gracilipes, *A. Gray, in Proc. Am. Acad.* iv. (1858–59) 353.—Ins. Pacific.
gracilis, *Franch. & Sebert, Not. Bois Nouv. Calédon.* 184. —N. Caled.
heterophylla, *Blume, Mus. Bot. Lugd. Bat.* i. 168.—*imperialis*, Miq. Fl. Ind. Bat. ii. 872 = auriculata.
khasiana, *Benth. in Journ. Linn. Soc.* i. (1857) 99.—Reg. Himal.
Kienaeg, *Blume, Mus. Bot. Lugd. Bat.* i. 173.—Java.
lanceolata, *Blume, Bijdr.* 1021.—Java.
laxiflora, Wall. Cat. n. 1599 = Wallichiana.
latifolia, *Miq. Fl. Ind. Bat.* ii. 869.—Sumatra.
Laschnaniti, *Blume, Rumphia*, ii. 30, in obs.—Java.
ligustrina, *Blume, l. c.* 88.—Malaya.
litoralis, *Blume, Bijdr.* 1021.—Java.
litoralis, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 168 = heterophylla.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 84.—Malaya.
malabarica, Blume, Rumphia, ii. 29 = obovata.
malabarica, *Wight, Ic.* t. 1317.—Ind. or.
malayana, *Mart. Nov. Gen. et Sp.* ii. 91.—Malaya.
minor, *Reinw. ex Blume, Bijdr.* 1021.—Java.
monantha, *Miq. Fl. Ind. Bat.* ii. 872.—Sumatra.
morindæfolia, *Blume, Rumphia*, ii. 28. t. 78.—Malaya ; Austral.
Moolleri, *Benth. Fl. Austral.* iv. 362.—Austral.
obovata, Griff. Notul. iv. 85 = malabarica.
obovata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 88.—Ind. or. ; Malaya.
obovato-spicata, Blume, Rumphia, ii. 29 = pauciflora.
oxyphylla, *Miq. Fl. Ind. Bat.* ii. 871.—Malaya.

FAGRÆRA :—
peregrina, Blume, Rumphia, ii. 34. t. 80 = fragrans.
picrophloeos, *Blume, l. c.* 36.—Borneo.
plumeriæflora, *DC. Prod.* ix. 29 —Ins. Philipp.
plumeriæfolia, Benth. in journ. Linn. Soc. i. (1857) 98 = grac.
racemosa, *Yach, in Roxb. Fl. Ind. ed. Carey*, ii. 85.—Malaya.
robusta, Blume, Mus. Bot. Lugd. Bat. i. 170 = morindæfolia.
rostrata, *Blume, Rumphia*, ii. 31. t. 76. f. 2.—N. Guinea.
scholaris, Blanco, Fl. Filip. ed. II. 98 = cordifolia.
species, *Blume, Rumphia*, ii. 35. t. 81.—Java.
splendens, *Blume, Mus. Bot. Lugd. Bat.* i. 168.—Borneo.
subreticulata, *Blume, l. c.* 171.—Java.
sumatrana, *Miq. Fl. Ind. Bat.* ii. 877.—Sumatra.
ternatana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 217.—Ins. Ternate.
tetragona, *Span. in Linnæa*, xv. (1841) 226.—Ins. Timor.
Thwaitesii, F. Muell. Fragm. ii. 137 = morindæfolia.
truncata, *Blume, Mus. Bot. Lugd. Bat.* i. 165.—Ins. Celebes.
tubuliosa, *Blume, l. c.* 167.—Sumatra.
valida, *Miq. Fl. Ind. Bat.* ii. 876.—Sumatra.
viridiflora, Seem. in Bonplandia, ix. (1861) 259 = grandiflora.
vitiensis, Seem. Viti, 459, nomen = Olea vitiensis.
volubilis, Jack, in Roxb. Fl. Ind. ed. Carey, ii. 86 = racemosa.
Wallichiana, *Benth. in Journ. Linn. Soc.* i. (1857) 96. —Ins. Penang.
zeylanica, Sesm. Fl. Vit. 165, in obs. = scq.
zeylanica, Murr. Syst. Veg. ed. XIV. 198 = cellanica.

FAGUETTIA, March. Rev. Anacard. 174. t. 2 (1869) *ANACARDIACEÆ*.
falcata, *March. l. c.* 175.—Madag.

FAGUS, [Tourn.] Linn. Syst. ed. I (1735). *CUPULIFERÆ*, Benth. & Hook. f. iii. 410.
CALUCECHINUS, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 19 (1853).
CALUSPARASSUS, Hombr. & Jacquinot, ex Decne. l. c. 20 (1853).
LOPHOZONIA, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 896.
MYRTILLOIDES, Banks & Soland. ex Hook. Ic. Pl. t. 639 (1844).
NOTHOFAGUS, Blume, Mus. Bot. Lugd. Bat. i. 307 (1850).
PHEGOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 138.
acuta, Dum.-Cours. Bot. Cult. ed. II. vi. 415 = sylvatica.
alba, Rafin. Fl. Ludov. 184 ; New Sylva, 80 = ferruginea.
alpina, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 69. t. 196, 198 *A.*—Patagon.
alpina, Poepp. ex A. DC. Prod. xvi. ii. 120 = antarctica.
americana, Sweet, Hort. Brit. ed. I. 370 = ferruginea.
antarctica, *Forst. f. in Comm. Gotting.* ix. (1789) 34.—Reg. Magellan.
apiculata, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 325.—N. Zel.
argentea, Blume, in Flora, vii. (1824) 291 = Castanopsis argentea.
asplenifolia, Dum.-Cours. Bot. Cult. ed. II. vi. 415 ; Rafin. New Fl. Am. iii. 82 = sylvatica.
betuloides, *Mirb. in Mém. Mus. Par.* xiv. (1827) 469. t. 25.—Reg. Magellan.
Blairii, *T. Kirk, in Trans. N. Z. Inst.* xvii. (1885) 397 ; 306.—N. Zel.
Carmichaelii, C. Moore, ex Benth. Fl. Austral. vi. 211 = Moorei.
Castanea, Linn. Sp. Pl. 997 = Castanea sativa.
chrysocoma, *Hook. f. in Hook. Ic. Pl.* t. 678 & t. 816. —N. Zel.
cochinchinensis, Lour. Fl. Cochinch. 571 = Castanea chinensis.
completifolia, Desf. Tabl. ed. II. 369 = sylvatica.
crenata, Blume, Mus. Bot. Lugd. Bat. i. 307 = Sieboldii.
cristata, Dum.-Cours. Bot. Cult. ed. II. vi. 415 = sylvatica.

FAGUS:—
Cunninghamii, *Hook. Journ. Bot.* ii. (1840) 169. t. 7. —Austral.
cupreo, Host. ex A. DC. in DC. Prod. xvi. ii. 119 = sylvatica.
Dombeyi, *Mirb. in Mém. Mus. Par.* xiv. (1827) 467. t. 24.—Chili.
dubia, Mirb. l. c. 471. t. 26 = betuloides.
echinata, Gilib. Exercit. ii. 396 = sylvatica.
ferruginea, [*Doyand. in*] *Ait. Hort. Kew.* ed. I. iii. 362.—Am. bor.
ferruginea, Siebold, in Verh. Batav. Gen. xii. (1830) 22 = Sieboldii.
Forsteri, Hook. Journ. Bot. ii. (1840) 156 = betuloides.
fusca, *Hook. f. in Hook. Ic. Pl.* t. 630, 631.—N. Zel.
glauca, *Phil. in Linnaea*, xxix. (1857–58) 48.—Chili.
glutinosa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 68. t. 194.—Chili.
grandifolia, Ehrh. Beitr. 22 = ferruginea.
Gunnii, *Hook. f. in Hook. Ic. Pl.* t. 881.—Tasman.
heterophylla, Raßn. Herb. 22 = ferruginea.
javanica, Blume, in Flora, vii. (1824) 295 = Castanopsis javanica.
laciniata, Hort. ex A. DC. in DC. Prod. xvi. ii. 119 = sylvatica.
latifolia, L'Hérit. ex Steud. Nom. ed. II. i. 624 = ferruginea.
lutea, Mœnch, ex C. Gay, Fl. Chil. i. 438 = Zanthoxylum Maya.
Menziesii, *Hook. f. in Hook. Ic. Pl.* t. 852.—N. Zel.
Montagnii, *Homb. & Jacquinot, Bot. Voy. Pole Sud*, t. 8. f. 11.—Chili.
Moorei, *F. Muell. Fragm.* v. 109.—Austral.
nana, Durol, ex Steud. Nom. ed. II. i. 624 = Castanea pumila.
nervosa, *Phil. in Linnaea*, xxix. (1857–58) 48.—Chili.
nigra, Raßn. Herb. 22; New Fl. Am. iii. 81 = ferruginea.
nitida, *Phil. in Linnaea*, xxix. (1857–58) 44.—Chili.
obliqua, *Mirb. in Mém. Mus. Par.* xiv. (1827) 465. t. 23.—Chili.
pendula, Dum.-Cours. Bot. Cult. ed. II. vi. 415 = sylvatica.
phillyraeoides, Blanco, Fl. Filip. ed. II. 508 = Castanopsis sumatrana.
procera, Phil. ex A. DC. in DC. Prod. xvi. ii. 120 = obliqua.
procera, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 69. t. 197. —Chili.
procera, Salisb. Prod. 391 = Castanea sativa.
pumila, Linn. Sp. Pl. 998 = Castanea pumila.
Pumilio, Poepp. & Endl. Nov. Gen. et Sp. ii. 68. t. 195 = antarctica.
purpurea, Desf. Tabl. ed. I. 214 = ferruginea.
purpurea, Dum.-Cours. Bot. Cult. ed. II. vi. 415 = sylvatica.
retusifolia, Raßn. Herb. 22; Atl. Journ. 177 = ferruginea.
salicifolia, Hort. ex A. DC. in DC. Prod. xvi. ii. 119 = sylvatica.
Sieboldii, *Endl. Gen. Suppl.* iv. ii. (1847) 29.—Japon.
Solandri, *Hook. f. in Hook. Ic. Pl.* t. 639 et t. 816.—N. Zel.
sylvatica, *Linn. Sp. Pl.* 998.—Europ.
sylvestris, Gaertn. Fruct. i. 182 = prœc.
sylvestris, Mœnch. Hist. Arbr. Am. Sept. ii. 170 = ferruginea.
uliginosa, Phil. ex A. DC. in DC. Prod. xvi. ii. 120 = antarctica.
waldiviana, Phil. in Linnaea, xxxiii. (1864–65) 226 = obliqua.
variegata, Hort. ex A. DC. in DC. Prod. xvi. ii. 119 = sylvatica.
virginiana, Hort. ex Wesmael, in Bull. Congr. Bot. Brux. (1864) 261 = ferruginea.

FAHRENHEITIA, Reichb. f. & Zoll. in Linnaea, xxviii. (1856) 600 = Ostodes, Blume (Euphorb.).
colitua, Reichb. f. & Zoll. l. c.—Java.

FAKELOBA, Raßn. Spira Tellur. 69 (1888) = Anthyllis, Linn. (Legumin.).
cretica, Raßn. l. c. = A. Hermanniae.

FALCARIA, Riv. ex Rapp. Fl. Jen. ed. Hall. 279 (1745); Host, Fl. Austr. i. 381 (1827). *UMBELLIFERÆ*, Benth. & Hook. *f.* i. 892.
CASTAMBA, Bess. Enum. Pl. Volh. 98 (1821).
DREPANOPHYLLUM, Hoffm. Gen. Umb. ed. I. 109 (1814), partim.
PRIONITIS, Adans. Fam. ii. 499 (1763).
carvifolia, *C. A. Mey. Beitr. Pfl. Russl.* i. 14.—Rossia.
falkurica, Dec. Prod. iv. 110 = Apium linearæ.
filcarifolia, DC. l. c. = Carum filcarifolium.
glauca, Dulac, Fl. Hautes-Pyr. 351 = vulgaris.
[*graveolens*, *DC. Prod.* iv. 110.—Java.
graveolens, *DC. l. c.*—Java.
latifolia, Koch, Syn. Fl. Germ. ed. II. 313 = Carum latifolium.
pastinacifolia, Reichb. f. Ic. Fl. Germ. xxxi. 12. t. 1958 = Carum latifolium.
Rivini, Host, Fl. Austr. i. 381 = vulgaris.
Sautiyage, Reichb. f. Ic. Fl. Germ. xxxi. 13 = Carum Bauhini.
sioides, Aschers. Fl. Brandenb. 241 = vulgaris.
vulgaris, *Bernh. Syst. Verz. Erf.* 176.—Europ.; Orient.

FALCATA, J. F. Gmel. Syst. 1181 (1791) = Amphicarpæa, Ell. (Legumin.).
caroliniana, J. F. Gmel. l. c. = A. monoica.

FALCATULA, Brot. Phyt. Lusit. 160. t. 88 (1801) = Trigonella, Linn. (Legumin.).
falcatrifolium, Brot. l. c. = T. ornithopodioides.
fœnigræca, Four. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 360 = T. monspeliaca.

FALCKIA, Thunb. Nov. Gen. 17 (1781) = **Falkia**, Linn. f. (Convolv.).

FALCONERA, Salisb. Gen. Pl. Fragm. 36 (1866) = Albuca, Linn. (Liliac.).
fastigiata, Salisb. l. c. = A. fastigiata.
spiralis, Salisb. l. c. = A. spiralis.
viscosa, Salisb. l. c. = A. viscosa.

FALCONERIA, Wight, Ic. v. 20 (1859) = FALCONERIA, Royle = Sapium, P. Br. (Euphorb.).

FALCONERIA, Hook. f. Ic. Pl. t. 1488 (1888).
SCROPHULARINEÆ.
himalaica, *Hook. f. l. c.*—Reg. Himal.

FALCONERIA, Royle, Illustr. Bot. Himal. 354. t. 84 a (1839) = Sapium, P. Br. (Euphorb.).
insignis, Royle, l. c. = S. insigne.
malabarica, Wight, Ic. t. 1866 = S. insigne.
Wallichiana, Royle, Illustr. Bot. Himal. 354 = S. insigne.

FALDERMANNIA, Trautv. in Bull. Sc. Acad. Pétersb. vi. (1840) 185 = Staphorea, Linn. (Rubiac.).
pauciflora, Trautv. l. c. vii. (1841) 21 = Z. tenuior.
taurica, Trautv. l. c. vi. (1840) 186 = Z. taurica.

FALIMIRIA, Bess. ex Reichb. Consp. 54 (1828) = Gentzinia, Besser. (Gramin.).
bromoides, Bess. ex Steud. Nom. ed. II. i. 635 = G. geminiflora.
scoarciata, Bess. ex Steud. l. c. = G. geminiflora.

FALKEA, Kom. ex Steud. Nom. ed. II. i. 626, 194 (1840) = Begonia, Linn.
tenera, Kom. ex Steud. l. c. = B. tenera.

FALKIA, Linn. f. Suppl. 80 (1781). *CONVOLVULACEÆ*, Benth. & Hook. f. ii. 876.
FALCKIA, Thunb. Nov. Gen. 17 (1781).
oblonga, *Bernh. in Flora*, xxvii. (1844) 890.—Afr. austr.
repens, *Linn. f. Suppl.* 211.—Afr. austr.

FALLOPIA, Adans. Fam. ii. 274, 277 (1763) = Brunnichia, Banks (Polygon.).

FALLOPIA, Lour. Fl. Cochinch. 625 (1790) = Cissampelos, Linn. (Tiliac.).
nervosa, Lour. l. c. 626 = C. micrantha.

FALLUGIA, Endl. Gen. 1246 (1840). *ROSACEÆ*, Benth. & Hook. f. i. 618.
mexicana, Walp. Rep. ii. 16 = paradoxa.
paradoxa, *Endl. Gen.* 1246.—N. Mexic.

FALONA, Adans. Fam. ii. 496 (1763) = Cynosurus, Linn. (Gramin.).

FALOPIA, Steud. Nom. ed. I. 333 (1821) = FALLOPIA, Adans. = Brunnichia, Banks (Polygon.).

FAMAREA, Vitm. Summa, i. 257 (1790), sphalm. = **Faramea**, Aubl. (Rubiac.).

FANNINIA, Harv. Gen. S. Afr. Pl. ed. II. 225 (1868). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 740.
caloglossa, *Harv. l. c.*—Afr. austr.

FARADAYA, F. Muell. Fragm. v. 21 (1865). *VERBENACEÆ*, Benth. & Hook. f. ii. 1154.
amicorum, Seem. in Journ. Bot. iii. (1865) 258 = Clerodendron amicorum.
ocellifolia, *Seem. l. c.*—Ins. Fiji.
papuana, *Scheff. in Ann. Yard. Buitenz.* i. (1876) 42. —N. Guinea.
spicuidis, *F. Muell. Fragm.* v. 21.—Austral.
vitiensis, *Seem. in Journ. Bot.* iii. (1865) 258.—Ins. Fiji.

FARAMEA, Aubl. Pl. Gui. i. 102. t. 40 (1775). *RUBIACEÆ*, Benth. & Hook. f. ii. 121.
ANTONIANA, Tuss. Fl. Antill. ii. 62. t. 16 (1818).
DARLUCA, Raßn. in Ann. Gén. Sc. Phys. vi. (1820) 87.
ENCOPEA, Presl, in Abh. Bohm. Ges. V. iii. (1845) 513 (Bot. Bemerk. 88).
FAMAREA, Vitm. Summa, i. 257 (1790).
HOMALOCLADOS, Hook. f. in Benth. & Hook. f. Gen. ii. 122 (1873).
NELREA, Raßn. New Fl. Am. iv. 55 (1836).
OMALOCLADOS, Hook. f. Ic. Pl. t. 1128 (1878).
POTIMA, Hedw. f. Gen. 148 (1806).
TETRAMERIUM, Gaertn. f. Fruct. iii. 90. t. 196 (1806).
acuminatissima, *Muell. Arg. in Flora*, lviii. (1875) 471.—Bras.
affinis, *Muell. Arg. l. c.* 469.—Bras.
albescens, DC. Prod. iv. 498 = Coussarea albescens.
albescens, Mart. in Flora, xxiv. (1841) ii. Beibl. 72 = Coussarea lilacina.
amazonica, *Muell. Arg. in Flora*, lviii. (1875) 470.—Bras.
angustifolia, *Spruce, ex Benth. & Hook. f. Gen.* ii. 121; *Muell. Arg. l. c.* 471.—Bras.
anisocalyx, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 38.—Peruv.
anisodonta, *Muell. Arg. in Flora*, lviii. (1875) 472.—Bras.
apodantha, *Muell. Arg. l. c.* 471.—Bras.
aristata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 460.—Bras.
australis, *Muell. Arg. in Flora*, lviii. (1875) 472.—Bras.
axilliflora, *DC. Prod.* iv. 497.—Bras.
axilliflora, Mart. in Flora, xxiv. (1841) ii. Beibl. 72 = Ixora bahiensis.
bahiensis, *Muell. Arg. in Flora*, lviii. (1875) 474.—Bras.
Blanchetiana, *Muell. Arg. l. c.* 471.—Bras.
brachyloba, *Muell. Arg. l. c.* 470.—Bras.
brachya, *Benth. in Linnaea*, xxiii. (1850) 452.—Bras.
calycidiora, *A. Rich. in Mém. Soc. Hist. Nat. Par.* vi. (1834) 176.—Guian.
campanulis, *Muell. Arg. in Flora*, lviii. (1875) 469.—Bras.
campestraria, *Muell. Arg. l. c.* 470.—Bras.
capillipes, *Muell. Arg. l. c.* 474.—Bras.
capitata, Benth. in Linnaea, xxiii. (1850) 450 = Coussarea capitata.
candata, *Muell. Arg. in Flora*, lviii. (1875) 471.—Bras.
candata, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 106.—Bras.
candelabrica, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 458. —Bras.
cauvilea, *DC. Prod.* iv. 497.—Bras.

FARAMEA:—
colorata, Benth. in Linnæa, xxiii. (1850) 452 = calycifloru.
contracta, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 851 = Cossæa contracta.
cornifolia, Benth. in Linnæa, xxiii. (1850) 450 = Cossæa cornifolia.
coronata, *Muell. Arg. in Flora*, lviii. (1875) 470.—Bras.
corymbosa, *Aubl. Pl. Gui.* i. 102.—Guiana.
corymbosa, Benth. in Hook. Journ. Bot. iii. (1841) 288 = Cossæa violacea.
corymbosa, Sieber, ex DC. Prod. iv. 463 = Myonima multiflora.
crassifolia, *Benth. in Hook. Journ. Bot.* iii. (1841) 288.—Guiana.
cuspidata, *Benth. Pl. Hartw.* 194.—N. Granat.
cyanea, *Muell. Arg. in Flora*, lviii. (1875) 473.—Bras.
diversifolia, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 136.—Bras.
egensis, *Muell. Arg. in Flora*, lviii. (1875) 470.—Bras.
Eichleri, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 109.—Bras.
erythrocarpa, *Griseb. Cat. Pl. Cub.* 134.—Cuba.
erythropoda, *Miq. in Linnæa*, xvii. (1844) 615.—Guiana.
fallax, *Muell. Arg. in Flora*, lviii. (1875) 474.—Bras.
filamentosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 156.—Bras.
filipes, Mart. ex Benth. in Linnæa, xxiii. (1850) 454 = cornifolia.
glandulosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 29.—Peruv.
Glaziovii, *Muell. Arg. in Flora*, lviii. (1875) 472.—Bras.
Godetiana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 153.—Bras.
gracilifiora, Mart. in Flora, xxiv. (1841) ii. Beibl. 72 = Cossæa gracilifiora.
guayaquilensis, *DC. Prod.* iv. 497.—Ecuador.
herbacea, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 176.—Guiana.
heterocalyx, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 158.—Bras.
heteromera, *Muell. Arg. in Flora*, lviii. (1875) 469.—Bras.
heterophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 151.—Bras.
hyacinthina, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 73.—Bras.
hydrangeæfolia, Benth. in Linnæa, xxiii. (1850) 451 = Cossæa hydrangeæfolia.
includens, *Muell. Arg. in Flora*, lviii. (1875) 469.—Bras.
interodens, *Muell. Arg. l. c.* 470.—Bras.
involucellata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 157.—Bras.
jambodes, *DC. Prod.* iv. 497.—N. Granat.
lagoënsis, *Muell. Arg. in Flora*, lviii. (1875) 474.—Bras.
latifolia, *DC. Prod.* iv. 497.—Bras.
leptobotrys, Benth. & Hook. f. Gen. ii. 121 = Cossæa leptobotrys.
leucocalyx, *Muell. Arg. in Flora*, lviii. (1875) 474.—Bras.
longiflora, Mart. in Flora, xxiv. (1841) ii. Beibl. 74 = Cossæa longiflora.
longifolia, *Benth. in Hook. Journ. Bot.* iii. (1841) 284.—Guian.
luciana, *Muell. Arg. in Flora*, lviii. (1875) 474.—Bras.
macra, *Muell. Arg. l. c.* 472.—Bras.
macrocalyx, *Muell. Arg. l. c.* 469.—Bras.
macrophylla, Mart. in Flora, xxiv. (1841) ii. Beibl. 75 = Cossæa macrophylla.
manulicoides, Muell. Arg. in Flora, lviii. (1875) 473 = pachyantha.
marginata, *Chem. in Linnæa*, ix. (1834) 221.—Bras.
martiana, Mart. in Flora, xxiv. (1841) ii. Beibl. 74 = Martiana.
Martiana, *Muell. Arg. in Flora*, lviii. (1875) 472.—Bras.
Meribal, DC. Prod. iv. 496 = Cossæa Martini.
mexicana, *Spruce, ex Benth. & Hook. f. Gen.* ii. 121, comm.—Peruv.
micrantha, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 136.—Bras.
microcalyx, *Muell. Arg. l. c.* 136.—Bras.

FARAMEA:—
mosantha, *Muell. Arg. in Flora*, lviii. (1875) 475.—Bras.
monovidensis, *DC. Prod.* iv. 497.—Reg. Argent.
Moritziana, *Benth. in Linnæa*, xxiii. (1850) 451.—N. Granat.
multiflora, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 176.—Guian.
nemoralis, Mart. in Flora, xxiv. (1841) ii. Beibl. 73 = Cossæa nemoralis.
Nettoana, *Muell. Arg. in Flora*, lviii. (1875) 473.—Bras.
nigrescens, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 73.—Bras.
nitida, *Benth. in Linnæa*, xxiii. (1850) 454.—Bras.
nodosa, Benth. l. c. 449 = Cossæa nodosa.
obtusifolia, *Muell. Arg. in Flora*, lviii. (1875) 472.—Bras.
occidentalis, *Muell. Arg. l. c.* 474.—Bras.
occidentalis, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 176 = odoratissima.
odoratissima, *DC. Prod.* iv. 496.—Am. austr.
oliganthа, *Muell. Arg. in Flora*, lviii. (1875) 474.—Bras.
oxycladia, *Muell. Arg. l. c.*—Bras.
pachyantha, *Muell. Arg. l. c.* 470.—Bras.
pachydictyon, *Muell Arg. l. c.* 471.—Bras.
paniculata, *Benth. in Linnæa*, xxiii. (1850) 451.—Guian.
panurensis, *Muell. Arg. in Flora*, lviii. (1875) 470.—Bras.
parviflora, *Muell. Arg. l. c.*—Bras.
pedicellaris, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 126.—Bras.
pendula, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 29.—Bras.
petiolaris, *Benth. in Linnæa*, xxiii. (1850) 450.—N. Granat.
platyclada, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 126.—Bras.
platyneura, *Muell. Arg. in Flora*, lviii. (1875) 470.—Bras.
platypoda, *Muell. Arg. l. c.* 472.—Bras.
Pohliana, *Muell. Arg. l. c.*—Bras.
porophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 116.—Bras.
pulchella, *Benth. & Hook. f. Gen.* ii. 121; *et ex Muell. Arg. in Flora*, lviii. (1875) 475.—Bras.
Purdieana, *Benth. in Linnæa*, xxiii. (1850) 458.—N. Granat.
quinqueflora, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 28. t. 284.—Peruv.
Riedeliana, *Muell. Arg. in Flora*, lviii. (1875) 471.—Bras.
rivularis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 108.—Bras.
salicifolia, *Presl, Symb. Bot.* ii. 24. t. 70.—Bras.
Schomburghiana, Benth. in Linnæa, xxiii. (1850) 450 = Cossæa violacea.
Sellowiana, *Benth. l. c.* 453.—Bras.
sertaliferа, *DC. Prod.* iv. 496.—Ind. occ.
sessiliflora, *Aubl. Pl. Gui.* i. 104.—Guian.
sessilifolia, *DC. Prod.* iv. 497.—Venezuela.
stenantha, *Muell. Arg. in Flora*, lviii. (1875) 473.—Bras.
stenopeciа, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 73.
stipulacea, *DC. Prod.* iv. 497.—Bras.
suaveolens, *Duchass. ex Griseb. in Bonplandia*, vi. (1858) 5.—Panama.
subsessilis, *Muell. Arg. in Flora*, lviii. (1875) 473.—Bras.
Tambelliana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 134.—Bras.
tenuiflora, *Muell. Arg. in Flora*, lviii. (1875) 474.—Bras.
tetragona, *Muell. Arg. l. c.* 472.—Bras.
tinguana, *Muell. Arg. l. c.* 471.—Bras.
torquata, *Muell. Arg. l. c.*—Bras.
truncata, *DC. Prod.* iv. 497.—Guiana.
trunculа, *Muell. Arg. in Flora*, lviii. (1875) 473.—Bras.
umbelliflora, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 134.—Bras.
urophylla, *Muell. Arg. in Flora*, lviii. (1875) 469.—Bras.

FARAMEA:—
vaginata, Benth. in Linnæa, xxiii. (1850) 454 = Irora pubescens.
vaginata, *Griseb. Cat. Pl. Cub.* 134.—Cuba.
videnaia, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 118.—Bras.
Warmingiana, *Muell. Arg. in Flora*, lviii. (1875) 472.—Bras.
xanthina, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 161.—Bras.

FAREINHETIA, Baill. Etud. Gén. Euph. 659 (1858) = FAHRENHEITIA, Reichb. f. & Zoll. = Ostodes, Blume (Euphorb.).

FARFARA, Gilib. Fl. Lituan. i. 177 (1781) = Tussilago, Linn. (Compos.).
radiata, Gilib. l. c. = T. Farfara.

FARFUGIUM, Lindl. in Gard. Chron. (1857) 4 = Senecio, Tourn. (Compos.).
grande, Lindl. l. c. = S. Kaempferi.
Kaempferi, Benth. Fl. Hongk. 191 = S. Kaempferi.

FARNESIA, Gasp. Desc. Nuov. Gen. (1838) = Acacia, Willd. 95 (Legumin.).
odora, Gasp. l. c. = A. Farnesiana.

FAROA, Welw. in Trans. Linn. Soc. xxvii. (1869) 45. t. 17. *GENTIANEAE*, Benth. & Hook. f. ii. 806.
salutaris, *Welw. l. c.* 46.—Afr. trop.

FAROBAEA, Schrank, ex Colla, Hort. Ripul. App. iii. 36, in note (1829); et iv. 19. t. 9 (1831) = Senecio, Tourn. (Compos.).
nemorensis, Schrank, ex Colla, l. cc. 36 et 21.—Hab.!
tropica, Colla, l. c. 36.—Hab.!

FARQUHARIA, Hilsenb. & Boj. ex Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 67 = Cratæva, Linn. (Capparid.).
excelsa, Hilsenb. & Boj. ex Boj. l. c.=C. excelsa.

FARSETIA, Turra, Farset. Nov. Gen. 5. t. 1 (1765).
CRUCIFERAE, Benth. & Hook. f. i. 72.
ACOSTON, Rafin. Sylva. Tellur. 121 (1838).
CYCLOCARPAEA, DC. in Mém. Mus. Par. vii. (1821) 232.
FISSIDIA, Medic. Pflanzeng. i. 90 (1792).
aegyptia, *Turra, Farset. Nov. Gen.* 5. t. 1.—Aegypt.
aegyptiaca, Desv. journ. Bot. iii. (1814) 307 = præc.
Boivinii, *Fourn. in Bull. Soc. Bot. Fr.* xi. (1864) 56.—Afr. trop.; Arab.; Ind. or.
Burtonae, *Oliver, in Hook. Ic. Pl.* t. 1310.—Arab.
cheiranthifolia, Desv. journ. Bot. iii. (1814) 178 = clypeata.
cheiranthoides, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 96 = aegyptia.
clypeata, [*R. Br.*] *l. c.*—Europ. austr.; Oriens.
dalmatica, Vis. in Flora, xii. (1829) i. Erg. 15; et in Fl. Dalmat. iii. 119. t. 82. f. 8 = triquetra.
deltoides, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 97 = Aubrietia deltoides.
depressa, *Kotschy, in Sitzb. Acad. Wien, Math. Nat.* lii. 1. (1866) 261.—Arab.
difusa, Desv. journ. Bot. iii. (1814) 178 = Heliophila diffusa.
Edgeworthii, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 147 = aegyptia.
elongata, Desv. journ. Bot. iii. (1814) 178 = Heliophila pectinata.
eriocarpa, *DC. Syst.* ii. 208.—Ae. Min.; Syria.
grandiflora, *Fourn. in Bull. Soc. Bot. Fr.* xi. (1864) 55.—Afr. trop.
Guentheriana, Andrz. ex DC. Syst. ii. 208 = Alyssum mutabile.
Hamiltonii, *Royle, Illustr. Bot. Himal.* 71.—Afr. bor.; Arab.; Ind. or.
incana, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 97 = Alyssum incanum.
Jacquemontii, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 148.—Ind. bor. occ.; Afghan.
libyca, Spreng. Syst. ii. 871 = Alyssum libycum.
linearis, Decne. ex Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 150 = Hamiltonii.

FARSETIA :—
linifolia, Andrz. ex DC. Syst. ii. 325 = Alyssum linifolium.
longisiliqua, *Decne. in Ann. Sc. Nat. Sér.* II. iv. (1835) 69.—Aegypt.
lunarioides, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 96. —Archip. Graec.
macrocarpa, *Boiss. in Ann. Sc. Nat. Sér.* II. xvii. (1842) 89.—Cappadoc.
macropteri, *Kotschy & Boiss. ex Fourn. in Bull. Soc. Bot. Fr.* xi. (1864) 60.—As. Min.
multicaulis, *Boiss. & Hohen. Diagn.* Ser. I. viii. 31.— Persia.
mutabilis, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 96 = Alyssum mutabile.
obliqua, Spreng. Syst. ii. 870 = Alyssum mutabile.
oblongata, *Presl, Bot. Bemerk.* 8.—Aegypt.
obovata, *Boiss. & Kotschy, Diagn.* Ser. II. v. 97.— Syria.
orbiculata, Spreng. Syst. ii. 870 = Alyssum orbiculatum.
ovalis, Boiss. Diagn. Ser. I. viii. 83 = aegyptia.
parviflora, Spreng. Syst. ii. 871 = Savignya aegyptiaca.
pendula, *Boiss. in Ann. Sc. Nat. Sér.* II. xvii. (1842) 89.—Persia.
peruviana, Spreng. Syst. ii. 871 (= *Berteroa peruviana*). —Peruv.
pinnata, Desv. Journ. Bot. iii. (1814) 178 = Heliophila Peltaria.
procurrata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 500.—Ins. Socotr.
ramosissima, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 42.—Afr. trop.
rostrata, *Schenk, Pl. Sp. Aegypt.* 42.—Egypt.; Syria.
sinuata, Roth, ex Stend. Nom. ed. II. i. 325 = Alyssum sinuatum.
spathulata, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 149.—Songaria.
stenoptera, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 42. —Abyss.
styloa, *T. Anders. in Journ. Linn. Soc. v. Suppl.* i. (1856) 1.—Arab.
styloa, *R. Br. in Denh. & Clapp. Trav. App.* 12.— Afr. trop.
suffruticosa, *DC. Syst.* ii. 287.—Persia.
triquetra, *DC. l. c.* 290.—Dalmat.
umbellata, *Boiss. Diagn.* Ser. I. vi. 13.—Persia.

FARTIS, Adans. Fam. ii. 37 (1763) = **Stizania**, Linn. (Gramin.).

FASCICULUS, Dulac, Fl. Hautes-Pyr. 345 (1867) = **Spergularia**, J. & C. Presl (Caryophyll.).
rubra, Dulac, l. c. = S. rubra.

FATERNA, Necuba, ex A. DC. in DC. Prod. viii. 327 (1844) = **Landolphia**, Beauv. (Apocyn.).
elastica, Sieber, ex A. DC. l. c. (= *Vahea madagascariensis*).—Madag.

FATIOA, DC. Prod. iii. 88 (1828) = **Lagerstroemia**, Linn. (Lythrar.).
nagadensis, DC. l. c. 89 = L. parviflora.

FATOUA, Gaudich. in Freyc. Voy. Bot. 509 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 368.
angera, Gaudich. l. c. 510 = pilosa.
cordata, Gaudich. l. c. = pilosa.
glabelifera, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 207 = pilosa.
japonica, Blume, Mus. Bot. Lugd. Bat. ii. t. 36 = pilosa.
lanceolata, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 493 = pilosa.
pilosa, *Gaudich. in Freyc. Voy. Bot.* 509.—As. or.; Ins. Malay. et Pacif.
scabra, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 207 = pilosa.
subcordata, Gaudich. Bot. Voy. Benite, t. 54. f. 2 = pilosa.

FATRAEA, Thou. Obs. Fl. Afr. 19 (1811) ; Juss. in Dict. Sc. Nat. xvi. 306 (1820) = **Turumiquia**, Lisa. (Combret.).

FATREA, Juss. in Ann. Mus. Par. v. (1804) 388 = prec.

FATSIA, Decne. & Planch. in Rev. Hortic. (1854) 105. *ARALIACEAE*, Benth. & Hook. f. i. 939.
ECHINOPANAX, Decne. & Planch. in Rev. Hortic. (1854) 105.
OPLOPANAX, Miq. in Ann. Mus. Lugd. Bat. i. 16 (1863).
RICHINOPHYLLUM, Pall. ex Ledeb. Fl. Ross. ii. 875 (1844–46).
TETRAPANAX, C. Koch, in C. Koch & Fint. Wochenschr. (1859) 371.
horrida, *Benth. & Hook. f. Gen.* i. 939.—Am. hor. occ.
japonica, *Decne. & Planch. in Rev. Hortic.* (1854) 105. —Japon.
Milnei, De Vriese, ex C. Koch & Fint. Wochenschr. ii. (1859) 371 = Dendropanax japonicum.
papyrifera, *Benth. & Hook. f. Gen.* i. 939.—China.

FAUCIBARBA, Dulac, Fl. Hautes-Pyr. 402 (1867) = **Calamintha**, Moench (Labiat.).
Acinos, Dulac, l. c. 408 = C. Acinos.
alpina, Dulac, l. c. = C. alpina.
Clinopodium, Dulac, l. c. = C. Clinopodium.
grandiflora, Dulac, l. c. 408 = C. grandiflora.
officinalis, Dulac, l. c. = C. officinalis.
parviflora, Dulac, l. c. 408 = C. Nepeta.
umbrosa, Dulac, l. c. 402 = C. officinalis.

FAUJASIA, Cass. in Bull. Soc. Philom. (1819) 80. *COMPOSITAE*, Benth. & Hook. f. ii. 443.
flexuosa, *Benth. & Hook. f. Gen.* ii. 443.—Ins. Mascar. pinifolia, *Cass. in Bull. Soc. Philom.* (1819) 80.—Ins. Mascar.
reticulata, *Benth. & Hook. f. Gen.* ii. 443.—Ins. Mascar.

FAULIA, Rafin. Fl. Tellur. ii. 84 (1836) = **Ligustrum**, Tourn. (Oleac.).
adorata, Rafin. l. c. = L. nepalense.
nepalensis, Rafin. l. c. = L. nepalense.

FAUREA, Harv. in Hook. Lond. Journ. Bot. vi. (1847) 578. t. 15. *PROTEACEAE*, Benth. & Hook. f. iii. 170.
discolor, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 64. —Afr. trop.
forficulifera, *Baker, in Journ. Linn. Soc.* xx. (1883) 245.—Madag.
saligna, *Harv. in Hook. Lond. Journ. Bot.* vi. (1847) 578. t. 15.—Afr. austr.
speciosa, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 63.—Afr. trop.

FAUSTULA, Cass. in Bull. Soc. Philom. (1818) 140 ; et in Dict. Sc. Nat. xvi. 251 (1820) = **Helichrysum**, Vaill. (Compos.).
reticulata, Cass. in Dict. Sc. Nat. xvi. 251 = H. reticulatum.

FAVONIUM, Gaertn. Fruct. ii. 421. t. 174 (1791) = **Didelta**, L'Hérit. (Compos.).
spinosum, Gaertn. l. c. = D. spinosa.

FAWCETTIA, F. Muell. Fragm. x. 93 (1877). *MENISPERMACEAE*.
tinasporoides, *F. Muell. l. c.*—Austral.

FAYA, Neck. Elem. ii. 102 (1790) = **Crenaea**, Aubl. (Lythrar.).

FAYA, Webb & Berth. Phyt. Canar. iii. 272 (1836–50) = **Myrica**, Linn.
fragifera, Webb & Berth. l. c. = M. Faya.

FAYANA, Rafin. Alsog. Am. 19 (1836) = **Myrica**, Linn.
asperiea, Rafin. l. c. = M. Faya.

FEAEA, Spreng. Syst. iii. 363 (1826) = **Salloa**, H. B. & K. (Compos.).
linearis, Spreng. l. c. 361 = S. linearis.
plantaginea, Spreng. l. c. = S. plantaginea.

FEDIA, Adans. Fam. ii. 128 (1763) = **Patrinia**, Juss. (Valer.).
intermedia, Monsen. Hort. Hath. i. 42 = P. intermedia.
saponaria, Vahl, Enum. ii. 28 = P. rupestris.
scabiosa, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1806) 304 = P. intermedia.

FEDIA :—
scabiosaefolia, Trev. in Nov. Act. Nat. Cur. xiii. (1826) 165 = P. scabiosaefolia.
serratadentata, Trev. Ind. Sem. Hort. Wratisl. (1830) = P. scabiosaefolia.
sibirica, Gaertn. Fruct. ii. 37 = P. sibirica.
villosa, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1806) 304 = P. villosa.

FEDIA, Gaertn. Fruct. ii. 86. t. 86. f. 3 (1791), partim ; Moench, Meth. 493 (1794). *VALERIANEAE*, Benth. & Hook. f. ii. 155.
MITREPHORA, Nock. Elem. i. 128 (1790).
Cornucopiae, *Gaertn. l. c.* 87. t. 86. f. 2, partim.—Reg. Mediterr.
gracilifora, Pisch. & Mey. in Ind. Sem. Hort. Petrop. vi. 60 = F. Cornucopiae.
graeca, Dufr. Hist. Valér. 56 = F. Cornucopiae.
incrassata, Moench, Meth. 496 = F. Cornucopiae.
scorpioides, Dufr. Hist. Valér. 55. t. 1 = F. Cornucopiae.

FEDIA, Gaertn. Fruct. ii. 86. t. 86. f. 3, (1791) partim = **Valerianella**, Tourn.
acaulis, Sav. in Mém. Soc. Nat. Mosc. iii. (1812) 251 ; v. (1817) 324 = Hohenackeria bupleurifolia.
Amarella, Lindh. ex Engelm. in A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 217 = V. Amarella.
Auricula, Mert. & Koch, Deutschl. Fl. i. 490 = V. rimosa.
brachystephana, Tenore, Rel. Viagg. 48 = V. brachystephana.
brevifolia, Rafin. Atl. Journ. 177.—Am. bor.
caerulea, Aikin, ex Eaton & Wright, N. Am. Bot. 469 = V. oliitoria.
campestralata, Stend. Nom. ed. I. 333 ; Presl, Fl. Sic. i. 28 = V. eriocarpa.
Caput-bovis, Poucel, Nouv. Mat. Fl. Atl. 72.—Alger.
carinata, Sav. in Mém. Soc. Nat. Mosc. v. (1817) 346 = V. carinata.
carnosa, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1804) 304 = Valeriana carnosa.
cephalophora, Ehrenb. ex Krok, in Act. Akad. Handl. Stockh. v. (1864) n. 1. 79 = V. coronata.
chenopodifolia, Pursh, Fl. Am. Sept. ii. 727 = V. chenopodifolia.
ciliata, Woods, in Trans. Linn. Soc. xvii. (1837) 427, 430 = V. hirsutissima.
ciliolata, Rafin. New Fl. Am. iv. 66.—Am. bor.
coronata, Gaertn. Fruct. ii. 87 = V. discoidea.
coronata, Vahl, Enum. ii. 30 = V. coronata.
costata, Sav. in Mém. Soc. Nat. Mosc. v. (1817) 344 = V. costata.
dasycarpa, Stev. l. c. ii. (1809) 178 ; v. (1817) 348 = V. dentata.
decipiens, Pomel, Nouv. Mat. Fl. Atl. 78.—Alger.
dentata, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1804) 304 = V. dentata.
dentata, Sieber, ex Betcke, Animadv. Bot. Valer. 22 = V. truncata.
discoidea, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1804) 305 = V. discoidea.
discoidea, Reichb. Fl. Crit. i. 84. t. 97 = V. platylobe.
eculeulata, Stokes, Bot. Mat. Med. i. 98 = V. oliitoria.
echinata, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1804) 304 = V. echinata.
eriocarpa, Roem. & Schult. Syst. i. 364 = V. eriocarpa.
eriocarpa, Sav. in Mém. Soc. Nat. Mosc. v. (1817) 349 = V. truncata.
eruca, Roem. & Schult. Syst. i. 366 = Hohenackeria bupleurifolia.
Fagopyrum, Torr. & Gray, Fl. N. Am. ii. 81 = V. chenopodifolia.
gibbosa, Cass. Fl. Sic. Prod. i. 98 = V. costata.
grandiflora, Wall. ex DC. Prod. iv. 634 = Nardostachys jatamansi.
hamata, Roem. & Schult. Syst. i. 363 = V. costata.
hemicles, Rafin. New Fl. Am. iv. 66.—Am. bor.
heterocarpa, Pomel, Nouv. Mat. Fl. Atl. 71.—Alger.
Jatamansi, Wall. ex DC. Prod. iv. 630 = Nardostachys jatamansi.
laciocarpa, Sav. in Mém. Soc. Nat. Mosc. v. (1817) 350 = V. laciocarpa.
laciocephala, Reichb. ex Woods, in Trans. Linn. Soc. xvii. (1837) 431 = V. coronata.
laciflora, Roem. & Schult. Syst. i. 362 = V. chenopodifolia.
ligulata, Rafin. New Fl. Am. iv. 66 = V. longiflora ?
Locusta, Rafin. in Fl. Cestr. i. 45 = V. oliitoria.

FEDIA:—
longifora, Torr. & Gray, Fl. N. Am. ii. 51 = Valerianella longiflora.
maxima, Roem. & Schult. Syst. i. 366 = Veronica maxima (= Veronica melissaefolia.)
microcarpa, Reichb. Ic. Pl. Crit. ii. 6. t. 114 = V. dentata.
minuta, Spreng. Syst. i. 144 = V. lasiocarpa.
mixta, Vahl, Enum. ii. 21 = V. dentata.
Mortonii, Spreng. Pugill. i. 4 = V. dentata.
muricata, Stev. ex Roem. & Schult. Syst. i. 366 = V. truncata.
nervata, Hochst. & Steud. ex Bolss. Fl. Orient. iii. 112 = V. obtusiloba.
Nuttallii, Torr. & Gray, Fl. N. Am. ii. 51 = V. Nuttallii.
oliteria, Gaertn. Fruct. ii. 86 = V. rimosa.
oliteria, Stev. in Mém. Soc. Nat. Mosc. ii. (1809) 178 = V. turgida.
oliteria, Vahl, Enum. ii. 19 = V. oliteria.
orientalis, Schlecht. in Linnaea, xvii. (1843) 196 = V. orientalis.
paniculata, Colla, Hort. Ripul. App. iii. 36 = V. oliteria.
patellaria, Salliv. ex A. Gray, Man. Bot. U. St. 183 = V. Woodsiana.
platyloba, Roem. & Schult. Syst. i. 365 = V. platyloba.
polystachia, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1804) 304 = Valerianopsis polystachya.
pterocles, Stev. ex Steud. Nom. ed. II. i. 626 = V. pterocloba.
puberula, Bertol. in Guss. Pl. Rar. Sic. (1826) 15 = V. dentata.
pumila, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1804) 305 = V. membranacea.
quadrangularis, Willd. ex Steud. Nom. ed. II. i. 626 = V. carinata.
radiata, Michx. Fl. Bor. Am. i. 18; Vahl, Enum. ii. 21 = V. radiata.
radiata, Torr. & Gray, Fl. N. Am. i. 35 = V. chenopodifolia.
rotata, Reichb. Ic. Fl. Germ. i. 93 = V. platyloba.
rugulosa, Hornem. ex Steud. Nom. ed. II. i. 626 = V. dentata.
rugulosa, Spreng. Pugill. ii. 2 = V. eriocarpa.
samolifolia, Bert. in 'Merc. Chil. (1829);' ex Bull. Férussac, xx. (1830) 111 = V. samolifolia.
seridens, Hort. Pest. ex Steud. Nom. ed. II. i. 626 = V. oxyrhyncha.
sicula, Guss. Fl. Sic. Prod. i. 25 = V. discoidea.
sphaerocarpa, Guss. i. c. 28 = V. membranacea.
stenocarpa, Engelm. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 216 = V. stenocarpa.
striata, Stev. in Mém. Soc. Nat. Mosc. ii. (1809) 177 = V. oliteria.
sulcata, Pomel, Nouv. Mat. Fl. Atl. 72.—Alger.
trichocarpa, D. Dietr. Syn. Pl. i. 169 = V. turgida.
tridentata, Reichb. Ic. Pl. Crit. f. 131 = V. rimosa.
tridentata, Stev. in Mém. Soc. Nat. Mosc. ii. (1809) 178; et v. (1817) 846 = V. membranacea.
trigonocarpa, Benth. ex Steud. Nom. ed. II. i. 626 = V. turgida.
triquetra, Hochst. & Steud. ex Shuttlew. in Flora, xx. (1837) 211 = V. chenopodifolia.
truncata, Reichb. Ic. Fl. Germ. ii. 7 = V. truncata.
turgida, Stev. in Mém. Soc. Nat. Mosc. v. (1817) 345 = V. turgida.
umbilicata, Salliv. in Am. Journ. Sc. xlii. (1842) 50 = V. Woodsiana.
uncinata, Willd. Enum. Hort. Berol. 51 (1809) = V. uncinata.
vesicaria, Mirb. in Hist. Nat. Pl. ed. II. xiii. (1804) 306 = V. vesicaria.
Woodsiana, Torr. & Gray, Fl. N. Am. ii. 52 = V. Woodsiana.

FEDIA, H. B. & K. Nov. Gen. et Sp. iii. 384 (1818) = Astrephia, Desv. (Valer.).
chenopodioides, H. B. & K. l. c. = A. chenopodifolia.
lasia, Hook. & Arn. Bot. Beech. Voy. 28 = A. laxa.

FEDTSCHENKOA, Regel, Fl. Nov. Fedsch. 8 (1868). *CRUCIFERAE.*
tentamina, *Regel & Schmalh. ex Regel, l. c.*—Turkest.

FEDIA, Hornem. Hort. Hafn. i. 47 (1813) = FEDIA, Gaertn. = Valerianella, Tourn. (Valerian.).

FEIDANTHUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 148 = **Astragalus**, Tourn. (Legumin.).
triastellus, Stev. l. c. = A. sviaticellus.

FELIPOA, Berg, in Linnaea, xxix. (1858) 258. *MYRTACEAE*, Benth. & Hook. f. i. 712.
ORTHOSTEMON, Berg, in Linnaea, xxvii. (1854) 440.
obovata, *Berg, in Mart. Fl. Bras.* xiv. 1. 616.—Bras.
Sellowiana, *Berg, l. c.* 615.—Bras.

FELICIA, Cass. in Bull. Soc. Philom. 165 (1818); et in Dict. Sc. Nat. xvi. 314 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 274.
AGATACHA, Cass. in Bull. Soc. Philom. (1815) 175.
AGATHAEA, Cass. in Dict. Sc. Nat. i. Suppl. 77 t. 89 (1816).
ASTEROSPERMA, Less. Syn. Comp. 389 (1832).
COLLOMIA, Sieber, ex Steud. Nom. ed. II. i. 398 (1840).
DETRIDIUM, Nees, Gen. et Sp. Ast. 255 (1833).
DETRIS, Adans. Fam. ii. 131 (1763), nomen prius.
ELPHEGRA, Less. Syn. Comp. 182 (1832).
MUNYCHIA, Cass. in Dict. Sc. Nat. xxxvii. 462 (1825).
POLYARRHENA, Cass. l. c. lvi. 172 (1826).
ab asinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 388.—Afr. trop.
affinis, *Nees, Gen. et Sp. Ast.* 309.—Afr. austr.
angustifolia, *Nees, l. c.* 213.—Afr. austr.
brachyglossa, Cass. in Dict. Sc. Nat. xxv. 97 = Aster Cymbalariae.
cafrorum, *Nees, Gen. et Sp. Ast.* 214.—Afr. austr.
cana, DC. Prod. v. 230 = angustifolia.
ciliaris, DC. l. c. 221 = Gymnostephium ciliare.
cotuloides, DC. l. c. 219 = fragilis.
Dregei, DC. l. c. 221 = Aster Dregei.
dubia, Cass. in Dict. Sc. Nat. xvi. 315 = Aster Candollii.
echinata, *Nees, Gen. et Sp. Ast.* 216.—Afr. austr.
erigeroides, DC. Prod. v. 219 = Aster erigeroides.
fascicularis, DC. l. c. 220 = muricata.
floridula, DC. l. c. 221 = Aster floridus.
fragilis, *Cass. in Dict. Sc. Nat.* xvi. 315.—Afr. austr.
fruticosa, Nichols. Dict. Gard. ii. 4 = Aster fruticosus.
hirsuta, DC. Prod. v. 220 = Aster confusus (in addend.).
hyssopifolia, Nees, Gen. et Sp. Ast. 211 = angustifolia.
imbricata, DC. Prod. v. 221 = Aster imbricatus.
Kraussii, *Sch. Bip. ex Walp. Rep.* ii. 956.—Afr. austr.
lasiocarpa, DC. Prod. v. 221 = Aster lasiocarpus.
macrorrhiza, DC. l. c. 219 = Aster macrorrhizus.
microsperma, DC. l. c. = Aster microspermus.
muricata, *Nees, Gen. et Sp. Ast.* 210.—Afr. austr.
natalensis, Sch. Bip. ex Walp. Rep. ii. 956 = Aster erigeroides.
Parodia, DC. Prod. v. 222 = echinata.
Rouvelii, *Nees, Ind. Sem. Hort. Vratisl.* (1850).—*Cf. Linnaea*, xxiv. (1851) 236.—Hab.?
reflexa, *DC. Prod.* v. 222.—Afr. austr.
retorta, DC. l. c. 221 = cafrorum.
Richardi, *Vatke, in Linnaea*, xxxix. (1875) 481.—Abyss.
rigidula, DC. Prod. v. 220 = angustifolia.
Schimperi, *Steud. & Hochst. ex Jaub. & Spach, Illustr.* iv. 86. t. 354.—Arabia.
tenella, Nees, Gen. et Sp. Ast. 208 = fragilis.
trinervia, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 60 = Aster erigeroides.
Zeyheri, *Nees, Gen. et Sp. Ast.* 215.—Afr. austr.

FELICIANA, Benth. & Hook. f. Gen. i. 717 (1868) sphalm. = seq.

FELICIANEA, Cambess. in St. Hil. Fl. Bras. Mer. ii. 275. t. 157 (1829) = **Myrrhinium**, Schott (Myrtac.).
rubriflora, Cambess. l. c. 276 = M. atropurpureum.

FENDLERIA, Steud. Syn. Pl. Gram. i. 419 (1855) = **Oryzopsis**, Michx. (Gramin.).
rhynchelytroides, Steud. l. c. 430.—N. Mexic.

FENELONIA, Rafin. Atl. Journ. 145 (1832) = Ornithogalum, Tourn. (Lillac.).
bracteata, Rafin. l. c. = O. longibracteatum.

FENICULUM, Gilib. Exercit. i. 220 (1792) = Foeniculum, Tourn. (Umbell.).

FENTZLIA, Reichb. Handb. 194 (1837) = FENZLIA, Benth. = **Gilia**, Ruiz & Pav. (Polemon.).

FENUGRAECUM, Adans. Fam. ii. 322 (1763) = **Trigonella**, Linn. (Legum.).

FENZLIA, Benth. in Bot. Reg. sub t. 1622 (1833) = **Gilia**, Ruiz & Pav. (Polemon.).
concinna, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 157 = G. dianthoides.
dianthiflora, Benth. in Bot. Reg. sub t. 1622 = G. dianthoides.
speciosa, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 157 = G. dianthoides.

FENZLIA, Endl. Atakt. 19. tt. 17, 18 (1833). *MYRTACEAE*, Benth. & Hook. f. i. 715.
obtusa, *Endl. l. c.* 19. t. 17.—Austral.
retusa, *Endl. l. c.* 20. t. 18.—Austral.

FERBERIA, Scop. Introd. 281 (1777) = **Althaea**, Tourn. (Malvac.).

FERDINANDA, Benth. & Hook. f. Gen. ii. 1047 (1876) = FERDINANDINA, Welw. = **Heterophragma**, DC. (Bignon.).

FERDINANDA, Lag. Gen. et Sp. Nov. 31 (1816) = **Saltuaria**, Pers. (Compos.).
angusta, Lag. l. c. 31. t. 2 = Z. angusta.
eminens, Lag. l. c. = Podachaenium paniculatum.
integrifolia, G. Don, in Sweet, Hort. Brit. ed. III. 856 = Z. angusta!
Lindenii, Sch. Bip. ex C. Koch, in Alig. Gartenz. (1858) 179.—Am. trop.
lutescens, DC. Prod. v. 558 = Z. angusta.
montagnaefolia, Sch. Bip. ex C. Koch, in Alig. Gartenz. (1858) 179 = Z. montagnaefolia.
oppositifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Calea scabriflora.
salutina, Cass. in Dict. Sc. Nat. xvi. 490 = Z. angusta.

FERDINANDEA, Pohl, in Flora, x. (1827) 158 = **Ferdinandusa**, Pohl (Rubiac.).
elliptica, Pohl, l. c. = Ferdinandusa elliptica.
ovalis, Pohl, l. c. = Ferdinandusa ovalis.
speciosa, Pohl, l. c. 154 = Ferdinandusa speciosa.

FERDINANDIA, Welw. ex Seem. Journ. Bot. iii. (1865) 330. t. 38 = seq.
suporba, Welw. ex Seem. l. c. (= *Ferdinandea superba*).—Afr. trop.

FERDINANDOA, Seem. Journ. Bot. viii. (1870) 280 = **Heterophragma**, DC. (Bignon.).
magnifica, Seem. l. c.—Afr. trop.
superba, Seem. l. c.—Afr. trop.

FERDINANDUSA, Pohl, Pl. Bras. Ic. ii. 8. t. 106 ad 108 (1831). *RUBIACEAE*, Benth. & Hook. f. ii. 40.
ASPIDANTHERA, Benth. in Hook. Journ. Bot. iii. (1841) 217.
FERDINANDEA, Pohl, in Flora, x. (1827) 158.
GOMPHOSIA, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 14 (Mon. Cinch. 98. t. 26 B).
andina, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 78.—Reg. Andin.
angustata, Wright, ex Griseb. Cat. Pl. Cub. 127 (Ferdinandea) (Rondeletiae sp.).—Cuba.
brachycarpa, Griseb. in Mem. Am. Acad. N. S. viii. (1862) 505 (Ferdinandea) (Rondeletiae sp.).—Cuba.
elliptica, Pohl, Pl. Bras. Ic. ii. 9. t. 106 = ovalis.
ovalis, *Pohl, l. c.* 10. t. 107.—Bras.
pubescens, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 78.—Bras.
rudgeoides, *Wedd. l. c.*—Guiana.
speciosa, *Pohl, Pl. Bras. ii.* 12. t. 108.—Bras.
stellata, Griseb. in Mem. Am. Acad. N. S. viii. (1862) 505 (Ferdinandea) (Rondeletiae sp.).—Cuba.

FERECUPPA, Dulac, Fl. Hautes-Pyr. 373 (1867) = **Tossia**, Linn. (Scrophularia.).
drupacea, Dulac, l. c. = *T. alpina*.

FEREIRA, Reichb. Consp. 182 (1828) = seq.

FEREIRIA, Vell. ex Vand. Fl. Lusit. 21. t. 1. f. 8 (1788); et in Roem. Script. 98. t. 6 = **Hillia**, Jacq. (Rubiac.).
Velloseana, Schult. f. Syst. vii. 68 = H. longiflora.

FERETIA, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 92. t. 1. f. 4. *RUBIACEAE*, Benth. & Hook. f. ii. 94.
apocabathea, *Delile, l. c.*—Abyss.
canthioides, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 116.—Afr. trop.

FERGUSONIA, Hook. f. in Hook. Ic. Pl. t. 1124 (1879). *RUBIACEAE*, Benth. & Hook. f. ii. 183.
tetracocca, Bedd. Hist. Pl. vii. 418 = seylanica.
Thwaitesii, Hook. f. in Hook. Ic. Pl. xii. 28 = seylanica.
seylanica, *Hook. f. l. c.* t. 1124.—Ind. or.

FERNANDEZIA, Lindl. Gen. et Sp. Orch. 307 (1838) = **Lockhartia**, Hook. (Orchid.).
acuta, Lindl. Bot. Reg. t. 1806 = L. acuta.
elegans, Lodd. Bot. Cab. t. 1314 = L. elegans.
longifolia, Lindl. Orch. Linden. 18.—Venezuela.
lunifera, Lindl. Bot. Reg. (1839) Misc. 91 = L. lunifera.
obtusa, Lindl. ex Linden, Illustr. Hortic. xxix. (1882) 51 = L. obtusa.
parthenocomos, Reichb. f. in Bot. Zeit. x. (1852) 839 = L. parthenocomos.
robusta, Batem. in Bot. Mag. t. 5592 = L. verrucosa.
robusta, Klotzsch, ex Reichb. f. in Walp. Ann. vi. 821 = L. lunifera.

FERNANDEZIA, Ruiz & Pav. Prod. Fl. Per. et Chil. 123. t. 27 (1794) = **Centropetalum**, Lindl. et Dickson, Lindl. (Orchid.).
conferta, Ruiz & Pav. Syst. Veg. 241.—Peruv.
denticulata, Ruiz & Pav. l. c. 240.—Peruv.
graminifolia, Ruiz & Pav. l. c. 241.—Peruv.
haematodes, Ruiz & Pav. l. c. 240.—Peruv.
laxa, Ruiz & Pav. l. c. = Dichaea laxa.
punctata, Ruiz & Pav. l. c. 239.—Peruv.
subdifera, Ruiz & Pav. l. c. 240.—Peruv.

FERNANDOA, Welw. ex Seem. Journ. Bot. iv. (1866) 193 = **FERDINANDOA**, Seem. = **Heterophragma**, DC. (Bignon.).

FERNELIA, Comm. ex Lam. Encyc. ii. 459 (1786); Juss. Gen. 199 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 93.
axillaris, Stend. Nom. ed. I. 334 = Manettia Lygistum.
bifora, Roem. & Schult. Syst. iii. 187 = Manettia Lygistum.
buxifolia, Gaertn. f. Fruct. iii. 68 = obovata.
buxifolia, *Lam. Encyc.* ii. 459.—Ins. Maurit.
mauritiana, J. F. Gmel. Syst. 344 = buxifolia.
obovata, Gaertn. f. Fruct. iii. 69. t. 191 = buxifolia.
obovata, *Lam. Illustr.* i. 367.—Ins. Maurit.
ovata, Ayres, ex Baker, Fl. Maurit. 163 = buxifolia.
pedunculata, Gaertn. f. Fruct. 3. t. 191. f. 3 = buxifolia.
uniflora, Roem. & Schult. Syst. iii. 187 = obovata.

FEROLIA, Anbl. Fl. Gal. Suppl. 7 (1775) = Parinarium, Juss. (Rosac.).
guianensis, Anbl. l. c.—Guian.
variegata, Lam. Encyc. ii. 459 = *guianensis*.

FERONIA, Correa, in Trans. Linn. Soc. v. (1800) 224. *RUTACEAE*, Benth. & Hook. f. i. 305.
JERONIA, Pritz. Ic. Ind. i. 579 (1855), sphalm.
Balsaphra, Stend. Nom. ed. II. i. 237 = Sterculia Balsaphra.
elephantum, *Correa, in Trans. Linn. Soc.* v. (1800) 225.—Ind. or.
lucida, *Teijsm. & Binn. Cat. Hort. Bog.* 230; *et ex Schegf. in Tijdschr. Naderl. Ind.* xxxi. (1870) 19. —Java.
pellucida, Roth, Nov. Sp. Pl. 334 = Aegle Marmelos.
ternata, Blanco, Fl. Filip. ed. II. 253 = Aegle decandra.

FERRANDIA, Gandich. in Freyc. Voy. Bot. 477 (1826) Cucurbitac. DC. (Momordicin.).
oleifera, Gandich. l. c. = C. Ferrandiana.

FERRARIA, Linn. Sp. Pl. ed. II. 1258 (1763). *IRIDEAE*, Benth. & Hook. f. iii. 693.
angustifolia, Sweet, Hort. Brit. ed. I. 396 = F. Ferrariola.
anthracina, Ker-Gawl. in Bot. Mag. t. 751 = F. Ferrariola.
atrata, *Ledd. Bot. Cab.* t. 1856.—Afr. austr.
blanda, Salisb. Prod. 42 = Moraea iridioides.
cathartica, Mart. Reise Bras. i. 548 (= *Lansbergia cathartica*).—Bras.
crocea, Salisb. Prod. 41 = Belamcanda punctata.
divaricata, *Sweet, Brit. Flow. Gard.* t. 192.—Afr. austr.
elegans, Salisb. Prod. 42 = Marica Northiana.
elongata, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.-June 1836) 179.—Reg. Argent.
Ferrariola, *Willd. Sp. Pl.* iii. 561.—Afr. austr.
ixioides, Willd. l. c. 562 = Libertia ixioides.
longiscapa, Salisb. Prod. 42 = Moraea plumaria.
lurida, Salisb. l. c. 41 = Trimezia lurida.
major, *Ecbl. Verz.* 18.—Afr. austr.
minor, Pers. Syn. i. 50 = F. Ferrariola.
obtusifolia, *Sweet, Brit. Flow. Gard.* t. 148.—Afr. austr.
ocellaris, Salisb. Prod. 41 = Moraea tricuspis.
ochroleuca, Salisb. l. c. 42 = Sisyrinchium striatum.
parviflora, Salisb. l. c. 42 = Elettheriza pilosta.
Pavonia, Linn. f. Suppl. 407 = Tigridia Pavonia.
pulchella, Salisb. Prod. 42 = Sisyrinchium angustifolium.
punctata, Pers. Syn. i. 50 = undulata.
purpurea, Mart. Reise Bras. i. 547 (= *Lansbergia purpurea*).—Bras.
pusilla, Link & Otto, Ic. Pl. Select. 125. t. 59 = Cypella pusilla.
Tigridia, Sims, Bot. Mag. t. 532 = Tigridia Pavonia.
tricuspis, Willd. Enum. Hort. Berol. 69 (= *Vieusseuxia variegata, glaucopis*).—Afr. austr.
tristis, Salisb. Prod. 42 = Moraea tristis.
undulata, *Sweet, Brit. Flow. Gard.* t. 161.—Afr. austr.
undulata, *Linn. Sp. Pl.* ed. II. 1258.—Afr. austr.
violacea, Salisb. Prod. 42 = Sisyrinchium Bermudianum.
viridiflora, Andr. Bot. Rep. t. 385 = F. Ferrariola.

FERREIRIA, Allem. in Trab. Soc. Viell. 26 (1845!). *LEGUMINOSAE*, Benth. & Hook. f. i. 556.
spectabilis, *Allem. l. c.*—Bras.

FERREOLA, Kom. ex Roxb. Pl. Corom. i. 35. t. 45 (1796) = **Ehretia**, Forst. (Ehretiac.).
buxifolia, Roxb. l. c. = M. buxifolia.
Ehretia, Roxb. Bot. Mat. Med. iv. 556 = M. buxifolia.
ellipticifolia, Stokes, l. c. = M. elliptica.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 448 = M. buxifolia.

FERRETIA, Pritz. Ic. Ind. i. 455 (1855), sphalm. = **Furcaea**, Delile (Rubiac.).

FERRIOLA, Roxb. Fl. Ind. iii. 790 (1832) = FERREOLA, Koen.

FERRUM-EQUINUM, Tourn. ex Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 370 = **Hippocrepis**, Linn. (Legumin.).
multisiliquum, Moench, Meth. 119 = H. multisiliquosa.
multisiliquosum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 371 = H. multisiliquosa.
uniflorum, Moench, Meth. 118 = H. unisiliquosa.
unisiliquosum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 371 = H. unisiliquosa.

FERULA, Tourn. ex Linn. Gen. ed. I. 75 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 917.
ERIOSTYRAPHIS, DC. Coll. Mém. v. 50 (1830); et Prod. iv. 175 (1830).
EURYANGIUM, Kunze, in Nouv. Mém. Soc. Nat. Mosc. xiii. (1871) 256.
FERULAGO, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 97.
HALMATOCAULIS, Tausch, in Flora, xvii. 847 (1834).
NARTHEX, Falc. in Trans. Linn. Soc. xx. (1847) 235.
SCORODOSMA, Bunge, in Mém. Sav. Etr. Acad. Pétersb. vii. (1854) 303.
SCRANTHOX, Ledeb. Ic. Pl. Ross. i. 30 (1829); Fl. Alt. i. 344 (1829).

FERULA:—
SUMBULUS, H. Reisech. in Jahrb. Pract. Pharm. xiii. (1846) 99.
ULOPTERA, Fenzl, in Flora, xxvi. (1843) 460.
XANTHOGALUM, Avé-Lall. in Fisch. & Mey. Ind. Sem. Hort. Petrop. viii. (1841) 74.
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 236. —Abyssin.
aciphylla, Bieb. ex Ledeb. Fl. Ross. ii. 308 = caspica.
adiana, *Bess. Cat. Hort. Crem.* (1816) 37.—Tauria.
alliacea, *Boiss. Fl. Orient.* ii. 995.—Persia.
anatolica, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 317.—As. Min.
angulata, *Schlecht. in Linnaea*, xvii. (1843) 125.—Kurdistan.
asmarica, *Sten. in Pall. N. Nord. Beitr.* vii. 295.—Russia.
ormosa, Pall. ex Ledeb. Fl. Ross. ii. 308 = caspica.
armena, *DC. Prod.* iv. 174.—Armen.
armena, *C. Koch, in Linnaea*, xvi. (1842) 355.—Armen.
Asa-foetida, Spreng. in Nene Schr. Halle, ii. (1813) 14 = Asse-foetida.
Asse-foetida, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 100 = silicosa.
Asse-foetida, *Linn. Sp. Pl.* 348.—Persia; Afghan.
Asse-foetida, Martyn, in Mill. Gard. Dict. ed. IX. n. 9 = persica.
aurea, Link, in Buch, Beschr. Canar. Ins. 152 = Feuccedanum aureum.
austriaca, Spreng. in Nene Schr. Halle, ii. (1813) 14 = Peucedanum austriacum.
Barrelieri, Ten. Fl. Neap. t. 133 = silicata.
Bessereana, Spreng. ex DC. Prod. iv. 177 = Peucedanum ruthenicum.
Blanchei, *Boiss. Fl. Orient.* ii. 987.—Syria.
brachylobe, *Boiss. & Reut. in Boiss. Voy. Bot. Espagne*, 738.—Hispan.
brevifolia, *Link, ex Schult. Syst.* vi. 592.—Lusitan.
cachrydes, Fisch. Hort. Gorenk. ed. II. (1812) 46 = caspica.
californica, *A. Gray, in Proc. Am. Acad.* vii. (1868) 348.—Am. bor. occ.
campestris, Bess. Enum. Pl. Volh. 44 = F. ferulago.
canescens, Lion. Sp. Pl. 347 = Ligusticum actaeifolium.
Candelabrum, Haldr. & Sart. in Boiss. Diagn. Ser. II. vi. 88 = glauca.
canescens, *Ledeb. Fl. Ross.* ii. 302.—Soongaria.
capillaris, *Link, in Spreng. Umbell. Sp.* 85.—Lusitan.
capillifolia, Stend. Nom. ed. II. i. 627 = prnec.
capillifolia, *Steudg. in Bull. Soc. Nat. Mosc.* (1854) t. 183.—Soongaria.
cervicifolia, Hook. & Arn. Bot. Beech. Voy. 348 = Peucedanum cervicifolium.
caspica, *Bieb. Fl. Taur. Cauc.* i. 220.—Reg. Casp. Ural.
caspica, Spreng. ex Ledeb. Fl. Ross. ii. 302 = nuda.
cassiophylla, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 396.—Turkestan.
communis, Gouan, Hort. Monsp. 140 = glauca.
communis, Hoffm. in Flora, xxxvi. (1853) 698 = Hoffmii.
communis, *Linn. Sp. Pl.* 246.—Reg. Mediterr.
decurrens, *Ledeb. ex Sweet, Hort. Brit.* ed. II. 250, nomen.—Sibir.
dissecta, A. Gray, in Proc. Am. Acad. vii. (1868) 348 = dissoluta.
dissecta, *Ledeb. Fl. Ross.* ii. 301.—Sibir. alpin.
dissoluta, *T. Wats. Bot. Calif.* i. 271.—Calif.
divaricatissima, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 388.—Turkestan.
Drudeanoides, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 47 = Polytaenia Nuttallii.
elegans, Spreng. in Nene Schr. Halle, ii. (1813) 14 = Peucedanum austriacum.
erubescens, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 316 = galbaniflua, rubricaulis.
Ferulago, *Linn. Sp. Pl.* 247.—Europ. austr.; Reg. Caucas.
fontanesiana, Hook. Fl. Bor. Am. i. 268 = Peucedanum foeniculaceum, millefolium, macrocarpum.
foeniculacea, Nutt. Gen. Am. i. 125 = Peucedanum foeniculaceum.
foetida, *Regel in Act. Hort. Petrop.* v. (1877) 392 (sphalm. 398).—Turkestan.

FERULA :—
fœtida, St. Leg. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = F. Asa-fœtida.
fœtidissima, Regel & Schmalh. i. c. 898 = Jaschkeana.
galbaniflua, Mill. Gard. Dict. ed. VIII. n 2.=F. Ferulago.
galbanifera, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 99.—Persia.
gummiflua, Bory & Chamb. Nouv. Fl. Pélop. 20 = nodosa.
gummifera, Boiss. Diagn. Ser. II. ii. 92 = galbaniflua.
Hermonis, *Boiss. Fl. Orient.* ii. 985.—Syria.
Heuffelii, *Griseb. ex Heuff. in Flora*, xxxvi. (1853) 693.—Europ. austr.
hirtella, *Boiss. Fl. Orient.* ii. 998.—Persia.
hispida, *Fryvald. in Flora*, xviii. (1835) 338.—Turcia.
Jaschkeana, *Vathe, Sem. Hort. Berol.* (1876) *App.* 2. —Himal. bor. occ. ; Turkestan.
karatavica, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 594.—Turkestan.
Karelini, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 206.—Persia ; Turkestan.
kokanica, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 593.—Turkestan.
laevigata, *Spreng. Umbell. Sp.* 88.—Afr. austr.
lancerotensis, *Parl. ex Hartwag, Geogn. Verh. Ins. Lanzar. Fuerst.* 145.—Ins. Canar.
Lehmanni, *Boiss. Fl. Orient.* ii. 992.—Reg. Casp. ; Turkestan.
linearis, *Cerv. ex Sweet, Hort. Brit.* ed. II. 591, nomen.—Mexic.
Linkii, *Webb & Berth. Phyt. Canar.* ii. 160.—Ins. Canar.
Lobeliana, Vis. in Nuov. Giorn. Bot. Ital. ii. (1870) 215 = Communis.
longifolia, *Fisch. Cat. Hort. Gorensk.* (1812) 46.—Rossia.
longifolia, Spreng. ex Ledeb. Fl. Ross. ii. 300 = tatarica.
longifolia, Willd. ex Ledeb. l. c. 248 = Trinia Kitaibelii.
lophoptera, *Benth. & Hook. f. Gen.* i. 817.—Persia.
lycia, *Boiss. Fl. Orient.* ii. 984.—Lycia.
macrocarpa, Hook. & Arn. Bot. Beech. Voy. 348 = Pseudanum macrocarpum.
macrocolea, *Boiss. Fl. Orient.* ii. 984.—Persia.
marathrophylla, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 347.—China.
meifolia, *Boiss. Fl. Orient.* ii. 986.—Syria.
meifolia, *Echl. & Zeyh. Enum.* 348.—Afr. austr.
meoides, *Linn. Sp. Pl.* 247.—Reg. Byzant.
meoides, Schousb. ex Bail, in journ. Linn. Soc. xvi. (1878) 479 = Elaeoselinum meoides.
Meyeri, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 207.—As. centr.
microcolea, *Boiss. Fl. Orient.* ii. 990.—Persia.
microlobe, *Boiss. l. c.* 989.—Persia.
microphylla, Bieb. ex Schur, Enum. Pl. Transs. 268 = sylvatica.
montana, Spreng. in Nene Schr. Halle, ii. (1813) 14 = Pseudanum austriacum.
monticola, Neilr. Nachtr. 199 = sylvatica.
monticola, *Nym. Syll. Suppl.* 26.—Graecia.
multifida, *A. Gray, in Proc. Am. Acad.* vii. (1868) 348.—Am. bor. occ.
myrrhaphila, Pleh. ex Rem. Enum. Pl. Volh. 44 = sylvatica.
Narthex, *Boiss. Fl. Orient.* ii. 994.—Baltisthan.
nebrodiana, Tenore, Fl. Nap. iii. 240. t. 121 = glauca.
Newberryi, *S. Watt. in Proc. Am. Acad.* xi. (1876) 148.—Am. bor. occ.
nodiflora, Jacq. Fl. Austr. App. t. 5 = F. Ferulago.
nodiflora, Lam. Sp. Pl. 247 = communis.
nodiflora, Pall. Reise, ii. 736 = Meyeri.
nodiflora, Rochel, Pl. Ban. Banat. 69 = sylvatica.
nodosa, *Kunth & Hook. f. Gen.* i. 918.—Europ.
nodosa, Pall. ex Ledeb. Fl. Ross. ii. 307 = longifolia.
nuda, *Spreng. Umbell. Sp.* 61. t. 7. f. 15.—Turkestan.
nudicaulis, Pall. ex Ledeb. Fl. Ross. ii. 302 = caspica.
nudicaulis, Nutt. Gen. Am. i. 126 = Pseudanum nudicaule.

FERULA :—
nudicaulis, *Spreng. Neue Entdeck.* ii. 149.—Ins. Sicil.
Nuttallii, DC. Prod. iv. 174 = Pseudanum nudicaule.
Nuttallii, Walp. Rep. ii. 411 = Pseudanum triternatum.
obtusifolia, Spreng. Syst. i. 914 = Malabaila obtusifolia.
Olgae, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 590, nomen ; *Pl. Nov. Fedsch.* 86.—Turkestan.
oopoda, *Boiss. Fl. Orient.* ii. 984.—Persia ; Beluchist.
Opopanax, Spreng. Umbell. Sp. 77 = Opopanax hispidum.
orientalis, Elchw. Skizze, 156 = caspica.
orientalis, *Linn. Sp. Pl.* 247.—As. Min.
orientalis, Schlecht. ex Ledeb. Fl. Ross. ii. 300 = tatarica.
ovina, *Boiss. Fl. Orient.* ii. 986.—Persia ; Beluchist.
pachyloba, *Fenzl, Pugill. Pl. Nov. Syr.* i. 18.—Cilicia.
Palmella, Hook. Fl. Bor. Am. i. 268 = Cymopterus glomeratus.
paniculata, Ledeb. Fl. Ross. ii. 305 = resinosa.
parviflora, Hook. & Arn. Bot. Beech. Voy. 348 = Pseudanum parvifolium.
pauciflora, *C. Koch, in Linnaea*, xvi. (1842) 358.—Armenia.
paucijuga, DC. Prod. iv. 171 = Johrenia Candollei.
pauniversis, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 591.—Turkestan.
persica, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 205 = Lehmanni !
persica, Sims, Bot. Mag. t. 2096 = galbaniflua, rubricaulis.
persica, *Willd. Sp. Pl.* i. 1413.—Reg. Caucas. ; Persia.
petiolaris, DC. Prod. iv. 178 = Pseudanum petiolare.
peucedanifolia, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1839) ii. 155 ; xv. (1842) ; 365 = Karelini.
peucedanifolia, Willd. ex Schult. Syst. vi. 592 = Meyeri.
pubens, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 98 = persica.
puberula, *Trautv. in Bull. Soc. Nat. Mosc.* xxxix. (1866) i. 823.—As. temp.
pubescens, Pall. ex Schult. Syst. vi. 598 = Cachrys odontalgica.
pumila, Pall. ex Schult. l. c. = nuda.
rablensis, Walf. in Jacq. Coll. iv. 313 = Pseudanum austriacum.
racemifera, Szov. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. 26 = Dorema glabrum.
resinosa, *Sievers, in Pall. N. Nord. Beitr.* vii. 260.—Sibir.
rigida, *Tenore, Fl. Neap. App.* iv. 15.—Italia.
rigidula, *Fisch. ex DC. Prod.* iv. 172.—Reg. Caucas. ; Persia.
rubricaulis, *Boiss. Diagn.* Ser. II. ii. 92.—Persia.
ruprechtii, Boiss. & Bal. i. c. Ser. II. vi. 86, in syn. = orientalis.
ruthenica, Spreng. Pugill. i. 38 = Pseudanum ruthenicum.
Sadleriana, *Ledeb. Fl. Ross.* ii. 30, in nota.—Hungar.
salsa, Ledeb. Fl. Alt. i. 344, in nota = Pseudanum redivivum.
saxatilis, Balb. Diagn. Ser. I. x. 39 = tingitana.
scabra, Fenzl, in Flora, xxvi. (1843) 461 = Hippomarathrum scabrum.
Schair, *Sieber, in Mém. Acad. Pétersb.* Sér. VII. iii. (1861) 8, 37. t. 7.—Turkestan.
Schlechtendalii, *Boiss. Fl. Orient.* ii. 985.—Kurdistan.
Schtschurowskiana, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 84.—Turkestan.
Scordoxema, Bentley & Trim. Med. Pl. t. 151 = foetida.
sesseloides, C. A. Mey. Verz. Pfl. Cauc. 126 = Peucedanum pauciflorum.
sibirica, Sadl. Fl. Com. Pest. i. 228 = Sadleriana.
sibirica, Schaug. in Pall. N. Nord. Beitr. vi. 111 = sonchifolia.
sibirica, Schlecht. ex Ledeb. Fl. Ross. ii. 307 = longifolia.
silvicola, Willd. Sp. Pl. i. 1411 = Meyeri.
slaskia, *Boiss. Diagn.* Ser. I. x. 40.—Arab.
sonchifolia, *Pall. ex Schult. Syst.* vi. 598.—Sibir. altaic.
stenocarpa, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 994.—Persia.
stricta, *Spreng. ex Schult. Syst.* vi. 592.—Afr. austr.
sulcata, *Desf. Fl. Atlant.* i. 253. t. 67.—Reg. Mediterr.
sulcata, Ledeb. Fl. Ross. ii. 299 = F. Ferulago.
Sumbul, *Hook. f. Bot. Mag.* t. 6196.—Turkestan.

FERULA :—
sylvatica, *Bess. Enum. Pl. Volh.* 44.—Europ. or. ; Reg. Caucas.
sylvatica, Orph. ex Nym. Consp. 284 = montanica, *Nym.*
sylvatica, Szov. ex Boiss. Fl. Orient. ii. 999 = pauciflora.
Szowitziana, *DC. Prod.* iv. 173.—Armen. ; Persia.
tatarica, *Fisch. ex Spreng. Pugill.* i. 37.—Ross. austr.
tenerrima, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 863.—Soongaria.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 708.—Reg. Himal.
thyrsiflora, *Stokh. & Sm. Fl. Graec. Prod.* i. 191.—Ins. Creta.
tingitana, *Linn. Sp. Pl.* 247.—Syria ; Afr. bor.
tolucensis, H. B. & K. Nov. Gen. et Sp. v. 12. t. 418 = Pseudanum tolucensis.
Tschurowskiana, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 593 = Schtschurowskiana.
tortuosa, Webb & Berth. Phyt. Canar. ii. 161 = Foeniculum tortuosum.
varia, Trautv. ex Bull. Soc. Nat. Mosc. xxxix. (1866) i. 825 = Pseudanum varium.
verticillata, *Marsh. in Ann. Gen. Wetter.* ii. (1811) 249 (Quid ?).—Europ.
villosa, Walt. Fl. Carol. 115 = Archangelica hirsuta.
wolgensis, Willd. ex Ledeb. Fl. Ross. ii. 302 = nuda.

FERULAGO, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 97 = Ferula, Linn. (Umbellif.).
angulata, Boiss. Fl. Orient. ii. 1005 = Ferula angulata.
asparagifolia, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 321.—As. Min.
athoa, Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 202. —Europ.
Auckerii, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 328.—As. Min.
Barrelieri, Guss. Fl. Sic. Syn. i. 354, in obs. = Ferula sulcata.
bicumbellata, Pomel, Nouv. Mat. Fl. Atl. 142.—Alger.
Bourgaei, Boiss. Fl. Orient. ii. 1000 = Ferula armena, *DC.*
brachylobe, Boiss. & Reut. ex Boiss. Voy. Bot. Espagne, ii. 732 = Ferula brachyloba.
bracteata, Boiss. & Hausskn. ex Boiss. Fl. Orient. ii. 1008.—Syria.
canduchorum, Boiss. & Hausskn. ex Boiss. l. c. 1005.—Assyria.
Cassia, Boiss. Diagn. Ser. I. x. 38.—As. Min. ; Syria.
cinerea, Boiss. l. c. vi. 90.—Persia.
communis, Pet. ex Nym. Consp. 284 = Ferula sulcata.
contracta, Boiss. & Hausskn. ex Boiss. Fl. Orient. ii. 1007.—Persia.
crassicosta, Pomel, Nouv. Mat. Fl. Atl. 143.—Alger.
Dodoneae, Kostel. Allg. Med.-Pharm. Flora, iv. 1158 = Ferula Ferulago.
frigida, Boiss. Diagn. Ser. I. x. 39.—Syria.
galbaniflora, Koch, Syn. Fl. Germ. ed. I. 302 = Ferula Ferulago.
geniculata, Boiss. in Ann. Sc. Nat. Ser. III. i. (1844) 324 = Ferula geniculata.
geniculata, Gmel. Fl. Sic. Syn. i. 354 = Ferula rigida.
granatensis, Boiss. Elench. 48 = Ferula granatensis.
humilis, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 322. —As. Min.
leptocarpa, Pomel, Nouv. Mat. Fl. Atl. 143.—Alger.
linearifolia, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 319 = Ferula angulata.
longifolia, Boiss. l. c. 321.—Cappadoc.
lophoptera, Boiss. l. c. 325 = Ferula lophoptera.
macrocarpa, Boiss. Fl. Orient. ii. 1002.—Mesopot.
macrocaulos, Boiss. & Bal. Diagn. Ser. II. vi. 82.—Phrygia.
meoides, Boiss. Fl. Orient. ii. 1004 = Ferula meoides.
monticola, Boiss. & Heldr. Diagn. Ser. II. ii. 91 = Ferula montícola.
monticola, Nym. Consp. 284, sub n. 7 = Ferula sylvatica.
nodiflora, Koch, in Nov. Act. Nat. Cur. xii. t. (1824) 98 = Ferula sulcata.
nodosa, Boiss. Diagn. Ser. I. x. 37 = Ferula nodosa.
Orphanidis, Boiss. & Heldr. l. c. Ser. II. ii. 91 = Ferula montícola.
oxyptera, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 320 = Ferula sylvatica.
pachyloba, Boiss. Fl. Orient. ii. 1006 = Ferula pachyloba.
parviflora, Pomel, Nouv. Mat. Fl. Atl. 211.—Alger.

FERULAGO:—
pauciflora, C. Koch, in Linnaea, xvi. (1842) 358 = Ferula pauciflora.
paucivittalata, Boiss. & Heldr. Diagn. Ser. I. x. 87.—As. Min.
platycarpa, Boiss. & Bal. l. c. II. vi. 32 = Ferula armena, *DC.*
Sadleri, Griseb. ex Nym. Consp. 284 = Ferula Sadleriana.
Sarteri, Boiss. Fl. Orient. ii. 999.—Ins. Cyclad.
scabra, Fenzl, in Flora, xxvi. (1843) 461.—Mesopot.
scabra, Fomel, Nouv. Mat. Fl. Atl. 143.—Alger.
setifolia, C. Koch, in Linnaea, xvi. (1842) 358 = Ferula pauciflora.
silaifolia, Boiss. Fl. Orient. ii. 1004.—Bithynia.
stellata, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 333. —Persia.
sulcata, Ledeb. Fl. Ross. ii. 299 = Ferula sulcata.
sylvatica, Reichb. Fl. Crit. iv. *f.* 555 = Ferula sylvatica.
sylvatica, Szov. ex C. Koch, in Linnaea, xvi. (1842) 358 = Ferula pauciflora.
syriaca, Boiss. Diagn. Ser. I. x. 88.—Syria.
Thirkeana, Boiss. Fl. Orient. ii. 1004.—Anatolia.
thyrsiflora, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 98 = Ferula thyrsiflora.
trachycarpa, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 319.—As. Min.
transiliensis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 80 = Ferula sylvatica.
trifida, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 318. —Persia.

FESTUCA, [Tourn.] Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1198.
AMPHIGENES, Janka, in Linnaea, xxx. (1859–60) 619.
CASTELLIA, Tineo, Pl. Rar. Sic. 17 (1817).
CATAPODIUM, Link, Hort. Berol. i. 64 (1827).
CHLOAENIA, Rafin. Neogenyt. 4 (1825).
DISTOMISCHUS, Dulac, Fl. Hautes-Pyr. 91 (1867).
DRYMONARTES, Ehrh. Beitz. iv. 147 (1789).
FESTUCARIA, Link, in Linnaea, xvii. (1843) 398.
GOUINIA, Fourn. ex Benth. & Hook. f. Gen. iii. 1178 (1883).
HELLERIA, Fourn. l. c. 1199 (1883).
LORETIA, Duval-Jouv. in Rev. Sc. Nat. Sér. II. ii. (1880) 88.
MICROPYRUM, Link, in Linnaea, xvii. (1843) 397.
MYGALURUS, Link, Enum. Hort. Berol. i. 92 (1821).
NARDURUS, Reichb. Nom. 89 (1841).—*Cf.* ej. Fl. Germ. Excurs. 19, in obs.
PROSPHYTES, Dulac, Fl. Hautes-Pyr. 67 (1867).
SCHEDONORUS, Beauv. Agrost. 99. t. 19. f. 9 (1812), partim.
SCLEROCHLOA, Reichb. Ic. Fl. Germ. i. 23. t. 58 (1854).
SCLEROPOA, Griseb. Spicil. Fl. Rumel. ii. 431 (1844).
SYNAPSIS, Dulac, Fl. Hautes-Pyr. 90 (1867).
VULPIA, C.C. Gmel. Fl. Bad. i. 8 (1805).
ZERNA, Panz. in Denkschr. Akad. Muench. 1818 (1814) 296, partim.
abbreviata, *Lindl. ex Fleisch. Syst. Vers. Ostsee-Prov. Phanerog.* 22.—Russia balt.
abyssinica, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 20, nomen; *et in A. Rich. Tent. Fl. Abyss.* ii. 482.—Abyss.
acanthophylla, *E. Desv. in C. Gay, Fl. Chil.* vi. 484.—Chili.
acerosa, C. Koch, in Linnaea, xix. (1847) 6 = punctoria.
acuminata, Gand. Agrost. Helv. ii. 287 = varia.
acutiflora, *Steud. Pl. Sest.* ex. III. 39.—Am. bor.
adscendens, Retz. in Vet. Acad. Handl. Stockh. (1769) 245 = *loliacea.*
aequiglumis, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 581.—Mexic.
aetnensis, Presl, Cyp. et Gram. Sic. 56 = Poa violacea.
aetnensis, Walp. Ann. i. 948 = ciliata.
affinis, Boiss. & Heldr. Diagn. Ser. II. iv. 127 = spectabilis.
agrestis, Lojac. Fl. Gall. ed. II. i. 87 = bromoides.
agrestis, Wulf. ex Steud. Nom. ed. II. i. 628 = ovina.
agrestium, *Le Pilego, in Mém. Soc. Linn. Par.* iv. (1826) 421.—"Terra Nova."
airoides, Lam. Encyc. ii. 464 = Poa nemoralis.
alaica, Kit. in Linnaea, xxxii. (1863) 813 = membranacea.
albida, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 10. —Ins. Madera.
alepplica, *Hochst. ex Steud. Syn. Pl. Gram.* 302.—Syria; Persia.
alaina, *Spreng. Syst.* i. 356.—Hab. ?
alopecuroides, Savi, Cose Bot. 59 = F. Alopecurus.
alopecuroides, Tenore, Fl. Nap. iv. 15 = F. Myurus.
Alopecurus, *Schousb. Vest. Marokko*, 40.—Reg. Mediterr. occ.
Alopecurus, Brongn. in Duperr. Voy. Bot. 82 = F. Arundo.
alpestris, Roem. & Schult. Syst. ii. 732 = varia.
alpestris, Wulf. Fl. Noric. Phan. ed. Fenzl & Graf, 151 = rubra.
alpigena, Hoppe, in Flora, i. (1818) 329 = rubra.
alpina, Host, Gram. Austr. iv. 36. t. 63 = varia.
alpina, Suter, Fl. Helv. i. 53 = ovina.
altaica, *Trin. in Ledeb. Fl. Alt.* i. 100.—Sibiria.
altissima, All. Auct. Fl. Pedem. 48 = sylvatica.
altissima, Boiss. Elench. 144 = procumbens.
ambigua, La Gall, Fl. Mantissa, 781 = F. Myurus.
americana, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 332 = bosciodes.
americana, Michx. ex Beauv. Agrost. 162 = Diarrhena americana.
amethystea, Pourr. ex Nym. Consp. 837 = rubra.
amethystina, All. Fl. Pedem. ii. 253 = varia.
amethystina, Delarb. Fl. Auvergne, ii. 694 = rubra.
amethystina, Host, Gram. Austr. ii. 53 = ovina.
amethystina, *Linn. Sp. Pl.* 74.—Europ. austr. : As. bor.
ammophyla, *Dum. Obs. Gram. Belg.* 105.—Belgium.
ampla, *Hack. Cat. Gram. Portugal*, 26.—Hispan.
amplifiora, Doell, in Mart. Fl. Bras. ii. iii. 116 = amplissima.
amplissima, *Rupr. in Bull. Acad. Brux.* ix. ii. (1842) 238.—Mexic.
ancepa, *Ard. in Linnaea*, xxxii. (1863) 814.—Hungar.
andicola, *H. B. & K. Nov. Gen. et Sp.* i. 153.—Ecuador.
angustata, *Griseb. in Goett. Abh.* xxiv. (1879) 288.—Ins. Argent.
anomala, *Hochst. in Flora*, xxxviii. (1855) 830.—Abyss.
antarctica, *Kunth, Rth. Gram.* i. 182.—Ins. Falkl.
antarctica, Spreng. Pugill. ii. 28 = flabellata.
anthoxanthia, Sm. in Trans. Linn. Soc. I. (1791) 115 = spadicea.
antucensis, *Steud. Syn. Pl. Gram.* 304.—Chili.
apennina, De Not. Prosp. Fl. Ligust. 56 = elatior.
aquatica, Bosc, ex Roem. & Schult. Syst. ii. 615 = Diplachne fasciularis.
arabica, Hort. ex Link, in Linnaea, xvii. (1847) 402 (= *Vulpia subciliata*).—Oraecia.
arctica, Schur, Enum. Pl. Transs. 799 = elatior.
arenaria, Dum. Obs. Gram. Belg. 108 = ovina.
arenaria, Kit. ex Steud. Nom. ed. II. i. 628 = ovina.
arenaria, *Lam. Tabl. Encyc.* i. 191.—Reg. Magellan.
arenaria, Batt. Dis. Suppl. R. Scud. (1856) 4 = rubra.
argaea, Boiss. & Bal. ex Boiss. Fl. Orient. v. 609 = Poa violacea.
argentea, Savi, in Uster, Ann. xxiv. (1800) 49 = Scleria cylindrica.
aristata, C. Koch, in Linnaea, xxi. (1848) 413 = gigantea.
articulata, De Not. ex Parl. Fl. Ital. i. 455 = elatior.
asundinacea, Lilljeh. Sv. Fl. ed. II. 47; ed. III. 70 = Graphephorum arundinaceum.
arundinacea, Schreb. Spicil. Fl. Lips. 57 = elatior.
arundinacea, *Vill. Prosp.* 17; *Hist. Fl. Dauph.* i. 262. —Europ.
Arundo, *Hook. f. Fl. Antarct.* 351.—Reg. Magellan.
Asheliana, Schult. Mant. ii. 294 = rubra.
aspera, Mart. & Koch, Deutschl. Fl. i. 672 = Bromus asper.
aspera, Pohr. Enczy. Suppl. ii. 635 = scabra.
aspera, Willd. ex Steud. Nom. ed. II. i. 628 = tolosensis.
asperula, *Phil. in Linnaea*, xxxiii. (1864–65) 296.—Chili.
aspertinsis, Hornem. ex Steud. Nom. ed. II. i. 626 = Poa nemoralis.
asperrima, Link, Hort. Berol. ii. 270 (an Festuceae sp. ?).—Persia.
atro-rubens, Wulf. Fl. Noric. Phan. ed. *Fenzl & Graf*, 151 = pallescens.
atrosanguinea, *Guss. Fl. Sic. Syn.* ii. 888.—Ital.

FESTUCA:—
aurata, Gand. Agrost. Helv. i. 234 = rubra.
auvata, Steud. Nom. ed. II. i. 624 = seq.
aurea, Lam. Fl. Fr. iii. 596 = spadicea.
australis, Nees, ex Steud. Syn. Pl. Gram. 304 = tenella.
australis, Schur, Enum. Pl. Transs. 796 = elatior.
austriaca, Hack. in Oestr. Bot. Zeitschr. xxviii. (1878) 349 = amethystina.
balearica, Gmelt. ex Zuccagni, in Roem. Collect. 195 = rubra ?
baltica, Homann, ex Mert. & Koch, Deutschl. Fl. i. 664; Fl. Posnn. i. 56 = rubra.
barbata, Lam. Dies. Dan. in Amoen. Acad. iii. 400 = Schismus marginatus.
barbata, Moench, Meth. Suppl. 63; Brol. Phyt. ii. 62 = F. Alopecurus.
barbata, Schrank, Prim. Fl. Salisb. 46 = rubra.
Barrelieri, Tenore, Prod. Fl. Nap. p. ix = Triticum Barrelieri.
Barthrei, Timb. in Bull. Soc. Hist. Nat. Toul. iv. (1870) 188 = rubra.
Beauguieriana, Schur, Enum. Pl. Transs. 787 = ovina.
Beckeri, Hack. in Bot. Centralb. viii. (1881) 405 = ovina.
Bellardi, Schott, ex Steud. Nom. ed. II. i. 628 = varia.
bellula, *Regel, in Act. Hort. Petrop.* vii. (1880) 594.—Turkestan.
Berteroniana, *Steud. Syn. Pl. Gram.* 311.—Chili.
biaristata, Ehrh. ex Boiss. Fl. Orient. v. 630 = inops.
bicornis, Schreb. ex Steud. Nom. ed. II. i. 628 = Glyceria acutiflora.
bifiora, *Steud. Syn. Pl. Gram.* 298.—Reg. Magellan.
Billardieri, Steud. l. c. 304 = Agropyrum scabrum.
blepharophora, Roem. & Schult. Syst. ii. 726 = Centotheca lappacea.
Boissieri, Janka, in Oestr. Bot. Zeitschr. xiv. (1864) 341 = montana.
bonariensis, Lam. Tabl. Encyc. i. 192 = Poa bonariensis.
borbonica, *Spreng. Syst.* i. 353.—Ins. Mascar.
Borderi, Hack. in Bot. Centralbl. viii. (1881) 406 = ovina.
borealis, Mert. & Koch, Deutschl. Fl. i. 664 = Graphephorum arundinaceum.
Borreri, Bab. in Trans. Linn. Soc. xvii. (1837) 565 = Glyceria Borreri.
bovinace, Kunth. & Sendtn. in Flora, xxxii. (1849) 764 = varia.
brachyphylla, Schult. Mant. ii. 246 = ovina.
bracteata, DC. Cat. Hort. Monsp. 111 (gen. dub.).—Hab. ?
brunnea, Facc. in Zeitschr. Ferdinand. Tirol. III. iv. (1854) 11 = Poa violacea.
brevifolia, R. Br. in Parry, 1st Voy. Suppl. p. cclxxxix = ovina.
brevifolia, Muhl. Desc. Gram. 167 = Diarrhena americanas.
brevisota, Steud. Syn. Pl. Gram. 317 = Triticum brevisetum.
Brinkmanni x, A. Br. Ind. Sem. Hort. Berol. (1861) App. 11.
brizoides, Spreng. Syst. i. 358 = Briza erecta.
brizoides, Wulf. Fl. Noric. Phan. ed. Fenzl & Graf, 141 = varia.
bromiformis, *Schleich. Cat. Pl. Helv.* ed. IV. 17, nomen.—Helvet.
bromoides, *Linn. Sp. Pl.* 76.—Reg. temp.
bromoides, Michx. Fl. Bor. Am. i. 66 = tenella.
Brotzi, Nym. Syll. 818 = hybrida.
Brotzi, Steud. Syn. Pl. Gram. 304 = F. Myurus.
Brownlanea, Steud. Syn. Pl. Gram. 304 = Agropyron scabrum.
Bryanii, F. Muell. Fragm. viii. 139 = Diplachne parviflora.
bulbosa, Bivona, Strp. Rar. Sic. Manip. iv. 7 = cserulescens.
caeruminis, Hensel. ex Ball, in Journ. Linn. Soc. xxii. (1885) 62, nomen = livida.
caerulata, Schur, Enum. Pl. Transs. 790 = amethystina.
caerulea, DC. Fl. Fr. iii. 48 = Molinia caerulea.
caerulescens, Desv. ex Hook. Mscrpt. Festuc. 337 = ovina.
caerulescens, *Desf. Fl. Atlant.* i. 97.—Reg. Mediterr.
caerulescens, Tenore, Fl. Nap. iii. 21 = glabra.
caespitosa, Sm. Engl. Bot. v. 1917 = ovina.
canaliculata, Desf. Fl. Atlant. i. 97 = Bromus madritensis.

FESTUCA :—
cespitosa, Roem. & Schult. Syst. ii. 782 = fabellata.
cespitosa, Willd. ex Spreng. Syst. i. 282 = borbonica.
cespitosula, Timb. in Bull. Soc. Hist. Nat. Toul. iii. (1869) 128 = ovina.
calabrica, *Huter, Porta & Rigo, Pl. Ital. Itin.* iii. (1877) n. 461 ; *ex Hack. Monogr. Fest.* 190.—Italia.
calamaria, Sm. Fl. Brit. i. 121 = sylvatica.
calcarea, Tausch, ex Stend. Nom. ed. II. i. 628 = ovina.
Calidasti, *Kunth, Rév. Gram.* i. 182.—Ecuador.
calycina, Loeß. Iter Hisp. 116 = Schismus marginatus.
cambrica, Huds. Fl. Angl. ed. II. i. 45 = rubra.
cana, Kit. ex Stend. Nom. ed. II. i. 629 = ovina.
canescens, Host, Fl. Austr. i. 155 = laxa.
canescens, Reichb. Ic. Fl. Germ. i. 26. t. 1541 = ovina.
capensis, Spreng. Syst. i. 252 = Lasiochloa ciliaris.
capillaris, Liljebl. Utk. Suec. ed. II. 46 = Glyceria distans.
capillaris, Wulf. Fl. Nor. Phan. ed. Fenzl & Graf. 144 = ovina.
capillata, Lam. Fl. Fr. iii. 597 = ovina.
capillata, Willd. ex Spreng. Syst. i. 256 = aliena.
capillifolia, Dufour, ex Roem. & Schult. Syst. ii. 735 = scabririma ?
captiata, Balb. ex Spreng. Syst. iv. Cur. Post. 36 = Dactylis capitata.
caprina, *Nees, Fl. Afr. Austr.* 448.—Afr. austr.
carolinizna, *Steud. Syn. Pl. Gram.* 312.—Carolina.
carpathica, F. G. Dietr. Vollst. Lexic. Gaertn. Nachtr. iii. 285.—Transsylv.
casapalensis, *Ball, in Journ. Linn. Soc.* xxii. (1885) 69.—Peruv.
caspica, Steud. Syn. Pl. Gram. 308 = Cutandia scleropoides.
cenisia, Hack. ex Boiss. Fl. Orient. v. 626 = sibirica.
cepacea, *Phil. in Linnaea*, xxxiii. (1864–65) 297.—Chili.
cernua, Schult. Oestr. Fl. ed. II. i. 248 = pulchella.
ciliaris, Heyne, ex Roem. & Schult. Syst. ii. 723 = Cenchrea lappacea.
ciliata, Danth. ex DC. Fl. Fr. iii. 55 = F. Myuros.
ciliata, Gouan, Hort. Monsp. 48, 547 = Brachypodium distachyum.
ciliata, Link, in Schrad. Journ. ii. (1799) 315 ; Brot. Fl. Lusit. i. 115 = F. Alopecuros.
ciliata, *Pers. Syn.* i. 94.—Europ. ; Oriens.
ciliata, Stend. Nom. ed. II. i. 629, sphalm. = *ciliaris*.
cinerea, DC. Fl. Fr. iii. 51 = rubra.
cinerea, Vill. Fl. Delphin. 8 ; Hist. Pl. Dauph. ii. 98 = ovina.
circinata, *Griseb. in Goett. Abh.* xxiv. (1879) 288.—Reg. Argent.
clandestina, *Muhl. Descr. Gram.* 162.—Am. bor.
clavata, *Mœnch, Meth.* 190.—Reg. Mediterr. occ.
Clementei, *Boiss. Elench.* 90.—Hispan.
coiron, *Steud. Syn. Pl. Gram.* 312.—Chili.
colorata, Schur, Enum. Pl. Transs. 788 = ovina.
Commersonii, Spreng. Syst. i. 256 = Brize subaristata.
commutata, *Steud. Syn. Pl. Gram.* 304.—Chili.
compacta, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 248. —Syria.
compressa, DC. Fl. Fr. vi. 368 = spadicea.
compressa, *Kit. in Linnaea*, xxxii. (1863) 811.—Hungar.
compressifolia, *Presl, Rel. Haenk.* i. 279.—Peruv.
confinis, *Vasey, in Bull. Torrey Bot. Club*, xi. (1884) 126.—Colorado.
consanitica, Timb. in Bull. Soc. Hist. Nat. Toul. iii. (1869) 130 = spadicea.
consolacia, Kunth, Rév. Gram. i. 129 = Glyceria distans.
Cookii, Hook. f. Fl. Antarct. 383. i. 129 = Poa Cookii.
cordata, Salzm. ex Stend. Nom. ed. II. i. 629 = ovina.
costata, *Nees, Fl. Afr. Austr.* 447.—Afr. austr.
crebra, Hack. ex Nym. Consp. 825 = elatior.
Crinum-ursi, Ramond, ex Schrad. Fl. Germ. i. 325 = varia.
cristata, C. C. Gmel. ex Roem. & Schult. Syst. ii. 660 = Trisetum Alopecurus.
cristata, Linn. Sp. Pl. 76 = Koeleria phleoides.
cristata, Vill. Hist. Pl. Dauph. ii. 96 = Koeleria cristata.
crocifolia, A. Kern. Sched. Fl. Exsicc. i. 106 = spectabilis.
cruciata, Steud. Nom. ed. I. 398, sphalm. = vida incisiana.
cuspidata, *Lag. ex* Willk. & Lange, Prod. Fl. Hisp. i. 92 = ovina.

FESTUCA :—
curvula, Gaudin, Agrost. Helv. i. 289 = ovina.
cyllenica, Boiss. & Heldr. Diagn. Ser. I. xiii. 58 = varia.
cynosuroides, Bal. ex Nym. Consp. 821 = gypsophila.
cynosuroides, Delile, Fl. Eg. Illustr. 107 = pectinella.
cynosuroides, *Desf. Pl. Atlant.* i. 82. t. 21.—Afr. bor.
dactyloides, Sibth. & Sm. Fl. Græc. Prod. i. 61 = Dactylis glomerata.
dasyantha, *E. B. & K. Nov. Gen. et Sp.* i. 154.—Ecuador.
debilis, *Schleich. Cat. Pl. Helv.* ed. IV. 17, nomen.—Helvet.
decidua, Bell. ex Sm. Engl. Bot. t. 2266 = sylvatica.
decipiens, [Claire.] Man. Herb. 34 = ovina.
decolorans, Mert. & Koch, Deutschl. Fl. i. 668 = elatior.
decumbens, Linn. Sp. Pl. 75 = Triodia decumbens.
delamarica, Kunth, Rév. Gram. i. 129 = Glyceria maritima.
delicatula, *Lag. in Varied. Cienc.* ii. iv. (1805) 39.—Lusit. ; Hisp.
delicatula, Savi, Cose Bot. 52 (= præc. ?).—Europ.
denudata, Dum. Obs. Gram. Belg. 104 = rubra ?
depauperata, Bertol. Fl. Ital. i. 620 = Poa Balbisii.
desertorum, *Phil. Pl. Atac.* 56.—Chili.
Desvauxii, *Phil. in Linnaea*, xxxiii. (1864–65) 295.—Chili.
diandra, Michx. Fl. Bor. Am. i. 67 = Diarrhena americana.
diandra, Moench, Meth. 191 = Brachypodium distachyum.
diaphana, Steud. Syn. Pl. Gram. 316 = Brachypodium flexum.
dichotoma, *Forsk. Fl. Aegypt. Arab.* 22.—Acgypt diffusa, *Dum. Obs. Gram. Belg.* 106.—Belgium.
digitata, Brouss. ex Horsem. Hort. Hafn. i. 96 = Diplachne fascicularis.
dimorpha, *Guss. Pl. Rar.* 84. t. 6.—Europ. austr.
dimorpha, Janka, in Oestr. Bot. Zeitschr. xvi. (1866) 101 = Brachypodium distachyum.
discolor, *Schur, in Verh. Siebenb. Ver. Naturw.* iii. (1852) 91.—Transsylv.
dissitiflora, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
distachyon, Roth, Catalect. fasc. i. 11 = Brachypodium distachyum.
distans, L. Kunth, Rév. Gram. i. 129 = Glyceria distans.
distichophylla, Michx. Fl. Bor. Am. i. 67 = Distichlis maritima.
divaricata, Clem. ex Willk. in Bot. Zeit. xviii. (1860) 130 = Cutandia scleropoides.
divaricata, Desf. Fl. Atlant. i. 80. t. 22 = Cutandia divaricata.
divaricata, *Roth, Catalect. fasc.* ii. 4.—Europ.
divaricata, Sieber, ex Steud. Nom. ed. II. i. 629 = Catandia maritima.
diversifolia, Bolss. & Bal. in Bull. Soc. Bot. Fr. iv. (1857) 306 = Poa diversifolia.
dives, F. Muell. Fragm. iii. 147 = Glyceria dives.
djimilensis, Boiss. & Bal. in Bull. Soc. Bot. Fr. xxi. (1874) 15 = rubra.
dolichophylla, *Ty & C. Presl, Rel. Haenk.* i. 258.—Peruv.
dominguensis, Lam. Tabl. Encyc. i. 189 = Leptochloa dominguensis.
donaciua, Wahlenb. Fl. Suec. i. 66 = Graphephorum arundinaceum.
Donax, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 9. —Ins. Madeir.
Dregeana, Mert. & Koch, Deutschl. Fl. i. 670 = montana.
dubia, *A. Rich. Tent. Fl. Abyss.* ii. 436.—Abyssin.
dumetorum, Linn. Sp. Pl. ed. II. 109 = rubra.
dumetorum, *Phil. in Linnaea*, xxxiii. (1864–65) 297.—Chili.
dura, Host, Gram. Austr. ii. 62. t. 87 = ovina.
dura, Vill. Hist. Pl. Dauph. ii. 94 = Sclerochloa dura.
Durandii, Clauson, in Billot, Annot. 163 = spadicea.
duriuscula, Linn. Sp. Pl. 74 = ovina.
duriuscula, Moris, ex Parl. Giorn. Bot. Ital. ii. (1844) 377 = Moriziana.
duriuscula, O. F. Muell. in Fl. Dan. t. 848 = rubra.
elatior, *Linn. Sp. Pl.* 75.—Europ. ; Ax. bor.
elatior, Uoria, Hort. Reg. Panorm. 90 = Aspelodesma tenax.
elegans, *Boiss. Elench.* 92.—Hispan.

FESTUCA :—
elongata, Ehrh. Beitr. vi. 58, 133 = *loliacea*, Huds.
eminens, *Kunth, Rév. Gram.* i. 182.—Ecuador.
eragrostoides, *Kroch. Pl. Siles. Suppl.* i. 171.—Europ.
eructa, Kunth, Enum. Pl. i. 378, in syn., sphalm. = Brize erecta.
eructa, *Ure, in Mem. Soc. Linn. Par.* iv. (1826) 601.—Reg. Magellan.
erecta, Wallr. Sched. Crit. 35 = Bromus erectus.
eriolepis, *E. Desv. in C. Gay, Fl. Chil.* vi. 438.—Chili.
Eshia, Lej. Fl. Spa, i. 52 = ovina.
Eshia, Ram. ex DC. Fl. Fr. iii. 52 = varia.
exaltata, Presl, Fl. Sic. p. xiv = montana.
expansa, Kunth, Rév. Gram. i. 129 = Sphenopus Gouani.
fagetina, Schur, Enum. Pl. Transs. 797 = montana.
fallax, Thuill. Fl. Par. ed. II. 80 = rubra.
fascicularis, Lam. Tabl. Encyc. i. 189 = Diplachne fascicularis.
fasciculata, Forsk. Fl. Aegypt. Arab. 22 = bromoides.
Fenzi, Lag. Gen. et Sp. Nov. 4 = elatior.
ferruginea, Reichb. Fl. Germ. Excurs. 140* = spadicea.
fibrosa, Griseb. Spicil. Fl. Rumel. ii. 435 = spadicea.
filifolia, Link, Enum. Hort. Berol. i. 448 = ovina ?
filiformis, Lam. Tabl. Encyc. i. 191 = Leptochloa filiformis, procera.
filiformis, *Ch. Sm. ex Link, in Buch, Beschr. Canar. Ins* 139.—Ins. Canar.
filiformis, *Steud. Syn. Pl. Gram.* 302.—Nepal.
fimbriata, *Nees, Agrost. Bras.* 472.—Reg. Argent.
flabellata, Lam. Encyc. ii. 322 = Poa flabellata.
flaccida, Schur, in Verh. Siebenb. Ver. Naturw. iv (1853) 89 = rubra.
flava, F. Muell. Sel. Pl. Indust Cult. 87 = Triodia sclerioides.
flavescens, Bell. in Mém. Acad. Turin, v. 1790, 179 ! (1793), 217 = varia.
Fleischeri, Steud. Nom. ed. II. i. 630 = spadicea.
flexa, Steud. Syn. Pl. Gram. 316 = Brachypodium flexum.
flexuosa, Willd. ex Kunth, Enum. Pl. i. 407 = quadridentata.
fluitans, Leers, ex Kunth, l. c. 406 = *loliacea*, Huds.
fluitans, Linn. Sp. Pl. 76 = Glyceria fluitans.
Forsica, Auct. vide *Festuca*.
foliata, *Steud. Syn. Pl. Gram.* 304.—Sicilia.
foliosa, Hook. f. Fl. Antarct. 39. t. 56 = Poa foliosa.
foliosa, *Schrad. ex Steud. Nom.* ed. II. i. 630, nomen.—Hab. ?
Fournieriana, *Hemsl. Biol. Centr. Am. Bot.* iii. 581.—Mexic.
francoula, *Rupr. in Bull. Acad. Brux.* ix. ii. (1842) 236.—Mexic.
frigida, Hack. in Bot. Centralb. viii. (1881) 406 = ovina.
fuegiana, *Hook. f. Fl. Antarct.* 380. t. 141.—Reg. Magellan.
fusca, Linn. Sp. Pl. ed. II. 109 = Diplachne fusca.
fusca, Mutel, Fl. Fr. iv. 105 = varia.
fusca, Vill. Hist. Pl. Dauph. ii. 98 = spadicea.
Gaudini, Kunth, Rév. Gram. i. 130 = ovina.
geniculata, Cav. in Anal. Cienc. Nat. vi. (1803) 150 ; Brot. Fl. Lusit. i. 118 = elatior.
geniculata, Pollini, Fl. Veron. i. 296 = Triticum gemmeum.
genuflexa, Krock. Fl. Siles. Suppl. i 172 = refracta.
Gibraltarica, Willd. ex Steud. Nom. ed. II. i. 630 = F. Alopecuros.
gigantea, Krock. Fl. Siles. Suppl. i. 174 = spadicea !
gigantea, *Vill. Hist. Pl. Dauph.* ii. 110.—Europ. ; Ax. bor. ; Afr. trop.
glaberrima, Schur, Enum. Pl. Transs. 791 = ovina.
glabra, Lightf. Fl. Scot. ii. 1085 = rubra.
glabra, *Spreng. Syst.* i. 258.—Am. bor.
glacialis, Miegev. in Bull. Soc. Bot. Fr. xxi. (1874) p. ix = ovina.
glauca, Lam. Encyc. ii. 459 = ovina.
glauca, Plan. ex Willk. & Lange, Prod. Fl. Hisp. i. 95 = rubra.
glaucescens, Hegetschw. Fl. Schw. 93 = rubra.
glaucescens, Roth, Nov. Pl. Sp. 78 = Koeleria glauca.
glomerata, All. Fl. Pedem. ii. 252 = Dactylis glomerata.
glomerata, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 248. —Syria.
glycericoides, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.

FESTUCA:—
gracilenta, Buchl. in Proc. Acad. Sc. Philad. 1862 (1863) 97 = microstachya.
gracilis, Bernh. Syst. Verz. Erf. 46 = Brachypodium pinnatum!
gracilis, *Kunth, Rév. Gram.* i. 129.—Gallia merid.!
gracilis, Moench, Meth. 191 = Brachypodium sylvaticum.
gracillima, *Hook. f. Fl. Antarct.* 383.—Calif.; Reg. Magellan.
granatensis, *Boiss. Elench.* 93.—Hispan.
graminicola, Willk. in Linnæa, xxv. (1852) 66 = cærulescens.
grandiflora, Lam. Tabl. Encyc. i. 191 = Arundinaria macrosperma.
grandiflora, Spreng. ex Steud. Nom. ed. II. i. 630 = Glyceria acutiflora.
grandiflora, Steud. Syn. Pl. Gram. 311 = livida.
graniflora, Boeming. ex Reichb. Fl. Germ. Excurs. 149 = ovina.
graniphalice, Schur, Enum. Pl. Transs. 790 = amethystina.
Granzowia, Trin. ex Steud. Nom. ed. II. i. 630 = sicula.
gypsophila, *Hack. in Oestr. Bot. Zeitschr.* xxvii. (1877) 47.—Hispan.
Hamiki, *Kunth, Enum. Pl.* i. 403.—Ferav.
Halleri, All. Fl. Pedem. ii. 258 = ovina.
Halleri, Pucc. in Parl. Giorn. Bot. Ital. i. 1. (1844) 191 = rubra.
hallerioides, Schur, Enum. Pl. Transs. 790 = ovina.
halmyris, Mabille, in Act. Linn. Soc. Bord. xxv. (1864) 61 = rubra.
hemipon. *Delile, ex Spreng. Syst.* iv. *Cur. Post.* 36.—Europ. austr.; Afr. bor.
Henriquezii, *Hack. in Bot. Centralb.* viii. (1881) 406, nomen; *Monog. Fest.* 126.—Lusitan.
hephaestophila, *Nees. ex Steud. Syn. Pl. Gram.* 310.—Mexic.
heptantha, C. Koch, in Linnæa, xxi. (1848) 410 = Poa persica.
heteromalla, Pourr. in Mém. Acad. Toul. iii. (1788) 319 = elatior.
heterophylla, Lam. Fl. Fr. iii. 600; Hasnk. in Jacq. Coll. ii. 93 = rubra.
heterophylla, Wahlenb. Veg. Helv. 18 = ovina.
hirsuta, DC. Fl. Fr. iii. 58 = Koeleria hirsuta.
hirsuta, Host, Gram. Austr. ii. 61. t. 85 = ovina.
hirsuta, *Nonch. Meth. Suppl.* 62.—Hab.?
hirta, Sesseu, Reise Istrien, 59 = Bromus erectus.
hispanica, Kunth, Rév. Gram. i. 129 = tenuiflora.
hispida, Savi, Fl. Pis. i. 117 = Koeleria hispida.
Hookeriana, *F. Muell. ex Hook. Fl. Tasm.* ii. 127. t. 165.—Anstral.
hordeiformis, Wulf. ex Steud. Nom. ed. I. 387 = bromoides.
Hostii, Kunth, Rév. Gram. i. 129 = Glyceria festucæformis.
Hostii, Schott, ex Roem. & Schult. Syst. ii. 722 = varia.
humilior, [*Nees, in*] *Mayen. Reise*, i. 406, nomen; *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 166.—Peruv.
hybrida, *Brot. Fl. Lusit.* i. 115.—Hispan.
Hystrix, *Boiss. Elench.* 80.—Hispan.
inermata, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 177 = amethystina.
incrassata, *Salem. ex Lois. Fl. Gall.* ed. II. i. 86.—Italia.
indica, Retz. Obs. iv. 21 = Diplachne indica.
indigesta, Boiss. Elench. 91 = ovina.
inermis, DC. Fl. Fr. iii. 49 = Bromus inermis.
insignis, Schur, Enum. Pl. Transs. 792 = rubra.
inops, *Delile, ex Steud. Nom.* ed. I. 387, nomen; *Boiss. Fl. Orient.* v. 630.—Aegypt.
inops, De Not. Report. Fl. Ligust. ii. 506 = ovina.
insularis, *Steud. Syn. Pl. Gram.* 313.—Calif.
intermedia, Roem. & Schult. Syst. ii. 715 = ovina.
interrupta, Desf. Fl. Atlant. i. 99 = elatior.
inundata, Moench, ex Roem. & Schult. Syst. ii. 790 = ovina.
invidiana, F. Muell. Veg. Chath. Isl. 59 = Triodia triflora.
Ivyana, *F. Muell. Veg. Chath. Isl.* 59 = Triodia triflora.
Jaxarty, *Vasey, Grass. U. St.* 43, nomen.—Am. bor. occ.
jubata, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimp. 6.—Ins. Asor.

FESTUCA:—
juncea, Moench, Meth. 190 = Agropyron junceum.
juncifolia, St. Am. Fl. Agen. 46 = rubra.
jurana, Gren. ex Nym. Consp. 836 = pulchella.
kergueleusis, *Hook. f. in Phil. Trans.* clxvii. (1879) 14.—Ins. Kerguel.
Kinglana, *Steud. Syn. Pl. Gram.* 316.—Ins. Norfolk.
Kirelowii, *Steud. l. c.* 306.—Soongar.
Klattebeliana, Schalt. Mant. ii. 296 = rubra.
Kunzei, *Regel, in Act. Hort. Petrop.* vii. (1880) 594.—Turkestan.
Lachenalii, Spenn. Fl. Friburg. iii. 1050 = F. Poa.
laevigata, Gaud. in Alpies, iii. (1906) 60 = ovina.
laevis, Hack. in Bot. Centralb. viii. (1881) 408 = ovina.
laevis, *Kit. in Linnæa*, xxii. (1863) 813.—Hunger.
laevis, Spreng. Syst. i. 356 = Lasiochloa laevis.
Lagascæ, Hort. Berol. ex Steud. Nom. ed. II. i. 630 = varia.
lanata, Spreng. Syst. i. 358 = Poa lanuginosa.
lancelata, Forsk. Fl. Aegypt. Arab. 22 = Cutandia lancelata.
lanuginosa, Scheele, in Flora, xxvii. (1844) 63 = elatior.
lasiantha, Schur. Enum. Pl. Transs. 788 = ovina.
latifolia, *DC. Cat. Hort. Monsp.* 111.—Hab.?
latifolia, Dulac, Fl. Hautes-Pyr. 92 = elatior.
latifolia, Host, Fl. Austr. i. 153 = sylvatica.
latifolia, Roth, Nov. Pl. Sp. 75 = Cenotheca lappacea.
latispicea, F. Muell. Fragm. viii. 127 = Glyceria latispicea.
laxa, Gaud. Agrost. Helv. i. 261 = elatior.
laxa, *Host, Gram. Austr.* iv. 25.—Europ. or.
laxa, Schleich. ex Steud. Nom. ed. II. i. 630 = varia.
laxiflora, *Phil. in Linnæa*, xxix. (1857–58) 98.—Chili.
laxiflora, *A. Rich. Fl. Cub. Fanerog.* ii. 318.—Cuba.
Lechleriana, *Steud. Syn. Pl. Gram.* 435.—Reg. Magellan.
Lemani, Bast. Ess. Fl. Maine-et-Loire, 36 = ovina.
leptothrix, *Trin. ex Steud. Nom.* ed. II. i. 630, nomen; *Doell, in Mart. Fl. Bras.* ii. 111. 115.—Bras.
leucochæta, Schur, Enum. Pl. Transs. 916, sphalm., vide *lasiantha*.
Leysseri, Moench, Enum. Pl. Hass. 41 = Bromus inermis.
Liebmanni, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 583.—Mexic.
ligustica, *Bertol. Amoen. Ital.* 8.—Italia.
linearis, Willk. Exercit. ex Fl. Hispan. 6 = F. Myuros.
Linnæana, Steud. Syn. Pl. Gram. 308 = bromoides.
lithuanica, Aschers. in Bot. Zeit. xxv. (1867) 160 = montana!
litorea, Hack. Monog. Fest. 153, sphalm. = elatior.
littoralis, *Labill. Nov. Holl. Pl.* i. 23. t. 37.—Austral.
littoralis, Sibth. & Sm. Fl. Græc. Prod. i. 61; Fl. Græc. t. 80 = Aeluropus lævis.
littoralis, Steud. Syn. Pl. Gram. 317 = Brachypodium ramosum.
littoralis, Wahlenb. in Nov. Act. Soc. Sc. Upsal. viii. (1821) 211, 238 = elatior.
livida, Schur, Enum. Pl. Transs. 787 = ovina.
livida, *Willd. ex Spreng. Syst.* i. 353.—Mexic.
loliacea, Gmel. Fl. Angl. ed. I. 38.—Europ.
loliacea, Lam. Encyc. ii. 462 = elatior.
loliiformis, F. Muell. Fragm. viii. 128 = Diplachne loliiformis.
loliiodes, Trin. ex Steud. Nom. ed. II. i. 630 = tenuiflora.
Lolium, *Salzmn, ex Cass. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 211.—Alger.
longearistata, Walp. Ann. i. 948 = Agropyron longearistatum.
longifolia, Hegetschw. Fl. Schw. 91 = rubra.
longiflolia, Vis. in Ann. Bot. i. ii. (1804) 145; Thuill. Fl. Par. ed. II. 60 = ovina.
longiseta, Brot. Fl. Lusit. i. 116 = bromoides.
longiseta, Hegetschw. Fl. Schw. 93 = rubra.
lubrica, Lapeyr. Hist. Abr. Pl. Pyr. 44 = varia.
lucorum, Schur, Enum. Pl. Transs. 797 = montana.
macrophylla, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 464.—Abyssin.
macrostachya, *Desf. Tabl. ed.* 1. 216, nomen.—Hab.?
madritensis, Desf. Fl. Atlant. i. 91 = bromoides.
magellanica, *Lam. Encyc.* ii. 461; *Poir.* i. 380.—Reg. Magellan.
maritima, DC. Fl. Fr. iii. 57 = Catapodia maritima.

FESTUCA:—
maritima, Linn. Sp. Pl. 75 = tenuiflora.
maritima, C. Sm. ex Link, in Buch, Beschr. Canar. Ins. 139 = Smithii.
media, *Kit. in Linnæa*, xxxii. (1863) 814.—Hungar.
megalura, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 188 = F. Myuros.
megaphylla, Schur, Enum. Pl. Transs. 789 = ovina.
megastachya, Hegenschw. Fl. Schw. 92 = rubra.
melangeni, Spreng. Syst. i. 353 = Lasiochloa hispida, membranacea, *Kit. in Linnæa*, xxxii. (1863) 813.—Hungar.
membranifolia, Boiss. ex Cos. Notes Crit. 183 = Catapodia scleropoides.
mexicana, Roem. & Schult. Syst. ii. 732 = Brachypodium mexicanum.
mexicana, Willd. ex Spreng. Syst. i. 356 = Willdenowiana.
Michelii, Kunth, Enum. Pl. i. 397 = Avellinia Michelii.
microstachya, F. Muell. Veg. Chath. Isl. 60 = Triodia microstachya.
microstachya, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 187.—Am. bor. occ.
minima, *A. Rich. Tent. Fl. Abyss.* ii. 486.—Abyssin.
minor, Schur, Enum. Pl. Transs. 795 = varia.
minuta, Hoffm. in Comm. Soc. Phys. Mosc. i. (1808) 48; ex Ledeb. Fl. Ross. iv. 404 = Schismus minutus.
misera, *Thunb. Fl. Jap.* 59.—Japon.
modesta, *Steud. Syn. Pl. Gram.* 314.—Nepal.
mollis, *Kunth, Rév. Gram.* i. 132.—Ecuador.
monandra, Ell. ex Steud. Nom. ed. II. i. 631 = F. Myuros.
monastachia, Poir. Voy. Barb. ii. 98 = Brachypodium distachyum.
montana, *Bieb. Fl. Taur. Cauc.* iii. 75.—Europ.; As. Min.
montana, Savi, Fl. Pis. i. 118 = Bromus erectus.
montana, Sternb. & Hoppe, in Denkschr. Bot. Ges. Regensb. i. ii. (1818) 96 = Poa hybrida.
montana, *Steud. Syn. Pl. Gram.* 302.—Hispan.
monticola, *Phil, in Anal. Univ. Chil.* (1873) 576.—Chili.
Montis-auvei, Delarb. Fl. Auvergne, ed. II. 696 = alpina.
Montis-celsi, Delarb. l. c. = rubra.
Moricciana, *Parl. Giorn. Bot. Ital.* ii. (1844) 377.—Sardinia.
mucronata, *Arnot & Dupuy, in Bull. Soc. Bot. Fr* xxix. (1882).—*Cf. Bibl.* 48.—Gallia.
mucronata, Forsk. Fl. Aegypt. Arab. 22 = Aeluropus arabica.
mucronata, C. Muell. in Walp. Ann. vi. 1033 = Brachypodium mucronatum.
multiculmis, Steud. Syn. Pl. Gram. 310 = tolucensis.
multiflora, Hoffm. Deutschl. Fl. ed. II. i. t. 50 = rubra.
multiflora, Presl, Cyp. et Gram. Sic. 37 = elatior.
multiflora, Steter, Fl. Helv. i. 56 = ovina.
multiflora, Walt. Fl. Carol. 81 = Diplachne fascicularis.
muralis, *Kunth, Syn. Pl.* i. 216.—Mexic.
muricata, Dur. ex Parl. Fl. Ital. i. 480 = tuberculosa.
mutica, Schleich. ex Hack. Monog. Fest. 152 = amethystina.
mutica, Wulf. Fl. Noric. Phan. ed. Fenzl & Graf, 145 = ovina.
myurosoides, Heyne, ex Roem. & Schult. Syst. ii. 800 = Tripogon bromoides.
Myuros, *Gouan, Hort. Monsp.* 49 = ciliata.
Myuros, *Linn. Sp. Pl.* 74.—Europ.; Am. et As. bor.
Myuros, Soyer-Willem. Obs. Pl. Fr. 128 = F. Alopecuros.
Myurus, H. B. & K. Nov. Gen. et Sp. i. 255 = muralis.
Myurus, Moench, Meth. Suppl. 62 = bromoides.
Myurus, Mckl. Dem. Gram. 146 = quadriflora.
nana, Steud. Syn. Pl. Gram. 428 = dichotoma.
nardifolia, *Griseb. in Goett. Abh.* xviii. (1879) 286.—Reg. Argent.
nebrodensis, Jan, Elench. 2 = Poa violacea.
Neesiana, *Steud. Nom.* ed. II. i. 631.—Afr. austr.
nemorum, Lejeun. Chlor. Lugd. 6 = rubra.
nemorum, Lejeun. in Abh. Naturf. Ges. Halle, i. (1783) 268 = rubra.
neogaea, *Steud. Syn. Pl. Gram.* 338.—"Terra nova."
nepalensis, Steud. l. c. 316 = Brachypodium nepalense.
nervosa, Hook. Fl. Bor. Am. ii. 253 = Poa cæspitosa.
nevadensis, Hack. in Bot. Centralb. viii. (1885) 407 = rubra.

FESTUCA:—
nigra, Gilib. Exercit. ii. 533 = ovina.
nigrescens, Lam. Encyc. ii. 449 = rubra.
nigricans, Steud. Syn. Pl. Gram. 311 = rigescens.
nitida, Kit. in Schult. Oestr. Fl. ed. II. i. 239 = rubra.
nobigens, *Jungh. in Nat. en Geneesk. Arch. Neerl. Indie*, ii. (1845) 51.—Java.
nubila, Steud. Syn. Pl. Gram. 315 = prasc.
nuda, *Schur, in Verh. Siebenb. Ver. Naturw.* iii. (1852) 91.—Transylv.
nutans, Host, Gram. Austr. iv. 35. t. 61 = pulchella.
nutans, Moench, Meth. 191 = Agropyron caninum.
nutans, *Spreng. Fl. Hal. Mant.* 84; *Muhl. Desc. Gram.* 168.—Am. bor.
nutans, Wahlenb. Fl. Carp. Principi. 38 = carpathica.
Nuttalliana, Kunth, Rév. Gram. i. 129 = Glyceria distans.
obovata, Kit. *ex* Roem. et Schult. Syst. ii. 584.—Hungar.
obtusa, Sprung. Fl. Hal. Mant. 84 = patens.
obtusata, Michx. ex Beauv. Agrost. 168 = Reboulea gracilis.
obtusiflora, Willd. ex Spreng. Syst. i. 356 = Diplachne dubia.
obtusiflula, Lag. ex Link, Enum. Hort. Berol. i. 95 = Brachypodium ramosum.
occidentalis, Hook. Fl. Bor. Am. ii. 249 = pauciflora.
ochroleuca, Timb. in Bull. Soc. Hist. Nat. Toul. iii. (1869) 139 = ovina.
octoflora, Walt. Fl. Carol. 81 = tenella.
olandica, Hack. ex Nym. Consp. 827 = rubra.
oligostachya, Schur, Enum. Pl. Transs. 754 = ovina.
Onurus, Willd. ex Spreng. Syst. i. 354 = muralis.
orawensis, Steud. Syn. Pl. Gram. 313 = caerulescens.
oraria, Dun. Obs. Gram. Belg. 105 = rubra.
orcgona, *Vasey, in Coult. Bot. Gaz.* ii. (1877) 126.—Am. bor. occ.
orgyalis, *Willd. ex Spreng. Syst.* i. 351, in syn.—Mexic.
orientalis, Kern. ex Hack. Monog. Fest. 154 = elatior.
oryzetorum, Pollini, Veron. Pl. Nov. 3 = elatior.
ovirtana, Lej. Fl. Spa, i. 57 = ovina.
ovina, Brot. Fl. Lusit. i. 114 = ampla.
ovina, *Linn. Sp. Pl.* 73.—Reg. temp. et arct.
pallens, Host, Gram. Austr. ii. 63 = ovina.
pallida, Gaud. in Alpina, iii. (1808) 56 = rubra.
paludosa, Gaud. Agrost. Helv. i. 329 = ovina.
palustris, Seegus, Reise Lothr. 72 = Glyceria distans.
pannonica, Wulf. ex Host, Gram. Austr. iv. 36. t. 62 = ovina.
panormitana, Guss. Fl. Sic. Syn. ii. 886 = bromoides.
paradoxa, *Desv. Opuse.* 105.—Hab.?
paradoxa, Ehrh. ex Steud. Nom. ed. II. i. 631 = spadicea.
parviflora, *Ell. Sketch.* i. 170.—Carolina.
parviglumis, Steud. Syn. Pl. Gram. 305 = pauciflora.
parvula, *Kunth, Rév. Gram.* i. 129.—Oriens.
patens, Steud. Syn. Pl. Gram. 298 = pectinella.
pectin, *Desf. Fl. Atlant.* i. 86.—Afr. bor.
pauciflora, Schleich. Cat. Pl. Helv. ed. IV. 17 = ovina.
pauciflora, *Thunb. Fl. Jap.* 52.—Japon.; Am. bor. occ.
pectinata, Labill. Nov. Holl. Pl. i. 21. t. 25 = Agropyron pectinatum.
pectinella, *Delile, Ind. Sem. Hort. Monsp.* (1836) 24; *ex Cass. & Dur. Expl. Alger.* 179.—Aegypt.
pendulina, Spreng. Syst. i. 356 = Bromus ciliatus.
pentastachya, Presl, Fl. Sic. p. xiv = Bromus pentastachyus.
persica, C. Koch, in Linnaea, xxi. (1848) 410 = Poa persica.
petraea, *Gaudin. ex Smith. Fl. Auser.* 20.—Ins. Azor.
phalaridoides, Lam. Tabl. Encyc. i. 190 = Dactylis glomerata, Koeleria phleoides.
phillitidea, Steud. Syn. Pl. Gram. 298 = Cutandia phillitidea.
phleoides, Vill. Fl. Delph. 7; Hist. Pl. Dauph. ii. 95 = Koeleria phleoides.
phoenicoides, All. Fl. Pedem. ii. 258 = Brachypodium ramosum.
phoenicoides, Linn. Mant. i. 28 = Brachypodium pinnatum.
phoenicoides, Rouillé, ex Roem. & Schult. Syst. ii. 699 = rubra.
Phoenix, Thrill. Fl. Par. ed. II. i. 58 = *intacea*, Huds.
Phoenix, Vill. Prosp. 17; Hist. Pl. Dauph. ii. 106 = elatior.
picta, J. F. Gmel. Syst. 187 = Panicum Crus-galli?

FESTUCA:—
picta, Kit. in Schult. Oestr. Fl. ed. II. i. 236 = rubra.
pilosa, Hall. f. ex Gaud. in Alpina, iii. (1808) 54 = Poa violacea.
pilosa, Willd. ex Spreng. Syst. i. 356 = Eragrostis mayparensis.
pinnata, Moench, Meth. 191; Huds. Fl. Angl. ed. II. i. 48 = Brachypodium pinnatum.
platyphylla, *Steud. Syn. Pl. Gram.* 428.—Reg. Magellan.
plebeia, R. Br. Prod. i. 178 = bromoides.
plicata, *Hack. in Oestr. Bot. Zeitschr.* xxvii. (1877) 48.—Hispan.
poliens, *Steud. Syn. Pl. Gram.* 317.—Hab.?
polyantha, Hall. f. ex Steud. Nom. ed. I. 338 = rubra.
polychroa, Trautv. in Act. Hort. Petrop. ii. (1873) 485 = Poa violacea.
polygama, C. Koch, in Linnaea, xxi. (1848) 409 = Poa persica.
polystachya, Michx. Fl. Bor. Am. i. 66 = Diplachne fascicularis.
Porcii, *Hack. in Bot. Centralb.* viii. (1881) 407, nomen; *Monog. Fest.* 147.—Transylv.
prasina, Dum. Obs. Gram. Belg. 102 = ovina?
pratensis, [Hoeck.] Verz. Aller Gew. Teutschl. 268 = ovina.
pratensis, Huds. Fl. Angl. ed. I. 37 = elatior.
pratensis, Schreb. Spicil. Fl. Lips. 57 = rubra.
procera, *H. B. & K. Nov. Gen. et Sp.* i. 154.—Ecuador.
procumbens, Kunth, Rév. Gram. i. 129 = Glyceria procumbens.
procumbens, Muhl. Desc. Gram. 160 = Diplachne fascicularis.
prostrata, Gmel. ex Steud. Nom. ed. II. i. 632 = Aelurops laevis.
Pseudairoides, Kcel. Desc. Gram. 270 = Brachypodium distachyum
Pseudo-dura, Steud. Syn. Pl. Gram. 306 = ovina.
Pseudo-duriuscula, *Steud. l. c.* 315.—Texas.
Pseudo-elatior, *Beitr. Flora Eichsch.* 21.—Hispan.
Pseudo-ixna, Schur, in Oestr. Bot. Zeitschr. viii. (1858) 29 = carpathica.
Pseudo-loliacea, Hack. Monog. Fest. 151, in syn. = elatior.
Pseudomyurus, Soyer-Willem. in Ann. Sc. Nat. Sér. I vii. (1826) 448 = F. Myuros.
Pseudo-nemorum, Schur, Enum. Pl. Transs. 794 = rubra.
Pseudo-nutans, Schur, l. c. 796 = pulchella.
Pseudo-ovina, Schur, l. c. 794 = rubra.
Pseudovina, Hack. ex Wiesb. in Oestr. Bot. Zeitschr. xxx. (1880) 126 = ovina.
pubescens, Willd. Enum. Hort. Berol. Suppl. 6 = rubra.
pubescens, Zea, ex Roem. & Schult. Syst. ii. 738 = F. Alopecurus.
Pucciniana, Parl. Fl. Ital. i. 440 = rubra.
pulchella, *Schrad. Fl. Germ.* i. 386. t. 5.—Europ.
pulchra, Schur, Enum. Pl. Transs. 785 = ovina.
pumila, Chaix, in Vill. Hist. Dauph. i. 316; ii. 102 = varia.
pumila, Willd. in Linnaea, xxv. (1852) 66 = ovina.
punctoria, *Sibth. & Sm. Fl. Graec. Prod.* i. 80.—Bithyn.
pungens, Kit. in Schult. Oestr. Fl. ed. II. i. 237 = varia.
pungens, Lam. Fl. Fr. iii. 596 = Brachypodium pinnatum.
pungens, Vahl, Symb. Bot. i. 10. t. 2 = Aelurops ambiens.
purpurascens, *Banks & Soland. ex Hook. f. Fl. Antarct.* 338. t. 140.—Reg. Magellan.
purpurea, F. Muell. Sel. Pl. Indust. Cult. 88 = Triplasis purpurea.
purpurea, Schreb. ex Steud. Nom. ed. II. i. 632 = Triodia cuprea.
pusilla, Banks & Soland. ex Hook. f. Fl. Antarct. 330 = Triodia antarctica.
pusilla, Bastd. in Proc. Acad. Sc. Philad. 1863 (1863) 96 = microstachya.

FESTUCA:—
pyrenaïca, Link, in Schrad. Journ. ii. (1799) 315 = bromoides.
pyrenaica, Rent. Cat. Sem. Hort. Genev. (1861) 4 = rubra.
quadridens, Poir. Encyc. Suppl. ii. 640 = Triodia cuprea.
quadridentata, *H. B. & K. Nov. Gen. et Sp.* i. 154.—Ecuador.
quadriflora, [Honck.]. Verz. Aller Gew. Teutschl. 268 = varia.
quadriflora, *Wall. Fl. Carol.* 31.—Am. bor.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* ii. 485.—Abyssin.
quiliensis, Sprung. Syst. i. 356 = Bromus unioloides.
quitensis, Willd. ex Kunth, Enum. Pl. i. 407 = Caldasii.
racemifera, Trin. ex Steud. Nom. ed. II. i. 632 = tenella.
racemosa, *Kit. in Linnaea*, xxxii. (1863) 218.—Hungar.
racemosa, Willd. ex Spreng. Syst. i. 352 = sadicola.
radicans, Steud. Syn. Pl. Gram. 309 = elatior.
ramosa, Gaus. Fl. Sic. Prod. i. 110 = Brachypodium pinnatum.
ramosa, Roth, Nov. Pl. Sp. 76 = Brachypodium ramosum.
rectiseta, Walp. Ann. i. 948 = Agropyron scabrum.
reflexa, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 98 = microstachya.
refracta, *Kroch. Fl. Siles. Suppl.* i. 178.—Europ.
remota, *Kit. in Linnaea*, xxxii. (1863) 218.—Hungar.
remotiflora, Steud. Syn. Pl. Gram. 315 = pauciflora?
repens, Knapp, Gram. t. 119 = rubra.
repens, *Trin. ex Steud. Nom.* ed. II. i. 632.—Hab.?
reptatrix, Linn. Sp. Pl. 108 = Diplachne fusca.
restituta, Steud. Syn. Pl. Gram. 314 = Schimperiana.
rhachiathra, Steud. l. c. 309 = Desmazeria castellana.
rhoetica, Suter, Fl. Helv. i. 56 = Poa violacea.
rhombifera, Kit. in Linnaea, xxxii. (1863) 214 = pulchella.
Richardsoni, *Hook. Fl. Bor. Am.* ii. 250.—Am. arct.
rigens, Trin. ex Steud. Nom. ed. II. i. 632 = rigida.
rigescens, *Kunth, Enum. Pl.* i. 408.—Peruv.
rigescens, Trin. ex Boiss. Fl. Orient. v. 685 = rigida.
rigida, Gaus. ex Steud. Nom. ed. II. i. 632 = Brachypodium distachyum.
rigida, *Kunth, Rév. Gram.* i. 129.—Europ. austr.; Afr. bor.
rigida, Roth, Catalect. fasc. i. 12 = Triticum asperum.
rigida-aspera, Steud. Syn. Pl. Gram. 317 = prasc.
rigidula, *Steud. Pl. Schimp. Abyss.* n. 814.—Abyssin.
rigurcos, Schur, Enum. Pl. Transs. 790 = ovina.
rivularis, Bolss. Elench. 90 = rubra.
robusta, *Kit. in Linnaea*, xxix. (1857–58) 98.—Chili.
Rochelii, Kit. l. c. xxxii. (1863) 218 = laevis.
Rochelii, *Steud. Nom.* ed. II. i. 632.—Banatus.
Rohlfsiana, *Cass. in Bull. Soc. Bot. Fr.* xix (1872) 83.—Afr. bor.
Rottboelii, DC. ex Steud. Nom. ed. II. i. 632 = Desmazeria loliacea.
rottboellioides, Kunth, Rév. Gram. i. 129 = Desmazeria loliacea.
rubens, Pers. Syn. i. 94 = Bromus rubens.
rubra, *Linn. Sp. Pl.* 74.—Reg. bor. temp. et arct.
rubra, With. Arr. Brit. Pl. ed. III. ii. 168 = ovina.
rupestris, *Otto, ex Sweet, Hort. Brit.* ed. II. 573, nomen.—Hab.?
rupestris, Roth, Enum. Pl. Phaen. Germ. i. 1. 311 = Brachypodium pinnatum.
rupicola, Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 223 = ovina.
sabulicola, Dufour, in Ann. Sc. Nat. Sér. I. v. (1825) 55 = rubra.
Salzmanni, *Boiss. ex Cosn. Notes Crit.* 181.—Hispan.
sancta, Janka, in Oestr. Bot. Zeitschr. xxi. (1871) 250 = Brachypodium sanctum.
Savii, Tenore, Fl. Nap. iv. 16 = F. Alopecurus.
saxatilis, Schur, Enum. Pl. Transs. 791 = ovina.
scaberrima, *Lange, in Kjoeb. Vidensk. Meddel.* (1860) 51.—Hispan.
scaberrima, *Steud. Syn. Pl. Gram.* 316.—Ind. or.
scabra, Labill. Nov. Holl. Pl. i. 22. t. 26 = Agropyrum scabrum.
scabra, Leg. Gen. et Sp. Nov. 4 = Brachypodium mexicanum.
scabra, *Vahl, Symb. Bot.* ii. 21.—Afr. austr.
scabra, Willd. ex Steud. Nom. ed. II. i. 632 = procera.
scabrella, *Torr. ex Hook. Fl. Bor. Am.* ii. 252.—Am. bor. occ.
scabriflora, *Steud. Syn. Pl. Gram.* 314.—Ins. Canar.

FESTUCA:—
scabriuscula, *Phil. in Linnaea*, xxix. (1857–58) 98.—Chili.
Schmidtkorii, Gaud. in Alpina, iii. (1868) 70 = pulchella.
schoenoides[†]*formis*, Schur, Enum. Pl. Transs. 796 = pulchella.
Schimperi, Steud. Syn. Pl. Gram. 316 = Triticum Schimperi.
Schimperiana, *A. Rich. Tent. Fl. Abyss.* ii. 488.—Abyssin.
Schkiehamii × , Grants. Fl. Ucherm. 240.—Europ.
sciaphila, Schur, Enum. Pl. Transs. 787 = ovina.
scirpifolia, *Kunth, Enum. Pl.* i. 408.—Pervv.
scirpoides, *F. Muell. Fragm.* viii. 129.—Austral.
sciuroa, Nutt. in Trans. Am. Phil. Soc. v. (1837) 147 = F. Myuros.
sciuroides, Roth, Bot. Abh. 43 = F. Myuros.
sclerophylla, Boiss. ex Bioch. Del. Sem. Hort. Heidelb. (1849) 4; Diagn. Ser. I. xiii. 59 = spectabilis.
scoparia, *Hook. f. Fl. Antarct.* 98.—N. Zel.; Ins. Antarct.
scoparia, Kern. ex Nym. Consp. 826 = varia.
segetum, Savi, Fl. Pis. i. 116 = Trisetum parviflorum.
septiflora, [Honck.] Vers. Aller Gew. Testischl. 268 = rubra.
serotina, Linn. Sp. Pl. ed. II. 111 = Diplachne serotina.
serrata, Spreng. Syst. i. 255 = Lasiochloa serrata.
setacea, Gmel. Fl. Sic. Syn. i. 88 = Thomasiana.
setacea, Poir. Encyc. Suppl. ii. 688 = tenella.
setifolia, *Steud. in Lechl. Bereb. Am. Austr.* 56.—Peruv.
sibirica, *Hack. ex Boiss. Fl. Orient.* v. 698.—Reg Caucas.; As. bor.
sicula, *Presl, Cyp. et Gram. Sic.* 86.—Sicili.; Sard.
sicula, Spreng. ex Steud. Nom. ed. II. i. 632 = clavata.
Sieberi, Tausch, in Flora, xx. (1837) 127 = spectabilis.
simensis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 486.—Abyssin.
simplex, Boiss. & Bal. Diagn. Ser. II. iv. 158 = elatior.
simplex, Homem. ex Steud. Nom. ed. II. i. 632 = incrassata.
siracusana, Jan, ex Steud. l. c. = ovina.
Smithii, *Kunth, Rév. Gram.* i. 133.—Ins. Canar.
spadicea, Gmelin. Fl. Scepus. n. 90; ex Roem. & Schult. Syst. ii. 719 = varia.
spadicea, *Linn. Syst. ed.* XII. 789.—Europ. austr.; Reg. Himal.
spadicea, Mielich. in Hoppe, Taschenb. (1801) 184 = pulchella.
spadicea, Moench, Meth. 190 = elatior.
spectans, Schreb. Spicil. Fl. Lips. 59 = Bromus inermis.
spectabilis, Gaud. & Godr. Fl. Fr. iii. 579 = spadicea.
spectabilis, Guss. ex Hack. Monog. Fest. 191 = calabrica.
spectabilis, *Vaw, Rlench.* 2.—Europ. austr.; Persia.
spectabilis, Schur, Enum. Pl. Transs. 796 = dimorpha.
Sphenopus, Trin. ex C. A. Mey. Vers. Pfl. Cauc. 22 = Sphenopus Gouani.
spicata, *Pursh, Fl. Am. Sept.* i. 58.—Am. bor.
spiciformis, Dulac, Fl. Hautes-Pyr. 94 = ovina.
spinosa, Linn. f. Suppl. 111 = Exagrostis spinosa.
splendens, Pourr. in Mém. Acad. Toul. iii. (1788) 319 = Koeleria setacea.
stenantha, Nym. Consp. 826 = ovina.
stipoides, Desf. Fl. Atlant. i. 90 = clavaza.
stolonifera, Mikger. in Ball. Soc. Bot. Fr. x. (1863) 87 = rubra.
stricta, Host, Gram. Austr. ii. 26. t. 36 = ovina.
stricta, Poir. Encyc. Suppl. ii. 686 = Bromus strictus.
stricta, Schleich. ex Gaud. in Alpina, iii. (1868) 59 = rubra.
strictifolia, Opiz, Natural. n. IX. (1825) 136 = ovina.
Stuartiana, *Steud. Syn. Pl. Gram.* 804.—Ins. Tasman. subcilista, *Steud. l. c.*—Graecia; Arabia.
subravescens, Schur, Enum. Pl. Transs. 785 = rubra.
subtilis, *Steud. Syn. Pl. Gram.* 317.—Hab.?
subulata, Bourg. Vég. Sticks, 55 = Melica acuminata.
subulifolia, *Benth. Pl. Hartw.* 263.—N. Granat.
sudetica, Tausch, ex Steud. Nom. ed. II. i. 633 = varia.
sulcata, Hack. in Bot. Centralb. viii. (1881) 405 = ovina.
supina, Schur, Enum. Pl. Transs. 784 = ovina.
sylvatica, Host, Gram. Austr. ii. 26. t. 36 = montana.
sylvatica, Huds. Fl. Angl. ed. I. 36 = Brachypodium sylvaticum.

FESTUCA:—
sylvatica, *Vill. Hist. Pl. Dauph.* ii. 106.—Europ.; As. Min.
syrtica, F. Muell. Fragm. viii. 130 = Glyceria stricta.
taurica, Kern. ex Hack. Monog. Fest. 104 = ovina.
Tauschii, *Steud. Syn. Pl. Gram.* 317.—Hab.†
tenax, Host, ex Link, Enum. Hort. Berol. i. 87 = elatior.
tenella, Chaub. ex Steud. Nom. ed. II. i. 633 = F. Poa.
tenella, *Mœnch, Meth. Suppl.* 68.—Hab.†
tenella, *Willd. Sp. Pl.* i. 419.—Am. bor.
Teneriffae, *Roth, Nov. Pl. Sp.* 76.—Ins. Teneriff.
Tenoriana, Steud. Syn. Pl. Gram. 317 = Triticum Barrelieri.
tenuicula, Link, Hort. Berol. i. 146 = F. Poa.
tenuicula, Nym. Syll. 418 = Avellinia tenuicula.
tenuiflora, *Schrad. Fl. Germ.* i. 245.—Reg. Mediterr.
tenuifolia, Sibth. Fl. Oxon. 44 = ovina.
tenuis, *Kitt. Taschenb* ed. II. 100.—Europ.
tenuis, *Nym. Syll.* 418; *Steud. Syn. Pl. Gram.* 305.—Italia.
tertia, *Steud. l. c.* 316.—Japon.
texana, *Steud. l. c.* 310.—Texas.
thalassica, Kunth, Rév. Gram. i. 129 = Glyceria maritima.
thermarum, *Phil. in Anal. Univ. Chil.* ii. (1862) 885.—Chili.
Thomasiana, *J. Gay, ex Mutel, Fl. Fr.* iv. 92.—Reg. Mediterr.
Thonini, *Steud. Syn. Pl. Gram.* 311.—Ind. occ.
Thunbergii, *Kunth, Rév. Gram.* i. 128.—Japon.
Tharberi, in Roths. Rep. Bot. U. St. Surv. vi. 292 = scabrella.
Tilingii, *Regel & Tiling, in Nouv. Mem. Soc. Nat. Mosc.* ii. (1859) 295.—As. or.
tolucensis, *H. B. & K. Nov. Gen. et Sp.* i. 158.—Mexic.
tolucensis, Rupr. in Bull. Acad. Brux. ix. ii. (1842) 286 = hephaestophila.
transsilvanica, Schur, Enum. Pl. Transs. 795 = rubra.
trichodes, Schult. Mant. ii. 650 = Eragrostis maypurensis.
trichophylla, Ducros, ex Gaud. Fl. Helv. i. 388 = rubra.
trifora, *Desf. Fl. Atlant.* i. 67.—Reg. Mediterr. occ.
trifora, Sm. Engl. Bot. t. 1918 = gigantea.
triloba, D. Dietr. Syn. Fl. i. 377 = trifolia.
triticea, Kunth, Enum. Pl. i. 325, sphalm. = scp.
triticoides, Lam. Tabl. Encyc. i. 191 = Distichlis maritima.
triticoides, Steud. Syn. Pl. Gram. 315 = littoralis.
tubervulata, Benth. in Journ. Linn. Soc. xix. (1881) 128, sphalm. = scp.
tuberculosa, *Coss. & Dur. Expl. Scient. Alg.* 189. t. 41.—Alger.
tunicata, *B. Desv. in C. Gay, Fl. Chil.* vi. 434.—Chili.
turfosa, Schur, Enum. Pl. Transs. 793 = rubra.
turkestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 273.—Turkestan.
Urchritsiana, Wiesb. in Oestr. Bot. Zeitschr. xxviii. (1878) 218 = elatior.
uligiuosa, Schur, Enum. Pl. Transs. 796 = gigantea.
ulochaeta, *Steud. Syn. Pl. Gram.* 305.—Boss.
umbrosa, *Kit. in Linnaea*, xxxii. (1863) 314, nomen.—Hungaria.
unifaria, *Dum. Obs. Gram. Belg.* 104.—Belgium.
uniglumis, Soland. in Ait. Hort. Kew. ed. I. i. 108 = bromoides.
unilateralis, Schrad. Cat. Hort. Gotting. (1814) = tenuiflora.
unioloides, Kunth, Rév. Gram. i. 129 = Demazeria sicula.
uniloides, Willd. Hort. Berol. i. 8. t. 8 = Bromus unioloides.
Uri, *Bishop, in Lond. Gard. Mag.* xiii. (1837) 258.—Hab.?
Urvilleana, *Steud. Syn. Pl. Gram.* 312.—Ins. Falkland.
vaginata, Rochel, Fl. Banat. Rar. Ind.=Rochelii.
vaginata, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 126 = ovina.
valentina, Balchb. ic. Fl. Germ. i. 27. t. 1247 = amethystina.
valentina, Schleich. ex Gaud. Agrost. Helv. i. 266 = ovina.
varia, *Haenke, in Jacq. Coll.* ii. 94.—Europ. mod.; As. Min.

FESTUCA:—
veraicolor, Presl, ex Kunth, Enum. Pl. i. 402 = rubra.
verricolor, Tausch, in Flora, iv. (1821) 559 = varia.
vestita, Walp. Ann. iii. 772 = dichotoma.
villiflora, *Steud. Syn. Pl. Gram.* 313.—Labrador.
villosa, Schweigg. in Hagen, Chlor. Borus. 85 = rubra.
villosa, Spreng. Syst. i. 255 = Lasiochloa villosa.
violacea, [Clairv.] Man. Herb. 24.—Europ.
violacea, Mert. & Koch, Fl. Deutschl. i. 546 = ovina.
violacea, Sm. ex Gaud. in Alpina, iii. (1868) 57 = rubra.
virgata, Heyne, ex Steud. Nom. ed. II. i. 633 = Cantothesa lapacea.
virgata, Lam. Tabl. Encyc. i. 189 = Leptochloa virgata.
viridis, Pena. ex Schult. Mant. ii. 397, in obs. 2 = rubra.
viscida, F. Muell. Veg. Chath. Isl. 50 = Triodia pungens.
vivipara, Sm. Brit. Fl. i. 114 = ovina.
vulpioides, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 89 = rubra.
vulpioides, *Steud. Syn. Pl. Gram.* 305.—Afr. austr.
Wightiana, *Steud. l. c.* 316.—Ind. or.
Willdenowiana, *Schult. Mant.* ii. 650.—Mexic.
Willemetii, Savi, Cens. Bot. 53 = bromoides.
xanthina, Roem. & Schult. Syst. ii. 721 = varia.

FESTUCARIA, Heist. Syst. 12 (1748) = **Festuca**, Linn. (Gramin.).
Heisteri, Fabr. ex Steud. Nom. ed. II. i. 633 = Glyceria fluitans.
paniculata, Link, in Linnaea, xvii. (1843) 393 = Festuca tenuiflora.
tenuicaulis, Link, l. c. = Festuca Poa.

FEUILLAEA, Gled. Syst. 368 (1764) = seq.

FEUILLEA, Gled. in Mém. Acad. Berl. v. (1749) 182 = seq.

FEUILLÉA, Linn. Gen. ed. I. 362 (1737). *CUCURBITACEAE*, Benth. & Hook. f. i. 840.
HYPANTHERA, Silva Manso, Cathart. Bras. 37 (1836).
NHANDIROBA, Plum. ex Adans. Fam. ii. 139 (1736).
alibi*cifolia*, *Cogn. in Mart. Fl. Bras.* vi. iv. 118.—Bras.
cordifolia, *Linn. Sp. Pl.* 1018.—Ind. occ.
cordifolia, Vell. Fl. Flum. x. t. 103 = trilobata.
deltoides, *Cogn. in Mart. Fl. Bras.* vi. iv. 119.—Bras.
hederacea, Poir. in Lam. Encyc. iv. 419 = cordifolia.
Tuvilla, H. B. & K. Nov. Gen. et Sp. ii. 124 = cordifolia.
Karsteni, *Cogn. in DC. Monog. Phan.* iii. 943.—Venezuela.
Marcgravii, Guib. Hist. Nat. Drog. Simpl. ed. IV. iii. 244 = trilobata.
monosperma, Vell. Fl. Flum. x. t. 108 = Sicydium monospermum.
Moorei, *Hook. f. Bot. Mag.* t. 6856.—Guiana.
Poaefolia, Vell. Fl. Flum. x. t. 104 = Anisosperma Passiflora.
podata, Sm. ex Sims, Bot. Mag. t. 2061 = Telfairia podata.
pseudata, Poir. in Lam. Encyc. iv. 418 = cordifolia.
scandens, Linn. Sp. Pl. ed. II. 1457 = cordifolia, trilobata.
tamnifolia, H. B. & K. Nov. Gen. et Sp. vii. 176. t. 646 = Sicydium tamnifolium.
trichosperma, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 566 = trilobata.
trilobata, Mart. ex Roem. Syn. Pepon. 114 = trilobata.
trilobata, *Linn. Sp. Pl.* 1014.—Am. trop.
trilobata, Reich. Syst. iv. 238 = cordifolia.
triphylla, Griseb. in Goett. Abh. xxiv. (1879) 126 = Alsodeia bracteosa.

FIALARIS, Rafin. Sylva Tellur. 136 (1838) = **Hyrtuna**, Link.
semibiflora, Rafin. l. c. = M. arundinacea.

FIBICHIA, Kosl: Gram. Gen. et Cens. 206 (1841) = Cynodon, Pers. (1806) (Gramin.).
semibiflora, Kosl. l. c. = C. Dactylon.

FIBIGIA, Medic. Pflanzeng. i. 90 (1792) = **Farsetia**, Turra (Crucif.).
clypeata, Medic. l. c.; Moench, Meth. 261 = Farsetia clypeata.
eriocarpa, Boiss. Fl. Orient. i. 258 = Farsetia eriocarpa.
lunarioides, Sibth. & Sm. Fl. Graec. vi. 695 = Farsetia lunarioides.
macrocarpa, Boiss. Fl. Orient. i. 258 = Farsetia macrocarpa.
macroptera, Boiss. l. c. 257 = Farsetia macrocarpa.
multicaulis, Boiss. & Hohen. ex Boiss. l. c. 259 = Farsetia multicaulis.
obovata, Boiss. l. c. 258 = Farsetia obovata.
pendula, Boiss. l. c. 260 = Farsetia pendula.
rostrata, Boiss. l. c. 257 = Farsetia rostrata.
suffruticosa, Sweet, Hort. Brit. ed. I. 467 = Farsetia suffruticosa.
umbellata, Boiss. Fl. Orient. i. 259 = Farsetia umbellata.

FIBRAUREA, Lour. Fl. Cochinch. 626 (1790). *MENISPERMACEAE*, Benth. & Hook. f. i. 29, 960.
chloroleuca, Miers, Contrib. iii. 43 = tinctoria.
fasciculata, Miers, l. c. 41 = tinctoria.
Haematocarpus, Hook. f. & Thoms. Fl. Ind. i. 204 = Haematocarpus Thomsoni.
laxa, *Miers, Contrib.* iii. 43.—Borneo.
recisa, *Pierre, Fl. For. Cochinch.* t. 111.—Cochinch.
tinctoria, *Lour. Fl. Cochinch.* 626.—Malaya.
tinosporoides, F. Muell. Fragm. x. 93 = Fawcettia tinosporoides.

FICARIA, [Dill.] Hall. Enum. Stirp. Helv. i. 321 (1742) = **Ranunculus**, Linn.
ambigua, Bor. Fl. Cent. Fr. ed. III. ii. 20 = R. Ficaria.
aperta, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 91 = R. Ficaria.
Beryi, Heldr. ex Nym. Consp. 7 = *peloponnesiaca*.
calthaefolia, Reichb. Fl. Germ. Excurs. 718 = R. Ficaria.
communis, Dum.-Cours. Bot. Calt. ed. II. iv. 445 = R. Ficaria.
fasciularis, C. Koch, in Linnaea, xv. (1841) 249 = R. edulis.
glaciatis, Flach. ex DC. Prod. i. 44 = Oxygraphis glacialis.
grandiflora, Robert, Cat. Toulon, 57, 112 = R. Ficaria.
Halleyi, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 93 = R. Ficaria.
intermedia, Schur, Enum. Pl. Transs. 14 = R. Ficaria.
nudicaulis, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 186 = R. Ficaria.
peloponnesiaca, Nym. Syll. 179.—Graecia.
polypetala, Gilib. Fl. Lituan. ii. 259 = R. Ficaria.
ranunculoides, Moench, Meth. 215 = K. Ficaria.
Roberti, F. Schultz, Arch. Fl. 346 = R. Ficaria.
rotundifolia, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 93 = R. Ficaria.
transsilvanica, Schur, Enum. Pl. Transs. 14 = R. Ficaria.
verna, Huds. Fl. Angl. ed. I. 214 = R. Ficaria.

FICHTEA, Sch. Bip. in Linnaea, x. (1835) 255 = **Microseris**, D. Don (Compos.).
Poeppigii, Sch. Bip. l. c.—Chili.

FICINDICA, St. Leg. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = **Cyuntia**, Tourn. (Cact.).

FICINIA, Schrad. in Comm. Gotting. vii. (1822) 143; (Anal. Fl. Cap. 43. t. 2. f. 3). *CYPERACEAE*, Benth. & Hook. f. iii. 1063.
Acrolepis, Schrad. in Comm. Gotting. vii. (1822) 143; (Anal. Fl. Cap. 42. t. 2. f. 6).
Chamaexiphium, Hochst. in Flora, xxvii. (1844) 108.
Pityolepis, Beauv. in Lestib. Essai Fam. Cyp. 33 (1819).
Sypophalium, Nees, in Linnaea, vii. (1832) 530.
Melancharis, Vahl, Enum. ii. 339 (1806).
Pleurachne, Schrad. in Comm. Gotting. vii. (1822) 147. t. 4. f. 3; (Anal. Fl. Cap. 47).
Schoenidium, Nees, in Linnaea, ix. (1834) 291.
Sickenahnia, Nees, l. c. 292 (1834).
acuminata, Kunth, Enum. Pl. ii. 256 = elongata, Kunthiana.
acuminata, *Nees, in Linnaea*, ix. (1834) 292.—Afr. austr.
Kuthiana, *Nees, l. c.* x. (1836) 176.—Afr. austr.

FICINIA:—
ambigua, *Steud. Syn. Pl. Cyp.* 135.—N. Zel.
anceps, *Nees, in Linnaea*, x. (1836) 179.—Afr. austr.
anomala, *Steud. Syn. Pl. Cyp.* 135.—N. Zel.
antarctica, Nees, in Linnaea, ix. (1834) 292 = Scirpus diabolicus.
aphylla, Nees, in Endl. Atact. 3. t. 19 = lateralis.
argyropa, *Nees, in Linnaea*, x. (1836) 177.—Afr. austr.
atrata, Nees, l. c. 167 = setiformis.
Berglera, *Kunth, Enum. Pl.* ii. 254.—Afr. austr.
Boluseli, Boeck. in Engl. Bot. Jahrb. v. (1884) 506 = lithosperma.
brachiata, *Boeck. in Linnaea*, xxxvii. (1871–73) 88.—Afr. austr.
brevifolia, *Nees, l. c.* ix. (1834) 292.—Afr. austr.
bulbosa, *Nees, l. c.* vii. (1832) 91.—Afr. austr.
Capitellum, *Nees, l. c.* x. (1836) 168.—Afr. austr.
ciliata, *Boeck. in Bremen, Abh.* vii. (1880) 88.—Afr. austr.
clandestina, *Boeck. in Linnaea*, xxxvii. (1871–73) 82.—Abyssin.
communata, *Kunth, Enum. Pl.* ii. 256.—Afr. austr.
comparbergensis, Drège, ex Steud. Syn. Pl. Cyp. 134 = sec.
compasbergensis, *Drège, in E. Mey. Zwei Pfl. Docum.* 185.—Afr. austr.
compositella, *Nees, in Linnaea*, x. (1836)172.—Afr. austr.
confera, Nees, l. c. 161 = scaniosa.
contexta, *Nees, l. c.* 292.—Afr. austr.
Drogeana, Kunth, Enum. Pl. ii. 256 = laevis.
Eckloniana, *Nees, in Linnaea*, ix. (1834) 292.—Afr. austr.
elongata, *Boeck. l. c.* xxxvii. (1871–73) 85.—Abyssin.
fascicularis, *Nees, l. c.* x. (1836) 178.—Afr. austr.
fastigiata, *Nees, l. c.* 170.—Afr. austr.
filamentosa, Nees, l. c. ix. (1834) 292 = tenuifolia.
filiformis, Nees in Linnaea, x. (1836) 178, partim = stoloniera.
filiformis, *Schrad. Anal. Cap. Cyp.* 46.—Afr. austr.
gracilis, *Schrad. l. c.* 44.—Afr. austr.
guttata, *Endl. Prod. Fl. Ins. Norfolk.* 23.—Ins. Norfolk.
involucrata, *Nees, in Linnaea*, ix. (1834) 292; x. (1836) 174.—Afr. austr.
involuta, *Nees, ll. cc.*—Afr. austr.
isidolea, *Nees, ll. cc.* 293, 180.—Afr. austr.
Kunthiana, *Boeck. l. c.* xxxvii. (1871–73) 67.—Afr austr.
lacinulata, *Nees, l. c.* x. (1836) 177.—Afr. austr.
laevis, *Nees, l. c.* ix. (1834) 292; x. (1836) 171.—Afr. austr.
lateralis, *Kunth, Enum. Pl.* ii. 254.—Afr. austr.
leiocarpa, *Nees, in Linnaea*, x. (1836) 171.—Afr. austr.
lithosperma, *Boeck. l. c.* xxxvii. (1871–73) 72.—Afr. austr.
Ludwigii, *Boeck. in Flora*, lxv. (1882) 15.—Afr. austr.
membranacea, *Kunth, Enum. Pl.* ii. 252.—Afr. austr.
monticola, *Kunth, l. c.* 261.—Afr. austr.
nuda, *Boeck. in Linnaea*, xxxvii. (1874) 383.—Afr. austr.
pallens, Nees, l. c. x. (1836) 169 = setiformis.
pallida, Nees, l. c. ix. (1834) 292 = setiformis.
paradoxa, *Nees, l. c.* x. (1836) 178.—Afr. austr.
paradoxa, Steud. Syn. Pl. Cyp. 135 = ambigua.
Poiretii, *Kunth, Enum. Pl.* ii. 255.—Afr. austr.
praecurva, *Nees, in Linnaea*, x. (1836) 167.—Afr. austr.
pravisia, Nees, l. c. ix. (1834) 292 = laevis.
pulchella, Kunth, Enum. Pl. ii. 261 = Scirpus pulchellus.
punctata, *Hochst. in Flora*, xxvii. (1844) 758.—Afr. austr.
pygmaea, *Boeck. in Linnaea*, xxxvii. (1871–73) 89.—Afr. austr.
quinquangularis, *Boeck. l. c.* 68.—Afr. austr.
radiata, *Kunth, Enum. Pl.* ii. 260.—Afr. austr.
ramosissima, *Kunth, l. c.* 262.—Afr. austr.
repens, *Kunth, l. c.* 258.—Afr. austr.
scariosa, *Nees, in Linnaea*, ix. (1834) 292; x. (1836) 182.—Afr. austr.
secunda, *Kunth, Enum. Pl.* ii. 262.—Afr. austr.
semibracteata, Nees, in Linnaea, ix. (1834) 292 = leiocarpa.
setiformis, *Schrad. Anal. Pl. Cap.* 45.—Afr. austr.
Shrubsoli, Nees, ex Kunth, Enum. Pl. ii. 256 = Eckloniana.
stoloniera, *Boeck. in Linnaea*, xxxvii. (1871–73) 90.—Afr. austr.
striata, Kunth, Enum. Pl. ii. 257 = setiformis, brevifolia.
sylvatica, *Kunth, l. c.* 254.—Afr. austr.

FICINIA:—
tenuifolia, *Kunth, Enum. Pl.* ii. 257.—Afr. austr.
tribracteata, *Boeck. in Linnaea*, xxxvii. (1871–73) 62.—Afr. austr.
trichodes, *Benth. & Hook. f. Gen.* iii. 1068.—Afr. austr.
tristachya, *Nees, in Linnaea*, ix. (1834) 292; x. (1836) 176.—Afr. austr.
truncata, *Schrad. Anal. Cap. Cyp.* 43. t. 2. f. 3.—Afr. austr.
Zeyheri, *Boeck. in Linnaea*, xxxvii. (1871–73) 58.—Afr. austr.

FICOIDES, [Tourn.] Rupp. Fl. Jen. ed. Hall. 164 (1748) = **Mesembryanthemum**, Linn. (Ficoid.).

FICULA, Heist. Syst. 8 (1748) = **Mesembryanthemum**, Linn. (Ficoid.).

FICUS, Tourn. ex Linn. Syst. ed. I (1735). *URTICACEAE*, Benth. & Hook. f. iii. 367, 1924.
Caprificus, Gasp. Nov. Gen. Fic. 6 (1844).
Covellia, Gasp. l. c. 10 (1844).
Cystogyne, Gasp. in Ann. Sc. Nat. Sér. III. iii. (1845) 345.
Eaoma, Booth, Cat. 113; ex Endl. Suppl. iv. ii. 64, in syn. (1849).
Krythrogyne, Vis. in Gasp. Ricerch. Caprif. 86 (1845).
Galoglychia, Gasp. Nov. Gen. Fic. 10 (1844).
Gonosyce, Rafin. Sylva Tellur. 58 (1838).
Macrophthalma, Gasp. Ricerch. Caprif. 88. t. 7 (1845).
Mastosyce, Necalistis, Oluntos, Perula, Rafin. Sylva Tellur. 58, 59 (1838).
Pharmacosycea, Miq. in Hook. Lond. Journ. Bot. vi. (1848) 64.
Plagiostigma, Zucc. in Abh. Akad. Muench. iv. i. (1845) 154.
Pogonotrophe, Miq. in Hook. Lond. Journ. Bot. vi. (1848) 72.
Rephesis, Rafin. Sylva Tellur. 59 (1838).
Sycomorphe, Miq. in Ann. Sc. Nat. Sér. III. i. (1844) 85.
Sycomorus, Gasp. Ricerch. Caprif. 86 (1845).
Synoecia, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 109.
Tengera, Gasp. Nov. Gen. Fic. 6 (1844).
Tremotis, Rafin. Sylva Tellur. 59 (1838).
Urostigma, Gasp. Nov. Gen. Fic. 7 (1844).
Varinga, Rafin. Sylva Tellur. 58 (1838).
Visiania, Gasp. Nov. Gen. Fic. 9 (1844).
abbreviata, *Wall. Catal.* n. 4573.—Singapore.
Abelii, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 281 = pyriformis.
abutilifolia, *Miq. l. c.* 328.—Abyss.; Afr. austr.
acamptophylla, *Miq. l. c.* 264.—Malaya.
accedens, *Steud. ex Miq. in Hook. Lond. Journ. Bot.* vi. (1847) 567, in syn.—Abyssin.
aculeata, *A. Cunn. ex Miq. l. c.* vii. (1848) 496.—Austral.
acuminata, Bech.-Ham. in Wall. Cat. n. 4478 = parietalis.
acuminata, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 21 = rostrata.
acuminata, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 538 = subulata.
acuminatissima, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 328.—pinifera.
acutilobe, Miq. l. c. 227. t. 5 a = heterophylla.
adenosperma, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 288.—Ins. Celebes.
adhaerens, Miq. Fl. Jungh. i. 55 = ramentacea.
adhatodaefolia, *Schott, ex Spreng. Syst. iv. App.* 409.—Bras.
adnascens, Wall. Cat. n. 4572 B = recurva.
aegrophylla, Hort. Schoenb. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 287 = Amottiana?
affinis, Griff. Notul. iv. 302; Ic. Pl. As. t. 553 = religiosa.
affinis, Blume, ex Steud. Nom. ed. II. i. 635 = Blumei.
affinis, Wall. Cat. n. 4594 = glabella.
Afzelii, *G. Don, in Lond. Hort. Brit.* 416, nomen.—Afr. occ.
Ayeelii, Hort. Berol. ex Kunth & Bouché, Ind. Sem Hort. Berol. (1848) 14 = eriobotryoides.

FICUS:—
africana, Hort. Berol. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 289 = ovata.
aggregata, Hort. ex Miq. l. c. 296 = arbutifolia.
aggregata, *Vahl, Enum.* ii. 191.—Ind. or.
alba, *Reinw. ex Blume, Bijdr.* 476.—Malaya.
albescens, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 228.—Reg. Mediterr.
albinervia, *Miq. Fl. Ind. Bat.* i. ii. 315.—Ins. Bonin.
Allmeeralso, Wight, Ic. t. 950, sphalm. = gibbosa.
alnifolia, Miq. Fl. Jungh. i. 61 = obtusa.
alpestris, Walp. Ann. i. 696, sphalm. = *sylvestris*.
alternans, *Wall. Cat.* n. 4555.—Siam.
Altimeralso, Kunth ex Miq. in Hook. Lond. Journ. Bot. vii. (1848) 225 = gibbosa.
altissima, *Blume, Bijdr.* 455.—Reg. Himal.; As. trop.; Malaya.
altuacea, *Blume, Bijdr.* 457 (altuaceo).—Java.
amara, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 15 = hispida.
amazonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Bras.
ambiphylla, Miq. l. c. 296 = retusa.
ambiguensis, Kostel. Allg. Med.-Pharm. Fl. ii. 406 = variegata.
americana, *Aubl. Pl. Gui.* ii. 952.—Guian.
Amhrettiana, Steud. Nom. ed. II. i. 626 = Erythroxylum burmanicum.
Ampelos, *Burm. f. Fl. Ind.* 226.—Ind. or.
Ampelos, Kom. ex Roxb. Hort. Beng. [108]; Fl. Ind. iii. 558 = gibbosa.
ampla, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 18 = infectoria.
amplissima, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 297, sphalm. = F. Arpazusa.
amplissima, Sm. in Rees, Cyel. xiv. n. 1 = F. Taiela.
ampullacea, Wight, ex Miq. in Hook. Lond. Journ. Bot. vii. (1848) 229 = quercifolia.
anabates, Voigt, Hort. Suburb. Calc. 286 = scandens.
anacardiifolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 18.—Hab.?
anadenocarpa, Wall. Cat. n. 4513 = quercifolia.
anculane, Miq. Pl. Jungh. i. 62 = subulata.
angulata, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 234 = gibbosa.
angulidens, Miq. Fl. Ind. Bat. i. ii. 310 = rostrata.
angustata, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 234.—Ind. or.
angustifolia, Blume, Bijdr. 462 = cuspidata.
angustifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Guian.
angustifolia, Roxb. Hort. Beng. [108]; Fl. Ind. iii. 554 = glaberrima.
anoulata, *Blume, Bijdr.* 448.—Burma; Malaya.
anonaefolia, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 272 = pisifera.
anthelmintica, *Mart. Syst. Mat. Med. Bras.* 88; *Reise*, iii. 1138.—Bras.
anthelmintica, Raensch. Nom. ed. III. 302 = septica.
anthelminthica, Rich. in DC. Ess. Méd. Pl. 267 = F. Radula.
antithenophylla, *Steud. ex A. Rich. Tent. Fl. Abyss.* ii. 272.—Abyssin.
aperta, *Wall. Cat.* n. 4552.—Siam.
apiculata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Ind. or.
apiculata, Miq. in Zoll. Syst. Verz. Ind. Archip. 92, 98 = falva, *Reinw.*
apiocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 289.—Malaya.
apocynoeoides, *Grif. Notul.* iv. 309.—Reg. Himal.
apodocephala, *Baker, in Journ. Linn. Soc.* xxi. (1886) 445.—Madag.
aquatica, Koen. ex Willd. Sp. Pl. iv. 1138 = heterophylla.
arborícola, Schult. Mant. i. 321 = dendrícola.
arbutifolia, *Link, Enum. Hort. Berol.* ii. 450.—Bras.
arbutifolia, Pers. Syn. ii. 610 = pertusa.
argentea, Blanco, Fl. Filip. ed. 1. 681 = F. Ribes.
argentea, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 290.—Afr. trop.
argutis, Wall. Cat. n. 4469 = chrysocarpa.
Arnottiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 287.—Ind. or.
Arpazusa, *Casar. Nov. Stirp. Bras. Dec.* 15.—Bras.
asperina, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 198 = tomentosa.

FICUS:—
aspera, Blanco, Fl. Filip. ed. 1. 676 = hirta.
aspera, *Forst. f. Pl. Esul.* 86.—Ins. N. Hebrid.
aspera, Hort. ex Steud. Nom. ed. II. i. 625 = exasperata.
asperifolia, Hook. ex Miq. in Hook. Lond. Journ. Bot. vii. (1848) 237 = pisifera.
asperifolia, *Miq. l. c.* 231.—Afr. trop.
asperior, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 291.—Ins. Amboin.
asperivaria, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 21 = obscura.
asperrima, *Roxb. Hort. Beng.* [108]; *Fl. Ind.* iii. 554.—Ind. or.
asperrima, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 402.—Java.
aspertula, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 361.—N. Caled.
assamica, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 226 = heterophylla.
atahbyhlla, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 227 = pedunculosa.
atrovirens, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 19 = tristis.
atrovirens, Schott, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 300 = oblongata.
atrox, *Mart. ex Miq. in Hook. Lond. Journ. Bot.* vi. (1847) 556, in syn.—Bras.
Attineeralso, Roxb. ex Wight, Ic. ii. text. 650 = gibbosa.
aurantiaca, *Grif. Notul.* iv. 304; *Ic. Pl. As.* t. 554. f. 11.—Malacca.
aurantiaca, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 17 = sapotaefolia.
aurantiacea, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 292 = glandulifera.
aurantiaca, Noronha, in Verh. Bat. Gen. v. (1790) ed. 1. Art. iv. 15 = obscura.
aurantiaca, Wall. Cat. n. 4465 = glaberrima.
aurata, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 291 = chrysocarpa.
aurea, *Nutt. Sylva*, ii. 4; ed. II. i. 154.—Florida.
auriculata, Lour. Fl. Cochinch. ii. 666 = F. Cunia.
aurita, *Blume, Bijdr.* 462.—Ins. Amboin.
australis, Willd. Sp. Pl. iv. 1138 = rubiginosa.
austro-caledonica, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 367.—N. Caled.
Avi-Avi, *Blume, Bijdr.* 446.—Java.
Avi-Avi, *Chapelier, ex Bot. Hort. Maurit.* 291.—Madag.
Ayresii, *Baker, Fl. Maurit.* 284.—Ins. Maurit.
baccata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 299.—Mexic.
Backhousii, *Miq. l. c.* 288.—Austral.
Balansaeana, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 363.—N. Caled.
balica, *Miq. Fl. Ind. Bat.* i. ii. 314.—Malaya.
bambaseoides, *Seem. Fl. Vit.* 250.—Ins. Fiji.
bandana, Miq. Fl. Ind. Bat. i. ii. 301 = F. Ampelos.
barbata, Wall. Cat. n. 4576 = villosa.
Barclayana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 296.—Ins. Fiji.
Barclayi, Seem. Fl. Vit. 250 = praec.
Baroni, *Baker, in Journ. Linn. Soc.* xx. (1883) 262.—Madag.
basidentata, *Miq. Fl. Ind. Bat.* i. ii. 314.—Archip. Malay.
Bechieri, Miq. in Journ. Bot. Néerl. (1861) 241 = orbicularis.
Beechyeana, Hook. & Arn. Bot. Beech. Voy. 371 = erecta.
benghalensis, *Linn. Sp. Pl.* 1059.—Ind. or.; Afr. trop.
Benjamina, *Linn. Mant.* 126.—As. trop.; Malaya.
Benjamina, Wall. Cat. n. 4508 = F. Taiela.
Benjamina, Willd. Sp. Pl. iv. 1148 = retusa.
Benjaminoides, Sabine, ex Miq. in Mart. Fl. Bras. iv. i. 101 = Salmasiana.
Bennettii, *Seem. Fl. Vit.* 250.—Ins. Fiji.
bibracteata, *Hort. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 295.—Hab.?
bicolor, Hook. ex Miq. in Hook. Lond. Journ. Bot. vii. (1848) 235 = alba.
bijendadeta, Blume, Bijdr. 475 = quercifolia.
bijendadeus, Steud. Nom. ed. II. i. 625 = quercifolia.
bijendadeus, Wall. Cat. n. 4460 = disticha.
biasia, Wall. l. c. n. 4564 = nemoralis.
Binnendijkii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Java.

FICUS:—
bistipulata, Gaill. Notul. iv. 298 = glaberrima.
blanda, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Hab.?
Blumei, *Steud. Nom.* ed. II. i. 625 (sp. dub.).—Java.
Boiei, Zipp. in Miq. Ann. Mus. Bot. Lugd. Bat. iii. 275 = subulata.
Bojeri, *Baker, Fl. Maurit.* 286.—Ins. Seychelles.
Bonplandiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 294.—Mexic.
Boonlusana, *Miq. l. c.* 290.—Bras.
botryspicloides, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 15.—Mexic.
botryocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 283.—Ins. Celebes.
brachyclada, *Baker, in Journ. Linn. Soc.* xx. (1883) 259.—Madag.
brachypoda, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 287 = cagenioides.
bracteata, *Wall. Cat.* n. 4496.—Malaya.
brasiliensis, Cels, ex Desf. Cat. Hort. Par. ed. III. 412 = Principa.
brasiliensis, Hort. Mon. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 298 = longifolia.
brasiliensis, *Link, Enum. Hort. Berol.* ii. 449.—Bras
Brasili, *R. Br. ex Sabine, in Trans. Hort. Soc.* v. (1824) 448.—Afr. trop.
Brasili, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 14 = Principa.
breviscapis, *Miq. Fl. Ind. Bat.* i. ii. 315.—Java.
brevifolia, *Nutt. Sylva*, ii. 8; ed. II. i. 158.—Florida.
brevipes, Pl. Jungh. 59 = obscura.
bulbosa, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 17 = sapotaefolia.
bullata, Roxb. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 286 = nervosa.
Burkei, *Miq. l. c.* 289.—Afr. austr.
caffra, Miq. l. c. 288 = lutea.
callicarpa, Aust. = kallicarpa.
callisphylla, *Blume, Bijdr.* 445.—Java.
calloza, *Willd. in Mém. Acad. Berl.* (1798) 102.—As. trop.
calosecura, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 106.—Burma.
calycipherma, Miq. in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 15 = Rumphii.
calyculata, *Mill. Gard. Dict.* ed. VIII. n. 11.—Mexic.
calyptrata, *Vahl, Enum.* ii. 186.—Afr. trop.
calyptrocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Bras.
cannabina, *Lour. Fl. Cochinch.* ii. 668.—Cochinch.
capensis, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 19 = cordata.
capensis, *Thunb. Diss. Pic.* 18.—Afr. austr.
caprarioides, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 94 = antithetophylla.
Capellona, *Risso, Fl. Nice*, 434; *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 296.—Europ.
caritose, *Yacq. Obs. Bot.* ii. 90.—Ind. occ.
Carica, *Linn. Sp. Pl.* 1059.—Europ.; Orient; Afr. bor. et austr.
caricoides, Roxb. Hort. Beng. 65; Fl. Ind. iii. 529 = palmata.
carophylla, Hort. Leyden. ex Miq. in Hook. Lond. Journ. Bot. vii. (1847) 493 = radicans.
casapiensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 296.—Peruv.
cassaefa, *P. Mantl. ex Benth. Fl. Austral.* vi. 177.—Austral.
castaneaefolia, *Roth, Nov. Pl. Sp.* 369.—Ind. or.
castaneaefolia, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 280 = ramentacea.
catappaefolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 14.—Ins. Martinic.
catarctarum, *Vieill. ex Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 365.—N. Caled.
Catesbaei, *Steud. Nom.* ed. II. i. 625.—Ind. occ.
caudata, *Grif. Notul.* iv. 302.—Assam.
caudata, Rohan, Bot. Mat. Med. iv. 256 = religiosa.
caudata, Wall. Cat. n. 4469 = clavata.
caulicinata, *Teijsm. in Journ. Bot.* xxiii. (1885) 243.—Zeylan.
caulobotrya, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 287.—Ind. or.
cauniocarpa, Miq. l. c. 296 = F. Tsiela.
Cawi, Hasse, ex Baker, in Journ. Linn. Soc. xx. (1883) 271 = Sycomorus.

FICUS :—
celebica, *Blume, Bijdr.* 471.—Malaya.
cerasifolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Hab. ?
cerasiformis, Desf. Cat. Hort. Par. ed. III. 413 = parietalis.
cerifera, Blume, in Ann. Sc. Nat. Sér. IV. iv. (1855) 333 = variegata.
cerifera, Jungh. Fl. Jav. i. 489 = variegata.
cerrifolia, *Schott, ex Spreng. Syst.* iv. *Add.* 409.—Bras.
ceylanica, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 208 = laevis.
Championi, Benth. in Hook. Kew Journ. vi. (1854) 76 = vasculosa.
Chanse, *Forsk. Fl. Aegypt. Arab.* 219.—Arabia.
chartacea, *Wall. Cat.* n. 4580.—Burma ; Malaya.
Chinchа, Roxb. Hort. Beng. [108] ; Fl. Ind. iii. 584 = clavata.
Chiriquiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 208.—Panama.
Chittagonga, Miq. l. c. 238 = glomerata.
chlorocarpa, Benth. Fl. Hongk. 330 = variegata.
chlorocarpa, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 234 = fulva, *Reinw.*
chloroleuca, Miq. Fl. Ind. Bat. I. ii. 294 = fulva, *Reinw.*
chrysocarpa, *Reinw. ex Blume, Bijdr.* 475.—Burma ; Malaya.
chrysocoma, Blume, Bijdr. 443 = pilosa.
chrysolepis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii 216.—Ins. Celebes.
chrysophthalma, *Miq. l. c.* 285.—Ind. or.
ciliolaea, Hort. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 17 ? = consanguinea.
ciliolaea, Hort. Berol. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 559 = terebrata ?
ciliolaea, *Link, Enum. Hort. Berol.* ii. 450.—Hab. ?
cinerascens, Thw. Enum. Fl. Zeyl. 266 = callosa.
cinerascens, *Wall. Cat.* n. 4535.—Burma.
citharexifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 232.—Africa ?
citrifolia, *Hort. ex Lam. Encyc.* ii. 494.—Mexic.
citrifolia, Mill. Gard. Dict. ed. VIII. n. 10 = Calebaci.
citrifolia, Willd. Sp. Pl. iv. 1137 = mysorensis.
clavigloeles, *Baker, in Journ. Linn. Soc.* xx. (1883) 266.—Madag.
clavata, *Wall. Cat.* n. 4495.—As. trop. ; Malaya.
clethraefolia, *Willd. Sp. Pl.* iv. 1143.—Venezuela.
cluniaefolia, Hort. ex Kunth & Bouché. Ind. Sem. Hort. Berol. (1846) 17 ? = consanguinea.
cluniaefolia, H. B. & K. ex Walp. Ann. i. 678 = obtusifolia ?
cluniaefolia, *Schott, in Spreng. Syst.* iv. *Add.* 409.—Bras.
cluniifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Ins. Philipp.
coccinella, Zipp. ex Miq. l. c. 280 = ramentacea.
coccolobifolia, *Ladd. ex Steud. Nom.* ed. II. i. 635 (Quid ?).—Ins. trop.
colombra, *Gasp. Ricrrch. Caprif.* 77.—Italia.
colosseа, *F. Muell. ex Benth. Fl. Austral.* vi. 163.—Austral.
columbarum, Hochst. ex Krauss, in Flora, xxviii. (1845) 88 = natalensis.
columnaris, *F. Muell. & C. Moore, in Proc. Acclim. Soc. Vict.* iii. (1874) 71.—Austral.
communis, Lam. Fl. Fr. iii. 502 = F. Carica.
comosa, Roxb. Pl. Corom. ii. 14. t. 125 = F. Benjamina.
conglobata, *H. B. & K. Nov. Gen. et Sp.* ii. 46.—Mexic.
compressicaulis, *Blume, Bijdr.* 439.—Java.
concentrica, Van Hass. in Miq. Choix Pl. Buitenz. t. 11 = parietalis.
conchlorum, Buch.-Ham. ex Wall. Cat. n. 4484 = Rumphii.
concinna, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 286 = glabella.
Condaravia, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 131 = retusa.
congesta, *Roxb. Hort. Beng.* 66 ; *Fl. Ind.* iii. 560.—Ins. Amboin.
conglomerata, Roxb. ll. cc. 66, 560 = F. Canis.
conjugata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 222.—Java.
connexiramea, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 17.—Mexic.
conocladia, *Baker, Fl. Maurit.* 286.—Ins. Mascar.
conocladus, *Blume, Bijdr.* 447.—Malaya.
Cooperi, *Hort. ex Regel, Ind. Sem. Hort. Petrop.* (1866) 86.—Am. trop.
copiosa, Steud. Nom. ed. II. i. 635 = F. Ribes.
cordata, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 14, 15 = elastica, syringaefolia.
cordata, *Thunb. Diss. Fic.* 8.—Afr. austr.
cordifolia, *Blume, Bijdr.* 438.—Java.
cordifolia, Roxb. Hort. Beng. 66 ; Fl. Ind. iii. 548 = Ramphii.
coriacea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 452.—Ind. or.
cornifolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 19.—Java.
coronata, Reinw. ex Blume, Bijdr. 470 = obscura.
coronata, Spin. ex Cella, Hort. Ripul. 56. t. 8 = exasperata, simplicifolia.
coronulata, *Miq. in Journ. Bot. Néerl.* (1861) 242.—Austral.
costaricensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 208.—Costa Rica.
costata, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 458.—Ins. Nicobar.
costigera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 206.—Ind. or.
cotinifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 49.—Mexic.
cotoneaefolia, Hort. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 299 = infectoria.
cotoneaefolia, Vahl, Enum. ii. 189 = mysorensis.
cotonifolia, Stokes, Bot. Mat. Med. iv. 355 = prasc.
courtillelensis, Baill. Hist. Pl. vi. 176 = Arnottiana.
coybana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 300.—Panama.
crassinervia, *Desf. Cat. Hort. Par.* ed. III. 418.—Jamaic.
crassinervia, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) = benghalensis.
crassiramea, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 286 = procera.
crenulata, *Hassk. Cat. Hort. Bog. Alt.* 76.—Java !
criniinervia, *Miq. Fl. Ind. Bat. Suppl.* 432.—Reg. Himal. ; Malaya.
crocata, *Mart. ex Miq. in Mart. Fl. Bras.* iv. i. 94, in syn.—Bras.
crustacea, Wall. Cat. n. 4501 = scandens.
Cumingii, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 235.—Philipp.
cuneo, Steud. Nom. ed. II. i. 635 = F. Canis.
cuneata, Blume, Bijdr. 468 = gibbosa.
cuneata, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 118.—Bras.
cuneata, *Wight, Ann. Mus. Bot. Lugd. Bat.* iii. 207.—Ins. Philipp.
cuneata, Wall. Cat. n. 4584 = Erythroxylum burmanicum.
cunefolia, Hook. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 297 = lepicarpa.
Cunia, *Buch.-Ham. ex Roxb. Hort. Beng.* 66 ; *Fl. Ind.* iii. 561.—As. trop.
Cunninghamii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Austral.
cupidatae, Desf. Cat. Hort. Par. ed. III. 413 = F. Benjamina.
cupidata, *Reinw. ex Blume, Bijdr.* 464.—Malaya.
cupidata, Zoll. & Mor. Syst. Verz. 77 = rostrata.
cupidata, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 234.—Ind. or.
cyclophylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Bras.
cyrtophylla, Wall. Cat. n. 4582 = obscura.
cystopoda, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Bras.
Daemona, [Koen.] Vahl, Enum. ii. 198 = bispida.
Daemonum, Zoll. & Mor. Syst. Verz. 77 = obscura.
Dahro, *Delile, in Ann. Sc. Nat.* Sér. II. xx. (1843) 94.—Abyssin.
Dalhousiae, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 285.—Ind. or.
daphniphylla, *Miq. l. c.* 300.—Bras.
deallbata, Hort. ex Hérincq, L'Horticr. Franç. (1869) 146 ; Linden, Illustr. Hortic. xvii. (1870) t. 4 = Cousapoa deallbata.
Decaisneana, Miq. Fl. Ind. Bat. I. ii. 312 = neq.
Decaisnei, *Steud. Nom.* ed. II. i. 636.—Ins. Timor.
decipiens, *Reinw. ex Blume, Bijdr.* 479.—Java.
Dekdekena, *A. Rich. Tent. Fl. Abyss.* ii. 268.—Abyssin.
delatorix, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 15 = alnifolia.
delicioso, *Gasp. Ricrrch. Caprif.* 78.—Italia.
deltoidea, Jack, in Malay. Misc. ii. (1822) vii. 71 = diversifolia.
dendrocida, *H. B. & K. Nov. Gen. et Sp.* ii. 46.—N. Granad.
dendrotrena, Spreng. Syst. iii. 780 = prasc.
dense, Miq. in Hook. Lond. Journ. Bot.-vii. (1848) 453 = nemoralis.
densifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 218.—Ins. Borbon.
densiverra, Miq. Fl. Ind. Bat. Suppl. 426 = chrysocarpa.
denticulata, Vahl, Symb. Bot. i. 53 = heterophylla.
denticulata, *Willd. Sp. Pl.* iv. 1139.—Sumatra.
depenes, *Benth. Fl. Austral.* vi. 172.—Austral.
depressa, Blume, Bijdr. 450 = pruniformis.
dialema, *Steud. Nom.* ed. II. i. 636.—China.
dicarpa, Blanco, Fl. Filip. ed. I. 682 = F. Benjamina.
dichrotrix, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 272, adn. = obscura.
dictyophleba, *F. Muell. in Benth. Fl. Austral.* vi. 170.—Austral.
dictyophylla, Wall. Cat. n. 4502 A, B, D = F. Benjamina.
difformis, Benth. Fl. Hongk. 337 = gibbosa.
difformis, *Lam. Encyc.* ii. 500.—Malaya.
dilatata, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 218 = retusa.
dimidiata, *Grisb. Fl. Brit. W. Ind.* 151.—Jamaic.
dimidiata, Wall. Cat. n. 4575 = sarmentacea.
diospyrifolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Hab. ?
diphylla, Wall. Cat. n. 4543 = fistulosa.
discolor, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 221.—Borneo.
dissecarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 268 = F. Dekdekena.
disticha, *Blume, Bijdr.* 458.—Java.
disticha, Thw. Enum. Fl. Zeyl. 266 = Thwaitesii.
diversifolia, *Blume, Bijdr.* 456.—Malaya.
diversifolia, Reinw. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 299 = rostrata.
diversiformis, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 441 = Thwaitesii.
Divoa, Miq. Choix Pl. Buitenz. t. 12 = villosa.
dodonaefolia, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 292 = gibbosa.
doliaria, *Mart. Syst. Mat. Med. Bras.* 88.—Bras.
domestica, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 282 = variegata.
dottata, *Gasp. Ricrrch. Caprif.* 77.—Italia.
drupacea, *Thunb. Diss. Fic.* 11.—Ind. or.
dubia, *Wall. Cat.* n. 4561.—Malaya.
dyctiophleba, *F. Muell. in Mig. Ann. Mus. Bot. Lugd. Bat.* iii. 218.—Austral.
ebenus, *Hort.—Cf. Gard. Chron.* (1869) 587.—Ind. or.
edulis, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 271.—N. Calcd.
ehretioides, *F. Muell. ex Benth. Fl. Austral.* vi. 171.—Austral.
elastica, *Roxb. Hort. Beng.* 65 ; *Fl. Ind.* iii. 541.—As. trop.
elegans, Hassk. Cat. Hort. Bog. Alt. 76 = toxicaria.
elegans, Hort. Berol. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 299 = sororia.
elegans, *Miq. l. c.* 288.—Afr. trop.
elegans, Steud. Nom. ed. II. i. 636 = foveolata.
ellipsoides, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 230 = Decaisnei.
ellipsoidea, F. Muell. ex Benth. Fl. Austral. vi. 164 = pilosa.
elliptica, Hook. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 582 = retusa ?
elliptica, *H. B. & K. Nov. Gen. et Sp.* ii. 46.—N. Granat.
elliptica, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 440 = dilatata.
elongata, Miq. l. c. 281 = heterophylla.
Emodi, Wall. Cat. n. 4515 = laevis.
enormis, *Mart. ex Miq. in Hook. Lond. Journ. Bot* vi. (1847) 544.—Bras.
erecta, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 294 = foveolata.
erecta, *Thunb. Diss. Fic.* 9.—Reg. Himal. ; China ; Japon.
eriobotrydies, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Hab. !
erythrosperma, *Miq. Ann. Mus. Bot Lugd. Bat.* iii. 226.—Ins. Celebes.
erythrostieta, *Miq. l. c.* 298.—Guian.

FICUS:—
cuspidifolia, *Hemsl. Biol. Centr. Am. Bot.* iii. 144.—Costa Rica.
cuspidoides, *F. Muell. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Austral
cuspidata, *Miq. l. c.* 299.—Bras.
cuspidyla, *Kurz, For. Fl. Brit. Burma*, ii. 445.—Burma; Malaya.
cuspidifolia, *Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 21.—Ins. Philipp.
cuspidata, Hort. Kew. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 291 = asperata.
exasperata, Roxb. Hort. Beng. 66; Fl. Ind. iii. 555 = pilifera.
exasperata, *Vahl, Enum.* iii. 197.—Afr. trop.
exotica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Java.
excelsa, Vahl, Enum. ii. 195 = gibbosa.
eximia, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 410.—Braz.
exculpta, [*Mast.*] *in Gard. Chron.* (1878) ii. p. viii (cf. 84).—Hab.?
Fadyeni, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 219.—Jamaic.
fagifolia, *Miq. l. c.* 299.—Bras.
fagines, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 20.—Am. bor.
falcata, Thunb. Diss. Fic. 8 = punctata.
fallax, Miq. Fl. Ind. Bat. i. 11. 306 = cuspidata.
fasciculata, *F. Muell. ex Benth. Fl. Austral.* vi. 177.—Austral.
fasciculosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 288.—Afr. trop.
fecunda, Blume, Cat. Gew. Buitenz. 36 = hispida.
ferruginea, *Descf. Cat. Hort. Par.* ed. III. 412.—Amer.
ferruginea, Hort. ex Miq. in Mart. Fl. Bras. iv. 2. 98 = rubiginosa.
Fieldingii, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 489 = nemoralis.
filiformis, *Blume, Bijdr.* 449.—Java.
Finlaysoniana, Wall. Cat. n. 4538 = pyriformis.
firma, Wall. l. c. n. 4564 A, B = globosa, Blume.
firmula, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 294 = Daemonat.
fistulosa, *Reinw. ex Blume, Bijdr.* 470.—Reg. Himal.; Malaya.
Fitalaisi, *Miq. Journ. Bot. Néerl.* i. (1861) 242.—Austral.
flavescens, Blume, Bijdr. 449 = annulata.
forcunda, Anct. = *fecunda*.
Fontanesii, *Spreng. Nom.* ed. II. i. 686.—Hab.?
formosana, *Maxim. in Bull. Acad. Pétersb.* 27 (1881) 546.—China.
Forskåhlei, *Vahl, Enum.* ii. 196.—Arab.
Forsteni, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 214.—Ins. Celebes.
foveolata, *Wall. Cat.* n. 4493.—Reg. Himal.; Burma; China.
foveolata, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 293 = sarmentacea.
fragilina, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 20.—Hab.?
Fraseri, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 283 = Cunninghamii.
fraterna, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 217 = glaberrima.
frigida, *Linden, ex Miq. l. c.* 291.—N. Granat.
fructicosa, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 533 = scandens.
fulva, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 14 = ferruginea.
fulva, *Reinw. ex Blume, Bijdr.* 473.—Burma; Malaya.
fulva, *Spreng. Syst.* iii. 779.—Bras.
fuscescens, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 296.—Mexic.
galactophora, *Tenore, in Atti Acc. Sc. Nap.* vi. (1851) 157; *et ex Gasp. Ricerch. Caprif.* 35, in syn.—Am. auct.
garcilasifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 218.—Ins. Timor.
gardneriaefolia, Hort. Berol. ex Miq. l. c. 299 = blanda.
gardneriaefolia, Hort. Lips. ex Miq. l. c. = wyrtifolia.
Gardneriana, *Miq. l. c.* 297.—Bras.
Gasparriniana, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 486.—Ind. or.

FICUS:—
Gelderii, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 216 = indica.
gemella, Wall. Cat. n. 4576 = nemoralis.
gemina, *Ruiz, ex Miq. in Mart. Fl. Bras.* iv. 1. 98, in syn.—Peruv.
geminifolia, Miq. in Zoll. Syst. Verz. Ind. Archip. 98, 90 = fistulosa.
geniculata, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 105.—As. trop.; Malaya.
geocoryphe, *Teijsm. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 281.—Ins. Celebes.
gesneriifolia, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 258.—Persia.
gibbosa, *Blume, Bijdr.* 466.—As. trop.; Malaya.
gigantea, *H. B. & K. Nov. Gen. et Sp.* ii. 48.—N. Granat.
gigantea, Noronha, In Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 15 = balica.
Gilsonii, *Miq. Fl. Ind. Bat. Suppl.* 498.—Sumatra.
glabella, *Blume, Bijdr.* 452.—Reg. Himal.; Burma.
glaberrima, *Blume, l. c.* 451.—As. trop.; Malaya.
glaberrima, Hort. Bogor. ex Miq. Fl. Ind. Bat. i. 11. 343 = Blumendijckii.
glabra, *Grif. Notul.* iv. 280.—Assam.
glabra, Vell. Fl. Flum. xi. t. 50 = eximia.
glabrata, H. B. & K. Nov. Gen. et Sp. ii. 47 = anthelmintica.
glandulifera, *Wall. Cat.* n. 4487.—Burma; Malaya.
glandulosa, Hort. ex G. Don, in Loud. Hort. Brit. 416 = exasperata.
glauca, Dam.-Cours. Bot. Cult. ed. II. vi. 357 = cordata.
glauca, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 299.—Mexic.
glaucescens, *Miq. l. c.* 300.—Mexic.
glaucophylla, Desf. Tabl. ed. I. 209 = cordata.
globosa, *Blume, Bijdr.* 449.—Burma; Malaya.
globosa, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 224.—Ital.
glomerata, Hort. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 293 = frangulifolia.
glomerata, *Roxb. Pl. Corom.* ii. 18. t. 128.—Ind. or.; Burma.
glumosa, *Delile, Cent. Pl. Méroé.* 63.—Abyssin.
glycicarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Mexic.
gnaphalocarpa, *Stend. ex Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 113.—Abyssin.
Gomelleira, *Hort. Monac. ex Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 18.—Bras.
Gonia, Buch.-Ham. in Trans. Linn. Soc. xv. (1826) 137 = myrosmoda.
Goolerees, Roxb. Hort. Beng. 65; Fl. Ind. iii. 538 = glomerata.
gossypina, Wall. Cat. n. 4488 = alba.
gracilis, Wall. l. c. n. 4573 (non Flora).—Ind. or.
Granatum, *Forst. f. Pl. Escul.* 37.—Ins. N. Hebrid.
grandibracta, *Mart. ex Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 70.—Bras.
grandifolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 18.—Hab.?
grandifolia, Wall. Cat. n. 4525 = parietalis.
grewioefolia, Blume, Bijdr. 475 = pilifera.
grewioefolia, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 21 = obscura.
grewioefolia, Zoll. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 286 = ramentacea.
Griffithii, *Miq. l. c.* 297.—Ind. or.
griseo, *Vahl, Enum.* ii. 194.—Hab.?
griseo, *Wall. Cat.* n. 4544.—Burma.
grossivenia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 227.—Borneo.
Grossularia, Buch.-Ham. ex Wall. Cat. sub n. 4514 = nervosa.
grossulariodes, Burm. f. Fl. Ind. 227 = heterophylla.
Gaspari, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 67; *Contrib.* 35.—Reg. Argent.
guatemalana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 296.—Guatemala.
guianensis, *Desv. in Ham. Prod. Pl. Ind. Occ.* 23.—Guian.
gummifera, *Bertol. Misc. Bot.* xx. 16.—Guatemala.
gummifera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Guayaquil.
gummiflua, Miq. l. c. 286 = variegata.
Gunnebaceæ, Blume, ex Decne. in Nouv. Ann. Mus. Par. iii. (1834) 404 = F. Benjamina.

FICUS:—
Hamiltoniana, Wall. Cat. n. 4545 A = pomifera.
Hampelos, Steud. Nom. ed. II. i. 686 = trichocarpa.
Hanceana, *Maxim. in Bull. Acad. Pétersb.* xxvii (1881) 558.—China.
hapalophylla, *Kuro, For. Fl. Brit. Burma*, ii. 461.—Ind. or.
Harlandi, *Benth. Fl. Hongk.* 330.—China.
Hartwegii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 299.—N. Granat.
Harveyi, *Seem. Fl. Vit.* 250.—Ins. Fiji.
Haulii, Blanco, Fl. Filip. ed. I. 684 = leucantatoma.
heterocarpa, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 538 = scandens.
Hennesana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 216.—Austral.
Hernandesii, *Miq. l. c.* 300.—Mexic.
heterophylla, *Linn. f. Suppl.* 442; *Lam. Encyc.* ii. 499.—As. trop.
heterophylaea, Blume, Bijdr. 466 = rostrata.
hexaropoda, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 282.—Ins. Moluce.
heterosela, *Sur. in Ann. Sc. Nat.* Sér. v. xiv. (1872) 277.—N. Caled.
hibiscifolia, Champ. ex Benth. in Hook. Kew Journ. vi (1854) 77 = hirta.
hirsuta, Hook. ex Miq. in Hook. Lond. Journ. vi. (1848) 451 = villosa.
hirsuta, Roxb. Hort. Beng. 65; Fl. Ind. iii. 536 = hirta.
hirsuta, *Vell. Fl. Flum.* xi. t. 49; *Schott, in Spreng. Syst.* iv. *Cur. Post.* 414.—Bras.
hirta, *Vahl, Enum.* ii. 201.—As. trop.; Malaya.
hispida, *Linn. f. Suppl.* 442.—As. et Austral. trop.
hispida, Kozb. ex Wall. Cat. n. 4491 = hirta.
hispidissima, Wight, ex Miq. in Hook. Lond. Journ Bot. vii. (1848) 239 = asperrima, *Roxb.*
Hochstetteri, *A. Rich. Tent. Fl. Abyss.* ii. 267.—Abyssin.
holosericea, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 410.—Bras.
Hookeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 215.—Reg. Himal.
Hookezi, *Sweet, Hort. Brit.* ed. I. 362.—Ind. occ.
Houttuijii, Kunth & Bouché, Ind. Sem. Hort. Berol (1846) 15 = macrophylla.
humilis, Reinw. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 294 = F. Ampelos.
humilis, Roxb. Hort. Beng. 65; Fl. Ind. iii. 535 = quercifolia.
Hunteri, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 235 = alba.
hypolenca, *Gasp. Ricerch. Caprif.* 79.—Italia.
hypophylla, Miq. Fl. Jungh. 60 = pilifera.
ichnopoda, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 229, 294 = pyriformis.
ilicina, *Send. in Miq. l. c.* 289.—Afr. austr.
impressa, Champ. ex Benth. in Hook. Kew Journ. vi. (1854) 76 = foveolata.
incisa, *Wall. Cat.* n. 4490.—Burma.
incisinota, Buch.-Ham. in Wall. l. c. n. 4486 A = lepicarpa.
inclinata, *Buch.-Ham. l. c.* 4486 B.—Silhet.
inconstans, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 283 = quercifolia.
inconstantissima, *Miq. Fl. Ind. Bat. Suppl.* 481.—Sumatra.
indecora, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 488 = orbiculata.
indecora, Miq. in Journ. Bot. Néerl. (1861) 242 = opposita.
indica, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 265 = populifolia.
indica, Blanco, Fl. Filip. ed. I. 681 = saxophila.
indica, Forsk. Fl. Aegypt. Arab. 179 = salicifolia.
indica, Herre, ex Rath, Nov. Pl. Sp. 891 = religiosa.
indica, Hochst. ex Wall. Cat. sub n. 4519 = F. Dahra.
indica, Linn. Sp. Pl. ed. II. 1514; Lam. Encyc. ii. 494 = myrosmoda.
indica, *Linn. l. c.* ed. I. 1060.—As. trop.; Malaya.
indica, Willd. Sp. Pl. iv. 1148 = F. Talela.
infectoria, *Roxb. Hort. Beng.* 66; *Fl. Ind.* iii. 551.—As. trop.; Malaya.
infectoria, Willd. Sp. Pl. iv. 1137 = F. Talela.
inflexa, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 288 = Sagittaria.
insigne, *Heer, in Neuen. Ac. Soc. Scaff.* xiii. (1872) ii. 205.—Boreal.

FICUS:—
insipida, Willd. Sp. Pl. iv. 1143 = anthelmintica.
insularis, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 485.—Ins. Loo Choo.
intermedia, *Delile, Cent. Pl. Mirof.* 84.—Aegypt.
intermedia, *Grisf. Notul.* iv. 395.—Ind. or.
intramarginalis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Costa Rica.
involucrata, *Blume, Bijdr.* 447.—Java.
involucrata, Lam. Encyc. ii. 497 = rubra.
involuta, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Nicaragua.
irregularis, *Miq. l. c.* 224.—Ins. Celebes.
irregularis, Steud. Nom. ed. II. i. 636 = gibbosa.
ischnopoda, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 229 = pyriformis.
Isiala, Pritz. Ic. Ind. 459, sphalm. = F. Tsiela.
stenides, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 315 = salicifolia.
isophylla, *Miq. l. c.* 218.—Afr. trop.
jacquinaefolia, *A. Rich. Fl. Cub. Fanerog.* ii 221.—Cuba.
jamaicensis, *Liebm. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 300.—Jamaica.
japonica, Blume, Bijdr. 440 = erecta.
javanica, Hort. Berol. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 324 = erecta.
javanica, *Reinw. ex Steud. Nom* ed. II. i. 636.—Java.
javensis, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 232 = heterophylla.
Johannis, Boiss. Diagn. Ser. I. vii. 96 = F. Carica
joponensis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 298 = yoponensis.
Junghuhniana, Miq. Pl. Jungh. 56 = parietalis.
kalkicarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 288, 289.—Malaya.
Karet, Baill. Hist. Pl. vi. 176 = mysorensis.
Korthalsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 215.—Borneo.
Kotschyana, *Miq. l. c.* 288.—Arab.
Kunthii, *Miq. l. c.* 296.—Bras.
laccifera, Blanco, Fl. Filip. ed. I 678 = leucantatoma.
laccifera, Roxb. Hort. Beng. 66; Fl. Ind. iii. 545 = altissima.
lachnocaulon, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 287 = platypoda.
Lacor, Buch.-Ham. in Trans Linn. Soc. xv. (1827) 150 = infectoria.
laeta, Decne. in Nouv. Ann. Mus. Par. iv. (1834) 495 = gibbosa, Decaisnei.
laevigata, Blanco, Fl. Filip. ed. I. 682 = cuneata, *Miq.*
laevigata, *Vahl, Enum.* ii. 193.—Ind. occ.
laevis, *Blume, Bijdr.* 437.—As. trop.; Malaya; Java.
laevis, Desf. Cat. Hort. Par. ed. III. 414 = sapotaefolia!
Lamberliana, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 286 = infectoria.
laminosa, Hardw. ex Roxb. Hort. Beng. [108]; Fl. Ind. iii. 531 = racemocarpa.
Lamponga, Miq. Fl. Ind. Bat. Suppl. 431 = lepidosa.
lanata, *Blume, Bijdr.* 441.—Java.
lanata, F. Muell. ex Benth. Fl. Austral. vi. 168 = leucotricha.
lanceolata, *Buch.-Ham. ex Roxb. Hort. Beng.* 66; *Fl. Ind.* iii. 557.—Reg. Himal.; Burma.
lancifolia, *Heck. & Arn. Bot. Beech. Voy.* 310.—Mexic.
lancifolia, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 462 = celebica.
lancifolia, Moench, Meth. 347 = indica.
langrera, Wall. Cat. n. 4327 = crinervia.
languinea, *Cauer. Nov. Stirp. Bras. Dec.* 48.—Bras.
lapathifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Mexic.
lasiocarpa, *Miq. Fl. Ind. Bat. Suppl.* 429.—Sumatra.
lasiophlebia, Miq. l. c. 427 = lasiocarpa.
lasiophylla, Link, Enum. Hort. Berol. ii. 449 = benghalensis.
lateriflora, *Vahl, Enum.* ii. 197.—Ins. Mascar.
latifolia, Hort. Berol. ex Kunth. & Bouché, Ind. Sem. Hort. Berol. (1846) 18 = grandifolia.
Latifolia, Salisb. Prod. 16 = F. Carica.
laurifolia, *Hort. ex Lam. Encyc.* ii. 495.—Ind. occ.
laurifolia, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 16 = gibosyrifolia.
Leichhardtii, Miq. Ann. Mus. Lugd. Bat. iii. 287 = platypoda.
lenticinosa, *Vahl, Enum.* ii. 188.—Ind. occ.
Leopoldi, *Van Houtte, ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 230.—Am. exotr.

FICUS:—
lepicarpa, *Blume, Bijdr.* 459.—Malaya.
lepidosa, *Wall. Cat.* n. 4541.—Ind. or.
Leprieurii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 219.—Afr. trop.
leptocarpa, Steud. Nom. ed. II. i. 636 = ramentacea.
leptoclada, *Benth. Fl. Austral.* vi. 172.—Austral.
leucantatoma, *Poir. Encyc. Suppl.* ii. 554.—Malaya.
leucatoma, Roem. & Schult. Syst. i. 501, sphalm. = praec.
leucocarpa, Gasp. Ricerch. Caprif. 77 = F. Carica.
leucocarpa, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 285 = infectoria.
leucocoma, Miq. Pl. Jungh. 54 = tricolor.
leucophlaea, Blume, Bijdr. 448 = leucantatoma.
leucoptera, *Miq. Fl. Jungh.* 62.—Ins. Celebes.
leucosticta, Spreng. Syst. iii. 779 = leucantatoma.
leucotoma, Miq. in Hook. Lond. Journ. vii. (1848) 468, in syn., sphalm. = leucantatoma.
leucotricha, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 285.—Austral.
leuconylon, Miq. Pl. Jungh. 61 = pisifera.
Lichtensteinii, Link, Enum. Hort. Berol. ii. 451 = capensis.
Liebmannians, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Panama.
ligustrina, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Mexic.
Lima, Koyen, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 293 = hispida.
littoralis, Blume, Bijdr. 455 = retusa.
Lobbii, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 233 = tontata.
Loddigesii, Heynh. Nom ed. 224 = quercifolia.
Logani, *G. Don, in Sweet, Hort. Brit.* ed. III. 607, nomen.—Venezuela.
longifolia, Buch.-Ham. in Wall. Cat. n. 4570, partim = indica.
longifolia, Buch.-Ham. l. c. 4570 A, B = obtusifolia.
longifolia, *Schott, in Spreng. Syst. iv. Add.* 409.—Bras.
longipes, *Baker, in Journ. Linn. Soc. xx.* (1883) 259.—Madag.
longipes, *Grisf. Notul.* iv. 397.—Assam.
longipes, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Mexic.
lucescens, Blume, Bijdr. 444 = infectoria.
lucescens, *Miq. Ann. Mus. Bot Lugd. Bat* iii. 264, obs. sumatrana.
lucida, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii 451.—Ind. or.
lucida, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 16 = nitida.
ludens, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 234.—Afr. trop.
ludens, Wall. Cat. n. 4579 = foveolata.
Luducca, Roxb. Hort. Beng. [108]; Fl. Ind. iii. 534 = foveolata.
Luschnathiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Bras.
lutea, *Vahl, Enum.* ii. 185.—Afr. trop.
lutescens, Desf. Cat. Hort. Par. ed. III. 413 = diversifolia.
macrocalyx, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 217 = involucrata.
macrocarpa, Blume, Bijdr. 459 = punctata.
macrocarpa, Gasp. Ricerch. Caprif. 78 = F. Carica.
macrocarpa, Hueg. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 287 = macrophylla.
macrocarpa, *Wight, ex Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 74.—Ind. or.
macrophylla, *Desf Tabl.* ed I. 209.—Austral.
macrophylla, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 15 = anacardiifolia.
macrophylla, Roxb. Hort. Beng. 66; Fl. Ind. iii. 556 = Roxburghii.
macropoda, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 150.—Ins. Andaman.
macropoda, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 442.—Ins. Philipp.
maculata, *Linn. Sp. Pl.* ed. II. 1515.—Ins. S. Doming.; Cochinch.
maduraua, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 222.—Java.
magnifolia, *F. Muell. Fragm.* iv. 50.—Australia.
magnoliaefolia, Blume, Bijdr. 448 = retusa.
mahabarica, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 457.—Ind. or.

FICUS:—
magnifolia, *Grisf. Notul.* iv. 395.—Assam.
Mensis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 261 = gibbosa, *Blume*.
Mappan, Miq. Fl. Ind. Bat. Suppl. 425 = alba.
marginalis, Buch.-Ham. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 293 = rostrata.
marginalis, Hunter, ex Miq. in Hook. Lond. Journ Bot. vii. (1848) 429 = urophylla.
marmorata, *Bey. ex Baker, in Journ. Linn. Soc.* xx. (1883) 257.—Madag.
martabanica, Wall. Cat. n. 4551 = lepidosa.
martinicensis, Hort. Berol. ex Walp. Ann. i. 676 = Kunthii.
martinicensis, Willd. Sp. Pl. iv. 1137 = laurifolia.
Martinii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 219.—Gutan.
Masoni, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 271.—Ins. Fiji.
Mathewsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Peruv.
mauritiana, *Lam. Encyc.* ii. 490.—Ins Mascar.
maxima, Mill. Gard. Dict. ed. VIII. n. 6 = laurifolia.
Maximiliana, *Mart. ex Miq. in Hook. Lond. Journ. Bot.* vi. (1847) 529, in syn.—Bras.
megapoda, *Baker, in Journ. Linn. Soc.* xxi. (1885) 444.—Madag.
melinocarpa, *Blume, Bijdr.* 460.—Java.
Melleri, *Baker, in Journ. Linn. Soc.* xx. (1883) 258.—Madag.
membranacea, *Wright, in Sauv. Fl. Cub.* 149.—Cuba.
menadana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 233.—Ins. Celebes.
mexicana, *Miq. l. c.* 299, 300.—Mexic.
microcantha, Miq. l. c. 221 = aculeata.
microcarpa, Blume, Bijdr. 442 = ramentacea.
microcarpa, Hort. Berol. ex Walp. Ann. i. 677 = cestrifolia.
microcarpa, Linn. f. Suppl. 442 = retusa.
microcarpa, *Vahl, Enum.* ii. 188.—Afr. trop.
microphylla, *Salzm. ex Miq. in Mart. Fl. Bras.* iv. i. 93.—Bras.
microstoma, *Wall. Cat.* n. 4566.—Malaya.
microtus, Miq. Fl. Ind. Bat. Suppl. 428 = pisifera.
Millettii, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 438 = seq.
Millettii, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 281 = pyriformis.
millingtonifolia, *Grisf. Notul.* iv. 396.—Assam.
Minahassae, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 231.—Ins. Celebes.
minima, Hort. ex Gard. Chron. (1880) ii. 716 = pumila.
modesta, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 263 = nervosa.
mollior, *F. Muell. ex Benth. Fl. Austral.* vi. 173.—Austral.
mollis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 283 = glomerata.
mollis, Vahl, Symb. Bot. i. 82 = tomentosa.
mollis, Willd. in Mém. Acad. Berl. (1798) 108 = hispida.
montana, Burm. f. Fl. Ind. 226 = quercifolia.
montana, Roxb. in Wall. Cat. n. 4574 = nervosa.
monticola, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 216.—Ind. or.
Moonlana, *King, ex Trim. Cat. Pl. Ceyl.* 84.—Zeylan.
Moorei, *Seem. Fl. Vit.* 249.—Ins. Pacif.
morifolia, Forsk. Fl. Aegypt. Arab. 179 = Forskâlaei.
morifolia, Husson, ex Boiss. Fl. Orient. iv. 1155 = palmata.
morifolia, Lam. Encyc. ii. 499 = laterifiora.
morifolia, *Vahl, Enum.* ii. 208.—Ind. or.
Motleyana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 238.—Borneo.
Mueller, *Miq. l. c.* 287.—Austral.
Muntii, *Link, Enum. Hort. Berol.* ii. 450.—Austral.
muriculata, Miq. in Zoll. Syst. Verz. Ind. Archip. 93, 98 = F. Ribes.
Murrayana, Hort. ex Miq. in Mart. Fl. Bras. iv. i. 95 = Princepa.
mutabilis, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 289.—N. Caled.
myriocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 230.—Ins. Amboin.
myrtifolia, *Link, Enum. Hort. Berol.* ii. 450.—Hab.?
mysorensis, *Heyne, ex Roth, Nov. Pl. Sp.* 390.—Ind. or.

FICUS :—
myrsæfolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 18.—Mexico.
natalensis, *Hochst. in Flora*, xxviii. (1845) 88.—Afr. austr.
nautarum, *Baker, Fl. Maurit.* 285.—Ins. Seychelles.
neapolitana, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 223.—Italia.
neglecta, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 494 = F. Benjamina.
nemoralis, *Wall. Cat.* n. 4517.—Reg. Himal.
nepalensis, Blanco, Fl. Filip. ed. II. 474 = F. Benjamina.
nepalensis, Spreng. Syst. iii. 729 = foveolata.
nervifolia, Hort. ex Walp. Ann. i. 676 = Kunthii.
nerifolia, *Reinw. ex Steud. Nom.* ed. II. i. 637.—Java.
nerifolia, *A. Rich. Tent. Fl. Abyss.* ii. 267.—Abyss.
nervosa, *Heyne, ex Roth, Nov. Sp. Pl.* 388 —As trop ; Malaya.
neurophila, F. Muell. ex Miq. Ann. Mus. Lugd. Bat. iii 296 = glabella.
Neumanni, *Cels, ex Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 14.—Hab. ?
nimphæifolia, Cerv. in Suppl. Gaz. Lit. Mexico (2 Jul. 1794), 4 = nymphææfolia.
nipponica, Franch. & Sav. Enum. Pl. Jap. i. 436 = foveolata.
nitida, *Blume, Bijdr.* 455.—Java.
nitida, Heyne, ex Roth, Nov. Pl. Sp. 387 = F. Benjamina.
nitida, Hook. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 582 = Hookeri, *Sweet*
nitida, Thunb. Diss. Fic. 10 = retusa.
nitidifolia, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 266.—N. Caled.
nivea, Blume, Bijdr. 476 = alba.
nodosa, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 245.—Ins. Amboin.
nodulosa, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 298 = obtusa.
Noronhæ, *Oliver, in Hook. Ic. Pl.* t. 1922.—Ins. Fernand. Noronh.
Notobor, Buch.-Ham. ex Wall. Cat. sub n. 4502 = F. Benjamina.
Novæ-Guineæ, *Zipp. in Flora*, xii. (1829) 282, nomen —N. Guin.
nuda, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 288 = F. Benjamina.
nymphææfolia, *Mill. Gard. Dict.* ed. VIII. n. 9.—Am. trop.
nymphoides, *Thumb. Diss. Fic.* 7.—Am. austr.
obliqua, *Forst. f. Prod.* 77.—Ins. Pacif.
obliqua, Miq. in Zoll. Syst. Verz. Ind. Archip. 92, 98 = melinocarpa.
oblongata, *Link, Enum. Hort. Berol.* ii. 449.—Afr. austr.
oblongifolia, *D. Don, Prod. Fl. Nep.* 61.—Nepal.
obovata, *Grif. Notul.* iv. 390 ; *Ic. Pl. Ax.* t. 502. f. 1. —Assam.
obovata, Hook. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 579 = glabella.
obscura, *Blume, Bijdr.* 474.—Reg. Himal. ; Malaya.
obtecta, Wall. Cat. n. 4505 = villosa.
obtusa, *Hassk. Cat. Hort. Bog. Alt.* 75.—Malaya.
obtusata, *Link, Enum. Hort. Berol.* ii. 450.—Hab. ?
obtusidens, Miq. Fl. Jungh. 59 = rostrata.
obtusifolia, H. B. & K. Nov. Gen. et Sp. ii. 49 = Bonplandiana.
obtusifolia, *Roxb. Hort. Beng.* [103] ; *Fl. Ind.* iii. 546.—Reg. Himal. ; Malaya.
obtusiuscula, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 300. —Bras.
ochroleuca, *Griseb. Fl. Brit. W. Ind.* 151.—Jamaic.
Oerstedtiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 299. —Panama.
Oldhami, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 242.—Ins. Formosa.
oligodon, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 284 = pomifera.
oligoneura, *Miq. l. c.* 288.—Sumatra.
oligosperma, Miq. Fl. Jungh. 55 = ramentacea.
omissa, Wall. Cat. n. 4508 = globosa, *Blume*.
opposita, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 426.—Austral.
oppositifolia, Willd. Sp. Pl. iv. 1151 = hispida.
orbicularis, *A. Cunn. ex Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 426.—Austral.

FICUS :—
organensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 299. —Bras.
ottoniæfolia, *Miq. l. c.* 288.—Afr. trop.
ovalis, *Miq. l. c.* 298.—Costa Rica.
ovata, D. Don, Prod. Fl. Nep. 61 = scandens ?
ovata, *Vahl, Enum.* ii. 185.—Afr. trop.
ovoidea, Jack, in Malay. Misc. ii. (1822) vii. 71 = diversifolia.
ovoidea, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 20.—Ind. or.
oxyphylla, Miq. in Zoll. Syst. Verz. Ind. Archip. 98 = erecta.
pachycarpa, Gasp. Ricerch. Caprif. 78 = F. Carica.
Padana, Burm. f. Fl. Ind. 226 : toxicaria.
padifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 47.—Mexic.
palæocrya, Steud. Nom. ed. II. i. 687 = retusa.
pallida, *Vahl, Enum.* ii. 174.—N. Granat.
pallida, Wall. Cat. n. 4567 = retusa.
palmata, *Forsk. Fl. Aegypt. Arab.* 179.—Afr. trop. ; Arab ; Ind. or.
palmata, Roxb. Hort. Beng. [108] ; Fl. Ind. iii. 529 = alba.
paludosa, *Perr. in Allm. Soc. Linn. Par.* iii. (1825) 117.—Java.
palustris, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 15 = anacardiifolia.
Pancheriana, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 274.—N. Caled.
panduræfolia, Hort. Par. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 546 = hirsuta
panduræformis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 299 = heterophylla.
pandurata, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 229.—China.
panifica, *Delile, l. c.* Sér II. xx. (1843) 94.—Abyss.
papyrifera, Griff. Notul. iv. 394 ; Ic. Pl. Ax. t. 554 = F. Benjamina.
paradoxa, Blume, Bijdr. 467 = gibbosa.
parasnæa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Bras.
parasitica, *Koen. ex Willd. in Mém. Acad. Berl.* 1798 (1802) 102 = gibbosa.
Parcelli, *Veitch, ex Capt. & Marchal, Pl. Ornem.* iii. (1874) t. 47.—Ins. Pacific.
parietalis, *Blume, Bijdr.* 463.—Malaya.
parietaria, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 15 = heterophylla ?
Parkeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 300.—Guian.
parvifolia, Miq. l. c. 286 = glabella.
Payapa, Blanco, Fl. Filip. ed. I. 688 = retusa.
pedicellata, Wall. Cat. n. 4486, partim = laevis.
pedunculara, *Wall. l. c.* n. 4528.—Burma.
pedunculata, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 450.—Ind. occ.
pedunculata, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 426.—Italia.
pedunculosa, *Miq. in Hook. l. c.* 442.—Ins. Philipp.
Peepul, Griff. Notul. iv. 388 = religiosa.
peguensis, Buch.-Ham. ex Wall. Cat. n. 4560 = altissima.
pellucido-punctata, Griff. Notul. iv. 394 = indica.
peltata, *Blume, Bijdr.* 488.—Java.
peltata, Link, Enum. Hort. Berol. ii. 450 = F. Benjamina.
peracuta, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 306, Index = pyrandromia.
Pereng, Steud. Nom. ed. II. i. 687 = gibbosa.
perforata, Linn. Fil. Suppl. n. 146 ; Amoen. Acad. viii. 365 = repens.
perforata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 300.—Bras.
pergamenea, *Gal. ex Miq. l. c.* 291.—Mexic.
perglobocarpella, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Am. trop. ?
persica, *Boiss. Diagn.* Ser. I. vii. 97.—Persia.
pertusa, Bory, ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 558 = urceolata.
pertusa, *Linn. f. Suppl.* 442.—Guian.
peruviana, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 423 = gibbosa.
petiolaris, *H. B. & K. Nov. Gen. et Sp.* ii. 49.—Mexic.
petiolata, Reinw. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 287 = superba.

FICUS :—
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 271.—Abyss.
petrophila, Hask. Cat. Hort. Bog. Alt. 75 = anaophila
phæopoda, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 293 —Java.
philippensæ, *Hort. ex Hérincq, L'Hortic. Franç.* (1869) 244.—Ins. Philipp.
philippinensis, Miq. in Hook. Lond. Journ. Bot. vii (1848) 435 = Decaisnei
phlebophylla, Miq. Fl. Ind. Bat. Suppl. 480 = parietalis.
phyllosperma, *Noronha, in Verh. Batav. Gen.* v (1790) ed. I. Art. iv. 15.—Java.
phyrolaccæfolia, *Desf. Cat. Hort. Par.* ed. III. 413. —Hab. ?
picta, Noronha, in Verh. Batav. Gen. v. (1790) ed I Art. iv. 15 = F. Benjamina.
pilosa, *Herb. Russ. in Wall. Cat.* n. 4588 = hispida.
pilosa, *Reinw ex Blume, Bijdr.* 446.—Burma ; Malaya ; Austral.
Pinkiana, *F. Muell. in Wing, South. Sc. Record*, ii (1882) 273.—Austral.
piperifolia, Miq Ann. Mus. Bot. Lugd. Bat. iii. 298 = obtusa.
pisifera, Hook. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 581 = retusa.
pisifera, *Wall. Cat.* n. 4504.—Malaya.
pisiformis, Hort. Berol. ex Miq. in Hook. Lond Journ. Bot. vii. (1848) 458 = lutescens.
pisocarpa, *Blume, Bijdr.* 464.—Malaya.
planicostata, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Hab. ?
platycaula, Miq. Fl. Ind. Bat. i. ii. 316 = obtusa.
platyphylla, *Delile, Cent. Pl. Hérof.* 62.—Aegypt.
platyphylla, *A. Cunn. ex Miq. in Hook. Lond. Journ Bot.* vi. (1847) 561, in syn.—Austral.
plerocarpa, *F. Muell Fragm.* viii. 246.—Austral.
podophylla, *Baker, in Journ. Linn. Soc.* xxi. (1885) 444.—Madag.
Pohliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Bras.
polita, Hort. in Miq. Ann. Mus. Lugd. Bot. Bat. iii. 299 = myrtifolia.
polita, *Vahl, Enum.* ii. 182.—Afr. trop.
politoria, Lam. Encyc. ii. 500 = F. Ampelos.
politoria, *Lour. Fl. Cochinch.* ii 667.—Cochinch.
politoria, Moon, Cat. Ceyl. Pl. 74 = asperrima, *Roxb*
polycarpa, Buch.-Ham. in Wall. Cat. n. 4545 = pomifera.
polycarpa, Roxb. Hort. Beng. 66 ; Fl. Ind. iii. 556 = F. Ribes.
polycarpa, *Jacq. Obs. Bot.* ii. 29.—Venezula.
polymorpha, *Gasp. Ricerch. Caprif.* 77.—Italia.
polyphlebia, *Baker, in Journ. Linn. Soc.* xx. (1884) 262.—Madag.
polypus, Schiede, ex Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 412. = arbutifolia.
pomifera, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 106 = kalkcarpa.
pomifera, *Wall. Cat.* n. 4547.—Reg. Himal. ; Malaya
Populnaster, Desf. Cat. Hort. Par. ed. III. 412 = Amottiana ?
populifolia, Desf. Tabl. ed. I. 309 = populnea.
populifolia, *Vahl, Symb. Bot.* i. 82.—Arab.
populifolia, Wall. Cat. n. 4482 A, = Amottiana.
populiformis, Schott, ex Miq. Ann. Mus. Bot. Lugd Bat. iii. 287 = Arnottiana ?
populnea, Decne. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 280 = asaophila.
populnea, Kunth & Bouché, in Ann. Sc. Nat. Sér. III. vii. (1847) 286 = Arnottiana ?
populnea, *Willd. Sp. Pl.* iv. 1141.—Am. austr.
Portiana, *Regel, Gartenfl.* (1862) 280.—Mexic.
praecox, Gasp. in Guss. Fl. Sic. Syn. ii. 880 = F. Carica.
Princeps, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 14.—Bras.
prinoides, *Cham. & Schlecht. in Linnæa*, vi. (1831) 387 = ligustrina.
prinoides, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1149. —N. Granat.
Pritchardii, *Seem. Fl. Vit.* 282.—Ins. Fiji.
procera, *Reinw. ex Blume, Bijdr.* 448.—Java.
procera, Salisb. Prod. 396 = benghalensis.
prolixa, *Forst. f. Prod.* 77.—Ins. Societat.
pruriens, Vieill. & Deplanche, Ins. Nouv. Calléd. 114 = retusa.

FICUS :—
prominens, Wall. Cat. n. 4537 = hispida.
pronigena, Hort. Schoenb. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 209 = sapotaefolia.
prostrata, *Buch.-Ham. ex Wall. Cat.* n. 4536, partim. —Reg. Himal.
prostrata, Wall. l. c., partim = F. Ribes.
protensa, *Hemsl. Biol. Centr. Am.* iii. 147.—Panama.
Protesa, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 250.—N. Caled.
prunifornis, *Blume, Bijdr.* 451.—Malaya
pruniformis, Hort. Bogor. ex Miq. Fl. Ind. Bat. i. 1. 317 = variabilis.
Pseudo-Benjamina, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 286 = F. Talela.
Pseudo-Carica, *Miq. in Hook Lond. Journ. Bot.* vii. (1848) 226.—Abyss.
Pseudo-radula, *Miq. Ann. Mus. Bot. Lugd Bat.* iii. 299.—Mexic.
Pseudo-rubra, *Miq. l. c.* 264.—Java.
Pseudo-Sycomorus, Decne. in Ann. Sc. Nat. Ser. II. ii. (1854) 242 = palmata.
Pseudo-Talela, *Trim. Cat. Pl. Cryl.* 84.—Zeylan.
psychotrioefolia, Miq. Ann. Mus. Lugd. Bat. iii. 286 = Cunninghamii.
puberula, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 19. n. 45 = hirsuta.
puberula, Miq. Ann. Mus. Lugd. Bat. iii 287.—Austral.
pubescens, Heyne, ex Roth, Nov Sp. Pl. 387 = tomentosa.
pubigera, *Kurz, For. Fl. Burma*, ii. 450 (sp. dub.).—Burma.
pubigera, Wall. Cat. n 4518 = foveolata.
pubinervia, *Blume, Bijdr.* 458.—Java.
pubinervis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 496 = Decaisnei.
pulchella, *Schott, ex Spreng. Syst.* iv 410 —Bras.
pulchra, Wall. Cat. n. 4571 (non Firl sp.).—Singapore.
pumila, *Linn. Sp Pl.* 1060.—Japon ; China.
pumila, Thunb. Diss. Fic. 9 = erecta.
punctata, Heyne, ex Wall. Cat. n. 4569 = infectoria.
punctata, Lam. Encyc. ii. 495 = aggregata.
punctata, *Thunb. Diss. Fic.* 9.—Malaya.
punctulata, *Thunb. l. c.* = Ind. or.
punctata, Steud. Nom. ed. II. i. 637, sphalm. = aggregata.
pungens, *Reinw. ex Blume, Bijdr.* 476.—Java
purpurascens, *Blume, l. c.* 471.—Java.
purpurascens, *Desf. Cat. Hort. Par.* ed. III 412.—Hab. ?
pyrifolia, Burm. f. Fl. Ind. 226 = erecta.
pyrifolia, Desf. Cat. Hort. Par. ed. III. 413 = Fontanesii.
pyrifolia, Lam. Encyc. ii. 495 = rubra.
pyrifolia, Salisb. Prod. 16 = F. Benjamina.
pyriformis, *Hook. & Arn. Bot. Beech. Voy.* 216.—Reg. Himal. ; China.
pyrrhocarpa, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 106 = racemocarpa.
pyrrhopoda, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 293 = obtusa.
quercifolia, Blume, Bijdr. 468 = rostrata.
quercifolia, *Roxb. Hort. Beng.* 65 ; *Fl. Ind.* iii. 534.—Burma ; Malaya.
racemifera, Roxb. ll. cc. 86, 560 = variegata.
racemigera, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 257.—N. Caled.
racemosa, Linn. Sp. Pl 1060 = hispida ?
racemosa, Wall. Cat. n. 4549 = glomerata.
racemosa, Willd. Sp. Pl. iv. 1146 = infectoria.
radiata, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 494 = leucantatoma.
radicans, Casar. in Att. Ters. Rim. Sc. Ital. (1841) 315 = F. Arpazusa.
radicans, *Desf. Cat. Hort. Par.* ed. III. 413.—Hab. ?
radicans, Roxb. Hort. Beng. 65 ; Fl. Ind. iii. 536 = rostrata.
Radula, *Willd. Sp. Pl.* iv. 1144.—Venezuela.
ramosa, Wall. Cat. n. 4556 = rubra.
ramentacea, *Roxb. Hort. Beng.* [108] ; Fl. Ind. iii. 546.—Reg. Himal. ; Malaya.
ramiformis, Roxb. ll. cc. 86 ; 561 = leucantatoma.
ramiflora, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 436 = rostrata.
ramiflora, *Desf. Cat. Hort. Par.* ed. III. 346, 412. —Hab. ?

FICUS :—
recurva, *Blume, Bijdr.* 457.—Malaya.
reflexa, Buch.-Ham. in Wall. Cat n. 4523 = retusa.
reflexa, *Thunb. Diss. Fic.* 11.—Ind. or.
regia, Miq. Ann. Mus. Bot. Lugd Bat. iii. 280 = Roxburghii, pomifera.
Regina, Hort. ex Miq. in Hook. Lond. Journ. Bot. vii. (1848) 228 = F. Carica.
Reinwardtis, Link & Otto, Ic. Pl. Rar. i. 6. t. 31 = falva, *Reinw.*
relapsana, Decne. ex Miq. Ann Mus. Bot. Lugd. Bat. iii. 287 = Rumphii.
religiosa, Forsk. Fl. Aegypt. Arab. 180 = populifolia.
religiosa, *Linn. Sp. Pl.* 1059.—Ind. or.
Remblas, Miq. Fl. Jungh. 60 = pisifera.
renitens, Miq. Fl. Ind. Bat. i. 11. 316 = variabilis.
repens, Hook. ex Miq. in Hook. Lond. Journ. Bot. vii (1848) 465 = hispida.
repens, Hort.—Cf. Gard. Chron. (1880) ii. 716 = pumila.
repens, Hort. Mus. ex Miq. in Hook. Lond. Journ. Bot. vii. (1848) 229 = quercifolia.
repens, *Kettl. in Ges. Naturf. Fr. Neue Schr.* iv (1808) 206.—Ind. or.
repens, Willd. Sp. Pl. iv. 1149 = heterophylla
reticulata, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 294 = laevigata.
reticulata, *Thunb. Diss. Fic.* 12.—Ind. or.
reticulosa, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 436 = gibbosa.
retusa, *Linn. Mant.* 129.—As. trop. ; Malaya.
retusa, Wall. Cat. n. 4530 = diversifolia.
rhizocarpa, *Teysm. & De Vriese, Cat. Hort. Bog.* 83, nomen.—Malaya.
rhododendrifolia, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 18 = Kunthii
rhododendrifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Reg. Himal.
rhomboidalis, *Vahl, Enum.* ii. 199.—Ind or.
rhynchophylla, Steud. Nom. ed. II. i. 637 = religiosa.
Ribes, *Reinw ex Blume, Bijdr.* 463.—Burma ; Malaya.
ribesioides, Wall. Cat. n. 4522 = recurva.
Riedlei, *Teysm. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 228.—Malaya.
rigescens, Miq. l. c. 278 = ramentacea.
rigida, Blume, Bijdr 465 = gibbosa.
rigida, Roxb.-Hau. in Wall. Cat. n. 4527, 4535 = infectoria.
rigida, Desf. Cat. Hort. Par. ed. III. 412 = Neumanni.
rigida, *Buch. in Malay. Misc.* ii. (1822) vii. 72.—Malaya.
rigida, *Miq. Ann. Sc. Nat.* Sér. V. xiv. (1872) 265.—N. Caled.
riparia, *Hochst. ex Miq. in Hook. Lond. Journ. Bot. vii.* (1848) 114.—Abyss.
Rolandri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Am. trop.
rostrata, Hort. Bogor. ex Miq. l. c. 278 = radicans.
rostrata, *Lam. Encyc.* ii. 498.—Reg. Himal. ; Malaya.
rotundifolia, *Roxb. Hort. Beng.* [108] ; *Fl. Ind.* iii. 555.—Ind. or.
Roxburghii, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 456 = hirta.
Roxburghii, *Wall. Cat.* n. 4508.—Reg. Himal. ; Burma.
rubescens, Blume, Bijdr. 458 = indica.
rubescens, Hort. Bogor. ex Miq. Fl. Ind. Bat. i. 11. 352 = annulata.
rubiginosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 289. —Afr. trop.
rubifolia, Griff. Notul. iv. 399 = heterophylla.
rubiginosa, *Desf. Tabl. ed.* I. 309.—Austral.
rubra, Blume, Bijdr. 458 = F. Pseudo-rubra.
rubra, Roth, Nov. Sp. 391 = retusa.
rubra, *Vahl, Enum.* ii. 191.—Ind. or.
rubricaulis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 496 = F. Ampelos.
rubrinervis, *Link, Enum. Hort. Berol.* ii. 448.—Bras.
rudis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 292.—Ins. Celebes.
rudis, Pers. Syn. ii. 609 = F. Ampelos.
rufa, *Schott, in Spreng. Syst.* iv. *Add.* 410.—Bras.
rufescens, Vahl, Enum. 208 = heterophylla.

FICUS :—
rufipila, Miq. Fl. Jungh. 57 = parietalis.
ruyssa, *G. Don, in Loud. Hort. Brit.* 416, nomen.—Afr. trop.
rugosa, *Miq. in Hook. Lond. Journ. Bot.* vii. (1848) 224.—Ital.
Rumphii, *Blume, Bijdr.* 457.—Reg. Himal. ; Malaya
rupestris, *Blume, l. c.* 467.—Java.
rupestris, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 130 = mysorensis.
saemocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 282 —Reg. Himal.
sagittata, *Vahl, Symb. Bot.* i. 88.—Ind. or.
salicifolia, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 431 = akkimensis.
salicifolia, *Vahl, Symb. Bot.* i. 82.—Arab.
salicina, F. Muell. Fragm. iv. 49 = coronulata.
saligna, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 268 = scrocarpa.
Salzmannians, *Miq Ann. Mus. Bot. Lugd. Bat.* iii. 299 —Bras.
Sancti-Crucis, *Miq. l. c.*—Mexic.
sapida, *Miq. l. c.* 298.—Costa Rica.
sapotaefolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 17.—Hab. ?
sapotoides, *Baker, Fl. Maurit.* 284.—Ins. Mascar.
sarmentosa, Buch.-Ham. ex Wall. Cat. n. 4533 = scandens.
Sausureana, *DC. in Mém. Soc. Phys. Genève.* ix (1841) 65.—Am. austr.
saxatilis, Blume, Bijdr 460 = rostrata.
saxatilis, Miq. in Zoll. Syst. Verz. 92. n. 876 = pisifera.
saxophila, *Blume, Bijdr.* 437.—Java ; Timor.
scaberrima, Blume, l. c. 474 = albacura.
scabra, *Forst. f. Prod.* 76.—N. Hebrid.
scabra, Hort. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 291 = simifolia.
scabra, Jacq. Hort. Schoenb. iii. 36 = hispida.
scabra, Willd. in Mém. Acad. Berl. (1798) t. 2 = exasperata.
scabrella, Roxb. Hort. Beng. [108] ; Fl. Ind. iii. 532 = heterophylla.
scabrifolia, *A Rich. Sert Astrol.* t. 17.—Austral.
scandens, Buch.-Ham. in Trans. Linn. Soc. xv. (1826) 149 = infectoria.
scandens, Hort. Van Houtte, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 300 = volubilis.
scandens, Lam. Encyc. ii. 498 = pumila.
scandens, *Roxb. Hort. Beng.* 65, *Fl. Ind.* iii. 536 — Reg. Himal ; Burma.
Schimperi, Hochst. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 555 = Hochstetteri.
Schimperiana, *Hochst. ex Miq. l. c.* 554, in syn.—Abyss.
Schottii, Hort. Berol. ex Kunth & Bouche, Ind. Sem. Hort. Berol. (1846) 14 = catappaefolia.
Schumacheri, *Griseb. Fl. Brit. W. Ind.* 151.—Ind. occ.
Schweinfurthii, *Miq. ex Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xvii. (1868) 688.—Afr. trop.
sclerocarpa, Griff. Notul. iv. 397 = Roxburghii.
sclerocoma, *Miq. Pl. Jungh.* 58.—Java.
sclerophylla, Roxb. Fl. Ind. iii. 546 = gibbosa.
sclerophylla, Griff. Ic. Pl. As. t. 558 = sclerocarpa, Roxburghii.
sclerotera, Miq. Pl. Jungh. 68 = calloza.
scobina, *Brath. Fl. Austral* vi. 176.—Austral.
Segorine, *Miq. Ann. Mus. Bot Lugd. Bat.* iii. 300.—Nicaragua.
semicordata, *Miq. l. c.* 296.—Ins. Celebes.
senegalensis, *Miq. l. c.* 280.—Afr. trop.
septica, Burm. f. Fl. Ind. 226 = leucantatoma ?
septica, *Forst. f. Prod.* 76.—N. Hebrid.
septica, Hook. ex Miq. in Hook. Lond. Journ. Bot. vii. (1848) 436 = insularis.
septica, *Lour. Fl. Cochinch.* ii. 667.—Cochinch.
septica, Royen, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 275 = calloza.
serpyllifolia, *Blume, Bijdr.* 448.—Java.
serraria, *Miq. Fl. Ind. Bat. Suppl.* 428.—Sumatra.
serrata, *Forsk. Fl. Aegypt. Arab.* 179 (= sec. ?).—Arab.
serrata, *Linn. Syst.* ed. X. 1315.—Arab.
setacea, *Ladd. Bot. Cab.* t. 1188.—'India.'
setifera, Steud. Nom. ed. II. i. 638 = hirta.
setosa, Blume, Bijdr. 477 = hirta.
setosa, *Hook. & Arn. Bot. Beech. Voy.* 216. t. 49.—China
sideroxylifolia, Griff. Notul. iv. 389 = diversifolia.

FICUS :—
Sieboldi, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 199 = erecta.
sikkimensis, *Miq. l. c.* iii. 225.—Reg. Himal.
silhetensis, *Miq. l. c.* 223.—Reg. Himal.
silvestris, Risso, Fl. Nice, 484 = F. Carica !
simplicissima, Lam. Encyc. ii. 498 = hispida.
simplicissima, *Lour. Fl. Cochinch.* ii. 667.—Cochinch.
sinuata, *Thunb. Diss. Fic.* 12.—Ind. or.
sinuosa, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 332 = quercifolia.
Smithii, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 372.—Ins. Fiji.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 96.—Ins. Socotr.
sondana, *Miq. Fl. Ind. Bat.* i. 11. 305.—Malaya.
Sonderi, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 295.—Malaya.
sorocoroides, *Baker, in Journ. Linn. Soc.* xx. (1883) 258.—Madag.
sororia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1840) 16.—Hab. ?
Spanogheana, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 451 = recurva.
spathulata, Miq. l. c. 441 = diversifolia.
speciosa, [Fisch.] Hort. Gorenk. ed. 1. (1808) 22, nomen = sallcifolia.
spectabilis, Kunth & Bouché, in Ann. Sc. Nat. Sér. III. vii. (1847) 255 = ovata.
sphaeridiophora, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 295.—Guian.
sphaerophylla, *Baker, in Journ. Linn. Soc.* xxi. (1885) 443.—Madag.
splendens, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 14.—Hab. ?
squamellosa, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 287 = macrophylla.
squamosa, Roxb. Hort. Beng. [103] ; Fl. Ind. iii. 531 = saemocarpa.
stenocarpa, *F. Muell. ex Benth. Fl. Austral.* vi. 174.—Austral.
stictocarpa, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 284 = leucantatoma.
stipulacea, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 15 = pumila !
stipulata, Moon, Cat. Ceyl. Pl. 6 = Thwaitesii.
stipulata, Thunb. Diss. Fic. 8 = pumila.
stipulata, Wall. Cat. n. 4574 = punctata.
stipulosa, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 287 = infectoria.
Storckii, *Seem. Fl. Vit.* 251.—Ins. Fiji.
striata, Roth, Nov. Sp. Fl. 387 = F. Benjamina.
stricta, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 266.—Java.
strigosa, Blume, Bijdr. 441 = recurva.
strigosa, Hort. Bogor. ex Miq. Fl. Ind. Bat. i. 1. 318 = obtusa.
stupenda, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Java.
subspicilata, *Miq. l. c.* 298.—Bras.
subcalyptrata, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 15 = lucescens.
subcordata, *Blume, Bijdr.* 440.—Java.
subcordata, Hort. Bogor. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 295 = callosa.
subcuneata, *Miq. l. c.* 285.—Ins. Ceram.
subdenticulata, Miq. Fl. Ind. Bat. i. 11. 323 = obscura.
subglabra, *F. Muell. Fragm.* ix. 152.—Austral.
subsymphaefolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 295.—Hab. ?
subsoliqua, Miq. l. c. 225 = glabella.
subspaca, Miq. Fl. Ind. Bat. i. 11. 320 = variegata.
subsubcularis, Miq. Fl. Ind. Bat. Suppl. 425 = fulva, *Reinw.*
suborna, Buch.-Ham. in Wall. Cat. n. 4557 = elastica.
subpandurae formis, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 335 = heterophylla.
subpodunculata, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 217 = glabella.
subpyriformis, Miq. l. c. 229 = pyriformis.
subracemosa, Blume, Bijdr. 469 = variegata.
subrepanda, *Wall. Cat.* n. 4568 B.—Reg. Himal.
subrigida, Miq. Fl. Ind. Bat. Suppl. 433 = ramentacea.
subsetulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 295.—Borneo.
subtriplinervis, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 67.—Bras.
subulata, *Blume, Bijdr.* 460.—Malaya ; China.

FICUS :—
suffocans, *Banks, ex Griseb. Fl. Brit. W. Ind.* 150.—Jamaic.
sulcipes, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Mexic.
sumatrana, *Miq. l. c.* 287.—Sumatra.
sundaica, Blume, Bijdr. 450 = indica.
superba, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 200.—Java ; Japon.
superstitiosa, Link, Enum. Hort. Berol. ii. 449 = rellosa.
Sur, *Forsk. Fl. Aegypt. Arab.* 180.—Arabia.
surinamensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 219.—Guiana.
Suringari, *Carr. in Rev. Hortic.* (1866) 388.—Ins. Amboin.
sycomoroides, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 280 = variegata.
Sycomorus, *Linn. Sp. Pl.* 1059.—Afr. bor.
sylvatica, Regel, Cat. Pl. Hort. Alexakov. 58 = seq. ?
sylvestris, Hort. Berol. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 587 = splendens.
symphyllifolia, Spreng. Syst. iii. 784 = hispida.
syringaefolia, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 35.—Venezuela.
Taaf, *Forsk. Fl. Aegypt. Arab.* 219.—Arabia.
Tabing, Miq. Fl. Ind. Bat. Suppl. 430 = parietalis.
Tadjam, Miq. Fl. Jungh. 92 = platlera.
Taeda, Hort. Berol. ex Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 14 = elastica.
Tampang, *Miq. Fl. Ind. Bat. Suppl.* 425.—Sumatra.
tecoloides, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 299.—Mexic.
tenax, *Blume, Bijdr.* 440.—Java.
tengerensis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 283 = fistulosa.
tenuiramis, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 21 = cuspidata.
terebrata, Hort. Berol. ex Walp. Ann i. 677 = cestrifolia.
terebrata, Link, Enum. Hort. Berol. ii. 450 = arbutifolia.
terebrata, *Willd. Sp. Pl.* iv. 145.—Ins. Mascar.
terminaloides, Griff. Notul. iv. 393 = infectoria
terminalis, Heyne, ex Roth, Nov. Pl. Sp. 392 = gibbosa.
ternatana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 296.—Ins. Ternate.
terragena, Comm. ex Bory, Voy. ii. 148 = mauritiana.
Tetangit, Miq. Fl. Ind. Bat. Suppl. 432 = apiocarpa.
Teysmanniana, Miq. Fl. Ind. Bat. i. 11. 319 = alnoides.
theophrastoides, *Seem. Fl. Vit.* 252.—Ins. Fiji.
Thomaes, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 290.—Hab. !
Thomsoni, Miq. l. c. 215 = glaberrima.
Thonningiana, *Miq. in Hook. Niger Fl.* 522.—Afr. trop.
Thonningii, Blume, Rumphia, ii. 18 = microcarpa.
Thunbergii, Maxim. in Bull. Acad. Péterab. xxvii. (1881) 552 = foveolata.
Thwaitesii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 229.—Zeylan.
tiliaefolia, *Baker, in Journ. Linn. Soc.* xxi. (1885) 443.—Madag.
timoremola, Decne. in Nouv. Ann. Mus. Par. ii. (1834) 495 = superba.
timorensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 286.—Ins. Timor.
tinctoria, *Forst. f. Prod.* 76.—Ins. Societ.
Tjasela, Hochst. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 588 = F. Dekdekena.
Tjaskal, Hook. ex Miq. Fl. Ind. Bat. i. 11. 309 = rigida.
Tjaskela, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 287 = F. Tjaskela.
Tjiela, Miq. l. c. 286 = F. Tsiela.
Toka, *Forsk. Fl. Aegypt. Arab.* 219.—Arabia.
tomentosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297.—Java.
tomentosa, *Roxb. ex Willd. Sp. Pl.* iv. 1136.—Ind. or.
tomentosa, Herb. Madr. ex Wall. Cat. n. 4499 = myocomata.
Tondana, Miq. Fl. Ind. Bat. i. 11. 305 = platlera.
tonsa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 285.—Ins. Celebes.
Tortosana, Miq. l. c. 292 = heterophylla.
torulosa, Wall. Cat. n. 4550 = chartacea.
toxica, Thunb. Diss. Fic. 14.—toxicaria.
toxicaria, Herb. Madr. ex Miq. in Hook. Lond. Journ. Bot. vii. (1848) 461 = hispida.

FICUS :—
toxicaria, *Linn. Mant.* 305.—Malaya.
trachycarpa, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 450 = clavata.
trachycoma, Miq. in Zoll. Syst. Verz. Ind. Archip. 92 = glaberrima.
trachylela, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 268.—N. Caled.
trachyphylla, *Fenzl, in Flora*, xxvii. (1844), 311, nomen. —Afr. trop.
trematicarpa, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 224 = Decaisnei.
Tremblas, Miq. Fl. Jungh. 61, partim = pisifera.
trichocarpa, *Blume, Bijdr.* 448.—Java ; Ins. Timor.
trichocarpa, Herb. Par ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 288 = glomerata.
trichocarpa, Zipp. ex Miq. l. c. 228 = obtusa.
trichophlebia, *Baker, in Journ. Linn. Soc.* xxi. (1885) 445.—Madag.
trichopoda, *Baker, l. c.* xx. (1883) 261.—Madag.
trichopharra, *Baker, l. c.*—Madag.
tricolor, *Miq. Fl. Jungh.* 53.—Java.
tridentata, *Fenzl, in Flora*, xxvii. (1844) 311, nomen.—Afr. trop.
trigona, *Linn f. Suppl.* 441.—Guian.
trigonata, *Linn. Pl. Surin.* n. 145 ; *Amoen. Acad.* viii. 265 (- pranc.!).—Guian.
trilepis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 228 = nemoralis.
triloba, Buch.-Ham ex Wall. Cat. n. 4491 = hirta.
Trimeni, *King, ex Trim. in Journ. Bot.* xxiii. (1885) 242.—Zeylan.
trinervia, Reinw. ex Miq. Ann. Mus Bot Lugd. Bat. iii. 265 = glabella.
tripinnervis, Buch.-Ham. ex Wall. Cat. n. 4533 = scandens.
tristis, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Hab. ?
truncata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 292.—Borneo.
truncata, Vahl, Symb. Bot. i. 38 = heterophylla.
trymatocarpa, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 292 = tetrasperma.
Tsiela, *Roxb. Hort. Beng.* 66 ; *Fl. Ind.* iii. 549.—Ind. or.
Tsjakela, *Burm. f. Fl. Ind.* 227.—Ind. or.
tuberculata, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 283, in syn. = fistulosa.
tuberculata, Roxb. Hort. Beng. [103] ; Fl. Ind. iii. 554 = gibbosa.
tuberculata, Wall. Cat. n. 4539 = saemocarpa.
turbinata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 298.—Mexic.
turbinata, Willd. Sp. Pl. iv. 1141 = scabra.
Tweediana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 299.—Bras.
tylophylla, *Hassk. Cat. Hort. Bog. Alt.* 75.—Hab. ?
uimifolia, Hort. Bogor. ex Miq. Fl. Ind. Bat. i. 1. 315 = brevicapis.
uimifolia, *Lam. Encyc.* ii. 499.—Malaya.
umbellata, *Vahl, Enum.* ii. 189.—Afr. trop.
umbonata, *Reinw. ex Blume, Bijdr.* 454.—Malaya ; Ind. or.
umbonata, Wall. Cat. n. 4548 = silhetensis.
umbrifera, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 18 = F. Awi-Awi, *Blume.*
umbrosa, Salisb. Prod. 16 = indica.
undulata, Buch.-Ham. in Trans. Linn. Soc. xv. (1827) 138 = nervosa.
undulata, *Desf. Tabl. ed.* II. 289, nomen.—Ind. occ.
uniglandulosa, Wall. Cat. n. 4479 = roxburghii.
urnigera, Miq. in Zoll. Syst. Verz. Ind. Archip. 92, 98 = recurva.
urophylla, *Wall. Cat.* n. 4483.—Reg. Himal. ; Burma.
urticifolia, *Roxb. Hort. Beng.* [103] ; *Fl. Ind.* iii. 553.—Bengal.
vaguna, Roxb. ii. cc. 66 ; 587 = laevis.
vaguna, Wall. Cat. n. 4562 = ramentacea.
valida, Blume, Bijdr. 449 = annulata.
validinervis, *F. Muell. ex Benth. Fl. Austral.* vi. 166.—Austral.
Vallis-Choudae, *Delile, in Ann. Sc. Nat. Sér. II.* xx. (1843) 94.—Abyss.
Varenya, Buch.-Ham. ex Wall. Cat. n. 4546 = altissima.
variabilis, *Miq. Fl. Ind. Bat.* i. 11. 310.—Java.
variegata, *Blume, Bijdr.* 459.—Reg. Himal. ; Malaya.
variolosa, *Lindl. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 473.—China ; Malaya.

FICUS:—
vasculosa, *Wall. Cat.* n. 4482.—Burma; China; Malaya.
vasta, Forsk. Fl. Aegypt. Arab. 179 = populifolia.
velutina, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1141. —N. Granat.
venosa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 451 = F. Tsjakela.
venosa, Wall. Cat. n. 4515 D, F = infectoria.
venosa, Willd. Hort. Berol. 36. t. 36 = leucantatoma.
venusta, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846) 16.—Cuba.
vermifuga, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 300. —Bras.
verrucosa, *Hemsl. Biol. Centr. Am. Bot.* iii. 148.— Nicaragua.
verrucosa, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 295 = nemoralis.
verrucosa, Vahl, Enum. ii. 192 = septica.
versicolor, *Bur. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 256.—N. Caled.
vesca, F. Muell. ex Miq. in Journ. Bot. Néerl. (1861) 248 = glomerata.
vestita, Desf. Cat. Hort. Par. ed. III. 346, 413 = pumila.
vestita, *Wall. Cat.* n. 4500.—Ind. or.
Vieillardiana, *Bur. in Ann. Sc. Nat* Sér. V. xiv. (1872) 258.—N. Caled.
villipes, Miq. in Hook. Lond. journ. Bot. vii. (1848) 451 = recurva.
villosa, *Blume, Bijdr.* 441.—Malaya.
virens, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 451 = lanifolia.
virgata, Reinw. ex Blume, Bijdr. 454 = subulata.
virgata, Roxb. Hort. Beng. 65; Fl. Ind. iii. 530 = palmata.
virginiana, Hort. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 298 = ferruginea.
viscifolia, Hort. Berol. ex Kunth & Bouché, Ind Sem. Hort. Berol. (1846) 30 = diversifolia.
vitellina, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 288 = puberula.
vitiensis, *Seem. Fl. Vit.* 250.—Ins Fiji.
Vogeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 295. —Afr. trop.
Vogelii, *Miq. l. c.* 288.—Afr. trop.
volkameriæfolia, Wall. Cat. n. 4542 = lepicarpa.
volubilis, *Van Houtte, ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 300.—Hab.?
Vrieseana, *Miq. l. c.* 284.—Java.
Warscewiczii, *Miq. l. c.* 296.—Guatemala.
Wassa, Roxb. Hort. Beng. 65; Fl. Ind. iii. 539 = difformis.
Webbiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 297. —N. Caled.
Wightiana, Benth. Fl. Hongk. 327 = glabella.
Wightiana, Wall. Cat. n. 4540 = infectoria.
Wrightii, Benth. Fl. Hongk. 329 = foveolata.
xanthoneura, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 293 = obtusa.
xanthophylla, Stend. ex Miq. in Hook. Lond. Journ. Bot. vi. (1847) 554 = lutea.
xiphocospis, *Baker, in Journ. Linn. Soc.* xx. (1883) 260.—Malaya.
xylophylla, *Wall. Cat.* n. 4558.—Malaya.
yoponensis, *Desv. in Ann. Sc. Nat.* Sér. II. xviii. (1843) 310.—Venezuela.
zeylanica, Trim. Cat. Pl. Ceyl. 54 = laevis.
Zollingeriana, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 264 = obscura.

FIDELIA, Sch. Bip. in Flora, xvii. (1834) 482 = **Leontodon**, Linn. (Compos.).
balbisioides, Sch. Bip. l. c. = L. hispidulum.
Robouliana, Pomel, Nouv. Mat. Fl. Atl. 269.—Alger.
tricholepis, Pomel, l. c. 270 = L. hispidulum.

FIKERRA, Opiz, Seznam, 44 (1852) = **Chaerophyllum**, Linn. (Umbellif.).
nodosa, Opiz, l. c. = C. nodosum.

FIDLERIA, Reichb. Nom. 206 (1841); Ic. Fl. Germ. xvi. 42. t. 246 (1844) = **Turritis**, Scop. (Cruciferæ).
thyrsoidea, Reichb. Ic. Fl. Germ. xvi. 42. t. 246 = T. thyrsoidea.

FIELDIA, A. Cunn. in Field, N. S. Wales, 363. t. 2 (1826), *GESNERACEAE*, Benth. & Hook. f. ii. 1012.
BASILOPHYTA, F. Muell. First Gen. Rep. 16 (1853).
australis, *A. Cunn. l. c.* 364.—Austral.

FIELDIA, Gaudich. in Freyc. Voy. Bot. 424. t. 36 (1826) = **Stauropsis**, Reichb. (Orchid.).
gigantea, Reichb. f. Xenia Orch. ii. 89 = S. gigantea.
lancifolioides, Gaudich. in Freyc. Voy. Bot. 424. t. 36 = Vanda Batemanni.

FIERAURERA, Reichb. Consp. 86 (1828), sphalm. = **Fibraurea**, Lour. (Menisp.).

FIGARAEA, Viv. Fl. Aegypt. Dec. 11 (1830) = **Neurada**, Linn. (Rosac.).
aegyptiaca, Viv. l. c. 12 = N. procumbens.

FIGUIERREA, Montr. in Mém. Acad. Lyon, x. (1860) 220. *INC. SED.*, Benth. & Hook. f. ii. 29.
fragrans, *Montr. l. c.*—N. Caled.

FILAGINELLA, Opiz, Seznam, 44 (1852) = **Gnaphallum**, Linn. (Compos.).
luteo-alba, Opiz, l. c. = G. luteo-album.
nuda, Opiz, l. c. = G. uliginosum.
uliginosa, Opiz, l. c. = G. uliginosum.

FILAGINOPSIS, Torr. & Gray, Fl. N. Am. ii. 263 (1841) = **Evax**, Gaertn. (Compos.).
Drummondi, Torr. & Gray, l. c. = E. multicaulis.
multicaulis, Torr. & Gray, l. c. = E. multicaulis.

FILAGO, Linn. Syst. ed. I (1735); Sp. Pl. ed. I. 927 (1753), ex parte, et sect. plur. *COMPOSITAE*, Benth. & Hook. f. ii. 299.
ACHARTERIUM, Bluff & Fingerh. Comp. Fl. Germ. ii. 345 (1825).
GIFOLA, Cass. in Bull. Soc. Philom. (1819) 142.
IMPIA, Bluff & Fingerh. Comp. Fl. Germ. ii. 342 (1825).
LOGFIA, Cass. in Bull. Soc. Philom. (1819) 143.
OGLIFA, Cass. l. c. (1819).
XERETIUM, Bluff & Fingerh. Comp. Fl. Germ. ii. 343 (1825).
abyssinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 482.—Abyssin.
acaulis, Krock. Fl. Siles. ii. 498. t. 41 = Gnaphalium supinum.
affinis, Tineo, ex Nym Consp. 384 = spathulata.
alpestris, Presl, Delic. Prag. 100 = arvensis.
angustifolia, Schur, Enum. Pl. Transs. 917, Index = arvensis.
apiculata, G. E. Sm. in Phytolog. (1846) 575 = germanica.
arizonica, *A. Gray, in Proc. Am. Acad.* viii. (1878) 652.—Calif.
arvensis, *Linn. Sp. Pl. Add.*—Europ.; As. bor.
arvensis, Wahlenb. Fl. Suec. 560 = minima.
caespitosa, Rafin. ex DC. Prod. vi. 247 = germanica.
californica, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 406.—Am. bor. occ.
Candolleana, Parl. in Giorn. Tosc. Sc. Med. Fis. Nat. i. (1840) 180 = spathulata.
canescens, Jord. Obs. Fl. Crit. iii. (1846) 202 = germanica.
cephalioides, Link, ex Stend. Nom. ed. II. i 638 = Gnaphalium japonicum.
Clementei, Willk. in Bot. Zeit. v. (1847) 859 = marcotica.
cossyrensis, *Tenore, ex Lojac. in Natural. Sicil.* iv. (1885) 108.—Sicil.
Cupaniana, *Parl. in Giorn. Tosc. Sc. Med. Fis. Nat.* i. (1840) 182.—Sicil.
dasycarpa, *Grisb. in Gott. Abh.* xxiv. (1879 185.— Reg. Argent.
depressa, *A. Gray, in Proc. Am. Acad.* xix. (1883) 3. —Calif.
desertorum, *Pomel, Nouv. Mat. Fl. Atl.* 46.—Alger.
dichotoma, Gilib. Fl. Lituan. i. 181 = arvensis.
Duriaei, *Coss. ex Lange, in Kjoeb. Vidensk. Meddel.* (1861) 70.—Afr. bor.
eriocephala, Cass. Fl. Raz. Ste. t. 69 = germanica.
filiformis, Lam. Fl. Fr. ii. 61 = gallica.

FILAGO:—
floribunda, Cos. & Kral. ex Boiss. Fl. Orient. iii. 247 = marcotica.
fuscescens, *Pomel, Nouv. Mat. Fl. Atl.* 44.—Alger.
gallica, *Linn. Sp. Pl. Add.*—Europ.; As. bor.; Orient. germanica, *Linn. l. c.* ed. II. 1311.—Europ.; Orient.
heterantha, Rafin. Glorn. ii. fasc. xii. 170; Guss. Fl. Sic. ii. 864 = Cupaniana.
iodolepis, *Brth. Fl. Norm.* ed. IV. 168.—Gallia.
Fauxiana, Cos. & Germ. ex Ann. Sc. Nat. Sér. II. xx. (1843) 284. t. 13. fg. C. 1–8 = spathulata.
Kaltenbachii, Sch. Bip. ex Nym. Consp. 384 = germanica.
Lagopus, Steph. ex Willd. Sp. Pl. iii. 1897 = arvensis
lanuginosa, Requien, ex DC. Prod. vi. 247 = germanica.
lasiocarpa, *Griseb. in Gott. Abh.* xix. (1874) 180.— Reg. Argent.
leontopodioides, Willd. Phyt. 12 = Leontopodium sibiricum.
Leontopodium, Linn. Sp. Pl. 1212 = Leontopodium alpinum.
Leontopodium, Pall. Reise, iii. 265 = Leontopodium sibiricum.
lutescens, Jord. Obs. Pl. Crit. iii. (1846) 201 = germanica.
marecotica, *Delile, Fl. Egypt.* 274.—Aegypt.
maritima, Hill, Veg. Syst. iii. 141 = Diotis candidissima.
microcephala, *Pomel, Nouv. Mat. Fl. Atl.* 45.— Alger.
micropodioides, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 71.—Hispan.; Ins. Sicil.
minima, *Fries, Novit. Fl. Suec.* 228.—Europ.
mixta, Heluby, in Oestr. Bot. Zeitschr. xxi. (1871) 262 = germanica.
montana, DC. Prod. vi. 248 = minima.
montana, Linn. Sp. Pl. add. = arvensis
multicaulis, Lam. Fl. Fr. ii. 59 = Micropus erectus.
neglecta, *DC. Prod.* vi. 248.—Europ.
obovata, *Pomel, Nouv. Mat. Fl. Atl.* 45.—Alger.
panciulata, Gilib. Fl. Lituan. i. 181; Moench, Meth. 577 = arvensis.
parvula, Torr. & Gray, Fl. N. Am. ii. 432 = californica.
pilularis, Link, Handb. i. 720 = Gnaphalium uliginosum.
prolifera, *Pomel, Nouv. Mat. Fl. Atl.* 47.—Alger.
prostrata, DC. in Wight, Contrib. 22 = Gnaphalium pulvinatum.
prostrata, Pall. Pl. Nov. 11 = spathulata.
pygmaea, Linn. Sp. Pl. 927 = Evax umbellata.
pyramidata, *Linn. Sp. Pl.* 1199.—Hispan.; Orient.
pyramidata, C. A. Mey. Verz. Pfl. Cauc. 78 = germanica.
pyramidata, Vill. Hist. Pl. Dauph. iii. 194 = spathulata
ramosissima, Lange, Ind. Sem. Hort. Haun. (1856) 24 = marcotica.
recta, Link, Handb. i. 720 = Gnaphalium sylvaticum.
repens, *Schreb, in Linnaea*, xxii. (1849) 164 (as genera?).—Texas.
robusta, *Pomel, Nouv. Mat. Fl. Atl.* 45.—Alger.
rotundata, Moench, Meth. 577 = germanica.
spathulata, *Presl, Delic. Prag.* 99.—Europ.; Afr. bor.; Orient.
stellata, Lam. Fl. Fr. ii. 58 = Leontopodium alpinum.
subspicata, Bor. Fl. Centr. Fr. ed. II. n. 1057 = germanica.
supina, Lam. Fl. Fr. ii. 60 = Micropus supinus.
sylvatica, Link, Handb. i. 720 = Gnaphalium sylvaticum.
tenuifolia, Presl, Delic. Prag. 101 = gallica.
texana, *Scheele, in Linnaea*, xxii. (1849) 164 (as genera?).—Texas.
uliginosa, Link, Handb. i. 720 = Gnaphalium sylvaticum.
vulgaris, Lam. Fl. Fr. ii. 61 = germanica.

FILAGO, Loefl. in Linn. Gen. ed. V. 397 (1754) = **Evax**, Gaertn. (Compos.).
acaulis, Linn. Syst. ed. XII. 580 = E. umbellata.
atlantica, Ball, in Journ. Bot. xi. (1873) 264 = E. Heldreichii.
congesta, Guss. ex DC. Prod. vi. 246 = E. exigua.
discolor, Guss. l. c. v. 459 = E. discolor.
exigua, Sibth. & Sm. Fl. Graec. t. 923 = E. exigua.
pygmaea, Cav. Ic. i. 23. t. 36 = E. exigua.

FILETIA, Miq. Fl. Ind. Bat. ii 822 (1856). *ACANTHACEAE*, Benth. & Hook. f. ii. 1101.
costulata, *Miq. l. c.* 823.—Sumatra.

FILICIRNA, Rafin Fl. Tellur. iii. 37 (1836) = **Drosera**, Linn.
filiformis, Rafin. l. c. = D. filiformis.
leionema, Rafin. l. c. = D. leionema.
tenuifolia, Rafin. l. c. = D. filiformis.

FILICIUM, Thw. Enum. Pl. Zeyl. 403 (1864). *BURSERACEAE*, Benth. & Hook. f. i. 325.
PTERIDOPHYLLUM, Thw. in Hook. Kew Journ. vi. (1854) 65. t. 1.
decipiens. *Thw. l. c.*—Ind. or.

FILIPENDULA, Tourn. ex Linn. Gen. ed. I. 145 (1737) = **Spiræa**, Linn. (Rosac.).
angustiloba, Maxim. in Act. Hort. Petrop. vi. (1879) 250 = S. angustiloba.
hexapetala, Gilib. Fl. Lituan. ii. 287 = S. Filipendula.
kamtschatica, Maxim. in Act. Hort. Petrop. vi. (1879) 248 = S. camtschatica.
lobata, Maxim. l. c. 251 = S lobata.
multijuga, Maxim. l. c 247.—Japon.
palmata, Maxim. l. c. 250 = S. palmata. *Pall.*
pubescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi (1868) 370 = S Filipendula.
purpurea, Maxim. in Act. Hort. Petrop. vi. (1879) 248 = S. palmata, *Thunb*
Ulmaria. Maxim. l. c. 251 = S. Ulmaria.
vestita, Maxim. l. c. 248 = S. vestita.
vulgaris, Moench, Meth. 683 = S. Filipendula.

FILLAEA, Guill. & Perr. Fl. Seneg. Tent. i. 242. t. 55 (1833) = **Erythrophloeum**, Afzel. (Legumin.)
Brucei, Fresen. in Flora, xxii. (1839) 54 = Pterolobium lacerans.
suaveolens, Guill. & Perr. Fl. Seneg. Tent. i 242 t. 55 = E. guineense.

FIMBRIARIA, A. Juss. in St. Hil. Fl Bras. Mer. iii. 63. t 173 (1838) = **Schwannia**, Endl. (Malpig.).
cujabensis, Grisseb. in Linnæa, xxiii. (1850) 189 = S. elegans.
elegans, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 63. t. 173 = S. elegans.
januioides, A. Juss. in Walp. Rep. v. 353 = Janusia amazonica.
muricata, A. Juss. l. c. 354 = Janusia muricata.

FIMBRILLARIA, Cass. in Bull. Soc. Philom. (1818) 81; et in Dict. Sc. Nat. xvii. 54 (1820) = **Conyza**, Linn. (Compos.).
baccharoides, Cass. in Dict. Sc. Nat. xvii. 54 = C. ivæfolia, *Less.*
tubifera, Cass. l. c. 55 = C. laurifolia.

FIMBRISTEMMA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 320. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 793.
CALLAZOLEPIUM, Karst. Fl. Columb. i. 128. t. 165 (1858).
gonocloioides, *Turcz. l. c.*—N. Granat.
Warscewiczii, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 356.—Guatemala.

FIMBRISTILIS, Ritg. in Marb. Schrift. ii. (1881) 120 = **Fimbristylis**, Vahl (Cyperac.).

FIMBRISTIMA, Rafin. Fl. Tellur ii. 46 (1836) = **Aster**, Tourn. (Compos.).
flexuosa, Rafin. l. c. = A. tenuifolius.
maritima, Rafin. l. c. = A. Tripolium?
sparsiflora, Rafin. l. c. = A. tenuifolius?
squamosa, Rafin. l. c.—Hab.?

FIMBRISTYLES, Schult. Mant. ii. 2 (1824) = seq.

FIMBRISTYLIS, Vahl, Enum. ii. 285 (1806). *CYPERACEAE*, Benth. & Hook. f. iii. 1048.
ABILGARDIA, Vahl, Enum. ii. 296 (1806).
BULBOSTYLIS, Kunth, Enum. Pl. ii. 206 (1837).
CAMPYLOSTACHYS, E. Mey. Zwei Pfl. Docum. 83 (1848).

FIMBRISTYLIS:—
ECHINOLYTRUM, Desv. Journ. Bot. i. (1806) 20. t. 1.
GUSNOMEA, Presl. Rel. Haenk. i. 153. t. 33 (1828).
IRIA, Hedw. f. Gen. 360 (1906).
MISCHOSPORA, Boeck. in Flora, xliii. (1860) 113.
ONCOSTYLIS, Nees, in Mart. Fl. Bras. ii. i. 80 (1842).
PODOSTYLIS, Bertol. Fl. Ital. i. 512 (1833).
PSEUDO-CYPERUS, Steud. in Flora, xxxiii. (1850) 229, nomen; Syn. Pl. Cyp. 108, in nota (1855).
TRICHELOSTYLIS, Lesch. Ias. Fam. Cyp. 40 (1819).
abbreviata, Boeck. in Flora, xli. (1858) 601 = junciformis.
abyssinea, Steud. Syn. Pl. Cyp. 127 = tetragona.
aciortiva, *Steud. l. c.* 111.—Madag.
acicularis, R. Br. Prod. 226 = acuminata.
acuminata, F. Muell. Fragm. viii. 274 = rhyticarpa.
acuminata, *Vahl, Enum.* ii. 285.—Ind. or.; China; Austral.
adventitia, *Ces. in Linnæa*, xxxii. (1863) 261. t. 2.—Italia.
æstivalis, *Vahl, Enum.* ii. 288.—Aa. Austral. et Amb. trop.
affinis, Hook. & Arn. Bot. Beech. Voy. 72 = diphylla.
albescens, Steud. Syn. Pl. Cyp. 107 = polytrichoides.
albicans, *Nees, in Wight, Contrib.* 100.—Ind. or.
ambigua, Steud. Syn. Pl. Cyp. 117 = diphylla.
amblyphylla, Steud. Syn. Pl. Cyp. 116 = complanata.
ancepa, Steud. l. c. 113 = complanata.
andamandica, Kurz, Andaman Rep. App. B 21 = ferruginea.
andongensis, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 158.—Afr. trop.
androgyna, *R. Br. Prod.* 226.—Austral.
angularis, *Lesh. Enum. Hort. Beral.* 1. 289.—Ind. or.
annua, *Link, Handb.* i. 108, in obs.—Europ. austr.
annua, Roem. & Schult. Syst. ii. 95 = diphylla.
anomala, *Boeck. in Flora*, xliii. (1860) 240.—Texas.
aphylla, F. Muell. *Fragm.* ix. 4 = Arthrostylis aphylla.
aphylla, Steud. Syn. Pl. Cyp. 114 = quinquangularis.
aphyllanthoides, *Welw. ex Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 151.—Afr. trop.
apus, S. Wats in Bot. Calif. ii. 224 = Vahlii.
arenaria, *Nees, in Linnæa*, x. (1836) 146.—Afr. austr.
argentea, *Vahl, Enum.* ii. 294.—Ind. or.
Baldwiniana, *Torr. in Ann. Lyc. N. York*, iii. (1836) 344 = diphylla.
barbata, *Benth. Fl. Austral.* vii. 321.—Austral.
Barteri, *Boeck. in Linnæa*, xxxvii. (1871-73). 83.—Afr. trop.
benghalensis, Roem. & Schult. Syst. ii. 94 = miliacea.
biflora, Boeck. in Linnæa, xxxviii. (1874) 398 = cyperoides.
bisplicata, *Nees & Meyen, in Wight, Contrib.* 97.—Ind. or.; China.
bismaloides, Boeck. in Flora, xli. (1858) 603 = spathacea.
Boeckeleri, Steud. Syn. Pl. Cyp. 113 = quinquangularis, complanata.
Boivinianus, Steud. l. c. 110 = diphylla.
borbonica, *Steud. l. c.* 107.—Ins. Borbon.
brachyiorne, F. Muell. Fragm. i. 199 = cæspitosa.
brachyphylla, J. & C. Presl, Rel. Haenk. i. 251 = junciformis.
brachyphylla, Schult. Mant. iii. 530 = diphylla.
brevicalmis, Boeck. in Flora, xli. (1858) 643 = Vahlii.
brevifolia, R. Br. Prod. 223 = ferruginea.
brevifolia, J. & C. Presl, Rel. Haenk. i. 192 = junciformis.
brevifolia, Wall. ex Roem. & Schult. Syst. Mant. ii. 54 = diphylla.
brizoides, Sm. ex Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 78 = diphylla.
bromeliæfolia, Spreng. f. Tent. 2 = Diplasia karatæfolia.

FIMBRISTYLIS:—
Brownii, Benth. Fl. Austral. vii. 308 = leptoclada.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 144.—Japon.
Burchellii, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 98.—Afr. austr.
burmifolia, Vidal, Phan. Cuming. Philipp. 14, 156 = setacea.
cæsia, *Miq. Fl. Ind. Bat.* iii. 215.—Java.
cæspitosa, *R. Br. Prod.* 226.—Austral.
calcarata, Steud. Syn. Pl. Cyp. 117 = diphylla.
Candelabrum, Link. Jahrb. i. III. 80 = diphylla.
capillacea, Hochst. ex Steud. Syn. Pl. Cyp. 111 = capillaris.
capillaris, *A. Gray, Man. Bot. N. U. St.* ed. V. 567.—Am. bor.
capitata, *R. Br. Prod.* 223.—Austral.
capitata, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 288 (= *Isolepis rigescens*).—Ins. Maurit.
capitata, Zoll. Syst. Verz. Ind. Archip. ii. 61 = spathacea
cardiocarpa, *F. Muell. Fragm.* i. 194.—Austral.
cardiocarpa, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 154.—Afr. trop.
carinata, Nees, in Linnæa, ix. (1834) 290 = dichotoma
cariensis, Steud. Syn. Pl. Cyp. 113 = diphylla.
castanea, *Vahl, Enum.* ii. 292.—Am. bor.
cephalophora, F. Muell. Fragm. i. 196 = capitata.
Cephalotes, *Steud. Syn. Pl. Cyp.* 120.—Bras.
chaetorrhiza, Kunth, Enum. Pl. ii. 240 = junciformis.
chaetorrhiza, Thw. Enum. Pl. Zeyl. 349, partim = asperrima.
ciliolata, Steud. Syn. Pl. Cyp. 100 = spathacea.
cincta, Nees, in Wight, Contrib. 98 = diphylla.
cinerea, *Ridl. in Journ. Linn. Soc.* xx. (1883) 386.—Madag.
cinnamometorum, Hance, l. c. xlii. (1872) 132 = fusca.
cinnamometorum, Kunth, Enum. Pl. ii. 229 = cyperoides.
Cioniana, *Savi, in Mem. Valdarnes.* iii. (1842) 98.—Italia.
circinnata, Steud. Syn. Pl. Cyp. 116 = diphylla.
cœrcitata, *Schweinf. in Beck, Fl. U. St.* 428 = capillaris.
coleotricha, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii 508.—Abyssin.
collina, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 154.—Afr. trop.
comata, *Nees, in Wight, Contrib.* 102 = squarrosa.
communis, Kunth, Enum. Pl. ii. 234 = diphylla.
communis, Seem. Fl. Vit. 318 = complanata.
complanata, Benth. Fl. Hongk. 398 = Thomasonii.
complanata, *Link, Hort. Beral.* i. 292.—Amphig. trop.
compressa, *Boeck. in Linnæa*, xxxviii. (1874) 387.—Ind. or.
compressa, Roem. & Schult. Syst. ii. 100 = ferruginea.
compressa, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 307.—Ins. Philipp.
conferta, *A. Rich. Fl. Cub. Fanerog.* ii. 291.—Cuba.
confinis, Steud. Syn. Pl. Cyp. 115 = diphylla.
congesta, Torr. in Ann. N. York, iii. (1836) 345 = Vahlii.
connectens, Thw. Enum. Pl. Zeyl. 349 = complanata.
conifera, Reichb. ex Kunth, Enum. Pl. ii. 206 (= *Isolepis conifera*).—Guian.
consanguinea, Kunth, Enum. Pl. ii. 228 = complanata.
contexta, *Kunth, l. c.* 245.—Ar. austr.
coryoocarpa, *F. Muell. Fragm.* i. 197.—Austral.
crassipes, *Boeck. in Flora*, xli. (1858) 602.—Ind. or.
crassipes, *Boeck. in Linnæa*, xxxviii. (1874) 393.—Mexic.
curvifolia, Steud. Syn. Pl. Cyp. 116 = diphylla.
cylindrica, Vahl, Enum. ii. 296 = spathacea.
cylindrocarpa, Kunth, Enum. Pl. ii. 223 = tetragona.
cylindrostachya, Steud. Syn. Pl. Cyp. 119 = complanata, ferruginea.
cymosa, *R. Br. Prod.* 226.—Austral.; Polynes.
cyperoides, *R. Br. l. c.* 226.—Aa. et Austral. trop.
cyperoides, F. Muell. Fragm. ix. 11 = complanata, stieroceryx.
cyrtophylla, Miq. Fl. Ind. Bat. iii. 225 = ferruginea.
Dallachyi, F. Muell. ex Benth. Fl. Austral. vii. 309 = dictena.
decaphylla, Miq. Fl. Ind. Bat. iii. 227 = setacea.
debilis, *F. Muell. Fragm.* i. 198.—Austral.
debilis, Steud. Syn. Pl. Cyp. 109 = diphylla.
decora, Nees & Meyen, in Wight, Contrib. 101 = setacea.
decumbens, Boeck. in Flora, lviii. (1875) 116 = setacea.

FIMBRISTYLIS:—
denudata, *R. Br. Prod.* 237.—Austral.
depauperata, R. Br. l. c = diphylla.
depauperata, Nees, in Linnæa, ix. (1834) 290 = Nemäl.
dichotoma, Hook. f. Handb. N. Zeal. Fl. 308 = velata.
dichotoma, *Vahl, Enum.* ii. 287.—Europ austr.; Afr. bor.
Dichrichenii, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 505. —China.
diffusula, Steud. Syn. Pl. Cyp. 116 = diphylla.
digitata, *Boeck. in Flora*, lxi. (1878) 35.—Ind. or.
diphylla, *Vahl, Enum.* ii. 289.—Amphig. trop. et temp. dispacea, *Benth. & Hook. f. Gen.* iii. 1049.—Ind. or.; Afr. trop.
disticha, *Boeck. in Linnæa*, xxxviii. (1874) 393.— Ind. or.; Cochinch.
divaricata, *Ham. Prod. Pl. Ind. Occ.* 14.—Porto Rico.
Dregeana, Kunth, Enum Pl. ii. 232 = diphylla.
Drummondii, Boeck. in Flora, xli. (1858) 608 = castanea.
Drummondii, Boeck. in Linnæa, xxxvii. (1871–73) 21 = anomala.
Ecklonii, Nees, in Linnæa, ix. (1834) 290; x. (1836) 143 = squarrosa.
efoliata, Steud. Syn. Pl Cyp. 109 = globulosa.
elata, R. Br. Prod. 237 = denudata.
elongata, *R. Br.* l. c. 232 = diphylla.
elongata, Sieber, ex Presl, in Oken, Isis, xxi (1828) 268 = pilosa.
emarginata, Wight, in Wall. Cat. n. 3500 = millacea.
Engleotis, *Hance, in Journ. Linn. Soc.* xiii. (1873) 132.—China.
exigua, Boeck. in Engl. Bot. Jahrb. v. (1884) 506 = Scirpus cernua
exilis, *Rorn. & Schult. Syst.* ii. 98.—Afr. trop; Am. austr.
falcata, Kunth. Enum. Pl. ii. 239 = junciformis.
falcifolia, *Boeck. in Linnæa*, xxxvii (1871–73) 25.— Afr. trop
ferruginea, *Hook. & Arn. Bot. Beech. Voy.* 312.— Mexic.
ferruginea, *Vahl, Enum.* ii. 291.—Amphig. trop.
filifolia, *Boeck. in Linnæa*, xxxvii. (1871–73) 32.— Ind. or.
filiformis, Kunth, Enum Pl. ii. 221 = paucifiora.
filiformis, Thonn. ex Kunth, l. c. 267 = Lipocarpha filiformis.
firmula, Steud. Syn. Pl. Cyp. 111 = autumnalis.
flaccida, Steud l. c. 113 = millacea.
flaccidula, Zoll Syst. Verz. Ind. Archip. i. 61 = millacea.
flexuosa, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 155.—Afr. trop.
foliosa, Link, Hort. Berol. i. 288 = diphylla.
Forsteri, Steud. Syn. Pl. Cyp. 111 = autumnalis.
fulvescens, *Thw. Enum. Pl. Zeyl.* 434.—Zeylan.
furva, *R. Br Prod.* 238.—Austral.
fuscata, Steud. Syn. Pl. Cyp. 113 = diphylla.
fuscescens, Steud. Nom. ed. II. i. 610 = diphylla.
gaimardioides, Steud. in Lechl. Berb. Am. Austr. 56 = Scirpus semsubterraneus.
geminata, Kunth, Enum. Pl. ii. 247 = autumnalis.
glabra, Hochst. ex Steud. Syn. Pl. Cyp. 111 = tenera.
glauca, Vahl, Enum. ii. 288 = diphylla.
globulosa, *Kunth, Enum. Pl.* ii. 231.—Ind. or.; Malaya; China.
glomerata, Nees, in Linnæa, ix. (1834) 290 = spathacea.
Gorringiana, Steud. Syn. Pl. Cyp. 118 = diphylla.
gracilenta, *Hance, in Journ. Bot.* vi. (1868) 89.— China.
gracilis, Arn. ex Boeck. in Linnæa, xxxviii. (1874) 388 = compressa
gracilis, R. Br. Prod. 237 = diphylla.
gracilis, *Trim. Cat. Pl. Ceyl.* 102.—Zeylan.
Griffithiana, Steud. Syn. Pl. Cyp. 110 = aestivalis.
Griffithii, Boeck, in Flora, xlill. (1860) 241 = aestivalis.
Harnåei, A. Dietz. Syst. ii. 161 = junciformis.
Hamiltonii, *Steud. Syn. Pl. Cyp.* 119.—Porto Rico.
Hanaensea, Boeck. in Linnæa, xxxviii. (1874) 394 = rigidula.
Heyneri, Boeck. in Flora, xlill. (1860) 244 = diphylla.
Hildebrandtii, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 156 = exilis.
hirsuta, Hochst. ex Steud. Syn. Pl. Cyp. 116 = Pollocarya hispidula.
hirta, Kunth & Schak. Syst. ii. 99 = squarrosa.
hirtula, Mem. Prod. Fl. Ind. Occ. 14 = Hamiltonii.

FIMBRISTYLIS:—
hirtula, Vahl, Enum. ii. 286 = diphylla.
hispidula, Kunth, Enum. Pl. ii. 227 = exilis.
Hookeriana, *Boeck. in Linnæa*, xxxvii. (1871–73) 22. —Ind. or.
humilensis, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 154 = Burchellii.
Humboldtii, J. & C. Presl, Rel. Haenk. i 190 = diphylla.
hybrida, Rich. ex Steud. Nom. ed. II. i. 640 = diphylla.
inconstans, Steud. Syn. Pl. Cyp. 107 = castanea.
insignis, *Thw. Enum. Pl. Zeyl.* 349.—Zeylan.; Malaya, China.
Involubilis, *Steud. Syn Pl. Cyp.* 111.—Am. austr.
japonica, Sieb. & Zucc. ex Steud. l. c. 107 = subbispicata.
juncea, Boeck. in Linnæa, xxxvii. (1871–73) 4 = poly trichoides.
juncea, *R. Br. ex Roem. & Schult. Syst.* ii. 102.—Ins. Pacif.
junciifolia, J. & C. Presl, Rel. Haenk. i. 190 = diphylla.
junciformis, *Kunth, Enum. Pl.* ii. 239.—Ind. or., Madag; Ins. Philipp
junciformis, Munro, in Seem. Bot. Voy Herald, 422 = spathacea.
juncoepchala, *Boeck. in Flora*, xli. (1858) 600.—Bras
Kamphoeveneri, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 506.—Ins. Nicobar
Kirkmaieri, Steud. Syn. Pl. Cyp. 118 = complanata
Kraussiana, Hochst in Flora, xxviii. (1845) 707 = complanata.
Kunthiana, *Ridl. in Trans. Linn. Soc.* Ser. II ii. (1884) 151.—Afr trop.
laevissima, Steud. Syn. Pl. Cyp. 117 = spathacea.
lanata, *Ker & Schult Syst.* ii. 92.—Venezuela
latifolia, Kunth, Enum. Pl. ii. 249 = junciformis.
laxa, Vahl, Enum. ii. 292 = diphylla.
leiocarpa, Maxim. Prim. Fl. Amur 301 = aestivalis
leiocarpa, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 144 = schoenoides.
leptoclada, *Benth. Fl. Hongk.* 398.—Hongkong; Hainan; Zeylan.
leptostachya, Steud. Syn. Pl. Cyp. 118 = autumnalis.
leucocoles, *Benth. Fl. Austral.* vii. 304.—Austral
leucostachya, *Boeck. in Linnæa*, xxxvii. (1874) 385 — Austral.
leucostachya, *Roem & Schult Syst.* ii. 92.—Venezuela
Lichtensteiniana, *Hemsl. Hot. Voy. Challenger*, i. 11. 86.—Ins. S. Helena.
limosa, *Poepp. & Kunth, in Kunth, Enum Pl.* ii. 225.—Am. austr.
Linniana, Steud Syn. Pl. Cyp. 112, sphalm. = exilis
littoralis, Gaudich. in Freyc. Voy. Bot. 513 = millacea.
lomentocarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii 504 = ferruginea.
longiculmis, *Steud. Syn. Pl. Cyp.* 110.—Afr. trop.; Madag.
longispica, *Steud. Syn. Pl. Cyp.* 118.—China; Japon.
Ludwigii, Steud. in Flora, xli. (1858) 189 = Scirpus Ludwigii.
macassarensis, *Steud. Syn. Pl. Cyp.* 109.— Malaya.
macra, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 150.—Afr. trop.
macrantha, Boeck. in Linnæa, xxxviii. (1874) 388 = monostachya.
macrostachya, *Boeck. l. c.* 386.—Austral.
madagascariensis, *Boeck. in Bremen, Abh.* vii. (1880) 38.—Madag.
malaccana, Boeck. in Flora, xli. (1858) 597 = paucifiora.
marginata, Labill. Sert. Austro-Caled. i 16 = ferruginea.
margaritana, Steud. Syn. Pl. Cyp. 107 = junces.
mariana, Gaudich. in Freyc. Voy. Bot. 513 = diphylla, cymosa.
Mortii, Boeck. in Linnæa, xxxvii. (1871–73) 24 = bahiensis.
mauritiana, Nees, ex Kunth, Enum. Pl. ii. 235 = diphylla.
mauritiana, Tausch, ex Presl, in Oken, Isis, xxi. (1828) 268 = ferruginea.
megastachya, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 156.—Afr. trop.
melanocephala, *Ridl. l. c.* 151.—Afr. trop.
melanostachya, Bronga. in Duperr. Voy. Bot. 188 = Elecocharis melanostachya.
Metzii, Steud. Syn. Pl. Cyp. 117 = diphylla.
mesostachya, Link, Hort. Berol. i. 290 = diphylla.

FIMBRISTYLIS:—
Michelisana, Reichb. Fl. Germ. Excurs. 75 = Scirpus Michelisanus.
Michelii, Bulami, Dodecanth. 34 = squarrosa.
microcarya, *F. Muell. Fragm.* i. 200.—Austral.
microstachya, Steud. Syn. Pl. Cyp. 113 = autumnalis.
miliacea, *Vahl, Enum.* ii. 287.—Amphig. trop.
mollis, *Kunth, Enum. Pl.* ii. 246.—Reg. Himal.
monandra, *F. Muell. Fragm.* i. 195.—Austral.
monandra, Koem. & Schult. Syst. Mant. ii. 59 = barbata.
Monnii, Steud. Syn. Pl. Cyp. 120 = obtusifolia.
monostachya, *Hassk. Pl. Jav. Rar.* 61.—Amphig. trop.
monticola, *Hochst. ex Steud. Syn. Pl. Cyp.* 111.— Ind. or.
mucronata, Boeck. in Linnæa, xxxvii. (1871–73) 41 = Barteri.
mucronata, Vahl, Enum. ii. 293 = Scirpus littoralis.
multifolia, Boeck. in Linnæa, xxxvii. (1874) 397 = cymosa.
multistriata, Boeck. in Flora, xliii. (1860) 243 = castanea.
muricatula, Steud Syn. Pl. Cyp. 113 = tenera.
muriculata, Benth. ex Hook. Niger Fl. 594 = tenera.
nana, Roth, Nov. Pl. Sp. 25 = squarrosa.
neglecta, *Hemsl. Bot. Voy. Challenger*, i. 11. 86.— Ins. S. Helena.
Neesii, *Kunth, Enum. Pl.* ii. 247, nomen.—Hab. ?
Nelsonii, *F. Muell. Fragm.* ix. 79.—Austral.
nesiotis, *Hemsl. Bot Voy. Challenger*, i. 11. 180.—Ins S. Trinit.
nigrescens, *Steud. Syn Pl. Cyp.* 118.—Guiana.
nigro-lurulnea, *Thw. Enum Pl Zeyl* 434.—Ind. or
nitidula, Roem. & Schult. Syst. Mant. ii. 61 = diphylla.
Novae-Britanniae, *Boeck. in Engl. Bot. Jahrb.* v. 1884) 93.—N. Britannia.
nuda, Boeck. in Flora, lviii. (1875) 110 = setacea.
nukahiwensis, Steud Syn. Pl. Cyp. 117 = diphylla.
nutans, *Vahl, Enum.* ii. 285.—As. et Austral. trop.
obtusangula, *F. Muell.* Fragm. i. 198 = rara.
obtusifolia, *Kunth, Enum. Pl.* ii. 240.—Afr. et Am. trop.
obtusifolia, Nees, ex Presl, in Oken, Isis, xxi. (1828) 268 = diphylla.
ochreata, Boeck. in Flora, xli. (1858) 599 = ferruginea.
oligostachys, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 508 = exilis.
ontrophes, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 155.—Afr. trop.
ovalis, Nees, in Wight, Contrib. 98 = diphylla.
oxylepis, Steud. Syn. Pl. Cyp. 110 = tenera.
oxyrhachis, Miq. Fl. Ind. Bat. Suppl. 601 = polytrichoides.
oxystachya, *F. Muell. Fragm.* i. 195.—Austral.
pallescens, Link, Hort Berol. i. 291 = diphylla.
pallescens, Nees, in Wight, Contrib. 101 = duchotoma.
parva, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 152.—Afr. trop.
parviflora, R. Br. Prod. 227 = diphylla.
pauciflora, *R. Br. l. c.* 235.—As. et Austral. trop.
paucispicata, F. Muell. Fragm. i. 197 = ferruginea.
paupercula, *Boeck. in Linnæa*, xxxviii. (1874) 396.— Ind. or.
pentaptera, *Kunth, Enum. Pl.* ii. 229.—Zeylan.
pentastachya, Boeck. in Flora, xl. (1857) 36 = diphylla.
philippica, Steud. Syn. Pl. Cyp. 116 = diphylla.
Pierotii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 145.— Ind. or.; Japon.
pilosa, *Vahl, Enum.* ii. 290.—Afr. trop.
planiculmis, Boeck. l. c. 391 = Arthrostylis aphylla.
planifolia, *Steud. Syn. Pl. Cyp.* 118.—Ins. Carolin.
platystachya, Boeck. in Linnæa, xxxviii. (1874) 390 = Schultzii.
podocarpa, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 77.—As. trop.
Pohlana, Steud. Syn. Pl. Cyp. 119 = diphylla.
polymorpha, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 141 = diphylla.
polytrichoides, *R. Br. Prod.* 226.—Geront. trop.
Presiii, Kunth, Enum. Pl. ii. 228 = exilis.
prolifera, *Steud. Syn. Pl. Cyp.* 120.—Guiana.
propinqua, R. Br. Prod. 227 = squarrosa.
pterigosperma, *R. Br. l. c.* 236.—Austral.
puberula, Vahl, Enum. ii. 289 = diphylla.
pubescens, Link, Jahrb. i. 111. 80 = diphylla.

FIMBRISTYLIS:—
pubigera, Link, Hort. Berol. i. 289 = dichotoma.
pulchella, *Trim. Cat. Pl. Zeyl.* 109.—Zeylan.
pumila, Benth. in Hook. Lond. journ. Bot. ii. (1843) 289 = pauciflora.
punctata, *R. Br. Prod.* 226.—Austral.
punctiscula, *Nees, in Linnaea*, iv. (1834) 90, nomen.—Hab.?
pusilla, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 506.—Abyssin.
pycnostachya, Hance, in Journ. Bot. xv. (1877) 338 = nigrobrunnea.
pygmaea, Vahl, Enum. ii. 285 = Scirpus cernuus.
quadrangularis, A. Dietr. ex Steud. Syn. Pl. Cyp. 114 = millacea.
quaternella, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 182.—Afr. trop.
quinquangularis, Benth. Fl. Austral. vii. 317 = corynocarya.
quinquangularis, *Kunth, Enum. Pl.* ii. 229.—Geront. trop.
rara, Benth. Fl. Austral. vii. 316 = salbundia, quinquangularis.
rara, *R. Br. Prod.* 227.—Austral.
retusa, Thw. Enum. Pl. Zeyl. 349 = leptoclada.
Reunionis, Steud. Syn. Pl. Cyp. 114 = spathacea.
rhyticarya, *F. Muell. Fragm.* i. 215.—Austral.
Richeana, Steud. Syn. Pl. Cyp. 114 = castanea.
rigida, *Kunth, Enum. Pl.* ii. 231 = spathacea.
rigidula, *Nees, in Wight, Contrib.* 99.—Ind. or.; China.
rigidula, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 150 = obtusifolia.
Roxburghii, A. Dietr. Sp. Pl. ii. 162 = ferruginea.
Royeniana, Nees, in Wight, Contrib. 99 = diphylla.
Rudgeana, Steud. Syn. Pl. Cyp. 115 = complanata.
rufa, Boeck. in Engl. Bot. Jahrb. v. (1884) 93 = rara.
salbundia, *Kunth, Enum. Pl.* ii. 230.—Ind. or. et Austral.
salbundia, Thw. Enum. Pl. Zeyl. 349 = pentaptera.
sansibariensis, Boeck. in Flora, lxiii. (1880) 437 = longiculmis.
scabberrima, *Nees, in Wight, Contrib.* 102.—Ind. or.
scabra, Kunth, Enum. Pl. ii. 245 = complanata.
scabra, Roem. & Schult. Syst. Mant. ii. 59 = acuminata.
scabrida, *Schum. & Thonn. Beskr. Guin. Pl.* 82.—Afr. trop.
Scherardi, Bubani, Dodecanth. = diphylla.
Schimperiana, *Boeck. in Flora*, xli. (1858) 600.—Abyssin.
schoenicides, *Vahl, Enum.* ii. 286.—As. et Austral. trop.
Schomburgkiana, *Steud. Syn. Pl. Cyp.* 111.—Guiana.
Schultzii, *Boeck. in Linnaea*, xxxvii. (1874) 891.—Austral.
Schweinfurthiana, *Boeck. in Flora*, lxii. (1879) 565.—Afr. trop.
seckelinsis, Baker, Fl. Maurit. 418 = diphylla.
separanda, *Steud. ex Yard. in Mém. Soc. Sc. Nat. Cherb.* v. (1857) 299, nomen.—Ins. Marquesas.
series, *R. Br. Prod.* 228.—As. et Austral. trop.
serrulata, Vahl, Enum. ii. 285 = diphylla.
setacea, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 289.—Malaya; Austral.
setifolia, *A. Rich. Tent. Fl. Abyss.* ii. 507.—Abyssin.
Sieberiana, Kunth, Enum. Pl. ii. 237 = ferruginea.
Sieboldii, *Miq. Cat. Mus. Bot. Lugd. Bat.* 116, nomen.—Japon.
similis, Steud. Syn. Pl. Cyp. 112 = diphylla.
solidifolia, *F. Muell. Fragm.* i. 198.—Austral.
sororia, Kunth, Enum. Pl. ii. 246 = diphylla.
spadicea, *Vahl, Enum.* ii. 294.—Am. austr.
Spartium, *Ham. Prod. Pl. Ind. Occ.* 14.—Porto Rico.
spathacea, *Roth, Nov. Pl. Sp.* 24.—Amphig. trop.
speciosa, Rohde, ex Spreng. Pugill. i. 5 = spadicea.
sphaerocephala, Benth. Fl. Austral. vii. 306 = monandra.
spiralis, *R. Br. Prod.* 226.—Austral.
spirostachya, F. Muell. ex Benth. Fl. Austral. vii. 311 = diphylla.
squamulosa, Hochs. ex Boeck. in Linnaea, xxxvii. (1873-73) 15 = diphylla.
squarrosa, *Vahl, Enum.* ii. 289.—Amphig. temp. et trop.

FIMBRISTYLIS:—
squarrulosa, *F. Muell. Fragm.* i. 216.—Austral.
stenia, Rich. ex Spreng. Pugill. i. 6 = ferruginea.
Stauntoni, *Debeaux & Franch. in Act. Linn. Soc. Bord.* xxxi. (1877) t. 3; et xxxii. (1878) 33.—China.
sterilis, Nees, in Linnaea, iv. (1834) 290 = spadicea.
striata, Ledeb. Sert. Austro-Caled. i. 16. f. 2 = diphylla.
stricta, R. Br. Prod. 228 = diphylla.
strictior, Steud. Syn. Pl. Cyp. 119 = Schomburgkiana.
subaphylla, *Boeck. in Flora*, lxii. (1879) 565.—Afr. trop.
subcristata, *Benth. Fl. Austral.* vii. 314.—Austral.
subbispicata, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 75.—As. trop.
subbulbosa, Benth. Fl. Austral. vii. 305 = schoenoides.
subbulbosa, Boeck. in Flora, xli. (1858) 598 = polytrichoides.
subalteralis, Steud. Syn. Pl. Cyp. 114 = castanea.
substrictastachya, Boeck. in Linnaea, xxxvii. (1871-72) 50 = nigrobrunnea.
substristachya, Steud. Syn. Pl. Cyp. 109 = diphylla.
sulcata, Roem. & Schult. Syst. Mant. ii. 52 = Elecharis sulcata.
tenella, Schnlt. l. c. 59.—Bras.
tenera, *Roem. & Schult. Syst. Mant.* ii. 57.—Geront. trop.
tenuicula, *Boeck. in Linnaea*, xxxviii. (1874) 385.—Ind. or.
tenuifolia, Thw. Enum. Pl. Zeyl. 424 = monticola.
tenuis, *Roem & Schult. Syst.* ii. 94.—Venezuela.
tenuissima, Steud. Syn. Pl. Cyp. 109 = barbata.
tertia, Steud. ex Jard. in Mém. Soc. Sc. Nat. Cherb. v. (1857) 299, nomen = diphylla.
tetragona, *R. Br. Prod.* 226.—As. et Austral. trop.
tetragona, A. Dietr. Sp. Pl. ii. 152 = millacea.
tetragona, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 141 = quadrangularis.
textilis, Roxb. in Beatson, St. Hel. Tracts, 309 = Scirpus nodosus.
thermalis, *S. Wats. in Bot. King's Exped.* 360.—Calif.
Thomsoni, *Boeck. in Linnaea*, xxxvii. (1871-73) 37.—Ind. or.; China.
Thunbergiana, *Boeck. l. c.* xxxviii. (1874) 395.—Afr. trop.
Thwaitesii, Boeck. l. c. xxxvi. (1871-73) 34 = insignis.
tomentosa, Vahl, Enum. ii. 290 = diphylla.
Torresiana, Gaudich. in Freyc. Voy. Bot. 413 = globulosa.
torta, Kunth, Enum. Pl. ii. 241 = junciformis.
trachycarya, F. Muell. Fragm. i. 199 = salbundia.
tricholepis, Miq. Fl. Ind. Bat. iii. 319 = acstivalis.
trifida, *Trim. Cat. Pl. Zeyl.* 102.—Zeylan.
trifolia, Nees, in Linnaea, iv. (1834) 290 = diphylla.
trifurca, Nees, in Mart. Fl. Bras. ii. 1. 74 = diphylla.
trigastrocarya, *F. Muell. Fragm.* i. 194.—Austral.
trispicata, Steud. Syn. Pl. Cyp. 107 = ferruginea.
tristachya, R. Br. Prod. 227 = ferruginea.
tristachya, Nees, ex Kunth, Enum. Pl. ii. 235, partim = quinquangularis.
tristachya, *Thw. Enum. Pl. Zeyl.* 434.—Ind. or.; Afr. trop.
uliginosa, *Hochst. ex Steud. Syn. Pl. Cyp.* 109.—Ind. or.
umbellaris, Vahl, Enum. ii. 291 = globulosa.
umbellata, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 73 = spadicea.
umbellato-capitata, Steud. Syn. Pl. Cyp. 120 = obtusifolia.
Urvilleana, *Steud. l. c.* 112.—Hab.?
Vahlii, *Link, Hort. Berol.* i. 287.—Am. trop.
variabilis, R. Br. Prod. 226 = diphylla.
velata, R. Br. l. c. 227 = squarrosa.
velutina, Franch. in Bull. Soc. Bot. Fr. xxvi. (1879) 88 = setacea.
verrucosa, J. & C. Presl, Rel. Haenk. i. 190 = diphylla.
vestita, *Hemsl. Biol. Centr. Am. Bot.* 460.—Mexic.
vestita, Steud. Syn. Pl. Cyp. 113 = exilis.
vexata, Steud. l. c. 316 = Ficinia angustifolia.
villosa, Roem. & Schult. Syst. ii. 98 = diphylla.
villosissima, *Steud. Syn. Pl. Cyp.* 119.—Bras.
Vincentii, Steud. l. c. 109 = Vahlii.
Wightiana, Nees, in Wight, Contrib. 99 = spathacea.
Xyridis, *R. Br. Prod.* 226.—Austral.
yprieides, Arn. ex Thw. Enum. Pl. Zeyl. 346 = tetragona.

FIMBROLINA, Rafin. Sylva Tellur. 71 (1838) = Beckeria, Linn. (Acanthac.).
incarnata, Rafin. l. c. = B. incarnata.

FIMBRYSTYLIS, D. Dietr. Syn. Pl. i. 129 (1839) = Fimbristylis, Vahl (Cyperac.).

FINCKEA, Klotzsch, in Linnaea, xii. (1838) 287 = Griesebachia, Klotzsch (Ericac.).
brunieades, Klotzsch, l. c. 288 = G. brunieades.
eriocephala, Klotzsch, l. c. = G. eriocephala.

FINDLAYA, Bowdich, Madeira, 258 (1825) (an Primulac.?).
alba, Bowdich, l. c.—Ins. Madera.

FINDLAYA, Hook. f. in Benth. & Hook f. Gen. ii. 589 (1876). *VACCINIACEÆ*, Benth. & Hook. l. l. c.
apophysata, *Hook. f. l. c.*—Ind. occ.

FINGALIA, Schrank, in Syll. Ratisb. i. (1824) 87 = Eleutheranthe, Poit. (Compos.).
hexagona, Schrank, l. c. = E. ovata.

FINGERHUTHIA, Nees, ex Lehm. Cat. Sem. Hort. Hamb. (1834):—Cf. Linnaea, x. (1836) Litt. 112. *GRAMINEÆ*, Benth. & Hook. f. iii. 1193.
LASIOTRICHOS, Lehm. Cat. Sem. Hort. Hamb. (1834).—Cf. Linnaea, x. (1836) Litt. 113.
affghanica, *Boiss. Fl. Orient.* v. 569.—Afghan.
africana, Aitch. in Journ. Linn. Soc. xix. (1882) 193 = affghanica.
africana, *Lehm. Cat. Sem. Hort. Hamb.* (1834).—Afr. austr.
capensis, Nees, ex Lehm. l. c. (1835) = africana.
cilitia, Nees, Fl. Afr. Austr. i. 136 = africana.
sesleriaeformis, *Nees, l. c.*—Afr. austr.

FINLAYSONIA, Wall. Pl. As. Rar. ii. 48. t. 162 (1831). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 743.
GURUA, Buch.-Ham. ex Wight, Contrib. 65 (1834). obovata, *Wall. l. c.*—Ind. or.; Malaya.

FINTELMANNIA, Kunth, Enum. Pl. ii. 362 (1837). *CYPERACEÆ*, Benth. & Hook. f. iii. 1068.
restioides, *Kunth, l. c.* 363.—Bras.
setifera, *Ridl. in Journ. Linn. Soc.* xx. (1883) 337.—Madag.

FIORINIA, Parl. Fl. Ital. i. 232 (1848) = Aira, Linn. (Gramin.).
pulchella, Parl. l. c. 233 = A. intermedia.

FIRENSIA, Scop. Introd. 157 (1777) = Cordia, Linn. (Boragin.).
fusca, Rafin. Sylva Tellur. 40.—Hab.?
hirsuta, Rafin. l. c. = C. nodosa.
lutea, Rafin. l. c. = C. rotundifolia.

FIRENZIA, DC. Prod. ix. 471 (1845) = prsec.

FIRKEA, Rafin. Sylva Tellur. 163 (1838) = Clusia, Linn. (Guttif.).
pedunculata, Rafin. l. c. 164 = C. pedicellata?
rosea, Rafin. l. c. 163 = C. rosea.
sessilis, Rafin. l. c. 164 = C. sessilis.
venosa, Rafin. l. c. 163 = C. venosa.

FIRMIANA, Marsigli, in Saggi Padov. i. (1786) 106, cum ic. = Sterculia, Linn.
chinensis, Medic. ex Steud. Nom. ed. I. 514, in syn. = S. platanifolia.
colorata, R. Br. in Benn. Pl. Jav. Rar. 235 = S. colorata.
diversifolia, A. Gray, in Bot. U. St. Expl. Exped. i. 185 = S. diversifolia, *Seem.*
heterophylla, Pasq. Cat. Orto Bot. Nap. 46 = Cola heterophylla.
platanifolia, Schott & Endl. Melet. 33 = S. platanifolia.

FISCHERA, Spreng. Pl. Umb. Prod. 27 (in Neue Schr. Naturf. Ges. Halle, ii.) (1813) = Platycnos, Bunge (Umbellif.).
canescens, Spreng. l. c. = Azorella canescens.
compressa, Spreng. l. c. (= *Sickera compressa*).—Austral.

FISCHERA:—
lancelata, Sm. in Rees, Cyclop. xxxix. n. 1 (= *Siebera Billardiera*).—Austral.
lanceolata, Spreng. Pl. Umb. Prod. 27 = Trachymene lanceolata.
linarifolia, Spreng. l. c. = Trachymene linearis.
linearis, Sm. in Rees, Cycl. Suppl. (= *Siebera linearifolia*).—Austral.
ovata, Spreng. Pl. Umb. Prod. 27 = Trachymene ovalis.

FISCHERA, Sw. in Mém. Soc. Nat. Mosc. v. (1817) 14. t. 1 = **Leiophyllum**, Hedw. f. (Umbellif.).
buxifolia, Sw. l. c. 17 = L. buxifolium.

FISCHERIA, DC. Cat. Hort. Monsp. 112 (1813). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 750.
acuminata, *Decne. in DC. Prod.* viii. 600.—Bras.
adenophylla, *Fourn. in Mart. Fl. Bras.* vi. iv. 301.—Bras.
calycina, *Decne. in DC. Prod.* viii. 600.—Am. trop.
cincta, Griseb. Fl. Brit. W. Ind. 421 = Gonolobus cinctus.
havanensis, *Decne. in DC. Prod.* viii. 601.—Cuba.
heterophylla, *Hemsl. Biol. Centr. Am. Bot.* ii. 320.—Nicaragua.
Hilariana, *Fourn. in Mart. Fl. Bras.* vi. iv. 299.—Bras.
macrocarpa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 68. t. 278.—Peruv.
Martiana, *Decne. in DC. Prod.* viii. 601.— Nicaragua.
multiflora, *Decne. l. c.*—Bras.
peruviana, *Decne. l. c.*—Peruv.
polytricha, *Decne. l. c.* 600.—Bras.
propinqua, *Decne. l. c.*—Bras.
Riedelii, *Fourn. in Mart. Fl. Bras.* vi. iv. 299.—Bras.
rotundifolia, *Decne. in DC. Prod.* viii. 601—Bras.
scandens, *DC. Cat. Hort. Monsp.* 112.—Am. trop.
stellata, *Fourn. in Mart. Fl. Bras.* vi. iv. 301.—Bras.
Warmingii, *Fourn. l. c.* 299.—Bras.

FISQUTIA, Gaudich. Bot. Voy. Bonite, t. 4, 5 (1844–53) = **Pandanus**, Linn.
macrocarpa, Gaudich. l. c. t. 4 = P. foetidus.
militaris, Gaudich. l. c. t. 5 = P. militaris.
ornata, Gaudich. l. c. t. 5 = P. ornatus.
ovata, Gaudich. l. c. t. 4 = P. ovatus.

FISSENIA, R. Br. ex Endl. Gen. Suppl. ii. 76 (1842) sphalm. = **Kissenia**, R. Br. (Lossac.).
arabica, R. Br. ex Presl, Bot. Bemerk. 78 = K. spathulata.
capensis, R. Br. ex Walp. Rep. ii. 238 = K. spathulata.
mentzeloides, R. Br. ex Harv. Thes. Cap. i. 61 = K. spathulata.

FISSICALYX, Benth. in Journ. Linn. Soc. v. (1861) 79. *LEGUMINOSAE*, Benth. & Hook. f. i. 562.
Fendleri, *Benth. l. c.*—Venezuela.

FISSILIA, Comm. ex Juss. Gen. 260 (1789) = **Olax**, Linn.
psittacorum, Blanco, Fl. Filip. ed. I. 98 = O. imbricata.
psittacorum, Lam. Illustr. Gen. i. 102. t. 98 = O. scandens.

FISSISTIGMA, Griff. Notul. ii. 706 (1854) = **Melodorum**, Dun. (Anonac.).
scandens, Griff. l. c. = M. Griffithii.

FITCHIA, Hook. f. in Hook. Lond. Journ. Bot. iv. (1845) 640 t. 23. *COMPOSITAE*, Benth. & Hook. f. ii. 506.
nutans, *Hook. f. l. c.*—Ins. Elizabeth.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 49.—Ins. Tahiti.

FITCHIA, Meissn. in Hook. Kew Journ. vii. (1855) 75 = **Orevillea**, R. Br. (Proteac.).

FITTONIA, E. Coem. in Fl. des Serres, xv. (1865) 185. *ACANTHACEAE*, Benth. & Hook. f. ii. 1137.
argyroneura, *E. Coem. l. c.* xvi. (1866–67) 108.—Peruv.
gigantea, *Linden, in Rev. Hortic.* (1866) 186.—Peruv. Verschaffeltii, *E. Coem. in Fl. des Serres*, xv. (1865–65) 184.—Peruv.

FITZALANIA, F. Muell. Fragm. iv. 33 (1863) = **Uvaria**, Linn. (Anonac.).
heteropetala, F. Muell. l. c. 34 = U. heteropetala.

FITZGERALDIA, F. Muell. Fragm. vi. 1 (1867). *ANONACEAE*.
nitrastigma, *F. Muell. l. c.*—Austral.

FITZGERALDIA, F. Muell. in Wing, South Sc. Record (1892), 56, in obs. = **Lyperanthus**, R. Br. (Orchid.).
Forrestii, F. Muell. l. c. = L. Forrestii.

FITZROYA, Hook. f. in Journ. Hort. Soc. vi. (1851) 264. *CONIFERAE*, Benth. & Hook. f. iii. 425.
CUPRESSTELLATA, Senills, [Nelson] Pinaceae, 60 (1866).
DIELMA, Hook. f. Fl. Tasman. i. 258. t. 98 (1859). Archeri, *Benth. & Hook. f. Gen.* iii. 426.—Tasman.
patagonica, *Hook. f. ex Lindl. in Journ. Hort. Soc.* vi. (1851) 264.—Reg. Magellan.

FIVA, Steud. Nom. ed. II. i. 642 (1840) = FIWA, J. F. Gmel. = **Litsea**, Lam. (Laurin.).

FIVALDIA, Walp. Rep. ii. 587, sphalm. (1843) = FRIWALDIA, Endl. = **Microglossa**, DC. (Compos.).

FIWA, J. F. Gmel. Syst. 745 (1791) = **Litsea**, Lam. (Laurin.).
japonica, J. F. Gmel. l. c. = L. japonica.

FLABELLARIA, Cav. Dis. 436. t. 264 (1790). *MALPIGHIACEAE*, Benth. & Hook. f. i. 259.
paniculata, *Cav. l. c.*—Afr. trop.

FLACOURTIA, [Comm.] L'Hérit. Stirp. Nov. 59 (1785). *BIXINEAE*, Benth. & Hook. f. i. 128.
RHAMNOPSIS, Reichb. Consp. 188 (1828).
STIGMAROTA, Lour. Fl. Cochinch. 633 (1790).
amara, Span. in Linnaea, xv. (1841) 166 = Xylosma fragrans!
aulakocarpa, *Hassk. in Versl. en Med. Akad. Wetensch.* iv. (1855) 138.—Java.
Benthami, Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 291 = Xylosma Benthami.
calophylla, Benth. ex Eichl. in Mart. Fl. Bras. xiii. i. 45 = Xylosma calophyllum.
Campbelliana, *Roxb. Hort. Beng.* 73, nomen.—Sumatra.
campylocarpa, *Miq. Fl. Ind. Bat. Suppl.* 388.—Sumatra.
cataphracta, Blume, Bijdr. 55 = F. Rukam.
cataphracta, *Roxb. ex Willd. Sp. Pl.* iv. 880.—Ind. or.; Malaya.
cataphracta, Sieber, ex Presl, Bot. Bemerk. 11 = Hilsenbergii!
cataphracta, Wall. Cat. n. 6774 C = Xylosma controversum.
celastrina, H. B. & K. Nov. Gen. et Sp. vii. 239 = Xylosma nitidum.
chinensis, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 219 = Xylosma racemosum.
cinerea, H. B. & K. ex Hemsl. Biol. Centr. Am. Bot. i. 57 = Xylosma cinereum.
cordata, *H. B. & K. Nov. Gen. et Sp.* vii. 241.—Peruv.
cordifolia, Roxb. Hort. Beng. 73, nomen = sepiaria.
coriifolia, Blanco, Fl. Filip. ed. II. 559 = Scolopia rhinanthera.
crenata, Wall. Cat. n. 6679 = Scolopia crenata.
digyna, Benth. ex Eichl. in Mart. Fl. Bras. xiii. i. 447 = Xylosma digynum.
edulis, Schum. & Thonn. Beskr. Guin. Pl. 450 = flavescens.
elegans, *Tul. in Ann. Sc. Nat. Sér. III.* vii. (1847) 294.—N. Granat.
ferox, Wall. ex Volgt, Hort. Suburb. Calc. 84 = Xylosma longifolium.
flavescens, *Willd. Sp. Pl.* iv. 880.—Afr. trop.
flexuosa, H. B. & K. Nov. Gen. et Sp. vii. 239 = Xylosma flexuosum.
Gambecola, *Clos, in Ann. Sc. Nat. Sér. IV.* viii. (1857) 219.—Afr. trop.
heterophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 321.—Java.
Hilsenbergii, *Presl, Bot. Bemerk.* 11.—Ins. Maurit.

FLACOURTIA:—
hirtiuscula, *Oliver, Fl. Trop. Afr.* i. 121.—Afr. trop.
inermis, Miq. ex Hook. f. & Thoms. in Fl. Brit. Ind. i. 192 = montana, *Y. Grah.*
inermis, *Roxb. Hort. Beng.* 73; *Fl. Ind.* 883.—Ins. Moluce.
inermis, Wall. Cat. n. 6673 G, H = Scolopia rhinanthera.
Jangomas, Steud. Nom. ed. I. 843 = cataphracta.
japonica, Hort. ex Lavall. Arbor. Segrez. 41 = Idesia polycarpa.
japonica, Walp. Rep. i. 205 = Xylosma racemosa.
lucida, Saliab. Prod. 386 = F. Ramontchi.
mollis, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 192.—Burma.
montana, *Y. Grah. Cat. Bomb. Pl.* 10.—Ind. or.
montana, *Zipp. ex Span. in Linnaea*, xv. (1841) 166, nomen.—Ins. Timor.
nitida, Benth. in Hook. Kew Journ. iii. (1851) 119 = Xylosma Benthami.
nivea, Moon, Cat. Ceyl. Pl. 70 = Fluggea Leucopyrus.
obcordata, Roxb. Hort. Beng. [106]; Fl. Ind. iii. 885 = sepiaria.
obtusa, Hochst. ex Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 236 = Aberia abyssinica.
obtusata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 34 = Aberia abyssinica.
parviflora, Blanco, Fl. Filip. ed. II. 560 = Scolopia Dayzanthera.
Perrottetiana, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 218 = F. Ramontchi.
prunifolia, Benth. Pl. Hartw. 160 = Xylosma Benthami.
prunifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 240. t. 654.—N. Granat.
quintuplinervis, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 553 = inermis.
racemosa, *Presl, Rel. Haenk.* ii. 98.—Calif.
racemosa, Sieb. & Zucc. Fl. Jap. i. 169. t. 88 = Xylosma racemosum.
Ramontchi, *L'Hérit. Stirp. Nov.* 59. t. 80.—Ind. or.; Malaya: Madag.
rhamnoides, Burch. ex DC. Prod. i. 256 = Dovyalis zizyphoides.
rotundifolia, *Clos, in Ann. Sc. Nat. Sér. IV.* viii. (1857) 218.—Java.
rotundifolia, Roxb. Hort. Beng. 73 = sepiaria.
Rukam, *Zoll. & Mor. Syst. Verz. Zoll.* 38.—Malaya.
sapida, Roxb. Pl. Corom. i. 49. t. 69 = F. Ramontchi.
sapida, Wall. Cat. n. 6675 = Scolopia crenata.
sepiaria, *Roxb. Pl. Corom.* i. 48. t. 68.—Ind. or.; Malaya.
spiculiflora, *Tul. in Ann. Sc. Nat. Sér. III.* vii. (1847) 292 (spiciflera).—N. Granat.
stigmarota, Wall. Cat. n. 6678, partim = Scolopia Roxburghii.
sumatrana, *Planch. ex Hook. f. Fl. Brit. Ind.* i. 192.—Burma: Malaya.
timorensis, Zipp. ex Span. in Linnaea, xv. (1841) 166 = Xylosma fragrans!
tomentella, *Miq. Fl. Ind. Bat. Suppl.* 105.—Sumatra.
tomentosa, *Presl, Rel. Haenk.* ii. 98.—Am. trop.
velutina, *Tul. in Ann. Sc. Nat. Sér. III.* vii. (1847) 295.—N. Granat.
Vogelii, *Hook. f. in Hook. Niger Fl.* 220.—Afr. trop
Wightiana, Wall. Cat. n. 6672 (Scepae sp.).—Ind. or.

FLACURTIA, Juss. Gen. 391 (1789) = **Flacourtia**, Comm. (Bixin.).

FLADERMANNIA, Endl. Gen. Suppl. i. 1400 (1841) = FALDERMANNIA, Trautv. = **Einiphora**, Linn. (Rubiac.).

FLAGELLARIA, Linn. Diss. Dass. 18 (1747); Amoen. Acad. i. 396 (1749). *FLAGELLARIEAE*, Benth. & Hook. f. iii. 860.
angustifolia, *Wall. Cat.* n. 5199.—Geront. trop.
aramiflora, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 16 (= *Dryniopsermum uvens* !).—Java.
ascendens, *Seem. Fl. Vit.* 315.—Ins. Fiji.
elegans, *Seem. l. c.*—Ins. Fiji.
gigantea, *Hook. f. in Hook. Ic. Pl.* t. 1439.—Ins. Fiji.

FLAGELLARIA :—
guineensis, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 181.—Afr. trop.
indica, *Linn. Sp. Pl.* 388.—Ind. or. ; Ins. Malay.
Loureiri, *Steud. Syn. Pl. Cyp.* 214.—Cochinchin.
minor, *Blume, ex Schult. f. Syst.* vii. 1498.—Malaya.
plicata, *Hook. f. in Kew Journ.* vii. (1855) 300.—N. Caled.
repens, Lour. Fl. Cochinch. 213 = Pothos Loureiri.

FLAGELLUM, Baill. Hist. Pl. vii. 448 (1880).
RUBIACEAE.
triflorum, *Baill. l. c.* 444.—Madag.

FLAMARIA, Rafin. New Fl. Am. ii. 71 (1836) = **Russelia**, Jacq. (Scrophul.).
coccinea, Rafin. l. c. = R. sarmentosa.

FLAMMARA, Hill, Veg. Syst. xvi. 37 (1770) = **Anemone**, Tourn. (Ranunc.).

FLAMMULA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 224 = **Ranunculus**, Linn.
Lingua, Fourr. l. c. = R. Lingua.
Ranunculus, Fourr. l. c. = R. Flammula.

FLAVERIA, Juss. Gen. 386 (1789). *COMPOSITAE*, Benth. & Hook. f. ii. 407.
BROTERA, Spreng. in Schrad. Journ. ii. (1802) 186. t. 5.
NAUENBERGIA, Willd. Sp. Pl. iii. 2893 (1803).
VERMIFUGA, Ruiz & Pav. Prod. Fl. Per. 114. t. 24 (1794).
angustifolia, *Pers. Syn.* ii. 489.—Mexic.
australasica, *Hook. in Mitch. Journ. Trop. Austral.* 118.—Austral.
bonariensis, DC. Prod. v. 635 = F. Contrayerba.
capitata, Juss. ex Sm. in Rees, Cycl. xiv. n. 1 = F. Contrayerba.
chilensis, J. F. Gmel. Syst. 1269 = angustifolia.
chlorsefolia, *A. Gray, Pl. Fendl.* 88.—Texas ; Mexic.
Contrayerba, *Pers. Syn.* ii. 489.—Am. trop.
humilima, *Sch. Bip in Linnaea*, xxxiv. (1865–66) 529, nomen.—Bolivia.
integrifolia, Moç. & Sesse, ex DC. Prod. v. 635 = angustifolia.
linearis, *Lag. Gen. et Sp. Nov.* 33. n. 407.—Ind. occ.
longifolia, *A. Gray, Pl. Fendl.* 88.—Mexic.
maritima, H. B. & K. Nov. Gen. et Sp. iv. 255 = linearis.
peruviana, J. F. Gmel. Syst. 1549 = F. Contrayerba.
peruviana, Juss. Gen. 187 = Piqueria artemisioides.
radicans, Moç. & Sesse, ex DC. Prod. v. 635 = angustifolia.
repanda, *Lag. Gen. et Sp. Nov.* 33.—Texas ; Mexic.
spicata, Sm. in Rees, Cycl. xiv. n. 2 = Piqueria artemisioides.
tenuifolia, Nutt. in Journ. Acad. Phil. N. S. vii. (1834) 81 = linearis.
trinervata, Baill. Hist. Pl. viii. 55 = repanda.

FLAVIA, Heist. Syst. 12 (1748) (Quid ?).

FLAVICOMA, Rafin. Fl. Tellur. iv. 68 (1836) = **Schaueria**, Nees (Acanthac.).
capitata, Rafin. l. c. = S. flavicoma.
paniculata, Rafin. l. c. = S. calycotricha.

FLEISCHERIA, Steud. & Hochst. ex Endl. Gen. 497 (1838) = **Sonchoneura**, Linn. (Compos.).
Gerogapon, Hochst. & Steud. ex Boiss. Fl. Orient. iii. 789 = S. ciata.

FLEISCHERIA, Steud. in Lehm. Pl. Preiss. i. 336 (1844) = **Sida**, Linn. (Malvac.).
pubens, Steud. l. c. 337 = S. calyxhymenia.

FLEISCHMANNIA, Sch. Bip. in Flora, xxxiii. (1850) 417. *COMPOSITAE*, Benth. & Hook. f. ii. 248.
rhodostyla, *Sch. Bip. l. c.*—Mexic.
Schaffneri, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 101.—Mexic.
ureasefolia, *Benth. & Hook. f. Gen.* ii. 248.—Mexic.

FLEMINGIA, Roxb. ex Wall. Cat. sub n. 4361 et 4364 (1830 ?) = **Canacora**, Lam. (Gentian.).
decussata, Roxb. ex Wall. l. c. n. 4364 = C. decussata.
virgata, Roxb. ex Wall. l. c. n. 4361 = C. diffusa.

FLEMINGIA, Roxb. ex Ait. Hort. Kew. ed. II. iv. 349 (1812). *LEGUMINOSAE*, Benth. & Hook. f. i. 544.
LEPIDOCOMA, Pangh. Reise durch Java, 388 (1846).
LOUREIRA, Jaume St. Hil. in Bull. Soc. Philom. (1812) 198 (Lacosa).
MILLINGTONIA, Roxb. ex Wight & Arn. Prod. 241, in syn. (1824).
MOGHANIA, Jaume St. Hil. in Desv. Journ. Bot. i. (1813) 61.
OSTRYODIUM, Desv. l. c. 119. t. 6 (1813).
affinis, *Presl, Rel. Bemarch.* 59.—Ins. Trinit.
abrupta, Wall. Cat. n. 5755 = strobilifera.
angustifolia, Roxb. Hort. Beng. [98] ; Fl. Ind. iii. 341 = congesta.
Bhotiya, Buch.-Ham. in Wall. Cat. n. 5746 = congesta.
Blancoana, Llanos, Fragm. 80 = lineata.
bracteata, Wight, Ic. t. 368 = strobilifera.
capitata, Buch.-Ham. ex Wall. Cat. n. 5749 = congesta.
capitata, Zoll. in Miq. Fl. Ind. Bat. i. 1. 166 = involucrata.
Chappar, *Buch.-Ham. ex Wall. Cat.* n. 5757.—Ind. or. ; Burma.
chlorostachys, Wall. l. c. n. 5756 = strobilifera.
congesta, *Roxb. ex Ait. Hort. Kew.* ed. II. iv. 349.—As trop.
Cuminginana, *Benth. in Miq. Pl. Jungh.* 245.—Ins. Philipp.
faginea, *Benth. & Hook. f. Gen.* i. 544 ; *Baker, in Oliver, Fl. Trop. Afr.* ii. 230.—Afr. trop.
ferruginea, Wall. Cat. n. 5750 = congesta.
fruticosa, Wall. l. c. n. 5764 = strobilifera.
Grahamiana, *Wight & Arn. Prod.* 242.—Ind. or.
guineensis, *G. Don, Gen. Syst.* ii. 309.—Afr. trop.
Hallaya, Buch.-Ham. ex Wall. Cat. n. 5440 = Spathoblus Roxburghii.
Helferiana, *Presl, Bot. Bemerk.* 140.—Ind. or. ; Burma.
Horsfieldii, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 165.—Java.
involucrata, *Benth. in Miq. Pl. Jungh.* 246.—As. et Austral. trop.
Lamentii, Hance, in Journ. Bot. xvi. (1878) 10 = congesta.
latifolia, Benth. in Miq. Pl. Jungh. 246 = congesta.
lineata, *Benth. ex Ait. Hort. Kew.* ed. II. iv. 350.—As. et Austral. trop.
macrophylla, Blume, ex Miq. Fl. Ind. Bat. i. 1. 163 = congesta.
nana, Roxb. ex Ait. Hort. Kew. ed. II. iv. 350 = congesta.
nana, Wall. Cat. n. 5748 B = Wallichii.
oblongifolia, *Baker, in Oliver, Fl. Trop. Afr.* ii. 230.—Afr. trop.
paniculata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 390 = congesta.
paniculata, *Wall. Cat.* n. 5759.—Reg. Himal. ; Burma.
parviflora, *Benth. Fl. Austral.* ii. 269.—Austral.
parviflora, Buch.-Ham. ex Wall. Cat. n. 5752 = lineata.
pauciflora, *Benth. Fl. Austral.* ii. 269.—Austral.
Phurzia, Buch.-Ham. ex Wall. Cat. n. 5758 = paniculata.
polysperma, Moon, Cat. Ceyl. Pl. 84 = Pycnospora nervosa.
procumbens, Roxb. Hort. Beng. 56 ; Fl. Ind. iii. 338 = congesta.
procumbens, Wight, Ic. t. 987 = vestita.
prostrata, Roxb. f. ex Roxb. Hort. Beng. 56 ; Fl. Ind. iii. 338 = congesta.
pycnocarpa, Benth. in Linnaea, xxiv. (1851) 648 = Grahamiana.
rhodocarpa, Baker, in Oliver, Fl. Trop. Afr. ii. 231 = congesta.
Rothiana, DC. Prod. ii. 351 = Rhynchosia rufescens.
semialata, Roxb. Hort. Beng. 56 ; Fl. Ind. iii. 340 = congesta.
semialata, Wall. Cat. n. 5746 G = Wallichii.
sericans, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 188 = Wallichii.
stricta, *Roxb. Hort. Beng.* 56 ; *Fl. Ind.* iii. 342.—Ind. or. ; China.
stricta, Wall. Cat. n. 5745 E = congesta.
strobilifera, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 350. —Ind. or. ; Malaya.
Teysmanniana, *Miq. Fl. Ind. Bat.* i. 1. 1083.—Sumatra.

FLEMINGIA—
trinervia, Desf. Tabl. ed. II. 269 = congesta.
tuberosa, *Dalzell, in Hook. Kew Journ.* ii. (1850) 34. —Ind. or.
vestita, *Benth. ex Baker, in Hook. Fl. Brit. Ind.* ii. 230.—Reg. Himal.
Wallichii, *Wight & Arn. Prod.* 242.—Ind. or. ; Burma.
Wightiana, R. Grah. in Wall. Cat. n. 5751 = congesta.

FLEMINGIA, Roxb. ex Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 202 = **Thunbergia**, Linn. f. (Acanthac.).
coccinea, Buch.-Ham. ex Nees, in DC. Prod. xi. 61 = T. coccinea.
grandiflora, Roxb. ex Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 202 = T. grandiflora.
racemosa, Herb. Madr. ex Wall. Cat. n. 7685 = T. grandiflora.

FLEMMINGIA, Walp. Rep. i. 790 (1842) = **Flemingia**. Roxb. (Legumin.).

FLESSERA, Adans. Fam. ii. 192 (1763) = **Lophanthus**, ? Adans. (Labiat.).

FLEURA, Stend. Nom. ed. II. i. 643 (1840), sphalm = **Fleurya**, Gaudich. (Urtic.).

FLEUROTIA, Reichb. Nom. 90 (1841) = **Siebera**, J. Gay (Compos.).

FLEURYA, Gaudich. in Freyc. Voy. Bot. 497 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 382.
FLEURYOPSIS, Opiz, in Lotos, iii. (1853) 240.
KERMULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2.
SCHYCHOWSKYA, Endl. in Ann. Wien. Mus. i. (1836) 167. t. 18.
aestuans, *Gaudich. in Freyc. Voy. Bot.* 497.—Am trop.
bulbifera, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 122 = Laportea bulbifera.
canadensis, Benth. in Hook. Niger Fl. 517 = Laportea canadensis.
capensis, *Wedd. in Ann. Sc. Nat.* Sér. IV. (1854) i. 188.—Afr. austr.
Caraccllana, Wedd. l. c. Ser. III. xviii. (1852) 304 = aestuans.
cordata, Gaudich. in Freyc. Voy. Bot. 498 = aestuans.
corylifolia, Gaudich. l. c. = aestuans.
cuneata, Wedd. in DC. Prod. xvi. 1. 73 = umbellata.
cymosa, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 118.—Malaya.
divaricata, Gaudich. in Freyc. Voy. Bot. 497 = Laportea canadensis.
glandulosa, Wedd. in Ann. Sc. Nat. Sér. III. xviii (1852) 305 = aestuans.
glechomaefolia, Miq. in Zoll. Syst. Verz. Ind. Archip. 103, 106 = Fatoua pilosa.
globulifera, Miq. l. c. = Fatoua pilosa.
glomerata, Gaudich. in Freyc. Voy. Bot. 496 = interrupta.
glomerata, Griseb. Cat. Pl. Cub. 58 = cuneata.
grossa, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 188. —Afr. austr.
humuifolia, Kunth, Ind. Sem. Hort. Berol. (1846) 11 = Cerastiana heterophylla.
ingrata, Miq. in Zoll. Syst. Verz. Ind. Archip. 103, 106 = aestuans.
interrupta, *Gaudich. in Freyc. Voy. Bot.* 497.—Geront. trop.
luzida, Blume, Mus. Bot. Lugd. Bat. ii. f. xxi = aestuans ?
mitis, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 188 = capensis.
paniculata, Gaudich. in Freyc. Voy. Bot. 497 = ruderalis.
peduncularia, Wedd. in DC. Prod. xvi. 1. 75 = capensis.
petiolata, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 496 = aestuans.
photinifolia, Kunth, Ind. Sem. Hort. Berol. (1846) 11 = Laportea photinifolia.
podocarpa, *Wedd. in DC. Prod.* xvi. 1. 76.—Afr. trop.
racemosa, *Gaudich. in Freyc. Voy. Bot.* 497.—Bourb. ? ruderalis, *Gaudich. l. c.*—Malaya.

FLEURYA :—
rugosissima, *Miq. Fl. Ind. Bat. Suppl.* 412.—Sumatra.
scabra, Miq. in Zoll. Syst. Vers. Ind. Archip. 108, 106 = Fatoua pilosa.
spicata, Gaudich. in Freyc. Voy. Bot. 497 = interrupta.
umbellata, *Weddl. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 304.—Cuba.

FLEURYOPSIS, Opiz, in Lotos, iii. (1853) 240 = **Fleurya**, Gaudich. (Urtic.).
petiolata, Opiz, l. c. = Fleurya aestuans.

FLEXULARIA, Rafin. in Journ. Phys. lxxxix. (1819) 105 (Gramin.).
rompresa, Rafin. l. c. (Quid ?).—Am. bor.

FLINDERSIA, R. Br. in Flind. Voy. ii. 595. t. 1 (1814). *MELIACEAE*, Benth. & Hook. f. i. 340.
OXLEYA, A. Cunn. in Hook. Bot. Misc. i. (1830) 246. t. 54.
STRZELECKYA, F. Muell. in Hook. Kew Journ. ix. (1857) 308.
amboinensis, *Poir. Encyc. Suppl.* iv. 650.—Ins. Molucc.
australis, *R. Br. in Flind. Voy.* ii. 595. t. 1.—Austral.
australis, F. Muell. Fragm. iii. 26 = Bennettiana.
Bennettiana, *F. Muell. ex Benth. Fl. Austral.* i. 389.—Austral.
Bourjotiana, *F. Muell. Fragm.* ix. 138.—Austral.
Brayleyana, *F. Muell. l. c.* v. 148.—Austral.
Fournieri, *Lench. & Sebert, Not. Bois Nouv. Calléd.* 228.—N. Caled.
Hillsiana, *F. Muell. Fragm.* x 94.—Austral.
Leichardtii, *C. DC. in DC. Monog. Phan.* i. 781.—Austral.
maculosa, *F. Muell. in Journ. Pharm. Soc. Vict.* ii. 14.—Austral.
Oxleyana, *F. Muell. Fragm.* i. 65.—Austral.
papuana, *F. Muell. Papuan Pl.* i. 84.—N. Guin
Pimentelliana, *F. Muell. Fragm.* ix. 132.—Austral.
Schottiana, *F. Muell. l. c.* iii. 35.—Austral.
Strzeleckiana, F. Muell. l. c. i. 65 = maculosa.

FLIPANTA, Rafin. Fl. Tellur. iii. 92 (1836) = **Salvia**, Tourn. (Labiat.).
ovata, Rafin. l. c. = S. Verbenaca.

FLOERKEA, Spreng. Anleit. ii. 11. 528 (1818) = **Ademonphora**, Pisch. (Campan.).
marsupiflora, Spreng. l. c. = A. marsupiflora.

FLOERKEA, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 448. *GERANIACEAE*, Benth. & Hook. f. i. 276.
Douglasii, Baill. Hist. Pl. v. 20 = Limnanthes Douglasii.
lacustris, Pers. Syn. i. 398 = proserpinacoides.
palustris, Nutt. Gen. Am. i. 238 = proserpinacoides.
proserpinacoides, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 449.—Am. bor.
uliginosa, Muhl. Cat. 36 = proserpinacoides.

FLOMOSIA, Rafin. Fl. Tellur. iv. 89 (1836) = **Verbascum**, Tourn. (Scrophul.).
condensata, Rafin. l. c. = V. phlomoides.
nemorosa, Rafin. l. c. = V. phlomoides.
phlomoides, Rafin. l. c. = V. phlomoides.

FLORESTINA, Cass. in Bull. Soc. Philom. (1815) 178. *COMPOSITAE*, Benth. & Hook. f. ii. 406.
LEPIDOPAPPUS, [Mog. & Sesse,] ex DC. Prod. v. 655 (1836).
callosa, DC. Prod. v. 655 = Polypteris callosa.
pedata, *Cass. in Bull. Soc. Philom.* (1818) 176.—Mexic.
tripteris, *DC. Prod.* v. 655.—Mexic.

FLORINDA, Noronha, ex Endl. Gen. 1086, in syn. (1841) = **Polycardia**, Juss. (Celastrin.).
anthophylla, Noronha, ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 101 = P. phyllanthoides.

FLORISCOPA, F. Muell. Census, 128 (1882), err. typ. = **Floscopa**, Lour. (Commelin.).

FLORKEA, Rafin. in Med. Repos. N. York, v. (1808) 351 = **Floerkea**, Willd. (Geraniac.).

FLOSCOPA, Lour. Fl. Cochinch. 192 (1790). *COMMELINACEAE*, Benth. & Hook. f. iii. 852.
DITHYROCARPUS, Kunth, in Ber. Akad. Berl. (1841) 245; Enum. Pl. iv. 76 (1843).
africana, *C. B. Clarke, in DC. Monog. Phan.* iii. 267. —Afr. trop.
axillaris, *C. B. Clarke, l. c.* 268.—Afr. trop.
capensis, Hassk. Commel. Ind. 166 = glomerata.
flavida, *C. B. Clarke, in DC. Monog. Phan.* iii. 269.—Afr. trop.
glabrata, *Hassk. Commel. Ind.* 166.—Am. austr.
glomerata, *Hassk. l. c.*—Afr. austr.
Hemiltonii, Hassk. l. c. = scandens.
hirsuta, Hassk. l. c. 165 = glabrata.
Mannii, *C. B. Clarke, in DC. Monog. Phan.* iii. 268.—Afr. trop.
Meyeriana, Hassk. Commel. Ind. 167 = scandens.
pannsculata, Hassk. in Miq. Pl. Jungh. 151 = scandens, Aneilema conspicuum.
peruviana, *Hassk. ex C. B. Clarke, in DC. Monog. Phan.* iii. 270.—Peruv.
petiolata, Hassk. Commel. Ind. 167 = scandens.
rivularis, *C. B. Clarke, in DC. Monog. Phan.* iii. 267. —Abyssin.
robusta, *C. B. Clarke, l. c.* 271.—Peruv.
rufa, Hassk. in Miq. Pl. Jungh. 151 = scandens.
scandens, *Lour. Fl. Cochinch.* 193.—As. et Austral.
Schweinfurthii, *C. B. Clarke, in DC. Monog. Phan.* iii. 269.—Afr. trop.
undulata, Hassk. Commel. Ind. 166 = scandens.

FLOSCUCULI, Opiz, Seznam, 44 (1852) = **Lychnis**, Linn. (Caryophyll.).
pratense, Opiz, l. c. = L. Flos-cuculi.

FLOTOVIA, Spreng. Syst. iii. 359 (1826) = **Chuquiraga**, Juss. (Compos.).
Candolleana, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 453 = C. Candolleana.
diacanthoides, Less. Syn. Comp. 95 = C. leucoxylon.
divaricata, Hieron. lc. Pl. Argent. 24.—Reg. Argent.
Doniana, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 453 = C. Doniana.
excelsa, DC. Prod. vii. 12 = C. excelsa.
ferox, Wedd. Chlor. And. i. 5.—Reg. Andin.
flagellans, Casar. Nov. Stirp. Bras. Dec. 86 = C. Regnellii.
floribunda, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 455 = C. floribunda.
fodinarum, Gardn. l. c. 453 = C. fodinarum.
glabra, Spreng. Syst. iii. 506 = C. glabra.
hispida, DC. Prod vii. 11 = C. hispida.
Hystrix, Wedd. Chlor. And. i. 6 = C. Hystrix.
imbricata, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 454 = C. reticulata.
lanceolata, Less. in Linnaea, v. (1830) 251 = C. paniculata.
latifolia, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 454 = C. latifolia.
leiocephala, Wedd. Chlor. And. i. 6.—Peruv.
leptacantha, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 138 = C. leptacantha.
Lessingiana, Gardn. l. c. 452 = C. Sprengeliana.
multiflora, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 383 = C. glabra.
orthacantha, DC. Prod. vii. 11 = C. orthacantha.
paniculata, DC. l. c. = C. paniculata.
quinquenervis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 137 = C. glabra.
Regnellii, Sch. Bip. in Linnaea, xxii. (1849) 569 = C. Regnellii.
spinescens, Less. l. c. v. (1830) 251 = C. spinescens.
Sprengeliana, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 452 = C. Sprengeliana.
synacanthe, Sch. Bip. in Linnaea, xxii. (1849) 569 = C. synacantha.
tomentosa, Spreng. Syst. iii. 506 = C. paniculata.
vagans, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 455 = C. vagans.
verians, Gardn. l. c. 454 = C. glabra.
vepreculata, DC. Prod. vii. 11 = C. vepreculata.

FLOTOVIA, Endl. Esch. 247 (1841) = FLOTOVIA, Spreng. = Chuquiraga, Juss. (Compos.).

FLOURENSIA, Cambess. in Jacquem. Voy. Bot. 27. t. 29 (1844) = **Thylacospermum**, Fenzl (Caryophyll.).
caespitosa, Cambess. l. c. = T. rupifragum.

FLOURENSIA, DC. Prod. v. 592 (1836), pro parte. *COMPOSITAE* [Benth. & Hook. f. ii. 376].
Bescriana, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 270.—Chili.
campestris, *Griseb. in Goett. Abh.* xix. (1874) 154.—Reg. Argent.
cernua, *DC. Prod.* v. 592.—Mexic.
corymbosa, DC. l. c. 592 = Viguiera Poeppigii.
hispida, *Phil. in Anal. Univ. Chil.* (1870) ii. 186.—Chili.
laurifolia, *DC. in Deless. Ic. Sel.* iv. t. 85.—Mexic.
riparia, *Griseb. in Goett. Abh.* xxiv. (1879) 196.—Reg. Argent.
thurifera, DC. Prod. v. 592 = Hellanthus thurifer.
tortuosa, *Griseb. in Goett. Abh.* xix. (1874) 154.—Reg. Argent.

FLOX, Adans. Fam. ii. p. (22) (1763) = **Lychnis**, Linn. (Caryophyll.).

FLUEGGEA, Rich. in Schrad. Neues Journ. ii. (1807) 8. t. 1 = **Ophiopogon**, Ker-Gawl. (Haemodorac.).
anceps, Rafin. Fl. Tellur. iv. 18 = O. japonicus.
angulata, Rafin. l. c. = O. japonicus !
Griffithii, Baker, in Journ. Linn. Soc. xvii. (1879) 502. —Assam.
intermedia, Kunth, Enum. Pl. v. 306 = O. intermedius.
Jaburan, Kunth, l. c. 308 = O. Jaburan.
Jacquemontiana, Kunth, l. c. 304 = praec.
japonica, Rich. in Schrad. Neues Journ. ii. (1807) 9 = O. japonicus.
prolifera, Baker, in Journ. Linn. Soc. xvii. (1879) 502.—Penang.
Wallichiana, Kunth, Enum. Pl. v. 308 = O. japonicus.

FLUEGGEA, Willd. Enum. Hort. Berol. 1013 (1809) = seq.

FLUEGGIA, Benth. & Hook. f. Gen. iii. 678 (1883) = seq.

FLUGEA, Rafin. Fl. Tellur. iv. 17 (1836) = seq.

FLUGGEA, Willd. Sp. Pl. iv. 757 (1805) [recte 1806]. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 276.
CHORISANDRA, Wight, Ic. vi. 18. t. 1994 (1853).
abyssinica, Baill. Etud. Gén. Euph. 593 = microcarpa.
Acidothamnus, Griseb. in Goett. Nachr. (1865) 164 = Securinega Acidothamnus.
angulata, Baill. Adansonia, i. (1860–61) 80 = microcarpa.
arborescens, Boj. Hort. Maurit. 278 = microcarpa.
comorensis, Boj. l. c. = microcarpa.
eglandulosa, Baill. Etud. Gén. Euph. 593 = Phyllanthus erythroxyloides.
elliptica, Baill. l. c. 592 = Securinega elliptica.
elliptica, Buch.-Ham. ex Wall. Cat. n. 7938 *B* = microcarpa.
flexuosa, Muell. Arg. in Linnaea, xxxiv. (1865–66) 76 = Securinega flexuosa.
glabra, Baak. Incl. ad DC. Prod. iv. 157, sphalm. = Phyllanthus Perrieanus.
Hilariana, *Baill. Adansonia*, v. (1864–65) 846.—Bras. javanica, *Blume, Bijdr.* 580.—Java.
leucophylla, Wall. Cat. n. 7915 = microcarpa.
Leucopyrus, Dalz. & Gibs. Bomb. Fl. 236 = microcarpa.
Leucopyrus, *Wight. Sp. Pl. iv.* 757.—Ind. or.
major, Baill. Etud. Gén. Euph. 593 = Phyllanthus indicus.
Meinechia, Muell. Arg. in Linnaea, xxxiv. (1865–66) 76 = Securinega elliptica.
melanthesioides, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 490 = microcarpa.
microcarpa, *Blume, Bijdr.* 580.—Geront. trop.
obovata, *Baill. Adansonia*, ii. (1861–62) 41.—Afr. trop.
obovata, Buch.-Ham. in Wall. Cat. n. 7926 = microcarpa.
ovalis, Baill. Etud. Gén. Euph. 593 = microcarpa.
phyllanthoides, *Baill. l. c.* 592.—Reg. Himal.
senensis, *Klotzsch, in Peters, Reise Mossamb. Bot* 106.—Afr. trop.
serrata, *Miq. Fl. Ind. Bat.* i. ii. 856.—Java.

FLUGGEA :—
sinensis, Baill. Etud. Gén. Euph. 592, nomen = microcarpa.
suffruticosa, Baill. l. c. = Securinega ramiflora.
trichogyna, Baill. l. c. 592 = Securinega trichogynia.
virosa, Baill. l. c. = microcarpa.
virosa, Buch.-Ham. ex Wall. Cat. n. 7928 B = microcarpa.
virosa, Dalz. & Gibs. Bomb. Fl. 286 = F. Leucopyrus.
Wallichiana, Baill. Etud. Gén. Euph. 592 = F. Leucopyrus.
xerocarpa, A. Juss. Tent. Euph. 106 = F. Leucopyrus.

FLUMINIA, Fries, Summa Veg. Scand. 247 (1846) = **Graphephorum**, Desv. (Gramin.).
arundinacea, Fries, l. c. = G. arundinaceum.

FLUNDULA, Rafin. Fl. Tellur. ii. 96 (1836) = **Hoeskia**, Dougl. (Legumin.).
comosa, Rafin. l. c. = H. bicolor.

FLUSTULA, Rafin. Sylva Tellur. 116 (1838) = **Vernonia**, Schreb. (Compos.).
tomentosa, Rafin. l. c. = V. arborea.

FLUVIALIS, Mich. ex Adans. Fam. ii. 472 (1763) = **Naias**, Linn.
flexilis, Pers. Syn. ii. 530 = N. flexilis.
indica, Pers. l. c. = N. indica.
minor, Pers. l. c. = N. flexilis.

FOCKEA, Endl. Iconog. t. 91 (1838) ; Nov. Stirp. Dec. 17 (1839). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 773.
CHYMOCORMUS, Harv. in Hook. Lond Journ. Bot. i. (1842) 28.
capensis, *Endl. il. cc.*—Afr. austr.
glabra, *Decne. in DC. Prod.* viii. 545.—Afr. austr.

FOENICULUM, Tourn. ex Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 902.
OXODIA, Wight & Arn. Prod. 375 (1834).
asiaticum, Mill. Gard. Dict. ed. VIII. n. 5 = vulgare.
capense, DC. Prod. iv. 142 = Carum capense.
capillaceum, Gilib. Fl. Lituan. ii. 40 = vulgare.
Carvi, Link, Enum. Hort. Berol. i. 324 = Carum Carvi.
divaricatum, Griseb. Spicil. Fl. Rumel. i. 358 = piperitum.
dulce, Mill. Gard. Dict. ed. VIII. n. 2 = vulgare.
Kraussianum, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 532 = Pimpinella caffra.
luteum, *Fisch. ex Sweet, Hort. Brit.* ed. II. 247, nomen. —Persia.
multiradiatum, C. Koch, in Linnaea, xix. (1847) 38 (certe non Foeniculum).—As. Min.
officinale, All. Fl. Pedem. ii. 25 = vulgare.
Panmorium, DC. Prod. iv. 142 = vulgare.
prucedanoides, Benth. & Hook. f. Gen. i. 902 = Silaus carvifolius.
piperitum, *Sweet, Hort. Brit.* ed. I. 187.—Europ. austr.
rigidum, Brot. ex Steud. Nom. ed. II. i. 643 = vulgare.
segetum, Presl, Fl. Sic. Prod. i. 26 = Ridolphia segetum.
tortuosum, *Benth. & Hook. f. Gen.* i. 902.—Ins. Canar.
virescens, Benth. & Hook. f. Gen. i. 902 = Silaus carvifolius.
vulgare, *Mill. Gard. Dict.* ed. VIII. n. 1.—Europ.

FOENUM-GRAECUM, [Tourn.] Rupp. Fl. Jen. ed. Hall. 268 (1745) = **Trigonella**, Linn. (Legum.).
officinale, Moench, Meth. 142 = T. Foenum-graecum.
prostratum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 260 = T. prostratum.
sativum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 388 = T. Foenum-graecum.

FOERSTERIA, Scop. Introd. 98 (1777) = **Breynia**, Forst. (Euphorb.).

FOETATAXUS, Sanlis. [Nelson], Pinaceae, 167 (1866) = **Torreya**, Arn. (Conif.).
montana, [Nelson,] l. c. = T. taxifolia.
Myristica, [Nelson,] l. c. 168 = T. californica.
nucifera, [Nelson,] l. c. = T. nucifera.
taxifolia, C. Koch, Dendrol. ii. ii. 100 = T. taxifolia.

FOETIDIA, Comm. ex Lam. Encyc. ii. 457 (1786). *MYRTACEAE*, Benth. & Hook. f. i. 724.
borbonica, J. F. Gmel. Syst. 792 = mauritiana.
mauritiana, *Lam. Encyc.* ii. 457 ; *Illustr.* t. 419.—Ins. Maurit.
obliqua, *Blume, Mus. Bot. Lugd. Bat.* i. 145.—Madag.
retusa, *Blume, l. c.* 144.—Madag.

FOLIANTHERA, Rafin. Sylva Tellur. 120 (1838) = **Strychnodendron**, Marr. (Legumin.).
guianensis, Rafin. l. c. = S. guianense.

FOLIS, Dulac, Fl. Hautes-Pyr. 190 (1867) = **Tesedalia**, R. Br. (Crucif.).
nudicaulis, Dulac, l. c. = T. nudicaulis.

FOLLICULIGERA, Pasq. Cat. Ort. Bot. Nap. 46 (1867) = **Trigonella**, Linn. (Legumin.).
cornuta, Pasq. l. c. = T. cornuta.
graveolens, Pasq. l. c. 47 = T. Foenum-graecum.
procumbens, Pasq. l. c. = T. prostrata.

FOLOMFIS, Rafin. Sylva Tellur. 97 (1838) = **Miconia**, Ruiz & Pav. (Melastom.).
fragilis, Rafin. l. c. = M. Acinodendrum.

FOMETICA, Rafin. Sylva Tellur. 75 (1838) = **Heritiera**, Dryand. (Sterculiac.).
punctata, Rafin. l. c. = H. Fomes.

FONKIA, Phil. in Linnaea, xxx. (1859–60) 198 = **Gratiola**, Rupp. (Scrophularin.).
uliginosa, Phil. l. c. 199 = G. peruviana ?

FONNA, Adans. Fam. ii. 214 (1763) = **Phlox**, Linn. (Polemon.).

FONTAINEA, Heckel, Thèse Inaug. Montpell. (1870). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 801.
Pancheri, *Heckel, l. c.*—N. Caled.

FONTANELLA, Klok, ex Bess. Prim. Fl. Gallic. ii. 368 (1808) = **Isopyrum**, Linn. (Ranunc.).
tertiaria, Klok, ex Bess. l. c. = I. thalictroides.

FONTANESIA, Labill. Pl. Syr. Dec. i. 9. t. 1 (1791). *OLEACEAE*, Benth. & Hook. f. ii. 676.
DESFONTAINESIA, Hoffingg. Verz. Pfl. 56 (1824).
chinensis, *Hance, in Journ. Bot.* vii. (1879) 136.—China.
Fortunei, Carr. in Rev. Hortic. Sér. IV. viii. (1859) 43 = phillyraeoides.
phillyraeoides, *Labill. Pl. Syr. Dec.* i. 9. t. 1.—Sicil. ; As. Min. ; Syria.

FONTENELLA, Walp. Rep. i. 524 (1842) = seq.

FONTENELLEA, St. Hil. & Tul. in Ann. Sc. Nat. Sér. II. xvii. (1842) 141. t. 7 = **Quillaja**, Molina (Rosac.).
brasiliensis, St. Hil. & Tul. l. c. = Q. Selloviana.

FOQUIERA, Hemsl. Biol. Centr. Am. Bot. i. 81 (1879) = **Fouquieria**, H. B. & K. (Tamaric.).

FORBESIA, Eckl. Verz. 4 (1827) = **Curculigo**, Gaertn. (Amaryllid.).
angustifolia, Eckl. l. c. 5 = C. plicata.
plicata, Eckl. l. c. 4 = C. plicata.

FORCHHAMMERIA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 98. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 278.
aptocarpa, *Radlk. Meth. Bot. Syst.* (1883) 54.—Mexic. pallida, *Liebm. in Kjoeb. Vidensk. Meddel.* (1853) 94. —Mexic.

FORBSTERA, Pursh, Encyc. Suppl. i. 132 (1810) ; ii. 664 (1811). *OLEACEAE*, Benth. & Hook. f. ii. 676.
ADELIA, Michx. Fl. Bor. Am. ii. 233. t. 48 (1803).
BIGELOVIA, Sm. in Rees, Cycl. xxxix (1819).
BORYA, Willd. Sp. Pl. iv. 711 (1805).
PIPTOLEPIS, Benth. Pl. Hartw. 29 (1840).
acuminata, *Pohr. Encyc. Suppl.* ii. 664.—Am. bor.
angustifolia, *Torr. Bot. Mex. Bound.* 168.—Am. bor. occ.
autumnalis, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) ? = ligustrina.

FORESTIERA :—
cassinoides, *Poir. Encyc. Suppl.* ii. 655.—Ind. occ.
cuneifolia, Rafin. New Fl. Am. iii. 91 = ligustrina ?
distichoides, Rafin. l. c. 90 = Buckleya distichophylla.
glauca, Rafin. l. c. = ligustrina ?
heterophylla, Rafin. l. c. = Buckleya distichophylla.
Fauxpulmiana, Dédr. ex Lange, Ind. Sem. Hort. Haun. (1858).—*Cf.* Linnaea, xxix. (1857–58) 737 = porulosa.
ligustrina, A. Gray, Man. N. U. St. Bot. ed. II. 358 = acuminata.
ligustrina, *Poir. Encyc. Suppl.* ii. 664.—Am. bor. neo-mexicana, *A. Gray, in Proc. Am. Acad.* xii. (1877) 68.—N. Mexic.
nitida, Rafin. New Fl. Am. iii. 91 = acuminata.
phillyreoides, *Torr. Bot. Mex. Bound.* 167.—Mexic.
porulosa, *Poir. Encyc. Suppl.* ii. 664.—Ind. occ.
prinoides, *Rafin. New Fl. Am.* iii. 92.—Am. bor.
pubescens, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 177 = ligustrina.
reticulata, *Torr. Bot. Mex. Bound.* 168.—Texas.
retusa, *Rafin. New Fl. Am.* iii. 92.—Am. bor.
rhamnifolia, *Griseb. Cat. Pl. Cub.* 169.—Cuba.
sphaerocarpa, *Torr. Bot. Mex. Bound.* 168.—Texas.
verrucosa, Rafin. New Fl. Am. iii. 90 = ligustrina ?

FORFICARIA, Lindl. Gen. et Sp. Orch. 362 (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 681.
graminifolia, *Lindl. l. c.*—Afr. austr.

FORGEROUXA, Neck. Elem. ii. 124 (1790) = **Rhamnus**, Linn.
repens, Rafin. Sylva Tellur. 32 = R. pumila.

FORGEROUXIA, Steud. Nom. ed. II. i. 643 (1840) = prec.

FORGEROUXIA, Rafin. Sylva Tellur. 32 (1838) = prec.

FORGESIA, Comm. ex Juss. Gen. 164 (1789). *SAXIFRAGEAE*. Benth. & Hook. f. i. 648.
DEFFORGIA, Lam. Illustr. ii. 71. t. 125 (1793).
borbonica, Pers. Syn. i. 194 = racemosa.
racemosa, *J. F. Gmel. Syst.* 858.—Ins. Borbon.

FORGETINA, Bocquill. ex Baill. Adansonia, vii. (1866) 49 = **Sloanea**, Linn. (Tiliac.).
guianensis, Bocquill. ex Baill. l. c. = S. cuneifolia ?

FORNASINIA, Bertol. Misc. Bot. viii. 18. t. 1 (1849) = **Millettia**, Wight & Arn. (Legumin.).
ebenifera, Bertol. l. c. 19.—Abyssin.

FORNELIA, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179, sphalm. = TORNELIA, Gutier. = **Monstera**, Adans. (Aroid.).

FORNEA, Steud. Nom. ed. II. i. 644 (1840) = seq.

FORNEUM, Adans. Fam. ii. 112 (1763) = **Andryala**, Linn. (Compos.).

FORNICARIA, Rafin. Sylva Tellur. 116 (1838) = **Salmea**, DC. (Compos.).
scandens, Rafin. l. c. = S. Eupatoria.

FORNICIUM, Cass. in Bull. Soc. Philom. (1819) 98 ; et in Dict. Sc. Nat. xvii. 349 (1820) = **Centaurea**, Linn. (Compos.).
rhaponticoides, Cass. ll. cc. 98 ; 350 = C. salina.

FORRESTIA, Rafin. in Med. Repos. N. York, II. iii. (1806) 423 = **Commelina**, Linn. (Rhaeo.).
thyrsoidea, Rafin. l. c. = C. thyrsiflora ?

FORRESTIA, A. Rich. Sert. Astrol. i. t. 1 (1834). *COMMELINACEAE*, Benth. & Hook. f. iii. 850.
AMISCHOTOLYPE, Hassk. in Flora, xlvi. (1863) 291.
glabrata, *Hassk. in Flora*, xlvii. (1864) 630.—Ind. or. Griffithii, *C. B. Clarke, in DC. Monog. Phan.* iii. 226. —Malacca.
hispida, Bot. Mag. t. 5425 = Hookeri.
hispida, *A. Rich. Sert. Astrol.* t. 1.—N. Guinea.
Hookeri, *Hassk. in Flora*, xlvii. (1864) 629.—Ind. or.
marginata, *Hassk. l. c.* 630.—Java.
mollis, *Hassk. l. c.* 628.—Singapore.
nigricans, Steud. Nom. ed. II. i. 644 = hispida.
rostrata, Hassk. in Flora, xlvii. (1864) 631 = marginata.

FORSAKHLIA, Ball. in Journ. Linn. Soc. xvi. 288 (1877), err. typ. = **Forskohlea**, Linn. (Urtic.).

FORSGARDIA, Vell. Fl. Flum. 152 (1825); iv. t. 13 = **Combretum**, Linn.
laxum, Vell. ll. cc.—Bras.

FORSHOHLEA, Betsch, Tab. t. 178 (1802), sphalm.; FORSKAELEA, Scop. Introd. 251 (1777); FORSKAHLEA, Agardh, Aphor. 204 (1832); FORSKÄHLIA, Webb & Berth. Phyt. Canar. iii. 266 (1836–50); FORSKALEA, Juss. Gen. 408 (1789); FORSKOEHLEA, Reich. Gen. 238 (1778); FORSKOELEA, Brongn. Enum. Gen. 99 (1843) = seq.

FORSSKOEHLEA, Linn. Mant. i. 79 (1767). *URTICACEÆ*. Benth. & Hook. f. iii. 313.
CAIDREJA, Forsk. Fl. Aegypt Arab. 83 (1775).
angustifolia, *Rets. Obs.* iii. 31.—Ins. Teneriff.
candida, *Linn. f. Suppl.* 240.—Afr. austr.
Cavaniana, Webb, Otia Hisp. 49. t. 45 = tenacissima.
fruticosa, Willd. Enum. Hort. Berol. 950 = angustifolia.
latifolia, Rets. Obs. iii. 31 = tenacissima.
procridifolia, *Webb, in Hook. Niger Fl.* i. 179.—Afr. trop.
scabra, Rets. Obs. iii. 31 = candida.
tenacissima, *Linn. Mant.* 72.—Geront. trop.
urticoides, Wight, Ic. t. 1982 = Drogueria diffusa.
viridis, *Ehrenb. ex Desf. Cat. Hort. Par.* ed. III. 347. —Arab.

FORSKOLEA, Linn. Dis. Opob. (1764); FORSKOLIA, Wight, Ic. vi. 11 (1853); FORSSKÅLEA, Linn. Amoen. Acad. vii. 75 (1769) = præc.

FORSTERA, Linn. f. in Nov. Act. Soc. Sc. Upsal. iii. (1780) 184. t. 9. *STYLIDIEÆ*, Benth. & Hook. f. ii. 685.
ATHECIA, Gaertn. Fruct. i. 141 (1788).
FORSTERIA, Neck. Elem. iii. 134 (1790).
arctiastrifolia, Hombr. & Jacquinot, Bot. Voy. Astrol. et Zél. (Dicot. Phan.). t. 16 = Phyllachne clavigera.
bellidifolia, *Hook. Ic. Pl.* t. 851.—Ins. Tasman.
Bidwillii, *Hook. f. Fl. N. Zel.* i. 155.—N. Zel.
clavigera, Hook. f. Fl. Antarct. 38. t. 28 = Phyllachne clavigera.
glabra, Gaertn. Fruct. i. 141. t. 28. f. 8 = sedifolia.
muscifolia, Willd. Sp. Pl. iv. 148 = Phyllachne uliginosa.
sedifolia, *Linn. f. Suppl.* 407.—N. Zel.
tenella, *Hook. f. Fl. N. Zel.* i. 154.—N. Zel.
uliginosa, Hombr. & Jacquinot, Bot. Voy. Astrol. et Zél. (Dicot. Phan.) t. 16 = Phyllachne uliginosa.

FORSTERIA, Neck. Elem. iii. 134 (1790) = **Forstera**, Linn. f. (Stylid.).

FORSTERIA, Steud. Nom. ed. I. 344 (1821) = FOERSTERIA, Scop. = **Breynia**, Forst. (Euphorb.).

FORSTRONIA, G. F. W. Mey. Prim. Fl. Esseq. 138 (1818). *APOCYNACEÆ*, Benth. & Hook. f. ii. 710.
SYRINGOSMA, Mart. ex Reichb. Consp. 134 (1828).
THYRSANTHUS, Benth. in Hook. Journ. Bot. iii. (1841) 345.
Acouci, *A. DC. in DC. Prod.* viii. 439.—Guian.
acuminata, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 99.— Bras.
adenobasis, *Muell. Arg. in Linnæa*, xxx. (1859–60) 412.—Guian.
affinis, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 100. t. 80. —Bras.
Alexandri, *Griseb. Fl. Brit. W. Ind.* 413.—Ind. occ.
australis, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 108.— Bras.
Benthamiana, *Muell. Arg. l. c.* 106.—Bras.
brachysta, *Muell. Arg. l. c.*—Bras.
brasiliensis, *A. DC. in DC. Prod.* viii. 436.—Bras.
corylifolia, *Griseb. Cat. Pl. Cub.* 172.—Cuba.
coerymbosa, *G. F. W. Mey. Prim. Fl. Esseq.* 134.—Guian.
difformis, A. DC. in DC. Prod. viii. 437 = Trachelospermum difforme.
diospyrifolia, *Muell. Arg. in Linnæa*, xxx. (1859–60) 413.—Guian.
divaricata, *Miers, Apocyn. S. Am.* 247.—Bras.
embelioides, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 97. —Bras.
floribunda, *G. F. W. Mey. Prim. Fl. Esseq.* 135, in obs.; *Decne. in DC. Prod.* viii. 437.—Jamaic.

FORSTRONIA:—
floribunda, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 96.— Bras.
Gardneri, *Muell. Arg. l. c.* 100.—Bras.
glabrescens, *Muell. Arg. l. c.* 102.—Bras.
gracilis, *Muell. Arg. l. c.* 101. in nota.—Guian.
guyanensis, *Muell. Arg. in Linnæa*, xxx. (1859–60) 414.—Guian.
lagoënsis, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 108.—Bras.
laurifolia, *A. DC. in DC. Prod.* viii. 438.—Am. trop.
leptocarpa, A. DC. l. c. = brasiliensis.
linearis, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 107.— Bras.
Luschnathii, *Muell. Arg. l. c.* 98.—Bras.
macrophylla, *Muell. Arg. in Linnæa*, xxx. (1859–60) 411.—Guian.
meridionalis, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 98. —Bras.
minutiflora, *Muell. Arg. l. c.* 99.—Bras.
montana, *Muell. Arg. l. c.* 101.—Bras.
multinervia, *A. DC. in DC. Prod.* viii. 437.—Bras.
obtusiloba, *Muell. Arg. in Linnæa*, xxx. (1859–60) 413.—Guian.
ovalifolia, *Miers, Apocyn. S. Am.* 248.—Ind. occ.
Pavonii, *A. DC. in DC. Prod.* viii. 438.—Am. trop.†
pilosa, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 99.— Bras.
protensa, *Miers, Apocyn. S. Am.* 248.—Bras.
pubescens, *A. DC. in DC. Prod.* viii. 436.—Bras.
refracta, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 97.— Bras.
Riedelii, *Muell. Arg. l. c.* 103.—Bras.
rotundiuscula, *Miers, Apocyn. S. Am.* 248.—Bras.
rufa, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 100.—Bras.
Schomburgkii, *A. DC. in DC. Prod.* viii. 438.—Guian.
Sellowii, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 101.— Bras.
spicata, *G. F. W. Mey. Prim. Fl. Esseq.* 133.—Guian.
thyrsoidea, *Muell. Arg. in Mart. Fl. Bras.* vi. t. 105. —Bras.

FORSTEROPSIS, Sond. in Lehm. Pl. Preiss. i. 398 (1845) = **Stylidium**, Sw.
Preissii, Sond. l. c. = S. Preissii.

FORSYTHIA, Vahl, Enum. i. 39 (1805). *OLEACEÆ*, Benth. & Hook. f. ii. 675.
RANGIUM, Juss. in Dict. Sc. Nat. xxiv. 200 (1822).
Fortunei, Lindl. in Gard. Chron. (1854) 412 = suspensa.
Mala-Elengi, Dennst. Schluess. Hort. Malab. 31 = Linociera cotinifolia.
suspensa, *Vahl, Enum.* i. 39.—Japon.; China.
viridissima, *Lindl. in Journ. Hort. Soc.* i. (1846) 226.—China.

FORSYTHIA, Walt. Fl. Carol. 154 (1788) = **Decumaria**, Linn. (Saxifrag.).
scandens, Walt. l. c. = D. barbara.

FORSYTHIOPSIS, Baker, in Journ. Linn. Soc. xx. (1883) 218. *ACANTHACEÆ*.
Baroni, *Baker, l. c.* 219.—Madag.

FORTUNAEA, Lindl. in Journ. Hort. Soc. i. (1846) 150 = **Platycarya**, Sieb. & Zucc. (Jugland.).
chinensis, Lindl. l. c. = P. strobilacea.

FORTUNEA, Polt. in Rev. Hortic. Sér. II. v. (1846) 262 = præc.

FORTUYNIA, Shettl. ex Boiss. in Ann. Sc. Nat. Ser. II. xvi. (1841) 379; xvii. (1842) 177. *CRUCIFERÆ*, Benth. & Hook f. i. 100.
Aucheri, *Shuttl. ll. cc.* 379, 178.—Persia.
Bongei, *Boiss. Fl. Orient.* i. 402.—Persia; Afghan.
Garcini, *Shuttl. ex Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 178.—Persia.

FORVSTHIA, Franch. & Sav. Enum. Pl. Jap. i. 312 (1875), sphalm. = **Forsythia**, Vahl (Oleac.).

FOSCARENA, Hedw. f. Gen. 388 (1808) = seq.

FOSCARENIA, Vell. ex Vand. Fl. Lusit. 7 (1788); in Roem. Script. 78 (1796) = **Bandia**, Houst. (Rubiac.).
spinosa, Roem. & Schult. Syst. ill. 99 = R. armata.

FOSSELINA, Medic. in Usteri, Ann. Bot. viii. (1794) 87 = seq.

FOSSELINIA, Scop. Introd. 318 (1777) = **Clypeola**, Linn. (Crucif.).
Jonthlaspi, All. Fl. Pedem. i. 246 = C. Jonthlaspi.

FOTERGHILLIA, Dum. Anal. Fam. 35 (1829), sphalm = **Fothergilla**, Murr. (Hamamel.).

FOTHERGILLA, Aubl. Pl. Gui. i. 440. t. 175 (1775) = **Miconia**, Ruiz & Pav. (Melast.).
mirabilis, Aubl. l. c. = M. Fothergilla.

FOTHERGILLA, Murr. Syst. Veg. ed. XIII. 418 (1774). *HAMAMELIDEÆ*, Benth. & Hook. f. i. 666.
ARROPHARS, Rafin. Sylva Tellur. 99 (1838).
alnifolia, Linn. f. Suppl. 267 = Gardeni.
Gardeni, *Murr. Syst. Veg.* ed. XIII. 418.—Am. bor.
involucrata, Falc. in Proc. Linn. Soc. i. (1839) 18 = Parrottia Jacquemontiana.
latifolia, *Prits. Ic. Ind. Bot. Suppl.* 125 (lc. ined.) (Quid?).
major, Lodd. Bot. Cab. t. 1520 = Gardeni.

FOTHERGILLIA, Spreng. Gen. i. 445 (1830) = præc.

FOUGERIA, Moench, Meth. Suppl. 243 (1802) = **Baltimora**, Linn. (Compos.).
tetragona, Moench, l. c. = B. recta.

FOUGEROUXIA, Cass. in Dict. Sc. Nat. liv. 461 (1829) = **Baltimora**, Linn. (Compos.).
alba, DC. Prod. v. 510 = B. recta.
recta, DC. l. c. = B. recta.

FOUHA, Pomel, Mat. Fl. Atl. 2 (1860) = **Colchicum**, Tourn. (Liliac.).
bulbocodioides, Pomel, l. c. = C. montanum.

FOUILLOYA, Benth. & Hook. f. Gen. iii. 950 (1883), sphalm. = seq.

FOULLIOYA, Gaudich. Bot. Voy. Bonite, t. 26 (1844†) = **Pandanus**, Linn.
graminifolia, Hort. ex Wendl. Ind. Palm. Cyclanth. Pand. Cycad. (1854) 47 = P. pygmaeus.
maritima, Gaudich Bot. Voy. Bomite, t. 26 = P. Vandermeeschi.
racemosa, Gaudich. l. c. = P. racemosus.

FOUQUIERA, Spreng. Syst. iii. 568 (1825) = seq.

FOUQUIERIA, H. B. & K. Nov. Gen. et Sp. vi. 61. t. 527 (1823). *TAMARISCINEÆ*, Benth. & Hook. f. i. 161.
BRONNIA, H. B. & K. Nov. Gen. et Sp. vi. 88. t. 538 (1823).
PHILETAIRIA, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 383.
formosa, *H. B. & K. Nov. Gen. et Sp.* vi. 88. t. 527.— Mexic.
spinosa, *Torr. in Emory, Notes Mil. Reconnois.* 147. t. 8.—Mexic.
splendens, *Engelm. in Wisliz. Tour North Mexic.* 98.* —N. Mexic.

FOURCROEA, Haw. Suppl. Pl. Succ. 42 (1819) = **Furcræa**, Vent. (Amaryllid.).

FOURCROYA, Spreng. Anleit. ii. t. 238 (1817) = præc.

FOVEOLARIA, Meisn. Pl. Gen. Comm. 28 (1836) = **Sloanea**, Linn. (Tiliac.).

FOVEOLARIA, Ruiz & Pav. Prod. 87. t. 9 (1794), partim; A. DC. in DC. Prod. viii. 271 (1844). *STYRACEÆ*, (Benth. & Hook. f. ii. 669, 670).
cordata, Ruiz & Pav. Syst. Veg. 99 = Styrax cordatum.
crocea, Ruiz & Pav. Fl. Per. iv. t. 393 = Styrax oblongum.
ferruginea, *Ruiz & Pav. Syst. Veg.* 100.—Peruv.
oblonga, Ruiz & Pav. l. c. = Styrax oblongum, Pavonii.
ovata, Ruiz & Pav. l. c. = Styrax ovatum.

FOXIA. Parl. Nuov. Gen. e Sp. 17 (1854) = **Hyacinthus**, Tourn. (Liliac.).
spicata, Parl. l. c. 18 = H. spicatus.

FRACASTORA, Adans. Fam. ii. 505 (1763) = **Sideritis**, Tourn. (Labiat.).

FRADINIA, Pomel, Nouv. Mat. Fl. Atl. 51 (1874) = **Micromischus**, Cos. (Compos.).
Geslini, Pomel, l. c. 52 = M. Geslini.
pedunculata, Pomel, l. c. 51 = Anacyclus pedunculatus.

FRAGA, Lapeyr. Hist. Abr. Pl. Pyr. 287 (1813) = **Potentilla**, Linn. (Rosac.).
micrantha, Four. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 270 = P. Fragariastrum.
sterilis, Lapeyr. Hist. Abr. Fl. Pyr. 287 = P. Fragariastrum.
Vaillantii, Lapeyr. l. c. = P. splendens.

FRAGARIA. [Tourn.] Linn. Syst. ed. 1 (1735). *ROSACEÆ*, Benth. & Hook. f. i. 620.
DUCHESNEA, Sm. in Trans Linn. Soc. x. (1811) 379.
abnormis, Tratt. Rosac. Monog. iii. 166 = vesca.
aborriva, Duchesne, in Lam. Encyc. ii. 534 = collina.
alba, Crantz, Stirp. Austr. ed. I. fasc. ii. 19; ed. II. 78 = Potentilla alba.
allena, *Weihe, in Flora*, xi. (1828) 80.—Hab.?
alpina, Stend. Nom. ed. I. 344 = vesca.
Ananassa, Duchesne, Hist. Nat. Frais. 190 = chiloënsis, grandiflora.
Anasrina, Crantz, Stirp. Austr. ed. I. fasc. ii. 9; ed. II. 71 = Potentilla Anasrina.
argentea, Crantz, Inst. ii. 177 = Potentilla argentea.
arguta, Lindl. in Wall. Cat. n. 1237 = indica.
aurea, Crantz, Stirp. Austr. ed. I. fasc. ii. 17; ed. II. 77 = Potentilla aurea.
bifera, Duchesne, in Lam. Encyc. ii. 534 = collina.
bifurca, Crantz, Inst. ii. 177 = Potentilla bifurca.
bonariensis, Pers. Syn. ii. 58 = chiloensis?
botryformis, Duchesne, in Lam. Encyc. ii. 532 = vesca.
brevifolia, Ser. in DC. Prod. ii. 570 = collina.
breviscapa, Tratt. Rosac. Monog. iii. 168 = Potentilla micrantha.
californica, *Cham. & Schlecht. in Linnæa*, ii. (1827) 20.—Calif.
californica, Newberry, in Pacif. Railr. Rep. vi. 73 = chiloënsis.
calycina, Loisel. Fl. Gall. I. 299 = collina?
calyculata, Duchesne, in Lam. Encyc. ii. 533 = chiloënsis.
campestris, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 176 = collina.
canadensis, Crantz, Inst. ii. 178 = Potentilla canadensis.
canadensis, Michx. Fl. Bor. Am. i. 299 = virginiana.
carolinensis, Duchesne, in Lam. Encyc. ii. 539 = chiloënsis.
caucasica, Crantz, Stirp. Austr. ed. I. fasc. ii. 90; ed. II. 78 = Potentilla Clusiana.
cerino-alba, Jord. & Four. Brev. Pl. Nov. fasc. i. 13 = collina.
chilensis, Molina, Sagg. Chile, 184 = seq.
chiloënsis, *Duchesne, Hist. Nat. Frais.* 165; *Mill. Gard. Dict.* ed. VIII. n. 4.—Am. bor.
chiloënsis, Georgi, Beschr. Russ. Reich. III. iv. 1038 = collina.
chrysantha, Zoll. & Mor. Syst. Verz. Zoll. 7 = indica.
collina, *Ehrh. Beitr.* vii. 26.—Europ.; As. bor.
collina, Maxim. Prim. Fl. Amur. 90 = elatior.
collinsiana, Jord. & Four. Brev. Pl. Nov. fasc. i. 14 = collina.
consobrina, Jour. & Four. l. c. 13 = collina.
Crantzii, Crantz, Inst. ii. 178 = Potentilla verna.
Daltoniana, *J. Gay, in Ann. Sc. Nat.* Sér IV. viii. (1857) 204.—Reg. Himal.
dicica. Stend. Nom. ed. I. 344 = elatior.
drymophila, Jord. & Four. Brev. Pl. Nov. fasc. i. 15 = collina.
dubia, Crantz, Stirp. Austr. ed. I. fasc. ii. 24; ed. II. 81 = Potentilla norvegica?
dubia, Duchesne, in Lam. Encyc. ii. 534 = collina.
dulcis, Sommerf. ex Homem. in Fl Dan. t. 1389 = collina.
dumetorum, Jord. ex Nym. Consp. 223 = collina.

FRAGARIA :—
ægaellis, Duchesne, Hist. Nat. Frais. 119 = vesca.
elatior, *Ehrh. Beitr.* vii. 23.—Europ.
elatius, Wight & Arn. Prod. 300 = ellgerrensis.
fragariastes, Crantz, Inst. ii. 177 = Potentilla fragaricides.
fruticosa, Crantz, l. c. 176 = Potentilla fruticosa.
grandiflora, Crantz, l. c. 179; Lam. Fl. Fr. iii. 112 = Potentilla grandiflora.
grandiflora, *Ehrh. Beitr.* vii. 25.—Gulan.
grandiflora, Thuill. Fl. Par. ed. II. 264 = collina?
Grayana, Vilm. ex J. Gay, in Ann. Sc. Nat. Sér. IV. viii. (1857) 302 = virginiana.
grossa, Sabin. Prod. 363 = chiloensis.
Hagenbachiana, Lange, ex Koch, in Flora, xxv. (1842) 583 = collina.
hirta, Crantz, Inst. ii. 177 = Potentilla hirta.
hispida, Duchesne, in Lam. Encyc. ii. 535 = collina.
hortensis, Duchesne, Hist. Nat. Frais. 113 = vesca.
hybrida, Duchesne, in Lam. Encyc. ii. 540 = virginiana.
incana, Lam. Fl. Fr. iii. 113 = Potentilla subcaulis.
indica, *Andr. Bot. Rep.* t. 479.—Ind. or.; Malaya; China.
indica, Wall. Cat. n. 1236, partim = Potentilla axilliflora.
lucida, Vilm. ex J. Gay, in Ann. Sc. Nat. Sér. IV. viii. (1857) 201 = californica.
ludoviciana, *Kuſn. Fl. Ludov.* 150.—Am. bor.
magna, Thuill. Fl. Par. ed. II. 254 = elatior.
majusfea, Ser in DC. Prod. ii. 570 = collina.
malaiyana, Roxb. Hort. Beng. 39; Fl. Ind. ii. 520 = indica.
media, Schur, Enum. Pl. Transs. 186 = collina.
mexicana, Schlecht. in Linnæa, xiii. (1839) 265 = vesca.
micrantha, Tratt. Rosac. Monog. iii. 169 = Potentilla micrantha.
minor, Duchesne, in Lam. Encyc. ii. 531 = vesca.
monophylla, Duchesne, Hist. Nat. Frais. 194 = vesca.
monspeliensis, Crantz, Inst. ii. 179 = Potentilla norvegica.
moschata, Duchesne, Hist. Nat. Frais. 145 = elatior.
multifida, Crantz, Inst. ii. 176 = Potentilla multifida.
multiplex, Duchesne, in Lam. Encyc. ii. 532 = vesca.
muricata, Linn. Sp. Pl. 495 = elatior, vesca.
neglecta, *Linden, in Bull. Soc. Nat. Mosc.* xxxviii. (1865) ii. 220.—Ross. austr.
nemorosa, Salisb. Prod. 363 = vesca.
nigra, Duchesne, in Lam. Encyc. ii. 535 = collina.
nigra, Gueldenst. ex Ledeb. Fl. Ross. ii. 65 (Rubi sp. 7). —Ross.
nilagirica, Zenk. Pl. Ind. Dec. I. 7 = indica.
nilgerrensis, *Schlecht. ex J. Gay, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 206.—Ind. or.
nitida, Crantz, Inst. ii. 179 = Potentilla nitida.
nivea, Crantz, l. c. = Potentilla nivea.
norvegica, Crantz, l. c. = Potentilla norvegica.
nubicola, Lindl. in Wall. Cat. n. 1238 = vesca.
nuda, Pers. in Linn. Syst. ed. XV. 511, in obs. = vesca.
odora, Salisb. Prod. 363 = virginiana.
opaca, Crantz, Inst. ii. 178 = Potentilla opaca.
palustris, Crantz, Stirp. Austr. ed. I. fasc. ii. 11; ed. II. 73 = Potentilla palustris.
parviflora, Lam. Fl. Fr. iii. 113 = Potentilla norvegica.
pendula, Duchesne, in Lam. Encyc. ii. 535 = collina.
pentaphyllum, Crantz, Inst. ii. 179 = Potentilla reptans.
pimpinelloides, Crantz, l. c. 177 = Potentilla pimpinelloides.
pinnatifolia, Stokes, Bot. Mat. Med. iii. 149 = Potentilla palustris.
plataneoides, *Ser. ex DC. Prod.* ii. 570.—Am. bor.?
praecox, *Kit. ex Schult. Oester. Fl.* ed. II. ii. 82.—Hungar.
praecox, Salisb. Prod. 363 = Potentilla Fragariastrum.
pratensis, Duchesne, in Lam. Encyc. ii. 536 = elatior.
recta, Crantz, Stirp. Austr. ed. I. fasc. ii. 22; ed. II. 79 = elatior, vesca.
reptans, Crantz, Inst. ii. 179 = Potentilla reptans.
reversa, *Kit. in Linnæa*, xxxii. (1863) 595.—Hungar, roscifiora, *Baulay, in Bull. Soc. Bot. Fr.* xviii. (1871) 98.—Europ.
Roxburghii, Wight & Arn. Prod. 300 = indica.
rubens, Crantz, Stirp. Austr. ed. I. fasc. ii. 14; ed. II. 75 = Potentilla opaca?
rubricaulis, Jord. & Four. Brev. Pl. Nov. fasc. i. 16 = collina.

FRAGARIA :—
rupestris, Crantz, Stirp. Austr. ed. I. fasc. ii. 10; ed. II. 73 = Potentilla rupestris.
scabra, Schur, Enum. Pl. Transs. 186 = elatior.
semperflorens, Duchesne, Hist. Nat. Frais. 49 = vesca.
semperirens, Duchesne, ex Steud. Nom. ed. I. 344 = vesca.
sericea, Crantz, Inst. ii. 176 = Potentilla sericea.
sericea, Dougl. ex Hook. Fl. Bor. Am. i. 185 = chiloënsis.
serotina, Raſn. Atl. Journ. 152 = virginiana?
sicca, Gilib. Fl. Lituan. ii. 248 = Potentilla Fragariastrum.
sikkimensis, Kurz, in Journ. As. Soc. Beng. xliv. (1876), ii. 206 = Daltoniana.
silvestris, Duchesne, Hist. Nat. Frais. 61 = vesca.
Soyeriana, Jord. & Four. Brev. Pl. Nov. fasc. i. 15 = collina.
sterilis, Bieb. Fl. Taur. Cauc. i. 408 = Potentilla micrantha.
sterilis, Linn. Sp. Pl. 495 = Potentilla Fragariastrum.
stipularis, Crantz, Inst. ii. 176 = Potentilla stipularis.
subacaulis, Crantz, l. c. 180 = Potentilla subacaulis.
succulenta, Gilib. Fl. Lituan. ii. 247 = vesca.
suecica, Jord. & Four. Brev. Pl. Nov. fasc. i. 18 = collina.
sundaica, Blume, Bijdr. 1106 = Potentilla Kleiniana.
supina, Crantz, Stirp. Austr. ed. I. fasc. ii. 10; ed. II. 73 = Potentilla supina.
sylvestris, Duchesne, in Lam. Encyc. ii. 531, vide *silvestris*.
sylvulicola, Jord. & Four. Brev. Pl. Nov. fasc. i. 14 = collina.
Thomasiana, Jord. & Four. l. c. = collina.
tinctu, Duchesne, in Lam. Encyc. ii. 539 = chiloënsis.
Tormentilla, Crantz, Stirp. Austr. ed. I. fasc. ii. 23; ed. II. 80 = Potentilla Tormentilla.
umbellifera, F. Schultz, ex Nym. Consp. 222 = collina.
unifolia, Stend. Nom. ed. I. 345 = vesca.
valderia, Crantz, Inst. ii. 179 = Potentilla valderia.
verna, Crantz, Stirp. Austr. ed. I. fasc. ii. 13; ed. II. 74 = Potentilla verna.
vesca, Benth. Pl. Hartw. 309 = californica.
vesca, *Linn. Sp. Pl.* 494.—Reg. temp.
vesca, Lour. Fl. Cochinch. 332 = elatior.
vesca, Walt. Fl. Carol. 150 = virginiana.
villosa, Crantz, Stirp. Austr. ed. II. 75. t. 1. f. 2 = Potentilla rubens.
virginiana, *Duchesne, Hist. Nat. Frais.* 204; *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. bor.
viridis, Duchesne, l. c. 135 = collina.
vulgaris, Ehrh. Beitr. vii. 21 = vesca.

FRAGARIASTRUM, Heist. Syst. 7 (1748); Schur, Enum. Pl. Transs. 187 (1866) = **Potentilla**, Linn. (Rosac.).
album, Schur, l. c. = P. alba.
caulescens, Schur, l. c. 188 = P. caulescens.
Clusianum, Schur, l. c. 187 = P. Clusiana.
micranthum, Schur, l. c. = P. micrantha.
petiolatum, Schur, l. c. 188 = P. caulescens.
sterile, Schur, l. c. 187 = P. Fragariastrum.

FRAGARIOPSIS, A. St. Hil. Morphol. 496 (1840) = **Plukenetia**, Linn. (Euphorb.).
discolor, Baill. Etud. Gén. Euph. 498 = *scandens*.
polyandra, Baill. l. c. = *scandens*.
scandens, A. St. Hil. Morphol. 496.—Bras.
Warmingii, Muell. Arg. in Mart. Fl. Bras. xi. ii. 388. —Bras.

FRAGERIA, Delile, ex Steud. Nom. ed. II. i. 645 (1840) = **Leucoseris**, Lag. (Compos.).

FRAGMOSA, Raſn. Fl. Tellur. ii. 50 (1836) = **Comyns**, Linn. (Compos.).
alpina, Raſn. l. c. — Hab.?
aspera, Raſn. l. c. = C. cachetonoides?
globella, Raſn. l. c. — Hab.?
nudicaulis, Raſn. l. c. — Hab.?
pumila, Raſn. l. c. = C. pumila.
uniflora, Raſn. l. c. = C. uniflora.

FRAGOSA, Rafs & Pav. Prod. 43. t. 84 (1794) = **Anacyclus**, Lam. (Unbellif.).
arvicoides, H. B. & K. Nov. Gen. et Sp. v. 87. t. 434 = A. arvicoides.

FRAGOSA :—
bilobo, Schlecht. in Linnæa, xxviii. (1856) 478.—Peruv.
cladorhiza, Ruiz & Pav. Fl. Per. iii. 27. t. 249 = Azorella cladorhiza.
corymbosa, Ruiz & Pav. l. c. t. 250 = A. corymbosa.
crenata, Ruiz & Pav. l. c. t. 249 = A. crenata.
multifida, Ruiz & Pav. l. c. = A. multifida.
reniformis, Ruiz & Pav. l. c. 26. t. 249 = A. reniformis.
spinosa, Ruiz & Pav. l. c. iii. 27 = A. spinosa.

FRAGROSA, Hedw. f. Gen. 191 (1806) = præc.

FRANCA, Mich. ex Adans. Fam. ii. 255 (1763) = **Frankenia**, Linn.
corymbosa, Vis. in Mem. Ist. Venet xvi. (1871) 158 = Frank. hirsuta.
ericifolia, Vis. l. c. = Frank. ericifolia.
fruticulosa, Vis. l. c. = Frank. pauciflora.
hispida, Vis. l. c. = Frank. hirsuta.
laevis, Vis. l. c. = Frank. hirsuta.
microphylla, Vis. l. c. = Frank. microphylla.
mollis, Vis. l. c. = Cressa cretica.
nodiflora, Vis. l. c. = Frank. nodiflora.
Nothria, Vis. l. c. = Frank. hirsuta.
pauciflora, Vis. l. c. = Frank. pauciflora.
pulverulenta, Vis. l. c. = Frank. pulverulenta
revoluta, Vis. l. c. = Frank. hirsuta.
tetrapetala, Vis. l. c. = Frank. tetrapetala.
thymifolia, Vis. l. c. = Frank. hirsuta.
velutina, Vis. l. c. = Frank. hirsuta.

FRANCASTORA, Steud. Nom. ed. II. ii. 580, in syn. (1841) = FRACASTORA, Adans. = **Sideritis**, Tourn. (Labiat.).

FRANCHETIA, Baill. in Bull. Soc. Linn. Par. i. (1885) 477 *EUPHORBIACEAE.*
sphæranthe, *Baill. l. c.*—Madag.

FRANCISCEA, Pohl, Pl. Bras. Ic. i. 1. t. 1 ad 7 (1827) = **Brunfelsia**, Linn. (Solanac.).
acuminata, Pohl, l. c. t. 1 = B. acuminata.
angusta, Regel, Cat. Pl. Hort. Aksakov. 59, nomen.—Hab ?
australis, Miers, Illustr. ii. 74 = B. Hopeana.
bahiensis, Miers, l. c. = B. bahiensis.
calycina, Hook. Bot. Mag. t. 4553 = B. pauciflora.
capitata, Miers, Illustr. ii. 73 = B. capitata.
confertiflora, T. Moore, Gard. Mag. Bot. ii. (1851) 73 = B. pauciflora.
confertiflora, Pohl, Pl. Bras. Ic. i. 6. t. 5 = B. ramosissima.
divaricata, Pohl, Pl. Bras. Ic. i. 6. t. 6 = B. ramosissima.
elegans, Regel, Cat. Pl. Hort. Aksakov. 59, nomen.—Hab ?
eximia, Lem. Jard. Fleur. iii. (1852–53) t. 249 = *macrantha*.
eximia, Scheidw. ex Moore & Ayres, Gard. Mag. Bot. i. (1850) 16 ; ii. 177.—Bras
gracilis, W. Bakt. in Lond. Hort Brit. Suppl. iii 551, nomen.—Hab ?
grandiflora, Miers, Illustr. ii. 74 = B. grandiflora.
Hopeana, Hook. Bot. Mag. t. 3929 = B. Hopeana.
hydrangeæformis, Pohl, Pl. Bras. Ic. i. 7 = B. hydrangeæformis.
latifolia, Pohl, l. c. 8. t. 9 = B. latifolia.
Lindeniana, Planch. in Belg. Hortic. xv. (1865) 100. —Bras.
macrantha, Lem. Jard. Fleur. iii. (1853) t. 249.—Bras.
macrophylla, Cham. & Schlecht. in Linnæa, ii. (1827) 603 = B. macrophylla.
maritima, Miers, Illustr. ii. 74 = B. maritima.
mutabilis, Hort. ex Regel, in Gartenfl. (1854) 154 = B. acuminata.
obovata, Miers, Illustr. ii. 74 = B. obovata.
pauciflora, Cham. & Schlecht. in Linnæa, ii. (1827) 606 = B. pauciflora.
Publiana, Hort. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 260 = B. ramosissima.
ramosissima, Pohl, Pl. Bras. Ic. i. 5. t. 4 = B. ramosissima.
tenuiflora, Pohl, l. c. 9. t. 1 = B. Hopeana.

FRANCISIA, Endl. Gen. 1226 (1840) = **Darwinia**, Rudge (Myrtac.).

FRANCOA, Cav. in Anal. Cienc. Nat. iv. (1801) 236. *SAXIFRAGEAE*, Benth. & Hook. f. i. 640.
appendiculata, Cav. l. c. 237.—Chili.
appendiculata, A. Juss. in Ann. Sc. Nat. Sér. I. iii. (1824) 196 = sonchifolia.
glabrata, DC. Prod. vii. 772 = ramosa.
ramosa, *D. Don, in Edinb. N. Phil. Journ.* (Oct.–Dec. 1828) 52.—Chili.
rupestris, Poepp. ex Kunze, in Flora, xiv. (1831) 374 = sonchifolia.
sonchifolia, *Cav. Ic.* vi 77, in nota.—Chili.
sonchifolia, Lodd. Bot Cab. t. 1864 = appendiculata.

FRANCOEURIA, Cass. in Dict. Sc. Nat. xxxiv. 44 (1825) = **Pulicaria**, Gaertn. (Compos.).
crispa, Cass. l. c. xxxvii. 374 = P. crispa.
diffusa, Shuttl. ex Brunn. in Flora, xxiii. (1840) t. Beibl. 72 = P. crispa.
lacinata, Cass. & Duc. in Bull. Soc. Bot. Fr. iv. (1857) 181.—Algeria.

FRANGULA, Tourn. ex Hall. Enum. Helv. i. 164 (1742); et Mill. Gard. Dict. ed. VI. (1752) = **Rhamnus**, Linn.
Alnus, Mill. i. c. ed. VIII. n. 1 = R. Frangula.
americana, Mill. l. c. n. 4 = R. alnifolia.
californica, A. Gray, Gen. Illustr. ii. 178 = R. californica.
caroliniana, A. Gray, l. c. = R. caroliniana.
chrysophylla, Relsack, in Mart. Fl. Bras. xi. t. 91.—Bras.
crenata, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 82 = R. crenata.
fragilis, Salis. Fl. Ludov. 97 = R. caroliniana.
latifolia, Mill. Gard. Dict. ed. VIII. n. 2 = R. alpina.
pentapetala, Gilib. Exercit. i. 280 = R. Frangula.
polymorpha, Reissek, in Mart. Fl. Bras. xi i 91 = R. sectipetala.
Purshiana, Cooper, in Pacif. Rail. Rep. xii. 29, 57 = R. Purshiana.
rotundifolia, Mill. Gard. Dict. ed. VIII. n. 3 = R. pumila.
rupestris, Schur, Enum. Pl. Transs. 142 = R. rupestris.
saxatilis, Cramer, Disp. Syst. Pl. 41 = R. saxatilis.
sphærocarpa, Griseb. Fl. Brit. W. Ind. 99 = R. sphærospermus.
vulgaris, Reichb. Fl. Germ. Excurs. 488 = R. Frangula.
Wulfenii, Reichb. l. c. = R. rupestris.

FRANKA, Steud. Nom. ed. II. i. 645, in syn. (1840) = seq.

FRANKENIA, Linn. Gen. ed. I. 129 (1737). *FRANKENIACEAE*, Benth. & Hook. f. i. 141.
BEATSONIA, Roxb. in Beats. St. Hel. App. 300 (1816).
FRANCA, Mich. ex Adans Fam. ii. 255 (1763)
HYPERICOPSIS, Boiss. Diagn. Ser. I. vi. (1845) 25.
MERETHO, Rafin Fl. Tellur. ii. 93 (1836).
NOTHRIA, Berg, Desc. Pl. Cap. 171 (1767).
STREPTIMA, Rafin. Fl. Tellur. ii. 93 (1836).
asperx, *Phil. Fl. Atac.* 10.—Chili.
Auckeri, Jaub. & Spach, Illustr. ii. 105. t. 187 = hirsuta.
Beatsonia, Schult. f. Syst. vii. 70 = portulacæfolia.
Bertoroana, *C. Gay, Fl. Chil.* i. 347.—Chili.
Boissieri, *Reut. ex Boiss. Voy. Bot. Espagne*, ii. 721.—Hispan.
bractesta, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1854) ii. 366.—Austral.
campestris, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 480.—Chili.
canescens, Presl, Delic. Prag. 61 = pulverulenta.
canescens, Willd. ex Schult. f. Syst. vii. 68 = hirsuta.
capitata, Webb & Berth. Phyt. Canar. i. 131 = hirsuta.
cespitosa, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) minus 28.—Madera.
chilensis, *Presl, in Schult. f. Syst.* vii. 1618.—Chili.
cineres, *A. DC. in Linnæa*, xxv. (1852) 576.—Austral.
corymbosa, Desf. Fl. Atlant. i. 315. t. 98 = hirsuta.
cymbifolia, Hook. lc. Fl. t. 965 = Wilsonia humilis.
Drummondii, *Benth. Fl. Australi.* i. 152.—Austral.
erecta, *C. Gay, Fl. Chil.* i. 346.—Chili.
ericæfolia, Salisb. Prod. 214 = hirsuta.

FRANKENIA :—
ericifolia, *C. Sm. in Buch, Beschr. Canar. Ins.* 154, nomen ; *DC. Prod.* i. 360.—Ins. Canar.
ericoides, Pers. Syn. i. 393 = hirsuta.
farinosa, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 236.—Chili.
floribunda, *Pomel, Nouv. Mat. Fl. Atl.* 343.—Alger.
florida, Phil. in Anal. Univ. Chil. (1872) 676 = chilensis.
fruticulosa, DC. Prod. i. 350 = pauciflora.
glabrata, *Phil. in Anal. Univ. Chil.* (1872) 677.—Chili
glomerata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii (1854) ii. 366.—Austral.
glomerulata, Coss. Notes Crit. ii. 30 = Boissieri.
grandiflola, *Cham. & Schlecht. in Linnæa*, i. (1826) 36.—Am. bor. occ.
Kvæbvi, Cham. & Schlecht. in Linnæa, i. (1826) 36 = hirsuta.
laevis, Habl. Taur. 153 ; ex Bieb. Fl. Taur. Cauc. i. 287 = pulverulenta.
laevis, Linn. Sp. Pl. 331 = hirsuta.
laevis, F. Muell. Fl. Vict. i. 82 = pauciflora.
leptis, Presl, in Schult. f. Syst. vii. 1620 = grandifolia.
micrantha, *C. Gay, Fl. Chil.* i 247.—Chili.
microphylla, *Cav. Ic. vi.* 76.—Am. austr.
mollis, Bieb. Fl. Taur. Cauc. iii. 276 = Cressa cretica.
neglecta, *Schult. f. Syst.* vii. 1618.—Oriens.
Nicolitiana, *C. Gay, Fl. Chil.* i. 348.—Chili.
nodiflora, *Lam. Tabl. Encyc.* ii. 396.—Afr. austr.
Nothria, Thunb. Prod. Pl. Cap. 58 = hirsuta.
pallida, Boiss. Diagn. Ser. II. i. 61 = hirsuta.
Palmeri, *S Wats. in Proc. Am. Acad.* xi. (1876) 124.—Calif.
parvula, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 368.—Austral.
pauciflora, *DC. Prod.* i. 350.—Austral.
persica, *Yaub. & Spach, Illustr.* ii. 106.—Persia.
pilosa, Pall. Reise, iii. 317 = hirsuta.
portulacæfolia, *Spreng. Syst.* ii. 184.—Ins. St. Helena.
pulverulenta, *Linn. Sp. Pl.* 332.—Reg. Mediterr. ; Orient ; Ind. or.
punctata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 367.—Austral.
Reuteri, Boiss. Diagn Ser. II. i. 60 = hirsuta.
revoluta, Forsk. Fl. Aegypt. Arab. 75 = hirsuta.
scabra, Lindl. in Mitch. Journ. Trop. Austral. 305 = pauciflora.
serpyllifolia, Lindl. in Mitch. Journ. Trop. Austral. 306 = pauciflora.
strigosa, *Presl, Fl. Sic.* i. 141.—Creta ; Aegypt.
tetrapetala, *Labill. Nov. Holl. Pl.* i. 66. t. 114.—Austral.
thymifolia, Desf. Fl. Atlant. i. 316 = hirsuta.
triandra, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 237.—Bolivia.
velutina, Brouss. ex DC. Prod. i. 350 = hirsuta.
Vidali, *Phil. in Anal. Univ. Chil.* (1875) 187.—Chili.
virens, Desf. Tabl. ed. I. 186, nomen ; Cat. Hort. Par. ed. III. 407 = pauciflora.
Webbii, Boiss. & Reut. Pugill. Pl. Nov. 16 = hirsuta.

FRANKIA, Bert. ex Steud. Nom. ed. II. i. 646 (1840) = **Phyllanthus**, Linn. (Euphorb.).
ramiflora, Bert. ex Steud. l. c. = P. Cicca.

FRANKIA, Steud. ex Steud. Nom. ed. II. i. 646 (1840) = **Gymnarrhena**, Desf. (Compos.).
Schimperi, Steud. l. c. = G. micrantha.

FRANKLANDIA, R. Br. in Trans. Linn. Soc. x. (1810) 157. *PROTEACEAE*, Benth. & Hook. f. iii. 174.
fucifolia, *R. Br. l. c.*—Austral.
armifolia, Steud. Nom. ed. II. i. 646, sphalm. = præc.
triandra, *Benth. Fl. Australi.* v. 277.—Austral.

FRANKLINA, J. F. Gmel. Syst. 810 (1791) = seq.

FRANKLINIA, Bartr. ex Marsh. Arbust. Am. 48 (1785) = **Gordonia**, Ellis (Ternstroem.).
Alatama, Rafin. Atl. Journ. 79 = G. pubescens.
Alatamaha, Marsh. Arbust. Am. 49 = G. pubescens.
americana, Marsh. ex Pers. Syn. ii. 259 = G. pubescens.

FKANKOA, Reichb. Handb. 230 (1837) = **Francoa**, Cav. (Saxifrag.).

FRANKOERIA, Steud. Nom. ed. II. i. 646 (1840) = FRANCOEURIA, Cass. = **Pulicaria**, Gaertn. (Compos.).

FRANQUEVILLEA, Zoll. ex Miq. Fl. Ind. Bat. iii. 586 (‘1855’) = **Hypoxis**, Linn. (Amaryllid.).

FRANQUEVILLIA, Saltab. ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 338 (1821) = **Microcala**, Hoffmgg. & Link (Gentian.).
minima, S. F. Gray, l. c. = M. filiformis.

FRANSERIA, Cav. Ic. ii. 78. t. 200 (1793). *COMPOSITAE.* Benth. & Hook. f. ii. 354.
GAERTNERIA, Medic. Phil. Bot. i. 45 (1789).
HEMIAMBROSIA, Delpino, Studi sopra Artemis, 57 (1871).
HEMIXANTHIDIUM, Delpino, l. c. 60 (1871).
XANTHIDIUM, Delpino, l. c. 62 (1871).
albicaulis, Torr. Pl. Frémont. 16 = dumosa.
ambrosioides, *Cav. Ic.* ii. 79. t. 200.—An. bor. occ.
artemisioides, *Willd. Sp. Pl.* iv. 574.—Peruv.
bipinnatifida, A. Gray, in Proc. Am. Acad. xi. (1876) 116 = camphorata.
bipinnatifida, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 344.—Am. bor. occ.
camphorata, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 199.—Calif.
Chamissonis, *Less. in Linnaea*, vi. (1831) 507.—Am. bor. occ.
chenopodifolia, *Benth. Bot. Voy. Sulph.* 26.—Calif.
cordifolia, *A. Gray, Syn. Fl. N. Am.* i. ii. 445.—Arizona.
runcifolia, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 345 = Chamissonis.
deltoides, *Torr. Pl. Fremont.* 15.—Arizona; Calif.
discolor, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 345.—Am. bor. occ.
dumosa, *A. Gray, in Rep. Prém. 2nd Exped.* 316.—Am. bor. occ.
ericoenira, *A. Gray, in Proc. Am. Acad.* vii. (1868) 356.—Am. bor. occ.
flexuosa, *A. Gray, l. c.* xx. (1885) 298.—Calif.
hispida, *Benth. Bot. Voy. Sulph.* 25.—Calif.
Hookeriana, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 345.—Am. bor. occ.
ilicifolia, *A. Gray, in Proc. Am. Acad.* xi. (1876) 77. —Am. bor. occ.
Lessingii, Meyen & Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 368 = bipinnatifida.
Meyeniana, Sch. Bip. in Bonplandia, iv. (1856) 54 = Ambrosia artemisioides.
montana, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 345 = Hookeriana.
pumila, Nutt. l. c. 344 = Ambrosia pumila.
tacorensis, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 368.—Peruv.
tenuifolia, *Haw. & Gray, in A. Gray, Pl. Fendl.* 80. —Am. bor. occ.
tumentosa, *A. Gray, l. c.*—Am. bor. occ.

FRAPPIERIA, Cordem. in Baill. Adansonia, x. (1871) 24 = **Faidia**, Jacq. (Compos.).
littoralis, Cordem. in Baill. l. c. 25.—Ins. Borb.
montana, Cordem. in Baill. l. c. 26.—Ins. Borb.
thermalis, Cordem. in Baill. l. c. 27.—Ins. Borb.

FRASERA, Walt. Fl. Carol. 87 (1788). *GENTIANEAE*, Benth. & Hook. f. ii. 817.
TESSARANTHIUM, Kellogg, in Proc. Calif. Acad. ii. (1863) 142.
albicaulis, *Griseb. Gen. et Sp. Gent.* 330.—Am. bor. occ.
albomarginata, *S. Wats. in Bot. King's Exped.* 280.—Am. bor. occ.
carolinensis, Hook. Fl. Bor. Am. ii. 66 = thyrsiflora.
carolinensis, *Walt. Fl. Carol.* 88.—Am. bor.
nitida, *Benth. Pl. Hartw.* 332.—Am. bor. occ.

FRASERA:—
paniculata, *Torr. in Pacif. Rail. Rep.* iv. 126.—Am. bor. occ.
Parryi, *Torr. Bot. Mex. Bound.* 156.—Am. bor. occ.
speciosa, *Dougl. ex Hook. Fl. Bor. Am.* ii. 66. t. 158.—Am. bor. occ.
thyrsiflora, *Hook. in Kew Journ.* iii. (1851) 288.—Am. bor. occ.
verticillata, *Rafin. Med. Fl.* i. 196.—Am. bor.
Walteri, Michx. Fl. Bor. Am. i. 97 = carolinensis.

FRAUNHOFERA, Mart. Nov. Gen. et Sp. iii. 85. t. 235 (1829). *CELASTRINEAE*, Benth. & Hook. f. i. 366.
multiflora, *Mart. l. c.* 86.—Bras.

FRAXIMA, Rafin. Fl. Tellur. iv. 83 (1836) = **Ipomoea**, Linn. (Convolv.).
ebracteata, Rafin. l. c. = I. ebracteata.
mindanensis, Rafin. l. c. = Jacquemontia hirsuta.
quinquefolia, Rafin. l. c. = I. quinquefolia.
sagittifolia, Rafin. l. c. = I. sagittata.
Sherardi, Rafin. l. c. = Convolvulus hirsutus.
tridentata, Rafin. l. c. = I. trilobiata.
umbellata, Rafin. l. c. = I. umbellata.

FRAXINELLA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 291 (1745) = **Dictamnus**, Linn. (Rutac.).
alba, Gaertn. Fruct. i. t. 69 = D. albus.
Dictamnus, Moench, Meth. 68 = D. albus.

FRAXINOIDES, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 198; Phil. Bot. ii. 38 (1791) = **Fraxinus**, Tourn. (Oleac.).
alba, Medic. Pfl. Anat. 290 = Fraxinus americana.
nigra, Medic. l. c. = Fraxinus nigra.

FRAXINUS, Tourn. ex Linn. Syst. ed. I (1735). *OLEACEAE*, Benth. & Hook. f. ii. 676.
APILIA, Rafin. Alsog. Am. 39 (1838).
APILIA, Rafin. New Fl. Am. iii. 93 (1836).
CALLICELLA, Kostel. Allg. Med.-Pharm. Fl. iii 1008 (1834).
FRASINOIDES, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 198; Phil. Bot. ii. 38 (1791).
LEPTALIX, Rafin. New Fl. Am. iii. 93 (1836).
MANNAPHORUS, Rafin. in Am. Monthly Mag. (1818) 178.
MELIOPSIS, Reichb. Nom. 135 (1841).
ORNANTHES, Rafin. New. Fl. Am. iii. 93 (1836).
ORNUS, Medic. Theor. Elem. ii. 275 (1790).
SAMARAPSIS, Rafin. New Fl. Am. iii. 93 (1836).
acuminata, Lam. Encyc. ii. 547 = americana.
alba, Marsh. Arbust. Am. 51; Bosc, in Mém. Instit. Fr. ix. 1808 (1811) 207 = americana.
albicans, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 4 = americana.
americana, *Linn. Sp. Pl.* 1057.—Am. bor.
angustifolia, Reut. ex Nym. Consp. 495 = excelsior.
angustifolia, *Vahl, Enum.* i. 32.—Tauris.
anomala, *Torr. ex S. Wats. in Bot. King's Exped.* 283. —Utah.
apetala, Lam. Fl. Fr. ii. 525 = excelsior.
appendiculata, Pers. Syn. ii. 605 = excelsior.
argentea, Loisel. Fl. Gall. iii. 227 = F. Ornus.
aria, Hort. ex Dum.-Cours. Bot. Cult. ed. II. ii. 585 = excelsior.
atrovirens, Desf. Hist. Arb. i. 104 = excelsior.
aurea, Willd. Enum. Hort. Berol. 1059 = excelsior.
australis, J. Gay, ex C. Koch, Dendrol. ii. 1. 247 = excelsior.
australis, Mont. ex Gren. & Godr. Fl. Fr. ii. 472 = oxyphylla.
Berlandieriana, A. DC. in DC. Prod. viii. 276 = viridis.
bifolia, Gren. & Godr. Fl. Fr. ii. 472 = oxyphylla.
Bosci, G. Don, Gen. Syst. iv. 55 = excelsior.
Bungeana, *A. DC. in DC. Prod.* viii. 275.—China.
canadensis, Gaertn. Fruct. i. 223 = americana.
canaliculata, *Rafin. Alsog. Am.* 85. (Quid?).—Am. bor.
cappadocica, Juss. ex Bosc, in Mém. Inst. Fr. ix. 1808 (1811) 214 = F. Ornus.
carolinensis, Mill. Gard. Dict. ed. VIII. n. 6; Lam. Encyc. ii. 548 = platycarpa.
caroliniana, Wangenh. ex DC, Prod. viii. 278 = americana.

FRAXINUS:—
caroliniana, Willd. ex Pursh, Fl. Am. Sept. i. 9 = viridis.
cerasifolia, Hoffmgg. Verz. Pfl. Nachtr. ii. 29 = pubescens.
chinensis, Roxb. Fl. Ind. ed. Carey, i. 150 = xanthoxyloides.
cineres, *Bosc, in Mém. Instit. Fr.* ix. 1808 (1811) 207. —Am. bor.
concolor, Mahl. Cat. Pl. Am. Sept. iii. 115. t. 11 = viridis.
cordata, *Rafin. Alsog. Am.* 34 (Quid?).—Am. bor.
coriacea, S. Wats. in Am. Natural. vii. (1873) 302 = platycarpa, americana.
coriariaefolia, Scheele, in Linnaea, xvii. (1843) 350 = excelsior.
crispa, Bosc, in Mém. Inst. Fr. ix. 1808 (1811), 214 = excelsior.
cubensis, *Griseb. Cat. Pl. Cub.* 170.—Cuba.
Curtisii, Vasey, Cat. Trees U. St. 20 = americana.
curvidens, Hoffmgg. Verz. Pfl. Nachtr. ii. 118 = platycarpa.
cuspidata, *Torr. Bot. Mex. Bound.* 156.—N. Mexic.
dentata, *Tausch, in Flora*, xxi. (1838) 761 (Quid?).—Sicily.
dimorpha, Coss. & Dur. in Bull. Soc. Bot. Fr. ii. (1855) 367 = xanthoxyloides.
diptetala, *Bosc, in Act. Soc. Linn. Bord. Voy.* 362. t. 87. —Calif.
discolor, Muhl. Cat. Pl. Am. Sept. 111 = americana.
diversifolia, Ait. Hort. Kew. iii. 445 = excelsior.
diversifolia, Rochel, ex Boiss. Fl. Orient. iv. 89 = F. Ornus.
elliptica, Bosc, in Mém. Instit. Fr. ix. 1808 (1811) 204 = pubescens.
epiptera, Michx. Fl. Bor. Am. ii. 256 = americana.
erosa, Pers. Syn. i. 604 = excelsior.
excelsa, Thunb. Fl. Jap. 23 = excelsior.
excelsior, Boiss. Voy. Bot. Espagne, 408 = angustifolia.
excelsior, Bové, ex A. DC. in DC. Prod. viii. 276 = oxyphylla.
excelsior, *Linn. Sp. Pl.* 1057.—Europ.; As. bor.; Orient.; Reg. Himal.
excelsior, Wall. Fl. Carol. 254 = platycarpa?
expansa, Willd. Berl. Wilde Baumz. ed. II. 150 = viridis.
floribunda, Bunge, ex A. DC. in DC. Prod. viii. 275 = Bungeana.
floribunda, *Wall. Cat.* n. 2886; *et in Roxb. Fl. Ind. ed. Carey*, i. 150.—Reg. Himal.
florifera, Scop. Fl. Carn. ed. II. ii. 382 = F. Ornus.
fungosa, Lodd. Cat. (1836) = excelsior.
fusca, *Bosc, in. Mém. Instit. Fr.* ix. 1808 (1811) 208. —Am. bor.
glauca, *Rafin. Alsog. Am.* 38 (Quid?).—Am. bor.
grandifolia, *Rafin. l. c.* 38.—Am. bor.
Greggii, *A. Gray, in Proc. Am. Acad.* xii. (1877) 63. —Am. bor. occ.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 605.—Reg. Himal.
halepensis, Asch. ex Steud. Nom. ed. II. i. 647 = rotundifolia.
heterophylla, Vahl, Enum. i. 33 = excelsior.
Hookeri, Wenzig, in Engl. Bot. Jahrb. iv. (1883) 179 = excelsior.
horizontalis, Dum.-Cours. Bot. Cult. ed. II. ii. 578 = excelsior.
implicata, Dum.-Cours. l. c. 577 = excelsior.
integrifolia, Moench, Verz. Ausl. Baume, 48 = excelsior.
japonica, Bunger, ex Wenzig, in Engl. Bot. Jahrb. iv (1883) 170 = Bungeana.
jaspidea, Dum.-Cours. Bot. Cult. ed. II. ii. 577 = excelsior.
juglandifolia, Lam. Encyc. ii. 548 = americana.
lacera, *Rafin. Fl. Ludov.* 82 (Quid?).—Am. bor.
lacerata, Dum.-Cours. Bot. Cult. ed. II. ii. 577 = excelsior.
lancea, Bosc, in Mém. Instit. Fr. ix. 1808 (1811) 209 = pubescens.
lanceolata, Borkh. Handb. Forsth. i. 696 = americana.
lanceifolia, Rafin. Alsog. Am. 35 = pruc.?
latifolia, Benth. Bot. Voy. Sulph. 25 = oregona.
lentiscifolia, Desf. Tabl. ed. I. 59; Hist. Arb. i. 102 = oxyphylla.
longicuspis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 189.—Japon.

FRAXINUS:—
longifolia, Bosc, in Mém. Instit. Fr. ix. 1808 (1811) 209 = pubescens.
macrophylla, Hoffmgg. Vers. Pfl. Nachtr. ii. 120 = americana.
mammifera, Steud. Nom. ed. I. 346, sphalm. = seq.
mandshurica, *Rupr. in Bull. Sc. Phys.-Math. Acad. Pétersb.* xv. (1857) 371.—Manchuria.
mannifera, Hort. ex Steud. Nom. ed. II. i. 647 = F. Ornus.
Marisetti, *Hook. f. Bot. Mag.* t. 6678.—China.
media, *Rafin. Alsog. Am.* 36 (Quid !).—Am. bor.
mexicana, Sweet, Hort. Brit. ed. II. 351 = cuspidata.
microphylla, Willd. Enum. Hort. Berol. Suppl. 69 = oxyphylla.
mille-lacum, C. Koch, in Linnæa, xxiii. (1850) 597 = F. Ornus.
mixta, Bosc, in Mém. Instit. Fr. ix. 1808 (1811) 213 = parvifolia.
mollis, *Rafin. Alsog. Am.* 34 (Quid ?).—Am. bor.
monophylla, Desf. Hist. Arb. i. 102 = excelsior.
Montagnei, Nym. Consp. 195 = oxyphylla.
montana, Salisb. Prod. 14 = *F.* Ornus.
Moorcroftiana, Brand. For. Fl. 304 = xanthoxyloides.
Moorcroftiana, Wall. Cat. n. 2834 = excelsior.
multifolia, *Rafin. Alsog. Am.* 37 (Quid !).—Am. bor.
nana, Pers. Syn. ii. 605 = excelsior.
nigra, *Bosc, in Mém. Instit. Fr.* ix. 1808 (1811) 203.—Am. bor.
nigra, Marsh. Arbust. Am. 51 = sambucifolia.
nigra, Pott, in Durol, Harbk. Baums. ed. II. i 398 = pubescens.
nigrescens, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 5 = platycarpa.
Novæ-Angliæ, Mill. Gard. Dict. ed. VIII. n. 5 = pubescens.
Novæ-Angliæ, Wangenh. ex DC. Prod. viii. 277 = americana.
Nuttallii, Buckl. in Proc. Acad. Sc. Philad. 1860 (1861) 444 = platycarpa.
obliqua, Tausch, in Flora, xvii. (1834) 521 = oxyphylla
oblongocarpa, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 4 = pubescens.
obovata, Blume, Mus. Bot. Lugd. Bat. i. 311 = Bungeana.
oregona, *Nutt. Am. Sylva*, iii. 59. t. 99.—Am. bor. occ.
Ornus, *Linn. Sp. Pl.* 1057.—Reg. Mediterr. ; Orient.
Ornus, Pall. ex Bieb. Fl. Taur. Cauc. ii. 451 = oxyphylla.
Ornus, Scop. Fl. Carn. ed. II. ii 281 = excelsior.
ovalis, Willd. Enum. Hort. Berol. Suppl. 69 = sec.
ovata, Bosc, in Mém. Instit. Fr. ix. 1808 (1811) 204 = pubescens.
oxycarpa, Willd. Sp. Pl. iv. 1100 = oxyphylla.
oxyphylla, *Bieb. Fl. Taur. Cauc.* ii. 450.—Tauria.
oxyphylla, Bourg. ex Nym. Consp. 195 = excelsior.
pallida, Bosc, in Mém. Instit. Fr. ix. 1808 (1811) 201 = platycarpa.
palustris, *Rafin. Alsog. Am.* 35 (Quid ?).—Am. bor.
paniculata, Mill. Gard. Dict. ed. VIII. n. 4 = F. Ornus.
pannosa, *Vent. ex Sprengl. Syst.* i. 95.—Am. bor.
parvifiora, Auch. ex DC. Prod. viii. 276 = oxyphylla.
parvifiora, Tenore, Syll. Fl. Neap. 11 = rotundifolia.
parvifolia, *Lam. Encyc.* ii. 546.—Calabr. ; Sicil.
parvifolia, Willd. Sp. Pl. iv. 1101 = excelsior.
pauciflora, Nutt. Am. Sylva, iii. 61. t. 100 = platycarpa.
pendula, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 445 = excelsior.
præcocissima, Marsh. Arbust. Am. 51 = pubescens.
pubescens, Lam. Encyc. i. 547 = oxyphylla.
pubescens, Poir. in Duham. Arb. ed. Nov. iv. 66 = excelsior.
quadrangulata, Michx. *Fl. Bor. Am.* ii. 255.—Am. bor.
raibocarpa, *Regel, in Act. Hort. Petrop.* viii. (1884) 665.—Turkestan.
repanda, *Rafin. Alsog. Am.* 37 (Quid ?).—Am. bor.
reticulata, *Rafin. l. c.* 34 (Quid ?).—Am. bor.
retusa, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 330.—Hongkong.
retusa, *Rafin. Alsog. Am.* 34 (Quid ?).—Am. bor.
rhynchophylla, Hance, in Journ. Bot. vii. (1869) 164 = chinensis.
Richardi, *Bosc, in Mém. Instit. Fr.* ix. 1808 (1811) 207.—Am. bor.
rostrata, Gmel. Fl. Bad. 374. t. 63 = oxyphylla.
rotundifolia, Lam. Encyc. ii. 546 = F. Ornus.
rotundifolia, *Mill. Gard. Dict.* ed. VIII. n. 2.—Italia ; Græc. ; As. Min.
rubicunda, *Bosc, in Mém. Instit. Fr.* ix. 1808 (1811) 212.—Am. bor.
rufa, *Bosc, l. c.* 208.—Am. bor.
salicifolia, Hort. ex Steud. Nom. ed. II. i. 647 = excelsior.
sambucifolia, *Lam. Encyc.* ii. 549.—Am. bor.
Schiedeana, *Cham. & Schlecht. in Linnæa*, vi. (1831) 391.—Mexic.
serratifolia, Michx. ex Steud. Nom. ed. II. i. 647 = platycarpa.
Sieboldiana, *Blume, Mus. Bot. Lugd. Bat.* i. 311.—Japonia.
simplicifolia, Willd. Berl. Wilde Baumz. 121 = excelsior.
sogdiana, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 590.—Turkestan.
Steudelii, Medic. Beobacht. (1782) 201 = excelsior.
striata, Dum.-Cours. Bot. Cult. ed. II. ii. 577 = excelsior.
striata, Spreng. Syst. i. 95 = sec.
strigata, *Bosc, in Mém. Instit. Fr* ix. 1808 (1811) 217.—Orient.
subpubescens, Pers. Syn. ii. 605 = pubescens.
subrillosa, Bosc, ex Pers. l. c. 604 = pubescens.
sylvatica, Gromow, ex Trautv. in Act. Hort. Petrop. ix. (1884) 11, nomen = excelsior !
syriaca, Boiss. Diagn. Ser. I. xi. 77 = oxyphylla.
tamaricifolia, Vahl, Enum. i. 52 = oxyphylla.
teres, *Rafin. Alsog. Am.* 36 (Quid ?).—Am. bor.
tetragona, Hort. Par. ex Duham. Arb. ed. Nov. iv. 64 = quadrangulata.
Theophrasti, Duham. ex Steud. Nom. ed. I. 346 = F. Ornus.
thryxantha, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = F. Ornus.
tomentosa, Michx. Arb. Am. Sept. iii. 113 = pubescens.
trifoliata, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 5 = viridis.
triptera, Nutt. Gen. Am. ii. 239 = platycarpa.
umbilata, *Rafin. Fl. Ludov.* 88 (Quid ?).—Am. bor.
urophylla, Wall. Cat. n. 2835 = floribunda.
velutina, Torr. in Emory, Notes Mil. Reconnois. 149 = pistaciæfolia.
verrucosa, Pers. Syn. ii. 604 ; Dum.-Cours. Bot. Cult. ed. II. ii. 577 = excelsior.
villosa, Hort. ex Dum.-Cours. l. c. 582 = americana.
viridis, *Bosc, in Mém. Instit. Fr.* ix. 1808 (1811) 209.—Am. bor.
viridis, *Michx. f. Syls. Am.* t. 120.—Am. bor.
xanthoxyloides, *Wall. Cat.* n. 2833.—Reg. Himal.

FREDERICIA, G. Don, Gen. Syst. iv. 227 (1837) = **Fridericia**, Mart. (Bignon.).

FREDOLIA, Coss. & Dur. ex Moq. & Coss. in Bull. Soc. Bot. Fr. ix. (1862) 801 = **Anabasis**, Linn. (Chenopod.).
arctioides, Coss. & Dur. ex Bunge, Anabas. Rev. 35 = *A.* arctioides.

FREEMANIA, Bot. ex DC. Prod. vi 218 (1837) = **Helichrysum**, Vaill. (Compos.).
hypnoides, Bot. ex DC. l. c. (= *Aphelexis hypnoides*). —Madag.
hypnoides, Steud. Nom. ed. II. i. 647 = præc.

FREEMANNIA, Steud. Nom. ed. II. i. 647 (1840) = præc.

FREESEA, Eckl. Verz. 30 (1827) = **Ixia**, Linn. etc (Irid.).
costata, Eckl. ex Baker, in Journ. Linn. Soc. xvi. (1877) 160 = Acidanthera tubulosa.
crispa, Eckl. Verz. 30 = Tritonia crispa.
flava, Eckl. l. c. = Tritonia flava.
longiflora, Eckl. l. c. = I. paniculata, *Delar.*
minuate-laterita, Eckl. l. c. = Tritonia crocata.
secunda, Eckl. l. c. = Tritonia secunda.

FREESIA, Benth. & Hook. f. Gen. iii. 704 (1883) = præc.

FREESIA, Klatt, in Linnæa, xxxiv. (1865–66) 672. *IRIDEAE*, Benth. & Hook. f. iii. 704.
Leichtlini, *Klatt, in Regel, Gartenfl.* (1874) 289.—Afr. austr.
odorata, Eckl. ex Klatt, in Linnæa, xxxiv. (1865–66) 672 = refracta.
refracta, *Klatt, l. c.* 673.—Afr. austr.
xanthospila, *Klatt, l. c.*—Afr. austr.

FREGEA, Reichb. f. in Bot. Zeit. x. (1852) 712 *ORCHIDEAE* [Benth. & Hook. f. iii. 590].
amabilis, *Reichb. f. l. c.*—Panama.

FREGIRARDIA, Dun. Ind. Sem. Monsp. (1849).—Cf Ann. Sc. Nat. Sér. III. xii. (1849) 366 = **Brachistus**, Miers (Solanac.).
angustifolia, Dun. in DC. Prod. xiii. 1. 506.—Ecuador.
ciliata, Dun. l. c. 505 = B. ciliatus.
dumetorum, Dun. l. c. 504 = B. dumetorum.
Dunaliana, Wedd. Chlor. And. ii. 100.—Peruv.
ervoloryna, Dun. in DC. Prod. xiii. 1. 503.—Bolivia.
leptoclada, Dun. l. c. 505.—Bras.
ligustrina, Dun. l. c. 507.—Mexic.
lindeni, Dun. l. c. 503.—Venezuela.
luteiflora, Dun. in Ind. Sem. Hort. Monsp. (1842).—Am. austr.
mollis, Dun. in DC. Prod. xiii. 1. 505 = B. mollis.
rhomboidea, Dun. l. c. 504 = B. rhomboideus.
riparia, Dun. l. c. 503 = B. riparius.
Spina-alba, Dun. l. c. 507.—Bras.
Vargasii, Dun. l. c.—Venezuela.

FREIRA, C. Gay, Fl. Chil. v. 366 (1849), sphalm. = seq.

FREIREA, Gaudich. in Freyc. Voy. Bot. 502 (1826) = **Paricteria**, Tourn. (Urtic.).
alsinæfolia, Gaudich. l. c. = P. alsinefolia.
alsinefolia, Schimp. ex Blume, Mus. Bot. Lugd. Bat. ii. 255 = P. debilis.
australis, Nees, in Lehm. Pl. Preiss. i. 636 = P. debilis.
erecta, Phil. Fl. Atac. 49 = P. debilis.
fernandeziana, Wedd. in Ann. Sc. Nat. Prod. xvi. 1. 236*, in syn. = P. debilis.
humifusa, C. Gay, Fl. Chil. v. 366 = P. debilis.
oxymifolia, Gaudich. in Freyc. Voy. Bot. 502 = P. mauritanica.
Studelii, Blume, Mus. Bot. Lugd. Bat. ii. 252 = P. debilis.

FREIREODENDRON, Muell. Arg. in DC. Prod. xv. II 244 (1866) = **Drypetes**, Vahl (Euphorb.).
sessiliflorum, Muell. Arg. l. c. 245 = D. sessiliflora.

FREMONTEA, Lindl. Veg. Kingd. 518 (1847) = seq.

FREMONTIA, Torr. in Emory, Rep. 95, 217. t. 3 (1845) = **Sarcobatus**, Nees (Chenop.).
vermicularis, Torr. l. c. = S. Maximiliani.

FREMONTIA, Torr. in Smithson, Contrib. vi. (Pl. Frémont.) 5. t. 2 (1854). *STERCULIACEAE*, Benth. & Hook. f. i. 219, 982.
californica, *Torr. l. c.* 6.—Calif.

FREMYA, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 373 = **Xanthostemon**, F. Muell. (Myrtac.).
auraniaca, Brongn. & Gris, l. c. 373.—N. Caled.
ciliata, Brongn. & Gris, l. c. 374.—N. Caled.

FREMYA :—
Deplanchei, Brongn. & Grls, in Bull. Soc. Bot. Fr. x. (1863) 378.—N. Caled.
elegans, Brongn. & Grls, l. c. 374.—N. Caled.
flava, Brongn. & Grls, l. c. 378.—N. Caled.
myrtifolia, Brongn. & Grls, l. c. 399.—N. Caled.
Pancheri, Brongn. & Grls, l. c. 373 = X. Pancheri.
pubescens, Brongn. & Grls, l. c.—N. Caled.
rubra, Brongn. & Grls, l. c. 373 = X. rubrum.
speciosa, Brongn. & Grls, l. c. 399.—N. Caled.
Vieillardi, Brongn. & Grls, l. c. 378.—N. Caled.

FRENELA, Mirb. in Mém. Mus. Par. xiii. (1825) 80, partim = Callitris, Vent. (Conif.).
Actinostrobus, F. Muell. Sel. Pl. Indust. Cult. 94 = Actinostrobus pyrmidalis.
arenaca, Hort. ex Endl. Syn. Conif. 38 = C. rhomboidea.
attenuata, A. Cunn. ex Benth. Fl. Austral. vi. 238 = C. rhomboidea.
australis, N. Br. ex Mirb. in Mém. Mus. Par. xiii. (1825) 74 = C. oblonga.
australis, Endl. Syn. Conif. 37 = C. rhomboidea.
Balansæ, Brongn. & Grls, in Bull. Soc. Bot. Fr. xvi. (1869) 237.—N. Caled.
calcarata, A. Cunn. ex Mirb. in Mém. Mus. Par. xiii. (1825) 74 = C. calcarata.
cancerra, Parl. in DC. Prod. xvi. II. 448 = C. robusta.
columellaris, F. Muell. Fragm. v. 198 = C. robusta.
crassivalvis, Miq. Stirp. Nov. Holl. Muell. Collect. 1 = C. robusta.
Drummondii, Parl. in Journ. Bot. i. (1863) 35 = C. Drummondii.
Endlicheri, Parl. in DC. Prod. xvi. II. 449 = C. calcarata.
ericoides, Hort. ex Endl. Syn. Conif. 38 = C. calcarata.
Fontanesii, Mirb. in Mém. Mus. Par. xiii. (1825) 74 = C. quadrivalvis.
Fothergilli, Hort. ex Endl. Syn. Conif. 38 = C. Gunnii.
fruticosa, Endl. l. c. 36 = C. calcarata.
glauca, Mirb. in Mém. Mus. Par. xiii. (1825) 74 = C. robusta.
Gulielmi, Parl. in DC. Prod. xvi. II. 449 = C. robusta.
Gunnii, Endl. Syn. Conif. 38 = C. oblonga.
Hugelii, Hort. ex Carr. Conif. 73 = C. Hugelii.
intratropica, F. Muell. ex Benth. Fl. Austral. vi. 237 = C. robusta.
juniperoides, Parl. in DC. Prod. xvi. II. 451 = C. arbores.
Macleyana, Parl. l. c. 446 = C. Macleyana.
macrostachya, Knight, ex Gord. Pinet. 86 = C. oblonga.
microcarpa, A. Cunn. ex Benth. Fl. Austral. vi. 237 = C. robusta.
Mooreï, Parl. in DC. Prod. xvi. II. 449 = C. robusta.
Muelleri, Parl. l. c. 450 = C. Muelleri.
Parlatorei, F. Muell. Fragm. v. 186 = C. Parlatorei.
prophinqua, A. Cunn. ex Mirb. in Mém. Mus. Par. xiii. (1825) 74 = C. robusta.
pyramidalis, [A. Cunn.] Sweet, Hort. Brit. ed. II. 473 = C. calcarata.
rhomboidea, Endl. Syn. Conif. 36 = C. rhomboides.
rhomboidea, Hort. ex Parl. in DC. Prod. xvi. II. 449 = C. calcarata.
rigida, Hort. ex Endl. Syn. Conif. 38.—Austral.
robusta, A. Cunn. ex Mirb. in Mém. Mus. Par. xiii. (1825) 74 = C. robusta.
Rori, Endl. Syn. Conif. 36 = C. Roei.
subcordata, Parl. Enum. Sem. Hort. Flor. (1863) 24 = C. Roei.
subumbellata, Parl. in DC. Prod. xvi. II. 447.—N. Caled.
sulcata, Parl. Enum. Sem. Hort. Flor. (1862) 38.—Hab.?
triquetra, Spach, Hist. Vég. Phan. xi. 345 = C. rhomboidea.
tuberculata, Mirb. in Mém. Mus. Par. xiii. (1825) 74 = C. robusta.
verticillis, Carr. Conif. ed. I. 75 = C. rhomboides.
Ventenatii, Mirb. in Mém. Mus. Par. xiii. (1825) 74 = C. rhomboides.
verrucosa, A. Cunn. ex Mirb. l. c. = C. robusta.

FREREA, Dalz. in Journ. Linn. Soc. viii. (1865) 10. t. 2. *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 783. Indica, *Dalz. l. c.*—Ind. or.

FRESNELIA, DC. Prod. v. 236 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 358.
fasciculata, *Baker, in Hook. Ic. Pl.* t. 1108.—Afr. austr.

FRESNIA :—
leptophylla, *DC. Prod.* v. 236.—Afr. austr.
scaposa, *DC. l. c.*—Afr. austr.
stephana, Steud. Nom. ed. II. i. 647, sphalm. = præc.

FRESIERA, Mirb. in Bull. Soc. Philom. (1813) 881 = FREZIERA, Sw. (Ternstr.).

FRESNELIA, Steud. Nom. ed. II. i. 646 (1840) = FRENELA, Mirb. = Callitris, Vent. (Conif.).

FREUCHENIA, Eckl. Verz. 14 (1827) = **Morsea**, Mill. (Irid.).
bulbifera, Eckl. l. c. = M. ramosa.

FREYCINETIA, Gaudich. in Ann. Sc. Nat. Sér. I. iii. (1824) 509. *PANDANEÆ*, Benth. & Hook. f. iii. 951.
JEZABEL, Banks, ex Salisb. Gen. Pl. Fragm. 5 (1866).
VICTORIPERREA, Gaudich. ex Decne. Bot. Voy. Astrol. et Zél. 83 (1845).
angustifolia, *Blume, Rumphia*, i. 159. t. 43.—Samatra.
angustifolia, Kurz, in Journ. Bot. v. (1867) 134 = Insonensis, pycnophylla.
arborea, *Gaudich. in Freyc. Voy. Bot.* 481. t. 41.—Ins. Sandvic.
Arnottii, *Gaudich. Bot. Voy. Bonite*, t. 85, 86.—Ins. Sandvic.
Banksii, *A. Cunn. in Hook. Comp. Bot. Mag.* ii. (1836) 377.—N. Zel.
Baueriana, *Endl. Prod. Fl. Norfolk.* 25.—Ins. Norfolk.
Beccarii, *Solms, in Ann. Yard. Buitenz.* ii. (1882) 100.—N. Guin.
Bennettii, Miq. Pl. Jungh. 167 = javanica.
Brownriana, Wall. Cat. n. 6660 = angustifolia.
celebica, *Solms, in Linnæa*, xlii. (1878–79) 108.—Ins. Celebes.
Cumingiana, *Gaudich. Bot. Voy. Bonite*, t. 88 et t. 60.—Ins. Philipp.
cylindraces, *Solms, in Linnæa*, xlii. (1878–79) 97.—N. Caled.
Dubreyniana, Gandich. Bot. Voy. Bonite, t. 87 = angustifolia.
demissa, *Benn. Pl. Jav. Rar.* i. 92.—Ins. Tahiti.
De Vriesei, *Solms, in Linnæa*, xlii. (1878–79) 96.—Ins. Celebes.
excelsa, *F. Muell. Fragm.* v. 39.—Austral.
Gaudichaudii, *Benn. Pl. Jav. Rar.* 81. t. 9.—Malaya; Austral.
graminea, *Blume, Rumphia*, i. 159.—Java.
graminifolia, *Solms, in Linnæa*, xlii. (1878–79) 90.—N. Caled.
imbricata, *Blume, Rumphia*, i. 157. t. 40.—Ins. Philipp.
inclinans, *Benn. Pl. Jav. Rar.* 82.—N. Zel.
insignis, *Blume, Rumphia*, i. 158. t. 42.—Ins. Philipp. javanica, *Blume, l. c.* 157. t. 41.—Java.
leucacantha, Miq. Ind. Sem. Hort. Amstelod. (1858) 4 = Pandanus pygmæus.
lucionensis, *Perrd. Spim. Bot.* 238.—Ins. Philipp.
marginata, *Blume, Rumphia*, i. 158.—N. Guinea.
Milnei, *Seem. in Bonplandia*, ix. (1861) 260, nomen; *Fl. Vit.* 283. t. 66.—Ins. Fiji.
nitida, Miq. Ind. Sem. Hort. Amstelod. (1858) 4 = Pandanus stenophyllus.
Pritchardii, *Seem. Fl. Vit.* 282.—Ins. Fiji.
pycnophylla, *Solms, in Linnæa*, xlii. (1878–79) 91.—Zeylan.
radicans, *Gaudich. in Freyc. Voy. Bot.* 492. t. 48.—Ins. Sandvic.
scandens, *Gaudich. l. c.* t. 42.—Ins. Molucc.
scandens, *Hook. & Arn. Bot. Beech. Voy.* 97.—Ins. Sandvic.
Scheffleri, *Solms, in Linnæa*, xlii. (1878–79) 98.—Java.
spectabilis, *Solms, l. c.* 88.—N. Caled.
sphærocephala, *Gaudich. Bot. Voy. Bonite*, t. 92.—Hab.?
Storckii, *Seem. in Bonplandia*, ix. (1861) 260, nomen; *Fl. Vit.* 283. t. 86.—Ins. Fiji.
strobilacea, *Blume, Rumphia*, i. 156. t. 39.—Ins. Amboin.
tenuis, *Solms, in Linnæa*, xlii. (1878–79) 87.—Sumatra.
Urvilleana, *Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél.* 83.—Ins. Timor.
Victoriperrea, *Solms, in Linnæan*, xlii. (1878–79) 103.—Ins. Tahiti.

FREYCINETIA :—
vitiensis, *Seem. in Bonplandia*, ix. (1861) 260, nomen; *Fl. Vit.* 282. t. 85.—Ins. Fiji.
Walkeri, *Solms, in Linnæa*, xlii. (1878–79) 92.—Zeylan.
Webbiana, *Gaudich. Bot. Voy. Bonite*, t. 87.—Hab.?

FREYERA, Reichb. Handb. 291 (1837) = **Chaerophyllum**, Linn. (Umbell.).
Biasolettiana, Reichb. l. c. (= *Biasolettia tuberosa*).—Europ.
congesta, Boiss. & Heldr. Diagn. Ser. II. vi. 88 (= *Biasolettia congesta*).—Græc.
cretica, Boiss. & Heldr. l. c. ii. 101 = C. creticum.
cynapioides, Griseb. Spicil. Fl. Rumel. i. 366 (= *Biasolettia tuberosa*).—Ital.; Dalmat.
divaricata, Boiss. & Orph. Diagn. Ser. II. ii. 102 (= *Biasolettia divaricata*).—Græc.
macrocarpa, Boiss. Diagn. Ser. II. ii. 101 (= *Biasolettia macrocarpa*).—Græc.
parnassica, Boiss. & Heldr. l. c. 102 (= *Biasolettia parnassica*).—Græc.
pumila, Boiss. Fl. Orient. iii. 897.—Græc.
stylosa, Boiss. Diagn. Ser. II. ii. 101.—Græc.
tuberosa, Reichb. Handb. 291 (= *Biasolettia tuberosa*) —Europ.

FREYERIA, Scop. Introd. 206 (1777) = **Linociera**, Sw. (Oleac.).

FREYLENIA, Brongn. Enum. Gen. 60 (1843) = seq.

FREYLINIA, Colla, Hort. Ripul. 56 (1823) *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 937.
cestroides, *Colla, l. c.*—Afr. austr.
densiflora, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 55.—Afr. austr.
lanceolata, G. Don, in Sweet, Hort. Brit. ed. III. 528 = cestroides.
longiflora, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 55.—Afr. austr.
oppositifolia, Colla, Hort. Ripul. 57 = cestroides.
rigida, G. Don, Gen. Syst. iv. 617 = undulata.
undulata, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 55.—Afr. austr.

FREZIERA, Sw. ex Willd. Sp. Pl. ii. 1179 (1799); et Fl. Ind. Occ. 971. t. 19 (1800). *TERNSTROEMIACEÆ*, Benth. & Hook. f. i. 188.
ERGTIUM, Sw. [Soland.] Prod. Veg. Ind. Occ. 85 (1788).—Cf. Fl. Ind. Occ. ii. 971.
LETTSOMIA, Ruiz & Pav. Prod. 77. t. 14 (1794).
angulosa, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 332.—Bolivia.
arbutifolia, *Planch. & Triana, l. c.* Sér. IV. xviii. (1862) 262.—N. Granat.
attenuata, Wall. Cat. n. 1451 = Pyrenaria attenuata.
Bonplandiana, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 335.—N. Granat.
calophylla, *Triana & Planch. l. c.* Sér. IV. xviii. (1862) 261.—N. Granat.
candicans, *Tul. l. c.* Sér. III. viii. (1847) 328.—N. Granat.
canescens, *Humb. & Bonpl. Pl. Aequin.* i. 27. t. 7.—Peruv.
cerrina, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 330.—Mexic.
chrysophylla, Humb. & Bonpl. Pl. Aequin. i. 31. t. 9 = sericea.
cordata, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 334.—N. Granat.
dioica, Macfad. Fl. Jamaic. i. 115 = Elæodendron dioicum.
discolor, Pœpp. ex Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 161 = Quiina Poeppigiana.
Dombeyana, *Tul. l. c.* viii. (1847) 329.—N. Granat.
elegans, Tul. l. c. 326 = Cleyera elegans.
floribunda, Pœpp. ex Tul. l. c. xi. (1849) 167 = Quiina floribunda.
hirsuta, Semm. Bot. Voy. Herald, 87 = sericea.
hirsuta, *Sw. in Rees, Cyclop.* xv. n. 4.—Ins. S. Vincent.
illicioides, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 327.—Jamaica.
integrifolia, Benth. Pl. Hartw. 6 = Cleyera integrifolia.
lanata, Tul. in Ann. Sc. Nat. Sér. III. viii. (1847) 334, in obs. = canescens.
longipes, *Tul. l. c.* 327.—N. Granat.
macrophylla, *Tul. l. c.* 330.—Mexic.

FREZIERA :—
nervosa, *Humb. & Bonpl. Pl. Aequin.* i. 81. t. 9.—Peruv.
Nimaninae, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 338.—Cuba.
ochnoides, Wall. Cat. sub n. 1460 = Cleyera ochnacea.
Perrottetiana, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 332.—Ins. Guadeloup.
reticulata, *Humb. & Bonpl. Pl. Aequin.* i. 81. t. 9.—Peruv.
rorairaensis, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 339.—Guiana.
salicifolia, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 124 = hirsuta.
serices, *Humb. & Bonpl. Pl. Aequin.* i. 29. t. 8.—Ecuador.
suberosa, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 330.—N. Granat.
theoides, Sw. Fl. Ind. Occ. ii. 972 = Cleyera theoides.
undulata, *Willd. Sp. Pl.* ii. 1179; *Sw. l. c.* 974.—Ind. occ.
yungasia, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 338.—Bolivia.

FRIDERICIA, Mart. in Nov. Act. Nat. Cur. xiii. (1827) Praef. 7. tt. A, B. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1052.
Gallesiana, *Mart. l. c.* 11. t. 13.—Bras.
speciosa, *Mart. l. c.*—Bras.

FRIEDERICHSTHALIA, A. DC. Prod. x. 173 (1846) = FRIEDRICHSTHALIA, Fenzl = Trichodesma, R. Br. (Borag.).

FRIEDERICIA, Reichb. Consp. 125 (1828) = **Fridericia**, Mart. (Bignon.).

FRIEDLANDIA, Cham. & Schlecht. in Linnaea, ii. (1827) 848 = Diplusodon, Pohl (Lythrar.).
albiflora, Mart. ex DC. Prod. iii. 94(b) = D. virgatus.
amoena, Cham. & Schlecht. in Linnaea, ii. (1827) 850 = D. virgatus.
angustifolia, Mart. ex DC. Prod. iii. 94(b) = D. hexander.
buxifolia, Cham. & Schlecht. in Linnaea, ii. (1827) 851 = D. buxifolius.
epilobioides, Mart. in DC. Prod. iii. 94(b) = D. epilobioides.
glaucescens, Mart. ex Koehne, in Mart. Fl. Bras. xiii. ii. 832 = D. glaucescens.
helianthemifolia, Mart. in DC. Prod. iii. 94(c) = D. helianthemifolius.
hexandra, Mart. ex DC. l. c. 94(a) = D. hexander.
hirsuta, Cham. & Schlecht. in Linnaea, ii. (1827) 852 = D. hirsutus.
hirtula, Cham. & Schlecht. l. c. = D. microphyllus.
leucantha, Mart. ex Koehne, in Mart. Fl. Bras. xiii. ii. 814 = D. virgatus.
lythroides, Mart. ex DC. Prod. iii. 94(c) = D. lythroides.
Myrsinites, Mart. ex DC. l. c. 94(b) = D. Myrsinites.
myrtifolia, Mart. ex Koehne, in Mart. Fl. Bras. xiii. ii. 814 = D. virgatus.
nitida, Mart. ex DC. Prod. iii. 94(a) = D. nitidus.
nummularifolia, Mart. ex Koehne, in Mart. Fl. Bras. xiii. ii. 823 = D. glaucescens.
parvifolia, Mart. ex DC. Prod. iii. 94(c) = D. parvifolius.
rotundifolia, Mart. ex DC. l. c. 94(c) = D. rotundifolius.
serpyllifolia, DC. ex Koehne, in Mart. Fl. Bras. xiii. ii. 834 = D. serpyllifolius.
stachyoides, Cham. & Schlecht. in Linnaea, ii. (1827) 853 = D. villosissimus.
thymifolia, Mart. ex DC. Prod. iii. 94 (b) = D. serpyllifolius.
vaccinifolia, Mart. l. c. = D. buxifolius.
verbenaefolia, Mart. l. c. 94(c) = D. hirsutus.
villosella, Mart. l. c. = D. microphyllus.

FRIEDRICHSTHALIA, Fenzl, in Endl. Nov. Stirp. Dec. 38 (1839) = **Trichodesma**, R. Br. (Borag.).
incana, Bunge, Del. Sem. Hort. Dorp. (1848) 7 = T. hirsutum.
physaloides, Fenzl, in Endl. Nov. Stirp. Dec. 53 = T. physaloides.
trichodesmoides, Bunge, Del. Sem. Hort. Dorp. (1848) 7 = T. calathiforme.

FRIESEA, Reichb. Nom. 209 (1841) = seq.

FRIESIA, DC. Prod. i. 520 (1824) = **Aristotelia**, L'Hérit. (Tiliac.).
chinensis, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 348 = Elaeocarpus chinensis.
fruticosa, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 46 = A. fruticosa.
peduncularis, DC. Prod. i. 520 = A. peduncularis.
racemosa, A. Cunn. in Ann. Nat. Hist. iv. (1840) 24 = A. racemosa.

FRIESIA, Spreng. Anleit. ii. 11. 885 (1818) = **Crotonopsis**, Michx. (Euphorb.).
argentea, Spreng. Syst. iii. 850 = C. linearis.

FRISCA, Spach, Hist. Vég. Phan. x. 460 (1841) = **Theaium**, Linn. (Santal.).

FRITILLARIA, [Tourn.] Linn. Syst. ed. I (1735) *LILIACEAE*, Benth. & Hook. f. iii. 817.
AMBLIRION, Rafin. in Journ. Phys. lxxxix. (1819) 102.
CORONA, Fisch. ex R. Grah. in Edinb. N. Phil. Journ. xxi. (1836) 157.
IMPERIALIS, Adans. Fam. ii. 48 (1763); Juss. Gen. 49 (1789).
KOROLKOWIA, Regel, Gartenfl. (1873) 161. t. 760.
LILIORHIZA, Kellogg, in Proc. Calif. Acad. ii. (1863) 46.
LYPERIA, Salisb. Gen. Pl. Fragm. 56 (1866).
MELORIMA, Rafin. Fl. Tellur. iv. 53 (1836).
MONOCOSIA, Salisb. Gen. Pl. Fragm. 56 (1866)
PETILIUM, Linn. Syst. ed. I (1735); Hort. Cliff. 119 (1737).
RHINOPETALUM, Fisch. ex Alex. in Edinb. N. Phil. Journ. viii. (1830) 19.
SUSIANA, Linn Syst. ed. I (1735).
THERESIA, C. Koch, in Linnaea, xxii. (1849) 232.
TOZZETTIA, Parl. Nuov. Gen. e Sp. 13 (1854).
acmopetala, Baker, in Bot. Mag. t. 6321 = Elwesii.
acmopetala, *Boiss. Diagn.* Ser. I. vi. 104.—Syria.
alba, Kellogg, in Proc. Acad. Calif. i. (1873) 45 = biflora.
alba, *Nutt. Gen. Am.* i. 222.—Am. bor.
altaica, Lam. ex Schult. f. Syst. vii. 397 = verticillata.
aquatica, Mill. Gard Dict. ed. VIII. n. 2 = pyrenaica.
armena, *Boiss. Diagn.* Ser. I. vi. 106.—Armenia.
assyriaca, *Baker, in Journ. Linn Soc.* xiv. (1874) 265. —Mesopotam.
atropurpurea, *Nutt. in Journ. Acad. Philad.* vii. (1834) 54.—Am. bor. occ.
aurea, Schott, in Oesterr. Bot. Wochenbl. iv. (1854) 137 = lutea.
autumnalis, Mill. Gard. Dict. ed. VIII. n. 10 = Eucomis undulata.
barbata, H. B. & K. Nov. Gen. et Sp. i. 288 = Calochortus flavus.
biflora, *Lindl. Bot. Reg.* sub t. 1663.—Calif.
bithynica, *Baker, in Journ. Linn. Soc.* xiv. (1874) 264. —Bithyn.
Boissieri, *Costa, Suplem. Fl. Catal.* 72.—Hispan.
bucharica, *Regel, in Act. Hort. Petrop.* viii. (1884) 652. —Bokhara.
Burnati, Hort. Backh. ex Gard. Chron. (1879) i. 685 = tubaeformis.
camschatcensis, *Ker-Gawl. in Bot. Mag.* sub t. 1216.—Am. or.
camtschatcensis, Sweet, Hort. Brit. ed. I. 427 = praec.
cantoniensis, Lour. Fl. Cochinch. i. 206 = Disporum pullum.
caucasica, Adam, in Web. & Mohr, Beitr. i. 51 = tulipifolia.
caussolensis, Goaty & Pons, ex Ardoino, Fl. Alp. Marit. 87 = tenella.
chamaerrutis, Turcz. ex Ledeb. Fl. Ross. iv. 148 = dagana.
chia, Boiss. ex Nym. Consp. 723 = Pinardi.
chrysea, *D. Don, Prod. Fl. Nep.* 51.—Reg. Himal.
collicola, Hance, in Journ. Bot. viii. (1870) 76 = verticillata.
collina, Adam, in Web. & Mohr, Beitr. i. 50 = lutea.
conica, *Boiss. Diagn.* Ser. I. vi. 105.—Graecia.
cornuta, *Hort. Goronh. ex Schult. f. Syst.* vii. 399, nomen = seq. ?
Cornuti, *Hort. Dorpat. ex Schult. f. l. c.* nomen.—Hab. ?

FRITILLARIA :—
coussolensis, Goaty & Pons, ex Nym. Consp. 722 = tenella.
crassifolia, *Boiss. & Huet, Diagn.* Ser. II. iv. 108.—As. Min.
cupres, *R. Grah. in Edinb. N. Phil. Journ.* (Jan. 1836) 192.—Mexic.?
dagana, *Turcz. in Flora*, xvii. (1834) 1. *Bröbl.* 25.—Sibiria.
dasyphylla, *Baker, in Gard. Chron.* (1875) 653.—Lycia.
delphinensis, Gren. ii. Gren. & Godr. Fl. Fr. iii 180 = tubaeformis.
Edwardi, *Regel, in Act. Hort. Petrop.* viii. (1884) 658. t. 2.—Bokhara.
Ehrharti, *Boiss. & Orph. Diagn.* Ser. II. iv. 106.—Archip. Graec.
Elwesii, *Boiss. Fl. Orient.* v. 181.—Lycia.
esculenta, Adam, in Web. & Mohr, Beitr. i. 52 = lutea.
esculenta, Nutt. ex Baker, in Journ. Linn. Soc. xiv. (1874) 271 = lanceolata.
Fleischeri, Steud. & Hochst. ex Kunth, Enum. Pl. iv. 248 = seq.
Fleischeriana, *Steud. & Hochst. ex Schult. f. Syst.* vii. 398.—As. Min.
Forbesii, *Baker, in Journ. Linn. Soc.* xiv. (1874) 264. —Lycia.
Gardneriana, *Wall. Cat. n.* 5080.—Reg. Himal.
Gawleri, Jaub. & Spach, Illustr. v. 110 = tenella.
gibbosa, *Boiss. Diagn.* Ser. I. vi. 107.—Persia; Afghan.
glaucescens, Willd. in Hort. Berol. (1818).—Cf. Kunth, Enum. Pl. iv. 250 = lutea.
graeca, *Boiss. & Sprun. Diagn.* Ser. I. vi. 104.—Graecia.
graminifolia, Stokes, Bot. Mat. Med. ii. 235 = F. Meleagris.
Grayana, Reichb. f. & Baker, in Journ. Bot. xvi. (1878) 268 = biflora.
Gussciardi, Heldr. & Sart. in Boiss. Diagn. Ser. II. iv. 102 = graeca.
Gutierrei-Waldemaris, Klotzsch, in Bot. Ergeb. Waldem. Reise, 53 = Roylei
Hermontis, Fenzl, ex Klatt, in Hamb. Gartenz. xvi. (1860) 442 = crassifolia.
heterophylla, Soland. ex Baker, in Journ. Linn. Soc. xiv. (1874) 258 = ruthenica.
hispanica, Boiss. & Reut. Diagn. Ser. II. iv. 101 = lusitanica.
Hookeri, Baker, in Journ. Linn. Soc. xiv. (1874) 269 = Lilium Hookeri.
Imperialis, *Linn. Sp. Pl.* 303.—Persia; Reg. Himal.
involucrata, *All. Auct. Fl. Pedem.* 24.—Italia.
involucrata, Reichb. Ic. Fl. Germ. x. 8 = tenella.
italica, Schult. f. Syst. vii. 393, in syn. = latifolia.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 158.—Japon.
kamtschatcensis, Fisch. ex Hook. Fl. Bor. Am. i. 181 = camtschatcensis.
kamtschatcensis, Torr. in Pacif. Rail. Rep. iv. 146 = biflora.
Karelinii, Baker, in Journ. Linn. Soc. xiv. (1874) 268 = gibbosa.
Kotschyana, Herb. in Lindl. Bot. Reg. (1844) Misc. 43 = latifolia.
kurdica, *Boiss. & Noë, Diagn.* Ser. II. iv. 103.—Kurdistan.
lanceolata, *Pursh, Fl. Am. Sept.* i. 230.—Am. bor. occ.
lanceolata, Torr. Bot. Mex. Bound. t. 61 = biflora
latifolia, Frivald ex Nym. Consp. 722 = graeca.
latifolia, *Willd. Sp. Pl.* ii. 93.—Reg. Caucas.; Persia.
leucantha, Fisch. ex Schult. f. Syst. vii. 397 = verticillata.
libanotica, *Baker, in Journ. Linn. Soc.* xiv. (1874) 270. —Syria.
liliacea, *Lindl. Bot. Reg.* sub t. 1663.—Calif.
lusophylla, Doumenjou, ex Nym. Consp. 721 = pyrenaica.
longifolia, Hill, Veg. Syst. xiii. 63 = Eucomis undulata.
longifolia, Stev. ex Ledeb. Fl. Ross. iv. 146 = melagroides.
lusida, Salisb. Prod. 237 = pyrenaica.
lusitanica, *Wikstr. in Vet. Acad. Handl. Stockh.* (1821) 252.—Lusitan.
lutea, *Mill. Gard. Dict.* ed. VIII. n. 4; *Bieb. Fl. Taur. Cauc.* i. 269.—As. Min.; Reg. Caucas.
lutea, Reichb. Ic. Fl. Germ. 10. t. 975 = F. Melagris.
lycia, Boiss. & Heldr. Diagn. Ser. I. xiii. 80 = acmopetala.

FRITILLARIA :—
macrandra, Baker, in Gard. Chron. (1875) i. 715 = Ehrharti.
macrophylla, D. Don, Prod. Fl. Nep. 51 = Lilium roseum.
major, Baker, in Journ. Linn. Soc. xiv. (1874) 254 = F. Melegris.
Melegris, Ambros. Fl. Tyr. Austr. i. 530 = tubaeformis.
Melegris, Bivona, Pl. Sic. ii. 26 = messanensis.
Melegris, Brot. Fl. Lusit. i. 520 = lusitanica.
Melegris, Falk, Beitr. ii. 158 = melegroides.
Melegris, Lapeyr. Hist. Abr. Fl. Pyr. 583 = pyrenaica.
Melegris, *Linn. Sp. Pl.* 304.—Europ. ; Reg. Caucas.
Melegris, Pall. ex Schult. f. Syst. vii. 391 = racemosa.
Melegris, Tenore, Fl. Nap. iii. 369 = tenella.
melegroides, *Patrin, ex Schult. f. Syst.* vii. 395.—Sibir.
messanensis, Bolss. Voy. Bot. Espagne, ii. 610 = lusitanica.
messanensis, *Raffin. in Desv. Journ. Bot.* iv. (1814) 272. —Reg. Mediterr.
minor, Ledeb. Fl. Alt. ii. 34 = melegroides.
minuta, *Boiss. & Noë, Diagn.* Ser. II. iv. 104.—Kurdist.
Moggridgei, Hort. ex Flor. Mag. N. S. t. 405 = tubaeformis.
montana, Hoppe, in *Flora*, xv. (1832) *ii. 476* = tenella.
montana, Sang. in Atti Acad. Pont. Lincei, Ser I. xv. (1862) 29 = F. Melegris.
multiflora, Kellogg, in Proc. Calif. Acad. i. (1855) 57 = parviflora.
multiscapidea, Kellogg, l. c. 46 = Erythronium purpurascens.
Munbyi, *Baker, in Gard. Chron.* (1877) i. 45.—Algeria.
mutica, Lindl. Bot. Reg. sub t. 1663 = lanceolata.
nana, Burm. f. Fl. Cap. Prod. 9 = Eucomis nana.
neglecta, Parl. Fl. Ital. ii. 415 = lusitanica.
nervosa, Unio Itin. (1829) ; ex Baker, in Journ. Linn. Soc. xiv. (1874) 261 = lusitanica.
nervosa, Willd. Enum. Hort. Berol. 364 = pyrenaica.
nigra, Mill. Gard. Dict. ed. VIII. n. 3 = pyrenaica.
nobilis, Salisb. Prod. 236 = persica.
obliqua, *Ker-Gawl. in Bot. Mag.* t. 857.—Graecia.
Olivieri, *Baker, in Journ. Linn. Soc.* xiv. (1874) 261. —Persia.
olympica, C. Koch, in Linnaea, xxii. (1849) 232 = pontica.
oranensis, *Pomel, Nouv. Mat. Fl. Atl.* 253 ; *Poir. in Rev. Hortic.* (1881) 80.—Algeria.
orientalis, Adam, in Web. & Mohr, Beitr. i. 50 = tenella.
orientalis, Willd. ex Ledeb. Fl. Ross. iv. 145 = lutea.
Orsiniana, Parl. Fl. Ital. ii. 411 = tenella.
oxypetala, *Royle, Illustr. Bot. Himal.* i. 388.—Reg. Himal.
pallida, Salisb. Prod. 237 = F. Melegris ?
pallidiflora, *Schrenk, Enum. Pl. Nov.* 5.—Sibir.
parviflora, Mart. Hort. Monac. (1838).—*Cf. Linnaea*, xiii. (1839) Litt. 102 = tulipifolia ?
parviflora, *Torr. in Pacif. Rail. Rep.* iv. 148.—Calif.
persica, S. G. Gmel. It. i. 66 = ruthenica.
persica, *Linn. Sp. Pl.* 304.—Armenia.
persica, Pall. ex Kunth, Enum. Pl. iv. 250 = latifolia.
Pinardi, *Boiss. Diagn.* Ser. I. vii. 106.—As. Min.
plantaginea, Pers. Syn. i. 359 = scq.
plantaginifolia, *Lam. Encyc.* ii. 550.—Oriens.
pluriflora, *Torr. ex Benth. Pl. Hartw.* 338.—Calif.
pontica, *Wahlenb. in Bergg. Reser, ii. Bihang*, 27.—Bysant. ; As. Min.
praecox, Hort. ex C. Koch, Ind. Sem. Hort. Berol. (1856) App. 10 = F. Melegris.
praecox, *Sweet, Hort. Brit.* ed. I. 426.—Europ. austr.
Przewalskii, *Maxim. in Trautv. Regel, Maxim. et Winkl. Dec. Pl. Nov.* 9.—As. or.
pterocarpa, Stocks, in Hook. Kew Journ. iv. (1852) 180 = gibbosa.
pudica, *Spreng. Syst.* ii. 64.—Am. bor. occ.
punctata, J. F. Gmel. Syst. 545 = Eucomis punctata.
purpurea, H. B. & K. Nov. Gen. et Sp. i. 288 = Calochortus Bonplandianus.
pyrenaea, Gren. in Acad. Sc. Besanç. Séances Publ. (1886) 147 = pyrenaica.
pyrenaica, Georgi, Beschr. Russ. Reich. III. iv. 900 = melegroides.
pyrenaica, Gmel. Cat. (1821) 26 = messanensis.
pyrenaica, Host, Fl. Austr. i. 496 = tenella.

FRITILLARIA :—
pyrenaica, *Linn. Sp. Pl.* 304.—Mont. Pyren.
pyrenaica, Pall. Reise, i. 147 = ruthenica.
pyrenaica, Sibth. & Sm. Fl. Graec. t. 338 = graeca.
pyrenaica, Urv. in Mém. Soc. Linn. Par. i. (1822) 294 = pontica.
pyrenica, Webb, ex Nym. Consp. 732 = lusitanica.
racemosa, Ker-Gawl. in Bot. Mag. t. 952 = tenella.
racemosa, Kunth, Enum. Pl. iv. 249, ex parte = pyrenaica.
racemosa, Mill. Gard. Dict. ed. VIII. n. 7 = tulipifolia.
racemosa, *Sm. in Rees, Cycl.* xv. n. 6 ; *Schult. f. Syst.* vii. 391.—Reg. Caucas.
recurva, *Benth. Pl. Hartw.* 340.—Am. bor. occ.
regia, Linn. Sp. Pl. 303 = Eucomis nana.
Reuteri, *Boiss. Diagn.* Ser. I. v. 57.—Persia.
Roxgayi, Boiss. & Blanche, ex Boiss. Fl. Orient. v. 180 = acmopetala.
rhodocanakis, *Orph. in Atti Congr. Bot. Fir.* 1874 (1877) 314, nomen ; *Baker, in Journ. Bot.* xvi. (1878) 232.—Archip. Graec.
Roylei, *Hook. Ic. Pl.* t. 860.—Reg. Himal.
ruthenica, *Wikstr. in Vet. Akad. Handl. Stockh.* (1821) 253.—Reg. Caucas.
scandens, Fisch. ex Schult. f. Syst. vii. 397 = verticillata.
Schliemanni, *Sintenis, in Illustr. Hortic.* xxxi. (1884) 106.—As. Min.
serotina, Schult. f. Syst. vii. *395*, in syn. = pyrenaica.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) i. 448.—As. centr.
Sibthorpiana, *Baker, in Journ. Linn. Soc.* xiv. (1874) 275.—Graecia.
stenophylla, Boiss. & Reut. Diagn. Ser. II. iv. 100 = lusitanica.
tardiflora, Lehm. ex Schult. f. Syst. vii. 399 = pyrenaica ?
tenella, *Bieb. Fl. Taur. Cauc.* i. 269.—Europ. ; Reg. Caucas.
tenella, *Reichb. Fl. Germ. Excurs.* 102.—Europ.
tenella, Wikstr. in Vet. Akad. Handl. Stockh. (1821) 256 = racemosa.
tenuifolia, Salisb. Prod. 236 = F. Melegris.
thessalica, Sprun. ex Nym. Consp. 732 = graeca.
Thomsoniana, D. Don, in Royle, Illustr. Bot. Himal. i. 388 = Lilium Thomsonianum.
Thunbergii, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 157 = verticillata.
tristis, Heldr. & Sart. in Boiss. Diagn. Ser. II. iv. 104 = oblqua.
tubaeformis, *Gren. & Godr. in Mém. Soc. Emul. Doubs* (1854) ; *ex eor. Fl. Fr.* iii. 180.—Europ. austr.
tulipifolia, Sprun. ex Nym. Consp. 732 = graeca.
tulipifolia, *Bieb. Fl. Taur. Cauc.* i. 270.—Reg. Caucas.
tulipifolia, Bory & Chaub. Fl. Pélop. 98 = conica.
umbellata, Mill. Gard. Dict. ed VIII. n. 5 = pyrenaica.
uniflora, *Maxim. in Trautv. Regel, Maxim. et Winkl. Dec. Pl. Nov.* 9.—Reg. Amur.
verticillata, Bieb. Fl. Taur. Cauc. i. 268 = ruthenica.
verticillata, Wall. Cat. n. 5079 A = cirrhosa.
verticillata, *Willd. Sp. Pl. ii.* 91.—Sibir.
viridis, Kellogg, in Proc. Calif. Acad. ii. (1868) 9 = lanceolata.
Waldemari, Pritz. Ic. Ind. ii. 136 = Gulielmia Waldemarii.
Wanjewi, *Regel, Gartenfl.* (1879) 253.—Turkestan.
Wolwowi, Regel, l. c. t. 993 = pracc.

FRITTILLARIA, Scop. Introd. 83 (1777) = pracc.

FRITSCHIA, Walp. Rep. ii. 126 (1843) = scq.

FRITZSCHIA, Cham. in Linnaea, ix. (1834) 897. *MELASTOMACEAE*, Benth. & Hook. f. i. 743.
anisostemon, *Cham. l. c.* 899.—Bras.
cuneata, *Cham. l. c.*—Bras.
integrifolia, *Cham. l. c.* 898.—Bras.
punctulata, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 279 = erecta.

FRIVALDIA, Endl. Gen. 884 (1838) = **Microglossa**, DC. (Compos.).
odoratula, *Zoll. & Mor. in Nat. en Genees. Arch. Nederl. Indie*, ii. (1845) 281 = M. volubilis.

FRIVALDZKIA, Reichb. Nom. 86 (1841) = pracc.

FROEBELIA, Regel, Gartenfl. i. (1852) 164. t. 16 = **Amoreuxia**, R. Br. (Spaerid.).
fasciculiflora, Regel, l. c. = A. fasciculiflora.

FROEHLICHIA, D. Dietr. Syn. Pl. i. 419, 454 (1839) = FROELICHIA, Vahl = **Coussarea**, Aubl. (Rosac.).

FROEHLICHIA, Endl. Each. 185 (1841) = **Froelichia**, Moench (Amarant.).

FROEHLICHIA, Pfeiff. Nom. i. 1278 (1874) = FROELICHIA, Wulf. = **Kobresia**, Willd. (Cyperac.).

FROELICHIA, Moench, Meth. 50 (1794). *AMARANTACEAE*, Benth. & Hook. f. iii. 41.
HOPLOTHECA, Spreng. Syst. iv. Cur. Post. 59 (1826)
LOPHOCARPUS, Link, Diss. Bot. Sïerin. (1795) 52
NINANGA, Raffin. Fl. Tellur. iii. 76 (1836), partim.
OPLOTHECA, Nutt. Gen. Am. ii. 78 (1818).
Drummondii, *Moq. in DC. Prod.* xiii. ii. 431.—Am bor.
floridana, *Moq. l. c.* 420.—Am. bor. ; Ind. occ.
gracilis, *Moq. l. c.*—Am. bor.
Humboldtiana, *Seub. in Mart. Fl. Bras.* v. i. 166.—Bras.
interrupta, Moq. in DC. Prod. xiii. ii. 421 = lanata.
lactea, Heynh. Nom. ii. 248 = Aerva sanguinolenta.
lanata, *Mornch, Meth.* 50.—Mexic.
lanata, Moq. in DC. Prod. xiii. ii. 422 = Humboldtiana.
Montsanna, Klotzsch, ex Seub. in Mart. Fl. Bras. v. i. 166 = tomentosa.
nudicaulis, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 193.—Ins. Galapag.
sericea, *Moq. in DC. Prod.* xiii. ii. 422.—Bras.
tomentosa, *Moq. in DC. l. c.* 421.—Bras.

FROELICHIA, Vahl, Eclog. Am. Praef. 3 (1796) = **Coussarea**, Aubl. (Rubiac.).
paniculata, Vahl, l. c. = C. Froelichia.
uniflora, Spreng. Syst. ii. 406 = Myonima multiflora.
violacea, Spreng. l. c. = Loranthus tetrandrus.

FROELICHIA, Wulf. ex Roem. & Schult. Syst. ii. 156, in syn. (1817) ; Fl. Nor. 729 (1858) = **Kobresia**, Willd. (Cyperac.).
caricoides, Wulf. ex Roem. & Schult. l. c. = K. scirpina.

FROLOVIA, Ledeb. ex DC. Prod. vi. 588 (1837) = **Saussurea**, DC. (Compos.).
lyrata, Ledeb. ex DC. l. c. = S. Frolovii.

FROPIERA, Boston, ex Hook. f. in Journ. Linn Soc v. (1861) 2. t. 1. *MYRTACEAE*, Benth. & Hook. i. 725.
PHILOXYLON, Thou. ex Tul. in Ann. Sc. Nat. Ser IV. vi. (1856) 138.
mauritiana, *Boston, ex Hook. f. in Journ. Linn. Soc.* v. (1861) 2.—Ins. Maurit.

FRORIEPIA, C. Koch, in Linnaea, xvi. (1842) 362 = **Carum**, Rupp. (Umbellif.).
nuda, C. Koch, l. c. = C. segetum.
subpinnata, Baill. Hist. Pl. vii. 220 = C. segetum.

FROSCULA, Raffin. Fl. Tellur. iv. 44 (1836) = **Dendrobium**, Sw. (Orchid.).
hispida, Raffin. l. c. = D. longicornu.

FROSTIA, Bert. ex Guill. in Ann. Sc. Nat. Sér. II. ii. (1834) 19 = **Apodanthes**, Poit. (Cytin.).
Berteri, Karst. in Nov. Act. Nat. Cur. xxvi. (1858) 923 = A. Berteri.
Blanchetii, Karst. l. c. = A. Blanchetii.
Callandrae, Karst. l. c. 921 = A. Callandra.
Saraticae, Bert. ex Steud. Nom. ed. II. i. 649 = A. Berteri.

FRUCTESCA, DC. in Mélang. Gen. 259 ; Comm. 368 (1839) = **Geartnera**, Lam. (Loganiac.).

FRUTESCA, DC. ex A. DC. in Prod. ix. 33 (1845) = pracc.

FUCHSIA, [Plum.] Linn. Syst. ed. I (1735) ; Gen. ed. I. 27 (1737). *ONAGRARIEAE*, Benth. & Hook. f. i. 796.
BARRESONIA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 178 ; Hist. Vég. Phan. iv. 401 (1835).
DORVALLA, Comm. ex Linn. Encyc. ii. 545 (1786).
ELLOBIUM, Lilja, in Linnaea, xv. (1841) 268.
ENCLIANDRA, Zucc. in Abh. Akad. Munch. ii. (1837) 385.
KIRSCHLEGERIA, Reichb. Handb. 836 (1837).

FUCHSIA:—
Lyciopsis, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 176.
Myrinia, Lilja, Fl. Sveriges, Suppl. i. 25 (1840).
Nahuatia, Schneev. Ic. 21 (1796).
Quelusia, Vand. Fl. Lusit. 28. f. 10 (1788); Roem. Script. Hist. Lusit. 101 (1796).
Schufia, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 177.
Skinnera, Forst. Char. Gen. 57. t. 29 (1776).
Spachia, Lilja, Tractgardtidning (1840) 62; ex Linnæa, xv. (1841) 262.
Thilkuhl, Molina, Sagg. Chile, ed. II. 146 (1810).
Tilco, Adans. Fam. ii. 498 (1763).
acynifolia, Scheidw. in Otto & Dietr. Allg. Gartenz. xv. (1847) 226 = parviflora.
affinis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 274 = integrifolia.
alpestris, *Gardn. in Bot. Mag.* t. 3999.—Bras.
alternans, [Moç. & Sesse,] ex DC. Prod. iii. 37 = thymifolia.
amœna, Hort. ex DC. Prod. i. c. = arborescens.
ampliata, *Benth. Pl. Hartw.* 178.—N. Granat.
apetala, *Ruiz & Pav. Fl. Per.* iii. 89. t. 292.—Peruv.
arborescens, *Sims, Bot. Mag.* t. 2620.—Mexic.
asyaxcensis, *H. B. & K. Nov. Gen. et Sp.* vi. 107.—Peruv.
bacillaris, *Lindl. Bot. Reg.* t. 1480.—Chili.
boliviana, *Carr. in Rev. Hortic.* (1876) 150.—Bolivia.
canescens, *Benth. Pl. Hartw.* 179.—N. Granat.
caracasana, *Field. & Gardn. Serr. Pl.* t. 29.—Venezuela.
chonotica, *Phil. in Linnæa*, xxviii. (1856) 687.—Chili.
coccinea, Curt. Bot. Mag. t. 97 = macrostemma.
coccinea, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii 8.—Bras.!
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 728.—N. Zel.
confertifolia, *Field. & Gardn. Serr. Pl.* t. 28.—Peruv.
conica, Lindl. Bot. Rep. t. 1062 = macrostemma.
corallina ×, Hort. ex Lynch, in Gard. Chron. (1883) ii. 604.
cordifolia, *Benth. Pl. Hartw.* 74.—Mexic.
corollata, Benth. l. c. 179 = petiolaris.
corymbiflora, *Ruiz & Pav. Fl. Per.* iii. 87. t. 325.—Peruv.
corymbosa, Prits. Ic. Ind. i. 469, sphalm. = præc.
curviflora, *Benth. Pl. Hartw.* 177.—Peruv.
cylindracea, Lindl. Bot. Reg. (1838) Misc. 54 = parviflora.
decussata, R. Grah. in Edinb. Phil. Journ. xi. (1824) 306, 401 = macrostemma.
decussata, *Ruiz & Pav. Fl. Per.* iii. 88. t. 328. f. b.—Peruv.
denticulata, *Ruiz & Pav. Fl. Per.* iii. 87. t. 325.—Peruv.
dependens, *Hook. Ic. Pl.* t. 65.—N. Granat.
discolor, Lindl. Bot. Reg. t. 1805 = macrostemma.
Dominicana ×, Hort. ex Fl. des Serres, Sér. I. x. (1854–55) 95. t. 1004.
elegans, 'Paxt,' ex Steud. Nom. ed. II. i. 640 = globosa.
elegans, Salisb. Ic. Stirp. Ic. Rar. 18 = coccinea.
Euclandra, *Steud. Nom.* ed. II. i. 649.—Mexic.
exoorticata, *Linn. f. Suppl.* 217.—N. Zel.
exoniensis ×, Paxt. Mag. Bot. x. (1843) 153 = *corallina*.
fulgens, *Moc. & Sesse, ex DC. Prod.* iii. 89.—Mexic.
globosa, *Lindl. Bot. Reg.* t. 1656.—Mexic.
gracilis, Lindl. l. c. t. 847 = macrostemma.
gracilis, [Moç. & Sesse,] ex DC. Prod. iii. 37 = microphylla.
Hartwegii, *Benth. Pl. Hartw.* 179.—N. Granat.
hirsuta, *Hemsl. in Journ. Bot.* xiv. (1876) 69.—Peruv.
hirtella, *H. B. & K. Nov. Gen. et Sp.* vi. 107.—N. Granat.
insignis, *Hemsl. in Journ. Bot.* xiv. (1876) 69.—Ecuador.
integrifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 278.—Bras.
intermedia, *Hemsl. Diag. Pl. Nov.* 14.—Mexic.
Kirkii, *Hook. f. ex T. Kirk, in Trans. N. Z. Inst.* i. (1868) 92.—N. Zel.
Lemoinei, *Wercz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 298.—Boliv.
longiflora, *Benth. Pl. Hartw.* 177.—Peruv.
loxensis, *H. B. & K. Nov. Gen. et Sp.* vi. 106.—Ecuador.

FUCHSIA:—
lycioides, Andr. Bot. Rep. t. 120 = rosea.
macrantha, *Hook. Bot. Mag.* t. 4383.—N. Granat.
macropetala, *Presl, Rel. Haenk.* ii. 28.—Peruv.
macrostemma, *Ruiz & Pav. Fl. Per.* iii. 88. t. 324. f. 6.—Chili; Fuegia.
macrostigma, *Benth. Pl. Hartw.* 129.—N. Granat.
magellanica, Lam. Encyc. ii. 564 = macrostemma.
membranacea, *Hemsl. in Journ. Bot.* xiv. (1876) 70.—Venezuela.
microphylla, *H. B. & K. Nov. Gen. et Sp.* vi. 103.—Mexic.
minutiflora, *Planch. & Linden, in Fl. des Serres*, Sér. I. viii. (1852–53) 7.—N. Granat.
minutiflora, *Hemsl. Diag. Pl. Nov* 14.—Mexic.
minutiflora, *Hemsl. l. c.* 15.—Mexic.
mixta, *Hemsl. l. c.*—Mexic.
montana, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 275.—Bras.
multiflora, Murr. Syst. ed. XIII. 299 = macrostemma.
nigricana, *Linden, in Fl. des Serres*, Sér. I. v. (1849) t. 481.—Venezuela.
Notarisii, *Lehm. in Hamb. Gartenz.* viii. (1852) 459.—Hab.!
ovalis, *Ruiz & Pav. Fl. Per.* iii 87. t. 323. f. a.—Peruv.
ovata, [Moç. & Sesse,] ex DC. Prod. iii. 37 = parviflora.
paniculata, Lindl. in Gard. Chron. (1856) 301 = arborescens.
parviflora, *Lindl. Bot. Reg.* t. 1048.—Mexic.
parviflora, Zucc. in Abh. Akad. Muench. ii. (1831–36) 337 = F. Encliandra.
pendula, Salisb. Prod 279 = macrostemma vel coccinea.
pendularifolia ×, Hort.—Cf. Gard. Chron. (1877) 1. 521
petiolaris, *H. B. & K. Nov. Gen. et Sp.* vi. 104.—Bras
pilosa, Field. & Gardn. Serr. Pl. t. 27 = ovalis.
procumbens, *R. Cunn. in Ann. Nat. Hist.* iii. (1839) 31.—N. Zel.
prostrata, Baill. in Bull. Soc. Linn. Par. i. (1880) 270, nomen = præc.!
pubescens, *Cambess. in St. Hil Fl. Bras. Mer.* ii. 275.—Bras.
pyrifolia, Presl, Symb. Bot. ii. 19 = integrifolia.
quinduensis, *H. B. & K. Nov. Gen. et Sp.* vi. 105.—Peruv.
racemosa, *Lam. Encyc.* ii. 565.—Ins. S Doming.
radicans, Miers, ex Lindl. Bot. Reg. (1841) Misc 78 = integrifolia.
recurvata, Niven, ex Hook. Bot. Mag. sub t. 3521 = macrostemma.
Riccartoni, Hort. ex Tillery, in Flor. et Pom. (1871) 217 = macrostemma!
robusta ×, Tengbergen, in Tuinb. Fl. ii. (1855) 345, cum tab.
rosea, *Ruiz & Pav. Fl. Per.* iii. 88.—Chili.
salicifolia, *Hemsl. in Journ. Bot.* xiv. (1876) 70.—N. Granat.
scabriuscula, *Benth. Pl. Hartw.* 177.—N. Granat.
serratifolia, *Ruiz & Pav. Fl. Per.* iii. 86. t. 323.—Peruv.
sessilifolia, *Benth. Pl. Hartw.* 176.—N. Granat.
simplicicaulis, *Ruiz & Pav. Fl. Per.* iii. 89. t. 322.—Peruv.
spectabilis, *Hook. Bot. Mag.* t. 4375.—Ecuador.
spinosa, *Presl, Rel. Haenk.* ii. 26. t. 51.—Chili.
splendens, Benth. Pl. Hartw. 61 = intermedia.
splendens, *Zucc. in Flora*, xv. (1832) ii. *Beibl.* 102.—Mexic.
Standishii ×, Paxt. Mag. Bot. xi. (1844) 33.
sylvatica, *Benth. Pl. Hartw.* 176.—N. Granat.
syringæflora, Carr. in Rev. Hortic. (1873) 311 = arborescens.
tenella, Hort. ex G. Don, in Loud. Hort. Brit. 155 = macrostemma.
tetradactyla, Lindl. in Journ. Hort. Soc. I. (1846) 304 = F. Encliandra.
thymifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 104.—Mexic.
triphylla, *Linn. Sp. Pl.* 1191; *H. B. & K. Nov. Gen. et Sp.* vi. 104.—Mexic.
umbrosa, *Benth. Pl. Hartw.* 176.—N. Granat.
vemusta, *H. B. & K. Nov. Gen. et Sp.* vi. 105.—N. Granat.
verrucosa, *Hartw. ex Benth. Pl. Hartw.* 178.—Peruv.
virgata, *Sweet, Hort. Brit.* ed. I. 465, nomen.—Mexic.

FUCHSIA, Sw. Prod. Veg. Ind. Occ. 69 (1788) = **Schradera**, Vahl (Rubiac.).
involucrata, Sw. l. c. = S. Cephalotes.

FUERNBOHRIA, C. Koch, in Linnæa, xvi. (1842) 356. Gen. dub. *UMBELL.*—Cf. Benth. & Hook. f. i. 877.
scitifolia, *C. Koch, l. c.*—Oriens.

FUGOSIA, Juss. Gen. 374 (1789). *MALVACEÆ*, Benth. & Hook. f. i. 208.
Alogyne, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 19.
Bombycospermum, Presl, Rel. Haenk. ii. 127. t. 71 (1831).
Cienfugosia, Cav. Dis. 174. t. 72. f. 2 (1786).
Elidurandia, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 450.
Redoutea, Vent. Hort. Cels, t. 11 (1800).
affinis, A. Juss. in St. Hil. Fl. Bras. Mer. i. 253 = phlomidifolia.
australis, *Benth. Fl. Austral.* i. 220.—Austral.
campestris, *Benth. in Hook. Journ. Bot.* iv. (1842) 166.—Guian.
cuneata, *Benth. Bot. Voy. Sulph.* 68.—Ecuador.
cuneiformis, *Benth. Fl. Austral.* i. 219.—Austral.
digitata, *Pers. Syn.* ii. 240.—Afr. trop.
Drummondii, *A. Gray, Pl. Wright.* i. 23.—Texas.
flavifiora, *F. Muell. Fragm.* v. 44.—Austral.
Gerrardii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 588.—Afr. austr.
hakeæfolia, *Hook. Bot. Mag.* t. 4261.—Austral.
heterophylla, *Spach, Hist. Vég. Phan.* iii. 397.—Am. austr.
latifolia, *Benth. Fl. Austral.* i. 221.—Austral.
lilacinus, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1426 = hakeæfolia.
phlomidifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* i. 253.—Bras.
populifolia, *Benth. Fl. Austral.* i. 221.—Austral.
punctata, *Benth. l. c.* 220.—Austral.
punctata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi (1858) 1. 196.—Bras.
retusa, *Turcz. l. c.* 197.—Venezuela.
Robinsoni, F. Muell. Fragm. ix. 126 = Gossypium Robinsoni.
sulfures, *St. Hil. Fl. Bras. Mer.* i. 252.—Bras.
thespesioides, *Benth. Fl. Austral.* i. 220.—Austral.
tripartita, *Steud. Nom.* ed. II. i. 649.—Am. austr.
triphylla, *Harv. in Harv. & Sond. Fl. Cap.* ii. 588.—Afr. austr.

FULCALDEA, Poir. Encyc. Suppl. v. 875 (1827) = **Barnadesia**, Mutis (Compos.).
laurifolia, Poir. l. c.—Peruv.
tomentosa, Lam. in Linnæa, v. (1830) 348 = Vernonia Monosis.

FUIRENA, Rottb. Descr. et Ic. Pl. 70. t. 19 (1778). *CYPERACEÆ*, Benth. & Hook. f. iii. 1058.
Vaginaria, Pers. Syn. i. 70 (1805).
arenosa, R. Br. Prod. 226 = glomerata.
aristulata, Steud. Syn. Pl. Cyp. 127 = simplex.
bahiensis, Nees, in Linnæa, ix. (1834) 288 = robusta.
canescens, Vahl, Enum. ii. 285 = umbellata.
capitata, Willd. ex Kunth, Enum. Pl. ii. 186 = umbellata.
Cephalotes, Schrad. in Goett. Anz. (1821) iii. 2071 = hirta.
chlorocarpa, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 159.—Afr. trop.
ciliaris, Nees, in Wight, Contrib. 93 = uncinata.
ciliaris, Roxb. Hort. Beng. 81; Fl. Ind. i. 180 = glomerata.
ciliata, Lepr. ex Steud. Syn. Pl. Cyp. 126 = glomerata.
cinerascens, *Boj. ex Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 161.—Afr. trop.; Madag.
coerulescens, *Steud. in Flora*, xii. (1829) 153.—Afr. austr.
complanata, Wright, in Sauv. Fl. Cub. 177 = umbellata.
cuspidata, Kunth, Enum. Pl. ii. 187 = Wallichiana.
Ecklonii, *Nees, in Linnæa*, x. (1836) 143.—Afr. austr.
eriolema, Nees, l. c. ix. (1834) 288 = hirta.
filifolia, Reichb. ex Kunth, Enum. Pl. ii. 209 (= *Elytrantha cuspidata*).—Afr. austr.
glabra, Ecklon ex Steud. Nom. ed. II. i. 650 = hirta.

FUIRENA:—
glabra, Hochst. in Flora, xxviii. (1845) 757 = microlepis.
glabra, *Kunth, Enum. Pl.* ii. 183.—Afr. austr.
glabrata, Steud. Syn. Pl. Cyp. 127 = robusta.
glomerata, Boj. Hort. Maurit. 283 = umbellata.
glomerata, *Lam. Illustr.* i. 160.—Geront. trop.
gracilis, *Kunth, Enum. Pl.* ii. 181.—Bras.
gracilis, Spreng. Nov. Prov. 46 = Rynchospora tenuis.
guianensis, Klotzsch, in Schomb. Fauna et Fl. Brit. Gui. 893, 1061, 1111 = umbellata.
haemaglottis, Steud. Syn. Pl. Cyp. 126 = umbellata.
hexachaeta, Schlecht. in Linnaea, xix. (1847) 69 = incompleta.
Hildebrandtii, Boeck. in Flora, lxv. (1882) 15 = glomerata.
hirta, *Vahl, Enum.* ii. 387.—Afr. austr.
hispida, Ell. Sketch, i. 79 = simplex.
incompleta, *Nees, in Mart. Fl. Bras.* ii. 1. 107.—Bras.
intermedia, Kunth, Enum. Pl. ii. 181 = hirta.
latifolia, Steud. Syn. Pl. Cyp. 126 = robusta.
leptostachya, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 168.—Afr. trop.
macrostachya, Boeck. in Engl. Bot. Jahrb. v. (1884) 507 = pachyrrhiza.
mauritiana, Nees, in Linnaea, ix. (1834) 288 = umbellata.
microlepis, *Kunth, Enum. Pl.* ii. 182, partim = gracilis.
microstachya, *Boeck. in Linnaea*, xxxviii. (1874) 398. —Guian.
mollicula, Kunth, Enum. Pl. ii. 182 = coerulescens, Ecklonii.
nana, *A. Rich. Tent. Fl. Abyss.* ii. 407.—Abyssin.
obtusiflora, Vahl, Eclog. Am. ii. 8 = simplex.
ochreata, Nees, in Linnaea, ix. (1834) 288 = glomerata.
pachyrrhiza, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 161.—Afr. trop.
paniculata, Linn. f. Suppl. 105 = umbellata.
pentagona, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 42.—Afr. trop.
pentagona, Wight & Arn. Contrib. 93 = umbellata.
pubescens, *Kunth, Enum. Pl.* ii. 182.—Afr. ; Corsica.
pumila, Spreng. Syst. i. 237 = squarrosa.
pygmaea, *Wiln. ex Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 160.—Afr. trop.
quinquangularis, Hassk. in Flora, xxv. (1842) 11. Belbl. 3 = umbellata.
rivularis, Nees, in Linnaea, ix. (1834) 288 = umbellata.
robusta, *Kunth, Enum. Pl.* ii. 185.—Bras.
Rottboellii, Nees, in Wight, Contrib. 94 = glomerata.
rubiginosa, Spreng. Fl. Hal. Mant. 39 = Cladium glomeratum.
Schiedeana, Kunth, Enum. Pl. ii. 183 = simplex.
schinophylla, Wright, in Sauv. Fl. Cub. 176 = robusta.
Schweinfurthiana, Boeck. in Flora, lxiii. (1880) 488 = leptostachya.
scirpoides, *Micks. Fl. Sor. Am.* i. 35.—Am. bor.
simplex, *Vahl, Eclog. Am.* ii. 8.—Am. austr.
squarrosa, *Micks. Fl. Sor. Am.* i. 37.—Am. bor.
striata, Llanos, Fragm. 21 = glomerata.
stricta, *Steud. Syn. Pl. Cyp.* 126.—Afr. trop.
tereticulmis, Presl. Rel. Haenk. i. 186 = umbellata.
Thouarsiana, Kunth, Enum. Pl. ii. 186 = umbellata.
Torreyana, Beck, Bot. N. & M. St. 420 = squarrosa.
umbellata, *Rottb. Descr. et Ic. Pl.* 70. t. 19. f. 3.—Amphig. trop. et subtrop.
uncinata, *Kunth, Enum. Pl.* ii. 184.—Ind. or.
uncinata, Wall. Cat. n. 5584 : et ex Thw. Enum. Pl. Zeyl. 347 = umbellata.
verticillata, Spreng. Nov. Prov. 47 = Lagenocarpus rigidus.
Wallichiana, *Kunth, Enum. Pl.* ii. 183.—Ind. or.
Weigelii, Spreng. f. Tent. Suppl. Syst. Veg. 4 = umbellata.
Welwitschii, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 161.—Afr. trop.

FULCHIRONIA, Lesch. in Desf. Cat. Hort. Par. ed. III. 39 (1829) = **Phoenix**, Linn. (Palm.).
senegalensis, Lesch. l. c. = P. spinosa.

FULLARTONIA, DC. Prod. v. 281 (1836) = **Doronicum**, Linn. (Compos.).
kumaonensis, DC. l. c. = D. Royleï.

FUMANA, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 359 = **Helianthemum**, Tourn. (Cistin.).
aciphylla, Boiss. Fl. Orient. i. 449.—As. Min.

FUMANA :—
arabica, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 359 = H. arabicum.
Arbuscula, Ball, in Journ. Bot. xi. (1873) 300 = H. calycinum.
calycina, Clauson, ex Ball, in Journ. Linn. Soc. xvi. (1877) 849 = H. calycinum.
canariensis, Rafin. Sylva Tellur. 132 = H. canariense.
cricifolia, Wallr. Beitr. Fl. Hercyn. 214 = H. Fumana ?
Jorretana, Rafin. Sylva Tellur. 132.—Hab. ?
Fontanesii, Clauson, ex Pomel, Mat. Fl. Atl. 10 = H. Fontanesii.
glutinosa, Boiss. Fl. Orient. i. 449 = H. glutinosum.
grandiflora, Jaub. & Spach, Illustr. iii. 73 = H. vulgare ?
hispidula, Losc. & Pard. Ser. Inc. Pl. Arag. ed. I. 12 = H. hirtum ?
laevipes, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 359 = H. laevipes.
minor, Mill. Icon. Comp. 76 = H. procumbens.
minor, Rafin. Sylva Tellur. 132 = H. Fumana.
montana, Pomel, Mat. Fl. Atl. 10 = H. montanum.
oligosperma, Boiss. & Kotschy, in Boiss. Fl. Orient. i. 448.—Syria.
pinifolia, Wallr. Beitr. Fl. Hercyn. 213.—Europ.
polifolia, Rafin. Sylva Tellur. 132 = H. polifolium.
procumbens, Gren. & Godr. Fl. Fr. i. 173 = H. procumbens.
scoparia, Pomel, Mat. Fl. Atl. 10.—Alger.
Spachii, Gren. & Godr. Fl. Fr. i. 174 = H. Fumana.
viscida, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 359 = H. glutinosum.
vulgaris, Spach, l. c. = H. Fumana.

FUMANOPSIS, Pomel, Mat. Fl. Atl. 9 (1860) = **Helianthemum**, Tourn. (Cistin.).
glutinosum, Pomel, l. c. = H. glutinosum.
laevipes, Pomel, l. c. = H. laevipes.
thymifolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 340 = H. glutinosum.

FUMARIA, Tourn. ex Linn. Syst. ed. I (1735). *PAPAVERACEAE*, Beuth. & Hook. f. i. 66.
DICOCAPNOS, Cham. & Schlecht. in Linnaea, i. (1826) 568.
PLATYCAPNOS, Bernh. in Linnaea, viii. (1833) 471.
RUPICAPNOS, Pomel, Mat. Fl. Atl. 16 (1850).
abyssinica, *Hornem. in Nov. Act. Soc. Sc. Upsal.* Ser. III. ii. (1856–58) 275.—Abyssin.
acaulis, Wulf. in Jacq. Coll. ii. 303 = Corydalis ochroleuca.
acaulescens, Peterm. ex Hammar, in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1856–58) 270 = Schlechteri.
aegyptiaca, Steinh. in Guill. Arch. Bot. i. (1838) 416 = densiflora.
affinis, Griff. Notul. iv. 719 = parviflora.
africana, Hammar, in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1856–58) 293 = media.
africana, *Lam. Encyc.* ii. 589.—Afr. bor.
agraria, Gren. & Godr. Fl. Fr. i. 67 = major.
agraria, *Lag. Gen. et Sp. Nov.* 21.—Reg. Mediterr.
alba, Mill. Gard. Dict. ed. VIII. n. 3 = Corydalis lutea.
albiflora, Kit. in Linnaea, xxii. (1868) 491 = Corydalis albiflora.
alexandrina, Ehrenb. ex V. Tenore, in Bull. Acc. Asp. Nat. i. (1842) 24 = judaica.
alexandrina, Gasp. in Rendic. Acc. Sc. Napoli, i. (1843) 51 = capreolata.
altaica, Ledeb. in Mém. Acad. Sc. Pétersb. v. (1812) 551 = Corydalis pauciflora.
amaryssia, *Boiss. & Heldr. ex Boiss. Fl. Orient.* i. 188. —Cyrenaica.
ambigua, Fall. ex Cham. & Schlecht. in Linnaea, i. (1826) 558 = Corydalis ambigua.
anatolica, *Boiss. Diagn.* Ser. I. viii. 14.—Europ. austr. ; As. Min.
angustifolia, Bieb. Fl. Taur. Cauc. ii. 146 = Corydalis angustifolia.
angustifolia, Gilib. Fl. Lituan. ii. 113 = officinalis.
apiculata, Hammar, in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1856–58) 287 = muralis.
apiculata, Lange, Ind. Sem. Hort. Haun. (1854) 28 ; in Ann. Sc. Nat. Sér. IV. ii. (1854) 371 = media.
aquilegifolia, Patz. ex DC. Syst. ii. 116 = Corydalis pauciflora.
arundanae, *Boiss. ex Willk. & Lange, Prod. Fl. Hisp.* iii. 850.—Hispan.

FUMARIA :—
asepala, *Boiss. Fl. Orient.* i. 135.—As. Min.
atlantica, *Coss. & Dur. in Willk. & Lange, Prod. Fl Hisp.* iii. 858, nomen.—Afr. bor.
aurea, Muhl. ex Willd. Enum. Hort. Berol. 740 = Corydalis aurea.
Bastardi, Bor. in Ducb. Rev. Bot. ii. (1847) 359 = media.
bicuritis, Vahl, ex Steud. Nom. ed. II. i. 650 = Dicentra formosa.
bicalcarata, Stokes, Bot. Mat. Med. iv. 9 = Dicentra Cucullaria.
bitornata, Pall. ex Steud. Nom. ed. II. i. 650 = Corydalis pauciflora.
Boissieri, *Haussk. in Flora*, lvi. (1873) 494.—Mesopotam.
Boraei, Jord. Cat. Jard. Bot. Grenoble (1849), 15 = media.
bracteata, Heldr. ex Boiss. Fl. Orient. i. 137 = pikerminea.
bracteata, Steph. ex Willd. Sp. Pl. iii. 858 = Corydalis bracteata.
bracteosa, *Pomel, Nouv. Mat. Fl. Atl.* 220.—Alger.
bulbosa, Gueldenst. ex Ledeb. Fl. Ross. i. 99 = Corydalis caucasica.
bulbosa, Linn. Sp. Pl. 699 = Corydalis tuberosa, fabacea.
bulbosa, Thunb. Fl. Jap. 277 = Corydalis decumbens.
Burchelli, DC. ex Steud. Nom. ed. II. i. 650 = officinalis.
caespitosa, *Losc. ex Willk. & Lange, Prod. Fl. Hisp* iii. 854.—Hispan.
calycina, Bab. in Trans. Bot. Soc. Edinb. i. (1841) 84 = densiflora.
capnoides, Linn. Sp. Pl. 700 = Corydalis lutea.
capreolata, Auch. ex Parl. Giorn. Bot. Ital. i. i. (1844) 143 = macrocarpa.
capreolata, Curt. Fl. Lond. ii. 145 = media.
capreolata, Delile, ex Parl. Giorn. Bot. Ital. i. i. (1844) 145 = agraria.
capreolata, Leight. Fl. Shropsh. 844 = muralis.
capreolata, *Linn. Sp. Pl.* 701.—Europ. ; Afr. bor.
carinata, Schur, Enum. Pl. Transs. 38 = Schleicheri.
caucasica, Hoffm. Hort. Mosq. (1808) n. 1394 ; ex Bieb. Fl. Taur. Cauc. iii. 438 = Corydalis angustifolia.
caudata, Lam. Encyc. ii. 589 = Corydalis longiflora.
cava, Mill. Gard. Dict. ed. VIII. n. 7 = Corydalis tuberosa.
Chavini, Reut. Cat. Pl. Vasc. Gen. ed. II. 10 = Vaillantii.
cilicica, *Haussk. in Flora*, lvi. (1873) 493.—Armenia.
cirrhata, Rohde, ex DC. Syst. iii. 134 = media.
clavicujata, Linn. Sp. Pl. 701 = Corydalis claviculata.
confusa, Jard. Cat. Jard. Dijon (1845), 18 = media.
corymbosa, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) 26. t. 6 = africana.
crassifolia, Desf. Fl. Atlant. i. 126 = Sarcocapnos crassifolia.
Cucullaria, Linn. Sp. Pl. 699 = Dicentra Cucullaria.
Cucullaria, Steller, in Pall. N. Nord. Beitr. ii. 300 = Dicentra lachenalioides.
cucullata, Linn. Syst. ed. X. 1152 = Dicentra Cucullaria.
decumbens, Thunb. in Nov. Act. Petrop. xii. (1801) 102 = Corydalis decumbens.
defensa, Heuff. in Flora, xxxvi. (1853) 619 = anatolica, parviflora.
densiflora, *DC. Cat. Hort. Monsp.* 118.—Gallia.
demiflora, DC. Syst. ii. 127 = parviflora.
demiflora, Delile, ex Nym. Consp. 27 = officinalis.
diffusa, Arn. ex Parl. Giorn. Bot. Ital. i. i. (1844) 127 = officinalis.
diffusa, Mœnch, Meth. Suppl. 215 = parviflora.
digitata, Lgl. Fl. Spa, ii. 69 = Corydalis bulbosa.
disjuncta, Jord. ex Nym. Consp. 27 = media.
Echlonlana, Sond. ex Harv. & Sond. Fl. Cap. i. 18 = officinalis.
elegans, Jord. ex Nym. Consp. 27 = capreolata.
enneaphylla, Linn. Sp. Pl. 700 = Sarcocapnos enneaphylla.
exrnia, Ker-Gawl. in Bot. Reg. t. 50 = Dicentra eximia.
fabacea, Bieb. Fl. Taur. Cauc. ii. 145 = Corydalis caucasica.
fabacea, Retz. Prod. Fl. Scand. ed. II. 167 = Corydalis fabacea.
flabellata, *Gasp. in Rend. Acc. Sc. Napoli*, i. (1842) 51. —Europ. austr.
flavula, Raf. in Med. Repos. N. York, v. (1808) 350 = Corydalis aurea.

FUMARIA:—
Fleischeri, Sond. ex Hammar, in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1856–58) 201 = macrocarpa.
formosa, Andr. Bot. Rep. t. 393 = Dicentra formosa.
formosa, Poir. Encyc. Suppl. vi. 684 = Dicentra eximia.
fungosa, Ait. Hort. Kew. iii. 1 = Adlumia cirrhosa.
gallicana, *Hausskn. in Flora*, lvi. (1873) 547.—Hispan. Gaillardoti, *Boiss. Fl. Orient.* i. 189.—Syria.
Gasparinii, Bab. in Trans. Bot. Soc. Edinb. i. (1841) 95 = officinalis.
glauca, Curt. Bot. Mag. t. 179 = Corydalis glauca.
glauca, Jord. Pugill. Fl. Nov. 8 = parviflora.
Gussonii, Boiss. Diagn. Ser. I. vii. 13 = media.
Halleri, Willd. Prod. Fl. Berl. 229 = Corydalis bulbosa.
Haldreichii, Boiss. Diagn. Ser. II. i. 16 = Thureti.
Impatiens, Pall. Reise, iii. 286 = Corydalis sibirica.
incisa, Thunb. in Nov. Act. Petrop. xii. (1801) 104 = Corydalis incisa.
intermedia, Ehrh. Beitr. vi. 146 = Corydalis fabacea.
intermedia, Gueldenst. ex Ledeb. Fl. Ross. i. 99 = Corydalis caucasica.
Jankae, *Hausskn. in Flora*, lvi. (1873) 491.—Hungar.
Jordani, Guss. Enum. Pl. Inar. 13. t. 2 = media.
judaica, *Boiss. Diagn.* Ser. I. viii. 15.—Syria; Aegypt.
Kralikii, Jord Cat. Jard. Dijon (1848), 19 = anatolica.
Laggeri, Janka, ex Nym. Consp. 28 = Schleicheri.
Laggeri, Jord. Pugill. Fl. Nov. 7 = Vaillantii.
lenensis, Pall. ex DC. Syst. ii. 122 = Corydalis paeoniaefolia.
leucanthe, Vir. Fl. Cors. Sp. Nov. 12 = parviflora.
Lichtensteinii, Schlecht. in Linnaea, i. (1826) 568 = officinalis.
littoralis, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 261 = densiflora.
Lobelii, *Tausch, in Flora*, xxii. (1839) 495.—Austria.
longiflora, Willd. Sp. Pl. iii. 860 = Corydalis longiflora.
longipes, *Coss. & Dur. in Bull. Soc Bot. Fr.* ii. (1855) 806.—Algeria.
lutea, Lam. Mant. ii. 258 = Corydalis lutea.
lutea, Thunb. Fl. Jap. 277 = Corydalis pallida.
macrocarpa, *Parl. Pl. Nov.* 5.—Graecia; Syria.
macrosepala, *Boiss. Elench.* 8.—Hispan.; Afr. bor. occ.
macrosepala, Schur, Enum. Pl. Transs. 88 = densiflora.
major, *Badaro, in Brugnat. Giorn. Fis.* II. ix. (1826) 72.—Europ. austr.
major, Roth, Tent. Fl. Germ. i. 300 = Corydalis tuberosa.
major, V. Tenore, in Bull. Acc. Asp. Nat. i. (1842) 24 = media.
malacitana, Haussk. in Flora, lvi. (1873) 548 = gaditana.
Marschalliana, Pall. in Nov. Act. Petrop. x. (1797) 315 = Corydalis Marschalliana.
media, DC. Syst. ii. 134 = major.
media, Fries, ex Nym. Consp. 27 = muralis.
media, Ledeb. Fl. Ross. i. 105 = densiflora.
media, *Loisel. in Desv. Journ. Bot.* ii. (1809) 357.—Europ.; Reg. Mediterr.
media, Montagne, ex Parl. Giorn. Bot. Ital. i. 1. (1844) 149 = capreolata.
media, Ortm. ex Nym. Consp. 28 = Schleicheri.
media, Reichb. Fl. Germ. Exqus. 697 = Wirtgeni.
megalocarpa, Boiss. & Spruu. Diagn. Ser. I. i. 68 = macrocarpa.
Meyeri, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) 1. 286, in textu.—Reg. Caucas.
micrantha, Lag. Gen. et Sp. Nov. 21 = densiflora.
microcarpa, *Boiss. Diagn.* Ser. II. v. 16, in obs.—Armenia.
microstachys, *Kralik, ex Hausskn. in Flora*, lvi. (1873) 529.—Aegypt.
minor, Roth, Tent. Fl. Germ. i. 300 = Corydalis bulbosa.
monizae, *y. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 268.—Ins. Cap. Verd.
macrocnemiaides, Schur, Enum. Pl. Transs. 88 = densiflora.
Mundtii, Boiss. & Reut. Pugill. Fl. Nov. 5 = media.
Mundtii, *Sprengl. Syst. iv. Cur. Post.* 264.—Afr. austr.
muralis, Sond. ex Koch, *Syn. Fl. Germ.* ed. II. 1017.—Europ.

FUMARIA:—
neapolitana, Vaill. ex Parl. Giorn. Bot. Ital. i. 1. (1844) 149 = capreolata.
nemorosa, *Lejac. in Natural. Sicil.* ii. (1883) 298.—Sicil.
nobilis, Linn. Syst. ed. XII. 469 = Corydalis nobilis.
nanaidies, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 806.—Afr. bor.
officinalis, Burm. f. Fl. Cap. Prod. 20 = Corydalis vesicaria.
officinalis, Chaub. Nouv. Fl. Pélop. 45 = densiflora.
officinalis, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 247 = Vaillantii.
officinalis, *Linn. Sp. Pl.* 700.—Geront. temp.
officinalis, Schimp. ex Hammar, in Nov. Act. Soc. Sc Upsal. Ser. III. ii. (1856–58) 275 = abyssinica.
officinalis, Webb & Berth. Phyt. Canar. i. 58 = capreolata.
officinarum, Neck. Delic. Gallo-Belg. ii. 298 = officinalis.
oxyloba, Boiss. Diagn. Ser. I. viii. 14 = macrocarpa.
paeoniaefolia, Steph. ex Willd. Sp. Pl. iii. 859 = Corydalis paeoniaefolia.
pallida, Salisb. Prod. 377 = Dicentra Cucullaria.
pallida, Thunb. in Nov. Act. Petrop. xii. (1801) 108 = Corydalis pallida.
pallidiflora, Jord. in Schultz, Arch. Fl. Fr. Allem. 305 = densiflora.
Parlatoriana, Kralik, ex Boiss. Fl. Orient. i. 136 = densiflora.
parviflora, *Lam. Encyc.* ii. 567.—Geront. temp.
parviflora, Sibth. & Sm. Fl. Graec. Prod. ii. 50 = densiflora.
parviflora, Wight & Arn. Prod. 18 = Vaillantii.
pauciflora, Steph. ex Willd. Sp. Pl. iii. 861 = Corydalis pauciflora.
peregrina, Rudolph, in Mém. Acad. Sc. Pétersb. i. (1806–8) 279. t. 19 = Dicentra lachenaliaeflora.
Petteri, Koch, Syn. Fl. Germ. ed. II. 485 = muralis.
Petteri, Reichb. Ic. Fl. Germ. iii. 1 = media.
Petteri, V. Tenore, in Bull. Acc. Asp. Nat. i. (1842) 25 = officinalis.
pikermiana, *Boiss. Diagn.* Ser. II. vi. 9.—Graecia.
platycalyx, *Pomel, Nouv. Mat. Fl. Atl.* 239.—Alger.
prehensilis, Ascher. in Verh. Bot. Ver. Brand. v. (1863) 223 = capreolata.
prehensilis, Kit. Ind. Pl. Hort. Pesth. (1812) 10; et in Linnaea, xxvii (1856) 493 = densiflora.
pulchella, Kralik, ex Walp. Ann. iii. 28 = anatolica.
pulchella, Salisb. Prod. 377 = officinalis.
pumila, Host, Fl. Austr. ii. 504 = Corydalis pumila.
racemosa, Thunb. in Nov. Act. Petrop. xii. (1801) 108 = Corydalis racemosa.
recognita, *Lacroix, in Bull. Soc. Bot. Fr.* vi. (1859) 851.—Europ.
recta, Mitch. Fl. Bor. Am. ii. 51 = Adlumia cirrhosa.
Reuteri, *Boiss. Diagn.* Ser. I. vii. 13.—Hispania.
Reuteri, Lange, in Kjoeb. Vidensk. Meddel. (1865) 65 = parviflora.
rosea, Steud. Nom. ed. I. 348 = Corydalis rosea, *Zeyh.*
rostellata, Knaf, in Flora, xxix. (1846) 290 = densiflora.
rupestris, Boiss. & Reut. Pugill. Fl. Nov. 4 = agraria.
rupestris, Nym. Consp 27 = gaditana.
rutifolia, Sibth. & Sm. Fl. Graec. Prod. ii. 49 = Corydalis rutaefolia.
sarcocapnoides, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 806.—Algeria.
sativa, Scop. ex Steud. Nom. ed. II. i. 650 = Corydalis nobilis.
saxatilis, *Clem. ex Lange, in Willk. & Lange, Prod. Fl. Hisp.* iii. 886, in obs.—Hispan.
saxicola, *Nym. Consp.* 28.—Hispan.
scandens. E. Mey. in Drège, Zwei Pfl. Docum. 186 = Mundtii.
Schangini, Pall. in Act. Petrop. vi. (1779) 267. t. 14 = Corydalis longiflora.
Schleicheri, *Soyer-Willem. Obs.* 17.—Europ.
Schrammii, Nym. Consp. 28 = Vaillantii.
semperviresa, Linn. Sp. Pl. 700 = Corydalis bulbosa.
sepium, Boiss. & Reut. Diagn. Ser. II. i. 16 = media.
serotina, Guss. in Ind. Sem. Hort. Boccad. Ital. i. t. (1844) 190 = media.
sibirica, Linn. f. Suppl. 314 = Corydalis sibirica.
sicula, Pisan. Cat. Hort. Panorm. (1816) = parviflora.
solida, Ehrh. Beitr, vi. 146 = Corydalis bulbosa.
speciosa, Jord. Grain. Jard. Grenoble (1848) 15 = capreolata.

FUMARIA:—
spectabilis, Bisch. Del. Sem. Hort. Heidelb. (1849) 4 = agraria.
spectabilis, Linn. Sp. Pl. 699 = Dicentra spectabilis.
spicata, Balb. ex Parl. Giorn. Ital. i. 1. (1844) 141 = Vaillantii.
spicata, Bert. ex Parl. l. c. 145 = agraria.
spicata, Gorter, Fl. Prov. Belg. 188 = parviflora.
spicata, *Linn. Sp. Pl.* 985.—Lusitan.
Sturmii, Opiz, Natural. x. (1825) 267 = densiflora.
suplua, *Janka, in Termesz. Fuzetek*, i. (1877) 80.—Transyl.
tenuiflora, Fries, in Bot. Notiser (1857), 51 = Wirtgeni.
tenuiflora, Janka, ex Nym. Consp. 28 = Vaillantii.
tenuifolia, Pers. Syn. ii. 270 = Dicentra lachenaliaeflora.
tenuifolia, *Roth, Catalecta, fasc.* ii. 89.—Europ.
tenuifolia, Symons, Syn. Pl. Brit. 200; Gaertn. Mey. & Scherb. Fl. Wett. iii. 14 = parviflora.
tenuiscta, *Ball, in Journ. Bot.* xi. (1873) 297; *et in Journ. Linn. Soc.* xvi. (1877) 318.—Marocco.
tenuisecta, Syme, Eng. Bot. ed. III. i. 118 = Vaillantii, parviflora.
tetragona, Stokes, Bot. Mat. Med. iv. 8 = Corydalis lutea.
Thureti, *Boiss. Diagn.* Ser. II. i. 15.—Graecia; Syria.
transsilvanica, Schur, Enum. Pl. Transs. 88 = densiflora.
turbinata, *Sm. ex DC. Syst.* ii. 132.—Oriens.
uniflora, Sieber, Reise Kreta, ii. 320 = Corydalis rutaefolia.
vagans, Jord. Grain. Jard. Grenoble (1849), 15 = media.
Vaillantii, *Loisel. in Desv. Journ. Bot.* ii. (1809) 358.—Geront. temp.
Vaillantii, Tenore, Fl. Nap. ii. 119 = capreolata.
vesicaria, Linn. Sp. Pl. 701 = Corydalis vesicaria.
Wirtgeni, *Koch, Syn. Fl. Germ.* ed. II. 1018.—Europ.

FUNASTRUM, Fourn. in Ann. Sc. Nat. Sér. VI. xiv. (1882) 388. *ASCLEPIADEAE.*
angustisimum, *Fourn. l. c.*—Ins. Galap.
subrutescens, *Fourn. l. c.* 389.—Bras.

FUNCKIA, Dennst. Schluss. Hort. Malab. 32 (1818) = **Lumnitzera**, Willd. (Combretac.).
Karabandel, Dennst. l. c. = L. racemosa.

FUNCKIA, Dum. Anal. Fam. 60 (1829) = **Funkia**, Spreng. (Liliac.).

FUNCKIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 19 = **Astelia**, Banks & Soland. (Liliac.).
magellanica, Willd. l. c. = A. pumila.

FUNIFERA, Leandr. ex C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 357. *THYMELAEACEAE.* Benth. & Hook. f. iii. 195.
Borgel, Fl. Flum. 150 (1825).
Nebbia, Mart. ex Meisan. in Mart. Fl. Bras. v. i. 67 (1855).
fascicularis, *Meisan. in Mart. Fl. Bras.* v. i. 68.—Bras. latifolia, *Fisch. & Mey. ex C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 357, nomen.—Hab.? utilis, *Leandr. ex C. A. Mey. l. c.*—Bras.

FUNIUM, Willem. in Usteri, Ann. Bot. xvIII. (1798) 26 = **Furcroea**, Vent. (Amaryllid.).
gigantum, Willem., l. c. = Furcroea gigantea.

FUNKIA, Benth. & Hook. f. Gen. iii. 751 (1883) = FUNCKIA, Willd. = **Astelia**, Banks & Soland. (Liliac.).

FUNKIA, Endl. Gen. 1182, in syn. (1841) = FUNCKIA, Dennst. = **Lumnitzera**, Willd. (Combretac.).

FUNKIA, Spreng. Anleit. ii. i. 246 (1817); Syst. ii. 40 (1825). *LILIACEAE*, Benth. & Hook. f. iii. 774.
Bryocles, Salisb. in Trans. Hort. Soc. i. (1812) 338.
Hosta, Tratt. Arch. Gew. i. 55 (1812).
Libertia, Dum. Comm. Bot. 9 (1822).
Niobe, Salisb. in Trans. Hort. Soc. i. (1812) 335.
Saussurea, Salisb. in Trans. Linn. Soc. viii. (1807) 11.

FUNKIA:—
alba, Sweet, Hort. Brit. ed. I. 409 = subcordata.
albomarginata, Hook. Bot. Mag. t. 3657 = lancifolia.
cærulea, Sweet, Hort. Brit. ed. I. 409 = ovata.
cordata, Siebold, ex Stend. Nom. ed. II. i. 651 = Sieboldiana.
cucullata, Siebold, ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 304 = glauca.
Fortunei, *Baker, in Gard. Chron.* vi. (1876) 36.—Japon.
glauca, *Siebold, ex Miq. in Verh. Kon. Akad. Wetensch.* II. iii. (1869) 303 ; *et in Arch. Néerl.* iv. (1869) 338.—Japon.
grandiflora, Sieb. & Zucc. in Fl. des Serres, Sér. I. ii. (1846) t. 158, 159 = subcordata.
japonica, Hort. ex Regel, Cat. Pl. Hort. Aksakov. 61 = grandiflora.
lanceæfolia, Dens. ex G. Don, in Loud. Hort. Brit. Suppl. i. 589 = lancifolia.
lancolata, Siebold, ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 300 = ovata.
lancifolia, *Spreng. Syst.* ii. 41.—Japon.
longipes, *Franch. & Sav. Enum. Pl. Jap.* ii. 529.—Japon.
maculata, Siebold, ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 303 = Sieboldiana.
marginata, Siebold, ex Miq. l. c. 303 = ovata.
ovata, *Spreng. Syst.* ii. 40.—Japon.
Sieboldiana, *Hook. Bot. Mag.* t. 3663.—Japon.
Sieboldii, Lindl. in Bot. Reg. (1839) t. 50 = præc.
sinensis, Siebold, ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 303 = Sieboldiana.
spathulata, Siebold, ex Miq. l. c. 300 = ovata.
subcordata, *Spreng. Syst.* ii. 41.—Japon.
umbellata, Hort.—Cf. Gard. Chron. (1868) 1040 = Sieboldiana ?
undulata, Otto & Dietr. Allg. Gartenz. i. (1833) 119 = lancifolia.
univittata, Siebold, ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 302 = Sieboldiana.
viridi-marginata, Siebold, ex Miq. l. c. 299 = ovata.
viridis, Siebold, ex Miq. l. c. 301 = Sieboldiana.

FURCARIA, Boiv. ex Baill. Etud. Gén. Euph. 56 (1858) = **Croton**, Linn. (Euphorb.).
Boiviniana, Baill. l. c. = C. Boivinianus.

FURCARIA, Kostel. Allg. Med.-Pharm. Flora, v. 1856 (1836) = **Ehretsena**, Linn. (Malvac.).
Cavanillesii, Kostel. l. c. 1857, in obs. = H. cannabina.
Roxburghii, Kostel. l. c. 1856 = H. furcatus.
surattensis, Kostel. l. c. = H. surattensis.

FURCRÆA, Vent. in Bull. Soc. Philom. i. (1793) 65 : reimpr. in Usteri, Ann. Bot. xix. (1796) 54. *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 729.
FOURCROYA, Haw. Syn. Pl. Succ. 78 (1812).
FOURCROYA, Spreng. Anleit. ii. i. 238 (1817).
FUNIUM, Willem. in Usteri, Ann. Bot. xviii. (1798) 26.
FURCROEA, Benth. & Hook. f. Gen. iii. 739 (1883).
FURCROYA, Rafin. Princ. Somiol. 31 (1814) ; Endl. Gen. 181 (1841).
agavephylla, Brot. ex Schult. Syst. vi. 731 = cubensis.
Alteni, Jacobi, in Abh. Schles. Ges. (1869) 175 = cubensis.
aspera, M. Roem. Syn. Enast. 293 = cubensis.
atroviridis, Jacobi & Goeff. in Jacobi, Vers. Syst. Ord. Agav. 273 = gigantea.
australis, Haw. Syn. Pl. Succ. 74 = Doryanthes excelsa.
Baillieti, *Jacobi, in Abh. Schles. Ges.* (1869) 166.—Hab. ?
Bedinghausii, *C. Koch, Wochenschr.* vi. (1863) 234 ; *Belg. Hortic.* (1863) t. 19.—Am. austr.
cantula, Haw. Syn. Pl. Succ. Suppl. 42 = Agave viripara.
Commelyni, *Kunth, Enum. Pl.* v. 843.—Am. trop.
cubensis, *Vent. in Bull. Soc. Philom.* i. (1793) 66 ; *Haw. Syn. Pl. Succ.* 75.—Am. trop.
Demoulinsiana, *Jacobi, Vers. Syst. Ord. Agav.* 297.—Mexic.
depauperata, *Jacobi, l. c.* 293.—Hab. ?
elegans, *Trel. Hort. Panorm.* 13. t. 4.—Mexic.
flavo-viridis, *Hook. Bot. Mag.* t. 5163.—Mexic.
fœtida, Haw. Syn. Pl. Succ. 78 = gigantea.

FURCRÆA:—
geminispina, *Jacobi, Vers. Syst. Ord. Agav.* 282.—Hab. ?
gigantea, *Vent. in Bull. Soc. Philom.* i. (1793) 66.—Am. trop.
Lindeni, *Jacobi.—Cf. Gard. Chron.* (1869) 887 ; *et in Illustr. Hortic.* xxi. (1874) t. 186.—N. Granat.
lipidensis, *Jacobi, in Abh. Schles. Ges.* (1869) 169.—Hab. ?
longæva, *Karw. & Zucc. in Nov. Act. Nat. Cur.* xvi. ii. (1833) 666. t. 48.—Mexic.
macra, Hort. Par. ex Jacobi, in Hamb. Gartenz. xxii. (1866) 412 = flagelliformis.
madagascariensis, *Haw. Syn. Pl. Succ. Suppl.* 42.—Madag.
pubescens, *Tod. Ind. Sem. Hort. Panorm.* (1877) 38 ; *Nuov. Sp. Fourcroya*, 9.—Hab. ?
rigida, Haw. Syn. Pl. Succ. 74 = Agave rigida.
rigida, Laudry, ex Jacobi, in Hamb. Gartenz. xxiii. (1867) 314 = Demoulinsiana.
Selloa, *C. Koch, Wochenschr.* iii. (1860) 22.—Am. austr.
sobolifera, Cels, ex Jacobi, in Hamb. Gartenz. xxiii. (1867) 314 = Demoulinsiana.
spinosa, Targ. Tozz. Obs. Bot. n. 16 = tuberosa.
stricta, *Jacobi, in Abh. Schles. Ges.* (1869) 171.—Am. trop.
tuberosa, *Ait. Hort. Kew.* ed. II. ii. 303.—Am. austr.
tuberosa, Kunth & Bouché, in Linnæa, xix. (1847) 831.—Beschoreria tuberosa.
undulata, *Jacobi, in Abh. Schles. Ges.* (1869) 170.—Mexic.
vallecultata, Jacobi, l. c. 175 = cubensis.
viridis, Hemsl. Biol. Centr. Am. Bot. iii. 353 = gigantea.

FURCROYA, Rafin. Princ. Somiol. 31 (1814) ; Endl. Gen. 181 (1841) = præc.

FURERA, Adans. Fam. ii. 193 (1763) = **Satureia**, Tourn. (Labiat.).

FURNROHRIA, Lindl. Veg. Kingd. 778 (1847) = **Fuernrohria**, C. Koch (Umbellif.).

FUSANUS, R. Br. Prod. 355 (1810). *SANTALACEÆ*, Benth. & Hook. f. iii. 225.
EUCARYA, Mitch. Three Exped. Austral. ii. 100 (1839).
LAURIPULA, Lour. Fl. Cochinch. 88 (1790).
MIDA, A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 376.
acuminatus, *R. Br. Prod.* 355.—Austral.
alternifolius, *R. Br. in Salt, Abyss. App.* 63, nomen.—Abyss.
compressus, Lam. Tabl. Encyc. iii. 485. t. 842 = Osyris compressa.
crassifolius, *R. Br. Prod.* 355.—Austral.
diversifolius, Miq. in Lehm. Pl. Preiss. i. 617 = persicarioides.
emarginatus, *Miq. l. c.*—Austral.
glaucus, Gaudich. ex A. DC. in DC. Prod. xiv. 685 = crassifolius.
forsterius, Blanco, Fl. Filip. ed. II. 53 = Viscum orientale.
pedatus, *Spreng. Syst.* i. 490.—Cochinch.
persicarioides, *F. Muell. ex Benth. Fl. Austral.* vi. 216.—Austral.
spicatus, *R. Br. Prod.* 355.—Austral.

FUSANUS, Murr. Syst. Veg. ed. XIII. 765 (1774) = **Colpoon**, Berg. (Santal.).
compressus, Murr. l. c. = C. compressum.

FUSIFILUM, Rafin. Fl. Tellur. ii. 27 (1836) = **Urgines**, Steinh. (Liliac.).
coarctatum, Rafin. l. c. = Excremis ramosa.
physodes, Rafin. l. c. = U. physodes.
pusillum, Rafin. l. c. = U. pusilla.

FUSSIA, Schur, Enum. Pl. Transs. 754 (1866) = **Aira**, Linn. (Gramin.).
capillaris, Schur, l. c. = A. capillaris.
caryophyllea, Schur, l. c. = A. caryophyllea.
præcox, Schur, l. c. = A. præcox.

FUSTICUS, Rafin. New Fl. Am. iii. 48 (1836) = **Maclura**, Nutt. (Urticac.).
glabra, *Tatacha*, *tinctoria*, *vera*, *xanthoxylon*, Rafin. l. c. iii. 44, 45 = M. tinctoria.

G

GABERTIA, Gaudich. in Freyc. Voy. Bot. 495 (1826) = **Grammatophyllum** Blume (Orchid.).
scripta, Gaudich. l. c. = Grammat. speciosum.

GABILA, Baill. Adansonia, x. (1871) 155. *MENISPERMACEÆ*.
longifolia, *Baill. l. c.* 156.—Ins Timor.

GAERDTIA, Klotzsch, in Monatsb. Berl. Acad. (Maer. 1854) 128 ; et Abh. Akad. Berl. (1854) 49. t. 8 A = **Begonia**, Linn.
argentea, Klotzsch, in Monatsb. Berl. Acad. (Maer. 1854) 128 = B. maculata.
Kunthiana, Klotzsch, l. c. = B. Kunthiana.
maculata, Klotzsch, l. c. = B. maculata.
stenobotrys, Klotzsch, l. c. nomen. = B. undulata.
undulata, Klotzsch, l. c. = B. undulata.

GAERTNERA, Lam. Illustr. ii. 278. t. 167 (1791). *LOGANIACEÆ*, Benth. & Hook. f. ii. 798.
ANDERSONIA, Schlecht. in Roem. & Schult. Syst. v. 21 (1819).
FRUCTESCA, DC. in Meisn. Gen. 259 ; Comm. 168 (1839).
PRISTIDIA, Thw. Enum. Pl. Zeyl. 149 (1859).
SYNKSIA, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 351.
acuminata, Benth. in Journ. Linn. Soc. i. (1857) 112 = Koenigii.
Arthrosoma, Steud. Nom. ed. II. i. 651 = calycina.
arenaria, *Baker, in Journ. Linn. Soc.* xx. (1883) 209. —Madag.
bifida, Boj. Hort. Maurit. 217 = psychotrioides.
cærulea, Boj. l. c. = Chasalia grandifolia.
calycina, *Boj. l. c.*—Ins. Maurit.
capitata, *Boj. l. c.* 216.—Ins. Maurit.
cardiocalyx, *Boiv. ex Baill. Adansonia*, xii. (1879) 238.—Madag.
cærulea, A. DC. in DC. Prod. ix. 35 = Chasalia grandiflora.
coriacea, Steud. Nom. ed. II. ii. 410, in syn. = edentata.
crassiflora, *Boj. Hort. Maurit.* 216.—Ins. Maurit.
cuneifolia, Boj. l. c. = Chasalia diversiflora ?
cymosa, Boj. l. c. 218 = Chasalia Boryana.
divaricata, Thw. Enum. Pl. Zeyl. 125 = Koenigii.
edentata, *Boj. Hort. Maurit.* 216.—Ins Maurit.
Gardneri, Thw. Enum. Pl. Zeyl. 203 = Walkeri.
grisea, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit Ind.* iv. 92.—Singapore.
hongkongensis, *Seem. Bot. Voy. Herald*, 384.—China.
incarnata, Boiv. ex Boj. Hort. Maurit. 217 = Chasalia divaricata.
inflexa, *Baill. Adansonia*, xii. (1879) 237.—Madag.
Junghuhniana, Miq. Fl. Ind. Bat. ii. 388 = Koenigii.
Koenigii, *Wight, Ic.* iv. t. 1313.—Ind. or. ; Malaya.
lanceolata, *Bentham, ex DC. Prod.* ix. 33.—Ins. Maurit.
longiflora, Gaertn. f. Fruct. iii. 58 = vaginata.
longifolia, *Boj. Hort. Maurit.* 216.—Ins. Maurit.
macroborya, *Baker, in Journ. Linn. Soc.* xx. (1883) 208.—Madag.
macrostipula, *Baker, l. c.* 207.—Madag.
obesa, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 98.—Singapore.
obovata, *Baker, in Journ. Bot.* xx. (1882) 218.—Madag.
occidentalis, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 383.—Afr. trop.
oxyphylla, *Benth. in Journ. Linn. Soc.* i. (1857) 112. —Singapore.
paniculata, *Benth. in Hook. Niger Fl.* 459.—Afr. trop.
parviflora, Boj. Hort. Maurit. 217 = psychotrioides.
pendula, *Boj. l. c.*—Ins. Maurit.
phanerophlebia, *Baker, in Journ. Linn. Soc.* xxi. (1885) 495.—Madag.
phyllocapsa, *Baker, l. c.* xx. (1882) 207.—Madag.
phyllostachya, *Baker, l. c.* xxi. (1885) 495.—Madag.
psychotrioides, *Baker, Fl. Maurit.* 281.—Ins. Maurit.
quadrisetis, A. DC. Prod. ix. 34 = psychotrioides.
rosea, *Thw. ex Benth. in Journ. Linn. Soc.* i. (1857) 112.—Zeylan.
rotundifolia, *Boj. Hort. Maurit.* 216.—Ins. Maurit.

GAERTNERA:—
Siebori, Presl, Bot. Bemerk. 77 = calycina ?
sphaerocarpa, *Baker*, *in Journ. Linn. Soc.* xx. (1883) 208.—Madag.
ternifolia, *Thw. Enum. Pl. Zeyl.* 202.—Zeylan.
Thwaesii, *Benth. in Bull. Soc. Linn. Par.* i. (1879) 209.—Madag.
thyrsiflora, Blume, Mus. Bot. Lugd. Bat. i. 174 = Koenigii.
truncata, A. DC. in DC. Prod. ix. 34 = edentata.
vaginata, Boj. Hort. Maurit. 216 = psychotrioides.
vaginata, *Lam. Tabl. Encyc.* ii. 225 t. 167; *Poir. Encyc. Suppl.* v. 685.—Ins. Maurit.
vaginata, Sieber, ex DC. Prod. ix. 35 = calycina.
viminea, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 91.—Singapore.
Walkeri, *Wight, Illustr.* ii. t. 156. b.—Zeylan.
Zollingeriana, *Miq. Fl. Ind. Bat.* ii. 882.—Sumatra.

GAERTNERA, Retz. Obs. vi. 24 (1791) = **Sphenocles**, Gaertn. (Campanulac.).
Pongati, Retz. l. c. = S. zeylanica.

GAERTNERA, Schreb. Gen. i. 290 (1789) = **Hiptage**, Gaertn. (Malpighiac.).
indica, J. F. Gmel. Syst. 685 = H. Madablota.
laurifolia, Wall. Cat. n. 7365 = H. parvifolia.
obtusifolia, Roxb. Hort. Beng. 32; Fl. Ind. ii. 369 = H. obtusifolia.
racemosa, Roxb. Pl. Corom. i. 19. t. 18 = H. Madablota.

GAERTNERIA, Medic. Phil. Bot i. 45 (1789) = **Franseria**, Cav. (Compos.).

GAERTNERIA, Neck. Elem. ii. 15 (1790) = **Gentiana**, Tourn.

GAGEA, Salisb. in Kon. & Sims, Ann. Bot. ii. (1806) 556. *LILIACEAE*, Benth. & Hook. f. iii. 819.
BULBILLARIA, Zucc. in Abh. Akad. Muench. iii. (1848) 229. t. 2.
HORNUNGIA, Bernh. in Flora, xxiii. (1840) 392.
ORNITHOGALUM, Link, Handb. i. 161 (1829).
PLICOSTIGMA, Turcz. in Trautv. Pl. Imag. Fl. Ross. ii. t. 2 (1844).
SOLENARIUM, Dulac, Fl. Hautes-Pyr. 113 (1867).
Alberti, *Regel, in Act. Hort. Petrop.* vi. (1879) 512.—China.
alexandrina, Boiss. Fl. Orient. v. 208 = reticularis.
amblopetala, *Boiss. & Heldr. Diagn.* Ser. I. vii. 107.—Greec. ; As. Min.
andegavensis, F. Schultz, in Flora, xiv. (1862) 459 = pygmaea.
anisanthos, C. Koch, in Linnaea, xxii. (1849) 230 = Liotardi.
arvensis, Dum. Fl. Belg. 140 ; Schult. f. Syst. vii. 547, 1705 = stellaris.
arvensis, Kar. & Kir. ex Ledeb. Fl. Ross. iv. 138 = pusilla.
aurea, C. Koch, in Linnaea, xxii. (1849) 229 = amblopetala.
Baumgarteniana, Schur, Enum. Pl. Transs. 667 = minima.
belgica, Dum. Fl. Belg. 139 = spathacea.
Billardieri, Kunth, Enum. Pl. iv. 242 = pygmaea.
bohemica, Gren. & Godr. Fl. Fr. iii. 195 = saxatilis.
bohemica, Schult. f. Syst. vii. 549, 1705 = pygmaea.
bonariensis, Schult. f. l. c. 558 = Nothoscordum Poiretii.
bracteata, Schult. f. l. c. 542 = Lloydia alpina.
bracteolaris, *Salisb. in Kon. & Sims, Ann. Bot.* ii. (1806) 556 (err. typ. 256).—Europ. ; Reg. Mediterr.
bulbifera, *Salisb. l. c.* 557.—Europ. or. ; As. trop.
buxbaumiana, Parl, Fl. Palerm. i. 379 (err. typ. 279) = bulbosa ?
callosa, Reichb. Fl. Germ. Excurs. 107 = minima.
callosa, *Schult. f. Syst.* vii. 554.—Reg. Caucas.
caucasica, *Stev. in Denkschr. Acad. Wien*, i. (1885) 15.—Reg. Transcauc.
cespitosa, Hausk. ex Boiss. Fl. Orient. v. 209 = damascena.
chlorantha, *Schult. f. Syst.* vii. 551.—Oriens.
chrysantha, *Schult. f. l. c.* 545.—Sicilia.
circinata, Lond. Hort. Brit. 184 = reticularis.
Clusiana, Schult. f. Syst. vii. 543 = pusilla.
commutata, C. Koch, in Linnaea, xxii. (1849) 227 = reticularis.

GAGEA:—
composita, Opiz, Seznam, 45 = bracteolaris ?
corsica, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 58 = saxatilis.
corsica, Tausch, ex F. Schultz, Arch. Fl. Fr. Allem. 18 = bulbosa.
damascena, *Boiss. & Gaill. Diagn.* Ser. II. iv. 105.—Syria.
divaricata, *Regel, in Act. Hort. Petrop.* vi. (1879) 510.—Turkestan.
dschungarica, *Regel, l. c.* 513.—Turkestan.
Durieui, *Parl. Fl. Ital.* ii. 426, in textu.—Algeria.
elegans, Wall. Cat. n. 5065 = fascicularis.
emarginata, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 851.—Sibir. alt.
equitans, Wallr. Beitr. Fl. Hercyn. 195 ; et in Linnaea, xiv. (1840) 562 = bracteolaris.
erubescens, Schult. f. Syst vii. 545 = Liotardi.
fasciicularis, *Salisb. in Kon. & Sims, Ann. Bot.* ii. (1806) 555.—Europ. ; As. bor.
fibrosa, *Schult. f. Syst.* vii. 552.—Afr. bor.
filiformis, *Kunth, Enum Pl. iv.* 237.—Sibir.
filiformis, Merckl. ex Herd. in Bull. Soc. Nat. Mosc. (1864) ii. 286 = rufescens.
fistulosa, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 180 ; Duby, Bot Gall. ed. II. i. 467 = Liotardi.
foliosa, Parl. Fl. Palerm. i. 378 (err. typ. 275) = chrysantha.
foliosa, *Schult. f. Syst.* vii. 1708.—Europ or. ; As. Min.
glacialis, *C. Koch, in Linnaea*, xxii. (1849) 228.—Oriens.
glauca, Sweet, Hort. Brit. ed. I. 418 = bracteolaris.
gracilis, Welw. ex Reichb. Ic. Fl. Germ. x. 18 = stellaris.
Granatelli, Boiss. Fl. Orient. v. 208 = reticularis.
Granatelli, *Parl Fl. Palerm.* i. 376 (err. typ. 276).—Sicil.
grandiflora, *Boiss. & Hurt, ex Tchihat. Asie Min. Bot.* iii. 584, nomen.—Armenia.
granulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 132, in textu.—Sibir. occ.
Guicciardii, Heldr. ex Nym. Consp 725 = stellaris.
hybrida, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 75.—Transsylv.
intermedia, Schlecht. ex Reichb. Ic. Fl. Germ. x. 18 = Liotardi.
Kotschyi, *in Denkschr. Acad. Wien*, i. (1885) 80.—Mesopotam.
mauritanica, Dur. ex Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 50 = foliosa.
minima, *Ker-Gawl. in* [*Quart.*] *Journ. Roy. Inst.* i. (1816) 180.—Europ. ; As. bor.
minutiflora, *Regel, in Act. Hort. Petrop.* iii. ii. (1875) 292.—Turkestan.
monticola, Payne, Palest. Exp. Soc. 124 ; ex Boiss. Fl. Orient. v. 209 = damascena.
Moorcroftiana, Wall. Cat. n. 5068 = fascicularis.
nebrodensis, *Nym. Syll.* 372.—Sicil.
nevadensis, Boiss. Elench. 85 = foliosa.
nivalis, Huet. ex Nym. Consp. 726 = foliosa.
Olgae, *Regel, in Act. Hort. Petrop.* iii. ii. (1875) 292.—Turkestan.
Ova, *Stapf, in Denkschr. Acad. Wien*, i. (1885) 16.—Persia.
pauciflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 113.—Dahuria.
peduncularis, Wall. Cat. n. 5066 = reticularis.
persica, *Boiss. Diagn.* Ser. I. vi. 106.—As. occ. ; Reg. Himal.
podolica, Schult. f. Syst. vii. 545 = pusilla ?
polymorpha, Boiss. Voy. Bot. Espagne, ii. 611 = foliosa.
polymorpha, F. Schultz, Arch. Fl. Fr. Allem. 17 = bracteolaris.
polyphylla, *Stev. in Bull. Soc. Nat. Mosc.* xxx. (1857) ii. 281.—Tauria.
pratensis, Dum. Fl. Belg. 140 ; Schult. f. Syst. vii. 536, 1702 = bracteolaris.
pulchella, Wall. Cat. n. 5064 = Lloydia alpina.
pusilla, *Schult. f. Syst.* vii. 548, 1708.—Europ.
pygmaea, *Salisb. in Kon. & Sims, Ann. Bot.* ii. (1806) 557.—Europ. ; As. Min.
pygmaea, Schult. f. Syst. vii. 547, 1705 = foliosa.
reflexa, Casm. ex Reichb. Ic. Fl. Germ. x. 18 = pusilla ?

GAGEA:—
reticularis, *Salisb. in Kon. & Sims, Ann. Bot.* ii. (1806) 557.—Oriens ; As. occ.
reticulata, Schult. f. Syst. vii. 542 = praec.
rigida, Boiss. & Sprun. Diagn. Ser. I. vi. 108 = reticularis.
rubicunda, *Meinsh. Fl. Ingr.* 352.—Ingria.
rubro-viridis, Boiss. & Kotschy, Diagn. Ser. II. iv. 106 = Lloydia rubro-viridis.
rufescens, *Regel, Ind. Sem. Hort. Petrop.* (1862) 28.—Hab. ?
sacculifera, *Regel, in Act. Hort. Petrop.* vi. (1879) 510.—Turkestan
sarmentosa, C. Koch, in Linnaea, xxii. (1849) 230 = reticularis.
saxatilis, Bor. ex Nym. Consp. 725 = pygmaea.
saxatilis, Parl. Fl. Ital. ii. 439, partim = nebrodensis.
saxatilis, *Schult. f. Syst.* vii. 549, 1705.—Bohemia.
Schreberi, Reichb. Fl. Germ. Excurs. 107 = bracteolaris.
serotina, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 180 = Lloydia alpina
setifolia, *Baker, in Journ. Linn. Soc.* xviii. (1880) 101.—Afghan.
Soleirolii, Mutel, Fl. Fr. iii. 298 = foliosa.
spathacea, *Salisb. in Kon. & Sims, Ann. Bot.* ii. (1806) 556 (err. typ. 256).—Europ.
stellaris, *Salisb. l. c.*—Europ. ; As. Min.
stenopetala, Reichb. Fl. Germ. Excurs. 107 = bracteolaris.
Sternbergii, Loud. Hort. Brit. 184 = minima.
stipitata, Merckl. ex Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 512 = persica.
striata, Sweet, Hort. Brit. ed. I. 418 = Lloydia alpina.
succedanea, *Griseb. & Schenk, in Wiegm. Archiv*, xviii. (1852) 858.—Banat.
sylvatica, Loud. Hort. Brit. 184 = fascicularis.
Sawitzi, Besa. ex Schult. f. Syst. vii. 550 = pygmaea.
taurica, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) ii. 83 = reticularis.
tauricola, *Schott & Kotschy, ex Tchihat. Asie Min. Bot.* ii. 584, nomen.—Cilicia.
Theobaldi x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 119.—Europ.
thrioides, C. A. Mey. ex Boiss. Fl. Orient. v. 210 = persica.
transbaicalensis, *Turcz. ex Steud. Nom.* ed. II. i. 652.—Sibir. baical.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 75 = fascicularis.
tranversalis, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) ii. 81 = bracteolaris.
trifora, Schult. f. Syst. vii. 551 = Lloydia trifora.
triphylla, C. Koch, in Linnaea, xxii. (1849) 228 = reticularis.
uniflora, Schult. f. Syst. vii. 553 = Tulipa uniflora.
villosa, Duby, Bot. Gall. ed. II. i. 467 = stellaris.

GAGERNIA, Klotzsch, in Schomb. Faun. et Fl. Gui. 1139 (1848), absque desc. = *OCHNACEAE*.
casiquiocensis, *Klotzsch, l. c.*—Guiana.

GAGIA, St. Lag. in Ann. Soc. Bot. Lyon, vii. 175, 303 (1881) = **Gagea**, Salisb. (Liliac.).

GAGNEBINA, Neck. Elem. ii. 453 (1790); DC. Mém. Lég. 428. t. 64. *LEGUMINOSAE*, Benth. & Hook. f. i. 591.
Commersonii, DC. Prod. ii. 482 = tamaricina.
pterocarpa, Baill. in Bull. Soc. Linn. Par. i. (1883) 554 = tamaricina.
tamaricina, *DC. Prod.* ii. 482.—Ins. Maurit.

GAGNEBINIA, Spreng. Gen. i. 351 (1830) = praec.

GAHNIA, Forst. Char. Gen. 51. t. 26 (1776). *CYPERACEAE*, Benth. & Hook. f. iii. 1066.
DIDYMONEMA, Presl, Diagn. Nov. Pl. Gen. (1829); Symb. Bot. i. t. 8 (1830).
EPIANDRIA, Presl, in Oken, Isis, xxi. (1828) 270.
HEXALEPIS, Boeck. in Flora, lviii. (1875) 118.
LAMPOCARYA, R. Br. Prod. 238 (1810).
MELACHNE, Schrad. ex Nees, in Linnaea, ix. (1834) 301.
MORELOTIA, Gaudich. in Freyc. Voy. Bot. 416. t. 28 (1826).
PRAECELLANTHUS, Stend. in Zoll. Syst. Verz. Ind. Archip. 61 (1854).

GAHNIA :—
Prittacoschoenus, Nees, in Lehm. Pl. Preiss. ii. 87 (1846), in obs.
Sydgantheus, Steud. Syn. Pl. Cyp. 153 (1855).
affinis, Steud. Syn. Pl. Cyp. 164 = Gaudichandii.
ancistrophylla, *Benth. Fl. Austral.* vii. 415.—Austral.
aromatica, Hook. f. Handb. N. Zeal. Fl. 306 = Gaudichandii.
aristata, *Benth. Fl. Austral.* vii. 416.—Austral.
aspera, *Spreng. Syst.* ii. 114.—Austral.; Ins. Pacif.
Banchryi, *R. Mann, in Proc. Am. Acad.* vii. (1867) 210.—Ins. Sandvic.
congesta, Boeck. in Linnæa, xxxviii. (1874) 353 = globosa.
decompositis, Benth. Fl. Austral. vii. 417 = Preissii.
deusta, *Benth. l. c.* 416.—Austral.
erythrocarpa, R. Br. Prod. 239 = psittacorum.
Gaudichandii, *Steud. Syn. Pl. Cyp.* 164.—N. Zel.; Ins. Sandvic.
globosa, *H. Mann, in Proc. Am. Acad.* vii. (1867) 210. —Ins. Sandvic.
gonyocarpa, Steud. Syn. Pl. Cyp. 164 = psittacorum.
Hectori, T. Kirk, in Trans. N. Z. Inst. ix. (1877) 551 = pauciflora.
javanica, *Mor. Syst. Veg. Zoll. Arch. Ind.* 98.—Malaya; Ins. Pacif.
lacera, *Steud. Syn. Pl. Cyp.* 164.—N. Zel.
lanigera, *Benth. Fl. Austral.* vii. 415.—Austral.
leptostachya, Boeck. in Linnæa, xxxviii. (1874) 354.—Ins. Sandvic.
leucocarpa, R. Br. Prod. 239 = psittacorum.
leucosperma, Nees, in Linnæa, ix. (1834) 301 = psittacorum.
melanocarpa, *R. Br. Prod.* 239.—Austral.
melanocarpa, Hook. f. Fl. Tasm. ii. 97 = G. Radula.
microstachya, *Benth. Fl. Austral.* vii. 414.—Austral.
mucronata, Boeck. in Linnæa, xxxviii. (1874) 342 = globosa.
pauciflora, *T. Kirk, in Trans. N. Z. Inst.* i. (1868) 94. —N. Zel.
polyphylla, *Benth. Fl. Austral.* vii. 415.—Austral.
Preissii, *Nees, in Lehm. Pl. Preiss.* ii. 87.—Austral.
procera, *Forst. Char. Gen.* 52.—N. Zel.
psiocaulon, Boeck. in Linnæa, xxxviii. (1874) 345 = psittacorum.
psittacorum, *Labill. Nov. Holl. Pl.* i. 89. t. 115.—Austral.
psittacorum, Nees, ex Benth. Fl. Austral. vii. 414 = Sieberi.
psittacorum, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 270 = tristis.
Radula, *Benth. Fl. Austral.* vii. 417.—Austral.
ramosensis, Steud. Syn. Pl. Cyp. 164 = sapera.
rigida, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 551. —N. Zel.
schoenoides, *Forst. f. Prod.* 98.—Ins. Societ.
setifolia, *Steud. Syn. Pl. Cyp.* 164.—N. Zel.
Sieberi, *Boeck. in Linnæa*, xxxviii. (1874) 343.—Austral.
Sieberiana, *Kunth, Enum. Pl.* ii. 382 = psittacorum.
tetragonocarpa, *Boeck. l. c.* 347.—Austral.
trifida, *Labill. Nov. Holl. Pl.* i. 89. t. 116.—Austral.
tristis, *Nees, in Linnæa*, ix. (1834) 301, nomen.—China; Malaya.
Urvilleana, Kunth, Enum. Pl. ii. 382 = psittacorum.
Wichurai, Boeck. in Linnæa, xxxviii. (1874) 348 = tristis.
xanthocarpa, *Hook. f. Handb. N. Zeal. Fl.* 306.—N. Zel.

GAJADENDRON, G. Don, Gen. Syst. iii. 481 (1834) = **Loranthus**, Linn.
eugenioides, G. Don, l. c. = L. eugenioides.
laurifolium, G. Don, l. c. = L. laurifolius.
nitidum, G. Don, l. c. 482 = L. nitidus.
punctatum, G. Don, l. c. = L. punctatus.
puracense, G. Don, l. c. = L. puracensis.
Tagua, G. Don, l. c. 481 = L. Tagua.

GAIDENDRON, Endl. Ench. 299 (1841) = prasc.

GAILLARDA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 176 = seq.

GAILLARDIA, Fouger. in Mém. Acad. Sc. Par. 1786 (1788) 5. t. 1, 2. *COMPOSITÆ*, Benth. & Hook. f. ii. 414.
Agassizia, A. Gray & Engelm, in Journ. Bost. Soc. Nat. Hist. vi. (1857) 229.

GAILLARDIA :—
Calonnea, Buchoz, Ic. (1786) t. 196.
Cercostylon, Less. Syn. Comp. 289 (1832).
Galardia, Lam. Encyc. ii. 590 (1786).
Guentheria, Spreng. Syst. iii. 556 (1826).
Othake, Rafin. New Fl. Am. iv. 73 (1836).
Polletherus, Rafin. in Am. Monthly Mag. (1818) 266.
Polypteris, Less. in Linnæa, vi. (1831) 518.
Vialaria, L'Hérit. Dis. (1788 ?).
acaulis, *A. Gray, in Proc. Am. Acad.* x. (1874) 78.—Am. bor. occ.
acaulis, Pursh, Fl. Am. Sept. ii. 742 = Actinella acaulis.
alternifolia, Steud. Nom. ed. II. i. 652, aphalm. vide *Galardia alternifolia*.
amara, Rafin. Fl. Ludov. 69 = pulchella.
amblyodon, *Y. Gay, in Ann. Sc. Nat.* Sér. II. xii. (1839) 62.—Texas.
aristata, *Pursh, Fl. Am. Sept.* ii. 573.—Am. bor.
arizonica, *A. Gray, Syn. Fl. N. Am.* i. ii. 353.—Arizon.
aromatica, Baill. Hist. Pl. viii. 307 = Cephalophora aromatica.
bicolor, Ell. Sketch, ii. 449 = lanceolata.
bicolor, Lam. Encyc. ii. 590; Illustr. t. 708 = pulchella.
bicolor, Pursh, Fl. Am. Sept. ii. 572; Hook. Fl. Bor. Am. i. 313 = aristata.
connata, *A. Gray, in Proc. Am. Acad.* xviii. (1882–83) 109; xix. (1883) 84.—Mexic.
Drummondii, DC. Prod. v. 552 = pulchella.
fimbriata, Michx. Fl. Bor. Am. ii. 142 = Helenium fimbriatum.
glauca, Baill. Hist. Pl. viii. 307 = Cephalophora glauca.
lanceolata, *Michx. Fl. Bor. Am.* ii. 142.—Am. bor.
lobata, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 459 = pulchella.
megapotamica, *Baker, in Mart. Fl. Bras.* vi. iii. 276. —Bras.
mexicana, *A. Gray, in Proc. Am. Acad.* xviii. (1882–83) 109.—Mexic.
odorata, *Lindh. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 280.—Am. bor. occ.
picta, Sweet, Brit. Flow. Gard. N. S. iii. t. 267 = pulchella.
pinnatifida, *Torr. in Ann. Lyc. N. York*, ii. (1828) 214.—Am. bor. occ.
pulchella, *Fouger. in Mém. Acad. Sc. Par.* 1786 (1788) 5.—Am. bor.
Roemeriana, Scheele, in Linnæa, xxii. (1849) 161 = Actinella scaposa.
Rosell, *Regel, Gartenfl.* (1875) 289.—Calif.
rustica, Cass. in Dict. Sc. Nat. xviii. 20 = aristata.
scabiosoides, Griseb. in Goett. Abh. xxiv. (1879) 199 = megapotamica.
scabrata, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 459 = amblyodon.
simplex, *Scheele, in Linnæa*, xxii. (1849) 160.—Am. bor. occ.
spathulata, *A. Gray, in Proc. Am. Acad.* xii. (1877) 59. —Am. bor. occ.
tonalensis, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 40.—Reg. Argent.
tuberculata, Scheele, in Linnæa, xxii. (1849) 349 = simplex.

GAILLIONIA, Endl. Gen. 539 (1838) = seq.

GAILLONIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 158. t. 15. f. 3. *RUBIACEÆ*, Benth. & Hook. f. ii. 144.
Choulettia, Pomel, Nouv. Mat. Fl. Atl. 81 (1874).
Jaubertia, Guill. in Ann. Sc. Nat. Sér. II. xvi. (1841) 60.
Aschers, *Jeneb. & Spach, in Ann. Sc. Nat.* Sér. II. xx. (1843) 67.—Arab.; Persia.
Bragtiad, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 158.—Persia.
calycopteris, *Jeneb. & Spach, Illustr.* i. 147. t. 80.—Arab.; Nubia.
cruciancellolides, *Jeneb. & Spach, l. c.* 149. t. 81.—Persia.
erianthe, *Jeneb. & Spach, l. c.* 145. t. 78.—Persia; Afghan.
humifusa, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xx. (1843) 68 = hymenostephana.
hymenostephana, *Jeneb. & Spach, Illustr.* i. 148. t. 79. —Arab.; Belochist.
incana, Jaub. & Spach, l. c. 148. t. 76 = Bragtiad.

GAILLONIA :—
Oliveri, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 158.—Persia.
puberula, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 886.—Ins. Socotr.
Rebaudiana, *Cass. in Bull. Soc. Bot. Fr.* ii. (1855) 250. —Alger.
Richardiana, Jaub. & Spach, Illustr. i. 143. t. 75 = Bragtiad.
rigida, *Rich. in Mém. Hist. Nat. Par.* v. (1834) 299. t. 15.—Hab.?
Seowitzii, *DC. Prod.* iv. 574 (Sowitsii).—Armenia; Persia.
thymoides, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 887.—Ins. Socotr.
tinctoria, *Balf. f. l. c.* 886.—Ins. Socotr.

GAIMARDA, Juss. in Dict. Sc. Nat. xlv. 272 (1827) = seq.

GAIMARDIA, Gaudich. in Freyc. Voy. Bot. 418. t. 30 (1826). *CENTROLEPIDEÆ*, Benth. & Hook. f. iii. 1026.
Alepyrum, Hieron. Centrolep. 103 (1878).
australis, *Gaudich. in Freyc. Voy. Bot.* 419. t. 2. f. 1.—Reg. Magellan.
ciliata, *Hook. f. Fl. Antarct.* 88.—Ins. Antarct.
pallida, *Hook. f. l. c.* 86.—Ins. Antarct.
setacea, *Hook. f. Fl. N. Zel.* i. 267.—N. Zel.

GAIODENDRON, Endl. Gen. 801, in syn. (1839) = GAIADENDRON, G. Don = **Loranthus**, Linn.

GAISSENIA, Rafin. in Med. Repos. N. York, v. (1808) 351 = **Trollius**, Linn. (Ranunc.).
erna, Rafin. l. c. = T. americanus.

GAJATI, Adans. Fam. ii. 328 (1763) = **Aeschynomene**, Linn. (Legumin.).

GAKENIA, Heist. Syst. 9 (1748) (Quid ?).

GALACTANTHUS, Lem. in Orb. Dict. v. 763 (1849) = **Galanthus**, Linn. (Amaryll.).

GALACTEA, Wight, Ic. ii. 5 (1841) = **Galactia**, P. Br. (Legumin.).

GALACTELLA, Rafin. Fl. Tellur. ii. 44 (1836) = GALATELLA, Cass. = **Aster**, Tourn. (Compos.).

GALACTIA, P. Br. Hist. Jamaic. 298 (1756). *LEGUMINOSÆ*. Benth. & Hook. f. i. 535.
Betencourtia, Sc. Hil. Voy. Brés. i. 376 (1833).
Collaea, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 96.
Heterocarpaea, Schecle, in Linnæa, xxi. (1848) 467.
Leucodictyon, Dals. in Hook. Kew Journ. ii. (1850) 264.
Odonia, Bertol. Luesb. Herb. 55 (1822).
Sweetia, DC. Prod. ii. 251 (1825).
angustifolia, Kunth, Mimos. t. 56 = Jussiæana.
angustifolia, Spreng. Syst. iii. 257 = Martii.
argentifolia, *S. Moore, in Journ. Bot.* xv. (1877) 291.—Afr. trop.
Benthamiana, *Micheli, in Kjoeb. Vidensk. Meddel.* (1878) 81.—Bras.
barbertana, DC. Prod. ii. 388 = tenuiflora.
brachycodon, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 176.—Ind. occ.
brachypoda, *Torr. & Gray, Fl. N. Am.* i. 288.—Am. bor.
brachyrachyx, *Benth. in Ann. Wien. Mus.* ii. (1838) 126.—Mexic.
brevistyla, Schlecht. in Linnæa, xii. (1838) 268 = tenuiflora.
canescens, *Benth. in Ann. Wien. Mus.* ii. (1838) 126. —Texas.
coriacea, Nees, in Nov. Act. Nat. Cur. xii. (1824) 80 = Camptosema coriaceum.
cubensis, *H. B. & K. Nov. Gen. et Sp.* vi. 439.—Ind. occ.
cuspidata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 87.—Ins. Moluce.
debilis, Mannk. in Handl. & Da Vriese, Tijdschr. v. (1849) 148 = Sinclairia vestita.
discrepitula, Eng. Bot. Mamft. 93 = Tanaensis labialis.
dubia, *DC. Prod.* ii. 386.—Cosmop. trop.

GALACTIA :—
dumetorum, *Benth. in Ann. Wien. Mus.* ii. (1838) 127. —Bras.
Elliottii, *Nutt. Gen. Am.* ii. 117.—Am. bor.
emarginata, *Desv. in Ann. Sc. Nat. Sér.* I. ix. (1826) 414.—Ind. occ.
filiformis, *Benth. in Ann. Wien. Mus.* ii. (1838) 127. —Am. trop.
filiformis, Wall. Cat. nn. 5510, 5519 = tenuiflora.
flavifiora, *Michels, in Mém. Soc. Phys. Genèv.* xxviii. (1888) n. VII. 24.—Reg. Argent.
floridana, *Torr. & Gray, Fl. N. Am.* i. 288.—Florida.
glabella, *Michx. Fl. Bor. Am.* ii. 62.—Am. bor.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* vi. 428. —Ecuador.
gracillima, *Benth. in Mart. Fl. Bras.* xv. i. 142.—Am. austr.
graminea, Heyne, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 192 = tenuiflora.
heterophylla, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 171.—Texas.
hirta, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 190.—Mexic.
hispidula, Benth. in Linnæa, xxii. (1849) 514 = Calopogonium galactoides.
Jussiæana, *Kunth, Mimos.* 196. t. 55.—Venezuela.
latiifolia, *Desv. in Ann. Sc. Nat. Sér.* I. ix. (1826) 414.—Am. austr.
leucocarpa, *Desv. l. c.*—Ins. S. Doming.
Lockhartì, *Griseb. Fl. Brit. W. Ind.* 194.—Ind. occ.
longiflora, *Arn. in Wight & Arn. Prod.* 206.—Ind. occ.
longifolia, *Benth. in Ann. Wien. Mus.* ii. (1838) 127; *Wight, Ic.* t. 493.—Ind. or.
Lorentzii, *Grisel. in Gott. Abh.* xix. (1874) 124.—Reg. Argent.
lucida, A. R. Grah. in Wall. Cat. n. 5511 = tenuiflora.
Macreti, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 120 = mollis.
marginalis, *Benth. in Ann. Wien. Mus.* ii. (1838) 126. —Mexic.
Martii, *DC. Prod.* ii. 238.—Bras.
megalophylla, F. Muell. Fragm. ix. 67 = Lamprolobium megalophyllum.
mexicana, Spreng. Syst. iv. Cur. Post. 238 = Colognania angustifolia.
mollis, *Michx. Fl. Bor. Am.* ii. 61.—Am. bor.
monophylla, *Grisel. Cat. Pl. Cub.* 75.—Cuba.
mucronata, Klotzsch, in Peters, Reise Mossamb. Bot. 39 = tenuiflora.
Mueller, *Benth. Fl. Austral.* ii. 255.—Austral.
Nessii, *DC. Prod.* ii. 238.—Bras.
obcordata, Boiv. ex Baill. in Bull. Soc. Linn. Par. i. (1888) 382 = Teramnus obcordatus.
Odonia, *Griseb. Cat. Pl. Cub.* 75.—Cuba.
oxyphylla, *Benth. in Mig. Pl. Jungh.* 233.—Burma.
parvifolia, *A. Rich. Ess. Fl. Cub.* 414.—Cuba.
pauciflora, *Benth. in Ann. Wien. Mus.* ii. (1838) 127. —Austral.
pendula, *Pers. Syn.* ii. 302.—Am. trop.
Perteriana, Benth. in Mart. Fl. Bras. xv. i. 143, sphalm. = *Berteriana*.
pilosa, Nutt. Gen. Am. ii. 116 = mollis.
pinnata, Mubl. Cat. 67 = Elliottii.
pinnata, Pers. Syn. ii. 302 = Barbieria polyphylla.
prostrata, *Benth. in Ann. Wien. Mus.* ii. (1838) 127. —Austral.
pulchella, Spreng. Syst. iv. Cur. Post. 239 = Colognania pulchella.
purpurea, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1844) 190 = Colognania pulchella.
Purshii, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 413 = glabella.
radicata, *DC. Prod.* ii. 238.—Mexic.
rotundifolia, Blume, ex Miq. Fl. Ind. Bat. i. i. 221 = Shuteria rotundifolia.
Sagræi, Decb. & Walp. in Linnæa, xviii. (1850) 739 = tenuiflora.
scriceo, Boj. Hort. Maurit. 93 = Teramnus labialis.
sericea, *Pers. Syn.* ii. 302.—Ins. Mascar.
sessiliflora, *Torr. & Gray, Fl. N. Am.* i. 289.—Am. bor.
simplicifolia, Dalz. in Hook. Kew Journ. iii. (1851) 209 = Crotalaria Dalzellii.
simplifolia, Spreng. Syst. iv. Cur. Post. 239 = Colognania procumbens.
spiciformis, *Torr. & Gray, Fl. N. Am.* i. 287.—Am. bor.
stenophylla, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 182.—Am. austr.
tenuiflora, Willd ex Wight & Arn. Prod. 206 = dubia.
tephrodes, *A. Gray, Pl. Wright.* ii. 84.—N. Mexic.
terminiflora, Blanco, Fl. Filip. ed. II. 411 = Milletia splendens.
texana, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 170.—Am. bor. occ.
tomentosa, *Spreng. Syst.* iii. 257.—Bras.
tuberosa, *DC. Prod.* ii. 238.—Mexic.
tuberosa, Hook. & Arn. Bot. Beech. Voy. 415 = Calopogonium galactoides.
velutina, *Benth. in Ann. Nat. Hist.* iii. (1839) 437.— Guian.
vicioeformis, *Spreng. Syst.* iii. 257.—Bras
villosa, Eaton & Wright, N. Am. Bot. 248 = mollis.
villosa, Wight & Arn. Prod. 207 = tenuiflora.
Weddelliana, *Benth. in Mart. Fl. Bras.* xv. i. 142.— Bras.
Wrightii, *A. Gray, Pl. Wright.* i. 44.—Mexic.

GALACTION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = præc.
sericeum, St. Lag. l. c. = Galactia sericea.

GALACTITES, Moench, Meth. 558 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 470.
LUPSIA, Neck. Elem. i. 71 (1790), nomen prius.
australis, *Sweet, ex Steud. Nom.* ed. II. i. 653 (Quid !). —Austral.
Duriaei, *Spach, in Duch. Rev. Bot.* i. (1845) 368.— Hispan.
elegans, Nym. Consp. 411 = tomentosa.
flavo-spinosa, *Klatt, in Nov. Act. Nat. Cur.* xli. ii. (1880) 402.—Ind. or.
mutabilis, *Dur. in Duch. Rev. Bot.* i. (1845) 362.— Algeria.
pycnantha, Dur. ex Willk. & Lange, Prod. Fl. Hisp. ii. 200 = Duriæi.
tomentosa, *Moench, Meth.* 558; *DC. Fl. Fr.* iii. 110.— Reg. Mediterr.

GALACTODENDRON, Reichb. Consp. 83 (1828) = seq.

GALACTODENDRUM, Kunth, ex Humb. Relat. Hist. ii. 108, in nota (1819) = Brosimum, Sw. (Urticac.).
chrysophyllifolium, Hem. ex Desv. in Ann. Sc. Nat. Sér. II. xviii. (1842) 312 = Ficus anthelmintica.
ovalifolium, Ham. ex Desv. l. c. = Ficus anthelmintica.
utilis, H. B. & K. Nov. Gen. et Sp. vii. 136 = B. Galactodendron.

GALACTOGLYCHIA, Miq. in Hook. Lond. Journ. Bot vi. (1847) 526 = GALOGLYCHIA, Gasp. = Ficus, Tourn. (Urticac.).

GALANGA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 9; ed. II. 65 (1827) = Alpinia, Linn. (Scitamin.).
officinalis, Salisb. in Trans. Hort. Soc. i. (1812) 281 = A. Galanga.

GALANTHUS, Linn. Syst. ed. I (1735). *AMARYLLIDEAE*. Benth. & Hook. f. iii. 719.
ACROCORION, Adans. Fam. ii. 57 (1763).
ERANGELIA, Rœsalm. ex Linn. Sp. Pl. ed. I. 288 (1753).
GALACTANTHUS, Lem. in Orb. Dict. v. 763 (1849).
NIVARIA, Heist. Syst. ii. 19 (1746).
autumnalis, All. Auct. Fl. Pedem. 33 = Leucojum hyemale.
Clusii, Fisch. ex Steud. Nom. ed. II. i. 653 = nivalis.
Elwesii, *Hook. f. Bot. Mag.* t. 6166.—As. Min.
graecus, *Orph. ex Boiss. Fl. Orient.* v. 145.—Ins. Chios.
Imperati, *Bertol. Fl. Ital.* iv. 5.—Italia.
latifolius, *Salisb. Gen. Pl. Fragm.* 95; *Rupr. in Regel, Gartenfl.* (1866) 190. t. 878.—Mag. Caucas.
montanus, Schur, Enum. Pl. Transs. 658 = nivalis.
nivalis, Falk, Betr. ii. 156 = plicatus.
nivalis, *Linn. Sp. Pl.* 288.—Europ.
Olgæ, *Orph. ex Boiss. Fl. Orient.* v. 146.—Græcia.
plicatus, *Bieb. Fl. Taur. Cauc. Suppl.* 255.—Reg. Caucas.
plicatus, Gaus. Fl. Rar. 140 = Imperati.
plicatus, Hohen. Enum. Pl. Elisabethpol. 226 = nivalis.

GALANTHUS :—
Redoutei, Rupr. in Regel, Gartenfl. (1863) 177, in textu = nivalis.
reflexus, Herb. ex Lindl. Bot. Reg. (1845) Misc. 85 = nivalis.
vernus, All. Fl. Pedem. ii. 255 = Leucojum vernum.

GALAPAGOA, Hook. f. in Proc. Linn. Soc. i. (1845) 277; et Trans. Linn. Soc. xx. (1847) 196 = Coldenia, Linn. (Borragin.).
Darwinii, Hook. f. in Trans. Linn. Soc. xx. (1847) 196 = C. dichotoma.
fusca, Hook. f. l. c. 197.—Ins. Galapag.

GALARDIA, Lam. Encyc. ii. 590 (1786) = Gaillardia, Fouger. (Compos.).

GALARHAEUS, Baill. Etud. Gén. Euph. 284 (1858) sphalm. = seq.

GALARHOEUS, Haw. Syn. Pl. Succ. 148 (1812) = Euphorbia, Linn.
alepicus, Haw. l. c. 151 = E. alepicus.
androsemifolius, Haw. l. c. 146 = E. lucida.
Apios, Haw. l. c. 149 = E. Apios.
carnuliciae, Haw. l. c. 148 = E. carnulicia.
coralloides, Haw. l. c. 146 = E. coralloides.
corollatus, Haw. l. c. = E. corollata.
creticus, Haw. l. c. 149 = E. Characias.
decussatus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 256 = E. Lathyris.
dulcis, Haw. Syn. Pl Succ. 147 = E. dulcis.
emarginatus, Haw. l. c. 145 = E. hyberna.
epithymoides, Haw. l. c. 147 = E. epithymoides.
flaviconus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 149 = E. flaviconus.
genistoides, Haw. Syn. Pl. Succ. 144 = E. genistoides.
heliscopius, Haw. l. c. 152 = E. helioscopia.
hybernus, Haw. l. c. 145 = E. hyberna.
imbricatus, Haw. l. c. = E. imbricata.
junceus, Haw. l. c. 150 = E. aleppica.
Lathyris, Haw. l. c. 148 = E. Lathyris.
linifolius, Haw. l. c. 150 = E. spinosa.
literatus, Haw. l. c. 151 = E. platyphyllos.
micranthus, Haw. l. c. 152 = E. stricta.
Myrsinites, Haw. l. c. 144 = E. Myrsinites.
nicaeensis, Haw. l. c. = E. nicaeensis.
orientalis, Haw. l. c. 148 = E. orientalis.
palustris, Haw. l. c. 145 = E. palustris.
Paralias, Haw. l. c. 144 = E. Paralias.
pilosus, Haw. l. c. 148 = E. pilosa.
Pithyusa, Haw. l. c. 147 = E. Pithyusa.
platyphyllus, Haw. l. c. 151 = E. platyphyllos.
pubescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 149 = E. pubescens.
rigidus, Haw. Syn. Pl. Succ. 150 = E. spinosa.
salicifolius, Haw. l. c. 148 = E. salicifolia.
serratus, Haw. l. c. 149 = E. serrata.
spathulaefolius, Haw. l. c. 146 = E. spathulaefolia.
spinosus, Haw. l. c. 149 = E. spinosa.
strictus, Haw. l. c. 151 = E. platyphyllos.
verrucosus, Haw. l. c. 148 = E. pubescens.

GALARRHAEUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 149 = præc.

GALARRHOEUS, Reichb. Fl. Germ. Excurs. 755 (1832) = præc.

GALASIA, Koch, Syn. Fl. Germ. ed. I. 436 (1837) = GELASIA, Cass. = Soorsonera, Linn. (Compos.).

GALASIA, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 310 = Microseris, D. Don (Compos.).
borealis, Sch. Bip. l. c. = Leontodon borealis.
lanceolata, Sch. Bip. l. c. = M. lanceolata.
scapigera, Sch. Bip. l. c. = M. scapigera.

GALATEA, Cass. in Bull. Soc. Philom. (1818) 165; et in Dict. Sc. Nat. xviii. 56 (1820) = Aster, Tourn. (Compos.).
albiflora, Cass. in Dict. Sc. Nat. xviii. 58 = A. acris.
canescens, Cass. l. c. 57 = A. canus.
intermedia, Cass. l. c. 58 = A. acris.
pauciflora, Cass. l. c. 56 = A. dracunculoides.
punctata, Cass. l. c. 57 = A. acris.
rigida, Cass. l. c. 58 = A. trinervis.

GALATEA, Herb. in Bot. Mag. sub t. 2118 (1890) = **Nerine**, Herb. (Amaryll.).
rosea, Herb. l. c. = N. sarniensis.
sarniensis, Herb. ex Steud. Nom. ed. II. i. 654 = N. sarniensis.

GALATEA, Salisb. in Trans. Hort. Soc. i. (1812) 310, absque char. = **Heuchera**, Herb. (Irid.).
plicata, Baker, in Journ. Linn. Soc. xvii (1877) 101 = E. plicata.
vesperina, Salisb. in Trans. Hort. Soc. i. (1812) 310 = E. plicata.

GALATELLA, Cass. in Dict. Sc. Nat. xxxvii. 468 (1825) = **Aster**, Tourn. (Compos.).
acris, Nees, Gen. et Sp. Aster. 171 = A. trinervis.
acris, F. Schultz, Arch. Fl. 310 = A. acris.
alhiflora, DC. Prod. v. 255 = A. acris.
aragonensis, Nees, Gen. et Sp. Aster. 167 = A. aragonensis
armena, Boiss. Fl. Orient. iii. 161.—Armenia.
asperrima, Nees, Gen. et Sp. Aster. 173 = A. trinervius.
biflora, Nees, l. c. 159 = A. dracunculoides.
cana, Nees, l. c. 163 = A. canus.
canescens, DC. Prod. v. 256 = A. canus.
dahurica, DC. l. c. = A. dahuricus.
desertorum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 376 = A. dracunculoides.
dracunculoides, Nees, Gen. et Sp. Aster. 164; DC. Prod. v. 256 = A. dracunculoides.
fastigiata, Nees, Gen. et Sp. Aster. 169 = A. Hauptii.
glandulosa, Nees, l. c. 174 = Olearia glandulosa.
graminifolia, Hook. Fl. Bor. Am. ii. 15 = Erigeron hysopifolius.
Hauptii, Lindl. ex DC. Prod. v. 256 = A. Hauptii.
hyssopifolia, Nees, Gen. et Sp. Aster. 160 = A. acris.
inscripta, Nees, l. c. 162 = A. acris.
intermedia, Bunk, Ind. ad DC. Prod. ii. 104 = A. acris.
juncea, Lindl. ex DC. Prod. v. 257 = A. altaicus.
leptophylla, Nees, Gen. et Sp. Aster. 167 = A. leptophyllus, *Desf.*
linifolia, Nees, l. c. 170 = A. acris.
Linoyris, Reichb. f. Ic. Fl. Germ. xvi. 8 = A. Linoyris.
lusitanica, Nees, Gen. et Sp. Aster. 165 = A. aragonensis.
Meyendorffii, Regel & Maack, in Regel, Tent. Fl. Ussur. 81 = Heteropappus hispidus.
nemoralis, Hook. Fl. Bor. Am. ii. 15 = Erigeron hysopifolius.
nemoralis, Nees, Gen. et Sp. Aster. 173 = A. nemoralis.
nudiflora, Lindl. ex DC. Prod. v. 257 = A. Hauptii !
obtusifolia, Lehm. Del. Sem. Hort. Hamb. (1837); ex Linnæa, xii. (1838) 111. 83 = Sericocarpus solidagineus.
punctifera, Nees, Gen. et Sp. Aster. 164, in syn. = A. dracunculoides.
platylepis, Nees, ex Torr. & Gray, Fl. N. Am. ii. 108 = Sericocarpus rigidus.
punctata, Nees, Gen. et Sp. Aster. 161 = A. acris.
rigida, DC. Prod. v. 256 = A. trinervis.
Sorrentini, Tod. Ind. Sem. Hort. Panorm. (1857) 42; Linnæa, xxix. (1857-58) 744.—Sicilia.
squamosa, DC. Prod. v. 257 = A. Hauptii.
strigosa, Weinm. in Bull. Soc. Nat. Mosc. xxiii. (1850) 1. 646.—Rossia.
tenuifolia, Lindl. ex DC. Prod. v. 257 = A. Hauptii.
villosa, Reichb. f. Ic. Fl. Germ. xvi. 8 (= *Linoyris villosa*).—Europ. ; Caucas. ; Sibir.

GALATHEA, Liebm. Ind. Sem. Hort. Haun. (1855) 26. —Cf. Linnæa, xxviii. (1856) 362 = **Marion**, Ker-Gawl. (Irid.).
coerulea, Liebm. l. c. = M. coerulea.
speciosa, Liebm. l. c.—Bras.

GALATHEA, Steud. Nom. ed. II. i. 654 (1840) = GALATEA, Herb. = **Nerine**, Herb. (Amaryll.).

GALATHEA, Steud. Nom. ed. II. i. 654 (1840) = GALATEA, Salisb. = **Heuchera**, Herb. (Irid.).

GALATHENIUM, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 449 = **Lactuca**, Tourn. (Compos.).
elongatum, Nutt. l. c. 449 = L. canadensis.
floridanum, Nutt. l. c. = L. hirsuta, floridana.
graminifolium, Nutt. l. c. = L. graminifolia.
integrifolium, Nutt. l. c. = L. integrifolia.
ludovicianum, Nutt. l. c. 444 = L. ludoviciana.
macrophyllum, Nutt. l. c. 448 = L. macrophylla.
multiflorum, Nutt. l. c. = L. leucophæa.
salicifolium, Nutt. l. c. = L. integrifolia, graminifolia.
sanguineum, Nutt. l. c. 444 = L. hirsuta.

GALAX, Linn. Sp. Pl. ed. I. 200 (1753). *DIAPENSIACEÆ*, Benth. & Hook. f. ii. 690.
HELVEDERE, Clayt. ex Gronov. Fl. Virgin. 25 (1739).
BLANDFORDIA, Andr. Bot. Rep. t. 343 (1804).
ERYTHRORHIZA, Michx. Fl. Bor. Am. ii. 34. t. 36 (1803).
SOLENANDRIA, Beauv. ex Vent. Jard. Malm. t. 69 (1803).
aphylla, *Linn. Sp. Pl.* 200.—Am. bor.
rotundifolia, Pursh, Fl. Am. Sept. ii. 446 = aphylla.

GALAXIA, Thunb. Nov. Gen. 50 (1782). *IRIDEÆ*, Benth. & Hook. f. iii. 693.
aurea, *Nois. ex Schult. Mant.* i. 326, nomen.—Hab. ?
ciliata, Pers. Syn. i. 41 = ovata.
graminea, *Thunb. Nov. Gen.* 51.—Afr. austr.
graminifolia, *Harv. Gen. S. Afr. Pl.* ed. II. 376.—Afr. austr.
gracilis, Andr. Bot. Rep. t. 164 = ovata.
ixiæflora, [DC. in] Red. Lil. i. t. 41 = Ixia monadelpha.
magellanica, Steud. Nom. ed. I. 349 = Tapeinia magellanica.
minuta, Ker-Gawl. in Kon. & Sims, Ann Bot. i. (1805) 241 = Pauridia hypoxidioides.
mucronifolia, Salisb. in Trans. Hort. Soc. i. (1812) 315 = ovata.
multiflora, Spreng. Neue Entdeck. i. 251 = ovata.
narcissoides, Willd. Sp. Pl. iii. 583 = Solenomelus biflorus.
obscura, Cav. Diss. 341 = Tapeinia magellanica.
obtusa, Salisb. in Trans. Hort. Soc. i. (1812) 315 = ovata.
ovata, *Thunb. Nov. Gen.* 51.—Afr. austr.
plicata, Jacq. Ic. Pl. Rar. ii. 8. t. 292; Coll. Suppl. 80 = Lapeyronsia fasciculata.
purpurea, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 385 = ovata.
ramosa, [DC. in] Red. Lil. i. sub t. 41 = Ixia monadelpha.
versicolor, Salisb. in Trans. Hort. Soc. i. (1812) 315 = ovata.
violacea, Eckl. Verz. 17 = ovata.

GALBANON, Adans. Fam. ii. 499 (1763) = **Polylophium**, Boiss. (1844) (Umbellif.).

GALBANOPHORA, Neck. Elem. i. 175 (1790) = **Seseli**, Linn. (Umbellif.).

GALBANUM, D. Don, in Phil. Mag. lx. (1823) 46; et in Trans. Linn. Soc. xvi. (1833) 603 = Polylophium, Boiss. (1844) (Umbellif.).
officinale, D. Don, in Trans. Linn. Soc. xvi. (1833) 603. —Orient.

GALE, Tourn. ex Adans. Fam. ii. 346 (1763) = **Myrica**, Linn.
belgica, Dum. Fl. Belg. 10 = M. Gale.

GALEANA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 19 (1824). *COMPOSITÆ*, Benth. & Hook. f. ii. 406.
bastata, *La Llave, l. c.*—Mexic.

GALEANDRA, Lindl. in Bauer, Illustr. Orch. Pl. Gen. t. 8 (1838 ?). *ORCHIDEÆ*, Benth. & Hook. f. iii. 526.
CORYDANDRA, Reichb. Nom. 83 (1841).
angorensis, Reichb. f. in Linnæa, xx. (1847) 680, sphalm. = sq.
anjouensis, *Reichb. f. in Walp. Ann.* vi. 650.—Ins. Comor.

GALEANDRA :—
barbata, *Lem. Illustr. Hortic.* iii. (1856) *Misc.* 86, 89. —Bras.
Baueri, *Lindl. in Bauer, Illustr. Orch. Pl. Gen.* t. 8. —Guiana ; Mexic.
Beyrichii, *Reichb. f. in Linnæa*, xxii. (1849) 854.—Venezuela.
Bonlinwongo, *Reichb. f. in Bot. Zeit.* x. (1852) 935.—Afr. trop.
cristata, Lindl. Bot. Reg. (1844) Misc. 72 = dives.
curvifolia, *Rodrig. Orch. Nov.* i. 85.—Bras.
Devoniana, *Schomb. ex Lindl. Sert. Orch.* t. 37.—Am. austr.
dives, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 98.—N. Granat.
euglossa, Reichb. f. in Bot. Zeit. x. (1852) 935 = Eulophia euglossa.
extinctoria, Lindl. Gen. et Sp. Orch. 187 = Eulophia gracilis.
gracilis, Lindl. l. c. = Eulophia gracilis.
graminoides, *Rodrig. Orch. Nov.* i. 88.—Bras.
Harveyana, *Reichb. f. in Gard. Chron.* (1888) ii. 716. —Hab. ?
hysterantha, *Rodrig. Orch. Nov.* i. 86.—Bras.
juncea, *Lindl. Sert. Orch.* sub t. 37.—Guiana ; Bras.
juncoides, *Rodrig. Orch. Nov.* i. 87.—Bras.
lacustris, *Rodrig. l. c.* 86.—Bras.
lagotana, *Reichb. f. & Warm. in Otia Bot. Hamb.* 88.—Bras.
longibracteata, *Lindl. in Journ. Linn. Soc.* vi. (1862) 138 (err. typ. 881).—Afr. trop.
minax, *Reichb. f. in Gard. Chron.* (1874) i. 786.—Hab. ?
montana, *Rodrig. Orch. Nov.* ii. 175.—Bras.
nivalis, [*Mast.*] *in Gard. Chron.* (1882) i. 587.—Am. trop.
Petersii, Reichb. f. in Linnæa, xx. (1847) 679 = Eulophia Petersii.
Quartiniana, Reichb. f. in Walp. Ann. iii. 552 = Eulophia Quartiniana.
Stangeana, *Reichb. f. in Bonplandia*, iv. (1856) 323. —Bras.
villosa, *Rodrig. Orch. Nov.* i. 85.—Bras.
viridis, *Rodrig. l. c.* ii. 176.—Bras.

GALEARIA, Heist. Syst. 7 (1748) (Quid ?).

GALEARIA, Presl, Symb. Bot. i. 49 (1830) = **Trifolium**, Tourn. (Legumin.).
Bonanni, Presl, l. c. 50 = T. Bonanni.
Cupani, Presl, l. c. = T. physodes.
fragifera, Presl, l. c. = T. fragifera.
resupinata, Presl, l. c. = T. resupinata.
tomentosa, Presl, l. c. = T. tumens.
tumens, Presl, l. c. = T. tumens.

GALEARIA, Zoll. & Mor. Syst. Verz. Zoll. 19 (1845-46). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 287.
BENNETTIA, R. Br. in Benn. Pl. Jav. Rar. 249. t. 50 (1852).
CREMOSTACHYS, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 259.
affinis, *Miq. Fl. Ind. Bat.* i. 12. 480.—Malaya.
angustifolia, *Miq. l. c. Suppl.* 476.—Sumatra.
aristifera, *Miq. l. c.* 471.—Sumatra.
elliptica, *Miq. l. c.* 469.—Sumatra.
Finlaysoni, *Miq. l. c.* i. 11. 480.—Malaya.
fulva, *Miq. l. c.* 480.—Malaya.
Jackiana, *Miq. l. c.*—Malaya.
pedicellata, *Zoll. & Mor. Syst. Verz. Zoll.* 19.—Malaya.
phleboscarpa, *Miq. Fl. Ind. Bat.* i. 11. 480.—Malaya.
sessilis, *Zoll. & Mor. Syst. Verz. Zoll.* 19.—Malaya.
splendens, *Miq. Fl. Ind. Bat. Suppl.* 469.—Sumatra.
sumatrana, *Miq. l. c.*—Sumatra.
Wallichii, *Kurz, For. Fl. Brit. Burm.* ii. 407.—Burma.

GALEARIS, Rafin. Fl. Tellur. ii. 39 (1836) = **Habenaria**, Willd. (Orchid.).
biflora, Rafin. l. c. = H. bifolia !
spectabilis, Rafin. l. c. = Orchis spectabilis.

GALEATA, Wendl. Bot. Beobacht. 36 (1798) (Quid ?).
ferruginea, Wendl. l. c.—Ind. or.

GALEDRAGON, S. F. Gray, Nat. Arr. Brit. Pl. ii. 675 (1821) = Dipsacus, Linn.
pilosus, S. F. Gray, l. c. = D. pilosus.

GALEDUPA, Lam. Encyc. ii. 594 (1786) = **Pongamia**, Vent. (Leguminˢ).
arborea, Roxb. Hort. Beng. 58 = P. glabra.
bijuga, Roxb. l. c.—Hab. ?
elliptica, Roxb. l. c. 53; Fl. Ind. iii. 242 = Derris elliptica.
frutescens, Blanco, Fl. Filip. ed. I. 559 = Derris scandens.
grandiflora, Herb. Jus. ex Vent. Jard. Malm. sub t. 28 = P. grandiflora.
indica, Lam. Encyc. ii. 594 = P. glabra.
macrocarpa, Roxb. Hort. Beng. 58 = Mucuna macrocarpa ?
maculata, Blanco, Fl. Filip. ed. I. 559 = P. glabra.
marginata, Roxb. Hort. Beng. 58; Fl. Ind. iii. 241 = Derris canelfolia.
megasperma, Roxb. Hort. Beng. [98], nomen = Mucuna macrocarpa ?
Piscidia, Roxb. l. c. 53; Fl. Ind. iii. 240 = Millettia Piscidia.
Pongam, Raeusch. Nom. ed. III. 203 = P. glabra.
Pongam, Blanco, Fl. Filip. ed. I. 558 = Gliricidia maculata ?
Pungum, J. F. Gmel. Syst. 1086 = P. glabra.
uliginosa, Roxb. Hort. Beng. 58; Fl. Ind. iii. 243 = Derris uliginosa.

GALEGA, Tourn. ex Linn. Gen. ed. I. 220 (1737).
LEGUMINOSÆ, Benth. & Hook. f. i. 496.
ACCOROMBONA, Endl. Gen. Pl. Suppl. i. 1427 (1841).
CALLOTROPIS, G. Don. Gen. Syst. ii. 228 (1832).
africana, Mill. Gard. Dict. ed. VIII. n. 2 = officinalis.
ambigua, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 121 = Tephrosia ambigua.
apollinea, Delile, Fl. Aegypt. 288. t. 38 = Tephrosia apollinea.
arborescens, Herb. Madr. ex Wall. Cat. n. 5627 = Tephrosia candida.
argentea, Lam. Encyc. ii. 599 = Tephrosia villosa.
astragaloides, R. Br. ex Benth. Fl. Austral. ii. 208 = Tephrosia astragaloides.
Barba-Jovis, Burm. f. Fl. Ind. 172 = Tephrosia villosa.
Berterii, Spreng. ex DC. Prod. ii. 263 = Poitæa galeoides.
bicolor, *Boiss. & Haussk. ex Rogel, Ind. Sem. Hort. Petrop.* (1867) 37; *Boiss. Fl. Orient.* ii. 191.—Mesopot.
biflora, Poir. Encyc. Suppl. iii. 699 = Tephrosia biflora.
bituba, Sweet, Hort. Brit. ed. I. 124 = officinalis.
binata, Spreng. ex Steud. Nom. ed. II. i. 654 = Lathyrus tomentosus.
capensis, Jacq. Coll. ii. 292; Thunb. Prod. Pl. Cap. 134 = Tephrosia capensis.
caribæa, Jacq. Select. Am. 212 = Tephrosia caribæa.
chrysophylla, Steud. Nom. ed. I. 350 = Tephrosia chrysophylla, *Pursh*.
cinerea, Linn. Syst. ed. X. 1172 = Tephrosia cinerea.
cornuta, Linn. f. Suppl. 335 = Dalea Muriali.
cornuta, Rottl. ex Wight & Arn. Prod. 213 = Tephrosia purpurea.
Colensia, Buch.-Ham. in Trans. Linn. Soc xiii. (1822) 547 = Tephrosia purpurea.
Colutea, Burm. f. Fl. Ind. 172 = Indigofera viscosa.
Colutea, Willd. Sp. Pl. iii. 1246 = Tephrosia villosa.
coronillæfolia, Desf. Tabl. ed. I. 194 = Tephrosia coronillæfolia.
crocea, R. Br. ex Benth. Fl. Austral. ii. 205 = Tephrosia crocea.
davurica, Pall. Reise, iii. 743 = Astragalus dahuricus.
diffusa, Roxb. Hort. Beng. 57; Fl. Ind. iii. 337 = Tephrosia purpurea.
dominguensis, Willd. Sp. Pl. iii. 1249 = Tephrosia purpurea.
dubia, Balb. ex DC. Prod. ii. 258 = Galactia dubia.
dubia, Jacq. Coll. ii. 182 = Lessertia diffusa.
enneliana, *Rottl. ex Spreng. Nachr.* i. *Bot. Gart. Halle*, 28 (Quid ?).—Hab. ?
falcata, Thunb. Prod. Pl. Cap. 134 = Tephrosia falcata.
filifolia, Thunb. Fl. Cap. 809 = Argyrolobium filiforme.
filiformis, Jacq. Coll. ii. 848 = Galactia dubia.
filiformis, Thunb. Prod. Pl. Cap. 133; Willd. Sp. Pl. iii. 1239 = Argyrolobium filiforme.
florida, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 423 = Tephrosia chrysophylla, *Pursh*.
frutescens, Mill. Gard. Dict. ed. VIII. n. 3 = Tephrosia frutescens ?

GALEGA:—
fruticosa, Linn. Syst. ed. X. 1172 = Tephrosia senticosa.
genistoides, Thunb. Prod. Pl. Cap. 133 = Cyclopia genistoides.
grandiflora, Vahl, Symb. Bot. ii. 84 = Tephrosia grandiflora.
Hayneii, Willd. Enum. Hort. Berol. Suppl. 51 = Tephrosia tinctoria.
heterophylla, Fisch. ex DC. Prod. ii. 253 = Tephrosia tinctoria.
Heyneana, Roxb. Hort. Beng. 57; Fl. Ind. iii. 334 = Tephrosia tinctoria.
Heyneii, Steud. Nom. ed II. i. 655 = Tephrosia tinctoria.
hirta, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 546 = Tephrosia Hookeriana.
hispidula, Michx. Fl. Bor. Am. ii. 68 = Tephrosia hispidula.
humilis, Thunb. Prod. Pl. Cap. 134 = Tephrosia pallens.
incana, Roxb. Hort. Beng. 57; Fl. Ind. iii. 335 = Tephrosia villosa.
juncea, R. Br. ex Benth. Fl. Austral. ii. 208 = Tephrosia juncea.
lancæfolia, Roxb. Hort. Beng. 57; Fl. Ind. iii. 336 = Tephrosia purpurea.
lanceolata, Hornem. ex Steud. Nom. ed. II i. 655 = Tephrosia senticosa.
linearis, Willd. Sp. Pl. ii. 1249 = Tephrosia linearis.
litteralis, Forst. f. Prod. 52 = Tephrosia purpurea.
litteralis, Linn. Syst ed. XII. 365 = Tephrosia cinerea.
longifolia, Jacq. Coll. ii. 349 = Galactia tenuiflora.
lupinifolia, Burch. ex DC. Prod. ii. 255 = Tephrosia lupinifolia.
maxima, Linn. Syst. ed. X. 1172 = Tephrosia purpurea mimosoides, *Willd. Sp. Pl.* iii. 1249 = Tephrosia mimosoides.
monsana, Georgi, Beschr. Russ. Reich. III. iv. 1181; Schult. Obs. Bot. 154 = orientalis.
mucronata, Thunb. Prod. Pl. Cap. 134 = Lessertia mucronata.
nervosa, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 423 = Tephrosia tinctoria.
oblongata, R. Br. ex Benth. Fl. Austral. ii. 205 = Tephrosia oblongata.
ochroleuca, Jacq. Coll. i. 79 = Vicia ochroleuca.
ochroleuca, Spreng. in Schrad. Journ. (1800) ii. 197 = prec. ?
officinalis, *Linn. Sp. Pl.* 714.—Europ.; As occ. orientalis, *Linn. Encyc.* ii. 596.—Reg. Caucas.
pallens, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 70 = Tephrosia pallens.
parviflora, Salisb. Prod. 338 = Tephrosia stricta.
patula, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) ii. 144.—Tauria.
peucifolia, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 122 = Tephrosia spicata.
pentaphylla, Roxb. Hort. Beng. 57; Fl. Ind. iii. 334 = Tephrosia senticosa.
Perriniana, Spreng. Neue Entdeck. ii. 161 = Tephrosia Perriniana.
persica, Pers. Syn. ii. 328 = officinalis.
pinnata, Thunb. Prod. Pl. Cap. 134 = Rhynchosia ferulæfolia.
piscatoria, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 71 = Tephrosia purpurea.
porrecta, R. Br. ex Benth. Fl. Austral. ii. 206 = Tephrosia porrecta.
procera, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 547 = Tephrosia purpurea.
prostrata, Koen. ex Wight & Arn. Prod. 213 = Tephrosia purpurea.
prostrata, Wendl. Bot. Beobacht. 56 = prec. ?
pubescens, Linn. Encyc. ii. 597 = Vicia ochroleuca.
pulchella, Scop. Delic. Insub. i. 8 = Tephrosia stricta.
pumila, Lam. Encyc. ii. 599 = Tephrosia purpurea.
purpurea, Linn. Sp. Pl. 1172 = Tephrosia purpurea.
pusilla, Thunb. Prod. Pl. Cap. 133 = Tephrosia pusilla.
reticulata, R. Br. ex Benth. Fl. Austral. ii. 205 = Tephrosia reticulata.
rosea, Lam. Encyc. ii. 599 = Tephrosia grandiflora.
senticosa, Linn. Sp. Pl. ed. II. 1068 = Tephrosia senticosa.
sericea, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 545 = Tephrosia purpurea.
sericea, Lam. Encyc. ii. 596 = seq. ?
sericea, Thunb. Prod. Pl. Cap. 134 = Argyrolobium pumilum.

GALEGA:—
Singapou, Buchoz, Hist. Règ. Vég. x. t. 4 = Tephrosia toxicaria.
spicata, Walt. Fl. Carol. 188 = Tephrosia spicata.
spinosa, Linn. f. Suppl. 335 = Tephrosia spinosa.
striata, Thunb. Prod. Pl. Cap. 134 = Lessertia pulchra.
stricta, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 70 = Tephrosia stricta.
ternata, Thunb. Prod. Pl. Cap. 134 = Aspalathus ferruginea.
tinctoria, Lam. Encyc. ii. 598; Roxb. Fl. Ind. iii. 336 = Tephrosia purpurea.
tinctorea, Linn. Syst. ed. X. 1172 = Tephrosia tinctoria.
tomentosa, Vahl, Symb. Bot. ii. 84 = Tephrosia tomentosa.
totta, Thunb. Prod. Pl. Cap. 134 = Tephrosia totta.
toxicaria, Sw. Prod. Veg. Ind. Occ. 108 = Tephrosia toxicaria.
trapezicarpa, Steud. Nom. ed. I. 350 = Indigofera senegalensis.
tricolor, Hook. Exot. Fl. t. 185 = officinalis.
trifoliata, Thunb. Prod. Pl. Cap. 134 = Argyrolobium setosum.
uniflora, Steud. Nom. ed. I. 350 = Tephrosia uniflora.
villosa, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 547 = Tephrosia Hookeriana.
villosa, Linn. Syst. ed. X 1172 = Tephrosia villosa.
villosa, Michx. Fl. Bor. Am. ii. 67 = Tephrosia spicata.
virginiana, Linn. Syst. ed. X. 1172 = Tephrosia virginiana.
virginiana, J. F. Gmel. Syst. 1552, Ind. = præc.
vulgaris, Lam. Fl. Fr. ii. 654 = officinalis.

GALENA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 177 = seq.

GALENIA, Linn. Syst. ed. I (1735). *FICOIDEÆ*, Benth. & Hook. f. i. 854.
KOLLENIA, Fenzl, Symb. Bot. ii. 24. t. 14 (1830).
SIALODES, Eckl. & Zeyh. Enum. 329 (1837).
TEPHRAS, E Mey. ex Harv. & Sond. Fl. Cap. ii. 477 (1862).
affinis, *Sond. in Harv. & Sond. Fl. Cap.* ii. 476.—Afr. austr.
africana, *Linn. Sp. Pl.* 359.—Afr. trop.
asiatica, Burm. f. Fl. Ind. 88 = Salvadora persica.
colesioides, Spreng. Nov. Prov. 18 = Microtea mayparensis.
collina, *Walp. Rep.* ii. 282.—Afr. austr.
crystallina, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 477.—Afr. austr.
Dregeana, *Fenzl, ex Harv. & Sond. l. c.*—Afr. austr.
Eckloniis, Walp Rep. ii. 283 = glauca.
fruticosa, *Sond. in Harv. & Sond. Fl. Cap.* ii. 477.—Afr. austr.
glauca, *Walp. Rep.* ii. 282.—Afr. austr.
hottentottica, Fenzl, ex Harv. & Sond. Fl. Cap. ii. 478 = africana.
hermanniæfolia, *Walp. Rep.* ii. 282.—Afr. austr.
heterophylla, Fenzl, ex Harv. & Sond. Fl. Cap. ii. 475 = spathulata.
hispidissima, *Fenzl, ex Harv. & Sond. l. c.* 478.—Afr. austr.
humifusa, Fenzl, ex Harv. & Sond. l. c. 476 = collina.
linearis, Thunb. Prod. Pl. Cap. 27 = africana.
pallens, *Walp. Rep.* ii. 282.—Afr. austr.
papulosa, Fenzl, ex Harv. & Sond. Fl. Cap. ii. 477 = pruinosa.
papulosa, *Sond. l. c.* 475.—Afr. austr.
portulacacea, *Fenzl. ex Harv. & Sond. l. c.*—Afr. austr.
procumbens, *Linn. f. Suppl.* 237.—Afr. austr.
pruinosa, *Sond. in Harv. & Sond. Fl. Cap.* ii. 477.—Afr. austr.
salsoloides, Fenzl, ex Harv. & Sond. l. c. 478 = fruticosa.
sarcophylla, *Fenzl, ex Harv. & Sond. l. c.* 475.—Afr. austr.
secunda, *Sond. l. c.* 474.—Afr. austr.
sericea, Fenzl, ex Harv. & Sond. l. c. 478 = fruticosa.
spathulata, *Fenzl, ex Harv. & Sond. l. c.* 475.—Afr. austr.
squamulosa, *Fenzl, ex Harv. & Sond. l. c.* 477.—Afr. austr.
tenuifolia, Salisb. Prod. 338 = africana.

GALEOBDOLON, Adans. Fam. ii. 190 (1763) = **Lamium**, Tourn. (Labiat.).
amplexicaule, Nordmann. 394 = L. amplexicaule.
Galeopsis, Curt. Fl. Lond. fasc. iv. t. 40 = L. Galeobdolon.
luteum, Huds. Fl. Angl. ed. II. i. 258 = L. Galeobdolon.
montanum, Reichb. Fl. Germ. Excurs. 860 = L. Galeobdolon.
sylvestre, Wibel, Prim. Fl. Werth. 139 = L. Galeobdolon.
vulgare, Pers. Syn. ii. 122 = L. Galeobdolon.

GALEOGLOSSUM, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 31 = **Prescottia**, Lindl. (Orchid.).
prescottioides, A. Rich. & Gal. l. c. 31 = P. Galeottii.

GALEOLA, Lour. Fl. Cochinch. ii. 520 (1790). *ORCHIDEÆ*, Benth. & Hook. f. iii. 589.
CYNTORIA, Blume, Bijdr. 396. t. 6 (1826).
ERIAXIS, Reichb. f. in Linnæa, xli. (1877) 68.
ERYTHRORCHIS, Blume, Rumphia, i. 200. t. 70 (1835).
HÆMATORCHIS, Blume, l. c. iv. t. 200 B (1848).
LEDGERIA, F. Muell. Fragm. i. 238 (1859).
POGOCHILUS, Falc. in Hook. Journ. Bot. iv. (1842) 73.
altissima, *Reichb. f. Xenia Orch.* ii. 77.—Malaya.
cassythioides, *Reichb. f. l. c.*—Austral.
foliata, *F. Muell. Fragm.* viii. 81.—Austral.
Humblotii, *Reichb. f. in Flora*, lxviii. (1885) 378.—Ins. Comor.
Hydra, *Reichb. f. Xenia Orch.* ii. 77.—Reg. Himal.; Malaya.
javanica, *Benth. & Hook. f. Gen.* iii. 590.—Malaya.
Kuhlii, *Reichb. f. Xenia Orch.* ii. 78.—Malaya.
Ledgeri, *F. Muell. in Fitzg. Austral. Orch.* ii. (1885) ii.—Austral.
Lindleyana, *Reichb. f. Xenia Orch.* ii. 78.—Reg. Himal.
nudifolia, *Lour. Fl. Cochinch.* ii. 520 (Cranichis nudifolia, *Pers.*).—Cochinch.
rigida, *Benth. & Hook. f. Gen.* iii. 590.—N. Caled.
septentrionalis, *Reichb. f. Xenia Orch.* ii. 76.—Japon.

GALEOPSIS, Linn. Syst. ed. I (1735). *LABIATAE* Benth. & Hook. f. ii. 1209.
LADANUM, Dill. ex Linn. Syst. ed. I (1735).
TETRAHIT, Adans. Fam. ii. 190 (1763).
abyssinica, Hort. ex Steud. Nom. ed. II. i. 655 = Plectranthus punctatus.
acuminata, Reichb. Fl. Germ. Excurs. 923 = G. Tetrahit.
alpicola, Nym. Consp. 576 = G. Tetrahit.
angustifolia, Ehrh. ex Hoffm. Deutschl. Fl. ed. II. i. 11. 8 = G. Ladanum.
arvatica, Jord. in Billot, Ann. Fl. Fr. Allem. 180 = G. Ladanum.
arvensis, Salisb. Prod. 82 = G. Ladanum.
Bertoli, Perrier, ex Nym. Consp. 576 = G. Ladanum.
bifida, Boenn. Prod. Fl. Monast. 178 = G. Tetrahit.
calycina, Schoenh. ex Steud. Nom. ed. II. i. 655 = G. Ladanum.
canescens, Schult. Obs. Bot. 108 = G. Ladanum.
cannabina, Pollich, Hist. Pl. Palat. ii. 148 = ochroleuca.
cannabina, Roth, Tent. Fl. Germ. i. 254 = versicolor.
cannabina, Willd. Sp. Pl. iii. 93 = G. Tetrahit.
carpetana, *Willk. in Flora*, xxxv. (1852) 282.—Hispan.
dubia, Leers, Fl. Herborn. 133 = ochroleuca.
elegans, Bor. Fl. Cent. Fr. ed. III. ii. 356, in obs. = ochroleuca.
Filholiana, Timb. in Bull. Soc. Bot. Fr. i. (1854) 214 = G. Ladanum.
Gacogravi, Nym. Consp. 576 = G. Tetrahit.
Galeobdolon, Linn. Sp. Pl. ed. I. 580 = Lamium Galeobdolon.
glabra, Des Etangs, in Bull. Soc. Bot. Fr. xxiii. (1876) 204 = angustifolia ?
glandulosa, C. Koch, in Linnæa, xxi. (1848) 681 = G. Ladanum.
giaucescens, *Rent. ex Ardoino, Fl. Alpes Mar.* 299.—Europ. austr.
grandiflora, Bieb. Fl. Taur. Cauc. ii. 461 = G. Ladanum.
grandiflora, Roth, Tent. Fl. Germ. i. 254 = ochroleuca.
grandiflora, Suter, Fl. Helv. ii. 18 = G. Tetrahit.

GALEOPSIS:—
hirsuta, Linn. Sp. Pl. 580 = Stachys hirta.
hispanica, Mill. Gard. Dict. ed. VIII. n. 6 = Stachys hirta.
hispida, Thunb. Prod. Pl. Cap. 96 = Stachys Thunbergii.
intermedia, Vill. Prosp. 21 = G. Ladanum.
Ladanum, *Linn. Sp. Pl.* 579.—Europ.; Reg. Caucas.
Lavarnbergwi, Martr. Fl. Tarn. 565 = G. Tetrahit.
latifolia, Hoffm. Deutschl. Fl. ii. 8 = G. Ladanum.
leucantha, Nym. Consp. 576 = G. Tetrahit.
lutea, Gilib. Chlor. Orosh. 38 = Lamium Galeobdolon.
maculosa, Lam. Encyc. ii. 601 = Plectranthus punctatus.
magnifolia, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 16 (Quid ?).—Java.
Michelianae, Torre, Fl. Ital. Prod. 66 = G. Tetrahit ?
micrantha, S. F. Gray, Nat. Arr. Brit. Pl. ii. 734 = G. Ladanum ?
moutana, *Rafin. New Fl. Am.* ii. 21.—Am. bor.
neglecta, *Schult. ex Steud. Nom.* ed. II. i. 655.—Hab. ?
nepetaefolia, *Timb. in Billotia.* i. (1864) 19.—Mont. Pyren.
nodosum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 378 = G. Tetrahit.
ochroleuca, *Lam. Encyc.* ii. 600.—Europ.
orientalis, Mill. Gard. Dict. ed. VIII. n. 5 = Stachys spicata.
parviflora, Lam. Encyc. ii. 600 = G. Tetrahit.
Poelii x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 102.—Europ.
praecox, Jord. in Billot, Ann. Fl. Fr. Allem. 21 = G. Tetrahit.
pratensis, Vill. Hist. Pl. Dauph. ii. 388 = ochroleuca.
pubescens, Bess. Prim. Fl. Galic. ii. 27 = G. Tetrahit.
pubescens, Griseb. Spicil. Fl. Rumel. ii. 125 = versicolor.
pyrenaica, *Bartl. Ind. Sem. Hort. Gotting.* (1848).—Europ. austr.
Reichenbachii, Rent. in Compt.-Rend. Soc. Hallér. (1858) 57 = G. Tetrahit.
Reuterii, Reichb. f. Ic. Fl. Germ. xxviii. 17 = G. Ladanum.
segetum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 377 = G. Ladanum.
segetum, Relch. Fl. Moeno-Franc. ii. 8 = ochroleuca.
speciosa, Mill. Gard. Dict. ed. VIII. n. 3 = versicolor.
subalpina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 59 = versicolor.
Tetrahit, *Linn. Sp. Pl.* 579.—Europ.; As. bor.
unicolor, Fries, Summ. Veg. Scand. 14 = versicolor.
urticaefolia, Salisb. Prod. 83; Tausch, in Lotos, ii. (1852) 69 = G. Tetrahit.
Verloti, Jord. in Billot, n. 1946.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 606 = G. Tetrahit.
versicolor, *Curt. Fl. Lond. fasc.* vi. t. 38.—Europ.; Reg. Caucas.
versicolor, Spenn. Fl. Frib. ii. 394 = G. Tetrahit.
villosa, Huds. Fl. Angl. ed. II. i. 256 = ochroleuca.
Wulfeniana Schlecht. Fl. Berol. i. 320 = G. Tetrahit.

GALEOPSIS, Moench, Meth. 397 (1794) = **Stachys**, Tourn. (Labiat.).
fusca, Moench, l. c. = S. alpina.

GALEOTTIA, Nees, in DC. Prod. xi. 311 (1847) = **Glockeria**, Nees (Acanth.).
gracilis, Nees, l. c. = Glockeria gracilis.

GALEOTTIA, A. Rich. in Ann. Sc. Nat. Sér. III. iii. (1845) 25 = **Epygeotalum**, Hook. (Orchid.).
Beaumontii, Lindl. & Paxt. Flow. Gard. i. (1850–51) 12.—Bras.
grandiflora, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 25 = Z. grandiflorum.

GALEOTTIA, Rupr. ex Gal. in Bull. Acad. Brux. ix. ii. (1842) 247 = **Zeugites**, P. Br. (Gramin.).

GALERA, Blume, Bijdr. 415. t. 8. (1825) = **Hypogynm**, S. G. Gmel. (Orchid.).
nutans, Blume, l. c. = E. nutans.
roxes, Blume, Mus. Bot. Lugd. Bat. ii. 168 = E. nutans.

GALIANTHE, Griseb. ex Lorentz, Veget. Nord. Entre-Rios, 73 (1878), nomen; et in Goett. Abh. xxiv. (1879) 166 = **Spermacoce**, Dill. (Rubiac.).
chlorroloides, Griseb. ll. cc. 73; 167 (= *Borreria centranthoides*).—Bras.
fastigiata, Griseb ll. cc. 73; 156.—Reg. Argent.
verbenoides, Griseb. in Goett. Abh. xxiv. (1879) 157 (= *Borreria verbenoides*).—Bras.

GALIASTRUM, Heist. ex Fabric. Hort. Helmst. ed. I. 108 (1759) = **Mollugo** ?, Linn. (Ficoid.).

GALILEA, Parl. Fl. Palerm. i. 297 (1845) = **Cyperus**, Mich.
mucronata, Parl. l. c. 299 = C. schoenoides.

GALIMBIA, Endl. Ench. 879 (1841) sphalm. = PALIMBIA, Bess. = **Peuscdanum**, Tourn. (Umbellif.).

GALINIERA, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 92. t. 1. *RUBIACEAE*, Benth. & Hook. f. ii. 91.
PTYCHOSTOMA, Hochst. in Flora, xxvii. (1844) 1. 23.
caffeoides, *Delile, in Ann. Sc. Nat.* Sér. II. xx. (1843) 93.—Abyss.

GALINSOGA, Ruiz & Pav. Prod. Fl. Per. 110. t. 24 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 390.
VARGASIA, DC. Prod. v. 676 (1836).
VIGOLINA, Polr. Encyc. viii. 618 (1808.
WIBORGIA, Roth, Cat. Bot. fasc. ii. 112 (1800).
alliaeocarpa, Spreng. Syst. iii. 579 = carcasana.
angustifolia, Spreng. Nene Entdeck. ii. 138 = Calea angustifolia.
balbisioides, Steud. Nom. ed. II. i. 656 = Tridax balbisioides.
brachiata, Spreng. Syst. iii. 579 = Calea brachiata.
brachycephana, *Regel, Ind. Sem. Hort. Turic.* (1846) 2.—Hab. ?
calva, Sch. Bip. in Linnæa, xxxiv. (1865–66) 529, nomen = parviflora, Jaegeria hirta.
carcasana, *Sch. Bip l. c. et in Bull. Soc. Bot. Fr.* xii. (1865) 80 = Am. austr.
dentata, Reichb. ex Steud. Nom. ed. II. i. 656 = Tridax trilobata ?
discolor, Spreng. Nov. Prov. 19; et Syst. iii. 577 = Montanoa arborescens.
filiformis, *Hemsl. Diag. Pl. Nov.* 34.—Mexic.
hispida, *Benth. Bot. Voy. Sulph.* 119.—Mexic.
laciniata, *Rets. in Hoffm. Phytogr. Bluetter*, i. (1803) 46.—Hab. ?
linearifolia, Spreng. Syst. iii. 579 = Gutierreria linearifolia.
Mandonii, Sch. Bip. in Linnæa, xxxiv. (1865–66) 529, nomen = Jaegeria hirta.
megapotamica, *Spreng. Syst.* iii. 580.—Bras.
oblongifolia, *DC. Prod.* v. 677.—Peruv.
Pacajigo, *Ruiz, ex Roem. Hort. Turic. fide Steud. Nom.* ed. II. i. 656.—Peruv. ?
parviflora, *Cav. Ic.* iii. 41. t. 281.—Am. austr.
quadriradiata, Ruiz & Pav. Syst. Veg. 198 = parviflora.
quinqueradiata, Ruiz & Pav. l. c. = parviflora.
reniuoa, Hook. & Arn. Bot. Beech. Voy. i. 82 = Gutierreria linearifolia.
serrata, Spreng. Syst. iii. 579 = Calea axillaris.
trilobata, Cav. Ic. iii. 42. t. 283 = Tridax trilobata.
uniflora, *Spreng. Syst.* iii. 580.—Reg. Argent.
unxiolcs, *Griseb. in Goett. Abh.* xxiv. (1879) 198. —Reg. Argent.
urticaefolia, Benth. Bot. Voy. Sulph. 119 = Sabazia urticaefolia.

GALINSOGEA, H. B. & K. Nov. Gen. et Sp. iv. 252. t. 386 (1820) = **Tridax**, Linn. (Compos.).
balbisioides, H. B. & K. l. c. 253. t. 386 = T. balbisioides.

GALINSOGEA, Willd. Sp. Pl. iii. 2286 (1803) = **Galinsoga**, Ruiz & Pav. (Compos.).

GALINSOGAEA, Zucc. in Flora, iv. (1821) 612 = prsec.

GALINSOGEOPSIS, Sch. Bip. in Seem. Bot. Herald, 306 (1857) = **Parthenium**, A. Gray (Compos.).
apiianthoides, Sch. Bip. l. c. 307 = P. apiianthoides.

GALINSOJA, Roth, Catalect. fasc. iii. 78 (1806) = **Galinsoga**, Ruiz & Pav. (Compos.).

GALINZOGA, Dum. Fl. Belg. 72 (1827) = prsec.

GALION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113, 186 = **Galitum**, Llan. (Rubiac.).
aparinum, St. Lag. l. c. = G. Aparine.
molluginum, St. Lag. l. c. = G. Mollugo.

GALIOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 398 = **Asperula**, Linn. (Rubiac.).
glauca, Fourr. l. c. = A. galloides.

GALIOPSIS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 79 = **Galeopsis**, Linn. (Labiat.).

GALIPEA, Aubl. Pl. Gui. ii. 662. t. 269 (1775).
RUTACEÆ, Benth. & Hook. f. i. 385.
Costa, Vell. Fl. Flum. 19 (1829); t. t. 48 (1827).
Endostephium, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 227.
Sciuris, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 150, 163.
Ticorea, St. Hil. in Bull. Soc. Philom. (1822) 132.
aromatica, *Spreng. Syst. iv. Cur. Post.* 91.—Guian, Arrabidæ, *Stend. Nom. ed.* II. i. 656.—Bras.
bracteata, *Schult. Mant. i. Add.* t. 195.—Bras.
Candolleana, St. Hil. in Bull. Soc. Philom. (1822) 132 = Cusparia Candolleana.
corymbosa, Spreng. Syst. iv. Cur. Post. 91 = Cusparia febrifuga.
cuneifolia, St. Hil. ex DC. Prod. i. 731 = Cusparia cuneifolia.
Cusparia, St. Hil. ex l. c. = Cusparia febrifuga.
elegans, St. Hil. Fl. Bras. Mer. i. 90 = Cusparia elegans.
fissa, *Mig. Stirp. Surin. Sel.* 64.—Guian.
Fontaneziana, St. Hil. in Bull. Soc. Philom. (1822) 132 = Cusparia Fontanesiana.
Gaudichaudiana, St. Hil. Fl. Bras. Mer. i. 89 = Cusparia Gaudichaudiana.
grandifolia, *Engl. in Mart. Fl. Bras.* xii. ii. 98.—Bras.
heterophylla, St. Hil. in Bull. Soc. Philom. (1823) 131 = Cusparia heterophylla.
jasminiflora, Engl. in Mart. Fl. Bras. xii. ii. 97 = multiflora.
Laxiostemon, St. Hil. ex DC. Prod. i. 731 = Cusparia silvestris.
laxiflora, *Engl. in Mart. Fl. Bras.* xii. ii. 99.—Bras.
longifolia, *St. Hil. in Bull. Soc. Philom.* (1823) 131.—Bras.
macrophylla, St. Hil. in Mém. Mus. Par. x. (1823) 282 = Cusparia macrophylla.
Martiana, St. Hil. Fl. Bras. Mer. i. 90 = Cusparia Martiana.
Mikaniana, *St. Hil. in Bull. Soc. Philom.* (1823) 131.—Bras.
multiflora, *Schult. Mant.* i. *Add.* t. 194.—Bras.
odoratissima, Lindl. Bot. Reg. t. 1490 = Cusparia odoratissima.
officinalis, *Hancock, in Trans. Med. Bot. Soc. Lond.* i. (1839) 25.—Am. austr.
oppositifolia, *Spruce, ex Engl. in Mart. Fl. Bras.* xii. ii. 105.—Bras.
Ozaena, DC. in Mém. Mus. Par. ix. (1822) 149. t. 10 = Raputia Ozaena.
ovata, St. Hil. & Tul. in Ann. Sc. Nat. Sér. II. xvii. (1842) 139 = Cusparia ovata.
pentagyna, St. Hil. in Mém. Mus. Par. x. (1823) 281 = Cusparia pentagyna.
pentandra, St. Hil. in Bull. Soc. Philom. (1823) 132 = Cusparia pentandra.
pentaphylla, *Macfad.* Fl. Jamaic. i. 196 = Esenbeckia pentaphylla.
ramiflora, Spruce, ex Engl. in Mart. Fl. Bras. xii. ii. 112 = Cusparia ramiflora.
resinosa, St. Hil. ex DC. Prod. i. 731 = Rania resinosa.
Riedeliana, *Regel, Gartenfl.* (1866) 355.—Bras.
ruiziana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 306.—Bras.
simplicifolia, *Schult. Mant. i. Add.* i. 196.—Bras.
toxicaria, Spruce, ex Engl. in Mart. Fl. Bras. xii. ii. 116 = Cusparia toxicaria.
trifoliata, *Aubl. Pl. Gui.* ii. 662. t. 269.—Guian.

GALISONGEA, Willd. Sp. Pl. iii. 2226 (1800) = **Galinsoga**, Ruiz & Pav. (Compos.).

GALIUM, Linn. Gen. ed. I. 24 (1737). *RUBIACEÆ*, Benth. & Hook. f. ii. 149.
Aparine, [Tourn.] Linn. Gen. ed. I. 24 (1737).
Asperea, Moench, Meth. 641 (1794).
Cruciata, Tourn. ex Adans. Fam. ii. 144 (1763).
Eysellia, Neck. Elem. i. 199 (1790).
Mesopeplus, Schrenk, in *Bull. Sc. Acad. Péterab.* xi. (1844) 115.
Rubiola, Tourn. ex Adans. Fam. ii. 144 (1763).
Trichogalium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 396.
abruptorum, *Pomel, Nouv. Mat. Fl. Atl.* 76.—Alger.
acuphyllum, Willk. & Cost. in Linnæa, xxx. (1859 60) 101 = fruticescens.
aculeatissimum, *Kit. ex Kanitz, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 648.—Hungar.
acuminatum, *Ball, in Journ. Bot.* xi (1873) 334.—Marocco.
acutissimum, A. Gray, in Proc. Am. Acad. vii. (1868) 350 = stellatum.
acutum, *Edgew. in Trans. Linn. Soc.* xx. (1846) 61.—Reg. Himal.
adnexens, *Boiss. & Bal. Diagn.* Ser. II. v. 106.—Cilic.; Syria.
adnexens, Gillb. Fl. Lituan. i. 13 = G. Aparine.
adnexens, Hort. Vindob. ex DC. Prod. iv. 608 = spurium.
aetnense, Presl, Delic. Prag. 120 = seq.
aetnicum, Biv.on, Stirp. Rar. Sicil. iv. 21 = corrudæfolium.
affine, Roth, Enum. Pl. Phæn. Germ. i. 1. 474 = verum!
affine, Schult. Mant. iii. 172 = leiddum.
aggregatum, Polr. Encyc. Suppl. ii. 690 = glomeratum.
agreste, Wallr. Sched. Crit. 50 = G. Aparine.
alboroseum, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 462 [bis].—Tasmen.
albo-lutescens, Gillb. Fl. Lituan. i. 8 = uliginosum.
album, Forsk. Fl. Aegypt. Arab. p. xx = erectum *vel* G. Mollugo.
album, Mill. Gard. Dict. ed. VIII. n. 7 = G. Mollugo.
album, Willd. Sp. Pl. i. 598 = brevifolium, *Sibth. & Sm.*
alepicum, Boiss. Diagn. Ser. I. iii. 50 = nigricans.
alpestre, Gaud. ex Roem. & Schult. Syst. iii. 225 = anisophyllum.
alpestre, Stev. ex Fisch. Cat. Hort. Gorenk. (1812) 48 = brachyphyllum.
alpicola, Jord. Obs. Pl. Crit. iii. (1846) 131 = obliquum.
alpicola, *Schrebr, in Linnæa*, xvii. (1843) 541.—Helvet.
alpinum, Schur, Enum. Pl. Transs. 288 = vernum.
altissimum, Gillb. Fl. Lituan. i. 11 = Asperula galioides.
Amani, Kotschy, ex Boiss. Fl. Orient. iii. 54 = erectum.
amathymbolicum, *Eckl. & Zeyh. Enum.* 370.—Afr. austr.
ambiguum x, Gren. & Godr. Fl. Fr. ii. 20.—Gallia.
ambiyophyllum, *Schrenk, Enum. Pl. Nov.* 96.—Sibir. alt.
andinum, *Phil. in Linnæa*, xxxiii. (1864–65) 98.—Chili.
Andrewsii, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 587.—Am. bor. occ.
anglicum, Huds. Fl. Angl. ed. II. 69 = parisiense.
angulosum, *A. Gray, in Proc. Am. Acad.* xi. (1876) 74.—Calif.
angustifolium, *Nutt. ex Torr. & Gray, Fl. N. Am.* ii. 22.—Calif.
anisophyllon, *Vill. Prosp.* 20; *Hist. Pl. Dauph.* ii. 317 [bis].—Europ.
anomalum, Hort. ex Reichb. f. Ic. Fl. Germ. xvii. 89 = Asperula mollugnoides.
antarcticum, *Hook. f. Fl. Antarct.* 308 [bis].—Reg. Magell.
Aparine, Aitch. in Journ. Linn. Soc. xix. (1882) 167 = tricorne.
Aparine, *Linn. Sp. Pl.* 108.—Reg. bor. temp. et Magell.
Aparine, Wall. Cat. n. 6218 = hirtiflorum, G. Mollugo.
aparinoides, Lange, Ind. Sem. Hort. Haun. (1854) 24 = spurium.
aparinoides, *Forsk. Fl. Aegypt. Arab.* 30.—Arab.
aparinoides, C. Koch, in Linnæa, xvii. (1843) 38 = spurium.

GALIUM:—
aphyllum, [*Pieri*!] *in Ionios Anthol.* iv. 940.—Ins. Corcyra.
apiculatum, Roem. & Schult. Syst. iii. 222 = leucocarpum.
apiculatum, *Sibth. & Sm. Fl. Græc. Prod.* i. 91.—Græcia.
apiculatum, Sprun. ex Nym. Consp. 328 = aureum.
approximatum x, Gren. & Godr. Fl. Fr. ii. 20.—Gallia.
apricum, *Sibth. & Sm. Fl. Græc. Prod.* i. 90.—Archip. Græc.
apricum, Sieber, ex Stend. Nom. ed. II. i. 657 = Asperula humifusa.
apricum, Vell. Fl. Flum. 43: t. t. 111 = radicosum.
aquaticum, Willd. ex Schult. Mant. iii. 158 = uliginosum.
arenarium, *Loisel. Fl. Gall.* 85; ed. II. i. 110.—Gallia.
arcticides, *Boiss. Diagn.* Ser. I. iii. 42.—Ass. Min.
argenteum, Reichb. ex Nym. Consp. 327 = anisophyllum.
argenteum, Vill. Prosp. 20; Hist. Pl. Dauph. ii. 318 [bis] = sylvestre.
aristatum, *Linn. Sp. Pl.* ed. II. 152.—Europ.
aristatum, Sm. Eng. Fl. i. 208 = erectum.
arkansanum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 86.—Am. bor. occ.
articulatum, Lam. Tabl. Encyc. i. 260 = rubioides.
articulatum, *Roem. & Schult. Syst.* iii. 250.—Oriens.
ascendens, Willd. ex Spreng. Syst. i. 390 = mexicanum.
Aschersonii, *S. Schwein. in Linnæa*, xx. (1847) 701.—Mexic.
asparagifolium, Boiss. Diagn. Ser. II. vi. 91 = aureum.
asparagifolium, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 327 = ochroleucum.
asperifolium, Wall. in Roxb. Fl. Ind. ed. Carey, i. 881 = G. Mollugo.
aspertimum, *A. Gray, Pl. Fendl.* 60.—Am. bor. occ.
asperuloides, Edgew. in Trans. Linn. Soc. xx. (1846) 61 = trifloriun.
asperum, [Horsk.] Verz. Aller Gew. Teutschl. 423 = G. Aparine!
asperum, Schreb. Spicil. Fl. Lips. 8 = sylvestre.
asperum, *Thunb. Prod. Pl. Cap.* 30.—Afr. austr.
asprellum, *Michx. Fl. Bor. Am.* i. 78.—Am. bor.
atherodes, *Spreng. Syst.* iv. *Cur. Post.* 39.—Bras.
atlanticum, Ball, in Journ. Bot. xi. (1873) 334 = sylvestre.
atlanticum, *Pomel, Nouv. Mat. Fl. Atl.* 76.—Alger.
atrovirens, Lapeyr. Suppl. Hist. Fl. Pyr. 22 = sylvaticum.
Aucheri, *Boiss. Diagn.* Ser. I. iii. 49.—Persia.
aureum, *Vis. in Orto Bot. Padov.* (1842) 194, in not.; *Fl. Dalmat.* iii. 6.—Europ. ox.
aurienae, Fourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 310 = boreale.
australe, *DC. Prod.* iv. 608.—Austral.
austriacum, Jacq. Fl. Austr. i. 51 = sylvestre.
asyriflorum, F. Müell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 113 = Gaudichaudi.
axillare, Presl, ex Gum. Fl. Sic. Prod. i. 178 = parisiense.
baldense, Spreng. Pugill. i. 10 = helveticum.
Barrelieri, Salzm. in Flora, iv. (1821) 107 = ellipticum.
Bauhini, Roem. & Schult. Syst. iii. 218 = vernum.
bellanense, Kotschy, ex Boiss. Fl. Orient. iii. 54 = erectum.
Benthamianum, Walp. Rep. vi. 17 = Relbunium tuberculosum.
bericum, Terra, in Giorn. Ital. Venez. i. (1765) 119 = aristatum.
bermudense, Linn. Sp. Pl. 105 = pilosum.
bermudianum, Linn. Syst. ed. XII. 117 = seq.
Bernardi, *Gren. & Godr. Fl. Fr.* ii. 23.—Corsica.
Biafræ, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 245.—Afr. trop.
bicolor, Schleich. Cat. Fl. Helv. ed. IV. 17 = sylvestre.
Bielzii, Schur, Enum. Pl. Transs. 285 = erectum.
biflorum, Hochst. ex Boiss. Fl. Orient. iii. 69 = spurium.
biflorum, *S. Wats. in Bot. King's Exped.* 184. t. 14.—Am. bor. occ.
bigeminum, *Griseb. in Goett. Abh.* xix. (1874) 160.—Reg. Argent.
blepharophorum, Roem. & Schult. Syst. iii. 249 = Valantia hispida.

GALIUM:—
Bloomeri, A. Gray, in Proc. Am. Acad. vi. (1864–5) 538 = multiflorum.
Boccosi, All. Fl. Pedem. i. 6 = anisophyllum.
Bocconi, DC. Prod. iv. 594 = sylvestre.
Bocconi, J. F. Gmel. Syst. 238 = anisophyllum?
Boccosi, Tenore, ex Steud. Nom. ed. II. i. 657 = erectum.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 350.—Calif.
borrale, Lapeyr. ex DC. Prod. iv. 596 = G. Mollugo, boreale, *Linn. Sp. Pl.* 108.—Reg. bor. temp.
borreli, Walt. Fl. Carol. 85 = circaezans.
Boryanum, *Waldst. Kop.* ii. 454.—Graecia.
Bourgeaeanum, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 60, nomen.—Afr. bor.
Bovei, *Boiss. & Kral. Pugill. Pl. Nov.* 50.—Algeria.
brachiatum, Muhl. Cat. 1. 15 = circaezans.
brachiatum, Pursh. Fl. Am. Sept. i. 103 = triflorum.
brachycarpon, Stend. Nom. ed. II. i. 657, sphalm. vide *trachycarpum*.
brachyphyllum, *Schult. Mant.* iii. 180.—Reg. Caucas.
brachypodum, Jord. Obs. Pl. Crit. iii. (1846) 180 = obliquum.
brachypodum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 280.—Manchuria.
bractеatum, *Boiss. Diagn.* Ser. I. iii. 51.—Mesopot.
Brandegei, *A. Gray, in Proc. Am. Acad.* xii. (1877) 58.—N. Mexic.
brasiliense, *Vell. Fl. Flum.* 43 ; t. t. 109.—Bras.
brasiliense, *Wernm, in Oestr. Bot. Zeitschr.* xxix. (1879) 216.—Bras.
Brittoni, Le Jolis, ex Nym. Consp. 328 = verum.
brevifolium, *Phil. in Linnaea*, xxxiii. (1864–65) 98.—Chili.
brevifolium, *Sibth. & Sm. Fl. Graec. Prod.* i. 95.—As. Min.
brevifolium, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 253 = brachyphyllum.
Broteranum, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 15.—Hispan.
brunneum, *Munby, Fl. Algér.* 16.—Algeria.
bryophyllum, Goldb. ex DC. Prod. iv. 602 = triflorum.
Bungei, Steud. Nom. ed. II. i. 657 = gracile.
cabrerense, Auct. vide cespitosum.
californicum, *Hook. & Arn. Bot. Beech. Voy.* 349. —Calif.
caninianum, Schult. Mant. iii. 186 = recurvum.
campestatum, Vill. Prosp. 19 ; Hist. Pl. Dauph. ii. 326 = Asperula galioides.
campestre, Duby, Bot. Gall. ed. II. i. 248 = Soleirolii.
campestre, Hochst. ex Boiss. Fl. Orient. iii. 89 = spurium.
campestre, Schousb. ex Willd. Enum. Hort. Berol. 152 = glomeratum.
camporum, Pohl, ex DC. Prod. iv. 604 (Relbunii sp.).—Bras.
canescens, *H. B. & K. Nov. Gen. et Sp.* iii. 386.—Ecuador.
canum, *Req. ex DC. Prod.* iv. 602.—Syria.
capense, *Thunb. Prod. Pl. Cap.* 30.—Afr. austr.
capillare, Cav. Ic. ii. 73. t. 191 ; Sibth. & Sm. Fl. Graec. Prod. i. 94 = setaceum.
capillare, Decne. in Ann. Sc. Nat. Sér. II. ii. (1884) 269 = Decaisnei.
capillipes, Reichb. Fl. Germ. Excurs. 847 = silvaticum.
capitatum, *Bory & Chaub. Nouv. Fl. Pélop.* 10.—Graecia.
cappadocicum, *Boiss. Diagn.* Ser. I. iii. 48.—Cappadoc. ; Syria.
capsiricense, *Ysamb. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* vi. (1858–84) 155.—Europ.
caripense, *H. B. & K. Nov. Gen. et Sp.* iii. 387.—Venezuela.
carium, *Stapf, in Denkschr. Acad. Wien*, i. (1885) 107.—As. Min.
carolinianum, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 429 = hispidulum.
caspicum, *Stev. in Nouv. Mém. Soc. Nat. Mosc.* i. (1829) 276.—Persia.
cassium, *Boiss. Diagn.* Ser. I. x. 66.—As. Min. ; Syria.
caucasicum, Lag. Gen. et Sp. Nov. 9 = verum.
caudatum, Boiss. Diagn. Ser. I. iii. 49 = brevifolium, *Sibth. & Sm.*
cerastocarpon, *Boiss. l. c.* 35.—Cappadoc.
cerastophyllioides, *Hook. f. Fl. Brit. Ind.* iii. 209.—Reg. Himal.

GALIUM:—
ceratopodum, *Boiss. Diagn.* Ser. I. iii. 45.—Arab. ; Assyr.
cespitosum, Lam. Tabl. Encyc. i. 363 = pusillum.
cespitosum, *Ram. in Mém. Mus. Par.* xiii. (1825) 266. —Mont. Fyren.
chaetophorum, *Griseb. in Goett. Abh.* xix. (1874) 160. —Reg. Argent.
chamaeparines, Willk. & Cost. in Linnaea, xxx. (1859–60) 102 = parisiense.
Chamissonis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 368.—Chili.
cherovense, Roem. & Schult. Syst. iii. 250 = *G. Cruciata*.
chilense, Hook. f. Fl. Antarct. 302 [bis] = chonoense.
chinense, *Spreng. f. Tent. Suppl.* 7.—Mongol.
chloranthum, Griseb. Fl. Insul. i. 149 = pedemontanum.
chloroleucum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 27.—Persia.
chlorophyllum, Baill. & Timb. in Mém. Acad. Tool. Sér. V. vi. (1862) 235 = sylvestre.
chonoense, *Hook. f. Fl. Antarct.* 303 [bis].—Chili.
ciliare, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 461 [bis].—Tasman.
ciliatum, D. Don, Prod. Fl. Nep. 138 = hirtiflorum.
ciliatum, Fourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 310 = ellipticum.
ciliatum, Ruiz & Pav. Fl. Per. i. 59 = Relbunium ciliatum.
ciliatum, Schrank, Prim. Fl. Salisb. 51 = sylvestre.
ciliiferum, *Boiss. Diagn.* Ser. II. v. 105.—Cilicia.
cinereum, *All. Fl. Pedem.* i. 6. t. 77.—Europ. austr.
circaezoides, Roem. & Schult. Syst. iii. 256 = circaezans.
circaezans, *Michx. Fl. Bor. Am.* i. 80.—Am. bor.
circaceum, Boiss. Diagn. Ser. I. vi. 69 = aureum.
Clausonis, *Pomel, Nouv. Mat. Fl. Alt.* 77.—Alger.
Cleyeri, Michx. Fl. Bor. Am. i. 78 = tridium.
Clusianum, Timb. in Bull. Soc. Bot. Fr. ix. (1862) 613 = Jordani.
collinum, Jord. Obs. Pl. Crit. iii. (1846) 185 = Jordani.
Colmeiroi, *Ehrenb. ex Boiss. Fl. Orient.* iii. 81.—Aegypt.
cometerhizon, Lapeyr. Suppl. Hist. Pl. Pyr. 154 = megalospermum.
commutatum, Jord. Obs. Pl. Crit. iii. (1846) 149 = sylvestre.
concatenatum, *Coss. Notes Pl. Crit.* 38.—Fort. Hercul.
concinnum, Boiss. Fl. Orient. iii. 65 = G. Lycium.
concinnum, *Torr. & Gray, Fl. N. Am.* ii. 28.—Am. bor.
confertum, *Royle, ex Hook. f Fl. Brit. Ind.* iii. 208.—Reg. Himal.
congestum, Jord. Obs. Pl. Crit. iii. (1846) 172 = constrictum.
conglobatum, Schousb. ex Bail, in Journ. Linn. Soc. xvi. (1878) 486 = concatenatum.
conzanguineum, Boiss. Diagn. Ser. I. vi. 69 = verum.
constrictum, *Chaub. in St. An. Fragm. Voy. Pyrén.* 208 ; *Fl. Agen.* 67.—Europ.
cordatum, Koem. & Schult. Syst. iii. 259 = articulatum.
cornaceum, *Bunge, in Ledeb. Fl. Alt.* i. 186.—Sibir. Altaic.
cornigerum, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 67.—As. Min.
coronatum, *Sibth. & Sm. Fl. Graec. Prod.* i. 96.—Graecia ; Oriens.
corrudaefolium, C. Koch, in Linnaea, xxiv. (1851) 466 = aureum.
corrudaefolium, *Vill. Prosp.* 20 ; *Hist. Pl. Dauph.* ii. 520.—Europ. austr.
corrudaefolium, Willk. in Flora, xxxv. (1852) 640 = erectum.
corsicum, Spreng. Syst. iv. Cur. Post. 39 = Soleirolii.
corsicum, *Teusch, in Flora*, xiv. (1831) 221.—Corsica.
corymbosum, Ruiz & Pav. Fl. Per. i. 59 = Rubia corymbosa.
cotinoides, *Chem. & Schlecht. in Linnaea*, iii. (1828) 227.—Chili.
Crespianum, *Rodr. in Anal. Soc. Esp. Hist. Nat.* viii. (1879) 55.—Ins. Balear.
creticum, Boiss. & Heldr. Diagn. Ser. I. x. 66 = constrictum.
crispulo-pilosum, C. Koch, ex Boiss. Fl. Orient. iii. 57 = orientale.
cristatum, Jaub. & Spach, Illustr. ii. 115 = Mariscarpaea vaillantioides.

GALIUM:—
crocеum, Ruiz & Pav. Fl. Per. i. 59 = Rubia crocea.
Cruciata, *Scop. Fl. Carn.* ed. II. i. 100.—Europ. ; As. bor.
cruciatum, Sm. Angl. Bot. t. 148 = praec.
cryptanthum, *Hemsl. in Hook. Ic. Pl.* t. 1469.—Reg. Himal.
Cucullaria, Roem. & Schult. Syst. iii. 259 = Callipeltis Cucullaria.
curtum, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 463 [bis].—Tasman.
cuspidatum, Muhl. Cat. ed. I (1813), 16 ; ed. II. 15 ; ex Ell. Sketch, i. 107 = triflorum.
cuspidatum, *Raffin. in Med. Repos. N. York*, v. (1808) 359, nomen.—Am. bor.
cuspidulatum, *Miq. Fl. Ind. Bat.* ii. 840.—Java.
cyllenium, *Boiss. & Heldr. Diagn.* Ser. II. ii. 117.—Graecia.
cynanchicum, Soop, Fl. Carn. ed. II. i. 101 = Asperula cynanchica.
daharicum, Turcz. in Bull. Soc. Nat. Mosc. xviii. (1845) i. 319 = asprellum.
dasycarpon, *Kunze, ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 857, nomen.—Peruv.
dasycarpum, *Hochst. in Schweinf. Beitr. Fl. Aethiop.* 135.—Afr. trop.
dasycarpum, *Nees, in Nenwied, Reise Nord-Am.* ii. 437.—Am. bor. occ.
debile, Banks & Soland. ex Hook. f. Fl. Antarct. 303 [bis] = asterocarpum.
debile, Desv. Obs. Pl. Angers, 134 = constrictum.
debile, *Hoffegg. & Link, Fl. Port.* ii. 48.—Lusitan.
Decaisnei, *Boiss. Diagn.* Ser. I. vi. 69.—Syria ; Persia.
decipiens, Ehrh. Beitr. v. 177 = rotundifolium.
decipiens, Jord. Obs. Pl. Crit. iii. (1846) 178 = parisiense.
decipiens, Roth, Enum. Pl. Phaen. Germ. i. t. 478 = sylvestre.
decolorans, Gren. & Godr. Fl. Fr. ii. 19.—Europ. delicatulum, *Boiss. & Hohen. Diagn.* Ser. I. x. 66.—Persia.
demissum, *Boiss. l. c.* iii. 40.—Graecia.
densiflorum, *Ledeb. Fl. Alt.* i. 187.—Sibir. altaic.
densum, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 461 [bis] = australe.
denticulatum, *Harvl. ex DC. Prod.* iv. 612.—Mexic.
dichotomum, *Linn. Ind. Sem. Hort. Hamb.* (1833) 7. —Hab.?
diffusum, *Cav, in C. Gay, Fl. Chil.* iii. 180.—Chili.
diffusum, D. Don, in Hook. Fl. Scot. i. 53 = erectum.
diffusum, Gilib. Fl. Lituan. i. 7 = palustre.
diffusum, Schrad. ex Link, Enum. Hort. Berol. i. 134 = boreale.
digenerum x, Kern. in Oestr. Bot. Zeitschr. xxvi. (1876) 118.—Europ.
diplopriun, *Boiss. & Hohen. Diagn.* Ser. I. x. 66.—Persia.
divaricatum, *Lam. Encyc.* ii. 580.—Reg. Mediterr.
divaricatum, Pall. ex Ledeb. Fl. Ross. ii. 416 = asperuloides.
dumetorum, Jord. Pugill. Pl. Nov. 78 = G. Mollugo.
dumosum, *Boiss. Diagn.* Ser. I. iii. 44.—As. Min.
Dregeanum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 38. —Afr. austr.
effusum, Boiss. Diagn. Ser. I. iii. 35 = longifolium.
Ehrenbergii, *Boiss. Fl. Orient.* iii. 52.—Syria.
elatum, J. F. Gmel. Syst. 235 = sylvestre.
elatum, Thuill. Fl. Par. ed. II. 76 = G. Mollugo.
elegans, Wall. in Roxb. Fl. Ind. ed. Carey, i. 382 = rotundifolium.
ellipticum, *Willd. Enum. Hort. Berol. Suppl.* 9, nomen ; *Harnem. Suppl. Hort. Hafn.* 17.—Reg. Mediterr. occ.
elongatum, *Presl, Delic. Prag.* 119.—Europ. ; Afr. bor. ; Oriens.
eminens x, Gren. & Godr. Fl. de Fr. ii. 19.—Europ.
ephedroides, *Willd. in Linnaea*, xxv. (1852) 80.—Hispan. ; Afr. bor.
erectum, Don, ex Schult. Mant. iii. 179 = aristatum.
erectum, Hoffm. Deutschl. Fl. ed. II. i. t. 73 = sylvestre.
erectum, *Huds. Fl. Angl.* ed. I. 56.—Europ. ; Afr. bor. eriocidos, *Lam. Encyc.* ii. 583.—Am. austr.
eriocarpum, *Bartl. ex DC. Prod.* iv. 600.—Chili.
eriocephalum, Chaix, ex Schult. Mant. iii. 186 = setaceum.
erythrosanlon, *Celenia, in Transa. N. Z. Inst.* xvi. (1884) 382.—N. Zel.
erythrorhizon, *Boiss. & Reut. Pugill. Pl. Nov.* 51.—Hispan.

GALIUM :—
erythrorhizum, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 118 = australe.
exile, *Hook. f. Fl. Brit. Ind.* iii. 207.—Reg. Himal.
expansum, Thunb. Prod. Pl. Cap. 30 = capense.
Fendleri, *A. Gray, Pl. Fendl.* 60.—N. Mexic.
filicaule, *Willd. ex Schult. Mant.* iii. 172.—Carolina.
filiforme, *Roem. & Schult. Syst.* iii. 252.—Ins. Teneriffa.
filiformе, Willd. ex Ledeb. Fl. Ross. ii. 418 = tenuissimum.
firmum, Tausch, in Flora, xiv. (1831) 222 = aureum.
flaccidum, Salisb. Prod. 58 = G. Mollugo.
flavescens, *Borb. in Ahnd. Kösl.* xl. 7 (1874) 266; *ex Kern. Sched.* iii. 106.—Transsylv.
flavum, Wallr. Beitr. Fl. Hercyn. 178; et in Linnaea, *x* xiv. (1840) 548 = *schroderianum*, Wolf.
fleurotii, Jord. Grain. Jard. Grenoble, (1849) 15 = Jordani.
floribundum, Bory & Chaub. Nouv. Fl. Pélop. 10 = aureum.
floribundum, Jacquem. ex Hook. f. Fl. Brit. Ind. iii. 206 = acutum.
floribundum, *Sibth. & Sm. Fl. Graec. Prod.* i. 94.—Graec.; As. Min.
floribundum, Urv. in Mém. Soc. Linn. Par. i. (1822) 275 = setaceum.
floridum, Salisb. Prod. 58 = verum.
fluminense, *Vell. Fl. Flum.* 48; i. t. 110.—Bras.
foliosum, *Mumby, ex Burnat & Barbey, Voy. Bot. Baléares*, 28.—Alger.
Fontanesianum, *Pomel, Nouv. Mat. Fl. Atl.* 75.—Alger.
fontinale, C. Koch, in Linnaea, xxiv. (1851) 469 = palustre.
fragile, *Krock, Fl. Siles. Suppl.* i. 210.—Europ.
fragile, Pourr. Chlor. Hisp. 512; ex DC. Prod. iv 610 = mutale.
fruticescens, *Cav. Ic.* iii. 8. t. 206.—Hispan.
fruticosum, Sieber, ex Boiss. Fl. Orient. iii. 60 = constrictum.
fruticosum, *Willd. Sp. Pl.* i. 585.—Ins. Creta.
fuegianum, *Hook. f. Fl. Antarct.* 302 [bis].—Reg. Magellan.
fuscum, *Mart. & Gal. in Bull. Acad. Brux* xi. (1844) 126.—Mexic.
Galeottianum, *Walp. Rep.* vi. 17.—Mexic.
Gardneri, *Walp. l. c.*—Bras.
garipense, *Sond. in Harv. & Sond. Fl. Cap. Cap.* iii. 37.—Afr. austr.
Gaudichaudii, *DC. Prod.* iv. 607.—Austral.
geminiflorum, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimpr. 19.—Ins. Madera.
geminiflorum, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 126.—Mexic.
geminiflolium, *F. Muell. in Trans. Vict. Inst.* (1855) 147.—Austral.
geniculatum, Roem. & Schult. Syst. iii. 215 = rubioides.
Gerardi, Vill. Prosp. 19 = erectum.
ghilanicum, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 53.—Persia.
gibraltaricum, *Schott, in Oken, Isis*, xxi. (1828) 821.—Gibraltar.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 264.—Chili.
glaberrimum, *Hemsl. Diagn. Pl. Nov.* 53.—Mexic.
glabrum, Hoffm. Deutschl. Fl. ed. II. i. i. 72 = sylvestre.
glabrum, Roehl. ex Stend. Nom. ed. I. 858 = verum.
glabrum, *Thunb. Prod. Pl. Cap.* 30.—Afr. austr.
glaucum, Linn. Sp. Pl. 107 = Asperula galloides.
glaucum, Ced. in Fl. Dan. t. 609 = sylvaticum.
glaucum, Suter, Fl. Helv. i. 91 = aristatum.
glomeratum, *Desf. Fl. Atlant.* i. 128. t. 40.—Reg. Mediterr. occ.
glomeratum, Vill. ex Roem. & Schult. Syst. iii. 216 = palustre.
Gmelini, [Honck.] Verz. Aller Gew. Teutschl. 425 (Gmelin).—Europ.
Goldbachianum, Steud. Nom. ed. II. i. 658, sphalm. = podiachicum.
Goldbachicum, DC. Prod. iv. 610, sphalm. = podiachicum.
gracile, *Bunge, Enum. Pl. Chin. Bor.* 35.—China.
gracile, Wallr. Sched. Crit. 57 = parisiense.
gracilescens, Jord. ex Billot, Annot. 318 = Jordani.
graecum, *Linn. Mant.* i. 38.—Graec.; As. Min.
grandiense, *Willd. ex Schult. Mant.* iii. 198.—Hispan.
grandiflorum, [Clairv.] Man. Herb. 41 = Asperula galloides.

GALIUM:—
grandiflorum, Ehrenb. ex Boiss. Fl. Orient. iii. 52 = Pestalozzae.
grandiflorum, Gillh. Fl. Lituan. i. 7 = uliginosum.
granulatum, Roem. & Schult. Syst. iii. 249 = saccharatum.
gratum, Wall. Cat. n. 6211 = trifforum.
grusinum, *Trautv. in Act. Hort. Petrop.* vii. (1880) 461.—Reg. Caucas.
Haenkeanum, *Endl. ex Walp. Rep.* ii. 459.—Chili.
Halleri, Roem. & Schult. Syst. iii. 218 = vernum.
Halleri, Suter, Fl. Helv. i. 88 = Asperula galioides.
hamatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 345.—Abyss.
Hamiltonii, Spreng. Syst. iv. Cur. Post. 39 = rotundifolium.
Hanryi, F. Schultz, Arch. Fl. 309 = Jordani.
harcynicum, Weigel, Obs. 25, sphalm. vide = *hercynicum*.
helderi, Hoffmgg. & Link, Fl. Port. ii. 47 = uliginosum.
helveticum, *Weigel, Obs.* 24.—Europ.
hercynicum, Weigel, l. c. 25 = saxatile.
hierosolymitanum, *Lunn. Amoen. Acad.* iv. 451.—Syria.
hierosolymitanum, Thore, Chlor. Land. 40 = arenarium.
himalayense, Klotzsch, in Bot. Ergeb. Waldem. Reise, 88 = acutum.
hirsutum, Gilib. Fl. Lituan. i. 10 = uliginosum.
hirsutum, Lapeyr. Suppl. Hist. Pl. Pyr. 25 = papillosum.
hirsutum, Nees, Hor. Phys. Berol. 118. t. 22 = Necsianum.
hirsutum, *Ruiz & Pav. Fl. Per.* i. 50.—Peruv.
hirtiflorum, *Req. ex DC. Prod.* iv. 600.—Reg. Himal.
hirtum, Hegetschw. Fl. Schw. 125 = rubrum.
hirtum, *Lam. Encyc.* ii. 583.—Bras.
hispidulum, *Michx. Fl. Bor. Am.* i. 70.—Am. bor.
hispidum, Gaertn. Fruct. i. 109. t. 24. f. 1 = Valantia hispida.
hispidum, Pourr. ex Nym. Consp. 330 = parisiense.
hispidum, Pursh, Fl. Am. Sept. i. 104 = hispidulum.
hispidum, Roth, Tent. Fl. Germ. ii. 134 = spurium.
hispidum, Schrad. Spicil. Fl. Germ. 14 = sylvestre.
hispidum, Willd. Enum. Hort. Berol. 154 = G. Aparine.
Hoffmanni, Fl. Necro. Danb. ex Steud. Nom. ed. II. i. 658 = sylvestre.
horridum, Eckl. & Zeyh. Enum. 370 = G. Aparine.
horridum, *Thunb. in Hoffmgg. Phytog. Blaetter*, i. (1808) 16.—Afr. austr.
humifusum, Beb. Fl. Taur. Cauc. i. 104 = Asperula humifusa.
humile, *Chem. & Schlecht. in Linnaea*, iii. (1828) 226. —Bras.
hungaricum ×, Kern. in Oestr. Bot. Zeitschr. xxvi. (1876) 119.—Europ.
Ilutert ×, Kern. l. c.—Europ.
hypadenium, *S. Schau. in Linnaea*, xx. (1847) 700.—Mexic.
hypnoides, *Cles, in C. Gay, Fl. Chil.* iii. 188.—Chili.
hypnoides, *Vill. Hist. Pl. Dauph.* ii. 328.—Europ.
hypotrichium, A. Gray, in Proc. Am. Acad. vi. (1864-65) 538 = multiflorum.
hyrcanicum, *C. A. Mey. Verz. Pfl. Cauc.* 58.—Reg. Caucas.
hyssopifolium, Hoffm. Deutschl. Fl. ed. II. i. i. 71 = boreale.
iberum, D. Dietr. Syn. Pl. i. 464 = G. Mollugo.
ibicinum, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 70.—Persia.
imbricatum, Jord. Obs. Pl. Crit. iii. (1846) 141 = Jordani.
incanum, Sibth. & Sm. Fl. Graec. Prod. i. 91 = orientale.
incarnatum, Gilib. Fl. Lituan. i. 8 = palustre.
inconspicuum, *Phil. in Linnaea*. xxxiii. (1864-65) 99. —Chili.
incurvum, Fleischer, ex Tausch, in Flora, xviii. (1835) 554 = tenuissimum.
incurvum, Sibth. & Sm. Fl. Graec. Prod. i. 92 = aureum.
indicum, *Heyne, in Wall. Cat.* n. 6214.—Ind. or.
infestum, Waldst. & Kit. Pl. Rar Hung. iii. 224 = spurium.
inflatum, Gilib. Fl. Lituan. i. 11 = sylvaticum.
innocuum, *Miq. Fl. Ind. Bat.* ii. 541.—Java.
insubricum, Gaud. Fl. Helv. i. 421 = G. Mollugo.
intercedens ×, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 296.—Europ.
intermedium, Eichw. Skizze, 158 = sylvaticum.
intermedium, Mérat, Fl. Par. ed. III. ii. 256 = G. Aparine.

GALIUM:—
internedium, *Schult. Obs. Bot.* 22.—Europ.
intertextum, Jord. Obs. Pl. Crit. iii. (1846) 142 = Jordani.
intricatum, Endl. ex Walp. Rep. ii. 459 = latoramosum.
intricatum, *Lowe, Man. Fl. Mad.* i. 398.—Ins. Cap. Verd.
intricatum, *Reut. in Mém. Soc. Phys. Genèv.* viii. (1839) 304.—Graecia.
involucratum, H. B. & K. Nov. Gen. et Sp. iii. 335 = Kelbuium ciliatum.
jalapense, *Schlecht. in Linnaea*, ix. (1834) 591.—Mexic.
javanicum, Blume, Bijdr. 948 = rotundifolium.
jenense, *Kotzsky, in Sitzb. Acad. Wien. Math -Nat Isl. Abth.* i. (1866) 257.—Arab.
jesoense, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 112.—Japon.
Jordani, *Loret & Barr. Fl. Montp.* 301.—Gallia.
judaicum, *Boiss. Diagn. Ser.* I. x. 69.—Syria.
juncivum, Heldr. ex Boiss. Fl. Orient. iii. 60 = constrictum.
junceum, Schreb. in Nov. Act. Nat. Cur. iii. (1767) 478 = gracceum.
juncivum, Sibth. & Sm. Fl. Graec. Prod. i. 91 = fruticosum.
jungermannioides, *Boiss. Diagn.* Ser. I. iii. 48.—Syria.
jungbuhnianum, *Miq. Fl. Ind. Bat.* ii. 540.—Java.
Fussiaer, Lapeyr. Hist. Abr. Pl. Pyr. 68 = sylvestre.
Fusseri, Vill. Prosp. 20; Hist. Pl. Dauph. ii. 328 = tenue.
kamtschaticum, *Steller, ex Schult. Mant.* iii. 186.—Kamtschat.; Am. bor.
khorasanense, *Grif. Notul.* iv. 262.—Afghan.
Kitaibelianum, Schult. Mant. iii. 198 = sylvaticum.
Kotschyanum, Boiss. Fl. Orient. iii. 51 = leiophyllum.
Kuetzingii, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 109 = verum.
kurdicum, *Boiss. & Hohen. Diagn.* Ser. I. iii. 87.—Persia.
laconicum, *Boiss. & Heldr. l. c.* vi. 66.—Graecia.
laetum, Jord. Obs. Pl. Crit. iii. (1846) 138 = obliquum.
laeve, Thuill. Fl. Par. ed. II. 77 = sylvestre.
laevigatum, Bartl. ex DC. Prod. iv. 591 = Relbunium laevigatum.
laevigatum, Linn. Sp. Pl. ed. II. 1667 = aristatum.
lancolatum, *Torr. & A. Gray, Fl. N. Am.* i. 388.—Am. bor.
languinosum, *Lam. Tabl. Encyc.* i. 361.—Am. bor.
Lapeyrousianum, Jord. Obs. Pl. Crit. iii. (1846) 154 = sylvestre.
lappaceum, *Ruiz & Pav. Fl. Per.* i. 59.—Peruv.
lappaceum, Salisb. Prod. 59 = G. Aparine.
laricocarpum, Boiss. Diagn. Ser. I. iii. 42 = orientale.
laricocarpum, *Phil. in Linnaea*, xxxiii. (1864-65) 99. —Chili.
latifolium, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 133 = rotundifolium.
latifolium, *Michx. Fl. Bor. Am.* i. 70.—Caroln.
latoramosum, *Cles, in C. Gay, Fl. Chil.* iii. 187.—Chili.
leiophyllum, *Boiss. & Hohen. Diagn.* Ser. I. iii. 86.—As. Min.; Persia.
leucocarpum, *DC. Prod.* iv. 612.—Peruv.
leucophaeum, Gren. & Godr. Fl. Fr. ii. 28 = obliquum.
leucotrichum, *Hemsl. Diagn. Pl. Nov.* 53.—Mexic.
Linchii, J. F. Gmel. Syst. 340 = rotundifolium.
lineariifolium, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) n. vii. 152.—China.
linifolium, Mill. Gard. Dict. ed. VIII. n. 8; Lam. Encyc. ii. 578 = aristatum.
litnifolium, Rochel, Fl. Banat. Rar. Ind. = sylvaticum ? lithospermifolium, *Fisch. ex Steud. Nom.* ed. II. i. 659, nomen.—Kossia.
litigiosum, DC. Fl. Fr. iv. 368 = parisiense.
litorale, *Guss. Fl. Sic. Prod.* i. 172.—Sicilia.
Littelii, Oakes, in Howey, Mag. Hortic. vii. (1841) 179 = kamtschaticum.
lividum, Jacquem. ex Hook. f. Fl. Brit. Ind. iii. 207 = G. Mollugo
longicaule, Rafin. Fl. Ludov. 75 = trifforum.
longiflorum, Gueldst. Reise, i. 427 = Asperula mollugiuoides.
longifolium, *Griseb. Spicil. Fl. Rumel.* ii. 157.—Bysant.; As. Min.
lucidum, All. Fl. Pedem. i. 5 = corrudaefolium.
lucidum, Bieb. Fl. Taur. Cauc. i. 104 = G. Mollugo.
lucidum, Koch, Syn. Fl. Germ. ed. I. 838 = erectum.

GALIUM :—
lutecruciatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = G. Cruciata.
luteolum, Jord. Obs. Pl. Brit. iii. (1846) 128 = obliquum.
luteovertic illatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = verum.
lutetum, Lam. Fl. Fr. iii. 381 = verum.
Lychnn. *Boiss Diagn.* Ser. I. x. 71.—As. Min.
magellanicum, *Hook. f. Fl. Antarct.* 302 [bis].—Reg Magell.
magellanus, Tenore, Relag. Viagg. Abruzz. 48 = pusillum.
Marchandi, Roem. & Schult. Syst. iii. 528 = papillosum.
margaricense, A. Gray, in Proc. Am. Acad. xiii. (1878) 371 = Bolanderi.
margaritaceum, *A. Kern. in Zeitschr. Ferdinand. Tirol* III. xv. (1870) 252.—Europ. austr.
maritimum, *Linn. Mant.* i. 38.—Europ. ; Orient.
maritimum, Thunb. Prod. Pl. Cap. 80 = tomentosum.
matrisylva, Wigg. Prim. Fl. Holsat. 13 = Asperula odorata.
Matthewii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 80.—Am. bor. occ.
maximum, Moris, Stirp. Sard. Elench. i. 55 = elongatum.
mediterraneum, DC. Prod. iv. 596 = Soleirolii.
megalanthum, Boiss. Diagn. Ser. I. vi 67 = leiophyllum.
megalospermum, *All. Fl. Pedem.* i. 9 ; *Vill. Hist. Pl. Dauph.* ii. 819 [bis].—Gallia.
megalospermum, DC. Fl. Fr. iv. 249, partim = arenarium.
megapotamicum, Spreng. Syst. iv. Cur. Post. 39 = hirtum.
melanantherum, Boiss. Diagn. Ser. I. vi. 68 = aureum.
mexicanum, *H. B. & K. Nov. Gen. et Sp.* iii. 337.—Mexic.
micranthum, Pursh, Fl. Am. Sept. i. 108 = asprellum ?
micranthum, Urv. in Mém. Soc. Linn. Par. i. (1822) 372 = recurvum.
microcarpon, Frivald. ex Griseb Spicil. Fl. Rumel. ii. 162 = intricatum.
microcarpum, Vahl, Symb. Bot. ii. 30 = setaceum.
microphyllum, *A. Gray, Pl. Wright.* i. 80.—Am. bor. occ.
micropermum, *Desf. Fl. Atlant.* i. 130.—Alger. ; Sardin.
miltorrhizum, Hance, in Journ. Bot. vi. (1868) 114 = gracile.
minimum, Gilib. Fl. Lituan. i. 10 = verum.
minimum, Koch, & Schult. Syst. iii. 262 = murale.
munzi, Gilib. Fl. Lituan. i. 9 = uliginosum.
minutiflorum, Brot. Fl. Lusit. i. 151 = parisiense.
minutulum, *Jord. Obs. Pl. Crit.* iii. (1846) 182.—Gallia.
minutum, *Linn. Sp. Pl.* 106.—Rossia.
mite, *Boiss. & Hohen. Diagn.* Ser. I. iii. 37.—As. Min.
molle, Gilib. Exerc. i. 92 = G. Mollugo.
Mollugo, *Linn. Sp. Pl.* 107.—Europ. ; As. temp.
Monachini, *Boiss. & Heldr. Diagn.* Ser. I. x. 67.—Graecia.
monanthos, Bartl. ex DC. Prod. iv. 591 = Rubia monantha.
montanum, Huds. Fl. Angl. ed. I. 56 = saxatile.
montanum, *Linn. Sp. Pl.* ed. II. 155.—Gallia.
montanum, Pollich, Hist. Pl. Palat. i. 154 = Asperula galioides.
montanum, Stend. Nom. ed. II. i. 659, sphalm. = vide *monanthos* !
montanum, Vill. Hist. Pl. Dauph. ii. 817 [bis] = sylvestre.
montanum, With. Bot. Arr. Brit. Pl. ed. III. ii. 187 = palustre.
monticolum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 36. —Afr. austr.
montium, J. F. Gmel. Syst. 327, sphalm. = *montanum*, Huds.
Morisii, Spreng. Syst. iv Cur. Post. 338 = rubrum.
mucronatum, Lam. Encyc. ii. 581 = obliquum, rubrum.
mucronatum, E. Mey. ex Harv. & Sond. Fl. Cap. iii. 37 = mucroniferum.
mucronatum, Kuiz & Pav. Fl. Per. i. 60 = leucocarpum.
mucronatum, Thunb. Prod. Pl. Cap. 30 = capense.
mucroniferum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 37. —Afr. austr.
multicaule, Schur, in Verh. Siebenb. Ver. Naturw. lv. (1854) 38 = anisophyllum.
multicaule, Wallr. Sched. Crit. 58 = sylvestre.
multiflorum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 97. f. 37.—Am. bor. occ.

GALIUM :—
Mungieri, Boiss. & Heldr. in Boiss. Fl. Orient. iii. 72 = divaricatum.
munale, *All. Fl. Pedem.* i. 8. t. 77. f. 1.—Reg. Mediterr.
murale, Bieb. Fl. Taur. Cauc. i. 105 = verticillatum.
muricum, *Boiss. & Reut. Diagn.* Ser. II. ii. 114.—Hispan.
muscoriforme, Boiss. Diagn. Ser. I. iii. 44 = canum.
muscoides, Lam. Encyc. ii. 580 = pyrenaicum.
mutabile, *Boss. Enum. Pl. Vols.* 42.—Ross. austr.
myrioneurum, Jord. Obs. Pl. Crit. iii. (1846) 126 = obliquum.
nanaquense, Eckl. & Zeyh. Enum. 369 = capense.
nebulosum, Boiss. Diagn. Ser. I. iii. 49 = floribundum.
Neesinum, *Reg ex DC. Prod.* iv. 606.—Ins. Teneriff.
negiectum, Le Gall, ex Gren. & Godr. Fl. Fr. ii. 22 = G. Mollugo.
nemorosum, Wierzsb. ex Nym. Consp. 324 = G. Mollugo.
nereosum, Lam. Fl. Fr. iii. 378 = boreale.
nevadense, Boiss. & Reut. Diagn. Ser. II. ii. 115 = sylvestre !
Niewerthi, *Franch. & Sav. Enum. Pl. Jap.* ii. 393.—Japon.
nigricans, *Boiss. Diagn.* Ser. I. iii. 48.—Syria ; Persia.
nigricans, *Clos, in C. Gay, Fl. Chil.* iii. 188.—Chili.
nitidulum, Thuill. Fl. Par. ed. II. 76 = sylvestre.
nitidum, Sieber, ex Stend. Nom. ed. II. i. 659 = corrudaefolium.
nitidum, Willd. ex Schult. Mant. iii. 168 = sylvaticum.
nolitangere, *Ball, in Journ. Bot.* xi. (1873) 335.—Maroc.
nouletianum, Bail. & Timb. in Mém. Acad. Toul. Sér. V. vi. (1862) 288 = papillosum.
nudiflorum, Viv. Fl. Cors. Prod. App ii. 3 = Soleirolii.
numidicum, *Pomel, Nouv. Mat. Fl. Atl.* 293.—Alger.
nutans, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 20 = virgatum.
Nuttalli, *A. Gray, Pl. Wright.* i. 80.—Calif.
obliquum, *Vill. Prosp.* 19 ; *Hist. Pl. Dauph.* ii. 320 [bis].—Gallia.
obliquum, Wolf. ex Nym. Consp. 329 = rubrum.
oblongifolium, Heldr. ex Nym. l. c. 328 = aureum.
obovatum, *H. B. & K. Nov. Gen. et Sp.* iii. 334.—Am. austr.
obovatum, Schlecht. & Cham. in Linnaea, v. (1830) 184 = jalapense.
obscurum, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = sylvestre.
obetipum, *Schlecht. in Linnaea*, lx. (1834) 592.—Mexic.
obtusiflorum, *Wallr. Beitr. Fl. Hercyn.* 178 ; *et in Linnaea*, xiv. (1840) 548.—Europ.
obtusum, Bigel. Fl. Bost. ed. II. 55 = trifidum.
ochroleucum, *Kit. ex Schult. Oestr. Fl.* ed. II. i. 805. —Hungar.
ochroleucum, Rochel, Pl. Banat. Rar. Ind. = verum.
ochroleucum x, Wolf. ex Schweigg. & Koert. Fl. Erlang. 86.—Europ.
odoratum, Scop. Fl. Carn. ed. II. i. 105 = Asperula odorata.
oligotrichum, *Borb. in Magyar Növénf. Lapok*, viii. (1884) 132.—Europ.
olympicum, *Boiss. Diagn.* Ser. I. iii. 41.—Europ. or.
opacum, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = sylvestre.
orientale, *Boiss. Diagn.* Ser. I. iii. 38.—Graec. ; As. Min.
orinocnse, Spreng. Syst. i. 897 = Rubia orinocensis.
orizabense, *Hemsl. Diagn. Pl. Nov.* 54.—Mexic.
ovale, Ruiz & Pav. Fl. Per. i. 59 = Rubia ovalis.
ovalifolium, Schott f. ex Schult. Mant. iii. 152 = ellipticum.
ovatum, Moench, Meth. 485 = saxatile.
palatinum, F. Schultz, ex Nym. Consp. 328 = *decolorans*.
pallens, Thuill. Fl. Par. ed. II. 77 = obliquum.
pallescens, Jan, Elench. 2. n. 70 = erectum.
pallidiflorum, Schur, Enum. Pl. Transs. 281 = verum.
pallidulum, Jord. Pugill. Pl. Nov. 80 = corrudaefolium !
pallidum, Presl, Delic. Prag. 121 = cinereum.
palustre, Bieb. Fl. Taur. Cauc. i. 108 = elongatum.
palustre, *Linn. Sp. Pl.* 105.—Europ. ; As. bor.
palustre, Poirr. ex Willd. & Lange, Prod. Fl. Hisp. ii. 323 = constrictum.
pamphylicum, *Boiss. & Heldr. Diagn.* Ser. I. x. 69.—As. Min.

GALIUM :—
pandoanum, Stend. Nom. ed. II. i. 659, sphalm. vide *pundoanum*.
Pansericanum, Schult. Mant. iii. 173 = filicaule.
papillosum, Heuff. in Flora, xl. (1857) 663 = aristatum.
papillosum, *Lapeyr. Hist. Abr. Pl. Pyr.* 66.—Mont Pyren.
paradoxum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 281.—Reg. Himal. ; Japon.
paralysne, *Vell. Fl. Flum.* 44 ; i. t. 112.—Bras.
parisifolium, Eschsch. ex Ledeb. Fl. Ross. iii. 412 = obovatum.
parisiense, *Linn. Sp. Pl.* 108.—Europ. ; Orient.
parisiense, Pall. Tabl. Taur. 46 = Asperula humifusa.
parviflorum, D. Don, Prod. Fl. Nep. 138 = G. Mollugo
parviflorum, Maxim. Prim. Fl. Amur. 472 = G. Aparine
parviflorum, Moench, Meth. 486 = gracile.
parviflorum, Rafin. in Med. Repos. N. York, v. (1808) 360 = concinnum !
parvifolium, Gaud. ex Roem. & Schult. Syst. iii. 246 = parisiense.
paschale, *Forsk. Fl. Aegypt. Arab.* 208.—Turc.
paucifiorum, Bunge, Enum. Pl. Chin. Bor. 35 = G. Aparine.
pauciflorum, Kotschy, Ins. Cyp. 260 = recurvum.
pauciflorum, Willd. ex DC. Prod. iv. 592 = Rubia orinocensis.
Paulitschum, F. Schultz, Arch. Fl. 158 = G. Mollugo.
pedemontanum, *All. Auct. Fl. Pedem.* 2.—Europ austr. ; Oriens.
pendeliflorum, *Boiss. Diagn.* Ser. I. iii. 46.—Graecia.
penicillatum, Bolss. l. c. x. 68 = nigricans.
pennsylvanicum, Bart. Comp. Fl. Philad. i. 83 = trifidum.
pennsylvanicum, Muhl. Cat. 15 ; ex Schult. Mant. iii. 183 = asprellum.
pentandrum, Gilib. Fl. Lituan. i. 10 = verum.
peplidiflolium, *Boiss. Diagn.* Ser. I. iii. 46.—As. Min.
peregrinum, *Pourret, Fl. Cher-et-Loire*, 260.—Gallia.
Perralderii, *Coss. in Bull. Soc. Bot. Fr.* ix. (1862) 172 —Algeria.
persicum, DC. Prod. iv. 606 = coronatum.
Pestalozzae, *Boiss. Diagn.* Ser. I. x. 65.—Syria.
Petrae, *[Oliver,] ex Hart, in Trans. Roy. Irish Acad.* xxvii. (1885) 488.—Palaestin.
petraeum, *Coss. & Dur. ex Munby, Cat. Pl. Algér* 15, nomen.—Algeria.
petrum, Schur, Enum. Pl. Transs. 284 = recurvum.
petrosum, Schur, l. c. = lucidum.
philistaeum, Boiss. Diagn. Ser. I. x. 71 = hierosolymitanum.
physocarpum, *Ledeb. Fl. Ross.* ii. 410.—Rossia.
pilliferum, *H. B. & K. Nov. Gen. et Sp.* iii. 337.—N. Granat.
pilosum, *Ait. Hort. Kew.* ed. I. i. 145.—Am. bor.
pisiferum, *Boiss. Diagn.* Ser. I. x. 67.—Syria.
plebeium, Boiss. & Heldr. Diagn. Ser. II. ii. 116 = sylvestre.
podlechianum, *Boss. Prim. Fl. Gal.* ii. 387.—Gallcia.
pogonanthum, Franch. & Savat. Enum. Pl. Jap. i. 213. ii. 393 = gracile.
Polretianum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 484.—Maroc.
poliganmum, Gilib. Fl. Lituan. i. 13 = verum.
polycarpum, Lam. ex Ledeb. Fl. Ross. ii. 402 = Asperula hamifusa.
polymorphum, *Knaf, in Flora*, xxix. (1846) 295.—Bohem.
pratense, *Scheele, in Linnaea*, xvii. (1843) 840.—German.
procumbens, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. i. 151 = saxatile.
productum, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1838) 29.—Ins. Madera.
prolliferum, *A. Gray, Pl. Wright.* ii. 67.—Am. bor. occ.
propinquum, A. Cunn. in Ann. Nat. Hist. ii. (1839) 208 = umbrosum.
Prostii, Jord. Obs. Pl. Crit. iii. (1846) 128 = obliquum.
prostratum, Hall. f. ex Roem. & Schult. Syst. iii. 238 = helveticum.
provinciale, Lam. Encyc. ii. 581 = erectum.
pruinosum, *Boiss. Elench.* 57.—Hispan.
pruinosum, C. Koch, in Linnaea, xxiv. (1851) 466 = aureum.
Pseudosparine, *Griseb, in Gerst. Abh.* vi. (1864) 125.—Reg. Magellan.
Pseudo-aristatum, Schur, Enum. Pl. Transs. 282 = sylvaticum.

GALIUM:—
Pseudo-cinereum, Schur, Enum. Pl. Transs. 285 = petrense.
Pseudo-rubioides, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 82 = rubioides.
Pseudo-verum, Schur, Enum. Pl. Transs. 281 = verum.
pubens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 350. —Calif.
pubescens, Req. ex DC. Prod. iv. 595 = pusillum.
pubescens, Schleich. ex DC. l. c. 596 = sylvestre.
pubescens, Schrad. ex Reichb. Fl. Germ. Excurs 210 = erectum.
pulchellum, Salisb. Prod. 58 = saxatile.
pulchellum, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 106.—Persia.
pulvinatum, *Boiss. Elench.* 87.—Hispan.
pumilum, Lam. Encyc. ii. 580 = pusillum.
pumilum, Murr. Prod. Gotting. 44 = sylvestre.
punctatum, Pers. Syn. i. 128 = pilosum.
puncticulatum, Michx. Fl. Bor. Am. i. 80 = pilosum.
pundsanum, Wall. Cat. n. 7291 = rotundifolium.
purpureum, DC. Fl. Fr. iv. 251 = rubrum
purpureum, *Linn. Sp. Pl.* 108.—Europ. austr.
purpureum, Wulf. Fl. Carol. 87 = pilosum.
pusillum, *Endl. ex Walp. Rep.* ii. 459.—Chili.
pusillum, *Linn. Sp. Pl.* 106.—Gall. austr.
pusillum, Sm. Engl. Bot. t. 74 = sylvestre.
pusillum, Vill. Hist. Pl. Dauph. ii. 324. t. 8 = helveticum.
pycnophyllum, Heldr. ex Boiss. Fl. Orient. iii. 64 = spicilatum.
pygmaeum, DC. Prod. iv. 606 = articulatum.
pyrenaicum, *Gouan, Illustr.* 5. t. 1. f. 4.—Mont. Pyren.
pyrenaicum, Sibth. & Sm. Fl. Graec. Prod. i. 92 = olympicum.
quinquefidum, *Andrz. Rys. Botan.* ii. (1830) 84.—Podolia.
radicosum, *Steud. Nom.* ed. II. i. 660.—Bras.
recurvum, *Req. ex DC. Prod.* iv. 609.—Archip. Graec.
reflexum, Pohl, ex DC. l. c. 604 = hirtum.
reflexum, Presl, Delic. Prag. 128 = pedemontanum.
Kelbun, *Clos, in C. Gay, Fl. Chil.* iii. 186.—Chili.
Requienianum, Wight & Arn. Prod. 443 = rotundifolium.
retrorsum, DC. Prod. iv. 605 = pedemontanum.
Richardianum, *Endl. ex Walp. Rep.* ii. 459.—Chili.
rigidum, *Ait. Hort. Kew.* ed. I. i. 144.—Ins. Azores.
rigidum, Vill. Hist. Pl. Dauph. ii. 319 = erectum.
rivale, Griseb. Spicil. Fl. Rumel. ii. 156 = Asperula Aparine.
rivulare, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 15.—Hispan.
rosellum, *Boiss. & Reut. Pugill. Pl. Nov.* 52.—Hispan.
Rothrockii, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 208.—Am. bor. occ.
rotundatum, Griseb. Spicil. Fl. Rumel. ii. 157 = Asperula laevigata.
rotundifolium, Herb. Linn. ex DC. Prod. iv. 582 = Asperula laevigata.
rotundifolium, *Linn. Sp. Pl.* 108.—Europ.; As. temp.
rotundum, Thunb. Fl. Jap. 89 = gracile.
rubidum, Jord. Obs. Pl. Crit. iii. (1846) 121 = rubrum.
rubioides, Brot. Fl. Lusit. i. 148 = Broterianum.
rubioides, *Linn. Sp. Pl.* 108.—Europ.; As. bor.
rubioides, Pollich, Hist. Pl. Palat. i. 160 = boreale.
rubioides, Suter, Fl. Helv. i. 88 = verum.
rubioides, Willd. ex Ledeb. Fl. Ross. ii. 411 = valantioides.
rubioides, Hill, Veg. Syst. vii. 21 = rubioides.
rubrum, *Linn. Sp. Pl.* 107.—Europ. austr.
rubrum, Pollich, Hist. Pl. Palat. i. 155 = parisiense.
rubrum, Scop. Fl. Carn. ed. II. i. 108 = purpureum.
rugosum, *DC. ex W. Bast. in Loud. Hort. Brit. Suppl.* ii. 686, nomen.—Hab.?
rupestre, DC. Prod. iv. 608 = aureum.
rupestre, Vis. in Flora, xii. (1829) i. Erg. 5 = Asperula scutellaris.
rupestre, *Walp. Rep.* vi. 17.—Bras.
rupicolum, Bertol. Fl. Ital. ii. 108 = helveticum.
rupicolum, Bor. Fl. Cent. Fr. ed. III. ii. 606 = palustre.
rupicolum, *Pomel, Nouv. Mat. Fl. Atl.* 74.—Alger.
ruricolum, Jord. Pugill. Pl. Nov. 81 = parisiense.
ruthenicum, Willd. Sp. Pl. i. 597 = verum.
sachsenstum, *All. Fl. Pedem.* i. 9.—Reg. Mediterr.
Sadleri, Frivald. ex Hampe, in Flora, xx. (1837) 238 = pyrenaicum.
saturejaefolium, Trevir. in Ges. Naturf. Fr. Berl. Mag. vii. (1815) 146 = elongatum.

GALIUM:—
saxatile, Herb. Johann. ex Vest, in Flora, iv. (1821) 587 = Vestianum.
saxatile, *Linn. Sp. Pl.* 106.—Europ.
scabrrimum, Vahl, ex Hornem Hort. Hafn. i. 185 = G. Aparine.
scabridum, Jord. Obs. Pl. Crit. iii. (1846) 136 = Jordani.
scabridum, Willk. in Flora, xxxv. (1852) 541 = sylvestre.
scabrum, Jacq. Fl. Austr v. 10 = sylvestre.
scabrum, *Linn. Sp. Pl.* 108.—Europ austr.
scabrum, Spreng. Fl. Halens. 82 = parisiense.
scabrum, With. Bot. Arr. Brit Pl. ed. III i. 190 = G. Mollugo.
scandens, *Phil. in Anal. Univ. Chil.* (1872) 731.—Chili.
Schlumbergeri, *Boiss. Fl. Orient.* iii. 51.—Syria.
Schultessii, Vest, in Flora, iv. (1821) 580 = aristatum.
Scopoli, Vill. Hist. Pl. Dauph. i. 304 = vernum.
segetum, C. Koch, in Linnaea, xvii. (1843) 38 = spurium.
Sellowianum, *Walp. Rep.* ii. 459.—Bras.
septentrionale, Roem. & Schult. Syst. iii. 253 = boreale
serotinum, *Boiss. & Heldr. Diagn.* Ser I. x 71.—As Min.
serotinum. *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 284.—Algeria.
serpyllioides, *Royle, ex Hook. f. Fl. Brit. Ind.* iii. 207.—Reg. Himal.
sessiliflorum, Willd. ex Schult. Mant. iii. 179 = Relbunium ciliatum.
setaceum, *Lam. Encyc.* ii. 584.—Reg. Mediterr.; As. temp.
Sibthorpii, Roem. & Schult. Syst. iii. 244 = setaceum
Sieberi, *Tausch, in Flora*, xviii. (1835) 356.—Corsica.
silvaticum, Bellil & Timb. in Mem. Acad. Toul Scr. V. vi. (1862) 239 = sylvestre.
simense, Fresen. in Mus. Senckenb. ii. (1837) 165 = spurium.
simense, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 344.—Abyss.
sinaicum, *Boiss. Fl. Orient.* iii. 63.—Arab.
Smithii, G. Don, Gen. Syst. iii. 660 = verticillatum
Soleirolii, *Duby, in Mém. Soc. Linn. Par.* vi. (1827) 399.—Ins. Corsica.
songaricum, *Schrenk, Enum. Pl. Nov.* 57.—Sibir. altaic.
sororium, Hance, in Walp. Ann. ii. 784 = G. Aparine.
spinulosum, Mérat, Nouv. Fl. Par. 58 = uliginosum.
spinulosum, Kafin. Prèc. Découv. (1814) 40 = asperellum.
splendens, Hornem. Suppl. Hort. Hafn. 17 = erectum.
Spruneri, Jord. Obs Pl. Crit. iii. (1846) 116 = aureum.
spurium, All. Fl. Pedem. i. 5 = parisiense.
spurium, Huds. Fl. Angl. ed. I. 57 = tricorne.
spurium, *Linn. Sp. Pl.* 106.—Europ.; Orient.
spurium, Spreng. Fl. Halens 82 = uliginosum.
squalidum, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 462 [bis] = australe.
stellariaefolium, Franch. & Sav. Enum. Pl. Jap. ii. 392 = parodoxum.
stellatum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 97. f. 26.—Am. bor. occ.
stenophyllum, Schur, Enum. Pl. Transs. 283 = aristatum.
Sternbergii, Knaf, ex Nym. Consp. 324 = *ochroleucum*.
stoloniferum, C. Koch, in Linnaea, xix. (1847) 82 = orientale.
strictum, Torr. Cat. Pl. N. York, 23 = boreale.
strigosum, *Thunb. in Nov. Act. Soc. Sc. Upsal.* vii (1818) 141.—Japon.
suaveolens, Lapeyr. Suppl. Hist. Pl. Pyr. 23 = megalospermum.
suaveolens, Wahlb. Fl. Lapp. 48 = trifforum.
suberosum, Griseb. Spicil. Fl. Rumel. ii. 161 = Asperula involucrata.
suberosum, *Sibth. & Sm. Fl. Graec. Prod.* i. 91.—Ins. Creta.
sublutaceum, Gilib. Fl. Lituan. i. 10 = verum.
subtrifidum, *Reinw. ex Blume, Bijdr.* 944.—Java.
subvelutinum, C. Koch, in Linnaea, xxiv. (1851) 465 = leiophyllum.
subvillosum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 88. —Afr. austr.
sudeticum, Tausch, in Flora, xviii. (1835) 347 = aniso-phyllum.

GALIUM:—
suffruticosum, *Hook. & Arn. in Hook. Bot Misc.* iii. (1833) 368.—Chili
suffruticosum, Nutt. ex Torr. & Gray, Fl N. Am. i. 21 = Nuttallii.
supinum, Lam. Fl. Fr. iii. 379 = sylvestre.
sylvaticum, Bess. Prim. Fl. Galic. i. 219 = aristatum.
sylvaticum, *Linn. Sp. Pl.* ed. II. 155.—Europ
sylvestre, *Polich, Hist. Pl. Palat.* i. 151.—Europ.
syriacum, *Boiss. Diagn.* Ser. I. ix. 47.—Syria.
tarmense, Spreng. Syst. i. 387 = leucocarpum.
tataricum, *Trevir. in Ges. Naturf. Fr. Berl. Mag.* vii. (1815) 146.—Astrakan.
tauricum, Roem. & Schult. Syst. iii 250 = coronatum
taurinum, Scop. Fl. Carn. ed. II. i. 101 = Asperula taurina
Tendae, Reichb. f. Ic. Fl. Germ. xvii. 97. t. 1194 = helveticum.
tenellum, *Clos, in C. Gay, Fl. Chil.* iii. 188.—Chili.
tenellum, *Jord. Obs. Pl. Crit.* iii. (1846) 130.—Europ
tenerrimum, Schur, Enum. Pl. Transs. 279 = G. Aparine
tenerum, Pers. Syn. i. 128 = parisiense.
teucrum, Schleich. Cat Pl. Helv ed. IV. 18; Gaud Fl. Helv. i. 442 = spurium.
tenue, *Vill. Prosp.* 20; *Hist. Pl. Dauph.* ii. 322.—Europ.
tenuicaule, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 206.—N. Zel.
tenuicaule, Jord. Pugill. Pl. Nov. 81 = divaricatum.
tenuicaule, Krock. Fl. Siles Suppl. i. 214 = saxatile?
tenuifolium, All. Fl. Pedem. i. 6 = corrudaefolium.
tenuissimum, *Bieb. Fl. Taur. Cauc.* i. 104.—Orient.; Reg. Himal.
tetragonum, *Griseb, in Goett. Abh.* xix. (1874) 161.—Reg. Argent
texanum, Scheele, in Linnaea, xxi. (1848) 597 = virgatum.
texense, *A Gray, in Proc. Am. Acad.* xix. (1883) 50. —Texas.
Thunbergianum, Eckl. & Zeyh. Enum. 369 = rotundifolium.
thymifolium, *Boiss. & Heldr.* Ser. I. vi. 67.—Greec
Timeroyi, Jord. Obs. Pl. Crit iii (1846) 138 = Jordani.
tinctorium, Linn. Sp. Pl. 108 = trifidum.
tinctorium, Scop. Fl. Carn. ed. II. i. 101 = Asperula tinctoria.
tmoleum, *Boiss. Diagn* Ser I. iii. 89.—Lydia.
tolosianum, *Boiss. & Kotschy, in Boiss Fl. Orient.* iii. 53.—As. Min.
tomentosum, *Thunb. Fl. Cap.* i. 551.—Afr. austr.
Torreyi, Bigel. Fl. Bost. ed. II. 56 = lanceolatum.
trachyanthum, Boiss. & Hohen. Diagn. Ser. I. iii. 50 = hierosolymitanum.
trachycarpum, Clarke, Trav. ii. 260 = sparinoides?
trachyspermum, A. Gray, in Perry, Jap. Exp. ii. 313 = gracile.
transcaucasicum, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 52.—Reg. Caucas.
transsilvanicum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 33 = aristatum.
truncatum, Roem. & Schult. Syst. iii. 269, in syn. aphalm.=Asperula tinctoria.
trichocarpon, Willd. ex Spreng. Syst. i. 392 = canescens.
trichocarpum, *DC. Prod.* iv. 600.—Chili.
trichophorum, Griseb. Spicil. Fl. Rumel. ii. 161 = intricatum.
trichophorum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 372.—Sibir. altaic.
trichophyllum, All. Auct. Fl. Pedem. i = pusillum.
trichophyllum, Wulf. in Roem. Arch. iii. 319 (err. typ. 329) = verum.
tricorne, *Stokes, in With. Bot. Arr. Brit. Pl.* ed. II. i. 153.—Europ.; Orient.
trifidum, *Linn. Sp. Pl.* 105.—Reg. bor. temp.
trifidum, Urv. in Mém. Soc. Linn. Par. iv. (1826) 612 = antarcticum.
trifforum, *Michx. Fl. Bor. Am.* i. 80.—Reg. bor. temp.
trifforum, Rochel, ex Steud. Nom. ed. I. 355 = tricorne.
triscarium, Roem. & Schult. Syst. iii. 238 = setnicum.
trinervo, Moench, Meth. 486 = boreale.
trinervium, Gilib Fl. Lituan. i. 19 = boreale.
trinioides, *Pomel, Nouv. Mat Fl. Atl.* 77.—Alger.
Tsukinoanis, Hansk. Cat. Hort. Bog. Alt. 106 = strigosum?

GALIUM:—
tuberculatum, Preal, Delic. Prag. 120 = verum.
tuberosum, *Lour. Fl. Cochinch.* 79.—China.
tunetanum, Bourg. ex Nym. Consp. 328 = verum.
tunetanum, *Lam. Encyc.* ii. 583.—Afr. bor. occ.
tyrolense, Willd. Enum. Hort. Berol. 153 = erectum.
uliginosum, *Linn. Sp. Pl.* 106.—Europ.; As. bor.
uliginosum, Pall. ex Bieb. Fl. Taur. Cauc. i. 102 = Asperula Aparine.
uliginosum, Thunb. Fl. Jap. 58 = strigosum.
umbellatum, Lam. Encyc. ii. 579 = sylvestre.
umbrosum, *Soland. ex Forst. Prod.* 89, nomen.—N. Zel.
uncinatum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 484 = G. Aparine.
uncinatum, Lichtenst. ex Bartl. & Wendl. f. Beitr. ii. 13 = glabrum.
uncinulatum, *DC. Prod.* iv. 600.—Mexic.
uncinulatum, A. Gray, in Bost. Journ. Nat. Hist. vi. (1857) 215 = texense.
undulatum, Stend. Nom. ed. II. i. 661, sphalm. vide *uncinatum*, Lichtenst.
uniflorum, *Michx. Fl. Bor. Am.* i. 79.—Am. bor.
uropetalum, *Hemsl. Biol. Centr. Am. Bot.* ii. 66.—Mexic.
Urvillei, Req. ex DC. Prod. iv. 607 = setaceum.
vaginx, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 461 [bis] = Gaudichaudi.
Vaillantii, DC. Fl. Fr. iv. 268 = G. Aparine.
Valantia, Gaertn. Mey. & Scherb. Fl. Wett. i. 188 = G. Cruciata.
Valantia, [Web. in] Wigg. Prim. Fl. Holsat. 12 = tricorne.
valantioides, *Bieb. Fl. Taur. Cauc.* i. 102.—Reg. Caucas.
valdivianum, *Phil. in Linnæa*, xxviii. (1856) 655.—Chili.
valentinum, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 95.—Hispan.
venustum, *Jord. Obs. Pl. Crit.* iii. (1846) 117.—Ins. Corsica.
verrucum, Hook. f. Fl. Brit. Ind. iii. 209 = cryptanthum.
vernum, *Scop. Fl. Carn.* ed. II. i. 99. t. 2.—Europ.; As. temp.
vernum, Sieber, ex Tausch, in Flora, xviii. (1835) 856 = Sieberi.
verosimile, Roem. & Schult. Syst. iii. 234 = verum.
verrucosum, Sm. Engl. Fl. i. 204 = saccharatum.
verticillatum, *Danth. ex Lam. Encyc.* ii. 585.—Europ. austr.; Oriens.
verticillisforum, Fourr. ex Lapeyr. Suppl. Hist. Pl. Pyr. 24 = verticillatum.
verum, *Linn. Sp. Pl.* 107.—Europ.; As. temp.
Vestianum, *Schult. Mant.* iii. 164.—Europ.
vestitum, *D. Don, Prod. Fl. Nep.* 133.—Reg. Himal.
vexans, *Reichb. f. Ic. Fl. Germ.* xvi. 92 = Valantia muralis.
Villarsii, Gand. ex Roem. & Schult. Syst. iii. 230 = sylvestre!
Villarsii, Req. ex Guer. Vasd. ed. II. 250; ex DC. Fl. Fr. vi. 497 = megalospermum.
villosum, Lam. Encyc. ii. 583 = maritimum.
virgatum, *Nutt. ex Torr. & Gray, Fl. N. Am.* ii. 20.—Am. bor.
virgultorum, *Lamotte, Prod.* ii. 360.—Gallia.
viridiflorum, *Boiss. & Reut. Pugill. Pl. Nov.* 51.—Hispan.
viridulum, Jord. Pugill. Pl. Nov. 79 = corrudæfolium.
viscosum, *Vahl, Symb. Bot.* ii. 39.—Afr. bor.
vulcanicum, Boiss. & Hohen. Diagn. Ser. I. x. 70 = nigricans.
vulgare, S. F. Gray, Nat. Arr. Bot. Pl. ii. 483 = G. Mollugo.
Walteri, J. F. Gmel. Syst. 327 = circæzans.
Wirtgenii, F. Schultz, Arch. Fl. 201 = verum.
Witheringii, Sm. Fl. Brit. 174 = palustre.
wittbergense, *Sond. in Harv. & Sond. Fl. Cap.* iii. 37.—Afr. austr.
Wrightii, *A. Gray, Pl. Wright.* i. 80.—Am. bor. occ.
xylorrhizum, Boiss. & Hoet, Diagn. Ser. II. ii. 115 = leiophyllum.
zacynthicum, Rent. in Mém. Soc. Phys. Genèv. vii. (1836) 304 = intricatum.

GALIZIOLA, Rafin. Sylva Tellur. 167 (1838) = Ardisia, Sw. (Myrsin.).

GALLAPAGOA, Pritz. Ic. Ind. ii. 129 (1866) = GALAPAGOA, Hook. f. = Coldenia, Linn. (Borag.).

GALLARIA, Schrank, ex Endl. Gen. 1217 (1841) = Medinalla, Gaudich. (Melastom.).

GALLASIA, Mart. ex DC. Prod. iii. 182 (1828) = Miconia, Ruiz & Pav. (Melastom.).
pepericarpa, Mart. ex DC. l. c. = M. pepericarpa.

GALLESIA, Casar. Nov. Stirp. Bras. Dec. 48 (1842). *PHYTOLACCACEÆ*, Benth. & Hook. f. iii. 88.
Gomeziana, Moq. in DC. Prod. xiii. II. 8 = G. Scorododendrum.
Scorododendron, *Casar. Nov. Stirp. Bras. Dec.* 44.—Bras.

GALLESIOA, M. Roem. Syn. Hesper. 83, 45 (1846) = Clausena, Burm. f. (Rutac.).
gravoleus, M. Roem. l. c. 45 = C. excavata.
pentaphylla, M. Roem. l. c. = C. pentaphylla.

GALLINSOGA, Juuss ex Hill. Expos. i. 417 (1805) = Galinsoga, Ruiz & Pav. (Compos.).

GALLITRICHUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134 = Salvia, Tourn. (Labiat.).
anglicum, Jord. & Fourr. Ic. Pl. Eur. ii. 19 = S. Verbenaca.
ornale, Jord. & Fourr. l. c. 18 = S. Verbenaca.
Candelle, Timb. in Mém. Acad. Toul. Sér. VII. ii. (1870) 288 = S. Candollei.
clandestinum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134 = b. Verbenaca.
Clusii, Timb. in Mém. Acad. Toul. Sér. VII. ii. (1870) 287 = S. pratensis.
dichroanthum, Jord. & Fourr. Ic. Pl. Eur. ii. 18 = S. Verbenaca.
Fuschii, Timb. in Mém. Acad. Toul. Sér. VII. ii. (1870) 243 = S. Fuschii.
Hermanni x, Timb. l. c. 244.—Gallia.
hormineum, Timb. l. c. 241 = S. Verbenaca.
maculatum, Jord. & Fourr. Ic. Pl. Eur. ii. 19 = S. Verbenaca.
palidiflorum, Jord. & Fourr. l. c. 18 = S. Verbenaca.
pratense, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134 = S. pratensis.
stypholophyllum, Jord. & Fourr. Ic. Pl. Eur. ii. 17 = S. Verbenaca.
vosulatum, Jord. & Fourr. l. c. = S. Verbenaca.
rubellum, Jord. & Fourr. l. c. 20 = S. Verbenaca.
stereocaulon, Jord. & Fourr. l. c. 19 = S. Verbenaca.
verbenacum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134 = S. Verbenaca.
virgatum, Jord. & Fourr. Ic. Pl. Eur. ii. 18 = S. Verbenaca.

GALLIUM, [Tourn.] Linn. Syst. ed. I (1735) = Galium, Linn. (Rubiac.).

GALLOA, Hassk. Cat. Hort. Bog. Alt. 172 (1844) = Cocculus, DC. (Menisperm.).
trinervia, Hassk. l. c. = C. laurifolius.

GALOGLYCHIA, Gasp. Nov. Gen. Fic. 10 (1844) = Ficus, Tourn. (Urtic.).
mauritanensis, Gasp. l. c. = F. laurifolia.
Saussureana, Gasp. l. c. = F. Saussureana.
Tenoreana, Gasp. l. c.—Hab.?

GALOPHTHALMUM, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 7. t. 2 = Blainvillea, Cass. (Compos.).
brasiliensis, Nees & Mart. l. c. = B. bahiensis.

GALOPINA, Thunb. Nov. Gen. i. 3 (1781). *RUBIACEÆ*, Benth. & Hook. f. ii. 189.
OXYPHORUS, Eckl. & Zeyh. Enum. 264 (1837).
aspera, Sond. in Harv. & Sond. Fl. Cap. iii. 26 = G. Oxyspermum.
circæoides, *Thunb. Nov. Gen.* i. 3.—Afr. austr.
hirsuta, E. Mey. ex Harv. & Sond. Fl. Cap. iii. 26 = G. Oxyspermum.
Oxyspermum, *Stend. Nom.* ed. II. i. 662.—Afr. austr.
tomentosa, Hochst. in Flora, xxvii. (1844) 555 = G. Oxyspermum.

GALORDIA, Raeusch. Nom. ed. III. 251 (1797) = Gaillardia, Fougr. (Compos.).
alternifolia, Raeusch. l. c. = Gaillardia pulchella.

GALORHOEUS, Endl. Gen. 1108 (1841) = GALARHOEUS, Haw. = Euphorbia, Linn.

GALPHIMIA, Cav. ic. v. 61. t. 489 (1799). *MALPIGHIACEÆ*, Benth. & Hook. f. i. 254.
THRYALLIS, Linn. Sp. Pl. ed. II. 554 (1762).
angustifolia, *Benth. Bot. Voy. Sulph.* 9.—Am. bor. occ.
biglandulosa, Pohr. Encyc. Suppl. iv. 7 = glandulosa.
brasiliensis, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 71.—Bras.
Cavea, Spreng. Syst. ii. 385 = Caucanthus arabicus.
chrysophylla, Spreng. Syst. ii. 385 = Byrsonima chrysophylla.
glandulosa, *Cav. in Anal. Hist. Nat.* i. (1799) 87.—Mexic.
glandulosa, H. B. & K. Nov. Gen. et Sp. v. 172. t. 459 = G. Humboldtiana.
glauca, *Cav. in Anal. Hist. Nat.* i. (1799) 87.—Mexic.
glauca, Hort ex Bartl. in Linnæa, xiii. (1839) 552 = sec.
gracilis, *Bartl. l. c.*—Mexic.
grandiflora, *Bartl. l. c.* 554.—Mexic.
hirsuta, *Cav. in Anal. Hist. Nat.* i. (1799) 87.—Mexic. Humboldtiana, *Bartl. in Linnæa*, xiii. (1839) 555.—Mexic.
latifolia, *Bartl. l. c.* 553.—Mexic.
linifolia, A. Gray, Gen. Illustr. ii. 196. t. 173 = angustifolia.
longifolia, H. B. & K. Nov. Gen. et Sp. v. 173 = Lophanthera Kunthiana.
mollis, Hort. ex Bartl. in Linnæa, xiii. (1839) 558 = hirsuta.
mollis, H. B. & K. Nov. Gen. et Sp. v. 173 = Tetrapterys Cotonaster.
multicaulis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 322.—Mexic.
paniculata, *Bartl. in Linnæa*, xiii. (1839) 556.—Mexic.
splendens, Mackoy, ex A. Juss. in Walp. Rep. v. 184 = gracilis.

GALPHINÍA, Poir. in Dict. Sc. Nat. xviii. 116 (1820) = præc.

GALSTRONEMA, Stend. Nom. ed. II. i. 662 (1840) = GASTRONEMA, Herb. = Cyrtanthus, Ait. (Amaryllid.).

GALTONIA, Decne. in Fl. des Serres, xxiii. (1880) 82. *LILIACEÆ*, Benth. & Hook. f. iii. 809.
candicans, *Decne. l. c.* 83.—Afr. austr.
clavata, [*Mast. in*] *Gard. Chron.* (1884) ii. 401.—Afr. austr.
Princeps, *Decne. in Fl. des Serres*, xxiii. (1880) 83.—Afr. austr.

GALUMPITA, Blume, Mus. Bot. Lugd. Bat. ii. 78. 1852 (1856) = Gironniera, Gaudich. (Urtic.).
cuspidata, Blume, l. c. = Girono. reticulata.

GALURUS, Spreng. Anleit. ii. ii. 364 (1817) = Acalypha, Linn. (Euphorb.).
spicijlorus, Spreng. Syst. i. 862 = A. spicijlora.

GALVANIA, Vand. Fl. Lusit. 15. t. 1. f. 7 (1788) = Pergobotria, Linn. (Rubiac.).
Vandellii, Spreng. Neue Entdeck. ii. 151 = P. Margravii.
Vellosii, Roem. & Schult. Syst. v. 198 = P. Margravii.

GALVESIA, J. F. Gmel. Syst. 927 (1791) = Galvezia, Domb. (Scrophul.).

GALVESIA, Pers. Syn. i. 445 (1805) = GALVEZIA, Ruiz & Pav. = Pitavia, Molina (Rutac.).

GALVEZIA, Domb. ex Juss. Gen. 119 (1789). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 984.
AGASSIZIA, Chav. Monog. Antirrh. 180. t. 11 (1846).
SACCULARIA, Kellogg, in Proc. Calif. Acad. ii. (1863) 17.
fruticosa, *Y. F. Gmel. Syst.* 927.—Peruv.
Hawesiæ, Domb. ex Chav. Monog. Antirrh. 180 = fruticosa.
punctata, *Ruiz & Pav. Fl. Per.* iv. t. 344.—Peruv.

GALVEZIA, Ruiz & Pav. Syst. Veg. 97 (1794) = **Piavia**, Molina (Ruiz.).
punctata, Ruiz & Pav. l. c. — P. punctata.
spicata, Bert. ¹ Merc. Chil. 642.—Cf. Bull. Férussac, xx. (1830) 111 = Ercilla volubilis.

GAMA, La Llave, in La Naturaleza, vii. (1885) Apénd. 71 = **Matricaria** ? Llan. (Compos.).
angulata, La Llave, l. c.—Mexic.

GAMANTHUS, Bunge, Anabas. Rev. 76 (Mém. Acad. Pétersb. Sér. VI. iv. (1862) n. 11) = **Halanthium**, C. Koch (Chenop.).
barbellatus, Bunge, in Act. Hort. Petrop. vi. (1879) 448.—Turkestan.
communis, Bunge, Anabas. Rev. 76.—Afghan.
gamocarpus, Bunge, l. c. 77. t. 1 f. 14.—Afghan.
sviuus, Bunge, in Act. Hort. Petrop. vi. (1879) 447.—Turkestan.
pilosus, Bunge, Anabas. Rev. 79. t. 1.—Reg. Casp.

GAMARIA, Rafin. Fl. Tellur. iv. 49 (1836) = **Disa**, Sw. (Orchid.).
cornuta, Rafin. l. c. = D. comuta.

GAMAZYGIS, Pritz. lc. Ind. i. 475 (1865) = GAMOZYGIS, Turcz. = **Angianthus**, Wendl. (Compos.).

GAMBELIA, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 149 = **Antirrhinum**, Linn. (Scrophul.).
speciosa, Nutt. l. c. = A. speciosum.

GAMBLEA, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 739 (1879). *ARALIACEAE.*
ciliata, *C. B. Clarke, l. c.* 740.—Reg. Himal.

GAMELYTHRUM, Nees, in Hook. Lond. Journ. Bot. ii (1843) 415 = **Amphipogon**, R. Br. (Gramin.).
denudatum, Nees, in Lehm. Pl. Preiss. ii. 101 = A. turbinatus.
turbinatum, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 415 = A. turbinatus.

GAMELYTRUM, Steud. Syn. Pl. Gram. 198 (1854) = GAMELYTHRUM, Nees = **Amphipogon**, R. Br. (Gramin.).

GAMOCARPHA, DC. Prod. v. 2 (1836) = **Boopis**, Juss. (Calycer.).
angustifolia, Phil. in Anal. Univ. Chil. ii. (1862) 879. —Chili
dentata, Phil. l. c. (1872) 785.—Chili.
Gillesii, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 281.—Chili.
lignlata, Miers, l. c. 282.—Chili.
Poeppigii, DC. Prod. v. 2 = B. alpina.
polycephala, Phil. in Anal. Univ. Chil. (1856) ii. 329. —Chili.
spuria, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 282.—Chili.

GAMOCHAETA, Wedd. Chlor. And. i. 151 (1856) = **Gnaphalium**, Linn. (Compos.).
americana, Wedd. l. c. = Gn. purpureum.
capitata, Wedd. l. c. 158 = Gn. capitatum ?
humilis, Wedd. l. c.—Bolivia.
serpyllifolia, Wedd. l. c. 159 = Gn. serpyllifolium.
sylvatica, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 93 = Gn. sylvaticum.

GAMOCHILUM, Walp. in Linnæa, xiii. (1839) 509 = **Argyrolobium**, Eckl. & Zeyh. (Legumin.).
obcordatum, Walp. l. c. 510 = A. obcordatum.
sericum, Walp. l. c. = A. sericeum.
umbellatum, Walp. l. c. 509 = A. umbellatum.

GAMOCHILUS, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 341 = **Hedychium**, Koen. (Scitamin.).
speciosus, Lestib. l. c. = H. Gardnerianum.

GAMOCHLAMYS, Baker, in Gard. Chron. (1876) 164 = **Spathanthemum**, Schott (Aroid.).
heterandra, Baker, l. c. = S. heterandrum.

GAMOGYNE, N. E. Br. in Journ. Bot. xx. (1882) 346. *AIZOIDEAE*, Benth. & Hook. f. iii. 905.
Burbidgei, *N. E. Br. l. c.* 196.—Borneo.

GAMOLEPIS, Less. Syn. Comp. 251 (1832). *COMPOSITAE*, Benth. & Hook. f. iii. 492.
JACQUEMONTIA, Belang. Voy. Ic. t. 10 (1836).
PHILOTHAMNUS, DC. Prod. vi. 41 (1837).
PSILOTHONNA, E. Mey. ex DC. Prod. vi. 40 (1867).
annua, *Less. Syn. Comp.* 251.—Afr. austr.
brachypoda, *DC. Prod.* vi. 40.—Afr. austr.
chrysanthemoides, *DC. l. c.*—Afr. austr.
debilis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 157.—Afr. austr.
ericoides, *Less. Syn. Comp.* 251.—Afr. austr.
eurlopoides, *DC. Prod.* vi. 41.—Afr. austr.
hebecarpa, *DC. l. c.*—Afr. austr.
laxa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 158.—Afr. austr.
leiocarpa, *Harv. l. c.* 608.—Afr. austr.
munita, *Less. Syn. Comp.* 251.—Afr. austr.
pectinata, *Less. l. c.*—Afr. austr.
pinnatipartita, *Harv. in Harv. & Sond. Fl. Cap.* iii. 607.—Afr. austr.
polytrichoides, *Harv. l. c.*—Afr. austr.
Tagetes, DC. Prod. vi. 40 = annua.
tagetis, St Lag. in Ann Soc. Bot. Lyon, vii. (1880) 126 = prace.
trifurcata, *Less. Syn. Comp.* 251.—Afr. austr.

GAMOPLEXIS, Falc. in Proc. Linn. Soc. i. (1847) 320; et in Trans. Linn. Soc. xx. (1847) 293. t. 18 = **Gastrodia**, R. Br. (Orchid.).
orobanchoides, Falc. ll. cc.—Reg. Himal.

GAMOZYGIS, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 75 = **Angianthus**, Wendl. (Compos.).
flexuosa, Turcz. l. c. 76 = A. amplexicaulis.

GAMPSOCARES, Stev. in Bull. Soc. Nat. Mosc. xxv. (1852) i. 541, in textu = **Ranunculus**, Tourn.
Pinardi, Stev. l. c. 542 = R. Pinardi.

GANDOLA, Moq. in DC. Prod. xiii. ii. 224 (1849) = **Ulluous**, Lozan. (Chenop.).
peruviana, Moq. l. c. 225 = U. tuberosus.
tuberosa, Moq. l. c. 224 = U. tuberosus.

GANDOLA, Rumph. ex Linn. Sp. Pl. ed. II. 890 (1762) = **Basella**, Kheede (Chenop.).
nigra, Rafin. Sylva Tellur. 60 = B. rubra.

GANDRILOA, Stend. Nom. ed. II. i. 662 (1840) = **Chenopodium**, Linn.
atriplicoides, Stend. l. c. = C. polyspermum.

GANGILA, Bernh. in Linnæa, xvi. (1842) 42 = **Sesamum**, Linn. (Pedalin.).
pentaphylla, Bernh. l. c.—Afr. austr.

GANITRUS, Gaertn. Fruct. ii. 271. t. 139 (1791) = **Elaeocarpus**, Linn. (Tiliac.).
obtusus, Rafin. Sylva Tellur. 59 (Ganitrum) = E. integer.
Roxburghii, Wight, Illustr. 88 = E. Ganitrus.
sphaerica, Gaertn. Fruct. ii. 271. t. 139 = E. Ganitrus.

GANJA, Reichb. Handb. 308 (1837) = **Corchorus**, Linn. (Tiliac.).

GANOPHYLLUM, Blume, Mus. Bot. Lugd. Bat. i. 230 (1850). *BURSERACEAE*, Benth. & Hook. f. i. 326.
falcatum, *Blume, l. c.*—Malacca.

GANOSMA, Decne. in DC. Prod. viii. 625 (1844), sphalm. = **Aganosma**, D. Don (Apocyn.).

GANSBLUM, Adans. Fam. ii. 490 (1763) = **Strophila**, DC. (Crucif.).

GANYMEDES, Salisb. in Trans. Hort. Soc. i. (1812) 353 = **Narcissus**, Tourn. (Amaryll.).
alba, Haw. Narcis. Monog. 5 = N. triandrus.
capax, Herb. Amaryll. 308 = N. triandrus.
cernuus, Haw. Suppl. Pl. Succ. 130 = N. Pseudonarcissus.
cernuus, Salisb. in Trans. Hort. Soc. i. (1812) 353 = N. triandrus.
concolor, Haw. Suppl. Fl. Succ. 131 = N. triandrus.
effusus, Salisb. in Trans. Hort. Soc. i. (1812) 354 = N. triandrus.
Linnaei, Kunth, Enum. Pl. v. 719 = N. triandrus.

GANYMEDES :—
nutans, Haw. Suppl. Pl. Succ. 131 = Narcissus triandrus.
ochroleucus, Haw. l. c. 206 = N. triandrus.
pulchellus, Salisb. in Trans. Hort. Soc. i. (1812) 34 = N. triandrus.
reflexus, Herb. Amaryll. 308 = N. triandrus.
striatulus, Haw. Suppl. Pl. Succ. 132 = N. triandrus.
triandrus, Haw. l. c. 130 = N. triandrus.

GARACIUM, Gren. & Godr. Fl. Fr. ii. 327, in syn. (1850) = **Lactuca**, Tourn. (Compos.).
alpinum, Gren. & Godr. l. c. = L. alpina.

GARALEUM, Sch. Bip. in Walp. Rep. ii. 954 (1843), sphalm. = **Garuleum**, Cass. (Compos.).

GARAPATICA, Karst. Fl. Colomb. i. 57. t. 28 (1858) = **Alibertia**, A. Rich. (Rubiac.).
edulis, Karst. l. c. = A. edulis.

GARBERIA, A. Gray, in Proc. Acad. Sc. Philad. (1879) 379. *COMPOSITAE.*
fruticosa, *A. Gray, l. c.*—Florida.

GARCIA, Rohr, in Skriv. Nat. Hist. Selsk. Kjobenh. ii. (1792) 217. t. 9. *EUPHORBIACEAE*, Benth. & Hook. f. ii. 392.
nutans, *Rohr, l. c.*—Mexic.

GARCIANA, Lour. Fl. Cochinch. 14 (1790) = **Phillydrum**, Banks.
cochinchinensis, Lour. l. c. 15 = P. lanuginosum.

GARCILASSA, Poepp. & Endl. Nov. Gen. et Sp. iii. 45. t. 251 (1842). *COMPOSITAE*, Benth. & Hook. f. ii. 882.
rivularis, *Poepp. & Endl. l. c.* 46.—Peruv.

GARCINIA, Linn. Gen. ed. I. 343 (1737). *GUTTIFERAE*, Benth. & Hook. f. i. 174.
BIWALGIA, Scop. Introd. 223 (1777).
BRINDONIA, Thou. in Dict. Sc. Nat. v. 339 (1804).
CAMPOGIA, Linn. Fl. Mart. Burs. (1745); Diss. Diss. 18 (1747); Amoen. Acad. i. 171, 402 (1749).
CLUSIANTHEMUM, Vieill. in Bull. Soc. Linn. Normand. ix. (1866) 338.
CODDAMPULI, Adans. Fam. ii 445 (1763).
DISCOSTIGMA, Hask. in Flora, xxv. (1842) ii. Beibl. 33.
HEBRADENDRON, R. Grah. in Hook. Comp. Bot. Mag. ii. (1836) 199. t. 27.
MANGOSTANA, Gaertn. Fruct. ii. 105. t. 105 (1791).
ORYCARPUS, Lour. Fl. Cochinch. 647 (1790).
RHINOSTIGMA, Miq. Fl. Ind. Bat. Suppl. 495 (1860).
TERPNOPHYLLUM, Thw. in Hook. Kew Journ. vi. (1854) 70. t. 2 C.
acuminata, Hort. Bogor. ex Pierre, Fl. For. Cochinch. fasc. v. 36 = calophylla.
acuminata, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 355 = G. Morella.
acuminata, Wall. Cat. n. 4871 A = Ternstroemia penangiana (B = Anonac. sp.).
affinis, Wall. Cat. n. 4864 = anomala, cornea.
affinis, Wight & Arn. Prod. 101 = G. Cambogia.
amboinensis, Spreng. Syst. ii. 448 = cochinchinensis.
amplexicaulis, *Vieill. ex Pierre, Fl. For. Cochinch.* fasc. v. 20.—N. Caled.
Anderson, *Hook. f. ex T. Anders. in Hook. f. Fl. Brit. Ind.* i. 270.—Malacca.
anomala, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 329.—Ind. or.
apetala, *Pierre, Fl. For. Cochinch. fasc.* v. 35.—Borneo.
Arthurgordoni, Horne, ex Baker, in Journ. Linn. Soc. xx. (1883) 361 = Gardneria Gordoni.
atroviridis, *Griff. ex T. Anders. in Hook. f. Fl. Brit. Ind.* i. 269.—Assam.
australis, *Montr. in Mém. Acad. Lyon*, x. (1860) 188. —N. Caled.
Baillonii, *Pierre, Fl. For. Cochinch. fasc.* v. 18.—Borneo.
Balansae, *Pierre, l. c. fasc.* 'vi.' 7.—N. Caled.
balica, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 209.—Ins. Bali.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 494.—Malaya.
Barteri, *Oliver, Fl. Trop. Afr.* i. 166.—Afr. trop.

GARCINIA :—
beccariana, *Pierre, Fl. For. Cochinch.* t. 58.—Cochinch.
Beccarii, *Pierre, l. c. fasc.* v. 26.—Borneo.
Benthami, *Pierre, l. c.* t. 55.—Cochinch.
Blumeiconea, Roxb. ex Wall. Cat. n. 4858 = paniculata.
Binnendijkii, *Pierre, Fl. For. Cochinch. fasc.* v. 7.—Malaya.
Binucao, *Choisy, Gutlif. Ind.* 84.—Ind. or.
Blancoi, *Pierre, Fl. For. Cochinch. fasc.* v. 15.—As. or.
Blumei, *Pierre, l. c.* 32.—Cambodia.
Bobe-rona, Choisy, Guttif. Ind. 36 = paniculata.
Boerlagii, *Pierre, Fl. For. Cochinch. fasc.* v. 36.—Malaya.
borneensis, *Pierre, l. c.* 27.—Borneo.
brasiliensis, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 34.—Bras.
brevipes, *Pierre, Fl. For. Cochinch. fasc.* v. 8.—Borneo.
brevirostris, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 358 ; et in Flora, liii. (1870) 241 = eugeniifolia.
Calleryi, *Pierre, Fl. For. Cochinch. fasc.* v. 15.—Ins. Philipp.
calophylla, *Pierre, l. c.* 36.—Malacca ; Java ?
calycina, *Kurz, in Journ. Bot.* xiii. (1875) 324.—Ins. Nicobar.
Cambogia, *Desr. in Lam. Encyc.* iii. 701.—Ind. or.
Cambogia, Roxb. Hort. Beng. 42 ; Fl. Ind. ii. 621 = G. Cowa.
cambogioides, Royle, Mat. Med. Ind. ed. III. 339 = G. Morella.
celebica, Desr. in Lam. Encyc. iii. 700 = indica.
celebica, Linn. Syst. ed. XII. 926 = cornea.
Choisyiana, *Wall. Cat.* n. 4870.—Malaya.
cladostigma, *Pierre, Fl. For. Cochinch. fasc.* v. 30.—Java ?
cochinchinensis, *Choisy, in DC. Prod.* i. 561.—China.
collina, *Vieill. ex Panch. & Sebert, Not. Bois Nouv. Caléd.* 228.—N. Caled.
concarpa, Wight, Ic. t. 1. 121 ; Illustr. i. 126 = G. Cambogia.
covallina, *Vieill. in Bull. Soc. Linn. Normand.* ix. (1865) 388.—N. Caled.
cornea, *Linn. in Murr. Syst.* ed. XIII. 868.—Ind. or. ; Malaya.
cornea, Roxb. ex Sm. in Rees, Cyclop. xv. n. 5 = G. Cowa.
corymbosa, *Wall. Cat.* n. 4859.—Ind. or.
Cowa, *Roxb. Hort. Beng.* 42 ; *Fl. Ind.* ii. 622.—Ind. or.
Cumingiana, *Pierre, Fl. For. Cochinch. fasc.* v. 11.—Ins. Philipp.
cuneifolia, *Pierre, l. c.* 35.—Borneo.
cymulosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 208.—Malaya.
dasdalanthera, *Pierre, Fl. For. Cochinch. fasc.* v. 40.—Ins. Celebes.
Delpyana, *Pierre, l. c.* t. 65.—Cambodia.
densiflora, *Pierre, l. c. fasc.* v. 25.—N. Caled.
Descrousseauxii, *Pierre, l. c.* 34.—Borneo.
diolca, *Blume, Bijdr.* 215.—Java.
dioica, Herb. Lugd. Bot. ex Pierre, Fl. For. Cochinch. fasc. v. 26 = calophylla.
dinica, Sm. in Rees, Cycl. xv. n. 5 = G. Cowa.
diospyrifolia, *Pierre, Fl. For. Cochinch. fasc.* v. 37.—Borneo.
dives, *Pierre, l. c.* 35.—Ins. Philipp.
dryobalanoides, *Pierre, l. c.* 7.—Borneo.
dulcis, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 88.—Ins. Andaman.
duodecandra, *Pierre, Fl. For. Cochinch. fasc.* v. 28.—Ins. Philipp.
echinocarpa, *Thw. in Hook. Kew Journ.* vi. (1854) 71.—Zeylan.
elliptica, *Choisy, in DC. Prod.* i. 561.—Malaya.
elliptica, Wall. Cat. n. 4869 = G. Morella.
eugeniifolia, *Wall. l. c.* 4873.—Malaya.
fabrilis, *Miq. Fl. Ind. Bat. Suppl.* 496.—Sumatra.
fasciculoris, *Wall. Cat.* n. 4858.—Ind. or.
ferrea, *Pierre, Fl. For. Cochinch.* t. 57.—Cochinch.
floribunda, *Miq. Stip. Surin. Sal.* 39.—Guian.
fulva, *Pierre, Fl. For. Cochinch. fasc.* v. 6.—Malacca.
fusca, *Pierre, l. c.* t. 67.—Siam.
Ganalchaudii, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 327 = G. Morella.
Gerrardi, *Harv. Gen. S. Afr. Pl.* ed. II. 26.—Afr. austr.

GARCINIA :—
gracilis, *Pierre, Fl. For. Cochinch.* t. 63.—Cochinch.
Grahami, *Pierre, l. c. fasc.* v. 23.—Borneo.
grandifolia, *Pierre, l. c.* 7.—Malaya ?
Griffithii, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 266. —Malaya.
Gutta, Roxb. Hort. Beng. 42 ; Wall. Cat. n. 4866 = G. Cambogia.
Gutta, Wight, Illustr. i. 126 = G. Morella.
Hanburyi, *Hook. f. in Journ. Linn. Soc.* xiv. (1875) 485.—Malaya.
Harmandii, *Pierre, Fl. For. Cochinch.* t. 60.—Cambodia.
Hasskarlii, *Pierre, l. c. fasc.* v. 8.—Sumatra.
heterandra, *Wall. Cat.* n. 4856.—Burma.
Hombroniana, *Pierre, Fl. For. Cochinch. fasc.* v. 12.—Singapore.
Hookeri, *Pierre, l. c.* 27.—Malacca.
Horafieldiana, *Pierre, l. c.* 23.—Java.
huillensis, *Welw. in Oliver, Fl. Trop. Afr.* i. 167.—Afr. trop.
indica, *Choisy, in DC. Prod.* i. 561.—Ind. or.
javanica, *Blume, Bijdr.* 215.—Java.
javanica, Hort. Bogor. ex Pierre, Fl. For. Cochinch. fasc v. 30 = cladostigma.
Jawoera, *Pierre, l. c.* 37.—Java.
Jelinckii, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 118.—Ins. Nicobar.
Keenania, *Pierre, Fl. For. Cochinch. fasc.* v. 8.—Ind. or.
Kernaniana, Pierre, l. c. t. 92. f. G = praec.
Kingii, *Miq. Fl. Ind. Bat. Suppl.* 493.—Ins. Banca.
Kola, Heckel.—Cf. Bull. Soc. Bot. Fr. xxx. (1883) Bibl. 150 = Cola acuminata.
Korthalsii, *Pierre, Fl. For. Cochinch. fasc.* v. 39.—Borneo.
Krawang, *Pierre, l. c.* 37.—Borneo.
Kurzii, *Pierre, l. c.* 14.—Ins. Andaman.
Kydia, Roxb. Fl. Ind. ii. 623 = G. Cowa.
Kydiana, Roxb. Hort. Beng. 42 = G. Cowa.
Lampong, *Miq. Fl. Ind. Bat. Suppl.* 494.—Sumatra.
lanceaefolia, *Roxb. Hort. Beng.* 42 ; *Fl. Ind.* ii. 623. —Reg. Himal.
Lanessanii, *Pierre, Fl. For. Cochinch.* t. 70.—Cambodia.
laterifloru, Blume, Bijdr. 214 = G. Morella.
latissima, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 209.—Java.
leucandra, *Pierre, Fl. For. Cochinch. fasc.* v. 27.—Sumatra.
linearis, *Pierre, l. c.* 8.—Borneo.
Livingstonei, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 263. —Afr. trop.
lobulata, Wall. Cat. n. 4868 A = G. Cowa, G. Morella.
longifolia, *Blume, Bijdr.* 214.—Java.
loniceroides, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 264.—Ind. or.
Loureiri, *Pierre, Fl. For. Cochinch.* t. 66.—Cochinch.
lucens *Pierre, l. c. fasc.* v. 37.—Borneo.
macrophylla, T. Anders. ex Hook. f. Fl. Brit. Ind. i. 270 = Andersoni.
macrophylla, Mart. in Flora, xxiv. (1841) II. Beibl. 35 = Rheedii.
macrophylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 208. —Sumatra.
madagascariensis, *Baill. ex Pierre, Fl. For. Cochinch. fasc.* v. 5.—Madag.
Mahingayi, *Hook. f. ex T. Anders. in Hook. f. Fl. Brit. Ind.* i. 267.—Malaya.
malabarica, Desr. in Lam. Encyc. iii. 701 = Diospyros malabarica.
malaccensis, *Hook. f. ex T. Anders. in Hook. f. Fl. Brit. Ind.* i. 261.—Malacca.
Mangorua, *Linn. Sp. Pl.* 448.—Ins. Moluco.
Mannii, *Oliver, Fl. Trop. Afr.* i. 167.—Afr. trop.
Mansoniana, Klotzsch. ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 326 = cornea.
Mellori, *Baker, in Journ. Linn. Soc.* xx. (1883) 92.—Madag.
mergensis, *Wight, Ic.* t. 114 ; *Illustr.* i. 124.—Burma ; Malaya.
microcarpa, *Pierre, Fl. For. Cochinch. fasc.* v. 39.—Borneo.
microstigma, *Kurz, in Journ. Bot.* xiii. (1875) 324.—Ins. Nicobar.
minahassensis, *Pierre, Fl. For. Cochinch. fasc.* v. 29.—Ins. Celebes.
Miquelii, *Pierre, l. c.* 24.—Borneo.
monosperma, *Benj. & C. F. Schmidt, Offic. Gew.* iv. t. 26 d.—Ind. or.

GARCINIA :—
Morella, *Desr. in Lam. Encyc.* iii. 701.—Ind. or., Malaya.
Moselleyana, *Pierre, Fl. For. Cochinch. fasc.* v. 10.—Ins. Philipp.
Mottleyana, *Pierre, l. c.* 30.—Borneo.
multiflora, *Champ. ex Benth. in Hook. Kew Journ.* iii. Ins. (1851) 310.—Ins. Hongkong.
Mungotia, *Deplanch. ex Pierre, Fl. For. Cochinch. fasc.* v. 35.—N. Caled.
myristicifolia, *Pierre, l. c.* 26.—Borneo.
neglecta, *Vieill. in Bull. Soc. Linn. Normand* ix (1865) 387.—N. Caled.
nervosa, Miq. Ann. Mus. Bot. Lugd. Bat. i. 208 = Andersoni.
nigricana, *Pierre, Fl. For. Cochinch. fasc.* v. 30.—Borneo.
nigro-lineata, *Planch. ex T. Anders. in Hook. f. Fl. Brit. Ind.* i. 268.—Malacca.
nitida, *Pierre, Fl. For. Cochinch. fasc.* v. 18.—Borneo.
oblongifolia, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 311.—Ins. Hongkong.
Oliveri, *Pierre, Fl. For. Cochinch.* t. 64.—Cambodia.
ovalifolia, Hook. f. ex T. Anders. in Hook. f. Fl. Brit. Ind. i. 269 = spicata.
ovalifolia, *Oliver, Fl. Trop. Afr.* i. 166.—Afr. trop.
oxyedra, *Miq. Fl. Ind. Bat. Suppl.* 494.—Sumatra.
oxyphylla, *Miq. l. c.* 495.—Sumatra.
oxyphylla, Planch. & Triana, in Ann. Sc. Nat. Sér. IV xiv. (1860) 342 = lanceaefolia.
Pancheri, *Pierre, Fl. For. Cochinch. fasc.* v. 25.—N. Caled.
paniculata, *Roxb. Hort. Beng.* 42 ; *Fl. Ind.* ii. 626.—Reg. Himal.
Papilla, Wight, Ic. tt. 960, 961 = G. Cambogia.
parviflora, *Benth. in Hook. Lond. Journ. Bot.* ii (1843) 370.—Guian.
parvifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 208.—Malaya.
pauciflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 92 —Madag.
pedicellata, *Seem. Viti*, 433 ; *Fl. Vit.* 11.—Ins. Fiji.
pedunculata, *Roxb. Hort. Beng.* 43 ; *Fl. Ind.* ii. 625.—Reg. Himal.
penangiana, *Pierre, Fl. For. Cochinch. fasc.* v. 37.—Ins. Penang.
petiolaris, *Pierre, l. c.* 3.—Borneo.
pictorius, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 209.—Ins. Ceram.
pictoria, Roxb. Hort. Beng. 43 ; Fl. Ind. ii. 627 = G Morella.
Planchoni, *Pierre, Fl. For. Cochinch.* t. 61.—Cochinch.
polyantha, *Oliver, Fl. Trop. Afr.* i. 166.—Afr. trop.
porrecta, Laness. Mem. Gen. Garcin. 21 = cornea.
Pseudo-guttifera, *Seem. Fl. Vit.* 11.—Ins. Fiji.
punctata, *Oliver, Fl. Trop. Afr.* i. 167.—Afr. trop.
purpurea, Roxb. Hort. Beng. 42 ; Fl. Ind. ii. 624 = indica.
purpurea, Wall. Cat. n. 4862 = lanceaefolia.
quadrifaria, *Baill. ex Pierre, Fl. For. Cochinch. fasc.* v. 4.—Afr. trop.
quaesita, *Pierre, l. c.* 22.—Zeylan.
Rheedii, *Pierre, l. c.* 34.—Borneo.
Rheediana, Choisy, Guttif. Ind. 35 = paniculata.
Riedeliana, *Pierre, Fl. For. Cochinch. fasc.* v. 13.—Ins. Celebes.
rigida, *Miq. Fl. Ind. Bat. Suppl.* 493.—Sumatra.
rostrata, *Hassk. ex Hook. f. in Journ. Linn. Soc.* xiv. (1875) 485.—Borneo ; Malaya.
Roxburghii, Wight, Illustr. i. 125 = G. Cowa.
Rumphii, *Pierre, Fl. For. Cochinch. fasc.* v. 18.—Ins. Banca.
salakensis, *Pierre, l. c.* 36.—Java.
sarawhensis, *Pierre, l. c.* 6.—Borneo.
Schefferi, *Pierre, l. c.* t. 59.—Cambodia.
Schomburgkiana, *Pierre, l. c. fasc.* v. 33.—Guian.
sessilis, *Seem. Viti*, 433 ; *Fl. Vit.* 10.—Ins. Fiji.
sisygiifolia, *Pierre, Fl. For. Cochinch. fasc.* v. 26.—Borneo.
Smeathmanni, *Oliver, Fl. Trop. Afr.* i. 168.—Afr. trop.
speciosa, *Wall. Pl. As. Rar.* iii. 87 ; *Cat.* n. 4855.—Burma.
spectabilis, Hook. f. Fl. Brit. Ind. i. 261, sphalm. = speciosa.
spectabilis, *Pierre, Fl. For. Cochinch. fasc.* v. 3.—Borneo.
spicata, *Hook. f. in Journ. Linn. Soc.* xiv. (1875) 486. —Malaya.

GARCINIA:—
stigmacantha, *Pierre, Fl. For. Cochinch. fasc.* v. 29.—Borneo.
stipulata, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 267.—Reg. Himal.
subtilinervia, *F. Muell. Papuan Pl.* i. 85.—N. Guinea.
succifolia, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 208.—Burma.
sulcata, Stokes, Bot. Mat. Med. iii. 11 = G. Cambogia.
terpnophylla, *Thw. Enum. Pl. Zeyl.* 406.—Zeylan.
tetrandra, *Pierre, Fl. For. Cochinch. fasc.* v. 20.—Ins. Philipp.
Teysmanniana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 7.—N. Guinea.
Thorelii, *Pierre, Fl. For. Cochinch.* t. 62.—Cambodia.
Thwaitesii, *Pierre, l. c. fasc.* v. 4.—Zeylan.
timorensis, *Zipp. ic.* 24; *ex Span. in Linnaea*, xv. (1841) 178.—Ins. Timor.
tranrcorica, *Bedd. Fl. Sylv.* t. 178.—Ind. or.
trechostigma, *Pierre, Fl. For. Cochinch. fasc.* v. 31.—Malaya.
Treubii, *Pierre, l. c.* 7.—Sumatra.
Trianii, *Pierre, l. c.* 18.—Borneo.
umbellifera, Wall. Cat. n. 4864 = G. Cowa.
vemulose, *Choisy, Guttif. Ind.* 24.—Ind. or.
Vieillardii, *Pierre, Fl. For. Cochinch. fasc.* v. 12.—N. Caled.
Vilersiana, *Pierre, in Bull. Soc. Linn. Par.* i. (1882) 348; *Fl. For. Cochinch.* t. 71.—Siam.
vitiensis, *Seem. Fl. Vit.* 10.—Ins. Fiji.
Wallichiana, Klotzsch, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 256 = G. Morella.
Wallichii, *Choisy, in Mém. Soc. Phys. Genève* xii. (1851) 417 = G. Cowa.
Wightii, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 265.—Ind. or.
Xanthochymus, *Hook f. ex T. Anders. l. c.* 269.—Ind. or.; Malaya.
xylosperma, *Pierre, Fl. For. Cochinch. fasc.* v. 39.—Ins. Celebes.
zeylanica, Roxb. Hort. Beng. [92]: Fl. Ind. ii. 621 = G. Cambogia.

GARDENIA, Adans. Fam. ii. p. (20) = seq.

GARDENIA, Ellis, in Phil. Trans. li. ii. (1761) 935.
RUBIACEÆ, Benth. & Hook. f. ii 89.
BERGKIAS, Sonner. Voy. 48. t. 17 (1776).
CAQUIPIRIA, J. F. Gmel. Syst. 651 (1791).
HYPERACANTHUS, E. Mey. in Drège, Zwei Pfl. Docum. 198 (1843).
PIRINGA, Juss. in Mém. Mus. Par. vi. (1820) 899.
ROTHMANNIA, Thunb. in Vet. Acad. Handl. Stockh. 65 (1776), cum ic.
SALNHERGIA, Neck. Elem. i. 262 (1790).
YANGAPA, Rafin. Sylva Tellur. 20 (1838).
Abbreviatæ, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 104.—Afr. trop.
aculeata, [Soland. in] Ait. Hort. Kew. ed. I. i. 295 = Randia aculeata.
acuminata, G. Don, Gen. Syst. iii. 499 = Randia acuminata.
acuminata, *Korth. in Nederl. Kruidk. Arch.* ii. ii. (1851) 199.—Borneo.
africana, *Regel, Cat. Pl. Hort. Absakov.* 62, nomen (Quid?).—Afr. trop.
amœna, *Sims, Bot. Mag.* t. 1904.—China.
angustifolia, *Ledd. Bot. Cab.* t. 813.—Ind. or.?
anisophylla, Jack, in Roxb. Fl. Ind. ed. Carey, ii. 461 = Randia anisophylla.
anisophylla, *Wall. Cat.* n. 8399.—Ind. or.
Annæ, *E. P. Wright, in Trans. Roy. Irish Acad.* xxiv. (1871) 875.—Ins. Seychell.; Afr. trop.
appendiculata, Stokes, Bot. Mat. Med. i. 494 = G. Thembergia.
arborea, Roxb. Hort. Beng. 15; Fl. Ind. i. 708 = gummifera.
armata, Bartl. ex DC. Prod. iv. 387 = Randia tetracantha.
armata, Sw. Prod. Veg. Ind. Occ. 51 = Randia armata.
arnensis, *Monstr. in Mém. Acad. Lyon*, x. (1860) 215.—N. Caled.
assimilis, *Afzel. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 103.—Afr. trop.
atrosperma, *Regel, Cat. Pl. Hort. Absakov.* 63, nomen (Quid?).—Hab.?

GARDENIA:—
Aubryi, *Viaill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 65.—N. Caled.
australis, *Regel, Cat. Pl Hort. Absakov.* 62, nomen (Quid?).—Hab.?
Benthamiana, F. Muell. Fragm. ix. 180 = Randia Benthamiana.
bignoniæflora, Welw. Apont. 585. n. 13 = Amaralia bignoniæflora.
bispinosa, *Regel, Cat. Pl. Hort. Absakov.* 62, nomen (Quid?).—Hab.?
Blumeana, DC. Prod. iv. 388 = campanulata.
brasiliensis, *Spreng. Syst.* i. 763.—Bras.
Brighamii, *H. Mann, in Proc. Am. Acad.* vii. (1867) 171.—Ins. Sandvic.
buffalina, Poir. Encyc. Suppl. ii. 708 = Genipa buffalina.
bullata, *Vell. Fl. Flum.* 103; ill. t. 12.—Bras.
Burka, Buch.-Ham. in Wall. Cat. n. 8298 = Wendlandia paniculata.
calycina, G. Don, Gen. Syst. iii. 497 = Sherbournia follosa.
calyculata, Roxb. Hort. Beng. 85; Fl. Ind. i. 704 = latifolia?
campanulata, *Roxb. l. c.* 15, 710.—Reg. Himal.; Burma.
carinata, Griff. Notul. iv. 261 = coronaria.
carinata, Thw. Enum. Pl. Zeyl. 159 = coronaria?
carinata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 560.—Malaya.
chartacea, *F. Muell. Pl. Fitsalan.* 12.—Austral.
chinensis, *Carey, ex Voigt, Hort. Suburb. Calc.* 379 (= seq. ?).—China.
chinensis, Spreng. Syst. i. 762 = Randia sinensis.
citriodora, Hook. Bot. Mag. t. 4987 = Mitrostigma axillare.
clusiæfolia, Jacq. Coll. App. 87. t. 4. f. 3 = Genipa clusiifolia.
coccinea, G. Don, in Edinb. Phil. Journ. xi. (1824) 343 = Mussænda elegans.
concolor, D. Dietr. Syn. Pl. i. 797 (= *Gardeniola concolor*).—Bras.
cornifolia, *DC. Prod. iv.* 384.—N. Granat.
coronaria, *Buch.-Ham. in Syme, Embassy Ava*, iii. ed. II. 307.—Burma.
corymbosa, Keichb. Hort. Bot. ii. 2. t. 106 = Olostyla corymbosa.
costata, Roxb. Hort. Beng. 15; Fl. Ind. i. 704 = coronaria.
crassicaulis, Salisb. Parad. Lond. t. 45 = G. Thunbergia.
cribita, *Afzel. Stirp. Guin. Medic. Sp. Nov.* (1829) 18. n. 5.—Afr. trop.
cruciata, *Regel, Cat. Pl. Hort. Absakov.* 62, nomen (Quid?).—Hab.?
cuneata, R. Br. in Wall. Cat. n. 8263 = turgida.
cuneata, *Kurz, For. Fl. Brit. Burma*, ii. 41.—Burma.
curvata, *Tréjem. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 404.—Ins. Bantam.
daphnoides, Hance, in Walp. Ann. ii. 796 = Diplospora viridiflora.
dasycarpa, *Kurz, For. Fl. Brit. Burma*, ii. 42.—Burma.
dasycarpa, Kurz, in Joam As. Soc. Beng. xli. (1872) ii. 219 = Randia tomentosa.
densa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 559 = Randia tetrasperma.
Devoniana, *Lindl. in Gard. Chron.* (1846) 663.—Afr. trop.
Dewea, Buch.-Ham. in Wall. Cat. n. 8292 = turgida.
dumetorum, Retz. Obs. ii. 14 = Randia dumetorum.
dumosa, Salisb. Prod. 65 = pasc.
edulis, *Monstr. in Mém. Acad. Lyon*, x. (1860) 216.—N. Caled.
edulis, *F. Muell. Fragm.* i. 54.—Austral.
edulis, Poir. Encyc. Suppl. ii. 708 = Alibertia edulis.
edulis, *Wall. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 65.—N. Caled.
enneandra, Koen. ex Roxb. Fl. Ind. ed. Carey, ii. 552 = latifolia?
erythrocladæ, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 211.—Burma.
esculenta, Stokes, Bot. Mat. Med. i. 495; Poir. Encyc. Suppl. ii. 708 = Randia edulis.
fagifolia, *Willd. ex Roem. & Schult. Syst.* v. 245.—Am. austr.
fasciculata, Roxb. Hort. Beng. 15; Spreng. Syst. iv. Cur. Post. 84 = Randia fasciculata.
fernandensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 105.—Afr. trop.

GARDENIA:—
ferox, Cham. & Schlecht. in Linnæa, iv. (1829) 198 = Randia ferox.
ferrea, *Vell. Fl. Flum.* 103; ill. t. 113.—Bras.
Fitzalani, *F. Muell. Rep. Burdt. Exp.* 12.—Austral.
flava, Poir. Encyc. Suppl. ii. 708 = Genipa flava.
floribunda, Roxb. Hort. Beng. 15 = Randia dumetorum.
florida, Linn. Sp. Pl. ed. II. 305 = jasminoides.
follosiana, Herb. Madr. ex Wight & Arn. Prod. 399 = Randia latifolia.
formosa, *Cham. & Schlecht. in Linnæa*, iv. (1829) 200.—Bras.
Forsteniana, *Meq. Ann. Mus. Bot. Lugd. Bat.* iv. 236.—Ins Celebes.
fragrans, Roxb. Pl. Corom. ii. 19. t. 187 = Randia malabarica.
frondosa, Lam. Encyc. ii. 608 = Mussænda frondosa.
fucata, *R. Br. ex Benth. Fl. Austral.* iii. 410.—Austral.
Genipa, Sw. Prod. Veg. Ind. Occ. 51 = Genipa americana.
Gerrardiana, *Hare. & Sond. Fl. Cap.* iii. 6.—Afr. austr.
glabra, R. Br. in Wall. Cat. n. 8258 = Randia dumetorum.
globosa, *Hockst. in Flora*, xxv. (1842) 287.—Afr. austr.
glutinosa, Teljsm. & Binn. Cat. Hort. Bog. 119 = tubifera.
Gordoni, *Baker, in Journ. Linn. Soc.* xx. (1883) 361.—Ins. Fiji.
Gorrii, *Horne, ex Baker, l. c.* 362.—Ins. Fiji.
grandiflora, *Laur. Fl. Cochinch.* 147.—Cochinch.
grandiflora, Siebold, ex Zucc. Fl. Jap. Fam. Nat. ii. 55. n. 604 = jasminoides.
grandis, *Korth. in Nederl. Kruidk. Arch.* ii. ii. (1851) 192.—Borneo.
Grievei, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 361.—Ins. Fiji.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 118.—Malacca.
gummifera, *Linn. f. Suppl.* 164.—Ind. or.
heterocalyta, Koen. ex Blume, Mus. Bot. Lugd. Bat. i. 164 = Fagræa cellanica.
heterophylla, *Montr. in Mém. Acad. Lyon*, x. (1860) 216.—N. Caled.
hexagona, *Lém. Illustr. Hortic.* xvi. (1869) *Misc.* 92.—Bras.
hexandra, *Willd. in Roem. & Schult. Syst.* v. 243.—Am. austr.
Hillii, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 362.—Ins. Fiji.
hirta, *F. Muell. Fragm.* vii. 46.—Austral.
horrida, Spreng. Syst. i. 762 = Randia horrida.
Humboldtiana, Roem. & Schult. Syst. v. 243 = Randia Humboldtiana.
hygrophila, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 212.—Burma.
inermis, F. G. Dietr. Vollst. Lexik. Gaertn. iv. 285 = gummifera.
ixoræfolia, *R. Br. in Wall. Cat.* n. 8262.—Burma, jamaicensis, *Spreng. Syst.* i. 761.—Jamaic.
Jardinei, *F. Muell. ex Benth. Fl. Austral.* iii. 410.—Austral.
jasminiflora, D. Dietr. Syn. Pl. i. 797 = Heinsia jasminiflora.
jasminiflora, Zipp. ic. t. 4; ex Span. in Linnæa, xv. (1841) 815 = Randia maculata.
jasminoides, *Ellis, in Phil. Trans.* li. ii. (1761) 935.—China.
jasminoides, Retz. Obs. ii. 13 = scandens.
Jovis-tonantis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 101.—Afr. trop.
Kalbreyeri, *Hiern, in Journ. Bot.* xvi. (1878) 97.—Afr. trop.
kongumensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 104.—Afr. trop.
latifolia, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 110 = Randia dumetorum.
latifolia, *[Soland. in] Ait. Hort. Kew.* ed. I. i. 294.—ined. or.
Leschenaultii, D. Dietr. Syn. Pl. i. 797 = Plectronia Leschenaultii.
longiflora, [Dryand. in] Ait. Hort. Kew. ed. II. i. 368 = Randia macrantha.
longiflora, Rulz & Pav. Fl. Per. ii. 67 = Randia Ruiziana.
longiflola, *G. Don, Gen. Syst.* iii. 499.—Afr. trop.

GARDENIA:—
longipina, Roxb. Hort. Beng. 15, nomen = Randia dumetorum.
longispina, Wall. Cat. n. 8277 D = campanulata, Randia tetrasperma.
longistyla, Hook. Bot. Mag. t. 4322 = Macrosphyra longistyla.
lucens, *Panch. & Sebert, Not. Bois Nouv. Calédon.* 182. —N. Caled.
lucida, *Roxb. Hort. Beng.* 15; *Fl. Ind.* i. 707.—Ind. or.; Burma.
lutea, *Fresen. in Mus. Senckenb.* ii. (1837) 167.—Abyssin.
Macgillivrsei, *Benth. Fl. Austral.* iii. 409.—Austral.
macrantha, Roem. & Schult. Syst. v. 237 = Randia macrantha.
macrocarpa, Carey, ex Voigt, Hort. Suburb. Calc. 379 = G. Thunbergia.
macroptera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 286. —Borneo.
madagascariensis, Lam. Encyc. ii. 608 = Randia madagascariensis.
malleifera, Hook. Bot. Mag. t. 4807 = Randia malleifera.
Manganjae, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 102.—Afr. trop.
maritima, Vahl, ex DC. Prod. iv. 388 = Randia Mussaendae.
Maruba, Siebold, in Blume, Bijdr. 1015 = jasminoides.
medicinalis, Vahl, ex Schum. & Thonn. Beskr. Guin. Pl. 148 = G. Thunbergia.
megasperma, *F. Muell. Fragm.* i. 54.—Austral.
Merianae, Poir. Encyc. Suppl. ii. 708 = Genipa Merianae.
micrantha, Thunb. Diss. Gard. 17 = Randia parviflora.
microcarpa, *Bartl. ex DC. Prod.* iv. 384.—Ins. Philipp.
microcarpa, Hochst. in Flora, xxv. (1842) 237 = Randia rudis.
mitis, Steud. Nom. ed. I. 856 = Randia latifolia.
montana, Roxb. Hort. Beng. 85; Fl. Ind. i. 709 = turgida.
multiflora, Willd. Sp. Pl. i. 1231 = Randia longiflora.
Mussaenda, Linn. f. Suppl. 168 = Randia Mussaenda.
mutabilis, *Reinw. ex Blume, Bijdr.* 1016.—Ins. Celebes.
Nexum-papata, Roxb. ex Wight & Arn. Prod. 428 = Plectronia didyma.
Neuberia, *Eckl. & Zeyh. Enum.* 360.—Afr. austr.
nitida, *Hook. Bot. Mag.* t. 4348.—Afr. trop.
nutans, Roxb. Hort. Beng. 15; Wall. Cat. n. 8290 = Randia dumetorum.
oblongifolia, Poir. Encyc. Suppl. ii. 708 = Genipa oblongifolia.
obovata, F. G. Dietr. Vollst. Lexik. Nachtr. iii. 441 = Randia obovata.
obovata, Spreng. Syst. i. 761 = Randia pubescens.
obscura, *Vidal, Phan. Cuming. Philipp.* 18, 119.—Ins. Philipp.
obtusifolia, *Roxb. Hort. Beng.* 15, nomen.—Burma.
ochracea, *F. Muell. Fragm.* i. 55.—Austral.
octomera, Hook. Bot. Mag. t. 5410 = Randia octomera.
oxycarpa, R. Br. in Wall. Cat. n. 8261 = sessiliflora.
palmosa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 241.—Afr. trop.
pallida, *Lindl. Bot. Reg.* t. 1952.—Am. trop.
Pandacaki, J. F. Gmel. Syst. 434 = Randia malabarica.
Pandaki, Vahl, ex DC. Prod. iv. 377 = prsec.
pannea, *Lindl. Bot. Reg.* t. 1952.—Am. austr.
parviflora, H. B. & K. Nov. Gen. et Sp. iii. 408. t. 293 = Randia tetrandra.
parviflora, Poir. Encyc. ii. 708 = Plectronia didyma.
parviflora, Smeathm. ex DC. Prod. iv. 393 = Pouchetia africana.
parviflora, *Vell. Fl. Flum.* 108; iii. t. 14.—Bras.
parviflora, Wall. Cat. n. 8256 = Randia fasciculata.
parvifolia, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 441 = Catesbaea parvifolia.
patula, Horsf. ex Willd. in Roem. & Schult. Syst. v. 244 = Randia macrantha.
Pavetta, Roxb. ex Wight & Arn. Prod. 401 = Stylocoryne Webera.
pentagonioides, *Sonn. Fl. Vit.* 122.—Ins. Fiji.
picturum, Hassk. in Flora, xxvii. (1844) 284 = jasminoides.
platixylon, *Vieill. ex Panch. & Sebert, Not. Bois Nouv. Calédon.* 183.—N. Caled.
pomifera, Wall. Cat. n. 8296 = Randia uliginosa.
propinqua, R. Br. in Wall. l. c. 8260 = Randia dumetorum.

GARDENIA:—
propinqua, *Lindl. Bot. Reg.* t. 975.—Ind. or.!
Psendopsidium, *Blanco, Fl. Filip.* ed. I. 497.—Ins. Philipp.
pubescens, Roth, Nov. Pl. Sp. 151 (non Gardenia).—Ind. or.
pulchella, G. Don, in Edinb. Phil. Journ. xi. (1824) 343 = Heinsia jasminiflora.
pulcherrima, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 312 = Randia exaltata.
pyriformis, *A. Cunn. ex Benth. Fl. Austral.* iii. 409. —Austral.
racemulosa, *Korth. in Nederl. Kruidk. Arch.* ii. ii. (1851) 192.—Borneo.
radicans, Thunb. Fl. Jap. 109. t. 20; Diss. Gard. 12. t. 1. f. 1 = jasminoides.
Randia, Sw. Prod. Veg. Ind. Occ. 51 = Randia aculeata.
Reinwardtiana, *Blume, Bijdr.* 1013.—Malaya.
Remyi, *H. Mann, in Proc. Am. Acad.* vii. (1867) 171. —Ins. Sandvic.
resinifera, Korth. in Nederl. Kruidk. Arch. ii. ii. (1851) 191 = tubifera.
resinifera, Roth, Nov. Pl. Sp. 150 = lucida.
resinifua, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 102.—Afr. trop.
resinosa, *F. Muell. Fragm.* i. 54.—Austral.
rigida, Buch.-Ham. in D. Don, Prod. Fl. Nep. 188 (= *Meteoraria rigida*).—Nepal.
rigida, Wall. Cat. n. 8257 = Plectronia parviflora.
rigida, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 570 = Randia fasciculata.
Rothmannia, *Linn. f. Suppl.* 165.—Afr. austr.
rotunda, D. Dietr. Syn. Pl. i. 797 (= *Scopsoathamnus rotunda*).—Bras.
rotundifolia, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 422 = Randia rotundifolia.
Sagraeana, A. Rich. Fl. Cub. Fanerog. ii. 10 = Scolosanthus crucifer.
scandens, *Thunb. Diss. Gard.* 17. t. 2. f. 5.—China.
Schiedeana, *Cham. ex Steud. Nom.* ed. II. i. 688 (Quid ?).—Mexic.
Schomburgkii, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 408.—Java; Ins. Batam.
Sellowiana, *Cham. & Schlecht. in Linnaea*, iv. (1829) 137.—Bras.
sessiliflora, *Wall. Cat.* n. 8291; *C. B. Clarke, in Journ. Linn. Soc.* xvii. (1878) 161.—Burma.
sessilis, F. Muell. Fragm. vii. 47 = Randia sessilis.
sessilis, *Vell. Fl. Flum.* 109; iii. t. 11.—Bras.
Sherbournise, Hook. Bot. Mag. t. 4044 = Sherbournia follosa.
Somersetii, Spreng. Syst. i. 762 = Randia parviflora.
speciosa, *Hook. f. Fl. Brit. Ind.* iii. 117.—Singapore.
speciosa, Roxb. ex Wight & Arn. Prod. 422 = Goettardia speciosa.
speciosa, Salisb. Prod. 68 = G. Thunbergia.
spinosa, Billb. ex Benzl. in Vet. Akad. Handl. Stockh. 1854 (1856) 132 = Randia nitida.
spinosa, Linn. f. Suppl. 164 = Randia dumetorum.
spinosa, Thunb. Diss. Gard. 16 = Randia fasciculata, dumetorum.
Stanleyana, Hook. ex Lindl. Bot. Reg. (1845) t. 47; Hook. Bot. Mag. t. 4185 = Randia maculata.
stellaris, Salisb. Prod. 64 = Randia latifolia.
stipularis, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 211 = Randia dumetorum.
stipulosa, F. Muell. Fragm. vii. 47 = Randia stipulosa.
stipulosa, *Zoll. & Mor. Syst. Verz. Zoll.* 60.—Java.
Storckii, *Oliver, in Hook. Ic. Pl.* t. 1448.—Ins. Fiji.
suaveolens, *Vell. Fl. Flum.* 102; iii. t. 9.—Bras.
suavis, Wall. Cat. n. 8274 = obtusifolia.
suffruticosa, *R. Br. ex Benth. Fl. Austral.* iii. 410.—Austral.
sulcata, *Gaertn. f. Fruct.* iii. 79. t. 194.—Hab.!
taitensis, DC. Prod iv. 380 = Randia tahitensis!
tenuiculata, *Hook. f. Fl. Brit. Ind.* iii. 119.—Malacca.
ternifolia, Schum. & Thonn. Beskr. Guin. Pl. 147 = G. Thunbergia.
tetracantha, Lam. Encyc. ii. 609 = Randia armata.
tetracantha, Willd. ex Roem. & Schult. Syst. v. 244 = Randia Humboldtiana.
tetrandra, Bedd. Ic. Pl. Ind. Or. i. 4. t. 90 = Byrsophyllum tetrandrum.
tetrasperma, Roxb. Hort. Beng. 85; Fl. Ind. i. 709 = Randia tetrasperma.
Thunbergia, *Linn. f. Suppl.* 162.—Afr. austr.
Tinneana, Kotschy & Peyritsch, in Bot. Zeit. xxiii. (1865) 178 = G. Thunbergia.

GARDENIA:—
tomentosa, Blume, ex DC. Prod. iv. 379 = Randia tomentosa.
triacantha, DC. l. c. 382 = G. Thunbergia.
trichoclada, *Miq. Fl. Ind. Bat.* ii. 231.—Malaya.
trimeris, Vell. Fl. Flum. 109; iii. t. 10 = Strychnos trijuinervia.
tubifera, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 562.—Malaya.
tubiflora, Andr. Bot. Rep. t. 188 = Oxyanthus tubiflorus.
turgida, *Roxb. Hort. Beng.* 15; *Fl. Ind.* i. 711.—Ind. or.; Burma.
uliginosa, Retz. Obs. ii. 14 = Randia uliginosa.
urceliformis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 104. —Afr. trop.
Urvillei, *Montr. in Mém. Acad. Lyon*, x. (1860) 214. —N. Caled.
verticillata, Lam. Encyc. ii. 607 = G. Thunbergia.
vinosa, D. Dietr. Syn. Pl. i. 797 (= *Scopsoathamnus rotunda*).—Bras.
vitiensis, *Seem. in Bonplandia*, ix. (1861) 256; x. (1862) 86; *Fl. Vit.* 122.—Ins. Pacif.
Vogelii, *Hook. f. in Hook. Ic. Pl.* t. 793-3.—Afr. trop.
volubilis, *Lour. Fl. Cochinch.* 148.—China.
Whitfieldii, Lindl. Bot. Reg. (1845) sub t. 47 = Randia malleifera.
zanguebarica, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 105. —Afr. trop.

GARDENIOLA, Cham. in Linnaea, ix. (1834) 247 = **Allbertia**, A. Rich. (Rubiac.).
concolor, Cham. l. c.—Bras.

GARDENIOPSIS, Miq. in Ann. Mus. Bot. Lugd. Bat. iv. 250, 262 (1868). *RUBIACEAE*, Benth. & Hook. f. ii. 116.
longifolia, *Miq.* l. c.—Sumatra.

GARDINIA, Bert. in Merc. Chil. (1829).—Cf. Bull. Férussac, xx. (1830) 111; et Am. Journ. Sc. xxiii. (1833) 81 = **Brodiaea**, Sm. (Liliac.).
purpurascens, Bert. ll. cc. (= *Milla violacea*).—Chili.
violacea, Bert. ex Steud. Nom. ed. II. i. 664 = prsec.

GARDNERA, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 318 (1824) = seq.

GARDNERIA, Wall. ex Roxb. Fl. Ind. ed Carey, i. 400 (1820). *LOGANIACEAE*, Benth. & Hook. f. ii. 798.
angustifolia, *Wall. l. c.* 318; *et Cat.* n. 817.—Reg. Himal.; Japon.
fragraeacea, F. Muell. Fragm. vi. 180 = Fagraea Muelleri.
glabra, Wall. ex D. Don, Prod. Fl. Nep. 132 = angustifolia.
nutans, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 165 = angustifolia.
ovata, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 400; *Cat.* n. 816.—Ind. or.
Wallichii, Wight, ex Wall. Pl. As. Rar. iii. 49. t. 281 = ovata.

GARDOQUIA, Ruiz & Pav. Prod. 86. t. 17 (1794). *LABIATAE*, Benth. & Hook. f. ii. 1189.
DIONEILIS, Raf. Fl. Tellur. iii. 85 (1836).
RASTENBOCKIA, Endl. l. c. 84 (1836).
RIZOA, Cav. in Anal. Cienc. Nat. iii. (1800) 138.
acutifolia, *Benth. in DC. Prod.* xii. 256.—Peruv.
affinis, *Benth. in Linnaea*, xi. (1837) 343, nn. 11080, 11083.—Hab.!
argentea, *H. B. & K. Nov. Gen. et Sp.* ii. 313.—Peruv.
betonicoides, Lindl. Bot. Reg. (1838) Misc. 86 = Cedronella mexicana.
breviflora, *Benth. Lab. Gen. et Sp.* 401.—Peruv.
capitata, Spreng. Syst. iv. Cur. Post. 228 = Camila fasciculata.
chilensis, Benth. in Hook. & Arn. Bot. Beech. Voy. 58 = Cilliesii.
discolor, *H. B. & K. Nov. Gen. et Sp.* ii. 313.—N. Granat.
elegans, *H. B. & K.* l. c. 315.—Peruv.
elliptica, *Ruiz & Pav. Syst. Veg.* 149.—Peruv.
fasciculata, *Benth. Pl. Hartw.* 243.—N. Granat.
foliolosa, *Benth. in DC. Prod.* xii. 256.—N. Granat.

GARDOQUIA :—
Gilliesii, *K. Crok. in Edinb. N. Phil. Journ.* (July-Sept. 1881) 277.—Chili.
glabrata, *H. B. & K. Nov. Gen. et Sp.* ii. 213.—Ecuador.
grandiflora, *H. B. & K. l. c.* 214.—Ecuador.
Hellerî, *Peyr. in Linnaea*, xxx. (1859–60) 34.—Mexic.
Hookeri, Benth. Lab. Gen. et Sp. 401 = Calaminthe coccinea.
incana, Herb. Willd. ex Benth. in Linnæa, xi. (1837) 331 = grandiflora.
incana, *Ruiz & Pav. Syst. Veg.* 148.—Peruv.
Jamesoni, *Benth. Lab. Gen. et Sp.* 404.—Peruv.
mexicana, *Benth. Pl. Hartw.* 50.—Mexic.
micromerioides, *Hemsl. Biol. Centr. Am. Bot.* ii. 550.—Mexic.
microphylla, *H. B. & K. Nov. Gen. et Sp.* ii. 211.—Ecuador.
multiflora, *Ruiz & Pav. Syst. Veg.* 149.—Chili.
obovata, *Ruiz & Pav. l. c.* 150.—Peruv.
obovata, Spreng. ex Benth. in DC. Prod. xii. 289 = Brunfelsia ramosissima.
origanoides, Reichb. ex Benth. l. c. (Verbenaces.)—Ins. Trinit.
pilosa, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 341.—Ins. Fiji.
pulchella, *H. B. & K. Nov. Gen. et Sp.* ii. 215.—Ecuador.
quitensis, Herb. Willd. ex Benth. in Linnæa, xi. (1837) 332 = grandiflora.
racemosa, Spreng. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 170 = Hedeoma multiflora.
revoluta, *Ruiz & Pav. Syst. Veg.* 149.—Peruv.
rugosa, *Benth. Lab. Gen. et Sp.* 399.—Peruv.
salvaefolia, *Colla, in Mem. Acc. Torin.* xxxix. (1836) 2. t. 48.—Chili.
sericea, *Presl, ex Benth. Lab. Gen. et Sp.* 402.—Peruv.
speciosa, *Regel, Cat. Pl. Hort. Absankov.* 62.—Hab. ?
spicata, Spreng. Syst. iv. Cur. Post. 225 = Cunila spicata.
striata, *Ruiz & Pav. Syst. Veg.* 148.—Peruv.
tarifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 212.—N. Granat.
thymoides, *H. B. & K. l. c.* 214.—Ecuador.
tomentosa, *H. B. & K. l. c.*—Ecuador.

GAREILASSA, Walp. Rep. ii. 601 (1843), sphalm. = Garellassa, Poepp. & Endl (Compos.).

GARHADIOLUS, Jaub. & Spach, Illustr. iii. 119. t. 234, 235 (1849) = **Rhagadiolus**, Juss. (Compos.).
angulosus, Jaub. & Spach, l. c. 120 = R. Hedypnois.
hamosus, Boiss. & Hausskn. ex Boiss. Fl. Orient. iii. 753.—Syria.
Hedypnois, Jaub. & Spach, Illustr. iii. 120 = R. Hedypnois.
papposus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 135.—Armen. ; Afghan.

GARIDELIA, Spreng. Anleit. ii. 11. 735 (1818) = seq.

GARIDELLA, Tourn. ex Linn. Gen. ed. I. 131 (1737) = **Nigella**, Linn. (Ranunc.).
anethifolia, Salisb. Prod. 374 = N. Nigellastrum.
nigellæformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 138 = seq.
Nigellastrum, Linn. Sp. Pl. 425 = N. Nigellastrum.
nigelliformis, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 103 = præc.
segetalis, Lam. Illustr. ii. 500. [t. 279. f. 2]—Syria.

GARNIERIA, Brongn. & Gris, in Bull. Soc. Bot. Fr. xviii. (1871) 189. *PROTEACEÆ*, Benth. & Hook. f. iii. 176.
spathulaefolia, *Brongn. & Gris, l. c.*—N. Caled.

GARNOTIA, Brongn. in Duperr. Voy. Coq. Bot. 132. t. 21 (1829). *GRAMINEÆ*, Benth. & Hook. f. iii. 1118.
Bergiana, Endl. Gen. Suppl. iii. 87 (1843).
Miquelia, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) (Pl. Meyen.) 177.
coerulescens, *Thw. Enum. Pl. Zeyl.* 363.—Ind. or.
dryuda, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1863) 223.—China.
fusoria, *Thw. Enum. Pl. Zeyl.* 363.—Zeylan.
miquelia, *Thw. l. c.*—Zeylan.

GARNOTIA :—
patula, *Munro, ex Benth. Fl. Hongk.* 416.—China.
polypogonoides, *Munro, ex Oliver, in Hook. Ic. Pl.* t. 1481.—Ind. or.
scoparia, *Thw. Enum. Pl. Zeyl.* 363.—Zeylan.
stricta, *Brongn. in Duperr. Voy. Coq. Bot.* 133. t. 21.—Ins. Hongkong.

GAROSMUS, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. viii. (1748) 211, App. (Quid ?).

GARRETIA, Welw. Apont. 587 (1858). *MELIACEÆ*.
anthotoca, Welw. l. c.—Afr. trop.

GARRIELIA, Gandich. Bot. Voy. Bonite, t. 115 (1844–52) = **Dyckia**, Schult. (Bromeliac.).
enchelioides, Gandich. l. c. = D. altissima.

GARRYA, Dougl. ex Lindl. Bot. Reg. t. 1686 (1834). *CORNACEÆ*, Benth. & Hook. f. i. 951.
Fadyenia, Endl. Gen. Suppl. ii. 80 (1842).
buxifolia, *A. Gray, in Proc. Am. Acad.* vii. (1868) 349.—Calif.
elliptica, *Dougl. ex Lindl. Bot. Reg.* t. 1686.—Am. bor. occ.
Fadyena, *Hook. Ic. Pl.* t. 333.—Jamaic.
flavescens, *S. Wats. in Am. Natural.* vii. (1873) 301.—Am. bor. occ.
Fremontii, *Torr. in Pacif. Rail. Rep.* iv. 136.—Am. bor. occ.
laurifolia, *Benth. Pl. Hartw.* 14.—Mexic.
Lindheimeri, *Torr. in Pacif. Rail. Rep.* iv. 136.—Texas.
Lindleyana, Hartw. ex A. DC. in DC. Prod. xvi. 1. 487 = laurifolia.
Lindleyana, *A. Murr. Oreg. Circ.* 1.—Am. bor. occ.
macrophylla, *Benth. Pl. Hartw.* 50.—Mexic.
oblonga, *Benth. l. c.* 51.—Mexic.
ovata, *Benth. l. c.* 14.—Mexic.
Veatchii, *Kellogg, in Proc. Calif. Acad.* v. (1873) 40.—Calif.
Wrightii, *Torr. in Pacif. Rail. Rep.* iv. 136.—Am. bor. occ.

GARUGA, Roxb. Hort. Beng. 33 (1814) ; Pl. Corom. iii. 5. t. 208 (1819) ; Fl. Ind. ii. 400 (1832). *BURSERACEÆ*, Benth. & Hook. f. i. 323.
Kunthii, Dennst. Schlüssl. Hort. Malab. 30 (1818).
floribunda, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 177.—Ins. Timor.
Pegantra, Engl. in Mart. Fl. Bras. xii. ii. 288 = Thyrsodium giganteum.
guianensis, Engl. l. c. = Thyrsodium guianense.
javanica, *Blume, Bijdr.* 1165.—Java.
Kenghar, Buch.-Ham. in Trans. Linn. Soc. xvii. (1835) 191 = pinnata.
Khasiana, *Buch.-Ham. l. c.*—Ind. or.
madagascariensis, *DC. Prod.* ii. 81.—Madag.
mollis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 475.—Ins. Philipp.
Pyarhad, Buch.-Ham. in Trans. Linn. Soc. xvii. (1835) 191 = pinnata.
pinnata, *Roxb. Hort. Beng.* 33 ; *Pl. Corom.* iii. 5. t. 208 ; *Fl. Ind.* ii. 400.—Ind. or. ; Malaya.
Schomburgkiana, Engl. in Mart. Fl. Bras. xii. ii. 287 = Thyrsodium Schomburgkianum.
Spruceana, Engl. l. c. 286 = Thyrsodium Spruceanum.

GARUGANDRA, Griseb. in Goett. Abh. xxiv. (1879) 96. *ANACARDIACEÆ*.
amorphoides, *Griseb. l. c.*—Reg. Argent.

GARULIUM, Cass. in Bull. Soc. Philom. (1819) 172 ; et in Dict. Sc. Nat. xviii. 169 (1820). *COMPOSITÆ*, Benth. & Hook. f. ii. 263.
album, *S. Moore, in Journ. Bot.* xvi. (1878) 183.—Afr. austr.
bipinnatum, *Less. Syn. Comp.* 194.—Afr. austr.
corniculum, Steud. Nom. ed. II. ii. 386, in syn. = viscosum.
latifolium, *Harv. in Harv. & Sond. Fl. Cap.* iii. 93.—Afr. austr.
pinnatifidum, DC. Prod. v. 309 = viscosum.
viscosum, *Cass. in Bull. Soc. Philom.* (1819) 172 ; *et in Dict. Sc. Nat.* xviii. 168.—Afr. austr.

GARULIUM, Bartl. Ord. 143 (1830) = præc.

GARUMHIUM, Blume, in Flora, viii. (1825) t. 103 err. typ. = CARUMBIUM, Reinw. = **Homalanthus**, A. Juss. (Euphorb.).

GASLONDIA, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 96. *MYRTACEÆ*, Benth. & Hook. f. i. 1006.
amphoricarpa, *Vieill. l. c.* 97.—N. Caled.

GASOUB, Adans. Fam. ii. 249 (1763) = **Mesembryanthemum**, Linn. (Ficoid.).

GASPARINIA, Bertol. Fl. Ital. iii. 614 (1839) = **Sfilaus**, Benth. (Umbell.).
virescens, Bertol. l. c. 615 = S. carvifolius.

GASPARINIA, Endl. Gen. 1408 (1841) = **Aeginetia**, Linn. (Orobanch.).
mirabilis, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 7 = A. Centronia.

GASSOLOMA, D. Dietr. Syn. Pl. ii. 793 (1840) = **Geissoloma**, Lindl. (Penæac.).

GASTERANTHOPSIS, Cerst. in Vidensk. Selsk. Skr. V. v. (1861) 129 (Gesn. Centr. Am. 55. t. 10. f. 1) = **Besleria**, Linn. (Gesneric.).
glabra, Oerst. ll. cc. = B. glabra.
hirsuta, Oerst. ll. cc. = B. hirsuta.

GASTERANTHUS, Benth. Pl. Hartw. 233 (1846) = **Besleria**, Plum. (Gesner.).
pendulus, Hanst. in Linnæa, xxvi. (1853) 216. t. 2. f. 58 = B. pendula.
pilosus, Benth. Pl. Hartw. 233 = B. calcarata.
quitensis, Benth. l. c. = B. quitensis.

GASTERIA, Duval, Pl. Succ. Hort. Alenç. 6 (1809) ; Haw. Syn. Pl. Succ. 85 (1812). *LILIACEÆ*, Benth. & Hook. f. iii. 775.
acinacifolia, *Haw. Suppl. Pl. Succ.* 49.—Afr. austr.
angustifolia, Haw. l. c. = disticha.
apicroides, *Baker, in Journ. Linn. Soc.* xviii. (1880) 197.—Afr. austr.
Bayfieldi, *Baker, l. c.*—Afr. austr.
bicolor, *Haw. in Phil. Mag.* (1826) 275.—Afr. austr.
brevifolia, *Haw. Syn. Pl. Succ.* 89.—Afr. austr.
candicans, *Haw. in Phil. Mag.* (1827) 360.—Afr. austr.
carinata, *Duval, Pl. Succ. Hort. Alenc.* 6 ; *Haw. Syn. Pl. Succ.* 87.—Afr. austr.
cheilophylla, *Baker, in Journ. Linn. Soc.* xviii. (1880) 199.—Afr. austr.
colubrina, *N. E. Br. in Gard. Chron.* (1877) ii. 88.—Afr. austr.
conspurcata, Haw. in Phil. Mag. (1827) 352 = disticha.
crenatifolia, Haw. l. c. 350 = nigricans.
Croscheri, *Baker, in Journ. Linn. Soc.* xviii. (1880) 196.—Afr. austr.
decipiens, *Haw. in Phil. Mag.* (1827) 356.—Afr. austr.
denticulata, Haw. Suppl. Pl. Succ. 50 = disticha.
dicta, *N. E. Br. in Gard. Chron.* (1876) ii. 68.—Afr. austr.
disticha, *Haw. in Phil. Mag.* (1827) 351.—Afr. austr.
ensifolia, Haw. l. c. (1825) 382 = acinacifolia.
excavata, *Haw. l. c.* (1827) 358.—Afr. austr.
excelsa, *Baker, in Journ. Linn. Soc.* xviii. (1880) 195.—Afr. austr.
fasciata, Haw. in Phil. Mag. (1827) 349 = nigricans.
formosa, Haw. l. c. = picta.
fuscopunctata, *Baker, in Journ. Linn. Soc.* xviii. (1880) 195.—Afr. austr.
glabra, *Haw. Syn. Pl. Succ.* 87.—Afr. austr.
gracilis, *Baker, in Journ. Linn. Soc.* xviii. (1880) 193.—Afr. austr.
intermedia, Haw. Syn. Pl. Succ. 89 = verrucosa.
lactepunctata, *Haw. in Phil. Mag.* (1827) 356.—Afr. austr.
lævis, Haw. l. c. 354 = Aloë lævis.
latifolia, Haw. Syn. Pl. Succ. 87 = exasperata.
latifolia, Hort. Kew. ex Baker, in Journ. Linn. Soc. xviii. (1880) 188 = nigricans.
Lingua, Link, Handb. i. 191 = Aloë Lingua, *Willd. limita*, Haw. in Phil. Mag. (1827) 360 = candicans.
longifolia, Duval, Pl. Succ. Hort. Alenç. 6 ; Haw. Syn. Pl. Succ. 88 = disticha.

GASTERIA:—
maculata, *Haw. in Phil. Mag.* (1827) 348.—Afr. austr.
marmorata, *Baker, in Journ. Linn. Soc.* xvii. (1880) 194.—Hab.?
marmorata, Haw. in Phil. Mag. (1827) 350 = Aloë formosa.
mollis, *Haw. Rev. Pl. Succ* 46, 203.—Afr. austr.
nigricans, *Duval, Pl. Succ. Hort. Alenç.* 6; *Haw. Syn. Pl. Succ.* 86.—Afr. austr.
nitens, *Haw. Suppl. Pl. Succ.* 48 = acinaciformis.
nitida, *Haw. in Phil. Mag.* (1827) 358.—Afr. austr.
obliqua, Duval, Pl. Succ. Hort. Alenç. 6; Haw. Syn. Pl. Succ. 85 = maculata.
obtusa, *Haw. in Phil. Mag.* (1827) 358.—Afr. austr.
obtusifolia, *Haw. l. c.* 350.—Afr. austr.
pallescens, *Baker, in Journ. Linn. Soc.* xviii. (1880) 190.—Afr. austr.
parva, Haw. in Phil. Mag. (1827) 35 = carinata.
parvifolia, *Baker, in Journ. Linn. Soc.* xviii. (1880) 193.—Afr. austr.
Peacockii, *Baker, l. c.* 195.—Afr. austr.
pethamensis ×, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 193.
picta, *Haw. in Phil. Mag.* (1827) 348.—Afr. austr.
planifolia, *Baker, in Journ. Linn. Soc.* xviii. (1880) 188.—Afr. austr.
platyphylla, Baker, l. c. 186 = nigricans.
pluripunctata, Haw. in Phil. Mag. (1827) 359 = acinaciifolia.
porphyrophylla, *Baker, in Journ. Linn. Soc.* xviii. (1880) 190.—Afr. austr.
pulchra, *Haw. Syn. Pl. Succ.* 86.—Afr. austr.
repens, *Haw. Rev. Pl. Succ.* 48.—Afr. austr.
retusa, *Haw. in Phil. Mag.* (1827) 350.—Afr. austr.
retusifolia, Haw. l. c. 351, in note = obtusifolia.
spiralis, *Baker, in Journ. Linn. Soc.* xviii. (1880) 189.—Afr. austr.
squarrosa, *Baker, l. c.* 197.—Afr. austr.
strigata, Haw. in Phil. Mag. (1827) 356 = carinata.
strigosa, Hort. Peacock, ex Baker, in Journ. Linn. Soc. xviii. (1880) 193 = subcarinata.
subcarinata, *Haw. Suppl. Pl. Succ.* 49; *et in Phil. Mag.* (1827) 357.—Afr. austr.
subnigricans, Haw. in Phil. Mag. (1827) 358 = nigricans.
subverrucosa, *Haw. l. c.* 354.—Afr. austr.
sulcata, *Haw. l. c.* 353.—Afr. austr.
trigona, *Haw. l. c.* 358.—Afr. austr.
undata, Haw. l. c. 357 = subcarinata.
variolosa, *Baker, in Saund. Refug. Bot.* t. 347.—Afr. austr.
venusta, Haw. in Phil. Mag. (1827) 359 = acinaciifolia.
verrucosa, *Duval, Pl. Succ. Hort. Alenç.* 6; *Haw. Syn. Pl. Succ.* 89.—Afr. austr.
Zeyheri, *Baker, in Journ. Linn. Soc.* xviii. (1880) 190.—Afr. austr.

GASTEROLYCHNIS, Rupr. in Beitr. Pfl. Russ. Reich. ii. 24 (1845) = **Lychnis**, Tourn. (Caryophyll.).
alpiflora, Rupr. l. c. vii. 30 = L. apetala †
angustiflora, Rupr. l. c. ii. 24.—Reg. Ural.
involucrata, Rupr. l. c.—Reg. Ural.
macropetala, Rupr. l. c. vii. 30 = L. apetala †
tristis, Rupr. l. c. 31 = L. tristis.
uralensis, Rupr. l. c. 30.—Reg. Ural.
Vahlii, Rupr. l. c. ii. 24 = L. alpina.

GASTERONEMA, Ldd. ex Stend. Nom. ed. II. i. 665 (1840) = GASTRONEMA, Herb. = Cyrtanthus, Ait. (Liliac.).
pallidum, Ldd. ex Stend. l. c. = C. pallidus.

GASTONIA, Comm. ex Lam. Encyc. ii. 610 (1786); Juss. Gen. 217 (1789). *ARALIACEAE*, Benth. & Hook. f. i. 941.
aculeata, Hort. ex Seem. in Journ. Bot. ii. (1864) 246 = Euphocarpus aculeatus.
Cumingiana, Hort. ex Seem. l. c. = Brassaiopsis speciosa.
cutispongia, *Lam. Encyc.* ii. 610.—Ins. Maurit.
denteta, Hort. ex Seem. in Journ. Bot. ii. (1864) 246 = Brassaiopsis speciosa.
duplicata, *Thou. ex Baill. Adansonia*, xii. (1878) 166.—Madag.
emirnensis, *Baker, in Journ. Linn. Soc.* xxi. (1884) 350. —Madag.
capterohides, *Trijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 410.—Malaya.

GASTONIA:—
lleptapleurum, *Baill. Adansonia*, xii. (1878) 166.—Madag.
longifolia, Hort. ex Seem. in Journ. Bot. ii. (1864) 246 = Brassaiopsis speciosa.
Naluga, Lam. Encyc. ii. 611 = Lees sambucina.
Naluga, Spreng. Syst. ii. 440 = præc.
oahuensis, A. Gray, Bot. U. St. Expl. Exped. i. 726 = Triplasandra oahuensis.
palmata, Roxb. Hort. Beng. 33; Fl. Ind. ii. 407 = Trevesia palmata.
paniculata, D. Dietr. Syn. Pl. ii. 1628 = Polyscias paniculata.
papuana, *Miq. Ann. Mus. Bot Lugd. Bat.* i. 5.—N. Guin.
repanda, D. Dietr. Syn. Pl. ii. 1628 = Polyscias repanda.
saurauioides, Roeper, in Bot. Zeit. vi. (1848) 257. t. 1. —Cf. Seem. Heder. 103 (Polyaciadis sp.).—Hab.?
saurauioides, Hook. Hort. Beng. 90; Fl. Ind. ii. 408 = Trevesia palmata.
simplicifolia, Seem. in Journ. Bot. iii. (1865) 75 = Osmoxylon amboinense.
spongicus, Pers. Syn. ii. 28 = cutispongia.
sundaica, Baill. Hist. Pl. vii. 161 = Trevesia sundaica.

GASTRANTHUS, Moritz, ex Benth. & Hook. f. Gen. ii. 1107 (1876). *ACANTHACEAE*, Benth. & Hook. f. i. c.
Schiechtendalii, *Moritz, i. c.; et in Hook. Ic. Pl.* t. 1210. —Venezuela.

GASTRANTHUS, F. Muell. Fragm. vi. 130 (1868) = **Parsonsia**, K. Br. (Apocyn.).
ventricosa, F. Muell. l. c. = P. ventricosa.

GASTRIDIUM, Beauv. Agrost. 21. t. 6. f. 6 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1161.
australe, *Beauv. l. c.*—Europ.; Reg. Mediterr.
laxum, Boiss. & Reut. Pugill. Pl. Nov. 126 = australe.
lendigerum, Gaud. Fl. Helv. i. 176 = australe.
littorale, Dur. ex Parl. Fl. Ital. i. 178 = Triplachne nitens.
muticum, *Guenth. in Spreng. Nov. Prov.* 90.—Reg. Mediterr.
nitens, Coss. & Dur. ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 229 = Triplachne nitens.
scabrum, Presl, Fl. Sicul. p. xliv, nomen = muticum.
siculum, Tin. Gram. Unifl. 169 = muticum.
triaristatum, Dur. in Duch. Rev. Bot. ii. (1846) 424 = Triplachne nitens.
ventricosum, Spreng. Neue Entdeck. ii. 36 = australe.

GASTRIDIUM, Blume, Fl. Jav. Praef. p. vii (1828) = GRASTIDIUM, Blume = **Dendrobium**, Sw. (Orchid.).

GASTRILIA, Rafin. Fl. Tellur. iv. 105 (1836) = **Thymelaea**, Endl. (Thymel.).
geminata, Rafin. l. c. = T. dioica.
umbellata, Rafin. l. c. = Daphnopsis Swartzii.

GASTROCARPHA, D. Don, in Trans. Linn. Soc. xvi. (1830) 281 = **Moscharia**, Ruiz & Pav. (Compos.).
runcinata, D. Don, l. c. 282 = M. pinnatifida.

GASTROCHILUS, D. Don, Prod. Fl. Nep. 32 (1825) = **Sacocalabium**, Blume (Orchid.).
calceolaria, D. Don, l. c. = S. calceolare.

GASTROCHILUS, Wall. Pl. As. Rar. i. 22. t. 24, 25 (1829). *SCITAMINEAE*, Benth. & Hook. f. iii. 642.
affinis, *Wall. ex Voigt, Hort. Suburb. Calc.* 572, nomen. —Assam.
Yenhiasii, Wall. l. c. = longiflora.
longiflora, *Wall. Pl. As. Rar.* i. 22. t. 25.—Reg. Himal.; Malaya.
parvula, *Wall. Cat.* n. 6590.—Burma.
pulcherrima, *Wall. Pl. As. Rar.* i. 22. t. 24.—Burma; Penang.

GASTROCOPTYLIS, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 406 (Not. Lehm. 239). *BORAGINEAE*, Benth. & Hook. f. ii. 855.
hispida, *Bunge, li. cc.*—Reg. Aral.

GASTRODIA, R. Br. Prod. 330 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 517.
EPIPHANES, Blume, Bijdr. 421. t. 4 (1825).
GAMOPLEXIS, Falc. in Trans. Linn. Soc. xx. (1847) 298. t. 18.
antennifera, Blume, Mus. Bot. Lugd. Bat. ii. 174 = Cunninghamii.
Cunninghamii, *Hook. f. Fl. N. Zel.* i. 251.—N. Zel.
elata, *Blume, Mus. Bot. Lugd. Bat.* ii. 174.—Reg. Himal.; Ins. or.
gracilis, *Blume, l. c.*—Japon.
Hasseltii, *Blume, l. c.* 176.—Malaya.
javanica, *Endl. Gen.* 312.—Java.
orobanchoides, F. Muell. Phyt. New Hebr. 22 = Cunninghamii?
pallens, F. Muell. l. c. = Didymoplexis pallens.
sesamoides, *Benth. & Hook. f. Gen.* iii. 617.—Reg. Himal.
sesamoides, *R. Br. Prod.* 33.—Austral.
verrucosa, *Blume, Mus. Bot. Lugd. Bat.* ii. 175.—Sumatra.

GASTROGLOTTIS, Blume, Bijdr. 397 (1825) = **Liparis**, Rich. (Orchid.).
montana, Blume, l. c.—Java.

GASTROLOBIUM, R. Br. in Ait. Hort. Kew. ed. II. iii. 18 (1811). *LEGUMINOSAE*, Benth. & Hook. f. i. 470.
acutum, Benth. in Lindl. Swan River App. 14 = Oxylobium acutum.
axillare, Meisn. in Bot. Zeit. xiii. (1855) 29 = Oxylobium reticulatum.
bidens, *Meisn. l. c.*—Austral.
bilobum, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iii. 16.—Austral.
Brownii, *Meisn. in Lehm. Pl. Preiss.* i. 71.—Austral.
Callistachys, *Meisn. l. c.* ii. 216.—Austral.
calycinum, *Benth. in Lindl. Swan River App.* 13.—Austral.
corallinum, Benth. l. c. = Oxylobium spectabile.
corymbosum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 272 = bilobum.
crassifolium, *Benth. Fl. Austral.* ii. 105.—Austral.
crenulatum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 273 = pyramidale.
cuneatum, Henfr. in Gard. Comp. i. (1855) 49 = velutinum?
Drummondii, Meisn. in Lehm. Pl. Preiss. i. 69 = oxylobioides.
alachistum, *F. Muell. Fragm.* ix. 67.—Austral.
emarginatum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 273 = velutinum.
epacridioides, *Meisn. in Lehm. Pl. Preiss.* i. 72.—Austral.
grandiflorum, *F. Muell. Fragm.* iii. 17.—Austral.
hamulosum, *Meisn. in Lehm. Pl. Preiss.* ii. 218.—Austral.
Henfreyi, Lem. Jard. Fleur. iii. (1853) in nota ad t. 224 = ovalifolium.
Hookeri, Meisn. in Lehm. Pl. Preiss. i. 71 = Brownii.
Huegelii, Henfr. in Moore & Ayres, Gard. Mag. Bot. i. (1850) 161 = Pultenæa ternata.
lilioifolium, *Meisn. in Lehm. Pl. Preiss.* i. 67.—Austral.
Lehmanni, *Meisn. l. c.* 70.—Austral.
lineare, Meisn. in Bot. Zeit. xiii. (1855) 30 = G. Callistachys.
microcarpum, *Meisn. in Lehm. Pl. Preiss.* i. 70.—Austral.
obovatum, *Benth. in Lindl. Swan River App.* 14.—Austral.
ovalifolium, *Henfr. in Gardn. Comp.* i. (1852) 41.—Austral.
ovalifolium, Lem. Jard. Fleur. iii. (1853) t. 294 = pyramidale.
oxylobioides, *Benth. in Lindl. Swan River App.* 13. —Austral.
parvifolium, *Benth. l. c.*—Austral.
pilosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 1. 274.—Austral.
polycephalum, Turcz. l. c. = pyramidale.
polystachyum, *Meisn. in Lehm. Pl. Preiss.* ii. 217.—Austral.
Preissii, Meisn. l. c. i. 68 = spinosum.
pulchellum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 1. 274.—Austral.
pycnostachyum, *Benth. Fl. Australi.* ii. 103.—Austral.

GASTROLOBIUM :—
pyramidale, *T. Moore, in Proc. Linn. Soc.* ii. (1852) 202.—Austral.
reticulatum, *Benth. Fl. Austral.* ii. 99.—Austral.
retusum, Lindl. Bot. Reg. t. 1647 = Oxylobium retusum.
rotundifolium, *Meissn. in Lehm. Pl. Preiss.* ii. 216.—Austral.
scorzifolium, *F. Muell. Fragm.* x. 35.—Austral.
spathulatum, *Benth. in Lindl. Swan River App.* 14.—Austral.
spinosum, *Benth. l. c.* 13.—Austral.
stenophyllum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 1. 275.—Austral.
stipulare, *Meisan. in Lehm. Pl. Preiss.* ii. 218.—Austral.
tricuspidatum, *Meissn. l. c.* i. 66.—Austral.
trilobum, *Benth. in Lindl. Swan River App.* 14.—Austral.
truncatum, *Benth. Fl. Austral* ii. 99 —Austral.
velutinum, *Lindl. & Paxt. Flow. Gard.* iii. (1852-53) 76. f. 270.—Austral.
verticillatum, Mœnch in Bot. Zeit. xiii. (1855) 25 = ilicifolium.
villosum, *Benth. in Lindl. Swan River App.* 13.—Austral.

GASTROLYCHNIS, Fenzl, ex Reichb. Nom. 206 (1841) = **Lychnis**, Tourn. (Caryophyll.).

GASTROMERIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. sub t. 75 (1830) = **Melasma**, Berg. (Scrophularin.).
phyzaloides, D. Don, l. c. = M. hispidum.

GASTRONEMA, Herb. App. 30 (1821) = **Cyrtanthus**, Ait. (Liliac.).
clavatum, Herb. l. c. = C. uniflorus.
helictum, M. Roem. Syn. Ensat. 52 = C. helictus.
pallidum, Lodd. Bot. Cab. t. 1808 = C. pallidus.
sanguineum, Lindl. in Journ. Hort. Soc. iii. (1848) 315 = sanguineus.
spirale, Eckl. & Zeyh. ex Kunth, Enum. Pl. v. 537 = C. spiralis.
uniflorum, Sallsb. Gen. Pl. Fragm. 189 = C. uniflorus.

GASTROPODIUM, Lindl. in Ann. Nat. Hist. Ser. I. xv. (1845) 107 = **Diothonea**, Lindl. (Orchid.).
violaceum, Lindl. l. c.—N. Granat.

GASTROSTYLUM, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 277, in nota (1845?) = seq.

GASTROSULUM, Sch. Bip. Tanacet. 29 (1844) = **Matricaria**, Linn. (Compos.).
annuum, Sch. Bip. l. c. 31.—Afr. austr.
dichotomum, Sch. Bip. l. c. = M. dichotoma.
praecox, Sch. Bip. l. c. 30 = M. praecox.
tenellum, Sch. Bip. l. c. = M. tenella.

GATESIA, Bertol. Misc. Bot. vii. 30. t. 1 (1843) = **Petalostemon**, Michx. (Legumin.).
alabamensis, Bertol. l. c. = P. corymbosus.

GATESIA, A. Gray, in Proc. Am. Acad. xiii. (1878) 365. *ACANTHACEAE.*
lasteviens, *A. Gray. l. c.*—Am. bor.

GATTENHOFFIA, Neck. Elem. i. 29 (1790) = **Dimorphotheca**, Moench (Compos.).

GATTENHOFIA, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 418 = **Iris**, Tourn.
verna, Medic. l. c. = I. verna.

GATYONA, Cass. in Bull. Soc. Philom. (1816) 168; et in Dict. Sc. Nat. xviii. 184 (1830) = **Crepis**, Vaill. (Compos.).
Bergeri, Sch. Bip. in Flora, xxii. (1839) 21 = C. Bergeri.
Dioscoridis, Dum. Fl. Belg. 60; Reichb. Fl. Germ. Excurs. 264 = C. Dioscoridis.
globuliflora, Cass. in Dict. Sc. Nat. xviii. 185 = C. Dioscoridis.
leiocarpa, Sch. Bip. in Flora, xxii. (1839) 21 = C. leiocarpa.
Pseudochondr. Vis. in Mem. Ist. Venet. xxi. (1860-62) 667.—Hungr.

GAUDICHAUDIA, H. B. & K. Nov. Gen. et Sp. v. 156. t. 445 (1821). *MALPIGHIACEAE*, Benth. & Hook. f. i. 261.
TRITOMOPTERIS. Juss ex Steud. Nom. ed. II. i. 665 (1840). In syn.
acuminata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 597.—Mexic.
albida, *Chaus. & Schlecht. in Linnaea*, v. (1830) 217.—Mexic.
Arnottiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 594.—Mexic.
brevipes, *A. Juss. l. c.* 597.—Mexic.
congestiflora, *A. Juss. l. c.* 590.—Mexic.
cynanchoides, *H. B. & K. Nov. Gen. et Sp.* v. 156. t. 455.—Mexic.
discolor, Griseb. in Linnaea, xiii. (1839) 187 = Camarea discolor.
Enrico-Martinezii, *Bárcena, Desc. Nuev. Pl. Mexic.* (1878) 4.—Mexic.
filipendula, *A. Juss. in Arch. Mus. Par* iii. (1843) 594.—Mexic.
guaranitica, St. Hil. in Bull. Soc Philom. (1823) 132 = Janusia guarnitica.
humilis, Benth. Pl. Hartw. 6 = Aspicarpa Hartwegiana.
Karwinskiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 592.—Mexic.
linearifolia, St. Hil. in Bull. Soc. Philom. (1823) 132 = Camarea junces.
mollis, *Benth. Pl. Hartw.* 6 —Mexic.
mucronata, A. Juss. in Arch. Mus. Par. iii. (1843) 596 = filicifolia.
oxyota, *A. Juss. l. c.*—Mexic.
pentandra, *A. Juss. l. c.* 598.—Mexic.
podocarpa, *A. Juss. l. c.* 596.—Mexic.
Schiedeana, *A. Juss. l. c.* 591.—Mexic.
sericea, St. Hil. in Bull. Soc. Philom. (1823) 132 = Camarea pulchella.
suffruticosa, Mohl, ex A. Juss. in Arch. Mus. Par. iii. (1843) 597 = Camarea filicifolia.
triphylla, Mohl, ex A. Juss. l. c. = Camarea triphylla.
Webbiana, *A. Juss. l. c.* 595.—Mexic.

GAUDINA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 172 = seq.

GAUDINIA, Beauv. Agrost. 95 (1812), partim. *GRAMINEAE*, Benth. & Hook. f iii. 1160.
ARTHROSTACHYA, Link, Hort. Berol. i. 151 (1827).
CYLICHNIUM, Delac, Fl. Hautes-Pyr. 68 (1867).
FALIMIRIA, Bess. ex Reichb. Consp. 54 (1828).
avenacea, Beauv. Agrost. Expl. Planches, 13 = fragilis.
Blehnetrinii, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 67 = Ventenata macra.
fragilis, *Beauv. Agrost.* 95.—Reg. Mediterr.
fragilis, Steuds, ex Ledeb. Fl. Ross. iv. 416 = Ventenata macra.
geminiflora, *Y. Gay, ex Kunth, Enum. Pl.* i. 308, in syn.—Hab. ?
planiculmis, Beauv. Agrost. 95 = Avena planiculmis.
tenuis, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 67 = Ventenata avenacea.

GAUDINIA, J. Gay, in Linnæa, xvii. (1829) 412 = **Linnæum**, Linn. (Ficoid.).
diffusa, J. Gay, l. c. = L. linifforum.
portulacifl., J. Gay, l. c. = L. linifforum.
viscosa, J. Gay, l. c. = L. viscosum.

GAULTERIA, Adans. Fam. ii. 165 (1763) = seq.

GAULTHERIA, Kalm, ex Linn. Diss. Chen. 20 (1751); in Amoen. Acad. iii. 14 (1756). *ERICACEAE*, Benth. & Hook. f. ii. 582.
BROSAEA, [Plum.] Linn. Gen. ed. I. 336 (1737).
SHALLONIUM, Rafin. in Am. Monthly Mag. (1818) 266.
Acosta, *Stud. Nom.* ed. II. i. 665.—Cochinch.
acuminata, *Chem. & Schlecht, in Linnaea*, v. (1830) 196.—Mexic.
adenothrix, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 44.—Japon.
alpina, Mart. ex Meisn. in Mart. Fl. Bras. vii. 150 = Pernettya myrtilloides.
anastomosans, *H. B. & K. Nov. Gen. et Sp.* iii. 285.—Am. austr.
antarctica, Hook. f. Fl. Antarct. ii. t. 116 = microphylla.
antipoda, *Forst. f. Prod.* 34. n. 196.—Tasman.; N. Zel.
bandongensis, Zoll. ex Miq. Fl. Ind. Bat. ii. 1055 = leucocarpa.
brachybotrys, *DC. Prod.* vii. 595.—Peruv.
bractesta, *G. Don, Gen. Syst.* iii. 840.—Ecuador.
buxifolia, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 422.—Venezuela.
cæspitosa, *Poepp. & Endl. Nov. Gen. et Sp.* i. 25. t. 37.—Chili.
candida, *DC. Prod.* vii. 597.—Bras.
ciliata, Cham. & Schlecht. in Linnæa, v. (1830) 126 = Pernettya ciliaris.
coccinea, *H. B. & K. Nov. Gen. et Sp.* iii. 284.—Venezuela.
Colensoi, Hook. Fl. N. Zel. i. 162 = rupestris.
conferta, *Benth. Pl. Hartw.* 219.—N. Granat.
cordata, Hort. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 312 = Maclenia cordata.
cordata, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 540 = hirtiflora.
cordifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 285. t. 261.—N. Granat.
cordifolia, Kœusch. Nom. ed. III 124 = Epigæa cordifolia.
crenulata, *Kurz, in Journ. Bot.* xi. (1873) 295.—Ind. or.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 184.—Ins. Philipp.
dependens, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 267 = antipoda.
discolor, Nutt. ex Hook. Bot. Mag. t. 5084 = Diplycosia discolor.
elliptica, *Cham. & Schlecht. in Linnæa*, viii. (1833) 502.—Bras.
epigeoides, *Domb. ex Wedd. Chlor. And.* ii. 173.—Ecuador.
erecta, Banks & Soland. ex Hook. f. Fl. N. Zel. 161 = antipoda.
erecta, *Vent. Hort. Cels.* t. 5.—Peruv.
eriophylla, Mart. ex Meisn. in Mart. Fl. Bras. vii. 151 = ferruginea.
fagifolia, *Hook f. Fl. N. Zel.* i. 162.—N. Zel.
ferruginea, *Cham. & Schlecht. in Linnæa*, i. (1826) 524.—Bras.
fluviatilis, A. Cunn. ex DC. Prod. vii. 594 = antipoda.
foliolosa, *Benth. Pl. Hartw.* 143.—Peruv.
formosa, *Remy, in Ann. Sc. Nat. Sér. III.* viii. (1847) 231.—Chili.
fragrans, D. Don, Prod. Fl. Nep. 151 = fragrantissima.
fragrantissima, *Wall. in As. Res.* xiii. (1820) 397.—Ind. or.; Malaya.
furiens, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 261 = Arbutus furiens.
glabra, *DC. Prod.* vii. 596.—Peruv.
glaucifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 281.—Mexic.
Griffithiana, *Wight, in. Calc. Journ. Nat. Hist.* viii. (1847) 176.—Reg. Himal.
heterophylla, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 9 = Diplycosia heterophylla.
hirsuta, Gardn. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 458 = fragrantissima.
hirtula, Mart. & Gal. in Bull. Acad. Belg. ix. (1842) 540 = Pernettya pilosa.
hirtiflora, *Benth. Pl. Hartw.* 66.—Mexic.
hispida, *R. Br. Prod.* 558.—Austral.; N. Zel.
hispida, Spreng. Syst. ii. 388 = Gaylussacia Pseudo-Gaultheria.
hispidula, Mohl. Cat. 44 = Chiogenes serpyllifolia.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 458.—Reg. Himal.
humilis, Sallsb. Prod. 389 = procumbens.
ignescens, Lem. Jard. Fleur. iii. (1852-53) t. 265 = ferruginea.
insipida, *Benth. Pl. Hartw.* 225.—Mexic.
Itatiaiae, Wawra, in Oestr. Bot. Zeitschr. xxxi. (1881) 380 = Agarista Itatiaiae.
laevigata, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 541 = nitida.
lanceolata, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 267.—Tasman.
lanigera, *Hook. Ic. Pl.* t. 66.—Am. austr.
latifolia, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 9 = Diplycosia latifolia.

GAULTHERIA:—
Leschenaultii, DC. Prod. vii. 593 = fragrantissima.
leucocarpa, *Blume, Bijdr.* 856.—Java.
leucocarpa, Kunze, ex DC. Prod. vii. 597 = Pernettya leucocarpa.
Lindeniana, *Planch. in Fl. des Serres*, Sér. I. v. (1849) 501 f.—Venezuela.
linearifolia, DC. Prod. vii. 597 = Gaylussacia pinifolia.
longiflora, *Morr. in Ann. Soc. Hort. Gand*, i. (1845) t. 48.—Mexic.?
Lowii, *Regel, Ind. Sem. Hort. Petrop.* (1856) 80.—Hab.?
loxensis, *Benth. Pl. Hartw.* 143.—Peruv.
luzonica, *A. Gray, in Proc. Am. Acad.* v. (1861) 324.—Ins. Philipp.
microphylla, *Hook. f. Fl. Antarct.* ii. 327 (t. 116 *antarctica*).—Reg. Magellan.
mucronata, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 281 = Pernettya rigida.
mucronata, *Phil. in Linnaea*, xxix. (1857–58) 12.—Chili.
mucronata, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 322.—Solivia.
Myrsinites, *Hook. Fl. Bor. Am.* ii. 35.—Am. bor. occ.
myrsinoides, H. B. & K. Nov. Gen. et Sp. iii. 283 = Pernettya myrsinoides.
myrtilloides, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 281.—Chili.
myrtilloides, Poepp. & Endl. Nov. Gen. et Sp. i. 23. t. 41 = Poeppigii.
myrtilloides, Schlecht. in Linnaea, i. (1826) 523 = Pernettya myrtilloides.
nitida, Benth ex Seem. Bot. Voy. Herald, 316 = glaucifolia.
nitida, *Benth. Pl. Hartw.* 45.—Mexic.
Nummulariae, DC. Prod. vii. 592 = nummulariioides.
nummularioides, *D. Don, Prod. Fl. Nep.* 150.—Reg. Himal.; Java.
odorata, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 425.—Am. trop.
oppositifolia, *Hook. f. Fl. N. Zel.* i. 162. t. 48.—N. Zel.
organensis, *Meisn. in Mart. Fl. Bras.* vii. 158.—Bras.
ovalifolia, Wall. Cat. n. 1523 = fragrantissima.
ovata, *DC. Prod.* vii. 526.—Mexic.
ovatifolia, *A. Gray, in Proc. Am. Acad.* xix. (1883) 85.—Am. bor. occ.
petraea, *Wedd. Chlor. And.* ii. 174.—N. Granat.
pichinchensis, *Benth. Pl. Hartw.* 252.—Ecuador.
pilosa, Zoll. & Mor. in Nat. in Genesak. Arch. Neêrl. Indie, ii. (1846) 9 = Diplycosia pilosa.
Poeppigii, *DC. Prod.* vii. 593.—Chili.
procumbens, *Linn. Sp. Pl.* 395.—Am. bor.
punctata, Blume, Bijdr. 856 = fragrantissima.
punctata, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 281 = vernalis.
purpurascens, *H. B. & K. Nov. Gen. et Sp.* iii. 282.—N. Granat.
pyrolaefolia, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 457.—Reg. Himal.
pyroloides, Hook. f. & Thom. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 30 = praec.
ramosissima, *Benth. Pl. Hartw.* 220.—N. Granat.
Rengifoana, *Phil. in Anal. Univ. Chil.* (1854) reimp. 6.—Chili.
repens, Blume, Bijdr. 857 = nummulariioides.
repens, Rafn. Med. Fl. i. 302 = procumbens.
reticulata, *H. B. & K. Nov. Gen. et Sp.* iii. 284.—Ecuador.
rigida, *H. B. & K. l. c.* 286.—Venezuela.
Rorsimae, *Klotzsch, ex Meisn. in Mart. Fl. Bras.* vii. 158.—Guian.
rufescens, *DC. Prod.* vii. 595.—Peruv.
rupestris, *R. Br. Prod.* 559.—N. Zel.
saxicola, *Wedd. Chlor. And.* ii. 174.—Bolivia.
scabra, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 424 = odorata.
secunda, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 321.—N. Granat.
serpyllifolia, Purah, Fl. Am. Sept. i. 268 = Chiogenes serpyllifolia.
Shallon, *Purah, l. c.* 268. t. 12.—Am. bor. occ.
sphagnicola, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Epigaea cordifolia.
Sprengelii, G. Don, Gen. Syst. iii. 840 = Gaylussacia Pseudo-Gaultheria.
strigosa, *Benth. Pl. Hartw.* 221.—N. Granat.

GAULTHERIA:—
tollimensis, *Wedd. Chlor. And.* ii. 178.—N. Granat.
tomentosa, *H. B. & K. Nov. Gen. et Sp.* iii. 287.—Ecuador.
trichocalycina, *DC. Prod.* vii. 595.—Mexic.
trichophylla, *Royle, Illustr. Bot. Himal.* 360. t. 68. f. 2.—Reg. Himal.
triquetra, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 199.—Japon.
vaccinioides, *Griseb. ex Wedd. Chlor. And.* ii. 176.—Peruv.
venusta, *DC. Prod.* vii. 597.—Bras.
vernalis, *Kunze, ex DC. l. c.* 594.—Chili.
vestita, *Benth. Pl. Hartw.* 224.—N. Granat.

GAULTHIERA, Reichb. Handb. 206 (1837) = praec.

GAULTHIERIA, Klotzsch, in Linnaea, xxiv. (1851) 17 = praec.

GAURA, Lam. Illustr. t. 281. f. 3 (1793 !) = **Lechea**, Kalm (Cistin.).
racemulosa, Lam. Tabl. ii. 423 = L. racemulosa.

GAURA, Linn. Diss. Chen. 47 (1751); Amoen. Acad. ii. 26 (1756). *ONAGRARIEAE*, Benth. & Hook. f. i. 792.
GAURIDIUM, Spach, Hist. Vég. Phan. iv. 879 (1835).
SCHIZOCARYA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835). 170, 283.
STENOSIPHON, Spach, l. c. 170 (1835).
angustifolia, Herb. Willd. ex Stend. Nom. ed. II. i. 666 = coccinea.
angustifolia, *Michx. Fl. Bor. Am.* i. 226.—Am. bor.
australis, *Griseb. in Gott. Abh.* xxiv. (1879) 130.—Reg. Argent.
biennis, *Linn. Sp. Pl.* 347.—Am. bor.
biennis, Walt. Fl. Carol. 128 = angustifolia.
bracteata, *Ser. in DC. Prod.* iii. 45.—Mexic.
canescens, Hort. ex Ser. l. c. = mutabilis !
chinensis, *Lour. Fl. Cochinch.* 235.—China.
coccinea, Bertol. Misc. Bot. ii. 17 = filipes.
coccinea, *Nutt. in Fraser, Cat.* (1813); *et Gen. Am.* i. 249.—Am. bor.
crispa, *D. Dietr. Syn. Pl.* i. 1298.—Mexic.
decorticans, Hook. & Am. Bot. Beech. Voy. 343 = Oenothera gaurseflora.
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 519.—Am. bor.
ephlobia, [*Moç. & Sesse,*] *ex Ser. in DC. Prod.* iii. 45.—Mexic.
epilobioides, H. B. & K. Nov. Gen. et Sp. vi. 93 = coccinea.
exalata, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 44 = longiflora.
filipes, *Spach, in Nouv. Ann. Mus. Par.* iv. (1835) 279.—Am. bor.
fruticosa, Jacq. Coll. i. 142 = angustifolia.
fruticulosa, *Benth. Bot. Voy. Sulph.* 15.—Calif.
glabra, Lehm. in Hook. Fl. Bor. Am. i. 209 = coccinea.
heterandra, Torr. in Facif. Rail. Rep. iv. 87 = Heterogaura californica.
hexandra, Orteg. Hort. Matr. Dec. 14 = tripetala.
hirsuta, Scheele, in Linnaea, xxi. (1848) 580 = Lindheimeri !
hispida, *Benth. Pl. Hartw.* 288.—Mexic.
humilis, D. Dietr. Syn. Pl. ii. 1297 = Gayophytum humile.
laxa, Loefl. It. Hispan. 248 = Combretum Loeflingii.
leucocarpa, *Comien, ex Steud. Nom.* ed. II. i. 666.—Texas.
Lindheimeri, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 217.—Texas.
linifolia, Nutt. in James, Exped. Rock. Mount. ii. 255 = Stenosiphon virgatus.
longiflora, *Spach, in Nouv. Ann. Mus. Par.* iv. (1835) 280.—Am. bor.
macrocarpa, *Rothrock, in Proc. Am. Acad.* vi. (1864–65) 353.—Am. bor.
marginata, Lehm. in Hook. Fl. Bor. Am. i. 208 = coccinea.
Michauxii, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 279 = filipes.
micrantha, D. Dietr. Syn. Pl. ii. 1297 = parviflora.
mollis, *H. B. & K. Nov. Gen. et Sp.* vi. 93. t. 529.—Mexic.
mollis, Nutt. ex Torr. in Ann. Lyc. N. York, ii. (1828) 200 = parviflora.

GAURA:—
multicaulis, *Rafin. Atl. Journ.* 146.—Am. bor.
mutabilis, *Cav. Ic.* iii. 80. t. 285.—Mexic.
odorata, *Lag. Gen. et Sp. Nov.* 14.—Mexic.
oenotheriflora, Zuccagni, in Roem. Collect. 140 = mutabilis !
parviflora, *Dougl. in Hook. Fl. Bor. Am.* i. 208.—Am. bor. occ.
parviflora, Torr. in Ann. Lyc. N. York, ii. (1828) 201 = coccinea.
Pitcheri, Picker. ex Torr. & Gray, Fl. N. Am. i. 517 = biennis.
primiveris, *A. Gray, Pl. Wright.* ii. 58.—Mexic.
Roemeriana, Scheele, in Linnaea, xxi. (1848) 579 = Drummondii.
sinuata, [Moç. & Sesse] ex Ser. in DC. Prod. iii. 45 = bracteata.
sinuata, *Nutt. ex Ser. l. c.* 44.—Am. bor. occ.
spicata, [Moç. & Sesse] l. c. = bracteata.
suffruticosa, [Moç. & Sesse] l. c. = coccinea.
suffulta, *Engelm. in A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 190.—Am. bor. occ.
triangulata, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 464 = tripetala.
tripetala, *Cav. Ic.* iv. 66. t. 396.—Mexic.
tripetala, Herb. Willd. ex Stend. Nom. ed. II. i. 666 = coccinea.
undulata. Desf. Tabl. ed. II. 196 = angustifolia.
villosa, *Torr. in Ann. Lyc. N. York*, ii. (1828) 200.—Am. bor. occ.

GAUREA, Reichb. Gen. 198 (1778) = **Guarea**, Allem. (Meliac.).

GAURIDIUM, Spach, Hist. Vég. Phan. iv. 879 (1835) = **Gaura**, Linn. (Onagrar.).
Kunthii, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 276 = Gaura mollis.
molle, Spach, Hist. Vég. Phan. iv. 880 = Gaura mollis.
mutabile, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 274 = Gaura mutabilis.

GAUROPSIS, Presl, Epim. Bot. 219 (1849). *ONAGRARIEAE*.
lancifolia, *Presl, l. c.*—Mexic.

GAUSSIA, H. Wendl. in Goett. Nachr. (1865) 327 *PALMAE*, Benth. & Hook. f. iii. 913.
Ghiesbreghti, *H. Wendl. in Kerch. Palm.* 245.—Ind. occ.
portoricensis, *H. Wendl. l. c.* 345, nomen.—Ind. occ.
Princeps, *H. Wendl. in Gnett. Nachr.* (1865) 328.—Cuba.
vinifera, *H Wendl. in Kerch. Palm.* 245.—Ind. occ.

GAUTIERA, Rafn. Med. Fl. i. 202 (1828) = **Gaultheria**, Kalm (Ericac.).

GAVARRETIA, Baill. Adansonia, i. 185. t. 7 (1860–61). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 316.
terminalis, *Baill. l. c.* 186.—Bras.

GAVESIA, Walp. Ann. ii. 650 (1852) = GATESIA, Bertol. = **Petalostemon**, Michx. (Legumin.).

GAVILLEA, Poepp. & Endl. ex Stend. Nom. ed. II. i. 666 (1840) = **Chlorasa**, Lindl. (Orchid.).

GAVNIA, Scop. Introd. 76 (1777) sphalm. = **Gahnia**, Forst. (Cyperac.).

GAYA, Gand. Fl. Helv. ii. 389 (1828) = **Ligusticum**, Linn. (Umbellif.).
multicaulis, Schur, Enum. Pl. Trans. 259 = L. simplex.
pyrenaica, Boiss. Voy. Bot. Espagne, 250, partim = Seseli granatense.
pyrenaica, Gand. Fl. Helv. ii. 390, in obs. = Seseli annum.
simplex, Gand. l. c. 389 = L. simplex.

GAYA, H. B. & K. Nov. Gen. et Sp. v. 266. t. 475, 476 (1821). *MALVACEAE*, Benth. & Hook. f. i. 208.
affinis, *A. Rich. Ess. Fl. Cub.* 157.—Cuba.
asarea, *A. St. Hil. Fl. Bras. Mer.* i. 198.—Bras.
canescens, H. B. & K. Nov. Gen. et Sp. v. 269 = Sida cardicana.

GAYA:—
distichа, *Presl, Rel. Haenk.* i. 466.—Mexic.
Gandichaudiana, *St. Hil. Fl. Bras. Mer.* i. 193.—Bras.
hermannioides, *H. B. & K. Nov. Gen. et Sp.* v. 288. t. 475.—Mexic.
nutans, Sweet, Hort. Brit. ed. II. 64; Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 201 = Sida nutans.
occidentalis, Griseb. Fl. Brit. W. Ind. 79 = affinis.
occidentalis, Sweet, Hort. Brit. ed. II. 64 = subtriloba.
subtriloba, *H. B. & K. Nov. Gen. et Sp.* v. 270. t. 476. —Mexic.

GAYA, Spreng. Syst. i. 535 (1825) = **Seringia**, J. Gay (Sercul.).
parviflora, Spreng. ex DC. Prod. ix. 15 = Strychnos brasiliensis.
platyphylla, Spreng. Syst. i. 971 = Ser. platyphylla.

GAYLUSSACIA, H. B. & K. Nov. Gen. et Sp. iii. 375. t. 257 (1818). *VACCINIACEAE*, Benth. & Hook. f. ii. 572.
DECACHAENA, Torr. & Gray, ex Lindl. Veg. Kingd. 758 (1847).
DECAMERIUM, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1842) 259.
LUSSACIA, Spreng. Syst. ii. 294 (1825).
acicularis, Mart. ex Meisn. in Mart. Fl. Bras. vii. 147 = pinifolia.
adenochaeta, *DC. Prod.* vii. 559.—Bras.
affinis, Mart. ex Meisn. in Mart. Fl. Bras. vii. 147 = pallida.
amoena, *Cham. in Linnaea*, viii. (1833) 501.—Bras.
amoena, Mart. ex Meisn. in Mart. Fl. Bras. vii. 154 = G. Vitis-Idaea.
angulata, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 131.—Bras.
angustifolia, *Cham. in Linnaea*, viii. (1833) 499.—Bras.
angustifolia, Dunal, ex Meisn. in Mart. Fl. Bras. vii. 144 = thymeleoides.
baccata, C. Koch, Dendrol. ii. i. 93 = resinosa.
brachycera, *Torr. & Gray, ex A. Gray, Chlor. Bor. Am.* 54.—Am. bor.
bracteata, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 130.—Bras.
brasiliensis, *Meisn. in Mart. Fl. Bras.* vii. 129.—Bras.
buxifolia, Cham. & Schlecht. in Linnaea, i. (1826) 528 = decipiens.
buxifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 276. t. 257. —N. Granat.
canescens, *Meisn. in Mart. Fl. Bras.* vii. 143.—Bras.
caracasana, *DC. Prod.* vii. 559.—Venezuela.
cardinalis, Mart. ex Meisn. in Mart. Fl. Bras. vii. 136 = salicifolia.
Chamissonis, *Meisn. l. c.* 146.—Bras.
ciliosa, *Meisn. in Mart. Fl. Bras.* vii. 146.—Bras.
cordata, Hemsl. Biol. Centr. Am. Bot. ii. 174 = sec. *cordifolia*, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 529 = Vaccinium cordatum.
crenata, *D. Don, ex Dunal, in DC. Prod.* vii. 557.—Peruv.
crenulata, Mart. ex Meisn. in Mart. Fl. Bras. vii. 133 = pulchra.
decipiens, *Cham. in Linnaea*, viii. (1833) 500.—Bras.
dentata, *Cham. l. c.* 498.—Bras.
dependens, *G. Don, Gen. Syst.* iii. 859.—Peruv.
discolor, *Dunal, in DC. Prod.* vii. 557.—Bras.
dumosa, *Torr. & Gray, in A. Gray, Chlor. Am. Bor.* 50. —Am. bor.
elliptica, Mart. ex Meisn. in Mart. Fl. Bras. vii. 131 = brasiliensis.
elliptica, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 9 = Vaccinium ellipticum.
fascicularis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 131.—Bras.
ferruginea, Meisn. in Mart. Fl. Bras. vii. 158 = Gaultheria erguensis.
frondosa, *Torr. & Gray, ex Torr. Fl. N. York*, i. 449. —Am. bor.
Gardneri, *Meisn. in Mart. Fl. Bras.* vii. 137.—Bras.
hirtella, Torr. & Gray, ex Torr. Fl. N. York, i. 448. t. 97 = dumosa.
hispida, *DC. Prod.* vii. 559.—Bras.
hispida, Steud. Nom. ed. II. i. 666 = G. Pseudo-Gaultheria.
imbricata, *Pohl, Pl. Bras. Ic.* ii. 40. t. 136 = brasiliensis.
GAYLUSSACIA:—
incana, Cham. in Linnaea, i. (1826) 536 = thymeleoides.
incurvata, Griff. Notul. iv. 800; Ic. Pl. As. t. 506 = Pentapterygium rugosum.
lanceolata, Blume, Bijdr. 861 = Vaccinium acuminatissimum.
ledifolia, *Mart. ex Meisn. in Mart. Fl. Bras.* vii. 145.—Bras.
leptobotrys, *Dunal, in DC. Prod.* vii. 560.—Bras.
macrostachia, Mart. ex Meisn. in Mart. Fl. Bras. vii. 142 = reticulata.
Martii, *Meisn. l. c.* 138.—Bras.
microphylla, *G. Don, Gen. Syst.* iii. 859.—Peruv.
montana, Mart. ex Meisn. in Mart. Fl. Bras. vii. 145 = hispida.
myrtifolia, Cham. in Linnaea, viii. (1833) 494 = brasiliensis.
myrtifolia, Mart. ex Meisn. in Mart. Fl. Bras. vii. 133 = pulchra.
myrtilloides, Cham. in Linnaea, viii. (1833) 498 = hispida.
nitida, *Mart. ex Meisn. in Mart. Fl. Bras.* vii. 134.—Bras.
obscura, DC. Prod. vii. 559 = pallida.
octosperma, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 130.—Bras.
oleaefolia, *Dunal, in DC. Prod.* vii. 559.—Bras.
pallida, *Cham. in Linnaea*, viii. (1833) 499.—Bras.
pallida, Dunal, ex DC. Prod. vii. 558 = virgata.
parvifolia, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 130.—Bras.
pinifolia, *Cham. & Schlecht. in Linnaea*, i. (1826) 536. —Bras.
Pseudo-Gaultheria, *Cham. & Schlecht. l. c.* 535.—Bras.
Pseudo-Vaccinium, *Cham. & Schlecht. l. c.* 530.—Bras.
Pseudo-Vaccinium, Dunal, ex Meisn. in Mart. Fl. Bras. vii. 136 = salicifolia.
Pseudo-Vaccinium, Dunal, in DC. Prod. vii. 557 = Vauthieri.
pulchra, Miq. ex Meisn. in Mart. Fl. Bras. vii. 160 = Leucothoe revoluta.
pulchra, *Pohl, Pl. Bras. Ic.* ii. 41. t. 127.—Bras.
resinosa, *Torr. & Gray, ex A. Gray, Chlor. Bor. Am.* 100.—Am. bor.
reticulata, *Mart. ex Meisn. in Mart. Fl. Bras.* vii. 142.—Bras.
retusa, *Mart. ex Meisn. l. c.* 132.—Bras.
rhodendra, St. Lag. in Ann. Sc. Nat. Lyon, vii. (1880) 126 = sec.
Rhododendron, *Cham. & Schlecht. in Linnaea*, i. (1826) 533.—Bras.
Riedelli, *Meisn. in Mart. Fl. Bras.* vii. 139.—Bras.
rigida, *Casar. Nov. Stirp. Bras. Decad.* iv. 40.—Bras
rugosa, *Cham. & Schlecht. in Linnaea*, i. (1826) 534. —Bras.
salicifolia, *Cham. & Schlecht. l. c.* 532.—Bras.
Sellowiana, Klotzsch, ex Meisn. in Mart. Fl. Bras. vii. 131 = brasiliensis.
serrata, Griff. Itin. Not. 37; Notul. iv. 800 = Pentapterygium rugosum.
serrata, Lindl. in Royle, Illustr. Bot. Himal. 257. t. 79 = Vaccinium serratum.
subcordata, *Dunal, in DC. Prod.* vii. 557.—Peruv.
thymeleoides, *Meisn. in Mart. Fl. Bras.* vii. 144.—Bras.
ursina, *Torr. & Gray, in A. Gray, Chlor. Bor. Am.* 49. t. 10.—Am. bor.
Vaccinium, Steud. Nom. ed. II. i. 666 = G. Pseudo-Vaccinium.
Vauthieri, *Meisn. in Mart. Fl. Bras.* vii. 138.—Bras.
venusta, Mart. ex Meisn. l. c. 131 = brasiliensis.
villosa, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 130.—Bras.
virgata, *Mart. ex Meisn. in Mart. Fl. Bras.* vii. 141. —Bras.
Vitis-Idaea, *Mart. ex Meisn. l. c.* 154.—Bras.

GAYOPHYTUM, A. Juss. in Ann. Sc. Nat. xxv. (1832) 16. t. 4. *ONAGRARIEAE*, Benth. & Hook. f. i. 789.
caesium, Torr. & Gray, Fl. N. Am. i. 514 = racemosum.
diffusum, *Torr. & Gray, l. c.* 513.—Am. bor. occ.
humile, *Fuss. in Ann. Sc. Nat. Sér. I.* xxv. (1832) 18. —Chili.
micranthum, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 311 = humile.
GAYOPHYTUM:—
minutum, *Phil. in Linnaea*, xxviii. (1856) 687.—Chili.
Nuttallii, Torr. & Gray, Fl. N. Am. i. 514 = racemosum.
pumilum, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 198.—Am. bor. occ.
racemosum, *Torr. & Gray, Fl. N. Am.* i. 514.—Am. bor. occ.
ramosissimum, *Torr. & Gray, l. c.* 513.—Am. bor. occ.
strictum, A. Gray, in Proc. Am. Acad. vii. (1868) 340 = Oenothera Torreyi.

GAVTANIA, Mnent. in Bot. Zeit. i. (1843) 730 = **Pimpinella**, Riv. (Umbellif.).
Chemeyana, Mnent. l. c. = P. eriocarpos.
daucoides, Boiss. Fl. Orient. ii. 867, in syn. = P. eriocarpos.

GAZANIA, Gaertn. Fruct. ii. 451. t. 173 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 459.
MELANCHRYSUM, Cass. in Bull. Soc. Philom. (1817) 12.
MERIDIANA, Hill, Veg. Syst. ii. 121** (1761).
MORNNIA, Neck. Elem. i. 9 (1790).
MUSINIA, Willd. Sp. Pl. iii. 2363 (1803).
araneosa, DC. Prod. vi. 519 = Lichtensteinii.
arctotoides, *Less. Syn. Comp.* 48.—Afr. austr.
arminioides, *DC. Prod.* 511.—Afr. austr.
Burchellii, *DC. l. c.* vi. 514.—Afr. austr.
caespitosa, *Bolus, in Journ. Linn. Soc.* xviii. (1881) 263.—Afr. austr.
canescens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 478.—Afr. austr.
ciliaris, DC. Prod. vi. 513 = pinnata.
coronopifolia, *DC. l. c.* 511.—Afr. austr.
diffusa, Spreng. Syst. iii. 606 = Gorteria diffusa.
Forbesiana, DC. Prod. vi. 508 = Cryptostemma Forbesianum.
grandis, DC. l. c. 509 = pinnata.
heterochaeta, *DC. l. c.* 511.—Afr. austr.
heterophylla, Willd. ex Steud. Nom. ed. II. i. 667 (Quid?)—Am. bor.
hirtella, DC. Prod. vi. 511 = G. Pavonia.
humilis, E. Mey. ex DC. l. c. 518 = variana.
integrifolia, Spreng. Syst. iii. 606 = Hirpicium integrifolium.
intrusa, E. Mey. ex DC. Prod. vi. 511 = othonnites.
jurineaefolia, *DC. l. c.* 511.—Afr. austr.
Kraussii, *Sch. Bip. in Flora*, xxvii. (1844) 777.—Afr. austr.
Krebsiana, *Less. Syn. Comp.* 44.—Afr. austr.
leptophylla, *DC. Prod.* vi. 513.—Afr. austr.
leucolaena, DC. l. c. 509 = uniflora.
Lichtensteinii, *Less. Syn. Comp.* 50.—Afr. austr.
linearifolia, *DC. Prod.* vi. 513.—Afr. austr.
longifolia, *Less. Syn. Comp.* 48.—Afr. austr.
longiscapa, *DC. Prod.* vi. 518.—Afr. austr.
macronata, *DC. l. c.*—Afr. austr.
multiceps, *DC. l. c.* 508.—Afr. austr.
nivea, *Less. Syn. Comp.* 49.—Afr. austr.
othonnites, *Less. l. c.* 48.—Afr. austr.
othonnites, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = prasc.
oxyloba, *DC. Prod.* vi. 513.—Afr. austr.
Pavonia, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 140.—Afr. austr.
pectinata, Spreng. Syst. iii. 607 = pinnata.
pinnata, DC. Prod. vi. 509 = G. Pavonia.
pinnata, *Less. Syn. Comp.* 43.—Afr. austr.
pygmaea, *Sond. in Linnaea*, xxiii. (1850) 69.—Afr. austr.
rigens, *Meusch, Meth. Suppl.* 209.—Afr. austr.
scabra, DC. Prod. vi. 509 = pinnata.
serrata, DC. l. c. 510 = pinnata.
serrulata, *DC. l. c.* 513.—Afr. austr.
speciosa, Less. Syn. Comp. 42 = pinnata.
splendens × *hirta*, Hort. Angl. ex Illustr. Bonq. ii. t. 29; Lem. Illustr. Hortic. vii. (1860) t. 235.
stoechadifolia, Hort.—Cf. Gard. Chron. (1883) ii. 77 = longiscapa.
subdipinnata, *DC. Prod.* vi. 508.—Afr. austr.
subulata, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 140.—Afr. austr.
subulata, Sch. Bip. in Flora, xxvii. (1844) 777 = Kraussii, longiscapa.
tenuifolia, *Less. Syn. Comp.* 50.—Afr. austr.
uniflora, *Sims, Bot. Mag.* t. 2270.—Afr. austr.
variana, *DC. Prod.* vi. 513.—Afr. austr.

GEANTHIA, Rafin. in Med. Repos. N. York, v. (1808) 350 = **Weldenia** ? Schult. f. (Commelinac.).
colchicoides, Rafin. l. c. = W. candida ?

GEANTHUS, Phil. Desc. Pl. Nuev. Chil. (1884) reimpr. 7. *LILIACEAE*.
humilis, *Phil. l. c.* 8.—Chili.

GEANTHUS, Reinw. in Blume, Cat. Gew. Buitenz. 29 (1823); et in Syll. Ratisb. ii. (1828) 5 = **Amomum**, Linn. (Scitamin.).
coccineus, Reinw. in Blume, l. c. = A. coccinea.
minor, Reinw. l. c. (= *Donacodes minor*).—Java.
Pisingi, Reinw. l. c. = Alpinia Pisinga.
speciosus, Reinw. l. c. = Hettaria speciosa.

GEARUM, N. E. Br. in Journ. Bot. xx. (1882) 196. t. 231. f. 1. *AROIDEAE*, Benth. & Hook. f. iii. 987.
brasiliense, *N. E. Br. l. c.*—Bras.

GEBLERA, Andrz. ex Bess. in Flora, xvii. (1834) i. Beibl. 18 = **Crepis**, Vaill. (Compos.).
prenanthoides, Andrz. l. c. = C. baicalensas.

GEBLERA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. (1835) 28 = **Securinega**, Juss. (Euphorbiac.).
suffruticosa, Fisch. & Mey. l. c. = S. ramiflora.
sungurensis, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 357.—As. centr.

GEBOSCON, Rafin. Cat. 14 (1824) = **Allium**, Tourn. (Liliac.).
obliquum, Rafin. Fl. Tellur. ii. 19 = A. obliquum.
striatum, Rafin. Cat. 14 = Nothoscordum striatum.

GEBRIA, Blume, Cat. Gew. Buitenz. 18 (1823); et Bijdr. 124 (1825) = **Bruya**, Thunb. (Ternstroem.).
angustifolia, Blume, Bijdr. 125 = E. angustifolia.
glabra, Blume, l. c. = E. japonica.
obovata, Blume, l. c. = E. obovata.
serrata, Blume, Cat. Gew. Buitenz. 111; et Bijdr. 124 = E. acuminata.

GEERIA, Neck. Elem. ii. 241 (1790) = **Paullinia**, Linn. (Sapindac.).

GEIGERA, Less. Syn. Comp. 199 (1832) = **Geigeria**, Griessel. (Compos.).

GEIGERA, Lindl. Veg. Kingd. 471 (1847) = **Geijera**, Schott (Rutac.).

GEIGERIA, Griessel. in Linnaea, v. (1830) 411. *COMPOSITAE*, Benth. & Hook. f. ii. 337.
ARASCHCOOLIA, Sch. Bip. ex Flora, xxiv. (1841) Intell. 144.
DIPLOSTEMMA, Steud. & Hochst. in DC. Prod. vii. 75 (1838).
DIRONIUM, Willd. ex Schlecht. in Linnaea, v. (1830) 411, in nota.
GEIGERA, Less. Syn. Comp. 199 (1832).
ZEYHERIA, Spreng. i. Tent. Suppl. 26 (1828).
acaulis, *Benth. & Hook. f. ex Yatky, in Oastr. Bot. Zeitschr.* xxv. (1875) 227.—Afr. trop.
africana, *Griessel. in Linnaea*, v. (1830) 411.—Afr. austr.
alata, *Benth. & Hook. f. ex Oliver, Fl. Trop. Afr.* iii. 368.—Afr. trop.
aspera, *Harv. in Harv. & Sond. Fl. Cap.* iii. 126.—Afr. austr.
brevifolia, *Harv. l. c.*—Afr. austr.
Burkei, *Harv. l. c.*—Afr. austr.
passerinoides, *Harv. l. c.* 125.—Afr. austr.
pectidea, *Harv. l. c.* 127.—Afr. austr.
protensa, *Harv. l. c.* 125.—Afr. austr.
Zeyheri, *Harv. l. c.* 126.—Afr. austr.

GEIJERA, Schott, Rutac. 7. t. 4 (1834). *RUTACEAE*, Benth. & Hook. f. i. 396.
COATESIA, F. Muell. Fragm. iii. 26 (1862).
canliflora, *Baill. Adansonia*, x. (1871-72) 321.—N. Caled.
latifolia, Lindl. in Mitch. Journ. Trop. Austral. 236 = salicifolia.
Muelleri, *Benth. Fl. Austral.* i. 364.—Austral.
parviflora, *Lindl. in Mitch. Journ. Trop. Austral.* 102. —Austral.
pendula, Lindl. l. c. 251 = parviflora.
salicifolia, *Schott, Rutac.* 7. t. 4.—Austral.

GEISARINA, Rafin. Alsog. Am. 5 (1838) = **Myrica** ? Linn.
verrucosa, Rafin. l. c.—Am. bor.

GEISELERIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 254 = **Croton**, Linn. (Euphorbiac.).
chamaedryfolia, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 47 = C. chamaedryfolius.
chamaedryfolia, Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 418 = C. glandulosus.
glandulosa, Klotzsch, in Wiegm. Archiv, vii. (1841) 254 = C. glandulosus.

GEISELERIA, Kunth, in Abh. Berl. Akad. (1842) 3 = ANTICLEA, Kunth = **Zygadenus**, Michx. (Liliac.).

GEISENIA, Endl. Gen. 848, in syn. (1839) = GAISSENIA, Rafin. = **Trollius**, Linn. (Ranunc.).

GEISOLOMA, Lindl. Veg. Kingd. 578 (1847) = **Geissoloma**, Lindl. (Penaeac.).

GEISORRHIZA, Reichb. Nom. 44 (1841) = **Geissorhiza**, Ker-Gawl. (Irid.).

GEISSANTHUS, Hook. f. in Benth. & Hook. f. Gen. ii. 642 (1876). *MYRSINEAE*, Benth. & Hook. f. i. c.

GEISSAPSIS, Baker, in Hook. f. Fl. Brit. Ind. ii. 141 (1876), sphalm. = seq.

GEISSASPIS, Wight & Arn. Prod. 217 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 516.
cristata, *Wight & Arn. l. c.* 218.—Ind. or.
lupulina, *Planch. ex Baker, in Oliver, Fl. Trop. Afr.* ii 155.—Afr. trop.
tenella, *Benth. in Flora*, xxxii. (1849) 559.—Ind. or.

GEISSOIS, Labill. Sert. Austro-Caled. 50. t. 50 (1825). *SAXIFRAGEAE*, Benth. & Hook. f. i. 650.
Benthami, *F. Muell. Fragm.* v. 180.—Austral.
Biagiana, F. Muell. l. c. 16 = Weinmannia Biagiana.
Denhami, *Seem. Fl. Vit.* 109.—N. Hebrides.
hirsuta, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 71.—N. Caled.
montana, *Pieill. ex Brongn. & Gris, l. c.* 71.—N. Caled.
pruinosa, *Brongn. & Gris, l. c.* 70.—N. Caled.
racemosa, *Labill. Sert. Austro-Caled.* 50. t. 50.—N. Caled.
rubifolia, F. Muell. Fragm. ii. 82 = Weinmannia rubifolia.
tenuata, *A. Gray, Bot. U. St. Expl. Exped.* i. 679. t. 86.—Ins. Fiji.

GEISSOLOMA, Lindl. ex Kunth, in Linnaea, v. (1830) 678. *PENAEACEAE*, Benth. & Hook. f. iii. 205.
laterifforum, Kunth, l. c.; Drège, ex A. DC. in Prod. xiv. 491, 492 = Endonemma Thunbergii.
marginatum, *Kunth, l. c.*—Afr. austr.
tomentosum, *Kunth, l. c.*—Afr. austr.

GEISSOMERIA, Lindl. Bot. Reg. t. 1045 (1827). *ACANTHACEAE*, Benth. & Hook. f. ii. 1103.
PACHYLOENEMIS, Mart. ex Nees, in DC. Prod. xi. 286 (1847).
SALPIXANTHA, Hook. Bot. Mag. t. 4158 (1845).
brasiliensis, *Nees, in Mart. Fl. Bras.* ix. 88.—Bras.
castrifolia, *Nees, l. c.* 80.—Bras.
cincinnata, *Nees, l. c.* 81.—Bras.
coccinea, *T. Anderso. ex Griseb. Fl. Brit. W. Ind.* 454. —Ind. occ.
distans, *Nees, in Mart. Fl. Bras.* ix. 85.—Bras.
fulgida, *Sweet. Hort. Brit.* ed. II. 430.—Ind. occ.
longiflora, *Lindl. Bot. Reg.* t. 1045.—Bras.
longiflora, Salzm. ex Nees, in DC. Prod. xi. 287 = macrophylla.
macrophylla, *Nees, in Mart. Fl. Bras.* ix. 80.—Bras.
nitida, *Nees, l. c.*—Bras.
pubescens, *Nees, l. c.* 82.—Bras.
Schottiana, *Nees, l. c.*—Bras.

GEISSOPAPPUS, Benth. in Hook. Lond. Journ. Bot. ii. (1840) 44. *COMPOSITAE*, Benth. & Hook. f. ii. 396.
SCHOMBURGKIA, DC. Prod. vii. 293 (1838).
TRINCHINETTIA, Endl. Gen. 1383 (1841).
calcoides, *Benth. in Hook. Journ. Bot. ii.* (1840) 44.—Guian.
gentianoides, *Baker, in Mart. Fl. Bras.* vi. 111. 279.—Bras.
polycephalus, *Baker, l. c.*—Bras.

GEISSORHIZA, Ker-Gawl. in Bot. Mag. t. 672 (1803). *IRIDEAE*, Benth. & Hook. f. iii. 703.
ROCHEA, Salisb. in Trans. Hort. Soc. i. (1812) 322.
WEIHEA, Eckl. Verz. 22 (1827).
abyssinica, R. Br. in Salt, Abyss. App. 64 = Lapeyronaia abyssinica.
abyssinica, Klatt, in Linnaea, xxxiv. (1865-66) 716 = Hesperantha Petitiana.
alberta, E. Mey. in Drège, Zwei Pfl. Docum. 157 = Gladiolus debilis.
alpina, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 223 —Afr. trop.
anemonaeflora, Klatt, in Linnaea, xxxiv. (1865-66) 697 = Iris anemonaeflora.
arenaria, Eckl. Top. Verz. 21 = imbricata.
aurea, Eckl. l. c. = humilis.
Bojeri, *Baker, in Journ. Bot.* xiv. (1876) 239.—Madag.
bractesta, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 201.—Afr. austr.
Brehmii, Eckl. Top. Verz. 21 = humilis.
candida, *Eckl. l. c.* nomen.—Afr. austr.
ciliaris, *Ker-Gawl. in Bot. Mag.* sub t. 672.—Afr. austr.
corrugata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 201.—Afr. austr.
cyanea, *Eckl. Top. Verz.* 20.—Afr. austr.
erecta, *Baker, in Journ. Bot.* xiv. (1876) 238.—Afr. austr.
excisa, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1806) 224.—Afr. austr.
filifolia, *Baker, in Journ. Bot.* xiv. (1876) 238.—Afr. austr.
flava, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 202.—Afr. austr.
foliosa, *Klatt, in Linnaea*, xxxiv. (1865-66) 658.—Afr. austr.
furva, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1806) 224.—Afr. austr.
geminata, *E. Mey. in Drège, Zwei Pfl. Docum.* 167.—Afr. austr.
grandis, *Hook. f. Bot. Mag.* t. 5877.—Afr. austr.
hirta, *Ker-Gawl. in Bot. Mag.* sub t. 672.—Afr. austr.
humilis, *Ker-Gawl. l. c.*—Afr. austr.
imbricata, *Ker-Gawl. l. c.*—Afr. austr.
imbricata, E. Mey. ex Baker, in Journ. Linn. Soc. xvi. (1877) 98 = secunda.
inflexa, *Ker-Gawl. in Bot. Mag.* sub t. 672.—Afr. austr.
juncea, *A. Dietr. Sp. Pl.* ii. 587.—Afr. austr.
Larochei, Lond. Hort. Brit. 16 = Rochenais.
latifolia, *Baker, in Journ. Linn. Soc.* xvi. (1877) 94. —Afr. austr.
longituba, Klatt, in Linnaea, xxxv. (1867-68) 388 = Hesperantha longituba.
lutea, *Eckl. Top. Verz.* 21.—Afr. austr.
minima, *Baker, in Journ. Bot.* xiv. (1876) 239. —Afr. austr.
monantha, Sweet, Hort. Brit. ed. II. 508 = Rochenais.
monanthos, *Eckl. Top. Verz.* 21.—Afr. austr.
nana, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1868) 201.—Afr. austr.
nigro-montana, *Eckl. Verz.* 21.—Afr. austr.
obtusata, *Ker-Gawl. in Bot. Mag.* t. 672.—Afr. austr.
ornithogaloides, *Klatt, in Linnaea*, xxxiv. (1865-66) 656.—Afr. austr.
purpureo-lutea, *Baker, in Journ. Bot.* xiv. (1876) 238. —Afr. austr.
quadrangula, *Ker-Gawl. Irid. Gen.* 88.—Afr. austr.
quinquangularis, Eckl. ex Klatt, in Linnaea, xxxiv. (1865-66) 624 = ciliaris.
radiana, Ker-Gawl. ex Steud. Nom. ed. II. i. 668 = Rochenais.
ramosa, *Ker-Gawl. in Bot. Mag.* sub t. 672.—Afr. austr.
remosa, Klatt, in Linnaea, xxxiv. (1865-66) 657 = secunda.

GEISSORHIZA, — *recurvifolia*, Klatt, in Linnaea, xxxiv. (1865-66) 655 = sublutea.
Rochenna, Sweet, Hort. Brit. ed. I. 389 = seq.
Rochennia, *Ker-Gawl. in Bot. Mag.* t. 672.—Afr. austr.
romuleoides, Eckl. Top. Verz. 21 = ornithogaloides.
rosea, Eckl. l. c. 20 = ciliaris.
sabulosa, *Klatt, in Trans. S. Afr. Phil. Soc.* iii (1885) 208.—Afr. austr.
scillaris, *Ker-Gawl. in Bot. Mag.* sub t. 672.—Afr. austr.
secunda, Houtt. Handleid. xii. t. 78. f. 1 = Rochensis.
secunda, *Ker-Gawl. in Bot. Mag.* sub t. 672 —Afr. austr.
setacea, *Baker, in Journ. Linn. Soc.* xvi. (1877) 95.—Afr. austr.
setacea, Ker-Gawl. in Bot. Mag. sub t. 672 = Rochensis.
setifolia, Eckl. Top. Verz. 22 = secunda.
striata, *Eckl. l. c.* 21.—Afr. austr.
sublutea, Ker-Gawl. in Bot. Mag. sub t. 672 = Romulea sublutea.
tenuis, *Eckl. Top. Verz.* 22.—Afr. austr.
tulipifera, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 390.—Afr. austr.
vaginata, Sweet, Brit. Flow. Gard. t. 188 = inflexa.
Wrightii, *Baker, in Journ. Bot.* xiv. (1876) 288.—Afr. austr.

GEISSORIZA, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 390 = praec.

GEISSORRHIZA, D. Dietr. Syn. Pl. i. 126 (1839) = praec.

GEISSOSPERMUM, Allem. Diss. (1845), cum ic. *APOCYNACEAE*, Benth. & Hook. f. ii. 707.
laeve, Miers, Apocyn. S. Am. 84 = Vellosii.
Martianum, *Miers, l. c.*—Bras.
ramiflorum, Miers, l. c. 85 = Apidosperma ramiflorum.
sericeum, *Benth. & Hook. f. Gen.* ii. 707.—Guiana.
Solandri, *Miers, Apocyn. S. Am.* 85.—Bras.
Vellosii, *Allem. Diss.* (1845), cum ic.—Bras.

GEITONOPLESIUM, A. Cunn. in Bot. Mag. t. 3131 (1832). *LILIACEAE*, Benth. & Hook. f. iii. 768.
CALOOA, Salisb. Gen. Pl. Fragm. 67 (1865).
LEUCHIAGA, R. Br. Prod. 281 (1810).
angustifolium, C. Koch, Ind. Sem. Hort. Berol. (1854) App. 10 = cymosum.
asperum, A. Cunn. in Bot. Mag. t. 3131 = cymosum.
cymosum, *A. Cunn. l. c.*—Austral.; Ins. Pacif.
humile, Hasak. Cat. Hort. Bog. Alt. 31 (Asparagi sp.).—Japon.
montanum, Kunth, Enum. Pl. v. 113 = cymosum.
scandens, Hasak. Cat. Hort. Bog. Alt. 31 (Asparagi sp.).—Japon.

GELA, Lour. Fl. Cochinch. 282 (1790) = **Acronychia**, Forst. (Rutac.).
lanceolata, Lour. l. c. = A. laurifolia.
oblongifolia, A. Cunn. ex Sweet, Hort. Brit. ed. III. 91 = Ximenia oblongifolia.

GELASIA, Cass. in Bull. Soc. Philom. (1818) 33 = **Soorconera**, Tourn. (Compos.).
decurrentum, Less. in Linnaea, ix. (1834) 184 = Aster dracunculoides.
Yucquini, Cass. in Dict. Sc. Nat. xxv. 32 = S. hirsuta.
pauciflora, Less. in Linnaea, ix. (1834) 169 = Aster dracunculoides.
villosa, Cass. in Dict. Sc. Nat. xviii. 286 = S. villosa.

GELASINE, Herb. in Bot. Mag. t. 2779 (1840), *IRIDEAE*, Benth. & Hook. f. iii. 695.
azurea, *Herb. l. c.*—Bras.
grandiflora, Herb. l. c. sub t. 2779 (Nemastylidis sp.).—Chili.
longispatha, Herb. in Benth. Fl. Hartw. 53 = Sphenostigma longispatha.
nuda, Herb. in Bot. Mag. sub t. 2779 = Calydorea nuda.
pumehuta, Herb. l. c. = Nemastylis multiflora.
purpurea, Herb. l. c. (Nemastylidis sp.).—Peruv.
texana, Herb. l. c. = Calydorea texana.

GELEZNOVIA, Benth. & Hook. f. Gen. i. 298 (1862), sphalm. = seq.

GELEZNOWIA, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 12. *RUTACEAE*, Benth. & Hook. f. i. 293.
SANDARIDA, J. Drumm. in Hook. Kew Journ. vii. (1855) 53.
calycina, *Benth. Fl. Austral.* i. 345.—Austral.
IMAGTOCARPA, *Benth. l. c.* 347.—Austral.
verrucosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxii (1849) ii. 13.—Austral.

GELONIUM, Roxb. in Willd. Sp. Pl. iv. 881 (1805). *EUPHORBIACEAE*, Benth. & Hook. f. ii. 324.
CRRATOPHORUS, Sond. in Linnaea, xxiii. (1850) 120.
ERYTHROCARPUS, Blume, Bijdr. 604 (1825).
SUREGADA [Roxb.] = Jones' ex Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 206, nomen prius.
adenophorum, *Muell. Arg. in DC. Prod.* xv. ii. 1128. —Madag.
aequoreum, *Hance, in Journ. Bot.* iv. (1866) 173.—Ins. Formosa.
africanum, *Muell. Arg. in DC Prod.* xv. ii. 1129.—Afr. austr.
angustifolium, Muell. Arg. l. c. 1128 = lanceolatum.
biferium, *Roxb. ex Willd. Sp. Pl.* iv. 881.—Malaya. Boivinianum, *Muell. Arg. in DC. Prod.* xv. ii. 1129.—Madag.
fasciculatum, Roxb. Hort. Beng. 73; Fl. Ind. iii. 832 = multiflorum
glomeruliatum, Hassk. Cat. Hort. Bog. Alt. 287 = bifarium.
lanceolatum, *Willd. Sp. Pl.* iv. 882.—Ind. or.
laurinum, *Muell. Arg. in DC. Prod.* xv. ii. 1129.—Madag.
multiflorum, *A. Juss. Tent. Euph.* 111. t. 10.—Ind. or.; Malaya.
obtusum, Miq. Fl. Ind. Bat. Suppl. 452 = bifarium.
oxyphyllum, *Miq. l. c.*—Sumatra.
spicatum, *Hassk. Cat. Hort. Bog. Alt.* 287.—Java.
zanzibaricum, *Muell. Arg. in DC. Prod.* xv. ii. 1130. —Afr. trop.

GELONIUM, Gaertn. Fruct. ii. 271. t. 139 (1791) = **Ratonia**, DC. (Sapindac.).
cupaniioides, Gaertn. l. c. 272 = Cupania cupaniioides.

GELPKEA, Blume, Mus. Bot. Lugd. Bat. i. 86. t. 35 (1849) = **Eugenia**, Linn. (Myrtac.).
pendula, Blume, l. c. = E. pendula.
stipularia, Blume, l. c. = E. stipularis.

GELSEMINUM, Pursh, Fl. Am. Sept. i. 184 (1814) = seq.

GELSEMIUM, Juss. Gen. 150 (1789). *LOGANIACEAE*, Benth. & Hook. f. ii. 789.
JEFFERSONIA, Brick. in Med. Rep. N. York, i. (1800) 555.
LEPTOPHASIS, Blume, Mus. Bot. Lugd. Bat. i. 240. t. 24 (1849).
MEDICIA, Gardn. in Hook. Kew Journ. i. (1849) 324.
elegans, *Benth. in Journ. Linn. Soc.* i. (1857) 90.—China.
lucidum, Poir. Encyc. Suppl. ii. 714 = sempervirens.
nitidum, Michx. Fl. Bor. Am. i. 120 = sempervirens sempervirens, *Ait. Hort. Kew.* ed. II. ii. 64.—Am. bor.

GEMBANGA, Blume, in Flora, viii. (1825) 580, 678 = **Corypha**, Linn. (Palm.).
rotundifolia, Blume, l. c. = C. umbraculifera.

GEMELLA, Hill, Veg. Syst. ii. 70** (1761) = **Coreopsis** ! Linn. (Compos.).

GEMELLA, Lour. Fl. Cochinch. 648 (1790) = **Schmidelia**, Linn. (Sapindac.).
trifoliata, Lour. l. c. 649 = S. Gemella.

GEMELLARIA, Pinell, ex Antoine, Bromel. 44 (1884) = **Karatas**, Mill. (Bromeliac.).
Innocentii, Pinell, l. c. = K. Innocentii.

GEMINARIA, Rafin. Cat. 14 (1824) = **Phyllanthus**, Linn. (Euphorbiac.).
obovata, Rafin. l. c. = P. carolinensis.

GEMMARIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 = **Conyza** ! Linn. (Compos.).

GEMMARIA, Salisb. Gen. Pl. Fragm. 127 (1866) = **Hessea**, Herb. (Amaryllid.).
gemmata, Salisb. l. c. = H. gemmata.

GENDARUSSA, Nees, in Wall. Pl. As. Rar. iii. 76 (1832) = **Justicia**, Linn. (Acanthac.).
abyssinica, Nees, in DC. Prod. xi. 780 = J. glabra.
Adhatoda, Steud. Nom. ed. II. i. 668 = Adhatoda Vasica.
arenaria, Steud. l. c. = J. trinervia.
argyrostachya, Steud. l. c. = J. nigrostachya.
Betonica, Nees, ex Steud. Nom. ed. II. i. 668; et in Linnaea, xv. (1841) 366 = J. capensis.
cordata, Hochst. ex Nees, in DC. Prod. xi. 879 = J. cordata.
cuneata, Nees, ex Steud. Nom. ed. II. i. 668; et in Linnaea, xv. (1841) 367 = J. cuneata.
debilis, Nees, in Linnaea, xvi. (1842) 302 = J. debilis.
decussata, Nees, in Wall. Pl. As. Rar. iii. 104 = J. decussata.
densiflora, Hochst in Flora, xxviii. (1845) 71.—Afr. austr.
dichotoma, Hochst. ex Nees, in DC. Prod. xi. 500 = J. glabra.
diosmophylla, Nees, l. c. 781 = Adhatoda diosmophylla.
diversiformis, Nees, l. c. 399, 781 = J. glauca.
foliolosa, Presl, Bot. Bemerk. 95 = Adhatoda cuneata.
furcata, Nees, in Wall. Pl. As. Rar. iii. 105 = J. furcata.
hyssopifolia, Webb & Berth. Phyt. Canar. iii. 159 = J. hyssopifolia.
incana, Nees, in Linnaea, xv. (1841) 367 = Adhatoda incana
leptantha, Nees, l. c. 372 = Adhatoda leptantha.
Linaria, Nees, ex T. Anders. in Journ. Linn. Soc. vii. (1864) 49 = J. Linaria.
lithospermifolia, Presl, Bot. Bemerk. 95 = Adhatoda lithospermifolia.
micrantha, Wall. ex Voigt, Hort. Suburb. Calc. 489 = J. micrantha.
mollis, Hochst. in Flora, xxviii. (1845) 71 (Adhatodæ sp.).—Afr. austr.
mollis, Presl, Bot. Bemerk. 95 = J. mollis.
Neesiana, Nees, in Wall. Pl. As. Rar. iii. 105 = J. Neesiana.
nigerrhensis, Steud. Nom. ed. II. i. 668 = J. nigerrhensis.
odora, Nees, ex Steud. l. c.; Presl, Bot. Bemerk. 95 = J. odora.
orchioides, Nees, ex Steud. l. c.; et in Linnaea, xv. (1841) 369 = J. orchioides.
orientalis, Nees, in Wall. Pl. As. Rar. iii. 104 = J. glauca.
palustris, Hochst. in Flora, xxiv. (1841) i. Intell. 43 = J. palustris !
patula, Nees, in Linnaea, xv. (1841) 367 = Adhatoda patula.
pedunculosa, Presl, Bot. Bemerk. 95 = Dianthera americana.
petiolaris, Presl, l. c. = J. petiolaris.
prostrata, Steud. Nom. ed. II. i. 668, sphalm. = seq.
protracta, Nees, in Linnaea, xv. (1841) 371 = Adhatoda protracta.
prunellæfolia, Hochst. in Flora, xxviii. (1845) 71 (Adhatodæ sp.).—Afr. austr.
pygmæa, Nees, in Linnaea, xv. (1841) 369 = Adhatoda pygmæa.
quadrifaria, Nees, in Wall. Pl. As. Rar. iii. 105 = J. quadrifaria.
ramosissima, Steud. Nom. ed. II. i. 668 = J. Petonica.
Schimperiana, Hochst. in Flora, xxiv. (1841) i. Intell. 24 = Adhatoda Schimperiana.
sessilis, Presl, Bot. Bemerk. 95 = Dianthera sessilis.
sumatrana, Miq. Fl. Ind. Bat. ii. 832 = J. ptychostoma.

GENDARUSSA :—
tranquebariensis, Nees, in Wall. Pl. As. Rar. iii. 105 = justicia tranquebariensis.
tridentata, Presl, Bot. Bemerk. 95 = Ramusia tridentata.
trinervia, Nees, ex Steud. Nom. ed. II. i. 668 = J. trinervia.
variegata, Hochst. ex Nees, in DC. Prod. xi. 385 = Adhatoda variegata.
vasculosa, Nees, in Wall. Pl. As. Rar. iii. 104 = J. vasculosa.
ventricosa, Nees, in Wall. l. c. = J. ventricosa.
virgata, Wall. Cat. n. 7176 = Dianthera virgata.
vulgaris, Nees, in Wall. Pl. As. Rar. iii. 104 = J. Gendarussa.
wissadensis, D. Dietr. Syn. Pl. i. 69, sphalm. = seq.
wynaadensis, Nees, in Wall. Pl. As. Rar. iii. 104 = J. wynaadensis.

GENERSICHIA, Heuff. in Flora, xxvii. (1844) ii. 528 = Carex, Linn. (Cyperac.).
obtusata, Heuff. l. c. = C. obtusata.

GENESPHYLA, Rafin. Sylva Tellur. 92 (1838) = seq.

GENESPHYLLA, L'Hérit. Sert. Angl. 29 (1788) = Phyllanthus, Linn. (Euphorb.).
asplenifolia, L'Hérit. l. c. = P. latifolius.
speciosa, Rafin. Sylva Tellur. 92 = P. speciosus.

GENETYLLIS, DC. Prod. iii. 209 (1828) = Darwinia, Rudge (Myrtac.).
affinis, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 155 = D. diosmoides.
agathosmoides, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 322 = D. vestita.
alpestris, Lindl. in Mitch. Three Exped. ii. 178 = Lhotskya genetylloides.
citriodora, Endl. in Enum. Pl. Hueg. 47 = D. citriodora.
diosmoides, DC. Prod. iii. 209 = D. diosmoides.
Drummondii, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 155 = D. Drummondii.
fimbriata, Kipp. in Journ. Linn. Soc. i. (1857) 49 = D. fimbriata.
fuchsioides, Hort. ex Lescuy. in Hérinog. Hortic. Franç. (1864) t. 15 = D. Hookeriana.
helichrysoides, Meisn. in Journ. Linn. Soc. i. (1857) 37 = D. helichrysoides.
Hookeriana, Meisn. l. c. = D. Hookeriana.
hypericifolia, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 323 = D. Hookeriana.
isjotyla, Turcz. l. c. = D. Meisneri ?
macrostegia, Hook. Bot. Mag. t. 4860 = D. Hookeriana.
macrostegia, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 18 = D. macrostegia.
Meisneri, Kipp. in Journ. Linn. Soc. i. (1857) 49 = D. Meisneri.
micropetala, F. Muell. Fragm. i. 12 = D. micropetala.
oederioides, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 18 = D. oederoides.
oxylepis, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 324 = D. Meisneri.
pauciflora, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 17 = Chamælaucium ciliatum.
pimeloides, F. Muell. Fragm. ii. 169 = D. citriodora.
pinifolia, Schau. Monog. Myrt. Xerocarp. i. 84 = D. pinifolia.
purpurea, Schau. l. c. 27 = D. purpurea.
sanguinea, Meisn. in Journ. Linn. Soc. i. (1857) 38 = D. sanguinea.
Schaumanni, F. Muell. Fragm. i. 12 = D. Schaumanni.
speciosa, Meisn. in Journ. Linn. Soc. i. (1857) 36 = D. speciosa.
spuorosa, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 323 = D. fimbriata.
thymoides, Schau. Monog. Myrt. Xerocarp. i. 83 = D. thymoides.
tulipifera, Schau. ex Hook. Bot. Mag. t. 4856 = D. macrostegia.
vestita, Endl. in Enum. Pl. Hueg. 47 = D. vestita.
virescens, Meisn. in Journ. Linn. Soc. i. (1857) 38 = D. virescens.

GENIANTHUS, Hook. f. Fl. Brit. Ind. iv. 15 (1883).
ASCLEPIADEÆ.
crassifolius, *Hook. f. l. c.* 16.—Silhet.
laurifolius, *Hook. f. l. c.*—Ind. or.
Maingayi, *Hook. f. l. c.*—Singapore.

GENIOSPORUM, Wall. ex Benth. in Bot. Reg. sub t. 1300 (1830). *LABIATÆ*, Benth. & Hook. f. ii. 1172.
axillare, Benth. in Wall. Pl. As. Rar. ii. 18 = Melissa axillaris.
elongatum, *Benth. Lab. Gen. et Sp.* 21.—Ind. or.
gracile, Benth. l. c. = prostratum.
holochilum, *Hance*, in *Journ. Bot.* xvii. (1879) 13.—China.
madagascarense, *Benth. Lab. Gen. et Sp.* 20.—Madag.
Pallisoti, *Benth. l. c.* 22.—Afr. trop.
parviflorum, Benth. in Wall. Pl. As. Rar. ii. 18 ; et Cat. n. 2750 = Mesona Wallichiana.
prostratum, *Benth. in Bot. Reg.* sub t. 1300 ; *et in Wall. Pl. As. Rar.* ii. 18.—Ind. or.
strobiliferum, *Wall. Pl. As. Rar.* ii. 18.—Reg. Himal.
thymifolium, *Benth. Lab. Gen. et Sp.* 21.—Madag.

GENIOSTEMON, Engelm. & Gray, in Proc. Am. Acad. xvi. (1881) 104. *GENTIANEÆ.*
Coulteri, *Engelm. & Gray, l. c.*—Mexic.
Schaffneri, *Engelm. & Gray, l. c.*—Mexic.

GENIOSTEPHANUS, Fenzl, in Flora, xxvii. (1844) 312, absque desc. = Trichilia, Linn. (Meliac.).
tomentosus, Fenzl, l. c. = T. emetica.

GENIOSTOMA, Forst. Char. Gen. 24. t. 12 (1776). *LOGANIACEÆ*, Benth. & Hook. f. ii. 791.
ANASSER, Just. Gen. 150 (1789).
ASPILORIUM, Soland. ex A. Cunn. in Ann. Nat. Hist. i. (Sept. 1838) 45.
HÆMOSPERMUM, Reinw. in Blume, Bijdr. 1018 (1825).
LASIOSTOMUM, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. (1850) 239.
acuminatum, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 816 = Urophyllum streptopodium.
angustifolium, Bouton, ex DC. Prod. ix. 28 = borbonicum !
asterium, *A. Gray, in Proc. Am. Acad.* iv. (1860) 321.—Ins. Society.
australianum, *F. Muell. Fragm.* v. 19.—Austral.
Balansaianum, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 248.—N. Caled.
borbonicum, *Spreng. Syst* i 588.—Ins. Mascar.
brasiliense, Spreng. l. c. = Haynaldia thapsoides.
celastrinum, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 247.—N. Caled.
cordifolium, *Bot. Hort. Maurit.* 215.—Ins. Maurit.
crassifolium, *Benth. in Journ. Linn. Soc.* i. (1857) 96.—Ins. Philipp.
Cumingianum, *Benth. l. c.* 97.—Ins. Philipp.
Cyttandræ, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 320.—Ins. Sandvic.
densiflora, *Baill. l. c.* 264.—N. Caled.
erythrosperma, *Baill. l. c.* 263.—N. Caled.
fagraeoides, *Benth. in Journ. Linn. Soc.* i. (1857) 96.—Mexic.
febrifugum, Spreng. Syst. i. 588 = Strychnos Pseudo-Quina.
foetens, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 247.—N. Caled.
Haemospermum, Steud. Nom. ed. II. i. 669 = rupestre.
lanceolatum, Boj. Hort. Maurit. 215 = borbonicum.
Lasiostemon, *Blume, Mus. Bot. Lugd. Bat.* i. 239.—Sumatra.
ligustrifolium, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 47.—N. Zel.
micranthum, *DC. Prod.* ix. 27.—Ins. Marian.
microphyllum, *Sonn. in Bonplandia*, x. (1862) 37, nomen ; *et Fl. Vit.* 104.—Ins. Fiji.
montanum, *Zoll. & Mor. Syst. Verz. Zoll.* 58.—Java.
Novæ-Caledoniæ, *Vieill. ex Baill. in Bull. Soc. Linn. Par.* i. (1880) 263.—N. Caled.
oboratum, Boj. ex Benth. in Journ. Linn. Soc. i. (1857) 96 = borbonicum.
ovatum, Boj. Hort. Maurit. 215 = borbonicum.
Pancheri, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 264.—N. Caled.
parviflorum, Boj. Hort. Maurit. 215 = pedunculatum.
pedunculatum, *Boj. l. c.*—Ins. Maurit.
petiolosum, *C. Moore & F. Muell. Fragm.* vii. 28 : ix. 193.—Ins. Dom. Howe.

GENIOSTOMA :—
phyllanthoides, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 264.—N. Caled.
Remyana, *Baill. l. c.* 240.—Ins. Sandvic.
reticulata, *Blume, Mus. Bot. Lugd. Bat.* i. 239.—Java.
rupestre, *Forst. Char. Gen.* 24. t. 12.—N. Hebrid.
rupestre, A. Rich. & Less. Voy. Astrol. Bot. 207 = ligustrifolium.
spinescens, *Spreng. Syst.* i. 588.—Bras.
thymeleaeoides, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 263.—N. Caled.
vestitum, *Baill. l. c.* 248.—N. Caled.
Vieillardi, *Baill. l. c.* 264.—N. Caled.

GENIPA, [Tourn.] Linn. Syst. ed. I (1735). *RUBIACEÆ*, Benth. & Hook. f. ii. 90.
THEVENOTIA et THEVETIA, Linn. Gen. ed. I. 350 (1737).
americana, *Linn. Syst.* ed. X. 931.—Am. austr.
arborea, Baill. Hist. Pl. vii. 374 = Gardenia gummifera.
barbata, *Frost, Symb. Bot.* ii. 13. t. 60.—Bras.
brasiliana, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v (1834) 244.—Bras.
brasiliensis, Baill. Hist. Pl. vii. 374 = Gardenia brasiliensis.
buffalina, *Lour. Fl. Cochinch.* 149.—Cochinch.
campanulata, Baill. Hist. Pl. vii. 371 = Gardenia campanulata.
caprnsii, Baill. l. c. 379, 388 = Burchellia capensis.
Caruto, *H. B. & K. Nov. Gen. et Sp.* iii. 407.—Am. austr.
cloalifolia, *Grosh. Fl. Brit. W. Ind.* 317.—Ind. occ.
dumetorum, Baill. Hist. Pl. vii. 372 = Randia dumetorum.
edulis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 107 = Alibertia edulis.
esculenta, Lour. Fl. Cochinch. 149 = Randia edulis.
ferax, *Lour. l. c.*—China.
florida, Baill. Hist. Pl. vii. 367, 379 = Gardenia jasminoides.
grandiflora, Baill. l. c. 374 = Gardenia grandiflora.
grandifolia, Pers. Syn. i. 198 = americana.
gummifera, Baill. Hist. Pl. vii. 379 = Gardenia gummifera.
humilis, *Vell. Fl. Flum.* 94 ; ii. t. 142.—Bras.
Jovis-tonantis, Baill. Hist. Pl. vii. 380 = Gardenia Jovis-tonantis.
labiata, Smeathm. ex DC. Prod. iv. 391 = Bertiera Pomalium.
longiflora, Baill. Hist. Pl. vii. 379 = Randia longiflora.
malleifera, Baill. l. c. = Randia malleifera.
Merianæ, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guian.
Mussaendæ, Baill. Hist. Pl. vii. 379 = Randia Mussaendæ.
oblongifolia, *Ruiz & Pav. Fl. Per. et Chil.* ii. 67. t. 220.—Peruv.
Pavetta, Baill. Hist. Pl. vii. 379 = Stylocoryne Webera. pubescens, *DC. Prod.* iv. 379.—Cuba.
radicans, Baill. Hist. Pl. vii. 379 = Gardenia jasminoides.
reunifera, Baill. l. c. 380 = Gardenia lucida.
Rothmannia, Baill. l. c. 379 = Gardenia Rothmannia.
Rutenbergiana, *Baill. ex Vathe, in Bremen, Abh.* ix. (1885) 118.—Afr. trop.
striffora, *DC. Prod.* iv. 378.—Bras.
Thunbergia, Baill. Hist. Pl. vii. 308 = Gardenia Thunbergia.

GENIPELLA, Rich. ex DC. Prod. iv. 443 (1830) = Alibertia, Rich. (Rubiac.).

GENISTA, Linn. Syst. ed. I (1735). *LEGUMINOSÆ*, Benth. & Hook. f. i. 482.
ARGELASIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 357.
ASTEROCYTISUS, Schur, ex Fuss, Fl. Transs. 154 (1866).
AVORNELA, Rafin. Sylva Tellur. 24 (1838).
BOELIA, Webb, Otia Hisp. 20. t. 15, 16 (1858).
CHAMÆSPARTIUM, Adans. Fam. ii. 321 (1763).
CORNIOLA, Presl, in Abh. Boehm. Ges. Wiss. V. iii. (1846) 566 [Bot. Bemerk. 136 (1844)].
COROTHAMNUS, Presl, l. c. (1846) 567 [l. c. 137 (1844)].
DENDROSPARTUM, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 123.
DRYMOSPARTUM, Presl, in Abh. Boehm. Ges. Wiss. V. iii. (1846) 567 [Bot. Bemerk. 138 (1844)].

GENISTA :—

Echinosparton, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 268.

Euteline, Rafin. Sylva Tellur. 24 (1838).

Genistella, Tourn. ex Rupp. Fl. Jen. ed. Hall. 266 (1745).

Genistoides, Moench, Meth. 132 (1794).

Gonocytisus, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 158.

Listera, Adans. Fam. ii. 321 (1763).

Lugaion, Rafin. Sylva Tellur. 22 (1838).

Lygos, Adans. Fam. ii. 321 (1763).

Phyllonotrys, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195.

Pterospartum, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 440 (1880).

Retama, Boiss. Voy. Bot. Espagne, 143 (1839)

Saltzwedelia, Gaertn. Mey. & Scherb. Fl. Wett. ii. 480 (1800).

Scorpius, Moench, Meth. 134 (1794).

Spartidium, Pomel, Nouv. Mat. Fl. Atl. 178 (1874).

Spartium, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 285. t. 16.

Sybsone, Griseb. Spicil. Fl. Rumel. i. 5 (1843).

Voglera, Gaertn. Mey. & Scherb. Fl. Wett. ii. 480 (1800).

acanthoclada, *DC. Mém. Lég.* vi. 208.—Graec.; As. Min.

acutifolia, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = Spartium junceum.

aegyptiaca, *Spreng. Nov. Prov.* 20.—Aegypt.

aethnensis, *DC. Prod.* ii. 150.—Aetna.

alba, Lam. Encyc. ii. 628 = Cytisus albus.

albida, Bess. Enum. Pl. Volh. 98 = tinctoria.

albida, *Willd. Sp. Pl.* iii. 942.—Ross.; As. Min.

algarbiensis, Brot. Fl. Lusit. ii. 89 = hirsuta.

alpicola, Schur, Enum. Pl. Transs. 145 = tinctoria.

alpina, Spach, in Ann. Sc. Nat. Sér. III. iii. (1844) 247 = acanthoclada.

alpina, Spach, l. c. iii. (1845) 157 = Laburnum alpinum!

Amarelle, *Lepech. Reise Prov. Russ. Reich.* iii. 42.—Kossia.

americana, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 157 = Spartium junceum.

amsanctica, Auct. vide anxantica.

anatolica, *Boiss. Diagn.* Ser. I. ii. 8.—As. Min.; Syria.

anclatrocytis, *Spach, in Ann. Sc. Nat.* Sér. III. iii. (1845) 105.—Lusitan.

andicola, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 178.—Chili.

anglica, *Lam. Sp. Pl.* 710.—Europ.

angulata, Lam. Encyc. ii. 624 = parviflora.

angulata, Poir. Encyc. Suppl. v. 688; DC. Prod. ii. 152 = Cytisus scoparius.

antiochia, Boiss. Diagn. Ser. I. ix. 4 = patula.

anxantica, Griseb. Spicil. Fl. Rumel. i. 3 = elatior.

anxantica, *Tenore, Fl. Nap. Prod.* 41.—Ital.

apetala, *Spach, in Ann. Sc. Nat.* Sér. III. iii. (1845) 157.—Hab.!

aphylla, DC. Prod. ii. 149 = Eremosparton aphyllum.

aprutia, Presl, Bot. Bemerk. 136 = Micheliî.

arborea, Spreng. Syst. iii. 220 = Cytisus arboreus.

arborescens, Mill. ex Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 154 = Brya Ebenus.

arcuata, Koch, Syn. Fl. Germ. ed. I. 154 = sylvestris.

argentea, Noulet, Fl. Bass. Sous-Pyr. 146; et in Ann. Sc. Nat. Sér. II. ix. (1838) 130 = Argyrolobium Linnaeum.

aristata, *Presl, Delic. Prag.* 84.—Sicilia.

armata, Poir. Encyc. Suppl. ii. 721 = Wiborgia spinescens.

armeniaca, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 118 = albida.

asphaltoides, Boiss. Voy. Bot. Espagne, 141 = baetica.

asparagoides, *Lam. Encyc.* ii. 620 (asparagoides).—Afr. bor.

atlantica, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 366.—Maroc.

Ascheri, *Boiss. Diagn.* Ser. I. ii. 7.—Cappadoc.

baetica, *Spach, in Ann. Sc. Nat.* Sér. III. iii. (1845) 118.—Hispan.

balearica, Fourr. in Willk. & Lange, Prod. Fl. Hisp. iii. 634 = tinctoria.

barbara, *Munby, It. Algér.* 74.—Alger.

Barmudiali, *Granito, in Mem. Acad. Cienc. Madr.* ii. (1850) 461.—Hispan.

barbuvicina, *Lange, Pl. Nov. Hisp. Fasc.* i. t. 1. t.—Hispan.

GENISTA :—

biflora, Poir. Encyc. Suppl. ii. 717; DC. Prod. ii. 146 = Cytisus Fontanesii.

Bivonae, *Presl. Delic. Prag.* 83.—Sicil.

Boissieri, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 358.—Hispan.

Bourgaei, Spach, ex Nym. Consp. 152 = florida.

bracteolata, Link, Enum. Hort. Berol. ii. 224 (= Cytisi sp.).—Hab.?

brachypoda, Willd. ex Spreng. Syst. iii. 176 = stylosa.

britannica, Hort. ex Reichb. Fl. Germ. Excurs. 521 = anglica.

Broteri, *Poir. Encyc. Suppl.* ii. 720.—Lusitan.

Brugnieri, Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 248 = acanthoclada.

**cafifolia*, Burm. f. Fl. Cap. Prod. 21 = Cyclopia latifolia.

caespitosa, C. Koch, in Linnaea, xix. (1847) 60 = lydia.

calycoda, Schur, in Verh. Naturf. Ver. Brunn, xv. 11. (1877) 166 = tinctoria.

campestris, Janka, in Linnaea, xxx. (1859–60) 562 = tinctoria.

canariensis, Bory & Chaub. Nouv. Fl. Pélop. 46 = Cytisus candicans.

canariensis, DC. Prod. ii. 145 = Cytisus monspessulanus.

canariensis, Linn. Sp. Pl. 709 = Cytisus canariensis.

candicans, Linn. Cent. I. 22; Amoen. Acad. iv. 284 = Cytisus candicans.

cantabrica, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 149 = tridentata.

capensis, Spach, l. c. 155 = Rafnia opposita.

capitata, Spach, l. c. 157 = clavata.

capitellata, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iii. (1856) 672.—Alger.

cappadocica, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 245.—Cappadocia.

carinalis, *Griseb. Spicil. Fl. Rumel.* i. 3.—Graec.; Macedon.

carpetana, *Lrestche, ex Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 287.—Hispan.

Cassia, Boiss. Diagn. Ser. I. ix. 3 = anatolica.

cephalantha, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 254.—Alger.

cinerascens, Lange, in Kjoeb. Vidensk. Meddel. (1865) 168 (Pugill. 858) = seq.

cinerea, *DC. Fl. Fr.* iv. 494.—Europ. austr. occ.

cinerea, Reichb. Fl. Germ. Excurs. 521 = Villarsi

cinerea, Willk. ex Lange, Notes Fl. 153 = florida.

cirtensis, *Pomel, Nouv. Mat. Fl. Atl.* 177.—Alger.

clavata, *Poir. Encyc. Suppl.* ii. 717.—Marocc.

comosa, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 132 = humifusa.

congesta, *Poir. Encyc. Suppl.* v. 688; *DC. Prod.* ii. 149.—Ins. Teneriffa.

contaminata, Poir. Encyc. Suppl. ii. 721 = Aspalathus corymbosa.

coriacea, *Kit. in Linnaea*, xxxii. (1863) 605.—Hungar.

corsica, *DC. Fl. Fr. Suppl.* 548.—Ins. Corsica.

cotinifolia, Burm. f. Fl. Cap. Prod. 21 = Rafnia elliptica!

crebripina, *Pomel, Nouv. Mat. Fl. Atl.* 176.—Alger.

cretica, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = Anthyllis Hermanniae.

Cumingii, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 178 = Anarthrophyllum Cumingii.

Cupani, *Guss. Adnot. Cat. Pl. Boccad.* 9.—Sicilia.

cuspidata, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 157 = hirsuta.

cuspidata, Steud. Nom. ed. II. i. 669 = seq.

cuspidosa, DC. Prod. ii. 147 = Lebeckia pilulobus.

cytisoides, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 155 = Lebeckia cytisoides.

dalmatica, Bartl. & Wendl. f. Beitr. ii. 74 = aristata.

dalmatica, Tenore, Syll. Fl. Neap. 344 = Micheliî.

dasycarpa, *Ball, in Journ. Linn. Soc.* xvi. (1878) 308. —Maroc.

derfjetana, Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 270 = Tournefortii.

decumbens, Durande, Fl. Bourg. i. 299 = Cytisus decumbens.

decumbens, Willd. Sp. Pl. iii. 941 = Cytisus procumbens.

defoliata, Lam. Fl. Fr. iii. 619 = monosperma.

Delarbrei, Lecoq & Lam. Cat. Pl. Cent. Fr. 125 = tinctoria.

GENISTA :—

delphinensis, Verl. in Bull. Soc. Stat.-Isère, xiv. (1872) 77 = sagittalis.

densa, Poir. Encyc. Suppl. ii. 721 = Lebeckia densa.

depressa, *Bieb. Fl. Taur. Cauc. Suppl.* 400.—Tauria.

depressa, Griseb. Spicil. Fl. Rumel. i. 2 = Friwaldskyi.

depressa, Tenore, Fl. Nap. v. 96. t. 171. f. 2 = humilis.

desiderata, DC. Prod. ii. 152 = Anarthrophyllum desideratum.

diffusa, Tenore, Fl. Nap. ii. 129 = humilis.

diffusa, Willd. Sp. Pl. iii. 942 = Cytisus decumbens.

dilcolor, *Webb, ex Lowe, Man. Fl. Mad.* i. 126.—Ins. Canar.

dispersa, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 155 = Cytisus albus.

divaricata, Link, Enum. Hort. Berol. ii. 224 = Adenocarpus intermedius.

dracunculoides, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 139 = patula.

Durieui, *Spach, l. c.* iii. (1844) 271.—Algeria.

Echinus, Spach, l. c. 249 = acanthoclada.

Echinus, Willd. ex Spreng. Syst. iii. 221 = aspalathoides.

elata, Wender. in Linnaea, xv. (1841) Litt. 100 = tinctora.

elatior, *Koch, Syn. Fl. Germ.* ed. II. 441.—Europ. austr. or.

elegans, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 178. t 103 = Anarthrophyllum elegans.

elliptica, Kit. in Linnaea, xxxii. (1863) 606 = Cytisus procumbens.

elliptica, *Steud. Nom.* ed. II. i. 669.—Hispan

ephedroides, Fourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 484 = cinerea.

ephedroides, *DC. Mem. Lég.* vi. 210. t. 36.—Ins. Sardin.

equisetiformis, *Spach, in Ann. Sc. Nat.* Sér. III. iii. (1845) 148.—Hispan.

erictorum, Hoffmgg. ex Spreng. Syst. iii. 221 = triscanthos.

erinacea, Gillib. Bot. Prat. ii. 239; ex DC. Prod. ii. 146 = horrida.

erinocarpa, Kunze, in Flora, xxix. (1846) 737 = Cytisus Kunzeanus.

erioclada, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 264.—Alger.

exaltata, Link, ex DC. Prod. ii. 151 = polygaloefolia.

falcata, *Brot. Phyt. Lusit.* 153. t. 55.—Lusitan.

fasciata, *Decne. in Ann. Sc. Nat.* Sér. iv. (1855) 200.—Syria.

ferox, *Poir. Encyc. Suppl.* ii. 708.—Afr. bor.

filifolia, Licht. ex Walp. in Linnaea, xv. (1839) 460 = Viminaria denudata.

filiramea, *Pomel, Nouv. Mat. Fl. Atl.* 318.—Alger.

florida, Asso, Syn. Pl. Arag. 94 = cinerea.

florida, *Linn. Syst.* ed. X. 1157.—Reg. Mediterr. occ.

foliosa, Link, Enum. Hort. Berol. ii. 224 = Adenocarpus anagyrus.

fragrans, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 155 = Cytisus fragrans.

friedrichsthaliana, Presl, Bot. Bemerk. 136 = acanthoclada.

Friwaldskyi, *Boiss. Fl. Or.-rst.* ii. 45.—Rumelia.

frutescens, Schloss. & Vukot. Syll. Fl. Croat. 124 = tinctoria.

Gasparrini, Presl, Fl. Sic. Praef. p. xix = ephedroides.

Gayana, A. Gray, Bot. U. St. Expl. Exped. i. 397 (= Anarthrophyllum sp.).—Chili.

genuensis, Pers. Syn. ii. 287 = triangularis.

germanica, Brot. Fl. Lusit. ii. 90 = Tournefortii.

germanica, *Linn. Sp. Pl.* 710.—Europ.

gibraltarica, *DC. Prod.* ii. 148.—Fret. Hercul.

gibraltarica, Lowe, in Journ. Linn. Soc. v. (1860) 87 = tridens.

glabra, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 157 = Cytisus scoparius.

Godetii, Spach, l. c. 118 = albida.

gracilis, Poir. Encyc. Suppl. ii. 715 = virgata, *DC.*

gracilis, Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 256 = carinalis.

grandiflora, Spach, l. c. iii. (1845) 155 = Cytisus grandiflorus.

gymnoptera, Duby, ex Nym. Consp. 152 = aspalathoides.

Haenseleri, Boiss. Elench. 20 = Cytisus Haenseleri.

Halleri, Reyn. ex DC. Prod. ii. 152 = Cytisus decumbens.

GENISTA :—
herbacea, Lam. Fl. Fr. ii. 616 = sagittalis.
heteracantha, Schloss. & Vukot. Syll. Fl. Croat. 125 = germanica.
heterophylla, DC. Prod. ii. 152 = Cytisus heterophyllus.
hirsuta, Moench, Meth. 144 = Cytisus scoparius.
hirsuta, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 155 = Cytisus Millerl.
hirsuta, Tenore, Fl. Nap. ii. 130 = Michelii.
hirsuta, Tineo, ex Spach, in Ann. Sc. Nat. Sér. III. ii (1844) 260 = aristata.
hirsuta, *Vahl, Symb. Bot. i.* 51.—Hispan.
hispanica, Host [Fl. Austr. ii. 318 ?] ; ex Nym. Consp. 151 = aristata.
hispanica, *Linn. Sp. Pl.* 711.—Europ. occ.
hispanica, Wulf. in Jacq. Coll. ii. 165 = sylvestris.
holopetala, Reichb. ex Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 241 = radiata.
horrida, Boiss. ex Spach, l. c. 258 = Boissieri.
horrida, *DC. Fl. Fr.* iv. 500.—Europ. occ.
humifusa, Bout. ex Willk. & Lange, Prod. Fl. Hisp. iii. 437 = micrantha.
humifusa, Dicks. ex Nym. Consp. 158 = tinctoria.
humifusa, *Linn. Syst. ed.* X. 1157.—Europ. ; As. Min.
humifusa, Thore, Chlor. Land. 208 = pilosa.
humifusa, Vill. Prosp. 41 = Villarsii.
humifusa, Willd. Sp. Pl. iii. 942 = albida.
humifusa, Wulf. in Jacq. Coll. ii. 169 = Cytisus decumbens.
humilis, Poir. Encyc. Suppl. ii. 721 = Lebeckia humilis.
humilis, *Tenore, Syll. Fl. Neap.* 345.—Europ. austr.
hungarica, Kern. in Oester. Bot. Zeitschr. (1868) 140 = tinctoria.
Hystrix, *Lange, Pl. Nov. Hisp. fasc.* i. t. 1. 2.—Hispan.
incerta, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 3 = carinalis.
incubacea, Schur. Enum. Pl. Transs. 145 = tinctoria.
inermis, Gillb. Fl. Lituan. ii. 77 = tinctoria.
infesta, *G. Don, Gen. Syst.* ii. 150.—Sicilia.
inops, Boiss. & Bal. Diagn. Ser. II. vi. 42 = Jaubertii.
interrupta, Steud. Nom. ed. II. i. 670 = tinctanthoa.
involucrata, *Spach, in Ann. Sc. Nat.* Sér. III. iii. (1845) 120.—As. Min.
ischiocalda, *Pomel, Nouv. Mat. Fl. Alt.* 318.—Alger.
italica, *Lodd. ex G. Don, in Loud. Hort. Brit.* 280.—Italia.
januensis, Viv. Elench. Fl. Hort. Dinegro, 19 = triangularis.
Jaubertii, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 242.—As. Min.
Jussei, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 247 = Alhagi maurorum.
juncea, Scop. Fl. Carn. ed. II. ii. 50 = Spartium junceum.
juniperina, Meyen, Reise, i. 315 = Anarthrophyllum juniperinum.
juniperina, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 272.—Marocc.
lamprophylla, Spach, l. c. iii. (1845) 126 = lydia.
lanigera, Spach, l. c. 155 = Calycotome villosa.
lanuginosa, Spach, l. c. ii. (1844) 264 = hirsuta.
laricifolia, Burm. f. Fl. Cap. Prod. 21. (Quid ?).—Afr. austr.
lasiantha, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 147 = tridentata.
lasiocarpa, Spach, l. c. 185 = mantica.
leptoclada, *J. Gay*, l. c. Sér. II. vi. (1836) 129 = florida.
leptophylla, *Spach, l. c.* Sér. III. iii. (1845) 127.—Bithyn.
libanotica, *Boiss. Diagn.* Ser. I. ix. 3.—Syria.
linifolia, Linn. Sp. Pl. ed. II. 405 = Cytisus linifolius.
linifolia, Vill. Fl. Delph. 74 = cinerea.
liparioides, Boiss. Diagn. Ser. I. ii. 6 = Cytisus orientalis.
Lobelii, DC. Fl. Fr. iv. 499 = sapalathoides.
Lobelii, Urv. in Mém. Soc. Linn. Par. i. (1822) 341 = acanthoclada.
lucida, *Cambess. in Mém. Mus. Par.* xiv. (1827) 231. t. 14.—Ins. Balear.
lugdunensis, Jord. Pugill. 52 = horrida.
lusitanica, Boiss. Voy. Bot. Espagne, 140 = Boissieri.
lusitanica, Desf. Tabl. ed. 1. 184, fide DC. Fl. Fr. iv. 500 = horrida.
lusitanica, *Linn. Sp. Pl.* 711.—Lusiten.
lusitanica, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 155 = Cytisus grandiflorus.

GENISTA :—
lydia, *Boiss. Diagn.* Ser. I. ii. 8.—Europ. or. ; As. Min.
madagascariensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 195.—Madag.
madrensis, *Lowe, Man. Fl. Mad.* i. 123.—Ins. Madera.
mantica, *Pollini, Cat. Sem. Vernon.* (1814) 18 ; *Spreng. Pugill.* ii. 78.—Ital.
marginata, Bess. ex Ledeb. Fl. Ross. i. 517 = tinctoria.
Mayeri, Janka, in Oestr. Bot. Zeitschr. ix. (1859) 61 = ovata.
melia, *Boiss. Diagn.* Ser. I. ix. 2.—Archip. Graec.
microcalycina, Spach, in *Ann. Sc. Nat.* Sér. III. ii. (1844) 259.—Europ. austr.
micrantha, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 3 = carinalis.
micrantha, *Ortog. Hort. Matr. Des.* 68. t. 10.—Hispan.
microcephala, *Coss. & Dur. in Ann. Sc. Nat. Sér.* IV. bv. (1855) 248.—Algeria.
microphylla, *DC. Prod.* ii. 146.—Ins. Canar.
microphylla, Moris, Stirp. Sard. Elench. 13 = Moriali.
microphylla, Webb & Berth. Phyt. Canar. ii. 42 = conqusta.
minor, Guillon, ex Verl. in Bull. Soc. Bot. Fr. xxiii. (1876) p. lxxx = sagittalis.
minor, Lam. Fl. Fr. ii. 615 = anglica.
mollis, *DC. Prod.* ii. 145.—Marocc.
monosperma, Delile, Fl. Aegypt. 21 = G. Raetam.
monosperma, *Lam. Encyc.* ii. 616.—Reg. Mediterr. occ.
monosperma, Link, in Buch, Beschr. Canar. Ins. 156 ; Lindl. Bot. Reg. t. 1918 = rhodorhizoides.
Monthyon, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 119 = albida.
Morisii, *Colla, Herb. Pedem.* ii. 65.—Ins. Sardin.
multiarctesta, *Tausch, in Flora*, xxi. (1838) i. *Beibl.* 78.—Kals. ?
multicaulis, *Lam. Encyc* ii. 617.—Ins. Balear.
multiflora, (Hort. Petrop.) Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 153 = Cytisus albus.
murcica, *Coss. Notes Crit.* 101.—Hispan.
myrianthe, *Ball. in Journ. Bot.* xi. (1873) 303.—Marocc.
myrtifolia, Burm. f. Fl. Cap. Prod. 21 (Quid ?).—Afr. austr.
nervata, Hoppe, ex Griseb. Spicil. Fl. Rumel. i. 4 = anxantica.
nervata, Kit. ex DC. Prod. ii. 151 = ovata.
nervosa, Kit. in Linnaea, xxiii. (1850) 605 = ovata.
nitens, *Steud. Nom.* ed. II. i. 670.—Ins. Teneriff.
nodosa, *Tausch, in Flora*, xxi. (1838) 748.—Austria.
nubigena, Link, in Buch, Beschr. Canar. Ins. 156 = Cytisus fragrans.
nuda, *Steud. Nom.* ed. II. i. 670.—Mauritia.
numidica, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 244.—Algeria.
obtusiramea, *J. Gay, l. c.* Sér. II. vi. (1836) 344.—Hispan.
odorata, Moench, Meth. 144 = Spartium junceum.
odoratissima, Pourr. in Willk. & Lange, Prod. Fl. Hisp. iii. 437 = micrantha.
odoratissima, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 155 = Spartium junceum.
Oliveri, Spach, l. c. ii. (1844) 266 = anstolices.
oretana, Webb, ex Willk. & Lange, Prod. Fl. Hisp. iii. 434 = florida.
orientalis, Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 267 = anatolica.
osmarensis, *Coss. in Bull. Soc. Bot. Fr.* xx. (1873) 245.—Alger.
ovata, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = Kalinia cuneifolia.
ovata, *Waldst. & Kit. Pl. Rar. Hung.* i. 86. t. 84.—Europ. or.
oxycedrus, *Pomel, Nouv. Mat. Fl. Alt.* 319.—Alger.
Palvae, *Lowe, Man. Fl. Mad.* i. 125.—Ins. Mader.
parviflora, Brot. Fl. Lusit. ii. 87 = broteri.
parviflora, *Potr. Encyc. Suppl.* ii. 715 ; *DC. Prod.* ii. 145.—As. Min.
parviflia, G. Don, Gen. Syst. ii. 150 = Moriali.
patens, DC. Prod. ii. 148 = Cytisus patens.
patula, *Bieb. Fl. Taur. Cauc.* ii. 148.—As. Min.
pedunculata, L'Hérit. Stirp. Nov. 184 = Cytisus decumbens.
peloponnensiaca, Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 249 = acanthoclada.

GENISTA :—
pendulina, Lam. Encyc. ii. 623 ; Poir. Voy. Barb. ii. 208 = Cytisus arboreus.
Perrymondi, Loisel. Fl. Gall. ed. II. ii. 105 = mantica.
persica, Poir. Encyc. Suppl. ii. 720 = Crotalaria furfuracea.
Pestalozzae, Boiss. Diagn. Ser. I. ix. 4 = albida.
Philippi, Lindl. in Gard. Chron. (1841) 365 = canariensis.
pilocarpa, *Link, Enum. Hort. Berol.* ii. 223.—Hab. ? *pilosa*, Ledeb. Fl. Ross. i. 516 = albida.
pilosa, *Linn. Sp. Pl.* 710.—Europ. austr.
pinastrifolia, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = Lebeckia sepiaria.
Pollinii, Spreng. ex Pollini, Fl. Veron. ii. 458 = mantica.
polyanthus, *De Roem. ex Willk. in Linnaea*, xxv. (1852) 20.—Hispan.
polygalaefolia, *DC. Prod.* ii. 151.—Lusitan.
polygaloaphylla, Brot. Fl. Lusit. 86 = praec.
polytricha, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 132 = ptilophylla.
Pomeli, *Maréz & Vigin. Cat. Pl. Baltar.* 70.—Ins. Balear.
pontica, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 127 = lydia.
pratensis, Pollini, Fl. Veron. ii. 457 = tinctoria.
procera, Poir. Encyc. Suppl. v. 689 = Cytisus proceras.
procumbens, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 3 = Frivaldskyi.
procumbens, Waldst. & Kit. ex Willd. Sp. Pl. iii. 940 = Cytisus procumbens, *Spreng.*
prostrata, Lam. Encyc. ii. 618 = Cytisus decumbens.
pseudopilosa, *Coss. Notes Crit.* 102.—Reg. Mediterr. occ.
pteroclada, *Boiss. ex Spach, in Ann. Sc. Nat. Sér.* III. iii. (1845) 154.—Syria.
ptiliophylla, *Spach, l. c.* 131.—Bithyn.
pubescens, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 4 = anxantica.
pubescens, Lange, ex Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 137 = tinctoria.
pulchella, Vis. in Flora, xiii. (1830) 51 = Villarsii.
pulverulenta, Flech. & Mey. in Ann. Sc. Nat. Sér. III. iv. (1845) 86. t. 1 = albida.
pungens, Poir. Encyc. Suppl. ii. 720 = Lebeckia pungens.
purgans, Linn. Syst. ed. X. 1157 ; Lam. Fl. Fr. ii. 618 = Cytisus purgans.
quadriflora, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 388.—Alger.
radiata, *DC. Fl. Fr.* iv. 500, in syn. = horrida.
radiata, Friedr. ex Presl, Bot. Bemerk. 136 = acanthoclada.
radiata, *Scop. Fl. Carn.* ed. II. ii. 51.—Europ. austr.
Raetam, *Forsk. Fl. Aegypt. Arab.* 214.—Reg. Mediterr. or.
ramosissima, Boiss. Voy. Bot. Espagne, 142 = cinerea.
ramosissima, *Poir. Encyc. Suppl.* ii. 715.—Alger.
repens, Lam. Fl. Fr. ii. 618 = pilosa.
Retama, Nichols. Dict. Gard. ii. 59 = G. Raetam.
retamoides, *Spach, ex Coss. Notes Crit.* 154.—Hispan.
rhodopkon, Webb, ex Dellile, Cat. Sem. Hort. Monsp. (1836) 95 = sec.
rhodopnos, Webb & Berth. Phyt. Canar. ii. 37 = Cytisus canariensis.
rhodorhizoides, *Webb & Berth. l. c.* 54.—Ins. Canar.
rigens, *Presl, Delic. Prag.* 84.—Sicilia.
rigida, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 178 = Anarthrophyllum rigidum.
rostrata, Poir. Encyc. Suppl. ii. 719 = triacanthos.
rupestris, Schur, Enum. Pl. Transs. 145 = tinctoria.
sagittalis, *Linn. Sp. Pl.* 710.—Europ.
Saharae, *Coss. in Bull. Soc. Bot. Fr.* ii. (1855) 347.—Alger.
Sakellaridis, *Boiss. & Orph. in Boiss. Diagn.* Ser. II. vi. 42.—Graecia.
saldituna, *Pomel, Nouv. Mat. Fl. Alt.* 320.—Alger.
Salzmanni, DC. Mém. Lég. vi. 311 = sapalathoides.
sarcina, *Pomel, Nouv. Mat. Fl. Alt.* 174.—Alger.
scandens, Lour. Fl. Cochinch. ii. 428 = Butea Loureiri.
scariosa, Viv. Ann. Bot. i. 175 = triangularis.
scoloponderis, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 148 = tridentata.

GENISTA :—
scolopendrina, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 469 = praec.
scoparia, Lam. Fl. Fr. ii. 619 = Cytisus scoparius.
scoparia, Vill. Hist. Pl. Dauph. iii. 420 = cinerea.
scorpia, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 208 = G. Scorpius.
scorpioides, Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 276 = tinctorthos.
scorpioria, St. Lag. in Ann. Soc. Bot. Lyon, vi. (1880) 126 = sec.
Scorpius, *DC. Fl. Fr.* iv. 408.—Reg. Mediterr.
Scorpius, Georgi, Reise, ii. 778 (Quid ?).—Rossia.
Scorpius, Sibth. & Sm. ex Boiss. Fl. Orient. ii. 41 = melia.
sepiaria, Lam. Encyc. ii. 619 = Lebeckia sepiaria.
sericea, Alach. Fl. Jadr. 169, partim = Villarsia.
sericea, Poir. Encyc. Suppl. ii. 721 = Lebeckia sericea.
serices, *Wulf. in Jacq. Coll.* ii. 167.—Austria.
sessilifolia, *DC. Mém. Lég.* vii. 208.—As. Min.
sibirica, Falk, Beitr. ii. 228 = tinctoria.
sibirica, Hohen. ex Boiss. Fl. Orient. ii. 44 = patula.
sibirica, *Linn. Mant.* ii. 571.—Sibiria.
Sigerisma, Fuss, Fl. Transc. 149 = tinctoria.
sophoroides, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = Hypocalyptus obcordatus.
Spachiana, *Webb, in Hook. Bot. Mag.* t. 4195.—Ins. Canar.
spartioides, *Spach, in Ann. Sc. Nat. Sér.* III. ii. (1844) 248.—Afr. austr.
spathulata, Spach, l. c. iii. (1845) 126 = lydia.
sphacelata, Decne. l. c. Sér. II. iv. (1835) 360 = acanthocalda.
sphaerocarpa, *Lam. Encyc.* ii. 616 ; *Spach, in Ann. Sc. Nat.* Sér. III. iii. (1845) 156.—Reg. Mediterr.
spicata, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 559 = Arthrosolen spicatus.
spiniflora, Lam. Fl. Fr. ii. 614 = G. Scorpius.
spinosa, Gilib. Fl. Lithuan. ii. 80 = germanica.
spinosa, Pollini, Fl. Veron. ii. 462 = Calycotome spinosa.
spinulosa, *Pomel. Nouv. Mat. Fl. Atl.* 178.—Alger.
splendens, *Webb & Berth. Phyt. Canar.* ii. 42.—Ins. Canar.
stenopetala, *Webb & Berth. l. c.* 39.—Ins. Canar.
stenophylla, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 166 = tinctoria.
stenoptera, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 148 = tridentata.
striata, *Hill, Veg. Syst.* xiii. 68.—Lusitan.
stylosa, *G. Don, Gen. Syst.* ii. 151.—Lusitan.
subcapitata, Pané. ex Nym. Consp. 158 = Sakellariadia.
subsecunda, Schur, Enum. Pl. Transs. 144 = Cytisus procumbens.
supranubia, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = Cytisus fragrans.
sylvestris, *Scop. Fl. Carn. ed.* II. ii. 53.—Austria.
syriaca, Boiss. & Blanche, Diagn. Ser. II. ii. 8 = Cytisus syriacus.
Tchihatchewi, Boiss. in Tchihat. Asie Min. Bot. I. 6 = sessilifolia.
tenella, Willk. in Bot. Zeit. v. (1847) 426 = micrantha.
Tenorii, Steud. Nom. ed. II. i. 671 = humilis.
tenuifolia, *Lamk. Nat.* 169.—Europ.
teretifolia, *Willd. in Flora*, xxxiv. (1851) 617.—Hispan.
ternifolia, Lag. Gen. et Sp. Nov. 22 = Anthyllis Genistae.
tetragona, Bess. Enum. Pl. Volh. ii. 73 = depressa.
tetragona, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 3 = scarica.
tetragona, Vill. ex Verl. in Bull. Soc. Stat. Iebre, xiv. (1872) 77 = sagittalis.
thebaica, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = Crotalaria thebaica.
tinctoria, *Linn. Sp. Pl.* 710.—Europ. ; Reg. Caucas.
tomentella, Boiss. & Noé, Diagn. Ser. II. ii. 9 = albida.
tomentosa, Poir. Encyc. Suppl. ii. 719 = Cytisus palens.
Tournefortii, *Spach, in Ann. Sc. Nat.* Sér. III. ii. (1844) 269.—Lusitan.
transsilvanica, Schur, Enum. Pl. Transs. 144 = lydia.
GENISTA :—
triacanthos, *Brot. Fl. Lusit.* ii. 80 ; *Phyt.* 130. t. 54. —Reg. Mediterr. occ.
triangularis, Baumg. Enum. Stirp. Transs. ii. 320 = lydia.
triangularis, Hort. ex Tausch, in Flora, xxi. (1838) 747 = scariosa.
triangularis, *Willd. Sp. Pl.* iii. 989.—Banatus.
tricuspidata, *Desf. Fl. Atlant.* ii. 188. t. 188.—Afr. bor. occ.
tricuspidata, Salzm. ex Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 274 = gibsilatarica.
tridens, *Cav. in Anal. Cienc. Nat.* iv. (1801) 59.—Marocc.
tridentata, *Linn. Sp. Pl.* 710.—Reg. Mediterr. occ.
trifoliata, *Jamka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 248.—Europ.
triquetra, *Lam. Encyc.* ii. 622.—Hispan.?
triquetra, *L'Hérit. Stirp. Nov.* 183. t. 85.—Corsic.
trugurina, Schur, ex Nym. Consp. 153 = lydia.
triquetra, Waldst. & Kit. Pl. Rar. Hung. ii. 165. t. 153 = triangularis.
trispermo, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = aethiensis.
ulicina, *Spach, l. c.* ii. (1844) 268.—Afr. bor.
umbellata, Cloe, in C. Gay, Fl. Chil. ii. 56 = Anarthrophyllum umbellatum.
umbellata, *Poir. Encyc. Suppl.* ii. 716.—Afr. bor. ; Corsic.
umbellata, Salzm. ex DC. Prod. ii. 147 = aspalathoides.
umbellata, Webb, It. Hisp. 58 = equisetiformis.
undulata, *Link, Handb.* ii. 150.—Lusit. ; Afr. bor.
valentina, *Cav. Ind. Nam.* ed. II. i. 671.—Hispan.
Vepres, *Pomel, Nouv. Mat. Fl. Atl.* 319.—Alger.
versicolor, Boiss. ex Steud. Nom. ed. II. i. 671 = asparthoides.
versicolor, Wall. Cat. n. 5922 = Caragana pygmaea.
versiflora, *Tausch, in Flora*, xxi. (1838) 746.—Hab.?
Villarsiana, Jord. Obs. Pl. Crit. vi. (1847) 86 = sec.
Villarsii, *Clementi, in Atti Teras Rium. Sc. Ital.* 517. —Dalmat.
villosa, Lam. Fl. Fr. ii. 615 = germanica, hispanica.
villosa, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 156 = Calycotome villosa.
virgata, Blasoletto, ex Reichb. Fl. Germ. Excurs. 519, in obs = sibirica ?
virgata, Hoppe, ex Reichb. l. c. 520, in obs. = ovata.
virgata, *Link, Enum Hort. Berol.* ii. 228 ; *DC. Prod.* ii. 148.—Ins. Madeir.
virgata, Lam. Encyc. ii. 616 = Eremosparton aphyllum.
virgata, Willd. Berl. Baumz. 159 = clatior.
viscosa, Willd. Sp. Pl. iii. 987 = Adenocarpus anagyrus.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 595 = scoparia.
Webbii, Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 252 = monosperma.
Wehwitschii, *Speach, l. c.* 262.—Lusitan.
Winkleri, *Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 286.—Hispan.

GENISTELLA, Tourn. ex Rupp. Fl. Jen. Hall. 266 (1745) = **Genista**, Linn. (Legumin.).
flagelliformis, Pourr. ex Colm. in Willk. & Lange, Prod. Fl. Hisp. iii. 485 = Genista florida.
florida, Pourr. ex Colm. l. c. 486 = Genista florida.
polymorpha, Pourr. ex Colm. l. c. 488 = Genista ramosissima.
racemosa, Moench, Meth. 138 = Genista sagittalis.
ramosissima, Pourr. ex Colm. in Willk. & Lange, Prod. Fl. Hisp. iii. 488 = Genista ramosissima.
verticillata, Pourr. ex Colm. l. c. 484 = Genista cinerea.

GENISTOIDES, Moench, Meth. 132 (1794) = **Genista**, Linn. (Legumin.).
elata, Moench, l. c. = Genista sibirica.
linifolia, Moench, Meth. Suppl. 44 = Cytisus linifolius.
tinctoria, Moench, Meth. 138 = Genista tinctoria.
tuberculata, Moench, l. c. = Genista pilosa.

GENLISEA, Benth. & Hook. f. Gen. iii. 701 (1883) = GENLISIA, Reichb. = **Aristea**, Ait. (Irid.).

GENLISEA, A. St. Hil. Voy. Distr Diam. ii. 428 (1833). *LENTIBULARIEAE*, Benth. & Hook. f. ii 988.
DORRIENIA, Endl. Gen. Suppl. ii. 65, in syn (1842).
africana, *Oliver, in Journ. Linn. Soc.* ix. (1867) 145. —Afr. trop.
aurea, *A. St. Hil. Voy. Distr. Diam.* ii. 429.—Bras.
bilobata, *Benj. in Mart. Fl. Bras.* x. 254.—Bras.
filiformis, *A. St. Hil. Voy. Distr. Diam.* ii. 430.—Bras.
luteo-viridis, *Wright, in Sauv. Fl. Cub.* 90.—Cuba.
minor, *A. St. Hil. Voy Distr. Diam.* ii. 430.—Bras.
ornata, *Mart. ex Benj. in Mart. Fl. Bras.* x. 252 t. 21. f. 2.—Bras.
pusilla, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 11. —Bras.
pygmaea, *A. St. Hil. Voy. Distr. Diam.* ii. 431.—Bras.
reflexa, *Benj. in Mart. Fl. Bras.* x. 254.—Bras.
repens, *Benj. l. c.* 253.—Bras.
violacea, *St. Hil Voy. Distr. Diam.* ii. 429.—Bras.

GENLISIA, Reichb. Consp. 60 (1828) = **Aristea**, Ait. (Irid.).
corymbosa, Reichb. Fl. Exot. t. 24 = A. corymbosa (in Addend.).

GENNARIA, Parl. Fl. Ital. iii. 404 (1858) = **Habenaria**, Willd (Orchid.).
diphylla, Parl. l. c. 405 = H. cordata.

GENOPLESIUM, R. Br. Prod. 319 (1810) = **Prasophyllum**, R. Br. (Orchid.).
Baueri, R. Br. l. c. = P. rufum.

GENORIA, Pers. Syn. ii. 9 (1807) = **Ginora**, Linn. f. (Lythrar.).

GENOSIKIS, Labill Nov. Holl. Pl. i. 13. t. 9 (1804) = **Patersonia**, R. Br (Irid.).
fragilis, Labill. l. c. = P. glauca.
glabrata, F. Muell. Fragm. vii. 35 = P. glabrata.
juncea, F. Muell. l. c. 33 = P. juncea.
lanata, F. Muell. l. c. 35 = P. lanata.
longifolia, F. Muell. l. c. = P. longifolia.
Maxwelli, F. Muell. l. c. 34 = P. Maxwelli.
occidentalis, F. Muell. l. c. 31 = P. occidentalis.
pygmaea, F. Muell. l. c. 33 = P. pygmaea.
rudis, F. Muell. l. c. 35 = P. rudis.
sericea, F. Muell. l. c. = P. sericea.
umbrosa, F. Muell. l. c. 32 = P. umbrosa.
xanthina, F. Muell. l. c. 33 = P. xanthina.

GENTIANA, Tourn. ex Linn. Syst. ed. I (1735). *GENTIANEAE*, Benth & Hook. f. ii. 815.
ALOITIS, Rafin. Fl. Tellur. iii. 21 (1836).
AMARELLA, Rafin. l. c. 20 (1836).
ANTHOPOGON, Neck. Elem. ii. 12 (1790).
ASTERIAS, Borckh. in Roem. Archiv, i. 1. 25 (1796).
BILAMISTA, Rafin. New Fl. Am. iv. 92 (1836).
CALATHIANA, Delarb. Fl. Auv. ed. II. 28 (1800).
CHARLOTHILUS, Neck. Elem. ii. 14 (1790).
CHIOPHILA, Rafin. Fl. Tellur. iii. 28 (1836).
CIMINALIS, Adans. Fam. ii. 204 (1763).
COILANTHA, Borckh. in Roem. Archiv, i. 1. 25 (1796).
CROSSOPETALON, Adans. Fam. ii. 224 (1763).
CROSSOPETALUM, Koth, Enum. Pl. Phaen. Germ. i. i. 515 (1827).
CRUCIATA, Gilib. Fl. Lituan. i. 86 (1781).
CUTTERA, Rafin. in Med. Repos. N. York, v. (1808) 352.
DAMISTEPMA, Rafin. Fl. Tellur. iii. 19 (1836).
DASYSTEPHANA, Adans. Fam. ii. 202 (1763) ; Borckh. in Roem. Archiv, i. 1. 25 (1796).
DENCKEA, Rafin. in Med. Repos. N. York, v. (1808) 352.
DICARDIOTIS, Rafin. Fl. Tellur. iii. 21 (1836).
DIPLOMA, Rafin. l. c. 19 (1836).
ERICOLA, Renealm. ex Borckh. in Roem. Archiv, i. 1. 26 (1796).
EUDOXIA, D. Don, ex G. Don, Gen. Syst. iv. 302 (1836).
EYSTYHALIA, Renealm. ex Borckh. in Roem. Archiv, i. 1. 28 (1796).
GAERTNERIA, Neck. Elem. ii. 15 (1790).
GENTIANELLA, Moench, Meth. 462 (1794).

GENTIANA:—

Glyphospermum, G. Don, Gen. Syst. iv. 195 (1836).
Gonipia, Rafin. Fl. Tellur. iii. 23 (1836).
Heteroclita, Rafin. l. c. 27 (1836).
Hippion, F. W. Schmidt, in Roem. Archiv, t. 1. 9. t. 2 ad 5 (1706).
Lymnanna, Casseb. & Theob. Fl. Wett. ed. II. 32 (1847).
Leimaniya, Rafin. New Fl. Am. iv. 98 (1836).
Lexipyrium, Dulac, Fl. Hautes-Pyr. 449 (1867).
Monoteria, Rafin. Fl. Tellur. iii. 49 (1836).
Opsantha, Delarb. Fl. Auv. ed. II. 80 (1800).
Pneumonanthe, Gleditsch. in Mém. Acad. Berl. v. (1751) 180.
Pogosolephis, Rafin. Fl. Tellur. iii. 26 (1836).
Psalina, Rafin. l. c. 25 (1836).
Rassia, Neck. Elem. ii. 12 (1790).
Ricoila, Renealm. ex Rafin. Fl. Tellur. iii. 28 (1836).
Sebbokia, Neck. Elem. ii. 13 (1790).
Selatium, G. Don, Gen. Syst. iv. 196 (1836).
Spiragyne, Neck. Elem. ii. 14 (1790).
Tetrorhiza, Renealm. ex Rafin. Fl. Tellur. iii. 25 (1836).
Thylacitis, Renealm. ex Adans. Fam. ii. 504, 539 (1763).
Thymophora, Neck. Elem. ii. 13 (1790).
Tretorhiza, Adans. Fam. ii. 503 (1763).
Ulostoma, G. Don, Gen. Syst. iv. 198 (1836).
Varasia, Phil. Fl. Atac. 35. t. 5 (1860).
Xolemia, Rafin. Fl. Tellur. iii. 23 (1836).

abscondita, Zenker, Pl. Ind. 11. t. 13 = quadrifaria.
acaulis, *Linn. Sp. Pl.* 229.—Europ. t. As. bor.
acaulis, Patrin, in Pall. N. Nord. Beitr. vi. 183 = verna.
achalensis. *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 378.—Reg. Argent.
acuta, Michx. Fl. Bor. Am. i. 177 = G. Amarella.
acutiflora, Schult. Syst. vi. 156, in syn. = verna.
adscendens, Pall. Fl. Ross. i. 11. 106. t. 94 = decumbens.
adaurgens, *Cerv. ex Griseb. Gen. et Sp. Gent.* 286.—Mexic.
aestiva, Schult. Syst. vi. 156 = verna.
affinis, *Griseb. Gen. et Sp. Gent.* 289 —Am. bor. occ.
aggregata, Bunge, ex Griseb. in DC. Prod. ix. 100 = G. Amarella.
alba, *Muhl. Cat.* ed. II. 39.—Am. bor.
albens, Thunb. Prod. Pl. Cap. 48 = Sebaea aurea.
albescens, *Favre, in Bull. Soc. Murith.* v. (1875) 145, 251.—Europ.
albiflora, Lam. Tabl. Encyc. i. 483 = frigida.
albiflora, *Schur, in Verh. Siebenb. Ver. Naturw.* iii. (1852) 88.—Transsylv.
aleutica, Cham. & Schlecht. in Linnaea, i. (1826) 175 = aurea.
algida, Pall. Fl. Ross. i. 11. 107. t. 95 = frigida.
algida, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 249 = gelida.
alopecuroides, Lam. Encyc. ii. 643 = Erythraea alopecuroides.
alpina, Adam, in Hoffmgg. Hort. Mosq. (1808) n. 1487 = gelida.
alpina, Vill. Prosp. 22; Hist. Pl. Dauph. ii. 526 = acaulis.
alsinoides, *Franch. in Bull. Soc. Bot. Fr.* xxxi. (1884) 374.—China.
altaica, *Laxm. in Nov. Comm. Acad. Petrop.* xvii. (1774) 598.—Sibir.
Amarella, All. Fl. Pedem. i. 97 = campestris.
Amarella, Froel. Gent. Diss. 86 = germanica.
Amarella, Jacq. Enum. Stirp. Vindob. 42 = austriaca.
Amarella, *Linn. Sp. Pl.* 330.—Amphig. bor.
Amarella, Pall. Fl. Ross. i. 11. 104 = caucasica.
Amarella, Sm. in Sibth. & Sm. Fl. Graec. Prod. i. 174 = obtusifolia.
amarelloides, Michx. Fl. Bor. Am. i. 175 = quinquefolia.
ambigens, Gaud. in Ebel, Man. Voy.; et in Fl. Helv. ii. 290 = obtusifolia.
amoena, *G. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 115.—Reg. Himal.
amoena, *Wedd. Chlor. And.* ii. 70.—Ecuador.
Andersonii, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 486 = capitata.
Andrewsii, *Griseb. Gen. et Sp. Gent.* 287.—Am. bor.
angulosa, Bieb. Fl. Taur. Cauc. i. 197 = verna.
angustifolia, *Michx. Fl. Bor. Am.* i. 177.—Am. bor.
angustifolia, Sturm, Deutsch. Fl. Heft 24 = Froelichii.
angustifolia, Vill. Hist. Pl. Dauph. ii. 526 = acaulis.
aperta, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 500.—China.

GENTIANA:—

aphylla, Jacq. Enum. Pl. Carib. 17 = Voyria aphylla.
aprica, Decne. in Jacquem. Voy. Bot. 112. t. 118 = decomfida.
aquatica, Froel. Gent. Diss. 77 = prostrata, squarrosa.
aquatica, *Linn. Sp. Pl.* 229.—Reg. bor. et arct.
aquatica, Laur. Fl. Cochinch. 129 = indica.
aquatica, Pall. Fl. Ross. i. 11. 109. t. 97 = humilis.
aquatica, Reichb. Fl. Germ. Excurs. 425 = nivalis.
aquatica, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 255 = squarrosa.
aquatica, Thunb. Fl. Jap. 115 = Thunbergii.
arenaria, Adam, ex Griseb. Gen. et Sp. Gent. 285 = gelida.
arenophila, *Griseb. l. c.* 251.—Am. arct.
arcuata, Griseb. l. c. 282 = Jamesoni.
argentea, *Royle, ex D. Don, in Transc. Linn. Soc.* xvii. (1837) 512.—Reg. Himal. ; China.
aristata, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 497.—China.
armerioides, *Griseb. in Lechl. Berb. Am. Austr.* 58.—Peruv.
arvernensis, *Hort. Backh. ex Gard. Chron.* (1883) ii. 40.—Hab. ?
ascendens, Poir. ex Steud. Nom. ed. II. i. 672 = decumbens.
asclepiadea, *Linn. Sp. Pl.* 227.—Europ. austr. ; Reg. Caucas.
aspera, Hegetschw. Fl. Schw. 210 = obtusifolia.
astata, *Bunge, ex Griseb. in DC. Prod.* ix. 98.—Sibir.
attenuata, *Griseb. l. c.* 98.—Peruv.
aurea, *Linn. Syst.* ed. X. 951.—Amphig. bor.
aurea, Thunb. Fl. Cap. ii. 171 = Sebaea aurea.
auriculata, *Pall. Fl. Ross.* i. 11. 102. t. 92.—Sibir. or. ; Ins. Aucl.
austriaca, *A. & J. Kern. Sched. Fl. Exsicc.* ii. 128.—Austria.
autumnalis, Prey. & Zirk. ex Bab. in Journ. Linn. Soc. ix. (1871) 318 = G. Amarella ?
axillaris, *Rafin. Med. Fl.* i. 213.—Am. bor.
axillaris, Reichb. Pl. Crit. ii. 18. t. 130 = G. Amarella.
aznrea, Bunge, in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 230 = aurea.
Baconi, Edgew. in Hook. Lond. Journ. Bot. iv. (1845) t. 10 = capitata.
barbata, Froel. Gen. Diss. 114 = detonsa.
barbellata, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 216 t. 2.—Am. bor. occ.
barbata, *Linn. Sp. Pl.* 229.—Europ.
bavarica, Pall. Fl. Ross. ii. 110 = verna.
bavarica, Zoëga, Fl. Island. 235 = detonsa.
bellidifolia, Hook. f. in Hook. Ic. Pl. t. 635 = montana.
bicolor, *Wedd. Chlor. And.* ii. 310.—Peruv.
Bieberatenii, Bunge, in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 247. t. 10. f. 1 = obtusifolia.
Bigelovii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 56. —N. Mexic.
biloba ×, DC. Fl. Fr. iii. 653.—Gallia.
Boissieri, *Schott & Kotschy, ex Boiss. Fl. Orient.* iv. 75.—As. min.
borealis, *Bunge, in Nouv. Mém. Soc. Nat. Mosc.* i. (1829) 251. t. 10. f. 2.—Reg. Himal. ; Am. bor. occ. bornensis, *Hook. f. Ic. Pl.* t. 1440 B.—Borneo.
Boryi, *Boiss. Elench.* 65.—Hispan.
brachystelata, Bunge, in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 225 = detonsa.
brachyphylla, *Vill. Prosp.* 23; *Hist. Pl. Dauph.* ii. 536.—Helvet.
brevidens, *Franch. & Sav. Enum. Pl. Jap.* i. 325.—Japon.
bromifolia, *Griseb. in Goett. Abh.* xix. (1874) 208.—Reg. Argent.
buccoviriensis, Herbich, Select. Pl. Rar. Galic. Bucov. 10 = utriculosa.
Bungei, *Mig. Ann. Mus. Bot. Lugd. Bat.* iii. 124.—Japon.
Burseri, *Lapeyr. Hist. Abr. Pl. Pyr.* 193.—Mont. Pyren.
Burseri-punctata, Griseb. Gen. et Sp. Gent. 301 = Burseri.
Cachalalaoen, Molina, Sagg. Chile, 147 = Erythraea chilensis.
cachemirica, *Decne. in Jacquem. Voy. Bot.* 111. t. 117. —Reg. Himal.
carnilea, Auct. vide = *caerulea*.
carneiserone, Gill. ex Wedd. Chlor. And. ii. 66, in obs. = diffusa ?
caespitosa, R. Grah. in Edinb. N. Phil. Journ. (July-Sept. 1830) 367 = glauca.

GENTIANA:—

caespitosa, *Mert. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 870.—Mexic.
calycina, *Boiss. & Heausb. ex Boiss. Fl. Orient.* iv. 75.—As. Min.
calycina, Lam. Encyc. ii. 638 = Sabbatia calycosa.
calycosa, A. Gray, in Pacif. Rail. Rep. vi. 86 = Newberryi.
calycosa, *Griseb. Gen. et Sp. Gent.* 292.—Am. bor. occ.
calycularis, *Lex. in La Llave & Lex. Nov. Veg. Descr.* fasc i. 18 (calliculatis).—Mexic.
campanata, [Clairv.] Man. Herb. 74 = acaulis.
campanulata, Jacq. Fl. Austr. v. 43 = punctata.
campanulata, Reyn. ex Griseb. Gen. et Sp. Gent. 212 = *hybrida*.
campanuloides, Willd. ex Schult. Syst. vi. 184 = rupicola.
Campbelli, *Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél.* 93.—Ins. Aucki.
campestris, All. Fl. Pedem. i. 97 = germanica.
campestris, Generx. Elench. 284 = G. Amarella.
campestris, *Linn. Sp. Pl.* 231.—Europ.
canaliculata, Royle, ex G. Don, Gen. Syst. iv. 182 = Jaeschkea lutlepalis.
Canchalagua, Palau, Pant. Pract. Bot. viii. 22; ex Mem. Acad. Med. Madr. i. (1797) 408 = *G. Cachalalaoen* ?
capitata, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 136. —Reg. Himal.
carinata, *Griseb. Gen. et Sp. Gent.* 304.—Reg. Himal.
carinthiaca, Froel. Gent. Diss. 108 = Pleurogyne carinthiaca.
carnica, Weiw. ex Reichb. Fl. Germ. Excurs. 865 = Froelichii.
carpathica, Kit. in Schult. Oestr. Fl. ed. II. i. 443 = nivalis.
Catesbaei, Andr. Bot. Rep. t. 418 = Andrewsii.
Catesbaei, Walt. Fl. Carol. 109 = G. Saponaria.
caucasica, *Bieb. Fl. Taur. Cauc.* i. 198.—Reg. Caucas.
cauliscens, Lam. Encyc. ii. 638 = acaulis.
Centaurium, Kalm, Fl. Fennic. n. 153; ex Ledeb. Fl. Ross. iii. 50 = Erythraea linariifolia.
Centaurium, Linn. Sp. Pl. 229 = Erythraea ramosissima.
Cephalotes, Edgew. in Hook. Lond. Journ. Bot. iv. (1845) 687 = capitata.
cerastioides, *H. B. & K. Nov. Gen. et Sp.* iii. 169.—Peruv.
cerina, *Hook. f. Fl. Antarct.* 54. t. 36.—Ins. Auckland.
cernua, *H. B. & K. Nov. Gen. et Sp.* iii. 170.—Ecuador.
cespitosa, Willd. ex Schult. Syst. vi. 155 = sedifolia.
Charpentieri ×, Thom. ex Hegetschw. Fl. Schw. 204, 988.—Europ.
chelonioides, Willd. ex Schult. Syst. vi. 155 = dianthoides.
chimborazensis, Willd. ex Schult. l. c. = sedifolia.
Chirayita, Roxb. ex Flem. in As. Res. xi. (1810) 167 = Swertia Chirata.
chlodes, Brot. Fl. Lusit. i. 276 = Erythraea confarta.
chloraefolia ×, Nees, in Nov. Act. Nat. Cur. ix. (1818) 171.—Europ.
ciliata, Gunner, Fl. Norveg. ii. 88; Pall. Fl. Ross. i. 11. 101 = detonsa.
ciliata, *Linn. Sp. Pl.* 231.—Europ. ; As. Min.
ciliata, Moq. & Sesse, ex G. Don, Gen. Syst. iv. 179 = macrantha.
cistiflora, Willd. ex Schult. Syst. vi. 184 = cerastioides.
clausa, Rafin. Med. Fl. i. 210 = Andrewsii.
clavata, Steud. Nom. ed II. i. 672 = ochroleuca.
Clusii, Perr. & Song. in Bull. Soc. Hist. Nat. Savoie, 1858 (1854) 185 = acaulis.
coarctata, Willd. ex Schult. Syst. vi. 184 = foliosa.
cocuisa, *Ruiz & Pav. ex G. Don, Gen. Syst. iv.* 196. —Peruv.
coerulea, Moq. & Sesse, ex Griseb. in DC. Prod. ix. 112 = Sessei.
coerulea, Ruiz & Pav. ex Griseb. Gen. et Sp. Gent. 285 = linifiora.
collina, Adam, ex Griseb. l. c. 245 = cancasica.
collina, With. Bot. Arr. Brit. Fl. ed. III. ii. 223 = Campanula glomerata.
Collisiana, *Rafin. Med. Fl.* i. 213.—Am. bor.
Columnae, Tenore, Fl. Nap. iii. 248 = G. Amarella.
compacta, Hegetschw. Fl. Schw. 210 = germanica.
concinna, *Hook. f. Fl. Antarct.* 53. t. 35.—Ins. Auckl.
conferta, *Flack. ex Smeet, Hort. Brit.* ed. II. 388.—Sibir. altaic.

GENTIANA:—
congesta, Willd. ex Schult. Syst. vi. 184 = corymbosa.
connata, Willd. ex Schult. i. c. = Eustoma silenifolium.
consobrina, Schott & Kotschy, ex Boiss. Fl. Orient. iv. 71 = aurea.
contorta, Falc. ex C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 488 = depressa.
contorta, *Royle, Illustr. Bot. Himal.* 278. t. 69.—Reg. Himal.
cordifolia, C. Koch, in Linnæa, xxiii. (1850) 585 = septemfida.
coronata, *Royle, Illustr. Bot. Himal.* 278. t. 68. f. 1.—Reg. Himal.
corymbosa, *H. B. & K. Nov. Gen. et Sp.* iii. 171.—N. Granat.
cosmantha, *Griseb. in Goett. Abh.* xix. (1874) 209.—Reg. Argent.
crassa, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) 11. 235.—Burma.
crassifolia, *Griseb. Gen. et Sp. Gent.* 227.—Am. austr.
crinita, *Prosl. Gent. Diss.* 112.—Am. bor.
crispata, Vis. in Flora, xlii. (1830) 50; Fl. Dalmat. ii. 258 = obtusifolia.
crtica, Ehrh. ex Griseb. Gen. et Sp. Gent. 244 = germanica.
crossolacma, *Wedd. Chlor, And.* ii. 55.—Peruv.
Cruciata, *Linn. Sp. Pl.* 281.—Europ.; As. bor.
cuspidata, *Griseb. Gen. et Sp. Gent.* 224.—Peruv.
cuspidata, Wall. ex C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 485 = quadrifaria.
dahurica, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 68 = decumbens.
David, *Franch. Pl. Davidian.* i. (1884) 211.—China.
decemfida, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 127. —Reg. Argent.
decumbens, *Linn. f. Suppl.* 174.—Reg. Himal.; Sibir.
Delavayi, *Franch. in Bull. Soc. Bot. Fr.* xxxi. (1884) 377.—China.
delicata, *Hance, in Journ. Bot.* xxi. (1883) 324.—China.
depressa, Bourg. ex Nym. Consp. 499 = G. Pneumonantha.
depressa, *D. Don, Prod. Fl. Nep.* 125.—Reg. Himal.
depressa, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 49 = phlogifolia.
detonsa, *Rottb. in Kiob. Selsk. Skr.* x. (1770) 485. t. 1. f. 3.—Amphig. bor.
dianthoides, *H. B. & K. Nov. Gen. et Sp.* iii. 170.—Peruv.
dichotoma, Pall. Fl. Ross. i. 11. 110 = tenella.
diemenxis, Griseb. Gen. et Sp. Gent. 224 = montana.
diffusa, *H. B. & K. Nov. Gen. et Sp.* iii. 172.—Ecuador.
diffusa, Vahl, Symb. Bot. iii. 47 = Canscora diffusa.
dilatata, Griseb. Gen. et Sp. Gent. 223 = primulæfolia.
diluta, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 388 = Swertia diluta.
discolor, Hoffmgg. ex Reichb. Fl. Germ. Excurs. 865 = verna.
dshimilensis, Walp. Ann. iii. 85, vide *dshimilensis*.
disstiflolia, *Griseb. Gen. et Sp. Gent.* 229.—Peruv.
djimilensis, Boiss. Fl. Orient. iv. 73, vide *dshimilensis*.
dodecapetala, Griseb. Gen. et Sp. Gent. 125 = Sabbatia chloroides.
Dombeyana, *Wedd. Chlor. And.* ii. 62.—Peruv.
Donii, *Griseb. in DC. Prod.* ix. 93.—Peruv.
Douglasiana, *Bong. in Mém. Acad. Pétersb.* ii. (1833) 158.—Am. bor. occ.
dschungarica, *Regel, in Act. Hort. Petrop.* vi. (1879) 384.—Turkestan.
dshimilensis, C. Koch, in Linnæa, xxiii. (1850) 588 = pyrenaica.
dubia, *Thunb. Prod. Pl. Cap.* 48.—Afr. austr.
Dumeetilii ×, Stein, ex Dalla Torre, Atl. Alpenfl. 171.—Europ.
Elliottea, *Raflu. Med. Fl.* i. 212 (Quid ?).—Am. bor.
Elliottii, *Chapm. Fl. S. U. St.* 356.—Am. bor.
elongata, Hænk. in Jacq. Coll. ii. 88 = bavarica.
Elwesii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 115. —Reg. Himal.
ericoides, *Griseb. Gen. et Sp. Gent.* 221.—Peruv.
exneeides, Linn. Sp. Pl. ed. ii. 332 = Belmontia cordata.
exaltata, Linn. f. c. 231 = Eustoma exaltatum.
exeisa, Presl, in Flora, xi. (1828) 366 = scaalis.
fakena, *Turcz. ex Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 404.—Sibir.

GENTIANA:—
Falconeri, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 483 = Microcorallana.
fastigiata, *Benth. Pl. Hartw.* 144.—Peruv.
fastigiata, *Franch. in Bull. Soc. Bot. Fr.* xxxi. (1884) 372.—China.
Festiowi, *Regel & Winkler, in Gartenfl.* (1882) 3.—Turkestan.
fibrisecnta, Schur, Enum. Pl. Transs. 457 = phlogifolia.
filamentosa, *Griseb. in DC. Prod.* ix. 94.—Peruv.
filiformis, Linn. Sp. Pl. 281 = Microcala filiformis.
filiformis, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 559 = Cicendia pusilla.
fimbriaspilica, C. Koch, in Linnæa, xxiii. (1850) 584 = septemfida.
fimbriata, Andr. Bot. Rep. t. 509 = crinita.
fimbriata, Vahl, Symb. Bot. iii. 46 = G. Saponaria.
fimbriata, Vitm. Summa Pl. ii. 117 = tenella.
fimbriata, Willd. ex Schult. Syst. vi. 175 = ciliata.
firma, Kern. in Oestr. Bot. Zeitschr. xxii. (1872) 56 = acaulis.
flava, Mayer, ex Schult. Oestr. Fl. ed. II. i. 445 = G. Amarella.
flava, Mœnch, ex Loisel. in Mém. Soc. Linn. Par. vi. (1827) 408 = germanica.
flawescens, Schur, Enum. Pl. Transs. 461 = G. Amarella.
flawescens, Vis. ex Griseb. Gen. et Sp. Gent. 244 = germanica.
flavida, A. Gray, in Am. Journ. Sc. Ser. II. i. (1846) 80 = alba.
floribunda, D. Don, Prod. Fl. Nep. 127 = Swertia Chirata.
floribunda, Willd. ex Schult. Syst. vi. 185 = linifiora.
florida, *Griseb. in Goett. Abh.* xxiv. (1879) 286.—Reg. Argent.
foliosa, *H. B. & K. Nov. Gen. et Sp.* iii. 170.—Peruv.
Fortuni, Hook. Bot. Mag. t. 4776 = scabra.
Forwoodii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 86.—Am. bor. occ.
Fremontii, Torr. in Frém. Rep. 94 = humilis.
frigida, *Haenke, in Jacq. Coll.* ii. 18.—Europ. or.; As. bor.; Am. bor. occ.
Froelichii, *Jan, ex Reichb. Fl. Germ. Excurs.* 865.—Carniol.
fruticulosa, *Domb. ex Wedd. Chlor. And.* ii. 71.—Peruv.
Galandeii, *Hirron. in Bot. Acad. Nac. Córdova*, iv. (1881) 373.—Reg. Argent.
Gaudini ×, Thom. ex Hegetschw. Fl. Schw. 205 (= seq. ?).—Europ.
Gaudinxana ×, Thom. ex Koch, Syn. Fl. Germ. ed. II. 580.—Europ.
Gayi, Griseb. in DC. Prod. ix. 106 = acaulis.
Gebleri, Ledeb. ex Bunge, in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 211. t. 5 = decumbens.
gelida, *Bieb. Fl. Taur. Cauc.* i. 196.—As. Min.
Gerardi, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 192 = Erythræa Centaurium.
germanica, Neilr. Fl. Wien, 319 = austriaca.
germanica, *Willd. Sp. Pl.* i. 1846.—Europ.; Reg. Caucas.
glacialis, Thom. ex Vill. Hist. Pl. Dauph. ii. 532 = tenella.
glauca, *Pall. Fl. Ross.* i. 11. 104. t. 98. f. 2.—Sibir.; Am. bor. occ.
gracilis, Chatn. & Schlecht. in Linnæa, i. (1826) 176, in nota = propinqua.
gracilis, *H. B. & K. Nov. Gen. et Sp.* iii. 168.—Ecuador.
gracilis, Nees, in Nov. Act. Nat. Cur. ix. (1818) 176 = G. Amarella.
gracillima, Bertol. in Misc. Bot. xv. 19 = Apteria setacea.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 169.—Peruv.
grandiflora, Lam. Fl. Fr. ii. 396 = acaulis.
grandiflora, Lam. in Nov. Comm. Acad. Petrop. xvii. (1774) 526. t. 6. f. 1 = altaica.
Griesbachii, Hook. f. in Hook. Ic. Pl. t. 686 = montana.
guayaquilensis, *Griseb. in DC. Prod.* ix. 93.—Ecuador.
Hartwegi, *Benth. Pl. Hartw.* 47.—Mexic.
Herrediana, *Raimondi, ex Wedd. Chlor. And.* ii. 309. —Peruv.
heterocalix, Linn. Mant. ii. 560 = Canscora sessilifora.
heterophylla, Rafin. Med. Fl. 211 = ochroleuca.
heterosepala, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 215. t. 8.—Am. bor. occ.
hexamera, *Phil. in Anal. Univ. Chil.* (1873) 510.—Chili.
Hitquius, *Griseb. Gen. et Sp. Gent.* 219.—Ecuador.

GENTIANA:—
Hladnikiana, Host, ex Koch, Syn. Fl. Germ. ed. I. 488 = Froelichii.
holosteoides, Schott & Kotschy, ex Boiss. Fl. Orient. iv. 71 = aurea.
Hookeri, *Arnott. in Trans. N. Z. Inst.* xiii. (1881) 340.—N. Zel.
Hookeri, *Griseb. Gen. et Sp. Gent.* 362.—Ecuador.
Hugelii, *Griseb. in DC. Prod.* ix. 108.—Reg. Himal.
humilis, Rochel, Pl. Banat. Rar. 61. f. 47 = nivalis.
humilis, Salisb. Prod. 137 = acaulis.
humilis, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 258. —As. et Am. bor.
hybrida, *Raflu. in Med. Repos. N. York*, v. (1808) 353, nomen.—Am. bor.
hybrida, *Schleich. ex DC. Fl. Fr.* iii. 451.—Europ.
hybrida, Vill. in Roem. Collect. 180 = Barrelieri.
hyssopifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 172.—Ecuador.
hyssopifolia, Linn. ex Spreng. Syst. i. 589 = Enicostema littorale.
imberbis, *Griseb. in Goett. Abh.* xix. (1874) 209.—Reg. Argent.
imbricata, *Prosl. Gent. Diss.* 74.—Europ.
imbricata, Schleich. ex DC. Fl. Fr. vi. 427 = bavarica.
imbricata, Willk. ex Nym. Consp. 499 = pumila.
immaculata, Pers. Syn. i. 284 = punctata.
incarnata, Sims, Bot. Mag. t. 1856 = ochroleuca.
incurva, *Hook. Bot. Misc.* ii. (1831) 228.—Peruv.
indica, *Steud. Nom.* ed. II. i. 674.—China.
Infelix, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 111. —Reg. Himal.
inflata, *Griseb. Gen. et Sp. Gent.* 232.—Ecuador.
intermedia, Sims, Bot. Mag. t. 2308 = ochroleuca.
involucrata, Rottb. in Kiob. Skr. Selsk. x. (1770) 434 = aurea.
islandica, J. F. Gmel. Syst. 463 = Pleurogyne rotata.
islandica, Herb. Joannet, ex Schult. Syst. vi. 171 = tenella.
Jaeschkei, Kurz, in Journ. Bot. v. (1867) 241 (sphalm. *Taeschke*) = Jaeschkea gentianoides.
jakutensis, Bunge, ex Griseb. in DC. Prod. ix. 118 = macrophylla.
Jamesoni, *Hook. Ic. Pl.* t. 61.—Ecuador.
Jamesonii, Griseb. Gen. et Sp. Gent. 219 = Hookeri, *Griseb.*
japonica, *Maxim. in Bull. Acad. Pétersb.* xx. (1875) 433.—Japon.
japonica, Schult. Syst. vi. 174 = Swertia diluta.
Jirasekii, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 195 = verna.
Karelinii, Griseb. in DC. Prod. ix. 106 = aquatica.
Kaufmannxana, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 381.—Turkestan.
kentrokensis, Pers. Syn. i. 286 = quinqueflora.
Kesselringi, *Regel, in Act. Hort. Petrop.* vii. (1880) 548.—Turkestan.
Kochiana, Perr. & Song. in Bull. Soc. Hist. Nat. Savoie, 1858 (1854 ?), 180 = acaulis.
Koenigii, Gmner, Fl. Norv. ii 102 = tenella.
Korroo, *Royle, Illustr. Bot. Himal.* 278. t. 68. f. 2.—Reg. Himal.
Laengstii, Hausm. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 378 = *Charpentieri*.
lacvigata, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 369.—Mexic.
lancoolata, *Griseb. in DC. Prod.* ix. 102.—Mexic.
lanciifolia, Rafin, Danm. og Holst. Fl. ii. 317 = G. Amarella.
lasiopoda, Schur, Enum. Pl. Transs. 457 = phlogifolia.
lasiocaulis, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 568 = quadrifaria.
Ledebourii, Reichb. Fl. Germ. Excurs. 436 = aquatica.
leptina, *Raflu. ex Steud. Nom.* ed. i. 462, nomen.—Hab. ?
lignea, Murr. ex Stend. l. c. ed. II. i. 674 = Enicostema littorale.
limosiflolides, *H. B. & K. Nov. Gen. et Sp.* iii. 167.—Ecuador.
linariæfolia, Lam. Encyc. ii. 641 = Erythræa linariifolia.
linariifolia, Lam. Fl. Fr. ii. 398 = G. Pneumonanthe.
linearis, *Prosl. Gent. Diss.* 37.—Am. bor.
linoiata, *Franch. in Bull. Soc. Bot. Fr.* xxxi. (1884) 375.—China.
linqulata, Agardh, in Phys. Sällsk. Årsb. (1824) 22 = G. Amarella.

GENTIANA:—
linifora, *H. B. & K. Nov. Gen. et Sp.* iii. 171.—Peruv.
linifolia, Salisb. Prod. 137 = G. Pneumonanthe.
linifolia, Willd. ex Schult. Syst. vi. 184 = graminea.
livonica, Eschh. ex Griseb. Spic. Fl. Rumel. et Bith. 241 = G. Amarella.
Loderi, *Hook. f. Ic. Pl.* t. 1440 A.—Reg. Himal.
longiflora, Lam. Tabl. Encyc. i. 485 = altaica.
Loureirii, Griseb. in DC. Prod. ix. 108 = indica.
lutea, Krock. Fl. Siles. t. 885 = asclepiadea.
lutea, *Linn. Sp. Pl.* 227.—Europ.; As. Min.
lutea, Ruiz & Pav. ex G. Don, Gen. Syst. iv. 181 = pedunculata.
macrantha, *Griseb. in DC. Prod.* ix. 102.—Mexic.
macrocalyx, *Lxx. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 18.—Mexic.
macrocarpophora, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = G. Pneumonanthe.
macrophylla, Bertol. Fl. Ital. iii. 79 = Burseri.
macrophylla, *Pall. Fl. Ross.* i. ii. 108. t. 98.—Sibir.
magellanica, *Gaudich. in Freyc. Voy. Bot.* 449.—Reg. Magell.
marginata, Griseb. Gen. et Sp. Gent. 278 = carinata.
marginata, Turcz. ex Bess. in Flora, xvii. (1834) i. Beibl. 19 = aurea.
marginata, Wall. Cat. n. 4391 = capitata.
maritima, La Pylaie, ex Griseb. in DC. Prod. ix 101 = detonsa.
maritima, Linn. Mant. i. 55 = Erythræa maritima Mendesii, *Griseb. Gen. et Sp. Gent.* 292.—Calif. mexicana, *Griseb. l. c.* 243.—Mexic.
Meyeniana, *Griseb. l. c.* 217.—Peruv.
Meyenii, Griseb. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 48 = præc.
micans, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 112.—Reg. Himal.
micrantha, *Aitch. ex Hemsl. in Journ. Linn. Soc.* xix. (1882) 177.—Afghan.
microcalyx, *Lemmon, ex A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 222.—Am. bor. occ.
microphylla, Griseb. in DC. Prod. ix. 105 = sedifolia.
minima, *Phil. in Linnæa*, xxix. (1857–58) 13.—Chili.
minima, Vill. Hist. Pl. Dauph. ii. 526 = nivalis.
minutissima, *Beiss. Fl. Orient.* iv. 71.—Afghan.
Mitsnofana, *Schult. Syst.* vi. 186.—Japon.
modesta, *Phil. in Linnæa*, xxix. (1857–58) 109.—Chili.
monnieroides, Benth. Pl. Hartw. 327 = nummularifolia.
montana, *Forst. f. Prod.* 21.—Austral.
montana, Nees, in Nov. Act. Nat. Cur. ix. (1818) 170 = obtusifolia.
Moorcroftiana, Griseb. in DC. Prod. ix. 96, pro parte = Jaeschkea latisepala.
Moorcroftiana, *Wall. Cat.* n. 4390.—Reg. Himal.
multicaulis, *Gill. ex Griseb. Gen. et Sp. Gent.* 225.—Chili.
multiflora, *Griseb. in Goett. Abh.* xxiv. (1879) 236.—Reg. Argent.
Naegeli x, Brongg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 108.—Europ.
nana, Wulf. in Jacq. Misc. i. 161. t. 18. f. 3 = tenella.
Newberryi, *A. Gray, in Proc. Am. Acad.* xi. (1876) 84.—Am. bor. occ.
nigrescens, Schur, Enum. Pl. Transs. 462 = *chloræfolia*.
nigricans, *Lam. Encyc.* ii. 645.—Hab.?
nikoënsis, Franch. & Sav. Enum. Pl. Jap. i. 322; ii. 449 = frigida.
nitida, *Griseb. Gen. et Sp. Gent.* 216.—Peruv.
nivalis, *Linn. Sp. Pl.* 229.—Reg. bor. et arct.
nivalis, Poir. Encyc. Suppl. ii. 733 = tenella.
Novæ-Zelandiæ, *Armst. in Trans. N. Z. Inst.* iv. (1872) 291.—N. Zel.
nubigena, *Edgew. in Trans. Linn. Soc.* xx. (1846) 85.—Reg. Himal.
nudicaulis, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 285.—Reg. Himal.; Burma.
nummulariæfolia, Benth. Pl. Hartw. 228 = selaginifolia.
nummulariifolia, *Griseb. Gen. et Sp. Gent.* 228.—Ecuador.
nutans, Bunge, in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 223 = prostrata.
obliqua, Nees, in Nov. Act. Nat. Cur. ix. (1818) 166 = germanica.

GENTIANA:—
obtusifolia, Fries, Nov. Fl. Suec. 76 = G. Amarella.
obtusifolia, *Willd. Sp. Pl.* i. 1347.—Europ. med.; As. Min.
ochroleuca, *Froel. Gent. Diss.* 35.—Am. bor.
ochroleuca, Sims, Bot. Mag. t. 1651 = alba.
octaflora, Vitm. Summa Pl. ii. 114 = Endicostema littorale.
Oederi, Soland. ex Bab. in Journ. Linn. Soc. xi. (1871) 318 = tenella.
Olgæ, *Regel, ex Schmalh. in Regel, Pl. Nov. Fedsch.* 56.—Turkestan.
oligosperma, Griseb. in DC. Prod. ix. 94 = Jaeschkea gentianoides.
Olivieri, Griseb. Gen. et Sp. Gent. 276 = decumbens.
oppositifolia, Zawad. ex Griseb. l. c. 245 = cancasica.
orbiculata, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = bavarica.
orbiculata, Heyne, ex Wall. Cat. n. 4395 = quadrifaria.
oregana, *Engelm. ex A. Gray, Syn. Fl. N. Am.* ii. i. 122.—Am. bor. occ.
ornata, *Wall. Cat.* n. 4386.—Reg. Himal.
Ottonis, *Phil. in Linnæa*, xxxiii. (1864–65) 177.—Chili.
ovalis, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 269.—Mexic.
ovata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 458 = Sebæa ovata.
paliens, Thunb. Mus. Acad. Upa. xi. (1792) 180, nomen (Sebææ sp.).—Hab.?
palustris, All. Fl. Pedem. i. 100 = Swertia perennis.
palustris, Lam. Tabl. Encyc. i. 486 = Erythræa pulchella.
paniculata, Lam. Fl. Fr. ii. 290 = Swertia perennis.
pannonica, Schleich. Cat. (1821); ex Gand. Fl. Helv. ii. 275 = *Gaudiniana*.
pannonica, *Scop. Fl. Carn.* ed. II. i. 182.—Europ.
papillosa, *Franch. in Bull. Soc. Bot. Fr.* xxxi. (1884) 374.—China.
Parryi, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 218. t. 10.—Am. bor. occ.
patagonica, *Griseb. Gen. et Sp. Gent.* 287.—Reg. Magell.
Pavonii, *Griseb. in DC. Prod.* ix. 94.—Peruv.
Pearcei, *Phil. in Anal. Univ. Chil.* i. (1861) 65.—Chili.
pedicellata, Wall. Cat. n. 4394 = quadrifaria.
pedunculata, *D. Don, ex G. Don, Gen. Syst.* iv. 181.—Peruv.
pedunculosa, Willd. ex Schult. Syst. vi. 185 = linifolia.
pedunculata, Domb. ex Griseb. in DC. Prod. ix. 92 = dissitiflora.
pedunculata, Royle, ex G. Don, Gen. Syst. iv. 182 = tenella.
pendula, *Griseb. Gen. et Sp. Gent.* 232.—Ecuador.
pensylvanica, *Hill, Veg. Syst.* viii. 59 (Quid?).—Am. bor.
perfoliata, Linn. Sp. Pl. 232 = Chlora perfoliata.
peruviana, *Griseb. in Goett. Abh.* xxiv. (1874) 236.—Reg. Argent.
peruviana, Lam. Encyc. ii. 642 = Erythræa chilensis.
phlogifolia, *Schott & Kotschy, in Bot. Zeit.* ix. (1851) 161.—Transsylv.
phyllocalyx, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 116.—Reg. Himal.; China.
Piaselezii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 498.—China.
pinifolia, *Ruiz & Pav. ex G. Don, Gen. Syst.* iv. 182.—Peruv.
pinnatifida, *Griseb. Gen. et Sp. Gent.* 291.—Ins. Sitcha.
plebeja, Cham. in Linnæa, i. (1826) 181 = G. Amarella.
pleurogynoides, Griseb. Gen. et Sp. Gent. 236 = montana.
plicata, Willd. ex Schult. Syst. vi. 184 = spathacea.
Pneumonanthe, *Linn. Sp. Pl.* 228.—Europ.; As. bor.
Pneumonanthe, Michx. Fl. Bor. Am. i. 176 = linearis.
pneumonanthoides, Schur, Enum. Pl. Transs. 457 = G. Pneumonanthe.
pneumonanthoides, *Wender. ex Steud. Nom. ed.* II. i. 675, nomen.—Hab.?
podocarpa, *Griseb. in Goett. Abh.* xix. (1874) 210.—Am. austr.
pœcantha, Fisch. ex Griseb. Gen. et Sp. Gent. 211 = Pleurogyne carinthiaca.

GENTIANA:—
Porphyrio, J. F. Gmel. Syst. 462 = angustifolia.
portensis, Brot. Fl. Lusit. i. 278 = Erythræa Massoni.
pratensis, Froel. Gent. Diss. 38 = G. Amarella.
primulæfolia, *Franch. in Bull. Soc. Bot. Fr.* xxxi. (1884) 375.—China.
primulæfolia, *Griseb. Gen. et Sp. Gent.* 221.—Peruv.
procumbens, Stend. Nom. ed. II. i. 768, in syn. = prostrata.
propinqua, *Richardz. in Frankl. Narr.* 1st. *Journ.* 734.—Sibir.; Am. bor.
prostrata, *Haenke, in Jacq. Coll.* ii. 66. t. 17.—Reg bor. et austr.
prostrata, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv (1842) 408 = aquatica.
prostrata, Schleich. ex Cham. & Schlecht. in Linnæa, i. (1826) 184 = verna.
prostrata, Wahlenb. Veg. Helv. 47 = bavarica.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 503.—China.
Pseud-Amarella, Stev. ex Herd. in Act. Hort. Petrop. i. (1872) 437 = aurea.
Pseudo-Pneumonanthe, Schult. Syst. vi. 146 = linearis.
puberula, *Michx. Fl. Bor. Am.* i. 176.—Am. bor.
pudica, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 497.—China.
pulchella, Salisb. Prod. 137 = G. Amarella.
pulchella, Sw. in Vet. Acad. Handl. Stockh. (1783) 84. t. 3. f. 3, 9 = Erythræa ramosissima.
pulla, *Griseb. in Goett. Abh.* xix. (1874) 209.—Reg Argent.
Pulmonaria, Turcz. in Flora (1834) i. Beibl. 19, nomen; et in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 317 = tenella.
pumila, Gouan, Fl. Monsp. 473 = Erythræa maritima.
pumila, Griff. Ic. Pl. Asiat. t. 884 = quadrifaria.
pumila, *Jacq. Enum. Vindob.* 41, 218.—Europ.
pumila, Suter, Fl. Helv. i. 142 = verna.
punctata, Georgi, Reise, i. 208 = frigida.
punctata, Jacq. Obs. Bot. ii. 17 = pannonica.
punctata, *Linn. Sp. Pl.* 227.—Europ. med.
punctata, Pall. Reise, iii. 317 = decumbens.
punctata, Vill. Hist. Pl. Dauph. ii. 523 = Burseri.
punices, *Wedd. Chlor. And.* ii. 70.—Peruv.; Bolivia.
purpurascens x, Beck, Ind. ad DC. Prod. iii. 195.—Europ.
purpurea, *Linn. Sp. Pl.* 227.—Europ.
purpurea, Schrank, Baier. Fl. i. 509 = pannonica.
purpurea, Vill. Hist. Pl. Dauph. ii. 528 = punctata.
purpurea, Wall. Fl. Carol. 109 = angustifolia.
pusilla, Lam. Encyc. ii. 645 = Cicendia pusilla.
pygmæa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 111.—Reg. Himal.
pygmæa, *Regel & Schmalh. in Regel, Pl. Nov. Fedsch.* 54.—Kokan.
pyramidalis, Nees, in Nov. Act. Nat. Cur. ix. (1818) 164 = obtusifolia.
pyramidalis, Willd. ex Griseb. Gen. et Sp. Gent. 239 = G. Amarella.
pyrenaica, *Linn. Mant.* i. 55.—Europ.; As. Min.
quadrangularis, Lam. Encyc. ii. 645 = Microcala quadrangularis.
quadrifaria, *Blume, Bijdr.* 847.—Ind. or.; Malaya; China.
quadrifolia, Linn. Sp. Pl. ed. II. 1671 = Chlora quadrifolia.
quinqueflora, A. Gray, in Torr. Bot. Mex. Bound. 157, partim = Wrightii.
quinqueflora, *Hill, Veg. Syst.* viii. 57; *Lam. Encyc.* ii. 643.—Am. bor.
quinquefolia, Linn. Sp. Pl. 230 = quinqueflora.
quinquefolia, Oeder, in Fl. Dan. t. 344 = aurea.
quinquepartita, Domb. ex Griseb. in DC. Prod. ix. 90 = cuspidata.
radicata, *Griseb. Gen. et Sp. Gent.* 229.—Peruv.
Raimondiana, *Wedd. Chlor. And.* ii. 210.—Peruv.
ramosissima, Brot. Fl. Lusit. i. 276 = Erythræa tenuiflora.
ramosissima, *Phil. in Anal. Univ. Chil.* (1879) 311.—Chili.
ramosissima, Vill. Fl. Delph. 23; Hist. Pl. Dauph. ii. 530 = Erythræa ramosissima.
ramulosa, *Tinzirow, in Bull. Soc. Murith.* i. (1868) 28.—Europ.
rapunculoides, Willd. ex Schult. Syst. vi. 185 = diffusa.

GENTIANA :—
recurvata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 110.—Reg. Himal.; China.
Renardi, *Regel, in Trautv. Regel, Maxim. & Winkl. Decas. Pl. Nov.* 7.—Turkest.
rhaetica, *A. & J. Kern. Sched. Fl. Exsicc.* ii. 124.—Europ.
rigida, *Rafin. Med. Fl.* i. 212.—Am. bor.
Rima, *G. Don, Gen. Syst.* iv. 181.—Peruv.
riparia, Buch.-Ham. ex Wall. Cat. n. 4892 = decemfida.
riparia, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 706 [bis] = aquatica.
rivularis, Thunb. Mus. Acad. Upa. xi. (1792) 180, nomen (Sebaea sp.).—Hab. ?
robusta, *King, ex Hook. f. Ic. Pl.* t. 1439.—Reg. Himal.
Romanzovii, Ledeb. ex Bunge, in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 215 = frigida.
Rostani, Reut. in Bull. Soc. Dauph. (1877) 119 = bavarica.
rotata, Domb. ex Wedd. Chlor. And. ii. 63 = Dombeyana.
rotata, Froel. Gent. Diss. 105 = Pleurogyne rotata.
rotata, Willd. Sp. Pl. i. 1851 = Pleurogyne carinthiaca.
rotundifolia, Hoppe, ex Griseb. Gen. et Sp. Gent. 265 = bavarica.
Royleana, Wall. Cat. n. 4893 = decemfida.
rubicunda, *Pterocarp. in Bull. Soc. Bot. Fr.* xxxi. (1884) 378.—China.
rubra, [Clairv.] Man. Herb. 73 = *hybrida*.
rubra, Uspenski, Desc. Jekatherinenb. 77 (Erythraeae sp. ?).—Sibir. ural.
rubricaulis, Schwein. in Long, Exped. Winnep. ii. 384 = ochroleuca.
Ruizii, *Griseb. in DC. Prod.* ix. 96.—Peruv.
rupicola, Benth. Pl. Hartw. 228 = cernstioides.
rupicola, Griseb. ex Wedd. Chlor. And. ii. 53 = primulaefolia.
rupicola, *H. B. & K. Nov. Gen. et Sp.* iii. 167.—Ecuador.
Kurihiana, Cham. & Schlecht. in Linnaea, i. (1826) 176 = propinqua.
sabauda, Boiss. & Reut. ex Reichb. Ic. Fl. Germ. xvii. 101. t. 1900 = acaulis.
salpinx, *Griseb. in Linnaea*, xxii. (1849) 44.—Mexic
Saponaria, Froel. Gent. Diss. 32 = Andrewsii.
Saponaria, *Linn. Sp. Pl.* 228.—Am. bor.
Saponaria, Walt. Fl. Carol. 109 = ochroleuca.
saponariifolia, Stokes, Bot. Mat. Med. ii. 51 = G. Saponaria.
saxicola, *Griseb. Gen. et Sp. Gent.* 216.—Peruv.
saxifragoides, *H. B. & K. Nov. Gen. et Sp.* iii. 168.—Peruv.
saxosa, Forst. f. in Vet. Akad. Handl. Stockh. xxxvii. (1777) 184 = montana.
scabra, *Bunge, Vers. Alt.* 21.—As. or.
scandens, Lour. Fl. Cochinch. 171 = Paederia foetida.
Sceptrum, *Griseb. Gen. et Sp. Gent.* 293.—Am. bor. occ.
schistocalyx, C. Koch, in Linnaea, xxiii. (1850) 585 = asclepiadea.
scilloides, Linn. f. Suppl. 175 = Erythraea Massoni.
scopulorum, *Wedd. Chlor. And.* ii. 67.—Bolivia.
sedifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 173.—Ecuador.
selaginifolia, *Griseb. in Linnaea*, xxii. (1849) 42.—N. Granat.
Selatium, *Griseb. in DC. Prod.* ix. 93.—Peruv.
semifida, Hoffingg. ex Reichb. Fl. Germ. Excurs. 428 = pannonica.
septemfida, *Pall. Fl. Ross.* i. ii. 101. t. 92. f. 3.—Orient.; As. bor.
serpentaria, Rafin. Ann. Nat. 13 = ochroleuca.
serpyllifolia, Lam. Encyc. ii. 640 = bavarica.
Serra, *Franch. in Bull. Soc. Bot. Fr.* xxxi. (1884) 376.—China.
serrata, Gunner, Fl. Norveg. ii. 101 = detonsa.
serrata, Lam. Fl. Fr. ii. 294 = verna, bavarica.
Sessei, *Griseb. in DC. Prod.* ix. 113.—Mexic.
sessilis, Linn. Sp. Pl. 231 (= *Chlora sessilis*).—Reg. Argent.
setiflora, Bunge, in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 242 = propinqua.
setigera, *A. Gray, in Proc. Am. Acad.* xi. (1876) 84.—Calif.
sexfida, Schult. Syst. vi. 156, in obs. = verna.
Shastiana, *Rafin. Med. Fl.* i. 211.—Am. bor.

GENTIANA :—
sikkimensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 114.—Reg. Himal.; China.
simplex, *A. Gray, in Pacif. Rail. Rep.* v. 87. t. 16.—Am. bor. occ.
spathacea, *H. B. & K. Nov. Gen. et Sp.* iii. 178.—Mexic.
spathulata, Bartl. ex Reichb. Pl. Crit. i. 78 = obtusifolia.
spathulata, *Pursger, Krit. Vergl. Gouv. Mosk. Wildwachs. Pfl.* 15. t. 11. f. 3.—Russia.
spicata, Linn. Sp. Pl. 230 = Erythraea spicata.
spuria, Lebert, ex Griseb. Gen. et Sp. Gent. 301 = *Gaudiniana*.
squarrosa, *Ledeb. in Mém. Acad. Pétersb.* v. (1812) 527.—Reg. Himal.; As. bor.
stellarioides, *Griseb. Gen. et Sp. Gent.* 225.—Peruv.
Stellersana, Cham. & Schlecht. in Linnaea, i. (1826) 188 = Pleurogyne carinthiaca.
stipitata, Edgew. in Trans. Linn. Soc. xx. (1846) 84 = cachemirica.
Stolicska, Kurz, ex C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 488 = aurea.
strumifera, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 502.—China.
striata, *Maxim. l. c.*—China.
strycta, Klotzsch, in Bot. Ergeb. Waldem. Reise, 90 = detonsa.
stricta, Willd. ex Schult. Syst. vi. 185 = diffusa.
Stuartiana, *A. & J. Kern. Sched. Fl. Exsicc.* ii. 122.—Austria.
stylophora, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 118.—Reg. Himal.; China.
subulata, Domb. ex Wedd. Chlor. And. ii. 71 = fruticulosa.
sulcata, Willd. Sp. Pl. i. 1851 = Pleurogyne rotata.
suecica, Griseb. Gen. et Sp. Gent. 283 = diffusa.
sylvestris, Bray, ex Griseb. l. c. 241 = G. Amarella.
Tauschex, Kurz, in Journ. Bot. v. (1867) 241 = *Jaeschkei*.
tenella, *Asara, Voy.* ii. 491.—Am. austr.
tenella, *Roth. in Kioh. Sbr. Selsh.* x. (1770) 486.—Reg. bor. et arct.
tenuiflora, Jan, ex Nym. Consp. 500 = G. Amarella.
tenuis, Griseb. Gen. et Sp. Gent. 250 = G. Amarella.
tergloensis, Hacq. Pl. Carn. 9. t. 2. f. 5 = imbricata.
ternifolia, *Franch. in Bull. Soc. Bot. Fr.* xxxi. (1884) 377.—China.
tetragona, Mayer, Samml. Phys. Aufs. i. (1791) 194 = G. Amarella.
tetragona, Roth, Tent Fl. Germ. ii. i. 290; Lam. Tabl. Encyc. i. 488 = tenella.
Thomasiana, Gaud. ex Nym. Consp. 498, sphalm. = seq.
Thomasii, Gillabox, ex Vill. in Room. Collect. 169 = *hybrida*.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 108.—Reg. Himal.
Thunbergii, *Griseb. in DC. Prod.* ix. 108.—As. or.
Thunbergii, Sieb. & Zucc. Fl. jap. Fam. Nat. ii. 84 = Zollingeri.
thyrsoides, *Hook. Bot. Misc.* ii. (1831) 227.—Peruv.
tianschanica, *Regel, Sert. Tianach.* 61.—As. centr.
tibetica, *King, ex Hook. f. Ic. Pl.* t. 1441.—Reg. Himal.; China.
Torreyana, *Rafin. Med. Fl.* 212.—Am. bor.
trichostemma, *Wedd. Chlor. And.* ii. 56.—Ecuador.
triflora, *Pall. Fl. Ross.* i. ii. 108.—Sibir. or.
tristriata, Turcz. in Flora, (1834) i. Beibl. 19 = tenella.
tubiflora, *Wall. Cat.* n. 4888.—Reg. Himal.
uliginosa, Willd. Sp. Pl. i. 1847 = G. Amarella.
umbellata, Bleb. Fl. Taur. Cauc. iii. 188 = aurea.
umbellata, Rutz & Pav. ex G. Don, Gen. Syst. iv. 187 = Rafini.
unalaccensis, Cham. ex Bunge, in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 240 = seq.
unalaschcensis, Cham. in Linnaea, i. (1826) 175, in nota = aurea.
uniflora, Georgi, Reise, i. 208 = verna.
uniflora, Willd. Sp. Pl. i. 1846 = germanica.
utriculosa, *Linn. Sp. Pl.* 229.—Europ. med.
utriculosa, Wahlenb. Veg. Helv. 48 = verna.
vaginalis, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 469 = Lagasia latifolia.
vaginalis, *Griseb. Gen. et Sp. Gent.* 215.—Peruv.

GENTIANA :—
vaginalis, Griseb. ex Wedd. Chlor. And. ii. 53 = primulaefolia.
Vaillantii, *F. W. Schmidt, in Mayer, Samml. Phys. Aufs.* i. (1791) 191.—Europ.
variabilis, *Rupr. Sert. Tianach.* 60.—As. centr.
ventricosa, *Griseb. Gen. et Sp. Gent.* 259.—Canada.
vernana, *Wall. Cat.* n. 4889.—Reg. Himal.
verna, *Linn. Sp. Pl.* 228.—Europ.; Reg. Caucas.
verticillaris, Ketz. Obs. ii. 15 = Endostema littorale.
verticillata, Linn. Syst. ed. X. 952 = Endostema littorale.
vertucillata, *Wedd. Chlor. And.* ii. 209.—Peruv.
villosa, Linn. Sp. Pl. 229 = ochroleuca.
violacea, *Rusz & Pav. ex G. Don, Gen. Syst.* iv. 195, in syn.—Peruv.
virdis, *Griseb. in Linnaea*, xxii. (1849) 43.—Venezuela.
viscosa, Ait. Hort. Kew. ed. I. i. 321 = Ixanthus viscosus.
volubilis, D. Don, Prod. Fl. Nep. 120 = Crawfurdia japonica.
Wagiana, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 20, nomen.—Rossia.
Waldjevi, *Regel & Schmalh. l. c.* vi. (1879) 384.—Turkest.
Weschniakowi, *Regel, l. c.* viii. (1884) 687.—Turkest.
Wislizenii, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 215. t. 7.—Arizona.
Wrightii, *A. Gray, Syn. Fl. N. Am.* ii. i. 118.—Arizona.
yunnanensis, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1884) 376.—China.
zeylanica, Griseb. Gen. et Sp. Gent. 274 = quadrifaria.
Zollingerii, *Fawcett, in Journ. Bot.* xxi. (1883) 183.—China; Japon.

GENTIANELLA, Moench, Meth. 482 (1794) = **Gentiana**, Tourn.
barbata, Bercht. & Presl, Rostl. i. Gentian. 22 = Gentiana detonsa.
ciliata, Borckh. in Roem. Arch. i. i. 29 = Gentiana ciliata.
crinita, Bercht. & Presl, Rostl. i. Gentian. 21 = Gentiana detonsa.
Crucata, Moench, Meth. Suppl. 188 = Gentiana Cruciata.
dentosa, Bercht. & Presl, Rostl. i. Gentian. 22, sphalm. = seq.
detonsa, G. Don, Gen. Syst. iv. 179 = Gentiana detonsa.
fimbrata, Delarb. Fl. Auv. ed. II. 31 = Gentiana ciliata.
macrantha, D. Don, ex G. Don, Gen. Syst. iv. 179 = Gentiana macrantha.
macrophylla, G. Don, ex Steud. Nom. ed. II. i. 676 = Gentiana macrantha.
serrata, Borckh. in Roem. Arch. i. i. 29 = Gentiana detonsa.
tetrandra, Moench, Meth. 482 = Gentiana campestris.

GEOBINA, Rafin. Fl. Tellur. iv. 49 (1836) = **Goodyera**, R. Br. (Orchid.).

GEOCALLIS, Horan. Prod. Monog. Scitam. 33 (1862) = **Amomum**, Linn. (Scitamin.).
fasciculata, Horan. l. c. = A. fasciculatum.

GEOCARVUM, Coss. Notes Crit. 112 (1851) = **Carum**, Rupp. (Umbellif.).
capillifolium, Coss. l. c. 113 = C. tenuifolium.
pumilum, Nym. Syll. 166 (*Biasolettia pumila*).—Graecia.
tenuifolium, Coss. in Bourg. in Bull. Soc. Bot. Fr. ii. (1855) 648 = C. tenuifolium.

GEOCHORDA, Cham. & Schlecht. in Linnaea, iii. (1828) 11. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 982.
cuneata, *Cham. & Schlecht. l. c.*—Bras.

GEOCOCCUS, J. Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 52. *CRUCIFERAE*, Benth. & Hook. f. i. 83.
pusillus, *J. Drumm. ex Harv. l. c.*—Austral.

GEODORUM, Jacks. in Andr. Bot. Rep. t. 626 (1810).
ORCHIDEÆ, Benth. & Hook. f. iii. 538.
CISTELLA, Blume, Bijdr. 398. t. 58 (1825).
ORTMANNIA, Opiz, in Flora, xvii. (1834) 592.
OTANDRA, Salisb. in Trans. Hort. Soc. i. (1812) 298, sbeque char.
appendiculatum, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 357.—Assam.
attenuatum, *Griff. l. c.* 358.—Burma.
bicolor, Voigt, Hort. Suburb. Calc. 628 = Eulophia herbacea.
bractеatum, Voigt, l. c. 629 = Eulophia bracteosa ?
candidum, Wall. Cat. n. 7374 = pallidum.
citrinum, *Jacks. in Andr. Bot. Rep.* t. 626.—Ind. or.
dilatatum, *R. Br. in Ait. Hort. Kew.* ed. II. v. 207.—Ind. or.
Duperreanum, *Pierre, in Rev. Hortic.* (1892) 501.—Cochinch.
fucatum, *Lindl. Bot. Reg.* t. 1657.—Zeylan.
javanicum, *Lindl. Fol. Orch. Geod.* 2.—Java.
laxiflorum, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 356.—Assam.
longifolium, Voigt, Hort. Suburb. Calc. 628 = Cymbidium cyperifolium.
pallidum, *D. Don, Prod. Fl. Nep.* 31.—Reg. Himal.
pictum, *Lindl. Gen. et Sp. Orch.* 175.—Austral.
pictum, Link, Ic. Pl. Hort. Reg. Berol. t. 14 = Eulophia maculata.
plicatum, Voigt, Hort. Suburb. Calc. 629 = Phaius callosus.
purpureum, *R. Br. in Ait. Hort. Kew.* ed. II. v. 207.—Ind. or.
ramentaceum, Voigt, Hort. Suburb. Calc. 628 = Eulophia campestris.
rariflorum, *Lindl. Fol. Orch. Geod.* 2.—Assam.
semicristatum, *Lindl. l. c.*—Malaya.

GEOFFRAEA, Linn. Sp. Pl. ed. II. 1043 (1763).
LEGUMINOSÆ, Benth. & Hook. f. i. 551.
UMARI, Adans. Fam. ii. 342 (1763).
acutifolia, Stokes, Bot. Mat. Med. iv. 46 = Andira inermis.
Bredemeyeri, H. B. & K. Nov. Gen. et Sp. vi. 379 = superba.
discolor, *Benth. in Hook. Journ. Bot.* ii. (1840) 69.—Bras.
inermis, Sw. Prod. Veg. Ind. Occ. 106 = Andira inermis.
obtusifolia, Stokes, Bot. Mat. Med. iv. 46 = Andira retusa.
pubescens, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 121 = Andira retusa.
racemosa, Poir. Encyc. viii. 188 = Andira excelsa.
retusa, Poir. l. c. 182 = Andira retusa.
Sieberi, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 134.—Ins. Trinitat.
spinosa, *Jacq. Enum. Pl. Carib.* 28 ; *Linn. Sp. Pl.* ed. II. 1043.—Ind. occ.
spinulosa, Mart. Reise Bras. ii. 788 = Andira vermifuga.
superba, *Humb. & Bonpl. Pl. Aequin.* ii. 69. t. 100.—Am. austr.
surinamensis, Bondt, Cort. Geoff. Surinam, 9 = Andira retusa.
tomentosa, *Poir. Encyc.* viii. 182.—Afr. trop.
vermifuga, Mart. Reise Bras. ii. 788 = Andira vermifuga.
violacea, *Pers. Syn.* ii. 378.—Guiana.

GEOFFROEA, Jacq. Enum. Pl. Carib. 7 (1763) = **Geoffraea**, Linn. (Legumin.).

GEOFFROYA, Murr. Syst. ed. XIII. 556 (1774) = **Geoffraea**, Linn. (Legumin.).

GEOHERPUM, Willd. ex Schult. Mant. iii. 18, 350 (1827) = **Mitchella**, Linn. (Rubiac.).
alsinefolium, Willd. ex Schult. l. c. 350 = M. ovata.

GEOLOBUS, Rafin. New Fl. Am. i. 81 (1836) = **Voandzeia**, Thou. (Legumin.).
flavus, Rafin. l. c. 85 = V. subterranea.

GEOMITRA, Becc. Malesia, i. (1878) 260. tt. 10, 11.
BURMANNIACEÆ, Benth. & Hook. f. iii. 436.
clavigera, *Becc. l. c.* 251.—Malaya.
episcopalis, *Becc. l. c.* 250.—Malaya.

GEONOMA, Willd. in Mém. Acad. Berl. (1804) 37.
PALMÆ, Benth. & Hook. f. iii. 913.
GYNESTUM, Poit. in Mém. Mus. Par. ix. (1822) 387.
VOUAY, Aubl. Pl. Gui. ii. App. 99 (1775), nomen prius.

GEONOMA :—
acaulis, *Mart. Hist. Nat. Palm.* ii. 18. tt. 4, 19.—Bras.
acutifora, Mart. l. c. 10. t. 9 = bacuifera.
amazonica, *Kerst. ex H. Wendl. in Kerch. Palm.* 245.—Bras.
ambigua, *Spruce, in Journ. Linn. Soc.* xi. (1871) 111.—Am. trop.
antiocuensis, *Hort. ex H. Wendl. in Kerch. Palm.* 245, nomen.—N. Granat.
Appuniana, *Spruce, in Journ. Linn. Soc.* xi. (1871) 106.—Am. trop.
Aricuna, Barb.-Rodr. Enum. Palm. Nov. Prot. App. 40 = Schottiana.
armilliaris, *Mart. Hist. Nat. Palm.* ii. 17. t. 18.—Fluv. Amazon.
aspidifolia, *Spruce, in Journ. Linn. Soc.* xi. (1871) 113.—Am. trop.
bacuifera, *Kunth, Enum. Pl.* iii. 233.—Guiana.
barbigera, *Barb.-Rodr. Palmiers*, 45.—Bras.
Barthii, *Engel. in Linnaea*, xxxiii. (1864–65) 688.—N. Granat.
bifurca, *Drude & H. Wendl. in Mart. Fl. Bras.* iii. ii. 504.—Bras.
bijuga, *Kerst.-Rodr. Enum. Palm. Nov.* 10.—Bras.
binervia, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 88.—Costa Rica.
Blanchetiana, *H. Wendl. ex Drude, in Mart. Fl. Bras.* iii. ii. 504.—Bras.
Bluntii, *Hort.—Cf. Gard. Chron.* (1881) i. 766.—Hab. ?
brachystachis, *Barb.-Rodr. Enum. Palm. Nov.* 10.—Bras.
brevispatha; *Barb.-Rodr. l. c. Prot. App.* 41.—Bras.
Bronguartii, *Mart. Palmet. Orb.* 24. t. 15. f. 2.—Bras.
caespitosa, *H. Wendl. ex Drude, in Mart. Fl. Bras.* iii. ii. 506.—Bras.
calophylla, *Barb.-Rodr. Palmiers*, 48.—Bras.
camana, *Trail, in Journ. Bot.* xiv. (1876) 324.—Bras.
Capanema, Barb.-Rodr. Enum. Palm. Nov. 9.—Bras.
Carderi, *Hort.—Cf. Gard. Chron.* (1877) i. 48.—N. Granat.
caudescens, *H. Wendl. ex Drude, in Mart. Fl. Bras.* iii. ii. 504.—Bras.
chelidura, *Spruce, in Journ. Linn. Soc.* xi. (1871) 111, 117.—Am. trop.
compacta, *Linden, Illustr. Hortic.* xxviii. (1881) 81, nomen.—Venezuela.
conigua, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 112.—Costa Rica.
corallifera, Hook. ex Morr. in Belg. Hortic. v. (1855) 168 = Chamaedorea Ernesti-Augusti.
cuneata, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 104.—Costa Rica.
Demarestii, Prits. Ind. Ic. 496, sphalm. = Desmarestii.
densa, *H. Wendl. in Linnaea*, xxviii. (1856) 338.—N. Granat.
densiflora, *Spruce, in Journ. Linn. Soc.* xi. (1871) 112.—Am. trop.
Desmarestii, *Mart. Palmet. Orb.* 28. t. 11. f. 2.—Bras.
deversa, *Kunth, Enum. Pl.* iii. 231.—Guian.
disodon, Spruce, in Journ. Linn. Soc. xi. (1871) 110 = multiflora.
dulcis, Wright, ex Griseb. Cat. Pl. Cub. 222 (Calyptrogynes sp.).—Cuba.
edulis, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 100.—Costa Rica.
elegans, *Mart. Hist. Nat. Palm.* ii. 144. t. 18 A.—Bras.
erythrospadicea, Barb.-Rodr. Enum. Palm. Nov. ed. II. Prot. App. 41 = trinervia.
falcata, *Barb.-Rodr. Enum. Palm. Nov.* 10.—Bras.
Fendleriana, *Spruce, in Journ. Linn. Soc.* xi. (1871) 108.—Am. trop.
fenestrata, *H. Wendl. in Kerch. Palm.* 245 = Malortiea gracilis.
ferruginea, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 110.—Costa Rica.
fiscellaria, *Mart. ex Drude, in Mart. Fl. Bras.* iii. ii. 496.—Bras.
flaccida, *H. Wendl. ex Spruce, in Journ. Soc.* xi. (1871) 108.—Costa Rica.
frigida, *Linden, Illustr. Hortic.* xxviii. (1881) 81.—N. Granat.
furcifrons, Drude, in Mart. Fl. Bras. iii. ii. 503 = praece.
Gamotiana, *Glas. ex Drude, l. c.* 496.—Bras.

GEONOMA :—
Ghiesbreghtiana, Lindl. & H. Wendl. in Linnaea, xxviii. (1856) 849 = Calyptrogyne Ghiesbreghtiana.
glauca, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 36 = Calyptrogyne glauca.
gracilis, Linden & André, Illustr. Hortic. xxi. (1874) t. 189 = Rhodelaea.
gracilis, *H. Wendl. in Kerch. Palm.* 245.—Costa Rica.
hexasticha, Spruce, in Journ. Linn. Soc. xi. (1871) 110 = multiflora.
Hoffmanniana, *H. Wendl. ex Spruce, l. c.* 106.—Am. trop.
imperialis, *Linden, Illustr. Hortic.* xxvii. (1881) 81.—N. Granat.
interrupta, *Mart. Hist. Nat. Palm.* ii. 8. t. 7.—Bras.
Irana, *Linden, Illustr. Hortic.* xxvii. (1881) 81.—Venezuela.
Jussieuana, *Mart. Palmet. Orb.* 24. t. 15. f. 1.—Bolivia.
lacerata, *Hort. ex Floral Mag.* viii. (1860) t. 446.—Am. centr.
laxiflora, *Mart. Hist. Nat. Palm.* ii. 12. t. 11.—Bras
leptospadix, *Trail, in Journ. Bot.* xiv. (1876) 327.—Bras.
Lindeniana, *H. Wendl. in Linnaea*, xxviii. (1856) 831.—N. Granat.
linearifolia, *Karst. l. c.* 411.—N. Granat.
longepetiolata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 38.—Costa Rica.
longe-vaginata, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 108.—Costa Rica.
longipes, *Hort. ex H. Wendl. in Kerch. Palm.* 245, nomen.—Hab. ?
macroclona, *Drude, in Mart. Fl. Bras.* iii. ii. 486.—Bras.
macrospatha, Spruce, in Journ. Linn. Soc. xi. (1871) 106 = bacillaris.
macrostachya, *Mart. Hist. Nat. Palm.* ii. 19. t. 20.—Bras.
magnifica, *Linden & H. Wendl. in Linnaea*, xxviii. (1856) 335.—Mexic.
margaritioides, Engel, in Linnaea, xxxiii. (1864–65) 682 = undata.
Margyraffia, *Engel, l. c.* 685.—N. Granat.
Martiana, *H. Wendl. in Linnaea*, xxviii. (1856) 342.—Bras.
martinicensis, *Mart. Palmet. Orb.* 28.—Ins. Martinic.
maxima, *Kunth, Enum. Pl.* iii. 232.—Guiana.
membranacea, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 106.—Guatemala.
metensis, *Karst. in Linnaea*, xxvii. (1856) 409.—Venezuela.
mexicana, *Liebm. ex Mart. Hist. Nat. Palm.* iii. 316.—Mexic.
microspadix, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 110.—Costa Rica.
microspatha, *Spruce, l. c.* 108.—Bras.
multiflora, *Mart. Hist. Nat. Palm.* ii. 7. tt. 4–6.—Am. austr.
multiflora, Wallace, Palm-trees Amaz. 68. t. xxiii = maxima.
negrensis, Spruce, in Journ. Linn. Soc. xi. (1871) 118 = multiflora.
obovata, *H. Wendl. ex Spruce, l. c.* 104.—Costa Rica.
Olfersiana, *Klotzsch, ex Drude, in Mart. Fl. Bras.* iii. ii. 506.—Bras.
oligoclona, *Trail, in Journ. Bot.* xiv. (1876) 325.—Bras.
Orbigniana, *Mart. Palmet. Orb.* 22. t. 11. f. 1.—Bolivia.
oxycarpa, *Mart. l. c.* 30.—Ins. S. Doming.
palmiris, *Barb.-Rodr. Enum. Palm. Nov.* 11.—Bras.
paniculigera, *Mart. Hist. Nat. Palm.* ii. 11. t. 10.—Bras.
paraënsis, *Spruce, in Journ. Linn. Soc.* xi. (1871) 112 = multiflora.
paraguassuia, *Karst. in Linnaea*, xxvii. (1856) 410.—Paraguay.
pauciflora, *Mart. Hist. Nat. Palm.* ii. 13. t. 13.—Bras.
peruana, *Spruce, in Journ. Linn. Soc.* xi. (1871) 112.—Am. trop.
pilosa, *Barb.-Rodr. Palmiers*, 43.—Bras.
pinnatifrons, *Willd. Sp. Pl.* iv. 893.—Am. trop.
platycuala, *Drude, in Mart. Fl. Bras.* iii. ii. 490.—Bras.
Plecana, *Mart. Palmet. Orb.* 28.—Venezuela.
Plumeriana, *Liebm. ex H. Wendl. in Kerch. Palm.* 246.—Ins. S. Doming.
Plumieriana, *Mart. Palmet. Orb.* 24.—Ins. S. Doming.

GEONOMA:—
Poeppigiana, *Mart. Palmet. Orb.* 35.—Peruv.
Pohliana, *Mart. Hist. Nat. Palm.* ii. 149. t. 6 A.—Bras.
Polteauana, *Kunth, Enum. Pl.* iii. 283.—Guiana.
Portesana, *H. Wendl. in Linnaea*, xxviii. (1856) 840.—Bras.
Princeps, *Linden, Illustr. Hortic.* xxviii. (1881) 31.—Peruv.
procumbens, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 106.—Costa Rica.
pulchella, *Hort. ex H. Wendl. Cycl. Pand. Cycad.* 28; *Linden, Illustr. Hortic.* xxviii. (1881) 31.—N. Granat.
pulchra, *Engel, in Linnaea*, xxviii. (1864–65) 686.—N. Granat.
pulchra, H. Wendl. ex Spruce, in Journ. Linn. Soc. xi. (1871) 105 = Pholidostachys pulchra.
pumila, *H. Wendl. in Linnaea*, xxviii. (1856) 331.—N. Granat.
Purdiaeana, *Spruce, in Journ. Linn. Soc.* xi. (1871) 109.—N. Granat.
pycnostachys, *Mart. Hist. Nat. Palm.* ii. 16. t. 17.—Am. trop.
ramosa, *Engel, in Linnaea*, xxxiii. (1864–65) 684.—N. Granat.
rectifolia, *Wallace, Palm Trees Amaz.* 67. t. xxv.—Bras.
Riedeliana, *H. Wendl. in Kerch. Palm.* 245.—Bras.?
rodeiensis, *Barb.-Rodr. Palmiers*, 42.—Bras.
rubescens, *H. Wendl. ex Drude, in Mart. Fl. Bras.* iii. ii. 491.—Bras.
rupestris, *Barb.-Rodr. Palmiers*, 47.—Bras.
Sagu, Spruce, in Journ. Linn. Soc. xi. (1871) 109 = Vaga.
Schomburgkiana, *Spruce, i. c.* 111.—Am. trop.
Schottiana, *Mart. Hist. Nat. Palm.* ii. 143. t. 11 A.—Bras.
Seemanni, *Hort. ex Floral Mag.* viii. (1869) t. 428.—Am. centr.
simplicifrons, Herb. Kew. ex Spruce, in Journ. Linn. Soc. xi. (1871) 107 (Quid?).—Am. austr.
simplicifrons, *Willd. Sp. Pl.* iv. 594.—Venezuela; Bras
speciosa, *Barb.-Rodr. Enum. Palm. Nov.* 9.—Bras.
spicigera, C. Koch, Wochenschr (1858) 244 = Calyptrogyne spicigera.
Spixiana, *Mart. Hist. Nat. Palm.* ii. 15. t. 15, 16.—Am. austr.
Spruceana, *Trail, in Journ. Bot.* xiv. (1876) 328.—Bras.
stricta, Kunth, Enum. Pl. iii. 282 = pycnostachys.
Swartzii, *Griseb. Cat. Pl. Cub.* 222.—Cuba.
sysanthera, *Mart. Hist. Nat. Palm.* ii. 18. t. 18.—Peruv.
Tamandua, *Trail, in Journ. Bot.* xiv. (1876) 328.—Bras.
tapajotensis, Trail, l. c. 324 = acaulis.
tomentosa, *Barb.-Rodr. Palmiers*, 44.—Bras.
trifurcata, *Orrst. in Kjoeb. Vidensk. Meddel.* (1858) 94.—Nicaragua.
trigonostyla, *Barb.-Rodr. Palmiers*, 46.—Bras.
trijugata, Barb.-Rodr. Enum. Palm. Nov. 12 = paniculigera.
trinervis, *Drude & H. Wendl. in Mart. Fl. Bras.* iii. ii. 492.—Bras.
tuberculata, *Spruce, in Journ. Linn. Soc.* xi. (1871) 113.—Bras.
uliginosa, *Barb.-Rodr. Enum. Palm. Nov.* 11.—Bras.
undata, *Klotzsch, in Linnaea*, xx. (1847) 453.—N. Granat.
vaga, *Griseb. & H. Wendl. in Fl. Brit. W. Ind.* 517.—Ind. occ.
ventricosa, *Engel, in Linnaea*, xxxiii. (1864–65) 688.—N. Granat.
Verdugo, *Linden. Illustr. Hortic.* xxviii. (1881) 31.—N. Granat.
Verschaffeltii, Hort. Linden. ex H. Wendl. in Kerch. Palm. 246, nomen = Acanthophoenix crinita?
versiformis, *H. Wendl. ex Spruce, in Journ. Linn. Soc.* xi. (1871) 109.—Costa Rica.
Wallisii, *Hort. Linden. ex H. Wendl. in Kerch. Palm.* 246, nomen.—N. Granat.
Weddelliana, *H. Wendl. ex Drude, in Mart. Fl. Bras.* iii. ii. 494.—Bras.
Wendlandii, *Hort.—Cf. Gard. Chron.* (1878) t. 440.—Hab.?
Wildenowii, Klotzsch, in Bot. Zeit. iv. (1846) 113 = simplicifrons.
Wittigiana, *Glas. ex Drude, in Mart. Fl. Bras.* iii. ii. 496.—Bras.
xanthocoma, *Hort. Linden, ex H. Wendl. in Kerch. Palm.* 246.—Ecuador.

GEOPHILA, Berger, Fl. Basses-Pyren. ii. 184 (1808) = **Merendera**, Ram. (Liliac.).
pyrenaica, Berger, l. c. = M. Bulbocodium.

GEOPHILA, D. Don, Prod. Fl. Nep. 136 (1825). *RUBIACEAE*, Benth. & Hook. f. ii. 127.
Afzelii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 221.—Afr. trop.
cordata, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 393.—Porto Rico.
cordata, Miq. in Linnaea, xvii. (1843) 72 (= *Mapouria cordata*).—Bras.; Guian.
cordifolia, *Miq. Stirp. Surin. Sel.* 176.—Guian.
dichondraefolia, A. Cunn. in Ann. Nat. Hist. ii (1839) 208 = Nertera grandiflora, dichondraefolia.
diversifolia, DC. Prod. iv. 537 = reniformis.
Gerrardi, *Baker, in Journ. Linn. Soc.* xxi. (1885) 418.—Madag.
gracilis, *DC. Prod.* iv. 537.—Peruv.
hirsuta, *Benth. in Hook. Niger Fl.* 422.—Afr. trop.
hirta, *Korth. in Nederl. Kruidk. Arch.* ii. ii. (1851) 247.—Borneo.
Hostmanniana, Miq. Stirp. Surin. Sel. 176 = cordifolia.
hydrocotyloides, Zipp. ex Span. in Linnaea, xv. (1841) 319 = reniformis.
involucrata, *Schnizl. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 222.—Afr. trop.
lancistipula, *Hiern, l. c.* 221.—Afr. trop.
macropoda, *DC. Prod.* iv. 537.—Peruv.
obvallata, *F. Delr. in Kjoeb. Vidensk. Meddel.* (1854) 186.—Afr. trop.
reniformis, Cham. & Schlecht. in Linnaea, iv. (1829) 187 = Psychotria herbacea.
reniformis, *D. Don, Prod. Fl. Nep.* 136.—Cosmop. trop.
uniflora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 221.—Afr. trop.
uniflora, Span. in Linnaea, xv. (1841) 319 = reniformis.
violacea, *DC. Prod.* iv. 537.—Am. austr.
violaefolia, *DC. l. c.*—N Granat.: Zollingeriana, *Miq. Fl. Ind. Bat.* ii. 312.—Java!

GEORCHIS, Lindl. in Wall. Cat. n 7379 (1831); in Bot. Reg. sub t. 1618 (1833); et Gen. et Sp. Orch. 496 (1840) = **Goodyera**, R. Br. (Orchid.).
bifida, Reichb. f. in Bonplandia, v. (1857) 35 = Goodyera bifida.
bifora, Lindl. in Wall. Cat. n. 7379.—Reg. Himal.
calva, Lindl. in journ. Linn. Soc. i. (1857) 184.—Java.
cordata, Lindl. Gen. et Sp. Orch. 496 = Goodyera cordata.
foliosa, Lindl. l. c.—Ind. or.
rubicunda, Reichb. f. in Bonplandia, v. (1857) 35 = Goodyera rubicunda.
Schlechtendaliana, Reichb. f. l. c. 36 = Goodyera Schlechtendaliana.
subregularis, Reichb. f. in Linnaea, xli. (1877) 61.—N. Caled.
viridiflora, F. Muell. Fragm. viii. 29 = Goodyera viridiflora.
vittata, Lindl. in journ. Linn. Soc. i. (1857) 184.—Ind. or.

GEORGIA, Spreng. Anleit. ii. ii. 567 (1818) = **Dahlia**, Cav. (Compos.).
bipinnata, Spreng. Syst. iii. 611 = Cosmos bipinnatus.
variabilis, Spreng. l. c. 610 = D. variabilis.

GEORGINA, Willd. Sp. Pl. iii. 2134 (1803) = **Dahlia**, Cav. (Compos.).
astrantiaefolia, Sweet, Hort. Brit. ed. II. 310 = D. astrantiaefolia.
Cervantesii, Sweet, Brit. Flow. Gard. Ser. II. i. t. 22 = D. coccinea.
coccinea, Willd. Sp. Pl. iii. 2134 = D. coccinea.
crocata, Sweet, Brit. Flow. Gard. t. 382 = D. coccinea.
frustranea, DC. in Ann. Mus. Par. xv. (1810) 310 = D. coccinea.
purpurea, Willd. Sp. Pl. iii. 2134 = D. variabilia.
rosea, Willd. l. c. = D. variabilis.
scaphigera, A. Dietr. in Otto & Dietr. Allg. Gartenz. i. (1833) 197 = D. scapigera.
superflua, DC. in Ann. Mus. Par. xv. (1810) 310 = D. variabilis.
variabilis, Willd. Hort. Berol. i. 28 = D. variabilis.

GERACIUM, Reichb. in Moesl. Handb. ed. II. ii. 1367 (1828) = **Crepis**, Vaill. (Compos.).
aureum, Reichb. Fl. Germ. Excurs. 259 = C. aurea.
chondrilloides, Reichb. l. c. 260 = C. Jacquini.
crocticum, Reichb. in Moesl. Handb. ed. II. ii. 1368 = C. succisaefolia.
cydoniaefolium, Reichb. l. c. 1367 = Hieracium cydoniaefolium.
incarnatum, Reichb. l. c. 1367 = C. incarnata.
lapsanoides, Reichb. l. c. 1367, in nota = C. lampsanoides.
molle, Schnz, Enum. Pl. Transs. 277 = C. succisaefolia.
paludosum, Reichb. in Moesl. Handb. ed. II. ii. 1368 = C. paludosa.
parviflorum, Reichb. l. c. 1367 = C. incarnata.
praemorsum, Reichb. l. c. = C. praemorsa.
prenanthoides, Reichb. l. c. 1369 = Hieracium prenanthoides.
succisaefolium, Reichb. l. c. 1368 = C. succisaefolia.
tingitanum, Reichb. l. c. 1367, in nota = C. tingitana!
viscidulum, Griseb. & Schenk, in Wiegm. Archiv. (1852) i. 349 = C. viscidula.

GERAEA, Torr. & Gray, in Proc. Am. Acad. i. (1847) 48 = **Encelia**, Adans. (Compos.).
canescens, Torr. & Gray, l. c. 49 = E. eriocephala.

GERANION, St. Lag, in Ann. Soc. Bot. Lyon, vii. (1880) 126 = seq

GERANIUM, [Tourn.] Linn. Syst. ed. I (1735) *GERANIACEAE*, Benth. & Hook. f. i. 272
RAMPHOCARPUS, Neck. Elem. ii. 488 (1790).
ROBERTIUM, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1837) 99.
aboriivum, De Not. ex Cesati, in Bibl. Ital. xci. (1888) 249 = molle.
abrotanifolium, Linn. f. Suppl. 304 = Pelargonium abrotanifolium.
absinthoides, Poir. Encyc. Suppl. ii. 741 = Erodium absinthoides.
acaule, DC. Prod. i. 649, in syn. = Pelargonium longifolium.
acaule, Linn. Syst. ed. X. 1143 = Erodium romanum.
acaule, Thunb. Prod. Pl. Cap. 116 = Pelargonium acaule.
acaule, *Willd. ex H. B. & K. Nov. Gen. et Sp.* v. 231.—Ecuador.
acerifolium, Cav. Dias. 243 = Pelargonium acerifolium.
acetosum, Dum.-Cours. Bot. Cult. ed. II. v. 18 = Pelargonium scandens.
acetosum, Linn. Sp. Pl. 676 = Pelargonium acetosum.
Ackneriauum, *Schur, Enum. Pl. Transs.* 136.—Transylv. (cult.).
aconitifolium, Eichw. Skizze, 170 = pratense?
aconitifolium, Knaf, ex Nym. Consp. 137 = sylvaticum.
acontitifolium, L'Hérit. Geran. t. 40 = rivulare.
acrocarpum, *Ledeb. Fl. Ross.* i. 471.—Soongaria.
aculeatum, J. F. Gmel. Syst. 1024 = Pelargonium aculeatum.
aculeolatum, *Oliver, Fl. Trop. Afr.* i. 291.—Afr. trop.
acuminatum, Thunb. Prod. Pl. Cap. 116 = Pelargonium acuminatum.
adulterinum, Thunb. l. c. 115 = Pelargonium adulterinum.
asthiopicum, Lam. Encyc. ii. 662 = Erodium cicutarium.
affine, *Ledeb. Fl. Alt.* iii. 299.—Sibir. altaic.
affine, Poir. Encyc. Suppl. ii. 757 = Pelargonium spathulatum.
affine, Wight & Arn. Prod. 188 = nepalense.
africanum, Cav. Dias. 249 = Pelargonium odoratissimum.
agrarium, *Willd. ex H. B. & K. Nov. Gen. et Sp.* v. 231.—Peruv.
albanum, *Bieb. Fl. Taur. Cauc.* ii. 127.—Reg. Caucas.; Persia.
albicans, *St. Hil. Fl. Bras. Mer.* i. 108.—Bras.
albiflorum, Hook. Fl. Bor. Am. i. 116. t. 40 = Richardsoni.
albiflorum, *Ledeb. Fl. Alt.* iii. 280.—Sibiria.
album, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1837) 139 = molle.
alsoides, Burm. f. Spec. Geran. 32 = Pelargonium grossularioides.

GERANIUM:—
alchemilloides, Linn. Sp. Pl. 678 = Pelargonium alchemilloides.
alnifolium, Poir. Encyc. Suppl. ii. 758 = Pelargonium alnifolium.
alpestre, Schur, Enum. Pl. Transs. 136 = sylvaticum.
alpinum, Burm. f. Spec. Geran. 30 = Erodium alpinum.
alpinum, *Kit. ex Kanits, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 581.—Hungar.
altaicum, *Herb. ex G. Don, in Loud. Hort. Brit.* 276, nomen.—Sibir. altaic.
alternans, Poir. Encyc. Suppl. ii. 756 = Pelargonium alternans.
althæoides, Linn. Sp. Pl. 679 = Pelargonium althæoides.
amethystinum, Lcdeb. ex Nordm. in Bull. Sc. Acad. Pétersb. ii. (1837) 814 = ibericum.
ancep, Poir. Encyc. Suppl. ii. 748 = Pelargonium anceps.
anemonæfolium, *L'Hérit. Geran.* t. 36.—Ins. Fortunat.
anemonoides, Thunb. Prod. Pl. Cap. 112 = Monsonia filia.
angelina, *Nois. ex Steud. Nom.* ed. I. 364, nomen.—Hab.?
angulatum, *Curt. Bot Mag.* t. 203.—Thracia.
angulosum, Mill Gard. Dict. ed. VIII. n. 22 = Pelargonium acerifolium.
angustifolium, Gilib. Fl. Lituan. ii. 176 = dissectum.
angustifolium, Thunb. Prod. Pl. Cap 113 = Pelargonium angustifolium.
anthemifolium, Poir. Encyc. Suppl. ii. 742 = Erodium anthemidifolium.
aphanoides, Thunb. Prod. Pl. Cap. 113 = Pelargonium aphanoides.
apiifolium, Andr. Geran. ii. = Pelargonium apioides.
appendiculatum, Linn f. Suppl 304 = Pelargonium appendiculatum.
apricum, *Phil. in Linnæa*, xxviii. (1856) 676.—Chili.
arabicum, *Forsk. Fl. Ægypt. Arab.* 124.—Arab.
arachnoideum, *St. Hil. Fl. Bras. Mer.* i. 102.—Bras.
arborescens, Desf. Fl. Atlant. ii. 110 = Erodium arborescens.
arboreum, *A. Gray, Bot. U. St. Expl. Exped.* i. 315. t. 31.—Ins. Sandvic.
ardens, Andr. Geran. i. = Pelargonium ardens.
arlescens, *Andr. i. c.* ii.—Hab.?
arjunense, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie*, ii. (1845) 585.—Java.
arduinum, Linn. Sp. Pl. ed. II. 952 = Erodium arduinum.
arenarium, Burm. f. Spec. Geran. 48 = Pelargonium arenarium.
arenicolum, *Steud. in Flora*, xxxix. (1856) 439.—Reg. Magellan.
argenteum, Bory & Chaub. Nouv. Fl. Pélop. 44 = subcaulescens.
argenteum, Genera. Elench. Scepus. n. 643; ex Wahlenb. Fl. Carpat. 308 = sylvaticum.
argenteum, *Linn. Cent.* ii. 25; *Amœn. Acad.* iv. 324. —Europ.
argentinum, Lucé. Fl. Osil. 286 = molle.
armenum, *Boiss. Fl. Orient.* i. 878.—Armenia.
Arnottianum, Steud. Nom. ed. II. i. 677 = nepalense.
articulatum, Cav. Diss. 293 = Pelargonium articulatum.
asperifolium, Andr. Geran. i = Pelargonium asperifolium.
asperum, Poir. Encyc. Suppl. ii. 759 = Pelargonium quercifolium.
asphodeloides, Bory & Chaub. Nouv. Fl. Pélop. 44 = peloponnesiacum.
asphodeloides, *Burm. f. Spec. Geran.* 28.—Ital.; Græc.; Ax. Min.
asphodeloides, Sibth. & Sm. Fl. Græc. Prod. ii. 40 = subcaulescens.
asplenioides, Desf. Fl. Atlant. ii. 109 = Erodium asplenioides.
astragalifolium, Cav. Diss. 257 = Pelargonium astragalifolium.
atlanticum, *Boiss. Diagn.* Ser. I. i. 59.—Algeria.
atrum, Moench, Meth. 686 = carolinianum.
atrum, Poir. Encyc. Suppl. ii. 746 = Pelargonium atrum.
auriculatum, Poir. l. c. 745 = Pelargonium longifolium.
auritum, Linn. Sp. Pl. 679 = Pelargonium auritum.

GERANIUM:—
australe, Nees, in Lehm. Pl. Preiss. i. 162 = dissectum.
australe, Poir. Encyc. Suppl. ii. 754 = Pelargonium australe.
ayavacense, Steud. Nom. ed. II. i. 677 = ayavacense.
Backhousianum, Regel, in Act. Hort. Petrop. ii. (1873) 492 = Grevilleanum.
baicalense, Turcz. ex Bess. in Flora, xvii. (1834) i. Beibl. 9 = albiflorum.
balsameum, Poir. Encyc. Suppl. ii. 754 = Pelargonium balsameum.
barbatum, Andr. Bot. Rep. t. 303 = Pelargonium aristatum, setigerum.
barbatum, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. iv. 142 = davuricum.
Barringtonii, Poir. Encyc. Suppl. ii. 751 = Pelargonium Barringtonii.
batrachioides, Cav. Diss. 231 = sylvaticum.
batrachyoides, Hochst. & Steud. ex Steud. Nom. ed. II. i. 677 = Endressi.
Bauæarterianum, Schur, Enum. Pl. Transs. 138 = pumilum.
Besufortiæ, *Andr. Geran.* i.—Hab.?
Beaufortianum, Poir. Encyc. Suppl. ii. 748 = Pelargonium Beaufortianum.
Berterianum, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 45.—Chili.
betonicum, Burm. f. Spec. Geran. 32 = Pelargonium myrrhifolium.
betulinum, Linn. Sp. Pl 679 = Pelargonium betulinum.
bicolor, Jacq. Hort. Vindob. iii. 28 = Pelargonium bicolor.
bicolor, Royle, Illustr. Bot. Himal. 149 = ocellatum.
bifforum, *Cav. Diss.* 251.—Afr. austr.
bifolium, Burm. f. Spec. Geran. 52 = Pelargonium bifolium.
bifolium, Maxim. Prim. Fl. Amur. 467 = dahuricum.
bifolium, Patrin, ex DC. Prod. i. 642 = Pseudosibiricum.
Bigosa, *Nois. ex Steud. Nom.* ed. II. i. 677, nomen.—Hab.?
bipinnatum, Cav. Diss. 273 = Erodium cicutarium.
Blandfordium, Andr. Geran. ii. = Pelargonium Blandfordianum.
blattarium, Poir. Encyc. Suppl. ii. 748 = Pelargonium ovale.
bohemicum, Krock. Fl. Siles. ii. 11. 120 = divaricatum.
bohemicum, *Linn. Cent.* ii. 25; *Amœn. Acad.* iv. 323.—Bohemia.
Botrys, Cav. Diss. 218 = Erodium Botrys.
brasiliense, *Prog. in Mart. Fl. Bras.* xii. 11. 522.—Bras.
brevicaule, Hook. Journ. Bot. i. (1834) 252 = sessiliflorum.
brevipes, L'Hérit. ex DC. Prod. i. 639 = sessiliflorum.
Bricenum, Sweet, Hort. Brit. ed. I. 74, nomen = Robertianum?
Broadleyæ, Andr. Geran. i = Pelargonium Broadleyæ.
bruttium, *Gasp. in Rendic. Acc. Nap.* i. (1842) 49.—Calabria.
bubonifolium, Andr. Bot. Rep. t. 328 = Pelargonium bubonifolium.
bulbosum, *Opiz, ex Steud. Nom.* ed. II. i. 677, nomen.—Hab.?
caerulastrum, Schur, Enum. Pl. Transs. 136 = pratense.
caerulicorpuscrum, Gilib. Fl. Lituan. ii. 176 = sylvaticum.
caeruleum, Gilib. l. c. 174 = pratense.
caeruleum, Moench, Meth. 684 = bohemicum.
caeruleum, Patrin, ex DC. Prod. i. 642 = Pseudosibiricum.
cæspitosum, *Yamez, ex Torr. in Ann. Lyc. N. York*, ii. (1828) 178.—N. Mexic.
cæspitosum, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 815.—Peruv.
californ, *Eckl. & Zeyh. Enum.* 58.—Afr. austr.
campanile, Schnegin, in Pall. N. Nord. Beitr. vi. 119 = Pseudo-sibiricum.
canariense, Poir. Encyc. Suppl. ii. 752 = Pelargonium canariense.
canariense, *Rent.*—*Cf. Linnæa*, xxix. (1857–58) 733. —Teneriffa.
canescens, *L'Hérit. Geran.* t. 36.—Afr. austr.
canescens, *Wedd. Chlor. And.* ii. 286.—Peruv.
capillare, Cav. Diss. 289 = Pelargonium capillare.
capitatum, Linn. Sp. Pl. 678 = Pelargonium capitatum.
carnosum, Linn. Cent. ii. 20 = Pelargonium carnosum.
carolinianum, *Linn. Sp. Pl.* 683.—Am. bor.

GERANIUM:—
carolinum, Crantz, Inst. ii. 156 = præc.
cataractarum, *Cos. Notes Crit.* 99.—Hispan.
caucalifolium, Poir. Encyc. Suppl. ii. 760 = Pelargonium caucalifolium.
ceratophyllum, Poir. l. c. 756 = Pelargonium ceratophyllum.
chærophyllum, Cav. Diss. 226 = Erodium cicutarium.
chamædrifolium, Poir. Encyc. Suppl. ii. 747 = Pelargonium chamædrifolium.
chamædroides, Cav. Diss. 197 = Erodium chamædryoides.
Chelidonium, Houtt. Handleid. x. 8. t. 61. f. 1 = Pelargonium Chelidonium.
chilloense, *Willd. ex H. B. & K. Nov. Gen. et Sp.* v 231.—Ecuador.
chium, Burm. f. Spec. Geran. 32 = Erodium chium.
chorense, Royle, Illustr. Bot. Himal. 150 = ocellatum.
ciconium, Linn. Cent. i 21 = Erodium ciconium.
cicutarium, Linn. Sp. Pl. 680 = Erodium cicutarium.
ciliatum, Andr. Bot. Rep. t. 247 = Pelargonium concavifolium.
ciliatum, Cav. Diss. 234 = Pelargonium parnassicides.
ciliatum, Willd. ex Spreng. Syst. iii. 71 = cucullatum.
cinereum, Lapeyr. Hist. Abr. Pl. Pyr. 394 = cinereum.
cinereum, Boiss. & Reut. ex Willk. & Lange, Prod. Fl. Hisp. iii. 525 = subargenteum.
cinereum, *Cav. Diss.* 204. t. 89. f. 1.—Mont. Pyren.
circinatum, *Kit. in Linnæa*, xxiii. (1868) 570.—Hungar.
citriodorum, Cav. Ic. i. 6 = Pelargonium acerifolium.
coelestinum, Schur, Enum. Pl. Transs. 136 = pratense.
collinum, *Steph. ex Willd. Sp. Pl.* iii. 705.—Europ. or.; As. bor.
columbinum, *Linn. Sp. Pl.* 683.—Europ.; As. bor.
commutatum, *Steud. in Flora*, xxxix. (1856) 439.—Chili.
compar, R. Br. in Salt, Abyss. App. 65, nomen = simense.
concavifolium, Poir. Encyc. Suppl. ii. 752 = Pelargonium concavifolium.
condensatum, Poir. l. c. 757 = Pelargonium condensatum.
conduplicatum, Poir. l. c. 751 = Pelargonium conduplicatum.
consanguineum, Poir. l. c. 759 = Pelargonium consanguineum.
contortum, Eckl. & Zeyh. Enum. 59 = ornithopodum.
cordifolium, Andr. Geran. i. = Pelargonium conduplicatum.
cordifolium, Cav. Diss. 240 = Pelargonium cordatum.
Core, Kostel. Alig. Med. Pharm. Flora, v. 1900 = rotundifolium.
Core-Core, Steud. in Flora, xxxix. (1856) 438 = præc.
coriandrifolium, Linn. Sp. Pl. ed. II. 949 = Pelargonium myrrhifolium.
cornutifolium, Andr. Bot. Rep. t. 305 = Pelargonium coronillæfolium.
coronopifolium, Andr. l. c. t. 338 = Pelargonium coronopifolium.
corsicum, Poir. Encyc. Suppl. ii. 743 = Erodium corsicum.
cortusæfolium, Andr. Bot. Rep. t. 191 = Pelargonium cortusæfolium.
Cotyledonis, Linn. Mant. ii. 569 = Pelargonium Cotyledonis.
crassicaule, Poir. Encyc. Suppl. ii. 752 = Pelargonium crassicaule.
crassifolium, Cav. Diss. 219 = Erodium alnifolium.
crassifolium, Desf. Fl. Atlant. ii. 111 = Erodium hirtum.
crassifolium, Forsk. Fl. Ægypt. Arab. 130 = Erodium glaucophyllum.
crassipes, Hook. ex A. Gray, in Bot. U. St. Expl. Exped. i. 309 = sessiliflorum.
crassum, Poir. Encyc. Suppl. ii. 741 = Erodium crassifolium.
cratægifolium, Roth, Bot. Abh. 50. t. 9 = Pelargonium glutinosum.
cratægifolium, Thunb. Prod. Pl. Cap. 115 = Pelargonium cratægifolium.
crenatum, Andr. Geran. ii = Pelargonium crenatum.
crenatum, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 354.—Mexic.
crenophilum, Boiss. Diagn. Ser. I. viii. 117 = asphodeloides.

GERANIUM:—
crispum, Berg. Desc. Pl. Cap. 176 = Pelargonium crispum.
cristatum, Stev. in Mém. Soc. Nat. Mosc. iv.(1813) 50 = albanum.
cristmifolium, Poir. Encyc. Suppl. ii. 750 = Pelargonium crithmifolium.
cucullatum, *H. B. & K. Nov. Gen. et Sp.* v. 231.—Am. austr.
cucullatum, Linn. Sp. Pl. 677 = Pelargonium cucullatum.
cunestrum, *Hook. Ic. Pl.* t. 198.—Ins. Sandvic.
cynosbatifolium, Poir. Encyc. Suppl. ii. 758 = Pelargonium malvæfolium.
dahuricum, *DC. Prod.* i. 642.—Dahuria.
daucifolium, Merr. in Comm. Gotting. (1780) 18 = Pelargonium flavum.
daucifolium, Thunb. Prod. Pl. Cap. 116 = Pelargonium daucifolium.
declinatum, *Mœnch, Meth. Suppl.* 97.—Hab.?
decumbens, Balb. Cat. Hort. Taur. (1813) App. i. 11 = Pelargonium triste.
delicatulum, *Tenore & Guss. ex Tenore, Fl. Nap.* v. p. xii.—Italia.
delphinifolium, Poir. Encyc. Suppl. ii. 752 = Pelargonium delphinifolium.
dentatum, Andr. Geran. i. = Pelargonium dentatum.
denticulatum, Poir. Encyc. Suppl. ii. 755 = Pelargonium denticulatum.
dilatum, *H. B. & K. Nov. Gen. et Sp.* v. 230.—Peruv.
difusum, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1887) 121 = columbinum.
dipetalum, Poir. Encyc. Suppl. ii. 744 = Pelargonium dipetalum.
diphyllum, L'Hérit. ex DC. Prod. i. 646 = Erodium laciniatum.
dissectum, *Linn. Cent.* i. 21.—Europ.; As. bor.; Austral.
divaricatum, *Ehrh. Beitr.* vii. 164.—Europ.; As. bor. et occ.
divaricatum, Loisel. Fl. Gall. ed. II. ii. 91 = bohemicum.
divaricatum, Thunb. Prod. Pl. Cap. 115 = Pelargonium divaricatum.
diversifolium, Poir. Encyc. Suppl. ii. 750 = Pelargonium glaucum.
Donianum, Hook. f. & Thoms. in Hook. f. *Fl. Brit. Ind.* i. 431 = polyanthes.
Donianum, Sweet, Geran. iv. sub t. 388 = collinum.
dubium, Chaix, in Vill. Hist. Pl. Dauph. i. 327 = pusillum.
duplicatum, *Kit. in Linnæa*, xxxii. (1863) 567.—Hungar.
echinatum, Andr. Bot. Rep. t. 158 = Pelargonium saepefolium.
echinatum, Thunb. Prod. Pl. Cap. 114 = Pelargonium aculeatum.
elatum, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1887) 138 = pusillum.
elegans, Andr. Bot. Rep. t. 28 = Pelargonium elegans.
elegans, Dum.-Cours. Bot. Cult. ed. II. v. 9 = Pelargonium betulinum.
ellipticum, Thunb. Fl. Cap. ed. Schult. 595 = Pelargonium ellipticum.
elongatum, Cav. Diss. 283 = Pelargonium tabulare.
elongatum, Salisb. Prod. 310 = incanum.
emarginatum, Linn. f. Suppl. 306 = Monsonia ovata.
eminense, *Hils. & Bojer ex Hook. f. in Journ. Linn. Soc.* vi. (1862) 6; vii. (1864) 135.—Madag.
Endressii, *J. Gay, in Ann. Sc. Nat. Sér.* I. xxvi. (1832) 238.—Mont. Pyren.
ensatum, Thunb. Prod. Pl. Cap. 118 = Pelargonium ensatum.
erianthum, *DC. Prod.* i. 641.—Sibiria; Am. bor. occ.
erianthum, Lindl. in Bot. Reg. (1841) *Misc.* 64; (1842) t. 29 = incisum.
erlostemon, D. Don, Prod. Fl. Nep. 208 = Grevilleanum.
eriostemon, *Fisch. ex DC. Prod.* i. 641.—Dahuria.
eriostemum, Poir. Encyc. Suppl. ii. 759 = Pelargonium ovale.
expansum, Thunb. Prod. Pl. Cap. 119 = Pelargonium expansum.
exstipulatum, Cav. Diss. 253 = Pelargonium exstipulatum.
fœtidum, Ascr. = *foetidum*.
fallax, *Steud. in Flora*, xxxix. (1856) 439.—Peruv.
fasciculatum, *Presl. in Verh. Zool.-Bot. Ges. Wien*, vi. (1856) 492.—Servia.

GERANIUM:—
fastigiatum, Gliem. ex Bab. in Journ. Linn. Soc. xi. (1871) 306 = sylvaticum.
favosum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 71.—Abyssin.
ferulaceum, Burm. f. Spec. Geran. 52 = Pelargonium ferulaceum.
Ficaria, Poir. Encyc. Suppl. ii. 745 = Pelargonium Ficaria.
fimbratum, Burm. f. Fl. Cap. Prod. 18 = Pelargonium cucullatum.
flanfolium, Andr. Bot. Rep. t. 378 = Pelargonium fissifolium.
flavum, Burm. f. Fl. Cap. Prod. 19 = Pelargonium flavum.
flexuosum, E. Mey. in Drège, Zwei Pfl. Docum. 187 = ornithopodon.
flexuosum, Thunb. Prod. Pl. Cap. 116 = Pelargonium flexuosum.
floribundum, Andr. Bot. Rep. t. 420 = Pelargonium floribundum.
foetidum, Gilib, Fl. Lituan. iii. 178 = Robertianum.
formosum, Poir. Encyc. Suppl. ii. 748 = Pelargonium formosissimum.
fragile, Andr. Bot. Rep. t. 37 = Pelargonium triparitum.
fragrans, Poir. Encyc. Suppl. ii 751 = Pelargonium fragrans.
Fremontii, *Torr. ex A. Gray, Pl. Fendl.* 26.—Am. bor. occ.
Freyeri, Griseb. Spicil. Fl Rumel. i. 126 = nodosum.
frigidum, *Hochst. in A. Rich. Tent. Fl. Abyss.* i. 116. —Abyssin.
fruticosum, Cav. Diss. 263 = Pelargonium fruticosum.
fulgidum, Linn. Sp. Pl. 676 = Pelargonium fulgidum.
furcatum, Kit. in Linnæa, xxxii. (1863) 567 = duplicatum.
furcatum, Schur, Enum. Pl. Transs. 138 = dissectum.
fuscatum, Poir. Encyc. Suppl. ii. 753 = Pelargonium saniculæfolium.
fuscum, Linn. Mant. i. 97 = phæum.
gefolium, Desf. Fl. Atlant. ii. 108 = Erodium hymenodes.
gibbosum, Linn. Sp. Pl. 677 = Pelargonium gibbosum.
glaberrimum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 116. —As. Min.
glandulosum, Cav. Diss. 271 = Erodium macradenum.
glandulosum, Lehm. ex Eckl. & Zeyh. Enum. 58 = canescens.
glaucoides, Andr. Geran. i. = Pelargonium glaucum.
glaucophyllum, Linn. Sp. Pl. 679 = Erodium glaucophyllum.
glaucum, Burm. f. Spec. Geran. 47 = Pelargonium glaucum.
glaucum, Cav. Diss. 237 = Pelargonium ovale.
glomeratum, Andr. Geran. ii = Pelargonium australe.
glutinosum, Jacq. Coll. i. 85 = Pelargonium glutinosum.
gracile, *Engelm. in A. Gray, Pl. Fendl.* 27.—Mexic.
gracile, *Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb.* ii. (1837) 314.—Reg. Caucas.
gracile, Schrank, in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 308 = Schrenkianum.
gracilipes, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 113.—N. Granat.
grandiflorum, Andr. Bot. Rep. t. 19 = Pelargonium grandiflorum.
grandiflorum, Cav. Diss. 194 = Monsonia tenuifolia.
grandiflorum, Edgew. in Trans. Linn. Soc. xx. (1846) 42 = palustre?
grandiflorum, Gilib. Fl. Lituan. ii 174 = sanguineum.
grandiflorum, Gueldenst. Reise, i. 426 = ibericum.
grandiflorum, Linn. Sp. Pl. 683 = Grielum tenuifolium.
gratissimum, *Nees. ex Steud. Nom.* ed. II. i. 678, nomen.—Hab.?
gratum, Poir. Encyc. Suppl. ii. 758 = Pelargonium gratum.
gravcolens, Stokes, Bot. Mat. Med. ii. 496 = Robertianum.
Grevilleæ, Andr. Geran. i, cum ic. = Pelargonium conspicuum.
Grevilleanum, *Wall. Pl. As. Rar.* iii. 4.—Reg. Himal.
grossularioides, Linn. Sp. Pl. 679 = Pelargonium grossularioides.
gruinum, Linn. l. c. 680 = Erodium gruinum.
guttatum, Desf. Fl. Atlant. ii. 113 = Erodium oxyrhinchum.
gymnocaulon, DC. Prod. i. 641 = ibericum.

GERANIUM:—
haemanthum, Willd. ex Spreng. Syst. iii. 74 = asphodeloides.
hederæfolium, Dum.-Cours. Bot. Cult. ed. II. v. 22 = Pelargonium saniculæfolium.
hedernum, Andr. Geran. i = Pelargonium lateripes.
heliotropioides, Cav. Diss. 220 = Monsonia heliotropioides.
hermannicefolium, Berg. Desc. Pl. Cap. 177 = Pelargonium crispum.
hermannioides, *Burm. f. Fl. Cap. Prod.* 19.—Afr. austr.
Hernandesii, *Moc. & Sesse, ex DC. Prod.* i. 640.—Mexic.
heterogynum, Poir. Encyc. Suppl. ii. 747 = Pelargonium heterogynum.
heterophyllum, Andr. Geran. ii = Pelargonium Andrewsii.
heterophyllum, Poir. Encyc. Suppl. ii. 746 = Pelargonium heterophyllum.
heterophyllum, Thunb. Prod. Pl. Cap 113 = Pelargonium variifolium.
heterotrichon, *Sm. ex Royle, Illustr. Bot. Himal.* 150, nomen.—Reg. Himal.
himalayense, Klotzsch, in Bot. Ergeb. Waldem. Reise, 122 = pratense
hirsutum, Burm. f. Spec. Geran. 50 = Pelargonium atrum.
hirtum, Burm. f. l. c. 48 = Pelargonium tenuifolium.
hirtum, Forsk. Fl. Aegypt. Arab. 123 = Erodium hirtum.
hirtum, Willd. ex Spreng. Syst. iii. 72 = He nandesii, mexicanum
hispidum, Linn. f. Suppl. 304 = Pelargonium hispidum.
holosericeum, *Willd. ex Spreng. Syst.* iii. 72.—Am. austr.
Homeyanum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi (1863) i. 591 = dissectum.
Hookerianum, Walp. Rep. i. 450 = Richardsoni.
Humboldtii, *Spreng Syst.* iii. 70.—*Am. austr.*
humile, Cav. Diss. 302 = pusillum.
hybridum, Linn. Mant. i. 97 = Pelargonium hybridum.
hymenodes, Andr. Bot. Rep. t. 413 = Erodium hymenodes.
hypoleucum, Benth. Pl. Hartw. 166 = acaule.
ibericum, *Cav. Diss.* 209. t. 124. f. 1.—Reg. Caucas.
ignescens, Andr. Geran. i. = Pelargonium ignescens.
impar, Steud. Nom. ed. II. i. 678, sphalm. vide *compar*.
incanum, *Burm. f. Spec. Geran.* 28.—Afr. austr.
incanum, E. Mey. in Drège, Zwei Pfl. Docum. 187 = sericeum, *Harv.*
incarnatum, Linn. Syst. ed. X. 1142 = Erodium incarnatum.
incisum, Andr. Bot. Rep. t. 67 = Pelargonium incisum.
incisum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 20.—Am. bor. occ.
increassatum, Andr. Bot. Rep. t. 246 = Pelargonium incrassatum.
inodorum, G. Don, Gen. Syst. i. 721 = Robertianum.
inodorum, Poir. Encyc. Suppl. ii. 752 = Pelargonium australe.
inquinans, Linn. Sp. Pl. 676 = Pelargonium inquinans.
intermedium, *Colla, in Mem. Acc. Torin.* xxxvii (1834) 46.—Chili.
Irbyanum, *Andr. Geran.* i.—Hab.?
japonicum, *Franch. & Sav. Enum. Pl. Jap.* ii. 305.—Japon.
Kotschyi, *Boiss. Diagn.* Ser. I. vi. 30.—Persia.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* ii. 306.—Japon.
lacerum, *Andr. Geran.* ii.—Hab.?
lacerum, Poir. Encyc. Suppl. ii. 749 = Pelargonium myrrhifolium.
lacinistatum, Andr. Bot. Rep. tt. 131, 269 = Pelargonium longifolium, penniforme.
laciniatum, Cav. Diss. 228 = Erodium laciniatum.
lactum, Leleb. Fl. Alt. iii. 298 = dahuricum?
laevigatum, Burm. ex DC. Prod. i. 640 = anemonæfolium.
laevigatum, Linn. f. Suppl. 306 = Pelargonium laevigatum.
laevigatum, Royle, Illustr. Bot. Himal. 150 = lucidum.
Lambertî, Sweet, Geran. iv. t. 338 = Grevilleanum?
lanatum, Thunb. Prod. Pl. Cap. 114 = Pelargonium cordatum.

GERANIUM :—
lancastrinse, Mill. Gard. Dict. ed. VIII. n. 4 = sanguineum.
lanceolatum, Cav. Diss. 285 = Pelargonium glaucum.
lanuginosum, Jacq. Hort. Schoenb. ii. 8 = carolinianum.
lanuginosum, *Lam. Encyc.* ii. 656.—Numidia.
lasiopus, *Boiss. & Heldr. Diagn.* Ser. I. viii. 117.—As. Min.
lateripes, Poir. Encyc. Suppl. ii. 758 = Pelargonium lateripes.
latistipulatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 117.—Abyssin.
Lauginianum, Nimmo, in J. Grah. Cat. Pl. Bomb. Add. post Ind. = Monsonia senegalensis.
Lebelii, Bor. Fl. Cent. Fr. i. 324 = Robertianum.
leiocaulon, Ledeb. Fl. Ross. i. 470 = molle.
lenticulum, *Rafin. New Fl. Am.* ii. 84.—Am. bor.
leucanthum, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 177, nomen.—Rossia.
leucanthum, *Griseb. in Goett. Abh.* xix. (1874) 108.—Reg. Argent.
libanoticum, *Boiss. & Blanche, in Boiss. Diagn.* Ser. II. v. 78.—Syria.
libanoticum, Schenk, Pl. Sp. Aegypt. 39 = tuberosum.
Lindenianum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 418.—Venezuela.
Lindleyanum, Royle, Illustr. Bot. Himal. 150 = Robertianum.
linaare, Andr. Bot. Rep. t. 198 = Pelargonium lineare.
linariifolium, DC. Fl. Fr. vi. 629, in nota = tuberosum.
littoreum, Cav. Diss. 322 = Erodium maritimum.
littoreum, Poir. Encyc. Suppl. ii. 744 = Erodium littoreum.
lividum, L'Hérit. Geran. t. 39 = phaeum.
lobatum, Burm. f. Spec. Geran. 44 = Pelargonium atrum, carneum, lobatum.
Londrei, Fisch. ex Link, Enum. Hort. Berol. ii. 196 = collinum.
Londrii, Fisch. ex Schrank, in Syll. Ratisb. ii. (1838) 67 = praec.
longicaule, *Thunb. Prod. Pl. Cap.* 116.—Afr. austr.
longiflorum, Poir. Encyc. Suppl. ii. 745 = Pelargonium longiflorum.
longifolium, Burm. f. Spec. Geran. 50 = Pelargonium longifolium.
longipedatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = acq.
longipes, DC. in Mém. Soc. Phys. Genèv. i. (1822) 442 = collinum.
lucanum, Gasp. ex Nym. Consp. 138 = molle.
lucidum, *Linn. Sp. Pl.* 682.—Europ.; As. bor.
ludovicianum, *Rafin. Fl. Ludov.* 89.—Am. bor.
lugubre, Salisb. Prod. 309 = macrorrhizum.
lupinoides, *Burm. f. Spec. Geran.* 43.—Afr. austr.
lutetum, Andr. Bot. Rep. t. 423 = Pelargonium luteum.
macrorrhizum, *Linn. Sp. Pl.* 680.—Europ. or.
macrostylum, Boiss. Diagn. Ser. I. i. 56 = tuberosum.
maculatum, *Linn. Sp. Pl.* 681.—Am. bor.
magellanicum, *Hook. f. Fl. Antarct.* 251.—Reg. Magellan.
malacoides, Linn. Sp. Pl. 680 = Erodium malachoides.
malapoides, Desf. Fl. Atlant. ii. 113 = Erodium alnifolium.
malacrum, Wahlenb. Fl. Suec. 452 = rotundifolium.
malvaeflorum, *Boiss. & Reut. Pugill. Pl. Nov.* 27.—Hispan.; Marocc.
malvaefolium, Schleich. Cat. 25 ; ex DC. Fl. Fr. v. 650 = molle.
malvaefolium, Scop. Fl. Carn. ed. II. ii. 37 = pusillum.
marginatum, Cav. Diss. 320 = Pelargonium marginatum.
maritimum, Burm. f. Spec. Geran. 46 = Erodium maritimum.
maritimum, Ucria, Hort. Reg. Panorm. 387 = Erodium laciniatum.
maacatense, *Boiss. Diagn.* Ser. I. i. 56.—Arabia.
Maximowiczii, *Regel & Maack, in Regel, Tent. Fl. Ussur.* 59.—Sibir. or.
mediterraneum, Jord. Pugill. Pl. Nov. 40 = Robertianum.
melananthos, Poir. Encyc. Suppl. ii. 747 = Pelargonium atrum.
melananthum, Andr. Bot. Rep. t. 209 = Pelargonium atrum.
menthaefolium, Nocca, ex Steud. Nom. ed. I. 564, 602 = Pelargonium tomentosum.

GERANIUM :—
mexicanum, Hort. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. ix. 74 = nepalense.
mexicanum, *H. B. & K. Nov. Gen. et Sp.* v. 230.—Mexic.
microphyllum, *Hook. f. Fl. Antarct.* 8. t. 5.—N. Zel.
Minaae, Tin. Fl. Rar. Sicil. (1846) 25 = pyrenaicum.
miniatum, Andr. Geran. ii = Pelargonium hybridum.
minimum, Cav. Diss. 260 = Pelargonium minimum.
minimum, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1887) 122 = dissectum.
minutiflorum, Jord. Cat. Dijon (1848) ; ex ej. Pugill. Pl. Nov. 39 = Robertianum.
modestum, Jord. Gr. Jard. Bot. Grenoble (1849) 16 ; ex Walp. Ann. ii. 284 = Robertianum.
molle, Gaertn. Meyer & Scherb. Fl. Wett. ii. 489 = pyrenaicum.
molle, *Linn. Sp. Pl.* 682.—Europ. ; As. bor. ; Orient. *Monsonia*, Thunb. Prod. Pl. Cap. 119 = Monsonia pilosa.
monstrum, Poir. Encyc. Suppl ii. 759 = Pelargonium monstrum.
montanum, *Y. & Gmel. Syst.* 1016.—Sibir.
montanum, Koch. ex O. Mucil. in Nov. Act. Nat. Cur. iv. (1770) 209, nomen = phaeum.
moschatum, Burm. f. Spec. Geran. 29 = Erodium moschatum.
mosqueras, Gold. in Mém. Soc. Nat. Mosc. v. (1817) 153 = Robertianum.
multicaule, Poir. Encyc. Suppl. ii. 749 = Pelargonium multicaule.
multiceps, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 417.—Venezuela.
multifidum, D. Don, Prod. Fl. Nep. 207 = collinum.
multifidum, Jacq. ex Steud. Nom. ed. II. i. 678 = Pelargonium multicaule.
multifidum, Patrin, ex DC. Prod. i. 645 = Erodium Stephanianum.
multifidum, *Sweet, Geran.* iii. t. 245.—Afr. austr.
multiflorum, *A. Gray, Bot. U. St. Expl. Exped.* i. 311. —Ins. Sandvic.
multinervum, *Benth. Pl. Hartw.* 166.—Ecuador.
murcicum, Cav. Diss. 272 = Erodium chium.
myrrhifolium, Linn. Sp. Pl. 677 = Pelargonium myrrhifolium.
napelliifolium, Schur, Enum. Pl. Transs. 186 = pratense.
neapolitanum, Nym. Consp. Index, 949 = pratense.
nemorosum, Tenore, Prod. Fl. Nap. p. lxx = asphodeloides.
nepalense, *Sweet, Geran.* i. t. 12.—Ind. or. ; China.
nervifolium, Poir. Encyc. Suppl. ii. 747 = Pelargonium nervifolium.
nobile, Walp. Rep. i. 458, in syn. sphalm. = Pelargonium nobile.
nodosum, *Linn. Sp. Pl.* 681.—Europ. austr.
notatum, *Andr. Geran.* ii.—Hab. ?
notkium, Pour. Encyc. Suppl. ii. 758 = Pelargonium atrum.
nowum, Winterl. Ind. Hort. Pest. f. 2 ; ex Roth, Catalect. ii. 76 = divaricatum.
numidicum, Poir. Voy. Barb. ii. 201 = Erodium cicutarium.
obtusilobum, Poir. Encyc. Suppl. ii. 759 = Pelargonium obtusilobum.
ocellatum, *Jacquem. Voy. Bot.* 33. t. 38.—Reg. Himal.
Ochsenii, *Phil. in Linnaea*, xxviii. (1856) 676.—Chile.
odoratissimum, Linn. Sp. Pl. 679 = Pelargonium odoratissimum.
oderatum, Thunb. Prod. Pl. Cap. 114 = praec.
Oenotherae, Linn. f. Suppl. 305 = Pelargonium Oenotherae.
omphalodeum, Lange, Ind. Sem. Hort. Haun. (1865) '67 = trilophum.
omphalodeum, Lange, l. c. (1867) App. 8 = favosum.
Onasi, *French. & Sav. Enum. Pl. Jap.* ii. 203.—Japon.
Orcades, Pans. Elench. Crta-Gora, 17 = asphodeloides.
orientale, Mill. Gard. Dict. ed. VIII. n. 13 = asphodeloides.
ornithopodum, *Eckl. & Zeyh. Enum.* 89.—Afr. austr.
ovale, Burm. f. Fl. Cap. Prod. 19 ; Thunb. Prod. Fl. Cap. 113 = Pelargonium ovale.
ovatifolium, *A. Gray, Bot. U. St. Expl. Exped.* i. 314. —Ins. Sandvic.

GERANIUM :—
ovatum, Cav. Diss. 288 = Pelargonium ovale, carinatum.
ovatum, Jacq. Coll. ii. 855 = Pelargonium Cenotherae.
oxalidifolium, Andr. Bot. Rep. t. 800 = Pelargonium oxalidifolium.
oxaloides, Burm. f. Spec. Geran. 51 = Pelargonium oxaloides.
oxoniense, *Andr. Geran.* i.—Hab. ?
oxyrhinchum, Poir. Encyc. Suppl. ii. 743 = Erodium oxyrhinchum.
pallens, Andr. Geran. ii = Pelargonium stipulaceum.
pallens, Bieb. Fl. Taur. Cauc. ii. 138 = Asphodeloides.
pallidum, Poir. Encyc. Suppl. ii. 750 = Pelargonium pallidum.
pallidum, Royle, ex Edgew. & Hook. f. in Fl. Brit. Ind. i. 491 = nepalense.
pallidum, Salisb. Prod. 310 = columbinum.
pallidum, Sweet, ex Steud. Nom. ed. II. i. 679 = eriostemon.
palmatiscum, Dulac, Fl. Hautes-Pyr. 240 = Robertianum.
palmatum, Cav. Diss. 316 = anemonaefolium.
palmatum, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1887) 122 = dissectum.
palustre, *Linn. Cent.* ii. 25 ; *Amoen. Acad.* iv. 328.—Europ.; As. bor.
papilionaceum, Linn. Sp. Pl. 676 = Pelargonium papilionaceum.
parilium, Willd. ex H. B. & K. Nov. Gen. et Sp. v. 231 = aquavaccine.
parviflorum, Andr. Geran. ii = Pelargonium parviflorum.
parviflorum, Curt. Fl. Lond. fasc. vi. t. 46 = pusillum.
parviflorum, Willd. Enum. Hort. Berol. 716 = dissectum.
parvulum, Scop. Delic. Insub. i. 8 = Erodium chamaedryoides.
pastinacaefolium, Mill. Gard. Dict. ed. VIII. n. 37 = Pelargonium triste.
patagonicum, *Hook. f. Fl. Antarct.* 252 = dissectum.
patens, Royle, ex Edgew. & Hook. f. in Fl. Brit. Ind. i. 491 = nepalense.
patulum, Vill. Hist. Pl. Dauph. iii. 371 = phaeum.
pedatum, *Rafin. New Fl. Am.* ii. 84.—Am. bor.
pedunculare, Willd. ex Spreng. Syst. iii. 71 = potentillaefolium.
pedunculatum, *Royle, Illustr. Bot. Himal.* 150, nomen. —Reg. Himal.
peloponesiacum, *Boiss. Diagn.* Ser. II. i. 110.—Graecia.
peltatum, Linn. Sp. Pl. 678 = Pelargonium peltatum.
penicillatum, Poir. Encyc. Suppl. ii. 758 = Pelargonium betulinum.
peruviforme, Poir. l. c. 757 = Pelargonium peruviforme.
pertagynum, Engelm. in Wisliz. Tour N. Mexic. 90 = Richardsoni.
pentandrum, Gilib. Fl. Lituan. ii. 178 = Erodium cicutarium.
peraense, Heds. Fl. Angl. ed. I. 265 = pyrenaicum.
Perreymondi, *Shuttlew. ex Barb. in Bull. Soc. Dauph.* (1881) 324.—Europ.
petraeum, Gouan, Illustr. 48. t. 21. f. 1 = Erodium petraeum.
Petroselinum, L'Hérit. ex Webb & Berth. Phyt. Canar. i. 92 = Erodium cicutarium.
phaeum, *Linn. Sp. Pl.* 681.—Europ.
phaeum, Sebast. & Maur. Fl. Rom. Prod. 225 = reflexum.
phaeum, Siev. in N. Nord. Beitr. vii. 229 = eriostemon.
philonotum, DC. Prod. i. 639 = dissectum.
pictum, Andr. Bot. Rep. t. 168 = Pelargonium pulchellum.
pilosum, Cav. Diss. 278 ; Andr. Bot. Rep. t. 269 = Pelargonium pilosum.
pilosum, Forst. f. Prod. 91 = dissectum.
pilosum, ? Th.' ex Nym. Consp. 140 = Erodium cicutarium.
pimpinellaefolium, Moench, Meth. 382 = Erodium cicutarium.
pinguc, Thunb. Prod. Pl. Cap. 116 = Pelargonium pinguc.
pinnatifidum, Burm. f. Spec. Geran. 50 = Pelargonium carneum.
pinnatifidum, Moench, Meth. 382 = Erodium romanum.

GERANIUM:—
pinnatifidum, Murr. ex Steud. Nom. ed. II. i. 679 = Pelargonium quercifolium.
pinnatifidum, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1887) 126 = rotundifolium.
pinnatum, Cav. Diss. 257 = Pelargonium pinnatum.
pinnatum, Linn. Sp. Pl. 677 = Pelargonium astragalifolium.
platypetalum, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 58 = Ibericum.
plicatum, Thunb. Prod. Pl. Cap. 116 = Pelargonium plicatum.
polyanthes, *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 431.—Reg. Himal.
potentillaefolium, *DC. Prod.* i. 639.—Mexic.
potentilloides, Bonpl. ex Wedd. Chlor. And. ii. 284 = acaule.
potentilloides, Klotzsch, in Bot. Ergeb. Waldem. Reise, 123 = rotundifolium.
potentilloides, L'Hérit. ex DC. Prod. i. 639 = dissectum.
potentilloides, Willd. ex Spreng. Syst. iii. 70 = Humboldtii.
praecox, Cav. Diss. 272 = Erodium primulaceum.
praemorsum, Andr. Bot. Rep. t. 150 = Pelargonium quinatum.
pratense, *Linn. Sp. Pl.* 681.—Europ.; As. bor.
procumbens, Andr. Bot. Rep. t. 264 = Pelargonium procumbens.
proliferum, Burm. f. Spec. Geran. 51 = Pelargonium barbatum.
prolificum, Linn. Sp. Pl. ed. II. 949 = Pelargonium astragalifolium.
propinquum, Sallab. Prod. 310 = rotundifolium.
prostratum, Cav. Diss. 196 = sanguineum.
proximum, *Steud. Nom.* ed. II. i. 679.—Chili.
Pseudo-molle, *Pantel. Adnot. Fl. Hercegov.* 121.—Europ.
Pseudo-pustillum, Schur, in Oestr. Bot. Zeitschr. xviii. (1868) 317 = pusillum.
Pseudo-sibiricum, Maxim. Prim. Fl. Amur. 71 = dahuricum.
Pseudo-sibiricum, *F. Mayer, in Boehm. Absh.* (1786) 288.—Japon.
Pseudo-villosum, Schur, Enum. Pl. Transs. 921 = villosum.
pulsatemon, Ledeb. Fl. Ross. i. 465 = armenum.
pubescens, *Thunb. Prod. Pl. Cap.* 113.—Afr. austr.
pubescens, Willd. ex H. B. & K. Nov. Gen. et Sp. v. 281 = chilloënse.
pulchellum, Pohr. Encyc. Suppl. ii. 758 = Pelargonium pulchellum.
pulverulentum, Cav. Diss. 272 = Erodium Cavanillesii.
pulverulentum, Desf. Fl. Atlant. ii. 111 = Erodium laciniatum.
pumilum, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1887) 121 = pyrenaicum.
pumilum, Poir. Encyc. Suppl. ii. 751 = Pelargonium scandens.
punctatum, Andr. Bot. Rep. t. 60 = Pelargonium punctatum.
punctatum, Cav. ex Steud. Nom. ed. I. 865 = Erodium incarnatum.
pungens, Poir. Encyc. Suppl. ii. 759 = Pelargonium spinosum.
purpurascens, Poir, l. c. 757 = Pelargonium longifolium.
purpureo-caeruleum, Ledeb. Fl. Ross. i. 465, sphalm. = sylvaticum.
purpureum, Gilib. Fl. Lituan. iii. 175 = palustre.
purpureum, Vill. Fl. Delphin. 73; Hist. Pl. Dauph. i. 373; iii. 374 = Robertianum.
pusillum, *Burm. f. Spec. Geran.* 27.—Europ.; Reg. Caucas. et Himal.
Pylzowianum, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 453, 466.—China.
pyrenaicum, *Burm. f. Spec. Geran.* 27.—Europ.; As. Min.
quercifolium, Linn. f. Suppl. 306 = Pelargonium quercifolium.
quinatum, Poir. Encyc. Suppl. ii. 758 = Pelargonium quinatum.
quinquelobum, Lam. Encyc. ii. 671 = Pelargonium heterogamum.
quinquelobum, *Wedd. Chlor. And.* ii. 286.—Ecuador.
quinquevulnerum, Andr. Bot. Rep. t. 114 = Pelargonium quinquevulnerum.
radiatum, Andr. l. c. t. 222 = Pelargonium radiatum.
radicans, DC. Prod. i. 650 = napulense.

GERANIUM:—
radicatum, Bieb. Fl. Taur. Cauc. ii. 184 = tuberosum.
radicatum, Lapeyr. Hist. Abr. Pl. Pyr. 390, in syn. = Erodium macradenum.
Radula, Cav. Diss. 262 = Pelargonium Radula.
Radula, Roth, Bot. Abh. 51. t. 10 = Pelargonium asperifolium.
Reii, Lindl. Syn. Brit. Fl. 67 = lucidum.
ramossissimum, Cav. Diss. 260 = Pelargonium ramosissimum.
ranunculifolium, Schur, ex Nym. Consp. 187 = sylvaticum.
ranunculoides, *Burm. f. Spec. Geran.* 17.—Africa.
rapaceum, Linn. Syst. ed. X. 1141 = Pelargonium rapaceum.
rapulum, *St. Hil. & Naud. in Ann. Sc. Nat. Sér. II.* xviii. (1842) 35.—Brasa.
rectum, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) i. 459.—Scoongar.; Reg. Himal.
reflexum, Andr. Bot. Rep. t. 224 = Pelargonium reflexum.
reflexum, *Linn. Mant.* ii. 257.—Italia.
refractum, *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 428.—Reg. Himal.
refulgens, *Andr. Geran.* ii.—Hab.?
Reichardii, Murr. in Comm. Gotting. (1780) 11. t. 3 = Erodium chamaedryoides.
Reinii, *Franch. & Sav. Enum. Pl. Jap.* ii. 804.—Japon.
Renardi, *Trautv. in Trautv. Regel, Maxim. & Winkl. Decas. Pl. Nov.* 5.—Reg. Caucas.
reniforme, Andr. Bot. Rep. t. 108 = Pelargonium reniforme.
reticulatum, Nocca, Pl. Select. t. 5 = Pelargonium denticulatum.
retrorsum, L'Hérit. ex DC. Prod. i. 644 = dissectum.
revolutum, Andr. Bot. Rep. t. 354 = Pelargonium revolutum.
revolutum, Jacq. Coll. i. 84 = Pelargonium Radula.
ribifolium, Dum.-Cours. Bot. Cult. ed. II. v. 19 = Pelargonium zonale.
ribifolium, Poir. Encyc. Suppl. ii. 754 = Pelargonium fulgidum.
Richardsonii, *Fisch. & Trautv. Ind. Sem. Hort. Petrop.* iv. 37.—Am. bor. occ.
rigidum, Poir. Encyc. Suppl. ii. 755 = Pelargonium crispum.
rivale, Vill. Hist. Pl. Dauph. i. 40 = seq.
rivale, *Vill. Prosp.* 40, *Hist. Pl. Dauph.* iii. 372.—Europ.
Robertianum, *Linn. Sp. Pl.* 681.—Reg. bor. temp.
romanum, Burm. f. Spec. Geran. 30 = Erodium romanum.
Rosa, Hort. ex DC. Prod. i. 674 = Pelargonium capitatum.
roseo-caeruleum, Gilib. Fl. Lituan. ii. 176 = columbinum.
roseum, Andr. Bot. Rep. t. 173 = Pelargonium condensatum.
roseum, *L'Hérit. ex Steud. Nom.* ed. I. 365, nomen.—Hab.?
rotundifolium, Ledeb. Fl. Alt. iii. 282 = acrocarpum.
rotundifolium, *Linn. Sp. Pl.* 683.—Europ.; As. bor.
rotundifolium, Pollich, Hist. Pl. Palat. ii. 265 = pusillum.
rubellum, *Moench, Meth. Suppl.* 95.—Hab.?
rubens, *Andr. Geran.* ii.—Hab.?
rubens, *Thunb. Prod. Fl. Cap.* 113.—Afr. austr.
rubifolium, Lindl. Bot. Reg. (1840) t. 67 = pratense.
rugosum, Andr. Geran. ii = Pelargonium rugosum.
rupestre, Pourr. in Mém. Acad. Toul. iii. (1788) 319 = Erodium supracanum.
ruplculum, *Wedd. Chlor. And.* ii. 285.—Bolivia.
ruthenicum, Ucechritz, in Oestr. Bot. Zeitschr. xxiii. (1873) 335 = sibiricum.
rutilans, Ehrh. Beitr. vii. 164 = anemonaefolium.
sanguineum, *Linn. Sp. Pl.* 682.—Europ.; Reg. Caucas.
saxatile, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 177.—Sibir. altaic.
scabrum, Linn. Amoen. Acad. iv. 261 = Pelargonium scabrum.
scandens, Poir. Encyc. Suppl. ii. 750 = Pelargonium scandens.
Schledeanum, *Schlecht. in Linnaea*, x. (1836) 258.—Mexic.
Schrenkianum, *Trautv. ex Becher, in Bull. Soc. Nat. Mosc.* lvii. (1882) i. 98.—Scoongar.
scopulicolum, Jord. ex Nym. Consp. 188 = Robertianum.

GERANIUM:—
Seemanni, *Peyr. in Linnaea*, xxx. (1859–60) 66.—Mexic.
selinum, Andr. Bot. Rep. t. 239 = Pelargonium rapaceum.
semiglabrum, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 180 = Robertianum.
semiorbiculare, Hort. ex Steud. Nom. ed. II. i. 679 = rotundifolium.
semitrilobum, Poir. Encyc. Suppl. ii. 755 = Pelargonium semitrilobum.
senecioides, Poir. l. c. 749 = Pelargonium senecioides.
sericeum, *Harv. in Harv. & Sond. Fl. Cap.* i. 267.—Afr. austr.
sericeum, *Willd. ex Spreng. Syst.* iii. 70.—Peruv.
serratum, Thunb. Prod. Pl. Cap. 113 = Pelargonium serratum.
sessiliflorum, *Cav. Diss.* 198. t. 77. f. 2.—Austral.; Chili.
setosum, Thunb. Fl. Cap. ed. Schult. 817 = Pelargonium setiferum.
sibbaldioides, Benth. Pl. Hartw. 166 = cuculatum.
sibiricum, *Linn. Sp. Pl.* 683.—Europ. or.; As. bor.
sidaefolium, Thunb. Prod. Pl. Cap. 114 = Pelargonium sidoides.
Sieboldi, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 452, 458.—Japon.; Manchuria.
simense, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 116.—Abyssin.
simile, Jord. ex Nym. Consp. 188 = Robertianum.
Sonderianum, C. Muell. in Walp. Ann. vii. 485, sphalm. = Lindenianum.
spathulatum, Andr. Bot. Rep. t. 152 = Pelargonium spathulatum.
speciosum, Steud. Nom. ed. II. i. 679, sphalm. = spinosum.
speciosum, Thunb. Prod. Pl. Cap. 112 = Monsonia speciosa.
spinosum, Burm. f. Spec. Geran. 16 = Sarcocaulon Burmanni.
splendens, Poir. Encyc. Suppl. ii. 759 = Pelargonium ignescens.
spurium, Poir. l. c. 758 = Pelargonium spurium.
stenopetalum, Andr. Geran. ii = Pelargonium leptopetalum.
stenopetalum, Poir. Encyc. Suppl. ii. 750 = Pelargonium zonale.
Stephanianum, Pohr. l. c. 741 = Erodium Stephanianum.
Steveni, Pohr. l. c. 745 = Erodium Steveni.
stipulaceum, Linn. f. Suppl. 306 = Pelargonium stipulaceum.
stipulare, Kuntze, in Flora, xxix. (1846) 698 = molle.
straminum, *Trinius & Ruprecht. in Ann. Sc. Nat. Sér. V.* xvii. (1873) 112.—N. Granat.
striatum, [Clairv.] Man. Herb. 158 = rivulare.
striatum, *Linn. Amoen. Acad.* iv. 283.—Europ. austr.
strictum, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1887) 124 = rotundifolium.
strigosum, *Burm. f. Fl. Cap. Prod.* 19.—Afr. austr.
suaveolens, Poir. Encyc. Suppl. ii. 747 = Pelargonium suaveolens.
subargenteum, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* iii. 525.—Hispan.
subcaulescens, *L'Hérit. ex DC. Prod.* i. 640.—Europ. or.; As. Min.
subcorrulcum, Schleich. Cat. 25; ex DC. Prod. i. 641 = phaeum.
subdivaricatum, Schur, in Verh. Naturf. Ver. Brunn, xv. 11. (1877) 160 = divaricatum.
submolle, *Steud. in Flora*, xxxix. (1856) 486.—Chili.
subnudicaule, *Purze. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 418.—Venezuela.
subrotundum, Rhrh. ex Hoffm. Deutschl. Fl. ed. II. i. 65 = rotundifolium.
sylvaticum, *Linn. Sp. Pl.* 681.—Europ.; As. bor.
sylvaticum, Saulcy, Cat. Pl. Palest. 5 = libanoticum.
sylvaticum, Urtis, Hort. Reg. Panorm. 227 = striatum.
sylvestre, Olaf. & Pov. ex Bab. in Journ. Linn. Soc. xi. (1871) 300 = sylvaticum?
tabulare, Burm. f. Spec. Geran. 37 = Pelargonium saniculaefolium.
tabulare, Linn. Sp. Pl. ed. II. 947 = Pelargonium tabulare.
tataricum, Pohr. Encyc. Suppl. ii. 741 = Erodium tataricum.
tauricum, *Rupr. Fl. Cauc.* 268, in textu.—Tauria.
tectum, Thunb. Prod. Pl. Cap. 116 = Pelargonium fulgidum.

GERANIUM :—
tenellum, Andr. Geran. ii = Pelargonium tenellum.
tenuifolium, Poir. Encyc. Suppl. ii. 757 = Pelargonium tenuifolium.
terebinthinaceum, Ait. ex Murr. in Comm. Gotting. vii. (1786) 88. t. 4 = Pelargonium quercifolium.
terebinthinaceum, Cav. Diss. 250 = Pelargonium graveolens.
ternatum, Linn. f. Suppl. 306 = Pelargonium ternatum.
tetragonum, Linn. f. i. c. 305 = Pelargonium tetragonum.
Thunbergii, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 28 = nepalense.
tomentosum, Andr. Bot. Rep. t. 115 = Pelargonium canum.
tomentosum, Thunb. Prod. Pl. Cap. 116 = Pelargonium tomentellum.
tordyloides, Desf. Fl. Atlant. ii. 107 = Erodium tordyloides.
tottum, Thunb. Prod. Pl. Cap. 115 = Pelargonium tottum.
Traversii, *Hook. f. Handb. N. Zeal. Fl.* 726.—N. Zel.
triangulare, Forsk. Fl. Aegypt. Arab. 123 = Erodium reflexum.
trichostomum, Poir. Encyc. Suppl. ii. 759 = Pelargonium ovale.
tricolor, Andr. Geran. i = Pelargonium tricolor.
tricuspidatum, *Andr. l. c.* ii.—Hab.?
tricuspidatum, Lam. Encyc. ii. 671 (Pelargonii sp.).—Afr. austr.?
trifidum, Burm. f. Spec. Geran. 52 = Pelargonium trifidum.
trifolium, *Cav. Diss.* 223.—Afr. bor.
trigonum, Scop. Delic. Insub. i. 12 = Pelargonium tetragonum.
trilobatum, Steud. Nom. ed. II. i. 680 = Erodium hymenodes.
trilobum, C. Koch, in Linnaea, xv. (1841) 716 = gracile, *Ledeb.*
trilobum, Thunb. Prod. Pl. Cap. 114 = Pelargonium trilobum.
trilophum, *Boiss. Diagn.* Ser. I. vi. 80.—Persia.
triphyllum, Poir. Encyc. Suppl. ii. 746 = Pelargonium triphyllum.
triste, Linn. Sp. Pl. 676 = Pelargonium triste.
Tuberaria, *Jacquem. Voy. Bot.* 38. t. 87.—Reg. Himal.
tuberosum, Boiss. Voy. Bot. Espagne, 119 = malvaeflorum.
tuberosum, *Linn. Sp. Pl.* 680.—Reg. Mediterr.; As. Min.; Persia.
Ulassovianum, Fisch. ex Reichb. Hort. Bot. i. 19, vide Wlassovianum.
umbrosum, Waldst. & Kit. Pl. Rar. Hung. ii. 181 = pyrenaicum.
undulatum, Andr. Bot. Rep. t. 292 = Pelargonium ovalifolium, undulatum.
undulatum, Andr. l. c. t. 317 = Pelargonium longifolium.
uniflorum, *Pacho, Voy.* t. 96.—Afr. bor.
valde-pilosum, Schur, ex Nym. Consp. 187 = pratense.
variabile, Moench, Meth. 288 = sibiricum.
variegatum, Linn. f. Suppl. 305 = Pelargonium variegatum.
variegatum, *Wender. Sem. Hort. Marb.* (1824); *ex Steud. Nom.* ed. II. i. 680.—Hab.?
varium, L'Hérit. Geran. t. 87 = cinereum.
velutinum, *Turcz. in Bull. Soc. Linn. Nat. Mosc.* xxxi. (1858) i. 417.—Venezuela.
venosum, Pers. Syn. ii. 335 = angulatum.
verbascifolium, Andr. Geran. ii = Pelargonium ovale.
verbascifolium, Steud. Nom. ed. II. i. 680 = praec.
vernicolor, Linn. Cent. Pl. i. 21 = striatum.
Villarsianum, Jord. in Cat. Gren. (1849); ex ej. Pugill. Pl. Nov. 38 = Robertianum.
villosum, *Andr. Geran.* ii.—Hab.?
villosum, Mill. Gard. Dict. ed. VIII. n. 38 = Pelargonium triste.
villosum, Tenore, Prod. Fl. Nap. p. lxi = molle.
villosum, Thunb. Prod. Pl. Cap. 116 = Pelargonium villiferum.
virgineum, Poir. Encyc. Suppl. ii. 757 = Pelargonium longifolium.
viscidulum, Fries, Novit. Fl. Suec. ed. II. 216 = rotundifolium.
viscosissimum, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. Suppl. 18 = incisum.
viscosum, Gilib. Fl. Lituan. ii. 177 = rotundifolium.

GERANIUM :—
viscosum, Mill Gard. Dict. ed. VIII. n. 20 = Erodium viscosum.
viscosum, Scop. Delic. Insub. ii. 27 = Pelargonium glutinosum.
vitifolium, Linn. Sp. Pl. 678 = Pelargonium vitifolium.
vulgare, *Guridenzi. Reise*, i. 56 (quid?).—Ross.
Wallichianum, *D. Don, in Sweet, Geran.* t. t. 90.—Reg. Himal.
Wilfordi, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 488.—Manchuria.
Winterli, Roth, Catalect. fasc. ii. 78 = divaricatum.
Wlassovianum, *Fisch. ex Link, Enum. Hort. Berol.* ii. 197.—Sibir.
yedoense, *Franch. & Sav. Enum. Pl. Jap.* ii. 805.—Japon.
zonale, Linn. Sp. Pl. 678 = Pelargonium zonale.

GERARDA, St. Lag. in Ann. Soc. Bot. Lyon, viii. 176 (1881) = seq.

GERARDIA, Linn. Gen. ed. I. 181 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 972.
AGALINIS, Rafin. New Fl. Am. ii. 61 (1836).
DASISTOMA, Rafin. in Journ. Phys. lxxiix. (1819) 98.
OTOPHYLLA, Benth. in DC. Prod. x. 512 (1846).
PAGESIA, Rafin. Fl. Ludov. 49 (1817), an huc referenda?
PANCTENIS, Rafin. New Fl. Am. ii. 60 (1836).
TOMANTHERA, Rafin. l. c. 65 (1836).
VERGULARIA, Reiz & Pav. Prod. 92. t. 19 (1794).
Afzelia, Michx. Fl. Bor. Am. ii. 20 = Seymeria tenuifolia.
angustifolia, *Mart. Nov. Gen. et Sp.* iii. 12. t. 206.—Bras.
aphylla, *Nutt. Gen. Am.* ii. 47.—Am. bor.
araucana, *Phil. in Anal. Univ. Chil.* (1884) reimpr. 8. —Chili.
aspera, *Dougl. ex Benth. in DC. Prod.* x. 517.—Am. bor.
auriculata, *Michx. Fl. Bor. Am.* ii. 48.—Am. bor.
brachyphylla, *Chem. & Schlecht. in Linnaea*, iii. (1828) 15.—Bras.
brasiliensis, Spreng. Syst. ii. 806 = Brunfelsia ramosissima.
caesarea, Cham. & Schlecht. in Linnaea, iii. (1828) 17 = Esterhazya splendida.
caviniifolia, Pers. Syn. ii. 154 = Seymeria tenuifolia.
commixta, *Cham. & Schlecht. in Linnaea*, iii. (1828) 12.—Bras.
Chyrze, *Steud. Nom.* ed. II. i. 680.—Hab.?
crinita, Eddy, ex Spreng. Syst. ii. 807 = maritima.
cuneifolia, Pursh, Fl. Am. Sept. ii. 422 = Herpestis nigrescens.
dasystacha, *Cham. & Schlecht. in Linnaea*, v. (1830) 104.—Mexic.
Dasystoma, Steud. Nom. ed. II. i. 680 = flava.
delphiniifolia, Linn. Cent. Pl. ii. 21 = Sopubia delphiniifolia.
densiflora, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 206.—Am. bor. occ.
Digitalis, *Benth. l. c.*—Bras.
digitata, *Spreng. Syst.* ii. 808.—Bras.
divaricata, *Chapm. Fl. S. U. St.* 299.—Florida.
domingensis, *Spreng. Syst.* ii. 807.—Ins. S. Doming.
Dregeana, Hochst. in Flora, xxv. (1842) 240 = Sopubia Dregeana.
erecta, Walt. Fl. Carol. 170 = tenuifolia.
fasciculata, Ell. Sketch, ii. 115 = purpurea.
filicaulis, *Chapm. Fl. S. U. St.* 299.—Am. bor.
filifolia, *Nutt. Gen. Am.* i. 48.—Am. bor.
filiformis, Rafin. ex Benth. in DC. Prod. x. 518 = tenuifolia.
filiformis, Schum. & Thonn. Beskr. Guin. Fl. 372 = Sopubia filiformis.
flava, Benth. Linn. ex A. Gray, Syn. Fl. N. Am. ii. i. 391 = quercifolia.
flava, *Linn. Sp. Pl.* 610.—Am. bor.
fruticosa, Pursh, Fl. Am. Sept. ii. 422. t. 18 = Pentstemon Menziesii.
genistifolia, *Cham. & Schlecht. in Linnaea*, iii. (1828) 15.—Bras.
glauca, Eddy, Cat. ex Spreng. Syst. ii. 807 = quercifolia.
glutinosa, Benge, Enum. Pl. Chin. Bor. 49 = Rehmannia chinensis.
glutinosa, Linn. Sp. Pl. 611 = Adenosma grandiflorum.

GERARDIA :—
guidioides, Cham. & Schlecht. in Linnaea, iii. (1828) 16 = Esterhazya splendida.
grandiflora, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 206.—Am. bor.
Greggii, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 181.—N. Mexic.
heterophylla, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 180.—Texas.
Heynеana, Benth. in Wall. Cat. n. 3891 = Sopubia delphiniifolia.
hispidula, *Mart. Nov. Gen. et Sp.* iii. 13. t. 207.—Bras.
indegrifolia, A. Gray, Man. Bot. N. U. St. ed. I. 307 = laevigata.
intermedia, Porter, ex A. Gray, Syn. Fl. N. Am. ii. i. 393 = purpurea.
japonica, *Thunb. Fl. Jap.* 251.—Japon.
laciniata, Mart. & Gal. in Bull. Acad. Brux. xii. 1 (1845) 26 = Seymeria virgata.
laevigata, *Rafin. Ann. Nat.* (1820) 18.—Am. bor.
lanceolata, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 207.—Peruv.
leucanthera, *Rafin. Fl. Ludov.* 50. (sp. dub.).—Am. bor.
linarioides, *Cham. & Schlecht. in Linnaea*, iii. (1828) 12.—Bras.
linifolia, Benth. in Hook. Comp. Bot. Mag. i. (1835) 209 = Pluknetii.
linifolia, *Nutt. Gen. Am.* ii. 47.—Am. bor.
longifolia, Benth. in Hook. Comp. Bot. Mag. i. (1835) 206 = aspera.
longifolia, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 180.—Am. bor. occ.
macrodonta, *Cham. & Schlecht. in Linnaea*, viii. (1833) 26.—Bras.
macrophylla, Benth. in Hook. Comp. Bot. Mag. i. (1835) 205 = Seymeria macrophylla.
maritima, *Rafin. in Med. Repos. N. York*, v. (1808) 361.—Am. bor.
Mettauеri, Wood, Class-Book (1861) 530 = divaricata
nigrina, Linn. f. Suppl. 278 = Melasma scabrum.
nigrum, *Thunb. Prod. Pl. Cap.* 106.—Afr. austr.
nuda, Wood, ex A. Gray, Syn. Fl. N. Am. ii. i. 395 = filicaulis.
obtusifolia, Benth. in Hook. Comp. Bot. Mag. i. (1835) 211 = Harveya obtusifolia.
orobanchoides, Lam. Encyc. ii. 690 = Alectra capensis.
parviflora, Benth. in Wall. Cat. n. 3888 = Leptorhabdos Benthamiana.
parviflora, Chapm. Fl. S. U. St. 300 = Skinneriana.
patula, *Chapm. ex A. Gray, Syn. Fl. N. Am.* ii. i. 391.—Am. bor.
pectinata, Torr. ex Benth. in Hook. Comp. Bot. Mag. i. (1835) 206 = G. Pedicularia.
Pedicularia, *Linn. Sp. Pl.* 611.—Am. bor.
pedunculosa, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 209.—Peruv.
Pluknetii, *Ell. Sketch*, ii. 114.—Am. centr.
prostrata, H. B. & K. Nov. Gen. et Sp. ii. 348 = Silvia prostrata.
purpurea, Bert. ex Mart. Nov. Gen. et Sp. ii. 15 = domingensis.
purpurea, *Linn. Sp. Pl.* 610.—Am. bor.
quercifolia, *Pursh, Fl. Am. Sept.* ii. 425. t. 19.—Am. bor.
ramosissima, Benth. in DC. Prod. x. 515 = linarioides.
rigida, *Gill. ex Benth. in Hook. Comp. Bot. Mag.* i. (1835) 206.—Chili.
rupestris, Rafin. Atl. Journ. 158 = Stenandrium rupestre.
scabra, Linn. f. Suppl. 279 = Graderia scabra.
scabra, Wall. Cat. n. 3389 = Sopubia trifida.
scabrosa, Rafin. Fl. Tellur. iv. 67 = Stenandrium scabrosum.
serpyllifolia, H. B. & K. Nov. Gen. et Sp. ii. 348 = Silvia serpyllifolia.
serrata, Torr. ex Benth. in DC. Prod. x. 521 = grandiflora.
sessilifora, *Vahl, Symb. Bot.* iii. 79.—Afr. austr.
setacea, *Walt. Fl. Carol.* 170.—Am. bor.
Skinneriana, *Wood, Class-Book* (1847); *ex A. Gray, Syn. Fl. N. Am.* ii. i. 394.—Am. bor.
Sopubia, Benth. in Hook. Comp. Bot. Mag. i. (1835) 210 = Sopubia trifida.
spiciflora, Engelm. in Bost. Journ. Nat. Hist. v. (1845) 287 = aspera.

GERARDIA:—
stemodiaefolia, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 289 = Alonaea incisifolia.
stricta, Benth. Scroph. Ind. 49 = Sopubia stricta, *G Don*.
strictiflora, *A. Gray, Syn. Fl. N. Am.* ii. 1. 294.—Am. bor occ.
strictiflora, *Benth. in Hook. Comp. Bot. Mag.* i. (1885) 209.—Texas.
tenuifolia, *Vahl, Symb. Bot.* iii. 79.—Am. bor.
tuberosa, Linn. Sp. Pl. 601 = Stenandrium rupestre.
tubulosa, Linn. f. Suppl. 279 = Rhamphicarpa tubulosa.
villosa, Muhl. ex Rafin. New Fl. Am. ii. 59 = flava.
virgata, Benth. in Hook. Comp. Bot. Mag. i. (1885) 205 = Seymeria decurva.
virgata, H. B. & K. Nov. Gen. et Sp. ii. 344 = Seymeria virgata.
Wallichiana, Stend. Nom. ed. II. i. 680 = Sopubia trifida.
Wrightii, *A. Gray, in Torr. Bot. Mex. Bound.* 118.—N. Mexic.

GERARDIANELLA, Klotzsch, in Peters, Reise Mossamb. Bot. 229 t. 36 (1861) = **Micrargeria**, Benth. (Scrophularin.).
scopiformis, Klotzsch, l. c.—Afr. trop.

GERASCANTHOS, Steud. Nom. ed. II. i. 681 (1840) = seq.

GERASCANTHUS, P. Br. Hist. Jamaic. 170. t. 29. f. 3 (1756) = **Cordia**, Linn. (Boraginⁿ.).
aspera, Mart. in Flora, xxi. (1838) ii. Beibl. 86 = C. insignis.
excelsa, Mart. l. c. = C. excelsa.
glabrata, Mart. l. c. 87 = C. glabrata.
scabra, Rafin. Sylva Tellur. 40 = C. scabra.
viviutina, Freseu. in Mart. Fl. Bras. viii. 5 = C. cujabensis.
vulgaris, Mart. in Flora, xxi. (1838) ii. Beibl. 86 = C. Gerascanthus.

GERASCHANTHUS, Lindl. Veg. Kingd. 629 (1847) = prec.

GERBERA, J. F. Gmel. Syst. 852 (1791) = GERBERIA, Scop. = **Quararibæa**, Aubl. (Malvac.).

GERBERA, Gronov. in Linn. Cor. Gen. 16 (1787). *COMPOSITAE*, Benth. & Hook. f. ii. 497.
ANANDRIA, Siegesb. ex Linn. Amoen. Acad. i. 243. t. 11 (1746).
APHYLLOCAULON, Lag. Amen. Nat. i. 85 (1811).
ATASITES, Neck. Elem. i. 7 (1790).
BERNIERA, DC. Prod. vii. 18 (1838).
CHAPTALIA, Royle, Illustr. Bot. Himal. t. 59. f. 2 (1839).
CLEISTANTHIUM, Kunze, in Bot. Zeit. ix. (1851) 850.
EPICLINANTHUM, Sch. ex DC. Prod. vii. 18 (1838).
IDICIUM, Neck. Elem. i. 28 (1790).
LASIOPUS, Cass. in Bull. Soc. Philom. (1817) 152.
LEIBNITZIA, Cass. in Dict. Sc. Nat. xxv. 429 (1822).
LEPTICA, E. Mey. ex DC. Prod. vii. 16 (1838).
OREOSERIS, DC. l. c. 17 (1838).
PERDICIUM, Linn. Fl. Afr. Rar. 22 (1762).
PARDIUM, Burm. f. Fl. Cap. Prod. 26 (1768).
abyssinica, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) i. *Intell.* 27, nomen.—Abyssin.
albicans, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia albicans.
amabilis, Hance, in Walp. Ann. ii. 947 = plinselloides.
ambigua, *Sch. Bip. in Flora*, xxvii. (1844) 780.—Afr. austr.
Anandria, *Sch. Bip. l. c.* 782.—Sibir. ; Japon.
aranzosa, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia aranzosa.
asplenifolia, *Spreng. Syst.* iii. 676.—Afr. austr.
aurantiaca, *Sch. Bip. in Flora*, xxvii. (1844) 781.—Afr. austr.
Bellidiastrum, Benth. in Journ. Linn. Soc. xiii. (1873) 348 = G. Anandria.
bicolor, *Sch. Bip. in Seem. Bot. Voy. Herald*, 313.—N. Granat.
Bojeri, *Sch. Bip. in Flora*, xxvii. (1844) 780.—Madag.
brasiensis, *Sch. Bip. in Seem. Bot. Voy. Herald*, 313.—Reg. Argent.
brevipes, *Sch. Bip. l. c.*—Bras.
Burmanni, *Cass. in Dict. Sc. Nat.* xviii. 461.—Afr. austr.

GERBERA:—
Candollei, *Sch. Bip. in Flora*, xxvii. (1844) 780.—Madag.
chilensis, *Sch. Bip. in Seem. Bot. Voy. Herald*, 313.—Chili.
cordata, *Less. in Linnaea*, v. (1830) 297.—Afr. austr.
crocea, Sch. Bip. in Flora, xxvii. (1844) 780 = ambigua.
coronopifolia, *Cass. in Dict. Sc. Nat.* xviii. 461.—Afr. austr.
crenata, [Ker-Gawl.] in Bot. Reg. t. 855 = Mairia crenata.
dentata, Sch. Bip. in Seem. Bot. Voy. Herald. 313 = Chaptalia dentata.
discolor, *Sond. in Harv. & Sond. Fl. Cap.* iii. 522.—Afr. austr.
Ehrenbergii, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia Ehrenbergii.
ferruginea, *DC. Prod.* vii. 15.—Afr. austr.
hirsuta, *Less. in Linnaea*, v. (1830) 298.—Arab.; China.
hirsuta, Spreng. ex DC. Prod. vii. 16 = tomentosa.
humilis, Sch. Bip. in Flora, xxvii. (1844) 779 = plinselloides.
hypochaeroidoides, *Baker, in Journ. Linn. Soc.* xx. (1883) 193.—Madag.
integralis, *Sond. in Harv. & Sond. Fl. Cap.* iii 520.—Afr. austr.
integrifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia integrifolia.
kokanica, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 58.—Kokan.
Kraussii, *Sch. Bip. in Flora*, xxvii. (1844) 781.—Afr. austr.
Kunzeana, *A. Br. & Aschers. in Cat. Sem. Hort. Berol.* (1871) *App.* 3.—Reg. Himal.
Lagascæ, *Cass. in Dict. Sc. Nat.* xviii. 462 (sp. dub.).—Hab. ?
lanuginosa, *Sch. Bip. in Flora*, xxvii. (1844) 780.—Reg. Himal.
leiocarpa, *Sch. Bip in Seem. Bot. Voy. Herald*, 313.—Cuba
leucochrix, *Harv. in Harv. & Sond. Fl. Cap.* iii. 521.—Afr. austr.
Linnaei, Cass. in Dict. Sc. Nat. xviii. 460 = asplenifolia.
lyrata, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia lyrata.
macrocephala, *Less. in Linnaea*, v. (1830) 295.—Afr. austr.
macrophylla, Benth. ex Hook. f. Fl. Brit. Ind. iii. 391 = nepalensis.
microcephala, Less. in Linnaea, v. (1830) 298 = tomentosa ?
natalensis, *Sch. Bip. in Flora*, xxvii. (1844) 778.—Afr. austr.
nepalensis, *Sch. Bip. l. c.* 780.—Reg. Himal.
nervosa, *Sond. in Linnaea*, xxiii. (1850) 70.—Afr. austr.
nivea, C. B. Clarke, Comp. Ind. 249 = Kunzeana.
nivea, *Sch. Bip. in Flora*, xxvii. (1844) 780.—Reg. Himal.
nutans, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia nutans.
oblonga, Sch. Bip. l. c. = Chaptalia oblonga.
ovalifolia, DC. Prod. vii. 17 = plinselloides.
ovalis, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia ovalis.
plinselloides, *Cass. in Dict. Sc. Nat.* xviii. 461.—Afr. austr.
plantaginea, *Harv. in Harv. & Sond. Fl. Cap.* iii. 522.—Afr. austr.
podophylla, *Baker, in Journ. Linn. Soc.* xviii. (1881) 278.—Madag.
pumila, *Sch. Bip. in Seem. Bot. Voy. Herald*, 313.—Ind. occ.
pusilla, Sch. Bip. in Flora, xxvii. (1844) 780 = lanuginosa.
rotundifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia rotundifolia.
Schimperi, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) i. Intell. 27 = plinselloides.
Seemannii, Sch. Bip. l. c. = Chaptalia sinuata.
Sellot, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia Seemannii.
semifoscularis, *Benth. & Hook. f. Gen.* ii. 497.—Afr. austr.
sinuata, Less. in Linnaea, v. (1830) 292 = ferruginea.
sinuata, Spreng. Syst. iii. 756 = Burmanni.

GERBERA:—
spathulata, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia spathulata.
tomentosa, *DC. Prod.* vii. 16.—Afr. austr.
viridiflora, Walp. Rep. vi. 315 = seq
viridifolia, *Sch. Bip. in Flora*, xxvii. (1844) 780.—Afr. austr.
Walteri, Sch. Bip. in Seem. Bot. Voy. Herald, 313 = Chaptalia tomentosa.
Wrightii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 521.—Afr. austr.

GERBERIA, Linn. Phil. Bot. 29 (1751) = **Gerbera**, Gronov. (Compos.).

GERBERIA, Scop. Introd. 286 (1777) = **Quararibæa**, Aubl. (Malvac.).
recursior, J. F. Gmel Syst. 692 = Q. guianensis.

GERBERIA, Stell. ex Choisy, in DC. Prod. xii. 24 (1848) = **Lagotis**, Gaertn. (Scrophul.).
Stelleri, Lam. ex Choisy, l. c. 25 = L. glauca.

GERDARIA, Presl, [Bot. Bemerk. 91 (1844)] in Abh. Boehm. Ges. V. iii. (1845) 521 = **Sopubia**, Buch.-Ham. (Scrophularin.).
Dregeana, Presl, ll. cc. = S. Dregeana.

GERMAINEA, Benth & Hook. f. Gen. iii. 1136 (1858) = seq.

GERMAINIA, Balansa & Poitrasa. in Bull. Soc. Hist. Nat. Toul. vii. (1873) 344 = **Anthistiria**, Linn. f. (Gramin.).
capillata, Balansa & Poitrasa. l. c.—Cochinch.

GERMANEA, Lam. Encyc. ii. 690 (1786) = **Plectranthus**, L'Hérit. (Labiat.).
crassifolia, Poir. Encyc. ii. 764 = Coleus spicatus ?
Forskohlei, Poir. l. c. = Coleus barbatus.
galeata, Poir. l. c. 763 = Coleus galeatus.
maculosa, Lam. Encyc. ii. 691 = P. punctatus.
mutisiflora, Poir. Encyc. ii. 763 = Coleus scutellarioides.
parviflora, Poir. l. c. 764 = P. parviflorus.
punctata, Poir. l. c. 763 = P. punctatus.
rotundifolia, Poir. l. c. = P. rotundifolius.
urticifolia, Lam. Encyc. ii. 692 = P. fruticosus.

GERMANIA, Hook. f. Fl Brit. Ind. iv. 625 (1885) = **Germanea**, Lam. = **Plectranthus**, L'Hérit. (Labiat.).

GERMARIA, Presl, Epim. Bot. 221 (1849) = **Pygeum**, Gaertn. (Rosac.).
latifolia, Presl, l. c. = P. latifolium.

GERONTOGEA, Cham. & Schlecht. in Linnaea, iv. (1829) 184 = **Oldenlandia**, Linn. (Rubiac.).
bifora, Cham. & Schlecht. l. c. 185 = O. corymbosa.
capensis, Cham. & Schlecht. l. c. 184 = O. capensis.
corymbosa, Cham. & Schlecht. l. c. = O. corymbosa.
*Drège*ana, Cham. & Schlecht. l. c. v. (1830) 169 = O. Deppeana.
fortida, Cham. & Schlecht. l. c. iv. (1829) 184 = O. foetida
herbacea, Cham. & Schlecht. l. c. = O. corymbosa.
lactea, Cham. & Schlecht. l. c. = O. lactea.
microthaca, Cham. & Schlecht. l. c. v. (1830) 169 = O. microthaca.
racemosa, Cham. & Schlecht. l. c. iv. (1829) 184 = O. paniculata.
umbellata, Cham. & Schlecht. l. c. = O. umbellata.

GEROPOGON, Linn. Sp. Pl. ed. II. 1109 (1763) = **Tragopogon**, Tourn. (Compos.).
australe, Spreng. Syst. iii. 663 = T. crocifolium.
calycibetum, Jacq. Hort. Schoenb. t. 108 = Scorzonera hirsuta.
glabrum, Linn. Sp. Pl. ed. II. 1109 = T. crocifolius.
hirsutum, All. ex Sm. in Rees, Cyclop. xvi ; DC. Prod. vii. 134 = Scorzonera hirsuta.
hirsutum, Georgi, Beschr. Russ. Reich. iv. 1202 = T. floccosus.
hirsutum, Linn. Sp. Pl. ed. II. 1109 = T. crocifolius.
hirsutum, Urv. in Mém. Soc. Linn. Par. i. (1822) 254 = T. porrifolius.
hybridum, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 472 = T. hybridum.

GEROSTEMUM, Stend. Nom. ed. II. i. 702 (1840) = GONOSTEMUM, Haw. = **Stapelia**, Linn. (Asclepiad.).
divaricatum, Stend. l. c. = S. divaricata.

GERRARDANTHUS, Harv. ex Benth. & Hook. Gen. i. 840 (1867). *CUCURBITACEAE*, Benth. & Hook. f. l. c.
macrorhiza, *Harv. ex Benth. & Hook. f. l. c.*—Afr. austr.
megarrhiza, Decne. & Harv. in Gen. S. Afr. Pl. ed. II. 127 = macrorhiza.
parviflora, *Cogn. in DC. Monog. Phan.* iii. 986.—Afr. trop.
portentosa, Naud. ex Dur. Cat. Sem. Hort. Bord. (1866) (1867) 26 = macrorhiza.
tomentosa, *Hook. f. Bot. Mag.* t. 6694.—Afr. austr.
Trimenii, *Cogn. in DC. Monog. Phan.* iii. 987.—Afr. trop.

GERRARDIANA, Oliver, in Hook. Ic. Pl. t. 1075 (1870). *SAMYDACEAE*.
foliosa, *Oliver, l. c.*—Afr. austr.

GERSINIA, Neraud, in Gaudich. Freyc. Voy. Bot. 27 (1826) = **Bulbophyllum**, Thou. (Orchid.).

GERUMA, Forsk. Fl. Aegypt. Arab. 62 (1775). *INC. SED.*—Cf. Benth. & Hook. f. i. 887.
alba, *Forsk. l. c.*—Arab.
subtriloba, Blanco, Fl. Filip. ed. I. 182 = Melochia corchorifolia.

GERYONIA, Schrank, ex Hoppe, in Flora, i. (1818) 229 = **Sastirago**, Linn.
cordifolia, Schrank, ex Steud. Nom. ed. II. i. 681 = S. cordifolia.
crassifolia, Schrank, ex Hoppe, in Flora, i. (1818) 229 = S. crassifolia.

GESNERA, Plum. ex Adans. Fam. ii. 157 (1763); Schreb. Gen. ii. 404 (1791) = **Gesneria**, Linn.

GESNERIA, Linn. Gen. ed. I. 179 (1737), partim; Mart. Nov. Gen. et Sp. iii. 27 (1829) (Gesnera) [Benth. & Hook. f. ii. 1008].
ALAGOPHYLLA, Rafin. Fl. Tellur. ii. 83 (1836).
CORYTHOLOMA, Decne. in Rev. Hortic. (1845) 466.
DICAEA, Decne. l. c. (1845).
ORCHANCHE, Vell. Fl. Flum. 254 (1825); vi. tt. 60–79 (1827), partim.
RECHSTEINERIA, Regel, in Flora, xxxi. (1848) 247.
acaulis, Hort. ex Hanst. in Mart. Fl. Bras. viii. 357 = tuberosa.
acaulis, Linn. Syst. ed. X. 1110 = Pentarhaphia Sloanei.
acaulis, Moç. & Sesse, in DC. Prod. viii. 11. 586 = Achimenes tenella.
adenantha, Fisch. ex Hanst. in Linnaea, xxix. (1857–58) 541 = Semeanula ternifolia.
affinis, Hort. Petrop. ex Hanst. in Mart. Fl. Bras. viii. 379 = bulbosa.
aggregata, [*Ker-Gawl.*] *in Bot. Reg.* t. 229.—Bras.
allagophylla, *Mart. Nov. Gen. et Sp.* iii. 36.—Bras.
atrosanguinea, Ann. Fl. (1836) 275, ex DC. Prod. vii. 529 = Lindleyi.
aurantiaca, *Hanst. Ind. Sem. Hort. Berol.* (1861) *App.* 8.—N. Granat.
barbata, Nees & Mart. in Nov. Act. Nat. Cur. xi. i. (1823) 48 = Ligeria barbata.
betonicaefolia, Hort. ex Hanst. in Mart. Fl. Bras. viii. 355 = allagophylla.
bibracteata, *D. Cameron, ex Loud. Hort. Brit. Suppl.* ii. 687, nomen.—Hab.?
Blasii, *Regel, in Schweiz. Zeit. Gaertn.* iv. (1846) 159.—Bras.
brevifolia, Lindl. in Journ. Hort. Soc. iii. (1848) 165 = Isoloma breviflorum.
bulbosa, Hook. Bot. Mag. t. 3886 = magnifica.
bulbosa, [*Ker-Gawl.*] *in Bot. Reg.* t. 243.—Bras.
calycina, Sieber, ex Steud. Nom. ed. II. i. 680, 200 = Besleria lutea.
calycina, Sw. Prod. Veg. Ind. Occ. 90 = Pentarhaphia calycina.

GESNERIA:—
canescens, *Mart. Nov. Gen. et Sp.* iii. 36.—Bras.
carassana, *Otto & Dietr. Allg. Gartens.* vi. (1838) 346.—Venezuela.
cardinalis, *Lehm. in E. Otto, Hamb. Gartens.* vi. (1850) 454.—Hab.?
chelonioides, *H. B. & K. Nov. Gen. et Sp.* ii. 292.—N. Granat.
cochlearis, Molina, Sagg. Chile, ed. II. 140 = Mitraria coccinea?
cinnabarina, Linden, in Belg. Hortic. vii. (1857) 197; Hook. Bot. Mag. t. 5086 = Nasgelia cinnabarina.
Clausseniana, *Brongn. ex Decne. in Rev. Hortic.* Sér. III. i. (1847) 365.—Bras.
coccinea, Hort. ex Hanst. in Mart. Fl. Bras. viii. 378 = rutila.
cochlearis, *Hook. Bot. Mag.* t. 3767.—Bras.
confertifolia, *Hanst. in Mart. Fl. Bras.* viii. 360.—Bras.
Cooperi, *Paxt. Mag. Bot.* i. (1835) 224.—Bras.
Cooperiana, Hort. Woburn. ex Lehm. in Otto, Hamb. Gartens. vi. (1850) 454 = prasc.?
cordata, *W. Bast. ex Loud. Hort. Brit. Suppl.* iii. 554, nomen.—Hab.?
corsicana, *Paxt. Mag. Bot.* xvi. (1849) 194.—Am. austr.
corymbosa, Balb. & Bert. ex DC. Prod. vii. 525 = Pentarhaphia exserta.
corymbosa, Sieber, ex Steud. Nom. ed. II. i. 681 = Episcia melittifolia.
corymbosa, Sw. *Prod. Veg. Ind. Occ.* 89.—Jamaic.
Cramsiana, Hort. Woburn. ex Pentarhaphia Cranioiaria.
crenulata, A. DC. in DC. Prod. vii. 524 = Rhytidophyllum crenulatum.
cuneifolia, Moç. & Sesse, in DC. l. c. 526 = Pentarhaphia cuneifolia.
cynocephala, *De Ven, ex Morr. Hortic. Belg.* iii. (1836) 320.—Bras.
dentata, *Hornsch. in Otto & Dietr. Allg. Gartens.* ii. (1834) 354.—Hab.?
Deppeana, Schlecht. & Cham. in Linnaea, v. (1830) 110 = Isoloma Deppeanum.
discolor, *Lindl. Bot. Reg.* (1841) t. 65. *Misc.* 45.—Bras.
Donkelaariana, Lem. Jard. Fleur. iv. (1854) t. 382 = seq.
Donklarii, *Hort. ex Hook. Bot. Mag.* t. 5070.—N. Granat.?
Douglasii, *Lindl. in Trans. Hort. Soc.* vii. (1828) 62.—Bras.
Douglasii, Mart. Nov. Gen. et Sp. iii. 35 = maculata.
egregia, Hort. ex Lem. Illustr. Hortic. iv. (1857) 129 = Heppiella naegelioides.
elatior, *H. B. & K. Nov. Gen. et Sp.* ii. 293.—Mexic.
elliptica, *Hook. Bot. Mag.* t. 4243.—N. Granat.
elongata, H. B. & K. Nov. Gen. et Sp. ii. 296 (= *Brachyloma elongatum*).—Ecuador.
elongata, Mart. & Gal. in Bull. Acad. Brux. ix. 11. (1842) 32 = Isoloma Deppeanum.
eriantha, Benth. Pl. Hartw. 298 = Isoloma erianthum.
erubescens, *Hanst. Ind. Sem. Hort. Berol.* (1861) *App.* 8.—N. Granat.
exgonia, [Moç. & Sesse,] ex DC. Prod. vii. 525 = Pentarhaphia pedunculosa.
exoniensis ×, Hort.—Cf. Gard. Chron. (1868) 1088; Nichols. Dict. Gard. ii. 66.
exserta, Sw. Prod. Veg. Ind. Occ. 90 = Pentarhaphia exserta.
fascialis, Hook. Bot. Mag. t. 3859 = seq.
fascialis, *Lindl. Bot. Reg.* t. 1785.—Bras.
fimbriata, Lam. Encyc. ii. 703 = Pentarhaphia Cranioiaria.
flaccuriifolia, Hort. ex Hanst. in Mart. Fl. Bras. viii. 355 = allagophylla.
flavescens, *Hanst. Ind. Sem. Hort. Berol.* (1861) *App.* 8.—Hab.?
fragilis, Poepp. & Endl. Nov. Gen. et Sp. iii. 7 = G. Scepturn.
Gardneri, Hook. Bot. Mag. t. 4121 = Hoottea leptopoda.
Gerardiana, Hort. ex Lem. in Fl. des Serres, Sér. I. ii. (Ann. 1846) t. 4 = Nasgelia Gerardiana.
Gerardiana, Kunth & Bouché, Ind. Sem. Hort. Berol. (1844) = Nasgelia Gerardiana.
Gollmeriana, *Hanst. l. c.* (1861) *App.* 7.—Venezuela.

GESNERIA:—
gracilis, Hanst. in Linnaea, xxxiv. (1865–66) 272 = punctata.
grandis, Hort. ex Hanst. in Mart. Fl. Bras. viii. 355 = allagophylla.
grandis, Sw. Prod. Veg. Ind. Occ. 89 = Rytidophyllum grande, Planchonianum.
guaranaefolia, *Benth. Pl. Hartw.* 147.—N. Granat.
guianensis, Benth. in Hook. Lond. Journ. Bot. v. (1846) 360 = Schomburgkiana.
Herbertiana, Lem. in Fl. des Serres, Sér. I. ii. (Ann 1846) t. 4 = Nasgelia Geroltiana.
hirsuta, Hort. Berol. ex Kunth, Ind. Sem. Hort. Berol. (1847) 13 = rubricaulis.
hirsuta, H. B. & K. Nov. Gen. et Sp. ii. 294 = Isoloma hirsutum.
hondensis, H. B. & K. l. c. 295 = Isoloma hondense.
hondensis, Morr. in Ann. Soc. Hort. Gand, ii. (1846) t. 84 = Pentarhaphia libanensis.
Hookeri, Decne. in Fl. des Serres, Sér. I. vii. (1851–52) 167 = maculata.
Hookeri, *Hanst. in Linnaea*, xxxiv. (1865–66) 276.—N. Granat.
Houttei, De Ven, ex Morr. in Hortic. Belg. iii. (1836) 320; Hanst. in Mart. Fl. Bras. viii. 374 = splendens.
Houttei, Dum. in Bull. Acad. Brux. iii. (1836) 362 = magnifica.
humilis, Linn. Sp. Pl. 619 = Pentarhaphia humilis.
ignorata, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 18 = Isoloma ignoratum.
inaequalis, Benth. Pl. Hartw. 230 = Isoloma inaequale.
incurva, Benth. Bot. Voy. Sulph. 131 = Isoloma incurvum.
laxiantha, Zucc. Pl. Nov. fasc. i. 300. n. 6 = Isoloma Deppeanum.
laterifia, *Lindl. Bot. Reg.* t. 1950.—Bras.
laterifia, Paxt. Mag. Bot. ix. (1842) 245 = cardinalis.
latifolia, *Mart. Nov. Gen. et Sp.* iii. 35.—Bras.
Leopoldi, *Schrädv. ex Planch. in Fl. des Serres*, Sér. I. vii. (1851–52) 167, 704.—Bras.
leptopus, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 129 = Hoottea leptopus.
libanensis, Morr. in Ann. Soc. Hort. Gand, ii. (1846) 361 = Pentarhaphia libanensis.
Lindeniana, Brongn. in Rev. Hortic. Sér. III. i. (1847) 368 = chelonioides.
Lindleyi, *Hook. Bot. Mag.* t. 3802.—Bras.
Linkiana, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 18 = Isoloma Linkianum.
lobulata, Hort. ex Hanst. in Mart. Fl. Bras. viii. 377 = macrorrhiza.
longiflora, H. B. & K. Nov. Gen. et Sp. ii. 296 = Isoloma longipes.
longiflora, [Moç. & Sesse] ex DC. Prod. vii. 536 = Achimenes longiflora.
longifolia, Lindl. Bot. Reg. (1841) Misc. 92 = Isoloma longifolium.
longipes, Benth. Pl. Hartw. 229 = Isoloma longipes.
longirostra, Mart. ex Hanst in Mart. Fl. Bras. viii. 380 = magnifica.
macrantha, Hort. Berol. ex Walp. Ann. ii. 1066 = cardinalis.
macrantha, *Sprengl. Syst.* ii. 820 (sp. dub.).—Bras.
macrorrhiza, *Dum. in Bull. Acad. Brux.* iii. (1836) 361.—Bras.
macrostachya, Lindl. Bot. Reg. t. 1202 = latifolia.
maculata, *Mart. Nov. Gen. et Sp.* iii. t. 215.—Bras.
maculata, *Moç. & Sesse, in DC. Prod.* vii. 532 (Quid?). —Mexic.
magnifica, *Otto & Dietr. Allg. Gartens.* i. (1833) 365.—Bras.
Marchi, *Walles, ex Hook. Bot. Mag.* t. 3744.—Bras.
melittifolia, *Brongn. in Rev. Hortic.* Sér. III. i. (1847) 368.—Mexic.
Merchii, Hanst. in Linnaea, xxxiv. (1865–66) 275 = Marchii.
Merckii, *Wendl. f. in Otto & Dietr. Allg. Gartens.* vi. (1838) 49.—Bras.
mollis, Hort. Berol. ex Kunth, Ind. Sem. Hort. Berol. (1848) 14 = Isoloma ignoratum.
mollis, H. B. & K. Nov. Gen. et Sp. ii. 295 = Isoloma mollis.
nitida, Hort. ex Hanst. in Mart. Fl. Bras. viii. 355 = allagophylla.
oblongata, Hort. Berol. ex Hanst. in Linnaea, xxxiv. (1865–66) 280 = Monmouthia Deppeana.

GESNERIA:—
oblongata, Hort. Young, ex Hook. Bot. Mag. sub t. 8735; Paxt. Mag. Bot. vi. (1839) 108 = elongata.
organa, Hort. ex Hanst. in Mart. Fl. Bras. viii. 394 = Houttea leptopus.
oxyphylla, *DC. Prod.* vii. 531.—Am. austr.
palustris, Hort. ex Hanst. in Mart. Fl. Bras. viii. 355 = allagophylla.
palustris, Hort. ex Regel, Gartenfl. iv. (1855) 247 = G. Scerptrum.
palustrienais, Hort. ex Hanst. in Mart. Fl. Bras. viii. 355 = allagophylla.
pardina, Hook. Bot. Mag. t. 4848 = Houttea pardina.
pendulina, *Lindl. Bot. Reg.* t. 1092.—Bras.
petiolaris, Benth. Bot. Voy. Sulph. 131 = Isoloma petiolare.
picta, Hook. Bot. Mag. t. 4481 = Isoloma pictum.
pilosa, Hort. ex Lem. in Fl. des Serres, Sér. I. iii. (1847) 223 = Columnea sarcostema.
polyantha, *DC. Prod.* vii 528.—Bras.
polyantha, Hook. Bot. Mag. t. 3995 = discolor.
prasinata, DC. Prod. vii. 532 = Paliavana prasinata.
prasinata, Ker-Gawl. in Bot. Reg. t. 428 = Rhytidophyllum prasinatum.
prasinata, Mart. Nov. Gen. et Sp. iii. 37 = Paliavana sericiflora.
pulchella, Hort. ex Hanst. in Mart. Fl. Bras. viii. 355 = allagophylla.
pulchella, Sw. Prod. Veg. Ind. Occ. 90 = Achimenes coccinea.
pumila, Sw. l. c. = Pentarhaphia pumila.
punctata, *Hort. ex Ann. Fl. et Pom.* (1837) 254.—N. Granat.†
purpurea, Hort. ex Lindl. & Paxt. Flow. Gard. ii. (1852–53) 17 = maculata, *Mart*
quadrifora, Lamb. ex Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 102 = Pentarhaphia parviflora.
quadrifolia, Warcz. ex Hanst. in Linnæa, xxix. (1857–58) 541 = Seemannia ternifora.
quaternifolia, Mog. ex DC. Prod. vii. 528 = Isoloma Deppeanum.
reflexa, *Knowl. & Westc. Fl. Cab.* ii. (1838) 65. t. 67.—Chili.
Regeliana, Warscz. ex Planch. in Fl. des Serres, Sér. I. ix. (1853–54) 213 = Sciadocalyx Warscewiczii.
rhynchocarpa, Benth. Bot. Voy. Sulph. 131 = Isoloma rhynchocarpum.
rubricaulis, Kunth & Bouché, Ind. Sem. Hort. Berol. (1847) 12 = Isoloma rubricaule.
rugata, *Scheid. in Otto & Dietr. Allg. Gartenz.* xv. (1847) 226 (sp. dub.).—Mexic.
rupestris, Mart. ex R. Grah. in Edinb. N. Phil. Journ. xxiv. (1838) 193 = tuberosa.
rupicola, *Mart. Nov. Gen. et Sp.* iii. 30. t. 218.—Bras.
rutila, *Lindl. Bot. Reg.* t. 1158.—Bras.
salviæfolia, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 139 = Houttea salviifolia.
Sartorii, *Liebm. Ind. Sem. Hort. Havn.* (1858) 20.—Mexic.
scabra, Sw. Prod. Veg. Ind. Occ. 89 = Pentarhaphia scabra.
sceptroides, *Hanst. in Ind. Sem. Hort Berol.* (1861) *App.* 7.—N. Granat.
Scerptrum, *Mart. Nov. Gen. et Sp.* iii. 32.—Bras.
Schiedeana, Hook. Bot. Mag. t. 4152 = Isoloma Schiedeanum.
Schomburgkiana, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1844).—Guiana.
Seemannii, Hook. Bot. Mag. t. 4504 = Isoloma Seemannii.
Selloi, Hanst. in Linnæa, xxxiv. (1865–66) 260 = Sellovii.
Selloi, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 897 = lateritia.
Sellovii, *Mart. Nov. Gen. et Sp.* iii. 36.—Bras.
silvatica, Anct. vide sylvatica.
spicata, Hort. ex Hanst. in Mart. Fl. Bras. viii. 355 = allagophylla.
spicata, Hort. ex Hanst. l. c. 371 = stricta.
spicata, *H. B. & K. Nov. Gen. et Sp.* ii. 393. t. 188 = Isoloma spicatum.
splendens, *Von Houtte, ex Hanst. in Mart. Fl. Bras.* viii. 374.—Bras.
splendidissima, Regel, Cat. Pl. Hort. Aksakov. 68 = prasc.
stachydifolia, *Benth. Pl. Hartw.* 280.—N. Granat.
stachydifolia, Hanst. in Linnæa, xxxiv. (1865–66) 264, *syphilis.* = prasc.
GESNERIA:—
stricta, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 280.—Bras.
Suttoni, *Booth, in Lindl. Bot. Reg.* t. 1627.—Bras.
sylvatica, *H. B. & K. Nov. Gen. et Sp.* ii. 393.—Bras.
tenella, Hort. ex Hanst. in Mart. Fl. Bras. viii. 366 = aggregata.
tenella, Moç. & Sesse, in DC. Prod. vii. 536 = Achimenes tenella.
tetraphylla, Hort. ex Hanst. in Mart. Fl. Bras. viii. 359 = Leopoldi.
tomentosa, Bert. ex DC. Prod. vii. 524 = Rhytidophyllum petiolare.
tomentosa, Jacq. Select. Am. 179. t. 178 = Rhytidophyllum crenulatum.
tomentosa, Linn. Sp. Pl. 612 = Rhytidophyllum tomentosum.
tomentosa, Moç. ex DC. Prod. vii. 528 = Isoloma Deppeanum.
Trianæi, Hérincq, L'Horticult. Franc. (1868) 30 = Isoloma Trianæi.
tribracteata, *Otto & Dietr. Allg. Gartenz.* ii. (1834) 194.—Bras.
trifora, Hook. Bot. Mag. t. 4342 = Isoloma trifolium.
trifora, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) reimpr. 2.—Mexic.
trifoliata, *Martens, in Linnæa*, xviii. (1844) 165.—Bras.
tuberosa, *Mart. Nov. Gen. et Sp.* iii. 29. t. 212.—Bras.
tubiflora, Cav Ic. vi. 61. t. 584 = Isoloma tubiflorum.
tubiflora, Endl. in Harting. Pæsel. Vindob. t. 94. t. 11 (= *Dolichodira tubiflora*).—Bras.
tubiflora, *Grisob. in Goett. Abh.* xxiv. (1879) 263.—Reg. Argent.
Tweediana, Hort. ex Hanst. in Mart. Fl. Bras. viii. 396 = Paliavana prasinata.
ulmifolia, H. B. & K. Nov. Gen. et Sp. ii. 394 = Heppiella ulmifolia.
umbellata, *Decne. in Fl. des Serres*, Sér. I. vii. (1851–52) 167.—Bras.
uniflora, *Moç. & Sesse, ex DC. Prod.* vii. 532 (sp. dub.).—Mexic.
Vargasii, *DC. Prod.* vii. 527.—Venezuela.
Vauthieri, *DC. l. c.* 530.—Bras.
velutina, Poepp. ex Hanst. in Linnæa, xxxiv. (1865–66) 290 = Moussonia purpurea.
velutina, Willd. ex DC. Prod. ix. 528 (= *Brachyloma elongatum*).—Ecuador.
ventricosa, Hort. Berol. ex Hanst. in Linnæa, xxix. (1857–58) 529 (= *Brachyloma ventricosum*).—Hab.?
ventricosa, Sw. Prod. Veg. Ind. Occ. 89 = Pentarhaphia longiflora.
Verdi, Hort. ex Hanst. in Mart. Fl. Bras. viii. 396 = Paliavana prasinata.
verticillata, *Cav. Ic.* vi. 62. t. 585.—Peruv.
verticillata, Hook. Bot. Mag. t. 2776 = maculata.
verticillata, Hort. ex Hanst in Mart. Fl. Bras. viii. 366 = aggregata.
vestita, Benth. in Lindl. Bot. Reg. (1845) Misc. 19 = Isoloma vestitum.
Warmingii, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 90.—Bras.
Warszewiczii, *Hanst. in Linnæa*, xxvi. (1853) 214.—Mexic.
xanthophylla, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 7.—Peruv.
zebrina, Paxt. Mag. Bot. viii. (1841) 96 = Naegelia zebrina.

GESNOUISIA, Gaudich. in Freyc. Voy. Bot. 502 (1826). *URTICACEÆ*, Benth. & Hook. f. iii. 392.
arborea, *Gaudich. l. c.*—Ins. Canar.
boehmerioides, Miq. in Mart. Fl. Bras. iv. 1. 194 = Phenax vulgaris.
filamentosa, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 207.—Ins. Canar.

GESNOUISIA, Steud. Nom. ed. II. i. 682 (1840), sphalm. = prasc.

GESSNERIA, Dum. Comm. Bot. 57 (1822) = Gesneria, Linn.

GESTROA, Becc. Malesia, i. (1877) 184. *VIOLARIÆ.*
candida, *Becc. l. c.*—Malaya.

GETHOSYNE, Salisb. Gen. Pl. Fragm. 72 (1866) = Asphodelus, Linn. (Liliac.).
acaulis, Salisb. l. c. = A. acaulis.

GETHYLLIS, Plum. ex Linn. Gen. ed. I. 181 (1737). *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 723.
ABAFUS, Adans. Fam. ii. 57 (1763).
PAPIRIA, Thunb. in Phys. Sallsk. Handl. i. (1776) 110.
acaulis, Blanco, Fl. Filip. ed. I. 260 = Hypoxis Franquevillei.
afra, *Linn. Sp. Pl.* 442.—Afr. austr.
Britteniana, *Baker, in Journ. Bot.* xxiii. (1885) 227.—Afr austr.
ciliaris, *Linn. f. Suppl.* 198.—Afr. austr.
lanceolata, Linn. f. l. c. = Apodolirion lanceolatum.
latifolia, *Masson, ex Baker, in Journ. Bot.* xxiii. (1885) 238.—Afr. austr.
longatyla, *Sobus, in Journ. Linn. Soc.* xviii. (1881) 206.—Afr. austr.
pilosa, Schum. & Thonn. Beskr. Guin. Pl. 172 = Curculigo gallabatensis.
plicata, Jacq. Hort. Schœnb. t. 42 = Curculigo plicata.
rosea, Eckl. Top. Verz. 4 = spiralis.
spiralis, Ait. Hort. Kew, ed. I. i. 437 = afra.
spiralis, *Linn. f. Suppl.* 198.—Afr. austr.
undulata, *Herb. Amaryll.* 186. t. 25. f. 5.—Afr. austr.
verticillata, *R. Br. Prod.* 290.—Afr. austr.
villosa, *Linn. f. Suppl.* 198.—Afr. austr.

GETHYRA, Salisb. in Trans. Hort. Soc. i. (1812) 282 = Reneslmia, Linn. (Scitamin.).
Cardamomum-medium, Sweet, Hort. Brit. ed. I. 390 = Amomum costatum.
linguiformis, Sweet, l. c. = Elettaria linguiformis.
occidentalis, Salisb. in Trans. Hort. Soc. i. (1812) 282 = Reneslmia jamaicensis.
punicea, Sweet, Hort. Brit. ed. I. 390 = Elettaria punicea.
tubulata, Sweet, l. c. = Alpinia tubulata.

GETHYUM, Phil. in Anal. Univ. Chil. (1873) 549. *LILIACEÆ*, Benth. & Hook. f. iii. 805.
atropurpureum, *Phil. l. c.*—Chili.

GETONIA, Banks & Soland. ex Benn. Pl. Jav. Rar. 122 (1838) = Cyrtandra, Forst. (Gesner.).

GETONIA, Roxb. Pl. Corom. i. 61. t. 87 (1795) = Calycopteris, Lam. (Combret.).
floribunda, Roxb. l. c. = C. floribunda.
nitida, Roth, Nov. Pl. Sp. 217 = C. floribunda.
nutans, Roxb. Hort. Beng. 33; Fl. Ind. ii. 428 = C. floribunda.

GETUONIS, Rafin. Fl. Tellur. ii. 20 (1836) = Allium, Tourn. (Liliac.).
vinealis, Rafin. l. c. = A. vineale.

GEUM, Linn. Syst. ed. I (1735). *ROSACEÆ*, Benth. & Hook. f. i. 619.
ADAMSIA, Fisch. ex Ser. in DC. Prod. ii. 558 (1825).
CARYOPHYLLATA, Tourn. ex Hall. Enum. Helv. i. 335 (1742).
SIEVERSIA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. 298 (1811).
STYLIPUS, Rafin. Neogenyt. 3 (1825).
adnatum, Wall. Cat. n. 712 = elatum.
agrimonioides, C. A. Mey. in Ind. Sem. Hort. Petrop. xi. Suppl. 29 = album?
agrimonioides, Pursh, Fl. Am. Sept. i. 351 = Potentilla arguta.
albiflorum, Scheutz, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 58, sphalm. = parviflorum.
album, *F. F. Genel. Syst.* 661.—Am. bor.
aleppicum, Jacq. Coll. i. 88 = strictum.
alpinum, Hornem. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 38 = brachypetalum.
alpinum, Mill. Gard. Dict. ed. VIII. n. 5 = montanum.
ambiguum, Schur, Enum. Pl. Transs. 183 = urbanum.
anemonoides, *Willd. Sp. Pl.* ii. 1117.—As. bor.; Am. arct.; Kamtschat.
antarcticum, Herb. ex Scheutz, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 30 = involucrosum.
atlanticum, Desf. Fl. Atlant. i. 403 = sylvaticum.
aurantiacum, *Fries, ex Scheutz, in Nov. Act. Soc. Sc. Upsal.* vii. ii. (1870) n. vi. 20.—Am. occ.†
Besseri, Fisch. ex Walp. Rep. ii. 47 = strictum.
Besserianum, Sweet, Hort. Brit. ed. II. 190 = strictum.

GEUM:—
bifolium, Brot. Fl. Lusit. ii. 353 = sylvaticum.
brachypetalum, *Ser. in Mém. Soc. Phys. Genève.* ii. (1824) 189.—Hab.?
bulgaricum, *Panč. Elem. Fl. Bulg.* 26.—Bulgaria.
calthifolium, *Sm. in Rees, Cycl.* v. n. 18.—Am. arct.
camtschaticum, Auct. vide *hamtschaticum*.
camtschaticum, Pall. ex Torr. & Gray, Fl. N. Am. i. 425 = calthifolium.
canadense, Jacq. Hort. Vindob. ii. 82 = album.
canadense, Murr. in Comm. Gotting. v. (1783) 88. t. 4 B = strictum.
capense, *Thunb. Prod. Pl. Cap.* 91.—Afr. austr.
carolinianum, Walt. Fl. Carol. 150 = album.
caryophyllata, Gilib. Fl. Lituan. ii. 238 = urbanum.
caryophyllatum, Pers. Syn. ii. 57 = urbanum.
caucasicum, Hort. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. Suppl. 81 = urbanum.
cercocarpoides, DC. Prod. ii. 554 = Fallugia paradoxa.
chamaedryfolium, Crantz, Stirp. Austr. ed. 1. fasc. ii. 7 = Dryas octopetala.
chiloense, *Balb. ex Ser. in DC. Prod.* ii. 551.—Chili.
cihatum, Pursh, Fl. Am. Sept. i. 352 = triflorum.
coccineum, Lindl. Bot. Reg. t. 1088 = chiloense.
coccineum, *Sibth. & Sm. Fl. Græc. Prod.* i. 354; *Fl. Græc.* t. 458.—Græcia; Ax. Min.
diversifolium, Hort. ex Schenk, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 29 = strictum.
dryadeides, DC. Prod. ii. 564 = Cowania mexicana.
Dryas, Crantz, Inst. ii. 175 = anemonoides.
dubium, Hornem. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 38 = *intermedium*.
elatum, *Wall. Cat.* n. 711.—Reg. Himal.
elegans, Poepp. ex Schenk, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 19 = chiloense.
Fischeri, Ikea ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 38 = strictum.
Francklf, *Steud. Nom.* ed. II. i. 682.—Am. bor.
geniculatum, *Michx. Fl. Bor.-Am.* i. 300.—Am. bor.
glaciale, *Fisch. in Mém. Soc. Nat. Mosc.* ii. (1809) 187. t. 11. f. 20.—Sibir. et Am. arct.
glutinosum, Pursh, ex Lehm. Pugill. iii. 25 = Potentilla argnta.
Grahami, Steud. Nom. ed. II. i. 682 = triflorum.
grandiflorum, C. Koch, in Linnæa, xix. (1847) 43 = coccineum.
grandiflorum, Steud. Nom. ed. II. i. 682 = chiloense.
haderaefolium, C. C. Gmel. Fl. Bad. ii. 460 = urbanum.
heboviticum, Schleich. ex Tratt. Rosac. Monog. iii. prænf. p. viii (Quid ?).—Europ.
heterocarpum, *Boiss. Voy. Bot. Espagne*, 201. t. 68 — Reg. Mediterr.
heterophyllum, Desf. Cat. Hort. Par. ed. III. 294 = virginianum.
heterophyllum, Hort. ex Link, Enum. Hort. Berol. ii. 65 = strictum.
hirsutum, Mahl. Cat. ed. I. (1818) 51 = virginianum.
hirtum, Wahlenb. Fl. Gothob. 60 = urbanum.
hispidum, Auct. Germ. ex Schenk, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 29 = strictum.
hispidum, *Fries, Fl. Halland.* 90.—Succ.; Hisp.
humile, Steud. Nom. ed. II. i. 682 = Rossii.
hybridum, Wulf. in Jacq. Misc. ii. 33 = rivale.
hyrcanum, *C. A. Mey. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 32.—Reg. Caucas.
ibericum, Besa. ex Schenk, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 26 = urbanum.
incisum, Dum. Fl. Belg. Prod. 97 = rivale?
inclinatum, *Schleich. Cat. Pl. Helv.* 46.—Helvetia.
inclinatum, Timb. ex Nym. Consp. 229 = sylvaticum.
intermedium, Ben. ex Bieb. Fl. Taur. Cauc. i. 411 = strictum.
intermedium ×, Ehrh. Beitr. vi. 148.—Europ.
intermedium, Tenore, ex Nym. Consp. 230 = molle.
intermedium, Willd. Hort. Berol. i. 69 = urbanum.
involucratum, *Yuss. ex Perz. Syn.* ii. 57.—Am. austr.
japonicum, Hort. ex Schenk, in Nov. Act. Soc. Sc. Upsal vii. ii. (1870) n. vi. 23 = strictum.
japonicum, *Thunb. Fl. Jap.* 230.—Japon.
hamtschaticum, Lam. ex Poir. Encyc. Suppl. i. 618 = anemonoides.
karstavicum, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 577.—Turkestan.
Kladitzianum, Peterm. Fl. Lips. Excurs. 282 = urbanum.
kokanikum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 34.—Turkestan.
lacinatum, Murr. in Comm. Gotting. v. (1783) 90. t. 2 = virginianum? *sed* Potentilla argnta?
laciniosum, Murr. in Comm. Gotting. v. (1783) 86 = album?
Lamanni, *Gaertn. Fruct.* i. 352. t. 74.—Sibir.
Lechlerianum, *Schlecht. in Linnæa*, xxviii. (1856) 465.—Chili.
lyratum, *Hort. Vind. ex Schenk, in Nov. Act. Soc. Sc. Upsal.* vii. ii. (1870) n. vi. 26. obs. 4.—Hab.?
macedonicum, Griseb. Relse Rumel. ii. 166, nomen = coccineum.
macrophyllum, Hook. Bot. Beech. Voy. 113 = strictum.
macrophyllum, *Wild. Enum. Hort. Berol.* 557.—Am. bor. occ.
magellanicum, Comm. ex Pers. Syn. ii. 57 = urbanum.
magellanicum, Lechler, ex Schenk, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 35 = parviflorum.
micropetalum, *Gasp. Notiz. Alc. Piante*, 11.—Ital.
molle, *Vis. & Panč. in Mem. Ist. Venet.* x. (1861) 429.—Serbia.
montanum, Gouan, ex Steud. Nom. ed. II. i. 682 = sylvaticum.
montanum, *Linn. Sp. Pl.* 501.—Europ. austr.
Mulleri, *Reych, ex Gal. in Journ. Hort. Prat. Sér.* II. i. 122. t. 12. f. 2.—Hab.?
mutans, Crantz, Stirp. Austr. ed. II. 70; Rafin. in Am. Monthly Mag. (1818) 344 = rivale.
nutans, *Hort. Par. ex Lam. Encyc.* i. 899.—Hab.?
obliquum, Steud. Nom. ed. I. 366; ed. II. i. 682 = Waldsteinia Moenchii.
ochroleucum, *Rafin. in Med. Repos. N. York*, v. (1808) 354, nom. n (Quid?).—Am. bor.
pallidum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 49.—? ieb.?
paradoxum, Steud. Nom. ed. II. i. 682 = Fallugia paradoxa.
parviflorum, *Sm. in Rees, Cycl.* v. n. 12.—Reg. Antarct.
parviflorum, Steud. & Hochst ex Steud. Nom. ed. II. i. 682 = Francklf.
Pechii, Pursh, Fl. Am. Sept. i. 352 = triflorum.
pictum, Hort. ex Schenk, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 38 = rivale.
pinnatifidum, *Rafin. in Med. Repos. N. York*, v. (1808) 354, nomen (Quid?).—Am. bor.
plumosum, Moç. & Sesse, ex D. Don, in Trans. Linn. Soc. xvi. (1833) 576 = Fallugia paradoxa.
ponticum, Pall. ex Lehm. Pugill. ix. 38 = Potentilla geoides.
Porteschlagianum, *Tratt. Ros.* iii. 116.—Europ.?
potentilloides, Ait. Hort. Kew. ed. I. ii. 219 = Laxmanni.
potentilloides, Pall. Ind. Taur. 52, nomen = Potentilla geoides.
Pseudo-molle, Pant. in Verh. Ver. Natur- u. Heilk. Presb. ii. (1871–72) 121 = molle.
pubescens, Hook. Fl. Bor. Am. i. 175 = triflorum.
pyrenaicum, Koch, Syn. Fl. Germ. ed. I. 209 = inclinatum.
pyrenaicum, *Mill. Gard. Dict.* ed. VIII. n. 8; *Willd. Sp. Pl.* ii. 1115.—Mont. Pyren.
pyrenaicum, Ram. in Bull. Soc. Philom. ii. (1800) 140. t. 10 = sylvaticum.
Quellyon, Hort. Edinb. ex Lindl. Bot. Reg. sub t. 1848 = chiloense.
radiatum, *Michx. Fl. Bor.-Am.* i. 300.—Am. bor.
radiatum, Pursh, Fl. Am. Sept. i. 359 = calthifolium.
Rafinesqueanum, Steud. Nom. ed. II. i. 683 = rivale.
ranunculoides, Ser. in Mém. Soc. Phys. Genève. ii. (1824) 138 = strictum.
renifolium, *F. Muell. in Hook. Kew Journ.* ix. (1857) 300.—Tasmania.
reptans, *Linn. Sp. Pl.* 501.—Europ.
rhaeticum ×, Brugg. in Jahresb. Naturf. Ges. Graub. ii. xxv. (1882) 66.—Europ.
rivale, Lamk. ex Nym. Consp. 230 = inclinatum.
rivale, *Linn. Sp. Pl.* 501.—Reg. bor. temp.
rivale, Tenore, Syll. in Add. et Emend. 528 = micropetalum.
rivulinentanum, Reichb. Fl. Germ. Excurs. 598 = *intermedium*.
robustum, Schur, Enum. Pl. Transs. 183 = urbanum.
Rossii, *Ser. in DC. Prod.* ii. 553.—Reg. arct.
rotundifolium, Fisch. in Mém. Soc. Nat. Mosc. ii. (1809) 187 = calthifolium.
Regeli, Wall. Cat. n. 713 = urbanum.
rubellum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 45.—Hab.?
rubifolium, Lej. in Lej. & Court. Comp. Fl. Belg. ii. 150; *et Rev. Fl. Spa*, 108 = urbanum.

GEUM:—
rubifolium, Schur, Enum. Pl. Transs. 184 = inclinatum
rugosum, Desf. Cat. Hort. Par. ed. III. 409 = strictum.
Sadleri, Privald. in Flora, xviii. (1835) 382 = coccineum.
Siebaldii, Hort. ex Schenk, in Nov. Act. Soc. Sc. Upsal. vii. ii. (1870) n. vi. 32. obs. 3 = japonicum.
Siewerti, Hort. ex Schenk, i. c. 26 = elatum.
sordidum, Salisb. Prod. 363 = urbanum.
spurium, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. *Suppl.* 38.—Hab.?
strictum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 217.—Reg. bor. temp.
sudeticum, Tausch, Hort. Canal. Enum. 90 = inclinatum.
sylvaticum, *Pourr. in Mém. Acad. Toul.* iii. (1788) 319.—Lusitan.; Afr. bor.
Thomasianum, Ser. in Mém. Soc. Phys. Genèv. ii. (1824) 140 = inclinatum.
tirolense ×, Kern. in Oestr. Bot. Zeitschr. xvii. (1867) 105.—Europ.
Tournefortii, Lapeyr. Hist. Abr. Pl. Pyr. 292 = pyrenaicum.
triflorum, *Pursh, Fl. Am. Sept.* ii. 736.—Am. bor.
triflorum, Torr. in Ann. Lyc. N. York, i. (1824) 195 = Rossii.
umbrosum, Boiss. Voy. Bot. Espagne, Suppl. 728 = heterocarpum.
umbrosum, Dum. Fl. Belg. Prod. 97 = urbanum?
uniflorum, *Y. Buch. in Trans. N. Z. Inst.* ii. (1870) 58.—N. Zel.
urbanum, A. Gray, Man. Bot. N. U. St. ed. IV. p. xcii = album.
urbanum, *Linn. Sp. Pl.* 501.—Reg. temp. bor.; Austral.; N. Zel.
vernum, *Torr. & Gray, Fl. N. Am.* i. 422.—Am. bor.
vicinum, Schur, Enum. Pl. Transs. 183 = urbanum.
Vidalii, *Franch. & Sav. Enum. Pl. Jap.* ii. 235.—Japon.
virginianum, Georgi, Beschr. Russ. Reich. iv. 1088 = strictum.
virginianum, *Linn. Sp. Pl.* 500.—Am. bor.
virginianum, Murr. in Comm. Gotting. v. (1783) 82 t. 3 = album.
virginianum, Spreng. Syst. ii. 542 = geniculatum.
Waldsteinia, Baill. Hist. Pl. i. 377 = Waldsteinia fragarioides.
Willdenowii ×, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. Suppl. 43 = *intermedium*.

GEUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 122 (1745) = Saxifraga, Tourn.
erectum, Moench, Meth. 64 = S. Geum.
rotundifolium, Moench, l. c. = S. rotundifolia.
umbrosum, Moench, l. c. 65 = S. umbrosa.

GHUNSIA, Blume, Cat. Gew. Buitenz. 11, 48 (1822); et Bijdr. 819 (1826). *VERBENACEÆ*, Benth. & Hook. f. ii. 1150.
Cunningiana, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 215.—Ins. Philipp.
farinosa, *Blume, Cat. Gew. Buitenz.* 48; *et Bijdr.* 819.—Java.
fastuosa, Rafin. Fl. Tellur. iv. 61 = Hypoëstes fastuosa.

GEUNSIA, Moç. & Sesse, ex DC. Prod. iii. 358 (1828) = Calandrinia, H. B. & K. (Portul.).
rosea, Moç. & Sesse, l. c. 359 = C. caulescens.

GEUNSIA, Neck. Elem. i. 351 (1790) = Dioliptera, Juss. *sed* Justicia, Linn. (Acanth.).

GEUNZIA, Neck. Elem. ii. 440 (1790) = Samyda, Linn.

GEVUINA, Molina, Sagg. Chil. 184 (1782). *PROTEACEÆ*, [Benth. & Hook. f. iii. 177.]
QUADRIA, Ruiz & Pav. Fl. Per. et Chil. Prod. 16. t. 28 (1794).
Avellana, *Molina, l. c.*—Chili.

GHESAKMBILLA, Adans. Fam. ii. 449 (1763) = Kimbalia, Juss. (Myrsin.).

GHIESBRECHTIA, Lindl. Veg. Kingd. 188 (1847) = GHIESBREGHTIA, Kich. & Gal. = Calanthe, R. Br. (Orchid.).

GHIESBREGHTIA, A. Gray, in Proc. Am. Acad. viii. (1872) 629. *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 926.
grandiflora, *A. Gray, l. c.* 630.—Mexic.

GHIESBREGHTIA, A. Rich. & Gal. in Ann. Sc. Nat Sér. III. iii. (1845) 28 = **Calanthe**, R. Br. (Orchid.).
calanthoides, A. Rich. & Gal. l. c. = *C.* mexicana

GHIESBREGHTIA, Lindl. Fol. Orch. (1854) = præc.

GHINIA, Schreb. Gen. 19 (1789) = **Tamonea**, Aubl. (Verbenac.).
mutica, Sw. Prod. Fl. Ind. Occ. 94; Willd. Sp. Pl. i. 114 = T. spicata.
spinosa, Willd. l. c. = T. curassavica.
Tamonea, J. F. Gmel. Syst. 37 = T. spicata.
verbenacea, Sw. Fl. Ind. Occ. iii. 1089 = T. curassavica.

GIBASIS, Rafin. Fl. Tellur. ii. 16 (1836) = **Tradescantia**, Linn. (Commel.).
pulchella, Rafin. l. c. = T. pulchella.

GIBBARIA, Cass. in Bull. Soc. Philom. (1817) 139; et in Dict. Sc. Nat. xviii. 526 (1820) = **Osteospermum**, Linn. (Compos.).
bicolor, Cass. in Dict. Sc. Nat. xviii. 527.—Afr. austr.

GIBSONIA, Stocks, in Proc. Linn. Soc. i. (1848) 367 = **Calligonum**, Linn. (Polygonac.)

GIESEKIA, Agardh, Aphor. 218 (1825) = **Giesekia**, Linn. (Ficoid.).

GIESECKIA, Reichb. Nom. 166 (1841) = præc.

GIESLERIA, Regel, in Flora, xxxii. (1849) 161 = **Isoloma**, Decne. (Gesnerac.).
picta, Regel, l. c. 182 = I. pictum.

GIFOLA, Cass. in Bull. Soc. Philom. (1819) 142 = **Filago**, Linn. (Compos.).
canescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1869) 94 = F. germanica.
germanica, Dum. Fl. Belg. 68 = F. germanica.
lutescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1869) 94 = F. germanica.
pyramidalis, Dum. Fl. Belg. 69 = F. spathulata.
spathulata, Reichb. f. Ic. Fl. Germ. xvi. 26 = F. spathulata.
vulgaris, Cass. in Bull. Soc. Philom. (1819) 143 = F. germanica.

GIGALOBIUM, R. Br. Hist. Jamaic. 362 (1756) = **Entada**, Adans. (Legumin.).

GIGANTABIES, Senilis [Nelson,] Pinaceæ, 77 (1866) = **Sequoia**, Endl. (Conif.).
taxifolia, Senilis [Nelson,] l. c. 78 = S. sempervirens.
Wellingtoniana, Senilis [Nelson,] l. c. 79 = S. gigantea.

GIGANTHEMUM, Welw. Apont. 585 (1858). *LEGUMINOSÆ.*
scandens, *Welw. l. c.*—Afr. trop.

GIGANTOCHLOA, Kurz, in Tijdschr. Nederl. Ind. xxvii. (1864) 226. *GRAMINEÆ*, Benth. & Hook. f. iii. 1211.
albo-ciliata, Kurz, For. Fl. Brit. Burma, ii. 555 (gen. al.).—Burma.
andamanica, *Kurz, l. c.* 556.—Ins. Andaman.
Apua, *Kurz, in Tijdschr. Nederl. Ind.* xxvii. (1864) 226.—Java.
ater, *Kurz, l. c.*—Java.
auriculata, *Kurz, For. Fl. Brit. Burma*, ii. 557.—Burma.
heterostachya, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 135.—Malacca.
macrostachya, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 251.—Burma.
maxima, Kurz, in Tijdschr. Nederl. Ind. xxvii. (1864) 226 = verticillata.
nigrociliata, *Kurz, l. c.*—Malaya.
robusta, *Kurz, Bamboos*, 60.—Malaya.
verticillata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 134.—Java.

GIGLIOLIA, Bocc. Malesia, i. 171 (1877). *PALMÆ*, Benth. & Hook. f. iii. 904.
insignis, *Becc. l. c.* 173.—Borneo.
subacaulis, *Becc. l. c.* 174.—Borneo.

GIGLIOLIA, Rodrig. Orch. Nov. i. 25 (1877). *ORCHIDEÆ.*
gracilis, *Rodrig. l. c.* 26.—Bras.
lithophila, *Rodrig. l. c.*—Bras.

GILBERTA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 192 = **Myriocephalus**, Benth. (Compos.).
tenuifolia, Turcz. l. c. 193 = M. gracilis.

GILESIA, F. Muell. Fragm. ix. 41 (1875) = **Hermannia**, Linn. (Sterculi.).
binifora, F. Muell. l. c. 42 = H. Gilesii.

GILIA, Ruiz & Pav. Prod. Fl. Per. 25. t. 4 (1794). *POLEMONIACEÆ*, Benth. & Hook. f. ii. 832.
AEGOCHLOA, Benth. in Bot. Reg. sub t. 1622 (1833).
BATANTHES, Rafin. Atl. Journ. 145 (1832).
BRICKELLIA, Rafin. in Med. Repos. N. York, v. (1808) 352.
DACTYLOPHYLLUM, Spach, Hist. Vég. Phan. ix. 108 (1840).
FENZLIA, Benth. in Bot. Reg. sub t. 1622 (1833).
HUGELIA, Benth. l. c. (1833).
IPOMERIA, Nutt. Gen. Am. i. 124 (1818).
IPOMOPSIS, Michx. Fl. Bor. Am. i. 141 (1803).
LEPTODACTYLON, Hook. & Arn. Bot. Beech. Voy. t. 80 (1841).
LEPTOSIPHON, Benth. in Bot. Reg. sub t. 1622 (1833).
LINANTHUS, Benth. l. c. (1833).
NAVARRETIA, Ruiz & Pav. Prod. Fl. Per. 20 (1794).
ROSSMÄSSLERA, Reichb. Nom. 118 (1841).
WELWITSCHIA, Reichb. Handb. 194 (1840).
achilleæfolia, *Benth. in Bot. Reg.* sub t. 1622.—Calif.
achilleæfolia, Lindl. in Bot. Reg. t. 1622 = multicaulis.
aggregata, *Spreng. Syst.* i. 626.—Am. bor. occ.
andicola, *Phil. Fl. Atac.* 36.—Chili.
androsace, *Steud. Nom.* ed. II. i. 683.—Calif.
arcuata, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 367.—Reg. Argent.
armeria, Benth. in Bot. Reg. sub t. 1622 = inconspicua.
atractyloides, *Steud. Nom.* ed. II. i. 683.—Calif.
aurea, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 155.—Am. bor. occ.
bella, *A. Gray, in Proc. Am. Acad.* xx. (1885) 301.—Calif.
Berteri, *DC. Not. 5ma Pl. Rar. Hort. Genbo.* 20. t. 5.—Chili.
Beyrichiana, Bouché, in Linnæa, xiii. (1839) 81 = coronopifolia.
Bigelovii, *A. Gray, in Proc. Am. Acad.* viii. (1870) 265.—Am. bor. occ.
Bolanderi, *A. Gray, l. c.* 263.—Calif.
Brandegei, *A. Gray, l. c.* xi. (1876) 85.—Am. bor. occ.
brevicula, *A. Gray, l. c.* xii. (1877) 79.—Calif.
Breweri, *A. Gray, l. c.* viii. (1870) 269.—Am. bor. occ.
caespitosa, *A. Gray, l. c.* xii. (1877) 80.—Am. bor. occ.
californica, *Benth. in DC. Prod.* ix. 316.—Calif.
calycytrata, Rivers, in Bailli. Adansonia, x. (1872) 241 = Anthodiscus peruanus.
campanulata, *A. Gray, in Proc. Am. Acad.* viii. (1870) 279.—Am. bor. occ.
capillaris, *Kellogg, in Proc. Calif. Acad.* v. (1873) 46.—Am. bor. occ.
capitata, *Sims, Bot. Mag.* t. 2698.—Am. bor. occ.
ciliata, *Benth. Pl. Hartw.* 326.—Am. bor. occ.
congesta, *Hook. Fl. Bor. Am.* ii. 75.—Am. bor. occ.
coronopifolia, *Pers. Syn.* i. 187.—Am. bor.
cotulæfolia, *Steud. Nom.* ed. II. i. 683.—Calif.
crassifolia, *Benth. in Bot. Reg.* sub t. 1622.—Chili.
crebrifolia, Nutt. in Journ. Acad. Philad. N.S. i. (1847) 156 = congesta.
dactylophylla, *Torr. in Ives, Rep. Colorado River, Bot.* 23.—Am. bor.
debilis, *S. Wats. in Am. Natural.* vii. (1873) 302.—Am. bor. occ.
demissa, *A. Gray, in Proc. Am. Acad.* viii. (1870) 263.—Am. bor. occ.
densiflora, *Benth. in DC. Prod.* ix. 316.—Calif.
densifolia, *Benth. l. c.* 311.—Am. bor. occ.
depressa, *M. E. Jones, ex A. Gray, in Proc. Am. Acad.* xvi. (1881) 106.—Am. bor. occ.
dianthifera, Benth. in DC. Prod. ix. 314, in syn. = seq.
dianthoides, *Endl. Atakt. Bot.* t. 29.—Calif.
dichotoma, *Benth. in DC. Prod.* ix. 314.—Am. bor. occ.
diffusa, *Phil. in Linnæa*, xxx. (1859–60) 197.—Chili.
divaricata, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 155 = Collomia gilioides.

GILIA:—
divaricata, *Torr. ex A. Gray, in Proc. Am. Acad.* viii. (1870) 270.—Calif.
Douglassii, Gill. ex Benth. in DC. Prod. ix. 310 = gossipifera.
Dunnii, *Kellogg, in Bull. Calif. Acad.* i. 11. (1885) 142. —Calif.
elongata, Steud. Nom. ed. II. i. 683 = densifolia.
erecta, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 368.—Reg. Argent.
eryngioides, Lehm. in Flora, xvi. (1833) 122 = G. Navarretia.
Fenzlia, Steud. Nom. ed. II. i. 683 = dianthoides.
filicaulis, *Torr. ex A. Gray, in Proc. Am. Acad.* viii. (1870) 270.—Calif.
filifolia, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 156.—Am. bor. occ.
filiformis, *Parry, ex A. Gray, in Proc. Am. Acad.* x. (1875) 76.—Am. bor. occ.
filipes, Benth. Pl. Hartw 325 = pusilla.
floccosa, *A. Gray, in Proc. Am. Acad.* viii. (1870) 272. —Am. bor. occ.
floribunda, *A. Gray, l. c.* 267.—Am. bor. occ.
floridana, G. Don, Gen. Syst. iv. 245 = coronopifolia.
foetida, *Gill. ex Benth in DC. Prod.* ix. 313.—Chili.
Gayana, *Wedd. Chlor. And.* ii. 82.—Chili.
glandulosa, Scheele, in Linnæa, xxi. (1848) 753 = rigidula.
glomeriflora, Juss. in Ann. Mus. Par. iii. (1804) 119 = Collomia Cavanillesiana.
glomeriflora, *Spreng. Syst.* i. 626.—Am. austr.
glutinosa, *Phil. in Linnæa*, xxx. (1859–60) 196.—Chili.
gossipifera, *Gill. ex Benth. in DC. Prod.* ix. 310.—Reg. Argent.
gracilis, Hook. Bot. Mag. t. 2994 = Collomia gracilis.
grandiflora, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 223.—Calif.
grandiflora, Steud. Nom. ed. II. i. 683 = densiflora.
Gunnisoni, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 129. t. 9.—Am. bor. occ.
Harknessii, *Curran, in Bull. Calif. Acad.* i. (1884) 12. —Am. bor. occ.
Haydeni, *A. Gray, in Proc. Am. Acad.* xi. (1876) 85.—Am. bor. occ.
heterodoxa, Greene, in Bull. Calif. Acad. i. n. 1. (1884) 10 = vaccifolia.
heterophylla, *Dougl. ex Hook. Bot. Mag.* t. 2895.—Am. bor. occ
Hookeri, Benth. in DC. Prod. ix. 316 = pungens.
Hugelia, Steud. Nom. ed. II. i. 683 = densifolia.
iberidifolia, *Benth. in Hook. Kew Journ.* iii. (1851) 291.—Am. bor. occ.
incisa, *Benth. in DC. Prod.* ix. 312.—Texas.
inconspicua, Sweet, Hort. Brit. ed. I. 286 = parviflora.
intermedia, *Phil. in Linnæa*, xxxiii. (1864–65) 181.—Chili.
intertexta, *Steud. Nom.* ed. II. i. 683.—Am. bor. occ.
Kennedyi, Porter, in Coult. Bot. Gaz. ii. (1877) 78 = Parryæ.
lacinlata, *Ruiz & Pav. Fl. Per.* ii. 17.—Peruv.; Chili.
lanata, *Walp. Ann.* i. 519.—Mexic.
languinosa, *Phil. in Linnæa*, xxix. (1857–58) 14.—Chili.
Larseni, *A. Gray, in Proc. Am. Acad.* xi. (1876) 84.—Calif.
latiflora, *A. Gray, Syn. Fl. N. Am.* ii. 1. 147.—Calif.
latifolia, *S. Wats. in Am. Natural.* ix. (1875) 347.—Am. bor. occ.
Lemmoni, *A. Gray, Syn. Fl. N. Am.* ii. 1. 394.—Calif.
leptomeria, *A. Gray, in Proc. Am. Acad.* viii. (1870) 278.—Am. bor. occ.
Leptosiphon, Steud. Nom. ed. II. i. 681 = densiflora.
leptotes, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 223. —Am. bor. occ.
leucocephala, *A. Gray, l. c.* viii. (1870) 270.—Calif.
Linanthus, Steud. Nom. ed. II. i. 681 = dichotoma.
Lindheimersana, Scheele, in Linnæa, xxi. (1848) 758 = incisa.
linearis, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 223. —Calif.
linifora, *Benth. in Bot. Reg.* sub t. 1622.—Am. bor. occ.
longiflora, G. Don, Gen. Syst. iv. 245 = Collomia longiflora.
longituba, Benth. Pl. Hartw. 324 = micrantha.
luteo, Steud. Nom. ed. II. i. 681 = micrantha.
lutescens, *Steud. l. c.*—Calif.

GILIA :—
Macombii, *Torr. ex A. Gray, in Proc. Am. Acad.* xx. (1885) 301.—Arizona.
micrantha, *Steud. Nom.* ed. II. i. 684.—Calif.
micromeria, *A. Gray, in Proc. Am. Acad.* viii. (1870) 279.—Am. bor. occ.
millefoliata, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 35 = multicaulis.
minima, *A. Gray, in Proc. Am. Acad.* viii. (1870) 269. —Am. bor. occ.
minutiflora, *Benth. in DC. Prod.* ix. 315.—Am. bor. occ.
nucronata, Lehm. Del. Sem. Hort. Hamb. (1832) = G. Navarretia.
multicaulis, *Benth. in Bot. Reg.* sub t. 1622.—Calif.
multiflora, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 104.—N. Mexic. ; Calif.
multiflora, Torr. Bot. Mex. Bound. 146, partim = Macombii.
Navarretia, *Steud. Nom.* ed. II. i. 684.—Chili.
nudicaulis, *A. Gray, in Proc. Am. Acad.* viii. (1870) 266.—Am. bor. occ.
Nuttallii, *A. Gray, l. c.* 267.—Am. bor. occ.
Orcuttii, *A. Gray, ex Parry, in Proc. Davenp. Acad.* iv. (1884) 46.—Calif.
Parryae, *A. Gray, in Proc. Am. Acad.* xii. (1877) 76.—Calif.
parviflora, *Spreng. Syst.* i. 626 —Am. bor. occ.
pharnaceoides, Benth in Bot. Reg. sub t. 1622 = liniflora.
pinnatifida, *Nutt. ex A. Gray, in Proc. Am. Acad.* viii. (1870) 276 —Am. bor. occ.
polycladon, *Torr. in Bot. Mex. Bound.* 147.—Texas.
prostrata, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 223.—Calif
pubescens, *Steud. Nom.* ed. II. i. 684.—Calif.
pulchella, Dougl. ex Benth. in Bot. Reg. sub t. 1622 = aggregata.
pumila, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 156.—Texas.
pungens, *Benth. in DC. Prod.* ix. 316.—Am. bor. occ
pungens, Dougl. ex Hook. in Bot. Mag. t. 2977 = squarrosa.
pusilla, *Benth. in Bot. Reg.* sub t. 1622.—Calif. ; Chili.
rigidula, *Benth. in DC. Prod.* ix. 312.—Am. bor. occ.
Schottii, *A. Gray, in Bot. King's Exped.* 267.—Am. bor. occ.
scopulorum, *M. E. Jones, in Bull. Torrey Bot. Club*, viii. (1881) 70.—Utah.
Sessei, G. Don, Gen. Syst. iv. 245 = heterophylla.
setosissima, *A. Gray, in Proc. Am. Acad.* viii (1870) 271.—Calif.
sinuata, Dougl. ex Benth. in DC. Prod. ix. 313 = parviflora.
spicata, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 156.—Am. bor. occ.
squarrosa, *Hook. & Arn. Bot. Beech. Voy.* 151.—Am. bor. occ.
stenothyrsa, *A. Gray, in Proc. Am. Acad.* viii. (1870) 276.—Am. bor. occ.
stricta, *Liebm. Ind. Sem. Hort. Haun.* (1853) 21.—Hab. ?
stricta, Scheele, in Linnæa, xxi. (1848) 755 = multicaulis.
subnuda, *Torr. ex A. Gray, in Proc. Am. Acad.* viii. (1870) 276.—Am. bor. occ
tenella, *Benth. Pl. Hartw.* 325.—Am. bor. occ.
tenerrima, *A. Gray, in Proc. Am. Acad.* viii. (1870) 277.—Am. bor. occ.
tenuiflora, *Benth. in Bot. Reg.* sub t. 1622.—Calif.
Thurberi, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 223.—Arizona.
tinctoria, Kellogg, in Bull. Calif. Acad. i. 111. (1885) 142 = linearis.
tomentosa, *Mart & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 273.—Mexic.
tricolor, *Benth. in Bot. Reg.* sub t. 1622.—Calif.
trifida, *Benth. in Hook. Kew Journ.* iii. (1851) 291.—Am. bor.
trifida, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 156 = spicata.
valdiviensis, *Griseb. in Gœtt. Abh.* vi. (1854) 121.—Chili.
Veatchii, *Parry, ex Greene, in Bull. Calif. Acad.* i. iv. (1885) 198.—Calif.
virgata, *Steud. Nom.* ed. II. i. 684.—Calif.
viscidula, *A. Gray, in Proc. Am. Acad.* viii. (1870) 271. —Calif.
Watsoni, *A. Gray, l. c.* 267.—Am. bor. occ.
Wrightii, *A. Gray, l. c.* 278.—Texas.

GILIBERTA, St. Lag. in Ann. Soc. Bot. Lyon, viii. 171 (1881) = seq.

GILIBERTIA, J. F. Gmel. Syst. 682 (1791) = **Quivisia**, Comm. (Meliac.).
decandra, J. F. Gmel. l. c. = Q. decandra.
heterophylla, J. F. Gmel. l. c. = Q. heterophylla.
oppositifolia, J. F. Gmel. l. c. = Q. oppositifolia.
ovata, J. F. Gmel. l. c. = Q. ovata.
rutilans, Sm. in Rees, Cycl. xvi. n. 4 = Q. ovata.

GILIBERTIA, Ruiz & Pav. Prod. Fl. Per. 50. t. 8 (1794). *ARALIACEAE*, Benth. & Hook. f. i 944.
GINANNIA, F. G. Dietr. Vollst. Lexik. Gaertn. iv. 857 (1804).
WANGENHEMIA, F. G Dietr. l. c. x. 536 (1810).
affinis, *March. in Mart. Fl. Bras.* xi. 1. 251.—Bras.
brasiliensis, *Seem. in Journ. Bot.* iii. (1865) 175.—Bras.
cuneata, *March. in Mart. Fl. Bras.* xi. 1. 250.—Bras.
dichrostachya, Boj. Hort. Maurit. 162 = Polyscias dichrooostachya.
heterophylla, *March. in Mart. Fl. Bras.* xi. 1. 246.—Bras.
Langeana, *March. in Bull Acad. Belg.* Sér. II. xlvii. (1879) 79.—Mexic.
Langsdorffi, *March. in Mart. Fl. Bras.* xi. 1. 248 — Bras.
monogyna, *March. l. c.* 247.—Bras.
Nalagu, DC. Prod. iv. 256 = Lees sambucina.
palmata, DC. l. c. = Trevesia palmata.
paniculata, DC. l. c. = Polyscias paniculata.
Pavoni, *March. in Mart. Fl. Bras.* xi. 1. 248 —Bras.
populifolia, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 77 —Mexic.
repanda, DC. Prod. iv. 256 = Polyscias repanda.
resinosa, *March. in Mart. Fl. Bras.* xi. 1. 249.—Bras.
saururoides, *DC Prod.* iv. 256.—Ins Molucc.
triloba, *March. in Mart. Fl. Bras.* xi. 1. 247.—Bras.
umbellata, *Ruiz & Pav. Fl. Per.* iii. 75. t. 312.—Peruv.

GILIPUS, Rafin. Sylva Tellur. 61 (1838) = **Cephalanthus**, Linn. (Rubiac.).
montanus, Rafin. l. c. = C. montanus.

GILLBEEA, F. Muell. Fragm. v. 17 (1865). *SAXIFRAGEAE*.
adenopetala, *F. Muell. l. c.*—Austral.

GILLENA, Adans. Fam. ii 166 (1763) = **Clethra**, Linn. (Ericac.).

GILLENIA, Moench, Meth. Suppl. 286 (1802). *ROSACEAE*, Benth. & Hook. f. i. 618.
stipulacea, *Nutt. Gen. Am.* i. 307.—Am. bor.
trifoliata, *Moench, Meth. Suppl.* 286.—Am. bor.

GILLENIA, Stend. Nom. ed. II. i. 684 (1840), aphalm. = GILLENA, Adans. = **Clethra**, Linn. (Ericac.).

GILLIA, Endl. Gen. Suppl. i. 1408 (1841), aphalm. = **Gilia**, Ruiz & Pav. (Polem.).

GILLIESIA, Lindl. in Bot. Reg. t. 992 (1826). *LILIACEAE*, Benth. & Hook. f. iii. 804.
Gaudichaudiana, *Kunth, Enum. Pl.* iv. 491.—Chili.
graminea, *Lindl. in Bot. Reg.* t. 992.—Chili.
monteria, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 27. t. 138.—Chili.

GILLONIA, A. Juss. in Orb. Dict. xi. 209 (1849), aphalm. = **Gillenia**, Moench (Rosac.).

GIMBERNATEA, Ruiz & Pav. Prod. Fl. Per. 138. t. 36 (1794) = **Terminalia**, Linn. (Combret.).
Calamansanay, Blanco, Fl. Filip. ed. II. 266 = T. blalata.
oblonga, Ruiz & Pav. Syst. Veg. 275 = T. oblonga.
obovata, Ruiz & Pav. l. c. 274 = T. obovata.

GINALLOA, Korth. in Verh. Batav. Gen. xvii. (1839) 260 (Loranth. 64). *LORANTHACEAE*, Benth. & Hook. f. iii. 215.
andamanica, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 209.—Ins. Andaman.

GINALLOA :—
Arnottiana, *Korth. in Verh. Batav. Gen.* xvii. (1839) 260 (Loranth. 64).—Malaya.
Cumingiana, *Benth. & Hook. f. Gen.* iii. 215.—Ins. Philipp.
Helferi, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 84.—Burma.
spathulifolia, *Oliver, in Journ. Linn. Soc.* vii. (1864) 108.—Zeylan.

GINANNIA, F. G. Dietr. Vollst. Lexik. Gaertn. iv. 857 (1804) = **Gilibertia**, Ruiz & Pav. (Araliac.).
umbellata, F. G. Dietr. l. c. Nachtr. iii. 488 = Gilibertia umbellata.

GINANNIA, Scop. Introd. 300 (1777) = **Palovea**, Aubl. (Legumin.).
guianensis, J. F. Gmel. Syst. 558 = P. guianensis.

GINGIDIUM, Forst. Char. Gen. 41. t. 21 (1776) = **Angelica**, Riv. (Umbellif.).
montanum, Forst. l. c. 42 = A. Gingidum.

GINGIDIUM, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 108 = **Actiphylla**, Forst. (Umbellif.).
antipodum, F. Muell. Veg. Chath. Isl. 18 (= *Anisotome antipoda*).—Ins. Chath.
Diefenbachii, F. Muell. l. c. 17.—Ins. Chath.
glaciale, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 108 = A. glacialis.
Monroi, F. Muell. Fragm. i. 16 = A. Monroi.
procumbens, F. Muell. l. c. 15 = A. procumbens.
simplicifolium, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 104 = A. simplicifolia.
squarrosum, F. Muell. Veg. Chath. Isl. 18, nomen = A. squarrosa.
Traversii, F. Muell. l. c. = A. Traversii.

GINGINSIA, DC. Prod. iii. 362 (1828) = **Pharnaceum**, Linn. (Ficoid.).
albens, DC. l. c. 363 = P. albens.
arenaria, DC. l. c. = P. reflexum.
brevicaulis, DC. l. c. = P. dichotomum.
conferta, DC. l. c. = P. incanum.
elongata, DC. l. c. = P. incanum.
microphylla, DC. l. c. = P. microphyllum.
pruinosa, G. Don, Gen. Syst. iii. 88 = P. pruinosum.
teretifolia, DC. Prod. iii. 363 = P. teretifolium.

GINKGO, Linn. Mant. ii. 313 (1771). *CONIFERAE*. Benth. & Hook. f. iii. 432.
PTEROPHYLLUS, Senilia [Nelson,] Pinaceae, 163 (1866)
SALISBURIA, Sm. in Trans. Linn. Soc. iii. (1797) 330.
biloba, *Linn. l. c.*—Japon.
macrophylla, Hort. ex C. Koch, Dendrol. ii. ii. 108 = biloba.

GINNANIA, M. Roem. Syn. Hesper. 79, 90 (1846) = **Quivisia**, Comm. (Meliac.).
oppositifolia, M. Roem. l. c 90 = Q. oppositifolia.
racemosa, M. Roem. l. c. = Q. decandra.

GINORA, Linn. Sp. Pl. ed. II. 642 (1762). *LYTHRACEAE*, Benth. & Hook. f. i. 780.
americana, *Linn. l. c.*—Ind. occ.
curvispina, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 349.—Cuba.
Diplusodon, *Koehne, l. c.* 350.—Cuba.
glabra, *Griseb. Cat. Pl. Cub.* 106.—Cuba.
nudiflora, Koehne, in Engl. Bot. Jahrb. iii. (1882) 351 = Autherylium nudiflorum.
Rohrii, Koehne, l. c. = Autherylium Rohrii.
spinosa, *Griseb. Cat. Pl. Cub.* 106.—Cuba.

GINORIA, Jacq. Enum. Pl. Carib. 5 (1760) = **Ginora**. Linn. (Lythrar.).
americana, Jacq. l. c. = Ginora americana.

GINORIA, [Moç. & Sesse,] ex DC. Prod. iii. 89 (1828) = **Hintmia**, Link & Otto (Lythrar.).
flava, [Moç. & Sesse,] ex DC. l. c. = H. salicifolia.
syphilitica, [Moç. & Sesse,] ex DC. l. c. = H. syphilitica.

GINSA, Stend. Nom. ed. II. ii. 349, in syn. (1841). aphalm. = seq.

GINSEN, Adans. Fam. ii. 102 (1763) = **Panax**, Linn. (Araliac.).

GIRARDINIA, Gaudich. in Freyc. Voy. Bot. 498 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 384.
adoensis, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 208 = condensata.
armata, Kunth, Ind. Sem. Hort. Berol. (1846) 11 = palmata.
bulbosa, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 208.—Abyss.
condensata, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 169.—Abyss.
cuspidata, *Wedd. in DC. Prod.* xvi. i. 103.—China.
crosa, Decne. in Jacquem. Voy. Bot. 161 = palmata.
furcalis, Blume, Mus. Bot. Lugd. Bat. ii. 159 = Obetia ficifolia.
heterophylla, Decne. in Jacquem. Voy. Bot. 151. t. 153 = palmata.
hibiscifolia, Miq. Pl. Jungh. 81 = palmata.
javanica, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 181 = palmata.
Leschenaultiana, Decne. in Jacquem. Voy. Bot. 152 = palmata.
palmata, *Gaudich. in Freyc. Voy. Bot.* 498.—Ind. or.; Malaya.
Vahlii, Blume, Mus. Bot. Lugd. Bat. ii. 158 = palmata.
vitifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 181 = palmata.
zeylanica, Decne. in Jacquem. Voy. Bot. 152 = palmata.

GIKEOUDIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 125; et Abh. Akad. Berl. (1854) 84. t. 7 C = **Begonia**, Linn.
asarifolia, Klotzsch, l. c. = B. hydrocotylifolia.
Barkeri, Klotzsch, l. c. = B. Barkeri.
cardiocarpa, Klotzsch, l. c. = B. cardiocarpa.
caroliniaefolia, Klotzsch, l. c. = B. caroliniaefolia.
carpinifolia, Klotzsch, l. c. = B. carpinifolia.
conchaefolia, Klotzsch, l. c. = B. conchaefolia.
crassicaulis, Klotzsch, l. c. = B. crassicaulis.
fibrillosa, Klotzsch, l. c. nomen.—Veragua.
fimbriata, Klotzsch, l. c. = B. fimbriata.
heracleifolia, Klotzsch. l. c. = B. heracleifolia.
hydrocotylefolia, Klotzsch, l. c. = B. hydrocotylifolia.
involucrata, Klotzsch, l. c. = B. involucrata.
laciniata, Klotzsch, in Abh. Akad. Berl. (1855) 85 = B. laciniata.
Lindleyana, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 126 = B. Lindleyan.
lobulata, Klotzsch, l. c. = B. lobulata.
macrophylla, Klotzsch, l. c. = B. macrophylla.
manicata, Klotzsch, l. c. = B. manicata.
multinervia, Klotzsch, l. c. = B. multinervia.
nelumbiifolia, Klotzsch, l. c. = B. nelumbiifolia.
Ottoniana, Regel, Gartenfl. (1859) 15 = B. Ottoniana.
pilifera, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 125 = B. pilifera.
plebeja, Klotzsch, l. c. = B. plebeja.
pruriseta, Klotzsch, l. c. = B. pruriseta.
punctata, Klotzsch, l. c. = B. heracleifolia.
rhizocaulis, Klotzsch, in Abh. Akad. Berl. (1855) 84 = B. rhizocaulis.
rotata, Klotzsch, l. c. 91 = B. caroliniaefolia.
sarcophylla, Klotzsch, l. c. 88 = B. sarcophylla.
schiedeifolia, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 126 = B. manicata.
sericonevra, Klotzsch, in Abh. Akad. Berl. (1855) 89 = B. sericoneura.
setosa, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 126, nomen = B. setosa.
setulosa, Klotzsch, in Abh. Akad. Berl. (1855) 91 = B. setulosa.
squarrosa, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 125 = B. squarrosa.
stigmosa, Klotzsch, l. c. = B. stigmosa.
strigillosa, Klotzsch, l. c. = B. strigillosa.
sycophylla, Klotzsch, in Abh. Akad. Berl. (1855) 92 = B. urophylla.
vitifolia, Klotzsch, l. c. 87 = B. vitifolia.
Warmscioldiana, Hort. Berol. ex A. DC. in DC. Prod. xv. i. 388 = B. conchaefolia.

GIRGENSOHNIA, Bunge, Rel. Lehm. Bot. 302; in Mém. Sav. Etr. Pétersb. vii. (1847) 478. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 72.
diptera, *Bunge, ll. cc.* 303 et 479.—Turkestan.
fruticulosa, *Bunge, Anabas. Rev.* 30.—Cappadocia.
gypsophiloides, Bunge, in Rel. Lehm. Bot. 303 = oppositiflora.
heteroptera, Bunge, l. c. = oppositiflora.
imbricata, *Bunge, Anabas. Rev.* 32.—Persia.
oppositiflora, *Fenzl, in Ledeb. Fl. Ross.* iii. 835.—Persia; Afghan.
Pallasii, Bunge, in Rel. Lehm. Bot. 303 = oppositiflora.

GIROA, Steud. Nom. ed. I. 867 (1821), sphalm. = **Guioa**, Cav.

GIRONNIERA, Gaudich. Bot. Voy. Bonite, t. 85 (1844–46). *URTICACEAE*, Benth. & Hook. f. iii. 356.
DICERA, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 67, 74 (1856) [1856].
GALUMPITA, Blume, l. c. 78 (1856) [1856].
HELMINTHOSPERMUM, Thw. in Hook. Kew Journ. vi. (1854) 301. t. 9.
NEMOSTIGMA, Planch. in Ann Sc. Nat. Sér. III. x. (1848) 265.
celtidifolia, *Gaudich. Bot. Voy. Bonite*, t. 85.—Ins. Philipp.
chinensis, *Benth. Fl. Hongk.* 325.—China.
costata, Miq. in Zoll. Syst. Verz. Ind. Archip. 88 = subaequalis.
cuspidata, Kurz, For. Fl. Burma, ii. 470 = reticulata.
lucida, *Kurz, l. c.*—Ins. Andamon.
nervosa, *Planch. in Ann. Sc Nat.* Sér. III. x. (1848) 886.—Malaya.
nitida, *Benth. Fl. Hongk.* 325.—China.
parvifolia, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 888.—Malaya.
reticulata, *Thw. Enum. Pl. Zeyl.* i. 268.—Ind. or.; Malaya.
reticulosa, Planch. in DC. Prod. xvii. 207, sphalm. = preec
rhamnifolia, *Blume, Mus. Bot. Lugd. Bat.* ii. 74.—N. Guinea.
subaequalis, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 889.—Ind. or.; Malaya.

GIRTANNERIA, Neck. Elem. ii. 121 (1790) = **Rhamnus**, Tourn.
alnifolia, Kafn Sylva Tellur. 28 = R. alnifolius.
frangulioides, Kafn. l. c. = R. Humboldtianus?

GISECHIA, Linn. Mant. ii. 562 (1771), err. typ. = **Giesekia**, Linn. (Ficoid.).

GISECKIA, Willd. Sp. Pl. i. 1547 (1797) = seq.

GISEKIA, Linn. Mant. ii. 554 (1771). *FICOIDEAE*, Benth. & Hook. f. i. 859.
KOELREUTERA, Murr. in Nov. Comm. Gotting. iii. (1773) 67. t. 2. f. 1.
MILTUS, Lour. Fl. Cochinch. 302 (1790).
sapera, *Klotzsch, in Peters, Reise Mossamb. Bot.* 186.—Afr. trop.
Burchellii, Moq. in DC. Prod. xiii. ii. 28 = pentadecandra.
congesta, *Moq. l. c.*—Afr. trop.
linariifolia, Fenzl, ex Moq. l. c. 27 = rubella.
linearifolia, Schum. & Thonn. Beskr. Guin. Pl. 167 = pharnaciodes.
Miltus, *Fenzl, Nov. Stirp. Dec. Vind.* x. 86.—Afr. trop.
molluginoides, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 162 = pharnaciodes.
occulta, Schult. Syst. vi. 735 = pharnaciodes.
pentadecandra, *E. Mey. ex Moq. in DC. Prod.* xiii. ii. 28.—Afr. austr.
pharnaciodes, *Linn. Mant.* ii. 562.—Ind. or.
rubella, *Hochst. ex Moq. in DC. Prod.* xiii. ii. 27.—Afr. trop.

GISSANTHE, Salisb. in Trans. Hort. Soc. i. (1812) 279 = **Costus**, Linn. (Scitamin.).
spiralis, Salisb. l. c. = C. spiralis.

GISSIPIUM, Medic. Beobacht. (1782) 198 = **Gossypium**, Linn. (Malvac.).

GISSONIA, Salisb. in Knight, Prot. 32 (1809) = **Leucodendron**, K. Br. (Proteac.).
collina, Knight, Prot. 33 = L. plumosum.
riparia, Knight, l. c.—Afr. austr.
sericea, Knight, l. c.—Afr. austr.

GITHAGO, Adans. Fam. ii. 255 (1763) = **Lychnis**, Linn. (Caryophyll.).
gracilis, Boiss. Fl. Orient. i. 661.—Ax. Min.
nicaeensis, Link, Diss. Bot. Suerin, (1795) 62 = L. Githago
segetalis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = L. Githago.
segetum, Link, Diss. Bot. Suerin, (1795) 62; Desf. Tabl. ed. I. 159 = L. Githago.

GITHOPSIS, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1842) 358. *CAMPANULACEAE*, Benth. & Hook. f. ii. 559.
calycina, Benth. Pl. Hartw. 321 = specularioides.
diffusa, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 221.—Calif.
pulchella, Vatke, in Linnaea, xxxvlii. (1874) 714 = specularioides.
specularioides, *Nutt. in Trans. Am. Phil. Soc.* N. S. viii. (1842) 358.—Am. bor. occ.

GIVOTIA, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 389. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 297.
GOVANIA, Wall. Cat. n. 7851 (1847?).
rottleriformis, *Griff. l. c.* 14.—Ind. or.

GLABKARIA, Linn. Mant. ii. 156 (1771); Miq. Fl. Ind. Bat. i. 1. 940 (1857) = **Litsea**, Lam. (Laurin.).
cuneata, Miq. l. c. 942 (= *Tetranthera cuneata*).—Malaya.
litoralis, Miq. l. c. = L. sebifera.
mollis, Miq. l. c. 941 (= *Tetranthera mollis*).—Java.
nitens, Miq. l. c. 943 (= *Tetranthera nitens*).—Java.
terax, Linn. Mant. ii. 278 = L. sebifera.
vestita, Miq. Fl. Ind. Bat. i. 1. 941 = L. tomentosa, *Blume*.

GLADIOLUS, Geertn. Fruct. i. 31. t. 11. f. 4 (1788) = **Antholyza**, Linn. (Irid.).

GLADIOLUS, [Tourn.] Linn. Syst. ed. I (1735). *IRIDEAE*, Benth. & Hook. f. iii. 709.
BALLONIPHYM, HYPTISA, OPHIOLYZA, RANISIA, SYMPHYDOLON, Salisb. Gen. Pl. Fragm. 142 (1866).
SYMAKROSORA, Sweet, Hort. Brit. ed. II. 501 (1839).
abbreviatus, Andr. Bot. Rep. t. 166 = Antholyza quadrangularis.
abyssinicus, *Brongn. in A. Rich. Tent. Fl. Abyss.* ii. 306.—Abyssin.
adenandriflorus, *Echl. Top. Vers.* 88.—Afr. austr.
aequinoctialis, Herb. in Lindl. Bot. Reg. (1842) Misc. 85 = Acidanthera aequinoctialis.
aethiopicus, *Drapiez, in Dict. Sc. Nat.* v. (1858) 129 (Quid?).—Afr. trop.
affinis, Pers. Syn. i. 45 = cuspidatus.
aghullensis, *Echl. Top. Vers.* 41.—Afr. austr.
alatus, Jacq. Ic. ii. 2. t. 259 = orchidiflorus.
alatus, *Linn. Pl. Rar. Afr.* 8; *Amoen. Acad.* vi. 2.—Afr. austr.
albidus, Jacq. Ic. ii. 5. t. 256; Coll. Suppl. 21 = blandus.
alepicus, Boiss. Diagn. Ser. I. xiii. 13 = atroviolaceus.
aletroides, Vahl, Enum. ii. 96 = Watsonia aletroides.
algoensis, Sweet, Hort. Brit. ed. I. 207 = alatus.
alopecuroides, Linn. Cent. Pl. ii. 8; Amoen. Acad. iv. 301 = Watsonia plantaginea.
alopecuroides, Pers. Syn. i. 46 = Watsonia spicata.
alpigenus, C. Koch, in Linnaea, xxi. (1848) 685 = illyricus?
amabilis, Salisb. Prod. 41 = Lapeyronsia juncea.
ambiguus, Roem. & Schult. Syst. i. 446 = hirsutus.
amoenus, Ehrh. ex Roem. & Schult. i. c. 485 = Tritonia rosea.
amoenus, Herb. Banks, ex Roem. & Schult. i. c. 409 = brevifolius.
amoenus, Salisb. Prod. 39 = Watsonia Meriana.
anceps, Linn. ex Baker, in Journ. Linn. Soc. xvi. (1877) 156 = Lapeyronsia Fabricii.
anceps, Linn. f. Suppl. 94, partim = Lapeyronsia comprecsa.
andongensis, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 269.—Afr. trop.

GLADIOLUS:—
Andrewsii, *Klatt, in Abh Naturf. Ges. Halle*, xii. (1882) 340.—Afr. austr.
angolensis, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 269.—Afr. trop.
angustifolius, Lam. Illustr. i. 119 = Babiana tubiflora.
angustifolius, *Regel, Ind. Sem. Hort. Petrop.* (1868) 30.—Hab.?
angustifolius, Salisb. Prod. 40 = angustus.
angustus, Herb. Linn. ex Baker, in Journ. Linn. Soc. xvi. (1877) 175, 176 = blandus, undulatus.
angustus, Jacq. ex Thunb. Fl. Cap. i. 182 = hastatus.
angustus, *Linn. Sp. Pl.* 37.—Afr. austr.
angustus, Thunb. Dis. Glad. 19 = undulatus.
Antholyza, Poir. Encyc. Suppl. ii. 795 = Antholyza nervosa.
aphyllus, Ker-Gawl. Irid. Gen. 184 = brevifolius.
arcuatus, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 338.—Afr. austr.
arenarius, *Baker, in Journ. Linn. Soc.* xvi. (1877) 178. —Afr. austr.
aristatus, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 206.—Afr. austr.
atropurpureus, *Baker, in Journ. Bot.* xiv. (1876) 336. —Afric.
atroviolaceus, *Boiss. Diagn.* Ser. I. xiii. 14.—Syria; Persia.
aurantiacus, *Klatt, in Linnaea*, xxxv. (1867–68) 378.—Afr. austr.
bengalensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 268.—Afr. trop.
bicolor, *Baker, in Journ. Linn. Soc.* xvi. (1877) 178. —Afr. austr.
bicolor, Thunb. Dias. Glad. 16 = Synnotia bicolor.
bifforus, *Klatt, in Trans. S. Afr. Phil. Soc.* iii. ii. (1885) 197.—Afr. austr.
biflorus, Roem. & Schult. Syst. i. 416 = birustus.
biflorus, Thunb. Dias. Glad. 10 = Solenanthus biflorus.
bimaculatus, Lam. Encyc. ii. 727 = involutus.
binervus, Sweet, Hort. Brit. ed. I. 397 = grandis.
blandus, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 64.— Afr. austr.
bockveldensis, *Herb. ex Sweet, Hort. Brit.* ed. II. 501, nomen.—Afr. austr.
Bornati, Ardoino, Fl. Alpes Marit. 363 = segetum.
Boucheanus, Schlecht. in Linnaea, vii. (1832) 488 = palustris.
Bowmanus × , Hort.—Cf. Gard. Chron. (1866) 1197, 1247.
brachyandrus, *Baker, in Bot. Mag.* t. 6463.—Afr. trop.
bracteatus, Thunb. Prod. Pl. Cap. 186 = Lapeyrousia fissifolia.
bractsolatus, Lam. Encyc. ii. 725 = Watsonia punctata.
brenchleyensis × , Hort. ex Baker, in Journ. Linn. Soc. xvi. (1877) 177.
brevicaulis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 267.—Afr. trop.
brevicollis, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 339.—Afr. austr.
brevifolius, *Jacq. Coll.* iv. 156.—Afr. austr.
Breynianus, Ker-Gawl. Irid. Gen. 185 = recurvus.
Burmanni, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 204 = Ixia Burmanni.
byzantinus, Bieb. Fl. Taur. Cauc. iii. 40 = segetum.
byzantinus, Cos. ex Ball, in Journ. Linn. Soc. xvi. (1878) 674 = illyricus.
byzantinus, *Mill. Gard. Dict.* ed. VIII. n. 3.—Reg. Mediterr.
caerulescens, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 267.—Afr. trop.
campanulatus, Andr. Bot. Rep. t. 188 = blandus.
canditus × , Hort. ex Baker, in Journ. Linn. Soc. xvi. (1877) 177.
capitatus, Linn. Sp. Pl. 37 = Aristea capitata.
cardinalis, *Curt. Bot. Mag.* t. 135.—Afr. austr.
cardinalis. Red. ex Steud. Nom. ed. II. i. 685 = Tritonia crocata.
carinatus, [Soland. in] Ait. Hort. Kew. ed. I. i. 64 = recurvus.
carneus, Andr. Bot. Rep. t. 240 = brevifolius.
carneus, *Burm. f. Pl. Cap. Prod.* 2.—Afr. austr.
carneus, Delar. Dis. 20 = blandus.
carneus, Herb. Banks, ex Roem. & Schult. Syst. i. 415 = birustus.
carneus, Jacq. Coll. ii. 370 = cuspidatus.
carneus, Klatt, in Linnaea, xxxii. (1863) 723 = Ecklonl.
GLADIOLUS:—
caryophyllaceus, Poir. Encyc. Suppl. ii. 795 = Watsonia humilis.
caryophyllens, Houtt. ex Roem. & Schult. Syst. i. 431, in syn. = Watsonia brevifolia.
caucasicus, Herb. in Lindl. Bot. Reg. (1842) Misc. 65 = segetum.
citrinus, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 340 = trichonemifolius.
coccineus, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 197 = Ixia speciosa, patens.
cochleatus, Baker, in Journ. Bot. xiv. (1876) 333 = unguiculatus.
cochleatus, *Sweet, Brit. Flow. Gard.* Ser. II. t. 140.— Afr. austr.
coerulescens, Baker, in Journ. Linn. Soc. xvi. (1877) 174, var. caerulescens.
colinus, Salisb. Prod. 40 = communis.
Coleilli × , Sweet, Brit. Flow. Gard. t. 155.
communis, Cav. Desc. 295 = illyricus.
communis, Krock. Fl. Siles. i. 50 = imbricatus.
communis, *Linn. Sp. Pl.* 36, partim.—Europ.; Oriens.
communis, Linn. l. c. partim = segetum.
communis, Thunb. Prod. Pl. Cap. 8 = carneus.
communis, Vahl, Enum. ii. 80, partim = byzantinus.
communitatus, Bouché, in Linnaea, xii. (1838) 484 = segetum.
concolor, Salisb. Parad. Lond. t. 8 = triatis.
Cooperi, Baker, in Hook. f. Bot. Mag. t. 6202 = psittacinus.
cordatus, Thunb. Prod. Pl. Cap. 185 = angustus.
corneus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 155. —Afr. trop.
corymbosus, *Burm. f. Fl. Cap. Prod.* 2.—Afr. austr.
cosmosus, *Bot. Top. Vers.* 88.—Afr. austr.
crassifolius, *Baker, in Journ. Bot.* xiv. (1876) 334.—Afr. austr.
crispiflorus, Herb. in Lindl. Bot. Reg. (1842) Misc. 65 = imbricatus.
crispus, Linn. f. Suppl. 94 = Tritonia crispa.
cristatus, Trew, Pl. Rar. 7. t. 24. f. 1 (Tritoniae sp.). —Afr. austr.?
crocatus, Pers. Syn. i. 44 = Tritonia crocata.
cruentus, *T. Moore, in Gard. Chron.* (1868) 1138.— Afr. austr.
Cunonia, Gaertn. Fruct. i. 31 = Antholyza Cunonia.
cuspidatus, *Jacq. Ic.* ii. 5. t. 257; *Coll. Suppl.* 17.— Afr. austr.
Daleni, *Van.-Geel, Sert. Bot.* ii. (1829) t. 18.—Afr. austr.
dalmaticus, Tausch, ex Reichb. Fl. Germ. Excurs. Add. p. lx. n. 576 B = segetum.
debilis, *Ker-Gawl. in Bot. Mag.* t. 2585.—Afr. austr.
decoratus, *Baker, in Journ. Bot.* xiv. (1876) 333.— Afr. trop.
denticulatus, Lam. Illustr. 118; Encyc. ii. 728 = Lapeyrousia Fabricii.
dichotomus, Thunb. Dias. Glad. 10 = permeabilis.
discolor, *Schrank, in Denkschr. Bot. Regensb.* ii. (1822) 207.—Afr. austr.
distichus, Roem. & Schult. Syst. i. 440 = Babiana disticha
dracocephalus, *Hook. f. Bot. Mag.* t. 5864.—Afr. austr.
Dregei, *Klatt, in Linnaea*, xxxii. (1863) 693.—Afr. austr.
bulbus, *Schl. Top. Vers.* 41, nomen.—Afr. austr.
dubius, Guss. Fl. Sic. Prod. Suppl. 8 = illyricus.
dubius, Parl. Fl. Panorm. i. 44 = spathaceus.
Ecklonii, *Lehm. in Del. S. m. Hort. Hamb.* (1835); *Ann. Sc. Nat.* Sér. II. vi. (1836) 107.—Afr. austr.
edulis, *Burch. ex Ker-Gawl. in Bot. Reg.* t. 169.—Afr. austr.
elatus, *Bulb. Cat. Hort. Taurin.* (1813) 98.—Afr. austr.
elongatus, Salisb. Prod. 40 = Babiana tubiflora.
elongatus, Thunb. Prod. Pl. Cap. 185 = grandis.
equitans, Thunb. l. c. 186 = alatus.
excelsus, Ker-Gawl. in Bot. Mag. t. 1665 = blandus.
excisus, Jacq. Hort. Schoenb. iv. 45 = Lapeyrousia juncea.
expallescens, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 210.—Afr. austr.
exscapus, Thunb. Prod. Pl. Cap. 184 = Acidanthera tubulosa.
Fabricii, Thunb. l. c. 186 = Lapeyrousia Fabricii.
falcatus, Linn. f. Suppl. 96 = Lapeyrousia Fabricii.
fasciatus, Roem. & Schult. Syst. i. 429 = vittatus.
festivus, Herb. in Bot. Reg. (1844) Misc. 89 = brevifolius.
GLADIOLUS:—
fissifolius, Jacq. Coll. iv. 164 = Lapeyrousia fissifolia.
fistulosus, Jacq. Hort. Schoenb. i. 8 = Watsonia spicata
flabellifer, *Tausch, in Flora*, xix. (1836) 421.—Afr austr.
flavus, [Soland. in] Ait. Hort. Kew. ed. I. i. 65 = Tritonia flava.
flexuosus, Linn. f. Suppl. 96 = Acidanthera tubulosa.
flexuosus, Thunb. Prod. Fl. Cap. 8 = Acidanthera flexuosa.
floribundus, Hort. Batav. ex Tausch, in Flora, xix (1836) 422 = flabellifer.
floribundus, *Jacq. Coll.* iv. 162.—Afr. austr.
formosus, *Klatt, in Linnaea*, xxxii. (1863) 692.—Afr austr.
formosus, Pers. Syn. i. 44 = striatus.
fragrans × , Herb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 177.
fragrans, Jacq. Hort. Schoenb. i. 7 = Babiana plicata.
galeatus, Burm. f. Fl. Cap. Prod. 2 = alatus.
galeatus, Jacq. Coll. iv. 157 = Synnotia galeata.
galicienis, Bess. Prim. Fl. Galic. i. 51 = imbricatus.
gandavensis × , Van Houtte, Cat. (1844); et Fl. des Serres (1846), t. 1.
Gandavi, Herb. in Journ. Hort. Soc. ii. (1847) 89 = prsec.
Garnieri, *Klatt, in Linnaea*, xxxvii. (1871–73) 511.— Madag.
Gawleri, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 341.—Afr. austr.
gisnaucaus, Thunb. Prod. Pl. Cap. 186 = Watsonia rosea.
gracilis, *Jacq. Coll.* iv. 159.—Afr. austr.
gracilis, Lichtst. ex Schult. Mant. i. 290 = scaber.
gramineus, Linn. f. Suppl. 95; Thunb. Dias. Glad. 22 = Melasphaerula graminea.
grandiflorus, Andr. Bot. Rep. t. 118 = floribundus.
grandis, *Thunb. Prod. Pl. Cap.* 185.—Afr. austr.
gregarius, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 268.—Afr. trop.
Guernsii, Kunze, in Linnaea, xx. (1847) 14 = blandus.
Guepini, Koch, in Flora, xxiii. (1840) 486 = segetum.
halophilus, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 14.— Graecia.
hastatus, *Thunb. Prod. Pl. Cap.* 185.—Afr. austr.
Haylockianus × , Herb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 177.
Herbertianus × , G. Don, in Sweet, Hort. Brit. ed. III. 666.
hirsutus, *Jacq. Coll.* iv. 161.—Afr. austr.
hirtus, Steud. Nom. ed. II. i. 686, in syn. sphalm. = prsec.
hmnilis, *Stapf, in Denkschr. Acad. Wien*, i. (1885) 81 —Mesopot.
hyalinus, *Jacq. Ic. Rar.* ii. t. 242.—Afr. austr.
hygrophilus, Boiss. & Huet, ex Baker, in Journ. Linn. Soc. xvi. (1877) 171 = imbricatus.
ignescens, *Boj. ex Baker, in Journ. Bot.* xiv. (1876) 334.—Madag.
illyricus, *Koch, ex Sturm, Deutschl. Fl.* Heft 83; *et Syn. Fl. Germ.* ed. II. ii. 606.—Reg. Mediterr. et Cancas.
imbricatus, *Linn. Sp. Pl.* 37.—Europ.; Orions.
imbricatus, Reut. ex Willk. & Lange, Prod. Fl. Hisp. i. 340 = illyricus.
inarimensis, Guss. Enum. Pl. Inar. 226. t. 14 = segetum.
incarnatus × , Hort. ex Baker, in Journ. Linn. Soc. xvi. (1877) 177.
inclinatus, [DC. in] Red. Lil. i. t. 44 = Babiana tubiflora.
indicus, Mill. Gard. Dict. ed. VIII. n. 4 = Ferraria undulata.
infestus, Blanco, in Att. Acc. Gioen. xix. (1842) 98 = segetum.
inflatus, *Thunb. Prod. Pl. Cap.* 185.—Afr. austr.
infundibuliformis, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 198 = Watsonia Meriana.
insignis × , Paxt. Mag. Bot. vii. (1840) 238.
involutus, *Burm. f. Fl. Cap. Prod.* 2.—Afr. austr.
iridifolius, Jacq. Coll. iv. 151 = Watsonia Meriana.
italicus, Gaud. Fl. Helvet. i. 96 = segetum.
italicus, Mill. Gard. Dict. ed. VIII. n. 2 = communis.
ixioides, Thunb. Fl. Cap. 208 = Tritonia rosea.
ixioides, Thunb. l. c. ed. II. 92 = Ixia paniculata.
Jacquini, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 198.—Afr. austr.
juncea, Burm. f. Fl. Cap. Prod. 2; Linn. f. Suppl. 94 = Lapeyrousia juncea.
Kotschyanus, *Boiss. Diagn.* Ser. I. xiii. 15.—Mesopot.
lacustus, Jacq. Coll. iv. 167 = Watsonia humilis.

GLADIOLUS:—
laccatus, Thunb. Prod. Pl. Cap. 186 = villosus.
lacteatus, Berm. f. Fl. Cap. Prod. 2 = Tritonia crispa.
lavis, Thunb. Prod. Pl. Cap. 184 = grandis.
Lamarckii, Roem. & Schult. Syst. I. 445 = villosus.
Lamida, *Klatt*, in Linnæa, xxxii. (1863) 708.—Afr. austr.
latifolius, Lam. Illustr. I. 49 = Babiana obtusifolia.
lanciformis, *Baker*, in *Trans. Linn. Soc.* Ser. II. i. (1878) 268.—Afr. trop.
laxus, Thunb. Fl. Cap. ed. Schult. 50 (= *Meristostigma laxa*).—Afr. austr.
Lemonia, Pourr. ex Steud. Nom. ed. I. 869; II. i. 686 = blandus.
leucanthus, Bouché, in Linnæa, xii. (1838) 483 = segetum!
liliaceicus, Boiss. & Kotschy, Diagn. Ser. II. iv. 92 = fimbricatus.
lilaceus, Houtt. Handleid. xii 55 = augustus, gracilis.
linnæus, Salisb. Prod. 48 = Tritonia linnæta.
Lomenia, J. F. Gmel. Syst. 112 (= *Lomenia borbonica*).—Ins. Borbon.
longicollis, Baker, in Journ. Bot. xiv. (1876) 182 = Acidanthera platypetala.
longiflorus, Andr. Bot. Rep. t 5 = Babiana tubata.
longiflorus, Herb. Linn. ex Baker, in Journ. Linn. Soc xvi. (1877) 160 = Acidanthera tubulosa.
longiflorus, Jacq. Ic. Rar. t. 269 = Tritonia pallida.
longiflorus, Linn. f. Suppl. 96; Thunb. Diss. Glad. 19 = Ixia paniculata.
Ludovicæ, Jan, ex Bertol. in Nov. Comm. Bonon. i. (1834) 74, in obs. = segetum.
Ludwigii, *Poppe, ex Baker*, in *Journ. Linn. Soc.* xvi. (1878) 175.—Afr. austr.
lunulatus, *Klatt*, in *Abh. Naturf. Ges. Halle*, xii. (1882) 342.—Afr. austr.
luridus, Hornem. Hort. Hafn. Suppl. 113 = trichonemifolius!
luridus, *Welw. ex Baker*, in *Trans. Linn. Soc.* Ser. II. i. (1876) 267.—Afr. trop.
luteolus, *Klatt*, in *Peters, Reise Mossamb. Bot.* 515.—Afr. trop.
luteus, Klatt, in Linnæa, xxxii. (1863) 694, partim = bicolor.
luteus, *Lam. Encyc.* ii. 725.—Madag.
Mac-Owanianus, Klatt, in Trans. S. Afr. Soc. iii. (1885) 139 = augustus.
maculatus, Sweet, Hort. Brit. ed. I. 397 = recurvus.
marginatus, Linn. f. Suppl. 95 = Watsonia marginata, humilis.
marmoratus, Lam. Encyc. ii. 727 = Lapeyrousia juncea.
marmoratus, Tausch, in Flora, xix. (1836) 421.—Afr. austr.
Marschallii, Poir. Encyc. Suppl. ii. 789 = imbricatus.
Melleri, *Baker*, in *Journ. Bot.* xiv. (1876) 334.—Afr. trop.
Merianellus, Thunb. Diss. Glad. 14 = Antholyza Merianella.
Merianus, Jacq. Coll. iv. 150 = Watsonia Meriana.
Merianus, Thunb. Prod. Fl. Cap. 7 = Watsonia alsteroides.
Milleri, *Ker-Gawl. in Bot. Mag.* t. 632.—Afr. austr.
ministus, *Eckl. Top. Verz.* 40.—Afr. austr.
minutiflorus, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1832) 196 = Watsonia plantaginea.
mitchamiensis x, Herb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 177.
mollis, *Vahl, Enum.* ii. 119.—Afr. austr.
monostachyus, Roem. & Schult. Syst. i. 418, adnot. = hastatus, etc.
montanus, *Linn. f. Suppl.* 95.—Afr. austr.
Mortieuanus, Steud. Nom. ed. II. i. 686 = ecq.
Mortonius, Herb. in Bot. Mag. t. 3680 = blandus.
mucronatus, [DC. in] Red. Liliac. iii. t. 142 = Babiana striata.
mucronatus, Jacq. Coll. iv. 169 = Babiana mucronata.
mucronatus, Lam. Illustr. i. 119 = hirsutus.
multiflorus, *Baker*, in *Trans. Linn. Soc.* Ser. II. i. (1876) 269.—Afr. trop.
nanoguensis, Ker-Gawl. in Bot. Mag. t. 593 = alatus.
nanus, Andr. Bot. Rep. t. 137 = Babiana nana.
natalensis, [Reinw.] Hort. Lugd. Bat. ex Lodd. Bot. Cab. t. 1786 = psittacinus.
neglectus, Schult. Obs. Bot. 14 = palustris.
nervosus, Lam. Encyc. ii. 724 = Babiana stricta.
Nervii, *Baker*, in *Journ. Bot.* xiv. (1876) 334.—Afr. trop.
Netterâtii, Pers. Fl. Ital. iii. 239 = communis.

GLADIOLUS:—
ochroleucus, *Baker*, in *Journ. Bot.* xiv. (1876) 182.—Afr. austr.
odorus, Salisb. Prod. 40 = recurvus.
odorus, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1832) 206 = Iris fragrans.
oppositiflorus, *Herb. in Bot. Reg.* (1842) *Misc.* 98.—Madag.
orchidiflorus, *Andr. Bot. Rep.* t. 241.—Afr. austr.
orchidiflorus, Pers. Syn. i. 44 = ixioides.
ornatus, Klatt, in Trans. S. Afr. Phil. Soc. iii. (1885) 198 = inflatus.
Orubaschæ, [DC. in] Red. Lil. t. 125 = brevifolius.
palæaceus, *Vahl, Enum.* ii. 118.—Afr. austr.
pallidus, Bouché, in Linnæa, xii. (1838) 482 = segetum!
palustris, *Jaud. Fl. Evlv.* i. 97.—Europ.
paniculatus, Pers. Syn. i. 45 = Lapeyrousia juncea.
Papilio, *Hook. f. Bot. Mag.* t. 7065.—Afr. austr.
papilionaceus, Lichtst. ex Roem. & Schult. Syst. I. 408 = alatus.
papilionaceus, Salisb. Prod. 41 = Antholyza Cunonia.
parviflorus, Jacq. Obs. Bot. iv. 2. t. 76 = montanus.
parviflorus, Schur, ex Nym. Comp. 703 = imbricatus.
parviflorus, Vahl, Enum. ii. 79 = Watsonia Lamarkii.
pauciflorus, Berdau, Fl. Cracov. 841 = imbricatus.
pectinatus, Soland. ex Baker, in Journ. Linn. Soc. xvi. (1877) 168 = Tritonia crispa.
pendulus, *Mund. Eckl. ex Steud. Nom.* ed. II. i. 687 (Quid?).—Afr. austr.
permeabilis, *Burm. f. Fl. Cap. Prod.* 2; *Delar. Diss.* 27. t. 2.—Afr. austr.
persicus, *Boiss. Diagn.* Ser. I. vii. 102.—Persia.
persicus, Boiss. & Hoet, ex Boiss. Fl. Orient. v. 142 = atroviolaceus.
pictus, Sweet, Hort. Brit. ed. I. 397 = blandus.
pilosus, Eckl. Top. Verz. 38 = villosus.
plantagineus, Pers. Syn. i. 46 = Watsonia plantaginea.
plicatus, Jacq. Ic. 2. t. 287 = Babiana distichia.
plicatus, Linn. Sp. Pl. ed. II. 58 = Babiana stricta.
plicatus, Thunb. Diss. Glad. 20 = Babiana plicata.
polystachyus, Andr. Bot. Rep. t. 66 = Lapeyrousia juncea.
polystachyus, Thunb. Diss. Glad. 24 = Babiana plicata.
Pottsii, McNab, ex Baker, in Gard. Chron. (1877) ii. 424 = Tritonia Pottsii.
præcox, Eckl. Top. Verz. 38 = Antholyza revolutum.
prætensis, A. Dietr. Fl. Borus. i. 48 = palustris.
propinquus x, Herb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 177.
psittacinus, *Hook. Bot. Mag.* t. 3032.—Afr. austr.
pterophyllus, Pers. Syn. i. 48 = grandis.
puberulus, *Baker*, in *Journ. Bot.* xiv. (1876) 333.—Afr. austr.
pudibundus x, Sweet, Brit. Flow. Gard. Ser. II. t. 176.
pulchellus, *Klatt*, in *Linnæa*, xxxii. (1863) 693.—Afr. austr.
pulchellus, *Salisb. Prod.* 39.—Hab.?
punctatus, Jacq. Coll. iv. 160 = recurvus.
punctatus, Roem. & Schult. Syst. i. 425 = Watsonia punctata.
punctatus, *Thunb. Fl. Cap. ed. Schult.* 44.—Afr. austr.
punctulatus, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1832) 216 = villosus!
punicus, Lam. Encyc. ii. 727 = villosus!
punicus, Vahl, Enum. ii. 114 = Babiana stricta.
purpureo-auratus, *Hook. f. Bot. Mag.* t. 5944.—Afr. austr.
purpureus, Vahl, Enum. ii. 114 = Babiana villosa.
pygmæus, Roem. & Schult. Syst. i. 441, adnot. = Babiana sulphurea.
pyramidalis, Burm. f. Fl. Cap. Prod. 2 = Antholyza aethiopica.
pyramidalis, Lam. Illustr. i. 118; Encyc. ii. 726 = Watsonia iridifolia.
pyramidatus, Andr. Bot. Rep. t. 325 = Watsonia rosea.
quadrangularis, Ker-Gawl in Bot. Mag. t. 567 = Antholyza quadrangularis.
Quartinianus, *A. Rich. Tent. Fl. Abyss.* ii. 306.—Abyss.
rachidiflorus, *Klatt*, in *Abh. Naturf. Ges. Halle*, xii. (1882) 339.—Afr. austr.
Raddeanus, Trautv. in Act. Hort. Petrop. iii. 11. (1875) 373 = imbricatus.
ramosus, Linn. Sp. Pl. 37 = Melasphærula graminea.
ramosus, Murr. Syst. ed. XIV. 86 = Moræa ramosa.

GLADIOLUS:—
ramosus x, Pant. Mag. Bot. vi. (1839) 99.
recurvatus, Baker, in Journ. Linn. Soc. xvi. (1877) 96. (err. typ.) = recurvus.
recurvus, Houtt. Handleid. xii. 49. t. 79. f. 1 = Antholyza revoluta.
recurvus, *Linn. Mant.* i. 28.—Afr. austr.
recurvus, Thunb. Diss. Glad. 9 = Hesperantha radiata.
recurvus, Willd. Enum. Hort. Berol. Suppl. 4 = gracilis.
reflexus, Lichtst. ex Roem. & Schult. Syst. i. 439 = Babiana plicata.
refractus, Jacq. Ic. ii. 4. t. 341; Coll. Suppl. 26 = Freesia refracta.
resupinatus, Pers. Syn. i. 45 = Freesia refracta.
Reuteri, Boiss. Pugill. Pl. Kar. 112 = Illyricus.
rigidus x, Herb. ex Baker, in Journ. Linn. Soc. xvi. (1879) 177.
ringens, Andr. Bot. Rep. t. 27. et t. 227 = recurvus.
ringens, Eckl. Top. Verz. 37 = inflatus.
ringens, Thunb. Prod. Pl. Cap. 186 = Babiana mucronata, sulphurea.
rosco-albus, Jacq. Hort. Schoenb. i. 7 = Watsonia Meriana.
roseus, Andr. Bot. Rep. t. 11 = hirsutus.
roseus, Jacq. Ic. ii. 5. t. 261; Coll. Suppl 22 = Tritonia rosea.
roseus, Willd. Enum. Hort. Berol. 60 = Ixia amœna.
rosiacus, Pers. Syn. i. 46 = imbricatus.
rubens, Vahl, Enum. ii. 98 = Watsonia punctata.
rubrocyanus, Vahl, l. c. = Babiana stricta.
rubromarginatus, Schrad. ex Roem. & Schult. Syst. i. 415 = hirsutus.
rudis, *Lichtst. ex Roem. & Schult. l. c.* 408.—Afr. austr.
sagittiflor, Salisb. Prod. 41 = blandus!
saltatorum, Baker, in Trans. Linn. Soc. xxix. (1875) 155 = Quartinianus!
sambucinus, Jacq. Hort. Schoenb. i. 7 = Babiana sambucina.
Saundersii, *Hook. f. Bot. Mag.* t. 5873.—Afr. austr.
scaber, *Spreng. & Link, Hahrb.* i. iii. 70.—Afr. austr.
Schimperianus, Steud. ex Baker, in Journ. Linn. Soc. xvi. (1877) 176 = Quartinianus.
secundus, Thunb. Prod. Pl. Cap. 186 = Babiana secunda.
securiger, [Soland. in] Ait. Hort. Kew. ed. I. i. 65 = Tritonia securigera.
segetalis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = ecq.
segetum, *Ker-Gawl. in Bot. Mag.* t. 719.—Reg. Mediterr.
sericeo-villosus, *Hook. f. Bot. Mag.* t. 5427.—Afr. austr.
serotinus, Welw. ex Boiss. & Reut. Pugill. Pl. Nov. 113 = illyricus.
setifolius, Eckl. Top. Verz. 37 = gracilis.
setifolius, Linn. f. Suppl. 96 = Lapeyrousia juncea.
silenoides, Jacq. Coll. iv. 168 = Lapeyrousia silenoides. similis, *Eckl. Top. Verz.* 40.—Afr. austr.
Sparmanni, Thunb. Fl. Cap. ed. Schult. 49 = Freesia refracta.
spathaceus, Linn. f. Suppl. 96 = Babiana spathacea.
spathaceus, *Parl. Fl. Ital.* iii. 262.—Ital.
speciosus, Eckl. Top. Verz. 41 = cardinalis.
speciosus, Thunb. Fl. Cap. 186 = alatus.
spicatus, *Klatt*, in *Linnæa*, xxxv. (1867-68) 377.—Afr. trop.
spicatus, Lam. Illustr. i. 120 = Watsonia Lamarkii.
spicatus, Linn. Sp. Pl. 37 = Watsonia spicata.
spilanthus, Spreng. ex Baker, in Journ. Linn. Soc. xvi. (1877) 173 = gracilis.
spiralis, Pers. Syn. i. 48 = tristis.
splendens, *Baker*, in *Journ. Bot.* xiv. (1876) 333.—Afr. austr.
splendens, Herb. in Lindl. Bot. Reg. (1843) Misc. 46 = Antholyza caffra.
splendens, Waiw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1876) 268 = Welwitschii.
spofforthianus x, Herb. ex Baker, in Journ. Linn Soc. xvi. (1879) 177.
Sprengelianus, Schult. Mant. i. 296 = Watsonia stricta.
stenophyllus, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1832) 212 = Babiana plicata.
stolonifcrus, Salisb. Prod. 41 = Antholyza æthiopica.
striatus, Andr. Bot. Rep. t. 111 = undulatus.
striatus, Herb. Banks, ex Roem. & Schult. Syst. i. 435 = Watsonia rosea.
striatus, *Jacq. Ic.* ii. 5. t. 260; *Coll. Suppl.* 28.—Afr. austr.
striatus, Soland. ex Baker, in Journ. Bot. xiv. (1876) 337 = Tritonia Bakeri.

GLADIOLUS:—
stricti̇florus, [Delile, in] Red. Liliac. vii. t. 399 = Watsonia humilis.
strictus, Jacq. Coll. iv. 170 = hyalinus.
strictus, *Jacq. Ic.* t. 260.—Afr. austr.
strictus, [Soland. in] Ait. Hort. Kew. ed. I. i. 63 = Babiana stricta.
suaveolens, Ker-Gawl. Irid. Gen. 136 = recurvus ?
subbiflorus, Boiss. ex Tchihat. Asie Min. Bot. ii. 519 = imbricatus.
subulatus, Vahl, Enum. ii. 97 = Watsonia punctata.
sulcatus, Lam. Illustr. i. 119 = mollis.
sulphureus, Jacq. Coll. iii. 258 = Babiana stricta.
superbus, *Fisch. ex Stend. Nom.* ed. II. i. 687.—Hab.?
tabularis, *Eckl. Top. Vers.* 88.—Afr. austr.
tabularis, Pers. Syn. i. 44 = montanus.
telifer, Stokes, Bot. Comment. i. 217 = angustus.
Templemanii, Klatt, in Trans. S. Afr. Phil. Soc. iii. (1885) 196 = bicolor.
tenellus, *Jacq. Coll.* iii. 255.—Afr. austr.
tenuiflorus, C. Koch, in Linnæa, xxi. (1848) 636 = illyricus.
tenuis, *Baker, in Journ. Bot.* xiv. (1876) 385.—Afr. austr.
tenuis, Bieb. Fl. Taur. Cauc. i. 29 = imbricatus.
tenuis, Salzm. ex Nym. Consp. 704 = palustris.
testaceus, Vahl, Enum. ii. 105 = Watsonia brevifolia.
Thomsoni, *Baker, in Journ. Bot.* xix. (1881) 179.—Afr. trop.
Thunbergii, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 494 = Acidanthera tubulosa.
Thunbergii, Eckl. Top. Verz. 87 = hastatus.
tigrinus, Eckl. ex Baker, in Journ. Linn. Soc. xvi. (1877) 178 = hastatus.
trichonemifolius, *Ker-Gawl. in Bot. Mag.* t. 1468.—Afr. austr.
tricolor, *Steepf. in Denkschr. Acad. Wien*, i. (1885) 92.—As. Min.
trimaculatus, Lam. Encyc. ii. 727 = angustus.
triphyllus, *Sibth. & Sm. Fl. Græc. Prod.* i. 25; *Fl. Græc.* t. 28. t. 38.—Ins. Cyprus.
triphyllus, Bertol. Mant. Fl. Alp. Apuan. 5 = palustris.
tristis, Herb. Linn. ex Baker, in Journ. Linn. Soc. xvi. (1877) 172 = grandis.
tristis, *Linn. Sp. Pl.* ed. II. 58.—Afr. austr.
tristis, Thunb. Diss. Gled. 11 = tenellus, etc.
triticeus, Thunb. Fl. Cap. 194 = Watsonia plantaginea.
tubatus, Jacq. Ic. ii. 6. t. 264; Coll. Suppl. 18 = Babiana tubata.
tubiflorus, Linn. f. Suppl. 96 = Babiana tubiflora.
tubulosus, Burm. f. Pl. Cap. Prod. 2 = Watsonia spicata.
tubulosus, Jacq. Coll. iv. 153 = Watsonia aletroides.
umbellatus, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 202.—Afr. austr.
undularis, *Jacq. Coll.* iii. 256.—Afr. austr.
undulatus, Linn. Mant. 97, partim = cuspidatus.
undulatus, Schneev. Ic. t. 19 = vittatus.
unguicularis, *Baker, in Journ. Linn. Soc.* xvi. (1877) 178.—Afr. trop.
uniflorus, Klatt, in Abh. Naturf. Ges. Halle, xii. (1869) 237 = alatus.
venosus, Willd. Enum. Hort. Berol. i. 58 = Tritonia linaria.
ventricosus, Lam. Encyc. ii. 727 = cuspidatus.
vermiculor, Andr. Bot. Rep. t. 19 = grandis.
villosiflavus, Soland. ex Baker, in Journ. Linn. Soc. xvi. (1877) 174 = villosus.
villosulus, Roem. & Schult. Syst. i. 444 = Babiana stricta.
villosus, Burm. f. Fl. Cap. Prod. 2 = Synnotia bicolor.
villosus, *Ker-Gawl. Irid. Gen.* 138.—Afr. austr.
villosus, Vahl, Enum. ii. 145 = Babiana stricta.
viridis, Klatt, in Trans. S. Afr. Phil. Soc. iii. (1885) 199 = vittatus.
violaceus, Pers. Syn. i. 43 = recurvus.
viperatus, Ker-Gawl. in Bot. Mag. t. 688 = orchidiflorus.
virescens, Thunb. Fl. Cap. 196 = orchidiflorus.
viridis, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 451 = Tritonia viridis.
Wertheimi, *Hort. Hafn.* ii. 950.—Afr. austr.
vittatus, Zuccagni, in Roem. Collect. 121 = undulatus.
vomerculus, *Ker-Gawl. Irid. Gen.* 149.—Afr. trop.
watsonioides, *Baker, in Journ. Linn. Soc.* xvi. (1885) 405.—Afr. trop.

GLADIOLUS:—
Watsonius, Thunb. Diss. Glad. 14 = Antholyza revoluta.
Welwitschii, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 268.—Afr. trop.
Willmorianus ×, Hort. Fl. des Serres, t. 830.
xanthospilus, [DC. in] Red. Liliac. iii. t. 194 = Freesia refracta.
Zeyheri, *Eckl. Top. Vers.* 88, nomen.—Afr. austr.

GLANDONIA, Griseb. in Mart. Fl. Bras. xii. t. 28 (1858). *MALPIGHIACEAE*, Benth. & Hook. f. i. 259.
macrocarpa, *Griseb. l. c.*—Bras.

GLANDULA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 374 = **Astragalus**, Tourn. (Legumin.).
leontina, Medic. l. c. = A. leontinus.

GLANDULARIA, J. F. Gmel. Syst. 920 (1791) = **Verbena**, Linn.
bipinnatifida, Nutt. in Journ. Acad. Sc. Philad. ii. (1821) 123 = V. bipinnatifida.
caroliniensis, J. F. Gmel. Syst. 920 = V. Aubletia.

GLANDULIFOLIA, Wendl. Collect. 85. t. 10 (1808) = **Adenocalymma**, Willd. (Rutac.).
umbellata, Wendl. l. c. 87. t. 10 = A. umbellata.
uniflora, Wendl. l. c. 89. t. 38 = A. uniflora.

GLAPHIRIA, Spach, Hist. Vég. Phan. iv. 106 (1835) = seq.

GLAPHYRIA, Jack, in Trans. Linn. Soc. xiv. (1823) 128 = **Leptospermum**, Forst. (Myrtac.).
nitida, Jack, l. c.—Malaya.
serices, Jack, l. c. 129.—Sumatra.

GLASTARIA, Boiss. in Ann. Sc. Nat. Sér. II. xvi. (1841) 382 = **Texiera**, Jaub. & Spach (Crucif.).
dejeana, Boiss. l. c. = T. glastifolia.

GLASTUM, Rupp. Fl. Jen. ed. Hall. 63 (1745) = **Isatis**, Linn. (Crucif.).

GLAUCENA, Vitm. Summa, ii. 441 (1789) = **Clausena**, Burm. f. (Rutac.).

GLAUCIDIUM, Sieb. & Zucc. in Abh. Akad. Muench. iv. 11. (1845) 184 (Fam. Nat. Fl. Jap. i. 76. t. 1). *RANUNCULACEAE*, Benth. & Hook. f. i. 7. palmatum, *Sieb. & Zucc. Il. cc.*—Japon.

GLAUCIUM, Tourn. ex Hall. Enum. Helv. i. 304 (1742). *PAPAVERACEAE*, Benth. & Hook. f. i. 53.
alepplcum, *Boiss. & Heusch. ex Boiss. Fl. Orient.* i. 122.—Cilicia.
arabicum, *Fresen. in Mus. Senckenb.* i. (1838) 174. t. 10.—Arab.
aurantiacum, Martr. in Flora, xxxix. (1856) 171 = corniculatum.
aureum, C. Koch, in Linnæa, xix. (1847) 49 = corniculatum.
calycinum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 375.—Persia.
cappadocicum, *Boiss. l. c.*—Cappadoc.
contortuplicatum, *Boiss. l. c.* 376.—Persia.
corniculatum, *Curt. Fl. Lond.* vi. t. 23.—Europ.; Reg. Mediterr.
elegans, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. (1835) 29.—Persia.
fimbrilligerum, *Boiss. Fl. Orient.* i. 120.—Turkest.; Afghan.
Fischeri, *Bernh. ex Steud. Nom.* ed. II. i. 687.—Persia.
flavum, *Crantz, Stirp. Austr.* ed. I. fasc. ii. 121; ed. II. 128.—Europ.; Reg. Mediterr.
fulvum, Sm. Exot. Bot. i. 11. t. 7 = flavum.
glaucum, Moench, Meth. 249 = flavum.
grandiflorum, *Boiss. & Huet, Diagn.* Ser. II. v. 15.—As. Min.; Persia.
hybridum, Dum.-Cours. Bot. Calt. ed. II. iv. 472 = Roemeria violacea.
intermedium, Link, in Buch, Beschr. Canar. Ins. 152 = corniculatum.
lutescarpum, *Boiss. Fl. Orient.* i. 122.—As. Min.; Persia.

GLAUCIUM:—
leptopodum, *Masim. in Bull. Acad. Pétersb.* xxiii. (1877) 210.—China.
littorale, Salisb. Prod. 377 = flavum.
luteum, Scop. Fl. Carn. ed. II. i. 369 = flavum.
mesopotamicum, Spreng. Syst. ii. 570 = Roemeria violacea.
oxylobum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat Mosc.* xii. (1860) 12. t. 5.—Persia.
persicum, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 192 = fimbrilligerum.
persicum, *DC. Syst.* ii. 97.—Persia.
phœnicewm, Crantz, Stirp. Austr. ed. I. fasc. ii. 133; ed. II. 141 = corniculatum.
pumilum, Boiss. in Ann. Sc. Nat. Sér. II. xvi. (1841) 374 = elegans.
refractum, J. Gay, ex Boiss. Fl. Orient. i. 122 = alepplcum.
refractum, Stev. ex DC. Syst. ii. 98 = Roemeria refracta.
rubrum, Sibth. & Sm. Fl. Græc. Prod. i. 257 = corniculatum.
Serpieri, *Heldr. in Regel, Gartenfl.* (1878) 323.—Græcia.
spurium, Stev. ex DC. Syst. ii. 98 = Roemeria violacea.
squamigerum, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 192 = elegans.
squamigerum, *Ker. & Kiv. in Bull. Soc. Nat. Mosc.* xv. (1842) 141.—As. centr.
tenue, *Regel & Schmalh. ex Regel, Fl. Nov. Fedsch* 2.—Turkest.
tricolor, Bernh. ex Spreng. Syst. App. 208 = corniculatum.
tricolor, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 411 = luteum.
trivalve, Moench, Meth. 249 = Roemeria violacea.
violaceum, Juss. Gen. 286 = Roemeria violacea.
vitellinum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 11. t. 4.—Persia.

GLAUCOIDES, Rupp. Fl. Jen. ed. Hall. 20 (1745) = **Glaux**, Tourn. (Primul.).

GLAUX, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 376 = **Astragalus**, Tourn. (Legumin.).
astragaloides, Medic. l. c. 377 = A. Cicer.

GLAUX, [Tourn.] Linn. Syst. ed. I (1735). *PRIMULACEAE*, Benth. & Hook. f. ii. 687.
GLAUCOIDES, Rupp. Fl. Jen. ed. Hall. 20 (1745).
atacamensis, *Phil. Fl. Atac.* 36.—Chili.
maritima, *Linn. Sp. Pl.* 207.—Reg. bor. temp.

GLAYPHYRIA, G. Don, Gen. Syst. iii. 871 (1831) sphalm. = GLAPHYRIA, Jack = **Leptospermum**, Forst. (Myrtac.).

GLAZIOSTELMA, Fourn. in Mart. Fl. Bras. vi. IV 227 (1886). *ASCLEPIADEAE*.
ovalifolium, *Fourn. l. c.*—Bras.

GLAZIOVA, Bur. in Adansonia, viii. (1868) 380. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1038.
GLAZIOVIA, Benth. & Hook. f. ii. 1038 (1876).

GLAZIOVA, Mart. ex H. Wendl. in Flor. & Pom. (1871) 118 = **Cocos**, Linn. (Palm.).
elegantissima, Hort. ex H. Wendl. l. c. = C. Weddelliana.
insignis, Hort. ex Drude, in Mart. Fl. Bras. iii. ii. 398 = C. insignis.
Martiana, Glaz. ex Drude, l. c. 397 = C. Weddelliana.

GLAZIOVIA, Benth. & Hook. f. Gen. ii. 1038 (1876) = **Glaziova**, Bur. (Bignon.).

GLEBIONIS, Cass. in Dict. Sc. Nat. xli. 41 (1826) = **Chrysanthemum**, Tourn. (Compos.).
Ronburghii, Cass. l. c. = C. coronarium.
segetum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 90 = C. segetum.

GLECHOMA, Linn. Syst. ed. I (1735) = **Nepeta**, Linn. (Labiat.).
arvensis, Linn. Sp. Pl. 578 = Stachys arvensis.
bolgica, Linn. l. c. = Stachys arvensis.
borealis, Salisb. Prod. 81 = N. Glechoma.
orecta, Buch.-Ham. ex Roxb. Hort. Beng. 44; Fl. Ind. iii. 7 = N. ruderalis.

GLECHOMA :—
grandiflora, DC. Fl. Fr. iii. 538 = Stachys corsica.
hederacea, Linn. Sp. Pl. 578 = Nepeta Glechoma.
heterophylla, Opiz, Natural. vii. (1824) 61 = N. Glechoma.
hindostana, Heyne, ex Roth, Nov. Sp. Pl. 258 = N. ruderalis.
hirsuta, Waldst. & Kit. Pl. Rar. Hung. ii. 124. t. 119 = N. Glechoma.
indica, Spreng. Syst. ii. 737 = N. ruderalis.
intermedia, Schrad. ex Benth. in DC. Prod. xii. 391 = N. Glechoma.
lamiifolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 58 = N. Glechoma.
lobulata, Kit. in Linnæa, xxxii. (1863) 484 = N. Glechoma ?
longicaulis, Dulac, Fl. Hautes-Pyr. 400 = N. Glechoma.
magna, Mérat, Fl. Par. 225 = N. Glechoma.
Marrubiastrum, Vill. Hist. Pl. Dauph. ii. 371 = Stachys arvensis.
micrantha, Boenn. ex Reichb. Fl. Germ. Excurs. 316 = N. Glechoma.
nivalis, Jacquem. ex Benth. Lab. Gen. et Sp. 737 = N. thibetica.
repens, Gilib. Fl. Lituan. i. 77 = N. Glechoma.
rigida, Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 215 = N. Glechoma.
rotundifolia, Rafin. Atl. Journ. 177 = N. Glechoma.
thibetica, Jacquem. ex Benth. Lab. Gen. et Sp. 737 = N. thibetica.

GLECHON, Spreng. Syst. iv. Cur. Post. 227 (1827). *LABIATAE*, Benth. & Hook. f. ii. 1192.
candida, Benth. in DC. Prod. xii. 250 = marifolia.
canescens, St. Hil. ex Benth. Lab. Gen. et Sp. 407 = marifolia.
ciliata, *Benth. l. c.* 408.—Bras.
marifolia, *Benth. l. c.* 407.—Bras.
myrtoides, *St. Hil. ex Benth. l. c.* 409.—Bras.
origanifolia, *Benth. l. c.* 408.—Bras.
ringens, *Benth. l. c.* 409.—Bras.
serpyllifolia, St. Hil. ex Benth l. c. 407 = marifolia
spathulata, *Benth. l. c.* 406.—Bras.
squarrosa, *Benth. l. c.* 408.—Bras.
thymoides, *Spreng. Syst. iv. Cur. Post.* 227.—Bras.

GLECHONION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = **Nepeta**, Linn.
hederaceum, St. Lag. l. c. = N. Glechoma.

GLECOMA, Linn. Sp. Pl. 578 (1753) = GLECHOMA, Linn. = **Nepeta**, Lann. (Lablat.).

GLEDITSCHIA, Clayton, ex Linn. Gen. ed. II. 480 (1742) (Gleditsia) ; Scop. Introd. 295 (1777). *LEGUMINOSAE*, Benth. & Hook. f. i. 568.
MELILOBUS, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) 215 App.
africana, *Welw. ex Benth. in Trans. Linn. Soc.* xxv. (1866) 804.—Afr. trop.
aquatica, Marsh. Arbust. Am. 95 = monosperma.
brachycarpa, Pursh, Fl. Am. Sept. i. 221 = triacanthos.
Bujotii, Neumann, in Rev. Hortic. Sér. II. iv. (1845–46) 205 = triacanthos ?
caroliniensis, Lam. Encyc. ii. 461 = monosperma.
caspica, *Desf. Hist. Arb.* ii. 247.—Reg. Casp.
chinensis, Auct. ex Steud. Nom. ed. II. i. 568 = sinensis.
elegans, Salisb. Prod. 338 = triacanthos.
ferox, Desf. Hist. Arb. ii. 247 = macracantha ?
fera, Hort. ex C. Koch, Dendrol. i. 9 = triacanthos.
Fontanesii, Spach, Hist. Vég. Phan. i. 95 = macracantha.
heterophylla, *Bunge, in Mem. Soc. Etr. Pétersb.* ii. (1835) 95.—China.
heterophylla, Rafin. Fl. Ludov. 99 = triacanthos.
horrida, Willd. Sp. Pl. iv. 1098 = sinensis.
indica, Pers. Syn. ii. 625 = sinensis.
inermis, Linn. *Syst. ed.* X. 1213 = Callisandra Houstoni.
inermis, Mill. Gard. Dict. ed. VIII. n. 2 = monosperma.
japonica, Lodd. Cat. ex W. Bext. in Loud. Hort. Brit. Suppl. ii. 638 = sinensis.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 54.—Japon.
jamaicensis, Lam. Encyc. ii. 466 = Acacia jamaicensis.
laxeta, Hort. Par. ex G. Don, in Loud. Hort. Brit. 413 = triacanthos.
latisiliqua, Lodd. Cat. ex G. Don, l. c. 414 = triacanthos.
macracantha, *Desf. Hist. Arb.* ii. 246.—China.

GLEDITSCHIA :—
melioba, Walt. Fl Carol. 254 = triacanthos.
monosperma, *Walt. l. c.*—Am. bor.
orientalis, Boec. ex DC. Prod. ii. 479 = macracantha.
polysperma, Stokes, Bot. Mat. Med. i. 238 = triacanthos.
sinensis, *Lam. Encyc.* ii. 465.—China.
spinosa, Marsh. Arbust. Am. 94 = triacanthos
triacantha, Gaertn. Fruct. ii. t. 146 = monosperma.
triacanthos, *Linn. Sp. Pl.* 1056 (exel. var.)—Am. bor.
xylocarpa, Hance, in Journ. Bot. xxii. (1884) 366 = sinensis.

GLEDITSIA, Linn. Gen. ed. II. 480 (1742) = præc.

GLEDITZIA, Jaume St. Hil. Exposs. ii. 202 (1805) = præc.

GLEHNIA, F. Schmidt, in Mig. Ann. Mus. Bot. Lugd. Bat. iii. 61 (1867) = **Phellopterus**, Benth. (Umbellif.).
littoralis, F. Schmidt, l. c. = P. littoralis.

GLENNIEA, Hook f. in Benth. & Hook. f. Gen. i 404 (1862). *SAPINDACEAE*, Benth. & Hook. f. i. c.
unijuga, *Hook. f. l. c.*—Zeylan.
zeylanica, Hook. f. ex Thw. Enum. Pl. Zeyl. 408 = unijuga.

GLIA, Sond. in Harv. & Sond. Fl. Cap. ii. 547 (1861–62) = **Annesorhiza**, Cham. & Schlecht. (Umbellif.).
gummifera, Sond. l. c. 548.—Afr. austr.

GLICIRRHIZA, Nocca, in Usteri, Ann. Bot. v. (1798) 53 = **Glycyrrhiza**, Tourn. (Legumin.).

GLINUS, Linn. It. Loefl. 145 (1758) = **Mollugo**, Linn.
Astrolasion, Zipp. ex Span. in Linnæa, xv. (1841) 207, nomen.—Ins. Timor.
Cambessedesii, Fenzl, in Ann. Wien. Mus. i. (1836) 358 = M. Spergula.
chrystallinus, Forsk. Fl. Aegypt. Arab. 95 = Aizoon canariense.
crystallinus, Auct. vide *chrystallinus*.
denticulatus, Fenzl, in Ann. Wien. Mus. i. (1836) 361 = M. Spergula.
dictamnoides, Burm. f. Fl. Ind. 113 ; Lam. Encyc. ii. 729 = M. hirta.
lothoides, Cranta, Inst. i. 139 = seq.
lotoides, Loefl. It. Hisp. 145 ; Linn. Sp. Pl. ed. II. 668 = M. hirta.
micranthus, Boiss. Diagn. Ser. I. x. 11 = M. hirta.
Millus, Kaesach. Nom. ed. III. 141 = Gisekia Millus.
Mollugo, Fenzl, in Ann. Wien. Mus. i. (1836) 359 = M. Spergula.
mozambicensis, Spreng. Syst. ii. 467 = Gisekia Millus.
mucronatus, Klotzsch, in Peters, Reise Mossamb. Bot. 570. t. 25 = Orygia decumbens.
ononoides, Burm. f. Fl. Ind. 113. t. 36. f. 2.—Ind. or.
oryzoides, F. Muell. Pl. Vict. i. 203 = M. orygioides.
parviflorus, Wall. Cat. n. 1519 = M. hirta.
Pauli-Wilhelmi, Hochst. in Flora, xxvii. (1844) Beibl. 4 = M. Pauli-Wilhelmi.
radiatus, Rohrb. in Mart. Fl. Bras. xiv. ii. 238 = M. radiata.
setiflorus, Forsk. Fl. Aegypt. Arab. 95 = M. hirta.
Spergula, Steud. Nom. ed. II. i. 668 = M. Spergula.
trianthemoides, Heyne, in Roth, Nov. Pl. Sp. 231 = Orygia decumbens.

GLYRICIDIA, H. B. & K. Nov. Gen. et Sp. vi. 393, in nota (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 499.
cubensis, *Wright, in Sauv. Fl. Cub.* 26.—Cuba.
latifolia, *Griseb. Cat. Pl. Cub.* 70.—Cuba.
maculata, *H. B. & K. Nov. Gen. et Sp.* vi. 393, in nota.—Guatemala.
platycarpa, *Griseb. Cat. Pl. Cub.* 70.—Cuba.
sepium, H. B. & K. Nov. Gen. et Sp. vi. 393, in nota ? = Lonchocarpus sepium.

GLISCHROCARYON, Endl. in Ann. Wien. Mus. ii. (1839) 309 = **Leucodendron**, Lindl. (Halorag.).
Rodi, Endl. l. c. 310 = L. Rodi.

GLISCHROCOLLA, A. DC. in DC. Prod. xiv. 490 (1856) = **Endonemma**, A. Juss. (Pennæc.).
Lessertiana, A. DC. l. c.—Afr. austr. ?

GLISSANTHE, Steud. Nom. ed. II. i. 688 (1840) = GISSANTHE, Salisb. = **Costus**, Linn. (Scitam.).

GLOBBA, Linn. Mant. ii. 148. n. 1247 (1771), partim. *SCITAMINEAE*, Benth. & Hook. f. iii. 640.
CERATANTHERA, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 341.
COLEBROOKIA, D. Don, ex Lestib. l. c. (1841).
HUSA, Koen. in Retz. Obs. iii. 49 (1783).
MANITIA, Gieske, Prael. Linn. 209 (1792).
SPHAEROCARPUS, J. F. Gmel. Syst. 9 (1791).
alba, *Zipp. ex Span. in Linnæa*, xv. (1841) 479, nomen.—Ins. Timor.
albo-bracteata, *N. E. Br. in Gard. Chron.* (1882) ii. 71.—Sumatra.
aphylla, *Miq. Fl. Ind. Bat. Suppl.* 614.—Sumatra
atracanensis, *Kurz, in Journ. As. Soc. Beng.* xxxix (1870) ii. 88.—Ind. or.
atrosanguinea, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 22.—Borneo.
aurantiaca, *Miq. Fl. Ind. Bat. Suppl.* 613.—Sumatra
bracteata, Heyne, ex Wall. Cat. n. 6582 = marantina. bulbifera.
bractelata, Horan. Prod. Scitam. 20, in obs. (Quid ?). —Hab. ?
bractrosa, Horan. l. c. = bulbifera.
bulbifera, *Roxb. in As. Res* xi. (1810) 358.—Ind. or. ; Malaya.
Careyana, Roxb. Hort. Beng. 2 ; Fl. Ind. i. 80 = sessiliflora.
ciliata, *Jack, in Malay. Misc.* ii. (1822) n. VII. p. 5.—Malaya.
coccinea, Hort. Veitch.—Cf. Gard. Chron. (1882) ii. 74 = albo-bracteata.
crispa, *Reta. Obs.* iii. 50 = Amomum echinatum.
cuspidata, Nees & Mart. in Nov. Act. Nat. Cur. xi (1823) 28 (Quid ?).—Bras.
dubia, Horan. Prod. Scitam. 20 (Quid ?).—Hab. ?
erecta, [DC. in] Red. Lil. t. 174 = Alpinia calcarata.
expansa, Wall. Cat. n. 6536 = versicolor, bracteolata.
Finlaysoniana, *Wall. i. c.* n. 6585.—Ind. or.
Hasseltii, *Boerl. in Hasselt & Boerl. Bydr. Middl. Sum.* 84.—Sumatra.
Hura, Roxb. in As. Res. xi. (1810) 359 = versicolor.
japonica, *Thunb. Fl. Yap.* 28.—Japon.
Koenigiana, Spreng. Syst. iv. Cur. Post. 8 = versicolor leucantha, *Miq. Fl. Ind. Bat. Suppl.* 612.—Sumatra.
maculata, Blume, Enum. Pl. Jav. 53 = pendula ?
manifera, Sims, Bot. Mag. xxb t. 1920 = Mantisia saltatoria.
maniformis, Horan. Prod. Scitam. 20, sphalm. = præc.
marantina, *Linn. Mant.* ii. 170.—Ins. Molucc.
marantinoides, Wight, Ic. vi. 15. t. 2001 = bulbifera.
monifera, Steud. Nom. ed. II. i. 688, sphalm. = *manifera*.
multiflora, *Wall. Cat.* n. 6587.—Ind. or.
multiflora, Wall. l. c. partim = pendula.
nutans, Linn. Mant. ii. 170 = Alpinia nutans.
oligantha, *Miq. Fl. Ind. Bat. Suppl.* 613.—Sumatra.
ophioglossa, *Wight, Ic.* vi. 16. t. 2002.—Ind. or.
orixensis, *Roxb. in As. Res.* xi. (1810) 358. t. 6. —Ind. or.
orixensis, Wall. Cat. n. 6585 = racemosa, ophioglossa, sessiliflora.
paniculoides, *Miq. Fl. Ind Bat. Suppl.* 614.—Sumatra.
parviflora, *Presl, Rel. Haenk.* ii. 115.—Ins. Luzon.
patens, *Miq. Fl. Ind. Bat. Suppl.* 613.—Sumatra.
pendula, *Roxb. in As. Res.* xi. (1810) 359.—Ind. or.
purpurea, Andr. Bot. Rep. t. 117 = Mantisia saltatoria.
racemosa, *Sm. Exot. Bot.* ii. 115. t. 117.—Reg. Himal.
radicalis, Roxb. in As. Res. xi. (1810) 359 = Mantisia saltatoria.
saltatoria, Roxc. Mon. Pl. ii. t. 16 ; Salt. t. 112 = Mantisia saltatoria.
Schomburgkii, *Hook. f. Bot. Mag.* t. 6293.—Siam.
secunda, Horan. Prod. Scitam. 20 (Quid ?).—Hab. ?
sessiliflora, *Sims, Bot. Mag.* t. 1498.—Ind. or.
spathulata, Roxb. Hort. Beng. 2 ; Fl. Ind. i. 83 = Mantisia spathulata.
strobilifera, Zoll. & Mor. Syst. Verz. Zoll. 84 = marantina.
subsessilis, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 29 (Quid ?).—Bras.
subulata, Roxb. Hort. Beng. 2 ; Fl. Ind. i. 81 = Mantisia saltatoria.

GLOBBA:—
uliginosa, *Miq. Fl. Ind. Bat. Suppl.* 613.—Ins. Banca.
wenformis, Linn. Mant. ii. 171 = Alpinia uviformis.
wolatina, Horan. Prod. Scitam. 20 = multiflora.
verticolor, *Sm. Exot. Bot.* t. 117.—Ind. or.; Malaya.

GLOBERIA, Rafin. Med. Fl. ii. 84 (1830) = **Liriope**, Lour. (Hasmodor.).
autumnalis, Rafin. Fl. Tellur. iv. 18 = L. spicata.
spicata, Rafin. Med. Fl. ii. 84 = L. spicata.

GLOBERIS, Rafin. Fl. Tellur. iv. 18 (1836) = praec.

GLOBIFERA, J. F. Gmel. Syst. 32 (1791) = **Micranthemum**, Michx. (Scrophul.).
umbrosa, J. F. Gmel. l. c. = M. orbiculatum.

GLOBULARIA, Tourn. ex Linn. Gen. ed. I. 358 (1737). *SELAGINEAE*, Benth. & Hook. f. ii. 1130.
ABOLARIA, Neck. Elem. i. 105 (1790).
CARRADORIA, A. DC. Prod. xii. 610 (1864).
alpina, Salisb. Prod. 52 = nudicaulis.
Alypa, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 126 = G. Alypum.
Alypum, Delile, Fl. Egypt. 5 = arabica.
Alypum, *Linn. Sp. Pl.* 95.—Reg. Mediterr.
amygdalifolia, *Webb, in Hook. Niger Fl.* 183.—Ins. Cap. Verd.
Apkyllanthes, Cranta, Inst. i. 350 = vulgaris ?
arabica, *Jaub. & Spach, Illustr.* iii. 75. t. 260.—Arab.; Egypt.
bellidifolia, Salisb. Prod. 52 = spinosa.
bellidifolia, Tenore, Prod. Fl. Nap. p. xi = nana.
bisnagarica, Linn. Sp. Pl. 96 = vulgaris.
caespitosa, Orteg. ex A. DC. in DC. Prod. xii. 612 = vulgaris.
collina, Salisb. Prod. 52 = vulgaris.
cordifolia, *Linn. Sp. Pl.* 96.—Europ. austr.; Tauria.
elongata, Hegetschw. Fl. Schw. 1.14 = vulgaris.
ericophala, *Pomel, Nouv. Mat. Fl. Atl.* 111.—Alger.
glauca, Balb. ex Steud. Nom. ed. II. i. 688 = incanescens.
ilicifolia, Willk. Rech. Gleb. 26 = spinosa.
incanescens, *Viv. Fl. Ital. Fragm.* i. 2. t. 3.—Europ. austr.
linifolia, Lam. Encyc. ii. 731 = vulgaris.
Linnaei, Roxy, in Bull. Soc. Bot. Fr. xxix. (1882) 350 = vulgaris.
longifolia, Ait. Hort. Kew. i. 130 = salicina.
macrantha, C. Koch, ex Walp. Rep. iv. 175 = trichosantha.
minima, Vill. Hist. Pl. Dauph. ii. 208 = cordifolia.
multicaulis, Tenore, ex Nym. Consp. 608 = vulgaris.
nana, *Lam. Encyc.* ii. 731.—As. or.
nudicaulis, Herb. Lian. ex Rolfe, in Journ. Linn. Soc. xx. (1883) 342 = vulgaris.
nudicaulis, *Linn. Sp. Pl.* 97.—Europ.
orientalis, *Linn. f. c.*—As. Min.
pallida, C. Koch, in Linnaea, xxii. (1849) 654 = trichosantha.
procera, Salisb. Prod 52 = salicina.
punctata, Lapeyr. Hist. Abr. Fl. Pyren. 57 = incanescens.
pungens, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 894 = vulgaris.
repens, Lam. Fl. Fr. ii. 825 = nana.
salicina, *Lam. Encyc.* ii. 732.—Ins. Mader.
saxatilis, Salisb. Prod. 52 = cordifolia.
spinosa, Lam. Encyc. ii. 731 = vulgaris.
spinosa, *Linn. Sp. Pl.* 96.—Hispan.
stygia, *Orph. in Boiss. Diagn.* Ser. II. iv. 60.—Graecia.
tenella, *Lange, in Kjoeb. Vidensk. Meddel.* (1868) 1.—Hispan.
trichocalyx, Steud. Nom. ed. II. i. 689 = arabica.
trichosantha, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 36.—As. Min.; Syria.
Turkish, Willk. Rech. Gleb. 26, in syn. = G. Alypum.
valentina, Willk. l. c. 21 = vulgaris.
virgata, Salisb. Prod. 52 = G. Alypum.
vulgaris, *Linn. Sp. Pl.* 96.—Europ. austr.; Reg. Caucas.
Willhommii, Nym. Syll. 140 = vulgaris.

GLOBULEA, Haw. Syn. Pl. Succ. 60 (1812) = **Crassula**, Dill.
atropurpurea, Haw. in Phil. Mag. (Sept. 1824) 189 = C. portulacea.
canescens, Haw. Syn. Pl. Succ. 61 = C. canescens.
capitata, Haw. Rev. Pl. Succ. 17 = C. capitata.
clavifolia E. Mey. ex Harv. & Sond. Fl. Cap. ii. 360 = C. clavifolia.
cultrata, Haw. Syn. Pl. Succ. 60 = C. cultrata.
hispida, Haw. in Phil. Mag. (Sept. 1824) 191 = C. hispida.
impressa, Haw. l. c. 188 = C. impressa, *D. Dietr.*
Linguia, Haw. l. c. 190 = C. Lingua.
Linguia, Haw. l. c. = C. Lingula.
mesembryanthemoides, Haw. l. c. 189 = C. mesembryanthemoides.
mollis, Haw. l. c. 191 = C. mollis.
nudicaulis, Haw. Syn. Pl. Succ. 61 = C. nudicaulis.
obvallata, Haw. l. c. 60 = C. obvallata.
paniculata, Haw. in Phil. Mag. (1825) 29 = C. paniculata.
radicans, Haw. l. c. (1824) 189; (1825) 27 = C. radicans.
subincana, Haw. l. c. (Sept. 1824) 190 = C. subincana.
sulcata, Haw. Rev. Pl. Succ. 18 = C. sulcata.

GLOCHERIA, Pritz. Ic. Ind. ii. 134 (1855), *err. typ.* = **Glochoeria**, Neck. (Acanthac.).

GLOCHIDINOPSIS, Steud. Nom. ed. II. i. 689 (1840) = GLOCHIDIONOPSIS, Blume = sec.

GLOCHIDION, Forst. Char. Gen. 113. t. 57 (1776). *EUPHORBIACEAE*, [Benth. & Hook. *f.* iii. 272.]
BRADLEIA, Cav. Ic. iv. 48. t. 371 (1797).
BRADLEJA, Banks, ex Gaertn. Fruct. ii. 197. t. 109 (1791).
GLOCHIDIONOPSIS, Blume, Bijdr. 588 (1825).
GLOCHISANDRA, Wight, Ic. v. ii. 28. t. 1905 (1850).
GYMNOLA, A. Juss. in Mém. Mus. Par. x. (1823) 335; et Euph. Tent. 17. t. 8 (1824).
LOBOCARPUS, Wight & Arn. Prod. 7 (1834).
acuminatum, *Muell. Arg. in Linnaea*, xxxii. (1863) 68.—Reg. Himal.
acuminatum, *Muell. Arg. l. c.* 59.—N. Caled.
andamanicum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 238.—Ins. Andaman.
arboreum, *Blume, Bijdr.* 584.—Java.
arboreum, *Wight, Ic.* t. 1907.—Ind. or.
Arnottianum, *Muell. Arg. in Linnaea*, xxxii. (1863) 60.—China.
asperum, *Bedd. Forest. Man. Bot.* 193.—Ind. or.
baladense, *Muell. Arg. in Linnaea*, xxxii. (1863) 59.—N. Caled.
bancanense, Miq. Fl. Ind. Bat. Suppl. 450 = zeylanicum.
Billardieri, *Baill. Adansonia*, ii. (1861-62) 241.—N. Caled.
Blumeanum, Muell. Arg. in Linnaea, xxxii. (1863) 65 = glanciun, *Blume*.
Bourgeoisii, *Muell. Arg. l. c.* 70.—N. Caled.
brachyloburn, *Muell. Arg. l. c.* 62.—Zeylan.
bupleuroides, *Muell. Arg. l. c.* 59.—N. Caled.
cacuminum, *Muell. Arg. l. c.* 60.—Java.
calciadicum, *Muell. Arg. l. c.* 63.—N. Caled.
calocarpum, *Kurz, in Journ. Bot.* xiii. (1875) 330.—Ins. Nicob.; Andaman.
canarium, Bedd. Forest. Man. Bot. 193 = zeylanicum.
canitiosum, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 341.—China.
catastachum, *Muell. Arg. in Linnaea*, xxxii. (1863) 70.—N. Caled.
Chamaecrotes, *Muell. Arg. l. c.*—N. Caled.
cinerascens, *Miq. Fl. Ind. Bat. Suppl.* 451.—Sumatra.
coccineum, *Muell. Arg. in Linnaea*, xxxii. (1863) 60.—Burma.
compressicaule, *Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 45.—Java.
concinnum, *Muell. Arg. in Linnaea*, xxxii. (1863) 62.—Polynes.
cordatum, *Seem. in Bonplandia*, ix. (1861) 280, nomen.—Ins. Fiji.
corticosum, *Thw. Enum. Pl. Zeyl.* 285.—Zeylan.
coriaceum, Thw. l. c. partim = brachyloburn.
cornutum, *Muell. Arg. in Linnaea*, xxxii. (1863) 71.—N. Caled.
Cumingii, *Muell. Arg. l. c.* 61.—Ins. Philipp.

GLOCHIDION:—
cyrtophyllum, *Miq. Fl. Ind. Bat.* i. ii. 378.—Sumatra.
cyrtostylum, *Miq. l. c.*—Java.
Daltoni, *Kurz, For. Fl. Brit. Burma*, ii. 344.—Reg. Himal.
dasyanthum, *Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 47.—Java.
dasyphyllum, Miq. Fl. Ind. Bat. Suppl. 451 = superbum.
dasystylum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 237.—Burma.
distichum, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 238.—China.
diversifolium, Bedd. Forest. Man. Bot. 193 = ellipticum.
eleutherostylum, *Muell. Arg. in Linnaea*, xxxii. (1863) 69.—Ins. Philipp.
ellipticum, *Wight, Ic.* t. 1606.—Ind. or.
ericocarpum, *Champ. ex Benth. in Hook. Kew Journ.* vi. (1854) 6.—Ins. Hongkong.
eriocarpum, *Zipp. ex Span. in Linnaea*, xv. (1841) 846, nomen.—Ins. Timor.
fagifolium, *Bedd. Forest. Man. Bot.* 193.—Ind. or.
Fagueti, *Muell. Arg. in Linnaea*, xxxii. (1863) 59.—N. Caled.
flavidum, *Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 46.—Java.
flexuosum, *F. Muell. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 128.—Japon.
Fortunei, Hance, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 238 = sinicum.
frondosum, Baill. Etud. Gén. Euph. 637, nomen = Phyllanthus frondosus.
fulvinaeanum, *Miq. Fl. Ind. Bat.* i. ii. 376.—Java.
Gardneri, *Thw. Enum. Pl. Zeyl.* 286.—Zeylan.
glanciflium, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 65.—Burma.
glaucagynum, Bedd. Forest. Man. Bot. 195 = Moonii.
glaucum, *Blume, Bijdr.* 587.—Java.
giaucum, *Muell. Arg. in Linnaea*, xxxii. (1863) 62.—N. Caled.
heterolobum, Muell. Arg. l. c. 68 = Billardieri.
Heyneanum, *Wight, Ic.* t. 1908.—Ind. or.
hirsutum, *Voigt, Hort. Suburb. Calc.* 158; *Muell. Arg. in Linnaea*, xxxii. (1863) 61.—Ind. or.
Hohenackeri, *Bedd. Forest. Man. Bot.* 193.—Ind. or.
hongkongense, *Muell. Arg. in Linnaea*, xxxii. (1863) 60.—Ins. Hongkong.
insulanum, *Muell. Arg. l. c.* 67.—Ins. Moluc.
Juzzuvanum, Thw. Enum. Pl. Zeyl. 285 = rigidum.
kanalense, *Baill. Adansonia*, ii. (1861-62) 241.—N. Caled.
Lambertianum, *Muell. Arg. in Linnaea*, xxxii. (1863) 60.—Ins. Moluco.
lanceolarium, *Voigt, Hort. Suburb. Calc.* 158; *Dais. & Gibs. Bomb. Fl.* 235.—Ind. or.
laurifolium, *Heynh. Nom.* ii. 266.—Hab. ?
leiostylum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 237.—Burma.
Lenormandi, *Muell. Arg. in Linnaea*, xxxii. (1863) 59.—N. Caled.
leptocaulon, Bedd. Forest. Man. Bot. 195 = Gardneri.
leucogynum, *Miq. Fl. Ind. Bat.* i. ii. 376.—Java.
littorale, Benth. Fl. Hongk. 314 = hongkongense.
littorale, *Blume, Bijdr.* 585.—Ind. or.; China.
Llanosi, *Muell. Arg. in Linnaea*, xxxii. (1863) 68.—Ins. Philipp.
lucidum, *Blume, Bijdr.* 584.—Java.
luteocoma, *Blume, l. c.* 583.—Java.
macrocarpum, *Blume, l. c.* 584.—Java.
macrochorion, *Muell. Arg. in Linnaea*, xxxii. (1863) 59.—N. Caled.
macrophyllum, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 491.—China.
macrophyllum, *Muell. Arg. in Linnaea*, xxxii. (1863) 61.—N. Caled.
majus, Baill. Etud. Gén. Euph. 637, nomen = talisense.
malabaricum, *Bedd. Forest. Man. Bot.* 194.—Ind. or.
Marcono, *Baill. Etud. Gén. Euph.* 637, nomen.—Ins. Tahiti.
marianum, *Muell. Arg. in Linnaea*, xxxii. (1863) 65.—Ins. Marian.
salcrophyllum, *Muell. Arg. l. c.* 66.—Guian.
molle, *Blume, Bijdr.* 586.—Java.
molle, Hook. & Arn. Bot. Beech. Voy. 210 = Arnottianum.
moluccanum, *Blume, Bijdr.* 587.—Ins. Moluco.

GLOCHIDION :—
montanum, *Thw. Enum. Pl. Zeyl.* 286.—Zeylan.
Moonii, *Thw. l. c.*—Zeylan.
multiloculare, *Voigt, Hort. Suburb. Calc.* 152; *Muell. Arg. in Linnaea*, xxxii. (1863) 59.—Ind. or.
neilgherrense, *Wight, Ic.* t. 1907.—Ind. or.
nemorale, *Thw. Enum. Pl. Zeyl.* 286.—Zeylan.
nepalense, *Kurz, For. Fl. Brit. Burma*, ii. 344.—Burma.
nitidum, Voigt, Hort. Suburb. Calc. 153; Dalz. & Gibs. Bomb. Fl. 235 = zeylanicum.
obliquum, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 481 = zeylanicum.
obovatum, *Steb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1843) 143.—Japon.
obscurum, *Blume, Bijdr.* 585.—Ind. or.; Malaya; China.
ornatum, *Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 46.—Sumatra.
ovalifolium, *Zipp. ex Span. in Linnaea*, xv. (1841) 346, nomen.—Ins. Timor.
ovatum, *Muell. Arg. l. c.* xxxii. (1863) 71.—Ins. Martinic.
Pancherianum, *Muell. Arg. l. c.*—N. Caled.
Perrottetianum, Bedd. Forest. Man. Bot. 194 = neilgherrense.
philippense, *Benth. Fl. Hongk.* 314.—Ins. Philipp.
pinnatum, Voigt, Hort. Suburb. Calc. 153 = obscurum.
Poeppigianum, *Muell. Arg. in Linnaea*, xxxii. (1863) 71.—Bras.
pubescens, *Zipp. ex Span. l. c.* xv. (1841) 346, nomen. —Ins. Timor.
pycnocarpum, *Bedd. Forest. Man. Bot.* 194.—Zeylan.
ramiflorum, *Forst. Char. Gen.* 114.—Ins. Pacif.
rigidum, *Muell. Arg. in Linnaea*, xxxii. (1863) 67.—Zeylan.
Roxburghianum, Muell. Arg. l. c. 61 = obscurum.
rubrum, *Blume, Bijdr.* 586.—Malaya; Ins. Philipp.
salicifolium, *Muell. Arg. in Linnaea*, xxxii. (1863) 59. —N. Caled.
Seemanni, *Muell. Arg. l. c.* 68.—Ins. Fiji.
sericeum, *Zoll. & Mor. in Nat. ex Geneesk. Arch. Nederl. Indie*, ii. (1845) 585.—Malaya.
sinicum, *Hook. & Arn. Bot. Beech. Voy* 210.—China.
sphaerogynum, *Kurz, For. Fl. Brit. Burma*, ii. 346.—Reg. Himal.
stellatum, *Bedd. Forest. Man. Bot.* 194.—Ind. or.
subcandens, *Kurz, For. Fl. Brit. Burma*, ii. 344.—Burma.
subcandens, *Zoll. & Mor. in Nat. ex Geneesk. Arch. Nederl. Indie*, ii. (1845) 584.—Java; Sumatra.
sumatranum, Miq. Fl. Ind. Bat. Suppl. 450 = subscandens, *Zoll. & Mor.*
superbum, *Baill. Etud. Gén. Euph.* 638.—Burma; Malaya.
symplocoides, Bedd. Forest. Man. Bot. 195 = montanum.
tahitense, *Baill. Etud. Gén. Euph.* 637, nomen.—Ins. Tahiti.
Thwaitesii, Muell. Arg. in Linnaea, xxxii. (1863) 66 = rigidum.
tomentosum, *Dals. in Hook. Kew Journ.* ii. (1851) 88. —Ind. or.
trichogynum, *Muell. Arg. in Linnaea*, xxxii. (1863) 66.—Ins. Philipp.
uniflorum, J. F. Gmel. Syst. 1555, sphalm. = ramiflorum.
vaccinifolium, *Muell. Arg. in Linnaea*, xxxii. (1863) 63.—Guian.
variana, *Miq. Fl. Ind. Bat. Suppl.* 450.—Sumatra.
velutinum, *Wight, Ic.* t. 1907, 2.—Ind. or.
Vespertilio, *Muell. Arg. in Linnaea*, xxxii. (1863) 59. —N. Caled.
Vieillardi, Muell. Arg. l. c. 70 = Pancherianum.
villosum, Miq. Fl. Ind. Bat. i. ii. 376 = molle.
Wallichianum, *Muell. Arg. in Linnaea*, xxxii. (1863) 67.—Ins. Penang.
Wrightii, *Benth. Fl. Hongk.* 313.—China.
zeylanicum, *A. Juss. Tent. Euphorb.* 107.—Ind. or.; Malaya.
Zollingeri, *Miq. Fl. Ind. Bat.* i. ii. 636.—Java.

GLOCHIDIONOPSIS, Blume, Bijdr. 588 (1825) = **Glochidion**, Forst. (Euphorb.).
sericea, Blume, l. c. (= *Phyllanthus sericeus*).—Malaya.

GLOCHIDOTHECA, Fenzl, in Russegg. Reise, i. ii. 870, in obs. (1842) = **Oxmelis**, Linn. (Umbellif.).
foeniculacea, Fenzl, l. c.—Oriens.

GLOCHISANDRA, Wight, Ic. v. ii. 28. t. 1905 (1852) = **Glochidion**, Forst. (Euphorb.).
acuminata, Wight, l. c. = Glochidion lanceolarium.

GLOCKERIA Nees, in DC. Prod. xi. 728 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1106.
GALEOTTIA, Nees, in DC. Prod. xi. 211 (1847).
glandulosa, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 141.—Mexic.
gracilis, *Nees, in DC. Prod.* xi. 728.—*Cf.* 811.—Mexic.
haematodes, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 141. t. 5. f. 21.—Mexic.
sessilifolia, *Oerst. l. c.*—Costa Rica.

GLOEOSPERMUM, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 128. *VIOLARIEAE*, Benth. & Hook. f. i. 119.
Martii, *Eichl. in Mart. Fl. Bras.* xiii. i. 389.—Bras.
sphaerocarpum, *Triana & Planch. in Ann Sc. Nat.* Sér. IV. xvii. (1862) 129.—N. Granat.
Sprucei, *Eichl. in Mart. Fl. Bras.* xiii. i. 390.—Bras.

GLOIOSPERMUM, Triana & Planch. ex Benth. & Hook f. Gen. i. 119 (1862), sphalm. = preec.

GLOMERA, Blume, Bijdr. 872. t. 68 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 516.
brasiliensis, *Rodr. Orch. Nov.* i. 147.—Bras.
erythrosma, *Blume, Bijdr.* 372. t. 68.—Java.
montana, *Reichb. f. ex Seem. in Bonpiandia*, ix. (1861) 260; *et in Linnaea*, xli. (1877) 77.—Ins. Fiji.

GLOMERARIA, Cav. Desc. 319 (1802) = **Amaranthus**, Linn.
bicolor, Cav. ex Moq. in DC. Prod. xiii. ii. 262 = A. melancholicus.
deflexa, Cav. Descr. 319 = A. deflexus.
viridis, Cav. l. c. = A. viridis.

GLONERIA, André, Illustr. Hortic. xviii. (1871) 76. t. 60 = **Psychotria**, Linn. (Rubiac.).
jasminiflora, André, l. c. = P. jasminiflora.

GLORIOSA, Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 830.
CLINOSTYLIS, Hochst. in Flora, xxvii. (1844) 26.
MENDONI, Adans. Fam. ii. 48, 576 (1763).
METHONICA, Juss. Gen. 43 (1789).
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 322.—Abyssin.
angulata, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 171 = superba.
caerulea, Ball. Gard. Dict. ed. VIII. n. 2 = simplex.
cirrhifolia, Stokes, Bot. Mat. Med. ii. 287 = superba.
Doniana, Schult. f. Syst. vii. 866 = superba.
nepalensis, G. Don, in Lond. Hort. Brit. 181 = superba.
Plantii, Lond. Encyc Pl. 1544 = simplex.
simplex, D. Don, Prod. Fl. Nep 51 = superba.
simplex, *Linn. Mant.* i. 62.—Afr. trop.
superba, *Linn. Sp. Pl.* 305.—As. trop.
virescens, Lindl. Bot. Mag. t. 2539 = simplex.

GLOSOCOMIA, D. Don, Prod. Fl. Nep. 158 (1825) = **Codonopsis**, Wall. (Campan.).
clematidea, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. v. 46 = C. ovata.
hortensis, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 209 = C. lanceolata.
lanceolata, Rupr. l. c. 228 = Campanumoea japonica.
ovata, Lindl. Bot. Reg. (1842) t. 3 = C. ovata.
purpurea, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 210 = C. purpurea.
rotundifolia, Rupr. l. c. = C. rotundifolia.
tenera, D. Don, Prod. Fl. Nep. 158 = C. thalictrifolia.
thalictrifolia, Wall. Cat. n. 1297 = C. thalictrifolia.
uzuvienzis, Maxim. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 209 = C. lanceolata?
viridis, Rupr. l. c. 210 = C. viridis.

GLOSSANTHUS, Klein, ex Benth. Scroph. Ind. 57 (1835) = **Klugia**, Schlecht. (Gesneriac.).
grandiflorus, Benth. in Wall. Cat. n. 6395 = Staurianthera grandiflora.
malabaricus, Klein, ex Benth. Scroph. Ind. 57 = K. Notoniana.
maximus, R. Br. in Benn. Pl. Jav. Rar. 121 = K. saxea.
Notoniana, R. Br. l. c. = K. Notoniana.
zeylanicus, R. Br. l. c. = K. zeylanica.

GLOSSARRHEN, Mart. Nov. Gen. et Sp. i. 21 (1824); et ex Ging. in DC. Prod. i. 290 (1824) = **Schweigeria**, Spreng. (Violar.).
floribundus, Mart. ll. oc. 22, 291 = S. fruticosa.
mexicanus, Steud. Nom. ed. II. i. 689 = S. mexicana.
parviflorus, Mart. ex Ging. in DC. Prod. i. 291 = seq.
pauciflorus, Mart. Nov. Gen. et Sp. i. 23 = S. fruticosa.

GLOSSASPIS, Spreng. Syst. iii. 675, 694 (1826) = **Glossula**, Lindl. (Orchid.).
antennifera, Reichb. f. in Linnaea, xxv. (1852) 225 = Glossula tentaculata.
tentaculata, Spreng. Syst. iii. 694 = Glossula tentaculata.

GLOSSOCALYX, Benth. in Hook. Ic. Pl. t. 1301 (1880). *MONIMIACEAE*, Benth. & Hook. f. iii. 144. brevipes, *Benth. l. c.* t. 1302.—Afr. trop.
longicaspis, *Benth. l. c.* t. 1301.—Afr. trop.

GLOSSOCARDIA, Cass. in Bull. Soc. Philom. (1817) 138; et in Dict. Sc. Nat. xix. 62 (1821). *COMPOSITAE*, Benth. & Hook. f. ii. 864.
Bosvallia, DC. in Wight, Contrib. 18 = linearifolia.
linearifolia, *Cass. in Dict. Sc. Nat.* xix. 62.—Ind. or.

GLOSSOCARYA, Wall. Cat. n. 1741 (1831). *VLACHENEAE*, Benth. & Hook. f. ii. 1158.
hemiderma, *Benth. & Hook. f. Gen.* ii. 1158.—Austral.
Linnaei, *Benth. & Hook. f. l. c.*, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—Zeylan.
mollis, *Wall. Cat.* n. 1741.—Burma.
pinnatifida, Steud. Nom. ed. II. i. 419, in syn. sphalm. = Glossogyne pinnatifida.

GLOSSOCENTRUM, Cneeg. in Linnaea, xx. (1847) 111 = **Miconia**, Ruiz & Pav. (Melast.).
collinum, Cneeg. l. c. = M. minutiflora.

GLOSSOCHILUS, Nees, in DC. Prod. xi. 83 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1092.
Burchellii, *Nees, l. c.*—Afr. austr.

GLOSSOCOMA, Endl Gen. 799, in syn. (1839) = GLOSSOMA, Schreb. = **Votomita**, Aubl. (Inc. sed.).

GLOSSOCOMIA, Reichb. Consp. 114 (1828) = **Codonopsis**, Wall. (Campan.).

GLOSSODIA, R. Br. Prod. 325 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 513.
Brunzoii, *Endl. Nov. Stirp. Dec.* 16.—Austral.
emarginata, *Lindl. Gen. et Sp. Orch.* 424.—Austral.
intermedia, *Fitzg. in Gard. Chron.* (1882) i. 462.—Austral.
major, *R. Br. Prod.* 326.—Austral.
minor, *R. Br. l. c.*—Austral.

GLOSSOGYNE, Cass. in Dict. Sc. Nat. li. 475 (1827). *COMPOSITAE*, Benth. & Hook. f. ii. 888.
DIODONTIUM, F. Muell. in Hook. Kew Journ. ix. (1857) 19.
GYNACTIS, Cass. in Dict. Sc. Nat. li. 475 (1827).
bidentidea, F. Muell. in Linnaea, xxv. (1853) 402 = tenuifolia.
brasiliensis, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 408 = Isostigma microcephalum.
chinensis, Less. Syn. Comp. 218 = Bidens pilosa.
filifolia, *F. Muell. ex Benth. Fl. Austral.* iii. 544.—Austral.
Kennedyi, *R. Br. ter. in Trans. N. E. Inst.* xv. (1883) 259.—N. Zel.
paucuculosa, DC. Prod. v. 582 = tenuifolia.
pinnatifida, *DC. in Wight, Contrib.* 19.—Ind. or.; Malaya.
retrofracta, *F. Muell. Fragm.* i. 51.—Austral.
tenuifolia, *Cass. in Dict. Sc. Nat.* li. 475.—Malaya; Austral.
tridentata, *Benth. & Hook. f. Gen.* ii. 369.—Ins. Philipp.

GLOSSOLOMA, Hanst. in Linnaea, xxvi. (1853) 209 = **Alloplectus**, Mart. (Gesneriac.).
tetragonum, Hanst. l. c. 209, 215. t. 2. f. 49 = A. tetragonus.

GLOSSOMA, Schreb. Gen. ii. 792 (1791) = **Votomita**, Aubl. (Inc. sed.).
acuminatum, Wall. ex Benn. Pl. Jav. Rar. 196 = Polycosma integrifolia.
arborescens, Willd. Sp. Pl. i. 664 = V. guianensis.
Votomita, Roem. & Schult. Syst. iii. 818 = V. guianensis.

GLOSSONEMA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 335. t. 12. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 748.
CONOMITRA, Fenzl, in Endl. Nov. Stirp. Dec. 65 (1839).
MASTIGOSTIGMA, Stocks, in Hook. Ic. Pl. t. 863 (1852).
BOVEANUM, *Decne. in Ann. Sc. Nat. Sér. II. ix.* (1838) 335. t. 12. f. D.—Arab.
Bovaanum, Hochst. & Steud. ex Decne. in DC. Prod. viii. 556 ~ nubicum.
linsare, *Decne. in DC. Prod.* viii. 555.—Afr. trop.
nubicum, *Decne. l. c.*—Afr. bor.
Revoili, *Franch. in Révoil, Çomali*, 40.—Afr. trop.
variaus, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 16.—Persia; Scindia.

GLOSSOPAPPUS, Kunze, in Flora, xxix. (1846) 748 = **Chrysanthemum**, Linn. (Compos.).
chrysanthemoides, Kunze, l. c. = C. macrotum.

GLOSSOPETALON, A. Gray, Pl. Wright. ii. 29. t. 12 B (1853). *CELASTRINEAE*, Benth. & Hook. f. i. 368.
nevadense, *A. Gray, in Proc. Am. Acad.* xi. (1876) 78. — Am. bor. occ.
spinescens, *A. Gray, Pl. Wright.* ii. 29.—Am. bor. occ.

GLOSSOPETALUM, Benth. & Hook. f. Gen. i. 368 (1862) = praec.

GLOSSOPETALUM, Schreb. Gen. i. 205 (1789) = **Goupia**, Aubl. (Celastr.).
glaberum, J. F. Gmel. Syst. 510; Willd. Sp. Pl. i. 1021 = G. glabra.
tomentosum, J. F. Gmel. l. c.; Willd. l. c. 1522 = G. tomentosa.

GLOSSOPHYLLUM, Fourr. in Ann. Soc. Linn. Lyon, xvi. (1868) 325 = **Ranunculus**, Tourn.
ophioglossifolium, Fourr. l. c. = R. ophioglossifolius.

GLOSSOSCHIMA, Walp. Rep. i. 375 (1842) = CLOSASCHIMA, Korth. = **Laplacea**, H. B. & K. (Ternstroem.).

GLOSSOSPERMUM, Wall. Cat. n. 1158 (1831) = **Melochia**, Linn. (Sterculiac.).
cordatum, Wall. l. c. n. 1155 = M. velutina.
quinquenalatum, Wall. l. c. n. 1154 = Columbia floribunda.
velutinum, Wall. l. c. n. 1153 = M. velutina.

GLOSSOSTEMON, Desf. in Mém. Mus. Par. iii. (1817) 288. t. 11. *STERCULIACEAE*, Benth. & Hook. f. i. 224.
Bruguieri, *Desf. l. c.* 239.—Persia.

GLOSSOSTEMUM, Steud. Nom. ed. I. 373 (1821) = praec.

GLOSSOSTEPHANUS, E. Mey. Comm. Pl. Afr. Austr. 217 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 759.
ONCINEMA, Arn. in Edinb. N. Phil. Journ. xvii. (1834) 261.
linearis, *E. Mey. Comm. Pl. Afr. Austr.* 218.—Afr. austr.

GLOSSOSTIGMA, Wight & Arn. in Nov. Act. Nat. Cur. xvii. (1836) 385. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 958.
PELTIMELA, Raf. Atl. Journ. 199 (1833), nomen prius.
TRICHOLOMA, Benth. in DC. Prod. x. 426 (1846).
Drummondi, *Benth. in DC. Prod.* x. 426.—Austral.
elatinoides, *Benth. ex Hook. f. Fl. N. Zel.* i. 189.—Austral.; N. Zel.
spathulatum, *Wight & Arn. in Nov. Act. Nat. Cur.* xvii. (1836) 385.—Ind. or.

GLOSSOSTYLIS, Cham. & Schlecht. in Linnaea, iii. (1828) 22 = **Alectra**, Thunb. (Scrophul.).
abyssinica, Hochst. ex A. DC. in DC. Prod. x. 545, in nota = Bartsia Trixago.
arvensis, Benth. Scroph. Ind. 49 = A. indica.
aspera, Cham. & Schlecht. in Linnaea, iii. (1828) 22 = A. brasiliensis.
asperrima, Hochst. ex Benth. in DC. Prod. x. 540 = A. asperrima.

GLOSSOSTYLIS:—
capensis, Benth. Scroph. Ind. 50 = Alectra melampyroides.
cordata, Hochst. ex Benth. in DC. Prod. x. 539 = A. indica.
parasitica, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 117 = A. parasitica.

GLOSSULA, Lindl. Bot. Reg. t. 862 (1824). *ORCHIDEAE*, Benth. & Hook. f. iii. 539.
GLOSSARPIS, Spreng. Syst. iii. 675, 694 (1826).
tentaculata, *Lindl. l. c.* (Cf. Habenaria lacertifera).—China.

GLOSSULA, Reichb. Handb. 178 (1837) = **Aristolochia**, Linn.

GLOTTES, Medic. Phil. Bot. i. 210 (1789) = GLOTTIS, Medic. = **Astragalus**, Tourn. (Legumin.).

GLOTTIDIUM, Desv. Journ. Bot. i. (1813) 119. t. 1 = **Sesbania**, Pers. (Legumin.).
floridanum, DC. Prod. ii. 266 = S. platycarpa.

GLOTTIS, Medic. in Vorles. Churpf. Phys. Oec. ii. (1787) 377 = **Astragalus**, Tourn. (Legumin.).
echinata, Medic. l. c. = A. pentaglottis.
epiglottis, Medic. l. c. = A. epiglottis.
pentaglottis, Medic. Phil. Bot. i. 210 = A. pentaglottis.

GLOXINIA, L'Hérit. Stirp. Nov. 149 (1785). *GESNERACEAE*, Benth. & Hook. f. ii. 998.
ALMANA, Raf. Fl. Tellur. iv. 68 (1836).
ESCHINA, Regel, in Flora, xxxii. (1849) 179, in syn.
EUCOLUM, Salisb. Prod. 98 (1796).
EUMOLPE, Decne. ex Jacques & Hérincq, Man. Gén. Pl. ii. 575 (1849?).
ORTHANTHE, Lem. in Illustr. Hortic. iii. (1856) sub t. 81.
PELTIPOMA, Hanst. in Linnaea, xxvi. (1853–55) 201.
SALISIA, Regel, in Bot. Zeit. ix. (1851) 894.
Amherstiana, DC. Prod. vii. 534 = Achimenes foliosa.
attenuata, *Hanst. in Linnaea*, xxvii. (1854) 716.—Bras.
arborea, *Calla, Hort. Ritsul.* 61.—Hab.?
bicolor, Poepp. ex Hanst. in Mart. Fl. Bras. viii. 419 = maculata.
calycina, Hort. ex Steud. Nom. ed. II. i. 690 = Pentarhaphia calycina.
caulescens, Lindl. Bot. Reg. t. 1127 = Sinningia speciosa.
digitaliflora, *Paxt. Mag. Bot.* x. (1843) 193.—Am. austr.?
discolor, Kunze, in Linnaea, xvi. (1842) 314 (= *Ligeria discolor*).—Bras.
dubia, *G. Don, Gen. Syst.* iv. 849.—Mexic.
fimbriata, *Brongn. in Rev. Hortic.* Sér. III. i. (1847) 264.—Mexic.
fimbriata, Hook. Bot. Mag. t. 4430 = glabrata.
floribunda, Walp. Ann. ii. 1067 = Sinningia floribunda.
fruticosa, *Dekntl. Mem. Sopr. Alc. Piante*, 3; *ex Walp. Rep.* ii. 730.—Bras.
Fyfiana ×, Hort. Angl. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 311.
gesneroides ×, Lem. l. c. ii. (Fév. 1846) t. 3.
glabrata, *Zucc. in Flora*, xv. (1832) ii. *Beibl.* 99.—Mexic.
gloxiniflora, Walp. Ann. i. 478 = Sinningia gloxiniflora.
gracilis, Mart. Nov. Gen. et Sp. iii. 64, t. 226 (= *Ligeria gracilis*).—Bras.
guttata, Mart. l. c. = Sinningia guttata.
gymnostoma, *Griseb. in Goett. Abh. xix.* (1874) 227.—Reg. Argent.
Hellerí, Hort. Nov. Gen. et Sp. iii. 64 = Sinningia Hellerí.
heterophylla, Poepp. & Endl. Nov. Gen. et Sp. iii. 9 = maculata.
hirsuta, Lindl. Bot. Reg. t. 1004; Lodd. Bot. Cab. t. 1296 (= *Ligeria hirsuta*).—Bras.
hypocyrtiiflora, *Hook. f. Bot. Mag.* t. 5695.—Reg. Argent.
ichthyostoma, Gardn. in Hook. Ic. Pl. t. 472 = Achimenes ichthyostoma.

GLOXINIA:—
immaculata, Mart. ex Hanst. in Mart. Fl. Bras. viii. 389 = Sinningia speciosa.
Lindleyi, Steud. Nom. ed. II. i. 690 = Pentarhaphia calycina.
macrophylla, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 96 (= *Ligeria macrophylla*).—Bras.
maculata, *L'Hérit. Stirp. Nov.* 149.—Am. austr.
Menziesiana, Young, ex Otto & Dietr. Allg. Gartenz. v. (1837) 394 = Sinningia speciosa.
Merkii, E. Otto, in Otto & Dietr. l. c. xvi. (1848) 242 = Sinningia speciosa.
micrantha, *Mart. & Gal. in Bull. Acad. Brux.* ix. 11. (1842) 35.—Mexic.
Micllieri, *Regel, Ind. Sem. Hort. Petrop.* (1865) 64.—Hab.?
multiflora, Mart. & Gal. in Bull. Acad. Brux. ix. 11. (1842) 36 = Naegelia multiflora.
pallidiflora, *Hook. Bot. Mag.* t. 4213.—N. Granat.
Paxtoniana, Paxt. Mag. Bot. xii. (1846) 267 = Sinningia speciosa.
picta, Vis. in Atti Ottava Riun. Sc. Ital. 559 = Tydaea picta.
punctata, Mart. & Gal. in Bull. Acad. Brux. ix. 11. (1842) 34 = Naegelia zebrina.
rubra, Paxt. Mag. Bot. vii. (1840) 271 = Sinningia speciosa.
Sarmentiana, *Gardn. ex Hook. Ic. Pl.* t. 378.—Bras.
Schottii, Mikan, Del. Fl. Bras. [21] = Sinningia Hellerí.
sinningiaeflora, Walp. Ann. ii. 1066, sphalm. = Sinningia gloxiniflora.
speciosa, Lodd. Bot. Cab. t. 28 = Sinningia speciosa.
suaveolens, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 468.—Guiana.
Teichleri ×, Hort. ex Lem. in Fl. des Serres, Sér. I. iii. (1847) 220.
Teuchleri ×, Lem. l. c. t. 345, verso.
tigridia, Ohlend. in Otto & Dietr. Allg. Gartenz. xiii. (1845) 376 = Campanea grandiflora.
trichotoma, Moench, Meth. Suppl. 194 = maculata.
trichantha, Miq. in Linnaea, xxii. (1849) 473 = maculata.
tubiflora, Hook. Bot. Mag. t. 3971 (= *Dolichodeira tubiflora*).—Bras.
variabilis, *Carr. in Rev. Hortic.* (1877) 70.—Hab.?
velutina, Mart. Nov. Gen. et Sp. iii. 64 = Sinningia velutina.
verticillata, Mart. & Gal. in Bull. Acad. Brux. ix. 11. (1842) 35 = Achimenes Andrieuxii.
villosa, Mart. Nov. Gen. et Sp. iii. 64 = Sinningia villosa.
Youngeana, Walp. Rep. vi. 401 = *Sinningia Youngeana*.

GLUMOSIA, Herb. in Lindl. Bot. Reg. (1843) Misc. 85 = **Sisyrinchium**, Linn. (Irid.).
palmifolium, Herb. l. c. = S. palmifolium.

GLUTA, Linn. Mant. ii. 293 (1771). *ANACARDIACEAE*, Benth. & Hook. f. i. 421.
STAGMARIA, Jack, Desc. Mal. Pl. iii. 12 (1823?).
SYNDEMIS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 314 (1824).
Benghas, Kurz, in Pegu Rep. A. 41 = tavoyana.
Benghas, *Linn. Mant.* ii. 293.—Java.
coarctata, Hook. f. Fl. Brit. Ind. ii. 22 = velutina.
elegans, *Hook. f. l. c.*—Ins. Penang.
longipetiolata, *Kurz, in Pegu Rep.* A. 41; *et in Journ. As. Soc. Beng.* xlv. (1876) 210.—Ins. Andaman.
orgyalis, Blanco, Fl. Filip. ed. II. 451 (Cleistanthi sp.).—Ins. Philipp.
tavoyana, *Hook. f. Fl. Brit. Ind.* ii. 22.—Burma.
Tourtour, March. Rév. Anac. 187 = G. Benghas.
travancorica, *Bedd. Fl. Syls.* t. t. 60.—Ind. or.
velutina, *Blume, Mus. Bot. Lugd. Bat.* i. 183.—Malaya.

GLUTAGO, Comm. ex Pelt. in Dict. Sc. Nat. xix. 57 (1821) = **Lorranthus**, Linn.
spicata, Raf. Sylva Tellur. 126 = L. spicatus.

GLUTINARIA, Raf. Fl. Tellur. iii. 98 (1836) = **Salvia**, Tourn. (Labiat.).
glutinosa, Raf. l. c. = S. glutinosa.

GLYASPERMUM, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, ii. (1845) 11 = **Pittosporum**, Soland.
ramiflorum, Zoll. & Mor. l. c. 12.—Malaya.

GLYCANTHES, Rafin. Sylva Tellur. 88 (1838) = **Columnea**, Linn. (Gesner.).
scandens, Rafin. l. c. = C. scandens.

GLYCE, Lindl. Syn. Brit. Fl. 26 (1839) = **Alysum**, Tourn. (Crucif.).
maritima, Lindl. l. c. = A. maritimum.

GLYCERIA, R. Br. Prod. 179 (1810). *GRAMINEAE*, Benth. & Hook. f. tii. 1197.
ATROPIS, Rupr. Fl. Samoj. 61 (1845).
DEVAUXIA, Beauv. ex Kunth, Enum. Pl. i. 367, in syn. (1838).
DIACHROA, Nutt. ex Steud. Nom. ed. II. i. 497 (1840).
EXYDRA, Endl. Fl. Poson. 119 (1830).
HYDROCHLOA, Hartm. Gram. Skand. 8 (1819).
PEROTERANTHE, Steud. Syn. Pl. Gram. 287 (1854).
PUCCINELLIA, Parl. Fl. Ital. i. 366 (1850).
acuminata, Schur, Enum. Pl. Transs. 781 = fluitans.
acutiflora, *Torr. ex Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68.—Am. bor.
airoides, A. Gray, in Proc. Acad. Sc. Philad. 1862 (1863) 336 = distans.
airoides, Reichb. in Moesal. Handb. ed. II. iii. 1827 = Catabrosa aquatica.
airoides, *Steud. Syn. Pl. Gram.* 287.—Japon.
algida, Hartm. Handb. ed. V. 294 = Phippsia algida.
angustata, *T. Fries, in Öfvers. Vet. Akad. Stockh.* xxvi. (1869) 139.—Reg. arct.
aquatica, J. & C. Presl, Fl. Cech. 25 = Catabrosa aquatica.
aquatica, *Wahlenb. Fl. Goth.* 18; *Sm. Engl. Fl.* i. 116.—Reg. bor. temp.
arctica. *Hook. Fl. Bor. Am.* ii. 248.—Am. arct.
arundinacea, Kunth, Rév. Gram. i. 118 = remota.
australasica, *Steud. Syn. Pl. Gram.* 286.—Austral.
Baumgartenvana, Schur, Enum. Pl. Transs. 781 (sp. deland.).
Borreri, *Bab. in Suppl. Engl. Bot.* t. 2797.—Europ.
brevifolia, Schult. Mant. ii. 387 = Diarrhena americana.
bulbosa, *Buchl. in Proc. Acad. Sc. Philad.* 1862 (1863) 95.—Am. bor.
canadensis, *Trin. in Mém. Acad. Pétersb. Sér.* VI. i. (1831) 366.—Am. bor.
Canbyi, *Scribner, in Bull. Torrey Bot. Club*, x. (1883) 77.—Am. bor.
capillaris, Wahlenb. Fl. Goth. 17 = distans.
caspia, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 68 = aquatica.
conferta, Fries, Mant. iii. 10 = Borreri.
convoluta, Fries, l. c. iii. 176 = distans.
declinata, Bréb. Fl. Norm. ed. III. 354 = fluitans.
delawarica, Heynh. Nom. i. 360 = maritima.
denticulata, Dum. Obs. Gram. Belg. 107 = fluitans.
distans, *Wahlenb. Fl. Upsal.* 36.—Europ.
dives, *F. Muell. ex Benth. Fl. Austral.* vii. 659.—Austral.
elongata, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68.—Am. bor.
expansa, *Crép. in Mém. Cour. Acad. Belg.* xviii. 8vo (1866), 206, 209.—Europ.
festucaeformis, Heynh. ex Retchb. Fl. Germ. Excurs. 45 = distans.
festucaeformis, Schur, Enum. Pl. Transs. 780 = maritima.
fluitans, *R. Br. Prod.* 179.—Reg. bor. temp.; Austral.
Fordeana, *F. Muell. Fragm.* viii. 180.—Austral.
fulva, Fries, Summ. Veg. Scand. 244 = Poa fulva.
glaucescens, *Benth. in Journ. Linn. Soc.* xix. (1881) 116.—Chili.
glumaris, *Ledeb. Fl. Ross.* iv. 392.—Sibir. or.
Gussonii, Nym. Syll. 422 = distans.
humilis, Heynh. Nom. i. 361 = Catabrosa humilis.
hybrida, Towns. in Trans. Bot. Soc. Edinb. iv. (1853) 27, in obs. = fluitans.
indegva, Dum. Obs. Gram. Belg. 107 = fluitans.
intermedia, Klinggraeff, Fl. Preuss. 491 = distans.
intricata, Crép. in Mém. Cour. Acad. Belg. xviii. 8vo (1866), 207 = distans?
isekyronensis, Steud. Syn. Pl. Gram. 287 = aquatica.
japonica, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 281.—Japon.
Kjellmanni, *Lange, ex Kjellm. & Lundstr. in Nordensk. Vega-Exped. Vetensk. Iakttt.* i. 378, 514.—Sibir.
latispicea, *F. Muell. Fragm.* viii. 137.—Austral.

GLYCHRIA:—
Lemmoni, *Vasey, Gram. U. St.* 43, nomen.—Am. bor. occ.
leptophylla, Steud. Syn. Pl. Gram. 286 = distans.
leptostachya, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 95 = aquatica.
laticeps, Godr. Fl. Lorr. iii. 168 = fluitans.
magellanica, *Benth. in Journ. Linn. Soc.* xix. (1881) 116.—Reg. Magell.
maritima, *Mert. & Koch, Deutschl. Fl.* i. 588.—Reg. bor. temp.
maritima, Roep. ex Nym. Consp. 831 = Borreri.
Michauxii, Kunth, Rév. Gram. i. 118, 348. t. 85 = nervata.
micrantha, Steud. Syn. Pl. Gram. 287 = Eragrostis interrupta.
microthera, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 96 = aquatica.
minuta, Steud. Syn. Pl. Gram. 287 = Festuca parvula.
montana, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 96 = distans.
multiflora, *Steud. Syn. Pl. Gram.* 285.—Chili.
Nevadensis, Steud. l. c. 286 = distans.
nemoralis, *Uechtr. & Koern. in Bot. Zent.* xxiv. (1886) 191.—Rosa.
neogaea, *Steud. Syn. Pl. Gram.* 285.—' Terra Nova.'
nervata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 365.—Am. bor.
norvegica, Sommer. in Vet. Acad. Handl. Stockh. (1832) 254 = remota.
obtusa, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 366 = Poa obtusa.
ochroleuca, Guss. Fl. Sic. Syn. i. 100 = Catabrosa aquatica.
pallida, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68.—Am. bor.
palustris, Lange, in Fl. Dan. t. 2582 = arctica.
pauciflora, *Y. & C. Presl, Rel. Haenk.* i. 257.—Sinus Nootka.
pedicellata, Towns. in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 105 = fluitans.
pendulina, Lacstad. ex Wahlenb. Fl. Suec. 1088 = Poa Laestadii.
pensylvanica, Heynh. Nom. i. 361 = Eatonia obtusata.
permixta, Guss. Fl. Sic. Syn. i. 99 = Borreri.
plicata, Fries, Mant. iii. 176 = fluitans.
procumbens, *Dum. Obs. Gram. Belg.* 145.—Europ.; Mesopot.
Pseudo-distans, Crép. in Mém. Cour. Acad. Belg. xviii. 8vo (1866) 209 = distans.
ramigera, *F. Muell. Fragm.* viii. 181; *ex Benth. Fl. Austral.* vii. 659.—Austral.
remota, *Fries, Mant.* iii. 5.—Europ. bor.; Reg. Caucas.
rigida, Sm. Engl. Fl. i. 119 = Festuca rigida.
scaberrima, *Nees, ex Steud. Syn. Pl. Gram.* 287.—Ind. or.
songarica, Schrenk, Enum. Pl. Nov. i = Poa soongarica.
spectabilis, Mert. & Koch, Deutschl. Fl. i. 586 = aquatica.
Sphenopus, Steud. Syn. Pl. Gram. 287 = Sphenopus Gossani.
spicata, Guss. Fl. Sic. Syn. ii. 784 = fluitans.
stricta, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 95 = Sporobolus asper.
stricta, *Hook. f. Fl. N. Zel.* 304.—Austral.; N. Zel.
subdistans, *Ledeb. Fl. Ross.* iv. 392.—Sibir. altaic.
subspicata, *Regel, in Act. Hort. Petrop.* vii. (1880) 624.—Turkestan.
tenella, *Lange, ex Kjellm. in Nordensk. Vega-Exped. Vetensk. Iakttt.* i. 318.—N. Zembla.
tentiflora, *Steud. Syn. Pl. Gram.* 286.—Sibir.
tenuiflora, Boiss. & Reut. Pugill. Pl. Nov. 127 = distans.
tenuifolia, Steud. Syn. Pl. Gram. 286 = Catabrosa tenuifolia.
tenuispica, Steud. l. c. 285 = stricta?
tonglensis, *C. B. Clarke, in Journ. Linn. Soc.* xv. (1876) 119.—Reg. Himal.
transsilvanica, Schur, Enum. Pl. Transs. 780 = maritima.
vaginata, *Lange, in Fl. Dan.* t. 2588.—Groenl.
Vahliana, *T. Fries, in Öfver. Vet. Akad. Stockh.* xvi. (1859) 140.—Reg. arct.
villosiflora, *T. Fries*, l. c. 139.—Groenl.
viridula, Heynh. Nom. i. 361 = Aira viridula.

GLYCERIA, Nutt. Gen. Am. i. 177 (1818) = **Hydrocotyle**, Linn. (Umbell.).
asiatica, Nutt. l. c. = H. asiatica.
repanda, Nutt. l. c. = H. repanda.
triflora, Nutt. l. c. = H. asiatica.

GLYCIA, Linn. Syst. ed. I (1735) = **Astragalus**, Tourn. (Legum.).

GLYCICARPUS, Benth. & Hook. f. Gen. i. 425 (1862) = GLYCYCARPUS, Dalz. = **Nothopegia**, Blume, (Anacard.).

GLYCIDERAS, DC. Prod. vii. 257 (1838) = **Glycoderas**, Cass. (Compos.).

GLYCINE, Linn. Gen. ed. I. 249 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 580.
BUJACIA, E. Mey. Comm. Pl. Afr. Austr. i. 127 (1835)
JOHNIA, Wight & Arn. Prod. 449 (1834).
KENNEDYELLA, Steud. Nom. ed. II. i. 845 (1840)
LEPTOCYAMUS, Benth. in Trans. Linn. Soc. xviii. (1839) 209, adnot.
LEPTOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 124.
NOTONIA, Wight & Arn. Prod. 207 (1834).
SOJA, Moench, Meth 153 (1794).
TRIEMDILIX, Rafin. New Fl. Am. i. 85 (1836).
Abrus, Linn. Sp. Pl. 753 = Abrus precatoris.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 212 = Teramnus labialis.
abyssinica, Jaub. ex DC. Prod. ii. 386 = Rhynchosia pubescens.
ambigua, *Yacq Fragm.* 84 (sp. dub.).—Hab.?
americana, *Tenore, ex Steud. Nom.* ed. II. i. 691 (Quid?).—Hab.?
adongensis, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 179.—Afr. trop.
angulata, Desv. Journ. Bot. i. (1814) 78 = Teramnus uncinatus.
angulosa, Muhl. ex Willd. Sp. Pl. iii. 1056 = Phaseolus diversifolius.
angustifolia, Jacq. Hort. Schoenb. ii. 55 = Rhynchosia calycina.
anonychica, Walp. in Linnaea, xiii. (1839) 532 = Teramnus labialis.
Apios, Linn. Sp. Pl. 753 = Apios tuberosa.
arborea, *Fisch. Hort. Gorenk.* ed. II. (1812) 70, nomen.—Hab.?
argentea, Thunb. Prod. Pl. Cap. 131 = Rhynchosia argentea.
atomaria, *Willd. ex Spreng. Syst.* iii. 196.—Am. austr.
atrorubra, *Regel, Cat. Pl. Hort. Absahor.* 65.—Hab.?
aurea, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 218 = Rhynchosia aurea.
axilliflora, Kotschy, in Sitzb. Acad. Wien, Math.-Nat. ii. Abth. ii. (1866) 366 = Teramnus labialis.
Backhousiana, *Regel, Cat. Pl. Hort. Absahor.* 65.—Hab.?
bifora, *Schum. & Thonn. Beshr. Guin. Pl.* 345.—Afr. trop.
biloba, *Lindl. Bot. Reg.* t. 1418.—Mexic.
bimaculata, Curt. Bot. Mag. t. 263 = Hardenbergia monophylla.
binaculata, *Moench, Meth. Suppl.* 46.—Hab.?
bituminosa, Linn. Sp. Pl. 754 = Fagelia bituminosa.
Borianii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 180.—Afr. trop.
Botrydium, Walp. Rep. i. 760 = Teramnus uncinatus.
brachybotria, *Hort. ex Pasq. Cat. Orto Bot. Nap.* 49, nomen.—Hab.?
bracteata, Linn. Sp. Pl. 754 = Amphicarpaea monoica.
Bujacia, Benth. in Ann. Wien. Mus. ii. (1838) 126 = javanica.
caerulea, Salisb. Prod. 335 = Wisteria speciosa.
cajanoides, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 334.—Ins. Philipp.
cana, Willd. Sp. Pl. iii. 1063 = Rhynchosia cana.
capitata, Heyne, in Roth, Nov. Pl. Sp. 346 = Rhynchosia aurea.
caribaea, Jacq. Coll. i. 67 = Rhynchosia phaseoloides?
carolinicana, Spreng. Syst. iii. 197 = Rhynchosia tomentosa, *Hook. & Arn.*
chinensis, Sims, Bot. Mag. t. 2063 = Wistaria chinensis.
clandestina, *Wendl. Beobacht.* 54.—Austral.
coccinea, Curt. Bot. Mag. t. 270 = Kennedya prostrata.
comosa, Linn. Sp. Pl. 754 = Amphicarpaea monoica.

GLYCINE:—
Comptoniana, Andr. Bot. Rep. t. 602 = Hardenbergia Comptoniana.
corallina, Saliab. Prod. 334 = Rhynchosia phaseoloides.
crinita, H. B. & K. Nov. Gen. et Sp. vi. 421 = Eriosema crinitum.
cylindrifiora, Wall. Cat. n. 5516 = Shuteria ferruginea.
dalbergioides, *Zipp. ex Span. in Linnæa*, xv. (1841) 190, nomen.—Ins. Timor.
debilis, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 54 = Teramnus labialis.
densiflora, Roth, Nov. Pl. Sp. 348 = Rhynchosia densiflora.
denuata, *Vahl, ex Schum. & Thonn. Beskr. Guin. Pl.* 348.—Afr. trop.
diffusa, H. B. & K. Nov. Gen. et Sp. vi. 420. t. 572 = Eriosema diffusum.
discolor, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 190 = Teramnus uncinatus.
discolor, *Willd. ex Spreng. Syst.* iii. 196.—Ind. or dolichoides, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 415.—Ins. Timor.
dubia, *Colla, Hort. Ripul App.* iii. 37.—Ins. Guadeloup.
elegans, Hochst. in Flora, xxiv. (1841) 1. Intell. 42 = Rhynchosia elegans.
ellipsica, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 190 = Teramnus uncinatus.
ellipsica, Sm. in Abbot, Hist. Nat. Georgia, t. 21 = Amphicarpæa monoica.
elongata, Roth, Nov. Pl. Sp. 347 = Rhynchosia aurea.
emarginata, Desv. Journ. Bot. i. (1814) 78 = Galactia emarginata.
erecta, Nutt. Gen. Am. ii. 114 = Rhynchosia tomentosa.
erecta, Thunb. Prod. Pl. Cap. 131 = Rhynchosia Chrysoscias.
falcata, *Benth. Fl. Austral.* ii. 243.—Austral.
ferruginea, R. Grah. in Wall. Cat. n. 5514 = Shuteria ferruginea.
filiformis, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 415.—Hab. ?
filiformis, Wall. Cat. n. 5510 = Teramnus labialis.
flexa, Hornem. Hort. Hafn. ii. 682 = Amphicarpæa monoica.
flaccida, Wall. Cat. n. 5517 = Argyrolobium flaccidum.
flexilis, R. Grah. in Wall. l. c. n. 5521 = Teramnus flexilis.
floribunda, Willd. Sp. Pl. iii. 1066 = Wistaria chinensis.
Fridericiana, Wehnn. in Flora, iv. (1821) 29 = Rhynchosia Fridericiana.
frutescens, Linn. Sp. Pl. 753 = Wisteria speciosa.
Galactia, Linn. Syst. ed. X. 1178 = Galactia pendula.
galactioides, H. B. & K. Nov. Gen. et Sp. vi. 427 = Calopogonium galactioides.
gamboenchia, Walp. in Linnæa, xiii. (1839) 583 = Teramnus labialis.
glabra, Spreng. Neue Entdeck. iii. 54 = Rhynchosia glandulosa.
glabrata, Steud. Nom. ed. II. i. 691 = Shuteria vestita.
glandulosa, Thunb. Prod. Pl. Cap. 131 = Rhynchosia glandulosa.
Grahamii, Wall. Cat. n. 5513 = Grona Grahami.
hedysaroides, *Willd. Sp. Pl.* iii. 1062.—Afr. trop.
heloela, Ell. in Journ. Acad. Sc. Philad. i. (1818) 326 = Phaseolus helvolus.
heterocarpa, Hegetschw. Comm. Bot. 9. t. 5 = Amphicarpæa monoica.
heterophylla, Thunb. Prod. Pl. Cap. 131 = Rhynchosia glandulosa.
hispida, *Mœnch. in Bull. Acad. Petersb.* xviii. (1873) 398; *Franch. & Sav. Enum. Pl. Jap.* i. 108.—Japon.
humifusa, Willd. Enum. Hort. Berol. 756 = Rothia trifoliata.
involucrata, Wall. Pl. As. Rar. iii. 22 = Shuteria vestita.
ixosperma, *Hoffmgg. Vers. Fl. Nachtr.* ii. 153.—Bras.
jamaicensis, *Guss. ex Colla, Hort. Ripul. App.* iii. 37.—Jamaic.
javanica, *Linn. Sp. Pl.* 754.—As. et Afr. trop.
javanica, Thunb. in Trans. Linn. Soc. ii. (1794) 340, partim = *G. Soja.*
labialis, Linn. f. Suppl. 325 = Teramnus labialis.
Lagascana, DC. Prod. ii. 241 = Calopogonium galactioides.
Lamarckii, H. B. & K. Nov. Gen. et Sp. vi. 424 = Rhynchosia minima.
lamprocarpa, A. Cunn. ex Benth. Fl. Austral. ii. 202 = Lamprolobium fruticosum.
lanciflola, *Lag. Gen. et Sp. Nov.* 24.—Ins. Canar.

GLYCINE:—
Lantzii, *Basill. in Bull. Soc. Linn Par.* i. (1883) 882. —Madag.
Latrobeana, *Benth. Fl. Austral.* ii. 244.—Austral.
leptocarpa, R. Grah. in Wall. Cat. n. 5515 = Rothia trifoliata.
leucosperma, Desv. Journ. Bot. i. (1814) 78 = Galactia dubia.
lignosa, Turpin, ex Pers. Syn. ii. 301 = Sweetia lignosa.
linearis, Heyne, ex Wall. Cat n. 5578 = Atylosia lineata.
littoralis, Vahl, ex DC. Prod. ii. 385 = Rhynchosia minima.
longicauda, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xvii. (1868) 658.—Afr. trop.
lucida, Blanco, Fl. Filip. ed. I. 578 = Dolichos Lablab.
lucida, Forst. f. Prod. 51 = Strongylodon ruber.
lucida, R. Grah. in Wall. Cat. n. 5511 = Galactia tenuiflora.
Lyallii, *Benth. in Journ. Linn. Soc.* viii. (1865) 266. —Madag.
macrophylla, H. B. & K. Nov. Gen. et Sp. vi. 426 = Rhynchosia reticulata.
macrophylla, *Thunb. in Schum. & Thonn. Beskr. Guin. Pl.* 348.—Afr. trop.
malacophylla, Spreng. Syst. iv. Cur. Post. 270 = Rhynchosia phaseoloides.
marmorata, K. Br. ex Benth. Fl. Austral. ii. 263 = Atylosia marmorata.
Memnonia, DC. Prod. ii. 386 = Rhynchosia Memonia.
mermisdes, Spreng. Syst. iv. Cur. Post. 270 = Rhynchosia menispermoides.
Meyeri, Benth. in Ann. Wien. Mus. ii. (1838) 126 = Teramnus labialis.
micans, *Vatke. & Baker, in Oliver, Fl. Trop. Afr.* ii. 179.—Afr. trop.
micrantha, Hochst. in Flora, xxiv. (1841) 1. Intell. 32 = javanica.
microptera, Zipp. ex Span. in Linnæa, xv. (1841) 195 = Rhynchosia minima.
minima, Willd. Enum. Hort. Berol. 756 = clandestina.
mollis, Hook. Exot. Fl. iii. t. 201 = Rhynchosia phaseoloides?
mollis, Wight & Arn. Prod. 208 = Teramnus labialis.
mollis, Willd. Sp. Pl. iii. 1062 = Rhynchosia mollis.
mollissima, Ell. Sketch, ii. 284 = Rhynchosia mollissima, *Shuttlew.*
monantha, *Zipp. ex Span. in Linnæa*, xv. (1841) 190, nomen.—Ins. Timor.
moniliformis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 211 = javanica.
Monniera, *DC. Prod.* ii. 242.—Afr. austr.
monoica, Linn. Sp. Pl. ed. II. 1028 = Amphicarpæa monoica.
monoica, Schkuhr, in Usteri, Ann. Bot. xii. (1794) 20. t. 2 = Amphicarpæa monoica.
monophylla, Burm. f. Fl. Ind. 161 = Eleiotis cordata.
monophylla, Linn. Mant. i. 101 = Hallia cordata.
monophylla, Nutt. Gen. Am. ii. 115 = Rhynchosia tomentosa, *Hook. & Arn.*
monosperma, Willd. ex Spreng. Syst. iii. 196 = Rhynchosia rufescens.
moniana, *Rafin. Atl. Journ.* 153 (sp dub.).—Am. bor.
moringæfolius, *Delile, in Caill. Voy. Méroé (Cent. Pl. Nub.)*.—Aegypt.
multijuga, Clemen. in Rev. Hortic. (1874) 300 = Wisteria chinensis.
mumularia, Linn. Mant. i. 271 = Rhynchosia Nummularia.
oblonga, Benth. Bot. Voy. Sulph. 84 = Teramnus volubilis.
oxyphylla, R. Grah. in Wall. Cat. n. 5522 = Galactia oxyphylla.
pallens, R. Grah. l. c. n. 5518 = Teramnus labialis.
parviflora, Lam. Encyc. ii. 728 = Teramnus labialis.
parviflora, Zoll. ex Miq. Fl. Ind. Bat. i. ii. 227 = Dumasia glaucescens.
pedunculoris, Muhl. Cat. 64 = Phaseolus helvolus.
pedunculosa, Rafin. in Med. Repos. N. York, v. (1808) 360 = Phaseolus diversifolius.
pentandra, Spreng. in Schrad. Journ. (1800) ii. 197 = Teramnus labialis.
pentandra, Spreng. Syst. iii. 235 = hedysaroides.
pentaphylla, *Delile, in Hook. Kew Journ. iv.* (1852) 344.—Ind. or.
Petitiana, *Schweinf. ex Oliver, Fl. Trop. Afr.* ii. 178.—Afr. trop.

GLYCINE:—
phaseoloides, Sw. Prod. Veg. Ind. Occ. 105 = Rhynchosia phaseoloides.
picta, Vahl, Symb. Bot. ii. 81 = Erioema violaceum.
pondicheriensis, Spreng. Syst. iii. 196 = Rhynchosia rufescens.
procatoria, Humb. & Bonpl. ex Willd. Enum. Hort Berol. 756 = Rhynchosia phaseoloides.
procumbens, Willd. ex Steud. Nom. ed. II. i. 691 = Rhynchosia diffusa.
prostrata, Burm. f. Fl. Cap. Prod. 21 (Quid ?).—Afr. austr.
pubescens, Balb. ex DC. Prod. ii. 386 = Rhynchosia pubescens.
pubescens, Bert. ex DC. l. c. 243 = Chaetocalyx pubescens.
pugiunculus, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 414 = Centrosema virginianum.
pulchella, H. B. & K. Nov. Gen. et Sp. vi. 422 = Eriosema pulchellum.
pulcherrima, Willd. ex Spreng. Syst. iii. 246 = Robinia aturensis.
punctata, Willd. Sp. Pl. iii. 1066 = Poiretia punctata
reflexa, Nutt. Gen. Am. ii. 115 = Rhynchosia minima.
remiformis, Pursh, Fl. Am. Sept. ii. 86 = Rhynchosia tomentosa.
repanda, *Spreng. Fl. Hal. Mant.* 47.—Hab. ?
reticulata, Sw. Prod. Veg. Ind. Occ. 105 = Rhynchosia reticulata.
retusa, Soland. ex Benth. Fl. Austral. ii. 247 = Hardenbergia retusa.
rhombea, Schum. & Thonn. Beskr. Guin. Pl. 346 = Rhynchosia minima.
rhombifolia, Herb. Madr. ex Wall. Cat. n. 5492 = Rhynchosia densiflora.
rhombifolia, Willd. Sp. Pl. iii. 1065 (err. typ. 1605) = Rhynchosia minima.
rosea, Forst. f. Prod. 50 = Canavalia obtusifolia.
rubicunda, Schneev. Ic. Pl. Rar. t. 28 = Kennedya rubicunda.
rufa, H. B. & K. Nov. Gen. et Sp. vi. 423 = Eriosema rufum.
rufa, Schum. & Thonn. Beskr. Guin. Pl. 344 = Eriosema glomeratum.
rufescens, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 222 = Rhynchosia rufescens.
sagittata, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 787 = Centrosema hastatum.
sarmentosa, Roth, Catalect. fasc. ii. 87 = Amphicarpæa monoica.
Schimperi, Hochst. & Steud. ex Steud. Nom. ed. II. i 691 = Rhynchosia Schimperi.
secunda, Thunb. Prod. Pl. Cap. 131 = Rhynchosia secunda.
senegalensis, DC. Prod. ii. 242 = Teramnus labialis.
sericea, *Benth. Fl. Austral.* ii. 245.—Austral.
sericea, *Willd. Sp. Pl.* iii. 1059.—Afr. trop.
simplicifolia, H. B. & K. Nov. Gen. et Sp. vi. 419 = Eriosema simplicifolium.
sinensis, Lindl. in Bot. Reg. t. 650 = Wistaria chinensis.
Soja, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1843) 119.—As. trop.
striata, *Sp. Pl. Ed. Wendeb.* i. 32.—Am. trop.
stricta, Hook. Comp. Bot. Mag. i. (1835) 22 = Galactia sessiliflora.
suaveolens, Linn. f. Suppl. 326 = Rhynchosia suaveolens.
sublobata, Schum. & Thonn. Beskr. Guin. Pl. 347 = Rhynchosia phaseoloides.
subterranea, Linn. Sp. Pl. ed. II. 1028 = Voandzeia subterranea.
suffulta, Wall. Cat. n. 5507 = Shuteria suffulta.
tabacina, *Benth. Fl. Austral.* ii. 244.—Austral.
tomentosa, R. Grah. in Wall. Cat. n. 5520 = Teramnus labialis.
tonsiflora, Klein, in Willd. Sp. Pl. iii. 1059 = Galactia tenuiflora.
tomentosa, *Benth. Fl. Austral.* ii. 245.—Austral.
tomentosa, Linn. Sp. Pl. 754 = Rhynchosia tomentosa.
totta, Thunb. Prod. Pl. Cap. 131 = Rhynchosia Totta.
trifolia, Burm. f. Fl. Ind. 166; Murr. Syst. Veg. ed. XIII. 549 = Phaseolus trilobus.
tuberosa, Salisb. Prod. 335 = Apios tuberosa.
umbellata, Muhl. ex Willd. Sp. Pl. iii. 1058 = Phaseolus helvolus.
unifora, Delz. in Journ. Linn. Soc. xiii. (1873) 146 = Dolichos biflorus.
usaramensis, *Engel & Mœsch, in Tent. Fl. Usamr.* 50.—As. bor. or.

GLYCINE:—
setulina, Bert. ex DC. Prod. ii. 289 = Odonia tomentosa.
vestita, R. Grah. in Wall. Cat. n. 5512 = Shuteria vestita.
villosa, Thunb. Fl. Jap. 288 = Dunbaria subrhombea.
vincentina, [Ker-Gawl.] in Bot. Reg. t. 709 = Chaetocalyx vincentina.
violacea, Schneev. Ic. Fl. Rar. t. 29 = Hardenbergia monophylla.
virens, Soland. ex Stend. Nom. ed. II. i. 692 = Hardenbergia monophylla.
virginica, Spreng. in Schrad. Journ. (1800) ii. 198 = Phaseolus diversifolius.
viscida, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 209 = Pseudarthria viscida.
viscosa, Moench, Meth. Suppl. 46 = Fagelia bituminosa.
viscosa, Roth, Nov. Pl. Sp. 349 = Rhynchosia viscosa.
warrenvis, Dalz. in Hook. Kew Journ. iii. (1851) 210 = Teramnus labialis.

GLYCIPHYLLA, Rafin. in Am. Monthly Mag. (1819) 192 = Chiogenes, Salisb. (Vacciniac.).
hispidula, Rafin. l. c. = C. serpyllifolia.

GLYCOSMA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 639 (1840) = Myrrhis, Scop. (Umbellif.).
ambiguum, A. Gray, in Proc. Am. Acad. viii. (1872) 386.—Am. bor. occ.
Bolanderi, A. Gray, l. c.=M. Bolanderi.
occidentale, Nutt. ex Torr. & Gray, Fl. N. Am. i. 639 = M. occidentalis.

GYCOSMIS, Corres, in Ann. Mus. Par. vi. (1805) 384. *RUTACEAE*, Benth. & Hook. f. i. 308.
CHIONOTRIA, Jack, in Malay. Misc. ii. (1822) vii. 53.
DIOXIPPE, M. Roem. Syn. Hesper. 33, 45 (1846).
MYOSPYRUM, Lindl. Veg. Kingd. ed. III. 617 (1853).
MYXOSPERMUM, M. Roem. Syn. Hesper. 31, 40 (1846).
africana, *Hook. f. ex Benth. in Hook. Niger Fl.* 256.—Afr. trop.
americana, Sagot, ex Oliver, in Journ. Linn. Soc. v. Suppl. ii. (1861) 37 = pentaphylla.
angustifolia, Lindl. in Wall. Cat. n. 6878 = pentaphylla.
arborea, DC. Prod. i. 536 = pentaphylla.
bilocularia, *Thw. Enum. Pl. Zeyl.* 45.—Zeylan.
chlorosperma, *Spreng. Syst. iv. Cur. Post.* 162.—Java.
chylocarpa, Wight & Arn. Prod. 93 = pentaphylla.
citrifolia, Lindl. in Trans. Hort. Soc. vi. (1826) 72 = reutaphylla.
cremulata, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 250 = Murraya cremulata.
cyanocarpa, *Spreng. Syst. iv. Cur. Post.* 161.—Java.
cymosa, *Zipp. ex Span. in Linnaea*, xv. (1841) 178, nomen.—Ins. Timor.
heterophylla, A. Rich. Ess. Fl. Cub. 231 = pentaphylla.
macrocarpa, Wight, Illustr. i. 109 = pentaphylla.
macrophylla, Lindl. in Wall. Cat. n. 6877 = pentaphylla.
macrophylla, *Miq. Fl. Ind. Bat.* i. ii. 522.—Java.
madagascariensis, *Correa, ex Risso Orang.* t. 100.—Madag.!
melanocarpus, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 20.—Hab.!
nitida, Wight & Arn. Prod. 93 = pentaphylla.
obtusa, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 211.—Java.
pentaphylla, *Correa, in Ann. Mus. Par.* vi. (1805) 386.—As trop.; Austral.
puberula, *Lindl. in Wall. Cat.* n. 6875.—Malaya.
pyriformis, *Zipp. ex Span. in Linnaea*, xv. (1841) 178, nomen.—Ins. Timor.
quinquefolia, Griff. Notul. iv. 695 = pentaphylla.
retusa, M. Roem. Syn. Hesper. 41 = pentaphylla.
sphæroidea, *Lindl. in Wall. Cat.* n. 6876.—Malaya.
simplicifolia, *Spreng. Syst. iv. Cur. Post.* 162.—Java.
singuliflora, *Kurz, in Journ. Bot.* xiv. (1876) 37.—Assam.
subrotundifolia, F. Muell. Fragm. i. 26 = Micromelum pubescens.
sinoposita, *Miq. Fl. Ind. Bat. Suppl.* 501.—Sumatra.
trifoliata, *Spreng. Syst. iv. Cur. Post.* 162.—Java.
triphylla, Wight, in Hook. Bot. Misc. iii. (1833) 298 = pentaphylla.

GLYCYCARPUS, Dalz. in Journ. Roy. As. Soc. Bombay, iii. (1849) 69 = Nothopegia, Blume (Anacard.).
edulis, Dalz. l. c. 70 = N. Colebrookiana.
racemosa, Dalz. in Hook. Kew Journ. ii. (1850) 89 = N. Colebrookiana.

GLYCYDERAS, Cass. in Dict. Sc. Nat. lix. 74 (1829). *COMPOSITAE*, Benth. & Hook. f. ii. 259.
GLYPHIA, Cass. in Bull. Soc. Philom. (1818) 141. lucida, *Cass. ex DC. Prod.* vii. 257.—Madag.

GLYCYPHYLLA, Spach, Hist. Vég. Phan. iv. 448 (1840) = GLYCIPHYLLA, Rafin. = Chiogenes, Salisb. (Vaccin.).

GLYCYPHYLLA, Stev. in Bull. Soc. Nat. Mosc. iv. 266 (1832) = Astragalus, Tourn. (Legum.).
galeiformis, Stev. l. c. = A. galeiformis.
glycyphylloides, Stev. l. c. = A. glycyphylloides.
labroderices, Stev. l. c. = A. secundus.
stipularis, Stev. l. c. = A. stipulatus.

GLYCYRRHIZA, Tourn. ex Linn. Cor. Gen. 18 (1787). *LEGUMINOSAE*, Benth. & Hook. f. i. 508.
CLIDANTHERA, R. Br. in Sturt, Exped. Centr. Austral ii. App. 78 (1849).
GLYCYRRHIZOPSIS, Boiss. Diagn. Ser. II. v. 82 (1856).
LIQUIRITIA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 367.
MERISTOTROPIS, Fisch. & Mey. Ind. Sem. Hort. Petrop. ix. 95 (1843).
aculeata, Forsk. Fl. Aegypt. Arab. 135 = Guilandina Bonduc.
aspera, Pall. Reise, i. 499 = asperrima.
asperrima, *Linn. f. Suppl.* 330.—Russia; As. centr.
astragalina, *Gill. in Hook. Bot. Misc.* iii. (1833) 183.—Chili.
brachycarpa, Boiss. Diagn. Ser I. ii. 33 = glabra.
bucharica, *Regel, in Act. Hort. Petrop.* viii. (1884) 697.—As. centr.
Dioscoridis, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 367 = echinata.
dubia, Bernh. ex Steud. Nom. ed. II. i. 692 = foetidissima.
echinata, Lepech. Reise, i. 214 = glabra.
echinata, *Linn. Sp. Pl.* 741.—Europ. austr.; Orient.
flavescens, *Boiss. Diagn.* Ser. I. vi. 33.—Cilicia; Syria.
foetida, *Desf. Fl. Atlant.* ii. 170. t. 199.—Afr. bor.
foetida, Jacq. Eclog. t. 51 = foetidissima.
foetidissima, *Tausch, in Flora*, xiv. (1831) 210.—Hab.!
Frearitis, Orph. ex Nym. Consp. 188 = echinata.
glabra, *Linn. Sp. Pl.* 742.—Reg. Mediterr.; Orient.
glabra, Torr. in Emory, Notes Mil. Reconnois. 408 = lepidota.
glandulifera, Waldst. & Kit. Pl. Rar. Hung. i. 20. t. 21 = glabra.
glutinosa, Nutt. in Torr. & Gray, Fl. N. Am. i. 298 = lepidota.
grandiflora, *Tausch, in Flora*, xiv. (1831) 209.—Hab.!
hirsuta, Hort. Prag. ex Tausch, l. c. = praec.
hirsuta, Linn. Sp. Pl. 742 = glabra.
hispida, Pall. Reise, iii. 754 = asperrima.
hybrida, *Gueldenst. Reise*, i. 195.—Russia.
Kansso, Siebold, in Verh. Batav. Gen. xii. (1830) 58 = Casia Tora.
laevis, Pall. Reise, i. 870 = glabra.
lepidota, *Pursh, Fl. Am. Sept.* ii. 480.—Am. bor. macedonica, *Boiss. & Orph. in Bull. Congr. Bot. Pétersb.* 1869 (1870) 135.—Graecia.
mediterranea, Vall. Fl. Flum. 317; vii. t. 145 = Periandra dulcis.
muricata, Georgi, Beschr. Russ. Reich. Nachtr. 294 = echinata.
officinalis, Lepech. Reise, i. 235 = glabra.
pallida, Boiss. Diagn. II. ii. 29 = glabra.
pallidiflora, *Maxim. Prim. Fl. Amur.* 79.—Reg. Amur.
pancifoliolata, *Hance, in Journ. Bot.* xx. (1882) 299.—China.
psoraloides, *Benth. Fl. Austral.* ii. 225.—Austral.
spinescens, *Sieber, ex Reichb. Fl. Germ. Excurs.* 513.—Europ.

GLYCYRRHIZA:—
squamulosa, *Franch. Fl. David.* i. (1884) 93.—China.
triphylla, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 39.—Reg. Casp.; Persia; Soongar.
undulata, *Ruiz & Pav. ex G. Don, Gen. Syst.* ii. 227.—Peruv.
uralensis, *Fisch. ex DC. Prod.* ii. 248.—Sibir.
violacea, Boiss. Diagn. Ser. II. ii. 28 = glabra.
viscida, *Turcz. ex Bess. in Flora*, xvii. (1834) i. *Beibl.* 10.—Sibir. baical.
viscosa, Turcz. ex Ledeb. Fl. Ross. i. 565 = glabra.
vulgaris, Gueldenst. ex Ledeb. l. c. = glabra.

GLYCYRRHIZOPSIS, Boiss. Diagn. Ser. II. v. 82 (1856) = Glycyrrhiza, Tourn. (Legum.).
flavescens, Boiss. l. c. = Glycyrrhiza flavescens.

GLYPHA, Lour. ex Endl. Gen. 506, in syn. (1838) = Sonsvola, Linn. (Goodeniac.).

GLYPHAEA, Hook. f. in Hook. Ic. Pl. t. 760 (1848). *TILIACEAE*, Benth. & Hook. f. i. 237.
grewioides, *Hook. f. l. c.*—Afr trop.
Montelrol, *Hook. f. Bot. Mag.* t. 5610.—Afr. trop.
tomentosa, *Mast. in Oliver, Fl. Trop. Afr.* i. 367.—Afr. trop.

GLYPHIA, Cass. in Bull. Soc. Philom. (1818) 141 = Glycyderas, Cass. (Compos.).
lucida, Cass. l. c. = Glycyderas lucida.

GLYPHOSPERMA, S. Wats. in Proc. Am. Acad. xvii. (1882-83) 164. *LILIACEAE*.
Palmeri, *S. Wats. l. c.*—Mexic.

GLYPHOSPERMUM, G. Don, Gen. Syst. iv. 195 (1838) = Gentiana, Tourn.
ornatum, G. Don, l. c.—Peruv.
violaceum, G. Don, l. c. = Gent. violacea.

GLYPTOPETALUM, Thw. in Hook. Kew Journ. viii. (1856) 267. t. 7 B. *CELASTRINEAE*, Benth. & Hook. f. i. 351.
grandiflorum, *Bedd. Ic. Pl. Ind. Or.* i. 21.—Zeylan.
sclerocarpum, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 613.—Burma.
zeylanicum, *Thw. in Hook. Kew Journ.* viii. (1856) 268.—Zeylan.

GLYPTOPLEURA, Eaton, in Bot. King's Exped. 207. t. 20 (1871). *COMPOSITAE*, Benth. & Hook. f. ii. 533.
marginata, *Eaton, l. c.*—Utah.
setulosa, *A. Gray, in Bot. Calif.* i. 431.—Utah.

GLYPTOSTROBUS, Endl. Syn. Conif. 89 (1847) = Taxodium, Rich. (Conif.).
heterophyllus, Endl. l. c. 70 = T. heterophyllum.
Horsfieldii, Hort. ex Carr. Conif. ed. II. 677 = Podocarpus cupressina.
pendulus, Endl. Syn. Conif. 71 = T. distichum.
pensilis, *C. Koch, Dendrol.* ii. ii. 191 = T. heterophyllum.

GLYSCOSMIS, D. Dietr. Syn. Pl. ii. 1372 (1840), sphalm. = Glycosmis, Corres (Rutac.).

GMELINA, Linn. Gen. ed. II. 526 (1742). *VERBENACEAE*, Benth. & Hook. f. ii. 1153.
arborea, *Roxb. Hort. Beng.* 46; *Fl. Ind.* iii. 84.—Ind. or.; Malaya.
asiatica, *Linn. Sp. Pl.* 626.—Ind. or.
asiatica, Wall. Cat. n. 1818, partim = villosa.
chinensis, *Benth. Fl. Hongk.* 272.—China.
coromandelina, Burm. f. Fl. Ind. 132 = asiatica.
elliptica, Sm. in Rees, Cyclop. xvi. n. 2 = villosa.
fasciculiflora, *Benth. Fl. Austral.* v. 65.—Austral.
Finlaysoniana, Wall. Cat. n. 6817 = G. Hystrix.
Hystrix, *Schult. ex Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 81.—As. trop. or.
indica, *Burm. f. Fl. Ind.* 132.—Java.
inermis, Blanco, Fl. Filip. ed. I. 498 = villosa.
Leichardtii, *F. Muell. ex Benth. Fl. Austral.* v. 65.—Austral.
lepidota, *Schaff. in Ann. Jard. Buitenz.* i. (1876) 41.—N. Guinea.
lobata, Gaertn. Fruct. i. 268 = asiatica.
macrophylla, *Benth. Fl. Austral.* v. 65.—Austral.

GMELINA:—
macrophylla, *Wall. Cat.* n. 1819.—Ind. or.
oblongifolia, *Roxb. Hort. Beng.* [95]; *Fl. Ind.* iii. 83. —Ind. or.
parvifora, Pers. Syn. ii. 142, sphalm. = seq.
parvifolia, Roxb. Fl. Corom. ii. 32. t. 162 = salatica.
philippensis, *Cham. in Linnaea*, vii. (1832) 109.— Ins. Philipp.
Rheedii, Hook. Bot. Mag. t. 4895 = arborea.
sinuata, *Hort. Herrenh. ex Link, Enum. Hort. Berol.* ii. 128.—Ind. or. ?
speciosissima, D. Don, Prod. Fl. Nep. 104 = Wightia gigantea.
Tacobonia, Buch. Ham. ex D. Don, Prod. Fl. Nep. 105 = Wightia gigantea.
vestita, *Wall. Cat.* n. 1820 (sp. dub.).—Ind. or.
villosa, *Roxb. Hort. Beng.* 46; *Fl. Ind.* iii. 86.—Ind. or.; Malaya.
vitiensis, Seem. Viti, 440 = Vitex vitiensis.

GMELINIA, Spreng. Gen. ii. 481 (1831) = prec.

GNAPHALION, St. Lag. in. Ann. Soc. Bot. Lyon, vii. (1880) 127 = **Gnaphalium**, Linn. (Compos.).
dyssolum, St. Lag. l. c. = Gnaphalium dysodes.

GNAPHALIUM, Linn. Gen. ed. I. 250 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 305.
BELLOA, Remy, in C. Gay, Fl. Chil. iii. 386. t. 88 (1849).
EUCHITON, Cass. in Dict. Sc. Nat. lvi. 215 (1828).
FILAGINELLA, Opiz, Seznam, 44 (1852).
GAMOCHAETA, Wedd. Chlor. And. i. 151 (1855).
MEROPE, Wedd. Chlor. And. i. 160. t. 24 (1855).
OMALOTHECA, Cass. in Dict. Sc. Nat. lvi. 218 (1828).
abyssinicum, Sch. Bip. in Bot. Zeit. iii. (1845) 174 = Helichrysum abyssinicum.
acaule, Sieber, ex Fisch. & Mey. in DC. Prod. vi. 245 = supinum.
Achillese, Sieber, ex DC. l. c. 267 = Petalacte coronata.
achilleoides, Lam. Encyc. ii. 758 = Helichrysum declinatum.
achilleoides, Sieber, ex DC. Prod. vi. 170 = Helichrysum expansum.
aciloge, DC. Prod. vi. 229 = repens.
acuminatissimum, Sch. Bip. in Bot. Zeit iii. (1845) 171 = Helichrysum oxyphyllum.
acuminatum, *Link, Enum. Hort. Berol.* ii. 312.—Afr. austr.
acutatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum acutatum.
adenocarpum, Sch. Bip. l. c. 170 = Helichrysum adenocarpum.
adhaerens, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 244.—N. Zel.
adnatum, Wall. Cat. n. 2948 = Anaphalis adnata.
adnatum, Willd. ex Spreng. Syst. iii. 459 = Helichrysum odoratissimum.
adolensis, Oliver, Fl. Trop. Afr. iii. 338, Index = Achyrocline Schimperi.
adscendens, Thunb. Prod. Pl. Cap. 150 = Helichrysum adscendens.
aegyptiacum, Pers. Syn. ii. 421 = Ifloga Fontanesii.
affine, D. Don, Prod. Fl. Nep. 175 = luteo-album.
affine, *Ure. in Mém. Soc. Linn. Par.* iv. (1826) 610.— Reg. Magell.
africanum, Hoffmgg. Verz. Pfl. 68 = Helichrysum orientale.
alatum, H. B. & K. Nov. Gen. et Sp. iv. 79 = Achyrocline alata.
alausense, Willd. ex Spreng. Syst. iii. 475 = Achyrocline alata.
albescens, *Sw. Prod. Veg. Ind. Occ.* 113.—Ind. occ.
albo-luteum, Roxb. Hort. Beng. 61; Fl. Ind. iii. 425 = luteo-album.
aldnatoides, *Remy, in C. Gay, Fl. Chil.* iv. 283.— Chili.
alienum, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 329 = Antennaria Geyeri.
alliodes, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum alliodes.
alpigenum, C. Koch, in Linnaea, xxiv. (1851) 854 = sylvaticum.
alpigenum, *F. Muell. ex Hook. f. Fl. Tasman.* i. 217. t. 69 A.—Austral.
alpinum, Asso, ex DC. Prod. vi. 270 = Antennaria dioica.

GNAPHALIUM:—
alpinum, Lightf. Fl. Scot. 470. t. 20. f. 2 = supinum.
alpinum, Linn. Sp. Pl. 856 = Antennaria alpina.
alpinum, Willd. Sp. Pl. iii. 1858 = Antennaria carpathica.
alsinoides, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum alsinoides.
alveolatum, Sch. Bip. l. c. 171 = Helichrysum trilineatum.
ambiguum, Guss. ex Tenore, Ind. Sem. Hort. Neap. (1830) 14 = Helichrysum rupestre.
ambiguum, Pers. Syn. ii. 417 = Helichrysum Lamarckii.
americanum, Mill. Gard. Dict. ed. VIII. n. 17 = purpureum.
ammophilum, Wall. Cat. n. 2943 = Ifloga Fontanesii.
amoenum, Hance, in Journ. Bot. vi. (1868) 174 = hypoleucum.
angustifolium, Lam. Encyc. ii. 746 = Helichrysum littoreum.
angustifolium, Loisel. Fl. Gall. ed. II. ii. 224 = Helichrysum angustifolium.
angustissimum, Mill. Gard. Dict. ed. VIII. n. 2 = Helichrysum angustifolium.
antarcticum, *Hook. f. Fl. Antarct.* ii. 310. t. 93.—Ins. Falkland.
antennarioides, *DC. Prod* vi. 224.—Peruv.
aphelexioides, Boj. ex DC. l. c. 207 = Helichrysum aphelexioides.
apiculatum, Labill. Nov. Holl. Pl. ii. 43. t. 188 = Helichrysum apiculatum.
appendiculatum, Schrank, ex DC. Prod. vi. 175 = Helichrysum grandiflorum.
appendiculatum, Thunb. Prod. Pl. Cap. 149 = Helichrysum appendiculatum.
apricum, *Vell. Fl. Flum.* viii. t. 101.—Bras.
aquaticum, Mill. Gard. Dict. ed. VIII. n. 5 = uliginosum.
arabicum, *y. F. Gmel. Syst.* 1216.—Arab.
arenarium, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum filagineum.
araucanum, *Phil. in Anal. Univ. Chil.* (1873) 502.— Chili.
arborescens, Lam. Amoen. Acad. vi. 98 = Anaxeton arboreum.
arborum, Linn. Sp. Pl. ed. II. 1191 = prec.
arenarium, Aubry, ex DC. Prod. vi. 183 = Helichrysum Stoechas.
arenarium, Linn. Sp. Pl. 854 = Helichrysum arenarium.
arenarium, Sibth. & Sm. Fl. Graec. ix. 45. t. 859 = Helichrysum graveolens.
arenarium, Thunb. Fl. Jap. 312 = luteo-album.
argenteum, Mill. Gard. Dict. ed. VIII. n. 14 = Helichrysum foetidum.
argenteum, Poir. Encyc. Suppl. ii. 802 = Helichrysum yuccaefolium.
argentiacum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum argenteum.
Argyrocoma, Sch. Bip. l. c. 169 = Helichrysum ericoides.
argyrophyllum, Sch. Bip. l. c. 171 = Helichrysum argyrophyllum.
argyrosphaerum, Sch. Bip. l. c. 170 = Helichrysum argyrosphaerum.
aristatum, DC. in Wight, Contrib. 21 = Anaphalis aristata.
arizonicum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 3.—Arizona.
armenium, Fisch. & Mey. ex DC. Prod. vi. 183 = Helichrysum armenium.
arnicoides, Lam. Encyc. ii. 762 = luteo-album.
arvense, Linn. Sp. Pl. 856 = Filago arvensis.
ascendens, Spreng. Syst. iii. 474 = Helichrysum adscendens.
asperum, *Pers. Syn.* ii. 420.—Peruv.
asperum, Thunb. Prod. Pl. Cap. 147 = Anaxeton asperum, hirsutum.
astericiflorum, Lam. Encyc. ii. 760 = Evax astericiflora.
asteroides, Balb. Cat. Hort. Taur. (1813) 38 = japonicum.
asteroides, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Ixiolaena viscosa.
atenuatum, *DC. Prod.* vi. 228.—Mexic.
auranticum, Bolss. Diagn. Ser. II. vi. 106 = Helichrysum lavandulaefolium.
aureofulvum, Berg. Desc. Fl. Cap. 257 = Helichrysum odoratissimum.

GNAPHALIUM:—
aureum, Gilib. *Fl.* Lituan. i. 179 = Helichrysum arenarium.
aureum, Hort. ex Spreng. Nov. Prov. 20 = Helichrysum apiculatum.
aureum, Houtt. Handleid. x. 690. t. 67. f. 3 = Helichrysum fulgidum.
auriculatum, Lam. Encyc. ii. 754 = Helichrysum crispum.
auriculatum, *A. Rich. Tent. Fl. Abyss.* i. 420.—Abyss.
auriculatum, *Schrank, in Denkschr. Akad. Munch* viii. (1824) 156 = Helichrysum stellatum.
auriculatum, Thunb. Prod. Pl. Cap. 151 = Helichrysum auriculatum.
auronitens, Sch. Bip. in Bot. Zeit. iii. (1845) 173 = Helichrysum auronitens.
australe, Fisch. ex Steud. Nom. ed. II. i. 693 (Quid?) —Austral.
axillare, *Remy, in C. Gay, Fl. Chil.* iv. 280.—Chili.
badium, *Wedd. Chlor. And.* i. 145.—Bolivia.
balearicum, Pourr. ex DC. Prod. vi. 182 = Helichrysum Lamarckii.
Banksii, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum bractestum.
Baxteri, Sch. Bip. l. c. = Helichrysum Baxteri.
Belangerianum, Sch. Bip. l. c. = Helichrysum leucocephalum.
bellidiflorum, Vir. Fl. Ital. Fragm. 16. t. 19 = Helichrysum frigidum.
bellidioides, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum bellidioides.
Berlandieri, *DC. Prod.* vi. 228.—Mexic.
Berterianum, DC. l. c 233 = purpureum.
bracteatum, Benth. & Hook. f. Gen. ii. 306 = Helichrysum bracteatum.
bicolor, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum bractestum.
Billotii, *F. Schultz, Arch. Fl. Fr. et Allem.* 127.— Europ.
bogotense, H. B. & K. Nov. Gen. et Sp. iv. 78 = Achyrocline bogotensis.
Bojerianum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum Bojerianum.
boreale, Salisb. Prod. 192 = Anaphalis margaritacea.
boreale, Turcz. ex DC. Prod. vi. 270 = Antennaria dioica.
Bourgovii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 3.—Mexic.
brachypterum, *DC. Prod.* vi. 226.—Mexic.
bractatum, Lam. Encyc. ii. 751 = Helichrysum rutilans.
bractestum, *Willd. Sp. Pl.* iii. 1892.—Reg. Argent.
brasilianum, Steud. Nom. ed. II. i. 22, in syn. = Blainvillea balbisiana.
brevicilium, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum semipapposum.
brevifolium, Lam. Encyc. ii. 744 = Metalasia fascicularis.
brevifolium, Thw. Enum. Pl. Zeyl. 166 = Anaphalis brevifolia.
Buchingeri, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop 285 = Helichrysum foetidum.
buchtormense, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum arenarium.
buddleioides, Sch. Bip. l. c. = Helichrysum buddleioides
Burchellii, Sch. Bip. l. c. = Helichrysum Burchellii.
Burmanni, Sch. Bip. l. c. 170 = Helichrysum panduratum.
Buxua, Buch.-Ham. in D. Don, Prod. Fl. Nep. 173 = Anaphalis araneosa.
canephoroum Lam. vide *cespitosum*.
caespitosum, J. & C. Presl, Delic. Prag. 98 = Helichrysum conglobatum.
calabrum, Tenore, Fl. Nep. v. 223 = Filago minima.
calicinum, Poir. Encyc. Suppl. ii. 807 = Chevreulia stolonifera.
californicum, DC. Prod. vi. 234 = decurrens.
callichrysum, Fisch. & Mey. ex DC. Prod. vi. 183 = Helichrysum callichrysum.
callunoides, Sch. Bip. in Bot. Zeit. iii. (1845) 173 = Helichrysum callunoides.
calycinum, Auct. vide *calicinum*.
campestre, Schrank, ex DC. Prod. vi. 229 = candidissimum.
canariense, Willd. ex Spreng. Syst. iii. 470 (= *Schizogyne sericea*).—Ins. Canar.
Candelabrum, Hornem. ex Steud. Nom. ed. II. i. 693, sphalm. ! = *calabrum* !

GNAPHALIUM:—
candicans, H. B. & K. Nov. Gen. et Sp. iv. 79 = Achyrocline satureioides.
candidissimum, Bieb. Tabl. Prov. Casp. 200 = Helichrysum candidissimum.
candidissimum, *Lam. Encyc.* ii. 754.—Afr. austr.
Candolleanum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Ixiolaena leptolepis.
canescens, *DC. Prod.* vi. 228.—Mexic.
canum, Wall. Cat. n. 3942 = Anaphalis Royleana.
capillaceum, Thunb. Prod. Pl. Cap. 152 = Helichrysum capillaceum.
capitatum, *Griseb. in Gentt. Abh.* xxiv. (1879) 186.—Reg. Argent.
capitatum, Lam. Encyc. ii. 744 = Metalasia capitata.
capitatum, Thunb. Prod. Pl. Cap. 148 = Metalasia Cephalotes.
capitellatum, Thunb. l. c. 150 = Helichrysum capitellatum.
carinatum, E. Mey. ex DC. Prod. vi. 196 = Helichrysum Burchellii.
carneum, Lam. Encyc. ii. 741 = Metalasia Cephalotes.
carneum, Schrank, in Denkschr. Akad. Muench. viii. (1824) 149 = Helichrysum stellatum.
carnulicum, Bernh. ex Steud. Nom. ed. II. i. 393 (Quid ?).—Europ.
carolinæ, Schrank, in Denkschr. Akad. Muench. viii. (1824) 159 = candidissimum.
carpathicum, Wahlenb. Fl. Suec. ii. 535 = Antennaria carpathica.
carpetanum, Bolas. & Reut. ex Willk. & Lange, Prod. Fl. Hisp. ii. 62 = sylvaticum.
carrotæ, Steud. Nom. ed. II. i. 393 = *carotæ*.
castaneum, Gilib. Fl. Lituan. i 179 = uliginosum.
catipes, DC. Prod. vi. 236 = Kaoulia catipes.
cauliflorum, Desf. Fl. Atlant. ii. 267 = Iloga Fontanesii.
celsoides, H. B. & K. Nov. Gen. et Sp. iv. 73. t. 329 = Achyrocline celsoides.
cephaloideum, Willd. Enum. Hort. Berol. 868 = japonicum.
Cephalotes, Thunb. Prod. Pl. Cap. 147 = Metalasia Cephalotes.
cerastioides, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum cerastioides.
cernuum, Thunb. Prod. Pl. Cap. 150 = Helichrysum parviflorum.
cespitosum, Lam. Encyc. ii. 742 = Helichrysum cespitosum.
Chamissonis, DC. Prod. vi. 233 = purpureum.
cheiranthemifolium, Pers. Syn. ii. 419 = cheiranthifolium.
cheiranthifolium, Bert. ex DC. Prod. vi. 225 = cymatoides.
cheiranthifolium, *Lam. Encyc.* ii. 752.—Reg. Argent.
chilense, Hook. & Arn. Bot. Beech. Voy. 81; Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum chilense.
chilense, Spreng. Syst. iii. 480 = Sprengelii.
chionanthum, Spreng. ex DC. Prod. vi. 218 = Anaphalis Royleana.
chionosphærum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum chionosphærum.
chlorochrysum, Sch. Bip. l. c. 170 = Helichrysum chlorochrysum.
chrysanthum, Sch. Bip. l. c. 171 = Helichrysum bracteatum.
chrysocephalum, Sch. Bip. l. c. 174 = Helichrysum cymosum.
Chrysocoma, Poir. Encyc. Suppl. ii. 807 = Iloga spicata.
Chrysocoma, Sch. Bip. in Bot. Zeit. iii. (1845) 174 = Helichrysum cymosum.
chrysocomum, Mill. Gard. Dict. ed. VIII. n. 9 = Helichrysum angustifolium.
cicatrisatum, *Vahl, in Hornem. Hort. Hafn.* ii. 801.—Afr. austr.
ciliatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum semipapposum.
cinerascens, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 82.—Austral.
cinereum, *y. F. Gmel. Syst.* 1216.—Arab.
cinnamomeum, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 286 = Anaphalis macrocenos.
cinnamomeum, Wall. Cat. n. 3944 = Anaphalis cinnamomea.
circinnatum, Hilsenb. & Boj. ex Less. in Linnæa, vi. (1831) 324 = Helichrysum mucronifolium.
circhusense, Boj. ex DC. Prod. vi. 198 = Helichrysum circhusense.
circhusii, Hook. & Arn. Bot. Beech. Voy. 81 = Gnaphalothalium.

GNAPHALIUM:—
citrinum, Lam. Fl. Fr. ii. 62 = Helichrysum Stœchas.
citrinum, Schrank, in Denkschr. Akad. Muench. viii. (1824) 152 = Helichrysum parviflorum.
coarctatum, Willd. Sp. Pl. iii. 1886 = spicatum.
cochinchinense, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum cochinchinense.
cochlearifolium, Sch. Bip. l. c. 171 = Helichrysum cochleariforme.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 154.—N. Zel.
collinum, *Labill. Nov. Holl. Pl.* ii. 44. t. 189.—Austral.
collinum, Salisb. Prod. 192 = Helichrysum Stœchas.
Commersonii, Spreng. Syst. iii. 472 = Laulia acutifolia.
comosum, Sch. Bip. in Bot. Zeit. iii. (1845) 173 = Helichrysum ericæfolium.
compressum, Poir. Encyc. Suppl. ii. 804 (= *Schisogyne obtusifolia*).—Ins. Canar.
concinnum, *A. Gray, in Proc. Am. Acad.* xv. (1880) 84.—Mexic.
concolorum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum erosum.
confertum, Benth. in Hook. Lond. Journ. Bot. i. (1842) 488 = hypoleucum.
confine, *Harv. in Harv & Sond. Fl. Cap.* iii. 268.—Afr. austr.
confusum, DC. Prod. vi. 222 = luteo-album.
congestum, Lam. Encyc. ii. 741 = Helichrysum felinum.
conglobatum, Lam. Fl. Fr. ii. 64 = luteo-album.
conglobatum, Viv. Fl. Lib. Spec. 55. t. 8. f. 5 = Helichrysum Fontanesii.
conicum, *Willd. Enum. Hort. Berol. Suppl.* 57.—Europ. austr.
connatum, Spreng. in Flora, xii. (1829) 7 = Leontonyx squarrosus.
conoideum, *H. B. & K. Nov. Gen. et Sp.* iv. 85.—Mexic.
conoideum, Lam. Encyc. ii. 755 = obtusifolium.
consanguineum, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 108.—Ins. Sandvic.
consanguineum, Homb. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zel. iii. = affine.
contortum, Ham. ex Spreng. Syst. iii. 479 = Anaphalis contorta.
conyzoides, *Thunb. Prod. Pl. Cap.* 148.—Afr. austr.
conyzoideum, Lam. Fl. Fr. ii. 63 = Phagnalon sordidum.
coquimbense, *Phil. in Linnæa*, xxix. (1857–58) 5.—Chili.
cordatum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum cordatum.
cordifolium, Boj. ex DC. Prod. vi. 208 = Helichrysum cordifolium.
coriaceum, E. Mey. ex DC. l. c. 167 = Helichrysum rotundatum.
coronatum, Linn. Syst. ed. X. 1210 = Petalacte coronata.
crassifolium, Guss. ex Tenore, Ind. Sem. Hort. Neap. (1830) 14 = Helichrysum Fontanesii.
crassifolium, Lam. Encyc. ii. 746 = Helichrysum Lamarckii.
crassifolium, Linn. Mant. i. 112 = Helichrysum crassifolium.
crassifolium, Lowe, ex DC. Prod. vi. 181 = Helichrysum oboonicum.
crassifolium, Willd. Sp. Pl. iii. 1865 = Helichrysum revolutum.
crassinerve, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum Krebsianum.
crispatulum, C. B. Clarke, Comp. Ind. 115 = pulvinatum.
crispatulum, *Delile, Fl. Egypte*, 267. t. 44. f. 3.—Aegypt. Ind. or.
crispum, Lam. Encyc. ii. 750 = Helichrysum revolutum.
crispum, Linn. Sp. Pl. ed. II. 1197 = Helichrysum crispum.
crispum, Linn. f. Suppl. 368 = Helichrysum undatum.
crispum, Pers. ex DC. Prod. vi. 197 = Achyrocline glumacea.
crispum, Pourr. ex DC. l. c. 182 = Helichrysum cespitosum.
cuneatum, Wall. ex DC. l. c. 270 = Anaphalis triplinervia.
cuneifolium, *y. F. Gmel. Syst.* 1216.—Arab.
cuneifolium, Wall. Cat. n. 3984 = Anaphalis cuneifolia.
Cunninghamii, DC. Prod. vi. 225 = japonicum.
Cupanianum, Guss. ex DC. l. c. 249 = Filago arvensis.

GNAPHALIUM:—
cylindricum, L'Hérit. ex DC. Prod. vi. 268 = Anaxeton recurvum.
cylindricum, Linn. Sp. Pl. ed. II. 1194 = Helichrysum cylindricum.
cylindriflorum, Linn. Pl. Rar. Afr. 19 = præc.
cylindrostachyum, Wall. Cat. n. 3931 = Pterocaulon cylindrostachyum.
cymatoides, *Kunze, ex DC. Prod.* vi. 225.—Chili.
cymosum, Linn. Sp. Pl. 855 = Helichrysum cymosum.
cynoglossoides, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 291 = Helichrysum buddleioides.
cynoglossoides, Trevir. in Nov. Act. Nat. Cur. xiii. i. (1826) 200 = Anaphalis triplinervis.
dasianthum, Steud. Nom. ed. II. i. 393, sphalm. = seq.
dasyanthum, Willd. Enum. Hort. Berol. 865 = Helichrysum maritimum.
decalvans, Forst. ex Hook. f. Fl. N. Zel. i. 188 = keriense !
dealbatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum dealbatum.
dealbatum, Thunb. Prod. Pl. Cap. 149 = candidissimum.
debile, Thunb. l. c. = prostratum.
decipiens, *DC. Prod.* vi. 236.—Hab. ?
declinatum, Linn. f. Suppl. 365 = Helichrysum declinatum.
decorum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum decorum.
decumbens, Lag. Gen. et Sp. Nov. 28. n. 357 = Helichrysum decumbens.
decumbens, Thunb. Prod. Pl. Cap. 148 (= *Trichogyne decumbens*).—Afr. austr.
decurrens, Buch.-Ham. in Wall. Cat. n. 2939 = Anaphalis araneosa.
decurrens, *Ives, in Am. Journ. Sc.* i. (1819) 380.—Am. bor.
decurrens, Linn. Syst. ed. X. 1211 = Conyza decurrens.
decurrens, Schrank, Pl. Rar. Hort. Monac. t. 84 = Pterocaulon virgatum.
dejectum, Jacq. Fragm. 65 = Helichrysum capitellatum ?
densum, Lam. Encyc. ii. 744 = Metalasia fasciculata, Cephalotes.
dentatum, Linn. Sp. Pl. 854 = Pentzia flabelliformis.
denudatum, Linn. f. Suppl. 364 = Helichrysum capillaceum ?
depressum, *Nutt. in Trans. Am. Phil. Soc. N. S.* vii. (1841) 405.—Am. austr.
depressum, Roxb. Hort. Beng. 61; Fl. Ind. iii. 425 = pulvinatum.
depressum, Steud. Nom. ed. I. 375 = luteo-album.
dichotomum, Salisb. Prod. 193 = Filago germanica.
dichotomum, *Willd. ex Spreng. Syst.* iii. 473.—Venezuela.
diffusum, Boj. ex DC. Prod. vi. 207 = Helichrysum eminense.
diffusum, E. Mey. ex DC. l. c. 175 = Helichrysum diffusum.
Dillenii, Schult. ex Steud. Nom. ed. II. i. 393 = purpureum.
diminutivum, *Phil. in Linnæa*, xxxiii. (1864–65) 167.—Chili.
dimorphum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 405 = Antennaria dimorpha.
dioicum, Linn. Sp. Pl. 850 = Antennaria dioica.
distantium, Hill, Herb. Brit. i. 36 = præc.
discmæfolium, Vent. Jard. Malm. 74 = Helichrysum dioamæfolium.
dioicoides, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum dioicoides.
discolor, Sch. Bip. l. c. 171 = Helichrysum appendiculatum.
discolor, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 83.—Peruv.
discolorum, Linn. Sp. Pl. ed. II. 1191 = Helichrysum imbricatum.
discolorum, Thunb. Prod. Pl. Cap. 149 = Helichrysum stellatum.
discolorum, Willd. Sp. Pl. iii. 1855 = Helichrysum felinum.
distans, Schrank, in Denkschr. Akad. Muench. viii. (1824) 160 = Metalasia distans.
Dioynanthus, Rafin. in Am. Monthly Mag. (1819) 195 = Antennaria plantaginea.
divaricatum, Berg. Desc. Pl. Cap. 250 = Helichrysum crispum.
divergens, Thunb. Prod. Pl. Cap. 148 = Metalasia divergens.
Dombeyanum, *DC. Prod.* vi. 235.—Am. austr.

GNAPHALIUM :—
dominguense, *Law. Encyc.* ii. 743.—Ins. S. Doming.
Dontianum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Astephalis canescfolia.
dorotheæflorum, Burch. ex DC. Prod. vi. 186 = Helichrysum umbraculigerum.
drakæformæ, Schrank, in Denkschr. Akad. Muench. viii. (1824) 162 = Helichrysum expansum.
dysodes, Spreng. Syst. iv. 776 = cheiranthifolium.
Echloianum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum cephaloideum.
Ehrenbergianum, *Sch. Bip. ex Klatt, in Linnæa*, xlii. (1878–79) 137.—Mexic.
Einsiedleanum, F. Schultz, in Arch. Fl. 311 = sylvaticum.
elatum, Lam. Encyc. ii. 742 = Helichrysum undatum.
elatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum elatum.
elegans. H. B. & K. Nov. Gen. et Sp. iv. 81 = domingense.
elegans, Presl, Fl. Sicul. p. xxix = pulchrum.
elegans, Salisb. Prod. 192 = Helichrysum orientale.
elegantissimum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum elegantissimum.
elichrysoides, Willd. ex Spreng. Syst. iii. 485 = antennarioides.
Elichrysum, Pall. Reise, iii. 587 = Helichrysum arenarium.
ellipticum, DC. in Wight, Contrib. 21 = Anaphalis oblonga.
elongatum, Hornem. Hort. Hafn. ii. 801 = Helichrysum orientale.
elongatum, Lam. Encyc. ii. 741 = Helichrysum fetidum.
emirnense, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum emirnense.
endeavourense, Sch. Bip. l. c. 171 = Helichrysum collinum.
erectum, *Vell. Fl. Flum.* viii. 98.—Bras.
ericæfolium, Salisb. Prod. 192 = Helichrysum ericæfolium.
ericoides, Lam. Encyc. ii. 745 = Helichrysum teretifolium.
ericoides, Linn. Pl. Rar. Afr. 19 ; Amoen. Acad. vi. 99 = Helichrysum ericæfolium.
Ernestianum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum Ernestianum.
erythractis, *Griseb. in Abh. Gostt.* xxiv. (1879) 186.—Reg. Argent.
evacifforme, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum sessile.
evacoides, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
excisum, Thunb. Prod. Pl. Cap. 151 = Helichrysum excisum.
eximium, Linn. Mant. ii. 573 = Helipterum eximium.
expansum, Thunb. Prod. Pl. Cap. 149 = Helichrysum expansum.
factidum, Burm. f. Fl. Cap. Prod. 95, vide *fœtidum*.
falcatum, Lam. Encyc. ii. 755 = purpureum.
fascicalatum, *Y. Buch. in Trans. N. Z. Inst.* lx. (1877) 590.—N. Zel.
fasciculatum, Lam. Encyc. ii. 745 = Metalasia muricata.
fasciculatum, Thunb. Prod. Pl. Cap. 148 = Metalasia fasciculata.
fastigiatum, Schrank, in Denkschr. Akad. Muench. viii. (1824) 148 = Helichrysum foliosum.
fastigiatum, Thunb. Prod. Pl. Cap. 148 = Metalasia muricata.
foliosum, Thunb. l. c. 149 = Helichrysum foliosum.
fernandezianum, *Phil. in Anal. Univ. Chil.* (1856) 167.—Ins. J. Fernand.
ferrugineum, Pers. Syn. ii. 430 = Achyrocline satureioides.
filaginem, *DC. Prod.* vi. 234.—Am. austr.
filaginoides, Boj. ex DC. Prod. vi. 219 = Stœnocline filaginoides.
filaginoides, Hook. & Arn. Bot. Beech. Voy. 359 = Filago californica.
Filago, Gueldenst. ex Ledeb. Fl. Ross. ii. 617 = Filago minima.
filicaula, *Hook. f. Handb. N. Zeal. Fl.* 153.—N. Zel.
filiforme, Liljebl. Svensk. Fl. ed. II. 389 ; ex Wahlenb. Fl. Lapp. 204 = supinum.
flaccidum, *Kunz, ex C. B. Clarke, Comp. Ind.* 290.—Ind. or. ; Burma.

GNAPHALIUM :—
flaccidum, Weinm. in Flora, iii. (1820) 610 = Achyrocline satureioides.
flaminianum, Sieber, ex Benth. Fl. Austral. iii. 624 = Helichrysum apiculatum.
flemmonum, E. Mey. ex DC. Prod. vi. 204 = Helichrysum marginatum.
floccosum, DC. l. c. 234 = indicum.
floridum, Poir. Encyc. Suppl. ii. 807 = Anaxeton arboreum.
fœtidum, Linn. Sp. Pl. 851 = Helichrysum fœtidum.
foliosum, *Phil. in Linnæa*, xxxii. (1864–65) 163.—Chili.
foliosum, Poir. Encyc. Suppl. ii. 804 = Helichrysum cespitosum.
follicilatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum appendiculatum.
formosissimum, Sch. Bip. l. c. 173 = Helichrysum elegantissimum.
Forskåhlii, J. F. Gmel. Syst. 1214 ; Racusch. Nom. ed. III. 238 = kurmeae.
fragrans, Sieber, ex DC. Prod. vi. 174 = Helichrysum stellatum.
frigidum, Poepp. ex Poepp. & Endl. Nov. Gen. et Sp. iii. 51 = Lucilia chilensis.
frigidum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum frigidum.
frigidum, *Wedd. Chlor. And.* i. 147.—Peruv.
fruticans, Linn. Mant. ii. 282 = Helichrysum fruticans.
fruticans, Schrank, in Denkschr. Akad. Muench. viii. (1824) 154 = Helichrysum foetidum.
fruticans, Willd. Sp. Pl. iii. 1851 = Helichrysum grandiflorum.
fruticosum, Forsk. Fl. Aegypt. Arab. 218 = Helichrysum cymosum.
fruticosum, Mill. Gard. Dict. ed. VIII. n. 20 = Helichrysum orientale.
fuchsii, Zuccagni, in Roem. Collect. 158 = Helichrysum fulgidum.
fulvescens, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum fulvescens.
fulvo-aureum, Sch. Bip. l. c. = Helichrysum microcephalum.
fulvum, Lam. Encyc. ii. 743 = Helichrysum stellatum.
fuscatum, Pers. Syn. ii. 431 = norvegicum.
fuscatum, Schur, Enum. Pl. Transs. 319 = sylvaticum.
fuscum, Lam. Encyc. ii. 757 = norvegicum.
fuscum, Oed. in Fl. Dan. t. 324 = sylvaticum.
fuscum, Scop. Fl. Carn. ed. II. ii. 152 = supinum.
fuscum, Vis. ex Nym. Consp. 388 = Hoppeanum.
gallicum, Lam. Sp. Pl. 857 = Filago gallica.
gallicum, Wohlf. Fl. Hal. Suppl. 162 = Filago montana.
garipinum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum garipinum.
Gaudichaudianum, *DC. Prod.* vi. 226.—Bras.
Gayanum, *Remy, in C. Gay, Fl. Chil.* iv. 225.—Chili.
Georgii, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum gracile.
gerberæfolium, Sch. Bip. l. c. 174 = Helichrysum gerberæfolium.
germanicum, Linn. Sp. Pl. 857 = Filago germanica.
glabratum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum brachteatum.
glabrescens, Schrank, in Denkschr. Akad. Muench. viii. (1824) 148 = Metalasia divergens.
glabrum, Eckl. ex DC. Prod. vi. 200 = Helichrysum nudifolium.
glanduliferum, *Griseb. in Gostt. Abh.* xxiv. (1879) 186.—Reg. Argent.
glandulifyrum, Sch. Bip. in Bot. Zeit. iii. (1845) 173 = Helichrysum glandulifyrum.
glandulosum, *Klatt, in Linnæa*, xlii. (1878) 139.—Peruv.
globosum, *Desf. Cat. Hort. Par.* ed. III. 299.—Austral.
globosum, Hornem. Hort. Hafn. 971 = japonicum.
globosum, Sch. Bip. in Bot. Zeit. iii. (1845) 174 = Helichrysum globosum.
globuliferum, E. Mey. ex DC. Prod. vi. 178 = Helichrysum serpyllifolium.
glomeratum, Hort. ex DC. l. c. 236 = japonicum.
glomeratum, Linn. Pl. Rar. Afr. 20 ; Amoen. Acad. vi. 99 = Leontonyx glomeratus.
glomeratum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 362.—Afr. austr.

GNAPHALIUM :—
glumaceum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Achyrocline glumacea.
glutinosum, Tenore, Cat. Sem. Hort. Neap. (1830) 14 = Helichrysum rupestre.
gossypinum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 403 = palustre.
gracile, Blume, Bijdr. 900 = luteo-album.
gracile, *H. B. & K. Nov. Gen. et Sp.* iv. 831.—Mexic.
gracillimum, *Porr. ex DC. Prod.* vi. 232.—Afr. trop.
grandicorpe, *Hook. f. Handb. N. Zeal. Fl.* 154.—N. Zel.
grandiflorum, Linn. Sp. Pl. 850 = Helichrysum grandiflorum.
grandiflorum, Willd. Sp. Pl. iii. 1851 = Helichrysum frutícans.
graveolens, Bieb. Fl. Taur. Cauc. ii. 299 = Helichrysum graveolens.
graveolens, Fenzl, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 316 = Helichrysum orientale.
graveolens, Henning, in Mém. Soc. Nat. Mosc. vi (1823) 79 = Helichrysum arenarium.
graveolens, H. B. & K. Nov. Gen. et Sp. iv. 81 = chrichanthifolium.
Gunnianum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum scorpioides.
gymnocephalum, DC. Prod. vi. 235 = collinum.
gymnocomum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum gymnocomum.
Hamiltonii, Wall. Cat. n. 2936 = Blumea hieracifolia.
hamulosum, E. Mey. ex DC. Prod. vi. 192 = Helichrysum hamulosum.
haplostephioides, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 139 = Chionolæna Arbuscula.
hebelepis, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum hebelepis.
helianthemifolium, Burm. f. Prod. Fl. Cap. 25 = Helichrysum serpyllifolium.
helianthemifolium, *Linn. Sp. Pl.* 851.—Afr. trop.
helianthemifolium, E. Mey. ex DC. Prod. vi. 206 = Helichrysum capitellatum.
helianthemifolium, Thunb. Fl. Cap 650 = Helichrysum stellatum.
helichrysoides, *Dall, in Journ. Bot.* xi. (1873) 364.—Marocc.
helichrysoides, *Wedd. Chlor. And.* i. 146.—Peruv.
helictropifolium, Lam. Encyc. ii. 748 = Helichrysum heliotropifolium.
hemisphæricum, Hornem. Hort. Hafn. ii. 802 = japonicum.
hemisphæricum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum hemisphæricum.
hermannioides, Sch. Bip. l. c. 169 = Helichrysum hermannioides.
heteranthum, Rafin. Specc. ii. 170 ; ex Bertol. Fl. Ital. ix. 163 = Filago Cupaniana.
heteroides, *Klatt, in Linnæa*, xlii. (1878–79) 137.—Mexic.
heterophyllum, Thunb. Prod. Pl. Cap. 149 = Helichrysum teretifolium.
heterotrichum, *Phil. in Linnæa*, xxix. (1857–58) 4.—Chili.
hirsutum, Thunb. Prod. Pl. Cap. 148 = Anaxeton hirsutum.
hirsutum, *Vell. Fl. Flum.* viii. t. 99.—Bras.
hirtum, *H. B. & K. Nov. Gen. et Sp.* iv. 82.—Mexic.
hispidum, Thunb. Prod. Pl. Cap. 148 = Elytropappus spicuiflosus.
Hochstetteri, Sch. Bip. in Bot. Zeit. iii. (1845) 174 = Achyrocline Hochstetteri.
Hookerianum, Sch. Bip. l. c. 171 = Helichrysum buddleioides.
Hoppeanum, *Koch, Syn. Fl. Germ.* ed. II. 389.—Europ.
horridum, Sch. Bip. in Bot. Zeit. iii. (1845) 173 = Helichrysum horridum.
humifusum, Paill. ex Nym. Consp. 388 = uliginosum.
humile, Thunb. Prod. Pl. Cap. 151 = Helichrysum appendiculatum.
humillimum, Spreng. Syst. iii. 476 = nanum.
hurdanicum, Wall. Cat. n. 2951 = Filago germanica.
hyberbrowa, Winch, ex DC. Prod. vi. 270 = Antennaria dioica.
hyemale, Walt. Fl. Carol. 208 = purpureum.
hypoleucum, *DC. in Wight, Contrib.* 21.—As. trop.
hypoleucum, Spreng. ex DC. Prod. vi. 271 = Anaphalis cinnamomea.

GNAPHALIUM :—

ignescens, Linn. Sp. Pl. 854 = Helichrysum arenarium.
Illaplelmum, *Phil. in Linnæa*, xxxiii. (1864–65).—Chili.
imbricatum, Linn. Sp. Pl. 856 = Helichrysum imbricatum.
incanum, *Burm. f. Fl. Cap. Prod.* 95.—Afr. austr.
incanum, H. B. & K. Nov. Gen. et Sp. iv. 80 = Kunthianum.
incarnatum, Bory, ex DC. Prod. vi. 178 = Helichrysum yuccæfolium.
incarnatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum rubellum.
indlenm, *Linn. Sp. Pl.* 852.—Geront. trop.
indicum, Thw. Enum. Pl. Zeyl. 166 = Anaphalis oblonga.
indutum, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 131.—Austral.
inneminatum, Wall. ex DC. Prod. vi. 278 = Anaphalis Notoniana.
inornatum, *DC. Prod.* vi. 235.—Mexic.
inselare, *Phil. in Bot. Zeit.* xiv. (1856) 645.—Ins. J. Fernand.
intermedium, Sch. Bip. l. c. iii. (1845) 170 = Helichrysum paniculatum.
intermedium, Wall. Cat. n. 2986 = Anaphalis nubigena.
intricatum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum intricatum.
involucratum, Forst. f. Prod. 291 = japonicum.
involucratum, Lam. Encyc. ii. 761 = bracteatum
italicum, Roth, in Roem. & Ust. Mag. iv. x. (1790) 19 = Helichrysum angustifolium.
itinerari, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) 1. Intell. 37.—Abyssin.
japonicum, *Thunb. Fl. Jap.* 311.—Cosmop.
javanicum, Reinw. ex Blume, Bijdr. 899 = Antennaria javanica.
jucundum, DC. Prod. vi. 222 = luteo-album.
Jungbuhnianum, *Miq. Fl. Ind. Bat.* ii. 91.—Java.
kerlense, *A. Cunn. ex Hook. f. Fl. N. Zel.* i. 188.—N. Zel.
Krausii, Sch. Bip. in Bot. Zeit. iii (1845) 178 = Helichrysum Krausii.
Krebsianum, Sch. Bip. l. c. 172 = Helichrysum Krebsianum.
Kunthianum, *Steud. Nom.* ed. II. i. 994.—Peruv.
kurmense, *Mart. in Denkschr. Akad. Muench.* vi. (1820) 198.—Arab.
lacteum, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 276.—Peruv.
laevissimum, Schur, Enum. Pl. Transs. 319 = ulliginosum.
Lagopus, Steph. ex Willd. Sp. Pl. iii. 1897 = Filago arvensis.
Lamarckii, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum Lamarckii.
Lambertianum, Sch. Bip. l. c. 171 = Helichrysum Lambertianum.
lanatum, Forst. f. Prod. 55 = japonicum.
lanciifolium, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum lanciifolium.
Landbeckii, *Phil. in Linnæa*, xxxiii. (1864–65) 165.—Chili.
lanuginosum, *H. B. & K. Nov. Gen. et Sp.* iv. 84.—Peruv.
lasiocalion, *Link, Enum. Hort. Berol.* ii. 317.—Afr. austr.
latifolium, Thunb. Prod. Pl. Cap. 153 = Helichrysum latifolium.
lavandulaceum, DC. Prod. vi. 227 = Chionolaena lavandulifolia.
lavandulæfolium, Willd. Sp. Pl. iii. 1868 = Helichrysum lavandulæfolium.
lavanduloides, Bcj. ex DC. Prod. vi. 207 = Helichrysum lavanduloides.
lavarum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum ericæfolium.
leptilimum, Gaertn. Fruct. ii. 391 = Diotis candidissima.
leiolepis, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum revolutum.
leiopodium, Sch. Bip.! c. = Helichrysum leiopodium.
Leontopodium, Sch. Bip. l. c. 169 = Helichrysum Leontopodium.
leontopodioides, Willd. Sp. Pl. iii. 1888 = Leontopodium alpinum.

GNAPHALIUM :—

Leontopodium, Linn. Sp. Pl. 855 = Leontopodium alpinum.
Leontopodium, Steud. Nom. ed. II. i. 694 = Leontopodium sibiricum.
leptolepis, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Isiolæna leptolepis.
leptophyllum, *DC. Prod.* vi. 236.—Mexic.
leptotrichum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum leptorhizum.
leucocephalum, *A. Gray, Pl. Wright. ii.* 99.—Am. bor. occ.
leucocephalum, *Phil. in Linnæa*, xxxiii. (1864–65) 166.—Chili.
leucophyllum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum crassifolium.
leucophyllum, *Schott & Kotschy, ex Boiss. Fl. Orient.* iii. 337.—Asa. Min.
leucopsideum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum leucopsideum.
leysseroides, Desf. Fl. Atlant. ii. 267 = Leyssera capillifolia.
linearc, Sch. Bip in Bot. Zeit. iii. (1845) 169 = Helichrysum ericæfolium.
littorale, Banks & Soland. ex Hook. f. Fl. Antarct. 310 = purpureum.
longifolium, Blume, Bijdr. 900 = Anaphalis longifolia.
longifolium, Sch. Bip. in Bot. Zeit. iii. (1845) 173 = Helichrysum longifolium.
Lorentii, Sch. Bip, l. c. 174 = Helichrysum armenium.
lorreus, Willd. ex Spreng. Syst. iii. 470 = Achyrocline celosioides.
lupulaceum, Lam. Encyc. ii. 752 = Helichrysum appendiculatum.
luteo-album, *Linn. Sp. Pl.* 851.—Cosmop.
luteo-fuscum, *Webb, in Hook. Niger Fl.* 143.—Ins. Cap. Verd.
luzulæfolium, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum luzulæfolium.
luteolides, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 149 = Achyrocline luteolides.
Lyallii, *Hook. f. Fl. N. Zel.* 187.—N. Zel.
lycopodioides, Urv. in Mém. Soc. Linn. Par. iv. (1826) 610 = Chevreulia lycopodioides.
Lyoopsis, *Pers. Syn.* ii. 429.—Reg. Magellan.
Macella, *Veil. Fl. Flum.* viii. t. 96.—Bras.
macilentum, E. Mey. ex DC. Prod. vi. 229 = candidissimum.
macranthum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum bracteatum.
macrocephalum, Sch. Bip. l. c. = Helichrysum bracteatum.
maculatum, Thunb. Prod. Pl. Cap. 149 = candidissimum.
madagascariense, Poir. Encyc. Suppl. ii. 804 = Helichrysum madagascariense.
magellanicum, *Sch. Bip. in Flora*, xxxviii. (1855) 117. —Reg. Magellan.
Mandonii, *Sch. Bip. in Linnæa*, xxxiv. (1865–66) 582, nomen.—Columbia.
marcescens, Wight, Ic. t. 1115 = Anaphalis marcescens.
margaritaceum, Linn. Sp. Pl. 850 = Anaphalis margaritacea.
marginatum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum marginatum.
marsilolium, Sch. Bip, l. c. 171 = Helichrysum marsilolium.
martinense, Linn. Mant. ii. 268 = Helichrysum martinense.
martinum, Schrank, in Denkschr. Akad. Muench. viii. (1820) 153 = Helichrysum odoratissimum.
martabanicum, Wall. Cat. n. 2950 = luteo-album.
medium, Vill. Prosp. 31 = norvegicum.
melanacme, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum melanacme.
melanophthalmum, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 99 = Helichrysum melanophthalmum.
melanospermoides, *Sch. Bip. in Benplandia*, iv. (1856) 54, nomen.—Peruv.
melanospermum, *Sch. Bip. in A. Rich. Tent. Fl. Abyss.* i. 481.—Abyssin.
membranum, Phil. in Anal. Univ. Chil. (1870) 184 = chelranthifolium.
metalasioides, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum ericæfolium.
microlæfolium, Sch. Bip. l. c. 171 = Helichrysum microlæfolium.
microanthum, *Thunb. Prod. Pl. Cap.* 149.—Afr. austr.
microcephalum, A. Gray, Pl. Wright. i. 124; ii. 99 = Wrightii.

GNAPHALIUM :—

microcephalum, *Nutt. in Trans. Am. Phil. Soc. N. S.* vii. (1841) 404.—Am. bor. occ.
microlepis, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum semipapposum.
microphyllum, Willd. Sp. Pl. iii. 1863 = Helichrysum angustifolium.
micropoides, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum micropoides.
millefolium, Herb. Willd. ex DC. Prod. vi. 190 = Helichrysum cylindricum.
millefolium, Linn. f. Suppl. 362 = Helipterum phlomoides.
minimum, Bieb. Fl. Taur. Cauc. ii. 301 = Filago arvensis.
minimum, Sm. Fl. Brit. ii. 873 = Filago minima.
modestum, Hook. Bot. Mag. t. 2710 = Helipterum gnaphaloides.
molle, Salisb. Prod. 192 = luteo-album.
molle, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helipterum incanum.
molle, Thunb. Prod. Pl. Cap. 150 = Helichrysum maritimum.
monanthon, Willd. ex DC. Prod. vi. 269 = Antennaria alpina.
monanthon, Willd. ex Spreng. Syst. iii. 470 = Helichrysum callichrysum?
monocephalum, Carp. ex Torr. & Gray, Fl. N. Am. ii. 310 = Antennaria plantaginea.
montanum, Huds. Fl. Angl. ed. II. ii. 362 = Filago minima.
montanum, Linn. Sp. Pl. 857; Hill, Veg. Syst. iii. 142 = Filago arvensis.
montanum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum splendidum.
montevidense, Spreng. Syst. iii. 475 = cheiranthifolium.
moschatum, A. Cunn. in DC. Prod. vi. 236 = Helipterum moschatum.
mucronatum, Berg. Desc. Pl. Cap. 269; Linn. Mant. ii. 283 = Helichrysum mucronatum.
multicaule, Lam. Encyc. ii. 748 = Helichrysum yuccæfolium.
multicaule, Willd. Sp. Pl. iii. 1888 = indicum.
multiceps, Wall. Cat. n. 2949 = luteo-album.
multiflorum, Poir. Encyc. Suppl. ii. 803 (= *Schizogyne obtusifolia*).—Ins. Canar.
multiflorum, Willd. Sp. Pl. iii. 1900 = Helichrysum crispum.
multinerve, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum quinquenerve.
muricatum, Linn. Sp. Pl. 859 = Metalasia aduncs.
muricatum, Thunb. Prod. Pl. Cap. 146 = Metalasia muricata.
muscoides, Desf. Fl. Atlant. ii. 267 = Lasiopogon lanatum.
mutabile, Rochel, Pl. Banat. Rar. Ind. = sylvaticum, supinum.
nanum, *H. B. & K. Nov. Gen. et Sp.* iv. 84.—Peruv.
nanum, Willd. Sp. Pl. iii. 1898 = Leontonyx psallius.
natalitium, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum natalitium.
neilgerryanum, DC. in Wight, Contrib. 21 = Anaphalis neilgerryana.
neglectum, Soyer-Willem. in Mém. Soc. Nanc. (1836) 45 = Filago neglecta.
nepalense, Hort. ex DC. Prod. vi. 270 = Anaphalis triplinervis.
nevadense, *Kellogg, in Proc. Calif. Acad.* v. (1873) 45. —Calif.
nilagiricum, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 287 = Anaphalis neilgerryana.
nitiacum, Reddi, ex Spreng. Syst. iii. 480 = indicum.
nitens, Willd. ex Spreng. l. c. = roseum.
nitidulum, Hook. f. Handb. N. Zeal. Fl. 154 = Raoulia Planchoni?
nitidum, *Hornem. Enum. Pl. Hort. Hafn.* (1807) 18. —Hab.?
niveum, Tenore, Prod. Fl. Nap. 48 = Leontopodium alpinum.
niveum, Juss. fide Cass. in DC. Prod. vi. 426 = Senecio Aspelina.
niveum, Linn. Sp. Pl. 852 = Helichrysum parvifloram.
niveum, Poir. Encyc. Suppl. ii. 808 = Helichrysum desiltatum.
niveum, Thunb. Prod. Pl. Cap. 150 = Helichrysum anomalum.
nodiflorum, Lam. Encyc. ii. 754 = Helichrysum nodiflorum.

GNAPHALIUM:—
norvegicum, Gunn. Fl. Norv. ii. 105.—Europ.; As. bor.; Am. bor.
norvegicum, Retz. Fl. Scand. Prod. 156 = sylvaticum.
notatum, Thunb. Prod. Pl. Cap. 150 = Helichrysum expansum.
notatum, Willd. ex DC. Prod. vi. 190 = Helichrysum cylindricum.
Natoniensum, Wall. Cat. n. 2952 = Anaphalis Natoniana.
Novæ-Zealandiæ, Sch Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum micranthum.
nubicum, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 149 = G. Unionis.
nubigenum, Wall Cat. n. 2985 = Anaphalis nubigena.
nudifolium, Linn. Pl. Rar. Afr. 19 = Helichrysum nudifolium.
nudum, Hoffm. ex J. F. Gmel. Syst. 1217 = uliginosum.
obconicum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum obconicum.
oblongum, Thw. Enum. Pl. Zeyl. 422 = Anaphalis oblonga.
obovatum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum obovatum.
obtusifolium, *Linn. Sp. Pl.* 851.—Am. bor.
obtusifolium, Willd. ex Spreng. Syst. iii. 478 = purpureum.
obvallatum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum obvallatum.
Oculus, Thunb. Prod. Pl. Cap. 151 = seq.
Oculus-cati, Linn. f. Suppl. 364 = Eriophaera Oculus-cati.
odoratissimum, Linn. Sp. Pl. 855 = Helichrysum odoratissimum.
odoratum, Thunb. Prod. Pl. Cap. 150 = Helichrysum rutilans.
odorum, Salisb. Prod. 192 = Helichrysum odoratissimum.
odorum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum apiculatum.
oliganthum, *Phil. in Linnæa*, xxxiii. (1864–65) 167.—Chili.
oligocephalum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum armenium.
olympicum, Sibth. ex Spreng. Syst. iii. 477 = Helichrysum Billardieri.
orbiculare, Thunb. Prod. Pl. Cap. 152 = Helichrysum serpyllifolium.
orientale, Linn. Sp. Pl. 858 = Helichrysum orientale.
orixense, Roxb. Hort. Beng. [101]: Fl. Ind. iii. 425 = luteo-album.
ovatum, Desf. Tabl. ed. I. 96 = Helichrysum hemisphæricum.
oxybolium, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum oxybolium.
oxyphyllum, *DC. Prod.* vi. 295.—Mexic.
oxyphyllum, Steetz, ex Grisch. Fl. Brit. W. Ind. 380, partim = domingense.
Pallasii, Spreng. Syst. iii. 470 = Helichrysum callichrysum.
pallens, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum undatum.
pallidum, Lam. Encyc. ii. 750 = luteo-album.
palustre, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 408.—Am. bor. occ.
panaricioides, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Leptorhynchus panaetioides.
panduræfolium, Schrank, in Denkschr. Akad. Muench. viii. (1824) 169 = Helichrysum auriculatum.
paniculatum, Berg. Desc. Pl. Cap. 256 = Helichrysum expansum.
paniculatum, Colla, in Mem. Acc. Tor. xxxviii. (1835) 17 = chrantifolium.
paniculatum, Thunb. Fl. Cap. 647 = Helichrysum cylindricum.
pannosum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 2.—Mexic.
pannosum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum pannosum.
papillosum, Poir. Encyc. Suppl. ii. 608 = Helichrysum bracteatum.
paronychioides, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum paronychioides.
parviflorum, *Cadenas, in Trans. N. Z. Inst.* xvi. (1884) 333.—N. Zel.
parviflorum, Lam. Encyc. ii. 748 = Helichrysum parviflorum.

GNAPHALIUM:—
parvulum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 262.—Afr. austr.
patulum, Berg. Desc. Pl. Cap. 249 = Helichrysum auriculatum.
patulum, Linn. Sp. Pl. 855 = Helichrysum patulum.
patulum, Thunb. Prod. Pl. Cap. 151 = micranthum, spathulatum.
pauciflorum, *DC. Prod.* vi. 239.—Afr. austr.
pedunculare, Linn. Mant. ii. 288 = Helichrysum rupestre.
pedunculare, Tenore, Ind. Sem. Hort. Nesp. (1830) 14 = Filago arvensis.
pedunculatum, *Benth. & Hook. f. ex Klatt, in Linnæa*, xlii. (1878-79) 143.—Chili?
pellitum, Hort. Gotting. ex DC. Prod. vi. 225 = Schraderi.
pellitum, *H. B. & K. Nov. Gen. et Sp.* iv. 80.—Peruv.
pendulum, Presl, Delic. Prag. 97 = Helichrysum rupestre.
perlanatum, Willd. Enum. Hort. Berol. 867 = purpureum.
pentzioides, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum pentzioides.
perfoliatum, Wall. Cat. n. 2987 = Anaphalis triplinervis.
perpetuum, *Phil. in Linnæa*, xxix. (1857–58) 6.—Chili.
peruvianum, Spreng. Syst. iii. 473 = asperum.
petiolatum, Linn. Sp. Pl. 854 = Helichrysum petiolatum.
petiolatum, Thunb. Prod. Pl. Cap. 152 = Helichrysum fruticans.
Petitianum, *A. Rich. Tent. Fl. Abyss.* i. 429.—Abyssin.
phaseolepis, *Phil. in Linnæa*, xxxiii. (1864–65) 168.—Chili.
phlomoides, Lam. Encyc. ii. 740 = Heliptertum phlomoides.
phylicæfolium, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum phylicæfolium.
pilosellum, Linn. f. Suppl. 364 = Helichrysum pedunculare.
pilulare, Wahlenb. Fl. Lapp. 205. t. 18 = uliginosum.
pilulifferum, Hort. ex DC. Prod. vi. 166 = Helichrysum dioniæfolium.
piptolepis, *Griseb. in Gortt. Abh.* xxiv. (1879) 186.—Reg. Argent.
Pisavira, Less. in Linnæa, vi. (1831) 227 = G. Viravira.
Planchonii, Hook. f. Fl. Tasman. i. 217. t. 62 = Raoulia Planchonii.
planifolium, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum argenteum?
plantagineum, Burm. f. Fl. Cap. Prod. 25 = Helichrysum latifolium.
plantagineum, Linn. Syst. ed. XII. 545 = Antennaria plantaginea.
plantaginifolium, Linn. Sp. Pl. 850 = præc.
Plantago, Boj. ex DC. Prod. vi. 200 = Helichrysum Plantago.
platypterum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum platypterum.
plebeium, Sch. Bip. l. c. 172 = Helichrysum plebeium.
plenum, Burm. f. Fl. Cap. Prod. 25 = Helichrysum stellatum.
plicatum, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1836) 387 = Helichrysum plicatum.
pluricaule, Poir. Encyc. Suppl. ii. 607 = indicum.
Poeppigianum, Benth. Fl. Hartw. 207 = domingense.
Poeppigianum, *DC. Prod.* vi. 227.—Peruv.
polifolium, Thunb. Prod. Pl. Cap. 151 = Helichrysum serpyllifolium.
Pollinii, *Vestd. Chlor. And.* i. 147.—Peruv.
polyanthus, Thunb. Prod. Pl. Cap. 147 = Metalasia auricasta, octoflora.
polyanthus, Thunb. l. c. 151 = Helichrysum crispum.
polycaulon, Pers. Syn. ii. 421 (policaulon) = indicum.
polycephalum, Michx. Fl. Bor. Am. ii. 127 = obtusifolium.
polycephalum, Spreng. ex DC. Prod. vi. 238 = domingense.
polycephalum, Wall. ex DC. l. c. v. 455 = Pterocaulon cylindrostachyum.
polycephalum, Willd. ex Spreng. Syst. iii. 478 = Helichrysum nodiflorum.
pompejanum, Tenore, Cat. Pl. Hort. Neap. (1819) App. ii. 78 = luteo-album.

GNAPHALIUM:—
populifolium, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum populifolium.
pratense, *Phil. in Linnæa*, xxxiii. (1864–65) 166.—Chili.
prostratum, Hook. f. Fl. N. Zel. 187 = Helichrysum prostratum.
prostratum, Hnet, ex Nym. Consp. 362 = uliginosum.
prostratum, Patrin, ex DC. Prod. vi. 154 = Helichrysum arenarium.
prostratum, Roxb. in Wall. Cat. n. 2955 = pulvinatum prostratum, *Thunb. Prod. Pl. Cap.* 150.—Afr. austr.
protexcolorum, Schrank, in Denkschr. Akad. Muench. viii. (1824) 156 = Helichrysum paniculatum.
protiiforme, *DC. Prod.* vi. 237 = Helichrysum protiiforme.
protoides, Lam. Encyc. ii. 742 = Helichrysum paniculatum.
protoides, Sieber, ex DC. Prod. vi. 173 = Helichrysum yuccæfolium.
psilophyllum, *Meyen & Walp. in Nov. Act. Nat Cur.* xix. *Suppl.* i. (1843) 275.—Chili.
pterocaulon, Franch. & Sav. Enum. Fl. Jap. ii. 456 = Anaphalis pterocaulon.
puberulum, *DC. Prod.* vi. 224.—Chili.
pulchellum, E. Mey. ex DC. Prod. vi. 190 = Helichrysum pulchellum.
pulchellum, Wall. Cat. n. 2945 = Leontopodium alpinum.
pulchrum, *Steud. Nom.* ed. II. i. 695.—Sicilia.
pulverulentum, Bové, ex DC. Prod. vi. 231 = pulvinatum.
pulvinatum, *Delile, Fl. Egypte*, 266. t. 44.—Aegypt.; Arab.; Ind. or.
pumilum, Willd. Enum. Hort. Berol. Suppl. 57 = Helichrysum teretifolium.
purpurascens, *DC. Prod.* vi. 225.—Mexic.
purpureum, *Linn. Sp. Pl.* 854.—Reg. trop. et subtrop.
pusillum, Hæncke, in Jirasek, Beob. Riesengeb. (1791) 92 = supinum.
pusillum, Thunb. Prod. Pl. Cap. 149 = Leontonyx pusillus.
pusillum, Vand. Fasc. Pl. 19 = sylvaticum?
pygmæum, Lam. Encyc. ii. 761 = Evax pygmæa.
pygmæum, Thunb. Prod. Pl. Cap. 150 = Helichrysum declinatum.
pyramidale, Berg. Desc. Pl. Cap. 255 = Helichrysum stellatum.
pyramidale, *Thou. Fl. Tristan D'Acugn.* 40.—Ins. Tristan d'Acugn.
pyramidatum, Willd. Sp. Pl. iii. 1805 = Filago germanica.
quinquenerve, Thunb. Prod. Pl. Cap. 152 = Helichrysum nudifolium.
quinqueplinerve, Buch.-Ham. ex DC. Prod. vi. 270 = Anaphalis triplinervis.
racemosum, *Bogusl. in Erman, Archiv*, vi. (1848) 61.—Rossia.
racemosum, Schrank, ex DC. Prod. vi. 178 = Helichrysum mucronatum.
radiatum, *Benth. Fl. Hartw.* 207.—N. Granat.
radiatum, J. F. Gmel. Syst. 1917 = Petalaxis coronata.
radicans, *Burm. f. Fl. Cap. Prod.* 25.—Afr. austr.
ramigerum, DC. Prod. vi. 222 = luteo-album.
ramosissimum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 178.—Calif.
ramosissimum, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
ramosum, Lam. Fl. Fr. ii. 85 = uliginosum.
ramosum, *Phil. in Linnæa*, xxxiii. (1864–65) 164.—Chili.
ramosum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum ramosum.
ramulosum, Sch. Bip. l. c. 169 = Helichrysum ramulosum.
rectum, C. A. Mey. Verz. Pfl. Cauc. 77. n. 641 = asperum.
rectum, Sm. Engl. Bot. t. 124 = sylvaticum.
recurvatum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum recurvatum.
recurvum, Lam. Encyc. ii. 746 = Anaxetes recurvum.
redolens, Forst. f. Prod. 91 (= *Monanthes redolens*).—N. Caled.
Regnellii, Sch. Bip. in Linnæa, xxx. (1859–60) 180 = chrantifolium.
Reinwardtianum, Miq. Fl. Ind. Bat. ii. 91 = luteo-album.

GNAPHALIUM:—

repandum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum scabrum.

repens, Houtt. Handleid. vi. 67. t. 81. f. 1 (= *Trichogyne reflexa*).—Afr. austr.

repens, *Linn. Mant.* ii. 288.—Afr. austr.

resedaefolium, *Tausch, in Flora*, xvi. (1833) 123.—Chili.

reticulatum, Spreng. Syst. iii. 471 = Helichrysum reticulatum.

retortum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum retortum.

retrorsum, Boj. ex DC. Prod. vi. 207 = Helichrysum retrorsum.

retusum, Lam. Encyc. ii. 758 = Facelis apiculata.

retusum, Poepp. ex DC. Prod. vi. 224 = purpureum.

revolutum, Thunb. Prod. Pl. Cap. 150 = Helichrysum revolutum.

rhodanthum, *Sch. Bip. in Senn. Bot. Voy. Herald*, 310.—Mexic.

Riedelianum, Klatt, in Linnæa, xlii. (1878–79) 115 = cheiranthifolium.

rigidulum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum rigidulum.

rigidum, Salisb. Prod. 192 = Anaxeton arborcum, Metalasia aduncа.

rigidum, Steud. Nom. ed. II. i. 695 = Helichrysum striatum.

rivulare, *Phil. ex Klatt, in Linnæa*, xxxiii. (1864–65) 191.—Chili.

Roseri, Boiss. & Heldr. Diagn. Ser. II. vi. 102 = Hoppeanum.

rosco-rufescens, Steud. Nom. ed. II. i. 695 = purpureum. roseum, *H. B. & K. Nov. Gen. et Sp.* iv. 81.—Mexic.

rosmarinifolium, Salzm. ex Nym. Comp. 882 = Helichrysum angustifolium.

rosmarinifolium, Sieber, ex Spreng. Syst. iii. 472 = Helichrysum diosmifolium.

rosum, Berg. Desc. Pl. Cap. 260 = Helichrysum rosum.

rotundifolium, Thunb. Prod. Pl. Cap. 152 = Helichrysum rotundifolium.

Roxburghianum, Voigt, Hort. Suburb. Calc. 421 = luteo-album.

rubellum, Thunb. Prod. Pl. Cap. 150 = Helichrysum rubellum.

rufescens, H. B. & K. Nov. Gen. et Sp. iv. 78 = Achyrocline rufescens.

rufescens, *Poir. Encycl. Suppl.* ii. 802.—Hispan.

rufum, Willd. ex Less. in Linnæa, vii. (1832) 280 = Achyrocline satureioides.

rugulosum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum rugulosum.

rupestre, Bory, ex DC. Prod. vi. 178 = Helichrysum conspicuum.

rupestre, Poir. ex Nym. Consp. 881 = Helichrysum decumbens.

rupestre, Rafin. Précc. Découv. 41 = Helichrysum rupestre.

rupicola, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum rupicolum.

Ruppelli, Fresen. in Mus. Senckenb. i. (1833) 79. t. 4. f. 1 = Ifloga Fontanesii.

rutidolepis, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum rutidolepis.

rutilans, Lam. Sp. Pl. 854 = Helichrysum rutilans.

salicifolium, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum salicifolium.

sandwicense, *Gaudich. in Freyc. Voy. Bot.* 466.—Ins. Sandvic.

sanguineum, Linn. Cent. Pl. i. 27 ; Amœn. Acad. iv. 78 = Helichrysum sangueinum.

satureioides, Lam. Encyc. ii. 747 = Achyrocline satureioides.

saturejæfolium, Berg. ex Moench, Meth. Suppl. 286 = Metalasia muricata !

saturejæfolium, Poepp. ex DC. Prod. vi. 220 = Achyrocline satureioides.

saxatile, Blume, Bijdr. 869 = Antennaria saxatilis.

saxatile, Linn. Sp. Pl. 857 = Phagnalon saxatile.

scabrum, Linn. f. c. 855 = Leontonyx tomentosus.

scabrum, Thunb. Prod. Pl. Cap. 150 = Helichrysum scabrum.

scaiosa, Sieber, ex DC. Prod. vi. 182 = Helichrysum siculum.

Schimperi, Sch. Bip. ex Oliver, Fl. Trop. Afr. iii. 341 = Achyrocline Schimperi.

Schimperi, Sch. *Bip. in Schomb. Faun. & Fl. Cui.* 1187, nomen.—Guian.

GNAPHALIUM:—

Schrader, *DC. Prod.* vi. 225.—Am. !

sclerochlænum, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 149. n. 778 = Achyrocline sclerochlæna.

scoparium, Schrank, in Denkschr. Akad. Muench. viii. (1824) 160 = Helichrysum teretifolium.

scorpioides, Poir. Encyc. Suppl. ii. 808 = Helichrysum scorpioides.

sedoides, *Klatt, in Linnæa*, xlii. (1878–79) 135.—Socotra.

Seemanni, Sch. Bip. in Seem. Bot. Voy. Herald, 309 = Chionolæna corymbosa.

selaginifolium, Boj. ex DC. Prod. vi. 217 (= *Apheloma selaginifolia*).—Madag.

semiamplexicaule, *DC. Prod.* vi. 228.—Mexic.

semidecurrens, Wall. Cat. n. 2947 = Anaphalis araneosa.

semipapposum, Labill. Nov. Holl. Pl. ii. 42. t. 187 = Helichrysum semipapposum.

sericeum, Berg. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 83 = indutum.

serphioides, Berg. Desc. Pl. Cap. 267 (= *Trichogyne serphioides*).—Afr. austr.

serotinum, *Reichb. f. ex Fl. Germ.* xvi. 80. t. 952.—Europ.

serpillifolium; Schrank, in Denkschr. Akad Muench. viii. (1824) 156 = ser. !

serpyllifolium, Berg. Desc. Cap. Pl. 250 = Helichrysum serpyllifolium.

serpyllifolium, *Remy, in C. Gay, Fl. Chil.* iv. 283.—Ins. J. Fernand.

serratum, Linn. Pl. Rar. Afr. 19 ; Amœn. Acad. vi. 98 = Helichrysum cymosum, maritimum.

serratum, Thunb. Prod. Pl. Cap. 149 = Helichrysum felinum.

sessile, Burm. f. Fl. Cap. Prod. 25 = Helichrysum expansum.

siculum, Spreng. Syst. iii. 476 = Helichrysum cespitosum.

siculum, Tenore, Syll. Fl. Neap. 425 = Helichrysum Stoechas.

siculum, Tineo, Cat Pl. Hort. Panorm. 278 = Helichrysum angustifolium.

Sieboldianum, Franch. & Sav. Enum. Pl. Jap. i. 241 = Leontopodium japonicum.

simillimum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum simillimum.

simplex, Boeuin. Prod. Pl. Monast. 249 = Filago arvensis.

simplex, A. Rich. & Less. Fl. Nov. Zel. 287 = collinum.

simplicicaule, Wall. Cat. n. 2946 = Anaphalis contorta.

simplicicaule, Willd. ex Spreng. Syst. iii. 461.—Venezuela.

Sinclairii, *Hook. f. Handb. N. Zeal. Fl.* 153.—N. Zel.

sinuatum, *Lour. Fl. Cochinch.* ii. 497.—Cochinch.

soldanginoides, Poir. Encyc. Suppl. ii. 805 = Blumea soldaginoides.

Sophiæ, Baldr. ex Bolss. Fl. Orient. iii. 226 = sylvaticum.

sordescens, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum sordescens.

sordidum, Linn. Sp. Pl. 853 = Phagnalon sordidum.

spadiceum, Gilib. Fl. Lituan, i. 180 = sylvaticum.

spadiceum, Lam. Encyc. ii. 753 = Helichrysum rubellum.

spathulatum, Delile, Fl. Egypte, 266. t. 44 = indicum.

spathulatum, Lam. Encyc. ii. 759 = purpureum.

spathulatum, Thunb. Prod. Pl. Cap. 151 = Leontonyx spathulatus.

spathulatum, Burm. f. Fl. Cap. Prod. 25 = Helichrysum crispum.

sphæciatum, H. B. & K. Nov. Gen. et Sp. iv. 86 = purpureum, Hoppeanum.

sphæricum, Perr. ex DC. Prod. vi. 230 = luteo-album.

sphæricum, Willd. Enum. Hort. Berol. ii. 867 = japonicum.

spicatum, Lam. Encyc. ii. 757 = purpureum.

spicatum, Mill. Gard. Dict. ed. VIII. n. 24 = Pterocaulon virgatum.

spicatum, Vahl, Symb. Bot. i. 70 = Ifloga Fontanesii.

spiciforme, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum spiciforme.

spiciforme, *Sch. Bip. in Flora*, xxxviii. (1855) 116.—Reg. Magellan.

spinosum, Sch. Bip. in Bot. Zeit. iii. (1845) 178 = Helichrysum ctirispinum.

splendens, Willd. ex Spreng. Syst. iii. 476 = viscosum.

splendidum, Thunb. Prod. Pl. Cap. 149 = Helichrysum splendidum.

GNAPHALIUM:—

Sprengelii, *Hook. & Arn. Bot. Beech. Voy.* 150.—Amer. bor. occ.

squamosum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum squamosum.

squarrosum, Linn. Sp. Pl. ed. II. 1196 = Leontonyx tomentosus.

squarrulosum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum semipapposum.

stachidifolium, Lam. Encyc. ii. 757 = purpureum.

Stæchas, Auct. vide *Stoechas*.

staehelinoides, Thunb. Prod. Pl Cap. 150 = Helichrysum lucilioides.

stellatum, Linn. Pl. Rar. Afr. 19 ; Amœn. Acad. vi. 98 = Helichrysum stellatum.

stenacladium, Schrank, in Denkschr. Akad. Muench viii. (1824) 161 = Anaxeton arboreum.

stenophyllum, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 344.—Afr. trop.

stenopterum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum stenopterum.

Steudelii, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) i. *Intell.* 27.—Abyss.

Stewartii, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind* iii. 289.—Keg. Himal.

stipitatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum elatum.

Stoechas, Burm. f. Fl. Cap. Prod. 25 = Helichrysum teretifolium !

Stoechas, Linn. Sp. Pl. 853 = Helichrysum Stoechas.

Stoechas, Sibth. & Sm. Fl. Græc. ix. 44. t. 857 = Helichrysum angustifolium.

Stoechas, Thunb. Fl. Cap. 647 = Helichrysum rosum.

stolonifirum, Sch. Bip. in Bot. Zeit. iii. (1845) 169 = Helichrysum stoloniferum.

stramineum, *H. B. & K. Nov. Gen. et Sp.* iv. 85.—Mexic.

striatum, Sch. Bip. in Bot. Zeit. iii. (1845) 170 = Helichrysum striatum.

strictum, *A. Gray, in Pacif. Rail. Rep.* iv. 110.—Am. bor. occ.

strictum, Lam. Encyc. ii. 747 = Helichrysum splendidum

strictum, Moench, Meth. 576 = sylvaticum.

strictum, Roxb. Hort. Beng. 61 ; Fl. Ind. iii. 424 = indicum.

strictum, Zeyh. ex DC. Prod. vi. 204 = Helichrysum anomalum.

strigosum, Thunb. Prod. Pl. Cap. 150 = Helichrysum odoratissimum.

suaveolens, Vell. Fl. Flum. viii. t. 100 = Pluches Quitoc.

subacaule, Krock, Fl. Siles Suppl ii. 209 = supinum !

subdecurrens, DC, in Wight, Contrib. 21 = Anaphalis oblonga.

subdecurrens, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum subdecurrens.

subglomeratum, Sch. Bip. l. c. 171 = Helichrysum subglomeratum.

subsigidum, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 245.—N. Zel.

subundulatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum acuminatum.

sufrutticosum, *Phil. in Linnæa*, xxxiii. (1864–65) 165.—Chili.

supinum, Hoppe, in Sturm, Deutschl. Fl Heft 88 = G. Hoppeanum.

supinum, *Linn. Syst. ed.* XII. 234.—Reg. bor. occ. et arct.

supracanum, Sibth. & Sm. Fl. Græc. ix. 47. t. 861 = Ifloga Fontanesii.

sylvaticum, *Linn. Sp. Pl.* 856.—Europ. ; Asa. bor.

sylvaticum, Sm. Engl. Bot. t. 918 = norvegicum.

tenellum, Wall. Cat. n. 2941 = Anaphalis contorta.

Tenerifæ, Spreng. Syst. iii. 468 (= *Schizogyne obtusifolia*).—Ins. Canar.

tenue, *H. B. & K. Nov. Gen. et Sp.* iv. 88.—Mexic.

tenuicaulum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum tenuicaulum.

tenuifolium, Salisb. Prod. 192 = Filago gallica.

tenuifolium, *Vahl, ex Spreng. Syst.* iii. 470.—Madag.

tephroides, Link, Enum. Hort. Berol. ii. 317 = Helichrysum teretifolium.

teretifolium, Linn. Sp. Pl. 854 = Helichrysum teretifolium.

Thomsoni, *Hook. f. Fl. Brit. Ind.* iii. 290.—Reg. Himal.

Thunbergii, Spreng. Syst. iii. 478 = pyramidale.

Timmius, Buch.-Ham. ex Spreng. l. c. 478 = Anaphalis cinnamomea.

GNAPHALIUM:—
tinctum, Thunb. Prod. Pl. Cap. 151 = Leontonyx tomentosus.
tomentosum, Hoffm. ex J. F. Gmel. Syst. 1217 = uliginosum.
Trawerii, *Hook. f. Handb. N. Zeal. Fl.* 154.—Austral.; N. Zel.
tricostatum, Sieber, ex DC. Prod. vi. 202 = Helichrysum cymosum.
tricostatum, Thunb. Prod. Pl. Cap. 151 = Helichrysum tricostatum.
trifidum, Thunb. l. c. 150 = luteo-album.
trilineatum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum trilineatum.
trinerve, *Forst. f. Prod.* 55; *DC. Prod.* vi. 236.—N. Zel.
triplinerve, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum triplinerve.
uliginosum, C. B. Clarke, Comp. Ind. 115 = Thomsoni.
uliginosum, *Linn. Sp. Pl.* 856.—Europ.; As. bor.
uliginosum, A. Rich. Tent. Fl. Abyss. i. 431 = Unionis.
ulophyllum, Hook. & Arn. Bot. Beech. Voy. 31 = cymatodes.
ulophyllum, E. Mey. ex DC. Prod. vi. 197 = Helichrysum Zeyheri.
umbellatum, Linn. f. Suppl. 363 = Erythropogon umbellatum.
umbellatum, Schrank, in Denkschr. Akad. Muench. vill. (1824) 161 = Metalasia muricata.
umbellatum, Willd. Sp. Pl. iii. 1861 = Helichrysum terretifolium.
umbraculigerum, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum umbraculigerum.
undatum, Thunb. Prod. Pl. Cap. 151 = Helichrysum undatum.
undulaefolium, Reichb. ex DC. Prod. vi. 202 = Helichrysum odoratissimum.
undulatum, *Linn. Sp. Pl.* 857.—Afr. austr.
undulatum, Pers. Syn. ii. 416 = brachypterum.
undulatum, Walt. Fl. Carol. 208 = Pterocaulon pycnostachyum.
uniflorum, *Lam. Encyc.* ii. 762.—Peruv.
uniflorum, Mill. Gard. Dict. ed. VIII. n. 3 = Phagnalon saxatile.
uniflorum, Pall. ex DC. Prod. vi. 269 = Antennaria alpina.
uniflorum, Pall. ex Willd. in Spreng. Syst. iii. 470 = Helichrysum callichrysum.
Unionis, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 27, nomen; *A. Rich. Tent. Fl. Abyss.* i. 431.—Abyss.
ustulatum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 404 = purpureum.
valdivianum, *Phil. in Linnaea*, xxix. (1857–58) 6.—Chili.
valdiviense, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Chili.
valparadisium, *Phil. in Linnaea*, xxix. (1857–58) 5.—Chili.
Vauthierianum, Sch. Bip. in Linnaea, xxx. (1859–60) 180 = Achyrocline alata.
velutinum, *Willd. ex Spreng. Syst.* iii. 468.—Hab.?
verbascifolium, Schrank, in Denkschr. Akad. Muench. viii. (1824) 150 = Helichrysum grandiflorum.
verticillatum, Linn. f. Suppl. 364 (= *Trichogyne verticillata*)—Afr. austr.
vestitum, Sch. Bip. in Bot. Zeit. iii. (1845) 172 = Helichrysum vestitum.
vestitum, Thunb. Prod. Pl. Cap. 148 = repens.
villosissimum, D. Don, Prod. Fl. Nep. 175 = Anaphalis araneosa.
Vira-Vira, *Molina, Sagg. Chil.* 149, 264.—Chili.
virgatum, Banks & Soland. ex Hook. f. Fl. N. Zel. 189 = japonicum.
virgatum, Linn. Amoen. Acad. v. 405 = Pterocaulon virgatum.
virgineum, Sibth. & Sm. Fl. Graec. ix. 46 = Helichrysum Billardieri.
viscidum, Blume, Bijdr., 900 = Anaphalis viscida.
viscosum, *H. B. & K. Nov. Gen. et Sp.* iv. 93.—Mexic.
viscosum, Poepp. ex DC. Prod. vi. 222 = cheiranthifolium.
Wahlenbergii, Sieber, ex Stend. Nom. ed. II. i. 696 = Antennaria carpathica.
Webbii, Sch. Bip. in Webb. & Berth. Phyt Canar. ii. 218 = Helichrysum gossypinum.
Wightianum, DC. in Wight, Contrib. 21 = Anaphalis Wightiana.

GNAPHALIUM:—
Wightianum, Thw. Enum. Pl. Zel. 166 = Anaphalis cinnamomea.
Wightii, Thw. l. c. 165 = Anaphalis Thwaitesii.
Willdenowii, Dietz. ex DC. Prod. vi. 236 = japonicum.
Willdenowii, Spreng. Syst. v. 814, Index = bractenatum.
Wirtgenii, Nym. Consp. 393 = uliginosum.
Wrightii, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 214.—Am. bor. occ.
xanthinoides, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum splendidum.
xeranthemoides, Sch. Bip. l. c. 169 = Helichrysum lanciifolium.
xerochrysum, Sch. Bip. l. c. 171 = Helichrysum xerochrysum.
yedoënse, *Franch. & Sav. Enum. Pl. Jap.* ii. 404.—Japon.
Youngii, *Hook. f. Handb. N. Zeal. Fl.* 152.—N. Zel.
yuccaefolium, Lam. Encyc. ii. 742 = Helichrysum yuccaefolium.
Zeyheri, Sch. Bip. in Bot. Zeit. iii. (1845) 171 = Helichrysum Zeyheri.
Zollingeri, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 82.—Java.

GNAPHALODES, A. Gray, in Hook. Kew Journ. iv. (1852) 238. *COMPOSITAE*, Benth. & Hook. f. ii. 291.
condensatum, *A. Gray, l. c.*—Austral.
crocinum, Sond. in Linnaea, xxv. (1852) 520 = uliginosum.
filifolium, *Benth. Fl. Austral.* iii. 578.—Austral.
uliginosum, *A. Gray, in Hook. Kew Journ.* iv. (1852) 238.—Austral.

GNAPHALODES, Tourn. ex Adans. Fam. ii. 118 (1763) = **Micropus**, Linn. (Compos.).
dendroides, Moench, Meth. 589 = M. supinus.
erecta, Moench, l. c. = M. erectus.

GNAPHALON, Lowe, Man. Fl. Mad. i. 438 (1858) = **Phagnalon**, Cass. (Compos.).
calycinum, Lowe, l. c. 441 = P. rupestre.
rupestre, Lowe, l. c. 439 = P. rupestre.
saxatile, Lowe, l. c. 438 = P. saxatile.

GNAPHALOPSIS, DC. Prod. vii. 258 (1838) = **Hymenolepis**, Cass. (Compos.).
micropoides, DC. l. c. = H. Gnaphalopsis.

GNEORUM, G. Don, Gen. Syst. ii. 62 (1832) orr. typ. = Cneorum, Linn. (Simarub.).

GNEPHOSIS, Cass. in Bull. Soc. Philom. (1820) 43; Dict. Sc. Nat. xix. 128 (1821). *COMPOSITAE*, Benth. & Hook. f. ii. 320.
CROSSOLEPIS, Less. Syn. Comp. 270 (1832); A. Gray, in Hook. Kew Journ. iii. (1851) 176.
CYATHOPAPPUS, F. Muell. Fragm. ii. 157 (1861).
LEPTOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 78.
NEMLEPOLIUM, A. Gray, in Hook. Kew Journ. iii. (1851) 160.
TRICHANTHODIUM, Sond. et F. Muell. in Linnaea, xxv. (1852) 489.
acicularis, *Benth. Fl. Australiae*, iii. 572.—Austral.
arachnoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 189.—Austral.
brevifolia, *Benth. Fl. Austral.* iii. 572.—Austral.
Burkittii, *Benth. l. c.* 570.—Austral.
cadonopappus, F. Muell. Fragm. ix. 2 = Angianthus cadonopappus.
cyathopappa, *Benth. Fl. Austral.* iii. 571.—Austral.
eriocarpa, *Benth. l. c.* 570.—Austral.
eriocephala, *Benth. l. c.* 573.—Austral.
leptoloba, *Benth. l. c.* 571.—Austral.
macrocephala, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 190.—Austral.
pygmaea, *Benth. Fl. Austral.* iii. 572.—Austral.
sphærophora, *Benth. l. c.* 570.—Austral.
tenuissima, *Cass. in Bull. Soc. Philom.* (1820) 43; *in Dict. Sc. Nat.* xix. 128.—Austral.

GNETUM, Linn. Mant. i. 13. n. 1276 (1767). *GNETACEAE*, Benth. & Hook. f. iii. 419.
ABUTUA, Loeu. Fl. Cochinch. 630 (1790).
ARTHOSTEMA, Neck. Elem. ii. 220 (1790).
THOA, Aubl. Pl. Gui. ii. 874. t. 320 (1775).

GNETUM:—
acutatum, *Miq. Fl. Ind. Bat. Suppl.* 588.—Sumatra.
africanum, *Welw. in Trans. Linn. Soc. xxvii.* (1869) 78.—Afr. trop.
amazonicum, *Tul. in Ann. Sc. Nat.* Sér. IV. x. (1858) 116.—Am. trop.
apiculatum, Griff. Notul. iv. 80 = neglectum.
Brunonianum, Griff. in Lindl. Veg. Kingd. 823. f. 164; et in Notul. iv. 80 = G. Gnemon.
cuspidatum, *Blume, Rumphia*, iv. 5.—Sumatra.
dioicum, Leyb. ex Tul. in Ann. Sc. Nat. Sér. IV. x. (1858) 115, 116 = Leyboldi, amazonicum.
edule, Blume, in Hoev. & De Vriese, Tijdschr. i. (1834) 161 = scandens.
funiculare, *Blume, l. c.* 162.—Java.
funiculare, Wight, Ic. t. 1955 = scandens.
Gnemon, *Linn. Mant.* i. 125.—Malaya.
gnemonoides, Brongn. in Duperr. Voy. Bot. 12 = scandens.
Griffithii, Parl. in DC. Prod. xvi. ii. 349 = G. Gnemon.
latifolium, *Blume, in Hoev. & De Vriese, Tijdschr.* i. (1834) 162.—Java.
latifolium, Parl. in DC. Prod. xvi. ii. 350 = scandens.
leptostachyum, *Blume, Rumphia*, iv. 5.—Borneo.
Leyboldi, *Tul. in Ann. Sc. Nat.* Sér. IV. x. (1858) 115.—Am. austr.
macrocarpum, *Becc. Malesia.* i. (1877) 182.—Borneo.
macropodum, *Kurz, in Journ. Bot.* xiii. (1875) 331.—Ins. Nicobar.
microcarpum, Blume, Rumphia, iv. 7 = neglectum.
microstachyum, *Spruce & Benth. ex Parl. in DC. Prod.* xvi. ii. 352.—Bras.
neglectum, *Blume, Rumphia*, iv. 6. t. 175. f. 2.—Burma; Malaya.
nigrum, *Carr. Conif.* ed. I. 545.—Ind. occ.
oodiflorum, *Brongn. in Duperr. Voy. Bot.* i. 12.—Guiana.
ovalifolium, Poir. Encyc Suppl. ii. 810 = G. Gnemon.
paniculatum, *Spruce, ex Benth. in Hook. Kew Journ.* viii. (1856) 357.—Bras.
polystachyum, Reinw. in Blume, Cat. Hort. Buitens. 106 = G. Gnemon.
pyrifolium, Miq. ex Parl. in DC. Prod. xvi. ii. 350 = scandens.
Rumphianum, *Becc. Malesia*, i. (1877) 182.—Malaya.
scandens, *Roxb. Hort. Beng.* 66; *Fl. Ind.* iii. 618.—Ind. or.; Malaya.
sylvestre, Brongn. in Duperr. Voy. Bot. 12 = G. Gnemon.
Thoa, *R. Br. in King, Narr. Surv. Austral.* ii. 555.—Guian.
Ula, Brongn. in Duperr. Voy. Bot. 12 = scandens.
urens, *Blume, in Hoev. & De Vriese, Tijdschr.* i. (1834) 162.—Guian.
venosum, *Spruce ex Benth. in Hook. Kew Journ.* viii. (1856) 358.—Bras.

GNIDIA, Linn. Phil. Bot. 32 (1751). *THYMELAEACEAE*, Benth. & Hook. f. iii. 197.
CANALIA, F. W. Schmidt, Neue u. Selt. Pfl. 30 (1793).
DESSENIA, Adans. Fam. ii. 285 (1763).
EPICEROXANTHA, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 580 (1856–57).
NECTANDRA, Berg. Descr. Pl. Cap. 131 (1767).
STRUTHIA, Royen, ex Linn. Gen. ed. II. 164 (1742).
THYMELINA, Hoffmgg. Verz. Pfl. 198 (1824).
acerosa, J. F. Gmel. Syst. 635 = juniperifolia.
acerifolia, *Willd. in Vet. Acad. Handl. Stockh.* (1818) 218.—Afr. austr.
albicans, *Meisn. in DC. Prod.* xiv. 584.—Afr. austr.
andromedaefolia, Eckl. & Zeyh. ex Meisn. l. c. 585 = candicans.
anomala, *Meisn. in Linnaea*, xiv. (1840) 485.—Afr. austr.
argentea, Eckl. & Meisn. in DC. Prod. xiv. 581 = anomala.
argentea, *Thunb. Prod. Pl. Cap.* 76.—Afr. austr.
argentea, Zeyh. ex Meisn. in DC. Prod. xiv. 594 = Lachnaea Meisneriana.
aurea, Stend. ex Meisn. in Linnaea, xiv. (1840) 481 = juniperifolia.
canescens, Burch. ex Meisn. l. c. 585 = serissa.
capitata, *Thunb. Prod. Pl. Cap.* 76.—Afr. austr.
filifera, Willdt. in Vet. Acad. Handl. Stockh. (1816) 214 = pinifolia.
Hegmanni, *Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv.* 583.—Afr. austr.

GNIDIA:—
carpitosa, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 564 = rodiflora.
caffra, Eckl. & Zeyh. ex Meisn. l. c. = strigillosa.
caffra, Meisn. l. c. 392 = Lasiosiphon caffer.
canifloro, *Meisn. in Linnaea*, xiv. (1840) 440.—Afr. austr.
capitata, Linn. f. Suppl. 224 = Lasiosiphon linifolius.
carinata, *Thunb. Prod. Pl. Cap.* 76.—Afr. austr.
Cephalotes, *Lichtenst. ex Meisn. in DC. Prod.* xiv. 581.—Afr. austr.
chrysophylla, *Meisn. l. c.* 584.—Afr. austr.
cornuta, Eckl. & Zeyh. ex Meisn. l. c. 560 = thesioides.
coriacea, *Meisn. in*, *Linnaea*, xiv. (1840) 454.—Afr. austr.
cuneata, Meisn. l. c. 427 = Lasiosiphon Meisneriana.
cyanea, *Burch. Trav.* i. 325, 327, nomen.—Afr. austr.
daphnaefolia, Boj. ex Decne. in Jacquem. Voy. Bot. 149 = Lasiosiphon Bojerianus.
daphnaefolia, Linn. f. Suppl. 225 = Lasiosiphon pubescens, madagascariensis.
daphnoides, Boj. ex Meisn. in DC. Prod. xiv. 596 = pruec.
decurrens, *Meisn. in Linnaea*, xiv. (1840) 451.—Afr. austr.
denudata, *Lindl. Bot. Reg.* t. 757.—Afr. austr.
Dregeana, Meisn. in Linnaea, xiv. (1840) 426 = Lasiosiphon Dregeanus.
elongata, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 561 = scabrida.
eriocephala, Meisn. in Denkschr. Bot. Ges. Regensb. iii. (1841) 292 = Lasiosiphon eriocephalus.
filamentosa, *Linn. f. Suppl.* 224.—Afr. austr.
flava, *Lindl. ex Steud. Nom.* ed. II. i. 697, nomen.—Afr. austr.
fulgens, *Welw. Apont.* 548.—Afr. trop.
geminiflora, *E. Mey. ex Meisn. in Linnaea*, xiv. (1840) 441.—Afr. austr.
genistaefolia, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 574 = Cryptadenia filicaulis.
glandulosa, Hayne, ex Meisn. in Linnaea, xiv. (1840) 429 = tomentosa.
glauca, Steud. Nom. ed. II. i. 697 = filamentosa.
grandiflora, Meisn. in DC. Prod. xiv. 582 = Cephalotes.
grandiflora, *Willd. Enum. Hort. Berol. Suppl.* 21, nomen.—Afr. austr.
grandis, Hort. ex Sims, Bot. Mag. t. 1658 = filamentosa.
Harveyana, *Meisn. in Linnaea*, xiv. (1840) 437.—Afr. austr.
Hugelii, Meisn. in DC. Prod. xiv. 592 = Lasiosiphon eriocephalus.
humilis, *Meisn. l. c.* 586.—Afr. austr.
imberbis, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 412 = carinata.
imbricata, *Linn. f. Suppl.* 225.—Afr. austr.
incana, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 588 = imbricata.
inconspicua, *Meisn. in Linnaea*, xiv. (1840) 433.—Afr. austr.
insularis, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 456 = Lasiosiphon eriocephalus.
involucrata, *Steud. ex A. Rich. Tent. Fl. Abyss.* ii. 234.—Abyssin.
juniperifolia, *Lam. Encyc.* ii. 765.—Afr. austr.
Krausciana, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 552 (err. typ. 452) = Lasiosiphon Krausii.
laevigata, Thunb. Prod. Pl. Cap. 76 = oppositifolia.
lanceolata, Lichtenst. ex Meisn. in DC. Prod. xiv. 588 = parviflora.
latifolia, Hort. ex Meisn. in Linnaea, xiv. (1840) 432 = oppositifolia.
linoides, *Wikstr. in Vet. Acad. Handl. Stockh.* (1818) 316.—Afr. austr.
macrophylla, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 553 (err. typ. 453) = Lasiosiphon macrocarpus.
Meyeri, *Meisn. in Linnaea*, xiv. (1840) 443.—Afr. austr.
microcephala, *Meisn. in DC. Prod.* xiv. 589.—Afr. austr.
microphylla, Meisn. in Linnaea, xiv. (1840) 428 = Lasiosiphon microphyllus.
monticola, Mtc. in Flora, xxxii. (1849) 657 = Lasiosiphon eriocephalus.
multiflora, *Burch. ex Meisn. in Linnaea*, xiv. (1840) 448.—Afr. austr.

GNIDIA:—
nana, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 588 = sericea.
nana, *Wikstr. in Vet. Acad. Handl. Stockh.* (1818) 316.—Afr. austr.
nodiflora, *Meisn. in Linnaea*, xiv. (1840) 458.—Afr. austr.
obtusifolia, Bartl. ex Meisn. l. c. 461 = obtusissima.
obtusissima, *Meisn. l. c.* 432.—Afr. austr.
occidentalis, *Regel, Cat. Pl. Hort. Akadow.* 65, nomen.—Hab.?
ochroleuca, Lodd. Bot. Cab. t. 1184 = pinifolia.
oppositifolia, Blanco, Fl. Filip. ed. I. 299 = philippinica.
oppositifolia, *Linn. Sp. Pl.* 358.—Afr. austr.
oppositifolia, Thunb. Prod. Pl. Cap. 76 = sericea.
ovalifolia, *Meisn. in Linnaea*, xiv. (1840) 455.—Afr. austr.
pachyphylla, Spreng. ex Meisn. l. c. 405 = Cryptadenia grandiflora.
pallida, *Meisn. l. c.* 442.—Afr. austr.
parviflora, *Meisn. l. c.* 458.—Afr. austr.
penicillata, *Lichtenst. ex Meisn. l. c.* 448.—Afr. austr.
pentandra, Thunb. Diss. Fructif. Sect. Pr. 19 = Lonchostoma obtusiflorum.
philippinica, *Meisn. in DC. Prod.* xiv. 592.—Ins. Philipp.
phylicifolia, Eckl. & Zeyh. ex Meisn. l. c. 588 = sericea.
pinifolia, *Linn. Sp. Pl.* 356.—Afr. austr.
pinifolia, Linn. f. Suppl. 225 = juniperifolia.
pinifolia, Wendl. Bot. Beobacht. 15. t. 2. f. 11 = carinata.
polygnaefolia, Lichtenst. ex Meisn. in Linnaea, xiv. (1840) 468 = parviflora.
polystachya, Berg. Desc. Pl. Cap. 128 = carinata.
prismatifolia, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 585 = scabra.
pubescens, Berg. Desc. Pl. Cap. 124 = tomentosa.
pulchella, Meisn. in Linnaea, xiv. (1840) 425 = Lasiosiphon pulchellus.
punctata, Lam. Encyc. ii. 765 = tomentosa.
racemosa, *Thunb. Prod. Pl. Cap. 76.*—Afr. austr.
radiata, Linn. Mant. i. 67; Wendl. Bot. Beobacht. 15. t. 2. f. 12 = pinifolia.
scabra, *Thunb. Prod. Pl. Cap.* 76.—Afr. austr.
scabrida, *Meisn. in Linnaea*, xiv. (1840) 446.—Afr. austr.
sericea, *Linn. Syst.* ed. XII. 372.—Afr. austr.
setosa, *Wikstr. in Vet. Acad. Handl. Stockh.* (1818) 315.—Afr. austr.
simplex, Andr. Bot. Rep. t. 70 = carinata.
simplex, Linn. Mant. i. 67 = juniperifolia, subulata.
stipitaria, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 457 = Lasiosiphon eriocephalus.
Sonderiana, *Meisn. in DC. Prod.* xiv. 587.—Afr. austr.
Spermanni, Martyn, in Mill. Gard. Dict. ed. IX. n. 10 = juniperifolia.
spariflora, *Bartl. ex Meisn. in Linnaea*, xiv. (1840) 493.—Afr. austr.
splendens, Meisn. l. c. 428 = Lasiosiphon splendens.
stricta, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 560 = thesioides.
stricta, *Wikstr. in Vet. Acad. Handl. Stockh.* (1818) 315.—Afr. austr.
strigillosa, *Meisn. in Linnaea*, xiv. (1840) 459.—Afr. austr.
stypheloides, *Meisn. l. c.* 458.—Afr. austr.
subcordata, *Meisn. l. c.* 480.—Afr. austr.
subulata, *Lam. Encyc.* ii. 765.—Afr. austr.
taxifolia, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 585 = canifloro.
tenella, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 554 (err. typ. 454) = albicans.
tennifiora, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 577, 579 = Lachnaea Burchellii, diosmoides.
thesioides, *Meisn. in Linnaea*, xiv. (1840) 457.—Afr. austr.
tomentosa, Eckl. ex Meisn. in Linnaea, xiv. (1840) 392 = Lasiosiphon sathyliodes.
tomentosa, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 584 = albicans.
tomentosa, Hook. Bot. Mag. t. 2761 = denudata.
tomentosa, *Linn. Sp. Pl.* 355.—Afr. austr.
triplinervia, Meisn. in Linnaea, xiv. (1840) 429 = Lasiosiphon triplinervis.

GNIDIA:—
vesiculosa, Eckl. & Zeyh. ex Meisn. in DC. Prod. xiv. 559 = Arthrosolen ornatus.
villosa, Meisn. l. c. 594 = Lasiosiphon Burchellii.
virescens, Eckl. & Zeyh. ex Meisn. l. c. 585 = denudata.
virescens, *Wikstr. in Vet. Acad. Handl. Stockh.* (1818) 317.—Afr. austr.
viridis, Berg. Desc. Pl. Cap. 125 = juniperifolia.
Wickstroemiana, *Meisn. in Linnaea*, xiv. (1840) 434.—Afr. austr.

GNIDIUM, G. Don, in Lond. Hort. Brit. 107 (1830) sphalm. = CNIDIUM, Cuss. = Selinum, Linn. (Umbellif.).

GNOTERIS, Rafin. Sylva Tellur. 76 (1838) = **Hyptis**, Jacq. (Labiat.).
cordata, Rafin. l. c. = H. suaveolens.
villosa, Rafin. l. c. = H. suaveolens.

GOCHNATEA, Steud. Nom. ed. II. i. 697 (1840) = seq.

GOCHNATIA, H. B. & K. Nov. Gen. et Sp. iv. 19. t. 309 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 490.
PENTAPHORUS, D. Don, in Trans. Linn. Soc. xvi. (1830) 295.
cinerea, *Griseb. in Goett. Abh.* xix. (1874) 195.—Reg Argent.
cordata, *Less. in Linnaea*, v. (1830) 268.—Bras.
corymbosa, *Less. l. c.* = Onoseris corymbosa.
discolor, *Baker*, in *Mart. Fl. Bras.* vi. iii. 350.—Bras. fasciculare, *D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 108.—Chili.
foliolosa, D. Don, ex Hook. & Arn. l. c. in texts = rigida?
genistoides, Hook. & Arn. l. c. 109 = Cyclolepis genistoides.
glomerifora, *A. Gray, in Proc. Am. Acad.* xix. (1883) 57.—Mexic.
glutinosa, *D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 108.—Chili.
glutinosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 334.—Peruv.
hypoleuca, *A. Gray, in Proc. Am. Acad.* xix. (1883) 57.—Texas.
ilicifolia, Less. in Linnaea, v. (1830) 261 = Anastraphis ilicifolia.
polymorpha, Herb. Berol. ex DC. Prod. vii. 28 = Moquinia polymorpha.
pyrifolia, *D. Don, ex Hook. & Arn. in Hook. Comp Bot. Mag.* i. (1835) 109.—Chili.
revoluta, D. Don, ex Hook. & Arn. l. c. 109 = Nardophyllum revolutum.
rigida, *D. Don, ex Hook. & Arn. l. c.* 108.—Chili.
rotundifolia, *Less. Syn. Comp.* 102.—Bras.
spectabilis, Less. in Linnaea, v. (1830) 261 = Leucomeris spectabilis.
vernonioides, *H. B. & K. Nov. Gen. et Sp.* iv. 20.—Bras.

GODETIA, Spach, Hist. Vég. Phan. iv. 386 (1835) = Oenothera, Linn. (Onagrar.).
albescens, Lindl. Bot. Reg. (1841) Misc. 61 = Oe. albescens.
ambigua, Phil. in Anal. Univ. Chil. ii. (1865) 327.—Chili.
amoena, G. Don, in Sweet, Hort. Brit. ed. III. 387; Lilja, in Linnaea, xv. (1841) 265 = Oe. amoena.
Aranetii, Walp. Rep. ii. 26 = Oe. decumbens.
biloba, S. Wats. in Bot. Calif. i. 231 = Oe. biloba.
Bottae, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 393 = Oe. Bottae.
Cavanillesii, Spach, Hist. Vég. Phan. iv. 390 = Oe. tenella.
dasycarpa, Phil. in Linnaea, xxx. (1859–60) 187.—Chili.
decumbens, Spach, Hist. Vég. Phan. iv. 388 = Oe. decumbens.
epilobioides, S. Wats. in Bot. Calif. i. 231 = Oe. epilobioides.
Gayana, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 391 = Oe. Gayana.
grandiflora, Lindl. Bot. Reg. (1841) Misc. 61 = Oe Whitneyi.
hispidula, S. Wats. in Bot. Calif. i. 231 = Oe. hispidula

GODETIA :—
Lehmanniana, Spach, Hist. Vég. Phan. iv. 301 = Oenothera amoena.
lepida, Lindl. Bot. Reg. t. 1849 = Oe. decumbens.
Lindleyana, Spach, Hist. Vég. Phan. iv. 302 = Oe. amoena.
macrantha, Lilja, Tradg. Tidning. (1840) ; ex cod. in Linnæo, xv. (1841) 236 = Oe. amoena.
Nivertiana, Gouj. in Rev. Hortic. (1872) 481 = Oe. amoena.
purpurea, G. Don, in Sweet, Hort. Brit. ed. III. 237 ; et S. Wats. in Bot. Calif. i. 229 = Oe. purpurea.
quadrivulnera, Spach, Hist. Vég. Phan. iv. 389 = Oe. quadrivulnera.
Romanzowii, Spach, l. c. 390 = Oe. Romanzowii
rubicunda, Lindl. Bot. Reg. t. 1856 = Oe. amoena.
sulfurea, Phil. in Anal. Univ. Chil. (1872) 712.—Chili.
tenella, Stend. Nom. ed. II. i. 697 ; et S. Wats. in Bot. Calif. i. 230 = Oe. tenella.
tenuifolia, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 290 = Oe. tenuifolia.
viminea, Spach, Hist. Vég. Phan. iv. 388 = Oe. viminea.
vinosa, Lindl. Bot. Reg. t. 1880 = Oe. amoena.
Whitneyi ×, Hort.—*Cf.* Flor. Mag. N. S. t. 227.
Whitneyi, T. Moore, in Flor. & Pomol. (1871) 101 = Oe. Whitneyi.
Willdenowiana, Spach, Hist. Vég. Phan. iv. 387 = Oe. purpurea.
Williamsoni, Dur. & Hilg. in Facif. Rail. Rep. v. 7 = Oe. Williamsoni.

GODIA, Stend. Nom. ed. II. ii. 607, in syn. (1841), sphalm. = GOLIA, Adans. = **Soldanella**, Linn. (Primulac).

GODIAEUM, Boj. Hort. Maurit. 282 (1837), sphalm. = **Codiaeum**, Rumph. (Euphorb.).

GODINELLA, Lestib. f. Bot. Belg. ii. 194 (1827) = **Lysimachia**, Linn. (Primul.).

GODMANIA, Hemsl. Diagn. Pl. Nov. 85 (1879). *BIGNONIACEAE*.
macrocarpa, *Hemsl. l. c.*—Panama.

GODOVIA, Pers. Syn. i. 467 (1805) = seq.

GODOYA, Ruiz & Pav. Prod. 58. t. 11 (1794). *OCHNACEAE*, Benth. & Hook. f. i. 319.
antioquiensis, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 598.—N. Granat.
gemmifera, Mart. Nov. Gen. et Sp. i. 119 = Blastemanthus gemmiflorus.
oblonga, Ruiz & Pav. Prod. 58 = obovata.
obovata, *Ruiz & Pav. Syst. Veg.* 101.—Peruv.
spathulata, *Ruiz & Pav. Prod.* 58.—Peruv.
splendida, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 599.—N. Granat.

GODWINIA, Seem. in Journ. Bot. (1869) 314. t. 96, 97 = **Dracontium**, Linn. (Aroid.).
Gigas, Seem. l. c. 315. t. 96, 97 = D. Gigas.

GOEBELIA, Bunge, ex Boiss. Fl. Orient. ii 628 (1872) = **Sophora**, Linn. (Legumin.).
alopecuroides, Bunge, ex Boiss. l. c. 628, 629 = S. alopecuroides.
pachycarpa, Bunge, ex Boiss. l. c. 629 = S. pachycarpa.

GOEPPERTIA, Griseb. in Journ. Linn. Soc. vi. (1862) 141. *GENTIANEAE*, Benth. & Hook. f. ii. 808.
gracilis, *Wright, ex Griseb. Cat. Pl. Cub.* 180.—Cuba.
volubilis, *Griseb. in Journ. Linn. Soc.* vi. (1866) 141. —Cuba.

GOEPPERTIA, Nees, in Linnæa, vi. (1831) 837 (genus spurium).
bicolor, Nees, l. c. = Maranta bicolor.
blanda, Nees, l. c. = Calathea blanda.
episcata, Nees, l. c. = Inchnosiphon episcatus.
zebrina, Nees, l. c. = Calathea zebrina.

GOEPPERTIA, Nees, Syst. Laurin. 365 (1836) = **Aydendron**, Nees (Laurin.).
anomala, Nees, l. c. 370.—Bras.
argentea, Meisn. in DC. Prod. xv. i. 174 = A. argenteum.

GOEPPERTIA :—
cantagallana, Meisn. in DC. Prod. xv. i. 173.—Bras.
caudata, Meisn. l. c. 175.—Bras.
chrysophylla, Meisn. l. c. 174.—Bras.
dyscolantha, Nees, in Linnæa, xxi. (1848) 514.—Peruv.
geminiflora, Meisn. in DC. Prod. xv. i. 175.—Bras.
hirsuta, Nees, Syst. Laurin. 366 = Cryptocarya hirsuta.
longiflora, Nees, l. c. 368.—Peruv.
macrocalyx, Meisn. in DC. Prod. xv. i. 174.—Bras.
multiflora, Miq. Stirp Surin. Sel. 208 = Ampelodaphne dasyantha.
panicularis, Nees, Syst. Laurin. 368.—Bras.
polyantha, Meisn. in DC. Prod. xv. i. 176.—Bras.
reflexiflora, Nees, in Linnæa, xxi. (1848) 514.—Guian.
serrulata, Nees, Syst. Laurin. 369 = Aydendron sericeum.
Spruceii, Meisn. in DC. Prod. xv. i. 172.—Bras.

GOETHEA, Nees, in Flora, iv. (1821) 304 ; Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 91. t. 8. *MALVACEAE*, Benth. & Hook. f. i. 208.
Sect. HOLOMITA, Schrad. in Goett. Gel. Anz. (1821) 717.
alnifolia, *Garcke, in Eichl. Fahrb. Berl.* i. (1881) 222, in syn.—Bras.
cauliflora, *Nees, in Flora*, iv. (1821) 304.—Bras.
Mackoyana, *Hook. f. Bot. Mag.* t. 6427.—Bras.
semperflorens, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 99 = Pavonia semperflorens.
strictiflora, *Hook. Bot. Mag.* t. 4677.—Bras.

GOETZEA, Reichb. Consp. 304 (1828) = **Rothia**, Pers. (Legumin.).

GOETZEA, Wydl. in Linnæa, v. (1830) 423. t. 8. *SOLANACEAE*, Benth. & Hook. f. ii. 1044.
amoena, Griseb. Cat. Pl. Cub. 191, partim = Espadea amoena.
elegans, *Wydl. in Linnaea*, v. (1830) 423. t. 8.—Ins. Porto Rico.

GOETZIA, Miers, in Trans. Linn. Soc. xxvii. (1870) 192 = prsec.

GOHORIA, Neck. Elem. i. 172 (1790) = **Ammi**, Tourn. (Umbellif.).

GOLDBACHIA, DC. Syst. ii. 576 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 101.
laevigata, *DC. l. c.* 577.— Ax. Min. ; Persia ; Ind. or.
tetraptera, Ledeb. Ind. Sem. Hort. Dorpat. (1822) 9 ; Fl. Ross. i. 215 = laevigata.
torulosa, DC. Syst. ii. 577 = laevigata.

GOLDBACHIA, Trin. in Spreng. Neue Entdeck. ii. 81 (1821) = **Arundinella**, Raddi (Gramin.).
Mikani, Trin. l. c. = A. brasiliensis.

GOLDENIA, Raeusch. Nom. ed. III. 43 (1797), sphalm. = **Coldenia**, Linn. (Boragin.).

GOLDFUSSIA, Nees, in Wall. Pl. As. Rar. iii. 75, 87 (1832) = **Strobilanthes**, Blume (Acanth.).
anisophylla, Nees, l. c. 88 = S. anisophyllus.
bibracteata, Nees, in DC. Prod. xi. 174 = S. bibracteatus.
bicepa, Nees, in Wall. Pl. As. Rar. iii. 88 ; et Cat. n. 7251 = S. capitatus.
bracteata, Nees, l. c. 88 = S. quadrangularis.
capitata, Nees, l. c. = S. capitatus.
colorata, Mor. Syst. Veg. Zoll. 47 = Ruellia colorata.
coloreata, Nees, in Wall. Pl. As. Rar. iii. 89 = S. coloratus.
crinita, Nees, in DC. Prod. xi. 176 = S. coloratus.
Cusia, Nees, in Wall. Pl. As. Rar. iii. 88 = S. flaccidifolius.
Dalhousiana, Nees, in DC. Prod. xi. 174 = S. Dalhousianus.
decurrens, Wight, Ic. t. 1533 = S. decurrens.
discolor, Nees, in DC. Prod. xi. 172 = S. discolor.
divaricata, Nees, in Wall. Pl. As. Rar. iii. 89 = S. divaricatus.
Edgeworthiana, Nees, in DC. Prod. xi. 173 = S. angustifrons ?
extensa, Nees, in Wall. Pl. As. Rar. iii. 88 = S. extensus.
filiformis, Nees, in DC. Prod. xi. 176 = S. filiformis.
flexuosa, Nees, in Wall. Pl. As. Rar. iii. 86 = S. pentstemonoides.
glomerata, Nees, l. c. = S. glomeratus.

GOLDFUSSIA :—
isophylla, Nees, in Wall. Pl. As. Rar. iii. 88 ; et Cat. n. 7162 = Strobilanthes isophyllus.
lamifolia, Nees, l. c. 88 = S. lamifolius.
Leschenaultiana, Nees, in DC. Prod. xi. 172 = S. foliosus, Zenkerianus.
Myrtilla, Nees, l. c. 174 = S. lanceolatus.
nutans, Nees, in Wall. Pl. As. Rar. iii. 88 = S nutans.
paniculata, Nees, in Hook. Comp. Bot. Mag. i. (1836) 318 = S. alatus, *Blume*.
pentastemonoides, Nees, in Wall. Pl. As. Rar. iii. 86 = S. pentstemonoides.
sessilis, Nees, in DC. Prod. xi. 172 = S. rhombifolius.
Thomsoni, Hook. Bot. Mag. t. 5119 = S. Wallichii.
tristis, Wight, Ic. t. 1508 = S. tristis.
Zenkeriana, Nees, in DC. Prod. xi. 172 = S. Zenkerianus.
Zenkeriana, Wight, Ic. t. 1517 = S. ciliatus ?

GOLIA, Adans. Fam. ii. 231 (1763) = **Soldanella**, Tourn. (Primul.).

GOLOWNINIA, Maxim. in Bull. Acad. Pétersb. iv. (1862) 252 = **Crawfurdia**, Wall. (Gentian.).
japonica, Maxim. l. c. = C. fasciculata.

GOMARA, Adans. Fam. ii. 248 (1763) = **Crassula**, Dill.

GOMARA, Ruiz & Pav. Prod. 93. t. 19 (1794). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 941.
racemosa, *Ruiz & Pav. Syst. Veg.* 103.—Peruv.

GOMARIA, Spreng. Gen. ii. 487 (1831) = prsec.

GOMESA, R. Br. in Bot. Mag. t. 1748 (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 565.
MATURNA, Rafin. Fl. Tellur. ii. 99 (1836).
chrysostoma, *Hoffmgg. Vera. Orch.—Cf. Bot. Zeit.* i (1843) 332.—Bras.
crispa, *Klotzsch, ex Reichb. f. in Bot. Zeit.* x. (1852) 772.—Bras.
Fischeri, *Regel, Ind. Sem. Hort. Petrop.* (1856) 21 ; *et in Ann. Sc. Nat.* Sér. IV. vi. (1856) 376.—Hab. ?
foliosa, *Klotzsch, ex Reichb. f. in Bot. Zeit.* x. (1852) 772.—Bras.
intermedia, *Hoffmgg. Vera. Orch.—Cf. Bot. Zeit.* i. (1843) 332.—Bras.
laxiflora, Klotzsch, ex Reichb. f. in Bot. Zeit. x. (1852) 772 = chrysostoma.
panciflora, *Hoffmgg. Vera. Orch.—Cf. Bot. Zeit.* i. (1843) 332.—Bras.
planifolia, *Klotzsch, ex Reichb. f. in Bot. Zeit.* x. (1852) 772.—Bras.
reclinata, *Hoffmgg. Vera. Orch.—Cf. Bot. Zeit.* i. (1843) 332.—Bras.
recurva, *R. Br. in Bot. Mag.* t. 1748.—Bras.
recurva, Lodd. Bot. Cab. t. 660 = planifolia.
sessilis, *Kodrig. Orch. Nov.* i. 143.—Bras.
stricta, Spreng. Syst. iii. 730 = Rodriguezia stricta.
tenuiflora, Lodd. Bot. Cab. t. 806 = Notylia punctata.
undulata, *Hoffmgg. Vera. Orch.—Cf. Bot. Zeit.* i. (1843) 332.—Bras.
viresccns, *Hoffmgg. ii. cc.*—Bras.

GOMESIA, La Llave, Reg. Trin. Mex. (1832) 40 ; ex DC. Prod. vii. 1. 262 (1838), homon (Compos.) (Quid ?).—Cf. Benth. & Hook. f. ii. 228.

GOMESIA, Spreng. Syst. iii. 729 (1826) = **Gomesa**, R. Br. (Orchid.).

GOMEZA, Lindl. Orch. Scel. 15 (1826) = prsec.

GOMEZIA, Bartl. Ord. Nat. 56 (1830) = prsec.

GOMEZIA, Benth. & Hook. f. Gen. ii. 933 (1873) = GOMESIA, La Llave (Compos.).

GOMIDESIA, Berg, in Linnæa, xxvii. (1854) 6 = **Myrcia**, DC. (Myrtac.).
amplexicaulis, Berg, in Mart. Fl. Bras. xiv. i. 13 = M. amplexicaulis.
anacardiifolia, Berg, l. c. 15 = M. anacardiifolia.
Aubletiana, Berg, l. c. 27.—Bras.
buxifolia, Berg, l. c. 25 = M. buxifolia.
Eryciblana, Berg, l. c. 23.—Bras.

GOMIDESIA :—
Blanchetiana, Berg, in Mart. Fl. Bras. xiv. 1. 14.—Bras.
Browniana, Berg, l. c. 18 = Myrcia Brownïana.
Cambessedeana, Berg, l. c. 24.—Bras.
Candolleana, Berg, l. c. 17.—Bras.
Casarettiana, Berg, l. c. 20.—Bras.
Chamissoeana, Berg, l. c. 11.—Bras.
Clausseniana, Berg, l. c. 28.—Bras.
crocea, Berg, l. c. 533.—Bras.
cricalyx, Berg, l. c. 25 = M. sericalyx.
Fensliana, Berg, l. c. 20.—Bras.
Freyreisiana, Berg, l. c. 19.—Bras.
Gardneriana, Berg, l. c. 22.—Bras.
Gaudichaudiana, Berg, l. c. 27.—Bras.
Haenkeana, Berg, l. c. 515.—Bras.
Hartwegiana, Berg, l. c. 22.—Bras.
hebepetala, Berg, l. c. 18.—Bras.
Hilariana, Berg, l. c. 26.—Bras.
Hookeriana, Berg, l. c. 18.—Bras.
Jacquiniana, Berg, l. c. 17.—Bras.
Klotzschiana, Berg, l. c. 534.—Bras.
Kunthiana, Berg, l. c. 28 = M. Kunthiana !
Langsdorffii, Berg, l. c. 531.—Bras.
Lindeniana, Berg, in Linnæa, xxix. (1857–58) 908.—Cuba.
Linkiana, Berg, in Mart. Fl. Bras. xiv. 1. 16 = M. Linkiana.
magnifolia, Berg, l. c. 531.—Bras.
Martiana, Berg, l. c. 12.—Bras.
Mikaniana, Berg, l. c. 21.—Bras.
Miqueliana, Berg, l. c. 24.—Bras.
Poeppigiana, Berg, l. c. 14.—Bras.
Pohliana, Berg, l. c. 16.—Bras.
Raddiana, Berg, l. c. 26.—Bras.
Regeliana, Berg, l. c. 535.—Bras.
reticulata, Berg, l. c. 15 = M. reticulata ? *Cambess.*
Riedeliana, Berg, l. c. 536.—Bras.
Schaueriana, Berg, l. c. 18.—Bras.
Sellowiana, Berg, l. c. 21.—Bras.
Sonderiana, Berg, l. c. 533.—Bras.
spectabilis, Berg, l. c. 1 = M. spectabilis.
Sprengeliana, Berg, l. c. 21.—Bras.
Springiana, Berg, l. c. 18.—Bras.
Spruceana, Berg, l. c. 534.—Bras.
Widgreniana, Berg, l. c. 26.—Bras.
Willdenowiana, Berg, l. c. 19.—Bras.

GOMIDEZIA, Benth. & Hook. f. Gen. i. 716 (1865) = prec.

GOMORTEGA, Ruiz & Pav. Prod. Fl. Per. 92. t. 10 (1794). *INC. SED.*, Benth. & Hook. f. iii. 149.
ADENOSTEMUM, Pers. Syn. i. 467 (1805).
KEULIA, Molina, Sagg. Chil. ed. II. 159 (1810).
nitida, *Ruiz & Pav. Fl. Per.* iv. t. 397 ; *Syst. Veg.* 108. —Peruv.

GOMOSIA, Lam. Encyc. ii. 769 (1786) = GOMOZIA, Mutis = **Nertera**, Banks & Soland. (Rubiac.).

GOMOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 87 = **Trichtmium**, R. Br. (Amarant.).
tomentosa, Turcz. l. c. = T. obovatum.

GOMOZIA, Mutis, ex Linn. f. Suppl. 17 (1781) = **Nertera**, Banks & Soland. (Rubiac.).
americana, Mirb. in Hist. Nat. Pl. xiii. [1805] 384 = N. depressa.
grenadensis, Mutis, ex Linn. f. Suppl. 129 = N. depressa.

GOMPHANDRA, Wall. Cat. nn. 3718, 7204 (1829) ; et in Lindl. Nat. Syst. ed. II. 489 (1836). *OLACINEÆ*, Benth. & Hook. f. i. 350.
SYMONURUS, Blume, Bijdr. 648 (1825), partim.
elliptica, *Mast. in Hook. f. Fl. Brit. Ind.* i. 586.—Malaya.
australiana, *F. Muell. Doc. Intercol. Exhib.* 38 ; *Fragm.* vi. 8, 253.—Austral.
axillaris, *Wall. Cat. n.* 3718.—Ind. or.
capitulata, *Becc. Malesia*, i. (1877) 111.—Malaya.
coriacea, Wight, Illustr. i. 108 = polymorpha.
cuspidata, *Mast. in Hook. f. Fl. Brit. Ind.* i. 587.—Burma.
Cumingiana, *Villar, in Blanco, Fl. Philipp.* ed. III. *Men. App.* 46.—Ins. Philipp.
laxiflora, *Vidal, Phan. Cuneing. Philipp.* 25, 103.—Ins. Philipp.

GOMPHANDRA :—
pensangiana, *Wall. Cat.* n. 7204.—Burma ; Malaya.
polymorpha, *Wight, Ic.* t. 958.—Ind. or.
prasina, *Becc. Malesia*, i. (1877) 110.—Malaya.
tomentella, *Mast. in Hook. f. Fl. Brit. Ind.* i. 587.—Burma.

GOMPHIA, Schreb. Gen. i. 291 (1789). *OCHNACEÆ*, Benth. & Hook. f. i. 318.
CHRYSOPHILUM, Willd. ex H. B. & K. Nov. Gen. et Sp. vi. 14 (1823).
CORREIA, Vell. in Roem. Script. 106. t. 6 (1796).
JABOTAPITA, Adans. Fam. ii. 364 (1763).
MESIA, Gaertn. Fruct. i. 344 (1788).
OURATEA, Aubl. Pl. Gui. i. 397. t. 152 (1775), nomen prius.
PHILOMACHA, Noronha, ex Thou. Gen. Mad. 17 (1806).
URATRA, J. F. Gmel. Syst. 677 (1791).
WOLKENSTEINIA, Regel, Gartenfl. xiv. (1865) 131. t. 471.
acuminata, *DC. in Ann. Mus. Par.* xvii. (1811) 419.—Bras.
arenaria, Pohl, Pl. Bras. Desc. ii. 120 = cuspidata.
affinis, *Hook. f. in Hook. Niger Fl.* 274.—Afr. trop.
alaternifolia, *A. Rich. Ess. Fl. Cub.* 339.—Cuba.
amplexicaulis, *O. Hoffm. Sert. Madag.* 9.—Madag.
anceps, *Baker, in Journ. Linn. Soc.* xxi. (1884) 330. —Madag.
angulata, *DC. in Ann. Mus. Par.* xvii. (1811) 415.—Madag.
angustifolia, *Vahl, Symb. Bot.* ii. 49.—Ind. or. ; Malaya.
aquatica, *H. B. & K. Nov. Gen. et Sp.* vi. 14.—Venezuela.
arguta, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1099, nomen.—Guian.
atropurpurea, *Schnizl. Ic.* iv. t. 248.—Hab. ?
australiana, F. Muell. Fragm. v. 39 = Brackenridgia australiana.
axillaris, *Oliver, Fl. Trop. Afr.* i. 322.—Afr. trop.
Blanchetiana, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 19.— Am. austr.
brachyandra, Planch. l. c. 18 = parvifolia.
bracteata, Wawra, Bot. Ergeb. Maxim. Bras. 29 = parvifiora.
calophylla, *Hook. f. in Hook. Niger Fl.* 274.—Afr. trop.
Candolleana, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 4 = guyannensis
caracasana, *Planch. l. c.* 9.—Venezuela.
carcinodes, *DC. in Ann. Mus. Par.* xvii. (1811) 421.—Gulan.
cassinefolia, *DC. l. c.*—Bras.
castaneæfolia, *DC. l. c.* 417.—Bras.
ceylanica, Spreng. Syst. ii. 313 = angustifolia.
Claessenii, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 16.—Bras.
cocchee, *Mart. ex Engl. in Mart. Fl. Bras.* xii. II. 311.—Bras.
conduplicata, *Klotzsch, ex Engl. l. c.* 344.—Bras.
confertiflora, *Pohl, Pl. Bras. Descr.* ii. 117.—Bras.
congesta, *Oliver, Fl. Trop. Afr.* i. 322.—Afr. trop.
crassifolia, *Pohl, Pl. Bras. Descr.* ii. 123.—Bras.
crenata, Spruce, ex Engl. in Mart. Fl. Bras. xii. II. 319 = acuminata.
curvata, *St. Hil. Fl. Bras. Mer.* i. 68.—Bras.
cuspidata, *St. Hil. l. c.* 67.—Bras.
decora, *Hort. in Rev. Hortic.* (1879) 311.—Bras.
decorata, Lem. Jard. Fleur. iv. t. 415 = oliveæformis.
deltoides, *Baker, in Journ. Linn. Soc.* xviii. (1881) 266.—Madag.
dentifera, Spruce, ex Engl. in Mart. Fl. Bras. xii. II. 319 = acuminata.
dependens, *DC. in Ann. Mus. Par.* xvii. (1811) 415.—Madag.
diantherula, Pohl, ex Engl. in Mart. Fl. Bras. xii. II. 309 = castaneæfolia.
divaricata, Pohl, ex Engl. l. c. 324 = semicrrata.
Duparquetiana, *Baill. Adansonia*, ix. (1868–70) 77.—Afr. trop.
dura, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1099, nomen.—Guian.
elliptica, *A. Rich. Ess. Fl. Cub.* 340.—Abyssin.
elongata, *Oliver, Fl. Trop. Afr.* i. 321.—Afr. trop.
erythrocalyx, Spruce, ex Engl. in Mart. Fl. Bras. xii. II. 326 = inundata.
Fieldingiana, *Gardn. in Field. & Garden. Sert. Pl.* t. 35.—Bras.

GOMPHIA :—
Finlaysoniana, *Wall. Cat.* n. 6392.—Ind. or.
fusca, Schum. & Thonn. Beskr. Guin. Pl. 216 = reticulata.
floribunda, *St. Hil. Fl. Bras. Mer.* i. 64.—Bras.
gigantophylla, *Erhard, in Flora*, xxxii. (1849) 250.—Bras.
glaberrima, *Beauv. Fl. Owar.* ii. 22. t. 71.—Afr. trop.
glaucescens, *St. Hil. Fl. Bras. Mer.* i. 68.—Bras.
glomerata, *Pohl, Pl. Bras. Descr.* ii. 122.—Bras.
grandiflora, *DC. in Ann. Mus. Par.* xvii. (1811) 420.—Bras.
grandifolia, *Planch. in Hook. Lond. Journ. Bot.* vi (1847) 4.—Bras.
guianensis, Aucr. vide guyannensis.
Guildingi, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 13 = nitida.
guyannensis, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 108.—Am. trop.
hexasperma, *St. Hil. Fl. Bras. Mer.* i. 61.—Bras.
Hookeri, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 8.—Ins. Penang.
humilis, Mart. ex Engl. in Mart. Fl. Bras. xii. II. 330 = spectabilis.
humilis, *St. Hil. Fl. Bras. Mer.* i. 66.—Bras.
ilicifolia, *DC. in Ann. Mus. Par.* xvii. (1811) 418.—Ins. S. Doming.
inundata, *Spruce, ex Engl. in Mart. Fl. Bras.* xii. II. 326.—Bras.
isophylla, *Carcke, in Linnæa*, xxii. (1849) 57.—Guian.
steudes, Erhard, in Flora, xxxii. (1849) 251 = stipulacea.
Jabotapita, *Sw Fl. Ind. Occ.* ii. 740.—Am. trop.
jamaicensis, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 11 = guyannensis.
Jurgensenii, *Planch. l. c.*—Mexic.
laevigata, Vahl, Symb. Bot. ii. 49 = obtusifolia.
lanceolata, *Baker, in Journ. Linn. Soc.* xxi. (1884) 330.—Madag.
lanceolata, *Pohl, Pl. Bras. Desc.* ii. 117.—Bras.
latifolia, Erhard, in Flora, xxxii. (1849) 248 = Fieldingiana.
laurifolia, *Sw. Fl. Ind. Occ.* ii. 741 —Ind. occ.
linearis, A. Gray, Bot. U. St. Expl. Exped. i. 359. t. 41 = stipulacea !
longifolia, *DC. in Ann. Mus. Par.* xvii. (1811) 417.—Ind. occ.
lucens, *H. B. & K. Nov. Gen. et Sp.* vii. 249.—N. Granat.
lucidula, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) t. 575.—Bras.
macranthera, Erhard, in Flora, xxxii. (1849) 248 = cassinefolia.
macrocarpa, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 2 = reticulata.
macrophylla, Oliver, Fl. Trop. Afr. i. 321 = reticulata.
macrostachys, Mart. ex Erhard, in Flora, xxxii. (1849) 246 = Fieldingiana.
Magdalenae, *Triana & Planch. in Ann. Sc. Nat*, Sér. IV. xviii. (1862) 278.—N. Granat.
magnoliaefolia, *Zipp. ex Span. in Linnæa*, xv. (1841) 186, nomen.—Ins. Timor.
mole'aefolia, DC. in Ann. Mus. Par. xvii. (1811) 416 = angustifolia.
Mannii, *Oliver, Fl. Trop. Afr.* i. 321.—Afr. trop.
membranacea, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 274.—N. Granat.
mexicana, *Humb. & Bonpl. Pl. Aequin.* ii. 21. t. 74.—Mexic.
micrantha, Hook. f. in Journ. Linn. Soc. vi. (1862) 8 = reticulata.
microdonta, *Benth. in Hook. Kew Journ.* iii. (1851) 271.—Bras.
Miersii, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 5.—Bras.
multiflora, *Pohl, Pl. Bras. Descr.* ii. 121.—Bras.
multiflora, Walp. Ann. i. 181 = Ochna multiflora.
nana, *St. Hil. Fl. Bras. Mer.* i. 66.—Bras.
nervosa, *St. Hil. l. c.* 65.—Bras.
nitida, DC. in Ann. Mus. Par. xvii. (1811) 419 = guyanensis.
nitida, Herb. Madr. ex Wall. Cat. sub n. 2802 = angustifolia.
obtusifolia, *DC. in Ann. Mus. Par.* xvii. (1811) 416.—Madag.
odora, *Poepp. ex Engl. in Mart. Fl. Bras.* xii. II. 334. —Bras.
oleaefolia, *St. Hil. Pl. Rem. Bris.* i. 194.—Bras.
oliveæformis, *St. Hil. Fl. Bras. Mer.* i. 67.—Bras.

GOMPHIA:—
ovalis, *Pohl, Pl. Bras. Descr.* ii. 118.—Bras.
ovata, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1189, nomen.—Guian.
parviflora, *DC. in Ann. Mus. Par.* xvii. (1811) 490.—Bras.
parviflora, Herb. Berol. ex Walp. Ann. i. 183 = Sellowii.
parvifolia, *St. Hil. Fl. Bras. Mer.* i. 65.—Bras.
pendula, *Poepp. ex Engl. in Mart. Fl. Bras.* xii. 11. 339.—Bras.
peruviana, *Baker, in Journ. Linn. Soc.* xxi. (1884) 330.—Madag.
pervicaria, St. Hil. Fl. Bras. Mer. i. 62 = semiserrata.
pinetorum, *Wright, ex Griseb. Cat. Pl. Cub.* 37.—Ind. occ.
Pohlii, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 5 = multiflora.
polita, *Presl, Bot. Bemerk.* 32.—Bras.
polyantha, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 274.—N. Granat.
pubescens, *St. Hil. & Tul. l. c.* Sér. II. xvii. (1842) 137.—Bras.
pulchella, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 7.—Bras.
pycnostachya, *Mart. ex Erhard, in Flora*, xxxii. (1849) 246.—Bras.
pyrifolia, *Griseb. Fl. Brit. W. Ind.* 105.—Ins. Trinit.
racemosa, Stend. ex Engl. in Mart. Fl. Bras. xii. 11. 344 = olivaeformis.
reticulata, *Beauv. Fl. Owar.* ii. 22. t. 72.—Afr. trop.
revoluta, *Wright, ex Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 108.—Cuba.
rotundifolia, *Field. & Gardn. Sert. Pl.* t. 34.—Bras.
rufidula, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 20 = Blanchetiana.
rupununiensis, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1189, nomen; *et ex Engl. in Mart. Fl. Bras.* xii. 11. 318.—Guian.
salicifolia, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 137.—Bras.
Schomburgkiniana, Klotzsch, in Abh. Berl. Acad. (1857) 238 = reticulata.
Schomburgkii, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 21.—Guian.
Sellowii, *Planch. l. c.* 5.—Bras.
semiserrata, *Mart. & Nees, in Nov. Act. Nat. Cur.* xii. (1824) 41.—Bras.
sepiaria, Mart. ex Engl. in Mart. Fl. Bras. xii. 11. 345 = cuspidata.
serratula, Pohl, Pl. Bras. Descr. ii. 119 = hexasperma.
spectabilis, *Mart. ex Engl. in Mart. Fl. Bras.* xii. 11. 330.—Bras.
squamosa, *DC. in Ann. Mus. Par.* xvii. (1811) 416.—Ind. occ.
stipelacea, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 6.—Bras.
stipulata, Planch. l. c. = prec.
snaveolens, *St. Hil. Fl. Bras. Mer.* i. 63.—Bras.
subscandens, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 14.—Bras.
subcrlutina, Planch. l. c. 17 = oleaefolia.
subverticillata, Erhard, in Flora, xxxiii. (1849) 254 = verticillata.
suffruticosa, *Bot. Hort. Maurit.* 67.—Madag.
sumatrana, *Jack, in Malay. Misc.* i. (1820) n. v. 20.—Malaya.
surinamensis, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 13 = subscandens.
Theophrasta, Planch. in Linden, Cat. n. 14 (1859) 8 = gigantophylla.
tomentella, Pohl, ex Engl. in Mart. Fl. Bras. xii. 11. 325 = oleaefolia.
Turnerae, Hook. f. in Hook. Niger Fl. 278 = reticulata.
umbellata, *Hook. f. ex A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 535, nomen.—Ins. Penang.
vaccinioides, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 137.—Bras.
Vauvae, *Spruce, ex Engl. in Mart. Fl. Bras.* xii. 11. 326.—Bras.
venosa, Sprang. Syst. iv. Cur. Post. 168 = nervosa.
verticillata, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 21.—Bras.
Vogelii, *Hook. f. in Hook. Niger Fl.* 272.—Afr. trop.
zeylanica, DC. in Ann. Mus. Par. xvii. (1811) 413 = angustifolia.

GOMPHICHIS, Lindl. Gen. et Sp. Orch. 446 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 504.
goodyeroides, *Lindl. l. c.* 447.—Peruv.
valida, *Reichb. f. Xenia Orch.* iii. 20.—Bolivia.

GOMPHIMA, Rafin. Fl. Tellur. ii. 10 (1836) = **Monochoria**, Presl (Pontederiaceae).
vaginalis, Rafin. l. c. = M. vaginalis.

GOMPHIPUS, Rafin. Fl. Tellur. iv. 76 (1836) = **Stemodia** ?, Linn. (Scrophul.).

GOMPHOCARPUS, R. Br. in Mem. Wern. Soc. i. (1809) 37. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 758.
ACERATES, Ell. Sketch, i. 316 (1817).
KERBELL, Harv. Gen. S. Afr. Pl. ed. II. 238 (1868).
PACHYCARPUS, E. Mey. Comm. Pl. Afr. Austr. 209 (1837).
POLYOTUS, Nutt. in Trans. Am. Phil. Soc. N. S. v. 160.—Mexic.
arborescens, *R. Br. in Mem. Wern. Soc.* i. (1809) 38.—Afr. austr.
asper, *Decne. in DC. Prod.* viii. 561.—Afr. austr.
asperifolius, *Meisen. in Hook. Lond. Journ. Bot.* ii. (1843) 448.—Afr. austr.
brasiliensis, *Fourn. in Mart. Fl. Bras.* vi. iv. 208.—Bras.
brevicuspis, *D. Dietr. Syn. Pl.* ii. 900.—Afr. austr.
campanulatus, *Harv. Thes. Cap.* i. 61.—Afr. austr.
chironioides, Decne. in DC. Prod. viii. 562 = Listanthoides.
concolor, *Decne. l. c.* 568.—Afr. austr.
cordifolius, *Benth. ex A. Gray, in Bot. Calif.* i. 477.—Calif.
corniculatus, *D. Dietr. Syn. Pl.* ii. 901.—Afr. austr.
cornutus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 294.—Afr. austr.
coronarius, *Decne. in DC. Prod.* viii. 562.—Afr. austr.
crinitus, *Bertol. f. in Mem. Acad. Sc. Bologn.* iii. (1851) 258.—Afr. trop.
crispus, *R. Br. in Mem. Wern. Soc.* i. (1809) 38.—Afr. austr.
cristatus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 292.—Afr. austr.
dealbatus, *Decne. in DC. Prod.* viii. 563.—Afr. austr.
diploglossus, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) 1. 228.—Afr. austr.
dregeanophanus, Hochst. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 110 = robustus.
eminens, *Harv. Thes. Cap.* ii. 60. t. 196.—Afr. austr.
eustegioides, *D. Dietr. Syn. Pl.* ii. 901.—Afr. austr.
expansus, *D. Dietr. l. c.*—Afr. austr.
filiformis, D. Dietr. l. c. 900 = Asclepias filiformis (in Addend.).
flexuosus, *D. Dietr. l. c.* 901.—Afr. austr.
fruticosus, *R. Br. in Mem. Wern. Soc.* i. (1809) 38.—Reg. Mediterr.
Gerrardi, *Harv. Thes. Cap.* ii. 59. t. 192.—Afr. austr.
globus, *D. Dietr. Syn. Pl.* ii. 900.—Afr. austr.
glabertimus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 110.—Afr. trop.
gracilis, D. Dietr. Syn. Pl. ii. 901 (= *Lagarinthus gracilis*).—Afr. austr.
grandiflorus, *Decne. in DC. Prod.* viii. 563.—Afr. austr.
hastatus, *E. Mey. Comm. Pl. Afr. Austr.* 201.—Afr. austr.
humilis, *Decne. in DC. Prod.* viii. 561.—Afr. austr.
hypoleucus, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 232.—Arizona.
interruptus, D. Dietr. Syn. Pl. ii. 901 = Schizoglossum interruptum.
involucratus, D. Dietr. l. c. 900 = Xysmalobium involucratum.
laevigatus, Buch.-Ham. ex Decne. in DC. Prod. viii. 565, 646 = Brachystoma laevigatum.
lanatus, E. Mey. Comm. Pl. Afr. Austr. 202 = tomentosus, *Burch.*

GOMPHOCARPUS:—
linearis, *D. Dietr. Syn. Pl.* ii. 901.—Afr. austr.
linoides, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 296.—Afr. trop.
listanthoides, *Decne. l. c.* 295.—Afr. trop.
longifolius, *Sprang. Syst.* i. 840.—Am. bor.
longipes, Oliver, in Trans. Linn. Soc. xxix. (1875) 111. t. 76, sphalm. = Asclepias macrantha.
mater, D. Dietr. Syn. Pl. ii. 901 (= *Listanthus macer*).—Afr. austr.
macrogiossus, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) 1. 259.—Afr. austr.
M'Kenii, *Harv. Thes. Cap.* ii. 60. t. 198.—Afr. austr.
marginatus, *Decne. in DC. Prod.* viii. 560.—Afr. austr.
multicaulis, *D. Dietr. Syn. Pl.* ii. 901.—Afr. austr.
multiflorus, *Decne. in DC. Prod.* viii. 560.—Afr. austr.
navicularis, *D. Dietr. Syn. Pl.* ii. 901.—Afr. austr.
niveus, Boj. ex Baker, Fl. Maurit. 228 = cornutus.
nutans, *Klotzsch, in Peters, Reise Mossamb. Bot.* 275.—Afr. trop.
oxyropis, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) 1. 259.—Afr. austr.
patifolius, Baker, in Saund. Refug. Bot. t. 254 (Xysmalobii sp.).—Afr. austr.
pauciflorus, Hochst. & Steud. ex Steud. Nom. ed. II. i. 698 = Glossonema Boveanum.
pauciflorus, Klotzsch, in Peters, Reise Mossamb. Bot. 276 = Asclepias macrantha.
pedunculatus, Decne. in DC. Prod. viii. 558 = Asclepias macrantha.
peltigerus, *D. Dietr. Syn. Pl.* ii. 901.—Afr. austr.
physocarpus, *E. Mey. Comm. Pl. Afr. Austr.* 202.—Afr. austr.
pulchellus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 325.—Afr. trop.
purpurascens, A. Gray, in Proc. Am. Acad. x. (1874) 76 = Schizonotus purpurascens.
purpurascus, *A. Rich. Tent. Fl. Abyss.* ii. 38.—Abyssin.
reflectens, *Decne. in DC. Prod.* viii. 568.—Afr. austr.
revolutus, *D. Dietr. Syn. Pl.* ii. 901.—Afr. austr.
rigidus, *Decne. in DC. Prod.* viii. 563.—Afr. austr.
robustus, *A. Rich. Tent. Fl. Abyss.* ii. 40.—Abyssin.
rubiolides, *Kotschy & Peyr. Pl. Tinn.* 29.—Afr. trop.
scaber, *Harv. Thes. Cap.* ii. 58. t. 192.—Afr. austr.
Schimperi, Presl, Bot. Bemerk. 103 = fruticosus.
semilunatus, A. Rich. Tent. Fl. Abyss. ii. 39 = physocarpus.
sessilis, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 325 = Xysmalobium sessile.
setosus, *R. Br. in Mem. Wern. Soc.* i. (1809) 38.—Arab.
sinicus, Hochst. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 110 = fruticosus.
sinuatus, Boiss. Diagn. Ser. I. xi. 80 = fruticosus.
stenophyllus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 110.—Afr. trop.
tenuis, D. Dietr. Syn. Pl. ii. 900 (= *Lagarinthus tenuis*).—Afr. austr.
tomentosus, *Burch. Trav.* i. 548.—Afr. austr.
tomentosus, *A. Gray, in Bot. Calif.* i. 477.—Calif.
truncatus, *D. Dietr. Syn. Pl.* ii. 901.—Afr. austr.
truncatus, Harv. Thes. Cap. i. 42. t. 67 = peltigerus.
undulatus, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) 1. 259.—Afr. austr.
ventricosus, *Turcz. l. c.* 258.—Abyssin.
virgatus, D. Dietr. Syn. Pl. ii. 901 (= *Lagarinthus virgatus*).—Afr. austr.
viridiflorus, *Decne. in DC. Prod.* viii. 561.—Afr. austr.
viridiflorus, Sprang. Syst. i. 849 (= *Acerates viridiflora*).—Am. bor.
viridis, Sprang. l. c. = Asclepiodora viridis.
wakefildia, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 21 = Holostemma Rheedianum.

GOMPHOGYNE, Griff. Pl. Cantor. 96. t. 4 (1887?). *CUCURBITACEAE*, Benth. & Hook. f. i. 836.
TRICEROS, Griff. Notul. iv. 606 (1854).
cissiformis, *Griff. Pl. Cantor.* 96. t. 4.—Sing. Himal.
heterogynaea, *Kuntz, in Journ. As. Soc. Beng.* xl. (1871) ii. 68.—Burma.

GOMPHOLOBIUM, Sm. in Trans. Linn. Soc. iv. (1798) 220. *LEGUMINOSAE*, Benth. & Hook. f. i. 468.
aciculare, Reichb. in Exot. t. 249 = tomentosum.
amplexifolium, *Meisn. in Lehm. Pl. Preiss.* i. 84.—Austral.

GOMPHOLOBIUM:—
aristatum, *Benth. in Enum. Pl. Hueg.* 29.—Austral.
asphaltoides, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 72 = virgatum.
barbigerum, DC. Prod. ii. 105 = latifolium.
Baxteri, *Benth. Fl. Austral.* ii. 46.—Austral.
bartonioides, *Meisn. in Lehm. Pl. Preiss.* i. 37.—Austral.
capitatum, *A. Cunn. in Lindl. Bot. Reg.* t. 1568.—Austral.
Drummondii, Meisn. in Bot. Zeit. xiii. (1855) 25 = aristatum.
elliptieum, Labill. Nov. Holl. Pl. i. 106. t. 135 = Oxylobium ellipticum.
ericaefolium, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 72 = Burtonia conferta.
fimbriatum, Sm. Exot. Bot. i. 113 = latifolium.
foliolosum, Benth. in Mitch. Journ. Trop. Austral. 346 = Burtonia foliolosa.
furcellatum, Bonpl. Jard. Malm. 30. t. 11 = Jacksonia furcellata.
glabratum, *DC. Prod.* ii. 106.—Austral.
glaucescens, A. Cunn. in Field, N. S. Wales, 346 = grandiflorum.
grandiflorum, Andr. Bot. Rep. t. 642 = polymorphum.
grandiflorum, *Sm. Exot. Bot.* i. 7.—Austral.
Hendersonii, Paxt. Mag. Bot. xi. (1844) 108 = Burtonia Hendersonii.
heterophyllum, A. Cunn. ex G. Don, Gen. Syst. ii. 118 = Knightianum.
hirsutum, *Pexl. Mag. Bot.* xvi. (1849) 35.—Austral.
Huegelii, *Benth. in Enum. Pl. Hueg.* 29.—Austral.
Knightianum, *Lindl. Bot. Reg.* t. 1468.—Austral.
lanatum, A. Cunn. ex G. Don, Gen. Syst. ii. 118 = tomentosum.
latifolium, Labill. Nov. Holl. Pl. i. 105 = Huegelii.
latifolium, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 506.—Austral.
maculatum, Andr. Bot. Rep. t. 427 = grandiflorum.
marginatum, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 11.—Austral.
minus, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 505.—Austral.
nitidum, *Soland. ex Benth. Fl. Austral.* ii. 48.—Austral.
obcordatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 358.—Austral.
ovatum, *Meisn. in Lehm. Pl. Preiss.* i. 35.—Austral.
paduncular, DC. Prod. ii. 105 = polymorphum.
paduncular, Lodd. Bot. Cab. t. 1639 = Huegelii.
pinnatum, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 506.—Austral.
polymorphum, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 11.—Austral.
polymorphum, Sieber, ex Benth. Fl. Austral. ii. 48 = glabratum.
polyzygum, F. Muell. Fragm. iii. 29 = Burtonia polyzyga.
Preissii, *Meisn. in Lehm. Pl. Preiss.* i. 40.—Austral.
psoraleaefolium, Salisb. Parad. Lond. t. 6 = latifolium.
scabrum, Sm. in Trans. Linn. Soc. ix. (1808) 250 = Burtonia scabra.
setifolium, Sieber, ex Benth. in Ann. Wien. Mus. ii. (1838) 72 = grandiflorum.
Shuttleworthii, *Meisn. in Lehm. Pl. Preiss.* i. 39.—Austral.
sparsum, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 72 = polymorphum.
spinosum, Labill. Nov. Holl. Pl. i. 107. t. 136 = Jacksonia spinosa.
stenophyllum, F. Muell. Fragm. iii. 30 = Burtonia subulata.
subulatum, Benth. in Ann. Wien. Mus. ii. (1838) 72 = Burtonia subulata.
tenue, Lindl. Bot. Reg. t. 1615 = polymorphum.
tetrathecoides, Sieber, ex DC. Prod. ii. 106 = minus.
tomentosum, *Labill. Nov. Holl. Pl.* i. 106. t. 134.—Austral.
undulatum, *A. Cunn. ex Benth. in Ann. Wien. Mus.* ii. (1838) 72.—Austral.
venulosum, Lindl. Bot. Reg. t. 1574 = polymorphum.
venustum, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 12.—Austral.
verticillo, Lindl. Bot. Reg. (1839) t. 43. Misc. 45 = polymorphum.
virgatum, *Meisn. ex DC. Prod.* ii. 106.—Austral.
viridiflorum, *Meisn. in Lehm. Pl. Preiss.* i. 39.—Austral.

GOMPHOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xv. (1841) 587 = **Angelica**, Linn. (Umbell.).
ailsfordense, Turcz. l. c. 540.—Siberia.
Maximowiczii, F. Schmidt, in Maxim. Prim. Fl. Amur. 126 = A. Maximowiczii.
viridiflorum, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 741.—Sibir, baical.

GOMPHOSIA, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 14 (Monog. Cinch. 95. t. 26 B) = **Ferdinandusa**, Pohl (Rubiac.).
chlorantha, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 14.—Peruv.
cinchonoides, Wedd. l. c.—N. Granat.
lanciflora, Benth. in Hook. Kew Journ. v. (1853) 282 = *chlorantha*.
longiflora, Benth. l. c.—Bras.

GOMPHOSTEMA, Hassk. Cat. Hort. Bogor. Alt. 138 (1844) = seq.

GOMPHOSTEMMA, Wall. Cat. n. 2151 (1830 †); et Pl. As. Rar. ii. 12 (1831). *LABIATAE*, Benth. & Hook. f. ii. 1216.
GOMPHOSTEGIA, Reichb. Handb. 189 (1837).
acaule, *Kurz, ex Hook. f. Fl. Brit. Ind.* iv. 696.—Burma.
chinense, *Oliver, in Hook. Ic. Pl.* t. 1468.—China.
crinitum, *Wall. Cat.* n. 2159.—Burma; Malaya.
dichotomum, *Coll. & Mor. Syst. Vers. Zoll.* 54.—Java.
eriocephalum, *Benth. in Wall. Pl. As. Rar.* ii. 12.—Ind. or.
flavescens, Miq. Fl. Ind. Bat. ii. 987 = Cymaria acuminata.
floyeriana, Wall. Cat. n. 2152 = strobilinum.
insuave, *Hance, in Journ. Bot.* xxii. (1884) 231.—China.
javanica, *Benth. Lab. Gen. et Sp.* 650.—Java.
lucidum, *Wall. Cat.* n. 2156.—Reg. Himal.
macrophyllum, Miq. Fl. Ind. Bat. ii. 968 = Leonurus oblongifolius.
Masters*ii*, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 699.—Reg. Himal.
melissaefolium, *Wall. Cat.* n. 2157.—Reg. Himal.
multiflorum, Wall. l. c. partim = Masters*ii*.
membranifolium, Miq. Fl. Ind. Bat. ii. 988 = Phlomis rugosa.
multiflorum, Benth. in DC. Prod. xii. 552 = parviflorum.
niveum, *Hook. f. Fl. Brit. Ind.* iv. 697.—Bengal.
nutans, *Hook. l. c.*—Reg. Himal.
oblongum, *Wall. Cat.* n. 2154.—Burma.
oblongum, Wight, Ic. t. 1457 = eriocarpum.
ovatum, *Wall. Cat.* n. 2155.—Reg. Himal.
parviflorum, *Wall. l. c.* n. 2158.—Reg. Himal.; Malaya.
pedunculatum, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 696.—Reg. Himal.
petiolare, Miq. Fl. Ind. Bat. ii. 987 = Leonurus javanicus.
philippinum, *Benth. in DC. Prod.* xii. 551.—Ins. Philipp.
phlomoides, *Benth. Lab. Gen. et Sp.* 649.—Java.
strobilinum, *Wall. Cat.* n. 2151.—Ind. or.; Burma.
Thomsoni, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 698.—Reg. Himal.
verticillatum, *Benth. in Wall. Pl. As. Rar.* ii. 12.—Reg. Himal.
viride, Wall. Cat. n. 2158 = strobilinum.

GOMPHOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 53. *LOGANIACEAE*, Benth. & Hook. f. ii. 792.
incanum, *Oliver, in Hook. Ic. Pl.* t. 1472.—Afr. austr.
scoparioides, *Turcz. in Bull. Soc. Nat. Mosc.* xvi. (1843) 53.—Afr. austr.

GOMPHOSTYLIS, Rafin. Fl. Tellur. ii. 80 (1836) = **Eygadenus**, Michx. (Liliac.).
bracteata, Rafin. l. c. = Z. glaucus.
fuscata, Rafin. l. c. = Z. fuscus.
paniculata, Rafin. l. c. = Z. paniculatus.

GOMPHOSTYLIS, Wall. ex Lindl. Gen. et Sp. Orch. 43 (1830) = **Oceloglyna**, Lindl. (Orchid.).
candida, Wall. ex Lindl. l. c. = C. maculata.

GOMPHOTIS, Rafin. Sylva Tellur. 108 (1838) = **Thrypstonanne**, Endl. (Myrtac.).
acuicola, Rafin. l. c. = T. acuicola.

GOMPHRAENA, Jacq. Select. Am. 88 (1763) = seq.

GOMPHRENA, Linn. Gen. ed. I. 69 (1737). *AMARANTACEAE*, Benth. & Hook. f. iii. 40.
BRAGANTIA, Vand. Fasc. Pl. Nov. 6 (1771).
CHNOANTHUS, Phil. in Anal. Univ. Chili (1862), II. 605.
COLUPPA, Adans. Fam. ii. 268 (1763).
NIWANGA, Rafin. Fl. Tellur. iii. 76 (1836).
SCHULTESIA, Schrad. in Goett. Gel. Anz. i. (1821) 708.
WADAPUS, Rafin. Fl. Tellur. iii. 77 (1836).
XEROSIPHON, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 55.
acaulis, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 250 —Bolivia.
affinis, *F. Muell. ex Benth. Fl. Austral.* v. 264.—Austral.
aggregata, Hort. Monsp. ex Moq. in DC. Prod. xiii. II. 414 = perennis.
aggregata, Willd. Enum. Hort. Berol. 294 = Philoxerus aggregatus.
agrestis, *Mart. Nov. Gen. et Sp.* ii. 18. t. 114.—Bras.
albiflora, *Moq. in DC. Prod.* xiii. II. 392.—Venezuela.
amaranthoides, Roth, Nov. Pl. Sp. 182 = Iresine amarantoides.
angustiflora, *Mart. Nov. Gen. et Sp.* ii. 19. t. 121.—Bras.
angustifolia, *Vahl, Symb. Bot.* iii. 45.—Ind. or.
aphylla, *Pohl, ex Moq. in DC. Prod.* xiii. II. 416.—Bras.
arborescens, Balb. ex Moq. l. c. 353 = Altemanthera argentata.
arborescens, *Linn. f. Suppl.* 173.—Bras.
argentea, Hort. ex Moq. in DC. Prod. xiii. II. 410 = decumbens.
argentina, *Saub. in Mart. Fl. Bras.* v. i. 208.—Bras.
bicolor, Hort. ex Moq. in DC. Prod. xiii. II. 410 = decumbens.
bicolor, *Mart. in Nov. Act. Nat. Cur.* xiii. (1826) 300.—Peruv.
Blanchetii, *Moq. in DC. Prod.* xiii. II. 413.—Bras.
boliviana, *Moq. l. c.* 401.—Bolivia.
bonariensis, Gill. ex Moq. l. c. 406 = pulchella.
brachiata, *Hemsl. in Bonpl. ex Roem. & Schult. Syst.* v. 542.—Venezuela.
brachystylis, *F. Muell. Fragm.* iii. 124.—Austral.
brasiliana, Linn. Cent. Pl. ii. 13; Amoen. Acad. iv. 10 = Mogiphanes stramínea.
brasilienzis, Hort. Aven. ex Moq. in DC. Prod. xiii. II. 364 = Telanthera polygonoides.
brasilianus, Coll. ii. 278 = Telanthera dentata.
brasilienzis, Lam. Illustr. ii. 289. t. 180. f. 1 = *brasiliana*.
brasilienzis, Linn. Syst. ed. X. 949 = Mophigenes stramínea.
brevifiora, F. Muell. Fragm. iii. 125 = conica.
Brownii, *Moq. in DC. Prod.* xiii. II. 397.—Austral.
caespitosa, *Torr. in Bot. Mex. Bound.* 181.—N. Mexic.
callosa, *Sweet, Hort. Brit.* ed. III. 571.—Am. austr.
canescens, *R. Br. Prod.* i. 416.—Austral.
celosioides, *Mart. in Nov. Act. Nat. Cur.* xiii. (1826) 301.—Bras.
cinerea, Moq. in DC. Prod. xiii. II. 390 (Pfaffiae sp.). —Hab.?
Claussenii, *Moq. l. c.* 396.—Bras.
coccinea, *Lamk. Encyc. Pl.* 1543.—Mexic.
conferta, *Benth. Fl. Austral.* v. 267.—Austral.
conica, *Spreng. Syst.* i. 894.—Austral.
correntina, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 30; *Contrib.* 91.—Reg. Argent.
crassifolia, Spreng. Syst. i. 894 = Iresine crassifolia.
Crucis, Vahl, ex Moq. in DC. Prod. xiii. II. 362 = Telanthera Crucis.
cylindrica, *Schum. & Thonn. Beskr. Guin. Pl.* 158.—Afr. trop.
debilis, *Mart. Nov. Gen. et Sp.* ii. 7. t. 107.—Bras.
decipiens, *Saub. in Mart. Fl. Bras.* v. i. 211.—Bras.
decumbens, *Vacq. Hort. Schoenbr.* t. 483.—Mexic.
decumbens, Lag. Elench. (1816) 17 = bolivíana.
demissa, *Mart. Nov. Gen. et Sp.* ii. 6.—Bras.
dentata, Moench, Meth. Suppl. 278 = Telanthera dentata.
denudata, Moq. in DC. Prod. xiii. II. 388. (Pfaffiae sp.).—Bras.
desertorum, *Mart. Nov. Gen. et Sp.* ii. 3. t. 103.—Bras.

GOMPHRENA:—
diffusa, Pav. ex Moq. in DC. Prod. xiii. II. 410 = decumbens.
diffusa, *Spreng. Syst.* i. 824.—Amaral.
Dunaliana, *Moq. in DC. Prod.* xiii. II. 384.—Bolivia.
durinscula, *Moq. l. c.* 404.—Bras.
elata, A. Cunn. ex Moq. l. c. 282 = Ptilotus conicus.
elegans, *Mart. Nov. Gen. et Sp.* ii. 17. t. 119.—Bras.
elongata, Willd. in Roem. & Schult. Syst. v. 542 = Telanthera elongata.
erianthe, Moq. in DC. Prod. xiii. II. 386 = Hebanthe virgata.
eriophylla, *Mart. Nov. Gen. et Sp.* ii. 9. t. 109.—Bras.
eriopoda, Gill. ex Moq. in DC. Prod. xiii. II. 410 = globosa.
exaltata, Delile, ex Moq. l. c. 490 = Froelichia floridana.
exaltata, Hort. Par. ex Moq. l. c. 418 = Froelichia floridana ?
exaltata, Hort. Tolos. ex Moq. l. c. 414 = perennis.
fallax, *Seub. in Mart. Fl. Bras.* v. t. 220.—Bras.
flexides, Jacq. Select. Am. 88 = Alternanthera frutescens.
flexides, Linn. Sp. Pl. 225 = Telanthera ficoides.
flagelloides, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 850.—Mexic.
firma, F. Muell Fragm. iii. 123 = flaccida.
flaccida, *R. Br. Prod.* i. 416.—Austral.
flaccida, Spreng. ex Moq. in DC. Prod. xiii. II. 417 = angustifolia.
flava, *Linn. Sp. Pl.* 224.—Mexic.
flava, Fav. ex Moq. in DC. Prod. xiii. II. 410 = decumbens.
floridana, Spreng. Syst. i. 824 = Froelichia floridana.
fruticosa, Hort. ex Moq. in DC. Prod. xiii. II. 377 = Telanthera porrigens.
fulgens, Hort. ex Moq. in DC. l. c. = Telanthera porrigens.
fusca, Mart. ex Moq. l. c. 406 = scapigera.
Gardneri, *Moq. l. c.* 404.—Bras.
glabrata, Moq. l. c. 388 = Pfaffia glabrata.
glauca, Moq. l. c. 384 = Pfaffia glauca.
globosa, *Linn. Sp. Pl.* 224.—Gerontr. trop.
gnaphaloides, Vahl, in Roem. & Schult. Syst. v. 542 = Pfaffia gnaphaloides.
gnaphaloides, *Griseb. in Gott. Abh.* xxiv. (1879) 34. —Reg. Argent.
gracilis, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 850 = Telanthera gracilis.
graminea, *Moq. in DC. Prod.* xiii. II. 416.—Bras.
Haageana, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 297.—Mexic.
Hanbaxa, *Mart. in Nov. Act. Nat. Cur.* xiii. (1826) 299.—Peruv.
helichrysoides, Moq. in DC. Prod. xiii. II. 391 (Pfaffiae sp.).—Bras.
Hilariana, *Moq. l. c.* 393.—Bras.
hirsutissima, Mart. ex Moq. l. c. 406 = scapigera.
hirtula, Moq. l. c. 388 = Pfaffia hirtula.
hispida, *Linn. Sp. Pl.* ed. II. 296.—Ind. or.
holosericea, Moq. in DC. Prod. xiii. II. 386 = Hebanthe holosericea.
Humboldtiana, Roem. & Schult. Syst. v. 540 = Froelichia Humboldtiana.
humilis, *R. Br. Prod.* i. 416.—Austral.
hygrophila, *Mart. in Flora*, xxiv. (1841) II. Beibl. 66. —Bras.
incana, *Mart. Nov. Gen. et Sp.* ii. 11. t. 112.—Bras.
interrupta, Linn. Sp. Pl. 224 = Froelichia lanata.
irsutissima, *Moq. in DC. Prod.* xiii. II. 388.—Bras.
jubata, Moq. l. c. 389 = Pfaffia jubata.
lactea. DC. Cat. Hort. Monsp. 113 = Aerva sanguinolenta.
lanata, R. Br. Prod. 416 = Brownii.
lanata, H. B. & K. Nov. Gen. et Sp. ii. 202 = Froelichia Humboldtiana.
lanata, Poir. Encyc. Suppl. i. 815 = Pfaffia tomentosa.
lanciflola, *Fav. ex Moq. in DC. Prod.* xiii. II. 394.—N. Granat.
lanigera, *Pohl, ex Moq. l. c.* 406.—Bras.
lanuginosa, *Spanz. in Linnaea*, xv. (1841) 346.—Ins. Timor.
latifolia, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 849 = Iresine latifolia.
leptoclada, *Benth. Fl. Austral.* v. 255.—Austral.
leucocephala, *Mart. Nov. Gen. et Sp.* ii. 10. t. 110.— Bras.
ligulata, *Griseb. in Gott. Abh* xix. (1874) 80.—Reg. Argent.

GOMPHRENA:—
Lundii, *Moq. in DC. Prod.* xiii. II. 408.—Bras.
luzulaefolia, *Moq. l. c.* 384.—Bras.
macrocephala, *St. Hil. Fl. Us. Bras.* t. 92.—Bras.
macrorhiza, *Mart. Nov. Gen. et Sp.* ii. 6. t. 106.— Bras.
marginata, *Seub. in Mart. Fl. Bras.* v. t. 209.—Bras. Maitlandi, *F. Muell. Fragm.* iii. 124. t. 28.—Austral. Martiana, *Gill. ex Moq. in DC. Prod.* xiii. II. 400.— Chili.
Martinsiana, Steud. Nom. ed. II. i. 699 = angustiflora.
Meyeniana, *Wight. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 404.—Peruv.
microcephala, *Moq. in DC. Prod.* xii. II. 394.—Bras.
minima, Fav. ex Moq. l. c. 414 = leucocephala.
minutissima, Mart. ex Seub. in Mart. Fl. Bras. v. t. 211.—scapigera.
mollis, *Mart. Nov. Gen. et Sp.* ii. 11. t. 111.—Bras. Moquini, *Seub. l. c.* 209.—Bras.
mucronata, *Moq. in DC. Prod.* xiii. II. 418.—Bras.
nigricans, *Mart. Nov. Gen. et Sp.* ii. 12. tt. 113, 115. f. 1.—Bras.
nitida, *Rohr. in Wheeler, Rep. Bot. Geogr. Surv.* vi. 226.—Mexic.
officinalis, Mart. Reise Bras. i. 280 = arborescens.
oligocephala, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 850.—Bolivia.
paniculata, Moq. in DC. Prod. xiii. II. 385 = Hebanthe paniculata, *Benth. & Hook f.*
parviflora, *Benth. Fl. Austral.* v. 257.—Austral.
patula, Wendl. Bot. Beobacht. 48 = Mogiphanes stamines.
Pearcei, *Oliver, in Hook. Ic. Pl.* t. 1073.—Peruv.
perennis, *Linn. Sp. Pl.* 224.—Reg. Argent.
phagnaloides, *Griseb. in Gott. Abh.* xix. (1874) 82.— Reg. Argent.
pilosa, *Moq. in DC. Prod.* xiii. II. 395.—Mexic.
Pohlii, *Moq. l. c.* 408.—Bras.
Poiretiana, Roem. & Schult. Syst. v. 541 = Pfaffia tomentosa.
polygonoides, Linn. Sp. Pl. 225 = Telanthera polygonoides.
polygogon, Moq. ex Seub. in Mart. Fl. Bras. v. t. 196 = jubata.
procumbens, Pav. ex Moq. in DC. Prod. xiii. II. 370 = Telanthera obovata.
procumbens, Zuccagni, in Roem. Collect. 134 = Alternanthera proomnbens.
prostrata, *Natt. ex Spreng.* ed. I. 219 = decumbens.
prostrata, *Mart. Nov. Gen. et Sp.* ii. 5. t. 105.—Bras.
pulchella, *Mart. in Nov. Act. Nat. Cur.* xiii. (1826) 302. ; Reg. Argent.
pulverulenta, Moq. in DC. Prod. xiii. II. 386 = Hebanthe pulverulenta, *Benth. & Hook f.*
pulverulenta, Moq. l. c. partim = Hebanthe Hookeriana.
pumila, *Gill. ex Moq. l. c.* 400.—Reg. Argent.
pungens, *Seub. in Mart. Fl. Bras.* v. t. 201.—Bras.
purpurea, Hort. ex Moq. in DC. Prod. xiii. II. 377 = Telanthera porrigens.
pusilla, *Benth. Fl. Austral.* v. 256.—Austral.
Regeliana, *Seub. in Mart. Fl. Bras.* v. t. 216.—Bras.
reticulata, *Seub. l. c.* 194.—Bras.
Riedelii, *Seub. l. c.* 212.—Bras.
rodantha, *Moq. in DC. Prod.* xiii. II. 414.—Bras.
rosea, *Griseb. in Gott. Abh.* xix. (1874) 80.—Reg. Argent.
rubra, Hort. ex Moq. in DC. Prod. xiii. II. 409 = globosa.
rudis, *Moq. l. c.* 402.—Bras.
rupestris, *Nees, in Flora*, iv. (1821) 396, 397.—Bras.
scapigera, *Mart. Nov. Gen. et Sp.* ii. 14. tt. 116, 117. f. 1.—Bras.
Schlechtendaliana, *Mart. in Nov. Act. Nat. Cur.* xiii. (1826) 299.—Bras.
Schultesii, Mart. l. c. 302 = rupestris.
Sellos, Spreng. Syst. iv. Cur. Post. 104 = seq.
Sellowiana, *Mart. Nov. Gen. et Sp.* ii. 15. t. 117.— Bras.
sericantha, *Mart. l. c.* 8. t. 108.—Bras.
sericea, Hoffmgg. ex Roem. & Schult. Syst. v. 542 = Froelichia sericea.
sericea, *Moq. in DC. Prod.* xiii. II. 365.—Ecuador.
sericea, Spreng. Syst. i. 823 = Pfaffia sericea.
serrata, *Linn. Sp. Pl.* 224.—Am. austr.
serrata, Fav. ex Moq. in DC. Prod. xiii. II. 410 = decumbens.
sessilis, Linn. Sp. Pl. 225 = Alternanthera sessilis.

GOMPHRENA:—
Sonorae, *Torr. Bot. Mex. Bound.* 181.—N. Mexic.
spicata, Lam. Encyc. i. 130 = Froelichia lanata.
spicata, Moq. in DC. Prod. xiii. II. 387 = Hebanthe spicata.
stenophylla, Spreng. Syst. i. 823 = Pfaffia glauca.
strigosa, *Isert, ex Roem. & Schult. Syst.* v. 542.— Afr. trop.
suffruticosa, *Griseb. in Gott. Abh.* xxiv. (1879) 33.— Reg. Argent.
tenella, *Benth. Fl. Austral.* v. 256.—Austral.
tomentosa, Pohl, ex Moq. in DC. Prod. xiii. II. 391 = Pfaffia gnaphaloides.
trichotoma, Hort. ex Moq. l. c. 377 = Telanthera porrigens.
tripicata, Pav. ex Moq. l. c. 410 = decumbens.
tuberifera, *Torr. Bot. Mex. Bound.* 181.—N. Mexic.
tuberosa, Spreng. Syst. i. 823 = Pfaffia sericea.
tumida, Seidl, ex Opiz, Natural. n. ix. (1825) 154 = decumbens ?
umbellata, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 849.—Bolivia.
vaga, Mart. Nov. Gen. et Sp. ii. 17. t. 120 = Hebanthe vaga.
velutina, Moq. in DC. Prod. xiii. II. 389 = Pfaffia velutina.
vermicularis, Linn. Sp. Pl. 224 = Philoxerus vermicularis.
villosa, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 303 = perennis.
virgata, *Mart. Nov. Gen. et Sp.* ii. 16. t. 118.—Bras.
volubilis, Blanco, Fl. Filip. ed. I. 199 = Anredera spicata.
Wallichii, *Hort. Monsp. ex Moq. in DC. Prod.* xiii. II 418, nomen.—Hab. ?

GOMUTUS, Corres, in Ann. Mus. Par. ix. (1807) 268. t. 24. f. 1 = **Arenga**, Labill. (Palm.).
obtusifolius, Blume, Rumphia, ii. 181, in syn. = A. obtusifolia.
Rumphii, Corres, in Ann. Mus. Par. ix. (1807) 288 = A. saccharifera.
saccharifer, Spreng. Syst. ii. 624 = A. saccharifera.

GONATANDRA, Schlecht. in Linnaea, xxiv. (1851) 659 = **Campelia**, Rich. (Commelin.).
tradescantioides, Schlecht. l. c. = C. Zanonia.

GONATANTHUS, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Rar. i. 33. t. 14 (1840). *AROIDEAE*, Benth. & Hook. f. ii. 974.
cuprous, Hort. ex C. Koch, Wochenschr. iv. (1861) 141 = Alocasia cuprea.
Griffithii, Schott, Prod. Aroid. 143 = Steudnera colocasiaefolia.
ornatus, *Schott, in Oestr. Bot. Zeitschr.* (1858) 121.— Reg. Himal.
peltatus, Hort. ex Van Houtte, in Fl. des Serres, xxi (1875) 83 = Steudnera colocasiaefolia.
sarmentosus, *Klotzsch, in Link, Klotzsch & Otto, Ic Pl. Rar.* i. 33.—Reg. Himal.

GONATIA, Nutt. ex DC. Prod. x. 595 (1846) = **Graticola**, Rupp. (Scrophul.).
gratioloides, Nutt. l. c. = Grat. ebracteata.

GONATOCARPUS, Schreb. Gen. i. 86 (1789) = **Haloragis**, Forst.
micranthus, Willd. Sp. Pl. i. 690 = H. micrantha.

GONATOGYNE, Klotzsch, ex Msell. Arg. in Mart. Fl. Bras. xi. II. 18 (1872) = **Savia**, Willd. (Euphorb.).
brasiliensis, Msell. Arg. l. c. 14.—Bras.
lucens, Klotzsch, ex Baill. Adansonia, v. (1865) 345 = prec.

GONATOPUS, Engl. in DC. Monog. Phan. ii. 208 (1879) = **Zamioculcas**, Schott (Aroid.).
Boivini, Engl. l. c. 209 = Z. Boivini.

GONATOSTEMON, Regel, Gartenfl. xv. (1866) 259 = **Chirita**, Buch.-Ham. (Gesneric.).
Bowringianus, Regel, l. c. = C. urticaefolia.

GONEMA, Rafin. Syiva Tellur. 96 (1838) = **Cuneos**, DC. (Melastom.).
divaricata, Rafin. l. c. = Asplectrum divaricatum.
scabrosa, Rafin. l. c. = O. scabrosa.

GONGORA, Ruiz & Pav. Prod. 117. t. 25 (1794). *ORCHIDEAE*, Benth. & Hook. f. iii. 549.
ACROPERA, Lindl. Gen. et Sp. Orch. 179 (1833).
ármenicos, *Reichb. f. Xenia Orch.* i. 62.—Nicaragua.
aromatica, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 281.—Nicaragua.
stropurpurea, *Hook. Exot. Fl.* t. 178.—Ins. Trinit.
Boothiana, *Hort. ex Reichb. f. Xenia Orch.* i. 54 = odoratissima.
bufonis, *Lindl. Bot. Reg.* (1841) t. 2.—Bras.
bufonia, Lindl. l. c. (1847) t. 17, partim = quinquenervis.
cassides, *Reichb. f. in Bot. Zeit.* xxii. (1864) 298.—Guatemala.
Charontia, *Reichb. f. in Linnaea*, xli. (1877) 110.—N. Granat.
Donchelaeriana, Walp. Ann. iii. 546 = truncata.
fulva, Lindl. Bot. Reg. (1839) t. 51 = quinquenervis.
galeata, *Reichb. f. Xenia Orch.* i. 51.—Mexic.
Galeottiana, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 25.—Mexic.
gratulabunda, *Reichb. f. in Bot. Zeit.* xv. (1857) 157.—N. Granat.
grossa, *Reichb. f. in Gard. Chron.* (1877) t. 781.—Ecuador.
Heisterii, Hort. ex Reichb. f. Xenia Orch. i. 54 = atropurpurea.
histrionica, *Reichb. f. in Bonplandia*, ii. (1851) 92 — Hab. ?
irrorata, Hoffmgg. Verz. Orch. (1842) 53.—Cf. Bot. Zeit. i (1843) 833 = bufonia.
Jenischii, Hort. ex Reichb. f. Xenia Orch. i. 54 = odoratissima.
leucochila, Lem. in Fl. des Serres, Sér. I. i. (1845) 87. t. 87 = quinquenervia.
macrantha, Hook. Bot. Misc. ii. (1831) 151 = Coryanthes macrantha.
maculata, Lindl. Bot. Reg. t. 1616 = quinquenervis.
minax, *Reichb. f. in Bot. Zeit.* xxiv. (1866) 385.—Am. austr.
nigrita, *Lindl. in Bot. Reg.* (1839) *Misc.* 59.—Guian.
odoratissima, *Lem. in Fl. des Serres*, Sér. I. ii. (1847) 229.—N. Granat.
philippica, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 498 = Renanthera coccinea ?
pleiochroma, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 421.—N. Granat. ?
portentosa, *Linden & Reichb. f. in Gard. Chron.* (1860) 892.—N. Granat. ?
quadricornis, Hort. ex Reichb. f. Xenia Orch. i. 54 = odoratissima.
quinquenervis, *Ruiz & Pav. Syst. Veg.* 227.—Peruv.
retrorsa, *Reichb. f. in Bonplandia*, ii. (1854) 19.—Venezuela.
saccata, Reichb. f. Xenia Orch. i. 52 = Cirrhaea saccata.
Scaphephorus, *Reichb. f. & Warscz. in Bonplandia*, ii. (1854) 98.—Peruv.
Seideliana, *Reichb. f. in Bot. Zeit.* x. (1852) 637.—Hab. ?
Shepherdii, Hort. ex Reichb. f. Xenia Orch. i. 54 = odoratissima.
similis, *Reichb. f. in Gard. Chron.* (1861) ii. 812.—N. Granat. ?
speciosa, Hook. Bot. Mag. t. 2755 = Coryanthes speciosa.
stenoglossa, *Reichb. f. in Bonplandia*, ii. (1854) 93.—Hab. ?
superdua, *Reichb. f. Xenia Orch.* ii. 189.—Ecuador.
tricolor, *Reichb. f. in Bonplandia*, ii. (1854) 93.—Peruv.
truncata, *Lindl. Bot. Reg.* (1848) *Misc.* 42.—Mexic.
viridiflorapurpurea, Hook. in Bot. Mag. t. 2978 = Cirrhaea Loddigesii.
vitellina, Hort. ex Reichb. f. Xenia Orch. i. 54 = odoratissima.

GONGRODISCUS, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 502, 507. *SAPINDACEAE*.
parvifolius, *Radlk. l. c.* 507.—N. Caled.
sufferrugineus, *Radlk. l. c.*—N. Caled.

GONGROMEMA, Decne. in DC. Prod. viii. 624 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 770.
stramosum, *Decne. l. c.* 625.—Ind or.
Sulawesiana, Decne. l. c. 624 = nepalense.
Thwaitesii, *Decne. l. c.* 625.—Siam.
latifolium, *Benth. in Hook. Niger Fl.* 456.—Aft trop.

GONGRONEMA :—
micradenia, *Benth. & Hook. f. ex F. Muell. Census*, 94.—Austral.
nepalense, *Decne. in DC. Prod.* viii. 624.—Reg. Himal.
recurvifolium, *Walp. Ann.* iii. 62.—N. Guinea.
sagittatum, Decne. in DC. Prod. viii. 624 = nepalense.
ventricosum, *Hook. f. Fl. Brit. Ind.* iv. 34.—Reg. Himal.
Wallichii, *Decne. in DC. Prod.* viii. 624.—Malaya.

GONGROTHAMNUS, Steetz, in Peters, Reise Mossamb. Bot. 336 (1862). *COMPOSITAE*, Benth. & Hook. f. ii. 487.
divaricatus, *Steetz, l. c.* 342.—Afr. trop.
Hildebrandtii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 401.—Afr. trop.
soldaguifolius, *Oliver & Hiern, l. c.*—Afr. trop.

GONGYLOCARPUS, Cham. & Schlecht. in Linnaea, v. (1830) 557. *ONAGRARIEAE*, Benth. & Hook. f. i. 798.
rubricaulis, *Cham. & Schlecht. l. c.* 556.—Mexic.

GONGYLOLEPIS, Schomb. in Linnaea, xx. (1847) 759 = **Stiftia**, Mikan (Compos.).
Benthamiana, Schomb. l. c. 760 = S. Benthamiana.

GONIANTHES, Blume, Cat. Gew. Buitenz. 19 (1823) = **Burmannia**, Linn.
candida, Blume, l. c. 20.—Malaya.
nepalensis, Miers, in Trans. Linn. Soc. xviii. (1841) 537. t. 38.—Reg. Himal. ; China.
pusilla, Miers, l. c. = B. pusilla.
Wallichii, Miers, l. c.—China.

GONIANTHES, A. Rich. in Fl. Cub. Fanerog. ii. 10 (1858). *RUBIACEAE*.
Liadeniana, *A. Rich. l. c.*—Cuba.
Sagraeana, *A. Rich. l. c.* 11.—Cuba.

GONIATICUM, Stokes, Bot. Mat. Med. ii. 338 (1812) = **Polygonum**, Linn.
solitarium, Stokes, l. c. = P. orientale.

GONIOCARPUS, Kon. in Kon. & Sims, Ann. Bot. i. (1805) 546 = **Haloragis**, Forst.
citriodorus, A. Cunn. in Ann. Nat. Hist. iii. (1839) 80 = H. micrantha.
cordiger, Endl. ex Nees, in Lehm. Pl. Preiss. ii. 225 = H. cordigera.
depressus, A. Cunn. in Ann. Nat. Hist. iii. (1839) 80 = H. depressa.
micranthus, Kon. in Kon. & Sims, Ann. Bot. i. (1805) 546 = H. micrantha.
microcarpus, Thibaud, ex DC. Prod. iii. 66 = H. micrantha.
mucronatus, Nees, in Lehm. Pl. Preiss. ii. 225 = H. macronata.
nodulosus, Nees, l. c. i. 158 = H. nodulosa.
pithyoides, Nees, l. c. ii. 225 = H. pithyoides.
rubricaulis, Griff. Notul. iv. 688 = H. macrantha.
salsoloides, Reichb. ex Benth. Fl. Austral. ii. 485 = H. salsoloides.
scaber, Kon. & Sims, Ann. Bot. i. (1805) 547. t. 12. f. 6 = H. tetragyna.
serpyllifolius, Hook. f. in Hook. Ic. Pl. t. 290, 311 = H. depressa.
tenuifus, DC. Prod. iii. 66 = H. tetragyna.
tetragynus, Labill. Nov. Holl. Pl. t. 53 = H. tetragyna.
truncicolus, DC. Prod. iii. 66 = H. teucrioides.
trifidus, Nees, in Lehm. Pl. Preiss. i. 159 = H. trifida.
veruticosus, Hook. f. in Hook. Ic. Pl. tt. 290, 311 = H. depressa.

GONIOCAULON, Cass. in Bull. Soc. Philom. (1817) 84. *COMPOSITAE*, Benth. & Hook. f. ii. 473.
glabrum, *Cass. l. c. ; et in Dict. Sc. Nat.* xiv. 201.—Ind. or.
indicum, C. B. Clarke, Comp. Ind. 286 = glabrum.

GONIOCHETON, Blume, Bijdr. 176 (1825) = **Dysoxylum**, Blume (Meliac.).
arborescens, Blume, l. c. 177 = D. arborescens.

GONIOCHITON, Reichb. Handb. 313 (1837) = prec.

GONIODIUM, Kunze, ex Reichb. Consp. 82 (1828), nomen (Thymel. ?).

GONIOGYNA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 91 = **Heylandia**, DC. (Lagunin.).
hebecarpa, DC. l. c. 92 = H. latebrosa.
latebrosa, DC. l. c. = H. latebrosa.
leiocarpa, DC. l. c. = H. latebrosa.

GONIOGYNE, Benth. & Hook. f. Gen. i. 479 (1865) = prec.

GONIOMA, E. Mey. Comm. Pl. Afr. Austr. 188 (1837). *APOCYNACEAE*, Benth. & Hook. f. ii. 703.
Kamassi, *E. Mey. l. c.*—Afr. austr.

GONIOPOGON, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) t. 173 = **Calotia**, R. Br. (Compos.).
multicaule, Turcz. l. c. 174 = C. plumulifera.

GONIOSCHETON, G. Don, Gen. Syst. i. 668 (1831) = GONIOCHETON, Blume = **Dysoxylum**, Blume (Meliac.).

GONIOSCYPHA, Baker, in Journ. Linn. Soc. xiv. (1875) 581. t. 19. *LILIACEAE*, Benth. & Hook. f. iii. 778.
eucomoides, *Baker, l. c.*—Reg. Himal.

GONIOSTEMMA, Wight, Contrib. 62 (1834) = **Toxocarpus**, Wight & Arn. (Asclep.).
acuminata, Wight, l. c. = T. acuminatus.

GONIOLIMON, Boiss. in DC. Prod. xii. 632 (1848) = **Statice**, Tourn. (Plumbag.).
Becherianum, Janka, in Termész. Füzetek, vi. (1882) 169 = S. tataries !
Besserianum, Nym. Consp. 613 = S. tatarica.
callicomum, Boiss. in DC. Prod. xii. 633 = S. callicoma.
collinum, Boiss. l. c. = S. collina.
dalmaticum, Reichb. Ic. Fl. Germ. xvii. 61. t. 149 = S. dalmatica.
elatum, Boiss. in DC. Prod. xii. 634 = S. elata.
eximium, Boiss. l. c. = S. eximia.
graminifolium, Boiss. l. c. 633 = S. graminifolia.
orthocladum, Rupr. Sert. Tianach. 69.—Turkestan.
Sartorii, Boiss. Diagn. Ser. II. iv. 87 = S. Sartorii.
serbicum, Vis. in Mem. Ist. Venet. x. (1861) 440 = S. serbica.
Szowszowitzii, Herd. in Bull. Soc. Nat. Mosc. xli. (1868) t. 396 = S. Seweszowi.
speciosum, Boiss. in DC. Prod. xii. 634 = S. speciosa.
tataricum, Boiss. l. c. 632 = S. tatarica.

GONIOTHALAMUS, Hook. f. & Thoms. Fl. Ind. i. 105 (1855). *ANONACEAE*, Benth. & Hook. f. i. 26.
arucnala, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 26.—Malaya.
brevicuspis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 33.—Ins. Celebes.
caloncurus, *Miq. l. c.* 34.—N. Guinea.
cardiopetalus, *Hook. f. & Thoms. Fl. Ind.* 108.—Ind. or.
ceramensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 33.—Ins. Ceram.
costularis, *Miq. Fl. Ind. Bat. Suppl.* 372.—Sumatra.
disperrus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 34.—Borneo.
eunerus, *Miq. l. c.* 33.—Ins. Celebes.
fulvus, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 78.—Malacca.
Gardneri, *Hook. f. & Thoms. Fl. Ind.* i. 107.—Zeylan.
giganteus, *Hook. f. & Thoms. l. c.* 109.—Malaya.
Griffithii, *Hook. f. & Thoms. l. c.* 110.—Burma.
Hookeri, *Thw Enum. Pl. Zeyl.* 6.—Zeylan.
imbricatus, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 27.—N. Guinea.
lanceolatus, *Miq. Fl. Ind. Bat. Suppl.* 373.—Sumatra.
longitrorsis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 4.—N. Guinea.
macrophyllus, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 74.—Malacca.
macrophyllus, *Miq. Fl. Ind. Bat.* i. 11. 28.—Java.
malayanus, *Hook. f. & Thoms. Fl. Ind.* i. 107.—Malaya.
oxycarpus, Miq. Fl. Ind. Bat. Suppl. 372 (= *Guatteria oxycarpa*).—Sumatra.
reticulatus, *Thw. Enum. Pl. Zeyl.* 7, 298.—Zeylan.

GONIOTHALAMUS :—
rafus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 35.—Borneo.
salicinus, *Hook. f. & Thoms. Fl. Ind.* i. 106.—Zeylan.
sesquipedalis, *Hook. f. & Thoms. l. c.* 108.—Reg. Himal. ; Burma.
Simondi, *Hook. f. & Thoms. l. c.*—Reg. Himal.
Slingerlandtianus, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 341.—Malaya.
sumatranus, *Miq. Fl. Ind. Bat. Suppl.* 371.—Sumatra.
Tapis, *Miq. l. c.*—Sumatra.
Thomsoni, *Thw. Enum. Pl. Zeyl.* 7.—Zeylan.
Thwaitesii, *Hook. f. & Thoms. Fl. Ind.* i. 106.—Zeylan.
Walkeri, *Hook. f. & Thoms. l. c.* 109.—Zeylan.
Wightii, *Hook. f. & Thoms. l. c.* 106.—Ind. or.
wynaadensis, *Bedd. Ic. Pl. Ind. Or.* i. 13. t. 61.—Ind. or.

GONIOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 181 = **Trichinium**, R. Br. (Amarant.).

GONIPIA, Rafin. Fl. Tellur. iii. 23 (1836) = **Gentiana**, Tourn.
bicolor, Rafin. l. c. 24.—Hab. ?
linearis, Rafin. l. c. = Gent. linearis.
paucifolia, Rafin. l. c. 23.—Hab. ?
pulchella, Rafin. l. c. 24 = Erythraea ramosissima.
pumila, Rafin. l. c. = Gent. pumila.
rotundifolia, Rafin. l. c. = Gent. bavarica.
ricula, Rafin. l. c. 23 (Quid ?).—Steil.
ternifiora, Rafin. l. c. 24 = Gent. Amarella ?

GONISTUM, Rafin. Sylva Tellur. 85 (1838) = **Piper**, Linn.
unguiculatum, Rafin. l. c. = P. unguiculatum.

GONISTYLUS, Baill. Hist. Pl. vi. 128 (1875) = **Gonystylus**, Teijsm. & Binn. (Thymel.).

GONIURUS, Presl, Epim. Bot. 244 (1849) = **Pothos**, Linn. (Aroid.).
luzonensis, Presl, l. c. = P. luzonensis.

GONOCALYX, Planch. & Linden, ex Gard. Chron. (1856) 152 = **Ceratostemma** ?, Juss. (Vaccinac.).
pulcher, Planch. & Linden, l. c. ; et in Belg. Hortic. vi. (1856) 333.—N. Granat.

GONOCARPUS, Ham. Prod. Pl. Ind. Occ. 89 (1825) = **Cornbretum**, Linn.
Jacquini, Ham. l. c. = C. alternifolium.

GONOCARPUS, Thunb, Nov. Gen. 55 (1783) = **Haloragis**, Forst.
micranthus, Thunb. l. c. = H. micrantha.

GONOCARYUM, Miq. Fl. Ind. Bat. Suppl. 843 (1860). *INC. SED.*—Cf. Benth. & Hook. f. i. 845.
affine, *Becc. Malesia*, i. (1878) 256.—N. Guinea.
gracile, *Miq. Fl. Ind. Bat. Suppl.* 843.—Sumatra.
Griffithianum, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 155.—Burma.
macrocarpum, *Scheff. in Ann. Jard. Buitens.* i. (1876) 102, in obs.—Ins. Lepar.
pyriforme, *Scheff. l. c.* 100.—Ins. Molucc.
selebicum, *Becc. Malesia*, i. (1877) 194.—Ins. Celebes.
Teysmanniannum, *Scheff. in Ann. Jard. Buitens.* i. (1876) 98.—Malaya.

GONOCITRUS, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 228. *RUTACEAE.*
angulata, *Kurz*, l. c.—Ind. or.

GONOCYTISUS, Spach, in Ann. Sc. Nat. Sér. III. iii. (1845) 158 = Genista, Linn. (Legumin.).
angulatus, Spach, l. c. 158, 154 = Gen. parviflora.
pterocladus, Spach, l. c. 154 = Gen. pterocalda.

GONOGONA, Link, Enum. Hort. Berol. ii. 369 (1822) = **Goodyera**, R. Br. (Orchid.).
discolor, Link, l. c. = Good. discolor.
repens, Link, l. c. = Good. repens.

GONOHORIA, G. Don, Gen. Syst. i. 840 (1831), sphalm. = CONOHORIA, Aubl. = **Alsodeia**, Thou. (Violar.).

GONOKEROS, Rafin. Fl. Tellur. iv. 96 (1836) = **Seshima**, Tourn. (Dipsac.).
hungaricus, Rafin. l. c. (Quid ?).—Hungar.

GONOLOBIUM, Hedw. f. Gen. 182 (1806) = **Gonolobus**, Michx. (Asclep.).

GONOLOBUS, Michx. Fl. Bor. Am. i. 119 (1803). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 766.
CETHAMALIA, Decne. in DC. Prod. viii. 605 (1844).
albus, Poir. Encyc. Suppl. ii. 815 = Sarcostemma Dombeyanum.
altissimus, *Schult. Syst.* vi. 55.—N. Granat.
aristolochioides, *H. B. & K. Nov. Gen. et Sp.* iii. 208.—Venezuela.
asper, *Decne. in DC. Prod.* viii. 595.—Mexic.
Baldwynianus, *Sweet. Hort. Brit.* ed. II. 860.—Am. bor.
barbatus, *H. B. & K. Nov. Gen. et Sp.* iii. 209. t. 239.—Mexic.
bifidus, *Hemsl. Biol. Centr. Am. Bot.* ii. 330.—Mexic.
biflorus, *Nutt. ex Torr. Bot. Mex. Bound.* 165.—Am. bor. occ.
biflorus, *Rafin. New Fl. Am.* iv. 58.—Am. bor.
Billbergianus, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 138.—Panama.
carolinensis, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—Am. bor.
carolinensis, Nutt. Gen. Am. i. 168 = Baldwynianus.
chloranthus, *Schlecht. in Linnaea*, viii. (1833) 520.—Mexic.
cinctus, *Benth. & Hook. f. Gen.* ii. 750, in nota sub Fischeria.—Jamaic.
congestus, *Decne. in DC. Prod.* viii. 597.—Mexic.
crispiflorus, R. Br. in Mem. Wern. Soc. i. (1809) 35 = Fischeria scandens.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 98.—Cuba.
Cundurango, Triana, in Comp. Rend. Par. lxxiv. (1872) 853 = Maradenia Cundurango.
cyanchoides, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 351.—Am. bor. occ.
diadematus, *Ker-Gawl. Bot. Reg.* t. 252.—Mexic
discolor, Schult. Syst. vi. 63 = obliquus.
edulis, *Hemsl. in Biol. Centr. Am. Bot.* ii. 331.—Guatemala.
erianthus, *Decne. in DC. Prod.* viii. 593.—Mexic.
erioclados, *Benth. Pl. Hartw.* 216.—N. Granat.
erioperalus, Moric. ex Decne. in DC. Prod. viii. 541 = Sarcostemma elegans.
filiformis, Schult. Syst. vi. 65 = Sarcostemma Jacquini.
floridanus, Chapm. in Coulter, Bot. Gas. iii. (1878) 12 = hirsutus.
flocculosus, *Bertol. in Opuc. Scient.* iv. (1823) 225.—Ins. Guadaloup.
foetidus, *Griseb. in Gaett. Abh.* xix. (1874) 206.—Reg. Argent.
fraternus, *Schlecht. in Linnaea*, viii. (1833) 521.—Mexic.
fuscus, *Decne. in DC. Prod.* viii. 592.—Mexic.
ganglinosus, *Decne. l. c.* 597.—Bras.
gemiuiflorus, *Decne. l. c.* 595.—Bras.
glabet, *Decne. l. c.* 594.—Guian.
glaberrimus, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 955, nomen.—Guian.
glaucifolius, *Poepp. ex Decne. in DC. Prod.* viii. 598.—Peruv.
Glaziovii, *Fourn. in Mart. Fl. Bras.* vi. iv. 315.—Bras.
gracilis, *Decne. in DC. Prod.* viii. 596.—Mexic.
grandiflorus, Benth. Pl. Hartw. 24 = erianthus.
grandiflorus, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—Am. septr.
grandiflorus, Lindl. Bot. Reg. t. 1053 = chloranthus.
granulatus, Scheele, in Linnaea, xxi. (1848) 759 = laevis.
granulatus, Torr. Bot. Mex. Bound. 165 = reticulatus.
guianensis, Spreng. Syst. i. 845 = viridiflorus.
hastulatus, *A. Gray, in Proc. Am. Acad.* xii. (1877) 76.—Calif.
hirsutus, Lodd. Bot. Cab. t. 965 = Baldwynianus ?
hirsutus, *Michx. Fl. Bor. Am.* i. 119.—Am. bor.
hirsutus, Pursh, Fl. Am. Sept. i. 176 ; Nutt. Gen. Am. i. 168 = obliquus.
hirtus, R. Br. in Mem. Wern. Soc. i. (1809) 35 = obliquus.
hirtus, *Griseb. in Gaett. Abh.* xxiv. (1879) 284.—Reg. Argent.

GONOLOBUS :—
hispidus, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 298.—Am. austr.
Hystrix, *Decne. in DC. Prod.* viii. 593.—Bras.
laevis, *Michx. Fl. Bor. Am.* i. 119.—Am. bor.
lanatus, *Griseb. in Goett. Abh.* xxiv. (1879) 285.—Reg. Argent.
lanceolatus, *Decne. in DC. Prod.* viii. 598.—Mexic.
laslostomus, *Decne. l. c.* 593.—N. Granat.
ligustrinus, *Decne. l. c.* 596.—Bras.
littoralis, *Decne. l. c.* 596.—Mexic.
luridus, *Decne. l. c.* 592.—Mexic.
luteolus, *Decne. l. c.* 593.—Mexic.
macranthus, *Kunze, in Linnaea*, xx. (1847) 27.—Mexic.
macrocarpus, *Gaspary. in Ann. Cir.* iii. (1819) 122.—Hab. ?
macrophyllus, Chapm. Fl. S. U. St. i. 368 = suberosus.
macrophyllus, Decne. in DC. Prod. viii. 596 = obliquus
macrophyllus, Ell. Sketch. i. 327 = hirsutus.
macrophyllus, Michx. Fl. Bor. Am. i. 119 = laevis.
maritimus, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—N. Granat.
Martianus, Hook. Bot. Mag. t. 4472 = Fischeria Martiana.
martinicensis, *Decne. in DC. Prod.* viii. 595.—Ind. occ.
micranthus, *Hook. f. ex Hemsl. Bot. Voy. Challenger*, i. 11. 15.—Ins. Fernando-Noronha.
mollis, *Regel, Ind. Sem. Hort. Petrop.* (1856) 26 ; *et in Gartenfl.* (1857) 262.—Bras.
mucronatus, Sweet, Hort. Brit. ed. II. 860 = Tylophora asthmatica ?
nemorosus, *Decne. in DC. Prod.* viii. 596.—Mexic.
niger, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—Mexic.
nigrescens, *Schlecht. in Linnaea*, viii. (1833) 522.—Mexic.
nummularius, *Hemsl. Biol. Centr. Am. Bot.* ii. 332.—Mexic.
Nuttallianus, Spreng. Syst. i. 845 = laevis.
Nuttallii, Decne. in DC. Prod. viii. 598 = laevis.
obliquifolius, Colla, in Mem. Acc. Torin. xxxviii. (1835) 128. t. 87 = Oxypetalum saxatile.
obliquus, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—Am. bor.
obtusiflorus, *Decne. in DC. Prod.* viii. 594.—Bras.
obtusifolius, *Schult. Syst.* vi. 66.—Hab. ?
orthosioides, *Fourn. in Mart. Fl. Bras.* vi. iv. 315.—Bras.
Ottonis, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1855) 13 ; *ex Walp. Ann.* v. 502.—Cuba.
oxyanthus, *Torrey, in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 318.—Venezuela.
parviflorus, *Decne. in DC. Prod.* viii. 597.—Peruv.
parviflorus, *A. Gray, in Proc. Am. Acad.* xii. (1877) 79.—Am. bor. occ.
parvifolius, *Torr. in Bot. Mex. Bound.* 166.—Am. bor. occ.
petens, *Decne. in DC. Prod.* viii. 594.—Bras.
pauciflorus, *Spreng. Syst.* i. 846.—Ins. S. Doming.
pedunculatus, *Hemsl. Biol. Centr. Am. Bot.* ii. 332.—Mexic.
picturatus, *Hemsl. l. c.*—Guatemala.
pilosus, *Benth. Pl. Hartw.* 289.—Mexic.
planiflorus, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—N. Granat.
pogonanthus, Hemsl. Biol. Centr. Am. Bot. ii. 333, pro parte = Pherotrichis villosa.
productus, *Torr. in Bot. Mex. Bound.* 165.—Am. bor. occ.
prostratus, Baldw. in Ell. Sketch. i. 329 = pubiflorus.
prostratus, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—Mexic.
pubescens, *Griseb. Fl. Brit. W. Ind.* 420.—Jamaic.
pubiflorus, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 352, in textu.—Am. bor. occ.
pyrrhotrichus, *Decne. in DC. Prod.* viii. 594.—Bras.
racemosus, R. Br. in Mem. Wern. Soc. i. (1809) 35 = Roulinia Jacquini.
reflexus, *Hemsl. Biol. Centr. Am. Bot.* ii. 332.—Nicaragua.
reticulatus, *Engelm. ex A. Gray, in Proc. Am. Acad.* xii. (1877) 78.—Am. bor. occ.
retusus, *Decne. in DC. Prod.* viii. 594.—Bras.
rhamnifolius, *Griseb. Fl. Brit. W. Ind.* 420.—Jamaic.
riparius, *H. B. & K. Nov. Gen. et Sp.* iii. 208.—N. Granat.
rostratus, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—Ind. occ.

GONOLOBUS:—
rotatus, *Decne. in DC. Prod.* viii. 593.—Bras.
sagittifolius, *A. Gray, in Proc. Am. Acad.* xii. (1877) 77.—Texas.
Salvinii, *Hemsl. Biol. Centr. Am. Bot.* ii. 334.—Guatemala.
Schaffneri, *A. Gray, ex Hemsl. l. c.*—Mexic.
setosus, *Benth. Pl. Hartw.* 216.—N. Granat.
Shortii, *A. Gray, in Coult. Bot. Gaz.* viii. (1883) 191. —Am. bor.
sidaefolius, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 367.—Mexic.
stellatus, *Griseb. Fl. Brit. W. Ind.* 420.—Jamaica.
stelliflorus, *Fourn. in Mart. Fl. Bras.* vi. iv. 317.—Bras.
stenolobus, Decne. in DC. Prod. viii. 595 = Exolobus stenolobus.
stephanotrichus, *Griseb. Cat. Pl. Cub.* 177.—Cuba.
striatus, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 365.—Mexic.
suberosus, Bertol. Elench. 5 = floccosus.
suberosus, *R. Br. in Mem. Wern. Soc.* i. (1809) 35.—Am. bor.
tetragonus, *Decne. in DC. Prod.* viii. 594.—Bras.
tigrinus, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 520.—Cuba.
tiliaefolius, Decne. in DC. Prod. viii. 596 = laevis.
tingens, *Decne. l. c.* 592.—Mexic.
trifloras, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1 (1844) 365.—Mexic.
tristis, *Decne. in DC. Prod.* viii. 596.—Mexic.
undulatus, *R. Br. in Mem. Wern. Soc.* i (1809) 35.— N. Granat.
semiverius, Scheele, in Linnaea, xxi. (1848) 760 = Roulinia unifaria.
uniflorus, *H. B. & K. Nov. Gen. et Sp.* iii. 207. t. 288. —Mexic.
velutinus, *Schlecht. in Linnaea*, viii. (1833) 521.— Mexic.
viresceus, *Decne. in DC. Prod.* viii 596.—Mexic.
viridiflorus, Natt. Cat. Am. i. 163 = laevis.
viridiflorus, *Schult. Syst.* vi. 61.—Bras.
viridiflorus, Torr. in Ann. Lyc. N. York, ii. (1828) 219 = Philibertia cynanchoides.
viridis, Targ. in Mem. Mod. xx. (1829) 295 = viridiflorus.
Voquicilla, Colla, in Mem. Acc. Torin. xxxviii. (1835) 134. t. 86 = Oxypetalum saxatile.

GONOLOMA, Rafin. Sylva Tellur. 86 (1838) = Vitis, Tourn.
alata, Rafin. l. c. = V. rhombifolia.

GONONCUS, Rafin. Fl. Tellur. iii. 16 (1836) = Polygonum, Tourn.
undulatus, Rafin. l. c. = P. alpinum.

GONOPTERA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 1. 160 = Bulnesia, C. Gay (Zygophyll.).
chilensis, Turcz. l. c. 161 = B. chilensis.

GONOPYRUM, Fisch. & Mey. ex C. A. Mey. in Mém. Acad. Pétersb. Sér. VI. vi. (1840) 144 = Polygonella, Michx. (Polygon.).
americanum, Fisch. & Mey. l. c. = P. ericoides.

GONOSPERMUM, Less. Syn. Comp. 263 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 417.
HYMENOLEPIS, Cass. in Bull. Soc. Philom. (1817) 186, partim.
LUGOA, DC. Prod. vi. 14 (1837).
canariense, *Less. Syn. Comp.* 263.—Ins. Canar.
elegans, DC. Prod. vi. 34 = canariense.
fruticosum, DC. l. c. = canariense.
fruticosum, *Less. Syn. Comp.* 263.—Ins. Canar.
Gomarse, *C. Bolle, in Bonplandia*, vii. (1859) 296.— Ins. Canar.
multiflorum, DC. Prod. vi. 34, excl. syn. = fruticosum.
revolutum, *Rich. Bip. in Webb & Berth. Phyt. Canar.* ii. 292. t. 88.—Ins. Canar.

GONOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) ii. 309 = Pouzolzia, Gaudich. (Urticac.).
alternifolia, Turcz. l. c. 310.—Ins. Luzon.
ambigua, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 208 = P. Wightii.
aquatica, Miq. l. c. = P. pentandra.
brachystachya, Miq. l. c. = P. candata.

GONOSTEGIA :—
candata, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 208 = Pouzolzia candata.
concinna, Miq. l. c. = P. Wightii.
gracilis, Miq. l. c.—Ind. or.
hirta, Miq. l. c. = P. hirta.
integrifolia, Miq. l. c. = P. integrifolia.
nilghirensis, Miq. l. c. = P. Wightii.
oppositifolia, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) ii. 310 = P. pentandra.
parvifolia, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 208 = P. parvifolia.
pentandra, Miq. l. c. 209 = P. pentandra.
quinquenervia, Miq. l. c. 208 = P. hirta.
scabra, Miq. l. c. = P. scabra.
triandra, Miq. l. c. = P. parvifolia.

GONOSTEMMA, Spreng. Gen. i. 208 (1830) = seq.

GONOSTEMON, Haw. Syn. Pl. Succ. 27 (1812) = Stapelia, Linn. (Asclep.).
divaricatus, Haw. l. c. = S. divaricata.
pallidus, G. Don, Gen. Syst. iv. 117 = S. pallida.
strictus, Haw. Suppl. Pl. Succ. 12 = S. stricta.

GONOSUKE, Rafin. Sylva Tellur. 58 (1838) = Ficus, Tourn. (Urtic.).
demonum, hispida, scaber, Rafin. l. c. = F. hispida.

GONOTHECA, Blume, ex DC. Prod. iv. 429 (1830) = Oldenlandia, Linn. (Rubiac.).
Blumei, DC. l. c. = O. alata.

GONOTHECA, Rafin. in Med. Repos. N. York, v. (1808) 352 = Tetragonotheca, Dill. (Compos.).
helianthoides, Rafin. l. c. = T. helianthoides.

GONTARELLA, Gillib. ex Steud. Nom. ed. I. 380 (1821) = Isopyrum, Linn. (Ranunc.).
tertiaria, Gillib. ex Steud. l. c. = I. thalictroides.

GONUFAS, Rafin. Sylva Tellur. 124 (1838) = Celosia, Linn. (Amarant.).
paniculata, Rafin. l. c. = C. paniculata.

GONUS, Lour. Fl. Cochinch. 658 (1790) = Brucea, J. S. Mill. (Simarub.).
amarissa, Lour. l. c. = B. sumatrana.

GONYANERA, Korth. in Nederl. Kruidk. Arch. ii. 17. (1851) 188. *RUBIACEAE*, Benth. & Hook. f. ii 715.
glauca, *Korth. l. c.*—Sumatra.

GONYANTHES, Nees, in Ann. Sc. Nat. Sér. I. iii. (1834) 369 = GONIANTHES, Blume = Burmannia, Linn.

GONYCLISIA, Dulac, Fl. Hautes-Pyr. 191 (1867) = Cochlearia, Linn. (Crucif.).
saxatilis, Dulac, l. c. = C. saxatilis.

GONYSTYLUS, Teijsm. & Binn. in Bot. Zeit. xx. (1862) 265. *THYMELAEACEAE*, Benth. & Hook. f. iii. 201.
bancanus, Baill Hist. Pl. vi. 128 = Miquelianus.
Miquelianus, *Teijsm. & Binn. in Bot. Zeit.* xx. (1862) 265.—Malays.

GONZALAGUNEA, Reichb. Nom. 77 (1841) = seq.

GONZALAGUNIA, Ruiz & Pav. Prod. 12 (1794) = Gonsalea, Pers. (Rubiac.).
Cacocoypselum, Cham. & Schlecht. in Linnaea, iv. (1829) 196 = Gonzalea Cacocoypselum.
dependens, Ruiz & Pav. Fl. Per. i. 56 = Gonzalea pendula.
dicocca, Cham. & Schlecht. in Linnaea, iv. (1829) 194 = Gonzalea dicocca.
parviflora, Schlecht. & Cham. in Linnaea, vi. (1831) 414 = Gonzalea parviflora.

GONZALEA, Pers. Syn. i. 182 (1805). *RUBIACEAE*, Benth. & Hook. f. ii. 25.
BUENA, Cav. in Anal. Hist. Nat. ii. (1800) 278; Ic. vi. 50. t. 571 (1801).
DUGOMNA, Vahl, in West, Bidr. Ste-Croix, 269 (1793) nomen prius.
GONZALAGUNIA, Ruiz & Pav. Prod. 12. t. 3 (1794).
brachyantha, *A. Rich. Fl. Cub. Fanerog.* ii. 16.—Cuba.
Cacocoypselum, *Steud. Nom.* ed. II. i. 701.—Porto Rico.

GONZALEA :—
cornifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 416.— N. Granat.
dicocca, *Steud. Nom.* ed. II. i. 701.—Bras.
incanescens, *DC. Prod.* iv. 437.—Ins. Tobag.
leptantha, *A. Rich. Fl. Cub. Fanerog.* ii. 16.—Cuba.
Lygistum, Schult. Mant. iii. 133 = Mancetta Lygistum.
niven, *Bartl. ex DC. Prod.* iv. 436.—Mexic.
panamensis, *Pers. Syn.* i. 183.—Panama; Mexic.
panamensis, Spreng. Syst. i. 1. = splenta.
panamensis, Wydler, ex Cham. & Schlecht. in Linnaea, iv. (1829) 196 = G. Cococcypselum.
parviflora, *G. Don, Gen. Syst.* iii. 587.—Mexic.
pendula, *Pers. Syn.* i. 182.—Peruv.
Petesia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 504.—Jamaic.
pulverulenta, *Humb. & Bonpl. Pl. Aequin.* i. 228.— Peruv.
secunda, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 235.—Mexic.
sessiliflora, *Presl, Rel. Bemerk.* 80 = Bras.
spicata, *DC. Prod.* iv. 437.—Am. austr.
tomentosa, *Humb. & Bonpl. Pl. Aequin.* i. 225. t 64. —Peruv.

GOODALLIA, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 233. *THYMELAEACEAE*, Benth. & Hook. f. iii. 194.
guianensis, *Benth. l. c.*—Guian.

GOODENIA, Sm. in Trans. Linn. Soc. ii. (1794) 346 *GOODENOVEAE*, Benth. & Hook. f. ii. 588.
AILLVA, De Vriese, in Verh. Haarl. x. (1854) 75. t. 13.
BOUTONIA, Hort. Erfurt. ex Steud. Nom. ed. II. i. 219 (1840).
COLLEMA, Andrex. ex DC. Prod. vii. 519 (1839).
MONOCHILA, Spach, Hist Vég. Phan. ix. 237 (1840).
PICROPHYTA, F. Muell. in Linnaea, xxv. (1852) 431.
STEKHOVIA, De Vriese, in Verh. Haarl. x. (1854) 166. t. 20.
TETRAPHYLAX, De Vriese, l. c. 164 (1854).
acuminata, R. Br. Prod. 576 = ovata.
affinis, De Vriese, in Verh. Haarl. x. (1854) 137 = geniculata.
albida, Sm. in Trans. Linn. Soc. ii. (1794) 348 = Scaevola microcarpa.
albiflora, *Schlecht. in Linnaea*, xx. (1847) 599.— Austral.
amplexans, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 70.—Austral.
appendiculata, Jacq. Fragm. 62 = grandiflora.
armeriaefolia, De Vriese, in Lehm. Pl. Preiss. i. 412 = Diaspasis filifolia.
armeriaefolia, Sieber, ex DC. Prod. vii. 518 = stelligera.
Armitiana, *F. Muell. Fragm.* x. 110.—Austral.
Armstrongiana, *De Vriese, in Verh. Haarl.* x. (1854) 138. t. 34.—Austral.
arthrotricha, *F. Muell. ex Benth. Fl. Austral.* iv. 62.— Austral.
auriculata, *Benth. l. c.* 72.—Austral.
azurea, *F. Muell. Fragm.* i. 117.—Austral.
barbata, *R. Br. Prod.* 576.—Austral.
Barlisia, F. Muell. Fragm. iii. 140 = caerulea.
bellidifolia, *Sm. in Trans. Linn. Soc.* ii. (1794) 349.— Austral.
Brevardiana, Gandich. in Freyc. Voy. Bot. 460 = Calogyne Berardiana.
bicolor, *F. Muell. ex Benth. Fl. Austral.* iv. 80.— Austral.
Bonnyana, *F. Muell. Fragm.* vi. 236.—Austral.
caerulea, *R. Br. Prod.* 578.—Austral.
calcarata, *F. Muell. Fragm.* vi. 14.—Austral.
calendulacea, Andr. Bot. Rep. t. 23 = Scaevola suaveolens.
Chambersii, *F. Muell. Fragm.* i. 204.—Austral.
ciliifolia, A. Cunn. ex DC. Prod. viii. 516 = barbata.
cleytoniacea, *F. Muell. Census*, 58 (Cf. *Leptonisma*). —Austral.
concinna, *Benth. Fl. Austral.* iv. 76.—Austral.
coronopifolia, *R. Br. Prod.* 576.—Austral.
corynocarpa, *F. Muell. Fragm.* ii. 16.—Austral.
cyanea, F. Muell. l. c. i. 155 = pterigosperma.
cycloptera, *R. Br. in Sturt, Exped. Centr. Austral.* ii. *App.* 53.—Austral.
decurrens, *R. Br. Prod.* 575.—Austral.
disperma, *F. Muell. Fragm.* i. 118.—Austral.

GOODENIA:—
dubia, Spreng. Syst. i. 721 = Calogyne pilosa.
Eatoniana, *F. Muell. Fragm.* viii. 186.—Austral.
elongata, De Vriese, in Lehm. Pl. Preiss. i. 413 = filiformis.
elongata, *Labill. Nov. Holl. Pl.* i. 59. t. 75.—Austral.
enigma, F. Muell. Fragm. iii. 142 = Selliera enigma.
filiformis, *R. Br. Prod.* 578.—Austral.
flagellifera, De Vriese, in Mitch. Journ. Trop. Austral. 378 = glabra.
flexuosa, De Vriese, in Verh. Haarl. x. (1854) 126 = paniculata.
geniculata, *R. Br. Prod.* 577.—Austral.
geniculata, De Vriese, in Lehm. Pl. Preiss. i. 418 = Anthotium humile.
grandiflora, De Vriese, l. c. ii. 244 = Anthotium humile.
glabra, *R. Br. Prod.* 577.—Austral.
glandulifera, De Vriese in Verh. Haarl. x. (1854) 129 = Diaspasis filifolia.
glauca, *F. Muell. in Trans. Vict. Inst.* i. (1855) 40.—Austral.
gracilis, *R. Br. Prod.* 575.—Austral.
graminifolia, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 265 = humilis.
grandiflora, *Sims, Bot. Mag.* t. 890.—Austral.
Hasselilli, *F. Muell. Fragm.* vi. 10. t. 51.—Austral.
hederacea, *Sm. in Trans. Linn. Soc.* ii. (1794) 349.—Austral.
herpystica, *Schlecht. in Linnaea*, xx. (1847) 598.—Austral.
heterochila, *F. Muell. Fragm.* iii. 142.—Austral.
heteromera, *F. Muell. l. c.* i. 115.—Austral.
heterophylla, *Sm. in Trans. Linn. Soc.* ii. (1794) 349.—Austral.
heteroptera, *F. Muell. Fragm.* x. 43.—Austral.
hirsuta, *F. Muell. l. c.* iii. 35.—Austral.
hispida, *R. Br. Prod.* 577.—Austral.
humilis, *R. Br. l. c.* 575.—Austral.
incana, *R. Br. l. c.* 578.—Austral.
intermedia, Gaertn. f. Fruct. iii. 165 = rotundifolia ?
junciformis, De Vriese, in Lehm. Pl. Preiss. i. 418 = Anthotium humile.
laxigata, Curt. Bot. Mag. t. 287 = Scaevola microcarpa.
laevis, *Benth. Fl. Austral.* iv. 61.—Austral.
lamprosperma, *F. Muell. Fragm.* i. 116.—Austral.
lanata, *R. Br. Prod.* 577.—Austral.
lanceolata, Hort. ex Steud. Nom. ed. ii. i. 701 = paniculata.
latifolia, *Hort. ex Steud. l. c.*—Austral.
Layloniana, F. Muell. ex Benth. Fl. Austral. iv. 79, sphalm.—Cf. claytoniacea.
leptoclada, *Benth. Fl. Austral.* iv. 67.—Austral.
leptotheca, F. Muell. Fragm. vi. 13 = xanthotricha.
littoralis, R. Br. in Flind. Voy. ii. 561 = Selliera radicans.
longifolia, De Vriese, in Verh. Haarl. x. (1854) 127 = stelligera.
Macmillanii, *F. Muell. Fragm.* i. 119. t. 5.—Austral.
macrophylla, F. Muell. l. c. vi. 11 = Velleia macrophylla.
margarata, De Vriese, in Verh. Haarl. x. (1854) 148.—Austral.
melanoptera, F. Muell. Fragm. i. 115 = heterochila.
micropters, *F. Muell. l. c.* iii. 34.—Austral.
minutiflora, *F. Muell. l. c.* viii. 244.—Austral.
minutifolia, F. Muell. l. c. ix. 196, sphalm. = praec.
Mitchellii, *Benth. Fl. Austral.* iv. 71.—Austral.
mollis, R. Br. Prod. 577 = grandiflora.
mollissima, *F. Muell. ex Benth. Fl. Austral.* i. 78.—Austral.
Moorckeana, *F. Muell. Fragm.* viii. 56.—Austral.
nana, De Vriese, in Verh. Haarl. x. (1854) 122 = humilis.
Nicolaoni, *F. Muell. Fragm.* i. 208. t. 4.—Austral.
ovata, *Sm. in Trans. Linn. Soc.* ii. (1794) 347.—Austral.
paniculata, *Sm. l. c.* 346.—Austral.
phylicoides, *F. Muell. Fragm.* i. 206.—Austral.
pinifolia, *De Vriese, in Verh. Haarl.* x. (1854) 157. t. 30. f. 1–14.—Austral.
pinnatifida, *Schlecht. in Linnaea*, xxi. (1848) 450.—Austral.
primulacea, Schlecht. l. c. xx. (1847) 601 = geniculata.
pterigosperma, *R. Br. Prod.* 578.—Austral.
pterygosperma, De Vriese, in Verh. Haarl. x. (1854) 133 = caerulea.

GOODENIA:—
pubescens, Sieber, ex DC. Prod. vii. 514 = Scaevola microcarpa.
pulcherlia, Benth. in Enum. Pl. Hueg. 71 = filiformis.
pumila, *R. Br. Prod.* 578.—Austral.
purpurascens, *R. Br. l. c.* 578.—Austral.
pusilla, De Vriese in Verh. Haarl. x. (1854) 131 = tenella.
pygmaea, De Vriese, in Lehm. Pl. Preiss. i. 413 = Anthotium humile.
quadrilocularis, *R. Br. Prod.* 578.—Austral.
racemosa, *F. Muell. Fragm.* i. 114.—Austral.
radicans, Pers. Syn. i. 196 = Selliera radicans.
Ramelli, *F. Muell. Fragm.* iii. 20. t. 17.—Austral.
ramosissima, Sm. in Trans. Linn. Soc. ii. (1794) 349 = Scaevola hispida.
repens, Labill. Nov. Holl. Pl. i. 58. t. 76 = Selliera radicans.
rigida, Benth. in Enum. Pl. Hueg. 71 = caerulea.
rotundifolia, *R. Br. Prod.* 576.—Austral.
scaevolina, *F. Muell. Fragm.* i. 113.—Austral.
scapigera, *R. Br. Prod.* 578.—Austral.
sepalosa, *F. Muell. ex Benth. Fl. Austral.* iv. 72.—Austral.
sessiliflora, F. Muell. Fragm. iv. 145 = disperma.
spathuiata, De Vriese, in Verh. Haarl. x. (1854) 123 = bellidifolia.
spicata, F. Muell. Fragm. iii. 35 = viscida.
squarrosa, De Vriese, in Lehm. Pl. Preiss. i. 413 = Scaevola fascicularis.
stelligera, *R. Br. Prod.* 575.—Austral.
stenophylla, F. Muell. Fragm. i. 113 = Scaevola stenophylla.
Stobbiana, *F. Muell. l. c.* xi. 49.—Austral.
stolonifera, De Vriese, in Verh. Haarl. x. (1854) 135 = tenella ?
Strangfordii, *F. Muell. Fragm.* vi. 11. t. 52.—Austral.
stricta, Sm. in Trans. Linn. Soc. ii. (1794) 349 = Dampiera stricta.
strongylophylla, *F. Muell. Fragm.* vi. 12.—Austral.
strophiolata, *F. Muell. l. c.* i. 119.—Austral.
Taylori, F. Muell. l. c. iii. 141 = quadrilocularis.
tenella, Andr. Bot. Rep. t. 466 = Velleia trinervis.
tenella, *R. Br. Prod.* 577.—Austral.
tenuifolia, F. Muell. Fragm. ii. 111 = claytoniacea.
tenuiloba, *F. Muell. in Wing, South Sc. Record*, (Feb. 1885) 25.—Austral.
teretifolia, De Vriese, in Verh. Haarl. x. (1854) 130 = caerulea.
teucriifolia, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 70 = heterophylla.
trichophylla, *De Vriese, ex Benth. Fl. Austral.* iv. 67.—Austral.
varia, *R. Br. Prod.* 576.—Austral.
Vilmoriniae, *F. Muell. Fragm.* iii. 19. t. 16.—Austral.
viscida, *R. Br. Prod.* 578.—Austral.
xanthosperma, *F. Muell. Fragm.* x. 12.—Austral.
xanthotricha, *De Vriese, in Verh. Haarl.* x. (1854) 155.—Austral.

GOODIA, Salisb. Parad. Lond. t. 41 (1806). *LEGUMINOSAE*. Benth. & Hook. f. i. 474.
lotifolia, Hort. in Kew, Hortic. (1876) 200 = Argyrolobium Andrewsianum.
lotifolia, *Salisb. Parad. Lond.* t. 41.—Austral.
medicaginea, F. Muell. Second Gen. Rep. 11 = lotifolia.
polysperma, DC. in Mém. Soc. Phys. Genèv. ii. 11. 138; ex ej. Prod. ii. 117 = Argyrolobium Andrewsianum.
pubescens, *Sims, Bot. Mag.* t. 1810.—Austral.
retusa, Sweet, Hort. Brit. ed. I. 110, nomen. = lotifolia !
simplicifolia, Spreng. Syst. iv. Cur; Post. 267 = Hovea elliptica.
subpubescens, Sweet, Hort. Brit. ed. II. 196, nomen = lotifolia !

GOODIERA, Koch, Syn. Fl. Germ. ed. II. 602 (1844) = seq.

GOODYERA, R. Br. in Ait. Hort. Kew. ed. II. v. 197 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 609.
CIONISACCUS, Beck, Orch. Kohl & Hass. (1892 ?).
CORDYLESTYLIS, Falc. in Hook. Journ. Bot. iv. (1842) 74.
ELASMATIUM, Dalzc, Fl. Hentze-Fyr. 131 (1867).
EPIPACTIS, Hall. Enum. Stirp. Helv. i. 377 (1742).
GEOBINA, Raffn. Fl. Tellur. iv. 49 (1830).

GOODYERA:—
GRUECHIS, Lindl. in Wall. Cat. n. 7379 (1831); Bot. Reg. xix. (1833) t. 1618.
GONODONA, Link, Enum. Hort. Berol. ii. 369 (1829).
GOODIERA, Koch, Syn. Fl. Germ. ed. II. 602 (1844).
LEUCOSTACHYS, Hoffm. Preisv. Orch. (1843) 26.—Cf. Linnaea, xvi. (1842) Litt. 128.
PERAMIUM, Salisb. in Trans. Hort. Soc. i. (1812) 301
TUBSACA, Raffn. in Journ. Phys. lxxxix. (1819) 261.
bifida, *Blume, Orch. Archip. Ind.* 40.—Java.
cernua, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 80 = procera.
colorata, *Blume, Orch. Archip. Ind.* 37.—Java.
cordata, *Nichols. Dict. Gard.* ii. 81.—Reg. Himal.
Dawsonii, *Boxall, ex Newes, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 248.—Ins. Philipp.
discolor, Ker-Gawl. Bot. Reg. t. 271 = Haemaria discolor.
elongata, Lindl. Gen. et Sp. Orch. 494 = Hetaeria elongata.
fabellata, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 79 = Cheirostylis fabellata.
fumata, *Thw. Enum. Pl. Zeyl.* 314.—Zeylan.
fusca, *Lindl. in Wall. Cat.* n. 7395.—Reg. Himal.
gwyaneusis, Lindl. Gen. et Sp. Orch. 494 = Spiranthes Hostmanni.
hirsuta, Griff. Notul. iii. 393 = Physurus hirsutus.
hispida, *Lindl. in Journ. Linn. Soc.* i. (1857) 188.—Reg. Himal.
japonica, Blume, Orch. Archip. Ind. 38 = Schlechtendaliana.
lancifolia, *Franch. & Sav. Enum. Pl. Jap.* ii. 520.—Japon.
macranthe, *Maxim. in Fl. des Serres.* Sér. II. vii. (1867–68) 113.—Japon.
macrophylla, *Lowe, in Trans. Camb. Phil. Soc.* iv (1831) 13.—Ins. Madera.
margmata, Lindl. in Wall. Cat. n. 7394 = repens.
Maurevertii, *Blume, Orch. Archip. Ind.* 42.—Java.
Meniesii, *Lindl. Gen. et Sp. Orch.* 402.—Am. bor.
neglecta, *Ernst, Fl. Venez.* (1877) 359.—Venezuela.
nuda, *Thou. Orch. Iles Austr. Afr.* t. 29.—Ins. Mascar.
occulta, Thou. l. c. 28 = Platylepis goodyeroides.
Ordeana, *Boxall, ex Newes, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 248.—Ins. Philipp.
ovalifolia, Wight, Ic. t. 1780 = Hetaeria ovalifolia.
parviflora, *Blume, Orch. Archip. Ind.* 34. t. 22.—Malaya.
picta, *Boxall & Newes, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 248.—Ins. Philipp.
polygonoides, *F. Muell. Fragm.* viii. 29.—Austral.
procera, *Hook. Exot. Fl.* t. 99.—Ind. or.; China; Malaya.
pubescens, *R. Br. in Ait. Hort. Kew.* ed. II. v. 198.—Am. bor.
pusilla, *Blume, Orch. Archip. Ind.* 36.—Java.
querciola, Champ. Fl. S. U. St. 468 = Physurus quercifolia.
recurva, *Lindl. in Journ. Linn. Soc.* i. (1857) 188.—Reg. Himal.
repens, *R. Br. in Ait. Hort. Kew.* ed. II. v. 198.—Reg. bor. temp.
reticulata, *Blume, Orch. Archip. Ind.* 35. t. 9b.—Java.
rubens, *Blume, l. c.* 43.—Java.
rubicunda, *Lindl. in Bot. Reg.* (1839) *Misc.* 61.—Malaya.
rubrovens, *Boxall, ex Newes, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 248.—Ins. Philipp.
Schlechtendaliana, *Reichb. f. in Linnaea*, xxii. (1849) 861.—Japon.
secundiflora, Griff. Notul. iii. 393 (*Georchis foliosa*).—Ind. or.
secundiflora, *Lindl. in Journ. Linn. Soc.* i. (1857) 182.—Reg. Himal.
similis, *Blume, Orch. Archip. Ind.* 39.—Japon.
sinica, *Reichb. f. in Linnaea*, xviii. (1844) 409.—Mexic.
tesselata, Lodd. Bot. Cab. t. 952 = pubescens.
velutina, *Maxim. in Regel, Gartenfl.* (1867) 86.—Japon.
viridiflora, *Blume, Orch. Archip. Ind.* 41.—Java; Austral.
viscosa, *Reichb. f. in Linnaea*, xli. (1877) 61.—Ins. Borneo.
Weitziana, *Blume, Orch. Archip. Ind.* 42.—Java.
Wrightii, *Reichb. f. in Flora*, xlviii. (1865) 274.—Ind. occ.
Zollingeri, *Reichb. f. in Bonplandia*, x. (1862) 98.—Java.

GORCEIXIA, Baker, in Journ. Bot. xx. (1882) 225. *COMPOSITAE.*
decurrens, *Baker, l. c.*—Bras.

GORDONIA, Ellis, in Phil. Trans. lx. (1770) 518, t. 11. *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 186.
ANTHERISCHIMA, Korth. Verh. Nat. Gesch. Bot. 137. t. 27 (1840).
CARRIA, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 7.
DIPTEROSPERMUM, Griff. Notul. iv. 564 (1854).
FRANKLINIA, Bartr. ex Marsh. Arbust. Am. 48 (1785).
LACATHEA, Salisb. Parad. Lond. t. 56 (1806).
LASIANTHUS, Linn. Syst. ed. I (1735).
POLYSPORA, Sweet, Hort. Brit. ed. I. 61 (1826).
acuminata, Wall. Cat. n. 3664 = Pyrenaria acuminata; Adinandra acuminata.
acuminata, *Zoll. Syst. Verz. Ind. Archip.* 144, 148.—Java.
anomala, *Spreng. Syst.* iii. 126.—China.
axillaris, D. Dietr. Syn. Pl. iv. 868 = anomala.
barbinervis, Walp. Rep. i. 375 = Laplacea barbinervis.
brevifolia, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 162.—Borneo.
Chilauuna, Buch.-Ham. in D. Don, Prod. Fl. Nep. 226 = Schima Wallichii.
decandra, Roxb. Fl. Ind. ii. 573 = Ixonanthes reticulata.
dipterosperma, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) 11. 119.—Ind. or.
elliptica, *Gardn. in Calc. Journ. Nat. Hist.* vii. (1847) 448.—Zeylan.
excelsa, *Blume, Bijdr.* 130.—Reg. Himal.; Malaya.
floribunda, Wall. Cat. n. 1456 = Schima crenata.
Franklinii, L'Hérit. Stirp. Nov. 156 = pubescens.
grandis, *André, in Illust. Hortic.* xxvii. (1880) 60.—Hab.?
Haematoxylon, Sw. Fl. Ind. Occ. ii. 1199 = Laplacea Haematoxylon.
integerrima, *Teijsm. & Binn. Cat. Hort. Bog.* 204.—Java.
integrifolia, Roxb. Hort. Beng. 52; Fl. Ind. ii. 572 = Schima Wallichii.
javanica, Rolfsen, ex Hook. Bot. Mag. t. 4539 = Schima Noronhae.
Lasianthus, *Linn. Mant.* ii. 570.—Am. bor.
Lobbii, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 162.—Borneo.
Malingeyi, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 291.—Malacca.
marginata, *Endl. in Walp. Rep.* v. 134.—Borneo.
mollis, Wall. Cat. n. 1458 = Schima mollis.
obtusa, Roxb. Hort. Beng. 93; Fl. Ind. ii. 572 = Schima crenata.
obtusa, *Wall. Cat.* n. 1459.—Ind. or.
obtusifolia, Wight, Illustr. i. 99 = obtusa.
ovalis, *Korth. in Walp. Rep.* v. 134.—Sumatra.
parviflora, Hemsl. Diag. Fl. Nov. Mex. 4 = Symplocos coccinea.
parvifolia, Wight, Illustr. i. 99 = obtusa.
paucinervis, Wall. Cat. n. 4409 = Ixonanthes icosandra.
pauciflora, Blanco, Fl. Filip. ed. I. 549 = Homalium foetidum.
pubescens, *L'Hérit. Stirp. Nov.* 156.—Am. bor.
pyramidalis, Salisb. Prod. 386 = G. Lasianthus.
reticulata, Wall. Cat. n. 3663 bis, et n. 7070 = Adinandra integerrima.
semiserrata, *Spreng. Syst.* iv. *Cur. Post.* 260, 406.—Brasa.
singuloriana, Wall. Cat. n. 1457 = excelsa.
speciosa, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 143; *Thw. Enum. Pl. Zeyl.* 46.—Zeylan.
superba, Hook. f. & Thoms. ex Dyer, in Hook. f. Fl. Brit. Ind. i. 289 = Schima khasiana.
tomentosa, Spreng. Syst. iv. Cur. Post. 260 = Laplacea tomentosa.
villosa, Macfad. Fl. Jamaic. i. 117 = Laplacea villosa.
Wallichii, DC. Prod. i. 528 = Schima Wallichii.
wrightiana, *Wight, Illustr.* i. 99.—Zeylan.

GORGONIDIUM, Schott, in Mtq. Ann. Mus. Bot. Lugd. Bat. i. 228 (1865). *ARGIDEAE*, Benth. & Hook. f. iii. 262.
calodictyon, Schott, l. c.—Arabia. Ind.

GORINKIA, J. & C. Presl, Fl. Čech. 140 (1819) = Conringia, Heist. (Crucif.).
campestris, J. & C. Presl, l. c. 141 = Brassica campestris.
orientalis, J. & C. Presl, l. c. = C. orientalis.

GOROSTEMUM, Steud. Nom. ed. II. i. 631, in syn. (1840), sphalm. = GONOSTEMON, Haw. = Stipellis, Link.

GORSKIA, Bolle, in Peters, Reise Mossamb. Bot. 15. t. 3 (1861) = Copaifera, Linn. (Legumin.).
conjugata, Bolle, l. c. = C. Gorskiana.

GORTERA, Hill, Veg. Syst. iii. 81 (1761) = Gorteria, Linn. (Compos.).

GORTERIA, Linn. Sp. Pl. ed. II. 1282 (1763), pro parte; Linn. Illustr. t. 702 (1789?) = Berkheya, Ehrh. (Compos.).
acaulis, Hort.—Cf. Gard. Chron. (1883) i. 78 = Haplocarpha Leichtlinii.
arachnoidea, Hort. Angl. ex Steud. Nom. ed. II. i. 702 (= sec. ?).—Afr. austr.
aranosa, Meerb. Afbeel. Gew. t. 40 = Didelta cernua.
asteroides, Linn. f. Suppl. 381 = B. fruticosa.
asteroites, Steud. Nom. ed. II. i. 702 = prec.
barbata, Linn. f. Suppl. 382 = B. carlinoides.
carthamoides, Lam. Encyc. iii. 2 = B. grandiflora.
cernua, Linn. f. Suppl. 383 = Didelta cernua.
ciliaris, Linn. Sp. Pl. ed. II. 1284 = Cullumia ciliaris.
ciliata, Thunb. Prod. Pl. Cap. 168 = Cullumia patula.
cruciata, Houtt. Nat. Hist. ii. t. 70; ex DC. Prod. vi. 506 = B. cruciata.
echinata, [Dryand. in] Alt. Hort. Kew. ed. I. iii. 254 = Didelta cernua.
fruticosa, Linn. Sp. Pl. ed. II. 1284 = B. grandiflora.
herbacea, Linn. f. Suppl. 381 = B. cynaroides.
heterophylla, Willd. Hort. Berol. ii. 98 = Gazania rigens!
hispida, Lichst. ex Less. Syn. Comp. 55 = Gazania Burchellii.
hispida, Linn. f. Suppl. 383 = Cullumia hispida.
ilicifolia, Linn. Encyc. iii. 2 = B. fruticosa!
incisa, Thunb. Prod. Pl. Cap. 162 = Gazania pinnata.
integrifolia, Thunb. l. c. 163 = Hirpicium integrifolium.
linearis, Thunb. l. c. 163 = Gazania subulata.
Louvreana, DC. Prod. v. 501 (sp. dub.).—China.
lyrato-pinnatifida, Poir. Encyc. Suppl. iii. 819 = Gazania pinnata.
mitis, Burm. f. Fl. Cap. Prod. 28 = Heterolepis mitis.
oppositifolia, Lam. Encyc. iii. 4 = B. obovata.
othonnites, Thunb. Prod. Pl. Cap. 163 = Gazania othonnites.
ovata, Vahl, ex Less. Syn. Comp. 375 = Relhania rotundifolia.
Pavonia, Andr. Bot. Rep. t. 523 = Gazania Pavonia.
pectinata, Thunb. Prod. Pl. Cap. 162 = Gazania pinnata.
personata, E. Mey. ex DC. Prod. vii. 36 = Dicoma capensis.
pinnata, Lam. Encyc. iii. 1 = sec. ?
pinnata, Thunb. Prod. Pl. Cap. 162 = Gazania pinnata.
rigens, Linn. Pl. Afr. Rar. 23; Amoen. Acad. vi. 105 = Gazania rigens.
rigens, Thunb. Fl. Cap. 698 = Gazania Pavonia.
rigida, Banks, ex DC. Prod. vi. 522 = Gazania rigens, pinnata.
setosa, Linn. Mant. ii. 287 = Cullumia setosa.
setosa, Lour. Fl. Cochinch. 507 = *Loureiriana.*
speciosa, Poir. Encyc. Suppl. ii. 618 = Gazania pinnata.
spectabilis, Salisb. Prod. 287 = Gazania rigens.
spinosa, Jacq. Hort. Schoenb. iii. 64. t. 371 = Cullumia sulcata.
spinosa, Linn. f. Suppl. 381 = B. obovata.
squarrosa, Berg. Descr. Pl. Cap. 301 = Cullumia squarrosa.
squarrosa, Linn. Pl. Rar. Afr. 23; Amoen. Acad. vi. 105 = Cullumia decurrens.
uniflora, Linn. f. Suppl. 382 = Gazania uniflora.

GORTERIA, Linn. Syst. ed. X. 1229 (1759). *COMPOSITAE*, Benth. & Hook. f. ii. 459.
ICTINUS, Cass. in Bull. Soc. Philom. (1818) 143; et in Dict. Sc. Nat. xxii. 259 (1821).
PERSONARIA, Lam. Illustr. t. 716 (1789).
affinis, DC. Prod. vi. 501 = diffusa.
calendulacea, *DC. l. c.*—Afr. austr.
corymbosa, *DC. l. c.*—Afr. austr.
diffusa, *Thunb. Prod. Pl. Cap.* 162.—Afr. austr.
lectinus, Cass. in Dict. Sc. Nat. xxxiii. 456 = diffusa.
personata, Hill, Veg. Syst. ii. 81 = Gazania rigens.
personata, *Linn. Syst. ed. X.* 1229.—Afr. austr.

GOSELA, Choisy, in DC. Prod. xii 92 (1848). *SELA GINEAE*, Benth. & Hook. f. ii. 1129.
Ecklesiana, *Choisy. l. c.*—Afr. austr.

GOSSAMPINUS, Buch.-Ham. in Trans. Linn. Soc. xv. (1826) 126 = Bridesdonia, DC. (Malvac.).
alba, Buch.-Ham. l. c. = E. anfractuosum.
rubra, Buch.-Ham. l. c. 128 = Bombax malabaricum.
Rumphii, Schott & Endl. Meletem. 35 = E. anfractuosum.

GOSSANIA, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 328 = Gouania, Jacq. (Rhamn.).

GOSSYPIANTHUS, Hook. Ic. Pl. t. 251 (1840). *AMARANTACEAE*, Benth. & Hook. f. iii. 39.
australis, Griseb. in Goett. Abh. xix. (1874) 8 = Guillemines australis.
lanuginosus, Moq. in DC. Prod. xiii. II. 237 = Guillemines lanuginosa.
rigidiflorus, *Hook. Ic. Pl.* t. 251.—Am. bor. occ.
tenuiflorus, *Hook. l. c.* sub t. 251.—Am. bor. occ.
tomentosus, Griseb. in Goett. Abh. xxiv. (1879) 25 = Gomphrena demissa.

GOSSYPIUM, Linn. Syst. ed. I (1735). *MALVACEAE*, Benth. & Hook. f. i. 209.
STURTIA, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 68 (1849).
THURBERIA, A. Gray, in Mem. Am. Acad. N. S. v. (1854) (Pl. Thurb.) 308
VILLOA, Tourn. ex Medic. Malv. 43 (1787).
acuminatum, Roxb. Hort. Beng. 51; Fl. Ind. iii. 186 = barbadense.
albescens, *amblosepalum*, *asiaticum*, *aureum*, *bicolor*, *cambayense*, *cinereum*, *conoreum*, *Cunninghamii*, *decurrens*, *divaricatum*, *fuscum*, *guyanense*, *isabelum*, *macrospermum*, *Nankin*, *niveum*, *pallens*, *purpureum*, *Rabinandium*, *rupestre*, *sarmentosum*, *speciosum*, *telinum*, *tenax*, *trichosperma*, *virens*, *virgatum*, Rafin. Sylva Tellur. 14–18, omnes formae coltae G. herbacei, barbadensis, arborei, etc.
albiflorum, Tod. Oss. Sp. Cotone, 28 = roseum.
albun, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 494 = herbaceum!
anomalum, *Wawra & Peyr. in Sert. Benguel.* 21.—Afr. trop.
arboreum, *Linn. Sp. Pl.* 693.—Reg. trop.
arborreum, Parl. Sp. Cot. 52. t. 1, partim = neglectum.
arboreum, Vell. Fl. Flum. vii. t. 20 = brasilicum.
australe, F. Muell. Fragm. i. 46 = Fugosia australis.
australense, Tod. Oss. Sp. Cotone, 19 = Sturtii.
barbadense, *Linn. Sp. Pl.* 693.—Trop. trop.
brasilicum, *Macfad. Fl. Jamaic.* i. 72.—Am. trop.
caespitosum, *Tod. Oss. Sp. Cotone*, 64.—Hab.!
Cavanillesianum, *Tod. l. c.* 63.—Am. trop.
cernuum, *Tod. l. c.* 31.—Hab.!
chinense, Fisch. & Otto, ex Steud. Nom. ed. II. i. 702 = herbaceum.
cotulatum, Tod. Prod. Gossyp. 7 = Fugosia latifolia.
croceum, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 494 = herbaceum.
Cunninghamii, Tod. Prod. Gossyp. 7 = Fugosia punctata. *Benth.*
Davidsoni, *Kellogg, in Proc. Calif. Acad.* v. (1873) 32.—Am. bor. occ.
drynarioides, *Sesse, Fl. Vit.* 22.—Ins. Sandvic.
eglandulosum, Cav. Diss. vi. 354 = herbaceum.
elatum, Salisb. Prod. 385 = herbaceum.
Figueri, *Tod. in Att. Soc. Agric. Sicil.* iv; *ex Tod. Rel. Cot. Colt.* 12.—Hab.!
flaviflorum, F. Muell. Fragm. ix. 127 = Fugosia flaviflora.
frutescens, Lastayr. Cot. 65. t. 2; ex Tod. Rel. Cot. Colt. 240 = barbadense.
frutioulosum, *Tod. Prod. Gossyp.* 9.—Mexic.
fuscum, Roxb. ex Wight & Arn. Prod. 448 = barbadense.
glabratum, *Tod. Oss. Sp. Cotone*, 49.—Hab.!
glabrum, Lam. Encyc. ii. 138 = barbadense.
glandulosum, Steud. Nom. ed. I. 381, sphalm. = *eglandulosum.*
gracile, Salisb. Prod. 385 = arboreum.
herbaceum, *Linn. Sp. Pl.* 693.—As. trop.
hirsutum, Cav. Diss. vi. 313. t. 167 = Cavanillesianum.
hirsutum, Linn. Sp. Pl. ed. II. 975 = herbaceum.
indicum, Lam. Encyc. ii. 134 = herbaceum.
intermedium, *Tod. Oss. Sp. Cotone*, 41.—As. or.

GOSSYPIUM:—
jamaicense, Macfad. Fl. Jamaic. i. 73 = barbadense?
jamphesicum, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 243.—Porto Rico.
javanicum, Blume, Bijdr. 74 = barbadense.
javanicum, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 488.—Ins. Moluce.
Kirkii, *Mast. in Journ. Linn. Soc.* xix. (1882) 214.—Afr. trop.
Klotzschianum, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1856) 238.—Ins. Galapag.
Labillarderianum, *Tod. Prod. Gossyp.* 11.—Ins. Waigiou.
lanceolatum, *Tod. l. c.* 9.—Mexic.
latifolium, Murr. in Comm. Gotting. (1776) 32. t. 1 = herbaceum.
leonimum, Medic. Beobacht. (1783) 199 = herbaceum.
macedonicum, Murr. in Comm. Gotting. (1776) 32 = herbaceum.
macranthum, *Tod. Prod. Gossyp.* 11.—Aegypt.
marrinum, Tod. Ons. Spc. Cotone, 33 = barbadense.
mexicanum, *Tod. Ind. Sem. Hort. Panorm.* (1867) 20, 31; *Prod. Gossyp.* 9.—Mexic.
micranthum, Cav. Diss. vi. 311; DC. Prod. i. 456 = herbaceum.
microcarpum, *Tod. in Att. Soc. Agric. Sicil. iv; esej. Rel. Cot. Coll.* 15.—Mexic.
molle, Mauri, ex Tenore, Cat. Hort. Nap. (1845) 86 = herbaceum.
Nanking, Meyen, Reise, ii. 328 = herbaceum.
neglectum, *Tod. Ons. Sp. Cotone*, 35.—Hab.?
nigrum, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 494 = barbadense, arboreum.
obtusifolium, Roxb. Hort. Beng. 51; Fl. Ind. iii. 183 = herbaceum.
oligospermum, Macfad. Fl. Jamaic. i. 74 = barbadense?
pallidum, *Montr. in Mém. Acad. Lyon*, x. (1860) 181.—N. Caled.
paniculatum, Blanco, Fl. Filip. ed. I. 589 = herbaceum?
parvifolium, Nutt. ex Seem. Fl. Vit. 23 = tomentosum.
perrone, Blanco, Fl. Filip. ed. I. 587 = barbadense?
peruvianum, Cav. Diss. vi. 313. t. 168 = barbadense.
populifolium, F. Muell. Fragm. ix. 127 = Fagosia populifolia.
prostratum, *Schum. & Thonn. Beskr. Guin. Pl.* 311.—Afr. trop.
puberulum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 128.—Afr. trop.
pubescens, *Spreng. ex De Vriese, in Nederl. Kruidk. Arch.* i. (1848) 33.—Guian.
punctatum, Guill. & Perr. Fl. Seneg. Tent. i. 62 = herbaceum.
punctatum, Schum. & Thonn. Beskr. Guin. Pl. 309 = barbadense.
punicum, Fenzl, in Jacq. Eclog. ii. 7. t. 184 = sanguineum.
purpurascens, Poir. Encyc. Suppl. ii. 369 = arboreum.
racemosum, Poir. l. c. 370 = barbadense.
religiosum, Linn. Syst. ed. XII. 642 = herbaceum.
religiosum, Farl. Sp. Cotoni Firens. (1866) 54. t. 4 = barbadense.
Rhorii, *Tod. Prod. Gossyp.* 10.—Hab.?
Robinsoni, *F. Muell. Fragm.* ix. 126.—Austral.
roseum, *Tod. Ons. Sp. Cotone*, 25.—Hab.?
Roxburghii, Tod. l. c. 45 = neglectum.
Reyleanum, Tod. l. c. 40, 41 = intermedium.
rubicundum, Roxb. ex Wight & Arn. Prod. 55 = arboreum.
rubrum, Forsk. Fl. Aegypt. Arab. 88 = arboreum.
rufum, *Scop. Delic. Insub.* iii. 70.—China.
sanderiense, Farl. Sp. Cotoni Firens. (1866) 57. t. 6 = tomentosum.
sanguineum, *Hassk. in Hoev. & De Vriese, Tijdschr.* xii. (1845) 98.—Java?
senarense, Fenzl, ex Mast. in Oliver, Fl. Trop. Afr. i. 211 = anomalum.
siamense, Tenore, in Att. Ist. Incorr. Nap. vi. (1840?) 189 = herbaceum.
sinense, Finch. Hort. Gorenk. ed. II. 56, nomen = herbaceum.
Stocksii, *Mast. in Hook. f. Fl. Brit. Ind.* i. 346.—Sindh.
strictum, Medic. Beobacht. (1783) 195 = herbaceum.
Sturtii, *F. Muell. Fragm.* iii. 6.—Austral.
suffruticosum, *Bertol. Diss. Pl. Nov. et Byss. Antiq.* 6.—Ind. or.

GOSSYPIUM:—
taitense, *Parl. Sp. Cotoni Firens.* (1866) 59. t. 6. f. A.—Ins. Tahiti.
thespesioides, F. Muell. Fragm. ix. 27 = Fagosia thespesioides.
Thurberi, *Tod. Prod. Gossyp.* 7.—Mexic.
tomentosum, *Nutt. ex Seem. Fl. Vit.* 23.—Ins. Sandvic.
trinervidatum, Lam. Encyc. ii. 286 = herbaceum.
vitifolium, Lam. l. c. 285 = barbadense.
vitifolium, Roxb. Hort. Beng. 51; Fl. Ind. iii. 186 = herbaceum.
Welchettianum, Tod. Rel. Cot. Colt. 119 = Sturtii?
Wightianum, *Tod. Ons. Sp. Catone*, 47.—Ind. or.

GOTHOFREDA, Vent. Choix, 36. t. 60 (1808) = **Oxypetalum**, R. Br. (1809) (Asclepiad.).
cordifolia, Vent. l. c. = O. riparium.

GOUANA, Linn. Sp. Pl. ed. II. 1663 (1763) = seq.

GOUANIA, Jacq. Stirp. Select. Am. 268 (1763). *RHAMNEAE*, Benth. & Hook. f. i. 385.
NARGELIA, Zoll. & Mor. Verz. 90 (1846).
RETINARIA, Geertn. Fruct. ii. 197 (1791).
acalyphoides, *Reiss. in Mart. Fl. Bras.* xi. i. 108.—Bras.
- alnifolia, *Reiss. l. c.* 106.—Peruv.
aphrodes, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 132.—Madag.
aptera, *DC. Prod.* ii. 80.—Am. trop.
aptera, Spreng. ex Reiss. in Mart. Fl. Bras. xi. i. 106 = alnifolia.
australiana, *F. Muell. Fragm.* iv. 144.—Austral.
bancana, *Teijsm. & Binn. Cat. Hort. Bog.* 221, nomen.—Ins. Bancа.
Blanchetiana, *Miq. in Linnaea*, xxii. (1849) 797.—Am. trop.
Brandisii, Kurz, in Flora, liv. (1871) 280, in nota = microcarpa.
brasiliensis, *Hoffmgg. Vers. Pfl. Nachtr.* ii. 126.—Bras.
casnecens, *Rich. ex Poir. Encyc. Suppl.* ii. 820.—Guian.
chrysophylla, *Reiss. in Mart. Fl. Bras.* xi. i. 108.—Bras.
columnsfolia, *Reiss. l. c.* 107.—Bras.
cordifolia, Raddi, in Mem. Mod. xviii. Fis. (1820) 351 = Reisackia amilacina.
cornifolia, *Reiss. in Mart. Fl. Bras.* xi. i. 107.—Bras.
corylifolia, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 351.—Bras.
crenata, *Lam. Encyc.* iii. 5.—Am. austr.
cyclocarpa, *Sm. in Rees, Cycl.* xvi. n. 3.—Ind. occ.
dasyantha, *Miq. Anal. Bot. Ind.* iii. 6.—Ind. or.
denticulata, *Sm. in Rees, Cycl.* xvi. n. 4.—Ind. or.
discolor, *Benth. in Hook. Kew Journ.* iv. (1852) 12.—Bras.
domingensis, Aubl. Pl. Gui. ii. 904 = striata.
domingensis, Blanco, Fl. Filip. ed. I. 196 = microcarpa.
domingensis, *Linn. Sp. Pl.* ed. II. 1668.—Ind. occ.
ericarpa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 132.—Ins. Comor.
frangulaefolia, *Radlk. in Sitzb. Math.-Phys. Acad. Munch.* xvii. (1878) 892.—Venezuela.
frangulaefolia, Willd. ex Reiss. in Mart. Fl. Bras. xi. i. 105 = Blanchetiana.
glabra, Jacq. Select. Am. 264 = domingensis.
glabriuscula, Stokes, Bot. Mat. Med. i. 496 = domingensis.
glandulosa, *Boia. ex Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 129.—Madag.
Hillii, *F. Muell. Fragm.* viii. 168.—Austral.
hypochroa, *Reiss. in Mart. Fl. Bras.* xi. i. 106.—Bras.
incisa, Vahl, Symb. Bot. iii. 101 = mauritiana.
inornata, *Reiss. in Mart. Fl. Bras.* xi. i. 109.—Bras.
integrifolia, Kurz, in Journ. As. Soc. Beng. xl. (1872) ii. 49 = microcarpa.
integrifolia, *Lam. Encyc.* iii. 5.—Hab.?
integrifolia, Meyen, Reise, ii. 180 = Meyeni.
javanica, *Miq. Fl. Ind. Bat.* i. 1. 649.—Java.
lamellata, Wall. Cat. n. 6878 = Helinus lamellatus.
latifolia, *Reiss. in Mart. Fl. Bras.* xi. i. 108.—Bras.
lucifera, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 130.—Madeira.
leptonachya, *DC. Prod.* ii. 80.—Ind. or.; Malaya.

GOUANIA:—
leptostachya, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 49 = microcarpa.
linosta, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 131.—Madag.
longipetala, *Hemsl. in Oliver, Fl. Trop. Afr.* i. 383.—Afr. trop.
madagascariensis, Rich. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 129 = glandulosa.
mortificensis, Poir. Encyc. Suppl. ii. 820 = pubescens.
mauritiana, *Lam. Encyc.* iii. 4.—Ins. Mascar.
Meyeni, *Steud. Nom.* ed. II. i. 708.—Ins. Sandvic.
microcarpa, *DC. Prod.* ii. 80.—Ind. or.; Malaya.
mollis, *Reiss. in Mart. Fl. Bras.* xi. i. 104.—Bras.
myricocarpa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 133.—Madag.
napalensis, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 417.—Reg. Himal.
obtusifolia, *Vent. ex Brongn. in Ann. Sc. Nat.* Ser. i. x. (1827) 379.—Am. austr.
orbiculoris, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 323 = Meyeni.
paniculata, *Spreng. Neue Entdech.* iii. 49.—Ins. S. Doming.
pannigera, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 134 = speciosa.
petiolaris, *Reiss. in Mart. Fl. Bras.* xi. i. 108.—Bras.
pubescens, *Lam. Illustr.* t. 845. f. 1; ex Poir. Encyc. Suppl. ii. 819.—Ind. or.
pyrifolia, *Reiss. in Mart. Fl. Bras.* xi. i. 110.—Bras.
Reisackia, DC. Prod. ii. 40 = tiliaefolia.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 282.—Ins. Fiji.
riparia, *Reiss. in Mart. Fl. Bras.* xi. i. 104.—Bras.
rumicina, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1873) 383.—N. Granat.
sericea, Hillenh. ex Presl, Bot. Bemerk. 39 = speciosa
Sieberiana, Schlecht. ex Presl, l. c. = tiliaefolia?
smilacina, Sm. in Rees, Cycl. xvi. n. 5 = Reisackia amilacina.
Stadtmanni, Soyer-Willem. in Uuterl, Ann. Bot. xviii. (1796) 58 = tiliaefolia.
stipularis, [*Mq. & Sesse,*] *ex DC. Prod.* ii. 39.—Mexic.
striata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 113.—Guian.
sulcata, *Roj. Hort. Maurit.* 71.—Madag.
tiliaefolia, *Lam. Encyc.* iii. 4.—Ins. Macaar.
tiliaefolia, Rottb. ex DC. Prod. ii. 40 = microcarpa.
tiliaefolia, Roxb. Pl. Corom. i. 67. t. 98 = leptostachya.
timorensis, *Zipp. ex Span. in Linnaea*, xv. (1841) 187.—Ins. Timor.
tomentosa, *Yacq. Select. Am.* 268.—Ind. occ.
trichodonta, *Reiss. in Mart. Fl. Bras.* xi. i. 108.—Peruv.
ulmifolia, *Hook.* & *Arn. in Hook. Bot. Misc.* iii. (1833) 174.—Chili.
ulmifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1873) 383.—N. Granat.
urticaefolia, *Reiss. in Mart. Fl. Bras.* xi. i. 110.—Bras.
velutina, *Reiss. l. c.* 105.—Guian.
virgata, *Reiss. l. c.* 104.—Bras.
vitifolia, *Spreng. Syst. U. St. Expl. Exped.* i. 283.—Ins. Sandvic.

GOUAREA, Hedw. f. Gen. 274 (1806) = **Guarea**, Allem. (Meliac.).

GOUDENIA, Vent. Tabl. ii. 473 (1799) = **Goodenia**, Sm.

GOUDOTIA, Decne. in Ann. Sc. Nat. Sér. III. iv. (1845) 84. t. 4 = **Dietschia**, Nees (Juncac.).
tolimensis, Decne. l. c. 85 = D. tolimensis.

GOUFFEIA, Robil. & Cast. ex DC. Fl. Fr. vi. 409 (1815) = **Arenaria**, Rupp. (Caryophyll.).
arenarioides, Robil. & Cast. l. c. = A. Gouffeia.
crassinervis, Cambess. in Jacquem. Voy. Bot. t. 30 = A. holosteoides.
holosteoides, C. A. Mey. Verz. Pfl. Cauc. 217 = A. holosteoides?
sinica, Fenzl, Verhe. Akad. zool. p. 27 = Buffonia tenuifolia.

GOUGHIA, Wight, Ic. v. 22. t. 1878, 1879 (1852) = **Daphniphyllum**, Blume (Euphorb.).
Griffithiana, Wight, l. c. = D. laurinum.
himalayensis, Benth. in Hook. *Kew Journ. vi. (1854)* 9 = D. himalayense.
laurina, Benth. l. c. = D. laurinum.
nilgherrensis, Wight, Ic. v. t. 1878, 1879 = D. glaucescens.

GOUINIA, Fourn. ex Benth. & Hook. f. Gen. iii. 1178 (1883) = **Festuca**, Tourn. (Gramin.).
polygama, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 551 = F. Fournieriana.

GOULDIA, A. Gray, in Proc. Am. Acad. iv. (1860) 319. *RUBIACEAE*, Benth. & Hook. f. ii. 77.
axillaris, *Wawra, in Flora*, lvii. (1874) 297.—Ins. Sandvic.
Romanzoffensis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 319.—Ins. Sandvic.
sandvicensis, *A. Gray, l. c.*—Ins. Sandvic.

GOUFIA, Aubl. Pl. Gui. i. 395. t. 146 (1775). *CELASTRINEAE*, Benth. & Hook. f. i. 369.
GLOSSOPETALUM, Schreb. Gen. i. 205 (1789).
GUPIA, Jaume St. Hil. Expos. ii. 267 (1805).
glabra, *Aubl. Pl. Gui.* i. 396. t. 146.—Guiana.
tomentosa, *Aubl. l. c.*—Guiana.

GOURLIEA, Gill. ex Hook. Bot. Misc. iii. (1833) 207. t. 106. *LEGUMINOSAE*, Benth. & Hook. f. i. 556.
chilensis, *Clos, in C. Gay, Fl. Chil.* ii. 218.—Chili.
decorticans, *Gill. ex Hook. Bot. Misc.* iii. (1833) 208. —Chili.

GOVANA, All. Misc. Taur. v. 98 (1778) = **Gouania**, Jacq. (Rhamn.).

GOVANIA, Raddi, in Mem. Mod. xviii. Fis. (1820) 394, 395 = prec.

GOVANIA, Wall. Cat. n. 7851, nomen (1847 ?) = **Grivola**, Gmel. (Euphorb.).
nivea, Wall. l. c. = Givotia rottleriformis.

GOVENIA, Lindl. in Lodd. Bot. Cab. t. 1709 (1831). *ORCHIDEAE*, Benth. & Hook. f. iii. 549.
EUCHEMIS, Lindl. Gen. et Sp. Orch. 161 (1833).
alba, *A. Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 25.—Mexic.
Andrieuxii, *Reichb. f. in Bot. Zeit.* x. (1852) 835.—Mexic.
barbata, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 5.—Peruv.
brevilabris, *Hemsl. Biol. Centr. Am. Bot.* iii. 249.—Mexic.
capitata, *Lindl. Bot. Reg.* sub t. 1795.—Mexic.
deliciosa, *Reichb. f. in Bot. Zeit.* x. (1852) 836.—Mexic.
fausta, *Lindl. Bot. Reg.* (1848) *Misc.* 70.—Venezuela.
Gardneri, *Hook. Bot. Mag.* t. 5660.—Bras.
lagenophora, *Lindl. Bot. Reg.* (1839) *Misc.* 46.—Mexic.
liliacea, *Lindl. l. c.* sub t. 1795; *et* (1838) t. 18.—Mexic.
limbata, Griseb. Fl. Brit. W. Ind. 628 (sp. delend.).
mutica, *Reichb. f. in Bot. Zeit.* x. (1852) 836.—Mexic.
pacifica, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 134.—Mexic.
quadriplicata, *Reichb. f. Beitr. Orch. Centr. Am.* 75.—Costa Rica.
sulphurea, *Reichb. f. in Gard. Chron.* (1865) ii. 70.—Paraguay.
superba, *Lindl. in Lodd. Bot. Cab.* t. 1709.—Mexic.
tinguia, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 5.—Peruv.
utriculata, *Lindl. Bot. Reg.* (1839) *Misc.* 47, in obs.—Ind. occ.; Mexic.

GOVINDOOIA, Wight, Ic. vi. 24. t. 2000 (1852) = **Tropidia**, Lindl. (Orchid.).
navicula, Wight, l. c.—Ind. or.

GOVINDOVIA, C. Mosll. in Walp. Ann. vi. 158 (1861) = prec.

GRABOWSKIA, Schlecht. in Linnaea, vii. (1832) 71. *SOLANACEAE*, Benth. & Hook. f. ii. 899.
PURDANTHUS, Reds. Sylva. Tellur. 53 (1838).
boerhaaviaefolia, Lindl. Bot. Reg. t. 1985 = Lindleyl.
boerhaaviaefolia, *Schlecht. in Linnaea*, vii. (1832) 72. —Peruv.
cuneifolia, *Dunal, in DC. Prod.* xiii. i. 20.—Bras.
distichа, *Walp. Rep.* iii. 118.—Peruv.
duplicata, *Arn. in Linnaea*, xi. (1837) 485.—Bras.
Lindleyi, *Sendt. in Mart. Fl. Bras.* x. 157.—Bras.
Meyeniana, *Walp. Rep.* iii. 118.—Peruv.
obscura, Dunal, in DC. Prod. xiii. i. 20, sphalm. = seq.
obtusa, *Arn. in Linnaea*, xi. (1837) 485.—Peruv.
Schlechtendalii, *Sendt. in Mart. Fl. Bras.* x. 156.—Bras.

GRABOWSKYA, Endl. Gen. 645 (1839) = prec.

GRACILEA, Koen. ex Rottl. in Ges. Nat. Fr. Neue Schr. iv. (1803) 218 = **Pommereulla**, Linn. f. (Gramin.).
nutans, Koen. l. c. = P. monoeca.

GRADERIA, Benth. in DC. Prod. x. 521 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 970.
ROPURIA, Presl, Bot. Bemerk. 91 (1844), nomen prius.
fruticosa, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 84.—Ins. Socotra.
scabra, *Benth. in DC. Prod.* x. 521.—Afr. austr.

GRAEFFRA, Scem. Fl. Vit. 27. t. 6 (1865). *TILIACEAE*, Benth. & Hook. f. i. 236.
calycolata, *Seem. l. c.*—Ins. Fiji.

GRAEFFENRIEDA, D. Dietr. Syn. Pl. ii. 1875, 1481 (1840) = **Graffenrieda**, DC. (Melast.).

GRAELLSIA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 172. *CRUCIFERAE*, Benth. & Hook. f. i. 72.
saxifragaefolia, *Boiss. l. c.* 173.—Persia.

GRAEMIA, Hook. Exot. Fl. t. 189 (1827) = **Cephalophora**, Cav. (Compos.).
aromatica, Hook. l. c. = C. aromatica.
aromatica, Poepp. ex DC. Prod. v. 501 = Podanthus Mitiqui.

GRAEVIA, Neck. Elem. ii. 258 (1790) = **Grewia**, Linn. (Tiliac.).

GRAFFENRIEDA, DC. Prod. iii. 105 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 750.
CYSNOPODIUM, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 52.
anomala, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 71.—N. Granat.
brachycentra, Griseb. in Mem. Ac. Acad. N. S. viii. (1861) 186 = Tetraxygia brachycentra.
caryophyllea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 69.—Bras.
chrysandra, *Triana, l. c.* 70.—Cuba.
conostegioides, *Triana, l. c.* 68.—N. Granat.
emarginata, *Triana, l. c.* 71.—Peruv.; N. Granat.
excelsa, DC. Prod. iii. 106 = Centronia excelsa.
floribunda, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 71.—Peruv.
intermedia, *Triana, l. c.*—Peruv.
jucunda, Mart. Nov. Gen. et Sp. iii. 145. t. 276 = Miconia jucunda.
latifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 71.—Am. trop.
laurina, *Triana, l. c.* 70.—Bras.
linifolia, *Triana, l. c.*—Peruv.
micranthoides, *Naud. in Ann. Sc. Nat. Sér.* III. xviii. (1852) 116.—Bras.
Moritziana, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 70.—Venezuela.
obliqua, *Triana, l. c.* 71.—Guian.
obtusa, Griseb. Cat. Pl. Cub. 103, partim = Miconia obtusa.
ovalifolia, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 117. t. 3. f. 4 = Waddellii.
patens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 70. —Bras.
rotundifolia, *DC. Prod.* iii. 106.—Venezuela.
sessilifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 70.—Guian.
Waddellii, *Naud. in Ann. Sc. Nat. Sér.* III. xviii. (1852) 117.—Bras.

GRAFFENRIEDERA, Reichb. Consp. 174 (1828) = prec.

GRAFFENRIEDIA, Spreng. Gen. i. 841 (1830) = prec.

GRAFIA, Reichb. Handb. 219 (1837) = **Pleurospermum**, Hoffm. (Umbellif.).

GRAHAMIA, Gill. in Hook. Bot. Misc. iii. (1833) 231. *POLYPLACACEAE*, Benth. & Hook. f. i. 157.
XERANTHUS, Miers, Trav. Chil. ii. 529 (1826), nomen.
bractеata, *Gill. in Hook. Bot. Misc.* iii. (1833) 232.—Chili.

GRAHAMIA, Spreng. Syst. iv. Cur. Post. 297, 299 = **Cephalophora**, Cav. (Compos.).
aromatica, Spreng. l. c. 299 = C. aromatica.

GRAMMERUM, Desv. Opusc. 61 (1831). *GRAMINEAE*.
convolutum, *Desv. l. c.*—Chili.

GRAMMADENIA, Benth. Pl. Hartw. 218 (1846). *MYRSINEAE*, Benth. & Hook. f. ii. 644.
linearis, *Benth. l. c.*—Guiana.
marginata, *Benth. l. c.*—N. Granat.
parasitica, *Griseb. Fl. Brit. W. Ind.* 393.—Ind. occ.

GRAMMANGIS, Reichb. f. in Hamb. Gartens. xvi. (1860) 590; et Xenia Orch. ii. 17 (1878 ?). *ORCHIDEAE*, Benth. & Hook. f. iii. 537.
Ellisii, *Reichb. f. ll. c.*—Madag.
falcigera, *Reichb. f. in Flora*, lxviii. (1885) 541.—Madag.!
Hottoni, Benth. & Hook. f. Gen. iii. 588 = Cymbidium Huttoni.
pardalina, *Reichb. f. in Flora*, lxviii. (1885) 541.—Madag.!

GRAMMANTHES, DC. Prod. iii. 392 (1828). *CRASSULACEAE*, Benth. & Hook. f. i. 658.
VAUANTHES, Haw. Revis. Pl. Succ. 18 (1821), nomen prius.
caesia, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 331 = gentianoides.
chloraefora, DC. Prod. iii. 392 = gentianoides.
chloraefolia, Steud. Nom. ed. II. i. 703 = prec.
depressa, Eckl. & Zeyh. Enum. 308 = gentianoides.
filiformis, Eckl. & Zeyh. l. c. = gentianoides.
flava, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 331 = gentianoides.
gentianoides, *DC. Prod.* iii. 393.—Afr. austr.
retroflexa, Sweet, Hort. Brit. ed. II. 223 = gentianoides.
sebaoides, Eckl. & Zeyh. Enum. 308 = gentianoides.

GRAMMARTHEON, Reichb. in Moesal. Handb. ed. II. ii. 1508 (1838) = seq.

GRAMMARTHRON, Cass. in Bull. Soc. Philom. (1817) 82 = Dovonieuxn, Linn. (Compos.).
bilignatum, Cass. in Dict. Sc. Nat. xix. 295 = D. Claus.
oppositifolium, Cass. l. c. 296 = Arnica nudicaulis.
scorpioides, Cass. l. c. 294 = D. grandiflorum.

GRAMMATOCARPUS, Presl, Symb. Bot. i. 59. t. 28 (1831). *LOASEAE*, Benth. & Hook. f. i. 806.
SCYPHANTHUS, D. Don, in Sweet, Brit. Flow. Gard. Ser. I. t. 238 (1828), nomen prius.
Cumingii, *Presl, Epim. Bot.* 246.—Chili.
volubilis, *Presl, Symb. Bot.* i. 60.—Chili.

GRAMMATOPHYLLUM, Blume, Bijdr. 377. t. 20 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 529.
GASERTIA, Gaudich. in Freyc. Voy. Bot. 425 (1826).
PAXTONIA, Wight, Ic. t. 1750 (1851).
elegans, *Reichb. f. in Gard. Chron.* (1882) ii. 776.—Ins. Pacif.
elegans, Walp. Ann. in. 1028 ind., sphalm. = Cymbidium elegans.
Ellisii, Lindl. in Bot. Mag. t. 5179 = Grammangis Ellisii.

GRAMMATOPHYLLUM :—
festivum, Lindl. & Part. Flow. Gard. ii. (1851-52) 180 = speciosum.
Fenellianum, *Reichb. f. Xenia Orch.* ii. 16.—Ins. Amboin.
Finlaysonianum, Lindl. in Wall. Cat. n. 7561 = Brombeadia palustris.
gigantæum, Blume, ex Reichb. f. in Linnæa, xli. (1877) 107, in obs. sphalm. ! = speciosum.
macranthum, Reichb. f. Xenia Orch. ii. 16 = speciosum.
multiflorum, *Lindl. Bot. Reg.* (1838) *Misc.* 46.—Ins. Philipp.
paludosum Griff. Notul. iii. 344 = Dipodium paludosum.
pantherinum, *Reichb. f. in Gard. Chron.* (1878) i. 793.—N. Guinea.
pantherinum, Zipp. ex Blume, Rumphia, iv. 50 = Vanda lissochiloides.
pubescens, Reichb. f. in Walp. Ann. vi. 680, sphalm. = sco.
pulchrum, Span. in Linnæa, xv. (1841) 477, nomen = Vanda insignis.
Reemplerianum, *Reichb. f. in Gard. Chron.* (1877) i. 340.—Madag.
Rumphianum, *Mig. Ann. Mus. Bot. Lugd. Bat.* iv. 219.—Ins. Moluce.
scandens, Griff. Notul. iii. 345 = Dipodium pictum.
scriptum, Blume, Rumphia, iv. 48 = speciosum.
speciosum, *Blume, Bijdr.* 378.—Malaya.
Wallisii, *Reichb. f. in Linnæa*, xli. (1877) 107.—Ins. Philipp.

GRAMMATOTHECA, Presl, Prod. Monog. Lobel. 43 (1836) = **Lobelia**, Linn. (Campan.).
Bergiana, Presl, l. c. 44 = L. Bergiana.
Dregeana, Presl, l. c. — L. Bergiana.
Eckloniana, Presl, l. c. = L. Bergiana.
erinoides, Sond. in Harv. & Sond. Fl. Cap. iii. 532 = L. Bergiana.
Meyeriana, Presl, Prod. Monog. Lobel. 44 = L. Bergiana.
Mundtiana, Presl, l. c. = L. Bergiana.

GRAMMICA, Lour. Fl. Cochinch. 170 (1790) = **Ouvouta**, Linn. (Convolv.).
americana, Des Moul. in Bull. Soc. Bot. Fr. i. (1854) 298 = C. americana.
aphylla, Lour. Fl. Cochinch. 171 = C. chinensis.
arabica, Des Moul. in Bull. Soc. Bot. Fr. i. (1854) 298 = C. arabica.
Cheyssoana, Des Moul. l. c. = C. australis.
obtusiflora, Berthiot, in Ellotia, i. (1864) 15 = C. australis.
suaveolens, Des Moul. in Bull. Soc. Bot. Fr. i. (1854) 296 = C. suaveolens.

GRAMMOCARPUS, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 16 = **Trigonella**, Linn. (Leguminos.).
cæruleus, Schur, l. c. = T. cæruleus.
procumbens, Schur, Enum. Fl. Transs. 152 = T. Bemerinus.

GRAMMOPETALUM, C. A. Mey. ex Meissn. in Linnæa, xxx. (1859-60) 514 = **Trinia**, Hoffm. (Umbellif.).
Lobelowii, C. A. Mey. l. c. = T. Kitaibelii.

GRAMMOSCIADIUM, DC. Coll. Mém. v. 62 (1829); et Prod. iv. 232 (1830) = **Chærophyllum**, Linn. (Umbellif.).
Aucheri, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 67. —Armen.
daucoides, DC. Coll. Mém. v. 62. t. 2. f. K.—As. Min.
Haussknechtii, Boiss. Fl. Orient. ii. 901.—Syria.
Kotschyi, Boiss. ex Boiss. l. c. 899 = *daucoides*.
longilobum, Boiss. & Hausskn. ex Boiss. l. c. 900.—Persia.
macrodon, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 67.—Armen.
nodosum, DC. Coll. Mém. v. 63 = C. macrospermum.
platycarpum, Boiss. & Hausskn. ex Boiss. Fl. Orient. ii. 901.—Persia.
pterocarpum, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 66.—As. Min.
scabridum, Boiss. l. c. 66.—Armenia.
Szowitzii, Boiss. l. c. 67 = *daucoides*.

GRANADILLA, [Tourn.], Rapp. Fl. Jen. ed. Hall. 166 (1745) = **Passiflora**, Linn.
capsularis, Medic. Malv. 96 = P. capsularis.
cærulea, Medic. l. c. = P. cærulea.
fœtida, Gaertn. Fruct. i. 389. t. 80 = P. fœtida.
Hondala, Gaertn. l. c. ii. 450 = Modecca palmata.
incarnata, Medic. Malv. 96 = P. incarnata.
laurifolia, Medic. l. c. 97 = P. laurifolia.
lutea, Medic. l. c. 96 = P. lutea.
normalis, Gaertn. Fruct. ii. 479 = P. perfoliata.
quadrangularis, Medic. Malv. 97 = P. quadrangularis.
rubra, Moench, Meth. Suppl. 15 = P. rubra.
serratifolia, Medic. Malv. 97 = P. serratifolia.
suberosa, Gaertn. Fruct. ii. 480 = P. suberosa.
Vespertilio, Moench, Meth. Suppl. 14 = P. Vespertilio.

GRANATUM, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 183 = **Punica**, Tourn. (Lythrar.).
punicum, St. Lag. l. c. = P. Granatum.

GRANDIDIERA, Jaub. in Bull. Soc. Bot. Fr. xiii. (1866) 467. *BIXINEÆ*, Benth. & Hook. f. i. 971. Bolvini, *Jaub. l. c.*—Afr. trop.

GRANGEA, Adans. Fam. ii. 121 (1763). *COMPOSITÆ*, Benth. & Hook. f. ii. 261.
PYRARDA, Cass. in Dict. Sc. Nat. xli. 120 (1826).
Adansonii, Cass. in Dict. Sc. Nat. xix. 304 = maderaspatana.
aegyptiaca, DC. Prod. v. 378 = maderaspatana.
bicolor, Loud. Hort. Brit. 354 = Dichrocephala latifolia.
cernuanoides, Cass. in Dict. Sc. Nat. xix. 307 = maderaspatana.
chinensis, Loud. Hort. Brit. 354 = Centipeda orbicularis.
cuneus, Link, Enum. Hort. Berol. ii. 344 = Cotula dioeca.
cuneifolia, Poir. Encyc. Suppl. ii. 825 = Centipeda orbicularis.
decumbens, Desf. Tabl. ed. I. 95 = Centipeda orbicularis.
dissecta, Boj. ex DC. Prod. v. 372 = Dichrocephala latifolia.
galamensis, Cass. in Dict. Sc. Nat. xix. 307 = maderaspatana !
lanceolata, Poir. Encyc. Suppl. ii. 826 = Eclipta erecta.
latifolia, Lam. ex Poir. l. c. = Dichrocephala latifolia.
maderaspatana, *Vahl, in Bremen, Abh.* ix. (1885) 126.—Madag.
maderaspatana, *Poir. Encyc. Suppl.* ii. 825.—As. et Afr. trop.
minima, Poir. l. c. = Centipeda orbicularis.
minima, *Zipp. ex Span. in Linnæa*, xv. (1841) 922, nomen.—Ins. Timor.
minuta, Poir. Encyc. Suppl. ii. 825 = Centipeda orbicularis.
muscosa, Buch.-Ham. ex Wall. Cat. sub n. 3236 = Cotula hemisphærica.
procumbens, DC. Prod. v. 378 = maderaspatana.
sonchifolia, Loud. Hort. Brit. 354 = Dichrocephala latifolia.
Sphærocladus, C. Koch, in Bot. Zeit. i. (1843) 41, in obs. = maderaspatana.

GRANGERIA, Comm. ex Juss. Gen. 340 (1789). *ROSACEÆ*, Benth. & Hook. f. i. 607.
borbonica, *Lam. Encyc.* iii. 21.—Ins. Mascar.
brasiliensis, Hoffmgg. ex Mart. & Zucc. in Abh. Akad. Münch. ii. (1829-30) 378 = Hirtella ciliata.
lucidifolia, Sm. in Rees, Cyclop. xvi. n. 1 = borbonica.
madagascariensis, *O. Hoffmgg. Sert. Madag.* 15.—Madag.
parœus, *Boiv. ex Baill. Adansonia*, viii. (1868) 200.—Madag.

GRANIERA, Mand. & Weld. ex Benth. & Hook. f. Gen. i. 799 (1867) = **Abatia**, Ruiz & Pav. (Samyd.).

GRANITIA, Boiss. Diagn. Ser. I. vi. 79 (1846). *CONVOLVULACEÆ*, Benth. & Hook. f. ii. 882.
PERRALDERIA, Coss. in Bull. Soc. Bot. Fr. vi. (1859) 294; et in Ann. Sc. Nat. Sér. IV. xviii. (1862) 208. t. 12.
arachnoidea, *Boiss. Diagn.* Ser. I. vi. 79.—Persia.
Auchæri, *Boiss. l. c.*—Persia.
coronopifolia, *Benth. & Hook. f. Gen.* ii. 882.—Algeria.
discoidea, *Bunge, ex Boiss. Fl. Orient.* iii. 214.—Persia.

GRANTIA, Griff. ex Voigt, Hort. Suburb. Calc. 692 (1845); et Notul. iii. 283 (1851) = **Wolffia**, Horkel (Lemnac.).
globosa, Griff. l. c. 229, 699 = W. Michelii.
microscopica, Griff. ii. 22. = W. microscopica.

GRAPHEPHORUM, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189. *GRAMINEÆ*, Benth. & Hook. f. iii. 1197.
DUPONTIA, R. Br. in Parry, 1st Voy. Suppl. 290, (err. typ. 190) (1824).
FLUMINIA, Fries, Summ. Veg. Scand. 247 (1846).
SCOLOCHLOA, Mert. & Koch, in Roehl. Deutschl. Fl. i. 874, 893 (1823).
altijugum, *Fourn. in Bull. Soc. Bot. Fr.* xxiv. (1877) 162.—Mexic.
arundinaceum, *Aschers. Fl. Brandenb.* 852.—Europ.
densiflorum, *Fourn. in Bull. Soc. Bot. Fr.* xxiv. (1877) 182.—Mexic.
Elliottii, *Kunth, Rév. Gram.* i. 80.—Am. bor.
festuceoides, *A. Gray, in Proc. Am. Acad.* v. (1862) 191.—Europ.; Am. bor.
Fischeri, *A. Gray, l. c.*—Reg. Arct.
flexuosum, *Thurb. in Proc. Acad. Sc. Philad.* 1868 (1864) 78.—Am. bor. occ.
fulvum, *A. Gray, in Proc. Am. Acad.* v. (1862) 191. —Europ.; Am. bor.
meliceides, Beauv. Agrost. 77. t. 15. f. 8 = seq.
melicoidum, *Desv. in Nouv. Bull. Soc. Philom.* ii. (1810) 189.—Am. bor.
pendulinum, *A. Gray, in Proc. Am. Acad.* v. (1861) 191.—Europ.; Am. bor.
psilosanthum, *Fourn. in Bull. Soc. Bot. Fr.* xxiv. (1877) 182.—Europ. bor.
Wolfii, *Coult. Man. Rocky Mount. Bot.* 423.—Am. bor. occ.

GRAPHIOSA, Alef. in Bonplandia, ix. (1861) 128 = **Lathyrus**, Tourn. (Legumin.).
inconspicua, Alef. l. c. = L. inconspicuus.

GRAPHISTEMMA, Champ. ex Benth. & Hook. f. Gen. ii. 760 (1876). *ASCLEPIADEÆ*, Benth. & Hook. f. l. c.
pictum, *Champ. (in Herb. Kew.)*.—China.

GRAPTOPHYLLUM, Nees, in Wall. Pl. As. Rar. iii. 76, 102 (1832). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1118.
EARLIA, F. Muell. Fragm. iii. 159 (1863).
MARAMA, Rafin. Fl. Tellur. iv. 69 (1836).
Earlii, *F. Muell. Fragm.* vi. 87.—Austral.
hortense, *Nees, in Wall. Pl. As. Rar.* iii. 102.—As. trop.
ilicifolium, *F. Muell. ex Benth. Fl. Austral.* iv. 559.—Austral.
mediterraneum, Linden & André, ex Linden, Cat. n. 87 (1873) 4 = Aphelandra mediasaxata.
pictum, Griff. Notul. iv. 139 = hortense.
siphonostemum, *F. Muell. Fragm.* vi. 87, in obs.—Ins. Fiji.
spiniferum, *F. Muell. l. c.* xi. 17.—Austral.

GRASTIDIUM, Blume, Bijdr. 888 (1825) = **Dendrobium**, Sw. (Orchid.).
acuminatissimum, Blume, l. c. = D. acuminatissimum.
rugosum, Blume, l. c. = D. rugosum:
salaccense, Blume, l. c. = D. salaccense.

GRATTOLA, [Rupp.] Linn. Gen. ed. I. 288 (1737). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 968.
DIBELINIA, Neraud, in Freyc. Voy. Bot. 98 (1826).
FONKIA, Phil. in Linnæa, xxx. (1859-60) 196.
GONATIA, Nutt. ex DC. Prod. x. 396 (1846).
NIBORA, Rafin. Fl. Ludov. 26 (1817).
SOPHRONANTHE, Benth. in Lindl. Introd. Nat. Syst. ed. II. 445 (1836).
aconitata, Franch. ex Benth. in DC. Prod. x. 403 = virginiana.
acuminata, Vahl, Enum. i. 92 = sphærocarpa.
acuminata, Walt. Fl. Carol. 61 = Herpestis nigrescens.
alata, Komb. Hort. Beng. [80]; Fl. Ind. i. 127 = Torenia alata.
amara, Roxb. ii. 44, 135 = Curanga amara.
anagallidea, Miehx. Fl. Bor. Am. i. 8 = Hyssanthes gratioloides.
Anagallis, Beyr. ex Benth. in DC. Prod. x. 404 = viscosa.
angustifolia, Cliffi. Fl. Linnæ. i. 128 = officinalis.

GRATIOLA:—
aphylla, Roth, Nov. Pl. Sp. 7 = Dopatrium nudicaule.
aromatica, Pers. Syn. i. 14 = Limnophila gratissima.
aspera, Roth, Nov. Pl. Sp. 11 = Vandellia crustacea.
attenuata, Spreng. Syst. i. 39 = Ilysanthes gratioloides.
aurea, *Muhl. Cat. ed.* I. (1818) ; *Le Conte, in Ann. Lyc. N. York*, i. (1824) 106.—Am. bor.
betonicifolia, J. F. Gmel. Syst. 1557, sphalm. = *veronicifolia*.
brachycarpa, Walp. Rep. iv. 440 (Index) sphalm. = Ilysanthes capensis.
brevifolia, *Raftn. Atl. Journ.* 276 (Quid !).—Am. bor.
Broteri, Nym. Consp. 536 = officinalis.
caroliniensis, Le Conte, in Ann. Lyc. N. York, i. (1824) 105 = sphaerocarpa.
caroliniensis, Pursh, Fl. Am. Sept. i. 12, in syn. = virginiana.
cataracta, *Raftn. in Am. Monthly Mag.* (1811) 356, nomen (Quid !).—Am. bor.
chamaedrifolia, Lam. Encyc. iii. 27 = Limnophila gratissima.
Chamaedrys, *Roth, Nov. Pl. Sp.* 10.—Ind. or.
ciliata, Colsm. Prod. Desc. Grat. 14 = Bonnaya veronicaefolia.
ciliata, Koem. ex Vahl, Enum. i. 97 = Bonnaya reptans.
cordifolia, Colsm. Prod. Desc. Grat. 15 ; Vahl, Enum. i. 97 = Vandellia pedunculata.
crassifolia, Lag. Gen. et Sp. Nov. 1 = linifolia.
cuneifolia, Roxb. Hort. Beng. 80 ; Fl. Ind. i. 142 = Dopatrium nudicaule.
debilis, Heyne, in Wall. Cat. n. 8946 = Ilysanthes rotundifolia.
dilatata, Muhl. Cat. ex Spreng. Syst. i. 39 = Ilysanthes gratioloides.
domingensis, Gaertn. f. Fruct. iii. 189 = Ilysanthes gratioloides.
Drummondii, *Benth. in DC. Prod.* x. 405.—Am. bor.
dubia, Linn. Sp. Pl. 17 = Ilysanthes gratioloides.
crucosa, *Benth. ex A. DC. in DC. Prod.* x. 595.—Am. bor. occ.
floridana, *Nutt. in Journ. Acad. Philad.* vii. (1834) 108.—Am. bor.
glabra, Walp. Rep. iii. 287, sphalm. = peruviana.
glandulifera, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 245.—N. Zel.
gnaneifolia, Hornem. Enum. Pl. Hort. Hafn. (1807) 19 = Mazus rugosus.
gracilis, *Benth. in DC. Prod.* x 402.—Texas.
grandiflora, Retz. Obs. iv. 8 ; Roxb. Fl. Corom. ii. 42. t. 179 = Bonnaya veronicaefolia.
grandiflora, Willd. ex G. Don, Gen. Syst. iv. 587 = Bonnaya brachiata.
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 278.—Assam.
hispidula, *Comm. ex Vahl, Enum.* i. 94.—Java.
Hookeri, Walp. Rep. iii. 286 = Stemodia lobelioides.
hyssopifolia, Salisb. Prod. 103 = Ilysanthes hyssopioides.
hyssopioides, Blanco, Fl. Filip. ed. I. 11 = Bonnaya reptans.
hyssopioides, Linn. Mant. ii. 174 = Ilysanthes hyssopioides.
hyssopioides, Lour. Fl. Cochinch. 22 = Morgania hyssopioides.
hyssopioides, Roxb. Fl. Corom. ii. 16. t. 128 = Dopatrium lobelioides.
inaequalis, Walt. Fl. Carol. 61 = Herpestis nigrescens.
integrifolia, Roxb. Hort. Beng. 4 ; Fl. Ind. i. 137 = Vandellia crecta.
invandata, Kit. ex Spreng. Pugill. i. 1 = Vandellia crecta.
involucrata, Roxb. Hort. Beng. [80] ; Fl. Ind. i. 137 = Epithema Roxburghii.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 117.—Japon.
juncea, Roxb. Fl. Corom. ii. 16. t. 129 = Dopatrium junceum.
latifolia, R. Br. Prod. 435 = peruviana.
linearifolia, Herb. Madr. ex Wall. Cat. sub n. 8862 = Bonnaya tenuifolia.
linifolia, Hoffmggy. & Link, Fl. Port. i. 225. t. 31 = officinalis.
lisianthoides, *Vahl, Enum.* i. 96.—Lusitan.
lobelioides, Retz. Obs. iv. 7 = Dopatrium lobelioides.
lucida, Heyne, ex Wall. Cat. sub n. 8935 = Limnophila conferta.
lucida, Willd. Sp. Pl. i. 108 = Vandellia crustacea.
lutea, Reich. in Mod. Repos. N. York, III. ii. (1811) 333 = officinalis.
marginata, Colsm. Prod. Desc. Grat. 11 = Bonnaya veronicaefolia.

GRATIOLA:—
megalocarpa, Ell. Sketch, i. 16 (sp. falsa).
micrantha, *Franch. & Sav. Enum. Pl. Jap.* ii. 456.—Japon.
micrantha, Nutt. in Am. Journ. Sc. v. (1822) 287 = Scoparia dulcis.
minima, Roth. Nov. Pl. Sp. 8 = Bonnaya oppositifolia.
missouriana, Beck, in Am. Journ. Sc. x. (1826) 258 = virginiana.
Monnieri, Linn. Syst. ed. X. 851 = seq.
Mounieri, Linn. Cent. Pl. ii. n. 126 ; Amoen. Acad. iv. 306 = Herpestis Monnieria.
montana, Rottl. ex DC. Prod. ix. 268 = Didymocarpus Rottleriana.
nana, *Benth. in DC. Prod.* x. 404.—Austral.
neglecta, Torr. Cat. Pl. N. York, 89 = virginiana.
Nibora, *Steud. Nom.* ed. II. i. 704.—Am. bor.
nudicaulis, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 198 = Dopatrium nudicaule.
nummularifolia, *Thou. ex Link, Jahrb.* i. III. 51, nomen.—Hab. ?
ocymifolia, Vahl, Enum. i. 96 = Vandellia crecta *vel* multiflora.
officinalis, Brot. Fl. Lusit. i. 15 = linifolia.
officinalis, *Linn. Sp. Pl.* 17.—Europ.
officinalis, Michx. Fl. Bor. Am. i. 6 = virginiana.
oppositifolia, Retz. Obs. iv. 8 ; Roxb. Fl. Corom ii. 30. t. 155 = Bonnaya oppositifolia.
oppositifolia, Willd. ex G. Don, Gen. Syst. iv. 587 = Bonnaya brachiata.
origanifolia, Vahl, Enum. i. 99 = Vandellia diffusa.
parviflora, Roxb. Fl. Corom. iii. 8 = Ilysanthes parviflora.
parviflora, Willd. ex Schlecht. & Cham. in Linnaea, v. (1830) 107 = Herpestes obovata.
pedunculata, *R. Br. Prod.* 435.—Austral.
peruviana, *Linn. Sp. Pl.* 17.—Am. trop. ; Austral. ; N. Zel.
peruviana, Walt. Fl. Carol. 62 = pilosa.
pilosa, *Michx. Fl. Bor. Am.* i. 7.—Am. bor.
portulacacea, Weinm. in Syll. Ratisb. i. (1824) 222 = Herpestes Monnieria.
pubescens, R. Br. Prod. 435 = peruviana.
pulegifolia, Vahl, Enum. i. 98 = Bonnaya veronicaefolia.
pumila, F. Muell. in Linnaea, xxv. (1852) 431 = peruviana.
punctata, Colsm. Prod. Desc. Grat. 7 = Limnophila serrata.
pusilla, *Torr. ex Benth. in DC. Prod.* x. 402.—Am. bor.
pusilla, Willd. Sp. Pl. i. 105 = Vandellia scabra.
quadridentata, Michx. Fl. Bor. Am. i. 6 = ramosa.
racemosa, Roxb. Hort. Beng. 43 ; Fl. Ind. i. 138 ; Roth, Nov. Pl. Sp. 9 = Bonnaya veronicaefolia.
ramosa, Herb. Madr. ex Wall. Cat. sub n. 8867 = Ilysanthes parviflora.
ramosa, *Wall. Fl. Carol.* 61.—Am. bor.
repens, Sw. Prod. Veg. Ind. Occ. 14 = Herpestis repens.
repens, Sweet, ex Steud. Nom. ed. II. i. 704 = micranthemum Nuttallii.
repens, Willd. ex Schlecht. & Cham. in Linnaea, v. (1830) 107 = Herpestes obovata.
reptans, Roxb. Hort. Beng. 4 ; Fl. Ind. i. 140 = Bonnaya reptans.
rigida, *Raftn. Atl. Journ.* 276 (Quid ?).—Am. bor.
riparia, Kaftn. ex Benth. in DC. Prod. x. 419 = Ilysanthes gratioloides.
rostrata, *Thou. ex Link, Jahrb.* i. III. 50, nomen.—Hab. ?
rotundifolia, Linn. Mant. 174 = Ilysanthes rotundifolia.
Roxburghiana, Koem. & Schult. Syst. i. 128 = Bonnaya veronicaefolia.
ruelioides, Colsm. Prod. Desc. Grat. 12 = Bonnaya reptans.
rugosa, *Lour. Fl. Cochinch.* 23.—Cochinch.
salsuginea, Lag. Gen. et Sp. Nov. 1 = linifolia.
serrata, Roxb. Hort. Beng. 4 ; Fl. Ind. i. 140 = Bonnaya brachiata.
sexdentata, *A. Cunn. in Ann. Nat. Hist.* i. (1838) 466.—Austral. ; N. Zel.
sphaerocarpa, *Ell. Sketch*, i. 14.—Am. bor.
stricta, *Lour. Fl. Cochinch.* 23.—Cochinch.
strigosa, *Roth, Nov. Pl. Sp.* 12.—Ind. or.
subulata, *Baldw. ex Benth. in DC. Prod.* x. 405.—Am. bor.
tenuifolia, Colsm. Prod. Desc. Grat. 6 = Bonnaya tenuifolia.

GRATIOLA:—
tetragona, Ell. Sketch, i. 15 = Ilysanthes gratioloides.
tetragona, Hook. Bot. Mag. t. 3184 = Stemodia lobelioides.
tetrandra, Stokes, Bot. Mat. Med. i. 86 = Herpestis Monnieria.
trifida, Vahl, Symb. Bot. Pl. i. 104 = Limnophila gratioloides.
uliginosa, *Phil. f. Cat. Pl. Vasc. Chil.* 289.—Chili.
verbenaefolia, Colsm. Prod. Desc. Grat. 9 = Bonnaya veronicaefolia.
veronicifolia, Retz. Obs. iv. 8 = Bonnaya veronicaefolia.
violacea, *Maxim. in Bull. Acad. Pétersb.* xx (1875) 440.—Japon.
virginiana, *Linn. Sp. Pl.* 17 (Planta virginiana).—Am. bor.
virginiana, Linn. l. c. partim = Limnophila gratioloides.
virginiana, Walt. Fl. Carol. 62 = sphaerocarpa.
virginica, Ell. Sketch, i. 12 = viscosa.
virginica, Lam. Encyc. iii. 27 = ramosa.
virginica, Linn. Syst. ed. XII. 62 = sphaerocarpa.
viscosa, Hornem. Enum. Pl. Hort. Hafn. (1807) 19 = Vandellia hirsuta.
viscosa, *Schwein. ex Le Conte, in Ann. Lyc. N. York*, i. (1824) 106.—Am. bor.

GRAUMUELLERA, Reichb. *Consp.* 43 (1828) = Cymodocea, Kon. (Naiad.).
antarctica, Reichb. l. c. = C. antarctica.

GRAVENHORSTIA, Nees, in Lindl. Introd. Nat. Syst. ed. II. 489 (1836) = Leonhostoma, Wikstr. (Bruniac.).
fastigiata, Nees, l. c. = L. monostylis.

GRAVESIA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 333. *MELASTOMACEAE*, Benth. & Hook. Hook. f. i. 755.
bertoloides, *Naud. l. c.*— Madag.
guttata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 78 —Madag.
pedunculata, *Triana, l. c.*—Madag.

GRAVIA, Steud. Nom. ed. II. i. 705 (1840) = GRAFIA, Reichb. = **Pleurospermum**, Hoffm. (Umbellif.).

GRAYA, Arn. ex Steud. Nom. ed. II. i. 705 (1840) = Andropogon, Linn. (Gramin.).
scylanica, Arn. l. c. = A. Pseudograya.

GRAYA, Endl. Gen. Suppl. i. 1376 (1841) = **Grayia**, Hook. & Arn. (Chenop.).

GRAYA, Nees, ex Steud. Syn. Pl. Gram. 119 (1854) = **Isachne**, R. Br. (Gramin.).
elegans, Nees, l. c. = I. pulchella.

GRAYIA, Hook. & Arn. Bot. Beech. Voy. 387 (1841) *CHENOPODIACEAE*, Benth. & Hook. f. iii. 54.
Brandegii, *A. Gray, in Proc. Am. Acad.* xi. (1876) 101.—Am. bor. occ.
polygaloides, *Hook. & Arn. Bot. Beech. Voy.* 387.—Am. bor. occ.
spinosa, Moq. in DC. Prod. xiii. 11. 119 = polygaloides.

GREENEA, Wight & Arn. Prod. 408 (1834). *RUBIACEAE*, Benth. & Hook. f. ii. 81.
RHOMBOSPORA, Korth. in Nederl. Kruidk. Arch. ii. (1851) 114.
Jackii, *Wight & Arn. Prod.* 404.—Malaya.
latifolia, *Trijtm. & Binn. Cat. Hort. Bog.* 116.—Ins. Amboin.
macrophylla, *Trijtm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 268.—Ins. Moluccc.
tetramera, Miq. Fl. Ind. Bat. ii. 157 = Hedyotis macrophylla.
Wightiana, *Wight & Arn. Prod.* 404.—Burma.

GREENELLA, A. Gray, in Proc. Am. Acad. xvi. (1861) 61. *COMPOSITAE*.
arizonica, *A. Gray, l. c.* 82.—Arizona.
discoidea, *A. Gray, l. c.* xix. (1883) 3.—Calif.

GREENIA, Nutt. in Trans. Am. Phil. Soc. v. (1837) 142 = **Thurberia**, Benth. (Gramin.).
arkansana, Nutt. l. c. = T. arkansana.

GREENOVIA, Webb, in Webb & Berth. Phyt. Canar. i. 198. t. 36 (1836-40) = **Sempervivum**, Rupp. (Crassulac.).
Aizoon, Bolle, in Bonplandia, vii. (1859) 242.—Ins. Canar.
aurea, Webb & Berth. Phyt. Canar. i. 199 = S. aureum.
diplocycla, Webb, ex Bolle, in Bonplandia, vii. (1859) 242.—Ins. Canar.
dodrantalis, Webb & Berth. Phyt. Canar. i. 200 = S. dodrantale.
gracilis, Bolle, in Bonplandia, vii. (1859) 242.—Ins. Canar.

GREENWAYA, Gieske, Fragl. Linn. 199, 206 (1792) = **Amomum**, Linn. (Scitamin.).
Lenaeus, Gieske, ll. cc. = A. Lecouvre.
scyphiferus, Gieske, ll. cc. = A. scyphiferum.

GREEVESIA, F. Muell. in Hook. Kew Journ. viii. (1856) 8 = **Pavonia**, Cav. (Malvac.)
cleiocalyx, F. Muell. l. c. = P. hastata.

GREGGIA, Engelm. in Wisliz. Tour N. Mexic. 114 (1848) = **Cowania**, D. Don (Rosac.).
purpurea, Engelm. ex Hemsl. Biol. Centr. Am. Bot. i. 374 = C. plicata.
rupestris, Engelm. in Wisliz. Tour N. Mexic. 114 = Fallugia paradoxa.

GREGGIA, Gaertn. Fruct. i. 168. t. 35 (1788) = **Eugenia**, Linn. (Myrtac.).
aromatica, Gaertn. l. c. = E. Greggia.

GREGGIA, A. Gray, in Smithson Contrib. iii. (Fl. Wright. i.) 8. t. 1 (1852). *CRUCIFERAE*, Benth. & Hook. f. i. 80.
camporum, *A. Gray, l. c.* 9.—Am. bor. occ.
linearifolia, *S. Wats. in Proc. Am. Acad.* xviii. (1882-83) 191.—Am. bor. occ.
montana, Griseb. in Goett. Abh. xix. (1874) 72 = Draba greggoides.

GREGIA, Carr. in Rev. Hortic. (1880) 255, sphalm. = **Greigia**, Regel (Bromel.).

GREGORIA, Duby [Bot. Gall. i. 383 (1828); et] in DC. Prod. viii. 45 (1844) = **Dionysia**, Fenzl (Primulac.).
arctioides, Duby, l. c. 46 = D. arctioides.
Aucheri, Duby, l. c. = D. odora.
caespitosa, Duby, l. c. = D. caespitose.
listeri. S. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = Douglasia Vitaliana.
Michauxii, Duby, in DC. Prod. viii. 46 = Dionysia Michauxii.
Vitaliana, Duby, Bot. Gall. i. 383 = Douglasia Vitaliana.

GREIGIA, Reg. Ind. Sem. Hort. Petrop. (1864) 13; et Gartenfl. xiv. (1865) 187. t. 474. *BROMELIACEAE*, Benth. & Hook. f. iii. 660.
Landbeckii, *Phil. f. Cat. Pl. Vasc. Chil.* 278.—Chili.
sphacelata, *Regel, Ind. Sem. Hort. Petrop.* (1864) 13; *et Gartenfl.* xiv. (1865) 187.—Chili.

GRENIERA, J. Gay, in Ann. Sc. Nat. Sér. III. iv. (1845) 87 = **Arenaria**, Rupp. (Caryophyll.).
Douglasii, J. Gay, l. c. = A. Douglasii.
tenella, J. Gay, l. c. = A. tenella.

GRENVILLEA, Sweet, Geran. sub t. 262 (1825) = **Pelargonium**, L'Hérit. (Geraniac.).
conspicua, Sweet, l. c. = P. conspicuum.

GRESLANIA, Balansa, in Bull. Soc. Bot. Fr. xix. (1872) 319. *GRAMINEAE*, Benth. & Hook. f. iii. 1213.
circinata, *Balansa, l. c.* 320.—N. Caled.
montana, *Balansa, l. c.*—N. Caled.
rivularis, *Balansa, l. c.*—N. Caled.

GREUIA, Stokes, Bot. Mat. Med. iii. 204 (1812) = **Grewia**, Linn. (Tiliac.).

GREWIA, Baill. in Bull. Soc. Linn. Par. i. (1884) 490. *SAXIFRAGACEAE?* madagascariensis, *Baill. l. c.*—Madag.

GREVIA, Linn. Hort. Upsal. 279 (1748) = **Grewia**, Linn. (Tiliac.).

GREVILLEA, R. Br. in Trans. Linn. Soc. x. (1810) 167. *PROTEACEAE*, Benth. & Hook. f. iii. 180.
ANADENIA, R. Br. in Trans. Linn. Soc. x. (1810) 166.
FITCHIA, Meisn. in Hook. Kew Journ. vii. (1855) 75.
LYSANTHE, Salisb. in Knight, Prot. 116 (1809).
LYSANTHES, Endl. Gen. 840 (1837).
MANGLESIA, Endl. Nov. Stirp. Dec. 25 (1839).
MOLLOYA, Meisn. in Hook. Kew Journ. vii. (1855) 383.
PLAGIOPODA, Spach, Hist. Vég. Phan. x. 402 (1841).
PTYCHOCARPA, Spach, l. c. (1841).
STYRANGEA, Meisn. in Hook. Kew Journ. vii (1855) 66.
STYLOURUS, Salisb. in Knight, Prot. 115 (1809).
acanthifolia, *A. Cunn. in Field, N. S. Wales*, 238.—Austral.
acerosa, *F. Muell. Fragm.* i. 186.—Austral.
aciculariis, A. Cunn. ex Meisn. in DC. Prod. xiv. 364 = diversifolia.
aciculariis, Schult. Mant. iii. 278 = juniperina.
acifolia, Sieber, ex Spreng. Syst. iv. Cur. Post. 46 = juniperina.
aciphylla, Sieber, ex Benth. Fl. Austral. v. 469 = juniperina.
acrobotrya, *Meisn. in Hook. Kew Journ.* vii. (1855) 74.—Austral.
acuaria, *F. Muell. ex Benth. Fl. Austral.* v. 453.—Austral.
acuminata, R. Br. in Trans. Linn. Soc. x. (1810) 178 = mucronulata.
acuminato, A. Cunn. ex Meisn. in DC. Prod. xiv. 358 = agrifolia.
agrifolia, *A. Cunn. ex R. Br. Prot. Nov.* 24.—Austral.
alpestris, Meisn. in Hook. Kew Journ. iv. (1852) 187 = alpina.
Alphonsiana, *F. Muell. l. c.* lx. (1857) 22.—Austral.
alpina, *Lindl. in Mitch. Three Exped. Austral.* ii. 179.—Austral.
amplexans, *F. Muell. ex Benth. Fl. Austral.* v. 488.—Austral.
amplifica, F. Muell. ex Meisn. in DC. Prod. xiv. 355 = australis.
anethifolia, *R. Br. Prot. Nov.* 21.—Austral.
angulata, *R. Br. l. c.* 24.—Austral.
angustata, R. Br. l. c. 24 = polystachya.
annulifera, *F. Muell. Fragm.* iv. 55.—Austral.
apiciloba, *F. Muell. l. c.* x. 45.—Austral.
Aquifolium, *Lindl. in Mitch. Three Exped. Austral.* ii. 178.—Austral.
Aquifolium, Meisn. in Lehm. Pl. Preiss. i. 51 = monticola.
arenaria, *R. Br. in Trans. Linn. Soc.* x. (1810) 179.—Austral.
argyrophylla, *Meisn. in Hook. Kew Journ.* vii. (1855) 75.—Austral.
armigera, *Meisn. l. c.* iv. (1852) 186.—Austral.
asparagoides, *Meisn. l. c.*—Austral.
aspera, *R. Br. in Trans. Linn. Soc.* x. (1810) 172.—Austral.
asplenifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 176.—Austral.
asplenifolia, *Knight & Salisb. Prot.* 120 (= prasc. ?).—Austral.
attenuata, A. Cunn. ex Meisn. in DC. Prod. xiv. 358.
australis, *R. Br. in Trans. Linn. Soc.* x. (1810) 171.—Austral.
autumnalis, Lhotsk. ex Benth. Fl. Austral. v. 440 = floribunda.
Baxteri, *R. Br. in Trans. Linn. Soc.* x. (1810) 176.—Austral.
Barklyana, *F. Muell. ex Benth. Fl. Austral.* v. 486.—Austral.
Baueri, *R. Br. in Trans. Linn. Soc.* x. (1810) 173.—Austral.
Baueri, *R. Br. Prot. Nov.* 23.—Austral.
Behrii, Schlecht. in Linnaea, xx. (1847) 585 = lilicifolia.
berberifolia, *Sweet, Hort. Brit.* ed. II. 446, nomen.—Austral.
biformis, Meisn. in Lehm. Pl. Preiss. ii. 258 = stenocarpa.
biglandulifera, *R. Br. Prot. Nov.* 23.—Austral.
biternata, *Meisn. in Lehm. Pl. Preiss.* i. 249.—Austral.

GREVILLEA:—
Blandaleti, F. Muell. Fragm. v. 90 = Rompala Blandaleti.
blechnifolia, A. Cunn. ex Benth. Fl. Austral. v. 485 = Caleyi.
brachyantha, Lindl. Swan River App. 81 = quercifolia.
brachystachya, *Meisn. in Lehm. Pl. Preiss.* ii. 254.—Austral.
brachystylis, *Meisn. l. c.* i. 238.—Austral.
bracteosa, *Meisn. l. c.*—Austral.
brevicuspis, *Meisn. l. c.* 256.—Austral.
brevifolia, *F. Muell. ex Benth. Fl. Austral.* v. 467.—Austral.
Brownei, *Meisn. in Lehm. Pl. Preiss.* i. 237.—Austral.
buxifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 174.—Austral.
Caleyi, *R. Br. Prot. Nov.* 22.—Austral.
callisteris, Meisn. in Hook. Kew Journ. iv. (1852) 187 = Dryandri.
Candollesiana, *Meisn. in Lehm. Pl. Preiss.* ii. 241.—Austral.
canescens, R. Br. Prot. Nov. 18 = arenaria.
capillaris, Meisn. in DC. Prod. xiv. 356 = didymobotrya.
capitellata, *Meisn. in Hook. Kew Journ.* iv. (1852) 187.—Austral.
cardwellia, A. Cunn. ex Benth. Fl. Austral. v. 456 = Cunninghamii.
ceratophylla, R. Br. in Trans. Linn. Soc. x. (1810) 177 = refracta.
Chrysodendron, Auct. = seq.
Chrysodendrum, *R. Br. l. c.* 176.—Austral.
chrysophaea, F. Muell. First Gen. Rep. 17 = floribunda.
cinerea, *R. Br. in Trans. Linn. Soc.* x. (1810) 173.—Austral.
cinerea, A. Cunn. in Field, N. S. Wales, 239 = arenaria.
cinerea, Lodd. Bot. Cab. t. 857 = mucronulata.
circiifolia, *Meisn. in Lehm. Pl. Preiss.* ii. 253.—Austral.
coccinea, Meisn. in Hook. Kew Journ. viii. (1856) 76 = concinna.
collina, Knight, ex Sweet, Hort. Brit. ed. II. 446 = buxifolia.
commutata, *F. Muell. Fragm.* vi. 207.—Austral.
concinna, *R. Br. in Trans. Linn. Soc.* x. (1810) 172.—Austral.
confertifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 93.—Austral.
crithmifolia, *R. Br. Prot. Nov.* 23.—Austral.
ctenophylla, Meisn. in Hook. Kew Journ. iv. (1852) 186 = pectinata.
Cunninghamii, *R. Br. Prot. Nov.* 23.—Austral.
cynanchicarpa, *Meisn. in Hook. Kew Journ.* vii. (1855) 75.—Austral.
Dallachiana, F. Muell. First Gen. Rep. 17 = alpina.
daphnoides, Sieber, ex Schult. Mant. iii. 281 = Baueri.
dehnxa, *F. Muell. Census*, 140, nomen.—Australia.
depauperata, R. Br. Prot. Nov. 21 = Brownii.
Depiancheii, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1866) 39.—N. Caled.
didymobotrya, *Meisn. in Hook. Kew Journ.* iv. (1852) 186.—Austral.
diffusa, A. Cunn. ex Meisn. in DC. Prod. xiv. 356 = capitellata.
diffusa, Sieber, ex Meisn. l. c. 356 = serices.
dimidiata, *F. Muell. Fragm.* iii. 146.—Austral.
dimorpha, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 21 = oleoides.
dispneata, *F. Muell. Fragm.* vi. 206.—Australia.
divaricata, *R. Br. Prot. Nov.* 20.—Austral.
diversifolia, *Meisn. in Lehm. Pl. Preiss.* i. 247.—Austral.
Drummondii, *Meisn. l. c.* 236.—Austral.
Dryandri, *R. Br. in Trans. Linn. Soc.* x. (1810) 175.—Austral.
dubia, R. Br. l. c. 169 = serices.
elaeagnifolia, Meisn. in Hook. Kew Journ. iv. (1852) 187 = lilicifolia.
Endlicheri, *F. Muell. Papuan. Pl.* ii. 9.—N. Guin.
Endlicherianа, *Meisn. in Lehm. Pl. Preiss.* i. 246.—Austral.
erectiflora, *F. Muell. Fragm.* v. 44.—Austral.
ericifolia, *R. Br. Prot. Nov.* 20.—Austral.
erinacea, *Meisn. in Hook. Kew Journ.* vii. (1855) 74.—Austral.

GREVILLEA :—
eriobotrys, *F. Muell. Fragm.* x. 44.—Austral.
eriostachys, *Lindl. Swan River App.* 86.—Austral.
eryngioides, *Benth. Fl. Australe.* v. 476.—Australia.
Exul, *Lindl. in Journ. Hort. Soc.* vi. (1851) 278.—N. Caled.
fasciculata, *R. Br. Prot. Nov.* 20.—Austral.
ferruginea, R. Grah. in Mannd, Botanist, t. 158 = floribunda.
ferruginea, Sieber, ex Schult. Mant. iii. 280 = montana.
filifolia, *Meisn. in Lehm. Pl. Preiss.* i. 547.—Austral.
flexuosa, *Meisn. l. c.* 553.—Austral.
Flindersii, *A. Cunn. ex Steud. Nom.* ed. II. i. 705.—Austral.
floribunda, *R. Br. Prot. Nov.* 19.—Austral.
Forsteri, *Hort.—Cf. Gard. Chron.* (1874) 60; (1881) 11. 336.—Hab.?
Gaudichaudii, *R. Br. ex Gaudich. in Freyc. Voy. Bot.* 448. t. 46.—Austral.
gibbosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 177.—Austral.
glabella, R. Br. Prot. Nov. 20 = divaricata.
glabrata, *Meisn. in Lehm. Pl. Preiss.* i. 549.—Austral.
glauca, Knight, Prot. 121 = gibbosa.
Goodii, *R. Br. in Trans. Linn. Soc.* x. (1810) 174.—Austral.
hakeoides, *Meisn.in Lehm. Pl. Preiss.* ii. 252.—Austral.
halmaturina, Tate, in Trans. Roy. Soc. S. Austral. vi. (1883) 42 = parviflora.
haplantha, *F. Muell. ex Benth. Fl. Australe.* v. 451.—Austral.
heliosperma, *R. Br. in Trans. Linn. Soc.* x. (1810) 176.—Austral.
heterochroma, *Brongn. in Bull. Soc. Bot. Fr.* xii. (1865) 40.—N. Caled.
heterophylla, A. Cunn. ex R. Br. Prot. Nov. 24 = refracta.
Hewardiana, Meisn. in Hook. Kew Journ. iv. (1852) 186 = concinna.
Hilliana, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 72.—Austral.
Hookeriana, *Meisn. in Lehm. Pl. Preiss.* i. 546.—Austral.
Hugelii, *Meisn. l. c.* 548.—Austral.
humifusa, A. Cunn. ex Benth. Fl. Austral. v. 436 = laurifolia.
ilicifolia, *R. Br. Prot. Nov.* 21.—Austral.
ilicifolia, A. Cunn. ex Meisn. in DC. Prod. xiv. 380 = angulata.
indusia, F. Muell. First Gen. Rep. 17 = G. Aquifolium.
insignis, *Kippist, ex Meisn. in Hook. Kew Journ.* vii (1855) 76.—Austral.
integrifolia, *Meisn. in DC. Prod.* xiv. 385.—Austral.
intricata, *Meisn. in Hook. Kew Journ.* vii. (1855) 74.—Austral.
juncea, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 87, sphalm. = acq.
juncifolia, *Hook. in Mitch. Journ. Trop. Austral.* 341.—Austral.
juniperina, *R. Br. in Trans. Linn. Soc.* x. (1810) 174.—Austral.
lanigera, *A. Cunn. ex Meisen. in DC. Prod.* xiv. 368.—Austral.
Latrobei, Meisn. in Lehm. Pl. Preiss. i. 539 = ericifolia.
laurifolia, *Sieber, ex Meisen. in DC. Prod.* xiv. 352.—Austral.
lavandulacea, *Schlecht. in Linnaea*, xx. (1847) 586.—Austral.
Lawrenciana, *Hort. ex Regel, Cat. Pl. Hort. Absehov.* 65.—Austral.
leiophylla, *F. Muell. ex Benth. Fl. Austral.* v. 471.—Austral.
Leucanniana, Meisn. in Hook. Kew Journ. v. (1852) 187 = concinna.
leptobotrys, *Meisen. in Lehm. Pl. Preiss.* ii. 256.—Austral.
Leucadendron, *A. Cunn. ex R. Br. Prot. Nov.* 25.—Austral.
leucopteris, *Meisen. in Hook. Kew Journ.* vii. (1855) 76.—Austral.
Lindleyana, Meisn. in Lehm. Pl. Preiss. i. 542 = Whitei.
linearis, *R. Br. in Trans. Linn. Soc.* x. (1810) 170.—Austral.
linearis, Sieber, ex Meisn. in DC. Prod. xiv. 356 = ilicifolia.

GREVILLEA :—
linearis, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 87 = striata.
lobata, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 22 = ilicifolia.
longifolia, R. Br. Prot. Nov. 22 = saplenifolia.
longifolia, F. Muell. Fragm. i. 136 = G. Leucadendron.
longistyla, *Hook. in Mitch. Journ. Trop. Austral.* 343.—Austral.
lorea, R. Br. in Trans. Linn. Soc. x. (1810) 177 = Hakea lorea.
macrostachya, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 36.—N. Caled.
macrostylis, *F. Muell. Fragm.* i. 137.—Austral.
Mangleti, Hort. ex Benth. Fl. Austral. v. 436 = glabrata.
manglesioides, *Meisn. in Lehm. Pl. Preiss.* i. 547.—Austral.
Martinii, F. Muell. Fragm. iv. 139 = polybotrya.
Meissneri, *Montr. in Mém. Acad. Lyon*, x. (1860) 248.—N. Caled.
Meisneriana, F. Muell. in Linnaea, xxvi. (1853–55) 857 = fasciculata.
micrantha, Meisn. l. c. 358 = parviflora.
mimosoides, *R. Br. in Trans. Linn. Soc.* x. (1810) 177.—Austral.
Miquelliana, *F. Muell. in Trans. Vict. Inst.* i. (1855) 129.—Austral.
Mitchelli, Hook. in Mitch. Journ. Trop. Austral. 265 = G. Chrysodendrum.
montana, *R. Br. in Trans. Linn. Soc.* x. (1810) 172.—Austral.
monticola, *Meisen. in Lehm. Pl. Preiss.* ii. 259.—Austral.
mucronifolia, *A. Cunn. ex Steud. Nom.* ed. II. i. 705.—Austral.
mucronulata, *R. Br. in Trans. Linn. Soc.* x. (1810) 178.—Austral.
Mueller, *Benth. Fl. Austral.* v. 479.—Austral.
myrtacea, Sieber, ex Schult. Mant. iii. 280 = mucronulata.
myrtillifolia, A. Cunn. ex Benth. Fl. Austral. v. 444 = Banksii.
negiecta, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 87 = venusta.
nematophylla, *F. Muell. Fragm.* i. 136.—Austral.
neriifolia, *Meisen. in Hook. Kew Journ.* iv. (1852) 186.—Austral.
nudana, Meisn. l. c. 187 = divaricata.
obtusa, R. Br. Prot. Nov. 25 = G. Leucadendron.
obtusifolia, *R. Br. l. c.* 19.—Austral.
obtusifolia, *Meisen. in Hook. Kew Journ.* iv. (1852) 187.—Austral.
occidentalis, *R. Br. in Trans. Linn. Soc.* x. (1810) 178.—Austral.
oleoides, *Sieber, ex Schult. Mant.* iii. 277.—Austral.
oligantha, F. Muell. Fragm. vi. 206 = paniculata.
orophila, F. Muell. First Gen. Rep. 17 = alpina.
ornithopoda, *Meisen. in Lehm. Pl. Preiss.* ii. 256.—Austral.
oxytigma, *Meisen. l. c.* t. 540.—Austral.
paniculata, *Meisen. l. c.* 550.—Australia.
paradoxa, *F. Muell. Fragm.* vi. 246.—Australia.
parallela, Knight, Prot. 121 = polystachya.
parviflora, *R. Br. in Trans. Linn. Soc.* x. (1810) 171.—Austral.
parviflora, F. Muell. First Gen. Rep. 17 = micrantha.
patentiloba, *F. Muell. Fragm.* i. 137.—Austral.
patuliflora, *R. Br. in Trans. Linn. Soc.* x. (1810) 171.—Austral.
pectinata, *R. Br. Prot. Nov.* 23.—Austral.
pedunculosa, F. Muell. Fragm. i. 135 = sudiflora.
petrophiloides, *Meisen. in Lehm. Pl. Preiss.* ii. 257.—Austral.
phylicoides, *R. Br. in Trans. Linn. Soc.* x. (1810) 174.—Austral.
phyllocalyx, A. Cunn. ex Meisn. in DC. Prod. xiv. 369 = sphacelata.
Pinaster, *Meisen. in Hook. Kew Journ.* vii. (1855) 76.—Austral.
pinifolia, Hook. ex Meisn. in DC. Prod. xiv. 383 = G. Chrysodendrum.
pinifolia, *Meisen. in Hook. Kew Journ.* iv. (1852) 186.—Austral.
pinnatisecta, *F. Muell. ex Benth. Fl. Austral.* v. 478.—Australia.
platyophylla, F. Muell. Fragm. vi. 208 = striostachya.
plumifolia, Lodd. Bot. Cab. t. 1737 = sericea.

GREVILLEA :—
platypoda, *F. Muell. Fragm.* vi. 205.—Austral.
plurijuga, *F. Muell. l. c.* iv. 84.—Austral.
podalyriaefolia, Sweet, Hort. Brit. ed. II. 446 = mucronulata.
podocarpifolia, *Sweet, l. c.* nomen.—Austral.
polybotrya, *Meisen. in Hook. Kew Journ.* iv. (1852) 185.—Austral.
polybotrya, F. Muell. l. c. ix. (1857) 33 = polystachya.
polystachya, *R. Br. in Trans. Linn. Soc.* x. (1810) 177.—Austral.
Preissei, *Meisen. in Lehm. Pl. Preiss.* i. 543.—Austral.
pteridifolia, Knight, Prot. 121 = G. Chrysodendrum.
pterosperma, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 22.—Australia.
pubescens, *R. Grah. in Edinb. N. Phil. Journ.* i. (1826) 172.—Austral.
pubescens, Hook. Exot. Fl. t. 216 = Baueri.
pulchella, *Meisen. in Lehm. Pl. Preiss.* i. 558.—Austral.
pungens, *R. Br. in Trans. Linn. Soc.* x. (1810) 175.—Austral.
punices, *R. Br. l. c.* 169.—Austral.
pyramidalis, *A. Cunn. ex Meisn. in DC. Prod.* xiv. 381.—Austral.
pyriformis, Hort. ex Meisn. l. c. 393, 422 = Xylomelum pyriforme.
quercifolia, *R. Br. Prot. Nov.* 23.—Austral.
ramosissima, *Meisen. in DC. Prod.* xiv. 388.—Austral.
ramulosa, F. Muell. l. c. 362 = lavandulacea.
refracta, *R. Br. in Trans. Linn. Soc.* x. (1810) 176.—Austral.
repens, *F. Muell. in Linnaea*, xxvi. (1853–55) 355.—Austral.
rigens, A. Cunn. ex Meisn. in DC. Prod. xiv. 374 = Dryandri.
rigidissima, F. Muell ex Meisn. l. c. 350 = Hugelii.
riparia, R. Br. in Trans. Linn. Soc. x. (1810) 170 = linearis.
riparia, Sieber, ex Schult. Mant. iii. 278 = rosmarinifolia robusta, *A. Cunn. in R. Br. Prot. Nov.* 24.—Austral.
rosea, Lindl. in Paxt. Flow. Gard. ii. (1852) 91. n. 386. t. 56 = lavandulacea.
rosmarinifolia, *A. Cunn. in Field, N. S. Wales*, 393.—Austral.
rubiginosa, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 40.—N. Caled.
rudis, *Meisen. in Hook. Kew Journ.* vii. (1855) 73.—Austral.
saccata, *Benth. Fl. Austral.* v. 450.—Austral.
scabra, *Meisn. in Lehm. Pl. Preiss.* i. 541.—Austral.
scabrella, Meisn. in Hook. Kew Journ. iv. (1852) 187 = ericifolia.
segmentosa, F. Muell. Fragm. iii. 145 = leiocpteris.
serices, *R. Br. in Trans. Linn. Soc.* x. (1810) 170.—Austral.
sericea, Lhotsky, ex Meisn. in DC. Prod. xiv. 369 = sphacelata.
sericea, Lodd. Bot. Cab. t. 850 = punices.
sericeistachya, Meisn. in Hook. Kew Journ. iv. (1852) 186 = pteropetma.
Seymourias, Sweet, ex Meisn. in DC. Prod. xiv. 354 = oleoides.
Shuttleworthiana, *Meisen. in Lehm. Pl. Preiss.* ii. 258.—Austral.
silaufolia, Donn, Hort. Cantab. ed. IV. 26 = Lomatia silaifolia.
singuliflora, *F. Muell. Fragm.* vi. 92.—Austral.
sinuata, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 40.—N. Caled.
spariflora, *F. Muell. Fragm.* vi. 206.—Austral.
sphacelata, *R. Br. in Trans. Linn. Soc.* x. (1810) 174.—Austral.
sphacelata, A. Cunn. ex Benth. Fl. Austral. v. 440 = floribunda.
stenobotrys, *F. Muell. Fragm.* ix. 3.—Austral.
stenocarpa, *F. Muell. ex Benth. Fl. Austral.* v. 485.—Austral.
stenomera, *F. Muell. Fragm.* iv. 35.—Austral.
Sternbergiana, Hort. ex Benth. Fl. Australia. v. 477 = crithmifolia.
Straugea, *Benth. l. c.* 459.—Austral.
striata, *R. Br. in Trans. Linn. Soc.* x. (1810) 177.—Austral.
stricta, R. Br. l. c. 170 = linearis.
Stuartii, Meisen. in Linnaea, xxvi. (1853) 357 = eastrala.
Sturtii, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 86 = juncifolia.

GREVILLEA :—
Stuartii, A. Cunn. ex Meisn. in DC. Prod. xv. 379 = illicifolia.
stylosa, Knight & Salisb. ex Loud. Encyc. Pl. ed. I. 84 = cineres ?
sulphurea, A. Cunn. in Field, N. S. Wales, 329 = juniperina.
Synaphaea, *R. Br. Prot. Nov.* 23.—Austral.
tenuiflora, *Meisn. in Lehm. Pl. Preiss.* i. 554.—Austral.
tenuifolia, R. Br. in Trans. Linn. Soc. x. (1810) 171 = sautralis.
tenuifolia, *Meisn. in Lehm. Pl. Preiss.* ii. 255.—Austral.
tetragonoloba, Meisn. in Hook. Kew Journ. iv. (1852) 186 = Hookeriana.
Thelemanniana, R. Br. ex Meisn. in DC. Prod. xiv. 373 = Wilsoni.
Thelemanniana, *Hueg. in Endl. Nov. Stirp. Decad.* i. 6.—Austral.
thyrsoides, *Meisn. in Hook. Kew Journ.* vii. (1855) 77.—Austral.
trachytheca, *F. Muell. Fragm.* vi. 207.—Austral.
Treveriana, *F. Muell. l. c.* ix. 128.—Austral.
tridentifera, *Meisn. in Lehm. Pl. Preiss.* i. 549.—Austral.
trifida, *Meisn. l. c.* 553.—Austral.
trifurcata, *Sweet, Hort. Brit.* ed. II. 446, nomen.—Austral.
trilobe, *Meisn. in Hook. Kew Journ.* vii. (1855) 74.—Austral.
trinervis, *R. Br. Prot. Nov.* 18.—Austral.
trinervia, F. Muell. Fragm. iii. 146 = Hakea trinerva.
tripartita, *Meisn. in Hook. Kew Journ.* iv. (1852) 186.—Austral.
triscela, F. Muell. First Gen. Rep. 17 = Hakea purpurea.
trifernata, *R. Br. Prot. Nov.* 21.—Austral.
umbelliflata, *Meisn. in Lehm. Pl. Preiss.* ii. 252.—Austral.
umbratica, A. Cunn. ex Meisn. in DC. Prod. xiv. 381 = robusta.
uniflora, *A. Cunn. ex Meisn. l. c.* 358.—Austral.
variabilis, Lindl. in Mitch. Three Exped. Austral. ii. 179 = G. Aquifolium.
velutina, A. Cunn. ex Benth. Fl. Austral. v. 458 = refracta.
venusta, *R. Br. in Trans. Linn. Soc.* x. (1810) 176.—Austral.
venusta, A. Cunn. ex Meisn. in DC. Prod. xiv. 381 = robusta.
vestita, *Meisn. in Lehm. Pl. Preiss.* i. 548.—Austral.
Victoriae, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 107.—Austral.
Vieillardi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 41.—N. Caled.
Wickhamii, *Meisn. in Hook. Kew Journ.* iv. (1852) 187.—Austral.
Wilsoni, *A. Cunn. in Wilson, Voy. Round World* (1835) 278.—Austral.

GREVILLIA, Knight, Prot. 130 (1809) = prec.

GREWIA, Linn. Syst. ed. I. (1735). *TILIACEAE*, Benth. & Hook. f. i. 283.
Arsis, Lour. Fl. Cochinch. 385 (1790).
Balmeda, Nocca, Hort. Ticin. 21 (1804).
Chadara, Forsk. Fl. Aegypt. Arab. 105 (1775).
Fallopia, Lour. Fl. Cochinch. 385 (1790).
Grarvia, Neck. Elem. ii. 358 (1790).
Mallococca, Forst. Char. Gen. 77. t. 39 (1776).
Microcos, Burm. ex Linn. Disc. Diss. 20 (1747); Amoen. Acad. i. 130 (1749).
Omphacarpus, Korth. Verh. Nat. Gesch. Bot. 192. t. 43 (1847 ?).
Sabali, Adans. Fam. ii. 305 (1763).
Triderma, Rafin. Sylva Tellur. 146 (1838).
Vincentia, Boj. in Hook. Bot. Misc. i. (1830) 293. t. 62.
Vinticena, Steud. Nom. ed. II. ii. 709 (1841).
Vitigena, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 87.
abutilifolia, *Vent. ex Juss. in Ann. Mus. Par.* iv. (1804) 93.—Ind. or.; Malaya.
acuminata, *Bedd. in Madr. Journ. Sc. Ser. III.* i. (1864) 88.—Ind. or.
acuminata, Juss. in Ann. Mus. Par. iv. (1804) 91 = scabrida.

GREWIA :—
afzelii, Hasak. Cat. Hort. Bog. Alt. 207 = paniculata.
africa, Lindl. in Trans. Hort. Soc. vi. (1826) 266 = G. Microcos.
africana, *Mast. in Oliver, Fl. Trop. Afr.* i. 253.—Afr. trop.
africana, Mill. Gard. Dict. ed. VIII. n. 2 = mollis ?
Amicorum, *Steud. Nom.* ed. II. i. 706.—Ins. Amicor.
ancolana, *Miq. Pl. Jungh.* 301.—Sumatra.
angolana, *Welw. ex Mast. in Oliver, Fl. Trop. Afr.* i. 251.—Afr. trop.
angustifolia, Wall. Cat. n. 1106 = polygama.
apetala, *Juss. in Ann. Mus. Par.* iv. (1804) 98. t. 49. f. 3.—Java.
Araria, Buch.-Ham. in Wall. Cat. n. 6308 = salvifolia.
arborea, Lam. Encyc. iii. 45 = excelsa.
arborea, Roxb. ex Rottl. in Ges. Nat. Fr. Neue Schr. iv. (1803) 205 = tiliaefolia.
arbutifolia, Pers. Syn. ii. 66 = abutilifolia.
asiatica, *Linn. Mant.* i. 122.—Ind. or.
asiatica, Wall. Cat. n. 1089 partim = abutilifolia.
aspera, Roxb. Hort. Beng. 42; Fl. Ind. ii. 591 = abutilifolia.
begoniifolia, *Roxb. ll. cc.* [98]; 592.—Ins. Amboin.
begoniifolia, Wall. Cat. p. 257 n. 1098 = G. Microcos.
betulaefolia, Juss. in Ann. Mus. Par. iv. (1804) 92 = populifolia.
bicolor, Juss. l. c. 90 = salvifolia.
bicolor, Roth, Nov. Pl. Sp. 240 = excelsa.
biflora, G. Don, Gen. Syst. i. 548 = carpinifolia.
bilobe, *Don, l. c.* 548.—China.
biloba, Wall. Cat. n. 1107 = polygama.
bilocularis, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 504.—Ins. Socotra.
bipartita, Wall. Cat. p. 287 n. 1113 B = pilosa.
Blumei, Hasak. in Hoev. & De Vriese, Tijdschr. xii. (1846) 130 = paniculata.
Blumei, *Steud. Nom.* ed. II. i. 706.—Java.
bracteata, Miq. ex Mast. in Hook. f. Fl. Brit. Ind. i. 383 = columnaris.
bractenea, *Roth, Nov. Pl. Sp.* 243.—Ind. or.
breviflora, *Benth. Fl. Austral.* i. 270.—Austral.
bridelisefolia, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 544.—Madag.
calfra, *Linn. in Hook. Lond. Journ. Bot.* ii. (1843) 53.—Afr. austr.
calophylla, *Kurz, in Andaman Report App.* B. iii.; *ex Mast. in Hook. f. Fl. Brit. Ind.* i. 392.—Ins. Andaman.
calvata, *Baker, in Journ. Bot.* xx. (1882) 48.—Madag.
cana, *Sond. in Linnaea*, xxiii. (1850) 20.—Afr. austr.
canescens, *A. Rich. Tent. Fl. Abyss.* i. 86.—Abyss.
carpinifolia, *Juss. in Ann. Mus. Par.* iv. (1804) 91. t. 51. f. 1.—Ind. or.; Afr. trop.
carpinifolia, Roth, Nov. Pl. Sp. 245, partim = emarginata.
carpinifolia, Roxb. Hort. Beng. 43; Fl. Ind. ii. 587 = pilosa.
carpinifolia, Wall. Cat. n. 1093 = columnaris.
Carrea, Bach.-Ham. in Wall. l. c. n. 1095 B = scabrophylla.
caudata, Wall. l. c. n. 1089 = Leptonychia glabra.
celastrifolia, *Juss. in Ann. Mus. Par.* iv. (1804) 93.—Java.
Chadara, Lam. Encyc. iii. 44 = populifolia.
cinerea, A. Rich. Tent. Fl. Abyss. i. 86 = salvifolia.
columnaris, *Sm. in Rees, Cyclop.* xvii. n. 5.—Ind. or.; Malaya; Afr. trop.
communis, DC. Prod. i. 511 = pilosa.
coriacea, Cambess. ex Mast. in Hook. f. Fl. Brit. Ind. i. 393 = mollifera.
coriacea, *Mast. in Oliver, Fl. Trop. Afr.* i. 252.—Afr. trop.
corylifolia, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 96. t. 20 = villosa.
crenata, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 248 = populifolia.
cubensis, *Urban, in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 282.—Cuba.
Cumingiana, *Turcz. l. c.* 281.—Ins. Philipp.
cuneifolia, *Baker, in Journ. Linn. Soc.* xxi. (1884) 236.—Madag.
cuneifolia, *Juss. in Ann. Mus. Par.* iv. (1804) 90.—Madag.
cyclopetala, *Wawra & Peyr. in Sitzb. Acad. Wien, Math. Nat.* xxxvIII. (1860) 556.—Afr. trop.

GREWIA :—
Dawnie, Gaertn. Fruct. ii. 113 = laevigata, salvifolia, Alangium Lamarckii.
denticulata, *Wall. ex Voigt, Hort. Suburb. Calc.* 128, nomen.—Ind. or.
didyma, Roxb. Hort. Beng. [98]; Fl. Ind. ii. 591 = laevigata.
diplocarpa, *Thw. Enum. Pl. Zeyl.* 31.—Zeylan.
discolor, Fresen. in Mus. Senckemb. ii. (1837) 159 = salvifolia.
disperma, Rottl. ex Spreng. Syst. ii. 579 = laevigata.
echimulata, Delile, Cent. Pl. Méroé, 82 = villosa.
ectasicarpa, *S. Moore, in Journ. Bot. xv.* (1877) 67.—Afr. trop.
elastica, Royle, Illustr. Bot. Himal. 104. t. 22 = asiatica.
emarginata, *Wight & Arn. Prod.* 79.—Ind. or.
ericarpa, *Juss. in Ann. Mus. Par.* iv. (1804) 98.—Java.
eriopoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 281.—Ins. Philipp.
Erythraea, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xvii. (1868) 671.—Afr. trop.
excelsa, *Vahl, Symb. Bot.* 35.—As. et Afr. trop.
ferruginea, Hochst. in Flora, xxiv. (1841) i. Intell. 29, nomen; et ex A. Rich. Tent. Fl. Abyss. i. 87 = scabrida.
fibrocarpa, *Mast. in Hook. f. Fl. Brit. Ind.* i. 391.—Malacca.
flava, *DC. Cat. Hort. Monsp.* 113.—Afr. austr.
flavescens, Juss. in Ann. Mus. Par. iv. (1804) 91 = pilosa.
floribunda, *Mast. in Oliver, Fl. Trop. Afr.* i. 252.—Afr. trop.
floribunda, Wall. ex Voigt, Hort. Suburb. Calc. 128 = Columbia floribunda.
florida, *Miq. Fl. Ind. Bat. Suppl.* 405.—Sumatra.
Forbesii, *Harv. ex Mast. in Oliver, Fl. Trop. Afr.* i. 250.—Afr. trop.
furfuracea, Salisb. Prod. 367 = orientalis.
glabra, Blume, Bijdr. 118 = laevigata.
glabra, Mast. in Hook. f. Fl. Brit. Ind. i. 393, in syn. sphalm. = G. Microcos.
glabrescens, *Benth. Fl. Hongk.* 42.—Ins. Hongkong.
glandulosa, *Vahl, Symb. Bot.* i. 34.—Ins. Maurit.
globuliflora, *Mast. in Hook. f. Fl. Brit. Ind.* i. 391.—Malacca.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 544.—Madag.
grandiflora, *Baker, in Journ. Bot.* xx. (1882) 47.—Madag.
Grevei, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 543.—Madag.
guazumaefolia, *Juss. in Ann. Mus. Par.* iv. (1804) 89. t. 48. f. 3.—Java.
helicterifolia, Wall. Cat. n. 1090 C = hirsuta.
helicterifolia, Wall. l. c. n. 1090, partim = polygama.
hermannioides, *Harv. in Harv. & Sond. Fl. Cap.* i. 226.—Afr. austr.
heterochita, Roxb. Hort. Beng. [98]; Fl. Ind. ii. 590 = Leptonychia glabra.
heterophylla, *A. Rich. Tent. Fl. Abyss.* i. 90.—Abyssin.
heterotricha, *Mast. in Hook. f. Fl. Brit. Ind.* i. 385.—ined. or.
Heynei, Steud. Nom. ed. II. i. 706 = villifera.
hirsuta, Roxb. Hort. Beng. [98]; Fl. Ind. ii. 587 = helicterifolia.
hirsuta, Sm. in Rees, Cycl. xvii. n. 9 = tomentosa.
hirsuta, *Vahl, Symb. Bot.* i. 34.—As. trop.
hirsuta, Wall. Cat. p. 287. n. 1090 C = polygama.
hispida, *Harv. in Harv. & Sond. Fl. Cap.* i. 226.—Afr. austr.
humilis, *Wall. Cat.* n. 1110.—Burma.
inaequalis, Blume, Bijdr. 118 = villaefolia.
inaequilatera, *Garcke, in Peters, Reise Mossamb. Bot.* 134.—Afr. trop.
involucrata, *Blume, Bijdr.* 116.—Java.
involucrata, Wight, Ill. Wall. Cat. p. 287, n. 1096 C = emarginata, pilosa.
lactea, *Delile, ex Steud. Nom.* ed. II. i. 706.—Afr. trop.
laevigata, Heyne, ex Steud. l. c. = excelsa.
laevigata, *Vahl, Symb. Bot.* i. 34.—As. et Afr. trop.
Latifolia, Bach.-Ham. in Wall. Cat. p. 287. n. 1095 C = laevigata.
lanceifolia, J. Grah. Cat. Pl. Bomb. 21 = polygama.

GREWIA :—
lanceæfolia, Roxb. Hort. Beng. [98] ; Fl. Ind. ii. 586 = populifolia.
lanceolata, *Baker, in Journ. Linn. Soc.* xx. (1883) 104. —Madag.
lanceolata, Herb. Heyne, ex Mast. in Hook. f. Fl. Brit. Ind. i. 391 = polygama.
lanceolata, *Miq. Anal. Bot. Ind.* iii. 4.—Ind. or.
lasiocarpa, *E. Mey. in Drège, Zwei Pfl. Docum.* 181, nomen ; *et Harv. & Sond. Fl. Cap.* i. 226.—Afr. austr.
lasiflora. *G. Don, Gen. Syst.* i. 549.—Afr. trop.
latifolia, *Mast. in Hook. f. Fl. Brit. Ind.* i. 392.—Malacca.
latifolia, *F. Muell. ex Benth. Fl. Austral.* i. 271.—Austral.
laurifolia, *Hook. ex Mast. in Hook. f. Fl. Brit. Ind.* i. 392.—Malaya.
lepidopetala, *Garcke, in Peters, Reise Mossamb. Bot.* 135.—Afr. trop.
longifolia, Buch.-Ham. in Wall. Cat. n. 1090 E = hirsuta.
macrophylla, *Baker, in Journ. Linn. Soc.* xxi. (1884) 336.—Madag.
macrophylla, G. Don, Gen. Syst. i. 549 = abutilifolia.
malacocarpa, *Mast. in Oliver, Fl. Trop. Afr.* ii. 258.—Afr. trop.
Mallococca, Blanco, Fl. Filip. ed. II. 310 = laevigata.
Malococca, *Linn. f. Suppl.* 409.—Ins. Amicor.
mantillensis, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 310.—Ins. Philipp.
megalocarpa, *Beauv. Fl. Ow.* ii. 69. t. 102.—Afr. trop.
membranacea, A. Rich. Tent. Fl. Abyss. i. 90 = populifolia.
mexicana, DC. Prod. i. 510 = Belotia grewiæfolia.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 311.—Ins. Philipp.
micrantha, *Boj. ex Mast. in Oliver, Fl. Trop. Afr.* i. 244.—Madag.
micrococca, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = seq.
Microcos, *Linn. Syst.* ed. XII. 602.—As. trop.
microphylla, *Beriol. Hort. Bonon. Pl. Nov.* ii. 12.—Ins. Cap. Verd.
microstemma, *Wall. Cat.* n. 1109.—Burma.
mollis, *Juss. in Ann. Mus. Par.* iv. (1804) 91.—Afr. trop.
Mollococca, Roxb.-Ham. in Wall. Cat. p. 237. n. 1086 F = laevigata.
montana, Koen. ex Wight & Arn. Prod. 78 = hirsuta.
monticola, *Sond. in Linnæa*, xxiii. (1850) 20.—Afr. austr.
Muenterii, Walp. Rep. i. 389 = G. Microcos.
multiflora, Blanco, Fl. Filip. ed. II. 309 = umbellata.
multiflora, *Juss. in Ann. Mus. Par.* iv. (1804) 89. t. 47. —Malaya ; Austral.
nana, Wall. Cat. n. 1102 = sapida.
nitida, *Juss. in Ann. Mus. Par.* iv. (1804) 90.—China.
obliqua, Blume, ex Steud. Nom. ed. II. i. 706 = Blumei.
obliqua, *Juss. in Ann. Mus. Par.* iv. (1804) 92.—Ind. or.
obliqua, Roxb. Hort. Beng. [98] ; Fl. Ind. ii. 590 = scabrophylla.
oblongifolia, *Blume, Bijdr.* 114.—Java.
obtecta, Wall. Cat. n. 1114 = asiatica.
obtusa, Wall. l. c. n. 1108 = bractesta, orientalis, columnaris.
obtusifolia, Eckl. & Zeyh. Enum. 88 = occidentalis.
obtusifolia, *Wild. Enum. Hort. Berol.* 566.—Afr. austr.
occidentalis, *Linn. Sp. Pl.* 964.—Africa.
odorata, Blume, Bijdr. 116 = scabrida.
omphacarpa, *Miq. Fl. Ind. Bat.* i. ii. 204.—Borneo.
opaca, *Miq. l. c.* Borneo.
oppositifolia, *Buch.-Ham. ex Roxb. Hort. Beng.* 42 ; *Fl. Ind.* ii. 583.—Reg. Himal.
orbicularis, Lamb. ex G. Don, Gen. Syst. i. 550 = orbiculata.
orbiculata, G. Don, l. c. 551 = villosa.
orbiculata, *Zoll. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 206.—Ind. or.
orbiculata, Wall. Cat. n. 1085 D = pracc. ?
orbiculis, *F. Muell. ex Benth. Fl. Austral.* i. 272.—Austral.
orientalis, Cesalp. Frost. ii. 113. t. 106 = pilosa.
orientalis, *Linn. Sp. Pl.* 964.—As. et Austral. trop.

GREWIA :—
orientalis, Roxb. ex Wight & Arn. Prod. 79 = emarginata.
orientalis, Vahl, Symb. Bot. i. 34 = columnaris.
ovalifolia, Juss. in Ann. Mus. Par. iv. (1804) 90 = laevigata.
ovalifolia, Wight, in Wall. Cat. n. 1103 B, C = orientalis.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 405.—Sumatra.
palidula, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 36 = salvifolia.
paniculata, *Roxb. Hort. Beng.* [98] ; *Fl. Ind.* ii. 591.—Malaya.
parviflora, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 83.—China.
parvifolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 91.—Abyssin.
pedicellata, Roxb. Hort. Beng. [98] ; Fl. Ind. ii. 585 = umbellata.
persicæfolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 198. Ins. Fiji.
Petitiana, *A. Rich. Tent. Fl. Abyss.* i. 89.—Ins. Philipp.
pilosa, *Lam. Encyc. Suppl.* iii. 43.—Ind. or. ; Afr. trop.
pilosa, Roxb. Hort. Beng. 42 ; *Fl. Ind.* ii. 588 = hirsuta.
pilosa, Wall. Cat. n. 1112 et p. 287.—Cf. columnaris, orientalis, scabrophylla, scabrida, humilis.
pinnatifida, *Mast. in Oliver, Fl. Trop. Afr.* i. 258.—Afr. trop.
placatorum, *Hance, in Ann. Sc. Nat. Sér.* V. v. (1866) 208.—China.
platostigma, *F. Muell. Fragm.* viii. 4.—Austral.
polygama, *Roxb. Hort. Beng.* 42 ; *Fl. Ind.* ii. 588.—As. et Austral. trop.
polygama, *Wild. Enum. Hort. Berol. Suppl.* 88, nomen.—Hab. ?
polyphylla, *Wall. Cat.* n. 1116.—Ind. or.
polyppyrsa, *Baker, in Journ. Linn. Soc.* xx. (1883) 105.—Madag.
populifolia, *Vahl, Symb. Bot.* i. 33.—Ind. or. ; Afr. trop.
prunifolia, A. Gray, Bot. U. St. Expl. Exped. i. 199 = multiflora.
pubescens, *Beauv. Fl. Ow.* ii. 76.—Afr. trop.
pumila, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 227 = sapida.
racemosa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 16 = laevigata.
reticulata, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 246 = populifolia.
retusifolia, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 294 = humilis.
rhamnifolia, Roth, Nov. Pl. Sp. 244 = orientalis.
ribesiæfolia, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 246 = populifolia.
Richardiana, Hook. in Mitch. Journ. Trop. Austral. 384.—latifolia.
Richardiana, Walp. Rep. i. 389 = Amicorum.
rigida, Buch.-Ham. in Wall. Cat. n. 1088 D et p. 287 = populifolia.
Ritchiei, *Mast. in Hook. f. Fl. Brit. Ind.* i. 389.—Ind. or.
robusta, *Burch. Trav.* ii. 138.—Afr. austr.
Rothii, Dec. Prod. i. 509 = excelsa.
rotundifolia, Juss. in Ann. Mus. Par. iv. (1804) 92. t. 50 = orbiculata.
Roxburghii, G. Don, Gen. Syst. i. 548 = hirsuta.
rugosa, G. Don, l. c. 550 = scabrophylla.
rugosa, Wall. Cat. n. 1098 B = G. Microcos.
saligna, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 544. —Madag.
salutaris, *Spach, in Linnæa*, xv. (1841) 176.—Timor.
salvifolia, *Heyne, ex Roth, Nov. Pl. Sp.* 239.—Ind. or.
salvifolia, Linn. f. Suppl. 409 = Alangium Lamarckii.
salvifolia, Roxb. Hort. Beng. [98] ; Fl. Ind. ii. 587 = excelsa.
salvifolia, Wall. Cat. n. 1091 = hirsuta.
sapida, *Roxb. Hort. Beng.* 43 ; *Fl. Ind.* ii. 590.—Reg. Himal.
saxatilis, Buch.-Ham. in Wall. Cat. n. 1091 E et p. 287 = hirsuta.
scabra, *DC. Prod.* i. 512.—Ins. Amboin.
scabrella, *Benth. Fl. Austral.* i. 272.—Austral.
scabrida, *Wall. Cat.* n. 1113, partim.—Burma ; Malaya ; Afr. trop.
scabrida, Wall. l. c. partim = pilosa.

GREWIA :—
scabrophylla, *Roxb. Fl. Ind.* ii. 584.—Reg. Himal. ; Burma.
sclerophylla, Roxb. Hort. Beng. 42 = pracc.
sepiaria, Roxb. l. c. ; Fl. Ind. ii. 589 = multiflora.
serrata, Blanco, Fl. Filip. ed. I. 444 = columnaris.
serrulata, DC. Prod. i. 510 = multiflora.
sinuata, *Wall. Cat.* n. 1108.—Burma.
stenophylla, *Boj. in Proc. Soc. Nat. Hist. Maur.* (1842-46) 97.—Madag.
subæqualis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 544.—Madag.
subinæqualis, DC. Prod. i. 511 = astatica.
subinæqualis, Wall. Cat. n. 1087 et p. 237 = tiliæfolia.
subcordata, *Miq. Fl. Ind. Bat. Suppl.* 404.—Sumatra.
suffruticosa, Buch.-Ham. in Wall. Cat. n. 1102 et p. 287 = sapida.
sulcata, *Mast. in Oliver, Fl. Trop. Afr.* i. 252.—Afr. trop.
sulcata, Wall. Cat. n. 1096 = scabrophylla.
Sulcoria, Buch.-Ham. in Wall. Cat. n. 1090 D et p. 287 = hirsuta.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 69.—Ins. Tahiti.
tembensis, Rosa. in Mus. Senckenb. ii. (1837) 158 = carpinifolia.
terebinthinacea, *DC. Cat. Hort. Monsp.* 114.—Hab. ?
tetragastris, *R. Br. ex Mast. in Oliver, Fl. Trop. Afr.* i. 253.—Afr. trop.
tiliæfolia, A. Rich. Sert. Astrol. p. ix = Amicorum.
tiliæfolia, *Vahl, Symb. Bot.* i. 35.—As. et Afr. trop.
tomentosa, *Juss. in Ann. Mus. Par.* iv. (1804) 89.—Java.
tomentosa, Roxb. ex Wight & Arn. Prod. 78 = hirsuta.
trichocarpa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 89. —Abyssin.
trifora, *Walp. Rep.* v. 119.—Ins. Maurit.
trifolia, *Roxb. Hort. Beng.* [98], nomen (Quid ?).—Ind. or.
trinervata, *Baker, in Journ. Bot.* xx. (1882) 47.—Madag.
trinervia, E. Mey. in Drège, Zwei Pfl. Docum. 188 = occidentalis.
truncata, *Mast. in Oliver, Fl. Trop. Afr.* i. 244.—Afr. trop.
turbinata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 504.—Ins. Socotr.
ulmifolia, Boj. Hort. Maurit. 44 = glandulosa.
ulmifolia, Roxb. Hort. Beng. 43 ; Fl. Ind. ii. 591 = G. Microcos.
ulmifolia, Salisb. Prod. 367 = occidentalis.
umbellata, *Roxb. Hort. Beng.* 42 ; *Fl. Ind.* ii. 591.—Malaya.
umbellifera, *Bedd. Forest. Man. Bot.* 37.—Ind. or.
variabilis, Wall. Cat. n. 1087, partim = tiliæfolia.
velutina, *Franch. in Revoil, Comali*, 21.—Afr. trop.
velutina, *Vahl, Symb. Bot.* i. 35.—Arabia.
venusta, *Fresen. in Mus. Senckenb.* ii. (1837) 159. t. 10. —Abyssin.
verrucosa, Juss. in Ann. Mus. Par. iv. (1804) 90 = excelsa.
vestita, Wall. Cat. n. 1105 = asiatica.
villifera, *Sweet, Hort. Brit.* ed. II. 71.—Ind. or.
villosa, Heyne, ex Roth, Nov. Pl. Sp. 248 = villifera.
villosa, *Willd. in Ges. Nat. Fr. Neue Schr.* iv. (1803) 206.—Ind. or.
viminea, Wall. Cat. n. 1111 = polygama.
viridifora, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 89.—Malaya.
xanthopetala, *F. Muell. ex Benth. Fl. Austral.* i. 271. —Austral.

GREWIA, Hook. & Harv. in Proc. Dubl. Univ. Zool. Bot. Assoc. i. (1859) 188. t. 18. *SAPINDACEÆ*, Benth. & Hook. f. i. 1000.
Sutherlandii, *Hook. & Harv. l. c.*—Afr. austr.

GRIAS, Linn. Syst. ed. X. 1075 (1759). *MYRTACEÆ*, Benth. & Hook. f. i. 732.
Anbletana, *Miers, in Trans. Linn. Soc.* xxx. (1874) 299.—Guiana.
cauliflora, *Linn. Syst.* ed. X. 1075.—Ind. occ.
Fendleri, *Seem. Bot. Voy. Herald*, 126.—Panama.
peruviana, *Miers, in Trans. Linn. Soc.* xxx. (1874) 301.—Peruv.

GRIELUM, Linn. Gen. ed. VI. 578 (1764). *ROSACEAE*, Benth. & Hook. f. i. 626.
flagelliforme, E. Mey. in Drège, Zwei. Pfl. Docum. 188 = humifusum.
humifusum, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 305 = tenuifolium.
humifusum, *Thunb. Prod. Pl. Cap.* 113.—Afr. austr.
lactucatum, Gaertn. Fruct. i. 180. t. 36 = humifusum.
obtusifolium, E. Mey. in Drège, Zwei Pfl. Docum. 188 = sinuatum.
sinuatum, *Licht. ex Burch. Trav.* i. 286.—Afr. austr.
tenuifolium, *Linn. Gen. ed.* VI. 578.—Afr. austr.

GRIESEBACHIA, Endl. Gen. 759 (1839) = Griesebachia, Klotzsch (Ericac.).

GRIESELINIA, Endl. Gen. 1309, in syn. (1841) = Griselinia, Forst. f. (Cornac.).

GRIFFINIA, [Ker-Gawl. in] Bot. Reg. sub t. 444 et t. 511 (1820). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 726.
LIBONIA, Lem. Jard. Fleur. iii. (1853–55) t. 290.
Blumenavia, *C. Koch & Bouché, ex Carr. in Rev. Hortic.* (1867) 92.—Bras.
dryades, *M. Korn. Syn. Ensat.* 32.—Bras.
hyacinthina, [*Ker-Gawl. in*] *Bot. Reg.* sub t. 444. —Bras.
intermedia, *Lindl. in Bot. Reg.* t. 990.—Bras.
Liboniana, *Lem. Jard. Fleur.* iii. (1853) t. 290.—Bras.
ornata, *T. Moore, in Gard. Chron.* (1876) i. 266.—Bras.
parviflora, [*Ker-Gawl. in*] *Bot. Reg.* t. 511.—Bras.

GRIFFITHIA, Wight & Arn. Prod. 299 (1834) = Bandia, Houtt. (Rubiac.).
acuminata, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 178.—Borneo.
cuneata, Kurz, in Journ. Bot. xiii. (1875) 326 = R. longiflora.
eucantha, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 178.—Borneo; Java.
fragrans, Wight & Arn. Prod. 400 = R. malabarica.
Gardneri, Thw. Enum. Pl. Zeyl. 158 = R. Gardneri.
latifolia, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxiv. (1864) 23 = R. latifolia.
leucantha, Wslp. Ann. iii. 1029, Ind. aphalm. = *eucantha*.
longiflora, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 178 = R. longiflora.
oxygyna, Korth. l. c.—Sumatra.
palembanica, Miq. Fl. Ind. Bat. Suppl. 541.—Malaya.
Reinwardtiana, Hassk. ex Teijsm. & Binn. Cat. Hort. Bog. 118.—Java.
rugulosa, Thw. Enum. Pl. Zeyl. 159 = R. rugulosa.
stamensis, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 130 = R. longiflora.
speciosa, Bedd. Ic. Pl. Ind. Or. i. 8. t. 87 = R. speciosa.

GRIFFONIA, Baill. in Adansonia, vi. 188. t. 2 (1865) = Bandstraea, Welw. (Legumin.).
physocarpa, Baill. l. c. t. 2. f. 3–5 = B. tenuiflora.
simplicifolia, Baill. l. c. (1866) 197 = B. simplicifolia.

GRIFFONIA, Hook. f. in Benth. & Hook. f. Gen. i. 606 (1865). *ROSACEAE*, Benth. & Hook. f. i. 6.
Bartet, *Hook. f. ex Oliver, Fl. Trop. Afr.* ii. 373.—Afr. trop.
Bellayana, *Oliver, l. c.* 371.—Afr. trop.
Icondera, *Oliver, l. c.*—Afr. trop.
Mannii, *Oliver, l. c.* 373.—Afr. trop.
pallescens, *Oliver, l. c.*—Afr. trop.

GRIMALDIA, Schrank, in Allg. Bot. Biblioth. iv. (1805) 185; et in Denkschr. Akad. Muench. (1808) 108. t. 3 = Cassia, Linn. (Legumin.).
Abms, Link, Handb. ii. 141 = C. Absus.
acuergana, Schrank, in Denkschr. Akad. Muench. (1808) 110 = C. nicitiana.
bifoliata, Schrank, ex Steud. Nom. ed II. i. 707 = C. rotundifolia.
Chamaecrista, Schrank, ex Link, Handb. ii. 141 = C. Chamaecrista.
dorembosa, Schrank, in Denkschr. Akad. Muench. (1808) 111 = C. nicitiana.
ophira, Schrank, in Allg. Bot. Biblioth. iv. (1805) 187 = C Absus.

GRINDELIA, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 259. *COMPOSITAE*, Benth. & Hook. f. ii. 250.
AURELIA, Cass. in Bull. Soc. Philom. (1818) 175; et in Dict. Sc. Nat. xxxvii. 468 (1825).
CHRYSOPSTHALMUM, Phil. in Linnaea, xxix. (1858) 9.
DEMETRIA, Lag. Gen. et Sp. Nov. 30 (1816).
DONIA, [R. Br. in] Ait. Hort. Kew. ed. II. v. 82 (1813).
HRINIANA, Rafin. in Am. Monthly Mag. (1818) 268.
acerosa, *Bert. in Merc. Chil.* (1829); *ex Am. Journ. Sc.* xxiii. (1833) 84.—Chili.
andina, *Phil. in Linnaea*, xxxiii. (1864–65) 137.—Chili.
angustifolia, *DC. ex Dunal, in Mém. Mus. Par.* v. (1819) 51. t. 3.—Mexic.
angustifolia, H. B. & K. Nov. Gen. et Sp. iv. 309 = arguta.
anomala, DC. Prod. v. 316 = discoides.
arguta, A. Gray, Pl. Wright. ii. 81 = squarrosa.
arguta, *Schrad. ex DC. Prod.* v. 315.—Mexic.
arizonica, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 208.—Arizona.
brachystephana, *Griseb. in Gott. Abh.* xxiv. (1879) 176.—Reg. Argent.
buphthalmoides, *DC. Prod.* v. 316.—Bras.
cancrorum, *Bert. in Merc. Chil.* (1829); *ex Am. Journ. Sc.* xxiii. (1833) 84.—Chili.
cascasica, Spreng. ex Steud. Nom. ed. II. i. 707 = Astex cascasica.
ciliata, *Spreng. Syst.* iii. 575.—Mexic.
coronopifolia, Lehm. Ind. Sem. Hort. Hamb. (1828) 16 = Xanthocephalum centauroides.
costata, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 208.—Mexic.
cuneifolia, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 315.—Calif.
diffusa, *D. Don, ex Arn. in DC. Prod.* vii. 278.—Patagon.
diffusa, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 45 = pulchella.
discoides, *Hook. & Arn. l. c.*—Bras.
discoides, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 315 = nana.
Duvalii, Spreng. Syst. iii. 575 = angustifolia.
foliosa, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 45.—Chili.
fruticosa, *Dunal, in Mém. Mus. Par.* v. (1819) 52 (sp. dub.).—Hab. ?
globularifolia, *Griseb. in Gott. Abh.* xxiv. (1879) 178. —Reg. Argent.
glutinosa, Bert. in Merc. Chil. (1829) 646; ex DC. Prod. v. 316 = Aplopappus Berterii, macrocephalus.
glutinosa, *Dunal, in Mém. Mus. Par.* v. (1819) 49.—Am. bor. occ.
grandiflora, Hook. Bot. Mag. t. 4628 = squarrosa.
hirsuta, *Hort.—Cf. Gard. Chron.* (1862) ii. 140.—Hab. ?
hirsutula, *Hook. & Arn. l. c.*—Calif.
incisa, Spreng. Syst. iii. 575 = Aster incisus.
integrifolia, *DC. Prod.* v. 315.—Am. bor. occ.
inuloides, *Willd. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 261.—Am. bor. occ.
Lamberti, *G. Don, in Lond. Hort. Brit.* 350, nomen. —Mexic.
lanceolata, *Nutt. in Journ. Acad. Philad.* vii. (1834) 78.—Am. bor.
latifolia, Kellogg, in Proc. Calif. Acad. v. (1873) 36 = glutinosa.
microcephala, DC. Prod. v. 315 = inuloides.
microcephala, Rothr. in Wheeler, Rep. vi. 141 = arisonica.
nana, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 314.—Am. bor. occ.
nitida, *Bernh. ex Sweet, Hort. Brit. ed.* II. 396.—Mexic.
nuda, Wood, in Coult. Bot. Gaz. iii. (1878) 50 = squarrosa.
oregana, *A. Gray, Syn. Fl. N. Am.* i. ii. 118.—Am. bor. occ.
parifloa, M. E. Jones, in Bull. Torrey Bot. Club, ix. (1882) 31 = nana.
pectinata, *Baker, in Mart. Fl. Bras.* vi. 111. t.—Bras.
perucosa, Sch. Bip. in Leubl. Beob. Am. Amoz. 37 = spp.
peruviana, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.

GRINDELIA:—
puberula, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 45.—Reg. Argent.
pulchella, Nutt. in Journ. Acad. Phil. vii. (1834) 74 = inuloides.
pulchella, Bert. in Merc. Chil. (1829) 643; et in Am. Journ. Sc. xxiii. (1833) 84 = Aplopappus glutinosus, pulchellus.
pulchella, *Dunal, in Mém. Mus. Par.* v. (1819) 51. t. 2.—Chili.
rosinosa, *Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 45.—Chili.
robusta, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 314.—Calif.
rubricaulis, DC. Prod. v. 316 = hirsutula.
scorzonerifolia, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 45.—Reg. Argent.
sibirica, Spreng. ex Link, Enum. Hort. Berol. ii. 336 = Aster sibiricus.
spathulata, Bivona, ex Link, l. c. = inuloides.
speciosa, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 45 = squarrosa.
speciosa, *Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 118. t. 294.—Patagon.
squarrosa, *Dunal, in Mém. Mus. Par.* v. (1819) 50.—Am. bor. occ.
stricta, DC. Prod. vii. 278 = integrifolia.
subdecurrens, DC. l. c. v. 315 = squarrosa.
tenuna, Schecle, in Linnaea, xxi. (1848) 60 = squarrosa.
villosa, *D'ongl. !] ex Sweet, Hort. Brit. ed.* II. 299.—Am. bor. occ.
virgata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 314 = oregana.

GRINDELIOPSIS, Sch. Bip. in Bonplandia, vi. (1858) 256 = Xanthocephalum, Willd. (Compos.).
gymnospermoides, Sch. Bip. l. c. = X. gymnospermoides.

GRIPIDEA, Miers, in Trans. Linn. Soc. xxv. (1865) 227. t. 28 = Blumenbachia, Schrad. (Loasene).
asperata, Miers, l. c. 235.—Bras.
scabra, Miers, l. c.—Bras.

GRISCHOWIA, Karst. Ausw. Neue Gew. 15. t. 5 (1848) = Monochaetum, Nand. (Melastom.).
hirta, Karst. l. c. 16. t. 5 = M. hirtum.
meridensis, Klotzsch & Karst. l. c. = M. meridense.
sobechioides, Klotzsch & Karst. l. c. = M. umbellatum.

GRISEBACHIA, Klotzsch, in Linnaea, xii. (1838) 220. *ERICACEAE*, Benth. & Hook. f. ii. 593.
ACROETHNON, Klotzsch, in Linnaea, xii. (1838) 237.
COMACEPHALUS, Klotzsch, l. c. 224 (1838).
FINCKEA, Klotzsch, l. c. 237 (1838).
PILOPUS, Rafin. Fl. Tellur. ii. 102 (1836), nomen prius.
brunlades, *Benth. in DC. Prod.* vii. 702.—Afr. austr.
ciliaris, *Klotzsch, in Linnaea*, xii. (1838) 228.—Afr. austr.
Dregeana, *Benth. in DC. Prod.* vii. 701.—Afr. austr.
eriocephala, *Benth. l. c.* 702.—Afr. austr.
hirta, *Klotzsch, in Linnaea*, xii. (1838) 220.—Afr, austr.
hispida, *Klotzsch, l. c.*—Afr. austr.
incana, *Klotzsch, l. c.* 225.—Afr. austr.
involuta, *Klotzsch, l. c.* 227.—Afr. austr.
pinnoss, *Klotzsch, l. c.* 226.—Afr. austr.
serrulata, Benth. in DC. Prod. vii. 701 = pinnosa.
Thunbergii, *Esch, in Linnaea*, xxvi. (1856) [1855] 789. —Afr. austr.
vellerifiora, *Klotzsch, l. c.* xii. (1838) 227.—Afr. austr. Zayberiana, *Klotzsch, l. c.*—Afr. austr.

GRISSEBACHIA, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 177 = Howea, Becc. (Palm.).
Belmoreana, H. Wendl. & Drude, l. c. 202 = H. Belmoreana.
Forsteriana, H. Wendl. & Drude, l. c. 203 = H. Forsteriana.

GRISELEA, Linn. Gen. ed. VI. 578 (1764). *ROSACEAE*, Benth. & Hook. f. i. 626.
flagelliforme, E. Mey. in Drège, Zwei. Pfl. Docum. 188 = humifusum.
humifusum, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 305 = tenuifolium.
humifusum, *Thunb. Prod. Pl. Cap.* 113.—Afr. austr.
lactucatum, Gaertn. Fruct. i. 180. t. 36 = humifusum.
obtusifolium, E. Mey. in Drège, Zwei Pfl. Docum. 188 = sinuatum.
sinuatum, *Licht. ex Burch. Trav.* i. 286.—Afr. austr.
tenuifolium, *Linn. Gen. ed.* VI. 578.—Afr. austr.

GRISELINIA, Forst. f. Prod. 75 (1786).—*Cl.* Forst. Char. Gen. Ind. *CORNACEAE*, Benth. & Hook. f. i. 951.
Decostea, Ruiz & Pav. Prod. 130 (1794).
Pukateria, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 130.
Scutolia, Forst. Char. Gen. 189. t. 70 (1776).
littoralis, *Raoul, Choix Pl. N. Zél.* 22. t. 19.—N. Zel. Incida, *Forst. f. Prod.* 75.—N. Zel.

GRISELINIA, Soop. Introd. 308 (1777) = **Pterocarpus**, Linn. (Legum.).

GRISIA, Brongn. in Bull. Soc. Bot. Fr. xii. (1865) 404 = **Bikkia**, Reinw. (Rubiac.).
campanulata, Brongn. l. c. 406.—N. Caled.
fritillarioides, Brongn. l. c.—N. Caled.
macrophylla, Brongn. l. c. 405.—N. Caled.
nervifolia, Brongn. l. c. 405.—N. Caled.
retusiflora, Brongn. l. c. 407.—N. Caled.
tubiflora, Brongn. l. c.—N. Caled.

GRISLEA, Linn. Gen. 342 (1737). *LYTHRARIEAE*, Benth. & Hook. f. i. 777.
microphylla, Hochst. & Steud. in Flora, xxiv. (1841) 1. Introl. 30 = Woodfordia floribunda.
multiflora, A. Rich. Tent. Fl. Abyss. i. 284 = Woodfordia floribunda.
punctata, Buch.-Ham. ex Sm. in Rees, Cycl. xvii. n. 2 = Woodfordia floribunda.
secunda, *Linn. Sp. Pl.* 848.—Am. trop.
tomentosa, Roxb. Pl. Corom. i. 29. t. 31 = Woodfordia floribunda.
uniflora, A. Rich. Tent. Fl. Abyss. i. 281 = Woodfordia floribunda.

GRISOLLEA, Baill. Adansonia, iv. (1863–64) 217. *OLACINEAE*, Benth. & Hook. f. i. 997.
myrianthæ, *Baill. l. c.*—Afr. austr.

GROBYA, Lindl. Bot. Reg. t. 1740 (1835). *ORCHIDEAE*, Benth. & Hook. f. iii. 543.
Amherstiæ, *Lindl. l. c.*—Bras.
galeata, *Lindl. l. c. xxvi.* (1840) *Misc.* 83.—Bras.

GROELANDIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. 159 (1869) = seq.

GROENLANDIA, J. Gay, in Compt.-Rend. Par. xxxvii. (1854) 708 = **Potamogeton**, Tourn. (Naiad.).
densa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = P. densus.

GROMOVIA, Regel, Gartenfl. xiv. (1865) 289 = **Belopærone**, Nees (Acanthac.).
pulchella, Regel, l. c. = B. violacea.
violacea, Regel, l. c. = B. violacea.

GROMPHAENA, St. Hil. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = **Gomphrena**, Linn. (Amarant.).

GROWA, Lour. Fl. Cochinch. 459 (1790). *LEGUMINOSAE*, Benth. & Hook. f. i. 584.
Dalbellii, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 191.—Ind. or.
filicoides, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 223.—Burma.
Grahamii, *Benth. Fl. Hongk.* 238.—Burma.
rubens, Herb. Banks. ex Benth. & Hook. f. Gen. i. 535 = Desmodium polystachyum.
repens, *Lour. Fl. Cochinch.* 459.—Cochinch.

GRONE, Spreng. Syst. vii. 945 (1828) = præc.

GRONOPHYLLUM, Scheff. in Ann. Jard. Buitenz. i. (1876) 155, 158. *PALMAE*, Benth. & Hook. f. iii. 886.
ischiocarpum, *Scheff. l. c.* 158.—Malaya.
selebicum, *Becc. l. c. ii.* (1886) 89.—Ins. Celebes.

GRONOVIA, Blanco, Fl. Filip. 186 (1837) = **Illigera**, Blume (Combretac.).
ternata, Blanco, l. c. 187 = I. corymbosa.

GRONOVIA, Houst. ex Linn. Syst. ed. I (1735). *LOASEAE*, Benth. & Hook. f. i. 803.
Humboldtiana, Roem. & Schult. Syst. v. 402 = scandens.
Scandalaena, M. Roem. Syn. Pepon. 108 = scandens.
scandens, *Linn. Sp. Pl.* 956.—Am. austr.

GROSOWIDYA, Reichb. f. in Bot. Zeit. xxii. (1864) 297 = **Sarcochilus**, R. Br. (Orchid.).
elegans, Reichb. f. l. c. = S. Hystrix.

GROSSOSTYLIS, Pers. Syn. ii. 260 (1807), sphalm. = **Crossostylis**, Forst. (Rhisoph.).

GROSSULARIA, Tourn. ex Adans. Fam. ii. 243 (1763) = **Ribes**, Linn. (Saxifrag.).
acicularis, Spach, Hist. Vég. Phan. vi. 173 = R. acicularе.
adenophylla, Rupr. Sert. Tianach. 47 = R. heterotrichum.
Cynosbati, Mill. Gard. Dict. ed. VIII. n. 5 = R. Cynosbati.
diacantha, Mill. ex Steud. Nom. ed. II. i. 708 = R. diacanthum.
divaricata, Stend. l. c. ii. 457, in syn. = R. divaricatum.
glandulosa-setosa, Opiz, Seznam, 48 = R. Grossularia.
hirsuta, Mill. Gard. Dict. ed. VIII. n. 2 = R. Grossularia.
hirtella, Spach, Hist. Vég. Phan. vi. 180 = R. oxyacanthoides.
intermedia, Opiz, Seznam, 48 = R. Grossularia.
irrigua, S. Wats. Bibliog. Ind. N. Am. Bot. i. 338 = R. divaricatum.
nigra, Mill. ex Steud. Nom. ed. II. i. 708 = R. nigrum.
nivea, Spach, Hist. Vég. Phan. vi. 179 = R. gracile.
oxyacanthoides, Mill. Gard. Dict. ed. VIII. n. 4 = R. oxyacanthoides.
pubescens, Opiz, Seznam, 48 = R. Grossularia.
reclinata, Mill. Gard. Dict. ed. VIII. n. 1 = R. Grossularia.
rubra, Desp. Fl. Carn. ed. II. i. 167 = R. rubrum.
triflora, Spach, Hist. Vég. Phan. vi. 176 = R. rotundifolium.
Uva, Desp. Fl. Carn. ed. II. i. 166 = R. Grossularia.
Uva-crispa, Mill. Gard. Dict. ed. VIII. n. 3 = R. Grossularia.
vulgaris, Spach, Hist. Vég. Phan. vi. 174 = R. Grossularia.

GROTEFENDIA, Seem. Journ. Bot. ii. (1864) 247 = **Polycnias**, Forst. (Araliac.).
cuneata, Seem. l. c. = P. paniculata.
paniculata, Seem. l. c. = P. paniculata.
repanda, Seem. l. c. 248 = P. repanda.

GROUTIA, Guill. & Perr. Fl. Senegamb. Tent. 100. t. 22 (1833) = **Opilia**, Roxb. (Opiliac.).
celtidifolia, Guill. & Perr. l. c. 101. t. 22 = O. amentacea.

GRUBBIA, Berg. in Vet. Akad. Handl. Stockh. (1767) 34; et Desc. Fl. Cap. 90. t. 2 (1767). *SANTALACEAE*, Benth. & Hook. f. iii. 231.
Ophira, Linn. Mant. ii. 150 (1771).
Strobilocarpus, Klotzsch, in Linnæa, xiii. (1839) 350.
capensis, J. F. Gmel. Syst. 629 = rosmarinifolia.
hirsuta, *E. Mey. in Drège, Zwei Pfl. Docum.* 188, nomen; *et ex A. DC. Prod.* xiv. 618.—Afr. austr.
pinifolia, *Sond. in Harv. & Sond. Fl. Cap.* ii. 326.—Afr. austr.
rosmarinifolia, *Berg. in Vet. Akad. Handl. Stockh.* (1767) 86; *et Desc. Pl. Cap.* 90.—Afr. austr.
rosmarinifolia, Knaus, ex A. DC. Prod. xiv. 618 = stricta.
stricta, *A. DC. l. c.*—Afr. austr.

GRUENERA, Opiz, Seznam, 48 (1852) = **Salix**, Tourn.
amygdaloides, Opiz, l. c. = S. triandra.

GRUHLMANIA, Neck. Elem. i. 302 (1790) = **Spermacoce**, Linn. (Rubiac.).

GRUMELIA, Wight, Ic. iii. 16 (1845) = seq.

GRUMILEA, Gaertn. Fruct. i. 133. t. 28. f. 2 (1788) = **Psychotria**, Linn. (Rubiac.).
adenophylla, Miq. Fl. Ind. Bat. ii. 297 = P. adenophylla.
africa, Thw. Enum. Pl. Zeyl. 146 = P. Wightiana.
argentea, Miq. Fl. Ind. Bat. ii. 299 = P. argentea.
articulata, Hiern, in Oliver, Fl. Trop. Afr. iii. 216.—Afr. trop.
asperuloides, Miq. Fl. Ind. Bat. ii. 296 = P. malayana.
capensis, Sond. in Harv. & Sond. Fl. Cap. iii. 21 = P. capensis.

GRUMILEA—
congesta, Wight & Arn. Prod. 422 = Psychotria congesta, *Hook. f.*
cymosa, E. Mey. in Drège, Zwei Pfl. Docum. 188 = P. Eckloniana.
divaricata, Thw. Enum. Pl. Zeyl. 149 = Gaertnera Koenigii.
divergens, Miq. Fl. Ind. Bat. ii. 299 = P. divergens.
elongata, Wight, Ic. t. 1036 = P. elongata, *Hook. f.*
fimbricalyx, Miq. Fl. Ind. Bat. ii. 297.—Java.
Gardneri, Thw. Enum. Pl. Zeyl. 147 = P. Gardneri.
globosa, Nochst. in Flora, xxvii. (1844) 554 = P. capensis.
Junghuniana, Miq. Fl. Ind. Bat. ii. 296 = P. sarmentosa.
Kirkii, Hiern, in Oliver, Fl. Trop. Afr. iii. 216.—Afr. trop.
longifolia, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 60, nomen.—Ind. or.
longiflora, Bedd. in Trans. Linn. Soc. xxv. (1865) 218 = P. nanaallaya.
macrocarpa, Hiern, in Oliver, Fl. Trop. Afr. iii. 217.—Afr. trop.
micrantha, Hiern, l. c.—Afr. trop.
Moonii, Thw. Enum. Pl. Zeyl. 148 = P. Moonii.
morindoides, Miq. Fl. Ind. Bat. Suppl. 547 = P. leucocarpa.
nigra, Gaertn. Fruct. i. 133. t. 28 = P. Thwaitesii †
nudiflora, Thw. Enum. Pl. Zeyl. 147 = P. Thwaitesii.
patens, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 169 = P. congesta, *Hook. f.*
polycarpa, Miq. Fl. Ind. Bat. ii. 295 = P. sarmentosa.
psychotrioides, DC. Prod. iv. 495.—Afr. trop.
Rawsoni, Hook. & Arn. Bot. Beech. Voy. 193 = P. elliptica.
sphærocarpa, Hiern, in Oliver, Fl. Trop. Afr. iii. 218.—Afr. trop.
stenophylla, Thw. Enum. Pl. Zeyl. 147 = P. stenophylla.
subintegra, Wight & Arn. Prod. 432 = P. subintegra.
succulenta, Hiern, in Oliver, Fl. Trop. Afr. iii. 216 = P. succulenta.
sulphurea, Hiern, l. c. 218 = P. sulphurea, *Schweinf.*
sylvatica, Miq. Fl. Ind. Bat. ii. 298 = P. sylvatica.
Teymannziana, Miq. l. c. Suppl. 547.—Ins. Banca.
umbellata, Miq. l. c. ii. 295.—Sumatra.
venosa, Hiern, in Oliver, Fl. Trop. Afr. iii. 217.—Afr. trop.
viridiflora, Miq. Fl. Ind. Bat. ii. 298 = P. viridiflora.
Wightiana, Thw. Enum. Pl. Zeyl. 148 = P. Wightiana.

GRUNDLEA, Steud. Nom. ed. II. i. 706 (1840), sphalm. = præc.

GRUVELIA, A. DC. in DC. Prod. x. 119 (1846) = **Pectocarya**, DC. (Boragin.).
pusilla, A. DC. l. c. = P. pusilla.

GRYMANIA, Presl, Epim. Bot. 193 (1849) = **Parinarium**, Juss. (Rosac.).
polyandra, Presl, l. c. = Conepia Kunthiana.
salicifolia, Presl, l. c. = P. salicifolium.

GUACHAMACA, De Gros, in Med. Bot. Criollo, i. 295; ex Ernst, in Trans. Bot. Soc. Edinb. x. (1870) 450 = **Prestonia**, R. Br. (Apocyn.).
toxicaria, De Gros, l. c.—Venezuela.

GUACO, Llebm. in Forhandl. Skandin. Naturf. 1844 (1847) 908 = Aristolochia, Linn.
mexicana, Llebm. l. c.—Mexic.

GUADUA, Kunth, Syn. Pl. i. 292 (1822). *GRAMINEAE*, Benth. & Hook. f. iii. 1210.
ecclesia, *Rupr. ex Fourn. in Compt.-Rend. Par.* lxxiv. (1877) 198.—Mexic.
amplexifolia, *J. & C. Presl, Rel. Haenk.* i. 256.—Mexic.
angustifolia, *Kunth, Syn. Pl.* i. 293.—N. Granat.
capitata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 81. —Peru.
disica, Steud. Syn. Pl. Gram. 384 = macrostachya.
distorta, *Rupr. in Mém. Acad. Sc. Pétersb. Sér.* VI. v. (1839) 181.—Mexic.
ensulata, Doell, in Mart. Fl. Bras. ii. iii. 181 = Arthrostylidium longiflorum.
fascicularis, *Doell, l. c.* 186.—Guiana.

GUADUA:—
flabellata, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 558.—Mexic.
glomerata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 79.—Bras.
latouana, *Rupr. ex Hemsl. Biol. Centr. Am. Bot.* iii. 558.—Mexic.
intermedia, *Rupr. ex Hemsl. l. c.*—Mexic.
latifolia, *Kunth, Syn. Pl.* i. 254.—Am. austr.
macrostachya, *Rupr. in Mém. Acad. Sc. Pétersb.* Sér. VI. v. (1839) 129.—Guiana.
pallescens, Doell, in Mart. Fl. Bras. ii. III. 186 = Bambusa pallida.
paniculata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 85.—Bras.
paraguayana, *Doell, in Mart. Fl. Bras.* ii. III. 179.—Bras.
parviflora, J. & C. Presl, in Rel. Haenk. i. 257 = Arthrostylidium maculatum.
polyclados, *Doell, in Mart. Fl. Bras.* ii. III. 182.—Bras.
refracta, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 84.—Bras.
Tagoara, *Kunth, Enum. Pl.* i. 434.—Bras.
Trinii, *Rupr. in Mém. Acad. Sc. Pétersb.* Sér. VI. v. (1839) 130.—Bras.
Venezuelae, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 86.—Venezuela.
virgata, *Rupr. in Mém. Acad. Sc. Pétersb.* Sér. VI. v. (1839) 130.—Bras.

GUAGNEBINA, Vell. Fl. Flum. 45 (1825); i. t. 115 (1827) = **Manettia**, Mutis (Rubiac.).
axillaris, Vell. l. c. 46; i. t. 119 = M. albiflora?
congesta, Vell. l. c.; i. t. 120 = M. albiflora.
ignita, Vell. l. c. 45; i. t. 116 = M. cordifolia.
luteo-rubra, Vell. l. c. 46; i. t. 121 = M. luteo-rubra.
intexens, Vell. l. c. 45; i. t. 116 = M. ciliata?
mitis, Vell. l. c. 46; i. t. 118 = M. fimbriata.
suavis, Vell. l. c. 45; i. t. 117 = M. fimbriata?

GUAIACON, Adans. Fam. ii. 507 (1763) = seq.

GUAIACUM, Plum. ex Linn. Gen. ed. I. 140 (1737). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 347.
Abilo, Blanco, Fl. Filip. ed. I. 264 = Garuga pinnata.
afrum, Linn. Sp. Pl. 383 = Schotia speciosa.
angustifolium, Engelm. in Wisliz. Tour N. Mexic. 118 = Porlieria angustifolia.
arboreum, *Dc. Prod.* i. 707.—Am. trop.
bijugum, Stokes, Bot. Mat. Med. ii. 456 = officinale.
Breynii, *Spreng. Syst.* ii. 322.—Am. austr.
Coulteri, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 312.—Mexic.
dubium, *Forst. f. Prod.* 32.—Ins. Amicor.
guatemalense, *Planch. ex Hemsl. Biol. Centr. Am. Bot.* i. 159, nomen.—Mexic.
microphyllum, *Baill. Adansonia*, x. (1872) 315.—Bolivia.
multijugum, Stokes, Bot. Mat. Med. ii. 456 = sanctum.
officinale, *Linn. Sp. Pl.* 381.—Ind. occ.
parvifolium, *Planch. ex A. Gray, Pl. Wright.* i. 29.—Mexic.
sanctum, *Linn. Sp. Pl.* 383.—Am. trop.
Stoemei, Shuttl. ex A. Gray, Pl. Wright. i. 29 = verticale.
verticale, *Orteg. Hort. Matr. Dec.* 98.—N. Hisp.; Deming.
verticale, A. Rich. Ess. Fl. Cub. 321 = sanctum.

GUAIAVA, Tourn. ex Adans. Fam. ii. 88 (1763) = Psidium, Linn. (Myrtac.).
pyriformis, Gaertn. Fruct. i. 155. t. 33 = P. Guajava.

GUAJACUM, Plum. ex Linn. Syst. ed. I (1735) = Guaiacum, Plum. (Zygophyll.).

GUALTERIA, Scop. Introd. 194 (1777) = Gaultheria, Linn. (Ericac.).
brevipedata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = Gaultheria brevipes.

GUALTHERIA, J. F. Gmel. Syst. 697 (1791) = Gaultheria, Linn. (Ericac.).

GUANDIOLA, Steud. Nom. ed. I. 384 (1821), sphalm. = Guardiola, Humb. & Bonpl. (Compos.).

GUANIA, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 129 = Gouania, Jacq. (Rhamn.).

GUAPEA, Endl. Ench. 861, in syn. (1841) = seq.

GUAPEBA, Gomes, in Mem. Ac. Lisbos, iii. (1812) Mem. dos Corresp. t. 2 = seq.

GUAPEBEIRA, Gomes, in Mem. Ac. Lisbos, iii. (1812) Mem. dos Corresp. 19 = Lucuma, Juss. (Sapotac.).
brasiliensis, Steud. Nom. ed. II. i. 708 = seq.
laurifolia, Gomes, in Mem. Ac. Lisbos, iii. (1812) Mem. dos Corresp. 21 = L. laurifolia.

GUAPINA, Steud. Nom. ed. II. i. 708 (1840), sphalm. = seq.

GUAPIRA, Aubl. Pl. Gui. i. 808. t. 319 (1775). *INC. SED.*, an Verbenaceae?—Cf. Benth. & Hook. f. ii. 1137.
GYNASTRUM, Neck. Elem i. 224 (1790).
guianensis, *Aubl. l. c.*—Guian.

GUAPURIUM, Juss. Gen. 394 (1789) = **Eugenia**, Linn. (Myrtac.).
fruticosum, Spreng. Syst. ii. 479 = E. Guapurium.
herbaceum, Spreng. l. c. = Klaprothia mentzelioides.
peruvianum, Poir. in Dict. Sc. Nat. xx. 11 = E. Guapurium.

GUAPURUM, J. F. Gmel. Syst. 793 (1791) = prec.

GUARANIA, Wedd. ex Baill. Etud. Gén. Euph. 598 (1858) = **Richeria**, Vahl (Euphorb.).
Gardneriana, Baill. Adansonia, v. (1864–65) 348 = R. grandis.
laurifolia, Baill. l. c. = R. grandis.
longifolia, Baill. l. c. 349 = R. grandis.
purpurascens, Baill. l. c. 347 = R. grandis.
ramiflora, Baill. Etud. Gén. Euph. 598, nomen = R. grandis.
Spruceana, Baill. Adansonia, v. (1864–65) 348 = R. grandis.

GUARARIBEA, Cav. Diss. ii. App. (1785), err. typ. = Quararibea, Aubl. (Malvac.).

GUARDIOLA, Cerv. ex Humb. & Bonpl. Pl. Aequin. i. 148. t. 41 (1808). *COMPOSITAE*, Benth. & Hook. f. ii. 347.
TULOCARPUS, Hook. & Arn. Bot. Beech. Voy. 298. t. 68 (1841).
stipulicifolia, *A. Gray, Pl. Wright.* i. 111.—Mexic.
mexicana, *Humb. & Bonpl. Pl. Aequin.* i. 144. t. 41.—Mexic.
platyphylla, *A. Gray, Pl. Wright*, ii. 91.—N. Mexic.
Tulocarpus, *A. Gray, l. c.* 113.—Mexic.

GUARREA, Allem. ex Linn. Mant. ii. 150 (1771). *MELIACEAE*, Benth. & Hook. f. i. 385.
PLUMEA, Lunan, Hort. Jamaic. ii. 77 (1814).
RUAGEA, Karst. Fl. Columb. ii. 51. t. 126 (1862–69).
acuminata, Wall. Cat. n. 7262 = Dysoxylum proceruum.
afinis, *A. Juss. in Mém. Mus. Par.* xix. (1830) 338.—Guian.
africana, *Welw. ex C. DC. in DC. Monog. Phan.* i. 576.—Afr. trop.
alba, *C. DC. in Mart. Fl. Bras.* xi. 1. 189.—Bras.
alliacea, Blume, in Flora, vii. (1824) 290 = Dysoxylum alliaceum.
alliaria, Buch.-Ham. in Mem. Wern. Soc. vi. (1832) 305 = Dysoxylum Hamiltonii.
alternans, *C. DC. in Mart. Fl. Bras.* xi. 1. 196.—Bras.
amoris, Buch.-Ham. in Mem. Wern. Soc. vi. (1832) 306 = Dysoxylum binectariferum.
Amyris, Steud. Nom. ed. II. i. 709, sphalm. = prec.
apiiodora, *Baill. Adansonia*, x. (1871). 110.—N. Granat.
Audotii, A. Juss. in Mém. Mus. Par. xix. (1830) 240 = trichilioides.
axillaris, *Blume, in Flora*, vii. (1824) 290, nomen.—Java.
bijuga, *C. DC. in DC. Monog. Phan.* i. 567.—Am. trop.
binectariflora, Roxb. Hort. Beng. [36]; Fl. Ind. ii. 360 = Dysoxylum binectariferum.
Blanchetii, *C. DC. in Mart. Fl. Bras.* xi. 1. 187.—Am. trop.
brachystachya, [*Mey. & Sesse,*] *ex DC. Prod.* i. 624.—Mexic.

GUAREA:—
canaliflora, Reinw. ex Blume, in Flora, vii. (1824) 290; Spreng. Syst. iv. Cur. Post. 251 = Dysoxylum ramiflorum.
cerrana, Vell. Fl. Flum. 149; iv. t. 9 = Lessoniana.
Clausseniana, *C. DC. in Mart. Fl. Bras.* xi. 1. 198.—Am. trop.
coriacea, *C. DC. l. c.* 197.—Am. trop.
costata, *A. Juss. in Mém. Mus. Par.* xix. (1830) 241.—Guian.
cuspidata, *C. DC. in Mart. Fl. Bras.* xi. 1. 193.—Am. trop.
dasmaraas, *C. DC. in DC. Monog. Phan.* i. 549.—Am. trop.
densiflora, Poepp. & Endl. Nov. Gen. et Sp. iii. 40 = Kunthiana.
densiflora, Spreng. Syst. iv. Cur. Post. 251 = Dysoxylum densiflorum.
dispyhonia, Griff. Notul. iv. 508 = Dysoxylum proceruum.
excelsa, H. B. & K. Nov. Gen. et Sp. vii. 227 = humilis.
filiformis, *C. DC. in DC. Monog. Phan.* i. 566.—Am. trop.
foetida, *Blume, in Flora*, vii. (1824) 290, nomen.—Java.
Francavillana, *C. DC. in Mart. Fl. Bras.* xi. 1. 187.—Am. trop.
fulva, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 371.—N. Granat.
Gardneri, *C. DC. in Mart. Fl. Bras.* xi. 1. 193.—Am. trop.
gigantea, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 370.—N. Granat.
glabra, *Vahl, Eclog. Am.* iii. 8.—Am. austr.
glauca, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 372.—N. Granat.
Glaziovii, *C. DC. in Mart. Fl. Bras.* xi. 1. 188.—Am. trop.
Gobara, Buch.-Ham. in Mem. Wern. Soc. vi. (1832) 306 = Dysoxylum procerum.
Goladhara, Wall. Cat. n. 6884 = Dysoxylum binectariferum.
Gotadhara, Buch.-Ham. in Mem. Wern. Soc. vi. (1832) 307 = prec.
grandiflora, Steud. Nom. ed. II. i. 709 = seq.
grandifolia, *DC. Prod.* i. 624.—Ind. occ.
grandis, Wall. Cat. n. 6889 = Dysoxylum grande.
hirsuta, *C. DC. in DC. Monog. Phan.* i. 578.—Am. trop.
Hoffmannsiana, *C. DC. l. c.* 570.—Am. trop.
humilis, *Bert. ex DC. Prod.* i. 624.—Porto Rico.
japurensis, *C. DC. in Mart. Fl. Bras.* xi. 1. 199.—Bras.
Kegelii, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 539 = bijuga.
Kunthiana, *A. Juss. in Mém. Mus. Par.* xix. (1830) 241.—Guian.
Langsdorffiana, *C. DC. in Mart. Fl. Bras.* xi. 1. 192.—Am. trop.
Lessoniana, *A. Juss. in St. Hil. Fl. Bras. Mer.* ii. 84.—Bras.
Lindbergii, *C. DC. in Mart. Fl. Bras.* xi. 1. 192.—Am. trop.
longifoliola, *C. DC. l. c.* 197.—Am. trop.
Lophos, *Vell. Fl. Flum.* 149; iv. t. 8.—Bras.
macrobotrya, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 40.—Peruv.
macrocarpa, Blume, in Flora, vii. (1824) 290 = Dysoxylum macrocarpum.
macrophylla, Vahl, Eclog. Am. iii. 8 = grandifolia.
Martiana, *C. DC. in Mart. Fl. Bras.* xi. 1. 184.—Am. trop.
Mayumana, *C. DC. in A. DC. Monog. Phan.* i. 550.—Am. trop.
magontha, A. Juss. in Mém. Mus. Par. xix. (1830) 241 = grandifolia.
microphylla, Hook. fc. Fl. t. 139 = Elutheria microphylla.
Mikaniana, *C. DC. in Mart. Fl. Bras.* xi. 1. 194.—Am. trop.
mollis, Wall. ex Hiern, in Hook. f. Fl. Brit. Ind. i. 548 = Dysoxylum Hamiltonii.
Moellieri, *C. DC. in Mart. Fl. Bras.* xi. 1. 190.—Am. trop.
multiflora, *A. Juss. in Mém. Mus. Par.* xix. (1830) 241.—Bras.
multijuga, A. Juss. in St. Hil. Fl. Bras. Mer. ii. 83 = trichilioides.
oblonga, Wall. Cat. n. 7262 = Dysoxylum proceruum.

GUARREA :—
oblongifolia, Griff. Notul. iv. 508 = Carapa moluccensis.
obtusifolia, Lam. Encyc. iii. 6 = Trichilia pallida.
pallida, *C. DC. in Mart. Fl. Bras.* xi. 1. 185.—Am. trop.
paniculata, Roxb. Hort. Beng. [98] ; Fl. Ind. ii. 342 = Chisocheton paniculatus.
paniculata, Wall. Cat. n. 6883 = Dysoxylum Hamiltonii.
parsænia, *C. DC. in Mart. Fl. Bras.* xi. 1. 187.—Am. trop.
parvifolia, *C. DC. l. c.* 189.—Am. trop.
pedicellata, *C. DC. l. c.* 196.—Am. trop.
Perrottetiana, *A. Juss. in Mém. Mus. Par.* xix. (1830) 241, 286.—Ind. occ.
Perrottetii, Griseb. Fl. Brit. W. Ind. 181 = Perrottetiana.
petiolulata, *C. DC. in Mart. Fl. Bras.* xi. 1. 197.—Am. trop.
Poeppigii, *C. DC. in DC. Monog. Phan.* i. 568.—Am. trop.
Poeppigii, Triana & Planch. in Ann. Sc. Nat. Sér. V. xv. (1872) 373 = Kunthiana.
Pohlii, *C. DC. in Mart. Fl. Bras.* xi. 1. 195.—Am. trop.
porcata, M. Roem. Syn. Hesper. 118 = tuberculata.
procera, Wall. Cat. n. 1291 = Dysoxylum proceræum.
pubescens, *A. Juss. in Mém. Mus. Par.* xix. (1830) 241.—Guian.
pubiflora, *A. Juss. l. c.*—Bras.
punctata, *C. DC. in DC. Monog. Phan.* i. 575.—Am. trop.
purgans, A. Juss. in St. Hil. Fl. Bras. Mer. ii. 28 = Martiana.
purgans, St. Hil. Pl. Us. Bras. n. 71 = trichilioides, tuberculata.
purpurea, *C. DC. in DC. Monog. Phan.* i. 564.—Am. trop.
ramiflora, *Vent. Choix*, 41.—Porto Rico.
Richardiana, *A. Juss. in Mém. Mus. Par.* xix. (1830) 241, 291.—Guian.
Riedelii, *C. DC. in Mart. Fl. Bras.* xi. 1. 189.—Am. trop.
rosea, *C. DC. l. c.* 186. t. 64.—Am. trop.
Ruagea, *C. DC. in DC. Monog. Phan.* i. 577.—Am. trop.
rubra, *C. DC. in Mart. Fl. Bras.* xi. 1. 191.—Am. trop.
salicifolia, *C. DC. l. c.* 198.—Am. trop.
scabra, *A. Juss. in Mém. Mus. Par.* xix. (1830) 241.—Guian.
Schomburgkii, *C. DC. in DC. Monog. Phan.* i. 565.—Am. trop.
Selloana, *C. DC. in Mart. Fl. Bras.* xi. 1. 198.—Am. trop.
sericea, Spreng. Syst. iv. Cur. Post. 251 = Dysoxylum ramiflorum.
silvatica, *C. DC. in Mart. Fl. Bras.* xi. 1. 195. t. 57.—Am. trop.
sinuata, M. Roem. Syn. Hesper. 119 = trichilioides.
spicata, *C. DC. in Mart. Fl. Bras.* xi. 1. 194.—Am. trop.
spiciflora, *A. Juss. in St. Hil. Fl. Bras. Mer.* ii. 31.—Bras.
Sprucei, *C. DC. in Mart. Fl. Bras.* xi. 1. 196. t. 58.—Am. trop.
suberosa, *C. DC. l. c.* 196.—Am. trop.
surinamensis, Miq. ex C. DC. in DC. Monog. Phan. i. 543 = trichilioides.
Swartzii, DC. Prod. i. 624 = glabra.
Swartzii, Macfad. Fl. Jamaic. 173 = trichilioides.
Trianæ, *C. DC. in A. DC. Monog. Phan.* i. 578.—Am. trop.
trichilioides, *Linn. Mantt.* ii. 228.—Ind. occ.
trunciflora, *C. DC. in DC. Monog. Phan.* i. 571.—Am. trop.
tuberculata, *Vell. Fl. Flum.* 150 ; iv. t. 10.—Bras.
Vahliana, A. Juss. in Mém. Mus. Par. xix. (1830) 240 = glabra.
velutina, *A. Juss. l. c.* 241.—Bras.
verrucosa, *C. DC. in Mart. Fl. Bras.* xi. 1. 198.—Am. trop.
Warmingiana, *C. DC. l. c.* 194.

GUARIA, Dum. Anal. Fam. 48 (1829) = prec.

GUARIBUMA, Cass. in Dict. Sc. Nat. xxxiii. 468, 472 (1824) = Montana, Lian. f. (Compos.).
Acutalbum, Cass. ii. cc. = M. lanatum.
Cinereum, Cass. ii. cc. = M. subalatum.
recurvatum, Cass. ii. cc. = M. retroreum.
sinuatum, Cass. ii. cc. = M. sinuatum.
subspinosum, Cass. ii. cc. = M. subspinosum.

GUAROPSIS, Presl, Epim. Bot. 219 (1849) = Clarkia, Pursh (Onagrar.).
lancifolia, Presl, l. c.—Mexic.

GUATTERIA, Ruiz & Pav. Prod. 85. t. 17 (1794). *ANONACEÆ*, Benth. & Hook. f. i. 28.
CANANGA, Aubl. Pl. Gui. t. 307. t. 244 (1775).
Aberemoa, *Dunal, Monog. Anon.* 126.—Guian.
acutiflora, DC. Syst. i. 508 = Polyalthia Korinti ?
acutiflora, *Mart. Fl. Bras.* xiii. 1. 29.—Bras.
acutiflora, Wall. Cat. n. 6438 = Polyalthia Korinti, Alphonsea zeylanica.
acutifolia, Dunal, Monog. Anon. 184 = Polyalthia Korinti ?
acutifolia, Wight, ex Wall. Cat. n. 6438 = Alphonsea zeylanica.
amplifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 35.—Panama.
anonæfolia, A. DC. in Wall. Cat. n. 6435 (Polyalthiæ sp.).—Ind. or.
apodocarpa, *Mart. Fl. Bras.* xiii. 1. 30.—Bras.
australis, *St. Hil. Fl. Bras. Mer.* i. 37.—Bras.
Badajanska, Roxb. ex Wall. Cat. n. 6448 (Polyalthiæ sp.).—Ind. or.
Bambau, Miq. Fl. Ind. Bat. Suppl. 377 (Polyalthiæ sp.).—Sumatra.
Berteriana, *Spreng. Syst.* ii. 685.—Porto Rico.
bibracteata, *Hemsl. Diagn. Fl. Nov.* 1.—Mexic.
bifaria, A. DC. Mém. Anon. 41 = Polyalthia bifaria.
biglandulosa, Blume, Fl. Jav. Anon. 102. t. 51 = Polyalthia biglandulosa.
blepharophylla, *Mart. Fl. Bras.* xiii. 1. 38.—Bras.
Bragma, Blume, Bijdr. 20 (Polyalthiæ sp.).—Java.
brevipes, *DC. Syst.* i. 502.—Guian.
brevipetala, Miq. Fl. Ind. Bat. Suppl. 381 = Polyalthia macrophylla.
caffra, *Sond. in Harv. & Sond. Fl. Cap.* i. 9.—Afr. austr.
canangioides, Reichb. f. & Zoll. in Linnæa, xxix. (1857–58) 328 (Polyalthiæ sp.).—Ind. or.
Candolleana, *Schlecht. l. c.* ix. (1884) 325.—Bras.
caniflora, *Mart. Fl. Bras.* xiii. 1. 37.—Bras.
Cargadero, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 34.—N. Granat.
caudata, Wall. Cat. n. 6469 = Xylopia caudata.
cauliflora, *Mart. Fl. Bras.* xiii. 1. 35.—Bras.
cerasoides, Dunal, Monog. Anon. 127 = Polyalthia cerasoides.
centrifera, *Triana & Planch, in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 33.—N. Granat.
cinnamomea, Wall. Cat. n. 6444 = Polyalthia cinnamomea.
coffeoides, Thw. Enum. Pl. Zeyl. 10 = Polyalthia coffeoides.
cordata, Dunal, Monog. Anon. 129. t. 30 = Uvaria macrophylla.
Corinti, Steud. Nom. ed. I. 354 = *G. Korinti*.
costata, Hook. f. & Thoms. Fl. Ind. i. 143 = Polyalthia costata.
cuneiformis, Blume, Bijdr. 19 = Polyalthia cuneiformis.
decandra, *G. Don, Gen. Syst.* i. 106.—Peruv.
denticoma, *Mart. Fl. Bras.* xiii. 1. 32.—Bras.
diospyroides, *Baill. Adansonia*, viii. (1868) 269.—Mexic.
elongata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 455.—Guian.
eriantha, Reichb. f. & Zoll. in Linnæa, xxix. (1857–58) 328 (Polyalthiæ sp.).—Malaya.
ertopoda, *DC. Syst.* i. 508.—Peruv.
euphela, Miq. Fl. Ind. Bat. Suppl. 380 (Polyalthiæ sp.).—Sumatra.
excelsa, *Poepp. ex Mart. Fl. Bras.* xiii. 1. 38.—Peruv.
fasciculata, Wall. ex Voigt, Hort. Suburb. Calc. 16 = Polyalthia simiarum.
ferruginea, *St. Hil. Fl. Bras. Mer.* i. 36.—Bras.
flava, *St. Hil. Fl. Bras. Mer.* i. 37.—Bras.
flavicoma, Wall. Cat. n. 7295 = Diospyros flavicoma.
foliosa, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 460.—Guian.
fragrans, Dalz. in Hook. Kew Journ. iii. (1851) 306 = Polyalthia fragrans.
Galeottiana, *Baill. Adansonia*, viii. (1868) 268.—Mexic.
gigantea, Miq. Fl. Ind. Bat. i. 11. 49 (Polyalthiæ sp.).—Java.
glauca, *Ruiz & Pav. Syst. Veg.* i. 146.—Peruv.
globosa, A. DC. in Mém. Anon. 43 = Millusa Rexburghiana.

GUATTERIA :—
Gomeziana, *St. Hil. Fl. Bras. Mer.* i. 36.—Bras.
Gondotiana, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 33.—N. Granat.
grandifolia, Steud. Nom. ed. II. i. 709 = Polyalthia macrophylla ?
gynæcania, *Klotzsch, in annoml. Fauv. et Fl. Gui.* 1168, nomen.—Guiana.
heteropetala, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 460 = Heteropetalum brasiliense.
Hilariana, *Schlecht. in Linnæa*, ix. (1834) 324.—Bras.
hirsuta, *Ruiz & Pav. Syst. Veg.* i. 146.—Peruv.
Hookeri, *St. Hil. & Tuss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 132.—Bras.
hypoleuca, Miq. Fl. Ind. Bat. Suppl. 381 (Polyalthiæ sp.).—Sumatra.
imbricata, Blume, Fl. Jav. Anon. 94 (Polyalthiæ sp.).—Java.
incerta, Blume, l. c. 100 (Polyalthiæ sp.).—Java.
inundata, *Mart. Fl. Bras.* xiii. 1. 36.—Guiana.
Jenkinsii, Hook. f. & Thoms. Fl. Ind. i. 141 = Polyalthia Jenkinsii.
Jurgensenii, *Hemsl. Diagn. Fl. Nov.* 1.—Mexic.
Klotzschiana, *Mart. Fl. Bras.* xiii. 1. 32.—Bras.
Korinti, Dunal, Monog. Anon. 138 = Polyalthia Korinti.
laevigata, *Mart. Fl. Bras.* xiii. 1. 32.—Bras.
laterifiora, Blume, Bijdr. 20 ; Fl. Jav. Anon. 100 (Polyalthiæ sp.).—Java.
laurifolia, Dunal, Monog. Anon. 132. t. 32 (= *Oxandra laurifolia*).—Ind. occ.
laurifolia, J. Grah. Cat. Bomb. Pl. 4 = Bocagea Dalzellii.
laurina, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 32.—N. Granat.
littoralis, Blume, Fl. Jav. Anon. 99 (Polyalthiæ sp.).—Java.
longifolia, Wall. Cat. n. 6442 = Polyalthia longifolia.
longipes, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1863) 32.—N. Granat.
lucida, Boj. Hort. Maurit. 6 = Uvaria lucida.
lucida, *Presl, Rel. Haenk.* ii. 78.—Peruv.
luteo, *St. Hil. Fl. Bras. Mer.* i. 37.—Bras.
lutescens, Pohl, ex Mart. Fl. Bras. xiii. 1. 24 = Duguetia Pohliana.
macrantha, A. DC. Mém. Anon. 42 (Polyalthiæ sp.).—Ind. or.
macrantha, *Presl, Rel. Haenk.* ii. 78.—Mexic.
macrophylla, Blume, Bijdr. 19 = Polyalthia macrophylla.
macropus, *Mart. Fl. Bras.* xiii. 1. 28. t. 8.—Bras.
magnifica, *Ruiz & Pav. Fl. Per.* v. t. 480.—Peruv.
malabarica, Dunal, Monog. Anon. 134 = Uvaria zeylanica.
Martiana, *Schlecht. in Linnæa*, ix. (1834) 326.—Bras.
mayapurensis, *H. B. & K. Nov. Gen. et Sp.* v. 94.—Venezuela.
membranacea, A. DC. Mém. Anon. 41.—Ind. or.
micrantha, A. DC. l. c. 43 = Uvaria micrantha.
microcarpa, *Ruiz & Pav. Fl. Per.* v. t. 479.—Peruv.
montana, DC. Syst. i. 508 = Uvaria zeylanica.
multinervis, Wall. Cat. n. 6445 = Polyalthia cinnamomea.
nigrescens, *Mart. Fl. Bras.* xiii. 1. 31.—Bras.
nitida, A. DC. Mém. Anon. 41 = Polyalthia nitida.
odontopoda, *Mart. Fl. Bras.* xiii. 1. 33.—Bras.
oligocarpa, *Mart. l. c.* 33.—Bras.
Ouregou, *Dunal, Monog. Anon.* 136.—Guian.
ovalis, *Ruiz & Pav. Syst. Veg.* i. 146.—Peruv.
oxycarpa, Miq. Fl. Ind. Bat. i. 1. 49 (Polyalthiæ sp.).—Sumatra.
oxycarpa, Poepp. ex Mart. Fl. Bras. xiii. 1. 33 = blepharophylla.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 378.—Sumatra.
pallida, Blume, Bijdr. 20 ; Fl. Jav. Anon. 97 (Polyalthiæ sp.).—Java.
pallida, Hook. f. & Thoms. Fl. Ind. i. 143 = Polyalthia argentea.
Parvoana, Miq. Fl. Ind. Bat. i. 11. 48 (Polyalthiæ sp.).—Sumatra.
Pavonii, *G. Don, Gen. Syst.* i. 100.—Peruv.
pendula, *Ruiz & Pav. Syst. Veg.* i. 146.—Peruv.
persicæfolia, Hook. f. & Thoms. Fl. Ind. i. 140 = Polyalthia persicæfolia.

GUATTERIA:—
persicifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 38.—N. Granat.
pilosa, G. Don, Gen. Syst. i. 100 = Uvaria hirsuta.
pilosula, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 31.—N. Granat.
placentra, Blume, Bijdr. 21 = Bocagea placentra.
platyphylla, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 35.—N. Granat.
podocarpa, *DC. Syst.* i. 508.—Guian.
Poeppigiana, *Mart. Fl. Bras.* xiii. i. 35.—Bras.
pogonopus, *Mart. i. c.* 34.—Bras.
Pohliana, *Schlecht. in Linnaea*, ix. (1834) 321.—Bras.
polita, Wall. Cat. n. 6450 (Polyalthiæ sp.).—Ind. or. *Pseudod.* Miq. Fl. Ind. Bat. Suppl. 380 (Polyalthiæ sp.).—Sumatra.
prinoides, *Spreng. Syst.* ii. 635.—Ins. S. Doming.
psilopus, *Mart. Fl. Bras.* xiii. i. 37.—Bras.
ptercyca, *Benth. in Hook. Kew Journ.* v. (1853) 8.—Bras.
quindoensis, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 34.—N. Granat.
ramentsirma, Wall. Cat. n. 7294, 8006 = Popowia ramosissima.
rufa, Dunal, Monog. Anon. 129. t. 29 (Polyalthiæ sp.).—Ind. or.; Malaya.
rufa, Lindl. Bot. Reg. t. 836 = Uvaria microcarpa.
rufa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 35.—N. Granat.
Schlechtendaliana, *Mart. Fl. Bras.* xiii. i. 34.—Bras.
Schomburgkiana, *Mart. i. c.* 33.—Guian.
Sellowiana, *Schlecht. in Linnaea*, ix. (1834) 323.—Bras.
sempervirens, Dunal, Monog. Anon. 138 (Polyalthiæ sp.).—Ind. or.
senpaljodalis, Wall. Cat. n. 6446 = Goniothalamus sesquipedalis.
simiarum, Buch.-Ham. in Wall. l. c. 6440 = Polyalthia simiarum.
spatulata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 420 (Polyalthiæ sp.).—Java.
suberosa, Dunal, Monog. Anon. 128 = Polyalthia suberosa.
subsessilis, *Mart. Fl. Bras.* xiii. i. 29.—Bras.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 380 (Polyalthiæ sp.).—Sumatra.
Teysmanni, Miq. l. c. 378 (Polyalthiæ sp.).—Sumatra.
Toralak, Blume, Fl. Jav. Anon. 103 (Polyalthiæ sp.).—Java.
umbilicata, *Dunal, Monog. Anon.* 125. t. 28.—Am. austr.
uncinatiflia, A. DC. Mém. Anon. 41 = Oxymitra uncinatiflia.
velutina, A. DC. l. c. 42 = Miliusa velutina.
venulosissima, *Mart. Reise Bras.* iii. 1257.—Bras.
vestita, *Aletzsch. in Schenk. Fauna et Fl. Gaz.* 979, nomen.—Guian.
villosa, G. Don, Gen. Syst. i. 100 = Miliusa velutina.
villosissima, *St. Hil. Fl. Bras. Mer.* 38.—Bras.
virgata, Dunal, Monog. Anon. 131. t. 31 (= *Guandra virgata*).—Ind. occ.
viridiflora, *G. Don, Gen. Syst.* i. 109.—Peruv.
Willemoetiana, *DC. Prod.* i. 94.—Hab. ?

GUAYABA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 16 = Psidium, Linn. (Myrtac.).

GUAZAMAEA, Cassel, Nat. Fam. 149 (1810) = seq.

GUAZUMA, Plum. ex Adans. Fam. ii. 383 (1763). *STERCULIACEÆ*, Benth. & Hook. f. i. 225.
Bubroma, Schreb. Gen. 513 (1791).
Dirodolobium, Turcz. in Bull. Soc. Nat. Mosc. xxx. (1857) ii. 157.
Blumei, G. Don, Gen. Syst. i. 528 = tomentosa.
Burbroma, Tussac, Fl. Antill. iv. 69 = ulmifolia.
cristis, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 95.—Bras.
crinita, Steud. Nom. ed. II. i. 710, sphalm. = prase.
grandiflora, *G. Don, Gen. Syst.* i. 528.—Bras.
Javira, *G. Don. i. c.*—Bras.
parvifolia, *A. Rich. Ess. Fl. Cub.* 190.—Cuba.
polybotrya, *Cav. Ic.* iii. 51. t. 290.—Mexic.
ruosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 72.—Bras.
GUAZUMA:—
tomentosa, *H. B. & K. Nov. Gen. et Sp.* v. 23.—Am. trop.; Java.
ulmifolia, *Lam. Encyc.* iii. 52.—Am. trop.
ulmifolia, Wall. Cat. n. 1141 = tomentosa.
ulmifolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 72.—Bras.

GUBLERIA, Gaudich. Bot. Voy. Bonite, t. 104 (1844–52) = Nolana, Linn. (Convolv.).
baccata, Gaudich. l. c.—Hab. ?

GUEINZIA, Sond. ex Schott, Aroidese, 10 (1853) = Stylochiton, Lepr. (Aroid.).
natalensis, Sond. ex Schott, l. c. = S. natalense.

GUELDENSTAEDTIA, Fisch. in Mém. Soc. Nat. Mosc. vi. (1839) 170. *LEGUMINOSÆ*, Benth. & Hook. f. i. 606.
cuneata, Benth. in Royle, Illustr. Bot. Himal. 200 = Caragana cuneata.
Delavayi, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 5.—China.
diversifolia, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 462.—China.
Gmelini, *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) 210.—China.
himalaica, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 117.—Reg. himal.
latifolia, Fisch. ex G. Don. Gen. Syst. ii. 278 = Astragalus latifolius.
maritima, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 7.—China.
mirpourensis, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 118.—Ind. bor. occ.
monophylla, *Fisch. ex DC. Prod.* ii. 307.—Sibir. altaic.
multiflora, *Bunge, Enum. Pl. Chin. Bor.* 18.—Reg. Himal.; China.
Olivieriana, Fisch. ex Jaub. & Spach, Illustr. i. 94 (= *Chesneya Olivieri*).—Mesopot.
pauciflora, *Fisch. ex DC. Prod.* ii. 307.—Sibir. altaic.
stenophylla, *Bunge, Enum. Pl. Chin. Bor.* 18.—China.

GUELDENSTAEDTIA, Neck. Elem. ii. 204 (1790) = Burotia, Adans. (Chenop.).

GUENTHERA, Andrz. ex Bess. Enum. Pl. Volh. 88 (1822) = Stenodes, Tourn. (Crucif.).
elongata, Bess. l. c. = S. elongata.

GUENTHERA, Regel, Ind. Sem. Hort. Petrop. (1857) 49; et Gartenfl. (1858) 44 = Xanthocephalum, Willd. (Compos.).
viscosa, Regel, ll. cc. = X. gymnospermoides.

GUENTHERIA, Spreng. Syst. iii. 856 (1826) = Gaillardia, Fouger. (Compos.).
nagapatanica, Spreng. l. c. 449 = G. megapotamica.

GUEPINIA, Bast. Fl. Maine-et-Loire, Suppl. 35 (1812) = Tremella, R. Br. (Crucif.).
fluvialis, Bast. Fl. Fr. Suppl. 896 = T. sudicualis.
Lepidium, DC. l. c. = T. regularis.
sudicualis, Bast. Fl. Maine-et-Loire, Suppl. 35 = T. sudicualis.

GUERSENTIA, Rafin. Sylva Tellur. 158 (1838) = Chrysophyllum, Linn. (Sapot.).
angustifolia, Rafin. l. c. = C. sagotiflolium.
microcarpa, Rafin. l. c. = C. microcarpum.
oleoformis, Rafin. l. c. = C. Cainito ?

GUESMELIA, Walp. Ann. i. 841 (1849), sphalm. = Quesmelia, Gaudich. (Bromel.).

GUETTARDA, Linn. Sp. Pl. 991 (1753). *RUBIACEÆ*, Benth. & Hook. f. ii. 99.
Cadamba, Sonner. Voy. ii. 228. t. 128 (1782).
Dichrostylus, Willd. ex Roem. & Schult. Syst. v. 321 (1819).
Donkelaaria, Linn. Illustr. Hortic. ii. (1856) Misc. 72.
Halesia, P. Br. Hist. Jamaic. 266 (1756).
Hallesia, Scop. Introd. 141 (1777).
Lauxesia, Linn. Syst. ed. XII. 177 (1767).
Lauxiesia, Jacq. Enum. Pl. Carib. 7 (1760).
GUETTARDA:—
Matthiola, Plum. ex Linn. Gen. ed. I. 49 (1737).
Sardinia, Vell. Fl. Flum. 144 (1825); ill. t. 168 (1827).
Tezebraria, Sesse, ex DC. Prod. iv. 458 (1830).
Viviania, Rafin. Specs. i. 117 (1814).
acuminata, Sieber, ex DC. Prod. iv. 460 = Antirrhoea borbonica (in Addend.).
acutiflora, Bartl. ex DC. l. c. 506 = Psychotria acutiflora.
ambigua, Chapm. Fl. S. U. St. 178 = scabra.
amblyca, *DC. Prod.* iv. 455.—Ins. Guadaloup.
Angelica, *Mart. ex Muell. Arg. in Flora*, lviii. (1875) 450.—Bras.
Antirhoea, D. Dietr. Syn. Pl. i. 788 = Antirrhoea borbonica (in Addend.).
arguetas, *Lam. Encyc.* iii. 54.—Guian.
armata, Bartl. ex DC. Prod. iv. 457 = Chomelia filipes.
Aromatis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 27. t. 283.—Peruv.
Balansæana, *Baill. Adansonia*, xii. (1879) 238.—N. Caled.
barbinervis, Sieber, ex DC. Prod. iv. 460 = Antirrhoea borbonica (in Addend.).
Blanchetiana, *Muell. Arg. in Flora*, lviii. (1876) 449.—Bras.
Blodgettii, Shuttl. ex Chapm. Fl. S. U. St. 178 = elliptica.
bracteata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 506.—Cuba.
Brunonis, Wall. Cat. n. 6320 = Timonius Jambosella.
Burchelliana, *Muell. Arg. in Flora*, lviii. (1876) 450.—Bras.
calyptrata, *A. Rich. Fl. Cub. Fanerog.* ii. 21.—Cuba.
coccinea, Aubl. Pl. Gui. i. 317 = Isertia coccinea.
comosa, *Muell. Arg. in Flora*, lviii. (1876) 450.—Bras.
conferta, *Benth. Bot. Voy. Sulph.* 106.—Ins. Cocos.
cordata, *H. B. & K. Nov. Gen. et Sp.* iii. 420.—N. Granat.
coriacea, *Pers. Syn.* i. 200.—Ind. occ.
crispiflora, *Vahl, Eclog. Am.* i. 26. t. 6.—Am. trop.
dasiloata, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 323.—Mexic.
dependens, *DC. Prod.* iv. 458.—Peruv.
dichotoma, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 323.—Mexic.
dioica, D. Dietr. Syn. Pl. i. 788; Balli. Hist. Pl. vii. 377 = Antirrhoea borbonica (in Addend.).
Echinocephalus, *Wright, in Sauv. Fl. Cub.* 56.—Cuba.
elliptica, Spreng. Syst. i. 739 = mollis.
elliptica, *Sw. Prod. Veg. Ind. Occ.* 59.—Ind. occ.
eximia, *Baill. Adansonia*, xii. (1879) 241.—N. Caled.
ferruginea, *Griseb. Cat. Pl. Cub.* 121.—Cuba.
Finlaysoniana, Wall. Cat. n. 6232 = Timonius Finlaysonianus.
Fontanesii, *DC. Prod.* iv. 458.—Hab. ?
franquiacea, Sieber, ex DC. Prod. iv. 460 = Antirrhoea franquiacea.
fusca, *Planch. ex Baill. Adansonia*, xii. (1879) 243.—N. Caled.
grandiflora, *Steud. in Flora*, xxvi. (1843) 763.—Am. trop.
hevaensis, *DC. Prod.* iv. 455.—Cuba.
hornuos, *Blanco, Fl. Filip.* ed. II. 499.—Ins. Philipp.
hirsuta, *Pers. Syn.* i. 200.—Am. trop.
Hostmanneggii, *Muell. Arg. in Flora*, lviii. (1876) 450.—Bras.
heloiensos, *Wright, in Sauv. Fl. Cub.* 55.—Cuba.
hypoleuca, *Baill. Adansonia*, xii. (1879) 240.—N. Caled.
inconspicua, *Sauv. Fl. Vit.* 131.—Ins. Fiji.
iotasis, *Baill. Adansonia*, xii. (1879) 241.—N. Caled.
jasminiflora, *Blanco, Fl. Filip.* ed. I. 122.—Ins. Philipp.
Lindeniana, *A. Rich. Fl. Cub. Fanerog.* ii. 20.—Cuba.
longiflora, *Griseb. Fl. Brit. W. Ind.* 332.—Ind. occ.
lucida, Pers. Syn. i. 201 = Antirrhoea lucida.
macrantha, *Benth. in Hook. Journ. Bot.* iii. (1841) 328.—Guian.
macrocarpa, *Griseb. in Mem. Am. Acad. N. S.* viii. (1863) 507.—Cuba.
Mathiola, *Pers. Syn.* i. 200.—Am. trop.
membranacea, Sieber, ex DC. Prod. iv. 458 = stipulosa.

GUETTARDA :—
membrasacea, *Sw. Prod. Veg. Ind. Occ.* 59.—Ins. S. Doming.
microphylla, *Bartl. ex DC. Prod. iv.* 457.—Ins. Philipp.
Mutisiana, Wall. Cat. n. 6221 = Timonius Jambosella.
mollis, *DC. Prod.* iv. 456.—Ins. S. Doming.
myrtoides, *F. Muell. Fragm.* ix. 184.—Austral.
nouneana, *Baill. Adansonia*, xii. (1879) 248.—N. Caled.
ochreata, *Schlecht. in Linnaea*, xxviii. (1856) 496.—Peruv.
odorata, *Low. Tabl. Encyc.* ii. 219.—Am. austr.
palmicos, *Muell. Arg. in Flora*, lviii. (1875) 450.—Bras.
paniculata, *Bartl. ex DC. Prod.* iv. 457.—Peruv.
parviflora, *Vahl, Eclog. Am.* ii. 26.—Am. trop.
parvifolia, Sw. *Fl. Ind. Occ.* iii. 1958.—Ind. occ.
pedunculariis, Wall. Cat. n. 6222 = Timonius Finlaysoniana, T. Jambosella.
Pervilleana, *Baill. Adansonia*, xii. (1879) 319.—Ins. Seychelles.
platyphylla, *Muell. Arg. in Flora*, lviii. (1875) 449.—Bras.
platypoda, *DC. Prod.* iv. 456.—Bras.
Pohliana, *Muell. Arg. in Flora*, lviii. (1875) 450.—Bras.
polyandra, Blanco, Fl. Filip. ed. II. 500 = Symplocos racemosa.
polygama, Baill. Hist. Pl. vii. 377 = Timonius Rumphii.
polyphragmoides, F. Muell. Fragm. ii. 184 = Timonius Rumphii.
prostrata, Hemsl. Biol. Centr. Am. Bot. ii. 42, sphalm. = sco.
protracta, Bartl. ex DC. Prod. iv. 457 = Antirrhoea protracta.
pubescens, F. Muell. Fragm. ix. 183.—Austral.
ramuliflora, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 182.—Panama.
reniuca, *Pers. Syn.* ii. 200.—Am. austr.
reticulata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 506.—Cuba.
retusa, *Wright, in Sauv. Fl. Cub.* 66.—Cuba.
rhabdocalyx, *Muell. Arg. in Flora*, lviii. (1875) 450.—Bras.
rhamnoides, *Baill. Adansonia*, xii. (1879) 241.—N. Caled.
rigida, *A. Rich. Fl. Cub. Fanerog.* ii. 20.—Cuba.
rugosa, Sw. Prod. Veg. Ind. Occ. 59 = scabra.
scabra, *Lam. Tabl. Encyc.* ii. 218. t. 154. f. 3.—Am. trop.
sericea, *Muell. Arg. in Flora*, lviii. (1875) 450.—Bras.
speciosa, Asbl. Pl. Gui. i. 820 = argentea.
speciosa, Blanco, Fl. Filip. ed. II. 490 = Maries begoniaefolia.
speciosa, *Linn. Sp. Pl.* 991.—As. trop.
speciosa, Van Eeden, ex Steud. Nom. ed. II. i. 710 = Hernandia leucocephala.
splendens, *Baill. Adansonia*, xii. (1879) 289.—N. Caled.
Spruceana, *Muell. Arg. in Flora*, lviii. (1875) 449.—Bras.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 52.—Ins. Tahiti.
tenuiflora, F. Muell. Census, 75 = Antirrhoea tenuiflora.
tomentosa, Pers. Syn. i. 201 (= *Stenostomum tomentosum*).—Jamaic.
umbellata, *Sprengl. Neue Entdeck.* iii. 48.—Bras.
undulata, *Griseb. Cat. Pl. Cub.* 131.—Cuba.
uruguensis, *Chem. & Schlecht. in Linnaea*, iv. (1829) 183.—Bras.
Valenzuelana, *A. Rich. Fl. Cub. Fanerog.* ii. 20.—Cuba.
verruciularis, Blanco, Fl. Filip. ed. I. 738 = Anneslea crassipes.
verticillata, Baill. Hist. Pl. vii. 377 = Danais fragrans.
villosissima, *Cham. & Schlecht. in Linnaea*, iv. (1829) 183.—Bras.
viscosa, *Duchass. & Walp. in Linnaea*, xxiii. (1850) 754.—Ins. Guadeloup.
Wallichiana, *G. Don, Gen. Syst.* iii. 592.—Ind. or *spicata*, Blfb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 188 = mamillicera.
xylosteifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 290.—Venezuela.
zygophylla, *Wright, in Sauv. Fl. Cub.* 66.—Cuba.

GUETTARDELLA, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 197 = **Antirrhoea**, Comm. (Rubiac.).
chinensis, Champ. ex Benth. l. c.—Ins. Hongkong.
philippinensis, Benth. l. c. = A. philippinensis.
putaminosa, Benth. Fl. Austral. iii. 419.—Austral.
sandwicensis, H. Mann, in Proc. Am. Acad. vii. (1867) 170 = Chomelia sandwicensis.

GUETZLAFFIA, Walp. Ann. iii. 213 (1853) = GUTZLAFFIA, Hance = **Strobilanthes**, Blume (Acanth.).

GUEVINA, Juss. Gen. 424 (1789) = **Gevuina**, Molina, (Proteac.).

GUEVINIA, Hort. Par. ex Decne. in Rev. Hortic. Sér. II. iv. (1845–46) 425 = **Celastrus**, Linn.
Abeliana, Hort. Par. l. c. = C. mollis.

GUIABARA, Adans. Fam. ii. 277 (1763) = **Cocciolobis**, Linn. (Polygon.).

GUIBOURTIA, Benn. in Journ. Linn. Soc. i. (1857) 149 = **Copaifera**, Linn. (Legumin.).
copallifera, Benn. l. c. 150 = C. Galbourtiana.

GUICHENOTIA, J. Gay, in Mém. Mus. Par. vii. (1821) 448. t. 20. *STERCULIACEAE*, Benth. & Hook. f. i. 227.
DITOMOSTROPHE, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) ii. 498.
Sturtii, Lindl. Swan River App. 19 (1839).
ledifolia, *J. Gay, l. c.* 449. t. 20.—Austral.
mascrantha, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) ii. 500.—Austral.
micrantha, *Benth. Fl. Austral.* i. 259.—Austral.
Sarotes, *Benth. l. c.* 258.—Austral.
semihastata, *Benth. l. c.*—Austral.

GUIDONIA, P. Br. Hist. Jamaic. 249 (1756) = **Laetia**, Adans. (Hixin.).
Laetia, Loefl. ex Willd. Sp. Pl. ii. 1163 = L. spetala.

GUIDONIA, Plum. ex Adans. Fam. ii. 305 (1763) = **Pisonia**, Jacq. (Samyd.).
Cansiala, Kurz, in Journ As. Soc. Beng. xlvi. (1877) ii. 92 = C. tomentosa.
glomerata, Kurz, l. c. = C. glomerata.
spinosa, Griseb. Fl. Brit W. Ind. 24 = C. brevipes.
tomentosa, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 92 = C. tomentosa.
Vareca, Kurz, ex Kurz, l. c. = C. Vareca.

GUIENZIA, Scand. ex Benth. & Hook. f. Gen. iii. 969 (1883) = GUEINZIA, Scand. = **Stylochiton**, Lepr. (Aroid.).

GUIERA, Adans. ex Juss. Gen. 320 (1789). *COMBRETACEAE*, Benth. & Hook. f. i. 687.
undifora, Reichb. ex DC. Prod. iii. 18 = Combretum aculeatum.
senegalensis, *Lam. Tabl. Encyc.* ii. 486; *Poir. Encyc. Suppl.* ii. 851.—Afr. trop.

GUIINA, Crueg. in Linnaea, xx. (1847) 115 sphalm. = **Quiina**, Aubl. (Guttif.).

GUILANDIA, P. Br. Hist. Jamaic. 228 (1756) = sco.

GUILANDINA, Linn. Gen. ed. I. 124 (1737) = **Caesalpinia**, Linn. (Legumin.).
aculeata, Salisb. Prod. 297 = C. Bonducella.
axillaris, Lam. Encyc. i. 495 = Mezoneuron cucullatum, Pl. Sappan.
Bonduc, Alt. Hort. Kew. ed. II. iii. 32, partim = C. Bonducella.
Bondue, Linn. Sp. Pl. 381 = C. Bondue.
Bondueella, Linn. l. c. ed. II. 548 = C. Bonducella.
ciliata, Berg. ex Wikstr. in Vet. Akad. Handl. Stockh. (1836) 451.—Ind. occ.
dioica, Linn. Sp. Pl. 381 = Gymnocladus canadensis.
echinata, Spreng. Syst. ii. 327 = C. echinata.
gemina, Lour. Fl. Cochinch. 265.—Cochinch.
glabra, Mill. Gard. Dict. ed. VIII. n. 3 = C. Bondue.
inermis, Salisb. Prod. 297 = Gymnocladus canadensis.
macrocarpa, R. Grah. in Wall. Cat. n. 5804 = Mezoneurum cucullatum.
microphylla, DC. Cat. Hort. Monsp. 114.—Hab. ?
Moringa, Linn. Sp. Pl. 381 = Moringa pterygosperma.
Nuga, Linn. l. c. ed. II. 545 = C. Nuga.

GUILANDINA :—
oleosperma, Roxb. ex Wight & Arn. Prod. 281 = Caesalpinia digyna.
paniculata, *Lam.* Encyc. i. 495 = C. Nuga.
parvifolia, Stokes, Bot. Mat. Med. ii. 466 = C. Nuga.
pedicellata, Pritz. Ic. Ind. 515, sphalm. vide *Guilandina pedicellata*.
rotunda, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. IV. 16 = Caesalpinia Nuga *vel* C. Bonduc.
viridiflora, Teijsm. & Binn. in Tijdsch. Nederl. Ind. xxvii. (1864) 59.—N. Guinea.
Wallichiana, R. Grah. in Wall. Cat. n. 5805 = Sindora Wallichii.

GUILDINGIA, Hook. Bot. Misc. i. (1830) 122. t. 80 = **Mouriria**, Juss. (Melastom.).
guidicides, Hook. l. c. = M. rhizophoraefolia.

GUILELMA, Link, Handb. i. 259 (1829) = GUILIELMA, Mart. = **Bactris**, Jacq. (Palm.).

GUILFOYLIA, F. Muell. Fragm. viii. 38 (1873) = **Cadellia**, F. Muell. (Sinarub.).
monostylis, F. Muell. l. c. 34 = C. monostylis.

GUILIELMA, Mart. Palm. Fam. 21 (1824); ex Hist. Nat. Palm. ii. 81. tt. 66, 67 = **Bactris**, Jacq. (Palm.).
caribaea, H. Wendl. in Kerch. Palm. 246 = B. caribaea.
ciliata, H. Wendl. l. c. 246 = B. ciliata.
granatensis, Karst. in Linnaea, xxvii. (1856) 400 = B. granatensis.
insignis, Mart. Palmet. Orb. 71. t. 10. f. 8.—Am. austr.
Macana, Mart. l. c. 74.—Venezuela.
Piritu, Karst. in Linnaea, xxviii. (1856) 397 = B. Piritu.
speciosa, Mart. Hist. Nat. Palm. ii. 82. t. 66, 67 = B. Gasipaës.
tenera, Karst. in Linnaea, xxviii. (1856) 399 = B. tenera.
utilis, Oerst. in Kjoct. Vidensk. Meddel. (1856) 46 = B. utilis.

GUILLAUMIA, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 92. *SCITAMINEAE*, Benth. & Hook. f. iii. 640.
novo-ebudica, *F. Muell. Phyt. New Hebrid.* 20.—N. Hebrid.
purpurata, *Vieill. in Bull. Soc. Linn. Normand.* x. (1866) 93.—N. Caled.

GUILLEMINEA, H. B. & K. Nov. Gen. et Sp. vi. 40. t. 518 (1823). *AMARANTACEAE*, Benth. & Hook. f. iii. 96.
australis, *Hook. f. in Benth. & Hook. f. Gen.* iii. 37.—Reg. Argent.
densa, Moq. in DC. Prod. xiii. ii. 388 = Illecebroides.
Illecebroides, *H. B. & K. Nov. Gen. et Sp.* vi. 41. t. 518.—Ecuador.
Illecebrum, Spreng. Syst. iv. Cur. Post. 108 = Illecebroides.
lanuginosa, *Benth. & Hook. f. Gen.* iii. 37.—Texas.

GUILLEMINIA, Neck. Elem. ii. 182 (1790) = **Votomita**, Aubl. (Inc. Sed.).

GUILLEMINIA, Reichb. Consp. 161 (1828) = **Guillemina**, H. B. & K. (Amarant.).

GUILLIMIA, Reichb. Consp. 196 (1828) = GWILLIMIA, Kotti. = **Magnolia**, Linn.

GUILLONEA, Cass. Notes Crit. 109 (1851) = **Laserpitium**, Linn. (Umbellif.).
cenescyna, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 81 = L. scabrum.
scabra, Cass. Notes Crit. 110 = L. scabrum.

GUINDILIA, Gill. in Hook. Bot. Misc. iii. (1833) 170 = **Valenzuelia**, Bert. (Sapindac.).
trinervis, Gill. l. c. = V. trinervis.

GUIOA, Cav. Ic. iv. 49. t. 378 (1797) = **Cupania**, Linn. (Sapind.).
acutifolia, Raffk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 606.—Malaya; N. Guin.
bijuga, Raffk. l. c. 611 = C. Griffithiana ?

GUIOA:—
crenata, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 615.—N. Caled.
diplopetala, Radlk. l. c. 610 = Cupania regularis.
fusca, Radlk. l. c. 614.—N. Caled.
fusciduta, Radlk. l. c. 609 = C. fusciduta.
glauca, Radlk. l. c. 613 = C. glauca.
gracilis, Radlk. l. c. 609 = C. gracilis!
lasioneura, Radlk. l. c. 608.—Austral.
lenticellata, Cav. Ic. iv. 50 = C. lenticellata.
leptoneura, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 611.—Ins. Celebes.
membranifolia, Radlk. l. c. 614.—N. Guin.
microcephala, Radlk. l. c. 615.—N. Caled.
Minjaillen, Radlk. l. c. 609 = C. Minjaillen.
ovalis, Radlk. l. c. 610.—N. Caled.
patentinervis, Radlk. l. c.—Ins. Moluce.
pectinata, Radlk. l. c. 614.—N. Caled.
Perrottetii, Radlk. l. c. viii. (1878) 302; et ix. (1879) 614.—Ins. Philipp.
pleuropteris, Radlk. l. c. ix. (1879) 611 = C. pleuropteris.
pteropoda, Radlk. l. c. 615.—N. Guin.
pubescens, Radlk. l. c. viii. (1878) 302; et ix. (1879) 613 = C. pallidula.
regularis, Radlk. in Act. Congr. Bot. Amst. 1877 (1879) 82, 119 = C. regularis.
rhoifolia, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 608 = C. rhoifolia.
rigidinucula, Radlk. l. c. 614.—N. Guin.
semiglauca, Radlk. l. c. 608 = C. semiglauca.
squamosa, Radlk. l. c. viii. (1878) 303; et ix. (1879) 609 = C. Griffithiana!
sulfalcata, Radlk. l. c. ix. (1879) 609.—Ins. Navigat.
tenuis, Radlk. l. c. 609.—N. Guin.
villosa, Radlk. l. c. 613.—N. Caled.

GUIRAOA, Cosm. Notes Crit. iii. 97 (1851). *CRUCIFERAE*, Benth. & Hook. f. i. 100.
arvensis, *Cosm. l. c.* 98.—Hispan.

GUIREA, Steud. Nom. ed. II. i. 710 (1840), sphalm. = **Guiera**, Adans. (Combret.).

GUISOTIA, Cass. in Bull. Soc. Philom. (1827) 127; et in Dict. Sc. Nat. lix. 287, 247 (1829). *COMPOSITAE*, Benth. & Hook. f. ii. 389.
RAMTILLA, DC. in Wight, Contrib. 18 (1834).
VERLINGIA, Vis. in Nuov. Sagg. Acc. Sc. Padov. v. (1840) 269.
WERRINUWA, Heyne, Tracts on India, 49 (1814).
abyssinica, *Cass. in Dict. Sc. Nat.* lix. 248.—Afr. trop.
bidentoides, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 386.—Afr. trop.
discoidea, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 160 = Sclerocarpus discoideus.
oleifera, DC. Segt. Not. Pl. Rar. Jard. Genèv. 5 = abyssinica.
Schimperi, *Sch. Bip. in Walp. Rep.* vi. 153.—Abyss.
Schultzii, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 407. —Abyss.
villosa, *Sch. Bip. ex A. Rich. l. c.* 406.—Abyss.

GULDAENSTEDTIA, A. Juss. in Orb. Dict. vii. 270 (1849) = **Gualdenstaedtia**, Fisch. & Mey. (Leguminos.).

GULDENSTAEDTIA, Dum. Anal. Fam. 40 (1829) = praec.

GULIELMA, Spreng. Syst. ii. 141 (1825) = GUILIELMA, Mart. = **Bactris**, Jacq. (Palm.).

GULUBIA, Becc. in Ann. Jard. Buitenz. ii. (1885) 181. *PALMAE*.
costata, *Becc. l. c.* 184.—Malaya.
moluccana, *Becc. l. c.* 181.—Malaya.

GUMIFERA, Rafin. Sylva Tellur. 118 (1838) = **Acacia**, Willd. (Leguminos.).
arabica, *nilotica*, *vera*, Rafin. l. c. = A. arabica.

GUMILLAEA, Roem. & Schult. Syst. vi. 297 (1820) = seq.

GUMILLEA, Ruiz & Pav. Prod. 62. t. 7 (1794). *SAXIFRAGEAE*, Benth. & Hook. f. i. 651.
auriculata, *Ruiz & Pav. l. c.*; *et Fl. Per.* iii. 23.—Peruv.

GUMIRA, Rumph. ex Hask. in Flora, xxv. (1842) ii. Beibl. 36 = **Pterona**, Linn. (Verbenac.).
domestica, Hask. l. c. = P. foetida, parasitica.
foetida, Hask. Cat. Hort. Bog. Alt. 135 = P. foetida.
integrifolia, Hask. l. c. = P. corymbosa.
parasitica, Hask. l. c. = P. parasitica.

GUMSIA, Buch.-Ham. ex Wall. Cat. p. 157 (1829) = **Hiroldsana**, DC. (Sterculiac.).
chlorocarpa, Buch.-Ham. l. c. n. 1178 E = E. spectabilis.
hirta, Buch.-Ham. l. c. n. 1174 B = E. spectabilis.
tiliacea, Wall. Cat. n. 1176 D = E. spectabilis.

GUMTEOLIS, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 87 (1825) = **Contranthera**, R. Br. (Scrophul.).

GUNDELIA, [Tourn.] Linn. Gen. ed. I. 242 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 461.
GUNDELSHEIMERA, Cass. in Dict. Sc. Nat. lvii. 344 (1828).
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = Tournefortii. Tournefortii, *Linn. Sp. Pl.* i. 814.—Syria; As. Min.; Persia.

GUNDELSHEIMERA, Cass. in Dict. Sc. Nat. lvii. 344 (1828) = prec.

GUNDLACHIA, A. Gray, in Proc. Am. Acad. xvi. (1881) 100. *COMPOSITAE*.
domingensis, *A. Gray, l. c.*—Ins. S. Doming.

GUNISANTHUS, A. DC. in DC. Prod. viii. 219 (1844) = **Diospyros**, Linn. (Ebenac.).
mollis, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 88 = D. martabanica.
pilosula, A. DC. in DC. Prod. viii. 220 = D. pilosula.

GUNNERA, Linn. Mant. i. 16 (1767). *HALORAGEAE*, Benth. & Hook. f. i. 676.
DYSEMONE, Soland. ex Forst. f. in Comm. Gotting. iv. (1789) 45, nomen.
GUNNEROPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 193.
MILLIGANIA, Hook. f. in Hook. Ic. Pl. t. 929 (1840).
MIRANDRA, Comm. ex Juss. Gen. 395 (1789).
MISANDROPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 192.
PAMPE, Molina, Sagg. Chil. 148 (1782).
PERPENSUM, Burm. f. Fl. Cap. Prod. 30 (err. typ. 36) (1768).
PSEUDO-GUNNERA, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 193.
Berteroi, *Phil. in Anal. Univ. Chil.* (1872) 714.—Chili.
bracteata, *Steud. ex Benn. Pl. Jav. Rar.* 75.—Chili.
brephogea, *Linden & André, Illustr. Hortic.* xix. (1872) 567. t. 111.—N. Granat.
calthaefolia, Sallab. Prod. 11 = perpensa.
chilensis, *Lam. Encyc.* iii. 61.—Chili.
commutata, *Blume, Mus. Bot. Lugd. Bat.* ii. 100.—Chili.
cordifolia, *Hook. f. Fl. Tasman.* 125.—Tasmania.
densiflora, *Hook. f. Handb. N. Zeal. Fl.* 68.—N. Zel.
erosa, Blume, in Verh. Batav. Gen. x. (1825) 81 = macrophylla.
fallaciandra, Hook. Ic. Pl. t. 489, 490 = magellanica.
glabra, *Phil. in Bot. Zeit.* xiv. (1856) 643.—Ins. J. Fernand.
Hamiltonii, *T. Kirk, ex W. Hem. in Trans. N. Z. Inst.* xvii. (1885) 292.—N. Zel.
insignis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 189; et *DC. in DC. Prod.* xvi. ii. 597.—Chili.
insularis, Phil. in Bot. Zeit. xiv. (1856) 642 = bracteata.
integrifolia, Blume, Mus. Bot. Lugd. Bat. ii. 171 = magellanica.
lobata, *Hook. f. Fl. Antarct.* ii. 374.—Reg. Magellan.
macrophylla, *Blume, Bijdr.* 518.—Java.
magellanica, *Lam. Encyc.* iii. 61.—Reg. Magellan.
manicata, *Linden, Illustr. Hortic.* xx. (1873) 156.—Bras.
monoica, *Raoul, in Ann. Sc. Nat. Sér. III.* ii. (1844) 117.—N. Zel.
peltata, *Phil. in Bot. Zeit.* xiv. (1856) 643.—Ins. J. Fernand.
perpensa, *Linn. Mant.* i. 121.—Ad. austr.
petaloidea, *Gaudich. in Freyc. Voy. Bot.* 512.—Ins. Sandvic.

GUNNERA:—
pilosa, Kunth, Syn. Pl. i. 376 = chilensis.
plicata, Vahl, Enum. i. 209 = magellanica.
prorepens, *Hook. f. Fl. N. Zel.* i. 96.—N. Zel.
reniformis, C. Gay, ex Blume, Mus. Bot. Lugd. Bat. ii. 108 = magellanica.
scabra, Phil. in Bot. Zeit. xiv. (1856) 643 = bracteata.
scabra, Ruiz & Pav. Fl. Peruv. i. 29. t. 44. f. a = chilensis.
semichylla, Mirb. Hist. Nat. Pl. ed. II. x. (1808) 141 = Frankea sonchifolia.
strigosa, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 322. —N. Zel.
tinctoria, Mirb. Hist. Nat. Pl. ed. II. x. (1808) 141 = chilensis.

GUNNEROPSIS, Oerst. in Kjoeb. Vidensk. Meddel. 198 (1857) = **Gunnera**, Linn. (Halorag.).
petaloidea, Oerst. l. c. = G. petaloidea.

GUNNIA, Lindl. Bot. Reg. sub t. 1899 (1834) = **Sarcochilus**, R. Br. (Orchid.).
australis, Lindl. l. c. = S. parviflorus.
picta, Lindl. Bot. Reg. (1838) Misc. 45 = S. olivaceus.

GUNNIA, F. Muell. Rep. Babb. Exped. 9 (1858). *FICOIDEAE*, Benth. & Hook. f. i. 856.
Drummondii, *Benth. Fl. Austral.* iii. 237.—Austral. septifraga, *F. Muell. Rep. Babb. Exped.* 9.—Austral.

GUNTHERA, Steud. Nom. ed. II. i. 711 (1840), sphalm. = GUENTHERA, Andrz. = **Brassica**, Tourn. (Crucif.).

GUNTHERIA, Benth. & Hook. f. Gen. ii. 414 (1873), sphalm. = GUENTHERIA, Spreng. = **Gaultheria**, Forget. (Compos.).

GUPA, Jeanne St. Hil. Expos. ii. 267 (1805) = **Goupia**, Aubl. (Celastr.).

GURANIA, Cogn. in Bull. Soc. Bot. Belg. xiv. (1875) 239. *CUCURBITACEAE*.
acuminata, *Cogn. Diagn. Cucurb.* i. 31 [*in Mém. Cour. Acad. Belg.* 8vo, xxvii. (1877)].—Bras.
Andreana, *Cogn. in Bull. Acad. Belg.* Sér. II. xlix. (1880) 194.—Ecuador.
Arabidae, Cogn. Diagn. Cucurb. i. 17 = multiflora.
Balfourana, *Cogn. in DC. Monog. Phan.* iii. 714.—Peruv.
bevriflora, *Cogn. Diagn. Cucurb.* i. 40.—Bras.
Candolleana, *Cogn. l. c.* 42.—Bras.
capitata, *Cogn. l. c.* 18.—Peruv.
cissoides, *Cogn. l. c.*—Bras.
coccinea, *Cogn. l. c.* 42.—Panama.
costaricensis, *Cogn. l. c.* 37.—Costa Rica.
diversifolia, *Cogn. l. c.* 42.—Bras.
Dumortieri, *Cogn. l. c.* 41.—Bras.
Duraudii, *Cogn. in Bull. Soc. Bot. Belg.* xvii. (1878) 296, in nota.—Guian.
erianthe, *Cogn. Diagn. Cucurb.* i. 16.—Mexic.
Francavillana, *Cogn. l. c.* 40.—Bras.
guianensis, *Cogn. l. c.* 18.—Bras.
inaequalis, *Cogn. l. c.* 32.—Bras.
Kegelians, *Cogn. l. c.* 16.—Bras.
Klotzschiana, *Cogn. l. c.* 28.—Bras.
lanata, *Cogn. l. c.* 26.—Peruv.
Levyana, *Cogn. l. c.*—Nicaragua.
ligulosa, *Cogn. l. c.* 27.—Bras.
Lindiana, *Cogn. l. c.* 43.—Bras.
macrophylla, *Cogn. l. c.* 28.—N. Granat.
Makoyana, *Cogn. l. c.* 17.—Guatemala.
Martiana, *Cogn. l. c.* 24.—Bras.
multiflora, *Cogn. l. c.* 16.—Bras.
multiflora, Cogn. in Mart. Fl. Bras. vi. pt. 49. t. 14 = subumbellata.
neo-granatensis, *Cogn. Diagn. Cucurb.* i. 34.—N. Granat.
ovata, *Cogn. l. c.* 26.—Bras.
parviflora, *Cogn. l. c.* 36.—Peruv.
Paulista, *Cogn. l. c.* 28.—Bras.
Pseudo-spinulosa, *Cogn. l. c.* 26.—Bras.
reticulata, *Cogn. l. c.* 25.—Bras.
rudgiifa, *Cogn. l. c.* 24.—Bras.
Sagotiana, *Cogn. l. c.* 29.—Guian.
Sarmentiana, *Cogn. l. c.* 25.—Panama.
Schwackeana, *Cogn. l. c.* 30, in obs.—Bras.
sinuata, *Cogn. l. c.* 19.—Bras.
speciosa, *Cogn. l. c.*—Peruv.

GURANIA :—
spinulosa, *Cogn. Diagn. Cucurb.* i. 17.—Bras.
Spruceana, *Cogn. l. c.* = 36.—Ecuador.
subumbellata, *Cogn. l. c.* 17.—Bras.
subumbellata, Cogn. in Mart. Fl. Bras. vi. IV. 56 = Durandii.
sylvatica, *Cogn. Diagn. Cucurb.* i. 25.—Bras.
trialata, *Cogn. l. c.* 29.—Bras.
tricuspidata, *Cogn. l. c.* 35.—Bras.
tuberosa, *Cogn. l. c.* 30.—Ecuador.
velutina, *Cogn. l. c.* 37.—Bras.
villosa, *Cogn. l. c.* 35.—Bras.
Wagneriana, *Cogn. l. c.* 17.—N. Granat.
Wawrsi, *Cogn. l. c.* 39.—Bras.

GURLTIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 138 ; et Abh. Akad. Berl. (1855) 56. t. 4 A = **Begonia**, Linn.
Bouchéana, Klotzsch, in Ann. Sc. Nat. Sér. IV. vi (1856) 351 = B. Bouchéana.
Meyeri, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 138 = B. tomentosa.
rigida, Klotzsch, in Abh. Akad. Berl. (1855) 57 = B. rigida.
strigulosa, Hassk. Hort. Bog. Desc. 328 = B. strigulosa.
tomentosa, Klotzsch, in Abh. Akad. Berl. (1855) 57 = B. tomentosa.

GUROA, Steud. Nom. ed. II. i. 711 (1840) = seq.

GURUA, Buch.-Ham. ex Wight, Contrib. 65 (1834) = **Pinlayscina**, Wall. (Asclepiad.).
obovata, Buch.-Ham. l. c. = P. obovata.

GUSMANIA, Remy, in C. Gay, Fl. Chil. iv. 12 (1849) = **Erigeron**, Linn. (Compos.).
chilensis, Remy, l. c. = E. Remyanum.

GUSMANNIA, Juss. in Dict. Sc. Nat. xx. 99 (1820) = **Guzmania**, Ruiz & Pav. (Bromeliac.).

GUSSONEA, J. & C. Presl, Rel. Haenk. i. 183. t. 33 (1828) = **Pimbristylis**, Vahl (Cyperac.).
cyperoides, J. & C. Presl, l. c. = P. cyperoides.
pauciflora, Brongn. Voy. Coq. Bot. t. 34 = P. pauciflora.

GUSSONEA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 67 = **Angrascum**, Bory (Orchid.).
aphylla, A. Rich. l. c. 68 = A. aphyllum.
cornuta, Ridl. in Journ. Bot. xxiii. (1885) 310 = A. cornutum.
exilis, Ridl. in Journ. Linn. Soc. xxi. (1885) 493 = *Microcoelia exilis*).—Madag.
Gilpinae, Ridl. l. c. 491 = A. Gilpinae.
globulosa, Ridl. l. c. = A. Guyonianum.
physophora, Ridl. l. c. 492 = A. physophorum.

GUSSONIA, D. Dietr. Syn. Pl. ii. 925 (1840), sphalm. = **Cussonia**, Thunb. (Araliac.).

GUSSONIA, Spreng. Gen. ii. 664 (1831) = GUSSONEA, Rich. = **Angrascum**, Bory (Orchid.).

GUSSONIA, Spreng. Neue Entdeck. ii. 119. t. 2 (1821) = **Sebastiania**, Spreng. (Euphorb.).
concolor, Spreng. l. c. 120 = Actinostemon caribaeus.
cuneata, Klotzsch, in Schomb. Fauna et Fl. Gui. 1184, nomen.—Guian.
discolor, Spreng. Neue Entdeck. ii. 119 = Exoecaria discolor.
grandifolia, Klotzsch, in Schomb. Fauna et Fl. Gui. 1184, nomen.—Guian.
serrulata, Miq. in Linnaea, xix. (1847) 446 = S. Gaudichaudii.

GUSTAVIA, Linn. Amoen. Acad. viii. 266. t. 5 (1785). *MYRTACEAE*, Benth. & Hook. f. i. 731.
PEREIRA, Ambl. Fl. Gui. 467 (1775).
SPALLANZANIA, Neck. Elem. ii. 79 (1790).
TESCHMEYERIA, Scop. Introd. 287 (1777).
alata, *Span. in Linnaea*, xv. (1841) 304.—Java.
augusta, J. F. Gmel. Syst. 1038, sphalm. = augusta.
augustifolia, *Benth. Bot. Voy. Sulph.* 99.—N. Granat.
axillaris, *Miers, in Trans. Linn. Soc.* xxx. (1874) 179. —Ind. occ.
augusta, DC. Prod. iii. 260 = speciosa.
augusta, *Linn. Amoen. Acad.* viii. 266.—N. Granat.

GUSTAVIA :—
augusta, Ruiz & Pav. Fl. Per. v. 117. t. 485 = Kuitiana.
brasiliana, *DC. Prod.* iii. 290.—Bras.
brasiliensis, *Morr. in Belg. Hortic.* xvi. (1866) 201.—Bras.
calycaris, *Miers, in Trans. Linn. Soc.* xxx. (1874) 185. —Bras.
conferta, Spruce, ex Berg, in Mart. Fl. Bras. xiv. i. 471 = augusta.
dubia, *Berg, in Linnaea*, xxvii. (1854) 446.—N. Granat.
fastuosa, Berg, in Mart. Fl. Bras. xiv. i. 478 = pterocarpus.
fastuosa, Mart. ex DC. Prod. iii. 290 = brasiliana.
fastuosa, Spruce, ex Berg, in Mart. Fl. Bras. xiv. i. 471 = augusta.
fastuosa, *Willd. Sp. Pl.* iii. 487.—Guian.
globosa, *Span. in Linnaea*, xv. (1841) 304, nomen.—Java.
gracillima, *Miers, in Trans. Linn. Soc.* xxx. (1874) 180.—N. Granat.
hexapetala, Sm. in Rees, Cycl. xvii. n. 2 = fastuosa.
insignis, Linden, ex Hook. Bot. Mag. t. 5069 = superba.
insignis, Willd. ex Berg, in Mart. Fl. Bras. xiv. i. 471 = augusta.
laciniosa, *Miers, in Trans. Linn. Soc.* xxx. (1874) 175. —Bras.
latifolia, *Miers*, l. c. 182.—Panama.
longifolia, *Poepp. ex Berg, in Mart. Fl. Bras.* xiv. i. 472.—Peruv.
Macrogravians, *Miers, in Trans. Linn. Soc.* xxx. (1874) 188.—Bras.
meisocarpa, Gaertn. Fruct. ii. 364. t. 188 = augusta.
Poeppigiana, *Berg, in Mart. Fl. Bras.* xiv. i. 471.—Bras.
pterocarpa, *Poit. in Mém. Mus. Par.* xiii. (1825) 158. t. 6.—Guian.
pubescens, *Kuiz, ex Berg, in Linnaea*, xxvii. (1854) 448.—Ecuador.
pulchra, *Miers, in Trans. Linn. Soc.* xxx. (1874) 182. —Am. trop.
Kuitiana, *Berg, in Linnaea*, xxvii. (1854) 444.—Peruv.
speciosa, Benth. Pl. Hartw. 176 = latifolia.
speciosa, *DC. Prod.* iii. 289.—N. Granat.
superba, *Berg, in Linnaea*, xxvii. (1854) 444.—N. Granat.
tetrapetala, Raeusch. Nom. ed. III. 202 = Griss Antillensis.
tetrapetala, Stokes, Bot. Mat. Med. iii. 549 = augusta.
Theophrasta, *Linden, Illustr. Hortic.* xx. (1873) 74.—Hab. ?
urcolata, Poit. in Mém. Mus. Par. xiii. (1825) 156. t. 5 = augusta.
valida, DC. Prod. iii. 290 = Planchonia littoralis.
verticillata, *Miers, in Trans. Linn. Soc.* xxx. (1874) 181.—N. Granat.

GUTENBERGIA, Sch. Bip. in Gedenkb. iv. Jubelf. Buchdr. (1840) 139. t. 4. *COMPOSITAE*, Benth. & Hook. f. ii. 324.
abyssinica, Sch. Bip. in Walp. Ann. ii. 808 = Ruepellii.
cordifolia, *Benth. ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 89. t. 56.—Afr. trop.
longifolia, *Steetz, in Peters, Reise Mossamb. Bot.* 349. —Afr. trop.
macrocephala, *Oliver & Hiern, in Fl. Trop. Afr.* iii. 264.—Afr. trop.
nigritiana, *Oliver & Hiern, l. c.* 264.—Afr. trop.
polycephala, *Oliver & Hiern, in Journ. Linn. Soc.* xv. (1876) 95.—Afr. trop.
Peterii, *Steetz, in Peters, Reise Mossamb. Bot.* 348. —Afr. trop.
Rueppellii, *Sch. Bip. in Gedenkb. iv. Jubelf. Buchdr.* (1840) 140. t. 4.—Abyssin.

GUTENBERGIA, Walp. Ann. i. 374 (1849) = GUTTENBERGIA, Zoll. = **Morinda**, Linn. (Rubiac.).

GUTHNICKIA, Regel, in Flora, xxxii. (1849) 179 = **Achimenes**, P. Br. (Gesneрac.).
atrosanguinea, Hort. ex Regel, in Cat. Hort. Turic. (1861).—Cf. Linnaea, xxiv. (1851) 196 = A. foliosa.
cardinalis, Hanst. in Linnaea, xxvi. (1853) 219. t. 1. f. 6 = A. cardinalis.

GUTHNICKIA :—
foliolosa, Hanst. in Linnaea xxvii. (1849) 712 = Achimenes foliosa.
foliosa, Liebm. in Vidensk. Selsk. Skr. V. v. (1861) 97 = A. foliosa.
mimuloflora, Regel, in Flora, xxxii. (1849) 179 = A. amoena.

GUTTERIA, Bolus, in Hook. Ic. Pl. t. 1161 (1876). *PASSIFLOREAE*.
capensis, *Bolus, l. c.*—Afr. austr.

GUTTERREIA, Lag. Gen. et Sp. Nov. 30 (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 250.
BRACHYACHYRIS, Spreng. Syst. iii. 574 (1826).
BRACHYRIS, Nutt. Gen. Am. ii. 163 (1818).
HEMIACHYRIS, DC. Prod. v. 313 (1836).
ODONTOCARPHA, DC. l. c. 71 (1836).
Alamani, A. Gray, Pl. Wright. i. 91 = Xanthocephalum Alamani.
baccharoides, *Sch. Bip. in Flora*, xxxviii. (1855) 115. —Reg. Magellan.
Berlandieri, *A. Gray, in Proc. Am. Acad.* xv. (1880) 81.—Mexic.
californica, *Torr. & Gray, Fl. N. Am.* ii. 198.—Am. bor. occ.
Coulteri, *Hemsl. Biol. Centr. Am. Bot.* ii. 111.—Mexic.
cupressiformis, *Sch. Bip. in Flora*, xxxviii. (1855) 114. —Reg. Magellan.
divaricata, *Torr. & Gray, Fl. N. Am.* ii. 193 = Euthamiae.
eriocarpa, *A. Gray, Pl. Wright.* i. 94.—Am. bor. occ.
Euthamiae, *Torr. & Gray, Fl. N. Am.* ii. 198.—Am. bor. occ.
Gilliesii, *Griseb. in Goott. Abh.* xix. (1874) 173.—Reg. Argent.
glutinosa, *Sch. Bip. in Flora*, xxxviii. (1855) 115 ; *Benth. & Hook. f. Gen.* ii. 250.—Mexic.
gymnospermoides, A. Gray, Pl. Wright. ii. 79 = Xanthocephalum gymnospermoides.
Haenkei, *Sch. Bip. in Flora*, xxxviii. (1855) 115.—Mexic.
laricifolia, *D. Don, in Hook. Comp. Bot. Mag.* ii. (1836) 51.—Chili.
lecifolia, *Griseb. in Goott. Abh.* xix. (1874) 173.—Reg. Argent.
Linchneriana, Scheele, in Linnaea, xxii. (1849) 351 = Amphiachyris dracunculoides.
linearifolia, A. Gray, in Bot. Calif i. 302 = californica.
linearifolia, *Lag. Gen. et Sp. Nov.* 30.—Mexic.
microcephala, A. Gray, Pl. Fendl. 74, adnot. = Euthamiae.
microphylla, Durand & Hilg. in Journ. Acad. Sc. Philad. iii. (1855-58) 40 = Euthamiae.
Nezsena, *Sch. Bip. in Flora*, xxxviii. (1855) 113.—Mexic.
paniculata, *A. Gray, Pl. Wright.* ii. 78.—Am. bor. occ.
repens, *Griseb. in Goott. Abh.* xix. (1874) 173.—Reg. Argent.
sphaerocephala, *A. Gray, Pl Fendl.* 73.—Am. bor. occ.
texana, *Torr. & Gray, Fl. N. Am.* ii. 194.—Am. bor. occ.
Wrightii, A. Gray, Pl. Wright. ii. 78 = Xanthocephalum Wrightii.

GUTTENBERGIA, Zoll. & Mor. in Nat. en Genees. Arch. Nederl. Indie, ii. (1845) 2 = **Morinda**, Linn. (Rubiac.).
umbellata, Zoll. & Mor. l. c. 3 = M. polyneura ?

GUTZLAFFIA, Hance, in Hook. Kew Journ. i. (1849) 142 = **Strobilanthes**, Blume (Acanthac.).
apricа, Hance, l. c. = S. apricus.

GUYONIA, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 149. *MELASTOMACEAE*, Benth. & Hook. f. i. 745.
ciliata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 448.—Afr. trop.
tenella, *Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 150. t. 6.—Afr. trop.

GUZMANIA, Ruiz & Pav. Fl. Per. iii. 87. t. 261 (1802). *BROMELIACEAE*, Benth. & Hook. f. iii. 669.
comosa, *Bert. ex Steud. Nom.* ed. II. i. 711 (= tricolor ?).—Porto Rico; Jamaica.
Devansayana, *E. Morr. in Belg. Hortic.* xxxiii. (1883) 113. t. 8–9.—Ecuador.
erythrolepis, *Brongn. ex Planch. in Fl. des Serres*, Sér. II. i. (1856) 25. t. 1089.—Cuba.
Melinoni, Hort.—Cf. Gard. Chron. (1883) ii. 727; Morr. in Regel, Gartenfl. (1885) 116 (Melinoki) (= Carguatae sp.).—Guiana.
picta, Lem. Jard. Fleur. iv. (1854) Misc. 60 = Karatas fulgens.
sympaganthera, Beer, Bromel. 103 = tricolor.
tricolor, *Ruiz & Pav. Fl. Per.* iii. 88. t. 261.—Am. trop.

GUZMANNIA, Phil. f. Cat. Pl. Vasc. Chil. 166 (1881) = GUSMANIA, Remy = **Brigeron**, Linn. (Compos.).

GWILLIMIA, Rottl. ex DC. Syst. i. 455 (1818) = **Magnolia**, Linn.
indica, Rottl. l. c. 458 = M. pumila.

GYAS, Salisb. in Trans. Hort. Soc. i. (1812) 299 = **Bletia**, Ruiz & Pav. (Orchid.).
florida, Salisb. l. c. = B. florida.
humilis, Salisb. l. c. 300 = B. hyacinthina.
verecunda, Salisb. l. c. 299 = B. verecunda.

GYAXIS, Salisb. Gen. Pl. Fragm. 131 (1866) = **Haemanthus**, Linn. (Amaryllid.).
punicea, Salisb. l. c. = H. punicea.

GYBIANTHUS, Pritz. Ic. Ind. i. 516 (1855) sphalm. = **Cybianthus**, Mart. (Myrsin.).

GYMNACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836) = **Sclerocalyx**, Nees (Acanthac.).
petiolaris, Nees, l. c. = S. mexicanus.

GYMNACANTHUS, Oerst. in Kjoeb. Videnskab. Meddel. (1854) 126 = **Ruellia**, Plum. (Acanth.).
campestris, Oerst. l. c. = R. campestris.
geminiflora, Oerst. l. c. 127 = R. geminiflora.
humilis, Oerst. l. c. = R. humilis.
Schlechtendalianus, Oerst. l. c. 126 = R. Schlechtendaliana.

GYMNACHAENA, Reichb. ex DC. Prod. vi. 264 (1837) = **Perrotriche**, Cass. (Compos.).
brunioides, Reichb. ex DC. l. c. = P. tortilis.

GYMNACTIS, Pfeiff. Nom. i. ii. 1520 (1874) = **Glossogyne**, Cass. (Compos.).

GYMNADENIA, R. Br. in Ait. Hort. Kew. ed. II. v. 191 (1813) = **Habenaria**, Willd. (Orchid.).
acuta Reichb. f. Otia Bot. Hamb. 32.—Cambogia.
affinis, Reichb. f. l. c. 33 = H. affinis.
albida, Rich. in Mém. Mus. Par. iv. (1818) 57 = H. albida.
angustifolia, Spreng. Syst. iii. 693 = Orchis angustifolia, *Bieb.*
aniziobula, Peterm. Deutschl. Fl. ii. 548 = H. conopsea, *Benth.*
bifolia, Meyer, Chlor. Hannov. 540 = H. bifolia.
Boryana, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 26 = Cynorchis Boryana.
bracteata, Presl, ex Steud. Nom. ed. II. i. 711 = H. viridis.
chloranthа, Ambros. Fl. Tir. Austr. i. 705 = H. bifolia.
Chusua, Lindl. in Wall. Cat. n. 7058 = Orchis Chusua.
conigera, Reichb. Fl. Germ. Excurs. 121 = H. conopsea, *Benth.*
Commersoniana, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 26 (= *Peristylus Commersonianus*).—Ins. Borbon.
conica, Lindl. Gen. et Sp. Orch. 280.—Am. bor.
conopea, Auct. Gall.—Cf. Gren. Fl. Ch. Jurass. 751; Franch. & Sav. Enum. Pl. Jap. ii. 39 = conopsea.
conopsea, R. Br. in Ait. Hort. Kew. ed. II. v. 191 = H. conopsea, *Benth.*
conopsea, Ledeb. Fl. Ross. iv. 65, sphalm. vide *conigera*.

GYMNADENIA :—
cucullata, Rich. in Mém. Mus. Par. iv. (1818) 57.—Podol.; Sibir.
cylindrostachya, Lindl. Wall. Cat. n. 7056 (= *Platanthera Orchidis*).—Ind. or.
denudata, A. Dietr. in Allg. Gartenz. vii. (1839) 170 = Habenaria conopsea, *Benth.*
diphylla, Link, Handb. i. 248 = H. cordata.
elata, Lindl. Gen. et Sp. Orch. 277.—Numidia.
fastigiata, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 23 = Cynorchis fastigiata.
flava, Lindl. Gen. et. Sp. Orch. 279 = H. integra.
flexuosa, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 25 (= *Peristylus flexuosus*).—Ins. Mascar.
Friwaldii, Reichb. Ic. Fl. Germ. xiii. t. 454 = sec.
Friwaldskyana, Hampe, in Flora, xx (1837) 290.—Rumelia.
Galeandra, Reichb. f. Otia Bot. Hamb. 33 = H. Galeandra.
gracilis, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 207.—Japon.
gracillima, Schur, in Oestr. Bot. Zeitschr. xxi. (1871) 44 = H. conopsea ! *Benth.*
graminifolia, Reichb. f. in Bot. Zeit. xxxvi. (1878) 75.—Japon.
Helfers, Reichb. f. in Flora, lv. (1872) 276.—Ind. or.; Burma.
humilis, Lindl. Gen. et Sp. Orch. i. 276.—Italia.
hybrida, Schur, Enum. Pl. Transs. 644 = H. conopsea, *Benth.*
hyperborea, Link, Handb. i. 242 = H. hyperborea.
interrupta, Peterm. Fl. Bienitz, 80 = H. conopsea, *Benth.*
Keiskei, Franch. & Sav. Enum. Pl. Jap. ii. 518 = H. Keiskei.
Keiskei, Maxim. in Bull. Soc. Nat. Mosc. liv. (1879) i. 61.—Japon.
lepida, Reichb. f. Otia Bot. Hamb. 51.—Ins. Loochoo.
longifolia, Lindl. in Wall. Cat. n. 7060 = H. longifolia.
longiscа, Durand, in Journ. Acad. Sc. Philad. Ser. II. ii. (1855) 101 = Herminium congestum.
Lyallii, Steud. Nom. ed. II. i. 712 = Cynorchis flexuosa.
macrantha, Lindl. Gen. et Sp. Orch. 279.—Afr. trop.
nigra, Reichb. f. in Bonplandia, iv. (1856) 321 = H. nigra.
nivea, Ingelm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1845) 236 = H. nivea.
nivea, Lindl. Gen. et Sp. Orch. 280 (= *Platanthera Orchidis*).—Ind. or.
obcordata, Reichb. f. Otia Bot. Hamb. 32 = H. Galeandra.
odoratissima, A. Rich. in Mém. Mus. Par. iv. (1818) 57 = H. odoratissima.
Orchidis, Lindl. Gen. et Sp. Orch. 278 (= *Platanthera Orchidis*).—Reg. Himal.
ornithis, Spreng. Syst. iii. 693 = seq.
ornithis ß, A. Rich. in Mém. Mus. Par. iv. (1818) 57 = H. conopsea, *Benth.*
pauciflora, Lindl. Gen. et Sp. Orch. 280.—Dahuria.
Pinguicula, Reichb. f. & S. Moore, in Journ. Bot. xvi. (1878) 185 = H. Pinguicula.
plantaginea, Lindl. in Wall. Cat. n. 7058 = H. plantaginea.
Pseudo-conopsea, Gren. ex Nym. Consp. 895 = H. conopsea, *Benth.*
puberula, Lindl. in Wall. Cat. n. 7059 = Orchis Chusua.
purpurascens, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 27 = Cynorchis purpurascens.
purpurea, Phil. ex Nym. Consp. 895 = H. conopsea, *Benth.*
rosellata, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 25 = H. nasuta.
rupestris, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 206.—Japon.
Schweinfurthii ß, Kern. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 213.—Europ.
secunda, Lindl. in Wall. Cat. n. 7054 (= *Peristylus laricifolius*).—Ind. or.
sibirica, Turcz. ex Lindl. Gen. et Sp. Orch. 277.—Sibir. bales.
spathulata, Lindl. l. c. 280 = H. spathulata.
squamata, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 23 = Cynorchis squamosa.

GYMNADENIA :—
Strumpfii ×, Aschers. in Oestr. Bot. Zeitschr. xv. (1865) 179.—Europ.
suaveolens, Reichb. f. in Bonplandia, iv. (1856) 321 = Habenaria odoratissima.
tenuiflora, Lindl. in Wall. Cat. n. 7055 = H. lacertifera.
tenuis, Lindl. l. c. n. 7057 = H. stenostachya.
transilvanica, Schur, Enum. Pl. Transs. 644 = H. conopsea, *Benth.*
tridentata, Lindl. Gen. et Sp. Orch. 277 = H. tridentata.
trispicula, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 24.—Ins. Maurit.
trophiaeformis, Reichb. f. in Journ. Bot. xiv. (1876) 209.—China.
uniflora, Steud. Nom. ed. II. i. 712 = Cynorchis uniflora.
Vidalii, Franch. & Sav. Enum. Pl. Jap. ii. 512.—Japon.
violacea, Lindl. Gen. et Sp. Orch. 278 (= *Platanthera Orchidis*).—Ind. or.
viridis, A. Rich. in Mém. Mus. Par. iv. (1818) 57 = H. viridis.
Wahlenbergii, Afzel. ex Reichb. Ic. Fl. Germ. xiii. 114. t. 423 = H. conopsea, *Benth.*

GYMNAGATHIS, Schau. in Linnaea, xvii. (1843) 242 = **Malaleuca**, Linn. (Myrtac.).
teretifolia, Schau. l. c. 248 = M. teretifolia.

GYMNALYPHA, Griseb. in Bonplandia, vi. (1858) 2 = **Acalypha**, Linn. (Euphorb.).
Jacquinii, Griseb. l. c. = A. villosa.

GYMNAMBLOSIS, Pfeiff. Nom. i. 1520 (1874), sphalm. = GYNAMBLOSIS, Torr. = **Croton**, Linn. (Euphorb.).

GYMNANDRA, Pall. Reise, iii. 710 (1776) = **Lagotis**, Gaertn. (Selagin.).
altaica, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 393. t. 4. f. 2 = L. glauca.
armena, Boiss. Diagn. Ser. I. iv. 75 = L. stolonifera.
borealis, Pall. Reise, iii. 710 = L. glauca.
cashmeriana, Royle, ex Benth. Scroph. Ind. 47 = L. glauca.
dentata, Willd. in Ges. Naturf Fr. Berl. Mag. v. (1811) 394. t. 9. f. 4 = L. glauca.
elongata, Willd. l. c. 395. t. 10. f. 7 = L. glauca.
globosa, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 89. t. 7. f. 1 = L. globosa.
Gmelini, Cham. & Schlecht. in Linnaea, ii. (1827) 561 = L. glauca.
gracilis, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 394. t. 9. f. 5 = L. glauca.
integrifolia, Willd. l. c. 392. t. 9. f. 1 = L. glauca.
Korolkowi, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 627 = L. Korolkowi.
leucantha, Royle, ex Benth. Scroph. Ind. 47 = L. glauca.
longiflora, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1843) 417 = L. glauca.
Merchii, Cham. ex Choisy, in DC. Prod. xii. 25 = L. glauca.
minor, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 393. t. 9. f. 3 = L. glauca.
ovata, Willd. l. c. 395. t. 10. f. 8 = L. glauca.
Pallasii, Cham. & Schlecht. in Linnaea, ii. (1827) 564 = L. glauca.
reniformis, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 396. t. 10. f. 9 = L. glauca.
rubra, Dougl. in Hook. Fl. Bor. Am. ii. 108 = Synthyris rubra.
spectabilis, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 80 = L. spectabilis.
Stelleri, Cham. & Schlecht. in Linnaea, ii. (1827) 563 = L. glauca.
Stelleri, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 559 = L. glauca.
stolonifera, C. Koch, in Linnaea, xvii. (1843) 259 = L. stolonifera.

GYMNANDROPOGON, [Munro] Duthie, in Atkins. Ges. N. W. Ind. 686 (1878) = **Andropogon**, Linn. (Gramin.).
convolutus, [Munro] Duthie, l. c. = A. annulatus.
hirtus, [Munro] Duthie, l. c.—Ind. or.

GYMNANTHELIA, Anderss. in Schweinf. Beitr. Fl. Aethiop. 299 (1867), nomen = **Andropogon**, Linn. (Gramin.).
communis, Schweinf. & Aschers. l. c. = A. communatus.
connata, Schweinf. & Aschers. l. c. = A. Schoenanthus.
lanigera, Anderss. l. c. = A. laniger.
Martini, Anderss. l. c. = A. Schoenanthus.
nervata, Schweinf. & Aschers. l. c. = A. Schoenanthus.
proxima, Anderss. l. c. = A. Iwarancusa.
sennaarensis, Schweinf. & Aschers. l. c. = A. Iwarancusa.

GYMNANTHEMUM, Cass. in Bull. Soc. Philom. (1817) 10 = **Vernonia**, Schreb. (Compos.).
abyssinicum, Sch. Bip. ex Walp. Rep. ii. 948 = V. amygdalina.
acuminatum, Steetz, in Peters, Reise Mossamb. Bot 336 = V. arborea ?
affine, Steetz, l. c. 338 = V. scandens ?
amygdalinum, Sch. Bip. ex Walp. Rep. ii. 948 = V. amygdalina.
angustifolium, Benth. in Hook. Niger Fl. 429 = V. Benthamiana.
blandum, Steetz, in Peters, Reise Mossamb. Bot. 339 = V. blanda.
Bollenium, Steetz, l. c. 336.—Afr. trop.
congestum, Cass. in Dict. Sc. Nat. xx. 110 = V. trifloriculosa.
cupuiere, Cass. l. c. 109 = V. senegalensis.
dendigulense, Walp. Rep. ii. 948 = V. indica.
divergens, Sch. Bip. ex Walp. l. c. = V. divergens.
extensum, Steetz, in Peters, Reise Mossamb. Bot. 337 = V. extensa.
fimbrilliferum, Cass. in Dict. Sc. Nat. xx. 109 (= *Decaneurum fimbrilliferum*).—Ins. Borbon.
frutescens, Sch. Bip. ex Walp. Rep. ii. 946 (Centratherae sp. ?).—Java.
glabrum, Sch. Bip. ex Walp. l. c. = Eupatorium glabrum.
grande, Sch. Bip. ex Walp. l. c. (= *Decaneurum grande*).—Madag.
Metalonum, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 334 = V. divergens.
molle, Sch. Bip. ex Hook. f. l. c. 228 = Centratherum Ritchiei.
molle, Sch. Bip. ex Walp. Rep. ii. 948 = Centratherum molle.
obovatum, Gandich. Voy. Uran. Bot. 471 = V. scandens.
phyllolaenum, Sch. Bip. ex Walp. Rep. ii. 949 (phyllolaenum) = Centratherum phyllolaenum.
quercifolium, Steetz, in Peters, Reise Mossamb. Bot. 334 = V. senegalensis.
reticulatum, Sch. Bip. ex Walp. Rep. ii. 948 = Centratherum reticulatum.
scandens, Steetz, in Peters, Reise Mossamb. Bot. 338 = V. scandens.
scariosum, Sch. Bip. ex Walp. Rep. ii. 949 = V. scariosa. *Arn.*
senegalense, Sch. Bip. ex Walp. l. c. 948 = V. senegalensis.
silhetense, Sch. Bip. ex Walp. l. c. = V. bracteata.
vagans, Steetz, in Peters, Reise Mossamb. Bot. 338 = V. scandens.

GYMNANTHERA, R. Br. in Mem. Wern. Soc. i. (1809) 58. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 742.
CYTIXYLON, Llanos, Fragm. 59 (1851).
DICHOLEPIS, Blume, Mus. Bot. Lugd. Bat. i. 145, t. 24 (1850).
nitida, *R. Br. Prod.* 464.—Austral.
pedunculata, *Viller, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 133.—Malaya; Ins. Philipp.

GYMNANTHES, Sw. Prod. Veg. Ind. Occ. 95 (1788). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 337.
actinostemoides, *Muell. Arg. in Linnaea*, xxxii. (1863) 108.—Mexic.
angustifolia, Muell. Arg. l. c. 99 = Sebastiania angustifolia.
bahiensis, Muell. Arg. l. c. 103 = Sebastiania bahiensis.
brachyclada, Muell. Arg. l. c. 98 = Sebastiania brachyclada.
brasiliensis, Muell. Arg. l. c. 104 = Sebastiania brasiliensis.
brevifolia, Muell. Arg. l. c. = Sebastiania brevifolia.
concolor, Muell. Arg. l. c. 103 = Actinostemon carlhaena.

GYMNANTHES:—
discolor, Baill. Etud. Gén. Euph. 531 = Sebastiania marginata.
discolor, Muell. Arg. in Linnaea, xxxii. (1863) 103 = Sebastiania discolor.
elliptica, *Sw. Prod. Veg. Ind. Occ.* 96.—Ind. occ.
Gaudichaudii, Muell. Arg. in Linnaea, xxxii. (1863) 96 = Sebastiania Gaudichaudii.
glandulosa, *Muell. Arg. l. c.* 106.—Bras.
granatensis, Muell. Arg. l. c. 107 = Sebastiania granatensis.
guyanensis, Muell. Arg. l. c. 102 = Sebastiania guyanensis.
hybolenca, *Benth. in Hook. Kew Journ.* vi. (1854) 325.—Bras.
jacobinensis, Muell. Arg. in Linnaea, xxxii. (1863) 106 = Sebastiania jacobinensis.
Klotzschiana, Muell. Arg. l. c. 98 = marginata.
ligustrina, Muell. Arg. l. c. 96 = Sebastiania ligustrina.
longipes, *Muell. Arg. l. c.* xxxiv. (1865–66) 216 — Mexic.
lucida, *Sw. Prod. Veg. Ind. Occ.* 96.—Ind. occ.
marginata, *Baill. Etud. Gén. Euph.* 531.—Bras.
multiramea, Muell. Arg. in Linnaea, xxxii. (1863) 97 = Sebastiania multiramea.
nervosa, Muell. Arg. l. c. 102 = Sebastiania nervosa.
obtusa, Baill. Etud. Gén. Euph. 530 = elliptica.
pachystachys, Baill. l. c. 531 = Sebastiania pachystachys.
pallida, *Muell. Arg. in Linnaea*, xxxii. (1863) 106.—Cuba.
Pavoniana, Muell. Arg. l. c. = Sebastiania Pavoniana.
polyandra, *Benth. & Hook. f. Gen.* iii. 337.—Ind. occ.
pterocladon, Muell. Arg. in Linnaea, xxxii. (1863) 107 = Sebastiania pteroclada.
rigida, Muell. Arg. l. c. 99 = Sebastiana rigida.
riparia, *Klotzsch, in Wiegm. Archiv*, vii. (1841) 182.—Mexic.
Schlechtendaliana, Muell. Arg. in Linnaea, xxxii. (1863) 100 = riparia.
Schottiana, Muell. Arg. l. c. 96 = Sebastiania Schottiana.
serrata, Baill. Etud. Gén. Euph. 531 = Sebastiania serrata.
stipulacea, Muell. Arg. in Linnaea, xxxii. (1863) 96 = Sebastiania stipulacea.
Treculiana, Muell. Arg. l. c. xxxiv. (1865–66) 216 = Sebastiania Treculiana.
trinervia, Muell. Arg. l. c. xxxii. (1863) 101 = Sebastiania trinervia.
Widgreni, Muell. Arg. l. c. 97 = Sebastiania Widgreni.
ypanemensis, Muell. Arg. l. c. 100 = Sebastiania ypanemensis.

GYMNANTHUS, Endl. Gen. Suppl. v. 87 (1850) = praec.

GYMNANTHUS, Jungh. in Hoer. & De Vriese, Tijdschr. vii. (1840) 808 = **Trochodendron**, Sieb. & Zucc. (Magnol.).
paradoxus, Jungh. l. c.—Japon.

GYMNARRHEA, Steud. Nom. i. 385 (1821), sphalm. = seq.

GYMNARRHENA, Desf. in Mém. Mus. Par. iv. (1818) 1. t. 1. *COMPOSITAE*, Benth. & Hook. f. ii. 397.
CRYPTADIA, Lindl. in Endl. Gen. Pl. Suppl. i. 1381 (1841).
FRANKIA, Steud. Nom. ed. II. i. 646 (1840).
Balsamea, Coss. & Dur. ex Coss. & Kralik, in Bull. Soc. Bot. Fr. iv. (1857) 160 = microcephala.
microcephala, *Desf. in Mém. Mus. Par.* iv. (1818) 2. t. 1.—Afr. bor.; Arab.; Persia.

GYMNELAEA, Spach, Hist. Vég. Phan. viii. 256 (1839) = Olea, Tourn.

GYMNEMA, R. Br. in Mem. Wern. Soc. i. (1809) 33. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 769.
acuminatum, *Wall. Tent. Fl. Nep.* 50.—Ind. or.; Malaya.
affine, *Decne. in DC. Prod.* viii. 622.—Ind. or.; Chile.
albidum, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 276.—Ins. Timor.

GYMNEMA:—
attenuatum, *Wall. Tent. Fl. Nep.* 50.—Reg. Himal.
aurantiacum, Wall. ex Hook. f. Fl. Brit. Ind. iv. 63 = Leptadenia reticulata.
brevifolium, *Benth. Fl. Austral.* iv. 343.—Austral.
columnare, Wall. ex Wight, Contrib. 46; Cat. n. 8193 = Gongronema nepalense.
crenatum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 278.—Afr. trop.
Decaisneanum, Wight, Ic. t. 1271 = hirsutum.
elegans, *Wight & Arn. in Wight, Contrib.* 46.—Ind. or.
erianthum, *Decne. in DC. Prod.* viii. 632.—N. Caled.
Finlaysonii, Wight, Contrib. 46 = Gongronema Finlaysonii
geminatum, R. Br. Prod. 462 = sylvestre.
glabrum, *Wight, Contrib.* 45.—Burma.
glaucum, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 323.—Singapore.
hirsutum, Wall. Tent. Fl. Nep. 50 = Tylophora hirsuta.
hirsutum, *Wight & Arn. in Wight, Contrib.* 44.—Ind. or.
humile, *Decne. in DC Prod.* viii. 621.—Abyssin.
inodorum, *Decne. i. c.* 551, nomen.—China.
lactiferum, *R. Br. in Mem Wern. Soc.* i. (1809) 33.—Ind. or ; Malaya.
latifolium, *Wall. ex Wight, Contrib.* 45.—Ind. or.; Burma.
litorale, *Blume, Mus. Bot. Lugd. Bat.* i. 149.—Java.
M'Kenii, Harv. Gen. S. Afr. Pl. ed. II. 259 = subvolubile.
macranthum, *Hook. f. in Hook. Ic. Pl.* t. 1486.—Reg. Himal.
macrocarpum, *A. Rich. Tent. Fl. Abyss.* ii. 43.—Abyss.
Maingayi, *Hook. f. Fl. Brit. Ind.* iv. 81.—Malacca.
malayanum, Grif. Notul. iv. 56 = lactiferum.
melicida, Edgew. in Journ. As. Soc. Beng. xxi. (1852) 174 = sylvestre.
micradenia, *Benth. Fl. Austral.* iv. 344.—Austral.
molle, *Wall. ex Wight, Contrib.* 45.—Burma.
montanum, *Hook. f. Fl. Brit. Ind.* iv. 81.—Ind. or.
Muelleri, *Benth. Fl. Austral.* iv. 343.—Austral.
nepalense, Wall. Tent. Fl. Nep. 49 = Gongronema nepalense.
nepaulense, J. Grah. Cat. Pl. Bomb. 120 = Anodendron paniculatum.
nitens, Blume, Mus. Bot. Lugd. Bat. i. 149 = lactiferum.
nitidum, *Benth. in Hook. Niger Fl.* 456.—Afr. trop.
parviflorum, Decne. in DC. Prod. viii. 621, in syn. sphalm. = seq.
parvifolium, Wall. Tent. Fl. Nep. 50 = affine.
pergularioides, *Wight & Gardn. ex Thw. Enum. Pl. Zeyl.* 128, in syn.—Zeylan.
pleiadenium, *F. Muell. Fragm.* xi. 77.—Austral.
pubigerum, Hort. Calc. ex Decne. in DC. Prod. viii. 609 = Tylophora exilis.
recurvifolium, *Blume, Mus. Bot. Lugd. Bat.* i. 150.—N. Guinea.
rotundatum, *Thw. Enum. Pl. Zeyl.* 197.—Zeylan.
rufescens, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 277.—Madag.
sagittatum, Wall. Cat. n. 8194 ; Tent. Fl. Nep. 50 = Gongronema nepalense.
sagittatum, Wall. l. c. 8194 B = Tylophora Wallichii.
Spartium, Wall. Cat. sub n. 8199 = Leptadenia Spartium.
stenolobs, *Hook. f. Fl. Brit. Ind.* iv. 82.—Zeylan.
stenophyllum, *A. Gray, in Proc. Am. Acad.* v. (1862) 326.—Austral.; Ins. Fiji.
subandum, *A. Gray, l. c.*—Ins. Fiji.
subvolubile, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 277.—Afr. trop.
sylvestre, *R. Br. in Mem. Wern. Soc.* i. (1809) 33.—Ind. or. ; Afr. trop. ; Austral.
tenacissimum, Spreng. Syst. i. 844 = Marsdenia tenacissima.
Thomsoni, *Hook. f. Fl. Brit. Ind.* iv. 82.—Reg. Himal.
tingens, *Spreng. Syst.* i. 844.—Ind. or.
trinerve, *R. Br. Prod.* 462.—Austral.
Wallichii, Wight, Contrib. 46 = Gongronema Wallichii et Tylophora Wallichii.
zeylanicum, Decne. in DC. Prod. viii. 622 = lactiferum.

GYMNEMA, Endl. Gen. 891, in syn. (1837) = GYNEMA, Rafin. = **Pluchea**, Cass. (Compos.).

GYMNOBALANUS, Nees & Mart. in Linnæa, viii. (1833) 86 = **Cocotes**, Aubl. (Laurin.).
Castelnæanus, Nees, Syst. Laurin. 488 = Nectandra Willdenowiana.
Femdleri, *Meisn. in DC. Prod.* xv. i. 142.—Venezuela.
guianensis, Nees, Syst. Laurin. 483 = O. guianensis?
latifolia, Meisn. in DC. Prod. xv. i. 141 = Nectandra polita.
minarum, Nees & Mart. in Linnæa, viii. (1833) 88 = Nectandra pauamophila.
pernoides, Meisn. in DC. Prod. xv. i. 141 = O. puberula.
punctata, Meisn. l. c. = Nectandra punctata.
Regnellii, Meisn. in Kjoeb. Vidensk. Meddel. (1870) 188 = Nectandra grandiflora.
reticulata, Meisn. l. c.—Bras.
Spruceri, Meisn. in DC. Prod. xv. i. 140.—Bras.

GYMNOBOTHRYS, Wall. ex Baill. Etud. Gén. Euph. 596 (1858) = **Sapium**, P. Br. (Euphorb.).
lucida, Wall. ex Baill. l. c. 597 = S. insigne.

GYMNOCALYCIUM, Pfeiff. Cat. Hort. Schel. ex Pfeiff. & Otto, Abbild. Cacteen, ii. t. 12 (1844) = **Echinocostus**, Link & Otto (Cact.).
denudatum, Pfeiff. ex Foerst. Handb. Cact. ed. I. 289 = E. denudatus.
gibbosum, Pfeiff. l. c. 291 = E. gibbosus.
reductum, Pfeiff. & Otto, Abbild. Cacteen, ii. t. 12 = E. gibbosus, Cereus reductus.
villosum, Pfeiff. ex Foerst. Handb. Cact. ed. I. 297 = E. polyrhaphis.

GYMNOCAMPUS, Lesch. ex Pfeiff. Nom. i. 1522 (1874) = GYNOCAMPUS, Lesch. = **Levenhookia**, R. Br. (Stylid.).

GYMNOCARPON, Pers. Syn. i. 262 (1805) = seq.

GYMNOCARPOS, Forsk. Fl. Aegypt. Arab. 65 (1775). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 17.
decandrum, *Forsk. l. c.*—Afr. trop.
fruticosum, Pers. Syn. i. 262 = decandrum.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 503.—Mongolia.

GYMNOCARPUM, DC. Prod. iii. 359 (1828) = praec.

GYMNOCARPUS, Thou. ex Baill. Etud. Gén. Euph. 595 (1858) = **Uapaca**, Baill. (Euphorb.).

GYMNOCAULIS, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 180 = **Aphyllon**, Mitch. (Orobanch.).
bifora, Nutt. l. c. = A. uniflorum.
fasciculata, Nutt. l. c. = A. fasciculatum.
uniflora, Nutt. l. c. = A. uniflorum.

GYMNOCAULUS, Phil. in Linnæa, xxviii. (1856) 705 = **Calyoera**, Cav.
viridiflorus, Phil. l. c. 706 = C. viridiflora.

GYMNOCHAETA, Steud. Syn. Pl. Cyp. 156 (1855) = **Bolboxus**, Linn. (Cyperac.).
Drummondii, Steud. l. c. = S. Drummondii.

GYMNOCHAETE, Benth. & Hook. f. Gen. iii. 1063 (1883), sphalm. = praec.

GYMNOCHILUS, Blume, Orch. Archip. Ind. 107. t. 29 (1858). *ORCHIDEAE*, Benth. & Hook. f. iii. 606.
nudum, *Blume, l. c.* 108.—Ins. Mascar.
recurvum, *Blume, l. c.* 109. t. 32. f. 2.—Madag.

GYMNOCLADUS, Lam. Encyc. i. 733 (1783). *LEGUMINOSAE*, Benth. & Hook. f. i. 568.
arabica, Lam. l. c. = Moringa aptera.
canadensis, *Lam. l. c.*—Am. bor.
chinensis, *Baill. in Bull. Soc. Linn. Par.* i. (1878) 34.—China.
dioica, C. Koch, Dendrol. I. 5 = canadensis.
Williamsii, Hance, in Journ. Bot. xxii. (1884) 366 = Gleditschia sinensis.

GYMNOCLINE, Cass. in Bull. Soc. Philom. (1816) 199; et in Dict. Sc. Nat. xx. 119 (1821) = **Chrysanthemum**, Linn. (Compos.).
achilleaefolia, C. Koch, in Linnæa, xxiv. (1851) 840 = C. achilleaefolium.
argyrophylla, C. Koch, l. c. (= *Pyrethrum myriophyllum*).—Reg. Caucas.

GYMNOCLINE:—
armenum, C. Koch, in Linnæa, xxiv. (1851) 840 = Chrysanthemum millefoliatum?
cedretorum, Schott & Kotschy, ex Tchihat. Asie Min. Bot. ii. 278.—As. Min.
chilophylla, C. Koch, in Linnæa, xxiv. (1851) 839 = C. millefoliatum.
leucocephala, Cass. in Dict. Sc. Nat. xx. 120 = C. macrophyllum.
macrophylla, Bieff & Fingerh. Fl. Germ. ii. 389 = C. macrophyllum.
myriophylla, C. Koch, in Linnæa, xxiv. (1851) 839 (= *Pyrethrum myriophyllum*).—Reg. Caucas.
Sawitzii, C. Koch, l. c. 840 (= *Pyrethrum polyphyllum*).—Armenia.
Vaillantii, Cass. in Dict. Sc. Nat. xx. 121 = C. achilleaefolium.
xanthocephala, Cass. l. c. = C. achilleaefolium.

GYMNOCOCCA, Fisch. & Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. x. 46 (1846) = **Pimelea**, Banks & Soland. (Thymel.).
arenaria, Fisch. & Mey. l. c. 47 = P. arenaria.
drupacea, Fisch. & Mey. l. c. 46 = P. drupacea.
microcarpa, Fisch. & Mey. l. c. 47 = P. Urvilleana.

GYMNOCORONIS, DC. Prod. v. 106 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 289.
altenuata, DC. l. c. = spilanthoides.
latifolia, *Hook. & Arn. Bot. Beech. Voy.* 296.—Mexic.
spilanthoides, *DC. Prod.* vii. 266.—Reg. Argent.
subcordata, DC. l. c. v. 106 = spilanthoides.

GYMNODES, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 172 = **Lunula**, DC. (Juncac.).
campestris, Fourr. l. c. = L. campestris.
multiflora, Fourr. l. c. 178 = L. erecta.
spicata, Fourr. l. c. = L. spicata.
sudetica, Fourr. l. c. = L. nigricans.

GYMNODISOUS, Less. in Linnæa, xiv. (1831) 95; et Syn. Comp. 89 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 452.
capillaris, *Less. in Linnæa*, xiv. (1831) 95.—Afr. austr.
linearifolius, *DC. Prod.* vi. 469.—Afr. austr.
orbiculifolius, *Sch. Bip. in Flora*, xxvii. (1844) 768. —Afr. austr.

GYMNOGYNE, F. Didr. Ind. Sem. Hort. Hamb. (1850). —Cf. Linnæa, xxix. (1857-58) 787 = Boehmeria, Jacq. (Urtic.).
elongata, F. Didr. l. c. = B. elongata.

GYMNOGYNE, Steetz, in Lehm. Pl. Preiss. i. 481 (1845) = Cotula, Linn. (Compos.).
cotuloides, Steetz, l. c. 482 = C. Gymnogyne.

GYMNOLAEMA, Benth. in Benth. & Hook. f. Gen. ii. 740 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. i. c.
Newii, *Benth. in Hook. Ic. Pl.* t. 1186.—Afr. trop.

GYMNOLEIMA, Decne. in Jacquem. Voy. Bot. 122 (1844) = Lithospermum, Tourn. (Borag.).
fruticiorum, Decne. l. c. = L. fruticosum.
graminifolium, Decne. l. c. = L. graminifolium.
oleifolium, Decne. l. c. = L. oleaefolium.
rosmarinifolium, Decne. l. c. = L. rosmarinifolium.

GYMNOLOMA, Ker-Gawl. in Bot. Reg. t. 669 (1822) = seq.

GYMNOLOMIA, H. B. & K. Nov. Gen et Sp. iv. 217, t. 378, 374 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 268.
GYMNOPSIS, DC. Prod. v. 561 (1836), partim.
HELIOMERIS, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 171.
MIRASOLIA, Sch. Bip. ex Benth. & Hook. f. ii. 267 (1873).
calva, *A. Gray, in Proc. Am. Acad.* xix. (1883) 6, in obs.—Mexic.
cheracoensis, *Benth. & Hook. f. Gen.* ii. 269.—Mexic.
connata, *Sprengl. Syst.* iii. 610.—Bras.
diversifolia, Pohl, ex Baker, in Mart. Fl. Bras. vi. iii. 268 = Calea diversifolia.

GYMNOLOMIA:—
encelioides, *A. Gray, in Proc. Am. Acad.* xix. (1883) 4.—Calif.
enallofila, *Benth. & Hook. f. Gen.* ii. 269.—Mexic.
flava, *Hemsl. Biol. Centr. Am. Bot.* ii. 161.—Mexic.
Ghiesbreghtii, *Hemsl. l. c.* 162.—Mexic.
Greggii, *A. Gray, in Proc. Am. Acad.* xv. (1880) 36.—Mexic.
hondensis, *H. B. & K. Nov. Gen. et Sp.* iv. 218.—N. Granat.
Kunthiana, *Baker, in Mart. Fl. Bras.* vi. iii. 172.—Bras.
latebractosta, *Hemsl. Biol. Centr. Am. Bot.* ii. 162.—Mexic.
maculata, Ker-Gawl. in Bot. Reg. t. 669 = Wulfia stenoglossa.
microcephala, *Less. in Linnæa*, v. (1830) 158.—Mexic.
multiflora, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 162.—Mexic.
ovata, *A. Gray, in Proc. Am. Acad.* xix. (1883) 4.—Mexic.
Parkinsonii, *Hemsl. Biol. Centr. Am. Bot.* ii. 168.—Mexic.
patens, *A. Gray, in Proc. Am. Acad.* v. (1862) 182.—Mexic.
pinnatiloba, *Benth. & Hook. f. Gen.* ii. 269.—Mexic.
platylepis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 5. —Mexic.
Porteri, *A. Gray, l. c.* xii. (1877) 59.—Am. bor.
quinata, *Benth. & Hook. f. Gen.* ii. 264.—Ecuador.
rudbeckioides, *H. B. & K. Nov. Gen. et Sp.* iv. 219. t. 374.—Am. centr.
squarrosa, *Benth. & Hook. f. Gen.* ii. 269.—Mexic.
subdecurens, *Benth. & Hook. f. Gen.* i. 264.—Mexic.
tenella, *H. B. & K. Nov. Gen. et Sp.* iv. 218.—N. Granat.
tenuifolia, *Benth. & Hook. f. Gen.* ii. 264.—Mexic.
triloba, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 217.—Arizona.
triplinervia, *H. B. & K. Nov. Gen. et Sp.* iv. 219.—N. Granat.
villosa, Pohl, ex Baker, in Mart. Fl. Bras. vi. iii. 262 = Calea villosa.

GYMNOMESIUM, Schott, in Oestr. Bot. Wochenbl. v. (1855) 17 = **Arum**, Linn. (Aroid.).
pictum, Schott, l. c. = A. pictum.

GYMNONYCHIUM, Bartl. in Linnæa, xvii. (1843) 353. t. 11 = **Agathosma**, Willd. (Rutac.).
pubescens, Bartl. l. c. 354. t. 11 = A. pubescens.

GYMNOPENTELA, Benth. in Benth. & Hook. f. Gen. ii. pars 1ma Add. 587; et 1285 (1873). *COMPOSITAE*, Benth. & Hook. f. i. c.
bifurcata, *Benth. l. c.*; *et in Hook. Ic. Pl.* t. 1155.—Afr. austr.

GYMNOPETALUM, Arn. in Hook. Journ. Bot. iii. (1841) 278. *CUCURBITACEAE*, Benth. & Hook. f. i. 822.
SCOTANTHUS, Nasd. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 172.
TRIPODANTHERA, M. Roem. Syn. Pepon. 48 (1846).
calycularium, Miq. Fl. Ind. Bat. Suppl. 582 = Curcurbita moschata.
ceylonicum, *Arn. in Hook. Journ. Bot.* iii. (1841) 278. —Zeylan.
cochinchinense, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 57.—As. trop.
heterophyllum, Kurz, in Journ. Bot. xiii. (1875) 326 = quinquelobum!
Horsfieldii, Miq. Fl. Ind. Bat. i. 1. 680 = Thladiantha cordifolia.
integrifolium, Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 58 = Trichosanthes integrifolia.
japonicum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 82 = Trichosanthes japonica.
lanceolatum, *Miq. Fl. Ind. Bat.* i. 1. 680.—Java.
piperifolium, Miq. l. c. = Thladiantha cordifolia.
quinquelobatum, Cogn. in A.DC. Monog. Phan. iii. 998, sphalm. = seq.
quinquelobum, *Miq. Fl. Ind. Bat.* i. 1. 681.—Java.
scplemlobum, Miq. l. c. 679 = Beninccsa cerifera.
tuberosum, Cogn. in DC Monog. Phan. iii. 998 = cepiendum.
Wehiei, *Cogn. l. c.* 994.—Cochinch.

GYMNOPETALUM :—
Wight&iii, *Arn. in Hook. Journ. Bot.* iii. (1841) 278.—Ind. or.
Wight&i, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 611, partim = ceylonicum.
ceylanicum, Walp. Rep. ii. 198, in syn. = ceylonicum.

GYMNOPHYTON, Clos, in C. Gay, Fl. Chil. iii. 102. t. 32 (1847) = **Asteriscolum**, Cham. & Schlecht. (Umbellif.).
flexuosum, Clos, l. c. 104 = A. flexuosum.
foliosum, Phil. Fl. Atac. 24.—Chili.
Kingi, Phil. in Anal. Univ. Chil. (1872) 727.—Chili.
polycephalum, Clos, in C. Gay, Fl. Chil. iii. 103. t. 32. f. 1 = A. polycephalum.
robustum, Clos, l. c. 104 = A. robustum.
spinescens, Phil. in Linnaea, xxxiii. (1864–65) 91.—Chili.
spinosissimum, Phil. Fl. Atac. 24.—Chili.

GYMNOPOGON, Beauv. Agrost. 41. t 9. f. 3 (1812).
GRAMINEAE, Benth. & Hook. f. iii. 1167.
ANTHOPOGON, Nutt. Gen. Am. i. 81 (1818).
BIATHERIUM, Desv. Opusc. 72 (1831).
DICHAETARIA, Nees, in Steud. Syn. Pl. Gram. 145 (1855).
brevifolius, *Trin. Gram. Unifl.* 238.—Am. bor. et austr.
digitatus, Nees, ex Steud. Nom. ed. II. i. 713 = Cynodon elongatus.
distichophyllus, *Steud. Syn. Pl. Gram.* 218.—Texas.
elongatus, Nees, ex Steud. Nom. ed. II. i. 713 = Cynodon elongatus.
fastigiatus, Nees, Agrost. Bras. 430 = brevifolius.
filiformis, *Griseb. Fl. Brit. W. Ind.* 588.—Ind. occ.
follosus, *Nees, Agrost. Bras.* 426.—Am. trop.
laevis, Nees, l. c. 428 = Muehlenbergia spicata.
longifolius, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 560.—Mexic.
mollis, *Nees, Agrost. Bras.* 427.—Bras.
pallulans, *Doell, in Mart. Fl. Bras.* ii. 111. 82.—Bras.
racemosus, *Beauv. Agrost.* 41. t. 9. f. 3.—Am. bor. et austr.
rigidus, *Doell, in Mart. Fl. Bras.* ii. 111. 80.—Bras.
rigidus, *Thw. Enum. Pl. Zeyl.* 372.—Zeylan.
scoparius, Trin. Gram. Unifl. 237 = racemosus.
Virletti, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 560.—Mexic.

GYMNOPSIS, DC. Prod. v. 561 (1836) = **Gymnolomia**, H. B. & K. (Compos.).
Clausseni, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 111. 188 = Wedelia paberula.
connata, Steud. Nom. ed. II. i. 718 = Gymnolomia connata.
costa-ricensis, Benth. ex Oerst. in Kjoeb. Videnak. Meddel. (1859) 90 = Gymnolomia radbeckioides.
dentata, DC. Prod. v. 561 = Sclerocarpus dentatus.
divaricata, Benth. Bot. Voy. Sulph. 116 = Sclerocarpus divaricatus.
eunoviaides, DC. Prod. v. 562.—Bras.
fruticosa, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 291 = Aspilia fruticosa.
glaberrima, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 111. 246 = Bidens Riedelii.
helianthoides, DC. Prod. v. 562.—Reg. Argent.
hondensis, DC. l. c. = Gymnolomia hondensis.
Kunthiana, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 292 = Gymnolomia Kunthiana.
Mandonii, Sch. Bip. in Linnaea, xxxiv. (1865–66) 528, nomen.—Bolivia.
microcephala, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 292 = Eleutheranthera ovata.
microcephala, Steud. Nom. ed. II. i. 718 = Gymnolomia microcephala.
microthamna, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 111. 204 = Aspilia fruticosa.
Pohlii, Sch. Bip. ex Baker, l. c. 201 = Aspilia Pohlii.
radbeckioides, DC. Prod. v. 562 = Gymnolomia radbeckioides.
Schiedeana, DC. l. c. 561 = Sclerocarpus Schiedeanus.
tenella, DC. l. c. 562 = Gymnolomia tenella.
triplinervia, DC. l. c. = Gymnolomia triplinervia.
uniserialis, Hook. in. Fl. t. 145 = Sclerocarpus uniseriatus.
verbesinoides, DC. Prod. v. 561.—Ins. Trinit.

GYMNOPSIS :—
vulcanica, Steetz, in Seem. Bot. Voy. Herald, 157 = Gymnolomia radbeckioides.
Weddelliii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. 111. 184 = Wedelia macrodonta.
wedelioides, DC. Prod. v. 562.—Ind. occ.

GYMNOPYRENIUM, Dulac, Fl. Hautes-Pyr. 816 (1867) = **Cotoneaster**, Medic. (Rosac.).
Pyracantha, Dulac, l. c. = Crataegus Pyracantha.
tomentosum, Dulac, l. c. = Cot. tomentosa.
vulgare, Dulac, l. c. = Cot. integerrima.

GYMNOREIMA, Endl. Gen. Suppl. iii. 77 (1843), sphalm. = GYMNOLEIMA, Decne. = **Lithospermum**, Tourn. (Boragin.).

GYMNOSCHOENUS, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 47 = **Mesomelaena**, Nees (Cyperac.).
adustus, Nees, l. c. = M. sphaerocephala.
anceps, Nees, l. c. = M. anceps.
sphaerocephalus, Nees, l. c. = M. sphaerocephala.

GYMNOSCIADIUM, Hochst. in Flora, xxvii. (1844) 20 = **Pimpinella**, Riv. (Umbellif.).
pimpinelloides, Hochst. l. c. = P. Gymnosciadium.

GYMNOSIPHON, Blume, Enum. Pl. Jav. i. 29 (1827). *BURMANNIACEAE*, Benth. & Hook. f. iii. 458.
BENITZIA, Karst. in Linnaea, xxviii. (1856) 420.
CYMBOCARPA, Miers, in Proc. Linn. Soc. i. (1840) 61. t. et in Trans. Linn. Soc. xviii. (1840) 453. t. 28. f. 4.
PTYCHOMBRIA, Benth. in Hook. Kew Journ. vii. (1855) 14.
aphyllum, *Blume, Enum. Pl. Jav.* f. i. 29.—Java.
cornutum, *Benth. & Hook.f. Gen.* iii. 458.—Bras.
cymosum, *Benth. & Hook. f. l. c.*—Bras.
divaricatum, *Benth. & Hook. f. l. c.*—Bras.
papuanum, *Becc. Malesia*, i. (1878) 241.—N. Guin.
refractum, *Benth. & Hook. f. Gen.* iii. 458.—Bras.

GYMNOSPERMA, Less. Syn. Comp. 194 (1832), partim. *COMPOSITAE*, Benth. & Hook. f. ii. 249.
SELLOA, Spreng. Nov. Prov. 86 (1818).
corymbosum, *DC. Prod.* v. 312.—Mexic.
eriocarpa, Walp. Ann. v. 191, sphalm. = Gutierrhiza eriocarpa.
glutinosum, Less. Syn. Comp. 194 = Baccharis glutinosa.
multiflorum, DC. Prod. v. 312 = corymbosum.
radiatum, DC. l. c. = Flaveria linearis.
oppositifolium, DC. l. c. = Flaveria longifolia.
scoparium, DC. l. c. = corymbosum.

GYMNOSPERMIUM, Spach, Hist. Vég. Phan. viii. 66 (1889) = **Leontice**, Linn. (Berberid.).
altaicum, Spach, l. c. 67 = L. altaica.

GYMNOSPORIA, Benth. & Hook. f. Gen. i. 365 (1862). *CELASTRINEAE* [Benth. & Hook. f. i. c. 368, 997].
ENCENTRUS, Presl, Bot. Bemerk. 38 (1844).
POLYACANTHUS, Presl, l. c. (1844).
acuminata, *Hook. f. ex M. Laws. in Fl. Brit. Ind.* i. 610.—Ind. or.
berberidacea, *Baker, in Journ. Linn. Soc.* xx. (1883) 130.—Madag.
brachystachya, *Baker, in Journ. Bot.* xx. (1882) 49.—Madag.
caulinodes, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 176.—Ins. Canar.
crataegina, *Baker, in Journ. Linn. Soc.* xx. (1883) 130.—Madag.
cressa, *Seem. Pl. Vit.* 41, in nota.—Ins. Marques.
divaricata, *Baker, in Journ. Bot.* xx. (1882) 50.—Madag.
diversifolia, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 499.—Ins. Loo-choo.
Dryandri, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 176.—Ins. Canar.
emarginata, *Thw. Enum. Pl. Zeyl.* 409.—Ind. or.
europaea, Masf. in Anal. Soc. Esp. Hist. Nat. x. (1881) 176 = montana.
Falconeri, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 630.—Reg. Himal.
fruticosa, *Thw. Enum. Pl. Zeyl.* 409.—Zeylan.

GYMNOSPORIA :—
Gibsoni, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 302.—Ind. or.
Harlandi, *Hance, in Journ. Bot.* iv. (1866) 171.—Cochinch.
Hayneana, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 630.—Ind. or.
leptopus, *Baker, in Journ. Linn. Soc.* xx. (1883) 121.—Madag.
montana, *Benth. Fl. Austral.* i. 403.—Geront. trop. neglecta, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 619.—Ind. or.
oblanceolata, *M. Laws. l. c.*—Burma.
ovata, *M. Laws. l. c.*—Ind. or.
paniculata, *Baker, in Journ. Linn. Soc.* xx. (1883) 121.—Madag.
philippinensis, *Vidal, Phan. Cuming. Philipp.* 58, 108.—Ins. Philipp.
puberula, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 619.—Ind. or.
Rothiana, *M. Laws. l. c.* 620.—Ind. or.
Koyleana, *M. Laws. l. c.*—Ind. bor. occ.
rufa, *M. Laws. l. c.*—Ind. or.
rugulosa, *M. Laws. l. c.*—Ind. or.
salicifolia, *M. Laws. l. c.* 619.—Assam.
Thomsoni, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 302.—Reg. Himal.
trigyna, *Baker, Fl. Maurit.* 50.—Ins. Maurit.
vitiensis, *Seem. Fl. Vit.* 40.—Ins. Fiji.
Wallichiana, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 621.—Ind. or.

GYMNOSTACHIUM, Reichb. Handb. 190 (1837) = Gymnostachyum, Nees (Acanth.).

GYMNOSTACHYS, R. Br. Prod. 387 (1810). *AROIDEAE*, Benth. & Hook. f. iii. 1000.
anceps, *R. Br. l. c.*—Austral.

GYMNOSTACHYUM, Nees, in Wall. Pl. As. Rar. iii. 76 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1099.
CRYPTOPHRAGMIUM, Nees, in Wall. Pl. As. Rar. iii. 76, 94 (1832).
ODONTOSTIGMA, Zoll. in Nat. en Genesk. Arch. Nederl. Indie, ii. (1846) 578.
PETRACANTHUS, Nees, in DC. Prod. xi. 97 (1847).
SARCANTHERA, Rafin. Fl. Tellur. iv. 84 (1836).
affine, *Nees, in DC. Prod.* xi. 94.—Ins. Philipp.
alatum, Wight, Ic. iv. t. 1494 = febrifugum.
andrographioides, T. Anders. in Journ. Linn. Soc. ix. (1867) 504 = Andrographis tenuiflora.
canescens, *T. Anders. l. c.* 505.—Ind. or.
cerylanicum, *Arn. & Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 311.—Zeylan.
Cuminghianum, *Nees, in DC. Prod.* xi. 94.—Ins. Philipp.
febrifugum, *Benth. in Flora*, xxxii. (1849) 558.—Ind. or. glabrum, *T. Anders. in Journ. Linn. Soc. ix.* (1867) 506.—Ind. or.
hirsutum, *T. Anders. in Thw. Enum. Pl. Zeyl.* 233.—Zeylan.
javanicum, *Nees, in DC. Prod.* xi. 93.—Java.
latifolium, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 505.—Ind. or.
leptostachyum, *Nees, in Wall. Pl. As. Rar.* iii. 106.—Burma.
longifolium, T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 509 = canescens.
ovatum, T. Anders. ex Bedd. Ic. Pl. Ind. Or. i. 60 = Andrographis ovata.
paniculatum, *T. Anders. in Thw. Enum. Pl. Zeyl.* 232.—Zeylan.
Parishii, T. Anders. in Journ. Linn. Soc. ix. (1867) 504 = Andrographis tenuiflora.
Pearcei, Hort. Veitch, in Gard. Chron. (1867) 458 = Verschaffelii ?
polyanthum, *Wight, Ic. iv.* t. 1494.—Ind. or.
sanguinolentum, *T. Anders. in Thw. Enum. Pl. Zeyl.* 232.—Zeylan.
serrulatum, T. Anders. in Journ. Linn. Soc. ix. (1867) 505 = canescens.
Thwaitesii, *T. Anders. in Thw. Enum. Pl. Zeyl.* 232.—Zeylan.
tomentosum, *T. Anders. in Journ. Linn. Soc. ix.* (1867) 505.—Ind. or.
Verschaffelii, *Lem. Illustr. Hortic.* x. (1863) t. 372.—Bras.
venustum, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 506.—Reg. Himal.

GYMNOSTECHYUM, Spach, Hist. Vég. Phan. ix. 146 (1840) = **Gymnostachyum**, Nees (Acanthac.).

GYMNOSTEPHIUM, Less. Syn. Comp. 185 (1832). *COMPOSITAE*. Benth. & Hook. f. ii. 266.
Heteractis, DC. Prod. vi. 468 (1837).
angustifolium, *Harv. in Harv. & Sond. Fl. Cap.* iii. 67.—Afr. austr.
ciliare, *Harv. l. c.* 68.—Afr. austr.
corymbosum, *Harv. l. c.*—Afr. austr.
fruticosum, *DC. Prod.* v. 300.—Afr. austr.
gracile, *Less. Syn. Comp.* 185.—Afr. austr.
hirsutum, *Less. l. c.*—Afr. austr.

GYMNOSTICHUM, Schreb. Beschr. Graes. ii. 127 t. 47 (1810) = **Asperella**, Humb. (Gramin.).
californicum, Boland. ex Thurb. in S. Wats. Bot. Calif. ii. 327 = A. californica (in Addend.).
gracilis, Hook. f. & t. N. Zel. 312. t. 70.—N. Zel.
Hystrix, Schreb. Beschr. Graes. ii. 127 = A. Hystrix.
majus, Heynh. Nom. i. 371 = A. Hystrix.

GYMNOSTILLINGIA, Muell. Arg. in Linnaea, xxxii. (1863) 89 = **Stillingia**, Garden (Euphorb.).
acutifolia, Muell. Arg. l. c. 90 = S. acutifolia.
loranthacea, Muell. Arg. in Mart. Fl. Bras. xi. ii. 541. —Bras.
macrantha, Muell. Arg. in Linnaea, xxxii. (1863) 90 = S. macrantha.

GYMNOSTYLES, Juss. in Ann. Mus. Par. iv. (1804) 258. t. 16 = **Boliva**, Ruiz & Pav. (Compos.).
alata, Spreng. Syst. iii. 500 = S. sessilis.
anthraifolia, A. Juss. in Ann. Mus. Par. iv. (1804) 262. t. 61 = S. anthemifolia.
Barchlayana, Steud. Nom. ed. II. i. 713 = S. sessilis.
chilensis, Spreng. Syst. iii. 500 = S. sessilis.
lanceolata, Spreng. l. c. = S. nasturtiifolia.
minuta, Spreng. l. c. = Cotula pygmaea.
Mutisii, Spreng. l. c. = S. Mutisii.
nasturtiifolia, A. Juss. in Ann. Mus. Par. iv. (1804) 262. t. 16. f. 2 = S. nasturtiifolia.
parthenifolia, [Moç. & Sesse,] ex DC. Prod. v. 530 (= *Euphrosyne parthenifolia*).—Mexico.
pedunculata, Moç. ex DC. l. c. vi. 148 = S. pediculata.
peruviana, Spreng. Syst. iii. 500 = S. pedicellata.
pterosperma, A. Juss. in Ann. Mus. Par. iv. (1804) 262. t. 16. f. 3 = S. sessilis.
stoloniféra, A. Juss. l. c. = S. nasturtiifolia.

GYMNOSTYLIS, Rafin. in Am. Monthly Mag. (1818) 268 = **Ptuchea**, Cass. (Compos.).
marilandica, Rafin. l. c. = P. camphorata.

GYMNOTERPE, Salisb. Gen. Pl. Fragm. 108 (1866) = **Tapeinanthus**, Herb. (Amaryllid.).
humile, Salisb. l. c. = T. humilis.

GYMNOTHECA, Decne. in Ann. Sc. Nat. Sér. III iii. (1845) 100. t. 5. *PIPERACEAE*, [Benth. & Hook. f. iii. 126].
chinensis, *Decne. l. c.*—China.

GYMNOTHRIX, Spreng. Anleit. ii. t. 154 (1817) = seq.

GYMNOTRIX, Beauv. Agrost. 59. t. 13. f. 6 (1812) = **Pennisetum**, Pers. (Gramin.).
adsurgens, Hochst. ex Steud. Nom. ed. II. i. 713 = P. adcense.
alopecuroides, Nees, ex Steud. Syn. Pl. Gram. 102 = P. alopecuroides.
bambusiformis, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 507 = P. bambusiforme.
caudata, Schrad. in Goett. Gel. Anz. (1821) iii. 2073 = P. macrourum.
cenchroides, Roem. & Schult. Syst. ii. 499 = P. compressum.
cenchroides, Wight, ex Steud. Syn. Pl. Gram. 102 = P. alopecuroides.
chilensis, Desv. in C. Gay, Fl. Chil. vi. 251. t. 74.—Chili.
cladodes, Hochst. ex Steud. Syn. Pl. Gram. 104 = P. glaucifolium.
complanata, Nees, in Bonplandia, iii. (1855) 83 = P. complanatum.
compressa, Brongn. in Duperr. Voy. Coq. Bot. 105 = P. compressum.

GYMNOTRIX :—
crinita, H. B. & K. Nov. Gen. et Sp. i. 112 = Peniscetum Humboldtianum.
distachya, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 509 = P. tristachyon.
dominguensis, Presl, ex Schult. Mant. ii. 284 = Setaria Venenatii.
elegans, Buese, in Miq. Pl. Jungh. 368.—Java.
geniculata, Schult. Mant. ii. 284 = P. Sieberi.
gigantea, Walp. Ann. iii. 723 = P. giganteum. *A. Rich. glabra*, Hochst. in Flora, xxiv. (1841) i. Intell. 19, nomen; et ex Steud. Syn. Pl. Gram. 104 = P. glabrum.
glaucifolia, Walp. Ann. iii. 722 = P. glaucifolium.
Grisebachiana, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 508 = P. mexicanum.
hordeiformis, Nees, in Linnaea, vii. (1832) 276 = P. compressum.
hordeoides, Kunth, Rév. Gram. i. 48 = Panicum hordeoides.
humilis, Walp. Ann. iii. 722 = *P.* humile.
japonica, Kunth, Enum. Pl. i. 158 = P. compressum.
latifolia, J. & C. Presl, Rel. Haenk. i. 316 = P. Preslii.
latifolia, Schult. Mant. ii. 601 = P. latifolium.
macrostachya, Brongn. in Duperr. Voy. Coq. Bot. 104 = P. macrostachyum.
mexicana, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 508 = P. mexicanum.
nervosa, Nees, Agrost. Bras. 277 = P. nervosum.
nigricans, J. & C. Presl, Rel. Haenk. i. 316 = P. nigricans.
nitens, Anders. in Peters, Reise Mossamb. Bot. 552.—Afr. austr.
nubica, Hochst. in Flora, xxv. (1842) i. Beibl. 133, nomen; et ex Steud. Syn. Pl. Gram. 118 = Reckera nubica.
paspora, Nees, ex Steud. l. c. 102 = P. pauperum.
peruviana, Doell, in Mart. Fl. Bras. ii. ii. 302 = P. peruvianum.
Preslii, Kunth, Enum. Pl. i. 160 = P. Preslii.
purpurascens, Schrad. in Goett. Gel. Anz. (1821) iii. 2072 = P. alopecuroides.
purpurascens, Sw. ex Steud. Nom. ed. II. i. 713 = P. compressum.
Quartinianus, Walp. Ann. iii. 723 = P. Quartinianum.
rigida, Griseb. in Goett. Abh. xix. (1874) 263.—Reg. Argent.
riparia, Walp. Ann. iii. 722 = P. riparium.
riparioides, Walp. l. c. 723 = P. riparioides.
Schimperi, Hochst. in Flora, xxiv. (1841) i. Intell. 19, nomen; et in A. Rich. Tent. Fl. Abyss. ii. 381 = P. Schimperi.
sphacelata, Nees, Fl. Afr. Austr. 68.—Afr. austr.
Thouarsii, Steud. Nom. ed. I. 356 = seq.
Thuarii, Beauv. Agrost. 60. t. 13 = P. compressum.
tristachya, H. B. & K. Nov. Gen. et Sp. i. 113 = P. tristachyon.
uniseta, Nees, Fl. Afr. Austr. 68 = P. unisetum.

GYMNOXIS, Hochst. Nom. ed. II. i. 620, in syn. (1840) sphalm. = GYMNOPSIS, DC. = **Gymnolomia**, H. B. & K. (Compos.).

GYMOSTYLES, Willd. in Ges. Nat. Fr. Berl. Mag. i. (1807) 141, sphalm. = GYMNOSTYLES, Juss. = **Solliva**, Ruiz & Pav. (Compos.).

GYNACTIS, Cass. in Dict. Sc. Nat. ii. 475 (1827) = **Glossogyne**, Cass. (Compos.).

GYNAECOCEPHALUM, Hassk. Cat. Hort. Bogor. Alt. 78 (1844) = GYNOCEPHALUM, Blume = **Phytocrene**, Wall. (Olacin.).

GYNAECOPACHYS, Hassk. Cat. Hort. Bog. Alt. 115 (1844) = GYNOPACHIS, Blume = **Randia**, Linn. (Rubiac.).

GYNAECOTROCHUS, Hassk. Cat. Hort. Bog. Alt. 218 (1844) = **Gynotrochea**, Blume (Rhizoph.).

GYNAECURA, Hassk. Cat. Hort. Bog. Alt. 108 (1844) = **Gynura**, Cass. (Compos.).

GYNAION, A. DC. in DC. Prod. ix. 468 (1845) = **Cordia**, Linn. (Borag.).
ovatifolium, A. DC. l. c. = C. ventita.

GYNAMBLOSIS, Torr. in Rep. Marcy Exped. 295 (1858) = **Croton**, Linn. (Euphorb.).
maritima, Baill. Recul. Obs. Euph. 272 = C. maritimus.
monanthogynus, Torr. in Rep. Marcy Exped. 295 = C. monanthogynus.

GYNAMPSIS, Rafin. Fl. Tellur. iii. 5 (1836) = **Downingia**, Torr. (Campanular.).

GYNANDRIRIS, Parl. Nuov. Gen. e Sp. 49 (1854) = **Iris**, Tourn.
monophylla, Klatt, in Linnaea, xxxiv. (1865–66) 578 = I. Sisyrinchium.
Sisyrinchium, Parl. Nuov. Gen. e Sp. 52 = I. Sisyrinchium.

GYNANDROPSIS, DC. Prod. i. 237 (1824). *CAPPARIDEAE*, Benth. & Hook. f. i. 106.
Roerpera, F. Muell. in Hook. Kew Journ. ix. (1857) 15.
Sinapistrum, Medic. Phil. Bot. i. 108 (1789).
adenocarpa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 316.—Ecuador.
affinis, Blume, Bijdr. 51 = pentaphylla.
aurantiaca, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 315.—Venezuela.
brachycarpa, *DC. Prod.* i. 238.—Peruv.
Candelabrum, Sweet, Hort Brit. ed. I. 468 = Cleome Candelabrum.
coccinea, *Benth. Pl. Hartw* 160.—N. Granat.
densiflora, *Benth. l. c.*—N. Granat.
denticulata, DC. Prod. i. 238 = pentaphylla.
glandulosa, *Presl, Bot. Bemerk.* 8.—Aegypt.
grandiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 314.—Mexic.
heterotricha, DC. Prod. i. 238 = pentaphylla.
hispidula, *DC. l. c.*—Peruv.
macrophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 314.—Venezuela.
Muelleri, *Benth. Fl. Austral.* i. 91.—Australasia.
muricata, Schrad. Cat. Sem. Hort. Gotting. (1824); ex Schult. f. Syst. vii. 95 = Cleome muricata.
ophitocarpa, DC. in Mém. Soc. Phys. Genèv. v. (1832) 144 t. 2 = Cleome muricata.
palmipes, DC. Prod. i. 238 = pentaphylla.
pentaphylla, *DC. l. c.*—Cosmop. trop.
phoenicea, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii (1854) ii. 316.—N. Granat.
pulchella, Lindl. in Trans. Hort. Soc. vii. (1830) 65 = Cleome pulchella.
sessilifolia, DC. Prod. i. 237 = Cleome sessilifolia.
suaica, Miq. in Journ. Bot. Néerl. i. (1861) 128 = pentaphylla.
speciosa, *DC. Prod.* i. 238.—Mexic.
trichopus, *Benth. Bot. Voy. Sulph.* 64.—Ecuador.
triphylla, DC. Prod. i. 237 = pentaphylla.
viscida, Bunge, Mém. Sav. Etr. Pétersb. ii. (1835) 81 = pentaphylla.
viscosa, Walp. Rep. ii. 764, sphalm. pro *Polanisia viscosa* = Cleome graveolens.

GYNANTHISTROPHE, Poit. ex DC. Prod. ii. 424 (1825) = **Swartzia**, Schreb. (Legumin.).

GYNAPHANES, Steetz, in Peters, Reise Mossamb. Bot. 457 (1864) = **Ipomœa**, Cass. (Compos.).
australis, Steetz, l. c. 458 = E. Cunninghamii.

GYNAPTEINA, Spach, Hist. Vég. Phan. viii. 113 (1839) = **Keptapleurum**, Gaertn. (Araliac.).

GYNASTRUM, Neck. Elem. i. 224 (1790) = **Guapira**, Aubl. (Incert. sed.).

GYNATRIX, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 84 = **Plagianthus**, Forst. (Malvac.).
pulchella, Alef. l. c. 86 = P. pulchellus.

GYNEMA, Rafin. Fl. Ludov. 68 (1817) = **Ptuchea**, Cass. (Compos.).
argyrea, Rafin. l. c.—Am. bor.
balsamica, Rafin. l. c. = P. camphorata.
dentata, Rafin. ex DC. Prod. v. 429 = P. camphorata.
parviflora, Rafin. Fl. Ludov. 64.—Am. bor.
viscida, Rafin. Ann. Nat. (1820) 15 = Ptuchea camphorata.

GYNERIUM, Humb. & Bonpl. Pl. Aequin. ii. 119. t. 115 (1809). *GRAMINEAE*, Benth. & Hook. f. iii. 1178.
argenteum, *Nees, Agrost. Bras.* 462.—Bras.
attacamense, *Phil. in Linnaea*, xxxiii. (1864–65) 289. —Chili.
jubatum, *Lemoine, ex Carr. in Rev. Hortic.* (1878) 449.—Ecuador.
modestum, *Doell, in Mart. Fl. Bras.* ii. 111. 240.—Bras.
Nereii, Meyen, Reise, i. 380, 407 = G. Quila.
parviflorum, *Nees, in Mart. Agrost. Bras.* ii. 463.—Am. trop.
procerum, Beauv. Agrost. 164 = saccharoides.
pygmaeum, [*Nees, in*] *Meyen, Reise*, i. 484, nomen.—Peruv.
Quila, *Nees & Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 163.—Chili.
saccharoides, *Humb. & Bonpl. Pl. Aequin.* ii. 112. t. 115.—Am. austr.
sagittatum, Beauv. Agrost. 188 = saccharoides.
speciosum, [Nees, in] Meyen, Reise, i. 407; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 153 = argenteum.
seelandicum, Steud. Syn. Pl. Gram. 198 = Arundo conspicua.

GYNESTUM, Polt. in Mém. Mus. Par. ix. (1822) 387 = **Geonoma**, Willd. (Palm.).
acaule, Polt. l. c. 391. t. 20 = Geon. Poiteanana.
baculiforme, Polt. l. c. 389. t. 17 = Geon. baculifera.
dieversum, Polt. l. c. 390 = Geon. diversa.
maximum, Polt. l. c. 388. t. 16 = Geon. maxima.
strictum, Polt. l. c. 391. t. 19 = Geon. pycnostachys.

GYNETERIA, Spreng. Anleit. ii. 11. 899 (1818) = GYNHETERIA, Willd. = **Tessaria**, Ruiz & Pav. (Compos.).

GYNETRA, Rafin. Sylva Tellur. 165 (1838) = **Tetracera**, Linn. (Dilleniac.).
volubilis, Rafin. l. c. = T. volubilis.

GYNHETERIA, Willd. in Ges. Naturf. Fr. Berl Mag. i. (1807) 140 = **Tessaria**, Ruiz & Pav. (Compos.).
dentata, Spreng. Syst. iii. 492 = T. integrifolia.
incana, Spreng. Neue Entdeck. ii. 135 = T. integrifolia.
salicifolia, Willd. ex Less. Syn. Comp. 208 = T. integrifolia.

GYNIZODON, Rafin. Fl. Tellur. iv. 40 (1836) = **Miltonia**, Lindl. (Orchid.).
Russelianum, Rafin. l. c. = M. Russeliana.

GYNOCAMPUS, Lesch. ex DC. Prod. vii. 338 (1839) = **Levenhookia**, R. Br. (Stylid.).

GYNOCARDIA, R. Br. in Roxb. Fl. Corom. iii. 95. t. 299 (1819). *BIXINEAE*, Benth. & Hook. f. i. 129.
CHAULMOOGRA, Roxb. Hort. Beng. 105 (1814); Fl. Ind. iii. 835 (1832).
CHILMORIA, Buch-Ham. in Trans. Linn. Soc. xiii. (1822) 500.
odorata, *R. Br. in Roxb. Pl. Corom.* iii. 95.—Ind. or.

GYNOCEPHALA, Benth. & Hook. f. Gen. i. 854 (1862), sphalm. = seq.

GYNOCEPHALIUM, Endl. Gen. 281 (1837), sphalm. = seq.

GYNOCEPHALUM, Blume, Bijdr. 483 (1825) = **Phytocreme**, Wall. (Oliacin.).
bracteatum, Trécul, in Ann. Sc. Nat. Sér. III. viii. (1847) 149 = P. bracteata.
giganteum, Trécul, l. c. = P. gigantea.
luzoniense, Llanos, ex Baill. Adansonia, x. (1872) 282 = P. luzoniensis.
macrophyllum, Blume, Bijdr. 483 = P. macrophylla.
oblongum, Trécul, in Ann. Sc. Nat. Sér. III. viii. (1847) 148 = P. oblonga.
palmatum, Trécul, l. c. = P. palmata.

GYNOCHTHODES, Blume, Bijdr. 992 (1826) (Gynochodes). *RUBIACEAE*, Benth. & Hook. f. ii. 119.
corinces, *Blume, l. c.*—Java.
macrophylla, *Korz, in Journ. As. Soc. Beng.* xli. (1872) ii. 314.—Ins. Andaman. ; Malaya.
sublanceolata, *Miq. Fl. Ind. Bat. Suppl.* 548.—Malaya.

GYNODON, Rafin. Fl. Tellur. ii. 18 (1836) = **Allium**, Tourn. (Liliac.).
cernuum, Rafin. l. c. = A. cernuum.
Ellioti, Rafin. l. c. = A. cernuum ?
rupestre, Rafin. l. c. = A. paniculatum.

GYNOISA, Rafin. Fl. Tellur. iv. 75 (1836) = **Ipomoea**, Linn. (Convolv.).
carolina, Rafin. l. c. = I. commutata.

GYNOMPHIS, Rafin. Sylva Tellur. 98 (1838) = **Miconia** ?, Ruiz & Pav. (Melastom.).
argentea, Rafin. l. c. = M. argentea ?

GYNOON, A. Juss. in Mém. Mus. Par. x. (1823) 335; et Euph. Tent. 17. t. 8 (1824) = **Glochidion**, Forst. (Euphorb.).
Heyneanum, Wight & Arn. in Edinb. N. Phil. Journ. (Apr. 1833) 300 = Gloch. Heyneanum.
hirsutum, Wight, Ic. t. 1909 = Gloch. Moonii.
Justricanum, Wight, l. c. v. 11. 29 = Gloch. rigidum.
rigidum, A. Juss. Euph. Tent. 107. t. 3 = Gloch rigidum.
tetandrum, D. Dietr. Syn. Pl. v. 868, sphalm. = seq.
triandrum, Wight & Arn. in Edinb. N. Phil. Journ. (Apr. 1833) 300 = Gloch. rigidum.

GYNOPACHIS, Blume, Cat. Gew. Buitenz. 48 (1823) = **Randia**, Linn. (Rubiac.).
abnormis, Korth. in Nederl. Kruidk. Arch. ii. ii. (1851) 182.—Borneo.
acuminata, Blume, Cat. Gew. Buitenz. 48.—Java.
attenuata, Korth. in Nederl. Kruidk. Arch. ii. ii. (1851) 182.—Sumatra.
axillaris, Miq. Fl. Ind. Bat. ii. 221 = R. densiflora.
corymbosa, Blume, Cat. Gew. Buitenz. 48.—Java.
deflectens, Miq. Fl. Ind. Bat. ii. 222.—Sumatra.
oblongata, Miq. l. c. 221 = R. densiflora.
rufidula, Miq. l. c.—Sumatra.
tomentosa, Blume, Bijdr. 984.—Java.
Zippeliana, Scheff. in Ann. Jard. Buitenz. i (1876) 28. —N. Guin.

GYNOPACHYS, Blume, Bijdr. 988 (1825) = prec.

GYNOPLEURA, Cav. Ic. iv 51 (1791). *PASSIFLOREAE*, Benth. & Hook. f. i. 810.
angustifolia, Walp. Rep. v. 775, sphalm. = linearifolia.
carcules, *Prest. Rel. Haenk.* ii. 46.—Chili.
coronata, *M. Roem. Syn. Pepon.* 211.—Chili.
dilatata, Walp. Rep. ii. 238, sphalm. = humilis.
fasciculata, *M. Roem. Syn. Pepon.* 210.—Chili.
humilis, *M. Roem. l. c.* 211.—Chili.
linearifolia, *Cav. Ic.* iv. 52.—Chili.
paniculata, *M. Roem. Syn. Pepon.* 211.—Chili.
solanoides, *M. Roem. l. c.*—Chili.
tenuifolia, M. Roem. l. c. = Malesherbia tenuifolia.
tubulosa, Cav. Ic. iv. 52 = Malesherbia thyrsiflora.

GYNOPOGON, Forst. Char. Gen. 35. t. 18 (1775) = **Alyxia**, Banks (1810) (Apocyn.).
Alyxia, Forst. f. Prod. 19 = A. Gynopogon.
breviflorum, Kurz, For. Fl. Brit. Burma, ii. 177.—Burma.
Hunteeii, Roxb. Hort. Beng. 19, nomen.—Ins. Penang.
lanceolatum, Kurz, For. Fl. Brit. Burma, ii. 177 = Hunteria corymbosa.
pugioniformis, A. Cunn. ex Steud. Nom. ed. II. i. 714 = A. ruscifolia.
scandens, Forst. Char. Gen. 36 = A. scandens.
stellatum, Forst. l. c. = A. stellata.

GYNOSTEMMA, Blume, Bijdr. 23 (1825). *CUCURBITACEAE*, Benth. & Hook. f. i. 839.
ENKYLIA, Griff. Fl. Cantor. 26. t. 4 (1837 ?); et in Journ. As. Soc. Beng. xxiii. (1854) 645.
PENTALOBIA, Zoll. & Mor. Syst. Verz. Zoll. 81 (1846).
Blumei, Hassk. Cat. Hort. Bog. Alt. 311 = pedatum, simplicifolium.
cissoides, Benth. & Hook. f. Gen. i. 839 = pedatum.
hederaefolium, *Cogn. in DC. Monog. Phan.* iii. 916. —Ins. Timor.
integrifolium, *Cogn. l. c.*— Ins. Philipp.
laxum, Cogn. l. c. 914 = Wightianum.
pedatum, *Blume, Bijdr.* 23.—As. trop.
simplicifolium, *Blume, l. c.* 24.—Java.
Wightianum, *Benth. & Hook. f. Gen.* i. 839.—Ind. or. ; Malaya.

GYNOTROCHES, Blume, Bijdr. 218 (1825). *RHIZOPHOREAE*, Benth. & Hook. f. i. 681.
DRYPTOPETALUM, Arn. in Ann. Nat. Hist. i. (1838) 372.
GYNACOTROCHUS, Hassk. Cat. Hort. Bog. Alt. 218 (1844).
axillaris, *Blume, Bijdr.* 219.—Malaya.
Dryptopetalum, Blume, Mus. Bot. Lugd. Bat. i. 127 = axillaris.
membranifolia, *Miq. Anal. Bot. Ind.* iii. 7.—Ind. or.
micrantha, *Blume, Mus. Bot. Lugd. Bat.* i. 127.—Malaya.
reticulata, A. Gray, Bot. U. St. Expl. Exped i. 607 = axillaris.

GYNOXIS, Reichb. Consp. 106 (1828) = seq.

GYNOXYS, Cass. in Dict. Sc. Nat. xlviii. 455 (1827). *COMPOSITAE*, Benth. & Hook. f. ii. 450.
SCROBICARIA, Cass. in Dict. Sc. Nat. xlviii. 456 (1827).
albiflora, *Wedd. Chlor. And.* i. 78.—N. Granat.
alternifolia, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 581, nomen.—Bolivia.
asteroirichia, *Sch. Bip. l. c.* 529.—Bolivia.
auriculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii. 86.—Peruv.
baccharoides, *Cass. in Dict. Sc. Nat.* xlviii. 455.—N. Granat. ; Ecuador.
Berlandieri, DC. Prod. vi. 326 = Senecio Berlandieri.
bazifolia, *Cass. in Dict. Sc. Nat.* xlviii. 455.—Ecuador.
cordifolia, *Cass. l. c.* 456.—Peruv.
Cummingii, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel (1852) 106 = Senecio calocephalus, *Hemsl.*
fragrans, Hook. Bot. Mag. t. 4511 = Senecio Skinneri.
fuliginosa, *Cass. in Dict. Sc. Nat.* xlviii. 455.—N. Granat.
glabrata, Less. Syn. Comp. 390 = Senecio Swartzii.
Haenkei, DC. Prod. vi. 326 = Senecio kermesinus.
heterophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) ii 85.—Ecuador.
hirsuta, *Wedd. Chlor. And.* i. 79.—N. Granat.
ilicifolia, *Wedd. l. c.* 80, in obs.—N. Granat.
incana, *Less. Syn. Comp.* 390.—Jamaic.
lacinsata, Less. l. c. = Senecio Swartzianus.
laurifolia, *Cass. in Dict. Sc. Nat.* xlviii. 455.—Ecuador.
Lindenii, Sch. Bip. ex Wedd. Chlor. And. i. 76 = baccharoides.
longifolia, *Wedd. l. c.* 79.—Peruv.
lucida, Less. Syn. Comp. 390 = Senecio lucidus.
Mandonii, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 80.—Bolivia.
Moritziana, *Sch. Bip. ex Wedd. Chlor. And.* i. 79.—Venezuela.
Oerstedii, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1859) 107.—Am. centr.
pendula, *Sch. Bip. ex Wedd. Chlor. And.* i. 78.—N. Granat.
Peopppigii, *DC. Prod.* vi. 326.—Peruv.
prenanthifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 207.—Peruv.
pulchella, *Cass. in Dict. Sc. Nat.* xlviii. 455.—Ecuador.
repanda, *Wedd. Chlor. And.* i. 77.—Bolivia.
scabra, *Benth. Bot. Voy. Sulph.* 121.—Ecuador.
Sinclairi, *Benth. l. c.* 120.—N. Granat.
verrucosa, *Wedd. Chlor. And.* i. 77.—N. Granat.
violacea, *Sch. Bip. ex Wedd. l. c.*—Venezuela.

GYNURA, Cass. in Dict. Sc. Nat. xxxiv. 391 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 445.
CRASSOCEPHALUM, Moench. Meth. 516 (1794), nomen prius.
CREMOCEPHALUM, Cass. in Dict. Sc. Nat. xxxiv. 390 (1825).
GYNAECURA, Hassk. Cat. Hort. Bog. Alt. 108 (1844).
affinis, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 201 = sarmentosa.
amplexicaulis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 408.—Afr. trop.
angulosa, *DC. Prod.* vi. 298.—Ind. or. ; Burma.
angulosa, Hance, in Journ. Bot. xxi. (1883) 322 = bicolor.
aurantiaca, Benth. in Hook. Niger Fl. 437 = vitellina.
aurantiaca, *DC. Prod.* vi. 300.—Java.

GYNURA:—
auranticaca, Sch. Bip. ex Miq. Fl. Ind. Bat. ii. 99 = densiflora.
auriculata, *Cass. Opusc. Phyt.* iii. 100.—China.
bicolor, *DC. Prod.* vi. 299.—Malaya.
bulbosa, Hook. & Arn. Bot. Beech. Voy. 194 = auriculata.
carnosula, *Zoll. & Mor. in Nat. en Geneesk. Arch. Ned. Indie*, ii. (1845) 265.—Java.
cernua, *Benth. in Hook. Niger Fl.* 437.—Afr. trop.
crepidioides, *Benth. l. c.* 435.—Afr. trop.
densiflora, *Miq. Fl. Ind. Bat.* ii. 99.—Java.
densifolia, *Miq. l. c.*—Java.
dichotoma, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 1. 202.—Java.
divaricata, *DC. Prod.* vi. 301.—China.
divaricatifolia, Sch. Bip. ex Aschers. in Schweinf. Beitr. Fl. Aethiop. 156 = crepidioides.
dubia, *Cass. in Dict. Sc. Nat.* xxxiv. 392.—Ind. or.
ecalyculata, DC. Prod. vi. 298 = Emilia sonchifolia.
Finlaysoniana, DC. l. c. 299 = sarmentosa.
foetens, DC. l. c. 300 = nepalensis.
glabrata, Hook. f. Fl. Brit. Ind. iii. 326 = auriculata.
haematophylla, *DC. Prod.* vi. 299.—Hab.?
hispida, Thw. Enum. Pl. Zeyl. 166 = G. Pseudo-China.
Lobbiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 202.—Java.
lycopersicifolia, *DC. Prod.* vi. 300.—Ind. or.
lyrata, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 124, 127.—Java.
microcephala, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 323 = crepidioides.
minuta, *Welw. Apont.* 586.—Afr. trop.
mollis, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 124, 127.—Java.
nepalensis, *DC. Prod.* vi. 300.—Reg. Himal.; Burma.
nitida, *DC. in Wight, Contrib.* 24; *et Prod.* vi. 299.—Ind. or.
nudiculis, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 351 = G. Pseudo-China.
ovalis, DC. Prod. vi. 300 = auriculata.
pinnatifida, *DC. l. c.* 301.—China.
polycephala, Benth. in Hook. Niger Fl. 437 = crepidioides.
Pseudo-China, Benth. Fl. Hongk. 189 = auriculata.
Pseudo-China, *DC. Prod.* vi. 299.—As. trop.; Austral.
purpurascens, DC. l. c. = bicolor?
sagittaria, *DC. l. c.* 301.—Ind. or.
sarcobasis, *DC. l. c.* 300.—Madag.
sarmentosa, *DC. l. c.* 298.—Malaya.
scabra, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 201 = sarmentosa.
simplex, Dals. & Gibs. Bomb. Fl. 180 = nitida, angulosa.
sinuata, DC. Prod. vi. 301 = G. Pseudo-China.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 536.—Sumatra.
vitellina, *Benth. in Hook. Niger Fl.* 438.—Afr. trop.
Walkeri, Wight, Ic. t. 1122 = Notonia Walkeri.
Zollingeriana, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126, nomen.—Java.

GYPOTHAMNIUM, Phil. Fl. Atac. 27. t. 3 (1860).
COMPOSITAE, Benth. & Hook. f. ii. 486.
pinifolium, *Phil. l. c.*—Chili.

GYPSOCALLIS, Salisb. ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 396 (1821) = Ehrtea, Tourn.
Actaea, G. Don, Gen. Syst. iii. 809 = E. peltata.
acutangula, G. Don, l. c. = E. Ilgetia.
articularis, G. Don, l. c. 808 = E. articularis.
Bedfordiana, G. Don, l. c. = E. turgida.
cassia, G. Don, l. c. = E. imbricata.
carnea, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 163 = E. carnea.
cistifolia, G. Don, Gen. Syst. iii. 804 = E. cistifolia.
concava, G. Don, l. c. 805 = E. bicolor.
distachyota, G. Don, l. c. 803 = E. diotaeflora.
dumosa, G. Don, l. c. 802 = E. dumosa.
eriocephala, Benth. in DC. Prod. vii. 617, in syn. sphalm. = E. villosa.
expromptu, G. Don, Gen. Syst. iii. 804 = Simochellus depressus.
fusilla, G. Don, l. c. 801 = E. racemosa.
formosa, G. Don, l. c. 803 = E. imbricata.
fragrans, G. Don, l. c. = E. fragrans.
hispida, G. Don, l. c. 801 = E. racemosa.
hortensialis, G. Don, l. c. = E. cumulifera.
imbricata, G. Don, l. c. 803 = E. imbricata.

GYPSOCALLIS:—
intersexta, G. Don, Gen. Syst. iii. 803 = Erica empetrifolia.
lanata, G. Don, l. c. 801 = E. lanata.
lasciva, G. Don, l. c. 803 = E. lasciva.
leucanthera, G. Don, l. c. 801 = E. leucanthera.
Leddigesii, G. Don, l. c. 803 = E. bicolor.
longipedunculata, G. Don, l. c. = E. dumosa.
lyrigera, G. Don, l. c. 804 = E. nigrita.
manniphlora, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 163 = E. verticillata.
mediterranea, D. Don, l. c. = E. mediterranea.
melanthera, G. Don, Gen. Syst. iii. 803 = E. melanthera.
microstoma, G. Don, l. c. 801 = E. nudiflora.
multiflora, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 163 = E. multiflora.
nigricans, G. Don, Gen. Syst. iii. 802 = E. nigrita.
nigrita, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 163 = E. nigrita.
nudiflora, D. Don, l. c. = E. nudiflora.
paniculata, G. Don, Gen. Syst. iii. 803 = E. paniculata.
patens, G. Don, l. c. 802 = E. patens.
peltata, G. Don, l. c. 803 = E. peltata.
pilosa, G. Don, l. c. 801 = E. villosa.
Pohlmannii, G. Don, l. c. 803 = E. Lachnaea.
procumbens, G. Don, l. c. 801 = E. procumbens.
purpurascens, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 163 = E. mediterranea.
Rastellum, G. Don, Gen. Syst. iii. 801 = E. penicilliflora.
rostella, G. Don, l. c. 803 = E. penicilliflora.
squamaeflora, G. Don, l. c. = E. imbricata.
stamínea, G. Don, l. c. 802 = E. stamínea.
stylosa, G. Don, l. c. 803 = E. imbricata.
thalictriflora, G. Don, l. c. 802 = E. stamínea.
tiarasflora, G. Don, l. c. = E. tiaraeflora.
umbellata, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 163 = E. umbellata.
vagans, S. F. Gray, Nat. Arr. Brit. Pl. ii. 398 = E. vagans.
villosa, G. Don, Gen. Syst. iii. 802 = E. villosa.

GYPSOPHILA, Linn. Diss. Chen. 41 (1761); Amoen. Acad. iii. 28 (1756). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 146.
ACOSMIA, Benth. in Wall. Cat. n. 844 (1828 ?).
ANKYROTAENIUM, Fenzl, in Bot. Zeit. i. (1843) 393.
BANFFYA, Baumg. Enum. Stirp. Transs. i. 385 (1816).
DICHOGLOTTIS, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 25 (1835).
GYPSOPHYTUM, Ehrh. Beitr. iv. 149 (1789).
HETEROCHROA, Bunge, in Ledeb. Fl. Alt. ii. 131 (1830).
KATAKEIS, Medic. Malv. 111 (1787).
LANARIA, Adans. Fam. ii. 255 (1763).
PETRORHAGIA, Link, Handb. ii. 255 (1831).
PSAMMOPHILA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 345.
ROKEJEKA, Forsk. Fl. Aegypt. Arab. 90 (1775).
TIMAROSIA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 138. t. 33 (1862).
acerosa, *Boiss. Fl. Orient.* i. 558.—Afghan.
acutifolia, *Fisch. Cat. Hort. Gorenk.* (1812) 59.—Reg. Caucas.
acutifolia, *Stev. ex Spreng. Nov. Prov.* 21.—Rossia acutr.
adenophora, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 85.—Persia.
adenotricha, *Wacq. Hort. Vindob.* ii. 65. t. 136.—Orient.
aggregata, Linn. Sp. Pl. 406 = Arenaria tetraquetra.
agrestis, Pers. Syn. i. 492 = muralis.
alpestris, Jord. & Fourr. Brev. Fl. Nov. fasc. i. 9 = repens.
alpina, Habl. in J. G. Gmel. It. iv. 178 = Tunica stricta.
alsinoides, *Bunge, in Arb. Naturf. Ver. Riga*, i. 11. (1847) 179.—Persia; Baluchist.
altissima, Bieb. Fl. Taur. Cauc. i. 319 = scutiflora.
altissima, *Linn. Sp. Pl.* 407.—Sibir.
altissima, Sibth. & Sm. Fl. Graec. Prod. i. 280 = Arrostii.
anatolica, *Boiss. & Heldr. Diagn.* Ser. I. viii. 57.—As. Min.
arborea, *Medic. Beobachtt.* (1782) 160.—Hab.?
arenaria, Waldst. & Kit. Pl. Rar. Hung. i. 40. t. 41 = fastigiata.
arenicola, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 240 = Tunica Saxifraga.

GYPSOPHILA:—
aretioides, *Boiss. Diagn.* Ser. I. i. 9.—Persia.
armeniaca, Ser. in DC. Prod. i. 353 = Tunica Sibthorpii.
Arrostii, *Guss. Pl. Rar.* 160.—Europ. austr.; As. Min.
arvensis, Borckh. ex Stend. Nom. ed. I. 386 = muralis.
ascendens, *Medic. Beobachtt.* (1782) 165.—Hab.?
asperula, Dul. ex Nym. Consp. 160 = Tunica Saxifraga.
atropurpurea, Boiss. & Heldr. Diagn. Ser. I. viii. 58 = frankenioides.
Aucheri, *Boiss. l. c.* i. 12.—As. Min.; Syria.
australis, A. Gray, Bot. U. St. Expl. Exped. i. 112 = tubulosa.
bellidifolia, *Boiss. Diagn.* Ser. I. i. 11.—Arab.; Beluchist.
Boissieri, *Vis. in Mem. Ist. Venet.* xv. (1870) 16.—As. Min.
brachypetala, *Trautv. in Act. Hort. Petrop.* ii. (1873) 471.—Armenia.
Bungeana, *D. Dietr. Syn. Pl.* ii. 1542.—Sibir. altaic.
campestris, *Pall. Reise*, iii. 682; *ex Ledeb. Fl. Ross.* i. 300 (Quid?).—Rossia.
capitata, *Bieb. Fl. Taur. Cauc.* i. 321.—Reg. Caucas.
capitata, Ledeb. Fl. Ross. i. 300 = globulosa.
capituliflora, *Rupr. Sert. Tiensch.* 40.—As. med.
cappadocica, Boiss. & Bal. Diagn. Ser. II. vi. 26 = sphaerocephala.
carlcifolia, *Boiss. l. c.* Ser. I. i. 14.—Persia.
cerastoides, *D. Don, Prod. Fl. Nep.* 218.—Reg. Himal.
collina, Stev. ex Ser. in DC. Prod. i. 353 = fastigiata.
compressa, Desf. Fl. Atlant. i. 343. t. 97 = Tunica compressa.
compressa, Nym. Consp. 160 = Tunica graminea.
cordifolia, Fenzl, ex Boiss. Fl. Orient. i. 546 = ruscifolia.
cretica, Griseb. Spicil. Fl. Rumel. i. 184 = Tunica illyrica.
cretica, Sibth. & Sm. Fl. Graec. Prod. i. 280 = Tunica cretica.
curvifolia, *Fenzl, Pugill. Pl. Nov. Syr.* 10.—As. Min.
dahurica, Turcz. ex Stend. Nom. ed. II. i. 715 = Gmelini.
Damascena, *Boiss. Diagn.* Ser. I. viii. 57.—Syria; Persia.
davurica, *Turcz. ex Fenzl, in Ledeb. Fl. Ross.* i. 294.—Sibir. baical.
desertorum, *Fenzl, in Ledeb. Fl. Ross.* i. 295 (1846).—Sibir. altaic.
dianthoides, Sibth. & Sm. Fl. Graec. Prod. i. 280 = Tunica dianthoides.
dichotoma, Bess. Prim. Fl. Gal. i. 372 = Gmelini.
diffusa, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1839) 146 = elegans.
diotica, *Spreng. Syst.* iv. *Cur. Post.* 178.—Reg. Argent.
dubia, *Willd. Enum. Hort. Suppl.* 23.—Hungar.
effusa, Tausch, in Flora, xiii. (1830) i. 244 = paniculata.
elata, *Wender. ex Steud. Nom.* ed. II. i. 715, nomen.—Hab.?
elegans, *Bieb. Fl. Taur. Cauc.* i. 319.—As. Min.
erectiuscula, Jord. & Fourr. Brev. Fl. Nov. fasc. i. 9 = repens.
erinacea, *Boiss. Fl. Orient.* i. 557.—Afghanist.
eriocalyx, *Boiss. Diagn.* Ser. I. i. 13.—As. Min.
eubonensis, Sibth. & Boiss. Fl. Orient. i. 519 = Tunica Saxifraga.
fasciculata, *Margot & Reut. in Mém. Soc. Phys. Genèv.* vii. (1839) 261. t. 1.—Graecia.
fastigiata, *Linn. Sp. Pl.* 407.—Europ.
fastigiata, Willk. Ic. Fl. Hisp. i. 24 = hispanica.
filiformis, *Walls. Beitr. Fl. Hercyn.* 199; *et in Linnaea*, xiv. (1840) 540.—Europ.
filifera, Bien. ex Boiss. Fl. Orient. i. 544 = G. Rokejeka.
floribunda, *Turcz. ex Ledeb. Fl. Ross.* i. 278.—Persia; Soongar.
frankenioides, *Boiss. Diagn.* Ser. I. i. 10.—As. Min.; Syria.
fruticulosa, *Boiss. Fl. Orient.* i. 556.—As. Min.
Gasparrini, *Guss. Fl. Sic. Syn.* i. 474.—Sicil.
glandulosa, *Walp. Rep.* ii. 774; *Boiss. Diagn.* Ser. I. viii. 59, in obs.—Reg. Caucas.
glauca, *Stev. ex Ser. in DC. Prod.* i. 353.—Reg. Caucas.
globulosa, *Stev. in Flora*, xv. (1832) ii. *Beibl.* 24.—Caucas.

GYPSOPHILA:—
glomerata, Adams, in Web. & Mohr, Beitr. i. 54 = capitata.
glomerata, *Pall. ex Bieb. Fl. Taur. Cauc.* i. 321.—Tauria.
Gmelini, *Bunge, in Ledeb. Fl. Alt.* ii. 128.—Sibir. altaic.
graminea, Sibth. & Sm. Fl. Graec. Prod. i. 279 = Tunica graminea.
grandiflora, *Poir. Encyc. Suppl.* ii. 874.—Hab.?
Haynaldiana, Janka, ex Nym. Consp. 100 = Tunica illyrica.
hirsuta, *Spreng. Syst.* ii. 373.—Graec.; Syria; Arab.
hispanica, *Willd. in Flora*, xxxv. (1852) 589 ; *et Iber. Hab.* 119.—Hispan.
hispida, *Boiss. Diagn.* Ser. I. i. 11.—Cappadoc.
Honigbergeri, *Boiss. Fl. Orient.* i. 558.—Afghan.
illyrica, Nym. Syl. 287 = Tunica illyrica.
illyrica, Sibth. & Sm. Fl. Graec. Prod. i. 281 = Tunica Sibthorpii.
imbricata, *Rupr. Fl. Cauc.* 176.—Reg. Caucas.
intricata, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 238.—Turkestan.
Jaligua, Walp. Rep. ii. 776, sphalm. = saligna.
latifolia, *Fisch. ex Steud. Nom.* ed. II. i. 715, nomen. —Sibir.
lepidioides, *Boiss. Diagn.* Ser. I. i. 14.—Cappadoc.
libanotica, *Boiss. l. c.* 13.—As. Min. ; Syria.
linearifolia, *Boiss. Fl. Orient.* i. 550.—As. Min. ; Reg. Casp.
melampoda, *Bien. ex Boiss. l. c.* 551.—Persia.
Meyeri, *Rupr. Fl. Cauc.* 178.—Reg. Caucas.
microphylla, *Fenzl, in Ledeb. Fl. Ross.* i. 291.—Soongar.
minuartioides, *Boiss. Fl. Orient.* i. 558.—As. Min.
montana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 501.—Ins. Socotr. ; Arab.
multiculis, Poir. Encyc. Suppl. iii. 874 = Tunica Saxifraga.
muralis, *Linn. Sp. Pl.* 408.—Europ. ; As bor.
nana, *Bory & Chaub. Nouv. Fl. Pélop.* 26.—Graec.
nebulosa, Boiss. Diagn. Ser. I. viii. 58 = Arrostii.
ocellata, Sibth. & Sm. Fl. Graec. Prod. i. 281 = hirsuta.
ochroleuca, Sibth. & Sm. l. c. = Tunica ochroleuca.
Oldhamiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 187. —Japon.
olympica, *Boiss. Diagn.* Ser. I. viii. 55 —As. Min.
ortegioides, *Boiss. Fl. Orient.* i. 552.—As. Min.
pachygona, D. Dietr. Syn. Fl. ii. 1543 = Tunica pachygona.
paniculata, *Linn. Sp. Pl.* 407.—Europ. ; As. bor.
parviflora, Moench, Meth. 66 = paniculata.
Patrini, Ser. in DC. Prod. i. 353 = Gmelini.
perfoliata, DC. Prod. i. 352 = trichotoma.
perfoliata, *Linn. Sp. Pl.* 408.—Hispan.
permixta, Guss. Suppl. Fl. Sic. Prod. 120 = Tunica Saxifraga.
petraea, Fenzl, in Ledeb. Fl. Ross. i. 291 = Bungeana.
petraea, Reichb. Fl. Germ. Excurs. 801 = transylvanica.
picta, *Boiss. Fl. Orient.* i. 554.—Armenia.
pilulifera, *Boiss. & Heldr. Diagn.* Ser. I. viii. 56.—Lycia.
polyzala, *Fenzl, ex Boiss. Fl. Orient.* i. 542.—Persia.
porrigens, *Boiss. l. c.* 557.—As. Min. ; Persia.
prostrata, All. Fl. Pedem. ii. 78 = repens.
prostrata, Boeber, in Pall. N. Nord. Beitr. vi. 261 = Gmelini.
prostrata, *Linn. Sp. Pl.* 1108.—As. Med.
pubescens, *G. Don, Gen. Syst.* i. 383.—Italia.
pubescens, *Ledeb. ex Steud. Nom.* ed. II. i. 715, nomen. —Sibir.
pulposa, Gilib. Fl. Lituan. ii. 158 = fastigiata.
purpurea, Gilib. l. c. 154 = muralis.
ramosissima, Fisch. & Mey. ex Boiss. Fl. Orient. i. 550 = Szovitsii.
repens, Bieb. Fl. Taur. Cauc. i. 318 = Stevensi.
repens, *Linn. Sp. Pl.* 407.—Europ. ; As. bor.
reticulata, Hochst. in Lorent, Wanderungen, 241 = vaccifolia.
rigida, Linn. Sp. Pl. 408 = Tunica Saxifraga.
Rokejeka, *Delile, Fl. Egypte*, 231. t. 29. f. 1.—Aegypt. ; Mesopot.
rupestris, Turcz. ex Ledeb. Fl. Ross. i. 292 = Gmelini.
rusciifolia, *Boiss. Diagn.* Ser. I. i. 12.—As. Min. ; Persia.
saxatilis, Jord. & Fourr. Brev. Pl. Nov. Fasc. i. 9 = repens.
saxatilis, Stev. ex Ser. in DC. Prod. i. 352 = trichotoma.

GYPSOPHILA:—
saligna, *Schrad. in Linnaea*, x. (1836) *Litt.* 110.—Europ.
Saxifraga, Linn. Syst. ed. X. 1028 = Tunica Saxifraga.
scabra, Schult. ex Stend. Nom. ed. I. 396 ; ed. II. i. 716 = Tunica Saxifraga.
scorzonerifolia, Ser. in DC. Prod. i. 352 = perfoliata.
Scovitzii, Walp. Rep. ii. 774, sphalm. = Szovitsii.
sedifolia, *Kurz, in Flora*, lv. (1872) 386.—Reg. Himal.
serotina, Hayne, in Dreves & Hayne, Bot. Bildt. t. 152 ; et ex Willd. Enum. Hort. Berol. 464 = muralis.
serpyllifolia, *Boiss. & Heldr. Diagn.* Ser. I. viii. 59.—As. Min.
sibeuoides, *Rupr. Fl. Cauc.* 180, in textu.—Reg. Caucas.
somalensis, Franch. in Révoil, Comali, 14 = montana.
spathulaefolia, *Fenzl, in Ledeb. Fl. Ross.* i. 290.—Reg. Casp.
spergulaefolia, Boiss. Fl. Orient. i. 559 = Boissieri.
spergulifolia, *Griseb. Spicil. Fl. Rumel.* i. 188.—Europ. austr.
sphaerocephala, *Fenzl, ex Tchihat. Asie Min. Bot.* i. 205.—As. Min.
squarrosa, Tausch, in Flora, xiv. (1831) i. 214 = scutifolia, *Pisch.*
Stevensi, *Rouy. Cat. Hort. Crem.* (1816) 64 ; *ex Link, Enum. Hort. Berol.* i. 417.—Reg. Transcauc
Stevensi, *Fisch. ex Schrank, Pl. Rar. Hort. Monac.* i. 32.—Reg. Caucas.
Stevensi, Hohen. ex Ledeb. Fl. Ross. i. 296 = scutifolia, *Fisch.*
Stewartii, *Thoms. ex Edgew. & Hook. f. in Fl. Brit. Ind.* ii. 116.—Afghan.
stricta, Bunge, in Ledeb. Fl. Alt. ii. 129 = Tunica stricta.
Struthium, Asso, Syn. Stirp. Arag. 52 = hispanica.
Struthium, Fenzl, in Ledeb. Fl. Ross. i. 295 = fastigiata.
Struthium, *Linn. Syst.* ed. X. 1028.—Hispan.
Struthium, Pall. Reise, ii. 676 = Gmelini.
subcapitata, Rupr. Fl. Cauc. 177 = tenuifolia.
Szovitsii, *Fisch. & Mey. ex Fenzl, in Ledeb. Fl. Ross.* i. 289.—Reg. Caucas.
tatarica, Gueldenst. Reise, ii. 205 = paniculata.
tenella, *Poir. Encyc. Suppl.* ii. 874.—Hab.?
tenuifolia, *Biebl. Fl. Taur. Cauc.* i. 319.—Reg. Caucas.
thessalica, Ser. in DC. Prod. i. 354 = Gmelini.
thessala, Nym. Syll. 288 = Tunica thessala.
thymifolia, Sibth. & Sm. Fl. Graec. Prod. i. 282 = hirsuta.
tibetica, Hook. f. & Thoms. ex Edgew. & Hook. f. in Fl. Brit. Ind. i. 217 = sedifolia.
tomentosa, Linn. Cent. Pl. i. 11 = perfoliata.
transylvanica, *Spreng. Syst.* iv. *Cur. Post.* 179.—Transyly.
trichopoda, Boiss. Diagn. Ser. I. i. 19 = linearifolia.
trichotoma, *Wender. Ind. Sem. Hort. Mark.* (1825) ; *ex Linnaea*, xi. (1837) *Litt.* 92.—Reg. Caucas. ; Turkest.
triquetra, Ledeb. Fl. Ross. i. 293 = Gmelini.
tubulosa, *Boiss. Diagn.* Ser. I. i. 11.—As. Min. (Austral. Inguilin.).
uralensis, *Less. in Linnaea*, ix. (1834) 172.—Sibir. ural.
Vaccaria, Sibth. & Sm. Fl. Graec. Prod. i. 279 = Saponaria Vaccaria.
velutina, D. Dietr. Syn. Pl. ii. 1542 = Tunica velutina.
venusta, *Fenzl, Pugill. Pl. Nov. Syr.* 9.—Syria.
violacea, *Fenzl, in Ledeb. Fl. Ross.* i. 291.—Manchur.
virgata, *Boiss. Diagn.* Ser. I. i. 13.—Persia.
viscosa, *Murr. in Comm. Gotting.* (1783) 9. t. 3.—Syria.
Wiedemanni, *Boiss. Fl. Orient.* i. 541.—Galatia.

GYPSOPHYLA, B. Juss. ex Juss. Gen. p. lxviii (1789) = prasc.

GYPSOPHYTUM, Adans, Fam. ii. 256 (1763) = **Arenaria** i, Rupp. (Caryophyll.).

GYPSOPHYTUM, Ehrh. Beitr. iv. 149 (1789) = **Gypsophila**, Linn. (Caryophyll.).

GYPTIS, Cass. in Bull. Soc. Philom. (1818) 189 ; et in Dict. Sc. Nat. xx. 177 (1821) = **Eupatorium**, Linn. (Compos.).
Aeschenoides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 366 = Symphyopappus viscosus.

GYPTIS :—
Commersonii, Cass. in Dict. Sc. Nat. xx. 178 = Eupatorium Bacasanum.
oblongifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras vi. ii. 383 = E. oblongifolium.
prucedanifolia, Sch. Bip. ex Baker, l. c. 315 = E. ormtophyllum.
pinnatifida, Cass. in Dict. Sc. Nat. xx. 178 = E. erodifolium.

GYRANDRA, Griseb. in DC. Prod. ix. 44 (1845) = **Erythraea**, Rich. (Gentian.).
chironioides, Griseb. l. c. = E. macrantha.
chironioides, Griseb. in Seem. Bot. Voy. Herald, 318 = E. madrensis.
speciosa, Benth. Bot. Voy. Sulph. 127. t. 45 = E. tenuifolia.

GYRANDRA, Moq. in DC. Prod. xiii. ii. 40 (1849) = **Tersonia**, Moq. (Phytolac.).

GYRANDRA, Wall. Cat. n. 8020 (1847 ?) = **Daphnidiphyllum**, Blume (Euphorb.).
laurina, Wall. l. c. = D. laurinum.

GYRENIA, Knowles & Westc. ex Loud. Hort. Brit. Suppl. ii. 689 (1839) = **Milla**, Cav. (Liliac.).
biflora, Knowles & Westc. ex Loud. l. c. = M. biflora.

GYRINOPS, Gaertn. Fruct. ii. 276. t. 140 (1791). *THYMELAEACEAE*, Benth. & Hook. f. iii. 200.
LACHNOLRPIS, Miq. Ann. Mus. Bot. Lugd. Bat. i. 132 (1863).
Wallia, *Gaertn. l. c.*—Zeylan.

GYRINOPSIS, Decne. in Ann. Sc. Nat. Sér. II. xix. (1843) 41. t. 1. *THYMELAEACEAE*, Benth. & Hook. f. iii. 200.
Cumingiana, *Decne. l. c.*—Ins. Philipp.

GYROCARPUS, Jacq. Select. Am. 282. t. 178 (1763). *COMBRETACEAE*, Benth. & Hook. f. i. 689.
acuminatus, Meisn. in DC. Prod. xv. i. 248 = americanus.
americanus, *Jacq. Select. Am.* 282.—Cosmop. trop.
anatocus, Willd. Sp. Pl. iv. 982 = americanus.
Yacquini, Gaertn. Fruct. ii. 92 = americanus.
lobatus, Blanco, Fl. Filip. ed. II. 54 = americanus.
pendulus, Blanco, l. c. 55 = Engelhardtia spicata.
rugosus, R. Br. Prod. 405 = americanus.
sphenopterus, R. Br. l. c. = americanus.

GYROCEPHALUM, Reichb. Nom. 67 (1841) = GYNOCEPHALUM, Blume = **Phytocrene**, Wall. (Olacin.).

GYROMIA, Nutt. Gen. Am. i. 238 (1818) = **Medeola**, Linn. (Liliac.).
virginica, Nutt. l. c. = M. virginica.

GYROSTACHIS, Pers. Syn. ii. 511 (1807) = **Spiranthes**, Rich. (1818) (Orchid.).
aestivalis, Dum. Fl. Belg. 124 = S. aestivalis.
amoena, Blume, Orch. Archip. Ind. 129 = S. australis.
australis, Blume, l. c. 128 = S. australis.
autumnalis, Dum. Fl. Belg. 124 = S. autumnalis.
pusilla, Blume, Orch. Archip. Ind. 131 = S. pusilla.

GYROSTELMA, Foum. in Mart. Fl. Bras. vi. tv. 302 (1855). *ASCLEPIADEAE*.
oxypetaloides, *Fourn. l. c.*—Bras.

GYROSTEMON, Desf. in Mém. Mus. Par. vi. (1820) 10. t. 6. *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 36.
CYCLOTHECA, Moq. in DC. Prod. xiii. ii. 37 (1849). *acaciaeformis*, F. Muell. in Linnaea, xxv. (1852) 489 = Codonocarpus cotinifolius.
angustifolius, Schnitzl. Ic. Fam. Nat. iii. t. 208 = Tersonia brevipes.
attenuatus, Hook. Bot. Misc. i. (1830) 243 = Codonocarpus australis.
Blanci, Llanos, Fragm. 74 = Phyllanthus philippinensis.
brachystigma, *F. Muell. ex Benth. Fl. Austral.* v. 146. —Austral.
brevipes, Hook. ex Moq. in DC. Prod. xiii. ii. 39, 40 = Tersonia brevipes.
cotinifolius, Desf. in Mém. Mus. Par. viii. (1822) 116 = Codonocarpus cotinifolia.

GYROSTEMON:—
Cyclotheca, *Benth. Fl. Austral.* v. 146.—Austral.
pungens, Lindl. in Mitch. Three Exped. Austral. ii. 121 = Codonocarpus cotinifolia.
pyramidalis, F. Muell. in Linnæa, xxv. (1852) 688 = Codonocarpus pyramidalis.
ramulosus, *Desf. in Mém. Mus. Par.* vi. (1820) 17 t. 6.—Austral.
ramulosus, Lehm. Pl. Preiss. i. 243 = Tersonia brevipes.
ramulosus, Schlecht. in Linnæa, xx. (1847) 693 = G. Cyclotheca.

GYROSTEPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 11.76 = Chthonocephalus, Steetz (Compos.).
rhizocephalum, Turcz. l. c. 77 = C. pygmæus.

GYROTAENIA, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 174. *URTICACEAE*, Benth. & Hook. f. iii. 384.
cephalantha, Wedd. in DC. Prod. xvi. 1. 195, in syn. = trinervata.
mytilocarpa, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 174.—Cuba.
spicata, *Wedd. in DC. Prod.* xvi. 1. 99.—Jamaica.
trinervata, *Wedd. l. c.* 100.—Ins. S. Doming.

GYROTHECA, Salisb. in Trans. Hort. Soc. i. (1812) 327 = Leachanthes, Ell. (1817) (Hæmodorac.).
tinctoria, Salisb. l. c. = L. tinctoria.

GYTONANTHUS, Raf. in Ann. Gén Sc. Phys. vi. (1820) 88 = **Patrinia**, Juss. (Valerian.).

H

HAAGEA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 125; et in Abh. Acad. Berl. (1855) 108. t. 9 C = **Begonia**, Linn.
dipetala, Klotzsch, ll. cc. 125; 104 = B. malabarica.

HAASIA, Nees, Syst. Laurin. 372 (1836) = **Dehaasia**, Blume (Laurin.).
bornensis, Meissn. in DC. Prod. xv. 1. 61.—Borneo.
caesia, Miq. Fl. Ind. Bat. i. 1. 981 = D. caesia.
cuneata, Nees, Syst. Laurin. 378 = D. cuneata.
cyriopoda, Miq. Fl. Jungh. 177.—Java.
elongata, Nees, Syst. Laurin. 377 = D. elongata.
firma, Miq. Fl. Ind. Bat. i. 1. 928 = D. firma.
incrassata, Nees, Syst. Laurin. 376 = D. microcarpa ?
media, Miq. Fl. Ind. Bat. i. 1. 920, in syn. = Beil. achmiedia Madag.
media, Nees, Syst. Laurin. 375 = D. media.
microcarpa, Nees, l. c. 373 = D. microcarpa.
nitida, Meissn. in DC. Prod. xv. 1. 61 (Alsodaphnidis sp.).—Borneo.
oppositifolia, Meissn. l. c. = Beilschmiedia oppositifolia.
paradoxa, Miq. Fl. Ind. Bat. i. 1. 981 = D. paradoxa.
pauciflora, Miq. l. c. = D. pauciflora.
pedunculariis, Nees, Syst. Laurin. 376 (Alsodaphnidis sp.).—Ind. or.
pedunculariis, Teijsm. & Binn. ex Miq. Fl. Ind. Bat. i. 1. 930 = Beilschmiedia Madag.
spectabilis, Miq. l. c. 930.—Sumatra.
squarrosa, Miq. l. c. 929 = D. squarrosa.
subcaesia, Miq. l. c. Suppl. 361.—Sumatra.
undulata, Teijsm. & Binn. ex Miq. Fl. Ind. Bat. i. 1. 920 = Beilschmiedia undulata.
Wightii, Nees, Syst. Laurin. 676 = Beilschmiedia Wightii.

HAASLUNDIA, Schum. & Thonn. Beskr. Guin. Pl. 15 (1827) = **Hoslundia**, Vahl (Labiat.).

HAASTIA, Hook. f. Handb. N Zeal. Fl. 155 (1864). *COMPOSITAE*, Benth. & Hook. f. ii. 284.
Lognii, *T. Buch. in Trans. N. Z. Inst.* xiv. (1882) 260.—N. Zel.
pulvinaris, *Hook. f. Handb. N. Zeal. Fl.* 156.—N. Zel.
recurva, *Hook. f. l. c.*—N. Zel.
Sinclairii, *Hook. f. l. c.*—N. Zel.

HABENARIA, Willd. Sp. Pl. iv. 44 (1805). *ORCHIDEAE*, Benth. & Hook. f. iii. 624.
Ayr, Lindl. Gen. et Sp. Orch. 326 (1835).
Barlaea, Reichb. f. in Linnæa, xli. (1877) 54.
Benthamia, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1829) 87.
Bicchia, Parl. Fl. Ital. iii. 396 (1858).
Bilabrella, Lindl. Bot. Reg. sub t. 1701 (1834).
Blephariglottis, Rafin. Fl. Tellur. ii. 38 (1836).
Bonatea, Willd. Sp. Pl. iv. 43 (1805).
Centrochilus, J. C. Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 485. t. 18.
Chorisiglossum, Hartm. Handb. Scand. Fl. ed. I. 329 (1820).
Conopidium, Wallr. in Linnæa, xiv. (1840) 147.
Derosmeria, Reichb. f. Poll. Orch. Comm. 29 (1852).
Digomphotis, Rafin. Fl. Tellur. ii. 37 (1836).
Diplectraden, Rafin. l. c. 60 (1836).
Dissorhynchium, J. C. Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 484. t. 18.
Ectatiicus, S. F. Gray, Nat. Arr. Brt. Pl. ii. 198 (1821).
Galearis, Rafin. Fl. Tellur. ii. 39 (1836).
Gennaria, Parl. Fl. Ital. iii. 404 (1858).
Gymnadenia, R. Br. in Ait. Hort. Kew. ed. II. v. 191 (1813).
Leucorchis, E. H. F. Mey. Preuss. Pfl. Gatt. 50 (1839).
Lindblomia, Fries, in Lindbl. Bot. Notiser (1843) 134.
Lysias, Salisb. in Trans. Hort. Soc. i. (1812) 288.
Macrocentrum, Phil. Sert. Mendoc. ii. 42 (1871).
Mycosa, Blume, Bijdr. 403. t. 1 (1825).
Myosticma, Blume, Mus. Bot. Lugd. Bat. ii. 189 (1856).
Montolivara, Reichb. f. Otia Bot. Hamb. 107 (1881).
Nemisanthes, Rafin. Fl. Tellur. ii. 61 (1836).
Neotinea, Reichb. f. Poll. Orch. Comm. 29 (1852).
Nigritella, Rich. in Mém. Mus. Par. iv. (1818) 48.
Pectelis, Rafin. Fl. Tellur. ii. 37 (1836).
Peristylus, Blume, Bijdr. 404. t. 30 (1825).
Perularia, Lindl. Bot. Reg. sub t. 1701 (1834).
Platanthera, Rich. in Mém. Mus. Par. iv. (1818) 48.
Ponerorchis, Reichb. f. in Linnæa, xxv. (1852) 227.
Rorippacrocharis, Reichb. f. Otia Bot. Hamb. 104 (1881).
Sieberia, Spreng. Anleit. ii. 332 (1817).
Synmeria, Nimmo, in J. Grah. Cat. Pl. Bomb. (pag. uit. absque numero) (1839).
Tinea, Bivona, in Giorn. Sc. Sicil. (1838) 149.
Tulotis, Rafin. Fl. Tellur. ii. 37 (1836).
abortiva, *Lindl. Gen. et Sp. Orch.* 326.—Peruv.
achnanthe, *Reichb. f. in Linnæa*, xxii. (1849) 812.—Am. trop.
aculifera, *Wall. ex Lindl. Gen. et Sp. Orch.* 325.—Ind. or. ; China.
acuminata, *Thw. ex Trim. Cat. Pl. Cyl.* 91.—Ceylan.
aestivalis, *Rodrig. Orch. Nov.* ii. 258.—Bras.
affinis, D. Don, Prod. Fl. Nep. 25 = goodyeroides.
affinis, *Wight, ic.* t. 1707.—Ind. or.
alata, *Hook. Exot. Fl.* t. 169.—Ind. occ.
albida, *R. Br. in Ait. Hort. Kew.* ed. II. v. 198.—Am. bor.
Allemanii, *Rodrig. Orch. Nov.* ii. 254.—Bras.
alta, *Rafi. in Journ. Linn. Soc.* xxi. (1885) 509.—Madag.
Amphorchis, Spreng. Syst. iii. 689 = Cynorchis squamosa.
Anaphysema, *Reichb. f. in Flora*, l. (1867) 101.—Afr. trop.
anaplectron, *Reichb. f. & Warm. in Otia Bot. Hamb.* 81.—Bras.
angulosa, *Rodrig. Orch. Nov.* ii. 258.—Bras.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* i. 330.—N. Granat.
anisoptera, *Reichb. f. Otia Bot. Hamb.* 101.—Afr. trop.
antennifera, *A. Rich. in Ann. Sc. Nat.* Sér. II. xiv. (1840) 268.—Abyssin.
aphylla, *R. Br. Prod.* 312, in adnot.—Arab.
aphylla, *Rodrig. Orch. Nov.* i. 158.—Bras.
arachnoides, *Thou. Fl. Ile Afr. Austr.* t. 18.—Madag.
aranefera, Rodrig. Orch. Nov. i. 159 = anaplectron.
aranifera, *Lindl. Gen. et Sp. Orch.* 313.—Reg. Argent.

HABENARIA:—
arenaria, *Lindl. Gen. et Sp. Orch.* 317.—Afr. austr.
aristata, *Trim. Cat. Pl. Cyl.* 91.—Zeylan.
aristulifera, *Reichb. f. in Bonplandia*, iii. (1855) 213 —Ins. Philipp.
armata, *Reichb. f. l. c. ii.* (1854) 10.—N. Granat.
armatissima, *Reichb. f. Otia Bot. Hamb.* 98.—Afr. trop.
arnhemica, F. Muell. ex Benth. Fl. Austral. vi. 395 = graminea.
arvensis, Rodrig. Orch. Nov. i. 163 = hexaptera.
attenuata, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 221.—Afr. trop.
autumnalis, *Poepp. & Endl. Nov. Gen. et Sp.* i. 44.—Peruv.
barbata, *Wight, in Wall. Cat.* n. 7084.—Ind. or.
biconvis, *Lindl. Gen. et Sp. Orch.* 309.—Cuba.
bidentata, Poepp. ex Steud. Nom. ed. II. i. 216 = alata.
bifora, *Rodrig. Orch. Nov.* i. 151.—Bras.
bifolia, *R. Br. in Ait. Hort. Kew.* ed. II. v. 193.—Europ.
bimaculata, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 506.—Madag.
Blephariglottis, *Hook. Exot. Fl.* t. 87.—Am. bor.
boliviana, *Reichb. f. in Bonplandia*, iii. (1855) 213.—Bolivia.
Boltoni, *Harv. Thes. Cap.* t. 55.—Afr. austr.
Bonatea, *Reichb. f. Otia Bot. Hamb.* 101, in obs.—Afr. austr.
Bongeanium, *Reichb. f. l. c.* 58.—Afr. trop.
borealis, Cham. in Linnæa, iii. (1828) 28 = dilatata.
brachycerea, *Phil. in Anal. Univ. Chil.* (1861) t. 59.—Chili.
brachyceras, Spreng. Syst. iii. 692 = seq.
brachyceratita, *Willd. Sp. Pl.* iv. 44.—Jamaic.
brachyphylla, *Astch. in Journ. Linn. Soc.* xix. (1882) 186.—Afghan.
bracteata, R. Br. in Ait. Hort. Kew. ed. II. v. 192 = viridis.
braciosa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 292.—Abyssin.
bractescens, *Lindl. Gen. et Sp. Orch.* 308.—Reg. Argent.
brevidens, *Lindl. l. c.* 314.—Bras.
brevifolia, *Greene, in Coult. Bot. Gaz.* vi. (1881) 218.—Am. bor.
breviloba, *Trim. Cat. Pl. Cyl.* 91.—Zeylan.
candida, Dals. in Hook. Kew Journ. ii. (1850) 262 = subpubens.
carasjensis, *Dals. l. c.*—Ind. or.
carinata, *Spun. in Linnæa*, xv. (1841) 478.—Ins. Timor.
cassidea, *Reichb. f. in Walp. Ann.* i. 797.—Afr. austr.
cataphysema, *Reichb. f. in Flora*, l. (1867) 101.—Afr. trop.
Cephalotes, *Lindl. Gen. et Sp. Orch.* 322.—Ind. or.
ceratopetala, *A. Rich. in Ann. Sc. Nat.* Sér. II. xiv. (1840) 267.—Abyssin.
chirensis, *Reichb. f. Otia Bot. Hamb.* 99.—Afr. trop.
chlorantha, Bab. in Trans. Linn. Soc. xvii. (1836) 462 = bifolia.
chlorantha, *Spreng. Syst.* iii. 691.—Ins. Mascar.
chlorantha, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 86 = viridiflora.
chlorina, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 140.—Burma.
chloroleuca, Ridl. in Journ. Bot. xxiii. (1885) 219 = bifolia.
chlorotica, *Reichb. f. in Flora*, xlviii. (1865) 178.—Afr. trop.
Chorisiana, *Cham. in Linnæa*, iii. (1828) 31.—Unalaschk.
Chusua, Benth. in Journ. Linn. Soc. xviii. (1881) 355 = Orchis Chusua.
ciliaris, *R. Br. in Ait. Hort. Kew.* ed. II. v. 194.—Am. bor.
ciliosa, *Lindl. in Ann. Nat. Hist. iv.* (1840) 314.—Afr. austr.
cirrhata, *Reichb. f. in Flora*, xlviii. (1865) 180.—Madag.
citrina, *Thou. Orch. Isles Afr. Austr.* t. 16.—Ins. Borbon.
clavata, *Reichb. f. in Flora*, xlviii. (1865) 180.—Afr. austr.
clavellata, Spreng. Syst. iii. 689 = tridentata.
clypeata, *Lindl. Gen. et Sp. Orch.* 311.—Mexic.
commelinifolia, *Wall. ex Lindl. l. c.* 326.—Ind. or.
concopsea, *Benth. in Journ. Linn. Soc.* xviii. (1880) 354.—Europ. ; As. bor.

HABENARIA:—
conopsea, *Reichb. f. in Bonplandia*, ii. (1854) 10.—N. Granat.
Cooperi, *S. Wats. in Proc. Am. Acad.* xii. (1877) 276.—Am. bor. occ.
cordata, *R. Br. Prod.* 312.—Europ.; Am. bor.
cornuta, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 208.—Afr. austr.
cornuta, *Span. in Linnaea*, xv. (1841) 478.—Ins. Timor.
corydophora, *Reichb. f.* l. c. xli. (1877) 80.—Peruv.
corymbosa, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 141.—Burma.
crassicornis, *Lindl. Gen. et Sp. Orch.* 311.—Mexic.
crassifolia, *A. Rich. in Ann. Sc. Nat.* Sér. II. xv. (1841) 72.—Ind. or.
cruciformis, *Lindl. Gen. et Sp. Orch.* 323.—Ind. or.
cristata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 194.—Am. bor.
crocea, *Rafin. Atl. Journ.* 119.—Am. bor.
crocea, *Schweinf. ex Reichb. f. in Otia Bot. Hamb.* 57.—Afr. trop.
crucifera, *Reichb. f. & Warm. l. c.* 80.—Bras.
cryptophila, *Rodrig. Orch. Nov.* i. 154.—Bras.
cryptostyla, *Reichb. f. Otia Bot. Hamb.* 50.—Ins. Taiti.
cubitalis, *R. Br. Prod.* 313, in adnot.—Ind. or.
cucullata, *Hooff. Cat. Pl. Koursk.* 56; *ex Ledeb. Fl. Ross.* iv. 66.—Europ.
culicina, *Reichb. f. & Warm. in Otia Bot. Hamb.* 79.—Bras.
cultelliformis, *Rodrig. Orch. Nov.* i. 155.—Bras.
cultrata, *A. Rich. Tent. Fl. Abyss.* ii. 297.—Abyssin.
curvilabra, Rodrig. Orch. Nov. i. 162 = leptoceras.
cyclochila, *Franch. & Sav. Enum. Pl. Jap.* ii. 516.—Japon.
debilis, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 221.—Afr. trop.
decaptera, *Reichb. f. in Flora*, lxv. (1882) 531.—Afr. trop.
decipiens, *Wight, Ic.* v. i. 14.—Ind. or.
decorata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 294.—Abyssin.
dense, *Wall. ex Lindl. Gen. et Sp. Orch.* 326.—Reg. Himal.
densiflora, *Reichb. f. in Walp. Ann.* i. 797.—Afr. austr.
dentiflora, Schur, Enum. Pl. Transs. 645 = transsilvanica.
denticulata, *Reichb. f. in Linnaea*, xix. (1847) 876.—Ind. or.
depauperata, *Kraenzl. in Bremen, Abh.* vii. (1882) 269.—Madag.
dichopetala, *Thw. Enum. Pl. Zeyl.* 309.—Zeylan.
difusa, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 28.—Guatemala.
digitata, *Lindl. Gen. et Sp. Orch.* 307.—Ind. or.
dilatata, *A. Gray, Man. Bot. N. U. St.* ed. V. 500.—Am. bor.
dilatata, Hook. Exot. Fl. t. 95 = hyperborea.
dilatata, S. Wats. in Bot. King's Exped. 340 = leucostachys.
diphylla, *Dalz. in Hook. Kew Journ.* ii. (1850) 262.—Ind. or.
disoides, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 511.—Madag.
distans, *Griseb. Cat. Pl. Cub.* 270.—Cuba.
distantiflora, *A. Rich. Tent. Fl. Abyss.* i. 291.—Abyssin.
dives, *Reichb. f. in Flora*, i. (1867) 117.—Afr. austr.
dolichocaccras, Rodrig. Orch. Nov. i. 163 = bractescens.
dolichostachya, *Thw. Enum. Pl. Zeyl.* 309.—Zeylan.
Dregeana, *Lindl. in Ann. Nat. Hist.* iv. (1840) 314.—Afr. austr.
elegans, *Boland. Cat. Pl. St. Franc.* 29; *ex S. Wats. in Bot. Calif.* ii. 182.—Calif.
Elliottii, Beck, Bot. N. & M. St. 348 = integra.
elliptica, *Wight, Ic.* t. 1706.—Ind. or.
elongata, *R. Br. Prod.* 313.—Austral.
endotrya, *Miq. in Journ. Bot. Néerl.* i. (1861) 92.—China.
enatifolia, Lindl. Gen. et Sp. Orch. 321 = pectinata.
entomanthe, *Lindl. l. c.* 311.—Mexic.
epipactidea, *Reichb. f. in Flora*, i. (1867) 100.—Afr. trop.
epiphylla, *Reichb. f. Otia Bot. Hamb.* 79.—Bras.
ericaeflora, *Reichb. f. in Linnaea*, xxii. (1849) 813.—Am. trop.

HABENARIA:—
exaltata, *Rodrig. Orch. Nov.* i. 156.—Bras.
falciformis, Bolus, in Journ. Linn. Soc. xix. (1882) 340 = tetraspeda.
falcigera, *Reichb. f. in Linnaea*, xli. (1877) 100.—Ins. Philipp.
filiformis, *Lindl. Gen. et Sp. Orch.* 318.—Afr. trop.
filiformis, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 504.—Madag.
fimbriata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 193.—Am. bor.
fimbriata, *Wight, Ic.* t. 1712.—Ind. or.
fusa, *R. Br. Prod.* 312, in adnot.—Am. bor.
fusa, A. Rich. = herbiola.
flexuosa, *Lindl. Gen. et Sp. Orch.* 311.—Mexic.
flexuosa, *Reichb. f. in Flora*, xlviii. (1865) 180.—Bras.
floribunda, *Lindl. Gen. et Sp. Orch.* 316.—Peruv.
foetida, S. Wats. in Bot. King's Exped. 341 = Herminium congestum.
foliosa, *Reichb. f. in Flora*, xlviii. (1865) 180.—Afr. austr.
foliosa, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 71 = digitata.
formosana, Bab. in Trans. Linn. Soc. xvii. (1836) 463 = bifolia.
furcifera, *Lindl. Gen. et Sp. Orch.* 319.—Reg. Himal.
gabonensis, *Reichb. f. in Bot. Zeit.* x. (1852) 934.—Afr. trop.
galeandra, *Benth. Fl. Hongk.* 363.—Ind. or.; China.
Garberi, *Porter, in Coult. Bot. Gaz.* v. (1880) 135.—Am. bor.
geniculata, *D. Don, Prod. Fl. Nep.* 25.—Reg. Himal.; Burma.
Gerardiana, Lindl. in Wall. Cat. n. 7081 = peutinata.
Germaini, *Phil. in Linnaea*, xxix. (1857–58) 46.—Chili.
Gervasii, Reichb. f. Otia Bot Hamb. 97 = tridens.
Ghiesbreghtiana, *Hemsl. Biol. Centr. Am. Bot.* iii. 805.—Mexic.
gigantea, D. Don, Prod. Fl. Nep. 24 = Susannae.
glabra, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 71 = Herneana.
glaucophylla, Rodrig. Orch. Nov. i. 160 = epiphylla.
Gnoma, *Rodrig. l. c.* ii. 250.—Bras.
Godefroyi, *Reichb. f. Otia Bot. Hamb.* 38.—Camboj. goodyeroides, Hook. Bot. Mag. t. 3397 = Wightii.
Gourlieana, *Gill. ex Lindl. Gen. et Sp. Orch.* 309.—Chili.
gracilis, *Colebr. in Hook. Exot. Fl.* t. 135.—Ind. or.
gracilis, *Lindl. Gen. et Sp. Orch.* 312.—Bras.
gracilis, Reichb. f. in Flora, xlviii. (1865) 180 = longecalcarata.
gracilis, *S. Wats. in Proc. Am. Acad.* xii. (1877) 276.—Am. bor.
gracilicaspe, Rodrig. Orch. Nov. i. 155 = brevidens.
graminea, *Lindl. Gen. et Sp. Orch.* 318.—Ind. or.
graminea, Naves, in Blanco, Fl. Philipp. ed. III. Nov. App. 251 = muricata, *Vidal.*
graminea, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 78 = viridiflora.
graminea, *Spreng. Syst.* iii. 690.—Madag.; Ins. Mascar.
grandiflora, *Lindl. in Wall. Cat.* n. 7083.—Ind. or.
grandiflora, *Torr. ex Beck, Bot. N. & M. St.* 349.—Am. bor.
grandis, *Benth. ex Ridl. in H. O. Forbes, Wand. Nat.* 519.—Ins. Timor.
Guilelmini, *Reichb f. in Linnaea*, xix. (1847) 875.—Bras.
Gutngangae, *Reichb. f. in Flora*, xlviii. (1865) 179.—Afr. trop.
hamata, *Rodrig. Orch. Nov.* i. 162.—Bras.
hamigera, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 380. t. 20 = furcifera.
helleborina, *Nichols. Dict. Gard.* ii. 307.—Afr. trop.
heloides, *Reichb. f. in Linnaea*, xxii. (1849) 813.—Am. trop.
Henschenians, *Rodrig. Orch. Nov.* i. 157.—Bras.
heptadactyla, *Reichb. f. in Linnaea*, xxii. (1849) 812.—Guian.
herbiola, *R. Br. in Ait. Hort. Kew.* ed. II. v. 193.—Am. bor.
hexaptera, *Lindl. Gen. et Sp. Orch.* 316.—Bras.
Heyneana, *Lindl. l. c.* 320.—Ind. or.
Hildebrandtii, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 503.—Madag.
Hilsenbergii, *Ridl. l. c.* 509.—Madag.

HABENARIA:—
hircina, *Reichb. f. in Flora*, i. (1867) 100.—Afr. trop.
hirtella, *Spreng. Syst.* iii. 690.—Jamaic.
hispida, Sprung. f. Tent. Suppl. 27 = Holothrix parviflora.
Hookeri, Lindl. Gen. et Sp. Orch. 286 = secq.
Hookeriana, *Torr. ex A. Gray, in Ann. Lyc. N. York*, iii. (1836) 229.—Am. bor.
huillensis, *Reichb. f. in Flora*, xlviii. (1865) 179.—Afr. trop.
Humblotii, *Reichb. f. l. c.* lxviii. (1885) 585.—Ins. Comor.
humilior, *Reichb. f. Otia Bot. Hamb.* 100.—Abyssin.
hurumensis, Spreng. Syst. iii. 688 = hyperborea.
hydrophila, *Rodrig. Orch. Nov.* i. 158.—Bras.
hyperborea, *R. Br. in Ait. Hort. Kew.* ed. II. v. 198.—Europ.; Am. bor.
ichneumones, *Lindl. Gen. et Sp. Orch.* 313.—Afr. trop.
imbricata, *Lindl. l. c.*—Bras.
imerinensis, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 505.—Madag.
incarnata, *Lyall, ex Lindl. Gen. et Sp. Orch.* 328.—Madag.
incisa, *Spreng. Syst.* iii. 692.—Am. bor.
intacta, *Benth. in Journ Linn. Soc.* xviii. (1881) 354.—Europ.; Afr. bor.
integra, *Spreng. Syst.* iii. 689.—Am. bor.
intermedia, *D. Don, Prod. Fl. Nep.* 24.—Reg. Himal.
japonica, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 410.—Japon.
Jerdoniana, Wight, Ic. t. 1715 = diphylla.
johannensis, *Rodrig. Orch. Nov.* ii. 261.—Bras.
josephensis, *Rodrig. l. c.* 257.—Bras.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 207.—Japon.
Kilimanjari, *Reichb. f. Otia Bot. Hamb.* 119.—Afr. trop.
lacera, *R. Br. Prod.* 312.—Am. bor.
lacertifera, *Benth. Fl. Hongk.* 362.—Ind. or.; China.
lacinnata, Dalz. in Hook. Kew journ. ii. (1850) 261 = digitata.
lactiflora, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 28.—Mexic.
laevigata, *Lindl. in Ann. Nat. Hist.* iv. (1840) 315.—Afr. austr.
lanciflora, *A. Rich. in Mém. Soc. Hist. Nat. Par.* iv. (1828) 19.—Ins. Maurit.
latifolia, *H. B. & K. Nov. Gen. et Sp.* i. 331.—N. Granat.
latifolia, Lindl. Gen. et Sp. Orch. 321 = Lindleyana, *Steud.*
laxiflora, *Poepp. & Endl. Nov. Gen. et Sp.* i. 46.—Peruv.
Leprieuri, *Reichb. f. in Linnaea*, xix. (1847) 876.—Guian.
leptoceras, *Hook. Bot. Mag.* t. 3736.—Bras.
leptoloba, *Benth. Fl. Hongk.* 362.—China.
leucophaea, *A. Gray, Man. Bot. N. U. St.* ed. V. 502.—Am. bor.
leucosantha, *Rodrig. Orch. Nov.* i. 151.—Bras.
leucostachys, *S. Wats. in Bot. Calif.* ii. 184.—Am. bor. occ.
limosa, *Hemsl. Biol. Centr. Am. Bot.* iii. 805.—Mexic.
Lindeni, *Lindl. Orch. Lindex.* 25.—Venezuela.
Lindleyana, *Steud. Nom.* ed. II. i. 717.—Malaya.
Lindleyana, *Wight, Ic.* t. 923.—Travancor.
linearifolia, Maxim. Prim. Fl. Amur. 269 = sagittifera.
Linguella, Lindl. Gen. et Sp. Orch. 325 = scutaria.
lingulicuris, *Reichb. f. in Bonplandia*, ii. (1854) 10.—N. Granat.
linifolia, *Presl, Rel. Haenk.* i. 91.—Peruv.
Lobbii, *Reichb. f. in Linnaea*, xli. (1877) 80.—Ind. or.
longebracteata, *Hochst. ex Seub. Fl. Aaor.* 25.—Ins. Azor.
longecalcarata, *A. Rich. in Ann. Sc. Nat.* Sér. III. xv. (1841) 71. t. 5.—Ind. or.
longicauda, *Hook. Bot. Mag.* t. 2967.—Guiana.
longicorniculata, J. Grah. Cat. Pl. Bomb. 202 = longecalcarata.
longicornis, Lindl. Gen. et Sp. Orch. 322, sphalm. = secq.
longicornu, *Lindl. in Wall. Cat.* n. 7027.—Ind. or.
longifolia, *Buch.-Ham. ex Lindl. Gen. et Sp. Orch.* 324.—Reg. Himal.

HABENARIA:—
lacilla, *Wall. ex Lindl. Gen. et Sp. Orch.* 319.—Burma.
latea, Benth. in journ. Linn. Soc. xviii. (1881) 854 = Perrotettiana.
maculata, *Reichb. f. in Flora*, xlviii. (1865) 180.—Guiana.
macrandra, *Lindl. in Journ. Linn. Soc.* vi. (1862) 139.—Afr. trop.
macrantha, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 294.—Abyssin.
macroceras, Spreng. Syst. iii. 692 = seq.
macrocerastilis, *Willd. Sp. Pl.* iv. 44.—Costa Rica.
macrophylla, *Goldie, in Brewst. Edinb. Phil. Journ.* vi. (1822) 381.—Am. bor.
macrostachya, *Lindl. Gen. et Sp. Orch.* 307.—Ind. or.
maculosa, *Lindl. l. c.* 309.—Am. trop.
malacophylla, *Reichb. f. Otia Bot. Hamb.* 97.—Afr. austr.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 222. —Afr. trop.
margaritacea, *Linden, in Illustr. Hortic.* xxix. (1882) 52, nomen.—N. Granat.
marginata, *Colebr. in Hook. Exot. Fl.* t. 156.—Ind. or.
martialis, *Reichb. f. Otia Bot. Hamb.* 99.—Afr. trop.
mascarenensis, *Spreng. Syst.* iii. 690.—Ins. Mascar.
maxillaris. *Lindl. in Hook. Journ. Bot.* i. (1834) 5.—Am. trop.
Mechowii, *Reichb. f. in Flora*, lxv. (1882) 532.—Afr. trop.
megaceras, Rodrig. Orch. Nov. i. 161 = obtusa.
Melvillii, *Ridl. in Journ. Bot.* xxiii. (1885) 170.—Bras.
membranacea, *Lindl. Gen. et Sp. Orch.* 313.—Afr. trop.
mesochasyla, *Griseb. Fl. Brit. W. Ind.* 544.—Ind. occ.
mexicana, *Linden, in Illustr. Hortic.* xxix. (1882) 52, nomen.—Mexic.
Michaeli, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 281.—Am. bor. occ.
Michauxii, *Nutt. Gen. Am.* ii. 189.—Am. bor.
microantha, *Hochst. ex Seub. Fl. Asor.* 25.—Ins. Azor.
microchila, Reichb. f. in Flora, xlviii. (1865) 180 = arenaria.
microchila, Wall. ex Lindl. Gen. et Sp. Orch. 302 = cubitalis.
microcerae, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 221.—Afr. trop.
microstylina, *Reichb. f. Xenia Orch.* iii. 17.—Bolivia.
Miersiana, *Champ. ex Benth. in Hook. Kew Journ.* vii. (1855) 87.—Hongkong.
Milnei, *Reichb. f. Otia Bot. Hamb.* 96.—Afr. austr.
minutiflora, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 508.—Madag.
misera, *Ridl. l. c.*—Madag.
modesta, *Dals. in Hook. Kew Journ.* ii. (1850) 262. —Ind. or.
modestissima, *Reichb. f. in Linnæa*, xxii. (1849) 811. —Am. trop.
montana, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 73 = longicorna.
montana, *Rodrig. Orch. Nov.* i. 154.—Bras.
montana, Wight, Ic. t. 1714, partim = decipiens.
montevidensis, *Lindl. Gen. et Sp. Orch.* 314.—Urug.
montevidensis, *Spreng. Syst.* iii. 692.—Urug.
muricata, *Rodrig. Orch. Nov.* ii. 255.—Bras.
muricata, *Vidal, Phan. Cuming. Philipp.* 80, 82, 88, 151.—Ins. Philipp.
mutica, *Spun. in Linnæa*, xv. (1841) 477.—Ins. Timor.
mystacina, *Lindl. Gen. et Sp. Orch.* 312.—Bras.
nasuta, *Reichb. f. & Warm. in Otia Bot. Hamb.* 80.—Bras.
natalensis, *Reichb. f. l. c.* 97.—Afr. austr.
nemorosa, *Rodrig. Orch. Nov.* i. 160.—Bras.
neuropetala, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 207. —Japon.
nigra, *R. Br. in Ait. Hort. Kew.* ed. II. v. 192.—Knrop.
niven, *Spreng. Syst.* iii. 689.—Am. bor.
novemfida, *Lindl. in Benth. Pl. Hartw.* 94.—Guatemala.
Novo-Ebudarum, *F. Muell. Phyt. New Hebr.* 22.—N. Hebr.
nuda, *Lindl. Gen. et Sp. Orch.* 312.—Bras.
nutans, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 507.—Madag.
nyikana, *Reichb. f. Otia Bot. Hamb.* 100.—Afr. trop.

HABENARIA:—
obsoleta, *Spreng. Syst.* iii. 689.—Am. bor.
obtusa, *Lindl. Gen. et Sp. Orch.* 315.—Bras.
obtusata, *Richards. in Frankl. Narr. 1st Journ.* 750. —Am. bor.
occultans, *Welw. ex Reichb. f. in Flora*, xlviii. (1865) 178.—Afr. trop.
ochroleuca, *R. Br. Prod.* 313.—Austral.
odontopetala, *Reichb. f. in Linnæa*, xviii. (1844) 407. —Mexic.
odoratissima, *Franch. Fl. Cher-et-Loire*, 378.—Europ.
Oerstedii, *Reichb. f. in Bonplandia*, iii. (1855) 213.—Nicaragua.
orangana, *Reichb. f. Otia Bot. Hamb.* 101.—Afr. austr.
orbiculata, Hook. Exot. Fl. t. 145 = Hookeriana.
ornithiodes, *Rodrig. Orch. Nov.* i. 162.—Bras.
ornithopoda, Reichb. f. in Linnæa, xx. (1847) 696 = laevigata.
ovalifolia, *Wight, Ic.* t. 1708.—Ind. or.
Palvaeana, *Reichb. f. Otia Orch.* iii. 17.—Bolivia.
paludosa, *Lindl. in Journ. Linn. Soc.* vi. (1862) 139. —Afr. occ.
paludosa, Reichb. f. in Flora, xlviii. (1865) 178 = Protesarum.
paludosa, *Rodrig. Orch. Nov.* i. 153.—Bras.
papillosa, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 504. —Madag.
paranaensis, *Rodrig. Orch. Nov.* ii. 249.—Bras.
parviflora, *Lindl. Gen. et Sp. Orch.* 317.—Peruv.
parviflora, *Lindl. l. c.* 314.—Ins. Philipp.
pauciflora, *Reichb. f. in Bonplandia*, ii. (1854) 10, sub n. 3.—Bras.
pauciflora, *Rodrig. Orch. Nov.* i. 157.—Bras.
pauciflora, *Lindl. Gen. et Sp. Orch.* 317.—Chili.
pectinata, *D. Don, Prod. Fl. Nep.* 24.—Reg. Himal.
pedicellaris, *Reichb. f. Otia Bot. Hamb.* 100.—Abyss.
pedicellata, *S. Watts. in Proc. Am. Acad.* xii. (1877) 276.—Am. bor. occ.
pelorioides, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 128.—Burma.
peltastes, *Reichb. f. Otia Bot. Hamb.* 99.—Afr. trop.
pentadactyla, *Lindl. Gen. et Sp. Orch.* 307.—Reg. Argent.
perpusilla, *A. Gray, Bot. N. U. St.* ed. V. 503.—Am. bor.
perbella, *Reichb. f. Otia Bot. Hamb.* 98.—Abyss.
peristylioides, *A. Rich. in Ann. Sc. Nat.* Sér. II. xiv. (1841) 276.—Abyss.
peristyloides, Wight, Ic. v. 18. t. 1702 = stenostachya.
Perrottetiana, *A. Rich. in Ann. Sc. Nat.* Sér. II. xv. (1841) 71.—Ind. or.
petaloides, *Lindl. Gen. et Sp. Orch.* 316.—Bras.
petromedusa, *Webb, in Hook. Niger Fl.* 180.—Ins. Cap. Virid.
phycnoclectra, *Reichb. f. in Linnæa*, xlii. (1877) 17.—N. Hebrid.
picnostachia, *Rodrig. Orch. Nov.* ii. 250.—Bras.
Pinguicula, *Benth. in Journ. Linn. Soc.* xviii. (1881) 355.—China.
plantaginea, *Lindl. Gen. et Sp. Orch.* 323.—Ind. or.
Plataniflora, *Reichb. f. in Linnæa*, xli. (1877) 55.—Urug.
platyanthera, *Reichb. f. l. c.* xxii. (1849) 860.—Abyss.
platyphylla, *Spreng. Syst.* iii. 690.—Ind. or.
plectostachya, *Reichb. f. & S. Moore, in Journ. Bot.* xvi. (1878) 186.—Afr. trop.
pleistadenia, *Reichb. f. in Journ. Linn. Soc.* xxi. (1885) 406.—Afr. trop.
polypodantha, *Reichb. f. Otia Bot. Hamb.* 97.—Afr. austr.
posaoensis, *Reichb. f. in Bonplandia*, iii. (1855) 213.—Ins. Philipp.
praealta, *Spreng. Syst.* iii. 691.—Ins. Borbon.
pratensis, *Reichb. f. in Linnæa*, xxii. (1849) 813.—Bras.
pretensa, *Lindl. Gen. et Sp. Orch.* 316.—Afr. trop.
promensis, Lindl. in Wall. Cat. n. 7083 = marginata.
propinquior, Reichb. f. Beitz. Syst. Pfl. 58 = xantha.
Protesarum, *Reichb. f. Otia Bot. Hamb.* 97.—Bras.
Pseudocytites, *Reichb. f. & Warm. l. c.* 79.—Bras.
psychodes, Spreng. Syst. iii. 698 = fusa ?
psycodes, A. Gray, Man. Bot. N. U. St. ed. V. 502 = fusa ?
pterocarpa, *Thw. Enum. Pl. Zeyl.* 309.—Zeylan.
pubescens, *Lindl. Gen. et Sp. Orch.* 323.—Reg. Himal.
pumilio, *Porpp. Progm. Syn.* 21.—Chili.

HABENARIA:—
purpurea, *Thou. Orch. Isles Austr. Afr.* t. 17.—Madag.
pusilla, *Reichb. f. Otia Bot. Hamb.* 98.—Cochinch.
pyramidalis, Lindl. in Ann. & Mag. Nat. Hist. xv. (1845) 386 = strictissima.
quadrata, *Lindl. Gen. et Sp. Orch.* 316.—Bras.
Quartiniana, *A. Rich. in Ann. Sc. Nat.* Sér. II. xiv. (1840) 271.—Abyss.
racemosa, *Rafin. Ann. Nat.* 15.—Am. bor.
radiata, *Spreng. Syst.* iii. 693.—Japon.
raribra, *A. Rich. in Ann. Sc. Nat.* Sér. II. xv. (1841) 70.—Ind. or.
reflexa, *Blume, Bijdr.* 408.—Java.
Reichenbachiana, *Rodrig. Orch. Nov.* i. 159.—Bras.
repens, *Nutt. Gen. Am.* ii. 190.—Am. bor.
replicata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 296. —Abyss.
replicata, *A. Rich. Fl. Cub. Funerog.* ii. 250.—Cuba.
retusa, *Riddell, Suppl. Cat. Ohio Pl.* 17.—Am. bor.
retusa, *Rodrig. Orch. Nov.* ii. 253.—Bras.
rhodochelia, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1856) 248.—China.
rhynchocarpa, *Trim. Cat. Pl. Ceyl.* 91.—Zeylan.
Richardiana, *Wight, Ic.* t. 1713.—Ind. or.
robusta, N. E. Br. in Gard. Chron. (1885) ii. 807 = H. Bonates.
robusta, *Welw. ex Reichb. f. in Flora*, l. (1867) 102.—Afr. trop.
rodeiensis, *Rodrig. Orch. Nov.* ii. 256.—Bras.
rostellifera, *Reichb. f. Otia Bot. Hamb.* 94.—Cochinch.
rostrata, *Wall. ex Lindl. Gen. et Sp. Orch.* 325.—Burma.
rotundifolia, Lindl. Gen. et. Sp. Orch. 306 = grandiflora, Lindl.
rotundifolia, *Richards. in Frankl. Narr. 1st Journ.* 750; *Spreng. Syst.* iii. 692.—Am. bor.
Rumphii, *Lindl. Gen. et Sp. Orch.* 320.—Ins. Amboina.
rupestris, *Poepp. Endl. Nov. Gen. et Sp.* i. 45.—Peruv.
rupicola, Rodrig. in Rev. Engenh. iii. (1881) 144 = montana.
Rutenbergiana, *Kraenzl. in Bremen, Abh.* vii. (1882) 258.—Madag.
sagittifera, *Reichb. f. in Bot. Zeit.* iii. (1845) 884.—Japon ; Manchur.
salaccensis, *Blume, Bijdr.* 408. t. 18.—Malaya.
santensis, *Rodrig. Orch. Nov.* ii. 253.—Bras.
Sartor, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 662.—Bras.
Saundersiae, *Harv. Thes. Cap.* ii. 29.—Afr. austr.
scaptrodes, *Reichb. f. in Bonplandia*, iv. (1856) 210. —Hab. ?
Schimperiana, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 295.—Abyss.
Schizochærophana, Cham. in Linnæa, iii. (1838) 29 = Herminium congestum.
Schizochilus, J. Grah. Cat. Pl. Bomb. 252 = crinifera.
Schomburgkii, *Lindl. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 673.—Guiana.
Schwackii, *Rodrig. Orch. Nov.* ii. 254.—Bras.
Schweinfurthii, *Reichb. f. Otia Bot. Hamb.* 98.—Abyss.
secunda, *Lindl. Gen. et Sp. Orch.* 307.—Bras.
secundiflora, *Rodrig. Orch. Nov.* ii. 252.—Bras.
setacea, *Lindl. Gen. et Sp. Orch.* 312.—Bras.
setiscala, *Lindl. in Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 673.—Guiana.
setifera, *Lindl. in Ann. Nat. Hist.* iv. (1839) 361.—Mexic.
Sheboldiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 208. —Japon.
Sigillum, *Thou. Orch. Afr.* t. 20.—Ins. Mauric.
simensis, *Reichb. f. in Linnæa*, xxii. (1849) 859.—Afr. trop.
simplifima, *Reichb. f. Xenia Orch.* iii. 18.—Bolivia.
simplex, *Kraenzl. in Bremen, Abh.* vii. (1882) 260. —Madag.
sochensis, *Reichb. f. Otia Bot. Hamb.* 100.—Afr. trop.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 96.—Ins. Socotra.
sparsiflora, *S. Watts. in Proc. Am. Acad.* xii. (1877) 276.—Am. bor. occ.
spathacea, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. ii. (1846) 99.—Mexic.
spathulata, *Benth. in Journ. Linn. Soc.* xviii. (1881) 355.—Reg. Himal.
spatulifolia, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 140.—Burma.

HABENARIA:—
species, *Poepp. & Endl. Nov. Gen. et Sp.* i. 46.—Peruv.
spectabilis, Spreng. Syst. iii. 689 = Orchis spectabilis.
spiralis, *A. Rich. in Mém. Soc. Hist. Nat. Par.* iv. (1838) 21.—Ins. Borbon.
spiralis, *Trim. Cat. Pl. Ceyl.* 91.—Ind. or. ; Zeylan.
Spiranthes, *Reichb. f. in Flora*, xlviii. (1865) 178.—Afr. trop.
Spiranthes, *Reichb. f. & Warm. in Otia Bot. Hamb.* 81.—Bras.
stenochila, *Lindl. in Journ. Linn. Soc.* vi. (1862) 139.—Afr. trop.
stenodon, *Reichb. f. in Linnaea*, xli. (1877) 17.—*N.* Hebrid.
stenopetala, *Lindl. Gen. et Sp. Orch.* 319.—Reg. Himal.
stenopetala, Lindl. l. c. 324 = rhynchocarpa.
stenostachya, *Benth. Fl. Hongk.* 362.—Ind. or. ; China.
Steudneri, *Reichb. f. Otia Bot. Hamb.* 101.—Nubia.
stricta, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 510.—Madag.
strictissima, *Reichb. f. in Linnaea*, xviii. (1844) 407.—Mexic.
stylites, *Reichb. f. & S. Moore, in Journ. Bot.* xvi. (1878) 136.—Afr. trop.
suaveolens, *Dalz. in Hook. Kew Journ.* ii. (1850) 268.—Ind. or.
subcalcarata, *Rodrig. Orch. Nov.* i. 159.—Bras.
subpubens, *A. Rich in Ann. Sc. Nat. Sér.* II. xv. (1841) 75.—Ind. or.
superflua, *Reichb. f. in Seem. Fl. Vit.* 293.—Ins. Fiji.
supervacanea, *Reichb. f. in Seem. l. c.*—Ins. Fiji.
Susannæ, *R. Br. Prod.* 312 ; *Blume, Bijdr.* 402.—Ind. or. ; Malaya ; Chana.
Sutteri, Reichb. f. in Linnæa, xxv. (1852) 229 = diphylla.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 38.—Ins. Tahiti.
tenerrima, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 505.—Madag.
tentaculata, *Reichb. f. Otia Bot. Hamb.* 24.—Borneo.
tentaculigera, *Reichb. f. in Flora*, l. (1867) 191.—Afr. trop.
tenuior, *N. E. Br. in Gard. Chron.* (1885) ii. 307.—Afr. austr.
tenuis, Grift. in Calc. Journ. Nat. Hist. iv. (1844) 379 = viridiflora, *R. Br.*
ternatea, *Reichb. f. in Linnæa*, xli. (1877) 50.—Ins. Ternate.
tetramera, Bolus, in Journ. Linn. Soc. xix. (1882) 340 = tetrapetala.
tetrapetala, *Reichb. f. in Flora*, xlviii. (1865) 180.—Afr. austr.
thomana, *Reichb. f. Otia Bot. Hamb.* 97.—Afr. trop.
Thomasii, *Reichb. f. in Journ. Linn. Soc.* xxi. (1885) 404.—Afr. trop.
Thurberi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 389.—Arizona.
timorensis, *Ridl. in Forbes, Nat. Wand. East. Archip.* 319.—Ins. Timor.
tipuliflora, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 139.—Ind. or.
tipuloides, *Benth. in Journ. Linn. Soc.* xviii. (1881) 356.—Kamtschatka.
tomentella, *Reichb. f. in Flora*, lxviii. (1885) 536.—Ins. Comor.
tradescantifolia, *Reichb. f. in Seem. Fl. Vit.* 293.—Ins. Fiji.
transvaalvica, *Schur, Enum. Pl. Transs.* 645.—Transvaly.
trichocorna, *Rodrig. Orch. Nov.* ii. 256.—Bras.
trichosantha, *Wall. Cat.* n. 7058.—Burma.
trilobata, *A. Rich. Fl. Cub. Fanerog.* ii. 249.—Cuba.
tridactyla, *A. Rich. Tent. Fl. Abyss.* i. 291.—Abyssin.
tridactylites, *Lindl. Gen. et Sp. Orch.* 318.—Ins. Canar.
trideus, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 208.—Afr. austr.
tridentata, *Hook. Exot. Fl.* t. 81.—Am. bor.
triflda, *H. B. & K. Nov. Gen. et Sp.* i. 330.—N. Granat.
trifloris, *D. Don, Prod. Fl. Nep.* 28.—Reg. Himal.
trinervia, Wight, Ic. t. 1701 = digitata.
triptera, *Reichb. f. in Linnæa*, xxii. (1849) 814.—Mexic.

HABENARIA:—
truncata, *Lindl. Gen. et Sp. Orch.* 311.—Madag.
uliginosa, *Reichb. f. in Linnæa*, xxii. (1849) 812.—Bras.
umbraticola, *Rodrig. Orch. Nov.* i. 156.—Bras.
umalaschensis, S. Watz. in Proc. Am. Acad. xii. (1877) 277 = Herminium congestum.
uniflora, Dalz. in Hook. Kew Journ. iii. (1851) 344 = rariflora.
uniflora, D. Don, Prod. Fl. Nep. 25 = triflora ?
uniflora, Griff. Ic. Pl. Asiat. t. 388. f. 2 = Diplomeris pulchella.
vaginata, *A. Rich. in Ann. Sc. Nat. Sér.* II. xiv. (1840) 269.—Abyssin.
Vaupellii, *Reichb. f. & Warm. in Otia Bot. Hamb.* 79.—Bras.
vesicalosa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* iv. (1828) 19.—Ins. Borbon.
vidua, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 140.—Burma.
virescens, Spreng. Syst. iii. 688 = herbiola.
viridiflora, *R. Br. Prod.* 312, in adnot.—Ind. or.
viridiflora, *Span. in Linnæa*, xv. (1841) 478.—Ins. Timor.
viridis, *R. Br. in Ait. Hort. Kew.* ed. II. v. 192.—Europ. ; Am. bor.
viridis, Cham. in Linnæa, iii. (1828) 21 = bracteata.
vulcanica, *Hemsl. Biol. Centr. Am. Bot.* iii. 306.—Mexic.
Walteri, *Reichb. f. Otia Bot. Hamb.* 98.—Afr. trop.
Warmingii, *Reichb. f. & Warm. l. c.* 80.—Bras.
Welwitschii, *Reichb. f. in Flora*, xlviii. (1865) 179.—Afr. trop.
Wightii, *Trim. Cat. Pl. Ceyl.* 91.—Ind. or.
xanthantha, *F. Muell. Fragm.* vii. 16.—Austral.
zambesina, *Reichb. f. Otia Bot. Hamb.* 96.—Afr. trop.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 85.—Java.

HABERLEA, Frivald. in Magyar Tud. Tars. Evkon. ii. (1835) 249. t. 1. *GESNERACEÆ*, Benth. & Hook. f. ii. 1025.
Heidreichii, Boiss. Diagn. Ser. II. iii. 141 = Ramondia Heidreichii.
rhodopensis, *Frivald. in Magyar Tud. Tars. Evkon.* ii. (1835) 249.—Graecia.

HABERLEA, Pohl, ex Baker, in Mart. Fl. Bras. vi. 11. 341 (1876) = **Eupatorium**, Tourn. (Compos.).
affinis, Pohl, ex Baker, l. c. 343 = E. kleinioides.
dentata, Pohl, ex Baker, l. c. = E. kleinioides.
divaricata, Pohl, ex Baker, l. c. 344 = E. urticæfolium.
pinnata, Pohl, ex Baker, l. c. 199 = Lomatorona artemisiifolia.
sperguloides, Pohl, ex Baker, l. c. 341 = E. capillare.

HABERLIA, Dennst. Schluss. Hort. Malab. 30 (1818) = **Odina**, Roxb. (Anacard.).
grandis, Dennst. l. c. = O. Wodier.

HABERSHAMIA, Rafin. Neogenyt. 2 (1825) = **Herpestis**, Gaertn. f. (Scrophul.).
cuneifolia, Rafin. l. c. = H. Monnieria.

HABLANTHERA, Hochst. in Bot. Zeit. ii. (1844) 853 = HAPLANTHERA, Hochst. = **Buttya**, Harv. (Acanth.).

HABLITZIA, Bieb. in Mém. Soc. Nat. Mosc. v. (1817) 24. *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 49.
tamnoides, *Bieb. l. c.*—Reg. Caucas.

HABLITZLIA, Reichb. Nom. 164 (1841) = præc.

HABLIZIA, Spreng. Syst. i. 824 (1825) = præc.

HABLIZLIA, Pritz. Ic. Ind. ii. 139 (1855) = præc.

HABRACANTHUS, Nees, in DC. Prod. xi. 312 (1847). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1106.
haematodes, Nees, l. c. = Glockeria haematodes.
sanguineus, *Nees, l. c.*—Mexic.
sylvaticus, *Nees. l. c.*—Mexic.

HABRANTHUS, Herb. in Bot. Mag. t. 2464 (1824) ; et in Bot. Reg. t. 1845 (1830) = **Hippeastrum**, Herb. (Amaryll.).
advenus, Herb. in Bot. Mag. sub t. 2685 = Hipp. advenum.
Andersoni, Herb. ex Lindl. Bot. Reg. t. 1845 = Zephyranthes Andersoniana.
andicola, Herb. Amaryll. 168.—Chili.
angustus, Herb. in Bot. Mag. t. 2689 = Hipp. bifidum.
Bagnoldi, Herb. in Bot. Reg. t. 1396 = Hipp. Bagnoldi.
Bagnoldianus, Herb. Amaryll. 162 = præc.
Bertersi, M. Roem. Syn. Ensat. 103 = *robustum*.
Berteroanus, Phil. in Linnæa, xxix. (1857-58) 66 = Hipp. Berteroanum.
bifidus, Herb. in Bot. Mag. t. 2597 = Hipp. bifidum.
Boothianus, Herb. Amaryll. 124 = Zephyranthes gracilifolius.
cearensis, Herb. in Bot. Mag. sub t. 3779.—Bras.
chilensis, Herb. Amaryll. 163 = Hipp. chilense.
concolor, Lindl. in Proc. Hort. Soc. (1838) 6 = Zephyranthes concolor.
flavus, Phil. in Linnæa, xxix. (1857-58) 66.—Chili.
fulgens, Hook. f. Bot. Mag. t. 5863.—Hab. ?
Gillesianus, M. Roem. Syn. Ensat. 88 = Hipp. Bagnoldi.
glaioloides, Hieron. in Bol. Acad. Nac. Córdova, iv. (1881) 70.—Reg. Argent.
gracilifolius, Herb. in Bot. Mag. t. 2464 = Zephyranthes gracilifolia.
hesperius, Herb. Amaryll. 161 = Hipp. advenum.
intermedius, Herb. l. c. 160 = Hipp. bifidum.
kermesinus, Herb. l. c. 159 = Hipp. bifidum.
lineatus, Phil. in Anal. Univ. Chil. (1878) 542 = Hipp. lineatum.
litoralis, M. Roem. Syn. Ensat 96 = Hipp. bifidum.
lorifolius, Herb. Amaryll. 160, in obs.—Am. austr.
maculatus, Herb. l. c. 167.—Chili.
maculosus, Phil. in Linnæa, xxvi. (1854) 11. 406 = Hipp. advenum.
miniatus, D. Don, ex Sweet, Brit. Flow. Gard. Ser. II. vi. t. 213 = Hipp. advenum.
montanus, Phil. in Anal. Univ. Chil. (1878) resmp. 66 = Hipp. montanum.
memoralis, Herb. Amaryll. 159 = Hipp. bifidum.
nubilis, Herb. in Lindl. Bot. Reg. (1844) Misc. 88 = Hipp. bifidum.
pallidus, Lodd. Bot. Cab. t. 1760.—Chili.
parvulus, Pritz. Ic. Ind. 522 = Zephyranthes Andersoniana.
pedunculosus, Herb. Amaryll. 161. t. 26. f. 3 = Hipp. bifidum.
phycelloides, Herb. in Bot. Reg. t. 1417 = Hipp. phycelloides.
pratensis, Herb. in Bot. Mag. t. 3961 = Hipp. pratense.
pulcher, Herb. Amaryll. 161. t. 26. f. 1 = Hipp. bifidum.
pumilus, Lodd. Bot. Cab. t. 1771 = Hipp. roseum.
punctatus, Herb. Amaryll. 403. t. 47. f. 1 = Hipp. Bagnoldi.
robustus, Herb. l. c. 166.—Bras.
roseus, Sweet, Brit. Flow. Gard. Ser. II. t. 107 = Hipp. roseum.
spathaceus, Herb. in Bot. Mag. t. 2597 = Hipp. bifidum.
speciosus, Herb. Amaryll. 158 = Hipp. pratense.
splendens, Renjifo, in Renjifo & Phil. Descr. Alg. Pl. Nuev. (1884) reimpr. 6.—Chili.
sylvaticus, Herb. Amaryll. 166.—Bras.
tenuanus, Herb. ex Steud. Nom. ed. II. i. 717 = Zephyranthes tenuana.
versicolor, Herb. in Bot. Mag. t. 2465 = Zephyranthes versicolor.

HABRANTUS, Dum. Fam. Anal. 58 (1829) = præc.

HABROBIA, Fenzl, in Bot. Zeit. i. (1843) 323. *ILLECEBRACEÆ*, Benth. & Hook. f. iii. 19.
spinuliflora, *Fenzl, l. c.* 328.—Syria ; Persia.

HABROTHAMNUS, Endl. Gen. 967 (1839) = **Cestrum**, Linn. (Solanac.).
auranticaus, Regel, Ind. Sem. Hort. Turic. (1850) 4.—Cf. Linnæa, xxiv. (1851) 231 = C. Regelii.
aurantiacus, Seem. in Bonplandia, vii. (1859) 15 = C. aurantiacum.
corymbosus, Endl. ex Meisn. Gen. Comm. 185 ; et ex Walp. Rep. iii. 125 = C. Endlicheri.
cyaneus, Lindl. Bot. Reg. (1844) Misc. 73 = Iochroma tubulosum.
elegans, Brongn. in Fl. des Serres, Sér. I. ii. (Févr. 1846) t. 9 = C. elegans.

HABROTHAMNUS :—
elegans, Scheidw. ex Walp. Rep. iii. 192 = Cestrum fasciculatum.
fasciculatus, Benth. Pl. Hartw. 49 = C. Hartwegi.
fasciculatus, Brongn. in Rev. Hortic. Sér. II. iv. (1845–46) 26 = C. fasciculatum.
Hartwegi, Seem. in Bonplandia, vii. (1859) 15 = C. Hartwegi.
Hueglii, Regel, in Bot. Zeit. ix. (1851) 598 = C. elegans.
Newelli, Hort. Veitch. ex Gard. Chron. (1877) i. 424.—Hab.?
paniculatus, Mart. & Gal. in Bull. Acad. Brux. xii. (1845) 148 = C. sylvaticum.
porphyreus, Seem. in Bonplandia, vii. (1859) 15 = C. porphyreum.
purpureus, Lindl. Bot. Reg. (1844) Misc. 12. t. 43 = C. fasciculatum.
vosmo, Miers, Illustr. S. Am. Pl. ii. 81 = C. roseum.
stellatus, Lem. in Fl. des Serres, Sér. I. ii. (Févr. 1846) sub t. 8.—Hab.?
tomentosus, Benth. Pl. Hartw. 49 = C. Benthami.

HABROZIA, Lindl. Veg. Kingd. 638 (1847) = **Habrosia**, Fenzl (Illecebr.).

HABSBURGIA, Mart. Syst. Mat. Med. Bras. 36, nomen tantum (1843) = **Skytanthus**, Meyen (Apocyn.).
comans, Mart. l. c. (= *Neriandra Martiana*).—Bras.

HABSIA, Steud. Nom. ed. II. i. 719 (1840), sphalm. = **Guettarda**, Linn. (Rubiac.).
arborescens, Steud. l. c. = G. argentea.

HABZELIA, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 207 (Mém. Anon.) = **Xylopia**, Linn. (Anonac.).
acuta, Mig. Fl. Ind. Bat. i. ii. 87.—Java.
æthiopica, A. DC. Mém. Anon. 31 = X. æthiopica.
aromatica, A. DC. l. c. 32 = Unona discolor.
cubensis, DC. ex Steud. Nom. ed. II. i. 717 = X. cubensis.
discreta, A. DC. Mém. Anon. 32 = Unona discreta.
ferruginea, Hook. f. & Thoms. Fl. Ind. i. 124 = X. ferruginea.
obtusifolia, A. DC. l. c. 42 = X. obtusifolia, *A. Rich. oxyantha*, Hook. f. & Thoms. Fl. Ind. i. 124 = X. oxyantha.
undulata, A. DC. Mém. Anon. 32 = X. undulata.

HACHENBACHIA, D. Dietr. Syn. Pl. i. 127 (1839) = **Hagenbachia**, Nees & Mart. (Haemodor.).

HACHETTIA, Baill. in Bull. Soc. Linn. Par. i. (1880) 229. *BALANOPHOREÆ*, Benth. & Hook. f. iii. 1221.
austro-caledonica, *Baill. l. c.* 230.—N. Caled.

HACKELA, Pohl, ex Flora, viii. (1825) 163 = HACKELIA, Pohl = **Curtia**, Cham. & Schlecht. (Gentian.).

HACKELIA, Opiz, in Bercht. Okon. Fl. Boehm. ii. 11. 146 (1838) = **Echinospermum**, Sw. (Borag.).
deflexa, Opiz, l. c. 147; Seznam, 49 = E. deflexum.

HACKELIA, Pohl, ex Griseb. in DC. Prod. ix. 56 (1845) = **Curtia**, Cham. & Schlecht. (Gentian.).
stellata, Pohl, ex Griseb. l. c. = C. gentianoides.

HACQUETIA, Neck. Elem. i. 182 (1790). *UMBELLIFERÆ*, Benth. & Hook. f. i. 879.
Dondia, Spreng. Pl. Umb. Prod. 91 (1813).
Dondisia, Reichb. Consp. 144 (1828).
bractogama, DC. Prod. iv. 068 (= *Posoa coriacea*).—Reg. Andin.
Epipactis, *DC. l. c.* 55.—Europ.

HADESTAPHYLUM, Dennst Schlüssl. Hort. Malab. 80 (1818) = **Holigarna**, Buch.-Ham. (Anacard.).
cauticum, Dennst. l. c. = H. longifolia.

HAECKERIA, F. Muell. in Linnæa, xxv. (1852) 406 = **Humea**, Sm. (Compos.).
cassiniæformis, F. Muell. l. c. = Humea cassiniacea.
oxathamnoides, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 45 = Humea oxathamnoides.

HAEMADICTION, Steud. Nom. ed. II. i. 718 (1840) = seq.

HAEMADICTYON, Lindl. in Trans. Hort. Soc. vi. (1826) 70 = **Prestonia**, R. Br. (Apocynac.).
acutifolium, Benth. ex Muell. Arg. in Mart. Fl. Bras. vi. i. 167.—Bras.
amazonicum, Benth. ex Muell. Arg. l. c. 166.—Bras.
amœnum, A. DC. in DC. Prod. viii. 428 = P. annularis.
asperum, Muell. Arg. in Mart. Fl. Bras. vi. i. 169.—Bras.
bracteosum, Muell. Arg. l. c. 168.—Bras.
calycinosum, Miers, Apocyn. S. Am. 280.—Peruv.
calycinum, Lindl. ex Miers, l. c. 259 = *Gaudichaudii*.
cayennense, A. DC. in DC. Prod. viii. 427.—Guian.
circinale, G. Don, Gen. Syst. iv. 88 = Echites circinalis.
contortum, Mart. & Gal. in Bull. Acad. Brux. xi. (1844) 360 = P. contorta.
denticulatum, Miers, Apocyn. S. Am. 257 = *macronewron*.
exsertum, A. DC. in DC. Prod. viii. 426.—N. Granat.
Gaudichaudii, A. DC. l. c.—Bras.
glabratum, A. DC. l. c. 427 = P. glabrata.
grandiflorum, A. DC. l. c.—Peruv.
grandiflorum, A. DC. l. c. 426 = Odontadenia speciosa.
lagoinse. Muell. Arg. in Kjoeb. Vidensk. Meddel. (1869) 115.—Bras.
macrocarpum, Muell. Arg. in Mart. Fl. Bras. vi. i. 169.—Bras.
marginatum, Benth. in Hook. Journ. Bot. iii. (1841) 260.—Guian.
megacarpon, Muell. Arg. in Mart. Fl. Bras. vi. i. 170.—Bras.
membranaceum, Muell. Arg. l. c. 167.—Bras.
mexicanum, A. DC. in DC. Prod. viii. 428.—Mexic.
molle, A. DC. l. c. 427 = P. mollis.
mutans, A. DC. l. c. 426 = *venosum*.
ovatum, Miers, Apocyn. S. Am. 256 = *Gaudichaudii*.
pallidum, A. DC. in DC. Prod. viii. 428.—Peruv.
pilosiflorum, Muell. Arg. in Linnæa, xxx. (1859–60) 435.—Am. trop.
parvifloram, Benth. Fl Hartw. 355.—N. Granat.
refulgens, Linden & André, Illustr. Hortic. xviii. (1871) 14. t. 49.—Peruv.
Riedelii, Muell. Arg. in Mart. Fl. Bras. vi. i. 170.—Bras.
schiedeanum, Muell. Arg. in Linnæa, xxx. (1859–60) 435 = P. schiedeana.
solanifolium, Muell. Arg. in Mart. Fl. Bras. vi. i. 171.—Bras.
subrectum, G. Don, Gen. Syst. iv. 88.—Ind. or.
tomentellum, Benth. Bot. Voy. Sulph. 126.—Am. trop.
trifidum, Poepp. & Endl. Nov. Gen. et Sp. iii. 67. t. 275.—Bras.
venosum, Lindl. in Trans. Hort. Soc. vi. (1825) 71.—Ind. occ.
Warmingii, Muell. Arg. in Kjoeb. Vidensk. Meddel. (1869) 116.—Bras.

HAEMANTHUS, [Tourn.] Linn. Syst. ed. I (1735). *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 720.
Diacles, Salisb. Gen. Pl. Fragm. 120 (1866).
Gyaxis, Salisb. l. c. 121 (1866).
Leucodermis, Kaſn. Fl. Tellur. iv. 19 (1836).
Melicum, Salisb. Gen. Pl. Fragm. 120 (1866).
Nerissa, Salisb. l. c. 121 (1866).
Perinema, Kaſn. Fl. Tellur. iv. 20 (1836).
Scadoxus, Raſn. l. c. 19 (1836).
Serena, Raſn. l. c. 20 (1836).
abyssinicus, Herb. Amaryll. 222 = multiflorus.
albiflos, *Jacq. Hort. Schœnb.* i. 31.—Afr. austr.
albo-maculatus, *Baker, in Gard. Chron.* (1878) i. 202.—Afr. austr.
amaryllioides, *Jacq. Hort. Schœnb.* iv. 5. t. 408.—Afr. austr.
angolensis, *Welw. in Journ. Bot. xvi.* (1878) 194.—Afr. trop.
angustifolius, Salisb. Prod. 217 = pubescens.
arabicus, M. Roem. Syn. Ensat. 42 = multiflorus.
Arnotii, *Baker, in Gard. Chron.* (1878) ii. 492.—Afr. trop.
Baurii, *Hort. Kew.—Cf. Gard. Chron.* (1885) ii. 813.—Afr. austr.
brevifolius, Herb. Amaryll. 224. t. 90. f. 3 = canescens.
caliconus, *Durch. ex M. Roem. Syn. Ensat.* 43.—Afr. austr.
capensis, *Linn. Prael. ed. Gleditsch*, 278.—Afr. austr.

HAEMANTHUS:—
carinatus, Linn. Sp. Pl. ed. II. 418 = coccineus.
carneus, [*Ker-Gawl.*] *in Bot. Reg.* t. 509.—Afr. austr.
cauticus, *Molina, Sagg. Chile*, ed. II. 199, 284.—Chili.
cernuiflorus, *Y. Bell, ex Drapiez, Herb. Amat.* iv. (1830) t. 237.—Afr. austr.
ciliaris, Linn. Sp. Pl. ed. II. 418 = Buphane disticha.
cinnabarinus, *Decne. in Fl. des Serres*, Sér. II. ii. (1857) 57. t. 1195.—Afr. trop.
coarctatus, Jacq. Hort. Schœnb. i. 30 = coccineus.
coccineus, *Linn. Sp. Pl.* 325.—Afr. austr.
coerulescens, Steud. Nom. ed. II. i. 72, in syn. = Hypoxis stellatus.
colchiciflos, Salisb. Prod. 217 = multiflorus.
concolor, *Herb. Amaryll.* 238.—Afr. austr.
crassipes, *Jacq. Hort. Schœnb.* iv. 7. t. 412.—Afr. austr.
cruentatus, *Schum. & Thonn. Beskr. Guin. Pl.* 168.—Afr. trop.
deformis, *Hook. f. Bot. Mag.* t. 5908.—Afr. austr.
delagoensis, Herb. Amaryll. 233 = multiflorus.
dubius, H. B. & K. Nov. Gen. et Sp. i. 281 (= *Collania dubia*).—Ecuador.
falcatus, Thunb. Prod. Pl. Cap. 58 = Ammocharis falcata.
filiflorus, *Baker, ex Hiern, in Journ. Bot.* xvi. (1878) 194.—Afr. trop.
grandifolius, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 96.—Ins. Socotra.
guttatus, Banks, ex Kunth, Enum. Pl. v. 605 = Buphane ciliaris.
hirsutus, *Baker, in Gard. Chron.* (1878) i. 756.—Afr. austr.
Hookerianus, *Herb. Amaryll.* 404. t. 46. f. 7.—Afr. austr.
humilis, *Jacq. Hort. Schœnb.* iv. 6. t. 411.—Afr. austr.
hyalocarpus, *Jacq. l. c.* 5. t. 409.—Afr. austr.
hydrophilus, Thunb. Mus. Ups. ex Schult. f. Syst. vii. 892 = Crinum campanulatum.
incarnatus, *Burch. ex Herb. Amaryll.* 237.—Afr. austr.
insignis, Hook. Bot. Mag. t. 4745 = magnificus.
intermedius, M. Roem. Syn. Ensat. 86 = albiflos.
Kalbreyeri, Baker, in Gard. Chron. (1878) ii. 202 = multiflorus.
Katherinæ, *Baker, l. c.* (1877) i. 656.—Afr. austr.
Lamberianus, *Schult. f. Syst.* vii. 891.—Afr. austr.
lanceæfolius, *Jacq. Hort. Schœnb.* i. 31. t. 60.—Afr. austr.
latifolius, Salisb. Prod. 216 = coccineus.
lescanthus, *Miq. in Journ. Bot. Néerl.* i. (1861) 38.—Afr. austr.
maculatus, *Jacq. ex Loud. Hort. Brit.* 119.—Afr. austr.
magnificus, *Herb. in Bot. Reg.* (1841) *Misc.* 71.—Afr. austr.
Mannii, *Baker, in Bot. Mag.* t. 6364.—Guian.
moschatus, *Jacq. Hort. Schœnb.* iv. 6. t. 410.—Afr. austr.
multiflorus, *Martyn, Monog. cum Ic. ; et ex Willd. Sp. Pl.* ii. 25.—Afr. austr.
natalensis, *Hook. Bot. Mag.* t. 5378.—Afr. austr.
obliquus, Donn, Hort. Cantab. ed. II. 38 = rotundifolius.
orbicularis, Donn, l. c. = rotundifolius.
orchidiflorus, Salisb. Prod. 217 = puniceus.
orientalis, Thunb. Prod. Pl. Cap. 59 = Brunsvigia gigantea.
pubescens, Blanco, Fl. Filip. ed. I. 253 = Crinum asiaticum.
pubescens, *Linn. f. Suppl.* 193.—Afr. austr.
Puntilo, *Jacq. Hort. Schœnb.* i. 32. t. 61.—Afr. austr.
puniceus, *Linn. Sp. Pl.* 326.—Afr. austr.
quadrifoliatus, Jacq. Hort. Schœnb. i. 30. t. 56 = pubescens.
Redoutéanus, M. Roem. Syn. Ensat. 38 = puniceus.
reniens, Hort. Angl. ex Link, Enum. Hort. Berol. i. 300 = canescens.
rotularis, *Baker, in Gard. Chron.* (1877) i. 656.—Afr. trop.
rotundifolius, [*Ker-Gawl.*] *in Bot. Mag.* t. 1618.—Afr. austr.
Rouyeri, Hort.—*Cf. Flor. Mag.* N. S. (1876) t. 168 = magnificus.
saponius, *Baker, in Gard. Chron.* (1877) i. 656.—Afr. trop.

HAEMANTHUS:—
sanguineus, *Jacq. Hort. Schoenbr.* iv. 4. t. 407.—Afr. austr.
sarmentosis, Thunb. Prod. Pl. Cap. 58 = Nerine sarmentosa.
sinuatus, *Thunb. Mus. Ups. ex Schult. f. Syst.* vii. 893.—Afr. austr.
spiralis, [Soland. in] Ait. Hort. Kew. ed. I. i. 405 = Carpolyza spiralis.
stellatus, Steud. Nom. ed. II. i. 71, in syn. = Hypoxis stellata.
strigosus, Herb. Amaryll. 284. t. 30. f. 2 = carneus.
tenellus, Steud. Nom. ed. I. 398, in syn. = Carpolyza spiralis.
tenuiflorus, Herb. in Bot. Mag. t. 3870 = multiflorus.
tigrinus, *Jacq. Hort. Schoenbr.* i. 29. t. 56.—Afr. austr.
toxicarius, Linn. f. ex Ait. Hort. Kew. ed. I. i. 405; Thunb. Prod. Pl. Cap. 59 = Buphane disticha.
triphathinus, Hort. ex M. Roem. Syn. Kossat. 87 = pubescens.
undulatus, *Herb. Amaryll.* 283. t. 30. f. 1.—Afr. austr.
undulatus, Thunb. Prod. Pl. Cap. 58 = Nerine undulata.
vaginatus, Thunb. Fl. Cap. ii. 254 = seq.
vaginatus, Thunb. Prod. Pl. Cap. 188 = Strumaria truncata.
viridescens, Herb. Amaryll. 285 = albiflos.
zebrinus, *Herb. l. c.* 287.—Afr. austr.

HAEMARIA, Lindl. Orch. Scel. 9 (1826). *ORCHIDEAE*, Benth. & Hook. f. iii. 601.
DICHROPHYLA, Rafin. Fl. Tellur. iv 39 (1836).
LUDISIA, A. Rich. in Dict. Clas. Hist. Nat. vii. 437 (1826).
MYODA, Lindl. in Wall. Cat. n. 7390 (1832); et Bot. Reg. xix. sub t. 1618 (1833).
argyroneura, Miq. Choix, t. 16. f. 1 = Dossinia marmorata.
discolor, *Lindl. Gen. et Sp. Orch.* 490.—China.

HAEMARTHRIA, Munro, in Journ. Linn. Soc. vi. (1862) 55 = HEMARTHRIA, R. Br. = **Bottboellia**, Linn. f. (Gramin.).

HAEMATOBANCHE, Presl, Epim. Bot. 249 (1849) = **Hyobanche**, Linn. (Scrophul.).
sanguinea, Presl, t. c. = Hyobanche sanguinea.

HAEMATOCARPUS, Miers, in Ann. Nat. Hist. Sér. III. xix. (1867) 194. *MENISPERMACEAE*, Benth. & Hook. f. i. 964.
comptus, Miers, t. c. 197 = Thomsonii.
Thomsoni, *Miers, l. c.*—Reg. Himal.

HAEMATOLEPIS, Presl, Epim. Bot. 238 (1849) = **Cytinus**, Tourn. (Balanoph.).
sanguinea, Presl, t. c. = C. dioicus.

HAEMATORCHIS, Blume, Rumphia, iv. t. 200 B (1848) = **Galeola**, Lour. (Orchid.).
altissima, Blume, l. c. = G. altissima.

HAEMATOSPERMUM, Wall. Cat. n. 7053 (1832) = **Homonoia**, Lour. (Euphorb.).
cuneatum, Wall. l. c. n. 7054 = Hom. retusa.
ferrifolium, Wall. l. c. n. 7052 = Hom. riparia.
riparium, Wall. l. c. n. 7935 = Hom. riparia.
salicinum, Baill. Etud. Gén. Euph. 298 = Hom. riparia.

HAEMATOSTAPHIS, Hook. f. in Trans. Linn. Soc xxiii. (1860) 169. t. 25. *ANACARDIACEAE*, Benth. & Hook. f. i. 422.
Barteri, *Hook. f. l. c.*—Afr. trop.

HAEMATOSTROBUS, Endl. Gen. 76, in adnot. (1836) = **Phoenicia**, Vahl (Balanoph.).

HAEMATOXYLUM, Scop. Introd. 335 (1777) = seq.

HAEMATOXYLON, Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 567.
Brasiletto, *Kunth. Fl. Columb.* ii. 97. t. 114.—N. Granat.
campechianum, *Linn. Sp. Pl.* 384.—Am. trop.

HAEMATOXYLUM, Linn. Gen. ed. I. 342 (1737) = prec.

HAEMAX, E. Mey. Comm. Pl. Afr. Austr. 228 (1837) = **Astephanus**, R. Br. (Asclep.).
Dregei, E. Mey. l. c. = A. Dregei.
Massoni, E. Mey. l. c. = A. Massoni.

HAEMOCARPUS, *Noronha, ex Thou. Gen. Nov. Madag.* 15 (1806) = **Haronga**, Thou. (Hyperic.).
axillaris, Spreng. Syst. iii. 833 = Psorospermum axillare.
corymbosus, Spreng. l. c. = Psorospermum citrifolium.
cymosus, Spreng l. c. = Psorospermum discolor.
paniculatus, Spreng. l. c. = Har. madagascariensis.

HAEMOCHARIS, Salisb. Parad. Lond. sub t. 56 (1806) = **Lapiacea**, H. B. & K. (1821) (Ternstroem.).
barbinervis, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 146 = L. barbinervis.
Hormatoxylum, Salisb. Parad. Lond. sub t. 56 = L. Haematoxylum.
intermedia, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 146 = L. intermedia.
parviflora, Choisy, l. c. = L. parviflora.
quinoderma, Choisy, l. c. 147 = L. quinoderma.
semserrata, Mart. & Zucc. Nov. Gen. et Sp. i. 107. t. 61 = L. semiserrata.
speciosa, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 148 = L. speciosa.
tomentosa, Mart. & Zucc. Nov. Gen. et Sp. i. 106. t. 67 = L. tomentosa.
villosa, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 146 = L. villosa.

HAEMODORON, Reichb. Consp 212b (1828) = **Clavanche**, Hoffmgg. & Link (Orobanch.).

HAEMODORUM, Sm. in Trans. Linn. Soc iv. (1798) 218. *HAEMODORACEAE*, Benth. & Hook. f. iii. 673.
hirsutissima, *Spreng. Syst.* iv. *Cur. Post.* 24.—Bras.
brevicaule, *F. Muell. Fragm.* i. 84.—Austral.
brevisepalum, *Benth. Fl. Austral.* vi. 420.—Austral.
coccineum, K. Br. Prod. 300 = corymbosum.
corymbosum, *Vahl, Enum.* ii. 179.—Austral.
distichophyllum, *Hook. Ic. Pl.* t. 866.—Tasman.
edule, Lehm. Pl. Preiss. ii. 15 = spicatum.
ensifolium, *F. Muell. Fragm.* i. 84.—Austral.
laxum, *R. Br. Prod.* 300.—Austral.
leptostachyum, *Benth. Fl. Austral.* vi. 423.—Austral.
paniculatum, *Lindl. Swan River App.* 44.—Austral.
parviflorum, *Benth. Fl. Austral.* vi. 423.—Austral.
planifolium, *R. Br. Prod.* 300.—Austral.
polycephalum, Endl. in Lehm. Pl. Preiss. ii. 16 = simplex.
simplex, *Lindl. Swan River App.* 44.—Austral.
simulans, *F. Muell. Fragm.* vii. 117.—Austral.
spurafforum, *F. Muell. l. c.*—Austral.
spicatum, *R. Br. Prod.* 300.—Austral.
strictum, Endl. in Lehm. Pl. Preiss. ii. 15 = paniculatum.
subviens, *F. Muell. Fragm.* i. 63.—Austral.
tenuifolium, *A. Cunn. ex Benth. Fl. Austral.* vi. 423.—Austral.
teretifolium, *R. Br. Prod.* 300.—Austral.

HAEMOSPERMUM, Reinw. in Blume, Bijdr. 1018 (1825) = **Geniostoma**, Forst. (Logan.).
arboreum, Reinw. ex Blume, l. c. = G. Haemospermum.
molaccanum, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 240 = G. Laslostemon.

HAENCKEA, *Juss. in Dict. Sc. Nat. xx. 204* (1821) = HAENKEA, Ruiz & Pav. = **Schoepfia**, Schreb. (Olacin.).

HAENELIA, Walp. Rep. ii. 974 (1843) = **Amellus**, Tourn. (Compos.).
capensis, Walp. l. c. = A. Lychnitis.

HAENIANTHUS, Griseb. in Mem. Am Acad. N. S viii. (1863) (Pl. Wright. Cub. ii.) 518 = **Linociera**, Sw. (Oleac.).
incrassatus, Griseb. Fl. Brit. W. Ind. 408.—Ind occ.
salicifolius, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 518.—Cuba.

HAENKAEA, Usteri, Ann. Bot. vi. (1795) 117 = HAENKEA, F. W. Schmidt = **Adenandra**, Willd. (Rutac.).

HAENKEA, Ruiz & Pav. Fl. Per. Ill. t. 231 (1802) = **Schoepfia**, Schreb. (Olacin.).
flexuosa, Ruiz & Pav. l. c. = S. fraxinea.
macrocarpa, Steud. Nom. ed. I. 170, in syn. = Celastrus multiflorus.
multiflora, Ruiz & Pav. Syst. Veg. 65 = Celastrus multiflorus.

HAENKEA, Ruiz & Pav. Prod. 86. t. 6 (1794) = **Maytenus**, Feuill. (Celastr.).

HAENKEA, Salisb. Prod. 174 (1796) = **Portulacaria**, Jacq. (Portul.).
crassifolia, Salisb. l. c. = P. afra.

HAENKEA, F. W. Schmidt, Neue n. Selt. Pfl. 19 (1793) = **Adenandra**, Willd. [1809] (Rutac.).
punctata, F. W. Schmidt, l. c. = A. Schmidtiana.
villosa, F. W. Schmidt, l. c. 21 = A. calycina.

HAENSELERA, Boiss. ex DC. Prod. vii. 83 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 505.
elatior, Willk. in Linnaea, ix. (1852) 41 = grantenesis.
grantensis, *Boiss. ex DC. Prod.* vii. 83.—Hispan.

HAENSELERA, Lag. Gen. et Sp. Nov. 18 (1816) = **Physosperma**, Cusson (Umbell.).
damascenia, Lag. l. c. = P. aquilegifolium.

HAENSELERIA, Reichb. Nom. 96 (1841) = prec.

HAENSLERA, Steud. Nom. ed. II. i. 718 (1840) = prec.

HAGAEA, Vent. Tabl iii. 240 (1799) = **Polycarpaea**, Lam. (Caryophyll.).
aizoaefolia, Bivona, Stirp. Rar. Sic. Man. iii. 7 = Polycarpon aizoefolium.
aristata, Sweet, Hort. Brit. ed. I. 176 = P. corymbosa.
diffusa, Sweet, l. c. = P. Teneriffae.
gnaphaloides, Pers. Syn. i. 262 = P. nivea.
latifolia, Sweet, Hort. Brit. ed. I. 176 = P. latifolia.
polycarpodes, Bivona, Stirp. Rar. Sic. Man. ii. [9] = Polycarpon peploides.
Teneriffae, Pers. Syn. i. 262 = P. Teneriffae.

HAGEA, Pers. Syn. i. 262 (1805) = prec.

HAGEA, Poir. in Dict. Sc. Nat. xii. 253 (1818) = **Brayera**, Kunth (Rosac.).

HAGENBACHIA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 18. t. 2. *HAEMODORACEAE*, Benth. & Hook. f. iii. 674.
brasiliensis, *Nees & Mart. l. c.* 19. t. 2.—Bras.

HAGENIA, J. F. Gmel. Syst. 613 (1791) = **Brayera**, Kunth (Rosac.).
abyssinica, J. F. Gmel. l. c. = B. anthelmintica.

HAGENIA, Moench, Meth. 61 (1794) = **Saponaria**, Linn. (Caryophyll.).
filiformis, Moench, l. c. = S. porrigens.

HAGIDRYAS, Griseb. Reise Rumel. i. 311 (1841) = **Prunus**, Tourn. (Rosac.).
prostrata, Griseb. l. c. = P. prostrata.

HAGIOSERIS, Boiss. Diagn. Ser. I. xi. 85 (1849) = **Picris**, Linn. (Compos.).
amalecitana, Boiss. l. c. 86 = *galilaea*.
galilaea, Boiss. l. c.—Syria.

HAHNIA, Medic. Geschich. 81 (1793) = **Pyrus**, Linn. (Rosac.).
arbutifolia, Medic. l. c. 82 = P. arbutifolia.
Aria, Medic. l. c. 81 = P. Aria.
Chamaemespilus, Medic. l. c. 82 = P. Chamaemespilus.
pinnatifida, Medic. l. c. = P. pinnatifida.
torminalis, Medic. l. c. 81 = P. torminalis.

HAKEA, Schrad. Sert. Hannov. 27. t. 17 (1797). *PROTEACEAE*, Benth. & Hook. f. iii. 181.
CONCHIUM, Sm. in Trans. Linn. Soc. iv. (1798) 215.
MERCKLINIA, Regel, Ind. Sem. Hort. Petrop. (1856) 25; Gartenfl. (1857) 185.
acanthophylla, Link, Enum. Hort. Berol. i. 118 = ceratophylla.

HAKEA:—
acicularis, *Knight, Prot.* 107; *R. Br. in Trans. Linn. Soc.* x. (1810) 181.—Austral.
adnata, *R. Br. Prot. Nov.* 28.—Austral.
ambigua, *Meisn. in Lehm. Pl. Preiss.* ii. 260.—Austral.
amplexicaulis, *R. Br. in Trans. Linn. Soc.* x. (1810) 184.—Austral.
angustifolia, Hort. Angl. ex Meisn. in DC. Prod. xiv. 315 = ulicina.
arborescens, *R. Br. in Trans. Linn. Soc.* x. (1810) 187. —Austral.
attenuata, R. Br. l. c. 188 = varia.
attenuata, Meisn. in DC. Prod. xiv. 406 = auriculata.
auriculata, *Meisn. in Hook. Kew Journ.* vii. (1855) 116.—Austral.
Baxteri, *R. Br. Prot. Nov.* 28.—Austral.
bifrons, Meisn. in DC. Prod. xiv. 400 = microcarpa.
bipinnatifida, *R. Br. Prot. Nov.* 28.—Austral.
Boucheana, Kunth, Ind. Sem. Hort. Berol. (1844) 9.— Cf. Linnæa, xvii. (1844) 499 = trifurcata.
brachypetera, *Meisn. in Hook. Kew Journ.* iv. (1852) 306, nomen; *et in DC. Prod.* xiv. 396.—Austral.
brachyphylla, F. Muell. First Gen. Rep. 17 = acicularis.
Brownii, *Meisn. in Lehm. Pl. Preiss.* i. 269.— Austral.
Candolleana, *Meisn. l. c. ii.* 262.—Austral.
canescens, Link, Enum. Hort. Berol. i. 118 = cinerea.
carinata, F. Muell. ex Meisn. in Linnæa, xxvi. (1853–55) 840 = ulicina.
ceratophylla, *R. Br. in Trans. Linn. Soc.* x. (1810) 184.—Austral.
cervina, Dum.-Cours. Bot. Cult. ed. II. vii. 110 = ceratophylla.
chordophylla, *F. Muell. in Hook. Kew Journ.* ix. (1857) 23.—Austral.
cinerea, *R. Br. in Trans. Linn. Soc.* x. (1810) 186.— Austral.
circumalata, *Meisn. in Hook. Kew Journ.* vii. (1855) 114.—Austral.
clavata, *Labill. Nov.-Holl. Pl.* i. 31. t. 41.—Austral.
commutata, *F. Muell. Fragm.* v. 26.—Austral.
conchifolia, Hook. Ic. Pl. t. 433 = cucullata.
corymbosa, *R. Br. Prot. Nov.* 28.—Austral.
cuneata, *Meisn. in Lehm. Pl. Preiss.* i. 275.— Austral.
cyanophylla, *Meisn. l. c.* 270.—Austral.
crassinervia, Meisn. l. c. 278 = petiolaris.
cristata, *R. Br. Prot. Nov.* 28.—Austral.
cucullata, *R. Br. l. c.* 30.—Austral.
Cunninghamii, *R. Br. l. c.* 26.—Austral.
cyclocarpa, *Lindl. Swan River App.* 36.—Austral.
cyclоptera, *R. Br. in Trans. Linn. Soc.* x. (1810) 182. —Austral.
dactyloides, *Cav. in Anal. Hist. Nat.* i. (1799) 215.— Austral.
decurrens, R. Br. Prot. Nov. 27 = acicularis.
decurrens, A. Cunn. ex Meisn. in DC. Prod. xiv. 321 = Conospermum longifolium.
denticulata, R. Br. Prot. Nov. 28 = glabella.
Drummondi, G. Don, ex Meisn. in DC. Prod. xiv. 409 = cucullata.
drupacea, R. Br. in Trans. Linn. Soc. x. (1810) 221; Koem. & Schult. Syst. iii. 436 = suaveolens!
echinata, *Hort. Madrap. ex Sweet, Hort. Brit.* ed. I. 491, nomen (Quid?).—Austral.
Edwisana, *Tate, in Trans. Roy. Soc. S. Austral.* vii. (1884) 70.—Austral.
elliptica, *R. Br. in Trans. Linn. Soc.* x. (1810) 187.— Austral.
Epiglottis, *Labill. Nov. Holl. Pl.* i. 30. t. 40.— Austral.
erianthа, *R. Br. Prot. Nov.* 29.—Austral.
crinacea, *Meisn. in Lehm. Pl. Preiss.* i. 259.— Austral.
eucalyptoides, Meisn. l. c. 273 = laurina.
falcata, *R. Br. Prot. Nov.* 29.—Austral.
ferruginea, Lodd. Bot. Cab. t. 1301 = dactyloides.
ferruginea, *Sweet, Fl. Austral.* t. 45.—Austral.
flabelliifolia, *Meisn. in Hook. Kew Journ.* vii. (1855) 116.—Austral.
flexilis, R. Br. in Trans. Linn. Soc. x. (1810) 180 = nodosa.
flexilis, *F. Muell. in Linnæa*, xxvi. (1853–55) 359.— Austral.
floridа, *R. Br. in Trans. Linn. Soc.* x. (1810) 183.— Austral.
florulenta, Meisn. in Hook. Kew Journ. vii. (1855) 116 = saligna.
Francisiana, F. Muell. Fragm. i. 20 = multilineata?
Fraseri, *R. Br. Prot. Nov.* 28.—Austral.
gibbosa, *Cav. in Anal. Hist. Nat.* i. (1799) 214.— Austral.
gibbosa, Labill. ex Meisn. in DC. Prod. xiv. 403 = suaveolens.
Gilberti, Kipp. ex Meisn. in Hook. Kew Journ. vii. (1855) 115 = sulcata.
glabella, *R. Br. Prot. Nov.* 28.—Austral.
glabra, Schrad. Sert. Hannov. 27 = pugioniformis.
glauca, Knight, Prot. 106 = pugioniformis.
grammatophylla, F. Muell. Fragm. vi. 214 = multilineata.
heterophylla, Hook. Ic. Pl. t. 427 = varia.
Hookeriana, *Meisn. in Hook. Kew Journ.* iv. (1852) 308, nomen; *et in DC. Prod.* xiv. 412.—Austral.
ilicifolia, R. Br. in Trans. Linn. Soc. x. (1810) 184 = varia.
incrassata, *R. Br. Prot. Nov.* 29.—Austral.
intermedia, Hook. Ic. Pl. t. 445 = varia.
latriciata, *R. Br. Prot. Nov.* 27.—Austral.
Kippistiana, *Meisn. in Hook. Kew Journ.* vii. (1855) 115.—Austral.
laciniosa, F. Muell. Fragm. iv. 49 = ceratophylla.
lanigera, Tenore, Fl. Nap. i. 23. t. 6 = gibbosa.
lasiantha, *R. Br. Prot. Nov.* 29.—Austral.
lasiocarpha, R. Br. l. c. 27 = varia.
latifolia, F. Muell. Fragm. vi. 219 = adnata.
laurina, *R. Br. Prot. Nov.* 29.—Austral.
Lehmanniana, *Meisn. in Lehm. Pl. Preiss.* i. 257.— Austral.
Leucadendron, Meisn. l. c. 272 = incrassata.
leucocephala, D. Dietr. Syn. Pl. i. 581 = seq.
leucoptera, *R. Br. in Trans. Linn. Soc.* x. (1810) 180. —Austral.
ligustrina, Knight, Prot. 108 = oleifolia.
linearis, *R. Br. in Trans. Linn. Soc.* x. (1810) 183.— Austral.
linearifolia, *Hort. Petrop. ex Regel, Gartenfl.* (1857) 155.—Austral.?
litsocarpa, Regel, l. c. in obs. sphalm. = seq.
lissocarpha, *R. Br. Prot. Nov.* 27.—Austral.
lissosperma, R. Br. in Trans. Linn. Soc. x. (1810) 180 = acicularis.
longicuspis, Hook. in Mitch. Journ. Trop. Austral. 397 = leucoptera.
longifolia, A. Cunn. ex R. Br. Prot. Nov. 26 = Conninghamii.
longifolia, *Dum.-Cours. Bot. Cult.* ed. II. ii. 423.— Austral.
loranthifolia, *Meisn. in Lehm. Pl. Preiss.* i. 274.— Austral.
lorea, *R. Br. Prot. Nov.* 25.—Austral.
macrocarpa, *A. Cunn. ex R. Br. l. c.* 30.—Austral.
macrocarpa, Hort. Belved. ex Koem. & Schult. Syst. iii. Mant. 378 = microcarpa?
marginata, *R. Br. in Trans. Linn. Soc.* x. (1810) 185. —Austral.
megalosperma, *Meisn. in Hook. Kew Journ.* vii. (1855) 117.—Austral.
Meisneriana, *Kipp. ex Meisn. l. c.* 114.—Austral.
microcarpa, *R. Br. in Trans. Linn. Soc.* x. (1810) 182. —Austral.
Milligani, Meisn. in Hook. Kew Journ. iv. (1852) 307, nomen; et in DC. Prod. xiv. 395 = H. Epiglottis.
minimoides, A. Cunn. ex Meisn. in DC. Prod. xiv. 416 = saligna.
Mitchellii, Meisn. in Hook. Kew Journ. iv. (1852) 306, nomen; et in DC. Prod. xiv. 393 = microcarpa?
mixta, Lindl. Swan River App. 36 = trifurcata.
mollifolia, *Meisn. in Lehm. Pl. Preiss.* i. 261.— Austral.
myricophylla, Ait. ex G. Don, in Sweet, Hort. Brit. ed. III. 768 = ceratophylla.
myrtoides, *Meisn. in Lehm. Pl. Preiss.* i. 277.— Austral.
nervosa, Knight, Prot. 108 = dactyloides.
neurophylla, *Meisn. in Hook. Kew Journ.* vii. (1855) 117.—Austral.
nitida, *R. Br. in Trans. Linn. Soc.* x. (1810) 184.— Austral.
nodosa, *R. Br. l. c.* 179.—Austral.
nodosa, Meisn. in Lehm. Pl. Preiss. i. 255 = communis.
obliqua, *R. Br. in Trans. Linn. Soc.* x. (1810) 180.— Austral.
obliqua, Lodd. Bot. Cab. t. 1689 = acicularis.
obtusa, *Meisn. in Hook. Kew Journ.* iv. (1852) 309, nomen; *et in DC. Prod.* xiv. 411.—Austral.
Oldfieldii, *Benth. Fl. Australe.* v. 530.—Austral.
oleifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 185.— Austral.
orthorrhyncha, *F. Muell. Fragm.* vi. 214.—Austral.
pachyphylla, Sieber, ex Spreng. Syst. iv. Cur. Post. 46 = propinqua.
Pampliniana, *Kipp. ex Meisn. in Hook. Kew Journ.* vii. (1855) 115.—Austral.
pandanicarpa, *R. Br. Prot. Nov.* 29.—Austral.
parilis, Knight, Prot. 106 = pugioniformis.
patula, R. Br. Prot. Nov. 27 = microcarpa.
pectinata, Dum.-Cours. Bot. Cult. ed. II. ii. 424; Colla, in Mem. Acc. Tor. xxxl. (1827) 320 = suaveolens.
pedunculata, *F. Muell. in Melb. Chemist* (July 1888). —Austral.
petiolaris, *Meisn. in Lehm. Pl. Preiss.* i. 277.— Austral.
petrophiloides, Hort. ex Meisn. in DC. Prod. xiv. 423 = lissocarpha.
petrophioloides, Mackay, ex Sweet, Hort. Brit. ed. I. 491, nomen (= præc.?).—Austral.
pilulifera, Lindl. Swan River App. 36 = Grevillea oxystigma.
pinifolia, Knight, Prot. 107 = gibbosa.
pinnata, Koem. & Schult. Syst. iii. 444 = suaveolens.
piriformis, Anct. vide *pyriformis*.
platysperma, *Meisn. in DC. Prod.* xiv. 413.—Austral.
plurinervia, *F. Muell. ex Benth. Fl. Australe.* v. 523.— Austral.
Preissii, *Meisn. in Lehm. Pl. Preiss.* i. 257.—Austral.
propinqua, *A. Cunn. in Field, N. S. Wales*, 337.— Austral.
prostrata, R. Br. in Trans. Linn. Soc. x. (1810) 184 = glabella.
pubescens, Hort. Cels, ex Schult. Mant. iii. 291 = Banksia integrifolia.
pubescens, Schrad. Sert. Hannov. 27, in obs. = gibbosa.
pugioniformis, *Cav. in Anal. Hist. Nat.* i. (1799) 213. —Austral.
purpurea, *Hook. in Mitch. Journ. Trop. Australe.* 348. —Austral.
pycnobotrya, F. Muell. Fragm. v. 72 = nitida.
pycnoneura, *Meisn. in Hook. Kew Journ.* vii. (1855) 117.—Austral.
pyriformis, Cav. in Anal. Hist. Nat. i. (1799) 217 = Xylomelum pyriforme.
recurva, *Meisn. in Hook. Kew Journ.* iv. (1852) 307, nomen; *et in DC. Prod.* xiv. 394.—Austral.
repanda, R. Br. Prot. Nov. 30 = ferruginea.
rhombalis, *F. Muell. Fragm.* x. 90.—Austral.
Roei, *Benth. Fl. Australe.* v. 499.—Austral.
rostrata, *F. Muell. First Gen. Rep.* 17; *et ex Meisn. in Linnæa*, xxvi. (1853–55) 359.—Austral.
rubricaulis, Cels, ex Colla, Hort. Ripul. 63; App. i. 114 = Stenocarpus saligna.
rugosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 179.— Austral.
rusciifolia, *Labill. Nov. Holl. Pl.* i. 30. t. 39.—Austral.
ruscifolia, Steud. Nom. ed. II. i. 403, sphalm. in syn. = præc.
salicifolia, Sweet, ex Meisn. in DC. Prod. xiv. 416 = saligna.
saligna, *Knight, Prot.* 108; *R. Br. in Trans. Linn. Soc.* x. (1810) 185.—Austral.
scoparia, Meisn. in Lehm. Pl. Preiss. i. 256 = sulcata.
semiglauca, F. Muell. First Gen. Rep. 17; *et ex Meisn. in Linnæa*, xxvi. (1853–55) 359 = nodosa.
serices, *Hort. Ups. ex Regel, Suppl. Ind. Sem. Hort. Petrop.* 1866 (1867) 29.—Austral.
serices, Schrad. Sert. Hannov. 27, in obs. = acicularis.
smilacifolia, *Meisn. in Lehm. Pl. Preiss.* i. 267.— Austral.
stenocarpa, *R. Br. Prot. Nov.* 29.—Austral.
stenocarpoides, *F. Muell. ex Benth. Fl. Australe.* v. 511.—Austral.
stenophylla, Bast. ex Meisn. in DC. Prod. xiv. 416 = ulicina.
stenophylla, *A. Cunn. ex R. Br. Prot. Nov.* 30.— Austral.
striata, F. Muell. ex Meisn. in Linnæa, xxvi. (1853–55) 360 = leucoptera.

HAKEA :—
strumosa, *Meisn. in Hook. Kew Journ.* iv. (1852) 208, nomen; *et in DC. Prod.* xiv. 406.—Austral.
suaveolens, *R. Br. in Trans. Linn. Soc.* x. (1810) 189. —Austral.
subsulcata, *Meisn. in Lehm. Pl. Preiss.* i. 555.— Austral.
subulata, *Hort. Mackay, ex Sweet, Hort. Brit.* ed. I. 491; *A. Cunn. ex Steud. Nom.* ed. II. i. 719, nomen. —Austral.
sulcata, *R. Br. in Trans. Linn. Soc.* x. (1810) 130.— Austral.
suicata, Meisn. in Lehm. Pl. Preiss. ii. 260 = Metaneriana.
tenuiflora, *Dum-Cours. Bot. Cult.* ed. I. v. 106; *Roem. & Schult. Syst.* iii. 425.—Austral.
tephrosperma, R. Br. Prot. Nov. 26 = leucoptera.
tephrosperma, Meisn. in Hook. Kew Journ. iv. (1852) 207 = Kippistiana.
tricostata, Hook. ic. Pl. t. 435 = cinerea.
tricrura, Lindl. Swan River App. 36 = trifurcata.
trifornis, Lindl. l. c. = amplexicaulis.
trifurcata, *R. Br. in Trans. Linn. Soc.* x. (1810) 133.— Austral.
trinervis, *Meisn. in Hook. Kew Journ.* iv. (1852) 209, nomen; *et in DC. Prod.* xiv. 414.—Austral.
trinervis, *F. Muell. Fragm.* iii. 146.—Austral.
tuberculata, R. Br. Prot. Nov. 28 = varia.
ulicina, *R. Br. l. c.* 29.—Austral.
undulata, *R. Br. in Trans. Linn. Soc.* x. (1810) 135.— Austral.
varia, *R. Br. l. c.* 133.—Austral.
verrucosa, *F. Muell. Fragm.* v. 25; vi. 218.—Austral.
Victoria, Drumm. in Bot. Mag. lxxiv. Comp. 3 = cucullata.
virgata, R. Br. Prot. Nov. 26 = leucoptera.
vittata, *R. Br. in Trans. Linn. Soc.* x. (1810) 132.— Austral.

HALANTHIUM, C. Koch, in Linnæa, xvii. (1843) 813. *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 74.
GAMANTHUS, Bunge, Anabas. Rev. 76 (1852).
PHYSOGYTON, Jaub. & Spach, Illustr. i. t. 135 (1846). Abichii, *Bunge, Pl. Abich.* 18.—Reg. Caucas.
Auckeri, Moq. in DC. Prod. xiii. ii. 204 = rarifolium.
Kulpianum, *Bunge, Anabas. Rev.* 80.—Armenia.
lanatum, Bunge, Pl. Abich. 18 = rarifolium.
lilacinum, Bunge, ex Boiss. Fl. Orient. iv. 982 = rarifolium.
mamamense, *Bunge, Anabas. Rev.* 82.—Persia.
molle, Bunge, ex Boiss. Fl. Orient. iv. 981 = mamamense.
purpureum, *Bunge, Anabas. Rev.* 85.—Persia.
rarifolium, *C. Koch, in Linnæa*, xvii. (1843) 814.— Persia.
rigidum, Bunge, ex Boiss. Fl. Orient. iv. 981 = mamamense.
robustum, Bunge, ex Boiss. l. c. 983 = rarifolium.
spicatum, Bunge, ex Boiss. l. c. = rarifolium.

HALARCHON, Bunge, Anabas. Rev. 75. t. 1 (1852). *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 76.
vesiculosum, *Bunge, l. c.*—Afghan.

HALEA, Torr. & Gray, Fl. N. Am. ii. 304 (1841) = **Tetragonotheca**, Linn. (Compos.).
ludoviciana, Torr. & Gray, l. c. = T. ludoviciana.
repanda, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 458 = T. ludoviciana.
texana, A. Gray, Pl. Fendl. 83 = T. texana.

HALECUS, Rumph. ex Rafin. Sylva Tellur. 62 (1838) = **Croton**, Linn. (Euphorbiac.).
amaritimensis, Rafin. l. c. = C. mauritianus.
perus, Rafin. l. c. = C. Tiglium †

HALENEA, Wight, Ic. iv. ii. 8 (1848) err. typ. = seq.

HALENIA, Borckh. in Roem. Arch. i. t. 25 (1796) *GENTIANEÆ*, Benth. & Hook. f. ii. 817.
EBALENIUS, Griseb. Obs. Gentian. 36 (1836).
TETRAGONANTHUS, S. G. Gmel. Fl. Sibir. iv. 118 (1769).
alata, *Hemsl. Biol. Centr. Am. Bot.* ii. 351.—Mexic.
aphyllata, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 371 = decumbens.
asclepiadea, Griseb. Gen. et Sp. Gent. 326 = Pavoniana.
Brachyodonea, Griseb. l. c. 325 = deflexa.
brevicornis, *G. Don, Gen. Syst.* iv. 177.—Ecuador.
decumbens, *Benth. Pl. Hartw.* 67.—Mexic.
deflexa, *Griseb. Gen. et Sp. Gent.* 324.—Am. bor. Dombeyana, *Wedd. Chlor. And.* ii. 76.—Peruv.
elata, *Wedd. l. c.* 78.—Venezuela.
elliptica, *Don, in Trans. Linn. Soc.* xvii. (1837-539.—Reg. Himal.
elongata, *D. Don, ex G. Don, Gen. Syst.* iv. 177.— Nepal.
Fischerii, R. Grah. in Edinb. N. Phil. Journ. (Apr.-June 1830) 174 = sibirica.
gentianoides, *Wedd. Chlor. And.* ii. 78.—N. Granat.
gracilis, *Griseb. Gen. et Sp. Gent.* 327.—Am. austr.
hexantha, Griseb. l. c. 325 = deflexa.
hypericoides, *Griseb. l. c.* 328.—N. Granat.
inæqualis, *Wedd. Chlor. And.* ii. 78.—Venezuela.
longicornu, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 370.—Mexic.
major, *Wedd. Chlor. And.* ii. 79.—N. Granat.
Michauxiana, G. Don, Gen. Syst. iv. 177 = deflexa.
multiflora, *Benth. Pl. Hartw.* 24.—Mexic.
nudicaulis, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 371.—Mexic.
nutans, *Mart. & Gal. l. c.*—Mexic.
parviflora, *G. Don, Gen. Syst.* iv. 177.—Mexic.
pauciflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 352.— Mexic.
Pavoniana, *G. Don, Gen. Syst.* iv. 177.—Am. austr.
Perrottetii, *Griseb. in DC. Prod.* ix. 129.—Ind. or.
pinifolia, *G. Don, Gen. Syst.* iv. 177.—Peruv.
plantaginea, *Griseb. Gen. et Sp. Gent.* 327.—Mexic.
Purdieana, *Wedd. Chlor. And.* ii. 76.—N. Granat.
Rothrockii, *A. Gray, in Proc. Am. Acad.* xi. (1876) 84.—Arizona.
Schiedeana, *Griseb. Gen. et Sp. Gent.* 327.—Mexic.
sibirica, *Borckh. in Roem. Arch.* i. t. 25.—Sibir.

HALESIA, P. Br. Hist. Jamaic. 205 (1756) = **Guettarda**, Linn. (Rubiac.).

HALESIA, Linn. Syst. ed. X. 1044 (1759). *STYRACEÆ*, Benth. & Hook. f. ii. 669.
PTEROSTYRAX, Sieb. & Zucc. Fl. Jap. i. 94. t. 47 (1835).
carolina, Linn. Syst. ed. X. 1044 = tetraptera.
corymbosa, *Nichols. Dict. Gard.* ii. 109.—Japon.
diptera, *Linn. Sp. Pl.* ed. II. 636.—Am. bor.
hispida, *[Mast. in] Gard. Chron.* (1884) ii. 176. f. 34. —Japon.
parviflora, Lindl. in Bot. Reg. t. 952 = Styrax americanum †
parviflora, Michx. Fl. Bor. Am. ii. 40 = tetraptera.
reticulata, Buckl. in Proc. Acad. Sc. Philad. 1860 (1861) 444 = diptera.
stenocarpa, *C. Koch, in C. Koch & Fint. Wochenschr.* (1858) 191.—Am. bor.
ternata, Blanco, Fl. Filip. ed. I. 299 = Illigera dubia.
tetraptera, *Linn. Sp. Pl.* ed. II. 636.—Am. bor.

HALFORDIA, F. Muell. Fragm. v. 43 (1865). *RUTACEÆ*, Benth. & Hook. f. i. 992.
drupifera, *F. Muell. l. c.*—Austral.
sclerotyla, *F. Muell. l. c.* vii. 142.—Austral.

HALGANIA, Gaudich. in Freyc. Voy. Bot. 448. t. 59 (1826). *BORAGINEÆ*, Benth. & Hook. f. ii. 842.
anagalloides, *Endl. in Ann. Wien. Mus.* ii. (1838) 204. —Austral.
andromedoides, Behr. & F. Muell. ex F. Muell. Fragm. i. 209 = lavandulacea.
Behrens, Oldfield & F. Muell. ex F. Muell. l. c. = littoralis.
corymbosa, *Lindl. Swan River App.* 40.—Austral.
cyanea, Lindl. l. c. = strigosa.
integerrima, *Endl. in Ann. Wien. Mus.* ii. (1838) 205. —Austral.
lavandulacea, *Endl. l. c.*—Austral.
Lehmanniana, Sond. in Lehm. Pl. Preiss. ii. 238 = Preissiana.
littoralis, *Gaudich. in Freyc. Voy. Bot.* 449. t. 59.— Austral.
Preissiana, *Lehm. Pl. Preiss.* i. 347.—Austral.
sericiflora, *Benth. Fl. Australiæ*, iv. 408.—Austral.
solanacea, *F. Muell. in Hook. Kew Journ.* ix. (1857) 21.—Austral.
strigosa, *Schlecht. in Linnæa*, xx. (1847) 614.— Austral.
tuberculosa, Schlecht. l. c. 615 = strigosa.

HALIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 175 = **Halesia**, Linn. (Styrac.).

HALIANTHUS, Fries, Fl. Halland. 75 (1817) = **Aronaria**, Rupp. (Caryophyll.).
peploides, Fries, l. c. = A. peploides.

HALIBREXIA, Phil. in Linnaea. xxxiii. (1864-65) 208 = **ALIBREXIA**, Miers = **Dolia**, Lindl. (Convolv.).

HALIMIUM, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 365 = **Helianthemum**, Toorn. (Cistin.).
alyssoides, Lamotte, Prod. 113 = Helianth. guttatum.
atriplicifolium, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 366 = Helianth. atriplicifolium.
Clusii, Spach, l. c. 365 = Cistus Clusii.
eriocephalum, Willk. ic. et Desc. Pl. ii. 62. t. 105 = Helianth. eriocephalum.
formosum, Willk. l. c. 59. t. 101 = Helianth. formosum.
halimifolium, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 717 = Helianth. halimifolium.
heterophyllum, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 366 = Helianth. ocymoides.
hirsutissimum, Willk. ic. et Desc. Pl. ii. 64. t. 106 = Helianth. hirsutissimum.
lasianthum, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 366 = Helianth. alyssoides
lepidotum, Spach, l. c. = Helianth. halimifolium.
Libanotis, Lange, in Kjoeb. Vidensk. Meddel. (1865) 90 (Pugill. 285) = Helianth. Libanotis.
multiflorum, Willk. ic. et Desc. Pl. ii. 67. t. 108 = Helianth. halimifolium †
occidentale, Willk. l. c. 59. t. 103 = Helianth. occidentale.
ocymoides, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 715 = Helianth. ocymoides.
racemiferum, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 366 = Helianth. Libanotis.
umbellatum, Spach, l. c. = Helianth. umbellatum.

HALIMOCNEMIS, C. A. Mey. in Ledeb. Fl. Alt. i. 381 (1829). *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 74.
HALIOTIS, Bunge, Anabas. Rev. 73. t. 1 (1852).
alternifolia, Moq. in DC. Prod. xiii. ii. 198 = Petrosimonia crassifolia.
Belangeri, Moq. l. c. = Heliantiium rarifolium.
brachiata, C. A. Mey. in Ledeb. Fl. Alt. i. 381 = Petrosimonia brachiata.
caspica, Moq. Chenop. Enum. 156 = Nanophytum caspicum.
crassifolia, C. A. Mey. in Ledeb. Fl. Alt. i. 385 = Petrosimonia crassifolia.
gamacrpa, Moq. Chenop. Enum. 155 (= *Gamanthus gamacrpa*).—Persia.
glauca, C. A. Mey. in Ledeb. Fl. Alt. i. 385 = Petrosimonia crassifolia.
hirsuta, C. Koch, in Linnæa, xvii. (1849) 192 = malacophylla.
hispida, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 260 = Halocharis hispida.
Hohenacheri, Presl, Bot. Bemerk. 109 = Petrosimonia brachiata.
juniperina, C. A. Mey. in Ledeb. Fl. Alt. i. 386 = Nanophyton caspicum.
Karelini, *Moq. in DC. Prod.* xiii. ii. 196.—Reg. Casp.; Soongar.
Kirilowii, Fenzl, ex Ledeb. Fl. Ross. iii. 844 = Salsola affinis.
Komarowyi, Stchegl. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 328. t. 5. f. 5 = Halanthium rarifolium.
Kulpiana, C. Koch, in Linnæa, xvii. (1843) 813 = Halanthium Kulpianum.
longifolia, *Bunge, in Act. Hort. Petrop.* v. (1877) 643 —Turkest.
macrantha, Bunge, in Ledeb. Fl. Ross. iii. 843 = macranthera.
macrantha, Kar. in Bull. Soc. Nat. Mosc. (1839) 168 = Nanophytum caspicum.
macranthera, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 478.—Turkestan.
malacophylla, Kar. & Kir. in Bull. Soc. Nat. Mosc. xiv. (1841) 722 = Salsola affinis.
malacophylla, *C. A. Mey. in Ledeb. Fl. Alt.* i. 387.— Reg. Caucas.; Sibir.
microphylla, Turcz. ex Ledeb. Fl. Ross. iii. 816 = Salsola gemmascens.
mollissima, *Bunge, Anabas. Rev.* 71.—Persia.

HALIMOCNEMIS :—
monandra, C. A. Mey. in Ledeb. Fl. Alt. i. 384 = Petrosimonia monandra.
obtusifolia, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 79 = Halogeton obtusifolius.
occidentalis, Nutt. ex Moq. in DC. Prod. xiii. 11. 379, 109 = Nitrophila occidentalis.
oppositifolia, Elchw. Casp. 154 = Petrosimonia crassifolia.
pilifera, Moq. in DC. Prod. xiii 11. 195 = pilosa.
pilosa, *Moq. Chenop. Enum.* 152.—Persia.
pubescens, Ledeb. ex Sweet, Hort. Brit. ed. III. 376 (Quid ?).—Sibir.
purpurea, Moq. Chenop. Enum. 153 = Halanthium purpureum.
pycnantha, C. Koch, in Linnaea, xxii. (1849) 193 = Halanthium Kulpianum.
sclerantha, Schrenk, ex Ledeb. Fl. Ross. iii. 846 = Karelinia.
sclerosperma, *C. A. Mey. in Ledeb. Fl. Alt* i. 387.—Tauria; As. med.
sibirica, C. A. Mey. l. c. 382 = Petrosimonia sibirica.
Smirnowii, *Bunge, in Act. Hort. Petrop.* v. (1877) 644 —Turkestan.
spicata, Presl, Bot. Bemerk. 109 (Quid ?).—Aegypt.
squarrosa, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 360 = Petrosimonia squarrosa.
sulphurea, Moq. Chenop. Enum. 152 = Halocharis sulphurea.
tomentosa, Moq. l. c. 151 = Salsola canescens, Nosea Tournefortii.
triandra, Moq. in DC. Prod. xiii. 11. 197 = Petrosimonia Volvox.
villosa, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 474 = pilosa.
villosa, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 484. —As. med.
Volvox, C. A. Mey. in Ledeb. Fl. Alt. i. 383 = Petrosimonia Volvox.

HALIMOCNEMUM, Lindem. in Bull. Soc. Nat. Mosc. lv. (1880) 11. 182 = **Haloxemum**, M. Bieb. (Chenop.).

HALIMODENDRON, Fisch. ex DC. Mém. Lég. 288 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 602.
HALODENDRON, DC. Prod. ii. 269 (1825)
argenteum, *Fisch. in DC. Prod.* ii. 269.—Reg. Caucas.; Persia; As. bor.
caspiadatum, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xviii. (1842) 239 = argenteum.
emarginatum, Jaub. & Spach, l. c. = argenteum.
speciosum, Carr. in Rev. Hortic. (1876) 30 = argenteum.
subviscosum, G. Don, Gen. Syst. ii. 244 = argenteum.
triflorum, Link, Handb. ii. 201 = argenteum.

HALIMOLOBOS, Tausch, in Flora, xix. (1836) 410. *CRUCIFERAE*.
patula, *Tausch, l. c.*—Hab. ?
stricta, Tausch, l. c. = Arabis lasioloba.

HALIMUS, Linn. Syst. ed. I (1735) = **Atriplex**, Linn. (Chenop.).
australis, T. Nees, Gen. Pl. Monochl. n. 64, in nota = A. portulacoides.
borealis, T. Nees, l. c. = A. portulacoides.
pedunculatus, Wallr. Sched. Crit. 117 = A. pedunculata.
portulacoides, Dum. Fl. Belg. 20 = A. portulacoides.
portulacoides, Schur, Enum. Pl. Transs. 578 = A. pedunculata.

HALLACKIA, Harv. Thes. Cap. ii. 2. t. 102 (1863) = **Huttonaea**, Harv. (Orchid.).
fimbriata, Harv. l. c. = H. fimbriata.

HALLERA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 171 = seq.

HALLERIA, Linn. Coroll. Gen. 10 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 986.
abyssinica, *Vaub. & Spach, Illustr.* v. 66. t. 459, 460. —Abyssin.
elliptica, *Linn. Sp. Pl.* 625.—Afr. austr.
elliptica, Thunb. Prod. Pl. Cap. 98 = lucida.
ligustrifolia, *Baker, in Journ. Linn. Soc.* xvIII. (1881) 278.—Madag.

HALLERIA:—
lucida, *Linn. Sp. Pl.* 625.—Afr. austr. et trop.
ovata, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 54. —Afr. austr.
tetragona, *Baker, in Journ. Linn. Soc.* xx. (1883) 214. —Madag.

HALLESIA, Scop. Introd. 141 (1777) = HALESIA, P. Br. = **Guettarda**, Linn. (Rubiac.).

HALLIA, Jaume St. Hil. in Bull. Soc. Philom. (Dec. 1813) 193 = **Alysicarpus**, Neck. (Legumin.).
buplevrifolia, Jaume St. Hil. l. c. = A. buplevrifolius.
glumacea, Wall. ex Wight & Arn. Prod. 284 = A. glumaceus.
salicifolia, Jaume St. Hil. in Bull. Soc. Philom. (Dec. 1813) 195 = A. longifolius ?
vaginalis, Jaume St. Hil. l. c. = A. vaginalis.

HALLIA, Thunb. in Schrad. Journ. i. (1799) 818. *LEGUMINOSAE*, Benth. & Hook. f. i. 528.
alata, *Thunb. l. c.* 819.—Afr. austr.
angustifolia, DC. Prod. ii. 128 = virgata.
argentea, Spreng. Syst. iii. 195 = Ammodendrum Sieversii.
Asarina, *Thunb. in Schrad. Journ.* i. (1799) 822.—Afr. austr.
Bojeriana, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 482. —Madag.
connexa, Burch. ex Harv. & Sond. Fl. Cap. ii. 232 = Asarina.
coridata, *Thunb. in Schrad. Journ.* i. (1799) 821.—Afr. austr.
filiformis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 232.—Afr. austr.
flaccida, *Thunb. in Schrad. Journ.* i. (1799) 819.—Afr. austr.
glauca, Zipp. ex Span. in Linnaea, xv. (1841) 190 = Indigofera albicans.
hirta, Willd. Sp. Pl. iii. 1169 = Heylandia latebrosa.
imbricata, *Thunb. in Schrad. Journ.* i. (1799) 822.—Afr. austr.
juncea, Otto, ex Steud. Nom. ed. II. i. 720 = Lespedeza juncea.
linearis, Buch.-Ham. ex Wall. Cat. n. 5752 = Flemingia lineata.
monophylla, Desv. in DC. Prod. ii. 128 = Heylandia latebrosa.
rurcifolia, Bot. Mag. ex Steud. Nom. ed. II. i. 720 = imbricata ?
sagittaria, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 408 = Desmodium sagittatum.
Sororia, Willd. Sp. Pl. iii. 1170 = Eleiotis Sororia.
trifoliata, Roth, Nov. Pl. Sp. 362 = Rhynchosia trifoliata.
virgata, *Thunb. in Schrad. Journ.* i. (1799) 820.—Afr. austr.

HALMIA, M. Roen. Syn. Rosifl. 101, 184 (1847) = **Crataegus**, Tourn. (Rosac.).
cornifolia, M. Roem. l. c. 185 = C. tomentosa.
flabellata, M. Roem. l. c. 186 = C. coccinea.
lobata, M. Roem. l. c. = C. grandiflora, *C. Koch.*
punctata, M. Roem. l. c. 184 = C. tomentosa.
tomentosa, M. Roem. l. c. 185 = C. tomentosa.

HALMYRA, Herb. Amaryll. 202 (1837) = **Panoratium**, Linn. (Amaryllid.).
stellaris, Parl. Nov. Gen. Monocot. 28 = P. illyricum.

HALOCHARIS, Bieb. ex DC. Prod. vi. 665 (1837) = **Centaurea**, Linn. (Compos.).
altaica, Fisch. ex Stend. Nom. ed. II. i. 720 = Leuzea salina.
carthamoides, Bieb. ex DC. Prod. vi. 666 = C. carthamoides.

HALOCHARIS, Moq. in DC. Prod. xiii. 11. 201 (1849). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 75.
Abichii, C. Koch, in Linnaea, xxiv. (1851) 93 = Halanthium reticulum.
clavata, *Bunge, Anabas. Rev.* 64.—Afghan.
gamocarpus, Moq. in DC. Prod. xiii. 11. 202 (= *Gamanthus gamocarpus*).—Persia; Afghan.
hispida, *Bunge, Anabas. Rev.* 62.—Orenb.
Kulpiana, C. Koch, in Linnaea, xxiv. (1851) 92, in obs. = Halanthium Kulpianum.

HALOCHARIS:—
pycnantha, C. Koch, in Linnaea, xxiv. (1851) 91 = Halanthium reticulum.
sulphurea, *Moq. in DC. Prod.* xiii. 11. 201.—Orient.
vesiculosa, Moq. l. c. 202 = Halarchon vesiculosus.
violacea, *Bunge, Anabas. Rev.* 63.—Afghan.

HALOCHLOA, Griseb. in Goett. Abh. xxiv. (1879) 285. *GRAMINEAE*.—Cf. Benth. & Hook. f. Gen. iii. 1096.
scoroas, *Griseb. l. c.*—Reg. Argent.

HALOCNEMON, Spreng. Syst. i. 19 (1825) = seq.

HALOCNEMUM, Bieb. Fl. Taur. Cauc. iii. 3 (1819). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 64.
HALINOCNEMUM, Lindem. in Bull. Soc. Nat. Mosc. lv. (1880) 11. 182.
HALOCNEMON, Spreng. Syst. i. 19 (1825).
americanum, Gill. ex Moq. in DC. Prod. xiii. 11. 150 = Heterostachys Ritteriana.
amplexicaule, Tineo, ex Ung.-Sternb. in Atti Congr. Bot. Firenze, 1874 (1876) 326 = Halopeplis amplexicaulis.
arabicum, Spreng. Syst. i. 19 = Kalidium caspicum.
australasicum, Moq. Chenop. Enum. 110 = Salicornia australis.
caspicum, Bieb. Fl. Taur. Cauc. iii. 3 = Halostachys caspica.
caspicum, Tausch, in Flora, xv. (1832) 737 = Kalidium caspicum.
cinereum, F. Muell. Fragm. i. 140 = Salicornia cinerea.
cruciatum, *Tod. Ind. Sem. Hort. Panorm.* (1872) 40; *et in Nuov. Giorn. Bot.* (1878) 159.—Ins.
foliatum, Spreng. Syst. i. 19 = Kalidium foliatum.
fruticosum, *D. Dietr. Sp. Pl.* i. 88; *Link. Handb.* ii. 414.—Europ. austr.; Afr. bor.
glaucum, Presl, Bot. Bemerk. 109 = Salicornia fruticosa.
monandrum, Georgi, Beschr. Russ. Reich. iv. 664 = Ofaiston pauciflorum.
moniliforme, Tausch, in Flora, xv. (1832) 738, in obs. = Halopeplis pygmaea.
nodulosum, Spreng. Syst. i. 19 = Halopeplis amplexicaulis.
Ritterianum, Moq. Chenop. Enum. 109 = Heterostachys Ritterianus.
strobilaceum, *Birb. Fl. Taur. Cauc.* iii. 3.—Taur. Casp.
strobilaceum, Moris, Fl. Sard. iii. 368 = cruciatum.

HALODENDRON, DC. Prod. ii. 269 (1825) = **Halimodendron**, Fisch. (Legumin.).

HALODENDRON, Roem. & Schult. Syst. iii. 485 (1818) = HALODENDRUM, Thou. = **Aviocennia**, Linn. (Verben.).
Thouarsii, Roem. & Schult. l. c. = A. officinalis.

HALODENDRUM, Thou. Gen. Nov. Madag. 8 (1806) = praec.

HALODULA, Benth. & Hook. f. Gen. iii. 1018 (1883). sphalm. = seq.

HALODULE, Endl. Gen. Suppl. i. 1368 (1841). *NAIADACEAE*, Benth. & Hook. f. iii. 1019.
australis, *Miq. Fl. Ind. Bat.* iii. 237.—Mare rubr.; Malaya.
emarginata, Boiss. Fl. Orient. v. 24 = australis.
Wrightii, *Aschers. in Sitzb. Ges. Naturf. Fr. Berl.* (1868) 19.—Cuba.

HALOGETON, C. A. Mey. in Ledeb. Fl. Alt. i. 378 (1829). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 75.
AGATHOPHORA, Bunge, in Mém. Acad. Pétersb. Sér. VII. iv. (1862) n. 11 (Anabas. Rev.) 92.
MICROPEPLIS, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 474 (Rel. Lehm. 286).
acutiflorus, *Bunge, in Mém. Soc. Etr. Pétersb.* vii. (1847) 427.—Silesia.
alopecuroides, *Moq. Chenop. Enum.* 161.—Aegypt.; Arab.
anachnoides, *Moq. in DC. Prod.* xiii. 11. 205.—Siberia.
articulatus, *Bail. in Journ. Linn. Soc. xvi.* (1878) 647. —Hisp.; Maroco.; Aegypt.
Ballotoides, *Aschers. in Journ. Linn. Soc. xiv.* (1874) 317, nomen.—Bermuda.

HALOGETON:—
cinerosa, Moq. Chenop. Enum. 159 = Seidlitzia florida.
georgiana, Moq. in DC. Prod. xiii. ii. 205 = Salsola verrucosa.
glomerata. *C. A. Mey. in Ledeb. Fl. Alt.* i. 378.—Reg. Casp.: As. bor.
Gmelinianus, Bal. ex Boiss. Fl. Orient. iv. 954 = Salsola spissa.
malacophylla, *Bunge, Lehm. Reliq. Bot.* (1847) 302.—Turkestan.
monandrus, C. A. Mey. in Ledeb. Fl. Alt. i. 397 = Ofaiston pauciflorum.
obtusifolius, *Fenzl, ex Ledeb. Fl. Ross.* iii. 684.—Sibir. alt.
Olivieri, Moq. Chenop. Enum. 159 = Salsola spissa.
oppositifolius, *C. A. Mey. in Ledeb. Fl. Alt.* i. 378.—Rossia: Sibir.
oppositifolius, Fenzl, in Ledeb. Fl. Ross. iii. 833 = Salsola brachiata.
paronychioides, *Less. in Linnaea*, ix. (1834) 198.—Reg. Casp.
purpureus, Fenzl, in Ledeb. Fl. Ross. iii. 833, in nota = Halanthium purpureum.
rariflorus, Fenzl, l. c. 834 = Halanthium rariflorum.
sativus, *Moq. Chenop. Enum.* 158.—Algeria.
spinosissimus, C. A. Mey. Verz. Pfl. Cauc. 159 = Noaea spinosissima.
tamariscifolia, C. A. Mey. in Ledeb. Fl. Alt. i. 378 = Salsola tamariscifolia.
tetragonus, Moq Chenop. Enum. 159 = Salsola tetragona.
tetrandrus, Moq. l. c. 160 = Salsola tetragona.
tibeticus, Bunge, Anabas. Rev. 94 = glomeratus.
Tournefortii, *Jaub. & Spach, Illustr.* ii. 43. t. 132.—Persia.

HALOLACHNA, Endl. Gen. 1087 (1840) = Hololachna, Ehrenb. (Tamariscin.).

HALOPEPLIS, Bunge, in Linnaea, xxviii. (1856) 678. *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 64. amplexicaulis, *Ung.-Sternb. ex Cesati, Passer. & Gibelli, Comp. Fl. Ital.* 271.—Reg. Mediterr.
Gillesii, *Griseb. in Goett. Abh.* xxiv. (1879) 88.—Reg. Argent.
nodulosa, Bunge, in Linnæa, xxviii. (1856) 678 = amplexicaulis.
patagonica, Ung.-Sternb. in Atti Congr. Bot. Firenze, 1874 (1876) 280 = Spirostachys patagonica.
perfoliata, *Bunge, ex Schweinf. & Aschers. in Fl. Aethiop.* 289, nomen; *et ex Ung.-Sternb. l. c.* 229.—Aegypt.: Arab.
pygmæa, *Bunge, ex Ung.-Sternb. Syst. Salic.* 105.—Ross.: Soongar.
Ritteriana, Bunge, ex Ung.-Sternb. l. c. 100 = Heterostachys Ritteriana.
songarica, Bunge, ex Ung.-Sternb. l. c. 105 = pygmæa
strobilacea, Cesati, Passer. & Gibelli, Comp. Fl. Ital. 271 = Halocnemum strobilaceum.

HALOPETALUM, Stend. Nom. ed. II. i. 188, in syn. (1840) = **Banisteria**, Linn. (Malpig.).

HALOPHILA, Thou. Gen. Nov. Madag. 2 (1806). *HYDROCHARIDEÆ*, Benth. & Hook. f. iii. 455.
BARKANIA, Ehrenb. & Hempr. in Abh. Acad. Berl. (1823) 429.
Baillonii, *Aschers. in Neumayer, Anleit. Wissensch. Beob.* 367.—Ind. occ.
Beccarii, *Aschers. in Nuov. Giorn. Bot.* iii. (1871) 302. —Borneo.
Engelmanni, *Aschers. in Neumayer, Anleit. Wissensch. Beob.* 368.—Florida.
Kotschyana, Fenzl, ex Aschers. in Linnæa, xxxv. (1867–68) 173 = ovata.
Lennopsis, Miq. Fl. Ind. Bat. iii. 280 = ovata.
madagascariensis, Steud. Nom. ed. II. f. 720 = ovata.
major, Miq. Fl. Ind. Bat. iii. 280 = ovata.
ovalis, Hook. f. Fl. Tasm. ii. 48 = sq.
ovata, *Gaudich. in Freyc. Voy. Bot.* 480. t. 40. f. 1.—As. trop.; Austral.
spinulosa, *Aschers. in Neumayer, Anleit. Wissensch. Beob.* 368; *Benth. Fl. Australi.* vii. 183.—Austral.
stipulacea, *Aschers. in Sitzb. Ges. Naturf. Fr. Berl.* (1867) 8.—Geront. trop.

HALORAGIDÆ, Fourt. Char. Gen. 81. t. 81 (1776). *HALORAGEÆ*, Benth. & Hook. f. i. 674.
CERCODIA, Murr. in Comm. Gotting. iii. (1780) 1. t. 1.

HALORAGIS:—
GONATOCARPUS, Schreb. Gen. i. 86 (1789).
GONOCARPUS, Thunb. Nov. Gen. iii. 85 (1788).
LINOCIRIA, Neck. Elem. iii. 866 (1790).
acanthocarpa, *Brongn. in Duperr. Voy. Coq. Bot.* t. 70.—Austral.
aculeolata, *Benth. Fl. Australi.* ii. 477.—Austral.
acutangula, *F. Muell. in Trans. Vict. Inst.* i. (1855) 125.—Austral.
aggregata, *T. Buch. in Trans. N. Z. Inst.* iv. (1872) 224.—N. Zel.
alata, *Sacp. Mise.* ii. 339; *Jc. Pl. Rar.* i. 7. t. 69.—Austral.; N. Zel.
alternifolia, Walp. Rep. ii. 99 = alata.
aspera, Lindl. in Mitch. Journ. Trop. Austral. 306 = ceratophylla.
asperula, *Nova. ex Colla, Hort. Ripul.* 68.—Afr. austr.
ceratophylla, *Zahlbr. ex Endl. Atakta Bot.* 16. t. 15. —Austral.
Cercodia, Ait. Hort. Kew. ed. I. ii. 87 = alata.
citricolora, Walp. Rep. ii. 99 = micrantha.
confertifolia, *F. Muell. Fragm.* x. 58.—Austral.
conferta, *Fenzl, in Enum. Pl. Hueg.* 45.—Austral.
cyathiflora, *Fenzl, l. c.* 44.—Australi.
depressa, *Walp. Rep.* ii. 99.—Austral.; N. Zel.
digyna, *Labill. Nov. Holl. Pl.* i. 101. t. 128.—Austral.
distichia, Jack, in Malay. Misc. ii. (1822) n. VII. 19 = Anisophyllea distichia.
elata, *A Cunn. ex Fenzl, in Enum. Pl. Hueg.* 45.—Austral.
elata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 475 = tecuroides.
filiformis, A. Gray, Bot. U. St. Expl. Exped. i. 628 = heterophylla.
foliosa, *Benth. Fl. Australi.* ii. 477.—Austral.
glauca, *Lindl. in Mitch. Journ. Trop. Austral.* 91.—Austral.
Gonocarpus, Spreng. Syst. ii. 261 = tetragyna.
Gossei, *F. Muell. Fragm.* viii. 161.—Austral.
Gunnii, Hook. f. Fl. Tasm. i. 120 = tecuroides.
heterophylla, *Brongn. in Duperr. Voy. Coq. Bot.* t. 68. —Austral.
hexandra, *F. Muell. Fragm.* iii. 31.—Austral.
incana, Walp. Rep. ii. 99 = tetragyna.
intricata, *Benth. Fl. Australi.* ii. 481.—Austral.
ivaefolia, Salisb. Prod. 276 = alata.
Lambertii, *Mentr. in Mém. Acad. Lyon*, x. (1860) 199. —N. Caled.
lancelata, *R. Br. ex Benth. Fl. Australi.* ii. 482.—Austral.
leptothrix, F. Muell. Fragm. iii. 32 = scanthocarpa.
micrantha, *R. Br. ex Sieb. & Zucc. Fl. Jap. Fam. Nat.* i. 25.—As. or.; Austral.; N. Zel.
monosperma, *F. Muell. in Proc. Linn. Soc. N. S. W.* x. (1885) 197.—Austral.
montana, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 475 = depressa.
mucronata, *Benth. Fl. Australi.* ii. 475.—Austral.
nodulosa, *Walp. Rep.* v. 672.—Austral.
odontocarpa, *F. Muell. Fragm.* i. 108.—Austral.
oligantha, Am. in Nov. Act. Nat. Cur. xviii. (1836) 386 = Serpicula zeylanica.
oligantha, Wight & Arn. Prod. 388 = Myriophyllum intermedium.
paniculata, *R. Br. ex Benth. Fl. Australi.* ii. 481.—Austral.
pinnatifida, *A. Gray, Bot. U. St. Expl. Exped.* i. 627. —Austral.
pinnatifida, Hook. f. Fl. Tasm. i. 119 = ceratophylla.
pithyoides, *Benth. Fl. Australi.* ii. 476.—Australi.
platycarpa, *Benth. l. c.* 478.—Austral.
prostrata, *Forst. Char. Gen.* 82. t. 81.—N. Caled.
pusilla, *R. Br. ex Benth. Fl. Australi.* ii. 481.—Austral.
racemosa, *Labill. Nov. Holl. Pl.* i. 100. t. 128.—Austral.
rotundifolia, *Benth. Fl. Australi.* ii. 480.—Austral.
rudis, *Benth. l. c.* 480.—Austral.
salsoloides, *Benth. l. c.* 485.—Australi.
scabra, Benth. Fl. Hongk. 139 = tetragyna.
scoparia, *Fenzl, in Enum. Pl. Hueg.* 45.—Austral.
scordioides, *Benth. Fl. Australi.* ii. 485.—Austral.
serpyllifolia, Walp. Rep. ii. 99 = depressa.
Serra, *Brongn. in Duperr. Voy. Coq. Bot.* t. 69.—Austral.
stricta, *R. Br. ex Benth. Fl. Australi.* ii. 482.—Austral.

HALORAGIS:—
tenella, Brongn. in Duperr. Voy. Coq. Bot. t. 68. f. B = micrantha.
tenuifolia, *Benth. Fl. Australi.* ii. 477.—Australi.
Tetragonia, L'Hérit. Stirp. Nov. 82 = alata.
tetragyna, *Hook. f. Fl. N. Zel.* i. 62.—As. or.; Austral.; N. Zel.
tetrandra, *Schott, in Spreng. Syst.* iv. *App.* 405.—Bras.
tenuicoides, *A. Gray, Bot. U. St. Expl. Exped.* i. 625. Austral.
trichostachya, *Benth. Fl. Australi.* ii. 481.—Austral.
trifida, *Walp. Rep.* v. 672.—Austral.
trigonocarpa, *F. Muell. Fragm.* x. 84.—Austral.
uniflora, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 548. —N. Zel.
verrucosa, Walp. Rep. ii. 99 = depressa.

HALOSCHOENUS, Nees, in Linnæa, ix. (1834) 296 = Eyncbospora, Vahl (Cyperac.).
albidus, Nees, in Mart. Fl. Bras. ii. 1. 122 = R. albida.
capillaris, Nees, in Linnæa, ix. (1834) 296 = R. tenuis.
caracasanus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 249 = R. tenuis.
contractus, Nees, in Mart. Fl. Bras. ii. t. 123 = R. micrantha.
elatior, Nees, l. c. 120 = R. stricta.
emaciatus, Nees, l. c. 121 = R. tenuis.
hispidulus, Nees, l. c. 124.—Bras.
nardifolius, Nees, l. c. 122 = R. nardifolia.
pygmæus, Nees, in Linnæa, ix. (1834) 296 = R. micrantha.
rigidus, Nees, ex Kunth, Enum. Pl. ii. 300 = R. polyphylla.
riparius, Nees, in Mart. Fl. Bras. ii. t. 120 = R. tenuis.
sabulosus, Nees, in Linnæa, ix. (1834) 296 = R. tenuis.
sparsus, Nees, l. c. = R. micrantha.
stoloniferus, Nees, in Mart. Fl. Bras. ii. t. 120 = R. stoloniferæ.
tenuius, Nees, l. c. 123 = R. tenella.

HALOSCIAS, Fries, Summ. Veg. Scand. 180 (1846) = **Ligusticum**, Linn. (Umbellif.).
scoticum, Fries, l. c. = L. scoticum.

HALOSTACHYS, C. A. Mey. in Bull. Soc. Nat. Mosc. (1838) 361. *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 63.
caspica, *C. A. Mey. l. c.* (caspia).—Reg. Casp.; Turkestan.
nodulosa, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 361 = Halopeplis amplexicaulis.
occidentalis, S. Wats. in Bot. King's Exped. 398 = Spirostachys occidentalis.
patagonica, Moq. in DC. Prod. xiii. ii. 148 = Spirostachys patagonica.
perfoliata, Moq. l. c. = Halopeplis amplexicaulis, Halopeplis perfoliata.
Ritteriana, Moq. l. c. = Heterostachys Ritteriana.
songarica, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 361 = Halopeplis pygmæa.

HALOSTEMMA, Benth. & Hook. f. Gen. iii. 1056 (1883), sphalm. = **Mapania**, Aubl. (Cyperac.).

HALOTHAMNUS, Jaub. & Spach, Illustr. ii. 50. t. 136 (1846) = **Salsola**, Linn. (Chenopod.).
Bottæ, Jaub. & Spach, l. c. = S. Bottæ.

HALOTHAMNUS, F. Muell. Fl. Vict. i. 158 (1862) = **Flagellanthus**, Forst. (Malvac.).
microphyllus, F. Mue l. l. c. 159 = P. microphyllus.

HALOTIS, Bunge, Anabas. Rev. 78. t. 1 (1862) = **Halimocnemis**, C. A. Mey. (Chenopod.).
occulta, Bunge, l. c. 74.—Persia.

HALOXYLON, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 488. *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 70.
Ammodendron, *Bunge, l. c.* 469.—Turkest.
articulatum, *Bunge, l. c.*—Reg. Mediterr. occ.; Syria: Arab.

HALOXYLON :—
floridum, *Fenzl, in Ledeb. Fl. Ross.* iii. 819.—Reg. Casc.
Griffithii, *Boiss. Fl. Orient.* iv. 950.—Afghan.
multiflorum, *Bunge, ex Boiss. l. c.* 949.—Afghan.
persicum, Bunge, ex Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 169 = H. Ammodendron.
ramosissimum, *Benth. & Hook. f. Gen.* iii. 70.—Syria.
recurvum, *Bunge, ex Boiss. Fl. Orient* iv. 949.—Beluchist. ; Ind. or. ; China.
Regelii, *Bunge, in Bull. Acad. Pétersb.* xxv. (1879) 568.—Turkestan.
salicornicum, *Bunge, ex Boiss. Fl. Orient.* iv. 949.—Persia ; Afghan.
Schmittianum, *Pomel, Nouv. Mat. Fl. Atl.* 334.—Alger.
scoparium, *Pomel, l. c.* 335.—Alger.
Stocksii, Hook. f. in Benth. & Hook. f. Gen. iii 70 = recurvum.
subulifolium, *Bunge, in Act. Hort. Petrop.* vi. (1879) 488.—Turkestan.
Thomsoni, *Bunge, ex Boiss. Fl. Orient.* iv. 950, in obs. —Tibet. occ.

HALYMUS, Wahlenb. Fl. Suec. ii. 662 (1826) = HALIMUS, Linn. = **Atriplex**, Tourn. (Chenopod.).

HAMADRYAS, Comm. ex Juss. Gen. 382 (1789). *RANUNCULACEÆ*, Benth. & Hook. f. i. 6.
amicola, Hook. lc. Pl. t. 137 = Anemone integrifolia.
argentea, *Hook. f. Fl. Antarct.* 227.—Ins. Falkland.
Kingii, *Hook. f. l. c.* 228.—Reg. Magellan.
magellanica, *Lam. Encyc.* iii. 97.—Reg. Magellan.
tomentosa, DC. Syst. i. 227 = magellanica.
tomentosa, Hook. f. Fl. Antarct. i. 85 (sphalm.) = argentea.

HAMALIUM, Hemsl. Biol. Centr. Am. Bot. ii. 187 (1881) sphalm. = HAMULIUM, Cass. = **Verbesina**, Linn. (Compos.).

HAMAMELIS, Gronov. ex Linn. Gen. ed. II. 54 (1742). *HAMAMELIDEÆ*, Benth. & Hook. f. i. 667.
TRILOPUS, Mitch. in Act. Nat. Cur. viii. (1746) App. 219.
afra, Burm. f. Fl. Cap. Prod. 4 (Quid ?).—Afr. austr.
androgyna, Walt. Fl. Carol. 255 = virginiana.
arborea, Ottolander, ex Mast. in Gard. Chron. (1874) i. 187 = japonica.
caroliniana, Walt. ex Stend. Nom. ed. I. 388 = virginiana.
chinensis, R. Br. in Abel, Narr. Journ. Chin. 375 = Loropetalum chinense.
coryliifolia, Moench, Meth. 278 = virginiana.
dentata, Rafin. New Fl. Am. iii. 17 = virginiana.
dioica, Walt. Fl. Carol. 255 = virginiana.
ætiensis, Rafin. New Fl. Am. iii. 16 = Fothergilla Gardeni ?
Ayenalis, Rafin. l. c. = virginiana.
japonica, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1843) 198.—Japon.
macrophylla, Pursh, Fl. Am. Sept. i. 116 = virginiana.
monnica, Walt. Fl. Carol. 255 = Fothergilla Gardeni ?
nigra, Rafin. New Fl. Am. iii. 16 = virginiana.
parvifolia, Rafin. Med. Bot. i. 280 = virginiana.
persica, DC. Prod. iv. 266 = Parrotia persica.
riparia, Rafin. New Fl. Am. iii. 16 = virginiana.
rotundifolia, Rafin. l. c. = virginiana.
virginiana, *Linn. Sp. Pl.* 124.—Am. bor.
virginica, Linn. Syst. ed. XII. 129 = præc.
Zuccariniana, Ottolander, ex Mast. in Gard. Chron. (1874) i. 187 = japonica.

HAMARIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 264 = **Astragalus**, Tourn. (Legumin.).
uncinata, Fourr. l. c. = A. hamosus.

HAMARIA, Kunze, ex Reichb. Consp. 212* (1828) 'Rosac.' = (Quid ?).

HAMATOLOBIUM, Fenzl, Pugill. Pl. Nov. Syr. 8 (1842) = **Hammatolobium**, Fenzl (Legumin.).

HAMATRIS, Salisb. Gen. Pl. Fragm. 11 (1866) = **Dioscorea**, Linn.
triphylla, Salisb. l. c. 12 = D. triphylla.

HAMBERGERA, Scop. Introd. 106 (1777) = **Casoneta**, Aubl. (Combretac.).

HAMELIA, Jacq. Enum. Pl. Carib. 2 (1760) ; Select. Am. 71 (1763). *RUBIACEÆ*, Benth. & Hook. f. ii. 75
DUHAMELIA, Pers. Syn. i. 208 (1805).
LONICERA, Adans. Fam. ii. 157 (1763).
TANGARACA, Adans. l. c. 148 (1763).
TEPESIA, Gaertn. f. Fruct. iii. 72. t. 192 (1805).
appendiculata, Gaertn. f. Fruct. iii. 64 = axillaris.
axillaris, *Sw. Prod. Veg. Ind. Occ.* 46.—Ind. occ.
breviflora, *A. Rich. Fl. Cub. Fanerog.* ii. 17.—Cuba.
chrysantha, Jacq. Coll. iii. 304 = lutea.
chrysantha, Sieber, ex Steud. Nom. ed. II. i. 721 = patens.
chrysantha, *Sw. Prod. Veg. Ind. Occ.* 46.—Ind. occ.
coccinea, Sw. l. c. = patens.
connata, Wall. ex DC. Prod. iv. 388 = Leycesteria formosa.
cupres, *Griseb Fl. Brit. W. Ind.* 320.—Jamaic.
erecta, Jacq. Enum. Pl. Carib. 16 = patens.
glabra, Lam. Encyc. iii. 65 = Amaioua guianensis.
grandiflora, L'Hérit. Sert. Angl. 4. t. 7 = ventricosa.
hamquœus, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 288.—Mexic.
latifolia, Reichb. ex DC. Prod. iv. 449 = patens.
lucida, Desf. Tabl. ed. II. 135 = axillaris.
lutea, *Hook, ex Sm. in Rees, Cycl.* xvii. n. 4.—Am. trop.
Mangillio, Steud. Nom. ed. I. 388 = Myrsine Mangillio.
nodosa, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 284.—Mexic.
odorata, Willd. ex Roem. & Schult. Syst. v. 267 = suaveolens.
patens, *Vorz. Enum. Pl. Carib.* 16 ; *Select. Am.* 72.—Am. trop.
pauciflora, Willd. ex Spreng. Syst. i. 765 = ventricosa.
rostrata, *Bartl. ex DC. Prod.* iv. 449.—Mexic.
sessiliflora, Willd. Sp. Pl. i. 981 = Amaioua guianensis
sphærocarpa, *Ruiz & Pav. Fl. Per.* ii. 69. t. 221.—Peruv.
suaveolens, *H. B. & K. Nov Gen. et Sp.* iii. 414.—Am. austr.
ventricosa, *Sw. Prod. Veg. Ind. Occ.* 46.—Am. austr.
verticillata, [Moç. & Sesse,] ex DC. Prod. iv. 449 = patens.
xorullensis, *H. B. & K. Nov. Gen. et Sp.* iii. 414.—Mexic.

HAMELINIA, A. Rich. Fl. Nouv. Zél. 158. t. 24 (1832) = **Astelia**, Banks (Liliac.).
veratroides, A. Rich. l. c. = A. veratroides.

HAMELLIA, Linn. Sp. Pl. ed. II. 246 (1762) = **Hamellia**, Jacq. (Rubiac.).

HAMILTONIA, Harv. Gen. Pl. Cap. ed. I. 298 (1838) = Colpoon, Berg (Santalac.).
capensis, Harv. l. c. (*Rhoicarpus capensis*, A. DC.). —Afr. austr.

HAMILTONIA, Muhl. ex Willd. Sp. Pl. iv. 1114 (1805) = **Pyrularia**, Michx. (Santal.).
oleifera, Muhl. ex Willd. l. c. 1130 = P. pubera.

HAMILTONIA, Roxb. Hort. Beng. 15 (1814) ; Fl. Ind. i. 554 (1832). *RUBIACEÆ*, Benth. & Hook. f. ii. 75.
SPERMADICTYON, Roxb. Corom. Pl. iii. 29. t. 286 (1819).
dulina, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 187 = suaveolens.
fruticosa, D. Don, l. c. = Leptodermis lanceolata.
inodora, Hamk. Cat. Pl. Hort. Beng. Alt. 111 = Leptodermis lanceolata.
lanceolata, Balll. Hist. Pl. vii. 278 = Leptadenia lanceolata.
myrsinoides, Wight & Arn. Prod. 428 = suaveolens.
oblonga, Franch. Pl. Davidian. i. 154 = Leptodermis oblonga.
pilosa, *Roxb. Hort. Beng.* 35 ; *Fl. Ind.* i. 555.—Ins. Moluc.
propinqua, Deene. in Jacquem. Voy. Bot. 83. t. 91 = suaveolens.
scabra, D. Don, Prod. Fl. Nep. 187 = suaveolens.
spectabilis, Carr. in Rev. Hortic. (1872) 191 = suaveolens !

HAMILTONIA :—
suaveolens, D. Don, Prod. Fl. Nep. 187 = Leptodermis lanceolata.
suaveolens, *Roxb. Hort. Beng.* 15 ; *Fl. Ind.* i. 554.—Ind. or. ; China.

HAMILTONIA, Spreng. Syst. i. 831 (1825), partim = **Comandra**, Nutt. (Santalac.).
elegans, Reichb. f. Ic. Fl. Germ. f. 1169 = C. elegans.
sarmentosa, Spreng. Syst. i. 831 = C. livida.
umbellata, Spreng. l. c. = C. umbellata.

HAMMATOCAULIS, Tausch, in Flora, xvii. (1834) 347 = **Ferula**, Tourn. (Umbellif.).
crotica, Tausch, l. c. = F. nodosa !

HAMMATOLOBIUM, Fenzl, Illustr. Pl. Syr. i. t. 1 (1843). *LEGUMINOSÆ*, Benth. & Hook. f. i. 509.
LUDOVICIA, Coss. in Bull. Soc. Bot. Fr. iii. 674 (1856).
Kremerianum, *C. Muell. in Walp. Ann.* vii. 744.—Alger.
lotoides, *Fenzl, Illustr. Pl. Syr.* i. t. 1.—Syria.

HAMOLOCENCHRUS, Scop. Introd. 78 (1777) = HOMALOCENCHRUS, Mieg. = **Leersia**, Sw. (Gramin.).

HAMOSA, Medic. in Vorles. Churpf. Phys. Oec. ii. (1787) 376 = **Astragalus**, Tourn. (Legumin.).
Astragalus, Medic. l. c. = A. hamosus.
glycyphylla, Medic. l. c. = A. glycyphyllus.
trimestris, Medic. l. c. = A. trimestris.

HAMPEA, Schlecht. in Linnæa, xi. (1837) 371. *MALVACEÆ*, Benth. & Hook. f. i. 211.
integerrima, *Schlecht. l. c.* 372.—Mexic.
thespesioides, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 188.—N. Granat.

HAMULIA, Rafin. Fl. Tellur. iv. 109 (1836) = **Utricularia**, Linn. (Lentib.).
alba, Rafin. l. c. = U. alba.
flava, Rafin. l. c. = U. flava.

HAMULIUM, Cass. in Bull. Soc. Philom. (1820) 173 ; et in Dict. Sc. Nat. xx. 260 (1821) = **Verbesina**, Linn. (Compos.).
alatum, Cass. in Dict. Sc. Nat. xx. 261 = V. alata.
Wrightii, Sch. Bip. in Bonplandia, ix. (1861) 365 = V. Wrightii, *Griseb.*

HANBURIA, Seem. in Bonplandia, vi. (1858) 293. *CUCURBITACEÆ*, Benth. & Hook. f. i. 836.
NIETOA, Schaffn. in La Naturaleza, iii. (1876) 343. mexicana, *Seem. l. c.*—Mexic.

HANCEA, Seem. Bot. Voy. Herald, 409. t. 96 (1857) = **Mallotus**, Lour. (Euphorb.).
brachiana, Seem. l. c. = M. Hookerianus.
muricata, Benth. Fl. Hongk. 306 = M. Furetianus.

HANCORNIA, Gomez, in Mem. Acad. Lisboa, iii. (1812) 3. Mem. Corresp. 51. *APOCYNACEÆ*, Benth. & Hook. f. ii. 693.
arborescens, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. i. 23 (= *Zschokkia arborescens*).—Bras.
floribunda, Poepp. & Endl. Nov. Gen. et Sp. iii. 70. t. 279 = Lacmellia floribunda.
Gardneri, Miers, Apocyn. S. Am. 12 = speciosa.
gracilis, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. i. 21 (= *Zschokkia gracilis*).—Bras.
laxa, A. DC. in DC. Prod. viii. 326 = Ambelania laxa.
macrophylla, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. i. 18 = Ambelania quadrangularis.
microcarpa, Spruce, ex Muell. Arg. l. c. 23 (= *Zschokkia microcarpa*).—Bras.
monosperma, Spruce, ex Muell. Arg. l. c. 23 (= *Zschokkia monosperma*).—Bras.
pubescens, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 56 = speciosa.
ramosissima, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. i. 21 (= *Zschokkia ramosissima*).—Bras.
speciosa, *Gomez, in Mem. Acad. Lisboa*, iii. (1812) *Mem. Corresp.* 53.—Bras.

HANDSCHIA, Pohl, in Flora, viii. (1825) 1. 183, *nomen* (Quid?).

HANGUANA, Blume, Enum. Pl. Jav. 15 (1827) = **Susum**, Blume (Flagellar.). *Kasimtu*, Blume, l. c. = S. anthelmiticum.

HANNAFORDIA, F. Muell. Fragm. ii. 9 (1860). *STERCULIACEAE*, Benth. & Hook. f. i. 227. Bisillili, *F. Muell. l. c.* x. 95.—Austral. quadrivalvis, *F. Muell. l. c.* ii. 9.—Austral. Shaneili, *F. Muell. l. c.* vi. 175.—Austral.

HANNOA, Planch. in Hook. Lond. Journ. Bot. v. (1846) 566. *SIMARUBEAE*, Benth. & Hook. f. i. 308. Schweinfurthii, *Oliver, in Hook. Ic. Pl.* t. 1256.—Afr. trop. undulata, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 567.—Afr. trop.

HANSALIA, Schott, in Oestr. Bot. Zeitschr. (1858) 82 = **Amorphophallus**, Blume (Aroid.). *grata*, Schott, l. c.—Afr. trop.

HANSENIA, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 754 = **Ligusticum**, Linn. (Umbellif.). *mongholica*, Turcz. l. c.—Mongol.

HANSTEINIA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 142. *ACANTHACEAE*, Benth. & Hook. f. ii. 1107. gracilis, *Oerst. l. c.* 148.—Costa Rica.

HAPALANTHUS, Jacq. Enum. Pl. Carib. 1 (1760) = **Callisia**, Linn. (Commelin.). *repens*, Jacq. l. c. 12 = C. repens.

HAPALE, Schott, in Oestr. Bot. Wochenbl. (1857) 85 = sec.

HAPALINE, Schott, Gen. Aroid. 85 (Hapale); Gen. Aroid. 44 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 977. Benthamiana, *Schott, ll. cc.* 86; 45.—Reg. Himal.

HAPALOCARPUM, Miq. Fl. Ind. Bat. ii. 1. 617 (1856) = **Ammannia**, Houst. (Lythrar.). *indicum*, Miq. l. c. 618 = A. baccifera. *vesicatorium*, Miq. l. c. = A. baccifera.

HAPALOCHLAMYS, Reichb. Nom. 90 (1841) = APALOCHLAMYS, Cass. = **Cassinia**, R. Br. (Compos.).

HAPALOSA, Edgew. & Hook. f. in Fl. Brit. Ind. i. 245 (1874) = sec.

HAPALOSIA, Wall. Cat. n. 6962; ex Wight & Arn. Prod. 358 (1834) = **Polycarpon**, Linn. (Caryophyll.). *Loeflingiae*, Wall. Cat. l. c. = P. Loeflingiae.

HAPALOSTEPHIUM, D. Don, ex Sweet, Hort. Brit. ed. I. 210 (1827); et in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 307 = **Crepis**, Vaill. (Compos.). *alpestre*, Sweet, Hort. Brit. ed. III. 413 = C. alpestris. *australasica*, Sweet, l. c. ed. I. 210 = C. blattarioides. *crocifolium*, Sweet, l. c. ed. III. 413 = C. hieracioides. *grandiflorum*, G. Don, in Loud. Hort. Brit. 326 = C. grandiflora. *hieracioides*, Sweet, Hort. Brit. ed. III. 413 = C. hieracioides. *lyratum*, G. Don, ex Loud. Hort. Brit. 326 = C. lyrata. *macrophyllum*, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 308 = C. macrophylla. *paludosum*, D. Don, l. c. = C. paludosa. *pilosum*, D. Don, ex G. Don, in Loud. Hort. Brit. 326 = C. blattarioides. *pyrenaicum*, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 307 = C. blattarioides. *sibiricum*, D. Don, l. c. 308 = C. sibirica. *spicatum*, G. Don, in Loud. Hort. Brit. 326 = C. blattarioides.

HAPALUS, Endl. Gen. 460 (1836) = APALUS, DC. = Blennospermas, Less. (Compos.).

HAPLACHNE, Presl, Rel. Haenk. i. 284. t. 38 (1830) = **Dimeria**, R. Br. (Gramin.). *pilosissima*, Presl, l. c. 285. t. 38 = D. pilosissima.

HAPLANTHERA, Hochst. in Flora, xxvi. (1843) 71 = **Ruttya**, Harv. (Acanthac.). *speciosa*, Hochst. l. c. 72.—Abyssin.

HAPLANTHUS, Nees, in Wall. Pl. As. Rar. iii. 77 (1839). *ACANTHACEAE*, Benth. & Hook. f. ii. 1099. hygrophiloides, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 508.—Burma. *nilgherrensis*, Wight, Ic. iv. t. 1356 = tentaculatus. *plumosus*, T. Anders. in Journ. Linn. Soc. ix. (1867) 504 = tentaculatus. *tener*, Nees, in Wall. Pl. As. Rar. iii. 115 = Andrographis tenuiflora. tentaculatus, *Nees, in DC. Prod.* xi. 513.—Ind. or. verticillaris, *Nees, l. c.*—Ind. or.

HAPLOCARPHA, Less. in Linnaea, vi. (1831) 90. *COMPOSITAE*, Benth. & Hook. f. ii. 457. DAMATRIS, Cass. in Bull. Soc. Philom. (1817) 139; et in Dict. Sc. Nat. xii. 471 (1818). lanata, *Less. l. c.*—Afr. austr. Leichtlinii, *N. E. Br. in Gard. Chron.* (1883) i. 78. —Afr. austr.? *Lessingii*, DC. Prod. vi. 494 = lanata. lyrata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 465.—Afr. austr. scaposa, *Harv. l. c.*—Afr. austr. *Thunbergii*, Hort. ex N. E. Br. in Gard. Chron. (1883) i. 78 = Leichtlinii. Thunbergii, *Less. Syn. Comp.* 87.—Afr. austr.

HAPLOCARYA, Phil. in Linnaea, xxxiii. (1864–65) 208 = APLOCARYA, Lindl. = **Dolia**, Lindl. (Convolv.).

HAPLOCHILUS, Endl. Ench. 118 (1841); Gen. Suppl ii. 20 (1842) = **Zeuxine**, Lindl. (Orchid.). *affinis*, D. Dietr. Syn. Pl. v. 173.—Ind. or. *flavus*, D. Dietr. l. c. 172 = Z. flava. *longifolia*, D. Dietr. l. c. 173 = Z. longilabria. *nervosus*, D. Dietr. l. c. 172 = Z. nervosa. *parviflorus*, Blume, Orch. Archip. Ind. 110.—Malaya. *regius*, D. Dietr. Syn. Pl. v. 173; Blume, l. c. 111.—Malaya.

HAPLOCLATHRA, Benth. in Journ. Linn. Soc. v. (1860) 54. *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 189. leiantha, *Benth. l. c.*—Am. trop. paniculata, *Benth. l. c.*—Bras.

HAPLOCOELUM, Radlk. in Sitzb. Math. Phys. Acad. Muench. viii. (1878) 336. *SAPINDACEAE*. inopleum, *Radlk. l. c.* 337.—Afr. trop.

HAPLODESMIUM, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 189. t. 5 = **Chaetolepis**, Miq. (Melastom.). *Lindenianum*, Naud. l. c. = C. Lindeniana.

HAPLOESTES, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) (Pl. Fendl.) 109. *COMPOSITAE*, Benth. & Hook. f. ii. 441. Greggii, *A. Gray, l. c.*—Mexic.

HAPLOPHIUM, Endl. Gen. 712 (1839) = **Aplophium**, Cham. (Bignon.).

HAPLOPAPPUS, Endl. Gen. 385 (1838) = **Aplolappappus**, Cass. (Compos.).

HAPLOPETALON, A. Gray, Bot. U. St. Expl. Exped. i. 508. t. 76 (1854). *RHIZOPHOREAE*, Benth. & Hook. f. i. 680. AGATEA, Rich, ex A. Gray, Bot. U. St. Expl. Exped. i. 509 (1854). multiflorum, *Bull. Adansonia*, iii. (1862–63) 40.—N. Caled. Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 508. t. 76. —Ins. Sandvic. Semannii, *A. Gray, in Bonplandia*, x. (1862) 36.—Ins. Fiji.

HAPLOPETALUM, Miq. Fl. Ind. Bat. i. 1. 596 (1855) = *prec.*

HAPLOPHYLLUM, Reichb. Fl. Germ. Excurs. 766 (1832) = APLOPHYLLUM, A. Juss. = **Ruta**, Linn.

HAPLOPHYTON, A. DC. in DC. Prod. viii. 413 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 723. cimicidum, *A. DC. l. c.*—Mexic.

HAPLORHUS, Engl. in Bot. Jahrb. i. (1881) 419. *ANACARDIACEAE*. peruviana, *Engl. l. c.*—Peruv.

HAPLOSCIADIUM, Hochst. in Flora, xxvii. (1844) 29 = **Trachydium**, Lindl. (Umbellif.). *abyssinicum*, Hochst. l. c. = T. abyssinicum.

HAPLOSTELIS, Reichb. Nom. 86 (1841) = sec.

HAPLOSTELLIS, Endl. Gen. 219 (1837) = APLOSTELLIS, Thou. = **Pogonia**, Juss. (Orchid.).

HAPLOSTEMMA, Endl. Gen. Suppl. iii. 75 (1843) = **Vincetoxicum**, Rupp. (Asclep.). *arahucus*, Endl. ex Decne. in DC. Prod. viii. 565 = V. fruticulosum.

HAPLOSTEMUM, Endl. Gen. 119 (1836) = APLOSTEMON, Kaßn. = **Scirpus**, Linn. (Cyperac.).

HAPLOSTEPHIUM, Mart. ex DC. Prod. v. 78 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 384. Passerina, *Mart ex DC. l. c.*—Bras. ramosissimum, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 375.—Bras.

HAPLOSTICHIA, Phil. in Linnaea, xxx. (1859–60) 198 = **Senecio**, Tourn. (Compos.). *stolonifera*, Phil. l. c. = S. xosteraefolia.

HAPLOSTIGMA, F. Muell. Fragm. viii. 100 (1873) = **Loxocarya**, R. Br. (Restiac.).

HAPLOSTYLIS, Nees, in Linnaea, ix. (1834) 295 = **Rynchospora**, Vahl (Cyperac.). *albiceps*, Nees, in Mart. Fl. Bras. ii. 1. 129 = R. albiceps. *arenariifolia*, Nees, l. c. 126 = R. arenariifolia. *bahamensis*, Nees, in Linnaea, ix. (1834) 295 = R. pterocarpa. *barbata*, Nees, l. c. = R. pterocarpa. *elatior*, Nees, in Mart. Fl. Bras. ii. 1. 125 = R. elatior. *hirta*, Nees, l. c. = R. hirta. *Humboldtii*, Nees, in Linnaea, ix. (1834) 295 = R. pterocarpa. *mexicana*, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 249 = R. mexicana. *Meyeni*, Nees, in Linnaea, ix. (1834) 295 = R. Wallichii. *minor*, Nees, in Mart. Fl. Bras. ii. 1. 129.—Bras.

HAPLOTAXIS, Endl. Gen. 468 (1838) = APLOTAXIS, DC. = **Saussurea**, DC. (Compos.).

HAPPIA, Neck. Elem. ii. 126 (1790) = **Tococa**, Aubl. (Melastom.).

HAPTOPHYLLUM, Vis. & Panč. in Mem. Ist. Venet. xv. (1870) 14, sphalm. = APLOPHYLLUM, Juss. = **Ruta**, Linn.

HAQUETIA, D. Dietr. Syn. Pl. i. 562 (1839) = **Hacquetia**, Neck. (Umbellif.).

HARDENBERGIA, Benth. in Enum. Pl. Hueg. 40 (1837). *LEGUMINOSAE*, Benth. & Hook. f. i. 580. Comptoniana, *Benth. l. c.* 41.—Austral. *cordata*, Benth. l. c. = monophylla. *digitata*, Lindl. Bot. Reg. (1840) Misc. 68 = Comptoniana. *Huegelii*, Benth. in Enum. Pl. Hueg. 41 = Comptoniana. *Lindleyi*, Meisn. in Lehm. Pl. Preiss. i. 94 = Comptoniana. *macrophylla*, Benth. in Enum. Pl. Hueg. 41 = Comptoniana. *Makoyana*, Lem. Illustr. Hortic. v. (1858) t. 179 = Comptoniana. monophylla, *Benth. Enum. Pl. Hueg.* 41.—Austral. *ovata*, Benth. l. c. = monophylla. retusa, *Benth. Fl. Austral.* ii. 367.—Austral.

HARDWICKIA, Roxb. Hort. Beng. 39 (1814); Fl. Corom. iii. 6. t. 209 (1819). *LEGUMINOSÆ*, Benth. & Hook. f. i. 586.
binata, *Roxb. ll. cc.*—Ind. or.
Mannii, *Oliver, Fl. Trop. Afr.* ii. 316.—Afr. trop.
pinnata, *Roxb. Hort. Beng.* 38; *Fl. Ind.* ii. 425.—Ind. or.
tropæiformis, R. Grah. in Wall. Cat. n. 5809 = binata.

HARGASSERIA, C. A. Mey. in Bull. Acad. Phys.-Math. Pétersb. i. (1843) 355 = **Daphnopsis**, Mart. (Thymel.).
cestrifolia, Endl. Gen. Pl. Suppl. iv. ii. 68 = D. castrifolia.
cubana, A. Rich. Fl. Cub. Fanerog. ii. 198.—Cuba.
mexicana, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 359 = D. Bonplandii.
occidentalis, Griseb. in Goett. Abh. vii. (1857) 219 = D. Swartzii.
salicifolia, Endl. Gen. Pl. Suppl. iv. ii. 68 = D. salicifolia.
Schiedeana, Endl. l. c. = D. Bonplandii.
tinifolia, Endl. l. c. = D. tinifolia.

HARINA, Buch.-Ham. in Mem. Wern. Soc. v. (1826) 317 = **Wallichia**, Roxb. (Palm.).
caryotoides, Buch.-Ham. l. c. = W. caryotoides.
caryotoides, Griff. Palms Brit. Ind. 175 = W. densiflora.
densiflora, Walp. Ann. iii. 1092, Index, sphalm. = W. densiflora.
nana, Griff. Palms Brit. Ind. 176. t. 238 a, b = Didymosperma nanum.
oblongifolia, Griff. l. c. 175. t. 237 a, b, c = W. densiflora.
Rumphiana, Mart. ex Miq. Fl. Ind. Bat. iii. 29 = seq.
Rumphii, Mart. Hist. Nat. Palm. (ed. prior, i. 189) ii. 214, sub n. 104 = Drymophloeus oliveæformis.
Wallichia, Steud. Nom. ed. II. i. 722 = W. caryotoides.

HARIOTA, Adans. Fam. ii. 348 (1763) = **Rhipsalis**, Gaertn. (Cact.).
alternata, Lem. Hort. Univ. i. t. 50 = R. paradoxa.
Cassytha, Cels, ex Foerst. Handb. Cact. ed. I. 458 = R. Cassutha.
cribrata, Lem. Illustr. Hortic. iv. (1857) Misc. 12 = R. sagilonis?
floccosa, Cels, ex Foerst. Handb. Cact. ed. I. 458 = R. floccosa.
funalis, Cels, l. c. 457 = R. funalis.
mesembrianthemoides, Lem. Cact. Aliq. Nov. Desc. 29 = R. mesembryanthemoides.
pentaptera, Lem. ex Foerst. Handb. Cact. ed. I. 458 = R. pentaptera.
prismatica, Lem. Illustr. Hortic. x. (1863) Misc. 84.—Bras.
sagilonis, Lem. Cact. Aliq. Nov. Desc. 29 = R. sagilonis.
salicornioides, DC. Mém. Cact. 28 = R. salicornioides.

HARLANDIA, Hance, in Walp. Ann. ii. 648 (1852) = **Malothria**, Linn. (Cucurbit.).
bryoniæfolia, Hance, l. c. = M. heterophylla.

HARMALA, Tourn. ex Adans. Fam. ii. 348 (1763) = **Peganum**, Linn. (Rutac.).
dauurica, Crantz, Inst. ii. 468 = Ruta dahurica.
multifida, All. Fl. Pedem. ii. 101 = P. Harmala.
Peganum, Crantz, Inst. ii. 468 = P. Harmala.

HARMOGIA, Schau. in Linnæa, xvii. (1843) 238 = **Baeckea**, Linn. (Myrtac.).
Baueriana, Schau. l. c. = B. densifolia.
coryophylla, F. Muell. Fragm. ii. 80 = B. coryophylla.
crenulata, Miq. in Nederl. Kruidk. Arch. iv. (1859) 148 = B. crenatifolia.
crispifora, F. Muell. Fragm. ii. 81 = B. crispiflora.
Cunninghamii, Schau. in Linnæa, xvii. (1843) 236 = B. Cunninghamii.
densifolia, Schau. l. c. = B. densifolia.
leptophylla, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 330.—Austral.
ovalifolia, F. Muell. Fragm. ii. 82 = B. ovalifolia.
parviflora, Turcz. in Bull. Phys.-Math. Pétersb. x. (1852) 330.—Austral.
parvula, Schau. in Linnæa, xvii. (1843) 238 = B. virgata.
pentandra, F. Muell. Fragm. ii. 81 = B. pentandra.

HARMOGIA:—
pinifolia, Schau. in Linnæa, xvii. (1843) 238.—Austral.
propinqua, Schau. l. c. = B. densifolia.
serpyllifolia, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 330.—Austral.
umbellata, F. Muell. Fragm. ii. 81 = B. virgata.
virgata, Schau. in Linnæa, xvii. (1843) 238 = B. virgata.

HARNIERIA, Solms, in Sitzb. Ges. Naturf. Fr. Berl. (1864) 21 = **Justicia**, Houst. (Acanthac.).
dimorphocarpa, Solms, l. c. 110 = J. hetrocarpa.

HARONGA, Thou. Gen. Nov. Madag. 15 (1806). *HYPERICINEÆ*, Benth. & Hook. f. i. 167.
ARONGANA, Pers. Syn. ii. 91 (1807).
HARMOCARPUS, Noronha, ex Thou. Gen. Nov. Madag. 15 (1806).
HARONGANA, Choisy, Prod. Mon. Hyper. 38 (1821).
axillaris, G. Don, Gen. Syst. i. 614 = Psorospermum axillare.
crenata, Steud. Nom. ed. II. i. 722 = Psorospermum cordiæfolium.
febrifuga, Steud. l. c. = Psorospermum febrifugum.
lanceolata, Choisy, ex G. Don, Gen. Syst. i. 614 = Psorospermum citrifolium.
madagascariensis, *Choisy, Prod. Mon. Hyper.* 34.—Madag.
mollissima, Steud. Nom. ed. II. i. 722 = Psorospermum cuspidatum.
paniculata, Lodd. ex Steud. l. c. = madagascariensis.
pubescens, Steud. l. c. = madagascariensis.
revoluta, Choisy, ex DC. Prod. i. 543 = Psorospermum discolor, *Spach*.

HARONGANA, Choisy, Prod. Mon. Hyper. 38 (1821) = præc.

HARPACHAENA, Bunge, Del. Sem. Hort. Dorp. (1845) = **Acanthoosphalus**, Kar. & Klr. (Compos.).
amplexifolia, Bunge, l. c. = A. amplexifolius.

HARPACHNE, Hochst.—Cf. Flora, xxiv. (1841) i. Intell. 20, nomen; et ex A. Rich. Tent. Fl. Abyss. ii. 431 (1850) = **Eragrostis**, Beauv. (Gramin.).
Schimperi, Hochst. ll. cc. = E. Schimperi.

HARPAECARPUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 389 = **Madia**, Molina (Compos.).
exigua, A. Gray, in Torr. Bot. Mex. Bound. 101 = M. filipes.
madarioides, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 389 = M. filipes.

HARPAGONELLA, A. Gray, in Proc. Am. Acad. xi. (1876) 88. *BORAGINEÆ*, Benth. & Hook. f. ii. 846.
Palmeri, *A. Gray, l. c.*—Calif.

HARPAGONIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 = Dysoxylum I, Blume (Meliac.).

HARPAGOPHYTUM, DC. ex Meisn. Gen. 298 (1840). *PEDALINEÆ*, Benth. & Hook. f. ii. 1057.
UNCARIA, Burch. Trav. i. 536 (1822).
Burchellii, *DC. in Deless. Ic. Sel. Pl.* v. 89. t. 40.—Afr. austr.
leptocarpum, *Decne. in Ann. Sc. Nat.* Sér. V. iii. (1865) 329.—Madag.
procumbens, *DC. Prod.* ix. 257.—Afr. austr.
Zeyheri, *Decne. in Ann. Sc. Nat.* Sér. V. iii. (1865) 329.—Afr. austr.

HARPALIUM, Cass. in Bull. Soc. Philom. (1816) 141 = **Helianthus**, Linn. (Compos.).
asperum, Cass. in Dict. Sc. Nat. xxv. 436 = Helianthus asperi, incanus.
mexicanum, Walp. in Linnæa, xiv. (18) 806 = Helianthus mexicanus.
rigidum, Cass. in Dict. Sc. Nat. xxv. 360 = Helianthus rigidus.
scrotinum, DC. Prod. v. 594.—Am. trop.
tracheliense, Cass. in Dict. Sc. Nat. xxv. 436 = Helianthus tracheliensis.

HARPALYCE, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 308 = **Prevanthus**, Vaill. (Compos.).
alba, D. Don, l. c. = P. alba.
altissima, D. Don, l. c. = P. altissima.
cordata, D. Don, l. c. = P. altissima.
crepidina, D. Don, l. c. = P. crepidinea.
racemosa, D. Don, l. c. = P. racemosa.
rubicunda, D. Don, in Sweet, Hort. Brit. ed. III. 417 = P. Serpentaria.
Serpentaria, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 306 = F. Serpentaria.
simplex, D. Don, l. c. = P. virgata.
virgata, D. Don, l. c. = P. virgata.

HARPALYCE, [Moç. & Sesse,] ex DC. Mém. Lég. 496 (1825). *LEGUMINOSÆ*, Benth. & Hook. f. i. 494.
arborescens, *A. Gray, in Proc. Am. Acad. Sc.* v. (1861) 178.—Mexic.
brasiliana, *Benth. in Hook. Journ. Bot.* iii. (1841) 210.—Bras.
cubensis, *Griseb. Cat. Pl. Cub.* 71.—Cuba.
formosa, *DC. Prod.* ii. 528.—Mexic.
Hilariana, *Benth. in Mart. Fl. Bras.* xv. i. 51.—Bras.
minor, *Benth. l. c.* 52.—Bras.

HARPANEMA, Decne. in DC. Prod. viii. 496 (1844). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 748.
acuminatum, *Decne. l. c.*—Madag.

HARPECHLOA, Kunth, Rév. Gram. i. 92 (1830) (Harpochloa); Enum. Pl. i. 274 (1888). *GRAMINEÆ*, Benth. & Hook. f. iii. 1164.
capensis, *Kunth, l. c.*—Afr. austr.

HARPELEMA, Jacq. f. Eclog. Pl. ii. 6. t. 129 (1844) = **Rothia**, Pers. (Legumin.).
Dillwynia, Jacq. f. l. c. = R. trifoliata.

HARPEPHORA, Endl. Gen. Pl. Suppl. i. 1382 (1841) = **Aspilia**, Thou. (Compos.).
Endlicheriana, Walp. Rep. ii. 622.—Afr. trop.

HARPEPHYLLUM, Bernh. ex Krauss, in Flora, xxvii. (1844) 849. *ANACARDIACEÆ*, Benth. & Hook. f. i. 1001.
caffrum, *Bernh. ex Krauss, l. c.*—Afr. austr.
serratum, *Sond. ex Walp. Ann.* ii. 237.—Afr. austr.

HARPOCARPUS, Endl. Gen. Pl. Suppl. iii. 70 (1848) = **Acanthoosphalus**, Kar. & Klr. (Compos.).
amplexifolius, Endl. ex Walp. Rep. ii. 666 = A. amplexifolius.

HARPOCHILUS, Nees, in Mart. Fl. Bras. ix. 146. t. 24 (1847). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1116.
Neesianus, *Mart. ex Nees, l. c.*—Bras.
phacoesprus, *Nees, l. c.* 147.—Bras.

HARPOCHLOA, Kunth, Rév. Gram. i. 92 (1830) = **Harpechloa**, Kunth (Gramin.).

HARPULIA, G. Don, Gen. Syst. i. 669 (1831) = seq.

HARPULLIA, Roxb. Hort. Beng. 36 (1814); Fl. Ind. i. 645 (1832). *SAPINDACEÆ*, Benth. & Hook. f. i. 407.
BLANCOA, Blume, Rumphia, iii. 151 (1847).
DONATOPHORUS, Zipp. in Bijdr. Natuurk. Wetensch. v. (1850) 181.
OTONYCHIUM, Blume, Rumphia, iii. 180 (1847).
STREPTOTHOMA, Thw. in Hook. Kew Journ. vi. (1854) 293. t. 9 A.
THANATOPHORUS, Walp. Ann. ii. 218 (1851).
TINA, Blume, Bijdr. 235 (1825).
alata, *F. Muell. Fragm.* ii. 103.—Austral.
angustifolia, *Benth. in Stizenb. Math. Phys. Acad. Munech.* iii. (1879) 899.—N. Cale.
austro-caledonica, *Baill. Adansonia*, xi. (1876) 249.—N. Caled.
Blancoi, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 89 = cupaniöides.

HARPULLIA:—
confusa, Blume, Rumphia, iii. 176 = cupanioides.
cupanioides, *Roxb. Hort. Beng.* 66; *Fl. Ind.* i. 645.—Ind. or.
fraxinifolia, *Blume*, *Rumphia*, iii. 177.—Ins. Timor.
frutiosa, *Blume*, l. c. 179.—N. Guin.
Hillii, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 26.—Austral.
imbricata, Thw. Enum. Pl. Zeyl. 56 = cupanioides.
jugitanidifolia, Blume, Rumphia, iii. 177 = cupanioides.
Leichhardtii, *F. Muell. ex Benth. Fl. Austral.* i. 470.—Austral.
madagascariensis, *Radlk. in Sitzb. Math.-Phys. Acad. Munench.* viii. (1878) 273; ix. (1879) 519, 546.—Madag.
pendula, *Planch. ex F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 93.—Austral.
ramiflora, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 124.—Ins. Arn.
rupestris, Blume, Rumphia, iii. 175 = cupanioides.
thaustophora, *Blume*, *l. c.* 178.—N. Guin.
Wadsworthii, F. Muell. Fragm. iv. 1 = Cupaniopsis Wadsworthii.

HARRACHIA, Jacq. f. Eclog. Pl. i. 33. t. 21 (1816) = **Crossandra**, Salisb. (Acanthac.).
macrothyrsus, Mart. ex Nees, in Mart. Fl. Bras. ix. 85 = Aphelandra prismatica.
montana, Mart. ex Nees, l. c. 87 (= *Lagochilium montanum*).—Bras.
repanda, Mart. ex Nees, l. c. 86 = Aphelandra repanda.
speciosa, Jacq. Eclog. Pl. i. 33. t. 21 = Crossandra undulaefolia.
spinosa, Hook. ex Nees, in DC Prod. xi. 281 = Crossandra undulaefolia.

HARRERA, Macfad. Fl. Jamaic. ii. 60 (1837) = **Tetrasygia**, Rich. (Melastom.).
geracanthoides, Macfad. l. c. 61 = T. angustiflora.
paduncularis, Macfad. l. c. 101.—Jamaic.
sparsifolia, Macfad. l. c. 63.—Jamaic.
trinervis, Macfad. l. c. 63 = T. angustiflora.

HARRISONIA, R. Br. ex A. Juss. in Mém. Mus. Par. xii. (1825) 517. t. 28. *SIMARUBEAE*, Benth. & Hook. f. i. 314.
EBELINGIA, Reichb. Consp. 199 (1828).
LASIOLEPIS, Benn. Pl. Jav. Rar. 209. t. 42 (1838).
abyssinica, *Oliver*, *Fl. Trop. Afr.* i. 311.—Afr. trop.
Bennetii, A. W. Benn. in Hook. f. Fl. Brit. Ind. i. 519 = paucijuga.
Browniana, Gaudich. in Freyc. Voy. Bot. 479 = seq.
Brownii, *A. Juss. in Mém. Mus. Par.* xii. (1825) 540.—Ins. Timor.
paucifora, *Oliver*, *Fl. Trop. Afr.* i. 312, in obs.—As. trop.

HARRISONIA, [Hook. in] Sims, Bot. Mag. t. 2699 (1826) = **Maradenia**, R. Br. (Asclep.).
lonicereoides, [Hook.] l. c.=M. lonicereoides.

HARRISONIA, Neck. Elem. i. 84 (1790) = **Xeranthemum**, Linn. (Compos.).

HARTIANA, Rafin. Neogenyt. 2 (1825) = **Anemone**, Tourn. (Ranunc.).
caroliniana, Rafin. l. c.=A. decapetala.
tenella, Rafin. l. c.=A. decapetala.

HARTIGHAEA, Reichb. *Nom.* 212 (1841), sphalm.=seq.

HARTIGHSEA, A. Juss. in Mém. Mus. Par. xix. (1830) 327. t. 4 = **Dysoxylum**, Blume (Meliac.).
acuminata, Miq. Fl. Ind. Bat. Suppl. 504 = D. arborescens.
alliaria, Arn. ex Wight & Arn. Prod. 121 = D. Hamiltonii.
angustifolia, Miq. Fl. Ind. Bat. Suppl. 504 = Aglaia angustifolia.
Billardieri, A. Juss. in Mém. Mus. Par. xix. (1830) 328.—N. Caled.
cauliflora, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 411 = D. Cumingianum.
confusa, Miq. Fl. Ind. Bat. Suppl. 505 = D. costulatum.

HARTIGHSEA:—
excelsa, A. Juss. in Mém. Mus. Par. xix. (1830) 328 = Dysoxylum excelsum.
Forsteri, A. Juss. l. c.=D. Forsteri.
Fraseriana, A. Juss. l. c.=D. Fraseranum.
Gobara, Wight & Arn. ex Voigt, Hort. Suburb. Calc. 136=D. procerum.
Lessertiana, A. Juss. in Mém. Mus. Par. xix. (1830) 328=D. Lessertianum.
mollissima, A. Juss. l. c.=D. mollissimum.
Patersoniana, Endl. Prod. Fl. Norfolk. 79 = D. Patersoni.
ramiflora, Griff. Notul. iv. 501 = D. Griffithii.
rufa, A. Rich. Sert. Astrol. 29 = D. rufum.
schiaochitoides, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 412 = D. schizochitoides.
spectabilis, A. Juss. in Mém. Mus. Par. xix. (1830) 328 = D. spectabile.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 505 = D. arborescens.
trichocarpa, F. Muell. ex C. DC. in DC. Monog. Phan. i. 508=D. Muelleri.

HARTIGIA, Miq. in Linnaea, xviii. (1844) 284. t. 7 = **Miconia**, Ruiz & Pav. (Melastom.).
barbigera, Miq. l. c. 293 = M. racemosa.
millefiora, Miq. l. c. 286 = M. milleflora.
minutiflora, Miq. l. c.=M. milleflora.
oblongifolia, Miq. l. c. 288 = M. ciliata.
spectabilis, Miq. l. c. 286 = M. racemosa.

HARTIGSEA, Steud. Nom. ed. II. i. 722 (1840) = HARTIGHSEA, A. Juss. = **Dysoxylum**, Blume (Meliac.).

HARTMANNIA, DC. Prod. v. 693 (1836) = **Hemisandra**, DC. (Compos.).
ciliata, DC. l. c. 694 = Layia chrysanthemoides.
corymbosa, DC. l. c.=Hemiz. corymbosa.
fasciculata, DC. l. c. 693 = Hemiz. fasciculata.
flemerula, Nutt. in Trans. Am. Phil. Soc. N. S vii. (1841) 391 = Hemiz. fasciculata.
pungens, Hook. & Arn. Bot. Beech. Voy. 357 = Hemiz. pungens.

HARTMANNIA, Spach, Hist. Vég. Phan. iv. 370 (1835) = **Oenothera**, Linn. (Onagrar.).
affinis, Spach, ex Steud. Nom. ed. II. i. 722 = O. virgata
epilobifolia, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 364 = O. epilobiifolia.
gaurides, Spach, Hist. Vég. Phan. iv. 371 = O. roseo.
Kunthiana, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 362 = O. punnatifida.
macrantha, Spach, Hist. Vég. Phan. iv. 372 = O. tetraptera.
parviflora, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 362 = O. fruticosa.
roseo, G. Don, in Sweet, Hort. Brit. ed. III. 286 = O. roseo.
targuensis, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 361 = O. targuensis.
virgata, Spach, l. c. 361 = O. virgata.

HARTOGIA, Hochst. in Flora, xxvii. (1844) 805 = **Cassinopsis**, Sond. (Olacin.).
ilicifolia, Hochst. l. c.=C. capensis.

HARTOGIA, Linn. Syst. ed. X. 939 (1759) = **Agathosma**, Willd. (Rutac.).
betulina, Berg. Desc. Pl. Cap. 67 = Barosma betulina.
capensis, Linn. Syst. ed. X. 939 = A. hispida.
ciliaris, Linn. l. c. ed. XII. 625 = A. pubescens, *Willd. ciliata*, Berg. Desc. Pl. Cap. 68 = A. imbricata.
hispida, Walp. Rep. iv. 448 = A. hispida.
imbricata, Linn. Mant. i. 224 = A. imbricata.
lanceolata, Linn. Syst. ed. XII. 625 = A. pubescens, *Willd.*
pulchella, Berg. Desc. Pl. Cap. 69 = Barosma pulchella.
uniflora, Berg. l. c. 71 = Adenandra uniflora.
villosa, Berg. l. c. 70 = Adenandra cuspidata.

HARTOGIA, Linn. f. Suppl. 16 (1781). *CELASTRINEAE*, Benth. & Hook. f. i. 368.
SCHREBERA, Schreb. in Nov. Act. Soc. Sc. Upsal. i. (1778) t. 5. f. 1.—Cf. p. 91.
angustifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 601.—Afr. austr.
capensis, *Linn. f. Suppl.* 138.—Afr. austr.
multiflora, Eckl. & Zeyh. Enum. 127 = capensis.

HARTOGIA:—
riparia, Eckl. & Zeyh. l. c.=capensis.
Thea, E. Mey. in Drège, Zwei Pfl. Docum. 189 (= *Methysophyllum glaucum*).—Afr. austr.
triflobocarpa, *Baker*, *in Journ. Linn. Soc.* xx. (1883) 119.—Madag.

HARTWEGIA, Lindl. Bot. Reg. sub t. 1970 (1837). *ORCHIDEAE*, Benth. & Hook. f. iii. 528.
Gemma, *Reichb. f. in Gard. Chron.* (1878) ii. 8.—Am. centr.
purpurea, *Lindl. Bot. Reg.* sub t. 1970.—Mexic.

HARTWEGIA, Nees, in Nov. Act. Nat. Cur. xv. (1831) ii. 372 = Chlorophytum, Ker-Gawl. (Liliac.).
comosa, Nees, l. c. 373 = C. Sternbergianum.
Nimmonii, Dalz. ex Baker, in Journ. Linn. Soc. xv. (1876) 324 = C. glaucum.

HARUNGANA, Lam. Illustr. t. 645 (1808!) = **Haronga**, Thou. (Hyper.).
crenata, Poir. Encyc. Suppl. iv. 721 = Psorospermum cordiaefolium.
madagascariensis, Poir. in Dict. Sc. Nat. xx. 307 = Haronga madagascariensis.
pubescens, Poir. Encyc. Suppl. iv. 721 = Haronga madagascariensis.

HARVEYA, Hook. lc. Pl. t. 118, 351 (1837). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 967.
AULAYA, Harv. Gen. Pl. Afr. Austr. ed. I. 249 (1838).
capensis, *Hook. Ic. Pl* t. 118.—Afr. austr.
comorensis, *Vatke*, *in Linnaea*, xliii. (1880–82) 309.—Ins. Comor.
lactea, *Presl*, *Bot. Bemerk.* 91.—Afr. austr.
lutea, *Steud. Nom.* ed. II. i. 723.—Afr. austr.
obtusifolia, *Vatke*, *in Breman, Abh.* ix. (1885) 180.—Madag.
parviflora, *Hook. ex Steud. Nom.* ed. II. i. 723; *Reut. in DC. Prod.* xi. 33.—Afr. austr.
pratensis, *Presl*, *Bot. Bemerk.* 91.—Afr. austr.
purpurea, *Harv. ex Hook. Ic. Pl.* t. 351.—Afr. austr.
scariatna, *Hook. ex Steud. Nom.* ed. II. i. 723.—Afr. austr.
speciosa, *Bernh. in Flora*, xxvii. (1844) 881.—Afr. austr.
spectabilis, *Hook. ex Steud. Nom.* ed. II. i. 723; *Reut. in DC. Prod.* i. 36.—Afr. austr.
squamosa, *Steud.* l. c.—Afr. austr.
tubata, Hook. ex Steud. l. c.; Reut. in DC. Prod. i. 38. = Cycnium tubatum.
tubulosa, *Harv. Gen. S. Afr. Pl.* 249, nomen.—Afr. austr.
tubaghensis, *Presl*, *Bot. Bemerk.* 91.—Afr. austr.
varia, *Hook. ex Steud. Nom.* ed. II. i. 723; *Reut. in DC. Prod.* i. 38.—Afr. austr.

HARWAYA, Steud. Nom. ed. II. i. 723 (1840) = prasc.

HASSELQUISTIA, Linn. Cent. Pl. i. 9 (1755); Amoen. Acad. iv. 270 (1759) = **Tordylium**, Linn. (Umbellif.).
aegyptiaca, Linn. ll. cc.=T. aegyptiacum.
cordata, Jacq. Hort. Vindob. ii. 91 = T. cordatum.
cordifolia, Salisb. Prod. 160 = T. cordatum.
lanata, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 347 = T. lanatum.
pinnata, Moench, Meth. 78 = T. aegyptiacum.
syriaca, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 346 = T. syriacum.

HASSELTIA, Blume, Bijdr. 1045 (1826) = **Kiekxia**, Blume (Apocyn.).
arborea, Blume, l. c. 1046 = K. arborea.

HASSELTIA, H. B. & K. Nov. Gen. et Sp. vii. 231. t. 651 (1825). *TILIACEAE*, Benth. & Hook. f. i. 285.
floribunda, *H. B. & K. l. c.* 232.—N. Granat.
pubescens, *Benth. Pl. Hartw.* 164.—N. Granat.
pyramidalis, Hemsl. Diagn. Pl. Nov. 4 = Banara mexicana.

HASSKARLIA, Endl. Adansonia, i. (1860) 51. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 308.
didymostemon, *Baill. l. c.* 53.—Afr. trop.

HASSKARLIA, Meisn. Gen. Comm. 348 (1843) = **Turpinia**†, Vent. (Sapindac.).

HASSKARLIA, Walp. Ann. i. 758 (1849) = **Pandahus**, Linn.
globosa, Walp. l. c. = P. utilis.
leucocaustica, Walp. l. c. = P. odoratissimus.

HASTIFOLIA, Ehrh. Beitr. iv. 148 (1789) = **Scutellaria**, Linn. (Labiat.).

HASTINGIA, Koen. ex Sm. Exot. Bot. ii. 41. t. 80 (1806) = **Holmskioldia**, Retz. (Verben.).
coccinea, Sm. l. c. = Holmsk. sanguinea.
sceudens, Roxb. Hort. Beng. [86]; Fl. Ind. iii. 66 = Holmsk. sanguinea.

HASTINGSIA, S. Wats. in Proc. Am. Acad. xiv. (1879) 217, 242. *LILIACEAE*, Benth. & Hook. f. iii. 785.
alba, *S. Wats. l. c.* 242.—Am. bor. occ.
bracteosa, *S. Wats. l. c.* xx. (1885) 377.—Am. bor. occ.

HAUSSKNECHTIA, Boiss. Fl. Orient. ii. 960 (1872). *UMBELLIFERAE*.
etymistica, *Boiss. l. c.*—Persia.

HAUSSMANNIA, F. Muell. Fragm. iv. 148 (1864). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1041.
jucunda, *F. Muell. l. c.*—Austral.

HAUSTRUM, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

HAUYA, [Moç. & Sesse,] ex DC. Mém. Onagr. 2. t. 1 (1828). *ONAGRARIEAE*, Benth. & Hook. f. i. 791.
Barcenae, *Hemsl. Diagn. Pl. Nov.* 18.—Mexic.
californica, *S. Wats. in Proc. Am. Acad.* xx. (1885) 366.—Calif.
cornuta, *Hemsl. Diagn. Pl. Nov.* 13.—Guatemala.
elegans, [*Moç. & Sesse,*] *ex DC. Prod.* iii. 36.—Mexic.

HAVETIA, H. B. & K. Nov. Gen. et Sp. v. 203. t. 469 (1821). *GUTTIFERAE*, Benth. & Hook. f. i. 171.
flavida, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 366.—Guian.
flexilis, Spruce, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 246 = Havetiopsis Martii.
laurifolia, *H. B. & K. Nov. Gen. et Sp.* v. 203. t. 462. —N. Granat.
obovata, *Spruce, ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 250.—Bras.
octandra, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 11.—Peruv.

HAVETIOPSIS, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 246. *GUTTIFERAE*, Benth. & Hook. f. i. 179.
ORDEMATOPUS, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 249.
caryophylloides, *Planch. & Triana, l. c.* 246.—N. Granat.
flavida, *Planch. & Triana, l. c.* 247.—Guian.
flexilis, *Planch. & Triana, l. c.* 246.—Bras.
hippocrateoides, *Planch. & Triana, l. c.* 247.—Peruv.
Martii, *Planch. & Triana, l. c.* 246.—Bras.

HAWORTHIA, Duval, Pl. Succ. Hort. Alenç. 7 (1808). *LILIACEAE*, Benth. & Hook. f. iii. 777.
affinis, *Baker, in Journ. Linn. Soc.* xviii. (1880) 213. —Afr. austr.
albicans, *Haw. Syn. Pl. Succ.* 91.—Afr. austr.
altilinea, *Haw. in Phil. Mag.* (1824) 301.—Afr. austr.
angolensis, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 353.—Afr. trop.
angustifolia, *Haw. in Phil. Mag.* (1825) 383.—Afr. austr.
arachnoides, *Duval, Pl. Succ. Hort. Alenç.* 7.—Afr. austr.
arachnoidea, Haw. Syn. Pl. Succ. 96 = pruec.
aristata, Haw. Suppl. Pl. Succ. 51 = altilinea.
aspera, Haw. Syn. Pl. Succ. 90 = Apicra aspera.
asperiuscula, *Haw. Suppl. Pl. Succ.* 60.—Afr. austr.
asperula, *Haw. in Phil. Mag.* (1824) 300.—Afr. austr.
attenuata, *Haw. Syn. Pl. Succ.* 97.—Afr. austr.
attenuata, *Haw. Syn. Pl. Succ.* 92.—Afr. austr.

HAWORTHIA:—
bilineata, *Baker, in Journ. Linn. Soc.* xviii. (1880) 213.—Afr. austr.
Bolusii, *Baker, l. c.* 215.—Afr. austr.
brevis, Haw. Suppl. Pl. Succ. 57 = margaritifera.
chloracantha, *Haw. Revis. Pl. Succ.* 57.—Afr. austr.
clariperla, Haw. in Phil. Mag. (1826) 186 = attenuata.
coarctata, *Haw. l. c.* (1824) 301.—Afr. austr.
concava, Haw. Revis. Pl. Succ. 98 = cymbiformis.
concinna, Haw. Suppl. Pl. Succ. 59 = viscosa.
Cooperi, *Baker, in Saund. Refug. Bot.* t. 282.—Afr. austr.
corallina, Hort. Peacock, ex Baker, in Journ. Linn. Soc. xviii. (1880) 205 = margaritifera.
cordifolia, *Haw. Suppl. Pl. Succ.* 60.—Afr. austr.
curta, Haw. l. c. = tortuosa.
cuspidata, *Haw. l. c.* 51.—Afr. austr.
cymbiformis, *Duval, Pl. Succ. Hort. Alenç.* 7; *Haw. Syn. Pl. Succ.* 98.—Afr. austr.
denticulata, *Haw. Revis. Pl. Succ.* 58.—Afr. austr.
distincta, N. E. Br. in Gard. Chron. (1876) ii. 130. f. 30 = venosa.
erecta, Haw. Revis. Pl. Succ. 55 = margaritifera.
expansa, Haw. Syn. Pl. Succ. 91 = rigida.
fascista, *Haw. Revis. Pl. Succ.* 54.—Afr. austr.
foliolosa, Haw. Syn. Pl. Succ. 99 = Apicra foliolosa.
glabrata, *Baker, in Journ. Linn. Soc.* xviii. (1880) 206. —Afr. austr.
glauca, *Baker, l. c.* 208.—Afr. austr.
granata, Haw. Suppl. Pl. Succ. 57 = margaritifera.
Greenii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 202. —Afr. austr.
hybrida, *Haw. Revis. Pl. Succ.* 51.—Afr. austr.
icosiphylla, *Baker, in Journ. Linn. Soc.* xviii. (1880) 207.—Afr. austr.
imbricata, Haw. Syn. Pl. Succ. 98 = Apicra spiralis.
indurata, Haw. Revis. Pl. Succ. 49 = viscosa.
insteirtirae, *Haw. Suppl. Pl. Succ.* 43.—Afr. austr.
laevis, Haw. Revis. Pl. Succ. 52 = albicans.
limpida, Haw. Suppl. Pl. Succ. 51 = altilinea.
major, Duval, Pl. Succ. Hort. Alenç. 7 = margaritifera.
margaritifera, *Haw. Suppl. Pl. Succ.* 55.—Afr. austr.
maxima, Duval, Pl. Succ. Hort. Alenç. 7 = margaritifera.
minima, *Baker, in Journ. Linn. Soc.* xviii. (1880) 215. —Afr. austr.
minima, Haw. Syn. Pl. Succ. 99 = Aloë granata.
minor, Duval, Pl. Succ. Hort. Alenç. 7 = margaritifera.
mirabilis, *Haw. Syn. Pl. Succ.* 95.—Afr. austr.
mucronata, Haw. Suppl. Pl. Succ. 50 = altilinea.
multifaria, Haw. in Phil. Mag. (1824) 300 = mirabilis.
multiperlata, Steud. Nom. ed. II. i. 725 = margaritifera.
mutica, Haw. Revis. Pl. Succ. 55 = retusa.
nigra, *Baker, in Journ. Linn. Soc.* xviii. (1880) 202.— Afr. austr.
nigricans, Haw. in Phil. Mag. (1824) 301 = Gasteria nigricans.
nitida, Loud. Hort. Brit. 138 = Gasteria nitida.
obtusa, Haw. in Phil. Mag. (1825) 382 = cymbiformis.
pallida, *Haw. Rev. Pl. Succ.* 56.—Afr. austr.
papillosa, *Haw. Suppl. Pl. Succ.* 58.—Afr. austr.
parva, Haw. in Phil. Mag. (1824) 301 = tessellata.
Peacockii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 202.—Afr. austr.
pellucens, Haw. Syn. Pl. Succ. 96 = translucens.
pentagona, Haw. l. c. 97 = Apicra pentagona.
pillifera, *Baker, in Journ. Linn. Soc.* xviii. (1880) 214.—Afr. austr.
planifolia, Haw. in Phil. Mag. (1825) 382 = cymbiformis.
polyphylla, *Baker, in Journ. Linn. Soc.* xviii. (1880) 213.—Afr. austr.
polyphylla, Haw. ex Salm-Dyck, Hort. Dyck. 12 = Aloë granata.
Pseudo-rigida, Haw. Suppl. Pl. Succ. 62 = subrigida.
Pseudo-tortuosa, Haw. l. c. 59 = viscosa.
pumila, Duval, Pl. Succ. Hort. Alenç. 7; Haw. Syn. Pl. Succ. 96 = atrovirens.
Radula, *Haw. l. c.* 93.—Afr. austr.
ramifera, Haw. Revis. Pl. Succ. 52 = albicans.
recurva, *Haw. Syn. Pl. Succ.* 94.—Afr. austr.
Reinwardti, *Haw. Revis. Pl. Succ.* 53.—Afr. austr.
reticulata, *Haw. Syn. Pl. Succ.* 94.—Afr. austr.
retusa, *Duval, Pl. Succ. Hort. Alenç.* 7; *Haw. l. c.* 93.—Afr. austr.

HAWORTHIA:—
rigida, *Haw. Revis. Pl. Succ.* 49.—Afr. austr.
rugosa, *Baker, in Journ. Linn. Soc.* xviii. (1880) 206. —Afr. austr.
scabra, *Haw. Suppl. Pl. Succ.* 58.—Afr. austr.
semiglabrata, *Haw. l. c.* 55.—Afr. austr.
semimargaritifera, Haw. l. c. 53 = margaritifera.
setata, *Haw. l. c.* 54.—Afr. austr.
sordida, *Haw. Revis. Pl. Succ.* 51.—Afr. austr.
spiralis, Duval, Pl. Succ. Hort. Alenç. 7; Haw. Syn. Pl. Succ. 97 = Apicra spiralis.
spirella, Haw. l. c. 97 = Apicra pentagona.
subattenuata, *Baker, in Journ. Linn. Soc.* xviii. (1880) 205.—Afr. austr.
subfasciata, *Baker, l. c.* 204.—Afr. austr.
subregularis, *Baker, in Saund. Refug. Bot.* t. 282.— Afr. austr.
subrigida, *Baker, in Journ. Linn. Soc.* xviii. (1880) 201.—Afr. austr.
subtortuosa, Sweet, Hort. Brit. ed. II. 230 = viscosa.
subulata, *Baker, in Journ. Linn. Soc.* xviii. (1880) 206.—Afr. austr.
tessellata, *Haw. in Phil. Mag.* (1824) 300.—Afr. austr.
Tisleyi, *Baker, in Journ. Linn. Soc.* xviii. (1880) 208. —Afr. austr.
torquata, Haw. in Phil. Mag. (1827) 128 = viscosa.
tortella, Haw. Suppl. Pl. Succ. 61 = tortuosa.
tortuosa, *Haw. Syn. Pl. Succ.* 90.—Afr. austr.
translucens, *Haw. Suppl. Pl. Succ.* 62.—Afr. austr.
triangularis, *Duval, Pl. Succ. Hort. Alenç.* 7.—Afr. austr. ?
turgida, *Haw. Suppl. Pl. Succ.* 22.—Afr. austr.
venosa, *Haw. Revis. Pl. Succ.* 44.—Afr. austr.
virescens, Haw. l. c. 53 = albicans.
viscosa, *Haw. Syn. Pl. Succ.* 90.—Afr. austr.
vittata, *Baker, in Saund. Refug. Bot.* t. 283.—Afr. austr.

HAXTONIA, A. Cunn in Ann. Nat. Hist. ii. (1839) 127.
argophylla, Caley, ex D. Don, l. c. = O. argophylla.
foliosa, Hort. Bonn. ex DC. Prod. v. 270 = O. ramosissima.
furfuracea, A. Cunn. in Ann. Nat. Hist. Ser. I. ii (1839) 127 = O. furfuracea.
phlegeppha, Don, in Edinb. N. Phil. Journ. (1831) 278 = O. stellulata.
stellulata, D. Don, l. c. = O. stellulata.
tomentosa, D. Don, l. c. = O. dentata.
viscosa, D. Don, l. c. = O. viscosa.

HAYA, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 408. *PARONYCHIEAE*.
obovata, *Balf. f. l. c.* 409.—Ins. Socotra.

HAYECKA, Pohl, in Flora, viii. (1825) 183, nomen (Incertae sedis).

HAYLOCKIA, Herb. in Bot. Reg. t. 1871 (1830). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 725.
mesochlon, M. Roem. Syn. Ensat. 47 = Zephyranthes mesochlos.
pusilla, *Herb. in Bot. Reg.* t. 1871.—Reg. Argent.

HAYNALDIA, Kanitz, in Magyar Növent. Lapok, i. (1877) 8. *CAMPANULACEAE*.
exalbata, *Kanitz, l. c.* 4.—Bras.
Hilairiana, *Kanitz, in Mart. Fl. Bras.* vi. IV. 143.— Bras.
organensis, *Kanitz, in Magyar Növent. Lapok*, i. (1877) 4.—Bras.
thapsioides, *Kanitz, l. c.*—Bras.
uranoscoma, *Kanitz, l. c.*—Bras.

HAYNALDIA, Schur, Enum. Fl. Transs. 807 (1866) = **Agropyron**, J. Gaertn. (Gramin.).
villosa, Schur, l. c. = A. villosum.

HAYNEA, Reichb. Consp. 202 (1828) = **Modiola**, Moench (Malvac.).

HAYNEA, Schum. & Thonn. Beskr. Guin. Pl. 406 (1827). *URTICACEAE*, Benth. & Hook. f. iii. 361.
ovalifolia, *Schum. & Thonn. l. c.*—Ins. trop.

HAYNEA, Willd. Sp. Pl. iii. 1787 (1800) = **Pancovinia**, Ambl. (Compos.).
edulis, Willd. l. c. = P. edulis.
pedunculata, *Spreng. Syst. iii.* 381 = Vernonia pedunculata.

HEARNIA, F. Muell. Fragm. v. 55 (1865). *MELIACEÆ*, Benth. & Hook. f. i. 994.
Beccariana, *C. DC. in DC. Monog. Phan.* i. 629.—Malaya.
Cumingiana, *C. DC.* l. c.—Malaya.
elliptica, *C. DC.* l. c. 628.—Malaya.
glaucescens, *C. DC.* l. c. 631.—Malaya.
lancifolia, *C. DC.* l. c. 630.—Malaya.
macrophylla, *C. DC.* l. c. 631.—Malaya.
sapindina, *F. Muell. Fragm.* v. 55.—Australi.
sarawakana, *C. DC. in DC. Monog. Phan.* i. 632.—Borneo.
villosa, *C. DC.* l. c.—Malaya.

HEBANTHE, Mart. Nov. Gen. et Sp. ii. 42. tt. 140-145 (1826). *AMARANTACEÆ*, Benth. & Hook. f. iii. 41.
TROMMSDORFFIA, Mart. Nov. Gen. et Sp. ii. 40. t. 139 (1826).
argenteo-sericea, *Benth. & Hook. f. Gen.* iii. 42.—Bras.
aurata, *Benth. & Hook. f.* l. c.—Bras.
decipiens, *Hook. f.* l. c. 41.—Persia.
gulanensis, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 933, nomen.—Guiana.
holosericea, *Mart. in Flora*, xxi. (1838) 11. *Beibl.* 65.—Bras.
Hookeriana, *Hemsl. Biol. Centr. Am. Bot.* iii. 19.—Mexic.
mollis, *Hemsl.* l. c. 20.—Mexic.
Palmeri, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 144.—Am. bor. occ.
paniculata, *Benth. & Hook. f. Gen.* iii. 41.—Bras.
paniculata, *Mart. Nov. Gen. et Sp.* ii. 43.—Bras.
parviflora, *Benth. Bot. Voy. Sulph.* 156.—Porto Rico.
pulverulenta, *Benth. & Hook. f. Gen.* iii. 41.—Mexic
pulverulenta, *Mart. Nov. Gen. et Sp.* ii. 46.—Bras.
spicata, *Mart.* l. c. 44. t. 141.—Bras.
subnuda, *Hemsl. Biol. Centr. Am. Bot.* iii. 20.—Mexic.
vaga, *Benth. & Hook. f. Gen.* iii. 42.—Bras.
virgata, *Mart. Nov. Gen. et Sp.* ii. 45.—Bras.

HEBE, Comm. ex Juss. Gen. 105 (1789) = **Veronica**, Tourn. (Scroph.).
magellanica,] F. Gmel. Syst. 27 = V. elliptica.

HEBEA, Hedw. f. Gen. 24 (1806) = **Gladiolus**, Tourn. (Irid.).
alata, Eckl. Top. Verz. 42 = G. alatus.
bicolor, Eckl. l. c. = G. bicolor.
formosa, Eckl. l. c. = G. formosus.
gulzeta, Eckl. l. c. 41 = G. alatus.
orchidiflora, Eckl. l. c. 43 = G. arenarius.
pulchella, Eckl. ex Baker, in Journ. Linn. Soc xvi. (1877) 178 = G. pulchellus.
ramosa, Eckl. Top. Verz. 43 = G. montanus.
speciosa, Stend. Nom. ed. II. i. 724 = G. speciosus.
tabularis, Eckl. Top. Verz. 43 = G. montanus.
Zeyheri, Eckl. l. c. 42 = G. permeabilis.

HEBEANDRA, Bonpl. in Gen. Naturf. Fr. Berl. Mag. ii. (1808) 40 = **Monnina**, Ruiz & Pav. (Polygal.).
calastroides, Bonpl. l. c. 41 = M. rupestris.
cestrifolia, Bonpl. l. c. = M. cestrifolia.
crassifolia, Bonpl. l. c. 42 = M. crassifolia.
cuneymoides, Bonpl. l. c. = M. xalapensis.
fastigiata, Bonpl. l. c. 43 = M. parviflora.
hirta, Bonpl. l. c. = M. pilosa.
latifolia, Bonpl. l. c. = M. latifolia.
ligustrina, Bonpl. l. c. 40 = M. ligustrifolia ?
macrocnata, Bonpl. l. c. = M. nemorosa.
myrtilloides, Bonpl. l. c. = M. myrtilloides.
pedifolia, Bonpl. l. c. = M. phytolaccæfolia.
parviflora, Bonpl. l. c. 44 = M. parviflora.
phyllirzeoides, Bonpl. l. c. 41 = M. obtusifolia.
prinoides, Bonpl. l. c. 44 = M. ligustrifolia.
pubescens, Bonpl. l. c. 43 = M. pubescens.
revoluta, Bonpl. l. c. 41 = M. revoluta.

HEBEANTHE, Reichb. Nom. 145 (1841) = **Hebanthe**, Mart. (Amarant.).

HEBECLADUS, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 231. *SOLANACEÆ*, Benth. & Hook. f. ii. 896.
EUCOLIA, Rafin. Sylva Tellur. 56 (1838), nomen prius.
UEUCOMIA, Rafin. l. c. (1838).

HEBECLADUS:—
asperus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 232.—N. Granat.
bicolor, *Miers*, l. c.—Peruv.
bifloras, *Miers*, l. c.—Peruv.
granulosus, *Miers*, l. c. vii. (1848) 352.—N. Granat.
hirsellus, *Miers*, l. c. iv. (1845) 228.—Bras.
intermedius, Miers, l. c. = bifloras.
lanceolatus, Miers, l. c. = viridiflorus.
nitidus, *Miers*, l. c. vii. (1848) 352.—N. Granat.
sinuosus, *Miers*, l. c.—Peruv.
Turneri, Miers, l. c. iv. (1845) 228 = umbellatus.
umbellatus, *Miers*, l. c. 232.—Peruv.
ventricosus, *Baker, in Saund. Refug. Bot.* t. 208.—Peruv.
viridiflorus, *Miers, in Hook. Lond. Journ. Bot* iv. (1845) 231.—N. Granat.

HEBECLINIUM, DC. Prod. v. 186 (1836) = **Eupatorium**, Tourn. (Compos.)
atrorubens, Lem. Illustr. Hortic. ix. (1862) t. 310.—Mexic.
brevipeticulatum, Sch. Bip. ex Klatt, in Leopoldina, xx. (1884) 90 = E. brevipeticulatum.
Ehrenbergii, Sch. Bip. ex Hemsl. Biol. Centr. Am Bot. ii. 94 = E. Ehrenbergii.
hecatenthum, DC. Prod. v. 186 = E. hecatenthum.
ianthinum, Hook. Bot. Mag. t. 4574 = E. ianthinum.
Liebmannianum, Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot. ii. 96 = E. Liebmannii, *Hemsl.*
macrocephalum, Benth. Pl. Hartw. 42 = E. Ehrenbergii.
macrophyllum, DC. Prod. v. 186 = E. macrophyllum.
megalophyllum, Lem. Illustr. Hortic. ix. (1862) t. 310 = E. megalophyllum.
paniculatum, Carr. in Rev. Hortic. (1877) 330 = E. ianthinum !
tripicanum, Hook. & Arn. Bot. Beech. Voy. 434 = E. tupicanum.
tetragonum, Benth. Pl. Hartw. 198 = E. guadalupense.
Urolepis, DC. Prod. v. 186 = E. hecatenthum.
vitifolium, Sch. Bip. ex Klatt, in Leopoldina, xx (1884) 90 = E. vitifolium.

HEBECOCCA, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 146 = **Omphalea**, Linn. (Euphorb.).
panamensis, Beurl. l. c. = O. diandra.

HEBECOCCUS, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 246. *SAPINDACEÆ*.
ferrugineus, *Radlk. l. c.* 309 —Java.

HEBELIA, C. C. Gmel. Fl. Bad. ii. 117 (1806) = **Tofieldia**, Huds. (Liliac.).
alternifolia, C. C. Gmel. l. c. = T. calyculata.
collina, C. C. Gmel. l. c. 118 = T. calyculata.

HEBENSTREITIA, Linn. in Murr. Syst. ed. XIII. 476 (1774) – seq.

HEBENSTRETIA, Linn. Gen. ed. I. 248 (1737). *SELAGINEÆ*. Benth. & Hook. f. ii. 1127.
POLYCENIA, Choisy, in Mém. Soc. Phys. Genève. ii. 11. (1823) 91.
alba, Jacq. Eclog. Pl. ii. t. 151 = Dischisma ciliatum.
alchemilloides, Link, Enum. Hort. Berol. ii. 124 = pinac.
augusta, Pritz. Ic. Ind. 526, sphalm. = robusta.
aurea, Andr. Bot. Rep. t. 352 = dentata.
capensis, Hort. Berol. ex Choisy, in DC. Prod. xii. 6, ? = Dischisma clandestinum.
capitata, Thunb. Prod. Pl. Cap. 103 = Dischisma capitatum.
chamædrifolia, Link, Enum. Hort. Berol. ii. 125 = Dischisma erinoides.
ciliata, Berg. Desc. Pl. Cap. 154 ; Linn. Mant. ii. 420 = Dischisma ciliatum.
comosa, *Hochst. in Flora*, xxviii. (1845) 70.—Afr. austr.
cordata, *Linn. Mant. ii.* 420.—Afr. austr.
dentalis, *Choisy, in DC. Prod.* xii. 4.—Afr. austr.
dentata, *Linn. Sp. Pl.* 889.—Afr. bor. et austr.
discolora, *E. Mey. Comm. Pl. Afr. Austr.* 249.—Afr. austr.
dura, *Choisy, in DC. Prod.* xii. 4.—Afr. austr.
erinoides, Berg. ex Link, Enum. Hort. Berol. ii. 125, in obs. = Dischisma ciliatum.
erinoides, Thunb. Prod. Pl. Cap. 103 = Dischisma erinoides.

HEBENSTRETIA:—
fastigiata, Stend. Nom. ed. II. i. 724 = seq.
fastigiosa, *Varasa, Pl. Nov. Cap.* 14.—Afr. austr.
frutescens, Hort. Berol. ex Choisy, in DC. Prod. xii. 4 = dentata.
frutiosa, *Linn. f. Suppl.* 287 ; *Sims, Bot. Mag.* t. 1970.—Afr. austr.
glabra, Walp. Rep. iv. 161, sphalm. = Selago geniculata.
hamulosa, *E. Mey. Comm. Pl. Afr. Austr.* 249.—Afr. austr.
hispida, Lam. Encyc. iii. 78 = Dischisma ciliatum.
integrifolia, Linn. Sp. Pl. 629 = dentata.
macra, *E. Mey. Comm. Pl. Afr. Austr.* 248.—Afr. austr.
parviflora, *E. Mey. l. c.* 249.—Afr. austr.
polystachya, Herb. Linn. ex Rolfe, in Journ. Linn. Soc. xx. (1883) 349 = Selago polystachya.
pulchella, Sallsb. Prod. 93 = dentata.
ramosissima, *Varasa, Pl. Nov. Cap.* 14.—Afr. austr.
repens, *Varasa, l. c.* 15.—Afr. austr.
robusta, *E. Mey. Comm. Pl. Afr. Austr.* 240.—Afr. austr.
scabra, Hort. Berol. ex Choisy, in DC. Prod. xii 5 = dentata.
scabra, Sieber, ex Choisy, l. c. 11 = Selago geniculata
scabra, *Thunb. Prod. Pl. Cap.* 103.—Afr. austr.
scabrida, *Burm. f. Fl. Cap. Prod.* 17.—Afr. austr.
spicata, Thunb. Prod. Pl. Cap. 103 = Dischisma spicatum.
tenera, *Spreng. ex Walp. Rep.* iv. 143.—Afr. austr.
tenuifolia, E. Mey. Comm. Pl. Afr. Austr. 246 = crassifolia.
tenuifolia, Schrad. Cat. Hort. Gotting. ex Reichb. Hort Bot. ii. 13. t. 133 = dentata.
virgata, *E. Mey. Comm. Pl. Afr. Austr.* 249.—Afr austr.

HEBENPETALUM, Benth. in Benth. & Hook. f. Gen i. 244 (1862). *LINEÆ*, Benth. & Hook. f. i. c. humirifolia, *Benth. l. c.* 245.—Guian.
latifolia, *Benth. l. c.*—Bras.

HEBERDENIA, Banks, ex Vent. Choix, n. 5, verso (1808) = **Myrsine**, Linn.
excelsa, Banks, ex Roem. & Schult. Syst. iv. 508 = M. Heberdenia.

HEBOKIA, Rafin. Alsog. Am. 47 (1838) = **Eusaphis**, Steb. & Zucc. (Sapindac.).
japonica, Rafin. l. c. = E. staphyleoides.

HEBRADENDRON, R. Grah. in Hook. Comp. Bot. Mag. ii. (1836) 199. t. 27 = **Garcinia**, Linn. (Guttif.).
cambogioides, R. Grah. l. c. = G. Morella.
Choisyanum, Choisy, in Mém. Soc. Phys. Genèv. xii (1851) 419 = G. Choisyana.
elliptica, R. Grah. in Hook. Comp. Bot. Mag. ii. (1836) 200 = G. elliptica.
eugenifolium, Choisy, in Mém. Soc. Phys. Genèv. xii. (1851) 419 = G. eugenifolia.
Wallichii, Choisy, l. c. = G. heterandra.

HECABE, Rafin. Fl. Tellur. iv. 44 (1836) = **Phalus**, Lour. (Orchid.).
lutea, Rafin. l. c. = P. maculatus.

HECALE, Rafin. Fl. Tellur. fl. 79 (1836) = **Wahlenbergia**, Schrad. (Campanulac.).
lobelioides, Rafin. l. c. = W. lobelioides.

HECASTE, Soland. ex Schum. in Skr. Naturf. Selsk. iii (1793) 10 = **Bobartia**, Linn. (Irid.).
juncea, Soland. l. c. = B. indica.

HECASTOCLEIS, A. Gray, in Proc. Am. Acad. xvii. (1881–82) 220. *COMPOSITÆ*.
Shockleyi, *A. Gray, l. c.* 221.—Am. bor. occ.

HECASTOPHYLLUM, H. B. & K. Nov. Gen. et Sp. vi 367 (1823) = **Bonetaphyllum**, P. Br. (Legumin.).

HECATANDRA, Rafin. Sylva Tellur. 120 (1838) = **Acacia**, Tourn. (Legumin.).
Oxyodras, Rafin. l. c. = A. Oxyodras.
suaveolens, Rafin. l. c. = A. suaveolens.

HECATEA, Thou. Hist. Vég. Isles Fr. 27. t. 5 (1804) = **Omphalea**, Linn. (Euphorb.).
alternifolia, Willd. Sp. Pl. iv. 514 = O. biglandulosa.
biglandulosa, Pers. Syn. ii. 556 = O. biglandulosa.
oppositifolia, Willd. Sp. Pl. iv. 513 = O. biglandulosa.

HECATERIUM, Kunze, ex Reichb. Handb. 281 (1837) = praec. †

HECATONIA, Lour. Fl. Cochinch. 302 (1790); DC. Syst. i. 227 (1818) = **Ranunculus**, Tourn.
aconitifolia, Schur, Enum. Pl. Transs. 18 = R. aconitifolius.
alpestris, Schur, l. c. = R. alpestris.
crenata, Schur, l. c. = R. crenatus.
glacialis, Schur, l. c. 13 = R. glacialis.
palustris, Lour. Fl. Cochinch. 303 = R. sceleratus.
pilosa, Lour. l. c. = R. pensylvanicus.
platanifolia, Schur, in Verh. Naturf. Ver. Bruenn, xv. 11. (1877) 20 = R. aconitifolius.
sceleruta, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 325 = R. sceleratus.
Transfellneri, Schur, Enum. Pl. Transs. 13 = R. alpestris.

HECATOUNIA, Poir. in Dict. Sc. Nat. xx. 327 (1821) = praec.

HECATRIS, Salisb. Gen. Pl. Fragm. 66 (1866) = **Aspa-ragus**, Linn. (Liliac.).

HECETTA, Klotzsch, in Otto & Dietr. Allg. Gartenz. iii. (1885) 401. *BROMELIACEAE*, Benth. & Hook. f. iii. 667.
argentea, *Baker, in Bot. Mag.* sub t. 6554, nomen; *et in Hemsl. Biol. Centr. Am. Bot.* iii. 317.—Mexic.
Besseriana, *Versch. Cat.* n. 18 (1874).—Mexic.
cordylinoides, Baker, in Bot. Mag. t. 6554 = stenopetala.
Ghiesbreghtii, Lem. Illustr. Hortic. x. (1863) t. 378 = glomerata.
glomerata, *Zucc. in Abh. Akad. Muench.* iii. (1843) 240. t. 6.—Mexic.
pitcairnaefolia, Hort. Berol. ex Veriot, in Rev. Hortic. (1868) 211 = Rhodostachys pitcairnaefolia.
Schottii, *Baker, in Hemsl. Biol. Centr. Am. Bot.* iii. 318.—Mexic.
stenopetala, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* iii. (1885) 402.—Mexic.
texensis, *S. Wats. in Proc. Am. Acad.* xx. (1885) 374. —Texas.

HECKERIA, Kunth, in Linnaea, xiii. (1839) 554 = **Piper**, Linn.
peltata, Kunth, l. c. 555 = P. peltatum.
scutata, Kunth, l. c. 567 = P. peltatum.
sidaefolia, Kunth, l. c. 559 = P. subpeltatum.
speciosa, Kunth, l. c. 558 = P. peltatum.
subpeltata, Kunth, l. c. 571 = P. subpeltatum.
umbellata, Kunth, l. c. 559 = P. umbellatum.

HECKERIA, Rafin. Sylva Tellur. 165 (1838) = **Lidsea**, Lam. (Laurin.).
glomerata, Rafin. l. c. = L. glomerata.

HECTOREA, DC. Prod. v. 95 (1836) = **Chrysopsis**, Nutt. (Compos.).
villosissima, DC. l. c. = C. graminifolia †

HECTORELLA, Hook. f. Handb. N. Zeal. Fl. 27 (1864). *PORTULACEAE*, Benth. & Hook. f. i. 979.
caespitosa, *Hook. f. l. c.*—N. Zel.
elongata, *Y. Buch. in Trans. N. Z. Inst.* xvi. (1884) 395.—N. Zel.

HECURABA, DC. Prod. v. 665 (1836). *COM-POSITAE*, Benth. & Hook. f. ii. 400.
scorzoneraefolia, *DC. l. c.*—Mexic.

HEDAROMA, Lindl. Swan River App. 7. t. 2 B (1839) = **Darwinia**, Rudge (Myrtac.).
bicalvatum, Baines, in Gard. Chron. (1874) i. 601 = D. fimbriata.
Hookerii, Baines, l. c. = D. Hookeriana.
latifolium, Lindl. Swan River App. 7. t. 2 B = D. citriodora.
pinifolium, Lindl. l. c. = D. pinifolia.
thymoides, Lindl. l. c. = D. thymoides.
tulipiferum, Lindl. in Gard. Chron. (1854) 823 = D. macrostegia.

HEDBOMA, Pers. Syn. ii. 181 (1807). *LABIATAE*, Benth. & Hook. f. ii. 1168.
scinoides, *Schreb., in Linnaea*, xxii. (1849) 592.—Texas.
arbanasana, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 138 = Calamintha Nuttallii.
bracteolata, Nutt. Gen. Am. ii. Add. = Micromeria bracteolata.
brevifora, *Grisob. in Lechl. Berb. Am. Austr.* 58, nomen.—Perur.
clilata, *Benth. in DC. Prod.* xii. 245.—Texas.
ciliata, Nutt. in Jqarn. Acad. Philad. N. S. i. (1847) 138 = Drummondii.
coerulescens, *Benth. in DC. Prod.* xii. 245.—N. Granat.
costata, *Hemsl. Biol. Centr. Am. Bot.* ii. 547.—Mexic.
dentata, *Torr. Bot. Mex. Bound.* 130.—Am. bor. occ.
Drummondii, *Benth. Lab. Gen. et Sp.* 568.—Am. bor. occ.
Gilliesii, *Benth. l. c.* 567.—Chili.
glabra, Nutt. Gen. Am. i. 16 = Calamintha Nuttallii.
glabra, Pers. Syn. ii. 181 (err. typ. 151) = Calamintha glabella †
graveolens, *Chapm. ex A. Gray, in Proc. Am. Acad.* viii. (1873) 357.—Am. bor.
hirta, Nutt. Gen. Am. i. 15 = hispida.
hispida, *Pursh, Fl. Am. Sept.* ii. 414.—Am. bor.
hysopifolia, *A. Gray, in Proc. Am. Acad.* xi. (1876) 98.—Arizona.
incana, Torr. Bot. Mex. Bound. 130 = Poliomintha incana.
Instiana, *Wawra, in Oestr. Bot. Zeitschr.* xxxi. (1881) 70.—Bras.
Mandoniana, *Wedd. Chlor. And.* ii. 148.—Bolivia.
micrantha, Kegel, Gartenfl. (1864) 357 = Mosla cocymoides.
mollis, Torr. Bot. Mex. Bound. 129 = Poliomintha mollis.
multiflora, *Benth. Lab. Gen. et Sp.* 567.—Bras.
napalensis, Benth. l. c. 566 = Mosla cocymoides.
Palmeri, *Hemsl. Biol. Centr. Am. Bot.* ii. 548.—Mexic.
piperita, *Benth. Lab. Gen. et Sp.* 780.—Mexic.
pipertta, A. Gray, in Proc. Am. Acad. viii. (1873) 366 = thymoides.
pilcata, *Torr. Bot. Mex. Bound.* 130.—Texas.
pulegioides, *Benth. Lab. Gen. et Sp.* 567.—Bras.
pulegioides, *Perr. Syn.* ii. 181.—Am. bor.
purpurea, Kellogg, in Proc. Calif. Acad. v. (1873) 52 = Micromeria purpurea.
rotundifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 549.—Mexic.
serpylloides, Torr. in Pacif. Rail. Rep. iv. 128 = Poggonia serpylloides.
tenella, *Hemsl. Biol. Centr. Am. Bot.* ii. 549.—Mexic.
thymoides, *A. Gray, Syn. Fl. N. Am.* ii. 1. 362.—Am. bor. occ.
thymoides, Pers. Syn. ii. 181 (err. typ. 151) = Thymus Serpyllum.

HEDERA, Tourn. ex Linn. Gen. ed. I. 56 (1737). *ARALIACEAE*, Benth. & Hook. f. i. 946.
CISSODENDRON, F. Muell. Census, 61 (1882).
HELIX, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 224.
IRVINGIA, F. Muell. Fragm. v. 17 (1865).
KISSODENDRON, Seem. in Journ. Bot. iii. (1865) 201.
PEDERA, Neck. Elem. i. 158 (1790).
acerifolia, DC. Prod. iv. 264 = Oreopanax acerifolium.
aculeata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 187 = Brassiopsis aculeata.
acuminata, Wight, Ic. t. 1062 = Polyscias acuminata.
acutifolia, DC. Prod. iv. 258 (Quid?).—Am. austr.
aesculifolia, Wall. Cat. n. 4913 = Brassiopsis aesculifolia.
alaris, Schlecht. in Linnaea, ix. (1834) 609 = Dendropanax alaris.
angularis, DC. Prod. iv. 263 = Oreopanax angulare.
Aralia, Jacq. in Wall. Cat. n. 4925 = Trevesia palmata.
arborea, Sw. Fl. Ind. Occ. i. 518 = Dendropanax arboreum.
arborea, Walk. Fl. Carol. 109 = Vitis arborea.
argentata, DC. Prod. iv. 263 = Oreopanax argentatum.
argentata, Sprang. Syst. i. 952 = Oreopanax acerifolium.
aromatica, Blume, in Flora, vii. (1824) 291; DC. Prod. iv. 265 (= *Aralia aromatica*).—Java.
sanatralica, *F. Muell. Fragm.* iv. 120.—Austral.
avicenniaefolia, DC. Prod. iv. 263 = Oreopanax avicenniaefolium.
canariensis, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 171.—Ins. Canar.

HEDERA:—
capitata, Sm. Ic. Pict. Rar. i. t. 4 = Oreopanax capitatum.
catalpaefolia, DC. Prod. iv. 263 = Oreopanax capitatum.
cheirophylla, DC. l. c. 264 = Oreopanax cheirophyllum.
chrysocarpa, Walsh, in Trans. Hort. Soc. vi. (1826) 42 = H. Helix.
cochleata, Sweet, Hort. Brit. ed. I. 193 = Panax cochleatum.
colchica, C. Koch, in C. Koch & Fint. Wochenschr. (1859) 78, 406 = H. Helix.
communis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 491 = H. Helix.
corymbosa, Choisy, in DC. Prod. iv. 262 = Hydrocarpus corymbosa.
crassifolia, A. Gray, Bot. U. St. Expl. Exped. i. 718 = Pseudopanax crassifolium.
crassinervia, DC. Prod. iv. 263 = Oreopanax crassinervium.
cumanensis, DC. l. c. = Oreopanax cumanense.
cuneata, DC. l. c. 262 = Gilibertia cuneata.
cunefolia, *Griseb. Cat. Pl. Cub.* 117.—Cuba.
discolor, DC. Prod. iv. 263 = Oreopanax discolor.
disperma, DC. l. c. 265 = Macropanax oreophilum.
diversifolia, Stokes, Bot. Mat. Med. i. 456 = H. Helix.
donerailensis, Hort. ex C. Koch, Dendrol. i. 680 = H. Helix.
Echinops, Cham. & Schlecht. in Linnaea, v. (1830) 174 = Oreopanax Echinops.
elata, Buch.-Ham. in D. Don, Prod. Fl. Nep. 187 = Heptapleurum elatum.
emarginata, Moon, Cat. Pl. Ceyl. 18 = Heptapleurum emarginatum.
exaltata, Thw. Enum. Pl. Zeyl. 132 = Heptapleurum exaltatum.
ferruginea, DC. Prod. iv. 264 = Sciadophyllum ferrugineum.
ferruginea, Wall. Cat. n. 4909 = Trevesia palmata.
floribunda, DC. Prod. iv. 264 = Oreopanax floribundum.
floribunda, Wall. Cat. n. 4912 = Brassiopsis speciosa.
fragrans, D. Don, Prod. Fl. Nep. 187 = Pentapanax Leschenaultii.
frondosa, Salisb. Prod. 144 = Oreopanax capitatum.
Gaudichaudi, A. Gray, Bot. U. St. Expl. Exped. i. 719 = Panax Gaudichaudii.
glauca, Wall. Cat. n. 4921 = Pentapanax parasiticum.
glomerulata, DC. Prod. iv. 265 = Brassiopsis speciosa.
Hainla, Buch.-Ham. in D. Don, Prod. Fl. Nep. 187 = Brassiopsis Hainla.
Helix, *Linn. Sp. Pl.* 202.—Europ.; Afr. bor.; As. temp.
Helix, Lowe, Fl. Mader. 376 = canariensis.
heptaphyllum, Jungh. ex Miq. Fl. Ind. Bat. i. 1. 752 (= *Agalma rugosum*).—Java.
heterophylla, Wall. Cat. n. 4919 = Heptapleurum heterophyllum.
hypoglauca, Hance, in Walp. Ann. ii. 734 = Vitis cantonensis.
Ischiana, G. Don, Gen. Syst. iii. 894 = Arthrophyllum diversifolium.
japonica, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 507 = Dendropanax japonicum.
jatrophaefolia, *DC. Prod. iv.* 264.—N. Granat.
latifolia, Wight & Arn. Prod. 376 = Polyscias pinnata.
Leschenaultii, Wight & Arn. l. c. 377 = Pentapanax Leschenaultii.
Lessonii, A. Gray, Bot. U. St. Expl. Exped. i. 719 = Pseudopanax Lessonii.
lobata, Gilib. Fl. Lituan. ii. 184 = H. Helix.
lucescens, Seem. Journ. Bot. ii. (1864) 306 = seq.
lutescens, DC. Prod. iv. 265 = Heptapleurum octophyllum.
macrophylla, Wall. Cat. n. 4918 = Aralia cachemirica †
mayumaa, *Sprues, in Journ. Linn. Soc.* v. (1861) 9 (Quid †).—Bras.
media, Wall. Cat. n. 4924 = Heptapleurum Cephalotes †
multiflora, DC. Prod. iv. 262 = Oreopanax capitatum.
Mutisiana, DC. l. c. = Oreopanax Mutisianum.
nodosa, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 181 = Polyscias nodosa.
nutans, Sw. Prod. Veg. Ind. Occ. 61 = Dendropanax nutans.
obovata, Wight, Ic. t. 1011/13 = Heptapleurum stellatum.
obtusiloba, DC. Prod. iv. 263 = Oreopanax obtusilobum.
ovata, Wall. Cat. n. 4911 = Arthrophyllum diversifolium.
palmata, Wall. l. c. n. 4925 = Trevesia palmata.

HEDERA:—
paraslica, D. Don, Prod. Fl. Nep. 188 = Pentapanax paraslicum.
parviflora, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 129 = Dendropanax parviflorum.
pendula, Sw. Prod. Veg. Ind. Occ. 51 = Dendropanax pendulus.
platanifolia, DC. Prod. iv. 263 = Oreopanax platanifolium.
platyphylla, A. Gray, Bot. U. St. Expl. Exped. i. 730. t. 91 = Panax platyphyllum.
poltarum, Bertol. Fl. Ital. iii. 688 = *H. Helix.*
poetica, Salisb. Prod. 143 = *H. Helix.*
polyacantha, Wall. Pl. As. Rar. ii. 82 = Brassaiopsis palmata.
pretea, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 129 = Dendropanax parviflorum.
quinquefolia, Linn. Sp. Pl. 202 = Vitis hederacea.
racemosa, Wight, Ic. t. 1015 = Heptapleurum racemosum.
ramiflora, DC. Prod. iv. 262 = Gilibertia cuneata.
resinosa, Spruce, in Journ. Linn. Soc. v. (1861) 9; Benth. ex March, in Mart. Fl. Bras. xi. i. 249 = Gilibertia resinosa.
reticulata, DC. Prod. iv. 264 = Oreopanax reticulatum.
rhombea, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1848) 209 = *H. Helix.*
rhombifolia, Rupr. ex Boiss. Fl. Orient. ii. 1091 = *H. Helix.*
Rostgurniana, Hort. ex Boiss l. c. = *H. Helix.*
rostrata, Wight, Ic. t. 1018/14 = Heptapleurum rostratum.
rugosa, DC. Prod. iv. 265 (= *Agalma rugosum*).—Java.
scandens, DC. l. c. 264 = Brassaiopsis Hainla.
Sciadophyllum, Sw. Fl. Ind. Occ. i. 519 = Sciadophyllum Browniì.
senticosa, *Rupr. in Bull. Soc. Phys.-Math. Acad. Petersb.* xv. (1856) 134.—Hab.?
septemnervia, DC. Prod. iv. 263 = Oreopanax capitatum.
serrata, Wall. Cat. n. 4915 = Macropanax oreophilum.
sessiliflora, Benth. Fl. Hartw. 146 = Oreopanax sessiliflorum.
simillima, DC. Prod. iv. 265 (= *Agalma simillimum*).—Java.
soliusmii, Steud. Nom. ed. II. i. 735 = Sciadophyllum ferrugineum.
squarrosa, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 301 (= *Agalma rugosum*).—Java.
subcordata, Wall. Cat. n. 4917 = Pentapanax subcordatum.
terebinthinacea, Vahl, Symb. Bot. iii. 42 = Heptapleurum stellatum.
terebinthinacea, Wall. Cat. n. 4920 = Heptapleurum venulosum.
tomentosa, Boch.-Ham. ex D. Don, Prod. Fl. Nep. 187 = Heptapleurum impressum.
trifoliata, Wight & Arn. Prod. 377 = Pentapanax Leschenaultii.
triloba, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 105 = Gilibertia triloba.
turbacensis, DC. Prod. iv. 263 = Oreopanax turbacense.
umbelliflora, DC. l. c. 262 = Osmoxylon umbelliferum.
umbraculifera, Sweet, Hort. Brit. ed. I. 103 = Polyscias nodosa.
undulata, Wall. Cat. n. 4916 = Macropanax undulatum.
unifolia, Vell. Fl. Flum. 77; ii. t. 75 (Quid?).—Bras.
Vahlii, Thw. Enum. Pl. Zeyl. 192 = Heptapleurum stellatum.
venosa, Wall. Cat. n. 4928 = Heptapleurum venulosum.
verticillata, Zipp. ex Span. in Linnaea, xv. (1841) 208 = Heptapleurum verticillatum.
Wallichiana, Steud. Nom. ed. II. i. 735 = Oreopanax floribundum.
xalapensis, DC. Prod. iv. 264 = Oreopanax xalapense.

HEDERANTHUM, Steud. Nom. ed. II. ii. 830, in syn. (1841) = **Phrynema**, Linn. (Campan.).

HEDEROPSIS, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 739 (1879). *ARALIACEÆ.*
Malingeyi, *C. B. Clarke, l. c.*—Malacca.

HEDICHIUM, Ring. in Mark. Schrift. ii. (1831) 132 = Hedychium, Koen. (Scitam.).

HEDISARUM, Neck. Delic. Gallo-Belg. ii. 311 (1768) = Hedysarum, Tourn. (Leguminos.).

HEDIOSMUM, Poir. in Dict. Sc. Nat. xx. 330 (1821) = **Hedyosmum**, Sw. (Chloranth.).

HEDONA, Lour. Fl. Cochinch. 286 (1790) = **Lychnis**, Linn. (Caryophyll.).
sinensis, Lour. l. c. = L. coronata.

HEDRAEANTHUS, Griseb. Spicil. Fl. Rumel. ii. 292 (1844) = EDRAIANTHUS, A. DC. = **Wahlenbergia**, Schrad. (Campan.).

HEDRAIANTHERA, F. Muell. *Fragm. v. 58 (1865). MELIACEÆ.* [Benth. & Hook. f. i. 998].
porphyropetala, *F. Muell. l. c.*—Austral.

HEDRAIOPHYLLUM, Less. ex Steud. Nom. ed. II. i. 697, in syn. (1840) = **Goohnattia**, H. B. & K. (Compos.).

HEDRAIOSTYLUS, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 141 = **Plukenetia**, Linn. (Euphorb.).
corniculatus, Hassk. Cat. Hort. Bog. Alt. 234 = P. corniculata.
glaberrimus, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 141 = P. corniculata.

HEDRANTHUS, Rupr. in Bull Acad. Pétersb. xi. (1857) 208 = EDRAIANTHUS, A. DC. = **Wahlenbergia**, Schrad. (Campan.).
Owerinianus, Rupr. l. c. (= *E. Owerinianus*).—Reg. Caucas.

HEDUSA, Rafin. Sylva Tellur. 101 (1838) = **Dissotis**, Benth. (1849) (Melastom.).
grandiflora, Rafin. l. c. = D. grandiflora.

HEDWIGIA, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 495 = **Commelina**, Linn.
africana, Medic. l. c. = C. africana.

HEDWIGIA, Sw. Prod. Veg. Ind. Occ. 4, 62 (1788); Fl. Ind. Occ. ii. 670. t. 13 (1800). *BURSERACEÆ, Benth. & Hook. f. i. 326.*
CAPROXYLON, Tuss. Fl. Antill. iv. t. 20 (1827).
KNORREA, [Moc. & Sesse,] ex DC. Prod. ii. 80 (1825).
SCHWARGRICHENIA, Reichb. Consp. 147 (1828).
TETRAGASTRIS, Gaertn. Fruct. ii. 130. t. 109 (1791).
balsamifera, Engl. in Mart. Fl. Bras. xii. ii. 286. t. 60 = panamensis.
balsamifera, *Sw. Prod. Veg. Ind. Occ.* 62.—Ins. S. Doming.
Houmanni, *Engl. in DC. Monog. Phan.* iv. 97.—Guian.
mexicana, *DC. Prod.* ii. 80.—Mexic.
panamensis, *Engl. in Bot. Jahrb.* i. (1881) 42 = Panama.
rhoifolia, Benth. in Hook. Kew Journ. iii. (1851) 17 = Crepidospermum rhoifolium.
simplicifolia, Spreng. Neue Entdeck. iii. 24 = Trichilia simplicifolia.
Tussaccii, Walp. Rep. i. 559 = balsamifera.

HEDYCAPNOS, Planch. in Fl. des Serres, viii. (1852–53) 198. t. 820 = **Dicentra**, Bernh. (Papav.).
spectabilis, Planch. l. c. = D. spectabilis.

HEDYCARIA, Murr. Syst. ed. XIV. 894 (1784) = **Hedyoarya**, Forst. (Monim.).

HEDYCARIX, Raoul, Choix, 30 (1846), sphalm. = praec.

HEDYCARPUS, Jack, in Trans. Linn. Soc. xiv. (1823) 118 = **Bacoucurea**, Lour. (Euphorb.).
javanicus, Miq. Fl. Ind. Bat. i. ii. 859 = B. javanica.
lanceolatus, Miq. l. c. = B. lanceolata.
malayanus, Jack, in Trans. Linn. Soc. xiv. (1823) 118. = B. samarra.

HEDYCARYA, Forst. Char. Gen. 127. t. 64 (1776). *MONIMIACEÆ*, Benth. & Hook. f. iii. 141.
CRINONIA, Banks, ex Tul. in Arch. Mus. Par. viii. (1856) 405.
angustifolia, *A. Cunn. in Ann. Nat. Hist.* Ser. I. i. (1838) 215.—Austral.
arborea, *Forst. Char. Gen.* 128. t. 64.—N. Zel.
australasica, A. DC. in DC. Prod. xvi. ii. 673 = angustifolia.

HEDYCARYA:—
Badaunii, *Baill. Adansonia*, ix. (1868–70) 132.—N. Caled.
bengalensis, Roxb. ex Walp. Ann. iv. 113 = arborea.
Cunninghami, Tul. in Arch. Mus. Par. viii. (1855–56) 408 = angustifolia.
cupulata, *Baill. Adansonia*, ix. (1868–70) 132.—N. Caled.
dentata, Forst. f. Prod. 71 = arborea.
dorstenioides, *A. Gray, in Journ. Bot.* iv. (1866) 83 —Ins. Fiji.
hirsuta, Spreng. Pugill. ii. 91 = arborea.
macrophylla, A. Cunn. in Ann. Nat. Hist. Ser. I. i (1838) 215 = angustifolia.
Pseudomorus, F. Muell. in Trans. Phil. Inst. Vict. ii (1858) 62 = angustifolia.
racemosa, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 45 = Palmeria racemosa.
scabra, A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 216 = arborea.

HEDYCHION, Hassk. Cat. Hort. Bog. Alt. 50 (1844) = seq.

HEDYCHIUM, Koen. in Retz. Obs. iii. 73 (1788). *SCITAMINEÆ*, Benth. & Hook. f. iii. 642.
GAMOCHILUS, Lestib. in Ann. Sc. Nat. Sér. II. xv (1841) 341.
acuminatum, Rosc. Scitam. t. 47 = spicatum.
album, Buch.-Ham ex Wall. in Hook. Kew Journ. v (1853) 325 = coronarium.
angustifolium, Roxb. Hort. Beng. 1; Fl. Ind. i. 13 = coccineum.
aurantiacum, Rosc. Scitam. t. 61 = coccineum.
aurantiacum, Wall. Cat. n. 6551 = speciosum.
barbatum, Wall. l. c. n. 6544 = stenopetalum.
candidum, *Wall. ex Horan. Prod. Scitam.* 26, nomen (sp. dub.).—Ind. or.
carneum, *Carey, ex Herb. in Bot. Mag.* t. 2687; *Lodd. Bot. Cab.* t. 698.—Ind. or.
carneum, Rosc. Scitam. t. 57 = coccineum.
cernuum, Wight, Ic. t. 3011 = venustum.
chrysoleucum, Hook. Bot. Mag. t. 4516 = coronarium.
coccineum, *Buch.-Ham. ex Sm. in Rees, Cyclop.* xvii. n. 5; *Lodd. Bot. Cab.* t. 705.—Nepal.
coccineum, Wall. in Hook. Kew Journ. v. (1853) 373, partim = stenopetalum.
coronarium, *Koen. in Retz. Obs.* iii. 73.—Ind. or.
densiflorum, *Wall. Cat.* n. 6553.—Ind. or.
elatum, *R. Br. in Bot. Reg.* t. 226.—Nepal.
elatum, Horan. Prod. Scitam. 25, partim = stenopetalum.
ellipticum, *Sm. in Rees, Cyclop.* xvii. n. 2.—Reg. Himal.
fastigiatum, Wall. Cat. n. 6540 = ellipticum.
flavescens, Carey, ex Rosc. Scitam. t. 50 = coronarium
flavescens, Lodd. Bot. Cab. t. 738; Lindl. in Journ. Hort. Soc. vii. (1852) 281 = spicatum?
flavum, Roxb. Hort. Beng. 1; Fl. Ind. i. 31 = coronarium.
formosum, *Wall. ex Horan. Prod. Scitam.* 26, nomen (sp. dub.).—Ind. or.
Gandasulium, Buch.-Ham. ex Wall. in Hook. Kew Journ. v. (1853) 325 = coronarium.
Gardnerianum, *Griff. Notul.* iii. 419.—Ind. or.
Gardnerianum, *Rosc. Scitam.* t. 62.—Reg. Himal.
Gardnerianum, Wall. ex Spreng. Syst. iv. Cur. Post. 6 = speciosum.
gigauteum, *Wall. ex Horan. Prod. Scitam.* 26, nomen (sp. dub.).—Ind. or.
glaucum, Rosc. Scitam. t. 58 = gracile.
Gomezianum, *Wall. Cat.* n. 6548.—Ind. or.
gracile, *Roxb. Hort. Beng.* 1; *Fl. Ind.* i. 14.—Ind. or.
grandiflorum, *Regel, Cat. Pl. Hort. Akakow.* 68.—Hab.?
gratum, *Wall. ex Horan. Prod. Scitam.* 26, nomen (sp. dub.).—Ind. or.
Griffithianum, *Wall. in Hook. Kew Journ.* v. (1853) 309.—Ind. or.
Hasseltii, *Blume, Enum. Pl. Yav.* i. 56.—Java.
helvolum, Steud. Nom. ed. II. i. 736, sphalm. = *trilobum.*
heteromallum, Ker-Gawl. Bot. Reg. t. 767 = thyrsiflorum.
Horsfieldii, *R. Br. ex Wall. in Hook. Kew Journ.* v (1853) 376.—Java.
intermedium, *Blume, Enum. Pl. Yav.* 57.—Java.
jasminiflorum, Horan. Prod. Scitam. 26, nomen, sphalm. = luteo-virescens.

HEDYCHIUM :—
lanatum, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 57. —N. Guinea.
linguiatum, *Hassk. in Hoev. & De Vriese, Tijdschr.* v. (1838–39) 357.—Java.
longifolium, Roxb. Scitam. t. 59 = coccineum.
luteum, *Hort. Angl. ex Link, Enum. Hort. Berol.* i. 443.—Hab.?
maximum, Roxb. Scitam. t. 58 = coronarium.
palembanicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 101.—Java.
pallidum, Regel, Ind. Sem. Hort. Petrop. (1856) 23; Gartenfl. (1857) 266 = Gardnerianum.
peregrinum, *N. E. Br. in Gard. Chron.* (1883) i. 268. —Madag.
Prophetae, Buch.-Ham. ex Wall. in Hook. Kew Journ. v. (1853) 325 = coronarium.
Roscoei, Wall. ex A. Dietr. Sp. Pl. i. 37 (Roscoi) = speciosum, coccineum.
Roxburghii, *Blume, Enum. Pl. Jav.* i. 57.—Java.
Roxburghii, Siebold, ex Lindl. in Journ. Hort. Soc. vii. (1852) 281. in syn. = spicatum!
scaposum, Nimmo, in J. Grah. Cat. Pl. Bomb. 205 = Monolophus scaposus.
Sieboldii, Wall. in Hook. Kew Journ. v. (1853) 371 = spicatum!
simile, *Blume, Enum. Pl. Jav.* i. 57.—Java.
Spanoghenanum, *Wall. in Hook. Kew Journ.* v. (1853) 375.—Java.
speciosum, Wall. in Roxb. Fl. Ind. i. 18 = speciosum.
spicatum, Lodd. Bot. Cab. t. 659 = coronarium.
spicatum, *Sm. in Rees, Cyclop.* xvii. n. 3.—Reg. Himal.
squarrosum, Buch.-Ham. ex Wall. in Hook. Kew Journ. v. (1853) 372 = coccineum
stenopetalum, *Lodd. Bot. Cab.* t. 1902.—Ind. or.
sulphureum, Wall. in Hook. Kew Journ. v. (1853) 325 = coronarium.
sumatranum, *Yach, in Malay. Misc.* ii. (1822) n. VII. 1.—Malaya.
sumatranum, Miq. Fl. Ind. Bat. Suppl. 273 = palembanicum.
tavoyanum, *Horan. Prod. Scitam.* 26 (javoyanum), nomen.—Ind. or.
tenuiflorum, Wall. Cat. n. 6546 C = villosum.
thyrsiforme, *Sm. in Rees, Cyclop.* xvii. n. 4.—Ind. or.
Torache, Buch.-Ham. ex Wall. in Hook. Kew Journ. v. (1853) 327 = thyrsiforme.
trilobum, Wall. Cat. n. 6554 = spicatum.
urophyllum, Lodd. Bot. Cab. t. 1785 = coronarium.
venustum, *Wight, Ic.* t. 2012.—Ind. or.
villosum, *Wall. in Roxb. Fl. Ind.* i. 12.—Ind. or.

HEDYCHLOA, Rafin. Ann. Nat. 16 (1820) = **Kyllingа**, Rottb. (Cyperac.).
fragrans, Rafin. l. c. = K. pumila.

HEDYCREA, Schreb. Gen. 160 (1789) = **Licania**, Aubl. (Rosac.).
incana, J. F. Gmel. Syst. 498; Willd. Sp. Pl. i. 1210 = L. incana.

HEDYOSMON, Spreng. Gen. ii. 696 (1831) = **Hedyosmum**, Sw. (Chloranth.).

HEDYOSMOS, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 211 = **Zimphora**, Linn. (Rubiac.).

HEDYOSMUM, Sw. Prod. Veg. Ind. Occ. 84 (1788). *CHLORANTHACEAE*, Benth. & Hook. f. iii. 135.
TAFALLA, Ruiz & Pav. Prod. 136. t. 29 (1794).
acuifolium, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 306.—Brza.
angustifolium, *Solms, in DC. Prod.* xvi. i. 485.—Peruv.
arborescens, Cordem. ex Baill. Adansonia, iii. (1863–63) 304 = Spruceі.
arborescens, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 176 = Griesebachii.
arborescens, *Sw. Prod. Veg. Ind. Occ.* 84.—Ind. occ.
Artocarpus, *Solms, in DC. Prod.* xvi. i. 485.—Mexic.
boliviarum, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 306.—Bolivia.
Bonplandianum, *H. B. & K. Nov. Gen. et Sp.* viii. 165. t. 684.—Am. austr.
Bonplandianum, Mart. Syst. Mat. Med. Bras. 100 = brasiliense.
brasiliense, *Mart. Fl. Bras.* iv. i. 2. —Brza.
calloso-serratum, *Oerst. in Kjoeb. Vidensak. Meddel.* (1856) 40.—Am. centr.
cumbalense, *Karst. Fl. Columb.* ii. tt. 128, 129 —N. Granat.
Dombeyanum, *Solms, in DC. Prod.* xvi. i. 482.—Peruv.
elegans, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 308.—Ins. Guadaloup.
glabratum, Cordem. l. c. 308 = Gondotianum.
glabratum, *H. B. & K. Nov. Gen. et Sp.* vii. 164.—Am. austr.
glaucum, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 303.—Peruv.
glaucum, *Solms, in DC. Prod.* xvi. i. 485.—Peruv.
Gondotianum, *Solms, l. c.* 482.—N. Granat.
Griesebachii, *Solms, l. c.* 479.—Cuba.
hirsutum, *H. B. & K. Nov. Gen. et Sp.* vii. 164.—N. Granat.
integrum, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 302.—Peruv.
laciniatum, *Solms, in DC. Prod.* xvi. i. 485.—Peruv.
latifolium, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 303.—Peruv.
Lechleri, *Solms, in DC. Prod.* xvi. i. 484.—Peruv.
Mandoni, *Solms, l. c.* 480.—Bolivia.
mexicanum, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 307.—Mexic.
nutans, *Sw. Prod. Veg. Ind. Occ.* 84.—Ind. occ.
parvifolium, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 308.—N. Granat.
Pseudo-andromeda, *Solms, in DC. Prod.* xvi. i. 481.—Venezuela.
racemosum, Cordem. ex Baill. Adansonia, iii. (1863–63) 304 = integrum.
racemosum, *D. Don, in Edinb. Rev.* iii. 482; *ex Solms, in DC. Prod.* xvi. i. 485.—Peruv.
scabrum, *Solms, l. c.* 480.—Peruv.
Spruceі, *Solms, l. c.* 483.—Peruv.
Weddellianum, *Cordem. ex Baill. Adansonia*, iii. (1863–63) 306.—Brza.

HEDYOTIS, Linn. Diss. Dasa. 7 (1747); Amoen. Acad. i. 117 (1749). *RUBIACEAE*, Benth. & Hook. f. iii. 56.
DIMETIA, Meisn. Gen. 160 (1838).
DIPLOPHRAGMA, Meisn. l. c. (1838).
DIPYCNOPORA, Reinw. ex Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 167.
EDRISSA, Endl. Gen. Suppl. ii. 53 (1842) (EDRISIA, Rafin. §).
LEPTOPETALUM, Hook. & Arn. Bot. Beech. Voy. 295 (1841).
MACRANDRIA, Meisn. Gen. 160 (1838).
METABOLUS, Blume, Bijdr. 890 (1826).
PELTOSPERMUM, Benth. in Hook. Niger Fl. 400 (1849).
SCLEROCOCCUS, Barel. Ord. 210 (1830).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 368 = Oldenlandia abyssinica.
accedens, *Miq. Fl. Ind. Bat.* ii. 258.—Ins. Bangka.
acerosa, Gray, Pl. Wright. i. 81 = Houstonia acerosa.
acuminata, Stend. Nom. ed. II. i. 726 = Kadua acuminata.
acutangula, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 171.—Hongkong.
Adscensionis, *DC. Prod.* iv. 419.—Ins. Adscens.
affinis, Roem. & Schult. Syst. iii. 194 = Oldenlandia diffusa.
affinis, Wight & Arn. Prod. 411 = Anotis Leschenaultiana.
alata, Kocn. ex Wall. Cat. n. 6196 = Oldenlandia alata.
albo-nervia, *Bedd. Ic. Pl. Ind. Or.* t. 1. t. 3.—Ind. or.
alsinifolia, K. Br. in Wall. Cat. n. 878 = Oldenlandia corymbosa.
alsinifolia, Zipp. ex Span. in Linnaea, xv. (1841) 317 = Oldenlandia paniculata.
amatymbica, Stend. Nom. ed. II. i. 726 (Oldenlandiae sp.).—Afr. austr.
ambrosioides, *Fotsh. in Act. Lit. Univ. Hafn.* i. (1778) 256.—Guian.
americana, Jacq. Enum. Pl. Carib. 13 = rupestris.
amplifiora, *Hance, in Journ. Bot. xvii.* (1879) 11.—China.
andamanica, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 211.—Ins. Andaman.
angustifolia, Cham. & Schlecht. in Linnaea, iv. (1829) 156 = umbellifera.
angustifolia, *Miq. Fl. Ind. Bat.* ii. 102.—Ins. Philipp.
angustifolia, Wall. Cat. n. 6906 = Oldenlandia brachiata.

HEDYOTIS :—
approximata, R. Br. in Wall. Cat. n. 852 = umbellifera.
arborea, *Roxb. in Beatson, Tracts St. Helena, App.* 310.—Ins. S. Helena.
arborescens, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 17 = Calicarpa lanata?
arenariodes, *Steud. Nom.* ed. II. i. 726.—Madag.
argentea, Wall. Cat. n. 858 = congesta.
argyta, *R. Br. in Wall. Cat.* n. 854.—Burma.
articularis, *R. Br. l. c.* n. 854.—Ind. or.
aspera, Bedd. Ic. Pl. Ind. Or. t. 32 = Oldenlandia gracilis.
aspera, Heyne, ex Roth, Nov. Pl. Sp. 94 = Oldenlandia aspera.
asperuloides, *Benth. Bot. Voy. Sulph.* 19. t. 18.—Am. trop.
attenuata, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 216 = Oldenlandia brachiata.
Auricularia, *Linn. Sp. Pl.* 101.—As. et Austral. trop.
Auricularia, Wall. Pl. Carol. 85 = Oldenlandia glomerata.
axillaris, Gardn. ex Thw. Enum. Pl. Zeyl. 148 = nodulosa.
barbata, *Korth. in Nederl. Kruidk. Arch.* ii. 11. (1851) 160.—Borneo.
Beddomei, *Hook. f. Fl. Brit. Ind.* iii. 53.—Ind. or.
bicornuta, *Bedd. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 405.—Ins. Socotr.
biflora, Homem. Hort. Hafn. 130; Roth, Nov. Pl. Sp. 99 = Oldenlandia corymbosa.
biflora, Lam. Tabl. Encyc. i. 272; Spreng. Pugill. ii. 31 = Oldenlandia biflora.
Blumeana, *Steud. Nom.* ed. II. i. 726 = Oldenlandia paniculata.
boerhaavioides, Hance, in Journ. Bot. viii. (1870) 78 = Anotis boerhaavioides.
Boyeri, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 282 = Oldenlandia Boyeri.
borneensis, *Miq. Fl. Ind. Bat.* ii. 184.—Borneo.
borrerioides, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 171 = uncinella.
Boscii, DC. Prod. iv. 430 = Oldenlandia Boscii.
bowaridoides, Sccm. Bot. Voy. Herald, 296. t. 64 = Houstonia bowaridoides.
brachiata, Miq. ex Hook. f. Fl. Brit. Ind. iii. 60 = coerulea.
brachiata, Wight, in Wall. Cat. n. 6901 = Oldenlandia brachiata.
brachyloba, Sond. in Linnaea, xxiii. (1850) 50 (Oldenlandiae sp.).—Afr. austr.
brachypoda, R. Br. in Wall. Cat. n. 874 = Oldenlandia diffusa.
bracteosa, *Hance, in Journ. Bot. xxiii.* (1885) 333.—China.
Burmanniana, R. Br. in Wall. Cat. n. 866, partim = Oldenlandia corymbosa.
Burmanniana, Schult. Mant. iii. 132 = Oldenlandia paniculata.
buxifolia, *Bedd. in Madr. Journ.* Ser. III. i. (1868) 49. —Ind. or.
cachiensis, *Korst. in Linnaea*, xxx. (1859–60) 145.—N. Granat.
caerulea, Hook. Fl. Scr. Am. i. 286 = Houstonia caerulea.
caerulea, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 160 = venitis.
caerulea, *Wight & Arn. Prod.* 412.—Ind. or.
caerulescens, *F. Muell. Progrs.* iv. 38.—Austral.
campbiana, Walp. Rep. vi. 56 = Oldenlandia Schimperi.
campa, Stend. Nom. ed. II. i. 726 = Oldenlandia caffra.
calycina, Wall. Cat. n. 875 = Anotis calycina.
calycosa, Shuttl. ex A. Gray, Pl. Wright. i. 81, in note = Houstonia purpurea.
campanuliflora, Hook. Bot. Mag. t. 5340 = Coccocypselum campanuliforme.
capensis, Lam. Tabl. Encyc. i. 271 = Oldenlandia capensis.
capillipes, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 72 = Anotis Rheedii.
capitata, Bedd. Ic. Pl. Ind. Or. i. 44. t. 191 = Beddomei.
capitata, Heyne, ex Wall. Cat. n. 870 = Oldenlandia umbellata.
capitata, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 159 = borneensis.
capitata, *Lam. Encyc.* iii. 80.—Ind. or.
capitata, Walp. Rep. vi. 56 = Hentreyiana.

HEDYOTIS :—

capitata, Wight, ex Wall. Cat. p. 214, n. 876 = Anotis quadrifoliacaria.
capitellata, *Wall. Cat.* n. 837.—Malaya; China.
capitulifloræ, Miq. Fl. Ind. Bat. ii. 158 = ventita.
capituligera, *Hance, in Journ. Bot.* xvii. (1879) 12.—China.
caracasana, H. B. & K. Nov. Gen. et Sp. iii. 398 = Mallostoma caracasana.
carnosa, Dalz. in Hook. Kew Journ. ii. (1850) 135 = Anotis carnosa.
carnosa, *Korth. in Nederl. Kruidk. Arch.* ii. 11. (1851) 161.—Borneo.
carolinæana, Raeusch. Nom. ed. III. 88 = Houstonia purpurea.
centranthoides, Steud. Nom. ed. II. i. 727 = Kadua centranthoides.
cephalantha, *Wedd. Chlor. And.* ii. 46.—N. Granat.
cephalophora, R. Br. in Wall. Cat. n. 842 = uncinella.
Cephalotes, Hochst. in Flora, xxvii. (1844) 558 (Oldenlandiæ sp.).—Afr. austr.
Cervantesii, *H. B. & K. Nov. Gen. et Sp.* iii. 390.—Mexic.
Chamissoniana, Steud. Nom. ed. II. i. 727 = Gouldia sandwicensis.
chilœnsis. *Phil. in Linnæa*, xxviii. (1856) 697.—Chili.
chlorophylla, Hochst. in Flora, xxvii. (1844) 558 (Oldenlandiæ sp.).—Afr. austr.
ciliicaulis, *Miq. Fl. Ind. Bat.* ii. 170.—Ins. Sumbawa.
ciliolata, Torr. ex Spreng. Syst. iv. Cur. Post. 40 = Houstonia purpurea.
ciliolata, Walp. Ann. ii. 773 = Oldenlandia ciliolata.
cinereo-viridis, *Thw. Enum. Pl. Zeyl.* 149.—Zeylan.
coarctata, Willd. ex Roem. & Schult. Syst. iii. 527 = Mallostoma thymifolia.
coccinea, Walp. Rep. ii. 501 = Oldenlandia coccinea.
coerulea, Anct. vide *cærulea*.
commutata, *Schult. Mant.* iii. 134.—Porto Rico.
conferta, Ruiz & Pav. Fl. Per. i. 57 = Mallostoma conferta.
congesta, *R. Br. in Wall. Cat.* n. 844.—Malava.
connata, Hook. f. Fl. Brit. Ind. iii. 62, err. typ. = coronata.
consatyla, Gaudich. in Freyc. Voy. Bot. 471. t. 94 = Kadua Mendiziana.
consanguinea, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 231.—China.
Cookiana, Steud. Nom. ed. II. i. 727 = Kadua Cookiana.
cordata, Steud. l. c. = Cocciocyselum umbellatum.
cordata, *Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 177.—Japon.
cordifolia, Spreng. Pugill. ii. 26 = Rondeletia virgata.
coriacea, Sm. in Rees, Cycl. xvii. n. 11 = Kadua Menziesiana.
coronata, *Wall. Cat.* n. 856.—Burma.
corymbosa, Herb. Madr. ex Wall. Cat. n. 6302 = pruinosa.
corymbosa, Lam. Tabl. Encyc. i. 272 = Oldenlandia corymbosa.
corymbosa, Spreng. Pugill. ii. 27 = fruticosa.
costata, R. Br. in Wall. Cat. n. 849 = H. Auricularia.
costata, Kurz, in Journ. As. Soc. Beng. xiv. (1876) 11. 135 = ventita.
crassifolia, Bartl. ex DC. Prod. iv. 427 = Oldenlandia crassifolia.
crassifolia, *Blume, Bijdr.* 971.—Java.
crassifolia, Raña. Fl. Ladov. 77 (Houstoniæ sp.).—Am. bor.
Crateagoneura, *Spreng. Pugill.* ii. 85.—Malaya.
Crouchiana, *F. Muell. Fragm.* x. 85.—Austral.
cyanantha, Kurz, in Journ. As. Soc. Beng. xiv. (1876) 11. 136 = cærulea.
cyanescens, *Thw. Enum. Pl. Zeyl.* 148.—Zeylan.
cymosa, R. Br. in Wall. Cat. n. 885 = Anotis Leschenaultiana.
cymosa, *Thw. Enum. Pl. Zeyl.* 149.—Zeylan.
cymosa, Spreng. Pugill. ii. 30 = Oldenlandia lactea.
cynanchica, Steud. Nom. ed. II. i. 727 (= *Kohautia cynanchica*).—Afr. austr.
debilis, Roem. & Schult. Syst. iii. 200 = Oldenlandia debilis.
decumbens, Hochst. in Flora, xxvii. (1844) 552 = Oldenlandia decumbens.
Dekenavriana, Steud. Nom. ed. II. i. 727 = gracilis.
deltoidea, Wight & Arn. Prod. 410 = Anotis Leschenaultiana.
densiflora, *Baj. Hort. Mauric.* 107, nomen.—Afr. trop.

HEDYOTIS :—

dentelloides, *Steud. Nom.* ed. II. i. 727.—Bras.
Deppeana, Steud. l. c. = Oldenlandia Deppeana.
depressa, Roem. & Schult. Syst. iii. 200 = Oldenlandia corymbosa ?
diantha, Schult. Mant. iii. 136 = Oldenlandia corymbosa.
dichotoma, Cav. Ic. vi. 58 = racemosa.
dichotoma, Koen. ex Roth, Nov. Pl. Sp. 98 = Oldenlandia dichotoma.
dichotoma, A. Rich. Tent. Fl. Abyss. i. 361 = Oldenlandia Heynii.
diffusa, Spreng. Syst. i. 413 = Oldenlandia dichotoma.
diffusa, Willd. Sp. Pl. i. 566 = Oldenlandia diffusa.
discolor, Spreng. Syst. i. 411 = Rondeletia discolor.
effusa, *Hance, in Journ. Bot.* xvii. (1879) 11.—China.
elatinoides, *Benth. Fl. Austral.* iii. 405.—Austral.
elegans, *Wall. Cat.* n. 837.—Burma.
elongata, R. Br. in Wall. l. c. n. 865 = Oldenlandia aspera.
ericoides, *Willd. ex Roem. & Schult. Syst.* iii. 527.—Am. austr.
evena, *Thw. Enum. Pl. Zeyl.* 140, 419.—Zeylan.
extensa, R. Br. in Wall. Cat. n. 869 = Oldenlandia diffusa.
fascicularis, *Bertol. Misc. Bot.* x. 24.—Am. bor.
filiformis, Ruiz & Pav. Fl. Per. i. 57 = Mallostoma filiformis.
fimbriata, *Wall. Cat.* p. 213, n. 851.—Reg. Himal.
Finlaysoniana, Wall. l. c. n. 6189 = capitellata, Anotis Wightiana.
flaccescens, Thw. Enum. Pl. Zeyl. 141 = Lessertiana.
floribunda, *Karst. in Linnæa*, xxx. (1859–60) 146.—N. Granat.
foetida, Dalz. in Hook. Kew Journ. ii. (1850) 134 = Anotis foetida.
fœtida, Spreng. Pugill. ii. 26 = Oldenlandia fœtida.
Frankii, *Presl, Bot. Bemerk.* 86.—Am. bor.
fruticosa, *Linn. Sp. Pl.* 150.—Ind. or.
fruticosa, [Moç. & Sesse,] ex DC. Prod. iv. 365 = Bouvardia triphylla.
fruticosa, Walp. Rep. iv. 450. Ind. sphalm. = Houstonia angustifolia.
fugax, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 232 = Oldenlandia decumbens.
fulva, *Hook. f. Fl. Brit. Ind.* iii. 58.—Reg. Himal.
fusca, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 134 = Oldenlandia gracilis.
galioides, F. Muell. Fragm. iv. 36 = Oldenlandia galioides.
galioides, Wall. Cat. n. 866 = H. Wallichii.
Gardneri, *Thw. Enum. Pl. Zeyl.* 142.—Zeylan.
geminiflora, Sond. in Linnæa, xxiii. (1850) 51 (Oldenlandiæ sp.).—Afr. austr.
geminulata, *Rafin. New Fl. Am.* ii. 21.—Am. bor.
gracilenta, Roth. Hort. Beng. 88; Fl. Ind. i. 364 = H. Auricularia.
gentianoides, Endl. Iconog. Gen. Pl. i. 89 = Houstonia cærulea.
Gerrardi, Sond. in Harv. & Sond. Fl. Cap. iii. 9 = Oldenlandia caffra.
globella, R. Br. in Wall. Cat. n. 886 = nitida.
Gluhea, *Hook. f. & Thoms. ex Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* iii. 62.—Reg. Himal.; Malaya.
globosa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 360 = Oldenlandia globosa.
glomerata, Ell. Sketch, i. 135 = Oldenlandia glomerata.
glomerata, Walp. Rep. ii. 501 = Kadua glomerata.
goreensis, DC. Prod. iv. 421 = Oldenlandia trinervia.
gracilis, *DC. l. c.* 419.—N. Caled.
gracilicaulis, Wall. in Roxb. Fl. Ind. ed. Carey, i. 371 = Oldenlandia gracilis.
graminicola, Kurz, in Journ. Bot. xiii. (1875) 296 = Oldenlandia corymbosa.
grandiflora, Lam. f. Suppl. 119 = Oldenlandia stricta.
grandiflora, A. Rich. Tent. Fl. Abyss. i. 365 = Oldenlandia noctiflora.
grandiflora, Steud. Nom. ed. II. i. 727 = Oldenlandia grandiflora.
Gryl, *Hook. f. in Benth. & Hook. f. Gen.* ii. 57.—Ins. Bonin.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 58.—Reg. Himal.
Haleï, Torr. & Gray, Fl. N. Am. ii. 42 = Oldenlandia Haleï.
Hartwegiana, *Wedd. Chlor. And.* ii. 45.—N. Granat.
Helferi, *Hook. f. Fl. Brit. Ind.* iii. 58.—Burma.
herbacea, Bert. ex Schult. Mant. iii. 134 = commutata.
herbacea, Forsk. Fl. Aegypt. Arab. p. cv = Oldenlandia stricta.

HEDYOTIS :—

herbacea, Linn. Sp. Pl. 102 = Oldenlandia Heynii ?
herbacea, Osbeck, Voy. ed. Angl. 4 = Oldenlandia corymbosa.
Heynii, Bedd. Ic. Pl. Ind. Or. t. 33 = Oldenlandia diohotoma.
Heynii, R. Br. in Wall. Cat. n. 867 = Oldenlandia Heynii.
hirsuta, Lam. Encyc. iii. 79 = H. Auricularia.
hirsuta, *Spreng. Pugill.* ii. 85.—Reg. Himal.; Java; Japon.
hirsutissima, *Bedd. in Made. Journ. Sc.* Ser. III. i. (1864) 49.—Ind. or.
hirtula, Harv. ex Sond. in Harv. & Sond. Fl. Cap. iii. 12 (Oldenlandiæ sp.).—Afr. austr.
hispida, Heyne, ex Wall. Cat. n. 886 = nitida.
hispida, *Retz. Obs.* iv. 28.—Ind. or.; As. trop.; Malaya.
hispida, Roth, Nov. Pl. Sp. 95 = Oldenlandia hispida.
Hookeriana, Steud. Nom. ed. II. i. 727 = Kadua glomerata.
Horneriana, Miq. Fl. Ind. Bat. ii. 158 = barbata.
Humboldtiana, Steud. Nom. ed. II. i. 727 = Mallostoma nitida.
humifusa, Engelm. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 218 = Houstonia humifusa.
hypnoides, *H. B. & K. Nov. Gen. et Sp.* iii. 389.—Ecuador.
hyssopifolia, Cav. Ic. vi. 54 = Mallostoma hyssopifolia.
inamœna, *Thw. Enum. Pl. Zeyl.* 143, partim.—Zeylan.
inamœna, Thw. l. c. partim = cinereo-viridis.
indica, Roem. & Schult. Syst. iii. 195 = Oldenlandia umbellata.
ingrata, Wall. Cat. n. 868 = Anotis ingrata.
intermedia, Wight & Arn. Prod. 415 = Oldenlandia corymbosa.
iodoneura, *Miq. Fl. Ind. Bat.* ii. 181.—Java.
juniperifolia, Ruiz & Pav. Fl. Per. i. 57 = Mallostoma juniperifolia.
lactea, Willd. Enum. Hort. Berol. 149 = Oldenlandia lactea.
lacustris, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 80.—Bras. lœvigata, *Miq. Fl. Ind. Bat.* ii. 178.—Ins. Philipp.
lances, *Thunb. ex Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 161.—China.
lancolata, Poir. Encyc. Suppl. iii. 14 = Houstonia purpurea.
lancifolia, Dalz. in Hook. Kew Journ. ii. (1850) 135 = Anotis lancifolia.
lancifolia, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 72 = Oldenlandia lancifolia.
Lapeyrousii, *DC. Prod. iv.* 430.—Ins. Vanikoro.
laricifolia, Cav. Ic. vi. 54 = Mallostoma lariclifolia.
lasiantha, Nutt. ex A. Gray, Pl. Wright. i. 81 = Houstonia angustifolia.
latifolia, Dalz. in Hook. Kew Journ. ii. (1850) 138 = Anotis Rheedii.
latifolia, *Reinw. ex De Vriese, Pl. Ind. Bat. Reinw.* 159.—Java.
latifolia, Walp. Rep. vi. 65 = Oldenlandia latifolia, *Mart. & Gal.*
Lawsoniæ, Benth. ex Hook. Lond. Journ. Bot. i. (1842) 486 = acutangula.
Lawsoniana, Wight, Ic. t. 1026 (Planta dubia).—Ind. or.
Lawsoniæ, *Wight & Arn. Prod.* 407.—Ind. or.
laxa, *D. Don, Prod. Fl. Nep.* 135 (Spermacoces sp. !).—Nepal.
laxiflora, Walp. Ann. ii. 773 = Oldenlandia macrophylla.
lenticinosa, Bedd. Ic. Pl. Ind. Or. t. 2. t. 6 = pruinosa.
leptostela, A. Gray, in Mém. Am. Acad. N. S. vi. (1858–59) 394 = Gray.
leptopetala, *Hook. f. in Benth. & Hook. f. Gen.* ii. 57.—Mexic.
Leschenaultiana, Hook. f. Fl. Brit. Ind. iii. 61, sphalm. vide *Lechenaultii*.
Leschenaultiana, Wight & Arn. Prod. 411 = Anotis Leschenaultiana.
Lechenaultii, DC. Prod iv. 422 = stylosa.
Lessertiana, *Arn. in Nov. Act. Nat. Cur.* xviii. i. (1836) 339.—Zeylan.
leucophylla, Spreng. Syst. i. 411 = Rondeletia leucophylla.
linariifolia, R. Br. in Wall. Cat. n. 870 = Oldenlandia umbellata.
Lindleyana, Hook. ex Wight & Arn. Prod. 409 = hirsuta.

HEDYOTIS :—
linearis, Stend. Nom. ed. II. i. 738 = Oldenlandia Heynii !
lineata, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 865.—Reg. Himal.; Burma.
linearis, Wall. Cat. n. 6198 = H. Auricularia.
linifolia, Willd. ex Roem. & Schult. Syst. iii. 526 = Oldenlandia dichotoma.
linoides, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 184, sphalm. = Oldenlandia linoides.
logeniicides, *Benth. Fl. Hongk.* 169.—Hongkong.
longidens, *Hance, in Journ. Bot.* xx. (1882) 289.—China.
longiflora, Spreng. Pugill. ii. 27 = Rondeletia pilosa.
longiflora, Stend. Nom. ed. II. i. 738 (= *Kohautia longiflora*).—Afr. austr.
longiflora, Walp. Rep. ii. 497 = Houstonia triflora.
longifolia, Hook. f. Bot. Am. i. 286 = Houstonia purpurea.
longifolia, Schum. & Thonn. Beskr. Guin. Pl. 70 = Oldenlandia corymbosa.
Macraei, *Hook. f. Fl. Brit. Ind.* iii. 50.—Zeylan.
macrocephala, K. Br. in Wall. Cat. n. 846, sphalm. = monocephala, *Wall.*
macrophylla, Lepr. & Perr. ex DC. Prod. iv. 427 = Oldenlandia macrophylla.
macrophylla, Thw. Enum. Pl. Zeyl. 149 = Thwaitesii. macrophylla, *Wall. Cat.* n. 841.—Burma; Malaya.
macrostemon, *Hook. & Arn. Bot. Beech. Voy.* 192.—China.
Maingayi, *Hook. f. Fl. Brit. Ind.* iii. 49.—Malacca.
marginata, Thw. ex Hook. f. Fl. Brit. Ind. iii. 58 = Leucertiana.
maritima, Linn. f. Suppl. 119 - Peplidium humifusum.
maritima, Wall. Cat. n. 6192 = Oldenlandia stricta, Shuteri.
media, Cav. Ic. vi. 58 = racemosa.
membranacea, *Thw. Enum. Pl. Zeyl.* 143.—Zeylan.
Menziesiana, Stend. Nom. ed. II. i. 738 = Kadua Menziesiana.
merguensis, Hook. f. in Benth. & Hook. f. Gen. ii. 57 = coronata.
micrantha, Hochst. ex Hiern, in Oliver, Fl. Trop. Afr. iii. 60 = Oldenlandia Heynii.
microphylla, DC. Prod. iv. 433, in syn. = Anotis serpens.
microphylla, *H. B. & K. Nov. Gen. et Sp.* iii. 389.—Peruv.
microthoca, Stend. Nom. ed. II. i. 738 = Oldenlandia microthoca.
minima, Heyne, ex Wall. Cat. n. 877 = Oldenlandia paniculata.
minima, Torr. & Gray, Fl. N. Am. ii. 38 = Houstonia patens, minima.
minuta, Phil. ex Benth. & Hook. f. Gen. ii. 58 = Oldenlandia uniflora.
mitrascmoides, *F. Muell. Fragm.* iv. 37.—Austral.
mollis, *Wall. Cat.* n. 859.—Ins. Penang.
monanthos, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 339 = Oldenlandia monanthos.
monocephala, *R. Br. in Wall. Cat.* p. 213.—Ind. or.
monocephala, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 236.—Java.
monosperma, Wight & Arn. Prod. 410 = Anotis monosperma.
macronata, *Benth. Bot. Voy. Sulph.* 19. i. 18.—Calif.
multicaulis, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 59 = H. Auricularia.
multiflora, Cav. Ic. vi. 58 = Oldenlandia paniculata.
muscosa, *St. Hil. Voy. Distr. Diam.* i. 206.—Bras.
mutica, *Wedd. Chlor. And.* ii. 48.—N. Granat.
myosuroides, Heyne, ex Wall. Cat. n. 882 = Anotis monosperma.
natalensis, Hochst. in Flora, xxvii. (1844) 552 (Oldenlandiae sp.).—Afr austr.
Nessiana, Arn. in Nov. Act. Nat. Cur. xvii. i. (1836) 841 = stricta.
nervosa, Lam. Encyc. iii. 79 = H. Auricularia, hispida.
stricta, H. B. & K. Nov. Gen. et Sp. iii. 392 = Mallostema stricta.
asitica, *Wight & Arn. Prod.* 412.—Ind. or.; Burma.
nodiflora, Wall. Cat. n. 855 = macrophylla.
nodulosa, *Arn. in Nov. Act. Nat. Cur.* xvii. i. (1836) 840.—Zeylan.
nodulosa, Thw. Enum. Pl. Zeyl. 148, partim = Allaeophania decipiens, Arnottii.
nodicaulis, Wight & Arn. Prod. 416 = Oldenlandia nudicaulis.
HEDYOTIS :—
nummularia, Arn. in Nov. Act. Nat. Cur. xvii. i. (1836) 841 = Anotis nummularia.
nummulariformis, Arn. l. c. = Anotis nummularia.
obscura, *Thw. Enum. Pl. Zeyl.* 141.—Zeylan.
orbiculata, Wall. Cat. n. 6191 = Oldenlandia trinervia.
ovalis, *Walp. Ann.* ii. 769.—Borneo.
ovata, *Thunb. ex Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 161.—China.
ovata, Thw. ex Hook. f. Fl. Brit. Ind. iii. 52 = obscura.
ovatifolia, Cav. Ic. vi. 52 = Oldenlandia nudicaulis.
oxyphylla, Wall. Cat. n. 6193 = Anotis oxyphylla.
paniculata, Lam. Encyc. iii. 79; Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 184 = Oldenlandia paniculata.
paniculata, Rottl. ex Wight & Arn. Prod. 417 = Oldenlandia aspera.
paradoxa, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 135.—Ins. Andaman.
parietarioides, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 235.—Ins. Moluce.
Parryi, *Hance, in Ann. Sc. Nat. Sér.* IV. xviii. (1862) 221.—China.
parviflora, Walp. Ann. ii. 772 = Oldenlandia virgata, parviflora.
parvifolia, *Presl, Bot. Bemerk.* 86.—Bras.
pauciflora, *Bartl. ex DC. Prod.* iv. 419.—Ins. Philipp.
pentasperma, Hochst. in Flora, xxvii. (1844) 552 = Oldenlandia macrophylla.
pentandra, Schum. & Thonn. Beskr. Guin. Pl. 71 = Oldenlandia macrophylla.
perpusilla, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 359 = Oldenlandia uniflora.
pilosa, Phil. ex Benth. & Hook. f. Gen. ii. 58 = Oldenlandia uniflora.
pilosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 80. t. 285. —Chili.
pinifolia, *Wall. Cat.* n. 850.—Ind. or.; Malaya.
plantaginea, Walp. Ann. i. 876 = seq.
plantaginifolia, Arn. in Nov. Act. Nat. Cur. xvii. i. (1836) 840 = verticillata.
plicata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 17 = Saurauja bracteosa !
polycarpa, R. Br. in Wall. Cat. n. 885 = scandens.
polycladia, *Muell. Fragm.* viii. 146.—Austral.
polygonoides, Wall. Cat. n. 872 = Oldenlandia diffusa !
procumbens, Wall. l. c. 861 = H. Auricularia.
prostrata, *Korth. in Nederl. Kruidk. Arch.* ii. ii. (1851) 160.—Burma; Malaya.
pruinosa, *Wight & Arn. Prod.* 408.—Ind. or.
pterita, Blume, Bijdr. 972 = Oldenlandia alata.
pterospora, *F. Muell. Fragm.* iv. 40.—Austral.
puberula, R. Br. in Wall. Cat. n. 884 = Oldenlandia umbellata.
pulvinata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 406.—Ins. Socotra.
pumila, Linn. f. Suppl. 119 = Oldenlandia crystallina.
pumila, Willd. ex Roem. & Schult. Syst. iii. 526 = Cervantesii.
pygmaea, Roem. & Schult. Syst. iii. 526 = Cervantesii.
quadrangularis, Miq. Fl. Ind. Bat. ii. 180 = tetrangularis.
quadricularis, Thw. Enum. Fl. Zeyl. 144 = Anotis quadricularis.
Quartiniana, A. Rich. Tent. Fl. Abyss. i. 362 = Oldenlandia grandiflora.
quinquenervia, *Thw. Enum. Pl. Zeyl.* 141.—Zeylan.
racemosa, Dennst. Schlüssel. Hort. Malab. 36; Wall. Cat. n. 875, partim = Anotis Rheedii.
racemosa, Lam. Encyc. iii. 80; Wight & Arn. Prod. 414 = Oldenlandia paniculata.
radicans, Baril. ex DC. Prod. iv. 424 = brachypoda.
radicans, *Miq. Fl. Ind. Bat.* ii. 161.—Ins. Philipp.
ramosa, Blume, Bijdr. 973 = Oldenlandia corymbosa.
ramosissima, Blume, l. c. 973 = Oldenlandia paniculata.
ramosissima, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 135 = Oldenlandia diffusa.
ramosissima, Spreng. Pugill. ii. 31 = Oldenlandia ramosissima.
recurva, Benth. in Hook. Lond. Journ. Bot. i. (1842) 495 = macrostemon.
HEDYOTIS :—
recurva, *Walp. Ann.* ii. 769.—Borneo.
repens, Bot. Hort. Mauric. 187 = Oldenlandia Sieberi.
repens, *Clos, in C. Gay, Fl. Chil.* iii. 208.—Chili.
repens, G. Don, Gen. Syst. iii. 526 = Oldenlandia trinervia !
repens, Lam. Tabl. Encyc. i. 371 = Dentella repens.
Rheedii, Wight & Arn. Prod. 409 = Anotis Rheedii.
rhizophylla, *Thw. ex Trim. in Journ. Bot.* xxii. (1884) 308.—Zeylan.
Richardiana, Arn, in Nov. Act. Nat. Cur. xvii. i. (1836) 840 = Anotis Richardiana.
rigida, *Miq. Fl. Ind. Bat.* ii. 181.—Java.
rigida, Walp. Ann. ii. 773 = Oldenlandia rigida.
riparia, Stend. Nom. ed. II. i. 738 = Oldenlandia capensis !
Romanoffiana, Stend. Nom. ed. II. i. 848, in syn. = Gouldia Romanoffiensis.
roses, *Rafin. Fl. Ludov.* 77.—Am. bor.
rotundifolia, Buch.-Ham. in Wall. Cat. n. 6190 = Oldenlandia nudicaulis.
rotundifolia, Spreng. Pugill. ii. 33 = Oldenlandia trinervia.
rotundifolia, Torr. & Gray, Fl. N. Am. ii. 39 = Houstonia rotundifolia.
rubra, A. Gray, Pl. Fendl. 61 = Houstonia rubra.
rugosa, *Korth. in Nederl. Kruidk. Arch.* ii. ii. (1851) 160.—Java.
rupestris, Sw. Prod. Veg. Ind. Occ. 29 = Rachicallis rupestris.
rupestris, Walp. Rep. iv. 450, Ind. sphalm. = Houstonia angustifolia.
rupicola, Sond. in Harv. & Sond. Fl. Cap. iii. 12 (Oldenlandiae sp.).—Afr. austr.
sabulosa, Stend. Nom. ed. II. i. 738 = Oldenlandia capensis !
Salsmanni, Stend. l. c. = Oldenlandia Salzmanni.
sarcophylla, Stend. l. c. = Oldenlandia crassifolia.
scaberula, *Hook. f. Fl. Brit. Ind.* iii. 50.—Burma.
scabra, *Wall. Cat.* n. 880.—Burma.
scabrida, Stend. Nom. ed. II. i. 738 = Oldenlandia corymbosa.
scandens, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 864.—Reg. Himal.; Burma.
scoparia, R. Br. in Wall. Cat. n. 861 = Oldenlandia nudicaulis.
Schimperi, Presl, Bot. Bemerk. 85 = Oldenlandia Schimperi.
Schlechtendahliana, Stend. Nom. ed. II. i. 738 = Kadua cordata.
scleranthoides, F. Muell. Fragm. iv. 89 = Oldenlandia sclerantholdes.
secunda, *Spreng. ex Schult. Mant.* iii. 133.—Porto Rico.
senegalensis, A. Rich. Tent. Fl. Abyss. i. 362 = Oldenlandia abyssinica.
senegalensis, Stend. Nom. ed. II. i. 738 = Oldenlandia senegalensis.
serpens, H. B. & K. Nov. Gen. et Sp. iii. 390 = Anotis serpens.
serpyllacea, *Schlecht. in Linnaea*, ix. (1834) 590.—Mexic.
serpyllifolia, Poir. Encyc. Suppl. iii. 14 = Oldenlandia trinervia.
serpyllifolia, Torr. & Gray, Fl. N. Am. ii. 39 = Houstonia serpyllifolia.
setifera, Stend. Nom. ed. II. i. 738 = Oldenlandia cæspitosa.
setosa, Rudd & Prv. Fl. Par. i. 57 = Mallostoma setosa.
Smithii, Walp. Rep. ii. 501 = Kadua Menziesiana.
sperguiacea, Stend. Nom. ed. II. i. 738 = Oldenlandia nudicaulis.
sperguloides, A. Rich. Tent. Fl. Abyss. i. 361 = Oldenlandia corymbosa !
spermacoces, Domb. ex DC. Prod. iv. 550 = Spermacoce corymbosa.
spermacocoides, *F. Muell. Fragm.* viii. 146.—Austral.
stellarioides, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1883) 384.—Ins. Socotra.
stenophylla, Torr. & Gray, Fl. N. Am. ii. 41 = Houstonia angustifolia.
stipitata, R. Br. in Wall. Cat. n. 6195; n. 868 A = sinuata.
stipulata, R. Br. l. c. n. 6195, partim = Anotis ingrata.
Stuckii, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iii. 57 = Oldenlandia Stuckii.
stricta, Sm. in Rees, Cycl. xvii. n. 22 = Oldenlandia Heynii.

HEDYOTIS:—
stricta, Wall. in As. Res. xiii. (1820) 369 = Oldenlandia gracilis.
strigulosa, Bartl. ex DC. Prod. iv. 437 = Oldenlandia strigulosa.
strumosa, A. Rich. Tent. Fl. Abyss. i. 364 = Oldenlandia strumosa.
stylosa, *R. Br. in Wall. Cat.* n. 858.—Ind. or.
subcarnosa, Wall. l. c. n. 860 = prostrata.
subcordata, *Dennst. Schlüssel. Hort. Malab.* 36.—Ind. or.
subtilis, Miq. Anal. Bot. Ind. iii. 12 = Anotis foetida.
subulata, Walp. Ann. ii. 772 = Oldenlandia subulata.
swertioides, *Hook. f. Fl. Brit. Ind.* iii. 51.—Ind. or.
tenella, Hochst. in Flora, xxvii. (1844) 553 (Oldenlandiae sp.).—Afr. austr.
tenella, Miq. ex Hook. f. Fl. Brit. Ind. iii. 74 = Anotis foetida.
tenelliflora, *Blume, Bijdr.* 971.—Reg. Himal.; Malaya.
tenuiflora, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 74 = Anotis foetida.
tenuifolia, Sm. in Rees, Cycl. xvii. n. 19; Spreng. Pugill. ii. 30 = Oldenlandia tenuifolia.
tenuifolia, Steud. Nom. ed. II. i. 729, sphalm. = Oldenlandia corymbosa.
tetragularis, *Wulff. Ann.* ii. 769.—Borneo.
thesiifolia, *A. St. Hil. Voy. Distr. Diam.* i. 397.—Bras.
thesioides, Willd. ex H. B. & K. Nov. Gen. et Sp. iii. 391 = Mallostoma thymifolia.
Thomasei, *Hook. f. Fl. Brit. Ind.* iii. 63.—Ind. or.
thuioides, Willd. ex Roem. & Schult. Syst. iii. 527 = Mallostoma thymifolia.
Tkwaitesiana, Hance, in Journ. Bot. vi. (1868) 298 = Anotis Thwaitesiana.
Thwaitesii, *Hook. f. Fl. Brit. Ind.* iii. 54.—Zeylan.
thymifolia, Presl, Bot. Bemerk. 85 (Oldenlandiae sp.).—Afr. austr.
thymifolia, Ruiz & Pav. Fl. Per. i. 56. t. 88 = Mallostoma thymifolia.
tilleacea, *F. Muell. Fragm.* iv. 29.—Austral.
trachymoides, *F. Muell. l. c.* 40.—Austral.
travancorica, *Bedd. Ic. Pl. Ind. Or.* i. 2.—Ind. or.
trichoglossa, *Baker, in Journ. Linn. Soc.* xxi. (1885) 409.—Madag.
trichopoda, A. Rich. Tent. Fl. Abyss. i. 360 = Oldenlandia corymbosa ?
triflora, Herb. Madr. ex Wall. Cat. p. 214. n. 867 = Oldenlandia Heynii.
trinervia, Roem. & Schult. Syst. iii. 197 = Oldenlandia trinervia.
tuberosa, Sw. Obs. Bot. 126. t. 1. f. 2 = Lucya tuberosa.
tubularis, R. Br. in Wall. Cat. n. 8576 = Anotis quadrilocularis, lancifolia.
ulmifolia, Wall. in Roxb. Fl. Ind. ed. Carey, i. 370 = linearis.
umbellata, Lam. Tabl. Encyc. i. 372 = Oldenlandia umbellata.
umbellata, Walt. Fl. Carol. 85 = Houstonia purpurea.
Uncinella, *Hook. & Arn. Bot. Beech. Voy.* 192.—Reg. Himal.; China.
uniflora, DC. Prod. iv. 421 = Oldenlandia uniflora.
uniflora, Lam. Tabl. Encyc. i. 371 = Oldenlandia glomerata.
urophylla, Wall. Cat. n. 6197 = Anotis urophylla.
Vachellii, *Benth. Fl. Hongk.* 148.—China.
vaginata, *Blume, ex DC. Prod.* iv. 421.—Ins. Molucc.
venosa, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 160 = H. Auricularia.
verrucicarpella, Steud. Nom. ed. II. i. 729 = Houstonia rotundifolia.
verticillaris, *Wall. Cat. n. 6188*; *Wight & Arn. Prod.* 409.—Ind. or.
verticillata, Lam. Tabl. Encyc. i. 371 = H. Cruciagounon.
verticillata, Walp. Ann. ii. 772 = Oldenlandia verticillata.
vestita, *R. Br. in Wall. Cat.* n. 847.—Reg. Himal.; Malaya.
villosa, Wight & Arn. Prod. 450 = Anotis Leschenaultiana.
virgata, Willd. Sp. Pl. i. 567 = Oldenlandia virgata.
virginica, Spreng. Pugill. ii. 84 = Oldenlandia glomerata.
viscida, *Bedd. Ic. Pl. Ind. Or.* i. 1.—Ind. or.
volubilis, R. Br. in Wall. Cat. n. 840 = scandens.
Walkeri, Arn. in Nov. Act. Nat. Cur. xviii. i. (1836) 330 = scandens.
Wallichii, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) pt. 2. 36.—Burma; Malaya.

HEDYOTIS:—
Wallichii, Walp. Rep. ii. 498 = hispida.
wauensis, Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr. iii. 64 = Oldenlandia wauensis.
Wightiana, Wall. Cat. n. 6194 = Anotis Wightiana.
Willdenowiana, Steud. Nom. ed. II. i. 729 = Oldenlandia Willdenowiana.
xanthochroa, *Hance, in Journ. Bot.* xxiii. (1885) 324.—China.
Zangebariae, *Roem. & Schult. Syst.* iii. 192.—Afr. trop.

HEDYPHYLLA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) 11. 143 = **Astragalus**, Tourn. (Lagumin.).
recta, Stev. l. c.—A. glycyphylloides.
vulgaris, Stev. l. c.—A. glycyphylloides.

HEDYPNOIS, [Tourn.] Scop. Fl. Carn. ed. II. ii. 99 (1772) = **Leontodon**, Linn. (Compos.).
arenaria, DC. Prod. vii. 52 = Rhagadiolus arenarius.
autumnalis, Huds. Fl. Angl. ed. II. ii. 341 = Leontodon autumnalis.
biennis, Huds. l. c. 342 = Crepis biennis.
cappica, Homem. Hort. Hafn. Suppl. 91.—Reg. Casp.
coronopifolia, Tenore, Cat. Hort. Neap. (1819) 60 = Rhagadiolus Hedypnois, *All.*
crepidiformis, *Reichb. Fl. Germ. Excurs.* 250 = Rhagadiolus Hedypnois, *All.*
cretica, Willd. Sp. Pl. iii. 1617 = Rhagadiolus Hedypnois, *All.*
furfuracea, Reichb. Fl. Germ. Excurs. 250.—Europ.
glabra, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 279 = Hypochoeris glabra.
globulifera, Lam. Fl. Fr. ii. 107 = Rhagadiolus Hedypnois, *All.*
hieracioides, Huds. Fl. Angl. ed. II. ii. 340 = Picris hieracioides.
hirta, Sm. Fl. Brit. ii. 824 = L. hirtus.
hispida, Huds. Fl. Angl. ed. II. ii. 342 = L. hispidus.
hispida, Sm. Fl. Brit. ii. 823 = L. hastilis.
lacinifiora, Viv. Fl. Lib. Spec. 52. t. 17. f. 2 = Rhagadiolus Hedypnois, *All.*
mauritanica, Schousb. ex Link, Handb. i. 810 = Rhagadiolus Hedypnois, *All.*
mauritiana, Willd. Sp. Pl. iii. 1616 = Rhagadiolus Hedypnois, *All.*
minutissima, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 372 = Rhagadiolus Hedypnois, *Fisch. & Mey.*
monspeliensis, Willd. Sp. Pl. iii. 1618 = Rhagadiolus Hedypnois, *All.*
paludosa, Scop. Fl. Carn. ed. II. ii. 100 = Taraxacum palustre.
pendula, Willd. Sp. Pl. iii. 1618; DC. Prod. vii. 82 = Rhagadiolus pendulus.
persica, Fisch. Hort. Gorenk. (1812) 34; Bieb. Fl. Taur. Cauc. iii. 539 = Rhagadiolus Hedypnois, *Fisch. & Mey.*
polymorpha, DC. Prod. vii. 81 = Rhagadiolus Hedypnois, *All.*
pygmaea, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 208.—Hispan.
pyrenaica, Vill. Prosp. 33; Hist. Pl. Dauph. iii. 78 = L. pyrenaicus.
radiata, Gaertn. Fruct. ii. 378. t. 160. f. 8 = Hyoseris radiata.
rhagadioloides, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 278 = Rhagadiolus Hedypnois, *All.*
sabulorum, Pomel, Nouv. Mat. Fl. Atl. 17.—Algeria.
scabra, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 279 = Tyroseris scabra.
Sprengelii, Schult. ex Steud. Nom. ed. II. i. 728 = Arnoseris pusilla.
Taraxaci, Huds. Fl. Angl. ed. II. ii. 341 = L. autumnalis.
Taraxacum, Scop. Fl. Carn. ed. II. ii. 99 = Taraxacum officinale.
tectorum, Huds. Fl. Angl. ed. II. ii. 341 = Crepis tectorum.
Tournefortii, Cam. in Dict. Sc. Nat. xx. 337 = Rhagadiolus Hedypnois, *All.*
tubaeformis, Tenore, Fl. Neap. ii. 179. t. 73 = Rhagadiolus Hedypnois, *All.*

HEDYSARUM, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 225 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 510.
ECHINOLOBIUM, Desv. Journ. Bot. i. (1813) 158. t. 5.
SULLA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 372.

HEDYSARUM:—
abyssinicum, Hoffmgg. Verz. Pfl. 217 = Desmodium Scalpe.
aculeolatum, *Munby, ex Boiss. Diagn. Ser. II. v. 92.*—Algeria.
acuminatum, Michx. Fl. Bor. Am. ii. 72 = Desmodium acuminatum.
adhaerens, Poir. Encyc. Suppl. v. 15 = Desmodium ormocarpoides.
adhaerens, Vahl, Symb. Bot. ii. 82 = Desmodium uncinatum.
adscendens, Sw. Prod. Veg. Ind. Occ. 106 = Desmodium adscendens.
aequidentatum, Sibth. & Sm. Fl. Graec. Prod. ii. 84 = Onobrychis cretica.
Aitkini, Eaton, Man. Bot. ed. VI. 166 = Desmodium canescens.
Aitkinianum, Steud. Nom. ed. II. i. 729, sphalm. vide *Desmodium Aitkinianum.*
Alasanni, Spreng. Syst. iv. Cur. Post. 291 = Desmodium Alasanni.
alatum, Roxb. Hort Beng. 56; Fl. Ind. iii. 348 = Desmodium triquetrum.
albens, Waldst. & Kit. Pl. Rar. Hung. ii. 115. t. 111 = Onobrychis alba.
algerience, *Pomel, Nouv. Mat. Fl. Atl.* 196.—Alger.
Alhagi, Lerch. in Nov. Act. Nat. Cur. v. (1772) App. 167; Forsk. Fl. Aegypt. Arab. 126 = Alhagi camelorum.
Alhagi, Linn. Sp. Pl. 745 = Alhagi maurorum.
Alhagi, Sibth. & Sm. Fl. Graec. viii. 15. t. 720 = Alhagi graecorum.
alopecuroides, Roxb. Hort. Beng. 57 = Uraria Lagopus.
alpinum, Linn. Sp. Pl. 750 = obscurum, sibthorpium.
alpinum, Richards. in Frankl. Narr. 1st Journ. 745 = boreale.
altaicum, Fisch. ex DC. Prod. ii. 343 = polymorphum.
altaicum, Schangin, in Fall. N. Nord. Beitr. vi. 68 = obscurum.
Ammoxylon, Spreng. Syst. iv. Cur. Post. 298 = Hremostiera Ammoxylon
ancistrocarpum, Ledeb. Ind. Hort. Derp. Suppl. (1823); ex DC. Prod. ii. 332 = Desmodium incanum.
angustifolium, H. B. & K. Nov. Gen. et Sp. vi. 517 = Desmodium angustifolium.
angustifolium, Thom. Cat. Pl. Suias. (1818) 17, nomen; et in Gaud. Fl. Helv. iv. 570 = Onobrychis arenaria.
anomalum, *DC. Prod.* ii. 341.—Hab.?
Aparines, Link, Enum. Hort. Berol. ii. 247 = Desmodium uncinatum.
arboreum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 248 = Uraria Lagopus.
arboreum, Roxb. Hort. Beng. [98]; Fl. Ind. iii. 361 = Desmodium umbellatum.
Arbuscula, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 465.—Mongolia.
arenarium, Kit. in Schult. Oestr. Fl. ed. II. ii. 368 = Onobrychis armaria.
argentatum, Linn. f. Suppl. 338 = argenteum.
argenteum, Claus, in Goebel, Reise, ii. 270 = argyrophyllum, *Ledeb.*
argenteum, Lam. ex Pall. Reise, ii. 748 = grandiflorum.
argenteum, *Murr. Syst.* ed. XIII. 562.—Reg. Caucas.
argenteum, Willd. Sp. Pl. iii. 1205 = candidum.
argyrophyllum, *Fisch. ex Steud. Nom.* ed. II. i. 729, nomen.—Sibir. alt.
argyrophyllum, *Ledeb. Fl. Ross.* i. 699.—Kossia.
armenum, Boiss. ex Tchihat. Asie Min. Bot. i. 99. t. 5 = obscurum.
articulatum, Roxb. Fl. Ind. iii. 355 = Desmodium diffusum.
asperum, Poir. Encyc. vi. 408 = Desmodium asperum.
astragaloides, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 146.—Reg. Himal.
atlanticum, *Pomel, Nouv. Mat. Fl. Atl.* 197.—Alger.
atomarium, *Boiss. Diagn. Ser.* I, ix. 111.—Syria.
atropatanum, *Bunge, ex Boiss. Fl. Orient.* ii. 623.—Persia.
Aucheri, *Boiss. Diagn. Ser.* I. ii. 87.—Cappadoc.
aureum, Herb. Madr. ex Wight & Arn. Prod. 229 = Rhynchosia densiflora.
aureum, Retzl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 218 = Rhynchosia aurea.
auriculatum, Link, Enum. Hort. Berol. ii. 249 = Desmodium triquetrum.
auriculatum, Roxb. Hort. Beng. 57; Fl. Ind. iii. 355 = Desmodium diffusum.

HEDYSARUM:—
australe, Willd. Sp. Pl. iii. 1183 = Desmodium umbellatum.
axillare, Sw. Prod. Veg. Ind. Occ. 107 = Desmodium axillare.
barbatum, Linn. Syst. ed. X. 1170 = Desmodium barbatum.
barbatum, Walt. Fl. Carol. 184 = Desmodium ciliare ?
Baxineri, *Boiss. Fl. Orient.* ii. 691.—Cappadoc.
Bergerianum, Spreng. Pugill. ii. 74 = Onobrychis Bergeriana.
biarticulatum, Linn. Sp. Pl. 747 = Desmodium biarticulatum.
biarticulatum, [Moç. & Sesse,] ex DC. Prod. ii. 880 = Desmodium infractum.
bicolor, Buch.-Ham. ex Wall. Cat. n. 5719 = Desmodium lasiflorum.
bicolorum, Poir. Encyc. vi. 448 = Adesmia bicolor.
biflorum, Willd. Sp. Pl. iii. 1900 = Desmodium biflorum.
biflorum, Willem. in Usteri, Ann. Bot. xviii. (1796) 49 = Desmodium triflorum !
bifolium, Vell. Fl. Flum. 318; vii. t. 147 = Zornia diphylla.
bijugum, Klein, ex Wight & Arn. Prod. 218 = Geissaspis cristata.
boreale, *Nutt. Gen. Am.* ii. 110.—Am. bor.
borreale, Richards. in Frankl. Narr. 1st Journ. 745 = Mackenzii.
Boveanum, *Bunge, ex Basiner, in Mém. Sav. Etr. Pétersb.* vi. (1851) 64.—Alger.
Bovei, Boiss. & Rent. Pugill. Pl. Nov. 40 = præc.
brachypterum, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1836) 92.—China.
brachypterum, *G. Don, Gen. Syst.* ii. 301.—Sibir. altaic.
bracteatum, Herb. Madr. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 159 = Alysicarpus longifolius.
bracteosum, Roxb. Hort. Beng. 56; Fl. Ind. iii. 351 = Flemingia strobilifera.
bractestum, Sieber, ex Steud. Nom. ed. II. i. 730 = Zornia diphylla.
bractreum, Michx. Fl. Bor. Am. ii. 73 = Desmodium cuspidatum.
brahuicum, *Boiss. Fl. Orient.* ii. 516.—Afghan.
Brantii, *Traust. & Mey. in Middend. Reise (Fl. Ochot.* 29).—Sibir.
brasilianum, Poir. Encyc. vi. 448 = Aeschynomene brasiliana.
brevifolium, Willd. Enum. Hort. Berol. Suppl. 51, sphalm. = leucophyllum.
Buhseanum, Boiss. in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 74 = formosum.
bupleurifolium, Herb. Wight, ex Wall. Cat. n. 5765 = Alysicarpus rugosus.
bupleurifolium, Linn. Sp. Pl. 745 = Alysicarpus bupleurifolius.
bupleurifolium, Sieber, ex DC. Prod. ii. 358 = Alysicarpus vaginalis.
Buschmianum, Georgi, Beschr. Russ. Reich. III. iv. 1161 = Onobrychis vaginalis.
Buschmianum, Pall. in Nov. Act. Petrop. x. (1797) 816, nomen = Onobrychis Pallasii.
Buschmii, Bieb. Fl. Taur. Casc. ii. 179 = Onobrychis radiata.
cachemirianum, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 146.—Reg. Himal.
caespitosum, Poir. Encyc. vi. 452 = Desmodium adscendens.
cajanifolium, H. B. & K. Nov. Gen. et Sp. vi. 525 = Desmodium cajanifolium.
callichroum, *Boiss. Diagn.* Ser. I. ii. 89.—Ax. Min.
callithrix, *Bunge, ex Boiss. Fl. Orient.* ii. 523.—Persia.
Calvertii, Boiss. Diagn. Ser. II. v. 91 = nitidum.
canadense, Linn. Sp. Pl. 748 = Desmodium canadense.
canditium, *Bieb. Fl. Taur. Casc.* ii. 176.—Tauria.
canescens, *Auct. Pl. Orient.* ii. 774.—Colaus.
canescens, Linn. Sp. Pl. 748 = Desmodium canescens.
canescens, Mill. Gard. Dict. ed. VIII. n. 7 = Desmodium incanum.
canescens, Nutt. in Torr. & Gray, Fl. N. Am. i. 357 = Mackenzii.
canescens, Willd. ex DC. Prod. ii. 330 = Desmodium retusifolium, *DC.*
canum, J. F. Gmel. Syst. 1194; Lunan, Hort. Jamaic. i. 305 = Desmodium incanum.
capense, Burm. f. Fl. Cap. Prod. 93 = Wiborgia obcordata.

HEDYSARUM:—
capense, F. G. Dietr. Volist. Lexik. Gaertn. iv. 527 = Rhynchosia pubescens.
capitatum, Burm. f. Fl. Ind. 167 = Desmodium capitatum.
capitatum, *Desf. Fl. Atlant.* ii. 177.—Reg. Mediterr.
cappadocium, *Boiss. Diagn.* Ser. I. ii. 87.—Cappadoc.
Caput-galli, Linn. Sp. Pl. 751 = Onobrychis Caput-galli.
carneum, *Ledb. Ind. Cat.* t. 312.—Reg. Caucas.
carnosum, Desf. Fl. Atlant. ii. 177 = pallidum.
carpathicum, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 779 = Onobrychis carpathica.
caucasicum, Bieb. Fl. Taur. Casc. ii. 178 = obscurum.
caudatum, Thunb. Fl. Jap. 286 = Desmodium laburnifolium.
celsiifolium, Salisb. Prod. 341 = Flemingia strobilifera.
Cephalotes, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 264.—Turkestan.
Cephalotes, Roxb. Hort. Beng. 57; Fl. Ind. iii. 360 = Desmodium Cephalotes.
cercifolium, Steud. Nom. ed. I. 392 = Uraria cercifolia.
chætocarpum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 22.—Turkestan.
ciliare, Muhl. ex Willd. Sp. Pl. iii. 1196 = Desmodium ciliare.
ciliatum, Roxb. ex Wight & Arn. Prod. 220 = Smithia sensitiva.
ciliatum, Thunb. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 48 = Rhynchosia puberula.
cinereo-sericeum, Basin. in Mém. Sav. Etr. Pétersb. vi. (1851) 66 = Basineri.
cinereum, H. B. & K. Nov. Gen. et Sp. vi. 526 = Desmodium cinereum.
circinatum, Willd. Sp. Pl. iii. 1218 = Onobrychis radiata.
circinnatum, Hoffm. Hort. Mosq. (1808) n. 1537; ex Ledeb. Fl. Ross. i. 712 = Onobrychis vaginalis.
clandestinum, Willd. ex Steud. Nom. ed. II. i. 730 = Zornia diphylla.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 197.—Alger.
coccineum, Zipp. ex Miq. Fl. Ind. Bat. i. i. 306 = Indigofera hirsuta.
cochinchinensis, *Hort. Par. ex Schrank, Pl. Rar. Hort. Monac.* t. 23.—Cochinch.
cornuto-violaceum, G. F. W. Mey. Prim. Fl. Esseq. 245 = Desmodium barbatum.
collinum, Roxb. Hort. Beng. 56; Fl. Ind. iii. 349 = Desmodium gangeticum.
colinum, Salisb. Prod. 341 = Onobrychis vicisefolia.
comosum, Vahl, Symb. Bot. ii. 86 = Uraria crinita.
concinnum, Spreng. Syst. iv. Cur. Post. 292 = Desmodium concinnum.
confertum, Bieb. Fl. Taur. Casc. ii. 180; iii. 484 = Onobrychis vicisefolia.
confertum, Desf. Fl. Atlant. ii. 178 = humile.
congestum, Kotil. ex Wall. Cat. n. 5723 = Desmodium Cephalotes.
conglomeratum, Poir. Encyc. vi. 416 = Lespedeza capitata.
conicum, Poir. l. c. 419 = Desmodium capitatum.
coningatum, Willd. Sp. Pl. iii. 1178 = Zornia diphylla.
conjunctum, Weinm. in Syll. Ratisb. ii. (1828) 175 = Desmodium incanum.
consanguineum, DC. Prod. ii. 348 = obscurum.
controversum, Cranta, Stirp. Austr. ed. II. fasc. v. 425 = obscurum.
cordatum, Jacq. Hort. Schoenb. iii. 25 = Hallia cordata.
coriaceum, Poir. Encyc. vi. 418 = Desmodium ciliare ?
cornutum, Linn. Sp. Pl. ed. II. 1060 = Onobrychis cornuta.
coronarium, *Linn. Sp. Pl.* 750.—Italia.
coronatum, Poir. Encyc. vi. 439 = Onobrychis ornata.
corsicum, Balb. Cat. Hort. Taur. (1813) 29; App. i. 12 = capitatum.
cretaceum, *Fisch. ex DC. Prod.* ii. 348.—Russia.
cretaceum, Pall. in Nov. Act. Petrop. x. (1797) 315, nomen; Georgi, Beschr. Russ. Reich. III. iv. 1161 = canditium.
creticum, Steud. Nom. ed. I. 392 = Onobrychis cretica.
crinitum, *Boiss. Diagn.* Ser. I. ii. 88.—Persia.
crinitum, Linn. Mant. i. 108; Burm. f. Fl. Ind. 169. t. 56 = Uraria crinita.

HEDYSARUM:—
crivitum, Steud. Nom. ed. I. 392 = Onobrychis crinita.
Crista-galli, Marr. Syst. ed. XIII. 568 = Onobrychis Crista-galli.
Crista-galli, [Soland. in] Russell, Aleppo, ed. II. ii. 260. t. 12 = Onobrychis Cnestaeriana.
cristatum, *Turra, Fervet. Nov. Gen.* 13.—Creta.
cryptopterum, L'Hérit. ex DC. Prod. ii. 347 = Onobrychis Michauxii.
cumanense, H. B. & K. Nov. Gen. et Sp. vi. 528 = Desmodium cuspidatum.
cuneifolium, Roth, Nov. Pl. Sp. 357 = Taverniera cuneifolia.
cuspidatum, Muhl. ex Willd. Sp. Pl. iii. 1198 = Desmodium cuspidatum.
cylindricum, Poir. Encyc. vi. 400 = Alysicarpus vaginalis.
daghestanicum, *Rupr. ex Boiss. Fl. Orient.* ii. 514.—Reg. Caucas.
dasycarpum, Turcz. in Bull. Soc. Nat. Mosc. (1840) 68 = Mackenzii.
dasyrrhizum, *Fisch. Hort. Gorenk.* ed. II. (1812) 73.—Dahuria.
decumbens, Dennst. Nom. Bot. 217 = capitatum.
deltoides, Poir. Encyc. Suppl. v. 18 = Desmodium latifolium.
deltoideum, Schum. & Thonn. Beskr. Guin. Pl. 361 = Desmodium latifolium.
dentatum, Spreng. Syst. iii. 218 = Adesmia dentata.
denticulatum, *Regel, Pl. Nov. Fedsch.* 22.—Turkestan.
depressum, Willd. Enum. Hort. Berol. Suppl. 51, nomen = capitatum.
dichotomum, Willd. Sp. Pl. iii. 1180 = Desmodium diffusum.
diffusum, Roxb. Hort. Beng. 57; Fl. Ind. iii. 357 = Desmodium lasiflorum.
diffusum, Vell. Fl. Flumn. 320; vii. t. 153 = Aeschynomene falcata.
diffusum, Willd. Sp. Pl. iii. 1180 = Desmodium diffusum.
dioicum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 244 = Desmodium dioicum.
diphyllum, Linn. Sp. Pl. 747 = Zornia diphylla.
distortum, *Aubl. Pl. Gui.* ii. 774.—Guiana.
divergens, Muhl. ex Willd. Sp. Pl. iii. 1196 = Lespedeza violacea.
divergens, Roxb. Hort. Beng. 57, nomen = Desmodium divergens.
diversifolium, Poir. Encyc. vi. 408 = Desmodium diversifolium.
dolicholoides, *Hort. Lind, ex Steud. Nom.* ed. II. i. 730, nomen (Quid ?).—Hab. ?
dametorum, *Boiss. & Heldr. Diagn.* Ser. I. ix. 111.—Ax. Min.
Ebenus, Raeusch. Nom. ed. III. 215 = Ebenus Sibthorpii.
Ecastaphyllum, Linn. Syst. ed. X. 1169 = Ecastaphyllum Browne.
echinatum, Gilib. Fl. Lituan. ii. 109 = Onobrychis vicisefolia.
echinatum, *Guss. Ind. Sem. Hort. Boccad.* (1825); *ex Linnæa*, iv. (1829) *Litt.* 37.—Ital.
echinatum, *Schweigg. Enum. Pl. Hort. Regiom.* 267.—Corsica.
echinocarpum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 410.—Sibir.
Echinus, Poir. Encyc. Suppl. v. 19 = Onobrychis Caput-galli ?
edule, Steph. ex Gleim, in Act. Hort. Petrop. iv. (1876) 54 = alpinum.
elatum, Steud. Nom. ed. II. i. 730, sphalm. = Desmodium asperum.
elegans, *Boiss. & Huet, Diagn.* Ser. II. ii. 37.—Armenia.
elegans, Lour. Fl. Cochinch. ii. 450 = Desmodium elegans.
elliptica, Zipp. ex Miq. Fl. Ind. Bat. i. i. 303 = Desmodium umbellatum.
elongatum, *Fisch. ex Ledb. Bot. Cab.* t. 1402.—Russia.
elymaticum, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* ii. 517.—Persia.
emarginatum, Poir. Encyc. vi. 412 = Desmodium emarginatum.
erectum, Petrin, ex DC. Prod. ii. 345 = Onobrychis arenaria.
erectum, Vell. Fl. Flum. 318; vii. t. 149 = Desmodium lasiocarpum ?

HEDYSARUM:—
erinaceum, Poir. Encyc. vi. 396 = Indigofera echinata.
eriophorum, Pourr. ex Desv. Journ. Bot. iii. (1814) 82 = Onobrychis eriophora.
erythrinaefolium, Juss. ex Poir. Encyc. vi. 408 = Desmodium erythrinaefolium.
erythrolecum, *Schost & Kotschy, ex Boiss. Fl. Orient.* ii. 515.—As. Min.
esculentum, *Ledeb. in Denkschr. Bot. Ges. Regensb.* iii. (1841) 113.—Sibir. or.
exaltatum, Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 103 = obscurum.
exiguum, Herb. Russ. ex Wall. Cat. n. 5497 = Rhynchosia minima.
falcatum, Poir. Encyc. vi. 448 = Aeschynomene falcata.
Falconeri, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 146.—Reg. Himal.
Fedtschenkoanum, *Regel, Pl. Nov. Fedsch.* 28.—Turkestan.
flaccidum, *Weinm. in Syll. Ratisb.* ii. (1828) 174.—Nepal.
flavescens, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 21.—Turkestan.
flavum, *Rupr. Sert. Tianach.* 44.—As. centr.
flexuosum, Balb. ex DC. Prod. ii. 341 = anomalum.
flexuosum, *Linn. Sp. Pl.* 750.—As. Min.
floribundum, D. Don, Prod. Fl. Nep. 244 = Desmodium floribundum.
Fontanesii, Boiss. Elench. 66 = humile.
formosum, *Fisch. & Mey. ex Basin. in Mém. Sav. Etr. Pétersb.* vi. (1851) 69.—Persia.
fruticescens, Linn. Sp. Pl. 748 = Lespedeza violacea.
fruticescens, Pall. *Reise*, iii. 321 = fruticosum.
fruticescens, Willd. Sp. Pl. iii. 1193 = Lespedeza capitata.
fruticosum, Habl. Taur. 121; ex Bieb. Fl. Tanr. Cauc. ii. 177 = tauricum.
fruticosum, *Pall. Reise*, iii. 753.—Sibir.
fruticosum, Vell. Fl. Flum. 319; vii. t. 154 = Aeschynomene Sellei.
fruticulosum, *Desv. in Mém. Soc. Linn. Par.* iv. (1826) 219.—Madag.
fruticulosum, Schum. & Thonn. Beskr. Guin. Pl. 368 = Desmodium mauritianum.
gangeticum, Linn. Sp. Pl. 746 = Desmodium gangeticum.
gemellum, Willd. ex Vog. in Linnaea, xii. (1838) 61 = Zornia diphylla.
geminiflorum, [Moç. & Sesse,] ex DC. Prod. ii. 330 = Desmodium Alamani.
Gibsoni, J. Grah. Cat. Pl. Bomb. 49 = Taverniera nummularia.
gigantenm, Jacque St. Hil. ex Desv. Journ. Bot. i. (1813) 123 = Desmodium asperum.
glabellum, Michx. Fl. Bor. Am. ii. 73 = Desmodium glabellum.
glabrum, Mill. Gard. Dict. ed. VIII. n. 12 = Desmodium glabrum.
glabrum, Steud. Nom. ed. I. 392 = Onobrychis viciaefolia.
glaucescens, Ledeb. Fl. Ross. i. 704 = Ibericum.
glomeratum, F. G. Dietr. Vollst. Lexik. Gaertn. iv. 534 = capitatum.
glomeratum, Roxb. Hort. Beng. 56; Fl. Ind. iii. 347 = Alysicarpus rugosus.
glumaceum, Vahl, Symb. Bot. ii. 106 = Alysicarpus glumaceus.
glutinosum, Mühl. ex Willd. Sp. Pl. iii. 1198 = Desmodium acuminatum.
glutinosum, Rottl. ex DC. Prod. ii. 337 = Rhynchosia viscosa.
Gmelini, Ledeb. in Mém. Acad. Sc. Pétersb. v. (1812) 551 = polymorphum.
graminenm, Buch.-Ham. ex Wall. Cat. n. 5762 = Alysicarpus rugosus.
gramineum, Retz. Obs. v. 26 = Alysicarpus bupleurifolius.
grandiflorum, Roxb. ex Wall. Cat. n. 5702 = Alysicarpus rugosus.
grandiflorum, Rumph. ex Wall. l. c. n. 5761 = Alysicarpus bupleurifolius.
grandiflorum, Sess. Examn. Pl. Volk. 29 = argenteum.
grandiflorum, DC. Prod. ii. 340 = setiferum.
grandiflorum, *Pall. Reise*, ii. 748.—Iber.; Caucas.; Ural.
grandiflorum, Schangin, in Pall. N. Nord. Beitr. vi. 45 = polymorphum.

HEDYSARUM:—
grandiflorum, Walt. Fl. Carol. 185 = Desmodium acuminatum.
grandifolium, Hornem. Hort. Hafn. ii. 700 = Desmodium grandifolium.
granulatum, Schum. & Thonn. Beskr. Guin. Pl. 362 = Desmodium triflorum.
granuliferum, *Spreng. Fl. Hal. Mant.* 48.—Afr. trop.
guianense, Aubl. Pl. Gui. ii. 774 = Desmodium spirale.
gyrans, Linn. f. Suppl. 332 = Desmodium gyrans.
gyroides, Roxb. Hort. Beng. 57 = Desmodium gyroides.
hamatum, Burm. f. Fl. Ind. 167 = Stylosanthes macrostata.
hamatum, Heyne, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 157 = Uraria prunellaefolia.
hamatum, Linn. Syst. ed. X. 1170 = Stylosanthes procumbens, viscosa.
Hamiltonii, Spreng. Syst. iii. 818 = Alhagi maurorum.
hamatum, Roxb. Hort. Beng. 57; et ex Wight & Arn. Prod. 222 = Uraria hamosa.
herbacenm, Lapeyr. Hist. Abr. Pl. Pyr. 426 = Onobrychis supina.
heterocarpon, Forst. f. Prod. 51 = Desmodium heterocarpum (in Addend.).
heterocarpon, Linn. Sp. Pl. 747 = Desmodium polycarpum.
heterocarpon, *Thunb. Fl. Jap.* 287 (*err. typ.* 289) = Desmodium Thunbergii.
heterophyllum, Willd. Sp. Pl. iii. 1201 = Desmodium heterophyllum, *DC.*
hirsutum, [J. F. Gmel. Syst. 1122 = canescens.
hirtum, Jacq. ex Steud. Nom. ed. I. 392 = Lespedeza capitata.
hirtum, Linn. Sp. Pl. 748 = Lespedeza hirta.
hirtum, Vell. Fl. Flum. 319; vii. t. 151 = Aeschynomene brasiliana ?
hispidulum, Spreng. Syst. iii. 818 = Adesmia hispidula.
holosericum, Boiss. & Hausskn. ex Boiss. Fl. Orient. ii. 585 = pannosum.
horridum, Steud. Nom. ed. I. 392 = Onobrychis horrida.
Huetii, *Boiss. Fl. Orient.* ii 521.—Armen.
humifusum, Pall. ex Steud. Nom. ed. I. 392 = candidum.
humifusum, Muhl. Cat. 69 = Desmodium humifusum, ochroleucum, rotundifolium, *DC.*
humile, Habl. Taur. 152; ex Bieb. Fl. Taur. Cauc. ii. 176 = candidum.
humile, *Linn. in Loefl. It.* 293.—Europ.
iberium, *Bieb. Fl. Taur. Cauc.* ii. 177.—Reg. Caucas.
ibericum, Willd. ex Cham. in Linnaea, vi. (1831) 547 = polymorphum.
ignescens, Zipp. ex Miq. Fl. Ind. Bat. i. r. 254 = Desmodium Scalpe.
imbricatum, Linn. f. Suppl. 330 = Hallia imbricata.
Immaculatum, *Forsk. Fl. Aegypt. Arab.* 214.—Arab.
inaequale, Noronha, in Verh. Batav. Gen. v. (1790), ed. I. Art. iv. 17 = Crotalaria retusa ?
incanum, Ledeb. Fl. Ross. i. 703 = varium.
incanum, Sw. Prod. Veg. Ind. Occ. 107 = Desmodium incanum.
incanum, Thunb. Fl. Jap. 289 = seq.
incanum, *Willd. Sp. Pl.* iii. 1209.—Japon.
incarnatum, Desv. in Mém. Soc. Linn. Par. iv. (1827) 311, sphalm. = prasc.
incurvatum, Hornem. Hort. Hafn. ii. 702 = Onobrychis laxiflora.
infractum, Spreng. Syst. iv. Cur. Post. 290 = Desmodium infractum.
Intactum, *Pomel, Nouv. Mat. Fl. Atl.* 198.—Alger.
intortum, Mill. Gard. Dict. ed. VIII. n. 11 = Desmodium uncinatum.
inundatum, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 761 = obscurum.
japonicum, *Basin. in Mém. Sav. Etr. Pétersb.* vi. (1851) 81.—Japon.
junceum, Linn. Sp. Pl. ed. II. 1068 = Lespedeza juncea.
juncenm, Walt. Fl. Carol. 185 = Lespedeza reticulata.
Kolenatii, C. Koch, in Linnaea, xxiv. (1851) 97 = setiferum.
Kotschyi, *Boiss. Diagn.* Ser. I. ix. 110.—Assyria.
kumaonense, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 145.—Ind. or.
laburnifolium, Poir. Encyc. vi. 422 = Desmodium laburnifolium.

HEDYSARUM:—
laeve, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 464.—Mongolia.
laevigatum, Nutt. Gen. Am. ii. 109 = Desmodium laevigatum.
lagenarium, Roxb. Hort. Beng. 57; Fl. Ind. iii. 365 = Aeschynomene aspera.
lagocephalum, Link, Enum. Hort. Berol. ii. 248 = Desmodium barbatum.
lagopodioides, Forst. f. Prod. 51 = Desmodium lagopodioides (in Addend.).
lagopodioides, Linn. Sp. Pl. 1198 = Uraria lagopoides.
lagopoides, Burm. f. Fl. Ind. 168. t. 53 = Uraria lagopoides.
lanceolatum, Schum. & Thonn. Beskr. Guin. Pl. 360 = Desmodium gangeticum.
lanceum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 16 = Uraria crinitum.
lanuginosum, *Desf. Tabl.* ed. I. 197, nomen.—Orient.
lappaceum, Forsk. Fl. Aegypt. Arab. 136 = Taverniera lappacea.
lasiocarpum, Beauv. Fl. Ow. i. 82. t. 18 = Desmodium latifolium.
lasiocarpum, Ledeb. Ind. Sem. Hort. Dorpat. (1824) Suppl. 4 = neglectum.
latebrosum, Linn. Mant. ii. 270 = Heylandia latebrosa.
latifolium, Herb. Madr. ex *Wall. Cat. n.* 5717 = Desmodium laxiflorum.
latifolium, Roxb. Hort. Beng. 57; Fl. Ind. iii. 350 = Desmodium latifolium.
latisiliquum, Poir. Encyc. vi. 432 = Chaetocalyx latisiliqua.
laxiflorum, *Benth ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 146 —Reg. Himal.
laxum, *Boiss. Diagn.* Ser. II. vi. 66.—Cappadoc.
laxum, *Pomel, Nouv. Mat. Fl. Atl.* 199.—Alger.
laxum, Spreng. Syst. iv. Cur. Post. 292 = Desmodium podocarpum.
Ledebourii, Spreng. Syst. v. 334 = neglectum.
Lehmannianum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 278.—Turkestan.
leiocarpum, Spreng. Syst. iii. 816 = Desmodium leiocarpum.
Lespedeza, Poir. Encyc. vi. 415 = Lespedeza repens.
leucocladum, *Boiss. & Heldr. Diagn.* Ser. I. ix. 112.—As. Min.
leucophyllum, *Link, Enum. Hort. Berol.* ii. 248, in obs.—Hab.?
ligneum, *Trautv. Cat. Pl. Lomon.* 12.—Mongolia.
Lindleyi, *Mart. Acava.* 25. t. 16.—Brasa.
lineare, *Lour. Fl. Cochinch.* 452.—Cochinch.
lineare, Rottl. ex Spreng. Syst. iii. 820 = Desmodium laxiflorum.
lineatum, Linn. Syst. ed. X. 1170 = Flemingia lineata.
lineatum, Michx. Fl. Bor. Am. ii. 73 = Desmodium lineatum.
linifolium, Linn. f. Suppl. 331 = Indigofera linifolia.
longifolium, Rottl. ex Spreng. Syst. iii. 819 = Alysicarpus longifolius.
Lourreiri, Spreng. Syst. iv. Cur. Post. 292 = Lourea obcordata.
lutescens, Poir. Encyc. vi. 417 = Desmodium lutescens.
lydium, Boiss. Diagn. Ser. I. ii. 33 = varium.
Mackenzii, *Richards. in Pranhl. Narr. 1st Journ.* 745.—Am. bor.
Mackeie, Buch.-Ham. ex Wall. Cat. n. 5721 = Desmodium Cephalotes.
macrophyllum, Steud. Nom. ed. I. 393 = Desmodium macrophyllum.
maculatum, Linn. Sp. Pl. 746 = Desmodium gangeticum.
madagascariense, *Desv. in Mém. Soc. Linn. Par.* iv. (1826) 220.—Madag.
malacophyllum, Link, Enum. Hort. Berol. ii. 247 = Desmodium malacophyllum.
marilandicum, Linn. Sp. Pl. 748, pro parte = Desmodium marilandicum.
marilandicum, Willd. Sp. Pl. iii. 1189 = Desmodium Dillenii.
mauritanicum, *Pomel, Nouv. Mat. Fl. Atl.* 196.—Afr. bor.
mauritanicum, Willd. Sp. Pl. iii. 1185 = Desmodium mauritanicum.
mauritianum, *Cass. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 246.—Afr. bor.
maximum, Sweet, Hort. Brit. ed. I. 180 = Desmodium uncinatum.

HEDYSARUM:—
micranthos, Poir. Encyc. vi. 446 = Aeschynomene micrantha.
microcalyx, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 147. —Reg. Himal.
microcarpum, Dufour, ex DC. Prod. ii. 345 = Onobrychis saxatilis.
microphyllum, Thunb. Fl. Jap. 284 = Desmodium parvifolium.
microphyllum, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 778.—Sibir.
microphyllum, *Bunge, ex Boiss. Fl. Orient.* ii. 538.—Persia.
molle, Vahl, Symb. Bot. ii. 83 = Desmodium molle.
molliculum, H. B. & K. Nov. Gen. et Sp. vi. 519 = Desmodium molliculum.
mongolicum, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 781, in obs.—Mongolia.
monilifernm, Linn. Mant. i. 102 = Alysicarpus monilifer.
monilifernm, Russ. ex Wall. Cat. n. 5772 = Alysicarpus vaginalis.
monuliforme, *Burm. f. Fl. Ind.* 168.—Ind. or.
montanum, Pers. Syn. ii. 324 = Onobrychis viciaefolia.
mucronatum, Blume, Cat. Gew. Buitens. 92 = Desmodium cajanifolium.
mucronatum, Sessé, ex DC. Prod. ii. 330 = Desmodium Alamani.
multijugum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 464.—Mongolia.
muricatum, Jacq. ic. Pl. Rar. iii. 13. t. 568; Coll. Suppl. 147 = Adesmia muricata.
Nalla-heschina, Roxb. ex Wight & Arn. Prod. 216 = Aeschynomene sesnoides.
Naudinianum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iii. (1856) 675.— Algeria.
neglectum, Lesch. Fl. Alt. iii. 340 = obscurum.
Nili-tuli, Roxb. Hort. Beng. 57; Fl. Ind. iii. 365 = Aeschynomene indica.
nigrum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. 1 Art. iv. 16 (Quid ?).—Java.
nitidum, *Willd. Sp. Pl.* iii. 1205.—Armenia.
nudiflorum, Linn. Sp. Pl. 749 = Desmodium nudiflorum.
nudum, Rottl. ex DC. Prod. ii. 335 = Rhynchosia minima.
nummularifolium, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 102 = Taverniera nummularia.
nummularifolium, Linn. Sp. Pl. 746 = Indigofera echinata, Alysicarpus vaginalis.
nutans, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.-June 1828) 173.—Ind. or.
obcordatum, Poir. Encyc. vi. 425 = Lourea obcordata.
oblongifolium, Bert. ex DC. Prod. ii. 332 = Desmodium axillare ?
obscurum, Georgi, Beschr. Russ. Reich. Nachtr. 295 = polymorphum.
obscurum, *Linn. Syst.* ed. X. 1171.—Europ. ; As. bor.
obtusum, Muhl. ex Willd. Sp. Pl. iii. 1190 = Desmodium ciliare.
obtusum, Pursh, Fl. Am. Sept. ii. 482 = Desmodium marilandicum.
ochroleucum, Moench, Meth. 118 = Desmodium grangeticum.
Oliverii, Spreng. Syst. iv. Cur. Post. 290 = Taverniera nummularia.
Onobrichis, Neck. Delic. Gallo-Belg. ii. 311 = Onobrychis viciaefolia.
onobrychioides, Winkl. Liter. Pfl. Verz. Fl. Balt. 104 = sec. ?
Onobrychis, Linn. Sp. Pl. 751 = Onobrychis viciaefolia.
Onobrychis, [Soland. in] Russell, Aleppo, ed. II. ii. 260. t. 11 = Onobrychis Kotschyana.
orbiculatum, Rottl. ex Wight & Arn. Prod. 281 = Desmodium Rottleri.
ornocarpoides, Desv. ex DC. Prod. ii. 327 = Desmodium ornocarpoides.
Ornocarpum, Steud. Nom. ed. II. i. 731, sphalm. = prec.
ornatum, Willd. Sp. Pl. iii. 1215 = Onobrychis ornata.
ovalifolium, Schum. & Thonn. Beskr. Guin. Pl. 359 = Alysicarpus rugosus.
ovalifolium, Vahl, ex DC. Prod. ii. 353 = Alysicarpus vaginalis.
ovatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 16 = Desmodium umbellatum.
ovalifolium, Bush.-Ham. ex Benn. Pl. Jav. Rar. 156 = Desmodium Rottleri.

HEDYSARUM:—
oxyphyllum, Spreng. Syst. iv. Cur. Post. 292 = Desmodium oxyphyllum.
palavicium, Hochst. in Lorent, Wanderungen, 347 = varium.
paleaceum, Poir. Encyc. vi. 418 = Desmodium canescens.
Pallasii, Willd. Sp. Pl. iii. 1214 = Onobrychis Pallasii.
pallidum, Bivona, Sic. Pl. Cent. ii. 29 = Sibthorpii.
pallidum, *Desf. Fl. Atlant.* ii. 177.—Afr. bor.
pallidum, Schleich. Cat. Pl. Helv. ed. III. 16 = Onobrychis arenaria.
paniculatum, Linn. Sp. Pl. 749 = Desmodium paniculatum.
paniculatum, Michx. Fl. Bor. Am. ii. 74 = Desmodium strictum.
pannosum, *Boiss. Fl. Orient.* ii. 525.—Syria.
papale, *Pomel, Nouv. Mat. Fl. Atl.* 196.—Alger.
papilliosum, *Boiss. Fl. Orient.* ii. 514.—Persia.
papposum, L'Hérit. ex DC. Prod. ii. 319 = Adesmia papposa.
parviflorum, Spreng. Syst. iii. 318 = Desmodium parvifolium.
patens, Roxb. Hort. Beng. 57; Fl. Ind. iii. 362 = Desmodium polycarpum.
pauciflorum, Nutt. Gen. Am. ii. 109 = Desmodium pauciflorum.
pedunculare, Cav. in Anal. Cienc. Nat. iv. (1801) 75 = Onobrychis pedunculariu.
pedunculatum, Mill. Gard. Dict. ed. VIII. n. 17 = Desmodium pedunculatum.
pendulum, Pou. Enc. v. 449 = Adesmia bicolor.
pennigerum, Lamb. ex DC. Prod. ii. 319 = Adesmia papposa.
Perraudianum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iii. (1856) 739.—Alger.
Pestalozzae, *Boiss. Diagn.* Ser. I. ix. 112.—Lycia.
petraeum, Bieb. ex Willd. Sp. Pl. iii. 1217 = Onobrychis petraea.
phrygium, Boiss. & Bal. Diagn. Ser. II. vi. 66 = varium.
pictum, Jacq. Coll. ii. 262 = Uraria picta.
pilosum, Roxb. ex Wight & Arn. Prod. 228 = Desmodium retroflexum.
pilosum, Thunb. Fl. Jap. 288 (err. typ. 290) = Desmodium pilosum.
pinnatifolium, Poir. Encyc. vi. 447 = Adesmia muricata.
platyphyllum, *Basin. in Mém. Sav. Etr. Pétersb.* vi. (1851) 7.—Rossia.
plumosum, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 515.—Persia.
podocarpon, Spreng. Syst. iii. 317 = Desmodium podocarpum.
pogonocarpum, *Boiss. Diagn.* Ser. I. ii. 87.—As. Min. ; Syria.
polycarpum, Poir. Encyc. vi. 413 = Desmodium polycarpum.
polymorphum, *Lesleb. Fl. Alt.* iii. 338.—Sibir.
portericenses, Spreng. Syst. iii. 314 = Desmodium portoricense.
procumbens, *Mill. Gard. Dict.* ed. VIII. n. 10.—Jamaic.
procumbens, Roxb. Hort. Beng. 56; Fl. Ind. iii. 345 = Desmodium retroflexum.
procumbens, Schangin, in Pall. N. Nord. Beitr. vi. 33 = polymorphum.
procumbens, Vell. Fl. Flum. 319; vii. t. 150 = Desmodium barbatum.
prostratum, Linn. Mant. i. 102; Burm. f. Fl. Ind. 168. t. 46 = Indigofera endesphylla.
prostratum, Muhl. ex Willd. Sp. Pl. iii. 1200 = Lespedeza repens.
prostratum, Pall. ex DC. Prod. ii. 342 = polymorphum.
prostratum, Roxb. ex Wight & Arn. Prod. 209 = Pseudarthria viscida.
Pryeni, Spreng. Syst. iv. Cur. Post. 292 = Desmodium Wightii.
Pseud-Alhagi, Bieb. Fl. Taur. Cauc. ii. 174 = Alhagi camelorum.
ptolemaicum, Delile, Fl. Egypte, 285 = Onobrychis ptolemaica.
pulchellum, Linn. Sp. Pl. 747 = Desmodium pulchellum.
pumilum, Linn. Syst. ed. X. 1171 = Onobrychis pumila.
punctatum, Poir. Encyc. vi. 447 = Adesmia punctata.
punctatum, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 223 = Rhynchosia densiflora.
purpureum, *Gmelin-Lat. Reise*, i. 186 (Quid ?).—Russia.
purpureum, Mill. Gard. Dict. ed. VIII. n. 4 = Desmodium incanum.

HEDYSARUM:—
purpureum, Roxb. Hort. Beng. 57; Fl. Ind. iii. 358 = Desmodium polycarpum.
quinquangulatum, Roxb. Hort. Beng. [98]; Fl. Ind. iii. 355 = Desmodium diffusum.
racemiferum, J. F. Gmel. Syst. 1125 = Desmodium canescens.
racemosum, Anbl. Pl. Gui. ii. 774 = Desmodium adscendens.
racemosum, Thunb. Fl. Jap. 285 = Desmodium racemosum.
radiatum, Bieb. Fl. Taur Cauc. iii. 488 = Onobrychis vaginalis.
radiatum, Desf. in Ann. Mus. Par. xii. (1808) 112 = Onobrychis radiata.
ramosum, Willd. ex Steud. Nom. ed. II. i. 731 = Desmodium arenarium.
Rasoumowinnum, *Helm. & Fisch. ex DC. Prod.* ii. 342. —Reg. Casp.
recurvatum, Roxb. Hort. Beng. 57; Fl. Ind. iii. 358 = Desmodium laxiflorum.
reflexum, *Hoffmgg. Vers. Pfl. Nachtr.* ii. 198.—Hab. ?
reflexum, Reinw. ex Blume, Cat. Gew. Buitens. 93 = Desmodium umbellatum.
remifolium, Linn. Syst. ed. X. 1169 = sec.
reniforme, Linn. Sp. Pl. ed. II. 1051 = Desmodium reniforme.
reniforme, Lour. Fl. Cochinch. 447 = Lourea obcordata.
repandum, Vahl, Symb. Bot. ii. 82 = Desmodium Scalpe.
repens, Linn. Sp. Pl. 749 = Lespedeza repens.
reptans, Poir. Encyc. vi. 432 = Desmodium axillare.
reptans, Roxb. Hort. Beng. 57; Fl. Ind. iii. 354 = Desmodium heterophyllum, *DC.*
reticulatum, Mohl. ex Willd. Sp. Pl. iii. 1194 = Lespedeza reticulata.
retroflexum, Linn. Mant. i. 108 = Desmodium retroflexum.
retroflexum, Linn. Mant. iii. 516, sphalm. = *lineatum*.
retusum, D. Don, Prod. Fl. Nep. 243 = Desmodium polycarpum.
rhombifolium, Ell. Sketch, ii. 216 = Desmodium laevigatum.
rhombifolium, Roxb. Hort. Beng. 57, nomen = Desmodium rhomboideum.
riparium, Ell. Sketch. ii. 215 = Desmodium rigidum.
Roezlianum, *Prantl, Ind. Sem. Hort. Wratisl.* (1878) 8.—Calif.
roseum, Blume, ex Miq. Fl. Ind. Bat. i. 1. 248 = Desmodium gangeticum.
roseum, Sims, Bot. Mag. t. 996 = lauricum.
roseum, Steph. ex DC. Prod. ii. 342 = polymorphum.
rosmarinifolium, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 17 = Alysicarpus vaginalis.
Rottleri, Spreng. Syst. iii. 320 = Desmodium laxiflorum.
rotundifolium, *Boiss. & Noë, Diagn.* Ser. II. ii. 87.—Armenia.
rotundifolium, Michx. Fl. Bor. Am. ii. 72 = Desmodium rotundifolium, *DC.*
rotundifolium, Vahl, Symb. Bot. ii. 81 = Indigofera echinata.
Roxburghii, Spreng. Syst. iv. Cur. Post. 292 = Desmodium laxiflorum.
rubrum, Spreng. Syst. iii. 317 = Desmodium rubrum.
rufescens, Spreng. l. c. iv. Cur. Post. 292 = Desmodium rufescens.
rufescens, Willd. ex Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 222 = Rhynchosia rufescens.
rugosum, Lieber, ex DC. Prod. ii. 353 = Alysicarpus buplenrifolius.
rugosum, Willd. Sp. Pl. iii. 1172 = Alysicarpus rugosus.
ruthlecarpum, DC. Prod. ii. 341 = polymorphum.
sagittatum, Poir. Encyc. vi. 406 = Desmodium sagittatum.
salicifolium, Poir. l. c. 422 = Desmodium salicifolium.
salicna, Salisb. Prod. 341 = Desmodium gyrans.
sambucense, D. Don, Prod. Fl. Nep. 243 = Desmodium floribundum ?
sanctae, Linn. Syst. ed. X. 1171 = Onobrychis sanctae.
sanctae, Pall. Reise, ii. 727 (Quid ?).—Sibir. sud.
sandwicense, Ell. Sketch. ii. 217 = Desmodium canescens.

HEDYSARUM:—
scabrum, Moench, Meth. 118 = Desmodium canadense.
Scalpe, Comm. ex DC. Prod. ii. 334 = Desmodium Scalpe.
scandens, *Mill. Gard. Dict.* ed. VIII. n. 18.—Mexic.
scariosum, Kotzl. ex Spreng. Syst. iii. 319 = Alysicarpus rugosus.
scoparium, Fisch. & Mey. in Schrenk, Enum. Pl. Nov. 87 = Corethrodendron scoparium.
Scorpiurus, Sw. Prod. Veg. Ind. Occ. 107 = Desmodium Scorpiurus.
Semenowii, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) 11. 38.—*An.* centr.
sennoides, Willd. Sp. Pl. iii. 1207 = Ormocarpum sennoides.
sericeum, *Bieb. Fl. Taur. Cauc.* ii. 176.—Reg. Cauca.
sericeum, *Mill. Gard. Dict.* ed. VIII. n. 8.—Mexic.
sericeum, Thunb. Fl. Jap. 287 (err. typ. 289) = Lespedeza juncea.
sericeum, Vahl, Symb. Bot. ii. 83 = Ebenus pinnata.
serotinum, Willd. Enum. Hort. Berol. 777 = Desmodium serotinum.
sessilifolium, Poir. Encyc. vi. 414 = Lespedeza reticulata.
sessilifolium, Torr. ex M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 122 = Desmodium sessilifolium, tenuifolium.
setigerum, *Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 29.—Dahuria.
Severzowii, *Bunge, ex Regel, in Act. Hort. Petrop.* iii. (1875) 116.—Turkestan.
sibiricum, Ledeb. Fl. Ross. i. 707 = elongatum.
sibiricum, *Poir. Encyc. Suppl.* v. 17.—Sibir.
Sibthorpii, *Nym. Consp.* 197.—Reg. Mediterr.
sikkimense, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 145.—Reg. Himal.
siliquosum, Burm. f. Fl. Ind. 169 = Desmodium polycarpum.
singareense, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 893.—Mesopotam.
sinjolense, Bolss. Diagn. Ser. I. ii. 89 = callichroum.
sosgaricum, *Bong. Vers. Saisang-Nor. Pfl.* 30. t. 8.—Sibir.
sororium, Linn. Mant. ii. 270 = Eleiotis sororia.
Spartium, Burm. f. Fl. Ind. 166 = Taverniera nummularia.
spectabile, Boj. ex Sweet, Hort. Brit. ed. I. 479 = Desmodium polycarpum.
spinosissimum, DC. Fl. Fr. vi. 588 = capitatum.
spinosissimum, *Linn. Sp. Pl.* 750.—Reg. Mediterr.
spinosissimum, Sibth. & Sm. Fl. Graec. viii. 16 = Sibthorpii.
spirale, Sw. Prod. Veg. Ind. Occ. 107 = Desmodium spirale.
splendens, *Fisch. ex DC. Prod.* ii. 340.—Sibir.
squarrosum, Thunb. Prod. Pl. Cap. 132 = Eriosema squarrosum.
sternorhizum, Dufour, ex DC. Prod. ii. 346 = Onobrychis sternorhiza.
stipulaceum, Burm. f. Fl. Ind. 168 = Desmodium triflorum.
stipulaceum, Sessé, ex DC. Prod. ii. 330 = Desmodium tortuosum.
stolonifervum, Rich. ex Poir. Encyc. vi. 421 = Desmodium axillare.
striatum, Thunb. Fl. Jap. 289 = Desmodium striatum.
strictum, Pursh, Fl. Am. Sept. ii. 468 = Desmodium strictum.
strobilifervm, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 146. —Reg. Himal.
strobiliferum, Linn. Sp. Pl. 746 = Flemingia strobilifera.
styracifolium, Linn. Syst. ed. X. 1169 = Desmodium rutrobaxium.
styracifolium, Roxb. ex Wight & Arn. Prod. 235 = Desmodium gangeticum.
styracifolium, Roxb. Hort. Beng. 36; Fl. Ind. iii. 347 = Alysicarpus rugosus.
subjectaceum, Pisch. ex DC. Prod. ii. 343 = Eversmannia hedysaroides.
Suwonium, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 248 = Desmodium floribundum.
subjectorum, Boiss. Diagn. Ser. II. v. 91 = varium.
sudanum, H. B. & K. Nov. Gen. et Sp. vi. 534 = Desmodium canadense.
sujetum, Pall. in Nov. Act. Petrop. x. (1797) 316, nomen = vendidum.

HEDYSARUM:—
supinum, Sw. Prod. Veg. Ind. Occ. 106 = Desmodium incanum.
supinum, Vill. Prosp. 41; Hist. Pl. Dauph. iii. 394 = Onobrychis supina.
syriacum, Boiss. Diagn. Ser. I. ii. 90 = stenarium.
tanakense, *Fisch. Hort. Gorenk.* ed. II. (1812) 78.—Rossia.
tauricum, *Pall. in Nov. Act. Petrop.* x. (1797) 315, nomen; *Bieb. Cent. Pl. i.* t. 35.—Tauria.
tenellum, D. Don, Prod. Fl. Nep. 245 = Desmodium parvifolium.
tenellum, H. B. & K. Nov. Gen. et Sp. vi. 522 = Desmodium spirale.
terminale, Rich. in Act. Soc. Hist. Nat Par. i. (1792) 112 (err. typ. 105) = Desmodium molle.
tetraphyllum, Thunb. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 44 = Zornia tetraphylla.
tiliaefolium, D. Don, Prod. Fl. Nep. 244 = Desmodium tiliaefolium.
timoriense, Spreng. Syst. iv. Cur. Post. 290 = Desmodium timoriense.
Thunbergianum, Steud. Nom. ed. II. i. 732 = reflexum.
tomentosum, Thunb. Fl. Jap. 286 = Lespedeza villosa.
tortuosum, Sw. Prod. Veg. Ind. Occ. 107 = Desmodium tortuosum.
Tournefortii, Willd. Sp. Pl. iii. 1214 = Onobrychis Tournefortii.
triangulare, *Rets. Obs.* iii. 40.—Java.
trichocarpum, *Duyv. ex DC. Prod.* ii. 341.—Persia.
trichocarpum, Willd. Sp. Pl. iii. 1194 = Lespedeza trichocarpa.
triflorum, Linn. Sp. Pl. 749 = Desmodium triflorum.
trifoliatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv 16 (Quid?).—Java.
trigonum, Sw. Prod. Veg. Ind. Occ. 107 = Desmodium uncinatum.
trinerve, Buch.-Ham. in Wall. Cat. n. 5720 = Desmodium podocarpum.
trinervium, Roxb. ex Wall. l. c. n. 5747 = Flemingia congesta.
trinervium, *Willd. Enum. Hort. Berol. Suppl.* 51, nomen.—Hab.?
triphyllum, DC. Prod. ii. 332, in syn. = Desmodium uncinatum.
triquetrum, Linn. Sp. Pl. 746 = Desmodium triquetrum.
tuberculosum, Labill. Sert. Austr. Caled. 71. t. 72 = Desmodium polycarpum?
tuberosum, Roxb. ex Willd. Sp. Pl. iii. 1197 = Pueraria tuberosa.
turkestanicum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 21.—Turkestan.
umbellatum, Linn. Sp. Pl. 747 = Desmodium umbellatum.
umbellatum, Roxb. Hort. Beng. [98]; Fl. Ind. iii. 360 = Desmodium Cephalotes.
umbellatum, Walt. Fl. Carol. 184 = Lespedeza capitata?
uncinatum, Jacq. Hort. Schoenb. iii. 27 = Desmodium uncinatum.
uniflorum, Domb. ex DC. Prod. ii. 319 = Adesmia hispidula.
uniforum, *Lapeyr. Hist. Abr. Pl. Pyr.* 426.—Pyren.
vaginale, Linn. Sp. Pl. 745 = Alysicarpus vaginalis.
varians, Labill. Sert. Austr. Caled. 71. t. 73 = Desmodium varians.
variegatum, Portenschl. Enum. Pl. Dalm. 15; Host, Fl. Austr. ii. 320 = capitatum.
varium, Roth, Nov. Pl. Sp. 354 = Alysicarpus vaginalis.
varium, *Willd. Sp. Pl.* iii. 1206.—*An.* Min.
velutinum, Boiss. & Noé, Diagn. Ser. II. ii. 88 = polycarpum.
velutinum, Willd. Sp. Pl. iii. 1174 = Desmodium velutinum.
venosum, Desf. Fl. Atlant. ii. 179. t. 201 = Onobrychis venosa.
venosum, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 220 = Rhynchosia reticulata.
venustulum, H. B. & K. Nov. Gen. et Sp. vi. 519 = Desmodium barbatum.
venustum, Fisch. ex DC. Prod. ii. 342 = polymorphum.
vespertilionis, Salisb. Prod. 341 = Lourea vespertilionis.
vespertilionum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 16 (Quid?).—Java.
vespertilionis, Linn. f. Suppl. 331 = Lourea vespertilionis.

HEDYSARUM:—
viciaefolium, Turcz. in Bull. Soc. Nat. Mosc. (1840) 69 = sibiricum.
villosum, Mill. Gard. Dict. ed. VIII. n. 9 = Desmodium villosum. *DC.*
villosum, Willd Sp. Pl. iii. 1195 = Lespedeza villosa
violaceum, Forsk. Fl. Aegypt. Arab. 136 = Alysicarpus glumaceus.
violaceum, Linn. Sp. Pl. 749 = Lespedeza violacea, reticulata.
violaceum, Vell. Fl. Flum. 318; vii. t. 148 = Desmodium axillare.
virgatum, Buch.-Ham. ex Wall. Cat. n. 5761 = Alysicarpus bupleurifolius.
virgatum, Thunb. Fl. Jap. 288 (err. typ. 290) = Lespedeza virgata, *DC.*
virgulosum, Hill, Veg. Syst. xxi. 43 = scq.
virgulosum, Linn. Sp. Pl. 750 = Aeschynomene hispida.
viridiflorum, Burm. f. Fl. Ind. 167 = Desmodium Wightii.
viridiflorum, Linn. Sp. Pl. 748 = Desmodium triviridiflorum.
viridiflorum, Willd. Sp. Pl. iii. 1192 = Desmodium canescens.
viscidum, Linn. Sp. Pl. 747 = Pseudarthria viscida.
viscosum, DC. Prod. ii. 336, in syn. sphalm. = Pseudarthria viscida.
volgense, Fisch. ex DC. Prod. ii. 343 = cretaceum.
volitans, Roxb. ex Wight & Arn. Prod. 221 = Lourea vespertilionis.
volubile, Linn. Sp. Pl. 750 = Galactia mollis.
Wade, Vand. Fasc. Pl. 17 = Desmodium Wade.
Wallichianum, Spreng. Syst. iv. Cur. Post. 293 = Desmodium floribundum.
Wightii, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 146. —Reg. Himal.
HEDYSCEPE, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 178. *PALMAE*, Benth. & Hook. f. iii. 888.
Canterburyana, *H. Wendl. & Drude, l. c.* 204.—Ins. Dom. Howe.
HEDYSTACHYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophul.).
spicata, *Fourr.* l. c. = V. spicata.
HEERIA, Meisn. Gen. Comm. 56 (1837) = **Anaphrenium**, E. Mey. (Anacard.).
argentea, Meisn. l. c. = A. argenteum.
inermis, Meisn. l. c. = Sideroxylon inerme.
longifolia, Bernh. ex Krauss, in Flora, xxvii. (1844) 549 = Protorhus longifolia.
melanophlea, Meisn. l. c. = Myrsine melanophloea.
mucronata, Bernh. ex Krauss, in Flora, xxvii. (1844) 549 = A. abyssinicum.
mucronifolia, Engl. in A. DC. Monog. Phan. iv. 388, sphalm. = prace.
HEERIA, Schlecht. in Linnaea, xiii. (1839) 482. *MELASTOMACEAE*, Benth. & Hook. f. i. 740.
HETEROCENTRON, Hemsl. & Arn. Bot. Beech. Voy. 290 (1840).
SCHIZOCENTRON, Meisn. Gen. Comm. 355 (1842).
anisiflora, Hemsl. Biol. Centr. Am. Bot. i. 416, in syn. = macrostachya.
cuphoides, Benth. Bot. Voy. Sulph. 93. t. 33 = Pterogastra cuphoides.
elegans, *Schlecht. in Linnaea*, xiii. (1839) 482.—Mexic.
macrostachya, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 24.—Mexic.
procumbens, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 151 = elegans.
rosea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 24 —Mexic.
subtriplinervis, *Triana, l. c.*—Mexic.
undulata, *Triana, l. c.*—Mexic.
HEGEMONE, Bunge, ex Ledeb. Fl. Ross. i. 51 (1842) = **Trollius**, Linn. (Ranunc.).
lilacina, Bunge, l. c. = T. lilacinus.
HEGETSCHWEILERA, Heer & Regel, Cat. Sem. Hort. Turic. (1842); ex Bot. Zeit. i. (1843) 47 = Alysicarpus, Neck. (Legumin.).
paichelia, Heer & Regel, ll. cc. = A. vaginalis.

HEIMIA, Link & Otto, Ic. Pl. 63. t. 28 (1822) = **Nesaea**, Comm. (Lythrar.).
grandiflora, Hook. Ic. Fl. t. 554 = N. salicifolia.
linariaefolia, Hort. ex G. Don, in Loud. Hort. Brit. 479 = N. linariaefolia.
myrtifolia, Hort. Berol. ex Cham. & Schlecht. in Linnaea, ii. (1827) 347 = N. myrtifolia.
salicifolia, Link & Otto, Ic. Pl. 63. t. 28 = N. salicifolia.
syphilitica, DC. Prod. iii. 89 = N. syphilitica.

HEINEKENIA, Webb, ex Benth. & Hook. f. Gen. i. 491 (1868) = **Lotus**, Linn. (Legumin.).
peteriyacha, Webb, ex Masf. in Anal. Soc. Esp. Hist. Nat. x. (1881) 160 = L. Berthelotii.

HEINSIA, DC. Prod. iv. 890 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 77.
benguelensis. *Welw. ex Benth. & Hook. f. Gen.* ii. 77. —Afr. trop.
capensis, Bunk, ex Harv. Thes. Cap. i. 22 = Randia radis.
densiflora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 81.—Afr. trop.
jasminiflora, *DC. Prod.* iv. 890.—Afr. trop.
pubescens, *Klotzsch, in Abh. Acad. Berl.* (1857) 228. —Afr. trop.

HEINTZIA, Karst. Ausw. Gew. Venez. 34. t. 11 (1848); et in Otto & Dietr. Allg. Gartenz. xvii. (1849) 33 = **Alloplectus**, Mart. (Gesnerac.).
picta, Karst. ex Planch. in Fl. des Serres, vii. (1851–52) 201. tt. 718, 719.—Venezuela.
tigrina, Karst. Ausw. Gew. Venez. 34. t. 11 = A. tigrinus.

HEINTZIA, Steud. Nom. ed. II. i. 789 (1840) = HEINZIA, Scop. = **Dipteryx**, Schreb. (Legumin.).

HEINZELIA, Nees, in Mart. Fl. Bras. ix. 153 (1847) = **Chaetothylax**, Nees (Acanthac.).
longiflora, Nees, in DC. Prod. xi. 314 = Jacobinia longiflora.
lythroides, Nees, in Mart. Fl. Bras. ix. 154 = C. lythroides.
ovalis, Nees, l. c.—Bras.

HEINZELMANNIA, Neck. Elem. i. 371 (1790) = **Spigelia**, Linn. (Loganiac.).

HEINZIA, Scop. Introd. 301 (1777) = **Dipteryx**, Schreb. (1791) (Legumin.).
peregrina, J. F. Gmel. Syst. 509 = D. odorata ?

HEISTERA, Schreb. Gen. i. 291 (1789) = seq.

HEISTERIA, Jacq. Enum. Pl. Carib. 4 (1760); Am. Select. 126 (1763). *OLACINEAE*, Benth. & Hook. f. i. 846.
ACROLOBUS, Klotzsch, in Abh. Akad. Berl. (1856) 296. t. 3.
HESIODA, Vell. Fl. Flam. 185 (1825); iv. t. 140 (1827).
RHAPTOSTYLUM, Humb. & Bonpl. Pl. Aequin. ii. 139. t. 125 (1809).
acuta, *Engl. in Mart. Fl. Bras.* xii. II. 21.—Am. austr.
brasiliensis, *Engl. l. c.* 19.—Bras.
caniflora, *Sw. in Nees, Cycl.* xvii. n. 2.—Guiana.
celastrina, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 381.—N. Granat.
citriifolia, *Engl. in Mart. Fl. Bras.* xii. II. 18.—Am. austr.
coccinea, Engl. l. c. 19 = brasiliensis.
coccinea, *Vacq. Enum. Pl. Carib.* 20; *Select. Am.* 126. —Ins. Martinic.
coccinea, Poepp. ex Engl. in Mart. Fl. Bras. xii. II. 18 = cyanocarpa.
crassifolia, Mart. ex Engl. l. c. 16 = ovata.
cyanocarpa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 85. t. 240.—Bras.
densiflora, *Engl. in Mart. Fl. Bras.* xii. II. 17.—Am. austr.
flexuosa, *Engl. l. c.*—Am. austr.
guianensis, *Engl. l. c.* 15.—Am. austr.
Kappleri, *Sagot, ex Engl. l. c.* 14.—Guian.
laxiflora, *Engl. l. c.* 18.—Am. austr.
longifolia, *Spruce, ex Engl. l. c.* 14.—Bras.

HEISTERIA:—
macrophylla, *Oerst. in Kjoeb. Vidensah. Meddel.* (1856) 40.—Am. centr.
maynadiensis, *Spruce, ex Engl. in Mart. Fl. Bras.* xii. II. 15.—Bras.
microcarpa, *Spruce, ex Engl. l. c.* 21.—Bras.
nitida, *Engl. l. c.* 18.—Am. austr.
ovata, *Benth. in Hook. Kew Journ.* iii. (1851) 866.—Bras.
pallida, *Engl. in Mart. Fl. Bras.* xii. II. 16.—Am. austr.
parvifolia, *Sm. in Rees, Cycl.* xvii. n. 3.—Afr. trop.
pentandra, *Engl. in Mart. Fl. Bras.* xii. II. 20.—Am. austr.
Raddiana, *Benth. in Hook. Niger Fl.* 258.—Bras.
Rhaptostylum, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 381.—N. Granat.
salicifolia, *Engl. in Mart. Fl. Bras.* xii. II. 20.—Am. austr.
Spruceana, *Engl. l. c.* 15.—Am. austr.
subsessilis, *Benth. in Hook. Kew Journ.* iii. (1851) 867.—Bras.
tubicina, Poepp. & Endl. Nov. Gen. et Sp. iii. 85. t. 241 = Aptandra tubicina.

HEISTERIA, Linn. Gen. ed. I. 350 (1737) = **Muraltia**, Neck. (Polygal.).
mitior, Berg. Descr. Pl. Cap. 187 = M. stipulacea.
pungens, Berg. l. c. 185 = M. Heisteria.

HEKATEROSACHNE, Steud. Syn. Pl. Gram. 118 (1855) = **Oplismenus**, Beauv. (Gramin.).
elatior, Steud. l. c. = O. setarius.

HEKERIA, Endl. Gen. Suppl. i. 1874, in syn. (1841) = HECKERIA, Kunth = **Piper**, Linn.

HEKISTOCARPA, Hook. f. Ic. Pl. t. 1151 (1878). *RUBIACEAE*, Benth. & Hook. f. ii. 62.
minutiflora, *Hook. f. l. c.*—Afr. trop.

HEKORIMA, Kunth, Enum. Pl. iv. 208 (1843) = **Streptopus**, Michx. (Liliac.).
canadica, Kunth, l. c. 304 = S. simplex.
dichotoma, Kunth, l. c. = S. roseus.

HELADENA, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 321. *MALPIGHIACEAE*, Benth. & Hook. f. i. 358.
HELADRAIA, Lindl. Veg. Kingd. 890 (1847).
albiflora, *A. Juss. in Arch. Mus. Par.* iii. (1843) 349. —Bras.
australis, *A. Juss. l. c.* 348.—Am. austr.
biglandulosa, *A. Juss. l. c.* 349.—Bras.
bunchosioides, *A. Juss. l. c.* 348.—Bras.

HELADENIA, Reichb. Nom. 207 (1841) = prec.

HELADRAIA, Lindl. Veg. Kingd. 890 (1847) = prec.

HELANTHIUM, Engelm. ex Benth. & Hook. f. Gen. iii. 1005 (1883) = **Alnus**, Dill.

HELCIA, Lindl. Bot. Reg. (1845) Misc. 17 = **Trichoglottis**, Lindl. (Orchid.).
picta, Linden, in Illustr. Hortic. xxix. (1882) 52.—N. Granat.
sanguinolenta, Lindl. Bot. Reg. (1845) Misc. 17 = T. sanguinolenta.

HELDREICHIA, Boiss. in Ann. Sc. Nat. Sér. II. xvi. (1841) 381. *CRUCIFERAE*, Benth. & Hook. f. i. 91.
ZYGOPELTIS, Fenzl, ex Endl. Gen. Suppl. ii. 107 (1845).
Bornmuelleri, *Boiss. Fl. Orient.* i. 220.—As. Min.
bupleurifolia, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 381; xvii. (1842) 186.—As. Min.; Syria.
Kotschyi, *Boiss. l. c.*—Cilicia.
longifolia, *Boiss. l. c.* 381; 187.—Persia.
rotundifolia, *Boiss. l. c.* 381; 185.—Armenia.
silaifolia, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 178.—Afghan.

HELEASTRUM, DC. Prod. v. 269 (1836) = **Aster**, Tourn. (Compos.).
albens, DC. l. c. 264 = A. pterocaulos.
paludosum, DC. l. c. = A. paludosus.

HELEIOTIS, Hassk. Cat. Hort. Bog. Alt. 272, in syn. (1844) = **Phylacium**, Benn. (Legumin.).
scandens, Hassk. l. c. = P. scandens.

HELEMONIUM, Steud. Nom. ed. II. i. 788 (1840) = HELENOMOIUM, Willd. = **Heliopsis**, Pers. (Compos.).
canescens, Steud. l. c. = Heliopsis buphthalmoides.

HELENA, Haw. Monog. Narcis. 18 (1831) = **Narcissus**, Tourn. (Amaryll.).
crocco-cincta, Haw. l. c. 14 = N. poëticus.
gracilis, Haw. l. c. 13 = N. gracilis.
pianicorona, Haw. l. c. = N. gracilis.
pumila, Haw. l. c. 14 = N. gracilis ?
purpureo-cincta, Haw. l. c. 13 = N. poëticus.
tenuior, Haw. l. c. = N. gracilis.

HELENEUM, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 460 = **Helenium**, Linn. (Compos.).

HELENIA, Linn. Gen. ed. I. 260 (1737) = **Helenium**, Linn. (Compos.).
autumnalis, Gaertn. Fruct. ii. 438. t. 169 = Helenium autumnale.
decurrens, Moench, Meth. 589 = Helenium autumnale.
Douglasii, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 390 = Monolopia major.
linearifolia, Moench, Meth. Suppl. 240 = Helenium quadridentatum.

HELENIOPSIS, Baker, in Journ. Bot. xii. (1874) 278, sphalm. = **Heloniopsis**, A. Gray (Liliac.).

HELENIUM, Linn. Sp. Pl. ed. I. 886 (1753). *COMPOSITAE*, Benth. & Hook. f. ii. 418.
AMBLYOLEPIS, DC. Prod. v. 667 (1836).
BRASSAVOLA, Adans. Fam. ii. 127 (1763).
DUGALDIA, Cass. in Dict. Sc. Nat. lv. 270 (1828).
EPELETIOPSIS, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 414 (1873).
HELENIA, Linn. Gen. ed. I. 260 (1737).
LEPTOCARPHA, Endl. Gen. Suppl. i. 1383, in syn. (1841).
LEPTOPHORA, Rafin. in Am. Monthly Mag. (1819) 195.
LEPTOPODA, Nutt. Gen. Am. ii. 174 (1818).
MERODETRA, Rafin. Fl. Ludov. 14 (1817).
OXYLEPIS, Benth. Pl. Hartw. 87 (1841).
TETRODUS, Cass. in Dict. Sc. Nat. lv. 264, 272 (1828).
aestivale, *Walt. Fl. Carol.* 210.—Am. bor.
alatum, C. C. Gmel. ex Steud. Nom. ed. I. 394 = quadridentatum.
altissimum, Link, Ind. Sem. Hort. Berol. (1840) = autumnale.
amphibolum, *A. Gray, in Proc. Am. Acad.* ix. (1874) 202.—Mexic.
anceps, *Rafin. New Fl. Am.* iv. 81 (Quid ?).—Am. bor.
atropurpureum, Kunth & Bonpl. Ind. Sem. Hort. Berol. (1845) 12 = nudiflorum.
auctumnale, J. F. Gmel. Syst. 1244 = seq.
autumnale, *Linn. Sp. Pl.* 886.—Am. bor.
Bigelovii, *A. Gray, in Pacif. Rail. Rep.* iv. 107.—Calif.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 358.—Calif.
brachypodum, Wood, Am. Bot. & Flor. 182 = nudiflorum.
brevifolium, *Wood, l. c.*; *A. Gray, in Proc. Am. Acad.* ix. (1874) 204.—Am. bor.
canaliculatum, Steud. Nom. ed. II. i. 788 = Eriophyllum caespitosum.
californicum, Link, Ind. Sem. Hort. Berol. (1840) 21 = puberum.
canaliculatum, Lam. in Journ. Hist. Nat. ii. (1792) 218. t. 86 = autumnale.
commutatum, Link, Ind. Sem. Hort. Berol. (1846) 21 = autumnale.
Curtisii, *A. Gray, in Proc. Am. Acad.* ix. (1874) 204. —Am. bor.
decurrens, Vatke, Ind. Sem. Hort. Berol. (1872) App. = puberulum.
dichotomum, *Rafin. New Fl. Am.* iv. 81 (Quid ?).—Am. bor.
discoideum, *Rafin. l. c.* (Quid ?).—Am. bor.

HELENIUM:—
Douglasii, Hort. ex W. Bart. in Loud. Hort. Brit. Suppl. iii. 590 = Monolopia major.
elegans, *DC. Prod.* v. 667.—Am. bor. occ.; Mexic.
fimbriatum, *Wood, Am. Bot. & Flor.* 182; *A. Gray, in Proc. Am. Acad.* ix. (1874) 204.—Am. bor.
flexuosum, *Rafin. New Fl. Am.* iv. 81 (Quid!).—Am. bor.
grandiflorum, Gillh. Fl. Lituan. i. 204 = Inula Helenium.
grandiflorum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 384 = autumnale.
heterophyllum, DC. Prod. v. 667 = microcephalum.
Hoopesii, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 65.—Am. bor. occ.
incisum, Wood, Am. Bot. & Flor. 182 = Nuttallii.
integrifolium, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 237.—Mexic.
laciniatum, *A. Gray, in Proc. Am. Acad.* ix. (1874) 208.—Am. bor. occ.
lanatum, Spreng. Syst. iii. 574 = Eriophyllum caespitosum.
latifolium, Mill. Gard. Dict. ed. VIII. n. 2 = autumnale.
Leptopoda, Wood, Am. Bot. & Flor. 182 = Nuttallii?
longifolium, Sm. in Rees, Cycl. xvii. n. 2 = autumnale.
mexicanum, DC. Prod. v. 666 = amphibolum.
mexicanum, A. Gray, in Pacif. Rail. Rep. iv. 107 = puberulum.
mexicanum, *H. B. & K. Nov. Gen. et Sp.* iv. 299.—Mexic.
micranthum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 385 = nudiflorum.
microcephalum, M. A. Curt. ex A. Gray, Syn. Fl. N. Am. i. ii. 346 = elegans.
microcephalum, *DC. Prod.* v. 667.—Mexic.
Milleri, Schult. ex DC. Prod. v. 666 = quadridentatum.
montanum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 384 = autumnale.
nudiflorum, *Nutt. l. c.*—Am. bor.
Nuttallii, *A. Gray, in Proc. Am. Acad.* ix. (1874) 204.—Am. bor.
ooclinium, *A. Gray, l. c.* 202.—Mexic.
oppositifolium, Spreng. Syst. iii. 573 = Bahia oppositifolia.
parviflorum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 384 = nudiflorum.
puberulum, *DC. Prod.* v. 667.—Calif.
puberulum, Wood, Am. Bot. & Flor. 182 = vernale?
pubescens, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 227 = autumnale.
pubescens, Hook. & Arn. Bot. Beech. Voy. 355 = puberulum.
pumilum, Willd. Enum. Hort. Berol. Suppl. 60 = autumnale.
quadridentatum, Hook. Comp. Bot. Mag. i. (1835) 98 = nudiflorum.
quadridentatum, *Labill. in Act. Soc. Hist. Nat. Par.* i. (1792) 22. t. 4.—Am. bor.
quadripartitum, Link, Enum. Hort. Berol. ii. 888 = quadridentatum.
Roulia, *Perez. in Bull. Soc. Nat. Moac.* xxiv. (1851) i. 186.—Calif.
scorzonerasfolia, A. Gray, in Proc. Am. Acad. vii. (1866) 359 = Hecubaea scorzonerasfolia.
semiamplexum, Featherm. in Louisiana Univ. Rep. (1871) = nudiflorum.
serotinum, *Walt. Fl. Carol.* 210.—Am. bor.
stenchaetifolium, Spreng. Syst. iii. 574 = Eriophyllum stenchaetifolium.
tenuifolium, *Nutt. in Journ. Acad. Philad.* vii. (1834) 66.—Am. bor.
texanum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 460 = microcephalum.
Thurberi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 82.—Arizona.
trexilium, *Rafin. New Fl. Am.* iv. 89 (Quid!).—Am. bor.
trolliifolium, Spreng. Syst. iii. 573 = Bahia trolliifolia.
tuberiformum, DC. Prod. v. 666 = autumnale.
undulatum, *Mœnch, ex Steud. Nom.* ed. II. i. 733.—Calif.
vernum, Schrad. Ind. Sem. Hort. Gotting. (1831) ex DC. Prod. v. 666 = mexicanum.
vernale, *Walt. Fl. Carol.* 210.—Am. bor.

HELENIUM, Linn. Syst. ed. I (1735); Adans. Fam. ii. 138 (1763) = Inula, Linn. (Compos.).

HELENOMOIUM, Willd. ex DC. Prod. v. 551, in syn. (1836) = **Heliopsis**, Pers. (Compos.).
canescens, Willd. ex DC. l. c. = Heliopsis buphthalmoides.

HELEOCHARIS, Lestib. Ess. Cyperac. 41 (1819) = **Blasocharis**, R. Br. (Cyperac.).

HELEOCHLOA, Beauv. Agrost. 28 (1812) = **Sporobolus**, R. Br. (Gramin.).
juncea, Beauv. l. c. 24 = S. junceus.
phalaroides, Beauv. l. c.—Hab.?

HELEOCHLOA, Host, Gram. Austr. i. 23. t. 29, 30 (1801). *GRAMINEAE*, Benth. & Hook. f. iii. 1146.
PACHERA, Pourr. Chlor. Narb. ex Kunth, Enum. Pl. i. 23, in syn. (1888).
RHIZOCEPHALUS, Boiss. Diagn. Ser. I. v. 68 (1844); xiii. 43 (1853).
acutiglumis, *Boiss. Fl. Orient.* v. 676.—Syria.
alopecuroides, *Host, Gram. Austr.* i. 23. t. 29.—Reg. Mediterr.
alpicola, *Boiss. Fl. Orient.* v. 677.—Kurdist.
ambigua, *Benth. & Hook. f. Gen.* iii. 1146.—Lydia.
cruciancilloides, *Benth. & Hook. f. l. c.*—As. Min.
dianthra, Host, Fl. Austr. i. 77 = Crypsis aculeata.
dura, *Boiss. Fl. Orient.* v. 677.—Beluchist.
pygmaea, *Benth. & Hook. f. Gen.* iii. 1146.—As. Min.; Persia.
schoenoides, *Host, Gram. Austr.* i. 23. t. 30.—Reg. Mediterr.; Persia.

HELEOGITON, Schult. Mant. ii. 2 (1824) = **Scirpus**, Linn. (Cyperac.).
compressa, Peterm. Fl. Bienitz, 22 = S. Caritis.
fluitans, Link, ex Reichb. Ic. Fl. Germ. viii. t. 298 = S. lacustris.
glaucum, Reichb. in Moesal. Handb. ed. II. iii. 1807 = S. lacustris.
Leyanni, Reichb. Fl. Germ. Excurs. 77 = S. triqueter.
litorale, Reichb. in Moesal. Handb. ed. II. iii. 1808 = S. littoralis.
Rothii, Reichb. l. c. = S. pungens.
tabernaemontani, Peterm. Fl. Bienitz, 22 = S. Tabernaemontani.
triqueum, Reichb. in Moesal. Handb. ed. II. iii. 1807 = S. carinatus.
triquetrum, Reichb. l. c. 1808 = S. triqueter.

HELEONASTES, Ehrh. Beitr. iv. 147 (1789) = **Carex**, Linn. (Cyperac.).

HELEOPHILA, Schult. Mant. ii. 2 (1824) = seq.

HELEOPHYLA, Flora, iv. (1821) 25 = seq.

HELEOPHYLAX, Beauv. ex Lestib. Ess. Fam. Cyp. 41 (1819) = **Scirpus**, Linn. (Cyperac.).

HELEPTA, Rafin. Neogenyt. 3 (1825) = **Heliopsis**, Pers. (Compos.).
angustifolia, Rafin. l. c. = Heliopsis laevis?
grandiflora, Rafin. l. c. = Heliopsis laevis.
parviflora, Rafin. l. c. = Heliopsis laevis?

HELIA, Benth. & Hook. f. Gen. i. 905 (1869) = HELIX, M. Roem. = **Atalantia**, Corres (Rutac.).

HELIA, Mart. Nov. Gen. et Sp. ii. 123. t. 191 (1827) = **Linnanthus**, Aubl. (Gentian.).
brevifolia, Cham. in Linnaea, viii. (1833) 11 = L. brevifolius.
oblongifolia, Mart. Nov. Gen. et Sp. ii. 123 = L. oblongifolius.
spathulata, Mart. l. c. 124 = L. Martii.

HELIAMPHORA, Benth. in Proc Linn. Soc. i. (1840) 53; et Trans. Linn. Soc. xvii. (1840) 483. t. 29. *SARRACENIACEAE*, Benth. & Hook. f. i. 48.
nutans, *Benth. ll. cc.*—Guiana.

HELIANTHELLA, Torr. & Gray, Fl. N. Am. ii. 338 (1841). *COMPOSITAE*, Benth. & Hook. f. ii. 379.
saxifylla, *A. Gray, in Proc. Am. Acad.* xix. (1884) 9.—Am. bor. occ.

HELIANTHELLA:—
californica, *A. Gray, in Pacif. Rail. Rep.* iv. 108.—Calif.
Douglasii, *Torr. & Gray, Fl. N. Am.* ii. 334.—Am. bor. occ.
grandiflora, *Torr. & Gray, l. c.* 338.—Am. bor.
lanceolata, Torr. & Gray, l. c. 334 = uniflora.
latifolia, Scheele, in Linnaea, xxii. (1849) 160 = Viguiera helianthoides.
mexicana, *A. Gray, in Proc. Am. Acad.* xv. (1880) 37.—Mexic.
microcephala, *A. Gray, l. c.* xix. (1884) 10.—Am. bor. occ.
multicaulis, Eaton, in Bot. King's Exped. 170 = uniflora.
nevadensis, Greene, in Bull. Calif. Acad. i. n. iii. (1885) 89 = californica.
nudicaulis, *A. Gray, in Proc. Am. Acad.* xix. (1884) 9.—Am. bor. occ.
Parryi, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 65.—Am. bor. occ.
quinquenervis, *A. Gray, in Proc. Am. Acad.* xix. (1884) 10.—Am. bor. occ.
tenuifolia, *Torr. & Gray, Fl. N. Am.* ii. 333.—Am. bor.
uniflora, A. Gray, in Proc. Acad. Sc. Philad. 1863 (1864) 65 = quinquenervis.
uniflora, *Torr. & Gray, Fl. N. Am.* ii. 334.—Am. bor. occ.

HELIANTHEMOIDES, Medic. Phil. Bot. i. 95 (1789) = **Talinum**, Adans. (Portulac.).
patens, Medic. l. c. = T. patens.

HELIANTHEMON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = **Helianthemum**, Tourn. (Cistin.).
fumanum, St. Lag. l. c. = Helianthemum Fumana.
libanotum, St. Lag. l. c. = Helianthemum Libanotis.
polyanthum, St. Lag. l. c. = Helianthemum polyanthos.
tuberarium, St. Lag. l. c. = Helianthemum Tuberaria.

HELIANTHEMUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 445 (1821), sphalm. = **Helianthus**, Linn. (Compos.).

HELIANTHEMUM, Tourn. ex Hall. Enum. Stirp. Helv. i. 358 (1742). *CISTINEAE*, Benth. & Hook. f. i. 113.
ANTHELIS, Rafin. Chlor. Acta. 9 (1813).
CROCANTHEMUM, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 370.
FUMANA, Spach, l. c. 359 (1836).
FUMANOPSIS, Pomel, Mat. Fl. Atl. 9 (1860).
HALIMIUM, Dun. ex Spach, in Ann. Sc. Nat. Sér. II. vi. 365 (1836).
HETEROMERIS, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 290.
PRISTINA, Rafin. Sylva Tellur. 133 (1838).
PRISTUS, Neck. Elem. ii. 278 (1790).
RHODAX, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 363.
STEGITRIS, Rafin. Sylva Tellur. 132 (1838).
TAENIOSTEMA, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 289.
TRICHASTEROPHYLLUM, Willd. ex Link, in Jahrb. Gew. i. iii. (1820) 69.
TUBERARIA, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 364.
XOLANTHA, Rafin. Caratteri, 73 (1810).
XOLANTHES, Rafin. Sylva Tellur. 132 (1838).
acuminatum, Pers. Syn. ii. 79 = varietate.
acutifolium, Ehrenb. ex Willk. Ic. Pl. Hisp. ii. 98 = kahiricum.
aegyptiacum, Privald. ex Willk. l. c. = lodiolium.
aegyptiacum, *Mill. Gard. Dict.* ed. VIII. n. 23.—Reg. Mediterr.; Persia.
aelandicum, Boiss. Diagn. Ser. II. vi. 93, sphalm. = oelandicum.
albicans, Ehrenb. ex Willk. Pl. Hisp. ii. 101 = Lippii.
algarvense, Dun. in DC. Prod. i. 268 = scynoides.
Allionii, Tineo, Pl. Rar. Sicil. 48 = vineale.
alpestre, Dun. in DC. Prod. i. 276 = italicum.
alpestre, Reichb. Pl. Crit. i. 8 = vineale.
alpoides, Pers. ex Nym. Consp. 72 = guttatum.
alpoides, *Vent. Choix*, 30. t. 30.—Hispan.
ambiguum, *Pomel, Nouv. Mat. Fl. Atl.* 210.—Alger.
ambiguum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 66 [bis] = rupifragum.
Anderssonii ×, Swess, Cat. t. 69.
angustatum, *Pomel, Nouv. Mat. Fl. Atl.* 218.—Alger.

HELIANTHEMUM:—

angustifolium, Pers. Syn. ii. 79 = variabile.
annuum, Fisch. ex Steud. Nom. ed. II. i. 733 = villosum.
antkyrum, Stend. l. c. = umbellatum.
apenninum, Boiss. Fl. Orient. i. 445 = pulverulentum.
apenninum, DC. Fl. Fr. iv. 894 = polifolium.
apenninum, Mill. Gard. Dict. ed. VIII. n. 4 = variabile.
apertum, *Pomel, Nouv. Mat. Fl. Atl.* 219.—Alger.
arabicum, *Pers. Syn.* ii. 80.—Arabia.
arcuatum, Presl, Fl. Sicul. i. 128 = glaucum.
arenicola, *Chapm. Fl. S. U. St.* 36.—Am. bor.
argenteum. *Hemsl. Diagn. Pl. Nov.* 90.—Mexic.
artense, *Motelet. in Mém. Acad. Lyon*, x. (1860) 178.—N. Caled.
asperum, *Lag. ex Dun. in DC. Prod.* i. 283.—Hispan.
asperum, Duf. ex Willk. lc. Pl. Hisp. ii. 92 = intermedium.
astylum, *Moq. & Sesse, ex Dun. in DC. Prod.* i. 284.—Mexic.
atriplicifolium, *Willd. Enum. Hort. Berol.* 569.—Hispan.
aureum, Thil. ex Pers. Syn. ii. 79 = hirtum.
baeticum, Hort. Matr. ex Dun. in DC. Prod. i. 281 = hirtum.
barbatum, Pers. Syn. ii 79 = variabile.
barbatum, Sweet, Cist. i. 73 = vulgare.
Barrelieri, Tenore, Prod. Fl. Neap. 31 = thymifolium.
Bertterianum, Bertol. Fl. Ital. v. 887 = vulgare.
bicolor, Presl, Fl. Sicul. i. 128 = glaucum.
biseriale, *Pomel, Nouv. Mat. Fl. Atl.* 218.—Alger.
Bourgeanum, Nym. Syll. 229 = Cistus Bourgeanus.
brasilianse, *Pers. Syn.* ii. 77.—Bras.
breviper, Boiss. & Reut. Pugill. Nov. Pl. 18 = guttatum.
Brewerl, Planch. in Hook. Lond. Journ. Bot. iii. (1844) 618 = guttatum.
Broussonetii, *Dun. in DC. Prod.* i. 279.—Ins. Teneriff.
bupleurifolium, *Dun. l. c.* 270.—Hispan.; Lusitan.
cabiricum, Stend. Nom. ed. II. i. 753 = kahiricum.
calcareum, Jord. Gr. Jard. Grenoble (1849), 16 = polifolium.
calycinum, *Dun. in Mém. Acad. Montp. Sect. Sc.* (1847) 4.—Reg. Mediterr. occ.
calycinum, Willd. Enum. Hort. Berol. 570 = H. Fumana.
canadense, *Michx. Fl. Am.* i. 308.—Am. bor.
canariense, *Pers. Syn.* ii. 76.—Afr. bor. occ.; Ins. Canar.
candidum, Sweet, Cist. t. 25 = ocymoides.
canescens, Moench, Meth. Suppl. 75 = canariense.
canescens, Sweet, Cist. t. 51 = variabile.
canum, *Boiss. Fl. Orient.* i. 444.—Europ.; As. Min.
canum, Dun. in DC. Prod. i. 277 = viscale.
capitatum, *Nutt. ex Engelm. & Gray, in Bost. Journ. Nat. Hist* v. (1847) 212.—Am. bor. occ.
Caput-felis, *Boiss. Elench.* 16.—Hispan.
carolinianum, *Michx. Fl. Bor. Am.* i. 307.—Am. bor.
Chamaecistus, Mill. Gard. Dict. ed. VIII. n. 1 = vulgare.
cheiranthoides, Pers. Syn. ii. 76 = alyssoides.
ciliatum, Moris, Stirp. Sard. Elench. i. 6 = Morisianum.
ciliatum, *Pers. Syn.* ii. 76.—Hispan.
ciliatum, Pourr. ex Nym. l. c. = occidentale.
cinereum, *Pers. Syn.* ii. 76.—Hispan.
cistifolium, Mill. Gard. Dict. ed. VIII. n. 9 = variabile.
clandestinum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 797 = aegyptiacum.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 221.—Alger.
confertum, Dun. in DC. Prod. i. 276 = canariense.
confertum, Willk. lc. Pl. Hisp. ii. 102. t. 181 A = Lappii.
confusum, Sweet, Cist. t. 91 = pulverulentum, *Willk.*
contortum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 794 = villosum.
controversum, F. Schultz, Fl. Palat. 60 = pulverulentum, *Willk.*
Cordi, *Rupr. Fl. Cauc.* 141.—Reg. Caucas.
coriaceum, Pers. Syn. ii. 76 = Hibbertia coriacea.
corymbosum, *Michx. Fl. Bor. Am.* i. 307.—Am. bor.
corymbosum, Pursh, Fl. Am. Sept. 353 = canadense.
Coulteri, *S. Watts. in Proc. Am. Acad.* xvii. (1881–82) 353.—Am. bor. occ.
crassifolium, Dun. ex Nym. Consp. 75 = H. Sexte.
crenatifolium, *Pers. Syn.* ii. 77.—Hispan.
crispatum, Boiss. & Reut. Diagn. Ser. II. i. 51 = halimifolium.
crispulum, Nym. Consp. 73, spbalm. = praec.
croceum, Pers. Syn. ii. 79 = glaucum.
crocrum, Tenore, Syll. App. 602 = vulgare.

HELIANTHEMUM:—

cupreum, Sweet, Cist. t. 66 = variabile.
dagestanicum, *Rupr. Fl. Cauc.* 143.—Reg. Caucas.
declinatum, *Moench, Meth. Suppl.* 75.—Hab.?
denticulatum, Thib. ex Pers. Syn. ii. 76 = salicifolium.
desertorum, *Willk. lc. Pl. Hisp.* ii. 127.—Hispan.
dichotomum, *Pers. Syn.* ii. 77.—Hispan.
dichroum, Kunze, in Flora, xxix. (1846) 633 = vinesle.
discolor, *Pomel, Nouv. Mat. Fl. Atl.* 217.—Alger.
distachyon, *Moench, Meth. Suppl.* 76.—Lusitan.
dinerifolium, Sweet, Cist. t. 95 = variabile.
dubium, *Pourr. ex Nym. Consp.* 76.—Hispan.
echioides, *Pers. Syn.* ii. 77.—Hispan.
Ehrenbergii, Willk. lc. Pl. Hisp. ii. 99. t. 128 = Lippii.
ellipticum, Pers. Syn. ii. 78 = Lippii.
elongatum, Pers. l. c. 76 = alyssoides.
eremophilum, *Pomel, Nouv. Mat. Fl. Atl.* 850.—Alger.
ericoides, Dun. in DC. Prod. i. 274 = H. Fumana.
ericoaulon, Dun. l. c. 271 = guttatum.
ericoephalum, *Pomel, Nouv. Mat. Fl. Atl.* 851.—Alger.
ericosepalum, *Sweet, Cist.* t. 76.—Europ. austr.
fasciculatum, *Mill. Gard. Dict.* ed. VIII. n. 22.—Afr. austr.
floribundum, *Pomel, Nouv. Mat. Fl. Atl.* 223.—Alger.
foetidum, Pers. Syn. ii. 79 = variabile.
Fontanesii, *Boiss. & Reut. Pugill. Pl. Nov.* 15.—Alger.
formosum, *Dun. in DC. Prod.* i. 268.—Lusitan.
fugacrum, Mill. Gard. Dict. ed. VIII. n. 19 = salicifolium.
fugax, Dun. in DC. Prod. i. 284 = salicifolium.
Fumana, *Mill. Gard. Dict.* ed. VIII. n. 6.—Reg. Mediterr.
fumanioides, Desf. ex Dun. in DC. Prod. i. 275 = praec.
fumanum, Moench, Meth. 333 = H. Fumana.
germanicum, Mill. Gard. Dict. ed. VIII. n. 2 = variabile.
getulum, *Pomel, Nouv. Mat. Fl. Atl.* 858.—Alger.
glabellum, Schur, Enum. Pl. Transs. 76 = italicum.
glabriflorum, *Kit. in Linnaea*, xxxii. (1863) 506.—Hungar.
glandulosum, *Presl, in Oken, Isis*, xxi. (1828) 275.—Ins. Creta.
glaucum, Pers. Syn. ii. 78.—Reg. Mediterr. occ.
glaucum, Sweet, Cist. t. 111 = canariense.
globulariaefolium, *Pers. Syn.* ii. 77.—Lusitan.
glomeratum, *Lag. ex Dun. in DC. Prod.* i. 276.—Mexic.
glutinosum, *Pers. Syn.* ii. 79.—Reg. Mediterr.
gorgoneum, *Webb. in Hook. Niger Fl.* 109.—Ins. Cap. Virid.
graecum, Boiss. & Heldr. Diagn. Ser. II. i. 52 = vulgare.
grandiflorum, DC. Fl. Fr. iv. 891 = vulgare.
Guirol, *Willk. in Linnaea*, xxxi. (1859–60) 86.—Hispan.
guttatum, *Mill. Gard. Dict.* ed. VIII. n. 18.—Reg. Mediterr.
halimifolium, *Pers. Syn.* ii. 75.—Reg. Mediterr.
halimioides, *Pomel, Nouv. Mat. Fl. Atl.* 221.—Alger.
Herrii × Bruegg, in Jahresb. Naturf. Ges. Graub. II. xxiii.-xxiv. (1880) 73.—Europ.
heterodoxum, Dun. in DC. Prod. i. 270 = guttatum.
hirtophyllum, Schur, Enum. Pl. Transs. 77 = vulgare.
hirtophyllum, Stend. Nom. ed. II. i. 734 = ocymoides.
hirsutissimum, *Presl, Rel. Haenk.* ii. 95.—Chili.
hirsutissimum, Willk. lc. Pl. Hisp. t. 106 = halimifolium.
hirsutum, Dun. in DC. Prod. i. 284 = variabile.
hirtum, *Mill. Gard. Dict.* ed. VIII. n. 14; *Pers. Syn.* ii. 79.—Hispan.
hispidum, Dun. in DC. Prod. i. 283 = hirtum.
hispidum, Tenore, Syll. 360 = polifolium.
hymettium, Boiss. & Heldr. Diagn. Ser. II. i. 52 = rubellum.
hyssopifolium, Tenore, Cat. Hort. Neap. (1819) 48 = polifolium.
ibericum, Desf. ex Stend. Nom. ed. II. i. 734 = arabicum.
imbricatum, Lag. ex Dun. in DC. Prod. i. 270 = guttatum.
incanum, Moench, Meth. Suppl. 76 = variabile.
incanopicnum, Thib. ex Pers. Syn. ii. 77 = guttatum.
inflatum, Moench, Meth. 333 = aegyptiacum.
intermedium, *Thib. ex Pers. Syn.* ii. 78.—Hispan.; Lusitan.
involucratum, Pers. l. c. 76 = alyssoides.

HELIANTHEMUM:—

italicum, Gren. & Godr. Fl. Fr. i. 171 = montanum.
italicum, *Pers. Syn.* ii. 76.—Europ. austr.
italicum, Tenore, Fl. Nap. iv. 214 = canum.
Jacquini, Willk. in Ardoino, Fl. Alp. Marit. 49 = Morisianum?
juniperinum, Lag. ex Dun. in DC. Prod. i. 275 = thymifolium.
juniperinum, Schloss. ex Nym. Consp. 76 = procumbens.
kahiricum, Delile, Fl. Egypte, 237.—Afr. bor.; Arab. Kotschyanum, *Boiss. Diagn.* Ser. II. i. 53.—Syria, Mesopot.
Kraeheri, Stend. Nom. ed. I. 394 = variabile.
laeve, Pers. Syn. ii. 78 = glutinosum.
laeve, Reichb. ex Nym. Consp. 76 = thymifolium.
laevigatum, Schur, Enum. Pl. Transs. 77 = vulgare.
laevipes, *Pers. Syn.* ii. 76.—Reg. Mediterr. occ.
laevipes, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 275 = glandulosum.
Lagnascae, Dun. in DC. Prod. i. 281 = hirtum.
lancolatum, Steud. Nom. ed. I. 394 = Hibbertia coriacea.
lanceolatum, Sweet, Cist. t. 100 = variabile.
Langei, Willk. lc. Pl. Hisp. ii. 73 = globulariaefolium.
lanuginosum, *Spreng. Syst.* ii. 589.—Afr. bor.
lasianthum, Pers. Syn. ii. 76 = alyssoides.
lasiocalycinum, Boiss. & Reut. Diagn. Ser. II. i. 50 = halimifolium.
lasiocarpum, Desf. ex Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 261 = ledifolium.
lavandulaefolium, *Mill. Gard. Dict.* ed. VIII. n. 18; *DC. Fl. Fr.* iv. 890.—Reg. Mediterr.
lavandulaefolium, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = Lippii.
lavandulaefolium, Tenore, Fl. Nap. i. 396 = polifolium.
ledifolium, *Mill. Gard. Dict.* ed. VIII. n. 20; *Moench, Meth.* 333.—Europ. austr.
ledifolium, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = retrofractum.
ledifolium, Spach, Hist. Vég. Phan. vi. 17 = villosum.
Ledon, Zeyh. ex Steud. Nom. ed. II. i. 734 = Cistus glaucus.
leptophyllum, *Dun. in DC. Prod.* i. 279.—Hispan.
leucanthum, *Ehrenb. ex Willk. lc. Pl. Hisp.* ii. 160.—Hispan.
levipes, Moench, Meth. 332, vide laevipes.
Libanotis, Dod. ex Willk. lc. Pl. Hisp. ii. 48. = Bourgeanus.
Libanotis, Gaza. Cat. Hort. Boccad. 30 = Cistus Clusii.
Lindisfa, *Willd. Enum. Hort. Berol.* 570.—Lusit.; Afr. bor.
libycum, *Pomel, Nouv. Mat. Fl. Atl.* 852.—Alger.
lignosum, Sweet, Cist. t. 46 = H. Tuberaria.
linarce, Pers. Syn. ii. 78 = variabile.
linariifolium, Moench, Meth. Suppl. 75 = variabile.
Lippii, *Pers. Syn.* ii. 78.—Reg. Mediterr.
lucidum, Herbert, Hort. Hafn. ii. 498 = variabile.
lunulatum, *DC. Fl. Fr.* iv. 816.—Alp. Pedem.
lusitanicum, Mill. Gard. Dict. ed. VIII. n. 16 = variabile.
lutcum, Hall. Fl. Wirceb. i. 350 = variabile.
macranthum, Schur, Enum. Pl. Transs. 76 = vulgare.
macranthum, Sweet, Cist. t. 108 = variabile.
macrocarpum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1877) 545 = guttatum.
majoranaefolium, DC. Fl. Fr. vi. 625 = hirtum.
marifolium, Hens. Enum. Pl. Velh. 23 = oelandicum.
marifolium, Mill. Gard. Dict. ed. VIII. n. 24; DC. Fl. Fr. iv. 817 = viscale.
maritimum, *Pomel, Nouv. Mat. Fl. Atl.* 220.—Alger.
medium, *Schur, in Verh. Siebenb. Ver. Naturw.* iii. (1852) 68.—Transsyl.
mendozanum, *Steud. Plant. Preiss.* ed. II. i. 795.—Mexic.
micranthum, Duf. ex Nym. Consp. 75 = intermedium.
microanthum, *Spreng. Syst.* iii. 886.—Afr. bor.
microcarpum, Cosson & Balan. Fl. Orient. i. 441 = ledifolium.
microphyllum, Sweet, Cist. t. 96 = ocymoides.
Millori, Sweet, l. c. t. 102 = variabile.
minutum, Schlecht. ex Willk. lc. Pl. Hisp. ii. 91 = salicifolium.
mollis, *Pers. Syn.* ii. 76.—Hispan.
montanum, *Vis. Pl. Dalmat.* iii. 146.—Dalmatia.
Morisianum, *Rouleï. Fl. Bad.* v. 874.—Ital.

HELIANTHEMUM :—

mucronatum, Dun. in DC. Prod. i. 274 = canariense.
multiflorum, Salzm. ex Dun. in Mém. Acad. Montp. (Sect.-Sc.) i. (1847) 3 = halimifolium.
multiflorum, Ziz, in Opiz, Natural. xi. (1836) 486 = vulgare!
multiple, Steud. Nom. ed. II. i. 735 = variabile.
mutabile, Moench, Meth. 234 = variabile.
mutabile, Willk. Ic. Pl. Hisp. ii. 115 = vulgare.
nebrodense, Heldr. in Guss. Fl. Sic. Syn. ii. 18 = vineale.
niloticum, Moench, Meth. 283 = ledifolium.
nitidum, *Clem. in Atti Tera. Riun. Sc. Ital.* 517.—Dalmat.
nudicaule, Dun. in DC. Prod. i. 279 = glaucum.
nummulariaefolium, Sprun. ex Willk. Ic. Pl. Hisp. ii. 141 = rubellum.
nummularium, Guss. Cat. Hort. Boccad. 30 = rubellum.
nummularium, Mill. Gard. Dict. ed. VIII. n. 12 = vulgare.
obcordatum, *Moq. & Sesse, ex Dun. in DC. Prod.* i. 284.—Mexic.
obovatum, Dun. l. c. 277 = montanum.
obovatum, Willk. Ic. Pl. Hisp. ii. 149 = Rossmaessleri.
obscurum, Pers. Syn. ii. 79 = vulgare.
obtusatum, *Pomel, Nouv. Mat. Fl. Atl.* 220.—Alger.
obtusifolium, *Dun. in DC. Prod.* i. 281.—Ins. Cyprus. occidentale, *Nym. Consp.* 72.—Hispan.
ocimifolium, Pourr. ex Nym. Consp. 72 = sco.
ocymoides, *Pers. Syn.* ii. 76.—Hispan. ; Lusitan.
oelandicum, DC. Fl. Fr. iv. 817 = italicum.
oelandicum, Fries, Novit. Fl. Suec. ed. II. 168 = canum.
oelandicum, Pers. Syn. ii. 77 = cinereum !
oelandicum, Sw. in Sv. Bot. vii. t. 499 = vineale.
oeiandicum, *Wahlenb. Fl. Suec.* 333.—Europ.
oleaefolium, Hort. ex Willk. Ic. Pl. Hisp. ii. 66 = lepidotum.
oligophyllum, *Dun. in DC. Prod.* i. 284.—Palaest.
origanifolium, *Pers. Syn.* ii. 76.—Hispan.
origanifolium, Pourr. ex Nym. Consp. 72 = ocymoides.
ovatum, Dun. in DC. Prod. i. 280 = vulgare.
pallens, *Kit. in Rochel, Pl. Banat. Rar. Ind.* nomen. —Banat.
paniculatum, Bolss. ex Willk. Ic. Pl. Hisp. ii. 140 = rubellum.
paniculatum, *Dun. in DC. Prod.* i. 278.—Hispan.
pannosum, *Boiss. Elench.* 15.—Hispan.
papillare, *Boiss. Voy. Bot. Espagne.* ii. 68. t. 14.—Reg. Mediterr. occ.
papillosum, Walp. Rep. i. 208 = prae.
papyraceum, *Sweet, ex Steud. Nom.* ed. II. i. 735 (Quid ?).—Europ.
patens, *Hensl. Diagn. Pl. Nov.* 20.—Mexic.
patulum, *Pomel, Nouv. Mat. Fl. Atl.* 354.—Alger.
pediforme, *Pomel. l. c.* 358.—Alger.
penicillatum, Thib. ex Dun. in DC. Prod. i. 277 = oelandicum.
pergamaceum, Pomel, Nouv. Mat. Fl. Atl. 350 = virgatum !
petiolatum, *Thib. ex Pers. Syn.* ii. 77.—Hispan.
piliferum, *Boiss. Elench.* 17.—Hispan.
piloselloides, Billot, ex Nym. Consp. 75 = vineale.
piloselloides, Dun. in DC. Prod. i. 284 = cinereum.
piloselloides, Pourr. ex Nym. Consp. 75 = paniculatum.
pilosum, Guir. ex Willk. Ic. Pl. Hisp. ii. 138 = Guirael.
pilosum, Mill. Gard. Dict. ed. VIII. n. 8 = ledifolium.
pilosum, Pers. Syn. ii. 79 = variabile.
planctagineum, Pers. l. c. 77 = guttatum.
polifolium, *DC. Fl. Fr.* v. 833.—Europ.
polifolium, Gren. & Godr. Fl. Fr. i. 170 = pulverulentum.
polifolium, Hook. & Arn. Bot. Beech. Voy. 410 = glomeratum.
polyfolium, Mill. Gard. Dict. ed. VIII. n. 11 ; Pers. Syn. ii. 80 = variabile.
polifolium, Tort. & Gray, Fl. N. Am. i. 151 = capitatum.
pollenis, *Kit. in Linnaea*, xxxii. (1863) 505.—Hungar.
polyanthum, *Pers. Syn.* ii. 78.—Afr. bor.
polygalaefolium, *Sweet, Cist.* t. 11.—Bess.
polymorphum, Pourr. ex Willk. Ic. Pl. Hisp. ii. 124 = apposum.
pomeridianum, *Dun. in Mém. Acad. Montp.* (*Sect. Sc.*) i. (1847) 8.—Alger.

HELIANTHEMUM :—

praecox, Salzm. ex Dun. in DC. Prod. i. 271 = guttatum.
procumbens, *Dun. l. c.* 275.—Tauria.
prostratum, *Pomel, Nouv. Mat. Fl. Atl.* 354.—Alger.
pulchellum, Sweet, Cist. t. 74 = cinereum.
pulverulentum, DC. Fl. Fr. iv. 823 = spaniinum.
pulverulentum, Pers. Syn. ii. 80 = variabile.
punctatum, Pers. l. c. 77 = guttatum.
punctatum, Willd. Enum. Hort. Berol. 570 = salicifolium.
racemosum, Dun. in DC. Prod. i. 281 = virgatum.
racemosum, Pers. Syn. ii. 79 = variabile.
ramulifolium, Mnhx. Fl. Bor. Am. i. 307 = canadense.
reflexum, Moench, Meth. 283 = variabile.
refractum, Frivald. ex Hampe, in Flora, xx. (1837) i. 227 = salicifolium.
retrofractum, Frivald. ex Nym. Consp. 78 = salicifolium.
retrofractum, *Pers. Syn.* ii. 78.—Hispan.
rhodanthum, Dun. in DC. Prod. i. 282 = variabile.
Rhodax, Steud. Nom. ed. II. i. 735 = cinereum.
Rochelii, *Kit. in Linnaea*, xxxii. (1863) 504.—Hungar.
rosetum, Barc. ex Nym. Consp. 74 = polifolium.
rosetum, DC. Fl. Fr. iv. 822 = vulgare.
rosetum, Ehrenb. ex Willk. Ic. Pl. Hisp. ii. 106 = vescarium.
roseum, *Mill. Gard. Dict.* ed. VIII. n. 17.—As. Min.
rosmarinifolium, Kunth, ex Willk. Ic. Pl. Hisp. ii. 135 = lavandulaefolium.
rosmarinifolium, Lag. ex DC. Prod. i. 267 = Ill. Libanotis.
rosmarinifolium, Mart. ex Steud. Nom. ed. II. i. 735 = variabile.
rosmarinifolium, Presl, Symb. Bot. i. 82 = Lippii.
rosmarinifolium, Purah, Fl. Am. Sept. ii. 364 = canadense.
Rossmaessleri, *Willk. in Linnaea*, xxx. (1859–60) 87. —Hispan.
rotatum, *Sweet, Hort. Brit.* ed. II. 42, nomen.—Hab ?
rotundifolium, Boiss. ex Willk. Ic. Pl. Hisp. ii. 141 = rubellum.
rotundifolium, Dun. in DC. Prod. i. 277 = paniculatum.
rubellum, Guir. ex Willk. Ic. Pl. Hisp. ii. 148 = paniculatum.
rubellum, Moench, Meth. Suppl. 74 = salicifolium.
rubellum, *Presl, Delic. Prag.* 35.—Reg. Mediterr. occ.
rufiooman, *Spreng. Syst.* iii. 889.—Afr. bor.
rugosum, Dun. in DC. Prod. i. 268 = alyssoides.
rugosum, Sweet, Cist. t. 65 = ocymoides.
rupicolum, Schur, Enum. Pl. Transs. 77 = vulgare.
rupifragum, *A. Kern. in Oestr. Bot. Zeitschr.* xvii. (1868) 18.—Europ. austr.
salicifolium, *Mill. Gard. Dict.* ed. VIII. n. 21 ; *Pers. Syn.* ii. 78.—Europ. austr. ; Oriens.
sampsuchifolium, Mill. l. c. n. 7 = variabile.
sanguineum, Host, ex Willk. Ic. Pl. Hisp. ii. 90 = salicifolium.
sanguineum, Lag. ex Dun. in DC. Prod. i. 273 = retroflactum.
Savii, Bertol. Amoen. Ital. 78 = arabicum.
scabrosum, Pers. Syn. ii. 76 = alyssoides.
scariosum, Duf. ex Willk. Ic. Pl. Hisp. ii. 125 = asperum.
scoparium, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 152. —Am. occ.
scorpioides, Coss. Notes Crit. ii. 29 = echioides.
Segnierii, Pollini, Viagg. Lag. Garda, 98 = vulgare.
semiglabrum, *Badaro, in Moretti, Bibl. Ital.* i. 14.—Sardin.
semiglabrum, Moris, Fl. Sard. i. 202 = Morisianum.
semistipulatum, Lag. ex Dun. in DC. Prod. i. 271 = guttatum.
sericeum, Pourr. ex Nym. Consp. 72 = alyssoides.
serpyllifolium, Mill. Gard. Dict. ed. VIII. n. 8 = vulgare.
Servat, Cambess. in Mém. Mus. Par. xiv. (1827) 216. t. 11 = origanifolium.
serratum, Mérat, Fl. Par. 206; ed. II. 391 = guttatum.
sessiliflorum, Pers. Syn. ii. 78 = Lippii.
Sexte, *Lag. ex Dun. in DC. Prod.* i. 278.—Hispan.
Sherardi, Steud. Nom. ed. II. i. 735 = roseum.
Sibthi x, Breegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 78.—Europ.
siculum, *Tineo, Pl. Rar. Sicil.* 44.—Sicilia.
sonnenbornum, *Schrenk, Enum. Pl. Nov.* 94.—Soongaria.
Spachii, Nym. Syll. 226 = H. Fumana.

HELIANTHEMUM :—

spartioides, *Presl, Rel. Haenk.* ii. 95.—Chili.
splendens, Pourr. ex Nym. Consp. 72 = H. Libanotis.
squamatum, *Pers. Syn.* ii. 78.—Hisp. ; Afr. bor.
Stabianum, Heldr. ex Willk. Ic. Pl. Hisp. ii. 114 = vulgare.
Stabianum, Tenore, Fl. Nap. iv. 318 = glaucum.
staechadifolium, Auct. vide *stoechadifolium*.
stellatum, Guss. Fl. Sic. Prod. Suppl. 178 = glaucum.
stellulatum, *Link, ex Steud. Nom.* ed. II. i. 735.—Hab ?
stipulare, *Ehrenb. ex Willk. Ic. Pl. Hisp.* ii. 160.—Hispan.
stoechadifolium, Pers. Syn. ii. 79 = glaucum.
stoechadifolium, Tenore, Fl. Nap. i. 298 = polifolium.
stoechadifolium, Willk. in Flora, xxxiv. (1851) 595 = lavandulaefolium.
stramineum, Sweet, Cist. t. 93 = variabile.
strictum, Pers. Syn. ii. 79 = virgatum.
strigosum, Fisch. Hort. Gorenk. ed II. (1812) 57 = cinereum.
sulphureum, Willd. Enum. Hort. Berol. Suppl. 39 = variabile.
surrejanum, Mill. Gard. Dict. ed. VIII. n. 15 = variabile.
syriacum, Boiss. Diagn. Ser. I. viii. 49 = umbellatum.
syriacum, Ehrenb. ex Willk. Ic. Pl. Hisp. ii. 164 = arabicum.
syrticum, *Spreng. Syst.* ii. 593.—Afr. bor.
tauricum, *Rupr. in Ind. Sem. Hort. Petrop.* (1851) ; *ex Bull. Phys-Math. Acad. Petersb.* xiv. (1856) 236.—Tauria.
tauricum, Sweet, Cist. t. 105 = variabile.
Tenerifae, *Coss. in Bull. Soc. Bot. Fr.* iii. (1856) 561. —Ins. Canar.
tenuicaulis, *Pomel, Nouv. Mat. Fl. Ital.* 222.—Alger.
tereifolium, Thib. ex Willk. Ic. Pl. Hisp. ii. 123 = hirtum.
ternifolium, Colm. & Willk. in Bot. Zeit. v. (1847) 236 = alyssoides.
thessalum, Boiss. & Orph. ex Orph. in Bull. Congr. Bot. Petersb. 1869 (1870) 136 = oelandicum.
Thibaudi, Pers. Syn. ii. 79 = lavandulaefolium.
thymifolium, *Pers l. c.*—Hispan. ; *Lusit.*
tomentosum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 663 ; Dun. in DC. Prod. i. 279 = vulgare.
tomentosum, Guss. Fl. Sic. Prod. Suppl. 177 = glaucum.
trisforum, Blanco, Fl. Filip. ed. II. 809 = Archytaea Vahli.
tripetalum, Miégev. in Bull. Soc. Bot. Fr. xiv. (1867) 144 = salicifolium.
tripetalum, Moq. & Sesse, ex Dun. in DC. Prod. i. 284 = Lechea Skinneri
Tuberaria, Hook. Bot. Mag. t. 4373 = globulariaefolia.
Tuberaria, *Mill. Gard. Dict.* ed. VIII. n. 10.—Reg. Mediterr.
tunetanum, *Coss. & Kralik, in Bull. Soc. Bot. Fr.* iv. (1857) 58.—Afr. bor.
umbellatum, *Mill. Gard. Dict.* ed. VIII. n. 5.—Reg. Mediterr.
umbellatum, Presl, Fl. Sic. i. 118 = Cistus Clusii.
variabile, *Spach, in Ann. Sc. Nat.* Sér. II. vi. (1836). 2, 362.—Reg. Mediterr.
variegatum, Sweet, Cist. t. 88 = variabile.
velutinum, Jord. Obs. Pl. Crit. iii. (1846) 25 = polifolium.
velutinum, *Pomel, Nouv. Mat. Fl. Atl.* 351.—Alger.
ventosum, *Boiss. Diagn.* Ser. I. viii. 50.—Arab.
vernutum, Sweet, Cist. t. 10 = variabile.
versicolor, Sweet, l. c. 26 = variabile.
verticillatum, Link, Handb. ii. 865 = umbellatum.
vescarium, *Boiss. Diagn.* Ser. I. viii. 50.—Aegypt. ; Palest.
villosissimum, *Pomel, Nouv. Mat. Fl. Atl.* 216.—Alger.
villosum, *Thib. ex Pers. Syn.* ii. 78.—Hispan. ; Armen.
vineale, *Pers. l. c.* 77.—Europ.
violaceum, *Langé, in Kjoeb. Vidensk. Meddel.* (1865) 91.—Hispan.
violaceum, Pers. Syn. ii. 78 = pilosum.
virens, Link, Handb. ii. 868 = *viride*.
virgatum, *Pers. Syn.* ii. 79.—Reg. Mediterr. occ.
virgatum, *Willk. Ic. Pl. Hisp.* ii. 106. t. 135.— Ins. Balear.
viride, Tenore, Prod. Fl. Nap. p. xxxi = H. Fumana.
viscarium, *Boiss. & Reut. Pugill. Pl. Nov.* 14.—Alger.
viscidulum, *Boiss. Elench.* 15.—Hispan.

HELIANTHEMUM:—
viscidulum, Stev. ex Dun. in DC. Prod. i. 275 = arabicum.
viscidum, Pomel, Nouv. Mat. Fl. Atl. 217 = glutinosum ?
Vivianii, *Pollini, Fl. Veron.* iii. 799.—Ital.
vulgare, *Gærtn. Fruct.* i. 371. t. 76.—Europ.
vulgare, Gmn. ex Nym. Consp. 74 = glaucum.

HELIANTHOSTYLIS, Baill. Adansonia, xi. (1875) 299. *URTICACEÆ*, Benth. & Hook. f. iii. 374. Spruce, *Baill. l. c.*—Am. trop.

HELIANTHUS, Linn. Syst. ed. I (1735). *COMPOSITÆ*, Benth. & Hook. f. ii. 376.
CHRYSIS, Renealm. ex DC. Prod. v. 586 (1836).
DIOMEDRA, Bertol. ex Colla, in Mem. Acc. Torin. xxxvii. (1835) 36. t. 81.
DISCOMELA, Raffin. Neogenyt. 3 (1825).
ECHINOMERIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. 256 (1841).
FLORENSIA, DC. Prod. v. 592 (1836).
HARPALIUM, Cass. in Bull. Soc. Philom. (1818) 141.
LINSECOMIA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 458.
VOSACAN, Adans. Fam. ii. 130 (1763).
altissimus, DC. Prod. v. 590 = strumosus.
altissimus, Linn. Sp. Pl. ed. II. 1278 = giganteus.
amplexicaulis, *DC. Prod.* v. 589.—Mexic.
angustifolius, *Linn. Sp. Pl.* 906.—Am. bor.
annuus, *Linn. l. c.* 904.—Am. bor.
apetalus, Lecoste, ex Benth. in Trans. Am. Phil. Soc. N. S. vii. (1841) 354 = H. Radula.
argenteus, *H. B. & K. Nov. Gen. et Sp.* iv. 221. t. 376. —Ecuador.
argophyllus, *Torr. & Gray, Fl. N. Am.* ii. 318.—Texas.
aristatus, Ell. Sketch. ii. 428 = Actinomeris nudicaulis.
asper, Hort. ex Steud. Nom. ed. II. i. 787 = rigidus.
atrorubens, Lam. Encyc. iii. 86 = lætiflorus.
atrorubens, *Linn. Sp. Pl.* 906.—Am. bor.
atrorubens, Michx. Fl. Bor. Am. ii. 140, partim = rigidus.
atrorubens, *H. B. & K. Nov. Gen. et Sp.* iv. 224.—Ecuador.
Besseriana, Benth. & Hook. f. Gen. ii. 376 = Flourensia Besseriana.
Bolanderi, A. Gray, in Proc. Am. Acad. vi. (1865) 544 = scaberrimus.
brasilicus, *Vell. Fl. Flum.* viii. t. 140.—Bras.
buddleiæformis, DC. Prod. v. 588 = Viguiera buddleiæformis.
buphthalmoides, Hort. ex DC. l. c. 551 = Heliopsis buphthalmoides.
californicus, *DC. l. c.* 589.—Am. bor. occ.
californicus, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 325 = Nuttallii.
calvus, *Sek. Bij. in Linnæa*, xxxiv. (1865–66) 528, nomen.—Bolivia.
canescens, Juss. ex DC. Prod. v. 591 = incanus.
canescens, Michx. Fl. Bor. Am. ii. 140 = mollis.
carolinianus, Hill, Veg. Syst. xvi. 56 = giganteus !
cernuus, Benth. & Hook. f. Gen. ii. 376 = Flourensia cernua.
ciliaris, *DC. Prod.* v. 587.—Mexic.
cinereus, *Torr. & Gray, Fl. N. Am.* ii. 324.—Texas.
cirrhoides, *Lehm. in Hamb. Gartens.* viii. (1853) 458. —Hab. ?
cochlachinensis, *Pers. Syn.* ii. 476.—Cochinch.
colossus, *Kunze, in Otto & Dietr. Allg. Gartens.* vii. (1839) 43.—Am. bor.
cordatus, Lam. ex DC. Prod. v. 551 = Heliopsis buphthalmoides.
cornifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 223.—Mexic.
corymbosus, Poepp. ex DC. Prod. v. 592 = Viguiera Poeppigii.
crassifolius, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 367 = rigidus.
crinitus, Steud. Nom. ed. II. i. 787 = giganteus.
cucumerifolius, Torr. & Gray, Fl. N. Am. ii. 319 = debilis.
deflexus, *A. Gray, Syn. Fl. N. Am.* i. ii. 380.—Calif.
debilis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 367.—Am. bor.
decapetalus, Darl. Fl. Cestr. ed. II. 483 = strumosus.
decapetalus, *Linn. Sp. Pl.* 905.—Am. bor.
dentatus, Cav. Ic. iii. 10. t. 220 = Viguiera dentata.
diffusus, *Moench, Meth. Suppl.* 259.—Hab. ?
diffusus, Sims, Bot. Mag. t. 2090 = rigidus.
divaricatus, *Linn. Sp. Pl.* 906.—Am. bor.

HELIANTHUS:—
divaricatus, Michx. Fl. Bor. Am. ii. 141 = parviflorus.
diversifolius, Ell. Sketch, ii. 428 = hirsutus.
doronicoides, *Lam. Encyc.* iii. 84.—Am. bor.
doronicoides, Torr. & Gray, Fl. N. Am. ii. 327, partim = tuberosus.
Douglasii, *Torr. & Gray, l. c.* 322.—Am. bor. occ.
Dumeilianus, M. A. Curt. in Am. Journ. Sc. xliv. (1843) 33 = occidentalis.
erythrocarpus, Bartl. in Linnæa, xiv. (1840) Litt. 125 = annuus.
esculentus, *Rottb. in Act. Litt. Univ. Hafn.* i. (1778) 296.—Guiana.
esculentus, *Warsc. in Otto & Dietr. Allg. Gartens.* xx. (1852) 293.—Bolivia.
excelsus, Willd. Sp. Pl. iii. 2248 = Viguiera excelsa.
exilis, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 545. —Calif.
floridanus, *A. Gray, ex Chapm. Fl. S. U. St. Suppl.* 529.—Florida.
frondosus, Linn. Cent. Pl. i. 28; Amœn. Acad. iv. 290 = decapetalus.
giganteus, Cav. Ic. iii. 10. t. 219 = Viguiera excelsa.
gigantеus, *Linn. Sp. Pl.* 905.—Am. bor.
gigantеus, Lour. Fl. Cochinch. 509 (sp. ignot.).—China.
Gigas, Michx. Fl. Bor. Am. ii. 141 = giganteus.
glutinosus, Hook. & Arn. Bot. Beech. Voy. 38 = thurifer.
gracilentus, *A. Gray, in Proc. Am. Acad.* xi. (1876) 77.—Calif.
gracilis, *Bertol. Misc. Bot.* vii. 41.—Am. bor.
grandiflorus, Juss. ex Pers. Syn. ii. 475 = decapetalus.
grandiflorus, Wender. ex Steud. Nom. ed. II. i. 787 = annuus.
grandiflorus, *Wood, Am. Bot. Flor.* 178.—Am. bor.
grosse-serratus, *Martius, Sel. Sem. Hort. Low.* (1839) ; *in Linnæa*, xiv. (1840) *Litt.* 128.—Am. bor. occ.
grosse-serratus, Pohl, ex Baker, in Mart. Fl. Bras. vi. iii. 174 = Wulffia stenoglossa.
herbaceus, *Vell. Fl. Flum.* viii. 145.—Bras.
heterophyllus, Willd. ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 314 = Viguiera Gilliesii.
heterophyllus, *Nutt. in Journ. Acad. Philad.* vii. (1834) 74.—Am. bor.
heterophyllus, Short, Cat. Kentuck. Pl. Suppl. 8 = occidentalis.
hirsutus, *Raffin. Ann. Nat.* (1820) 14.—Am. bor.
hispidus, Ell. Sketch, ii. 429 = hirsutus.
Hookeri, G. Don, in Loud. Hort. Brit. 358 = doronicoides.
Hookerianus, DC. Prod. v. 590 = Wyethia angustifolia.
incanus, *Pers. Syn.* ii. 475.—Peruv.
indicus, Linn. Mant. i. 117 = annuus.
integrifolius, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 366 = petiolaris.
laciniatus, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 84.—Mexic.
lætiflorus, *Pers. Syn.* ii. 476.—Am. bor. occ.
lævigatus, *Torr. & Gray, Fl. N. Am.* ii. 330.—Am. bor.
lævis, *Linn. Sp. Pl.* 906 = Heliopsis lævis.
lævis, Walt. Fl. Carol. 215 = strumosus !
lanceolatus, *Meyen, Reise*, i. 311.—Chili.
lanuginosus, Benth. & Hook. f. Gen. ii. 376 = Flourensia lanuginosa.
leiostemon, Raffin. ex DC. Prod. v. 588 = divaricatus.
leptolaciniatus, Dougl. in Bot. Reg. t. 1966 = annuus.
Lindheimerianus, Scheele, in Linnæa, xxii. (1849) 159 = debilis.
linearis, Cav. Ic. iii. 9. t. 218 = Viguiera linearis.
lingularifolius, *Bertol. Pl. Centtin.* 36.—Guatemala.
longifolius, Hook. Fl. Bor. Am. i. 313 = Wyethia angustifolia.
longifolius, *Pursh, Fl. Am. Sept.* ii. 571.—Am. bor.
macrospus, DC. Fl. Far. Jard. Gen. 7me Not. 8 = annuus.
macrophyllus, Willd. Hort. Berol. t. 76 = strumosus.
Maximowiczii, *Sch. Bip. in Linnæa*, xxiv. (1851–66) 525, nomen.—Bolivia.
Maximiliani, *Schrad. Ind. Sem. Hort. Gotting.* (1836) ; *ex Ann. Sc. Nat.* Sér. II. vi. (1836) 103.—Am. bor. occ.
membranifolius, *Poir. Encyc. Suppl.* iii. 12.—Guian.
mexicanus, *Humb. Bial. Centr. Am. Bot.* ii. 180.—Mexic.
microcephalus, *Sprang. Syst.* iii. 618.—Mexic.

HELIANTHUS:—
microcephalus, Torr. & Gray, Fl. N. Am. ii. 329 = parviflorus.
microphyllus, H. B. & K. Nov. Gen. et Sp. iv. 230 = Leighia microphylla.
missouriensis, Schwein. ex Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 366 = rigidus.
missuricus, Spreng. Nov. Prov. 21 = rigidus.
mollis, *Lam. Encyc.* iii. 85.—Am. bor.
mollis, Willd. Sp. Pl. iii. 2240 = strumosus.
multiflorus, Hook. Fl. Bor. Am. i. 318 (excl. syn.) = annuus.
multiflorus, Linn. Sp. Pl. 905 = decapetalus.
neglectus, Otto, ex A. Gray, Syn. Fl. N. Am. i. ii. 379 = strumosus *vel* lætiflorus.
Nuttallii, *Torr. & Gray, Fl. N. Am.* ii. 324.—Am. bor. occ.
occidentalis, *Riddell, Suppl. Cat. Ohio Pl.* 18.—Am. bor.
oleifer, Wall. Cat. n. 3194 = Guizotia abyssinica.
Oliveri, *A. Gray, in Proc. Am. Acad. xx.* (1885) 299. —Calif.
orgyalis, *DC. Prod.* v. 586.—Am. bor. occ.
ovatus, Lehm. Ind. Sem. Hort. Hamb. (1828) 16 = annuus.
Parishii, *A. Gray, in Proc. Am. Acad. xix.* (1883) 7.—Calif.
parviflorus, *Bernh. ex Spreng. Syst.* iii. 617.—Am. bor.
parviflorus, H. B. & K. Nov. Gen. et Sp. iv. 222. t. 378 = microcephalus.
patens, Lehm. Ind. Sem. Hort. Hamb. (1831) 8 = petiolaris.
pauciflorus, *Nutt. Gen. Am.* ii. 177.—Louisiana.
pedunculatus, Hort. ex Link, Enum. Hort. Berol. ii. 252 = rigidus !
peruvianus, *Klotzsch, in Otto & Dietr. Allg. Gartens.* iii. (1835) 389.—Peruv.
petiolaris, Hort. ex DC. Prod. v. 586 = annuus.
petiolaris, *Nutt. in Journ. Acad. Sc. Philad.* ii. (1821) 115.—Am. bor. occ.
pilosus, Tausch, in Flora, xi. (1828) 502 = doronicoides.
platycephalus, Cass. in Dict. Sc. Nat. xx. 352 = annuus.
polycephalus, Cass. l. c. = decapetalus.
praecox, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1847) 221 = debilis.
procumbens, *Pers. Syn.* ii. 475.—Peruv.
procumbens, *Raffin. Fl. Ludov.* 70 (Quid ?).—Am. bor.
prostratus, Willd. Sp. Pl. iii. 2243 = trachelifolius.
pubescens, Hook. Bot. Mag. t. 2778 = doronicoides.
pubescens, [Ker-Gawl. in] Bot. Reg. t. 594 = tomentosus.
pubescens, Vahl, Symb. Bot. ii. 92 = mollis.
pumilus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 366.—Am. bor. occ.
pumilus, Pers. Syn. ii. 475 = annuus.
quinquenervis, Hook. Lond. Journ. Bot. vi. (1847) 247 = Helianthella quinquenervis.
quinquenervius, *Cav. Ic.* iii. 36. t. 272.—Mexic.
Radula, *Torr. & Gray, Fl. N. Am.* ii. 321.—Am. bor.
ramosissimus, *Mill. Gard. Dict.* ed. VIII. n. 6.—Am. austr.
revolutus, *Meyen, Reise*, i. 311.—Chili.
rigidus, *Desf. Cat. Hort. Par.* ed. III. 184.—Am. bor. occ.
rivularis, Poepp. ex DC. Prod. v. 495 = Leptocarpha rivularis.
rugosus, *Meyen, Reise*, ii. 45.—Peruv.
rugosus, Schau. in Linnæa, xix. (1847) 738 = Viguiera rugosa.
salicifolius, *A. Dietr. in Otto & Dietr. Allg. Gartens.* ii. (1834) 387.—Am. bor.
sarmentosus, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 113 (err. typ. 105) = Wulffia stenoglossa.
scaber, Pohl, ex Baker, in Mart. Fl. Bras. vi. iii. 174. —Wulffia stenoglossa.
scaberrimus, *Benth. Bot. Voy. Sulph.* 26.—Calif.
scaberrimus, Ell. Sketch, ii. 429 = rigidus.
Schweinitzii, *Torr. & Gray, Fl. N. Am.* ii. 326.—Am. bor.
serotinus, Tausch, in Flora, xi. (1828) 504.—Am. bor.
silphioides, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 366 = strumosus.
spathulatus, A. Gray, Syn. Fl. N. Am. i. ii. 369 (err. typ.).—Mexic.
spurcifolius, Ell. Sketch, ii. 415 = strumosus.

HELIANTHUS:—
spathulatus, Ell. Sketch, ii. 421 = tomentosus.
speciosus, Hook. Bot. Mag. t. 2295 = Tithonia speciosa.
squarrosus, H. B. & K. Nov. Gen. et Sp. iv. 222. t. 377 = Viguiera linearis.
squarrosus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 367 = tomentosus.
striatus, *Rafin. Atl. Journ.* 17 (Quid ?).—Am. bor.
strumosus, *Linn. Sp. Pl.* 905.—Am. bor.
strumosus, Willd. Sp. Pl. iii. 2243 = decapetalus.
subtuberosus, Bourg. ex A. Gray, Syn. Fl. N. Am. i. ii. 274 = giganteus.
tenellus, *Rafin. Atl. Journ.* 17 (Quid ?).—Am. bor.
tenuifolius, Ell. Sketch, ii. 420 = decapetalus.
tephrodes, A. Gray, in Torr. Bot. Mex. Bound. 90 = Viguiera tephrodes.
tetragonus, Moç. ex DC. Prod. v. 576 = Actinomeris tetragona.
tetrapterus, Orteg. Hort. Matr. Dec. 74 = Actinomeris tetraptera.
tharifer, *Molina, Sagg. Chile*, 160.—Chili.
tomentosus, *Michx. Fl. Bor. Am.* ii. 141.—Am. bor.
trachelifolius, *Mill. Gard. Dict.* ed. VIII. n 7 ; *Willd. Sp. Pl.* iii. 2241.—Am. bor.
trachelifolius, Hook. Comp. Bot. Mag. i. (1835) 98 = parvifloras.
tricuspis, Ell. Sketch, ii. 422 = laeviflorus.
trilobatus, *Link, Enum. Hort. Berol.* ii. 252.—Mexic.
trinervis, Hort. Madr. ex DC. Prod. v. 585 = Viguiera excelsa.
trinervis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 18 (Quid ?).—Java.
tripunctus, Ortog. Hort. Matr. Dec. 76 = Viguiera helianthoides.
truncatus, Schwein. ex Ell. Sketch, ii. 416 = divaricatus.
tuxillensis, *H. B. & K. Nov. Gen. et Sp.* iv. 223.—Peruv.
tuberiformis, Nutt. Gen. Am. ii. 177 = annuus.
tuberiformis, Ortog. Hort. Matr. Dec. 181 = Tithonia tuberiformis.
tuberosus, *Linn. Sp. Pl.* 905.—Am. bor.
tuberosus, Parry, in Owen, Rep. Minnesot. Surv. 614 = giganteus.
uniflorus, Nutt. in Journ. Acad. Philad. vii. (1834) 87 = Helianthella uniflora.
verbenaloides, *H. B. & K. Nov. Gen. et Sp.* iv. 221.—Peruv.
vicetinus, *Turra, ex Zang. in Atti Soc. Sc. Nat. Milan*, xi. 1868 (1869) 889.—Hab. ?
virgatus, Lam. Encyc. iii. 85 = giganteus.

HELICANDRA, Hook. & Arn. Bot. Beech. Voy. 198, in nota (1838?) = Parsonsia, R. Br. (Apocyn.).
sinensis, Hook. & Arn. l. c. = P. Helicandra.

HELICANTHERA, Roem. & Schult. Syst. v. pp. x, 170 (1819) = HELIKANTHERA, Laur. = Loranthus, Linn.

HELICHROA, Rafin. Neogenyt. 8 (1825) = Echinacea, Moench (Compos.).
alba, *amœna*, *crocea*, *elatior*, *fusca* (*fuscata*, Atl. Journ. 150), *Linnæana*, *purpurea*, *uniflora*, Rafin. l. c. = E. purpurea.

HELICHRYSUM, Vaill. ex Linn. Gen. ed. I. 250 (1737) ; Gaertn. Fruct. ii. 404 (1791) *COMPOSITÆ*, Benth. & Hook. f. ii. 309.
Acanthocladium, F. Muell. Fragm. ii. 155 (1860–61).
Achillexis, D. Don, in Mem. Wern. Soc. v. (1826) 546.
Argyroglottis, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 84.
Argyrophanes, Schlecht. in Linnæa, xx. (1847) 896.
Cretocephalum, Walp. in Linnæa, xiv. (1840) 508.
Cladochaeta, DC. Prod. v. 245 (1837).
Conanthodium, A. Gray, in Hook. Kew Journ. iv. (1852) 272.
Ebenoidia, Cass. in Bull. Soc. Philom. (1818) 75 ; et in Dict. Sc. Nat. xiv. 393 (1819).
Eucretium, Tourn. ex Linn. Gen. ed. I. 250 (1737).
Eucullia, D. Don, in Mem. Wern. Soc. v. (1826) 548.

HELICHRYSUM:—
Faustula, Cass. in Bull. Soc. Philom. (1818) 140 ; et Dict. Sc. Nat. xvi. 251 (1820).
Freemania, Boj. ex DC. Prod. vi. 218 (1837).
Lawrencella, Lindl. Swan River App. 28 (1839).
Leriche, Less. Syn. Comp. 296 (1832).
Lerichiline, Cass. in Bull. Soc. Philom. (1818) 81.
Leucostemma, D. Don, in Mem. Wern. Soc. v. (1826) 540.
Manopappus, Sch. Bip. in Flora, xxvii. (1844) 677.
Ozothamnus, R. Br. in Trans. Linn. Soc. xii. (1816) 125.
Pentataxis, D. Don, in Mem. Wern. Soc. v. (1826) 550.
Petalolepis, Cass. in Bull. Soc. Philom. (1817) 138.
Swammerdamia, DC. Prod. vi. 164 (1837).
Trichandrum, Neck. Elem. i. 84 (1790), ex char.
Xanthochrysum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 199.
abyssinicum, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) i. *Intell.* 26, nomen ; *et in Bot. Zeit.* iii. (1845) 174.—Afr. trop.
aciculare, *Balf. f. in Proc. Roy Soc. Edinb.* xi. (1882) 839.—Ins. Socotr.
acuminatum, DC. Prod. vi. 188 = bracteatum.
acuminatum, Sweet, Hort. Brit. ed. I. 223 = Gnaphalium acuminatum.
acutatum, *DC. Prod.* vi. 186.—Afr. austr.
adenocarpum, *DC. l. c.* 180.—Afr. austr.
adenophorum, *F. Muell. in Trans. Vict. Inst.* i. (1855) 38.—Austral.
adnatum, *Benth. Fl. Austral.* iii. 628.—Austral.
adscendens, DC. Prod. vi. 185 = pilolepis.
adscendens, *Less. Syn. Comp.* 274.—Afr. austr.
affine, D. Don, ex Loud. Hort. Brit. Suppl. ii. 640 = Gnaphalium luteo-album.
affine, Less. Syn. Comp. 293 = argenteum.
alatum, Baill. Hist. Pl. viii. 302, 314 = Ammobium alatum.
albicans, A. Cunn. in Field, N. S. Wales, 359 = Helipterum incanum.
albicans, Sieber, ex DC. Prod. vi. 193 = elatum.
algarbiensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 65 = H. Stoechas.
alloides, *Less. Syn. Comp.* 299.—Afr. austr.
alnioides, *DC. Prod.* vi. 169.—Afr. austr.
alveolatum, DC. l. c. 192 = trilineatum.
ambiguum, *Presl, Fl. Sic.* p. xxix (Quid ?).—Sicilia.
ambiguum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 196 = Leptorhynchus ambiguus.
amorginum, *Boiss. & Orph. Diagn.* Ser. II. v. 110.—Ins. Cyclad.
amplexicaule, *Baker, in Journ. Linn. Soc.* xx. (1883) 185.—Madag.
anatolicum, Boiss. Diagn. Ser. I. iv. 11 = plicatum.
angustifolium, *DC. Fl. Fr. Suppl.* 467.—Reg. Mediterr. or.
angustifolium, Pers. Syn. ii. 415 = paniculatum.
angustifolium, Plan. Fl. Fan. Galleg. 270 = serotinum.
anomalum, *Less. Syn. Comp.* 303.—Afr. austr.
antennarioides, *F. Muell. ex Benth. Fl. Austral.* iii. 632. —Austral.
anthemoides, Sieber, ex Spreng. Syst. iii. 484 = Helipterum anthemoides.
aphelexioides, *DC. Prod.* vi. 206.—Madag.
apiculatum, *D. Don, in Mem. Wern. Soc.* v. (1824) 550.—Austral.
appendiculatum, *Less. Syn. Comp.* 308.—Afr. austr.
arachnoides, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 839.—Ins. Socotr.
arborum, D. Don, in Mem. Wern. Soc. v. (1824) 550 = Anaxeton arborosum.
arenarium, *Moench, Meth.* 576 ; *DC. Fl. Fr.* iv. 132.—Europ. ; Reg. Caucas.
arenicolum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 79 = Bryomorphe Zeyheri.
argenteum, Andr. ex Stend. Nom. ed. II. i. 788 (740 cum cit. falsa) = paniculatum.
argenteum, Schrank, in Denkschr. Akad. Muench. viii. 1821–22 (1824) 167 = retortum.
argenteum, *Thunb. Fl. Cap.* 662.—Afr. austr.
argyreum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 66 = Italicum.
argyrocaule, F. Muell. ex Sond. in Linnæa, xxv, (1852) 515 = dealbatum.
Argyroglottis, *Benth. Fl. Austral.* iii. 626.—Austral.

HELICHRYSUM:—
argyrophyllum, *DC. Prod.* vi. 186.— Afr. austr.
argyrolepis, *DC. l. c.* 174.—Afr. austr.
armenium, *DC. l. c.* 183.—As. Min. ; Persia.
artemisioides, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 287.—Persia.
asteroides, DC. Prod. vi. 194 = Ixiolaena viscosa.
Auctori, *Boiss. Diagn.* Ser. I. xi. 28.—As. Min. ; Persia.
aurantiacum, Boiss. & Huet, Diagn. Ser. II. vi. 105 = lavandulifolium.
auriculatum, E. Mey. in Drège, Zwei Pfl. Docum. 189 = stellatum.
auriculatum, *Less. Syn. Comp.* 311.—Afr. austr. et trop. aurunitzen, *Sch. Bip. in Flora*, xxvii. (1844) 680.—Afr. austr.
Ayersii, *F. Muell. Fragm.* viii. 67.—Austral.
baccharoides, *F. Muell. ex Benth. Fl. Austral.* iii. 633. —Austral.
Backhousii, *F. Muell. ex Benth. l. c.* 632.—Austral.
Banksii, A. Cunn. ex DC. Prod. vi. 188 = bracteatum.
barbellatum, Bunk, Ind. ad DC. Prod. ii. p. vi = cespitosum.
Barrelieri, Nym. Consp. 381 = H. Stoechas.
Baxteri, *A. Cunn. ex DC. Prod.* vi. 193.—Austral.
Beckleri, *F. Muell. ex Benth. Fl. Austral.* iii. 627.— Austral.
Beiengerianum, DC. Prod. vi. 184 = leucocephalum.
bellidioides, *Willd. Sp. Pl.* iii. 1911.—N. Zel.
bisfranum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 302.—Afr. trop.
bicolor, Lindl. Bot. Reg. t. 1814 = bracteatum.
bicolorum, A. Cunn. ex DC. Prod. vi. 215 = Helipterum incanum.
Bidwillii, *Benth. Fl. Austral.* iii. 627.—Austral.
Billardieri, *Boiss. & Blanche, Diagn.* Ser. II. v. 111 — Syria.
Blanchyrsklanum, *Steetz, ex Sond. in Linnæa*, xxv. (1852) 512.—Austral.
Boissieri, *Nym. Consp.* 381.—Hispan.
Bojerianum, *DC. Prod.* vi. 208.—Madag.
brachycladum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 63 = H. Stoechas.
bracteatum, *Andr. Bot. Rep.* sub t. 428 ; *et Willd. Enum. Hort. Berol.* 895.—Austral.
bracteolatum, *Benth. Fl. Austral.* iii. 630.—Austral.
brevicilium, DC. Prod. vi. 196 = sempapposum.
buchtormense, DC. l. c. 184 = arenarium.
buddleioides, *DC. ex Wight, Contrib.* 10.—Ind. or.
Bulsaenum, *Boiss. Fl. Orient.* iii. 287.—Persia.
bullatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 184. —Madag.
bupthalmoides, Sieber, ex DC. Prod. vi. 194 = scorpioides.
Burchellii, *DC. Prod.* vi. 196.—Afr. austr.
caespititium, *Sond. ex Harv. & Sond. Fl. Cap.* iii. 217. —Afr. austr.
caespitosum, Auct. vide cespitosum.
callichrysum, *DC. Prod.* vi. 183.—As. Min. ; Persia.
callicomum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 247. —Afr. austr.
callunoides, *Sch. Bip. in Flora*, xxvii. (1844) 677.—Afr. austr.
Calvertianum, *F. Muell. Fragm.* x. 108.—Austral.
candidissimum, *D. Don, ex G. Don, in Loud. Hort. Brit.* 841.—Reg. Caucas.
Candollеanum, Bunk, Ind. ad DC. Prod. ii. p. vi = Ixiolaena leptolepis.
canescens, Willd. Sp. Pl. iii. 1906 = Helipterum canescens.
capillaceum, *Less. Syn. Comp.* 275.—Afr. austr.
capitellatum, *D. Don, in Mem. Wern. Soc.* v. (1824) 550 ; *Less. Syn. Comp.* 305.—Afr. austr.
Casimirianum, Gandich. in Freyc. Voy. Bot. 466. t. 87 = Schouia Casimiana.
cassinioides, *Benth. Fl. Austral.* iii. 630.—Austral.
Catipes, *Harv. in Harv. & Sond. Fl. Cap.* iii. 223.— Afr. austr.
cephaloideum, *DC. Prod.* vi. 197.—Afr. austr.
Cephalotes, D. Don, ex G. Don, in Loud. Hort. Brit. 841 = Metalasia Cephalotes.
cerastioides, *DC. Prod.* vi. 171.—Afr. austr.
cespitosum, *DC. l. c.* 173. n. 28.—Ins. Maurit.
cespitosum, DC. l. c. 182. n. 70 = conglobatum.
chilense, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 236.—Chili.
chlorophyllum, Bolss. & Bal. Diagn. Ser. II. vi. 108 = callichrysum.

HELICHRYSUM:—
chionospharum, *DC. Prod.* vi. 174.—Afr. austr.
chlorochrysum, *DC. l. c.* 179.—Afr. austr.
chlorolepis, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 68 = serotinum.
chrysanthum, Pers. Syn. ii. 414 = bracteatum.
chrysocephalum, Sch. Bip. in Bot. Zeit. iii. (1845) 174 = cymosum.
Chrysocoma, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 434 = cymosum.
chrysophorum, Boiss. ex Boiss. Fl. Orient. iii. 280 = callichrysum.
ciliatum, DC. Prod. vi. 195 = semipapposum.
cinereum, *F. Muell. ex Benth. Fl. Austral.* iii. 629.—Austral.
cirrhosum, *Less. Syn. Comp.* 282.—Madag.
citrinum, Less. ex Harv. & Sond. Fl. Cap. iii. 269 = Helipterum citrinum.
citrinum, Nym. Consp. 381 = H. Stoechas.
citrispinum, *Delile, in Ann. Sc. Nat.* Sér. II. xx. (1843) 93.—Abyssin.
cladochaetum, *F. Muell. Fragm.* v. 199.—Austral.
cochinchinense, *Spreng. Syst.* iii. 482.—Cochinch.
cochleariforme, *DC. Prod.* vi. 185.—Afr. austr.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 781, *Index* (154 sub Gnaphalio).—N. Zel.
collinum, *DC. Prod.* vi. 190.—Austral.
collinum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 62 = H. Stoechas.
commixtum, C. Koch, in Linnaea, xxiv. (1851) 351 = lavandulaefolium !
commutatum, Sch. Bip. in Flora, xxvii. (1844) 677 = anomalum.
comosum, Sch. Bip. l. c. 678 = ericaefolium.
compactum, *Boiss. Diagn.* Ser. I. iv. 13.—As. Min.
concolorum, DC. Prod. vi. 206 = rosum.
concretum, *Baker, in Journ. Bot.* xx. (1882) 170.—Madag.
congestum, D. Don, in Mem. Wern. Soc. v. (1824) 550 = felinum.
congestum, *Mœnch, Meth. Suppl.* 326.—Hab. ?
conglobatum, *Stend. Nom.* ed. II. i. 736 ; *Boiss. Diagn.* Ser. I. xi. 81.—Reg. Mediterr.
conglomeratum, Mœnch, Meth. 576 = Gnaphalium luteo-album.
conicum, Sweet, Hort. Brit. ed. I. 228 = Gnaphalium conicum.
Cooperi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 281.—Afr. austr.
cordatum, *DC. Prod.* vi. 180.—Austral.
cordifolium, *DC. l. c.* 208.—Madag.
coriaceum, Harv. in Harv. & Sond. Fl. Cap. iii. 280 = rotundatum.
coriaceum, *Sond. in Harv. & Sond. l. c.* 289.—Afr. austr.
corymbosum, Pers. Syn. ii. 416 = fulgidum.
Cotula, Benth. in Enum. Pl. Hueg. 65 = Helipterum Cotula.
crassifolium, *D. Don, in Mem. Wern. Soc.* v. (1824) 550 ; *Less. Syn. Comp.* 282.—Afr. austr.
crassinervum, DC. Prod. vi. 199 = Krebsianum.
crispum, *D. Don, in Lond. Hort. Brit.* 341 ; *Less. Syn. Comp.* 310.—Afr. austr.
cryptomerioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 186.—Madag.
cuneifolium, *F. Muell. ex Benth. Fl. Austral.* iii. 632.—Austral.
Cunninghamii, *Benth. l. c.* 629.—Austral.
cylindricum, *D. Don. ex G. Don, in Lond. Hort. Brit.* 341 ; *Less. Syn. Comp.* 281.—Afr. austr.
cymosum, *D. Don, in Mem. Wern. Soc.* v. (1824) 550 ; *Less. Syn. Comp.* 309.—Afr. austr.
dasyanthum, Sweet, Hort. Brit. ed. I. 228 = maritimum.
Dasenportii, F. Muell. Fragm. iii. 32 = H. Lawrencella.
declinatum, *Labill. Nov. Holl. Pl.* ii. 45. t. 190.—Tasman.
declinatum, *Less. Syn. Comp.* 278.—Afr. austr.
decorum, *DC. Prod.* vi. 188.—Afr. austr.
decorum, Knaus, ex Harv. & Sond. Fl. Cap. iii. 282 = fulgidum.
decumbens, *Cambess. in Mém. Mus. Par.* xiv. (1827) 271.—Ins. Balear.
decumbens, Portensch. ex Walp. Rep. vi. 288 = Waltaia corymbosa.
decurrens, Mœnch, Meth. 576 = Gnaphalium undulatum.
decurrens, *F. Muell. Fragm.* viii. 46.—Austral.

HELICHRYSUM:—
deterrum, Stend. Nom. ed. II. i. 738 = leptolepis.
dichoon, F. Muell. ex Sond. in Linnaea, xxv. (1852) 512 = Blandowskianum.
diffusum, A. Cunn. ex DC. Prod. vi. 216 = Helipterum polygalifolium.
diffusum, *DC. l. c.* 175.—Afr. austr.
diosmaefolium, *Sweet, Hort. Brit.* ed. I. 228.—Austral.
dioscmifolium, Less. ex Steud. Nom. ed. II. i. 738 = prasc.
dioicoides, *DC. Prod.* vi. 208.—Madag.
diotophyllum, *F. Muell. Fragm.* v. 180.—Austral.
discolor, DC. Prod. vi. 197 = appendiculatum.
discolor, Steud. Nom. ed. II. i. 738 = seq.
discolorum, D. Don, ex Sweet, Hort. Brit. ed. I. 222 = felinum.
divaricatum, D. Don, ex G. Don, in Loud. Hort. Brit. 341 = crispum.
divergens, Less. in Linnaea, vi. (1831) 228 = Metalasia divergens.
diversifolium, Sch. Bip. in Flora, xxvii. (1844) 679 = squamosum.
Dockrelli, *Benth. Fl. Austral.* iii. 626.—Austral.
Dregeanum, *Harv. & Sond. Fl. Cap.* iii. 251.—Afr. austr.
dubium, Cass. in Dict. Sc. Nat. xx. 458 = Leptorhynchus squamatus.
Ecklonis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 254.—Afr. austr.
Eichwaldi, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 121. t. 12. f. 3 = arenarium.
elatum, *A. Cunn. in DC. Prod.* vi. 198.—Austral.
elegans, D. Don, Prod. Fl. Nep. 176 = Anaphalis cuneifolia !
elegans, Presl, Fl. Sic. p. xxix = Gnaphalium pulchrum.
elegantissimum, *DC. Prod.* vi. 179.—Afr. austr.
elongatum, Mœnch, Meth. 575 = orientale.
emirnense, *DC. Prod.* vi. 207.—Madag.
ericaefolium, *Less. Syn. Comp.* 314.—Afr. austr.
ericoides, Boj. ex DC. Prod. vi. 219 = Stenocline ericoides.
ericoides, D. Don, ex G. Don, in Lond. Hort. Brit. 341 = ericaefolium.
ericoides, *Pers. Syn.* ii. 415.—Afr. austr.
Ernestianum, *DC. Prod.* vi. 178.—Afr. austr.
erosum, Harv. in Harv. & Sond. Fl. Cap. iii. 251 = rosum.
erosum, Schlecht. in Linnaea, xx. (1847) 695 = rutidolepis.
Errerae, Tineo, Pl. Rar. Sicil. 27 = saxatile.
Erythropogon, Spreng. Fl. Hal. Mant. 50 = Metalasia unifora.
excisum, *Less. Syn. Comp.* 283.—Afr. austr.
eximium, Less. l. c. 319 = Helipterum eximium.
expansum, *Less. l. c.* 275.—Afr. austr.
fascicularum, Willd. Sp. Pl. iii. 1909 ; Schrank, in Denkschr. Akad. Muench. viii. 1821-22 (1824) 156 = sesamoides.
fastigiatum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 219. — n. sanct.
felinum, *Less. Syn. Comp.* 287.—Afr. austr.
ferrugineum, Less. Syn. Comp. 317 = Helipterum citrinum.
ferrugineum, Pers. Syn. ii. 414 = squamosum.
filagineum, *DC. Prod.* vi. 170.—Afr. austr.
filicaule, Hook. f. Fl. N. Zel. 140. t. 36 B = Gnaphalium filicaule.
filifolium, *F. Muell. Fragm.* iii. 134.—Austral.
filiforme, Less. Syn. Comp. 323 = Helipterum filiforme.
flaccidum, D. Don, ex G. Don, in Lond. Hort. Brit. 341 = Achyrocline flaccida.
flagellare, *Baker, in Journ. Linn. Soc.* xx. (1883) 183.—Madag.
flavissimum, DC. Prod. vi. 195 = spiculatum.
flexicaulum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 62 = H. Stoechas.
floribundum, A. Cunn. ex DC. Prod. vi. 217 = Helipterum floribundum.
foetidum, *Mœnch, Meth.* 575.—Afr. trop. et austr.
folliculatum, DC. Prod. vi. 197 = appendiculatum.
Fontanesii, *Cambess. in Mem. Mus. Par.* xiv. (1827) 270. t. 8.—Sicil.
formosissimum, Sch. Bip. in Bot. Zeit. iii. (1845) 173 = elegantissimum.
formosum, Sch. Bip. ex Klatt, in Ann. Sc. Nat. Sér. V. xviii. (1873) 272 = prasc.

HELICHRYSUM:—
fragrans, *Andr. ex Steud. Nom.* ed. II. i. 739.—Afr. austr.
frigidum, *Willd. Sp. Pl.* iii. 1908.—Ins. Corsica.
fruticans, *D. Don, in Mem. Wern. Soc.* v. (1824) 550.—Afr. austr.
fruticosum, Vatke, in Linnaea, xxxix. (1876) 491 = cymosum.
fulgidum, *Willd. Sp. Pl.* iii. 1904.—Afr. austr.
fulvellum, Harv. in Harv. & Sond. Fl. Cap. iii. 268 = rotundatum.
fulvescens, *DC. Prod.* vi. 207.—Madag.
galactosphaerum, Sch. Bip. in Flora, xxvii. (1844) 681 = rugulosum.
garieplnum, *DC. Prod.* vi. 174.—Afr. austr.
gerberaefolium, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 36 ; *ex Rich. Tent. Fl. Abyss.* i. 435.—Abyssin.
Gerrardi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 244.—Afr. austr.
Gilsoli, *F. Muell. Fragm.* x. 85.—Austral.
glabratum, DC. Prod. vi. 189 = bracteatum.
glanduliferum, Sch. Bip. ex Boiss. Fl. Orient. iii. 286 = arenarium.
glandulíferum, *Sch. Bip. in Bot. Zeit.* iii. (1845) 173. —Kurdistan.
glandulosum, Labill. Fl. Ross. ii. 605 = arenarium.
globiferum, *Boiss. Diagn.* Ser. I. xi. 80.—Persia.
globosum, *Sch. Bip. in Bot. Zeit.* iii. (1845) 174 ; *A. Rich. Tent. Fl. Abyss.* i. 425.—Abyssin.
glumaceum, DC. Prod. vi. 197 = Achyrocline glumacea.
glutinosum, *Benth. Fl. Austral.* iii. 621.—Austral.
glutinosum, A. Br. in Flora, xxiv. (1841) 1. 277 = foetidum.
glutinosum, Tenore, ex Nym. Consp. 381 = rupestre.
gnaphaloides, H. B. & K. Nov. Gen. et Sp. iv. 87 = Gnaphalium antennarioides.
gnaphaloides, Steud. Nom. ed. II. i. 739 = Helipterum gnaphaloides.
gossypinum, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 312.—Ins. Canar.
gracile, DC. Prod. vi. 181 = ramosum.
gracilipes, *Olivier & Hiern, in Oliver, Fl. Trop. Afr.* iii. 348.—Afr. trop.
grandiceps, *Hook. f. Handb. N. Zeal. Fl.* 781, *Index* (154 sub Gnaphalio).—N. Zel.
grandiflorum, *D. Don, in Mem. Wern. Soc.* v. (1824) 550.—Afr. austr.
grandiflorum, Eckl. ex DC. Prod. vi. 212 = Helipterum speciosissimum.
graveolens, Boiss. Fl. Orient. iii. 282 = lanatum.
graveolens, *Sweet, Hort. Brit.* ed. I. 228 ; *DC. Prod.* vi. 184.—Tauria.
Griffithii, *Boiss. Diagn.* Ser. II. vi. 104.—Afghan.
griseum, *Sond. in Linnaea*, xxiii. (1850) 65.—Afr. austr.
Gunnianum, Hook. Ic. Pl. t. 820 = scorpioides.
Gunnii, *F. Muell. ex Benth. Fl. Austral.* iii. 630.—Austral.
gymocomum, *DC. Prod.* vi. 209.—Afr. austr.
hamulosum, *DC. l. c.* 192.—Afr. austr.
hebelepis, *DC. l. c.* 186.—Afr. austr.
Heldreichii, *Reiss. Fl. Orient.* iii. 290.—Ins. Creta.
helianthemifolium, D. Don, ex G. Don, in Loud. Hort. Brit. 849 = serpyllifolium.
heliotropifolium, *DC. Prod.* vi. 209.—Ins. Masear.
hemisphaericum, *DC. l. c.* 181.—Afr. austr.
herbaceum, Sweet, Hort. Brit. ed. I. 228 = squamosum.
hermiaefolium, *DC. Prod.* vi. 170.—Afr. austr.
heterophyllum, Steud. Nom. ed. II. i. 739 = sesamoides.
hispanicum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 67 = serotinum.
Hochstetteri, Hook. f. in Journ. Linn. Soc. vi. (1862) 18 = Achyrocline Hochstetteri.
Hookerianum, Wight & Arn. ex DC. Prod. vi. 201 = luteiflorum.
horridum, *Sch. Bip. in Bot. Zeit.* iii. (1845) 173.—Abyssin.
Humboldtianum, Gaudich. in Freyc. Voy. Bot. 465. t. 86 = Helipterum Humboldtianum.
humile, *Less. Syn. Comp.* 323.—Afr. austr.
hypoleucum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 253. —Afr. austr.
Ignaeum, *D. Don, in Mem. Wern. Soc.* v. (1824) 550. —Europ. austr.
imbricatum, DC. Prod. vi. 191 = spiculatum.
imbricatum, *Less. Syn. Comp.* 279.—Afr. austr.

HELICHRYSUM:—
imbricatum, Thunb. Fl. Cap. ed Schult. 668 = Heliptérum canescens.
incanum, Hook. Bot. Mag. t. 2881 = Helipterum incanum.
incarnatum, DC. Prod. vi. 191 = rubellum.
intermedium, Less. Syn. Comp. 295 = paniculatum.
intricatum, *DC. Prod.* vi. 204.—Afr. austr.
inuloides, Boj. ex DC. l. c. 219 = Stenocline inuloides.
involucratum, *F. Muell. Fragm.* iii. 185.—Austral.
italicum, G. Don, in Loud. Hort. Brit. 842 = angustifolium.
Kempei, *F. Muell. in Melb. Chemist* (Jan. 1882).—Austral.
Kirkii, *Oliver & Hiern, in Trans. Linn. Soc.* xxix. (1873) 93, t. 61.—Afr. trop.
Kirkii, Vatke, in Oestr. Bot. Zeitschr. xxvii. (1877) 195 = Petersii.
Kotschyi, Boiss. Diagn. Ser. I. vi. 92 = armenium.
Krausii, *Sch. Bip. in Flora*, xxvii. (1844) 679.—Afr. austr.
Krebsianum, *Less. Syn. Comp.* 308.—Afr. austr.
lacteum, *Cass. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 866.—Algeria.
Lamarckii, *Cambess. in Mém. Mus. Par.* xiv. (1827) 269, t. 7.—Ins. Balear.
Lambertianum, *DC. Prod.* vi. 190.—Afr. austr.
lamprospermum, Sch. Bip. in Flora, xxvii. (1844) 681 = roseum.
lanatum, *DC. Prod.* vii. 298.—As. Min.
lanatum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 283.—Afr. austr.
lanatum, Schrank, in Denkschr. Akad. Muench. viii. 1821-22 (1824) 166 = vestitum.
lancifolium, *Willd. Sp. Pl.* iii. 1910.—Afr. austr.
lanuginosum, A. Cunn. ex DC. Prod. vi. 198 = elatum.
lanuginosum, Dulac, Fl. Hautes-Pyr. 514 = H. Stoechas.
lasiocaulon, Sweet, Hort. Brit. ed. I. 223 = Gnaphalium lasiocaulon.
latifolium, *Less. Syn. Comp.* 297.—Afr. austr.
lavandulifolium, *D. Don, in Mem. Wern. Soc.* v. (1824) 550; *DC. Prod.* vi. 184.—Armen.
lavandulaefolium, H. B. & K. Nov. Gen. et Sp. iv. 86 = Chiconlaena lavandulacea.
lavanduloides, *DC. Prod.* vi. 207.—Madag.
Lawrencella, *F. Muell. ex Benth. Fl. Austral.* iii. 616.—Austral.
laxum, E. Mey. ex DC. Prod. vi. 171 = ericaefolium.
ledifolium, *Benth. Fl. Austral.* iii. 630.—Austral.
leiolepis, DC. Prod. vi. 187 = revolutum.
leiopodium, *DC. l. c.* 200.—Afr. austr.
leontonychium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = seq.
Leontonyx, *DC. Prod.* vi. 169.—Afr. austr.
Leontopodium, *Hook. f. Fl. N. Zel.* 141. t. 37 B.—N. Zel.
lepidophyllum, *F. Muell. ex Benth. Fl. Austral.* iii. 633.—Austral.
leptolepidium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = seq.
leptolepis, *DC. l. c.* 170. n. 9.—Madag.
leptolepis, DC. Prod. vi. 194. n. 136 = Ixiolaena leptolepis.
leptophyllum, DC. in Wight, Contrib. 20 = Anaphalis leptophylla.
leptophyllum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 68 = serotinum.
leptorhizum, *DC. Prod.* vi. 189.—Afr. austr.
leucocephalum, *Ausfeld, in Regel, Gartenfl.* (1864) 97.—Austral.
leucocephalum, *Boiss. Diagn.* Ser. I. xi. 80.—Persia.
leucocephalum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Moscou, xii. (1860) 121 = Buhseanum.
leucophyllum, DC. Prod. vi. 175 = crassifolium.
leucopadeum, *DC. l. c.* 198.—Austral.
leucophaeum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 417.—Madag.
Libanotis, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 66 = serotinum.
linearc, DC. Prod. vi. 172, partim = ericaefolium.
linearifolium, *Sch. Bip. in Flora*, xxvii. (1844) 678.—Afr. austr.
linifolium, A. Cunn. ex DC. Prod. vi. 216 = Heliptérum anthemoides.
litoreum, *Guss. Fl. Sic. Syn.* ii. 468.—Italia.
longibracteolatum, Schrank, in Denkschr. Akad. Muench. viii. 1821-22 (1824) 168 = humile.

HELICHRYSUM:—
longifolium, *DC. Prod.* vi. 198.—Afr. austr.
Lorentii, Hochst. in Lorent, Wanderungen, 329 = armenium.
lucidum, Henck. Adumb. 5; Hornem. Hort. Hafn. ii. 804 = bracteatum.
lucilioides, *Less. Syn. Comp.* 290.—Afr. austr.
luteo-album, Reichb. in Moesal. Handb. ed. II. ii. 1460 = Gnaphalium luteo-album.
lutescens, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 66 = H. Stoechas.
luzulaefolium, *DC. Prod.* vi. 201.—Madag.
lycopodioides, *Benth. Fl. Austral.* iii. 634.—Austral.
MacIverii, *F. Muell. in Wing, South Sc. Record*, iii. (1883) 99.—Austral.
macranthum, Benth. in Enum. Pl. Hueg. 65 = bracteatum.
macrocephalum, A. Cunn. ex DC. Prod. vi. 188 = bracteatum.
madagascariense, *DC. l. c.* 206.—Madag.
Mangleni, Baill. Hist. Pl. viii. 314 = Heliptérum Mangleni.
Mannii, Hook. f. in Journ. Linn. Soc. vi. (1862) 12 = foetidum.
margaritaceum, Moench, Meth. 576 = Antennaria margaritacea.
marginatum, *DC. Prod.* vi. 180.—Afr. austr.
martiflorum, *DC. l. c.* 186.—Afr. austr.
maritimum, *D. Don, ex G. Don, in Loud. Hort. Brit.* 841; *Less. Syn. Comp.* 304.—Afr. austr.
maritimum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 62 = H. Stoechas.
melaleucum, Reichb. in Flora, xiii. (1830) 382 = melanophthalmum.
melanacme, *DC. Prod.* vi. 203.—Afr. austr.
melanophthalmum, *DC. l. c.* 177.—Ins. Madera.
metalasioides, DC. l. c. 171 = ericaefolium.
Meyeri, DC. l. c. 244, in syn. sphalm. = Zeyheri.
microcephalum, *DC. l. c.* 200.—Afr. austr.
micranthum, *A. Cunn. ex DC. l. c.* 189.—N. Zel.
microcephalum, *DC. l. c.* 207.—Madag.
microlepis, DC. l. c. 195 = scmipapposum.
microphyllum, Cambess. in Mém. Mus. Par. xiv. (1827) 272 = angustifolium.
microspides, *DC. Prod.* vi. 170.—Afr. austr.
Milliganii, *Hook. f. Fl. Tasm.* i. 214. t. 60.—Tasmania.
molle, A. Cunn. ex DC. Prod. vi. 194 = Heliptérum incanum.
Monizii, *Lowe, Man. Fl. Mad.* i. 481.—Ins. Madera.
monspelense, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 64 = H. Stoechas.
monstrosum, DC. Prod. vi. 186 = splendidum.
montevidense, Spreng. Syst. iii. 485 = Lucilia acutifolia.
mucronatum, *Less. Syn. Comp.* 295.—Afr. austr.
multiceps, DC. Prod. vi. 173 = crassifolium.
multinerve, DC. l. c. 199 = quinquenerve.
Mundtii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 248.—Afr. austr.
mutisiaefolium, *Less. in Linnaea*, vi. (1831) 224.—Madag.
natalitium, *DC. Prod.* vi. 201.—Afr. austr.
nebrodense, Heldr. in Flora, xxvii. (1844) 67 = rupestre.
nepalense, Spreng. Syst. iii. 485 = Anaphalis triplinervis.
Newii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 349.—Afr. trop.
Nimmoanum, *Oliver & Hiern, l. c.* 347.—Afr. trop.
nitens, *Oliver & Hiern, l. c.* 350.—Afr. trop.
nivcum, *Boiss. & Heldr. Diagn.* Ser. I. xi. 80.—As. Min.
niveum, R. Grah. in Bot. Mag. t. 3857 = bracteatum.
niveum, Less. Syn. Comp. 302 = perviflorum.
nodiflorum, *DC. Prod.* vi. 177.—Lusitan.
Noëanum, *Boiss. Diagn.* Ser. II. vi. 104.—Anatolia.
notatum, Sieber, Herb. Nom. ed. II. i. 395, in syn.=expansum.
Notonianum, DC. in Wight, Contrib. 20 = Anaphalis Notoniana.
nudifolium, *Less. Syn. Comp.* 299.—Afr. austr.
numidicum, *Pomel, Nouv. Mat. Fl. Alt.* 386.—Alger.
obconicum, *DC. Prod.* vi. 181.—Ins. Madera.
obcordatum, *F. Muell. ex Benth. Fl. Austral.* iii. 632.—Austral.
obovatum, *DC. Prod.* vi. 180.—Austral.
obtusifolium, *F. Muell. & Sond. in Linnaea*, xxv. (1852) 513.—Austral.

HELICHRYSUM:—
obvallatum, *DC. Prod.* vi. 174.—Afr. austr.
odoratissimum, *Sweet, Hort. Brit.* ed. I. 223; *Less. Syn. Comp.* 301.—Afr. austr.
odorum, DC. Prod. vi. 199 = spiculatum.
Oldfieldii, F. Muell. *Fragm.* iii. 134 = Heliptérum Cotula.
oligocephalum, DC. Prod. vi. 184 = armenium.
oligocephalum, *F. Muell. Fragm.* vi. 235.—Austral.
olonnense, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 65 = H. Stoechas.
oocephalum, *Boiss. Fl. Orient.* iii. 286.—Persia.
orientale, *Gaertn. Fruct.* ii. 404. t. 166.—Archip.; As. Min.
oxybellum, *DC. Prod.* vi. 171.—Afr. austr.
oxylepis, *F. Muell. Fragm.* i. 35.—Austral.
oxyphyllum, *DC. Prod.* vi. 199.—Afr. austr.
pachyrhizum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 222.—Afr. austr. et trop.
Pallasii, Ledeb. Fl. Ross. ii. 606 = callichrysum.
pallidum, DC. Prod. vi. 199 = undatum.
panaetioides, DC. l. c. 194 = Leptorhynchos panaetioides.
pandurae folium, Schrank, in Denkschr. Akad. Muench. viii. 1821-22 (1824) 169 = auriculatum.
paniculatum, *Willd. Sp. Pl.* iii. 1911.—Afr. austr.
pannosum, *DC. Prod.* vi. 204.—Afr. austr.
pannosissimum, Heldr. in Boiss. Diagn. Ser. I. xi. 81 = Heldreichii.
panormitanum, Tineo, in Guss. Fl. Sic. Syn. ii. 467 = rupestre.
papillosum, Labill. Nov. Holl. Pl. ii. 46. t. 192 = bracteatum.
parychioides, *DC. Prod.* vi. 171.—Afr. austr.
perviflorum, *DC. l. c.* 208.—Afr. austr.
parvulum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 65 = H. Stoechas.
patulum, *Baker, in Journ. Linn. Soc.* xx. (1883) 185.—Madag.
patulum, *D. Don, in Mem. Wern. Soc.* v. (1824) 550.—Afr. austr.
pedunculare, *DC. Prod.* vi. 198.—Afr. austr.
pendulum, Presl, Fl. Sic. p. xxix = rupestre.
pentzioides, *Less. Syn. Comp.* 283.—Afr. austr.
Petersii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 349.—Afr. trop.
petiolatum, *D. Don, in Mem. Wern. Soc.* v. (1824) 550.—Afr. austr.
phlomoides, Spreng. Syst. iii. 482 = Heliptérum phlomoides.
pholidotum, *F. Muell. ex Benth. Fl. Austral.* iii. 634.—Austral.
phylicaefolium, *DC. Prod.* vi. 207.—Madag.
Picardi, Boiss. & Reut. Diagn. Ser. II. vi. 103 = serotinum.
Pichleri, *W. Barbey, in Bull. Soc. Vaud. Sc. Nat.* xxi (1885) 96.—Ins. Sardoa.
pilosellum, Less. Syn. Comp. 297 = pedunculare.
Pinsapi, Schrank, in Denkschr. Akad. Muench. viii. 1821-22 (1824) 168 = humile.
plantagineum, St. Lag. in Ann. Soc. Bot. Lyon, vii (1880) 127 = seq.
Plantago, *DC. Prod.* vi. 200.—Madag.
platypterum, *DC. l. c.* 201.—Afr. austr.
plebeium, *DC. l. c.* 206.—Afr. austr.
plicatum, *DC. l. c.* 183.—As. Min.; Persia.
pluriceps, C. Koch, in Linnaea, xxiv. (1851) 848 = arenarium.
podolepideum, *F. Muell. Rep. Babb. Exp.* 18.—Austral.
polycephalum, Sch. Bip. in Flora, xxvii. (1844) 681 = capitellatum.
polygalaefolium, A. Cunn. ex DC. Prod. vi. 216 = Heliptérum polygalaefolium.
polyphyllum, Ledeb. Fl. Ross. ii. 605 = plicatum.
popolifolium, *DC. Prod.* vi. 190.—Afr. austr.
Porcari, *Tin. ex Lojac. in Natural. Sicil.* ii. (1883) 182.—Sicil.
proliferum, Willd. Sp. Pl. iii. 1905 = Phaenocoma prolifera.
prostratum, *Hook. f. Fl. Antarct.* 30. t. 21.—N. Zel.
proteiforme, *Baker, Fl. Maurit.* 167.—Ins. Mascar.
Pseudo-fasciculatum, *Schrank, in Denkschr. Akad Muench.* viii. 1821-22 (1824) 167.—Hab.?
psiliolepis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 225.—Afr. austr.
psychrophilum, Boiss. Diagn. Ser. I. xi. 38 = callichrysum.

HELICHRYSUM:—
pterochaetum, F. Muell. Rep. Babb. Exp. 14 = Helipterum pterochaetum.
pulchellum, *DC. Prod.* vi. 190.—Afr. austr.
pumilum, *Hook. f. Fl. Tasm.* i. 218. t. 60.—Tasmania.
Quartinianum, *A. Rich. Tent. Fl. Abyss.* i. 427.—Abyss.
quinquenerve, *Less. Syn. Comp.* 300.—Afr. austr.
radicans, Willd. Sp. Pl. iii. 1907 = retortum.
ramosissimum, Hook. in Mitch. Journ. Trop. Austral. 83 = spinulatum.
ramosum, *DC. Prod.* vi. 181.—Austral.
ramulosum, *DC. l. c.* 176.—Afr. austr.
recurvatum, *Willd. Sp. Pl.* iii. 1908.—Afr. austr.
rependum, DC. Prod. vi. 208 = scabrum.
reticulatum, *Less. ex Steud. Nom.* ed. II. i. 740.—Tasmania.
retortum, *Willd. Sp. Pl.* iii. 1907.—Afr. austr.
retrorsum, *DC. Prod.* vi. 207.—Madag.
retusum, *F. Muell. Fragm.* viii. 46.—Austral.
retusum, Sprang. Syst. iii. 494 = Facelis spicalata.
revolutum, *Less. Syn. Comp.* 305.—Afr. austr.
Rhinocerotis, Herb. Berol. ex Steud. Nom. ed. II. i. 552, in syn. = Elytropappus Rhinocerotis.
rigens, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 63 = H. Stoechas.
rigidulum, *DC. Prod.* vi. 198.—Austral.
rigidum, Ait. Hort. *Kew.* ed. II. v. 34 = striatum.
Rochetii, *Delile, in Rochet, Sec. Voy. Choa*, 342.—Afr. trop.
rosmum, Baill. Hist. Pl. viii. 314 = Helipterum roseum.
rosmarinifolium DC. Prod. vi. 166, in obs. = diosmifolium.
roxulatum, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 347.—Afr. trop.
rosum, *Less. Syn. Comp.* 306.—Afr. austr.
rotundatum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 280. —Afr. austr.
rotundifolium, *Less. Syn. Comp.* 277.—Afr. austr.
rubellum, *Less. l. c.* 280.—Afr. austr.
rugulosum, *Less. l. c.* 307.—Afr. austr.
rupestre, Boiss. Voy. Bot. Espagne, 325 = Boissieri.
rupestre, *DC Prod.* vi. 189.—Reg. Mediterr.
rupestre, Gaus. ex Nym. Consp. 331 = saxatile.
rupicolum, *DC. Prod.* vi. 190.—Austral.
rupicolum, *Pomel, Nouv. Mat. Fl. Atl.* 47.—Algeria.
rutidolepis, *DC. Prod.* vi. 194.—Austral.
rutilans, *D. Don, in Mem. Wern. Soc.* v. (1824) 550 ; *Less. Syn. Comp.* 276.—Afr. austr.
sabulosum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 64 = H. Stoechas.
salicifolium, *Bertol. Fl. Guatimal.* 33.—Guatemala.
sanguineum, *Kostel. Alig. Med.-Pharm. Fl.* ii. 709.—Syria.
saturejaefolium, Moench, Meth. Suppl. 336 = Metalasia muricata !
saxatile, *Moris, Fl. Sard.* 387. t. 62.—Ins. Sard. ; Sicil.
scabrum, DC. Prod. vi. 208 = Dregeanum.
scabrum, *Less. Syn. Comp.* 315.—Afr. austr.
scandens, Gaus. Fl. Sic. Syn. ii. 466 = siculum.
scorpioides, *Labill. Nov. Holl. Pl.* ii. 45. t. 191.—Austral.
scutellifolium, *Benth. Fl. Australi.* iii. 633.—Austral.
selaginoides, *F. Muell. ex Benth. l. c.* 634.—Austral.
sermaclavum, F. Muell. Fragm. ii. 156 = Leptorhynchos ambiguus.
semiferlle, *F. Muell. Rep. Babb. Exped.* 14.—Austral.
semipapposum, *DC. Prod.* vi. 195.—Austral.
serotinum, *Botan. Voy. Bot. Espagne*, ii. 229.—Europ.
serpyllifolium, *Less. Syn. Comp.* 277.—Afr. austr.
serpyllifolium, Pers. Syn. ii. 414 = Helipterum canescens.
sessamoides, *Willd. Sp. Pl.* iii. 1908.—Afr. austr.
sessile, *DC. Prod.* vi. 178.—Afr. austr.
setosum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 281.—Afr. austr.
siculum, Boiss. Fl. Orient. iii. 229 = conglobatum.
siculum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 67 = serotinum.
similllimum, *DC. Prod.* vi. 208.—Afr. austr.
simulans, *Harv. & Sond. Fl. Cap.* iii. 217.—Afr. austr.
Sinclairii, *Hook. f. Handb. N. Zeal. Fl.* 131, *Index* (153 sub Gnaphalio).—N. Zel.
Sonderi, F. Muell. in Seem. Journ. Bot. iv. (1866) 121 = Ixiolaena tomentosa.
sordescens, *DC. Prod.* vi. 178.—Afr. austr.

HELICHRYSUM:—
spathulatum, Moench, Meth. Suppl. 335 = Gnaphalium indicum.
speciosissimum, Willd. Sp. Pl. iii. 1904 = Helipterum speciosissimum.
speciosum, Thunb. Fl. Cap. ed. Schult. 664 = vestitum, Helipterum variegatum.
spectabile, *G. Don, ex Loud. Hort. Brit. Suppl.* iii. 861 (= sect. 9).—Austral.
spectabile, Lodd. Bot. Cab. t. 59 = humile.
sphaerocephalum, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1883) 839.—Ins. Socotr.
Spicori, *F. Muell. Fragm.* xi. 47.—Austral.
spiciforme, *DC. Prod.* vi. 171.—Afr. austr.
spinosum, Sch. Bip. in Bot. Zeit. iii. (1845) 178 = citrispinum.
spinosum, Willd. Sp. Pl. iii. 1908 = Dicoma diacanthoides.
spirale, Willd. l. c. 1903 = Helipterum variegatum.
splendens, Sims, Bot. Mag. t. 1773 = squamosum.
splendidum, *Less. Syn. Comp.* 286.—Afr. austr.
squamosum, *Thunb. Fl. Cap. ed. Schult.* 661.—Afr. austr.
squarrosum, *Baker, in Journ. Linn. Soc.* xx. (1883) 184.—Madag.
squarrulosum, DC. Prod. vi. 195 = semipapposum.
Staechas, DC. Fl. Fr. iv. 132 = H. Stoechas.
Staehelina, Thunb. Fl. Cap. ex Schult. 664 = Helipterum speciosissimum.
Staehelina, Willd. Sp. Pl. iii. 1910 = argenteum.
staehelinoides, Less. Syn. Comp. 321 = Helipterum gnaphaloides.
stellatum, *Less. l. c.* 279.—Afr. austr.
stenopterum, *DC. Prod.* vi. 201.—Afr. austr.
Stendelii, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 421 = Gnaphalium Steudelii.
stipitatum, F. Muell. Fragm. iii. 133 = Helipterum stipitatum.
stoechadaeae, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 197 = seq.
Stoechas, *DC. Fl. Fr.* iv. 132.—Reg. Mediterr. occ.
stoechadinum, D. Don, Prod. Fl. Nep. 176 = Anaphalis contorta.
stoloniferum, *Willd. Sp. Pl.* iii. 1907.—Afr. austr.
stramineum, Gaus. Fl. Sic. Syn. ii. 467 = saxatile.
striatum, *Willd. Sp. Pl.* iii. 1905.—Afr. austr.
strictum, Moench, Meth. 676 = Gnaphalium sylvaticum.
subdecurrens, *DC. Prod.* vi. 202.—Afr. austr.
subglomeratum, *Less. Syn. Comp.* 283.—Afr. austr.
subsilifolium, *Harv. in Harv. & Sond. Fl. Cap.* iii. 241. —Afr. austr.
subsilifolium, *F. Muell. Fragm.* iii. 134.—Austral.
suffruticosum, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1883) 840.—Ins. Socotr.
Sutherlandi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 218. —Afr. austr.
synciadum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 62 = H. Stoechas.
taxacertifrons, *Baker, in Journ. Linn. Soc.* xx. (1883) 188.—Madag.
Tardentii, Boiss. Diagn. Ser. I. xi. 29 = armenium.
tchabanicum, *C. Koch, in Linnaea*, xxiv. (1851) 849.—Armenia.
tenuiculum, *DC. Prod.* vi. 208.—Afr. austr.
tephrodes, Sweet, Hort. Brit. ed. I. 232 = teretifolium.
Tepperi, *F. Muell., in Wing, South St. Record*, ii. (1882) 1.—Austral.
teretifolium, *Sweet, Hort. Brit.* ed. I. 232 ; *Less. Syn. Comp.* 315.—Afr. austr.
thianschanicum, *Regel, in Act. Hort. Petrop.* vi. (1879) 307.—Turkestan.
Thomsoni, *F. Muell. Fragm.* viii. 45.—Austral.
tricostatum, *Less. Syn. Comp.* 310.—Afr. austr.
trilloseatum, *DC. Prod.* vi. 192.—Afr. austr.
trinervatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 182. —Madag.
triplinerve, *DC. Prod.* vi. 207.—Madag.
tschabanicum, C. Koch, in Linnaea, xxiv. (1851) 849 = tchabanicum.
uliginosum, Moench, Meth. 676 = Gnaphalium uliginosum.
umbellatum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 220. —Afr. austr.
umbracullgerum, *Less. Syn. Comp.* 284.—Afr. austr.
undatum, *Less. l. c.* 306.—Afr. austr.
undulatum, Labill. Fl. Rom. ii. 606 = armenium.
variegatum, Thunb. Fl. Cap. ed. Schult. 660 = Helipterum variegatum.

HELICHRYSUM:—
variegatum, Willd. Sp. Pl. iii. 1911 = canescens.
vermiculatum, Pers. Syn. ii. 415 = Metalasia uniflora.
vestitum, *Schrank, in Denkschr. Akad. Muench.* viii. 1821-23 (1824) 166 ; *Less. Syn. Comp.* 314.—Afr. austr.
vestitum, Willd. Sp. Pl. iii. 1908 = Helipterum variegatum.
virgatum, Willd. l. c. 1910 = Helipterum virgatum.
virgineum, DC. Prod. vi. 177 = Billardieri.
virgineum, *Griseb. Spicil. Fl. Rumel.* ii. 197.—Graecia.
viscosum, Sieber, ex DC. Prod. vi. 189 = bracteatum.
Wickstromii, *Tineo, ex Lojac. in Natural. Sicil.* ii. (1883) 185.—Sicil.
Wightii, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 291.—Ind. or.
xanthinum, DC. Prod. vi. 185 = splendidum.
xeranthemoides, DC. l. c. 178 = lancifolium.
xerochrysum, *DC. l. c.* 201.—Afr. austr.
xylocladum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 418.—Madag.
Youngii, *Hook. f. Handb. N. Zeal. Fl.* 131, *Index* (152 sub Gnaphalio).—N. Zel.
yuccaefolium, *Lam. Illustr.* t. 693. f. 2 ; *ex Poir. Tabl. Encyc.* iii. 271.—Ins. Maurit.
Zeyheri, *Less. Syn. Comp.* 309.—Afr. austr.

HELICIA, Lour. Fl. Cochinch. i. 83 (1790). *PROTEACEAE*, Benth. & Hook. f. iii. 179.
CASTRONIA, Noronhs, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2.
HELIOTOPHYLLUM, Blume, Bijdr. 652 (1825).
amboinensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 204. —Ins. Amboin.
attenuata, *Blume, in Ann. Sc. Nat.* Sér. II. i. (1834) 216.—Malaya.
australasica, *F. Muell. in Hook. Kew Journ.* ix. (1857) 22.—Austral.
Bennettiana, *Miq. Fl. Ind. Bat.* i. 1. 984.—Java.
castaneaefolia, Meisn. in DC. Prod. xiv. 441 = robusta.
ceylanica, *Gardn. in Calc. Journ. Nat. Hist.*, vii. (1847) 458.—Zeylan.
cochinchinensis, *Lour. Fl. Cochinch.* 83.—Cochinch.
conjunctiflora, F. Muell. Fragm. v. 38 = glabriflora.
Cumingiana, *Presl, Epim. Bot.* 246.—Ins. Philipp.
Darlingiana, F. Muell. Fragm. v. 24 = Darlingia spectatissima.
excelsa, *Blume, in Ann. Sc. Nat.* Sér. II. i. (1834) 219.—Ind. or. ; Burma.
ferruginea, *F. Muell. Fragm.* iii. 37.—Austral.
glabrata, *Meisn. in DC. Prod.* xiv. 442.—Burma.
glabriflora, *F. Muell. Fragm.* ii. 91.—Austral.
javanica, Blume, in Ann. Sc. Nat. Sér. II. i. (1834) 217 = robusta.
lancifolia, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 74.—Japon.
loranthoides, *Presl, Epim. Bot.* 247.—Ins. Philipp.
macrophylla, Wall. Cat. n. 5661 = robusta.
maccocana, *Blume, in Ann. Sc. Nat.* Sér. II. i. (1834) 216.—Ins. Moluccc.
nilagirica, *Bedd. in Madr. Journ. Sc. Ser.* III. i. (1864) 96.—Ind. or.
oblongifolia, *Benn. Pl. Jav. Rar.* 83.—Java.
obovata, *Benn. l. c.*—Java.
obtusidens, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 204.—Ins. Amboin.
ovata, *Benn. Pl. Jav. Rar.* 84.—Sumatra.
petiolaris, *Benn. l. c.*—Singapore.
philippinensis, *Meisn. in DC. Prod.* xiv. 441.—Ins. Philipp.
praealta, *F. Muell. Fragm.* iii. 37.—Austral.
pyrrhobotrys, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) pt. 2. 108.—Burma.
robusta, Villar, in Blanco, Fl. Philip. ed. III. Nov. App. 182 = philippinensis.
robusta, *Wall. Cat.* n. 3702.—Reg. Himal. ; Burma ; Malaya.
salicifolia, Presl, Epim. Bot. 247 = excelsa.
Scottiana, F. Muell. Fragm. iv. 107 = Xylomelum scottianum.
serrata, *Blume, in Ann. Sc. Nat.* Sér. II. i. (1834) 215. —Ins. Moluccn.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 364.—Sumatra.
terminalis, *Kurz, For. Fl. Brit. Burma*, ii. 238.—Burma.
torulifolia, F. Muell. Fragm. ii. 91 = Macadamia torulifolia.
verticillata, F. Muell. l. c. vi. 221 = Macadamia verticillata.

HELICIA:—
Youngiana, F. Muell. Fragm. iv. 84 = Macadamia Youngiana.
ceylanica, Gardn. ex Meisn. in DC. Prod. xiv. 436 = ceylanica.

HELICIA, Pers. Syn. i. 214 (1805) = **Loranthus**, Linn.
parasitica, Pers. i. c.—China.

HELICILLA, Moq. in DC. Prod. xiii. ii. 189 (1849). *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 71.
altissima, *Moq. l. c.* 170.—China.

HELICODEA, Lem. Illustr. Hortic. xi. (1864) t. 421 = **Billbergia**, Thunb. (Bromeliac.).
Baraquiana, Lem. l. c. = B. decora.
Leopoldii, Hort. Versch. ex Lem. l. c. sub t. 421 = B. Leopoldii.
zebrina, Hort. ex Nichols. Dict. Gard. i. 128 = B. zebrina.

HELICODICEROS, Schott, in Oestr. Bot. Wochenbl. iii. (1853) 369. *AROIDEÆ*, Benth. & Hook. f. iii. 968.
MEGOTIGEA, Rafin. Fl. Tellur. iii. 64 (1836), nomen prius.
crinitus, *Schott, l. c.*; *et Gen. Aroid.* t. 21.—Ins. Corsica.
muscivorus, Engl. in DC. Monog. Phan. ii. 605 = crinitus.

HELICONIA, Linn. Mant. ii. 147 (1767). *SCITAMINEÆ*, Benth. & Hook. f. iii. 655.
BIHAI, Adans. Fam. ii. 67 (1763).
HELICONIOPSIS, Miq. Fl. Ind. Bat. iii. 590, 1855 [1858].
acuminata, A. Rich. in Nov. Act. Nat. Cur. xv. Suppl. (1831) 26. t. 11 = H. Bihlia.
alba, Linn. f. Suppl. 167 = Strelitzia augusta.
Andreanii, Klotzsch, in Linnæa, xx. (1847) 465 = pittacorum.
angusta, Vell. Fl. Flum. 106; ill. t. 20 = pittacorum.
angustifolia, *Hook. Bot. Mag.* t. 4475.—Bras.
augusta, Salisb. Prod. 144 = Strelitzia augusta.
auranliaca, *Ghiesbr. in Lem. Illustr. Hortic.* lx. (1862) t. 332.—Mexic.
aurea-striata, Hort. ex Bull. Cat. (1881) 18; in Lem. Illustr. Hortic. xxix. (1882) t. 464 = H. Bihai!
austro-caledonica, Vieill. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 47 = H. Bihai.
Bailla, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guian.
Bihai, Vell. Fl. Flum. 106; ill. t. 19 = H. Bihai.
bicolor, Benth. in Mauud, Bot. iii. t. 101 = angustifolia.
bicolor, Klotzsch, in Linnæa, xx. (1847) 465 = hirsuta.
Bihai, *Linn. Mant.* ii. 211.—Am. austr.
Bihai, Linn. f. Suppl. 157 = Strelitzia augusta.
Bihai, J. Mill. Ic. tt. 5, 6 = Strelitzia Reginæ.
brasiliensis, *Hook. Exot. Fl.* t. 190.—Bras.
brevispatha, Hook. Bot. Mag. t. 5416 = aurantiaca.
bucciniata, Roxb. Hort. Beng. 19; Fl. Ind. i. 670 = H. Bihai.
cannoidea, A. Rich. in Nov. Act. Nat. Cur. xv. Suppl. (1831) 24. t. 9 = hirsuta.
caribæa, Lam. Encyc. i. 426 = H. Bihai.
dasyantha, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1854) 12.—Hab.?
densiflora, *Hort. Par. ex Verlot, in Rev. Hortic.* (1869) 374; *Cogn. & Marchal, Pl. Ornem.* ii. (1874) t. 46.—Guian.
discolor, Hort. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 186 = Calathea varians.
episcopalis, *Vell. Fl. Flum.* 107; ill. t. 22.—Bras.
farinosa, *Ruddi, in Mem. Mod.* xviii. *Fis.* (1820) 993. —Bras.
flava, *Yacq. Hort. Schoenb.* i. 25.—Am. trop.
flexuosa, *Presl, Rel. Haenk.* i. 105.—Am. trop.
glauca, *Pait. in Rev. Horticic.* (1869) 113.—Am. austr.
hirsuta, *Linn. f. Suppl.* 156.—Am. austr.
hirsuta, Schlecht. & Cham. in Linnæa, vi. (1831) 57 = Schiedeana.
humilis, *Yacq. Hort. Schœnb.* i. 28.—Am. austr.
indica, Lam. Encyc. i. 426 = H. Bihai.
latispatha, *Benth. Bot. Voy. Sulph.* 170.—Am. trop.
leucogramma, Hort. ex Horan. Prod. Scitam. 40 (Calathea sp. ?).—Hab.?
lingulata, *Ruiz & Pav. Fl. Per.* iii. 71.—Peruv.
lutea-fusca, Jacq. Hort. Schœnb. i. 26 = H. Bihai.

HELICONIA:—
marantifolia, G. Shaw, Chmel. Phys. 74. t. 38 = pittacorum.
Martiæ, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 69. —N. Granat.
meridensis, Klotzsch, in Linnæa, xx. (1847) 462 = latispatha.
metallica, *Planch. & Linden, in Bot. Mag.* t. 5315.—N. Granat.
nigrescens, Jacq. Hort. Schœnb. i. 25 = H. Bihai.
pendula, *Wawra, in Oestr. Bot. Zeitschr.* xiii. (1863) 8.—Bras.
pittacorum, *Linn. f. Suppl.* 158; *Andr. Bot. Rep.* t. 134.—Am. austr.
pittacorum, Sw. Obs. Bot. 98 = hirsuta.
pulverulenta, *Lindl. Bot. Reg.* t. 1648.—Am. austr.
Ravennæ, Willem. in Usteri, Ann. Bot. xviii. (1796) 22 = Ravenala madagascariensis.
Richardiana, Miq. in Linnæa, xviii. (1844) 70 = hirsuta.
rostrata, *Ruiz & Pav. Fl. Per.* iii. 71. t. 305.—Peruv.
Schiedeana, *Klotzsch, in Linnæa*, xx. (1847) 468.—Mexic.
Schomburgiana, Klotzsch, l. c. 464 = pittacorum.
Sevennesi, Van Houtte, Cat. (1875-76) 183 = H. Bihai!
speciosa, Hort. ex Horan. Prod. Scitam. 40 = brasiliensis.
Strelitzia, J. F. Gmel. Syst. 423 = Strelitzia Reginæ.
striata, Hort. Veitch, ex Fl. des Serres, xviii. (1880) t. 2416-17 = H. Bihai!
subulata, Ruiz & Pav. Fl. Per. iii. 70 = pittacorum.
tenuiflora, Rosm. & Schult. Syst. v. 691 = hirsuta.
Thalia, *Vell. Fl. Flum.* 106; ill. t. 21.—Bras.
triumphans, Linden, in Illustr. Hortic. xxix. (1881) t. 448 = H. Bihai!
vaginalis, Benth. Bot. Voy. Sulph. 171 = hirsuta.
variegata, Jacq. Hort. Schœnb. i. 25 = H. Bihai.
velleriger, *Poepp. Reise Chili*, ii. 295.—Peruv.
villosa, *Klotzsch, in Linnæa*, xx. (1847) 463.—N. Granat.
Wawræ, *Klotzsch, in Linnæa*, xx. (1847) 463.—N. Granat.
zebrina, *Pel. Fl. Hort. Alsahov.* 69 (Quid?).—Hab.?

HELICONIOPSIS, Miq. Fl. Ind. Bat. iii. 590, 1855 [1858] = **Heliconia**, Linn. (Scitam.).
amboinensis, Miq. l. c. = Heliconia Bihai.

HELICOPHYLLUM, Schott, Syn. Aroid. i. 22 (1856); Gen. Aroid. 20. t. 28-30 (1856). *AROIDEÆ*, Benth. & Hook. f. iii. 968.
EMMUSA, Schott, Syn. Aroid. 16 (1856).
Alberti, *Regel, in Act. Hort. Petrop.* viii. (1884) 683. —Turkestan.
angustatum, Schott, Syn. Aroid. 22 = crassipes.
Ascherl, Schott, Prod. Aroid. 118 = Rauwolfii.
crassifolium, *Engl. in DC. Monog. Phan.* ii. 597.—Sibir.
crassipes, *Schott, Syn. Aroid.* i. 22.—Palæstin.
Dracunculus, Schott, Prod. Aroid. 115 = crassipes.
Kotschyi, Schott, l. c. 114 = Rauwolfii.
Lehmanni, *Regel, in Act. Hort. Petrop.* vii. (1880) 549.—Turkestan.
Loftusii, Schott, in Oestr. Bot. Wochenbl. (1857) 268 = crassipes.
Luph, Schott, Prod. Aroid. 118 = Rauwolfii *vel* crassipes.
Olivieri, Schott, Syn. Aroid. 22 = Rauwolfii.
Rauwolfii, *Schott, Prod. Aroid.* 113.—Syria; sopotam.; Armen.
Russelianum, Schott, l. c. 118 = Rauwolfii.
spiculatum, Schott, l. c. 118 = crassipes.

HELICOSTYLIS, Trécul, in Ann. Sc. Nat. Sér. III. vii. (1847) 134. t. 5. *URTICACEÆ*, Benth. & Hook. f. iii. 373.
MAQUIRA, Aubl. Pl. Gui. Suppl. 36. t. 389 (1775).
affinis, Miq. in Mart. Fl. Bras. iv. i. 113.—Am. trop.
Poeppigiana, *Trécul, in Ann. Sc. Nat.* Sér. III. vii. (1847) 134.—Guiana.

HELICOTRICHUM, Bess. ex Reichb. Fl. Germ. Excurs. 1409. a. 852 (1832) = **Avena**, Linn. (Gramin.).
planiculme, Bess. ex Reichb. l. c. = A. planiculmis.

HELICTA, Cass. in Bull. Soc. Philom. (1818) 167 = **Borrichia**, Adans. (Compos.).
sarmentosa, Cass. l. c.—Madag.

HELICTA, Less. Syn. Comp. 221 (1832) = **Epallage**, DC. (Compos.).
alternifolia, Less. l. c. = Sphacophyllum madagascariense.
madagascariensis, Less. l. c. = E. dentata.

HELICTERES, Pluk. ex Linn. Syst. ed. I (1735). *STERCULIACEÆ*, Benth. & Hook. f. i. 230.
ALICTERES, Neck. Elem. ii. 804 (1790).
ANISORA, Rafin. Sylva Tellur. 74 (1838).
CAMAION, Rafin. l. c. 75 (1838).
ISORA, Adans. Fam. ii. 389 (1763); Schott, Meletem. 31 (1832).
METRORIUM, Schott, l. c. 29. t. 5 (1832).
NISORALIS, Rafin. Sylva Tellur. 74 (1838).
OPSOPARA, Neck. Elem. ii. 805 (1790).
ORTHOTRICHUM, Schott. Meletem. 31 (1832).
OUDEMANSIA, Miq. Fl. Jungh. i. 295 (1854).
OZOKETA, Rafin. Sylva Tellur. 75 (1838).
althææfolia, Lam. Encyc. iii. 88 = guazumæfolia *vel* jamaicensis.
altheæfolia, Benth. Bot. Voy. Sulph. 70 = barsensis.
angustifolia, *Linn. Sp. Pl.* 963.—Malaya; China.
apetala, Jacq. Enum. Pl. Carib. 30 = Sterculia Helicteres.
arborescens, Mill. Gard. Dict. ed. VIII. n. 3 = hirsuta.
aspera, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 212.—Bras.
barbadensis, *Jacq. Enum. Pl. Carib.* 30.—Ind. occ.
barsensis, Benth. Bot. Voy. Sulph. 70 = guazumæfolia.
baruensis, *Yacq. Enum. Pl. Carib.* 30.—Am. austr.
bractosa, Mart. in Flora, xxii. (1839) 1. Beibl. 15 = pentadon.
brasiliana, *Zucc. in Syll. Ratisb.* ii. (1832) 69.—Bras.
brasiliensis, Mikan, Delect. Fl. et Faun. Bras. fasc. iv. t. 21 = ovata.
brevior, Mill. Gard. Dict. ed. VIII. n. 2 = H. Isora.
brevispira, *A. Juss. in St. Hil. Fl. Bras. Mer.* i. 374. —Bras.
cana, *Benth. Fl. Austral.* i. 232.—Austral.
carpinifolia, Presl, Rel. Haenk. ii. 38 = guazumæfolia.
carthagenensis, Herb. Madr. ex Wall. Cat. n. 2687 = spicata!
carthagenensis, *Jacq. Enum. Pl. Carib.* 30.—Venezuela.
chrysocalyx, Miq. ex Mast. in Hook. f. Fl. Brit. Ind. i. 365 = H. Isora.
cordifolia, *Hoffmgg. Vers. Pfl. Nachtr.* ii. 131.—Bras.
corylifolia, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1825) 44.—Bras.
dentata, *F. Muell. ex Benth. Fl. Austral.* i. 232.—Austral.
dumosa, Salisb. Prod. 388 = jamaicensis.
elongata, *Wall. Cat.* n. 1846; *et ex Boj. Hort. Maurit.* 80.—Reg. Himal.; Burma.
ferruginata, Link, Enum. Hort. Berol. ii. 199 = ovata.
fetida, Rafin. Sylva Tellur. 75 (Alicteres) = carthagenensis.
frondosa, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guian.
furfuracea, *A. Rich. Ess. Fl. Cub.* 171.—Cuba.
Gardneriana, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 212.—Bras.
glabra, Klotzsch, in Schomb. Faun. et Fl. Gui. 1172, nomen.—Guian.
glabriuscula, *Wall. Cat.* n. 1185.—Burma.
grewiæfolia, *DC. Prod.* i. 477.—Ins. Moluce.
grewinæfolia, Wall. Cat. n. 1443 = spicata.
guazumæfolia, *H. B. & K. Nov. Gen. et Sp.* v. 304.—Am. austr.
harmsnii, Vell. Fl. Flum. ix. t. 101 = brevispira.
hirsuta, Buch.-Ham. in Wall. Cat. n. 2686 = spicata.
hirsuta, *Lour. Fl. Cochinch.* ii. 580.—Cochinch.
hispida, *Presl, Rel. Haenk.* ii. 140.—Ins. Philipp.
integerrima, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 302.—Borneo.
involucrata, Moric. Pl. Rar. Am. 100. t. 68 = corylifolia.
Isora, *Linn. Sp. Pl.* 963.—As. et Austral. trop.
Isora, Vell. Fl. Flum. ix. t. 102 = ovata.
jamaicensis, DC. ex Spreng. Syst. iii. 80 = ovata.
jamaicensis, *Yacq. Select. Am.* 235. t. 179. f. 99.—Ind. occ.

HELICTERES:—
jamaicensis, Lam. Encyc. vi. 87 = H. Isora.
javensis, *Blume, Bijdr.* 80.—Java.
lanata, *Kurz, For. Fl. Brit. Burma*, i. 143.—Siam.
lancæolata, DC. Prod. i. 476 = angustifolia.
longiflora, Wall. Cat. n. 1846 = spicata.
lanta, *Mart. in Flora*, xxii. (1839) 1. *Beibl.* 16.—Bras.
longifolia, Wall. Cat. n. 2687 = spicata †
macropetala, *A. Juss. in St. Hil. Fl. Bras. Mer.* i. 375.—Bras.
macrophylla, Wight, ex Wight & Arn. Prod. 60 = H. Isora.
mexicana, H. B. & K. Nov. Gen. et Sp. v. 305 = guazumæfolia.
microcarpa, *Span. in Linnæa*, xv. (1841) 173.—Malaya.
mollis, *Presl, Rel. Haenk.* ii. 139.—Mexic.
muscosa, *Mart. in Flora*, xxii. (1839) 1. *Beibl.* 14.—Bras.
oblonga, Wall. Cat. n. 1188, partim = spicata.
obtusa, *Wall. Cat.* n. 1184.—Burma; China.
ovata, *Lam. Encyc.* iii. 88.—Bras.
paniculata, Lour. Fl. Cochinch. 531 = Sterculia grandiflora.
pentandra, Linn. Mant. 294.—Guian.
pinnata, Blanco, Fl. Filip. ed. I. 384 = Odina speciosa.
plebeja, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 67.—Burma.
pronifera, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 111 = pentandra.
pulchella, *Wall. ex Bot. Hort Maurit.* 85.—Ind. or.
rotundata, *Griseb. Cat. Pl. Cub.* 29.—Cuba.
Rosenbrighii, G. Don, Gen. Syst. i. 506 = H. Isora.
Sacarolha, *A. Juss. in St. Hil. & Cambess. Pl. Us. Bras.* i. 64.—Bras.
salicifolia, *Presl, Rel. Haenk.* ii. 139.—Ins. Philipp.
semiglabra, *F. Muell. Fragm.* v. 43.—Austral.
semitriloba, *Bert. ex DC. Prod.* i. 476.—Ins. S. Doming.
spicata, *Cristr. ex Roxb. Hort. Beng.* [97], nomen.—Reg. Himal.; Malaya.
spinulosa, Wall. Cat. n. 1847 = glabriuscula.
sundaica, *G. Don, Gen. Syst.* i. 508.—Malaya.
tiliæfolia, *Dum.-Cours. Bot. Cult.* ed. II. v. 119.—Hab.?
tiliæfolia, Fisch. ex Weinm. in Syll. Ratisb. ii. (1828) 34 (= prior ?).—Bras.
trapezifolia, *A. Rich. Ess. Fl. Cub.* 170.—Cuba.
undulata, Lour. Fl. Cochinch. 531 = Sterculia lanceolata.
verbascifolia, Hort. Cels, ex Link, Enum. Hort. Berol. ii. 200 = ovata.
vestita, Wall. Cat. n. 1844 = spicata.
virgata, Wall. l. c. n. 1181 = angustifolia.
viscida, *Blume, Bijdr.* 79.—Java.
Vuardmæ, Mart. in Flora, xxii. (1839) 1. Beibl. 18 = altheæfolia.

HELICTONIA, Ehrh. Beitr. iv. 148 (1789) = Spiranthes, Rich. (Orchid.).

HELIE, M. Roem. Syn. Hesper. 42 (1846) = Atalantia, Corres. (Rutac.).
atalantioides, M. Roem. l. c. = A. buxifolia.
lanceolata, M. Roem. l. c. 43 = Glycosmis simplicifolia.

HELIETTA, Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 280. *RUTACEÆ*, Benth. & Hook. f. i. 801.
multiflora, *Engl. in Mart. Fl. Bras.* xii. ii. 185.—Bras.
parvifolia, *Benth. in Hook. Ic. Pl.* t. 1885.—Mexic.
Plasana, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 281.—N. Granat.

HELIGMA, Benth. & Hook. f. Gen. ii. 711 (1876) = seq.

HELIGME, Blume, Fl. Jav. Pracf. p. vii (1828) = Parsonsia, R. Br. (Apocyn.).
burmanica, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 406.—Java.
javanica, Blume, Bijdr. 1043 = P. spiralis.
Korthalsiana, Miq. Fl. Ind. Bat. ii. 489.—Sumatra.
Miquelianus, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 404.—Java.
Rheedii, Naves, in Blanco, Fl. Philipp. ed. III. t. 310 = P. Rheedii.
Rheedii, Wight, Ic. iv. t. 1302 = P. spiralis.
spiralis, Thw. Enum. Pl. Zeyl. 198 = P. spiralis.

HELINUS, E. Mey. ex Endl. Gen. 1102 (1840). *RHAMNEÆ*, Benth. & Hook. f. i. 385.
MYSTACINUS, Rafin. Sylva Tellur. 90 (1838) nonen phus.
arabica, *Yemb. & Spach, Illustr.* v. 81. t. 479.—Arab. brevipes, *Radlk. in Bremen, Abh.* viii. (1883) 385.—Afr. trop.
lanceolata, *Brand. For. Fl.* 574.—Reg. Himal.
Mystacinus, *E. Mey. ex Steud. Nom.* ed. II. i. 742; *Hemsl. in Oliver, Fl. Trop. Afr.* i. 385.—Afr. trop.
ovatus, *E. Mey. in Drège, Zwei Pfl. Documen.* 190, nomen; *ex Harv. & Sond. Fl. Cap.* i. 479.—Afr. austr.
scandens, A. Rich. Tent. Fl. Abyss. i. 159 = H. Mystacinus.

HELIOCARPOS, Linn. Gen. ed. I. 157 (1737) = seq.

HELIOCARPUS, Linn. Syst. ed. II. 25 (1740). *TILIACEÆ*, Benth. & Hook. f. i. 284.
MOLINA, Houst. ex Linn. Hort. Cliff. 211 (1737).
americanus, *Linn. Sp. Pl.* 468.—Mexic.
appendiculatus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 236.—Mexic.
arborescens, *Sesse. Bot. Voy. Herald*, 86.—N. Granat.
oblongifolius, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 578.—Bras.
popayanensis, *H. B. & K. Nov. Gen. et Sp.* v. 241.—N. Granat.
tomentosus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 235.—Mexic.
trichopodus, *Turcz. l. c.* 236.—Venesuel.

HELIOCARYA, Bunge, Heliocarya, Mosc. (1871) 4. *BORAGINEÆ*, Benth. & Hook. f. ii. 846.
monandra, *Bunge, l. c.*—Persia.

HELIOCHARIS, Lindl. Syn. 281 (1829) = **Blecocharis**, R. Br. (Cyperac.).

HELIOCHROA, A. Gray, Syn. Fl. N. Am. i. ii. 258 (1884) = HELICHROA, Rafin. = **Echinacea**, Moench (Compos.).

HELIOGENES, Benth. Pl. Hartw. 42 (1840) = **Agapippes**, DC. (Compos.).
longifolia, Benth. l. c. = A. bellidiflora.
Regiæ, Benth. l. c. = A. bellidiflora †

HELIOMERIS, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 171 = Gymnolomia, H. B. & K. (Compos.).
multiflora, Nutt. l. c. = G. multiflora.
simplex, Dur. in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 91 = seq.?
tenuifolia, A. Gray, Fl. Fendl. 84 = G. tenuifolia.

HELIONOPSIS, Franch. & Sav. Enum. Pl. Jap. ii. 529 (1879) = **Heloniopsis**, A. Gray (Liliac.).

HELIOPHILA, Burm. f. ex Linn. Sp. Pl. ed. II. 926 (1763). *CRUCIFERÆ*, Benth. & Hook. f. i. 81.
CARPOPODIUM, Eckl. & Zeyh. Enum. 13 (1835).
LEPTORMUS, Eckl. & Zeyh. Enum. 6 (1835).
ORMISCUS, Eckl. & Zeyh. l. c. 9 (1835).
ORTHOLIS, Spach, Hist. Vég. Phan. vi. 458 (1838).
PACHYSTYLUM, Eckl. & Zeyh. Enum. 13 (1835).
PRICIANA, Rafin. Sylva Tellur. 114 (1838).
SELENOCARDAMA, Eckl. & Zeyh. Enum. 10 (1835).
TRENTEPOHLIA, Roth, Catalect. ; ex ejusd. Nov. Pl. Sp. 235 (1821).
abrotanifolia, *Banks, ex DC. Syst.* ii. 690.—Afr. austr.
acuminata, *Steud. Nom.* ed. II. i. 742.—Afr. austr.
adinia, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 208.—Afr. austr.
aggregata, *Steud. Nom.* ed. II. i. 742.—Afr. austr.
amplexicaulis, *Linn. f. Suppl.* 296.—Afr. austr.
arabioides, Sims, Bot. Mag. t. 496 = pilosa.
arenaria, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 262. Afr. austr.
asurea, Willd. ex Sond. l. c. 209 = dissecta.
brachycarpa, *Meisen. in Hook. Lond. Journ. Bot.* i. (1842) 446.—Afr. austr.
brassicæfolia, *Eckl. & Zeyh. Enum.* 12.—Afr. austr.
burchellii, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 814 = Brassica cotyrhina.
caledonica, *Steud. Nom.* ed. II. i. 743; *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 205.—Afr. austr.
callosa, *DC. Syst.* ii. 696.—Afr. austr.
canescens, Willd. Sp. Pl. iii. 539 = Farsetia Jacquemontii.

HELIOPHILA :—
carnosa, *Steud. Nom.* ed. II. i. 742.—Afr. austr.
chamædrifolia, Stend. l. c., sphalm. = seq.
chamæmelifolia, *Burch. Trav.* i. 223, 235.—Afr. austr.
chamæmelifolia, Eckl. & Zeyh. Enum. 11 = abrotanifolia.
circæoides, Linn. f. Suppl. 298 = Chamira coronata.
clavicaulis, DC. Syst. ii. 695 = callosa.
cleomoides, Stend. Nom. ed. II. i. 800, sphalm. = glauca.
concatenata, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 214.—Afr. austr.
cornuta, *Sond. l. c.* 246. t. 28.—Afr. austr.
coronopifolia, Lam. Encyc. iii. 90, partim = pendula.
coronopifolia, *Linn. Sp. Pl.* ed. II. 927, partim.—Afr. austr.
coronopifolia, Thunb. Prod. Pl. Cap. 108 = pilosa.
crithmifolia, Eckl. & Zeyh. Enum. 12 = abrotanifolia.
crithmifolia, *Willd. Enum. Hort. Berol.* 652.—Afr. austr.
dentifera, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 219.—Afr. austr.
diffusa, *DC. Syst.* ii. 685.—Afr. austr.
digitata, Linn. f. Suppl. 296 = pilosa.
dissecta, Herb. Drège, ex Harv. & Sond. Fl. Cap. i. 41 = variabilis.
dissecta, *Thunb. Prod. Pl. Cap.* 108.—Afr. austr.
divaricata, *Banks, ex DC. Syst.* ii. 687.—Afr. austr.
divaricata, Eckl. & Zeyh. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 236 = stricta.
Dregeana, *Sond. l. c.* 260. t. 28. f. 2.—Afr. austr.
Eckloniana, *Sond. l. c.* 206. t. 20.—Afr. austr.
elata, *Sond. l. c.* 247.—Afr. austr.
elongata, DC. Syst. ii. 697 = linearifolia.
falcata, Eckl. & Zeyh. Enum. 11 = linearis.
fascicularis, *Banks, ex DC. Syst.* ii. 691.—Afr. austr.
fasciculoris, Eckl. & Zeyh. Enum. 12 = linearifolia.
filifolia, Thunb. Prod. Pl. Cap. 108 = linearifolia.
filifolia, Willd. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 212 = pusilla.
filiformis, Lam. Encyc. iii. 91 = longifolia.
filiformis, Linn. f. Suppl. 296 = Carpoponema filiformis.
fissulosa, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 209.—Afr. austr.
flacca, *Sond. l. c.* 238. t. 29. f. 1.—Afr. austr.
flava, Linn. f. Suppl. 297 = Brachycarpæa varians.
flornlenta, Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 268 = carnosa.
fonticulacea, *Ett. Br. in] Ait. Hort. Kew.* ed. II. iv. 100.—Afr. austr.
fornicivalvea, Eckl. & Zeyh. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 232 = tenuifolia.
frutescens, Lam. Encyc. iii. 91 = incana.
glabra, *Steud. Nom.* ed. II. i. 742.—Afr. austr.
glauca, *Burch. ex DC. Syst.* ii. 690.—Afr. austr.
glauca, Eckl. & Zeyh. Enum. 13 = virgata, linearis.
gracilis, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 230.—Afr. austr.
graminea, *DC. Syst.* ii. 697.—Afr. austr.
heterophylla, Thunb. Prod. Pl. Cap. 108 = abrotanifolia.
hirsuta, E. Mey. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 206 = pubescens.
incana, Burm. f. Fl. Ind. 178 = Farsetia Jacquemontii.
incana, [*Saland. in*] *Ait. Hort. Kew.* ed. I. ii. 397.—Afr. austr.
incana, Thunb. Fl. Cap. ed. Schult. 494 = pilosa.
incisa, Banks, ex DC. Syst. ii. 687 = pilosa.
inconspicua, E. Mey. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 227 = pectinata.
integrifolia, Linn. Sp. Pl. ed. II. 926 = pilosa.
juncea, Vahl, ex DC. Syst. ii. 691 = sinuosissima.
lanata, E. Mey. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 258 = crithmifolia.
latistiliqua, E. Mey, ex Sond. l. c. 234 = Thumbergii.
lepidioides, Link, Enum. Hort. Berol. ii. 176 (err. typ. 174) = diffusa.
linearifolia, *Burch. ex DC. Syst.* ii. 692.—Afr. austr.
linearis, *DC. Syst.* ii. 697.—Afr. austr.
linifiora, *DC. l. c.* 696.—Afr. austr.
longifolia, Eckl. & Zeyh. Enum. 13 = longifolia.
longifolia, *DC. Syst.* ii. 681.—Afr. austr.
longifolia, E. Mey. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 228 = affinis.
lyrata, Thunb. *Prod. Pl. Cap.* 108.—Afr. austr.
macrocarpona, *Burch. ex DC. Syst.* ii. 693.—Afr. austr.
macrostylis, *E. Mey, ex Sond. in Abh. Naturw. Hamb.* i. (1846) 225.—Afr. austr.

HELIOPHILA:—
maritima, *Eckl. & Zeyh. Enum.* 11.—Afr. austr.
Meyeri, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 226.—Afr. austr.
mollugines, *DC. Syst.* ii. 696.—Afr. austr.
monticola, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 216.—Afr. austr.
obliqua, E. Mey. ex Harv. & Sond. Fl. Cap. i. 58 = cuneata.
parviflora, Burch. ex DC. Prod. i. 233 = crithmifolia.
pectinata, *Burch. Trav.* i. 260.—Afr. austr.
pectinata, Eckl. & Zeyh. Enum. 11 = trifurca.
pectinata, E. Mey. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 236 = Meyeri.
Peltaria, *DC. Syst.* ii. 685.—Afr. austr.
pendula, E. Mey. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 40 = monticola.
pendula, Pall. ex Sond. l. c. 179 = Carponema filiformis.
pendula, *Willd. Sp. Pl.* iii. 529.—Afr. austr.
pennata, Sond. in Harv. & Sond. Fl. Cap. i. 41 = pendula.
pilosa, *Lam. Encyc.* iii. 90.—Afr. austr.
pinnata, Linn. f. Suppl. 297 = trifida.
pinnata, Vent. Jard. Malm. t. 113 = pendula.
pinnata, Willd. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 238 = crithmifolia.
platyuliqua, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 99. —Afr. austr.
platyuliqua, Eckl. & Zeyh. Enum. 12 = linearifolia.
platyuliqua, Meisn. in Hook. Lond. Journ. Bot. i. (1842) 462 = stylosa.
pubescens, *Burch. Trav.* i. 259.—Afr. austr.
pubescens, Eckl. & Zeyh. Enum. 11 = subulata.
pusilla, *Linn. f. Suppl.* 297.—Afr. austr.
refracta, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 234.—Afr. austr.
reticulata, *Eckl. & Zeyh. Enum.* 12.—Afr. austr.
rigidinscula, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 251. t. 27.—Afr. austr.
rivalis, *Burch ex DC. Syst.* ii. 682.—Afr. austr.
rostrata, Presl, Bot. Bemerk. 10 = pilosa.
sarcophylla, Meisn. in Hook. Lond. Journ. Bot. i. (1842) 468 = glauca.
scandens, *Harv. Thes. Cap.* ii. 43.—Afr. austr.
scoparia, *Burch. ex DC. Syst.* ii. 692.—Afr. austr.
scoparia, Herb. Drège, ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 246 = cuneata.
sesclifolia, *Burch. ex DC. Syst.* ii. 684.—Afr. austr.
sesclifolia, Eckl. & Zeyh. Enum. 11 = foeniculacea, crithmifolia.
sonchifolia, *DC. Syst.* ii. 681.—Afr. austr.
spathulata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 40 = amplexicaulis.
sphaerostigma, Kunze, Ind. Sem. Hort. Lips. (1846) = fistulosa.
stricta, Sims, Bot. Mag. t. 2526 = pilosa.
stricta, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 236.—Afr. austr.
stylosa, *Burch. ex DC. Syst.* ii. 693.—Afr. austr.
suaveissima, *Burch. ex DC. l. c.* 691.—Afr. austr.
suaveissima, Drège, ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 251 = rigidiuscula.
subulata, *Burch. ex DC. Syst.* ii. 691.—Afr. austr.
subulata, Drège, ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 251 = rigidiuscula.
succulenta, Banks, ex DC. Syst. ii. 692 = platyuliqua.
tenella, Banks, ex DC. l. c. 680 = dissecta.
tenuifolia, *Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 232.—Afr. austr.
tenuifolia, DC. Syst. ii. 680 = pusilla.
Thunbergii, *Steud. Nom.* ed. II. i. 748.—Afr. austr.
trifida, DC. Syst. ii. 688 = pinnata.
trifida, *Thunb. Prod. Pl. Cap.* 108.—Afr. austr.
trifurca, *Burch. ex DC. Syst.* ii. 686.—Afr. austr.
trifurca, Herb. Berol. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 239 = tenuifolia.
tripartita, Thunb. Prod. Pl. Cap. 108 = abrotanifolia.
variabilis, *Burch. ex DC. Syst.* ii. 683.—Afr. austr.
viminalis, *E. Mey. ex Sond. in Abh. Geb. Naturw. Hamb.* i. (1846) 231.—Afr. austr.
virgata, *Burch. ex DC. Syst.* ii. 693.—Afr. austr.
virgata, Eckl. & Zeyh. Enum. 13 = rigidiuscula.
virgata, Meisn. in Hook. Lond. Journ. Bot. i. (1842) 468 = stylosa.
Zeyheri, *Steud. Nom.* ed. II. i. 748.—Afr. austr.

HELIOPHTHALMUM, Rafin. Fl. Ludov. 72 (1817) = **Rudbeckia**, Linn. (Compos.).
cicutefolium, Rafin. l. c. 73 = R. cicutefolia.

HELIOPHYLA, Neck. Elem. iii. 79 (1790) = **Heliophila**, Burm. f. (Crucif.).

HELIOPHYLAX, Lestib. ex Steud. Nom. ed. II. i. 748 (1840) = HELSOPHYLAX, Beauv. = **Scirpus**, Linn. (Cyperac.).

HELIOPHYLLA, Scop. Introd. 317 (1777) = **Heliophila**, Burm. f. (Crucif.).

HELIOPHYTON, Benth. Pl. Hartw. 240 (1846) = seq.

HELIOPHYTUM, DC. Prod. ix. 551 (1845) = **Heliotropium**, Linn. (Boragin.).
anchusaefolium, DC. l. c. 554 = Heliotrop. anchusaefolium.
brasiliastachyum, DC. l. c.—Peruv.
chenopodiacrum, A. DC. in DC. l. c. 553 = Heliotrop. chenopodiaceum.
crispulum, Fresen. in Mart. Fl. Bras. viii. 45.—Bras.
elongatum, DC. Prod. ix. 555 = Heliotrop. elongatum.
erosum, DC. l. c. 552 = Heliotrop. erosum.
floridum, DC. l. c. 553 = Heliotrop. floridum.
fortiflorum, DC. l. c. = Heliotrop. fortiflorum.
glabriusculum, Torr. in Bot. Mex. Bound. 189 = Heliotrop. glabriusculum.
indicum, DC. Prod. ix. 556 = Heliotrop. indicum.
Ketschyi, Bunge, Rel. Bot. Lehm. 404 = Heliotrop. tuberculosum.
linearе, A. DC. in DC. Prod. ix. 555.—Afr. austr.
lithospermifolium, DC. l. c. 554.—Bras.
longiflorum, A. DC. in DC. l. c. 555 = Heliotrop. longiflorum.
macrostachyum, DC. l. c. 556 = Heliotrop. macrostachyum.
molle, Torr. in Bot. Mex. Bound. 188 = Tournefortia mollis.
monostachyum, DC. Prod. ix. 556 = Heliotrop. monostachyum.
nicotianaefolium, DC. l. c. 554 = Heliotrop. nicotianaefolium.
odorum, Fresen. in Mart. Fl. Bras. viii. 45.—Bras.
parviflorum, DC. Prod. ix. 553 = Heliotrop. parviflorum.
passerinoides, Klotzsch, in Schomb. Faun. & Fl. Gui. 1152, nomen.—Guian.
persicariaefolium, DC. Prod. ix. 556 = Heliotrop. monostachyum.
Petersii, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1852).—Afr. austr.
phylicoides, DC. Prod. ix. 554 = Heliotrop. phylicoides.
portoricense, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 297.—Ins. Porto Rico.
pterocarpum, DC. Prod. ix. 552 = Heliotrop. pterocarpum.
rosmarinifolium, Bert. ex DC. l. c. = Heliotrop. rosmarinifolium, *Bert.*
sidaefolium, DC. l. c. 553 = Heliotrop. sidaefolium.
tiaridioides, DC. l. c. 555 = Heliotrop. tiaridioide
volutinum, DC. l. c. 557 = Heliotrop. indicum.

HELIOPSIS, Pers. Syn. ii. 473 (1807). *COMPOSITAE*, Benth. & Hook. f. ii. 358.
ANDRIEUXIA, DC. Prod. v. 559 (1836).
HELENOMOIUM, Willd. ex DC. l. c. 551, in syn. (1836).
HELRPTA, Rafin. Neogenyt. 3 (1825).
KALLIAS, et CALLIAS, Cass. in Dict. Sc. Nat. xxiv. 327 (1822).
annua, *Hemsl. Biol. Centr. Am. Bot.* ii. 156.—Mexic.
Balsamorhiza, Hook. Fl. Bor. Am. i. 310 = Balsamorhiza terebinthacea.
buphthalmoides, *Dun. in Mém. Mus. Par.* v. (1819) 57.—Mexic.; Peruv.
buphthalmoides, A. Gray, Bot. Mex. Bound. 58 = parviflora.
canescens, D. Don, in Bot. Reg. t. 592 = scabra.
canescens, H. B. & K. Nov. Gen. et Sp. iv. 212 = buphthalmoides.

HELIOPSIS:—
dubia, Dun. in Mém. Mus. Par. v. (1819) 58 = Spilanthus Mutisii.
gracilis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii (1841) 353.—Am. bor.
helianthoides, Sweet, Hort. Brit. ed. I. 487 = laevis.
laevis, *Pers. Syn.* ii. 473.—Am. bor.
mollis, *Wender. Ind. Sem. Hort. Marb.* (1825); *ex Steud. Nom.* ed. II. i. 748.—Hab.?
paduls [patula?], *Wender. l. c.*; *ex Steud. l. c.*—Hab.?
parviolla, *A. Gray, in Pl. Wright.* ii. 86.—Mexic.
platyglossa, Cass. in Dict. Sc. Nat. xxiv. 332 = Gaudia abyssinica.
procumbens, *Hemsl. Biol. Centr. Am. Bot.* ii. 156.—Mexic.
rugosa, Hort. Berol. ex DC. Prod. v. 551 = bupthalmoides.
scabra, *Dun. in Mém. Mus. Par.* v. (1819) 54.—Am. bor.
stricta, *Wender. Ind. Sem. Hort. Marb.* (1825); *ex Steud. Nom.* ed. II. i. 748.—Hab.?
terebinthacea, Hook. Fl. Bor. Am. i. 310 = Balsamorhiza terebinthacea.

HELIOREOS, Rafin. Atl. Journ. 145 (1832) = **Fectis**, Linn. (Compos.).
angustifolius, Rafin. l. c. = P. angustifolia.

HELIOSCIADIUM, Bluff & Fing. Comp. Fl. Germ. i. 298 (1836) = HELOSCIADIUM, Koch = **Apium**, Linn. (Umbellif.).

HELIOSPERMA, Reichb. Nom. 206 (1841); et Ic. Fl. Germ. xvi. 71. t. 377 (1844) = **Silene**, Linn. (Caryophyll.).
alpestre, Reichb. Repert. 206 = S. alpestris.
chromodontum, Rohrb. Monog. Silene, 232 = S. chromodonta.
eriophorum, Juratzka, in Verh. Zool. Bot. Ges. Wien, viii. (1858) 87 = S. colonata ?
glutinosum, Reichb. ex Juratzka, l. c. Sitzb. 80 = S. quadridentata ?
macranthum, Panč. Elench. Crna-Gora, 11.—Montenegr.
monacherum, Vis. & Panč. in Mem. Ist. Venet. xii. (1864) 468 = S. quadridentata.
pudibundum, Griseb. Spicil. Fl. Rumel, i. 182 = S. quadridentata.
pusillum, Vis. Fl. Dalm. iii. 171 = S. quadridentata.
quadridentum, Griseb. Spicil. Fl. Rumel. i. 182, in obs. = S. quadrifida.
Tommasinii, Vis. Fl. Dalm. iii. 171.—Dalmat.
Veselskyi, Janka, in Bot. Zeit. xvi. (1858) 65.—Syria.

HELIOSPORA, Hook. f. Fl. Brit. Ind. iii. 127 (1880) = HELOSPORA, Jack = **Timonius**, Rumph. (Rubiac.).

HELIOTRICHUM, Bess. ex Schur, Enum. Pl. Transs. 762 (1866) = **Avena**, Tourn. (Gramin.).
planiculme, Bess. ex Schur, l. c. = A. planiculmis.

HELIOTROPIUM, [Tourn.] Linn. Syst. ed. I (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 843.
BOURJOTIA, Pomel, Nouv. Mat. Fl. Atl. 89 (1874).
BUCANION, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 568.
DIALION, Rafin. Sylva Tellur. 89 (1838).
ELIOPIA, Rafin. l. c. 90 (1838).
EUPLOCA, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 189.
HELIOPHYTUM, DC. Prod. ix. 551 (1845).
HIERANTHEMUM, Spach, Hist. Vég. Phan. iv. 31 (1840).
ORTHOSTACHYS, Spach, l. c. 32 (1840).
PENTACARVA, DC. Prod. ix. 539 (1845).
PERISTIMA, Rafin. Sylva Tellur. 89 (1838).
PIOCTONON, Rafin. l. c. 88 (1838).
PITPOCLAIMA, G. Don, Gen. Syst. iv. 364 (1837).
PRESLARA, Mart. Nov. Gen. et Sp. ii. 75. t. 164 (1827).
PRESLIA, Spreng. Syst. iv. Cur. Post. 336 (1827).
SARCANTHUS, Anderss. in Vet. Acad. Handl. Stockh. 1853 (1855) 209.
SCHLEIDENIA, Endl. Gen. 646 (1839).
SCHOBERA, Scop. Introd. 158 (1777).
SCORPIANTHES, Rafin. Sylva Tellur. 89 (1838).
SYNZISTACHYUM, Rafin. l. c. (1838).
TIARIDIUM, Lehm. Asperif. 15 (1818).

HELIOTROPIUM:—
abyssinicum, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 168.—Abyssin.
acutiflorum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 406.—Soongaria.
aegyptiacum, Lam. Ind. Sem. Hort. Hamb. (1830) 7 = pallens.
astracum, Jan, Elench. 3; Presl, ex DC. Prod. ix. 535 = Bocconi.
affghanum, Boiss. Fl. Orient. iv. 148 = undulatum.
albicaulon, *Schum. & Thonn. Beskr. Guin. Pl.* 87.—Afr. trop.
aqdense, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) 1. 297.—Persia.
aleppicum, *Botan. Diagn.* Ser. I. xi. 58.—Cilicia; Syria.
amblaniense, G. Don, Gen. Syst. iv. 361, sphalm. = urbainense.
amblyoum, *DC. Prod.* ix. 538.—Afr. austr.
amboinense, DC. l. c. 537, sphalm. = urbainense.
americanum, *Mill. Gard. Dict.* ed. VIII. n. 11.—Mexic.
amplexicaule, *Vahl, Symb. Bot.* iii. 21.—Bras.
anchusaefolium, *Poir. Encyc. Suppl.* iii. 23.—Reg. Argent.
angiospermum, Murr. Prod. Stirp. Gotting. 217 = parviflorum.
angustifolium, *Torr. in Bot. Mex. Bound.* 137.—Texas; Mexic.
anisephyllum, Beauv. Fl. Ow. ii. 62 = indicum.
anomalum, *Hook. & Arn. Bot. Beech. Voy.* 66.—Ins. Pacif.
apiculatum, E. Mey. ex DC. Prod. ix. 541 = ovalifolium.
arbainense, *Fresen. in Mus. Senckenb.* i. (1838) 168.—Aegypt.; Arab.
arborescens, Linn. Syst. ed. X. 918 = peruvianum.
arenarium, F. Muell. Fragm. vi. 112 = undulatum.
armenium, *Vatke, in Linnaea*, xliii. (1880–83) 219.—Afr. trop.
argenteum, Willd. ex Lehm. Asperif. 78 = lanatum.
argutioides, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 406.—Soongaria.
asperrimum, *R. Br. Prod.* 493.—Austral.
asperrimum, Auch. ex DC. Prod. ix. 537 = ramosissimum.
Aucheri, *DC. l. c.* 538.—Persia.
bacciferum, *Forsk. Fl. Aegypt. Arab.* 38.—Arab.
bacciferum, [Mog. & Sesse,] ex DC. Prod. ix. 551 = Tournefortia Hernandesii.
Baclei, *DC. l. c.* 546.—Afr. trop.
bahiense, DC. l. c. 544 = polyphyllum.
barbatum, *DC. l. c.* 541.—Bras.
biannulatum, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) 300.—Afghan.
bicolor, Hochst. & Steud. ex DC. Prod. ix. 546 = strigosum.
Bocconi, *Guss. Ind. Sem. Hort. Boccad.* (1835) 6; *Fl. Sic. Prod.* i. 305.—Sicil.; Graec.
Borei, *Botan. Diagn.* Ser. I. xi. 57.—As. Min.; Syria.
brachygype, *Benth. Fl. Australas.* iv. 396.—Austral.
bracteatum, *R. Br. Prod.* 493.—Ind. or.; Malaya; Austral.
brahuicum, Stocks, in Hook. Kew Journ. iv. (1852) 173 = dasycarpum.
brasilianum, *Roth, Nov. Pl. Sp.* 108.—Bras.
brevifolium, Wall. Cat. n. 914 = strigosum.
brevilimbe, *Boiss. Diagn.* Ser. I. xi. 59.—Persia.
Brocchianum, Vis. Fl. Aeg. et Nub. 8. t. 2. f. 1 = ovalifolium.
Burmanni, *Roem. & Schult. Syst.* iv. 736.—Afr. austr.
buxifolium, Wright, in Griseb. Cat. Pl. Cub. 211 = polyphyllum.
cabulicum, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) 1. 298.—Afghan.
calcareum, *Stocks, in Hook. Kew Journ.* iv. (1852) 174.—Baluchist.
californicum, Greene, in Bull. Calif. Acad. i. iv. (1885) 208 = convolvulaceum.
callosum, Sprang. Nov. Prov. 21 = lanosum.
campechianum, *Jl. B. & K. Nov. Gen. et Sp.* iii. 86.—Mexic.
campestre, *Griseb. in Goett. Abh.* xix. (1874) 234.—Reg. Argent.
canariense, Mill. Gard. Dict. ed. VIII. n. 5 = Bystropogon canariense.
canariense, Willd. ex Roem. & Schult. Syst. iv. 736 = erosum.
canescens, H. B. & K. Nov. Gen. et Sp. iii. 86 = inundatum.
canescens, *C. Koch, in Linnaea*, xix. (1847) 21.—As. Min.
canescens, Lehm. Asperif. 36 = barbatum?
canescens, Moench, Meth. 415 = europaeum.
canescens, Willd. ex Lehm. Asperif. 38 = Willdenowii.
capense, *Sw. ex Lehm. in Neue Schr. Naturf. Ges. Halle*, iii. 11. (1817) 18; *Asperif.* 50.—Afr. austr.
capense, *Thunb. Fl. Cap.* ii. 245.—Afr. austr.
capitatum, Mill. Gard. Dict. ed. VIII. n. 4 = Bystropogon canariense.
cardiochorum, C. Koch, in Linnaea, xxii. (1849) 627 = europaeum.
carnanicum, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) 1. 324.—Persia.
chenopodiaceum, *C. Gay, Fl. Chil.* iv. 458.—Chili.
chenopodioides, Willd. Enum. Hort. Berol. 175 = curassavicum.
chilense, *Bert. in Mercr. Chil.* 614.—*Cf. Bull. Fbrussac*, xx. (1830) 113.—Chili.
chorassanicum, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) 1. 301.—Persia.
chrysanthum, *Phil. in Anal. Univ. Chil.* (1862) 11. 401.—Chili.
cinerascens, Aitch. Cat. Punjab Pl. 94 = calcareum.
cinerascens, *Stocks, ex DC. Prod.* ix. 534.—Abyssin.
cinereum, R. Br. in Salt, Abyss. App. 62 = ovalifolium.
cinereum, H. B. & K. Nov. Gen. et Sp. iii. 89. t. 206 = inundatum.
circinatum, *Griseb. Spicil. Fl. Rumel.* ii. 78.—As. Min.; Mesopot.
citriodorum, *Lehm. Asperif.* 82.—Am. austr.
Clausseni, *DC. Prod.* ix. 543.—Bras.
commutatum, Roem. & Schult. Syst. iv. 29 = europaeum.
compactum, D. Don, Prod. Fl. Nep. 101 = strigosum.
conferti*florum*, *Boiss. & Hoh. Diagn.* Ser. II. iii. 132.—Persia.
confertifolium, *Torr. ex A. Gray, Syn. Fl. N. Am.* ii. 1. 184.—Mexic.
conocarpum, *F. Muell. ex Benth. Fl. Australas.* iv. 398.—Austral.
convolvulaceum, *A. Gray, in Mem. Am. Acad.* vi. (1857) 408.—Am. bor. occ.
cordifolium, Moench, Meth. 415 = indicum.
cordofanum, Hochst. ex DC. Prod. ix. 546 = strigosum.
coriaceum, *Lehm. in Nov. Act. Nat. Cur.* xiv. ii. (1829) 810.—Mexic.
coromandelianum, Koen. ex Retz. Obs. ii. 9 = ovalifolium.
corymbodelianum, Raddi, ex DC. Prod. ix. 533 = supinum.
corymbosum, *Ruiz & Pav. Fl. Per.* ii. 3. t. 107.—Peruv.
crassifolium, *Boiss. Diagn.* Ser. II. iii. 131.—Persia.
crassifolium, *Phil. in Anal. Univ. Chil.* (1873) 515.—Chili.
crenatum, *Franch. in Revoil, Comali*, 46.—Afr. trop.
crispatum, *F. Muell. ex Benth. Fl. Australas.* iv. 395.—Austral.
crispum, Desf. Fl. Atlant. i. 151. t. 41 = undulatum.
Cunninghamii, *Benth. Fl. Australas.* iv. 400.—Austral.
curassavicum, *Linn. Sp. Pl.* 130.—Cosmop. trop.
cyrtostachyum, Miq. Fl. Ind. Bat. ii. 924 = bracteatum.
dasycarpum, *Ledeb. ex Bichw. Pl. Nov. Casp.-Cauc.* 11. t. 5.—As. occ.; Ind. or.
decumbens, *Lehm. in Neue Schr. Naturf. Ges. Halle*, iii. ii. (1817) 16.—Venezuela.
demissum, Roem. & Schult. Syst. iv. 26, 733 = fruticosum.
denticulatum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 81.—Ins. Socotra.
denticulatum, *Boiss. & Hausskn. in Boiss. Fl. Orient.* iv. 145.—Persia.
deserticolum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 150 = transoxanum.
dieteriflorum, *Botan. Diagn.* Ser. I. xi. 58.—Persia.
dietrichianum, Stocks, ex Boiss. Fl. Orient. iv. 146 = H. Ophioglossum.
diversifolium, *F. Muell. ex Benth. Fl. Australas.* iv. 400.—Austral.
dolosum, De Not. Repert. Ligur. 219 = Richwaldi.
dubium, *R. Br. in Salt, Abyss. App.* 62.—Abyssin.
Ehrenbergii, *Steud. Nom.* ed. II. i. 744.—Europ. austr.; As. temp.; Austral.
HELIOTROPIUM:—
elachanthum, F. Muell. in Linnaea, xxv. (1852) 424 = Eritrichium australasicum.
ellipticum, *R. Br. in Salt, Abyss. App.* 62.—Abyssin.
ellipticum, Ledeb. in Eichw. Pl. Nov. Casp. Cauc. 10.—Eichwaldi.
elongatum, *Willd. ex Cham. in Linnaea*, iv. (1829) 485.—Bras.
epacrideum, *F. Muell. ex Benth. Fl. Australas.* iv. 396.—Austral.
erectum, Lam. Fl. Fr. ii. 381 = europaeum.
erectum, *Vell. Fl. Flum.* 69; ii. t. 42.—Bras.
eremobium, *Bunge, in Bull. Soc. Nat Mosc.* xlii. (1869) 1. 321.—Persia.
ericarpum, Delile, ex Lehm. Asperif. 55 = undulatum.
ericarpum, Schimp. ex DC. Prod. ix. 537 = ramosissimum.
erirtichioides, *Kotschy, in Sitzb. Acad. Wien, Math.-Nat. lit. Abth.* i. (1866) 259.—Orient.
erosum, *Lehm. in Neue Sch. Naturf. Ges. Halle*, iii. ii. (1817) 15.—Ins. Teneriff.
europaeum, Aitch. in Journ. Linn. Soc. xix. (1882) 178 = Richwaldi.
europaeum, Forsk. Fl. Aegypt. Arab. 38 = supinum.
europaeum, *Linn. Sp. Pl.* 130.—Europ. austr.; Afr. bor.; Orient; Austral.
europaeum, Pall. Ind. Taur. 47, nomen = mavolens.
fasciculatum, *R. Br. Prod.* 494.—Austral.
filaginoides, *Benth. Fl. Australas.* iv. 398.—Austral.
filiforme, *H. B. & K. Nov. Gen. et Sp.* iii. 86. t. 204.—Am. trop.
filosum, Lehm. ex DC. Prod. ix. 550, nomen.—Hab.? floridum, *C. Gay, Fl. Chil.* iv. 457.—Chili.
foetidum, Salisb. Prod. 113 = indicum.
foetidum, *Salzm. ex DC. Prod.* ix. 551, 553.—Bras.
foliatum, R. Br. Prod. 493 = bracteatum.
foliatum, Lehm. Pl. Preiss. ii. 338 = asperrimum.
foliosum, Willd. ex Roem. & Schult. Syst. iv. 737 = polyphyllum.
fruticosum, Forsk. Fl. Aegypt. Arab. 38 = strigosum.
fruticosum, *Linn. Syst.* ed. X. 918.—N. Granat.
globellum, R. Br. Prod. 494 = paniculatum.
glabriusculum, *A. Gray, in Proc. Am. Acad.* x. (1875) 50.—Mexic.
glandulosum, R. Br. Prod. 493 = europaeum.
glaucophyllum, Moench, Meth. Suppl. 147 = curassavicum.
glaucum, Salisb. Prod. 113 = curassavicum.
glomeratum, A. DC. in DC. Prod. ix. 550 = polyphyllum.
glutinosum, *Phil. Fl. Atac.* 38.—Chili.
gnaphalioides, Auct. = sq.
gnaphaloides, Linn. Syst. ed. X. 918 = Tournefortia gnaphaloides.
gracile, R. Br. Prod. 493 = ovalifolium.
gracile, R. Br. in Salt, Abyss. App. 62 = Saltianum.
gracillimum, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) 1. 294.—Persia.
grandiflorum, *Auch. ex DC. Prod.* ix. 542.—Armenia.
grandiflorum, Donn, Hort. Cantab. ed. VI. 42; Schrank, Pl. Rar. Hort. Monac. t. 2 = corymbosum.
Greggii, *Torr. in Bot. Mex. Bound.* 137.—Mexic.
Griffithii, *Boiss. Diagn.* Ser. II. iii. 131.—Afghan.
gymnocarpum, Borb. in Koeslem. xiv. (1876) 405 = europaeum.
Halam, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 150.—Persia.
Hausknechtii, Bunge, in Bull. Soc. Nat. Mosc. xlii. (1869) 1. 310 = circinatum.
heliophilum, *Meisn. in Flora*, xxi. (1838) ii. *Beibl.* 88.—Bras.
hirsutissimum, *Grauer, Dec.* 1.—Europ. austr.; As. Min.
hirsutissimum, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 167.—Afr. trop.
hirsutum, *Clarke, Trav.* ii. 481 = supinum.
hirtum, Lehm. in Neue Schr. Naturf. Ges. Halle, iii. ii. (1817) 10 = fruticosum.
hispidum, H. B. & K. Nov. Gen. et Sp. iii. 87 = fruticosum.
horminoides, Mill. Gard. Dict. ed. VIII. n. 3 = indicum.
Howardi, DC. Prod. ix. 549 = procumbens.
Humboldtianum, Roem. & Schult. Syst. iv. 737 = inundatum.
humifusum, *H. B. & K. Nov. Gen. et Sp.* iii. 85.—Cuba.
humile, R. Br. Prod. 497 = fruticosum.

HELIOTROPIUM :—
humile, *Lam. Illustr.* i. 393.—Ind. occ.
humile, Salisb. Prod. 113 = europæum.
humile, Schult. Obs. Bot. 27 = stavesolense.
humistratum, *Chem. in Linnæa*, iv. (1829) 462.—Bras.
imbricatum, *Griseb. Cat. Pl. Cub.* 211.—Cuba.
incanescens, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 89, nomen.—Rossia.
incanum, *Ruiz & Pav. Fl. Per.* ii. 2. t. 108.—Ecuador.
indicum, *Linn. Sp. Pl.* 180.—Cosmop. trop.
intermedium, Andrz. ex DC. Prod. ix. 535 = stavesolens.
inundatum, *Sw. Prod. Veg. Ind. Occ.* 40.—Am. calid.
japonicum, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 408.—Japon.
Kotschyanum, Bunge, in Bull. Soc. Nat. Mosc. xlii. (1869) i. 319 = Bovei.
Kovalevskyi, Stschég. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 296 = Szowitsii.
Krallikii, *Pomel, Nouv. Mat. Fl. Atl.* 89.—Alger.
kumense, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) i. 330.—Persia.
Kunzii, Lehm. Ic. et Desc. Stirp. 19 = ovalifolium.
lacunarifolium, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 20 = europæum.
lanatum, *H. B. & K. Nov. Gen. et Sp.* iii. 89.—N. Granat.
lanceolatum, Norohha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 18 = indicum.
lanceolatum, *Ruiz & Pav. Fl. Per.* ii. 4. t. 111.—Peruv.
laticarpum, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 38 = Eichwaldi.
latifolium, *Willd. ex Lehm. Asperif.* 33.—Venezuela.
lasiflorum, Kech. Nov. Pl. Sp. 109 = bracteatum.
Leavenworthii, Torr. ex A. Gray, Syn. Fl. N. Am. ii. 1. 185 = polyphyllum.
ligusticum, *Vatke, in Oestr. Bot. Zeitschr.* (1876) 167, in obs.—Afr. trop.
limbatum, *Benth. Pl. Hartw.* 20.—Mexic.
limbatum, Torr. Bot. Mex. Bound. 138 = confertifolium.
linearifolium, *Phil. Fl. Atac.* 38.—Chili.
lineatum, Delile, Fl. Egypt. 87 = luteum.
lineatum, *Vahl, Symb. Bot.* i. 18.—Aegypt.
linifolium, Lehm. Asperif. 35 = paniculatum.
lithospermoides, Willd. ex Roem. & Schult. Syst. iv. 737 = Anchusa tuberosa.
littorale, Stev. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 565 = Eichwaldi.
longiflorum, *Hochst. & Steud. ex DC. Prod.* ix. 555.—Arab.
longifolium, *Klotzsch, in Peters, Reise Mossamb. Bot.* 251.—Afr. trop.
longifolium, Norohha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 18 (Quid?).—Java.
longistylum, *Phil. in Anal. Univ. Chil.* (1873) 515.—Chili.
luteum, Aitch. Cat. Pl. Punj. 94 = dasycarpum.
lutetum, *Poir. Encyc. Suppl.* iii. 23.—Aegypt.
macrocarpum, Guss. Enum. Pl. Inarim. 214. t. 7 = Eichwaldi.
macrostachyum, *Hemsl. Biol. Centr. Am. Bot.* ii. 375.—Mexic.
malabaricum, Retz. Obs. iv. 24 = supinum.
mamanense, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) i. 308.—Persia.
marchionicum, *Decne. Bot. Voy. Vénus*, 21.—Ins. Marques.
marifolium, *Koen. ex Retz. Obs.* ii. 8.—Ind. or.
marocczanum, Lehm. Asperif. 36 = undulatum.
marocczanum, Raddi, Cat. 26 ; ex DC. Prod. ix. 537 = urbelianum.
mendocinum, *Phil. in Anal. Univ. Chil.* (1868) ii. 600.—Chili.
mesinense, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) i. 312.—Persia.
microcalix, Boiss. Fl. Orient. iv. 186 = microcalyx.
microcalyx, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 209.—Turkestan.
microcarpum, *Wats. Apont.* 291.—Arab. trop.
microchlyx, *Ruiz & Pav. Fl. Per.* ii. 2. t. 109.—Peruv.
microphyllum, *Sw. ex Wikstr. in Vet. Acad. Handl. Stockh.* (1827) 86.—Ins. Guadalup.
microstachyum, *Ruiz & Pav. Fl. Per.* ii. 2. t. 110.—Peruv.
microstylosum, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) i. 324.—Persia.

HELIOTROPIUM :—
monostachyum, *Cham. in Linnæa*, iv. (1829) 455.—Bras.
morocczanum, G. Don, Gen. Syst. iv. 360 = undulatum.
myosotoides, Champ. Fl. S. U. St. 380 = phyllostachyum.
myosotoides, Lehm. in Neue Schr. Naturf. Ges. Halle, iii. 11. (1817) 16 = bicolor.
myosotoides, [Soland. in] Russell, Aleppo, ed. II. ii. 245 = Cynbalcolæ myosotoides.
neglectum, Bunge, in Bull. Soc. Nat. Mosc. xlii. (1869) i. 312 = circinnatum.
nicotianæfolium, *Poir. Encyc. Suppl.* iii. 23.—Reg. Argent.
nigricans, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 81.—Ins. Socotr.
nilsoticum, A. DC. in DC. Prod. ix. 541 = ovalifolium.
Noëanum, *Boiss. Diagn.* Ser. II. iii. 132.—Persia.
nubicum, Bunge, in Bull. Soc. Nat. Mosc. (1869) i. 330 = undulatum.
oaxacanum, *DC. Prod.* ix. 548.—Mexic.
oblongifolium, Hornem. Hort. Hafn. Suppl. 19 = europæum.
oblongifolium, *Hart. & Gal. in Bull. Acad. Brux.* xi. ii. (1844) 386.—Mexic.
obovatum, D. Don, Prod. Fl. Nep. 101 = ovalifolium.
ocellatum, *Cham. in Linnæa*, iv. (1829) 463.—Bras.
odoratum, Moench, Meth. 415 = peruvianum.
odorum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 81.—Ins. Socotr.
odorum, Salisb. Prod. 119 = peruvianum.
odorum, Saov. ex Stev. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 567 = suaveolens.
Olgæ, Bunge, in Act. Hort. Petrop. iii. (1875) 114 = chorassanicum.
Ophioglossum, *Stocks, ex Aitch. Cat. Pl. Punj.* 94.—Beluchist.
opposititolium, *Ruiz & Pav. Fl. Per.* ii. 2. t. 108.—Peruv.
orientale, *Linn. Sp. Pl.* 131 ; *Houtt. Handleid.* vii. 418 (pro sub.).—Malaya.
Ottoni, *Lehm. Asperif.* 59.—Venezuela.
Ottonianum, Roem. & Schult. Syst. iv. 733 = præc.
ovalifolium, *Forsk. Fl. Aegypt.-Arab.* 35.—Geront trop.
pallens, *Delile, Fl. Afr. Calil.* 69. t. 64.—Afr. trop.
Palmeri, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 121.—Mexic.
paniculatum, *Br. Prod.* 494.—As. et Austral. trop.
paniculatum, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 148 = zeylanicum.
paniculatum, Roxb. Hort. Beng. [28] ; Fl. Ind. i. 465 = Roxburghii.
pannifolium, *Burch. ex Hemsl. Bot. Voy. Challenger*, i. ii. 76.—Ins. S. Helena.
parviflorum, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 167.—Arab.
parciflorum, *Griseb. Fl. Brit. W. Ind.* 486, 742 ; *Cat. Pl. Cub.* 212.—Cuba.
parcmychioides, *DC. Prod.* ix. 565.—Chili.
parvifolium, Blanco, Fl. Filip. ed. I. 80 = indicum.
parvifolium, *Linn. Mant.* ii. 201.—Am. trop.
parvifolium, Stev. ex Boiss. Fl. Orient. iv. 181 = Eichwaldi.
parvifolium, Edgew. in Journ. As. Soc. Beng. xvi. (1846) 1316 = strigosum.
patebilosum, *H. B. & K. Nov. Gen. et Sp.* iii. 87.—Am. austr.
pauciflorum, *R. Br. Prod.* 493.—Austral.
Pearcei, *Phil. in Anal. Univ. Chil.* (1861) i. 65.—Chili.
persicum, Aucb. ex DC. Prod. ix. 551 ; x. 72 = Moltkia angustifolia.
persicum, *Burm. f. Fl. Ind.* 41. t. 29.—Ind. or.
persicum, Lam. Encyc. iii. 94 = undulatum.
peruvianum, *Linn. Sp. Pl.* ed. II. 187.—Peruv.
phyllocladum, *Cham. in Linnæa*, iv. (1829) 460.—Bras.
phyllostachyum, *Torr. Bot. Mex. Bound.* 137.—Am. bor. occ.
pilosum, *Ruiz & Pav. Fl. Per.* ii. 3. t. 110.—Peruv.
pinnatum, Vahl, Symb. Bot. iii. 21 = Fiacella circinnata.
plebeium, Banks, ex Buch, Besch. Canar. Ins. 144 = arenar.
pleiopterum, *F. Muell. Progm.* ix. 121.—Austral.
polyphyllum, *Lehm. in Neue Schr. Naturf. Ges. Halle*, iii. 11. (1817) 9.—Am. trop.
polystachyum, Poir. Encyc. Suppl. iii. 23 = ovalifolium.

HELIOTROPIUM :—
procumbens, H. B. & K. Nov. Gen. et Sp. iii. 88 = inundatum.
procumbens, *Mart. Gard. Dict.* ed. VIII. n. 10.—Am. trop.
prostratum, *R. Br. Prod.* 494.—Austral.
pruinosum, Bunge, in Bull. Soc. Nat. Mosc. xlii. (1869) i. 310 = Bovei.
pteroscarpum, *Hochst. & Steud. ex DC. Prod.* ix. 552.—Arab.
pycnophyllum, *Phil. Fl. Atac.* 38.—Chili.
pygmæum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 252.—Afr. trop.
Radula, *Fisch. & Mey. ex Ledeb. Fl. Ross.* iii. 101.—Turkestan.
ramosissimum, Sieber, ex DC. Prod. ix. 536 = undulatum.
ramosum, Roxb. ex Wall. Cat. n. 7015 = bracteatum.
rariflorum, *Stocks, in Hook. Kew Journ.* iv. (1852) 174.—Beluchist.
repens, *Griseb. in Gœtt. Abh.* xix. (1874) 288.—Reg. Argent.
rigidifolium, *A. DC. in DC. Prod.* ix. 543.—Bras.
rigidulum, *DC. l. c.* ix. 540.—Bras.
rosmarinifolium, Bert. ex Steud. Nom. ed. II. i. 744 = stenophyllum.
rosmarinifolium, *Phil. in Anal. Univ. Chil.* (1873) 514.—Chili.
Rottleri, *Lehm. Asperif.* 66.—Ind. or.
rotundifolium, *Sieber, ex Lehm. Ic. et Desc. Stirp.* 19. t. 20.—Syria ; Palæstin.
Roxburghiana, Voigt, Hort. Suburb. Calc. 445 = seq.
Roxburghii, *Spreng. Syst. iv. Cur. Post.* 54.—Ind. or.
rugosum, *Mart. & Gal. in Bull. Acad. Brux.* xi. ii. (1844) 386.—Mexic.
rugosum, *Phil. Fl. Atac.* 38.—Chili.
rupestre, *Mart. & Gal. in Bull. Acad. Brux.* xi. ii. (1844) 386.—Mexic.
salicioides, *Cham. in Linnæa*, viii. (1833) 117.—Bras.
salsum, *Griseb. in Gœtt. Abh.* xix. (1874) 288.—Reg. Argent.
Saltianum, *Steud. Nom.* ed. II. i. 744.—Abyssin.
samoliifolium, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) i. 314.—Persia.
scabrum, Lehm. Asperif. 70 = marifolium.
scabrum, *Retz. Obs.* ii. 8.—Ind. or.
scandens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 18 = Tournefortia tetrandra.
scandens, *Vell. Fl. Flum.* 69 ; ii. t. 41.—Bras.
Schrenkianum, Ledeb. Fl. Ross. iii. 109 = dasycarpum.
sclerocarpum, *Phil. in Anal. Univ. Chil.* (1873) 515.—Chili.
scorpioides, H. B. & K. Nov. Gen. et Sp. iii. 89 = latifolium.
scorpioides, Willd. ex Roem. & Schult. Syst. iv. 737 = Myosotis grandiflora.
senense, *Klotzsch, in Peters, Reise Mossamb. Bot.* 253.—Afr. trop.
sericocarpum, Bunge, in Bull. Soc. Nat. Mosc. xlii. (1869) i. 322 = Aucheri.
serpyllioides, *Griseb. Cat. Pl. Cub.* 213.—Cuba.
sidæfolium, *Cham. in Linnæa*, iv. (1829) 460.—Bras.
Sieniten, DC. Prod. ix. 541 = *H. Syenites*.
simile, *Vatke, in Linnæa*, xliii. (1880–81) 317.—Afr. trop.
simplex, *Meyen, Reise*, i. 426.—Peruv.
smyrnaceum, Bunge, in Bull. Soc. Nat. Mosc. xlii. (1869) i. 311 = Bovei.
sogdianum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 402.—Turkestan.
somalense, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 166.—Afr. trop.
stenophyllum, *Hook. & Arn. Bot. Beech. Voy.* 33.—Chili.
Steudneri, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 167.—Afr. trop.
Stevenianum, Andrz. ex Hohen. Enum. Pl. Talusch. 76 = Eichwaldi.
strictum, Hemsl. Biol. Centr. Am. Bot. ii. 376 = strictum.
strictissimum, *Moric. Pl. Nouv. Am.* 146. t. 67.—Bras.
strictum, *H. B. & K. Nov. Gen. et Sp.* iii. 87.—Venezuela.
strictum, Ledeb. Fl. Ross. iii. 100 = Eichwaldi.
strigosum, *Willd. Sp. Pl.* i. 748.—Afr. trop. Beluchist. ; Austral.

HELIOTROPIUM:—
syligerum, *Treviso. in Act. Hort. Petrop.* vii. (1880) 484.—Reg. Caucas.
stylosum, Franch. in Révoil, Çomali, 45 = H. Ophioglossum.
stylosum, *Phil. in Bot. Zeit.* xxviii. (1870) 800.—Chili.
suaveolens, *Bieb. Fl. Taur.-Cauc.* iii. 116.—As. Min.; Persia.
subcaseosum, Stev. in Bull. Soc. Nat. Mosc. xxiv. (1851) 1. 564 = europæum.
submolle, *Klotzsch, in Otto & Dietr. Allg. Gartens.* xx. (1852) 89.—Ecuador.
subulatum, Hochst. ex DC. Prod. ix. 529, 551 = zeylanicum.
suffruticosum, *Pomel, Nouv. Mat. Fl. Atl.* 90.—Alger.
sultanense, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) 1. 316.—Persia.
supinum, *Linn. Sp. Pl.* 130.—Europ. austr.; Afr bor.; Orient.
supinum, Pall. ex Georgi, Beschr. Russ. Reich. iv. 749 = europæum.
supinum, Thunb. Prod. Fl. Cap. 88 = ambiguum.
Syenites, Spreng. Syst. i. 589 = ovalifolium.
synsystachyum, *Ruiz & Pav. Fl. Per.* ii. 3.—Peruv.
Szowitsii, *Stschig. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 1. 235.—Armen.
teheranicum, *Bunge, l. c.* xlii. (1869) 1. 309.—Persia.
tenellum, *Torr. in Marcy, Rep. Expl. Red River* (1853), 304. t. 14.—Am. bor.
tenue, Wall. Cat. n. 915 = paniculatum.
tenue, *Willd. ex Roem. & Schult. Syst.* iv. 737 = filiforme.
tenuiflorum, Guss. Fl. Sic. Syn. i. 216 = europæum.
tenuifolium, *R. Br. Prod.* 494.—Austral.
ternatum, *Vahl, Symb. Bot.* iii. 21.—Ind. occ.
tetranthum, *Lour. Fl. Cochinch.* i. 108.—China.
thymeloides, Walp. Ann. v. 548 = strigosum.
thymifolium, *Vahl, ex Lehm. in Nouv Schr. Naturf. Ges. Halle* iii. 11. (1817) 17; *Asperif.* 67.—Hab.? *thymoides*, Jaub. & Spach, Illustr. iv. 98 = strigosum.
tiaridicides, *Cham. in Linnæa*, iv. (1829) 458.—Bras.
tomentosum, Burm. f. Fl. Cap. Prod. 4 = Burmanni.
tomentosum, Poir. Encyc. Suppl. iii. 28 = ovalifolium.
transalpinum, *Vell. Fl. Fium.* 68; ii. t. 40.—Bras.
tranaxoxanum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 402.—Persia; Turkest.
trichostomum, *Bunge, in Bull. Soc. Nat. Mosc.* xlii. (1869) 1. 318.—Persia.
tuberculosum, *Boiss. Fl. Orient.* iv. 147.—Persia.
tuberosum, *E. Mey. ex DC. Prod.* ix. 537.—Afr. austr.
turcicum, Friv.ld. in Flora, xix. (1836) 436 = suaveolens.
umbellatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. xv. 18 (Quid?).—Java.
undulatum, Kral. ex Pomel, Nouv. Mat. Med. Atl. 89 = Kralikii.
undulatum, *Vahl, Symb. Bot.* i. 18.—As. occ.; Afr. trop.; Austral.
ventricosum, *R. Br. Prod.* 494.—Austral.
veronicifolium, *Griseb. in Goett. Abh.* xix. (1874) 282. —Reg. Argent.
vestitum, *Benth. Fl. Australi.* iv. 395.—Austral.
villosum, Presl, Fl. Sic. p. xxxiv = Bocconi.
villosum, Sieber. ex DC. Prod. ix. 541 = ovalifolium.
villosum, Willd. Sp. Pl. i. 741 = hirsutissimum, *Gmelin*.
virens, E. Mey. ex DC. Prod. ix. 538 = curassavicum.
viridiflorum, Lehm. Asperif. 30 = Tournefortia viridiflora.
vulgare, Rota, Prosp. Fl. Berg. 63 = europæum.
Willdenowii, G. Don, Gen. Syst. iv. 356.—Bras.
zeylanicum, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 151 = paniculatum.
zeylanicum, *Lam. Encyc.* iii. 94.—Zeylan.
zeylanicum, Wall. Cat. n. 3091 = bracteatum.

HELIPTERON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = seq.

HELIPTERUM, DC. Prod. vi. 211 (1837). *COMPOSITÆ*, Benth. & Hook. f. ii. 906.
Acroclinum, A. Gray, in Hook. Kew Journ. iv. (1852) 270.

HELIPTERUM:—
Anaxeton, Gaertn. Fruct. ii. 406 (1791); Cass. in Dict. Sc. Nat. xxvi. 52 (1823); xxxiv. 87 (1825).
Anisolepis, Steetz, in Lehm. Pl. Preiss. i. 446 (1845).
Argyranthes, Neck. Elem. i. 75 (1790).
Argyrocome, Gaertn. Fruct. ii. 410 (1791); Schrank, in Denkschr. Akad. Muench. viii. (1824) 146, 168.
Asthlma, R. Br. in Bot. Reg. sub t. 538 (1820).
Casimiola, F. Muell. Fragm. iii. 189 (1863).
Damironia, Cass. Dict. Sc. Nat. lvi. 224 (1828).
Dimorpholepis, A. Gray, in Hook. Kew Journ. iv. (1852) 327.
Duttonia, F. Muell. in Linnæa, xxv. (1852) 409.
Hyalospersma, Steetz, in Lehm. Pl. Preiss. i. 476 (1845).
Monenchanthes, A. Gray, in Hook. Kew Journ. iv. (1852) 229.
Pteropogon, DC. Prod vi. 245 (1837).
Rhodanthe, Lindl. in Bot. Reg. t. 1703 (1834).
Roccardia, Neck. Elem. i. 74 (1790).
Syncarpha, DC. in Ann. Mus. Par. xvi. (1810) 206. t. 5. f. 21.
Triptilodiscus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 66.
Xeranthemum, Neck. Elem. i. 83 (1790).
Xerodanthes, Lindl. Swan River App. 58 (1839).
albicans, DC. Prod. vi. 215 = incanum.
anactinum, F. Muell. Fragm. iii. 137 = heteranthum.
anthemoides, *DC. Prod.* vi. 216.—Austral.
argyreum, Sm. in Trans. Soc. Bot. Lyon, vii. (1880) 127 = seq.
argyropsis, *DC. Prod.* vi. 213.—Afr. austr.
bicolorum, DC. l. c. 215 = incanum.
brachyrrhynchum, Sond. in Linnæa, xxv. (1852) 517 = incanum.
Calvertianum, *F. Muell. Fragm.* x. 108.—Austral.
canescens, *DC. Prod.* vi. 212.—Afr. austr.
Casuinianum, DC. l. c. 216 = Schœnia Cassiniana.
Charsleyæ, *F. Muell. Fragm.* viii. 168.—Austral.
chlamydes, F. Muell. Linnæa, xxv. (1852) 418 = floribundum.
chlorocephalum, *Benth. Fl. Australi.* iii. 641.—Austral.
citrinum, *Harv. & Sond. Fl. Cap.* iii. 250.—Afr. austr.
citrinum, Steetz, in Lehm. Pl. Preiss. i. 474 = H. Cotula.
condensatum, *F. Muell. Fragm.* iii. 136.—Austral.
corymbiflorum, *Schlecht. in Linnæa*, xxi. (1848) 648. —Austral.
corymbosum, *Benth. Fl. Australi.* iii. 649.—Austral.
Cotula, *DC. Prod.* vi. 215.—Austral.
difusum, DC. l. c. 216 = polygalifolium.
dimorpholepis, *Benth. Fl. Australi.* iii. 650.—Austral.
discoideum, A. Gray, in Hook., Kew Journ. iv. (1854) 230 = heteranthum.
Dregeanum, *DC. Prod.* vi. 214.—Afr. austr.
elegantissimum, Steud. Nom. ed. II. i. 461, in syn. = canescens.
exiguum, *F. Muell. in Trans. Vict. Inst.* i. (1855) 89. —Austral.
eximium, *DC. Prod.* vi. 213.—Afr. austr.
fascicultatum, DC. l. c. 214 = Helichrysum sesamoides.
ferrugineum, DC. l. c. 213 = Helichrysum squamosum.
ferrugineum, *Harv. & Sond. Fl. Cap.* iii. 258.—Afr. austr.
filifolium, *DC. Prod.* vi. 215.—Afr. austr.
floribundum, *DC. l. c.* 217.—Austral.
Forrestii, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 278.—Austral.
Fraseheli, *F. Muell. l. c.*, iii. (1883) 24.—Austral.
fuscescens, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 80 = Podolhecia fuscescens.
glabrum, Sond. in Linnæa, xxiii. (1850) 65 = speciosissimum.
glutinosum, Hook. in Mitch. Journ. Trop. Austral. 361 = Helichrysum glutinosum.
gnaphaloides, *DC. Prod.* vi. 213.—Afr. austr.
gracile, *Benth. Fl. Australi.* iii. 646.—Austral.
Haigii, *F. Muell. Fragm.* x. 107.—Austral.
heteranthum, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 1. 188.—Austral.
heterophyllum, DC. Prod. vi. 214 = Helichrysum sesamoides.
Humboldtianum, *DC. l. c.* 216.—Austral.
humile, DC. l. c. 214 = Helichrysum humile.

HELIPTERUM:—
hyalospermum, *F. Muell. ex Benth. Fl. Austral.* iii. 644.—Austral.
incanum, *DC. Prod.* vi. 215.—Austral.
involucratum, *F. Muell. Fragm* iii. 136.—Austral.
Kendallii, *F. Muell. l. c.* viii. 165.—Austral.
laeve, *Benth. Fl. Australi.* iii. 649.—Austral.
largiflorens, F. Muell. Fragm. iii. 135 = Humboldtianum.
longibracteatum, Steud. Nom. ed. II. i. 799, in syn. = Helichrysum humile.
Mangleisii, *F. Muell. ex Benth. Fl. Australi.* iii. 640.—Austral.
Marquesthae, *F. Muell. Fragm.* xi. 48.—Afr. austr.
monenchanthoides, F. Muell. l. c. iii. 137 = splenatum.
moschatum, *Benth. Fl. Australi.* iii. 648.—Austral.
niveum, Steetz, in Lehm. Pl. Preiss. i. 475 = obtusifolium.
phlomoides, *DC. Prod.* vi. 213.—Afr. austr.
polycephalum, *Benth. Fl. Australi.* iii. 649.—Austral.
polygalæfolium, *DC. Prod.* vi. 216.—Austral.
polyphyllum, *F. Muell. Fragm.* i. 35.—Austral.
praecox, F. Muell. in Trans. Vict. Inst. i. (1855) 58 = H. Cotula.
pterochæstum, *Benth. Fl. Australi.* iii. 648.—Austral.
punctatum, DC. Prod. vi. 216 = anthemoides.
pusillum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 80 = H. Cotula.
pygmæum, *Benth. Fl. Australi.* iii. 647.—Austral.
Pyrethrum, *Benth. l. c.* 642.—Austral.
roseum, *Benth. l. c.* 640.—Austral.
rubellum, *Benth. l. c.* 641.—Austral.
Sandfordii, Hook. Bot. Mag. t. 5850 = Humboldtianum.
semisaudum, Sch. Bip. in Flora, xxvii. (1844) 692 = speciosissimum.
semisterile, F. Muell. Fragm. ii. 157 = H. Cotula.
senecioides, DC. Prod. vi. 214 = Helichrysum sesamoides.
simplex, Steetz, in Lehm. Pl. Preiss. i. 475 = H. Cotula.
speciosissimum, *DC. Prod.* vi. 211.—Afr. austr.
splenatum, *F. Muell. ex Benth. Fl. Australi.* iii. 647.—Austral.
spinosum, Harv. & Sond. Fl. Cap. iii. 621, Index = Dicoma discanthoides.
spinulosum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 1. 197 = ferrugineum.
sterilescens, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 274.—Austral.
stipitatum, *F. Muell. ex Benth. Fl. Australi.* iii. 643.—Austral.
strictum, *Benth. Fl. Australi.* iii. 646.—Austral.
Stuartianum, Sond. in Linnæa, xxv. (1852) 518 = floribundum.
tenellum, A. Gray, in Hook. Kew Journ. iv. (1852) 230 = Helichrysum filifolium.
tenellum, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 1. 198.—Austral.
Tieckenii, *F. Muell. Fragm.* viii. 227.—Australi.
variegatum, *DC. Prod.* vi. 212.—Afr. austr.
virgatum, *DC. l. c.* 212.—Afr. austr.
Zeyheri, *Sond. in Linnæa*, xxiii. (1850) 66.—Afr. austr.

HELISANTHERA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 87 = HELIXANTHERA, Lour. = Loranthus, Linn.

HELITTOPHYLLUM, Blume, Bijdr. 659 (1825) = Melicta, Lour. (Proteac.).
japonicum, Steud. Nom. ed. II. i. 748, sphalm. = seq.
javanicum, Blume, Bijdr. 659 = Helicia robusta.
parvifolium, Zipp. ex Miq. Fl. Ind. Bat. i. t. 967 = Helicia serrata.

HELIX, Dum. ex Steud. Nom. ed. II. i. 745 (1840) = Helixr, Tourn.

HELIX, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 234 = Hedera, Tourn. (Araliac.).

HELIXANTHERA, Lour. Fl. Cochinch. 168 (1790) = Loranthus, Linn.
parasitica, Lour. l. c.—China.

HELIXIRA, Steud. Nom. ed. II. i. 745 (1840) = seq.

HELIXYRA, Salisb. in Trans. Hort. Soc. i. (1812) 328 = Maytenus, Linn. (Ilicin.).
ferox, Salisb. l. c. = M. Ilex-Aquifolium.

HELLEBORASTER, Moench, Meth. 336 (1794) = **Helleborus**, Tourn. (Rannne.).
foetidus, Moench, l. c. = Helleborus foetidus.
viridis, Moench, l. c. = Helleborus viridis.

HELLEBORINE, Tourn. ex Hall. Enum. Stirp. Helv. i. 274 (1742); Pers. Syn. ii. 512 (1807) = **Serapias**, Linn. (Orchid.).
americana, Stend. Nom. ed. I. 398 = Bletia verecunda.
Corallorrhiza, F. W. Schmidt, Fl. Boëm. i. 79 = Corallorrhiza innata.
cordata, F. W. Schmidt, l. c. 81 = Listera cordata.
cordigera, Pers. Syn. ii. 512 = S. cordigera.
latifolia, Moench, Meth. Suppl. 311 = Epipactis palustris.
leucoglottis, Stend. Nom. ed. I. 398 = S. Lingua.
Lingua, Pers. Syn. ii. 512 = S. Lingua.
longifolia, Moench, Meth. 715 = S. longifolia.
longipetala, Tenore, Fl. Neap. Prod. 58 = S. Pseudocordigera.
Nidus-avis, F. W. Schmidt, Fl. Boem. i. 78 = Neottia Nidus-avis.
ovata, F. W. Schmidt, l. c. 80 = Listera ovata.
oxyglottis, Pers. Syn. ii. 512 = S. Lingua.
palustris, Schrank, Fl. Monac. ii. 190 = Epipactis palustris.
Pseudo-cordigera, Sebast. Rom. Pl. fasc. i. 14. t. 4. f. 1 = S. Pseudo-cordigera.
rubra, Schrank, Fl. Monac. ii. 119 = Cephalanthera rubra.
succulenta, F. W. Schmidt, Fl. Boëm. i. 78 = Neottia Nidus-avis.

HELLEBOROIDES, Adans. Fam. ii. 458 (1763) = **Eranthis**, Salisb. (Ranunculac.).

HELLEBORUS, Gueldenst. Reisen, ii. 196 (1791) = **Veratrum**, Tourn. (Liliac.).
albus, Gueldenst. l. c. = V. album.

HELLEBORUS, [Tourn.] Linn. Syst. ed. I (1735). *RANUNCULACEÆ*, Benth. & Hook. f. i. 7.
HELLEBORASTER, Moench, Meth. 336 (1794).
abchasicus, *A. Br. Ind. Sem. Hort. Berol.* (1858) *App.* 34.—Reg. Caucas.
altifolius, Stend. Nom. ed. II. i. 746 = niger.
angustifolius, Host, Fl. Austr. ii. 90 = viridis.
angustifolius, Walp. Rep. i. 481, sphalm. = *argutifolius*.
antiquorum, *A. Br. Ind. Sem. Hort. Berol.* (1858) *App.* 33.—Ithynia.
argutifolius, Viv. Fl. Cors. Sp. Nov. 8 = lividus.
atropurpureus, Schult. Oestr. Fl. ed. II. ii. 103 = seq.
atrorubens, Waldst. & Kit. Pl. Rar. Hung. iii. 301. t. 273 = odorus.
Baumgartenï, *Kov. ex Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 4.—Transsylv.
bengisicus, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 225 = foetidus.
Bocconi, Schur, Enum. Pl. Transs. 27 = Baumgartenï.
Bocconi, Tenore, Prod. Fl. Nap. App. iv. 26; ex ej. Fl. Nap. iv. Syll. 79 = viridis.
breviscalis, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 225 = viridis.
caucasicus, *A. Br. Ind. Sem. Hort. Berol.* (1858) *App.* 34.—Reg. Caucas.
colchicus, *Regel, Gartenfl.* (1856) 299.—Mingrel.
corsicus, Willd. Enum. Hort. Berol. Suppl. 40 = lividus.
cupreus, Host, Fl. Austr. ii. 87 = odorus.
cyclophyllus, *Boiss. Fl. Orient.* i. 61.—Graec.
dubius, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 225 = foetidus.
dumetorum, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 538 = viridis.
foetidus, Georgi, Beschr. Russ. Reich. 1071 = orientalis.
foetidus, *Linn. Sp. Pl.* 558.—Europ.
fumarioides, Lam. Encyc. iii. 99 = Isopyrum fumarioides.
grandiflorus, *C. Koch, ex Rupr. Fl. Cauc.* 32.—Reg. Caucas.
grandiflorus, Salisb. Prod. 374 = niger.
graveolens, Host, Fl. Austr. ii. 89 = odorus.
guttatus, *A. Br. Ind. Sem. Hort. Berol.* (1858) *App.* 33.—Reg. Caucas.
hemisphæricus, *Wendroc. ex Steud. Nom.* ed. II. i. 746.—Alsat.
hirsutus, Host, Fl. Austr. ii. 87 = odorus.
hybridus, Host, Fl. Austr. ii. 87 = odorus.
hyemalis, Linn. Sp. Pl. 557 = Eranthis hyemalis.
Ibericus, *Stev. ex Rupr. Fl. Cauc.* 287.—Reg. Caucas. Intermedius, *Guss. Pl. Rar. Sic.* 224. t. 4.—Ital.
intermedius, Host, Fl. Austr. ii. 88 = latifolius.
involucratus, Stokes, Bot. Mat. Med. iii. 271 = Eranthis hyemalis.
latifolius, *Mill. Gard. Dict.* ed. VIII. n. 6.—Europ. or.
laxus, Host, Fl. Austr. ii. 89 = viridis.
lividus, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 272.—Corsica.
monanthos, Moench, Meth. 313 = Eranthis hyemalis.
multifidus, Vis. in Flora, xli. (1829) t. Erg. 18 = viridis.
nemorosus, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 225, nomen = foetidus.
niger, *Linn. Sp. Pl.* 558.—Europ.
nivalis, Siev. ex Ledeb. Fl. Koss. i. 735 = Eranthis sibirica.
occidentalis, Rent. Cat. Grain. (1866); ex Bull. Soc. Bot. Fr. xvi. (1869) Bibl. 58 = viridis.
odorus, *Waldst. & Kit. ex Willd. Enum. Hort. Berol.* 592.—Hungar.
officinalis, Salisb. in Trans. Linn. Soc. viii. (1807) 305 = orientalis.
olympicus, *Lindl. Bot. Reg.* (1841) *Misc.* 54.—Bithynia.
orientalis, Garn. ex Nym. Consp. 17 = cyclophyllus.
orientalis, *Lam. Encyc.* iii. 96.—Graecia; An. Min.
pallidus, Host, Fl. Austr. ii. 90 = viridis.
ponticus, A. Br. in Ind. Sem. Hort. Berol. (1858) App. 34 = orientalis.
porphyromelas, A. Br. & Bouché, l. c. (1861) 13 = colchicus.
pumilus, Sallsb. Prod. 374 = Coptis trifolia.
purpurascens, Fusa, Pl. Transs. 29, partim = Baumgartenï.
purpurascens, Waldst. & Kit. Pl. Rar. Hung. ii. 105. t. 101 = odorus.
ranunculinus, Sm. Ic. fasc. ii. 37 = Trollius patulus.
rhodanicus, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 225, nomen = foetidus.
sibiricus, Spreng. Syst. ii. 658 = Eranthis sibirica.
thalictroides, Lam. Fl. Fr. iii. 315 = Isopyrum thalictroides.
tinctus, Stev. ex Rupr. Fl. Cauc. 287 = colchicus.
trifolius, Linn. Sp. Pl. 558 = Coptis trifolia.
trifolius, Mill. Gard. Dict. ed. VIII. n. 4 = lividus.
trilobus, Lam. Encyc. iii. 98 = Coptis trifolia.
triphyllus, Lam. l. c. 97 = lividus.
vaginatus, Kit. ex Stend. Nom. ed. II. i. 746 = viridis.
vernalis, D. Don, ex Loud. Hort. Brit. Suppl. ii. 640 = latifolius!
vesicarius, *Auch. ex Boiss. in Ann. Sc. Nat. Sér.* II. (1841) 357.—Syria.
viridiflorus, Stokes, Bot. Mat. Med. iii. 274 = viridis.
viridis, Bieb. Fl. Taur. Cauc. ii. 80 = caucasicus.
viridis, *Linn. Sp. Pl.* 558.—Europ.

HELLENIA, Retz. Obs. vi. 18 (1791) = **Costus**, Linn. (Scitamin.).
grandiflora, Retz. l. c. = C. speciosus.

HELLENIA, Willd. Sp. Pl. i. 4 (1797) = **Alpinia**, Linn. (Scitamin.).
abnormis, Lindl. in Trans. Hort. Soc. vii. (1830) 60 = A. chinensis!
alba, Willd. Sp. Pl. 5 = A. alba.
Allughas, Willd. l. c. 4 = A. Allughas.
aquatica, Willd. l. c. 5 = A. aquatica.
arctiflora, F. Muell. Fragm. viii. 35 = A. arctiflora.
bracteata, Blume, Enum. Pl. Jav. 60 = A. alata.
chinensis, Willd. Sp. Pl. i. 5 = A. chinensis.
corvulae, R. Br. Prod. i. 308 = A. caerulea.
gracilis, Hasskarl. in Flora, xlvii. (1864) 19 = A. gracilis.
melanocarpa, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxiv. (1862) 238.—Sumatra.
parviflora, Horan. Prod. Scitam. 36 = A. parviflora.
pubiflora, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 235.—N. Guin.
rubra, Mig. Fl. Ind. Bat. iii. 604 = A. rubra.
rufa, Presl, Rel. Haenk. 114. t. 21.—Ins. Philipp.
scabra, Blume, Enum. Pl. Jav. 60.—Java.

HELLERIA, Fourn. ex Benth. & Hook. f. Gen. iii. 1199 (1883) = **Festuca**, Linn. (Gramin.).
livida, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 582 = F. livida.

HELLERIA, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 89. t. 7 = **Vantanea**, Aubl. (Humiriac.).
obovata, Mart. l. c. 89 = V. obovata.
ovalifolia, A. Juss. in St. Hil. Fl. Bras. Mer. ii. 61 = V. obovata.

HELLMUTHIA, Stend. in Flora, xxxiii. (1850) 229, nomen; et Syn. Pl. Cyp. 90 (1855) = **Fiorinia**, Schrad. (Cyperac.).
restioides, Stend. ll. cc. = F. membranacea?

HELMENTIA, Janme St. Hil. Expos. i. 376 (1805) = HELMINTIA, Juss. = **Picris**, Linn. (Compos.).

HELMHOLTZIA, F. Muell. Fragm. v. 202 (1866). *PHILYDRACEÆ*, Benth. & Hook. f. iii. 841.
acorifolia, *F. Muell. l. c.* 203.—Austral.
glaberrima, *Caruel, in DC. Monog. Phan.* iii. 6.—Ins. Pacif.

HELMIA, Kunth, Enum. Pl. v. 414 (1850) = **Dioscorea**, Linn.
adenocarpa, Kunth, l. c. 425 = D. adenocarpa.
anomala, Kunth, l. c. 427 = D. anomala.
aspera, Kunth, l. c. 429 = D. aspera.
brachycarpa, Kunth, l. c. 413 = D. brachycarpa.
bulbifera, Kunth, l. c. 435 = D. sativa.
campestris, Kunth, l. c. 425 = D. campestris.
comargyrioea, Kunth, l. c. 426.—Guian.
convolvulacea, Kunth, l. c. 415 = D. convolvulacea
coriacea, Kunth, l. c. 422 = D. coriacea.
cuspidata, Kunth, l. c. 438 = D. cuspidata.
daemona, Kunth, l. c. 439 = D. daemona.
Dregeana, Kunth, l. c. 437.—Afr. austr.
dumetorum, Kunth, l. c. 436 = D. triphylla.
Ehrenbergiana, Hemsl. Biol. Centr. Am. Bot. iii. 360, sphalm. = seq.
Ehrenbergiana, Kunth, Enum. Pl. v. 433 = D. remotiflora.
fracta, Kunth, l. c. 420 = D fracta.
furcata, Kunth, l. c. 419 = D. furcata.
galipaenensis, Kunth, l. c. 417 = D. galipaenensis.
Griesbachii, Kunth, l. c. 432.—Bras.
hirsuta, Kunth, l. c. 436 = D. hirsuta, *Blume.*
Lacerdari, Kunth, l. c. 427 = D. Lacerdari.
monadelpha, Kunth, l. c. 421 = D. monodelpha.
Moritziana, Kunth, l. c. 423.—N. Granat.
multiflora, Kunth, l. c. 431 = D. multiflora, *Mart.*
pilosiuscula, Kunth, l. c. 434 = D. piloisiscula.
pallentackya, Kunth, l. c. 439.—Cuba.
racemosa, Klotzsch, in Otto & Dietr. Allg. Gartenz. xix. (1851) 393.—Am. austr.
scabra, Kunth, Enum. Pl. v. 430 = D. scabra.
Schomburgkiana, Kunth, l. c. 424 = D. Schomburgkiana!
syringaefolia, Kunth, l. c. 423 = D. syringaefolia!
tomentosa, Kunth, l. c. 429 = D. tomentosa, *Korn.*
trifoliata, Kunth, l. c. 417 = D. trifoliata.

HELMINTA, Willd. Sp. Pl. iii. 1607 (1800), sphalm. = seq.

HELMINTHIA, DC. Fl. Fr. iv. 57 (1805) = seq.

HELMINTHIA, Juss. Gen. 170 (1789) (Helmintia) = **Picris**, Linn. (Compos.).
aculeata, DC. Prod. vii. 132 = P. aculeata.
asplenioides, DC. l. c. = P. asplenioides.
Balansae, Cos. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 282, nomen; et in Bull. Soc. Bot. Fr. iii. (1866) 764.—Orient.
callosa, Pomel, Nouv. Mat. Fl. Atl. 14 (= *Dechera callosa*).—Alger.
comosa, Boiss. Elench. 62.—Hispan.
dichotoma, Dulac, Fl. Hautes-Pyr. 496 = P. echioides.
echioides, Gaertn. Fruct. ii. 368. t. 159. f. 2 = P. echioides.
glomerata, Pomel, Nouv. Mat. Fl. Atl. 13 (= *Dechera glomerata*).—Alger.
hamulosa, Trevir. in Nov. Act. Nat. Cur. xiii. (1826) 195 = P. echioides.

HELMINTHIA:—
hastanica, Welw. ex Schlecht. in Linnaea, xxvii. (1856) 315.—Lusitan.
montana, Pomel, Nouv. Mat. Fl. Atl. 14 (= *Dechera montana*).—Algar.
mucronata, Terrac. Fl. Valt. Syn. 111; et ex Nouv. Giorn. Bot. Ital. iii. (1871) 196 = P. scaiosta.
pratensis, Chev. Fl. Par. iii. 362 = P. schioides.
racemosa, Pomel, Nouv. Mat. Fl. Atl. 18 (= *Dechera racemosa*).—Algar.
rigida, Schult. ex Steud. Nom. ed. I. 806 = P. schioides.
spinosa, DC. Fl. Fr. iv. 58 = P. schioides.
Sprengeriana, Gaertn. ex DC. Prod. vii. 138 = P. Sprengeriana.
tuberculata, Moench, Meth. 589 = P. schioides.

HELMINTHION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = prænc.

HELMINTHOCARPUM, A. Rich. Tent. Fl. Abyss. i. 200. t. 36 (1847)). *LEGUMINOSÆ*, Benth. & Hook. f. i. 469.
abyssinicum, *A. Rich. l. c.*—Afr. trop.

HELMINTHOSPERMUM, Thw. in Hook. Kew Journ. vi. (1854) 301. t. 9 = Gironniera, Gaudich. (Urtic.).
globrescens, Thw. ex Pisach. in DC. Prod. xvii. 209 = G. reticulata.
scabridum, Thw. in Hook. Kew Journ. vi. (1854) 302 = G. subæqualis.

HELMINTHOTECA, Vaill. ex Juss. Gen. 170 (1789) = seq.

HELMINTIA, Juss. Gen. 170 (1789) = HELMINTHIA, Juss. = Picris, Linn. (Compos.).

HELMONTIA, Cogn. in Bull. Soc. Bot. Belg. xiv. (1875) 239. *CUCURBITACEÆ*.
leptantha, *Cogn. Diagn. Cucurb. fasc.* i. 20.—Bras. simplicifolia, *Cogn. l. c.* 44.—Bras.

HELODEA, Reichb. Nom. 211 (1841) = Elodea, Michx. (Hydrochar.).

HELODES, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 54 = prænc.

HELODIUM, Dum. Fl. Belg. 77 (1827) = Aptum, Tourn. (Umbell.).
inundatum, Dum. l. c. = A. inundatum.
nodiflorum, Dum. l. c. = A. nodiflorum.
repens, Dum. l. c. = A. nodiflorum.

HELOGYNE, Benth. Bot. Voy. Sulph. 20. t. 14 (1844) = Hofmeisteria, Walp. (Compos.).
fasciculata, Benth. l. c. = Hofmeisteria fasciculata.
urenifolia, Walp. Rep. iv. 457, ex Ind. sphalm. = Hofmeisteria urenifolia.

HELOGYNE, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 449. *COMPOSITÆ*, Benth. & Hook. f. ii. 289.
spaloides, *Nutt. l. c.*—Peruv.

HELONIAS, Linn. Dis. Chen. 17 (1751); Amoen. Acad. iii. 12 (1756). *LILIACEÆ*, Benth. & Hook. f. iii. 827.
Abalum, Adans. Fam. ii. 47 (1763).
Gomphostylis, Rafin. Fl. Tellur. ii. 80 (1836).
angustifolia, Michx. Fl. Bor. Am. i. 213 = Zygadenus angustifolius.
angustifolia, Nutt. in Trans. Am. Phil. Soc. v. (1837) 154 = Zygadenus Nuttallii.
anthericoides, Hoppe, ex Schult. *f. Syst.* vii. 1575 = Tofieldia calyculata.
asphodeloides, Linn. Sp. Pl. ed. II. 485 = Xerophyllum asphodeloides.
borealis, Willd. Sp. Pl. ii. 274 = Tofieldia calyculata.
bracteata, Sims, in Bot. Mag. t. 1703 = Zygadenus bracteatus.
bullata, *Linn. Sp. Pl.* 342.—Am. bor.
dioica, Pursh, Fl. Am. Sept. i. 248 = Chamælirium carolinianum.
dubia, Michx. Fl. Bor. Am. i. 213 = Schœnocaulon gracile.
erythrosperma, Michx. l. c. 213 = Zygadenus mucronicus.
glaberrima, Link, Enum. Hort. Berol. i. 248 = Zygadenus glaberrimus.
glaberrima, Sims, Bot. Mag. t. 1680 = Zygadenus Fremontii.
graminea, Ell. Sketch, i. 432 = Zygadenus leimanthoides.
graminea, [Ker-Gawl.] in Bot. Mag. t. 1899; Pursh, Fl. Am. Sept. ii. 738 = Stenanthium angustifolium.
gulseënsis, *Thunb. in Schum. & Thonn Beskr. Guin. Pl.* 182.—Afr. trop.
japonica, Schult. f. Syst. vii. 1567 = Chionographis japonica.
lanta, Ker-Gawl. in Bot. Mag. t. 808 = Zygadenus muscætoxicus.
lanceolata, Sims, Bot. Mag. sub t. 747, sphalm. = seq.
latifolia, Michx. Fl. Bor. Am. i. 213 = bullata.
longifolia, *Pitch. ex Schult. f. Syst.* vii. 1567, nomen. —Hab.?
lutea, [Ker-Gawl.] in Bot. Mag. t. 1062 = Chamælirium carolinianum.
minuta, Linn. Mant. ii. 225 = Hypoxis minuta.
nigra, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 164 = Veratrum nigrum.
officinalis, D. Don, in Edinb. N. Phil. Journ. (1833) 38 = Schœnocaulon officinale.
paniculata, Nutt. in Journ. Acad. Philad. vii. (1834) 57 = Zygadenus Nuttallii.
pumila, Jacq. Coll. ii. 260 = Chamælirium carolinianum.
scapigera, Stokes, Bot. Mat. Med. ii. 338 = bullata.
striata, *Rafin. Atl. Journ.* 147 (Quid?).—Am. bor.
tenax, Pursh, Fl. Am. Sept. i. 248 = Xerophyllum asphodeloides.
tenuifolia, Salisb. Prod. 250 = Xerophyllum asphodeloides.
tomentosa, Muhl. Cat. 87; et ex Schult. f. Syst. vii. 989 = Lophiola aurea.
viridescens, H. B. & K. Nov. Gen. et Sp. i. 267 = Zygadenus mexicanus.
virginica, Sims, Bot. Mag. t. 985 = Melanthium virginicum.
viridis, [Ker-Gawl.] in Bot. Mag. t. 1096 = Veratrum viride.

HELONIOPSIS, A. Gray, in Mem. Am. Acad. Ser. II. iv. (1855–59) 416. *LILIACEÆ*, Benth. & Hook. f. iii. 827.
Scoliokia, Miq. in Ann. Mus. Bot. Lugd. Bat. iii. 144 (1867).
breviscapa, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 486.—Japon.
grandiflora, Franch. & Sav. Enum. Pl. Jap. ii. 88, 539 = breviscapa.
japonica, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 487.—Japon.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 146, ex parte = breviscapa.
pauciflora, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 416.—Japon.
pauciflora, Maxim. in Bull. Acad. Pétersb. xi. (1867) 486 = breviscapa.
umbellata, *Baker, in Journ. Bot.* xii. (1874) 278.—Ins. Formosa.

HELOPHYLLUM, Hook. f. Handb. N. Zeal. Fl. 167 (1864) = Phyllachne, Forst. (Stylid.).
alsinifolium, Hook. f. l. c. = T. clavigera.
Colensoi, Hook. f. l. c. 168 = P. Colensoi.
rubrum, Hook. f. l. c.—N. Zel.

HELOPHYTUM, Eckl. & Zeyh. Enum. 286 (1836) = Tillæa, Mich. (Crassulac.).
filiforme, Eckl. & Zeyh. l. c. 289 = T. capensis.
fluitans, Eckl. & Zeyh. l. c. = T. capensis.
imbricatum, Eckl. & Zeyh. l. c. = T. capensis.
natans, Eckl. & Zeyh. l. c. 289 = T. capensis.
reflexum, Eckl. & Zeyh. l. c. 289 = T. capensis.

HELOPUS, Tris. Fund. Agrost. 108 (1820) = Brizochloa, H. B. & K. (Gramin.).
acrotrichus, Steud. Nom. ed. II. i. 747.—Ind. or.; Arab.
acuminatus, Steud. l. c. = E. polystachya.
annulatus, Hochst. ex Steud. Syn. Pl. Gram. 190 = *nubicus*.
arundinaceus, Nees, Agrost. Bras. 17 = E. annulatus.
barbatus, Nees, ex Steud. Nom. ed. II. i. 747 = *acrotrichus*.
barbatus, Tris. in Spreng. Neue Entdeck. ii. 46.—Ins. Borbon.

HELOPUS:—
bellodes, Hochst. ex Steud. Syn. Pl. Gram. 100.—Abyssin.
brachystachys, Trin. Sp. Gram. xxiv. t. 277 = E. distachys.
cognatus, Steud. Syn. Pl. Gram. 101 = E. polystachya.
glabrous, Steud. l. c. 98.—Japon.
gracilis, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 464 = E. distachys.
grandiflorus, Trin. Sp. Gram. xxiv. t. 278 = E. grandiflorus.
javanicus, Steud. Syn. Pl. Gram. 101.—Java.
junceus, C. Muell. in Bot. Zeit. xix. (1861) 314.—Am. bor.
Kunthii, Trin. ex Steud. Nom. ed. II. i. 747 = E. Kunthii.
laevis, Trin. in Spreng. Neue Entdeck. ii. 49 = E. annulatus.
Meyeni, Steud. Nom. ed. II. i. 747.—Peruv.
mollis, C. Muell. in Bot. Zeit. xix. (1861) 314.—Am. bor.
nubicus, Steud. Syn. Pl. Gram. 100.—Nubia.
obtusus, Steud. Nom. ed. II. i. 747.—Peruv.
pilosus, Trin. Fund. Agrost. 104 = E. polystachya.
polystachyus, Trin. ex Steud. Nom. ed. II. i. 747 = E. polystachya.
punctatus, Nees, Agrost. Bras. 18 = E. polystachya.
sanguinalis, Nees, l. c. 17 = Digitaria sanguinalis.
succinctus, Trin. ex Steud. Syn. Pl. Gram. 100.—Reg. Casp.
trichopus, Hochst. ex Steud. Syn. Pl. Gram. 100 = Panicum javanicum.
tridentatus, Trin. ex Steud. Nom. ed. II. i. 747; Syn. Pl. Gram. 100.—Am. austr.
villosus, Nees, ex Steud. ll. cc.—E. villosus.

HELOSCHIADIUM, Marsson, Fl. Werml. 88 (1859) = seq.

HELOSCIA, Dum. Anal. Fam. 34 (1829) = seq.

HELOSCIADIUM, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 135 = Aptum, Tourn. (Umbellif.).
australe, Bunge, in Lehm. Pl. Preiss. i. 294 = A. prostratum.
bulbosum, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 136 = Carum inundatum.
californicum, Hook. & Arn. Bot. Beech. Voy. 142 = Oenanthe sarmentosa.
capense, Walp. Rep. ii. 892 = Carum capense.
crassipedatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = seq.
crassipes, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 135 = A. crassipes.
deserticola, Phil. Fl. Atac. 26.—Chili.
Falcaria, Hegetschw. Fl. Schw. 262 = Falcaria vulgaris.
gracile, Clos, in C. Gay, Fl. Chil. iii. 124. t. 81. f. 2 = A. gracile.
Heynianum, DC. Prod. iv. 106 = Pimpinella Heyniana.
hybridum, Mérat, ex F. Schultz, in Bonplandia, ii. (1854) 236 = A. nodiflorum.
intermedium, DC. Prod. iv. 105 = Carum inundatum.
intermedium, Tenore, ex Nym. Consp. 300 = A. nodiflorum.
inundatum, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 136 = A. inundatum.
laciniatum, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 495.—Peruv.
lateriflorum, Hook. & Arn. Bot. Beech. Voy. 26 = *ranunculifolium*.
lateriflorum, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 136 = A. Ammi.
leptophyllum, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 495 = A. leptophyllum.
nodiflorum, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 136 = A. nodiflorum.
oppositifolium, Reum. Kwit. Slov. (1858) 178.—Hungar.
palustrum, F. Schultz, ex Nym. Consp. 300 = A. repens.
prostratum, Bunge, in Lehm. Pl. Preiss. i. 294 = A. procumbens.
pulcherrimum, DC. Prod. iv. 106 = Pimpinella diversifolia.
ranunculifolium, DC. l. c. 106.—Chili.
repens, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 136 = A. repens.
rupestre, Syme, ex F. Schultz, in Bonplandia, ii. (1854) 237 = A. nodiflorum.
Rœtz, DC. Prod. iv. 106 = A. graveolens.

HELOSCIADIUM :—
rutaceum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = A. graveolens.
simonae, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 324 = Pimpinella simensis.
stoloniforum, Nym. Syll. 160 = A. nodiflorum.
tenerum, DC. Prod. iv. 106 = Pimpinella tenera.
trifoliatum, DC. l. c. 106 = Pimpinella diversifolia.
Wallichianum, Miq. ex Hohen. in Bot. Zeit. vii. (1849) 775 = Pimpinella monoica.

HELOSERIS, Reichb. ex Steud. Nom. ed. II. i. 748 (1840) = **Senecio**, Tourn. (Compos.).
palustris, Reichb. ex Nym. Consp. 352 = S. palustris.

HELOSIS, Rich. in Mém. Mus. Par. viii. (1822) 416. t. 20. *BALANOPHOREAE*, Benth. & Hook. f. iii. 397.
CALDASIA, Mutis, ex Caldas, in Seman. Nuev. Gran. ii. (1810) 26.
LATRAPOPHILA, Lasnér. Seeman. ex A. St. Hil. in Ann. Sc. Nat. Sér. II. vii. (1837) 92.
aquatica, Mutis, ex Hook. f. in Trans. Linn. Soc. xxii. (1856) 59 = mexicana.
brasiliensis, Schott & Endl. Melet. 12 = guyannensis.
cayennensis, Spreng. Syst. iii. 755 = guyannensis.
guianensis, Aubl. = seq.
GUYANENSIS, *Rich. in Mém. Mus. Par.* viii. (1822) 416. t. 20.—Am. trop.
jamaicensis, Rich. l. c. 425 = Scyballum jamaicense.
mexicana, *Hchm. in Fork. Vidensk. Skand. Nat.* iv. (1844) 181.—Mexic.

HELOSPORA, Jacq. in Trans. Linn. Soc. xiv. (1824) 127. t. 4. f. 3 = Pimpinella, Rumph. (Rubiac.).
flavescens, Jacq. l. c. = T. Jamboselle.

HELOTHRIX, Nees, in Ann. Nat. Hist. Sér. I. vi. (1841) 48 = **Schoenus**, Linn. (Cyperac.).
pusilla, Nees, l. c. = S. axillaris.

HELVINGIA, Adans. Fam. ii. 345 (1763) = **Laetia**, Linn. (Bixin.).

HELWINGIA, Willd. Sp. Pl. iv. 716 (1805). *ARALIACEAE*, Benth. & Hook. f. i. 939.
hinalaica, *Hook. f. & Thoms. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 736.—Reg. Himal.
japonica, Morr. & Decne. in Bull. Acad. Brux. iii. (1836) 170 = ruscifolia.
polyflora, Spreng. Pugill. ii. 89 (non generis).—Bengal.
ruscifiora, *Willd. Sp. Pl.* iv. 716.—Japon.
ruscifolia, Steud. Nom. ed. I. 399 = prsec.

HELXINE, Linn. Syst. ed. I. (1735) = **Polygonum**, Tourn.
arifolia, Raffn. Fl. Tellur. iii. 94 = P. arifolium.
cilinode, Raffn. l. c. 10 = P. cilinode.
Convolvulus, Raffn. l. c. = P. Convolvulus.
dumetorum, Raffn. l. c. = P. dumetorum.
multiflorum, Raffn. l. c. = P. multiflorum.
sagittatum, Raffn. l. c. 10; 95 = P. sagittatum.
scandens, Raffn. l. c. 10 = P. scandens.

HELXINE, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 384. *URTICACEAE*, Benth. & Hook. f. iii. 398.
SOLEIROLIA, Gaudich. in Freyc. Voy. Bot. 504 (1826).
Soleirolii, *Req. l. c.*—Cors. ; Sardin.

HELYGIA, Blume, Bijdr. 1043 (1826) = **Parsonsia**, R. Br. (Apocyn.).
javanica, Blume, l. c. = P. spiralis.

HEMARTHRIA, R. Br. Prod. 207 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1181.
LOXOLAMARIA, Beauv. Agrost. 108. t. 21 (1819).
capensis, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. ii. (1833) 348 = compressa.
candidissima, Steud. Syn. Pl. Gram. 359 = compressa.
compressa, *R. Br. Prod.* 207.—Reg. calid.
corcondadensis, Steud. Syn. Pl. Gram. 358 = compressa.
fasciculata, Kunth, Rév. Gram. i. 158 = compressa.
foliata, Steud. Syn. Pl. Gram. 359 = compressa.
fragilissima, Steud. l. c. = compressa.
Hookeriana, Steud. l. c. 358 = compressa.

HEMARTHRIA :—
perforata, Kunth, Rév. Gram. i. 158 = Ophiurus perforatus.
persuiuna, Steud. Nom. ed. II. i. 748 = compressa.
protensa, *Steud. Syn. Pl. Gram.* 359.—Ind. or.
rugosa, Kunth, Rév. Gram. i. 158 = Rottboellia rugosa.
uncinata, R. Br. Prod. i. 207 = compressa.
vaginata, Buese, in Miq. Pl. Jungh. 354 = protensa.

HEMATODES, Raffn. Fl. Tellur. iii 92 (1836) = **Salvia**, Tourn. (Labiat.).
indica, Raffn. l. c. = S. indica.

HEMATOPHYLA, Raffn. Sylva Tellur. 71 (1838) = **Psychotria**, Linn. (Gesneracj.).
villosa, Raffn. l. c.—Hab. ?

HEMECYCLIA, Wight & Arn. in Edinb. N. Phil. Journ. xiv. (Apr. 1833) 297, sphalm.?—Cf. **Hemicyclia**, Wight & Arn.

HEMENAEA, Scop. Introd. 296 (1777), sphalm. = **Hymenaea**, Linn. (Legumin.).

HEMEROCALLIS, Murr. Syst. ed. XIV (1784) 830 = seq.

HEMEROCALLIS, Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 778.
alba, Andr. Bot. Rep. t. 194 = Funkia subcordata.
albo-marginata, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 366 = Funkia lancolata.
caerulea, Andr. Bot. Rep. t. 6 = Funkia ovata.
chinensis, Hort. ex Steud. Nom. ed. II. i. 748 = Funkia Sieboldiana.
cordata, Cav. Desc. 124 = Funkia subcordata.
cordata, Thunb. Fl. Jap. 143 = Lilium cordifolium.
cordifolia, Salisb. in Trans. Linn. Soc. viii. (1807) 11 = Lilium cordifolium.
crocea, Lam. Fl. Fr. iii. 267 = fulva.
cuculiata, Hort. ex Steud. Nom. ed. II. i. 748 = Sieboldiana.
disticha, Donn, Hort. Cantab. ed. VI. 98 = fulva.
Dumortierii, *Morr. in Hortic. Belge*, ii. (1844) 195. t. 48.—Japon.
flava, *Linn. Sp. Pl.* ed. II. 462.—Europ. ; As. temp.
flava, Suter, Fl. Helv. i. 206 = fulva.
fulva, *Linn. Sp. Pl.* ed. II. 462.—Europ. ; As. temp.
graminea, Andr. Bot. Rep. t. 244 = minor.
graminea, Schlecht. in Halle, Abh. I. iii. (1824) 15 = Dumortierii.
graminifolia, Schlecht. l. c. = minor.
japonica, Thunb. Fl. Jap. 143 = Funkia subcordata.
Kwanso, Hort. ex Regel, Gartenfl. (1866) 66. t. 500 = fulva.
lancifolia, Thunb. in Trans. Linn. Soc. ii. (1794) 335 = Funkia lancifolia.
Liliastrum, Linn. Sp. Pl. 324 = Paradisia Liliastrum.
Liliastrum, Linn. l. c. = flava.
longituba, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 152 = fulva.
lutea, Gaertn. Fruct. ii. 18. t. 88 = flava.
Middendorffii, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 94).—Reg. Amur.
minor, *Mill. Gard. Dict.* ed. VIII. n. 2.—As. bor.
plantaginea, Lam. Encyc. iii. 108 = Funkia subcordata.
pumila, Salisb. Gen. Pl. Fragm. 81 = minor.
rutilans, Hort. ex Baker, in Journ. Linn. Soc. xi. (1871) 359 = Dumortierii.
Sieboldiana, Lodd. Bot. Cab. t. 1869 = Funkia Sieboldiana.
Sieboldii, Hort. ex Baker, in Journ. Linn. Soc. xi. (1871) 359 = Dumortierii.
speciosa, Sweet, Hort. Brit. ed. II. 516, sphalm. = Hymenocallis speciosa.
undulata, Siebold, ex Otto & Dietr. Allg. Gartenz. i. (1833) 119 = Funkia lancifolia.

HEMESOTRIA, Raffn. in Ann. Gén. Sc. Phys. vi. (1820) 88 = **Astrephia**, Duf. (Valerian.).

HEMIACHYRIS, DC. Prod. v. 313 (1836) = **Guttiferae**, Lag. (Compos.).
glutinosa, St. Schaz. in Linnaea, xix. (1847) 784 = G. glutinosa.
texana, DC. Prod. v. 314 = G. texana.

HEMIADELPHIS, Nees, in Wall. Pl. As. Rar. iii. 75 (1832) = **Hygrophila**, R. Br. (Acanthac.).
polysperma, Nees, l. c. 80 = Hygrophila polysperma.

HEMIAGRAPHIS, T. Anders. in Journ. Linn. Soc. vii. (1864) 113 = **Hemigraphis**, Nees (Acanthac.).

HEMIAMBROSIA, Delpino, Studi Sopra Artemis. 57 (1871) = **Franseria**, Cav. (Compos.).
heterocephala, Delpino, l. c. = Ambrosia pumila.

HEMIANDRA, R. Br. Prod. 509 (1810). *LABIATAE*, Benth. & Hook. f. ii. 1218.
brevifolia, Benth. in Enum. Pl. Hueg. 79 = pungens.
candidissima, Rich. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 29 = Trembleya Lychnitis.
emarginata, Lindl. in Bot. Reg. (1841) Misc. 72 = pungens.
glabra, Benth. Lab. Gen. et Sp. 287 = pungens.
hirsuta, Benth. in Enum. Pl. Hueg. 79 = pungens.
incana, *Bartl. in Lehm. Pl. Preiss.* i. 257.—Austral.
juniperina, Bartl. l. c. 355 = pungens.
lstantha, *Benth. Fl. Austral.* v. 110.—Austral.
linearis, Benth. in Enum. Pl. Hueg. 79 = pungens.
longifolia, Bartl. in Lehm. Pl. Preiss. i. 356 = pungens.
pungens, *R. Br. Prod.* 509.—Austral.
rupestris, Hueg. Bot. Arch. t. 4 = pungens.

HEMIANDRINA, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 171. t. 28 = **Agelaea**, Soland. (Connarac.).
borneensis, Hook. f. l. c. = A. vestita.

HEMIANTHUS, Nutt. in Journ. Acad. Philad. i. (1817) 119. t. 6. f. 2 = **Micranthemum**, Michx. (Scrophul.).
adenander, Wright, ex Griseb. Cat. Pl. Cub. 185.—Cuba.
callitrichoides, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 522 = M. callitrichoides.
elatinoides, Griseb. Cat. Pl. Cub. 185 = M. elatinoides.
erosus, Griseb. l. c. 184.—Cuba.
micranthemoides, Nutt. in Journ. Acad. Philad. i. (1817) 119. t. 6. f. 2 = M. Nuttallii.
multiflorus, Wright, ex Griseb. Cat. Pl. Cub. 185.—Cuba.
punctatus, Wright, ex Griseb. l. c. 184.—Cuba.
reflexus, Wright, ex Griseb. l. c. 186 = M. reflexum.
tristeolus, Wright, ex Griseb. l. c. 185.—Cuba.

HEMIARRHENA, Benth. Fl. Austral. iv. 518 (1869). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 974.
plantaginea, *Benth. l. c.*—Austral.

HEMIBROMUS, Steud. Syn. Pl. Gram. 317 (1855) = **Glyceria**, R. Br. (Gramin.).
japonicus, Steud. l. c. = G. japonica.

HEMICAREX, Benth. in Journ. Linn. Soc. xviii. (1881) 367 = **Kobresia**, Willd. et **Schoenoxiphium**, Nees (Cyperac.).
curvirostris, C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 384 (Kobresiae sp.).—Reg. Himal.
Eckloni, Benth.—Cf. l. c. xviii. (1881) 367 = Schoenoxiphium Ecklonii.
filicina, C. B. Clarke, l. c. xx. (1883) 384 (Kobresiae sp.).—Reg. Himal.
Hookeri, Benth. l. c. xviii. (1881) 367 = Kobresia Hookeri.
laxa, Benth. l. c. 367 = Kobresia laxa.
linearis, Benth. l. c. = Carex linearis, Kobresia trinervis.
Meyeriana, Benth.—Cf. l. c. = Schoenoxiphium capense.
pygmaea, C. B. Clarke, l. c. xx. (1883) 383 (Kobresiae sp.).—Reg. Himal.
reticulariz, Benth. l. c. xviii. (1881) 367 = Kobresia reticularis.
Sickenanniana, Benth.—Cf. l. c. = Schoenoxiphium Sickenannianum.
Thunbergii, Benth.—Cf. l. c. = Schoenoxiphium Thunbergii.
trinervis, Benth. & Hook. f. ex C. B. Clarke, l. c. xx. (1883) 383 = Carex linearis, Kobresia trinervis.

HEMICARPHA, Nees & Arn. in Edinb. N. Phil. Journ. xvii. (1834) 268 = **Scirpus**, Tourn. (Cyperac.).
Drummondii, Nees, in Mart. Fl. Bras. ii. 1. 62 = S. reticranthus.

HEMICARPHA :—
Isolepis, Nees, ex Arn. in Edinb. N. Phil. Journ. xvii. (1834) 363 = Scirpus Isolepis.
nodo, Boeck. in Flora, xli. (1858) 604 = S. lenticularis !
occidentalis, A. Gray, in Proc. Am. Acad. vii. (1868) 391.—Calif.
Schraderi, Kunth, Enum. Pl. ii. 268 = S. Isolepis.
senegalensis, Steud. Syn. Pl. Cyp. 180 = S. Isolepis.
subquarrosa, Nees, in Mart. Fl. Bras. ii. 1. 61 = S. micrantha.

HEMICARPURUS, Nees, Del. Sem. Hort. Vrat. (1839) 4; et Linnæa, xiv. (1840) Litt. 167 = Pinzellia, Tenore (Aroideæ.).
fornicatus, Nees, ll. cc. 4; 168 = P. tuberifera.

HEMICARPUS, F. Muell. in Hook. Kew Journ. ix. (1857) 18 = Trachynema, Rudge (Umbellif.).
didiscoides, F. Muell. l. c. = T. Hemicarpus.
villosa, F. Muell. l. c. = T. villosa.

HEMICHÆNA, Benth. Pl. Hartw. 78 (1841). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 948.
fruticosa, *Benth. l. c.*—Mexic.

HEMICHARIS, Salisb. ex DC. Prod. vii. 510 (1839) = Bosevola, Link. (Goodeniac.).

HEMICHLÆNA, Schrad. in Goett. Gel. Anz. (1821) iii. 2066. *CYPERACEÆ*, Benth. & Hook. f. iii. 1053.
angustifolia, *Schrad. l. c.*—Afr. austr.
bulbosa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 509 = Cyperus blysmoides.
capillifolia, *Schrad. in Goett. Gel. Anz.* (1821) iii. 2066.—Afr. austr.
fasciculoris, Steud. Syn. Pl. Cyp. 2 = angustifolia.
longifolia, *Nees, in Linnæa*, ix. (1834) 283.—Afr. austr.

HEMICHORISTE, Nees, in Wall. Pl. As. Rar. iii. 76 (1832) = Justicia, Linn. (Acanthac.).
montana, Nees, l. c. 102 = J. montana.

HEMICHROA, R. Br. Prod. 409 (1810). *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 59.
diandra, *R. Br. Prod.* 409.—Austral.
mesembryanthema, *F. Muell. Fragm.* viii. 38.—Austral.
pentandra, *R. Br. Prod.* 409.—Austral.

HEMICICCA, Baill. Etud. Gén. Euph. 645 (1858) = Phyllanthus, Linn. (Euphorbiac.).
japonica, Baill. l. c. 646 = P. japonicus.

HEMICLIDIA, R. Br. Prot. Nov. 40 (1810) = Dryandra, R. Br. (Proteac.).
Baxteri, R. Br. l. c. = D. falcata.

HEMICRAMBE, Webb, in Ann. Sc. Nat. Sér. III. xvi. (1851) 348. *CRUCIFERÆ*, Benth. & Hook. f. i. 99.
fruticulosa, *Webb, l. c.*—Marocc.

HEMICYCLIA, Wight & Arn. in Edinb. N. Phil. Journ. xiv. (Apr. 1833) 297 [Hemecyclia]. *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 378.
ANAMA, Miq. Fl. Ind. Bat. Suppl. 410 (1860).
ASTYLIS, Wight, Ic. t. 1992 (1853).
PERIPLEXIS, Wall. Cat. n. 6022 (1831).
andamanica, *Kurz, in Andam. Rep.* (1870) 47 ; *et Journ. As. Soc. Beng.* xlii. (1873) ii. 240.—Ins. Andam.
australasica, *Muell. Arg. in DC. Prod.* xv. ii. 487.—Austral.
elata, *Bedd. Fl. Sylv.* ii. t. 270.—Ind. or.
Gardneri, *Thw. in Hook. Kew Journ.* vii. (1855) 271. —Zeylan.
lanceolata, *Thw. Enum. Pl. Zeyl.* 287.—Zeylan.
lasiogyna, *F. Muell. Fragm.* iv. 119.—Austral.
rhabdolacea, Hassk. in Versl. en Med. Akad. Wetensch. iv. (1855) 140 = Cyclostemon rhacodiscus.
sepiaria, *Wight & Arn. in Edinb. N. Phil. Journ.* xiv. (1833) 297.—Ind. or.
sumatrana, *Muell. Arg. in DC. Prod.* xv. ii. 488.—Sumatra.
venusta, *Thw. in Hook. Kew Journ.* vii. (1855) 272. —Ind. or.

HEMIDEMUS, Dum. Anal. Fam. 28 (1829) = Hemidesmus, R. Br. (Asclep.).

HEMIDESMA, Rafin. Sylva Tellur. Index, 741 = seq.

HEMIDESMAS, Rafin. Sylva Tellur. 119 (1838) = Neptunia, Lour. (Legumin.).
diffusus, Rafin. l. c. = N. plena.
plenus, Rafin. l. c. = N. plena.

HEMIDESMUS, R. Br. in Mem. Wern. Soc. i. (1809) 56. *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 740.
cordatus, Schult. Syst. vi. 126 = indicus.
indicus, *R. Br. in Mem. Wern. Soc.* i. (1809) 56.—Zeylan.
pubescens, *Wight & Arn. Contrib.* 63.—Ind. or.
Wallichii, Miq. ex Hook. f. Fl. Brit. Ind. iv. 5 = indicus.
Wallichii, Wight & Arn. Contrib. 63 = Atherolepsis Wallichii.

HEMIDIA, Rafin. in Am. Monthly Mag. (1819) 191 (Quid ?).

HEMIERIUM, Rafin. Fl. Tellur. ii. 27 (1836) = Lloydia, Salisb. (Liliac.), etc.
græcum, Rafin. l. c. = L. græca.
planifolium, Rafin. l. c. = Simethis bicolor.

HEMIEVA, Rafin. Fl. Tellur. ii. 70 (1836) = Saxifraga, Linn.
ranunculifolia, Rafin. l. c. = S. ranunculifolia.
Richardsoni, Rafin. l. c. = Boykinia Richardsoni.

HEMIGENIA, R. Br. Prod. 509 (1810). *LABIATÆ*, Benth. & Hook. f. ii. 1218.
ATHELANDRA, Lindl. Swan River App. 40. t. 5 (1839).
COLOBANDRA, Bartl. in Lehm. Pl. Preiss. i. 357 (1845).
argens, Bartl. in Lehm. Pl. Preiss. i. 360 = sericea.
barbata, *Bartl. l. c.*—Austral.
brachyphylla, *F. Muell. Fragm.* x. 19.—Austral.
canescens, *Benth. in DC. Prod.* xii. 566.—Austral.
canefolia, *Benth. Fl. Austral.* v. 118.—Austral.
curvifolia, *F. Muell. Fragm.* i. 210.—Austral.
diplanthera, *F. Muell. l. c.* vi. 111.—Austral.
Drummondii, *Benth. Fl. Austral.* v. 118.—Austral.
glabrescens, *Benth. in DC. Prod.* xii. 566.—Austral.
humilis, *Benth. l. c.* 567.—Austral.
incana, *Benth. l. c.* 566.—Austral.
lanata, Benth. l. c. 567 = canescens !
lelantha, *F. Muell. Fragm.* xi. 20.—Austral.
loganiacea, *F. Muell. l. c.* 19.—Austral.
longifolia, Benth. Lab. Gen. et Sp. 291 = Microcoris longifolia.
macrantha, *F. Muell. Fragm.* i. 210.—Austral.
microphylla, *Benth. in DC. Prod.* xii. 565.—Austral.
mollis, Benth. l. c. 566 = canescens.
obovata, *F. Muell. Fragm.* xi. 19.—Austral.
obtusa, *Benth. in DC. Prod.* xii. 567.—Austral.
parviflora, Bartl. in Lehm. Pl. Preiss. i. 359 = sericea.
pimelifolia, *F. Muell. Fragm.* vi. 113.—Austral.
platyphylla, *Benth. in DC. Prod.* xii. 566.—Austral.
podalyriæa, *F. Muell. Fragm.* vi. 113.—Austral.
polystachya, Benth. in DC. Prod. xii. 566 = incana.
pungens, *F. Muell. Fragm.* xi. 20.—Austral.
purpurea, *R. Br. Prod.* 509.—Austral.
ramulosissima, *Benth. in DC. Prod.* xii. 566.—Austral.
rigida, *Benth. l. c.*—Austral.
scabra, *Benth. Fl. Austral.* v. 117.—Austral.
sericea, *Benth. in Enum. Pl. Hongk.* 86.—Austral.
Sieberi, Benth. Lab. Gen. et Sp. 457 = purpurea.
subvillosa, Benth. in DC. Prod. xii. 566 = platyphylla ?
tereticaulis, *F. Muell. Fragm.* vi. 111.—Austral.
westringioides, *Benth. in DC. Prod.* xii. 568.—Austral.

HEMIGRAPHIS, Nees, in DC. Prod. xi. 733 (1847). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1090.
HEMIGRAPHIS, T. Anders. in Journ. Linn. Soc. vii. (1864) 118.
alternata, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 114.—Ind. or.
confusa, *T. Anders. l. c.* ix. (1867) 468.—Malaya.
crenata, Walp. Ann. iii. 1094, Index, sphalm. = latebrosa.

HEMIGRAPHIS :—
crispa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 118. —China.
Cumingiana, *Villar, in Blanco, Fl. Philipp.* ed. iii. *Nov. App.* 153.—Ins. Philipp.
Decaisneana, *T. Anders. in Journ. Agr. Hort. Soc. Ind.* N. S. i. (1868) 270.—Malaya.
decipiens, *Villar, in Blanco, Fl. Philipp.* ed. iii. *Nov. App.* 153.—Ins. Philipp.
Gam, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 461. —Ind. or.
ebracteolata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 494.—Ind. or.
elegans, T. Anders. in Journ. Linn. Soc. ix. (1867) 468 = latebrosa.
elegans, *Nees, in DC. Prod.* xi. 733.—Burma.
flaccida, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 494.—Burma.
flava, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 74. —Ind. or.
glandulosa, T. Anders. in Kurz, Andam. Rep. App B. 13 = Strobilanthes glandulosus.
glaucescens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 495.—Burma.
Griffithiana, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 468.—Burma.
hirsuta, *T. Anders. in Journ. Agr. Hort. Soc. Ind.* N. S. i. (1868) 270.—Malaya.
hirta, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 462. —Ind. or.
latebrosa, *Nees, in DC. Prod.* xi. 733.—Ind. or.
lusona, *Villar, in Blanco, Fl. Philipp.* ed. iii. *Nov. App.* 153 = Ins. Philipp.
parabolica, *Villar, l. c.*—Ins. Philipp.
Pavala, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 78 = latebrosa.
primulæfolia, *Villar, in Blanco, Fl. Philipp.* ed. iii. *Nov. App.* 153.—Ins. Philipp.
quadrifaria, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 463.—Burma.
repens, *Villar, in Blanco, Fl. Philipp.* ed. iii. *Nov. App.* 154.—Ins. Philipp.
reptans, *T. Anders. ex Hemsl. Bot. Voy. Challenger*, i. iii. 178.—Ins. Arou.
rhytiphylla, *Radlk. in Sitzb. Math. Phys. Acad. Muench.* xiii. (‘1884’) (1883) 279.—Ins. Philipp.
rhytiphylla, *Villar, in Blanco, Fl. Philipp.* ed. iii. *Nov. App.* 153.—Ins. Philipp.
rupestris, T. Anders. in Journ. Linn. Soc. ix. (1867) 463 = ebracteolata.
strigosa, *Villar, in Blanco, Fl. Philipp.* ed. iii. *Nov. App.* 153.—Ins. Philipp.
venosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 493.—Ind. or.

HEMIGYMNIA, Griff. in Calc. Journ. Nat. Hist. iii. (1843) 363 = Cordia, Linn. (Borragin.).
Macleodii, Griff. l. c. = C. Macleodii.

HEMIGYROSA, Blume, Rumphia, iii. 165 (1850). *SAPINDACEÆ*, Benth. & Hook. f. i. 395.
canescens, *Thw. Enum. Pl. Zeyl.* 56, 408.—Zeylan.
deficiens, *Bedd. Fl. Sylv.* t. 261.—Ind. or.
longifolia, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 671.—Malacca.
Perrottetii, Blume, Rumphia, iii. 165 (= *Guioa Perrottetii*).—Ins. Philipp.
Perrillei, *Blume, l. c.* 166.—Ins. Amboin.
trichocarpa, Thw. Enum. Pl. Zeyl. 56 = canescens.

HEMILEPIS, Kunze, Ind. Sem. Hort. Lips. (1836); ex Schlecht. in Adv. Zeit. x. (1852) 675 = Leucosidea, Linn. (Compos.).
Ehrenbergii, Kunze, ex Schlecht. l. c.—Hab. !

HEMIMERIS, Linn. f. Suppl. 45 (1781); Thunb. Nov. Gen. 74 (1794). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 951.
alsinoides, Linn. Enum. iii. 106 = montana.
diffusa, Linn. f. Suppl. 280 = Dissotis diffusa, etc.
latipes, *Hance, ex Benth. in DC. Prod.* x. 284, in obs. —Afr. austr.
macrophylla, Thunb. Prod. Pl. Cap. 108 = Dissotis macrophylla, *Benth.*
montana, *Linn. f. Suppl.* 280.—Afr. austr.
pedunculosa, Lam. Encyc. iii. 108 = Dissotis Burgæana.
villosa, *Linn. f. Suppl.* 280.—Afr. austr.

HEMIMERIS :—
sessilifolia. *Benth. in Hook. Comp. Bot. Mag.* ii. (1886) 16.—Afr. austr.
sinuata, Sm. in Rees. Cyclop. xvii. n. 4 = Diascia heterandra †
unilabiata, Thunb. Prod. Pl. Cap. 105 = Diascia unilabiata.

HEMIMERIS, Pers. Syn. ii. 162 (1807), partim; H. B. & K. Nov. Gen. et Sp. ii. 376 (1817) = **Alonsoa**, Ruiz & Pav. (Scrophul.).
acutifolia, Pers. Syn. ii. 162 = A. acutifolia.
caulialata, Pers. l. c. = A. linearis.
coccinea, Willd. Sp. Pl. iii. 283 = A. linearis.
hirsuta, Spreng. Syst. ii. 809 = A. hirsuta.
incisifolia, Pers. Syn. ii. 162 = A. incisifolia.
intermedia, Lodd. Bot. Cab. t. 1456 = A. incisifolia.
linearifolia, H. B. & K. Nov. Gen. et Sp. ii. 377 = A. linearifolia.
linearis, Sm. in Trans. Linn. Soc. vi. (1802) 97 = *A. linearis*.
Mutisii, H. B. & K. Nov. Gen. et Sp. ii. 376 = A. caulialata.
parviflora, H. B. & K. l. c. = A. caulialata.
procumbens, Pers. Syn. ii. 162 = A. procumbens.
urticifolia, Willd. Sp. Pl. iii. 283 = A. incisifolia.

HEMINEMA, Rafin. Fl. Tellur. ii. 17 (1836) = **Tradescantia**, Linn. (Commelin.).
multiflora, Rafin. l. c. = T. multiflora.

HEMIORCHIS, Ehrenb. ex Schweinf. Betr. Fl. Aethiop. 241 (1867) = **Lindenbergia**, Lehm. (Scroph.).
habeanrica, Ehrenb. ex Schweinf. l. c. = L. scutellarioides.

HEMIORCHIS, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 108. t. 8. *SCITAMINEAE*, Benth. & Hook. f. iii. 641.
burmanica, *Kurz, l. c.*—Burma.

HEMIPAPPUS, C. Koch, in Linnaea, xxiv. (1851) 340 = **Tauscetum**, Tourn. (Compos.).
canus, C. Koch, l. c. 341 = T. argenteum.
tsabellinus, C. Koch, l. c. = T. argenteum.

HEMIPHORA, F. Muell. Census, 103 (1882). *VERBENACEAE*.
Elderi, *F. Muell. l. c.*—Austral.

HEMIPHRACTUM, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 262 = **Vateria**, Linn. (Dipterocarp.).
oxyandrum, Turcz. l. c. 263 = V. acuminata.

HEMIPHRAGMA, Wall. in Trans. Linn. Soc. xiii. (1822) 611; et Tent. Fl Nep. 16. t. 8 (1824). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 969.
heterophyllum, *Wall. Il. cc.* 17; 613.—Neg. Himal.

HEMIPHUES, Hook. f. in Hook. Lond. Journ. vi. (1847) 460 [bis] = **Actinotus**, Labill. (Umbellif.).
sphæle, Hook. f. l. c. 471 [bis] = A. bellidioides.
bellidioides, Hook. f. l. c. 470 [bis] = A. bellidioides.
Novæ-Zelandiæ, Petrie, in Trans. N. Z. Inst. xii. (1880) 385.—N. Zel.
suffocata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 471 [bis] = A. bellidioides.
tridentata, Hook. f. l. c. = A. bellidioides.

HEMIPHYLACUS, S. Wats. in Proc. Am. Acad. xviii. (1883–83) 164, 165. *LILIACEAE*.
latifolius, *S. Wats. l. c.* 164.—Mexic.

HEMIPILIA, Lindl. Gen. et Sp. Orch. 296 (1835). *ORCHIDEAE*, Benth. & Hook. f. iii. 638.
calophylla, *Par. & Reichb. f. in Journ. Bot.* xii. (1874) 197.—Burma.
cordifolia, *Lindl. Gen. et Sp. Orch.* 296.—Reg. Himal.

HEMIPOGON, Decne. in DC. Prod. viii. 500 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 748.
abietoides, *Fourn. in Mart. Fl. Bras.* vi. iv. 196.—Bras.
acerosus, *Decne. in DC. Prod.* viii. 500.—Bras.
hispidus, *Fourn. in Mart. Fl. Bras.* vi. iv. 197.—Bras.

HEMIPOGON :—
peruvianus, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 368.—Peruv.
setaceus, *Decne. in DC. Prod.* viii. 509.—Bras.
Spruceî, *Fourn. in Mart. Fl. Bras.* vi. iv. 197.—Bras.

HEMIPTELEA, Planch. in Comp. Rend. Par. lxxiv. (1872) 131 = **Kelkova**, Spach (Urticac.).
Davidii, Planch. l. c.—China.

HEMIPTILIUM, A. Gray, in Torr. Bot. Mex. Bound. 105 (1858) = **Stephanomeria**, Nutt. (Compos.).
Bigelowii, A. Gray, l. c. = S. exigua.
Schottii, A. Gray, l. c. = S. Schottii.

HEMISACRIS, Steud. in Flora, xii. (1829) 490 = **Schizmus**, Beauv. (Gramin.).
gonatodes, Steud. l. c. = S. marginatus.

HEMISANDRA, Scheidw. in Bull. Acad. Brux. ix. i. (1842) 22 = **Aphelandra**, R. Br. (Acanthac.).
aurantiaca, Scheidw. l. c. 23 = A. aurantiaca.

HEMISCLERIA, Lindl. Fol. Orchid. (1853) = **Diothonea**, Lindl. (Orchid.).
nutans, Lindl. l. c.—Peruv.

HEMISCOLA, Rafin. Sylva Tellur. 111 (1838) = **Cleome**, Linn. (Capparid.).
aculeata, Rafin. l. c. = C. aculeata.
ornithopoides, Rafin. l. c. = C. brachycarpa †

HEMISODON, Rafin. Fl. Tellur. iii. 88 (1836) = **Leonotis**, Pers. (Labiat.).
Leonurus, Rafin. l. c. = L. Leonurus.

HEMISPADON, Endl. in Flora, xv. (1832) 885. t. 1 = **Indigofera**, Linn. (Legum.).
lanata, Endl. l. c. = I. pilosa.

HEMISPHACE, Opiz, Seman, 50 (1852) = **Salvia**, Tourn. (Labiat.).
verticillata, Opiz, l. c. = S. verticillata.

HEMISTEGIA, Rafin. Fl. Tellur. iii. 89 (1836) = **Salvia**, Tourn. (Labiat.).
mexicana, Rafin. l. c. 90 = S. mexicana.

HEMISTEIRUS, F. Muell. in Linnaea, xxv. (1852) 424 = **Triohinium**, R. Br. (Amaranthac.).
spiniorichoides, F. Muell. l. c. 425 = T. corymbosum.

HEMISTEMMA, Juss. ex Thou. Gen. Nov. Madag. 18 (1806) = **Hibbertia**, Andr. (Dilleniac.).
angustifolium, R. Br. in DC. Syst. i. 414 = Hibbertia angustifolia.
asperifolium, F. Muell. Fragm. i. 161 = Hibbertia furfuracea.
Auberti, DC. Syst. i. 413 = Hibbertia coriacea.
Banksii, R. Br. in DC. Syst. i. 414 = Hibbertia Banksii.
candicans, Hook. f. in Hook. Kew Journ. ix. (1857) 48 = Hibbertia candicans.
Commersonii, DC. Syst. i. 413 = Hibbertia coriacea.
denthatum, R. Br. ex DC. Syst. i. 413.—Austral.
ledifolium, A. Cunn. ex Benth. Fl. Austral. i. 22 = Hibbertia ledifolia.
Lenehenaultii, DC. Syst. i. 414 = Beyeria opaca.
revolutum, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 4 = Hibbertia microphylla.

HEMISTEMMA, Reichb. Consp. 116 (1828) = **Leucas**, R. Br. (Labiat.).

HEMISTEMON, F. Muell. Fragm. x. 78 (1876) = **Chlorasthes**, R. Br. (Verbenac.).
Bonnayi, F. Muell. l. c. = C. Bonnayana.

HEMISTEPHIA. Steud. Nom. ed. II. i. 749 (1840), sphalm. = HEMISTEPTA, Bunge.

HEMISTEPHUS, Drum. ex Harv. in Hook. Kew Journ. vii. (1855) 51 = **Hibbertia**, Andr. (Dillen.).
linearis, Drum. ex Harv. l. c. 53 = Hibbertia spicata.

HEMISTEPTA, Bunge, in Dorp. Jahrb. Litt. i. (1838) 223 = **Saussurea**, DC. (Compos.).
lyrata, Bunge, l. c. = S. affinis.

HEMISTEPTIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. (1835) 38 = prec.

HEMISTOMA, Ehrenb. ex Benth. in Wall. Pl. As. Rar. i. 60 (1830) = **Leucas**, R. Br. (Labiat.).
ovatum, Ehrenb. ex Benth. l. c. = L. urticæfolia.

HEMISTYLIS, Walp. Ann. i. 648 (1849) = seq.

HEMISTYLUS, Benth. Pl. Hartw. 123 (1848). *URTICACEAE*, Benth. & Hook. f. iii. 393.
boehmerioides, *Benth. l. c.*—Ecuador.
brasiliensis, *Wedd. ex Warm. in Kjoeb. Vidensk. Meddel.* (1878) 51.—Bras.
macrostachys, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 208.—Venezuela.
odontophylla, *Wedd. in DC. Prod.* xvi. i. 235*.—Venezuela.
velutina, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 526.—N. Granat.

HEMITHRINAX, Hook. f. in Benth. & Hook. f. Gen. iii. 980 (1883). *PALMAE*, Benth. & Hook. f. i. c.
compacta, *Hook. f. l. c.* 981.—Cuba.

HEMITOME, Nees, in DC. Prod. xi. 295 (1847), in obs. = **Stenandrium**, Nees (Acanth.).
ignea, Nees, l. c. 292, in syn. = S. igneum.

HEMITOMES, A. Gray, in Pacif. Rail. Rep. vi. 80. t. 12 (1855) = **Newberrya**, Torr. (Monotrop.).
congestum, A. Gray, l. c. 81 = N. congesta.

HEMITOMUS, L'Hérit. ex Desf. Tabl. ed. I. 66 (1804) = **Alonsoa**, Ruiz & Pav. (Scroph.).
fruticosus, L'Hérit. ex Desf. l. c.; et ed. II. 79 = A. linearis.
urticæfolius, L'Hérit. ex Desf. l. c. ed. II. 79 = A. urticæfolia.

HEMITRIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 79. *LORANTHACEAE*.
unifora, Rafin. l. c.—Hab. †

HEMIXANTHIDIUM, Delpino, Studi sopra Artemis. 60 (1871) = **Franseria**, Cav. (Compos.).
montanum, Delpino, l. c. = F. Hookeriana.

HEMIZONELLA, A. Gray, in Proc. Am. Acad. ix. (1874) 189. *COMPOSITAE*.
Durandi, *A. Gray, l. c.*—Calif.
minima, *A. Gray, l. c.*—Calif.
parvula, A. Gray, l. c. = Durandi.

HEMIZONIA, DC. Prod. v. 692 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 394.
CALYCADENIA, DC. Prod. v. 692 (1836).
HARTMANNIA, DC. l. c. 693 (1836).
OSMADENIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 391.
angustifolia, Benth. Pl. Hartw. 318 = corymbosa.
angustifolia, *DC. Prod. v.* 692.—Calif.
angustifolia, Durand, in Pacif. Rail. Rep. v. 10 = mollis.
balsamifera, Kellogg, in Proc. Calif. Acad. ii. (1863) 64 = corymbosa.
Cephalotes, Greene, in Bull. Torrey Bot. Club, ix. (1882) 110 = multiglandulosa.
ciliata, Torr. & Gray, Fl. N. Am. ii. 398 = Layia chrysanthemoides.
citrioloru, *A. Gray, Syn. Fl. N. Am.* i. ii. 207.—Calif.
Clevelandi, *Greene, in Bull. Torrey Bot. Club.* ix. (1882) 109.—Calif.
congesta, *DC. Prod. v.* 692.—Calif.
congesta, A. Gray, in Pacif. Rail. Rep. iv. 109 = Lagophylla congesta.
congesta, Hook. & Arn. Bot. Beech. Voy. 355 = angustifolia.
corymbosa, *Torr. & Gray, Fl. N. Am.* ii. 398.—Calif.
decumbens, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 175 = angustifolia.
Douglasii, *A. Gray, in Bot. Calif.* i. 366.—Calif.
Durandii, A. Gray, in Proc. Am. Acad. vi. (1864–65) 549 = Hemizonella Durandi.
fasciculata, *Torr. & Gray, Fl. N. Am.* ii. 397.—Calif.
filipes, Hook. & Arn. Bot. Beech. Voy. 356 = Lagophylla filipes.
Fitchii, *A. Gray, in Pacif. Rail. Rep.* iv. 109.—Calif.
floribunda, *A. Gray, in Proc. Am. Acad.* xi. (1876) 79. —Calif.

HEMIZONIA :—
Pronostii, A. Gray, l. c. ix. (1874) 191 = multiglandulosa.
fruticosa, *A. Gray*, *l. c.* xi. (1876) 79.—Calif.
glomerata, A. Gray, Syn. Fl. N. Am. i. ii. 309 = fasciculata.
Heermanni, *Greene*, *in Bull. Torrey Bot. Club*, ix. (1882) 15.—Calif.
hispida, Greene, l. c. 63 = Douglasii.
Kellogii, *Greene*, *l. c.* x. (1883) 41.—Calif.
Labbii, Greene, l. c. ix. (1882) 109 = fasciculata.
lasulacifolia, *DC. Prod. v.* 692.—Calif.
macradenia, *DC. l. c.* 693.—Calif.
macrocephala, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 175 = corymbosa.
minima, *A. Gray*, *in Proc. Am. Acad.* vi. (1864–65) 549.—Calif.
mollis, *A. Gray*, *in Bot. Calif.* i. 365.—Calif.
multiculis, Hook. & Arn. Bot. Beech. Voy. 355 = angustifolia ?
multiglandulosa, *A. Gray*, *in Bot. Calif.* i. 366.—Calif.
oppositifolia, Greene, in Bull. Torrey Bot. Club, ix. (1882) 110 = multiglandulosa.
paniculata, *A. Gray*, *in Proc. Am. Acad.* xix. (1883) 17.—Calif.
Parryi, *Greene*, *in Bull. Torrey Bot. Club*, ix. (1882) 16.—Calif.
parvifolia, Greene, l. c. = Heermanni.
parvula, A. Gray, in Proc. Am. Acad. vi. (1864–65) 549 = Hemizonella Durandi.
pauciflora, *A. Gray*, *in Bot. Calif.* i. 365.—Calif.
plumosa, A. Gray, in Proc. Am. Acad. ix. (1874) 192 = Blepharizonia plumosa (in Addend.).
pungens, *Torr. & Gray*, *Fl. N. Am.* ii. 299.—Calif.
ramosissima, Benth. Bot. Voy. Sulph. 30 = fasciculata.
rudis, Benth. l. c. 31 = lasulacifolia.
scrisea, Hook. & Arn. Bot. Beech. Voy. 356 = luxulacifolia.
spicata, Greene, in Bull. Torrey Bot. Club, ix. (1882) 16 = Douglasii.
squalida, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 308.—Ins. Galapag.
Streetsii, *A. Gray*, *in Proc. Am. Acad.* xii. (1877) 162.—Calif.
tenella, *A. Gray*, *l. c.* ix. (1874) 191.—Calif.
truncata, *A. Gray*, *in Bot. Calif.* i. 365.—Calif.
virgata, *A. Gray*, *in Torr. Bot. Mex. Bound.* 100.—Calif.
Wheeleri, *A. Gray*, *in Bot. Calif.* i. 617.—Calif.
Wrightii, *A. Gray*, *in Proc. Am. Acad.* xix. (1883) 17.—Calif.

HEMOLEPIS, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 396 (1866) = **Heliopsis**, Pers. (Compos.).
canescens, Hort. ex Vilm. l. c. = Heliopsis buphthalmoides.

HEMONACANTHUS, Nees, in DC. Prod. xi. 725 (1847) sphalm. = STEMONACANTHUS, Nees = **Ruellia**, Linn. (Acanth.).

HEMPRICHIA, Ehrenb. in Linnæa, iv. (1829) 396 = **Commiphora**, Jacq. (Burserac.).
erythræa, Ehrenb. l. c. = C. erythræa.

HEMYPHYES, Endl. Gen. Suppl. v. 2 (1850) = HEMIPHUES, Hook. f. = **Actinotus**, Labill. (Umbellif.).

HENANTHUS, Less. Syn. Comp. 196 (1832) = Piptonia, Llan. (Compos.).
fasciculatus, Less. l. c. = P. fasciculata.

HENCKELIA, Spreng. Anleit. ii. 409 (1817) = **Didymocarpus**, Wall. (Gesnerac.).
alata, A. Dietr. Sp. Pl. i. 578 = Bonnaya alata.
aromatica, Spreng. Syst. iv. Cur. Post. 18 = D. subalternans.
barbata, Spreng. l. c. = Chirita Horsfieldii.
bifolia, A. Dietr. Sp. Pl. i. 574 = Chirita bifolia.
caponis, A. Br. ex C. B. Clarke, in DC. Monog. Phan. v. 151 = Streptocarpus Rexii.
cinerea, Spreng. Syst. iv. Cur. Post. 14 = D. cinerea.
cornicalata, Spreng. l. c. 18 = D. cornicalata.
crinita, Spreng. l. c. = D. crinita.
elongata, Spreng. l. c. = Didissandra elongata.
fruticosa, Spreng. l. c. = Didymocarpus fruticosus.
grandiflora, A. Dietr. Sp. Pl. i. 578 = Bonnaya verscaefolia.
grandifolia, A. Dietr. l. c. 576 = Chirita macrophylla.

HENCKELIA :—
incana, Spreng. Syst. i. 98 = Didymocarpus Rottleriana.
macrophylla, Spreng. l. c. iv. Cur. Post. 14 = Chirita macrophylla.
oblonga, Spreng. l. c. 18 = D. oblonga.
oppositifolia, A. Dietr. Sp. Pl. i. 573 = Bonnaya oppositifolia.
plicata, Spreng. Syst. iv. Cur. Post. 14 = D. macrophylla.
primulæfolia, Spreng. l. c. 14 = D. aromatica.
pumila, A. Dietr. Sp. Pl. i. 574 = Chirita pumila.
racemosa, Spreng. Syst. iv. Cur. Post. 18 = D. racemosa.
reptans, Spreng. l. c. 14 = D. reptans.
Rexii, Heynh. Nom. i. 390; ii. 708 = Streptocarpus Rexii.
Roxburghiana, A. Dietr. Sp. Pl. i. 572 = Bonnaya reptans.
scrisea, *Wallichiana*, A. Dietr. l. c. 574 = Chirita urticæfolia.
villosa, Spreng. Syst. iv. Cur. Post. 13 = D. villosa.
Wallichiana, A. Dietr. Sp. Pl. i. 574 = Chirita urticæfolia.

HENDECANDRAS, Eschsch. in Mém. Acad. Pétersb. x. (1826) 387 = Croton, Linn. (Euphorb.).
divaricatus, Klotzsch, in Wiegm. Archiv, vii. (1841) 198 = C. nitraziæfolius.
glabrescens, Klotzsch, l. c. = C. nitraziæfolius.
longifolius, Klotzsch, l. c. = C. montevidensis.
maritimus, Klotzsch, l. c. 252 = C. maritimus.
montevidensis, Klotzsch, l. c. 198 = C. montevidensis.
multiflorus, Torr. in Frém. First Rep. 96 = C. texensis.
polymorphus, Klotzsch, in Wiegm. Archiv, vii. (1841) 198 = C. nitraziæfolius.
prunifolius, Eschsch. in Mém. Acad. Pétersb. x. (1826) 387 = C. californicus.
texensis, Klotzsch, in Wiegm. Archiv, vii. (1841) 252 = C. texensis.
velleriflora, Klotzsch, l. c. 198 = C. nitraziæfolius.

HENFREYA, Lindl. Bot. Reg. (1847) t. 31 = **Asystasia**, Blume (Acanthac.).
scandens, Lindl. l. c. = A. scandens.

HENICOSTEMMA, Endl. Gen. 605 (1838) = **Enicostema**, Blume (Gentian.).

HENISIA, Walp. Rep. ii. 517 (1843) = **Heinsia**, DC. (Rubiac.).

HENKELIA, Reichb. Consp. 134 (1828) = HENCKELIA, Spreng. = **Didymocarpus**, Wall. (Gesner.).

HENLEA, Griseb. in Abh. Ges. Goott. ix. (1860) 37. *MALPIGHIACEÆ* [Benth. & Hook. f. i. 356].
HENLEOPHYTUM, Karst. Fl. Colomb. i. 158 (1861).
echinata, *Griseb. in Abh. Ges. Goott.* ix. (1860) 37.—Cuba.

HENLEA, Karst. Fl. Columb. 86, in textu (1858); 157. t. 76 (1861) = **Rustia**, Klotzsch (Rubiac.).
macrocnemis, Klotzsch & Karst. ex Walp. Ann. ii. 788 = Cascarilla mascnensis.
rosea, Klotzsch & Karst. ex Walp. l. c. = Cascarilla mascnensis.
splendens, Karst. Fl. Columb. i. 157. t. 78.—N. Granat.
thibaudioides, Karst. in Linnæa, xxx. (1859–60) 151.—N. Granat.

HENLEOPHYTUM, Karst. Fl. Columb. i. 158 (1861) = **Henlea**, Griseb. (Malpig.).
plumigerum, Sauv. Fl. Cub. 16 = Henlea echinata.

HENNECARTIA, Poisson, in Bull. Soc. Bot. Fr. xxxii. (1885) 40; Etude Nouv. Gen. Hennec. 5. *MONIMIACEÆ*.
omphalandra, *Poisson*, *ll. cc.* 41; 5.—Paraguay.

HENNINGIA, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 518 = **Brunnera**, Bieb. (Liliac.).
angustifolia, J. Gay, ex Baker, in Journ. Linn. Soc. xv. (1876) 322 = E. angustifolius.
asiatopora, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 517 = E. asiaticus.
Auchericana, Boiss. Diagn. Ser. II. iv. 98 = E. Aucherianus.
persica, Regel, in Bull. Soc. Nat. Mosc. xli. (1868) i. 485 = E. persicus.
robusta, Regel, l. c. 487 = E. robustus.

HENONIA, Cass. & Dur. in Bull. Soc. Bot. Fr. ii. (1855) 346 = **Henophyton**, Cass. & Dur. (Crucifer.).
deserti, Cass. & Dur. l. c. 347 = Henophyton deserti.

HENONIA, Moq. in DC. Prod. xiii. ii. 337 (1849). *AMARANTACEÆ*, Benth. & Hook. f. iii. 24.
scoparia, *Moq. l. c.*—Madag.

HENONIX, Rafin. Fl. Tellur. ii. 13 (1836) = Scilla, Linn. (Liliac.).
japonica, Rafin. l. c. = S. japonica.

HENOONIA, Griseb. Cat. Fl. Cub. 166 (1866). *SAPOTACEÆ*, Benth. & Hook. f. ii. 662.
myrtifolia, *Griseb. l. c.* 167.—Cuba.

HENOPHYTON, Cass. & Dur. in Bull. Soc. Bot. Fr. ii. (1855) 695; iv. (1857) 469. *CRUCIFERÆ*, Benth. & Hook. f. i. 85.
HENONIA, Cass. & Dur. in Bull. Soc. Bot. Fr. ii. (1855) 346.
deserti, *Cass. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 469.—Alger.

HENRICEA, Lem. Ls. in Bull. Soc. Philom. (1824) 171 = **Swertia**, Linn. (Gentian.).
pharmacoarcha, Lem. Ls. l. c. = S. Chirayta.
spicata, Lem. Ls. (sp. ignot.).—Hab. ?

HENRICIA, Cass. in Bull. Soc. Philom. (1817) 11. *COMPOSITÆ*, Benth. & Hook. f. ii. 377.
agathacoides, *Cass. l. c.*—Madag.

HENRIETIA, Reichb. Consp. 174 (1828) = seq.

HENRIETTA, Macfad. Fl. Jamaic. ii. 76 (1837 ?) = seq.

HENRIETTEA, DC. Prod. iii. 173 (1828). *MELASTOMACEÆ*, Benth. & Hook. f. i. 766.
PHYLLOPUS, DC. Prod. iii. 177 (1828).
angustifolia, *Berg*, *ex Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 145.—Bras.
brasiliensis, Cogn. Nov. Stirp. Bras. Dec. 86 = succosa.
fissacarpa, Benth. Hist. Fl. vii. 84 = Henriettella fissacarpa.
grandifolia, Macfad. Fl. Jamaic. ii. 76 = ramiflora.
granulata, *Berg*, *ex Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 144.—Bras.
maronicensis, *Sagot*, *in Ann. Sc. Nat.* Sér. VI. xv. (1883) 280.—Guian.
Martii, *Naud. l. c.* III. xviii. (1852) 105.—Guian.
multiflora, *Naud. l. c.*—Guian.
oblongifolia, Naud. l. c. iv. (1845) 56 = Martii.
ovinacensis, Naud. l. c. 57 = Martii.
parviflora, Griseb. Cat. Pl. Cub. 95 = Henriettella parviflora.
Patriziana, *DC. Prod.* iii. 178.—Guian.
racemosa, *Macfad. Fl. Jamaic.* ii. 76.—Jamaic.
ramiflora, *DC. Prod.* iii. 178.—Jamaic.
Sagotiana, *Naud. ex Sagot*, *in Ann. Sc. Nat.* Sér. VI. xv. (1883) 280.—Guian.
Sagotiana, Naud. ex Triana, in Trans. Linn. Soc. xxvii. (1871) 143 = Loreya mespiloides.
stellaris, *Berg*, *ex Triana*, *l. c.* 146.—Bras.
succosa, Benth. in Hook. Journ. Bot. ii. (1840) 306 = multiflora.
succosa, *DC. Prod.* iii. 178.—Guian.
succosa, Griseb. Fl. Brit. W. Ind. 246 = ramiflora.
surinamensis, Miq. Stirp. Surin. Select. 45 = ramiflora.
trinervia, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 105 = ramiflora.

HENRIETTELLA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 107. *MELASTOMACEÆ*, Benth. & Hook. f. i. 769.
aggregata, *Triana*, *in Trans. Linn. Soc.* xxviii. (1872) 148.—Peruv.
fascicularis, *Triana*, *l. c.*—Ind. occ.
flexuosa, *Triana*, *l. c.*—Guiana.
Goudotiana, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 106.—N. Granat.
lateriflora, *Triana*, *in Trans. Linn. Soc.* xxviii. (1872) 144.—Ind. occ.; Venezuel.
Maschnyiana, *Triana*, *l. c.* 148.—Jamaic.
parviflora, *Triana*, *l. c.*—Cuba.
Patriziana, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 107 = Henriettea Patriziana.

HENRIETTELLA:—
punctata, Triana, in Trans. Linn. Soc. xxviii. (1871) 148.—Cuba.
Seemannii, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 108.—Panama.
ramiflcila, Triana, in Trans. Linn. Soc. xxviii. (1871) 148.—Ind. occ.; Venezuel.
trachyphylla, *Triana, l. c.* 144.—N. Granat.
trifora, *Triana, l. c.* 148.—Ind. occ.
umbellalibera, *Triana, l. c.*—Bras.
verrucosa, *Triana, l. c.* 144.—Peruv.

HENRINCQUIA, Benth. & Hook. f. Gen. ii. 1006 (1876), sphalm. = HERINCQUIA, Decne. = Pentaphaphita, Lindl. (Gomerac.).

HENRIQUESIA, Spruce, ex Benth. in Hook. Kew Journ. vi. (1854) 338. *RUBIACEAE*, Benth. & Hook. f. ii. 44.
nitida, *Spruce, ex Benth. in Trans. Linn. Soc.* xxii. (1859) 397.—Bras.
oblonga, *Spruce, ex Benth. l. c.*—Bras.
obovata, *Spruce, ex Benth. l. c.* 396.—Bras.
verticillata, *Benth. in Hook. Kew Journ.* vi. (1854) 338.—Bras.

HENRYA, Nees, in Benth. Bot. Voy. Sulph. 148. t. 49 (1844) = Tetramerium, Nees (Acanthac.).
Barclayana, Nees, l. c. 149. t. 49 = T. scorpioides.
insularis, Nees, l. c. = T. scorpioides.
scorpioides, Nees, l. c. 148. t. 49 = T. scorpioides.

HENSCHELIA, Presl, Rel. Haenk. ii. 81. t. 63 (1831) = Illigera, Blume (Combretac.).
lusoensis, Presl, l. c. = I. Meyeniana.

HENSLERA, Endl. Gen. 791 (1839) = HAENSELERA, Lag. = Physospermum, Cuss. (Umbellif.).

HENSLERA, Reichb. Nom. 96 (1841) = Haenslera, Boiss. (Compos.).

HENSLOVIA, A. Juss. in Orb. Dict. vi. 536 (1849) = HENSLOWIA, Wall. = Cryptaronia, Blume (Lythrar.).

HENSLOWIA, Blume, Mus. Bot. Lugd. Bat. i. 242. t. 48 (1850). *SANTALACEAE*, Benth. & Hook. f. iii. 228.
DENDROTROPHE, Miq. Fl. Ind. Bat. i. i. 779 (1855).
DUPRENOVA, Chatin, in Compt.-Rend. Par. li. (1860) 657.
TUPEIA, Blume, Fl. Jav. Loranth. tt. 27, 28 (1828 ?).
buxifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 244.—Malaya.
castaefolia, *Blume, l. c.*—Borneo.
erythrocarpa, *Kurz, in Journ. Bot.* xiii. (1875) 329.—Ins. Nicobar.
fruticosa, *Benth. in Hook. Kew Journ.* v. (1853) 194.—Hongkong.
grandiata, *Hook. f. & Thoms. ex A. DC. in DC. Prod.* xiv. 632.—Reg. Himal.
heterotricha, *Hook. f. & Thoms. ex A. DC. l. c.*—Reg. Himal.; Burma.
Lobbiana, *A. DC. l. c.* 631.—Malaya.
paniculata, *Miq. Fl. Ind. Bat.* i. 1. 716.—Sumatra.
pancifiora, *Blume, Mus. Bot. Lugd. Bat.* i. 244.—Java.
philippinensis, A. DC. in DC. Prod. xiv. 631 = *variana*.
Reinwardtiana, *Blume, Mus. Bot. Lugd. Bat.* i. 244.—Sumatra.
retusa, *Blume, l. c.* 243.—Java.
spicata, *Blume, l. c.* 244.—Malaya.
umbellata, *Blume, l. c.* 243.—Malaya.
variana, Blume, l. c. 244. t. 43.—Burma; Malaya.

HENSLOWIA, Lowe, ex DC. Prod. viii. 288 (1844) = Notaelaea, Vent. (Oleac.).

HENSLOWIA, Wall. Pl. As. Rar. iii. 13 (1832) = Cryptaronia, Blume (Lythrar.).
affinis, Planch. in Hook. Lond. Journ. Bot. iv. (1845) 477 = C. pubescens.
Cumingii, Planch. l. c. 476 = C. Cumingii.
glabra, Wall. Pl. As. Rar. iii. 14 = C. glabra.
Hookeri, Wall. Cat. n. 6946 = C. pubescens.
leptostachya, Planch. in Hook. Lond. Journ. Bot. iv. (1845) 476 = C. pubescens.
pubescens, Wall. Pl. As. Rar. iii. 14 = C. pubescens.

HEOCARPHUS, Phil. in Anal. Univ. Chil. (1861) 51, sphalm. = PLEOCARPHUS, D. Don = Jungia, Linn. f. (Compos.).

HEPATICA, Dill. ex Linn. Syst. ed. I (1735) = Anemone, Linn. (Ranunc.).
acutiloba, DC. Prod. i. 22 = A. acutiloba.
americana, Ker-Gawl. in Bot. Reg. t. 887 = A. Hepatica.
anemonoides, Vest, Man. Bot. 206 = A. Hepatica.
angulosa, DC. Syst. i. 217 = A. angulosa.
integrifolia, DC. l. c. = A. integrifolia.
multiloba, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 116 = A. angulosa.
nobilis, Schreb. Spicil. Fl. Lips. 89 = A. Hepatica.
transilvanica, Fuss, in Verh. Siebenb. Ver. Naturw. i. (1850) 83 = A. angulosa.
triloba, Choix, in Vill. Hist. Pl. Dauph. i. 338 = A. Hepatica.

HEPETIS, Sw. Prod. Veg. Ind. Occ. 56 (1788) = Pitcairnia, L'Hérit. (Bromeliac.).
angustifolia, Racusch. Nom. ed. III. 91 = P. angustifolia.
angustifolia, Sw. Prod. Veg. Ind. Occ. 56 = P. bromeliaefolia.
bromeliaefolia, Saliab. Prod. 247 = P. bromeliaefolia.
latifolia, Racusch. Nom. ed. III. 91 = P. latifolia.

HEPETOSPERMUM, Spach, Hist. Vég. Phan. vi. 188 (1838), sphalm. = Herpetospermum, Wall. (Cucurb.).

HEPHESTIONIA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 36 = Tibouchina, Aubl. (Melastom.).
Chamaecistus, Naud. l. c. (= *Pleroma Chamaecistus*). —Ins. Martinic.
strigosa, Naud. l. c. (= *Pleroma strigosum*).—Ind. occ.

HEPTIRILLA, Regel, Gartenfl. ii. (1853) 353. t. 70. *GESNERACEAE*, Benth. & Hook. f. ii. 999.
CHEIRISANTHERA, Hort. ex Regel, Gartenfl. ii. (1853) 353.
amplia, *Hanst. in Linnaea*, xxix. (1857–58) 514.—Ecuador.
atrosanguinea, *Regel, in Gartenfl.* (1853) 353.—N. Granat.
cordata, *Klotzsch & Hanst. in Linnaea*, xxix. (1857–58) 513.—Venezuela.
nasgueliedes ×, Lam. Illustr. Hortic. iv. (1857) t. 129.
ovata, *Klotzsch & Hanst. in Linnaea*, xxix. (1857–58) 512.—Venezuela.
pancifiora, *Hanst. l. c.* 514.—N. Granat.
repens, *Hanst. l. c.*—Ecuador.
stenoclyta, *Hanst. l. c.* 513.—N. Granat.
umbrifolia, *Hanst. l. c.*—Ecuador.
Warszewiczii, *Hanst. l. c.* 514.—Am. austr.

HEPTACA, Lour. Fl. Cochinch. 657 (1790) = Onooba, Forsk. (Ilicin.).
africana, Lour. l. c.—Afr. trop.
latifolia, Hardw. & Champ. in Hook. Kew Journ. i. (1849) 248 = Actinidia Championi.

HEPTALLON, Raf. Neogenyt. 1 (1825) = Croton, Linn. (Euphorb.).
aromaticum, Raf. l. c. = C. aromaticus, *Linn.*
capitatum, Raf. l. c. = C. capitatus.
elliptieum, Raf. l. c. = C. elliptieus.
graveolens, Raf. l. c.—Hab. ?

HEPTANTHUS, Griseb. Cat. Pl. Cub. 148 (1866). *COMPOSITAE*, Benth. & Hook. f. ii. 343.
brevipes, *Griseb. l. c.*—Cuba.
cochlearifolius, *Griseb. l. c.*—Cuba.
rauncunloides, *Griseb. l. c.*—Cuba.

HEPTAPLEURUM, Gaertn. Fruct. ii. 472. t. 178. (1791). *ARALIACEAE*, Benth. & Hook. f. i. 942.
ACTINOMORPHE, Miq. Comm. Phyt. 109 (1840).
AGALMA, Miq. Fl. Ind. Bat. i. 1. 751. t. 12 (1855).
ASTROPANAX, Seem. in Journ. Bot. iii. (1865) 176.
GYNOPHTHINA, Spach, Hist. Vég. Phan. viii. 118 (1838), partim.
PARASTROPIA, DC. Prod. iv. 265 (1830).
SCIADOPHYLLUM, Blume, Bijdr. 875 (1826).
UNGALA, Blume, Cat. Gew. Buitenz. 66 (1823).

HEPTAPLEURUM:—
abyssinicum, *Benth. & Hook. f. Gen.* i. 942; *Vathe, in Linnaea*, xl. (1876) 191.—Afr. trop.
acutangulum, Gaertn. Fruct. ii. t. 178. f. 3 = stellatum.
acuticuminum, *Seem. in Journ. Bot.* iii. (1865) 79.—Borneo.
apiculatum, *Seem. l. c.* 78.—Java.
avene, *Seem. l. c.* 79.—Sumatra.
Balkisi, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 80.—Afr. trop.
Barteri, *Hiern, l. c.*—Afr. trop.
biternatum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 731.—Malacca.
capitatum, Seem. in Journ. Bot. iii. (1865) 81 = Brassaia capitata.
caudatum, Vidal, Phan. Cuming. Philipp. 175 = H. Cephalotes.
Cephalotes, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 731.—Malacca.
confine, *Seem. in Journ. Bot.* iii. (1865) 79.—Ins. Celebes.
Coronas-sylvae, *Seem. l. c.* 78.—Java.
Cumingii, *Seem. l. c.* 81.—Ins. Philipp.
dipyrenum, *H. Mann, in Proc. Am. Acad.* vii. (1867) 168.—Ins. Sandvic.
divaricatum, *Seem. in Journ. Bot.* iii. (1865) 78.—Java.
elatum. *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 728.—Reg. Himal.
elatum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 80.—Afr. trop.
ellipticum, Seem. in Journ. Bot. iii. (1865) 78 = venulosum.
emarginatum, *Seem. l. c.* 80.—Zeylan.
euryhynchum, *Seem. l. c.* 78.—Java.
exaltatum, Seem. l. c. 80 = Wallichianum.
fastigiatum, *Seem. l. c.* 78.—Java.
fimbriatum, *F. Muell. Papuan Pl.* 89.—N. Guin.
glaucum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 728.—Reg. Himal.
glaucum, Kurz, For. Fl. Brit. Burma, i. 588 = elatum.
gracile, *Blume, ex Seem. in Journ. Bot.* iii. (1865) 78.—Borneo.
heterophyllum, *Seem. l. c.* 77.—Malaya.
hypoleucum, *Kurz, For. Fl. Brit. Burma*, i. 589.—Reg. Himal.; Burma.
impressum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 728.—Reg. Himal.
insularum, *Seem. in Journ. Bot.* iii. (1865) 80.—Ins. Philipp.
Junghuhnianum, *Seem. l. c.* 77.—Java.
kavalense, *H. Mann, in Proc. Am. Acad.* vii. (1867) 168.—Ins. Sandvic.
khasianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 730.—Reg. Himal.
longifolium, *Seem. in Journ. Bot.* iii. (1865) 79.—Java.
macrophyllum, Wall. Cat. n. 4918 = venulosum.
Mannii, *Benth. & Hook. f. Gen.* i. 942.—Afr. trop.
micranthum, *Seem. in Journ. Bot.* iii. (1865) 78.—Sumatra.
octophyllum, *Benth. & Hook. f. Gen.* i. 942.—China.
oxyphyllum, *Seem. in Journ. Bot.* iii. (1865) 80.—Sumatra.
parasiticum, *Seem. l. c.* 79.—Malaya.
pergameceum, *Hassk. Cat. Hort. Bog. Alt.* 165.—Jave.
petiolosum, *Seem. in Journ. Bot.* iii. (1865) 78.—Borneo.
politum, *Seem. l. c.*—Borneo.
polyboryum, *Seem. l. c.*—Java.
polyphyllum, *Seem. l. c.* 79.—Java.
racemosum, *Bedd. Fl. Sylv.* t. 214.—Ind. or.
rigidum, Hassk. Cat. Hort. Bog. Alt. 165 = longifolium.
rigidum, *Seem. in Journ. Bot.* iii. (1865) 79.—Java.
rostratum, *Bedd. Fl. Sylv.* ii. 122.—Ind. or.
scandens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 80.—Afr. trop.
scandens, *Seem. in Journ. Bot.* iii. (1865) 79.—Malaya.
schizophyllum, *Hance, l. c.* xv. (1877) 334.—China.
serratum, *Seem. l. c.* iii. (1865) 79.—Java.
singalangense, *Seem. l. c.* 78.—Sumatra.
stellatum, *Gaertn. Fruct.* ii. t. 178. f. 3 (= acutangulum in textu, p. 472).—Zeylan.
subavene, *Hassk. Cat. Hort. Bog. Alt.* 165.—Java.
subalatum, *Seem. in Journ. Bot.* iii. (1865) 78.—Malaya.

HEPTAPLEURUM :—
tomentosum, *Hassk. Cat. Hort. Bog. Alt.* 165; *Buch.-Ham. ex C. B. Clarke, in Journ. Linn. Soc.* xv. (1876) 151.—Malaya.
venulosum, *Seem. in Journ. Bot.* iii. (1865) 80.—As. et Austral. trop.
verticillatum, *Serm. l. c.* 80.—Ins. Timor.
Waimese, *Wawra, in Flora*, lvi. (1873) 159.—Ins. Sandvic.
Wallichianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 730.—Ind. or.
Wallichianum, Seem. in Journ. Bot. iii. (1865) 80 = khasianum.

HEPTAPTERA, Marg. & Reut. in Mém. Soc. Phys. Genèv. viii. (1838) 302. t. 5 = **Prangos**, Lindl. (Umbellif.).
colladonioides, Marg. & Reut. l. c. = P. colladonioides.

HEPTARINA, Rafin. Fl. Tellur. iii. 15 (1836) = **Polygonum**, Tourn.
orientalis, Rafin. l. c. = P. orientale.

HEPTARINIA, Rafin. New Fl. Am. iv. 46 (1836) = **Polygonum**, Tourn.
longifolia, Rafin. l. c. = Am. bor.

HEPTAS, Meisn. Gen. 298; Comm. 302 (1840) = **Herpestis**, Gaertn. (Scrophularin.).
repens, Meisn. Comm. 302 = Herpestis Monnieria.

HEPTEIRECA, Rafin. Sylva Tellur. 126 (1838) = **Cassia**, Tourn. (Legumin.).
glandulosa, Rafin. l. c. 127 = C. virgata.

HEPTRILIS, Rafin. Fl. Tellur. 89 (1836) = **Leucas**, R. Br. (Labiat.).
glabrata, Rafin. l. c. = L. glabrata.

HERACANTHA, Hoffmgg. & Link, Fl. Port. ii. 205 (1830) = **Carthamus**, Tourn. (Compos.).
armeniia, G. Don, ex Loud. Hort. Brit. 382 = C. flavescens.
creticа, Hoffmgg. & Link, Fl. Port. ii. 207 = C. leucocaulos.
glauca, Sweet, Hort. Brit. ed. I. 216 = C. glaucus.
lanata, Hoffmgg. & Link, Fl. Port. ii. 206 = C. lanatus.
taurica, Link, Enum. Hort. Berol. ii. 304 = C. lanatus.

HERACLEA, Hill, Veg. Syst. iv. 37 (1762) = **Centaurea**, Linn. (Compos.).

HERACLEUM, Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 921.
Barysoma, Bunge, Del. Sem. Hort. Dorp. (1839) 7.
Sphondylium, [Tourn.] Adans. Fam. ii. 100 (1763).
Stenotaenia, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 329.
Symphyoloma, C. A. Mey. Verz. Pfl. Casc. 127 (1831).
Tordyliopsis, DC. Prod. iv. 199 (1830).
Trigonocladium, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 344.
Wendia, Hoffm. Fl. Umbell. Gen. 126 (1814).
absinthifolium, Vest. Choriz. *Z. Zoolina orientalis.*
acuminatum, Schleich. Cat. Pl. Helv. ed. IV. 19 = H. Sphondylium.
aestivum, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 318 = H. Sphondylium.
alpinum, Georgi, Beschr. Russ. Reich. iv. 849 = ligusticifolium.
alpinum, *Linn. Sp. Pl.* 250.—Europ.
alpinum, Pollini, Fl. Veron. i. 339 = palmatum.
alpinum, Siev. in Pall. N. Nord. Beitr. vii. 247 (Quid?). —Sibir.
amanum, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* ii. 1045.—Syria.
amplifolium, Lapeyr. Hist. Abr. Pl. Pyr. 153 = pyrenaicum.
amplissimum, *Wender. Ind. Sem. Hort. Marb.* (1840). —*Cf. Linnaea*, xv. (1841) *Litt.* 101.—Hab. ?
angustatum, Bor. Fl. Cent. Fr. ed. III. ii. 769, in obs. = H. Sphondylium.
angustifolium, Bieb. Fl. Taur. Cauc. i. 223 = chorodannum.

HERACLEUM:—
angustifolium, Jacq. Enum. Vindob. 45, 217 = H. Sphondylium.
angustifolium, Linn. Mant. i. 57 = sibiricum.
anisactis, *Boiss. & Hohen. Diagn.* Ser. I. x. 41.—Persia.
apiifolium, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 332.—Armenia.
aquilegifolium, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 715.—Ind. or.
arcticum, *Rupr. in Beitr. Pfl. Russ. Reich.* ii. 37, in textu.—Kom. arct.
argenteum, *Boiss. & Bal. Diagn.* Ser. II. vi. 87.—As. Min.
armeniacum, Bor. ex Nym. Consp. 290 = H. Sphondylium.
asperum, *Bieb. Fl. Taur. Cauc. Suppl.* 234.—Reg. Caucas.
asperum, Mert. & Koch, Deutschl. Fl. iii. 274 = palmatum.
aureum, Sibth. & Sm. Fl. Graec. t. 282 = Malabaila aurea.
auritum, Bischoff, in Linnaea, xiv. (1840) Litt. 131 = lanatum.
australe, Hartm. Svensk Excurs. Fl. 89 = H. Sphondylium.
austriacum, Habl. Taur. 162 = sibiricum.
austriacum, *Linn. Sp. Pl.* 249.—Austr.; Hungar.
barbatum, Ledeb. Fl. Alt. i. 300 = lanatum.
basicum, *Fisch. Hort. Gorenk.* (1812) 46, nomen (Quid?).—Sibir.
bifarium, Fisch. ex Reichb. Fl. Germ. Excurs. 458 = flavescens.
bipinnatum, Gmel. ex DC. Prod. iv. 195 = minimum.
Brunii, Retzl. Nom. ed. I. 400 = sec. *Brance-Ursina*, Adl. Fl. Pedem. ii. 4 = H. Sphondylium.
brevivittatum, *Ledeb. Fl. Ross.* ii. 294.—Reg. Caucas.
brignolaefolia, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 297.—Turkestan.
Brunonis, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 718.—Reg. Himal.
burmanicum, *Kurs, in Journ. As. Soc. Beng.* xli (1872) ii. 309; et (1877) ii. 116.—Reg. Himal.; Burma.
cachemiricum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 712.—Reg. Himal.
canariense, Cholay, ex DC. Prod. iv. 190 = Astydamia canariensis.
candicans, *Wall. Cat.* n. 573.—Reg. Himal.
canescens, *Lindl. in Royle, Illustr. Bot. Himal.* 232. —Reg. Himal.
Carmeli, Hort. ex Wender. in Linnaea, xv. (1841) Litt. 101 = amplissimum.
Carmeli, *Labill. Ic. Pl. Syr. Dec.* v. 8.—Syria.
caspicum, *Hort. Berol. ex DC. Prod.* iv. 192.—Reg. Casp.
caucasicum, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 329 = pubescens.
ceylanicum, *Gardn. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 716.—Zeylan.
chorodanum, *DC. Prod.* iv. 194.—Reg. Caucas.
cinereum, Lindl. in Royle, Illustr. Bot. Himal. 232 = canescens.
Clausii, *Ledeb. Fl. Ross.* ii. 293.—Rossia.
concanense, *Dalz. in Hook. Kew Journ.* ii. (1850) 260. —Ind. or.
cordatum, *Presl, Delic. Prag.* 135.—Sicilia.
crenatifolium, *Boiss. Fl. Orient.* ii. 1045.—As. Min.
cuneifolium, *DC. Prod.* iv. 194.—Sibir.
cyclocarpum, *C. Koch, in Linnaea*, xvi. (1842) 361.—Reg. Caucas.
dactylites, Fisch. ex Hoffm. Umbellif. 124 = villosum.
delphinense, Jord. Pugill. 75 = H. Sphondylium.
dichotoma, *Benth. & Hook. f. Gen.* ii. 922.—Orient.
dissectum, Ledeb. Fl. Alt. i. 301 = lanatum.
diversifolium, Wall. Cat. n. 574 = Wallichii.
Douglasii, DC. Prod. iv. 198 = lanatum.
dubium, Tenore, Fl. Nap. t. 221 = setosum.
dulce, Fisch. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 26 = lanatum.
elegans, Jacq. Fl. Austr. t. 175 = H. Sphondylium.
eminens, Lange, Ind. Sem. Hort. Haun. (1861) 20 = platytaenium.
festuosum, Salisb. Prod. 164 = H. Sphondylium.
flavescens, *Bunge, Alsin. Gen. Comm. Strip. Transc.* i. 214. —Rossia; As. bor.
flavescens, Bess. Enum. Fl. Volh. 12 = sibiricum.
flavescens, Willd. Sp. Pl. i. 1481 = H. Panaces.

HERACLEUM:—
gigantеum, Fisch. Hort. Gorenk. (1812) 46 = villosum.
glaberrimum, Boiss. & Hohen. Diagn. Ser. I. x. 41 = persicum.
granatense, *Boiss. Elench.* 49.—Hispan.
grandiflorum, Dals. & Gibs. Bomb. Fl. 108 = concanense.
grandiflorum, *Stev. ex Bieb. Fl. Taur. Cauc.* iii. 227. —Reg. Caucas.
graveolens, Spreng. Umbellif. Prod. 12 = Pastinaca graveolens.
gummiferum, Willd. Enum. Hort. Berol. i. 58, 54 = pubescens, asperum.
hirsutum, Edgew. in Trans. Linn. Soc. xx. (1846) 57 = canescens.
Hookerianum, *Wight & Arn. Prod.* 373.—Ind. or.
humile, *Sibth. & Sm. Fl. Graec. Prod.* i. 198.—As. Min. ; Syria.
hypoleucum, *Vis. in Linnaea*, xviii. (1844) 512.—Ind. or.
incanum, *Boiss. & Huet, Diagn.* Ser. II. ii. 98.—Armen.
intermedium, Gaud. Fl. Helv. ii. 319 = pyrenaicum.
intermedium, Ledeb. Fl. Ross. ii. 328 = splitolium.
Jacquemontii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 719.—Reg. Himal.
jugatum, *Boiss. Fl. Orient.* ii. 1043.—Reg. Caucas.
laciniatum, Desf. Tabl. ed. I. 119 = H. Sphondylium.
laciniatum, Hornem. Hort. Hafn. i. 279 = villosum.
lanatum, *Michx. Fl. Bor. Am.* i. 166.—Am. bor.; As. occ.
lasiopetalum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 332.—Persia.
latifolium, Gilib. Fl. Lituan. ii. 7 = H. Sphondylium.
Lecokii, Gren. & Godr. Fl. Fr. i. 695 = sibiricum.
Lecoquii, Mart. Fl. Tam. 255 = prasc.
Lehmannianum, *Bunge, Del. Sem. Hort. Dorp.* (1849) 2.—Turkestan.
Leichtlini, *Hort. ex Vathe, in Ind. Sem. Hort. Berol.* (1876) *App.* 2.—Hab. ?
leucocarpum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 155.—Afghan.
ligusticifolium, *Bieb. Fl. Taur. Cauc.* i. 224; *Suppl.* 237.—Reg. Caucas.
longifolium, Bieb. l. c. 223 = chorodanum.
longifolium, Jacq. Fl. Austr. ii. 46 = H. Sphondylium.
longifolium, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 329 = grandiflorum.
marginatum, Fisch. ex Hoffm. Umbellif. 124 = villosum.
microcarpum, *Franch. Pl. Davidian.* i. (1844) 144.—China.
minimum, *Lam. Fl. Fr.* iii. 418.—Alp. Delphin.
Moellendorffi, Hance, in Journ. Bot. xvi. (1878) 12 = lanatum.
montanum, Georgi, ex Ledeb. Fl. Ross. ii. 327 = ligusticifolium.
montanum, Schleich. Cat. Pl. Helv. ed. IV. 19 = pyrenaicum.
nepalense, *D. Don, Prod. Fl. Nep.* 135.—Reg. Himal.
nubigenum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 713.—Reg. Himal.
obtusifolium, *Wall. Cat.* n. 577.—Reg. Himal.
occidentale, Bor. Fl. Cent. Fr. ed. III. ii. 767 = H. Sphondylium.
Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 58.—Kokan.
Orphanidis, *Boiss. Fl. Orient.* ii. 1041.—Macedon.
Orsinii, Guss. Fl. Rar. Sic. 183. t. 37 = palmatum.
palmatifidum, Fisch. & Avé-Lall. in Ledeb. Fl. Ross. ii. 327 = cyclocarpum.
palmatilobum, *DC. Prod.* iv. 192.—Hab. ?
palmatum, *Bunge, Enum. Strip. Transc.* i. 215.—Orens.
Panaces, DC. ex Nym. Consp. 289 = setosum.
Panaces, Gmeldenst. Reisen, i. 435 (Quid?).—Imeretia.
Panaces, Koch, Syn. Fl. Germ. ed. II. 288 = palmatum.
Panaces, *Linn. Sp. Pl.* 249.—Europ. austr.
Panaces, Pall. ex Habl. Taur. 149 = villosum.
Panaces, Steph. Fl. Mosq. n. 179 = H. Sphondylium.
Panaces, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 329 = pubescens.
Pastinaca, *Fenzl, in Russegg. Reis. ii.* 968.—As. Min.

HERACLEUM:—
pastinacifolium, *C. Koch, in Linnæa*, xvi. (1842) 360. —Armenia.
pedatum, *Wight, Ic.* t. 842.—Ind. or.
persicum, *Desf. Cat. Hort. Par.* ed. III. 452.—Oriens.
pimpinellifolium, *Rupr. Sert. Tianach.* 49.—Turkestan.
pimpinellifolium, Spreng. Umb. Prod. 12 = Malabaila pimpinellifolia.
Pinda, *Dalz. & Gibs. Bomb. Fl.* 107.—Ind. or.
pinnatum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 712.—Reg. Himal.
platyphyllum, Ram. ex DC. Prod. iv. 198 = pyrenaicum.
platyphyllum, Schloss. & Vukot. ex Nym. Consp. 289 = palmatum.
platysaemium, *Boiss. in Ann. Sc. Nat. Sér.* III. i. (1844) 231.—As. Min.
Pollinianum, Bertol. Fl. Ital. iii. 432 = palmatum.
pratense, Jord. ex F. Schultz, Arch. Fl. Fr. et Allem. 316 = H. Sphondylium.
propinquum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 155.—Afghan.
prostrâtiforme, Crantz, Stirp. Austr. ed. I. fasc. iii. 11; ed. II. 155 = H. Panaces, H. Sphondylium.
pubescens, *Bieb. Fl. Taur. Cauc. Suppl.* 225.—As. Min.
pulchellum, Pall. ex Ledeb. Fl. Ross. ii. 227 = liguisticifolium.
pulchrum, *Zumag. Fl. Pedem.* ii. 8 (Quid ?).—Italia.
pumilum, Vill. Prosp. 28; Hist. Pl. Dauph. ii. 640 = minimum.
pumilum, Viv. ex DC. Prod. iv. 190 = Malabaila pumila.
pusillum, Pall. ex Ledeb. Fl. Ross. ii. 227 = liguisticifolium.
pyrenaicum, Bieb. Fl. Taur. Cauc. i. 224 = villosum. pyrenaicum, *Lam. Encyc.* i. 403.—Europ.
rodolens, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 316 = setosum.
rigens, *Wall. Cat.* n. 575; *DC. Prod.* iv. 191.—Ind. or.
rosæum, Stev. in *Mém. Soc. Nat. Mosc.* iii. (1812) 260 = chordalanum.
setosum, *Lapeyr. Hist. Abr. Pl. Pyr.* 158.—Europ. austr.
sibiricum, *Linn. Sp. Pl.* 249.—Europ.; As. bor.
sifolium, Reichb. Fl. Germ. Excurs. 456 = austriacum.
simplicifolium, Herb. ex Nym. Consp. 289 = palmatum.
sisiarense, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 101 = pastinacifolium.
spathiphyllum, *C. Koch, ex Boiss. Fl. Orient.* ii. 1042. —As. Min.
speciosum, Hoffm. ex Link, Handb. i. 356 = sec. ?
speciosum, Ledeb. Hort. Dorpat. 77 = pubescens.
speciosum, Rudolph, in Cat. Hort. Gorenk. (1812) 46 = villosum.
Sphondylium, Bourg. ex Reut. in Bull. Soc. Bot. Fr. xi. (1864) 46 = grunatense.
Sphondylium. Clans, in Goebel, Reise, ii. 276 = Clausii.
Sphondylium, *Linn. Sp. Pl.* 249.—Europ.; As. bor.
Sphondylium, Pall. Reise, ii. 838; Chamm. & Schlecht. in Linnæa, i. (1826) 294 = lanatum.
Sphondylium, Ucria, ex Guss. Fl. Sic. Prod. i. 864 = cordatum.
Sprengelianum, Dalz. & Gibs. Bomb. Fl. 108 = rigens.
Sprengelianum, Thw. Enum. Pl. Zeyl. 131 = ceylanicum.
Sprengelianum, *Wight & Arn. Prod.* 372.—Ind. or.
stenophyllum, Jord. Grain. Jard. Gren. (1849) 16 = H. Sphondylium.
sublineare, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 713.—Reg. Himal.
subtilifolium, Stev. in Nouv. Mém. Soc. Nat. Mosc. iii. (1829) 279 = villosum.
succisum, Fries, Summa, 181; ex Nym. Consp. 290 = H. Sphondylium.
tauricum, Stev. ex Reichb. Fl. Germ. Excurs. 457 = H. Sphondylium.
testiculatum, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 43 = elegans.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 712.—Reg. Himal.

HERACLEUM:—
tomentosum, Dalz. & Gibs. Bomb. Fl. 108, 213 = Pimpinella tomentosa.
tomentosum, Sibth. & Sm. Fl. Græc. Prod. i. 292 = Zosima orientalis.
Tordylium, Spreng. Umbellif. Sp. 49 = Tordylium maximum.
trachyloma, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 29 = pubescens.
tragodes, Benth. & Hook. f. Gen. i. 924 = Zosima tragodes.
transsilvanicum, Schur, Enum. Pl. Transs. 267 = palmatum.
tuberosum, *Molina, ex Willd. Sp. Pl.* i. 1428.—Chili.
umbellatum, *Gueldenst. Reisen*, i. 422.—Imeretia.
umbonatum, *Boiss. Fl. Orient.* ii. 1042.—Reg. Caucas.
urninum, Wallr. Beitr. Fl. Hercyn. 190 = H. Sphondylium ?
verrucosum, Stev. ex Steud. Nom. ed. II. i. 750 (Quid ?). —Tauria.
vescum, Willd. ex Steud. l. c. ed. I. 400.—Hab. ?
villosum, *Fisch. ex Steud. l. c.; et ex Spreng. in Schult. Syst.* vi. 579.—Reg. Caucas.
Wallichii, *DC. Prod.* iv. 195.—Reg. Himal.
Wilhelmsii, Fisch. & Mey. Ind. Sem. Hort. Petrop. vii. 51 = pubescens.
zosimioides, *Benth. & Hook. f. Gen.* i. 922.—As. Min.

HERBERTIA, Sweet, Brit. Flow. Gard. t. 222 (1827). *IRIDACEÆ*. [Benth. & Hook. f. iii. 691].
ALOPHIA, Herb. in Bot. Mag. sub t. 3779 (1838).
CIPURA, Klotzsch, ex Klatt, in Abh. Naturf. Ges. Halle, xii. 262 (1852).
LARENTIA, Klatt, l. c. (1862).
TRIFURCIA, Herb. in Bot. Mag. sub t. 3779 (1838).
amœna, *Griseb. in Goett. Abh.* xxiv. (1879) 325.—Reg. Argent.
brasiliensis, *Baker, in Journ. Linn. Soc.* xvi. (1877) 134.—Bras.
caerulea, *Herb. in Bot. Mag.* t. 3862.—Texas ?
cærulea, Klatt, in Linnæa, xxxi. (1861–62) 554 = pulchella.
Drummondiana, *Herb. in Bot. Reg.* (1842) *Misc.* 65. —Texas.
Drummondiana, Klatt, in Linnæa, xxxi. (1861–62) 555 = brasiliensis.
erythrina, *Griseb. in Goett. Abh.* xix. (1874) 270.—Reg. Argent.
lineata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 268.—Chili.
pulchella, *Sweet, Brit. Flow. Gard.* t. 222.—Chili.
pusilla, *Sweet, Hort. Brit.* ed. II. 497.—Bras.
stricta, *Griseb. in Goett. Abh.* xxiv. (1879) 324.—Reg. Argent.
umbellata, *Klatt, in Linnæa*, xxxi. (1861–62) 555.—Bras.
unguicalata, *Baker, in Journ. Linn. Soc.* xvi. (1877) 134, nomen.—Bras.

HERBICHIA, Zawadski, in Guill. Arch. Bot. i. (1832) 450; et Enum. Pl. Galic. 198 (1835) = **Senecio**, Tourn. (Compos.).
abrotanifolia, Zawadski, ll. cc. 451, 198 = S. abrotanifolius.
carpatica, Steud. Nom. ed. II. i. 750 = S. abrotanifolius.

HERDERIA, Cass. in Dict. Sc. Nat. lx. 599 (1830). *COMPOSITÆ*, Benth. & Hook. f. ii. 382.
scaposa, *Benth. in Hook. Niger Fl.* 425.—Afr. trop.
truncata, *Cass. in Dict. Sc. Nat.* lx. 586.—Afr. trop.

HERETIERA, G. Don, Gen. Syst. i. 516 (1831) = **Heritiera**, Dryand. (Sercul.).

HERICINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 235 = **Rauwenulfia**, Tourn.
arvensis, Fourr. l. c. = R. arvensis.

HERINCQUIA, Decne. ex Jacques & Hérincq, Man. Pl. ii. 562 [1850] = **Pennisetaphis**, Lindl. (Gen. nerac.).
floribunda, Decne. l. c. = P. floribunda.

HERISSANTHIA, Steud. Nom. ed. II. i. 750 (1840) = sec.

HERISSANTIA, Medic. Phil. Bot. i. 90 (1789) = **Abutilon**, Tourn. (Malvac.).
crispa, Medic. l. c. = A. crispum.

HERITERA, Stokes, Bot. Mat. Med. ii. 565 (1812) = **Heritiera**, Dryand. (Sercul.).

HERITERIA, Dum. Anal. Fam. 46 (1829) = **Heritiera**, Dryand. (Sercul.).

HERITERIA, Dum. Anal. Fam. 58 (1829) = HERITIERA, J. F. Gmel. = **Lachnanthus**, Ell. (Hæmodorac.). (Sercul.).

HERITERIA, Schrank, Baier. Fl. i. 133, 629 (1789) = **Tofieldia**, Huds. (Liliac.).
anthericoides, Schrank, l. c. 629 = T. calyculata.

HERITIERA, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 546 (1789). *STERCULIÁCEÆ*, Benth. & Hook. i. 219.
ATUNA, Rafin. Sylva Tellur. 158 (1838).
BALANOPTERIS, Gaertn. Fruct. ii. 94. tt. 98, 99 (1791).
FOMETICA, Rafin. Sylva Tellur. 75 (1838).
acuminata, *Wall. ex Voigt, Hort. Suburb. Calc.* 103, nomen; *et ex Kurz, in Journ. Bot.* xii. (1874) 65.—Ind. or.
attenuata, Wall. Cat. n. 1140 = Brownlowia lanceolata.
dubia, *Wall. ex Kurz, in Journ. Bot.* xii. (1874) 65.—Ind. or.
Fischeri, *Regel & Rach, Ind. Sem. Hort. Petrop.* (1858) 48.—As. trop.
Fomes, [*Buch.-Ham. in*] *Syme, Embassy*, ed. II. iii. 319.—Ind. or.; Malaya.
Fomes, Wall. Cat. n. 1139, partim = littoralis.
grandis, Fisch. ex Steud. Nom. ed. I. 400 (Quid ?).—Hab. ?
lanceolata, A. Gray, Bot. U. St. Expl. Exped. i. 184, lapsu = Niotæ sp.
littoralis, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 546. —Geront. trop.
macrophylla, *Wall. ex Voigt, Hort. Suburb. Calc.* 108, nomen; *et ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 61.—Ind. or.; Burma.
minor, *Lam. Encyc.* iii. 229.—Ins. Maurit.
minor, Roxb. Hort. Beng. 50; Fl. Ind. iii. 142 = H. Fomes.
Papilio, *Bedd. Fl. Sylvat.* t. 218.—Ind. or.
spectabilis, Ball. Hist. Fl. iv. 62 = Octolobus spectabilis.
tinctoria, *Blanco, Fl. Filip.* ed. I. 653.—Ins. Philipp.
Tothila, Kurz, in Journ. As. Soc. Beng. xlii. (1874) ii. 118 = littoralis.
Vespertilio, Bedd. ex Kurz, in Journ. Bot. xii. (1874) 65 = H. Papilio.

HERITIERA, J. F. Gmel. Syst. 113 (1791) = **Lachnanthes**, Ell. (Hæmodorac.).
Gmelini, Michx. Fl. Am. Bor. i. 21 = L. tinctorium.
tinctorium, J. F. Gmel. Syst. 113 = L. tinctorium.

HERITIERA, Retz. Obs. vi. 17. t. 1 (1791) = **Alpinia**, Linn. (Schamin.).
alba, Retz. l. c. 18 = A. alba.
Allughas, Retz. l. c. 17 = A. Allughas.
aquatica, Retz. l. c. 18 = A. aquatica.
chinensis, Retz. l. c. = A. chinensis.

HERMANNIA, Linn. Syst. ed. I (1735). *STERCULIÁCEÆ*, Benth. & Hook. f. i. 232.
EURYNEMA, Endl. Gen. Suppl. ii. 93 (1842).
GILBSIA, F. Muell. Fragm. iv. 41 (1875).
TRICHANTHERA, Ehrenb. in Linnæa, iv. (1829) 401.
abrotanoides, *Schrad. Hort. Getting.* 17. t. 11.—Afr. austr.
alnifolia, Drège, ex Harv. & Sond. Fl. Cap. i. 189 = cuneifolia.
alnifolia, *Linn. Sp. Pl.* 674.—Afr. austr.
alnifolia, Wall. Cat. n. 1196 K = Sida Schimperiana.
althæifolia, *Linn. Sp. Pl.* 673; ed. II. 941.—Afr. austr.
althæoides, Eckl. & Zeyh. Enum. 43, partim = candicans.
althæoides, Hort. ex Link, Enum. Hort. Berol. ii. 179 = incana.

HERMANNIA :—
angularis, Eckl. & Zeyh. Enum. 45 = hyssopifolia.
angularis, *Jacq. Hort. Schoenb.* i. 68. t. 126.—Afr. austr.
angustifolia, *Otto & Dietr. Allg. Gartenz.* viii. (1840) 382.—Hab.?
anthemifolia, *Presl, Bot. Bemerk.* 22.—Afr. austr.
arabica, *Hochst. & Steud. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* vi. 51.—Arab.
argentea, Eckl. & Zeyh. Enum. 47 = pulverata.
argentea, *Sm. in Rees, Cycl.* xvii.—Afr. austr.
argyrata, Presl, Bot. Bemerk. 20 = decumbens.
aspera, Wendl. Bot. Beobacht. 52 = scabra.
aurea, Jacq. Hort. Schoenb. ii. 46 = althaeifolia.
betonicaefolia, *Eckl. & Zeyh. Enum.* 46.—Afr. austr.
bicornis, Eckl. & Zeyh. l. c. 49 = trifurca.
bifaria, *Linnæus in Act. Soc. Sc. Fenn.* x. (1871) 121.—Hab.?
bifora, Eckl. & Zeyh. Enum. 44 = scabra.
bipinnata, Burch. Trav. i. 310 = abrotanoides.
biserrata, Linn. f. Suppl. 302 = Mahernia biserrata.
boraginiflora, *Hook. Ic. Pl.* t. 597.—Afr. austr.
brachypetala, Harv. in Harv. & Sond. Fl. Cap. i. 202 = Mahernia tomentosa.
bracteosa, *Presl, Bot. Bemerk.* 21.—Afr. austr.
bryoniaefolia, *Burch. Trav.* ii. 258.—Afr. austr.
bryonifolia, Eckl. & Zeyh. Enum. 49 = diversistipula.
carnulea, Burm. f. Fl. Cap. Prod. 18 = trifurcata.
cardicans, *Ait. Hort. Kew.* ed. I. ii. 412.—Afr. austr.
candida, Salisb. Prod. 365 = prace.
candidissima, *Spreng. f. Tent. Suppl.* 18.—Afr. austr.
capensis, *L. f. Genel. Syst.* 1013.—Afr. austr.
Cavanillesiana, *Eckl. & Zeyh. Enum.* 45.—Afr. austr.
cernua, *Thunb. Diss. Herm.* 11.—Afr. austr.
chrysanthemifolia, *E. Mey. in Drège, Zwei Pfl. Docum.* 191, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 204.—Afr. austr.
chrysophylla, *Eckl. & Zeyh. Enum.* 44.—Afr. austr.
ciliaris, Linn. f. Suppl. 302 = Mahernia verticillata.
chytiaefolia, *Otto & Dietr. Allg. Gartenz.* viii. (1840) 382.—Afr. austr.
collina, Eckl. & Zeyh. Enum. 41 = decumbens.
comosa, *Burch. ex DC. Prod.* i. 498.—Afr. austr.
conglomerata, *Eckl. & Zeyh. Enum.* 44.—Afr. austr.
cordifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 193.—Afr. austr.
cornuofolia, Eckl. & Zeyh. Enum. 47 = tenuifolia.
coronopifolia, *Link, Enum. Hort. Berol.* ii. 161.—Afr. austr.
cuneifolia, *Jacq. Hort. Schoenb.* i. 66. t. 124.—Afr. austr.
decipiens, *E. Mey. in Drège, Zwei Pfl. Docum.* 191, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 192.—Afr. austr.
decumbens, Drège, ex Presl, Bot. Bemerk. 21 = Mahernia grandistipula.
decumbens, *Willd. Enum. Hort. Berol. Suppl.* 46, nomen; *Spreng. Neue Entdeck.* i. 299.—Afr. austr.
denudata, Eckl. & Zeyh. Enum. 43 = Prealiana.
denudata, *Linn. f. Suppl.* 301.—Afr. austr.
desertorum, *Eckl. & Zeyh. Enum.* 46.—Afr. austr.
diffusa, Linn. f. Suppl. 302 = Mahernia diffusa.
diffusa, E. Mey. in Drège, Zwei Pfl. Docum. 191 = Mahernia parviflora.
discolor, *Otto & Dietr. Allg. Gartenz.* viii. (1840) 314.—Afr. austr.
disermaefolia, Eckl. & Zeyh. Enum. 41 = althaeifolia.
disermaefolia, *Jacq. Hort. Schoenb.* i. 65. t. 131.—Afr. austr.
dissecta, Harv. in Harv. & Sond. Fl. Cap. i. 204 = chrysanthemifolia.
distichа, *Schrad. Sert. Hannov.* 28. t. 16.—Afr. austr.
diversifolia, Eckl. & Zeyh. Enum. 41 = althaeifolia.
diversistipula, *Presl, Bot. Bemerk.* 21.—Afr. austr.
Drogeana, *Presl*, l. c. 20.—Afr. austr.
dryadifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 191.—Afr. austr.
Echinoizoea, Presl, Bot. Bemerk. 20 = candicans.
ecosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 191, nomen.—Afr. austr.
exstipulata, *E. Mey.* l. c., nomen; *et ex Harv. & Sond. Fl. Cap.* i. 203.—Afr. austr.
falcata, Eckl. & Zeyh. Enum. 46 = flammea.
fasciculata, Baker, in Saund. Refug. Bot. t. 289 = linearifolia.
ferruginea, Hort. ex Link, Enum. Hort. Berol. ii. 179 = incana.

HERMANNIA :—
filifolia, Drège, ex Harv. & Sond. Fl. Cap. i. 195 = pallens.
filifolia, *Linn. f. Suppl.* 302.—Afr. austr.
filipes, *Harv. in Harv. & Sond. Fl. Cap.* i. 206.—Afr. austr.
flabelloides, Eckl. & Zeyh. Enum. 49 = candicans.
flammea, *Jacq. Hort. Schoenb.* i. 69. t. 139.—Afr. austr.
Flammula, *Harv. in Harv. & Sond. Fl. Cap.* i. 196.—Afr. austr.
floribunda, *Harv.* l. c. 201.—Afr. austr.
fragrans, Hort. ex Link, Enum. Hort. Berol. ii. 179 = althaeifolia.
garlepina, Eckl. & Zeyh. Enum. 49.—Afr. austr.
geniculata, Eckl. & Zeyh. l. c. 40 = Mahernia chrysantha.
Gerardii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 588.—Afr. austr.
Gilesii, *F. Muell. Fragm.* ix. 42, in obs.—Austral.
glabrata, Linn. f. Suppl. 301 = Mahernia glabrata.
gladiolifolia, *Link, Enum. Hort. Berol.* ii. 179.—Afr. austr.
glauca, *Hort. Herrenh. ex Otto & Dietr. Allg. Gartenz.* viii. (1840) 380.—Afr. austr.
glauca, Salisb. Prod. 366 = denudata.
glomerata, E. Mey. in Drège, Zwei Pfl. Docum. 191 = conglomerata.
grandiflora, *Jacq. Hort. Schoenb.* i. 67. t. 129.—Afr. austr.
grandifolia, Eckl. & Zeyh. Enum. 45.—Afr. austr.
grandiflora, Ait. Hort. Kew. ed. II. iv. 141 = Mahernia grandiflora.
grossulariifolia, Linn. Sp. Pl. 678 = Mahernia heterophylla.
halicacabo, DC. Prod. i. 498 = abrotanoides.
heterophylla, *Thunb. Diss. Herm.* 14.—Afr. austr.
hirsuta, Drège, ex Harv. & Sond. Fl. Cap. i. 189 = alnifolia.
hirsuta, Eckl. & Zeyh. Enum. 48 = prismatocarpa.
hirsuta, *Will. Gard. Dict.* ed. VIII. n. 8.—Afr. austr.
hirsuta, *Schrad. Sert. Hannov.* 10. t. 4.—Afr. austr.
hirta, Sparm. ex Steud. Nom. ed. II. i. 751 = scabra.
hispida, *Thunb. Diss. Herm.* 8.—Afr. austr.
hispidula, *Reichb. fc. Pl. Cult.* i. t. 68.—Afr. austr.
holoseriсea, *Jacq. Hort. Schoenb.* iii. 28. t. 292.—Afr. austr.
humilis, E. Mey. in Drège, Zwei Pfl. Docum. 191 = Mahernia oblongifolia.
humilis, *Thunb. Diss. Herm.* 14.—Afr. austr.
hyssopifolia, *Linn. Sp. Pl.* 674.—Afr. austr.
ignescens, *Presl, Bot. Bemerk.* 21.—Afr. austr.
imbricata, Eckl. & Zeyh. Enum. 48 = trifoliata.
incana, *Cav. Diss.* 328.—Afr. austr.
incana, Eckl. & Zeyh. Enum. 42 = candicans.
incana, Thunb. Fl. Cap. ed. Schult. 505 = holoseriсea.
incisa, *Willd. Sp. Pl.* iii. 599.—Hab.?
infusa, *Link & Otto in. Pl. Hor.* 85. t. 28.—Mexic.
involucrata, Cav. Diss. 328 = salvifolia.
Jourbertiana, *Harv. in Harv. & Sond. Fl. Cap.* i. 196.—Afr. austr.
Kirkii, *Mast. in Oliver, Fl. Trop. Afr.* i. 238.—Afr. trop.
latifolia, *Eckl. & Zeyh. Enum.* 44.—Afr. austr.
latifolia, Jacq. Hort. Schoenb. i. 64 = salvifolia.
lavandulifolia, *Linn. Sp. Pl.* 674.—Afr. austr.
lepidota, Buch. ex Krauss, in Flora, xxvii. (1844) 294 = pallens.
leucanthemifolia, *Otto & Dietr. Allg. Gartenz.* viii. (1840) 330.—Afr. austr.
leucanthemoides, *Presl, Bot. Bemerk.* 22.—Afr. austr.
leucocephala, *Presl*, l. c. 20.—Afr. austr.
linearifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 205.—Afr. austr.
linifolia, *Burm. f. Fl. Cap. Prod.* 18.—Afr. austr.
linifolia, E. Mey. ex Harv. & Sond. Fl. Cap. i. 206 = linearifolia.
macrophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 216.—Afr. austr.
melochioides, *Burch. ex DC. Prod.* i. 494.—Afr. austr.
micans, Schrad. Sert. Hannov. 11. t. 5 = salvifolia.
micropetala, *Harv. in Harv. & Sond. Fl. Cap.* i. 201.—Afr. austr.
modesta, *Planch. in Ann. Sc. Nat. Sér.* IV. iii. (1855) 292.—Afr. trop.
mollis, Willd. Enum. Hort. Berol. 692 = candicans.
mucronulata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 217.—Afr. austr.
multicaulis, E. Mey. in Drège, Zwei Pfl. Docum. 191 = Mahernia erodioides.
multifida, DC. Prod. i. 498 = abrotanoides.

HERMANNIA :—
multiflora, Drège, ex Harv. & Sond. Fl. Cap. i. 190 = pallens.
multifora, Eckl. & Zeyh. Enum. 46 = alnifolia.
multiflora, Jacq. Hort. Schoenb. i. 69. t. 128 = cuneifolia.
muricata, Eckl. & Zeyh. Enum. 48 = hirsuta, *Schrad.*
myrrhifolia, *Thunb. Diss. Herm.* 16.—Afr. austr.
nemorosa, Eckl. & Zeyh. Enum. 43 = candicans.
odorata, Ait. Hort. Kew ed. I. ii. 413 = lavandulaefolia.
orophila, *Eckl. & Zeyh. Enum.* 47.—Afr. austr.
pallens, *Eckl. & Zeyh. l. c.* 48.—Afr. austr.
paniculata, *Franch. in Rétouil, Cumali*, 19.—Afr. trop.
paniculata, E. Mey. in Drège, Zwei Pfl. Docum. 191 = scordifolia.
parviflora, E. Mey. l. c. 192 = Mahernia parviflora.
passerinaeformis, Eckl. & Zeyh. Enum. 47 = filifolia.
passerinoides, Harv. in Harv. & Sond. Fl. Cap. i. 195, sphalm. = prace.
patula, Harv. l. c. 187 = Prealiana.
pauciflora, *J. Wats. in Proc. Am. Acad.* xvii. (1881–82) 368.—Arizona.
paucifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 218 = chrysanthemifolia.
pinnata, Linn. Sp. Pl. 674 = Mahernia pinnata.
plicata, Ait. Hort. Kew. ed. I. ii. 411 = althaeifolia.
plicata, Eckl. & Zeyh. Enum. 43 = candicans.
plicata, E. Mey. in Drège, Zwei Pfl. Docum. 192 = leucophylla.
polymorpha, Eckl. & Zeyh. Enum. 46 = flammea.
praemorsa, Eckl. & Zeyh. l. c. 43 = cuneifolia.
praemorsa, Wendl. Bot. Beobacht. 51 = candicans.
pratensis, Eckl. & Zeyh. Enum. 41 = decumbens.
Prealiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 259.—Afr. austr.
prismatocarpa, *E. Mey. in Drège, Zwei Pfl. Docum.* 192, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 186.—Afr. austr.
procumbens, *Cav Diss.* 329.—Afr. austr.
procumbens, E. Mey. ex Harv. & Sond. Fl. Cap. i. 210, 211 = Mahernia humifusa, vesicaria.
prostrata, *E. Mey. in Drège, Zwei Pfl. Docum.* 192, nomen.—Afr. austr.
pulchella, Linn. f. Suppl. 302 = Mahernia pulchella.
pulverata, *Andr. Bot. Rep.* iii. t. 161.—Afr. austr.
pulverulenta, DC. Prod. i. 496 = prace.
punctata, *Presl, Bot. Bemerk.* 21.—Afr. austr.
purpurea, Hort. ex Otto & Dietr. Allg. Gartenz. viii. (1840) 313 = diserniaefolia.
Quartinianа, A. Rich. Tent. Fl. Abyss. i. 75 = Mahernia abyssinica.
quercifolia, Eckl. & Zeyh. Enum. 49 = denudata.
racemosa, E. Mey. in Drège, Zwei Pfl. Docum. 192 = garlepina.
rigida, *Harv. in Harv. & Sond. Fl. Cap.* i. 188.—Afr. austr.
rigida, Nocca, ex Steud. Nom. ed. II. i. 751 = distichа.
rotundata, E. Mey. in Drège, Zwei Pfl. Docum. 192 = Mahernia linearifolia.
rotundifolia, Jacq. Hort. Schoenb. i. 68 = distichа.
salicifolia, Harv. in Harv. & Sond. Fl. Cap. i. 199 = mucronulata.
salsifolia, Eckl. & Zeyh. Enum. 45 = mucronulata.
salvifolia, *Linn. f. Suppl.* 302.—Afr. austr.
Sandersoni, *Harv. in Harv. & Sond. Fl. Cap.* i. 200.—Afr. austr.
scabra, *Cav. Diss.* 331. t. 183. f. 2.—Afr. austr.
scabra, Jacq. Hort. Schoenb. i. 66. t. 127 = hirsuta, *Schrad.*
scoparia, *Harv. in Harv. & Sond. Fl. Cap.* i. 194.—Afr. austr.
scordifolia, Eckl. & Zeyh. Enum. 48 = Prealiana.
scordifolia, *Jacq. Hort. Schoenb.* i. 64. t. 120.—Afr. austr.
secundiflora, Eckl. & Zeyh. Enum. 48 = pallens.
spinosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 193, nomen; *et ex Harv. in Harv. & Sond. Fl. Cap.* i. 208.—Afr. austr.
stipulacea, *Lehm. ex Echl. & Zeyh. Enum.* 44.—Afr. austr.
stricta, *Harv. in Harv. & Sond. Fl. Cap.* i. 206.—Afr. austr.
suavis, *Presl, Bot. Bemerk.* 21.—Afr. austr.
sulcata, *Harv. in Harv. & Sond. Fl. Cap.* i. 200.—Afr. austr.
tenuifolia, *Sims, Bot. Mag.* t. 1248.—Afr. austr.
tenuifolia, *Presl, Bot. Bemerk.* 22.—Afr. austr.
texana, *A. Gray, Gen. Illustr.* ii. 56. t. 125.—Texas.

HERMANNIA :—
tigrensis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 74.—Abyss.
trifoliata, *Linn. Mant.* ii. 481.—Afr. austr.
trifurca, Eckl. & Zeyh. Enum. 47 = H. Flammula, angularis.
trifurca, *Linn. Pl. Rar. Afr.* 13; *Amoen. Acad.* vi. 90. —Afr. austr.
trifurcata, Linn. Sp. Pl. ed. II. 942 = prsec.
triquetrifolia, *Presl, Bot. Bemerk.* 21 = Mahernia grandistipula.
triphylla, *Cav. Diss.* 338.—Afr. austr.
triphylla, Drège, ex Harv. & Sond. Fl. Cap. i. 198 = ternifolia.
triphylla, Linn. Fl. Afr. Rar. 13; Amoen. Acad. vi. 91 = Lotononis polycephala?
velutina, *DC. Prod.* i. 495.—Afr. austr.
venosa, *Hort. Gotting. ex Otto & Dietr. Allg. Gartenz.* viii. (1840) 315.—Afr. austr.
venosa, *Thunb. Diss. Herm.* 12.—Afr. austr.
verrucosa, Eckl. & Zeyh. Enum. 48 = hirsuta, *Schrad.*
verticillata, Berg. Desc. Pl. Cap. 169 = Mahernia pinnata.
vesicoria, Cav. Diss. 331 = Mahernia vesicaria.
vestita, E. Mey. in Drège, Zwei Pfl. Docum. 192 = candidissima.
vestita, *Thunb. Diss. Herm.* 13.—Afr. austr.
Zeyheri, Walp. Ann. i. 108, sphalm. = soq.
Zeyheriana, *Presl, Bot. Bemerk.* 22.—Afr. austr.

HERMAS, Linn. Mant. ii. 168 (1771). *UMBELLIFERAE*, Benth. & Hook. f. i. 878.
australis, Spreng. ex Schult. Syst. vi. 880 = Panax australis.
capitata, *Linn. f. Suppl.* 295.—Afr. austr.
ciliata, *Linn. f. l. c.* 296.—Afr. austr.
depauperata, *Linn. Mant.* ii. 299.—Afr. austr.
gigantea, *Linn. f. Suppl.* 295.—Afr. austr.
minima, Eckl. & Zeyh. Enum. 354 = capitata.
quercifolia, Eckl. & Zeyh. l. c. 355 = quinquedentata.
quinquedentata, *Linn. f. Suppl.* 296.—Afr. austr.
*rudis*ima, Reichb. in Spreng. Syst. iv. Cur. Post. 118 = Lichtensteinia lacera.
wiltenbergensis, Eckl. & Zeyh. Enum. 354 = ciliata.
villosa, Thunb. in Nov. Act. Petrop. xiv. (1805) 581 = depauperata.

HERMBSTAEDTIA, Reichb. Consp. 164 (1828). *AMARANTACEAE*, Benth. & Hook. f. iii. 25.
BEZELIA, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 299.
HYPARRTE, Rafin. Fl. Tellur. iii. 43 (1836).
LANGIA, Endl. Gen. 804 (1837).
PELIANTHUS, E. Mey. ex Moq. in DC. Prod. xiii. 11. 246 (1849).
caffra, *Moq. in DC. Prod.* xiii. 11. 246.—Afr. austr.
elegans, *Moq. l. c.* 247.—Afr. austr.
glauca, *Reichb. ex Steud. Nom.* ed. II. i. 752.—Afr. austr.

HERMESIA, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 809 (1805) = **Alchornea**, Sw. (Euphorb.).
banksiaefolia, Spreng. Syst. ii. 246 = Callicoma serratifolia.
castaneaefolia, Humb. & Bonpl. ex. Willd. Sp. Pl. iv. 809 = A. castaneaefolia.
mexicana, Hook. & Arn. Bot. Beech. Voy. 309 = Bernardia mexicana.
salicifolia, Baill. Etud. Gén. Euph. 447 = A. castaneaefolia.

HERMESIAS, Loefl. It. 278 (1758) = Brownea, Jacq. (Legumin.).

HERMIDIUM, S. Wats. in Bot. King's Exped. 286. t. 32 (1871). *NYCTAGINEAE*, Benth. & Hook. f. iii. 6.
alipes, *S. Wats. l. c.*—Am. bor. occ.

HERMINTIERA, Guill. & Perr. Fl. Seneg. Tent. 201. t. 51 (1888). *LEGUMINOSAE*, Benth. & Hook. f. i. 515.
ARDISSONIA, Kotschy, in Oesterr. Bot. Zeitschr. viii. (1854) 134. t. 1.
elaphriorylon, *Guill. & Perr. l. c.*—Afr. trop.

HERMINIUM, Linn. Syst. ed. I (1735). *ORCHIDEAE*, Benth. & Hook. f. iii. 522.
ADELA, Lindl. Bot. Reg. sub t. 1701 (1836).
HERMINIUM :—
CHAMAERREPES, Spreng. Syst. iii. 709 (1826).
CHAMORCHIS, Rich. in Mém. Mus. Par. iv. (1818) 49.
CYBELE, Falc. in Lindl. Veg. Kingd. 558 (1847).
MONORCHIS, Ehrh. Beitr. iv. (1789) 147.
THERMIA, Falc. in Lindl. Veg. Kingd. 553 (1847).
alpinum, *Lindl. Bot. Reg.* sub t. 1499.—Europ.
angustifolium, *Benth. & Hook. f. Gen.* iii. 622.—Reg. Himal.
clandestinum, Gren. & Godr. Fl. Fr. iii. 299 = H. Monorchis.
congestum, *Lindl. in Wall. Cat.* n. 7068.—Reg. Himal.
constrictum, Lindl. Bot. Reg. sub t. 1499 = (*Peristylus constrictus*).—Reg. Himal.
cordatum, Lindl. l. c.=Habenaria cordata.
fallax, *Lindl. in Wall. Cat.* n. 7412.—Reg. Himal.
goodyeroides, Lindl. l. c. n. 7066 = Habenaria goodyeroides.
gramineum, *Lindl. l. c.* n. 7413.—Reg. Himal.
Hamiltonianum, Lindl. l. c. n. 7069 (= *Peristylus Hamiltonianus*).—Reg. Himal.
Josephi, *Reichb. f. in Flora*, lv. (1872) 276.—Reg. Himal.
latifolium, Lindl. Bot. Reg. sub t. 1499 = Habenaria chlorantha.
Lefebreanum, *Reichb. f. in Linnaea*, xxv. (1852) 282. —Abyssin.
longicrure, *Benth. & Hook. f. Gen.* iii. 622.—Japon.
Monorchis, *R. Br. in Ait. Hort. Kew.* ed. II. v. 191.—Europ.; As. bor.
natalense, *Reichb. f. Otia Bot. Hamb.* 108.—Afr. austr.
plantagineum, Lindl. in Bot. Reg. sub t. 1499 = Habenaria Wightii.
reniforme, *Lindl. in Wall. Cat.* n. 7067.—Reg. Himal.
spirale, Reichb. f. in Bonplandia, iii. (1855) 218 = Habenaria spiralis, *A. Rich.*
Steudelianum, *Reichb. f. Otia Bot. Hamb.* 109.—Abyssin.
unalaschense, Reichb. f. Ic. Fl. Germ. xiii. et xvi. 107. t. 65 = congestum.

HERMIONE, Salisb. in Trans. Hort. Soc. i. (1812) 257 = **Narcissus**, Tourn. (Amaryll.).
aequilimbe, Herb. Amaryll. 404 = N. Tazetta.
alba, Haw. Monog. Narciss. 8 = N. intermedius.
ambigena, Salisb. in Trans. Hort. Soc. i. (1812) 361 = N. Tazetta.
amoena, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 117 = N. Tazetta.
ancops, M. Roem. Syn. Enstat. 229 = N. papyraceus.
angustifolia, M. Roem. l. c. 234 = N. angustifolius.
antiphonensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 111 = N. Tazetta.
aporticorona, Haw. Monog. Narciss. 9 = N. Tazetta.
aureaincrena, Haw. l. c. 11 = N. Tazetta.
aurea, Jord. & Fourr. Ic. Pl. Eur. i. 65 = N. aureus.
autumnalis, M. Roem. Syn. Enstat. 231 = N. elegans.
Bertolonii, Jord. & Fourr. Ic. Pl. Eur. i. 64. t. 193 = N. Tazetta.
Biancae, Tod. ex Rent. in Linnaea, xxix. (1857–58) 746 = N. Tazetta.
bifrons, Salisb. in Trans. Hort. Soc. i. (1812) 359; Haw. Revis. Pl. Succ. 138 = N. intermedius.
brevifrons, Haw. Monog. Narcias. 11 = N. Tazetta.
bromiciformis, Herb. Amaryll. 234 = M. Tazetta?
calliche, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 119 = N. Tazetta.
calliopsis, Jord. & Fourr. l. c. 117 = N. Tazetta.
candida, M. Roem. Syn. Enstat. 216 = N. moschatus.
candidissima, M. Roem. l. c. 228 = N. papyraceus.
cerina, Haw. Syn. Pl. Succ. 229 = N. Tazetta.
chaetocrena, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 117 = N. Tazetta.
chlorotica, Jord. & Fourr. l. c. 116 = N. Tazetta.
chrysantha, Haw. Monog. Narcias. 12. n. 41 = N. Tazetta.
citrina, Haw. Revis. Pl. Succ. 143.—N. Tazetta.
compressa, Haw. l. c. 139 = N. intermedius.
concolor, M. Roem. Syn. Enast. 214 = N. homochroos.
contorta, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 113 = N. Tazetta.
corcyrensis, Herb. Amaryll. 333. t. 37. f. 2 = N. Tazetta.
corrugata, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 115 = N. Tazetta.
HERMIONE :—
crenularis, Salisb. in Trans. Hort. Soc. i. (1812) 363 = Narcissus Trewianus.
crenulata, Haw. Monog. Narcias. 10. n. 23 = N. Tazetta.
crispicorona, Haw. l. c. 11 = N. Tazetta.
croceo-cincta, M. Roem. Syn. Enast. 234 = N. croceocinctus.
cupularis, Salisb. in Trans. Hort. Soc. i. (1812) 361 = N. Tazetta.
Cypri, Haw. in Phil. Mag. (1831) 184 = N. Tazetta.
debilis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 114 = N. Tazetta.
decora, Haw. Monog. Narcias. 9 = N. Tazetta.
deficiens, Kunth, Enum. Pl. v. 750 = N. serotinus.
defloxicaulis, Haw. Monog. Narcias. 9 = N. Tazetta.
discolor, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 111 = N. Tazetta.
discreta, Jord. & Fourr. l. c. 113 = N. Tazetta.
dubia, Haw. Monog. Narcias. 12 = N. Tazetta.
elegans, Haw. l. c. 18 = N. elegans.
erodora, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 120 = N. Tazetta.
fistulosa, Haw. Syn. Pl. Succ. 230 = N. Tazetta.
flava, M. Roem. Syn. Enast. 220 = N. Jonquilla.
flaveola, Haw. Monog. Narcias. 8 = N. Tazetta.
flexiflora, Haw. l. c. 10 = N. Tazetta.
floriunda, Salisb. in Trans. Hort. Soc. i. (1812) 362; Haw. Revis Pl. Succ. 141 = N. Tazetta.
floxlactis, Haw. Monog. Narcias. 8 = N. Tazetta?
formosa, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 115 = N. Tazetta.
formosissima, Tod. Nouv. Gen. e Spec. 29 = N. Tazetta?
fulgida, Jord. & Fourr. Brev. Pl. Nov. fasc. ii.120 = N. Tazetta.
ganymedioides, Jord. & Fourr. l. c. 113 = N. Tazetta.
gracilis, M. Roem. Syn. Enast. 232 = N. gracilis.
grandicrenata, Biancs, ex Parl. Fl. Ital. iii. 154 = N. Tazetta.
grandiflora, Haw. Syn. Pl. Succ. 230 = N. Trewianus.
hololeucs, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 109 = N. Tazetta.
homochroos, M. Roem. Syn. Enast. 229 = N. jonquilloides.
insolita, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 114 = N. Tazetta.
intermedia, Haw. Monog. Narcias. 7 = N. Tazetta.
italica, Haw. Revis. Pl. Succ. 144 = N. Tazetta.
italica, Herb. Amaryll. 325 = N. Tazetta.
jasminea, Salisb. in Trans. Hort. Soc. i. (1812) 360 = N. Tazetta.
Jonquilla, Haw. Revis. Pl. Succ. 137 = N. Jonquilla.
jucunda, Jord. & Fourr. Brev. Pl. Nouv. fasc. ii. 115 = N. Tazetta.
juncifolia, Salisb. in Trans. Hort. Soc. i. (1812) 357 = N. Jonquilla.
latifolia, Haw. Revis. Pl. Succ. 146 = N. Tazetta.
leucotheia, Salisb. in Trans. Hort. Soc. i. (1812) 359 = N. seroticulia.
littoralis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 113 = N. Tazetta.
lobata, Jord. & Fourr. l. c. = N. Tazetta.
longiflora, M. Roem. Syn. Enast. 218 = N. longiflorus.
Luna, Haw. Revis. Pl. Succ. 143 = N. Tazetta.
lutea, Haw. Monog. Narcias. 10 = N. Tazetta.
luteola, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 118 = N. Tazetta.
mediterranea, Jord. & Fourr. l. c. 110 = N. Tazetta.
micrantha, Jord. & Fourr. l. c. 108 = N. Tazetta.
modesta, Jord. & Fourr. l. c. 113 = N. Tazetta.
monspeliensis, Jord. & Fourr. l. c. 110 = N. Tazetta.
multiflora, Haw. Monog. Narcias. 6 = N. Tazetta.
neapolitana, M. Roem. Syn. Enast. 230 = N. Tazetta.
neglecta, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 111 = N. Tazetta.
nivea, M. Roem. Syn. Enast. 230 = N. papyraceus.
nobilis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 118 = N. Tazetta.
nutans, M. Roem. Syn. Enast. 215 = N. nutans.
obliterata, Haw. Monog. Narcias. 13 = N. Broussonnetii.
obsoleta, Haw. Revis. Pl. Succ. 147 = N. elegans.
ochroleuca, M. Roem. Syn. Enast. 219 = N. Tazetta.
pallida, M. Roem. l. c. 217 = N. Tazetta.
papyracea, Schult. f. Syst. vii. 1783, in obs. = soq.
papyratia, Haw. Revis. Pl. Succ. 143 = N. papyraceus.

HERMIONE :—
petula, Haw. Monog. Narciss. 11 = Narcissus Tazetta.
perinlea, Haw. l. c. 8 = N. Tazetta.
planicorona, M. Roem. Syn. Ensat. 233 = N. gracilis.
polyantha, Haw. Monog. Narciss. 11; Jord. & Fourr. Ic. Pl. Eur. i. 69 = N. Tazetta.
praecox, Haw. l. c. 13. n. 43 = N. Tazetta.
pratensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 109 = N. Tazetta.
primulina, Haw. Syn. Pl. Succ. 209 = N. intermedius.
pulchella, M. Roem. Syn. Ensat. 215 = N. pulchellus.
pumila, M. Roem. l. c. 234 = N. pumilus.
purpuro-cincta, M. Roem. l. c. 234 = N. purpuro-cinctus.
radiata, M. Roem. l. c. 217 = N. intermedius.
rotularis, Haw. Monog. Narciss. 8 = N. rotularis.
saguentis, Tod. Ind. Sem. Hort. Panorm. (1858) 27; et in Ann. Sc. Nat. Sér. IV. xi. (1859) 879 = N. Tazetta.
serotina, Haw. Monog. Narcias. 13 = N. serotinus.
sertulina, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 118 = N. Tazetta.
sesiobata, M. Roem. Syn. Ensat. 223 = N. Tazetta.
similis, Salisb. in Trans. Hort. Soc. i. (1812) 358 = N. Jonquilla.
solaris, Haw. Monog. Narcias. 9 = N. Tazetta.
splendens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 119 = N. Tazetta.
stellaris, Salisb. in Trans. Hort. Soc. i. (1812) 359 = N. bifrons.
straminea, Haw. Monog. Narcias. 12 = N. Tazetta.
stylosa, Salisb. in Trans. Hort. Soc. i. (1812) 360 = N. Tazetta.
subalbida, Haw. Monog. Narcias. 12 = N. Tazetta.
subcornata, Haw. l. c. 10 = N. Tazetta.
sublutea, Haw. l. c. 8 = N. Tazetta.
subcalcatis, Haw. l. c. 11 = N. Tazetta.
sulphurea, Haw. l. c. 10 = N. Tazetta.
syracusana, Tod. Ind. Sem. Hort. Panorm. (1858) 27; et in Ann. Sc. Nat. Sér. IV. xi. (1859) 879.—Sicilia.
Tangieri, Pritz. Ic. Ind. 542 = N. Broussonnetii ?
Tazetta, Haw. Revis. Pl. Succ. 142 = N. Tazetta.
tegulaeflora, Salisb. in Trans. Hort. Soc. i. (1812) 362 = N. gracilis.
tenuiflora, Haw. in Phil. Mag. (1830) 188 = N. Tazetta.
tenuior, Haw. Revis. Pl. Succ. 145 = N. gracilis.
tennis, M. Roem. Syn. Ensat. 238 = N. gracilis.
teretiaculis, Haw. Revis. Pl. Succ. 140 = N. Tazetta.
Trewiana, Sweet, Brit. Flow. Gard. Ser. II. t. 116 = N. Tazetta.
trifida, Haw. Monog. Narcias. 9 = N. Tazetta.
tripartita, M. Roem. Syn. Ensat. 217 = N. tripartitus.
tubulosa, Tod. Ind. Sem. Hort. Panorm. (1858) 27; et in Ann. Sc. Nat. Sér. IV. xi. (1859) 880 = N. Tazetta ?
unicolor, Haw. Monog. Narcias. 12 = N. Tazetta.
vernata, Tod. Nuov. Gen. e Spec. 15 = N. Tazetta ?
virginea, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 109 = N. Tazetta.
viridiflora, Haw. Revis. Pl. Succ. 147 = N. viridiflorus.
Weinmanni, M. Roem. Syn. Ensat. 233 = N. triandrus.
xanthica, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 116 = N. Tazetta.

HERMODACTYLOS, Reichb. Consp. 64 (1828) = seq.

HERMODACTYLUM, Bartl. Ord. 52 (1830) = seq.

HERMODACTYLUS, Reichb. Nom. 47 (1841) = Colchicum, Tourn. (Liliac.).
Ritchii, R. Br. ex Steud. Nom. ed. II. i. 753 = C. montanum.

HERMODACTYLUS, Tourn. ex Mill. Gard. Dict ed. VI. (1752). *IRIDEÆ*, Benth. & Hook. f. iii. 687.
bicaphaceus, Sweet, Brit. Flow. Gard. Ser. II. sub t. 146 = tuberosus.
longifolius, Sweet, l. c. t. 146 = tuberosus.
repens, Sweet, l. c. sub t. 146 = tuberosus.
tuberosus, *Mill. Gard. Dict.* ed. VIII.—Reg. Mediterr.

HERMSTAEDTIA, Steud. Nom. ed. II. i. 752 (1840) = Hermbstaedtia, Reichb. (Amarant.).

HERMUPOA, Loefl. It. 307 (1758) = Steriphoma, Spreng. (Capparid.).
Loefflingiana, DC. Prod. i. 254 = S. paradoxum.

HERNANDEZIA, Hoffmgg. Verz. Pfl. Nachtr. i. 219 (1824) = Hernandia, Plum. (Laurin.).
leucocephala, Hoffmgg. l. c. Nachtr. iii. 41 = Hernandia leucomphala.
sonora, Hoffmgg. l. c. Nachtr. i. 219 = Hernandia sonora.

HERNANDIA, Plum. ex Linn. Gen. ed. I. 374 (1737). *LAURINEÆ*, Benth. & Hook. f. iii. 164.
Hernandezia, Hoffmgg. Verz. Pfl Nachtr. i. 219 (1824).
Hernandiopsis, Meisan. in DC. Prod. xv. i. 264 (1864).
Hertelia, Neck. Elem. ii. 345 (1790).
bivalvis, *Benth. Fl. Austral.* v. 214.—Austral.
cordigera, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 92.—N. Caled.
cubensis, *Grisob. in Mem. Am. Acad.* N. S. viii. (1861) 188.—Ind. occ.
guianensis, *Aubl. Pl. Gui.* ii. 848.—Guian.
incompleta, *Steud. Nom.* ed II. i. 752.—Hab. ?
Moerenhoutiana, *Guill. in Ann. Sc. Nat.* Sér. II. vii. (1837) 189.—Ins. Tahiti.
ovigera, Lam. Encyc. iii. 123 = guianensis.
ovigera, *Linn. Diss. Herb. Amboin. in Amoen. Acad.* iv. 125.—Ind. occ.
peltata, *Meisan. in DC. Prod.* xv. i. 263.—Geront. trop.
sonora, *Linn. Sp. Pl.* 981.—Ind. or.
sonora, Miq. Fl. Ind. Bat. i. i. 887 = peltata.
sonora, Zoll. Syst. Verz. Arch. Pfl. 117 = ovigera.

HERNANDIOPSIS, Meisan. in DC. Prod. xv. i. 264 (1864) = Hernandia, Linn. (Laurin.).
Vieillardi, Meisan. l. c. = Hernandia cordigera.

HERNIARIA, [Tourn.] Linn. Syst. ed. I. (1735). *ILLECEBRACEÆ*, Benth. & Hook. f. iii. 16.
alpestris, Aubry, ex DC. Prod. iii. 367 = glabra.
alpestris, Lam. Encyc. iii. 125 = alpina.
alpina, Jord. ex DC. Prod. iii. 368 = incana.
alpina, *Vill. Hist. Pl Dauph.* ii. 556.—Europ.
alsines-folia, Mill. Gard. Dict. ed. VIII. n. 8 = Polycarpon tetraphyllum.
americana, Nutt. in Am. Journ. Sc. v. (1822) 291 = Siphonychia americana.
annua, Lag. Gen. et Sp. Nov. 12 = cinerea.
arenaria, Kuntze, Taschen-Fl. Leipz. 224 = glabra.
argnea, *Boiss. Fl. Orient.* i. 741.—Cappadoc.
bactica, *Boiss. & Reut. Diagn.* Ser. II. i. 95.—Hispan.
Bourgi, Fisch. ex Hornem. Hort. Hafn. Suppl. 157 = incana.
Boissieri, J. Gay, in Duch. Rev. Bot. ii. (1846–47) 370 = alpina.
cachemiriana, *J. Gay, l. c.*—Cachemir.
capensis, Steud. Nom. ed. I. 401; Bartl. in Linnaea, vii. (1833) 624 = lenticulata.
canescens, *Kupr. Fl. Casuc.* 241.—Reg. Caucas.
ciliata, Bab. in Trans. Linn. Soc. xvii. (1837) 453 = glabra.
ciliata, [Clairv.] Man. Herb. 71 = alpina.
cinerea, *DC. Fl. Fr. Suppl.* 275.—Europ. austr.
diandra, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 296 = cinerea.
dichotoma, *DC. Prod.* iii. 368.—Hab. ?
erecta, Desf. Fl. Atlant. i. 214 = Paronychia suffruticosa.
eustachya, *Reichb. f. in Ber. Deutsch. Bot. Ges.* iii. (1885) 274.—Ins. Porto Rico.
flavescens, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) reimpr. 21 = cinerea.
Fontanesii, *Y. Gay, in Duch. Rev. Bot.* ii. (1846–47) 371.—Afr. bor. occ.
frigida, J. Gay, ex Kunze, in Flora, xxix. (1846) 650 = alpina.
fruticosa, Delile, Fl. Egypt. Illustr. 57 = hemistemon.
fruticosa, Desf. Fl. Atlant. i. 213 = Fontanesii.
fruticosa, Falk, Bekr. ii. 139 = polygama.
fruticosa, Gouan, Fl. Monsp. 358 = glabra.
fruticosa, *Linn. Cent. Pl.* i. 6 ; *Amoen. Acad.* iv. 269. Reg. Mediterr. occ.
glabra, *Linn. Sp. Pl.* 218.—Europ. ; As. bor.
hemistemon, *Y. Gay, in Duch. Rev. Bot.* ii. (1846–47) 371.—Afr. bor. ; Arab.
hirsuta, Bieb. Fl. Taur. Cauc. i. 179 = incana.
hirsuta, *Linn. Sp. Pl.* 218.—Europ. ; As. Min. ; Ind. or.
hirsuta, Urv. in Mém. Soc. Linn. Par. i. (1822) 284 = incana.

HERNIARIA :—
hirsuta, Viv. Fl. Lib. Spec. 15 = cinerea.
filecebroides, *C. Sm. in Tuckey, Congo*, 250 ; *et ex DC. Prod.* iii. 366.—Ins. Cap. Verd.
incana, Boiss. Voy. Bot. Espagne, 218 = baetica.
incana, *Lam. Encyc.* iii. 124.—Europ. ; Orient.
Joannensa, Schult. Syst. vi. 297 = Paronychia suffruticosa.
latifolia, *Lapeyr. Hist. Abr. Pl. Pyr.* 197.—Mont. Pyren.
lenticularis, Hill, Veg. Syst. xxvi. 9 = *lenticulata*, Linn. ?
lenticulata, All. Fl. Pedem. ii. 209 = alpina.
lenticulata, Huds. Fl. Angl. 94 = Polycnemum Rail.
lenticulata, Linn. Sp. Pl. 218 = incana, Cressa cretica.
lenticulata, *Thunb. Prod. Fl. Cap.* 48.—Afr. austr.
macrocarpa, Sibth. & Sm. Fl. Graec. Prod. i. 167 ; Fl. Graec. t. 258 = incana.
maritima, Link, ex Nym. Consp. 256 Add. = glabra.
microcarpa, Presl, Delic. Prag. 65 = glabra.
Millegrana, Bess. Cat. Hort. Cremn. (1816) 68 ; Link, Enum. Hort. Berol. i. 247 ; Pall. ex Spreng. f. Tent. Suppl. 178 = incana.
monsperlina, *Reichb. f. in Ber. Deutsch. Bot. Ges.* iii. (1885) 274.—Ins. Porto Rico.
multicaulis, *Kit. ex Kan. in Verh. Zool. Bot. Ges. Wien*, xii. (1862) 540.—Hungar.
nebrodensis, Jan, ex Nym. Consp. 256 Add. = glabra.
odorata, Andrz. ex J. Gay, in Duch. Rev. Bot. ii. (1846–47) 371 = polygama.
olympica, *Y. Gay, l. c.* 370.—Bithyn.
paniculata, Web. ex Hist. Hisp. 49 = Paronychia paniculata.
parnassica, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. i. 95.—Graec.
Payco, Molina, Sagg. Chile, 150 = Roubieva multifida.
permixta, Jan, ex Nym. Consp. 256 = parnassica.
polygama, *Y. Gay, in Duch. Rev. Bot.* ii. (1846–47) 371.—Soongaria.
polygonoides, Cav. Ic. ii. 27. t. 137 = Paronychia suffruticosa.
pyrenaica, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 224 = latifolia.
scabrida, *Boiss. Elench.* 42.—Hispan.
setigera, *Gill. in Hook. Bot. Misc.* iii. (1833) 337.—Chili.
sphacelata, Hochst. ex Webb, Fragm. Fl. Aethiop. 36 = hemistemon.
suffruticosa, Desf. Tabl. ed I. 45 = Paronychia suffruticosa.
verticillata, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 181 = fruticosa.
virescens, Salzm. ex DC. Prod. iii. 367 = cinerea.
vulgaris, Spreng. Syst. i. 929 = cinerea, glabra, hirsuta.

HERODIUM, Reichb. Ic. Fl. Germ. v. 21 nn. 4864–4870 (1841) = Erodium, L'Hérit. (Geraniac.).

HEROION, Rafin. Fl. Tellur. iv. 12 (1836) = Asphodeline, Reichb. (Liliac.).
filiformis, Rafin. l. c. = A. tenuior.

HEROTIUM, Steud. Nom. ed. II. i. 686, in syn. (1840) sphalm. = XEROTIUM, Bluff & Fingerh. = Filago, Linn. (Compos.).

HERPESTES, Kunth, Syn. Pl. ii. 124 (1823), sphalm. = seq.

HERPISTIS, Gaertn. f. Fruct. iii. 186. t. 214 (1805). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 951.
Ahisocalyx, Hance, in Walp. Ann. iii. 195 (1858).
Apozia, Willd. ex Steud. Nom. ed. II. i. 114 (1840).
Bhami, Adans. Fam. ii. 208 (1763).
Caconapea, Cham. in Linnaea, viii. (1833) 28.
Calytriflex, Rniz & Pav. Prod. Fl. Per. 96 (1794).
Cardiolophus, Griff. Notul. iv. 105 (1854).
Habbrzhamia, Rafin. Neogenyt. 2 (1825).
Hiptas, Meisn. Gen. 393 ; Comm. 202 (1840).
Hydrotrida, Willd. ex Steud. Nom. ed. II. i. 783 (1840).
Macuilzamia, Rafin. Neogenyt. 2 (1825).
Megardonia, Ruiz & Pav. Prod. Fl. Per. 95 (1794).
Mella, Vand. Fl. Lusit. & Bras. Spec. 43. f. 23 (1788).
Monniera, P. Br. Hist. Jamaic. 269 (1756).
Ranapalus, Kellogg, in Proc. Calif. Acad. vii. (1877) 118.
Ramaria, Cham. in Linnaea, viii. (1833) 20.

HERPESTIS:—
Septas, Lour. Fl. Cochinch. 392 (1790).
Septilia, Rafin. Fl. Tellur. iv. 66 (1836).
africana, Steud. Nom. ed. II. i. 753 = H. Monnieria.
amara, Benth. Scroph. Ind. 30 = Curanga amara.
amara, Span. in Linnæa, xv. (1841) 382 = floribunda.
amplexicaulis, *Pursh, Fl. Am. Sept.* ii. 418.—Am. bor.
amplexifolia, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1156, nomen.—Guian.
andicola, Gill. ex Benth. in DC. Prod. x. 374 = Melosperma andicola.
angulata, *Benth. l. c.* 395.—Bras.
arenaria, *Y. A. Schmidt, in Mart. Fl. Bras.* viii. 313. —Bras.
axillaris, *Benth. in DC. Prod.* x. 396.—N. Granat.
bacopoides, *Benth. l. c.* 399.—Bras.
Beccabunga, *Griseb. Cat. Pl. Cub.* 182.—Cuba.
Brownei, Steud. Nom. ed. I. 402; II. i. 753 = H. Monnieria.
Caconospora, Steud. l. c. ed. II. i. 753 = gratioloides.
cæspitosa, *Cham. in Linnæa*, viii. (1833) 33.—Bras.
callitrichoides, H. B. & K. Nov. Gen. et Sp. ii. 357 = Ilysanthes gratioloides.
calycina, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 57.—Afr. trop.
Calyptriplex, Steud. Nom. ed. II. i. 753 = H. Monnieria.
caprariicides, *H. B. & K. Nov. Gen. et Sp.* ii. 363.—N. Granat.
chamædrifolia, Steud. Nom. ed. II. i. 584, in syn. = seq.
chamædryoides, *H. B. & K. Nov. Gen. et Sp.* ii. 369. —Am. trop.
chrysantha, Cham. & Schlecht. in Linnæa, ii. (1827) 577 = chamædryoides.
colubrina, H. B. & K. Nov. Gen. et Sp. ii. 368 = chamædryoides.
connata, Spreng. Syst. iv. Cur. Post. 234 = Limnophila hypericifolia.
crenata, Beauv. Fl. Owar. t. 112 = H. Monnieria.
cubensis, Poepp. in Spreng. Syst. ii. 802 = chamædryoides.
cuneifolia, Pursh, Fl. Am. Sept. ii. 418 = H. Monnieria.
depressa, *Benth. in DC. Prod.* x. 395.—Bras.
diffusa, *Y. A. Schmidt, in Mart. Fl. Bras.* viii. 315.—Bras.
diffusa, Willd. ex Cham. & Schlecht. in Linnæa, iii. (1828) 7 = Stemodia parviflora.
divaricata, *Y. A. Schmidt, in Mart. Fl. Bras.* viii. 309. —Bras.
dominguensis, Spreng. Neue Entdeck. iii. 25 = stricta.
dubia, *Steud. Nom.* ed. II. i. 753; *D. Dietr. Syn. Pl.* iii. 519.—Reg. Argent.
elongata, Benth. in DC. Prod. x. 396 = stricta.
erecta, Spreng. Syst. ii. 801 = Bayrichia villosa.
flagellaris, *Cham. & Schlecht. in Linnæa*, ii. (1827) 576.—Bras.
flexilis, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1156, nomen.—Guian.
floribunda, *R. Br. Prod.* 442.—Geront. trop.
glechomoides, Spreng. Syst. iv. Cur. Post. 234 = Gerardia chamædryoides.
gracilis, *Benth. in DC. Prod.* x. 395.—Bras.
grandiflora, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 56.—Am. auct.
gratioloides, *Benth. l. c.* 57.—Am. trop.
Hamiltoniana, *Benth. Scroph. Ind.* 30.—As. et Afr. trop.
hermannioides, *Cham. in Linnæa*, viii. (1833) 34.—Bras.
humifusa, *Griseb. Cat. Pl. Cub.* 182.—Cuba.
imbricata, *Benth. in DC. Prod.* x. 397.—Guian.
javanica, Blume, Bijdr. 748 = Adenosma ovatum.
laucifolia, Wight, in Benth. Scroph. Ind. 30 = floribunda.
Imigura, *Cham. & Schlecht. in Linnæa*, ii. (1827) 573. —Bras.
laxiflorissima, Blume, Bijdr. 747 = Adenosma cæruleum.
laxiflora, *Benth. in DC. Prod.* x. 396.—Bras.
linearis, Spreng. Syst. ii. 802 = floribunda.
madagascariensis, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 56.—Madag.
marginata, Benth. in DC. Prod. x. 396 = laxiflora.
Matourea, Steud. Nom. ed. II. i. 753 = Stemodia pusilla.
micrantha, Ell. Sketch, ii. 105 = Micranthemum Nuttallii!
micrantha, Pursh, Fl. Am. Sept. ii. 418 = pusc.

HERPESTIS:—
micromonnieria, *Griseb. Cat. Pl. Cub.* 182.—Cuba.
Monnieria, *H. B. & K. Nov. Gen. et Sp.* ii. 366.—Cosmo.
monosticta, *Schlecht. in Linnæa*, xxiv. (1851) 694.—Bras.
montevidensis, *Spreng. Syst.* ii. 802.—Reg. Argent.
morannaia, H. B. & K. Nov. Gen. et Sp. ii. 367 = H. Monnieria.
myriophylloides, *Benth. in DC. Prod.* x. 398.—Bras.
nigrescens, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 56.—Am. bor.
obovata, *Poepp. ex Cham. & Schlecht. in Linnæa*, v. (1830) 107.—Cuba.
ovata, Benth. Scroph. Ind. 30 = Adenosma ovatum.
pedunculata, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 56.—Texas.
pedunculosa, Steud. Nom. ed. II. i. 753 = H. Monnieria.
pilosa, Benth. in Hook. Comp. Bot. Mag. ii. (1836) 57 = Mimulus pilosus.
pilosa, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 21.—Mexic.
polyadena, *Miq. in Linnæa*, xxii. (1849) 474.—Guian.
polycnemoides, Benth. in Hook. Comp. Bot. Mag. ii. (1836) 57 = stricta.
polygonoides, Benth. in Wall. Cat. n. 3897 = Torenia polygonoides.
prasiola, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 57.—Bras.
procumbens, Hort. Berol. ex Spreng. Nov. Prov. 22 = H. Monnieria.
pulcherrima, Griff. Notul. iv. 104 = Limnophila pulcherrima.
pusilla, Meisn. ex D. Dietr. Syn. Pl. iii. 519. n. 86 = serpyllioides.
pygmæa, Griff. Notul. iv. 104 = Limnophila micrantha, pygmæa.
radicata, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 56.—Patagonia.
Ranaria, *Benth. l. c.* 57.—Bras.
reptans, Benth. in DC. Prod. x. 399 = Myriophyllum proserpinacoides.
repens, *Cham. & Schlecht. in Linnæa*, v. (1830) 107. —Am. bor. et trop.
reptans, *Benth. in DC. Prod.* x. 395.—Bras.
rotundifolia, *Pursh, Fl. Am. Sept.* ii. 418.—Am. bor.
rugosa, Roth, Nov. Pl. Sp. 290 = Limnophila Roxburghii.
Salzmannii, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 58.—Bras.
scabra, Benth. l. c. 57 = laxiflora.
selaginoides, Steud. Nom. ed. II. i. 585, in syn. = Zalusianskya selaginoides.
semiserrata, Steud. l. c. 753 = gratioloides.
serpyllifolia, *Benth. in DC. Prod.* x. 396.—Bras.
serpyllioides, *Cham. & Schlecht. in Linnæa*, ii. (1827) 574.—Bras.
sessiliflora, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 58.—Mexic.
sessiliflora, *Voigt, Hort. Suburb. Calc.* 502 (Quid!). —Ind. or.
spathulata, Blume, Bijdr. 748 = H. Monnieria.
stellarioides, *Cham. in Linnæa*, viii. (1833) 32.—Bras.
stricta, *Schrad. in Link, Enum. Hort. Berol.* ii. 142.—Bras.
tenella, *Cham. & Schlecht. in Linnæa*, ii. (1827) 576. —Bras.
Thouningii, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 58.—Afr. trop.
tomentosa, Cham. & Schlecht. in Linnæa, v. (1830) 108 = Stemodia lanata.
Tweedlei, *Benth. in DC. Prod.* x. 397.—Reg. Argent.
vandellioides, *H. B. & K. Nov. Gen. et Sp.* ii. 367.—Mexic.

HERPETACANTHUS, Nees, in Mart. Fl. Bras. ix. 93. t. 19 (1847). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1102.
Schultzia, Nees, in Nov. Act. Nat. Cur. xi. (1828) 68.
decipiens, *Nees, in DC. Prod.* xi. 367.—Bras.
longiflorus, *Moric. Pl. Nouv. Am.* 159. t. 98.—Bras.
rascahensis, *Nees, in Mart. Fl. Bras.* ix. 95.—Bras.
macrophyllus, *Nees, l. c.*—Bras.
melancholicus, *Nees, l. c.* 98.—Bras.
rubiginosus, *Nees, l. c.* 94.—Bras.
Schultzii, *Nees, l. c.* 95.—Bras.

HERPETICA, Rafin. Sylva Tellur. 123 (1838) = **Cassia**, Tourn. (Legumin.).
alata, Rafin. l. c. = C. alata.

HERPETOSPERMUM, Wall. Cat. n. 6761 (1831!) *CUCURBITACEÆ*, Benth. & Hook. f. i. 834.
Kampinia, C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 129.
caudigerum, *Wall. Cat. n.* 6761.—Reg. Himal.
pedunculosum, C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 115 = caudigerum.

HERPOLIRION, Hook. f. Fl. N. Zel. i. 258 (1858). *LILIACEÆ*, Benth. & Hook. f. iii. 793.
capense, Bolus, in Journ. Linn. Soc. xviii. (1881) 395 = Nanolirion capense.
Novæ-Zelandiæ, *Hook. f. Fl. N. Zel.* i. 258.—N. Zel.; Tasman.
Tasmaniæ, Hook. f. l. c.; et Fl. Tasm. ii. 54 = Novæ-Zelandiæ.

HERPYSMA, Lindl. in Wall. Cat. n. 7389 (1831); et Bot. Reg. sub t. 1618 (1833). *ORCHIDEÆ*, Benth. & Hook. f. iii. 599.
longicaulis, *Lindl. in Wall. Cat.* n. 7389.—Reg. Himal.

HERPYZA, Sauv. Fl. Cub. 29 (1873). *LEGUMINOSÆ*.
grandiflora, *Sauv. l. c.*—Cuba.

HERRANIA, Goudot, in Ann. Sc. Nat. Sér. III. ii. (1844) 230. t. 5. *STERCULIACEÆ*, Benth. & Hook. f. i. 226.
Lighia, Schomb. in Rep. Brit. Assoc. (1844) 11. 71.
albiflora, *Goudot, l. c.*—N. Granat.
aspera, *Karst. in Linnæa*, xxvii. (1856) 446.—Am. trop.
laciniifolia, *Goudot, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvi. (1862) 337.—N. Granat.
Marise, *Goudot, in Ann. Sc. Nat.* Sér. III. ii. (1844) 233.—Bras.
pulcherrima, *Goudot. l. c.* 232.—N. Granat.

HERRARIA, Ritg. in Marb. Schr. ii. (1831) 132 = **Herreria**, Ruiz & Pav. (Liliac.).

HERRERA, Adans. Fam. ii. 158 (1763) = **Brithalis**, Linn. (Compos.).

HERRERIA, Ruiz & Pav. Prod. 48. t. 35 (1794). *LILIACEÆ*, Benth. & Hook. f. iii. 766.
Clara, Kunth, in Abh. Akad. Berl. (1843) 44.
brasiliensis, Hort. Berol. ex Kunth, Enum. Pl. v. 294 = H. Salsaparilha.
grandiflora, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 159.—Bras.
interrupta, Griseb. in Mart. Fl. Bras. iii. 1. 24 = H. Salsaparilha.
montevidensis, *Klotzsch, ex Griseb. l. c.*—Reg. Argent.
parviflora, Lindl. Bot. Reg. t. 1043 = H. Salsaparilha.
Salsaparilha, *Mart. Reise Bras.* i. 545.—Bras.
stellata, *Ruiz & Pav. Fl. Per.* iii. 69. t. 308.—Chili.
verticillata, Molina, Sagg. Chile, ed. II. 136 = stellata.

HERRMANNIA, Link & Otto, Ic. Pl. Rar. 55. t. 28 (1828) = **Hermannia**, Linn. (Stercul.).

HERSCHELIA, Bowdich, Excurs. Mader. 159 (1825) = **Physalis**, Linn. (Solanac.).
edulis, Bowdich, l. c. = P. edulis.

HERSCHELIA, Lindl. Gen. et Sp. Orch. 362 (1838) = **Disa**, Berg. (Orchid.).
barbata, Bolus, in Journ. Linn. Soc. xix. (1882) 336, 344 = D. barbata.
coelestis, Lindl. Gen. et Sp. Orch. 362 = D. graminifolia.
graminifolia, G. Don, in Sweet, Hort. Brit. ed. III. 651 = D. graminifolia.

HERSCHELLIA, Bartl. Ord. 194 (1830) = **Herschelia**, Bowdich. = **Physalis**, Linn. (Solanac.).

HERSILEA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 75 (1862) = **Setaria**, Tourn. (Compos.).
ramosa, Klotzsch, l. c. 76 = A. altaicus.
simplex, Klotzsch, l. c. 75 = A. molluscullus.

HERSILIA, Rafin. Fl. Tellur. iii. 88 (1836) = **Phlomis**, Linn. (Labiat.).
bilaba, Rafin. l. c. = *P. crinita*.

HERTELIA, Neck. Elem. ii. 345 (1790) = **Hernandia**, Linn. (Laurin.).

HERTIA, Neck. Elem. i. 8 (1790); Less. in Linnaea, vi. (1831) 94 = **Othonnopsis**, Jaub. & Spach (Compos.).
crassifolia, Less. l. c. = *O. chetrifolia*.

HESIODA, Vell. Fl. Flum. 185 (1825); iv. t. 140 (1827) = **Heisteria**, Jacq. (Olacin.).
perianthomega, Vell. Fl. Flum. 185; iv. t. 140.—Bras.

HESIODIA, Stend. Nom. ed. II. i. 758 (1840), sphalm. = *praec.*

HESIODIA, Moench, Meth. 391 (1794) = **Sideritis**, Tourn. (Labiat.).
hicolor, Moench, l. c. 392 = *S. montana*.
montana, Dum. Fl. Belg. 44 = *S. montana*.

HESPERALOË, Engelm. in S. Wats. Bot. King's Exped. 497 (1871). *LILIACEÆ*, Benth. & Hook. f. iii. 778.
Engelmanni, Kraskopf, ex Baker, in Journ. Linn. Soc. xviii. (1880) 231 = *yuccæfolia*.
yuccæfolia, *Engelm. in S. Wats. Bot. King's Exped.* 497.—Am. bor. occ.

HESPERANTHA, Ker-Gawl. in Kon. et Sims, Ann. Bot. i. (1805) 294. *IRIDEÆ*, Benth. & Hook. f. iii. 709.
acuta, Ker-Gawl. Irid. Gen. 91 = *longituba*.
angusta, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 295.—Afr. austr.
Baurii, *Baker, in Journ. Bot.* xiv. (1876) 182.—Afr. trop
bulbifera, *Baker, l. c.*—Afr. austr.
caricina, Klatt, in Abh. Natarf. Ges. Halle, xii. (1882) 295 = *radiata*.
ciliata, E. Mey. ex Klatt, l. c. 294 = *Geissorhiza ciliaris*.
cinnamomea, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 295.—Afr. austr.
crispa, *Eckl. Top. Verz.* 22.—Afr. austr.
cucullata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 293.—Afr. austr.
disticha, *Klatt, l. c.*—Afr. austr.
falcata, *Ker-Gawl. in Kon. & Sims, Ann. Soc.* i. (1805) 295.—Afr. austr.
flexuosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 294.—Afr. austr.
graminifolia, *Sweet, Hort. Brit.* ed. II. 503.—Afr. austr.
humilis, *Baker, in Journ. Bot.* xiv. (1876) 239.—Afr. austr.
hermenia, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 295.—Afr. austr.
latifolia, *Spreng. ex Stend. Nom.* ed. II. i. 758, nomen. —Afr. austr.
longituba, *Baker, in Gard. Chron.* (1877) i. 652; *Journ. Linn. Soc.* xvi. (1877) 96.—Afr. austr.
maritima, *Eckl. Top. Verz.* 23.—Afr. austr.
montana, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 293.—Afr. austr.
pallida, *Eckl. Top. Verz.* 23.—Afr. austr.
pentandra, Drège, ex Baker, in Journ. Linn. Soc. xvi. (1877) 168 = *Tritonia scillaris*.
Petitiana, *Baker, l. c.* 96.—Abyssin.
pilosa, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 295.—Afr. austr.
quinquangularis, Eckl. Top. Verz. 23 = *Geissorhiza ciliaris*.
radiata, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 295.—Afr. austr.
ramosa, *Eckl. Top. Verz.* 22.—Afr. austr.
rosea, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 296.—Afr. austr.
rubella, *Baker, in Journ. Bot.* xiv. (1876) 239.—Afr. austr.
setacea, Eckl. Top. Verz. 23 = *radiata*.
tenuifora, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 309 = *Petitiana*.
virginea, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 295 = *angusta*.

HESPERANTHES, S. Wats. in Proc. Am. Acad. xiv. (1879) 241 = **Antherium**, Linn. (Liliac.).
leptophylla, S. Wats. l. c. = *A. leptophyllum*.
scabrella, S. Wats. l. c. = *A. scabrellum*.
stenocarpa, S. Wats. l. c. = *A. stenocarpum*.
Torreyi, S. Wats. l. c. = *A. Torreyi*.

HESPERANTHUS, Salisb. in Trans. Hort. Soc. i. (1812) 321 = **Hesperantha**, Ker-Gawl. (Irid.).
tenuifolius, Salisb. l. c. = *Hesperantha radiata*.

HESPERELAEA, A. Gray, in Proc. Am. Acad. xi. (1876) 83. *OLEACEÆ*, Benth. & Hook. f. Gen. ii. 1040.
Palmeri, *A. Gray, l. c.*—Am. bor. occ.

HESPERETHUSA, M. Roem. Syn. Hesper. 31, 38 (1846) = **Lizmontia**, Linn. (Rutac.).
acidissima, M. Roem. l. c. 38 = *L. acidissima*.
ambigua, M. Roem. l. c. = *L. acidissima*.
crenulata, M. Roem. l. c. = *L. acidissima*.
mauritiana, M. Roem. l. c. = *L. mauritiana*.

HESPEREVAX, A. Gray, in Proc. Am. Acad. vii. (1868) 356, in obs. = **Evax**, Gaertn. (Compos.).
caulescens, A. Gray, l. c. = *E. caulescens*.

HESPERIS, Linn. Syst. ed. I (1735). *CRUCIFERÆ*, Benth. & Hook. f. i. 76.
CHALCANTHUS, Boiss. Fl. Orient. i. 156 (1867).
CLAUSIA, Trevisan, Nov. Sem. Hort. Casan. (1839).
DEILOSMA, Andrz. ex DC. Prod. iii. 450, 451, in syn. (1821); Spach, Hist. Vég. Phan. vi. 397 (1838).
KLADNIA, Schur, Enum. Pl. Transs. 58 (1866).
PLAGIOLOBA, Reichb. Nom. 183 (1841).
acris, Forsk. Fl. Aegypt. Arab. 118 = *Diplotaxis Sieberi*.
aerugines, Jord. Diagn. i. 136 = *laciniata*.
aestiva, Lam. Encyc. iii. 324 = *Matthiola incana*.
africana, Linn. Sp. Pl. 663 = *Malcolmia africana*.
alba, Mill. Gard. Dict. ed. VIII. n. 2 = *matronalis*.
albiflora, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 66 = *matronalis*.
Allionia, Lam. Fl. Fr. ii. 503 = *Sisymbrium Allionia*.
alpina, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 66.—Hungar.
alyssifolia, DC. Syst. ii. 447 = *Matthiola albicaulis*.
alyssoides, Pers. Syn. ii. 203 = *Malcolmia littorea*.
angustifolia, DC. Syst. ii. 457 = *Malcolmia torulosa*.
angustifolia, Lam. Encyc. iii. 323 = *Matthiola tristis*.
angustifolia, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 807 = *Erysimum linifolium*.
aprica, *Pair. Encyc. Suppl.* iii. 194.—Sibir.
arabidiflora, DC. Syst. ii. 454 = *Parrya Linnaeana*.
arcuata, Nocca, Fl. Sel. Hort. Ticin. i. 3. t. 2 = *Chorispora tenella*.
arenaria, Desf. Fl. Atlant. ii. 91 = *Malcolmia parvula*.
arenaria, Lag. Gen. et Sp. Nov. 20 = *Malcolmia patula*.
armena, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 68.—As. Min.
arvensis, Cav. Descr. 481 = *Moricandia arvensis*.
Aucheri, Bal. ex Fourn. in Bull. Soc. Bot. Fr. xiii. (1866) 388 = *Balansa*.
Aucheri, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 66 = *pendula*.
balansae, *Fourn. in Bull. Soc. Bot. Fr.* xiii. (1866) 388.—Anatolia.
bicuspidata, *Pair. Encyc. Suppl.* iii. 195.—As. Min.
bitumicosa, Willd. Enum. Hort. Berol. Suppl. 46 = *matronalis*.
brevicarpa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 67.—Armenia.
campicarpa, *Boiss. l. c.* 68.—Cilicia.
cappadocica, *Fourn. in Bull. Soc. Bot. Fr.* xiii. (1866) 351.—Cappadoc.
carnea, Poir. Encyc. Suppl. iii. 197 = *Heliophila platysiliqua*.
caucasica, *Rupr. Fl. Cauc.* 91.—*Rog. Caucas.*
chirinthoides, Patrin, ex DC. Syst. ii. 453 = *aprica*.
Chirinthus, Pers. Syn. ii. 203 = *aprica*.
chia, Lam. Encyc. iii. 325 = *Malcolmia chia*.
cinerea, Poir. Encyc. Suppl. iii. 196 = *Cheiranthus scoparius*.
contortuplicata, Bieb. Fl. Taur. Cauc. ii. 114 = *Malcolmia contortuplicata*.

HESPERIS:—
coronopifolia, Poir. ex Steud. Nom. ed. II. i. 754 = *Matthiola tristis*.
crenulata, DC. Syst. ii. 456 = *Malcolmia crenulata*.
crenulata, C. A. Mey. Verz. Pfl. Casc. 187 = *Malcolmia Meyeri*.
crucasca, *Adam, in Web. & Mohr, Beitr.* i. 62.—Rossia.
Cupaniana, *Guss. Fl. Sic. Syn.* ii. 186.—Sicilia.
cuspidata, DC. Syst. ii. 458 = *bicuspidata*.
dalmatica, Fourn. in Bull. Soc. Bot. Fr. xiii. (1866) 335 = *secundiflora*.
dauricula, *Ame, Fl. Iber.* vi. 495.—Hispan.
denticta, Linn. Sp. Pl. 664 = *Sisymbrium barrifolium*.
diffusa, Banks, ex DC. Syst. ii. 438 = *Brassica frutescens*.
diffusa, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 271 = *Malcolmia aegyptiaca*.
diffusa, Lam. Fl. Fr. ii. 504 = *Malcolmia africana*.
elata, Homem. Hort. Hafn. Suppl. 74 = *matronalis*.
elongata, Poir. Encyc. Suppl. iii. 197 = *Heliophila linearifolia*.
eroca, Lag. Gen. et Sp. Nov. 20 = *Malcolmia lacera*.
exgracea, Marsigli, ex Tenore, Prod. Fl. Nap. p. xxxvii = *matronalis*.
exigua, *Mill. Gard. Dict.* ed. VIII. n. 6 (Quid?).—Europ.
Farsetia, Poir. Encyc. Suppl. iii. 196 = *Farsetia aegyptia*.
fasciculata, Rafin. Specch. ii. 7 = *Matthiola fasciculata*.
fenestralis, Lam. Encyc. iii. 324 = *Matthiola fenestralis*.
flava, Georgi, Reise, i. 225 = *Erysimum altaicum*.
flava, *Kotschy & Boiss. in Boiss. Fl. Orient.* i. 287.—Kurdist.
fragrans, *Fisch. ex Sweet, Brit. Flow. Gard.* t. 61.—Sibiria.
frutescens, Poir. Encyc. Suppl. iii. 195 = *Cheiranthus semperflorens*.
glabra, *Boiss. & Noë, Diagn.* Ser. II. v. 22.—Armenia.
glabra, *Royle, Illustr. Bot. Himal.* 78.—*Reg. Himal.*
glabra, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 65 = *alpina*.
glandulosa, Pers. Syn. ii. 203 = *Dontostemon integrifolius*.
glutinosa, Vis. in Flora, xii. (1829) i. Erg. 16 = *secundiflora*.
graminea, Poir. Encyc. Suppl. iii. 197 = *Heliophila graminea*.
grandiflora, *Sims, Bot. Mag.* t. 2688.—Hab.?
heterophylla, Tenore, Fl. Nap. Prod. p. xxxix = *matronalis*.
hieracifolia, Vill. Hist. Pl. Dauph. iii. 817 = *laciniata*.
hispida, Schreb. ex Roth, in Usteri, Ann. Bot. xiv. (1795) 29 = *Malcolmia africana*.
Hoeberi, Ledeb. Fl. Ross. i. 174 = *Cheiranthus pygmaeus*.
hortensis, Pers. ex Steud. Nom. ed. I. 402 = *matronalis*.
humilis, *Boiss. Fl. Orient.* i. 231.—As. Min.
hyrcanica, Spreng. Syst. ii. 900.—Persia.
incana, Patrin, ex DC. Syst. ii. 210 = *Arabis incarnata*.
incrassata, Urv. in Mém. Soc. Linn. Par. i. (1822) 338 = *Malcolmia flexuosa*.
inodora, Aso, Syn. Pl. Arag. 87 = *Malcolmia africana*.
inodora, Gouan, Fl. Monsp. 167 = *Arabis scabrosa*.
inodora, Linn. Sp. Pl. 797 = *matronalis*.
Jana, Host, ex Flech. May. & Avé-Lall. Jad. Sem. Hort. Petrop. xi. 75 = *Malcolmia africana*.
Kladnii, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 65 = *alpina*.
Kotschyana, *Fenzl, Pugill. Pl. Nov. Syr.* 19.—Syria.
Kotschyi, Boiss. Diagn. Ser. II. v. 21 = *humilis*.
lanuginosa, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 136 = *Parrya macrocarpa*.
lasiura, *Georgi, Bescbr. Russ. Reich.* iv. 1134.—Russ. austr.
lasiura, Lam. Encyc. iii. 323 = *Malcolmia lasiura*.
lasiura, Sibth. ex DC. Syst. ii. 477 = *Matthiola sinuata*.
laxiuscula, *All. Fl. Pedem.* i. 273. t. 33. f. 1.—Europ. austr.
laxiuscula, Vis. Fl. Dalmat. 190 = *secundiflora*.

HESPERIS:—
laxa, Lam. Encyc. iii. 325 = Malcolmia africana.
leucantha, Schur, Enum. Pl. Transs. 52 = matronalis.
leucoclada, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 69.—Persia.
linearis, Poir. Encyc. Suppl. iii. 197 = Heliophila linearis.
linifolia, Pourr. ex Desf. Tabl. ed. I. 129 = Erysimum linifolium.
littorea, Lam. Encyc. iii. 323 = Malcolmia littorea.
longifolia, Poir. Encyc. Suppl. iii. 195 = Cheiranthus mutabilis.
lutea, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 383.—Japon.
lyrata, *Lam. Encyc.* iii. 325.—Sibir.
majuscula, *Fourn. in Bull. Soc. Bot. Fr.* xiii. (1866) 339.—Syria!
maritima, Lam. Fl. Fr. ii. 504 = Malcolmia maritima.
matronalis, *Linn. Sp. Pl.* 663.—Europ.; As. bor.
Menziesii, Hook. Fl. Am. Bor. i. 60 = Cheiranthus Menziesii.
microcalyx, *Fourn. in Bull. Soc. Bot. Fr.* xiii. (1866) 351.—Armenia.
minima, Torr. & Gray, Fl. N. Am. i. 90 = Cheiranthus pygmaeus.
monilliformis, Schur, in Oestr. Bot. Zeitschr. (1858) 23 = alpina.
multicaulis, *Boiss. Diagn.* Ser. I. vi. 10.—Persia.
nitens, Viv. Fl. Lib. Spec. 38. t. 5. f. 3 = Moricandia suffruticosa.
nivea, Baumg. Enum. Stirp. Transs. ii. 278 = matronalis.
oblongifolia, Schur, Enum. Pl. Transs. 52 = alpina.
obtusa, Moench, Meth. Suppl. 86 = matronalis.
odoratissima, Poir. Encyc. Suppl. iii. 196 = Matthiola odoratissima.
Oreinliana, *Tenore, Viag.* 78.—Europ. austr.
Pachypodium, Fourn. in Bull. Soc. Bot. Fr. xiii. (1866) 335 = Malcolmia africana.
pachyrhiza, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) 1. 113.—Soongaria.
Pallasii, Porter & Coulter, Fl. Colorad. 9 = Erysimum pumilum!
Pallasii, Torr. & Gray, Fl. N. Am. i. 667 = Cheiranthus pygmaeus.
paniculata, *Boiss. in Tchihat. Asie Min. Bot.* i. 331, nomen.—As. Min.
parviflora, DC. Fl. Fr. iv. 654 = Malcolmia parviflora.
parviflora, Poir. Encyc. Suppl. iii. 194 = Matthiola parviflora.
parviflora, Schur, Enum. Pl. Transs. 52 = alpina.
parvula, *Rets. in Hoffm. Phytog. Blatt.* i. (1808) 43.—Europ.
patula, Lag. ex DC. Syst. ii. 444 = Malcolmia patula.
pendula, *DC. Syst.* ii. 457.—As. Min.; Syria.
pendula, Murr. in Nov. Comm. Gotting. vi. (1776) 38. t. 6 = Arabis pendula.
persica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 64.—Persia.
petiolata, *Bieb. Fl. Taur. Cauc.* ii. 126.—Reg. Caucas.
pilosa, Poir. Encyc. Suppl. iii. 197 = Dontostemon pectinatus.
pinnata, Pers. Syn. ii. 203 = Dontostemon pectinatus.
pinnatifida, *Desf.* Choix, 63 = Malcolmia lacera.
pinnatifida, Michx. Fl. Bor. Am. ii. 31 = Thelypodium pinnatifidum.
podocarpa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii (1842) 65.—Syria.
provincialis, Linn. Sp. Pl. 664 = Matthiola tristis.
pulchella, DC. Syst. ii. 456 = Malcolmia pulchella.
pulmonarioides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 68.—Syria.
pumila, Poir. Encyc. Suppl. iii. 194 = Malcolmia parviflora.
punctulosa, Poir. l. c. 195 = Dontostemon pectinatus.
purpurascens, Jord. Diagn. i. 124 = laciniata.
pygmaea, DC. Syst. ii. 454 = Malcolmia pulchella.
pygmaea, Delile, Fl. Aegypt. Illustr. 67 = Malcolmia pygmaea.
pygmaea, Hook. Fl. Bor. Am. i. 60 = Cheiranthus pygmaeus.
quadrangula, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 67 = pulmonarioides.
remotiuscula, DC. Syst. ii. 455 = Malcolmia parviflora.
remotiuscula, *Desf. ex* Boiss. Fl. Orient. i. 233 = Malcolmia nana.

HESPERIS:—
ramosissima, *Desf. Fl. Atlant.* ii. 91. t. 161.—Afr. bor.; Sicil.
renifolia, *Boiss. & Hohen. Diagn.* Sér. I. viii. 22.—Persia.
repanda, Lag. Gen. et Sp. Nov. 20 = Erysimum linifolium.
Reuteriana, Boiss. & Huet, Diagn. Ser. II. v. 20 = bicuspidata.
rhamphicarpa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 66.—Taurus.
rigida, Sibth. ex DC. Syst. ii. 448 = Malcolmia chia.
rigida, Stev. in [Fisch.] Hort. Gorenk. ed. I. (1808) 83 = Malcolmia torulosa.
runcinata, Waldst. & Kit. Pl. Rar. Hung. ii. 220. t. 200 = matronalis.
saharitis, Boiss. & Noé, Diagn. Ser. II. i. 22 = pendula.
rupestris, Rafin. Specch. ii. 46 = Matthiola incana.
salina, Lam. Encyc. iii. 324 = Arabis incarnata.
salina, Pall. ex DC. Syst. ii. 426 = Chorispora tenella.
scabrida, *Boiss. in Ann. Sc. Nat.* Sér. III. xvii. (1842) 64.—Mesopotam.
scapigera, DC. Syst. ii. 454 = Parrya macrocarpa.
secundiflora, *Boiss. & Sprun. Diagn.* Ser. I. i. 70 — Graecia; As. Min.
semperflorens, Poir. Encyc. Suppl. iii. 196 = Cheiranthus semperflorens.
sibirica, Hohen. ex Boiss. Fl. Orient. i. 233 = matronalis.
sibirica, *Linn. Sp. Pl.* 663.—As. bor.
sinuata, Lam. Encyc. iii. 323 = Matthiola sinuata.
spectosa, Sweet, Brit. Flow. Gard. Ser. II. t. 135 = Parrya integerrima.
spectabilis, Jord. Diagn. i. 125 = laciniata.
Stevenliana, *DC. Syst.* ii. 429.—Graecia; As. Min.
suaveolens, Bess. ex Steud. Nom. ed. II. i. 754 = laciniata.
sylvestris, Crantz, Stirp. Austr. ed. I. fasc. i. 34; ed. II. 92 = matronalis.
tatarica, Pall. Reise, i. 497 = Matthiola tatarica.
taurica, Kotschy, ex Boiss. Fl. Orient. i. 232 = Kotschyana.
tenella, Hort. ex DC. Syst. ii. 436 = Chorispora tenella.
thymoides, *Boiss. Fl. Orient.* i. 234.—Armen.
trichocalycina, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) ii. 189 = aprica.
tricuspidata, Lam. Encyc. iii. 323 = Matthiola tricuspidata.
triloba, Lam. l. c. = Malcolmia littorea.
tristis, *Linn Sp. Pl.* ed. II. 927.—Europ. austr.
umbrosa, Herbich, Stirp. Rar. Bucov. 48 = matronalis.
uncinata, Steud. Nom. ed. II. i. 754, in syn., sphalm. = matronalis.
ungrolaelaria, *Boiss. Diagn.* Ser. II. v. 21.—Armenia.
unguiculata, Schrank, Pl. Rar. Hort. Monac. t. 77 = matronalis.
verna, Linn. Sp. Pl. 664 = Arabis verna.
verna, Pall. ex Ledeb. Fl. Ross. i. 172 = matronalis.
villosa, DC. Syst. ii. 449 = laciniata.
villosa, Poir. Encyc. Suppl. iii. 197 = Matthiola tricuspidata.
violacea, *Boiss. Diagn.* Ser. I. v. 80.—As. Min.
violaria, Lam. Encyc. iii. 328 = Matthiola incana.
Vitaliana, Fourn. in Bull. Soc. Bot. Fr. xiii. (1866) 338 = secundiflora.

HESPEROCALLIS, A. Gray, in Proc. Am. Acad. vi. (1866) 390. *LILIACEAE*, Benth. & Hook. f. iii. 777.
undulata, *A. Gray, l. c.* 391.—Mexic.

HESPEROCHIRON, S. Wats. Bot. King's Exped. 281. t. 80 (1871). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 829.
californicus, *S. Wats. l. c.*—Calif.
latifolius, *Kellogg, in Proc. Calif. Acad.* v. (1873) 44.—Calif.
pumilus, *Porter, in Hayden, Geol. Rep.* (1872) 768.—Am. bor. occ.

HESPEROCLES, Salisb. Gen. Pl. Fragm. 85 (1866) = **Hothoscordum**, Kunth (Liliac.).
fragrans, Salisb. l. c. = N. fragrans.

HESPEROCNIDE, Torr. in Pacif. Rail. Rep. iv. 139 (1856). *URTICACEAE*, Benth. & Hook. f. iii. 361.
sandwicensis, *Wedd. in DC. Prod.* xvi. 68.—Ins. Sandvic.
tenella, *Torr. in Pacif. Rail. Rep.* iv. 139.—Calif.

HESPEROMANNIA, A. Gray, in Proc. Am. Acad. vi. (1864-65) 554. *COMPOSITAE*, Benth. & Hook. f. ii. 459.
arborescens, *A. Gray, l. c.*—Ins. Sandvic.

HESPEROMELES, Lindl. Bot. Reg. sub t. 1956 (1837) = **Osteomeles**, Lindl. (Rosac.).
cordata, Lindl. l. c. = O. latifolia.
cuneata, Lindl. l. c.—Peruv.
ferruginea, Lindl. in Benth. Pl. Hartw. 129 = O. ferruginea.
glabrata, M. Roem. Syn. Rosif. 225 = O. glabrata.
heterophylla, Hook. fc. Fl. t. 846.—N. Granat.
latifolia, M. Roem. Syn Rosif. 225 = O. latifolia.
oblonga, Lindl. Bot. Reg. sub t. 1956.—N. Granat.
obtusifolia, Lindl. l. c. = O. glabrata.
pernettyoides, Wedd. Chlor. And. ii. 230 = O. pernettyoides.

HESPEROSCORDIUM, Baker, in Journ. Linn. Soc. xi. (1871) 378 = seq.

HESPEROSCORDON, Hook. Fl. Bor. Am. ii. 185 (1839) = seq.

HESPEROSCORDUM, Lindl. Bot. Reg. sub t. 1298 (1829) = **Brodiaea**, Sm. (Liliac.).
hyacinthinum, Lindl. l. c. t. 1639 = B. lactea.
lacteum, Lindl. l. c. t. 1639 = B. lactea.
Lemisii, Hook. Fl. Bor. Am. ii. 186 = B. lactea.
maritimum, Torr. in Pacif. Rail. Rep. iv. 148 = Muilla maritima.

HESPEROXIPHION, Baker, in Journ. Linn. Soc. xvi. (1877) 127 = **Cypella**, Herb. (Irid.).
peruvianum, Baker, l. c. = C. peruviana.
pusillum, Baker, l. c. = C. pusilla.

HESSEA, Berg. ex Schlecht. in Linnaea, i. (1826) 252 = Carpolyza, Salisb. (Amaryll.).
spiralis, Berg. ex Schlecht. l. c. = C. spiralis.

HESSEA, Herb. Amaryll. 289. t. 29 (1837). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 730.
GEMMARIA, Salisb. Gen. Pl. Fragm. 127 (1866).
IMHOFIA, Herb. App. 18 (1821).
PEIRITHANES, Salisb. Gen. Pl. Fragm. 118 (1866).
brevifiora, *Herb. Amaryll.* 289. t. 29. f. 6.—Afr. austr.
Burchelliana, *Benth. & Hook. f. Gen.* iii. 731.—Afr. austr.
crispa, *Kunth, Enum. Pl.* v. 632.—Afr. austr.
Dregeana, *Kunth, l. c.* 633.—Afr. austr.
filifolia, *Benth. & Hook. f. Gen.* iii. 731.—Afr. austr.
gemmata, *Benth. & Hook. f. l. c.*—Afr. austr.
stellaria, *Herb. Amaryll.* 289.—Afr. austr.
vaginata, Herb. l. c. = Strumaria truncata.

HETAERIA, Blume, Bijdr. 409. t. 14 (1825) (Etaeria); Fl. Jav. Praef. p. vii (1828). *ORCHIDEAE*, Benth. & Hook. f. iii. 493.
AETHERIA, Endl. Gen. 214 (18 7).
CREOCHILUS, Lindl. in Gard. Chron. (1854) 67.
KHAMPHIDIA, Lindl. in Journ. Linn. Soc. i. (1857) 181.
RHOMBODIA, Lindl. l. c. (1857) 181.
SALACISTIS, Reichb. f. Xenia Orch. i. 214. t. 87 (1857).
abbreviata, *Lindl. in Wall. Cat.* n. 7385; *Gen. et Sp. Orch.* 481.—Reg. Himal.
aginis, Lindl. in Wall. Cat. n. 7382 (Zeuxines sp.).—Ind. or.
albida, Blume, Bijdr. i. 410 = Vrydagyneia albida.
anomala, *Lindl. in Journ. Linn. Soc.* i. (1857) 185.—Reg. Himal.
cassythoea, *Lindl. in Hook. Lond. Journ. Bot.* i. (1842) 189.—Bras.
coelata, *Lindl. in Journ. Linn. Soc.* i. (1857) 185.—Ins. Banda.
cristata, *Blume, Orch. Archip. Ind.* 103.—Java.
elongata, *Lindl. in Wall. Cat.* n. 7384.—Malaya.

HETAERIA :—
elongata, Miq. Fl. Ind. Bat. iii. 726 = Vrydagzynia elongata.
flava, Lindl. in Wall. Cat. n. 7380 = Zeuxine flava.
forcipata, *Reichb. f. in Linnaea*, xli. (1877) 62.—Ins. Fiji.
fusca, *Lindl. Gen. et Sp. Orch.* 491.—Reg. Himal. Gardneri, *Benth. & Hook. f. Gen.* iii. 604.—Zeylan.
gracilis, Miq. Fl. Ind. Bat. iii. 726 = Vrydagzynia gracilis.
javanica, Blume, Bijdr. 410 = Cystorchis javanica.
lamellata, *Blume, Orch. Archip. Ind.* 106.—Java.
lanceolata, *Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 142.—Reg. Himal.
longifolia, *Benth. in Journ. Linn. Soc.* xviii. (1880) 546.—Reg. Himal.
micrantha, *Blume, Orch. Archip. Ind.* 103.—Java.
mollis, Lindl. in Journ. Linn. Soc. i. (1857) 184 (Zeuxines sp.).—Ind. or.
moulmeinensis, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 149.—Burma.
nervosa, Lindl. in Wall. Cat. n. 7381 = Zeuxine nervosa.
nuda, Miq. Fl. Ind. Bat. iii. 726 = Vrydagzynia nuda.
obliqua, *Blume, Orch. Archip. Ind.* 104.—Borneo.
oblongifolia, *Blume, Bijdr.* 410. t. 14; *Orch. Archip. Ind.* 102.—Java.
obscura, Miq. Fl. Ind. Bat. iii. 726 = Cystorchis obscura.
occulta, Lindl. Bot. Reg. (1838) Misc. 94 = Platylepis goodyeroides.
ovalifolia, *Benth. & Hook. f. Gen.* iii. 604.—Ind. or.
polyphylla, *Reichb. f. Otia Bot. Hamb.* 52.—Ins. Fiji.
purpurascens, *Blume, Orch. Archip. Ind.* 105.—Java.
purpurea, Miq. Fl. Ind. Bat. iii. 726 = Vrydagzynia purpurea.
roxaena, *Reichb. f. in Bonplandia*, v. (1857) 35.—Java.
rubens, *Benth. & Hook. f. Gen.* iii. 604.—Reg. Himal.; Burma.
rubicunda, Reichb. f. in Bonplandia, iii. (1855) 214 = Goodyera rubicunda.
tennis, *F. Muell. Census*, 111; *Vidal, Phan. Cuming. Philipp.* 81, 151.—Ins. Philipp.; Austral.
uncinata, Miq. Fl. Ind. Bat. iii. 727 = Vrydagzynia uncinata.
vaginalis, *Reichb. f. in Flora*, lxvii. (1885) 537.—Ins. Comor.
variegata, Miq. Fl. Ind. Bat. iii. 726 = Cystorchis variegata.
Whitmeei, *Reichb. f. in Journ. Bot.* xv. (1877) 133.—Ins. Samoa.

HETAERIA, Endl. Gen. 133 (1837) = **Pritzelia**, F. Muell. (Philyd.).
pygmaea, Endl. l. c. = P. pygmaea.

HETERACANTHA, Link, ex Steud. Nom. ed. II. i. 755 (1840) = **Carthamus**, Tourn. (Compos.).
lanata, Link, ex Steud. l. c. = C. lanatus.

HETERACEA, Steud. Nom. ed. II. i. 755 (1840) = **Heteraelia**, Fisch. & Mey. (Compos.).

HETERACHAENA, Fresen. in Mus. Senckenb. iii. (1839) 74. *COMPOSITAE*, Benth. & Hook. f. ii. 526.
massaensis, *Fresen. l. c.*—Abyssin.

HETERACHAENA, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Ind. ii. (1845) 577 = **Pimpinella**, Riv. (Umbellif.).
alpina, Zoll. & Mor. l. c. = P. javana.
javanica, Walp. Ann. i. 846, in syn. = P. javana.

HETERACHENE, Benth. in Hook. Ic. Pl. xiii. 40 t. 1250 (1877); Fl. Australas. vii. 634 (1878). *GRAMINEAE*, Benth. & Hook. f. iii. 1186.
Brownii, *Benth. ll. cc.* 40; 635.—Austral.
Gulliveri, *Benth. in Hook. Ic. Pl.* t. 1250.—Austral.

HETERACHTHIA, Kunze, in Bot. Zeit. viii. (1850) 1 = **Trachocentia**, Linn. (Cotemella.).
Gaudichaudiana, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 808 = T. elongata.
paichelia, Hassk. ex C. B. Clarke, l. c. 804 = T. amplexicaulis.
pulchella, Kunze, in Bot. Zeit. viii. (1850) 1 = T. pulchella.

HETERACIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. (1835) 28. *COMPOSITAE*, Benth. & Hook. f. ii. 510.
Steetziai, *Fisch. & Mey. l. c.* 30.—Armenia.

HETERACTIS, DC. Prod. vi. 468 (1837) = **Gymnosteblium**, Less. (Compos.).
falcata, DC. l. c. = G. ciliare.

HETERANDRA, Beauv. in Trans. Am. Phil. Soc. iv. (1799) 175 = **Heteranthera**, Ruiz & Pav. (Pontederiac.).
reniformis, Beauv. l. c. 177 = Heteranthera reniformis.

HETERANTHELIUM, Hochst. ex Jaub. & Spach, Illustr. iv. 24 (1850) = **Agropyron**, J. Gaertn. (Gramin.).
piliferum, Hochst. ex Jaub. & Spach, l. c. = A. orientale.

HETERANTHEMIS, Schott, in Oken, Isis (1818), 821 = **Chrysanthemum**, Linn. (Compos.).
hirta, Spreng. Nov. Prov. 9 = C. viscosum.
viscidi-hirta, Schott, in Oken, Isis (1818), 821. f. 5 = C. viscosum.

HETERANTHERA, Ruiz & Pav. Prod. 9. t. 2 (1794). *PONTEDERIACEAE*, Benth. & Hook. f. iii. 888.
Heterandra, Beauv. in Trans. Am. Phil. Soc. iv. (1799) 175.
Leptanthus, Michx. Fl. Bor. Am. i. 24. t. 5 (1803).
Lunania, Rafin. Med. Fl. ii. 106 (1830).
Schollera, Schreb. Gen. 726 (1791).
Thiekastina, Rafin. Fl. Tellur. iv. 121 (1836).
acuta, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 438; *Vahl, Enum.* ii. 42 = reniformis.
alismoides, H. B. & K. ex Link, Jahrb. i. iii. 78 = limosa.
allamoides, *Willd. ex Kunth, Enum. Pl.* iv. 122.—Venezuela.
callaefolia, *Reichb. ex Kunth, l. c.* 123.—Afr. trop.
cordata, *Vahl, Enum.* ii. 44.—Am. austr.
diversifolia, *Vahl, l. c.*—Guiana.
formosa, Mig. in Linnaea, xvii. (1843) 60 = Eichhornia speciosa.
graminea, *Vahl, Enum.* ii. 45.—Am. bor.
grandiflora, Klotzsch, in Schomb. Fauna et Fl. Brit. Gui. 1118 = Eichhornia pauciflora.
hydrocleiifolia, *Griseb. Cat. Pl. Cub.* 253.—Cuba.
Kotschyana, Fenzl, ex Schweinf. Beitr. Fl. Aethiop. 294, nomen = callaefolia.
limosa, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 439; *Vahl, Enum.* ii. 44.—Am. calid.
mexicana, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 168.—Mexic.
oblongifolia, *Mart. in Schult. f. Syst.* vii. 1148.—Bras.
paucalaris, *Benth. Pl. Hartw.* 25.—Mexic.
Potamogeton, *Soims, in DC. Monog. Phan.* iv. 521.—Afr. trop.
pulchella, Vahl, Enum. ii. 43 = reniformis.
reniformis, *Ruiz & Pav. Fl. Per.* i. 43. t. 71.—Am. calid.
retusifolia, Griseb. Cat. Pl. Cub. 253 = limosa.
Seubertiana, *Solms, in DC. Monog. Phan.* iv. 518.—Bras.
spicata, *Presl, Symb. Bot.* t. 16.—Cuba.
sosterasefolia, *Mart. Nov. Gen. et Sp.* i. 7. t. 8.—Bras.

HETERANTHIA, Nees & Mart. in Nov. Act. Nat. Cur. xi. i. (1823) 61. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 998.
Vrolikia, Spreng. Syst. iii. 149 (1826).
decipiens, *Nees & Mart. l. c.*—Bras.

HETERANTHUS, Bonpl. ex Cass. in Dict. Sc. Nat. xxi. 110 (1821) = **Perezia**, Lag. (Compos.).

HETERANTHUS, Dum. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 183 = **Wentenaria**, Kuhl. (Gramin.).
tenuis, Fourr. l. c. = V. avenacea.

HETERAPITHMOS, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 365. Gen. dub. *OLACIN.*—Cf. Benth. & Hook. f. Gen. i. 995.
floribunda, *Turcz. l. c.*—Bras.

HETERELYTRON, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 294 = **Anthisteiria**, Linn. (Gramin.).
scaberrima, Jungh. l. c. 295 = A. gigantea.

HETERISIA, Rafin. Fl. Tellur. ii. 69 (1836) = **Saxifraga**, Tourn.

HETEROCANSCORA, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 481 = **Canscora**, Lam. (Gentian.).
Schulteaii, C. B. Clarke, l. c. = C. Schultesii.

HETEROCARPAEA, Schoele, in Linnaea, xxi. (1848) 467 = **Galactia**, P. Br. (Legumin.).
texana, Scheele, l. c. 469 = G. canescens.

HETEROCARPUS, Phil. in Bot. Zeit. xiv. (1856) 641 = **Cardamine**, Tourn. (Crucif.).
fernandezianus, Phil. l. c. = C. chenopodiifolia.

HETEROCARPUS, Wight, Ic. vi. 29. t. 2067 (1853) = **Commelina**, Linn.
glaber, Wight, l. c. t. 2067 = C. glabra.
hirsutus, Wight, l. c. = C. hirsuta.
obliquus, Hassk. Commel. Ind. 7 = C. obliqua.
oligotrichus, Hassk. l. c. 7 = C. oligotricha.

HETEROCARYUM, A. DC. in DC. Prod. x. 144 (1846) = **Bohinospermum**, Sw. (Borag.).
candylopherum, A. DC. l. c. = K. candylopherum.
dinaricatum, Stocks, ex Boiss. Fl. Orient. iv. 258 = E. semilliforum.
laevigatum, A. DC. in DC. Prod. x. 145 = E. laevigatum.
minimum, A. DC. l. c. 144 = E. minimum.
pachypodum, A. DC. l. c. = E. Szovitsianum.
rigidum, A. DC. l. c. 145 = E. Szovitsianum.
Szovitsianum, A. DC. l. c. = E. Szovitsianum.

HETEROCENTRON, Hook. & Arn. Bot. Beech. Voy. 290 (1840) = **Heeria**, Schlecht. (Melastom.).
alpestre, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 156 = Heeria macrostachya.
Arsteadi, O. Berg, ex Cogn. in Mart. Fl. Bras. xiv. iii. 297, sphalm. = Tibouchina Oerstedii.
axillare, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 155 = Heeria macrostachya.
dicranantherum, Naud. l. c. 156 = Monochaetum dichranantherum.
glandulosum, Schrenk, in Regel, Gartenfl. (1856) 227. t. 169 = Heeria macrostachya.
Humboldtianum, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 156 = Monochaetum Humboldtianum.
macrostachyum, Naud. l. c. 155 = Heeria macrostachya.
muricatum, Hook. & Arn. Bot. Beech. Voy. 290 = Heeria subtriplinervia.
muricatum, Naud. in Bot. Mag. t. 5166 = Heeria rosea.
Oerstedii, O. Berg, ex Triana, in Trans. Linn. Soc. xxvii. (1871) 40 = Tibouchina Oerstedii.
rosea, A. Br. & Bouché, in Linnaea, xxv. (1852) 299 = Heeria rosea.
subtriplinervium, A. Br. & Bouché, l. c. = Heeria subtriplinervia.
undulatum, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 155 = Heeria undulata.

HETEROCENTRUM, Hemsl. Biol. Centr. Am. Bot. i. 418 (1880) = pmec.

HETEROCHAENIA, A. DC. in DC. Prod. vii. 441 (1839). *CAMPANULACEAE*, Benth. & Hook. f. ii. 554.
cmifolia, *A. DC. l. c.* 442.—Ins. Macaar.

HETEROCHAETA, DC. Prod. v. 323 (1836) = **Aster**, Tourn. &c. (Compos.).
asteroides, DC. l. c. = A. Heterochaeta.
diplostephioides, DC. l. c. = A. diplostephioides.
erigeroides, DC. l. c. = Erigeron alpinus.
gnaphalioides, DC. l. c. = Erigeron gnaphalioides.
hispida, DC. l. c. = Hysterionica jasionoides.

HETEROCHAETA :—
leucophylla, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 229 = Erigeron leucophyllus.
Pseudo-Erigeron, Bunge, l. c. 238.—Siberia.
pubescens, DC. Prod. v. 288 = Erigeron pubescens.
stricta, Benth. Pl. Hartw. 306 = Erigeron strictus †

HETEROCHLAMYS, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 61 = **Julocroton**, Mart. (Euphorb.).
quinquenervia, Turcz. l. c. 62 = J. argenteus.

HETEROCHLOA, Desv. Opuse. 66 (1831) = **Andropogon**, Linn. (Gramin.).
Alopecurus, Desv. l. c. = A. peduncularis, *Baker*.

HETEROCHLOA, Endl. Gen. Suppl. iii. 92 (1843) = seq.

HETEROCHROA, Bunge, in Ledeb. Fl. Alt. ii. 131 (1830) = **Gypsophila**, Linn. (Caryoph.).
desertorum, Bunge, Vers. Suppl. Fl. Alt. 29 = G. desertorum.
glandulosa, Boiss. Diagn. Ser. I. i. 15 = G. glandulosa.
microphylla, Schrenk, Enum. Pl. Nov. i. 92 = G. microphylla.
minuartioides, Jaub. & Spach, Illustr. i. 26 = G. minuartioides.
petraea, Bunge, in Ledeb. Fl. Alt. ii. 131 = G. Bungeana.
sperguloefolia, Jaub. & Spach, Illustr. i. 28 = G. sperguloefolia.
violacea, Walp. Rep. v. 81 = G. violacea.

HETEROCLADUS, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 152 = **Coriaria**, Linn.
caracasanus, Turcz. l. c. 158 = C. thymifolia.

HETEROCLITA, Rafin. Fl. Tellur. iii. 27 (1836) = **Gentiana**, Tourn.
dichotoma, Rafin. l. c. = G. tenella.

HETEROCODON, Nutt. in Trans. Am. Phil. Soc. N. S. II. viii. (1842) 255 = **Campanula**, Tourn.
minimum, *Kellogg*, in Proc. Calif. Acad. vii. (1877) 111 = Alchemilla arvensis.
rariflorum, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1842) 255 = Calif.

HETEROCOMA, DC. in Ann. Mus. Par. xvi. (1810) 191. t. 7. *COMPOSITAE*, Benth. & Hook. f. ii. 224.
albida, *DC. l. c.*—Bras.
bifrons, DC. l. c. = Chronopappus bifrons.
pedunculata, Steud. Nom. ed. I. 408 = Vernonia pedunculata.

HETEROCRAMBE, Coss. & Dur. Ind. Sem. Hort. Burdig. (1867) 25 = **Brassica**, Tourn. (Crucif.).
Aristida, Coss. & Dur. l. c. (= *Sinapis pubescens*).—Hab.?

HETERODENDRON, Spreng. Syst. ii. 356 (1825) = seq.

HETERODENDRUM, Desf. in Mém. Mus. Par. iv. (1818) 8. t. 3. *SAPINDACEAE*, Benth. & Hook. f. i. 407.
diversifolium, *F. Muell. Fragm.* i. 46.—Austral.
macrocalyx, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 119.—Austral.
microcalyx, *Radlk. l. c.*—Austral.
oleaefolium, *Desf. in Mém. Mus. Par.* iv. (1818) 8. t. 3. —Austral.

HETERODERIS, Boiss. Fl. Orient. iii. 798 (1875) = **Crepis**, Vaill. (Compos.).
pusilla, Boiss. l. c. 794.—Persia.
Stocksiana, Boiss. l. c.—Beluchist.

HETERODON, Meisn. Gen. 72 (1837) = **Bernelia**, Brongn. (Bruniac.).
fragarioides, Meisn. Gen. Comm. 52 (= *Berardia fragarioides*).—Afr. austr.
superbus, Meisn. l. c. = Bernelia superba.

HETERODONTA, Nutt. ex Benth. & Hook. f. Gen. ii. 385 (1876) = **Coreopsis**, Linn. (Compos.).

HETERODRABA, Greene, in Bull. Calif. Acad. i. 2. iii. (1885) 71. *CRUCIFERAE*.
unilateralis, *Greene, l. c.* 72.—Calif.

HETEROGAURA, Rothr. in Proc. Am. Acad. vi. (1864) 554. *ONAGRARIEAE*, Benth. & Hook. f. i. 793.
californica, *Rothr. l. c.*—Calif.

HETEROLAENA, C. A. Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. x. 67 (1845) = **Pimelea**, Banks (Thymel.).
decussata, C. A. Mey. in Bull. Phys. Math. Acad. Pétersb. iv. (1846) 74 = *P. tetragona*.
Hendersonii, C. A. Mey. l. c. 74 = P. rosea.
hispida, C. A. Mey. l. c. 73 = P. hispida.
incana, C. A. Mey. l. c. = P. nivea.
nivea, C. A. Mey. l. c. = P. nivea.
rosea, C. A. Mey. l. c. = P. rosea.
spectabilis, C. A. Mey. ex Fisch. Mey. & Avé-Lall. in Ind. Sem. Hort. Petrop. x. 48 = P. spectabilis.

HETEROLAENA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 948 (1878) = **Eupatorium**, Tourn. (Compos.).
alpestris, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 328 = E. serratum.
anethifolia, Sch. Bip. ex Baker, l. c. 330 = E. anethifolium.
conferta, Sch. Bip. ex Baker, l. c. 328 = E. serratum.
Gaudichaudiana, Sch. Bip. ex Baker, l. c. 329 = E. Gaudichaudianum.
intermedia, Sch. Bip. ex Baker, l. c. = E. intermedium.
involucrata, Sch. Bip. ex Baker, l. c. 331 = E. coriaceum.
stepingensis, Sch. Bip. ex Baker, l. c. 329 = E. intermedium.
leucodon, Sch. Bip. ex Baker, l. c. 330 = E. Gaudichaudianum.
mollissima, Sch. Bip. ex Baker, l. c. 331 = E. mollissimum.
nitida, Sch. Bip. ex Baker, l. c. 370 = Kaunia erythralina.
organis, Sch. Bip. ex Baker, l. c. 328 = E. serratum.
serrata, Sch. Bip. ex Baker, l. c. = E. serratum.

HETEROLATHUS, Presl, Bot. Bemerk. 131 (1844); in Abh. Boehm. Ges. Wiss. V. iii. (1845) 561 = **Aspalathus**, Linn. (Legumin.).
involucrata, Presl, Bot. Bemerk. 132 = A. undulata.
polycephala, Presl, l. c. = A. polycephala.
suaveolens, Presl, l. c. = A. suaveolens.

HETEROLEPIS, Cass. in Bull. Soc. Philom. (1820) 26. *COMPOSITAE*, Benth. & Hook. f. ii. 328.
HETEROMORPHA, Cass. in Bull. Soc. Philom. (1817) 19.
conyzoides, Bert. ex DC. Prod. vi. 416 = Senecio adenotrichius.
decipiens, *Cass. in Dict. Sc. Nat.* xxi. 120.—Afr. austr.
mitis, *DC. Prod.* vi. 497.—Afr. austr.
oreades, Steud. Nom. ed. II. i. 755, sphalm. = *conyzoides*.
peduncularis, *DC. Prod.* vi. 497.—Afr. austr.

HETEROLEPIS, Ehrenb. ex Boiss. Fl. Orient. v. 554 (1881) = **Chloris**, Sw. (Gramin.).
elegans, Ehrenb. l. c. = C. meccana.

HETEROLOMA, Desv. ex Reichb. Consp. 152 (1828) = **Adamia**, DC. (Legumin.).
ovalopycnoides, Desv. ex DC. Prod. ii. 319 = A. papposa.

HETEROLOPHUS, *Cass.* in Dict. Sc. Nat. i. 250 (1827) = **Centaurea**, Linn. (Compos.).
sibiricus, Cass. l. c. = C. Marschalliana.

HETEROMELES, M. Roem. Syn. Rosifl. 105 (1847) = **Photinia**, Lindl. (Rosac.).
arbutifolia, M. Roem. l. c. = P. arbutifolia.
Fremontiana, Decne. in Nouv. Ann. Mus. Par. Sér. I. x. (1874) 144 = P. arbutifolia.

HETEROMERA, Pomel, Nouv. Mat. Fl. Atl. 60 (1874) = **Chrysanthemum**, Tourn. (Compos.).
fuscata, Pomel, l. c. = Matricaria fuscata.
macrocarpa, Pomel, l. c.—Algar.

HETEROMERIS, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 290 = **Helianthemum**, Tourn. (Cistin.).
canadensis, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 370 = Helianth. canadense.
cymosa, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 291 = Helianth. canadense.
mexicana, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 370 = Helianthemum glomeratum.
Michauxii, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 291 = Helianth. canadense.
polyfolia, Spach, l. c. 291 = Helianth. capitatum.

HETEROMMA, Benth. in Benth. & Hook. f. Gen. ii. 386 (1873). *COMPOSITAE*.
decurrens, *Benth. (in Herb. Kew.).*—Afr. austr.

HETEROMORPHA, Cass. in Bull. Soc. Ph. lom. (1817) 12 = **Heterolepis**, Cass. (Compos.).
decipiens, Cass. l. c. = Heterolepis decipiens.

HETEROMORPHA, Cham. & Schlecht. in Linnæa, i. (1826) 256. t. 5. *UMBELLIFERAE*, Benth. & Hook. f. i. 887.
abyssinica, Hochst. in Flora, xxiv. (1841) 1. Intell. 28 = arborescens.
arborescens, *Cham. & Schlecht. in Linnæa*, i. (1826) 255.—Afr. austr.
collina, Eckl. & Zeyh. Enum. 342 = arborescens.
trifoliata, Eckl. & Zeyh. l. c. = arborescens.

HETEROMORPHA, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 278 = **Hypochoeris**, Linn. (Compos.).
crinita, Viv. ex Coss. l. c. = Hypoch. aethensis.

HETEROMYRTUS, Blume, Mus. Bot. Lugd. Bat. i. 76 (1849), in obs. = **Myrtus**, Tourn.
umbilicata, Blume, l. c. = M. umbilicata.

HETERONEMA, Reichb. Consp. 173 (1828) = HETERONOMA, DC. = **Arthrostema**, Ruiz & Pav. (Melastom.).

HETERONEURON, Hook. f. in Benth. & Hook. f. Gen. i. 768 (1867) = **Eoreya**, DC. (Melastom.).
nigricans, Hook. f. l. c. = L. nigricans.

HETERONOMA, DC. Prod. iii. 122 (1828) = **Arthrostema**, Ruiz & Pav. (Melastom.).
campanularе, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 158 = A. campanulare.
ciliatum, Naud. l. c. 154 = A. ciliatum.
coriaceum, Mart. Nov. Gen. et Sp. iii. 140.—Bras.
cubense, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 152 = A. fragile.
diversifolium, Benth. Bot. Voy. Sulph. 98 = A. campanulare.
diversifolium, DC. Prod. iii. 123 = A. ciliatum.
diversifolium, Link & Otto, Ic. Pl. Rar. 73. t. 37 = A. fragile.
Galeottianum, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 153 = A. fragile.
Gaudichaudianum, Naud. l. c. 152 = A. volubile.
latifolium, Naud. l. c. 153 = A. latifolium.
mexicanum, Benth. Bot. Voy. Sulph. 94 = Heeria subtriplinervia.
Pachyloma, Mart. Nov. Gen. et Sp. iii. 140. t. 278 = Pachyloma coriaceum.
subtriplinervium, DC. 3es Notice Pl. Genèv. 12 = Heeria subtriplinervia.
volubile, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 153 = A. volubile.

HETEROPANAX, Seem. Journ. Bot. iv. 297 (1866); Fl. Vit. 114. *ARALIACEAE*, Benth. & Hook. f. i. 946.
fragrans, *Seem. Fl. Vit.* 114.—Ins. Fiji.

HETEROPAPPUS, Less. Syn. Comp. 189 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 269.
decipiens, Maxim. Prim. Fl. Amur. 146 = hispidus
hispidus, *Less. Syn. Comp.* 189.—Japon.
incisus, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 182 = hispidus.
ringens, Sieb. & Zucc. l. c. 184 = hispidus.
Sampsoni, *Hance, in Journ. Bot.* v. (1867) 370.—China.
subserratus, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 183 = hispidus.

HETEROPETALUM, Benth. in Journ. Linn. Soc. v. (1861) 69. *ANONACEAE*, Benth. & Hook. f. i. 37.
brasiliense, *Benth. l. c.*—Bras.

HETEROPHRAGMA, DC. Prod. ix. 210 (1845). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1046.
FERDINANDIA, Welw. ex Seem. Journ. Bot. iii. (1865) 280. t. 28.
adenophyllum, *Seem. ex Benth. & Hook. f. Gen.* ii. 1047.—Ind. or.
chelonioides, Dals. & Gibs. Bomb. Fl. 160 = Stereospermum chelonioides.
Roxburghii, *DC. Prod.* ix. 210.—Ind. or.
suaveolens, Dals. & Gibs. Bomb. Fl. 161 = Stereospermum suaveolens.
sulfureum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 90.—Burma.

HETEROPHYLLAEA, Hook. f. Ic. Pl. t. 1134 (1876). *RUBIACEAE*, Benth. & Hook. f. ii. 37.
lancolata, *Griseb. in Goett. Abh.* xxiv. (1879) 154.—Reg. Argent.
pentulata, *Hook. f. Ic. Pl.* t. 1134.—Bolivia.

HETEROPHYLLEIA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) t. 591 = **Coriaria**, Linn.
coraczana, Turcz. l. c.—Cf. eund. l. c. xx. (1847) t. 158 = C. thymifolia.

HETEROPHYLLUM, Boj. ex Hook. Bot. Misc. i. (1830) 291 = **Buettneria**, Loefl.
ramosum, Boj. ex Hook. l. c. = B. heterophylla.

HETEROPLEURA, Sch. Bip. in Flora, xlv. (1862) 484 = **Hieracium**, Linn. (Compos.).
ambigua, Sch. Bip. l. c. 485 = H. Pendleri.
crepidisperma, Sch. Bip. l. c. = H. Crepidispermum.

HETEROPOGON, Pers. Syn. ii. 533 (1807). *GRAMINEAE*, Benth. & Hook. f. iii. 1133.
acuminatus, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 234.—Reg. trop.
Allionii, Roem. & Schult. Syst. ii. 835 = glaber.
arrhenobasis, *Anderss. in Schweinf. Beitr. Fl. Aethiop.* 310.—Abyss.
barbinensis, Miq. Fl. Ind. Bat. iii. 494 = hirtus.
concinnus, Thw. Enum. Fl. Zeyl. 368 = Andropogon caricosus.
contortus, Beauv. ex Roem. & Schult. Syst. ii. 836 = hirtus.
filifolius, Ness, Fl. Afr. Austr. 102 = Andropogon filifolius.
finnus, *V. & C. Presl, Rel. Haenk.* i. 334.—Reg. trop. et subtrop.
glaber, *Pers. Syn.* ii. 533.—Reg. temp. et trop.
hirsutus, Beauv. Agrost. 164 = hirtus.
hirtus, Anderss. in Schweinf. Beitr. Fl. Aethiop. 310 = pubescens.
hirtus, *Pers. Syn.* ii. 533.—Reg. trop.
hispidissimus, Hochst. ex Steud. Syn. Pl. Gram. 367 = hirtus.
Hochstetteri, Anderss. in Schweinf. Beitr. Fl. Aethiop. 310 = Andropogon fastigiatus.
Hohenackeri, Hochst. ex Miq. Anal. Bot. Ind. ii. 24 = hirtus.
insignis, *Thw. Enum. Pl. Zeyl.* 427.—As. trop.
inclyranthus, Miq. Fl. Ind. Bat. iii. 493 = insignis.
lianantherus, Miq. l. c. = insignis.
megapotamicus, Schult. Mant. ii. 668 = Trachypogon polymorphus.
melanocarpon, *Duthie, in Athins. Gas. N. W. Ind.* 639, nomen.—Ind. or.
melanocarpus, Benth. in Journ. Linn. Soc. xix. (1881) 71 = acuminatus.
messanensis, Guss. Fl. Sic. Syn. i. 164 = glaber.
pilosus, *Spreng. Nov. Prov.* 22.—Ind. or.
polystachyus, Ness, Agrost. Bras. 264 = hirtus.
polystachyus, Schult. Mant. ii. 460 = finnus!
polystictus, Hochst. in Flora, xxxix. (1856) 28 = acuminatus.
pubescens, *Anderss. in Schweinf. Beitr. Fl. Aethiop.* 310.—Reg. Mediterr.
Roxburghii, Arn. ex Ness, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 185 = hirtus.
Royleri, Ness, ex Steud. Syn. Fl. Gram. 367 = acuminatus.

HETEROPOGON:—
scrobiculatus, Fourn. ex Hack. in Mart. Fl. Bras. ii. iii. 268 = acuminatus.
secundus, J. & C. Presl, Rel. Haenk. i. 335 = Trachypogon polymorphus.
stipoides, J. & C. Presl, l. c. = Trachypogon polymorphus.
tenellus, Schult. Mant. ii. 460 = Andropogon tenellus.
tenuiculus, Miq. Fl. Ind. Bat. iii. 494 = Andropogon sericeus.
truncatus, Ness, Fl. Afr. Austr. 102 = Trachypogon polymorphus.
villosus, Ness, Agrost. Bras. 262 = Andropogon Nees.

HETEROPSIS, Kunth, Enum. Pl. iii. 59 (1841). *AROIDEAE*, Benth. & Hook. f. iii. 991.
oblíqua, Miq. ex Engl. in Mart. Fl. Bras. iii. ii. 112 = Monstera obliqua.
oblongifolia, *Kunth, Enum. Pl.* iii. 60.—Bras.
ovata, Miq. Del. Sem. Hort. Amstel. (1853); ex Walp. Ann. v. 895 = Monstera pertusa.
Riedeliana, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 90.—Bras.
salicifolia, *Kunth, Enum. Pl.* iii. 60.—Bras.
Spruceana, *Schott, Aroid.* i. 27. t. 60.—Bras.
surinamensis, Miq. Del. Sem. Hort. Amstel. (1853); ex Walp. Ann. v. 895 = Monstera pertusa.

HETEROPTERA, Stend. Nom. ed. II. i. 756 (1840), sphalm. = HEPTAPTERA, Marg. & Reut. = **Prangos**, Lindl. (Umbellif.).

HETEROPTERIS, H. B. & K. Nov. Gen. et Sp. v. 168 (1821). *MALPIGHIACEAE*, Benth. & Hook. f. i. 356.
acreoides, *Griseb. in Linnaea*, xiii. (1839) 228.—Bras.
acutifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 459. —Am. austr.
acuta, *Griseb. in Linnaea*, xiii. (1839) 257.—Bras.
affinis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 465.—Am. austr.
africana, *A. Juss. l. c.* 456.—Afr. austr.
albida, *Blume, Bijdr.* 226.—Java.
angustifolia, *Griseb. in Linnaea*, xiii. (1839) 223.—Bras.
anomala, *A. Juss. in Arch. Mus. Par.* iii. (1843) 485. —Bras.
anoptera, *A. Juss. l. c.* 459.—Bras.
apiculata, *Miq. Stirp. Surin. Sel.* 31.—Guian.
appendiculata, *DC. Prod.* i. 592.—Ins. S. Vincent.
argentea, H. B. & K. Nov. Gen. et Sp. v. 164 = Banisteria argentea.
argyrophaea, *A. Juss. in Arch. Mus. Par.* iii. (1843) 447.—Bras.
aureo-nitens, *Planch. & Linden, in Linden, Cat.* n. 8. (1853) 8.—N. Granat.
australis, A. Juss. ex Walp. Rep. v. 278 = syringaefolia.
bashkeiloides, *Griseb. in Linnaea*, xiii. (1839) 219.—Bras.
Beecheyana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 475.—Mexic.
Bertteroana, *A. Juss. l. c.* 476.—Ind. occ..
bicolor, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 28.—Bras.
biglandulosa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 464.—Guiana.
brachiata, *H. B. & K. Nov. Gen. et Sp.* v. 163, in obs. —Ind. austr.
brasiliensis, *Regel & Kcern. Ind. Sem. Hort. Petrop.* (1858) 47.—Bras.
byrsonimaefolia, A. Juss. in Arch. Mus. Par. iii. (1843) 464 = xanthophylla.
cacrulea, *H. B. & K. Nov. Gen. et Sp.* v. 163, in obs. —Ind. occ.
cœrulea, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 27 = anoptera.
campestris, *A. Juss. l. c.* 33.—Bras.
Candolleana, A. Juss. in Arch. Mus. Par. iii. (1843) 463 = suberosa.
carinata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 183.—Guian.
catlagarum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 448.—Bras.
chrysophylla, *H. B. & K. Nov. Gen. et Sp.* v. 163, in obs.—Am. austr.
cinerascens, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 181.—Guiana.

HETEROPTERIS:—
cochleosperma, *A. Juss. in Arch. Mus. Par.* iii. (1843) 457.—Bras.
cœrulea, Auct. = cærulea.
coleoptera, A. Juss. in Arch. Mus. Par. iii. (1843) 448 = acuta.
confertiflora, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 34.—Bras.
cordifolia, *Moric. Pl. Nouv. Am.* 108. t. 66.—Bras.
coriacea, *A. Juss. in Arch. Mus. Par.* iii. (1843) 467. —Bras.
cornifolia, *H. B. & K. Nov. Gen. et Sp.* v. 165.—N. Granat.
cotinifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 438.—Mexic.
crenulata, *Mart. ex Griseb. in Mart. Fl. Bras.* xii. i. 62.—Bras.
cringens, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 136.—Bras.
cristata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 181.—Guiana.
cynanchoides, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 141.—Bras.
discolor, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 32.—Bras.
diversifolia, A. Juss. ex Griseb. in Mart. Fl. Bras. xii. i. 63 = sceroides.
Duartenea, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 28. —Bras.
eglandulosa, *A. Juss. l. c.* 27.—Bras.
excalliscifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 468.—Bras.
falcifera, *A. Juss. l. c.* 440.—Bolivia.
floribunda, *H. B. & K. Nov. Gen. et Sp.* v. 166.—Mexic.
fulgens, R. Grah. in Edinb. N. Phil. Journ. (Jan.-Mar. 1838) 391 = nitida!
Gaynna, *A. Juss. in Arch. Mus. Par.* iii. (1843) 459.—Mexic.
glabra, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 157 = umbellata.
grandiflora, *A. Juss. in Arch. Mus. Par.* iii. (1843) 461.—Bras.
heliciua, *Griseb. in Mart. Fl. Bras.* xii. i. 67.—Bras.
hirsutoides, A. Juss. ex Griseb. l. c. 63 = venosa.
hirsutoides, Mart. Herb. Fl. Bras. n. 1173 = Hiraea chlorocarpa.
hypericifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 34.—Bras.
ilicifolia, *Griseb. in Linnaea*, xiii. (1839) 220.—Bras.
intermedia, Griseb. in Mart. Fl. Bras. xii. i. 62 = venosa.
Jussieui, Hook. f. in Hook. Niger Fl. 246 = africana.
Laurentii, *Regel, Ind. Sem. Hort. Petrop.* (1865) 50.—Am. austr.
laurifolia, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 536 = venosa.
laurifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 458. —Mexic.
Leschenaultiana, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 26.—Bras.
Lessertiana, A. Juss. in Arch. Mus. Par. iii. (1843) 462 = suberosa.
Lessertiana, Seem. Bot. Voy. Herald, 93 = floribunda.
Lindeniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 457.—Mexic.
Lindleyana, *A. Juss. l. c.* 446.—Bras.
longifolia, *H. B. & K. Nov. Gen. et Sp.* v. 166.—Mexic.
loniceraefolia, *Triana & Planch. in Ann. Sc. Nat. Sér. IV.* xvii. (1862) 324.—N. Granat.
macrostachys, *A. Juss. in Arch. Mus. Par.* iii. (1843) 450.—Afr. trop.
Martiana, *A. Juss. l. c.* 474.—Bras.
Mathewsiana, *A. Juss. l. c.* 454.—Peruv.
megaptera, *A. Juss. l. c.* 468.—Bras.
metallicolrox, A. Juss. l. c. 449 = acrea.
nervosa, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 26.—Bras.
nitida, *H. B. & K. Nov. Gen. et Sp.* v. 163, in obs.—Bras.
obtusata, *Griseb. in Linnaea*, xiii. (1839) 227.—Bras.
ochnaefolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 464.—Bras.
oleaefolia, Griseb. in Linnæa, xxii. (1849) 19 = Tetrapteris oleifolia.
oxiocoucha, *A. Juss. in Arch. Mus. Par.* iii. (1843) 468.—Venez.
pannosa, *Griseb. in Mart. Fl. Bras.* xii. i. 70.—Bras.

HETEROPTERIS :—
parvifolia, *DC. Prod.* i. 591.—Ind. occ.
parvifolia, Mart ex A. Juss. in Arch. Mus. Par. iii. (1843) 465 = ascalioniefolia.
patens, *A. Juss. l. c.* 487.—Bras.
pauciflora, *A. Juss. l. c.* 479.—Bras.
platypiera, *DC. Prod.* i. 593.—Am. trop.
pteropetala, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 31. t. 167.—Bras.
purpurea, *H. B. & K. Nov. Gen. et Sp.* v. 168, in obs. —Ind. occ.
racemosa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 445. —Bras.
Regnellii, Miq. in Linnæa, xxii. (1849) 548 = umbellata.
reticulata, *Griseb. in Mart. Fl. Bras.* xii. t. 71.—Bras.
rhopalæfolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 441.—Bras.
rubiginosa, *A. Juss. l. c.* 477.—Bras.
rufula, *Mart. ex A. Juss. l. c.* 476.—Bras.
saligna, *Mart. ex A. Juss. l. c.* 466.—Bras.
scandens, *Regel & Korn. Ind. Sem. Hort. Petrop.* (1858) 46.—Hab.!
sericea, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 29.—Bras.
sessilifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 480. —Bras.
Smeathmanni, DC. Prod. i. 592 = Acridocarpus Smeathmanni.
spectabilis, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 442.—Bras.
Spruceana, Mart. ex A. Juss. in Walp. Rep. v. 288 = discolor.
stanleæ, *Griseb. in Kjoeb. Vidensk. Meddel.* (1858) 46. —Am. cent.
suberosa, *Griseb. in Linnæa*, xiii. (1839) 229.—Bras.
sylvatica, *A. Juss. in Arch. Mus. Par.* iii. (1843) 469. —Bolivia.
syringæfolia, *Griseb. in Linnæa*, xiii. (1839) 224.—Bras.
ternstroemiæfolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 443.—Bras.
thyrsoides, *A. Juss. l. c.* 436.—Bras.
tomentosa, Hook. & Arn. Bot. Beech. Voy. 281 = Beecheyana.
tomentosa, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 31 = spectabilis.
trichanthera, *A. Juss. in Arch. Mus. Par.* iii. (1843) 488.—Bras.
trigoniæfolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 35.—Bras.
umbellata, *A. Juss. l. c.* 25.—Bras.
undulata, Tenore, Hort. Bot. Neap. (1840) : et ex C. Koch, Wochenschr. iii. (1860) 111 = umbellata.
venosa, *Griseb. in Linnæa*, xiii. (1839) 230.—Bras.
verbasciifolia, *Griseb. in Kjoeb. Vidensk. Meddel.* (1858) 186.—Bras.
Warmingiana, *Griseb. l. c.* 186.—Bras.
Wiedeana, A. Juss. in Arch. Mus. Par. iii (1843) 470 = Leschenaultiana.
Wydleriana, *A. Juss. l. c.* 452.—Ind. occ.
xanthophylla, *A. Juss. l. c.* 459.—Bras.

HETEROPTILIS, E. Mey. ex Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 534 = Selinum, Linn. (Umbellif.).
arenaria, E. Mey. ex Meisn. l. c. = S. suffruticosum.

HETEROPYXIS, Griff. Notul. iv. 594 ; Ic. (1854) t. 594 (Hetropyxis) = Boschia, Korth. (Malvac.).

HETEROPYXIS, Harv. Thes. Cap. ii. 18. t. 128 (1868). *LYTHRARIEÆ*, Benth. & Hook. f. i. 785. natalensis, *Harv. l. c.*—Afr. austr.

HETERORACHIS, Sch. Bip. in Flora, xxvii. (1844) 775, in syn. = Berchheya, Ehrh. (Compos.).
spinosissima, Sch. Bip. l. c. = B. palmata.

HETEROSCIADIUM, DC. Prod. iv. 93, in syn. (1830) = Petagnia, Guss. (Umbellif.).

HETEROSICYOS, Welw. in Trans. Linn. Soc. xxvii. (1869) 93 = Trochomeria, Hook. f. (Cucurb.).
polymorpha, Welw. l. c. 94 = T. polymorpha.
stenoloba, Welw. l. c. = T. stenoloba.

HETEROSMILAX, Kunth, Enum. Pl. v. 270 (1850). *LILIACEÆ*, Benth. & Hook. f. iii. 763.
OLIGOSMILAX, Seem. Journ. Bot. vi. (1868) 258. t. 83.
borneensis, *A. DC. Monog. Phan.* i. 43.—Borneo.

HETEROSMILAX :—
Gaudichaudiana, *A. DC. Monog. Phan.* i. 44.—China. indica, *A. DC. l. c.* 43.—Malaya.
japonica, *Kunth, Enum. Pl.* v. 270.—Japon.
sumatrensis, *A. DC. Monog. Phan.* i. 42.—Malaya.

HETEROSOMA, Guill. Arch. Bot. i. (1833) 550 = Heterotoma, Zucc. (Campanul.).

HETEROSPATHE, Schef. in Ann. Jard. Buitenz. i. (1876) 141, 162. *PALMÆ*, Benth. & Hook. f. iii. 906.
elata, *Scheff. l. c.* 162.—Ins. Amboin.

HETEROSPERMA, Cav. Ic. iii. 84. t. 267 (1794) = Heterospermum, Cav. (Compos.).

HETEROSPERMA, Tausch, in Flora, xvii. (1834) t. 857, *nomen* = Heteromorpha, Cham. & Schlecht. (Compos.).

HETEROSPERMUM, Cav. Ic. iii. 84. t. 267 (1794) (Heterosperma). *COMPOSITÆ*, Benth. & Hook. f. ii. 388.
MICRODONTA, Nutt. in Trans. Am. Phil Soc. N. S. vii. (1841) 869.
depressum, *Griseb. in Gott. Abh.* xix. (1874) 186.—Reg. Argent.
dicranocarpum, A. Gray, Pl. Wright. i. 109 = Dicranocarpus parviflorus.
diversifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 236 — Ecuador.
maritimum, H. B. & K. l. c. 245 = ovatum.
ovatifolium, Cav. Desc. 204 = ovatum.
ovatum, *Willd. Sp. Pl.* iii. 2129.—Am. austr.
pinnatum, *Cav. Ic.* iii. 84. t. 267 ; *Willd. Sp. Pl.* iii. 2129.—Mexic.
rhombifolium, *Griseb. in Goett. Abh.* xix. (1874) 187. —Am. austr.
tagetinum, *A. Gray, Pl. Fendl.* 87.—N. Mexic.
Xanti, *A. Gray, in Proc. Am. Acad.* v. (1861) 162.—Calif.

HETEROSTACHYS, Ung.-Sternb. in Atti, Congr. Bot. Firenze, 1874 (1876) 331, 373. *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 64.
SPINOSTACHYS, Ung.-Sternb. Vers. Syst. Salicorn. 100 (1866).
Ritteriana, *Ung.-Sternb in Atti, Congr. Bot. Firenze*, 1874 (1876) 352.—Am. trop.

HETEROSTALIS, Schott, in Oestr. Bot. Wochenbl. (1857) 381 = Typhonium, Schott (Aroid.).
diversifolia, Schott, l. c. = T. diversifolium.
flagelliformis, Schott, l. c. = T. cuspidatum.
foliosa, Schott, l. c. = T. diversifolium.
Huegeliana, Schott, l. c. = T. diversifolium.
podata, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 278 = T. fallax.

HETEROSTECA, Devx. in Nouv. Bull. Soc. Philom. ii. (1810) 188 ; Journ. Bot. iii. (1813) 68 = Boutelous, Lag. (Gramin.).
americana, Desv. Journ. Bot. iii. (1813) 68 = B. bromoides.
juncyformis, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = B. bromoides.

HETEROSTEGA, Kunth, in Mém. Mus. Par. ii. (1815) 73 = grass.

HETEROSTEGON, Schwein. ex Hook. f. in Trans. Linn. Soc. xx. (1847) 175, in nota = Boutelous, Lag. (Gramin.).
curtipendulus, Schwein. ex Hook. f. l. c. = B. racemosa.

HETEROSTEMMA, Wight & Arn. Contrib. 42 (1834). *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 774.
SYMPHYSICARPUS, Hassk. in Flora, xl. (1857) 101.
acuminatum, *Decne. in Ann. Sc Nat. Sér.* II. ix. (1838) 368.—Java.
alatum, *Wight & Arn. Contrib.* 42.—Ind. or.
cuspidatum, *Decne. in DC. Prod.* viii. 630.—Ins. Philipp.
Dalzellii, *Hook. f. Fl. Brit. Ind.* iv. 48.—Ind. or.
javanicum, *Hassk. in Flora*, xl. (1857) 101.—Malaya.
maxillense, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 865.—Ins. Philipp.

HETEROSTEMMA :—
stellatum, *Hook. f. Fl. Brit. Ind.* iv. 47.—Reg. Himal.
tanjorensis, *Wight & Arn. Contrib.* 42.—Ind. or.
urceolatum, *Dals. in Hook. Kew Journ.* iv. (1852) 295. —Ind. or.
Wallichii, Dals. & Gibs. Bomb. Fl. 152 = Dalzellii.
Wallichii, *Wight & Arn. Contrib.* 42.—Ind. or.

HETEROSTEMON, Desf. in Mém. Mus. Par. iv. (1818) 241. t. 12. *LEGUMINOSÆ*, Benth. & Hook. f. i. 578.
conjugatus, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 216.—Bras.
ellipticus, *Mart. ex Benth. l. c.* 215.—Bras.
impar, *Spruce, ex Benth. l. c.*—Bras.
mimosoides, *Desf. in Mém. Mus. Par.* iv. (1818) 248. t. 12.—Bras.
simplicifolia, Spruce, ex Benth. in Mart. Fl. Bras. xv. ii. 215 = ellipticus.

HETEROSTEMUM, Steud. Nom. ed. II. i. 756 (1840) = prace.

HETEROSTIGMA, Gandich. Bot. Voy. Bonite, t. 25 (1844-52) = Pandanus, Linn.
Heudelotianum, Gaudich. l. c. t. 26 = P. Heudelotianus.

HETEROSTYLUS, Hook. Fl. Bor. Am. ii. 171. t. 185 (1840) = Lilæa, Humb. (Naiad.).
graminea, Hook. l. c. = L. subulata.

HETEROTAENIA, Boiss. Voy. Bot. Espagne, ii. 267. t. 80 A. (1840) = Concopodium, Koch (Umbellif.).
arvensis, Coss. Notes Crit. 111.—Hispan.
Bourgæi, Coss. l. c. 110 = C. Bourgæi.
glaberrima, Boiss. Voy. Bot. Espagne, ii. 270 = *thalictrifolia*, Boiss. l. c. ii. 269. t.80.—Hispan.

HETEROTAXIS, Lindl. Bot. Reg. t. 1028 (1826) = Maxillaria, Ruiz & Pav. (Orchid.).
crassifolia, Lindl. l. c. = M. crassifolia.

HETEROTHALAMUS, Less. in Linnæa, v. (1830) 146. *COMPOSITÆ*, Benth. & Hook. f. ii. 387.
boliviensis, *Wedd. Chlor. Andd.* i. 179.—Bolivia.
bruniioides, *Less. in Linnæa*, vi. (1831) 504.—Bras.
nivalis, *Wedd. Chlor. Andd.* i. 179. t. 21 B.—Chili.
pascalioides, *Less. in Linnæa*, vi. (1831) 504.—Bras
spartioides, *Hook. & Arn. ex DC. Prod.* vii. 271.—Patagon.
trinervis, *Hook & Arn. in Hook. Journ. Bot.* iii. (1841) 48.—Reg. Argent.

HETEROTHECA, Cass. in Bull. Soc. Philom. (1817) 137. *COMPOSITÆ*, Benth. & Hook. f ii. 351.
CALYCIUM, Ell. Sketch, ii. 339 (1824).
DIPLOCOMA, D. Don, in Sweet, Brit. Flow. Gard. t. 246 (1838).
STELMANIS, Rafin. Fl. Tellur. ii. 47 (1836).
Chrysopsidis, DC. Prod. v. 317 = Lamarckii.
floribunda, Benth. Bot. Voy. Sulph. 34 = grandiflora.
grandiflora, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 315.—Am. bor. occ.
inuloides, *Cass. in Dict. Sc. Nat.* li. 460.—Mexic. Lamarckii, *Cass. l. c.* xxi. 130.—Am. bor.
Lamarckii, DC. Prod. v. 317 = Aplopappus divaricatus.
latifolia, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 459 = Lamarckii.
leptoglossa, DC. Prod. v. 317 = Lamarckii.
scabra, DC. l. c. = grandiflora.

HETEROTHRIX, Muell. Arg. in Mart. Fl. Bras. vi. 138. t. 40 (1860) = Richtera, Lian. (Apocyn.).
pycnantha, Muell. Arg. l. c. = E. pycnantha.
Vanheurckii, Muell. Arg. in Heurck, Obs. Bot. 164.—Bras.

HETEROTIS, Benth. in Hook. Niger Fl. 347 (1849) = Dissotis, Benth. (Melastom.).
enterantha, Benth. l. c. 349 = D. antennina.
capitata, Benth. l. c. 352 = Tristemma capitatum.
cornifolia, Benth. l. c. 351 = Tristemma neglectum.
laevis, Benth. l. c. 348 = D. decumbens.
plumosa, Benth. l. c. = D. plumosa.
prostrata, Benth. l. c. 349 = D. prostrata.
segregata, Benth. l. c. 350 = Tristemma segregatum.

HETEROTIS:—
thesiifolia, Benth. in Hook. Niger Fl. 351 = Tristemma thesiifolium.
triplinervia, Klotzsch, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 58 = Disactis prostrata.
Vogelii, Benth. in Hook. Niger Fl. 353 = Tristemma capitatum.

HETEROTOMA, Zucc. in Flora, xv. (1832) ii. Beibl. 100. *CAMPANULACEAE*, Benth. & Hook. f. ii. 553.
Myopsis, Presl, Prod. Mon. Lobel. 8 (1836).
arabidoides, *Benth. & Hook. f. Gen.* ii. 553.—Mexic. intermedia, *Hemsl. Biol. Centr. Am. Bot.* ii. 269.—Mexic.
lobelioides, *Zucc. in Flora*, xv. (1832) ii. *Beibl.* 101. Mexic.
macrocentron, *Benth. in Hook. Ic. Pl.* t. 1177.—Mexic
tenella, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 175.—Mexic.

HETEROTRICHUM, Bieb. Fl. Taur. Casc. iii. 551 (1819) = Saussurea, DC. (Compos.).
alpinum, Link, Enum. Hort. Berol. ii 298 = S. alpina.
arcticum, Bieb. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1869) ii. 43 = S. alpina.
atriplicifolium, Bieb. ex Herd. l. c. 19 = S. serrata.
dissectum, Bieb. ex Herd. l. c. 46 = S. japonica.
glabellum, Bieb. ex Mey. in DC. Prod. vi. 534 = S. serrata.
leptophyllum, Bieb. ex DC. l. c. 533 = S. salicifolium.
pulchellum, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 71 = S. japonica.
salsum, Bieb. Fl. Taur. Casc. iii. 551 = S. crassifolia.
Schangini, Bieb. ex DC. Prod. vi. 532 = S. pygmaea.
squalidum, Bieb. in Ledeb. Fl. Ross. ii. 656 = S. japonica.
Stelleri, Bieb. l. c. = S. elongata.

HETEROTRICHUM, DC. Prod. iii. 173 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 765.
Diplodonta, Karst. in Linnaea, xxviii. (1856) 439.
Octomeris, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1845) 378, partim.
angustifolium, *DC. Prod.* iii. 173.—Ins. S. Doming.
dubium, Macfad. Fl. Jamaic. ii. 65 = octonum.
hispidum, Griseb. in Goett. Abh. ix. (1861) 52 = octandrum.
lucidum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 184.—Venezuela.
macrodon, *Planch. in Hook. Bot. Mag.* t. 4421.—Venezuela.
niveum, DC. Prod. iii. 173 = patens.
novemnervium, DC. l. c. = Clidemia novemnervia.
octandrum, *Macfad. Fl. Jamaic.* ii. 66.—Ind. occ.
octonum, *DC. Prod.* iii. 173.—Am. trop.
patens, *DC. l. c.*—Ins. S. Doming.
Schlimii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 134.—Venezuela.
strigosum, *Triana, l. c.* 184.—Bras.
viscosum, Macfad. Fl. Jamaic. ii. 65 = octonum.

HETEROTROPA, Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1884) 314. t. 10 = Asarum, Linn. (Aristoloch.).
arifolia, A. Gray, in Hook. Lond. Journ. Bot. i (1842) 225 = A. arifolium.
asaroides, Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1884) 314 = A. albiwenium.
parviflora, Hook. f. Bot. Mag. t. 5380 = A. elegans.
virginica, A. Gray, in Hook. Lond. Journ. Bot. i. (1842) 225 = A. virginicum.

HETEROZYGIS, Bunge, Suppl. Fl. Alt. 82 (1836) (Heterozygia) = Tribulus, Linn. (Zygophyll.).

HETERYTA, Rafin. in Med. Repos. N. York, v. (1808) 353 = Phaseolia, Jus. (Hydrophyll.).
paleoneioides, Rafin. l. c.—Am. bor.

HETREPTA, Rafin. Fl. Tellur. iii. 88 (1836) = Leucosa, R. Br. (Labiat.).
lavandulifolia, Rafin. l. c. = L. linifolia.

HETTLINGERIA, Neck. Elem. ii. 124 (1790) = Rhamnus, Linn.

HEUCHERA, Linn. Syst. ed. I (1735). *SAXIFRAGEAE*, Benth. & Hook. f. i. 636.
Holochloa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 580 (1840).
Leptaxis, Rafin. Fl. Tellur. ii. 75 (1836).
Oreanthus, Rafin. ex Ser. Bull. Bot. i. (1830) 216.
Oreotyx, Rafin. Atl. Journ. 145 (1832); Fl. Tellur. ii. 74 (1836).
Tiarella, Rafin. Fl. Tellur. ii. 75 (1836).
acerifolia, Kafin. Med. Fl. i. 211. t. 49 = villosa.
americana, Georgi, Beschr. Russ. Reich. iv. 830 (Quid?).—Rossia.
americana, *Linn. Sp. Pl.* 226.—Am. bor.
Barbarossa, Presl, Rel. Haenk. ii. 56 = micrantha.
bracteata, *Ser. in DC. Prod.* iv. 52.—Am. bor. occ.
californica, Kellogg, in Proc. Calif. Acad. v. (1873) 58 = Tiarella unifoliata.
caulescens, Pursh, Fl. Am. Sept. i. 188 = villosa.
Cortusa, Michx. Fl. Bor. Am. i. 171 = americana.
Curtisii, *Torr. & Gray, ex A. Gray, in Am. Journ. Sc.* Ser. I. xlii. (1842) 15.—Am. bor.
cylindrica, *Dougl. in Hook. Fl. Bor. Am.* i. 236.—Am. bor. occ.
dichotoma, Murr. in Comm. Gotting. (1772) 84. t. 1 = Vahlia Oldenlandia.
divaricata, Fisch. ex DC. Prod. iv. 51 = glabra.
erubescens, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1871) App. 2 = micrantha.
foliosa, Rafin. Med. Fl. i. 244 = villosa.
glabella, Torr. & Gray, Fl. N. Am. i. 581 = cylindrica.
glabra, *Willd. ex Roem. & Schult. Syst.* vi. 216.—Am. bor. occ.
glauca, Rafin. Med. Fl. i. 244 = Rugelii.
grandiflora, Rafin. l. c. = pubescens.
Hallii, *A. Gray, in Proc. Acad. Sc. Philad.* 1868 (Aug.) 62.—Am. bor. occ.
hirtiflora, Torr. & Gray, Fl. N. Am. i. 582 = pilosissima.
hispida, Hook. & Arn. Bot. Beech. Voy. 347 = pilosissima.
hispida, *Pursh, Fl. Am. Sept.* i. 188.—Am. bor.
longipetala, *Moç. ex Ser. in DC. Prod.* iv. 52.—Mexic.
lucida, Schlecht. Ind. Sem. Hort. Hal. (1846) 8.—Cf. Linnaea, xxiv. (1851) 186 = americana.
macrophylla, Lodd. ex G. Don, in Loud. Hort. Brit. 99 (Quid?).—Am. bor.
Menziesii, Hook. Fl. Bor. Am. i. 237 = Tolmiea Menziesii.
micrantha, *Dougl. in Lindl. Bot. Reg.* t. 1302.—Am. bor. occ.
minutiflora, *Hemsl. Diagn. Fl. Nov.* 50.—Mexic.
obtusissima, *Regel & Rach, Ind. Sem. Hort. Petrop.* (1856) 31.—Am. bor.
orizabensis, *Hemsl. Diagn. Fl. Nov.* 50.—Mexic.
ovalifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 581 = cylindrica.
parviflora, Bartl. Ind. Sem. Hort. Gotting. (1838).—Cf. Linnaea, xiii. (1839) Litt. 96 = villosa.
parviflora, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 581. —Am. bor. occ.
pilosissima, Benth. Pl. Hartw. 311 = micrantha?
pilosissima, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 56.—Calif.
Portula, Michx. Fl. Bor. Am. i. 171 = americana.
pubescens, *Pursh, Fl. Am. Sept.* i. 187.—Am. bor.
pubescens, Rafin. Med. Fl. i. 243 = pubescens.
racemosa, *S. Wats. in Proc. Am. Acad.* xx. (1885) 365.—Am. bor. occ.
rosiflora, Rafin. Med. Fl. i. 244 = pubescens.
ribifolia, Fisch. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 62 = pubescens.
Richardsonii, R. Br. in Frankl. Narr. 1st Voy. 766. t. 29 = hispida.
rubescens, *Torr. in Stansb. Exped. Utah*, 388.—Am. bor. occ.
Rugelii, *Shuttl. ex Kunze, in Linnaea*, xx. (1847) 48. —Florida.
sanguinea, *Engelm. in Wisliz. Tour N. Mexico*, 107. —N. Mexic.
scapifera, Moench, Meth. 674 = americana.
squamosa, Rafin. Med. Fl. i. 244 = villosa.
undulata, *Regel & Rach. Ind. Sem. Hort. Petrop.* (1856) 32.—Am. bor.
villosa, *Michx. Fl. Bor. Am.* i. 172.—Am. bor.
viscida, Pursh, Fl. Am. Sept. i. 187 = americana.

HEUDELOTIA, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 150. t. 39 (1832) = Commiphora, Jacq. (Burser.).
africana, A. Rich. l. c. = C. africanum.

HEUDUSA, E. Mey. Comm. Pl. Afr. Austr. 158 (1835) = Lasthrioguna, Eckl. & Zeyh. (Legumin.).
candicans, Presl, Bot. Bemerk. 45 = L. candicans.
decipiens, E. Mey. Comm. Pl. Afr. Austr. 158 = L. parvifolia.

HEUFFELIA, Schur, Enum. Pl. Trans. 766 (1866) = Avena, Linn. (Gramin.).
australis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 183 = A. pratensis.
bromoides, Schur, Enum. Pl. Transs. 762 = A. bromoides.
compressa, Schur, l. c. = A. compressa.
convoluta, Schur, l. c. 763 = A. convoluta.
laevigata, Schur, l. c. 761 = A. pubescens.
lucida, Schur, l. c. = A. pubescens.
montana, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 183 = A. montana.
planiculmis, Schur, Enum. Pl. Transs. 762 = A. planiculmis.
pratensis, Schur, l. c. = A. pratensis.
pubescens, Schur, l. c. = A. pubescens.
Scheuchzeri, Schur, l. c. 763 = A. Scheuchzeri.
sempervirens, Schur, l. c. = A. sempervirens.
setacea, Schur, l. c. = A. setacea.

HEURCKIA, Muell. Arg. in Flora, liii. (1870) 166 = Baillonella, Linn. (Apocyn.).
semperflorens, Muell. Arg. l. c. 169.—N. Caled.
viridis, Muell. Arg. in Heurck, Obs. Bot. 205.—N. Caled.

HEURLINIA, Rafin. Sylva Tellur. 166 (1838) = Myrsine, Linn.
variabilis, Rafin. l. c. = M. variabilis.

HEURNIA, Spreng. Anleit. ii. 148 (1817) = Huernia, R. Br. (Asclepiad.).

HEVEA, Aubl. Pl. Gui. ii. 871. t. 335 (1775). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 290.
Micrandra, R. Br. in Benn. Pl. Jav. Rar. 237 (1844).
Siphonanthus, Schreb. ex Baill. Etud. Gén. Euph. 334 (1858).
Siphonia, Rich. in Schreb. Gen. 656 (1791).
apiculata, Baill. Adansonia, iv. (1863–64) 295 = lutea.
Benthamiana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 204.—Bras.
brasiliensis, *Muell. Arg. l. c.*—Bras.
discolor, *Muell. Arg. in DC. Prod.* xv. ii. 737.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 871.—Guian.
janeirensis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 706. —Bras.
lutea, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 204.—Bras.
membranacea, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 299.—Bras.
nitida, *Muell. Arg. l. c.* 301.—Bras.
paraensis, Baill. Adansonia, iv. (1863–64) 284, in syn. = discolor.
panciflora, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 203.—Bras.
parviflora, Anbl. Pl. Gui. iv. t. 335 = guianensis.
rigidifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 203.—Bras.
Spruceana, *Muell. Arg. l. c.* 204.—Bras.

HEWARDIA, Hook. Ic. Pl. t. 858 (1862). *LILIACEAE*, Benth. & Hook. f. iii. 287.
Isophysis, T. Moore, in Proc. Linn. Soc. ii. (1858) 212.
tasmanica, *Hook. l. c.*—Tasmania.

HEWETIA, Pritz. Ic. Ind. i. 544 (1885) = sec.

HEWITTIA, Wight & Arn. in Madr. Journ. Sc. Ser. I. v. (1837) 23. *CONVOLVULACEAE*, Benth. & Hook. f. ii. 872.
Ketmoria, Rafin. Fl. Tellur. iv. 78 (1836), nomen prius.
Palmia, Endl. Gen. 853 (1838).

HEWITTIA:—
Sanilum, Rafin. Fl. Tellur. iv. 70 (1836), nomen prius.
Shuteria, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 465 (Conv. or. 109).
acerifolia, Klotzsch, in Peters, Reise Mossamb. Bot. 243 = bicolor.
bicolor, *Wight & Arn. in Madr. Journ. Sc.* Ser. I. v. (1837) 22.—Geront. trop.
cordifolia, Steud. Nom. ed. II. i. 757 = Ipomaea linifolia.
hirta, Klotzsch, in Peters, Reise Mossamb. Bot. 243 = bicolor.

HEXABOLUS, Steud. Nom. ed. II. i. 757 (1840), sphalm. = **Hexalobus**, A. DC. (Anonac.).

HEXACADICA, Rafin. Sylva Tellur. 158 (1838) = **Hexadica**, Lour. (Inc. sed.).
corymbosa, Rafin. l. c. = *Hexadica cochinchinensis* (nom. delend.)

HEXACENTRIS, Nees, in Wall. Pl. As. Rar. iii. 78 (1832) = **Thunbergia**, Linn. f. (Acanth.).
acuminata, Nees, l. c. = T. coccinea.
coccinea, Nees, l. c. 78 = T. coccinea.
dentata, Nees, l. c. = T. coccinea.
lutea, Lindl. in Gard. Chron. (1854) 151 = T. lutea.
mysorensis, Wight, Ic. t. 871 = T. mysorensis.

HEXACHLAMYS, Berg, in Linnaea, xxvii. (1854) 345 = **Eugenia**, Linn. (Myrtac.).
humilis, Berg, l. c.—Reg. Argent.

HEXACTINA, Willd. ex Schlecht. in Schult. f. Syst. vii. 91 (1829) = **Amafoua**, Aubl. (Rubiac.).
corymbosa, Willd. l. c. = A. corymbosa.

HEXADENA, Rafin. Sylva Tellur. 92 (1838) = **Phyllanthus**, Linn. (Euphorb.).
angustifolia, Rafin. l. c. = P. angustifolium.

HEXADENIA, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 253; et in Abh. Akad. Berl. (1860) 106 = **Pedilanthus**, Neck. (Euphorb.).
macrocarpa, Klotzsch & Garcke, ll. cc. 253; 107 = P. macrocarpus.

HEXALEBESMIA, Brongn. in Ann. Sc. Nat. Sér. II. xvii. (1842) 44. *ORCHIDEAE*, Benth. & Hook. f. iii. 530.
Hexopia, Batem. ex Lindl. in Bot. Reg. (1840) Misc. 46, nomen; et (1844) Misc. 2.
bicornis, *Lindl. Bot. Reg.* (1844) *Misc.* 41.—N. Granat.
bifida, *Reichb. f. in Saund. Refug. Bot.* ii. sub t. 113. Mexic.
brachyphylla, *Reichb. f. Beitr. Orch. Centr. Am.* 89.—Costa Rica.
crurigera, *Lindl. Bot. Reg.* (1844) *Misc.* 2.—Guatemala.
fasciculata, *A. Br. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 45.—Mexic.
fusiformis, *Griseb. Fl. Brit. W. Ind.* 623.—Ind. occ.
Lindeniana, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 23 = fasciculata.
lurida, *Batem. Syn. Orch. in Orch. Mex. et Guat.*—Mexic.
micrantha, *Lindl. in Bot. Reg.* (1844) *Misc.* 2.—Panama.
monophylla, *Rudrig. Orch. Nov.* i. 80.—Bras.
Reedii, *Reichb. f. in Saund. Refug. Bot.* t. 113.—Bras.
rhodoglossa, *Reichb. f. in Bonplandia*, iv. (1856) 328.—Mexic.
sessilis, *Reichb. f. in Saund. Refug. Bot.* ii. sub t. 113; *et in Linnaea*, xlii. (1877) 84.—Mexic.
stenopetala, *Reichb. f. in Bonplandia*, iii. (1855) 231.—Am. austr.

HEXADICA, Lour. Fl. Cochinch. 562 (1790), nomen delendum.—*Cf.* Benth. & Hook. f. Gen. i. 981; iii. 256.
Hexacadica, Rafin. Sylva Tellur. 158 (1838).
cochinchinensis, Lour. l. c.—Cochinch.

HEXAGLOTTIS, Vent. Dec. Gen. Nov. 6 (1808). *IRIDEAE*, Benth. & Hook. f. iii. 692.
Plantia, Herb. in Bot. Reg. (1844) Misc. 89.
flexuosa, Sweet, Hort. Brit. ed. II. 498 = longifolia.
longifolia, *Vent. Dec. Gen. Nov.* 6.—Afr. austr.
virgata, *Sweet, Hort. Brit.* ed. II. 498.—Afr. austr.

HEXAGONOTHECA, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) 11. 505 = **Berrya**, Roxb. (Lillac.).
cordata, Turcz. l. c. = *B. Amomella*.

HEXALECTRIS, Rafin. Neogenyt. 4 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 498.
aphylla, *Rafin. l. c.*—Mexic.
squamosa, Rafin. Fl. Tellur. iv. 48 = aphylla.

HEXALEPIS, Boeck. in Flora, lviii. (1875) 118 = **Gahnia**, Forst. (Cyperac.).
scabrifolia, Boeck. l. c. = G. aspera.

HEXALEPIS, Rafin. Fl. Tellur. iv. 24 (1836) = **Thilandsia**, Linn. (Bromeliac.).
pittacina, Rafin. l. c. = T. pittacina.

HEXALOBUS, A. DC. in Mém. Soc. Phys. Genèv. v. (1833) 212 (Mém. Anon. 36. t. 5 A). *ANONACEAE*, Benth. & Hook. f. i. 24.
brasiliensis, *A. DC. in Fl. Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 138.—Bras.
calicarpus, Baill. in Bull. Soc. Par. i. (1882) 338 = Uvaria calicarpa.
crispiflorus, *A. Rich. Fl. Cub.* i. 43.—Afr. trop.
grandiflorus, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 468.—Afr. trop.
Jussiaeanus, *Baill. Adansonia*, viii. (1867–68) 348.—Hab.?
madagascariensis, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 213.—Madag.
senegalensis, *A. DC. l. c.*—Afr. trop.

HEXAMERIA, R. Br. in Benn. Pl. Jav. Rar. 26. t. 7 (1838) = **Podochilus**, Blume (Orchid.).
distichа, R. Br. l. c. = P. serpyllifolia.

HEXAMERIA, Torr. & Gray, in Torr. Rep. Fl. N. York, 137 (1839) [Geol. Surv.] = **Bohtnocystis**, Torr. & Gray (Cucurb.).
echinata, Torr. & Gray, l. c. = E. lobata.

HEXANTHUS, Lour. Fl. Cochinch. 195 (1790) = **Litsea**, Lam. (Laurin.).
umbellatus, Lour. l. c. = L. Hexanthus.

HEXAPHOMA, Rafin. Fl. Tellur. ii. 66 (1836) = **Asclepiastrum**, Tourn.
ferruginea, Rafin. l. c. = S. leucanthemifolia.
petiolaria, Rafin. l. c. 67 = S. rivularis.
pyrenaica, Rafin. l. c. = S. mucoides?
virgata, Rafin. l. c. = S. viscosa?

HEXAPLECTRIS, Rafin. Fl. Tellur. iv. 97 (1836) = **Aristolochia**, Tourn.
bicolor, Rafin. l. c.—Hab.?
fuscata, Rafin. l. c.—Hab.?

HEXAPTERA, Hook. Bot. Misc. i. (1830) 350. t. 73, 74. *CRUCIFERAE*, Benth. & Hook. f. i. 90.
cicutifolia, *Phil. in Anal. Univ. Chil.* (1862) ii. 442.—Chili.
cuneata, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 353. t. 74.—Chili.
frigida, *Phil. Fl. Atac.* 8.—Chili.
Junciaci, *Barn. in C. Gay, Fl. Chil.* i. 179. t. 5.—Chili.
linearis, *Barn. l. c.* 176. t. 5.—Chili.
littoralis, *Barn. l. c.* 177.—Chili.
pinnatifida, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 350. t. 72.—Chili.
pusilla, *Phil. in Anal. Univ. Chil.* (1872) 674.—Chili.
scapigera, *Phil. l. c.* (1862) 11. 359.—Chili.
spathulata, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 351. t. 73.—Chili.
violacea, *Phil. in Anal. Univ. Chil.* (1872) 673.—Chili.

HEXARRHENA, J. & C. Presl, Rel. Haenk. i. 326. t. 45 (1830) = **Hilaria**, H. B. & K. (Gramin.).
amchroides, J. & C. Presl, l. c.—Mexic.

HEXASEPALUM, Bartl ex DC. Prod. iv. 561 (1825). *RUBIACEAE*, Benth. & Hook. f. ii. 145.
angustifolium, *Bartl. ex DC. l. c.*—Mexic.

HEXASTEMON, Klotzsch, in Linnaea, xii. (1838) 220 = **Jacaranda**, D. Don (Bignon.).
lanatus, Klotzsch, l. c. = J. lanata.

HEXASTYLIS, Rafin. Fl. Tellur. iii. 78 (1836) = **Asarum**, Tourn. (Aristolochiac.).
arifolium, Rafin. l. c. = A. arifolium.

HEXATHECA, C. B. Clarke, in DC. Monog. Phan. v. 198 (1883). *GESNERACEAE*.
fulva, *C. B. Clarke, l. c.* 194.—Borneo.

HEXATHECA, Sond. ex F. Muell. Fragm. viii. 217 (1874) = **Lepilaena**, Drumm. (Naiad.).
australis, Sond. ex Benth. Fl. Austral. vii. 180 = L. australis.
Preissii, Sond. ex F. Muell. Fragm. viii. 217 = L. Preissii.

HEXEPTA, Rafin. Sylva Tellur. 164 (1838) = **Coffea**, Linn. (Rubiac.).
axillaris, Rafin. l. c. = C. Zanguebariae.
racemosa, Rafin. l. c. = C. racemosa, *Lour.*

HEXISEA, Lindl. in Hook. Journ. Bot. i. (1834) 7. *ORCHIDEAE*, Benth. & Hook. f. iii. 524.
Enothonara, Reichb. f in Bot. Zeit. x. (1852) 772.
bidentata, *Lindl. in Hook. Journ. Bot.* i. (1834) 8.—Panama.
imbricata, *Reichb. f. in Walp. Ann.* vi. 470.—Mexic.
opposititifolia, *Reichb. f. l. c.*—Mexic.
reflexa, *Reichb. f. in Linnaea*, xli. (1877) 131.—Guiana.

HEXODONTOCARPUS, Dulac, Fl. Hautes-Pyr. 467 (1867) = **Sherardia**, Linn. (Rubiac.).
arvensis, Dulac, l. c. = S. arvensis.

HEXONYCHIA, Salisb. Gen. Pl. Fragm. 88 (1866) = **Allium**, Tourn. (Lillac.).
stellatum, Salisb. l. c. 89 = A. stellatum.

HEXOPEA, Steud. Nom. ed. II. i. 757 (1840) = seq.

HEXOPIA, Batem. ex Lindl. in Bot. Reg. (1840), Misc. 46, nomen; et (1844), Misc. 2 = **Hexadesmia**, Brongn. (Orchid.).
crurigera, Batem. ex Lindl. l. c. = Hexad. crurigera.

HEXORIMA, Rafin. in Med. Rep. N. York, v. (1808) 351.—Cf. Desv. Journ. Bot. ii. (1809) 168 = **Streptopus**, Michx. (Liliac.).
dichotoma, Rafin. l. c. = S. roseus.

HEXORINA, Steud. Nom. ed. II. i. 757 (1840) = prec.

HEXOSTEMON, Rafin. Cat. 14 (1824) = **Lythrum**, Linn.
virgatum, Rafin. l. c. = L. virgatum.

HEXOTRIA, Rafin. in Am. Monthly Mag. iv. (1818) 265 = Ilex, Tourn.
lanceolata, Rafin. l. c. = I. lanceolata? *Chapm.*

HEXURIS, Miers, in Proc. Linn. Soc. ii. (1850) 73; et in Trans. Linn. Soc. xxi. (1853) 44 = **Triuris**, Miers.
Gardneri, Miers, ll. cc. = T. lutea.

HEYDERIA, C. Koch, Dendrol. ii. 11. 177 (1872) = **Liboscdrus**, Endl. (Conif.).
decurrens, C. Koch, l. c. 179 = L. decurrens.

HEYDIA, Dennst. Schlüssel. Hort. Malab. 30 (1818) = **Solenopyrum**, Arn. (Santalac.).
horrida, Dennst. l. c. = S. Wallichianum.

HEYDUSA, Walp. Ann. i. 904 (1848–49) = HEUDUSA, Presl = **Lathrogyne**, Eckl. & Zeyh. (Legumin.).

HEYFELDERA, Sch. Bip. in Flora, xxxvi. (1853) 35 = **Chrysopsis**, Nutt. (Compos.).
sericea, Sch. Bip. l. c. 36 = C. graminifolia.

HEYLANDIA, DC. Mém. Lég vi. 198. t. 84; Prod. ii. 138 (1825). *LEGUMINOSÆ*, Benth. & Hook. f. i. 479.
GONIOGYNA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 91.
cordifolia, R. Grah. in Wall. Cat. n. 5343 = Indigofera cordifolia.
hebecarpa, DC. Mém. Lég. vi. 200. t. 84 = latebrosa.
latebrosa, *DC. l. c.* 201.—Ind. or.
leiocarpa, DC. l. c. 200 = latebrosa.
Perrottetii, Spreng. Syst. iv. Cur. Post. 267 = Crotalaria Perrottetii.

HEYLYGIA, G. Don, Gen. Syst. iv. 79 (1837) = HELYGIA, Blume = **Parsonsia**, R. Br. (Apocyn.).

HEYMASSOLI, Aubl. Pl. Gui. i. 324. t. 125 (1775) = **Ximenia**, Plum. (Olacin.).
inermis, Aubl. l. c. 325 = X. inermis.
spinosa, Aubl. l. c. 324 = X. americana.

HEYMIA, Dennst. Schlüssel. Hort. Malab. 35 (1818) = **Dentella**, Forst. (Rubiac.).
rigida, Dennst. l. c. = D. repens.

HEYNEA, Roxb. Hort. Beng. 33 (1814); et in Bot. Mag. t. 1738 (1815). *MELIACEÆ*, Benth. & Hook. f. i. 386.
affinis, A. Juss. in Mém. Mus. Par. xix. (1830) 285, 376 = trijuga.
cochlochinensis, *Baill. Adansonia*, xi. (1873–76) 265. —Cochinch.
connaroides, Wight, ex Voigt, Hort. Suburb. Calc. 186 = trijuga.
fruticosa, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 432 = sumatrana.
multijuga, Blume, Bijdr. 168 = Dysoxylon Blumei.
pubescens, Hook. f. Fl. Brit. Ind. i. 718 = trijuga.
quinquejuga, *A. Juss. in Mém. Mus. Par.* xix. (1830) 283.—Java.
quinquijuga, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 391 = trijuga.
sumatrana, *Mig. Fl. Ind. Bat. Suppl.* 505.—Sumatra.
trifoliata, A. Juss. in Mém. Mus. Par. xix. (1830) 285, 376 = Walsura piscidia.
trijuga, *Roxb. Hort. Beng.* 33; *et in Sims, Bot. Mag.* t. 1738.—Ind. or.

HEYNICHIA, Kunth, Ind. Sem. Hort. Berol. (1844) Adnot. 5.—Cf. Linnæa, xviii. (1844) 508 = Trichilia, Linn. (Meliac.).
malleodes, Kunth, l. c. = T. elegans.

HEYNICKIA, C. DC. in DC. Monog. Phan. i. 680 (1876) = praec.

HIBANTHUS, D. Dietr. Syn. Pl. i. 575 (1839) = HYBANTHUS, Jacq. = Ionidium, Vent. (Violac.).

HIBBERTIA, Andr. Bot. Rep. t. 136, 472 (1800). *DILLENIACEÆ*, Benth. & Hook. f. i. 14.
BURTONIA, Salisb. ex DC. Syst. i. 425 (1818).
HEMISTEMMA, Juss. ex Thou. Gen. Nov. Madag. 18 (1806).
HEMISTEPHUS, Drum. & Harv. in Hook. Kew Journ. vii. (1855) 51.
OCHROLASIA, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 8.
PLEURANDRA, Labill. Nov. Holl. Pl. ii. 5. t. 149 (1806).
TRICHOPHANDRA, Brongn. & Gris, in Ann. Sc. Nat. Sér. V. ii. (1864) 146.
WARBURTONIA, F. Muell. Fragm. i. 229 (1858–59).
acerosa, *Benth. Fl. Austral.* i. 24.—Austral.
acicularis, *F. Muell. Pl. Vict.* i. 17.—Austral.
amplexicaulis, *Steud. in Lehm. Pl. Preiss.* i. 266.—Austral.
angustifolia, Benth. *Fl. Austral.* i. 21.—Austral.
angustifolia, F. Muell. Pl. Vict. i. 18 partim = fasciculata.

HIBBERTIA:—
angustifolia, Salisb. Parad. Lond. sub t. 73 = procumbens.
argentea, *Steud. in Lehm. Pl. Preiss.* i. 266.—Austral.
aspera, DC. Syst. i. 430 = Billardieri.
aspera, Steud. in Lehm. Pl. Preiss. i. 270 = hyperícoides.
astrophylla, Steud. l. c. = furfuracea.
aurea, *Steud. l. c.* 272.—Austral.
Banksii, *Benth. Fl. Austral.* i. 20.—Austral.
baxteriana, Steud. in Lehm. Pl. Preiss. i. 268 = Candollia pachyrrhiza.
Baudouinii, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 191.—N. Caled.
Benthamii, F. Muell. Census, 1 = angustifolia.
Billardieri, *F. Muell. Vict. Pl.* i. 14.—Austral.
bracteata, *Benth. Fl. Austral.* i. 25.—Austral.
bracteata, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 140.—Austral.
Brownei, *Benth. Fl. Austral.* i. 21.—Austral.
bujotvaillifolia, Lehm. Nov. Hort. Hamb. (1851) 7 = amplexicaulis.
camphoroma, A. Gray, Bot. U. St. Expl. Exped. i. 21 = fasciculata.
candicans, *Benth. Fl. Austral.* i. 21.—Austral.
canescens, Sieber, ex Spreng. Syst. iv. Cur. Post. 211 = linearis.
cinerascens, Steud. in Lehm. Pl. Preiss. i. 271 = hyperícoides.
cistifolia, *R. Br. in DC. Syst.* i. 431.—Austral.
communata, Steud. in Lehm. Pl. Preiss. i. 267 = montana.
confertifolia, Steud. l. c. = montana.
coriacea, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 322. —Madag.
corifolia, Sims, Bot. Mag. t. 2672 = pedunculata.
crassifolia, *Benth. Fl. Austral.* i. 25.—Austral.
crenata, Andr. Bot. Rep. t. 472 = grossulariæfolia.
cuneiformis, Sm. in Rees, Cycl. xvii. n. 2 = Candollea cuneiformis.
Cunninghamii, *Ait. ex Hook. Bot. Mag.* t. 3183.—Austral.
dealbata, *Benth. Fl. Austral.* i. 21.—Austral.
densiflora, F. Muell. Pl. Vict. i. 15 = sericea.
dentata, *R. Br. in DC. Syst.* i. 426.—Austral.
deperdita, Steud. in Lehm. Pl. Preiss. i. 268 = Candollea fasciculata.
desmophylla, *F. Muell. Fragm.* xi. 95.—Austral.
diffusa, *R. Br. in DC. Syst.* i. 429.—Austral.
dilatata, A. Cunn. ex Benth. Fl. Austral. i. 27 = diffusa.
discolor, Steud. l. c. 267 = montana.
distincta, Lehm. Nov. Hort. Hamb. (1851).—Cf. Linnæa, xxv. (1852) 309 = amplexicaulis.
echilifolia, *R. Br. ex Benth. Fl. Austral.* i. 31.—Austral.
ericæfolia, Hook. f. Fl. Tasman. i. 14. t. 3 = serpyllifolia.
fasciculata, *R. Br. ex DC. Syst.* i. 428.—Austral.
furfuracea, *Benth. Fl. Austral.* i. 23.—Austral.
glaberrima, *F. Muell. Fragm.* iii. 1.—Austral.
glandulosa, Schlecht. in Linnæa, xx. (1847) 626 = fasciculata.
glomerata, *Benth. Fl. Austral.* i. 34.—Austral.
glomerosa, *F. Muell. Census*, 2.—Austral.
Goyderi, *F. Muell. Fragm.* vii. 123.—Austral.
gracilliflora, *Benth. Fl. Austral.* i. 28.—Austral.
grossulariæfolia, *Salisb. Parad. Lond.* n. 73. t. 73.—Austral.
hastata, Steud. in Lehm. Pl. Preiss. i. 266 = Cunninghamii.
hellanthemoides, *F. Muell. Census*, 2.—Austral.
hermanniæfolia, *DC. Syst.* i. 431.—Austral.
hirsuta, *Benth. Fl. Austral.* i. 26.—Tasman.
Holtzei, *F. Muell. in Wing. South Sc. Record* (Nov. 1885).—Austral.
Huegelii, *F. Muell. Fragm.* xi. 95.—Austral.
humifusa, *F. Muell. Vict. Pl.* i. 16. t. 1.—Austral.
Huttii, *F. Muell. Fragm.* vii. 123.—Austral.
hypericoides, *Benth. Fl. Austral.* i. 28.—Austral.
inclusa, *Benth. l. c.* 24.—Austral.
lactucæfolia, Steud. in Lehm. Pl. Preiss. i. 267 = Cunninghamii.
lasiopus, *Benth. Fl. Austral.* i. 40.—Austral.
latifolia, Steud. in Lehm. Pl. Preiss. i. 269 = grossulariæfolia.
latifolia, Benth. *Fl. Austral.* i. 22.—Austral.
leptodephylla, F. Muell. Fragm. i. 217 = microphylla.
lepidota, *R. Br. ex DC. Syst.* i. 426.—Austral.

HIBBERTIA:—
leptopus, *Benth. Fl. Austral.* i. 41.—Austral.
limosa, *Steud. in Lehm. Pl. Preiss.* i. 266.—Austral.
linearis, *R. Br. ex DC. Syst.* i. 428.—Austral.
lineata, *Steud. in Lehm. Pl. Preiss.* i. 272.—Austral.
longifolia, *F. Muell. Fragm.* iv. 115.—Austral.
lucens, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 191 = salicifolia.
melhanoides, *F. Muell. Fragm.* iv. 116.—Austral.
microphylla, *Steud. in Lehm. Pl. Preiss.* i. 273.—Austral.
monogyna, R. Br. ex DC. Syst. i. 429 = diffusa.
montana, *Steud. in Lehm. Pl. Preiss* i. 270.—Austral
mucronata, *Benth. Fl. Austral.* i. 39.—Austral.
Muelleri, *Benth. l. c.* 21.—Austral.
Mylnei, *Benth. l. c.* 39.—Austral.
nitida, *Benth. l. c.* 25.—Austral.
nutans, *Benth. l. c.* 40.—Austral.
obcuneata, Salisb. Parad. Lond. sub n. 73 = Candollea cuneiformis.
oblongata, *R. Br. ex DC. Syst.* i. 431.—Austral.
obtusifolia, DC. l. c. 429 = linearis.
Ochrolasia, *Benth. Fl. Austral.* i. 33; *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 549.—Austral.
oenotheroides, *F. Muell. Fragm.* vii. 87.—Austral.
ovata, Steud. in Lehm. Pl. Preiss. i. 270 = montana.
pachyrrhiza, Steud. l. c. 269 = Candollea pachyrrhiza.
pallida, Steud. l. c. 272 = aurea.
pedunculata, *R. Br. ex DC. Syst.* i. 430.—Austral.
perfoliata, *Hueg. ex Endl. Enum. Pl. Hueg.* 3.—Austral.
pilosa, *Steud. in Lehm. Pl. Preiss.* i. 272.—Austral.
polystachya, *Benth. Fl. Austral.* i. 29.—Austral.
potentilleefolia, *F. Muell. ex Benth. l. c.* 40.—Austral.
procumbens, *DC. Syst.* i. 427.—Austral.
prostrata, Hook. journ. Bot. i. (1834) 246 = fasciculata.
proxima, Steud. in Lehm. Pl. Preiss. i. 271 = hyperícoides.
pungens, *Benth. Fl. Austral.* i. 40.—Austral.
Readii, *Hort. ex Benth. Fl. Austral. Hist. Pl.* i. 99.—Hab. ?
recurvifolia, *Benth. Fl. Austral.* i. 34.—Austral.
rhadinopoda, *F. Muell. Fragm.* xi. 93.—Austral.
rostellata, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 8.—Austral.
salicifolia, F. Muell. Fragm. i. 161 = Adrastæa salicifolia.
salicifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 549.—N. Caled.
saligna, *R. Br. ex DC. Syst.* i. 427.—Austral.
scabra, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 191.—N. Caled.
scabra, *R. Br. ex Benth. Fl. Austral.* i. 31.—Austral.
scandens, Dryand. in Kon. & Sims, Ann. Bot. ii. (1807) 525 = volubilis.
sericea, *Benth. Fl. Austral.* i. 26.—Austral.
serpyllifolia, *R. Br. ex DC. Syst.* i. 430.—Austral.
spicata, *F. Muell. Fragm.* ii. 1.—Austral.
squamosa, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 9 = Candollea exasperata.
stellaris, *Endl. in Enum. Pl. Hueg.* 3.—Austral.
stricta, *R. Br. ex DC. Syst.* i. 425.—Austral.
subsericea, Steud. in Lehm. Pl. Preiss. i. 269 = Candollea pedunculata.
subvaginata, *F. Muell. Fragm.* xi. 95.—Austral.
synandra, *F. Muell. l. c.* iv. 151.—Austral.
tasmanica, *Benth. Fl. Austral. Adansonia*, vi. (1865–66) 264.—Tasmania.
tenuiramea, Steud. in Lehm. Pl. Preiss. i. 268 = stellaris.
tenuifolia, *F. Muell. Fragm.* iv. 117.—Austral.
tomentosa, *R. Br. ex DC. Syst.* i. 432.—Austral.
trachyphylla, Steud. in Lehm. Pl. Preiss. i. 271 = hyperícoides.
unolnata, *F. Muell. Census*, 2.—Austral.
vaginata, *F. Muell. Fragm.* xi. 96.—Austral.
velutina, *R. Br. ex Benth. Fl. Austral.* i. 30.—Austral.
verrucosa, *Benth. l. c.* 22.—Austral.
vestita, *A. Cunn. ex Benth. l. c.* 31.—Austral.
virgata, *R. Br. ex DC. Syst.* i. 428.—Australasia.
virgata, Hook. Ic. Pl. i. 107 = fasciculata.
volubilis, *Andr. Bot. Rep.* t. 126.—Austral.

HIBISCOS, St. Leg. in Ann. Soc. Bot. Lyon, vii. (1880) 197 = seq.

HIBISCUS, Linn. Gen. ed. I. 207 (1737). *MALVACEAE*, Benth. & Hook. f. i. 207.
ABELMOSCHUS, Medic. Malv. 45 (1787).
AZANZA, [Moc. & Sesse,] ex DC. Prod. i. 458 (1824).
BAMIA, [Linn.] R. Br. ex Wall. Pl. As. Rar. i 39 (1830).
BOMBYCODENDRON, Zoll. ex. Hassk. in Flora, xxx. (1847) 660.
COTYLOPLECTA, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 14.
ERKENNIA, Alef. l. c. (1863).
FURCARIA, Kostel. Allg. Med.-Pharm. Fl. v. 1856 (1856).
HYMENOCALYX, Zenk. Pl. Ind. 8. t. 10 (1885).
KETMIA, Tourn. ex Burm. Thes. Zeyl. 228 (1737).
LAGUNA, Cav. Diss. 178. t. 71. f. 1 (1787).
LAGUNARIA, Reichb. Consp. 203 (1828).
MUNCHUSIA, Heist. ex Rafin. Sylva Tellur. 115 (1838).
PARITI, Adans. Fam. ii. 401 (1763).
PARITIUM, St. Hil. Fl. Bras. Mer. i. 255 (1827).
PENTASPERMUM, DC. ex Steud. Nom. ed. II. i. 758, in syn. (1840).
PETITIA, Neck. Elem. ii. 407 (1790).
POLYCHLAENA, Garcke, in Schweinf. Beitr. Fl. Aethiop. 56 (1867).
SABDARIFFA, Kostel. Allg. Med.-Pharm. Fl. v. 1857 (1856).
SOLANDRA, Murr. in Comm. Gotting. 1783–84 (1788) 21. t. 1.
TRIGUERA, Cav. Diss. 41, nec 107 (1786).
TRIONAEA, Medic. Malv. 45 (1787).
TRIONUM, Linn. Syst. ed. I (1735).
TRIPLOCHITON, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 18.

Abelmoschus, *Linn. Sp. Pl.* 696.—Geront. trop.
abutiloides, *Willd. Enum. Hort. Berol.* 736.—Am. austr.
abyssinicus, *Steud. Nom.* ed. II. i. 755.—Abyssin.
acerifolius, DC. Prod. i. 448 = collinus.
acerifolius, *Salisb. Parad. Lond.* t. 33 —China.
acetosaefolius, DC. *Prod.* i. 455.—Mexic.
acetosa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 17 = vulpinus.
achanioides, *Hemsl. Biol. Centr.-Am. Bot.* i. 121.—Mexic.
aculeatus, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 678 = heterophyllus.
aculeatus, G. Don, Gen. Syst. i. 480 = surattensis, cannabinus.
aculeatus, Roxb. Fl. Ind. iii. 206 = furcatus.
aculeatus, *Walt. Fl. Carol.* 177.—Am. bor.
acuminatus, Cav. Diss. iii. 144. t. 50. f. 2 = Kosteletzkya acuminata.
adoënsis, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 29. —Abyssin.
adscendens, G. Don, Gen. Syst. i. 482 = physaliodes.
aestuans, Rottl. ex Mast. in Hook. f. Fl. Brit. Ind. i. 345 = mutabilis.
aestuans, Wall. Cat. n. 1908 E = collinus.
aethiopicus, *Linn. Mant.* ii. 258.—Afr. austr.
affinis, *H. B. & K. Nov. Gen. et Sp.* v. 289.—Am. austr.
africanus, Mill. Gard. Dict. ed. VIII. n. 20; Roth, Beitr. i. 42 = H. Trionum.
agioxillos, *Vell. Fl. Flum.* 228; vii. t. 35.—Bras.
albus, Wall. Cat. n. 1876 E = H. Solandra.
althaeaefolius, Shuttlew. ex A. Gray, Fl. Wright. i. 23 = Kosteletzkya virginica.
amblycarpus, Hochst. in Webb, Fragm. Fl. Aethiop. 45 = obtusilobus.
ambogoënsis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 518.—Madag.
amoenus, *Link & Otto, Ic. Pl. Select.* i. 119.—Reg. Argent.
anemoneflorus, Zipp. ex Span. in Linnaea, xv. (1841) 169 = virgatus.
angulosus, *Steud. Nom.* ed. II. i. 756; *Mast. in Hook. f. Fl. Brit. Ind.* i. 341.—Ind. or.
angustifolius, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 152.—Reg. Argent.
antanosarum, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 517.—Madag.
appendiculatus, Stokes, Bot. Mat. Med. iii. 542 = surattensis.
apricus, *Vell. Fl. Flum.* 224; vii. t. 40.—Bras.
aquaeflorus, DC. Fl. Fr. Suppl. 597 = roseus.

HIBISCUS:—
arborens, *Desr. in Ham. Prod. Pl. Ind. Occ.* 49.—Ind. occ.
arenarius, Murr. in Comm. Gotting. vii. (1785) t. 3 = Pavonia arenaria.
arenarius, Scop. Delic. Insub. iii. 3 = Pavonia zeylanica.
aristaeoides, Garcke, in Bot. Zeit. vii. (1849) 849 = intermedius.
aristatus, Hort. ex Poir. Encyc. v. 108 = Pavonia spinifera.
armenicaus, *Bouché, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 408.—Armenia.
Arnottianus, *A. Gray, Bot. U. St. Expl. Exped.* i. 176. —Ins. Sandvic.
Arnottii, Griff. ex Mast. in Hook. f. Fl. Brit. Ind. i. 344 = H. Rosa-sinensis.
asterisatus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 60. —Abyssin.
arvensis, *Vell. Fl. Flum.* 224; vii. t. 38.—Bras.
ascendens, Walp. Rep. i. 810 = *adscendens*.
aspera, Hook. f. in Hook. Niger Fl. 238 = cannabinus.
asperatus, Wall. Cat. n. 1908 B = collinus.
asperifolius, Eckl. & Zeyh. Enum. 38 = aethiopicus.
atromarginatus, *Eckl. & Zeyh. l. c.*—Afr. austr.
atrovioiaceus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 516.—Madag.
attenuatus, *Bosse, in Verh. Ver. Bef. Gartenb.* iv. t. 1. —Am. bor.
axillaris, *Vell. Fl. Flum.* 228; vii. t. 37.—Bras.
Azanzae, *DC. Prod.* i. 454.—Mexic.
azureus, Boj. ex Baker, in Journ. Linn. Soc. xx. (1883) 98 = Pavonia macrotis.
bacciferus, Forst. f. Prod. 48 = Thespesia populnea.
bahamensis, *Mill. Gard. Dict.* ed. VIII. n. 14.—Ins. Baham.
Bamia, *Tussetti, in Mus. Firens.* ii. 57. t. 7.—Aegypt.
Bancroftianus, Ham. ex Griseb. Fl. Brit. W. Ind 85 = phoeniceus.
Barclaianus, *Macfad. Fl. Jamaic.* i. 70.—Ind. occ.
bantamensis, *Miq. Fl. Sumgh.* 282.—Java.
barbatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 17 = vulpinus.
Bartlingianus, *Steud. Nom.* ed. II. i. 758.—Ind. or.
batacensis, *Blanco, Fl. Filip.* ed. I. 544.—Ins. Philipp.
Beckeri, F. Muell. Fragm. ii. 112 = diversifolius.
berberidifolius, A. Rich. Tent. Fl. Abyss. i. 56 = diversifolius.
Berlandierianus, *Moric. Pl. Nouv. Am.* 22. t. 15.— Mexic.
Bersieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 510. —Madag.
bifurcatus, H. B. & K. Nov. Gen. et Sp. v. 292 = phoeniceus.
betulinus, DC. Prod. i. 452 = phoeniceus.
bicornis, *G. F. W. Mey. Prim. Fl. Esseq.* 231.— Guiana.
biflorus, *Spreng. f. Tent. Suppl.* 19.—Afr. austr.
bijurcatus, Blanco, Fl. Filip. ed. I. 545 = surattensis.
bifurcatus, *Cav. Diss.* iii. 146. t. 51. f. 1.—Am. austr.
bifurcatus, Roxb. Hort. Beng. 51 = furcatus.
Boivini, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 510. —Madag.
Boivini, *Bélanger, Voy. Ind. Or.* (absque num.).—Ind. or.
Bojerianus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 511.—Madag.
borbonicus, Link, Enum. Hort. Berol. ii. 216 = calycinus.
borbonicus, Wight, in Wall. Cat. n. 3694 = canescens.
Boryanus, *DC. Prod* i. 446.—Ins. Borbon.
Boryanus, Hook. & Arn. Bot. Beech. Voy. 79 = Arnottianus.
brachycalyx, F. Muell. Fragm. iii. 5 = microchlaenus.
brachysiphonius, *F. Muell. l. c.* i. 67.—Austral.
Brackenridgei, *A. Gray, Bot. U. St. Expl. Exped.* i. 175.—Ins. Sandvic.
bractenus, *Vell. Fl. Flum.* 229; vii. t. 38.—Bras.
bracteosus, [*Moc. & Sesse,*] *ex DC. Prod.* i. 455.— Mexic.
brasiliensis, *Linn. Sp. Pl.* ed. II. 977.—Bras.
brevipes, Garcke, in Bot. Zeit. vii. (1849) 853 = crassinervis.
Burkealli, D. Dietr. Syn. Pl. iv. 689 = ureus.

HIBISCUS:—
caerulescens, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 516.—Madag.
caesius, Garcke, in Bot. Zeit. vii. (1849) 850; et Peters, Reise Mossamb. Bot. 125 = physaliodes.
californicus, *Kellogg, in Proc. Calif. Acad.* iv. (1873) 292.—Calif.
callosus, Blanco, Bijdr. 67 = Thespesia Lampas.
calycinus, *Willd. Sp. Pl.* iii. 817.—Ins. Borbon.
calycosus, A. Rich. Tent. Fl. Abyss. i. 62 = calycinus.
calycularis, E. Mey. ex Harv. & Sond. Fl. Cap. i. 171 = Ludwigii.
calyphyllus, Cav. Diss. 253 = calycinus.
Cameroni, *Knowles & Westc. Fl. Cub.* iii. (1838) 149. —Madag.
campylosiphon, Turcz. in Bull. Soc. Nat. Mosc. xxxi (1858) i. 193 = Thespesia campylosiphon.
canarensis, Miq. ex Mast. in Hook. f. Fl. Brit. Ind. i. 341 = tetraphyllus.
cancellatus, Linn. f. Suppl. 311 = Pavonia cancellata.
cancellatus, *Roxb. Hort. Beng.* 51; *Fl. Ind.* iii. 201.— Reg. Himal.
canescens, *Heyne, in Wall. Cat.* n. 2698.—Ind. or.
cannabinus, *Linn. Syst.* ed. X. 1149.—Geront. trop.
cardiophyllus, *A. Gray, Pl. Wright.* i. 23.—Texas.
cardiophyllus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 509.—Madag.
carolinianus, *Michl. Cat.* 65.—Am. bor.
Cavanillesianus, *H. B. & K. Nov. Gen. et Sp.* v. 289.— Am. austr.
chinensis, DC. Prod. i. 455 = syriacus.
chinensis, Mast. in Hook. f. Fl. Brit. Ind. i. 342, in syn. sphalm. pro *Bamia chinensis* = seq.
chinensis, Roxb. Hort. Beng. 51; Fl. Ind. iii. 212 = H. Abelmoschus?
Chittle-Benda, Roxb. ex Wight & Arn. Prod. 47 = Pavonia odorata.
ciliaris, Presl, Rel. Haenk. ii. 133 = H. Abelmoschus *citiatus*, Boj. ex Baill. in Bull. Soc. Linn. Par. i. (1885) 541 = Kosteletzkya velutina.
circumatus, Willd. Enum. Hort. Berol. 735; DC. Prod. i. 454 = illiacus.
cisplatinus, *St. Hil. Fl. Bras. Mer.* i. 250.—Bras.
clandestinus, Cav. Ic. i. 1. t. 2 = micranthus.
clypeatus, *Linn. Syst.* ed. X. 1149.—Jamaic.
clypeatus, Walt. Fl. Carol. 177 = Kosteletzkya virginica.
Coatesii, *F. Muell. Fragm.* iii. 5.—Austral. ?
coccineus, *Walt. Fl. Carol.* i. 177.—Am. bor.
coelatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 17 (Quid ?).—Java.
Colleri, *Hort. ex Floral Mag.* N. S. (1876) t. 214.— Polynesia.
Collinianus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 237 = H. Trionum, Manihot, esculentus.
collinus, *Roxb. Hort. Beng.* 51; *Fl. Ind.* iii. 198.— Ind. or.
collinus, Voigt, in Syll. Ratisb. ii. (1828) 54 = præc. ?
columbinus, [Moc. & Sesse,] ex DC. Prod. i. 452 = spiralis.
columnaris, *Cav. Diss.* iii. 166. t. 59. f. 2.—Ins. Mascar.
commersonis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 515.—Ins. Comor.
congener, Schum. & Thonn. Beskr. Guin. Pl. 319 = cannabinus.
convolvuliflorus, *Baill. in Bull. Soc. Linn. Par.* i (1885) 516.—Madag.
Cooperi, Hort. ex Fl. des Serres, Sér. II. v. (1862–63) 109. t. 1656 = H. Rosa-sinensis.
cordatus, D. Dietr. Syn. Pl. iv. 689 = Kosteletzkya cordata.
cordatus, *Harv. in Harv. & Sond. Fl. Cap.* i. 172.— Afr. austr.
cordatus, Hochst. ex Webb, Fragm. Fl. Aethiop. 45 = rhabdotosperma.
cordifolius, Linn. f. Suppl. 309 = Pavonia Mutisii.
cordifolius, *Mill. Gard. Dict.* ed. VIII. n. 18.— Cuba.
cordobensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 193.—Afr. trop.
coriaceus, Heyne, ex Wight & Arn. Prod. 53 = Decaschistia crotonifolia.
corensopifolius, Miq. in Lehm. Pl. Preiss. i. 289 = Fugosia hakeaefolia.
corylifolius, *Presl, Rel. Haenk.* ii. 133.—Ecuador.
corymbosus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 57 —Abyssin.
costatus, *A. Rich. Ess. Fl. Cub.* 138.—Cuba.

HIBISCUS:—
Coulteri, *Harv. ex A. Gray, Pl. Wright.* i. 23.—Mexic.
crassinervius, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 61.—Afr. trop.
crenatus, *Sibth. ex Da Vriese, in Nederl. Kruidk. Arch.* i. (1846) 396.—Guiana.
crenatus, *Vell. Fl. Flum.* 282; vii. t. 82.—Bras.
crinitus, *Wall. ex G. Don, Gen. Syst.* i. 480.—Ind. or.
crotonifolius, Wall. Cat. n. 1901 = Decaschista crotonifolia.
cruentatus, *Schrank, ex Steud. Nom.* ed. II. i. 758.—Hab.!
cruentus, *Bertol. Fl. Guatimal.* 28. t. 10.—Guatemala.
cryptocarpos, A. Rich. Ess. Fl. Cub. 149 = H. Abelmoschus.
cubensis, A. Rich. l. c. 140 = H. Abelmoschus.
cucurbitaceus, St. Hil. Fl. Bras. Mer. i. 244 = H. Abelmoschus.
cucurbitinus, Burch. Trav. i. 278 = urens.
cuneifolius, Garcke, in Bot. Zeit. vii. (1849) 852 = H. Trionum.
cuneiformis, Hort. ex Poir. Encyc. v. 108 = Pavonia praemorsa.
cuneiformis, DC. Prod. i. 454 = Fugosia cuneiformis.
cuspidatus, Edgew. in Journ. As. Soc. Beng. xxi. (1858) 166 = vitifolius.
cyanopyrus, [*Mor. & Sesse,*] *ex DC. Prod.* i. 455.—Mexic.
cymosa, *Walt. Fl. Carol.* 179.—Am. bor.
1) Albertisii, *F. Muell. Papuan. Pl.* i. 56.—N. Guin.
decandrus, Medic. ex Steud. Nom. ed. II. i. 758 = Pavonia praemorsa.
decipiens, *St. Hil. Fl. Bras. Mer.* i. 247.—Bras.
Deniseni, *Hort. ex Flor. Mag.* N. S. (1876) t. 382.—Hab.!
denudatus, *Benth. Bot. Voy. Sulph.* 7. t. 3.—Am. bor. occ.
dictyocarpus, *Webb, Fragm. Fl. Aethiop.* 46.—Afr. trop.
digitatus, *Cav. Diss.* iii. 151. t. 70. f. 3.—Bras.
digitatus, Poir. Encyc. Suppl. iii. 320 = digitiformis.
digitiformis, *DC. Prod.* i. 454.—Hab.!
dimidiatus, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 61.—Bras.
Dioön, *DC. Prod.* i. 449.—Guian.
dissectus, Wall. Cat. n. 2696 = H. Trionum.
divaricatus, *R. Grah. in Edinb. N. Phil. Journ.* (July-Sept. 1840) 367.—Austral.
diversifolius, *Jacq. Ic. Pl. Rar.* t. 551; *Coll.* ii. 807.—Geront. trop.
domingensis, *Jacq. l. c.* tii. 850; *Coll.* iii. 218.—Ins. S. Doming.
dongolensis, Caill. ex Delile, Voy. Meroë Bot. 39 = lunariifolius.
Donii, Walp. Rep. i. 804 = Melochia corchorifolia.
Drummondii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 192.—Austral.
Drummondii, Voang. Fl. Texas, 186; ex S. Wats. Bibl. Ind. N. Am. Bot. 139 = Malvaviscus Drummondii.
ebracteatus, *Mast. in Oliver, Fl. Trop. Afr.* i. 206.—Afr. trop.
elatus, DC. Prod. i. 454 = tiliaceus.
elatus, *Sw. Fl. Ind. Occ.* ii. 1218.—Ind. occ.
Elliottiae, *Harv. in Harv. & Sond. Fl. Cap.* ii. 587.—Afr. austr.
Elliottii, F. Muell. Fragm. i. 220 = Drummondii.
elliptieus, E. Mey. in Drège, Zwei Pfl. Docum. 192 = aethiopicus.
Ellisii, *Baker, in Journ. Linn. Soc.* xx. (1883) 100.—Madag.
Elsworthii, *F. Muell. Fragm.* viii. 241.—Austral.
Endlicheri, Walp. Rep. i. 804 = Melochia corchorifolia.
erianthus, R. Br. in Salt, Abyss. App. 65 = crasinervis.
eriocarpus, DC. Prod. i. 452 = collinus.
eriopermus, Hochst. ex A. Br. in Flora, xxiv. (1841) i. 279 = crasinervis.
esculentus, *Linn. Sp. Pl.* 696.—Cosmop. trop.
Farragei, *F. Muell. Fragm.* viii. 241.—Austral.
fasciculatus, [*Mor. & Sesse,*] *ex DC. Prod.* i. 454.—Mexic.
ferox, *Hoch. Bot. Mag.* t. 4401.—N. Granat.
ferruginens, Boj. ex Baill. in Bull. Soc. Linn. Par. i. (1885) 811 = Bojerianus.
ferrugineus, *Cav. Diss.* iii. 162. t. 60. f. 1.—Madag.
ferruginosus, Oliver, Fl. Trop. Afr. i. Index, 471 = ferrugineus.

HIBISCUS:—
fastatis, Salisb. Prod. 383 = H. Rosa-sinensis.
fecerius, E. Mey. ex Harv. & Sond. Fl. Cap. i. 170 = calycinus.
fieifolius, Mill. Gard. Dict. ed. VIII. n. 15 = esculentus.
ficuloides, *Lindl. in Bot. Reg.* t. 988.—As. et Austral. trop.
ficulneus, Cav. Diss. iii. t. 51. f. 2 = diversifolius.
ficulneus, *Linn. Sp. Pl.* 696.—As. et Austral. trop.
Fitzgeraldi, *F. Muell. Fragm.* viii. 242.—Austral.
flabellatus, Desf. Tabl. ed. I. 147; Poir. Encyc. Suppl. iii. 320 = heterophyllus, *Vent.*
flagellaria, Walp. Rep. i. 805 = acq.
flagelliformis, *St. Hil. Fl. Bras. Mer.* i. 248.—Bras.
flavescens, Cav. Diss. iii. 164. t. 70. f. 3 = H. Abelmoschus.
flavus, Forsk. Fl. Aegypt. Arab. 126 = Pavonia arabica.
floccosus, *Mast. in Hook. f. Fl. Brit. Ind.* i. 348.—Malacca.
floridanus, Shuttlew. ex A. Gray, Pl. Wright. i. 23 = Bancroftanus.
floridus, Salisb. Prod. 383 = syriacus.
fluminensis, *Vell. Fl. Flum.* 283; vii. t. 84.—Bras.
fragilis, DC. Prod. i. 446 = liliiflorus.
fragrans, *Roxb. Hort. Beng.* [97]; *Fl. Ind.* iii. 195.—Reg. Himal.
fraterus, *Linn. Pl. Surin.* n. 90; *Amoen. Acad.* viii. 260.—Guian.
fugax, *Mart. ex Spreng. Syst.* iv. Cur. Post. 258.—Bras.
fulgens, Hort. ex W. Baxt. in Lond. Hort. Brit. Suppl. iii. 592 = H. Rosa-sinensis.
fulvus, *Nutt. in Torkel, Fl. Banat. Rar. Ind.*—Bannat.
furcatus, Wall. Cat. n. 1986 C = surattensis.
furcatus, *Willd. Enum. Hort. Berol.* 786; *Roxb. Hort. Ind.* 21; *Fl. Ind.* iii. 204.—Ceylon, trop.
furcellatus, *Lam. Encyc.* iii. 358.—Am. trop.
fuscus, Garcke, in Bot. Zeit. vii. (1849) 854 = gossypinus, *Thunb.*
fusiformis, Steud. Nom. ed. II. i. 759 = cancellatus.
gangeticus, Roxb. ex Wight & Arn. Prod. 49 = Thespesia Lampas.
gossypinus, *Willd. Enum. Hort. Berol. Suppl.* 49, nomen.—Ind. or.
Gouanii, Boj. Hort. Maurit. 28; et ex Hook. Bot. Mag. t. 3. = liliiflorus.
graminifolius, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 195 = Hasgelli.
gumniolen, *A. Cunn. ex Benth. Fl. Austral.* i. 215.—Australia.
Gibsoni, *Stocks, ex Harv. & Sond. Fl. Cap.* ii. 587; *et ex Mast. in Hook. f. Fl. Brit. Ind.* i. 339.—Geront. trop.
glabrifolius, *A. St. Hil. & Naud. in Ann. Sc. Nat. Sér.* II. xviii. (1842) 40.—Bras.
Goldsworthii, *F. Muell. Fragm.* xi. 90.—Austral.
gossypifolius, Mill. Gard. Dict. ed. VIII. n. 10 = H. Sabdariffa.
gossypinus, *Thunb. Prod. Pl. Cap.* 118.—Afr. austr.
gossypinus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 515.—Madag.
gossypinus, Eckl. & Zeyh. Enum. 38 = pusillus.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 515.—Madag.
grandiflorus, *Micke. Fl. Ber. Am.* ii. 46.—Am. bor.
grandiflorus, Salisb. Parad. Lond. t. 22 = heterophyllus, *Vent.*
grandifolius, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 61 = calycinus.
Greatheadi, *Mast. in Oliver, Fl. Trop. Afr.* i. 208.—Afr. trop.
Greveanus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 515.—Madag.
grewiaefolius, *Hassk. in Hoev. & De Vriese, Tijdschr.* xii. (1855) 53.—Malaya.
grewiaefolius, F. Villar, in Blanco, Fl. Philipp. ed. III. iv. 24 = Thespesia campylocalyon.
grossulariaefolius, Mlq. in Lehm. Pl. Preiss. i. 340 = Hasgelli.
guianensis, *Aubl. Pl. Gui.* ii. 708.—Guiana.
guineensis, DC. Prod. i. 454 = tiliaceus.
guineensis, G. Don, Gen. Syst. ii. 481 = lunariifolius.
habeseffinus, Giordano, Mem. Nov-Ibla, 6 = Fugosia hakeaefolia.
Hamabo, *Sieb. & Zucc. Fl. Jap.* i. 176.—Japon.

HIBISCUS:—
hamatus, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 177 = furcatus.
Harrisii, *Bosse, Handb. Blum.* ix. 344; *ex Garcke, in Bot. Zeit.* viii. (1850) 669.—Hab.!
hastaefolius, E. Mey. ex Harv. & Sond. Fl. Cap. i. 176 = H. Trionum.
hastatus, Cav. Diss. iii. 144. t. 50. f. 1 = Kosteletzkya hastata, *Garcke.*
hastatus, Linn. f. Suppl. 301 = tricuspis.
hastatus, Michx. ex DC. Prod. i. 451 = militaris.
Haynaldi, *F. Muell. in Wing, South Sx. Record*, iii. (March 1883).—Austral.
hebeatus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 174 = aethiopicus.
heptaphyllus, Dals. & Gibs. Bomb. Fl. 20 = radiatus.
herbaceus, *Vell. Fl. Flum.* 284; vii. t. 89.—Bras.
heterophyllus, Buch.-Ham. ex Wall. Cat. n. 2691 = varifolius.
heterophyllus, *Griff. Notul.* iv. 520.—Burma.
heterophyllus, *Vent. Yard. Malm.* 103. t. 103.—Austral.
heterotrichus, *DC. Prod.* i. 450.—Hab.!
heterotrichus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 173 = physaloides.
hirtus, Buch.-Ham. ex Roxb. Hort. Beng. 51; Fl. Ind. iii. 208 = rugosus.
hirtus, *Linn. Sp. Pl.* 694.—Ind. or.; Malaya.
hispidissimus, Griff. Notul. iv. 521 = furcatus.
hispidulus, *Spreng. Pugill.* ii. 78.—Hab.!
hispidus, D. Dietr. Syn. Pl. iv. 529 = Kosteletzkya hispida, *Presl.*
hispidus, Mill. Gard. Dict. ed. VIII. n. 21 = H. Trionum.
hispidus, Spreng. Syst. iii. 99 = Kosteletzkya hispidula.
Hornei, *Baker, Fl. Maurit.* 23.—Ins. Seychelles.
hostilis, *Walt. ex Mast. in Hook. f. Fl. Brit. Ind.* i. 342.—Burma.
Hasgelii, *Endl. Enum. Pl. Hueg.* 10.—Austral.
Humblotii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 517.—Madag.
Humboldtii, Hort. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 52 = hispidus!
Humboldtii, Schrank, ex Colla, Hort. Ripul. 349, 350 = H. Trionum!
hybridus, *F. G. Dietr. Vollst. Lexik. Gaertn.* ii. Nachtr. iv. 433.—Hab.!
hypoglaucus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 177 = surattensis.
immutabilis, *Dehn. Memor. Sopr. Alc. Piant.* 4; *ex Walp. Rep.* i. 807.—Hab.!
incanus, *Wendl. Bot. Beobacht.* 54; *Schrad. Sert. Hannov.* t. 24; *Willd. Sp. Pl.* iii. 807.—Am. bor.
ingratus, *Mlq. in Linnaea*, xix. (1847) 148.—Guiana.
insularis, *Endl. Prod. Fl. Norfolk.* 32.—Ins. Norfolc.
intermedius, *Bilanger, Voy. Ind. Or. Ic.* (absque num.) —Ind. or.
intermedius, Hochst. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 8 = micranthus.
intermedius, *A. Rich. Tent. Fl. Abyss.* i. 58.—Ind. or.; Afr. trop.
involucratus, Salisb. Prod. 384 = surattensis.
Jacquini, *Colla, Hort. Ripul. App.* iv. 24. t. 19 (*in Mem. Acc. Torin.* xxxv. (1831) 176).—Hab.!
japonicus, *Mlq. Ann. Mus. Bot. Lugd. Bat.* 19.—Japon.
jatrophaefolius, A. Rich. Tent. Fl. Abyss. i. 58 = vitifolius.
javanicus, Mill. Gard. Dict. ed. VIII. n. 7 = H. Rosa-sinensis.
javanicus, *Weinm. in Syll. Ratisb.* ii. (1828) 172.—Java.
Jerrodianus, *Past. Flor. Mag.* xiii. (1847) t. 1.—Brus.
kaiguhirensis, *Hofmgg. Verz. Pfl. Nachtr.* ii. 133.—Ind. or.
Kirkii, *Mast. in Oliver, Fl. Trop. Afr.* i. 199.—Afr. trop.
kitaibelifolius, *St. Hil. Fl. Bras. Mer.* i. 248.—Bras.
Kokia, *Hillob. ex Wawra, in Flora*, lvi. (1873) 174.—Ins. Sandvic.
Krausianus, Buch. ex Hochst. l. c. xxvii. (1844) 993 = pedunculatus.
Krichauffianus, *F. Muell. Rep. Bobb. Exped.* 7,—Austral.
lacniatus, Turg. ex Colla, Hort. Ripul. 349, 350 = roseus?

HIBISCUS:—
lasiol, Scop. Delic. Insub. iii. 36 = militaris.
lasseneideles, Edgew. in Journ. Linn. Soc. vi. (1862) 139 = punctatus.
Lambertianus, *H. B. & K. Nov. Gen. et Sp.* v. 291. t. 478.—Am. austr.
lampadatus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = seq.
Lampas, Cav. Diss. iii. 154. t. 56. f. 2 = Thespesia Lampas.
lanceolatus, *Gaudich. in Freyc. Voy. Bot.* 476.—Austral.
lasiocarpos, *Cav. Diss.* iii. 159. t. 70. f. 1.—Hab. ?
lasiococcus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 511.—Madag.
laziospermus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 175 = atromarginatus.
laurinus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 512. —Madag.
laveraroides, *Meric. Pl. Nouv. Am.* 23. t. 16.—Mexic.
laxiflorus, St. Hil. Fl. Bras. Mer. i. 245 = H. Abelmoschus.
leiospermus, *Harv. in Harv. & Sond. Fl. Cap.* i. 173. —Afr. austr.
lepidospermus, *Mast. in Oliver, Fl. Trop. Afr.* i. 197. —Afr. trop.
lepidospermus, *Miq. Fl. Ind. Bat.* i. ii. 159.—Java.
leptocalyx, Sond. in Linnæa, xxiii. (1850) 17 = Pavonia odorata.
leptocladus, *Benth. Fl. Austral*, i. 214.—Austral.
litacinus, Lindl. Bot. Reg. t. 2009 = Fugosia hakiæfolia.
lilliiflorus, *Cav. Diss.* iii. 154. t. 57. f. 1.—Ins. Mascar.
litiiflorus, Griff. ex Mast. in Hook. f. Fl. Brit. Ind. i. 344 = H. Rosa-sinensis.
Lindleyi, Wall. Pl. As. Rar. i. 4. t. 4 (*Lindleï*) = radiatus, linearis, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 40.—Bras.
longifolius, Roxb. Hort. Beng. 58; Fl. Ind. iii. 210 = esculentus.
longifolius, Willd. Sp. Pl. iii. 827 = H. Abelmoschus. Ludwigii, *Eckl. & Zeyh. Enum.* 39.—Afr. austr.
lunariifolius, *Willd. Sp. Pl.* iii. 811.—As. et Afr. trop.
Macleyanus, Walp. Rep. i. 306, sphalm. = seq.
MacLeayanus, Bancroft, ex Macfad. Fl. Jamaic. i. 71 = Bancroftianus.
macranthus, Hochst. in Flora, xxiv. (1841) t. Intell. 29 = lunariifolius.
macrocalyx, Garcke, in Bot. Zeit. vii. (1849) 888 = atromarginatus.
macrogonus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 510.—Madag.
macrophyllus, *Roxb. Hort. Beng.* 51; *ex Hornem. Hort. Hafn. Suppl.* 149.—Ind. or.; Burma; Malaya.
maculatis, E. Mey. ex Harv. & Sond. Fl. Cap. i. 171 = diversifolius.
maculatus, Bartl. Ind. Sem. Hort. Gotting. (1837) = Bartlingianus.
maculatus, *Lam. Encyc.* iii. 349.—Ins. S. Doming.
magnificus, F. Muell. Fragm. ii. 112 = divaricatus.
malacophyllus, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 503.—Ins. Socotr.
malacospermus, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 174.—Afr. austr.
malvoreus, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Act. iv. 17 = mutabilis.
Malvaviscus, Linn. Sp. Pl. 694 = Malvaviscus arboreus.
Manihot, *Linn. l. c.* 696.—Geront. trop.
Marchallianus, Hort. ex Flsch. & Mey. Ind. Sem. Hort. Petrop. ix. 76 = hispidus.
Margravius, A. Cunn. ex Benth. Fl. Austral. i. 213 = heterophyllus, *Vent.*
marmoratus, *Lem. Illustr. Hortic.* (1856) t. 82.—Mexic.
Martianus, *Zucc. in Linnæa*, xxiv. (1851) 198.—Mexic.
mauritianus, Spreng. Syst. iv. Cur. Post. 258 = fimlasiodes.
Mechowii, *O. Hoffm. in Linnæa*, xliii. (1880–82) 131.—Afr. trop.
Meisneri, Miq. in Lehm. Pl. Preiss. i. 240 = Huegelii.
membranaceus, *Cav. Diss.* iii. 154. t. 56. f. 2.—Mexic.
Meyeri, *Harv. in Harv. & Sond. Fl. Cap.* i. 173.—Afr. austr.

HIBISCUS:—
micans, *Cav. Diss.* iii. 167. t. 60. f. 2.—Java.
micranthus, *Linn. f. Suppl.* 308, 310.—Ind. or.; Afr. trop.
microcarpus, Garcke, in Bot. Zeit. vii. (1849) 836 = malacospermus.
microchlamys, *F. Muell. Fragm.* ii. 116.—Austral.
microphyllus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 178 = Meyeri.
microphyllus, Vahl, Symb. Bot. i. 50 = Pavonia arabica.
microphyton, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 511.—Madag.
militaris, *Cav. Diss.* vi. 352 t. 198. f. 2.—Am. bor.
modaticus, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 56 = vitifolius.
mollis, Ziz, ex Span. in Linnæa, xv. (1841) 169 = pandurifformis.
morifolius, Rottl. ex Wight & Arn. Prod. 47 = Pavonia odorata.
moschatus, Roxb. ex Wight & Arn. l. c. 58 = rugosus.
moschatus, Salisb. Prod. 384 = H. Abelmoschus.
Moschutos, *Linn. Sp. Pl.* 693.—Am. bor.
Muhamedis, *Webb, Fragm. Fl. Aethiop.* 46.—Afr. trop.
multifidus, Pæpp. Flow. Gard. vii. (1840) 108 = Fugosia hakeæfolia.
multiformis, St. Hil. *Fl. Bras. Mer.* i. 246.—Bras.
multiformis, Steud. Nom. ed. II. i. 759 = H. Abelmoschus.
multistipulatus, Garcke, in Bot. Zeit. vii. (1849) 849 = pandurifformis.
mutabilis, *Linn. Sp. Pl.* 694.—China.
natalitus, *Harv. in Harv. & Sond. Fl. Cap.* ii. 587.—Afr. austr.
neglectus, *Wright, in Sauv. Fl. Cub.* 13.—Cuba.
Normani, *F. Muell. Fragm.* iii. 4.—Austral.
Notho-Manihot, *F. Muell. l. c.* v. 57.—Austral.
obscurus, A. Rich. Tent. Fl. Abyss. i. 57 = vitifolius.
obtusatus, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 221 = cannabinus.
obtusifolius, Willd. Sp. Pl. iii. 829 = vitifolius.
odontiflorus, *Garcke, in Bot. Zeit.* vii. (1849) 837.—Afr. trop.; Beluchist.
ochroleucus, *Baker, in Journ. Bot.* xx. (1882) 46.—Madag.
odoratus, Roxb. ex Wight & Arn. Prod. 47 = Pavonia odorata.
oenotheroides, Webb, Fragm. Fl. Aethiop. 47 = articulatus.
orbicularis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 518.—Madag.
ovalifolius, Vahl, Symb. Bot. i. 50 = micranthus.
ovatus, Cav. Diss. iii. 148. t. 50. f. 3 = acthiopicus.
ovarionsis, Beauv. Fl. Ow. ii. 86. t. 117 = calycinus.
oxalidiformis, *Boj. Hort. Maurit.* 28, nomen; *et ex Baker, in Journ. Linn. Soc.* xx. (1883) 99.—Ins. Maurit.
oxyphyllus, [*Moç. & Sesse,*] *ex DC. Prod.* i. 455.—Mexic.
pallidus, Rafin. ex S. Wats. Ind. N. Am. Bot. 135 = H. Trionum.
palmatiflorus, *Baker, in Journ. Linn. Soc.* xxi. (1884) 234.—Madag.
palmatilobus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 509.—Madag.
palmatus, Cav. Diss. 168 = H. Manihot.
palmatus, *Forsk. Fl. Aegypt. Arab.* 126.—Arab.
palmatus, *Willd. Enum. Hort. Berol. Suppl.* 50, nomen.—Ind. or.
palustris, Linn. Sp. Pl. 693 = H. Moscheutos.
palustris, Thore, Chlor. Land. 296 = roseus.
pandurifformis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 517.—Madag.
pandurseformis, *Burm. f. Fl. Ind.* 151. t. 47. f. 2.—Ceylon, trop.
paniculatus, Domb. ex Poir. Encyc. v. 105 = Pavonia paniculata.
papilionaceus, Banks, ex Poir. l. c. 109 = Pavonia papilionacea.
papyriferus, Salisb. Prod. 384 = H. Manihot.
Parkeri, *Baker, in Journ. Bot* xx. (1882) 46.—Madag.
parviflorus, R. Br. in Salt, Abyss. App. 65 = abyssinicus.
parviflorus, *Weinm. in Flora*, iii. (1820) 610.—' Am.'

HIBISCUS:—
parvifolius, Hochst. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 6 = micranthus.
Patersonii, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 224 = Lagunaria Patersonii.
Patersonius, Andr. Bot. Rep. t. 286 = Lagunaria Patersonii.
pavoniformis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 515.—Madag.
pedunculatus, *Linn. f. Suppl.* 309; *Thunb. Prod. Pl. Cap.* 118; *Cav. Diss.* iii. 163. t. 66. f. 2.—Afr. austr.
pentacarpos, Linn. Sp. Pl. 697 = Kosteletzkya pentacarpa.
pentaphyllus, F. Muell. Fragm. ii. 13 = Gibsoni.
pentaphyllus, Roxb. Hort. Beng. 52; Fl. Ind. iii. 212 = H. Manihot.
pentaspermus, Bert. ex DC. Prod. i. 447 = Kosteletzkya pentasperma.
pentaspermus, Nutt. in Am. Journ. Sc. v. (1822) 208 = Kosteletzkya virginica.
pernambucensis, *Arruda, in Koster, Trav. Bras.* 487.—Bras.
pernambucensis, *Bertol. in Opusc. Scient.* iv. (1823) 229 (= præc. ?).—Bras.
Perrottetii, Steud. Nom. ed. II. i. 759 = quinquelobus.
persicifolius, *Eckl. & Zeyh. Enum.* 38.—Afr. austr.
petiolifolius, Stokes, Bot. Mat. Med. iii. 548 = H. Moscheutos.
petiolosus, *Miq. Fl. Ind. Bat.* i. ii. 159.—Java.
phoenicous, *Linn. f. Suppl.* 310.—Ind. or.
phoenicus, Willd. Sp. Pl. iii. 818 = hirtus.
phyllochlanus, *F. Muell. Fragm.* ix. 128.—Austral.
physalodes, *Guill. & Perr. Fl. Senag. Tent.* i. 82.—Afr. trop.
physodes, E. Mey. ex Harv. & Sond. Fl. Cap. i. 176 = H. Trionum.
pilosus, Roxb. ex Wall. Cat. n. 3902 = pandurseformis.
Pinonianus, *Gaudich. in Freyc. Voy. Bot.* 476. t. 100. —Austral.
Pinonianus, Miq. in Lehm. Pl. Preiss. i. 240 = Huegelii.
platanifolius, Sweet, Hort. Brit. ed. I. 51 = collinus.
platycalyx, *Mast. in Oliver, Fl. Trop. Afr.* i. 202.—Afr. trop.
platycstegius, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 194.—Java.
Poeppigii, *Garcke, in Eichl. Fairb.* i. (1881) 228, in syn.—Cuba.
Poeppigii, Garcke, in Jahresb. Nat. Ver. Halle (1850), 128 = Bancroftianus.
polygama, *Walt. Fl. Carol.* 179.—Carolin.
polymorphus, *Boj. in Desf. 6me Rapp. Soc. Hist. Nat.* 35.—Ins. Maurit.
ponticus, *Rupr. Fl. Cauc.* 251, in obs.—Reg. Caucas.
populifolius, Salisb. Prod. 383 = Thespesia populnea.
populneoides, Roxb. Hort. Beng. 51; Fl. Ind. iii. 191 = Thespesia populnea.
populneus, Linn. Sp. Pl. 694 = Thespesia populnea.
porophyllus, *Vell. Fl. Flum.* 250; vii. t. 28.—Bras.
praecox, Forsk. Fl. Aegypt. Arab. 125 = esculentus.
praemorsus, Linn. f. Suppl. 309 = Pavonia præmorsa.
Prestii, D. Dietr. Syn. Pl. iv. 829 = Kosteletzkya hastata!
procerus, *Rosk. ex Wall. Cat.* n. 3692.—Burma.
propinquus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 174 = æthiopicus.
prostratus, Roxb. Hort. Beng. 53; Fl. Ind. iii. 208 = fieuleus.
pruniifolius, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr* iii. 893.—Gaiac.
pruriens, Jacq. ex Steud. Nom. ed. I. 450 = Jacquinii.
pruriens, Roxb. Hort. Beng. 51; Fl. Ind. iii. 196 = lunariifolius.
Pseud-Abelmoschus, Blume, Bijdr. 70 = H. Abelmoschus.
Pseudo-Manihot, *DC. Prod.* i. 448.—Ins. Borbon.
Pseudo-palmatus, *Span. in Hook. Comp. Bot. Mag.* i. (1836) 344, nomen.—Ins. Timor.
pteropermoides, Wight, ex Wight & Arn. Prod. 58 = Decaschistia crotonifolia.
pulcherrimus, *Spegr. in Anal. Soc. Cient. Argent.* xv. (1883) 99.—Reg. Argent.
pulverulentus, *Griseb. in Goett. Abh.* xxiv. (1879) 49. —Reg. Argent.
pumilus, Roxb. Hort. Beng. [97]; Fl. Ind. iii. 208 = H. Solandra.
punctatus, A. Cunn. ex Benth. Fl. Austral. i. 221 = Fugosia punctata.

HIBISCUS:—
punctatus, *Dals. in Dals. & Gibs. Bomb. Fl.* 20.—Ind. or.
pungens, *Roxb. Hort. Beng.* 52; *Fl. Ind.* iii. 213.—Reg. Himal.
punicus, *Hort, ex Regel, Cat. Pl. Hort. Akashov.* 71.—Hab.?
purpureus, *Forsk. Fl. Aegypt. Arab.* 126.—Arab.
pusillus, Eckl. & Zeyh. Enum. 38 = H. Trionum.
pusillus, *Thunb. Prod. Fl. Cap.* 118.—Afr. austr.
pyramidatus, *Garcke, in Bot. Zeit.* vii. (1849) 861.—Hab.?
quinquelobatus, *Vell. Fl. Flum.* 283; vii. t. 86.—Bras.
quinquelobus, *G. Don, Gen. Syst.* i. 482.—Afr. trop.
racemosus, Herb. Willd. ex Steud. Nom. ed. II. i. 760 = Malvaviscus concinnus.
racemosus, Lindl. in Bot. Reg. t. 917 = lunariifolius.
radiatus, *Cav. Diss.* iii. 160. t. 54. f. 2.—As. et Austral. trop.
radiatus, Sieber, ex Steud. Nom. ed. II. i. 760 = cannabinus.
radiatus, *Willd. ex Roxb. Fl. Ind.* iii. 209.—Ind. or.; Malaya.
Rainerianus, *Vis. Ind. Sem. Hort. Patav.* (1841) 3.—Afr. bor.
ramosus, *D. Dietr. Syn. Pl.* iv. 838.—Afr. trop.
rhabdotosperma, *Garcke, in Bot. Zeit.* vii. (1849) 859.—Afr. trop.
rhodopetalus, *F. Muell. ex Benth. Fl. Austral.* i. 209.—Austral.
rhombifolius, Cav. Diss. iii. 154. t. 69. f. 3 = syriacus.
ribesifolia, Guill. & Perr. Fl. Seneg. Tent. i. 58 = physaloides.
ribesioides, Steud. Nom. ed. II. i. 760 = praec.
ribifolius, *A. Gray, in Proc. Am. Acad.* v. (1861) 354.—Calif.
Richardsonii, Lindl. in Bot. Reg. t. 875 = H. Trionum.
ricinifolius, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 171.—Afr. austr.
ricinifolius, Wall. Cat. n. 1915 = H. Abelmoschus.
ricinoides, Garcke, in Bot. Zeit. vii. (1849) 854 = ricinifolius.
ricinoides, Zipp. ex Span. in Linnaea, xv. (1841) 170 = tuncensis.
rigidus, Linn. f. Suppl. 310 = micranthus.
riparius, Pers. Syn. ii. 254 = militaris.
Rosa-malabarica, Ker-Gawl. in Bot. Reg. t. 337 = hirtus.
Rosa-sinensis, *Linn. Sp. Pl.* 694.—Geront. trop.
roseus, *Thore, in Loisel. Fl. Gall. ed.* I. ii. 434.—Gall.; Hispan.
rosiflorus, Stokes, Bot. Mat. Med. iii. 543 = H. Rosa-sinensis.
Rossii, Knowles & Westc. Flor. Cab. ii. (1838) t. 63 = Cameroni.
rostellatus, *Guill. & Perr. Fl. Seneg. Tent.* i. 55.—Afr. trop.
Roxburghii, Steud. Nom. ed. II. i. 760 = H. Abelmoschus!
rugosus, *Rouk. ex Steud. Nom.* ed. II. i. 760; *Mast. in Hook. f. Fl. Brit. Ind.* i. 342.—Ind. or.
rugosus, Seidel, ex Steud. Nom. ed. II. i. 760 = aculeatus.
Rutenbergii, *Garcke, in Bremen, Abh.* vii. (1881) 199.—Madag.
Sabdariffa, *Linn. Sp. Pl.* 695.—Geront. trop.
sagittatus, D. Dietr. Syn. Pl. iv. 839 = Kosteletzkya sagittata.
sagittifolius, A. Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 46 = H. Abelmoschus.
Sagraeanus, *Mercier, in Ser. Bull. Bot.* i. (1830) 169.—Cuba.
salicifolius, Linn. Pl. Surin. n. 92; Amoen. Acad. viii. 305 = Pavonia Typhalaea.
salviaefolius, *St. Hil. Fl. Bras. Mer.* i. 249.—Bras.
sanguineus, *Franch. in Rivoil, Çemali*, 15.—Afr. trop.
sanguineus, Griff. Notul. iv. 590 = H. Sabdariffa.
scaber, Lam. Encyc. iii. 350 = diversifolius.
scaber, Michx. Fl. Bor. Am. ii. 45 = acuminatus.
scandens, Dals. & Gibs. Bomb. Fl. 20 = intermedius.
scandens, *Roxb. Hort. Beng.* 51; *Fl. Ind.* iii. 200.—Reg. Himal.
schizopetalus, *Hook. f. Bot. Mag.* t. 5524.—Afr. trop.
scindicus, *Stocks, in Hook. Ic. Pl.* t. 809.—Ind. or.
Scotti, *Bolf. f. in Proc. Roy. Soc. Edinb.* xi. (1883) 908.—Ins. Socotr.
senegalensis, Cav. Diss. iii. 160. t. 68. f. 1 = Pavonia zeylanica.
senegalensis, Guill. & Perr. Fl. Seneg. Tent. i. 58 = pauciuriformis.
septemnervosus, Wight, in Wall. Cat. n. 2698 = canescens.
serratus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 175 = pusillus.
serratus, Wall. Cat. n. 1900 = vitifolius.
setifer, *Presl, Rel. Haenk.* ii. 128.—Am. trop.
setosus, Roxb. Hort. Beng. [97]; Fl. Ind. iii. 194 = macrophyllus.
setosus, Wall. Cat. n. 1909 = pauciuriformis.
setulosus, *Fr. Muell. Fragm.* i. 221.—Austral.
sideaeformis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 518.—Madag.
sideoides, Wight & Arn. Prod. 49 = lunariifolius.
similis, Blume, Bijdr. 78 = elatus.
simplex, *D. Dietr. Syn. Pl.* iv. 838.—Afr. trop.
simplex, Linn. Sp. Pl. ed. II. 977 = Sterculia platanifolia.
simplex, Roxb. Hort. Beng. 51 = collinus.
sinensis, Mill. Gard. Dict. ed. VIII. n. 2 = mutabilis.
sinuatus, Cav. Diss. iii. 147 t. 53. f. 2 = flexuosus.
smilacifolius, Shuttlew. ex A. Gray, Pl. Wright. i. 28 = Kosteletzkya virginica.
Solandra, *L'Hérit. Stirp. Nov.* 108. t. 49.—As. et Afr. trop.
solanifolius, F. Muell. Fragm. ii. 116 = Pinonianus.
somalensis, *Franch. in Révoil, Çomali*, 17.—Afr. trop.
Sororia, *Linn. Pl. Surin.* n. 91; *Amoen. Acad.* viii. 260.—Guian.
spathaceus, *Blume, Bijdr.* 72.—Java.
spathaceus, Bach.-Ham. in Wall. Cat. n. 1915 = H. Abelmoschus.
spathulatus, *Garcke, in Bot. Zeit.* vii. (1849) 840.—Am. austr.
spathulatus, *Gaudich. in Freyc. Voy. Bot.* 476.—Austral.
speciosus, [Soland. in] Ait. Hort. Kew. ed. I. ii. 456 = coccineus.
sphaerocarpus, *Bouché, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 402.—Hab.?
spicatus, *Cav. Diss.* iii. 153. t. 59. f. 1.—Hab.?
spinifex, Crantz, Inst. ii. 170 = seq.
spinifex, Linn. Syst. ed. X. 1149 = Pavonia spinifex.
spiralis, *Cav. Ic.* ii. 47. t. 162.—Mexic.
splendens, *Drap. ex R. Grah. in Edinb. N. Phil. Journ.* (Apr.–June, 1830) 175.—Australi.
stenanthus, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1883) 908.—Ins. Socotr.
stenophyllus, *Baker, in Journ. Bot.* xx. (1882) 47.—Madag.
sterculiaefolius, Steud. Nom. ed. II. i. 760 = quinquelobus.
stipularis, Salisb. Prod. 384 = tubulosus.
Storckii, *Seem. in Bonplandia*, ix. (1861) 254.—Ins. Fidji.
striatus, Cav. Diss. iii. 146. t. 54. f. 1 = domingensis.
strictus, Roxb. Hort. Beng. 52; Fl. Ind. iii. 206 = ficulneus.
strigosus, *Lindl. in Bot. Reg.* t. 860.—Am. austr.
strigosus, Schum. & Thonn. Beskr. Guin. Pl. 314 = vitifolius.
Sturtii, *Hook. in Mitch. Journ. Trop. Austral.* 368.—Austral.
suaveusis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 514.—Madag.
Suddarffa, Rottl. in Act. Lit. Univ. Hafn. i. (1778) 297, sphalm. = H. Sabdariffa.
suberbicalatus, Wall. Cat. n. 1906 = micranthus.
subtrilobatus, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 208 = calycinus.
sulphureus, *H. B. & K. Nov. Gen. et Sp.* v. 289.—Venezuela.
sulphureus, *Wall. ex Voigt, Hort. Suburb. Calc.* 120.—Ind. or.
surattensis, *Linn. Sp. Pl.* 696.—Geront. trop.
sylvaticus, *Benth. Pl. Hartw.* 114.—Mexic.
syriacus, *Linn. Sp. Pl.* 695.—Geront. calid.
tamping, *Mart. Pl. Nouv. Am.* 20. t. 14.—Mexic.
Taffairias v. G. Don, in Sweet, Hort. Brit. ed. II. 506; Maund. & Henal. The Botanist, v. t. 312.—Mauric.
ternatus, Cav. Diss. iii. 172. t. 64. f. 3 = H. Trionum.
ternatus, *Mast. in Oliver, Fl. Trop. Afr.* i. 206.—Afr. trop.

HIBISCUS:—
ternifloruus, Garcke, in Bot. Zeit. vii. (1849) 883 = Kosteletzkya adoënsis.
tetralocularis, Roxb. Hort. Beng. [97]; Fl. Ind. iii. 198 = Thespesia Lampas.
tetraphyllus, *Roxb. il. cc.* 52, 211.—Ind. or.
thespesianus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 511.—Madag.
thespesioides, R. Br. ex Benth. Fl. Austral. i. 220 = Pugoaia thespesioides.
tiliaceus, *Linn. Sp. Pl.* 694.—Amphig. trop.
tiliaefolius, Salisb. Prod. 383 = tiliaceus.
timorensis, *DC. Prod.* i. 448.—Ins. Timor.
tomentosus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Ind. occ.
tomentosus, *Stahl, Estud. Fl. Puerto-rico*, Foll. ii. 92 (= praec. ?).—Ins. Porto Rico.
tortuosus, Roxb. Hort. Beng. 51; Fl. Ind. iii. 192 = tiliaceus.
torulosus, Salisb. Prod. 383 = H. Solandra.
tricolor, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 20.—Austral.
tricuspis, *Banks, ex Cav. Diss.* iii. 152. t. 55. f. 2.—Ins. Pacif.
trideactylites, Lindl. in Mitch. Three Exped. Austral. i. 35 = H. Trionum.
tridentatus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 174 = malacospermus.
trilinearis, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 89.—Bras.
trilobatus, *Vell. Fl. Flum.* 281; vii. t. 29.—Bras.
trilobus, *Asbl. Pl. Gui.* ii. 708.—Guian.; Dominic.
trinitarius, Nocrna, in Act. Batav. Gen., v. (1790) ed. I. Art. iv. 17 = surattensis.
trioniceus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 127 = H. Trionum.
trionioides, G. Don, Gen. Syst i. 483 = seq.
Trionum, *Linn. Sp. Pl.* 697.—Geront. trop.
tripartitus, *Forsk. Fl. Aegypt. Arab.* 126.—Arab.
triactus, Bertol. Misc. Bot. xv. 19 = Jatropha urens.
triumfetaefolius, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 312 = calycinus.
truncatus, A. Rich. Ess. Fl. Cub. 144 = Bancroftianus.
truncatus, Roxb. Hort. Beng. 51; Fl. Ind. iii. 200 = vitifolius.
tubiformis, [Moç. & Sesse] ex DC. Prod. i. 447.—Mexic.
tuberculosus, Cav. Diss. iii. 161. t. 68. f. 2 = pandurae-formis.
tulipaefolius, *Hook. Ic. Pl.* tt. 707, 708.—Ind. occ.
tulipifolius, Regel, Cat. Pl. Hort. Akashov. 71 = praec.?
uncinellus, [*Moç. & Sesse*] *ex DC. Prod.* i. 449.—Mexic.
uncinatus, *DC. ex Spreng. Syst. iv. Cur. Post.* 258.—Hab.?
unidens, *Lindl. in Bot. Reg.* 878.—Bras.
uniflorus, E. Mey. ex Harv. & Sond. Fl. Cap. i. 176 = H. Trionum.
umilateralis, Cav. Diss. iii. 158. t. 67. f. 2. c = spiralis.
urens, *Linn. f. Suppl.* 309.—Afr. austr.
urticaefolius, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 89.—Bras.
variabilis, Garcke, in Bot. Zeit. vii. (1849) 841 = physaloides.
variatus, *Splitg. ex De Vriese, in Nederl. Kruidk. Arch.* i. (1846) 388.—Guian.
vastifolius, *Sieud. Nom.* ed. II. i. 760.—Hort. Calcutt.
velutinus, DC. Prod. i. 455 = pauciuriformis.
venustus, Blume, Bijdr. 71 = mutabilis.
verrucosus, Guill. & Perr. Fl. Seneg. Tent. i. 57 = cannabinus.
vesicolor, *Schum. & Thonn. Beskr. Guin. Pl.* 311.—Afr. trop.
vesicarius, Cav. Diss. iii. 171. t. 64. f. 2 = H. Trionum.
vestitus, Griff. Notul. iv. 618 = macrophyllus.
vestitus, Wall. Cat. n. 1924, partim = pungens.
Vidalianus, Naves, in Blanco, Fl. Philipp. ed. III. t. 338; *Nov. App.* 94 = Thespesia campylocalyx.
virgatus, *Blume, Bijdr.* 71.—Malaya.
virginicus, Linn. Sp. Pl. 697 = Kosteletzkya virginica.
virginicus, Walt. Fl. Carol. 177 = militaris.
vitifolius, *Linn. Sp. Pl.* 696.—Geront. trop.
vitifolius, Mill. Gard. Dict. ed. VIII. n. 3 = cannabinus.
Vriesianus, *Hassk. in Hoev. & De Vriese, Tijdschr.* v. (1838–39) 385.—Java.
vulpinus, *Salisb. ex Blume, Cat. Gev. Buitenz.* 38, nomen.—Java.
Wahlii, T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 8 (= *Cienfuegosia Wahlii*).—Aphak.
Wightianus, Wall. Cat. n. 2695 A, 2695 = cannabinus.

HIBISCUS:—
Wrayae, Lindl. Bot. Reg. (1840) Misc. 69 = Huegelii.
Xanthiana, Steud. Nom. ed. II. i. 760 = Urena Xanthium.
Y-unganu, *Gaudich. in Freyc. Voy. Bot.* 91.—Ins. Sandvic.
* *zeylanicus*, Linn. Sp. Pl. 697 = Pavonia zeylanica.
zonatus, *F. Muell. Fragm.* i. 221.—Austral.

HICKESEACHIA, F. Muell. in Wing, South Sc. Record, iii. (1883) 23. *PROTEACEAE.*
pinnatifolia, *F. Muell. l. c.*—Austral.

HICORIA, Rafin. Alsog. Am. 66 (1838) = seq.

HICORIUS, Rafin. Fl. Ludov. 109 (1817) = **Carya**, Nutt. (Jugland.).
alba, Rafin. Alsog. Am. 66 = C. alba.
amara, Rafin. Fl. Ludov. 109 = C. amara.
aquatica, Rafin. Alsog. Am. 66 = C. aquatica.
integrifolia, Rafin. Fl. Ludov. 109 = C. integrifolia.
maxima, Rafin. Alsog. Am. 66, 67 = C. tomentosa ?
microcarpa, Rafin. l. c. 66 = C. microcarpa.
Myristica, Rafin. l. c. = C. myristicaeformis.
olivaeformis, Rafin. Fl. Ludov. 109 = C. olivaeformis.
porcina, Rafin. l. c. = C. porcina.
scabra, Rafin. l. c.—Am. bor.
sulcata, Rafin. l. c. = C. sulcata.
tomentosa, Rafin. l. c. = C. tomentosa.

HIDALGOA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 15 (1824). *COMPOSITAE*, Benth & Hook. f. ii. 386.
Lexingii, DC. Prod. v. 511 = ternata.
ternata, *La Llave, in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 15.—Mexic.

HIDROCOTILE, Neck. Delic. Gallo-Belg. i. 136 (1768) = **Hydrocotyle**, Tourn. (Umbellif.).

HIDROSIA, E. Mey. Comm. Pl. Afr. Austr. 89 (1835) = **Rhynchosia**, Lour. (Legumin.).
bullata, E. Mey. l. c. = R. bullata.

HIERACEUM, Hoppe, Bot. Taschenb. (1791) 278 = **Hieracium**, Tourn. (Compos.).

HIERACHIUM, Hill, Hort. Kew. 43 (1769) = **Hypochoeris**, Linn. (Compos.).
laevigatum, Hill, l. c. = H. Achyrophorus.

HIERACIOIDES, (Vaill.) Moench, Meth. 547 (1794) = **Hieracium**, Tourn. (Compos.).
flexuosa, Moench, l. c. = Hieracium sabaudum.
staticifolia, Moench, Meth. Suppl. 223 = Hieracium staticifolium.
umbellatum, Moench, Meth. 547 = Hieracium umbellatum.

HIERACION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113, 127 = seq.

HIERACIUM, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 516.
APTANTHUS, Vis. Fl. Lib. Spec. 54. t. 7. f. 3 (1824).
ARACIUM, Neck. Elem. i. 49 (1790).
CHLOSCREPIS, Griseb. in Goett. Abh. v. (1853) 155; (Comm. Hierac. Gen. 75).
CHRYSOSPLENIUM, Fries, Epicris. Hierac. 153 (1862).
HETEROPLEURA, Sch. Bip. in Flora, xlv. (1862) 484.
HIERACIODES, [Vaill.] Moench, Meth. 547 (1794).
MANDONIA, Sch. Bip. in Linnaea, xxxiii. (1864–65) 767.
MIRGIA, Neck. Elem. i. 49 (1790).
PILOSELLA, Rupp. Fl. Jen. ed. Hall. 202 (1745); Sch. Bip. in Flora, xlv. (1862) 417, partim.
SCHLAGINTWEITIA, Griseb. in Goett. Abh. v. (1853) 156 (Comm. Hierac. Gen. 76).
SYNOTHECA, Mosn. Ess. Hierac. 71 (1829).
abietinum, Jord. ex Nym. Consp. 443 = vulgatum.
abietiolum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 417 = murorum.
abietinum, Rent. ex Boiss. Fl. Orient. iii. 672 = vulgatum.
abruptifolium, Vukot. in Rad Jugos. Akad. Zagreb. xxxiv. (1876) 16 = sabaudum.
absinthium, *Linn. in Linnaea*, v. (1830) 183.—Mexic.
abscondidum N., Hutet, ex Dalla Torre, Art. Alpenfl. 156.—Europ.
accipitrinum, *Uspensky, in Bull. Soc. Nat. Mosc.* vii. (1834) 378, nomen.—Russia.
accrocomum, *Naeg. & Peter, Hierac.* 709.—Europ.
acrothyrsum, *Naeg. & Peter, l. c.* 714.—Europ.
actinotum, *Naeg. & Peter, l. c.* 477.—Europ.
acuminatum, Jord. Gr. Jard. Gren. (1849) 17 = vulgatum.
acuminatum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 79 = Crepis paludosa.
acutatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 406 = vulgatum.
acutifolium, Hall. f. ex Steud. Nom. ed. II. i. 761 = villosum.
acutifolium, Vill. Précis Voy. Bot. 59. t. 3 = piloselloides, dentatum.
adenophorum, Scheele, in Linnaea, xxxii. (1863) 682 = corinthoides.
adriaticum, *Naeg. ex Freyn, in Verh. Zool.-Bot. Ges. Wien*, xxvii. (1878) 870.—Europ.
adscitum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 410 = murorum.
adulterinum, *Naeg. & Peter, Hierac.* 427.—Europ.
aestivale, *Jord. in Bor. Fl. Cent. Fr.* ed. III. ii. 391.—Europ.
aestivum, Bill. ex Nym. Consp. 441 = tridentatum.
aestivum, *Fries, Symb. Hierac.* 176.—Norveg.
aestivum, Grm. & Godr. Fl. Fr. ii. 387 = umbellatum.
affine, Tausch, in Flora, xi. (1828) 1. Erg. 70 = rigidum.
affine, *Zumag. Fl. Pedem.* i. 374.—Ital.
agerastolens, *Fries, Epicrisis Hierac.* 108.—Ital.
aggregatum, Backh. Mon. Hierac. 52 = argenteum.
agrarium, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875–76) 610 = anglicum.
agrestis, Bernh. Syst. Verz. Erf. 187 = Crepis tectorum.
alatum, *Lapeyr. Hist. Abr. Pl. Pyr.* 478.—Mont. Pyren.
albereanum, *Timb. in Mem. Acad. Toulouse*, Sér. VIII. i. ii. (1879) 194.—Europ.
albidum, Vill. Prosp. 36; Hist. Pl. Dauph. iii. 138 = lintybaceum.
albiflorum, *Hook. Fl. Bor. Am.* i. 298.—Am. bor. occ.
albinum, Fries, Epicris. Hierac. 108 = jurasum.
albipedicellatum, *Naeg. & Peter, Hierac.* 726.—Europ.
albulum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 419 = pallidum.
albacarensis, *Amo, Desc. Pl. Gran.* 2.—Hispan.
Allionii, Hall. f. in Roem. Arch. i. ii. 2 = Crepis alpestris.
Allionii, Mérat, ex Steud. Nom. ed. II. i. 761 = glaucum.
Allionii, Monn. Ess. Hierac. 15 = saxatile.
Allionii, Tausch, in Flora, xi. (1828) i. Erg. 58 = cymosum.
alpestre, Jacq. Fl. Austr. ii. 54 = Crepis alpestris.
alpicolum, *Schleich. Cat. Pl. Helv.* ed. IV. 19, nomen; *et ex Frost. in DC. Prod.* vii. 201.—Helvet.
alpicolum, Schur, Enum. Fl. Transs. 382 = H. Auricula.
alpicolum, Tausch, in Flora, xi. (1828) i. Erg. 55 = sphaerocephalum.
alpinum, All. Fl. Pedem. i. 212 = glandulíferum.
alpinum, Hoppe, in Sturm, Deutschl. Fl. Heft 37 = piliferum, H. Pilosella.
alpinum, *Linn. Sp. Pl.* 800.—Europ.; As. bor.; Greenl.
alpinum, Lucé, Fl. Osil. 362 = Leontodon hastilis.
alpinum, Pall. Reise, iii. 445 = Crepis chrysantha.
alpinum, Vill. ex Fries, Symb. Hierac. 49 = piliferum.
alpinissimum, *Krock. Fl. Silес. Suppl.* ii. 259.—Europ.
altissimum, *Lapeyr. Suppl. Hist. Pl. Pyr.* 135.—Mont. Pyren.
amaurocephalum, *Naeg. & Peter, Hierac.* 355.—Europ.
amaurophyllum, Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, iii. (1875–76) 601 = hirsutum.
ambiguum, Ehrh. Beitr. v. 178, nomen = dubium.
ambiguum, Hort. Landsb. ex Frost in DC. Prod. vii. 223 = politum.
ambiguum, Lapeyr. ex Froel. i. c. 212 = Candollei.
ambiguum, Schleich. ex Pers. Syn. ii. 375 = rigidum.
ambiguum, *Schult. Obs. Bot.* 165.—Hab. ?
ambiguum, Schur, Enum. Pl. Transs. 386 = subfuscum.
ambiguum, Spenn. Fl. Friburg. ii. 665 = piloselloides.
ambliopia, *Boiss. Fl. Orient.* iii. 874.—As. Min.
amphibolum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 392 = umbellatum.
amplexicaule, Bieb. Fl. Taur. Cauc. ii. 254 = cydoniaefolium.
amplexicaule, Georgi, Beschr. Russ. Reich. III. iv. 1213 = sabaudum.
amplexicaule, *Linn. Sp. Pl.* 803.—Europ. austr.
amplifolium, Schleich. Cat. Pl. Helv. ed. IV. 19 = cerinthoides.
analogum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 394 = tridentatum.
ancepa, *Ford. ex Martr. Fl. Tarn*, 447.—Gallia.
anchusaefolium, Bertol. Amoen. Ital. 402 = dentatum.
anchusaefolium, Spreng. Syst. iii. 642 = Liottardi.
andinum, *Phil. in Linnaea*, xxxiii. (1864–65) 125.—Chili.
andryaloides, *Vill. Prosp.* 35; *Hist. Pl. Dauph.* iii. 151. t. 29.—Europ.
anelmenum, *Naeg. & Peter, Hierac.* 687.—Europ.
anfractum, *Fries, in Oefver. Vet. Akad. Foerhandl. Stockh.* (1856) 148.—Scandinav.
anglicum, *Fries, Symb. Hierac.* 93.—Europ.
anglicum, Wimm. Fl. Schles. ed. III. 308 = pallescens.
angulare, Fries, in Oefver. Vet. Akad. Foerhandl. Stockh. (1856) 148 = caesium.
angustatum, Fries, ex Nym. Consp. 440 = crocatum.
angustatum, Lindeb. in Hartm. Handb. Skand. Fl. ed. XI. 44 = caesium.
angustatum, Tausch, in Flora, xi. (1828) i. Erg. 57 = cymosum.
angustifolium, DC. Fl. Fr. vi. 428 = glaciale.
angustifolium, C. C. Gmel. Fl. Bad. iii. 823 = vulgatum.
angustifolium, Hoppe, Bot. Taschenb. (1799) 180 = sphaerocephalum.
angustifolium, Lapeyr. ex Steud. Nom. ed. II. i. 761 = cerinthoides.
angustifolium, Ledeb. ex Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 67 = Ledebourii.
angustifolium, Schleich. ex Steud. Nom. ed. II. i. 761 = prenanthoides.
angustifolium, Schur, Enum. Pl. Transs. 390 = vulgatum.
angustum, Lindeb. in Hartm. Handb. Skand. Fl. ed. XI. 55 = crocatum.
anisocladum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 408 = vulgatum.
anisophyllum, *Boiss. Fl. Orient.* iii. 876.—Bithyn.
anisotrichum, Griseb. in Goett. Abh. v. (1858) 85 = Vahlii.
antarcticum, *Phil. in Anal. Univ. Chil.* (1873) 483.—Reg. Magell.
antarcticum, *Urv. in M ém. Soc. Linn. Par.* iv. (12) 608.—Reg. Magell.
anthrum, *Fries, Symb. Hierac.* 153.—Mexic.
apatelium, *Naeg. & Peter, Hierac.* 706.—Europ.
apiculatum, Tausch, in Flora, xx. (1837) 1. Beibl. 70 = nigrescens.
approximatum, Jord. Cat. Gr. Jard. Dijon (1848) 20 = vulgatum.
arachnoideum, *Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, i. (1872–73) 95.—Europ.
arbuscense, Timb. l. c. ii. (1874) 462 = alatum.
arcticum, Fr. Incr. DC. Prod. vii. 209 = gracile.
arcticum, Schur, in Oestr. Bot. Zeitschr. ix. (1861) 22 = transilvanicum, rotundatum.
arenarium, Sch. Bip. in Pollichia, xiii. (1855) 23 = caesium.
arenicola, *Gremli, Neue Beitr.* iii. 14.—Helvet.
argenteum, *Fries, Symb. Hierac.* 99.—Norveg.
argillaceum, Jord. Gr. Jard. Gren. (1849) 17 = vulgatum.
argutidens, Fries, ex Nym. Consp. 443 = fasciculare.
argutum, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 447.—Calif.
aridum, *Freyn, in Oestr. Bot. Zeitschr.* xxvi. (1876) 369.—Europ.
aristatum, Vim. Samma Fl. iv. 411 = Crepis blattaroides.
armeriaefolium, *Reichb. in Mossi. Handb.* ed. II. ii. 1357.—Helvet.
armericides, *Arv. in Bull. Soc. Dauph.* (1880) 288.—Europ.
armiceides, Grm. & Godr. Fl. Fr. ii. 371 = murorum.
armoricides, *Naeg. & Peter, Hierac.* 597.—Europ.
arragonense, Scheele, in Linnaea, xxxii. (1863) 667 = cerinthoides.

HIERACIUM :—
arrectarium, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 408 = vulgatum.
arrectum, *Gren. & Godr. Fl. Fr.* ii. 379 = cydoniæfolium.
arrectum, *Jord. in Bor. Fl. Cent. Fr.* ed. III. ii. 394. —Gallia.
arrhinophyllum, Vukot. in Rad Jagos. Akad. Zagreb. vii. (1869) 79 = sabaudum.
artefactum × , Naeg. & Peter, Hierac. 678.—Europ.
arvense, Scop. Fl. Carn. ed. II. ii 110 = Sonchus arvensis.
arvenense, *Arv. in Lamatte, Prod.* ii. 488.—Gallia.
Arvetii, *Verlot, Cat. Yard. Grenoble* (1879), 12.—Europ.
arvicola, *Naeg. & Peter, Hierac.* 606.—Europ.
Aschersonianum, Uechtritz, in Oestr. Bot. Zeitschr. xxii. (1872) 78 = olympicum.
asperatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 406 = vulgatum.
aspericaule, Jord. in Bor. l. c. 384, in obs. = tridentatum.
asperifolium, Schur, Enum. Pl. Transs. 383 = brachiatum.
aspernatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 400 = vulgatum.
aspernimum, D. Don, Prod. Fl. Nep. 165 = Lactuca Dalrympa !
aspernimum, Schur, Enum. Pl. Transs. 400 = umbellatum.
asperulum, *Peyrn, in Flora*, lxiv. (1881) 217.—Europ.
asperum, Bernh. Syst. Verz. Erf. 136 = Sonchus asper.
asperum, Bieb. ex Fries, Symb. Hierac. 44 = incanum.
asperum, Fries, l. c. 172 = tridentatum.
asperum, Schleich. ex Reichb. Fl. Germ. Excurs. 267 = ramosum.
asperum, Tausch, in Flora, xi. (1828) 1. Erg. 58 = prænitum.
asplenioides, Besit. Hort. Breit. 192 = Picris asplenioides.
aspreticolum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 418 = murorum.
asyngamicum, *Kern. ex Dalla-Torre, Atl. Alpenfl.* 135.—Europ.
atactum × , Naeg. & Peter, Hierac. 355.—Europ.
athoum, Griseb. ex Pant. in Verh. Vat. Natur. Heilk. Presb. ii. (1871-72) 53 = chalcidicum.
atlanticum, *Fries, Epicris. Hierac.* 53.—Alger.
atratum, Bab. Man. Brit. Bot. ed. III. 196 = nigrescens.
atratum, *Fries, Symb. Hierac.* 106.—Europ.
atricapillum, Hoppe, in Flora, xiv. (1831) 1. 184 = alpinum.
atrovirens, Froel. in DC. Prod. vii. 231 = murorum.
atrovirens, Guss. Fl. Sic. Syn. ii. 408 = Bourgæi.
attenuatum, Tausch, in Flora, xi. (1828) 1. Erg. 56 = dubium.
atticum, Nym. Consp. 453 = sabinum.
aurantellum, *Naeg. & Peter, Hierac.* 347.—Europ.
aurnaticum, *Linn. Sp. Pl.* 801.—Europ.
auratum, Fries, Symb. Hierac. 131 = canadense.
auricomum × , Bor. Fl. Cent. Fr. ed. III. ii. 406 = vulgatum.
aurosum, Hort. Genev. ex Schlecht. in Linnæa, xxv. (1852) 906 = Leontodon hastilis.
aurosum, Kit. ex Nym. Consp. 456 = Crepis Kitaibelii.
aurosum, Scop. Fl. Carn. ed. II. ii. 104 = Crepis aurea.
Auricula, Lees, Fl. Hasborn. ed. II. 174 = prætense.
Auricula, *Linn. Sp. Pl.* 800.—Europ. ; Reg. Casc.
Auricula, Vill. Hist. Pl. Dauph. iii. 100 = *bifurcum*.
Auricula, Willd. Sp. Pl. iii. 1564 = præsatum.
Auricula, Willk. ex Willk. & Lange, Prod. Fl. Hisp. ii. 353 = Vahlii.
auriculæfolium, Gremli, Excurs. Fl. Schw. ed. II. 261 = H. Pilosella.
auriculæfolium, Tausch, ex Koch, Syn. Fl. Germ. ed. I. 445 = *bifurcum*.
auriculæfolium, Willd. ex Froel. in DC. Prod. vii. 166 = Crepis incarnata.
auriculæforme, Fries, Symb. Hierac. 7 = *bifurcum*.
auriculatum, *Hornem. Hort. Hafn.* 768.—Hab. !
auriculatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 197, 218 = H. Auricula.
auriculatum, Willd. ex Spreng. Syst. iii. 644 = boreale.
auriculiforme, Pand. & Vis. in Mem. Ist. Venet. xxi. (1880-82) 584 = Nangelianum.
auriculoides, Lang, in Syll. Ratisb. i. (1824) 183 = præsaltum.
auriparvum, Loret, ex Nym. Consp. 445 = pallidum.
auriparvum, Timb. in Bull. Soc. Bot. Fr. v. (1858) 615 = lactucaceum.

HIERACIUM :—
aurimontanum, Schrank, Baier. Fl. ii. 382 = villosum.
auriotplon, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 393 = tridentatum.
aurosicum, *Serres, in Bull. Soc. Bot. Fr.* iii. (1856) 277.—Gallia.
aurulentum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 399 = vulgatum.
Auszerdorferi, Hausm. ex Nym. Consp. 441 = lapponicum !
australe, *Fries, Symb. Hierac.* 190.—Europ. austr.
austriacum, Hoffm. Deutschl. Fl. ed. II. i. 2. 119 = Crepis blattarioides.
autumnale, Griseb. in Goett. Abh. v. (1853) 183 = sabaudum.
avicolum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 401 = vulgatum.
Avilae, *H. B. & K. Nov. Gen. et Sp.* iv. 3.—Venezuela
badense, *Dicht. in Deutsche Bot. Monatsschr.* ii. (1884) 86.—Europ.
Balansae, *Boiss. Diagn.* Ser. II. v. vi. 119.—As. Min. ; Syria.
balkanum, *Uechtr. ex Panč. Elem. Fl. Bulg.* 46.—Europ.
balsamum, Asso, Syn. Stirp. Arag. 111 = amplexicaule.
banaticum, Heuff. ex Nym. Consp. 453 = brachiatum.
barbatum, Loisel. Fl. Gall. ii. 253 ; et ed. II. ii. 192 = dubium.
barbatum, Nutt. in Journ. Acad. Philad. vii. (1834) 70 = longiflum.
barbatum, *Tausch, in Flora*, xi. (1828) 1. *Erg.* 73.—Europ.
barbicaule, *Celak. in Arch. Naturw. Landesd. Boehm. lv.* (1881) 139.—Europ.
Barthianum, Schœele, in Linnæa, xxxi. (1861-62) 656 = compositum.
basifurcum, *Naeg. & Peter, Hierac.* 251.—Europ.
basiphyllum, Vukot. in Rad Jagos. Akad. Zagreb. vii. (1869) 68 = racemosum.
Bastardianum, Bor. Fl. Cent. Fr. ed. III. ii. 402 = vulgatum.
Bauhini, Bess. Prim. Fl. Gal. ii. 149 ; Schult. Obs. Bot. 164 = præsaltum.
Bauhini, *Schwarz. ex Schrank, in Denkschr. Bot. Ges Regensb.* i. (1822) 66.—Hab. !
Baumgartenianum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853)) 45 = porphyriticum !
bellidiforme, *Tausch. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse*, iii. (1875-76) 615, nomen.—Europ.
bellidifolium, *Froel. in DC. Prod.* vii. 209.—Carinth.
bellidifolium, Scheele, in Linnæa, xxxii. (1863) 666 = compositum.
bellum, Jord. ex Nym. Consp. 448 = vulgatum.
berneusis, Christ. in Mitthel. Bern. Naturf. Ges. (1860) (Hierac. Schw. 19) = villosum.
Besserianum, Host, Fl. Austr. ii. 406 = H. Auricula.
Besserianum, Spreng. Syst. iii. 639 = præsatum.
bicolor, Fries, ex Nym. Consp. 454 = aurantiacum.
bicolor, Scheele, in Linnæa, xxvi. (1861-62) 654 = olivaceum.
bidentatum, Tausch, in Flora, xi. (1828) 1. Erg. 74 = prænanthoides.
bifidum, Blytt, ex Lange, Suppl. Fl. Dan. t. 96 = murorum.
bifidum, Hornem. Hort. Hafn. 767 = Crepis biennis.
bifidum, Kit. ex Hornem. Hort. Hafn. ii. 761 = caninum.
bifidum, Koch, Syn. Fl. Germ. ed. II. 523 = rupicolum.
bifidum, Schleich. ex Fries, Symb. Hierac. 111 = murorum.
bifidum, Ser. ex Froel. in DC. Prod. vii. 230 = lasiophyllum, pallidum.
bifolium, *Arv. in Bull. Soc. Dauph.* (1880) 288.—Europ.
biforme, *Fries, Epicris. Hierac.* 118.—Europ. austr.
bifurcatum, Hort. ex Tausch, in Flora, xi. (1828) 1. Erg. 54 = dubium.
bifurcum × , Bieb. Fl. Taur. Cauc. ii. 251.—Reg. Caucas.
bifurcum, Fries, ex Nym. Consp. 454 = cerinthe.
bifurcum, Froel. in DC. Prod. vii. 200 = stoloniflorum.
bifurcum, Schur, Enum. Pl. Transs. 361 = brachiatum.
bifurcum, Tenore, Syll. Fl. Neap. 400 = H. Pilosella.
bihariense, *Kern. in Oestr. Bot. Zeitschr.* xiii. (1863) 246.—Europ.
bitense, F. Schultz, Fl. Pfals, 276 = *bifurcum*.
bituriginse, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 386 = boreale.
Blanci, Serres, in Bull. Soc. Bot. Fr. ii. (1856) 225 = Pseudo-Cerinthe.

HIERACIUM :—
blattarioides, Linn. Sp. Pl. 804 = Crepis blattarioides.
blattarioideum, Arv. in Bull. Soc. Dauph. (1880) 289 = jurassum.
Blyttianum, *Fries, Cat. Hort. Ups.* (1858) ; *ex ej. Epicris. Hierac.* 21.—Norveg.
Bocconei, *Griseb. in Goett. Abh.* v. (1853) 115.—Helv. ; Tyrol.
bohemicum, *Fries, Epicris. Hierac.* 47.—Silesia.
Boissieri, Huet, ex Nym. Consp. 450 = villosum.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 365.—Calif.
boliciense, Sch. Bip. in Bonplandia, ix. (1861) 174 = Crepis boliviensis.
bombycinum, *Boiss. & Reut. ex Fries, Epicris. Hierac.* 50.—Hispan.
Bordiri, Sch. Bip. ex Nym. Consp. 448 = mixtum.
borrelii, Costa, Fl. Catal. 189 = sabaudum.
boreale, *Fries, Nov. Fl. Suec.* ed. I. 77 ; ed. II. 261.—Europ. ; Orient ; As. bor.
borreoli, Kittel, ex Walp. Ann. ii. 1012 = lycopsifolium.
borreoli, Lacstad. ex Fries, Symb. Hierac. 184 = crocatum.
borreoli, E. Mey. ex Fries, l. c. 186 = canadense.
Borreanum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 392 = tridentatum.
Borreri, *Syme, Engl. Bot.* ed. III. v. 212.—Brit.
bourgeoisianum, Jord. in Bor. Fl. Cent. Fr. ed. III ii. 412 = murorum.
Bourgæi, *Boiss. Diagn.* Ser. II. iii. 102.—Europ.
brachiatum, *Bert. ex DC. Fl. Fr.* vi. 442.—Europ.
brachycomum, *Naeg. & Peter, Hierac.* 255.—Europ.
brachyphyllum, Nym. Consp. 455 = floribundum.
bractestum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 72 = Crepis blattarioides.
bracteolatum, Fries, Epicris. Hierac. 71 = Crepis athoa.
bracteolatum, Link, Enum. Hort. Berol. ii. 268 = umbellatum.
bracteolatum, *Sibth. & Sm. Fl. Græc. Prod.* ii. 135.—Sieli. ; Græc.
bractreum, Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875-76) 604 = scapigerum.
Brandegei, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 194.—Calif.
Braunii, F. Schultz, in Arch. Fl. Fr. et Allem. 18 = prænanthoides.
brevidentatum, Jord. in Bor. Fl. Centr. Fr. ed. III. ii. 397 = vulgatum.
brevifolium, Froel. in DC. Prod. vii. 225 = umbellatum.
brevifolium, Tausch, in Flora, xi. (1828) 1. Erg. 71 = latifolium.
brevipes, Jord. in Bor. Fl. Centr. Fr. ed. III. ii. 412 = murorum.
brevipilum, Greene, in Bull. Torrey Bot. Club, ix. (1882) 54 = Fendleri.
brevicaspum, DC. Fl. Fr. vi. 439 = glaciale.
brevicaspum, Griseb. Spicil. Fl. Rumel. ii. 371 = prætense.
Brewerі, A. Gray, in Proc. Am. Acad. vi. (1864-65) 553 = horridum.
brigantiacum, Jord. ex Nym. Consp. 450 = glaucum.
Brutori, *Frost, in DC. Prod.* vii. 235.—Lusitan.
bructerum, Fries, Symb. Hierac. 73 = alpinum.
Buschii x, Thuem. in Bonplandia, vi. (1858) 157. —Europ.
buglossoides, *Arv. in Bull. Soc. Dauph.* (1880) 286.—Europ.
bulbosum, Willd. Sp. Pl. iii. 1563 = Crepis bulbosa.
bupleurifolium, Lapeyr. ex Steud. Nom. ed. II. i. 762 = cerinthoides.
bupleurifolium, Tausch, in Flora, xi. (1828) 1. Erg. 74 = prænanthoides.
buplevroides, Bellard. ex Willd. Sp. Pl. iii. 1561 = Reuxnum, *Waldst. & Kit.*
buplevroides, *C. C. Gmel. Fl. Bad.* iii. 817.—Europ.
buplevroides, Lapeyr. ex Steud. Nom. ed. I. 406 = dubium.
burnaefolium, Fries, Symb. Hierac. 116 = ramosum.
burnaefolium, Froel. ex Nym. Consp. 443 = fuscinlare.
caesulum, *Arv. in Bull. Soc. Dauph.* (1880) 286.—Europ.
caesarianum × , Naeg. & Peter, Hierac. 763.—Europ.
caesium, *Fries, Symb. Hierac.* 112.—Europ.
campitosum, Dum. Fl. Belg. Prod. 62 = prætense.
calabricum, Bivet, ex Nym. Consp. 447 = Sartorianum.
calabricum, *Naeg. & Peter, Hierac.* 692.—Europ.
calanthes × , Naeg. & Peter, l. c. 350.—Europ.

HIERACIUM :—
calcareum, Bernh. ex Hornem. Hort. Hafn. ii. 762 = saxetanum.
calcareum, Willd. ex Froel. in DC. Prod. vii. 216 = rupestre.
calcigenum, Rehm. in Oestr. Bot. Zeitschr. xxiii. (1873) 213 = vulgatum.
caledon, Stend. Nom. ed. II. i. 763, sphalm. in syn. = dubium.
calenduliflorum, Backh. Mon. Hierac. 23 = eximium.
californicum, Froel. in DC. Prod. vii. 235 = Malacothrix saxatilis.
callianthum, Arv. in Bull. Soc. Dauph. (1880) 262 = amplexicaule.
callicomum ×, Naeg. & Peter, Hierac. 761.—Europ.
callimorphum, *Naeg. & Peter, l. c.* 896.—Europ.
calocephalum, Scheele, ex Willk. & Lange, Prod. Fl. Hisp. ii. 268 = neo-cerinthe.
caladon, Tausch, in Flora, xi. (1828) i. Erg. 56 = dubium.
calomastix ×, Naeg. & Peter, Hierac. 657.—Europ.
calophyllum, Uechtritz, in Oestr. Bot. Zeitschr. xxiv (1874) 106 = lanatum.
calophyton, *Naeg. & Peter, Hierac.* 422.—Europ.
calocicas ×, Naeg. & Peter, l. c. 762.—Europ.
calycinum, *Arv. in Bull. Soc. Dauph.* (1880) 282.—Europ.
calycinum, Hornem. ex Reichb. Fl. Germ. Excurs. 261 = H. Pilosella.
calyculatum, Hornem. Hort. Hafn. ii. 775 = saxetanum.
Camerarii, Collay, in Bull. Soc. Bot. Fr. vi. (1860) 796 = H. Pilosella.
canadense, Froel. in DC. Prod. vii. 218 = vulgatum.
canadense, Hort. Goett. ex Froel. l. c. 226 = sabaudum.
canadense, *Michx. Fl. Bor. Am.* ii. 86.—Am. bor.
canalicans, Tausch. in Flora, xi. (1828) i. Erg. 65 = pallidum.
candidissimum, Pourr. ex Nym. Consp. 446 = rupicolum.
candidum, Scheele, in Linnaea, xxxii. (1863) 678 = saxatile.
Candollei, *Froel. in DC. Prod.* vii. 212.—Europ.
Candollii, Monn. Ess. Hierac. 28 = glaciale.
canescens, *Link, Enum. Hort. Berol.* ii. 288.—Europ.
canescens, Schleich. Cat. Pl. Helv. ed. III. 17 = vulgatum.
canaticum, Arv. in Bull. Soc. Dauph. (1880) 289 = cydoniaefolium.
canum, *Naeg. & Peter, Hierac.* 426.—Europ.
capense, *Linn. Pl. Afr. Rar.* 17; *Amoen. Acad.* vi. 96.—Afr. austr.
capillacrum, All. Fl. Pedem. i. 214 = glaucum.
capitatum, Hort. ex Stend. Nom. ed. II. i. 762 = cerinthoides.
capsirlense, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, vi. (1888-84) 184.—Europ.
Carionii, Bor. Fl. Cent. Fr. ed. III. ii. 382 = sabaudum.
carneum, *Greene, in Coult. Bot. Gas.* vi. (1881) 184.—Am. bor. occ.
carolinianum, Fries, Symb. Hierac. 145 = marianum.
carpathicum, *Bess. Prim. Fl. Galic.* ii. 154.—Europ.
carpaticum, Griseb. in Goett. Abh. v. (1853) 114 = bohemicum.
carpetanum, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 266 = maculentum.
castellanum, Boiss. & Rent. Diagn. Pl. Nov. Hisp. 20 = *bifurcum*.
caucasicum, Fries, Epicris. Hierac. 46 = hispidum.
caulophyllum, *Bertol. Misc. Bot.* iv. 12.—Am. bor.
celsicaule, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 402 = vulgatum.
cephaloides, *Arv. in Bull. Soc. Dauph.* (1880) 287.—Europ.
cerasiphyllum, Desf. Cat. Hort. Par. ed. III. 145 = umbellatum.
cerinthoides, Backh. Mon. Hierac. 36 = anglicum.
cerinthoides, Boeb. in Pall. N. Nord. Beitr. vi. 259 (Quid ?).—Ross. med.
cerinthoides, Gaud. Fl. Helv. v. 119 = compositum.
cerinthoides, Gouan, Illustr. 58. t. 25 = Neo-Cerinthe.
cerinthoides, *Linn. Sp. Pl.* 908.—Europ.
cerinthoides, Monn. Ess. Hierac. 49 = prenanthoides.
cerinthoides, Vill. Fl. Delphin. 86; Hist. Fl. Dauph. iii. 110 = Pseudo-cerinthe.
cerinum, *Fries, Symb. Hierac.* 10.—Norweg.
cernuum, Friwald. ex Boiss. Fl. Orient. iii. 878 = sparsum.

HIERACIUM :—
cernuum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 62 = Crepis neglecta.
chalcidicum, *Boiss. & Heldr. Diagn.* Ser. II. iii. 104. — Graecia.
chalicophilum, Jord. ex Billot, Annot. 119 = saxatile.
charadriorum, Jord. ex Billot, l. c. = caesium.
cheiranthifolium, Balll. Hist. Fl. viii. 294 = Andryala ragusina.
cheirifolium, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 866.—Persia.
chenacaule, Vukot. ex Nym. Consp. 436 = sabaudum.
cherleuse, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 397 = vulgatum.
chilense, *Less. in Linnaea*, vi. (1831) 100.—Chili.
chlorasfolium, Arv. in Bull. Soc. Dauph. (1880) 282 = saxetanum.
chlorocephalum, Wimm. ex Nym. Consp. 447 = bohemicum.
chlorophyllum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 398 = vulgatum.
chlorosporum, Gren. & Godr. Fl. Fr. ii. 368 = saxetanum.
chlorospermum, Froel. in DC. Prod. vii. 222 = crinitum.
chloroticum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 386, in syn. = boreale.
chondrillaefolium, Fries, Epicris. Hierac. 87 = chondrilloides.
chondrilloides, Linn. Sp. Pl. 801 = Crepis jacquini.
chondrilloides, *Vill. Hist. Pl. Dauph.* iii. 114.—Europ.
chrysanthum, Backh. Monog. Hierac. 34 = nigrescens.
chrysanthum, Ledeb. Fl. Alt. iv. 129 = Crepis chrysantha.
chrysocephaloides, *Norrl. ex Wainio, in Meddel. Soc. Faun. et Fl. Fenn.* iii. 65.—Fennia.
cibinense, Schur, Enum. Pl. Transs. 399 = boreale.
cichoriifolium, Host Barn. ex Froel in DC. Prod. vii. 203 = pratense.
cidonifolium, Vill. Prosp. 34 = cydoniaefolium.
ciliatum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 77 = andryaloides.
ciliatum, Willd. Sp. Pl. iii. 1585 = Picris Sprengeriana.
ciliolatum, *Serres, in Bull. Soc. Bot. Fr.* iii. (1856) 276. — Gallia.
ciliosum, Pané, ex Nym. Consp. 442 = eximium.
cincinnatum, Fries, Epicris. Hierac. 181 = boreale.
Cineraria, *Naeg. & Peter, Hierac.* 664.—Europ.
cinerascens, Jord. Gr. Rec. Jard. Gren. (1849) 17 = laciophyllum.
cinereum, Doell, Rhela. Fl. 524 = H. Pilosella.
cinereum, Tausch, in Flora, ii. (1819) ii. 463 = setigerum, *Tausch*; echioides.
cinerum, Schur, Enum. Pl. Transs. 881 = brachiatum.
cirritum, *Arv. in Bull. Soc. Dauph.* (1880) 287.—Europ.
cladocraum ×, Vukot. in Rad Jugos. Akad. Zagreb. xxxiv. (1876) 18.—Europ.
cladophyllum, Bull. Soc. Bot. Fr. xxi. (1874), Rev. Bibl. 227, sphalm., vide *calophyllum*.
cochleare, *Huter, ex Dalla Torre, Atl. Alpenfl.* 149.—Europ.
coerulcum, Scop. Fl. Carn. ed. II. ii. 111 = Sonchus alpinus.
collinum, Besa. Prim. Fl. Galic. ii. 148 = *bifurcum*.
collinum, Bohm. ex Fries, Symb. Hierac. 16 = sueicum.
collinum, Bot. Norveg. ex Fries, l. c. 17 = floribundum.
collinum, Froel, in DC. Prod. vii. 208, partim = glomeratum.
collinum, Gochn. Tent. Med. Bot. Cich. 17. t. 1 = praecultum.
collinum, Kostel. ex Froel. in DC. Prod. vii. 200 = H. Pilosella.
collinum, Tausch, in Flora, xi. (1828) i. Erg. 58; Fries, Symb. Hierac. 29 = dubium.
collinum, Willd. ex Froel. in DC. Prod. vii. 203 = pratense.
Columnae, Tenore, Syll. Fl. Nap. 398 = Crepis Columnae.
comatulum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 410 = murorum.
comatum, *Fries, Symb. Hierac.* 148.—Mexic.
Commersonii, *Monn. Ess. Hierac.* 43.—Reg. Argent.
commiticum, Jord. Cat. Gr. Jard. Dijon (1848) 20 = vulgatum.
commutatum, *Beck, ex Lindeb. in Blytt, Norges Fl.* ii. 677.—Europ.

HIERACIUM :—
commutatum, J. Gay, ex Monn. Ess. Hierac. 48 = pulmonarioides.
commutatum, *Lindeb. in* Hartm. Handb. Skand. Fl. ed. X. 24 = nigrescens.
Comperianum, *Andrz. Rys Botan.* ii. (1830) 27.—Ross. austr. occ.
compositum, *Lapeyr. Hist. Abr. Pl. Pyr.* 476.—Mont. Pyren.
concinnum, Jord. Gr. Jard. Gren. (1849) 17 = boreale.
confertum, Jord. ex Nym. Consp. 443 = vulgatum.
confusum, Jord. l. c. 440, 443 = virosum, vulgatum.
conicum, Arv. in Bull. Soc. Dauph. (1880) 290 = lycopaifolium.
conjugatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 418 = murorum.
conseciatum, Jord. in Bor. l. c. 400 = vulgatum.
conspicuum, Jord. in Bor. l. c. 393 = boreale.
constrictum, *Arv. in Bull. Soc. Dauph* (1880) 289.—Europ.
contiguum, Juratz. ex Nym. Consp. 445 = canescens.
controversum, Timb. in Mém. Acad. Toulouse, Sér. IV. vi. (1856) 128 = sabaudum.
convenarum, *Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, ii. (1874) 462.—Europ.
conyzaefolium, Roth, Tent. Fl. Germ. ii. ii. 269 = Crepis biattaridoides ?
conyzaefolium, Gouan, Illustr. 59 = Crepis grandiflora.
conyzoideum, Lam. Fl. Fr. ii. 97 = Crepis grandiflora.
corconticum, *K. Knaf, ex Čelak. in Oestr. Bot. Zeitschr.* xxxiii. (1883) 79.—Europ.
cordatum, Scheele, ex Costa, Fl. Catal. 158 = Pseudo-Cerinthe.
cordifolium, Lapeyr. Suppl. Hist. Pl. Pyr. 128 = compositum.
cordifolium, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 73 = Crepis pygmaea.
coriaceum, *Mertr. Pl. Tirol.* 438.—Europ.
coriaceum, *Scheele, ex Willk. & Lange, Prod. Fl. Hisp.* ii. 269.—Hispan.
coronopifolium, Bernh. ex Hornem. Hort. Hafn. 970 = umbellatum.
corrugatum, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, vi. (1888-84) 194.—Europ.
corruscans, *Fries, Symb. Hierac.* 62.—Mont. Pyren.
corsicum, Hort. ex Stend. Nom. ed. II. i. 762 = cerinthoides.
corymbosum, Fries, Symb. Hierac. 185 = H. Eupatorium.
corymbosum, C. Koch, in Linnaea, xxiii. (1850) 680 = boreale.
corymbosum, Pers. Syn. ii. 374 = virosum.
corymbuliferum, Gmelin, Excurs. Fl. Schw. ed. II. 264 = praealtum.
Costae, Scheele, in Linnaea, xxxii. (1863) 686 = pyrenaeum.
cotonefolium, Froel. in DC. Prod. vii. 210 = vallisicum.
cotonefolium, Gerh. ex Froel. l. c. 226 = sabaudum.
cotonefolium, *Hort. Par. ex Lam. Encyc.* ii. 367.—Europ.
crassicaule, Stend. Nom. ed. II. i. 762 = sabaudum.
crassifolium, *Bernh. ex Hornem. Hort. Hafn. Suppl.* 185.—Hab ?
crassifolium, Hort. ex Stend. Nom. ed. II. i. 762 = villosum.
crassifolium, Schult. Oba. Bot. 165 = graminifolium.
crassisetum ×, Naeg. & Peter, Hierac. 502.—Europ.
crepidinerum, Froel. ex DC. Prod. vii. 242 = Nabalus crepidineus.
Crepidisparmum, *Fries, Symb. Hierac.* 146.—Mexic.
crinigerum, Fries, l. c. 94 = pallidum.
crinigerum, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. Math. Toulouse*, iii. (1875-76) 610.—Europ.
crinitum, Fries, ex Willk. & Lange, Prod. Fl. Hisp. ii. 265 = pyrenaicum.
crinitum, Jan, ex Froel. in DC. Prod. vii. 222 = symphitifolium.
crinitum, *Sibth. & Sm. Fl. Graec. Prod.* ii. 184.—Sicil.; As. Min.
crinitum, Tenore, ex Monn. Ess. Hierac. 42 = compositum.
crispatum, Jord. in Billot, Annot. 129 = andryaloides.
crispum, Dalac, Fl. Hautes-Pyr. 488 = mixtum.

HIERACIUM:—
crisaliense, Boiss. & Reut. ex Nym. Consp. 440 = prenanthoides.
croaticum, Lapeyr. Hist. Abr. Pl. Pyr. 475 = cerinthoides.
croaticum, Rostk. & Schmidt, Fl. Sedin. 322 = saxetanum.
croaticum, Schloss. ex Nym. Consp. 439 = boreale.
croaticum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 78 = racemosum.
croaticum, Waldst. & Kit Pl. Rar. Hung. iii. 242. t. 218 = Crepis succisaefolia.
crocatum, Bunge, ex Ledeb. Fl. Ross. ii. 825 = Crepis Pallasii.
crocatum, *Fries, Symb. Hierac.* 183.—Europ.; Reg. Himal.
crocatum, Wimm. Fl. Schles. ed. III. (1857) 315 = gothicum.
croceum, Lam. Encyc. ii. 360 = Crepis Pallasii.
crocetum, Jord. Gr. Jard. Gren. (1849) 18 = vulgatum.
Cserriasum, Baung. Enum. Stirp. Transs. iii. 20 = glabratum.
curvidens, Jord. Gr. Jard. Gren. (1849) 18 = boreale.
cuspidatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii 396 = tridentatum.
cuspidatum, *Lindeb. in Bot. Notiser* (1878), 184.—Europ.
cuspidatum, Scheele, ex Costa, Fl. Catal. 158 = Pseudo-Cerinthe.
cyclophyllum, Jeanb & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875-76) 609 = alatum.
cydonaefolium, Fries, Summa Veg. Scand. 7 = dovrense.
cydoniaefolium, Froel. in DC. Prod. vii. 212 = juranum.
cydoniaefolium, Tausch, in Flora, xi. (1828) 1. Erg. 75 = bohemicum.
cydoniaefolium, *Vill. Hist. Pl. Dauph.* iii. 107.—Europ.; Reg. Caucas.
cydoniaefolium, Willd. Enum. Hort. Berol. 824 = strictum.
cymiflorum, *Naeg. & Peter, Hierac.* 424.—Europ.
cymigerum, Reichb. in Moesal. Handb. ed. II. ii. 1384 = cymosum.
cymosiflorum, *Froel. in DC. Prod.* vii. 207.—Palatinat.
cymosum, Leers, Fl. Herborn. ed. II. 175 = praealtum.
cymosum, *Linn. Sp. Pl.* ed. II. 1126.—Europ. med.
cymosum, O. F. Muell. in Fl. Dan. t. 810 = glomeratum.
cymosum, Vill. Voy. 62. t. 4. f. 2 = sabinum.
cymosum, Willd. Sp. Pl. iii. 1566 = pratense.
cynoglossifolium, Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875-76) 611 = lasiophyllum.
cynoglossoides, *Arv. ex A. Gray, in Proc. Am. Acad.* xix. (1883) 68.—Am. bor. occ.
Cserriasum, Nym. Consp. 450, sphalm., vide *Cerriasum*.
dacicum, *Uechtritz, in Oestr. Bot. Zeitschr.* xxv. (1875) 314.—Transylv.
dannbiale, Pollich, Hist. Pl. Palat. ii. 385 = Leontodon hastilis.
dasyophalum, Froel. in DC. Prod. vii. 234 = Llottardi.
dasypodum, Jord. ex Nym. Consp. 448 = amplexicaule.
dasytrichum, *Arv. in Bull. Soc. Dauph.* (1880) 283.—Europ.
debile, *Fries, Epicris. Hierac.* 146.—Ecuador.
debile, Rehm. in Oestr. Bot. Zeitschr. xxiii. (1873) 184 = alpinum.
decipiens, *Monn ex Froel. in DC. Prod.* vii. 230.—Europ.
decipiens, Nym. Consp. 449 = anglicum.
decipiens, Tausch, in Flora, xi. (1828) 1. Erg. 66 = nigrescens.
decoloratum, *Fries, Symb. Hierac.* 21.—Europ.; Sibir.
decrescens, Knaf, ex Nym. Consp. 441 = gothicum.
Delasorii, Lagg. ex Nym. l. c. 450 = saxetanum.
densicapillum, *Naeg. & Peter, Hierac.* 446.—Europ.
densiflorum, Tausch, in Flora, xi. (1828) 1. Erg. 59 = H. Auricula †
dentatum, *Hoppe, in Sturm, Deutsch. Fl.* Heft 39.—Europ.
dentatum, Lagg. in Bull. Soc. Murith. ii. (1873) 72 = scorzonerifolium, *La Soie*.
denticulatum, Sm. Engl. Bot. t. 2122 = strictum.
densulatum, Lapeyr. Hist. Abr. Pl. Pyr. 473 = pyrenaicum.
densulatum, Rochel, ex Schult. Oesterr. Fl. ed. II. ii. 437 = bupleuroides.

HIERACIUM:—
depauperatum, Jord. Obs. Pl. Crit. vii. (1846) 38 = boreale.
depilatum, Almq. in Bot. Notiser (1867), 68 = linguilatum.
Desegliesi, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 392 = umbellatum.
Dewari, *Boswell, [Syme] in Trans. Soc. Bot. Edinb.* xiii. (1879) 211.—Brit.
diaphanoides, Lindeb. in Bot. Notiser (1882), 127 = murorum.
diaphanum, Arv. in Soc. Dauph. Exch. Pl. iv. (1877) 105 = vulgatum.
diaphanum, *Fries, Nov. Fl. Suec.* ed. I. 75; *ex ej. Mant.* ii. 45.—Scandinav.
dichotomum, Fries, ex Nym. Consp. 454 = cernuum.
dichotomum, *? F. Gmel. Syst.* 1179.—Sibir.
dichotomum, Koxb. ex DC. Prod. vii. 181 = Launaea asplenifolia.
dioreacoides, Vukot. in Rad Jugos. Akad. Zagreb. li (1880) 62 = vulgatum.
diffusum, Froel. in DC. Prod. vii. 222 = politum.
dimorphoides, *Norrl. ex Wainio, in Meddel. Soc. Faun. et Fl. Fenn.* iii. 65.—Fennia.
dimorphum, *Norrl. in Notis. Sällsk Faun. et Fl. Fenn.* II. viii. 182.—Fennia.
dinaricum, *Fries, Epicris. Hierac.* 106.—Europ.
dinothum × , Naeg. & Peter, Hierac. 508.—Europ.
discolor, Kit. in Linnaea, xxxii. (1863) 417 = stoloniflorum, *Waldst. & Kit.*
dissidens, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 408 = vulgatum.
dissitum, Verlot, ex Nym. Consp. 436 = boreale.
distractum, Fries, Epicris. Hierac. 72 = piloselloides.
divarcatum, Scheele, ex Willk. & Lange, Prod. Fl. Hisp. ii. 269 = prenanthoides.
divergens, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 411 = murorum.
divisum, Jord. Cat. Gr. Jard. Dijon (1848) 21 = vulgatum.
djemilense, *Boiss. & Bal. ex Boiss. Fl. Orient.* iii. 877.—As. Min.
Dollineri, Sch. Bip. ex Nym. Consp. 443 = fasciculare.
dovrense, Engl. ex Nym. l. c. = alsiacum.
dovrense, *Fries, Symb. Hierac.* 128.—Scandinav.
dovrense, Lindeb. ex Nym. Consp. 439 = truncatum.
dryadeum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 390 = umbellatum.
dryophilum, Jord. l. c. = umbellatum.
dubium, Chaix, in Vill. Hist. Pl. Dauph. i. 367; iii. 99; Leers, Fl. Herborn. ed. II. 174 = H. Auricula.
dubium, Hartm. Handb. Skand. Fl. ed. III. 185 = pratense.
dubium, Hoppe, ex Froel. in DC. Prod. vii. 201 = glaucocephalum.
dubium, *Linn. Sp. Pl.* 800.—Europ. bor.
dubium, Tausch, ex Reichb. Fl. Germ. Excurs. 261 = H. Pilosella.
dubium, Vahl, ex Steud. Nom. ed. II. i. 762 = praealtum.
dubium, Woodw. in With. Arr. Brit. Pl. ed. III. iii. 684 = stoloniflorum, *Waldst. & Kit.*
dumosum, *Jord. Gr. Jard. Gren.* (1849) 18.—Gallia.
dunense, Reyn. ex Hall, Fl. Belg. i. 566 = umbellatum.
duplex × , Naeg. & Peter, Hierac. 486.—Europ.
dycalum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 391 = umbellatum.
echioides, *Lamarck. Fl. Fasan.* 846.—Europ.; Aa. bor.
edituum, Jord. ex Nym. Consp. 438 = boreale.
elatum, *Fries, Symb. Hierac.* 167.—Europ. med.
elatum, Gren. & Godr. Fl. Fr. ii. 280 = juranum.
elatum, Lindeb. in Bot. Notiser (1878), 182 = decoloratum.
elegans, *Lindeb. l. c.* (1872) 121.—Europ.
elegans, Reichb. ex Nym. Consp. 451 = praealtum.
elisum, Arv. in Bull. Soc. Dauph. (1880) 287 = alpinum.
ellipticum, Jord. Cat. Gr. Jard. Dijon (1848), 21 = vulgatum.
elongatum, Bot. Helv. ex Fries, Symb. Hierac. 61 = valdepilosum.
elongatum, Lapeyr. Hist. Abr. Pl. Pyr. 476 = Neo-Cerinthe.
elongatum, Thom. ex Fries, Symb. Hierac. 62 = Candollei.
elongatum, Un. Itin. Easling. (1850); ex Froel. in DC. Prod. vii. 216 = compositum.

HIERACIUM:—
elongatum, Willd. ex Froel. in DC. Prod. vii. 229 = villosum.
emarginatum, *Schoele, in Linnaea*, xxvii. (1853) 680.—Europ.
eminens × , Naeg. & Peter, Hierac. 230.—Europ.
Engleri, Uechtritz, in Oestr. Bot. Zeitschr. xxi. (1871) 293 = alsiacum.
epimedium, *Fries, Epicris. Hierac.* 108.—Europ.
erianthum, *H. B. & K. Nov. Gen. et Sp.* iv. 4.—Am. austr.
ericaceum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 46 = transsilvanicum.
eriocephalum, *Wedd. Chlor. And.* i. 226.—Ecuador.
eriocrinthe, Fries, ex Nym. Consp. 448 = compositum.
eriophion, Schur, Enum. Pl. Transs. 395 = rotundatum.
eriophorum, Lapeyr. Hist. Abr. Pl. Pyr. 474 = sabaudum.
eriophorum, Schur, Enum. Pl. Transs. 395 = rotundatum.
eriophorum, *St. Amans, in Bull. Soc. Philom.* iii. (1809) 26. t. 2. f. 1.—Gallia.
eriophyllum, *Link, Enum. Hort. Berol.* ii. 286.—Hab.†
eriophyllum, Schur, Enum. Pl. Transs. 395 = rotundatum.
eriophyllum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 77 = lanatum.
eriophyllum, Willd. Enum. Hort. Berol. Suppl. 54 = villosum.
eriopsilon, Jord. in Billot, Annot. 128 = andryaloides.
eriopus, Boiss. & Heldr. Diagn. Ser. II. iii. 106 = crinitum.
erubescens, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 398 = vulgatum.
erythropodum, *Uechtr. ex Velen. in Oestr. Bot. Zeitschr.* xxxiii. (1883) 891.—Europ.
erythropermum, Greene, ex A. Gray, Syn. Fl. N. Am. i. ii. 429 = Fendleri.
euchaetium, *Naeg & Peter, Hierac.* 764.—Europ.
Eupatorium, *Griseb. in Goett. Abh.* v. (1853) 131.—Europ. bor.
eurylepium, *Naeg. & Peter, Hierac.* 247.—Europ.
euryops, *Knaf, in Oestr. Bot. Zeitschr.* xxv. (1875) 181 —Europ.
exclusum, *Rehm. l. c.* xxiii. (1873) 109.—Europ.
eximium, *Backh. Monogr. Hierac.* 90.—Europ.
exotericum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 417 = murorum.
extensum, Laebeck, ex Lindeb. in Hartm. Handb. Skand. Fl. ed. XI. 42 = pallidum.
fagicolum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 416 = murorum.
falcatum, Arv. in Bull. Soc. Dauph (1880) 282 = bupleuroides.
falcidens, Arv. ex Lamotte, Prod. ii. 481 = neocerinthe.
fallacinum, F. W. Schultz, Fl. Pfalz, 277 = H. Pilosella.
fallax, Froel. in DC. Prod. vii. 205 = praealtum.
fallax, Willd. Enum. Hort. Berol. 822 = dubium.
fallens, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 412 = murorum.
fariniferum, *Boiss. Fl. Orient.* iii. 866.—Armenia.
farinorum, Lam. Fl. Fr. ii. 94 = glaucum.
farinulentum, Griseb. in Goett. Abh. v. (1853) 144 = lasiophyllum.
farinulentum, Jord. Cat. Gr. Jard. Dijon (1848) 21 = pictum.
fasciculare, *Fries, Epicris. Hierac.* 100.—Europ.
fasciculatum, Pursh, Fl. Am. Sept. ii. 504 = canadense.
fasciculatum, Torr. ex Froel. in DC. Prod. vii. 217 = scabrum.
fastigiatum, Fries, Symb. Hierac. 119 = vulgatum.
fastigiatum, Tausch, in Flora, xi. (1828) 1. Erg. 59 = praealtum.
fastigiosum, Gremli, Excurs. Fl. Schw. ed. II. 268 = praealtum.
Favrati, Muret, ex Gremli, l. c. 278 = lycopodifolium.
Favrat, Wolf. in Bull. Soc. Murith. x. (1881) 21, nomen = prec.?
Fendleri, *Sch. Bip. in Bonplandia*, ix. (1861) 178.—Am. bor. occ.
fennicum, Norrl. ex Wainio, in Meddel. Soc. Faun. Fl. Fenn. iii. (1878) 65 = saxicum.

HIERACIUM :—

fontinum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 399 = vulgatum.

fictum, Jord. in Bor. l. c. 408 = vulgatum.

filiformum, Tausch, in Flora, xi. (1828) i. Erg. 59 = praealtum.

filiforme, *Lindeb. in Hartm. Handb. Skand. Fl.* ed. X. 21.—Europ.

filiforme, *Petr. Encyc. Suppl.* ii. 560.—Ins. Mauric.

finalense, *Naeg. & Peter, Hierac.* 269.—Europ.

finitimum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 398 = vulgatum.

firmatum, Jord. l. c. 394 = tridentatum.

firmum, Jord. Cat. Gr. Jard. Dijon (1848) 22 = tridentatum.

fissifolium, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 414 = murorum.

fistulosum, *Lindeb. in Hartm. Handb. Skand. Fl.* ed. XI. 57.—Europ.

flaccidum, *Fries, in Oefver. Vet. Akad. Forrhandl. Stockh.* (1856) 145.—Bras.

flagellare, Reichb. Fl. Saxon. 178 = stoloniflorum, *Waldst. & Kit.*

flagellare, Schur, Enum. Pl Transs. 381 = brachiatum.

flagellare, Willd. Enum. Hort. Berol. Suppl. 54 = H Pilosella.

flagelliformum, Rayaud, in Soc. Dauph. Exch. Pl. iv. (1877) 117 = boreale.

flagelliformum ×, Čelak. in Arch. Naturw. Landesd. Boehm. iv. (1881) 787.—Europ.

flagellosum, *Jord. ex Martr. Fl. Tarn.* 452.—Europ.

flammeum, *Lindeb. in Hartm. Handb. Skand. Fl.* ed. X. 20.—Europ.

flexicaule, Tausch, in Flora, xi. (1828) i. Erg. 66 = murorum.

flexuosum, Balb. ex Steud. Nom. ed. II. i. 763 = dentatum.

flexuosum, DC. Fl. Fr. vi. 436 = glabratum.

flexuosum, Gaud. Fl. Helv. v. 98 = cerinthoides.

flexuosum, J. F. Gmel. Syst. 1179 = incertum.

flexuosum, Hort. Taur. ex Froel. in DC. Prod. vii. 212 = speciosum.

flexuosum, Lapeyr. Hist. Abr. Pl. Pyren. 475 = Perusianum.

flexuosum, *Martr. Fl. Tarn*, 483.—Gallia.

flexuosum, Schleich. ex Steud. Nom. ed. I. 407 = villosum.

flexuosum, *Waldst. & Kit. ex Willd. Sp. Pl.* iii 1581 —Europ.

floccosum, Bab. Man. Brit. Bot. ed. V. 205 = flocculoseum.

floccosum, Schur, Enum. Pl. Transs. 388 = flexuosum, *Waldst. & Kit.*

floculosum, *Backh. Mon. Hierac.* 60.—Europ.

florentinum, All. Fl. Pedem. i. 218 = piloselloides.

florentinum, Lasch. ex Nym. Consp. 453 = H. Auricula.

florentinum, Spreng. Fl. Hal. 222. t. 10. f. 1 = praealtum.

floribundum, *Wimm. & Grab. Fl. Siles.* ii. 11. 204.—Europ. med.

floscullosum, *Hornem. Hort. Hafn.* 760.—Hab. ?

fluminense, *Kern. in Oestr. Bot. Zeitschr.* xxiv. (1874) 171.—Europ.

foecundum, Froel. ex Steud. Nom. ed. II. i. 763 = radicans.

foecundum, Jord. ex Nym. Consp. 443 = vulgatum.

foeniculaceum, Wulf. ex Steud. Nom. ed. II. i. 763 = Crepis foeniculacea.

foetidum, Sibth. & Sm. Fl. Graec. viii. 74. t. 799 = Crepis Sibthorpiana.

foetidum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 68 = Crepis foetida.

foetidum, Willd. Sp. Pl. iii. 1575 = Crepis foetens.

foliosum, Steud. Nom. ed. I. 407 = Gronovii.

foliosum, *Waldst. & Kit. ex Willd. Sp. Pl.* iii. 1589 ; *Fl. Bar. Hung.* ii. 155. t. 145.—Europ. or. ; Reg. Caucas.

Forsskahlei, Froel. in DC. Prod. vii. 223 = boreale.

fragile, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 158. —N. Zel.

fragile, *Jord. Obs. Pl. Crit.* vii. (1849) 84.—Gallia ; Hispan.

fraxini, Sch. Bip. Cich. 34 = H. Pilosella.

Freyeanum, *Velen. in Oestr. Bot. Zeitschr.* xxxiii. (1883) 389.—Europ.

Friesii, Fries, ex Nym. Consp. 440 = crocatum.

HIERACIUM :—

Friesii, Hartm. Handb. Skand. Fl. ed. IV. 257 = rigidum.

Friesii, Sch. Bip. in Bonplandia, ix. (1861) 226 = Schultzii.

frigidarium, *Naeg. & Peter, Hierac.* 656.—Europ.

frigidum, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) ii. 112 = Crepis chrysantha.

frigidum, *Wedd. Chlor. And.* i. 225. t. 42.—Venezuela ; N. Granad.

Fritzei, *F. Schultz, in Flora*, lv. (1872) 281.—Europ.

Friwaldii, Reichb. f. ic. Fl. Germ. xix. 94 = pannosum.

Froelichianum, Buek, Ind. ad DC. Prod. ii. p. vi = maculentum.

Froelschii, Fries, Epicris. Hierac. 146 = maculentum.

Froelichii, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875–76) 392 = cerinthoides.

fructiscrum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 384 = boreale.

frutticosum, Willd. Sp. Pl. iii. 1591 = Tolpis fruticosa.

fulgens, *Naeg. & Peter. Hierac.* 348.—Europ.

fulgidum, *Heynh. ex Reichb. f. ic. Fl. Germ.* xix. 55.—Hab. ?

fulgidum, *Kreck. Fl. Siles. Suppl.* ii 258.—Europ.

fulgidum, Wallr. ex Nym. Consp. 441 = gothicum.

fuliginatum, Huter, ex Nym. Consp. 452 = subnivale.

fuliginosum, Anderss. ex Fries, Symb. Hierac. 105 = stratum.

fulvicapillum, Link, Handb. i. 801 = H. Pilosella.

fulvipes, *Wedd. Chlor. And.* i. 224.—Bolivia.

furcatum, Hoppe, in Flora, xiv. (1831) i. 181 = sphaerocephalum.

furcellatum, Fries, Symb. Hierac. 102 = crocomiodes.

Furcellum, *Naeg. & Peter, Hierac.* 656.—Europ.

furcillatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 418 = murorum.

furfuricallum, *Jord. ex Matr. Fl. Tarn*, 489.—Europ.

fuscidulum, Jord. ex Nym. Consp. 439 = boreale.

fuscoatrum, *Naeg. & Peter, Hierac.* 315.—Europ.

fuscum, Fries, ex Nym. Consp. 453 = Blyttianum.

fuscum, Vill. Voy. 19. t. 1 = aurantiacum.

fuxianum, Schur, Enum. Pl. Transs. 384 = piloselloides.

galactinum, *Norrl. ex Wainio, in Meddel. Soc. Faun. et Fl. Fenn.* iii. 65.—Fennia.

gallicum, Jord. Gr. Jard. Gren. (1849) 19 = boreale.

Ganderi, Hausm. ex Nym. Consp. 448 = fasciculare.

Ganderi, Huter, ex Nym. l. c. 440 = cydoniaefolium.

Garckeanum, Aschers. Ind. Sem. Hort. Berol. (1869) 24 ; et (1872) App. 8 = tridentatum.

Garibaldium, *Fries, Epicris. Hierac.* 111.—Italia.

Gaudini, Christen. Hierac. Schweiz. 10 = dentatum.

Gaudryi, Boiss. & Orph. Diagn. Ser. II. iii. 105 = pannosum.

Gautieranum, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, vi. (1888–84) 201.—Europ.

gentile, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 415 = murorum.

georgicum, Fries, Epicris. Hierac. 106 = Crepis glabra germanicum, *Naeg. & Peter, Hierac.* 738.—Europ.

Gibsoni, Backh. Mon. Hierac. 47 = caesium.

gioberrimum, Spreng. Syst. iii. 643 = graminifolium.

glabratum, *Hoppe, ex Willd. Sp. Pl.* iii. 1562.—Europ.

glabratum, Schleich. ex Koch, Syn. Fl. Germ. ed. I. 454 = glanduliferum.

glabratum, Willd. ex Froel. in DC. Prod. vii. 228 = villosum.

glabrum, Hoppe, ex Steud. Nom. ed. II. i. 763 = villosum.

glabrum, *Kit. in Linnaea*, xxii. (1865) 433.—Europ.

glabrum, *Kreck. Fl. Siles.* ii. ii. 816.—Europ.

glabrum, Turcz. in Bull. Soc. Nat. Mosc. (1838) 96 ; xxi. (1848) ii. 108 = Tarraxacum glabrum.

glaciale, C. Koch, in Linnaea, xxiii. (1850) 677 = sphaerocephalum.

glaciale, *Reyn. ex Lachen. in Nov. Act. Helv.* i. (1787) 306.—Europ.

glaciallum, *Naeg. & Peter, Hierac.* 269.—Europ.

gladiatum, *Martr. Fl. Tarn*, 485.—Europ.

glanduliferum, *Hoppe, in Sturm, Deutschl. Fl.* Heft 29.—Europ.

glandulosum, Schult. Oba. Bot. 166 = Crepis grandiflora.

glareosum, Serres, in Bull. Soc. Bot. Fr. ii. (1855) 224 = rigidum, tridentatum.

glastifolium, Willd. ex Steud. Nom. ed. II. i. 763 = villosum.

glaucellum, Lindeb. in Hartm. Handb. Skand. Fl. ed. X. 37 = vulgatum.

HIERACIUM :—

glaucescens, Bess. Prim. Fl. Gall. ii. 150 = H. Auricula.

glaucescens, Froel. in DC. Prod. vii. 202 = praealtum.

glaucescens, Hort. Landsh. ex Froel. l. c. 203 = prattense.

glaucifolium, *Poepp. ex Froel. l. c.* 207.—Chili.

glaucinum, Jord. Cat. Gr. Jard. Dijon (1848) 22 = fragile, *Jord.*

glaucophyllum, Huet, ex Nym. Consp. 450 = glaucinum.

glaucophyllum, Scheele, in Linnaea, xxxii. (1863) 658 = Pseudo-Cerinthe.

glaucophyllum, Schur, Enum. Pl. Transs. 386 = subfuscum.

glaucopsis, Gren. & Godr. Fl. Fr. ii. 355 = saxicolum.

glaucovirens, *Jeanb. & Timb. in Bull. Soc. Phys. Nat. Toulouse*, vi. (1888–84) 191.—Europ.

glaucum, *All. Fl. Pedem.* i. 214.—Europ. austr.

glaucum, Alsch. Fl. Jadr. 202 ; ex DC. Prod. vii. 169 = Crepis heterogyna.

glaucum, Lej. Fl. Spa, ii. 129 = prasinum.

glaucum, Schleich. ex Moun. Eas. Hierac. 14 = graminifolium.

glaucum, Vill. Hist. Pl. Dauph. iii. 116 = politum.

glaucum, Wahlenb. Fl. Carp. 241 = bupleuroides.

globosum, Backh. Mon. Hierac. 25 = nigrescens.

globosum, *Desf. Cat. Hort. Par.* ed. III. 308.—Hab. ?

globuliferum, Tausch, ex Steud. Nom. ed. II. i. 763 = piloselloides.

glomeratum, Fries, Summ. Veg. Scand. 533 = dublum.

glomeratum, *Froel. in DC. Prod.* vii. 207.—Europ.

glumaceum, *Fries, Epicris. Hierac.* 141.—Ind. or.

glutinosum, *Linn. Sp. Pl.* 804.—Gallia.

Gmelini, Lam. i. c. 802 = Crepis Gmelini.

Gmelini, Schreb. ex Froel. in DC. Prod. vii. 188 = Crepis Schreberi.

Gmelini, Willd. ex Spreng. Syst. iii. 637 = Pterotheca hilida.

gnaphaloides, *Arv. in Bull. Soc. Dauph.* (1880) 280. —Europ.

Gochnati, C. A. Mey. Verz. Pfl. Cauc. 89 = pratense.

Gochnati, Spreng. Syst. iii. 639 = praealtum.

Godeti, Christen. ex Gremli, Beitr. 94 = Jacquini.

Godroniauum, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875–76) 388 = cerinthoides.

gombrense, Lagg. ex Christen. Hierac. Schweiz. 19 (*gombrene*) = anglicum.

gothicum, *Fries, Symb. Hierac* 121.—Scandinav.

Gouani, *Hegetschw. Fl. Schw.* 746.—Europ.

Gougetianum, Gren. & Godr. Fl. Fr. ii. 368 = anglicum.

gracile, Froel. in DC. Prod. vii. 231 = maculentum.

gracile, *Hook. Fl. Bor. Am.* i. 296.—Am. bor. occ.

gracillentum, Backh. Mon. Hierac. 24 = nigrescens.

gracilisepum, Sch. Bip. ex Nym. Consp. 446 = tomentosum.

graecum, *Boiss. & Heldr. Diagn.* Ser. II. iii. 101.—Graecia.

graminifolium, *Gaud. in Nat. Ana. Schw. Ges.* iv. (1820) 43.—Europ.

graminifolium, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 74 = porrifolium.

grandidentatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 383 = sabaudum.

grandiflorum, All. Fl. Pedem. i. 217 = Crepis grandiflora.

grandiflolium, Arv. in Soc. Dauph. Exch. Pl. iv. (1877) 108 = hemiplexum.

grandiflolium, *Sch. Bip. in Bull. Soc. Bot. Fr.* ix. (1862) 440.—Algeria.

granaticum, Sch. Bip. Cich. 38 = lastophyllum.

granulosa, Froel. in DC. Prod. vii. 219 = saxagranum.

Greenei, *A. Gray, in Proc. Am. Acad.* xix. (1883) 69. —Calif.

Gremlei, *Schenle, in Linnaea*, xxxii. (1863) 688. — Europ.

Grenieri, Timb. & Jeanb. in Bull. Soc. Bot. Fr. xi. (1864) p. lxxiii = cerinthoides.

Grisebachii, *A. Kern. Sched. Fl. Exsicc.* i. 63.— Europ.

Gronovii, *Linn. Sp. Pl.* 802.—Am. bor.

Gronovii, Willd. ex Froel. in DC. Prod. vii. 217 = venosum.

Guthnikianum, *Hegetschw. Fl. Schw.* 781.—Europ.

gymnocephalum, *Griseb. ex Pant. in Oestr. Bot. Zeitschr.* xxiii. (1873) 366.—Montenegro.

hadracaulon x, Nasg. & Peter, Hierac. 680.—Europ.
halimifolium, Froel. ex Fries, Epicris. Hierac. 186 = umbellatum.
Halleri, Hook. in Curt. Fl. Lond. v. 208. t. 215 = anglicum.
Halleri, Vill. Hist. Pl. Dauph. iii. 104 = alpinum.
hamadanense, *Heimerl, in Denkschr. Acad. Wiss*, l. (1888) 68.—Persia.
Hartwegii, Sch. Bip. ex Fries, Epicris. Hierac. 147 = erianthum.
Hausmanni, Reichb. f. Ic. Fl. Germ. xix. 58 = stoloniflorum, *Waldst. & Kit.*
Hausmanni, Sch. Bip. ex Nym. Consp. 439 = rigidum.
hedypnoides, *Linn. Sp. Pl.* 800.—Europ.
Heldreichii, *Boiss. Diagn.* Ser. II. iii. 102.—Graecia.
Helenium, *Dichtl. & Wiesb. in Deutsche Bot. Monatsschr.* ii. (1884) 66.—Austria.
helianthifolium, Froel. in DC. Prod. vii. 225 = amadense.
helveticum, *Bruegg. ex Dalla Torre, Atl. Alpenfl.* 149. —Europ.
helveticum, Suter, Fl. Helv. ii. 151 = glabratum.
hemipleucum, *Arv. in Soc. Dauph. Exch. Fl.* iv. (1877) 105; *Bull. Soc. Dauph.* (1880) 288.—Europ.
Herculis, Borb. ex Nym. Consp. 462 = sabinum.
heteranthum, Wall. Cat. n. 3358 = Lactuca Dubyaea.
heterodon, Arv. in Bull. Soc. Dauph. (1880) 285 = hamile.
heterodontum, Steud. Nom. ed. II. i. 763 = setigerum, *Tausch.*
heterogynum, Froel. ex Nym. Consp. 450 = stupposum.
heterophyllum, Bladon, in Phytol. ii. (1847) 961 = boreale.
heterophyllum, Retz. Obs. ii. 24 = Lactuca repens.
heterophyllum, Vukot, in Rad Jugos. Akad. Zagreb. vii. (1869) 69 = Crepis blennis.
heterogynum, *Arv. in Bull. Soc. Dauph.* (1880) 290. —Europ.
Heufleri, Janka, ex Nym. Consp. 452 = sabinum, pateuum.
Hinterhuberi, Sch. Bip. ex Griseb. in Goett. Abh. v. (1853) 88 = aurantiacum.
hirsutissimum, *Krock. Fl. Siles.* ii. 11, 806.—Europ.
hirsutulum, *Yord. ex Bor. in Mém. Soc. Acad. Maine-et-Loire*, vi. (1859) 110.—Europ.
hirsutum, *Bernh. ex Steud. Nom.* ed. I. 407, nomen; *Tausch, in Flora*, xi. (1828) 1. *Erg.* 72.—Am. bor.; Fyren.
hirsutum, Lapeyr. ex Fries, Symb. Hierac. 65 = phlomoides.
hirsutum, Schur, Enum. Pl. Transs. 399 = boreale.
hirsutum, Sesse & Moç. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 175 = strigosum.
hirtellum, *Lindeb. in Hartm. Handb. Skand. Fl.* ed. XI. 58.—Europ.
hirtum, *Froel. in DC. Prod.* vii. 213.—Europ.
hirtum, Polr. Encyc. Suppl. ii. 564 = scabrum.
hispidum, D. Don, Prod. Fl. Nep. 165 = Lactuca Dubyaea
hispidum, Forsk. Fl. Aegypt. Arab. 216 = boreale.
hispidum, Fries, Symb. Hierac. 168 = Bocconei.
hispidum, Vukot, in Rad Jugos. Akad. Zagreb. vii. (1869) 69 = Crepis setosa.
hololeion, *Maxim. Prim. Fl. Amur.* 182.—Reg. Amur.
holosericeum, Backh. Mon. Hierac. 19 = alpinum.
Hookeri, Steud. Nom. ed. II. i. 763 = gracile.
Hoppeanum, *Froel. in DC. Prod.* vii. 222.—Europ.
Hoppeanum, Schult. Oesterr. Fl. ed. II. ii. 428 = H. Pilosella.
Hoppeanum, Wallr. ex Nym. Consp. 444 = murorum.
horrens x, Nasg. & Peter, Hierac. 760.—Europ.
horridulum x, Nasg. & Peter, l. c. 761.—Europ.
horridum, *Fries, Epicris. Hierac.* 154.—Calif.
hortulanum, *Nasg. & Peter, Hierac.* 596.—Europ.
humifusum, Spreng. Pugill. ii. 76 = glaucum.
humile, *Jacq. Hort. Vindob.* iii. 2, in obs.; *Host, Syn. Fl. Austr.* 493.—Europ.
humile, Lapeyr. Hist. Abr. Pl. Pyr. 471 = amplexicaule.
Huteri, Hausm. ex Nym. Consp. 441 = picroides.
hybridum, *Chaix, in Vill. Hist. Pl. Dauph.* iii. 100,—Europ. med.
hybridum, Gand. Fl. Helv. v. 74 = brachiatum.
hybridum, Griseb. in Goett. Abh. v. (1853) 87 = sphaerocephalum.
hyoseridifolium, Vill. ex Pers. Syn. ii. 370 = Crepis hyoseridifolia.

hyperboreum, *Fries, Symb. Hierac.* 98.—Norveg.
hypenyrum, *Nasg. & Peter, Hierac.* 178.—Europ.
hypochoeridoneum, Arv. in Bull. Soc. Dauph. (1880) 287 = alpinum.
Hypochoeridis, Fries, Symb. Hierac. 118 = cassium.
Ignatianum, *Baker, in Mart. Fl. Bras.* vi. iii. 387.—Bras.
ilicetorum, Jord. ex Nym. Consp. 439 = umbellatum.
illyricum x, Nasg. & Peter, Hierac. 734.—Europ.
Illyricum, *Fries, Epicris. Hierac.* 71.—Europ.
imbricatum, *Lindeb. in Bot. Notiser* (1876), 153.—Europ.
imparum, Koel. ex Steud. Nom. ed. II. i. 764 = alpinum.
incanum, *Bieb. Fl. Taur. Cauc.* ii. 258.—Reg. Caucas.
incanum, Linn. Sp. Pl. 799 = Leontodon incanus.
incanum, Poll. Hist. Fl. Palat. ii. 353 = Leontodon hispidus.
incanum, Steph. Fl. Mosq. n. 523; ex Ledeb. Fl. Ross. ii. 780 = Leontodon hastilis.
incarnatum, Jacq. Coll. ii. 137 = Crepis incarnata.
incertum, *Froel. in DC. Prod.* vii. 226.—Sibiria.
incertum, Urv. in Mém. Soc. Linn. Par. iv. (1826) 808 = Erigeron Solivani.
incisifolium, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 409 = vulgatum.
incisum, *Hoppe, in Sturm, Deutschl. Fl.* Heft 89.—Europ.
incisum, Koch, Syn. Fl. Germ. ed. I. 457 = murorum.
incrassatum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 73 = Crepis montana.
indolatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 385 = boreale.
ineptum x, Nasg. & Peter, Hierac. 807.—Europ.
inops x, Nasg. & Peter, l. c. 674.—Europ.
inquinatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 404 = vulgatum.
insidiosum, Jord. ex Nym. Consp. 440 = cydoniaefolium.
insertum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 396 = tridentatum.
integrifolium, Hoppe, ex Willd. Sp. Pl. iii. 1566 = Crepis succisaefolia.
integrifolius, Lange, in Fl. Dan. t. 2661 = murorum.
intermedium, Hoppe, ex DC. Prod. vii. 150 = alpinum.
intermedium, Lapeyr. Hist. Abr. Pl. Pyr. 471 = humile.
intermedium, Rehm. in Oestr. Bot. Zeitschr. xxiii. (1873) 149 = pratense.
internodium, *Vest, in Flora*, iii. (1820) 5.—Styria.
intermedium, *Zumagg. Fl. Pedem.* i. 378.—Europ.
intersitum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 409 = vulgatum.
intybaceum, Hoppe, in Sturm, Deutschl. Fl. Heft 89 = amplexicaule.
intybaceum, *Jacq. Fl. Austr.* v. 52.—Europ.
Intybus, *Schmidt, ex Willd. & Lange, Prod. Fl. Hisp.* ii. 263 = Neo-Cerinthe.
inuloides, Sch. Bip. Cich. Suppl. ii. 146 = H. Espatorium.
inuloides, Tausch, in Flora, xx. (1837) 1. Beibl. 71 = boreale, crocatum.
iranense, Bornm. ex Oerst. in Kjoeb. Vidensk. Meddel. (1859) 113 = H. Lagopus.
iricum, *Fries, Symb. Hierac.* 60.—Europ.
irriguum, Griseb ex Nym. Consp. 439 = rigidum.
italicum, Fries, Symb. Hierac. 124 = crinitum.
Jacquinii, *Pacher, in Jahrb. Landesm. Kaernt.* xv. (1882) 179.—Europ.
jacobaeifolium, Froel. in DC. Prod. vii. 223 = tridentatum.
Jacquini, Vill. Hist. Pl. Dauph. iii. 128 = humile.
Janka, Boiss. ex Nym. Consp. 441 = olympicum.
Jankae, *Uechtritz. in Oestr. Bot. Zeitschr.* xxiii. (1873) 283.—Europ.
Jansu, Gren. ex Gren. & Godr. Fl. Fr. ii. 378 = murorum.
japonicum, *Franch. & Sav. Enum. Pl. Jap.* ii. 417.—Japon.
Jaubertianum, Bill. ex Nym. Consp. 444 = bifidum.
Jaubertianum, Timb. & Loret, in Bull. Soc. Bot. Fr. v. (1858) 807 = maculosum.
jovanicum, Burm. f. Fl. Ind. 174 = Prenanthes javanica.
Jovanbornati, Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1878-79) 613 = inclusum.
jubatum, *Fries, in Oefvers. Vet. Akad. Forhandl. Stockh.* (1856) 168.—N. Granat.

juncaeum, Berah. Syst. Verz. Erf. 137 = Chondrilla juncea.
junceum, *Fries, Symb. Hierac.* 144.—Mexic.
jurassum, *Fries, l. c.* 129.—Europ. austr.
jurassum, Rapin, Guide Bot. Canton du Vaud, 212 = dactylina.
jurasicum, Grieso. in Goett. Abh. v. (1858) 112 = jurassum.
kajanense, *Malmgr. in Notis. Sällsk. Fauna et Fl. Fenn. Förh.* Ser. II. *Haeft.* iii. (1861) 19, in nota.—Fennia.
kalkbergense, *Dichtl. in Deutsche Bot. Monats.* ii. (1884) 85.—Europ.
Kalmii, Linn. Sp. Pl. 804 (= *Pachylepis Kalmii*).—Am. bor.
Kalmii, Spreng. Syst. iii. 646 = canadense.
Kalmii, Symons, Syn. Pl. 172 = prenanthoides.
Kalsianum, Hutzr, ex Dalla Torre, Atl. Alpenfl. 155 = valdepilosum.
Kladnianum, Schur, Enum. Pl. Transs. 390 = vulgatum.
Knafianum, *Arv. in Bull. Soc. Dauph.* (1880) 284.—Europ.
kobrianum, Gorski, ex Fries, Epicris. Hierac. 23 = pratense.
Kochianum, *Yord. Gr. Yard. Gren.* (1849) 19.—Europ.
Kochii, *Dalla Torre, Atl. Alpenfl.* 148.—Europ.
Kokcilii, *Pacher, in Jahrb. Landesm. Kaernt.* xv. (1882) 174.—Europ.
Kotschyanum, *Heuff. in Flora*, xxxvi. (1853) 618.—Transsylv.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* ii. 418.—Japon.
kravarikense, *Vukot. in Rad Yugos. Akad. Zagreb.* lviii. (1881) 164.—Europ.
laceratum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 406 = vulgatum.
lacerum, *Rouy. ex Fries, Epicris. Hierac.* 86.—Helvetia.
Lachenalii, C. C. Gmel. Fl. Bad. iii. 632 = vulgatum.
Lachenalii, Hort. Landsh. ex Froel. in DC. Prod. vii. 221 = strictum.
laciniatum, Monn. Ess. Hierac. 63 = Andryala ragusina.
laciniatum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 72 = Crepis hyoseridifolia.
laciniosum, Jord. Cat. Gr. Jard. Dijon (1848), 22 = fragile, *Jord.*
Lactaris, Bertol. Amoen. Ital. 64 = umbellatum.
lactucaceum, *Froel. in DC. Prod.* vii. 223.—Ital.
lactucaceum, Schrank, Baier. Fl. ii. 321 = Crepis pratensis.
lactucaefolium, Arv. in Bull. Soc. Dauph. (1880) 289 = lycopsifolium.
Lactucella, Wallr. Sched. Crit. 408 = H. Auricula.
laevicaule, Jord. Cat. Gr. Jard. Dijon (1848), 23 = fragile, *Jord.*
laevigatum, Froel. in DC. Prod. vii. 230 = boreale.
laevigatum, Griseb. in Goett. Abh. v. (1853) 119 = canescens.
laevigatum, Reichb. Fl. Germ. Excurs. 267 = rigidum.
laevigatum, Sch. Bip. Cich. 37 = fasciculare.
laevigatum, Willd. Sp. Pl. iii. 1590 = saxifragum.
Laggeri, Jord. ex Nym. Consp. 446 = tomentosum.
Laggeri, Sch. Bip. ex Nym. l. c. 452 = sabinum.
Lagopus, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 176.—Mexic.
Lamottianum, *Arv. in Lamotte, Prod. Cent. Fr.* ii. 490.—Europ.
lamprophyllum, Schesel, in Linnaea, xxxi. (1861-62) 658 = Neo-Cerinthe.
lampsanoides, Gouan, Illustr. 57 = Crepis lampsanoides.
lampsanoides, Hort. ex DC. Prod. vii. 77 = Lapsana grandiflora.
Lampi, F. Schultz, in Bor. Fl. Cent. Fr. ed. III. ii. 395 = boreale.
lanatum x, Arv. in Bull. Soc. Dauph. (1880) 266. —Europ.
lanatum, Brot. Fl. Lusit. i. 319 = Broteri.
lanatum, Griseb. Spicil. Fl. Rumel. ii. 371 = pannosum.
lanatum, Vill. Prosp. 35; Hist. Pl. Dauph. iii. 120 = tomentosum.
lanatum, *Waldst. & Kit. ex Willd. Sp. Pl.* iii. 1566.—Europ.
lancealatum, Froel. in DC. Prod. vii. 231 = australe.
lanceolatum, Lapeyr. Hist. Abr. Pl. Pyr. 473 = subandum.

HIERACIUM :—
lancelotatum, Royle, ex Hook. f. Fl. Brit. Ind. iii. 400 = umbellatum.
lancelotatum, Schleich. Cat. Pl. Helv. ed. III. (1815) 17 = cymosilanifolium.
lancelotatum, Tausch, ex Fries, Symb. Hierac. 174 = rigidum.
lancelotatum, Vill. Hist. Pl. Dauph. iii. 123. t. 33 = strictum.
lancifolium, Jord. ex Nym. Consp. 438 = boreale.
lancifolium, *Vukot. in Rad Jugos. Akad. Zagreb.* lvii. (1881) 167.—Europ.
Langel, *Prize, Epicris. Hierac.* 57.—Mont. Pyren.
laniferum, *Cav. Ic.* iii. 18. t. 284.—Hispan.
lanigerum, Schur, Enum. Pl. Transs. 391 = murorum.
Lapeyrousii, Bab. in Suppl. Engl. Bot. t. 2916 = iricum.
Lapeyrousii, *Prod. in DC. Prod.* vii. 222.—Mont. Pyren.
lapponicum, *Fries, Symb. Hierac.* 170.—Lappon.
largum, *Fries, Epicris. Hierac.* 127.—Hungar.
Laschianum ×, Thaem. in Bonplandia, vi. (1858) 157. —Europ.
lasiophyllum, Bab. Man. Brit. Bot. ed. IV. 202 = pallidum.
lasiophyllum, Hillebr. ex Nelir. Diagn. Gefaesspfl. 79 = murorum.
lasiophyllum, *Koch, Syn. Fl. Germ.* ed. II. 522.—Europ.
latebrosum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 400 = vulgatum.
lathraeum, *Naeg. & Peter, Hierac.* 262.—Europ.
latibracteum ×, Peter, in Flora, lxiv. (1881) 124.— Europ.
latifolium, *Froel. ex Link, Enum. Hort Berol.* ii. 287. —Rossia; Pyren.
latifolium, Ledeb. Fl. Ross. ii. 854 = foliosum.
latisquamum, *Naeg. & Peter, Hierac.* 212.—Europ.
laurinum, Arv. in Bull. Soc. Dauph. (1880) 291 = lycopsifolium.
Lawerrellei ×, Timb. in Mém. Acad. Sc. Toulouse, Sér. VII. iv. (1877) 186.—Europ.
Lawsoni, Bab. Man. Brit. Bot. ed. II. 196 = anglicum.
Lawsonii, Hiytt, ex Fries, Symb. Hierac. 98 = pallescens.
Lawsonii, Retchb. in Moesl. Handb. ed. II. ii. 1389 = lanatum.
Lawsonii, Sm. Engl. Bot. t. 2088 = glabratum.
Lawsonii, Vill. Prosp. 36; Hist. Pl. Dauph. iii. 118 = saxatile.
laxiflorum, *Wallr. Beitr. Fl. Hercyn.* 283; *et in Linnaea*, xiv. (1840) 658.—Europ.
laxum, (*Cleriev.*) *Man. Herb.* 281.—Europ.
Lazergui, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875–76) 600 = hirsutum.
laticum, *Soies. & Bal. ex Boiss. Fl. Orient.* iii. 878. —As. Min.
Lecokianum, Arv. in Lamotte, Prod. ii. 485 = vulgatum.
Ledebourii, Gerh. ex Froel. in DC. Prod. vii. 224 = umbellatum, lapponicum.
Ledebourii, Mus. Petrop. ex Fries, Symb. Hierac. 178 = umbellatum.
Ledebourii, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 67.—Rossia.
legionense, Cosa. ex Willk. & Lange, Prod. Fl. Hisp. ii. 307 = Bourgaei.
Legrandianum, Arv. in Bull. Soc. Bot. Fr. xxxi. (1884) 186 = Neo-Cerinthe.
leiocephalum, Bartl. ex Griseb. in Goett. Abh. v. (1858) 102 = bupleuroides.
leiocephalum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 48 = porphyritifolium?
leiopogon, Gren. ex Verlot, in Bull. Soc. Stat. Isère, xiv. (1873) 396 = saxetanum.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1884) 70.—Arizona.
Leontodontis, Linn. Fasc. Suec. ii. App. 558 = Leontodon Taraxaci.
leptocephalum, *Benth. Pl. Hartw.* 137.—Peruv.
leptocephalum, Fries, ex Nym. Consp. 442 = juranum.
leptocephalum, Vukot. in Rad Jugos. Akad. Zagreb. xxxiv. (1876) 19 = rotundatum.
leptocladon, *Naeg. & Peter, Hierac.* 675.—Europ.
leptophyton, *Naeg. & Peter, l. c.* 642.—Europ.
leptopogon, Rhenib. ex Fries, Epicris. Hierac. 158 = Pieris Spomuriana.
Lesserianum, Wall. Cat. n. 2354 = Lactuca Lessertiana.
leucanthemum, *Wedd. Chlor. And.* i. 235.—N. Granat.

HIERACIUM :—
leucocephalum, *Rupr. ex Kaufm. Ind. Sem. Hort. Mosq.* (1868) 9.—Hab.?
leucocephalum, Schur, Enum. Pl. Transs. 225, Index, sphalm., vide *leucophylium*.
leucocephalum, Vukot. ex Nym. Consp. 454 = H. Pilosella.
leucopappum, Steud. Nom. ed. I. 407 = Crepis grandiflora.
leucophaeum, Gren. & Godr. Fl. Fr. ii. 354 = glaucum.
leucophyllum, Schur, Enum. Pl. Transs. 379 = H. Pilosella.
leucopogon, Gren. ex Bull. Soc. Bot. Fr. xix. (1872), Bibl. 87 = saxetanum.
leucotrichum, *Fries, in Oefver. Vet. Akad. Forhandl.* Stockh. (1856) 144.—Ross.
libanoticum, *Boiss. & Bal. ex Boiss. Fl. Orient.* iii. 870.—Syria.
ligusticum, De Not. ex Nym. Consp. 442 = crinitum.
ligusticum, *Fries, Symb. Hierac.* 74.—Europ.
Lindebergii, Nym. Consp. 445 = canescens.
linearc, Steud. Nom. ed. I. 407 = umbellatum.
linearifolium, Froel. ex Steud. Nom. ed. II. i. 764 = graminifolium.
linearifolium, Monn. Ess. Hierac. 64 = Andrysia rupestris.
Lingrenfeldii, Sch. Bip. ex Nym. Consp. 444 = murorum.
lingulatum, *Backh. Mon. Hierac.* 30.—Brit.
*lini*folium, Saelan, ex Lindeb. in Blytt, Norges Fl. ii. 662 = lapponicum.
Linnaeanum, Jord. ex Nym. Consp. 444 = murorum.
Liottardi, Koch, Syn. Fl. Germ. ed. I. 458 = Kochiana.
Liottardi, *Vill. Prosp.* 35; *Hist. Pl. Dauph.* iii. 121. —Gallia.
lividum, Arv. in Bull. Soc. Dauph. (1880) 283 = Neo-Cerinthe.
lixarense, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875–76) 588 = cerinthoides.
longifolium, *Hornem. Hort. Hafn. Suppl.* 155.—Hab.?
longifolium, Schleich. Cat. Pl. Helv. ed. III. (1815) 17 = cerinthoides.
longipes, C. Koch, in Linnaea, xxiii. (1850) 680 = praealtum.
longipetalum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 72 = Crepis grandiflora.
longipilum, *Torr. ex Hook. Fl. Bor. Am* i. 298.—Am. bor.
longiscapum, Bois. Fl. Orient. iii. 864 = sabinum.
Loreti, Fries, ex Nym. Consp. 450 = villosum.
Lortetiae, *Balb. Fl. Lyonn.* i. 1. 450.—Mons Pilatus.
Lucaeinianum, Schrenk, in Linnaea, xxxii. (1863) 668 = cerinthoides.
lozenae, *Benth. Pl. Hartw.* 137.—Ecuador.
luciferum, *Soies. ind. Sem. Hort. Boccad.* (1922) 6; *Fl. Sic. Syn.* ii. 405.—Sicilia.
lucorum, Schur, Enum. Pl. Transs. 399 = boreale.
luridum, Jord. ex Nym. Consp. 438 = boreale.
lutescens, Ex Favre, in Bull. Soc. Murith. v. (1875) 134 = picroides.
lychnioides, Arv. in Bull. Soc. Dauph. (1880) 286 = lycopsifolium.
Lychnitis, *Scheele, in Linnaea*, xxxi. (1861–62) 668.— Hispan.
lycopalifolium, *Froel. in DC. Prod.* vii. 224.—German.
lycopsifolium, Vill. Hist. Pl. Dauph. i. 28 = humile.
lyratum, Linn. Sp. Pl. 803 = Crepis lyrata.
lyratum, Schleich. ex Froel. in DC. Prod. vii. 170 = Crepis hyoseridifolia.
macedonicum, *Boiss. & Orph. ex Boiss. Fl. Orient.* iii. 872.—Macedon.
macilentum, *Fries, in Oefver. Vet. Akad. Forhandl. Stockh.* (1856) 147.—Helvetia.
macranthelium, *Naeg. & Peter, Hierac.* 473.—Europ.
macranthum, Lindeb. in Hartm. Handb. Skand. Fl. ed. X. 29 = norvegicum.
macranthum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 446 = umbellatum.
macranthum, Tenore, Fl. Nap. v. 190 = H. Pilosella.
macrocephalum, Huter, ex Dalla Torre, Atl. Alpenfl. 157 = picroides.
macrolepis, *Boiss. Fl. Orient.* iii. 878.—Reg. Caucas.
macrophyllum, Pursh, Fl. Am. Sept. ii. 504 = canadense.
macrophyllum, Scheele, in Linnaea, xxxii. (1863) 674 = Neo-Cerinthe.
macrophyllum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 69 = Crepis sibirica.

HIERACIUM :—
macrothyrsum ×, Naeg. & Peter, Hierac. 763.— Europ.
macrotrichum, *Boiss. Diagn. Ser. I. iv.* 29.—As. Min.
maculatum, Blas. ex Nym. Consp. 444 = murorum.
maculatum, *Schrank, Baier. Fl.* ii. 319.—Europ.
maculatum, Tausch, in Flora, xi. (1828) 1. Erg. 70 = rigidum.
maculatum, Tenedale, ex Turner & Dillwyn, Bot. Guide, 707 = vulgatum.
maculiparum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 408 = vulgatum.
madagascaricum, *DC. ex Froel. in DC. Prod.* vii. 218 —Madag.
magellanicum, *Sch. Bip. in Flora*, xxxvii. (1855) 122 —Reg. Magell.
magistri, Godr. ex Nym. Consp. 441 = tridentatum.
magyaricum, *Naeg. & Peter, Hierac.* 566.—Europ.
Mairanum, Mancel, ex Nym. Consp. 453 = subnivale
majellense, Huet, ex Nym. l. c. 449 = scorsonerae-folium, *Vill.*
marginatum, *Froel. in DC. Prod.* vii. 220.—Hab.?
marianum, *Willd. Sp. Pl.* iii. 1572.—Am. bor.
mariolenae, *Kouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 122.—Hispan.
murmorum, Janka, ex Nym. Consp. 446 = Jankae.
marmorum, Pant. & Vis. in Mem. Ist. Venet. xii. (1864) 468 = tomentosum.
Marschallianum, *Andrz. Rys. Bot.* ii. (1830) 27.— Podol.; Ucrania.
mauritianum, Willd. ex Froel. in DC. Prod. vii. 104 = Crepis mauritiana.
medullegense, *Dichtl. in Deutsche Bot. Monatsschr.* ii. (1884) 58.—Europ.
medioximum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 401 = vulgatum.
medium, Jord. Gr. Jard. Gren. (1849) 19 = murorum.
Medusae, *Retchb. f. in Bot. Zeit.* xx. (1862) 61 — Rumelia.
melanchaetum, Tausch, in Flora, xi. (1828) 1. Erg. 58 = praealtum.
melanistum ×, Naeg. & Peter, Hierac. 664.—Europ.
melanocadenum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 75 = murorum.
melanocephalum, Lindeb. in Bot. Notiser (1878) 184, nomen; et in Hartm. Handb. Skand. Fl. ed. XI. 48 = rigidum.
melanocephalum, Tausch, in Flora, xi. (1828) 1. Erg. 68 = alpinum.
melanotrichum, *Kern. in Oestr. Bot. Zeitschr.* xxiv. (1874) 170.—Europ.
melanotrychum, *Reut. in Compt.-Rend. Soc. Hallér.* (1852–53) 5; (1853–54) 25.—Europ.
mendax ×, Naeg. & Peter, Hierac. 679.—Europ.
Mertini, C. C. Gmel. Fl. Bad. iv. 891 = vulgatum.
mespilifolium, *Arv. in Bull. Soc. Dauph.* (1880) 282. —Europ.
mexicanum, *Less. in Linnaea*, v. (1830) 132.—Mexic.
Michelii, Tausch, in Flora, xi. (1828) 1. Erg. 60 = pilosellioides.
microdeuton, *Fries, Epicris. Hierac.* 149.—N. Granat.
micranthum, Huet, ex Nym. Consp. 453 = serrylli-folium.
microcephalum, Fries, Symb. Hierac. 174 = rigidum.
microcephalum, *Sch. Bip. in Bonplandia*, iv. (1856) 55. nomen.—Peruv.
micromegas, Fries, Epicris. Hierac. 25 = petraeum.
microtum, *Boiss. Fl. Orient.* iii. 878.—As. Min.
Milleri, *Link, Enum. Hort. Berol.* ii. 287.—Hab.?
minutiflorum, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, iii. (1875–76) 615, nomen.—Europ.
minutum, Jeanb. & Timb. l. c. 609 = anglicum.
mirabile, *Naeg. & Peter, Hierac.* 561.—Europ.
mixtum, *Lapeyr. ex Froel. in DC. Prod.* vii. 216.— Mont. Pyren.
mnoophorum, *Naeg. & Peter, Hierac.* 767.—Europ.
Molanum, Lindeb. in Bot. Notiser (1872) 118 = murorum.
monticolum, Kern. ex Nym. Consp. 441 = olympicum.
molle, Jacq. Fl. Austr. ii. 12 = Crepis succisaefolia.
molle, Lachenal, in Nov. Act. Helv. i. (1787) 293 = juranum.
molle, Pall. Tabl. Taur. 57 = echioides.
molle, Pursh, Fl. Am. Sept. ii. 508 = vulgatum.
mollitum, *Arv. in Bull. Soc. Dauph.* (1880) 282 — Europ.
monocense ×, F. Schultz, Arch. Fl. 5.—Europ.

HIERACIUM:—
monasteriale ×, Naeg. & Peter, Hierac. 501.—Europ.
montanum, Jacq. Fl. Austr. ii. 54 = Crepis montana.
montanum, Scop. Fl. Carn. ed. II. ii. 106 = Crepis incarnata.
montenegrinum, Pant. ex Nym. Consp. 446 = lanatum.
monticola, Jord. Gr. Jard. Gren. (1849) 30 = umbellatum.
montolearense, *Timb. & Jeanb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, ii. (1874) 247.—Europ.
moravicum, *Fryn. ex Oborny, in Verh. Naturf. Ver. Brunn.* xxii. 11. (1884) 590.—Europ.
Moriisianum, *Reichb. f. Ic. Fl. Germ.* xix. 67.—Italia.
Moritzianum, *Hegetschw. Fl. Schw.* 781.—Helvetia.
mosanum, Crép. Man. Fl. Belg. 138 = pallidum.
Mougeoti, Frœl. ex Koch, Syn. Fl. Germ. ed. I. 453 = cerinthoides.
Mougeoti, Godr. Fl. Lorr. ed. I. ii. 77 = decipiens.
mucronulatum, *Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, ii. (1874) 256.—Europ.
multicaule, Schaffn. ex Fries, Epicris. Hierac. 149 = mexicanum.
multiflorum, Gilib. Fl. Lituan. i. 236 = sabaudum.
multiflorum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 421 = H. Auricula.
multiflorum, Schleich. Cat. Pl. Helv. ed. III. 17 = sabinum.
murale, Salisb. Prod. 181 = murorum.
muricollum, *Fries, Epicris. Hierac.* 117.—Reg. Caucas. Muribianum, *Favrat, ex Payot, in Bull. Soc. Bot. Fr.* xxix. (1882) *Bibl.* 43.—Europ.
murorum, C. B. Clarke, Comp. Ind. 956 = vulgatum.
murorum, Frœl. in DC. Prod. vii. 213 = caesium.
murorum, *Linn. Sp. Pl.* 800.—Reg. bor. temp.
mycnotifolium, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Chili.
myriadenum, *Boiss. & Reut. ex Reichb. fe. Fl. Germ.* xix. 58.—Hispan.
myriantlium, Scheele, ex Nym. Consp. 438 = hirsutum.
myriophorum, Scheele, ex Nym. l. c. 448 = Pseudo-Cerinthe.
myriocephalum, *Panč. Elench. Crna-Gora*, 61.—Europ.
myriophyllum, *Scheele, in Linnæa*, xxvii. (1853) 660. —Hispan.
Naegelianum, *Panč. Elench. Crna-tora*, 57.—Europ.
naevuliferum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 404 = vulgatum.
nanum, Scheele, in Linnæa, xxxi. (1861–62) 648 = H. Auricula.
nanum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 70 = Crepis chrysantha.
neapolitanum, Huet, ex Nym. Consp. 450 = villosum.
neglectum, *Norrl. ex Wainio, in Meddel. Soc. Faun. et Fl. Fenn.* iii. (1878) 62.—Fennia.
nemogetum, *Jord. ex Pérard, in Bull. Soc. Bot. Fr.* xvii. (1870) 121.—Europ.
nemophilum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 396 = vulgatum.
nemorale, Jord. ex Nym. Consp. 443 = vulgatum.
nemorense, Jord. Cat. Gr. Jard. Dijon (1848) 23 = murorum.
nemorivagum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 385, in obs. = boreale.
nemorosum, Bess. Enum. Pl. Volh. 81 = vulgatum.
nemorosum, Dierb. Fl. Heidelb. ii. 262 = boreale.
nemorosum, Lapeyr. ex Frœl. in DC. Prod. vii. 223 = tridentatum.
nemorosum, Stend. Nom. ed. I. 468 = ramosum.
nemorum, Jord. ex Nym. Consp. 449 = decipiens.
nemorum, Nym. l. c. 444, in syn. = caesium.
Neo-Cerinthe, *Fries, Symb. Hierac.* 67.—Mont. Pyren.
Nestleri, *Koch, Syn. Fl. Germ.* ed. I. 449.—Europ. centr.
Nestleri, Lasch. Fl. Ross. ii. 849 = sabinum.
Nestleri, Vill. Voy. 63. t. 4. f. 1 = cynoxum.
nigrescens, *Willd. Sp. Pl.* iii. 1574.—Silesia.
nigrescens, Wimm. & Grab. Fl. Siles. ii. ii. 122 = stratum.
nigrinum, *Meinsh. Fl. Ingr.* 198.—Ingria.
nigricans, Schur, Enum. Pl. Transs. 397 = stratum.
nigrisarinum, *Naeg. & Peter, Hierac.* 362.—Europ.
nigrispa, *Lindeb. in Bot. Notiser* (1865),137.—Norveg.
nigroglandulosum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 175 = murorum.
nigrobilum, *Naeg. & Peter, Hierac.* 204.—Europ.
nitens, *Lindeb. in Bot. Notiser* (1872), 119.—Norveg. occ.

HIERACIUM:—
nitidum, Backh. Mon. Hierac. 59 = pallidum.
nitidum, Scheele, in Linnæa, xxxii. (1863) 655 = amplexicaule.
nivale, *Froel. in DC. Prod.* vii. 221.—Tyrol.
nivocappum, *Fries, Symb. Hierac.* 129.—Mexic.
nobile, Gren. & Godr. Fl. Fr. ii. 376 = pyrenaicum.
Norrlini, *Naeg. & Peter, Hierac.* 475.—Europ.
norvegicum, *Fries, Symb. Hierac.* 109.—Norveg.
Notarisii, Reichb. f. Ic. Fl. Germ. xix. 59 = cinitum.
nothagenicum ×, Naeg. & Peter, Hierac. 883.—Europ.
nudicaule, Stend. Nom. ed. I. 468 = Grenovii.
Oberleitneri, Sch. Bip. ex Nym. Consp. 447 = humile.
obliquum, Jord. Cat. Gr. Jard. Dijon (1848) 23 = boreale.
oblongifolium, Schur, in Oestr. Bot. Zeitschr. xi. (1861) 350 = rotundatum.
oblongifolium, Tausch, in Flora, xi. (1828) 1. Erg. 54 = dubium.
oblongum, Jord. Gr. Jard. Gren. (1849) 20 = caesium.
Obornyanum, *Naeg. & Peter, Hierac.* 711.—Europ.
obovatum, Lapeyr. Suppl. Hist. Pl. Pyr. 129 = Neo-Cerinthe.
obscurum, Lang, in Syll. Ratisb. i. (1824) 184 = brachiatum.
obscurum, Reichb. Pl. Crit. i. 92 = praealtum.
obscurum, Zollik. ex Gaud. Fl. Helv. v. 98 = villosum.
obtusifolium, Backh. Monog. Hierac. 56 = caesium.
obtusifolium, Stend. Nom. ed. II. i. 764 = murorum.
occitanicum, Jord. Obs. Pl. Crit. vii. (1849) 36 = boreale.
ochroleucum, Pané. ex Nym. Consp. 455 = Crepis viscidula.
ochroleucum, Schleich. Cat. Pl. Helv. ed. IV. (1821) 19 = glaucophaloides.
ocnodes ×, Naeg. & Peter, Hierac. 784.—Europ.
officinale, Bernh. Syst. Verz. Erf. 138 = Taraxacum officinale.
oligocladum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 414 = murorum.
oleosorum, Arv. in Bull. Soc. Dauph. (1880) 287 = olivarum.
oleraceum, Scop. Fl. Carn. ed. II. ii. 110 = Sonchus oleraceus.
oligocephalum, *Arv. in Bull. Soc. Dauph.* (1880) 286. —Europ.
olivaceum, *Gren. & Godr. Fl. Fr.* ii. 361.—Mont. Pyren.
olympicum, *Boiss. Diagn.* Ser. I. iv. 80.—Graecia; Bithyn.
onosmoides, *Fries, Symb. Hierac.* 102.—Norveg.
operinodes, Sch. Bip. ex Nym. Consp. 451 = bupleuroides.
opposititolium, Vukot. ex Nym. l. c. 439 = racemosum.
orarium, Lindeb. in Bot. Notiser (1872) 121 = dovrense.
oreades, *Fries, Symb. Hierac.* 100.—Suecia.
oreades, Heuff. in Flora, xxxvi. (1853) 617 = petraeum.
oreades, Wimm. ex Uechtr. in Oestr. Bot. Zeitschr. xxii. (1872) 277 = pallescens.
oreophilum, Heuff. ex Nym. Consp. 452 = petraeum.
Orini, Gren. in Oestr. Bot. Zeitschr. xxiv. (1874) 170 = lanatum.
orientale, Fries, Epicris. Hierac. 74 = pannosum.
originifolium, Frœl. in DC. Prod. vii. 233 = laniferum.
Orphanidis, Boiss. Fl. Orient. iii. 861 = pannosum.
orthotrichum, *Fries, Epicris. Hierac.* 146.—Chili.
ovalifolium, Jord. Obs. Pl. Crit. vii. (1849) 33 = murorum.
ovalifolium, Frœl. in DC. Prod. vii. 234, in syn. = umbellatum.
ovalifolium, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 76 = saxatile.
ovatum, Hort. Vind. ex Frœl. in DC. Prod. vii. 230; Schleich. ex Stend. Nom. ed. I. 468 = lasiophyllum, pallidum.
ovatum, Hull, Brit. Fl. 175 = villosum.
oxydon, *Fries, Epicris. Hierac.* 90.—Helvetia.
oxyephilum, *Rehm. in Oestr. Bot. Zeitschr.* xxiii. (1878) 110.—Europ.
Oxononis, F. Schultz, Arch. Fl. 262 = scorzonerifolium, *Vill.*
Pachuri, *Sch. Bip. ex Pacher, in Jahrb. Landesm. Kornt.* xv. (1862) 188.—Europ.
pachylodes, *Naeg. & Peter, Hierac.* 180.—Europ.
pallescens, Pană in Abh. Zool.-Bot. Ver. Wien, vi. (1856) 961 = Schultzianum.
pallescens, Sch. Bip. ex Nym. Consp. 446 = murorum.

HIERACIUM:—
pallescens, *Waldst. & Kit. Pl. Rar. Hung.* iii. 241. t. 217.—Europ. centr.
pallescens, Wittg. ex Nym. Consp. 445 = saxifragum.
pallidiflorum, Jord. ex Nym. l. c. 441 = picroides.
pallidiflorum, Reichb. Ic. Fl. Germ. xix. 74 = vulgatum.
pallidifolium, Jord. in Bor. Fl. Centr. Fr. ed. III. ii. 407 = vulgatum.
pallidifolium, Knaf, in Oestr. Bot. Zeitschr. xxii. (1872) 77 = bohemicum.
pallidiformi, Arv. Suppl. Mon. Hierac. 20 = pallidum.
pallidulum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 419 = murorum.
pallidum, *Bie. ex Guss. Fl. Sic. Syn.* ii. 404.—Norveg.
pallidum, Fries, ex Nym. Consp. 444 = bifidum.
pallidum, Lange, in Kjoeb. Videnak. Meddel. (1861) 104 = vapicalum.
pallidum, Tausch, in Flora, xi. (1828) 1. Erg. 67 = murorum.
pallidum, Willd. ex Frœl. in DC. Prod. vii. 231 = pallescens.
paludosum, Linn. Sp. Pl. 808 = Crepis paludosa.
Pamphilii, Arv. in Bull. Soc. Dauph. (1880) 282 = scorzonerifolium, *Vill.*
Panicii, Sch. Bip. ex Vis. in Mem. Ist. Venet. xv. (1870) 9 = Sartorianum.
panduriforme, Dulac, Fl. Hautes-Pyr. 486 = prenanthoides.
panduriforme, Timb. in Bull. Soc. Bot. Fr. xviii. (1871) 315 = neo-cerinthe.
paniculatum, Gilib. Fl. Lituan. i. 236 = Crepis paludosa.
paniculatum, Gueldenst. Reisen, i. 68 (Quid ?).—Ross. austr.
paniculatum, Hofm. ex Frœl. in DC. Prod. vii. 220 = saxetanum.
paniculatum, Jacq. Coll. Suppl. 148 = Crepis rigida.
paniculatum, *Linn. Sp. Pl.* 803.—Am. bor.
pannonicum, Jacq. Coll. v. 148 = Crepis rigida.
pannonicum, *Naeg. & Peter, Hierac.* 749.—Europ.
pannosum, *Boiss. Diagn.* Ser. I. iv. 82.—Graecia; Ar. Min.
paniculiflorum, *Naeg. & Peter, Hierac.* 662.—Europ.
Papperitzi, Reichb. f. Ic. Fl. Germ. xix. 99 = stupposum.
pappoleucum, Vill. Prosp. 36; Hist. Pl. Dauph. iii. 134 = Crepis grandiflora.
papyraceum, Sch. Bip. ex Nym. Consp. 440 = prenanthoides.
paradoxum, Nees, in Flora, v. (1822) 189 = dubium.
paradoxum, Schult. Obs. Bot. 164 = pratense.
paraguayum, *Naeg. & Peter, Hierac.* 658.—Europ.
paraguense, *Baker, in Mart. Fl. Bras.* vi. 111. 337.—Paraguay.
Parishi, Heuff. in Flora, xxxvi. (1853) 618 = piloselloides.
Parishii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 67. —Calif.
Parnassi, *Fries, Symb. Hierac.* 86.—Graecia.
partitum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 407 = vulgatum.
parviflorum, Schleich. Cat. Pl. Helv. ed. III. 17 = Crepis incarnata.
parviflorum, Stend. Nom. ed. II. i. 765 = sphaerocephalum.
parvifolium, Link, Enum. Hort. Berol. ii. 385, sphalm. = porrifolium.
parvulum, Tenore, ex Frœl. in DC. Prod. vii. 167 = Crepis Columnae.
patagonicum, *Hook. f. Fl. Antarct.* ii. 324.—Patagon.
patens, Berl. Ind. Sem. Hort. Gotting. (1850) 4.—Cf. Linnæa, xxiv. (1851) 207 = australe?
patulipes, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 409 = murorum.
patulum, Tausch, in Flora, xi. (1828) 1. Erg. 73 = prostratum.
paucifloram, Bernh. Syst. Verz. Erf. 138 = Lactuca muralis.
paucifoliatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 400 = vulgatum.
paucinucamum, Jord. in Bor. l. c. 403 = vulgatum.
Pavichii, Heuff. ex Nym. Consp. 451 = piloselloides.
pedunculare, Tausch, in Flora, xi. (1828) 1. Erg. 76 = sudeticum.
pedunculare, Wallr. ex Nym. Consp. 454 = H. Pilosella.
pedunculatum, Wallr. Sched. Crit. 406 = H. Pilosella.
Pelsterianum, Mérat, Nouv. Fl. Par. 306 = H. Pilosella.
pellophyllum, *Schur, Enum. Pl. Transs.* 387.—Transylv.

HIERACIUM:—
pellitum, *Fries, Epicris. Hierac.* 79.—Europ.
pellucidum, Wahlenb. Fl. Suec. ii. 514 = vulgatum.
pennsylvanicum, Fries, Symb. Hierac. 160 = Gronovii.
pentagenes × , Naeg. & Peter, Hierac. 358.—Europ.
pentaphylum × , Naeg. & Peter, l. c. 784.—Europ.
perclarum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 400 = vulgatum.
perfoliatum, Fresl. in DC. Prod. vii. 211 = prenanthoides.
permutatum, *Naeg. & Peter, Hierac.* 366.—Europ.
Perreymondi, *Jord. Obs. Fl. Crit.* vii. (1849) 42, in obs.—Gallia.
persecifolium, Fries, Symb. Hierac. 95 = pallidum.
persicum, Boiss. Diagn. Ser. I. xf. 60 = sablnum.
personatum, Fries, Epicris. Hierac. 45 = sudeticum.
peruanum, *Fries, Symb. Hierac.* 156.—Peruv.
Petrasinum, *Timb. in Bull. Soc. Bot. Fr.* xviii. (1871) 313.—Mont. Pyren.
petrogrum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 386 = umbellatum.
pestivense × , Simk. in Termesz. Füzetek, ii. (1878) 81.—Europ.
petiolare, Jord. Gr. Jard. Gren. (1849) 20 = fragile, *Jord.*
petraeum, *Friwald. in Flora*, xix. (1836) 436.—Macedon; Serv.
petraeum, Hoppe, ex Bluff & Fingerh. Comp. Fl. Germ. ii. 396 = amplexicaule.
petrogenes, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 418 = murorum.
petrophilum, Godet, ex Nym. Consp. 449 = scorzonerae-folium, *Vill.*
petrophilum, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, vi. (1883–84) 214.—Europ.
petrophilum, Schur, Enum. Pl. Transs. 392 = bifidum.
petroum, Hoppe, ex Nym. Consp. 448 = pulmonarioides.
petraum, Schur, Enum. Pl. Transs. 925 = stratum.
phlomoides, *Froel. in DC. Prod.* vii. 283.—Mont. Pyren.
Picdari, Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 170 = lanatum.
picridifolium, *Schleich. Cat. Pl. Helv.* ed. IV. 19, nomen. —Helvet.
picroides, Gaud. Fl. Helv. v. 115 = cydoniaefolium.
picroides, *Vill. Voy.* 22. t. 1. f. 3.—Helv.; Gall.
pictoriense, Sauré & Maill. in F. Schultz, Arch. Fl. Fr. et Allem. 385 = australe.
pictum, Jord. ex Nym. Consp. 444 = caesium.
picrum, *Pers. Syn* ii. 374 = saxatile.
pictum, Schleich. ex Hoppen. Hort. Hafn. 763 = humile.
picturatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 405 = vulgatum.
pienikense, *Rehm. in Oestr. Bot. Zeitschr.* xxiii. (1873) 110.—Europ.
pilatrae, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 407 = vulgatum.
pilliferum, *Hoppe, ex Fries, Symb. Hierac.* 49.—Europ.
pilipes, *Saelan, in Medd. Soc. Faun. et Fl. Fenn.* (1881) 188.—Europ.
pilocephalum, Link, Enum. Hort. Berol. ii. 286 = villosum.
Pilosella, *Linn. Sp. Pl.* 800.—Europ.; Afr. bor.
pilosellaeforme, Hoppe, in Sturm, Deutschl. Fl. Heft 87 = H. Pilosella.
pilosellifolium, *Naeg. & Peter, Hierac.* 707.—Europ.
pilosellinum, Sch. Bip. ex Nym. Consp. 454 = H. Pilosella.
piloselloides, Hartm. Handb. Skand. Fl. ed. III. 186 = hyperboreum.
pilosiloides, *Vill. Prosp.* 84; *Hist. Pl. Dauph.* iii. 100. t. 97.—Europ.
pilosiloides, Wallr. Sched. Crit. 410 = praealtum.
piloisissimum, *Friwald. in Flora*, xix. (1836) 436.—Rumel.
piloisissimum, Schrank, Beitr. Fl. ii. 318 = murorum.
pilosulum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 416 = murorum.
pilosum, Schleich. Cat. Pl. Helv. ed. III. 17 = dentatum.
pilosum, Willd. ex Stend. Nom. ed. II. i. 765 = Crepis blattarioides.
pinetorum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 414 = murorum.
pinetorum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 69 = Crepis chondrilloides.
pinnatifidum, Willd. Sp. Pl. iii. 1560 = Crepis pinnatifida.

HIERACIUM:—
pistoriense, *Naeg. & Peter, Hierac.* 601.—Europ.
Planchonianum, Timb. & Loret, in Bull. Soc. Bot. Fr. v. (1858) 508 = bifidum.
pleicephalum, *Schur, in Verh. Siebenb. Ver. Naturw.* ii. (1851) 171.—Transylv.
plericephalum, Amb. Fl. Tir. Mer. ii. 614 = brachiatum.
plerophyllum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 46 = rotundatum.
pleiatocephalum, *Baker, in Mart. Fl. Bras.* vi. III. 338. —Bras.
plicatum, Lindeb. in Bot. Notiser (1872) 121 = dovrense.
plumbeum, Fries, Symb. Hierac. 111 = bifidum.
plumosum, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 179 = villosum.
plumulosum, Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 170 = lanatum.
poliocephalum, *Naeg. & Peter, Hierac.* 277.—Europ.
poliotrichum, Wimm. ex Nym. Consp. 452 = cymosum.
pollinum, *Fries, Symb. Hierac.* 84.—Europ. austr.
pollosphaerum × , Naeg. & Peter, Hierac. 766.—Europ.
Pollichiae, Sch. Bip. & G. F. Koch, in Oestr. Bot. Wochenbl. iii. (1853) 247 = murorum.
polyanthemum, Vill. ex Steud. Nom. ed. II. i. 765 = alpinum.
polycephalum, *Velen. in Oestr. Bot. Zeitschr.* xxxiii. (1883) 390.—Europ.
polycladium × , Link. in Arch. Naturw. Landesk. Boehm. iv. (1881) 786.—Europ.
polycladon, Schur, Enum. Pl. Transs. 381 = *bifurcum*.
polycladum, Juratz. in Oestr. Bot. Wochenbl. vii. (1857) 58.
polynothum, *Naeg. & Peter, Hierac.* 281.—Europ.
polyodon, *Fries, Epicris. Hierac.* 67.—Afr. austr.
polyphyllum, *Bernh. ex Hornem. Hort. Hafn.* 765.—Hab.?
polyphyllum, Hort. Pesth. ex Prod. in DC. Prod. vii. 219.—speciosum.
polyphyllum, Schleich. Cat. Pl. Helv. ed. III. 17 = graminifolium.
polyphyllum, Willd. Enum. Hort. Berol. Suppl. 54 = bupleuroides.
polyschistum × , Naeg. & Peter. Hierac. 686.—Europ.
polytrichum, Ledeb. Fl. Alt. iv. 130 = Crepis polytricha.
polytrichum × , Naeg. & Peter. Hierac. 705.—Europ.
polytrichum, Schur, Enum. Pl. Transs. 394 = rotundatum.
pontanum, J. F. Gmel. Syst. 1177 = Crepis montana.
porphyritae, *F. W. Schultz, in Oestr. Bot. Zeitschr.* xxii. (1872) 310.—Europ.
porphyritacum, *Kern. l. c.* xiii. (1868) 247.—Europ.
porrectum, Christ, ex Nym. Consp. 450 = dentatum.
portenschlagiae, *Froel. Symb. Hierac.* 106.—Europ. austr.
porrifolium, Balb. ex Froel. in DC. Prod. vii. 220 = marginatum.
porrifolium, *Linn. Sp. Pl.* 802.—Austr.; Ital.
porrifolium, Vill. Hist. Pl. Dauph. iii. 115 = gleucum.
Pourretianum, *Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, ii. (1874) 66.—Europ.
praealtum, *Gaeke. Tent. Med.-Bot. Cich.* 17; *Vill. Voy.* 22. t. 2. f. 1.—Europ.; Oriens.
praealtum, Reichb. Pl. Crit. i. 45 = setigerum, *Tausch.*
praecelsum, *Rov. in Soc. Dauph. Exch. Pl.* iv. (1877) 116.—Europ.
praecox, Sch. Bip. in Pollichia, ix. (1851) 35, 44 = murorum.
praecox, Tausch, ex Nym. Consp. 451 = praealtum.
praecurrens, *Vukot. in Rad Jugos. Akad. Zagreb.* lviii. (1881) 164.—Europ.
praelongum, *Lindeb. in Bot. Notiser* (1866) 78.—Europ.
praemorsiforme, *Sch. Bip. in Bonplandia*, ix. (1861) 174, 237.—Mexic.
praemorsum, All. Fl. Pedem. i. 213 = cymosum.
praemorsum, Linn. Sp. Pl. 801 = Crepis praemorsa.
praemorsum, Gren. & Godr. ex Nym. Consp. 440 = H. Expatriatum.
praeniflorum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 418 = murorum.
praesanum, *Dum. Fl. Belg. Prod.* 62.—Belgium.
pratense, *Tausch, in Flora*, xl. (1838) i. *Erg.* 56.—Europ.; As. bor.
prenanthoides, Doell, Rhein. Fl. 528 = lycopsifolium.
prenanthoides, Hook. Fl. Bor. Am. i. 269 = canadense.
prenanthoides, Lej. Rev. Fl. Spa, 168 = sabaudum.
prenanthoides, Sm. Fl. Brit. 836; Engl. Fl. iii. 368; Lieben. in Fl. Dan. t. 2425 = strictum.
prenanthoides, *Vill. Prosp.* 85; *Hist. Pl. Dauph.* iii. 108.—Europ.; Reg. Caucas. et Himal.

HIERACIUM:—
primulaefolium, *Vis. Fl. Lib. Spec.* 68.—Sardinia.
Pringlei, *A. Gray, in Proc. Am. Acad.* xix. (1883) 69. —Arizona.
procerum, *Fries, Symb. Hierac.* 43.—Reg. Caucas.
proliferum, Hartm. Svensk Excurs. Fl. 113 = vulgatum.
prostratum, DC. Rapp. Voy. 78; *ex* ej. Fl. Fr. vi. 437 = eriophorum.
prostratum, Panz. ex Froel. in DC. Prod. vii. 225 = boreale.
prostratum, *Poir. Encyc. Suppl.* ii. 560.—Europ.
protractum, Fries, ex Nym. Consp. 442 = dovrense.
provinciale, Jord. Obs. Pl. Crit. viii. (1849) 41 = australe.
provincuale, F. Schultz, ex Nym. Consp. 442 = crinitum.
prunellaefolium, Gouan, Illustr. 67 = Crepis pygmaea.
pruscicum × , Naeg. & Peter, Hierac. 378.—Europ.
Pseud-albinum, *Uechtr.—Cf. Oestr. Bot. Zeitschr.* xxxv. (1885) 366, nomen.—Europ.
Pseud-effusum × , Naeg. & Peter, Hierac. 595.—Europ.
Pseud-eriophorum, Loret & Timb. in Bull. Soc. Bot. Fr. v. (1858) 616 = hirsutum.
Pseudo-Auricula, Schur, Enum. Pl. Transs. 382 = praealtum.
Pseudo-Auricula, Tausch, in Flora, vi. (1823) i. Erg. 53 = bracliatum, praetense.
Pseudo-bifidum, Schur, Enum. Pl. Transs. 392 = bifidophyllum.
Pseudo-bifurcum, *Porc. in Magyar Novényt. Lapok*, ix. (1885) 129, nomen.—Europ.
Pseudo-Blyttii, Norrl. in Notis. Sällsk. Faun. et Fl. Fenn. (1874) 437 = flammeum.
Pseudo-caloxon × , Naeg. & Peter, Hierac. 748.—Europ.
Pseudo-Cerinthe, *Koch, Syn. Fl. Germ.* ed. II. 525.—Helvetia.
Pseudo-dentatum, Bill. ex Nym. Consp. 450 = villosum.
Pseudo-gothicum, Arv. Suppl. Monog. Hierac. 29 = tridentatum.
Pseudo-laziophyllum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 46 = rotundatum.
Pseudo-murrum, Schur, l. c. x. (1859) 209 = rotundatum.
Pseudo-Pilosella, Tenore, Prod. p. lxxi = Tenoreanum.
Pseudo-Pyrenaeithes, Sennen, in Bull. Soc. Bot. Fr. ii. (1855) 223 = strictum.
Pseudopsis, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 390 = umbellatum.
Pseudo-ramosum, Schur, Enum. Pl. Transs. 891 = vulgatum.
Pseudo-Schmidtii, *Schur, l. c.* 393.—Transylv.
Pseudo-sciadium, Bor. Fl. Cent. Fr. ed. III. ii. 389 = umbellatum.
pubescens, *Kit. in Linnaea*, xxxii. (1863) 421.—Hungar.
pubescens, Walp. Ann. ii. 1019 (= *Aracium pubescens*). —As. Min.?
pulchellum, Godr. in Gren. & Godr. Fl. Fr. ii. 367, in obs. = Lloltardi.
pulchellum, *Lindeb. in Bot. Notiser* (1872), 120.—Europ.
pulchrum, Froel. ex Steud. Nom. ed. II. i. 765 = Crepis virens.
Pulmonaria, Wallr. Sched. Crit. 423, in obs. = vulgatum.
pulmonarioides, J. & C. Presl, Fl. Čech. 159 = bohemicum.
pulmonarioides, Spreng. Syst. iii. 640 = amplexicaule.
pulmonaroides, *Vill. Prosp.* 86; *Hist. Pl. Dauph.* iii. 133.—Europ.
pulmonarium, Sm. Engl. Fl. iii. 362 = saxatile.
pumilum, Hoppe, ex Willd. Sp. Pl. iii. 1563 = alpinum.
pumilum, Jacq. Fl. Austr. ii. 53. t. 189 = humile.
pumilum, Lapeyr. Hist. Abr. Pl. Pyr. 496 = glaciale.
pumilum, Linn. Mant. ii. 279 = Crepis pygmaea.
pumilum, Tenore, Viagg. Calab. 128 = Crepis Columnae.
purpurascens, Scheele, ex Willk. & Lange, Prod. Fl. Hisp. ii. 362 = saxatile.
purpureum, Willd. Sp. Pl. iii. 1560 = Pterotheca bifida.
pusillum, Hoppe, in Flora, xiv. (1831) 183 = sphaerocephalum.
pusillum, Pursh, Fl. Am. Sept. ii. 502 = Erigeron uniflorus.
pustulatum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 46 = murorum.
pygmaeum, Lam. Fl. Fr. ii. 100 = Crepis pygmaea.
pyrenaeorum, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875–76) 617 = prenanthoides.
pyrenaicum, *Jord. Obs. Pl. Crit.* vii. (1849) 37.—Mont. Pyren.
pyrenaicum, Linn. Sp. Pl. 804 = Crepis blattarioides.

HIERACIUM:—

pyrenaicum, Rochel, Pl. Banat. Rar. 69. t. 39. f. 59 = Crepis viscidula.

pyrrhanthes, *Naeg. & Peter, Hierac.* 330.—Europ.

quercoticolum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 397 = vulgatum.

quercetorum, Jord. in Bor. l. c. 386 = boreale.

quercetorum, *Vukot. in Rad Jugos. Akad. Zagreb.* lviii. (1881) 109.—Europ.

quinquelos x, Naeg. & Peter, Hierac. 385.—Europ.

racemosum, *Waldst. & Kit. ex Willd. Sp. Pl.* iii. 1588; *et Pl. Rar. Hung.* ii. 211. t. 193.—Europ. or.; Reg. Caucas.

Rackii, *Vukot. in Rad Jugos. Akad. Zagreb.* xxxix. (1877) 204.—Europ.

radicans, *Steud. Nom.* ed. II. i. 763, in syn. (Quid ?).—Hab. ?

radicale, Hort. Berol. ex Froel. in DC. Prod. vii. 202 = H. Auricula.

radicescis, Tausch, in Flora, xi. (1828) i. Erg. 55 = praealtum.

Radula, Fries, Epicris. Hierac. 40 = Balansae.

ramiferum, *Dalla Torre, Atl. Alpenfl.* 151.—Europ

Ramondi, Griseb. in Goett. Abh. v. (1853) 100 = Neo-Cerinthe.

ramosissimum, *Gremli, Neue Beitr.* iii. 18.—Helvet.

ramosum, Vill. ex Steud. Nom. ed. I. 409 = alpinum.

ramosum, Wahlenb. Fl. Carp. 345 = gothicum.

ramosum, *Waldst. & Kit. ex Willd. Sp. Pl.* iii. 1579. —Europ.

rapaefolium, *Gilib. Exerc. Schol. Univ. Viln.* (1772) 18; *ex Traute. in Act. Hort. Petrop.* viii. (1883) 550. —Europ.

rapunculoides, *Arv.-in Bull. Soc. Dauph.* (1880) 289. —Europ.

rarinarium, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 413 = murorum.

recreatum, Jord. in Bor. l. c. 415 = murorum.

reconditum, Jord. in Bor. l. c. 409 = vulgatum.

rectiflorum, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875-76) 691 = cerinthoides.

Reichenbachii, Verlot, Cat. Jard. Grenoble (1879), 11 = pulmonarioides.

Reichsteinii, Heldr. & Sart. in Boiss. Diagn. Ser. II. v. 117 = bracteolatum.

relicinum, *Fries, Epicris. Hierac.* 153.—Calif.

remotum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 419 = corinthodes.

repandum, *Schrank, in Denkschr. Akad. Muench.* 1813 (1814) 319.—Hab. ?

repens, S. F. Gray, Nat. Arr. Brit. Pl. ii. 421 = H. Pilosella.

repens, Willd. Sp. Fl. iii. 1564 = aurantiacum.

repens-multiformis, Gilib. Fl. Lituan. i. 240 = H. Pilosella.

reptans, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 74 = staticifolia.

reticulatum, Lindeb. in Bot. Notiser (1878), 185 = sirfcium.

retrodentatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 411 = murorum.

Retzii, Fries, Nov. (1819); ex ej. Summ. Veg. Scand. 338 = pallidum.

Retzii, Griseb. in Goett. Abh. v. (1853) 138, partim = bifidum.

Reuteri, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875-76) 468, 617 = juranum.

Reuteri, Reichb. f. Ic. Fl. Germ. xix. 66. t. 1498 = pulmonarioides.

Reuterianum, Boiss. Diagn. Ser. I. iv. 81 = sparsum.

rhaeticum, Fries, Epicris. Hierac. 46 = alpinum.

rhaeticum, Froel. ex Steud. Nom. ed. II. i. 765 = Crepis Jacquinii.

rhodanthum, *Fries, Epicris. Hierac.* 21.—Sibir.

rhodopeum, Griseb. in Goett. Abh. v. (1853) 91 = petraeum.

rhoeadifolium, Flach. ex Steud. Nom. ed. II. i. 765 = Crepis foetida.

rhoeadifolium, *Kern. ex Dalla Torre, Atl. Alpenfl.* 149.—Europ.

rhomboidale, Lapeyr. in Mém. Acad. Toul. i. (1782) 215. t. 18 = neo-cerinthe.

rigens, Jord. Gr. Jard. Gren. (1849) 20 = boreale.

rigidatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 394 = tridentatum.

rigidulum, Jord. in Bor. l. c. 386 = boreale.

rigidum, Fries, Epicris. Hierac. 138 = umbellatum ?

rigidum, Griseb. in Goett. Abh. v. (1853) 126 = tridentatum.

HIERACIUM:—

rigidum, *Hartm. Handb. Skand. Fl.* ed. I. 300.—Europ.

Rionii, Gremli, Neue Beitr. iii. 16 = caesium.

riphaeum, Uechtritz, in Oestr. Bot. Zeitschr. xxii. (1872) 41 = prenanthoides.

rivulare, *Krock. Fl. Silez. Suppl.* ii. 249.—Europ.

robustum, Fries, Symb. Hierac. 128 = foliosum.

robustum, *Martr. Fl. Tarn*, 488.—Gallia.

rodnense, *Porc. in Magyar Növénet. Lapok*, ix. (1885) 139.—Europ.

rohacsense, *Kit. in Linnaea*, xxxii. (1863) 422.—Hungar.

roseum, *Sch. Bip. in Bonplandia*, ix. (1861) 173, 174. —N. Granat.

rosulatum, Lindeb. in Bot. Notiser (1878) 182 = saxifragum.

Rotae x, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 112.—Europ.

Rothianum, Guerb. ex Froel. in DC. Prod. vii. 208 = praealtum.

Rothianum, Wallr. Sched. Crit. 417 = setigerum.

rotundatum, *Kit. in Schult. Oestr. Fl.* ed. II. ii. 439.—Europ.

rotundellum, *Jeanb. ex Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, iii. (1875-76) 596.—Europ.

rotundifolium, Monn. ex Steud. Nom. ed. II. i. 765 = murorum.

rotundifolium, Willd. ex Spreng. Syst. iii. 641 = pallescens.

rotundum, Dulac, Fl. Hautes-Pyr. 486 = murorum.

rosolanicum, *Rehm. in Oestr. Bot. Zeitschr.* xxiii. (1873) 151.—Europ.

rubellum x, Naeg. & Peter, Hierac. 360.—Europ.

rubellum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 68 = Crepis rubra.

rubescens, *Gilib Exerc. Bot. Schol. Univ. Viln.* (1772) 18; *ex Traute. in Act. Hort. Petrop.* viii. (1883) 551. —Europ.

rubescens, Jord. ex Willk. & Lange, Prod. Fl. Hisp. ii. 268 = caesium.

rubricatum, *Naeg. & Peter, Hierac.* 219.—Europ.

rubricaule, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875-76) 609 = alatum.

rubricaule, Dum. Fl. Belg. Prod. 62 = sabaudum.

rubriformе x, Naeg. & Peter, Hierac. 348.—Europ.

rubrostriatum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 68 = Crepis taraxacifolia.

rubrum, *Peter, in Flora*, lxiv. (1881) 126.—Europ.

rufculum x, Naeg. & Peter, Hierac. 360.—Europ.

rufiarbas, Pers. ex Herd. in Bull. Soc. Nat. Mosc. xlii. (1870) i. 202 = prenanthoides.

Rugelii, Arv. Spicil. Hierac. (1881) 11 = marianum.

runcinatum, Hablitz. ex Rchv. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 413, in textu = Crepis Marschallii.

runcinatum, James, ex Torr. in Ann. Lyc. N. York, ii. (1828) 206 = Crepis runcinata.

runcinatum, Turcz. ex Ledeb. Fl. Ross. ii. 827 = Crepis polytricha.

runcinatum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 77 = Kochianum.

rupestre, *All. Auct. Pedem.* 19. t. 1. f. 2.—Europ.

rupestre, Auct. ex Fries, Symb. Hierac. 96 = rupicolum.

rupestre, F. W. Schmidt, in Neue Abh. Boehm. Ges. i. (1791) 68 = pallidum.

rupicarpum, Schrank, ex Steud. Nom. ed. II. i. 765 = alpinum.

rupicolum, Jord. Cat. Gr. Jard. Dijon (1848) 24 = amplexicaule.

rupicolum, *Fries, Symb. Hierac.* 96.—Europ.

Ruprechtii, *Reiss. Fl. Orient.* iii. 861.—Reg. Caucas.

Rusbyi, *Greene, in Bull. Torrey Bot. Club*, ix. (1882) 94.—N. Mexic.

rutilum x, Naeg. & Peter, Hierac. 360.—Europ.

sabaudum, Ledeb. Fl. Ross. ii. 854 = boreale.

sabaudum, Ledeb. l. c. ii. 852, sphalm. = murorum.

sabaudum, *Linn. Sp. Pl.* 804.—Europ.

sabaudum, Linn. Fil. Suec. ed. II. 974 = rigidum.

sabaudum, O. F. Muell. in Fl. Dan. t. 872 = umbellatum.

sabaudum, Pall. Reise, ii. 397 = virosum.

sabaudum, Willd. ex Froel. in DC. Prod. vii. 206 = latifolium.

sabinum, *Sebast. & Mauri, Fl. Rom. Prod.* 270.—Europ. or.; Orient.

Saelani, Norrl. in Notis. Sällsk. Fenn. et Fl. Fenn. (1874) 496 = flammeum.

HIERACIUM:—

sagittaefolium, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 69 = Crepis nicaeensis.

sagittatum, *Hoffmgg. & Link, Fl. Port.* ii. 146.—Lusitan.

Sagodi, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 393 = tridentatum.

salebrorum, Jord. ex Nym. Consp. 445 = pallidum.

salicifolium, Fries, ex Nym. l. c. 440 = H. Eupatorium.

salicifolium, *Lindeb. in Hartm. Handb. Skand. Fl.* ed. XI. 56.—Europ.

sanctum, Georgi, Beschr. Russ. Reich. III. iv. 1212 = Pterotheca lobata.

sanctum, Linn. Cent. Pl. ii. 30; Amoen. Acad. iv. 328 = Pterotheca nemausensis.

sarmenticaule, Schur, Enum. Pl. Transs. 381 = *bifurcum*.

sarmentosum, Froel. in DC. Prod. vii. 202 = praealtum.

sarmentosum, Salisb. Prod. 181 = H. Pilosella.

Sartorianum, *Boiss. & Heldr. Diagn. Ser. I.* vii. 15.—Graecia; An. Min.

saucium, Jord. ex Nym. Consp. 445 = pallidum.

Saussuri, Sch. Bip. ex Nym. l. c. 454 = subcauliflorum, *Waldst. & Kit.*

saxatile, Balb. ex Fries, Symb. Hierac. 88 = Liottardi.

saxatile, Boiss. Voy. Bot. Espagne, 392 = phlomoides.

saxatile, Elchw. Skizze, 180 = vulgatum.

saxatile, Jacq. Obs. Bot. ii. 30 = glaucum, saxetanum.

saxatile, Schleich. ex Fries, Symb. Hierac. 101 = saxifragum.

saxatile, Thomas, ex Monn. Ess. Hierac. 14 = graminifolium.

saxatile, *Vill. Prosp.* 85; *Hist. Pl. Dauph.* iii. 118.—Europ.

saxetanum, *Fries, Epicris. Hierac.* 69.—Europ.

saxetorum, Jord. ex Nym. Consp. 448 = amplexicaule.

saxifragum, Bab. ex Syme, Engl. Bot. ed. III. v. 177 = linguistum.

saxifragum, *Fries, Symb. Hierac.* 100.—Suecia.

scabricaule, *Birch. Del. Sem. Hort. Heidelb.* (1847); *ex Walp. Ann.* i. 465.—Persia.

scabripes, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 415 = murorum.

scabriusculum, Schwein. in Long, Exped. Winnep. ii. 394 = canadense.

scabrum, Gand. in Nat. Anz. Schw. Ges. iv. (1890) 45 = ramosum.

scabrum, *Michx. Fl. Bor. Am.* ii. 86.—Am. bor.

scabrum, Willd. ex Froel. in DC. Prod. vii. 213 = inuloides.

scapigerum, *Boiss. ex Orph. & Heldr. in Diagn.* Ser. II. iii. 108.—Graecia.

*scapiger*um, Fries, Symb. Hierac. 96 = pallidum.

Schenkii, Willk. ex Fries, in Öfvers. Vet. Akad. Stockh. xxiii. (1866) 161 = prenanthoides.

Schlosseri, Reichb. f. Ic. Fl. Germ. xix. 94. t. 1556 = lanatum.

Schmidtii, Sond. ex Fries, Symb. Hierac. 110 = murorum.

Schmidtii, Tausch, in Flora, xi. (1828) i. Erg. 65 = lactophyllum, pallidum.

Schraderi, Gand. Fl. Helvet. v. 70 = villosum, flexuosum, *Waldst. & Kit.*

Schraderi, Hort. ex Fries, Symb. Hierac. 167 = elatum.

Schraderi, Schleich. ex DC. Fl. Fr. iv. 90 = pilliferum.

Schultzii, F. Schultz, in Arch. Fl. Fr. et Allem. 96; Arch. Fl. 254 = *bifurcum*.

Schultzianum, *Pomel. & Vis. in Mem. Ist. Veget.* xv. (1870) 9.—Europ.

Schulzii, *Fries, Epicris. Hierac.* 150.—Mexic.

sciadophorum, *Naeg. & Peter, Hierac.* 440.—Europ.

sciaphilum, *Dickst. in Deutsche Bot. Monatsb.* ii. (1884) 69.—Europ.

sciaphilum, Jord. ex Nym. Consp. 443 = vulgatum.

scoparium, *Fries, Epicris. Hierac.* 137.—Italia ?

scopulorum, Lapeyr. Suppl. Hist. Pl. Pyr. 124 = saxatile.

scorpioides, *Arv.-in Bull. Soc. Dauph.* (1880) 286. —Europ.

scorzonerifolium, *La Soie, in Bull. Soc. Murith.* ii. (1870) 73.—Helvet.

scorzonerifolium, *Vill. Prosp.* 85; *Hist. Pl. Dauph.* iii. 111.—Europ.

Scouleri, Hook. ex A. Gray, Syn. Fl. N. Am. i. ii. 439 = cynoglossoides.

Scouleri, *Hook. Fl. Bor. Am.* i. 296.—Am. bor. occ.

HIERACIUM:—
semiglaberatum, Vahl, ex Froel. in DC. Prod. vii. 213 = Commersonii.
semivestitum, Sch. Bip. ex Nym. Consp. 446 = lanatum.
semivillosum, Sendtn. ex Nym. l. c. 450 = glaucum.
semproniarum, *Wolf. ex Favrat*, in *Bull. Soc. Murith.* ix. (1880) 67.—Europ.
Sendtneri x, Brugg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 112.—Europ.
Sendtneri, Fries, Epicris. Hierac. 17 = brachiatum.
senescens, Backh. Mon. Hierac. 32 = nigrescens.
sericeum, Gren. & Godr. Fl. Fr. ii. 360 = phlomoides.
sericeum, Lapeyr. Hist. Abr. Pl. Pyr. 477 = Lapeyrousii.
sericeum, Reichb. Ic. Fl. Germ. xix. 65 = saxatile.
Seridis, Fries, ex Nym. Consp. 439 = lactucaceum.
serotinum, Host, Fl. Austr. ii. 419 = umbellatum.
serpyllifolium, *Fries, Epicris. Hierac.* 19.—Italia.
serratifolium, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 417 = murorum.
serratifolium, *Vukot. in Rad Jugos. Akad. Zagreb.* lviii. (1881) 164.—Europ.
sertiflorum, Lam. Fl. Fr. ii. 98 = Crepis praemorsa.
serum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 388 = boreale.
sessiliflorum, Friv. Evk. iv. t. 11 ; ex Boiss. Fl. Orient. iii. 875 = racemosum.
sessiliflorum, *Trinh. in Mém. Acad. Toulouse*, Sér. VII. vii. (1879) 467.—Europ.
setigerum, *Tausch, in Flora*, xi. (1828) i. *Erg.* 61.—Europ.
setigerum, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) ii. 190 (sp. dub).—Sibir.
setosum, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 45.—Transylv.
setosum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 68 = Crepis graveolens.
sibiricum, Lam. Encyc. ii. 368 ; Willd. Sp. Pl. iii. 1583 = Crepis sibirica.
siculum, Guss. Fl. Sic. Syn. ii. 404 = crinitum.
Sieberi, Tausch, in Flora, xi. (1828) i. Erg. 75 = pleiotheles.
Silesi, *Norrl. in Notis. Sällsk. Faun. et Fl. Fenn. Förh. N. S. Haeft.* viii. 135, in nota.—Fennia.
silesiacum, *Krause, in Jahresb. Schles. Ges.* (1850) 101.—Silesia, Sudet.
silhetense, DC. ex Froel. in DC. Prod. vii. 218 = Crepis silhetensis.
silvaticum, Tausch, ex Nym. Consp. 441 = gothicum.
silvestre, Tausch, in Flora, xi. (1828) i. Erg. 70 = boreale, gothicum.
similatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 411 = murorum.
simplex, DC. ex Stend. Nom. ed. II. i. 766 = villosum.
simplex, Tausch, ex Koch, Syn. Fl. Germ. ed. I. 448 = echioides.
simplex, *Timb. & Jeanb. in Bull. Soc. Bot. Fr.* xi. (1864) p. lxxiv.—Europ.
simplex, Vis. Fl. Lib. Spec. 50. t. 13. f. 4 = Leontodon Salzmanni.
sinuosum, *Fries, in Oefver. Vet. Akad. Forhandl. Stockh.* (1856) 147.—Europ.
solidagineum, *Fries, Epicris. Hierac.* 55.—Mont. Pyren.
Sommerfelti, Lindeb. in Bot. Notiser (1872), 119 = bifidum.
sonchifolium, *Bieb. Fl. Taur. Cauc.* ii. 252.—Reg. Caucas.
sonchifolium, Fisch. ex Froel. in DC. Prod. vii. 223 = tridentatum.
sonchifolium, Schrank, in Linnaea, xxxii. (1863) 858 = Pseudo-Cerinthe.
sonchoides, Arv. Suppl. Monog. Hierac. 8 ; et Addit. 10 = saxatile.
sordidum, *Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 292.—Reg. Argent.
spanochlorum, *Lindeb. in Bot. Notiser* (1868), 79.—Europ.
spaniifolium, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 77 = vulgatum.
sparsiflorum x, Naeg. & Peter, Hierac. 767.—Europ.
sparsum, *Fritsch. in Flora*, xix. (1836) 436.—Rumelia.
sparsum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 415 = murorum.
spathophyllum, *Naeg. & Peter, Hierac.* 886.—Europ.
spathulaefolium, Vukot. ex Nym. Consp. 464 = stoloniflorum, *Waldst. & Kit.*
spathulatum, Schenk, in Linnaea, xxxii. (1863) 666 = compositum.

HIERACIUM:—
speciosissimum, Willd. Enum. Hort. Berol. Suppl. 54 = dentatum.
speciosum, Gren. & Godr. Fl. Fr. ii. 359 = saxetanum.
speciosum, *Hornem. Hort. Hafn.* 764.—Europ.
spectosum, Salisb. Prod. 181 = auranticum.
spectabilum, *Yeand. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, vi. (1883–84) 185.—Europ.
sphaerocephalum, Blytt, ex Fries, Symb. Hierac. 10 = cernuum.
sphaerocephalum, *Froel. ex Reichb. in Mossl. Handb.* ed. II. ii. 1386.—Europ. ; As. Min.
sphaerophyllum, Vukot. ex Nym. Consp. 444 = fragile, *Froel.*
sphagnophilum, Ehrenb. ex Boiss. Fl. Orient. iii. 862 = praecultum.
spicatum, All. Fl. Pedem. i. 218 = cydoniaefolium.
spicatum, Gilib. Fl. Lituan. i. 238 = Crepis praemorsa.
spilophaeum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 404 = vulgatum.
spinosum, *Gilib. Exerc. Schol. Univ. Viln.* (1772) 18 ; *ex Traubv. in Act. Hort. Petrop.* viii. (1883) 558.—Europ.
spinulosum, Spreng. Syst. iii. 645 = Sonchus arvensis.
spontaneum x, Naeg. & Peter, Hierac. 439.—Europ.
Sprengerianum, Linn. Sp. Pl. 804 = Picris Sprengerianus.
spurcatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 404 = vulgatum.
spurium, Chaix, in Vill. Hist. Pl. Dauph. iii. 102 ; ex Froel. in DC. Prod. vii. 204 = hybridum.
squamosum, *S. G. Gmel. Itin.* i. 9 ; *ex Y. F. Gmel. Syst.* 1179.—Sibiria.
squarrosum, Ledeb. Fl. Ross. ii. 857 = praec.
staticifolium, *Vill. Prosp.* 35 ; *All. Fl. Pedem.* i. 214. —Europ.
stellatum, Tausch, in Flora, xi. (1828) i. Erg. 59 = pilosum.
stelligerum, *Froel. in DC. Prod.* vii. 214.—Gallia.
stellulipilum, *Naeg. & Peter, Hierac.* 276.—Europ.
stenocladium, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 393 = dentatum.
stenolepis, Lindeb. in Hartm. Handb. Skand. Fl. ed. XI. 45 = murorum.
stenolithum, Schur, Enum. Pl. Transs. 399 = boreale.
stenophyllum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 70 = Crepis alpestris.
Sternbergii, Froel. in DC. Prod. vii. 214 = pallidum.
Sternbergii, Hoppe, ex Hornem. Suppl. Hort. Hafn. 90 = Crepis succisaefolia.
sterrochaetium, *Naeg. & Peter, Hierac.* 498.—Europ.
stipiflorum, Jacq. Fl. Austr. iii. 51 = Chondrilla Pelidium.
stoloni-angustifolium, Gilib. Fl. Lituan. i. 239 = H. Auricula.
stoloniflorum, Bess. Enum. Pl. Volh. 75 = praecultum.
stoloniferum, Desf. Cat. Hort. Par. ed. III. 146 = sparrocephalum.
stoloniflorum, Vis. Fl. Ital. Fragm. 17. t. 20 = Crepis bulbosa.
stoloniflorum, Thuem. in Bonplandia, vi. (1858) 158 = H. Pilosella.
stoloniflorum, *Waldst. & Kit. Pl. Rar. Hung.* iii. 303. t. 278.—Europ.
stoloniflorum, *Wimm. Fl. Schles.* ed. II. (1844) 460.— Germania.
stoloni-latifolium, Gilib. Fl. Lituan. i. 240 = dubium.
stolonosum, *Hegetschw. Fl. Schw.* 779.—Helvet.
striatum, *Rafin. in Am. Monthly Mag.* (1811) 356, nomen.—Am. bor.
strictieaule, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 395 = tridentatum.
strictissimum, *Froel. in DC. Prod.* vii. 211.—Europ. centr.
strictum, *Fries, Symb. Hierac.* 164.—Scandinav.
strictum, Ledeb. Fl. Alt. iv. 123 = Crepis Bungei.
strictum, Opiz, Ackerm. ex Steud. Nom. ed. II. i. 706 (Quid ?).—Europ.
strictum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 68 = Crepis tectorum.
strigosum, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 175.—Mexic.
stupeum, *Fries, in Oefver. Vet. Akad. Forhandl. Stockh.* (1856) 146.—Mexic.
stuppeum, Fries, Epicris. Hierac. 66 = soc.
stupposum, *Reichb. Fl. Germ. Excurs.* 265.—Europ.
subalpinum, *Arv. in Bull. Soc. Dauph.* (1880) 238.— Europ.

HIERACIUM:—
subaudum, Pall. Fl. Ross. Index Tomi ii. pars i = virosum.
subauratum, Schur, Enum. Pl. Transs. 396 = subfuscum.
subcaesium, Fries, ex Nym. Consp. 444 = murorum.
subcandatum, *Lacc, ex Sazs. Phaner. Fl. Orsol's*, 59.—Ins. Oellia.
subdolum, Jord. ex Nym. Consp. 444 = murorum.
subfuscum, *Schur, Enum. Pl. Transs.* 396.—Transylv.
subhirsutum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 384 = boreale.
subluridum, *Arv. in Bull. Soc. Bot. Fr.* xxvi. (1879) *Bibl.* 39.—Europ.
submaculatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 406 = murorum.
submurorum, *Lindeb. in Blytt, Norges Fl.* ii. 643.— Europ.
subnigrescens x, Simk. in Termész. Füzetek, v. (1881) 52.—Europ.
subnivale, *Gren. & Godr. Fl. Fr.* ii. 356.—Mont. Pyren.
subnudum, Froel. in DC. Prod. vii. 218 = venosum.
subnudum, Gilib. Fl. Lituan. i. Coll. iii. 287 = murorum.
subnudum, Schleich. Cat. Pl. Helv. ed. IV. 19 = dentatum.
subolidum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 416 = murorum.
subprealtum, Lindeb. in Bot. Notiser (1868) 78 = praealtum.
subracemosum, Schur, Enum. Pl. Transs. 400 = umbellatum.
subradiatum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 387 = tridentatum.
subrude, *Arv. in Bull. Soc. Dauph.* (1880) 286.— Europ.
subsalinum, Jord. ex Nym. Consp. 439 = boreale.
substoloniflorum, *Naeg. & Peter, Hierac.* 816.— Europ.
subsulfureum, *Naeg. & Peter, l. c.* 176.—Europ.
subumbellatum, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 87 ; iv. (1853) 46.—Europ centr.
subvestitum, *Yeand. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, vi. (1883–84) 190.—Europ.
succisaefolium, All. Fl. Pedem. i. 215 = Crepis succisaefolia.
succloides, *Arv. in Bull. Soc. Dauph.* (1880) 287.— Europ.
sudeticum, *Sterub. in Denkschr. Bot. Ges. Regensb.* i. 11. (1818) 62, partim.—Europ. centr.
sudeticum, Sterub. l. c. 62, partim = bohemicum.
sudeticum, Tausch, ex Fries, Symb. Hierac. 70 = alpinum.
sudeticum, Tausch, ex Link, Handb. i. 808 = carpathicum.
suecicum, *Fries, Symb. Hierac.* 16.—Scandinav.
Sulliantii, Arv. Spicil. Hierac. (1881) 11 = paniculatum.
sulphureum, *Doell, Rhein. Fl.* 521.—Europ.
superbum x, Naeg. & Peter, Hierac. 680.—Europ.
sylvaticum, Balb. ex Froel. in DC. Prod. vii. 232 = Hoppeanum.
sylvaticum, Bertol. Fl. Ital. viii. 485 = crinitum.
sylvaticum, Gouan, Illustr. 56 ; Retz. Obs. i. 27 = saxatile.
sylvaticum, Jan, ex Froel. in DC. Prod. vii. 221 = lanceolatum.
sylvaticum, Lam. Fl. Fr. ii. 96 = vulgatum.
sylvaticum, Lapeyr. Hist. Abr. Pl. Pyr. 472 = sabaudum.
sylvaticum, Salisb. Prod. 181 = umbellatum.
sylvaticum, Wahlenb. Fl. Lapp. 197 ; Fl. Suec. 515 = boreale.
sylvaticum, Wallr. Sched. Crit. 422 = caesium.
sylvaticum, Acct. vide *silvestre*.
sylvicola, Jord. Gr. Jard. Gren. (1849) 21 = vulgatum.
sylvicoagrum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 417 = murorum.
sylvicolorum, Jord. in Bor. l. c. 418 = murorum.
symphytaceum, Arv. in Bull. Soc. Dauph. (1880) 290 = australe.
symphytifolium, *Froel. in DC. Prod.* vii. 232.— Sicilia.
syngynes, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 417 = murorum.
Tarasaci, Linn. Sp. Pl. ed. II. 1125 = Leontodon Taraxaci.

HIERACIUM :—
Taraxaci, Tim. ex Reichb. Fl. Germ. Excurs. 258 = Leontodon autumnalis.
taraxacifolium, Frœl. ex Stend. Nom. ed. II. i. 766 = Crepis Jacquini.
taraxaciformæ, Arv. in Bull. Soc. Dauph. (1880) 287. —Europ.
taraxacoides, Ehrenb. ex Fries, Epicris. Hierac. 158 = Taraxacum Asserani.
Taraxacoris, Georgi, Beschr. Russ. Reich. IH. iv. 1211 = Leontodon autumnalis.
tardans, *Peter*, in *Engl. Bot. Jahrb.* v. (1884) 256.—Helvet.
tardiusculum, *Naeg. & Peter, Hierac.* 255.—Europ.
Tatrae, Griseb. in Goett. Abh. v. (1858) 158 = bupleuroides.
taurinense, *Jord. Obs. Pl. Crit.* vii. (1849) 40.—Europ.
tayyetum, Boiss. & Heldr. Diagn. Ser. I. vii. 15 = pannosum.
taygetum, *Fries, Symb. Hierac.* 91.—Taygeti.
tectorum, Hornem. Hort. Hafn. 766 = Crepis tectorum.
tendinum, *Naeg. & Peter, Hierac.* 458.—Europ.
Tenoreanum, *Frœl. in DC. Prod.* vii. 200.—Ital.
tenue, Freyn, ex Velen. in Oestr. Bot. Zeitschr. xxxiii. (1883) 389 = Freynianum.
tenuifolium, Host. Fl. Austr. ii. 411 = barbatum.
tephrocephalum, Vukot. ex Nym. Consp. 454 = *bifurcum*.
tephrodes, *Naeg. & Peter, Hierac.* 641.—Europ.
tephroglaucum, *Naeg. & Peter, l. c.* 518.—Europ.
tetradymum x, Naeg. & Peter, l. c. 872.—Europ.
tetragenes x, Naeg. & Peter, l. c. 633.—Europ.
thapsiformæ, Uechtr. ex Asch. & Kan. Cat. Serb. 48 = thapsoides.
thapsoides, *Panč. Elench. Crna Gora.* 59.—Montenegro.
Thevenii, *Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, vi. (1883–84) 185.—Europ.
thlaspifolium, *Frœl. ex Stend. Nom.* ed. II. i. 766, nomen.—Europ.
Thomasii, Stand. l. c. sphalm. vide *Tomasinii*.
thuringiacum, *Bernh. Syst. Verz. Erf.* 187.—Europ.
thynoideum, *Fisch. ex Stend. Nom.* ed. I. 409, nomen. —Sibir.
thyrsoideum, Fries, Symb. Hierac. 141 = abeciasum !
tinctum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 405 = vulgatum.
tingitanum, Salzm. ex DC. Prod. vii. 158 = Crepis tingitana.
Tomasinii, Host, Fl. Austr. ii. 414 = lanatum.
tomentosum, *Alt. Fl. Pedem.* i. 216.—Europ.
tomentosum, Linn. Cent. Pl. i. 26 = andryaloides.
Tommasinii, Reichb. f. Ic. Fl. Germ. xix. 100. t. 1569 = stupposum.
tortifolium, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 396 = vulgatum.
tovarense, *Fries, Epicris. Hierac.* 153.—Venezuela.
Trachselianum, Christen. in Mittheil. Bern. Naturf. Ges. (1880) ; ex *cj.* Hierac. Schweiz. 15 = oxydon.
trachyculon, Vukot. ex Nym. Consp. 438 = sabaudum.
transilvanica, Arv. Suppl. Monog. Hier. 17 = vulgatum.
transilvanicum, *Schur, ex Fries, Epicris. Hierac.* 97. —Transayl.
transylvanicum, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 97 = rotundatum.
Triasii, Cambess. in Mém. Mus. Par. xiv. (1827) 268 = Crepis Triasii.
trichocephalum, Willd. Enum. Hort. Berol. Suppl. 55, nomen = glabratum.
trichodes, Griseb. ex Reichb. f. Ic. Fl. Germ. xix. 88 = crocatum.
trichodes, *Naeg. & Peter, Hierac.* 452.—Europ.
trichophyllum, Schur, Enum. Pl. Transs. 396, Index, sphalm. vide *trichocephalum*.
tricolor, Kit. in Linnæa, xxxii. (1863) 417 = stoloniflorum, *Waldst. & Kit.*
tridentatum, *Fries, Mant.* ii. 48, in syn.; *et Symb. Hierac.* 171.—Europ. ; As. Bor.
trigenes x, Naeg. & Peter, Hierac. 629.—Europ.
trinothum x, Naeg. & Peter, l. c. 768.—Europ.
triplex, *Naeg. & Peter, l. c.* 260.—Europ.
trisanguineum, *Kit. in Linnæa*, xxxii. (1863) 416.—Hungar.
triste, *Willd. ex Spreng. Syst.* iii. 640.—Unallasch. ; Am. bor.
truncatum, *Lindeb. in Bot. Notiser*, (1866) 69.—Europ.
tuberosum, Balb. ex Stend. Nom. ed. II. i. 766 = Leontodon tuberosus, *Linn.*

HIERACIUM :—
tuberosum, Brot. Fl. Lusit. i. 318 = Crepis bulbosa.
tubulosum, Lam. Encyc. ii. 367 = intybaceum.
tubulosum, Schleich. ex Stend. Nom. ed. II. i. 766 = alpinum.
turbinatum, Spreng. Syst. iii. 647 = Crepis blattarioides.
uliginosum, *G. F. Koch, in Pollichia*, x. (1852) 29, 30. —Europ.
uliginosum, Turcz. in Bull. Soc. Nat. Mosc. (1838) 96 = Crepis Bungei.
Umbella, *Naeg. & Peter, Hierac.* 462.—Europ.
umbellatum, *Linn. Sp. Pl.* 804.—Hab. bor. temp.
umbelliferum, *Naeg. & Peter, Hierac.* 785.—Europ.
umbelliforme, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 389 = umbellatum.
umbraticolum, Jord. l. c. 399 = vulgatum.
umbrosum, Jord. Cat. Gr. Jard. Dijon, (1848) 24 = vulgatum.
umbrosum, Schur, Enum. Pl. Transs. 392 = cæsium.
uncinatum, *Kit. in Linnæa*, xxxii. (1863) 424.—Hungar.
undulatum, *Ait. Hort. Kew.* ed. I. iii. 124.—Hispan.
undulatum, *Boiss. Fl. Orient.* iii. 867.—Thessalia.
uniflorum, Hoppe, in Flora, xiv. (1831) 189 = sphaerocephalum.
urens, *Fisch. ex Stend. Nom.* ed. I. 409, nomen.—Hab. ?
urticaceum, *Arv. in Bull. Soc. Dauph.* (1880) 285.—Europ.
Urvillei, *Sch. Bip. in Bonplandia*, ix. (1861) 174.—Bras.
ustulatum, Arv. in Bull. Soc. Dauph. (1880) 288 = glanduliferum.
vagans, Grm. & Godr. Fl. Fr. ii. 258 = glabratum.
vaginatum, Jeanb. & Timb. in Bull. Soc. Sc. Phys. & Nat. Toulouse, iii. (1875–76) 603 = alatum.
vagum, Jord. Gr. Jard. Grem. (1849) 21 = boreale.
Vahlii, *Frœl. in DC. Prod.* vii. 204.—Hispan.
Vaillantii, Tausch, in Flora, xi. (1828) i. Erg. 57 = cymosum.
valderianum, Fries, Symb. Hierac. 60 = pyrenaicum.
valdepilosum, *Vill. Prosp.* 24 ; *Hist. Pl. Dauph.* iii. 106.—Europ.
valderamosum, Schleich. ex Stend. Nom. ed. II. i. 766 = alpinum.
valletrianum, *Boutigny, ex Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, vi. (1883–84) 191, nomen.—Europ.
vallicolum, Jord. ex Nym. Consp. 439 = boreale.
vallisiacum, *Fries, Epicris. Hierac.* 122.—Helvetia.
Vancouverianum, Arv. Suppl. Hierac. 10 = albiflorum.
variegatum, Lam. Encyc. ii. 362 = Hypochoeris variegata.
vasconicum, *Jord. ex Martr. Fl. Tarn*, 452.—Gallia.
Vayredae, Arv. ex Vayr. in Anal. Soc. Esp. Hist. Nat. xi. (1882) 108 = saxatile.
vellerum, Schenk, ex Fries, in Ofvers. Vet. Akad. Stockh. xxiii. (1866) 160 = plumosum.
velutellum, *Naeg. & Peter, Hierac.* 272.—Europ.
velutinum, Hegetschw. ex Frœl. in DC. Prod. vii. 199 = H. Pilosella.
venetianum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 390 = umbellatum.
venetianum, *Naeg. & Peter, Hierac.* 636.—Europ.
venosum, *Linn. Sp. Pl.* 803.—Am. bor.
ventricosum, *Norrl. ex Wainio, in Meddel. Soc. Fauna et Fl. Fenn.* iii. (1878) 65.—Fennia.
verbascifolium, Pers. Syn. ii. 374 = tomentosum.
verbascifolium, Vis. ex Nym. Consp. 446 = lanatum.
verbenacum, Arv. in Lamotte, Prod. Cent. Fr. ii. 492 = umbellatum.
Verlotii, Jord. ex Nym. Consp. 444, 445 = cæsium, lasiophyllum.
vernale, Schur, Enum. Pl. Transs. 398 = lasiophyllum.
vernum, Steud. ex Nym. Consp. 444 = bifidum.
verruculatum, Link, Enum. Hort. Berol. ii. 287 = scabrum.
versicolor, Fries, in Oefver. Vet. Akad. Forhandl. Stockh. (1856) 149 = stoloniflorum, *Waldst. & Kit.*
versicolor, Wallr. Beitr. Fl. Hercyn. 233 = floribundum.
versutum, Frivald. ex Griseb. in Goett. Abh. v. (1858) 148 = chaldicolum.
versutum, Frivald. ex Nym. Consp. 446 = pannosum.
vestitum, Fisch. ex Hend. in Bull. Soc. Nat. Mosc. xlii. (1870) 1. 204 = tritte.
vestitum, Grm. & Godr. Fl. Fr. ii. 269 = rupicolum.
vicinum, Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse, iii. (1875–76) 606 = anglicum.

HIERACIUM :—
victum, Fries, ex Nym. Consp. 445 = pallidum.
Villarsii, *Jeanb. ex Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, iii. (1875–76) 594.—Europ.
Villarsii, F. Schultz, in Flora, xliv. (1861) 35 = *bifurcum*.
Villarsii, Serres, in Bull. Soc. Bot. Fr. ii. (1855) 225 = Pseudo-Cerinthe.
Villarsii, Vill. ex Steud. Nom. ed. I. 58, 409 = Leontodon hirtus.
villosum, Dicks. in Trans. Linn. Soc. ii. (1794) 288 = sudeticum !
villosum, Griseb. Spicil. Fl. Rumel. ii. 271 = chaldicolum.
villosum, *Jacq. Enum. Vindob.* 149, 271 ; *Linn. Sp. Pl.* ed. II. 1130.—Europ.
villosum, Lapeyr. Hist. Abr. Pl. Pyr. 476 = cerinthoides.
villosum, Lepech. Reise, i. 287 (Quid ?).—Ross.
villosum, Mönn. Enc. Hierac. 59 = piliferum.
villosum, Sm. Engl. Bot. t. 2379 = eximium.
vindobonense, *Wiesb. ex Dichtl. in Deutsche Bot. Monatsh.* ii. (1884) 66.—Austria.
Violletianum, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, iii. (1875–76) 586.—Europ.
virescens, *Schleich. Cat. Pl. Helv.* ed. III. 17 (Quid ?). —Helvet.
virescens, Sond. ex Koch, Syn. Fl. Germ. ed. II. 1027 = tridentatum.
virestorum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 395 = tridentatum.
Virga-aurea, Cass. in Ann. Sc. Nat. Sér. III. vii. (1847) 209. t. 12 = crinitum.
virgatum, Gerh. ex Fries, Symb. Hierac. 119 = vulgarum.
virgatum, Hort. ex Frœl. in DC. Prod. vii. 226 = sabaudum.
virgatum, Lapeyr. ex Pursh, Fl. Am. Sept. ii. 508 = canadense.
virgultorum, Jord. Cat. Gr. Jard. Dijon (1848) 24 = boreale.
viride, *Arv. in Bull. Soc. Dauph.* (1880) 287.—Europ.
viridicollum, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 416 = murorum.
viridissimum, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. Nat. Toulouse*, iii. (1875–76) 613.—Europ.
virosum, Frœl. in DC. Prod. vii. 226, partim = foliosum.
virosum, Grm. & Godr. Fl. Fr. ii. 287 = lactucaceum.
virosum, *Pall. Reise*, i. 501.—Europ. ; As. bor.
viscidatum, Tausch, ex Koch, Syn. Fl. Germ. ed. I. 447 = præsilium.
viscidatum, Vukot. in Rad Jugos. Akad. Zagreb. vii. (1869) 74 = intybaceum.
viscosum, *Arv. in Bull. Soc. Dauph.* (1880) 289.—Europ.
vitistipetalum, Vukot. ex Nym. Consp. 454 = stoloniflorum, *Waldst. & Kit.*
vivariense, Jord. in Bor. Fl. Cent. Fr. ed. III. ii. 387 = boreale.
vogesiacum, Mougeot, ex Fries, Symb. Hierac. 59 = decipiens.
vulcanicum, Griseb. in Bot. Zeit. viii. (1850) 688 = pallidum.
vulgare, Tausch, in Flora, xi. (1828) i. Erg. 58 = dubium.
vulgatum, *Fries, Nov. Fl. Suec.* ed. II. 258.—Reg. bor. temp.
vulgatum, Spreng. Syst. iii. 644 = humile.
Waldsteinii, Tausch, in Flora, xi. (1828) i. Erg. 65 = lanatum.
Warmingii, *Baker, in Mart. Fl. Bras.* vi. 111. 388.—Bras.
Wulfenianum, Hoppe, ex Hornem. Hort. Hafn. 766 = graminifolium.
Wilhemsii, Schneid, in Linnæa, xxxi. (1861–62) 655 = olivaceum.
Wimmeri, Uechtritz, in Oestr. Bot. Zeitschr. xxii. (1872) 277 = pallescens.
Wolfgangianum, Bess. ex Richw. Skisse, 150 = prætense !
Wolfianum, Favre, ex Favrat, in Bull. Soc. Murith. ix. (1880) 67 = rupicolum.
xanthopsilopsilum x, Naeg. & Peter, Hierac. 319.—Europ.
xystolepium, *Naeg. & Peter, l. c.* 231.—Europ.
Zielanum, Tausch, in Flora, xi. (1828) i. Erg. 62 = pseudicum.

HIERANTHEMUM, Spach, Hist. Vég. Phan. iv. 81 (1840) = **Heliotropium**, Tourn. (Boragin.).

HIERANTHES, Rafin. Sylva Tellur. 79 (1838) = **Stereospermum**, Cham. (Bignoniac.).
fragrans, Rafin. l. c. = S. chelonoides.

HIERICONTIS, Adans. Fam. ii. 421 (1763) = **Anastatica**, Linn. (Crucif.).

HIERNIA, S. Moore, in Journ. Bot. xvlii. (1880) 196.
ACANTHACEAE.
angolensis, *S. Moore*, *l. c.*—Afr. trop.

HIEROCHLOA, Beauv. Agrost. 62 (1812) = seq.

HIEROCHLOË, S. G. Gmel. Fl. Sibir. i. 100 (1747). *GRAMINEAE*, Benth. & Hook. f. iii. 1139.
ATAXIA, R. Br. in Parry, 1st Voy. App. 198 (1824).
DIMERIA, Endl. Gen. 81, in syn. (1836).
DIMESIA, Rafin. in Am. Monthly Mag. (1818) 175.
DISARRHENUM, Labill. Nov. Holl. Pl. ii. 82. t. 282 (1806).
SAVASTANA, Schrank, Baier. Fl. i. 100, 387 (1789).
TORRESIA, Ruiz & Pav. Prod. Fl. Per. 125 (1794).
alpina, *Roem. & Schult. Syst.* ii. 515.—Reg. bor. et arct. ; N. Zel.
altissima, Steud. Syn. Pl. Gram. 18 = utriculata.
antarctica, R. Br. Prod. 209 = redolens.
arctica, *Fries*, *Kel. Hæmk.* i. 252.—Sinus Nootka.
arenaria, *Steud. Syn. Pl. Gram.* 416.—Reg. Magellan.
australis, *Roem. & Schult. Syst.* ii. 514.—Europ.
borealis, Hook. f. Fl. N. Zel. i. 300 = alpina.
borealis, *Roem. & Schult. Syst.* ii. 513.—Reg. bor.
Brunonis, *Hook. f. Fl. Antarct.* 93.—Ins. D. Auckland.
Bungeana, *Trin. in Mém. Acad. Pétersb.* Sér. V. vi. 11. (1839) 82.—China.
dahurica, Trin. l. c. 80 = glabra.
Dregei, *Nees*, *Fl. Afr. Austr.* i. 9.—Afr. austr.
Eckloni, *Nees*, *l. c.* 7.—Afr. austr.
fragrans, Roem. & Schult. Syst. ii. 514 = borealis.
Fraseri, Hook. f. Fl. Tasm. ii. 108 = redolens.
glabra, *Trin. in Spreng. Neue Entdech.* ii. 66.—Davuria.
glauca, Gliem. ex Bab. in Journ. Linn. Soc. xi. (1870) 341 = borealis ?
latifolia, Kunth, Rév. Gram. i. 21 = Centotheca lappacea.
laxa, *R. Br. in Wall. Cat.* n. 8796.—Reg. Himal.
macrophylla, *Thurb. ex Boland. in Trans. Calif. Agr. Soc.* (1864) 68.—Calif.
magellanica, Hook. f. Fl. Antarc. ii. 375 = redolens.
mexicana, *Benth. in Journ. Linn. Soc.* xix. (1881) 77. —Mexic.
occidentalis, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863), 100 = borealis.
odorata, Beauv. Agrost. t. 19. *f.* 5 = australis.
odorata, Wahlenb. Fl. Ups. 32 = borealis.
orientalis, Fries & Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 224 = borealis.
parviflora, Presl, Cyp. et Gram. Sic. 37 = Holcus tenuis.
pauciflora, *R. Br. in Parry*, *1st Voy. Suppl.* 198.—Am. arct.
racemosa, Trin. in Mém. Acad. Pétersb. Sér. V. vi. 11. (1839) 79 = pauciflora.
rariflora, *Hook. f. Fl. Antarct.* 93.—Austral.
redolens, *R. Br. Prod.* 209 ; *Roem. & Schult. Syst.* ii. 514.—Reg. austr. temp.
repens, Beauv. Agrost. 62 = borealis.
submutica, F. Muell. in Trans. Vict. Inst. i. (1855) 48 = redolens.
tenuifolia, *Steud. in Lechl. Berl. Am. Austr.* 52.—Chili.
Tongo, *Nees*, *Fl. Afr. Austr.* i. 7.—Afr. austr.
utriculata, *Kunth*, *Rév. Gram.* i. 198, t. 8.—Chili.
vinealis, Schur, in Oestr. Bot. Zeitschr. x. (1859) 12 = borealis.

HIEROCHONTIS, Medic. Pflanzeng. 51 (1792) = **Anastatica**, Linn. (Crucif.).

HIEROCONTIS, Steud. Nom. ed. I. 409 (1821) = prec.

HIERONIA, Vell. Fl. Flum. 284 ; v. t. 116 (1825) = **Davilla**, Vand. (Dilleniac.).
scabra, Vell. Il. cc. = D. rugosa.

HIERONIMA, Allem. Dis. Rio Janeiro (1848), t. = **Hyeronima**, Allem. (Euphorb.).

HIERONYMA, Baill. Etud. Gén. Euph. 652 (1858) = prec.

HIESINGERA, Endl. Gen. Suppl. v. 47 (1850) = HISINGERA, Hellen. = **Xylosma**, Forst. (Bixin.).

HIGGENSIA, Steud. Nom. ed. II. i. 767 (1840) = HIGGINSIA, Pers. = **Hoffmannia**, Sw. (Rubiac.).

HIGGINSIA, Blume, Bijdr. 988 (1826) = **Petunga**, DC. (Rubiac.).
glomerata, Blume, l. c. 989 = P. glomerata.
longifolia, Blume, ex DC. Prod. iv. 399 = P. longifolia.
microcarpa, Blume, Bijdr. 988 = P. Roxburghii.

HIGGINSIA, Pers. Syn. i. 188 (1805) = **Hoffmannia**, Sw. (Rubiac.).
aggregata, Pers. l. c. (= *Euosmia aggregata*).—Peruv.
angustifolia, Bartl. ex DC. Prod. iv. 399.—Peruv.
discolor, Planch. in Fl. des Serres, v. (1849) 482 d = Hoffmannia discolor.
Ghiesbreghtii, Hook. Bot. Mag. t. 5383 (*Ghrisbechtii*) = Hoffmannia Ghiesbreghtii.
latifolia, Bartl. ex DC. Prod. iv. 399.—Peruv.
macrophylla, Planch. in Fl. des Serres, v. (1849) 482 d = Hoffmannia macrophylla.
mexicana, Link, Klotzsch & Otto, Ic. Pl. Rar. i. 58 = Hoffmannia mexicana.
obovata, Pers. Syn. i. 188.—Peruv.
psychotriaefolia, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 50 = Hoffmannia psychotriaefolia.
refulgens, Hook. Bot. Mag. t. 5346 = Hoffmannia refulgens.
regalis, Hook. l. c. t. 5280 = Hoffmannia regalis.
verticillata, Pers. Syn. i. 188.—Peruv.

HILACIUM, Steud. Nom. ed. II. i. 767 (1840), sphalm. = HYLACIUM, Beauv. = **Psychotria**, Linn. (Rubiac.).

HILAIRIA, DC. Prod. vii. 33 (1838) = **Onoseris**, DC. (Compos.).
paniculata, DC. l. c. = O. Isotypus.

HILARIA, H. B. & K. Nov. Gen. et Sp. i. 116. t. 37 (1815). *GRAMINEAE*, Benth. & Hook. f. iii. 1121.
HEXARRHENA, J. & C. Presl, Rel. Haenk. i. 326. t. 45 (1830).
PLEURAPHIS, Torr. in Ann. Lyc. N. York, i. (1824) 148. t. 10.
SYMBASIANDRA, Willd. ex Steud. Nom. ed. II. i. 767 (1840).
cenchroides, *H. B. & K. Nov. Gen. et Sp.* i. 117.—Mexic.
Jamesii, *Benth. in Journ. Linn. Soc.* xix. (1881) 62.—Am. bor. occ.
mutica, *Benth. l. c.*—Am. bor. occ.
rigida, *Vasey*, *Grass. U. St.* 16.—Am. bor. occ.
setigera, *Benth. in Journ. Linn. Soc.* xix. (1881) 62.—Am. bor. occ.

HILBERTIA, Thouin, ex Reichb. Consp. 219* (1828) (Quid ?).

HILDEBRANDTIA, Vatke, in Journ. Bot. xiv. (1876) 313 ; et in Monatsb. Akad. Berl. (1876) 864.
CONVOLVULACEAE, Benth. & Hook. f. ii. 1243.
africana, *Vatke*, *ll. cc.* 313 ; 865.—Afr. trop.

HILDEGARDIA, Schott & Endl. Melet. 33 (1832) = Sterculia, Linn.
Candolleana, Steud. Nom. ed. II. i. 767 = seq.
Candollei, Schott & Endl. Melet. 33 = S. populifolia.
macrophylla, Schott & Endl. l. c. = S. macrophylla.
populifolia, Schott & Endl. l. c. = S. populifolia.

HILLEBRANDIA, Oliver, in Trans. Linn. Soc. xxv. (1866) 361. t. 46. *BEGONIACEAE*, Benth. & Hook. f. i. 849.
sandwicensis, *Oliver*, *l. c.*—Ins. Sandvic.

HILLERIA, Vell. Fl. Flum. 47 ; t. t. 122 (1825) = **Mohlana**, Mart. (Phytol.).
elastica, Vell. Il. cc. = M. nemoralis.

HILLIA, Jacq. Enum. Pl. Carib. 3 (1760). *RUBIACEAE*, Benth. & Hook. f. ii. 29.
FERREIRIA, Vell. ex Vand. Fl. Lusit. 21. t. 1. f. 8 (1788) ; et in Roem. Script. 98. t. 6.
SALDANHA, Vell. Fl. Flum. 142 ; ii. t. 157, 158 (1825).
brasiliensis, *Cham. & Schlecht. in Linnaea*, iv. (1829) 201.—Bras.
longiflora, Blanco, Fl. Filip. ed. I. 235 = Loranthus ampullaceus.
longiflora, Hort. ex Lem. in Fl. des Serres, Sér. I. iii (1847) 188 = prasantha.
longiflora, Sw. Prod. Veg. Ind. Occ. 58 = parasitica.
parasitica, *Jacq. Enum. Pl. Carib.* 18.—Am. trop.
prasantha, Lem. in Fl. des Serres, Sér. I. iii. (1847) t. 188 = Fagraea zeylanica.
tetrandra, *Sw. Prod Veg. Ind. Occ.* 58.—Jamaica.
tubaeflora, *Cham. in Linnaea*, ix. (1834) 260.—Bras
tuxtlensis, [Moç. & Sesse,] ex DC. Prod. iv. 351 = tetrandra ?

HILSENBERGIA, Boj. Hort. Maurit. 42 (1837) = **Dombeya**, Cav. (Malvac.).
cannabina, Boj. l. c. = D. cannabina.

HILSENBERGIA, Tausch, ex Reichb. Consp. 117 (1828) = **Ilhretia**, Linn. (Borag.).
Ehretia, Tausch, ex Dunal, in DC. Prod. xiii. 478 = E. petiolaris.
rugosa, Dunal, l. c. = E petiolaris.

HIMANTHOGLOSSUM, Koch, Syn. Fl. Germ. ed. I. 689 (1837) = HIMANTOGLOSSUM, Spreng. = **Orchis**, Tourn.

HIMANTHOPHYLLUM, D. Dietr. Syn. Pl. ii. 1045, 1186 (1840) = HIMANTOPHYLLUM, Spreng. = **Clivia**, Lindl. (Amaryll.).

HIMANTOCHILUS, T. Anders. in Benth. & Hook. f. Gen. ii. 1117 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
sessiliflorus, *T. Anders. l. c.*—Afr. trop.

HIMANTOGLOSSUM, Spreng. Syst. iii. 675, 694 (1826) = Orchis, Tourn.
anthropophorum, Spreng. l. c. 694 = Aceras anthropophora.
caprinum, Spreng. l. c. = O. hircina.
cucullatum, Reichb. Fl. Germ. Excurs. 120 (= *Gymnadenia cucullata*).—Podol. ; Sibir.
formosum, C. Koch, in Linnaea, xxii. (1849) 287 = O. formosa.
hircinum, Spreng. Syst. iii. 694 = O. hircina.
parviflorum, Spreng. l. c. = Habenaria intacta.
satyriodes, Spreng. l. c. = O. satyriodes.
secundiflorum, Reichb. Fl. Germ. Excurs. 120 = Habenaria intacta.
viride, Reichb. l. c. = Habenaria viridis.

HIMANTOPHYLLUM, Spreng. Gen. i. 276 (1830) = **Clivia**, Lindl. (Amaryllid.).
Aitoni, Duch. in Journ. Soc. Imp. Hortic. v. (1850) 298 = C. nobilis.
cyrtanthiflorum x, Groenl. in Rev. Hortic. (1859) 258.
miniatum, Groenl. l. c. 125 = C. miniata.

HIMANTOSTEMMA, A. Gray, in Proc. Am. Acad. xx. (1885) 294. *ASCLEPIADACEAE*.
Pringlei, *A. Gray*, *l. c.*—Am. bor. occ.

HIMAS, Salisb. Gen. Pl. Fragm. 91 (1866) = **Lachenalia**, Jacq. (Liliac.).
angustifolia, Salisb. l. c. = L. contaminata.
hyacinthoides, Salisb. l. c. = L. contaminata.

HIMATANTHUS, Willd. ex Roem. & Schult. Syst. v. p. xiii. 231 (1819) = **Plumeria**, Linn. (Apocyn.).
rigida, Hoffmgg. ex Willd. l. c.—Bras.

HIMENANTHUS, Steud. Nom. ed. II. i. 768 (1840) sphalm. = seq.

HIMERANTHUS, Endl. Gen. 666 (1839). *SOLANACEAE*, Benth. & Hook. f. ii. 898.
erosus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 346.—Reg. Argent.
Lhotzkii, Steud. Nom. ed. II. i. 768 = runcinatus.
magellanicus, *Griseb. in Gotts. Abh.* vi. (1854) 128.—Reg. Magellan.
runcinatus, *Endl. Gen.* 666.—Reg. Argent.
tridentatus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 346.—Reg. Argent.

HINDSIA, Benth. in Lindl. Bot. Reg. (1844) Misc. 40. t. 40. *RUBIACEAE*, Benth. & Hook. f. ii. 87.
MACROSIPHON, Miq. in Linnaea, xix. (1847) 442.
longiflora, *Benth. in Lindl. Bot. Reg.* (1844) t. 40.—Bras.
ramosissima, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 106.—Bras.
violacea, *Benth. in Lindl. Bot. Reg.* (1844) *Misc.* 40. t. 40.—Bras.

HINGCHA, Roxb. Hort. Beng. 69 (1814) = HINGTSHA, Roxb. = **Ehnhydra**, Lour. (Compos.).

HINGHSTONIA, Steud. Nom. ed. II. i. 768 (1840) sphalm. = HINGSTONIA, Rafin. = **Verbesina**, Linn. (Compos.).

HINGSTONIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Verbesina**, Linn. (Compos.).
exaltata, Rafin. l. c. = V. occidentalis.

HINGTSHA, Roxb. Fl. Ind. iii. 448 (1832) = **Ehnhydra**, Lour. (Compos.).
repens, Roxb. l. c. = E. fluctuans.

HINTERHUBERA, Reichb. ex Nym. Consp. 66 (1878) = **Hutchinsia**, R. Br. (Crucif.).
Berengariana, Reichb. ex Nym. l. c. = Capsella procumbens.
pauciflora, Reichb. ex Nym. l. c. = Hutchinsia pauciflora.

HINTERHUBERA, Sch. Bip. in Flora, xxiv. (1841) 1. Intell. 42; xxv. (1842) 419 = **Chrysanthellum**, Rich. (Compos.).
abyssinica, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 150, nomen = Microlecane abyssinica.
Kotschyi, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) 1. Intell. 42 = C. indicum.

HINTERHUBERA, Sch. Bip. in Wedd. Chlor. And. i. 185. t. 39 (1858). *COMPOSITAE*, Benth. & Hook. f. ii. 275.
SCHAETZELLA, Sch. Bip. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 102.
columbica, *Sch. Bip. in Wedd. Chlor. And.* i. 186.—Venezuela.
ericoides, *Wedd. l. c.* 184.—Venezuela.
Lessigni, *Wedd. l. c.* 184.—Venezuela.

HIORTHIA, Neck. Elem. i. 97 (1790) = **Anacyclus**, Linn. (Compos.).
alexandrina, Less. Syn. Comp. 258 = Anacyclus alexandrinus.
aurea, Less. l. c. = Anthemis aurea.
orientalis, Less. l. c. = Anthemis montana.
valentina, Less. l. c. = Anacyclus valentinus.

HIORTIA, Juss. in Dict. Sc. Nat. xxi. 170 (1821) = prasc.

HIOSCIAMUS, Neck. Delic. Gallo-Belg. i. 122 (1768) = **Hyoscyamus**, Tourn. (Solan.).

HIPECOUM, Vill. Hist. Pl. Dauph. ii. 248 (1787) = **Hypecoum**, Linn. (Papav.).

HIPERICUM, Neck. Delic. Gallo-Belg. ii. 315 (1768) = **Hypericum**, Tourn.

HIPOCHAERIS, Nocca, in Usteri, Ann. Bot. v. (1795) 48 = **Hypochoeris**, Vaill. (Compos.).

HIPOCHOERIS, Neck. Delic. Gallo-Belg. ii. 332 (1768) = prasc.

HIPOCREPIS, Neck. Delic. Gallo-Belg. ii. 309 (1768) = **Hippocrepis**, Linn. (Legum.).

HIPPASTRUM, Herb. App. 21 (1821). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 794.
ASCHAMIA, Salisb. Gen. Pl. Fragm. 124 (1866).
AULICA, Rafin. Fl. Tellur. iv. 10 (1836).
CHONAIS, Salisb. Gen. Pl. Fragm. 138 (1866).
EUSARCOPE, Rafin. Fl. Tellur. iv. 11 (1836).
HABRANTHUS, Herb. Amaryll. 166 (1837), partim.
LAIS, Salisb. Gen. Pl. Fragm. 124 (1866).
LEOPOLDIA, Herb. in Bot. Mag. sub t. 2113, in nota (1819).
MYOSTEMMA, Salisb. Gen. Pl. Fragm. 135 (1866).
OMPHALISSA, Salisb. l. c. 134 (1866).
PHYCELLA, Lindl. in Bot. Reg. sub t. 998 (1825).
RHODOLIRION, Phil. in Linnaea, xxiii. (1857-58) 65.
RHODOPHIALA, Presl, Bot. Bemerk. 115 (1844).
TRISACARPIS, Rafin. Fl. Tellur. iv. 11 (1836).
acuminatum, M. Roem. Syn. Ensat. 141 = rutilum.
advenum, *Herb. App.* 21.—Chili.
ambiguum, Hook. Bot. Mag. t. 3542 = solandriifolium.
Anderssoni, *Baker, in Journ. Bot.* xvi. (1878) 82.—Uruguay.
andicolum, *Baker, l. c.*—Chili.
andinum, Baker, l. c. 84 = Herbertianum.
Andreanum, *Baker, in Gard. Chron.* (1880) ii. 424.—Am. trop.
anomalum, Lindl. ex Planch. in Fl. des Serres, Sér. I. v. (1849) sub t. 455, 456 = H. Cybister.
aulicum, *Herb. App.* 21; *Amaryll.* 135.—Bras.
Bagnoldi, *Baker, in Journ. Bot.* xvi. (1878) 83.—Am. austr.
bakianum, M. Roem. Syn. Ensat. 138 = rutilum.
barbatum, Herb. Amaryll. 138, 409. t. 21. f. 1 = equestre.
Batemanni x, Herb. l. c. 163.
Benthami, *Herb. l. c.*—Hab. ?
Berteroanum, *Baker, in Journ. Bot.* xvi. (1878) 83.—Chili.
bicolor, *Baker, l. c.*—Chili.
bifidum, *Baker, l. c.*—Reg. Argent.
brasilicum, M. Roem. Syn. Ensat. 138 = rutilum?
breviflorum, *Herb. Amaryll.* 137. t. 21. f. 4.—Reg. Argent.
Brookesii, *Herb. l. c.* 143.—Hab. ?
bulbulosum, Herb. l. c. 139 = rutilum.
calyptratum, *Herb. App.* 21.—Bras.
cearense, *Baker, in Journ. Bot.* xvi. (1878) 83.—Bras.
chilense, *Baker, l. c.*—Chili; Peruv.
concolor, Baker, l. c. = Zephyranthes concolor.
coruscum, Herb. App. 21 = rutilum.
Cybister, *Benth. & Hook. f. Gen.* iii. 725.—Bolivia.
decoratum, *Lem. Jard. Fleur.* iv. (1854) t. 388.—Bras.
equestre, *Herb. App.* 21.—Mexic.
formosissimum, Loud. Encyc. Pl. 1542 = Sprekelia formosissima.
franckeanum, *Baker, in Journ. Bot.* xvi. (1878) 82.—Reg. Argent.
fulgidum, Herb. App. 21 = rutilum.
glaucescens, Herb. Amaryll. 139 = rutilum.
gracilifolium, *Baker, in Journ. Bot.* xvi. (1878) 82.—Uruguay.
Grahami, *Herb. Amaryll.* 143.—Hab. ?
Gravisiae x, M. Roem. Syn. Ensat. 137.
Herbertianum, *Baker, in Journ. Bot.* xvi. (1878) 83.—Chili.
Houwtinnum, Karst. Fl. Columb. ii. 2. t. 102 = aulicum.
iguiscense, *Regel, Gartenfl.* (1874) 147.—N. Granat.
Jameseoni, *Baker, in Journ. Bot.* xvi. (1878) 83.—Reg. Argent.
latifolium, M. Roem. Syn. Ensat. 140 = rutilum.
Leopoldi, *Hort. ex Gard. Chron.* (1870) 723; *et Baker*, in *Journ. Bot.* xvi. (1878) 84.—Peruv.
linastum, *Baker, l. c.* 82.—Chili.
Martianaeum, M. Roem. Syn. Ensat. 134 = rutilum.
miniatum, *Herb. App.* 21.—Peruv.
modestum, *Baker, in Journ. Bot.* xvi. (1878) 83.—Chili.
montanum, *Baker, l. c.* 84.—Chili.
occidentale, M. Roem. Syn. Ensat. 132 = equestre.
organense, *Herb. in Bot. Mag.* sub t. 3808.—Bras.
pardinum, *Lem. Illustr. Hortic.* xiv. (1867) Misc. 44.—Peruv.
phycelloides, *Baker, in Journ. Bot.* xvi. (1878) 83.—Chili.
pratense, *Baker, l. c.* 84.—Chili.

HIPPASTRUM:—
procerum, *Lem. Illustr. Hortic.* xi. (1864) t. 408.—Bras.
proliferum, Herb. in Bot. Mag. sub t. 2475 = rutilum.
pronum, *C. Koch, in Wochenschr.* vii. (1864) 37.—Venezuela.
pulticaulum, *Herb. App.* 21.—Bras.
pulverulentum, Herb. in Bot. Mag. t. 2273 = rutilum.
pyrrochroum, Lem. Illustr. Hortic. xi. (1864) p. 490. t. 408 = equestre.
Reginae, *Herb. App.* 21.—Mexic.
regium, Herb. Amaryll. 139 = prasc.
reticulatum, *Herb. in Bot. Mag.* sub t. 2475.—Bras.
Rhodolirion, *Baker, in Journ. Bot.* xvi. (1878) 84.—Chili.
robustum, A. Dietr. in Walp. Ann. iii. 616 = aulicum.
Kortzi, *Baker, in Journ. Bot.* xvi. (1878) 85.—Bolivia.
roseum, *Baker, l. c.* 82.—Chili; Peruv.
rutilum, *Herb. App.* 21.—Bras.
Sweetianum, Herb. l. c. 140 = rutilum.
solandriiflorum, *Herb. l. c.* 31.—Bras.
spathaceum, Sims, Bot. Mag. t. 2315 = equestre.
spectabile, M. Roem. Syn. Ensat. 136 = Reginae.
splendens x, Herb. App. 52. t. 4.
stenopetalum, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 193.—Peruv.
striatifolium, Herb. in Bot. Mag. t. 2475 = reticulatum.
stylosum, *Herb. l. c.* t. 2276.—Guian.
subbarbatum, Herb. l. c. t. 2475 = rutilum.
sylvaticum, *Baker, in Journ. Bot.* xvi. (1878) 81.—Bras.
texanum, *Baker, l. c.* 82.—Am. bor. occ.
tubispathum, *Baker, l. c.*—Reg. Argent.
unguiculatum, M. Roem. Syn. Ensat. 133 = rutilum.
uniflorum, *Baker, in Journ. Bot.* xvi. (1878) 83.—Chili.
versicolor, *Baker, l. c.* 82.—Reg. Argent.
vittatum, *Herb. App.* 21.—Peruv.
Warscewiczianum, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 1.—Bolivia.

HIPPIA, Linn. Mant. ii. 158 (1767). *COMPOSITAE*, Benth. & Hook. f. ii. 484.
bicolor, Sm. in Rees, Cycl. xviii. n. 2 = Dichrocephala latifolia.
frutescens, *Linn. Mant.* ii. 261.—Afr. austr.
gracilis, *Less. in Linnaea*, vi. (1831) 213.—Afr. austr.
hirsuta, *DC. Prod.* vi. 144.—Afr. austr.
integrifolia, *Less. Syn. Comp.* 268.—Afr. austr.
integrifolia, *Linn. f. Suppl.* 389 (Quid?).—' India.'
odora, Salisb. Prod. 211 = frutescens.
repens, DC. Prod. vi. 144 = gracilis.
Vahliana, Steud. Nom. ed. II. i. 768 = integrifolia, *Less.*

HIPPIA, Linn. f. Suppl. 389 (1781) partim; H. B. & K. Nov. Gen. et Sp. iv. 301 (1820) = **Plagiocheilus**, Arn. (Compos.).
bogotensis, H. B. & K. l. c. = P. bogotensis.
minuta, Linn. f. Suppl. 389 = Cotula pygmaea.
minuta, Mutis, ex DC. Prod. vi. 148 = Soliva Mutisii.
pedunculoris, H. B. & K. Nov. Gen. et Sp. iv. 301 = P. pedunculoris.
stolonifera, Brot. Fl. Lusit. i. 372 = Soliva nasturtiifolia.

HIPPION, F. W. Schmidt, Fl. Boëm. ii. 16 (1793); et in Roem. Arch. i. 1. 9. t. 4 ad 5 (1796) = **Gentiana**, Tourn.
acaulescens, F. W. Schmidt, Fl. Boëm. ii. 20 = Gentiana verna.
alpinum, F. W. Schmidt, in Roem. Arch. i. 1. 10, 14 = Gentiana acaulis.
altaicum, F. W. Schmidt, l. c. = G. altaica.
Amarella, F. W. Schmidt, Fl. Boëm. ii. 26 = G. germanica.
aphyllum, F. W. Schmidt, in Roem. Arch. i. 1. 11 = Cyrtia aphylla.
aquaticum, F. W. Schmidt, l. c. 10 = G. aquatica.
aureum, F. W. Schmidt, l. c. 19, 17 = G. saxea.
auriculatum, F. W. Schmidt, Fl. Boëm. ii. 26 = G. campestris.
auriculatum, F. W. Schmidt, in Roem. Arch. i. 1. 11 = G. auriculata.

HIPPION :—
axillare, F. W. Schmidt, Fl. Boëm. ii. 29 = Gentiana Amarella.
bavaricum, F. W. Schmidt, l. c. 21 = G. bavarica.
Cachaishuan, F. W. Schmidt, in Roem. Arch. i. 1. 11 = Erythraea chilensis.
campestre, F. W. Schmidt, Fl. Boëm. ii. 25 = G. campestris.
ciliatum, F. W. Schmidt, l. c. 26 = G. ciliata.
cruciatum, F. W. Schmidt, l. c. 16 = G. cruciata.
dentosum, F. W. Schmidt, in Roem. Arch. i. 19 = G. ciliata.
dichotomum, F. W. Schmidt, l. c. 11 = G. tenella.
diffusum, Spreng. Syst. iv. Cur. Post. 340 = Curtia diffusa.
elongatum, F. W. Schmidt, in Roem. Arch. i. 1. 11, 19 = G. verna.
exacutum, F. W. Schmidt, l. c. 11 = Belmontia cordata.
filiforme, F. W. Schmidt, l. c. = Microcala filiformis.
Gentianella, F. W. Schmidt, Fl. Boëm. ii. 29 = G. Amarella.
heteroclita, F. W. Schmidt, in Roem. Arch. i. 1. 11 = Canacora sessiliflora.
imbricatum, F. W. Schmidt, Fl. Boëm. i. 18 = G. prostrata.
involueratum, F. W. Schmidt, in Roem. Arch. i. 1. 11 = G. aurea.
islandicum, F. W. Schmidt, l. c. = G. tenella.
lancifolium, Pers. Syn. i. 286 = G. Amarella.
longepedunculatum, F. W. Schmidt, in Roem. Arch. i. 1. 11, 21 = G. tenella.
macrophyllum, F. W. Schmidt, l. c. 10 = G. macrophylla.
nanum, F. W. Schmidt, l. c. 11, 19 = G. tenella.
nivale, F. W. Schmidt, l. c. 10, 18 = G. nivalis.
obtusifolium, F. W. Schmidt, Fl. Boem. ii. 27 = G. obtusifolia.
patulum, Spreng. Syst. iv. Cur. Post. 340 (= *Schuebleria patula*!)—Bras.
Porphyris, F. W. Schmidt, in Roem. Arch. i. 1. 11 = G. angustifolia.
prostratum, F. W. Schmidt, l. c. = G. prostrata.
pulchellum, F. W. Schmidt, l. c. 10, nomen = Erythraea ramosissima ?
pumilum, F. W. Schmidt, l. c. 10, 17 = G. prostrata.
pusillum, F. W. Schmidt, l. c. = G. imbricata.
pyrenaicum, F. W. Schmidt, l. c. 10, 14 = G. pyrenaica.
quinquefolium, F. W. Schmidt, l. c. 11 = G. quinquefolia.
scilloides, F. W. Schmidt, l. c. = Erythraea Massoni.
sessile, F. W. Schmidt, l. c. (= *Chlora sessilis*).—Reg. Argent.
sexfidum, F. W. Schmidt, l. c. 10, 15 = G. vera.
spicatum, F. W. Schmidt, l. c. 11 = Erythraea spicata.
tenellum, F. W. Schmidt, l. c. = G. tenella.
utriculosum, F. W. Schmidt, Fl. Boëm. ii. 22 = G. utriculosa.
vernum, F. W. Schmidt, l. c. 19 = G. verna.
viscosum, Spreng. Syst. i. 589 = Ixanthus viscosus.

HIPPION, F. W. Schmidt, in Roem. Arch. i. 1. (1796) § partim ; Spreng. Syst. i. 505 (1825), partim = Zinlocostema, Blume (1826) (Gentian.).
controversum, Spreng. Syst. iv. Cur. Post. 340 = E. littorale.
hyssopifolium, Spreng. l. c. i. 589 = E. littorale.
littorale, Miq. Fl. Ind. Bat. ii. 559 = E. littorale.
maritimum, F. W. Schmidt, in Roem. Arch. i. 1. 10 = E. littorale.
orientale, Dals. & Gibs. Bomb. Fl. 157 = E. littorale.
verticillare, F. W. Schmidt, in Roem. Arch. i. 1. 11 = E. littorale.
verticillatum, F. W. Schmidt, l. c. ; Spreng. Syst. i. 589 = E. littorale.

HIPPOBROMA, G. Don, Gen. Syst. iii. 717 (1834) = Isotoma, Lindl. (Campan.).
brevifolia, G. Don, in Sweet, Hort. Brit. ed. III. 767 = I. brevifolia.
longiflora, G. Don, Gen. Syst. iii. 717 = I. longiflora.

HIPPOBROMA, Lindl. Veg. Kingd. 885 (1847) = seq.

HIPPOBROMUS, Eckl. & Zeyh. Enum. 151 (1836). *SAPINDACEAE*, Benth. & Hook. f. i. 402.
alatus, *Eckl. & Zeyh. l. c.*—Afr. austr.

HIPPOCASTANUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 291 (1745) = Aesculus, Linn. (Sapind.).
Aesculus, Cav. Descr. 458 = Ae. Hippocastanum.
barbatum, Rafin. Alsog. Am. 70 = Ae. barbata.
carneum, Rafin. l. c. = Ae. glabra.
laevigatum, Rafin. l. c. = Ae. laevigata.
rubicundum, Rafin. l. c. = Ae. glabra.
vulgare, Gaertn. Fruct. ii. 285. t. 9 = Ae. Hippocastanum.

HIPPOCENTAUREA, Schult. Oesterr. Fl. ed. II. i. 388 (1814) = Erythraea, Rich. (Gentian.).
Centaurium, Schult. l. c. = E. Centaurium.
emarginata, Schult. l. c. 390 = E. ramosissima.
fragrans, Schult. l. c. = E. ramosissima ?
pulchella, Schult. l. c. 389 = E. ramosissima.
uliginosa, Schult. l. c. = E. linarifolia.

HIPPOCRATEA, Linn. Gen. ed. I. 368 (1737). *CELASTRINEAE*, Benth. & Hook. f. i. 369.
CURVEA, Triana, ex Miers, in Trans. Linn. Soc. xxvii. (1872) 370.
PERESKIA, Vell. Fl. Flum. 84 (1825) ; l. t. 81 (1827).
PRESTIMERA, Miers, in Trans. Linn. Soc. xxvii. (1872) 360.
ROMUALDEA, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvi. (1872) 370.
TYLODERMA, Miers, in Trans. Linn. Soc. xxvii. (1872) 413.
acapulcensis, Hook. & Arn. Bot. Beech. Voy. 280 = Tontelea Hookeriana.
acapulcensis, *H. B. & K. Nov. Gen. et Sp.* v. 136.—N. Granat.
acuminata, Hoffmgg. ex Link, in Jahrb. i. III. (1820) 68 = malpighiaefolia.
acutiflora, [*Mey. & Szyszl.*] *ex DC. Prod.* i. 568.—Mexic.
affinis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 108 = ovata.
aggregata, *Peyr. in Mart. Fl. Bras.* xi. i. 137.—Bras.
amligua, *Peyr. l. c.* 138.—Guian.
andongensis, *Welw. ex Oliver, Fl. Trop. Afr.* i. 370.—Afr. trop.
angulata, Griff. Notul. iv. 473 = Eucnymus Griffithii.
Anthodon, Pers. Syn. i. 41 = decussata.
apiculata, *Welw. ex Oliver, Fl. Trop. Afr.* i. 369.—Afr. trop.
apocynoides, *Welw. ex Oliver, l. c.* 368.—Afr. trop.
arborea, *Roxb. Hort. Beng.* 5 ; *Pl. Corom.* iii. 3. t. 205. —Reg. Himal.
Arnottiana, *Wight, Illustr.* i. 133.—Ind. or.
aspera, *Lam. Illustr.* i. 101.—Guian.
Aubletiana, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 382.—Am. trop.
barbata, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 28 = obtusifolia.
bilobicarpa, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 349.—Bras.
Bojeri, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 92. —Malacc.
Bonplandiana, *Peyr. in Sitzb. Wien. Acad.* lxx. *Abth.* i. (1875) 407.—Am. austr.
campestris, *Peyr. in Mart. Fl. Bras.* xi. i. 133.—Bras.
casinoides, *DC. Prod.* i. 569.—Ins. Timor.
cearensis, Miers, in Trans. Linn. Soc. xxvii. (1872) 345 = ovata.
celastroides, *H. B. & K. Nov. Gen. et Sp.* v. 136.—Mexic.
comosa, *Peyr. in Mart. Fl. Bras.* xi. i. 138 = flaccida.
comosa, *Sw. Prod. Veg. Ind. Occ.* 17.—Ins. S. Doming.
coptosiflora, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 341.—Am. trop.
coriacea, *Wright, ex Griseb. Cat. Pl. Cub.* 282.—Cuba.
Cumingii, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 624.—Ind. or.
decussata, *Peyr. in Mart. Fl. Bras.* xi. i. 136.—Bras.
diffusa, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 346. —Bras.
discolor, G. F. W. Mey. Prim. Fl. Esseq. 19 = ovata.
discolor, Seem. Bot. Voy. Herald, 90 = versicolor.
disperma, Vahl, Enum. ii. 26 = indica.
disperma, Wall. Cat. n. 4213 B = arborea.
divaricata, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 346.—Bras.
elliptica. *H. B. & K. Nov. Gen. et Sp.* v. 138.—Mexic.
emarginata, Rudge, Fl. Guian. 10. t. 8 = Salacia multiflora, *DC.*
ensomoides, Vahl, Enum. ii. 26 = indica.
erosomoides, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 91.—Madag.

HIPPOCRATEA :—
exsolas, *H. B. & K. Nov. Gen. et Sp.* v. 135.—Mexic
festiva, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 348. —Bras.
flaccida, *Peyr. in Mart. Fl. Bras.* xi. i. 132.—Bras.
flotibunda, *Benth. Bot Voy. Sulph.* 78.—Ins. Gorgona.
fuscescens, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 200.—Burma.
glaga, *Korth. in Flora*, xxxi. (1848) 580.—Borneo.
gracilifiora, *Welw. ex Oliver, Fl. Trop. Afr.* i. 371.—Afr. trop.
Grahamii, *Wight, Illustr.* i. 134.—Ind. or.
granateoula, *Peyr. in Mart. Fl. Bras.* xi. i. 137, in obs. —N. Granat.
grandiflora, Payen, Organogr. 163. t. 35. f. 84-44 = Pricnostemma scabridula.
grandiflora, Wall. Cat. n. 4213 = obtusifolia.
Hasseltiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 154. —N. Guinea.
Hilariana, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 345.—Bras.
indica, *Willd. Sp. Pl.* i. 1193.—Geront. trop.
infausta, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 352.—Bras.
integrifolia, *A. Rich. Ess. Fl. Cub.* 252.—Ins. Cuba.
integrifolia, *Seem. Bot. Voy. Herald*, 90.—Panama.
inundata, *Mart. ex Peyr. in Mart. Fl. Bras.* xi. i. 136. —Bras.
Kirkii, *Welw. ex Oliver, Fl. Trop. Afr.* i. 370.—Afr. trop.
Klapperiana, *Miq. in Linnaea*, xxvi. (1853) 220.—Guian.
laevigata, Rich. in Vahl, Enum. ii. 27 = ovata.
lanceolata, Buch-Ham. ex Wall. Cat. n. 4214 = Grahamii.
Lobbii, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 624.—Burma.
lonchophylla, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 348.—Peruv.
longipes, *Oliver, Fl. Trop. Afr.* i. 367.—Afr. trop.
longipetiolata, *Oliver, l. c.* 372.—Afr. trop.
macrantha, Korth. in Verh. Nat. Gesch. Bot. 187. t. 39 = obtusifolia.
macrophylla, *Vahl, Enum.* ii. 28.—Afr. trop.
madagascariensis, Boj. ex Tul. in Ann. Sc. Nat. Sér IV. viii. (1857) 99 = Bojeri.
madagascariensis, Lam. Illustr. i. 101 = Salacia madagascariensis.
Maingayi, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 625. —Malacc.
malpighiaefolia, *Rudge, Pl. Guian.* 10. t. 8.—Guian. ; Ind. occ.
megacarpa, *Peyr. in Mart. Fl. Bras.* xi. i. 137.—Bras.
meticrora, Miers, in Trans. Linn. Soc. xxvii. (1872) 352 = uniflora.
micrantha, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 108. —Bras.
multiflora, Lam. Illustr. i. 101 = Salacia multiflora, *DC.*
myriantha, *Oliver, Fl. Trop. Afr.* i. 369.—Afr. trop.
nicoharica, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 208.—Ins. Nicobar.
obcordata, *Lam. Illustr.* i. 100. t. 28. f. 1.—Am. austr.
oblongata, *Soland. ex Miers, in Trans. Linn Soc.* xxvii. (1872) 352.—Ind. occ.
obovata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Salacia multiflora, *DC.*
obtusifolia, R. Grah. ex Wight, Illustr. i. 134 = Grahamii.
obtusifolia, *Roxb. Hort. Beng.* 5 ; *Fl. Ind.* i. 166.—Geront. trop.
obtusifolia, Wall. Cat. n. 4211 = indica.
ovalifolia, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 352.—Cuba.
ovata, Griseb. Fl. Brit. W. Ind. 148 = pallidula.
ovata, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 173 = ovalifolia.
ovata, *Lam. Illustr.* i. 100. t. 28. f. 2.—Am. austr.
pallens, *Planch. ex Oliver, Fl. Trop. Afr.* i. 367.—Afr. trop.
pallidula, *Miers, in Trans. Linn. Soc.* xxvii. (1872) 348.—Ind. occ.
paniculata, *Vahl, Enum.* ii. 28.—Afr. trop.
parvifolia, *Oliver, Fl. Trop. Afr.* i. 368.—Afr. trop.
pauciflora, *DC. Prod.* i. 569.—Ins. Timor.
pentandra, *Grif. Notul.* iv. 472.—Burma.
Pereskia, *Steud. Nom.* ed. II. i. 769.—Bras.

HIPPOCRATEA:—
perspicua, *Miers, in Trans. Linn. Soc.* xxviii. (1872) 245.—Ferrv.
Flumieri, *Miers, l. c.* 356.—Ind. occ.
Plumieri, Rich. ex Peyr. in Mart. Fl. Bras. xi. 1. 130 = ovata.
Richardii, Steud. Nom. ed. II. i. 768 = acq.
Richardiana, Cambess. in St. Hil. Fl. Bras. Mer. ii. 109, in adnot. = obtusifolia.
Riedeliana, *Peyr. in Mart. Fl. Bras.* xi. 1. 132.—Bras.
rigida, Hamp. ex M. Laws. in Hook. f. Fl. Brit. Ind. i. 624 = obtusifolia.
rigida, *Span. in Linnaea*, xv. (1841) 178.—Ins. Timor.
rotundifolia, Hook. f. in Hook. Niger Fl. 279 = macrophylla.
salicifolia, *Korth. in Flora*, xxxi. (1848) 580.—Sumatra.
scandens, Griseb. Cat. Pl. Cub. 56 = Vahliana, *Peyr.*
scandens, Jacq. Select. Am. 9. t. 9 = oboordata.
scandens, Turp. Atlas, t. 192 = ovata.
Schimperiana, Hochst. & Steud. in Flora, xxiv. (1841) i. Intell. 28 = obtusifolia.
Schomburgkii, Klotzsch, in Schomb. Faun. & Fl. Gui. 1188, nomen; et ex Peyr. in Mart. Fl. Bras. xi. 1. 186 = aspera.
scutellata, Griseb. Fl. Brit. W. Ind. 148 = Vahliana, *Peyr.*
senegalensis, Lam. Illustr. i. 401 = Salacia senegalensis.
serrata, *Grisb. Notul.* iv. 473.—Burma.
serrulata, *Miers, in Trans. Linn. Soc.* xxviii. (1872) 244.—Am. trop.
serrulata, Pohl, ex Peyr. in Mart. Fl. Bras. xi. 1. 130 = ovata.
Swartziana, *Miers, in Trans. Linn. Soc.* xxviii. (1872) 238.—Ind. occ.
tenuiflora, *Mart. ex Peyr. in Mart. Fl. Bras.* xi. 1. 132. —Bras.
timorensis, *Span. in Linnaea*, xv. (1841) 178—Ins. Timor.
tortuosa, Wall. Cat. n. 6216 = obtusifolia.
uniflora, [*Moq. & Sesse*] *ex DC. Prod.* i. 567.—Mexic.
Urceolua, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 91.—Madag.
Vahliana, *Miers, in Trans. Linn. Soc.* xxviii. (1872) 237.—Am. trop.
Vahliana, *Peyr. in Mart. Fl. Bras.* xi. 1. 126.—Bras.
velutina, *Afzel. ex Spreng. Neue Entdeck.* iii. 226.—Afr. trop.
verrucosa, *Griseb. in Goett. Abh.* xxiv. (1879) 84.—Reg Argent.
verrucosa, *H. B. & K. Nov. Gen. et Sp.* v. 186.—N. Granat.
versicolor, *Miers, in Trans. Linn. Soc.* xxviii. (1872) 250.—Panama.
verticillata, Steud. Nom. ed. I. 410 = Salacia senegalensis.
viridis, *Ruiz & Pav. Fl. Per.* i. 44. t. 74. f. a.—Am. trop.
volubilis, Aubl. Pl. Gui. i. 80 = Ambletiana.
volubilis, Heyne, in Wall. Cat. n. 4215 = obtusifolia.
volubilis, Linn. Sp. Pl. 1191 = ovata.
volubilis, Sw. Obs. Bot. 28 = Swartziana.
Welwitschii, *Oliver, Fl. Trop. Afr.* i. 367.—Afr. trop.
Zippeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 158.—N. Guinea.

HIPPOCRATIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. 175 (1881) = prasc.

HIPPOCREPANDRA, Mnell. Arg. in Linnaea, xxxiv. (1865–66) 61 = **Monotaxis**, Bronga. (Euphorb.).
gracilis, Mnell. Arg. l. c. 62 = M. gracilis.
ericoides, Mnell. Arg. l. c. = M. grandiflora.
lurida, Mnell. Arg. l. c. 61 = M. lurida.
Neesiana, Mnell. Arg. l. c. 92 = M. grandiflora.

HIPPOCREPIS, Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 510.
FERUM-EQUINUM, Tourn. ex Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 370.
HIPPOCRIS, Rafin. Princ. Somiol. 27 (1814).
annua, Lag. Gen. et Sp. Nov. 28 = ciliata.
annularis, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 163 = bifora.
arenaria, *Scheele, in Linnaea*, xxi. (1848) 573.—Arab.
areolata, Desv. in Mém. Soc. Linn. Par. iv. (1827) 289 = ciliata.
atlantica, *Ball, in Journ. Bot.* xi. (1873) 307.—Marocc.

HIPPOCREPIS:—
atlantica, Dur. ex Munby, Cat. Pl. Algér. 10 = comosa.
balearica, Cav. ex Nym. Consp. 186 = valentina.
balearica, *Jacq Misc.* ii. 306.—Ins. Balear.
barbata, Lour. Fl. Cochinch. 459 = Desmodium polycarpum.
bicontorta, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 424 = cornigera, *Boiss.*
biflora, Jacq. ex Walp. Ann. ii. 407 = arenaria.
bifloris, *Spreng. Pugill.* ii. 78.—Graec.; As. occ.
Bourgaei, Nym. Consp. 186 = glauca.
ciliata, Del Campo, ex Willk. & Lange, Prod. Fl. Hisp. iii. 256 = scabra.
ciliata, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 178.—Reg. Mediterr.
comosa, Asso, Syn. Pl. Arag. 101 = glauca.
comosa, *Linn. Sp. Pl.* 744.—Europ.
comosa, Lucé. Fl. Ouil. 248 = Ornithopus perpusillus.
comosa, Willk. ex Scheele, in Linnaea, xxi. (1848) 574 = scabra.
constricta, *Kunze, Pugill. Pl. Prim.* 12.—Hab. ?
cornigera, *Boiss. Diagn.* Ser. I. ii. 102.—Afr. bor. occ.
cornigera, *Pari. Giorn. Bot. Ital.* i. 1. (1844) 307.—Arab.
dicarpa, Bieb. Fl. Taur. Cauc. iii. 460 = ciliata.
divaricata, Hochst. ex Boiss. Fl. Orient. ii. 185 = cornigera, *Boiss.*
eriocarpa, Boiss. Diagn. Ser. II. ii. 84 = squamata.
flexuosa, Zahlb. ex Host, Fl. Austr. ii. 345 = bifora.
Fontanesii, *Boiss. Elench.* 88.—Hispan.
glandulosa, *Presl, Fl. Sic.* i. p. xxiii.—Sicilia.
glauca, *Tenore, Fl. Nap.* ii. 155. t. 69.—Europ. austr.
helvetica, G. Don, in Loud. Hort. Brit. 308 = comosa.
humilis, Bisso, Fl. Filip. ed. I. 585 = Desmodium triflorum.
mimulosa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 18 = Aeschynomene indica.
minor, Munby, Fl. Algér. 80 = multisiliquosa.
monocarpa, Bieb. Fl. Taur. Cauc. iii. 480 = unisiliquosa.
multisiliquosa, *Linn. Sp. Pl.* 744.—Reg. Mediterr.
nautisiliquosa, Vill. Hist. Pl. Dauph. iii. 400 = ciliata.
perennis, Lam. Fl. Fr. ii. 687 = comosa.
prostrata, Boiss. Elench. 88 = comosa.
pusilla, *Turcz. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 347.—Sibir.
rhomboidea, Blanco, Fl. Filip. ed. I. 585 = Desmodium spirale.
Salzmanni, *Boiss. & Reut. Diagn.* Ser. I. ii. 101.—Reg. Medit. occ.
scabra, *DC. Prod.* ii. 312.—Reg. Medit. occ.
scabra, Salzm. ex Cos. Not. Crit. 106 = Salzmanni.
squamata, *Cosz. l. c.* 105.—Hispan.
unisiliquosa, *Linn. Sp. Pl.* 744.—Reg. Mediterr.
ustoriviæ, Georgi, Beschr. Rus. Reich. III. iv. 1178 = unisiliquosa.
valentina, *Boiss. Elench.* 88.—Hispan.
Willkommiana, Scheele, in Linnaea, xxi. (1848) 574 = scabra.

HIPPOCRIS, Rafin. Princ. Somiol. 27 (1814) = prasc.

HIPPODAMIA. Decne. in Rev. Hortic. Sér. III. ii. (1848) 464 = Solenophora, Benth. (Gesner.).
insignis, Decne. l. c. = S. insignis.

HIPPOGLOSSUM, Hasp. Handb. Orch. Kuhl. & Hasp. [t. 14] (1827) = **Ophrapetalum**, Lindl. (Orchid.).
umbellatum, Breda, l. c. = C. maculosum !

HIPPOGLOSSUM, Hartm. Handb. Scand. Fl. ed. II. (1820); ed. IV. 66 (1843) = **Mactenzia**, Roth (Borag.).
maritimum, Hartm. l. c. = M. maritima.

HIPPOMANE, Linn. Gen. ed. I. 368 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 333.
MANCINELLA, Tuss. Fl. Antil. iii. 91. t. 5 (1824).
MARCANILLA, Plum. ex Adans. Fam. ii. 364 (1763).
aucuparia, Cranz, Inst. i. 169 = Sapium aucuparium.
biglandulosa, Linn. Sp. Pl. ed. II. 1481 = Sapium aucuparium.
biglandulosa, Sw. Obs. Bot. 370 = Sapium Laurocerasus.
dioica, Roth, in Act. Lit. Univ. Hafn. i. (1778) 301 = H. Mancinella !
glandulosa, Linn. Sp. Pl. 1191 = Sapium aucuparium.
Manzanilla, Jacq. Select. Am. 260 = acq.
Mancinella, *Linn. Sp. Pl.* 1191.—Ind. occ.
spinosa, *Linn. l. c.* (sp. dub.).—Ind. occ.

HIPPOMANES, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 88 = prasc.

HIPPOMANICA, Molina, Sagg. Chil. 126 (1782) genus fabulosum = **Phase** spp. + Pemettya furiens, *Phil. f. incana*, Molina, l. c.—Chili.
sessilifolia, J. F. Gmel. Syst. 674 = *incana*.

HIPPOMARATHRUM, Hall. in Rupp. Fl. Jen. ed. II. 280 (1745) = **Sesseli**, Linn. (Umbellif.).
buchtormense, Link, Handb. i. 349 = S. buchtormense.
dichotomum, Link, l. c. = S. dichotomum.
leucospermum, Link, l. c. = S. leucospermum.
pelviforme, Gaertn. Mey. & Scherb. Fl. Wett. i. 413 = S. Hippomarathrum.
petraeum, Link, Handb. i. 349 = S. petraeum.
peucedanifolium, Link, l. c. = S. rigidum.
vulgare, Koehl. ex DC. Prod. iv. 144 = S. Hippomarathrum.

HIPPOMARATHRUM, Hoffmgg. & Link, Fl. Port. ii. 411 (1820). *UMBELLIFERAE*, Benth. & Hook. f. i. 888.
LOPHOCACHRYS, Koch, ex DC. Prod. iv. 238 (1830); Bertol. Fl. Ital. iii. 455 (1837).
SCHRANKIA, Fisch. & Mey. in Schrenk, Enum. Pl. Nov. 63 (1841).
TRACHYMARATHRUM, Turcz, in Flora, xvii. (1834) 844.
amplifolium, Ledeb. ex C. A. Mey. Vers. Pfl. Cauc. 131 = crispum.
Bocconi, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 74.—Sicilia.
Boissieri, *Reut. & Hausskn. ex Boiss. Fl. Orient.* ii. 838.—Syria
crassilobum, *Boiss. & Heldr. Diagn.* Ser. I. x. 56.—Graecia.
crispatum, *Pomel. Nouv. Mat. Fl. All.* 151.—Alger.
crispum, *Koch, in Nov. Act. Nat. Cur. xii.* (1824) 186. —As. Min.; Reg. Casp.
cristatum, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 75.—Europ. austr.
Fedtschenkoi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 608.—Turkestan.
Lilaeotis, *Koch, ex DC. Prod.* iv. 237.—Afr. bor.
pterochlaenum, Boiss. Fl. Orient. ii. 932 = cristatum.
pterochlaenum, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 74 = siculum !
pungens, *Boiss. ex Pomel, Nouv. Mat. Fl. All.* 152.—Afr. bor.
saravschanicum, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 608.—Turkestan.
scabrum, *Boiss. Fl. Orient.* ii. 933.—Syria.
siculum, *Heffmgg. & Link, Fl. Port.* ii. 411.—Reg. Mediterr.
vaginatum, *Benth. & Hook. f. Gen.* i. 888.—Sibir. altaic.

HIPPOPHAE, Linn. Syst. ed. I (1735). *ELAEAGNACEAE*, Benth. & Hook. f. iii. 204.
OLEASTER, Heist. Syst. 11, 16 (1748).
RHAMNOIDES, Tourn. ex Moench, Meth. 343 (1794).
argentea, Pursh, Fl. Am. Sept. i. 115 = Shepherdia argentea.
canadensis, Linn. Sp. Pl. 1024 = Shepherdia canadensis.
conferta, Wall. Cat. n. 6023 = salicifolia.
fasciculata, Wall. ex Steud. Nom. ed. II. i. 769.—Nepal.
Hitterala, Salisb. Prod. 71 = rhamnoides.
nepalensis, Hort. Angl. ex Steud. Nom. ed. II. i. 769 = salicifolia !
rhamnoides, *Linn. Sp. Pl.* 1023.—Europ.; As. temp.
salicifolia, *D. Don, Prod. Fl. Nep.* 68.—Nepal.
sibirica, Hort. ex Steud. Nom. ed. II. i. 770 = rhamnoides.
tibetana, Schlecht. in Linnaea, xxxii. (1863) 296 = rhamnoides.

HIPPOPHAES, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 88 = **Hippophaë**, Linn. (Elaeagn.).
rhamnoides, St. Lag. l. c. 197 = Hippophaë rhamnoides.

HIPPOPHAESTUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 448 (1821) = **Centaurea**, Linn. (Compos.).
vulgare, S. F. Gray, l. c. = C. Calcitrapa.

HIPPOSERIS, Cass. in Dict. Sc. Nat. xxxiii. 444 (1824) = **Crepsis**, DC. (Compos.).
acerifolia, Cass. l. c. = C. acerifolia.
salicifolia, Cass. l. c. = C. salicifolia.

HIPPOTHRONIA, Benth. Lab. Gen. et Sp. 120, in syn. (1833) = HYPOTHROMA, Schrank = Hyptis, Jacq. (Labiat.).

HIPPOTIS, Ruiz & Pav. Prod. Fl. Per. 88 (1794). *RUBIACEAE*, Benth. & Hook. f. ii. 70.

albiflora, *Karst. Fl. Columb.* i. t. 17.—N. Granat.

multiflora, Benth. ex Hook. f. in Ic. Pl. t. 1093 = Phitopis multiflora.

pyrifolia, Bert. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 360 = Psychotria pyrifolia.

trifora, Bert. in Ann. Sc. Nat. Sér. I. xxi. (1830) 347 = Coprosma trifora.

trifora, *Ruiz & Pav. Prod.* 33; *Fl. Per.* ii. 56.—Peruv.

HIPPOXYLON, Rafin. Sylva Tellur. 78 (1838) = Oroxylum, Vent. (Bignon.).

indica, Rafin. l. c. = O. indicum.

HIPPURIS, Linn. Syst. ed. I (1735). *HALORAGEAE*, Benth. & Hook. f. i. 675.

LIMNATRUCE, Adans. Fam. ii. 80 (1763).

Alopecuros, Bousch. ex Brouss. Elench. Pl. Hort. Monsp. 1804 (1805) 30, nomen = vulgaris.

Eschscholtzii, Cham. ex Ledeb. Fl. Ross. ii. 120 = vulgaris.

fluitans, Lilljebl. ex Hising. Fl. Fagervik, 32 = vulgaris.

fluviatilis, Hoffm. Deutschl. Fl. ed. II. i. 1 = vulgaris.

indica, Lour. Fl. Cochinch. 16 = Elecocharis plantagines.

lanceolata, Retz. Obs. iii. 7. t. 1 = vulgaris.

maritima, Hellen. in Usteri, Delect. Opusc. Bot. i. (1790) 11 = vulgaris.

montana, Ledeb. ex Reichb. Ic. Fl. Germ. i. 71. t. 86. f. 181 = vulgaris.

palustris, Gorter, Fl. Ingr. 1 = vulgaris.

polyphylla, Rafin. Fl. Ludov. 13 = vulgaris.

tetraphylla, Linn. f. Suppl. 81 = vulgaris.

verticillata, Gilib. Exercit. ii. 418 (err. typ. 518) = vulgaris.

vulgaris, *Linn. Sp. Pl.* 4.—Europ.: As. bor.: Am. bor. et austr.

HIPTAGE, Gaertn. Fruct. ii. 69. t. 116 (1791). *MALPIGHIACEAE*, Benth. & Hook. f. i. 258.

GAERTNERA, Schreb. Gen. i. 290 (1789).

MOLINA, Cav. Diss. ix. 435. t. 303 (1790).

SUCCONIA, Dennst. Schlüssel. Hort. Malab. 32 (1818).

acuminata, *Wall. Cat.* n. 1065.—Reg. Himal.

arborea, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 228.—Burma.

benghalensis, *Kurz, l. c.* xliii. (1874) ii. 136.—Burma.

candicans, *Hook. f. Fl. Brit. Ind.* i. 419.—Burma.

javanica, *Blume, Bijdr.* 224.—Java.

laurifolia, A. Juss. in Arch. Mus. Par. iii. (1843) 502 = parvifolia ?

Madablota, *Gaertn. Fruct.* ii. 169. t. 116.—As. trop.

myrtifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 267.—Ins. Fiji.

obtusifolia, *DC. Prod.* i. 583.—China.

parviflora, Wight, ex Hook. f. Fl. Brit. Ind. i. 419 = sericea.

parvifolia, *Wight & Arn. Prod.* 107.—Ind. or.

sericea, *Hook. f. Fl. Brit. Ind.* i. 419.—Malaya.

sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 512.—Sumatra.

trialata, Span. in Linnaea, xv. (1841) 179 = H. Madablota.

HIRAEA, Jacq. Enum. Pl. Carib. 4 (1760). *MALPIGHIACEAE*, Benth. & Hook. f. i. 260.

MARCAGRAVIA, Bert. ex Colla, Hort. Ripul. 86 (1824).

acuminata, [Mog. & Sesse,] ex DC. Prod. i. 586 = Gaudichaudia acuminata.

acutifolia, *H. B. & K. Nov. Gen. et Sp.* v. 169.—Venezuel.

affinis, *Miq, in Linnaea*, xix. (1847) 188.—Guian.

amblyura, *A. Juss. in Arch. Mus. Par.* iii. (1843) 555.—Bras.

anisopetala, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 16.—Bras.

argutea, *A. Juss. l. c.* 17. t. 168.—Bras.

Ashtonae, Moric. Pl. Nouv. Am. 107 = Salmasmiana.

Barclayana, *Benth. Bot. Voy. Sulph.* 78.—Am. centr.

Barretiae, *Bárcena, Not. Cient. Estad. Hidalg.* 86.—Mexic.

HIRAEA:—

bierosa, *Moric. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 559.—Bras.

Blanchetiana, *Moric. Pl. Nouv. Am.* 104.—Guian.

brachyptera, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1863) 382.—N. Granat.

bunchosioides, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 18 = Heladena bunchosioides.

candicans, A. Rich. in Arch. Mus. Par. iii. (1843) 455 = psilophylla.

Cephalocalyx, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 331.—N. Granat.

chlorocarpa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 560.—Bras.

chrysophylla, *A. Juss. l. c.* 572.—Guian.

cinerea, *Bunb. in Proc. Linn. Soc.* i. (1841) 109.—Bras.

complicata, *H. B. & K. Nov. Gen. et Sp.* v. 169.—Venezuela.

concava, Wall. Pl. As. Rar. i. 18 = Aspidopterys concava.

cordata, Heyne, ex Wall. l. c. = Aspidopterys cordata.

cordifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 19. t. 164.—Bras.

corniifolia, Willd. ex Griseb. in Mart. Fl. Bras. xii. t. 98, sphalm. = præc.

crassipes, *A. Juss. in Arch. Mus. Par.* iii. (1843) 568.—Bras.

crispa, A. Rich. l. c. 522 = Tetrapterys crispa.

cujabensis, *Griseb. in Mart. Fl. Bras.* xii. t. 100.—Bras.

cuneata, *Griseb. in Linnaea*, xlii. (1880) 246.—Bras.

cyclocarpa, Poepp. ex Griseb. in Mart. Fl. Bras. xii. t. 95 = elegans.

cyclopetala, Hook. & Arn. Bot. Beech. Voy. 380 = Gaudichaudia Arnottiana.

cyclopters, [*Mog. & Sesse,*] ex *DC. Prod.* i. 586.—Mexic.

dentulata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 565.—Bras.

divaricata, *H. B. & K. Nov. Gen. et Sp.* v. 169.—Venezuel.

elegans, *A. Juss. in Arch. Mus. Par.* iii. (1843) 549.—Peruv.

elliptica, Blume, Bijdr. 225 = Aspidopterys elliptica.

fagifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 567.—Guian.

Finlaysoniana, Wall. Cat. n. 7263 = Decaspermum paniculatum.

floribunda, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 22 = Heteropterys sceroides.

fluminensis, *Griseb. in Linnaea*, xlii. (1839) 243.—Bras.

fulgens, *A. Juss. in Arch. Mus. Par.* iii. (1843) 572.—Bras.

galphimioides, Grisch. ex Schomb. Faun. et Fl. Brit. Gui. 1096 = Tetrapterys marahamensis.

Gaudichaudiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 571.—Bras.

glabra, Lamk. Nouv Entdeck. ii. 154 = Tetrapterys punctulata.

glabriuscula, Wall. Cat. n. 6626. p. 260 = Aspidopterys Roxburghiana.

glaucescens, Lodd. ex G. Don, in Loud. Hort. Brit. 168 (Quid ?).—Ind. or.

grandifolia, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 136 = Tetrapterys marahamensis.

Greggii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 383.—Mexic.

heteropetala, A. Juss. in Arch. Mus. Par. iii. (1843) 557 = Tetrapterys heteropetala.

hippocrateoides, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 328.—N. Granat.

hirsuta, Wall. Pl. As. Rar. i. 18. t. 18 = Aspidopterys hirsuta.

Hookeriana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 570.—Ind. occ.

Houlettiana, A. Juss. l. c. 575 = Wiedeana.

hypericoides, Barch. ex DC. Prod. i. 585 = Triaspis hypericoides.

indica, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 448 = Aspidopterys Roxburghiana.

indica, Wall. Cat. n. 1057, partim = Aspidopterys nutans.

intermedia, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 21 = Heteropterys intermedia.

Jussieuana, *Miq. in Linnaea*, xix. (1847) 142.—Guian.

HIRAEA:—

Kunthiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 571.—N. Granat.

lanuginosa, Wall. Pl. As. Rar. i. 18 = Aspidopterys nutans.

laxiandra, A. Juss. in Arch. Mus. Par. iii. (1843) 557 = Tetrapterys mogorifolia.

laurifolia, A. Juss. l. c. 556 = Tetrapterys laurifolia.

laurifolia, Willd. ex Griseb. in Mart. Fl. Bras. xii. t. 80 = Tetrapterys mucronata.

leucopoala, Grisch. in Linnaea, xlii. (1839) 240 = Tetrapterys leucosepala.

lilacina, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 383.—Mexic.

longifolia, Rich. ex A. Rich. in Arch. Mus. Par. iii. (1843) 524 = Diplopterys Paralias.

macradina, *DC. Prod.* i. 585.—Am. austr.

macrodisca, *Planch. & Triana, in Ann. Sc. Nat. Sér.* IV. xvii. (1863) 326.—Am. austr.

macroptera, [Mog. & Sesse,] ex DC. Prod. i. 586 = septentrionalis ?

mandonellensis, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 117.—Panama.

maritima, *A. Juss. in Arch. Mus. Par.* iii. (1843) 562.—Bras.

mergurnis, Wight, Illustr. i. 139 (*merguiensis*) = Aspidopterys concava.

microphylla, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 16.—Bras.

Moricandiana, A. Juss. in Arch. Mus. Par. iii. (1843) 564 = cuneata.

mucronata, [Mog. & Sesse,] ex DC. Prod. i. 586 = Gaudichaudia filipendula.

mucronata, Rich. ex A. Rich. in Arch. Mus. Par. iii. (1843) 524 = Diplopterys Paralias.

multiflora, Griseb. in Linnaea, xlii. (1839) 239 = Tetrapterys mogorifolia.

multiflora, *Mtg. Stirp. Surin. Sel.* 88.—Guian.

multiradiata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 575.—Guian.

nepalensis, *Wall. ex Voigt, Hort. Suburb. Calc.* 170, nomen.—Nepal.

nitida, *H. B. & K. Nov. Gen. et Sp.* v. 169.—Venezuel.

nutans, Roxb. Hort. Beng. 84; Fl. Ind. ii. 447 = Aspidopterys nutans.

nutans, Wall. Pl. As. Rar. i. 18 = Aspidopterys Wallichii.

oblongifolia, *DC. Prod.* i. 585.—Am. austr.

obscura, Blume, Bijdr. 226 = Ryssopterys timoreusis.

odorata, Willd. Sp. Pl. ii. 743 = Triaspis odorata.

oleaefolia, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 136 = Tetrapterys olefolia.

orbiculata, Roxb. ex G. Don, Gen. Syst. i. 641 = Aspidopterys nutans.

ovata, Blume, Bijdr. 226 = Ryssopteris timoreusis.

ovatifolia, *H. B. & K. Nov. Gen. et Sp.* v. 169.—Venezuela.

oxyota, [Mog. & Sesse,] ex DC. Prod. i. 586 = Gaudichaudia oxyota.

oxyphylla, Wall. Cat. n. 7264 = Aspidopterys Roxburghiana.

panamensis, Griseb. in Mart. Fl. Bras. xii. t. 100 = Barclayana.

paniculata, Raeusch. Nom. ed. III. 131 = Flabellaria paniculata.

pauciflora, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 22 = Heteropteris paucifloru.

pinnata, Willd. Sp. Pl. ii. 749 = Flabellaria paniculata.

platypters, Griseb. in Linnaea, xlii. (1839) 244 = psilophylla.

platyrachis, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 328.—N. Granat.

poeppigiana, [Mog. & Sesse,] ex DC. Prod. i. 586 = Gaudichaudia podocarpa.

Poeppigiana, A. Juss. in Arch. Mus. Par. iii. (1843) 556 = Tetrapterys Poeppigiana.

polybotrya, *A. Juss. l. c.* 553.—Mexic.

prunifolia, *H. B. & K. Nov. Gen. et Sp.* v. 169.—Venezuela.

psilophylla, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 20.—Bras.

pubiflora, *A. Juss. l. c.* 14.—Bras.

reclinata, Blanco, Fl. Filip. ed. I. 378 (Hiptages sp.).—Ins. Philip.

rochiata, *Vaco. Enum. Pl. Carib.* 21.—N. Granat.

ruiziana, A. Juss. in Arch. Mus. Par. iii. (1843) 558 = Tetrapterys ruiziana.

Riedleyana, *A. Juss. l. c.* 569.—Guian.

HIRAEA:—
rigida, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 14.—Bras.
rosea, *Mig. Stirp. Surin. Sel.* 84.—Guian.
retusifolia, Kotth. Hort. Bang. [90]: Fl. Ind. ii. 448 = Aspidopterys nutans.
salicifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 18. —Bras.
Salzmanniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 566.—Bras.
schizoptera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 584.—Ins. S. Vincent.
septum, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 19. t. 165.—Bras.
septentrionalis, *A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 259.—Mexic.
sericea, *Engelm. ex A. Gray, Pl. Wright.* i. 87.— Mexic.
sericea, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 18 = Tetrapteryx sericea.
serrifolia, A. Juss. ex Mart. in Flora, xxiv. (1841) ii. Beibl. 104 = cujabensis.
Sinuiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 560. —Ind. occ.? Guian.
spicigera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 584.—Mexic.
Swartziana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 571.—N. Granat.
ternifolia, *A. Juss. l. c.* 575.—N. Granat.
tomentosa, Blume, Bijdr. 235 = Aspidopterys tomentosa.
violacea, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 327.—N. Granat.
Wiedeana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 574.—Bras.

HIRCULUS, Haw. Saxifr. Enum. 40 (1821) = **Saxifraga**, Linn.
flagellaris, Haw. l. c. = S. flagellaris.
propinquus, Haw. l. c. = S. Hirculus.
punctatus, Rafin. Fl. Tellur. ii. 69 = S. Hirculus.
ranunculoides, Haw. Saxifr. Enum. 40 = S. Hirculus.

HIRNELLIA, Cass. in Bull. Soc. Philom. (1820) 57 = **Myriocephalus**, Benth. (Compos.).
cotuloides, Cass. in Dict. Sc. Nat. vii. 229 (= Angianthus tomentosus ?).—Austral.

HIRPICIUM, Cass. in Bull. Soc. Philom. (1820) 27. *COMPOSITAE*, Benth. & Hook. f. ii. 460.
echinulatum, *Cass. l. c.*—Afr. austr.
Echinus, *Less. Syn. Comp.* 55.—Afr. austr.
integrifolium, *Less. l. c.* 53.—Afr. austr.

HIRSCHFELDIA, Moench, Meth. 264 (1794) = **Brassica**, Tourn. (Crucif.).
adpressa, Moench, l. c. = B. adpressa.
consobrina, Pomel, Mat. Fl. Afr. 12 = B. adpressa.
geniculata. Pomel, l. c. = B. adpressa.
heterophylla, Ann. Fl. Ther. vi. 484 = B. adpressa.
incana, Lowe, Man. Fl. Mad. i. 586 = B. adpressa.
inflexa, Presl, Fl. Sicil. i. 97 = B. incana.
integrifolia, Presl, l. c. = B. incana ?
varians, Pomel, Mat. Fl. Afr. 12 = B. varia ?

HIRTELLA, Linn. Hort. Cliff. 17 (1787). *ROSACEAE*, Benth. & Hook. f. i. 608.
CAUSEA, Scop. Introd. 210 (1777).
COMBUERNA, Ruiz & Pav. Prod. Fl. Per. 10. t. 2 (1794).
SPHENISTA, Rafin. Sylva Tellur. 90 (1838).
ZANZELA, Rafin. l. c. (1838).
acaynoensis, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 529.— Mexic.
aggregata, Poir. Encyc. Suppl. ii. 53 = hirsuta.
americana, Jacq. Select. Am. 8. t. 8 = triandra.
americana, *Linn. Sp. Pl.* 34.—Am. trop.
angustifolia, *Schott, in Spreng. Syst. iv. Cur. Post.* 341.—Bras.
apetala, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 308 = Mocquilea floribunda.
bicornis, *Mart. & Zucc. in Abh. Ahad. Muench.* i. (1832) 877.—Bras.
brachystachya, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 34.—Bras.
bracteata, *Mart. & Zucc. in Abh. Ahad. Muench.* i. (1832) 884.—Bras.
bractosa, *Steud. in Flora*, xxvi. (1843) 761.—Guian.
bullata, *Benth. in Hook. Journ. Bot.* ii. (1840) 214.— Guian.

HIRTELLA:—
conomensis, Spreng. Syst. iv. Cur. Post. 341 = Conepia canomensis.
castanea, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 528.— Mexic.
ciliata, *Mart. & Zucc. in Abh. Ahad. Muench.* i. (1832) 876.—Bras.
Cliffortia, *Vell. Fl. Flum.* 91 ; ii. 136.—Bras.
cognata, *Steud. in Flora*, xxvi. (1843) 761.—Guian.
corticosa, Mart. & Zucc. in Abh. Ahad. Muench. i. (1832) 883 = americana.
corymbosa, *Cham. & Schlecht. in Linnaea*, ii. (1827) 545.—Bras.
Cosmibuena, *Lem. Illustr.* ii. 114.—Peruv.
dodecandra, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 529.— Mexic.
elliptica, Presl, Symb. Bot. ii. 11 = triandra.
elongata, *Mart. & Zucc. in Abh. Ahad. Muench.* i. (1832) 884.—Bras.
eriandra, *Benth. in Hook. Journ. Bot.* ii. (1840) 217.— Am. trop.
filiformis, Presl, Symb. Bot. ii. 23 = americana.
floribunda, *Cham. & Schlecht. in Linnaea*, ii. (1827) 546.—Bras.
Gardneri, Benth. in Hook. Journ. Bot. ii. (1840) 216 = hebeclada.
glaberrima, Steud. in Flora, xxvi. (1843) 761 = Crudia oblique.
glandulosa, Spreng. Neue Entdeck. i. 308.—Bras.
Guainiae, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 31.—Bras.
hebeclada, *Moric. in DC. Prod.* ii. 530.—Bras.
hexandra, Willd. ex Roem. & Schult. Syst. v. 274 = americana.
hirsuta, *Lam. Illustr.* ii. 114.—Guian.
hispidula, *Miq. Stirp. Surin. Sel.* 27.—Guian.
indecora, Schott, ex Spreng. Syst. iv. App. 404 = triandra.
longifolia, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 31.—Bras.
macrophylla, Benth. *ex Hook. f. l. c.* 36.—Bras.
Martiana, *Hook. f. l. c.* 29.—Bras.
mollicoma, *H. B. & K. Nov. Gen. et Sp.* vii. 263.— N. Granat.
mollissima, Hemsl. Biol. Centr. Am. Bot. i. 366 = pracc.
nemorosa, *Hoffmgg. ex Roem. & Schult. Syst.* v. 274. —Bras.
nitida, Humb. & Bonpl. ex Roem. & Schult. l. c. = americana.
oblongifolia, DC. Prod. ii. 529 = americana.
octandra, Hoffmgg. ex Roem. & Schult. Syst. v. 274 = Moquilea Turiuva.
oleaefrontis, Poir. Encyc. Suppl. iii. 53 = Icacina senegalensis.
paniculata, Lam. Encyc. iii. 33 = triandra.
paniculata, Sw. Prod. Veg. Ind. Occ. 51 = hirsuta.
pendula, *Soland. ex Lam. Encyc.* iii. 134.—Am. austr.
peruviana, Pers. Syn. i. 250 = H. Coumbuena.
physophora, *Mart. & Zucc. in Abh. Ahad. Muench.* i. (1832) 874.—Bras.
pilosissima, *Mart. & Zucc. l. c.* 873.—Bras.
Pohlii, *Hook. f. in Mart. Fl. Bras.* xiv. ii. 33.—Bras.
polyandra, H. B. & K. Nov. Gen. et Sp. vi. 246 = Conepia Kunthiana.
porterioreisis, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 274 = rugosa.
prasina, *Segref. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 304.—Guian.
punctulata, Mlq. in Linnaea, xix. (1847) 499 = triandra.
racemosa, Lam. Encyc. iii. 133 = americana.
racemosa, Ruiz & Pav. Fl. Per. iii. 5 = H. Coumbuena.
Rouxiuae, *Klotzsch, in Schomb. Faun. et Fl. Brit. Gui.* 1108, nomen.—Guian.
rosea, [Moç. & Sesse,] ex DC. Prod. ii. 528 = americana.
rubra, Benth. in Hook. Journ. Bot. ii. (1840) 217 = ciliata.
rugosa, *Thunb. ex Pers. Syn.* i. 250.—Ind. occ.
scabrella, *Sprucc, ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 32.—Bras.
scabra, *Benth. in Hook. Journ. Bot.* ii. (1840) 218.— Guian.
scandens, Hoffmgg. ex Roem. & Schult. Syst. v. 274 = americana.
Sellowna, *Hook. f. in Mart. Fl. Bras.* xiv. ii. 29.— Bras.

HIRTELLA:—
silicia, *Griseb. Fl. Brit. W. Ind.* 229.—Ins. Trinitat.
Spruceі, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 31.—Bras.
strigulosa, *Steud. in Flora*, xxvi. (1843) 761.—Guian.
subscandens, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. ii. 29.—Bras.
tenticulata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 75.— Bras.
tomentella, *Schott, ex Spreng. Syst. iv. Cur. Post.* 341. —Bras.
triandra, *Sw. Prod. Veg. Ind. Occ.* 51.—Ind. occ.
violacea, *Steud. in Flora*, xxvi. (1843) 761.—Guian.
xanthoica, *Oliver, in Hook. Ic. Pl.* t. 1198.—Afr. trop.

HIRTELLIA, Dum. Anal. Fam. 40 (1829) = prace.

HIRTELLINA, Cass. in Dict. Sc. Nat. xlvii. 499, 511 (1827) t. 441 = **Stachelina**, Linn. (Compos.).
lanceolata, Cass. l. c. 441 = S. fruticosa.

HISBANCHE, Sparrm. ex Meisn. Gen. Comm. 275 (1848) = **Cytinus** ! Linn. (Legum.).

HISINGERA, Hellen. in Vet. Akad. Handl. Stockh. (1792) 29. t. 2 = **Xyloma**, Forst. (Lixin.).
Benthami, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 231, 232, 235 = X. Benthami.
calastrinea, Clos, l. c. 234 = X. nitidum.
ciliatifolia, Clos, l. c. 233 = X. ciliatifolium.
cinerea, Clos, l. c. = X. cinereum.
elegans, Clos, l. c. 234 = X. elegans.
elliptica, Clos, l. c. = X. ellipticum, *Hemsl.*
flexuosa, Clos, l. c. 235 = X. flexuosum.
grandifolia, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 332.—Ins. Philipp.
intermedia, Seem. Bot. Voy. Herald. 249 = X. intermedium.
japonica, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 60 = X. racemosum.
lucrea, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 225 = X. Benthami.
mexicana, Planch. ex Hemsl. Biol. Centr. Am. Bot. i. 57 = X. ellipticum.
nitida, Hellen. in Vet. Akad. Handl. Stockh. (1792) 32. t. 2 = X. nitidum.
nitida, Seem. Bot. Voy. Herald, 249 = X. Seemanni.
obovata, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 228 = X. obovatum.
Palluras, Clos, l. c. 237.—Hab.?
Prachie, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 332.—Bras.
prunifolia, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 227 = X. prunifolium.
puberula, Schlecht. ex Griseb. Fl. Brit. W. Ind. 21 = X. nitidum.
racemosa, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 169 = X. racemosum.
Rumea, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 225 = X. nitidum.
Salzmannii, Clos, l. c. 224 = X. Salzmanni.
spiculifera, Clos, l. c. 223 (*spiciflora*) = Flacourtia spiculifera.
Tulasnei, Clos, l. c. 234 = Flacourtia elegans, velutina.
Tweedieana, Clos, l. c. 236.—Bras.

HISPIDELLA, Barnad. ex Lam. Encyc. i. iii. 134 (1789). *COMPOSITAE*, Benth. & Hook. f. ii. 508.
SOLDEVILLA, Lag. Varied. de Clenc. ii. iv. (1806) 87.
Barnardesii, Cass. in Dict. Sc. Nat. xxi. 248 = hispanica.
hispanica, *Barnad. ex Lam. Encyc.* iii. 134.—Hispan.
Wolowitschi, Rupr. in Bull. Phys.-Math. Acad. Petersb. xiv. (1856) 256 = hispanica ?

HISSOPUS, Noces, in Usteri, Ann. Bot. v. (1792) 82 = **Hyssopus**, Tourn. (Labiat.).

HISUTSUA, DC. Prod. vi. 44 (1837) = **Boltonia**, L'Hérit. (Compos.).
canariensis, DC. l. c. = B. indica.
serrata, Hook. & Arn. Bot. Beech. Voy. 205 = B. indica.

HITCHENIA, Wall. in Trans. Med. Phys. Soc. Calc. vii. (1824) 215. *SCITAMINEAE*, Benth. & Hook. f. iii. 643.
Discnema, Voigt, Hort. Suburb. Calc. 566 (1845).
Careyana, *Benth. & Hook. f. Gen.* iii. 643.—Ind. or. glanca, *Wall. in Trans. Med. Phys. Soc. Calc.* vii. (1824) 215.—Burma.
Roxcoeana, Benth. & Hook. f. Gen. iii. 643 = Curcuma Roxcoeana.

HITCHINIA, Hook. f. Fl. Brit. Ind. iv. 78 (1885), lapsu = HUTCHINIA, Wight & Arn. = **Bouceroma**, Wight & Arn. (Asclep.).
indica, Hook. f. l. c. = B. Hutchinia.

HITCHINIA, Horan. Prod. Monog. Scit. 24 (1862) = **Hitchenia**, Wall. (Scitam.).

HITZERA, Klotzsch, in Peters, Reise Mossamb. Bot. 89 (1861) = **Commiphora**, Jacq. (Anacard.).
edulis, Klotzsch, l. c. = C. edulis.

HLADNICKIA, Meisn. Gen. 148 (1838) = HLADNIKIA, Reichb. = **Carum**, Linn. (Umbellif.).

HLADNICKIA, Steud. Nom. ed. II. i. 771 (1840) = seq.

HLADNIKIA, Koch, Syn. Fl. Germ. ed. I. 320 (1835) = **Pleurospermum**, Hoffm. (Umbellif.).
Cicutaria, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 73 = Eleutherospermum grandiflolium.
golacensis, Koch, Syn. Fl. Germ. ed. I. 320.—Europ. *Golaka*, Reichb. f. Ic. Fl. Germ. xxi. 96 = *golacensis*.

HLADNIKIA, Reichb. Ic. Fl. Crit. ix. 9. t. 825 (1851); et Fl. Germ. Excurs. 476 (1832) = Carum, Linn. (Umbell.).
pastinacifolia. Reichb. il. cc. = C. latifolium.

HOAREA, Sweet, Geran. t. 18 (1820) = **Pelargonium**, L'Hérit. (Geran.).
amoena, Sweet, Hort. Brit. ed. I. 76 = P. amoenum.
angustifolia, Eckl. & Zeyh. Enum. 61 = P. astragalifolium.
appendiculatum, Eckl. & Zeyh. l. c. 65 = P. appendiculatum.
astragalifolia, Eckl. & Zeyh. l. c. 64 = P. astragalifolium.
atra, Sweet, Geran. t. 72 = P. atrum.
atrofusca, Steud. Nom. ed. II. i. 771 = P. atrofuscum.
atrosanguinea, Sweet, Geran. t. 151 = P. atrosanguineum.
auriculata, Sweet, Hort. Brit. ed. I. 76 = P. auriculatum.
aurita, Eckl. & Zeyh. Enum. 60 = P. auritum.
barbata, Eckl. & Zeyh. l. c. 62 = P. barbatum.
bicolor, Sweet, Hort. Brit. ed. I. 76 = P. pilosum.
bifuga, Eckl. & Zeyh. Enum. 65 = P. bifugum.
bipinnatifida, Eckl. & Zeyh. l. c. 62 = P. bipinnatifidum.
blanda, Sweet, Hort. Brit. ed. I. 76 = P. pilosum.
bubonifolia, Sweet, l. c. 75 = P. bubonifolium.
campestris, Eckl. & Zeyh. Enum. 63 = P. campestre.
capitata, Sweet, Hort. Brit. ed. I. 76 = P. capitatum.
carinata, Sweet, Geran. t. 135 = P. nutans.
ciliata, Sweet, Hort. Brit. ed. I. 76; Eckl. & Zeyh. Enum. 61 = P. longifolium.
cocoinea, Sweet, Geran. t. 206 = P. hybridum.
coluteaefolia, Sweet, l. c. t. 311 = P. coluteaefolium.
Colvillii ×, Sweet, l. c. t. 260.
congesta, Sweet, l. c. t. 302 = P. congestum.
corymbiflora, Sweet, l. c. t. 18 = P. rapaceum.
depressa, Sweet, Hort. Brit. ed. I. 76 = P. longifolium.
dilata, Sweet, l. c. = P. dilutum.
dioica, Sweet, l. c. = P. atrum.
elegans, Sweet, Geran. t. 135 = P. Sweetianum.
ensata, Eckl. & Zeyh. Enum. 61 = P. ensatum.
erythrophylla, Eckl. & Zeyh. l. c. 60 = P. erythrophyllum.
cupatoriaefolia, Eckl. & Zeyh. l. c. 64 = P. eupatoriaefolium.
fasciculata, Eckl. & Zeyh. l. c. 62 = P. fasciculatum.
fasciculifolia, Eckl. & Zeyh. l. c. 64 = P. fasciculifolium.
fastuosa, Colville, ex Sweet, Hort. Brit. ed. I. 76 = P. fasciculatum.
fulgidifolia ×, Sweet, l. c. nomen

HOAREA:—
gracilis, Eckl. & Zeyh. Enum. 64 = Pelargonium gracile.
hedysarifolia, Sweet, Geran. t. 355 = P. hedysarifolium.
heterophylla, Eckl. & Zeyh. Enum. 63 = P. variifolium.
hirsuta, Sweet, Hort. Brit. ed. I. 76; Eckl. & Zeyh. l. c. = P. atrum.
incrassata, Sweet, l. c. 75 = P. incrassatum.
intermixta, Sweet, l. c. 76 = P. intermixtum.
intertexta, Steud. Nom. ed. II. i. 771 = P. intertextum.
Jenkinsoni, Colville, ex Sweet, Hort. Brit. ed. I. 76 = P. Jenkinsoni.
labyrinthica, Sweet, Geran. t. 276 = P. labyrinthicum.
laciniata, Sweet, Hort. Brit. ed. I. 75; Eckl. & Zeyh. Enum. 62 = P. laciniatum.
lanceifolia, Eckl. & Zeyh. l. c. 61 = P. lanceifolium.
Lewana, Sweet, Geran. t. 258 = P. Lecanum.
lessertiaefolium, Eckl. & Zeyh. Enum. 64 = P. astragalifolium.
lilacina, Sweet, Hort. Brit. ed. I. 76 = P. lilacinum.
linearis, Eckl. & Zeyh. Enum. 62 = P. lineare.
literata, Sweet, Hort. Brit. ed. II. 91, nomen.—Hab.?
longifera, Sweet, l. c. 76 = P. longiflolium.
longifolia, Sweet, l. c. = P. longifolium.
lutea, Sweet, l. c. = P. luteum.
magica, Colville, ex Sweet, l. c. nomen.—Hab.?
margarita, Sweet, l. c. nomen = P. marginatum.
melanantha, Sweet, Geran. t. 73 = P. atrum.
nervifolia, Sweet, Hort. Brit. ed. I. 76 = P. nervifolium.
nigricans, Colville, ex Sweet, l. c. nomen = P. nigrescens.
nivea, Sweet, Geran. t. 182 = P. niveum.
nummulariifolia, Sweet, Hort. Brit. ed. I. 76 = P. nummulariifolium.
nutans, Sweet, Geran. t. 135 = P. nutans.
ornithopifolia, Eckl. & Zeyh. Enum. 64 = P. ornithopifolium.
orobifolia, Sweet, Geran. t. 306 = P. orobifolium.
ovalifolia, Sweet, l. c. t. 106 = P. ovalifolium.
parnassioides, Eckl. & Zeyh. Enum. 60 = P. parnassioides.
patens, Sweet, Hort. Brit. ed. I. 76 = P. patens.
pectinata, Eckl. & Zeyh. Enum. 61 = P. reticulatum.
pennsiformis, Sweet, Hort. Brit. ed. I. 75; Eckl. & Zeyh. Enum. 62 = P. penniforme.
pilosa, Sweet, l. c. 76 = P. pilosum.
pilosellaefolia, Eckl. & Zeyh. Enum. 61 = P. pilosellaefolium.
pinnata, Eckl. & Zeyh. l. c. 64 = P. pinnatum.
prolifera, Eckl. & Zeyh. l. c. 62 = P. barbatum.
pulchella, Sweet, Hort. Brit. ed. I. 76 = P. pulchellum.
purpurascens, Sweet, l. c. 75; Eckl. & Zeyh. Enum. 62 = P. longifolium.
radiata, Eckl. & Zeyh. l. c. = P. radiatum.
radicata, Sweet, Geran. t. 174 = P. radicatum.
rapacea, Sweet, Hort. Brit. ed. I. 75; Eckl. & Zeyh. Enum. 65 = P. rapaceum.
reticulata, Sweet, Geran. t. 91 = P. reticulatum.
retusa, Sweet, Hort. Brit. ed. III. 110 = P. retusum.
revoluta, Eckl. & Zeyh. Enum. 60 = P. revolutum.
rhoradifolia, Eckl. & Zeyh. l. c. 63 = P. rhoradifolium.
rosea, Sweet, Geran. t. 302 = P. condensatum.
rubescens, Sweet, Hort. Brit. ed. I. 76 = P. rubescens.
salmifolia, Sweet, Geran. t. 189 = P. sellnifolium.
setosa, Sweet, l. c. t. 88; Eckl. & Zeyh. Enum. 63 = P. setosum.
sicyonifolia, Sweet, l. c. t. 292 = P. sicyonifolium.
strigosa, Eckl. & Zeyh. Enum. 61 = P. strigosum.
superba, Sweet, Hort. Brit. ed. I. 76 = P. superbum.
tenuifolia, Sweet, l. c. = P. tenuiflolium.
thiemiana, Eckl. & Zeyh. Enum. 63 = P. ochroleucum.
trifoliolata, Eckl. & Zeyh. l. c. 64 = P. astragalifolium.
triphylla, Sweet, Hort. Brit. ed. I. 76 = P. triphyllum.
undulaefiora, Sweet, Geran. t. 263 = P. undulaefiorum.
unguiculata, Sweet, Hort. Brit. ed. I. 76 = P. unguiculatum.
varia, Sweet, Geran. t. 166 = P. variam.
venosa, Sweet, l. c. t. 206 = P. venosum.
villosa, Sweet, Hort. Brit. ed. I. 76 = P. villosum.
violaefiora, Sweet, Geran. t. 138 = P. violaefiorum.
virginea, Eckl. & Zeyh. Enum. 61 = P. longifolium.

HOCHENWARTIA, Crantz, *Inst.* ii. 468 (1766) = **Rhododendron**, Linn. (Ericac.).
canadensis, Crantz, l. c. 469 = R. Rhodora.

HOCHSTETTERA, Spach, Hist. Vég. Phan. x. 28 (1841) = seq.

HOCHSTETTERIA, DC. Prod. vii. 287 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 492.
Schimperi, *DC. l. c.*—Arab.

HOCKEA, Lindl. Veg. Kingd. 636 (1847), err. typ. = **Fockea**, Endl. (Asclep.).

HOCKINIA, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 12. *GENTIANEAE*, Benth. & Hook. f. ii. 808.
montana, *Gardn. l. c.*—Bras.

HOCQUARTIA, Dum. Comm. Bot. 30 (1822) = **Aristolochia**, Linn.
macrophylla, Dum. l. c. = A. Sipho.
tomentosa, Dum. l. c. = A. tomentosa.

HODGKINSONIA, F. Muell. Fragm. ii. 182 (1861). *RUBIACEAE*, Benth. & Hook. f. ii. 106.
ovatiflora, *F. Muell. l. c.*—Austral.

HODGSONIA, Hook. f. & Thoms. in Proc. Linn. Soc. ii. (1855) 257. *CUCURBITACEAE*, Benth. & Hook. f. i. 820.
heteroclita, *Hook. f. & Thoms. l. c.*—Reg. Himal.; Burma; Malaya.
macrocarpa, Cogn. in DC. Monog. Phan. iii. 849 = heteroclita.

HODGSONIA, F. Muell. Fragm. ii. 95 (1860) = **Hodgsoniola**, F. Muell. (Liliac.).
junciformis, F. Muell. l. c. 96 = Hodgsoniola junciformis.

HODGSONIOLA, F. Muell. Fragm. ii. 176 (1861). *LILIACEAE*, Benth. & Hook. f. iii. 791
Hodgsonia, F. Muell. Fragm. ii. 95 (1860).
junciformis, *F. Muell. Fragm.* ii. 176.—Austral.

HOEFFNAGELIA, Neck. Elem. iii. 68 (1790) = **Trigonia**, Aubl. (Vochysiac.).

HOELSELIA, Juss. in Dict. Sc. Nat. xxi. 269 (1821) = seq.

HOELZELIA, Neck. Elem. iii. 62 (1790) = **Swartzia**, Schreb. (Legumin.).

HOEFTNERIA, Vatke, in Oestr. Bot. Zeitschr. xxix. (1879) 222. *LEGUMINOSAE*.
africana, *Vatke, l. c.*—Afr. trop.

HOFERIA, Scop. Introd. 194 (1777) = **Ternstroemia**, Linn. f.

HOFFMANNIA, Loefl. It. 194 (1758) = **Durante**, Linn. (Verben.).

HOFFMANNIA, Sw. Prod. Veg. Ind. Occ. 80 (1788). *RUBIACEAE*, Benth. & Hook. f. ii. 76.
Camptolobium, Lem. in Fl. des Serres, Sér. I. iii. (1847) sub t. 260.
Higginsia, Pers. Syn. i. 185 (1805).
Ohigginsia, Ruiz & Pav. Fl. Per. i. 55. t. 85 (1798).
affinis, *Hemsl. Diagn. Pl. Nov.* 31.—Costa Rica.
amicrorum, Spreng. Syst. i. 416 = Bikkia grandiflora.
canliflora, *Hemsl. Diagn. Pl. Nov.* 30.—Guatemala.
discolor, *Hemsl. Biol. Centr. Am. Bot.* ii. 36.—Mexic.
Ghiesbreghtii, *Hemsl. l. c.*—Mexic.
jamaicensis, Spreng. Syst. i. 418 = pedunculata.
lanceolata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 504.—Ind. occ.
lanticollata, *Hemsl. Diagn. Pl. Nov.* 30.—Mexic.
longepetiolata, *Palab. in Linnaea*, xli. (1877) 567.—Am. centr.
macrophylla, *Hemsl. Biol. Centr. Am. Bot.* ii. 37.—Guatemala.
mexicana, *Hemsl. l. c.*—Mexic.
pedunculata, *Sw. Prod. Veg. Ind. Occ.* 80.—Jamaic.
psychotrioides, *Griseb. Fl. Brit. W. Ind.* 321.—Ind. occ.; Mexic.
refulgens, *Hemsl. Biol. Centr. Am. Bot.* ii. 37.—Mexic.
regalis, *Hemsl. l. c.*—Mexic.
strigillosa, *Hemsl. Diagn. Pl. Nov.* 31.—Mexic.
tubiflora, *Griseb. Fl. Brit. W. Ind.* 321.—Ind. occ.

HOFFMANNSEGGIA, Willd. Enum. Hort. Berol. 445 (1809) = seq.

HOFFMANNSIA, Bronn, Diss. Legum. 180 (1822) = seq.

HOFFMANNSEGGIA, Cav. Ic. fv. 63. t. 592 (1797). *LEGUMINOSÆ*, Benth. & Hook. f. i. 567.
MELANOSTICTA, DC. Mém. Lég. 474. t. 69; Prod. ii. 485 (1825).
alpina, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 209 = falcaria.
andina, *Meyen, Trav. Chil.* ii. 582.—Chili.
brachyosrpa, *A. Gray, Pl. Wright.* i. 55.—Mexic.
Burchellii, *Benth. & Hook. f. Gen.* i. 567; *et ex Oliver, Fl. Trop. Afr.* ii. 353.—Afr. trop.
caudata, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 179.—Texas.
chiensis, Miers, ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 209 = falcaria.
demissa, Benth. ex A. Gray, Pl. Wright. i. 56 = stricta.
densiflora, *Benth. ex A. Gray, l. c.* 56.—Texas.
Doelli, *Phil. in Anal. Univ. Chil.* (1872) 710.—Chili.
drepanocarpa, *A. Gray, l. c.* 58.—N. Mexic.
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 393.—Texas.
falcaria, *Cav. Ic.* iv. 63.—Chili; Peruv.
gladiata, *Benth. in A. Gray, Pl. Wright.* i. 57.—Mexic.
glandulosa, *Willd. ex Vog. in Linnæa*, xi. (1837) 404.—Bras.
gracilis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 209.—Chili.
gracilis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 347.—N. Mexic.
humilis, *Hensl. Biol. Centr. Am. Bot.* i. 326.—Mexic.
Jamesii, *Torr. & Gray, Fl. N. Am.* i. 393.—Am. bor. occ.
melanosticta, *A. Gray, Pl. Wright.* i. 54.—Mexic.
microphylla, *Torr. in Bot. Mex. Bound.* 58.—Calif.
oxycarpa, *Benth. ex A. Gray, Pl. Wright.* i. 55.—N. Mexic.
pilosa, *G. Don, Gen. Syst.* ii. 434.—Peruv.
platycarpa, *Benth. ex A. Gray, Pl. Wright.* i. 57.—Mexic.
prostrata, *Lag. ex DC. Prod.* ii. 485.—Peruv.
stricta, *Benth. ex A. Gray, Pl. Wright.* i. 56.—Mexic.
striata, C. Muell. in Walp. Ann. iv. 593, sphalm. = prasc.
trifoliata, *Cav. Ic.* iv. 65.—Patagon.
viscosa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 209.—Peruv.

HOFMANNIA, Spreng. Anleit. ii. 11. 605 (1819) = **Hoffmannia**, Sw. (Rubiac.).

HOFMEISTERA, Reichb. f. Poll. Orchid. 30 (1852) = **Hofmeisterella**, Reichb. f. (Orchid.).
cumicroscopica, Reichb. f. l. c. = Hofmeisterella cumicroscopica.

HOFMEISTERELLA, Reichb. f. in Walp. Ann. iii. 568 (1852). *ORCHIDEÆ*, Benth. & Hook. f. iii. 570.
HOFMEISTERA, Reichb. f. Poll. Orchid. 30 (1852). cumicroscopica, *Reichb. f. in Walp. Ann.* iii. 568.—Peruv.

HOFMEISTHRIA, Walp. Rep. vi. 106 (1847). *COMPOSITÆ*, Benth. & Hook. f. ii. 343.
HELOGYNE, Benth. Bot. Voy. Sulph. 30. t. 14 (1844).
fasciculata, *Walp. Rep.* vi. 106.—Calif.
pluriseta, *A. Gray, in Pacif. Rail. Rep.* iv. 96. t. 6.—Calif.
urenifolia, *Walp. Rep.* vi. 106.—Calif.

HOHENACKERIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 38 (1835). *UMBELLIFERÆ*, Benth. & Hook. f. i. 866.
bupleurifolia, *Fisch. & Mey. l. c.*—Reg. Caucas.
polyodon, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 183.—Algeria.

HOHENBERGIA, Schult. f. Syst. vii. 71, 1951 (1830) = **Aechmea**, Ruiz & Pav. (Bromeliac.).
angustifolia, Baker, in Saund. Refug. Bot. post t. 284 = A. spicata.
angusta, E. Morr. Cat. (1873) 9 = A. angusta.
bilbergioides, Schult. f. Syst. vii. 1953.—Bras.
bracteata, Baker, in Saund. Refug. Bot. post t. 284 = A. bracteata.

HOHENBERGIA:—
bracteata, Beer, Bromel. 73.—Bras.
caniculata, Baker, in Saund. Refug. Bot. t. 284 = Aechmea caniculata.
calyculata, Baker, l. c. t. 286 = A. calyculata.
capitata, Schult. f. Syst. vii. 1952 = A. exsudans.
comata, Baker, in Saund. Refug. Bot. post t. 284.—Hab.?
cyathiformis, Beer, Bromel. 73 (*cyanthiformis*).—Bras.
distans, Baker, in Saund. Refug. Bot. post t. 284 = A. distans.
distichacantha, Baker, l. c. = A. distichantha.
erythrostachys, Brongn. in Rev. Hortic. Sér. IV. iv. (1855) 344 = A. glomerata.
exsudans, E. Morr. in Belg. Hortic. xxix. (1879) 253. t. 18 = A. exsudans.
fasciata, Schult. f. Syst. vii. 1958 = A. fasciata.
ferruginea, Carr. in Rev. Hortic. (1881) 487. f. 104 = A. augusta.
glomerata, Baker, in Saund. Refug. Bot. post t. 284 = A. glomerata.
laxiflora, Baker, l. c. = A. laxiflora.
Legrelliana, Baker, l. c. t. 285 = A. Legrelliana.
Martii, Baker, l. c. post t. 284 = A. spicata.
Melinonii, Baker, l. c. = A. Melinonii.
Mertensii, Baker, l. c. = A. Mertensii.
mucronisflora, Baker, l. c. = A. mucronisflora.
nudicaulis, Baker, l. c. = A. nudicaulis.
odora, Baker, l. c. = A. odora.
paniculigera, Baker, l. c. excl. syn. = A. paniculigera.
platynema, Baker, l. c. = A. platynema.
pyramidalis, Baker, l. c. = A. pyramidalis.
spicata, Baker, l. c. = A. spicata.
stelata, Schult. f. Syst. vii. 1953 = A. glomerata.
strobilacea, Schult. f. l. c. 1952 = Acanthostachys strobilacea.
terminalis, Beer, Bromel. 73.—Bras.

HOHENWARTHA, Vest, in Flora, iii. (1820) 1 = Carthamus, Linn. (Compos.).
gymnospyna, Vest, l. c. = C. lanatus.

HOHERIA, A. Cunn. in Ann. Nat. Hist. Sér. I. iii. (1839) 319. *MALVACEÆ*, Benth. & Hook. f. i. 204.
angustifolia, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 122 = populnea.
Lyallii, Hook. f. Fl. N. Zel. i. 31. t. 11 = Plagianthus Lyallii.
Cunninghamii, H. B. & K. Nov. Gen. et Sp. iii. 164 = L. glandulosa.
coccinea, Cav. Ic. iv. 44 = L. coccinea.
corrolata, Cav. l. c. t. 356 = L. corrulea.
longifolia, H. B. & K. Nov. Gen. et Sp. iii. 165 = L. glandulosa.
ilata, Hook. & Arn. Bot. Beech. Voy. Suppl. 441 = L. glandulosa.
floribunda, Mart. & Gal. in Bull. Acad. Brux. xii. 11. (1845) 275 = L. rupestris.
glandulosa, Cav. Ic. iv. 43 = L. glandulosa.
linearis, Sprong. Syst. i. 636 = Collomia linearis.
Loeselia, Sprung. l. c. = L. ciliata.
lupulina, Hook. & Arn. Bot. Beech. Voy. Suppl. 441 = L. involucrata.
mexicana, Lam. Encyc. iii. 134 = L. coccinea.
nepetæfolia, Cham. & Schlecht. in Linnæa, vi. (1831) 385 = L. glandulosa.
pumila, Mart. & Gal. in Bull. Acad. Brux. xii. 11. (1845) 275 = L. pumila.
ramosissima, Mart. & Gal. l. c. 278 = L. ramosissima.
scabra, Mart. & Gal. l. c. 274 = L. scabra.
scariosa, Mart. & Gal. l. c. = L. scariosa.
splenta, Willd. in Roem. & Schult. Syst. iv. 370 = L. glandulosa.
squarrosa, Eschsch. ex Benth. in DC. Prod. ix. 230 = Gilia squarrosa.

HOLACANTHA, A. Gray, in Mem. Am. Acad. Ser. II. v. (1855) (Pl. Nov. Thurb.) 310. t. 8. *SIMARUBEÆ*, Benth. & Hook. f. i. 310.
Emoryi, *A. Gray, l. c.*—Mexic.

HOLARGES, Ehrh. Beitr. iv. 148 (1789) = Drabe, Linn. (Crucif.).

HOLARGIDIUM, Turcz. in Bull. Soc. Nat. Mosc. (1838) 87; et xv. (1842) 265.—Turcz. Linn. (Crucif.).
Kuznetzowii, Turcz. l. c. 266.—Sibir.

HOLARRHENA, R. Br. in Mem. Wern. Soc. i. (1809) 62. *APOCYNACEÆ*. Benth. & Hook. f. ii. 706.
PHYSETOSTASIS, Hassk. in Versl. en Med. Kon. Akad. Wetensch. v. (1857) 99; et Flora, xl. (1857) 104.
affinis, *Hook. & Arn. Bot. Beech. Voy.* 196.—China.
africana, *A. DC. in DC. Prod.* viii. 414.—Afr. trop.
antidysenterica, *Wall. Cat.* n. 1679.—As. trop.
Codaga, G. Don, Gen. Syst. iv. 78 = antidysenterica.
febrifuga, *Klotzsch, in Peters, Reise Mossamb. Bot.* 277.—Afr. trop.
glabra, *Klotzsch, l. c.* 279.—Afr. trop.
landolphioides, A. DC. in DC. Prod. viii. 414 = Alafia landolphioides (in Addend.).
macrocarpa, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 130.—Java?; Ins. Philipp.
madagascariensis, *Baker, in Journ. Linn. Soc.* xxi. (1883) 434.—Madag.
malaccensis, Wight, Ic. t. 1298 = antidysenterica.
mitis, *R. Br. in Mem. Wern. Soc.* i. (1809) 62.—Zeylan.
ovata, *A. DC. in DC. Prod.* viii. 414.—Afr. trop.
pubescens, Wall. Cat. n. 1678 = antidysenterica.
tettensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 278.—Afr. trop.
villosa, Alt. ex Loud. Hort. Brit. Suppl. ii. 641 = antidysenterica?

HOLBOELLIA, Wall. in Hook. Bot. Misc. ii. (1831) 144. t. 76 = Lophoiepis, Decne. (Gramin.).
ornithocephala, Hook. l. c.—Ind. or.

HOLBOELLIA, Wall. Tent. Fl. Nep. 23. t. 16, 17 (1824). *BERBERIDEÆ*, Benth. & Hook. f. i. 42.
acuminata, Lindl. in Journ. Hort. Soc. i. (1846) 313 = latifolia.
angustifolia, Wall. Tent. Fl. Nep. 25. t. 17 = latifolia.
latifolia, *Wall. l. c.* 24. t. 16.—Reg. Himal.

HOLCUS, Linn. Syst. ed. I (1735). *GRAMINEÆ*, Benth. & Hook. f. iii. 1159.
aciocularis, *Resp. in Ann. Obs.* iv. t. 10; *ex Prits. Ic. Ind.* 1. 555.—Hab.?
aciculatus, Griff. Ic. Fl. Asiat. t. 141 = Chrysopogon aciculatus.
acutiflorus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 198 = lanatus.
albus, Steud. Nom. ed. I. 413 = Sorghum vulgare.
alpinus, Sw. ex Willd. Sp. Pl. iv. 937 = Hierochloë alpina.
annuus, Salisb. ex Trin. in Mém. Acad. Pétersb. Sér. V. vi. ii. (1840) 67 = setosus.
arcticus, Sommerf. in Mag. Naturvid. Ser. II. i. (1838) 286 = Hierochloë borealis?
Arduini, J. F. Gmel. Syst. 174 = Sorghum vulgare.
arenarius, All. Auct. Fl. Pedem. 46 = Koeleria villosa.
argenteus, Agardh, ex Roem. & Schult. Syst. ii. 656 = lanatus.
asper, Thunb. Prod. Fl. Cap. 20 = Danthonia papillosa.
atropurpureus, Wahlenb. in Svensk Bot. t. 147 = Deschampsia atropurpurea.
australis, Schrad. Fl. Germ. i. 253 = Hierochloë australis.
avenaceus, Scop. Fl. Cass. ed. II. ii. 276 = Arrhenatherum avenaceum.
avenaceus, Thunb. Prod. Fl. Cap. 20 = Danthonia cervicula.
bicristatus, [Web. in] Wigg. Prim. Fl. Hols. 73 = Gaudinia tenuis.
bicolor, Lisa, Mant. ii. 201 = Sorghum vulgare.
borealis, Schrad. Fl. Germ. i. 252 = Hierochloë borealis.
bulbosus, Schrad. l. c. 245 = Arrhenatherum avenaceum.

HOLCUS:—
caerulescens, Gandich. in Freyc. Voy. Bot. 411. t. 97 = Chrysopogon violascens.
cespitosus, *Boiss. Voy. Bot. Espagne*, 686. t. 173; *Elench.* 86.—Hispan.
cafer, Ardnini, in Saggi Padov. i. (1786) 119 = Sorghum vulgare.
Caffrorum, Thunb. Prod. Pl. Cap. 20 = Sorghum vulgare.
capillaris, Thunb. l. c. = Andropogon capillaris.
cernuus, Arduini, in Saggi Padov. i. (1786) 128; Willd. Sp. Pl. iv. 931 = Sorghum vulgare.
ciliatus, *Roxb. Hort. Beng.* 83; *Fl. Ind.* i. 318.—Ind. or *caerulescens*, Auct. vide *caerulescens*.
compactus, Lam. Encyc. iii. 140 = Sorghum vulgare.
cubanicus, Gleditsch, ex Schult. Mant. ii. 462 (Quid?). —Cuba.
cylindricus, *Brouss. Elench. Pl. Hort. Monsp.* 1804 (1805), 80, nomen; *ex Roem. & Schult. Syst.* ii. 857 (Quid?).—Hab.?
decolorans, Willd. Sp. Pl. iv. 931 = Sorghum halepense.
dersus, Peterm. in Flora, xxvii. (1844) 233 = mollis.
distachyos, Cat. Hort. Dorpat. (1814); ex Roem. & Schult. Syst. ii. 857 (Quid?).—Hab.?
Dochna, Forsk. Fl. Aegypt. Arab. 174 = Sorghum vulgare.
Dora, Mieg, in Act. Helv. viii. (1777) 125. t. 4. f. 3 = Sorghum vulgare.
Durra, J. F. Gmel. Syst. 178, sphalm. = *H. Durra*.
Durrissi, Steud. Syn. Pl. Gram. 15 = Gayanus.
Durra, Forsk. Fl. Aegypt. Arab. 174 = Sorghum vulgare.
elongatus, R. Br. Prod. 200 = Chrysopogon elongatus.
exiguus, Forsk. Fl. Aegypt. Arab. 174 = Sorghum halepense.
ferrugineus, Schrad. ex Cat. Hort. Dorpat. (1814); ex Roem. & Schult. Syst. ii. 857 = Sorghum vulgare.
fragrans, Willd. Sp. Pl. iv. 936 = Hierochloë fragrans.
fulvus, R. Br. Prod. 199 = Andropogon serratus.
Gayanus, *Boiss. Voy. Bot. Espagne*, 687. t. 178 C.— Hispan.
glaucus, Willd. in Willd. & Lange, Prod. Fl. Hisp i. 307 = lanatus.
grandiflorus, *Boiss. & Reut. Pugill. Pl. Nov.* 119.— Hispan.
Gryllus, R. Br. Prod. 199 = Chrysopogon Gryllus.
halepensis, Linn. Sp. Pl. 1047 = Sorghum halepense.
intermedius, Krock. Fl. Siles. Suppl. i. 137 = lanatus?
lanatus, *Linn. Sp. Pl.* 1048.—Europ.
latifolius, Linn. Syst. ed. X. 1305 = Centotheca lappacea.
laxus, Bosc, ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. 11. (1840) 57 = Sporobolus centrifugus.
laxus, Linn. Sp. Pl. 1048 = Uniola gracilis.
liburnicus, Scop. Fl. Carn. ed. II. ii. 276 = Andropogon distachyos.
mollis, *Linn. Syst.* ed. X. 1305.—Europ.
monticola, *Bigel. in New Engl. Med. Journ.* v. (1816) 334.—Am. bor.
muricus, Kunze, ex Willk. Pugill. Pl. Nov. 142 = Poa balbosa.
nervosus, Roxb. Hort. Beng. 82; Fl. Ind. i. 318 = Panicum sensitivum.
niger, Arduini, in Saggi Padov. i. (1786) 134; J. F. Gmel. Syst. 174 = Sorghum vulgare.
nigerrimus, Arduini, l. c. t. 5. f. 1; ex Roem. & Schult. Syst. ii. 857 = Sorghum vulgare.
nigricans, Hort. ex Steud. Nom. ed. II. i. 772 = Sorghum vulgare.
nitidus, Vahl, Symb. Bot. ii. 102 = Andropogon serratus.
Notarisii, *Nym. Syll.* 441; *Cat. Comp.* 65.—Italia.
odoratus, Linn. Sp. Pl. 1048 = Hierochloë australis, borealis.
odoratus, Michx. Fl. Bor. Am. i. 56 = Hierochloë fragrans.
pallidus, R. Br. Prod. 199 = Chrysopogon Gryllus.
parviflorus, R. Br. l. c. = Chrysopogon violascens.
pertusus, Linn. Mant. ii. 301 = Andropogon pertusus.
plumosus, R. Br. Prod. 200 = Andropogon australis.
pyramidalis, Hort. ex Steud. Nom. ed. II. i. 772 = Sorghum vulgare.
racemosus, *Forsk. Fl. Aegypt. Arab.* 176.—Arabia.
redolens, Forst. f. Prod. 91; Vahl, Symb. Bot. ii. 102 = Hierochloë redolens.
repens, Host, Gram. ii. 3. t. 3 = Hierochloë borealis.
Reuteri, *Boiss. Pugill. Pl. Nov.* 119.—Hispan.

HOLCUS:—
rigidus, *Hochst. ex Snb. Fl. Asor.* 17.—Ins. Azoric.
rubens, Guerin, Fruct. ii. 3 = Sorghum vulgare.
saccharatus, Griseb. Fl. Brit. W. Ind. 560, in obs. = Oplismenus undulatifolius.
saccharatus, Linn. Sp. Pl. 1047 = Sorghum vulgare.
Sorni, Steud. Synopsis. Pugill. i. 8 = Trisetum paniceum.
serratus, Thunb. Prod. Pl. Cap. 20 = Panicum serratum.
setifolius, Thunb. in Hoffmgg. Phytogr. Blaett. i. (1803) 6 = Andropogon setifolius.
setiger, De Not. ex Parl. Fl. Ital. i. 65 = Notarisii.
setiger, *Nees, in Linnaea*, vii. (1832) 276.—Afr. austr.
setiglumis, Boiss. & Kent. Diagn. Pl. Nov. Hisp. 27 = setosus.
setosus, *Trin. in Mém. Acad. Pétersb.* Sér. V. vi. 11. (1840) 87.—Hispan.; Afr. bor.
Sorghum, Linn. Sp. Pl. 1047 = Sorghum vulgare.
spicatus, Linn. Syst. ed. X. 1305 = Pennisetum typhoideum.
striatus, *Linn. Sp. Pl.* 1048.—Am. bor.
tenuis, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 136 = Gayanus.
tenuis, *Trin. in Spreng. Neue Entdeck.* ii. 75.—Sicilia.
tuberosus, Salzm. ex Trin. in Mém. Acad. Pétersb. Sér. V. vi. ii. (1840) 86 = lanatus.
vestitissimus, Flach. ex Trin. l. c. 88 = setosus.
virginicus, Steud. Nom. ed. II. i. 778 = Andropogon virginicus.

HOLDERLINIA, Neck. Elem. i. 106 (1790) = **Serruria**, Salisb. (Proteac.).

HOLIGARNA, Buch.-Ham. ex Roxb. Hort. Beng. 22 (1814); et Pl. Corom. iii. 79. t. 282 (1819). *ANACARDIACEAE*, Benth. & Hook. f. i. 423.
HADESTAPHYLUM, Dennst. Schluss. Hort. Malab. 30 (1818).
albicans, *Hook. f. Fl. Brit. Ind.* ii. 38.—Burma.
Arnottiana, *Hook. f. l. c.* 86.—Ind. or.
Beddomei, *Hook f. l. c.*—Ind. or.
ferruginea, *March. Rev. Anacard.* 171.—Ind. or.
Grahamii, *Hook. f. Fl. Brit. Ind.* ii. 37.—Ind. or.
Grahamii, Kurz, in journ. As. Soc. Beng. xli. (1872) ii. 305 = albicans.
Helferi, *Hook. f. Fl. Brit. Ind.* ii. 37.—Burma.
longifolia, *Buch.-Ham. ex Roxb. Hort. Beng.* 22; *Pl. Corom.* iii. 79. t. 82.—Ind. or.
longifolia, Span. in Linnaea, xv. (1841) 188 = Semecarpus longifolia.
longifolia, Wight & Arn. Prod. 169 = Arnottiana.
racemosa, Hohen. ex Engl. in DC. Monog. Phan. iv. 466 = Arnottiana.
racemosa, Roxb. Hort. Beng. 22; Fl. Ind. ii. 82 = Drimycarpus racemosus.

HOLLBOELLIA, Meisn. Gen. 418 [320] (1843) = HOLBOELLIA, Wall. = **Lopholepis**, Decne. (Gomphand.).

HOLLBOELLIA, Spreng. Syst. iv. Cur. Post. 343 (1827) = **Holboellia**, Wall. (Berberid.).

HOLLIA, Heynh. Nom. i. 850 (1840); et ii. 303 (1846) = **Noltea**, Reichb. (Rhamn.).
africana, Heynh. ii. cc. = N. africana.

HOLLIA, Heynh. Nom ii. 303 (1846) = **Chlorophytum**, Ker-Gawl. (Liliac.).
comosa, Heynh. l. c. = C. Sternbergianum.

HOLLISTERIA, S. Wats. in Proc. Am. Acad. xiv. (1879) 226. *POLYGONACEAE*, Benth. & Hook. f. iii. 94.
lanata, *S. Wats. l. c.*—Am. bor. occ.

HOLMSKIDIA, Dum. Anal. Fam. 22 (1829), sphalm. = seq.

HOLMSKIOLDIA, Retz. Obs. vi. 31 (1791). *VERBENACEAE*, Benth. & Hook. f. ii. 1156.
HARTINGIA, Koen. ex Sm. Exot. Bot. ii. 41. t. 80 (1806).
PLATUNIUM, Juss. in Ann. Mus. Par. vii. (1806) 76.
rubra, Pers. Syn. ii. 144 = sanguinea.
sanguinea, *Retz. Obs.* vi. 31.—Ind. or.
scandens, Sweet, Hort. Brit. ed. I. 253 = sanguinea.

HOLOCALYX, Michell, in Mém. Soc. Phys. Genèv. xxvii. (1883) n. vii. 41. *LEGUMINOSAE*.
Balansae, *Michell, l. c.*—Paraguay.

HOLOCARPA, Baker, in Journ. Linn. Soc. xxi. (1885) 414. *RUBIACEAE*.
veronicoides, *Baker, l. c.*— Madag.

HOLOCHEILUS, Cass. in Bull. Soc. Philom. (1818) 73 = **Trixis**, P. Br. (Compos.).
ochroleucus, Cass. l. c. = T. ochroleuca.

HOLOCHILOMA, Hochst. in Flora, xxiv. (1841) 871 = **Premna**, Linn. (Verbenac.).
reticosum, Hochst. l. c.—Afr. trop.

HOLOCHILUS, Dalz. in Hook. Kew Journ. iv. (1852) 290 = **Meba**, Forst. (Ebenac.).
micranthus, Dalz. l. c. 291 = M. micrantha.

HOLOCHLAMYS, Engl. in Becc. Malesia, i. (1883) 265. *AROIDEAE*.
Beccarii, *Engl. l. c.*—Malaya.

HOLOCHLOA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 580 (1840) = **Heuchera**, Linn. (Saxifr.).
cylindrica, Nutt. ex Torr. & Gray, l. c. = Heuch. cylindrica.
elata, Nutt. ex Torr. & Gray, l. c. = Heuch. cylindrica.
glabra, Nutt. ex Torr. & Gray, l. c. 581 = Heuch. cylindrica.
ovalifolia, Nutt. ex Torr & Gray, l. c. = Heuch. cylindrica.

HOLODISCUS, Maxim. in Act. Hort. Petrop. vi. (1879) 253 = **Spiraea**, Linn. (Rosac.).
argentea, Maxim. l. c. 254 = S. argentea.
discolor, Maxim. l. c. = S. discolor.

HOLOGAMIUM, Nees, in Edinb. N. Phil. Journ. xviii. (Jan. 1835) 185 = **Isohaemum**, Linn. (Gramin.).
nervosum, Nees, l. c. = I. laxum.

HOLOGRAPHIS, Nees, in DC. Prod. xi. 728 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1104.
Ehrenbergiana, *Nees, l. c.*—Mexic.

HOLOGYMNE, Bartl. Ind. Sem. Hort. Gotting.; ex Linnaea, xi. (1838) Litt. 81 = **Lasthenia**, Cass (Compos.).
californica, Bartl.—Cf. Linnaea, xii. (1838) Litt. 82 = L. glabrata.
Douglasii, C. A. Mey. Ind. Sem. Hort. Petrop. viii. 64 = Monolepis major.
glabrata, Bartl.—Cf. Linnaea, xii. (1838) Litt. 81; et xiv. (1840) Litt. 125 = L. glabrata.

HOLOLACHNA, Ehrenb. in Linnaea, ii. (1827) 278. *TAMARISCINEAE*, Benth. & Hook. f. i. 161.
Shawiana, *Hook. f. ex Hemd. & Hume, Yarkand*, 313.—As. temp.
songarica, *Ehrenb. in Linnaea*, ii. (1827) 278.—Sibiria.

HOLOLACHNE, Reichb. Consp. 186 (1828) = prec.

HOLOLEPIS, DC. in Ann. Mus. Par. xvi. (1810) 155, 189. t. 6 = **Vernonia**, Schreb. (Compos.).
involucrata, Sch. Bip. in Pollichia, xx.-xxi. (1863) 880 = V. pedunculosa.
pedunculata, DC. in Ann. Mus. Par. xvi. (1810) 190 = V. pedunculata.

HOLOPEIRA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42 = **Cocculus**, DC. (Menisp.).
auriculata, Miers, l. c. Ser. III. xix. (1867) 29 = C. auriculatus.
australis, Miers, l. c. = C. laurifolius.
fecunda, Miers, l. c. = C. hexagynus?
fusiformis, Miers, l. c. = C. laurifolius.
laevinervis, Miers, l. c. 28 = C. villosus.
laurifolia, Miers, l. c. 29 = C. laurifolius.
lonchophylla, Miers, l. c. 29 = C. Ferrandianus?
torrida, Miers, l. c. = C. villosus.
villosa, Miers, l. c. Ser. II. vii. (1851) 42 = C. villosus.

HOLOPETALA, Wight, Ic. vi. t. 1968 (1853), sphalm. = **Holopetalon**, Planch. (Urticac.).

HOLOPETALON, Reichb. Nom. 907 (1841) = **Banisteria**, Linn. (Malpig.).

HOLOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 51 = **Oligomeris**, Cambess. (Resed.).
Burchelii, Mneil. Arg. in Bot. Zeit. xiv. (1856) 39; et Monog. Resed. 215. t. 9. f. 187 = O. Burchelii.
pumilum, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 51 = O. capensis.
spathulatum, Turcz. l. c. xxvii. (1854) 18. 880 = O. spathulatum.

HOLOPHYLLUM, Less. Syn. Comp. 262 (1832) = **Athanasia**, Linn. (Compos.).
capitatum, Less. l. c. = A. capitata.
capitatum, Sch. Bip. in Flora, xxvii. (1844) 676 = A. dimorpha.
lanuginosum, DC. Prod. vi. 86 = A. canescens.
scabrum, DC. l. c. = A. capitata.

HOLOPLEURA, Regel & Schmalh. ex Regel, Pl. Nov. Fedsch. 28 (1889). *UMBELLIFERÆ*.
carioides, *Regel & Schmalh. l. c.*—Turkestan.

HOLOPTELEA, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 259. *URTICACEÆ*; Benth. & Hook. f. iii. 362.
integrifolia, *Planch. l. c.*—Ind. or.

HOLOPTOLAEA, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 266 = prec.

HOLOREGMIA, Nees, in Flora, iv. (1821) 800 = **Martynia**, Linn. (Pedalin.).
viscida, Nees, l. c. = M. spathacea.

HOLOSCHOENUS, Link, Hort. Berol. Desc. i. 293 (1827) = **Scirpus**, Linn. (Cyperac.).
alto-nitatus, Reichb. ex Nym. Consp. 765 = S. Holoschoenus.
atrovirens, Link, Hort. Berol. Desc. i. 293 = S. atrovirens.
australis, Reichb. in Flora, xiii. (1830) 499 = S. Holoschoenus.
brunneus, Steud. Nom. ed. II. i. 278 = S. brunneus.
elatior, Nees, in Hook. Journ. Bot. ii. (1840) 304 (Ryschoporæ sp. ?).—Am. austr.
exsertus, Reichb. in Flora, xiii. (1830) 501 = S. Holoschoenus.
fluviatilis, Reichb. l. c. 501 = S. Holoschoenus.
globiferus, A. Dietr. Sp. Pl. ii. 164 = S. Holoschoenus.
incurvatus, A. Dietr. l. c. 165 = S. incurvatus.
lævus, Opiz, Seznam, 38 = S. Holoschoenus.
Linnæi, Reichb. Fl. Germ. Excurs. 76 = S. Holoschoenus.
melanocarpus, Boeck. in Flora, xl. (1857) 87 (Quid?).—Mexic.
nodosus, A. Dietr. Sp. Pl. ii. 165 = S. nodosus.
panormitanus, Nym. Consp. 765 = S. Holoschoenus.
subarticulatus, A. Dietr. Sp. Pl. ii. 166 = S. articulatus.
Thunbergii, A. Dietr. l. c. 164 = S. Holoschoenus.
tuberculatus, Reichb. in Flora, xiii. (1830) 498 = S. Holoschoenus.
vulgaris, Link, Hort. Berol. Desc. i. 293 = S. Holoschoenus.

HOLOSEPALUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 352 = **Hypericum**, Tourn.
dubium, Fourr. l. c. = Hyp. dubium.
humifusum, Fourr. l. c. = Hyp. humifusum.
quadrangulum, Fourr. l. c. = Hyp. quadrangulum.

HOLOSETUM, Steud. in Flora, xxxiii. (1850), 228, nomen; et Syn. Pl. Gram. 118 (1855) = **Panicum**, Linn. (Gramin.).
philippicum, Steud. ll. cc. = P. semialatum.

HOLOSTEMMA, R. Br. in Mem. Wern. Soc. i. (1809) 43. *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 760.
Ada-Kodien, Schult. Syst. vi. 95 = Rheedianum.
Brunoniense, Royle, Illustr. Bot. Himal. 276. t. 66 = Rheedianum.
Candollеanum, Spreng. Syst. i. 851 = Fischeria scandens.

HOLOSTEMMA:—
chilense, *Phil. in Linnæa*, xxxiii. (1864–65) 174.—Chili.
fragrans, Wall. Pl. As. Rar. ii. 51 = Rheedianum.
laeve, Blume, Bijdr. 1055 = Cynanchum ovalifolium.
muricatum, Blume, Bijdr. 1054 (= *Cynactonum muricatum*).—Java.
pictum, Benth. in Hook. Kew Journ. v. (1853) 53 = Graphistemma pictum.
Rheedei, Wall. Pl. As. Rar. ii. 51; et Cat. n. 4409 = sec.
Rheedianum, *Spreng. Syst. i.* 851.—Ind. or.
tuberculatum, Blume, Bijdr. 1055 (*Cynodonum tuber culatum*, Blume, in Addend.).—Java.

HOLOSTEUM, Dill. ex Linn. Gen. ed. I. 376 (1737). *CARYOPHYLLEÆ*, Benth. & Hook. f. i. 148.
Meyera, Adans. Fam. ii. 257 (1763).
Alsine, Sw. Obs. 118 = Stellaria media.
ciliatum, Opiz, Natural. ix. (1825) 128 = umbellatum ?
cordatum, Linn. Sp. Pl. 88 = Drymaria cordata.
diandrum, Sw. Prod. Veg. Ind. Occ. 27 = Drymaria cordata.
dichotomum, C. Koch, in Linnæa, xv. (1841) 708 = Cerastium anomalum.
filiforme, Risso, ex Willk. & Lange, Prod. Fl. Hisp. iii. 829 = Cerastium quaternellum.
glandulosum, Bertol. Misc. Bot. i. 11 = linifolium.
glutinosum, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 52 = umbellatum.
Heufleli, Wierzbicki, in Flora, xxv. (1842) 264 = umbellatum.
hirsutum, *Linn. Sp. Pl.* 88 (Quid ?).—Ind. or.
imberbe, Gay, Monog. Holost. 37 = linifolium.
linifolium, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 39 = umbellatum.
linifloron, *Stev. ex Fisch. & Mey. l. c.* iv. 10.—As. Min. ; Persia.
marginatum, *C. A. Mey. ex Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 402.—As. Min. ; Persia.
montanum, Wight, ex Wall. Cat. n. 647 = Drymaria cordata.
mucronatum, Moç. & Sesse, ex DC. Prod. i. 393 = Drymaria pauciflora.
polygynum, C. Koch, in Linnæa, xv. (1841) 708 = linifolium.
Sibthorpii, Boiss. Fl. Orient. i. 710 = tenerrimum.
sperguloides, *Lehm. Fugill. Pl. Hort. Hamb.* i. 10.—Aegypt.
succulentum, Linn. Sp. Pl. 88 = Arenaria peploides.
tenerrimum, *Boiss. Diagn.* Ser. I. i. 53.—As. Min.
tetraphyllum, Thunb. Prod. Fl. Cap. 24 = Polycarpon tetraphyllum.
umbellatum, *Linn. Sp. Pl.* 88.—Europ.

HOLOSTIGMA, G. Don, Gen. Syst. iii. 716 (1834) = **Lobelia**, Linn. (Campan.).
dioicum, G. Don, l. c. = L. dioica.

HOLOSTIGMA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 365 = **Oenothera**, Linn. (Onograr.).
alyssoides, Hook. lc. Pl. t. 839 = Oe alyssoides.
argutum, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 168 = Oe. dentata.
Bottæ, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 366 = Oe. Bottæ.
cheiranthifolium, Spach, l. c. = Oe. cheiranthifolia.
heterophyllum, Spach, l. c. 364 = Oe. heterophylla, *Steud.*
micranthum, Spach, l. c. 365 = Oe. micrantha, *Presl.*
minutiflorum, Steud. Nom. ed. II. i. 774 = Oe. minutiflora.
paradoxum, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 354 = Oe. micrantha, *Presl.*
quercifolium, Spach, l. c. 352 = Oe. cheiranthifolia.
tenuifolium, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 168 = Oe. stenophylla.
virgatum, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 353 = Oe. dentata.

HOLOSTIGMATEIA, Reichb. Nom. 102 (1841) = prec.

HOLOSTYLA, Endl. Gen. 846 (1838) = **Clostyla**, DC. (Rubiac.).

HOLOSTYLIS, Duch. in Ann. Sc. Nat. Sér. IV. ii. (1854) 88. t. 5. *ARISTOLOCHIACEÆ*, Benth. & Hook. f. iii. 123.
reniformis, *Duch. l. c.*—Bras.

HOLOSTYLIS, Reichb. Nom. 77 (1841) = **Clostyla**, DC. (Rubiac.).

HOLOTHRIX, Rich. in Mém. Mus. Par. iv. (1818) 55. *ORCHIDEÆ*, Benth. & Hook. f. iii. 632.
Bucculina, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 209.
Monotris, Lindl. Bot. Reg. sub t. 1701 (1834).
Saccidium, Lindl. Gen. et Sp. Orch. 401 (1835).
Scopularia, Lindl. Bot. Reg. sub t. 1701 (1834).
Tryphia, Lindl. l. c. (1834).
aphylla, *Reichb. f. Otia Bot. Hamb.* 119.—Afr. austr.
arachnoides, *Reichb. f. l. c.* 107.—Afr. austr.
aspera, *Reichb. f. l. c.* 118.—Afr. austr.
brachylabris, *Sond. in Linnæa*, xix. (1847) 78.—Afr. austr.
Bronguiartiana, *Reichb. f. Otia Bot. Hamb.* 107.—Afr. austr.
Burchellii, *Reichb. f. l. c.* 119.—Afr. austr.
condensata, *Sond. in Linnæa*, xix. (1847) 76.—Afr. austr.
exilis, *Lindl. Gen. et Sp. Orch.* 288.—Afr. austr.
gracilis, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 207.—Afr. austr.
grandiflora, *Reichb. f. Otia Bot. Hamb.* 119.—Afr. austr.
Harveiana, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 208.—Afr. austr.
incurva, *Lindl. l. c.* 207.—Afr. austr.
Lindleyana, *Reichb. f. Otia Bot. Hamb.* 119.—Afr. austr.
Mac-Owaniana, *Reichb. f. l. c.* 108.—Afr. austr.
Monotris, *Reichb. f. l. c.* 119.—Afr. austr.
Mundii, *Sond. in Linnæa*, xix. (1847) 77.—Afr. austr.
orthoceras, *Reichb. f. Otia Bot. Hamb.* 119.—Afr. austr.
parviflora, *Reichb. f. l. c.*—Afr. austr.
parvifolia, *Lindl. Gen. et Sp. Orch.* 288.—Afr. austr.
pillosa, *Reichb. f. Otia Bot. Hamb.* 118.—Afr. austr.
præcox, *Reichb. f. l. c.* 108.—Afr. austr.
Schimperi, *Reichb. f. l. c.*—Abyssin.
Scopularia, *Reichb. f. l. c.* 119.—Afr. austr.
secunda, *Reichb. f. l. c.*—Afr. austr.
squamata, *Reichb. f. l. c.*—Afr. austr.
squamulosa, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 206.—Afr. austr.
tridentata, *Reichb. f. Otia Bot. Hamb.* 119.—Afr. austr.
uniflola, *Reichb. f. l. c.*—Afr. austr.
Vatkeana, *Reichb. f. in Journ. Bot.* xiv. (1876) 346.—Afr. trop.
villosa, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 207.—Afr. austr.

HOLOTOME, Endl. Gen. 766 (1839) = **Actinotus**, Labill. (Umbellif.).
leucocephala, Walp. Rep. ii. 389 = A. leucocephalus.
omnifertilis, F. Muell. Fragm. ii. 239 = A. omnifertilis.

HOLOZONIA, Greene, in Bull. Torrey Bot. Club, ix. (1882) 122 = **Lagophylla**, Nutt. (Compos.).
filipes, Greene, l. c. = L. filipes.

HOLTZENDORFFIA, Klotzsch & Karst. ex Nees, in DC. Prod. xi. 727 (1847) = **Exallia**, Linn. (Acanth.).
caracasana, Klotzsch & Karst. ex Nees, l. c. 728.—Venezuela.

HOLUBIA, Oliver, in Hook. Ic. Pl. t. 1475 (1884). *PEDALINEÆ*.
senensis, *Oliver, l. c.*—Afr. austr.

HOMAID, Adans. Fam. ii. 470 (1763) = **Biarum**, Schott (Aroid.).

HOMAIDA, Rafin. Fl. Tellur. iii. 66 (1836) = **Arisarum**, Tourn. (Aroid.).
proboscidea, Rafin. l. c. = A. proboscideum.

HOMALADENIA, Miers, Apocyn. S. Am. 163 (1878) = **Dipladenia**, A. DC. (Apocyn.).
brevifolia, Miers, l. c. 168 = D. polymorpha.
lineariifolia, Miers, l. c. 164 = D. polymorpha.
pastorum, Miers, l. c. 164 = D. polymorpha.
pedunculata, Miers, l. c. 165 = D. polymorpha.
puberula, Miers, l. c. = D. tenuifolia.
tenuifolia, Miers, l. c. 164 = D. tenuifolia.
vincæflora, Miers, l. c. 165 = D. vincæflora.

HOMALANTHUS, A. Juss. Tent. Euph. 50. t. 16 (1824) (Omalanthus). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 881.
CARUMBIUM, Reinw. in Syll. Ratisb. ii. (1828) 6.
DIERACHION, Regel, Gartenfl. xv. (1866) 100. t. 504.
IUANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 9; ed. II. 85 (1827).
GARUMBIUM, Flora, viii. (1825) 1. 103, err. typ.
WARTMANNIA, Muell. Arg. in Linnæa, xxxiv. (1865–66) 218.
fastuosus, *Villar. in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 196.—Ins. Philipp.
giganteus, *Zoll. & Mor. in Nat. en Geneesk. Arch. Ned. Indie*, ii. (1846) 384.—Java.
Leschenaultianus, *A. Juss. Tent. Euph.* 50. t. 16.—Ind. or.; Malaya.
nutans, *Guill. Esph. Tahit.* 35.—N. Caled.
pedicellatus, *Benth. in Hook. Lond. Journ Bot.* ii. (1843) 382.—Ins. Amicor.
populifolius, R. Grah. in Edinb. N. Phil. Journ. (Apr.–June 1827) 175.= Leschenaultianus.
stillingiifolius, *F. Muell. Fragm.* i. 82.—Austral.

HOMALIUM, Jacq. Enum. Pl. Carib. 5 (1760) *SAMYDACEÆ*, Benth. & Hook. f. i. 800.
ACOMA, Adans. Fam. ii. 510 (1763).
ASTRANTHUS, Lour. Fl. Cochinch. 221 (1790).
BLAKWELLIA, Comm. ex Juss. Gen. 343 (1789).
CORDYLANTHUS, Blume, Mus. Bot. Lugd. Bat. ii. 27. t. 3 (1852).
LAGUNGIA, Scop. Introd. 216 (1777).
LINSCHOTTIA, Comm. ex Juss. Gen. 343, in obs. (1789).
MARSHALLIA, J. F. Gmel. Syst. 826 (1791).
MYRIANTHEIA, Thou. Nov. Gen. Madag. 21 (1806).
NAPIMOGA, Aubl. Pl. Gui. i. 592 (1775).
NISA, Noronha, ex Thou. Nov. Gen. Madag. 24 (1806).
ODODECA, Rafin. Neogenyt. 3 (1825).
PYTHAGOREA, Lour. Fl. Cochinch. 243 (1775).
RACOUREA, Aubl. Pl. Gui. i. 592. t. 236 (1775).
TATTIA, Scop. Introd. 104 (1777).
VERMONETA, Comm. ex Juss. Gen. 343, in obs. (1789).
Abdessammadii, *Aschers. & Schweinf. in Sitzb. Ges. Naturf. Fr. Berl.* (1880) 120.—Afr. trop.
africanum, *Benth. in Journ. Linn. Soc.* iv. (1860) 35. —Afr. occ.
albiflorum, *O. Hoffm. Sert. Madag.* 18.—Madag.
alnifolium, F. Muell. Fragm. ii. 127 = vitiense.
angustifolium, *Sw. in Roer, Cycl.* xvii. n. 8.—Afr. trop.
austro-caledonicum, *Seem. Fl. Vit.* 95.—N. Calcd.
axillare, *Benth. in Journ. Linn. Soc.* iv. (1860) 34.— Madag.
Barandas, *Vidal, ex Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 94.—Ins. Philipp.
brachybotrys, *F. Muell. Fragm.* ii. 127.—Austral.
bracteatum, *Benth. in Journ. Linn. Soc.* iv. (1860) 37. —Ins. Philipp.
caryophyllaceum, Benth. *l. c.* 38.—Java.
ceylanicum, Bedd. Fl. Sylv. t. 210 = zeylanicum.
confertum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 341. —Madag.
Cordylanthus, *Benth. in Journ. Linn. Soc.* iv. (1860) 38.—Java.
cuneifolium, *Willd. ex Eichl. in Mart. Fl. Bras.* xiii. i. 508.—Bras.
dasanthum, *Spruce, ex Benth. in Journ. Linn. Soc.* iv. (1860) 36.—Bras.
fagifolium, *Benth. l. c.* 35.—China.
foetidum, *Benth. l. c.* 37.—Malaya.
foetidum, Benth. l. c. partim = Griffithianum.
foetidum, *Kurz, For. Fl. Brit. Burma*, i. 532.—Burma.
grandiflorum, *Benth. in Journ. Linn. Soc.* iv. (1860) 37.—Malaya.
grandiflorum, Naves, in Blanco, Fl. Philipp. ed. III. t. 448 = penangianum.
Griffithianum, *Kurz, in Journ. As. Soc. Beng.* xl. (1872) ii. 57.—Burma.
incanum, Pers. Syn. ii. 82 = Banara incana.
involucratum, *O. Hoffm. Sert. Madag.* 18.—Madag.
longifolium, *Benth. in Journ. Linn. Soc.* iv. (1860) 35. —Malaya.
longifolium, Benth. l. c. partim = propinquum.
longistylum, *Mast. in Oliver, Fl. Trop. Afr.* ii. 497.— Afr. trop.
luxuriosum, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 94.—Ins. Philipp.

HOMALIUM:—
micranthum, *O. Hoffm. Sert. Madag.* 18.—Madag.
microphyllum, *O. Hoffm. l. c.*—Madag.
minutiflorum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 208.—Burma.
Napimoga, *Spreng. Syst. iv. Cur. Post.* 210.—Guian.
nepalense, *Benth. in Journ. Linn. Soc.* iv. (1860) 34. —Nep. Himal.
obovale, *Miq. Fl. Ind. Bat. Suppl.* 384.—Sumatra.
obtusatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 466.—Cuba.
panayanum, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 94.—Ins. Philipp.
paniculatum, *Benth. in Journ. Linn. Soc.* iv. (1860) 34.—Ins. Maurit.
paniculatum, *O. Hoffm. Sert. Madag.* 18.—Madag.
Parkeri, *Baker, in Journ. Linn. Soc.* xx (1883) 150. —Madag.
parvifolium, *Hook. f. ex Benth. l. c.* iv. (1860) 37.— Borneo.
pedicellatum, *Spruce, ex Benth. l. c.* 36.—Ireas.
propinquum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 597.—Burma.
puberulum, *Klotzsch, ex Eichl. in Mart. Fl. Bras.* xiii. i. 507.—Guian.
racemosum, *Jacq. Enum. Pl. Carib.* 24.—Jamaic.
Racoubea, *Sw. Prod. Veg. Ind. Occ.* 86.—Guian.
ruiseanum, *Benth in Journ. Linn. Soc.* iv. (1860) 34. —Afr. austr.
Schlichii, *Kurz, For. Fl. Brit Burma*, i. 532.—Burma
senarium, [*Mog. & Sesse,*] *ex DC. Prod.* ii. 54.— Mexic.
spicatum, Lam. Encyc. i. 32 = H. Racoubea.
stipulaceum, *Wight, ex Mast. in Oliver, Fl. Trop. Afr.* ii. 496. Afr. trop.
sumatranum, *Miq. Fl. Ind Bat. Suppl.* 384.—Sumatra.
surinamensis, Steud. in Flora, xxvi. (1843) 756 = H. Racoubea
tetramerum, *Baker, in Journ Bot.* xx. (1882) 110.— Madag.
tomentosum, *Benth. in Journ. Linn. Soc.* iv. (1860) 34.—Burma; Malaya.
travancoricum, *Bedd. Fl. Sylv.* t. 211.—Ind or.
Vatkeanum, *O. Hoffm. Sert. Madag.* 19.—Madag.
vitiense, *Benth. in Journ. Linn. Soc.* iv. (1860) 36.— Ins. Fiji.
zeylanicum, *Benth. l. c.* 35.—Zeylan.

HOMALOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 353 (1840) = **Astragalus**, Tourn. (Legumin.).
brachycarpus, Nutt. l. c. 352 = A. caespitosus.
caespitosus, Nutt. l. c. = A. caespitosus.
campestris, Nutt. l. c. 351 = A. campestris.
canescens, Nutt. l. c. 352 = A. caespitosus.
decumbens, A. Gray, in Proc. Acad. Sc. Philad. (1863) 60 = A. campestris.
decumbens, Nutt. ex Torr. & Gray, Fl. N. Am. i. 351 = A. decumbens.
diffusus, Nutt. l. c. 350 = A. multiflorus.
junceus, Nutt. l. c. 351 = A. junceus.
multiflorus, Nutt. l. c. = A. multiflorus.
multiflorus, Torr. in Pacif. Rail. Rep. vii. 10 = A. Antiselli.
nigrescens, Nutt. ex Torr. & Gray, Fl. N. Am. i. 351 = A. multiflorus.
orthocarpus, Nutt. l. c. = A. junceus.
tenuifolius, Nutt. l. c. 352 = A. campestris.

HOMALOCALYX, F. Muell. in Hook. *Kew Journ.* ix. (1857) 309. *MYRTACEÆ*, Benth. & Hook. f. i. 699.
ericaeus, *F. Muell. l. c.*—Austral.
polyandrus, *F. Muell. ex Benth. Fl. Austral.* iii. 56; *et Fragm.* iv. 77.—Austral.

HOMALOCARPUS, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 348 = **Bowlesia**, Ruiz & Pav. (Umbellif.).
bowlesioides, Hook. & Arn. l. c. = B. nigripetala.

HOMALOCARPUS, Schur, Enum. Pl. Transs. 8 (1866) = **Anemone**, Linn. (Ranunculac.).
narcissiflorus, Schur, l. c. = A. narcissiflora.

HOMALOCENCHRUS, Mieg. ex Hall. Scirp. Helv. ii. 201 (1768) = **Leersia**, Schaod. (1788) (Gramin.).
oryzoides, Mieg, ex Pollich, Hist. Pl. Palat. i. 32 = L. oryzoides.

HOMALOCLADOS, Hook. f. in Benth. & Hook. f. Gen. ii. 123 (1873) = **Faramea**, Aubl. (Rubiac.).
colorata, Hook. f. l. c. = F. calycifora.
Sellowiana, Hook. f. l. c. = F. Sellowiana.

HOMALOCLINE, Reichb. Consp. 97 (1828) = OMALOCLINE, Cass. = **Crepis**, Linn. (Compos.).

HOMALODISCUS, Bunge, ex Boiss. Fl. Orient. i. 422 (1867) = **Ochradenus**, Delile (Resed.).
Aucheri, Boiss. l. c. = O. Aucheri.
major, Bunge, ex Boiss. l. c. 423 = O. Aucheri.
minor, Bunge, ex Boiss. l. c. = O. Aucheri.
Ochradeni, Boiss. l. c. 422.—Persia.

HOMALOLEPIS, Turcz. in Bull. Soc. Nat. Mosc xvi (1848) i. 575 = **Sinabis**, Aubl.
Blanchetii. Turcz. l. c. = S. Blanchetii.

HOMALOMENA, Schott, Melet. i. 20 (1832). *AROIDEÆ*, Benth. & Hook. f. iii. 982.
ADELONEMA, Schott, Prod. Aroid. 316 (1860).
CURMERIA, Linden & André, in Illustr. Hortic. xx. (1873) 45. t. 121.
CYRTOCLADON, Griff Notul. iii. 147 (1851)
HOMALONEMA, Endl. Gen. 288 (1837).
SPIROSPATHA, Rafin. Fl. Tellur. iv 8 (1836).
ZANTEDESCHIA, Spreng. Syst. iii 765 (1826), partim
alba, *Hasск. in Hort. & De Vriese, Tijdschr.* ix. (1842) 161.—Java.
aromatica, *Schott, Melet.* i. 20.—Ind or.; Malaya.
Beccariana, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 296.—Borneo.
bellula, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 126.—Java.
caerulescens, *Yungh. ex Schott, Prod. Aroid.* 310.— Java.
calyptrata, Kunth, Enum. Pl. iii. 57 = Schismatoglottis calyptrata.
cordata, Schott. Melet. i. 20 = aromatica.
cordata, Zoll. Syst. Verz. Pfl. 85. n. 559 = alba.
elegans, *Engl. in Bot. Jahrb.* i. (1881) 153.—Sumatra.
Gaudichaudii, Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 280 = aromatica.
insignis, *N. E. Br. in Illustr. Hortic.* xxxii. (1885) 93 t. 560.—Borneo.
major, Griff. Notul. iii 153 = minor.
minor, *Griff. l. c.* 152.—Malaya.
Miquelana, *Schott, in Miq. Ann. Mus. Bot. Lugd Bat.* i. 126.—Borneo.
occulta, Schott, Melet. i. 20 = aromatica.
ovata, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 296. —Borneo.
peltata, *Mast. in Gard. Chron.* (1877) 373. f. 46.— Peruv.; N. Granat.
picturata, *Regel, in Gartenfl.* (1877) 83; *Act. Hort. Petrop.* v. (1877) 269.—N. Granat.
Portes, Hort. Par.-ex Engl. in DC. Monog. Phan. ii. 385 = propinqua.
propinqua, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 280.—Borneo.
punctulata, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 296.—Borneo.
Roezelii, *Regel, Gartenfl.* (1877) 83; *Act. Hort. Petrop.* v. (1877) 270.—N. Granat.
rostratum, *Griff. Notul.* iii. 154 (roshalum).—Malaya.
rubescens, *Kunth, Enum. Pl.* iii. 57.—Ind. or.
rubescens, Miq. Fl. Ind. Bat. iii. 212 = rubra.
rubra, *Hasск. in Hort. & De Vriese, Tijdschr.* ix. (1842) 162.—Java.
sagittifolia, *Yungh. ex Schott, Prod. Aroid.* 311.— Java.
singaporense, *Regel, in Ind. Sem. Hort. Petrop.* (1869) 18.—Malaya.
sinuordata, *Engl. in Bot. Jahrb.* i. (1881) 153.— Borneo.
Wallichii, *Schott, in Bonplandia*, vii. (1859) 30.— Penang.
Wallisi, *Regel, in Gartenfl.* (1877) 83.—N. Granat.
Wendlandii, *Schott, Prod. Aroid.* 308.—Costa Rica.
Zollingeri, *Schott, in Bonplandia*, vii. (1859) 30.— Java.

HOMALONEMA, Endl. Gen. 288 (1837) = præc.

HOMALOSPERMUM, Schau. in Linnæa, xvii. (1843) 249 = **Leptospermum**, Forst. (Myrtac.).
firmum, Schau. l. c. = L. firmum.

HOMALOSTOMA, Stachégl. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 21 = **Andersonia**, R. Br. (Epacrid.).
simplex, Stachégl. l. c. = A. Homalostoma.

HOMALOTES, Endl. Gen. 487 (1838) = **Tanacetum**, Tourn. (Compos.).

HOMALOTHECA, Reichb. Nom. 89 (1841) = OMALOTHECA, Cass. = **Gnaphalium**, Linn. (Compos.).
supina, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 98 = G. supinum.

HOMANTHUS, H. B. & K. Nov. Gen. et Sp. iv. 12 (1820) = **Furaea**, Lag. (Compos.).
multiflorus, H. B. & K. l. c. 14 = P. multiflora.
pinnatifidus, H. B. & K. l. c. 308 = P. pinnatifida.
pungens, H. B. & K. l. c. 13 = P. pungens.

HOMBAK, Adans. Fam. ii. 408 (1763) = **Capparis**, Linn.

HOMBRONIA, Gaudich. Bot. Voy. Bonite, t. 22 (1844–52) = **Pandanus**, Linn.
calathiphora, Gaudich. in Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 38 = P. calathiphorus.
edulis, Gaudich. Bot. Voy. Bonite, t. 22. f. 17 = P. dubius.

HOMEOPLITIS, Endl. Gen. 107, in syn. (1836) = HOMOPLITIS, Trin. = **Pogonatherum**, R. Br. (Gramin.).

HOMERIA, Vent. Dec. Gen. Nov. 5 (1808). *IRIDEAE*, Benth. & Hook. f. iii. 692.
aurantiaca, Sweet, Hort. Brit. ed. II. 498 = collina.
collina, *Vent. Dec. Gen. Nov.* 5.—Afr. austr.
elegans, *Sweet, Hort. Brit.* ed. I. 395.—Afr. austr.
exalbata, Sweet, l. c. = collina.
flaccida, Sweet, l. c. = collina.
flexuosa, Baker, in Sandd. Refug. Bot. t. 24 = Hexaglottis longifolia.
flexuosa, Sweet, Hort. Brit. ed. I. 395 = Sisyrinchium cuspidatum.
linearis, *Sweet, Brit. Flow. Gard.* t. 178.—Afr. austr.
maculata, *Klatt, in Linnaea*, xxxiv. (1865–66) 627 — Afr. austr.
minuta, Sweet, Brit. Flow. Gard. t. 152 = collina.
ochroleuca, Sallsb. in Trans. Hort. Soc. i. (1812) 308 = collina.
parvifolia, Sweet, Hort. Brit. ed. II. 498 = linearis.
spicata, Sweet, l. c. ed. I. 395 = elegans.
virgata, Sweet, l. c. = Hexaglottis virgata.

HOMOCENTRIA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 308 = **Oxypora**, DC. (Melastom.).
vaguaz, Naud. l. c. t. 15. f. 4 = O. vagans.

HOMOCHAETE, Benth. in Hook. Ic. Pl. t. 1110 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 331.
conferta, *Benth. l. c.*—Afr. austr.

HOMOCHROMA, DC. Prod. v. 324 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 258.
Echlonis, *DC. l. c.*—Afr. austr.

HOMOCNEMIA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40 = **Stephania**, Lour. (Menispерm.).
Meyeriana, Miers, ex Harv. & Sond. Fl. Cap. i. 10 = S. Meyeriana.
umbellata, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40 = S. Meyeriana.

HOMOEANTHERUM, Steud. Nom. ed. II. i. 774 (1840) = HOMOEATHERUM, Nees = **Andropogon**, Linn. (Gramin.).

HOMOEANTHUS, Spreng. Syst. iii. 508 (1826) = HOMOIANTHUS, Bonpl. = **Furaea**, Lag. (Compos.).

HOMOEATHERUM, Nees, in Hook. & Arn. Bot. Beech. Voy. 289 (1836). = **Andropogon**, Linn. (Gramin.).
chinense, Nees, l. c. = A. apricus.

HOMOGLOSSUM, Salisb. in Trans. Hort. Soc. i. (1812) 335 = **Anthotysa**, Linn. (Irid.).
lucidior, Baker, in Journ. Linn. Soc. xvi. (1877) 161 = A. lucidior.
Merianella, Baker, l. c. = A. Merianella.
praecox, Salisb. in Trans. Hort. Soc. i. (1812) 325 = A. revolutus.
revolutum, Baker, in Journ. Linn. Soc. xvi. (1877) 161 = A. revolutus.

HOMOGYNE, Cass. in Bull. Soc. Philom. (1816) 198; et Dict. Sc. Nat. xxi. 412 (1821). *COMPOSITAE*, Benth. & Hook. f. ii. 439.
alpina, *Cass. l. c.* 413.—Europ.
discolor, *Cass. l. c.* 413.—Europ.
montana, C. Froel. Alpenfl. v. t. 5 = alpina?
sylvestris, *Cass. in Dict. Sc. Nat.* xxi. 413.—Austral.

HOMOIANTHUS, Bonpl. ex DC. in Ann. Mus. Par. xix. (1812) 65 = **Furaea**, Lag. (Compos.).
ambiguus, Cass. Opusc. ii. 107 = P. squarrosa.
cubatanensis, Less. ex DC. Prod. vii. 64 = P. cubatanensis.
Donianus, Kenny, in C. Gay, Fl. Chil. iii. 423 = P. recurvata.
echinulatus, Cass. in Dict. Sc. Nat. xxxviii. 458 = P. recurvata.
Gayanus, Remy, in C. Gay, Fl. Chil. iii. 419 = P. lyrata.
inermis, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 290 = P. inermis.
laevis, DC. Prod. vii. 64 = P. laevis.
linearis, DC. l. c. = P. linearis.
lyratus, Remy, in C. Gay, Fl. Chil. iii. 418 = P. lyrata.
magellanicus, DC. Prod. vii. 63 = P. lactucoides.
multiflorus, D. Don, in Trans. Linn. Soc. xvi. (1830) 211 = P. multiflora.
palustris, Phil. in Anal. Univ. Chil. (1865) ii. 816.—Chili.
pectinatus, Phil. l. c. (1862) ii. 836; et in Linnaea, xxxiii. (1864–65) 124.—Chili.
pinnatifidus, D. Don, in Trans. Linn. Soc. xvi. (1830) 209 = P. pinnatifida.
pristiphyllus, Remy, in C. Gay, Fl. Chil. iii. 419 = P. pristiphylla.
pungens, DC. Mém. Labiatif. t. 3. f. 5 = P. pungens.
rigidus, Phil. in Linnaea, xxviii. (1856) 719.—Chili.
scaber, Benth. Pl. Hartw. 186 = P. pungens.
squarrosus, DC. Prod. vii. 64 = P. squarrosa.
tomentosus, Poepp. ex DC. l. c. 60 = Leucheria Bridgesii.
viscosus, DC. l. c. 64 = P. viscosa.

HOMOIOCELTIS, Blume, Mus. Bot. Lugd. Bat. ii. 64. t. 84, 1852 [1856] = **Aphanantha**, Planch. (Urticac.).
aspera, Blume, l. c. = A. aspera.

HOMOLOSTYLES, Hook. et Wight, Contrib. 50 (1834) = **Tylophora**, R. Br. (Asclep.).
tenerrima, Wall. ex Wight, l. c. 51 = T. tenerrima.

HOMONOIA, Lour. Fl. Cochinch. 686 (1790). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 292.
HARMATOSPERMUM, Wall. Cat. nn. 7953–55 (1829?)
LUMANAJA, Blanco, Fl. Filip. 821 (1837).
SPATHIOSTEMON, Blume, Bijdr. 621 (1826).
javensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 200. —Java.
retusa, *Muell. Arg. l. c.*—Ind. or.
riparia, *Lour. Fl. Cochinch.* 687.— Cochinch.
symphyllacifolia, *Kurz, in Flora*, lvii. (1875) 92.—Reg. Himal.

HOMONOMA, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 369. *MELASTOMACEAE*.
aridum, *Bello, l. c.*—Ins. Porto Rico.

HOMOPAPPUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 280 = **Aplopappus**, Cass. (Compos.).
argutus, Nutt. l. c. 331 = A. racemosus.
glomeratus, Nutt. l. c. = A. racemosus.
insulcatus, Nutt. l. c. 332 = A. uniflorus.
multiflorus, Nutt. l. c. = A. lanceolatus.
paniculatus, Nutt. l. c. 331 = A. racemosus.
racemosus, Nutt. l. c. 332 = A. racemosus.
spathulatus, Nutt. l. c. = Solidago spathulata.
squarrosus, Nutt. l. c. = A. squarrosus.
uniflorus, Nutt. l. c. = A. uniflorus.

HOMOPLITIS, Trin. Fund. Agrost. 166 (1820) = **Pogonatherum**, Beauv. (Gramin.).
crinita, Trin. l. c. = P. saccharoideum.

HOMORANTHUS, A. Cunn. ex Schau. in Linnaea, x. (1836) 310. *MYRTACEAE*, Benth. & Hook. f. i. 697.
ENGOANTHES, A. Cunn. ex Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 191.
flavescens, A. Cunn. ex Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1842) 192 = virgatus.
virgatus, *A. Cunn. ex Schau. l. c.* 193.—Austral.

HOMOSTYLES, Wall. ex Hook. f. Fl. Brit. Ind. iv. 44 (1855) = **Tylophora**, R. Br. (Asclep.).
tenerrima, Wall. ex Hook. f. l. c. = T. tenerrima.

HOMOSTYLIUM, Nees, in Linnaea, xvlii. (1844) 513 = **Microglossa**, DC. (Compos.).
cabulicum, Nees, l. c. = M. albescens.

HOMOTROPIUM, Nees, in Mart. Fl. Bras. ix. 47. t. 2 (1847) = **Ruellia**, Linn. (Acanthac.).
erythrorhizum, Nees, l. c. 48.—Bras.

HONCKENEJA, Maxim. in Mém. Sav. Etr. Pétersb. ix. (1859) 56 = HONKENYA, Ehrh. = **Arenaria**, Rupp. (Caryoph.).

HONCKENEYA, Steud. Nom. ed. II. i. 771 (1840) = **Honckenya**, Willd. (Tiliac.).

HONCKENEYA, Steud. Nom. ed. II. i. 771 (1840) = HONKENYA, Ehrh. = **Arenaria**, Rupp. (Caryoph.).

HONCKENIA, Pers. Syn. i. 416 (1805) = **Honckenya**, Willd. (Tiliac.).

HONCKENIA, Rafin. in Am. Monthly Mag. (1818) 266 = seq.

HONCKENYA, Bartl. Ord. 308 (1830) = HONKENYA, Ehrh. = **Arenaria**, Rupp. (Caryoph.).

HONCKENYA, Willd. in Usteri, Delect. Opusc. Bot. ii. (1793) 200. t. 4. *TILIACEAE*, Benth. & Hook. f. i. 285.
CLAPPERTONIA, Meisn. Gen. 86 (1837).
HONKENYA, Willd. ex Cothen. Meth. Pl. Disp. 19 (1795).
ficifolia, *Willd. in Usteri, Delect. Opusc. Bot.* ii. (1793) 201. t. 4.—Afr. trop.
minor, *Baill. Adansonia*, x. (1872) 183.—Afr. trop.

HONCKNEYA, Spach, Hist. Vég. Phan. iv. 3 (1835) = **Honckenya**, Willd. (Tiliac.).

HONDBESSEN, Adans. Fam. ii. 158 (1763) = **Paederia**, Rupp. (Rubiac.).

HONKENEJA, Endl. Gen. 966 (1840) = seq.

HONKENYA, Ehrh. Beitr. ii. 180 (1788) = **Arenaria**, Rupp. (Caryophyll.).
oblongifolia, Torr. & Gray, Fl. N. Am. i. 176 = A. peploides.
peploides, Ehrh. Beitr. ii. 181 = A. peploides.

HONKENYA, Willd. ex Cothen. Meth. Pl. Disp. 19 (1795) = Honckenya, Willd. (Tiliac.).
Willdenowii, J. F. Gmel. Syst. 609 = Honckenya ficifolia?

HONORIUS, S. F. Gray, Nat. Arr. Brit Pl. ii. 177 (1821) = Ornithogalum, Tourn. (Liliac.).
autumn, S. F. Gray, l. c. = O. autumn.

HONOTTIA, Reichb. Consp. 178 (1828) = **Limnophila**, R. Br. (Scrophul.).

HOODIA, Sweet, Hort. Brit. ed. II. 269 (1830). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 788.
MONOTHYLACIUM, G. Don, Gen. Syst. iv. 116 (1837).
SCYTANTHUS, Hook. Ic. Pl. t. 645, 646 (1844).
Bainii, *Dyer, in Bot. Mag.* t. 6348.—Afr. austr.
Barklyi, *Dyer, in Journ. Linn. Soc.* xv. (1876) 253.—Afr. austr.
Carrei, *Decne. in DC. Prod.* viii. 668.—Afr. trop.
Gordoni, *Sweet, Hort. Brit.* ed. II. 269.—Afr. austr.

HOOKERA, Salisb. Parad. Lond. t. 98 (1808) = **Brodiæa**, Sm. (1811) (Liliac.).
coronaria, Salisb. l. c. = B. grandiflora.
pulchella, Salisb. l. c. sub t. 98; t. 117 = B. congesta.

HOOKIA, Neck. Elem. i. 70 (1790) = **Centaurea**, Linn. (Compos.).

HOOPESIA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 458 = **Cercodium**, Tul. *etc.* (Legum.).
arborea, Buckl. l. c. = C. texanum, Acacia flexicaulis, amentacea.

HOOREBECKIA, Steud. Nom. ed. II. i. 774 (1840) = seq.

HOOREBEKIA, Cornelius. in Mussch. Hort. Gand. 120 (1817); ex DC. Prod. v. 846 (1836) = **Aplopappus**, Cass. (Compos.).
chilensis, Cornelius. l. c. = A. Hoorebekia.

HOPEA, Linn. Mant. i. 14 (1767) = **Symplocos**, Linn. (Styrac.).
racemosa, Dals. & Gibs. Bomb. Fl. 140 = S. Beddomei.
spicata, Dals. & Gibs. l. c. = S. spicata.
ternifiora, Orteg. Hort. Matr. Dec. 181. t. 17.—N. Granat.
tinctoria, Linn. Mant. i. 105 = S. tinctoria.

HOPEA, Roxb. Hort. Beng. 42 (1814); Pl. Corom. iii 7. t 210 (1819). *DIPTEROCARPEÆ*, Benth. & Hook. f. i. 198.
NEESANDRA, Rafin. Sylva Tellur. 168 (1838).
PETALANDRA, Hassk. Hort. Bogor. 104 (1858).
aspera, *De Vriese, Minjak Tangkawang*, 28.—Borneo.
Balangeran, Korth. Verh. Nat. Gesch. Bot. 74. t. 7 = (Shorea sp.).—Malaya.
cernua, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 252.—Sumatra.
dealbata, *Hance, in Journ. Bot.* xv. (1877) 329.—Cambodia.
decandra, Buch.-Ham. ex Wight, Illustr. 88 = odorata.
discolor, *Thw. Enum. Pl. Zeyl.* 36.—Zeylan.
diversifolia, *Miq. Fl. Ind. Bat. Suppl.* 493.—Sumatra.
dryobalanoides, *Miq. l. c.* 492.—Sumatra.
eglandulosa, Roxb. Hort. Beng. 42; Fl. Ind. ii. 611 = Cyclostemon eglandulosum.
fagifolia, *Miq. Fl. Ind. Bat. Suppl.* 490.—Ins. Banca.
faginea, Hort. Calc. ex A. DC. in DC. Prod. xvi. ii. 582 = odorata.
fagifera, Wall. Cat. n. 968 = Vatica faginea.
floribunda, Wall. Cat. n. 964 = Shorea floribunda.
glabra, *Wight & Arn. Prod.* 85.—Ind. or.
grandifolia, *Miq. Fl. Ind. Bat. Suppl.* 490.—Sumatra.
grandiflora, Wall. Cat. n. 965 = Vatica grandiflora.
gratissima, Wall. l. c. n. 960 = Shorea gratissima.
Griffithii, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 60.—Burma.
jucunda, *Thw. Enum. Pl. Zeyh.* 403.—Zeylan.
lanceolata, De Vriese, Minjak Tangkawang, 32 = Vateria borneensis.
longifolia, Dyer, in Hook. f. Fl. Brit. Ind. i. 309 = Balanocarpus utilis.
lucida, *Thunb. Ic. Jap.* t. 14.—Japon.
macrophylla, De Vriese, Minjak Tangkawang, 28 = Shorea maritima.
malabarica, *Bedd. Ic. Pl. Ind. Or.* i. 42.—Ind. or.
Maranti, *Miq. Fl. Ind. Bat. Suppl.* 489.—Sumatra.
Mengarawan, *Miq. l. c.* 491.—Sumatra.
micrantha, Hance, in Journ. Bot. xiv. (1876) 242 = Pierrea.
micrantha, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 161.—Malaya.
modesta, *Thw. ex A. DC. in DC. Prod.* xvi. ii. 633.—Zeylan.
myrtifolia, *Miq. Fl. Ind. Bat. Suppl.* 493.—Sumatra.
oblongifolia, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 309.—Burma.
odorata, *Roxb. Hort. Beng.* 42; *Pl. Corom.* iii. 7. t. 210; *Fl. Ind.* ii. 609.—Burma.
parviflora, *Bedd. Fl. Sylv.* i. t. 7.—Ind. or.
philippinensis, *Dyer, in Journ. Bot.* xvi. (1878) 100.—Ins. Philipp.
Pierrei, *Hance, in Journ. Bot.* xv. (1877) 329, in nota.—Cambodia.
racophlœa, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 310.—Ind. or.
Sangal, *Korth. in Verh. Nat. Gesch. Bot.* 75.—Borneo.
Scaphula, Roxb. Hort. Beng. [98]; Fl. Ind. ii. 611 = Vatica Scaphula.
sclanica, Wight & Arn. Prod. 85 = Shorea sclanica.
semicta, *De Vriese, Minjak Tangkawang*, 93.—Ins. Celebes.
sericea, *Blume, Mus. Bot. Lugd. Bat.* ii. 85.—Borneo.
Singkawang, *Miq. Fl. Ind. Bat. Suppl.* 489.—Sumatra.
Siranda, *Miq. l. c.* 491.—Sumatra.
speciosa, Pritz. Ic. Ind. 555, sphalm. vide *Hoppea speciosa*.
splendida, *De Vriese, Minjak Tangkawang*, 28.—Borneo.
squamata, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 239 = Shorea squamata.
suavis, Wall. Cat. n. 959 = Shorea siamensis.
Tangili, *Blume, Mus. Bot. Lugd. Bat.* i. 35.—Malaya.
vasta, Wall. Cat. n. 962 = odorata.
Wightiana, Miq. ex Dyer, in Hook. f. Fl. Brit. Ind. i. 308 = odorata.
Wightiana, *Wall. Cat.* n. 6295.—Ind. or.

HOPEA, Vahl, Enum. i. 3 (1805) = **Hoppea**, Willd. (Gentian.).
dichotoma, Vahl, l. c. = Hoppea dichotoma.

HOPKIRKIA, DC. Prod. v. 660 (1836) = **Sohkuhria**, Roth (Compos.).
anthemoidea, DC. l. c. = S. Hopkirkia.

HOPKIRKIA, Spreng. Nov. Prov. 23 (1818) = **Salmea**, DC. (Compos.).
Eupatoria, Spreng. l. c. = S. Eupatoria.
fruticulosa, Spreng. Syst. iii. 444 (= *Armania frutescens*).—Am. austr.
hirsuta, Spreng. l. c. = S. hirsuta.
scandens, Spreng. l. c. 443 = S. Eupatoria.

HOPLISMENUS, Hassk. Cat. Hort. Bog. Alt. 16 (1844) = **Opliamenus**, Beauv. (Gramin.).

HOPLOPHYLLUM, DC. Prod. v. 73 (1836). *COMPOSITÆ*, Benth. & Hook. f. ii. 231.
ferox, *Sond. in Harv. & Sond. Fl. Cap.* iii. 54.—Afr. austr.
spinosum, *DC. Prod.* v. 74.—Afr. austr.

HOPLOPHYTUM, Beer, in Flora, xxxvii. (1854) 545; et Bromel. 129 (1857) = **Aechmea**, Ruiz & Pav. (Bromel.).
angustifolium, Beer, Bromel. 132 = A. spicata.
augustum, Beer, l. c. 136 = A. augusta.
aureum, Ann. in Illustr. Wien. Gart. Zeit. (1861) 97. t. 1 = Billbergia atro-rosea?
bracteatum, C. Koch, Wochenschr. iii. (1860) 306 = A. bracteata.
calyculatum, Morr. in Belg. Hortic. xv. (1865) 162 = A. calyculata.
coelesta, C. Koch, Ind. Sem. Hort. Berol. (1856) 6 = A. coelestis.
corruiescens. E. Morr. Cat. (1873) 9 = A. caerulescens.
comosum, Beer, Bromel. 150 = A. comata.
cyaneum, Beer, l. c. 131 (= *Nidularium cyaneum*).—Bras.
distichantoides, Beer, l. c. 136 = A. distichantha.
fasciatum, Beer, l. c. 129 = A. fasciata.
giganteum, E. Morr. in Belg. Hortic. xxiv. (1874) 174. —Hab.?
lanuginosum, Beer, Bromel. 138 = A. nudicaulis.
Lindeni, E. Morr. in Belg. Hortic. xv. (1865) 164 = A. Lindeni.
lingulatum, Beer, Bromel. 139 = A. linguiata.
Mertensii, Beer, l. c. 134 = A. Mertensii.
mucronifiora, Beer, l. c. = A. mucronifiora.
nudicaule, C. Koch, Ind. Sem. Hort. Berol. (1856) 6 = A. nudicaulis.
paniculatum, Beer, Bromel. 130 = A. paniculata.
platynema, Beer, l. c. 139 = A. platynema.
Platzmanni, E. Morr. in Belg. Hortic. xxv. (1875) 362, nomen = A. floribunda.
polystachium, Beer, Bromel. 137 = A. brasiliensis?
purpureo-roseum, Beer, l. c. 135 = A. suaveolens.

HOPLOPHYTUM :—
spicatum, Beer, Bromel. 134 = Aechmea spicata.
spicatum, Beer, l. c. 140 = A. nudicaulis.
suaveolens, Beer, l. c. 135 = A. suaveolens.
tetrastachyum, Beer, l. c. 137.—Bras.
unispicatum, Beer, l. c. 138 = A. nudicaulis.

HOPLOTHECA, Spreng. Syst. iv. Cur. Post. 52 (1827) = OPLOTHECA, Nutt. = **Froelichia**, Moench (Amarant.).

HOPPEA, Endl. Gen. 744, in syn. (1839) = HOPEA, Linn. = **Symplocos**, Linn. (Styrac.).

HOPPEA, Endl. Gen. 1014 (1840) = **Hopea**, Roxb. (Dipterocarp.).

HOPPEA, Reichb. in Flora, vii. (1824) 245 = **Senecio**, Tourn. (Compos.).
sibirica, Reichb. l. c. = S. Ligularia.
speciosa, Reichb. l. c. = S. Ligularia.

HOPPEA, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 434. *GENTIANEÆ*, Benth. & Hook. f. ii. 805.
HOPEA, Vahl, Enum. i. 3 (1805).
HOPPIA, Spreng. Anleit. ii. 11. 889 (1818).
dichotoma, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 435.—Ind. or.
fastigiata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 100.—Ind. or.

HOPPIA, Nees, in Mart. Fl. Bras. ii. t. 199. t. 80 (1842). *CYPERACEÆ*, Benth. & Hook. f. iii. 1069.
angustifolia, *Boeck. in Flora*, liv. (1871) 37.—Guian.
irrigua, *Nees, in Mart. Fl. Bras.* ii. t. 199.—Bras. microcephala, *Boeck. in Flora*, liv. (1871) 37.—Guian.

HOPPIA, Spreng. Anleit. ii. 11. 889 (1818) = **Hoppea**, Willd. (Gentian.).

HORANINOVIA, Fisch. & Mey. in Schrenk, Enum. Pl. Song. 10 (1841). *CHENOPODIACEÆ*, Benth. & Hook. f. iii. 70.
anomala, Moq. in DC. Prod. xiii. ii. 170 = juniperina.
juniperina, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 468 = ulicina.
juniperina, *C. A. Mey. in Schrenk, Enum. Pl. Song.* 11.—Turkest.; Sibir.
minor, *Schrenk, l. c.*—Sibir.
ulicina, *Fisch. & Mey. in Schrenk. l. c.* 10.—Turkest.; Sibir.

HORANTHES, Rafin. New Fl. Am. iii. 80 (1836) = **Leuchea**, Linn. (Cistin.).
arenaria, Rafin. Sylva Tellur. 133 (= *Anthelis arenaria*).—Am. bor.
podanisia, Rafin. l. c. (= *Anthelis podanisia*).—Am. bor.
tripetala, Rafin. New Fl. Am. iii. 80 = L. Skinneri.

HORANTHUS, Rafin. Sylva Tellur. 132 (1838) = præc.

HORAU, Adans. Fam. ii. 80 (1763) = **Lagunclaria**, Gaertn. (Combret.).

HORDEUM, [Tourn.] Linn. Syst. ed. I (1735). *GRAMINEÆ*, Benth. & Hook. f. iii. 1206.
CRITHSION, Rafin. in Journ. Phys. lxxxix. (1819) 108.
CRITHO. Rœm. et Schult. Syst. Veget. (1818) 5; et in Ann. Sc. Nat. Sér. III. xi. (1849) 358.
CUVIERA, Koel. Gram. 298 (1802).
ZEOCRITON, Beauv. Agrost. 114. t. 21. f. 2 (1812).
adpressum, Moench, Meth. 199 = Elymus sibiricus.
adscendens, *H. B. & K. Nov. Gen. et Sp.* i. 160.—Mexic.
Aegiceras, Royle, ex Walp. Ann. iii. 787 = vulgare.
ambiguum, *Doell, in Mart. Fl. Bras.* ii. iii. 232.—Reg. Argent. (e Creta ?).
andicola, *Griseb. in Goett. Abh.* xxiv. (1879) 285.—Reg. Argent.
andinum, *Trin. in Linnaea*, x. (1835–36) 304.—Reg. Andin.

HORDEUM:—
arenarium, Aschers. Fl. Brandenb. 874 = Elymus arenarius.
arenarium, Steud. Nom. ed. I. 418 = Arrhenatherum avenaceum.
Bertero-anum, *E. Desv. ex C. Gay, Fl. Chil* vi. 460. —Chili.
bifarium, *Roth, Nov. Pl. Sp.* 59.—Sicilia.
brevicornum, Presl, Bot. Bemerk. 120 = bulbosum.
brevisubulatum, Link, in Linnaea, xvii. (1843) 891 = secalinum.
bulbosum, *Linn. Cent. Pl.* ii. 8.—Reg. Mediterr.; Oriens.
bulbosum, Sieber, ex Kunth, Enum. Pl. i. 449 = Secale creticum.
caducum, *Munro, ex Aitch. in Journ. Linn. Soc.* xviii. (1880) 110, nomen.—Afghan.
capense, *Thunb. Prod. Pl. Cap* 28.—Afr. austr.
Caput-Medusæ, Loret & Barr. Fl. Monsp. 766 = Elymus Caput-Medusæ.
cordilignicum, Moench, Meth. 199 = Elymus virginicus.
chilense, *Brongn. in Duperr. Voy. Coq. Bot.* 54.—Chili.
chilense, Roem. & Schult. Syst. ii. 796 = maritimum.
ciliatum, Gilib. Exercit. ii. 520 = murinum.
ciliatum, Lam. Encyc. iv. 604 = bulbosum.
coeleste, Viborg, ex Kunth, Enum. Pl. i. 455 = vulgare.
comosum, *J. C. Presl, Rel. Haenk.* i. 327.—Chili.
complanatum, *Beauv. Agrost.* 115.—Europ. austr.
compressum, Bois. & Orph. ex Boiss. Fl. Orient. v. 681 = Lolium rigidum.
compressum, *Grisеb. in Gortt. Abh.* xix. (1874) 249.—Reg. Argent.
crinitum, Desf. Fl. Atlant. i. 113 = Elymus Caput-Medusæ.
cylindricum, Murr. Prod. Gotting. 48 = Elymus europæus.
cylindricum, *Steud. Syn. Pl. Gram.* 353.—Chili.
Decaisnei, Hort. Par. ex Boiss. Fl. Orient. v. 688 = ithaburense.
deficiens, *Steud. ex A. Br. Del. Sem. Hort. Frib.* (1848) 2.—Abyssin.
depauperatum, *Steud. Syn. Pl. Gram.* 353.—Chili.
distichon, *Linn. Sp. Pl.* 85.—(Stirps cult.).
divergens, *Nees & Meyen, ex Steud. Nom.* ed. II. i. 776.—Chili.
elymoides, Vest, ex Kunth, Enum. Pl. i. 442 = Elymus europæus.
euclaston, *Steud. Syn. Pl. Gram* 353.—Am. bor.
europæum, All. Fl. Pedem. ii. 260 = Elymus europæus.
flexuosum, *Steud. Syn. Pl. Gram.* 353.—Chili!
fragile, *Boiss. Diagn.* Ser. I. vii. 128.—As. Min.; Persia.
fragile, *Godr. in Mém. Acad. Montp. (Sect. Medic.).* I. (1853) 455.—Hab.!
geniculatum, All. Fl. Pedem. ii. 259 = maritimum.
geniculatum, Bieb. Fl. Taur. Cauc. i. 83 = secalinum, maritimum.
glaucum, *Steud. Syn. Pl. Gram.* 353.—Arab.
Gussoneanum, *Parl. Pl. Palerm.* 244, in obs.—Sicil.
halophilum, *Griseb. in Gortt. Abh.* xix. (1874) 249.—Reg. Argent.
heterostychyon, Beauv. Agrost. 114 = hexastichon.
hexastichon, *Linn. Sp. Pl* 85.—(Stirps cult.).
hexastichum, Auct. = pracc.
himalayense, *Schult. Mant.* ii. 481.—Reg. Himal.
hirsutum, *Bertol. Misc. Bot.* i. 11.—Mesopot.
Hystrix, Roth, Catalect. i. 23 = maritimum.
imbrbe, Arduini, ex Roem. & Schult. Syst. ii 798 = distichon.
inclusum, *Forsk. Fl. Aegypt. Arab.* p. xix.—Aegypt.
incrnium, Poir. in Lam. Encyc. iv. 605, sphalm. = pracc.
ithaburense, *Boiss. Diagn.* Ser. I. xiii. 70.—As. Min.; Oriens.
jubatum, DC. Fl. Fr. vi. 286 = Elymus Caput-Medusæ.
jubatum, *Linn. Sp. Pl.* 85.—Am. bor. et austr.
Kaufmanni, *Regel, in Act. Hort. Petrop.* vii. (1880) 578.—As. centr.
leporinum, Link, in Linnaea, ix. (1835) 138 = murinum.
leptostachys, *Griff. Itin. Notes*, 270.—Afghan.
lyratum, Boiss. ex Fl. Orient. v. 688 = bulbosum.
maximum, *Steud. Syn. Pl. Gram.* 352.—Dahuria.
macrolepis, *A. Br. Del. Sem. Hort. Frib.* (1848) 2, adn. —Abyssin.

HORDEUM:—
marinum, Huds. Fl. Angl. ed. II. i. 57 = maritimum.
maritimum, Roth, Tent. Fl. Germ. ii. i. 150 = secalinum.
maritimum, *With. Arr. Brit. Pl.* 172.—Europ.; Afr. bor.; Oriens.
maximum, Vill. Fl. Delphin. 10 = secalinum.
montanum, Schrank, Baier. Fl. i. 886 = Elymus europæus.
murinum, *Linn. Sp. Pl.* 85.—Europ.; Afr. bor.; Oriens.
muticum, *J. & C. Presl, Rel. Haenk.* i. 327.—Peruv.
nepalense, Sweet, Hort. Brit. ed. I. 444 = vulgare.
nigrum, Willd. Enum. Hort. Berol. 1037 = vulgare.
nodorum, Bleb. Fl. Taur. Cauc. 72 = violaceum.
nodosum, Linn. Sp. Pl. ed. II. 126 = secalinum.
nodosum, Ucria, Hort. R. Panorm. 64; Savi, Duc Cent. ii. 87 = bulbosum.
nudum, *Arduini, ex Schult. Mant.* ii. 487 (Stirps cult.).
patulum, Moench, Meth. 199 = Elymus canadensis.
perseum, *Forsk. Fl. Aegypt. Arab.* p. xix.—Arab.
pilosum, *Steud. Syn. Pl. Gram.* 353.—Chili.
plumosum, Gilib. Exercit. 520 = vulgare.
polystichon, Hall. f. in Nov. Comm. Gottng. vi. (1776) 5 = vulgare.
pratense, Huds. Fl. Angl. ed. II. 56 = secalinum.
Pseudo-murinum, Tapp. ex Koch, Syn. Fl. Germ. ed. II. 952 = murinum.
pubescens, Guss. Fl. Slc. Prod. i. 144 = maritimum.
publiflorum, *Hook. f. Fl. Antarct.* 388.—Reg. Magellan.
pusillum, Nutt. Gen. Am. i. 87 = secalinum.
Richlii, *Steud. Syn. Pl. Gram* 353.—Am. bor.
rigidum, Roth, Catalect. fasc. i. 24 = maritimum.
Rothii, *Lonk, in Linnaea*, xvii. (1843) 393.—German.
rubens, Willk. in Oestr. Bot. Zeitschr. xxv. (1875) 109 = murinum.
sativum, Pers. Syn. i. 106 = vulgare.
secalinum, Guss. Pl. Rar. fl, 88. t. 11 = Gussoneanum.
secalinum, Savi, Fl. Pis. i. 145 = maritimum.
secalinum, *Schreb. Spicil. Fl. Lips.* 148.—Reg. bor. temp.
sibiricum, Link, ex Steud. Nom. ed. II. i. 775 = secalinum.
Subcrinitum, Bess. ex Schult. Mant. ii. 662 = Secale creticum.
spontaneum, C. Koch, in Linnaea, xxi. (1848) 430 = ithaburense.
stenostachys, *Godr. in Mém. Acad. Montp. (Sect. Médic.).* i. (1853) 455.—Hab.!
strictum, Desf. Fl. Atlan. i. 113. t. 37 = bulbosum.
subfastigiatum, *Doell, in Mart. Fl. Bras.* ii. iii. 234.—Reg. Argent.
sylvaticum, Huds. Fl. Angl. ed. II. i. 57 = Elymus europæus.
tetrastichum, Stokes, Bot. Mat. Med. i. 166 = vulgare.
trifurcatum, *Yacq. ex Baill. in Bull. Soc. Bot. Fr.* i. (1854) 97.—Hab.!
utriculatum, *Bert. in Mercv. Chil.* 1829; *ex Am. Journ. Sc.* xxiii. (1833) 86.—Chili.
vaginatum, C. Koch, in Linnaea, xxi. (1848) 433 = murinum.
villosum, Moench, Meth. 199 = Elymus arenarius.
villosum, Steud. Syn. Pl. Gram. 352 = hirsutum
violaceum, *Boiss. & Hohen. Diagn.* Ser. I. viii. 70.—As. Min.
vulgare, *Linn. Sp. Pl.* 84 (Stirps cult.).
Wittii, Hack. in Oestr. Bot. Zeitschr. xxvii. (1877) 49 = Gussoneanum.
Zeocriton, *Linn. Sp. Pl.* 85 (Stirps cult.).

HORKELIA, Cham. & Schlecht. in Linnaea, ii. (1827) 26 – **Potentilla**, Linn. (Rosac.).
Bolanderi, A. Gray, in Proc. Am. Acad. vii. (1868) 388. —Cal.
californica, Cham. & Schlecht. in Linnaea, ii. (1827) 26 = P. multijuga.
capitata, Lindl. Bot. *Reg.* sub t. 1997.—Am. bor. occ.
capitata, Regel, Gartenfl. (1872) t. 711 = P. Haematochrus.
capitata, Torr. in Pacif. Rail. Rep. iv. 84 = P. multijuga.
congesta, *Dougl. in Hook. Bot. Mag.* t. 2880 = P. congesta.
congesta, Newberry, in Pacif. Rail. Rep. vi. 78 = P. congesta?
cuneata, Lindl. Bot. Reg. sub t. 1997 = P. multijuga.

HORKELIA:—
cuneata, Newberry, in Pacif. Rail. Rep. vi. 78 = Potentilla Haematochrus.
Douglasiana, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 3. 425 = P. multijuga.
fusca, Lindl. Bot. Reg. t. 1997 = P. Haematochrus?
Gordoni, Hook. Kew Journ. v. (1853) 341 = Ivesia Gordoni.
grandis, Hook. & Arn. Bot. Beech. Voy. 389 – P. multijuga.
hirsuta, Lindl. Bot. Reg. sub t. 1997 = P. congesta.
multifoliolata, Torr. in Sitgreaves, Rep. 169 = Ivesia Gordoni.
parviflora, Nutt. ex Torr. & Gray, Fl. N. Am. i. 425 = P. Haematochrus?
pilosa, Nutt. ex Torr. & Gray, l. c. 424 = P. congesta.
purpurascens, S. Wats. in Proc. Am. Acad. xi. (1876) 148.—Calif.
sericata, S. Wats. l. c. xx. (1885) 364.—Am. bor. occ
tenuiloba, A. Gray, in Proc. Am. Acad. vi. (1866) 529. —Calif.
Tilingi, Regel, in Act. Hort. Petrop. i. (1871) 153 = P. tridentata.
tridentata, Torr. in Pacif. Rail. Rep. iv. 84. t. 6 = P. tridentata.

HORKELIA, Reichb. ex Bartl. Ord. 76 (1830) = **Wolfia**, Horkel (Lemnac.).

HORMIDIUM, Lindl. ex Heynh. Nom. i. 880 (1840). *ORCHIDEÆ*, Benth. & Hook. f. iii. 524.
leiochelium, *Benth. & Hook. f. Gen.* iii. 524.—Cuba
miserum, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 218.—Mexic.
pulchellum, *Benth. & Hook. f. Gen.* iii. 524.—Mexic.
pygmæum, Benth. & Hook. f. l. c. = uniflorum.
serpens, *Benth. & Hook. f. l. c.*—Hab.!
Sophronitis, *Benth. & Hook. f. l. c.*—N. Granat.
uniflorum, *Heynh. Nom.* i. 880, nomen.—Mexic.

HORMINUM, [Tourn.] Linn. Coroll. Gen. 9 (1737) 750. *LABIATÆ*, Benth. & Hook. f. ii. 1198.
PASITA, Adans. Fam. ii. 169 (1763).
pyrenaicum, *Linn. Sp. Pl.* 596.—Europ. austr.
virginicum, Linn. l. c. = Salvia lyrata.

HORMINUM, Mill. Gard. Dict. ed. VIII. (1768) = **Salvia**, Tourn. (Labiat.).
caulescens, Orteg. Hort. Matr. Dec. 68 = Lepechinia spicata.
clinopodiifolium, Pers. Syn. ii. 132 = Mentha dahurica.
coccineum, Moench, Meth. 377 = S. coccinea.
coloratum, Moench, l. c. = S. Horminum.
disermas, Moench, l. c. = S. disermas.
foliosum, Burm. f. Fl. Cap. Prod. 16.—Afr. austr.
frutescens, Gmelenne. Reise, i. 260.—Reg. Cauc.
lyratum, Mill. Gard. Dict. ed. VIII. n. 2 = S. lyrata.
napifolium, Mill. l. c. n. 4 = S. napifolium.
nilolicum, Moench, Meth. 377 = S. nilotica.
parviflorum, Moench, Meth. 378 = S. bryantina.
pyrenaicum, Georgi, Beschr. Russ. Reich. III. v. 1096 = Calamintha ambrosa.
sativum, Mill. Gard. Dict. ed. VIII. n. 5 = S. Horminum.
suaveolens, Moench, Meth. 377 = S. spinosa.
sylvestre, S. F. Gray, Nat. Arr. Brit. Pl. ii. 388 = S. Verbenaca.
verbenaceum, Mill. Gard. Dict. ed. VIII. n. 1 = S. Verbenaca.
verticillatum, Mill. l. c. n. 3 = S. verticillata.
viride, Moench, Meth. 377 = S. viridis.

HORMOCARPUS, Spreng. Gen. ii. 594 (1831) = **Ormocarpum**, Beauv. (Legumin.).

HORMOGYNE, A. DC. in DC. Prod. viii. 176 (1844). *SAPOTACEÆ*, Benth. & Hook. f. ii. 656.
ostiniflis, *A. DC. l. c.*—Austral.

HORMOSCIADIUM, Endl. Gen. Suppl. v. 16 (1850) = **Ormosetadium**, Boiss. (Umbellif.).

HORNEA, Baker, Fl. Maurit. 89 (1877). *SAPINDACEÆ*.
mauritiana, *Baker, l. c.*—Ins. Maurit.

HORNEMANNIA, Benth. in DC. Prod. x. 498 (1846) = **Blythorpia**, Link. (Scrophular.)
pinnata, Benth. l. c. = S. pinnata.

HORNEMANNIA, Link & Otto, Ic. Pl. Sel. 9. t. 3 (1820 ?) = **Vandellia**, Linn. (Scrophular.).
ovata, Link & Otto, l. c. = V. crustacea.
prostrata, Jacq. f. Eclog. Pl. t. 150 = V. hirsuta.

HORNEMANNIA, Vahl, in Skrivt. Nat. Selsk. Kiobenh. vi. (1810) 120. *VACCINIACEAE*, Benth. & Hook. f. ii. 567.
ANDREUSIA, Dunal, in DC. Prod. vii. 560 (1839).
PEVREIRIA, Rich. ex Dunal, in DC. l. c. in syn. (1839).
SYMPHYSIA, Presl, Epist. ad Jacq. (1837).
TAUSCHIA, Preisaler, in Flora, xi. (1828) 48.
martinicensis, *Hook. f. in Benth. & Hook. f. Gen.* ii. 567.—Ins. Martinic.
racemosa, *Vahl, in Skrivt. Nat. Selsk Kiobenh.* vi. (1810) 121.—Ins Carib.'
smilacifolia, *Hook. f. in Benth. & Hook. f. Gen.* ii. 567 —Ins. Dominic.

HORNEMANNIA, Willd. Enum. Hort. Berol. 653 (1809) = **Mauss**, Lour. (Scrophular.).
bicolor, Willd. l. c. = M. rugosus.
goodeniaefolia, Hoffmgg. ex Steud. Nom. ed. II. i. 776 = M. rugosus.
viscosa, Willd. Enum. Hort. Berol. 653 = Vandellia hirsuta.

HORNERA, Jungh. in Hoev. & De Vriese, Tijdschr. vil. (1840) 314. Gen. Gub. *THYMELEARUM.*—Cf. Benth. & Hook. f. Gen. iii. 188.
glomerata, *Jungh. l. c* 316.—Japon.
umbellata, *Jungh. l. c.* 314.—Japon.

HORNERA, Neck. Elem. iii. 43 (1790) = **Mucuna**, Adans. (Legumin.).

HORNSCHUCHIA, Blume, Cat. Gew. Buitenz. 15 (1823) = **Oratoxylon**, Blume (Hypericin.).
hypericina, Blume, l. c. 16 = C. Hornschuchii.

HORNSCHUCHIA, Nees, in Flora, iv. (1821) 302; et in Denkschr. Bot. Ges. Regensb. ii. (1822) 160. *ANONACEAE*, Benth. & Hook. f. ii. 603 (Cf. i. 392).
Bryotrophe, *Nees, ll. cc.* 302; 161. t. 11.—Bras.
Myrtillus, *Nees, ll. cc.* 302; 162. t. 12.—Bras.

HORNSCHUCHIIA, Spreng. Neue Entdeck. iii. 64 (1822) (Quid ?).
brasiliensis, Spreng. l. c.—Bras.

HORNSTEDIA, Juss. in Dict. Sc. Nat. ii. Suppl. 20 (1816) = seq.

HORNSTEDTIA, Retz. Obs. vi. 18 (1791) = **Amomum**, Linn. (Scitamin.).
Leonurus, Retz. l. c. = A. Leonurus.
scyphifera, Steud. Nom. ed. II. i. 776 = seq.
Scyphus, Retz. Obs. vi. 18 = A. scyphiferum.

HORNUNGIA, Bernh. in Flora, xxiii. (1840) 392 = **Gagea**, Salisb. (Liliac.).

HORNUNGIA, Reichb. Ic. Fl. Germ. xii. t. 6. f. 4190 (1837) = **Hutchinsia**, R. Br. (Crucif.).
petraea, Reichb. l. c. = Hutchinsia petraea.

HORREOLA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 = **Blatostemma**, Forst. (Urtic.).

HORSFIELDIA, Pers. Syn. ii. 635 (1807) = HORSFIELDIA, Willd. = **Myristica**, Linn.

HORSFIELDIA, Blume, ex DC. Prod. iv. 87 (1830). *ARALIACEAE*, Benth. & Hook. f. i. 937.
SCHUBERTIA, Blume, Bijdr. 864 (1826).
amlensis, *Blume, ex DC. Prod.* iv. 87.—Java.
horrida, Soem. in Journ. Bot. v. (1867) 237 = Fatsia horrida.

HORSFIELDIA, Willd. Sp. Pl. iv. 872 (1805) = **Myristica**, Linn.
odorata, Willd. l. c. = M. Horsfieldii.

HORSTIA, Heist. Syst. 16 (1748) (Quid ?).

HORTA, Thunb. ex Steud. Nom. ed. II. i. 776 (1840) = **HOSTA**, Tratt. = **Funkia**, Spreng. (Liliac.).

HORTA, Vell. Fl. Flum. 48; i. t. 124 (1825) = **Clavija**, Ruiz & Pav. (Myrsin.).
spinosa, Vell. ll. cc. = C. ornata.

HORTENSIA, Comm. ex Juss. Gen. 214 (1789) = **Hydrangea**, Linn. (Saxifrag.).
japonica, J. F. Gmel. Syst. 722 = Hydrangea hortensia.
mutabilis, Schneev. Ic. Pl. Rar. 36 = Hydrangea hortensia.
opuloides, Lam. Encyc. iii. 136 = Hydrangea hortensia.
rosea, Desf. Tabl. ed. 1. 115 = Hydrangea hortensia.
speciosa, Pers. Syn. i. 505 = Hydrangea hortensia.

HORTIA, Vand. Fl. Lusit. 14 (1788); et in Roem. Script. 88 (1796). *RUTACEAE*, Benth. & Hook. f. i. 351.
arbores, *Engl. in Mart. Fl. Bras.* xii. ii. 183.—Bras.
brasiliana, *Vand. ex DC. Prod.* i. 732.—Bras.
coccinea, *Benth. ex Engl. in Mart. Fl. Bras.* xii. ii. 183.—Bras.
longifolia, *Benth. ex Engl. l. c.* 184.—Bras.

HORTONIA, Wight, ex Arn. in Mag. Zool. & Bot. ii (1838) 645. *MONIMIACEAE*, Benth. & Hook. f. iii. 146.
acuminata, Wight, Ic t. 1998 = floribunda.
floribunda, *Wight, ex Arn. in Mag. Zool & Bot.* ii. (1838) 645.—Zeylan.
ovalifolia, Wight, Ic. t. 1998 = floribunda.

HORTSMANIA, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Strp. Surin. Sel.) 167 = **Condylocarpon**, Desf. (Apocyn.).
ciliata, Benth. ex Muell. Arg. in Mart. Fl. Bras. vi. i. 66.—C. ciliatum.
myrtifolia, Miq. Strp. Surin. Sel. 168 = C. myrtifolium.
pubiflora, Benth. ex Muell. Arg. in Mart. Fl. Bras. vi. i. 67 = C. pubiflorum.

HORTSMANNIA, Pfeiff. Nom. i. 1670 (1874) = praec.

HOSACKIA, Dougl. ex Benth. in Bot. Reg. t 1257 (1829). *LEGUMINOSAE*, Benth. & Hook. f. i. 491.
ACMISPON, Rafin. Atl. Journ. 144 (1832).
ANISOLOTUS, Bernh. Sem. Hort. Erf. (1837); Linnaea, xii. (1838) Litt. 75.
DESMOLOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 324 (1840)
FLUNDULA, Rafin. Fl. Tellur. ii. 96 (1836).
SYRMATIUM, Vog. in Linnaea, x. (1836) 590.
angustifolia, *G. Don, Gen. Syst.* ii. 200.—Am. bor. occ.
argentea, Kellogg, in Proc. Calif. Acad. iii. (1863) 38 = argophylla.
argophylla, *A. Gray, in Mem. Acad. Am.* N. S. v. (1855) 316.—Calif.
argyraea, *Greene, in Bull Calif. Acad.* i. iv. (1885) 184.—Calif.
balsamifera, Kellogg, in Proc. Calif. Acad. ii. (1863) 136 = stipularis.
bicolor, *Dougl. ex Benth. in Lindl. Bot. Reg.* t. 1257. —Am. bor. occ.
brachycarpa, *Benth. Pl. Hartw.* 306.—Calif.
crassifolia, *Benth. in Trans. Linn. Soc.* xvii. (1837) 365. —Am. bor. occ.
crassifolia, Nutt. ex Torr. & Gray, Fl N. Am. i. 325 —Calif.
cytisoides, *Benth. in Trans. Linn. Soc.* xvii. (1837) 366.—Calif.
decumbens, *Benth. l. c.*—Am. bor. occ.
distichs, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 186.—Calif.
elata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 326 = Purshiana.
flexuosa, *Greene, in Bull. Calif. Acad.* i. iii. (1885) 82. —Calif.
floribunda, Nutt. ex Torr. & Gray, Fl. N. Am. i. 327 = Purshiana.
glabra, *Torr. in Bot. U. St. Expl. Exped.* ii. 274.—Am. bor. occ.
gracilis, *Benth. in Trans. Linn. Soc.* xvii. (1837) 365. —Am. bor. occ.
grandiflora, *Benth. l. c.* 366.—Calif.
Heermanni, C. L. Anders. Pl. Nevada, 119 = decumbens.
Heermanni, *Durand & Hilg. in Journ. Acad. Sc. Philad.* Ser. II. iii. (1855) 39.—Am. bor. occ.
incana, *Torr. in Pacif. Rail. Rep.* iv. 79.—Calif.
indica, K. Grah. in Wall. Cat. n. 5940 = Rothia trifoliata.
juncea, *Benth. in Trans. Linn. Soc.* xvii. (1837) 366.—Calif.
latiperulata, Durand & Hilg. in Pacif. Rail. Rep. v. (1855) 6 = oblongifolia.
Macraei, *Benth. ex A. Gray, Bot. U. St. Expl. Exped.* i. 400.—Chili.
macrantha, *Greene, in Bull. Calif. Acad.* i. iii. (1885) 81.—Calif.
macrophylla, Kellogg, in Proc. Calif. Acad. ii. (1863) 138 = stipularis.
maritima, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 327. —Calif.
maritima, Torr. Bot. Mex. Bound. 58 = strigosa.
mexicana, Benth. in Trans. Linn. Soc. xvii. (1837) 365 = angustifolia.
micrantha, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 324. —Calif.
microphylla, Nutt. ex Torr. & Gray, l. c. 326 = strigosa.
mollis, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 185. —N. Mexic.
mollis, Nutt. ex Torr. & Gray, Fl. N. Am. i. 327 = Purshiana.
nana, *S. Wats. in Proc. Am. Acad.* xx. (1885) 359.—Mexic.
nudiflora, Nutt. ex Torr. & Gray Fl N. Am. i. 326 = strigosa.
oblongifolia, *Benth. Pl. Hartw.* 305.—Calif.
ochroleuca, Nutt. ex Torr. & Gray, Fl. N. Am. i. 328 = grandiflora.
Ornithopus, Greene, in Bull. Calif. Acad. i. iv. (1885) 185 = argophylla ?
parviflora, *Benth. in Lindl. Bot. Reg.* sub t. 1257.—Am. bor. occ.
pilosa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 327 = Purshiana.
platycarpa, Nutt. ex Torr. & Gray, l. c. 323 = crassifolia.
procumbens, *Greene, in Bull. Calif. Acad.* i. iii. (1885) 82.—Am. bor. occ.
prostrata, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 325. —Calif.
puberula, *Benth. Pl. Hartw* 305.—N. Mexic.
puberula, A. Gray, Pl. Wright. ii. 42 = rigida.
Purshiana, *Benth. in Lindl. Bot. Reg.* sub t. 1257. —Am. bor. occ.
repens, *G. Don, Gen. Syst.* ii. 200.—Mexic.
rigida, *Benth. Pl. Hartw.* 305.—Am. bor. occ.
rubella, Nutt. ex Torr. & Gray, Fl. N. Am. i. 326 = strigosa.
scoparia, Torr. & Gray, l. c. 325 = glabra.
sericea, *Benth. in Trans. Linn. Soc.* xvii. (1837) 367. —Calif.
stipularis, *Benth. l. c.* 365.—Calif.
stolonifera, Lindl. Bot. Reg. t. 1977 = crassifolia.
strigosa, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 326.—Calif.
subpinnata, *G. Don, in Sweet, Hort. Brit.* ed. III. 165; *Torr. & Gray, Fl. N. Am.* i. 326.—Am. bor. occ.; Chili.
tomentosa, *Hook. & Arn. Bot. Beech. Voy.* 137.—Calif.
Torreyi, *A. Gray, in Proc. Am. Acad.* viii. (1878) 626.—Am. bor.
umbellulata, Hook. Fl. Bor. Am. i. 135 = Purshiana.
Veatchii, *Greene, in Bull. Calif. Acad.* i. iii. (1885) 83.—Calif.
Wrangeliana, G. Don, in Sweet, Hort. Brit. ed. III. 165; Torr. & Gray, Fl. N. Am. i. 326 = subpinnata.
Wrightii, *A. Gray, Pl. Wright.* ii. 42.—N. Mexic.

HOSAKIA, Steud. Nom. ed. II. i. 776 (1840) = praec.

HOSANGIA, Neck. Elem. i. 225 (1790) = **Maiota**, Aubl. (Melastomac.).

HOSEA, Dennst. Schlssl. Hort. Malab. 31 (1818) = (Quid ?).
malabarica, Dennst. l. c.—Ind. or.

HOSLUNDA, Roem. & Schult. Syst. i. 57, 207 (1817) = seq.

HOSLUNDIA, Vahl, Enum. ii. 212 (1805). *LABIATAE*, Benth. & Hook. f. ii. 1174.
HAASLUNDIA, Schum. & Thonn. Beskr. Guin. Pl. 15 (1827).
decumbens, *Benth. in DC. Prod.* xii. 54.—Afr. trop.
opposita, *Vahl, Enum.* i. 212.—Afr. trop.
oppositifolia, Benv. Fl. Owar. i. 58 = opposita.
verticillata, *Vahl, Enum.* i. 218.—Afr. trop.

HOSTA, Jacq. Hort. Schoenb. i. 60, 114 (1797) = Cornutia, Linn. (Verbenac.).
cærulea, Jacq. l. c. = C. pyramidata.
coerulea, Walp. Rep. iv. 80 = C. pyramidata.
grandiflora, A. Dietr. Sp. Pl. i. 252 = C. grandifolia.
grandifolia, Cham. & Schlecht. in Linnæa, v. (1830) 97 = C. grandifolia.
latifolia, H. B. & K. Nov. Gen. et Sp. ii. 248 = C. pyramidata.
longifolia, H. B. & K. l. c. 247 = C. longifolia.
odorata, Poepp. & Endl. l. c. iii. 68. t. 269 = C. longifolia.
pyramidata, A. Dietr. Sp. Pl. i. 253 = C. pyramidata.

HOSTA, Tratt. Arch. Gewaechs. i. (1812) 55 = Funkia, Spreng. (Lillac.).
cornuta, Tratt. l. c. in obs. = F. ovata.
japonica, Tratt. l. c. t. 89 = F. subcordata.
lancifolia, Tratt. l. c. in obs. = F. lancifolia.
plantaginea, Aschers. Fl. Brandenb. 728 = F. subcordata.

HOSTA, Vell. ex Pfeiff. Nom. i. 1670 (1874) = HORTA, Vell. = Clavija, Ruiz & Pav. (Myrsin.).

HOSTANA, Pers. Syn. i 148 (1805) = HOSTA, Jacq. = Cornutia, Linn. (Verbenac.).
cornuta, Pers. l. c. = C. pyramidata.

HOSTEA, Willd. Sp. Pl. i. 1374 (1797) = Mateles, Aubl. (Asclepiad.).
viridiflora, Willd. l. c. = M. latifolia, palustris.

HOSTIA, Moench, Meth. Suppl. 221 (1802) = Crepis, Linn. (Compos.).
fœtida, Moench, l. c. = C. foetida.
rubra, Cass. in Dict. Sc. Nat. xxi. 448 = C. rubra.

HOSTMANNIA, Planch. in Hook. Ic. Pl. t. 709 (1848) = Hivasia, DC. (Ochnac.).
ochnoides, Planch. l. c. = E. Hostmannia.

HOSTMANNIA, Stend. ex Nasd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 25 = Cornutia, DC. (Melastomac.).
lythrarioides, Stend. ex Nasd. l. c. 27 = C. lythrarioides.

HOTRIA, C. Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 216. t. 11 = Astilbe, Buch.-Ham. (Saxifrag.).
horbata, Hook. Bot. Mag. t. 3891 = A. japonica.
biternata, C. Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 218, in obs. = Tiarella biternata.
chinensis, Maxim. Prim. Fl. Amur. 120 = A. chinensis.
japonica, C. Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 217 = A. japonica.
Thunbergii, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1843) 191 = A. Thunbergii.

HOTTONIA, Boerh. ex Linn. Syst. ed. I (1735). *PRIMULACEAE*, Benth. & Hook. f. ii. 681.
BRYOGLADIUM, Dulac, Fl. Hautes-Pyr. 428 (1867).
indica, Linn. Syst. ed. X. 919 = Limnophila gratioloides.
inflata, *Ell. Sketch*, i. 281.—Am. bor.
litteralis, Lour. Fl. Cochinch. 105 = Erythraea cochinchinensis.
Millefolium, Gilib. Fl. Lituan. i. 86 = palustris.
palustris, *Linn. Sp. Pl.* 146.—Europ.
palustris, Pursh, Fl. Am. Sept. i. 136 = inflata.
serrata, Willd. Sp. Pl. i. 814 = Hydrilla *ovalifolia*!
sessiliflora, Vahl, Symb. Bot. ii. 36 = Limnophila sessiliflora.

HOTTONIA, Burm. Thes. Zeyl. 121. t. 55. f. 1 (1737) = Limnophila, R. Br. (Scrophular.).

HOTTUYNIA, Cram. Disp. 141 (1808) = Houttuynia, Thunb. (Piperac.).

HOULLETIA, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 37. *ORCHIDEAE*, Benth. & Hook. f. iii. 550.
Brockhurstiana, *Lindl. Bot. Reg.* (1841) *Misc.* 47.—Bras.
chrysantha, Linden & André, Illustr. Hortic. xviii. (1871) 168. t. 71 = Wallisii.
Landbergi, *Linden & Reichb. f. in Regel, Gartenfl.* (1855) 2.—Bras.
Lowiana, *Reichb. f. in Gard. Chron.* (1874) ii. 484.—N. Granat.
odoratissima, *Linden, ex Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 172.—N. Granat.
picta, *Linden & Reichb. f. in Regel, Gartenfl.* (1855) 2.—N. Granat.
stapeliæflora, Brongn. ex Lem. in Illustr. Hortic. (1859) t. 612, in syn. = seq.
stapeliiflora, *Brongn. ex Reichb. f. in Regel, Gartenfl.* (1855) 2.—Bras.!
tigrina, *Linden, ex Lindl. in Paxt. Flow. Gard.* iii. (1852–53) 171, 172.—N. Granat.
vittata, Lindl. Bot. Reg. (1841), Misc. 47 = Polycynis vittata.
Wallisii, *Linden & Reichb. f. in Gard. Chron.* (1869) 611.—N. Granat.

HOUMIRI, Aubl. Pl. Gui. i. 564 (1775) = Humiria, Jaume St. Hil. (1805).
balsamifera, Aubl. l. c. = Humiria balsamifera.
guianensis, Baill. Hist. Pl. v. 59.—Afr. trop.
guianensis, Baill. l. c. 54 = Humiria guianensis.

HOUMIRIA, Juss. Gen. 425 (1789) = prsec.

HOUNEA, Baill. in Bull. Soc. Linn. Par. i. (1881) 301. *PASSIFLORACEAE*.
madagascariensis, *Baill. l. c.*—Madag.

HOUSTONIA, Gronov. ex Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth. & Hook. f. ii. 60.
CRUSEMA, Rafin. ex Stend. Nom. ed. II. i. 776, in syn. (1840).
PAINTOS, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 267. vi. 81.
POIRETIA, J. F. Gmel. Syst. 368 (1791).
acerosa, *Benth. & Hook. f. Gen.* ii. 60; *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 208.—Mexic.
angustifolia, *Michx. Fl. Bor. Am.* i. 85.—Am. bor.
angustifolia, Pursh, Fl. Am. Sept. i. 106 = purpurea.
bouvardioides, *Benth. & Hook. f. Gen.* ii. 60.—Mexic.
bryoides, Willd. ex Roem. & Schult. Syst. iii. 527 = Hedyotis hypnoides.
caerulea, *Linn. Sp. Pl.* 105.—Am. bor.
canadensis, *Willd. ex Roem. & Schult. Syst.* iii. 527. *chlorantha*, Bertol. Elench. 8 = Bouvardia chlorantha.
ciliolata, Torr. Fl. N. Am. i. 178 = purpurea.
coerulea, Andr. Bot. Rep. t. 106 = Bouvardia triphylla.
coerulea, Auct. vide caerulea.
crocea, *Schnabl. Pl. Nov. Hort. Tubing.* (1835) 20; *ex Steud. Nom.* ed. II. i. 777.—Hab.?
fasciculata, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 208.—Mexic.
fruticosa, Rafin. ex A. Gray, Syn. Fl. Am. ii. i. 26 = angustifolia.
fruticosa, Willd. ex Roem. & Schult. Syst. iii. 527 = Declieuxia chiococcoides.
geniculata, *Rafin. New Fl. Am.* iv. 102.—Am. bor.
grandiflora, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 225 = caerulea.
heterophylla, *Rafin. l. c.* 227.—Am. bor.
hanmifusa, *A. Gray, in Proc. Am. Acad.* iv. (1858) 314.—Texas.
humifusa, Hensl. Biol. Centr. Am. Bot. ii. 30 = Wrightii.
latifolia, *Willd. ex Roem. & Schult. Syst.* iii. 527.—Am. bor.
Linnæi, Michx. Fl. Bor. Am. i. 85 = caerulea, purpurea.
longifora, *A. Gray, in Proc. Am. Acad.* iv. (1860) 314.—Mexic.
longifolia, Cavrin. Fruct. i. 226. t. 49. f. 3 = purpurea.
longipes, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 97.—N. Mexic.

HOUSTONIA:—
macrocephala, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 40 = purpurea.
microphylla, Willd. ex Roem. & Schult. Syst. iii. 527 = Anotis serpens.
minima, *Beck, in Am. Journ. Sc.* x. (1826) 262.—Am. bor. occ.
oblongifolia, *Rafin. in Ann. Gén. Sc. Phys.* v. (1820) 226.—Am. bor.
obtusifolia, *Rafin. l. c.*—Am. bor.
Palmeri, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 209.—Mexic.
patens, *Ell. Sketch*, i. 191.—Am. bor.
pubescens, Rafin. in Med. Repos. N. York, v. (1808) 361 = purpurea.
purpurea, *Linn. Sp. Pl.* 105.—Am. bor.
pusilla, J. F. Gmel. Syst. 236 = caerulea!
reticulata, *Rafin. New Fl. Am.* iv. 102.—Am. bor.
rotundifolia, *Michx. Fl. Bor. Am.* i. 85.—Am. bor.
rubra, *Cav. Ic.* v. 46. t. 474. f. 1.—Mexic.
rupestris, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 226 = angustifolia.
serpyllifolia, R. Grah. in Hook. Bot. Mag. t. 2832 = caerulea!
serpyllifolia, *Michx. Fl. Bor. Am.* i. 85.—Am. bor.
subviscosa, *A. Gray, in Proc. Am. Acad.* iv. (1858–59) 314.—Texas.
tenella, Pursh, Fl. Am. Sept. i. 106 = serpyllifolia.
tenuifolia, Nutt. Gen. Am. i. 127 = purpurea.
trifora, *A. Gray, in Proc. Am. Acad.* iv. (1858–59) 314.—Mexic.
umbellata, Walp. Rep. i. 496, in syn. = purpurea.
variana, Michx. Fl. Bor. Am. i. 86 = purpurea.
Wrightii, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 209.—Am. bor. occ.

HOUTOUVYNIA, Pers. Syst. 556 (1797) = Houttuynia, Thunb. (Piperac.).

HOUTTEA, Decne. in Rev. Hortic. Sér. III. iii. (1848) 469. *GESNERACEAE*, Benth. & Hook. f. ii. 1002.
VANHOUTTEA, Lem. in Bull. Soc. Hort. Orléans, i. (1846) 846.
Gardneri, *Decne. in Rev. Hortic. Sér.* III. ii. (1848) 469.—Bras.
Leptopus, Hanst. in Mart. Fl. Bras. viii. 393 = Gardneri.
pardina, *Decne. in Rev. Hortic. Sér.* III. ii. (1848) 469.—Bras.
salviifolia, *Hanst. in Mart. Fl. Bras.* viii. 395.—Bras.

HOUTTEA, Heynh. Nom. ii. 306 (1846) = Achimenes, P. Br. (Gesnerac.).
tubiflora, Heynh. l. c. (= *Dolichodeira tubiflora*).—Bras.

HOUTTINIA, Neck. Elem. iii. 291 (1790) (Hoottinia) = Calla, Linn. (Aroid.).

HOUTTOUVNIA, Batsch, Tab. 159 (1802) = seq.

HOUTTUYNIA, Thunb. Fl. Jap. 12 (1784). *PIPERACEAE*, Benth. & Hook. f. iii. 126.
ANEMIA, Nutt. in Ann. Nat. Hist. Ser. I. i. (1838) 136.
ANEMOPSIS, [Hook.] in Ann. Nat. Hist. Ser. I. i. (1838) 136, in note; Hook. & Arn. Bot. Beech. Voy. 390. t. 93.
POLYPARA, Lour. Fl. Cochinch. 61 (1790).
Belandieri, Benth. & Hook. f. Gen. iii. 128 = californica.
californica, *Benth. & Hook. f. ex Bot. Calif.* ii. 483.—Am. bor. occ.
cordata, *Thunb. Fl. Jap.* 234. t. 26.—Reg. Himal.; China; Japon.
fœtida, Hort. ex Loud. Hort. Brit. 87 = cordata!

HOUTTUYNIA, Hoott. Handleid. xii. 448. t. 55 (1778) = Acidanthera, Hochst. (Irid.).
capensis, Houtt. l. c.—Afr. austr.

HOUTUVNIA, Thunb. Fl. Jap. 231. t. 26 (1784) = prsec.

HOUTUYNIA, Thunb. in Vet. Acad. Handb. Stockh. iv. (1788) 149. t. 5 = Houttuynia, Thunb. (Piperac.).

HOVEA, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 275 (1812). *LEGUMINOSAE*, Benth. & Hook. f. i. 474.
PLAGIOLOBIUM, Sweet, Fl. Austral. t. 3 (1827).
PLATYCHILUS, Delaun. Herb. Amat. t. 187 (1815).
POIRETIA, Sm. in Trans. Linn. Soc. ix. (1808) 304.

HOVEA:—
scanthoclada, *F. Muell. Fragm.* iv. 15.—Austral.
scottiolis, *A. Cunn. ex G. Don, Gen. Syst.* ii. 196.—Austral.
apiculata, A. Cunn. ex G. Don, l. c. = longifolia.
Beckeri, F. Muell. in Linnæa, xxv. (1852) 891 = longifolia.
Celsi, *Bonpl. Jard. Malm.* t. 51.—Austral.
chorisemæfolia, *DC. Prod.* ii. 116.—Austral.
crispa, Lindl. Bot. Reg. (1839) Misc. 19 = trisperma.
elliptica, DC. Prod. ii. 115 = Celsi.
elliptica, Meisn. Gen. Fl. Preiss. i. 79 = trisperma.
heterophylla, *A. Cunn. ex Hook. f. Fl. Tasm.* i. 95. t. 16.—Austral.
ilicifolia, A. Cunn. ex Lindl. Bot. Reg. (1844) t. 58 = chorisemæfolia.
lanceolata, Sims, Bot. Mag. t. 1624 = longifolia.
lanigera, Lodd. ex Steud. Nom. ed. II. i. 777 = longifolia.
latifolia, Lodd. Bot. Cab. t. 30 = Celsi.
leiocarpa, Benth. in Mitch. Journ. Trop. Austral. 289 = longipes.
linearis, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 275.—Austral.
longifolia, [*R. Br. in*] *Ait. l. c.*—Austral.
longipes, *Benth. in Enum. Pl. Hueg.* 37.—Austral.
Mangliæi, Lindl. Bot. Reg. (1838) t. 62 = trisperma.
mucronata, A. Cunn. ex G. Don, Gen. Syst. ii. 126 = longifolia.
pungens, *Benth. in Enum. Pl. Hueg.* 37.—Austral.
purpurea, Sweet, Fl. Austral. t. 13 = longifolia.
racemulosa, Benth. in Lindl. Bot. Reg. (1842) Misc. 39 = longifolia.
ramulosa, A. Cunn. ex Lindl. l. c. (1843) sub t. 4 = longifolia.
rosmarinifolia, A. Cunn. in Field, N. S. Wales, 348 = longifolia.
spheroclena, Pxst. Mag. Bot. x. (1843) 108 = trisperma.
stricta, *Meisn. in Lehm. Pl. Preiss.* i. 79.—Austral.
trisperma, *Benth. in Enum. Pl. Hueg.* 37.—Austral.
ulicina, Meisn. in Bot. Zeit. xiii. (1855) 30 = pungens.
villosa, Lindl. Bot. Reg. t. 1513 = longifolia.

HOVENIA, Thunb. Nov. Gen. 7 (1781). *RHAMNEÆ*, Benth. & Hook. f. i. 378.
acerba, Lindl. in Bot. Reg. t. 501 = dulcis.
dulcis, *Thunb. Nov. Gen.* 8.—Reg. Himal.; China; Japon.
inæqualis, DC. Prod. ii. 40 = dulcis.
pubescens, Sweet, Hort. Brit. ed. I. 91 = dulcis.

HOVERDENIA, Nees, in DC. Prod. xi. 380 (1847). *ACANTHACEÆ*, Benth. & Hook. f. ii. 1116.
speciosa, *Nees, l. c.* 381.—Mexic.

HOWARDIA, Klotzsch, in Monatsb. Acad. Berl. (1859) 807 = **Aristolochia**, Linn.
angusicida, *barbata*, *bilabiata*, *bilobata*, *Bridgesii*, *caudata*, *Chamisonis*, *coriágera*, *costaricensis*, *cynanchifolia*, *deltoidea*, *Ehrenbergiana*, *emarginata*, *orientis*, *fimbriata*, *fortida*, *fragrantissima*, *gigantea*, *glaucescens*, *grandiflora*, *hians*, *inflata*, *macroura*, *maxima*, *obtusata*, *pandurata*, *peltata*, *pilosa*, *pubescens*, *punctata*, *Raja*, *reniformis*, *ringens*, *Ruiziana*, *ramiflora*, *sessilifolia*, *smilacina*, *tamnifolia*, *triangularis*, *trilobata*, *truncata*, *veraguensis*, Klotzsch, l. c. 807–822 = *hemonyma omnia* Aristolochiæ.
Benthamii, Klotzsch, l. c. 820 = A. gibbosa.
brasiliensis, Klotzsch, l. c. 808 = A. cymbifera.
brevifolia, Klotzsch, l. c. 816 = A. angustifolia.
cymbifera, Klotzsch, l. c. 809 = A. galeata.
fluminensis, Klotzsch, l. c. 812 = A. odora.
fusiformis, Klotzsch, l. c. 809 = A. grandiflora.
galeata, Klotzsch, l. c. 809 = A. brasiliensis.
gymnistoma, Klotzsch, l. c. 819 = A. maxima.
Galeottii, Klotzsch, l. c. 821 = A. maxima.
Hoffmanni, Klotzsch, l. c. = A. maxima.
Lansbergii, Klotzsch, l. c. 813 = A. sempervirens.
longifolia, Klotzsch, l. c. 816 = A. angustifolia.
macrodonia, Klotzsch, l. c. 815 = A. glandulosa.
Schomburghii, Klotzsch, l. c. 818 = A. dictyantha.
Selloviana, Klotzsch, l. c. 816 = A. triangularis.
serpens, Klotzsch, l. c. 817 = A. chilensis.

HOWARDIA :—
surinamensis, Klotzsch, in Monatsb. Acad. Berl. (1859) 818 = Aristolochia macrura.
trifida, Klotzsch, l. c. 817 = A. trilobata.
Warszewiczii, Klotzsch, ex Duch. in DC. Prod. xvi. 476 = A. pandurata.

HOWARDIA, Weidl in Ann. Sc. Nat. Sér. IV. i. (1854) 66. t. 10 = **Pogonopus**, Klotzsch (Kubiac.).
cavacaensis, Weidl. l. c. 71 = *P. exsertus*.
febrifuga, Weidl. l. c. 67.— Bolivia.
grandiflora, Wedd. l. c. 70.—N. Granat.
Richardii, Wedd. l. c. 69.—Bras.

HOWEA, Becc. Malesia, i. (1877) 41, 66. *PALMAE*, Benth. & Hook. f. iii. 904.
Gribbachia, H. Wendl. & Drude, in Linnæa, xxxix. (1875) 177.
Belmoreana, *Becc. Malesia*, i. (1877) 66.—Ins. Dom. Howe.
Forsteriana, *Becc. l. c.*—Ins. Dom. Howe.

HOWELLIA, A. Gray, in Proc. Am. Acad. xv. (1880) 48. *CAMPANULACEÆ*.
aquatilis, *A. Gray, l. c.*—Am. bor. occ.

HOWIEA, Becc. Malesia, i. (1877) 66 = **Howea**, Becc. (Palm.).

HOWITTIA, F. Muell. in Trans. Vict. Inst. i. (1855) 116; in Hook. Kew Journ. viii. (1856) 9. *MALVACEÆ*, Benth. & Hook. f. i. 203.
trilocularis, *F. Muell. ll. cc.*—Austral.

HOYA, R. Br. in Mem. Wern. Soc. i. (1809) 26. *ASCLEPIADEÆ*, Benth. & Hook. f. ii. 776.
Acanthostemma, Blume, Mus. Bot. Lugd. Bat. i. 9. t. 10 (1849).
Cathetostemma, Blume, l. c. 59. t. 18 (1849).
Centrostemma, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 371. t. 12.
Cyatocekas, Benn. Pl. Jav. Rar. 90. t. 21 (1838).
Otostemma, Blume, Mus. Bot. Lugd. Bat. i. 59. t. 11 (1849).
Plocostemma, Blume, Rumphia, iv. 30 (1848); et l. c. t. 14 (1849).
Pyrostelma, Wight, Contrib. 39 (1834).
Scholelia, Jacq. f. Eclog. Pl. i. 5. t. 2 (1811).
Spirlingia, Vahl, in Skrivt. Nat. Selsk. Kiobenh. vi. (1810). 118.
Telisma, Van Hass. ex Miq. Fl. Ind. Bat. ii. 928 (1857).
acuminata, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 53.—Reg. Himal.
acuta, Haw. Rev. Pl. Succ. 4 = parasitica.
africana, Decne. in DC. Prod. viii. 689 (= *Pterygocarpus abyssinicus*).—Afr. trop.
alba, Koxb. ex Wall. Alg. Med.-Pharm. Fl. iii. 1084 = Rumphii!
albens, Millers. ex Steud. Nom. ed. II. i. 777 = parasitica.
albiflora, *Zipp. ex Blume, Rumphia*, iv. 88.—Java.
angustifolia, *Traill, in Trans. Hort. Soc. vii.* (1830) 39 (sp. dub.).—China.
apiculata, *Schef. in Ann. Jard. Buitenz.* i. (1876) 37.—N. Guinea.
Ariadna, *Decne. in DC. Prod.* viii. 685.—Ins. Amboin.
Arnottiana, *Wight, Contrib.* 37.—Reg. Himal.
australis, *R. Br. ex Traill, in Trans. Hort. Soc.* vii. (1830) 38.—Austral.
barbata, Spreng. Syst. i. 843 = Tylophora barbata.
Barnesii, Horne, ex Baker, in Journ. Linn. Soc. xx. (1883) 369 = australis.
bella, *Hook. Bot. Mag.* t. 4402.—Burma.
bicarinata, A. Gray, in Proc. Am. Acad. v. (1861–62) 335 = australis.
Billardieri, Decne. in DC. Prod. viii. 686 = australis.
Brunoniana, Wight, Contrib. 37 = coriacea.
burmensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 141.—Ins. Burn.
campanulata, Blume, Bijdr. 1064 = Physostelma Wallichii.
carnosa, *R. Br. in Mem. Wern. Soc.* i. (1809) 27.—As. et Austral. trop.
carnosa, R. Br. Prod. 460 = australis.
caudata, *Hook. f. Fl. Brit. Ind.* iv. 60.—Malacca.
chinensis, Traill, in Trans. Hort. Soc. vii. (1830) 37 = carnosa.

HOYA :—
ciliata, *Teijsm. & Binn. Cat. Hort. Bog.* 129.—Ins. Baru.
cinnamomifolia, *Hook. Bot. Mag.* t. 4347.—Java.
clandestina, *Blume, Rumphia*, iv. 92.—Java.
coccinea, Hort. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 310 = cinnamomifolia.
cochinchinensis, *Schult. Syst.* vi. 52.—Cochinch.
collyrioides, *Teijsm. & Binn. Cat. Hort. Bog.* 130, nomen.—Ins. Celebes.
coriacea, *Blume, Bijdr.* 1061.—Malaya.
coriacea, Lindl. Bot. Reg. (1839) t. 18 = multiflora.
coriacea, Zoll. ex Miq. Fl. Ind. Bat. ii. 519 = Zollingeriana.
Corona-Ariadnes, Blume, Rumphia, iv. 81 = H. Ariadna.
coronaria, *Blume, Bijdr.* 1063.—Malaya.
crassifolia, Haw. Suppl. Pl. Succ. 8 = carnosa.
crassipes, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) i. 261.—Java.
Cunningiana, *Decne. in DC. Prod.* viii. 686.—Ins. Philipp.
Dalrympliana, F. Muell. Rep. Burdek. Exped. 16 = australis.
densifolia, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 261 = Cumingiana.
diptera, *Sonn. in Bonplandia*, ix. (1861) 257; *Fl. Vit.* 168.—Ins. Fiji.
diversifolia, *Blume, Bijdr.* 1064.—Burma; Malaya.
Edeni, *King, ex Hook. f. Fl. Brit. Ind.* iv. 58.—Reg. Himal.
elegans, Kostel. Alg. Med.-Pharm. Fl. iii. 1084 = Rumphii!
elliptica, *Hook. f. Fl. Brit. Ind.* iv. 58.—Malacca.
excavata, *Tæjsm. & Binn. in Tijdschr. Nederl. Ind* xxv. (1863) 406.—Malaya.
Finlaysonii, *Wight, Contrib.* 38.—Singapore.
flexuosa, Spreng. Syst. i. 843 = Tylophora flexuosa.
fraterna, *Blume, Rumphia*, iv. 89.—Java.
fusca, *Wall. Pl. As. Rar.* i. 68.—Reg. Himal.
globulifera, *Blume, Mus. Bot. Lugd. Bat.* i. 44.—N. Guinea.
globulosa, *Hook. f. in Gard. Chron.* (1882) i. 732.—Reg. Himal.
goniocloides, *Regel, in Act. Hort. Petrop.* viii. (1884) 642.—Hab.?
graddifolora, *Blume, ex Decne. in DC. Prod.* viii. 685.—Java.
grandiflora, Spreng. Syst. i. 843 = Tylophora grandiflora.
Griffithiana, Decne. ex Hook. f. Fl. Brit. Ind. iv. 58 = acuminata.
Griffithii, *Hook. f. l. c.* 59.—Reg. Himal.
gymnanthera, *Wight, Contrib.* 37.—Ind. or.
Haseltii, *Miq. Fl. Ind. Bat.* ii. 924.—Java.
Hookeriana, Wight, Contrib. 37 = parasitica.
imbricata, *Decne. in DC. Prod.* viii. 687.—Ins. Philipp.
imperialis, *Lindl. Bot. Reg.* (1846) sub t. 68.—Malaya.
Keysii, *F. M. Bailey, in Proc. Roy. Soc. Queensl.* i. (1884) 87.—Austral.
Lacuna, Buch.-Ham. in Wall. Cat. n. 3169 = Dregea volubilis.
lacunosa, *Blume, Bijdr.* 1063.—Malaya.
lanceolata, Lindl. in Donn, Cat. Hort. Cantab. ed. XI. 92 = parasitica.
lanceolata, *Wall. ex D. Don, Prod. Fl. Nep.* 130.—Reg. Himal.
lasiantha, *Korth. ex Blume, Rumphia*, iv. 80.—Borneo.
latifolia, *G. Don, Gen. Syst.* iv. 127.—Malaya.
laurifolia, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 372.—Ins. Timor.
laurifolia, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 120 = carnosa.
linearis, *Wall. in Wight, Contrib.* 36.—Reg. Himal.
Lobbii, *Hook. f. Fl. Brit. Ind.* iv. 54.—Reg. Himal.
longifolia, *Wall. in Wight, Contrib.* 36.—Reg. Himal.
lutea, *Decne. in DC. Prod.* viii. 685.—Ins. Amboin.
lutea, Kostel. Alg. Med.-Pharm. Fl. iii. 1083 = coronaria!
macrophylla, *Blume, Bijdr.* 1062.—Java.
macrophylla, Wight, Contrib. 38 = latifolia.
Malayeri, *Hook. f. Fl. Brit. Ind.* iv. 60.—Malacca.
maxima, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 406.—Ins. Celebes.

HOYA:—
micrantha, *Hook. f. Fl. Brit. Ind.* iv. 55.—Burma.
Metzkei, Teijsm. & Binn. in Nederl. Kruidk. Arch. iii. (1854) 899 = carnosa.
multiflora, *Blume, Cat. Gew. Buitenz.* 49; *Bijdr.* 10 4. —Malaya.
Nicholsoniæ, *F. Muell. Fragm.* v. 159.—Austral.
nicobarica, *R. Br. ex Traill, in Trans. Hort. Soc.* vii. (1830) 38.—Ins. Nicobar.
nummularia, *Decne. ex Hook. f. Fl. Brit. Ind.* iv. 55.—Reg. Himal.
obcordata, *Hook. f. l. c.* 56.—Reg. Himal.
obcordata, *Teijsm. & Binn. Cat. Hort. Bog.* 285, nomen.—Siam.
oblanceolata, *Hook. f. Fl. Brit. Ind.* iv. 57.—Reg. Himal.
obovata, *Decne. in DC. Prod.* viii. 685.—Ins. Molucc.
obtusifolia, *Wight, Contrib.* 38.—Burma; Malaya.
odoratissima, *Linden, ex Regel, Cat. Hort. Absehov.* 78.—Hab.?
opposita, G. Don, Gen. Syst. iv. 138 = Rumphii.
orbiculata, Wall. ex Wight, Contrib. 36 = diversifolia.
ovalifolia, Wall. Cat. n. 8160 B = revoluta.
ovalifolia, *Wight & Arn. in Wight, Contrib.* 37.—Ind. or.
pallida, Dals. & Gibs. Bomb. Fl. 152 = Wightii.
pallida, Lindl. Bot. Reg. t. 951 = parasitica.
paniculata, Spreng. Syst. i. 843 = Tylophora paniculata.
parasitica, *Wall. ex Traill, in Trans. Hort. Soc.* vii. (1830) 38.—Reg. Himal.; Malaya.
parviflora, *Wight, Contrib.* 37.—Burma.
pauciflora, Wight, Ic. t. 1269 = parviflora.
Paxtoni, Nichols. Dict. Gard. ii. 155 = bella.
pendula, Wight & Arn. in Wight, Contrib. 36 = Wightii.
pendula, *Wight, Ic. t.* 474.—Ind. or.
picta, Hort. in Rev. Hortic. (1853) 277, partim = carnosa.
picta, *Miq. Fl. Ind. Bat.* ii. 524.—Java.
pilosa, *Sonn. in Bonplandia*, xix. (1861) 257; *Fl. Vit.* 168.—Ins. Fiji.
planiflora, Wall. ex Hook. f. Fl. Brit. Ind. iv. 45 = Tylophora asthmatica.
polyneura, *Hook. f. l. c.* 54.—Reg. Himal.
polystachya, Blume, Mus. Bot. Lugd. Bat. i. 45 = latifolia.
Pottsii, *Traill, in Trans. Hort. Soc.* vii. (1830) 25.—China.
Præstorii, *Miq. Fl. Ind. Bat.* ii. 526.—Sumatra.
pruinosa, *Miq. l. c.* 525.—N. Guinea.
pubera, *Blume, Bijdr.* 1065.—Java.
purpurascens, *Teijsm. & Binn. Cat. Hort. Bog.* 285, nomen.—Sumatra.
purpurea, *Blume, Rumphia*, iv. 30.—N. Guinea.
purpureo-fusca, *Hook. Bot. Mag.* t. 4520.—Java.
recurvifolia, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 150 = Gymnema recurvifolia.
retusa, *Dals. in Hook. Kew Journ.* iv. (1852) 294.—Ind. or.
revoluta, *Wight, ex Decne. in DC. Prod.* viii. 686.—Malaya.
rotundifolia, Siebold, Cat. Pl. Jap. (1840) = carnosa.
Rumphii, *Blume, Bijdr.* 1065.—Ins. Molucc.
ruscifolia, *Decne. in DC. Prod.* viii. 689.—Ins. Philipp.
samoënsis, *Sonn. Fl. Vit.* 168.—Ins. Samoa.
serpens, *Hook. f. Fl. Brit. Ind.* iv. 55.—Reg. Himal.
Shepherdi, Short, ex Hook. Bot. Mag. t. 5269 = longifolia.
speciosa, Decne. in DC. Prod. viii. 684 = imperialis?
suavеolens, Hort. Amst. ex Miq. Fl. Ind. Bat. ii. 525 = laanceae.
subquaterna, *Miq. l. c.*—Java.
subquintuplinervia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 141.—Siam.
tenuifolia, *Griff. ex Hook. f. Fl. Brit. Ind.* iv. 54.—Reg. Himal.
Teysmanniana, Miq. Fl. Ind. Bat. ii. 522 = obtusifolia.
Thomsoni, *Hook. f. Fl. Brit. Ind.* iv. 61.—Reg. Himal.
trinervia, *Hort. Mach. ex Regel, Gartenfl.* (1865) 306.—As. trop.
trinervis, Traill, in Trans. Hort. Soc. vii. (1830) 26 = Pottsii.
undulata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 408.—Malaya.
vaccinioides, *Hook. f. Fl. Brit. Ind.* iv. 56.—Reg. Himal.

HOYA:—
variegata, Siebold, ex Morr. in Ann. Soc. Bot. Gand, ii. (1846) 401. t. 90 = carnosa
volubilis, Griff. Notul. iv. 40 = coronaria?
volubilis, Wight, Contrib. 35 = coronaria.
verticillata, *G. Don, Gen. Syst.* iv. 138.—Ind. or.
viridiflora, R. Br. in Mem. Wern. Soc. i. (1809) 26 = Dregea volubilis.
viridiflora, Griff. Notul. iv. 87; Ic. Pl. Asiat. t. 388 = pruna?
vitellina, *Blume, Mus. Bot. Lugd. Bat.* i. 45.—Java.
volubilis, Griff. Notul. iv. 39 = Dregea volubilis.
Wallichiana, Decne. in DC. Prod. viii. 685 = coronaria.
Wightiana, Thw. Enum. Pl. Zeyl. 199 = pauciflora.
Wightii, *Hook. f. Fl. Brit. Ind.* iv. 59.—Ind. or. Zollingeriana, *Miq. Fl. Ind. Bat.* ii. 518.—Java.

HUALANIA, Phil. in Linnæa, xxxiii. (1864–65) 18 = **Bredemeyera**, Willd. (Polygal.).
colletioides, Phil. l. c. 19.—Am. austr.

HUANACA, Cav. Ic. vi. 18. t. 528 (1801). *UMBELLIFERAE*, Benth. & Hook. f. i. 876.
DIPLARPUS, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 469.
LECHLERIA, Phil. in Linnæa, xxxiii. (1864–65) 93, 94.
POZOSIS, Hook. Ic. Pl. t. 859 (1852).
TRISCIDIUM, Phil. in Anal. Univ. Chil. (1861) 1 61.
acaulis, Cav. Ic. vi. 18. t. 528 = Cavanillesii.
andina, *Phil. in Linnæa*, xxxiii. (1864–65) 93.—Chili.
Bergii, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 250.—Reg. Argent.
Cavanillesii, *DC. Prod.* iv. 81.—Reg. Magellan.
cordifolia, *F. Muell. Census*, 63.—Tasman.
grænlandica, *DC. Prod.* iv. 81.—Mexic.
hydrocotyles, *F. Muell. Census*, 63.—Austral.

HUANUCA, Raña. Sylva Tellur. 54 (1888) = **Aconitus**, Schott (Solanac.).
spathulata, Raña. l. c. = A. spathulatus.

HUBERIA, DC. Prod. iii. 167 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 748.
annulata, DC. l. c. = Marumia annulata.
argentea, DC. l. c. 168 = Miconia squamulosa.
laurina, *DC. l. c.* 167.—Bras.
ovalifolia, *DC. l. c.*—Bras.
reminoa, Presl, Symb. Bot. i. 58. t. 27 = ovalifolia.
semiserrata, *DC. Prod.* iii. 167.—Bras.

HUBERTIA, Bory, Voy. i. 334. t. 14 (1804) = **Senecio**, Tourn. (Compos.).
Ambavilla, Bory, l. c. = S. Ambavilla.
convexoides, Bory, l. c. ii. 333 = S. Hubertia.
pinifolia, Juss. ex Pers. Syn. ii. 436; Sieber, ex DC. Prod. vi. 293 = Fanjasia pinifolia.
salicifolia, Juss. ex Pers. l. c. = S. salicifolius.
tomentosa, Bory, Voy. i. 335. t. 14 = S. Hubertia.

HUDSONIA, Linn. Mant. i. 11 (1767). *CISTINEAE*, Benth. & Hook. f. i. 114.
australis, Spreng. Syst. ii. 452 = Heliantheum brasiliense.
ericoides, *Linn. Mant.* i. 74.—Am. bor.
ericoides, Richards. in Frankl. Narr. 1st Journ. App. 722.—nomenclos.
maritima, Torr. ex Steud. Nom. ed. II. i. 778 = tomentosa.
monzana, *Nutt. Gen. Am.* ii. 5.—Am. bor.
Nuttallii, Sweet, Cist. t. 19 = ericoides.
tomentosa, *Nutt. Gen. Am.* ii. 5.—Am. bor.

HUDSONIA, Robins. ex Lunan, Hort. Jamaic. ii. 210 (1814) = **Terminalia**, Linn. (Combret.).
arborea, Lunan, l. c. = T. Hilariana.

HUEGELIA, Benth. in Bot. Reg. sub t. 1622 (1833) = **Gilia**, Ruiz & Pav. (Polem.).
densifolia, Benth. in Bot. Reg. sub t. 1622 = G. densifolia.
elongata, Benth. l. c.; et in Journ. Hort. Soc. ii. (1847) 211 = G. densifolia.
lanata, Lindl. t. c. iii. (1846) 74 = G. lanata.
tenuis, Benth. in Bot. Reg. sub t. 1622 = G. tenuissima.
virgata, Benth. l. c. = G. virgata.

HUEGELIA, R. Br. ex Endl. Gen. 1156 (1840). † Genus ignotum,' Benth. & Hook. f. i. 888.

HUEGELIA, Reichb. Consp. 144 (1828) = **Trachymene**, Rudge (Umbellif.).
cyanea, Reichb. Ic. Exot. t. 201 = T. caerulea.

HUEGUENINIA, Reichb. Ic. Fl. Germ. ii. 20 (1837) = HUGUENINIA, Reichb. = **Stæmbrium**, Tourn. (Crucif.).

HUENEFELDIA, Walp. in Linnæa, xiv. (1840) 307 = **Calotis**, R. Br. (Compos.).
angustifolia, Walp. l. c. 506.—Austral.
coronopifolia, Walp. l. c. 507 = C. glandulosa.

HUERNIA, R. Br. in Mem. Wern. Soc. i. (1809) 22. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 784.
DECODONTIA, Haw. Syn. Pl. Succ. 28, in obs. (1812).
barbata, *Haw. Syn. Pl. Succ.* 31.—Afr. austr.
brevirostra, *N. E. Br. in Gard. Chron.* (1877) i. 780.—Afr. austr.
campanulata, *R. Br. in Mem. Wern. Soc.* i. (1809) 22.—Afr. austr.
clavigera, *Haw. Suppl. Pl. Succ.* 10.—Afr. austr.
crispa, *Haw. Syn. Pl. Succ.* 31.—Afr. austr.
duodecimfida, Sweet, Hort. Brit. ed. II. 359 = tubata.
guttata, *R. Br. in Mem. Wern. Soc.* i. (1809) 22.—Afr. austr.
humilis, *Haw. Syn. Pl. Succ.* 30.—Afr. austr.
Hystrix, *N. E. Br. in Gard. Chron.* (1876) i. 795.—Afr. austr.
lentiginosa, *Haw. Syn. Pl. Succ.* 29.—Afr. austr.
occellata, *Schult. Syst.* vi. 9.—Afr. austr.
ocellata, *Hook. f. Bot. Mag.* t. 6858.—Afr. austr.
reticulata, *Haw. Syn. Pl. Succ.* 28.—Afr. austr.
Thureti, *Cels, ex Hérincq, L'Horticult. Franç.* (1866) 78.—Afr. austr.?
tubata, *Haw. Syn. Pl. Succ.* 30.—Afr. austr.
venusta, *R. Br. in Mem. Wern. Soc.* i. (1809) 22.—Afr. austr.

HUERNIOPSIS, N. E. Br. in Journ. Linn. Soc. xvii. (1878) 171. *ASCLEPIADEAE*.
decipiens, *N. E. Br. l. c.*—Afr. austr.

HUERTA, Jaume St. Hil. Expos. ii. 357 (1805) = seq.

HUERTEA, Ruiz & Pav. Prod. 34. t. 6 (1794). *ANACARDIACEAE*, Benth. & Hook. f. i. 426.
cubensis, *Griseb. Cat. Pl. Cub.* 66.—Cuba.
glandulosa, *Ruiz & Pav. Fl. Per.* iii. 4. t. 227.—Peruv.

HUERTIA, G. Don, Gen. Syst. ii. 77 (1832) = prunc.

HUETIA, Boiss. Diagn. Ser. II. ii. 108 (1856) = **Carum**, Rupp.
pumila, Boiss. l. c. 104 = C. alpinum.

HUFELANDIA, Nees, Progr. Gratul. 11 (1833); et Laurin. Expos. (1836) 167, 674 = **Beilschmiedia**, Nees (Laurin.).
caracasana, Klotzsch & Karst. ex Meisn. in Mart. Fl. Bras. v. ii. 206.—N. Granat.
latifolia, Nees, Syst. Laurin. 674.—Peruv.
pendula, Nees, Disp. Laurin. 28 = B. pendula.
sulcata, Nees, in Linnæa, xxi. (1848) 494.—N. Granat.
thomasea, Nees, Disp. Laurin. 28 = B. thomasea.
thyrsiflora, Nees, Syst. Laurin. 190.—Madag.
tomentosula, Klotzsch & Karst. ex Meisn. in DC. Prod. xv. 65.—Am. austr.

HUGELIA, DC. Prod. iv. 72 (1830) = HUEGELIA, Reichb. = **Trachymene**, Rudge (Umbellif.).

HUGELROEA, Steud. Nom. ed. II. i. 778 (1840) = **Sphærocroloban**, Sm. (Legumin.).
lanceolata, Steud. l. c. = S. linophyllum.

HUGONIA, Linn. Gen. ed. i. 134 (1737). *LINEAE*, Benth. & Hook. f. i. 243, 267.
PENICILLANTHEMUM, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 94.
UGONA, Adans. Fam. ii. (63) (1763).
Afzelii, *R. Br. ex Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 625.—Afr. trop.

HUGONIA:—
angolensis, *Welw. ex Oliver, Fl. Trop. Afr.* i. 271.—Afr. trop.
conata, *Miq. Illustr. Vi. Ind. Arch.* 67.—Sumatra.
ferruginea, *Wight & Arn. Prod.* 73.—Zeylan.
foliosa, *Oliver, Fl. Trop. Afr.* i. 271.—Afr. trop.
integerrima, Stokes, Bot. Mat. Med. iii. 520 = H. Mystax.
Jenkinsii, *F. Muell. Fragm.* v. 7.—Austral.
macrocarpa, *Welw. Apont.* 585. n. 21.—Afr. trop.
macrophylla, *Oliver, Fl. Trop. Afr.* i. 271.—*Afr.* trop.
mystacina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = seq.
Mystax, Cav. Diss. iii. 377. t. 38. *f.* 1 = serrata.
Mystax, *Linn. Sp. Pl.* 675.—Ind. or.
obovata, Buch.-Ham. in Trans. Linn. Soc. xiv. (1824) 205 = H. Mystax.
Planchoni, *Hook. f. in Hook. Ic. Pl.* t. 777.—Afr. trop.
platyscypta, *Welw. ex Oliver, Fl. Trop. Afr.* i. 272.—Afr. trop.
serrata, *Linn. Encycl.* iii. 149.—Ins. Mascar.
spicata, *Oliver, Fl. Trop. Afr.* i. 270.—Afr. trop.
sumatrana, *Miq. Illustr. Fl. Ind. Arch.* 68.—Sumatra.
tomentosa, *Can. Diss.* iii. 178. t. 28. f. 2.—Ins. Maurit.

HUGUENINIA, Reichb. Fl. Germ. Excurs. 691 (1832) = **Sisymbrium**, Tourn. (*Crucif.*).
tanacetifolia, Reichb. l. c. = S. tanacetifolium.

HUIDOBRIA, C. Gay, Fl. Chil. ii. 438. t. 98 (1846) = **Loasa**, Juss.
chilensis, C. Gay, l. c. ii. 440.—Chili.
fruticosa, Phil. Fl. Atac. 19.—Chili.

HULSEA, Torr. & Gray, in Pacif. Rail. Rep. vi. 76, 77 (1857). *COMPOSITAE*, Benth. & Hook. f. ii. 401.
algida, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 547. —Calif.
brevifolia, *A. Gray, l. c.* vii. (1868) 359.—Calif.
californica, *A. Gray, in Pacif. Rail. Rep.* vi. 77.—Calif.
callicarpa, S. Wats. ex A. Gray, Syn. Fl. N. Am. ii. 1. 342 = vestita.
heterochroma, *A. Gray, in Proc. Am. Acad.* vii. (1868) 359.—Calif.
nana, *A. Gray, in Pacif. Rail. Rep.* vi. 76. t. 18.—Am. bor. occ.
Parryi, A. Gray, in Proc. Am. Acad. xii. (1877) 59 = vestita.
vestita, *A. Gray, l. c.* vi. (1864–65) 547.—Calif.

HULTEMIA, Reichb. Nom. 168 (1841) = seq.

HULTENIA, Reichb. Handb. 248 (1837) = HULTHEMIA, Dum. = **Rosa**, Tourn.

HULTHEMIA, Blume, ex Miq. Fl. Ind. Bat. i. 1. 160 (1855) = **Abrus**, Linn. (Legumin.).
javanica, Blume, l. c. = A. pulchellus.

HULTHEMIA, Dum. Not. Hulth. 18 (1824) = **Rosa**, Tourn.
berberifolia, Dum. l. c. = R. berberifolia.

HULTHENIA, Bronga. Enum. Gen. 125 (1843) = praec.

HUMBERTIA, Lam. Encyc. ii. 258 (1788). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 869.
ENDRACHIUM, Juss. Gen. 133 (1789).
SMITHIA, J. F. Gmel. Syst. ii. 388 (1791).
TMOUINIA, Sm. Ic. Ined. t. 7 (1789).
arvicrensis, Comm. ex Lam. Encyc. ii. 256 = madagascariensis.
madagascariensis, *Lam. l. c.*; *et Illustr.* 449. t. 106.—Madag.

HUMBOLDIA, Reichb. Consp. 156 (1828) = **Humboldtia**, Vahl (Legumin.).

HUMBOLDTIA, Neck. Elem. ii. 16 (1790) = **Voyria**, Aubl. (Gentian.).

HUMBOLDTIA, Ruiz & Pav. Prod. 121. t. 27 (1794) = **Pleurothallis**, R. Br. (Orchid.).
acutiflora, Ruiz & Pav. Syst. Veg. 236 = Stelis acutiflora.
aspera, Ruiz & Pav. l. c. = Stelis aspera.
contorta, Ruiz & Pav. l. c. 236 = Stelis contorta.
cordata, Ruiz & Pav. l. c. 234 = P. cordata.

HUMBOLDTIA:—
emarginata, Pav. ex Lindl. Gen. et Sp. Orch. 6 = Physosiphon emarginatus.
lanceolata, Ruiz & Pav. Syst. Veg. 237 = Stelis lanceolata.
oblonga, Ruiz & Pav. l. c. 236 = Stelis oblonga.
oblonga, Willd. ex Lindl. Fol. Orch. Stelis, 9 = Stelis ecostata.
parviflora, Ruiz & Pav. Syst. Veg. 236 = Stelis micrantha.
polystachya, Ruiz & Pav. l. c. 234 = Stelis polystachya.
purpurea, Ruiz & Pav. l. c. 235 = Pleurothallis linearis.
revoluta, Ruiz & Pav. l. c. = Stelis revoluta.
spiralis, Ruiz & Pav. l. c. 237 = P. spiralis.
succosa, Pav. ex Lindl. Gen. et Sp. Orch. 5 = P. ruscifolia.

HUMBOLDTIA, Vahl, Symb. Bot. iii. 106 (1794). *LEGUMINOSAE*, Benth. & Hook. f. i. 579.
BATSCHIA, Vahl, Symb. Bot. iii. 89. t. 56 (1794).
HUMBOLDIA, Reichb. Consp. 156 (1828).
africana, *Baill. Hist. Pl.* ii. 95.—Afr. trop.
Brunonis, *Wall. Pl. As. Rar.* iii. 17. t. 228.—Ind. or.
laurifolia, *Vahl, Symb. Bot.* iii. 106.—Ind. or.
pentapetala, *R. Br. ex Steud. Nom.* ed. II. i. 778 (Quid ?).—Zeylan.
tripetala, Heyne, ex Wight & Arn. Prod. 285 = Brunonis.
unijuga, *Sedd. Fl. Syle.* i. t. 183.—Ind. or.
Vahliana, *Wight, Ic.* iv. t. 1607.—Ind. or.

HUMEA, Roxb. Hort. Beng. 42 (1814); Fl. Ind. ii. 640 (1832) = **Brownlowia**, Roxb. (Tiliac.).
elata, Roxb. li. cc. = E. elata.

HUMEA, Sm. Exot. Bot. i. t. 1 (1804). *COMPOSITAE*, Benth. & Hook. f. ii. 317.
AGASMERIS, Delaun. in Bon Jardin. (1805); ex DC. Prod. vi. 157.
CALOMERIA, Vent. Jard. Malm. t. 78 (1804).
HARCKERIA, F. Muell. in Linnaea, xxv. (1852) 406.
OXIPHORIA, Hort. ex Dum.-Cours. Bot. Cult. ed. II. iv. 205 (1811).
RAZUMOVIA, Spreng. ex Juss. in Dict. Sc. Nat. xlv. 526 (1826).
cassiniacea, *F. Muell. Fragm.* i. 17.—Austral.
elegans, *Sm. Exot. Bot.* i. t. 1.—Austral.
ozothamnoides, *F. Muell. Fragm.* i. 17.—Austral.
paniculata, *F. Muell. l. c.* iii. 137.—Austral.
squamata, *F. Muell. l. c.* xi. 86.—Austral.

HUMIRIA (HOUMIRI, Aubl. Pl. Gui. i. 564. t. 225) (1775); Jaume St. Hil. Expos. ii. 374 (1806); Pers. Syn. ii. 70 (1807). *HUMIRIACEAE*, Benth. & Hook. f. i. 247.
HOUMIRIA, Juss. Gen. 395 (1789).
HUMIRIUM, Rich. ex Mart. Nov. Gen. et Sp. ii. 142 (1846).
MYRODENDRUM, Schreb. Gen. 358 (1789).
WERNERIA, Scop. Introd. 278 (1777).
amplexicaulis, Mart. ex Urban, in Mart. Fl. Bras. xii. ii. 440 = balsamifera.
arenaria, Guill. in Baill. Adansonia, i. (1860–61) 208 = floribunda.
balsamifera, *Jaume St. Hil. Expos.* ii. 374.—Guian.
contracta, Moric. ex Urban, in Mart. Fl. Bras. xii. ii. 451 = Vantanea parviflora.
crassifolia, *Mart. Nov. Gen. et Sp.* ii. 148.—Bras.
cuspidata, *Benth. in Hook. Kew Journ.* v. (1853) 101. —Bras.
densiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 374.—Guian.
dentata, *Casar. Decad. Nov. Stirp. Bras.* iv. 89.—Bras.
elliptica, Klotzsch, ex Urban, in Mart. Fl. Bras. xii. ii. 438 = floribunda.
floribunda, *Mart. Nov. Gen. et Sp.* ii. 145.—Bras.
guianensis, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 374.—Guian.
laurina, Klotzsch, ex Urban, in Mart. Fl. Bras. xii. ii. 439 = floribunda.
macrophylla, *Benth. in Hook. Kew Journ.* v. (1853) 102.—Bras.
montana, *A. Juss. in St. Hil. Fl. Bras. Mer.* i. 99.—Bras.
multiflora, Pritz. Ic. Ind. 561 = floribunda.
oblongifolia, *Benth. in Hook. Kew Journ.* v. (1853) 102.—Bras.
obovata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 373.—Guian.

HUMIRIA:—
parviflora, A. Juss. in St. Hil. Fl. Bras. Mer. ii. 63 = floribunda.
parvifolia, A. Juss. l. c. 89 = floribunda.
subcrenata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 374.—Guian.
surinamensis, Miq. Stirp. Surin. Sel. 36 = floribunda.

HUMIRIUM, Rich. ex Mart. Nov. Gen. et Sp. ii. 142 (1826) = praec.

HUMULUS, Linn. Syst. ed. I (1735). *URTICACEAE*, Benth. & Hook. f. iii. 356.
LUPULUS, Tourn. ex Mill. Gard. Dict. ed. VI. (1752).
aculeatus, Nutt. in Journ. Acad. Philad. N. S. i (1847) 182 = japonicus.
americanus, Nutt. l. c. 181 = H. Lupulus.
cordifolius, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 133. —Japon.
japonicus, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 89.—Japon.
Lupulus, *Linn. Sp. Pl.* 1028.—Europ.; Am. bor.
Lupulus, Thunb. Fl. Jap. 113 = japonicus.
volubilis, Salisb. Prod. 176 = H. Lupulus.
vulgaris, Gilib. Hist. Pl. Europ. ii. 348 = H. Lupulus.

HUNEFELDIA, Lindl. Veg. Kingd. 710 (1847) = HUENEFELDIA, Walp. = **Calotis**, R. Br. (Compos.).

HUNEMANNIA, A. Juss. in Orb. Dict. ix. 455 (1849) = seq.

HUNNEMANIA, G. Don, Gen. Syst. i. 129, 135 (1831) = seq.

HUNNEMANNIA, Sweet, Brit. Flow. Gard. iii. 54. t. 276 (1828). *PAPAVERACEAE*, Benth. & Hook. f. i. 54.
fumariaefolia, *Sweet, l. c.*—Mexic

HUNTERIA, [Moç. & Sesse,] ex DC. Prod. v. 649 (1836) = Porophyllum, Vaill. (Compos.).
columbina, [Moç. & Sesse,] ex DC. l. c. = P. Linaria.

HUNTERIA, Roxb. Hort. Beng. 84 (1814); Fl. Ind. i. 695 (1832). *APOCYNACEAE*, Benth. & Hook. f. ii. 698.
atrovirens, G. Don, Gen. Syst. iv. 105 = Chilocarpus atrovirida.
atroviridis, Wall. Cat. n. 1614 = Chilocarpus atroviridis.
coriacea, Wall. Cat. n. 1610 = Alyxia coriacea.
corymbosa, *Roxb. Hort. Beng.* 84; *Fl. Ind.* i. 695.—Ind. or.; Penang.
cuspidata, Wall. Cat. n. 1609 = Melodinus orientalis.
eugeniaefolia, Wall. l. c. n. 1615 = Wrightia tomentosa.
fasciculoris, Wall. l. c. n. 1612 = Alyxia fasciculoris.
gracilis, Wall. l. c. n. 1613 = Alyxia gracilis.
lanceolata, Wall. l. c. n. 1611 = corymbosa.
Roxburghiana, Wight, Ic. iv. t. 1294 = corymbosa.
sundana, *Miq. Fl. Ind. Bat.* ii. 409.—Ins. Sunda.
zeylanica, Gardn. ex Thw. Enum. Pl. Zeyl. 191 = corymbosa.

HUNTLEYA, Batem. ex Lindl. in Bot. Reg. sub t. 1991 (1837); et (1839) t. 14 = **Zygopetalum**, Hook. (Orchid.).
albidorubea, Lem. Illustr. Hortic. xv. (1868) t. 556.—Bras.
candida, Hort. ex Walp. Ann. iii. 543 = Z. candidum.
cerina, Lindl. & Paxt. Flow. Gard. iii. (1852–53) 62. *f.* 268 = Z. cerinum.
fimbriata, Hort. Hamb. ex Reichb. f. in Bot. Zeit. x. (1852) 634 = Z. gramineum.
grandiflora, Lem. Illustr. Hortic. xv. (1868) Misc. 59. —N. Granat.
imbricata, Hort. Hamb. ex Reichb. f. in Bot. Zeit. x. (1852) 714 = Z. cochleare.
marginata, Hort. ex Reichb. f. l. c. 636 = Z. marginatum.
Melangris, Lindl. Bot. Reg. sub t. 1991 = Z. Melangris.
radicans, Hort. Hamb. ex Linden, Pescatores, t. (1850) sub t. 15 = Z. candidum.
sessiliflora, Batem. in Lindl. Bot. Reg. sub t. 1991.—Guian.
violacea, Lindl. Sert. Orchid. t. 26 = Z. violaceum.

HUONIA, Montr. in Mém. Acad. Lyon, x. (1860) 185 = **Acronychia**, Forst. (Rutac.).
laevis, Montr. l. c. 186.—N. Caled.

HURA, Koen. ex Retz. Obs. iii. 49 (1783) = **Globbe**, Linn. (Scitamin.).
Koenigii, Roem. & Schult. Syst. i. 20, in obs. = G. versicolor.
siamensium, Koen. ex Retz. Obs. iii. 49 = G. versicolor.

HURA, Linn. Gen. ed. I. 377 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 839.
brasiliensis, Willd. Enum. Hort. Berol. 997 = crepitans.
crepitans, *Linn. Sp. Pl.* 1008.—Am. austr. polyandra, *Baill. Etud. Gén. Euph.* 543.—Mexic.
senegalensis, Baill. Adansonia, i. (1860-61) 77 = crepitans.
strepens, Willd. Enum. Hort. Berol. 997 = crepitans.

HUSANGIA, Juss. in Dict. Sc. Nat. xxii. 59 (1821) = HOSANGIA, Neck. = **Maleta**, Aubl. (Melastom.).

HUSEMANNIA, F. Muell. in Wing. South Sc. Record, iii. (1883) 197. *MENISPERMACEAE*.
protense, *F. Muell. l. c.*—Austral.

HUSNOTIA, Fourn. in Mart. Fl. Bras. vi. iv. 211 (1885). *ASCLEPIADEAE*.
rotundifolia, *Fourn. l. c.*—Bras.

HUSSONIA, Boiss. Diagn. Ser. I. viii. 46 (1849) = **Smarthrocarpus**, Lahll. (Crucif.).
Aegiceras, Cosa. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 15, 469 = seq.
uncata, Boiss. Diagn. Ser. I. viii. 46.—Arab.; Afr. bor.

HUSZIA, Klotzsch, in Monatsb. Berl. Acad. (Mærz 1854) 121; et Abh. Akad. Berl. (1854) 17. t. 1 A = **Begonia**, Linn.
octopetala, Klotzsch, ll. cc. 121; 18 = B. octopetala.
rubricaulis, Klotzsch, Abh. Akad. Berl. (1854) 19 = B. rubricaulis.

HUTCHINIA, Wight & Arn. Contrib. 84 (1834) = **Bouvardia**, Wight & Arn. (Asclep.).
indica, Wight & Arn. l. c. = B. Hutchinia.

HUTCHINSIA, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 93 (1812). *CRUCIFERAE*, Benth. & Hook. *f.* i. 92.
ASTYLUS, Dulac, Fl. Hautes-Pyr. 188 (1867).
HINTERNUMERA, Reichb. ex Nym. Consp. 66 (1878).
HORNUNGIA, Reichb. Ic. Fl. Germ. ii. t. 6 (1882).
NASTURTIOLUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 602 (1821).
affinis, Gren. ex Schultz, Arch. Fl. Fr. Allem. 274 = alpina.
alba, Bunge, Del. Sem. Hort. Dorp. (1839); et in Linnaea, xiv. (1840) Litt. 119 = Smelowskia cinerea.
alpina, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 93.—Europ.
alpina, Cosa. ex Willk. & Lange, Prod. Fl. Hisp. iii. 779 = Noccaea Auerswaldii.
aragonensis, *Loac. in Loac. & Pard. Ser. Inconf.* ed. I. 10, in nota.—Hispan.
Auerswaldii, Willk. Sert. Fl. Hisp. 14 = alpina.
aurea, Fenzl, Pugill. Pl. Nov. Syr. 14 (= *Chrysochamea velutina*).—Syria.
austriaca, Hook. f. Fl. Tasman. i. 93. t. 4 = Capsella australis.
bipartita, Bunge, Del. Sem. Hort. Dorp. (1839).—Cf. Linnaea, xiv. (1840) Litt. 119 = Smelowskia cinerea.
brevicaulis, Hoppe, ex Sturm, Deutschl. Fl. Heft 65 = alpina.
brevistyla, DC. Syst. ii. 887 = Thlaspi rivale.
calycina, Desv. Journ. Bot. iii. (1814) 168 = Smelowskia calycina.
canescens, Schur, Enum. Pl. Transs. 71 = alpina.
capnoides, DC. Syst. ii. 886 = Thlaspi capnoides.
chinensis, Steud. Nom. ed. II. i. 779 = Nasturtium sinense.
chrysantha, Jaub. & Spach, Illustr. i. 108 (= *Chrysochamea velutina*).—Syria.
corymbosa, J. Gay, in Syll. Retisb. i. (1834) 169 = Thlaspi rotundifolium.

HUTCHINSIA:—
diffusa, Jord. Diagn. i. 235 = Capsella procumbens.
flexuosa, Ledeb. Fl. Ross. i. 764 = Smelowskia borysthena.
foliosa, *Weihe in Ann. Sc. Nat.* Sér. V. i. (1864) 384. —Bolivia.
grandiflora, Soleir. ex Nym. Consp. 65 = Lepidium humifusum.
hastulata, DC. Syst. ii. 888 = Thlaspi hastulatum.
Lereti, Jord. ex Nym. Consp. 66 = Capsella procumbens.
maritima, Jord. Diagn. i. 234 = Capsella procumbens.
pauciflora, Loret, Herb Lozère, 10 = Prostii.
pauciflora, *Nym. Syll.* 207.—Europ. austr.
pectinata, *Bunge, ex Ledeb. Fl. Ross.* i. 201.—Sibir. altaic.
petraea, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 82.—Europ.
procumbens, Desv. Journ. Bot. iii. (1814) 168 = Capsella procumbens.
procumbens, Lec. & Lamotte, Cat. Pl. Cent. Fr. 76 = Prostii.
Prostii, *Y. Gay, ex Loret, Herb. Lozère*, 10.—Gallia.
pumila, DC. Syst. ii. 886 = Thlaspi pumilum.
pusillima, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 384.—Bolivia.
pygmaea, Viv. ex Spreng. Syst. iv. Cur. Post. 240 = Thlaspi rivale.
reticulata, *Griseb. in Goett. Abh.* vi. (1854) 116.—Reg. Magellan.
rotundifolia, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 82 = Thlaspi rotundifolium.
sinensis, Desv. Journ. Bot. iii. (1814) 168 = Nasturtium sinense.
stayneoides, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 99.—As. centr.
spinuscarum, Jord. Diagn. i. 237 = Capsella procumbens.
stylosa, DC. Syst. ii. 887 = Noccaea stylosa.
tasmanica, Hook. fc. Fl. t. 848 = Thlaspi tasmanicum.
tibetica, *T. Thoms. in Hook. l. c.* t. 900.—Reg. Himal.
trinervia, DC. Syst. ii. 887 = Iberidella trinervia.

HUTSCHINIA, D. Dietr. Syn. Pl. i. 579 (1839) = HUTCHINIA, Wight & Arn. = **Bouvardia**, Wight & Arn. (Asclep.).

HUTTIA, Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 51 = **Pachycarpa**, R. Br. (Dilleniac.).
conspicua, Drumm. ex Harv. l. c. = P. conspicuum.

HUTTIA, Preiss, ex Hook. Bot. Mag. t. 3884 (1840) = **Calectasia**, R. Br. (Liliac.).
elegans, Preiss, ex Hook. l. c. = C. cyanea.

HUTTONAEA, Harv. Thes. Cap. ii. t. 101 (1863). *ORCHIDACEAE*, Benth. & Hook. f. iii. 523.
HALLACKIA, Harv. Thes. Cap. t. 102 (1863).
fimbriata, *Reichb. f. in Flora*, l. (1867) 116.—Afr. austr.
Hallackii, Bolus, in Journ. Linn. Soc. xx. (1882) 339 = fimbriata.
pulchra, *Harv. Thes. Cap.* ii. 1.—Afr. austr.

HUTTONIA, Bolus, in Journ. Linn. Soc. xix. (1882) 339 = preac.

HUTTUM, Adans. Fam. ii. 88 (1763) = **Barringtonia**, Forst. (Myrtac.).

HYACINTHELLA, Schur, in Oestr. Bot. Wochenbl. (1856) 237 = **Hyacinthus**, Tourn. (Liliac.).
leucophaea, Schur, l. c. = H. leucophaeus.
pallens, Schur, l. c. (1857) 225 = H. leucophaeus.

HYACINTHOIDES, Heist. Syst. 5 (1748). Quid ?

HYACINTHOIDES, Medic. in Usteri, Ann. Bot. ii. (1791) 9 = **Scilla**, Linn. (Liliac.).
racemosa, Medic. l. c. = C. hispanica.

HYACINTHORCHIS, Blume, Mus. Bot. Lugd. Bat. i. 43. t. 16 (1849) = **Cremastra**, Lindl. (Orchid.).
variabilis, Blume, l. c. = C. Wallichiana.

HYACINTHUS, [Tourn.] Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 812.
BARNETIPS, Saliab. Gen. Pl. Fragm. 17 (1866).
BELLEVALIA, Lapeyr. in Journ. Phys. lxv i. (1808) 425.
BORBOVA, Rafin. Fl. Tellur. iii. 52 (1836).
BRIMMURA, Saliab. Gen. Pl. Fragm. 26 (1866).
BUSSEQUIA, Saliab. l. c. 26 (1866).
FOXIA, Farl. Nuov. Gen. Monocot. 17 (1854).
HYACINTHELLA, Schur, in Oestr. Bot. Wochenbl. (1856) 237.
PERIBOEA, Kunth, Enum. Pl. iv. 399 (1843).
RHYTIDOLOBUS, Dulac, Fl. Hautes-Pyr. 113 (1867).
SARCOMPHALIUM, Dulac, l. c. (1867).
STRAMOWEJA, Bertol. in Mem. Soc. Ital. Mod. xxi. (1835) 2. t. 1.
aboriteus, Caval. Deux Nouv. Esp. (1846) 14 = trifoliatus.
albulus, Jord. Pugill. 140 = orientalis.
amethystinus, Lam. Encyc iii. 190 = Scilla hispanica.
amethystinus, *Linn. Sp. Pl.* 317.—Hisp.; Pyren.; Croat.
amethystinus, Pall. Reise, iii. 589 = Pallasianus.
amoenus, C. Muell. in Walp. Ann. vi. 110, sphalm. = *albulus*.
angustifolius, Medic. in Usteri, Ann. Bot. ii. (1791) 18 = amethystinus.
aporus, *Forsk. Fl. Aegypt. Arab.* p. cx (Quid ?).—Arab.
Aucheri, *Baker, in Journ. Linn. Soc.* xi. (1871) 481. —Syria.
azureus, Baker, l. c. 427 = ciliatus.
belgicus, Hort. ex Steud. Nom. ed. II. i. 779 = Scilla festalis.
biflolius, *Boutel. ex Cav. Anal. Cienc.* v. (1802) 14. —Afr. austr.
bifolius, Gouan, ex Schult. f. Syst. vii. 594 = Muscari commutatum.
botryoides, All. Fl. Pedem. i. 118 = Muscari racemosum.
botryoides, Linn. Sp. Pl. 318 = Muscari botryoides.
brachypodus, *Borb.—Cf. Oestr. Bot. Zeitschr.* xxxiii. (1883) 303.—Europ.
brevifolius, *Thunb. Prod. Pl. Cap.* 68.—Afr. austr.
brumalis, Haw. ex G. Don, in Loud. Hort. Brit. Suppl. i. 591 = orientalis ?
campanulatus, Mill. Gard. Dict. ed. VIII. n. 3 = Scilla festalis.
candicans, Baker, in Saund. Refug. Bot. t. 174 = Galtonia candicans.
capitatus, Baker, in Journ. Bot. xii. (1874) 8 = sessiliflorus.
carnosus, Saliab. Gen. Pl. Fragm. 25, sphalm. = comosus.
cernuus, Linn. Sp. Pl. 317 = Scilla festalis.
chilensis, Molina, Sagg. Chil. ed. II. 285 = Leucocoryne ixioides.
ciliaris, Poir. Encyc. Suppl. iii. 120 = Drimia ciliaris.
ciliatus, *Cyrill. Pl. Rar. Neap. Fasc.* ii. 28. t. 10.—Europ. or; Afr. bor.; Orient.
ciliatus, Fleischer, ex Baker, in Journ. Linn. Soc. xi. (1871) 499 = lineatus.
colchicoides, *Delile, in Laborde, Voy. Arab. Petr.* 82. —Arab.
commutatus, Tenore, Syll. 176 = Muscari commutatum.
comosus, Herb. Unio Itin. ex Steud. Nom. ed. II. i. 779 = fastigiatus.
comosus, Linn. Sp. Pl. 318 = Muscari comosum.
convallarioides, Drège, ex Baker, in Journ. Linn. Soc. xv. (1876) 264 = Eriospermum convallarioides.
convallarioides, Linn. f. Suppl. 204 = Rhadamanthus convallarioides.
corymbosus, *Linn. Mant.* ii. 238.—Afr. austr.
creticus, Nym. Syll. Suppl. 64 = Chionodoxa cretica.
cryptopodus, *Baker, in Journ. Linn. Soc. xx.* (1883) 276.—Madag.
curvifolius, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 125 = amethystinus.
curvulus, Jord. ex Nym. Consp. 731 = amethystinus.
dalmaticus, Avé-Lall. in Jnd. Sem. Hort. Petrop. xi. (1846) 71 = Muscari pallens.
dalmaticus, *Baker, in Journ. Linn. Soc.* xi. (1871) 498.—Dalmat.
damascenus, *Baker, l. c.* 483.—Syria.
dubius, *Gmel. Adnot. Cat. Pl. Suecod.* (1831) 99, 78; *Fl. Sic. Prod.* i. 426.—Reg. Mediterr. or.
elatus, Poir. Encyc. Suppl. iii. 120 = Drimia elata.

HYACINTHUS:—
exaristatus, Baker, in Journ. Bot. xii. (1874) 7 = nervosa.
fastigiatus, *Bertol. Ann. Stor. Nat.* iv. 62.—Corsica.
flexuosus, *Baker, in Journ. Bot.* xii. (1874) 8.—Syria.
flexuosus, Thunb. Prod. Pl. Cap. 62 = Scilla flexuosa.
fuliginosus, Pall. in Nov. Act. Petrop. x. (1797) 309 = Muscari comosum.
fulvus, Cav. in Anal. Cienc. Nat. iii. (1801) 47 = Dipcadi serotinum.
glaucus, Baker, in Journ. Linn. Soc. xi. (1871) 488 = Scilla Hohenackeri.
grisea, *Brouss. Bleneh. Pl. Hort. Monsp.* 1804 (1805) 80, nomen; *ex Schult. f. Syst.* vii. 588 (Quid?).—Heb.?
Haynei, *Baker, in Journ. Bot.* xii. (1874) 7.—Palæstin.
hispanicus, Lam. Encyc. iii. 191 = amethystinus.
hispidus, *Baker, in Journ. Linn. Soc.* xi. (1871) 429.—Cilicia.
juncifolius, Lam. Fl. Fr. iii. 271 = Muscari racemosum.
lanatus, Linn. Sp. Pl. 318 = Lanaria plumosa.
leichtlinioides, *Baker, in Journ. Linn. Soc.* xi. (1871) 427.—Afr. trop.
leucophæus, *Stev. ex Ledeb. Fl. Ross.* iv. 156.—Ross. austr.
linearis, *Steud. ex Schult. f. Syst.* vii. 584.—Aa. Min.
lividus, Pers. Syn. i. 375 = Dipcadi serotinum.
macrobotrys, *Baker, in Journ. Linn. Soc.* xi. (1871) 433.—Palæstin.
maritimus, Pers. Syn. i. 376 = Muscari maritimum.
medius, Poir. Encyc. Suppl. iii. 120 = Drimia media.
micranthus, *Baker, in Journ. Linn. Soc.* xi. (1871) 430.—Aa. Min.
modestus, Jord. & Fourr. Ic. Pl. Europ. ii. 7 = orientalis.
monanthostaphesis, Desf. ex Poir. Encyc. iii. 119 = Dipcadi serotinum.
monstrosus, Linn. Sp. Pl. 318 = Muscari comosum.
montanus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 125 = amethystinus.
Muscari, Linn. Sp. Pl. 317 = Muscari moschatum.
nanus, Schult. f. Syst. vii. 581 = Chionodoxa nana.
nervosus, *Bertol. Misc. Bot.* i. 20.—Mesopot.
nivalis, *Baker, in Journ. Linn. Soc.* xi. (1871) 430.—Syria.
Non-scriptus, Habl. Taur. 118; ex Bieb. Fl. Taur. Cauc. i. 248 = Muscari racemosum.
Non-scriptus, Linn. Sp. Pl. 316 = Scilla festalis.
Non-scriptus, Thuill. Fl. Par. ed. ii. i. 173 = Scilla hispanica.
nutans, S. F. Gray, Nat. Arr. Brit. Pl. ii. 177 = Scilla festalis.
Olivieri, Baker, in Journ. Bot. xii. (1874) 8 = ciliatus.
orchioides, Linn. Sp. Pl. 318 = Lachenalia orchioides.
orientalis, *Linn. l. c.* 317.—Reg. Mediterr.
Pallasianus, *Stev. in Bull. Soc. Nat. Mosc.* xxx. (1857) ii. 85, in textu.—Reg. Caucas.
pallens, *Bieb. Fl. Taur. Cauc.* i. 253 = Muscari pallens.
pallens, Vis. Fl. Dalmat. i. 150 = dalmaticus.
pallidiflorus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 126 = amethystinus.
paniculatus, Lam. Encyc. iii. 192 = Muscari comosum.
paradoxus, Baker, in Journ. Bot. xii. (1874) 8 = flexuosus.
paradoxus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 30.—Persia.
parviflorus, Pers. Syn. i. 375 = Muscari parviflorum.
patriciniusus, Pall. ex Baker, in Journ. Linn. Soc. xi. (1871) 428 = Pallasianus.
patulus, Bertol. Misc. Bot. i. 21 = ciliatus.
patulus, Desf. Tabl. ed. i. 96; ed. ii. 80 = Scilla hispanica.
persicus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 213.—Persia.
peruvianus, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 795 = Scilla peruviana.
Pouzolzii, J. Gay, ex Loisel. in Mém. Soc. Linn. Par. vi. (1827) 407 = fastigiatus.
præcox, Jord. Pugill. 141 = orientalis.
præcox, Lam. Fl. Fr. iii. 271 = Scilla festalis.
Princeps, Baker, in Saund. Refug. Bot. t. 175 = Galtonia Princeps.
provincialis, Jord. Pugill. 139 = orientalis.
Pseudo-Muscari, *Baker, in Journ. Linn. Soc.* xi. (1871) 432.—Persia.
purpureus, Gilib. Notul. ii. 243; *fc.* t. 276 = Scilla Hohenackeri.

HYACINTHUS:—
pusillus, Thunb. ex Lam. Encyc. iii. 194; Poir. Encyc. Suppl. iii. 120 = Drimia pusilla.
pycnanthus, *Baker, in Journ. Linn. Soc.* xi. (1871) 430.—Armenia.
pyrenaeus, Jord. & Fourr. Pl. Nov. Brev. fasc. ii. 125 = amethystinus.
racemosus, Linn. Sp. Pl. 318 = Muscari racemosum.
revolutus, Linn. f. Suppl. 204 = Scilla revoluta.
rigidulus, Jord. & Fourr. Ic. Fl. Eur. ii. 7 = orientalis.
romanus, Desf. Fl. Atlant. i. 303 = dubius.
romanus, *Linn. Mant.* ii. 224.—Reg. Mediterr.
sarmaticus, Pall. in Nov. Act. Petrop. x. (1797) 309 = ciliatus.
scilloides, *Bonato, ex Steud. Nom.* ed. II. i. (Quid?). —Italia.
serotinus, Forsk. Fl. Aegypt. Arab. 209 = Dipcadi erythraeum.
serotinus, Linn. Sp. Pl. 317 = Dipcadi serotinum.
sessiliflorus, Baker, in Journ. Linn. Soc. xi. (1870) 429, partim = nervosa.
sessiliflorus, *Vis. Fl. Lib. Spec.* 21. t. viii. f. 2.—Afr. bor.
spicatus, Moench, Meth. 632 = Scilla hispanica.
spicatus, *Sibth. & Sm. Fl. Græc. Prod.* i. 287.—Græcia.
staveselensis, Steud. Nom. ed. II. i. 779 = Muscari staveselense.
sulphureus, Poir. Encyc. Suppl. iii. 119 = Dipcadi serotinum.
syriacus, Baker, in Journ. Bot. xii. (1874) 8 = trifoliatus.
trifoliatus, *Tenore, Fl. Nap.* iii. 376.—Reg. Mediterr. *viridis*, Linn. Syst. ed. X. 984 = Dipcadi viride.
Webbianus, *Nym. Syll. Suppl.* 64.—Italia.

HYAENACNE, Benth. & Hook. f. Gen. iii. 280 (1880) = seq.

HYAENANCHE, Lamb. Descr. Cinch. (1797) 52. t. 10 = Toxiocoendrum, Thunb. (Enphorb.).
capensis, Pers. Syn. ii. 637 = T. capense.
globosa, Lamb. Descr. Cinch. 52. t. 10 = T. capense.

HYALA, L'Hérit. ex DC. Prod. iii. 378 (1828) = **Polycarpæa**, Lam. (Caryophyll.).

HYALAEA, Benth. & Hook. f. Gen. ii. 482 (1873) = HYALAEA, Jaub. & Spach = **Centaurea**, Linn. (Compos.).

HYALAENA, C. Muell. in Walp. Ann. v. 70 (1858) = HYALOLAENA, Bunge = **Bellinum**, Linn. (Umbellif.).

HYALEA, Jaub. & Spach, Illustr. iii. 19. t. 214–217, 292 (1847) = **Centaurea**, Linn. (Compos.).
leucoides, Jaub. & Spach, l. c. 21. t. 216 = C. pergamacea.
muscroïdes, Jaub. & Spach, l. c. 19. t. 214 = C. muscroïdes.
Olivieriana, Jaub. & Spach, l. c. 188. t. 292 = C. Olivieriana.
pergamacea, Jaub. & Spach, l. c. 20. t. 215 = C. pergamacea.
pervica, Jaub. & Spach, l. c. 23. t. 217 = C. Candolleana.
pulchella, C. Koch, in Linnæa, xxiv. (1851) 422 = C. pulchella.
stenophylla, Jaub. & Spach, Illustr. iii. 22. t. 217 = C. Candolleana.

HYALIS, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1836) 106. *COMPOSITAE*, Benth. & Hook. f. ii. 466.
APHYLLOCLADUS, Wedd. Chlor. And. i. 11. t. 3 (1856).
JOHAPHES, Phil. Fl. Atac. 27. t. 4 (1860).
argentea, *D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1836) 106.—Chili.

HYALIS, Salisb. in Trans. Hort. Soc. i. (1812) 317 = Exlis, Linn. (Irid.).
aulica, Salisb. l. c. 318 = I. scarlosa.
gracilis, Salisb. l. c. 317 = I. capillaris, linearis.
lanciflora, Baker, in Mart. Fl. Bras. vi. III. 368.—Bras.
latifolia, Salisb. in Trans. Hort. Soc. i. (1812) 317 = I. capillaris.

HYALIS:—
longiflora, Salisb. in Trans. Hort. Soc. i. (1812) 318 = Tritonia longiflora.
Lorentzii, Klotzsch, Hercn. Ic. Fl. Argent. 82.—Reg. Argent.
marginifolia, Salisb. in Trans. Hort. Soc. i. (1812) 318 = Tritonia scariosa.

HYALISMA, Champ. in Calc. Journ. Nat. Hist. vii. (1847) 466 = **Scilaphila**, Blume (Triurid.).
janthina, Champ. l. c. 467.—Ind. or.

HYALOCALYX, Roiſe, in Journ. Linn. Soc. xxi (1884) 257. *TURNERACEAE* setiferus, *Roiſe, l. c.* 258.—Madag.

HYALOCHLAMYS, A. Gray, in Hook. Kew Journ. iii. (1851) 101 = **Angianthus**, Wendl. (Compos.).
globifera, A. Gray, l. c. = A. globifer.

HYALOLAENA, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 804 (Lehm. Rel. Bot. 128) = **Bellinum**, Linn. (Umbellif.).
jaxartica, Bunge, ll. cc.—Aa. temp.
Semenowii, Regel & Herd. in Bull. Soc. Nat. Mosc. xxxix. (1866) ii. 114.—Aa. centr.

HYALOLEPIS, DC. Prod. vi. 149 (1837) = **Myriocephalus**, Benth. (Compos.).
occidentalis, F. Muell. Fragm. iii. 155 = M. rhizocephalus.
rhizocephala, DC. Prod. vi. 149 = M. rhizocephalus.

HYALOSERIS, Griseb. in Goett. Abh. xxiv. (1879) 212. *COMPOSITAE*.
cinerea, *Griseb. l. c.* 213.—Reg. Argent.
rubicunda, *Griseb. l. c.* 212.—Reg. Argent.
salicifolia, *Hieron. Ic. Pl. Argent.* 35.—Reg. Argent.
tomentella, *Hieron. l. c.* 38.—Reg. Argent.

HYALOSPERMA, Steetz, in Lehm. Pl. Preiss. i. 476 (1845) = **Helipterum**, DC. (Compos.).
glutinosum, Steetz, l. c. 477 = Helipt. hyalospermum.
strictum, Steetz, l. c. = Helipt. hyalospermum.
variabile, Sond. in Linnæa, xxv. (1852) 519 = Helipt. hyalospermum.

HYALOSPERMUM, Benth. & Hook. f. Gen. iii. 309 (1873) = prec.

HYALOSTEMMA, Wall. Cat. n. 6484 (1831) = **Miltuse**, Leschen. (Anomac.).
Roxburghianum, Wall. l. c. = M. Roxburghiana.

HYBANTHERA, Endl. Prod. Fl. Norf. 59 (1838) = **Tylophora**, R. Br. (Asclepiad.).
bigландulosa, Endl. l. c. = T. bigландulosa.
cordifolia, Link, Klotzsch & Otto, Ic. Pl. Hort. Berol. 109. t. 44.—China.
javanica, Hassk. Cat. Hort. Bog. Alt. 209.—Java.
villosa, Miq. Fl. Ind. Bat. ii. 490 = T. villosa.

HYBANTHUS, Jacq. Enum. Pl. Carib. 2 (1760) = **Ionidium**, Vent. (1803) (Violac.).
calycinus, F. Muell. Fragm. x. 81 = I. calycinum.
concolor, Spreng. Syst. i. 805 = I. concolor.
debilissimus, F. Muell. Fragm. xi. 4 = I. debilissimum.
enneaspermus, F. Muell. l. c. x. 83 = I. suffruticosum.
filiformis, F. Muell. Nat. Pl. Vict. i. 44 = I. filiforme.
floribundus, F. Muell. l. c. 45 = I. floribundum.
howaensis, Jacq. Enum. Pl. Carib. 17 = I. Jacquinianum.
mexicanus, Ging. ex DC. Prod. i. 312.—Mexic.
strictus, Spreng. ex Steud. Nom. ed. II. i. 780 = I. strictum.
Tatei, F. Muell. in Trans. Roy. Soc. S. Austral. iv. (1882) 102.—Austral.
Vernonii, F. Muell. Nat. Pl. Vict. i. 45 = I. Vernonii.

HYBERICUM, Schrank, Baier. Fl. ii. 399 (1789), sphalm. = **Hypericum**, Tourn.

HYBIDIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 399 = **Centranthus**, DC. (Valerian.).
Calcitrapa, Fourr. l. c. = C. Calcitrapa.

HYBISCUS, Dum. Anal. Fam. 46 (1829) = **Hibiscus**, Linn. (Malvac.).

HYBOTROPIS, E. Mey. ex Steud. Nom. ed. II. i. 780 (1840) = **Rafnia**, Thunb. (Legumin.).

HYBRIDELLA, Cass. in Bull. Soc. Philom. (1817) 19; et in Dict. Sc. Nat. xxii. 86 (1821) = **Eclusania**, Pers. (Compos.).
globosa, Cass. in Dict. Sc. Nat. xxii. 86 = Z. globosa.

HYDASTYLIS, Steud. Nom. ed. II. i. 780 (1840) = seq.

HYDASTYLUS, Dryand. ex Salisb. in Trans. Hort. Soc. i. (1812) 310 = **Sisyrinchium**, Linn. (Irid.).
californicus, Salisb. l. c. = S. californicum.

HYDATICA, Neck. Elem. ii. 887 (1790) = **Saxifraga**, Tourn.
cuneifolia, Rafin. Fl. Tellur. ii. 65 = S. cuneifolia.
foliolosa, Rafin. l. c. = S. stellaria.
Grewn, Rafin. l. c. = S. Grewn.
hirsuta, Rafin. l. c. = S. umbrosa.
stellaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 580 = S. stellaris.
umbrosa, Rafin. Fl. Tellur. ii. 65 = S. umbrosa.

HYDNOCARPUS, Gaertn. Fruct. i. 288. t. 60 (1788). *BIXINEÆ*, Benth. & Hook. f. i. 129.
MAROTTIA, Rafin. Sylva Tellur. 155 (1838).
alpina, *Wight, ic.* t. 942.—Ind. or.
cauasana, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 197.—Ins. Andaman; Malaca.
corymbosa, *Seem. in Journ. Bot.* iii. (1865) 203.—Ins. Canar.
glaucescens, *Blume, Rumphia*, iv. 22.—Sumatra.
heterocilita, Spreng. Syst. iv. Cur. Post. 94 = Alsodeia Roxburghii.
heterophylla, *Blume, Rumphia*, iv. 22.—Java.
inebriani, Vahl, Symb. Bot. iii. 100 = venenata.
inebriani, Wall. Cat. n. 6670 = Wightiana.
laseia, *Miq. Fl. Ind. Bat. Suppl.* 853.—Sumatra.
lanceolata, Spreng. Syst. iv. Cur. Post. 93 = Alsodeia lanceolata.
mollocana, *Spreng. l. c.*—Ins. Molocc.
obtusa, Presl, Epim. Bot. 213 = Aberia abyssinica.
octandra, *Thw. in Hook. Kew Journ.* vii. (1855) 197.—Zeylan.
polyandra, *Blanco, Fl. Filip.* ed. II. 545.—Ins. Philipp.
venenata, *Gaertn. Fruct.* i. 288. t. 60. f. 3.—Zeylan. Wightiana, *Blume, Rumphia*, iv. 22.—Ind. or.

HYDNOPHYTUM, Jack, in Trans. Linn. Soc. xiv. (1823) 124. *RUBIACEÆ*, Benth. & Hook. f. ii. 129.
LASIOSTOMA, Spreng. Syst. i. 422 (1825).
Albertisii, *Becc. Malesia*, ii. (1884) 124; (1885) 136.—N. Guinea.
amboinense, *Becc. l. c.* 138.—Ins. Amboin.
andamanense, *Becc. l. c.* ii. (1885) 155, 156.—Ins. Andaman.
Bunel, *Becc. l. c.* (1884) 125.—Malaya.
borneanum, *Becc. l. c.* ii. (1885) 168.—Borneo.
borneense, *Becc. l. c.* 128.—Borneo.
coriaceum, *Becc. l. c.*; (1885) 154, 158.—Borneo.
crassifolium, *Becc. l. c.* 124; (1885) 148.—N. Guinea.
formicarum, *Jack, in Trans. Linn. Soc.* xiv. (1823) 124.—Malaya.
formicarum, Kurz, For. Fl. Brit. Burma, ii. 8 = andamanense.
Gaudichaudii, *Becc. Malesia*, ii. (1884) 124; (1885) 139.—Ins. Moluce.
grandiflorum, *Becc. l. c.* 126; (1885) 171.—Ins. Fiji.
Guppyanum, *Becc. l. c.* ii. (1885) 129, 183.—Ins. Salom.
Horneanum, *Becc. l. c.* ii. (1884) 125; (1885) 168.—Ins. Fiji.
hirtum, Becc. l. c. ii. (1885) 131 = seq.
hujusc, *Becc. l. c.* ii. (1884) 123. t. 21.—Malaya.
lanceolatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 257.—N. Guinea.
longiflorum, *A. Gray, in Proc. Am. Acad.* iv. (1860) 63.—Ins. Pacif.
longistylum, *Becc. Malesia*, ii. (1885) 152.—Ins. Salom.
loranthifolium, *Becc. l. c.* ii. (1884) 124; (1885) 146.—N. Guinea.
microphyllum, *Becc. l. c.* 126; (1885) 174.—N. Guinea.
montanum, Blume, Bijdr. 966 = formicarum.
montanum, Scheff. in Ann. jard. Buitenz. i. (1876) 31 = seq.

HYDNOPHYTUM:—
Moseleyanum, *Becc. Malesia*, ii. 125; (1885) 150.—Ins. Admiralty.
normale, *Becc. l. c.* (1884) 123; (1885) 130.—N. Guin.
oblongum, *Becc. l. c.* 124; (1885) 140.—N. Hibern.
ovatum, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 257.—Ins. Ternate.
papuanum, *Becc. Malesia*, ii. (1884) 124; (1885) 147.—N. Guin.
petiolatum, *Becc. l. c.* 124; (1885) 144.—N. Guinea.
philippinense, *Becc. l.c.* 125; (1885) 149.—Ins. Philipp.
radicans, *Becc. l. c.* 128; (1885) 182.—N. Guinea.
selebicum, *Becc. l. c.* 125; (1885) 155, 157.—Ins. Celebes.
simplex, *Becc. l. c.* 128; (1885) 129.—Ins. Aru.
sumatranum, *Becc. l. c.* 124; (1885) 137.—Sumatra.
tenuiflorum, *Becc. l. c.* 125; (1885) 169.—Ins. Fiji.
tetrapterum, *Becc. l. c.* ii. (1884) 126; (1885) 173.—N. Guinea.
tortuosum, *Becc. l. c.* 124; (1885) 141.—N. Guin.
Wilkinsoni, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 365.—Ins. Fiji.
Wilsoni, *Horne, ex Baker, l. c.*—Ins. Fiji.
Zippelianum, *Becc. Malesia*, ii. (1885) 174.—N. Guinea.

HYDNORA, Thunb. in Vet. Acad. Handl. Stockh. (1775) 69. *CYTINACEÆ*, Benth. & Hook. f. iii. 190.
APHYTRIA, Linn. Diss. resp. Achar. Aphyt. 7 (1776); Amoen. Acad. vii. 310. t. 7 (1785).
abyssinica, *A. Br. in Schweinf. Beitr. Fl. Aethiop.* 217, nomen.—Abyssin.
Ackarii, Thunb. ex Hook. f. in DC. Prod. xvii. 109 = africana.
aethiopica, *Decne. in Bull. Soc. Bot. Fr.* xx. (1873) 77.—Afr. trop.
africana, *Thunb. in Vet. Akad. Handl. Stockh.* (1775) 69.—Afr. austr.
americana, R. Br. in Trans. Linn. Soc. xiv. (1845) 345 = Prosopanche Burmeisteri.
angolensis, *Decne. in Bull. Soc. Bot. Fr.* xx. (1873) 77.—Afr. trop.
bogotensis, *Becc. in Nouv. Giorn. Bot. Ital.* iii. (1871) 6.—Abyss.
Johannis, *Becc. l. c.* 5.—Abyss.
longicollis, Welw. in Trans. Linn. Soc. xxvii. (1869) t. 21 = africana.
triceps, *Delpy & E. Mey. in Nov. Act. Nat. Cur.* xvi. ii. (1833) 779.—Afr. austr.

HYDNOSTACHYON, Liebm. in Kjoeb. Vidensk. Meddel. (1849) 23 = **Spathiphyllum**, Schott (Aroid.).
brevirostre, Liebm. l. c. 25 = S. brevirostre.
cochlearispathum, Liebm. l. c. 24 = S. cochlearispathum.
longirostre, Liebm. l. c. = S. cochlearispathum.

HYDORA, Bess. in Flora, xv. (1832) ii. Beibl. 18, nomen. = UDORA, Nutt. = **Rhodea**, Michx. (Hydrochar.).
canadensis, Bess. l. c. = R. canadensis.
lithuanica, Bess. l. c. (= *Udora lithuanica*).—Europ.

HYDRANGEA, Gronov. ex Linn. Coroll. Gen. 7 (1737). *SAXIFRAGEÆ*, Benth. & Hook. f. i. 640.
CORMIDIA, Ruiz & Pav. Prod. 53. t. 85 (1794).
HORTENSIA, Comm. ex Juss. Gen. 314 (1789).
SARCOSTYLIS, Presl, ex Ser. in DC. Prod. iv. 15 (1830).
acuminata, Sieb. & Zucc. Fl. Jap. i. 110 = H. Hortensia.
acuta, Rafin. New Fl. Am. iii. 77 = arborescens.
alba, Reinw. ex Miq. Fl. Ind. Bat. i. 1. 721 = oblongifolia.
alternifolia, *Siebold, in Nov. Act. Nat. Cur.* xiv. ii. (1829) 692.—Japon.
alsiaefolia, *Wall. Tent. Fl. Nep.* 2. t. 50.—Reg. Himal.
amplifolia, Rafin. New Fl. Am. iii. 78 = arborescens.
angulata, Tausch, in Flora, xviii. (1834) 494 = quercifolia.
anomala, D. Don, Prod. Fl. Nep. 211 = Dichroa febrifuga.
arborescens, *Linn. Sp. Pl.* 397.—Am. bor.
aspera, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 211.—Reg. Himal.
Azinal, Siebold, in Nov. Act. Nat. Cur. xiv. ii. (1829) 689 = H. Hortensia.

HYDRANGEA:—
Belzonii, Sieb. & Zucc. Fl. Jap. i. 109 = H. Hortensia.
bracteata, Sieb. & Zucc. l. c. 176 = scandens, *Maxim. Buergeri*, Sieb. & Zucc. l. c. 111 = H. Hortensia.
chinensis, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. x. (1867) n. xvi. 7.—China.
cordata, Pursh, Fl. Bor. Sept. i. 309 = arborescens.
cordifolia, Sieb. & Zucc. Fl. Jap. i. 113 = scandens, *Maxim.*
cuspidata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 98.—Japon.
cyanerea, Nutt. in Hook. Bot. Mag. t. 5088 = robusta.
discolor, Rafin. New Fl. Am. iii. 79 = radiata.
florida, Sallab. Prod. 394 = H. Hortensia.
frutescens, Moench, Meth. 106 = arborescens.
glauca, Hort. ex Dum.-Cours. Bot. Cult. ed. II. vii. 369; Link, Enum. Hort. Berol. i. 411 = radiata.
glauca, Rafin. New Fl. Am. iii. 77 = arborescens.
heteromalla, D. Don, Prod. Fl. Nep. 211 = aspera.
heterophylla, Hort. Cels, ex Steud. Nom. ed. I. 416 = seq. †
heterophylla, Rafin. New Fl. Am. iii. 78 = arborescens.
hirta, *Siebold, in Flora*, xi. (1828) 757.—Japon.
Hortensia, *Siebold, in Nov. Act. Nat. Cur.* xiv. ii. (1829) 688.—Ins. or.
hortensis, Sm. Ic. Pict. 12 = H. Hortensia.
involucrata, *Siebold, in Nov. Act. Nat. Cur.* xiv. ii. (1829) 691.—Japon.
japonica, Siebold, l. c. 689 = H. Hortensia.
khasiana, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 75 = aspera.
laevigata, Hort. Angl. ex Steud. Nom. ed. I. 416 = arborescens †
Lobbii, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. x. (1867) n. xvi. 15.—Archip. Ind.
macrophylla, *Ser. in DC. Prod.* iv. 15.—Japon.
Moellendorffii, *Hance, in Journ. Bot.* xii. (1874) 177.—China.
mutabilis, Steud. Nom. ed. I. 416 = H. Hortensia.
Neesiana, *Steud. l. c.* ed. II. i. 780 = radiata.
nivea, Michx. Fl. Bor. Am. i. 268 = radiata.
oblongifolia, *Blume, Bijdr.* 930.—Java.
opuloides, Steud. Nom. ed. I. 416 = H. Hortensia.
Otaksa, Sieb. & Zucc. Fl. Jap. i. 105 = H. Hortensia.
paniculata, Rafin. New Fl. Am. iii. 77 = arborescens †
paniculata, *Sieb. in Nov. Act. Nat. Cur.* xiv. ii. (1829) 691.—Japon.
peruviana, *Meric. in DC. Prod.* iv. 14.—Peruv.
petiolaris, Sieb. & Zucc. Fl. Jap. i. 106 = scandens, *Maxim.*
pubescens, *Court. in Syll. Ratisb.* ii. (1838) 45.—Japon.
pubescens, Decne. in Fl. des Serres, iv. (1848) t. 378 = aspera.
pubescens, Nees, ex Steud. Nom. ed. II. i. 780 = radiata.
pubescens, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 722 = Dichroa pubescens.
quercifolia, *Bartram, Trav.* 382. t. 7; ed. II. 380, cum tab. 6.—Am. bor.
quinquefolia, D. Dietr. Syn. Pl. 1525 = præc.
radiata, Sm. Ic. Pict. 13 pag. averso = quercifolia.
radiata, *Walt. Fl. Carol.* 131.—Am. bor.
robusta, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 76.—Reg. Himal.
rotundifolia, Rafin. New Fl. Am. iii. 78 = arborescens.
scandens, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. x. (1867) n. xvi. 16.—Japon.
scandens, *Pepp. ex DC. Prod.* iv. 664.—Chili.
scandens, Ser. in DC. Prod. iv. 15 = virens.
serrata, *Ser. l. c.*—Japon.
serratifolia, *Phil. f. Fl. Vasc. Chil.* 97.—Chili.
Sieboldii, Siebold, in Nov. Act. Nat. Cur. xiv. ii. (1829) 692 = H. Hortensia.
stellata, Sieb. & Zucc. Fl. Jap. i. 112 = H. Hortensia.
stylosa, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 75.—Reg. Himal.
Thunbergii, *Siebold, in Nov. Act. Nat. Cur.* xiv. ii. (1829) 690.—Japon.
vestita, Wall. Tent. Fl. Nep. 2. t. 50 = aspera.
villosa, Rafin. Sallab. Prod. 394 = arborescens.
virens, *Siebold, in Nov. Act. Nat. Cur.* xiv. ii. (1829) 690.—Japon.
vulgaris, Michx. Fl. Bor. Am. i. 268 = arborescens.
Wenzeslaii, *Vahle, in Ind. Sem. Hort. Berol.* (1872) *App.* 6.—Hab.?

HYDRANGIA, Linn. Gen. ed. VI. 228 (1764) = præc.

HYDRANTHELIUM, H. B. & K. Nov. Gen. et Sp. vii. 202. t. 646 (1824). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 987.
Brauniî, *Ernst, in Vorgesin.* i. (1870) 189.—Venezuel.
callitrichoides, *H. B. & K. Nov. Gen. et Sp.* vii. 203. t. 646.—N. Granat.
egense, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 75. t. 287.—Bras.
humifusum, *Wright, in Sauv. Fl. Cub.* 100.—Cuba.

HYDRASTIS, Ellis, ex Linn. Syst. ed. X. 1088 (1759). *RANUNCULACEÆ*, Benth. & Hook. f. i. 7.
Warnera, Mill. Gard. Dict. ed. VIII. (1768).
canadensis, *Linn. Syst.* ed. X. 1088.—Am. bor.
canadensis, Pair. Encyc. Suppl. iii. 71 = Trautvetteria palmata.
carolinensis, Walt. Fl. Carol. 156 = Trautvetteria palmata.
japonica, *Siebold, ex Miq. Ann. Mus. Lugd. Bat.* iii. 205.—Japon.
trifolia, Rafin. New Fl. Am. ii. 34 = canadensis!

HYDRASTYLIS, Steud. Nom. ed. II. i. 780 (1840) = HYDASTYLUS, Salisb. = **Sisyrinchium**, Linn. (Irid.).

HYDRIASTELE, H. Wendl. & Drude, in Linnæa, xxxix. (1875) 208. *PALMÆ*, Benth. & Hook. f. iii. 885.
Wendlandiana, *H. Wendl. & Drude, l. c.* 209.—Austral.

HYDRILLA, Rich. in Mém. Inst. Par. xii. (1811) ii. 9, 61. *HYDROCHARIDEÆ*, Benth. & Hook. f. iii. 460.
Epignanthus, Blume, in Flora, viii. (1825) 679.
Hydrospondylus, Hassk. in Flora, xxv. (1842) Beibl. 38.
Leptanthes, Wight, ex Wall. Cat. sub n. 5048 (1831).
Serpicula, Linn. f Suppl. 416 (1781) partim.
alternifolia, Miq Illustr. Fl. Ind. Arch. 52 = verticillata.
angustifolia, Hassk. Pl. Jav. Rar. 117; Blume, Mus. Bot. Lugd. Bat. i. 82 = verticillata
dentata, Casp. in Bot. Zeit. xii. (1854) 56 = verticillata.
Dregeana, Freal, Bot. Bemerk. 113 = verticillata.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 271 = verticillata.
muscoides, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 79 = verticillata.
najadifolia, Zoll. & Moritzi, Syst. Verz. Zoll. 91 = verticillata.
ovalifolia, Rich. in Mém. Inst. Par. xii. (1811) ii. 76. t. 2 = verticillata.
Roxburghii, Steud. Nom. ed. II. i. 780 = verticillata.
verticillata, *Presl, Bot. Bemerk* 112.—Ind. or.
Wightii, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 79 = verticillata.

HYDROBRYUM, Endl. Gen. 1375, partim (1841). *PODOSTEMACEÆ*, Benth. & Hook. f. iii. 112.
Griffithii, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 104.—Reg. Himal.
griseum, Tul. l. c. 104 = Podostemon olivaceus.
lichenoides, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 103 = Podostemon micranthus.
nudum, Bisch. ex Krauss, in Flora, xxvii. (1844) 496 = Sphærothylax algiformis.
olivaceum, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 104 = Podostemon olivaceus.

HYDROCALYX, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 43 = **Juxuillosa**, Ruiz & Pav. (Solan.).
quindiuensis, Triana, l. c. = J. speciosa.

HYDROCARPUS, D. Dietr. Syn. Pl. i. 574 (1839), sphalm. = **Hydnocarpus**, Gaertn. (Bixin.).

HYDROCERA, Blume, Bijdr. 941 (1826). *GERANIACEÆ*, Benth. & Hook. f. i. 276.
Tetronia, G. Don, Gen. Syst. i. 749 (1831).
angustifolia, *Blume, Bijdr.* 941.—Ind. or.; Malaya.
madagascariensis, *Endl. ex Walp. Rep.* i. 477.—Madag.
trifera, Wight & Arn. Prod. 140 = angustifolia.

HYDROCERAS, Hook. f. & Thoms. in Journ. Linn. Soc. iv. (1860) 188, lapsu = prace.

HYDROCHAERIS, Gaertn. Mey. & Scherb. Fl. Wett. iii. 400, 432 (1801) = **Hydrochæris**, Linn.

HYDROCHARELLA, Spruce, ex Benth. & Hook. f. iii. 462, in nota (1883) = **Limnobium**, Rich. (Hydrocharid.).
schimsporo, Spruce, l. c. = L. Sinclairi!

HYDROCHARIS, Linn. Syst. ed. I (1735). *HYDROCHARIDEÆ*. Benth. & Hook. f. iii. 462.
asarifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 220 = H. Morsus-ranæ.
asiatica, Miq. Fl. Ind. Bat. iii. 239 = H. Morsus-ranæ.
batrachydegma, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 71 = H. Morsus-ranæ.
celiulosa, Buch.-Ham. in Wall. Cat. n. 5042 = H. Morsus-ranæ.
cordifolia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = H. Morsus-ranæ.
Morsus-ranæ, *Linn. Sp. Pl.* 1036.—Europ.; Aa temp.
rotundifolia, Gilib. Fl. Lituan. ii. 221 = H. Morsus-ranæ.
Spongia, Bosc, in Ann. Mus. Par. ix. (1807) 396. t. 30 = Limnobium Bosci.

HYDROCHLOA, Beauv. Agrost. 135. t. 24. f. 4 (1812). *GRAMINEÆ*, Benth. & Hook. f. iii. 1115.
carolinensis, *Beauv. l. c.*—Am. bor.

HYDROCHLOA, Hartm. Gram. Skand. 8 (1819) = **Glyceria**, R. Br. (Gramin.).
axcides, Hartm. l. c. = G. aquatica.
aquatica, Hartm. l. c. = G. aquatica.
arundinacea, Sweet, Hort. Brit. ed. I. 453 = G. remota.
corsulea, Hartm Gram. Skand. 8 = Molinia cærulea.
distans, Hartm. l. c. = G. distans.
fluitans, Hartm. l. c. = G. fluitans.
maritima, Hartm. l. c = G. maritima.

HYDROCLEIS, Reichb. Consp. 45 (1828) = seq.

HYDROCLEYS, Rich. in Mém. Mus. Par. i. (1815) 368. t. 18. *ALISMACEÆ*, Benth. & Hook. f. iii. 1009.
Vesputcia, *Parl. Nuov. Gen. e Sp.* Monocot. 55 (1854).
asarca, Schult. f. in Mart. Fl. Bras. iii. i. 118 = Commersonii.
Commersonii, *Rich. in Mém. Mus. Par.* i. (1815) 368. t. 18.—Bras.
Humboldtii, Endl. Gen. 129 = Commersonii.
Martii, *Seub. in Mart. Fl. Bras.* iii. i. 116. t. 16. f. 1.—Bras.
nymphoides, Buchen. in Bremen, Abh. ii. (1871) 2 = Commersonii.
parviflora, *Seub. in Mart. Fl. Bras.* iii. i. 117. t. 18. f. 1. —Bras.

HYDROCOTILE, Crantz, Inst. ii. 140 (1766) = seq.

HYDROCOTYLE, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERÆ*, Benth. & Hook. f. i. 872.
Centella, Linn. Pl. Rar. Afr. 28 (1760).
Chorocarpus, Nutt. Gen. Am. Corrigend. (1818).
Glycesia, Nutt. Gen. Am. i. 177 (1818).
Trisanthus, Lour. Fl. Cochinch. 175 (1790).
abbreviata, A. Rich. in Ann. Sc. Phys. iv. (1820) 188. t. 58. f. 19 = asiatica.
acutiloba, *A. Rich. i. c.* 175. t. 58. f. 5.—N. Granat.
acuminata, *Urban, in Mart. Fl. Bras.* xi. i. 378.—Bras.
acutifolia, *Ruiz & Pav. Fl. Per.* iii. 25. t. 248.—Peruv.
adscens, Hochst. in Flora, xxiv. (1841) i. Intell. 28 = ranunculoides.
affinis, Eckl. & Zeyh. Enum. 286 = H. Centella.
alata, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 213. t. 61. f. 28.—Austral.
alchemilloides, *A. Rich. l. c.* 193. t. 56. f. 31.—Peruv.
alpestris, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 105.—Bras.
alpina, *Eckl. & Zeyh. Enum.* 284.—Afr. austr.
alpina, Vell. Fl. Flum. 128; iii. t. 89 = quinquefolia.
amblyna, Pursh, Fl. Am. Sept. ii. 732 = Erigenia bulbosa.
americana, *Linn. Sp. Pl.* 234.—Am. bor.

HYDROCOTYLE:—
americana, Walt. Fl. Carol. 113 = ranunculoides.
apiifolia, A. Cunn. ex Hook. f. Fl. Tasman. i. 155 = Xanthosia dissecta.
artemisia, *Montr. in Mém. Acad. Lyon*, x. (1860) 212.—N. Caled.
asiatica, Bert. ex Urban, in Mart. Fl. Bras. xi. i. 273 = modesta.
asiatica, *Linn. Sp. Pl.* 234.—Reg. trop. et subtrop.
asiatica, Thunb. Fl. Cap. 303 = triantha.
Asterias, *Cham. & Schlecht. in Linnæa*, i. (1826) 861 (err. typ. 261).—Bras.
Barbarosa, *Cham. & Schlecht. l. c.* 860 (err. typ. 260). —Bras.
batrachoides, DC. Prod. iv. 667 = ranunculoides.
Batrachium, Hance, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 220 = rotundifolia.
bidentata, J. F. Gmel. Syst. 1568, sphalm. = tridentata.
biflora, Vell. Fl. Flum. 124; iii. t. 98 = asiatica.
bipinnata, Rafin. in Med. Repos. N. York, v. (1808) 353, nomen; Muhl. Cat. ed. II. 39 = Erigenia bulbosa.
blepharocarpa, *F. Muell. in Wing, South Sc. Record*, (July 1888).—Austral.
bonariensis, Lam. Encyc. iii. 147 = umbellata.
Bonplandi, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 192. t. 54. f. 7.—N. Granat.
brachystachya, DC. Prod. iv. 68 = hirsuta.
bracteata, *DC. l. c.* 668.—Afr. trop.
brasiliensis, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1849) 296 = asiatica.
brevipedata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = seq.
brevipes, DC. Prod. iv. 68 = asiatica.
bulbosa, Eaton & Wright, N. Am. Bot. 277 = Erigenia bulbosa.
bupleurifolia, A. Rich. in Ann. Sc. Phys. iv. (1820) 218. t. 67. f. 39 = H. Centella.
burmanica, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 60; (1877) ii. 113.—Burma.
caffra, Meisn. in Hook. Lond. Journ. Bot. ii. (1843) 523 = umbellata.
callicarpa, *Bunge, in Lehm. Pl. Preiss.* i. 283 — Austral.
calliophala, *Cham. in Linnæa*, viii. (1833) 329.—Bras.
calliodus, *Cham. & Schlecht. l. c.* i. (1826) 871.—Afr. austr.
capillaris, *F. Muell. Fragm.* iv. 178.—Austral.
capitata, Banks & Soland. ex A. Rich. in Ann. Sc. Phys. iv. (1820) 201 = compacta.
capitata, *Thou. Fl. Tristan d'Acugn.* 48. t. 12.—Ins. Trist. da Cunha.
capitata, Willd. in Schult. Syst. vi. 856 = Bonplandi.
Centella, *Cham. & Schlecht. in Linnæa*, i. (1826) 875 (err. typ. 275).—Afr. austr.
Chamaemorus, *Cham. & Schlecht. l. c.* 868 (err. typ. 268).—Chili.
chilensis, Phil. f. Cat. Pl. Vasc. Chil. 105 = Bonplandi.
chinensis, Linn. Sp. Pl 234 = Crantzia lineata.
citrioloa, *Ruiz & Pav. Fl. Per.* iii. 26.—Chili; Peruv.
communis, DC. Prod. iv. 59 = interrupta.
compacta, A. Rich. in Ann. Sc. Phys. iv. (1820) 201 = mochata.
composita, Pursh, Fl. Am. Sept. i. 190 = Erigenia bulbosa.
concinna, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 389.—N. Zel.
conferta, *Wight, Ic.* t. 1009.—Ind. or.
cordata, Walt. Fl. Carol. 113 = Limnanthemum trachyspermum.
cordifolia, Hook. f. in Hook. Ic. Pl. t. 803 = asiatica.
crassiuscula, *Tate, in Trans. Roy. Soc. S. Austral.* vii. (1885) 69.—Austral.
crinita, Thou. ex DC. Prod. iv. 65 = capitata.
cryptocarpa, *Speg. in Anal. Soc. Cient. Argent.* xv. (1883) 106.—Reg. Argent.
cuspidata, Willd. in Schult. Syst. vi. 896 = triantha.
Cusconi, *Montr. in Mém. Acad. Lyon*, x. (1860) 211. —N. Caled.
Cymbalaria, Benth. in Enum. Pl. Hueg. 53 = alata.
cymbalarifolia, Muhl. Cat. 80 = ranunculoides.
debilis, *Eckl. & Zeyh. Enum.* 286.—Afr. austr.
denudiora, DC. Prod. iv. 67 = laxiflora.
dentata, A. Rich. in Ann. Sc. Phys. iv. (1820) 192. t. 60. f. 22 = asiatica.

HYDROCOTYLE:—
diantha, *DC. Prod.* iv. 68.—Austral.
dichondrifolia, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1838) 211 = Novæ-Zealandiæ.
difformis, Eckl. & Zeyh. Enum. 338 = montana.
dissecta, *Hook. f. Fl. N. Zeal.* 84.—N. Zel.
dissecta, Rafin in Med. Repos. N. York, v. (1808) 353 (Quid?).—Am. bor.
Drapeana, *Sond. in Harv. & Sond. Fl. Cap.* ii. 580.—Afr. austr.
Dus, Vell. Fl. Flum. 125; iii. t. 96 = leucocephala.
elegans, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 198. t. 58. f. 52.—Austral.
elongata, *A. Cunn. ex Hook. f. Fl. N. Zel.* i. 84.—N. Zel.
emarginata, Vell. Fl. Flum. 125; iii. t. 95 = leucocephala.
erecta, Linn. f. Suppl. 177 = peltata ?
crientha, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 183. t. 56. f. 13.—Afr. austr.
falcata, Eckl. & Zeyh. Enum. 335 = H. Centella.
ficarifolia, Stokes, Bot. Mat. Med. ii. 57 = salatica.
ficarioides, Lam. Encyc. iii. 153 = repanda.
filicaulis, *Baker, in Journ. Linn. Soc. xxi.* (1884) 348. —Madag.
flexuosa, Eckl. & Zeyh. Enum. 338 = virgata.
fluctida, R. Br. ex A. Rich. in Ann. Sc. Phys. iv. (1820) 200 = heteromeria.
fluxuosa, *Eckl. & Zeyh. Enum.* 331.—Afr. austr.
fluitans, DC. Prod. iv. 69 = umbellata.
fluminensis, Vell. Fl. Flum. 124; iii. t. 90 = quinquelobata.
fusca, Eckl. & Zeyh. Enum. 336 = H. Centella.
Gaudichaudiana, DC. Prod. iv. 67 = hirta ?
geranifolia, *F. Muell. in Trans. Vict. Inst.* i. (1855) 136.—Austral.
geraniifolia, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 197. t. 59. f. 31.—Peruv.
glabra, *Thunb. Prod. Pl. Cap.* 49.—Afr. austr.
glabrata, Linn. f. Suppl. 176; Lam. Encyc. iii. 155 = H. Centella.
glechomoides, A. Rich. in Ann. Sc. Phys. iv. (1820) 180. t. 58. f. 17 = Dysopsis Guyana.
globata, *Blume, Bijdr.* 883.—Java.
globifera, Urban, in Mart. Fl. Bras. xi. t. 381 = seq.
globiflora, *Ruiz & Pav. Fl. Per.* iii. 25. t. 247.—Am. austr.
glochidiata, *Benth. Fl. Australiæ.* iii. 346.—Austral.
gracilentis, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 467 = peduncularis.
gracilis, *Ruiz & Pav. Fl. Per.* iii. 26. t. 247.—Peruv.
grammatocarpa, *F. Muell. Fragm.* ii. 138.—Austral.
grossulariodes, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 207. t. 62. f. 80.—Ins. Borbon.
grumosa, [Moç. & Sesse] ex DC. Prod. iv. 70 = Microplcura ranifolia.
gummifera, Lam. Encyc. iii. 156 = Azorella caespitosa.
gunnerifolia, *Wedd. Chlor. Andin.* ii. 186.—N. Granat.
hebecarpa, DC. Prod. iv. 68 = asiatica.
hederæfolia, *Bunch. Trav.* i. 46.—Afr. austr.
hermanniæfolia, *Eckl. & Zeyh. in S. Afr. Quart. Journ.* (1830) 375.—Afr. austr.
heterocrela, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 200. —N. Zel.
Heyneana, Wall. Cat. n. 5652 = javanica.
hirsuta, Blume, Bijdr. 884 = javanica.
hirsuta, *Sm. Prod. Veg. Ind. Oac.* 84; *Fl. Ind. Occ.* i. 560.—Ins. S. Doming.
hirta, *R. Br. ex A. Rich. in Ann. Sc. Phys.* iv. (1820) 204.—Austral.
hispida, D. Don, Prod. Fl. Nep. 158 = javanica.
hispidula, *Bunge, in Lehm. Pl. Preiss.* i. 283.—Austral.
hemalocarpa, F. Muell. Fragm. ii. 129 = verticillata.
Humboldtii, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 179. t. 55. f. 6.—Ecuador.
Humboldtii, Willd. ex DC. Prod. iv. 61 = Bonplandi.
humifusa, *Pohl, ex DC. Prod.* iv. 63.—Bras.
inconspicua, DC. Prod. iv. 69 = asiatica.
incrassata, Rafin. Fl. Ludov. 31 = umbellata.
incrassata, *Ruiz & Pav. Fl. Per.* iii. 26.—Peruv.
indecora, *DC. Prod.* iv. 668.—Chili.
indivisa, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 83 = asiatica.
intermixta, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 240.—N. Zel.

HYDROCOTYLE:—
interrupta, *Muhl. Cat.* 10.—Am. bor.
intertexta, R. Br. ex A. Rich. in Ann. Sc. Phys. iv. (1820) 202 = hirta.
inundata, Sm. Fl. Brit. i. 290 = Apium inundatum.
isolobs, *Wedd. Chlor. Andin.* ii. 189.—Ecuador.
javanica, *Thunb. Diss. Hydroc.* n. 17. t. 3; *ed. Pers.* ii. 415. t. 2.—Java.
javanica, Zoll. Syst. Verz. Ind. Archip. 41 = sundaica.
Langsdorffii, *DC. Prod.* iv. 60.—Bras.
lanuginosa, Eckl. & Zeyh. Enum. 337 = virgata.
latisecta, *Zoll. Syst. Vers. Ind. Archip.* 139, 140.—Java.
laxiflora, *DC. Prod.* iv. 61.—Austral.
lemnoides, *Benth. Fl. Australiæ.* iii. 345.—Austral.
leptostachya, A. Rich. in Ann. Sc. Phys. iv. (1820) 191. t. 57. f. 18 = hirsuta.
leptostachys, Spreng. Syst. i. 876 = asiatica.
leucocephala, *Cham. & Schlecht. in Linnæa*, i. (1826) 264.—Bras.
ligustrina, Spr. ex DC. Prod. iv. 71 = Crantzia lineata.
linearis, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 582 = H. Centella.
linearis, Michx. Fl. Bor. Am. i. 162 = Crantzia lineata.
linifolia, Linn. f. Suppl. 176 = H. Centella.
littoralis, Eckl. & Zeyh. Enum. 334 = hermanniæfolia.
lobocarpa, F. Muell. Fragm. iv. 178 = mediocginoides.
lunata, Lam. Encyc. iii. 152 = asiatica.
lurida, Hance, in Walp. Ann. ii. 690 = asiatica.
macrocarpa, A. Rich. in Ann. Sc. Phys. iv. (1820) 220. t. 67. f. 40 = virgata.
macrodonta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 157 = sec.
macrodus, Spreng. Neue Entdeck. ii. 292; Syst. i. 877 = hederæfolia.
macrophylla, Pohl, ex DC. Prod. iv. 61 = quinquelobata.
Mannii, Hook. f. in Journ. Linn. Soc. vii. (1864) 194 = hirta.
marchantioides, Clos, in C. Gay, Fl. Chil. iii. 67 = hirta.
medicaginoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 27.—Austral.
medicaus, *Cham. & Schlecht. in Linnæa*, v. (1830) 208.—Mexic.
microphylla, *A. Cunn. in Ann. Nat. Hist.* Ser. I. ii. (1838) 211.—N. Zel.
miaima, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 319.—Abyssin.
minuta, Pohl, in DC. Prod. iv. 66 = leucocephala.
modesta, *Cham. & Schlecht. in Linnæa*, i. (1826) 258 (err. typ. 258).—Reg. Argent.
mollissima, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 529 = hirsuta.
monopetala, Blanco, Fl. Filip. ed. I. 218 = Epithema triandrum.
monsuen, *Cham. & Schlecht. in Linnæa*, i. (1826) 374 (err. typ. 274).—Afr. austr.
montana, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 532 = H. Centella.
monticola, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 194.—Afr. trop.
moschata, *Forst. f. Prod.* 23.—N. Zel.
muscifera, Spreng. in Schult. Syst. vi. 351 = hederæfolia.
multicaulis, Pohl, ex DC. Prod. iv. 64 = leucocephala.
multifida, *A. Rich. in Ann. Sc. Phys.* iv. (1820) 208.—N. Zel.
multiflora, Ruiz & Pav. Fl. Per. iii. 24 = umbellata.
muricinata, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 28.—Austral.
muscosa, R. Br. ex A. Rich. in Ann. Sc. Phys. iv. (1820) 208. t. 61. f. 37 = triparitis.
natans, Cyrilli, Pl. Rar. Neap. Fasc. i. 20. t. 6 B = ranunculoides.
natans, *Thunb. Diss. Hydroc.* 3.—Reg. temp.
natans, Torrey, Bot. Mex. Bound. 69 = prolifera.
nepalensis, Hook. Exot. Bot. i. t. 30 = javanica.
nitidula, A. Rich. in Ann. Sc. Phys. iv. (1820) 200. t. 58. f. 28 = rotundifolia.
Novæ-Zealandiæ, *DC. Prod.* iv. 67.—N. Zel.
nummularioides, A. Rich. in Ann. Sc. Phys. iv. (1820) 176. t. 155. f. 9 = asiatica.
nutans, G. Don, in Lond. Hort. Brit. 106, sphalm. = natans.
pallida, DC. Prod. iv. 68 = asiatica.
pallida, Eckl. & Zeyh. Enum. 331 = crientha.

HYDROCOTYLE:—
parstyensis, *Vell. Fl. Flum.* 124; iii. t. 92. f. 11.—Bras.
pedicellata, *F. Muell. Fragm.* iv. 183.—Austral.
pedunculatis, *R. Br. ex A. Rich. in Ann. Sc. Phys.* iv. (1820) 202. t. 61. f. 36.—Austral.
peltata, Salisb. Prod. 152 = vulgaris.
pereniqua, Hance, in Walp. Ann. ii. 691 = rotundifolia.
petiolaris, DC. Prod. iv. 60 = umbellata.
pilifera, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 26.—Austral.
pinnatifida, *Sprong. Pugill.* ii. 49.—Hab. ?
plantaginea, Spreng. Neue Entdeck. i. 284 = H. Centella.
plebeia, *R. Br. ex A. Rich. in Ann. Sc. Phys.* iv. (1820) 184. t. 60. f. 23.—Austral.
plerantha, Cesati, in Linnæa, xi. (1837) 318 = vulgaris.
podantha, Molk. in Miq. Pl. Jungh. 89 = javanica.
Poeppigii, *DC. Prod.* iv. 667.—Chili.
polycephala, Wight & Arn. Prod. 356 = javanica.
polyrrhiza, DC. Prod. iv. 65 = leucocephala.
polystachya, A. Rich. in Ann. Sc. Phys. iv. (1820) 171 = umbellata.
prolifera, *Kellogg, in Proc. Calif. Acad.* i. (1854) 15.—Am. bor. occ.
prolifera, *Otto, ex Sweet, Hort. Brit. ed.* III. 295, nomen.—Bras.
pterocarpa, *F. Muell. in Trans. Vict. Inst.* i. (1855) 136.—Austral.
pulchella, R. Br. ex A. Rich. in Ann. Sc. Phys. iv. (1820) 199 = hirta.
puncticulata, Miq. Fl. Ind. Bat. i. t. 782 = rotundifolia.
pusilla, R. Br. ex Rich. in Ann. Sc. Phys. iv. (1820) 213 = alata.
pusilla, *A. Rich. l. c.* 167. t. 52. f. 2.—Reg. Argent.
pygmæa, *Wright, in Sauv. Fl. Cub.* 57.—Cuba.
quinquiloba, *Ruiz & Pav. Fl. Per.* iii. 25. t. 248.—Peruv.
quinqueradiata, Thou. ex DC. Prod. iv. 60 = umbellata.
racemosa, [Moc. & Sesse], ex DC. l. c. 70 = interrupta.
ranunculoides, Blume, Bijdr. 884 = rotundifolia.
ranunculoides, Heyne, ex Wall. Cat. n. 5653 = javanica.
ranunculoides, *Linn. f. Suppl.* 177.—Am. bor.
reniformis, *Spreng. Neue Entdeck.* i. 282 = crientha.
reniformis, Walt. Fl. Carol. 113 = repanda.
repanda, *Pers. Syn.* i. 302.—Am. bor.
rhombifolia, *F. Muell. Fragm.* vii. 147.—Austral.
rigescens, Eckl. & Zeyh. Enum. 337 = H. Centella.
rotundifolia, *Roxb. Hort. Beng.* 31; *Fl. Ind.* ii. 88.—As. et Afr. trop.
rotundifolia, Wall. Cat. n. 5654 = asiatica.
rugulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 27.—Austral.
rupestris, Eckl. & Zeyh. Enum. 334 = H. Centella.
sanicularifolia, Lam. Encyc. iii. 154 = Dipoda sanicularifolia.
sarmentosa, Salisb. Prod. 159 = asiatica.
scaposa, Steud. in Flora, xxvi. (1853–55) 768 = umbellata.
Schkuhriana, Reichb. Fl. Germ. Excurs. 432 = vulgaris.
scutellifera, *Benth. Fl. Australiæ.* iii. 343.—Austral.
sibthorpioides, Lam. Encyc. iii. 158 = rotundifolia.
sinensis, J. F. Gmel. Syst. 468 = Crantzia lineata.
Solandra, *Linn. f. Suppl.* 176.—Afr. austr.
Spananthe, Sm. ex DC. Prod. iv. 70 = Apium inundatum.
Spananthe, Willd. Sp. Pl. i. 1368 = Spananthe paniculata.
sphæroloba, *Wedd. Chlor. Andin.* ii. 189.—N. Granat.
spicata, Lam. Encyc. iii. 153 = hirsuta.
splendens, *Blume, Bijdr.* 884.—Java.
splendens, Zoll. ex Miq. Fl. Ind. Bat. i. t. 788 = rotundifolia.
Stella, Pohl, ex DC. Prod. iv. 61 = quinquelobata.
stellata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 158 = H. Asteriss.
striata, *Benth. Enum. Pl. Hong.* 58.—Austral.
strigosa, Bunb.-Ham. in Wall. Cat. n. 7219 = javanica.
sundaica, *Blume, Bijdr.* 883.—Malaya.
superposita, *Baker, in Journ. Linn. Soc.* xxi. (1884) 348.—Madag.
tabularis, Steud. Nom. ed. II. i. 782 = alpina.
tasmanica, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 467 = hirta.

HYDROCOTYLE:—
tenella, D. Don, Prod. Fl. Nep. 182 = rotundifolia.
tetragonocarpa, *Bunge*, *in Lehm. Pl. Preiss.* i. 294.—Austral.
Thunbergiana, Spreng. Neue Entdeck. i. 283 = asiatica.
tomentosa, Thunb. Prod. Pl. Cap. 49 = H. Solandra.
trachycarpa, *F. Muell. in Linnaea*, xxv. (1853) 394.—Austral.
tribotrys, Ruiz & Pav. Fl. Peruv. iii. 24 = umbellata.
trichophylla, Eckl. & Zeyh. Enum. 333 = virgata.
tridentata, *Linn. f. Suppl.* 176; *Thunb. Prod. Pl. Cap.* 49.—Afr. austr.
triflora, Ruiz & Pav. Fl. Per. iii. 24 = asiatica.
triflora, *Vell. Fl. Flum.* 125; iii. t. 94.—Bras.
triloba, *Thunb. Diss. Hydroc.* 6.—Afr. austr.
tripartita, *R. Br. ex A. Rich. in Ann. Sc. Phys.* iv. (1820) 209. t. 61. f. 25.—Austral.
tripartita, Hook. f. in Hook. Ic. Pl. t. 812 = callicarpa.
tussilaginifolia, *Baker*, *in Journ. Linn. Soc.* xx. (1883) 151.—Madag.
umbellata, *Linn. Sp. Pl.* 234.—Reg. trop.
umbellulata, Michx. Fl. Bor Am. i. 161 = umbellata.
uncinata, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 169 = villosa.
uniflora, *Colenso*, *in Trans. N. Z. Inst.* xvii. (1885) 229.—N. Zel.
vagans, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 468 = hirta.
valdiviana, *Phil. in Linnaea*, xxxiii. (1864–65) 88.—Chili.
varians, Pohl, in DC. Prod. iv. 61 = quinqueloba.
verticillata, Thunb. Diss. Hydroc. 5 = vulgaris.
verticillata, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) ii. 28.—Austral.
villosa, *Linn. f. Suppl.* 175.—Afr. austr.
villosa, Thunb. ex Harv. & Sond. Fl. Cap. ii. 532 = H. Centella.
virgata, *Linn. f. Suppl.* 176.—Afr. austr.
Volkmanni, *Phil. in Linnaea*, xxxiii. (1864–65) 88.—Chili.
vulgaris, Bory, ex A. Rich. in Ann. Sc. Phys. iv. (1820) 172 = umbellata.
vulgaris, Cham. & Schlecht. in Linnaea, i. (1826) 356 (err. typ. 256), partim = interrupta.
vulgaris, *Linn. Sp. Pl.* 234.—Europ.
vulgaris, Michx. Fl. Bor. Am. i. 257 = interrupta.
vulgaris, Thunb. Fl. Cap. 252 = verticillata.
Wightiana, Wall. Cat. n. 7230 = asiatica.
zeylanica, DC. Prod. iv. 67 = javanica.
Zollingeri, Molk. in Mtq. Pl. Jungh. 91 = rotundifolia.

HYDROGETON, Lour. Fl. Cochinch. 244 (1790) = **Potamogeton**, Tourn. (Naiad.).
heterophyllum, Lour. l. c. = P. octandrum.

HYDROGETON, Pers. Syn. i. 400 (1805) = **Aponogeton**, Thunb. (Naiad.).
fenestralis, Pers. l. c. = A. fenestrale.

HYDROLAEA, Dum. Anal. Fam. 25 (1829) = seq.

HYDROLEA, Linn. Sp. Pl. ed. II. 328 (1763). *HYDROPHYLLACEÆ*, Benth. & Hook. f. ii. 881.
Reichelia, Schreb. Gen. 300 (1789).
Saponea, Aubl. Pl. Gui. i. 285, t. 111 (1775).
Steris, Linn. Mant. i. 8 (1767).
affinis, *A. Gray*, *Man. Bot. N. U. St.* ed. V. 370.—Am. bor.
arayatensis, Blanco, Fl. Filip. ed. I. 211 = zeylanica.
auriculata, Moç. ex Choisy, in DC. Prod. x. 181, 184 = Wigandia Kunthii.
Bartrami, Herb. Mus. Brit. ex Choisy, Desc. Hydrol. 15 = caroliniana.
caracasana, Steud. Nom. ed. I. 417 = Wigandia caracasana.
caroliniana, *Michx. Fl. Bor. Am.* i. 177.—Am. bor.
ceilonica, F. Muell. Fragm. v. 192 = zeylanica.
congesta, Willd. ex Choisy, in DC. Prod. x. 181, 182 = Nama undulata.
corymbosa, *Ell. Sketch*, i. 296.—Am. bor.
crispa, Ruiz & Pav. Fl. Per. iii. 22 = Wigandia crispa.
decurrens, Moç. ex Choisy, in DC. Prod. x. 181 = Nama jamaicensis.
demosperata, *Klotzsch*, *in Schomb. Faun. et Fl. Gui.* 1154, nomen.—Guian.

HYDROLEA:—
dichotoma, Ruiz & Pav. Fl. Per iii. 22. t. 244 = Nama dichotoma.
diffusa, *Raeusch. Nom. ed.* III. 76; *et ex Choisy*, *in DC. Prod.* x. 181.—Afr. austr.
elatior, Schott, in Spreng. Syst. iv. Cur. Post. 404 = multiflora.
elegans, *A. W. Benn. in Journ. Linn. Soc.* xi. (1871) 271.—Bras.
exaltata, Schott, ex Steud. Nom. ed. II. i. 782 = multiflora.
extra-axillaris, *Morr. in Ann. Soc. Bot. Gand*, ii. (1846) 221. t. 79.—Hab.?
floribunda, *Klotzsch & Peyr. Pl. Tinn.* 22. t. ix B.—Afr. trop.
glabra, *Choisy*, *Desc. Hydrol.* 16.—Venezuela.
glabra, *Schum. & Thonn. Beskr. Guin. Pl.* 161.—Afr. trop.
glabra, Sm. in Rees, Cyclop. xxi. Art. Lychnis = elegans.
graminifolia, *A. W. Benn. in Journ. Linn. Soc.* xi. (1871) 277.—Afr. trop.
guineensis, Choisy, in Ann. Sc. Nat. Sér. II. i. (1834) 180 = glabra.
inermis, Lour. Fl. Cochinch i. 172 = zeylanica.
jamaicensis, Raeusch. Nom. ed III. 76 = Nama jamaicensis.
javanica, Blume, Bijdr. 725 = zeylanica.
latifolia, *Raffn. Atl. Journ.* 146.—Am. bor.
leptocaulis, Featherman, Louisiana Univ. Rep. (1871) = affinis.
ludoviciana, Featherman, l. c. = ovata.
macrosepala, *A. W. Benn. in Journ. Linn. Soc.* xi (1871) 277.—Afr. trop.
madagascariensis, *Choisy*, *Desc. Hydrol.* 17.—Madag.
megapotamica, *Spreng. Syst. iv. Cur. Post.* 114.—Bras.
mollis, Willd. ex Spreng. Syst. i. 866 = Wigandia caracasana.
monteviadensis, *Choisy*, *in DC. Prod.* x. 181.—Reg. Argent.
multiflora, *Choisy*, *l. c.* 180.—Bras.
nigricaulis, *Wright*, *ex Griseb. Cat. Pl. Cub.* 207.—Cuba.
ovata, *Nutt. in Trans. Am. Phil Soc.* N. S. v. (1837) 196.—Am. bor. et austr.
ovatifolia, Raffn. New Fl. Am. iv. 64 = ovata.
paludosa, *A. W. Benn. in Journ. Linn. Soc.* xi. (1871) 270.—Bras.
palustris, Raeusch. Nom. ed. III. 76 = caroliniana.
paniculata, Raffn. New Fl. Am. iv. 84 = caroliniana.
quadrivalvis, Walt. Fl. Carol. i. 109, 110 = caroliniana.
radians, Moç. ex Choisy, in DC. Prod. x. 181, 182 = Nama undulata.
rupincola, Moç. ex Choisy, l. c. 181 = Nama undulata.
scorpioides, Moç. ex Choisy, l. c. 182, 184 = Wigandia scorpioides.
spinosa, *Linn. Sp Pl.* 238.—Am. trop.
tenella, Moç. ex Choisy, in DC. Prod. x. 182, 183 = Nama origanifolia.
trigyna, Sw. Prod. Veg. Ind. Occ. 54 = spinosa.
urens, Ruiz & Pav. Fl. Per. iii. 21 = Wigandia urens.
verticillata, *Raffn. Fl. Ludov.* 51.—Am. bor.
violacea, Moç. ex Choisy, in DC. Prod. x. 182 = Nama sericea.
zeylanica, *Vahl*, *Symb. Bot.* ii. 46.—Reg. trop.

HYDROLIA, Thou. Gen. Nov. Madag. 9 (1806) = prasc.

HYDROLYTHRUM, Hook. f. in Benth. & Hook. f. Gen. i. 777 (1867). *LYTHRARIEÆ*, Benth. & Hook. f. i. 777.
Wallichii, *Hook. f. l. c.*; *et in Hook. Ic. Pl.* t. 1007.—Ind. or.

HYDROMESTES, Benth. & Hook. f. Gen. ii. 1103 (1876) = seq.

HYDROMESTUS, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 285 = **Aphelandra**, R. Br. (Acanthac.).
maculatus, Scheidw. l. c. = A. Hydromestus.

HYDROMISTRIA, Bartl. Ord. 74 (1830) = seq.

HYDROMYSTRIA, G. F. W. Mey. Prim. Fl. Esseq. 152 (1818) = **Limnobium**, Rich. (Hydrocharid.).
stolonifera, G. F. W. Mey. l. c. 153 = L. Bonell.

HYDROPELTIS, Michx. Fl. Bor. Am. i. 232. t. 22 (1803) = **Brasenia**, Schreb. (Nymphaeac.).
fulla, Salisb. in Kon. & Sims, Ann. Bot. ii. (1807) 74 = B. Schreberi.
purpurea, Michx. Fl. Bor. Am. i. 324. t. 29 = B. Schreberi.

HYDROPHACA, Steud. Nom. ed. II. i. 783 (1840) = seq.

HYDROPHACE, Hall. Enum. Stirp. Helv. iii. 66 (1768) = Lemna, Linn.

HYDROPHYLAX, Linn. f. Suppl. 120 (1781). *RUBIACEÆ*, Benth. & Hook. f. ii. 142.
Sarissus, Gaertn. Fruct. i. 118. t. 25 (1788).
carnosa, *Sond. in Harv. & Sond. Fl. Cap.* iii. 25.—Afr. austr.
madagascariensis, *Willd ex Roem. & Schult. Syst.* iii 527.—Madag.
maritima, *Linn. f. Suppl.* 126.—Ind. or.
pusilla, Pohl, ex DC. Prod. iv. 550 (= *Borreria eupatorioides*).—Bras.

HYDROPHYLLAX, Raffn. in Ann. Gén. Sc. Phys. vi (1820) 87 = praec.

HYDROPHYLLUM, Linn. Syst. ed. I (1735, *HYDROPHYLLACEÆ*, Benth. & Hook. f. ii. 926.
Deckmium, Raffn. Fl. Ludov. 84, in obs. (1817).
Viticella, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 220.
acrifolium, Salisb. Prod. 120 = canadense.
Aldea, Roem. & Schult. Syst. iv. 114 = Phacelia circinnata.
appendiculatum, *Michx. Fl. Bor. Am.* i. 134.—Am. bor.
canadense, *Linn. Sp. Pl.* ed. II. 208.—Am. bor.
capitatum, *Dougl. ex Benth. in Trans. Linn. Soc.* xvii. (1836) 273.—Am. bor. occ.
cochinchinense, Lour. ex Steud. Nom. ed. II. i. 788 (Quid ?).—Cochinch.
hispidum, *Riddell. Syn. Fl. W. St. Am.* 84.—Am. bor.
lineare, Pursh, Fl. Am. Sept. i. 134 = Phacelia Menziesii.
lobatum, Stokes, Bot. Mat. Med. i. 302 = canadense.
macrophyllum, *Nutt. in Journ. Acad. Philad.* vii. (1834) 111.—Am. bor.
magellanicum, *Lam. in Journ. Hist. Nat. Par.* i. (1792) 378.—Reg. Magellan.
occidentale, *A. Gray*, *in Proc. Am. Acad.* x. (1875) 314.—Am. bor. occ.
paniculatum, Poepp. ex A. DC. in DC. Prod. ix. 298. 289 = Phacelia circinata.
pinnatifidum, Moench, Meth. 428 = virginicum.
pusillum, Muhl. ex A. DC. in DC. Prod. ix. 289, 292 = Nemophila microcalyx.
spiraeaefolium, Salisb. Prod. 120 = virginicum.
triloburn, Raffn. Fl. Ludov. 83 = appendiculatum.
virginianum, Linn. Sp. Pl. 146 = seq.
virginicum, *Linn. Syst. ed.* XII. 152.—Am. bor.

HYDROPHYTUM, Flora, vi. (1823) Beibl. 96, sphalm. = **Hydrophytum**, Jack (Rubiac.).

HYDROPIPER, Buxb. ex Fourr. in Ann. Soc. Linn. Lyon. N. S. xvi. (1868) 349 = **Blatina**, Linn.
Fabri, Fourr. l. c. = E. macropoda.
hexandrum, Fourr. l. c. = E. hexandra.
majus, Fourr. l. c. = E. hexandra.

HYDROPITYON, Gaertn. f. Fruct. iii. 19. t. 168 (1805) = **Limnophila**, R. Br. (Scrophulariin.).
pedunculatum, Ser. in DC. Prod. i. 422 = L. gratioloides.
zeylanica, Gaertn. f. Fruct. iii. 19. t. 168. f. 2 = L. gratioloides.

HYDROPITYUM, Steud. Nom. ed. II. i. 788 (1840) = praec.

HYDROPYRUM, Link, Hort. Berol. i. 252 (1827) = **Eisantia**, Gronov. (Gramin.).
aculeatum, Link, l. c. = Z. aquatica.
fluitans, Kunth, Rév. Gram. i. 7 = Z. aquatica.
latifolium, Griseb. in Ledeb. Fl. Ross. iv. 466 = Z. aquatica.

HYDROPYXIS, Rafin. Fl. Ludov. 94 (1817); DC. Prod. iii. 364 (Geu. delendum). Benth. & Hook. f. i. 186.
palustris, Rafin. l. c.

HYDROSCHOENUS, Zoll. & Mor. Syst. Vers. Zoll. 95 (1846) = **Cyperus**, Michx.
bytlingioides, Zoll. & Mor. l. c. = C. Cephalotes.

HYDROSIA, A. Juss. in Orb. Dict. vii. 270 (1849) = HIDROSIA, E. Mey. = **Rhynchosia**, Lour. (Legumin.).

HYDROSME, Schott, in Oestr. Bot. Wochenbl. (1857) 389 = **Amorphophallus**, Blume (Aroid.).
angolensis, Schott, in Journ. Bot. iii. (1865) 85.—Afr. trop.
consimilis, Engl. in DC. Monog. Phan. ii. 325 = A. consimilis.
Eickleri, Engl. in Eichl. Jahrb. Berl. ii. (1888) 285.—Afr. trop.
Fontanesii, Schott, Prod. Aroid. 188 = A. Fontanesii.
grata, Engl. in DC. Monog. Phan. ii. 328.—Afr. trop.
Hildebrandtii, Engl. in Bot. Jahrb. i. (1881) 187.—Afr. trop.
leonensis, Engl. l. c. = A. leonensis.
maximo, Engl. in DC. Monog. Phan. ii. 328.—Afr. trop.
mossambicensis, Schott, Gen. Aroid. 33 = A. mossambicensis.
Prieuriana, Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 279 = A. consimilis.
Riviori, Engl. in Bot. Jahrb. i. (1881) 188 = A. Konjac.
Schweinfurthii, Engl. in DC. Monog. Phan. ii. 322.—Afr. trop.
Teuszii, Engl. in Regel, Gartenfl. (1884) 2; et Bot. Jahrb. v. (1884) Beibl. n. 6.—Afr. trop.

HYDROSPONDYLUS, Hassk. in Flora (1842), Beibl. 33 = **Hydrilla**, Rich. (Hydrocher.).
submersus, Hassk. l. c. = Hydrilla verticillata.

HYDROSTACHYS, Thou. Gen. Nov. Madag. 2 (1806). *PODOSTEMACEAE*. Benth. & Hook. f. iii. 115.
cristulata, *Wedd. in DC. Prod.* xvii. 89.—Afr. trop.
distichophylla, *A. Juss. in Deless. Ic. Sel.* iii. t. 94.—Madag.
Goudotiana, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 93.—Madag.
imbricata, *A. Juss. in Deless. Ic. Sel.* iii. t. 92.—Madag.
madagascariensis, Steud. Nom. ed. II. i. 788 = Goudotiana, imbricata, verruculosa.
multifida, *A. Juss. in Deless. Ic. Sel.* t. 93.—Madag.
natalensis, *Wedd. in DC. Prod.* xvii. 88.—Afr. austr.
plumosa, *A. Juss. ex Tul. in Arch. Mus. Par.* vi. (1852) 87.—Madag.
polymorpha, *Klotzsch, in Peters, Reise Mossamb. Bot.* 506. t. 52.—Afr. trop.
stolonifera, *Baker, in Journ. Linn. Soc.* xxi. (1885) 485.—Madag.
verruculosa, *A. Juss. in Deless. Ic. Sel.* iii. t. 91.—Madag.

HYDROSTIS, Reichb. in Moesal. Handb. ed. II. i. p. lx (1827) = **Hydrastis**, Ellis (Ranunc.).

HYDROTAENIA, Lindl. Bot. Reg. (1838) Misc. 69. *IRIDEAE* [Benth. & Hook. f. iii. 690].
PARDINIA, Herb. in Lindl. Bot. Reg. (1844) Misc. 66, in obs. partim.
Ehrenbergii, *Schlecht. in Linnaea*, xviii. (1844) 666.—Mexic.
lobata, *Herb. in Lindl. Bot. Reg.* (1844) *Misc.* 66.—Peruv.
Melengris, *Lindl. l. c.* (1838) *Misc.* 69.—Mexic.

HYDROTICHE, A. Juss. in Orb. Dict. xi. (1849) 459 = seq.

HYDROTRICHE, Zucc. in Abh. Acad. Muench. (1832) 808. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 951.
hottoniaeflora, *Zucc. l. c.* 309.—Madag.

HYDROTRIDA, Willd. ex Schlecht. & Cham. in Linnaea, v. (1830) 107 = **Herpestis**, Gaertn. f. (Scrophular.).
Beccabunga, Willd. l. c. = Herpestis obovata.

HYDROTROPHUS, C. B. Clarke, in Journ. Linn. Soc. xiv. (1873) 8. t. 1 = **Hyxa**, Noronha (Hydrochar.).
echinospermus, C. B. Clarke, l. c.—Ind. or.

HYERONIMA, Allem. Disa. in Trab. Vell. Rio Janeiro (1848). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 284.
SECRETANIA, Muell. Arg. in DC. Prod. xv. ii. 227 (1866).
SYLLAGINELLA, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 340.
alchorneoides, *Allem. Diss. in Trab. Vell. Rio Janeiro* (1848).—Bras.
Blanchetiana, *Tul. in Mart. Fl. Bras.* iv. i. 333.—Bras.
clusioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 157.—Porto Rico.
cubana, *Muell. Arg. in DC. Prod.* xv. ii. 270.—Cuba.
ferruginea, *Tul. in Mart. Fl. Bras.* iv. i. 334. t. 91.—Bras.
laxiflora, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 67.—Guian.
macrocarpa, *Muell. Arg. l. c.* 66.—Venezuela.
mollis, *Muell. Arg. in DC. Prod.* xv. ii. 269.—Bras.
oblonga, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 66.—Guian.
palida, *Muell. Arg. in DC. Prod.* xv. ii. 270.—Cuba.
scabra, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 66.—N. Granat.

HYGEA, Klotzsch, in Schomb. Faun. et Fl. Brit. Gui. 1147, nomen (1848). *ASCLEPIADEAE*.
subulata, *Klotzsch, l. c.* nomen.—Guian.

HYGEA, Hanst. in Linnaea, xxvi. (1858) 188, 209. *GESNERIACEAE*!
barbigera, *Hanst. l. c.*—Chili.

HYGROCHARIS, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 77 (1851) = **Nephrophyllum**, A. Rich. (Hydrochar.).
abyssinica, Hochst. ex A. Rich. l. c. = N. abyssinicum.

HYGROCHARIS, Nees, in Mart. Fl. Bras. ii. 1. 147, adnot. (1842) = **Eynohospora**, Vahl (Cyperac.).

HYGROPHILA, R. Br. Prod. 479 (1810). *ACANTHACEAE*, Benth. & Hook. f. ii. 1075.
ASTERACANTHA, Nees, in Wall. Pl. As. Rar. iii. 75 (1832).
BAHEL, Adans. Fam. ii. 210 (1763).
EBERLEA, Riddell, ex Nees, in DC. Prod. xi. 85 (1847).
HEMIADELPHIS, Nees, in Wall. Pl. As. Rar. iii. 75 (1832).
PHYSICHILUS, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 310.
POLYECHMA, Hochst. in Flora, xxiv. (1841) 876.
TENORIA, Dehnh. & Glord. Cat. Pl. Hort. Camald. ed. II. (1829) 124.
abyssinica, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 29.—Abyssin.
acutangula, Nees, ex Mart. in Flora, xxiv. (1841) 11. Beibl. 26 = costata.
angustifolia, *R. Br. Prod.* 479.—Geront. trop.
asperrima, Nees, in DC. Prod. xi. 90 = angustifolia.
balsamica, Rafin. Fl. Tellur. iv. 66 = Cardanthera balsamica.
barbata, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. trop.
barbata, Nees, ex Steud. Nom. ed. II. i. 783 = quadrivalvis.
brasiliensis, *Spreng. Syst. iv. Cur. Post.* 287.—Bras.
caerulea, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 24.—Afr. trop.
ciliaris, Nees, in DC. Prod. xi. 156, in syn. = angustifolia.
conferta, *Nees, in Mart. Fl. Bras.* ix. 21.—Bras.
costata, *Sinning, in Nees & Sinning, Amoen. Bot. Bonn.* ii. (1834) 7. t. 3.—Am. trop.
difformis, Blume, Bijdr. 804 = Ruellia difformis.
dimidiata, Nees, in Wall. Pl. As. Rar. iii. 81 = angustifolia.
formosa, Spreng. Syst. ii. 836 = Aphelandra formosa.

HYGROPHILA:—
glandulifera, *Nees, in Mart. Fl. Bras.* ix. 22.—Bras.
guianensis, *Nees, ex Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 634.—Guian.
Helodes, *Nees, in Mart. Fl. Bras.* ix. 23.—Bras.
hirsuta, *Nees, l. c.* 24.—Bras.
hispida, *Nees, in DC. Prod.* xi. 87.—Cuba.
illinoiensis, Wood, in Bull. Torrey Bot. Club, v. 41; ex A. Gray, Syn. Fl. N. Am. ii. 1. 327 = Ruellia strepens.
incana, *Nees, in DC. Prod.* xi. 91.—Ind. or.
lacustris, *Nees, l. c.* 88.—Am. centr.
lances, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 123.—Japon.
latifolia, *Nees, in Mart. Fl. Bras.* ix. 24.—Bras.
longifolia, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 78 = spinosa.
longifolia, *Nees, in Mart. Fl. Bras.* ix. 21.—Mexic.
lutea, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 22.—Afr. trop.
malabarica, Rafin. Fl. Tellur. iv. 66 = angustifolia.
micrantha, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 22.—Afr. trop.
oblongifolia, Nees, in Mart. Fl. Bras. ix. 21 = costata.
oblongifolia, Rafin. Fl. Tellur. iv. 66 = Ruellia ciliosa?
obovata, Griff. Notul. iv. 134 = phlomoides.
obovata, Nees, in Wall. Pl. As. Rar. iii. 81 = angustifolia.
obovata, Wight, Ic t. 1489 = quadrivalvis.
odora, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 22.—Afr. trop.
ovata, *Nees, in DC. Prod.* xi. 89.—Bras.
Pavala, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 78 = Hemigraphis latebrosa.
phlomoides, *Nees, in Wall. Pl. As. Rar.* iii. 80.—Ind. or.; Burma.
polysperma, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 456.—Ind. or.; Malaya.
portoricensis, *Nees, in DC. Prod.* xi. 92.—Porto Rico.
pubescens, *Nees, l. c.* 88.—Bras.
pustila, *Blume, ex Steud. Nom.* ed. II. i. 783.—Ins. Moluce.
quadrivalvis, *Nees, in Wall. Pl. As. Rar.* iii. 80.—Ind. or.; Malaya.
radicans, Nees, l. c. 81 = angustifolia.
rengasa, R. Br. ex Steud. Nom. ed. II. i. 783 = angustifolia.
rivularis, *Nees, in DC. Prod.* xi. 87.—Mexic.
salicifolia, Nees, in Wall. Pl. As. Rar. iii. 81 = angustifolia.
Schottiana, *Nees, in Mart. Fl. Bras.* ix. 22.—Bras.
senegalensis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 22.—Afr. trop.
Serpyllum, *T. Anders. l. c.* ix. (1867) 456.—Ind. or.
spinosa, *T. Anders. l. c.* vii. (1864) 22.—Ind. or.; Malaya.
Stocksii, *T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 407.—Ind. or.
stricta, Hassk. Cat. Hort. Bog. Alt. 148 = angustifolia.
uliginosa, *S. Moore, in Journ. Bot.* xviii. (1880) 197.—Afr. trop.
undulata, Blume, Bijdr. 803 = angustifolia.
undulata, Nees, in Wall. Pl. As. Rar. iii. 80, in syn. = quadrivalvis.

HYGRORHIZA, Benth. in Journ. Linn. Soc. xix. (1881) 55 = seq.

HYGRORYZA, Nees, in Edinb. N. Phil. Journ. xv. (Apr.–Oct. 1833) 350. *GRAMINEAE*, Benth. & Hook. f. iii. 1116.
POTAMOCHLOA, Griff. in Journ. As. Soc. Beng. v. (1836) 571. t. 22.
aristata, *Nees, in Edinb. N. Phil. Journ.* xv. (Apr.–Oct. 1833) 350.—Ind. or.
ciliata, Steud. Nom. ed. II. i. 783 = Zizania ciliata.

HYLACIUM, Beauv. Fl. Owar. ii. 83. t. 113 (1807) = **Psychotria**, Linn. (Rubiac.).
owariense, Beauv. l. c. = P. owariensis.

HYLAS, Bigel. ex DC. Prod. iii. 69, in obs. (1828) = **Myriophyllum**, Linn. (Halorag.).

HYLEBIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 247 = **Stellaria**, Linn. (Caryophyll.).
nemorum, Fourr. l. c. = S. nemorum.

HYLENAEA, Miers, in Trans. Linn. Soc. xxviii. (1872) 366 = **Hippocratea**, Linn. (Celastr.).
capillaeflora, Miers, l. c. 368 = Hippocratea comosa.
comosa, Miers, l. c. 367 = Hippocratea comosa.
jamaicensis, Miers, l. c. 369 = Hippocratea ovata.
multiflora, Miers, l. c. 368 = Salacia multiflora, *DC.*

HYLETHALE, Link, Handb. i. 788 (1829) = **Frenanthes**, Linn. (Compos.).
purpurea, Link, l. c. = F. purpurea.

HYLINE, Herb. in Bot. Mag. sub t. 3779 (1840). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 730.
Cardenasiana, *Herb. l. c.*—Bras.

HYLOCHARIS, Miq. Fl. Ind. Bat. Suppl. 319 (1860) = **Oxypora**, DC. (Melastom.).
macrophylla, Miq. l. c. = O. macrophylla.

HYLOCHARIS, Tilling, ex Regel & Tilling, in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 123 = **Clintonia**, Rafin. (Liliac.).
cyanocarpa, Tilling, l. c. = C. udensis.

HYLOCOCCUS, R. Br. ex Benth. Fl. Austral. vi. 92 (1873) = **Petalostigma**, F. Muell. (Euphorb.).
sericeus, R. Br. ex Benth. l. c. = P. quadriloculare.

HYLOGETON, Salisb. Gen. Pl. Fragm. 91 (1866) = **Allium**, Tourn. (Liliac.).
ursinum, Salisb. l. c. = A. ursinum.

HYLOGYNE, Salisb. in Knight, Prot. 126 (1809) = **Telopea**, R. Br. (Proteac.).
australis, Salisb. l. c. 127 = T. truncatum.
speciosa, Salisb. l. c. 126 = T. spectosissimum.

HYLOMECON, Maxim. in Mém. Sav. Etr. Acad. Pétersb. ix. (1858); (Prim. Fl. Amur.) 36. t. 3 = **Stylophorum**, Nutt. (Papaverac.).
vernalis, Maxim. l. c. = S. japonicum.

HYLOMENES, Salisb. Gen. Pl. Fragm. 26 (1866) = **Sotilla**, Linn. (Liliac.).
campanulata, Salisb. l. c. = S. campanulata.
non-scripta, Salisb. l. c. = S. festalis.

HYLONOME, Webb & Berth. Phyt. Canar. iii. 320 (1850), partim; Baker, in Journ. Linn. Soc. xiv. (1875) 561 = **Bohnia**, Didr. (Liliac.).
reticulata, Webb & Berth. l. c.; Baker, l. c. = B. reticulata.

HYLOPHILA, Lindl. in Wall. Cat. n. 7396 (1831); et Bot. Reg. sub t. 1618 (1833). *ORCHIDEAE*, Benth. & Hook. f. iii. 602.
DICROGYLIS, Blume, Orch. Archip. Ind. i. 116 (1858).
lanceolata, *Miq. Fl. Ind. Bat.* iii. 746.—Malaya.
mollis, *Lindl. in Wall. Cat.* n. 7396; *et Bot. Reg.* sub t. 1618.—Malaya.
purpurea, Miq. Fl. Ind. Bat. iii. 746 = Zeuxine purpurascens.

HYMANTHOGLOSSUM, Tod. Orch. Sic. 67 (1842) = HIMANTOGLOSSUM, Spreng. = **Orchis**, Linn.

HYMENACHNE, Beauv. Agrost. 48. t. 10. f. 8 (1812) = **Panicum**, Linn. (Gramin.).
amplexicaulis, Nees, Agrost. Bras. 276 = P. Myurus.
Boiviniana, Steud. l. c. 419.—Madag.
campestris, Nees, Agrost. Bras. 275, 274 = P. camporum.
camporum, Kunth, Enum. Pl. i. 88 = P. vilvoides.
fluviatilis, Nees, Agrost. Bras. 275 = P. vilvoides.
frondescens, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 496 = P. stoloniferum.
Gouini, Fourn. ex Hemsl. l. c. 492 (Quid?).—Mexic.
indica, Busse, in Miq. Fl. Jungh. 377 = P. indicum.
interrupta, Busse, l. c. = P. Myurus.
interrupta, Stend. Syn. Pl. Gram. 101 (an praec.?).—Java.
leptostachya, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 490 = P. leptostachyum.
montana, Griseb. in Goett. Abh. xxiv. (1879) 307.—Reg. Argent.
Myosurus, Nees, Agrost. Bras. 276 = P. Myosurus.

HYMENACHNE:—
Myurus, Beauv. Agrost. 49 = Panicum Myurus.
Myurus, Steud. Syn. Pl. Gram. 101 = Sporobolus ciliatus?
patula, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 492 (Quid?).—Mexic.
phalaroides, Nees, Agrost. Bras. 276 = P. phalaroides.
phleiformis, Fourn. ex Hemsl. Biol. Centr. Centr. Am. Bot. iii. 494 = P. phleiforme.
Pseudo-interrupta, C. Muell. in Bot. Zeit. xix. (1861) 333 = P. Myurus.
striata, Griseb. Fl. Brit. W. Ind. 554 = P. striatum.

HYMENAEA, Linn. Gen. ed. 1. 366 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 583
COURBARI, Adans. Fam. ii. 317 (1763).
Courbaril, Plum ex Endl. Gen. 1317 (1841).
TAVROUJOU, Jus. Gen. 351, in adnot. (1789).
animifera, Stokes, Bot. Mat. Med. ii. 449 = H. Courbaril.
Candolliana, *H. B. & K. Nov. Gen. et Sp.* vi. 328 t. 566.—Mexic.
chrysophylla, Willd. ex Vog. in Linnaea, xiii. (1839) 518 = Bauhinia speciosa.
confertifolia, *Mart. ex Hayne, Arzneigew.* xi. t. 8 — Bras.
confertifolia, Hayne, l. c. t. 9 = stilbocarpa.
cupulifera, Mart. ex Hayne, l. c. sub t. 15 = Martiana.
Courbaril, *Linn. Sp. Pl.* 1192.—Am. austr
eriogyne, *Benth. in Mart. Fl. Bras.* xv. ii. 237.—Bras.
floribunda, *H. B. & K. Nov. Gen. et Sp.* vi. 328. t. 567 —Guian.
Gaertneriana, D. Dietr. Syn. Pl. ii. 1473 = Trachylobium Hornemannianum.
Hayneana, D. Dietr. l. c. = verrucosa.
Hornemanniana, D. Dietr. l. c. = Trachylobium Hornemannianum.
Lamarchiana, D. Dietr. l. c. = verrucosa.
latifolia, Hayne, Arzneigew. xi. t. 7 = Peltogyne latifolia.
Martiana, *Hayne, l. c.* t. 15.—Bras.
obtusifolia, Willd. ex Hayne, l. c. xi. sub t. 7 = Peltogyne latifolia.
Oliveriana, *Hayne, l. c.* t. 14.—Bras.
pubescens, Mart. ex Hayne, l. c. sub t. 16 = Sellowiana.
resinifera, Salisb. Prod. 327 = H. Courbaril.
retusa, Willd. ex Hayne, Arzneigew. xi. subt. 12 = Candolliana.
rotundata, *Hayne, l. c.* t. 13.—Am. austr.
Sellowiana, *Hayne, l. c.* t. 16.—Bras.
splendida, *Vog. in Linnaea*, xi. (1837) 409.—Bras
stigonocarpa, *Mart. ex Hayne, Arzneigew.* xi. t. 18.—Bras.
stilbocarpa, *Hayne, l. c.* t. 11.—Bras.
venosa, *Vahl, Eclog. Am.* ii. 31.—Guian.
verrucosa, *Gaertn. Fruct.* ii. 306. t. 189. f. 7.—Madag.
verrucosa, Hornem. ex Hayne, Arzneigew. xi. sub t. 18 = Trachylobium Hornemannianum.

HYMENANDRA, A. DC. ex Spach, Hist. Vég. Phan. ix. 474 (1840). *MYRSINEAE*, Benth. & Hook. f. ii. 647.
Wallichii, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 84.—Reg. Himal.

HYMENANTHERA, R. Br. in Tuckey, Congo, 442 (1818). *VIOLARIEAE*, Benth. & Hook. f. i. 120.
SOLENANTHA, G. Don, Gen. Syst. ii. 89 (1832).
angustifolia, R. Br. ex DC. Prod. i. 315 = dentata.
Banksii, F. Muell. First Gen. Report, 9 = dentata.
crassifolia, *Hook. f. Fl. N. Zel.* i. 7.—N. Zel.
dentata, *R. Br. ex DC. Prod.* i. 315.—Austral.
latifolia, *Endl. Prod. Fl. Norfolk.* 127.—Ins. Norfolc.
oblongifolia, A. Cunn. in Hook. Lond. Journ. Bot. i. (1842) 194 = dentata.
Traversii, *Y. Buch. in Trans. N. Z. Inst.* xv. (1883) 389.—N. Zel.

HYMENANTHES, Blume, Bijdr. 862 (1826) = **Rhododendron**, Linn. (Ericac.).
japonica, Blume, l. c. = R. Metternichii.

HYMENANTHUS, D. Dietr. Syn. Pl. iii. 21 (1848) = prsec.

HYMENATHERUM, Cass. in Bull. Soc. Philom. (1817) 12. *COMPOSITAE*, Benth. & Hook. f. ii. 410.
ACIPHYLLAEA, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) (Pl. Fendl.) 91.
GNAPHALOPSIS, DC. Prod. vii. 258 (1838).
acerosum, *A. Gray, Pl. Wright.* i. 115.—Mexic.
aureum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 42 —Am. bor. occ.
Belenidium, DC. Prod. vii. 292 = tenuifolium.
Berlandieri, Benth. Pl. Hartw. 18 = Hartwegi.
Berlandieri, DC. Prod. v. 642 = pentachaetum.
boeberoides, *A. Gray, Pl. Wright.* i. 115.—Mexic.
Candollianum, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 320 = tenuifolium.
concinnum, *A. Gray, Syn. Fl. N. Am.* i. ii. 446, 453. —Arizona.
diffusum, *A. Gray, Pl. Wright.* i. 116.—Mexic.
gnaphaloides, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 90 = H. Gnaphalopsis.
Gnaphalopsis, *A. Gray, l. c.* 115.—Mexic
Greggii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 42 — Mexic
Hartwegi, *A. Gray, Pl. Wright.* i. 116, 117.—Mexic
Kunthii, *Less. Syn. Comp.* 286.—Chili.
Nesei, *DC. Prod.* iv. 642.—Chili.
neo-mexicanum, *A. Gray, in Proc. Am. Acad.* xix (1883) 40.—Mexic.
pentachaetum, *DC. Prod.* v. 642.—Mexic.
polychaetum, *A. Gray, Pl. Wright.* i. 116.—Mexic.
serratifolium, *Hemsl. Biol. Centr. Am. Bot.* ii. 221.—Mexic.
sessilifolium, *A. Gray, in Proc. Am. Acad* xix. (1885) 42.—Mexic.
tagetoides, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 88.—Texas.
tenuifolium, *Cass. in Dict. Sc. Nat.* xxii. 314.—Chili? Mexic.
tenuilobum, DC. Prod. v. 642 = tenuifolium
Thurberi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 41. —Mexic.
Treculii, *A. Gray, l. c.* 42.—Texas.
Wrightii, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 89.—Texas.

HYMENELLA, [Moç. & Sesse,] ex DC. Prod. i. 389 (1824). *CARYOPHYLLEAE*, [Benth. & Hook. f. i. 150]
TRIPLATEIA, Bartl. in Presl, Rel. Haenk. ii. 11. t. 50 (1830).
moehringioides, [*Moç. & Sesse,*] *ex DC. Prod.* i. 390.—Mexic.

HYMENESTHES, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 26. t. 6 = **Bourreria**, P. Br. (Boragin.).
nitida, Miers, l. c. = B. succulenta.

HYMENETRON, Salisb. Gen. Pl. Fragm. 128 (1866) = **Strumaria**, Jacq. (Amaryllid.).
lingusefora, Salisb. l. c. = S. truncata.
truncata, Salisb. l. c. = S. truncata.

HYMENIA, Griff. Notul. iv. 449 (1854) = **Hymenaea**, Linn. (Legumin.).

HYMENIDIUM, Lindl. in Royle, Illustr. Bot. Himal. 238 (1858) = **Pleurospermium**, Hoffm. (Umbellif.).
Brunonis, Lindl. l. c. = P. Brunonis.
densiflorum, Lindl. l. c. = P. densiflorum.
suaveolens, Klotzsch, in Bot. Ergeb. Waldem. Reise, 148 = P. densiflorum.

HYMENOCALLIS, Salisb. Trans. Hort. Soc. i. (1812) 388. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 734.
CHOLETIS, Herb. Amaryll. 219. f. 85 (1837).
ISMENE, Salisb. in Trans. Hort. Soc. i. (1812) 342.
HYMENOCALYX, Hoolk. in Rev. Hortic. (1869) 418.
NEMEPIODON, Rafin. Fl. Tellur. iv. 22 (1836).
SIPHOTOMA, Rafin. l. c. (1836).
TOMODON, Rafin. l. c. (1836).
acutifolia, Mi. Roem. Syn. Ensat. 174 = littoralis.
adnata, Herb. Amaryll. 218 = littoralis.
Amancaes, *Nichols. Dict. Gard.* ii. 165.—Peruv.
americana, M. Roem. Syn. Ensat. 176 = littoralis.
amœna, Herb. App. 44 = ovata.
Andreana, *Nichols. Dict. Gard.* ii. 165.—Ecuador.
angusta, Herb. App. 44 = caribaea.
arenaria, M. Roem. Syn. Ensat. 176 = littoralis.

HYMENOCALLIS:—
biflora, *C. Koch & Bouché, in C. Koch, Wochenschr.* ix. (1866) 370.—Bras.
bistubata, *Herb. in Bot. Reg.* (1844) *Misc.* 63.—Mexic.
Bonplandii, *Kunth, Enum. Pl.* v. 666.—N. Granat.
Borbhiana, De Vriese, in Nederl. Kruidk. Arch. i. (1846) 156 = seq.
Boschiana, Kunth, Enum. Pl. v. 854 = undulata.
calathina, *Nichols. Dict. Gard.* ii. 165.—Peruv.
caribæa, *Herb. App.* 44.—Ind. occ.
carolinianæ, Herb. l. c. = Pancratium maritimum.
caymanensis, *Herb. Amaryll.* 614.—Florida.
Choretis, Hemsl. Biol. Centr. Am. Bot. iii. 335 = glauca.
cinerascens, M. Roem. Syn. Enast. 171 = caribæa.
coronaria, Kunth, Enum. Pl. v. 855 = crassifolia.
crassifrons, Kunth, l. c. 677, sphalm. = seq.
crassifolia, *Herb. App.* 44.—Am. bor.
declinata, M. Roem. Syn. Enast. 171 = caribæa.
Dillenii, M. Roem. l. c. 174 = lacera.
disciformis, M. Roem. l. c. 178 = lacera.
distichæ, Herb. App. 44 = littoralis.
Dryanderi, M. Roem. Syn. Enast. 175 = littoralis.
eucharidifolia, *Baker, in Gard. Chron.* (1884) i. 709. —Am. trop.
expansa, *Herb. App.* 44.—Ind. occ.
formosa, M. Roem. Syn. Enast. 171 = speciosa.
fragrans, Salisb. in Trans. Hort. Soc. i. (1812) 340 = caribæa.
galvestonensis, *Baker*, (*in Herb. Kew.*)—Am. bor.
glauca, *M. Roem. Syn. Enast.* 173.—Mexic.
guianensis, Herb. App. 44 = tubiflora.
Harrisiana, *Herb. in Bot. Reg.* (1840) *Misc.* 35.—Mexic.
humilis, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 301. —Florida.
insignis, Kunth, Enum. Pl. v. 675 = littoralis.
lacera, *Salisb. in Trans. Hort. Soc.* i. (1812) 338.—Am. bor.
latifolia, M. Roem. Syn. Enast. 168 = Urceolina latifolia.
littoralis, *Salisb. in Trans. Hort Soc.* i. (1812) 338.—Am. austr.
lobata, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xi. (1843) 124.—Venezuela.
Macleana, *Nichols. Dict. Gard.* ii. 165.—Peruv.
macrostephana, *Baker, in Gard. Chron.* (1879) i. 430.—Hab.?
maritima, M. Roem. Syn. Enast. 176 = Pancratium maritimum.
mexicana, Herb. App. 44 = lacera.
Milleri, M. Roem. Syn. Enast. 177 = lacera.
Moritziana, *Kunth, Enum. Pl.* v. 668.—Venezuela.
obtusata, Griseb. ex Walp. Ann. iii. 618 = caribæa?
occidentalis, Kunth, Enum. Pl. v. 856 = crassifolia.
ornata, M. Roem. Syn. Enast. 169 = ovata?
ovalifolia, Herb. Amaryll. 212 = ovata.
ovata, *M. Roem. Syn. Enast.* 169.—Ind. occ.
Palmeri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 301. —Florida.
paludosa, Salisb. in Trans. Hort. Soc. i. (1812) 338 = lacera.
panamensis, Lindl. Bot. Reg. (1841) Misc. 67 = littoralis.
patens, Herb. App. 44 = caribæa?
pedalis, *Herb. l. c.*—Bras.
peruviana, M. Roem. Syn. Enast. 176 = littoralis.
petiolata, M. Roem. l. c. 168 = tubiflora.
quitensis, Stend. Nom. ed. II. i. 784 = seq.
quitoënsis, *Herb. App.* 44.—Ecuador.
repanda, *Otto & Dietr. Allg. Gartenz.* xi. (1843) 128. —Mexic.
rotata, Herb. App. 44 = lacera.
Ruizii, M. Roem. Syn. Enast. 177 = Pancratium maritimum.
senegambica, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 13.—Afr. trop.
sessilis, Salisb. in Trans. Hort. Soc. i. (1812) 339 = caribæa.
Skinneriana, *Herb. in Bot. Reg.* (1848) *Misc.* 45.—Guatemala.
Sloanei, M. Roem. Syn. Enast. 171 = caribæa.
speciosa, *Salisb. in Trans. Hort. Soc.* i. (1812) 340.—Ind. occ.
spiralis, Lee, ex Herb. App. 46 = ovata.
Staplesi, Sweet, Hort. Brit. ed. II. 513 = littoralis.
Staplesiana, M. Roem. Syn. Enast. 176 = præc.
tenuiflora, *Herb. App.* 44.—N. Granat.

HYMENOCALLIS:—
tubiflora, *Salisb. in Trans. Hort. Soc.* i. (1812) 341.—Guian.
undulata, *Herb. App.* 44.—Venezuela.
virescens, Nichols. Dict. Gard. ii. 165 = Macleana.

HYMENOCALYX, Houll. in Rev. Hortic. (1869) 418 = præc.

HYMENOCALYX, Zenk. Pl. Ind. 8. t. 10 (1835) = Hibiscus, Linn. (Malvac.).
variabilis, Zenk. l. c. 9. t. 10 = H. angulosus.

HYMENOCARDIA, Wall. Cat. n. 3549 (1831). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 285.
SAMAROPYXIS, Miq. Fl. Ind. Bat. Suppl. 464 (1860). acida, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 256. —Afr. trop.
Heudelotii, *Muell. Arg. in Flora*, xlvii. (1864) 518.—Afr. trop.
lyrata, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 256. —Afr. trop.
plicata, Kurz, in journ. As. Soc. Beng. xlii. (1873) ii. 239 = Coccoceras plicatum.
punctata, *Wall. Cat.* n. 3549.—Burma.
ulmoides, *Oliver, in Hook. Ic. Pl.* t. 1131.—Afr. trop.
Wallichii, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 256 = punctata.

HYMENOCARPOS, Savi, Fl. Pis. ii. 205 (1798). *LEGUMINOSÆ*, Benth. & Hook. f. i. 489.
CIRCINNUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 384.
circinnata, *Savi, Fl. Pis.* ii. 205.—Hispan.
nummularius, *Willd. Enum. Hort. Berol. Suppl.* 52, nomen; *G. Don, Gen. Syst.* iii. 373.—Reg. Mediterr.
radiatus, Link, Enum. Hort. Berol. ii. 267 = Trigonella radiata.

HYMENOCARPUS, Reichb. Consp. 148 (1828) = præc.

HYMENOCENTRON, Cass. in Dict. Sc. Nat. xliv. 87 (1826) = Centaurea, Linn. (Compos.).
dilutum, Cass. l. c. = C. diluta.

HYMENOCEPHALUS, Jaub. & Spach, Illustr. iii. 12, t. 209 (1847) = Centaurea, Linn. (Compos.).
rigidus, Jaub. & Spach, l. c.—Persia.

HYMENOCHAETA, Beauv. in Lestib. Ess. Fam. Cyp. 48 (1819) = Scirpus, Tourn. (Cyperac.).
grossa, Nees, in Wight, Contrib. 110 = S. grossus.
haematodes, Nees & Meyen, in Linnæa, ix. (1834) 298 = S. grossus.
maxima, Nees, l. c. = S. grossus.

HYMENOCHARIS, Salisb. in Trans. Hort. Soc. i. (1812) 276 = Zachnospithon, Korn. (Scitamin.).
obliqua, Salisb. l. c. = I. obliqua.

HYMENOCLEA, Torr. & Gray, ex Torr. in Emory, Mil. Reconnois. 149 (1848); et in Mem. Am. Acad. N. S. iv. (1849) 79. *COMPOSITÆ*, Benth. & Hook. f. ii. 354.
monogyra, *Torr. & Gray, ll. cc.* 148, 79.—Am. bor. occ.
Salsola, *Torr. & Gray, ll. cc.* 142, 79.—Calif.

HYMENOCNEMIS, Hook. f. in Benth. & Hook. f. Gen. ii. 193 (1878). *RUBIACEÆ*, Benth. & Hook. f. i. c.
madagascariensis, *Hook. f. l. c.* 193.—Madag.

HYMENOCRATER, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 39 (1835). *LABIATÆ*, Benth. & Hook. f. ii. 1201.
SESTINIA, Boiss. Diagn. Ser. I. v. 40 (1844).
Auckeri, Jaub. & Spach, Illustr. v. 61. t. 456 = bituminosa.
bituminosa, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 177 = calycina.
bituminosus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 39.—Persia.
calycinus, *Benth. in DC. Prod.* xii. 406.—Persia.
elegans, *Bunge, Lab. Pers.* 68.—Persia.
Hausknechtii, Boiss. & Reut. ex Boiss. Fl. Orient. iv. 677 = longiflorus.

HYMENOCRATER:—
incanus, *Bunge, Lab. Pers.* 63.—Persia.
longiflorus, *Benth. in DC. Prod.* xii. 407.—Kurdist.
macrophyllus, *Bunge, Lab. Pers.* 62.—Persia.
pallens, *Bunge, l. c.* 61.—Persia.
paniculatus, *Bunge, l. c.* 62.—Persia.
secundiflorus, Jaub. & Spach, Illustr. v. 63 = bituminosus.
sessilifolius, *Benth. in DC. Prod.* xii. 406.—Afghan.

HYMENODICTYON, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 148 (1824). *RUBIACEÆ*, Benth. & Hook. f. ii. 25.
KURRIA, Hochst. in Flora, xxv. (1842) 283.
bifranum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 42.—Afr. trop.
excelsum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 149.—Ind. or.; Malaya.
flaccidum, *Wall. l. c.* 152.—Reg. Himal.
Horsfieldii, Miq. Fl. Ind. Bat. ii. 154 = excelsum.
Kurria, *Hochst. in Flora*, xxvi. (1843) 71.—Afr. trop.
madagascarium, *Baill. Hist. Pl.* vii. 482.—Madag.
obovatum, *Wall in Roxb. Fl. Ind. ed. Carey*, ii. 153. —Ind. or.
obovatum, Wight, Ic. t. 80 = excelsum.
parvifolium, *Oliver, in Hook. Ic. Pl.* t. 1488.—Afr. trop.
thyrsiflorum, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 151 = excelsum.
timoriense, *Klotzsch, ex Walp. Rep.* vi. 63.—Ins. Timor.
utile, Wight, Ic. t. 1159 = excelsum.

HYMENODYCTION, DC. Prod. iv. 358 (1830) = præc.

HYMENOGYNE, Haw. Rev. Pl. Succ. 74, 192 (1821) = Mesembrianthemum, Dill. (Ficoid.).
glabra, Haw. l. c. 192 = M. glabrum.

HYMENOLAENA, DC. Prod. iv. 244 (1830) = Pleurospermum, Hoffm. (Umbellif.).
angelicoides, DC. l. c. 245 = P. angelicoides.
Benthami, DC. l. c. 246 = P. Benthami.
Brunonis, DC. l. c. 245 = P. Brunonis.
Candollei, DC. l. c. = P. Candollii.
dentata, DC. l. c. = P. dentatum.
erosa, Steud. Nom. ed. II. i. 784 = P. dentatum.
Govaniana, DC. Prod. iv. 246 = P. Govanianum.
latifolia, Lindl. in Royle, Illustr. Bot. Himal. 233 = P. Candollii.
Lindleyana, Klotzsch, in Bot. Ergeb. Waldem. Reise, 150 = P. stellatum.
nana, Osten-Sacken & Rupr. Sert. Tianach. 49 = P. stellatum.
obtusiscula, DC. Prod. iv. 246 = Trachydium obtusiusculum.
pimpinelliifolia, Osten-Sacken & Rupr. Sert. Tianach. 50 = P. Candollii.
pumila, DC. Prod. iv. 245 = P. pumilum.
rotundata, DC. l. c. = P. rotundatum.
stellata, Lindl. in Royle, Illustr. Bot. Himal. 233 = P. stellatum.

HYMENOLEPIS, Cass. in Bull. Soc. Philom. (1817) 188; et in Dict. Sc. Nat. xxii. 315 (1821) = Athanasia, Linn. (Compos.).
canariensis, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 294. tt. 97, 98 = Gonospermum canariense.
Dregeana, DC. Prod. vi. 85 = A. Dregeana.
elegans, Cass. in Dict. Sc. Nat. xxxiii. 61 = Gonospermum canariense.
incisa, DC. Prod. vi. 85 = A. incisa.
indivisa, Drège, ex Harv. & Sond. Fl. Cap. iii. 190 = A. indivisa.
leptocephala, Cass. in Dict. Sc. Nat. xxii. 315 = A. parviflora, *Thunb.*
leucoclada, DC. Prod. vi. 86 = A. leucoclada.
parviflora, DC. l. c. 85 = A. parviflora, *Thunb.*
pinnata, Less. Syn. Comp. 265 = A. parviflora, *Thunb.*
punctata, DC. Prod. vi. 85 = A. punctata.

HYMENOLOBIUM, Benth. ex Mart. in Flora, xx. (1837) ii. Beibl. 122 = Platymiscium, Vog. (Leguminos.).
hebrum, Benth. l. c. = P. floribundum.

HYMENOLOBIUM, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 84. *LEGUMINOSÆ*, Benth. & Hook. f. i. 548.
nittidum, *Benth. l. c.*—Bras.

HYMENOLOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 117 (1838) = **Capsella**, Moench (Crucif.).
divaricatus, Nutt. l. c. = C. divaricata.
erectus, Nutt. l. c. = C. divaricata.
procumbens, Nutt. l. c. in nota = C. procumbens.
pubens, A. Gray, Pl. Wright. t. 9; ii. 14 = C. pubens.

HYMENOLYTRUM, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 174. t. 22 (1842) = **Scleria**, Berg. (Cyperac.).
comosum, Nees, l. c. 175 = S. comosa.
cyperinum, Nees, l. c. = S. cyperina.
Martii, Nees, l. c. 176 = S. Martii.
Poeppigii, Nees, l. c. 175 = S. Poeppigii.
sileostris, Schrad. ex Nees, l. c. 176 = S. stipularis.

HYMENOMENA, Less. Syn. Comp. 133 (1832) = seq.

HYMENONEMA, Cass. in Bull. Soc. Philom. (1817) 84. *COMPOSITÆ*, Benth. & Hook. f. ii. 505.
asperum, DC. Prod. vii. 117 = Tournefortii.
Bergeri, Sch. Bip. ex Nym. Consp. 472 = Tournefortii
Fontanesii, Cass. in Dict. Sc. Nat. xxii. 317 = Tournefortii.
glauca, Hook. Fl. Bor. Am. i. 300 = Microseris laciniata ?
graveum, DC. Prod. vii. 116 = Tournefort..
laciniata, Hook. Fl. Bor. Am. i. 301 = Microseris laciniata.
laeviceum, *Boiss. & Heldr. ex Boiss. Fl. Orient* iii. 715.—Graecia.
Tournefortii, *Cass. in Dict. Sc. Nat.* xxii. 316.—Graecia.

HYMENOPAPPUS, L'Hérit. Diss. (1788). *COMPOSITÆ*, Benth. & Hook. f. ii. 402.
Kothia, Lam. Journ. Hist. Nat. Par. i. (1792) 16. t. 1.
anthemoides, Juss. in Ann. Mus. Par. ii. (1803) 426 = Hymenoxys anthemoides.
artemisiæfolius, *DC. Prod.* v. 658.—Texas.
corymbosus, *Torr. & Gray, Fl. N. Am.* ii. 272 — Texas.
Douglasii, Hook. Fl. Bor. Am. i. 316 = Chaenactis Douglasii.
Engelmannianus, *Kunth, Ind. Sem. Hort. Berol.* (1848) 15; *et in Ann. Sc. Nat.* Ser. III. xi. (1849) 229.—Texas.
filifolius, *Hook. Fl. Bor. Am.* i. 317.—Am. bor. occ.
flavescens, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 97.—Am. bor. occ.
glaucus, Spreng. Syst. iii. 449 = Cephalaphora aromatica, glauca.
integrifolius, Spreng. l. c. = Palafoxia fastigiata.
lanciformis, L'Hérit. ex DC. Prod. v. 658 = scabioseus.
lutens, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 374 = filifolius.
matricarioides, Spreng. Syst. iii. 450 = Ageratum matricarioides.
mexicanus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 29.—Mexic.
nevadensis, Kellogg, in Proc. Calif. Acad. v. (1873) 46 = Chaenactis nevadensis.
pedatus, Cav. ex Lag. Gen. et Sp. Nov. 28 = Florestina pedata.
robustus, Greene, in Bull. Torrey Bot. Club, ix. (1882) 53 = flavescens.
scabiosus, Stend. Nom. ed. II. i. 764, sphalm. = seq.
scabiosaeus, Dougl. ex Hook. Fl. Bor. Am. i. 316 = Chaenactis Douglasii.
scabiosaeus, *L'Hérit. Diss.* (1788); *Cass. in Dict. Sc. Nat.* lv. 395.—Am. bor.
tenuifolius, Dougl. ex Hook. Fl. Bor. Am. i. 317 = filifolius.
tenuifolius, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 272 = corymbosus.
tenuifolius, *Pursh, Fl. Am. Sept. ii. Suppl.* 742.—Am. bor.

HYMENOPHOLIS, Garden. in Hook. Lond. Journ. Bot. vii. (1848) 58 = **Oligandra**, Less. (Compos.).
imbricata, Garden. l. c. = O. lycopodioides.

HYMENOPHORA, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 281 = **Pituranthos**, Viv. (Umbellif.).
denudata, Viv. ex Coss. l. c. = P. denudatus.

HYMENOPHYSA, C. A. Mey. in Ledeb. Fl. Alt. iii. 180 (1831). *CRUCIFERÆ*, Benth. & Hook. f. i. 88.
fenestrata, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 107.—Persia.
macrocarpa, *Franch. l. c.* Sér. VI. xv. (1883) 288.—Turkestan.
pubescens, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 181.—Sibir.

HYMENOPOGON, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 156 (1824). *RUBIACEÆ*, Benth. & Hook. f. ii. 84.
asaamicus, *Hook. f. Fl. Brit. Ind.* iii 34.—Reg. Himal. parasiticus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 157. —Reg. Himal.

HYMENOPYRAMIS, Wall. Cat. n. 774 (1831). *VERBENACEÆ*, Benth. & Hook. f. ii. 1158.
brachiata, *Wall. l. c.*—Burma.

HYMENOSPERMUM, Benth. in Wall. Cat. n. 8968 (1831) = **Alcotra**, Thunb. (Scrophular.).
dentatum, Benth. l. c. = A. indica.

HYMENOSPORUM, R. Br. ex F. Muell. Fragm. vi. 77 (1860). *PITTOSPOREÆ*, Benth. & Hook f. i. 131.
flavum, *F. Muell. l. c.*—Austral.

HYMENOSPRON, Spreng. Syst. iv. Cur. Post. 282 (1827) = **Dicoelus**, H. B. & K. (Legumin.).
apurensis, Spreng. l. c. = D. apurensis
molle, Heynh. Nom. i. 408 = D. mollis.
rubrum, Spreng. Syst. iv. Cur. Post. 288 = D. Jacquinianus.
sericeum, Spreng. l. c. = D. sericeus.

HYMENOSTEMMA, Kunze, ex Willk. in Bot. Zeit. xxii. (1864) 258 = **Chrysanthemum**, Tourn. (Compos.).
Fontanesii, Willk. l. c. = C. glabrum.
paludosum, Pomel, Nouv. Mat. Fl. Atl. 292 = C. paludosum.
paludorum, Willk. in Bot. Zeit. xxii. (1864) 253 = C. glabrum.
Pseudanthemis, Kunze, ex Willk. l. c. = C. paludosum.

HYMENOSTEPHIUM, Benth. in Benth. & Hook. f. Gen. ii. 882 (1873). *COMPOSITÆ*, Benth. & Hook. f. i. c.
angustifolium, *Benth. in Hook. Ic. Pl.* sub t. 1154.—N. Granat.
mexicanum, *Benth. l. c.*—Mexic.

HYMENOSTIGMA, Hochst. in Flora, xxvii. (1844) 24 = **Morasa**, Linn. (Irid.).
Schimperi, Hochst. l. c. = M. diversifolia.
tridentatum, Hochst. l. c. 25 = M. diversifolia.

HYMENOTHECA, F. Muell. Fragm. i. 201 (1858-59) = **Codonocarpus**, A. Cunn. (Phytolac.).
pyramidalis, F. Muell. l. c. = C. pyramidalis.

HYMENOTHECA, Salisb. in Trans. Hort. Soc. i. (1812) 268 = **Ottelia**, Pers. (Hydrochar.).
latifolia, Salisb. l. c. = O. alismoides.

HYMENOTHECIUM, Lag. Gen. et Sp. Nov. 4 (1816) = **Aegopogon**, Willd. (Gramin.).
quinquestrium, Lag. l. c. = A. cenchroides.
tenuiflum, Lag. l. c. = A. geminiflorus.
triastrum, Lag. l. c. = A. cenchroides.
unistrum, Lag. l. c. = A. geminiflorus.

HYMENOTHRIX, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) [Fl. Fendl.) 109. *COMPOSITÆ*, Benth. & Hook. f. ii. 403.
Whitlavii, *A. Gray, l. c.*—N. Mexic.
Wrightii, *A. Gray, Pl. Wright.* ii. 97.—N. Mexic.

HYMENOXIS, Endl. Ench. 940 (1841) = seq.

HYMENOXYS, Cass. in Dict. Sc. Nat. lv. 378 (1828). *COMPOSITÆ*, Benth. & Hook. f. ii. 415.
anthemoides, *Cass. l. c.*—Reg. Argent.
californica, Hook. Bot. Mag. t. 3828 = Baeria coronaria.
calva, Torr. & Gray, Fl. N. Am. ii. 331 = Baeria anthemoides.
chrysanthemoides, DC. Prod. v. 661 = Helenium linifolium.
Haenkeana, *DC. Prod.* v. 661.—Peruv.
linearifolia, Hook. Ic. Pl. t. 146 = Actinella linearifolia.
mutica, Torr. & Gray, Fl. N. Am. ii. 331 = Baeria mutica.
odorata, DC. in Deless. Ic. Sel iv. t. 42 = Actinella odorata.
Tweedlei, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 323.—Reg. Argent.

HYOBANCHE, Linn. Mant. ii. 155 (1771). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 988.
Haematobanche, Presl, Epim. Bot. 249 (1850).
atropurpurea, *Bolus, in Hook. Ic. Pl.* t. 1486.—Afr. austr.
coccinea, Steud. Nom. ed. I. 418 = sanguinea.
sanguinea, *Linn. Mant.* ii. 253.—Afr. austr.

HYOCYAMUS, G. Don, Gen. Syst. iv. 471 (1837), sphalm. = **Hyoscyamus**, Tourn. (Solanac.).

HYOGETON, Steud. Nom. ed. II. i. 785 (1840), sphalm. = ILYOGETON, Endl. = **Vandellia**, Linn. (Scroph.).
alsinoides, Steud. l. c. = V. alsinoides.
scapiger, Steud. l. c. = V. scapigera.
subulatus, Steud. l. c. = V. subulata.

HYOPHORBE, Gaertn. Fruct. ii. 186. t. 120 (1791). *SOLANA PALMÆ*, Benth. & Hook. f. iii. 912.
Sublimia, Comm. ex Mart. Hist. Nat. Palm. iii. 164 (1836?).
amaricaulis, *Mart. Hist. Nat. Palm.* iii. 309.—Ins. Maurit.
Commersoniana, Mart. l. c. 164. t. 143. f. 1; t. 164 = Clarysalidocarpus lutescens.
indica, *Gaertn. Fruct.* ii. 186. t. 120.—Ins. Mascar.
Verschaffelti, *H. Wendl. in Lem. Illustr. Hortic.* xiii. (1866) t. t. 462.—Ins. Mascar.

HYOSCARPUS, Dulac, Fl. Hautes-Pyr. 444 (1867) = **Hyoscyamus**, Tourn. (Solanac.).
niger, Dulac, l. c. = Hyoscyamus niger.

HYOSCHYAMUS, Zumag. Fl. Pedem. i. 280 (1849) = seq.

HYOSCIAMUS, Neck. Delic. Gallo-Belg. i. 122 (1768) = seq.

HYOSCYAMUS, [Tourn.] Linn. Syst. ed. I (1735). *SOLANACEÆ*, Benth. & Hook. f. ii. 908.
Hyoscarpus, Dulac, Fl. Hautes-Pyr. 444 (1867).
agrestis, Kit. in Schult. Oestr. Fl. ed. II. i. 583 = niger.
albus, *Linn. Sp. Pl.* 180.—Europ. ; Orient.
angulatus, Griff. Notul. iv. 96; Ic. Pl. As. t. 412 = muticus.
arenarius, *Dun. in DC. Prod.* xiii. 550.—Ins. Asor.
aureus, All. Fl. Pedem. i. 104 = albus.
aureus, *Linn. Sp. Pl.* 180.—As. Min. ; Mesopot. ; Syria.
auriculatus, Tenore, Fl. Nap. iii. 222 = niger.
betaefolius, Lam. Encyc. iii. 329 = muticus.
bipinnatisectus, Boiss. Diagn. Ser. I. vii. 86 = Senecionis.
bohemicus, F. W. Schmidt, Fl. Boëm. iii. 31 = niger.
Camerarii, Finch. & Hey. Ind. Sem. Hort. Petrop. iv. 38 = reticulatus.
canariensis, Ker-Gawl. in Bot. Reg. t. 180 = albus.
cerstophyllus, *Fisch. ex Steud. Nom.* ed. I. 418.—Hab. ?
chlorantha, Wender. ex Steud. l. c. ed. II. i. 785 = Scopolina Hladnikiana !
Clusii, G. Don, in Sweet, Hort. Brit. ed. II. 582 = albus.
Daiera, Forsk. Fl. Aegypt. Arab. 45 = muticus.
eminens, *Kunze, Del. Sem. Hort. Lips.* (1847).—*Cf. Linnaea*, xxiv. (1851) 176.—Rossia.

HYOSCYAMUS:—
Faleslex, *Cos. in Bull. Soc. Bot. Fr.* xi. (1864) 166. t. 5.—Afr. bor.
grandiflorus, *French. in Révoil. Çomali*, 48.—Afr. trop.
insanus, Stocks, in Hook. Kew Journ. iv. (1852) 178 = muticus.
lethalis, Salisb. Prod. 131 = niger.
luridus, Salisb. l. c. = albus.
major, Mill. Gard. Dict. ed. VIII. n. 2 = albus.
micranthus, Ledeb. ex G. Don, Gen. Syst. iv. 472 = pusillus.
minor, Mill. Gard. Dict. ed. VIII. n. 4 = albus.
muticus, *Linn. Mant.* i. 45.—Aegypt.; As. occ.
niger, *Linn. Sp. Pl.* 179.—Europ.; As. occ.; Reg. Himal.
officinarum, Crantz, Inst. ii. 825 = niger.
orientalis, Bieb. Fl. Taur-Cauc. i. 164 = Physochlaina orientalis.
pallidus, Waldst. & Kit. ex Willd. Enum. Hort. Berol. i. 228 = niger.
persicus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 158 = niger.
physaloides, *Linn. Sp. Pl.* 180.—Sibir.
pictus, Roth, Nov. Fl. Sp. 119 = niger.
pinnatifidus, Schlecht. in Linnæa, xvii. (1843) 127 = reticulatus.
praealtus, Walp. Rep. iii. 21 = Physochlaina praealta. Pseudophysaloides, *Roth, Nov. Pl. Sp.* 119.—Sibir.
pungens, Griseb. Spicil. Fl. Rumel. ii. 52 = pusillus.
purpureus, Grisch. l. c. in nota = reticulatus.
pusillus, *Linn. Sp. Pl.* 180.—Aegypt.; As. bor. et occ.
reticulatus, *Linn. l. c.* ed. II. 257.—As. Min.; Aegypt.
Scopolia, Linn. Mant. i. 46 = Scopola carniolica.
Senecionis, *Willd. Enum. Hort. Berol.* i. 228.—Orient.
squarrosus, Griff. Notul. iv. 95; Ic. Pl. Asiat. t. 413 = reticulatus.
syspirensis, C. Koch, in Linnæa, xxii. (1849) 736 = niger.
varians, Vis. in Flora, xii. (1829) i. Erg. 7 = albus.
vernicosus, Lej. Fl. Spa, i. 116 = niger.
vulgaris, Neck. Delic. Gallo-Belg. i. 122 = niger.

HYOSERIS, Linn. Gen. ed. I. 239 (1737) partim. *COMPOSITÆ*, Benth. & Hook. f. ii. 507.
ACHYRASTRUM, Neck. Elem. i. 57 (1790).
APOSERIS, Neck. l. c. (1790).
THLIPSOCARPUS, Kunze, in Flora, xxix. (1846) 695.
TRINCIALTELLA, Adans. Fam. ii. 113 (1763).
aculeata, Pers. Syn. ii. 369 = Leontodon hastilis.
adspersa, Moench, Meth. 541 = scabra.
amplexicaulis, Michx. Fl. Bor. Am. ii. 87 = Krigia amplexicaulis.
angustifolia, Michx. l. c. = Krigia Dandelion.
arenaria, Schousb. Vextr. Marokko, 197 = Rhagadiolus arenarius.
aspera, Desf. Tabl. ed. I. 89 = scabra.
bætica, Sch. Bip. ex Nym. Consp. 474 = radiata.
bifora, Walt. Fl. Carol. 194 = Krigia amplexicaulis.
blechnoides, *Pomel, Nouv. Mat. Fl. Alg.* 271.—Alger.
calyculata, Jan, ex Guss. Fl. Sic. Syn. ii. i. 417 = radiata.
calyculata, Potr. Voy. Barb. ii. 229 = lucida.
canadensis, Fisch. ex Steud. Nom. ed. I. 418 (Quid?). —Am. bor.
capensis, Less. Syn. Comp. 180 = Hypochoeris glabra.
caroliniana, Walt. Fl. Carol. 194 = Krigia virginica.
cichoroides, Steud. Nom. ed. I. 418 = Leontodon hirtus.
crepoides, Presl. Fl. Sic. p. xxxi = Rhagadiolus pendulus?
cretica, Linn. Sp. Pl. 810 = Rhagadiolus Hedypnois, *All.*
exigua, Salisb. Prod. 132 = Arnoseris pusilla.
foetida, *Linn. Sp. Pl.* 808.—Europ.
glauca, Zeyh. ex Sch. Bip. in Nov. Act. Nat. Cur. xx. i. (1845) 161 = Hypochoeris Achyrophorus.
grumosa, Poir. Encyc. Suppl. iii. 82 = Leontodon taberosus.
Hedypnois, Linn. Sp. Pl. 809 = Rhagadiolus Hedypnois, *All.*
hirta, Balb. ex Willd. Sp. Pl. iii. 1614 = Crepis bursifolia.
hirta, Gaertn. Fruct. ii. 375 = Leontodon hirtus.
hispida, Schousb. Vextr. Marokko, 197 = Leontodon marocanus.

HYOSERIS:—
incrassata, Moench, Meth. Suppl. 218 = Rhagadiolus Hedypnois, *All.*
integrifolia, *Poir. Encyc. Suppl.* iii. 80.—Am. bor.
lucida, Anct. Europ. ex Nym. Consp. 474 = radiata.
lucida, *Linn. Mant.* i. 108.—Aegypt.
major, Walt. Fl. Carol. 194 = Krigia Dandelion.
maroccana, Pers. Syn. ii. 368, in syn. = Leontodon marocanus.
maura, Pers. l. c. 369 = Rhagadiolus Hedypnois, *All.*
microcephala, Cass. in Dict. Sc. Nat. xxii. 326 = scabra.
minima, Linn. Sp. Pl. 809 = Arnoseris pusilla.
minor, J. F. Gmel. Syst. 1132 = Krigia amplexicaulis.
monspeliensis, Pers. Syn. ii. 369 = Rhagadiolus Hedypnois, *All.*
montana, Michx. Fl. Bor. Am. ii. 87 = Krigia montana.
muricata, *Sprengl. in Schrad. Journ.* (1806) ii. 198.—Hab. ?
nana, *Poir. Encyc. Suppl.* iii. 82.—Ins. Majorca.
parviflora, Pers. Syn. ii. 369 = scabra.
pendula, Balb. in Pers. Syn. ii. 369 = Rhagadiolus pendulus.
prenanthoides, Willd. Sp. Pl. iii. 1615 = Krigia amplexicaulis.
pusilla, Regel, in Linnæa, xvi. (1842) 51, in syn. = Hypochoeris glabra.
pygmaea, Ait. Hort. Kew. ed. I. iii. 130 = Leontodon nudicaulis.
radiata, Gaertn. ex Steud. Nom. ed. I. 419 = scabra.
radiata, *Linn. Sp. Pl.* 808.—Reg. Mediterr.
radiata, Lucé, Fl. Cail. 264 = Leontodon hastilis.
ramosissima, Barton, Prod. Fl. Philad. 75 = Krigia virginica.
rhagadioloides, Linn. Sp. Pl. 809 = Rhagadiolus Hedypnois, *All.*
scabra, *Linn. l. c.*—Reg. Mediterr.
scabra, Moench, Meth. 541 = Rhagadiolus Hedypnois, *All.*
scabra, Pourr. in Willk. & Lange, Prod. Fl. Hisp. ii. 209 = radiata.
setulosa, Guss. ex DC. Prod. vii. 80 = lucida.
simplex, Mérat, Fl. Par. 310 = Hypochoeris glabra.
Sprengeli, Steud. Nom. ed. I. 419 = Arnoseris pusilla.
taraxacoides, Lam. Encyc. iii. 159 = Leontodon hirtus.
taraxacoides, Vill. Prosp. 33; Hist. Pl. Dauph. iii. 166 = Leontodon hispidus.
tenella, Nym. Consp. 474. Sch. Bip. in Linnæa, x. (1835) 255 (=*Schkea Poeppigii*).—Chili.
tenella, Thunb. Prod. Fl. Cap. 139 = Hypochoeris glabra.
tuberifera, Tenore, Fl. Nap. ii. 178. t. 78 = Rhagadiolus Hedypnois, *All.*
tuberosa, Savi, Fl. Pis. ii. 231 = Leontodon tuberosus.
virginica, Linn. Sp. Pl. 809 = Krigia virginica

HYOSICAMUS, Hill, Veg. Syst. vii. 51 (1765) sphalm. = Hyoscyamus, Tourn. (Solanac.).

HYOSPATHE, Mart. Hist. Nat. Palm. ii. 1. t. 1, 2 (1838). *PALMAE*, Benth. & Hook. f. iii. 898.
antiloguensis, *Hort. Linden, ex H. Wendl. in Kerch. Palm.* 247, nomen.—N. Granat.
chiriquensis, *Hort. Linden, ex H. Wendl. l. c.*—Costa Rica.
elegans, *Mart. Hist. Nat. Palm.* ii. 1. t. 1, 2.—Bras.
filiformis, *H. Wendl. ex Drude, in Mart. Fl. Bras.* iii. 11. 523.—Bras.
gracilis, H. Wendl. ex Drude, l. c. = filiformis.
montana, Mart. Palmet. Orb. t. 6. f. 1 = Chamaedorea concolorpa.
pubispina, Griseb. & H. Wendl. ex Grisb. Fl. Brit. W. Ind. 516 = Prestoëa pubigera.

HYPAELYPTUM, Vahl, Enum. ii. 288 (1806) = seq.

HYPAELYTRUM, Poir. in Dict. Sc. Nat. xxii. 340 (1821) = **Hypolytrum**, Rich. (Cyperac.).

HYPANTHERA, Silva Manso, Cathart. Bras. 37 (1836) = **Pavillea**, Linn. (Cucurb.).
Guapeva, Silva Manso, l. c. 38 = F. trilobata.

HYPAPHORUS, Hamk. Hort. Bogor. Desc. 197 (1858) = **Erythrina**, Linn. (Legumin.).
subumbrosus, Hassk. l. c. 198 = E. lithosperma.

HYPARETE, Rafin. Fl. Tellur. iii. 43 (1836) = **Hermbstaedtia**, Reichb. (Amarant.).
glauca, Rafin. l. c. 44 = Hermbstaedtia glauca.

HYPARGYRIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 371 = **Potentilla**, Linn. (Rosac.).
argentatum, *argentatum*, *canum*, *confine*, *decipiens*, *decumbens*, *demissum*, Fourr. l. c. = P. argentea.
hirtum, Fourr. l. c. 372 = P. hirta.
rectum, Fourr. l. c. 371.—Gallia.
rectum, Fourr. l. c. 372 = P. recta.
suberectum, Fourr. l. c. 371 = P. inclinata.
tenuilobum, Fourr. l. c. = P. argentea.

HYPARRHENIA, Anders. in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1855) 254 = **Andropogon**, Linn. (Gramin.).
anthistirioides, Anders. in Schweinf. Beitr. Fl. Aethiop. 300 = A. anthistirioides.
coleotricha, Anders. l. c. = A. coleotrichus.
comosa, Anders. l. c. = A. coleotrichus.
confinis, Anders. l. c. = A. confinis.
dissoluta, Anders. in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1856) 254 = Trachypogon polymorphus.
finitima, Anders. in Schweinf. Beitr. Fl. Aethiop. 300 = A. finitimus.
falcata, Anders. in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1855) 254 = A. bracteatus.
fulvicoma, Anders. in Schweinf. Beitr. Fl. Aethiop. 300 = A. rufus.
Gemisimia, Anders. l. c.—Afr. trop.
glabriuscula, Anders. l. c. = A. glabriusculus.
monathera, Anders. l. c. = A. monatherus.
multiplex, Anders. in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1855) 254.—Abyss.
papillipes, Anders. in Schweinf. Beitr. Fl. Aethiop. 300 = A. papillipes.
podotricha, Anders. l. c. = Heteropogon hirtus.
Pseudo-Cymbaria, Anders. in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1855) 254 = A. cymbarius.
quinquenlpta, Anders. l. c. = A. cymbarius.
reflexa, Anders. l. c. = A. bracteatus.
Ruprechti, Fourn. ex Hemsl. Biol. Centr. Am. Bot. ii. 509 = A. Ruprechti.
Schimperi, Anders. in Schweinf. Beitr. Fl. Aethiop. 300 = A. Schimperi.
Tamba, Anders. l. c. = A. lepidus.
umbrosa, Anders. l. c. = A. lepidus.

HYPECHUSA, Alef. in Bot. Zeit. xviii. (1860) 165 = **Viola**, Tourn. (Legumin.).
hircania, Alef. l. c. 166 = V. hircanica.
hybrida, Alef. l. c. = V. hybrida.
lutea, Alef. l. c. = V. lutea.
pannonica, Alef. l. c. = V. pannonica.
purpurascens, Alef. l. c. = V. pannonica.
sericarpa, Alef. l. c. = V. sericarpa.
tricolor, Alef. l. c. = V. Melanops.

HYPECOON, [Tourn.] Linn. Syst. ed. I (1735) = seq.

HYPECOUM, Tourn. ex Linn. Gen. ed. I. 32 (1737). *PAPAVERACEAE*, Benth. & Hook. f. i. 54.
CHIAZOSPERMUM, Bernh. in Linnæa, viii. (1833) 465.
MUNBOMILLA, Forsk. Fl. Aegypt. Arab. 122 (1775).
aequilobum, *Vis. Fl. Lib. Spec.* 7. t. 3. f. 3.—Cyrenaic.
arcuatum, Moench, Meth. 217 = procumbens.
armatum, Steud. Nom. ed. II. i. 786, sphalm. = praec.
caucasicum, Koch, ex Ledeb. Fl. Ross. i. 94 = pendulum.
chinense, *Franch. Pl. Davidian.* i. 27.—China.
dimidiatum, Delile, in Laborde, Voy. Arab. Pétr. 84 = procumbens.
Duriaei, *Pomel, Nouv. Mat. Fl. Alg.* 246.—Alger.
erectum, *Linn. Sp. Pl.* 794.—Sibir.
Gaillei, Cos. & Kral. in Bull. Soc. Bot. Fr. iv. (1857) 523, nomen = procumbens.
glaucescens, Gess. Pl. Rar. 79. t. 15 = procumbens.
gracile, Benth. Cat. Pl. Pyr. 91 = procumbens.
grandiflorum, *Benth. l. c.*—Reg. Mediterr. et Caucas.
imberbe, *Sibth. & Sm. Fl. Graec. Prod.* i. 107.—Reg. Mediterr. or.
leptocarpum, *Hook. f. & Thoms. Fl. Ind.* i. 276.—Reg. Himal.
littorale, Wall. in Jacq. Coll. ii. 305 = procumbens.
nudicaule, Lam. Fl. Fr. ii. 670 = procumbens.
parviflorum, *C. & W. Herb. Herbar. au Levant*, 115. t. 2.—Syria.
parviflorum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 141.—Songaria.
patens, Willd. Enum. Hort. Berol. i. 5 = imberbe.

HYPECOUM:—
pendulum, Lapeyr. Hist. Abr. Pl. Pyr. 76 = grandiflorum.
pendulum, *Linn. Sp. Pl.* 124.—Europ. ; Orient.
procumbens, S. G. Gmel. It. ii. 197 = pendulum.
procumbens, *Linn. Sp. Pl.* 124.—Reg. Medit. ; Arab.
litrogenum, Bertol. Misc. Bot. i. 12 = procumbens.
trilobum, *Trautv. in Act. Hort. Petrop.* ix. (1884) 366. —Turcomania.

HYPELATE. P. Br. Hist. Jamaic. 280 (1756). *SAPINDACEÆ*, Benth. & Hook. f. i. 408.
EXOTHEA, Macfad. Fl. Jamaic. i. 333 (1837).
cochinchinensis, DC. Prod. ii. 480, in syn. sphalm. (= *Hyperanthera cochinchinensis*).—Cochinch.
dentata, Cambess. in Mém. Mus. Par. xviii. (1829) 22 = Melicocca dentata.
diversifolia, Cambess. l. c. = Doratoxylon diversifolium.
geniculata, G. Don, Gen. Syst. i. 672 = Melicocca geniculata.
oblongifolia, Hook. Lond. Journ. Bot. iii. (1844) 7 = paniculata.
paniculata, *Cambess. l. c.*—Ind. occ.
trifoliata, Griseb. in Goett. Abh. vii. (1857) 188 = paniculata.
trifoliata, *Sw. Prod. Veg. Ind. Occ.* 61.—Ind. occ.

HYPELYTHRUM, D. Dietr. Syn. Pl. i. 180 (1839) = seq.

HYPELYTRUM, Poir. in Dict. Sc. Nat. xxxii. 340 (1821) = **Hypolytrum**, Rich. (Cyperac.).

HYPENANTHE, Blume, Mus. Bot. Lugd. Bat. i. 21 (1849) = **Medinilla**, Gaudich (Melastom.).
verucosa, Blume, l. c. = M. venosa.

HYPENIA, Mart. ex Benth. Lab. Gen. et Sp. 186 (1888) = **Hyptis**, Jacq. (Labiat.).
ocymoides, Mart. ex J. A. Schmidt, in Mart. Fl. Bras viii. 137 = Hyptis communis.
Salzmanni, Mart. ex J. A. Schmidt, l. c. = Hyptis Salzmanni.

HYPERACANTHUS, E. Mey. in Drège, *Zwei Pfl.* Docum. 198 (1848) = **Gardenia**, Ellis (Rubiac.).
floridus, E. Mey. l. c. = G. Neuberia.

HYPERANTHERA. Forsk. Fl. Aegypt. Arab. 67 (1775) = **Moringa**, Juss.
aptera, Steud. Nom. ed. II. i. 786 = M. aptera.
arborea, J. F. Gmel. Syst. 408 = M. pterygosperma.
cochinchinensis, Willd. Sp. Pl. ii. 587 (Caesalpinie sp. ?).—Cochinch.
decandra, Willd. l. c. 585 = M. pterygosperma.
dioica, Vahl, Symb. Bot. i. 31 = Gymnocladus canadensis.
monolynama, Vitm. Summa Pl. ii. 248 = Moringa aptera.
Moringa, Vahl, Symb. Bot. i. 30 = M. pterygosperma
peregrina, Forsk. Fl. Aegypt. Arab. p. cvii = M. aptera.
semidecandra, Vahl, Symb. Bot. i. 30 = M. aptera.

HYPERANTHUS, Harv. & Sond. Fl. Cap. iii. 6, in syn. (1864) = HYPERACANTHUS, E. Mey.

HYPERBAENA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 44 = **Pachygone**, Miers (Menisp.).
banisteriæfolia, Miers, l. c. Ser. III. xix. (1867) 98 = P. oblongifolia.
columbica, Miers, l. c. = P. colombica.
crebriflora, Miers, l. c.—Cuba.
cuneifolia, Miers, l. c.—Cuba.
dominguensis, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 50 = P. dominguensis.
gracilifiora, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 94 = P. dominguensis.
Hostmanni, Miers, l. c. Ser. II. vii. (1851) 44 = P. dominguensis.
longipaniculata, Miers, l. c. Ser. III. xix. (1867) 93.—Cuba.
mexicana, Miers, l. c. Ser. II. vii. (1851) 44 = P. dominguensis.
Mericandi, Miers, l. c. = P. dominguensis.
nemoralis, Miers, l. c. = P. dominguensis.
Prieriana, Miers, l. c. Ser. III. xix. (1867) 95.—Ind. occ.
reticulata, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 50 = P. dominguensis.

HYPERBAENA :—
retinervis, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 96.—Ind. occ.
rotundifolia, Miers, l. c.—Cuba.
Tweedieri, Miers, l. c. Ser. II. vii. (1851) 44 = Pachygone dominguensis.
validia, Miers, l. c. Ser. III. xix. (1867) 95 = P. dominguensis.

HYPERICINEA, Wall. Cat. n. 4827, etc. (1831) ; Steud. Nom. ed. II. i. 786 (1840)
angustata, Wall. Cat. n. 4830 = Buchanania florida.
dentata, Wall. l. c. n. 4832 = Ixonanthes icosandra.
lucida, Wall. l. c. n. 4827 = Buchanania acuminata.
macrocarpa, Wall. l. c. n. 4838 = Ixonanthes reticulata.
micrantha, Wall. l. c. n. 4829 · Tristania burmannica.
paucinervia, Wall. l. c. n. 4826 = Ixonanthes khasiana.
pimentifolia, Wall. l. c. n. 4828 = Tristania Wightiana.

HYPERICOIDES, Plum. ex Adans. Fam. ii. 448 (1763) = **Ascyrum**, Tourn. (Hyperac.).
Crux-Andreæ, Poir. Tabl. Encyc. iii. 201 = A. Crux-Andreæ.
perforata, Poir. l. c. = A. Hypericoides !

HYPERICON, J. F. Gmel. Syst. 1156 (1791) = **Hypericum**, Tourn.

HYPERICOPHYLLUM, Steetz, in Peters, Reise Mossamb. Bot. 498. t. 50 (1862) = **Jaumea**, Pers. (Compos.).
compositarum, Steetz, l. c. 499 = J. compositarum.

HYPERICOPSIS, Boiss. Diagn. Ser. I. vi. 25 (1846) = **Frazeria**, Linn.
persica, Boiss. Diagn. Ser. I. vi. 26 = F. persica.

HYPERICOPSIS, Opiz, Seznam, 58 (1852) = **Hypericum**, Tourn.
elegans, Opiz, l. c. = H. elegans.
hirsuta, Opiz, l. c. = H. hirsutum.
montana, Opiz, l. c. = H. montanum.
pulchra, Opiz, in Lotos, ii. (1852) 280 = H. pulchrum.

HYPERICUM, Tourn. ex Linn. Gen. ed. I. 281 (1737). *HYPERICINEÆ*, Benth. & Hook. f. i. 165.
ADENOSEPALUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 352.
ADENOTRIAS, Jaub. & Spach, Illustr. i. 76 (1842).
ANCISTROLOBUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 362.
ANDROSÆMUM, Tourn. ex Adans. Fam. ii. 444 (1763).
ASCYRIA, Choisy, Prod. Hyp. 38 (1821).
BRATHYDIUM, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 365.
BRATHYS, Mœnch, ex Linn. f. Suppl. 43 (1781).
CAMPYLOPUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 360.
CAMPYLOSPOROUS, Spach, l. c. 363 (1836).
DIOSAPHES, Spach, l. c. 365 (1836).
DROSOCARPIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 353.
ELODEA, Juss. Gen. 256, in obs. (1789) ; Vent. Tabl. iii. 144 (1799).
ELODES, Adans. Fam. ii. 444 (1763) ; Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 353.
EPISTIPIS, Raf. Fl. Tellur. iii. 78 (1836).
EREMANTHE, Spach, Hist. Vég. Phan. v. 421 (1836) ; et in Ann. Sc. Nat. Sér. II. v. (1836) 363.
EREMOPYRUM, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 365.
HOLOSEPALUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 352.
HYPERICOPSIS, Opiz, Seznam, 58 (1852).
IKOPHYLLUM, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367.
KNIPHOFIA, Adans. Fam. ii. 444 (1763).
KOMANA, Adans. l. c. (1763).
MARTIA, Spreng. Anleit. ii. ii. 788 (1818).
MYRIANDRA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 364.
NORYSCA, Spach, l. c. 363 (1836).
OLYMPIA, Spach, l. c. 369 (1836).
PANCALIUM, Ehrh. Beitr. iv. 147 (1789).
PERFORARIA, Choisy, Prod. Hyp. 44 (1821).
PETALANDRA, Raf. Fl. Tellur. iii. 80 (1836).
PLEURENDOPON, Raf. l. c. (1836).

HYPERICUM :—
PSOROPHYTUM, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 360.
RHEXYEURA, Vell. Fl. Flum. v. t. 119, 120 (1827).
ROSCYNA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 364.
SAROTHRA, Linn. Diss. Chen. 14 (1751) ; Amoen. Acad. iii. 11 (1787).
STREPTALON, Rafin. Fl. Tellur iii. 80 (1836)
THYMOPSIS, Jaub. & Spach, Illustr. i. 72 (1842).
TRIADENIA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 364.
TRIADENUM, Rafin. in Med. Repos. N. York, v. (1808) 352.
TRIDIA, Korth. in Hoev. & De Vriese, Tijdschr. iii. (1836) 17. t. 1.
TRIPENTAS, Casp. in Monatsb. Berl. Acad. 1857 (1858) 43 ; et in Pringsh. Jahrb. i. (1858) 497, in obs.
VISMIA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 356.
acerosum, H. B. & K. Nov. Gen. et Sp. v. 187. t. 457 = laricifolium.
aciculare, *H. B. & K. l. c.* 190.—Peruv.
acuminatum, Lam. Encyc. i. 150 = Vismia acuminata.
acutifolium, Ell. Sketch, ii. 26 = angulosum.
acutum, Mœnch, Meth. 138 = tetrapterum.
acutum, Wall. Cat. n. 4807 = cordifolium.
adenocarpum, Montbr. ex Spach, Hist. Vég. Phan. v. 395 = Montbretii.
adenocladum, *Boiss. Fl. Orient.* i. 809.—Europ. ; As. Min.
adenophorum, Wall. Cat. n. 4812 = elodeoides.
adenophyllum, Ledeb. Fl. Ross. i. 445 = olympicum.
adenotrichum, *Spach, Hist. Veg. Phan.* v. 405.—As. Min.
adpressum, *Barton, Comp. Fl. Philad.* ii. 15.—Am. bor.
aegyptium, Blanco, Fl. Filip. ed. I. 615 = Cratoxylon formosanum.
aegyptiacum, *Linn. Sp. Pl.* 784.—Aegypt.
aethiopicum, *Thunb. Prod. Pl. Cap.* 135.—Afr. austr.
affine, Steud. ex Oliver, Fl. Trop. Afr. i. 156 = Quartinianum.
afram, *Lam. Encyc.* iv. 166.—Reg. Mediterr.
Afzelii, Purdie, ex Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 579 = Psorospermum febrifugum ?
alatum, *Reta. Pugill.* 18.—Am. bor.
albens, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 136.—Ins. Teneriffe.
alpestre, Stev. ex Ledeb. Fl. Ross i. 451 = repens.
alpigenum, Kit. in Linnæa, xxxii. (1863) 550 = Richeri.
alpinum, Waldst. & Kit. Pl. Rar. Hung. iii. 294 = Richeri.
alternifolium, Labill. Ic. Pl. Syr. Dec. ii. 17 = Reaumuria Billardieri.
alternifolium, Vahl, Symb. Bot. ii. 85. t. 42 = Archytæa Vahlii.
amanum, *Boiss. Fl. Orient.* i. 804.—Syria.
amarisciunum, Steud. Nom. ed. II. i. 786, sphalm. = tamariscinum.
ambiguum, Ell. Sketch, ii. 30 = galioides.
ambiguum, Kit. in Linnæa, xxxii. (1863) 549 = quadrangulum.
amblysepalum, Hochst. in Lorent, Wanderungen, 841 = retusum.
amoenum, Pursh, Fl. Am. Sept. ii. 375 = aureum.
amplexicaule, Gilib. Fl. Lituan. ii. 304 = pulchrum.
amplexicaule, Lam. Encyc. iv. 147 = pyramidatum.
anagallidifolium, Boiss. Fl. Orient. i. 800 = tetrapterum.
anagallidifolium, Presl, Delic. Prag. 22 = elegans.
anagallidioides, Jaub. & Spach, Illustr. i. 47 = tetrapterum.
anagalloides, *Cham. & Schlecht. in Linnaea*, iii. (1828) 127.—Am. bor. occ.
androsæmifolium, Vill. Prosp. 44 = Richeri.
androsæmoides, Vill. Hist. Fl. Dauph. t. 44 = præc.
androsæmoides, *Wender. Ind. Hort. Mark.* (1856) 8. —*Cf. Linnæa*, xxix. (1857–58) 737.—Hab. ?
Androsæmum, *Linn. Sp. Pl.* 784.—Europ. ; As. Min.
anglicum, Bartol. Fl. Ital. viii. 310 = H. Androsæmum.
angulosum, *Michx. ex Willd. Sp. Pl.* iii. 1454.—Am. bor.
angustatum, Wall. ex Steud. Nom. ed. II. i. 787 = Buchanania florida.
angustifolium, Lam. Encyc. iv. 145 = lanceolatum.
angustifolium, Lowe, in Trans. Camb. Phil. Soc. iv. (1835) 58 = perforatum.

HYPERICUM :—
annulatum, *Moris, Stirp. Sard. Elench.* 9.—Ins. Sardin.
Apollinis, *Boiss. & Heldr. Diagn.* II. i. 105.—Graecia.
appressum, Steud. Nom. ed. II. i. 786 = adpressum.
apricum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 176.—Songaria.
arbuscosum, Vahl, Symb. Bot. ii. 85. t. 45 = Cratoxylon Blancoi.
arborescens, J. F. Gmel. Syst. 1156 = Vismia laurifolia.
aremariolides, *A. Rich. Ess. Fl. Cub.* 238.—Cuba.
armenum, *Jaub. & Spach, Illustr.* i. 65. t. 83.—Armenia.
articulatum, Lam. Encyc. iv. 159 = Elisea articulata.
ascyroides, Willd. Sp. Pl. iii. 1448 = pyramidatum.
Ascyron, *Linn. Sp. Pl.* 783.—Sibir.; Japon.
Ascyron, Mill Gard. Dict. ed. VIII. n. 7 = calycinum.
Ascyron, Siebold, ex Blume, Mus. Bot. Lugd. Bat. ii. 21 = japonicum.
aspalathoides, Willd. Sp. Pl. iii. 1451 = fasciculatum.
asperuloides, *Coerr. & Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 385.—Persia
asperulum, *Jaub. & Spach, Illustr.* i. 60.—Persia.
asperum, Ledeb. Ic. Pl. t. 17 = scabrum.
asperum, C. A. Mey. in Ledeb. Fl. Alt. iii. 266 = montanum.
assyriacum, *Boiss. Diagn.* Ser. I. viii. 116.—Mesopot.
athoum, *Boiss. & Orph. in Boiss. Fl. Orient.* i. 794.—Macedon.
atlanticum, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 56 = coadunatum.
atomarium, *Boiss. Diagn.* Ser. I viii. 114.—Graecia; As. Min.
attenuatum, *Choisy, Prod. Hyp.* 47. t. 6.—Sibir.; Dahur.
attenuatum, *Link, Enum. Hort. Berol.* ii. 275.—Hab.!
Aucheri, *Jaub. & Spach, Illustr.* i. 161. t. 61. t. 81.—Phrygia.
aureum, *Bartram, Trav.* 383; *ex Torr. & Gray, Fl. N. Am.* i. 76.—An. bor.
aureum, Lour. Fl. Cochinch. ii. 472 = chinense.
australe, *Tenore, Fl. Nap. Prod. App.* v. 25.—Reg. Mediterr.
avicularisfolium, *Jaub. & Spach, Illustr.* i. 69. t. 80.—As. Min.
axillare, Lam. Encyc. iv. 160 = galloides.
axillare, Michx. Fl. Bor. Am. ii. 81 = paludosum.
bacciferum, Lam. Fl. Fr. iii. 151 = H. Androsaemum.
bacciferum, Lin. Mant. ii. 277 = Vismia baccifera.
bacciferum, Moç. & Sesse, ex DC. Prod. i. 542 = Vismia guianensis.
baeticum, vide boeticum.
balearicum, *Linn. Sp. Pl.* 783.—Ins. Majorca.
barbatum, All. Fl. Pedem. ii. 46 = Richeri.
barbatum, *Jacq. Fl. Austr.* iii. 38.—Orient.
barbatum, *Linn. Amoen. Acad.* viii. 323.—Europ. or.
Bartramianum, Pursh, Fl. Am. Sept. ii. 374 = seq.
Bartramium, Mill. Gard. Dict. ed. VIII. n. 10 = Kalmianum.
Baumgartenianum, Schur, Enum. Pl. Transs. 138 = pulchrum.
bifarium, Choisy, Prod. Hyp. 38 = Tridesmis.
bifarum, Lam. Encyc. iv. 170 = Cratoxylon polyanthum.
bithynicum, *Boiss. Diagn.* Ser. I. viii. 119.—Bithynia.
blentinense, Bertol. Fl. Ital. viii. 340 = mutilum.
blentinense, Walp. Rep. i. 384, sphalm. = praec.
Bocconii, Tenore, ex Nym. Consp. 139 = perfoliatum.
boeticum, Boiss. Voy. Bot. Espagne, i. 114. t. 34 = undulatum.
Bonaparteae, Barton, Fl. N. Am. iii. 95 = adpressum.
bonariense, *Griseb. in Goett. Abh.* xxiv. (1879) 41.—Reg. Argent.
Bourgaei, Boiss. Fl. Orient. i. 796 = scabrum.
Bourgaei, Boiss. l. c. 811 = leptocaulon.
brachyphyllum, Steud. Nom. ed. II. i. 787 = fasciculatum.
bracteatum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 217; Wall. Cat. n. 4804 = cordifolium.
bracteatum, Kellogg, in Proc. Calif. Acad. ed. II. i. (1878) 67 = concinnum.
brasiliense, *Choisy, in DC. Prod.* i. 547.—Bras.
Brathydilium, Steud. Nom. ed. II. i. 786 = canadense.
Brathys, *Sm. Ic. Ined. Fasc.* ii. 41; *Lam. Encyc.* iv. 156.—N. Granat.
brevifolium, *Wall. Cat.* n. 4876.—Reg. Himal.

HYPERICUM :—
brevistylum, *Choisy, Prod. Hyp.* 51. t. 71.—Am. bor.
Buckleii, *M. A Curt. in Am. Journ. Sc.* xliv. (1843) 80.—Am. bor.
Buckleyi, S. Wats. Bibl. Ind. N. Am. Bot. 135 = prasc.
bupleuroides, *Griseb. in Wiegm. Archiv*, xviii. (1852) 299, in obs.—As. Min.
Burseri, DC. ex Choisy, in DC. Prod. i. 552, in syn. = fimbriatum.
Burseri, Spach, Hist. Vég. Phan. v. 397 = Richeri.
caespitosum, *Chamm. & Schlecht. in Linnaea*, iii. (1828) 136.—Chili.
caimanense, Steud. Nom. ed. I. 430 = *cayennense*.
calabricum, Spreng. Neue Entdeck. iii. 300 = barbatum.
callianthum, *Boiss Fl. Orient.* i. 800.—Persia.
callithyrsum, Coss. Notes Cent. 152 = hyposipholium.
calycatum, Jacquem. ex Dyer, in Hook. f. Fl. Brit. Ind. i. 256 = japonicum.
calycinum, *Linn. Mant.* i. 106.—Bithyn.; Reg. Caucas.
Cambessedesii, *Coss. ex Marès & Vigin. Cat. Pl. Balear.* 330. t. 8.—Ins. Balear.
campanulatum, Walt Fl. Carol. 191 = virginicum.
campestre, *Cham. & Schlecht. in Linnaea*, iii. (1828) 132.—Bras.
Campylopus, Steud. Nom. ed. II. i. 787 = rhodopeum.
camtschaticum, Steud. l. c. = kamtschaticum.
canadense, *Linn. Sp. Pl.* 785.—Am. bor.
canariense, Brouss. ex Webb & Berth. Phyt. Canar. i. 51 = grandiflorum.
canariense, Cambess. ex Marès & Vigin. Cat. Pl. Balear. 330 = Cambessedesii.
canariense, Hort. Par. ex Webb & Berth. l. c. 49 = platypetalum.
canarium, *Linn. Sp. Pl.* 784.—Ins. Canar.
canescens, Trev. Hyp. Animad. 10 = lusitanicum ?
capitatum, Choisy, Prod. Hyp. 57. t. 9 = helianthemoides.
caprifolium, *Cham. & Schlecht. in Linnaea*, iii. (1828) 125.—Bras.
caprifolium, *Boiss. Elench.* 26.—Hispan.
carassanum, *Willd. Sp. Pl.* iii. 1468.—Venezuela.
cardiophyllum, *Boiss. Fl. Orient.* i. 791.—Syria.
carinatum, *Griseb. in Goett. Abh.* xxiv. (1879) 41.—Reg. Argent.
carneum, Wall. Cat. n. 4820 = Cratoxylon polyanthum.
casium, *Boiss. Diagn.* Ser. I. viii. 111.—Syria.
cayanense, Lin. Mant. ii. 277 = seq.
cayanense, Jacq. Enum. Pl. Carib. 26 = Vismia cayennensis.
cemuum, *Roxb. Hort. Beng.* 59; *Fl. Ind.* iii. 400.—Reg. Himal.
Cervantesii, *Willd. ex Spreng. Syst.* iii. 847.—Mexic.
Chamaemyrtus, *Triana & Planch. in Ann. Sc. Nat. Sér.* IV. xviii. (1862) 298.—N. Granat.
Chamaeranium, Steud. Nom. ed. II. i. 787 = sphaerocarpum.
chilense, *C. Gay, Fl. Chil.* i. 355.—Chili.
chinense, Linn. Syst. ed. X. ii. 1184.—China.
chinense, Lour. Obs. Cochinch. v. 57 = Cratoxylon polyanthum.
chloraefolium, St. Hil. Fl. Bras. Mer. i. 329 = connatum.
Choisianum, Wall. Cat. n. 4805 = Hookerianum.
chrysanticum, *Webb, Fragm. Fl. Aethiop.* 94.—Afr. trop.
ciliatum, Lam. Encyc. iv. 170 = perfoliatum.
cistifolium, *Lam. l. c.* 168.—Am. bor.
coadunatum, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 153.—Reg. Mediterr. or.; Ins. Canar.
coccineum, Wall. Cat. n 4823 = Cratoxylon Blancoi.
cochinchinense, Lour. Fl. Cochinch. ii. 472 = Cratoxylon polyanthum.
collinum, *Cham. & Schlecht. in Linnaea*, v. (1830) 219.—Mexic.
connatatum, *Nolte, Novit. Fl. Holsat.* 69.—Europ.
concinnum, *Benth. Pl. Hartw.* 300.—Calif.
confertum, *Choisy, Prod. Hyp.* 55. t. 8.—As. Min.; Syria.
confertum, Moench, Meth. 129 = montanum.
connatum, *Lam. Encyc.* iv. 168.—Reg. Argent.
cordifolium, *Choisy, in DC. Prod.* i. 545.—Reg. Himal.
cordiforme, *A. St. Hil. Fl. Bras. Mer.* i. 330.—Bras.
coriaceum, *Blume, Bijdr.* 142.—Java.
coriaceum, Royle, *Illustr. Bot. Himal.* 191 = lysimachioides.
corifolium, Salisb. Prod. 389 = H. Coris.
Coris, Bieb. Desc. Prov. Casp. App. 210 = hyssopifolium.

HYPERICUM :—
Coris, *Linn. Sp. Pl.* 787.—Europ. austr.; Orient.
Coris, Sibth. & Sm. Fl. Graec. t. 777 = empetrifolium.
Coris, Walt. Fl. Carol. 190 = fasciculatum ?
coriscum, Steud. Nom. ed. II. i. 787 = tetrapterum.
corymbosum, Moench, Meth. Suppl. 41 = canariense.
corymbosum, *Muhl. ex Willd. Sp. Pl.* iii. 1457.—Am. bor.
crenulatum, *Boiss. Diagn.* Ser. II. v. 69.—Cilicia.
creticum, Hort. ex Link, Enum. Hort. Berol. ii. 276 = maritimum.
crispum, *Linn. Mant.* i. 106.—Reg. Mediterr.
Crux-Andreae, Crantz, Inst. ii. 540 = Ascyrum Crux-Andreae.
cryptopetalum, Boiss. & Bal. Diagn. Ser. II. vi. 39 = avicularisfolium
cymosum, Hochst. in Lorent, Wanderungen, 342 = scabrum.
debile, Salisb. Prod. 389 = canariense.
decandrum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) t. 389.—Ecuador.
decipiens, H. Wats. in Hook. Lond. Journ. Bot. iii. (1844) 588 = undulatum.
decorticans, Planch. ex Wedd. Chlor. And. ii. 272 = stenopetalum.
decumbens, *Peterm. Fl. Lips. Excurs.* 565.—Europ.
decussatum, Kunze, Del. Sem. Hort. Lips. i. 848.—Cf. Linnaea, xxiv. (1851) 199 = orientale.
dendesheimense, Sch. Bip. ex Trev. Hyp. Animad. 10 = perforatum.
delphicum, *Boiss. & Heldr. Diagn.* Ser. II. i. 106.—Graecia.
delphinense, Vill. Prosp. 44 = quadrangulum.
densiflorum, *Pursh, Fl. Am. Sept.* ii. 376.—Am. bor.
dentatum, Loisel. Fl. Gall. 499. t. 17 = perfoliatum.
denticulatum, *Ji. B. & K. Nov. Gen. et Sp.* v. 191. t. 458.—Mexic.
denticulatum, Wall. Cat. n. 4821 = angulosum.
dentiferum, *Steud. Nom.* ed. II. i. 787.—Ind. or.
denudatum, St. Hil. Fl. Bras. Mer. i. 336 = riqidum ?
Desetangsii, Lamotte, in Bull. Soc. Bot. Fr. xxi. (1874) 121 = quadrangulum ?
dichotomum, *Lam. Encyc.* iv. 167.—Ins. S. Doming.
dichotomum, *Phil. Fl. Atac.* 12.—Chili.
dichotomum, *Zumag. Fl. Pedem.* ii. 888.—Ital.
diomoides, *Griseb. Cat. Pl. Cub.* 40.—Cuba.
diversifolium, DC. Fl. Fr. Suppl. 631 = hyssopifolium.
dolabriforme, Steud. Nom. ed. II. i. 787 = sphaerocarpum.
dolabriforme, *Vent. Yard. Cels.* 45.—Am. bor.
Dodonaei, *Vill. Rè ex Steud. Nom.* ed. II. i. 787.—Italia.
Drummondii, Steud. Nom. ed. II. i. 787 = virginicum.
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 165.—Am. bor.
dubium, Leers, Fl. Herborn. 165 = quadrangulum.
dubium, Mauri, Rom. Pl. Cent. xiii. 27 = scabrum.
elatum, [*Doryand. in*] *Ait. Hort. Kew.* ed. I. iii. 104.—Am. bor.
electrocarpum, Maxim. in Bull. Acad. Pétersb. xii. (1868) 62 = Sampsoni.
elegans, Bertol. Fl. Ital. viii. 322 = perfoliatum.
elegans, *Steph. ex Willd. Sp. Pl.* iii. 1469.—Europ.; As. bor.
elegantissimum, Baumg. Enum. Stirp. Trans. ii. 391, in syn. = pulchrum.
elegantissimum, Crantz, Stirp. Austriac. ed. I. fasc. ii. 68 = montanum.
elliptícum, *Hook. Fl. Bor. Am.* i. 110.—Am. bor. occ.
elodeoides, *Choisy, in DC. Prod.* i. 551.—Reg. Himal.; Burma.
Elodes, *Huds. Fl. Angl.* ed. I. 292.—Europ.
elongatum, C. A. Mey. in Ledeb. Fl. Alt. iii. 267 = hyssopifolium.
emarginatum, Lam. Encyc. iv. 154 = virginicum.
empetrifolium, Kunzky, ex Jaub. & Spach, Illustr. i. 77 (= *Triadenia Kunzeyyeri*).—Syria.

HYPERICUM:—
empetrifolium, *Willd. Sp. Pl.* iii. 1452.—Graecia; As. Min.
enneandrum, Stokes, Bot. Mat. Med. iv. 97 = virginicum.
enodes, Neck. Delic. Gallo-Belg. ii. 319, sphalm. = H. Elodes.
erectum, Mass. ex Webb & Berth. Phyt. Canar. i. 51 = grandifolium.
erectum, *Thunb. Fl. Jap.* 296.—Japon.
ericoides, *Linn. Sp. Pl.* 785.—Europ. occ.
Erythraeae, Steud. Nom. ed. II. i. 787 = angulosum.
eugeniaefolium, Willd. ex Spreng. Syst. iii. 350 = Vismia acuminata, glabra.
euphorbiaefolium, Spach, Hist. Vég. Phan. v. 395 = Montbretia.
euphorbioides, A. St. Hil. Fl. Bras. Mer. i. 332. t. 69 = mutilum.
fallax, Grimm, in Nov. Act. Nat. Cur. iii. (1767) App. 362 = quadrangulum.
fasciculatum, *Lam. Encyc.* iv. 160.—Am. bor.
fasciculatum, Lapeyr. Hist. Abr. Pl. Pyr. 450 = hysopifolium.
fasciculatum, Michx. ex Willd. Sp. Pl. iii. 1452 = galloides.
fastigiatum, Ell. Sketch, ii. 31 = adpressum.
fastigiatum, *H. B. & K. Nov. Gen. et Sp.* v. 195.—Mexic.
ferrugineum, Banks, ex Pursh, Fl. Am. Sept. ii. 379 = hirsutum.
ferrugineum, Boiss. & Heldr. Diagn. Ser. II. i. 104 = Sprunerі.
filicaule, Hook. f. & Thoms. ex Dyer, in Hook. f. Fl. Brit Ind. i. 252 = Ascyrum filicaule.
fimbriatum, Lam. Encyc. iv. 148 = Richeri floribundum. [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 104.—Ins. Canar.
floribundum, Reichb. Ic. Exot. ii. 64. t. 95 = canariense.
floridum, Salisb. Prod. 389 = H. Androsaemum.
foetidum, Hook. f. & Thoms. ex Dyer, in Hook. f. Fl. Brit. Ind. i. 257 = Lalandii.
foliosum, Brouss. ex Webb & Berth. Phyt. Canar. i 45 = reflexum.
foliosum, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 104.—Ins. Azores.
foliosum, Jacq. Hort. Schoenb. iii. 27 = prolificum.
formosanum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 428.—Ins. Formosa.
formosum, A. Gray, Pl. Wright. ii. 17 = Scouleri.
formosum, *H. B. & K. Nov. Gen. et Sp.* v. 196. t. 460. —Mexic.
fragile, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. i. 108. —Graecia.
Fraseri, Steud. Nom. ed. II. i 788 = virginicum.
frondosum, Michx. Fl. Bor. Am. ii. 81 = aureum.
fulgidum, Rafin. Fl. Ludov. 68 = fasciculatum †
gallifolium, *Rupr. Fl. Casuc.* 245.—Reg. Caucas.
galloides, *Lam. Encyc.* iv. 161.—Am. bor.
galioides, Pursh, Fl. Am. Sept. ii. 376 = densiflorum.
Gebleri, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 864.— Sibir.
gheiwense, *Boiss. Fl. Orient.* i. 810.—Anatolia.
glandulosum, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 107.—Ins. Fortunat.
glandulosum, Gilib. Fl. Lituan. ii. 205 = montanum.
glaucum, Michx. Fl. Bor. Am. ii. 78 = myrtifolium.
goldiaefolium, *A. Rich. Tent. Fl. Abyss.* i. 98.— Abyssin.
goldioides, *Sesm. Bot. Voy. Herald*, 88.—Mexic.
Govanianum, Wall. Cat. n. 4815 = crnuum.
gracile, Boiss. Diagn. Ser. II. v. 70 = intermedium.
graminicum, *Forst. f. Prod.* 58.—N. Caled.
grandiflorum, Salisb. Prod. 389 = calycinum.
grandifolium, *Choisy, Prod. Hyp.* 38. t. 3.—Ins. Teneriff.
graveolens, *Buchl. in Am. Journ. Sc.* xlv. (1843) 174. —Am. bor.
Griffithii, *Hook. f. & Thoms. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 253.—Reg. Himal.
Grisebachii, Boiss. Fl. Orient. i. 815 = Richeri.
guianense, Aubl. Pl. Gui. ii. 784 = Vismia guianensis.
guineense, Lam. Amoen. Acad. viii. 321 = Vismia laeneala.
gymnanthum, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1847) 213 = mutilum.
gymopodioides, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 296.—N. Granat.

HYPERICUM:—
hakonense, *Franch. & Sav. Enum. Pl. Jap.* ii. 298.— Japon.
Hartwegii, *Benth. Pl. Hartw.* 127.—Peruv.
hecatophyllum, *Wright, in Sav. Fl. Cub.* 8.—Cuba.
hedyotifolium, Poir. Encyc. Suppl. iii. 700 = angulosum.
Heldreichii, Boiss. Diagn. Ser. I. viii. 111 = perfoliatum.
heliantheoides, *Boss. l. c.* 116.—As. Min.; Persia.
Helodes, St. Lag. in Ann. Soc. Bot Lyon, vi. (1880) 128 = H. Elodes.
helodeum, St. Lag. l. c = H. Elodes.
heterophyllum, *Vent. Yard. Cels.* t. 68.—Persia.
heterostylum, Parl. Fl. Ital. v. 550 = aegypticum.
hircinum, *Linn. Sp. Pl* 784.—Reg Mediterr.
hircinum, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 975 = inodorum.
hirsutum, Asso, Syn. Stirp. Arag. 107 = caprifolium.
hirsutum, *Linn. Sp. Pl.* 786 —Europ.; As. bor.
hirtellum, *Boiss. Diagn.* Ser. I. viii. 116.—Persia.
Hookerianum, *Wight & Arn. Prod.* 99.—Ind. or
horridum, Wall. Cat. n. 4822 = Cratoxylon polyanthum.
Humboldtianum, *Steud. Nom.* ed. II. i. 788.—Am. austr.
Humboldtii, Schlecht. ex Steud. l. c. = formosum.
humifusum, Linn. *Sp. Pl. 785* —Europ.; As. bor.
humifusum, Tenore, Fl. Nap. Prod. App. v. 25; ex e). Syll. 886 = australe.
Hypericum, Crantz, Inst. ii. 520 = Ascyrum Hypericoides.
hysopifolium, *Vill. Hist. Pl. Dauph.* iii. 805. t. 44.— Europ.; As. bor.
impunctatum, Hochst. ex Steud. Nom. ed. II. i. 788 = mutilum.
incertum, Steud. l. c. = angulosum.
indecorum, H. B & K. Nov. Gen. et Sp. v 198 = thesiifolium.
Inodorum, *Mill. Gard. Dict.* ed. VIII. n. 6; *Willd. Sp. Pl.* iii. 1449.—Graecia; As. Min.
intermedium, Bellynck, Fl. Namur. (1855) 31 = quadrangulum †
intermedium x, Retz. ex Nym. Consp. 138.—Europ.
Intermedium, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 95. —Abyssin.
involutum, Choisy, Prod. Hyp. 50 = gramineum.
Isophyllum, Steud. Nom. ed. II. i. 788 = Drummondii.
japonicum, *Thunb. Fl. Jap.* 295. t. 31.—As. temp. or.; Australas.
Jussiaei, Spach, in Jaub. & Spach, Illustr. i. 38. t. 118 = orientale.
javanicum, Blume, Bijdr. 141 = patulum.
juniperinum, H. B. & K. Nov. Gen. et Sp. v. 189 = H. Brathys.
Jussiaei, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 297.—N. Granat.
Kalmianum, Durol, Harbk. Baums. i. 310 = densiflorum.
Kalmianum, *Linn. Sp. Pl.* 788.—Am. bor.
Kalmianum, Vahl, Symb. Bot. i. 66 = seq.
Kalmii, Forsk. Fl. Aegypt. Arab. p. cxviii = revolutum.
kamtschaticum, *Ledeb. in Denkschr. Bot. Ges. Regensb.* iii. (1841) 181.—Kamtschat.
Kohlianum, Spreng. Fl. Hal. 214. t. 9 = elegans.
kotschyanum, *Boiss. Diagn.* Ser. I. i. 56.—As. Min.; Syria.
laeve, *Boiss. & Hausskn. in Boiss. Fl. Orient.* i. 796.— As. Min.; Syria.
lanigerum, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 106 = angulosum †
Lalandii, *Choisy, in DC. Prod.* i. 550.—Ind. or.; Afr. trop.
lanatum, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 296.—Guian.
lancealatum, *Lam. Encyc.* iv. 145.—Ins. Mascar.
lancealatum, Steud. Nom. ed. I. 420, 888 = Vismia glabra.
lanuginosum, *Lam. Encyc.* iv. 171.—As. Min.; Syria.
laricifolium, *Juss. in Ann. Mus. Par.* iii. (1804) 160. t. 16. f. 1.—Peruv.
Lasianthus, Linn. Sp. Pl. 783 = Gordonia Lasianthus.
latifolium, Aubl. Pl. Gui. ii. 787 = Vismia latifolia.
laurifolium, Willd. Sp. Pl. iii. 1440 = Vismia laurifolia.

HYPERICUM:—
lauriforme, Lam. Encyc. iv. 152 = Vismia laurifolia.
laurocerasifolium, Jacq. Obs. Bot. iii. 4 = Symplocos tinctoria.
laxiusculum, *A. St. Hil. Pl. Us. Bras.* t. 62.—Bras.
Leconteanum, Steud. Nom. ed. II. i. 788 = paludosum
ledifolium, Steud. l. c. 789 = undiflorum.
Leersii, C. C. Gmel. Fl. Bad. iii. 352 = quadrangulum.
leprosum, *Boiss. Diagn.* Ser. I. i. 58.—As. Min.
leptocladium, *Boiss. Fl. Orient.* i. 803.—Mesopotam.
leptophyllum, *Hochst. in Lorent, Wanderungen*, 342. —Oriens.
Leschenaultii, Choisy, in DC. Prod. i. 545 = Hookerianum.
leucoptychodes, Steud. ex A. Rich. Tent. Fl. Abyss. i. 96 = lanceolatum.
Lhotskyanum, *Steud. Nom.* ed. II. i. 788.—Bras.
ligustrinum, Banks, ex Pursh, Fl. Am. Sept. n. 375 = undiflorum.
limosum, *Griseb. Cat. Pl. Cub.* 20.—Cuba.
linarifolium, Bertol. Fl. Ital. viii. 328 = australe.
linarifolium, *Vahl, Symb. Bot.* i. 65.—Europ. occ.
linearifolium, Willd. Sp Pl iii. 1470 = linarifolium.
linearioides, Bosse, in Otto & Dietr. Allg. Gartenz. iii. (1835) 99 = repens.
lanoiatum, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 343 = perforatum.
linoides, *A. St. Hil. Fl. Bras. Mer.* i. 338.—Bras.
Liottardii, Vill Prosp. 44 = humifusum.
Lorentii, Hochst. in Lorent, Wanderungen, 342 = asperulum.
loxense, *Benth. Pl. Hartw.* 126.—Am. austr.
lucidum, Wall. ex Steud. Nom. ed. II. i. 788 = Ioxanthes icosandra †
Lungnanum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 218 = cordifolium.
lusitanicum, *Poir. Encyc. Suppl* iii. 702.—Lusitan.
lycopodioides, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 296.—N. Granat.
lydium, Boiss. Diagn. Ser. I i. 57 = hyssopifolium.
lysimachioides, Boiss. & Noé, l. c. Ser. II. i. 106 = callianthum.
lysimachioides, Wall. *Cat.* n. 4817.—Reg. Himal.
lythrifolium, Boiss. Fl. Orient. i. 800 = hyssopifolium.
macedonicum, Boiss. & Orph. Diagn. Ser. II. vi. 38 = barbatum.
macrocarpum, Michx. Fl. Bor Am. ii. 82 = pyramidatum.
macrocarpum, Wall. ex Steud. Nom. ed. II. i. 788 = Ixoxanthes reticulata.
maculatum, Crantz, Stirp. Austr. ed. II. fasc. ii. 98 = quadrangulum.
maculatum, Michx. Fl. Bor. Am. ii. 80 = corymbosum.
maculatum, Orsin. ex Nym. Consp. 132 = Richeri.
maculatum, *Walt. Fl. Carol.* 189.—Am. bor.
maculatum, Willd. ex Steud. Nom. ed. II. i. 788 = formosum.
madagascariense, *Steud. l. c.*—Madag.
maesicum, G. Don, Gen. Syst. i. 510 = *moesicum*.
martinum, *Sieber, Reise Kreta*, ii. 323.— Ins. Creta.
maximum, Peterm. Fl. Lips. Excurs. 568 = perforatum.
mexicanum, *Linn. f. Suppl.* 246.—Am. austr.
Michauxii, Poir. Encyc. Suppl. iii. 696 = galioides.
micranthum, Choisy, Prod. Hyp. 44. t. 5 = corymbosum.
micranthum, Wall. ex Steud. Nom. ed. II. i. 788 = Tristania burmanica.
microcalycinum, Boiss. & Heldr. Diagn. Ser. I. viii. 115 = hysopifolium.
microphyllum, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 341 = perforatum.
microsepalum, *A. Gray, in S. Wats. Bibliog. Ind. N. Am. Bot.* ii. 465.—Am. bor.
milleporum, *Willd. ex Spreng. Syst.* iii. 348.—Ins. Teneriff.
moesicum, Spreng. l. c. = aviculariaefolium.
monanthemum, *Hook. f. & Thoms. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 256.—Reg. Himal.
monogynum, Linn. Sp. Pl. ed. II. 1107; Thunb. Fl. Jap. 297 = chinense.
montanum, *Linn. Fl. Suec.* ed. II. 266.—Europ.
montanum, Urria, Hort. Reg. Panorm. 526 = perfoliatum.
Montbretii, *Spach, Hist. Vég. Phan.* v. 395.— Graecia.

HYPERICUM:—
moranense, *H. B. & K. Nov. Gen. et Sp.* v. 198.—Mexic.
multicaule, Lam. Encyc. iv. 178 = empetrifolium.
multiflorum, H. B. & K. Nov. Gen. et Sp. v. 194 = thesiifolium.
myricoides, *Phil. in Linnæa*, xxxiii. (1864–65) 31.—Chili.
mutilum, *Linn. Sp. Pl* 787.—Am. bor.
mutilum, Maxim. in Bull. Acad. Pétersb. xxvii. (1861) 486 = japonicum.
Mutisianum, H. B. & K. Nov. Gen. et Sp. v. 185 = mexicanum.
myrtanthum, *Cham. & Schlecht. in Linnæa*, iii. (1828) 123.—Bras.
myrtifolium, Boiss. Fl. Orient. i. 795 = adenotrichum.
myrtifolium, *Lam. Encyc.* iv. 180.—Am. bor.
myrtifolium, Spach, Hist. Vég. Phan. v. 399 = perfoliatum.
myrtifolioides, Fenzl, Pugill. Pl. Nov. Syr. 7 = cuneatum.
myscrense, *Heyne, in Wall. Cat.* n. 4808.—Ind. or.
nanum, *Pers. Encyc. Suppl.* iii. 699.—Syria.
napaulense, *Choisy, in DC. Prod.* i. 562.—Ind. or.
Naudinianum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 808.—Algeria.
neapolitanum, Tenore, Ind. Sem. Hort. Neap. (1829) 19 = tetrapterum.
nepalense, Hort. ex C. Koch, Dendrol. i. 497 = patulum.
nervifolium, Wall. Cat. n. 4824 = Cratoxylon neriifolium.
nervatum, Hance, in Walp Ann. ii. 168 = japonicum.
nervosum, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 151, 155.—Java.
nervosum, D. Don. Prod. Fl. Nep. 219 = elodeoides
neurocalycinum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 114.—As. Min.
nigricans, *Zucc. ex G. Don, Gen. Syst.* i. 612.—Hab.?
nitidum, Lam. Encyc. iv. 180 = fasciculatum.
Nordmanni, *Boiss. Fl. Orient.* i. 815.—Reg. Caucas.
Noryxca, Steud. Nom. ed. 11. i. 789 = myxcense.
nubigenum, Lowe, in Hook. Journ. Bot. i. (1834) 48 = linarifolium.
nudicaule, Walt. Fl. Carol. 190 = H. Sarothra.
nudiflorum, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 199 = clatifolium.
nudiflorum, *Michx. ex Willd. Sp. Pl.* iii. 1456.—Am. bor.
nummularioides, *Trauto. in Bull. Acad. Pétersb.* x. (1866) 894.—Reg. Caucas.
nummularium, *Linn. Sp. Pl.* 787.—Mont. Pyren.
Nuttallii, G. Don, Gen. Syst. i. 607 = setosum.
oblongifolium, Choisy, Prod. Hyp. 42. t. 4 = cernuum.
oblongifolium, Hook. Bot. Mag. t. 4949 = Hookerianum.
oblongifolium, Wall. Pl. As. Rar. t. 244 = patulum.
obovatum, Wall. ex Steud. Nom. ed. 11. i. 789 (Quid?). —Singapore.
obtusatum, *Fisch. & Lovillf, in Demid. Voy. Russ. Mér.* ii. 207.—Tauria.
obtusum, Dulac, Fl. Hautes-Pyr. 325 = Richeri.
obtusum, Moench, Meth. 129 = quadrangulum.
officinale, Gater. ex Steud. Nom. ed. 1. 421 = perforatum.
officinarum, Crantz, Stirp. Austr. ed. 1. fasc. ii. 66; ed. II. 99 = perforatum.
oliganthum, *Franch. & Sav. Enum. Pl. Jap.* ii. 299. —Japon.
Olivieri, *Boiss. Fl. Orient.* i. 803.—Mesopotam.
olympicum, Blanco, Fl. Filip. ed. I. 618 = clatifolium.
olympicum, *Linn. Sp. Pl.* 784.—Græcia; As. Min.
olympicum, Sm. Fl. Græc. Prod. ii. 115 = adenotrichum.
opacum, Torr. & Gray, Fl. N. Am. i. 168, 673 = clatifolium.
orientale, *Linn. Sp. Pl.* 785.—As. Min.
origanifolium, Urv. in Mém. Soc. Linn. Par. i. (1822) 514 = rhodopeum.
origanifolium, *Willd. Sp. Pl.* iii. 1467.—As. Min.
pallens, D. Don, Prod. Fl. Nep. 219 = napaulense.
pallens, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 270.—As. Min.
paludosum, *Choisy, Prod. Hyp.* 43.—Am. bor.
palustre, Salisb. Prod. 389 = H. Elodes.
paniculatum, *H. B. & K. Nov. Gen. et Sp.* v. 195. t. 459. —Mexic.
paniculatum, *Lam. Encyc.* iv. 181.—Hab.?
papillare, Boiss. & Heldr. Diagn. Ser. I. viii. 110 = leprosum.

HYPERICUM,—
parviflorum, Salisb. Prod. 389 = Ascyrum Crux-Andreæ.
parviflorum, *A. St. Hil. Fl. Bras. Mer.* i. 333.—Bras.
parvifurum, Willd. Sp. Pl. iii. 1456 = mutilum.
patulum, *Thunb. Fl. Jap.* 295.—Reg. Himal.; Chin.; Japon.
pauciflorum, *H. B. & K. Nov. Gen. et Sp.* v. 192.—Mexic.
pedicellatum, Endl. in Enum. Pl. Hueg. 19 = gramineum.
pedunculosum, Wall. ex Steud. Nom. ed. II. i. 789 = Ixonanthes khasiana.
Pelletierianum, *A. St. Hil. Fl. Bras. Mer.* i. 334. t. 70. —Bras.
pentandrum, *Blanco, Fl. Filip.* ed. I. 614.—Ins. Philipp.
Peplicoides, Comm. ex Lam. Encyc. iv. 145 = lanceolatum.
peplidifolium, *A. Rich. Tent. Fl. Abyss.* i. 95.—Abyssin.
perfoliatum, Ledeb. in Bull. Sc. Acad. Pétersb. ii. (1837) 314 = bupleuroides.
perfoliatum, *Linn. Syst.* ed. XII. 510.—Græcia; As. Min.
perfoliatum, Munby, in Bull. Soc. Bot. Fr. ii. (1855) 283 = Naudinianum
perforatum, *Linn. Sp. Pl.* 785.—Europ.; As. bor.; Am. bor.
Pestalozzæ, *Boiss. Diagn.* Ser. I. viii. 113.—As. Min.
petiolatum, Linn. Sp. Pl. ed. II. 1102 = Vismia petiolata.
petiolatum, Linn. f. Suppl. 345 = Vismia laurifolia.
petiolatum, Lour. Fl. Cochinch. 472 = Cratoxylon polyanthum.
petiolatum, Walt. Fl. Carol. 191 = paludosum.
petiolulatum, *Hook. f. & Thoms. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 255.—Reg. Himal.
philonotis, *Cham. & Schlecht. in Linnæa*, v. (1830) 219.—Mexic.
pilosum, Walt. Fl. Carol. 190 = setosum.
pimeleoides, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 299.—N. Granat.
pimentifolium, Walt. ex Steud. Nom. ed. II. i. 789 = Vismia Wightiana.
platypetalum, Steud. l. c. = canariense.
platypetalum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 388.—Ecuador.
platypsepalum, Walp. Rep. i. 386 = *platypetalum*.
polyanthemum, *Klotzsch, ex H. Reich. in Mart. Fl. Bras.* xii. i. 199.—Bras.
polygonifolium, *Rupr. Fl. Cauc.* 247.—Reg. Caucas.
polyphyllum, *Boiss. & Bal. Diagn.* Ser. II. v. 68.—Cilicia.
pratense, *Cham. & Schlecht. in Linnæa*, v. (1830) 218. —Mexic.
procumbens, Desf. ex Willd. Sp. Pl. iii. 1450 = floridiiforme.
prolificum, *Linn. Mant.* i. 106.—Am. bor.
prunifolium, Wall. Cat. n. 7276 = Cratoxylon prunifolium.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 481.—As. centr.
Pseudo-Brathys, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 387 = struthiolæfolium.
Pseudo-patulum, Hort.—Cf. Gard. Chron. (1882) ii. 149 = patulum.
Pseudo-perforatum, Bertol. Misc. Bot. iii. 14 = perforatum.
ptarmicæfolium, Spach, Hist. Vég. Phan. v. 404 = orientale.
pubescens, Boiss. Elench. 98 = tomentosum.
pulchellum, Willd. Enum. Hort. Berol. Suppl. 58; Wall. Cat. n. 4821 = Cratoxylon polyanthum.
pulchrum, Aubry, ex Steud. Nom. ed. I. 421 = linarifolium.
pulchrum, *Linn. Sp. Pl.* 786.—Europ.
pulchrum, Pall. Ind. Taur.; ex Bieb. Fl. Taur. Cauc. ii. 230 = elegans.
pulverulentum, Fenzl, Pugill. Pl. Nov. Syr. 7 = origanifolium.
punctatum, Lam. Encyc. iv. 184 = maculatum.
punctatum, Reichb. fc. Exot. Bot. 61. t. 88 = corymbosum.
punctatum, Willd. Sp. Pl. iii. 1464 = sviculariæfolium.
punctulatum, A. St. Hil. Fl. Bras. Mer. i. 334 = brasiliense.

HYPERICUM,—
punctulosum, Bertol. Misc. Bot. xiii. 18 = clatifolium.
pusillum, Choisy, Prod. Hyp. 50 = japonicum.
pyramidatum, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 108.—Am. bor.
quadrangulare, Sm. Engl. Bot. t. 870; O. F. Muell. in Fl. Dan. t. 840 = tetrapterum.
quadrangulare, Willd. Sp. Pl. iii. 1459 = quadrangulum.
quadrangulum, Colon. Rec. Bot. Calcic. 10 = undulatum.
quadrangulum, *Linn. Sp. Pl.* 785.—Europ.
quadratum, Stokes, Bot. Mat. Med. iv. 99 = quadrangulum.
quadrialatum, Wahlberg, Fl. Gothob. 75 = tetrapterum.
quadrifarium, Krock. Fl. Siles. Suppl. ii. 223 (Quid?). —Siles.
Quartinianum, *A. Rich. Tent. Fl. Abyss.* i. 97.—Abyssin.
quinquenervium, Walt. Fl. Carol. 190 = mutilum.
ramosissimum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 580.—Peruv.
ramosissimum, Hort. ex C. Koch, Dendrol. i. 497 = patulum.
ramosissimum, Ledeb. Fl. Ross. i. 449 = inodorum.
rariflorum, *Stev. ex Rupr. Fl. Cauc.* 298.—Reg. Caucas.
recognitum, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 36. t. 1 = rhodopeum.
reflexum, *Linn. f. Suppl.* 346.—Ins. Teneriffæ.
repandum, *Kit. & Rochel, Pl. Banat. Rar.* 26, nomen. —Banat.
repens, *Duby, ex Nym. Consp.* 133 = tetrapterum.
repens, Georgi, Beschr. Russ. Reich. iv. 1200 = humifusum.
repens, *Linn. Cent. i.* 26 (sp. dub.).—Europ.; Oriens.
repens, Forr. Voy. Barb. ii. 224 = australe.
repens, Sibth. Fl. Græc. t. 775 = trichocaulon.
repiatum, *Hook f. & Thoms. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 255.—Reg. Himal.
resinosum, Benth. Pl. Hartw. 165 = mexicanum.
reticulatum, Poir. Encyc. Suppl. iii. 694 = Vismia latifolia.
retusum, *Auch. in Jaub. & Spach, Illustr.* i. 53. t. 27. —Syria.
revolutum, *Vahl, Symb. Bot.* i. 66.—Arabia.
rhodopeum, *Frivald. in Flora*, xix. (1836) 486.—Rumel.; As. Min.
Richeri, Ledeb. Fl. Ross. i. 449 = Nordmanni.
Richeri, Rochel, Pl. Rar. Banat. 49. t. 12. f. 27 = barbatum.
Richeri, *Vill. Prosp.* 44; *Hist. Pl. Dauph.* iii. 501.—Europ. med.
rigidum, *A. St. Hil, Fl. Bras. Mer.* i. 336.—Bras.
Rochelii, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 299 = Richeri.
Roeperianum, Schimp. ex A. Rich. Tent. Fl. Abyss. i. 96 = Quartinianum.
romaninifolium, Choisy, in DC. Prod. i. 547 = myrtifolium.
romaninifolium, Klotz. ex Walp. Rep. i. 386 = fasciculatum.
romaninifolium, Lam. Encyc. iv. 169 = clatifolium.
rotundatum, Schur, in Verh. Naturf. Ver. Brunn, xv. ii. (1877) 159 = tetrapterum.
rostratum, Rafin. Fl. Ludov. 88 = densiflorum.
rubrum, Hochst. in Loresl, Wanderungen, 343 = læve.
rubrum, Wight, ex Dyer, in Hook. f. Fl. Brit. Ind. i. 285 = humifusum.
rufescens, *Klotzsch, ex H. Reich. in Mart. Fl. Bras.* xii. i. 194.—Bras.
rufescens, Lam. Encyc. iv. 180 = Vismia rufescens.
Rupellianum, Kunze, Del. Sem. Hort. Lips. (1847); ex Linnæa, xxiv. (1851) 177 = aureum.
rumelicum, *Boiss. Diagn.* Ser. I. viii. 113.—Macedon.
rupestre, *Yusk & Spach, Illustr.* i. 44.—Cilicia.
rutilum, D. Dietr. Syn. Pl. iv. 1295, sphalm. = mutilum.
sabinæfolium, *Trev. Hyper. Animad.* 15.—N. Granat.
Salicaria, Reichb. Fl. Crit. v. 53. t. 490. f. 683 = H. Ascyron.
salicifolium, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1848) 182 = chinense.
Sampsoni, *Hance, in Journ. Bot.* iii. (1865) 378.—China.
Sarothra, *Michx. Fl. Bor. Am.* ii. 79.—Am. bor.
saturejæfolium, Jaub. & Spach, Illustr. i. 55. t. 28 = scoparium.
scabrellum, *Boiss. Diagn.* Ser. II. v. 69.—As. Min.

HYPERICUM:—
scabridum, *Steud. Nom.* ed. II. i. 789.—Graecia.
scabrum, *Linn. Cent.* i. 26; *Amoen. Acad.* iv. 287.—As. Min.; Persia; Songar.
scaturiginum, Buch.-Ham. ex Wall. Cat. n. 4871 = japonicum.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 330.—Mexic.
Schimperi, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 99; *et ex A. Rich. Tent. Fl. Abyss.* i. 97.—Afr. trop.
Schlechteri, Hensl. in Flora, lxxvi. (1893) 628 = perforatum.
scopulorum, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 502.—Ins. Socotra.
Scouleri, *Hook. Fl. Bor. Am.* i. 111.—Am. bor. occ.
sebastaeum, Boiss. & Noé, Diagn. Ser. II. i. 107 = hyssopifolium.
Seliberianum, Klotzsch, ex H. Reich. in Mart. Fl. Bras. xii. i. 189 = rigidum.
Seplawini, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 434.—China.
septinerveum, Krock. Fl. Siles. Suppl. ii. 220 (Quid?). —Siles.
serpyllifolium, *Lam. Encyc.* iv. 176.—Syria.
sessiliflorum, Willd. ex Spreng. Syst. iii. 846 = myrtifolium.
sessilifolium, Aubl. Pl. Gui. ii. 787 = Vismia rufescens.
setosum, *Linn. Sp. Pl.* 787.—Am. bor.
setosum, Wall. Cat. n. 4814 = napaulense.
silenoides, H. B. & K. Nov. Gen. et Sp. v. 191 = thesiifolium.
silenoides, *Juss. in Ann. Mus. Par.* iii. (1804) 162. t. 16. f. 8.—Peruv.; N. Granat.
simplex, Michx. Fl. Bor. Am. ii. 80 = setosum.
sinaicum, *Hochst. ex Boiss. Fl. Orient.* i. 808 — Arabia.
sinense, J. F. Gmel. Syst. 1157 = *chinense*.
Skoulerı, Welp. Rep. i. 887, sphalm. = Scouleri.
songaricum, Ledeb. ex Spreng. Syst. iv. Cur. Post. 297 = perforatum.
Spachianum, *Steud. Nom.* ed. II. i. 789.—Hab.?
spuriflorum, *Schur, in Verh. Naturf. Ver. Brunn*, xv. 11. (1877) 158.—Transylv.
spathulatum, Steud. Nom. ed. II. i. 789 = prolificum.
speciosum, Wall. Cat. n. 4808 = cernuum.
spectabile, *Jaub. & Spach, Illustr.* i. 57. t. 29.—Syria.
sphaerocarpum, Barton, Comp. Fl. Philad. ii. 14 = elliptícum.
sphaerocarpum, *Michx. Fl. Bor. Am.* ii. 78.—Am. bor. Spruneri, *Boiss. Diagn.* Ser. I. viii. 112.—Graecia.
stellaroides, H. B. & K. Nov. Gen. et Sp. v. 196 = mutilum.
stenobotrys, Boiss. Diagn. Ser. I. viii. 115 = confertum.
stenopetalum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 355.—Venezuela.
stenopykilum, Opiz, Natural. n. ix. (1825) 158 = perforatum.
strictum, H. B. & K. Nov. Gen. et Sp. v. 190 = struthiolaefolium.
struthiolaefolium, *Juss. in Ann. Mus. Par.* iii. (1804) 161. t. 16. f. 2.—Peruv.
struthiolaefolium, Trev. Hyper. Animad. 15 = Humboldtianum.
stypheloides, *A. Rich. Ess. Fl. Cub.* 287.—Ind. occ.
suberosum, Salzm. ex Boiss. Voy. Bot. Espagne, ii. 115 = tomentosum.
sumatranum, *Miq. Pl. Jungh.* 395.—Sumatra.
tamaricinum, Cham. & Schlecht. in Linnaea, iii. (1828) 134 = Pallenianum.
targuense, H. B. & K. Nov. Gen. et Sp. v. 198 = thesiifolium.
tauricum, Hort. ex Ledeb. Fl. Ross. i. 450 = montanum.
tenellum, *Jacub. in Oestr. Bot. Zeitschr.* xxii. (1872) 175.—Europ.
tenellum, Kotschy, ex Boiss. Fl. Orient. i. 794 = cinereum.
tenellum, [Soland. ex] Clarke, Trav. ii. 644 = serpyllifolium.
tenellum, Tausch, in Flora, xiv. (1831) 211 = tetrapterum.
teneriusculum, *Hook. f. & Thoms. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 294.—Reg. Himal.
tenuifolium, Forsk, Fl. Aeg. Arab. Sept. ii. 877 = fasciculatum.
tenuifolium, *A. St. Hil. Fl. Bras. Mer.* i. 287.—Bras. meridionalium, *A. St. Hil. l. c.* 361.—Bras.
ternum, *A. St. Hil. Fl. Bras. Mer.* i. 330.—Bras.
tetrapetalum, Lam. Encyc. iv. 158 = Ascyrum amplexicaule.
tetrapterum, *Fries, Nov. Fl. Suec.* 256.—Europ.; As. bor.
thasium, *Griseb. Spicil. Fl. Rumel.* i. 227.—Macedon.
thesiifolium, *H. B. & K. Nov. Gen. et Sp.* v. 192.—Mexic.
Thunbergii, *Franch. & Sav. Enum. Pl. Jap.* ii. 300. —Japon.
thuyoides, *H. B. & K. Nov. Gen. et Sp.* v. 187. t. 456.—N. Granat.
thymbraefolium, *Boiss. & Nol, Diagn.* Ser. II. i. 107.—Anatolia.
thymifolium, H. B. & K. Nov. Gen. et Sp. v. 186. t. 455 = Humboldtianum.
thymifolium, [*Soland. in*] *Russell, Aleppo.* ed. II. ii. 270.—Cilicia.
thymopsis, *Boiss. Diagn.* Ser. II. i. 109.—Mesopotam.
tinctorum, Willd. ex Spreng. Syst. iii. 845 = laricifolium.
tomentosum, Durand, ex Steud. Nom. ed. I. 421 = H. Elodes
tomentosum, *Linn. Sp. Pl.* 786.—Reg. Mediterr.
tomentosum, Moris, Fl. Sardoa, i. 322. t. 21 = annulatum.
tortuosum, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 502.—Ins. Socotr.
Tournefortii, Spach, Hist Vég. Phan. v. 404 = orientale.
trachyphyllum, Griseb. Spicil. Fl. Rumel. i. 222 = leprosum.
transilvanicum, Celak, in Oestr. Bot. Zeitschr. xxiv. (1874) 188 = Richeri.
trichanthum, Boiss. & Sprun. Diagn. Ser. II. i. 105 = barbatum.
trichocaulon, *Boiss. & Heldr. l. c.* Ser. I. viii. 110. —Creta.
Tridesmis, *Steud. Nom.* ed. II. i. 790.—Ins. Bouton.
triflorum, Blume, Bijdr. 142 = Hookerianum.
triplinerve, *Vent. Navd. Cels.* 58. t. 58.—Am. bor.
triquetrifolium, Turra, Farset. Nov. Gen. 12 = crispum.
tubulosum, Walt. Fl. Carol. 191 = virginicum.
tymphaeostemum, Boiss. & Sprun. Diagn. Ser. I. i. 57 = hyssopifolium.
uliginosum, H. B. & K. Nov. Gen. et Sp. v. 194 = thesiifolium.
umbellatum, *Kern. in Oestr. Bot. Zeitschr.* xiii. (1863) 141.—Europ.
undulatum, *Schousb. ex Willd. Enum. Hort. Berol.* 810.—Hispan.
uniflorum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 109.—As. Min.
uralum, Buch.-Ham. ex D. Don, in Bot. Mag. t. 2875 = patulum.
velutinum, Boiss. Fl. Orient. i. 801 = Kotschyanum.
venosum, Lam. Encyc. iv. 146 = calycinum?
venustum, *Vensl, Pugill. Pl. Nov. Syr.* 7.—As. Min.
veronense, Schrank, ex Link, Enum. Hort. Berol. ii. 276 = perforatum.
verruculosum, Sallsb. Prod. 369 = balearicum.
verticillatum, Lam. Fl. Fr. iii. 149 = H. Coris.
verticillatum, *Thunb. Prod. Pl. Cap.* 188.—Afr. austr.
vesiculosum, *Griseb. Spicil. Fl. Rumel.* i. 226.—Graecia.
villosum, Crantz, Inst. ii. 620 = setosum.
villosum, Crantz, Stirp. Austr. ed. I. fasc. ii. 62; ed. II. 96 = hirsutum.
virgatum, Lam. Encyc. iv. 158 = angulosum.
virgatum, [*Soland. ex*] *Clarke, Trav.* ii. 262.—As. Min.
virginicum, *Linn. Syst.* ed. X. 1164.—Am. bor.
virginicum, Walt. Fl. Carol. 189 = corymbosum.
vulgare, Lam. Fl. Fr. iii. 151 = perforatum.
Walkeri, J. F. Gmel. Syst. 1159 = paludosum.
Webbii, Nym. Syll. 222 = aegyptícum.
Webbii, *Steud. Nom.* ed. II. i. 790.—Malta; Orient.
Wightianum, Wall. Cat. n. 4818 = napaulense.

HYPEROGYNE, Salisb. Gen. Pl. Fragm. 81 (1866) = **Paradisia**, Mazz. (Liliac.).

HYPERTELIS, E. Mey. ex Fenzl, in Ann. Wien. Mus. ii. (1839) 261 = **Pharnaceum**, Linn. (Ficoid.).
arenicola, Sond. in Harv. & Sond. Fl. Cap. i. 145.—Afr. austr.

HYPERTELIS:—
Bowkeriana, Sond. in Harv. & Sond. Fl. Cap. i. 145 — Afr. austr.
spergulacea, E. Mey. ex Fenzl, in Ann. Wien. Mus. ii. (1839) 263.—Afr. austr.
verrucosa, Fenzl, l. c. 262 = P. verrucosum.

HYPERUM, Presl, Epim. Bot. 211 (1849) = **Wendtia**, Meyen (Geraniac.).
trifidum, Presl, l. c. = W. gracilis.

HYPHAENE, Gaertn Fruct. ii. 18. t. 82 (1788) *PALMAE*, Benth. & Hook. f. iii. 940.
CUCIFERA, Delile, Fl. Egypte, 145 (1813).
DOMA, Lam. Illustr. i. 900 (1823).
DOUMA, [Polr.] in Nouv. Duham. iv. 47 (1809).
Argun, *Mart. Hist Nat. Palm.* iii. 227.—Nubia.
benguelensis, *Welw ex H. Wendl. in Bot. Zeit.* xxxix. (1881) 92.—Afr. trop.
compressa, *H. Wendl. l. c.* xxxvi. (1878) 116.—Afr. trop.
coriacea, *Gaertn. Fruct.* i. 28. t. 10. f. 2.—Afr. trop.
crinita, *Gaertn. l. c.* ii. 18.—Afr. austr.
Cucifera, *Pers. Syn.* ii. 628.—Afr. trop.
guineensis, Schum. & Thonn Beskr. Guin. Pl. 445 = thebaica.
macrosperma, *H. Wendl in Bot. Zeit.* xxxix. (1881) 92.—Afr. austr.
natalensis, Kunze, in Linnaea, xx. (1847) 15 = crinita.
Peteriana, Klotzsch, ex Mart. Hist. Nat. Palm. iii. 227 = crinita.
Schatan, *Boj Hort. Maurit.* 308.—Hab.?
thebaica, *Mart. Hist. Nat. Palm.* iii. 226. tt. 131–133 —Afr. trop.
turbinata, *H. Wendl. in Bot. Zeit.* xxxix. (1881) 92 — Afr. trop.
ventricosa, *Kirk, in Journ. Linn. Soc.* ix. (1867) 285. —Afr. trop.

HYPHIPUS, Rafin. Sylva Tellur. 125 (1838) = **Loranthus**, Linn.
bracteata, Rafin. l. c. = L. cupulifer.
trigona, Rafin. l. c. = L. americanus.

HYPHYDRA, Schreb. Gen. 666 (1791) = **Tonina**, Aubl. (Eriocaulonese).
amplexicaulis, Vahl, Symb. Bot. iii. 99 = T. fluviatilis.

HYPNOTICON, Reichb. Nom. 122 (1841) = seq.

HYPNOTICUM, Rod. ex Meisn. Gen. Comm. 184 (1840) = **Withania**, Pauq. (Solanac.).
frutescens, Rod. ex Boiss. Voy. Bot. Espagne, ii. 437 = W. frutescens.
somniferum, Rod. ex Boiss. l. c. 436 = W. somnifera.

HYPOBATHRUM, Blume, Bijdr. 1007 (1826). *RUBIACEAE*, Benth. & Hook. f. ii. 92.
albicaule, *Baill. Adansonia*, xii. (1878) 209.—Madag.
angolense, Baill. Hist. Pl. vii. 315 = Tricalysia angolensis.
Bolvinianum, *Baill. Adansonia*, xii. (1878) 208.—Madag.
canthioides, Baill. l. c. 211 = Fertia canthioides.
Commersonianum, *Baill. l. c.* 204.—Madag.
comorense, *Baill. l. c.* 210.—Ins. Comor.
elliptícum, Baill. l. c. 211 = Fertia spodanthera.
frutescens, *Blume, Bijdr.* 1007.—Java.
gravesolens, Baill. Adansonia, xii. (1878) 206 = Zygoon gravesolens.
Kirkii, Baill. l. c. 205 = Empogona Kirkii.
lagoense, *Baill. l. c.* 210.—Afr. trop.
lanceolatum, Baill. l. c. 206 = Krausia lanceolata.
leucocarpum, *Baill. l. c.* 205.—Madag.
myrtifolium, Baill. l. c. 204 = Nesciella myrtifolia
parviflorum, *Miq. Fl. Ind. Bat.* ii. 286.—Java.
racemosum, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 185 = Petunga Roxburghii.
strictum, Kurz, For. Fl. Brit. Burma, ii. 50 = Hyptianthera strícta.
viridiflorum, Baill. Adansonia, xii. (1878) 212 = Diplospora viridiflora.

HYPOBRICHIA, M. A. Curt. ex Torr. & Gray, Fl. N. Am. i. 479 (1840) = **Didiplis**, Rafin. (Lythrar.).
Nuttallii, Torr. & Gray, l. c. 480 = D. linearis.
Spruceana, Benth. ex Benth. & Hook. f. Gen. i. 777, in obs. = Rotala verticillaris.

HYPOCALYMMA, Endl. Gen. 1280 (1840).
MYRTACEAE, Benth. & Hook. f. i. 702.
angustifolium, *Schau. in Linnaea*, xvii. (1843) 241.—Austral.
asperum, Schau. l. c. = strictum.
boronizeum, *F. Muell. ex Benth. Fl. Austral.* iii. 95. —Austral.
ciliatum, Turcz. in Bull. Soc. Nat. Mosc. (1862) ii. 325 = xanthopetalum.
cordifolium, *Schau. in Linnaea*, xvii. (1843) 241; *et in Lehm. Pl. Preiss.* i. 112.—Austral.
cuneatum, Turcz. in Bull. Soc. Nat. Mosc. (1862) ii. 326 = xanthopetalum.
Cunninghamii, Schau. in Linnaea, xvii. (1843) 241 = strictum.
ericifolium, *Benth. Fl. Austral.* iii. 94.—Austral.
hypericifolium, *Benth. l. c.* 95.—Austral.
linifolium, *Turcz. in Bull. Soc. Nat. Mosc.* (1862) ii. 325.—Austral.
longifolium, *F. Muell. Fragm.* ii. 28.—Austral.
myrtifolium, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 332.—Austral.
strictum, *Schau. in Linnaea*, xvii. (1843) 241.—Austral.
suave, Lindl. Bot. Reg. (1844) Misc. 27 = angustifolium.
tetrapetalum, *Turcz. in Bull. Soc. Nat. Mosc.* (1862) ii. 326.—Austral.
xanthopetalum, *F. Muell. Fragm.* ii. 29.—Austral.

HYPOCALYMNA, Meisn. Gen. Comm. 354 (1843) = prec.

HYPOCALYPTUS, Thunb. Prod. Pl. Cap. ii. 124 (1800); Nov. Gen. xi. 151 (1800); partim. *LEGUMINOSAE*, Benth. & Hook. f. i. 484.
DUVALIA, Bonpl. Desc. Pl. Rar. Malm. 130. t. 53 (1813).
calyptratus, Thunb. Prod. Pl. Cap 192; Nov. Gen. 151 = Podalyria calyptrata.
cerasicus, *Thunb. ll. c.* 124; 153.—Afr. austr.
capensis, Thunb. ll. cc. 192; 154 = Virgilia capensis.
cordatus, Thunb. ll. cc. 192; 154 = Podalyria Burchellii.
glaucus, *Thunb. Nov. Gen.* 152.—Afr. austr.
obcordatus, *Thunb. Prod. Pl. Cap.* 124.—Afr. austr.
oxalidifolius, Baill. Hist. Pl. ii. 336 = Loddigesia oxalidifolia.
pedunculatus, Thunb. Prod. Pl. Cap. 192; Nov. Gen. 152 = Podalyria argentea.
sericeus, Thunb. ll. cc. 124; 152 = Podalyria sericea.

HYPOCARPUS, A. DC. in DC. Prod. viii. 245, 673 (1844) = **Xylosma**, Poepp. (Oiacin.).
Gardnerianus, A. DC. l. c. 246 = L. Gardnerianus.
pauciflorus, A. DC. l. c. = L. paucifloras.

HYPOCHAERIS, Linn. Hort. Cliff. 385 (1737) = seq.

HYPOCHOERIS, Linn. Gen. ed. I. 238 (1737).
COMPOSITAE, Benth. & Hook. f. ii. 519.
ACHYROPHORUS, Adans. Fam. ii. 113 (1763).
AGENORA, D. Don, in Edinb. N. Phil. Journ. (1828–29) 310.
AMELACHAENIUM, Turcz. ex DC. Prod. viii. 94 (1838).
ARACHNOPOGON, Berg, ex Haberle, Cat. Pesth.; ex Stend. Nom. ed. II. i. 118 (1840).
ARACHNOSPERMUM, Berg, ex Stend. l. c. (1840).
CYNOSERIS, Endl. in Bot. Zeit. i. (1843) 458.
FABERA, Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 192.
HETEROMORPHA, Viv. ex Cosa. in Bull. Soc. Bot. Fr. xii. (1865) 278.
HIERACHIUM, Hill, Hort. Kew. 43 (1769).
METABASIS, DC. Prod. vii. 97, 207 (1838).
ORROPHILA, D. Don, in Phil. Mag. xi. (1832) 338.
PITTOPOGON, Cass. in Bull. Soc. Nat. lxviii. 507 (1827).
POLALLITES, Cass. l. c. xliii. 42 (1826), partim.
ROBERTIA, DC. Fl. Fr. Suppl. 458 (1815).

HYPOCHOERIS:—
SERIOLA, Linn. Sp. Pl. ed. II. 1139 (1763).
TROMMSDORFIA, Bernh. Syst. Vers. Erf. 103 (1800).
Achyrophorus, *Linn. Sp. Pl.* 810.—Afr. bor.; Sicilia.
adscendens, Brot. Phytog. Lusit. i. 55. t. 25 = glabra.
aethnensis, Ball, in Journ. Linn. Soc. xvi. (1878) 542 = serioloides.
actinusis, *Benth. & Hook. f. Gen.* ii. 520; *Cos. Passer. & Gib. Comp. Fl. Ital.* 465.—Europ.
albicans, Urcla, in Roem. Arch. i. i. 70 = H. Achyrophorus.
alpicola, Schur, Enum. Pl. Transs. 227, Index = maculata.
alpina, Nym. Consp. 470, in syn. = maculata.
andina, *Griseb. in Goett. Abh.* xix. (1874) 198.—Reg. Andin.
aparsioides, *Hook. & Arn. Bot. Beech. Voy.* 28.—Chili.
Arachnites, Link, Handb. i. 792 = seq.
arachnoideus, Dc. Prod. vii. 90 = seq.
arachnoides, Poir. Encyc. v. 572 = glabra.
arenaria, *Gaudsch. in Ann. Sc. Nat. Sér.* I. v. (1825) 102.—Chili.
ascendens, Walp. Rep. vi. 340 = glabra.
aurantiaca, Turcz. ex DC. Prod. vii. 94 = grandiflora.
Balbisii, Mauri, Cent. Pl. Rom. xiii. 38; Loisel. Not. 124 = glabra.
Balbisii, Reichb. Fl. Germ. Excurs. 268 = radicata.
Barrhadii, Willd. ex Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 142 = glabra.
Bertersii, Colla, in Mem. Acc. Tor. xxxvili. (1835) 40. t. 33 = sparsioides.
brasiliensis, *Griseb. in Gontt Abh.* xxiv. (1879) 217.—Bras.
Candollei, Regel, in Linnaea, xvi. (1842) 64 = glabra.
canescens, Stev. in Mém. Soc. Nat. Mosc. iv. (1812) 60 = Picris laevis.
capensis, *Lin. Syn. Comp.* 130.—Afr. austr.
chrysantha, *Poepp. ex DC. Prod.* vii. 94.—Chili.
contexta, Wallr. in Linnaea, xiv. (1840) 663; et Beitr. Fl. Hercyn. 293 = glabra.
crenica, Tausch, in Flora, xii. (1829) 1. Erg. 36 = pinnatifida.
crotensis, *Benth. & Hook. f. Gen.* ii. 520; *Boiss. Fl. Orient.* iii. 784.—Creta; Sicilia.
dawurica, Pall. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) i. 99 = grandiflora.
dimorpha, Brot. Fl. Lusit. i. 332 = glabra.
dimorpha, Tenore, Fl. Nap. v. 201 = radicata.
dubia, *Mart. Cat. Hort. Erlang.* (1812).—Hab.?
Echegaryi, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 51.—Reg. Argent.
elata, *Griseb. in Goett. Abh.* xix. (1874) 198.—Reg. Andin.
Facchiniana, Ambr. Fl. Tyr. Austr. ii. 558 = uniflora.
foliosa, Vill. Fl. Delph. 58 = uniflora.
Gardneri, *Baker, in Mart. Fl. Bras.* vi. iii. 331.—Bras.
glabra, *Linn. Sp. Pl.* 810.—Europ.; Reg. Mediterr.
glabra, E. Mey. ex DC. Prod. vii. 91 = capensis.
glauca, Presl, Delic. Prag. 114 = H. Achyrophorus.
grandiflora, Gilib. Fl. Lituan. i. 232 = maculata.
grandiflora, *Lodd. Fl. Alt.* iv. 164.—Sibir. altaic.
helvetica, Wulf. in Jacq. Misc. ii. 25 = uniflora.
heterosperma, Schult. ex Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 142 = glabra.
hirsuta, DC. Prod. vii. 90 = glabra.
hirta, *Urcia, in Roem. Arch.* i. i. 70.—Sicilia.
hispida, Roth, in Usteri, Ann. Bot. xiv. (1795) 30 = glabra.
hispida, Willd. Hort. Berol. i. 46 = cretensis.
infesta, Salisb. Prod. 182 = radicata.
intermedia, Rchb. ex Reichb. Fl. Germ. Excurs. 266 = radicata.
intertexta, Petterm. in Flora, xxvii. (1844) 481 = glabra.
laevigata, Benth. & Hook. f. Gen. ii. 520; Cos. Passer. & Gib. Comp. Fl. Ital. 465 = H. Achyrophorus.
laziophylla, C. Koch, ex Nym. Consp. 470 = radicata.
leontodontoides, *Ball, in Journ. Bot.* xi. (1873) 271.—Marocc.
linearifolia, *Moris, in Mem. Acc. Tor.* Ser. II. ii. (1840) 308.—Europ.
macrorhiza, Gmelin. ex Spreng. Syst. iii. 669 = pinnatifida.
maculata, *Linn. Sp. Pl.* 810.—Europ.; As bor.
Meyeniana, *Griseb. in Goett. Abh.* xix. (1874) 198.—Peruv.; Bolivia.
minima, Cyrilli, Fl. Rar. Neap. Fasc. i. 29. t. 10; Desf. Fl. Atlant. ii. 238 = glabra.
montana, Hill, Veg. Syst. iii. 153 = glabra?
neapolitana, DC. Prod. vii. 91 = radicata.
odorata, *Benth. & Hook f. Gen.* ii. 519.—Chili.
petiolaris, *Griseb. in Goett. Abh.* xxiv. (1879) 218.—Reg. Argent.
picroides, *Griseb. l. c.* 217.—Reg. Argent.
pinnatifida, Aucher, ex DC. Prod. vii. 97 = cretensis.
pinnatifida, *Cyrilli, ex Tenore, Syll.* 406.—Italia.
platylepis, Boiss. Voy. Bot. Espagne, ii. 876 = radicata.
pontana, Linn. Sp. Pl. 810 = Crepis montana.
pusilla, Poir. Encyc. Suppl. iv. 530 = glabra.
radicata, Falk, Beitr. ii. 234, sphalm. = seq.
radicata, *Linn. Sp. Pl.* 810.—Europ.; Afr. bor.
runcinata, Gilib. Fl. Lituan. i. 226 = radicata.
rupestris, Poepp. ex Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 124 = spargioiorites.
salina, Gren. Fl. Chaine Jurass 456, in nota = radicata.
Salsmanniana, DC. Prod. vii. 91 = glabra.
Scorsonerae, *F. Muell. Sel. Pl. Indust. Cult.* 107.—Chili.
scrioloides, *Bertol. Misc. Bot* ii. 236.—Reg. Mediterr.
sessiliflora, *H. B. & K. Nov. Gen. et Sp.* iv. 2.—Ecuador.
sicula, *Ucria, in Roem. Arch.* i. i. 70.—Sicilia.
simplex, Mérat, Fl. Par. ed. I. 310 = glabra.
sonchoides, Bert. ex Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 117 = Scorzonerae.
sonchoides, *H. B. & K. Nov. Gen. et Sp.* iv. 2. t. 301. —Ecuador.
stellata, Gater. Fl. Montauh. 140 = glabra.
stigosa, *Soland. ex Lowe, Man. Fl. Mad.* i. 537.—Ins. Mader.
subalpina, Schur, Enum. Pl. Transs. 365 = radicata.
taraxacoides, *Loisel. in Mém. Soc. Linn. Par.* vi. (1827) 425.—Corsica.
taraxacifolia, Moench, Meth. Suppl. 224 = Crepis albida.
taraxacoides, *Benth. & Hook. f. Gen.* ii. 519.—Peruv.
taraxacoides, Pourr. ex Steud. Nom. ed. I. 422, 618 = Crepis albida.
Tenorii, Guss. Fl. Sicul. Syn. ii. 428 = radicata.
tenuiflora, *Boiss. Fl. Orient.* iii. 785.—Ins. Creta.
tenuifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 218.—Chili.
tenuifolia, Trev. ex Sch. Bip. in Nov. Act. Nat. Cur. xxi. i. (1845) 102 = taraxacifolia.
thracica, Stev. ex DC. Prod. vii. 90 = glabra.
tyrolensis, Schlw. ex DC. l. c. 93 = maculata.
uniflora, Hoffm. Deutschl. Fl. ed. II. ii. 128 = maculata.
uniflora, *Vill. Prosp.* 37.—Europ.
urens, Linn. Sp. Pl. 810 = serioloides
variegata, *Baker, in Mart. Fl. Bras.* vi. iii. 333.—Reg. Argent.

HYPOCISTIS, [Tourn.] Adans. Fam. ii. 76 (1763) = **Cytinus**, Lin.
lutea, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 148 = C. Hypocistis.
rubra, Fourr. l. c. = C. Hypocistis.

HYPOCYRTA, Mart. Nov. Gen. et Sp. iii. 43 (1829). *GESNERACEAE*, Benth. & Hook. f. ii. 1010.
aggregata, Mart. Nov. Gen. et Sp. iii. 51. t. 201 = Codonanthe aggregata.
barbata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 8.—Peruv.
brevicalyx, Hort. Veitch.—Cf. Gard. Chron. (1867) 458 = Isolonus hypocyrtiiformis.
calycodonta, Lem. in Fl. des Serres, Sér. I. ii (Août, 1846) = Alloplectus dichrous.
carnosa, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 178 = Codonanthe carnosa.
ciliosa, Mart. Nov. Gen. et Sp. iii. 52 = Episcia ciliosa.
crassifolia, *Focke, in Tijdschr. Nat. Wetens.* v. (1852) 199.—Guian.
discolor, Lindl. Bot. Reg. (1845) Misc. 19 = Alloplectus dichrous.
epidendra, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 8.—Peruv.
ferruginea, *Poepp. & Endl. l. c.*—Peruv.
gibbosa, *Poepp. & Endl. l. c.* t. 202.—Bras.

HYPOCYRTA :—
glabra, *Hook. Bot. Mag.* t. 4346.—Am. austr.
gracilis, *Hook. l. c.* t. 4581.—Bras.
gracilis, Mart. Nov. Gen. et Sp. iii. 50. t. 220 = Codonanthe gracilis.
hirsuta, *Mart. l. c.* 52. t. 222.—Bras.
leucostoma, Hook. Bot. Mag. t. 4310 = Besleria leucostoma.
macrocalyx, *Hanst. in Mart. Fl. Bras.* viii. 410.—Bras.
macrophylla, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 3.—Peruv.
Nummularia, *Hanst. in Linnaea*, xxxiv. (1865–66) 381.—Costa Rica.
perianthomega, *Tenore, in Parl. Giorn. Bot. Ital.* ii. (1847) fasc. VII–VIII. 200, 201.—Bras.
radicans, *Klotzsch & Hanst. in Mart. Fl. Bras.* viii. 409.—Bras.
scabrida. *Lem. in Rev. Hortic.* Sér. III. i. (1847) 364.—Bras.
Selloana, *Klotzsch & Hanst. in Mart. Fl. Bras.* viii. 409.—Bras.
sericea, *Hanst. l. c.* 413.—Bras.
strigillosa, *Mart. Nov. Gen et Sp.* iii. 52.—Bras.
villosa, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 280.—Bras.
villosa, Klotzsch & Hanst. in Mart. Fl. Bras. viii. 407 = Alloplectus villosus.

HYPODAEURUS, Hochst. in Flora, xxvi. (1844) 101 = **Anthephora**, Schreb. (Gramin.).
cenchroides, Hochst. l. c. = A. Hochstetteri.

HYPODEMA, Reichb. Nom. 56 (1841) = **Cypripedium**, Linn. (Orchid.).

HYPODEMATIUM, A. Rich. Tent. Fl. Abyss i. 348 (1847) = **Spermacoce**, Linn. (Rubiac.).
amplisatum, A. Rich. l. c. 349 = S. amplisata.
sphaerostigma, A. Rich. l. c. = S. amplisata.

HYPODEMATIUM, A. Rich. Tent. Fl. Abyss ii. 286. t. 83 (1850) = **Lisschcilus**, R. Br. (Orchid.).
abyssinicum, A. Rich. l. c.—Abyssin.

HYPODISCUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 450 (1836). *RESTIACEAE*, Benth. & Hook. f. iii. 1035.
BOECHIA, Kunth, Enum. Pl. iii. 448 (1841).
LEPIDANTHUS, Nees, in Linnaea, v. (1830) 665.
LEUCOPLEURUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 450 (1836).
albo-aristatus, Mast. in Journ. Linn. Soc. viii. (1865) 255 = duplicatus.
argenteus, *Mast. l. c.* x. (1869) 261.—Afr. austr.
aristatus, *Nees, in Lindl. Introd. Nat. Syst.* ed. II. 450.—Afr. austr.
binatus, *Mast. in Journ Linn. Soc.* x. (1869) 258.—Afr. austr.
duplicatus, *Hochst. in Flora*, xxviii. (1845) 388.—Afr. austr.
Necsii, *Mast. in Journ. Linn. Soc.* x. (1869) 260.—Afr. austr
nitidus, *Nees, ex Mast. l. c.* 259.—Afr. austr.
Oliverianus, *Mast. l. c.* 254.—Afr. austr.
rugosus, *Mast. l. c.* 255.—Afr. austr.
striatus, *Mast. l. c.* 258.—Afr. austr.
synchrolepis, *Mast. l. c.* viii. (1865) 255.—Afr. austr.
Willdenovia, *Mast. l. c.* x. (1869) 259.—Afr. austr.

HYPOELYTRUM, H. B. & K. Nov. Gen. et Sp. i. 218 (1815), sphalm. = **Hypolytrum**, Rich. (Cyperac.).

HYPOESTES, Soland. ex R. Br. Prod. 474 (1810). *ACANTHACEAE*, Benth. & Hook. f. ii. 1122.
acuminata, Hochst. ex T. Anders. in Journ. Linn. Soc. vii. (1864) 50 = trifora.
adoënsis, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 24, nomen; *et in A. Rich. Tent. Fl. Abyss.* ii. 162.—Abyssin.
adscendens, *Nees, in DC. Prod.* xi. 509.—Madag.
Alsine, Nees, l. c. = serpens.
anisophylla, *Nees, l. c.* 508.—Madag.
annua, *Hochst. & Steud. ex Steud. Nom.* ed. II. i. 791.—Arab.
antennifera, *S. Moore, in Journ. Bot.* xviii. (1880) 41.—Afr. trop.
aristata, *R. Br. Prod.* 474, in obs.—Afr. austr.
Bakeri, *Vatke, in Bremen, Abh.* ix. (1885) 133.—Madag.
Barteri, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 49.—Afr. trop.
Bojeriana, *Nees, in DC. Prod.* xi. 506.—Madag.
brachiata, *Baker, in Journ. Linn. Soc.* xx. (1883) 224.—Madag.
calaminthoides, *Baker, l. c.* 222.—Madag.
cancellata, *Nees, in DC. Prod.* xi. 505.—Afr. trop.
Cephalotes, *Link, Jahrb.* i. 111. 46.—Ind. or.
cernua, *Nees, in DC. Prod.* xi. 509.—Madag.
clivicola, E. Mey. ex Nees, l. c. 508 = verticillaris.
comorensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 223.—Madag.
contorta, Nees, in DC. Prod. xi. 156 = Peristrophe contorta.
corymbosa, *Baker, in Journ. Linn. Soc.* xx. (1883) 228.—Madag.
Cuminglanus, *Benth. & Hook f. Gen.* ii. 1122.—Ins. Philipp.
Decaisneana, Nees, in DC. Prod. xi. 508 = malacocnsis.
dicliptcroides, *Nees, l. c.* 510.—Madag.
elegans, *Nees, l. c.* 502.—Madag.
fascicularis, *Nees, l. c.* 504.—Madag.
fastuosa, *R. Br. Prod.* 474, in obs.—Ind or.?
fimbriata, E. Mey. in Drège, Zwei Pfl. Docum. 193 = Ruttya ovata.
flexibilis, *Nees, in DC Prod.* xi. 505.—Madag.
floribunda, *R. Br. Prod.* 474.—Austral.
Forskalei, *R. Br. l. c.*, in obs.—Arab.
glabrata, *Presl, Rel. Bemork.* 96.—Afr. austr.
glandulosa, *Hochst in Flora*, xxiv. (1841) i. *Intell* 24, nomen; *et ex Nees, in DC. Prod.* xi. 573.—Abyssin.
gracilis, *Nees, l. c.* 506.—Madag.
hirsuta, *Nees, l. c.* 504.—Madag.
inconspicua, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 18.—Ins. Mascar.
insularis, *T. Anders. l. c.* vii. (1862) 49.—Afr. trop.
involucrata, *Roem. & Schult. Syst.* i. 141.—Ind. or.
jasminoides, *Baker, in Journ. Linn. Soc.* xxi. (1885) 432.—Madag.
lanata, *Dalz. in Hook. Kew Journ.* ii. (1850) 343.—Ind. or.
lasioclada, *Nees, in DC. Prod.* xi. 503.—Madag.
lasiostegia, *Nees, l. c.* 504.—Madag.
latifolia, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 43; *et ex Nees, l. c.* 509.—Afr. trop.
laxiflora, Nees, l. c. 508 = malacocnsis.
longifolia, *Hort. Bogor. ex T. Anders. in Journ. Agr. Hort. Soc. Ind.* N. S. i. (1868) 283.—Hab.?
lonicerioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 225.—Madag.
maculosa, *Nees, in DC. Prod.* xi. 503.—Madag.
malacocnsis, *Wight, Ic.* iv. t. 1555.—Malaya.
menthaefolia, *E. Mey. in Drège, Zwei Pfl. Docum.* 193.—Afr. austr.
microphylla, *Nees, in DC. Prod.* xi. 504.—Abyssin.
mollis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 49.—Afr. trop.
mollissima, *Nees, in DC. Prod.* xi. 505.—Madag.
oxystegia, *Nees, l. c.*—Madag.
plumosa, E. Mey. in Drège, Zwei Pfl. Docum. 193 = aristata.
polymorpha, E. Mey. l. c. = verticillaris.
polyphylla, *Miq. Fl. Ind. Bat.* ii. 852.—Java.
populifolia, *Miq. l. c.*—Java.
pubescens, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 89.—Socotr.
pulchra, *Nees, in DC. Prod.* xi. 505.—Madag.
purpurea, *R. Br. Prod.* i. 474, in obs.—Malaya; China.
Richardi, *Nees, in DC. Prod.* xi. 502.—Madag.
rodriguesiana, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 18.—Ins. Mascar.
roseo, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 381.—Ins. Timor.
roseo, *Nees, in DC. Prod.* xi. 506.—Afr. trop.
sanguinolenta, *Hook. Bot. Mag.* t. 5511.—Madag.
saxicola, *Nees, in DC. Prod.* xi. 503.—Madag.
serpens, *R. Br. Prod.* 474, in obs.—Ins. Maurit.
simensis, *Hochst. ex Schweinf. Beitr. Fl. Aethiop.* 111.—Afr. trop.
sinica, *Miq. in Journ. Bot. Néerl.* i. (1861) 117.—China.
spicata, *Nees, in DC. Prod.* xi. 504.—Madag.
stachyoides, *Baker, in Journ. Linn. Soc.* xxi. (1885) 431.—Madag.
strobilifera, *S. Moore, in Journ. Bot.*, xviii. (1880) 40.—Afr trop.
teucrioides, *Nees, in DC. Prod.* xi. 508.—Madag.
Teysmanniana, *Miq. Fl. Ind. Bat.* ii. 853.—Java.
Thomsoniana, *Nees, in DC. Prod.* xi. 502.—Madag.
tomentosa, *Miq. Fl. Ind. Bat.* ii. 852.—Java.
trichochlamys, *Baker, in Journ. Linn. Soc.* xxi. (1885) 432.—Madag.
trifiora, *Roem. & Schult. Syst.* i. 141.—Reg. Himal.; Abyssin.
umbellata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 57 = Dicliptera micrantha.
uniflora, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 24, nomen.—Abyssin.
unilateralis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 431.—Madag.
verticillaris, *R. Br. Prod.* 474, in obs.—Afr. austr.
Wallichii, Nees, in Wall. Pl. As. Rar. iii. 114 = trifiora.
Zollingeriana, *Miq. Fl. Ind. Bat.* ii. 852.—Malaya

HYPOGLOTTIS, Four. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364 = **Astragalus**, Tourn. (Legumin.).
Glaux, Four. l. c. = A. Glaux.
purpurea, Four. l. c. = A. Hypoglottis.

HYPOGOMPHIA, Bunge, in Bull. Acad. Pétersb. xviii (1873) 80. *LABIATAE*, Benth. & Hook. f. ii. 1201.
nana, *Benth. & Hook. f. l. c.*—Afghan.
turkestana, *Bunge, in Bull. Acad. Pétersb.* xviii. (1873) 80.—Turkestan.

HYPOGON, Kaſin. Fl. Ludov. 148 (1817) = **Collinsonia**, Linn. (Labiat.).
anisatum, Kaſin. l. c. = C. anisata.

HYPOGYNIUM, Nees, Agrost. Bras. 364 (1829) = **Andropogon**, Linn. (Gramin.).
campestre, Nees, l. c. 305 = A. hypogynus.
spathiflorum, Nees, l. c. 349 = A. spathiflorus.

HYPOLAENA, R. Br. Prod. 251 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1035.
CALORPHUS, Labill. Nov. Holl. Pl. ii. 78. t. 228 (1806).
CALOSTROPHUS, F. Muell. Fragm. viii. (1874) 86, 88.
anceps, *Mast. in Journ. Linn. Soc.* x. (1869) 267.—Afr. austr.
aspera, Mast. in DC. Monog. Phan. i. 379 = Restio Loxocarya.
aspera, *Mast. in Journ. Linn. Soc.* x. (1869) 264.—Afr. austr.
Benthamii, *Mast. in DC. Monog. Phan.* i. 367.—Austral.
Burchellii, *Mast. in Journ. Linn. Soc.* x. (1869) 268.—Afr austr.
crispata, Mast. in DC. Monog. Phan. i. 379 = Restio crispatus.
densa, *Mast. l. c.* 366.—Austral.
Ecklouiana, *Nees, ex Mast. in Journ. Linn. Soc.* x. (1869) 268. t. 7.—Afr. austr.
elongata, *Mast. in DC. Monog. Phan.* i. 379.—Austral.
Esenbeckii, F. Muell. Fragm. viii. 86 = exsulca.
exsulca, *R. Br. Prod.* 251.—Austral.
fasciculata, *Mast. in DC. Monog. Phan.* i. 368.—Austral.
fastigiata, *R. Br. Prod.* 251.—Austral.
fastigiata, Nees, in Lehm. Pl. Preiss. ii. 69 = exsulca.
filiformis, *Mast. in Journ. Linn. Soc.* x. (1869) 267.—Afr. austr.
flexuosa, *Mast. in DC. Monog. Phan.* i. 365.—Austral.
gracilis, *Nees, ex Mast. in Journ. Linn. Soc.* x. (1869) 266.—Afr. austr.
gracillima, *Benth. Fl. Austral.* vii. 239.—Austral.
grandiuscula, F. Muell. Fragm. viii. 85 = exsulca.
impolita, *Mast. in Journ. Linn. Soc.* x. (1869) 264.—Afr. austr.
lateriflora, *Benth. Fl. Austral.* vii. 238.—Austral.
laxiflora, *Nees, in Linnaea*, v. (1830) 663.—Afr. austr.
longissima, Benth. Fl. Austral. vii. 238 = elongata.
pubescens, *Mast. in DC. Monog. Phan.* i. 365.—Austral.
pubescens, Nees, in Lehm. Pl. Preiss. ii. 69 = Restio Loxocarya.
tenuis, *Mast. in Journ. Linn. Soc.* x. (1869) 265.—Afr. austr.
ventita, *Mast. in DC. Monog. Phan.* i. 367.—Austral.
virgata, *Mast. in Journ. Linn. Soc.* x. (1869) 268.—Afr. austr.

HYPOLEPIS, Beauv. ex Lestib. Essai Cyp. 38 (1819) = **Fiotnia**, Schrad. (Cyperac.).
atrata, Nees, 16 Linnæa, viii. (1833) 98 = F. setiformis.
Capitellum, Nees, l. c. vii. (1832) 522 = F. setiformis.
composita, Nees, l. c. 525 = F. composita.
nigrescens, Nees, l. c. 522 = F. bracteata.
scariosa, Nees, l. c. 521 = F. bracteata.

HYPOLEPIS, Pers. Syn. ii. 598 (1807) = **Cytinus**, Linn.
sanguinea, Pers. l. c.—Afr. austr.

HYPOLOBUS, Fourn. in Mart. Fl. Bras. vi. iv. 311 (1885). *ASCLEPIADEÆ.*
Infractus, *Fourn. l. c.*—Bras.

HYPOLYTHRUM, Walp. Ann. i. 903 (1849) = seq.

HYPOLYTRUM, Rich. in Pers. Syn. i. 70 (1805). *CYPERACEÆ*, Benth. & Hook. f. iii. 1054.
ALBRIA, J. & C. Presl, Rel. Haenk. i. 184. tt. 34, 35 (1828).
BEKRA, Beauv. in Lestib. Essai Fam. Cyp. 48 (1819).
TRINCA, Roxb. Hort. Beng. (31) (1814), nomen.
TUNGA, Roxb. Fl. Ind. ed. Carey, i. 187 (1820), partim.
africanum, *Nees, in Linnæa*, ix. (1834) 288, nomen; *Steud. Syn. Pl. Cyp.* 182.—Afr. trop.
albidum, Willd. ex Kunth, Enum. Pl. ii. 266 = Lipocarpha argentea.
amplum, *Kunth, l. c.* 272.—Bras.
anomalum, *Steud. Syn. Pl. Cyp.* 183.—Ins. Borbon.
argenteum, H. B. & K. Nov. Gen. et Sp. i. 218 = Lipocarpha Humboldtiana.
argentinum, Vahl, Enum. Pl. ii. 283 = Lipocarpha argentea.
Aschersonianum, Boeck. in Flora, lxv. (1882) 26 (Mapaniae sp.).—Afr. bor.
Berterianum, Schult. Mant. ii. 183 = seq.
Berterii, Spreng. Neue Entdeck. i. 241 (= *Dichromena pusilla*).—Ind. occ.
borneense, Kurz, in Journ. Soc. Beng. xxxviii. (1869) 11. 74 = Mapania bancana.
caespitosum, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 68 = longifolium.
capillare, Schrad. ex Nees, l. c. 61 = Scirpus micranthus.
cariciforme, *Nees, in Linnæa*, ix. (1834) 288, nomen; *Steud. Syn. Pl. Cyp.* 182.—Guiana.
ceylanicum, Heyne, ex Nees, l. c. = Lipocarpha sphacelata.
compactum, Zoll. & Mor. Syst. Verz. Zoll. 98 (Mapaniae sp.).—Java.
compactum, *Nees & Mey. in Linnæa*, ix. (1834) 288. —Ins. Philipp.
corymbosum, Rich. ex Boeck. l. c. xxxvii. (1871-73) 125 = sylvaticum.
costatum, Hochst. ex Steud. Syn. Pl. Cyp. 183 = amplum.
costatum, Thw. Enum. Pl. Zeyl. 346 = Scirpodendron costatum.
dianthrum, A. Dietr. Sp. Pl. ii. 365 = latifolium.
dianthorum, Seem. Fl. Vit. 317, in nota = Schrad erianum.
disstiflorum, *Steud. Syn. Pl. Cyp.* 182.—Ins. Carolin.
ensifolium, Willd. ex Kunth, Enum. Pl. ii. 271, 272 = latifolium; mauritianum.
filiforme, Heyne, in Wall. Cat. n. 8444 C = Lipocarpha sphacelata.
filiforme, Vahl, Enum. ii. 284 = Lipocarpha filiformis.
floribundum, Nees, in Linnæa, ix. (1834) 288 (Mapaniae sp.).—Ins. Marian.
fuscum, *Nees, in Mart. Fl. Bras.* ii. i. 67.—Bras.
giganteum, Nees, in Wight, Contrib. 93, partim = Wightianum.
gigantinum, Wall. Cat. n. 8404 = latifolium.
Glaziovii, *Boeck. in Flora*, lxiii. (1880) 488.—Bras.
glomeratum, Rich. ex Kunth, Enum. Pl. ii. 270 = pungens.
glomeratum, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 64 = Lipocarpha Sellowiana.
gracile, Rich. ex Pers. Syn. i. 70 = Lipocarpha sphacelata.
humile, Boeck. in Linnæa, xxxvii. (1871-73) 128 (Mapaniae sp.).—Java.
iridifolium, Link, Jahrb. i. iii. 91 = Diplasia karatæfolia.
iridifolium, Willd. ex Kunth, Enum. Pl. ii. 272 = mauritianum.
irriguum, *Nees, in Mart. Fl. Bras.* ii. i. 66.—Bras.
laevigatum, Spreng. Syst. i. 233 = Lipocarpha argentea.
latifolium, Baker, Fl. Maurit. 423 = mauritianum.
latifolium, Boeck. in Linnæa, xxxvii. (1871) 129 = nemorum.
latifolium, *Rich. ex Pers. Syn.* i. 70.—As. et Austral. trop.
latifolium, *Seem. Fl. Vit.* 317.—Ins. Fiji.
laxum, Kunth, Enum. Pl. ii. 270 = longifolium.
Leprieurii, Nees, ex Steud. Syn. Pl. Cyp. 183 = Mapania africana.
longifolium, *Nees, in Linnæa*, ix. (1834) 288.—Guiana.
longifolium, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 59 = scapigerum.
longivaginatum, Thw. Enum. Pl. Zeyl. 346 (Mapaniae sp.).—Zeylan.
maculatum, Steud. Nom. ed. II. i. 791 = Lipocarpha Humboldtiana.
macranthum, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 507.—Afr. trop.
macrocephalum, Gaudich. in Freyc. Voy. Bot. 415 (Mapaniae sp.).—Ins. Molucc.!
macrophyllum, *Boeck. in Flora*, lxi. (1878) 142.—Ins. S. Doming.
macrophyllum, Miq. Fl. Ind. Bat. iii. 334, in obs. (Mapaniae sp.).—Ins. Molucc.
mauritianum, *Nees, in Linnæa*, ix. (1834) 288.—Ins. Maurit.
micranthum, Boeck. in Kjoeb. Vidensk. Meddel. (1871) 154 (Mapaniae sp.).—Bras.
microcephalum, R. Br. Prod. 220 = Lipocarpha microcephala.
multinervi, Hochst. ex Steud. Syn. Pl. Cyp. 183 = Calyptrocarya longifolia.
myrianthum, Miq. Fl. Ind. Bat. iii. 333 = latifolium.
nemorum, *Beauv. Fl. Owar.* ii. 12, 13. t. 67.—Afr. trop.
nemorum, Boj. Hort. Maurit. 385 = mauritianum.
nicaraguense, *Liebm. in Vidensk. Selsk. Skr.* v. ii. (1851) 285.—Nicarag.
pallidum, *Willd. ex Steud. Nom.* ed. II. i. 792.—Am. austr.
pandanophyllum, *F. Muell. Fragm.* ix. 16.—Austral.
proliferum, *Boeck. in Linnæa*, xxxvii. (1871-73) 126. —Hab.!
punctatum, Nees, in Mart. Fl. Bras. ii. i. 66 = sylvaticum.
pungens, *Vahl, Enum.* ii. 283.—Am. austr.
pycnocephalum, Benth. Bot. Voy. Sulph. 177 = Mapania pycnocephala.
pycnostachyum, Benth. in Journ. Linn. Soc. xv. (1877) 519, lapsu = praec.
rigens, Nees, in Mart. Fl. Bras. ii. i. 67 = pungens.
scabriramum, Boeck. in Flora, lxv. (1882) 26 (Mapaniae sp.).—Afr. bor.
scapigerum, *Nees, in Mart. Fl. Bras.* ii. i. 69.—Bras.
schoenoides, Nees, in Linnæa, ix. (1834) 288 = latifolium.
Schraderianum, *Nees, in Mart. Fl. Bras.* ii. i. 65.—Bras.
Sellowianum, Boeck. in Linnæa, xxxvii. (1871-73) 134 = amplum.
Sellowianum, Steud. Nom. ed. II. i. 792 = Lipocarpha Sellowiana.
senegalense, Rich. ex Pers. Syn. i. 70 = Lipocarpha argentea.
silvaticum, *Poepp. ex Kunth, Enum. Pl.* ii. 270.—Bras. *Soyauxii*, Boeck. in Flora, lxv. (1882) 25 (Mapaniae sp.).—Afr. trop.
sphacelatum, Vahl, Enum. ii. 283 = Lipocarpha sphacelata.
sphaerostachyum, *Boeck. in Kjoeb. Vidensk. Meddel.* (1871) 156.—Bras.
strictum, Poepp. ex Kunth, Enum. Pl. ii. 270 (Mapaniae sp.).—Bras.
sylvaticum, *Poepp. ex Kunth, l. c.*—Bras.
triceps, A. Dietr. Sp. Pl. ii. 368 = Lipocarpha sphacelata.
trinervium, Kunth, Enum. Pl. ii. 272 = latifolium.
umbellatum, Willd. ex Link, Jahrb. i. iii. 91 = pungens.
Wightianum, *Boeck. in Linnæa*, xxxvii. (1871-73) 130.—Ind. or.

HYPOPHAË, Medic. Pfl. Anat. 95 (1799) = **Hippophaë**, Linn. (Elæag.).

HYPOPHIALIUM, Nees, in Linnæa, vii. (1832) 580, in syn. = **Fictnia**, Schrad. (Cyperac.).
capillifolium, Nees, l. c. = F. ramosissima.

HYPOPHYLLANTHUS, Regel, Ind. Sem. Hort. Petrop. (1855) 51. *RUTACEÆ.*
Lindeni, *Regel, l. c.*—N. Granat.

HYPOPITHIS, Rafin. in Med. Rep. N. York, Ser. III. i. (1810) 297 = seq.

HYPOPITHYS, Scop. Fl. Carn. ed. II. i. 285 (1772) = seq.

HYPOPITYS, Dill. ex Adans. Fam. ii. 448 (1763). *MONOTROPEÆ*, Benth. & Hook. f. ii. 607.
MONOTROPA, Linn. Gen. ed. I. 111 (1737), partim.
dentata, Rafin. in Med. Rep. N. York, Ser. III. i. (1810) 297 = multiflora.
europæa, Nutt. Gen. Am. i. 271 = multiflora.
glabra, DC. Prod. vii. 780 = multiflora.
hypophegea, G. Don, Gen. Syst. iii. 866 = multiflora.
lanuginosa, Rafin. in Med. Rep. N. York, Ser. III. i. (1810) 297 : Nutt. Gen. Am. i. 271 = multiflora.
lutea, S. F. Gray, Nat. Arr. Brit. Pl. ii. 404 = multiflora.
Monotropa, Crantz, Inst. ii. 467 = multiflora.
multiflora, *Scop. Fl. Carn.* ed. II. i. 285.—Reg. bor temp.
racemosa, Rafin. in Med. Rep. N. York, Ser. III. i. (1810) 297 = multiflora.
secunda, Rafin. l. c. = multiflora.
striata, Rafin. l. c. = multiflora.
uniflora, Crantz, Inst. ii. 467 = Monotropa uniflora.

HYPOPOGON, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 246 = **Symplocos**, Linn. (Styrac.).
brevipes, Turcz. l. c. = S. coccinea.

HYPOPORUM, Nees, in Linnæa, ix. (1834) 303 = **Soleria**, Berg. (Cyperac.).
annulare, Nees, l. c. = S. annularis.
Baldwinii, Torr. in Ann. Lyc. N. York, iii. (1828-36) 382 = S. Baldwinii.
capillare, Nees, in Linnæa, ix. (1834) 303 = S. lithosperma.
capitatum, Nees, l. c. = S. Neesii.
compositum, Nees, in Mart. Fl. Bras. ii. i. 171 = S. composita.
diffusum, Nees, in Linnæa, ix. (1834) 303 = S. verticillata.
gracile, Torr. in Ann. Lyc. N. York, iii. (1828-36) 381 = S. gracilis.
hirtellum, Nees, in Linnæa, ix. (1834) 303 = S. hirtella.
humile, Nees, l. c. = S. hirtella.
interruptum, Nees, l. c. = S. interrupta.
interruptum, Torr. in Ann. Lyc. N. York, iii. (1828-36) 382 = S. hirtella.
leptostachyum, Nees, in Mart. Fl. Bras. ii. i. 172 = S. leptostachya.
lithospermum, Nees, in Wight, Contrib. 117 = S. lithosperma.
micrococcum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 255 = S. micrococca.
nutans, Nees, in Mart. Fl. Bras. ii. i. 170 = S. hirtella.
pergracile, Nees, in Edinb. N. Phil. Journ. (1834) 367 = S. pergracilis.
pleostachyum, Nees, in Mart. Fl. Bras. ii. i. 170 = S. pleostachya.
purpurascens, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 256 = S. Liebmanni.
purpurascens, Nees, in Linnæa, ix. (1834) 303 = S. lithosperma.
pygmæum, Nees, l. c. = S. pygmæa.
Sieberi, Nees, l. c. = S. lithosperma.
tenuilum, Nees, in Mart. Fl. Bras. ii. i. 171 = S. tenella.
variegatum, Nees, l. c. 173 = S. variegata.
verticillatum, Nees, in Bonplandia, iii. (1855) 87 = S. Liebmanni.
verticillatum, Nees, in Linnæa, ix. (1834) 303 = S. verticillata.
virgatum, Nees, in Mart. Fl. Bras. ii. i. 173 = S. virgata.

HYPOPTERON, Hassk. Cat. Hort. Bog. Alt. 154 (1844) = **Chirita**, Buch.-Ham. (Gesnerac.).

HYPOPTERYGIUM, Schlecht. in Linnaea, xvii. (1843) 685 = **Julianis**, Schlecht. (Anacard.).
adstringens, Schlecht. l. c. = ♩. adstringens.

HYPOPYTHIS, Refn. in Med. Repos. N. York, v. (1808) 352 = **Hypopitys**, Dill. (Monotr.).

HYPOTHRONIA, Schrank, in Syll. Ratisb. t. (1824) 85 = **Hyptia**, Jacq. (Labiat.).
undata, Schrank, l. c. = H. fasciculata.

HYPOXANTHUS, Rich. ex DC. Prod. iii. 180 (1828) = **Miconia**, Ruiz & Pav. (Melastom.).

HYPOXIDOPSIS, Steud. ex Baker, in Journ. Linn. Soc. xvii. (1879) 450 = **Iphigenia**, Kunth (Liliac.).
pumila, Steud. ex Baker, l. c. = I. indica.

HYPOXIS, Forsk. Fl. Aegypt. Arab. 74 = **Scilla** ♩, Linn. (Liliac.).

HYPOXIS, Linn. Syst. ed. X. 986 (1759). *AMAEVLLIDEAE*, Benth. & Hook. f. iii 717.
FRANQUEVILLEA, Zoll. ex Miq. Fl. Ind. Bat. iii. 586 (1856).
IANTHE, Salisb. Gen. Pl. Fragm. 44 (1866).
NIOBEA, Willd. ex Schult. f. Syst. vii. 762 (1830).
SPILOXENE, Salisb. Gen. Pl. Fragm. 44 (1866).
abyssinica, Hochst. in Flora, xxiv. (1841) i. Intell. 21 = villosa.
acuminata, *Eckl. Top. Verr.* 9.—Afr. austr.
affinis, Schult. f. Syst. vii. 774 = alba.
alba, *Linn. f. Suppl.* 198.—Afr. austr.
Andrewsii, *Baker, in Journ. Linn. Soc.* xvii. (1878) 104.—Afr. austr.
angolensis, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 266.—Afr. trop.
angustifolia, *Lam. Encyc.* iii. 182.—Ins. Borbon.
aquatica, *Linn. f. Suppl.* 197.—Afr. austr.
argentea, *Harv. ex Baker, in Journ. Linn. Soc.* xvii. (1878) 110.—Afr. austr.
Arnottii, *Baker, in Gard. Chron.* (1877) ii. 552.—Afr. austr.
aurea, Eckl. Top. Verr. 10 = stellata.
aurea, *Lour. Fl. Cochinch.* 200.—Cochinch.
Bauri, *Baker, in Journ. Bot.* xiv. (1876) 181.—Afr. austr.
bifora, Baker, l. c. = angustifolia.
brachystachya, Wight, Ic. t. 2046 = Curculigo Finlaysoniana.
breviscapa, H. B. & K. Nov. Gen. et Sp. i. 287 = decumbens.
cærulescens, [DC. in] Red. Lil. sub t. 169 = stellata.
canaliculata, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 265.—Afr. austr.
canescens, Fisch. & Mey. Ind. Sem. Hort. Petrop. x. 50 = villosa.
caricifolia, Salisb. Prod. 248 = decumbens.
carolinensis, Michx. Fl. Bor. Am. i. 188 = erecta.
colchicifolia, *Baker, in Gard. Chron.* (1884) ii. 649.—Afr. austr.
costata, *Baker, in Journ. Linn. Soc.* xvii. (1878) 119.—Afr. austr.
crassifolia, Pappe, ex Baker, l. c. 102 = alba.
cunnensis, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 266.—Afr. trop.
curculigoides, Wall. Cat. n. 6184 = aurea.
decumbens, Aubl. Pl. Gui. 304 = Curculigo scorzoneræfolia.
decumbens, Lam. Encyc. iii. 172 = villosa.
decumbens, *Linn. Pl. Sam. Pugill.* 11; *Syst.* ed. X. 986.—Am. trop.
dubia, Schult. f. Syst. vii. 775 = alba.
dulcis, Steud. ex Baker, in Journ. Linn. Soc. xvii. (1878) 124 = Curculigo orchioides.
Eckloniana, *Schult. f. Syst.* vii. 1775, *Index*.—Afr. austr.
elata, Hook. f. Bot. Mag. t. 5690 = hemerocallidea.
elata, Schult. f. Syst. vii. 778 = stellata.
elegans, Poir. Encyc. Suppl. iii. 113 = stellata.
elongata, H. B. & K. Nov. Gen. et Sp. i. 287 = decumbens.
erecta, *Linn. Syst.* ed. X. 986.—Am. bor.
Fabricia, Gaertn. Fruct. i. 33. t. 11 = villosa.
farinosa, Thunb. Fl. Jap. 83 = Aletris japonica.
fascicularis, Linn. Syst. ed. X. 986 = Colchicum montanum.
filifolia, *Eckl. Top. Verr.* 10.—Afr. austr.
filifolia, Ell. Sketch, 897 = juncea.

HYPOXIS—
filiformis, *Baker, in Journ. Linn. Soc.* xvii. (1878) 109.—Afr. austr.
flavescens, *Eckl. Top. Verr.* 9.—Afr. austr.
flavopetala, *Eckl. l. c.* 10.—Afr. austr.
Frankquevillea, Miq. Fl. Ind. Bat. iii. 586 = aurea.
Gawleri, Baker, in Journ. Linn. Soc. xvii. (1878) 101 = stellata.
geniculata, *Eckl. Top. Verr.* 9.—Afr. austr.
Gerrardi, *Baker, in Journ. Linn. Soc.* xvii. (1878) 110.—Afr. austr.
glabella, *R. Br. Prod.* 289.—Austral.
gracilis, Lehm. ex Schult. f. Syst. vii. 764 = decumbens.
graminea, Pursh, Fl. Am. Sept. i. 224 = erecta.
graminea, *Willd. ex Schult. f. Syst.* vii. 768.—Madag.
hemerocallidea, *Fisch. Mey. & Ave-Lall. Ind. Sem. Hort. Petrop.* viii. 64.—Afr. austr.
humilis, H. B. & K. Nov. Gen. et Sp. i. 286 = decumbens.
hygrometrica, *Labill. Nov. Holl. Pl.* i. 18. t. 108.—Austral.
iridifolia, *Baker, in Journ. Linn. Soc.* xvii. (1878) 117.—Afr. trop.
Jacquini, *Baker, in Gard. Chron.* (1877) ii. 552.—Afr. austr.
juncea, Eckl. Top. Verr. 10 = stellata.
juncea, *Sm. Spicil.* 15.—Am. bor.
Krausiana, *Buching. ex Krauss, in Flora*, xxviii. (1845) 811.—Afr. austr.
Krebsii, *Fisch. in Ind. Sem. Hort. Petrop.* xi. 72.—Afr. austr.
lanata, Eckl. ex Baker, in Journ. Linn. Soc. xvii. (1878) 118 = stellipila.
latifolia, *Hook. Bot. Mag.* t. 4817.—Afr. austr.
latifolia, Wight, Ic. t. 2044 = Curculigo Finlaysoniana.
laxa, *Eckl. Top. Verr.* 9.—Afr. austr.
leptocarpa, Benth. Fl. Austral. vi. 451 = occidentalis.
leptostachya, Wight, Ic. t. 2045 = Curculigo Finlaysoniana.
linearis, Andr. Bot. Rep. t. 171 = stellata.
longifolia, *Baker, in Bot. Mag.* t. 6035.—Afr. austr.
longifolia, *Eckl. Top. Verr.* 9.—Afr. austr.
Ludwigii, *Baker, in Journ. Bot.* xiv. (1876) 181.—Afr. austr.
lunulæfolia, [DC. in] Red. Lil. t. 260 = Curculigo plicata.
lunulæfolia, Eckl. Top. Verr. 10 = serrata.
marginata, *R. Br. Prod.* 289.—Austral.
membranacea, *Baker, in Journ. Linn. Soc.* xvii. (1878) 106.—Afr. austr.
mexicana, Schult. f. Syst. vii. 761 = decumbens.
microsperma, *Ave-Lall. in Ind. Sem. Hort. Petrop.* x. 50.—Afr. austr.
millifolia, *Baker, in Journ. Linn. Soc.* xvii. (1878) 105.—Afr. austr.
minor, D. Don, Prod. Fl. Nep. 53 = aurea.
minor, *Eckl. Top. Verr.* 10.—Afr. austr.
minor, Sesm. Bot. Voy. Herald. 419 = Curculigo orchioides.
minuta, Eckl. ex Schult. f. Syst. vii. 774 = Pauridia hypoxidoides.
minuta, *Linn. f. Suppl.* 197.—Afr. austr.
monanthos, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. 266.—Afr. austr.
multiceps, *Buching. ex Krauss, in Flora*, xxviii. (1845) 811.—Afr. austr.
nana, E. Mey. ex Baker, in Journ. Linn. Soc. xvii. (1878) 126 = Pauridia hypoxidoides.
natalensis, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 164.—Afr. austr.
obliqua, Andr. Bot. Rep. t. 198 = Andrewsii.
obliqua, Eckl. & Zeyh. ex Baker, in Journ. Linn. Soc. xvii. (1878) 102 = alba.
obliqua, Jacq. Coll. Suppl. 54 = villosa.
obtusa, *Burch.* [*ex Ker-Gawl. in*] *Bot. Reg.* t. 159.—Afr. austr.
occidentalis, *Benth. Fl. Austral.* vi. 451.—Austral.
orchioides, Kurz, in Mig. Ann. Mus. Bot. Lugd. Bat. iv. 177 = Curculigo orchioides.
ovata, *Linn. f. Suppl.* 197.—Afr. austr.
pallida, Salisb. Prod. 248 = erecta.
parviflora, Baker, in Gard. Chron. (1874) ii. 180 = villosa.
parvula, *Baker, in Journ. Linn. Soc.* xvii. (1878) 113.—Afr. austr.
pauciflora, Wight, Ic. t. 2046 = Curculigo Finlaysoniana.

HYPOXIS—
Pavonina, Salisb. Gen. Pl. Fragm. 44 = stellata.
Petitiana, A. Rich. Tent. Fl. Abyss. ii. 315 = villosa.
platypetala, *Baker, in Journ. Linn. Soc.* xvii. (1878) 105.—Afr. austr.
plicata, Jacq. Coll. Suppl. 55 = Curculigo veratrifolia.
plicata, Linn. f. Suppl. 197 = Curculigo plicata.
polystachya, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 266.—Afr. austr.
pratensis, R. Br. Prod. 289 = hygrometrica.
pumila, Lam. Encyc. iii. 184 = minuta.
pusilla, *Hook. f. Fl. Tasm.* ii. 26. t. 130 B.—Austral.
pusilla, H. B. & K. Nov. Gen. et Sp. i. 286 = decumbens.
pusilla, *Presl, Bot. Bemerk.* 116.—Afr. austr.
ramosa, Lodd. Bot. Cab. t. 1936 (Romuliæ sp.).—Afr. austr.
rhizophylla, Baker = *Molinaria rhizophylla*.—Ins. Mascar.
rigidula, *Baker, in Journ. Linn. Soc.* xvii. (1878) 116.—Afr. austr.
Rooperii, *T. Moore, Gard. Comp.* (1852) 65.—Afr. austr.
scabra, Lodd. Bot. Cab. t. 970 = villosa.
scabra, *Presl, Bot. Bemerk.* 116.—Afr. austr.
Schimperi, *Baker, in Journ. Linn. Soc.* xvii. (1878) 110.—Afr. austr.
Schnitzleinia, Hochst. in Flora, xxvii. (1844) 31 = (Velloziæ sp.).—Abyssin.
scorzoneræfolia, Lam. Encyc. iii. 188 = Curculigo scorzoneræfolia.
sericea, *Baker, in Journ. Linn. Soc.* xvii. (1878) 111.—Afr. austr.
serrata, *Linn. f. Suppl.* 197.—Afr. austr.
sessilis, Linn. Sp. Pl. ed. II. 439 = erecta.
setosa, *Baker, in Journ. Linn. Soc.* xvii. (1878) 118.—Afr. austr.
sinensis, Hochst. in Flora, xxvii. (1844) 32 = villosa.
sobolifera, Jacq. Coll. Suppl. 53 = villosa.
spathacea, Eckl. Top. Verr. 11 = alba.
spicata, Thunb. Fl. Jap. 136 = Aletris japonica.
stellata, *Linn. f. Suppl.* 197.—Afr. austr.
stellipila, [*Ker-Gawl.*] *in Bot. Reg.* t. 668.—Afr. austr.
subrigma, *Willd. ex Schult. f. Syst.* vii. 779.—Am. austr.
tabularis, Eckl. Top. Verr. 10 = stellata.
tenuifolia, *Eckl. l. c.*—Afr. austr.
tomentosa, Lam. Encyc. iii. 112 = villosa.
truncata, Pappe, ex Baker, in Journ. Linn. Soc. xvii. (1878) 126 = Pauridia hypoxidoides.
trichocarpa, Wight, Ic. t. 2045 = Curculigo Finlaysoniana.
tridentata, [DC. in] Red. Lil. sub t. 169 = stellata.
trifora, Harv. ex Baker, in Journ. Linn. Soc. xvii. (1878) 102 = minuta.
truncata, Thunb. ex Schult. f. Syst. vii. 773 = Pauridia hypoxidoides.
vaginata, Schlecht. in Linnaea, xx. (1847) 568 = glabella.
veratrifolia, Willd. Sp. Pl. ii. 109 = Curculigo veratrifolia.
villosa, Jacq. Coll. Suppl. 51 = Jacquini.
villosa, *Linn. f. Suppl.* 198.—Afr. austr.
violacea, *Schult. f. Syst.* vii. 779.—Arabia.
Zeyheri, *Baker, in Journ. Linn. Soc.* xvii. (1878) 112.—Afr. austr.

HYPPOCHAERIS, Bivona, Cent. Pl. ii. 31 (1806) = **Hypochoeris**, Linn. (Compos.).

HYPSELA, Presl, Prod. Monog. Lobel. 45 (1836). *CAMPANULACEAE*, Benth. & Hook. f. ii. 550.
FRATIA, Wedd. Chlor. And. ii. 9. t. 45 (1857), partim.
atacamensis, Phil. f. Cat. Pl. Vasc. Chil. 188 = Fratia longiflora.
longiflora, *Benth. & Hook. f. Gen.* ii. 550.—Reg. Magell.
longiflora, Phil. f. Cat. Pl. Vasc. Chil. 188 = Fratia longiflora.
reniformis, *Presl, Prod. Monog. Lobel.* 45.—Reg. Andin.

HYPSEOCHARIS, Remy, in Ann. Sc. Nat. Sér. III. viii. (1847) 238. *GERANIACEAE*, Benth. & Hook. f. i. 276.
pimpinellifolia, *Remy, l. c.* 239.—Peruv.

HYPSERPA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40 = **Limacia**, Lour. (Menisp.).
borneensis, Becc. Malesia, i. (1877) 149 = L. borneensis.
cuspidata, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40; et Ser. III. xiv. (1864) 365 = L. cuspidata.
funifera, Miers, l. c. 365.—Afr. trop.
heteromera, Miers, l. c. = L. borneensis.
nitida, Miers, ex Benth. in Hook. Kew Journ. iii. (1851) 258 = L. cuspidata.
paucifora, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 365 = L. cuspidata?
polyandra, Becc. Malesia, i. (1877) 148.—N. Guinea.
procurvicarata, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 365 = L. cuspidata.
propensa, Miers, l. c.—Borneo.
selebica, Becc. Malesia, i. (1877) 143.—Ins. Celebes.
Selwyni, F. Muell. Fragm. ix. 82 = Selwynia laurina.
trifiora, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 365 = L. microphylla.
unifora. Miers, l. c. = L. cuspidata.

HYPSIPODES, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 82 (1868–69). *MENISPERMACEAE.*
subcordatus, *Miq. l. c.*—Java.

HYPTIANDRA, Hook. f. in Benth. & Hook. f. Gen. i. 293 (1862). *SIMARUBEAE*, Benth. & Hook. f. i. 393, 990.
Bidwillii, *Hook. f. l. c.* 294.—Austral.

HYPTIANTHERA, Wight & Arn. Prod. 399 (1834). *RUBIACEAE*, Benth. & Hook. f. ii. 94.
macrocarpa, Thw. Enum. Pl. Zeyh. 187 = Nargedia macrocarpa
rhamnoides. Zoll. & Mor. Syst. Verz. Zoll. 60 = Plectronia horrida.
stricta, Wall. Cat. n. 8813 L = Lasianthus inconspicuus.
stricta, *Wight & Arn. Prod.* 399.—Ind. or.; Burma.

HYPTIS, Jacq. Coll. i. 101 (1786). *LABIATAE*, Benth. & Hook. f. ii. 1178.
BROTERA, Spreng. in Trans. Linn. Soc. vi. (1802) 151. t. 13.
GNOTERIS, Rafin. Sylva Tellur. 76 (1838).
HIPPOTHRONIA, Benth. Lab. Gen. et Sp. 130, in syn. (1833).
HYPOTHRONIA, Schrank, in Syll. Ratisb. i. (1824) 85.
MESOSPHAERUM, P. Br. Hist. Jamaic. 257 (1756).
RHAPHIODON, Schau. in Flora, xxvii. (1844) 345.
SCHAUERIA, Hassk. l. c. xxv. (1842) 11; Beibl. 35.
STAGONARRHEN, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 146 (1858).
actinocephala, *Griseb. Cat. Pl. Cub.* 212.—Cuba.
acuta, Benth. in Linnaea, vi. (1831) 82 = brevipes.
adamantium, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 91. —Bras.
adpressa, *A. St. Hil. ex Benth. l. c.* 84.—Bras.
affinis, *Benth. l. c.* 109.—Ecuador.
albida, *H. B. & K. Nov. Gen. et Sp.* ii. 219.—Mexic.
albipes, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 88. —Bras.
alpestris, *A. St. Hil. ex Benth. l. c.* 92.—Bras.
alsheastifolia, *Pohl, ex Benth. l. c.* 115.—Bras.
altissima, *A. St. Hil. ex Benth. l. c.* 135.—Bras.
alutacea, *Pohl, ex Benth. l. c.* 112.—Bras.
amethystoides, *Benth. in DC. Prod.* xii. 130.—Bras.
ammotropha, *Griseb. Cat. Pl. Cub.* 213.—Cuba.
angulosa, *Schott, ex Benth. Lab. Gen. et Sp.* 102.— Bras.
angustifolia, *Pohl, ex Benth. l. c.* 89.—Bras.
anomala, Benth. l. c. 118 = conferta.
aquatica, Pohl, ex Benth. l. c. 116 = carpinifolia.
arenipenula, *Benth. in DC. Prod.* xii. 93.—Bras.
arbores, *Benth. l. c.* 132.—Cuba.
arrnaria, Benth. l. c. 104 = linarioides.
asgrata, Pohl, ex Benth. Lab. Gen. et Sp. 86 = rugosa.
axilis, A. St. Hil. ex Benth. l. c. 116 = violacea.
aristata, *Benth. in DC. Prod.* xii. 111.—Bras.
arvensis, Poepp. ex Benth. Lab. Gen. et Sp. 712 = osmoscma.
aspera, Mart. & Gal. in Bull. Acad. Brux. xi. ii. (1844) 189 = spicata.
asteroides, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 93. —Bras.
atzoubens, *Poit. in Ann. Mus. Par.* vii. (1806) 466. t. 27. f. 3.—Am. trop.

HYPTIS:—
barbata, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 69.—Bras.
Beschkena, *Kunze, in Linnaea*, xx. (1847) 29.—Bras.
bisdentata, Pohl, ex Benth. Lab. Gen. et Sp. 87 = longipes.
Blanchetii, *Benth. l. c.* 712.—Bras.
brachyphylla, *Mart. ex Benth. l. c.* 96.—Bras.
brachystachya, *Pohl, ex Benth. l. c.* 127.—Bras.
brevipes, *Poit. in Ann. Mus. Par.* vii. (1806) 465.— Am. austr.
brunnescens, *Pohl, ex Benth. Lab. Gen. et Sp.* 88.— Bras.
caespitosa, *A. St. Hil. ex Benth. l. c.* 113.—Bras.
calida, *Mart. ex Benth. l. c.* 131.—Bras.
calophylla, *A. St. Hil. ex Benth. l. c.* 138.—Bras.
calycina, *Pohl, ex Benth. l. c.* 140.—Bras.
campanulata, *Benth. in DC. Prod.* xii. 127.—Bras.
camporum, *Benth. l. c.* 92.—Bras.
cana, *Pohl, ex Benth. Lab. Gen. et Sp.* 125.—Bras.
candida, Pohl, ex Benth. l. c. 85 = nervosa.
canescens, Benth. l. c. 120, partim = polystachya.
canescens, *H. B. & K. Nov. Gen. et Sp.* ii. 221.—Am. austr.
capitata, *Jacq. Coll.* i. 102; *Ic. Rar.* 3. t. 114.—Am. austr.
capreolifolia, *Pohl, ex Benth. Lab. Gen. et Sp.* 83.— Bras.
cardiophylla, *Pohl, ex Benth. l. c.* 84.—Bras.
carpinifolia, *Benth. l. c.* 115.—Bras.
Chamædrys, Willd. Sp. Pl. iii. 85 = Marsypianthes hyptioides.
Chamissonis, *Benth. in Linnaea*, vi. (1831) 570.—Bras.
Clausseni, *Benth. in DC. Prod.* xii. 131.—Bras.
clavellifera, *Benth. l. c.* 91.—Bras.
coccines, *Mart. ex Benth. Lab. Gen. et Sp.* 139.— Bras.
coerulea, *A. St. Hil. ex Benth. l. c.* 91.—Bras.
communis, *A. St. Hil. ex Benth. l. c.* 128.—Bras.
complicata, *A. St. Hil. ex Benth. l. c.* 111.—Bras.
concinna, *Benth. in DC. Prod.* xii. 126.—Bras.
conferta, *Pohl, ex Benth. Lab. Gen. et Sp.* 112.—Bras.
consperna, *Benth. in DC. Prod.* xii. 134.—Bras.
cordata, Pohl, ex Benth. Lab. Gen. et Sp 110 = remota.
coriacea, *Benth. in DC. Prod.* xii. 95.—Bras.
corymbosa, *Benth. l. c.* 112.—Bras.
crassifolia, *Mart. ex Benth. Lab. Gen. et Sp.* 94.— Bras.
crenata, *Pohl, ex Benth. l. c.* 93.—Bras.
crinita, *Benth. l. c.* 95.—Bras.
crocodilites, Lour. Fl. Cub. Fanerog. ii. 157, sphalm. = eriocauloides.
crispata, *Pohl, ex Benth. Lab. Gen. et Sp.* 139.—Bras.
cuneata, *Pohl, ex Benth. l. c.* 86.—Bras.
cymulosa, *Benth. in DC. Prod.* xiii. ii. 459.—Bras.
densiflora, *Pohl, ex Benth. Lab. Gen. et Sp.* 137.— Bras.
densifolia, *Pohl, ex Benth. l. c.* 100.—Bras.
desertorum, Pohl, ex Benth. l. c. 88 = brunnescens.
dicticocalyx, *Benth. in DC. Prod.* xii. 131.—Bras.
dictyoda, *Pohl, ex Benth. Lab. Gen. et Sp.* 91.—Bras.
dilatata, *Benth. in DC. Prod.* xii. 108.—Cuban.
discolor, Hort. Berol. ex Lehm. Ind. Sem. Hort. Hamb. (1830) 5 (Quid?).—Hab. ?
ditasoides, *Mart. ex Benth. Lab. Gen. et Sp.* 99.— Bras.
divaricata, *Pohl, ex Benth. l. c.* 82.—Bras.
diversifolia, *Benth. Pl. Hartw.* 144.—Peruv.
dubia, *Pohl, ex Benth. Lab. Gen. et Sp.* 128.—Bras.
duplicato-dentata, Pohl, ex Benth. l. c. 114 = vestita.
dyctioides, Benth. in DC. Prod. xii. 101 = dictyoda.
ebractesta, R. Br. in Ait. Hort. Kew. ed. II. iii. 391 = suaveolens.
elongata, *Benth. Lab. Gen. et Sp.* 88.—Peruv.
Emoryi, *Torr. in Pacs, Rep. Colorado River, Bot.* 20. Calif.
eriocalyx, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 131.— Bras.
eriocauloides, *A. Rich. Fl. Cub. Fanerog.* ii. 157 (err. typ. eriocaulolides).—Ins. Pinet.
eriocephala, *Benth. in DC. Prod.* xii. 124.—Peruv.
eriophylla, *Pohl, ex Benth. Lab. Gen. et Sp.* 96.— Bras.
excelsa, *Mart. & Gal. in Bull. Acad. Brux.* xi. ii. (1844) 193.—Mexic.
expansa, *Pohl, ex Benth. Lab. Gen. et Sp.* 123.—Bras.
fasciculata, *Benth. l. c.* 130.—Bras.

HYPTIS:—
fastigiata, Benth. Lab. Gen. et Sp. 130 = fasciculata.
ferruginea, *Benth. l. c.* 118.—Mexic.
ferruginosa, *Pohl, ex Benth. l. c.* 109.—Bras.
flabellata, Pohl, ex Benth. l. c. 122 = communis.
florida, *Benth. Bot. Voy. Sulph.* 150.—Ecuador.
foliosa, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 117. —Bras.
fragilifolia, *A. St. Hil. ex Benth. l. c.* 85.—Bras.
fruticosa, *Salem. ex Benth. l. c.* 123.—Bras.
Gaudichaudii, *Benth. in Linnaea*, vi. (1831) 77.— Bras.
glechomoides, *Weinm. Hort. Pawl.* 270 (Quid?).— Bras.
glandulosa, Sieber, ex Benth. Lab. Gen. et Sp. 244 = Salvia occidentalis.
glauca, *A. St. Hil. ex Benth. l. c.* 141.—Bras.
globifera, G. F. W. Mey. Prim. Fl. Esseq. 207 = brevipes.
glomerata, Hort. ex Presl, Bot. Bemerk. 101 = Marsypianthes glomerata.
glomerata, *Mart. ex Schrank, in Syll. Ratisb.* ii. (1828) 56.—Bras.
glutinosa, *Benth. in DC. Prod.* xii. 130.—Bras.
gonocephala, *Wright, ex Griseb. Cat. Pl. Cub.* 212.— Cuba.
goyazensis, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 92. —Bras.
graticolafolia, *A. St. Hil. ex Benth. l. c.*—Bras.
graveolens, Mart. ex Benth. in DC. Prod. xii. 126 = suaveolens.
graveolens, *Salem. ex Benth. Lab. Gen. et Sp.* 125.— Bras.
graveolens, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 62 = umbrosa.
halimifolia, *Mart. ex Benth. Lab. Gen. et Sp.* 94.— Bras.
heterantha, *Benth. in DC. Prod.* xii. 134.—Bras.
heterophylla, *Benth. l. c.* 101.—Bras.
Hilarii, *Benth. Lab. Gen. et Sp.* 90.—Bras.
hirsuta, *H. B. & K. Nov. Gen. et Sp.* ii. 218. t. 161.— Am. austr.
hispida, *Benth. in DC. Prod.* xii. 118.—Bras.
hispida, Poepp. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 83 = spicigera.
homalophylla, *Pohl, ex Benth. Lab. Gen. et Sp.* 107.— Bras.
humilis, *Benth. in DC. Prod.* xii. 118.—Bras.
hypoleuca, *Benth. l. c.* 135.—Bras.
imbricata, *Pohl, ex Benth. Lab. Gen. et Sp.* 98.— Bras.
inamœna, *Benth. l. c.* 104.—Bras.
incana, Willd. ex Stend. Nom. ed. II. i. 794 = tomentosa.
incisa, *A. St. Hil. & Mart. ex Benth. Lab. Gen. et Sp.* 87.—Bras.
infasta, Spreng. Syst. ii. 731 = Marsypianthes hyptioides.
insolora, Schrank, in Syll. Ratisb. ii. (1828) 56 = muricata.
interrupta, *Pohl, ex Benth. Lab. Gen. et Sp.* 77.—Bras.
involucrata, *Benth. l. c.* 108.—Mexic.
involucrata, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 190 = alutacea.
irregularis, *Benth. in DC. Prod.* xii. 127.—Bras.
Itatiaiae, *Wawra, in Oestr. Bot. Zeitschr.* xxxi. (1881) 70.—Bras.
laciniata, *Benth. in Hook. Journ. Bot.* ii. (1840) 49.— Guian.
lacunosa, *Pohl, ex Benth. Lab. Gen. et Sp.* 94.—Bras.
lacustris, *A. St. Hil. ex Benth. l. c.* 107.—Bras.
lagenaria, *A. St. Hil. ex Benth. l. c.* 98.—Bras.
lamifolia, *Weinm. in Syll. Ratisb.* i. (1824) 235, 226. —Bras.
lanata, *Pohl, ex Benth. Lab. Gen. et Sp.* 84.—Bras.
lancifolia, *Thonn. in Schum. & Thonn. Beschr. Guin. Pl.* 261.—Afr. trop.
lanceolata, *Poir. Encyc. Suppl.* iii. 114.—Bras.
lanifiora, *Benth. Bot. Voy. Sulph.* 42. t. 20.—Calif.
lantanæfolia, *Poit. in Ann. Mus. Par.* vii. (1806) 466. t. 29. f. 1.—Am. austr.
lappacea, *Benth. Lab. Gen. et Sp.* 103.—Bras.
lappulacea, *Mart. ex Benth. l. c.* 104.—Bras.
latifolia, *Mart. ex Benth. l. c.* 126.—Bras.
latifolia, J. A. Schmidt, in Mart. Fl. Bras. viii. 114 = linarioides.
laurifolia, *A. St. Hil. ex Benth. l. c.* 134.—Bras.
lavandulæfolia, *Pohl, ex Benth. l. c.* 82.—Bras.
lavendulæfolia, Stend. Nom. ed. II. i. 794 = pruec.

HYPTIS :—
lasiflora, *Mart. ex Benth. Lab. Gen. et Sp.* 189.—Bras.
leptoclada, *Benth. in DC. Prod.* xii. 100.—Bras.
leucocephala, *Mart. ex Benth. Lab. Gen. et Sp.* 89.—Bras.
leucophylla, *Pohl, ex Benth. l. c.* 184.—Bras.
lilacina, *Cham. & Schlecht. in Linnaea*, v. (1830) 101.—Mexic.
linarioides, *Pohl, ex Benth. Lab. Gen. et Sp.* 99.—Bras.
lipptioides, *Pohl, ex Benth. l. c.* 86.—Bras.
lobata, *A. St. Hil. ex Benth. l. c.* 97.—Bras.
longiflora, Pohl, ex Benth. l. c. 140 = reticulata.
longifolia, *Pohl, ex Benth. l. c.* 106.—Bras.
longipes, *A. St. Hil. ex Benth. l. c.* 88.—Bras.
lophantha, *Mart. ex Benth. l. c.* 76.—Bras.
Lorentziana, *O. Hoffm. in Linnaea*, xliii. (1880–82) 137.—Reg. Argent.
lucida, *Pohl, ex Benth. Lab. Gen. et Sp.* 109.—Bras.
Lundii, *Benth. in DC. Prod.* xii. 111.—Bras.
lurida, Spreng. Syst. ii. 781 = Marsypianthes hyptoides.
lutescens, *Pohl, ex Benth. Lab. Gen. et Sp.* 109.—Bras.
lythroides, *Pohl, ex Benth. l. c.* 118.—Bras.
macrantha, *A. St. Hil. ex Benth. l. c.* 140.—Bras.
macrocephala, *Mart. & Gal. in Bull. Acad. Brux.* xi. 11. (1844) 186.—Mexic.
macrochila, *Mart. ex Steud. Nom.* ed. II. i. 794.—Bras.
macrostachys, *Benth. in DC. Prod.* xii. 130.—Bras.
madagascariensis, Boj. Hort. Maurit. 251 = spicigera.
malacophylla, *Benth. in DC. Prod.* xii. 86.—Bras.
marathrosma, Benth. l. c. 140 = Lophantha anisata.
marginata, *J. A. Schmidt, in Mart. Fl. Bras.* viii. 108.—Bras.
martiola, *Benth. in DC. Prod.* xii. 126.—Bras.
marrubiaea, Benth. l. c. 105 = sinuata.
marrubiastra, Pohl, ex Benth. Lab. Gen. et Sp 102 = sinuata.
Martiusi, *Benth. l. c.* 95.—Bras.
melanosticta, *Griseb. Fl. Brit. W. Ind.* 488.—Ind. occ.
melissoides, *H. B. & K. Nov. Gen. et Sp.* ii. 320.—N. Granat.
melochioides, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 187.—Bras.
membranacea, *Benth. l. c.* 132.—Bras.
menthodora, Schrank, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 94 = lophantha.
micrantha, Pohl, ex Benth. Lab. Gen. et Sp. 120 = spicata.
microcephala, Bert. ex Benth. l. c. 81 = recurvata.
micrantha, *Pohl, ex Benth. l. c.* 82.—Bras.
Mikanii, J. A. Schmidt, in Mart. Fl. Bras. viii. 138 = umbrosa.
mimtiflola, *Griseb. Cat. Pl. Cub.* 213.—Cuba.
Moçiniana, *Benth. Lab. Gen. et Sp.* 139.—Mexic.
mollis, *Pohl, ex Benth. l. c.* 102.—Bras.
mollissima, *Benth. l. c.* 85.—Bras.
monticola, *Mart. ex Benth. l. c.* 111.—Bras.
multibracteata, *Benth. l. c.* 100.—Bras.
multiflora, *Pohl, ex Benth. l. c.* 96.—Bras.
multiaria, Benth. in DC. Prod. xii. 123 = spicata.
muricata, *Schott, ex Benth. Lab. Gen. et Sp.* 119.—Bras.
nepalensis, Lehm. ex Steud. Nom. ed. II. i. 794 (sp. ignot.).—Nepal.
nepetoides, Flach. ex Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 83 = pectinata.
nervosa, *Pohl, ex Benth. Lab. Gen. et Sp.* 85.—Bras.
nicaraguensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 84.—Nicaragua.
nigrescens, *Pohl, ex Benth. Lab. Gen. et Sp.* 111.—Bras.
nitida, Mart. & Gal. in Bull. Acad. Brux. xi. 11. (1844) 188 = Salvia nitida.
nitidula, *Benth. in DC. Prod.* xii. 91.—Bras.
nudiuscula, *Benth. Lab. Gen. et Sp.* 79.—Bras.
oblongifolia, *Benth. in DC. Prod.* xii. 125.—Mexic.
obscura, *Benth. l. c.* 99.—Bras.
obtusata, *Benth. Pl. Hartw.* 241.—Am. austr.
obtusiflora, *Presl, ex Benth. Lab. Gen. et Sp.* 107.—Peruv.
obtusifolia, *R. Br. in Tuckey, Voy. Congo App.* 478.—Afr. trop.
obvallata, *Sprang. ex Benth. Lab. Gen. et Sp.* 115.—Bras.

HYPTIS :—
odorata, *Benth. Lab. Gen. et Sp.* 81.—Peruv.
oppositiflora, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 83 = glomerata.
oppositifolia, J. A. Schmidt, in Mart. Fl. Bras. viii. 138 = glomerata.
orbiculata, *Pohl, ex Benth. Lab. Gen. et Sp.* 111.—Bras.
organoides, Pohl, ex Benth. l. c. 91 = velutina.
ovalifolia, *Benth. in DC. Prod.* xii. 87.—Bras.
ovata, *Pohl, ex Benth. Lab. Gen. et Sp.* 92.—Bras.
paludosa, *A. St. Hil. ex Benth. l. c.* 82.—Bras.
paniculata, *Benth. l. c.* 189.—Bras.
Parkeri, *Benth. l. c.* 106.—Am. austr.
parviflora, Mart. & Gal. in Bull. Acad. Brux. xi. 11. (1844) 186 = verticillata.
parvistylis, Pohl, ex Benth. Lab. Gen. et Sp. 118 = subrotunda.
pascuorum, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 118 = atrorubens.
Passerina, *Mart. ex Benth. Lab. Gen. et Sp.* 89.—Bras.
pauciflora, *Pohl, ex Benth. l. c.* 117.—Bras.
pectinata, *Pohl. in Ann. Mus. Par.* viii. (1806) 474. t. 30.—Cosmop. trop.
pedalipes, *Griseb. Cat. Pl. Cub.* 213.—Cuba.
pedunculata, *Benth. Lab. Gen. et Sp.* 87.—Bras.
perzica, Polt. in Ann. Mus. Par. vii. (1806) 471; Roth, Catalect. fasc. iii. 48 = pectinata.
petiolata, *Pohl, ex Benth. Lab. Gen. et Sp.* 101.—Bras.
petræa, *A. St. Hil. ex Benth. l. c.* 117.—Bras.
pilosa, *Benth. l. c.* 134.—Bras.
pistanifolia, *Mart. ex Benth. l. c.* 77.—Bras.
plectranthoides, *Benth. l. c.* 122.—Bras.
Plumieri, Polt. in Ann. Mus. Par. vii. (1806) 473 = suaveolens.
plumosa, *Benth. in DC. Prod.* xii. 94.—Bras.
Pohliana, *Fzacy. ex Benth. l. c.* 140.—Bras.
polyantha, *Polt. in Ann. Mus. Par.* vii. (1806) 470.—Peruv.
polystachya, *H. B. & K. Nov. Gen. et Sp. ii.* 321.—Am. trop.
procumbens, Schlecht. & Cham. in Linnæa, v. (1830) 101 = atrorubens.
protesoides, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 110.—Bras.
pruinosa, *Pohl, ex Benth. l. c.* 187.—Bras.
Pseudachamaedrys, Polt. in Ann. Mus. Par. vii. (1806) 469 = Marsypianthes hyptoides.
pubescens, *Benth. Lab. Gen. et Sp.* 130.—Mexic.
pulegioides, *Pohl, ex Benth. l. c.* 128.—Bras.
pumila, Pohl, ex Benth. l. c. 122 = communis.
punctata, *Mart. & Gal. in Bull. Acad. Brux.* xi. 11. (1844) 188.—Mexic.
Purdiei, *Benth. in DC. Prod.* xii. 125.—N. Granat.
pycnocephala, *Benth. l. c.* 102.—Bras.
racemosa, Pohl, ex Benth. l. c. 132 = spicata.
racemosa, Zuccagni, in Roem. Collect. 143 = pectinata.
racemulosa, *Mart. ex Benth. Lab. Gen. et Sp.* 126.—Bras.
radiata, H. B. & K. Nov. Gen. et Sp. ii. 318 = brevipes.
radiata, *Willd. Sp. Pl.* iii. 84.—Am. trop.
Radula, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 121 = conferta.
ramosa, *Pohl, ex Benth. Lab. Gen. et Sp.* 104.—Bras.
recurvata, *Polt. in Ann. Mus. Par.* vii. (1806) 467. t. 28. f. 2.—Am. trop.
reflexa, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 83.—Bras.
remota, *Pohl, ex Benth. l. c.* 110.—Bras.
reticulata, *Mart. ex Benth. l. c.* 141.—Bras.
rhabdocalyx, *Mart. ex Benth. l. c.* 133.—Bras.
rhomboidea, *Mart. & Gal. in Bull. Acad. Brux.* xi. 11. (1844) 188.—Mexic.
rhytidea, *Benth. Pl. Hartw.* 21.—Mexic.
rigida, *Pohl, ex Benth. Lab. Gen. et Sp.* 85.—Bras.
rostrata, Salzm. ex Benth. l. c. 121 = spicata.
rotundifolia, *Benth. l. c.* 111.—Bras.
rubicunda, *Pohl, ex Benth. l. c.* 118.—Bras.
rubiginosa, *Benth. l. c.* 106.—Bras.
rufula, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 138 = lutescens.
rugosa, *Benth. Lab. Gen. et Sp.* 86.—Bras.
salicina, *J. A. Schmidt, in Mart. Fl. Bras.* viii. 124.—Bras.
salviæfolia, *Pohl, ex Benth. Lab. Gen. et Sp.* 136.—Bras.

HYPTIS :—
Salzmanni, *Benth. Lab. Gen. et Sp.* 138.—Bras.
sauvallis, *A. St. Hil. ex Benth. l. c.* 90.—Bras.
scabra, *Benth. l. c.* 134.—Bras.
scoparia, *Polt. in Ann. Mus. Par.* vii. (1806) 475. t. 31. f. 2.—Am. austr.
scleiginifolia, *Mart. ex Benth. Lab. Gen. et Sp.* 87.—Bras.
Selloi, *Benth. l. c.* 89.—Bras.
sericea, *Benth. l. c.* 79.—Bras.
Sidæritis, *Mart. ex Benth. l. c.* 97.—Bras.
simplex, *A. St. Hil. ex Benth. l. c.* 138.—Bras.
sinuata, *Pohl, ex Benth. l. c.* 108.—Bras.
sordida, Pohl, ex Benth. l. c. 135 = scabra.
spicata, *Polt. in Ann. Mus. Par.* vii. (1806) 474. t. 28. f. 2.—Am. austr.
spicigera, *Lam. Encyc.* iii. 185.—Am. et Afr. trop.
spinulosa, *Benth. Lab. Gen. et Sp.* 129.—Mexic.
spiræacifolia, *Mart. ex Benth. l. c.* 95.—Bras.
stachyoides, Link, Enum. Hort. Berol. ii. 106 = Elsholtzia incisa.
stellulata, *Benth. Lab. Gen. et Sp.* 129.—Mexic.
stricta, *Benth. l. c.* 79.—Bras.
strictissima, Pohl, ex Benth. l. c. 99 = linarioides.
suaveolens, *Polt. in Ann. Mus. Par.* vii. (1806) 472. t. 29. f. 2.—Am. austr.
subrotunda, *Pohl, ex Benth. Lab. Gen. et Sp.* 118.—Bras.
subsessilis, Benth. in DC. Prod. xii. 92 = longipes.
subverticillata, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 197.—Ins. Galap.
sylvularum, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 119.—Bras.
Tafallæ, *Benth. l. c.* 132.—Peruv.
tenuiflora, *Benth. l. c.* 121.—Bras.
tephrodes, *A. Gray, in Proc. Am. Acad.* v. (1861) 164.—Mexic.
tetragona, *Pohl, ex Benth. Lab. Gen. et Sp.* 100.—Bras.
tomentosa, *Polt. in Ann. Mus. Par.* vii. (1806) 469.—Mexic.
trichphala, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 112.—Bras.
trifoliifolia, *Mart. ex Benth. l. c.* 93.—Bras.
Tweediei, *Benth. in DC. Prod.* xii. 110 (Tweedii).—Bras.
uliginosa, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 81.—Bras.
umbrosa, *Salzm. ex Benth. l. c.* 125.—Bras.
uncinata, *Benth. l. c.* 80.—Peruv.
undulata, Schrank, ex Denkschr. Bot. Ges. Regensb. ii. (1822) 51 = fasciculata?
urticoides, *H. B. & K. Nov. Gen. et Sp.* ii. 320.—Am. austr.
velutina, *Pohl, ex Benth. Lab. Gen. et Sp.* 90.—Bras.
vepretorum, *Mart. ex Benth. l. c.* 131.—Bras.
verbenacifolia, *Mart. ex J. A. Schmidt, in Mart. Fl. Bras.* viii. 110.—Bras.
veronicæfolia, *Pohl, ex Benth. Lab. Gen. et Sp.* 100.—Bras.
verticillata, *Jacq. Ic. Rar.* 11. t. 113; *Coll.* i. 101.—Ind. occ.
vesiculosa, *Benth. Lab. Gen. et Sp.* 131.—Am. austr.
vestita, *Benth. l. c.* 114.—Bras.
viejensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 84.—Nicaragua.
villi, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 13.—Venezuela.
villosa, *Pohl, ex Benth. Lab. Gen. et Sp.* 110.—Bras.
violacea, *Pohl, ex Benth. l. c.* 116.—Bras.
virgata, *Benth. l. c.* 77.—Bras.
viscidula, *Benth. in DC. Prod.* xii. 92.—Bras.
vitifolia, *Pohl, ex Benth. Lab. Gen. et Sp.* 138.—Bras.
vulcanica, *Seem. Bot. Voy. Herald*, 138.—Panama.
xanthiocephala, *Mart. ex Benth. Lab. Gen. et Sp.* 113.—Bras.

HYPTISSA, Salisb. Gen. Pl. Fragm. 142 (1866) = Gladiolus, Tourn. (Irid.).
ramea, Salisb. l. c. 143.—Hab.?

HYPUDAEURUS, Reichb. Nom. 37 (1841) = HYPODAEURUS, Hochst. = Anthephora, Schreb. (Gramin.).
cenchroides, Hochst. ex A. Br. in Flora, xxiv. (1841) 876 = A. Hochstetteri.

HYRTANANDRA, Miq. Pl. Jungh. 25 (1851) = **Pousolzia**, Gaudich. (Urtic.).
canulata, Miq. ex Bedd. in Trans. Linn. Soc. xxv. (1865) 234 = P. caudata.
ciliaris, Miq. ex Bedd. l. c.—Ind. or.
gracilis, Miq. Anal. Bot. Ind. iii. 20.—Ind. or.
hirta, Miq. Pl. Jungh. 26 = P. hirta.
javanica, Miq. l. c. 25 = P. pentandra.
lythroides, F. Muell. Fragm. vi. 253 = P. hirta.
macrophylla, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 57.—Ind. or.
pentandra, Miq. Pl. Jungh. 25; et Fl. Ind. Bat. i. ii. 261 = P. pentandra.
Pousolzia, F. Muell. Fragm. v. 194 = P. hirta.
scabra, Miq. ex Bedd. in Trans. Linn. Soc. xxv. (1865) 234 = P. scabra.
ternata, Miq. ex Bedd. l. c. = P. Wightii.
triandra, Miq. Fl. Ind. Bat. i. ii. 262 = P. parvifolia.
Walleriana, Thw. Enum. Pl. Zeyl. 261 = P. pentandra.

HYSSOPIFOLIA, Opiz, Seznam, 58 (1852) = **Lythrum**, Linn.
parvifiora, Opiz, l. c. = L. Hyssopifolia.

HYSSOPUS, [Tourn.] Linn. Syst. ed. I (1735). *LABIATEAE*, Benth. & Hook. f. ii. 1187.
alopecuroides, Fisch. ex Benth. in DC. Prod. xii. 252 = officinalis.
altissimus, Mill. Gard. Dict. ed. VIII. n. 3 = officinalis.
angustifolius, Bieb. Fl. Taur. Cauc. ii. 38 = officinalis.
anisatus, Nutt. Gen. Am. ii. 27 = Lophanthus anisatus.
aristatus, Godr. in Mém. Acad. Stanislas, Sér. III. (1850) 106 = officinalis.
bungeianus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 91 = officinalis.
bracteatus, C. C. Gmel. ex Steud. Nom. ed. I. 296, 423 = Elsholtzia cristata.
canescens, DC. ex Nym. Consp. 557 = officinalis.
cataractarius, Host. Par. ex Benth. Lab. Gen. et Sp. 403 = Lophanthus scrophulariaefolius
caucasicus, Spreng. ex Steud. Nom. ed. I. 423 = officinalis.
cinerascens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 92 = officinalis.
cristatus, Lam. Encyc. iii. 187 = Pogostemon paniculatus.
decumbens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 46 = officinalis.
discolor, Desf. Cat. Hort. Par. ed. III. 97 = Lophanthus anisatus.
Flascheri, Hort. ex Steud. Nom. ed. II. i. 795 = officinalis.
Forniculum, Spreng. Nov. Prov. 24 = Lophanthus anisatus.
hirsutus, Hill, Veg. Syst. xvi. 56 = officinalis ?
lophanthoides, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 110 = Plectranthus Gerardianus ?
Lophanthus, Linn. Sp. Pl. 569 = Lophanthus chinensis.
montanus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 90 = officinalis.
myrtifolius, Desf. Tabl. ed. I. 58 (*myrthifolius*) = officinalis.
nepetoides, Herb. Madr. ex Wall. Cat. n. 1530 = Pogostemon plectranthoides.
nepetoides, Linn. Sp. Pl. 569 = Lophanthus nepetoides.
ocymifolius, Lam. Encyc. iii. 187 = Elsholtzia cristata.
officinalis, *Linn. Sp. Pl.* 569.—Europ.; As. temp.
orientalis, Adam, ex Willd. Enum. Hort. Berol. 599 = officinalis.
polyclados, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 90 = officinalis.
pubescens, Jord. & Fourr. l. c. 91 = officinalis.
recticaulis, Jord. & Fourr. l. c. 90 = officinalis.
resupinatus, Moench, Meth. Suppl. 134 = Lophanthus chinensis.
ruber, Mill. Gard. Dict. ed. VIII. n. 2 = officinalis.
Schischkini, G. Don, ex Loud. Hort. Brit. 233 = officinalis.
scrophulariaefolius, Willd. Sp. Pl. iii. 48 = Lophanthus scrophulariaefolius
septemcrenatus, Ehrenb. ex Sweet, Hort. Brit. ed. II. 402 = sec. ?
septemfidus, Ehrenb. ex Sweet, l. c., nomen (Quid ?).—Aegypt.

HYSTERIA, Reinw. ex Hume, Cat. Gew. Buitenz. 99 (1823), nomen, = Corymbia, Thou. (Orchid.).
ovalifolia, Reinw. l. c. = C. corymbosa.

HYSTERICINA, Steud. Syn. Pl. Gram. 85 (1855) = **Echinopogon**, Beauv. (Gramin.).
alopecuroides, Steud. l. c. = E. ovatus.

HYSTERIONICA, Willd. in Ges. Naturf. Fr. Berl. Mag. I. (1807) 140. *COMPOSITAE*, Benth. & Hook. f. ii. 252.
NEJA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 78 (1831).
gracilis, *Benth. & Hook. f. Gen.* ii. 253.—Reg. Argent.
jasionoides, *Willd. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 140.—Reg. Argent.
linearifolia, *Baker, in Mart. Fl. Bras.* vi. 111. 13.—Bras.
montevidensis, *Baker, l. c.*—Reg. Argent.
nidoreliolides, *Baker, l. c.* 12.—Bras.
pinifolia, *Benth. & Hook. f. Gen.* ii. 253; *Baker, in Mart. l. c.*—Bras.
scaposa, Benth. & Hook. f. l. c. = Leucopsis scaposa.
squarrosa, Baill. Hist. Pl. viii. 38 = Grindelia squarrosa.
subvillosa, *Griseb. in Goett. Abh.* xix. (1874) 170.—Reg. Argent.

HYSTERONICA, Endl. Esch. 251 (1841) = prsec.

HYSTEROPHORUS, Adans. Fam. ii. 128 (1763) = **Parthenium**, Linn. (Compos.).

HYSTRIX, Moench, Meth. 294 (1794) = **Asperella**, Humb. (Gramin.).
patula, Moench, l. c. 295 = A. Hystrix.

I

IACARANDA, Nees, in R. Br. Verm. Schrift. v. 325 (1843) = **Jacaranda**, Juss. (Bignon.).

IALAPA, Crantz, Inst. ii. 365 (1766) = JALAPA, Tourn. = **Mirabilis**, Linn. (Nyctag.).

IANTHA, Hook. Exot. Fl. t. 118 (1825) = **Ionopsis**, Thunb. (Orchid.).
pallidiflora, Hook. l. c. = Ionopsis pallidiflora.

IANTHE, Salisb. *Parad. Lond.* Fl. Fragm. 44 (1866) = **Hypoxis**, Linn.
linearis, Salisb. l. c. = H. stellata.
ovata, Salisb. l. c. = H. ovata.
serrata, Salisb. l. c. = H. serrata.

IARAVAEA, Scop. Introd. 215 (1777) = **Miorlicia**, D. Don (Melastom.).

IASIONE, Moench, Meth. 518 (1794) = **Jasione**, Linn. (Campan.).

IATROPHA, Stokes, Bot. Mat. Med. iv. 447 (1812) = **Jatropha**, Linn. (Euphorb.).

IBATIA, Decne. in DC. Prod. viii. 599, 605 (1844) = **Lachnostoma**, H. B. & K. (Asclep.).
albiflora, Kanst. Fl. Columb. ii. 113. t. 160. f. 1.—N. Granat.
ciliata, Fourn. in Mart. Fl. Bras. vi. iv. 309.—Bras.
diversifolia, Fourn. l. c.—Bras.
fimbriata, Karst. Fl. Columb. ii. 113. t. 160. f. 2.—N. Granat.
lanosa, Fourn. in Mart. Fl. Bras. vi. iv. 309.—Bras.
maritima, Decne. in DC. Prod. viii. 599 = L. maritimum.
maritima, Griseb. Fl. Brit. W. Ind. 744, sphalm. = *muricata*.
mollis, Griseb. Cat. Pl. Cub. 177.—Cuba.
muricata, Griseb. Fl. Brit. W. Ind. 421 = Gonolobus muricatus.
quinquelobata, Fourn. in Mart. Fl. Bras. vi. iv. 308.—Bras.
Sellowna, Fourn. l. c. 309.—Bras.

IBBERTSONIA, Steud. Nom. ed. II. i. 800 (1840) = sec.

IBBETSONIA, Sims, Bot. Mag. t. 1259 (1810) = **Cyclopia**, Vent. (Legumin.).
genistoides, Sims, l. c. = C. genistoides.

IBERIDELLA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 188. *CRUCIFERAE*, Benth. & Hook. f. i. 92.
STENOPSIS, Jaub. & Spach, Illustr. i. t. 3 (1842).
Andersoni, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 177.—Reg. Himal.
cespitosa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 190.—Armenia.
heterophylla, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 95.—Reg. Caucas.
ovalifolia, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 189 = trinervia.
rotundifolia, *Hook. f. Bot. Mag.* t. 5749.—Europ.
sagittata, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 188.—Persia.
tenuis, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 95.—Persia.
trinervia, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 188.—Armenia.

IBERIS, Adans. Fam. ii. 422 (1763) = **Lepidium**, Tourn. (Crucif.).

IBERIS, Dill. ex Linn. Syst. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 92.
ARABIS, Adans. Fam. ii. 422 (1763).
acutiloba, Bertol. Misc. Bot. ii. 12. t. 9. f. 1 = odorata.
affinis, Jord. Adnot. Gr. Jard. Bot. Dijon (1848) 25 = pandurseformis.
amara, *Linn. Sp. Pl.* 649.—Europ.
amara, Pall. Ind. Taur.; ex Bieb. Fl. Taur. Cauc. ii. 103 = taurica.
amoena, Jord. Obs. Pl. Crit. vi. (1847) 46 = umbellata.
apricorum, *Giraudias, in Feuilles Jeun. Natural.* vi. (1876) 103.—Gallia.
arabica, Linn. Cent. Pl. i. 17; Amoen. Acad. iv. 276 = Aethionema cappadocicum.
Arbuscula, Spach, Hist. Veg. Phan. vi. 502 = sempervirens.
arvatica, Jord. Diagn. i. 288 = amara.
attica, Jord. Obs. Pl. Crit. vi. (1847) 42 = Jordani.
aurosica, Chaix, in Vill. Hist. Pl. Dauph. i. 349; iii. 389 = saxa.
aurosica, Coss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 771 = Lagascana.
badensis, Linn. Cent. Pl. i. 17 = Thlaspi montanum.
balansæ, *Jord. Diagn.* i. 274.—Algeria.
bellidifolia, Presl, Delic. Prag. 13 = Pruiti.
Benthamiana, Boiss. & Reut. Pugill. Fl. Nov. (1852) 12 = Bernardiana.
Bernardiana, *Gren. & Godr. Fl. Fr.* i. 138.—Mont. Pyren.
bicolor, *Reichb. Fl. Germ. Excurs.* 659.—Europ.
bicorymbifera, Gren. & Godr. Fl. Fr. i. 141 = pandurseformis.
boppardensis, Jord. Obs. Pl. Crit. vi. (1847) 62 = intermedia.
Bourgæi, Boiss. Diagn. Ser. II. i. 39 = pectinata.
Bourgæi, Jord. Diagn. i. 276 = contracta.
Boutelouæ, Willk. in Bot. Zeit. v. (1847) 225 = pectinata.
brachystyla, Fenzl, Pugill. Pl. Nov. Syr. 14 = Aethionema Iberideum.
Bubani, Deville, in Bull. Soc. Bot. Fr. vi. (1859) 69 = Bernardiana.
Bursa-pastoris, Crantz, Stirp. Austr. ed. I. fasc. i. 21; ed. II. 20 = Capsella Bursa-pastoris.
burzifolia, Berger. ex DC. Syst. ii. 892 = Teesdalia Iberis.
calycina, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 16 (Quid ?).—Java.
campestris, Wallr. Sched. Crit. 341 = Lepidium campestre.
candicans, Rafin. Act. Journ. 146 (Quid ?).—Am. bor.
Candolleana, Jord. Obs. Pl. Crit. vi. (1847) 37 = sana.
cappadocica, Willd. Sp. Pl. iii. 492 (= *Pilotrichum cappadocicum*).—Cappadoc.
camosa, *Willd. l. c.* 486.—Europ.
capsuloides, Pourr, in Mém. Acad. Toul. iii. (1788) 321 = carnosa.
capsuloides, Tenore, Prod. Fl. Nap. p. xxxvii = Tenorii.
capsuloides, Walf. in Jacq. Misc. ii. 89 = Thlaspi capsuloides.

IBERIS:—
cerastophylla, Reut. in Compt.-Rend. Soc. Hallér. (1853–54) 18 = pandurseformis.
chlorsefolia, Sibth. & Sm. Fl. Græc. Prod. ii. 10 = Aethionema chlorsefolium.
ciliata, *All. Fl. Pedem.* 16.—Europ.
ciliata, Willd. Sp. Pl. iii. 455; Bieb. Fl. Taur. Cauc. ii 109 = taurica.
cinerea, *Poir. Encyc. Suppl.* iii. 138.—Hispan.
collina, Jord. Obs. Fl. Crit. vi. (1847) 57 = intermedia.
commutata, Schott & Kotschy, ex Boiss. Fl. Orient. i. 333 = sempervirens.
conferta, *Lag. in Varied. Cienc.* ii. iv. (1806) 213.—Hispan.
Contejeani, Billot, Annot. 95 = intermedia.
contracta, *Pers. Syn.* ii. 186.—Hispan.
confolia, *Smert. Hort. Brit.* ed. I. 32.—Europ. austr.
coronaria, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 359 = umbellata.
corymbosa, Moench, Meth. 269 = umbellata.
crassifolia, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 770 = carnosa.
crenata, Lam. Encyc. iii. 218 = pinnata.
cretica, Linn. Sp. Pl. 649 = linifolia ?
cuneata, Moench, Meth. 269 = sempervirens.
decipiens, Jord. Diagn. i. 289 = amara.
deflexifolia, Jord. Pugill. 18 = Prostii.
delphinensis, Jord. Diagn. i. 286 = Prostii.
dentata, Moench, Meth. Suppl. 88 = gibraltarica.
divaricata, Tausch, in Flora, xiv. (1831) 218 = intermedia.
Durandii, Lorey & Duret, Fl. Côte d'Or, i. 68. t. 1 = Prostii.
elliptica, Salisb. Prod. 367 = sempervirens.
Forestieri, Jord. Cat. Recol. (1849) 21 = amara.
Garreviana, All. Fl. Pedem. i. 250. t. 40. f. 3 & t. 54. f. 2 = sempervirens.
Garreviana, Scop. Delic. Insub. i. 16. t. 7 = saxatilis.
Gastoni, Delacroix, in Bull. Soc. Bot. Fr. ii. (1855) 456 = sempervirens.
gibraltarica, *Linn. Sp. Pl.* 649.—Hispan. ; Marocc.
glauca, Lag. ex Willk. & Lange, Prod. Fl. Hisp. iii. 773 = conferta.
glaucescens, *Boiss. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 248.—Cilicia.
granatensis, Boiss. & Reut Pugill. Pl. Nov. 11 = Tenoreana.
Hegelmaieri, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 770.—Hispan.
hirsuta, Hort. Par. ex Willd. Enum. Hort. Berol. Suppl. 44 = pubescens.
hispidula, *Tenore, Ind. Sem. Hort. Neap.* (1838) 14.—Hab. ?
hortensis, Jord. Obs. Fl. Crit. vi. (1847) 46, 74. t. 1 = umbellata.
humifusa, *Lag. ex Link, Handb.* ii. 290 (Quid ?).—Hispan.
humilis, Presl, Delic. Prag. 19 = semperflorens.
integerrima, Moris, Stirp. Sard. Elench. i. 4 = Prostii.
integrifolia, Ledeb. Fl. Ross. i. 165 = taurica.
intermedia, *Guersenti, in Bull. Soc. Philom.* iii. (1811) 169. t. 21.—Gallia.
Jordani, *Boiss. Diagn.* Ser. II. i. 88.—An. Min.
jucunda, Schott & Kotschy, ex Boiss. Fl. Orient. i. 347 = Aethionema coridifolium.
Lagascana, *DC. Syst.* ii. 490.—Hispan.
Lamottii, Jord. Pugill. Pl. Nov. (1852) 13 = intermedia.
leptophylla, Jord. Diagn. i. 287 = linifolia.
linearifolia, DC. Syst. ii. 496 = Lepidium linifolium.
linifolia, *Linn. Syst.* ed. X. 1129.—Europ. centr.
linifolia, Lorey & Duret, Fl. Côte d'Or, i. 69, in syn. = Prostii.
linifolia, Nym. Consp. 80 = umbellata.
linifolia, Schkuhr, ex Steud. Nom. ed. II. i. 800 = amara.
Lorreyi, J. Gay, ex Lorey & Duret, Fl. Côte d'Or, i. 69, in syn. = Prostii.
lusitanica, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. xi. 73 = amara ?
lusitanica, Jord. Diagn. i. 276 = contracta.
maialis, Jord. Obs. Fl. Crit. vi. (1847) 52 = intermedia.
Martindi, Tineh.—Cf. Bull. Soc. Bot. Fr. xxx. (1883) Bibl. 186 = pandurseformis.
Moltkerii, Balh. Cat. Hort. Taur. (1813) 48 = ciliata.
montisserratis, Tineh. in Bull. Soc. Sc. Phys. & Nat. Tosc. ii. (1874) 258 = seq.
monteilvensis, *Tineh. l. c.* i. (1873–73) 275.—Gallia.

IBERIS:—
nana, *All. Auct. Fl. Pedem.* 15. t. 2. f. 1.—Europ.
nana, Boiss. Voy. Bot. Espagne, 64 = Tenoreana.
neglecta, *Tenore, Cat. Orto Nap.* (1846) 101.—Hab. ?
nudicaulis, Linn. Sp. Pl. 650 = Teesdalia nudicaulis.
numidica, Jord. Diagn. i. 290 = parviflora.
odorata, Boiss. Voy. Bot. Espagne, 55 = pectinata.
odorata, *Linn. Sp. Pl.* 649.—Græcia; Syria.
olympica, *Boiss. Diagn.* Ser. II. i. 37.—As. Min.
olympica, Sibth. ex DC. Syst. ii. 556 = Aethionema chlorsefolium.
oppositifolia, Pers. Syn. ii. 186 = Aethionema oppositifolium.
pandurseformis, *Pourr. in Mém. Acad. Toul.* iii. (1788) 320.—Europ. austr.
parviflora, Lam. Encyc. iii. 221 = Aethionema saxatile.
parviflora, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 262.—Afr. bor. occ.
pectinata, *Boiss. Diagn.* Ser. I. i. 75.—Hispan.
Petiti, Link, Handb. ii. 291, sphalm. = Prostii.
petræa, Jord. Obs. Fl. Crit. vi. (1847) 69 = Tenoreana.
pilosa, Desv. Journ. Bot. iii. (1814) 167 = Tenoreana.
Pinardi, Jord. Fragm. vi. 42 = taurica.
pinnata, *Linn. Cent. Pl.* i. 18; *Amœn. Acad.* iv. 278. —Europ. austr.
polita, Jord. Obs. Fl. Crit. vi. (1847) 51 = Prostii.
procumbens, *Lange, Ind. Sem. Hort. Haun.* (1861) 29; *Pl. Hisp. Hisp.* fasc. i. n. 1. 6.—Hispan.
Prostii, *Sayer-Willem. ex Godr. Fl. Lorr.* ed. I. i. 78.—Gallia.
Pruiti, *Tineo, Pugill. Pl. Sic.* i. 11. n. 9.—Sicilia.
pubescens, *Willd. Enum. Hort. Berol. Suppl.* 43.—Hab. ?
pulchra, Salisb. Prod. 367 = umbellata.
pumilio, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 269 = Thlaspi pumilum.
pyrenaica, Lapeyr. Hist. Abr. Pl. Pyr. 370 = Aethionema saxatile.
radicans, Lag. ex Nym. Consp. 80 = conferta.
Rayneyalli, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 5. —Hispan.
Raynevalii, Torrep. ex Willk. & Lange, Prod. Fl. Hisp. iii. 769 = umbellata.
repens, Lam. Fl. Fr. ii. 674 = Iberidella rotundifolia.
rhodocarpa, Willk. in Bot. Zeit. v. (1847) 235 = Tenoreana.
rotundifolia, Lam. Encyc. iii. 221 = carnosa.
rotundifolia, Linn. Sp. Pl. 649 = Iberidella rotundifolia.
ruderalis, Crantz, Stirp. Austr. ed. I. fasc. i. 21 = Lepidium ruderale.
rupifragilis, Lej Fl. Spa, ii. 58 = amara.
sabauda, Puget, in Bull. Soc. Bot. Fr. xiii. (1866) p. clxxv = amara.
saxatilis, Asso, Syn. Pl. Aragon. 84 = ciliata.
saxatilis, *Linn. Cent. Pl.* ii. 23; *Amœn. Acad.* iv. 321.—Europ. austr.
semperflorens, Burm. f. Prod. Fl. Cap. 17 = Alyssum spinosum.
semperflorens, *Linn. Sp. Pl.* 648.—Europ. austr.
sempervirens, Burm. f. Prod. Fl. Cap. 17 = Clypeola Jonthlaspi.
sempervirens, Lag. Gen. et Sp. Nov. 19 = subveluting.
sempervirens, *Linn. Sp. Pl.* 648.—Europ. austr. ; As. Min.
sempervirens, Pall. Ind. Taur. ; ex Ledeb. Fl. Ross. i. 166 = saxatilis.
serrulata, Duf. ex Nym. Consp. 81 = Lagascana.
serrulata, *Vis. Fl. Dalmat.* iii. 111.—Dalmat.
simplex, DC. Fl. Fr. Suppl. 597 = taurica.
spathulata, Berger. Phytonomat. ; ex DC. Fl. Fr. iv. 716 = carnosa.
spathulata, Lag. ex Willk. & Lange, Prod. Fl. Hisp. iii. 771 = Lagascana.
speciosa, Salisb. Prod. 367 = gibraltarica.
Sprunerl, *Jord. Obs. Pl. Crit.* vi. (1847) 48.—Græcia.
stricta, Jord. l. c. 50 = linifolia.
stylosa, Tenore, Prod. Fl. Nap. p. xxxvii = Nocæa stylosa.
subvelutina, *DC. Syst.* ii. 497.—Hispan.
taurica, *DC. l. c.* 499.—As. Min. ; Reg. Caucas.
Tenoreana, *DC. l. c.* 494. (Cf. Prostii).—Italia.
Tenorii, *Presl, Delic. Prag.* 14.—Italia.
Tenorii, Spreng. Syst. ii. 864 = Tenoreana.
tenuifolia, Presl, Delic. Prag. 14 = linifolia.
Timorayi, Jord. Obs. Pl. Crit. vi. (1847) 54 = intermedia.

IBERIS:—
umbellata, Georgi, Beschr. Russ. Reich. iv 1124 = taurica.
umbellata, *Linn. Sp. Pl.* 649.—Europ. austr.
umbellata, Webb, It. Hisp. 77; Boiss. Voy. Bot. Espagne, 54 = contracta.
vernicularis, Willd. Sp. Pl. iii. 454 = saxatilis.
Villarsii, Jord. Gr. Recolt. (1849) 22 = Prostii.
violacea, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 85.—Hab. ?
Violetti, Soyer-Willem. ex Godr. Fl. Lorr. ed. I. i. 73 = Prostii.
virginica, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii 13 = Lepidium virginicum.
Welwitschii, Boiss. Diagn. Ser. II. i. 39 = contracta.
Zanardinii, *Vis. Fl. Dalmat. Suppl.* 119.—Dalmat.

IBETTSONIA, Stend. Nom. ed. II. i. 459, in syn (1840) = IBBETSONIA, Sims = **Cyclopia**, Vent. (Legumin.).

IBIDIUM, Salisb. in Trans. Hort. Soc. i. (1812) 291 = **Spiranthes**, Rich. (Orchid.).
crystallisperum, Salisb. l. c. 292 = S. orchioides.
elatum, Salisb. l. c. 291 = S. elata.
speciosum, Salisb. l. c. = S. colorans.
spirale, Salisb. l. c. = S. autumnalis.

IBINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 = **Sauropus** ?, Blume (Euphorb.).

IBISCUS, Linn. Syst. ed. I (1735) = **Hibiscus**, Linn (Malvac.).

ICACINA, A. Juss. Desc. Nouv. Gen. (1821); et in Mém. Soc. Nat. Hist. Par. i. (1823) 174. t. 9. *OLACINEÆ*, Benth. & Hook. f. i. 852.
dubia, Macfad. Fl. Jamaic. i. 122 = Mappia racemosa.
grandifolia, *Mers, Contrib.* i. 56. t. 4.—Madag.
macrocarpa, *Oliver, Fl. Trop. Afr.* i. 357.—Afr. trop.
Mannii, *Oliver, l. c.*—Afr. trop.
mauritiana, Miers, Contrib. i. 56. t. 4 = Apodytes mauritiana.
senegalensis, *Juss. in Mém. Soc. Hist. Nat. Par.* i. (1823) 174. t. 9.—Afr. trop.
trichantha, *Oliver, Fl. Trop. Afr.* i. 358.—Afr. trop.

ICACO, Adans. Fam. ii. 305 (1763) = **Chrysobalanus**, Linn. (Rosac.).

ICACOREA, Aubl. Pl. Gui. ii. Suppl. i. t. 368 (1775) — **Ardisia**, Sw. (Myrsin.).
balsamifera, Lam. Illustr. i. 186; ex Roem. & Schult. Syst. iv. 508 = A. excelsa.
dentata, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 94 = A. semicrenata.
guianensis, Aubl. Pl. Gui. ii. Suppl. i. t. 368 = A. acuminata.
Humboldtii, A. DC. in DC. Prod. viii. 140 = A tetrandra.
principis, A. DC. l. c. 120 = A. semicrenata.
semicrenata, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 94 = A. semicrenata.
soplanea, Lam. ex Roem. & Schult. Syst. iv. 516 = A. humilis.

ICARANDA, Pers. Syn. ii. 178 (1807) = **Jacaranda**, Juss. (Bignon.).

ICHNANTHUS, Beauv. Agrost. 56. t. 12. f. 1 (1812). *GRAMINEÆ*, Benth. & Hook. f. iii. 1103.
NAVICULARIA, Raddi, Agrost. Bras. 33 (1829).
affinis, *Doell, in Mart. Fl. Bras.* ii. ii. 277.—Bras.
almadensis, Kunth, Enum. i. 185 = Panicum nemorale.
alsinoides, *Munro, ex Hemsl. Biol. Centr. Am. Bot.* iii. 500.—Nicaragua.
bambusiiformis, *Doell, in Mart. Fl. Bras.* ii. ii. 288.—Bras.
brevirachis, *Doell, l. c.* 294.—Bras.
calvescens, *Doell, l. c.* 285.—Bras.
candicans, *Doell, l. c.* 291.—Bras.
Cynotis, *Doell, l. c.* 284.—Bras.
fastigiatus, *Brongn. in Duperr. Voy. Coq. Bot.* 118. t. 18.—Bras.
Hoffmannseggii, *Doell, in Mart. Fl. Bras.* ii. ii. 287. —Am. trop.
inconstans, *Doell, l. c.* 284.—Am. trop.
leptophyllus, *Doell, l. c* 287.—Am. trop.

ICHNANTHUS:—
leiocarpus, *Kunth, Rév. Gram.* ii. t. 168.—Bras.
longiflorus, *Benth. in Journ. Linn. Soc.* xix. (1881) 45. —Bras.
Martianus, *Doell, in Mart. Fl. Bras.* ii. 11. 280.—Am. trop.
mexicanus, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 500.—Mexic.
Minarum, *Doell, l. c.* 294.—Am. trop.
nemorosus, *Doell, l. c.* 299.—Am. trop.
oplismenoides, *Doell, l. c.* 285.—Am. trop.
oplismenoides, *Munro, ex Benth. & Hook. f. Gen.* iii. 1104.—Bras.
pallescs, *Munro, ex Benth. Fl. Hongk.* 414.—Am. trop.
panicoides, *Beauv. Agrost.* 56. t. 12.—Am. austr.
petiolatus, *Doell, in Mart. Fl. Bras.* ii. 11. 278.—Bras.
Planotis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1885) 223.—Am. trop.
Riedelii, *Doell, in Mart. Fl. Bras.* ii. 11. 278 —Bras.
Ruprechtii, *Doell, l. c.* 293.—Bras.

ICHNOCARPUS, R. Br. in Mem. Wern. Soc. i. (1809) 61. *APOCYNACEÆ*, Benth. & Hook. f. ii. 717.
AGANOSMA, G. Don, Gen. Syst. iv. 77 (1837).
QUIRIVELIA, Poir. Encyc. vi. 43 (1804), nomen prius.
SPRINGIA, Muell. Arg. in Heurck, Obs. Pl. Nov. 143 (1871).
affinis, Hook. f. & Thoms. ex Benth. & Hook. f. Gen. ii. 717 = ovatifolius.
Afzeli, *Roem. & Schult. Syst.* iv. 899.—Afr. trop.
bantamensis, *Miq. Fl. Ind. Bat.* ii. 449.—Malaya.
dasycalyx, Miq. l. c. = frutescens.
fragrans, Wall. Cat. n. 1675 = Trachelospermum fragrans.
frutescens, *R. Br. in Mem. Wern. Soc.* i. (1809) 62.— Zeylan.
frutescens, Naves, in Blanco, Fl. Philipp. ed. III. t. 97 = Naves!.
frutescens, Vidal, Sinopa. t. 67. f. E = ovatifolius.
infastus, Buch.-Ham. ex Wall. Cat. n. 1675 = Trachelospermum fragrans.
latifolius, Buch.-Ham. ex Wall. l. c. n. 1682 = ovatifolius.
leptoclictyus, F. Muell. Fragm. vi. 118 = frutescens.
Loureiri, *Spreng. Syst.* i. 635.—Afr. trop.
moluccanus, *Miq. Fl. Ind. Bat.* ii. 448.—Ins. Moluco.
Navesii, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 312. —Ins. Philipp.
ovatifolius, *A. DC. in DC. Prod.* viii. 485.—Ins. Philipp.
ovatifolius, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 121 = Navesii.
publiflorus, *Hook. f. Fl. Brit. Ind.* iii. 670.—Reg. Himal.
radicans, Wall. Cat. n. 1676 = frutescens.

ICHTHYOSMA, Schlecht. in Linnæa, ii. (1827) 671. t. 8; iii. (1828) 194 = **Sarcophyte**, Sparrm. (Baldsoph.).
Wehdemanni, Steud. Nom. ed. II. i. 801 = seq.
Wehdemannii, Schlecht. in Linnæa, ii. (1827) 674 = S. sanguinea.
Weigmanni, Harv. Gen. S. Afr. Pl. 300 = præc.
Wiedemanni, Griff. in Trans. Linn. Soc. xix. (1845) 339 = præc.

ICHTHYOTHERA, [Klotzsch, ex] Bot. Zeit. ii. (1844) 600 = seq.

ICHTHYOTHERE, Mart. in Buchn. Rep. Pharm. xxxv. (1830) 195. *COMPOSITÆ*, Benth. & Hook. f. ii. 346.
LATREILLEA, DC. Prod. v. 504 (1836).
TORRENTIA, Vell. Fl. Flum. vii. t. 149 (1827).
agrestis, *Baker, in Mart. Fl. Bras.* vi. 111. 157.—Bras.
cearensis, Gardn. in Field. & Gardn. Sert. Fl. sub t. 9 = l. Cunabi.
Cunabi, *Mart. in Buchn. Rep. Pharm.* xxxv. (1830) 198.—Bras.
curvifolia, Moric. Pl. Nouv. Am. 150. t. 89 = l. Cunabi.
hirsuta, *Gardn. in Field. & Gardn. Sert. Pl.* sub t. 9. —Bras.

ICHTHYOTHERE:—
integrifolia, *Baker, in Mart. Fl. Bras.* vi. 111. 157.— Bras.
latifolia, *Baker, l. c.* 155.—Bras.
latifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 494 = l. Cunabi.
linearis, *Baker, in Mart. Fl. Bras.* vi. 111. 154.—Bras.
mollis, *Baker, l. c.* 156.—Bras.
peruviana, *Baker, l. c.* 158.—Bras.
rufa, *Gardn. in Field & Gardn. Sert. Pl.* t. 9.—Bras.
suffruticosa, *Gardn. l. c.* t. 8.—Bras.
ternifolia, *Baker, in Mart. Fl. Bras.* vi. 111. 156.— Bras.

ICHTYOTHERE, DC. Prod. v. 504 (1836) = Ichthyothere, Mart. (Compos.).

ICHTYOSMA, Steud. Nom. ed. II. i. 801 (1840) = ICHTHYOSMA, Schlecht. = **Sarcophyte**, Sparrm. (Balanoph.).

ICICA, Aubl. Pl. Gui. 387. tt. 130–135 (1775) = Protium, Burm. f. (Burserac.).
Abilo, Blanco, Fl. Filip. ed. II. 256 = Garuga floribunda.
Acuchini, J. F. Gmel. Syst. 619, sphalm. = P. Aracouchini.
acuminata, Poir. Encyc. Suppl. iii. 186; DC. Prod. ii. 78.—Guian.
altissima, Aubl. Pl. Gui. i. 342. t. 132 = P. altissimum.
amboynensis, Mart. Reise, i. 561; et in Linnæa, v. (1830) Litt. 42 = P. heptaphyllum, guianense.
Aracouchili, March. in Adansonia, viii. (1867) 51 = seq.
Aracouchini, Aubl. Pl. Gui. i. 343 = P. Aracouchini.
bengalensis, Voigt, Hort. Suburb. Calc. 149 = P. serratum.
Carana, H. B. & K. Nov. Gen. et Sp. vii. 84 = P. Carana.
Copal, A. Rich. Ess. Fl. Cub. 386 = P. guianense.
Copal, Schlecht. & Cham. in Linnæa, v. (1830) 601 = P. Copal.
crassifolia, Engl. in DC. Monog. Phan. iv. 81 = P. crassifolium.
cuspidata, H. B. & K. Nov. Gen. et Sp. vii. 84 = P. altissimum.
decandra, Aubl. Pl. Gui. i. 346 = P. decandrum.
decandra, Sagot, ex Engl. in DC. Monog. Phan. iv. 78 = P. Sabanum.
dentata, DC. Prod. ii. 78 = Clausena Willdenowii.
emarginata, Aubl. Pl. Gui. i. 345. t. 134 = P. decandrum.
Goudotiana, Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 273 = Crepidospermum Goudotianum.
guianensis, Aubl. Pl. Gui. i. 340. t. 131 = P. guianense.
guianensis, Triana & Planch. in Ann. Sc. Nat. Sér. V. xiv. (1872) 297, partim = P. heptaphyllum.
Hedwigia, A. Rich. Ess. Fl. Cub. 386.—Cuba.
heptaphylla, Aubl. Pl. Gui. i. 337. t. 130 = P. heptaphyllum.
heptaphylla, Griseb. Fl. Brit. W. Ind. 178 = P. guianense.
heterophylla, DC. Prod. ii. 77 = P. Arcouchini.
Hostmanni, Miq. Stirp. Surin. Sel. 66 = P. Hostmanni.
falcifolia, DC. Prod. ii. 77 = P. falcifolium.
indica, Wight & Arn. Prod. 177 = P. serratum.
inæqualis, Triana & Planch. in Ann. Sc. Nat. Sér. V. xiv. (1872) 299 = P. insigne.
leptophlœos, Mart. ex Engl. in DC. Monog. Phan. iv. 44 = Bursera leptophlœos.
leptostachya, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 473.—Mexic.
macrophylla, H. B. & K. Nov. Gen. et Sp. vii. 83 = P. macrophyllum.
martiana, Cuatr. ex Engl. in DC. Monog. Phan. iv. 70 = P. brasilienne.
obovata, Engl. in DC. l. c. 58, in syn. = P. Copal.
parviflora, Spruce, ex Engl. l. c. 74 = P. Aracouchini.
pentandra, Aubl. Pl. Gui. t. 135 = P. decandrum.
polybotrya, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 473 = P. polybotryum.
pubescens, Benth. in Hook. Kew Journ. iv. (1852) 16 = P. pubescens.
rufa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106. —Guian.
Salzmanni, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 473 = P. heptaphyllum.

ICICA:—
Schomburgkii, Klotzsch, in Schomb. Faun. et Fl. Gui. 1138, nomen.—Guian.
Seilowii, Engl. in DC. Monog. Phan. iv. 69, in syn. = Protium loleatha.
serrata, DC. Prod. ii. 77.—Mexic.
simplicifolia, Mart. ex Engl. in DC. Monog. Phan. iv. 68 = P. unifoliolatum.
Spruceana, Benth. in Hook. Kew Journ. iv. (1852) 16 = P. Spruceanum.
surinamensis, Miq. Stirp. Surin. Sel. 65 = P. heptaphyllum.
Tacamahaca, H. B. & K. Nov. Gen. et Sp. vii. 83 = P. heptaphyllum.
timorienis, DC. Prod. ii. 78.—Ins. Timor.
trianensis, March. ex Engl. in DC. Monog. Phan. iv. 74 = P. Arecouchini.
unifoliolata, Spruce, ex Engl. l. c. 68 = P. unifoliolatum.
viridiflora, Lam. Encyc. iii. 235 = P guianense.

ICICOPSIS, Engl. in Mart. Fl. Bras. xii. ii. 255 (1874) = Protium, Burm. f. (Burserac.).
brasiliensis, Engl. l. c. 257 = P. Blanchetti.
caudata, Engl. l. c. = P. Sagotianum.
ferruginea, Engl. l. c. 256 = P. ferrugineum.
insignis, Engl. l. c. 256 = P. insigne.
reticulata, Engl. l. c. 255 = P. reticulatum.
subserrata, Engl. l. c. 259 = P. subserratum.
tenuifolia, Engl. l. c. 255 = P. tenuifolium.

ICMA, Phil. in Anal. Univ. Chil. (1872) 740; Desc. Nuev. Pl. (1872) 8. *COMPOSITÆ*, Benth. & Hook. f. ii. 1227.
involucrata, Phil. l. c. 741; 88.—Chili.

ICOSANDRA, Phil. in Linnæa, xxix. (1857) 39. *LAURINEÆ*, Benth. & Hook. f. iii. 151.
rufescens, *Phil. l. c.* 40.—Chili.

ICOSINIA, Rafin. Sylva Tellur. 75 (1838) = **Sterculia**, Linn.
paniculata, Rafin. l. c. = S. grandiflora.

ICOSTEGIA, Rafin. Sylva Tellur. 168 (1838) = **Clusia**, Linn. (Guttif.).
flava, Rafin. l. c. = C. flava.

ICOTORUS, Rafin. in Ser. Bull. Bot. i. (1830) 216 = **Nesilia**, D. Don (Rosac.).
montanus, Rafin. l. c. = N. Torreyi.

ICTHYOCTONAM, Bolv. ex Baill. in Bull. Soc. Linn. Par. i. (1884) 440 = **Lonohoorpus**, H. B. & K. (Legumin.).
madagascariensis, Bolv. ex Baill. l. c. = L. ichthyoctonus.

ICTHYOTHERE, Baker, in Mart. Fl. Bras. vi. 111. 158 (1884) = **Ichthyothere**, Mart. (Compos.).

ICTINUS, Cass. in Bull. Soc. Philom. (1818) 149; et in Dict. Sc. Nat. xix. 284; xxii. 559 (1821) = **Gortaria**, Linn. (Compos.).
gorteriodes, Steud. Nom. ed. II. i. 801 = G. diffusa.
pilosellaides, Cass. in Bull. Soc. Philom. (1818) 143 = G. diffusa.

ICTODES, Bigel. Med. Bot. ii. 41. t. 24 (1818) = **Symplocarpus**, Salisb. (Aroid.).
fœtidus, Bigel. l. c. = S. fœtidus.

IDALIA, Rafin. Fl. Tellur. iv. 76 (1836) = **Convolvulus**, Tourn.
albiflora, Rafin. l. c. 77 = C. elongatus.

IDELERIA, Kunth, Enum. Pl. ii. 210 (1837) = **Tetraria**, Beauv. (Cyperac.).
capensis, Kunth, l. c. 211.—Afr. austr.
Nemei, Kunth, l. c. (= *Hymenochæ Kraussii*).—Afr. austr.

IDESIA, Maxim. in Bull. Acad. Pétersb. x. (1866) 455. *BIXINEÆ*, Benth. & Hook. f. i. 272.
POLYCARPA, Lindan, ex Carr. in Rev. Hortic. (1866) 290.
polycarpa, *Maxim. in Bull. Acad. Pétersb.* x. (1866) 455.—Japon.

IDESIA, Soop. Introd. 199 (1777) = **Ropourea**, Aubl. (Verbenac.!).

IDIANTHES, Desv. Fl. Anj. 199 (1827) = **Phaseolus**, Can. (Compos.).
pulchra, Desv. l. c. = P. lampsanoides.

IDICIUM, Neck. Elem. i. 38 (1790) = **Gerbera**, Gronov. (Compos.).

IDOTHEA, Kunth, Enum Pl. iv. 841 (1848) = **Drimia**, Jacq. (Liliac.).
ciliaris, Kunth, l. c. 848 = D. ciliaris.
drimioides, Kunth, l. c. 845 = Urginea pusilla.
elata, Kunth, l. c. 848 = D. elata.
humilis, Kunth, l. c. 844 = D. pusilla.
Ludwigii, Kunth, l. c. 681 = Scilla Ludwigii.
marginata, Kunth, l. c. 846 = Urginea marginata.
media, Kunth, l. c. 842 = D. media.
physodes, Kunth, l. c. 845 = Urginea physodes.
purpurascens, Kunth, l. c. 842 = D. purpurascens.
pusilla, Kunth, l. c. 844 = D. pusilla.
villosa, Kunth, l. c. 848 = D. villosa.

IDOTHEARIA, Presl, Bot. Bemerk. 114 (1844) = **Drimia**, Jacq. (Liliac.).
ciliaris, *elata*, *media*, *purpurascens*, *pusilla*, *villosa*, Presl, l. c. = omnes *Drimiae* specierum homonymae.
humilis, Presl, l. c. = D. pusilla.

IDRIA, Kellogg, in Proc. Calif. Acad. ii. (1863) 84 = **Fouquiera**, H. B. & K. (Tamarisc.).
columnaris, Kellogg, l. c. = F. spinosa !

IEBINE, Rafin. Fl. Tellur iv. 39 (1836) = **Liparis**, Rich. (Orchid.).
nervosa, Rafin. l. c. = L. nervosa.

IFDREGEA, Steud. Nom. ed. II. i. 801 (1840) = **Puocedanum**, Tourn. (Umbellif.).
capensis, Steud. l. c. = P. capense.
collina, Steud. l. c. = P. collinum.
montana, Steud. l. c. = P. Ecklonianum.
virgata, Steud. l. c. = P. capense.

IFLOGA, Cass. in Bull. Soc. Philom. (1819) 142; et in Dict. Sc. Nat. xxiii. 18 (1823). *COMPOSITAE*, Benth. & Hook. f. i. 899.
TRICHOGYNE, Less. in Linnaea, vi. (1831) 231.
canifolova, C. B. Clarke, Comp. Ind. 99 = Fontanesii.
Fontanesii, *Cass. in Dict. Sc. Nat.* xxiii. 14.—Afr. bor.; Ind. or.
obovata, *C. Bolle, in Bonplandia*, vii. (1859) 297.—Ins. Canar.
paroychioides, *Fenzl, in Flora*, xxii. (1849) 748.—Afr. austr.
polycnemoides, *Fenzl, l. c.*—Afr. austr.
Sibthorpii, Nym. Syll. 15 = Fontanesii.
spicata, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 310 = Fontanesii.
vertucillata, Fenzl, in Flora, xxii. (1849) 748 = polycnemoides.

IFUON, Rafin. Fl. Tellur. iv. 12 (1836) = **Ornithogalum**, Tourn. (Liliac.).
nutans, Rafin. l. c. = O. nutans.

IGNATIA, Linn. f. Suppl. 20 (1781) = **Strychnos**, Linn. (Loganiac.).
amara, Linn. f. l. c. 149 = S. Ignatii.

IGNATIANA, Lour. Fl. Cochinch. 126 (1790) = **Strychnos**, Linn. (Loganiac.).
philippica, Lour. l. c. = S. Ignatii.

IGUANARA, Relchb. Nom. 58 (1841) = seq.

IGUANURA, Blume, in *Bull. Néerl.* 66 (1838); et Rumphia, ii. 105. t. 117 (1843). *PALMAE*, Benth. & Hook. f. iii. 907.
SLACKIA, Griff. Notul. iii. 162 (1851).
bonensis, *Scheff. in Ann. Jard. Buitens.* i. (1876) 161.—Borneo.
geonomaeformis, *Mart. Hist. Nat. Palm.* iii. 229. t. 178.—Malaya.
leiocarpa, *Blume, in Bull. Néerl.* (1838) 66.—Sumatra.
remotiflora, *H. Wendl. in Bot. Zeit.* xvi. (1858) 63.—Borneo.
Wallichiana, [*Hook. f. in*] *Benth. & Hook. f. Gen.* iii. 908.—Malaya.

ILDEFONSIA, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 184. *SCROPHULARINEAE*, Benth. & Hook. f. iii. 953.
bibracteata, *Gardn. l. c.*—Bras.

ILDEFONSIA, Mart. ex Steud. Nom. ed. II. i. 802 (1840) = **Urtica**, Tourn.

ILEOCARPUS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40 = Elaphantia, Lour. (Menispermn.).
Schimperi, Miers, l. c. = S. hemandifolia.

ILEX, [Tourn.] Linn. Syst. ed. I (1735). *ILICINEAE*, Benth. & Hook. f. i. 856.
AGERIA, Rafin. Sylva Tellur. 46 (1838).
AQUIFOLIUM, [Tourn.] Adans. Fam. ii. 166 (1763).
ARINEMIA, Rafin. Sylva Tellur. 49 (1838).
BRAXYLIS, Rafin. l. c. 51 (1838).
CHOMELIA, Vell. Fl. Flum. i. t. 106 (1827).
ENNEPTA, Rafin. Sylva Tellur. 52 (1838).
HEXOTRIA, Rafin. in Am. Monthly Mag. iv. (1818) 265.
LABATIA, Soop. Introd. 197 (1777).
LEUCODERMIS, Planch. ex Benth. & Hook. f. Gen i. 357 (1862).
MACOUCOUA, Aubl. Pl. Gui. i. 88. t. 34 (1775).
OTHERA, Rumph. Nov. Gen. 56 (1743).
PAUTOLA, Ruiz & Pav. Prod. 13 (1794); Fl. Peruv. i. 54. t. 84 (1798).
PILEOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 276.
PRINODIA, Griseb. in Goett. Abh. vii. (1857) 224, in obs.
PRUNUS, Gronov. ex Linn. Coroll. Gen. 6 (1737).
PRSUDEHRETIA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 607.
SYNSTIMA, Rafin. Sylva Tellur. 46 (1838).
VENTERLLA, Moench, Meth. 74 (1794).
acrodonta, *Reiss. in Mart. Fl. Bras.* xi. i. 51.—Bras.
acuminata, Willd. Sp. Pl. i. 711 = I. Macoucona.
acutangula, Nees, in Flora, iv. (1821) 301 = Maytenus acuminata.
acutangula, Nees, l. c. 329 = Elaeodendron quadrangulatum.
aristnella, Lam. Encyc. iii. 147; et in Journ. Hist. Nat. Par. i. (1792) 416 = decidua.
affinis, *Gardn. in Hook. Ic. Pl.* i. 465.—Bras.
amara, Bonpl. ex Miers, in Ann. & Mag. Nat. Hist. Ser. III. viii. (1861) 395 = Symplocos lanceolata.
ambigua, *Chapm. Fl. S. U. St.* 269.—Am. bor.
ambigua, Ell. Sketch. ii. 705 = decidua.
ambigua, Torr. Fl. N. York, i. = montícola, *A. Gray*.
Amelanchier, *M. A. Curt. ex Chapm. Fl. S. U. St.* 270.—Am. bor.
americana, Lam. in Journ. Hist. Nat. Par. i. (1792) 416 = quercifolia ?
Andersonii, *Pasq. Cat. Ort. Bot. Nap.* 55.—Hab. ?
angustifolia, Salisb. Prod. 70 = I. Cassine.
angustifolia, Willd. Enum. Hort. Berol. i. 272 = I. Dahoon.
angustissima, *Reiss. in Mart. Fl. Bras.* xi. i. 74.—Bras.
anomala, Hook. & Arn. Bot. Beech. Voy. 111 = Byronia sandwicensis.
Apollinis, *Reiss. in Mart. Fl. Bras.* xi. i. 69.—Bras.
Aquifolium, *Linn. Sp. Pl.* 125.—Europ.; As. occ.
Aquifolium, Lour. Fl. Cochinch. 91 = Loureirei.
Aquifolium, Marsh. Arbust. Am. 68 = quercifolia.
arshinicosa, *Reiss. in Mart. Fl. Bras.* xi. i. 58.—Bras.
arguidens, Miq. in Verol. Med. Akad. Wetensch. II. ii. (1862) 84 = serrata.
asiatica, *Linn. Sp. Pl.* 125 (sp. dub.).—Ind. or.
asperula, *Mart. ex Reiss. in Mart. Fl. Bras.* xi. i. 48.—Bras.
aspuella, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 329.—China.
balearica, Desf. Hist. Arb. ii. 262 = I. Aquifolium.
Betschiana, *Goepp. Del. Sem. Hort. Vratisl.* (1852).—*Cf. Linnaea*, xxvi. (1853-55) 746.—Mexic.
Bonplandiana, Muenter, in Mitt. Naturw. Ver. Neu-Vorp. xiv. (1883) 158 = paraguensis.
brevicuspis, *Reiss. in Mart. Fl. Bras.* xi. i. 56.—Bras.
brevifolia, Muenter, in Mitt. Naturf. Ver. Neu-Vorp. xiv. (1883) 198 = ovalifolia, *G. F. W. Mey.*
brextaefolia, *Houtt. ex Goepp. Del. Sem. Hort. Vratisl.* (1852).—*Cf. Linnaea*, xxvi. (1853-55) 746.—Hab. ?
ILEX :—
Buergeri, *Miq. in Verh. Med. Akad. Wetensch.* II. ii. (1862) 84.—Japon.
bumeloides, *H. B. & K. Nov. Gen. et Sp.* vii. 71.—Ecuador.
buxifolia, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 132.—Bras.
buxifolia, Hance, in Journ. Bot. xiv. (1876) 364 = Hanceana.
canadensis, Hort. ex Decne. in Fl. des Serres, Sér. I. ix. (1853-54) 184 = I. Aquifolium.
canadensis, Marsh. Arbust. Am. 64 = quercifolia.
canadensis, Michx. Fl. Bor. Am. ii. 239 = Nemopanthes canadensis.
canariensis, *Poir. Encyc. Suppl.* iii. 67.—Ins. Canar.
capensis, *Harv. & Sond Fl. Cap.* i. 472.—Afr. austr.
carolinana, *Mill. Gard. Dict.* ed. VIII. n. 3 = I. Cassine.
Cassena, Michx. Fl. Bor Am. ii. 239 = I. Cassine.
Cassine, Linn. Sp. Pl. 125 = I. Cassine, I. Dahoon.
Cassine, *Walt. Fl. Carol.* 241.—Am. bor.
Cassine, Willd. Hort. Berol. i. 81 = I. Dahoon.
cassinoides, Dum.-Cours. Bot. Cult. ed. II. vi. 251 = I. Cassine.
cassinoides, Link, Enum. Hort. Berol. i. 148 = I. Dahoon.
castaneaefolia, Hort. ex Decne. in Fl des Serres, Sér. I. ix. (1853-54) 186 = I. Cassine.
celastroides, Klotzsch, ex Garcke, in Linnaea, xxii. (1849) 58, partim = I. Macoucoua.
cerasifolia, *Reiss. in Mart. Fl. Bras.* xi. i. 55.—Bras.
chamaedryfolia, *Reiss. l. c.* 78.—Bras.
chinensis, *Sims, Bot. Mag.* t. 2042.—China.
chrysocarpa, *Wender. Sem. Hort. Marb* (1825); *ex Steud. Nom.* ed. I. i. 402.—Hab. ?
cineres, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 327.—Hongkong.
cognata, *Reiss. in Mart. Fl. Bras.* xi. i. 68.—Bras.
condensata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 277.—Mexic.
congesta, *Reiss. in Mart. Fl. Bras.* xi. i. 56.—Guiana.
conocarpa, *Reiss. l. c.* 65.—Bras.
coriacea, Chapm. Fl. S. U. St. 270 = lucida.
cornuta, *Lindl. & Paxt. Flow. Gard.* i. (1850-51) 43.—China.
coronaria, *Reiss. in Mart. Fl. Bras.* xi. i. 51.—Bras.
costata, Blume, ex Maxim. in Mém. Acad. Pétersb. Sér. VII. xxiv. n. 11. (1881) 51 = macropoda.
crassifolia, Ait. ex Steud. Nom. ed. II. i. 802 = I. Aquifolium.
crassifolia, *Hook. Ic Pl.* t. 149.—Peruv.
crassifolia, Meerb. Pl. Select. Ic. t. 4 = I. Perado.
crassifolia, Willd. ex Schult. Mant. iii. 337 = scopulorum.
crenata, *Thunb. Fl. Jap.* 78.—Japon.
crepitans, Boupl. ex Miers, in Ann. & Mag. Nat. Hist. Ser. III. viii. (1861) 396 = Humboldtiana.
crispa, Siebold, Aaedr. & Volk. Toelicht. Ontdekk. Vries, 156, nomen = rugosa.
crocea, Drège, ex Presl, Bot. Bemerk. 37 = Elaeodendron capense.
crocea, Hort. ex Temore, in Atti Teraa Riun. Sc. Ital. (1841) 504 = Elaeodendron ilicifolium.
crocea, Thunb. Prod. Pl. Cap. 32 = Elaeodendron croceum.
cuisbensis, *Reiss. in Mart. Fl. Bras.* xi. i. 71.—Bras.
cumaneusis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 456.—Venezuela.
Cunninghana, *Kolfe, in Journ. Linn. Soc.* xxi. (1884) 306.—Ins. Philipp.
cuneifolia, *Hook. Ic. Pl.* t. 294.—Jamaic.
cuneifolia, Linn. Sp. Pl. 125 = Iodina cuneifolia.
Cunninghami, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1302 = dipyrena.
curitibensis, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vii. (1861) 393 = paraguensis.
cymosa, *Blume, Bijdr.* 1149.—Malaya.
Dahoon, *Walt. Fl. Carol.* 241.—Am. bor.
daphnephylloides, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 72 = Nyssa sessiliflora.
daphnogena, *Reiss. in Mart. Fl. Bras.* xi. i. 46.—Guiana.
daphnoides, *Reiss. l. c.* 58.—Bras.
decidua, *Walt. Fl. Carol.* 241.—Am. bor.
delicatula, Barton, Fl. Virg. 67 = Nemopanthes canadensis.
densifolia, *Miq. Pl. Junghh.* 396.—Java.

ILEX :—
denticulata, *Wall. Cat.* n. 4833.—Ind. or.
Dentoni, Hort. ex Loud. Encyc. Fl. Suppl. ii. 1302 = dipyrena.
dichotoma, Bernh. ex Hochst. in Flora, xxvii. (1844) 805 = Casimopsis capensis.
diminuta, *Reiss. ex Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 21.—Bras.
dioica, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 171. -Ind. occ.
diospyroides, *Reiss. in Mart. Fl. Bras.* xi. 1. 47.—Bras.
dipyrena, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 478.—Reg. Himal.
discolor, *Hemsl. Diagn. Pl. Nov.* 5.—Mexic.
diuretica, *Mart. ex Reiss. in Mart. Fl. Bras* xi. 1. 61.—Bras.
divaricata, *Mart. ex Reiss. l. c.* 57.—Bras.
Dodonæa, Linn. Sp. Pl. 125 = Comocladia ilicifolia.
domestica, Reiss. in Mart. Fl. Bras. xi. 1. 67 = paraguensis.
Doniana, *DC. Prod.* ii. 644.—Reg. Himal.
dumosa, *Reiss. in Mart. Fl. Bras.* xi. 1. 64.—Bras.
ebenacea, *Reiss. l. c.* 41.—Bras.
echinata, Mill. Gard. Dict. ed. VIII. n. 2 = I. Aquifolium.
elliptica, D. Don, Prod. Fl. Nep. 189 = Doniana.
elliptica, H. B. & K. Nov. Gen. et Sp. vii. 70 = I. Paltoria.
emarginata, *Thunb. Fl. Jap.* 78.—Japon.
emarginella, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 606 = Walkeri.
embelioides, *Hook. f. Fl. Brit. Ind.* i. 601.—Reg. Himal.
euryaeformis, *Reiss. in Mart. Fl. Bras.* xi. 1. 61.—Bras.
exandra, *Bello, in Anal. Soc. Hist. Nat. Esp.* x. (1881) 251.—Ins. Porto Rico.
excelsa, Wall. Cat. n. 4821 (errore *exsulca*) = Doniana.
exsulca, Wall. vide præc.
fasciculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 456.—Hab. ?
ferox, Ait. ex Steud. Nom. ed. II. i. 602 = I. Aquifolium.
fertilis, *Reiss. in Mart. Fl. Bras.* xi. 1. 42.—Bras.
flexuosa, E. Mey. in Drège, Zwei Pfl. Docum. 194 = Celastrus pedunculatus.
floribunda, *Reiss. ex Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 26.—Bras.
floridana, Lam. Illustr. i. 856. n. 1781 = I. Cassine.
formosana, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 46.—Ins. Formosa.
Fortunei, *Lindl. in Gard. Chron.* (1857) 868 (Fortuni). —China.
fragilis, *Hook. f. Fl. Brit. Ind.* i. 602.—Reg. Himal.
furcata, *Lindl. ex Goepp. in Regel, Gartenfl.* iii. (1854) 292.—Hab. ?
Gale, *Treana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 377.—N. Granat.
Gardneriana, *Wight, Ic.* t. 1917.—Ind. or.
gaultheriæfolia, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 209 = theæfolia.
geniculata, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 50.—Japon.
gigantea, Pursh, ex Miers, in Ann. & Mag. Nat. Hist. Ser. III. viii. (1861) 394 = fertilis.
glabra, *A. Gray, Man. Bot. N. U. St.* ed. II. 264.—Am. bor.
Glaziovii, *Warm. in Kjoeb. Vidensk. Meddel.* (1880) 369.—Bras.
Godajam, *Colebr. ex Wall. Cat.* n. 4829.—Reg. Himal.
Gongonha, Mart. Reise, ed. Angl. ii. 100 = Villaresia mucronata.
gracilliflora, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 328.—Hongkong.
grandis, Hort. Hannov. ex Decne. in Fl. des Serres, Sér. I. ix. (1853-54) 186 = Betscheriana.
grandis, *Reiss. in Mart. Fl. Bras.* xi. 1. 47.—Bras.
Griffithii, *Hook. f. Fl. Brit. Ind.* i. 601.—Ind. or.; Malaya.
Griseb achii, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 27.—Ind. occ.
Hanceana, *Maxim. l. c.* 53.—Hongkong.
heterophylla, G. Don, Gen. Syst. ii. 17 = I. Aquifolium.
hippocrateoides, *H. B. & K. Nov. Gen. et Sp.* vii. 71. —Am. austr.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. 11. 594.—Java.

ILEX :—
Humboldtiana, *Bonpl. ex Miers, in Ann. & Mag. Nat. Hist.* Ser. III. viii. (1861) 395.—Bras.
humirioides, *Reiss. in Mart. Fl. Bras.* xi. 1. 52.—Bras.
insignis, *Hook. f. Fl. Brit. Ind.* i. 599.—Reg. Himal.
integerrima, *Reiss. in Mart. Fl. Bras.* xi. 1. 45.—Bras.
integra, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 105 = Symplocos japonica.
Integra, *Thunb. Fl. Jap.* 77.—Japon.
intricata, *Hook. f. Fl. Brit. Ind.* i. 602.—Reg. Himal.
inundata, *Poepp. ex Reiss. in Mart. Fl. Bras.* xi. 1. 43. —Bras.
japonica, Thunb. Fl. Jap. 79 = Berberis nepalensis.
laevigata, *A. Gray, Man. Bot. N. U. St.* 307.—Am. bor.
lagunensis, *Warm. in Kjoeb. Vidensk. Meddel.* (1880) 368.—Bras.
lanceolata, *Champ. Fl. S. U. St.* 270.—Am. bor.
lanceolata, *Griseb. Cat. Pl. Cub.* 56.—Cuba.
lanceolata, Klotzsch, in Schomb. Faun. et Fl. Gui. 1188 = I. Martiniana.
latifolia, *Thunb. Fl. Jap.* 70.—Japon.
Laureola, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 377.—N. Granat.
laurifolia, Nutt. in Am. Journ. Sc. v. (1822) 289 = I. Dahoon.
laurina, *H. B. & K. Nov. Gen. et Sp.* vii. 71.—Am. austr.
laurina, Klotzsch, in Schomb. Faun. et Fl. Gui. 1185 = Gaphonogenes.
laxiflora, Lam. Encyc. ill. 147 = quercifolia.
Lepta, Spreng. Syst. i. 496 = Evodia triphylla.
leptacantha, *Lindl. & Paxt. Flow. Gard.* iii. (1852-53) 73.—China.
leucantha, Hassk. Hort. Bog. Desc. 354 = Celastrus articulatus.
ligustrifolia, G. Don, Gen. Syst. ii. 19 = I. Dahoon.
ligustrina, Ell. Sketch, ii. 208 = I. Dahoon.
ligustrina, Jacq. Coll. iv. 105 = I. Cassine.
lineata, E. Mey. in Drège, Zwei Pfl. Docum. 194 = Celastrus acuminatus.
Lobbiana, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 309. —Ins. Philipp.
longifolia, Nees, in Flora, iv. (1821) 301 = Pachystroma ilicifolium.
loranthoides, *Mart. ex Reiss. in Mart. Fl. Bras.* xi. 1. 47.—Bras.
Loureirei, *Steud. Nom.* ed. II. i. 602.—China.
lucida, E. Mey. ex Presl, Bot. Bemerk. 37 = Celastrus acuminatus.
lucida, *Torr. & Gray, ex S. Wats. Bibliog. Ind. N. Am. Bot.* i. 159.—Am. bor.
Lucumæfolia, *Reiss. ex Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 28.—Bras.
Lundii, *Warm. in Kjoeb. Vidensk. Meddel.* (1880) 369.—Bras.
lutescens, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 309. —Ins. Philipp.
Macrocosa, *Pers. Syn.* i. 152.—Guian.
macrophylla, Blume, Bijdr. 1150 = latifolia.
macrophylla, *Wall. Cat.* n. 4831.—Malaya.
macropoda, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 105. —Japon.
madagascariensis, *Lam. Encyc.* ill. 148.—Madag.
maderensis, Lam. l. c. 146 = I. Perado.
maderensis, Willd. Enum. Hort. Berol. Suppl. 8 = I. Aquifolium.
Malagayi, *Hook. f. Fl. Brit. Ind.* i. 605.—Ins. Penang.
malabarica, *Bedd. Fl. Sylv.* i. t. 148.—Ind. or.
Martiana, Walp. Rep. i. 840 = saq.
Martiniana, *Lamb. Pin.* ii. *App.* 8.—Guian.
Mate, A. St. Hil. Fl. Ken. i. p. xli = paraguensis.
medica, *Reiss. in Mart. Fl. Bras.* xi. 1. 62.—Bras.
memecylifolia, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 328.—Hongkong.
Mertensii, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 42.—Ins. Loochoo.
mexicana, *Black, ex Hemsl. Biol. Centr. Am. Bot.* i. 167.—Mexic.
mexicana, *Hort. Versch. ex Decne. in Fl. des Serres*, Sér. I. ix. (1853-54) 186.—Mexic.
micrantha, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 376.—N. Granat.
microcarpa, Lindl. & Paxt. Flow. Gard. i. (1850-51) 48 = rotunda.
microcarpa, Lindl. & Paxt. l. c. = rotunda.

ILEX :—
micrococca, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 39.—Japon.
microlonta, *Reiss. in Mart. Fl. Bras.* xi. 1. 41.—Bras.
microphylla, *Hook. Ic. Pl.* t. 189.—Persv.
microphylla, Spreng. ex DC. Prod. ii. 12 = Myginda microphylla.
minutiflora, *A. Rich. Ess. Fl. Cub.* 352.—Cuba.
mollis, *A. Gray, Man. Bot. N. U. St.* 306.—Am. bor.
montana, *Griseb. in Mem. Am. Acad.* N. S. viii (1861) 171.—Ind. occ.
monticola, *A. Gray, Man. Bot. N. U. St.* 306.—Am. bor.
monticola, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 111.—Malaya.
myriadenia, *Hance, in Journ. Bot.* xxi. (1883) 296.—China.
myricoides, *H. B. & K. Nov. Gen. et Sp.* vii. 71.—N. Granat.
Myrsinites, Pursh, Fl. Am. Sept. i. 119 = Pachystima Myrsinites.
myrtifolia, *Lam. Illustr.* i. 856. n. 1782.—Ind. occ.
myrtifolia, Walt. Fl. Carol. 241 = I. Dahoon.
neocaledonica, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 23.—N. Caled.
nepalensis, Spreng. Syst. iv. Cur. Post. 48 = excelsa.
nervosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 377.—N. Granat.
nigro-punctata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. viii. (1861) 397.—Bras.
nilagirica, Miq. ex Hook. f. Fl. Brit. Ind. i. 600 = denticulata.
nitida, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 27.—Ind. occ.
nummularia, *Franch. & Sav. Enum. Pl. Jap.* ii. 311. —Japon.
nummularia, *Reiss. in Mart. Fl. Bras.* xi. 1. 49.—Bras.
obcordata, *Sw. Fl. Ind. Occ.* i. 388.—Jamaic.
obtusata, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 378.—N. Granat.
occidentalis, *Macfad. Fl. Jamaic.* i. 204.—Ind. occ.
odorata, *Buch-Ham. in D. Don, Prod. Fl. Nep.* 189. —Reg. Himal.
Oldhami, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 105.—Japon.
opaca, [Soland. in] Ait. Hort. Kew. ed. I. i. 169 = quercifolia.
orbiculoris, Willd. ex Schult. Mant. iii. 338 = rupicola.
Orixa, Spreng. Syst. i. 496 = Celastrus Orixa.
Others, *Spreng. l. c.*—Japon.
ovalifolia, *Bonpl. ex Miers, in Ann. & Mag. Nat. Hist.* Ser. III. viii. (1861) 394.—Bras.
ovalifolia, *C. F. W. Mey. Prim. Esseq.* 91.—Guian.
ovata, *Goepp. Del. Sem. Hort. Vratisl.* (1852).—*Cf. Lwnnæa*, xxvi. (1853-55) 745.—Hab. ?
oxyphylla, *Miq. in Journ. Bot. Neerl.* i. (1861) 194.—China.
pachypoda, *Reiss. in Mart. Fl. Bras.* xi. 1. 71.—Bras.
Paltoria, *Pers. Syn.* i. 152.—Am. trop.
paltoriodes, *Reiss. in Mart. Fl. Bras.* xi. 1. 60.—Bras.
paniculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 456.—Guian.
paraguayensis, Hook. Bot. Mag. t. 3992 = Elaeodendron quadrangulatum.
paraguensis, *A. St. Hil. in Mém. Mus. Par.* ix. (1822) 351 (paraguariensis); *D. Don, in Lamb. Pin.* ii. *App.* 49. t. 11.—Paraguay.
parvifiora, *Benth. in Hook. Kew Journ.* iv. (1852) 11. —Bras.
pedunculatis, *F. Muell. Fragm.* vii. 105.—Austral.
peduncularis, *Reiss. in Mart. Fl. Bras.* xi. 1. 50.—Bras.
pedunculosa, *Miq. in Versl. Med. Akad. Wetensch.* II. ii. (1868) 88.—Japon.
Perado, Link, in Buch, Beschr. Canar. Ins. 127 = canariensis.
Perado, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 169.—Ins. Madera.
Perado, Webb & Berth. Phyt. Canar. ii. 126 = platyphylla.
Pernyi, *Franch. Pl. Davidian.* i. (1884) 99.—Mongol.
petiolaris, *Benth. in Hook. Kew Journ.* iv. (1852) 11. —Bras.
phillipinensis, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 308.—Ins. Philipp.

ILEX :—
phillyreæfolia, *Reiss. in Mart. Fl. Bras.* xi. 1. 60.—Bras.
phyllobolos, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 50.—Japon.
phyllyreæfolia, Hort. ex Decne. in Fl. des Serres, Sér. I. ix. (1853–54) 186 = I. Dahoon.
platyphylla, Hort. ex Booth, l. c. 184 = I. Aquifolium.
platyphylla, *Webb & Berth. Phyt. Canar.* ii. 135. t. 68.—Ins. Canar.
pleurostachys, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 456.—Bras.
polyphylla, *Benth. Pl. Hartw.* 167.—N. Granat.
prinoides, [Soland. in] Ait. Hort. Kew. ed. I. i. 169 = decidua.
prionites, Willd. Enum. Hort. Berol. Suppl. 8 = decidua.
psammophila, *Mart. ex Reiss. in Mart. Fl. Bras.* xi. 1. 42.—Bras.
Pseudobuxus, *Reiss. l. c.* 40.—Bras.
Pseudothea, *Reiss. l. c.* 64.—Bras.
Pseudo-vaccinium, *Reiss. ex Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 22.—Bras.
pubescens, *Hook. & Arn. Bot. Beech. Voy.* 176. t. 35.—China.
pubiflora, *Reiss. in Mart. Fl. Bras.* xi. 1. 53.—Bras.
purpurea, *Hassk. Cat. Hort. Bog. Alt.* 230.—Java.
pustulosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 378.—N. Granat.
quadripetala, *Medic. Beobacht.* (1782) 208.—Hab.?
quercifolia, *Meerb. Pl. Select. fc. t.* 5.—Am. bor.
ramulosa, Retin. Sylva Tellur. 44 = I. Cassine.
recurva, *Link, Enum. Hort. Berol.* i. 147.—Hab.?
Reevesiana, Fortune, in Gard. Chron. (1851) 5 = Fortunei.
Regnelliana, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 25.—Bras
religiosa, Barton, Fl. Virg. 66 = I. Cassine.
repanda, *Griseb. in Mém. Am. Acad.* N. S. viii. (1861) 172.—Ind. occ.
retusa, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1097, nomen.—Guian.
riparia, *Reiss. in Mart. Fl. Bras.* xi. 1. 43.—Bras.
rivularis, Gardn. in Hook. Ic. Pl. t. 465 = affinis.
rosmarinifolia, Lam. Illustr. i. 256 = I. Dahoon.
rotunda, Colebr. ex Wall. Fl. As. Rar. iii. 38 = I. Godajam.
rotunda, *Thunb. Fl. Jap.* 77.—Japon.
rugosa, *F. Schmidt, Reisen Amurl.* 122.—Ins. Sachalin.
rupicola, *H. B. & K. Nov. Gen. et Sp.* vii. 70.—Ecuador.
salicifolia, Jacq. Coll. v. 86. t. 2. f. 2 = Gymnosporia trigyna.
sapiiformis, *Reiss. in Mart. Fl. Bras.* xi. 1. 45.—Bras.
saponifolia, *Reiss. l. c.* 44.—Bras
Schomburgkii, *Klotzsch, in Schomb. Faun. et Fl Gui.* 1097, nomen.—Guian.
sclerophylla, *Hook. f. Fl. Brit. Ind.* i. 606.—Malacca.
scopularum, *Humb. & Bonpl. ex Schult. Mant.* iii. 387.—Ecuador.
scutiseformis, *Reiss. in Mart. Fl. Bras.* xi. 1. 50.—Bras.
Seberiti, *Panch. in Planch. & Sebert, Not. Bos Nouve. Calléd.* 285.—N. Caled.
sempervirens, Salisb. Prod. 70 = I. Aquifolium.
serrata, *Thunb. Fl. Jap.* 78.—Japon.
sessiliflora, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 378.—N. Granat.
sideroxyloides, *Griseb. in Goett. Abh.* vii. (1857) 224.—Ins. Dominic.
Sieboldii, *Miq. in Verol. Med. Akad. Wetensch.* II. ii. (1866) 84.—Japon.
sikkimensis, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 202.—Reg. Himal.
singapuriana, Wall. Cat. n. 6536 = cymosa.
Sieboldii, Spreng. Syst. i. 695 = Skimmia japonica.
sorbida, Retz. in Mart. Fl. Bras. xi. 1. 66 = paraguensis.
spicata, *Blume, Bijdr.* 1149.—Malaya.
spruceana, *Reiss. in Mart. Fl. Bras.* xi. 1. 59.—Bras.
subcordata, *Reiss. l. c.* 48.—Bras.
subpuberula, *Miq. in Verol. Med. Acad. Wetensch.* II. ii. (1866) 84.—Japon.
subtilis, *Miq.* l. c.—Japon.
Sugeroki, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 25.—Japon.
sulcata, *Wall. Cat.* n. 4390.—Burma.
symplocifolia, *Reiss. in Mart. Fl. Bras.* xi. 1. 68.—Bras.

ILEX :—
Tarajo, *Goepp. Del. Sem. Hort. Vratisl.* (1852)—*Cf. Linnæa*, xxvi. (1853–55) 749.—Hab.?
tenuifolia, Salisb. Prod. 70 = decidua.
theezefolia, *Hook. f. Fl. Brit. Ind.* i. 601.—Reg. Himal.; Burma.
theezans, Hoepl. ex Miers, in Ann. & Mag. Nat. Hist. Ser. III. viii. (1861) 389, 391 = paraguensis.
theezans, *Mart. Syst. Mat. Med. Veg. Bras.* 61.—Bras.
Thomsoni, *Hook. f. Fl. Brit. Ind.* i. 602.—Reg. Himal.
thyrsiflora, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1097, nomen.—Guiana.
tolucana, *Hemsl. Diagn. Pl. Nov.* 5.—Mexic.
trifora, *Blume, Bijdr.* 1150.—Japon.
truncata, Necs, in Flora, iv. (1821) 301 = Maytenus truncata.
tsuiiflensis, *Turcz. in Bull. Soc Nat. Mosc.* xxxi. (1858) 1. 455.—Venezuela.
umbellata, *Klotzsch, in Schomb. Faun. et Fl. Gui* 1098, nomen.—Guian.
uniflora, *Benth. Pl. Hartw.* 217.—N. Granat.
vaccinifolia, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1097, nomen.—Guian.
velutina, *Mart. ex Reiss. in Mart. Fl. Bras.* xi. 1. 53.—Bras.
venulosa, *Hook. f. Fl. Brit. Ind.* i. 602.—Reg. Himal
verticillata, *A. Gray, Man. Bot. N. U. St.* 307.—Am. bor.
vestita, Retz. in Mart. Fl. Bras. xi. 1. 54 = paraguensis.
viridis, *Champ. ex Benth. in Hook Kew Journ.* iv. (1852) 229.—Hongkong.
vismæfolia, *Reiss. in Mart Fl. Bras.* xi. 1 59.—Bras.
vitiensa, *A. Gray, Bot. U. St. Expl. Exped.* i. 295. t. 25.—Ins. Fiji.
vomitoria, [Soland. in] Ait. Hort. Kew. ed. I. i. 170 = I. Cassine.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 491 = I. Aquifolium.
Walkeri, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 603 = acq. I
Walkeri, *Wight & Gardn. ex Thw. Enum. Pl Zeyl.* 194.—Zeylan.
Wallichii, *Hook. f. Fl Brit. Ind.* i. 605.—Burma.
Wallichii, Steud. Nom. ed. II. i. 803 = macrophylla.
Watsoniana, Speach, Hist. Vég. Phan. ii. 429 = I. Dahoon.
Wightiana, Dals. & Gibs. Bomb. Fl. 143 = malabarica.
Wightiana, *Wall. Cat.* n. 4382.—Ind. or.
xeylanica, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. n. iii. (1881) 24.—Zeylan.

ILICIOIDES, Dum.-Cours. Bot. Cult. ed. I. iv. 127 (1804) = **Nemopanthus**, Rafin. (Ilicin.)

ILIOGETON, Benth. in DC. Prod. x. 412. in syn. (1846) = EUVOGETON, Endl. = **Vandellia**, Linn. (Scrophul.).

ILLA, Adans. Fam. ii. 446 (1763) = **Callicarpa**, Linn. (Verbenac.).

ILLAIREA, Lemé & C. Koch, in Verh. Berl. Gartenb. Ver. II. i. (1858) 397 = **Loasa**, Juss.
canarinoides, Lemé & C. Koch, l. c.—Am. austr.

ILLECEBRUM, Rupp. ex Linn. Coroll. Gen. 4 (1737); partim. *ILLECEBRACEÆ*, Benth. & Hook. f. Gen. 13.
Achyrantha, Linn. Sp. Pl. ed. II. 299 = Alternanthera Achyrantha.
alpinum, Vill. Hist. Pl. Dauph. i. 379 = Paronychia polygonifolia.
alsinæfolium, Pav. ex Moq. in DC. Prod. xiii. II. 389 = Guilleminea illecebroides.
alsinifolium, Scop. Delic. Insub. iii. 27 = Telanthera frutescens.
alsinifolium, Comm. ex Poir. Encyc. v. 25 = Polycarpon tetraphyllum.
alsinifolium, Linn. Mant. i. 51 = Paronychia alsinefolia.
angustifolium, Spreng. Syst. i. 818 = Alternanthera angustifolia.
arabicum, Bertol. Fl. Ital. ii. 784, in obs. = Paronychia desertorum.
arabicum, Linn. Mant. i. 51 = Paronychia arabica.
arctioides, Pourr. ex DC. Prod. iii. 371 = Paronychia arctioides.

ILLECEBRUM :—
argenteum, Pourr. in Mém. Acad. Toul. iii (1788) 321 = Paronychia argentea.
aristatum, Ait. Hort. Kew. ed. I. i. 290 = Polycarpæa aristata.
arragonicum, Vest, ex Roem. & Schult. Syst. v. 520 = Paronychia arragonica.
atrorubens, Thuill. ex Steud. Nom. ed. II. i. 803 = verticillatum.
atrovirens, Thuill. ex Steud. l. c., in syn. = præc.
bengalense, Linn. Mant. ii. 313 = Achyranthes bengalensis.
brachiatum, Linn. l. c. = Nothosæræa brachiata.
brasilianum, Steud. Nom. ed. I. 428 = Paronychia bonariensis.
canariense, Linn. f. Suppl. 426 = Paronychia canariensis.
canescens, Spreng Syst. i. 820 = Alternanthera canescens.
capitatum, Bieb. Fl. Taur. Cauc. i. 174 = Paronychia Cephalotes.
capitatum, Clem. ex Willk. & Lange, Prod. Fl. Hisp. iii. 157 = Paronychia areticides.
capitatum, Linn. Sp. Pl. 207 = Paronychia capitata.
capitatum, Sieb. Fl. Græc. i. 247 = Paronychia macrosepala.
Cephalotes, Bieb. Fl. Taur. Cauc. iii. 169 = Paronychia kurdica.
congestum, *Rud. ex Roem. & Schult. Syst.* v. 517.—Am. trop.
cucullatum, Moench, Meth. Suppl. 118 = Paronychia canariensis.
cymosum, Linn. Sp. Pl. 206 = Paronychia cymosa.
cymosum, Sm Fl. Græc. Prod. i. 163 = Paronychia echinata.
densum, Willd. in Roem. & Schult. Syst. v. 517 = Guillemínea illecebroides
denticulatum, Spreng. Syst. i. 820 = Alternanthera nodiflora.
dichotomum, Pers. Syn. i. 261 = Paronychia dichotoma.
diffusum, Willd. ex Roem. & Schult. Syst. v. 516 = Alternanthera pulchella.
divaricatum, [Soland. in] Ait. Hort. Kew. ed. I. i. 291 = Polycarpæa Teneriffæ.
dubium, Spreng. Syst. i. 818 = Iresine canescens.
echinatum, Desf. Fl. Atlant. i. 204 = Paronychia echinata.
ficoideum, Linn. Sp. Pl. ed. II. 300 = Telanthera ficoides.
flavescens, Spreng. Syst. i. 819 = Telanthera flavescens.
frutescens, Comm. ex Roem. & Schult. Syst. v. 523 = Paronychia alsinefolia.
frutescens, L'Hérit. Stirp. Nov. 75. t. 34 = Telanthera frutescens.
glabrum, Spreng. ex Moq. in DC. Prod. xiii. II. 361 = Alternanthera glabra.
glomeratum, *Hoffmgg. ex Roem. & Schult. Syst.* v. 516.—Bras.
gnaphaloides, Schousb. Vextr. Marok. 117 = Polycarpæa nivea.
gomphrenoides, Willd. ex Roem. & Schult. Syst. v. 516 = Telanthera gomphrenoides.
herniarioides, Pourr. in Mém. Acad. Toul. iii. (1788) 321 = Paronychia capitata.
Humboldtianum, Roem. & Schult. Syst. v. 517 = Telanthera canescens.
incanum, Rich. ex Moq. in DC. Prod. xiii. II. 362 = Telanthera Crucia.
ireninoides, Spreng. Syst. i. 818 = Gomphrena iresinoides
italicum, Vill. in Schrad. Journ. (1801) 413 = Paronychia argentea.
javanicum, Murr. Syst. ed. XIII. 206 = Aerva tomentosa.
Kapela, Hacq. Pl. Alp. Carn. 8. t 2. f. 1 = Paronychia imbricata.
lanatum, Hort. ex Moq. in DC. Prod. xiii. II. 299 = Aerva tomentosa.
lanatum, Hort. Par. ex Moq. l. c. 337 = Guillemínea lanuginosa.
lanatum, Spreng. Syst. ed. XIII. 206; Linn. Mant ii. 344 = Aerva lanata.
lanuginosum, Poir. ex Moq. in DC. Prod. xiii. II. 337 = Guillemínea lanuginosa.
latifolium, Steud. Nom. ed. I. 428 = Aerva tomentosa.
leucurum, Hort. Berol. ex Moq. in DC. Prod. xiii. II. 305 = Aerva leucura.
limense Hort. Par. ex Dum.-Cours. Bot. Cult. i. 346 = Telanthera frutescens.

ILLECEBRUM:—
linoevo, Thunb. Diss. Mus. Acad. Ups. xvii. App. 6 = Iresine linearis.
linnæifolium, Pers. Syn. i. 361 = Polycarpæa linæsrifolia.
longiflorum, Vill. in Schrad. Journ. (1801) 112. t. 4 = Paronychia imbricata.
lupulinum, Spreng. Syst. i. 930 = Telanthera lupulina.
maritimum, Spreng. Syst. iv. Cur. Post. 108 = Telanthera maritima.
maritimum, Vill. in Schrad. Journ. (1801) 412 = Paronychia capitata, argentea.
maritimum, Willd. ex Moq. in DC. Prod. xiii. ii. 368, 371 = Telanthera canescens.
mauritanicum, Bronss. ex Roem. & Schult. Syst. v. 516 = Paronychia argentea.
Monsoniæ, Linn. f. Suppl. 161 = Aerva Monsonia.
nanum, Spreng. Syst. i. 819 = Alternanthera nana.
narbonnense, Vill. in Schrad. Journ. (1801) 411 = Paronychia argentea.
niveum, Pers. Syn. i. 361 = Paronychia capitata.
obcordatum, Vahl, ex Steud. Nom. ed. I. 438 = Aerva tomentosa.
obliquum, Schum. & Thonn. Beskr. Guin. Pl. 142 (Quid ?).—Afr. trop.
Orva, Vahl, ex Roem. & Schult. Syst. v. 516 (Quid ?). —Aegypt.
paniculatum, Spreng. Syst. i. 819 = Telanthera paniculata.
Paronychia, Habl. Taur. 150 ; ex Bieb. Fl. Taur. Cauc. i. 173 = Paronychia Cephalotes.
Paronychia, Linn. Sp. Pl. 206 = Paronychia argentea.
peploides, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 517 (Quid ?).—Ins. S. Doming.
polygonifolium, Vill. Fl. Delph. 21 ; Hist. Pl. Dauph. ii. 557. t. 16 = Paronychia polygonifolia.
polygonoides, Linn. Sp. Pl. ed. II. 300 = Telanthera polygonoides.
pubescens, Stend. Nom. ed. I. 429 = Paronychia capitata.
pubescens, Willd. Enum. Hort. Berol. 371 = Aerva pubescens.
pulvernlentum, Spreng. Syst. iv. Cur. Post. 103 = Alternanthera pulvernlenta.
pungens, Spreng. Syst. i. 820 = Alternanthera Achyrantha.
ramosum, Hort. ex Moq. in DC. Prod. xiii. ii. 359 = Alternanthera Achyrantha.
sanguinolentum, Linn. Mant. ii. 244 = Aerva sanguinolenta.
sedifolium, Stend. Nom. ed. II. i. 808 = Paronychia sedifolia.
sericeum, Spreng. Syst. i. 819 = Gomphrena sericea.
serpyllifolium, Vill. Hist. Pl. Dauph. i. 328 (err. typ. 232) = Paronychia capitata.
sessile, Linn. Sp. Pl. ed. II. 300 = Alternanthera sessilis.
sessile, Pall. ex Moq. in DC. Prod. xiii. ii. 360 = Alternanthera sibirica.
sibiricum, Vent. ex Roem. & Schult. Syst. v. 516 = Alternanthera sibirica.
sinense, Hort. ex Moq. in DC. Prod. xiii. ii. 359 = Alternanthera Achyrantha.
spinosum, Spreng. Syst. i. 819 = Alternanthera spinosa.
spurium, Loefl. It. 118 = Loeflingia hispanica.
striatum, Pers. Syn. i. 361 = Polycarpon tetraphyllum.
subulatum, Pers. l. c. = Polycarpæa corymbosa.
suffruticosum, Linn. Sp. Pl. 206 = Paronychia suffruticosa.
tenellum, Hort. ex Moq. in DC. Prod. xiii. ii. 355 = Alternanthera tenella.
tenuifolium, Willd. Enum. Hort. Berol. Suppl. 12 = Paronychia tenuifolia.
thymifolium, Poirr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 155 = Paronychia cymosa.
tomentosum, Hort. ex Moq. in DC. Prod. xiii. ii. 299 = Aerva tomentosa.
triandrum, Buch.-Ham. ex Moq. l. c. 360 = Alternanthera sessilis.
triandrum, Llanos, Fragm. 61 = Alternanthera sessilis.
trunilense, Spreng. Syst. i. 819 = Telanthera truxillensis.
vermicularium, Linn. Sp. Pl. ed. II. 300 = Philoxerus vermicularis.
verticillatum, Barm. f. Fl. Ind. 66 = Portulaca quadrifida.
verticillatum, *Linn. Sp. Pl.* 206.—Europ. occ. ; Reg. Mediterr.
villosum, Stend. Nom. ed. I. 429 = Paronychia capitata.
villosum, Willd. ex Roem. & Schult. v. 516 = Telanthera villosa.

ILLICIUM. Linn. Syst. ed. X. 1050 (1759). *MAGNOLIACEÆ*, Benth. & Hook. f. i. 18.
CYMBOSTEMON, Spach, Hist. Vég. Phan. vii. 444 (1839).
anisatum, Bartr. ex Michx. Fl. Bor. Am. i. 99 (err. typ. 598) = parviflorum.
anisatum, *Guerin. Fructi.* i. 888. t. 69 (non Linn.). —China.
anisatum, *Linn. Syst.* ed. X. 1050.—Japon.
cambodianum, *Hance, in Journ. Bot.* xiv. (1876) 240. —Cambodia.
cambogianum, Pierre, Fl. Forest. Cochinch. t. 4 = præc.
floridanum, *Ellis, in Phil. Trans.* lx. (1770) 534. t. 19. —Am. bor.
Griffithii, *Hook. f. & Thoms. Fl. Ind.* i. 74.—Reg. Himal.
japonicum, Siebold, in Verh. Batav. Gen. xii. (1830) 50 = anisatum.
majus, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 40.—Burma.
parviflorum, *Michx. Fl. Bor. Am.* i. 226 (err. typ. 526). —Florida.
religiosum, Sieb. & Zucc. Fl. Jap. i. 6 = anisatum, *Linn.* ?
Sau-ki, Perr. in Mém. Soc. Linn. Par. iii. (1825) 121 = anisatum, *Linn.* ?
variegatum, Hort. ex Gard. Chron. (1861) 735 = anisatum, *Linn.*

ILLIGERA, Blume, Bijdr. 1158 (1826). *COMBRETACEÆ*, Benth. & Hook. f. i. 689.
CORYSANDRIA, Griff. in Proc. Linn. Soc. i. (1846) 281.
GRONOVIA, Blanco, Fl. Filip. ed. I. 186 (1837).
HENSCHELIA, Presl, Rel. Haenk. ii. 81. t. 63 (1831).
appendiculata, Blume, Bijdr. 1158 = corysandeia.
appendiculata, Vidal, Sinops. t. 48. f. F = Mayenlasa.
celebica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 315.— Ins. Celebes.
corysandeia, *Meisen. in DC. Prod.* xv. i. 251.—Burma.
dasyphylla, *Miq. Fl. Ind. Bat.* i. 1. 1094.—Sumatra.
dubia, *Span. in Linnæa*, xv. (1841) 187.—Ins. Timor.
khasiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 461.—Reg. Himal.
Kunzii, *C. B. Clarke, l. c.* 460.—Burma ; Malaya.
lucida, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 29.—Java.
obtusa, *Meisen. in DC. Prod.* xv. i. 251 (sp. dub.).— Ind. or.
Meyeniana, *Kunth, ex Walp. in Nov. Act. Nat. Cur.* xix. Suppl. i. (1843) 410.—Ins. Philipp.
pentaphylla, *Wight, in Trans. Linn. Soc.* xxvii. (1869) 26.—Afr. trop.
pulchra, *Blume, Bijdr.* 1154.—Java.
rhodantha, *Hance, in Journ. Bot.* xxi. (1883) 291.— China.

ILLIPE, F. Muell. Extra-trop. Pl. ed. Am. (1884) 181 = Bassia, Koenig (Sapotac.).
latifolia, F. Muell. l. c. = B. latifolia.

ILLUS, Haw. Monog. Narcis. 4 (1831) = Narcissus, Tourn. (Amaryll.).
cernuus, Haw. l. c. = N. cernuus.
triandrus, Haw. l. c. = N. triandrus.

ILMU, Adans. Fam. ii. 497 (1763) = Romulea, Maratti (Irid.).

ILOGETON, A. Juss. in Orb. Dict. xi. 459 (1849) = seq.

ILVOGETHOS, Hassk. Cat. Hort. Bog. Alt. 146 (1844) = seq.

ILVOGETON, Endl. Gen. 684 (1839) = Vandellia, Linn. (Scrophul.).
alsinoides, Endl. in Walp. Rep. iii. 297 = V. alsinoides.
nervosum, Hassk. Cat. Hort. Bog. Alt. 146 = V. nervosum.
scapigerum, Endl. in Walp. Rep. iii. 297 = V. scapigera.
sphærocarpum, Hassk. Cat. Hort. Bog. Alt. 146 = V. sphærocarpa.
subulatum, Endl. in Walp. Rep. iii. 297 = V. subulata.

ILYANTHES, Raf. Fam. Nat. 12 (1820). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 955.
alternifolia, Urban, in Berl. Deutsch. Bot. Ges. ii. (1884) 436 = Bonnaya alternifolia.
attenuata, Raf. ex Benth. in DC. Prod. x. 479 = riparia.

ILYSANTHES:—
capensis, *Benth. in DC. Prod.* x. 419.—Afr. austr.
clausa, Urban, in Berl. Deutsch. Bot. Ges. ii. (1884) 436 = Bonnaya clausa.
dilatata, Raf. ex Benth. in DC. Prod. x. 419 = riparia.
grandiflora, *Benth. l. c.* 418.—Am. bor.
gratioloides, Benth. l. c. 419 = riparia.
gratioloides, Regel, Fl. Ussur. 120. n. 359 = Vandellia erecta.
hyssopioides, *Benth. in DC. Prod.* x. 419.—Ind. or. ; China.
lobelioides, *Benth. Fl. Austral.* iv. 498.—Austral.
minima, *Benth. in DC. Prod.* x. 420.—Ind. or.
oblongifolia, *Baker, in Journ. Bot.* xx. (1882) 221.— Madag.
oppositifolia, Urban, in Berl. Deutsch. Bot. Ges. ii. (1884) 435 = Bonnaya oppositifolia.
parviflora, *Benth. in DC. Prod.* x. 419.—Ind. or. ; Afr. trop.
pusilla, Urban, in Berl. Deutsch. Bot. Ges. ii. (1884) 435 = Bonnaya pusilla.
refracta, *Benth. in DC. Prod.* x. 419.—Am. bor.
reptans, Urban, in Berl. Deutsch. Bot. Ges. ii. (1884) 436 = Bonnaya reptans.
riparia, *Rafin. Ann. Nat.* 13.—Am. bor. et trop.
rotundifolia, *Benth. in DC. Prod.* x. 420.—Ind. or. ; Ins. Mascar.
saginoides, *Franch. & Sav. Enum. Pl. Jap.* i. 346.— Japon.
saxicola, Chapm. Fl. S. U. St. 294 = refracta.
serrata, Urban, in Berl. Deutsch. Bot. Ges. ii. (1884) 436 = Bonnaya brachiata.
tenuifolia, Urban, l. c. 435 = Bonnaya tenuifolia.
trichotoma, Urban, l. c. = Bonnaya trichotoma.
veronicifolia, Urban, l. c. 436 = Bonnaya veronicæfolia.

ILVSANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 = præc.

ILYTHURIA, Raf. Fl. Tellur. iv. 51 (1836) = Clinogyne, Salisb. (Scitam.).
cannæformis, Rafin. l. c. = C. dichotoma.

IMANTINA, Hook. f. in Benth. & Hook. f. Gen. ii. 920 (1873). *RUBIACEÆ*, Benth. & Hook. f. i. c. Deplanchei, *Hook. f. l. c.*—N. Caled.

IMANTOPHYLLUM, Benth. & Hook. f. Gen. iii. 739 (1883) = seq.

IMATOPHYLLUM, Hook. Bot. Mag. t. 3856 (1838) = Clivia, Lindl. (Amaryll.).
Aiteni, Hook. Bot. Mag. t. 3856 = C. nobilis.
cyrtanthiflorum, Van Houtte, in Fl. des Serres, Sér. II. viii. (1869–70) 87. t. 1877 = C. miniata.
maximum, Guillon, in Kerr. Hortic. (1831) 140 = C. nobilis ?
miniatum, Hook. Bot. Mag. t. 4788 = C. miniata.

IMBRICARIA, Comm. ex Juss. Gen. 159 (1789). *SAPOTACEÆ*, Benth. & Hook. f. ii. 661.
BINECTARIA, Forsk. Fl. Aegypt. Arab. 89 (1775).
Binectaria, A. DC. in DC. Prod. viii. 201 = maximæ ?
berbonica, Gaertn. f. Fruct. iii. 128. t. 206 = maximæ.
Commersonii, G. Don, Gen. Syst. iv. 85 = Labramia Bojeri ?
coriacea, A. DC. in DC. Prod. viii. 200 = Labramia Bojeri.
fragrans, *Baker, in Oliver, Fl. Trop. Afr.* iii. 509.— Afr. trop.
macrocarpa, *Gaertn. f. Fruct.* iii. 124. t. 206.—Hab. ? malabarica, *Poir. Encyc.* iv. 484.—Ind. or.
maxima, *Poir. l. c.* 483.—Ins. Mascar.
media, *Bojer, ex Baker, Fl. Mauric.* 195.—Ins. Mauric.
obovata, Nees, ex Harv. Thes. Cap. i. 23 = Mimusops obovata.
Perrottii, *Montr. in Mém. Acad. Lyon*, x. (1860) 230. —N. Caled.
petiolaris, *A. DC. in DC. Prod.* viii. 200.—Ins. Mauric.

IMBRICARIA, Sm. in Trans. Linn. Soc. iii. (1797) 257 = Bacchus, Linn. (Myrtac.).
ciliata, Sm. l. c. = Micromyrtus calycophylla.
crenulata, Sm. l. c. 259 = B. crenulata.
tenella, Dryand. in Ker. & Sims, Ann. Bot. ii. (1807) 319 = B. crenulata.

IMHOFIA, Herb. App. 18 (1921) = **Hessea**, Herb. (Amaryll.).
Borziana, Kuntb, Enum. Pl. v. 627 = H. gemmata.
Burchelliana, Herb. Amaryll. 1900. t. 29. f. 5 = H. gemmata.
cinnamomea, M. Roem. Syn. Ensat. 39 = H. crispa.
corvacea, Salisb. Gen. Pl. Fragm. 118 = Nerine curvifolia
crispa, Herb. Amaryll. 290 = H. crispa.
filifolia, Herb. l. c. 290. t. 29. f. 6 = H. filifolia.
gemmata, Herb. l. c. 291 = H. gemmata.
glauca, Salisb. Gen. Pl. Fragm. 118 = Nerine curvifolia.
marginata, Herb. App. 18 = Nerine marginata.
rosea, Salisb. Gen. Pl. Fragm. 118 = Amaryllis Belladonna.
sarmentosa, Salisb. l. c. = Nerine sarniensis.
tenella, M. Roem. Syn. Ensat. 38 = Strumaria filifolia.
venusta, Salisb. Gen. Pl. Fragm. 118 = Nerine sarniensis.

IMPATIENS, Riv. ex Linn. Syst. ed. I (1735). *GERANIACEAE*, Benth. & Hook. f. i. 277.
BALSAMINA, Gaertn. Fruct. ii. 151. t. 118 (1791).
abyssinica, *Hook. f. ex Oliver, Fl. Trop Afr.* i. 302. — Afr. trop.
acaulis, *Arn. in Hook. Comp. Bot. Mag* i. (1835) 325. —Ind. or.
acuminata, *Benth. in Wall. Cat.* n. 4784.—Reg. Himal.
Akka, Bedd. in Madr. Journ. Sc. Ser. II. xx. (1859) 69. f. 11 = cornuta.
alba, *Houllet, in Rev. Hortic.* (1884) 143.—Hab.?
albida, Wight, in Madr. Journ. Sc. Ser. I. v. (1837) 7 = Henslowiana.
albo-flava, *Miq. Fl. Ind. Bat. Suppl.* 396.—Sumatra.
amphorata, *Edgew. in Trans. Linn. Soc.* xx. (1846) 30. —Reg. Himal.
amplexicaulis, *Edgew. l. c.* 37.—Reg. Himal.
anamallayensis, Bedd. in Madr. Journ. Sc. Ser. II xx. (1859) 68 = Gonghii.
angustiflora, *Hook. f. Fl. Brit. Ind.* i. 480.—Reg. Himal.
angustifolia, Blume, Cat. Gew. Buitens. 49 = Hydrocera triflora.
angustifolia, Steud. Nom. ed. II. i. 804 = I. Balsamina.
appendiculata, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 328.—Zeylan.
arcuata, Benth. in Wall. Cat. n. 4738 = I. Balsamina.
arguta, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 137.—Reg. Himal.; China.
Arnottiana, Miq. ex Hook. f. & Thoms. l. c. 122 = diversifolia
Arnottii, *Thw. Enum. Pl. Zeyl.* 67.—Zeylan.
assamensis, *Griff. Notul.* iv. 459. t. 576.—Assam.
aurea, *Wall. Cat.* 26.—Am. bor.
auriculata, *Wight, in Madr. Journ. Sc.* Ser. I. v. (1837) 8.—Ind. or.
baccifera, Roxb. ex Wight & Arn. Prod. 141 = Hydrocera angustifolia.
Ballardi, *Bedd. Ic. Pl. Ind. Or.* i. 44. t. 192.—Ind. or.
Balsamina, *Linn. Sp. Pl.* 938.—As. trop.
Baroni, *Baker, in Journ. Bot.* xx. (1882) 49.—Madag.
Beddomei, *Hook. f. Fl. Brit. Ind.* i. 442.—Malabar.
bella, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 135.—Reg. Himal.
bicolor, *Hook. f. l. c.* vi. (1862) 7.—Afr. trop.
bicolor, Royle, Illustr. Bot. Himal. 151. t. 38 = amphorata.
biconnata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 460.—Reg. Himal.
bifida, Thunb. Prod. Pl. Cap. 41 = flaccida.
bifora, *Walt. Fl. Carol.* 218.—Am. bor.
biglandulosa, Moon, Cat. Ceyl. Fl. 18 = Hookeriana.
bipartita, Arn. in Hook. Comp. Bot. Mag. i. (1835) 328 = flaccida.
Blumei, *Zoll. & Mor. Syst. Verz. Zoll.* 13.—Reg. Himal.; Songar.
borealis, Sweet, Hort. Brit. ed. II. 388 = I. Noli-tangere.
borneensis, *Miq. Illustr. Fl. Ind. Archip.* 96.—Borneo.
brachycentra, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 179.—As. centr.
brachystis, Colebr. ex Roxb. Fl. Ind. ed. Carey, ii. 459 = fimbriata.
brevicornu, Arn. in Hook. Comp. Bot. Mag. i. (1835) 321 = leptopoda.
brevisepala, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 187.—Afr. trop.
bulbosa, Moon, Cat. Ceyl. Fl. 16 = acaulis.

IMPATIENS:—
Burtoni, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 187. —Afr. trop.
calycina, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 463 = scabrida.
campanulata, *Wight, in Madr. Journ. Sc.* Ser. I. v. (1837) 11.—Ind. or.
candida, Lindl. Bot Reg. (1840), Misc. 85 = Roylei.
capensis, Boj. ex Baker, in Journ. Linn. Soc. xx. (1883) 115 = formosa.
capensis, *Meerb. Afbeel. Gew.* t. 10; *Thunb. Prod. Fl. Cap.* 41.—Afr. austr.
capillipes, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 135.—Burma.
Catharti, *Hook. f. Fl. Brit. Ind.* i. 478.—Reg. Himal.
caucasica, Stev. in Bull. Soc. Nat. Mosc. xxi. (1848) II. 276 = I. Noli-tangere.
celebica, *Miq. Illustr. Fl. Ind. Archip.* 94.—Ins. Celebes.
chinensis, *Linn. Sp. Pl.* 937.—Reg. Himal.; Malaya. chonceorea, *Hassk. Hort. Bog. Desc.* 146.—Java.
circaeoides, Turcz. in Bull. Soc. Nat Mosc. xxxvi (1863) I. 594 = Gonghii.
circaeoides, *Wall. Cat. n.* 4772 —Burma.
clavicornu, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 271.—Ind. or.
coccinea, Wall. Cat. n. 4732 = I. Balsamina.
cochleata, *Lour. Fl. Cochinch.* ii 512.—China
comorensis, *Baker, in Journ. Linn. Soc. xx.* (1883) 114.—Madag.
concinna, *Hook. f. Fl. Brit. Ind.* i. 449.—Malabar.
cordata, Wight, in Madr. Journ. Sc. Ser. I. v. (1837) 10 = reticula.
cordifolia, Herb. Madr. ex Wall. Cat. n. 4758 = scapiflora.
cornigera, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 323.—Ceylan.
cornigera, Hook. Bot. Mag. t. 4628 = glandulifera
cornuta, Linn. Sp Pl. 937 = I. Balsamina.
crenata, *Bedd. in Madr. Journ. Sc.* Ser. II xx (1859) 69. t. 7. f. 10.—Ind. or.
cristata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 456 = scabrida.
crispura, Edgew. ex Hook. f. & Thoms. in Journ Linn. Soc. iv. (1860) 148 = bicornuta.
cucullata, Gilib. Fl. Lituan. ii. 121 = I. Noli-tangere.
cuspidata, Wight & Arn. in Hook. Comp. Bot. Mag. i. (1835) 321 = latifolia.
cyclocorna, *Miq. Illustr. Fl. Ind. Archip.* 92.—Java.
cymbifera, *Hook. f. Fl. Brit. Ind.* i. 474.—Reg. Himal.
Dalzellii, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 123.—Ind. or.
dasysperma, *Wight, in Madr. Journ. Sc.* Ser. I. v. (1837) 7.—Ind. or.
Davidi, *Franch. Pl. Davidian.* i. (1884) 65.—Mongol.
debilis, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 271 = tenella?
Denisonii, *Bedd. in Madr. Journ. Sc.* Ser. III i. (1864) 41.—Ind. or.
depauperata, *Hook. f. Fl. Brit. Ind.* i. 481.—Reg. Himal.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 397.—Sumatra.
discolor, DC. Prod. i. 687 = scabrida.
discolor, *Wall. Cat.* n. 4767.—Reg. Himal.
diversifolia, *Heyne, in Wall. l. c.* 4749.—Ind. or.
Edgeworthii, *Hook. f. Fl. Brit. Ind.* i. 476.—Reg. Himal.
elata, Edgew. in Trans. Linn. Soc. xx. (1846) 41 = micranthemum.
elegans, *Bedd. in Madr. Journ. Sc.* Ser. II. xx. (1859) 67.—Ind. or.
elongata, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 324.—Zeylan.
emirnensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 115.—Madag.
erianthe, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 270 = Munroii.
esbenya, *Miq. Fl. Ind. Bat. Suppl* 397.—Sumatra.
fasciculata, Lam. Encyc. i. 859 = chinensis.
filicorna, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 6.—Afr. trop.
filiformis, Wight & Arn. Prod. 140 = inconspicua.
fimbriata, *Hook. Exot. Fl.* ii. t. 146.—Reg. Himal.
firmula, *Baker, in Journ. Linn. Soc.* xx. (1883) 114. —Madag.
flaccida, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 322. —Ind. or.

IMPATIENS:—
flagellifera, Hochst. ex A. Rich. Tent. Fl. Abyss. i 120 = tinctoria.
flavida, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 127.—Reg. Himal.
floribunda, Wight, in Madr. Journ. Sc. Ser. I. v. (1837) 7 = flaccida.
fruticosa, *Leschen. ex DC. Prod.* i. 687.—Ind. or.
fulva, Nutt. Gen. Am. i. 146 = biflora.
Gardneriana, Wight, ex Hook. f. Fl. Brit. Ind. i. 450 = concinna.
Gardneriana, *Wight, Ic.* t. 1050.—Ind. or.
gibbosa, Arn. in Hook. Comp. Bot. Mag. i. (1835) 321 = leptopoda.
gigantea, Edgew. in Trans. Linn. Soc. xx. (1846) 38 = sulcata.
glandulifera, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 322.—Zeylan.
glandulifera, Royle, Illustr. Bot. Himal. 151. t. 38, f. 3 = Roylei.
glandulifera, Lindl. Bot. Reg. (1840) t. 22 = Roylei
glauca, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 155.—Reg. Himal.
Gordoni, *Horne, in Baker, Fl. Maurit.* 38.—Ins. Mascar.
Gonghii, *Wight, Illustr. Ind. Bot.* i. 160.—Ind. or.
gracilis, Bedd. in Madr. Journ. Sc. Ser. II. xx. (1859) 68 = acaulis.
grandifora, Herb. Madr. ex Wall. Cat. n. 4759 = grandis
grandis, *Heyne, in Roxb. Fl. Ind. ed. Carey*, ii. 465.—Ind. or.
Griffithii, *Hook. f. & Thoms. in Journ. Linn Soc.* iv. (1860) 120.—Malaya.
Hamiltoniana, D. Don, Prod. Fl. Nep. 204 = scabrida.
Henslowiana, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 322.—Ind. or.
heterophylla, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 458 = chinensis.
hians, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 7.—Afr. trop.
hirsuta, *Blume, Cat. Gew. Buitens.* 50, nomen.—Java.
hispidula, Benth. in Wall. Cat. n. 4740 = puberula.
Hofmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 121 = discolor.
Hookeriana, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 324.—Zeylan.
Humblotiana, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 986.—Madag.
inconspicua, *Benth. in Wall. Cat.* n. 4741.—Ind. or.
insignis, *DC. Prod.* i. 688.—Reg. Himal.
Irvingii, *Hook. f. ex Oliver, Fl. Trop. Afr.* i. 300.—Afr. trop.
janthina, *Thw. Enum. Pl. Zeyl.* 68.—Zeylan.
japonica, *Franch. & Sav. Enum. Pl. Jap.* ii. 210.—Japon.
javensis, *Steud. Nom.* ed. II. i. 804.—Java.
Jerdoniae, *Wight, Ic.* t. 1602.—Ind. or.
Junghuhnii, *Miq. Illustr. Fl. Ind. Archip.* 99.—Sumatra.
Jurpis, *Buch.-Ham. in Wall. Cat.* n. 4761.—Reg. Himal.
Kilimanjari, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 398.—Afr. trop.
Kirkii, *Hook. f. ex Oliver, Fl. Trop. Afr.* i. 300.—Afr. trop.
Kleinii, *Wight & Arn. Prod.* 140.—Ind. or.
Korthalsii, *Miq. Illustr. Fl. Ind. Archip.* 100.—Sumatra.
laevigata, *Wall. Cat.* n. 4753.—Reg. Himal.
latiflora, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 139.—Reg. Himal.
latifolia, Hook. Bot. Mag. t. 2625 = flaccida.
latifolia, *Linn. Sp. Pl.* 937.—Ind. or.
latifolia, Wall. Cat. n. 4737 = leptopoda.
latifolia, Wight & Arn. Prod. 138 = lucida.
Lawii, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 122.—Ind. or.
laxiflora, *Edgew. in Trans. Linn. Soc.* xx. (1846) 40.—Reg. Himal.
Lechenaultii, Walp. Rep. i. 467 = Leschenaultii.
Lemaoni, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 154.—Afghan.
leptocerca, *DC. Prod.* i. 688.—Reg. Himal.
leptocerata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = prsec.

IMPATIENS:—
leptopoda, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 321.—Zeylan.
leptura, *Hook. f. Fl. Brit. Ind.* i. 467.—Ind. or.
Leschenaultii, *Wall. Cat.* n. 4739.—Ind. or.
leucantha, *Thw. Enum. Pl. Zeyl.* 67.—Zeylan.
ligulata, *Bedd. in Madr. Journ. Sc.* Ser. II. xx. (1859) 67.—Ind. or.
linearis, *Arn. in Hook. Comp. Bot. Mag.* i. (1835) 323. —Zeylan.
lobata, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 271 = tavoyana.
Lobbiana, Turcz. l. c. 270 = I. Balsamina.
longicornu, Strach. & Wint. ex Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1860) 147 = bicornuta.
longicornu, Wall. Cat. n. 4729 = amphorata, bicornuta.
longicornu, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 462 = leptocorea.
longiflora, Heyne, ex Hook. f. Fl. Brit. Ind. i. 447 = tenella.
longifolia, Wight, in Wall. Cat. n. 4734 = I. Balsamina.
longipes, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 151.—Reg. Himal.
lucida, *Heyne, in Wall. Cat.* n. 4738.—Ind. or.
lutea, Lam. Fl. Fr. ii. 668 = I. Noli-tangere.
Lyallii, *Baker, in Journ. Linn. Soc.* xx. (1883) 118. —Madag.
Mackeyana, *Hook. f. l. c.* vii. (1864) 186.—Afr. trop.
macrochila, Lindl. Bot. Reg. (1840) t. 8 = Roylei.
macrophylla, *Gardn. ex Hook. Bot. Mag.* t. 4663.— Zeylan.
macroptera, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 186.—Afr. trop.
maculata, Mihl. Cat. 26 = biflora.
maculata, *Wight, in Madr. Journ. Sc.* Ser. I. v. (1837) 12.—Ind. or.
madagascariensis, *Wight & Arn. Prod.* 140.— Madag.
malabarica, Klein, ex Wight & Arn. l. c. 139 = diversifolia.
malayensis, Griff. Notul. iv. 457. t. 576. f. 2 = I. Balsamina?
mamillensis, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 317.—Ins. Philipp.
Mannii, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 7.— Afr. trop.
Masterstana, *Stend. Nom.* ed. II. i. 804.—Ind. or.
micrantha, D. Don, Prod. Fl. Nep. 203 = racemosa.
micrantha, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 121. —Abyssin.
microanthemum, *Edgew. in Trans. Linn. Soc.* xx. (1846) 40.—Reg. Himal.
minutiflora, *Span. in Linnaea*, xv. (1841) 185.—Ins. Timor.
mishmiensis, *Hook. f. Fl. Brit. Ind.* i. 476.—Reg. Himal.
modesta, *Wight in Madr. Journ. Sc.* Ser. I. v. (1837) 13.—Ind. or.
mollis, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 461 = pubersula.
monzana, *Rafin. Atl. Journ.* 153 (Quid?).—Am. bor.
moschata, Edgew. in Trans. Linn. Soc. xx. (1846) 38 = Roylei.
multiflora, Wall. Cat. n. 4742 = tripetala.
Munroii, *Wight, Illustr. Ind. Bot.* i. 160.—Ind. or.
nautila, *Lour. Fl. Cochinch.* 512.—Cochinch.
myzorensis, Roth, ex Wall. Cat. n. 4743 = inconspicua.
myzorensis, *Roth, Nov. Pl. Sp.* 184.—Ind. or.
natans, Willd. Sp. Pl. i. 1175 = Hydrocera triflora.
nematocorea, *Miq. Illustr. Fl. Ind. Archip.* 97.— Java.
Noli-me-tangere, Crantz, Inst. ii. 430 = I. Noli-tangere.
Noli-tangere, D. Don, Prod. Fl. Nep. 204 = scabrida.
Noli-tangere, *Linn. Sp. Pl.* 938.—Europ. ; Sibir.
Noli-tangere, Michx. Fl. Bor. Am. i. 149, partim = aurea.
oborata, D. Don, Prod. Fl. Nep. 203 = leptocorea.
oppositifolia, *Linn. Sp. Pl.* 937.—Ind. or.
oppositifolia, Rottl. ex Wight & Arn. Prod. 138 = chinensis.
orchidoides, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 41.—Ind. or.
pachycentra, *Rupr. Fl. Cauc.* i. 265.—Reg. Caucas.
pallens, Edgew. in Trans. Linn. Soc. xx. (1846) 39 = amphorata.
pallida, Nutt. Gen. Am. i. 146 = aurea.

IMPATIENS:—
palpebrata, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 186.—Afr. trop.
paludosa, *Hook. f. Fl. Brit. Ind.* i. 480.—Reg. Himal.
palustris, Pers. Syn. i. 257 = I. Noli-tangere.
parasitica, Bedd. in Madr. Journ. Sc. Ser. II. xx. (1859) 68 = Jerdoniæ.
Parishii, *Hook. f. Fl. Brit. Ind.* i. 456.—Burma.
parviflora, *DC. Prod.* i. 687.—Turkestan. ; Sibir.
parviflora, *Bedd. in Madr. Journ. Sc.* Ser. II. xx. (1859) 68.—Ind. or.
pendula, *Heyne, in Wall. Cat.* n. 4744.—Ind. or.
pendulifloru, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 138, 70 = I. Noli-tangere.
Persii, *Teysm. ex Miq. Choix Pl. Buitens.* t. 5.— Sumatra.
Petersiella, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 594 = inconspicua.
phœnicea, *Bedd. in Madr. Journ. Sc.* Ser. II. xix. (1858) 176.—Ind. or.
picta, Knowl. & Westc. Flor. Cab. iii. (1849) 149. t. 138 = amphorata.
platypetala, *Lindl. Bot. Reg.* (1846) t. 53.—Java.
porrecta, *Wall. Cat.* n. 7375.—Reg. Himal.
puberula, *DC. Prod.* i. 687.—Reg. Himal.
pulcherrima, *Dalzell, in Hook. Kew Journ.* ii. (1850) 37.—Ind. or.
pulchra, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 139.—Reg. Himal.
pulviscula, *Bedd. in Madr. Journ. Sc.* Ser. II. xix. (1858) 176 = Congbii.
punctata, Wall. ex Hook. f. & Thoms. in Journ. Linn. Soc. iv. (1860) 136 = scabrida.
pusilla, Heyne, in Wall. Cat. n. 4745 = inconspicua.
pythoriche, *Miq. Fl. Ind. Bat. Suppl.* 397.—Sumatra.
racemosa, *DC. Prod.* i. 688.—Reg. Himal.
racemosa, D. Don, Prod. Fl. Nep. 203 = insigula.
racemosa, Hook. f. & Thoms. in Journ. Linn. Soc. iv. (1860) 147 = radiata.
racemosa, Wall. Cat. n. 4730 = racemosa, leptocorea.
racemulosa, *Wall. l. c.* n. 7374.—Reg. Himal.
radiata, *Hook. f. Fl. Brit. Ind.* i. 476.—Reg. Himal.
radicans, *Benth. in Wall. Cat.* n. 4763.—Reg. Himal.
radicans, *Zoll. & Mor. Syst. Verz. Zoll.* 14.—Java.
ramosiana, Bedd. in Hook. Kew Journ. iii. (1851) 231 = inconspicua.
repens, *Moon, Cat. Cryl. Pl.* 18 ; *et in Wight, Illustr. Ind. Bot.* i. 160.—Ind. or.
reticulata, *Wall. Pl. As. Rar.* i. 19. t. 19.—Burma.
Rheedii, Wight & Arn. Prod. 138 = tripetala?
rivalis, *Wight, in Madr. Journ. Sc.* Ser. I. v. (1837) 13. t. 8.—Ind. or.
rosea, Lindl. Bot. Reg. (1841) Misc. 6. t. 27 = I. Balsamina.
roseoviridifolia, Retz. Obs. v. 29 = oppositifolia, tenella.
Rothii, *Hook. f. in Oliver, Fl. Trop. Afr.* i. 302.— Afr. trop.
Roylenna, Payer, Organog. Comp. 84 = Roylei.
Roylei, Klotzsch, in Bot. Ergeb. Waldem. Reise Bot. 121 = amphorata.
Roylei, *Walp. Rep.* i. 478.—Reg. Himal.
rubescens, Benth. in Wall. Cat. n. 4747 = tomentosa.
Rutenbergii, *O. Hoffm. in Bremen, Abh.* vii. (1882) 335.—Madag.
Sakeriana, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 185.—Afr. trop.
salicifolia, *Baker, l. c.* xx. (1883) 115.—Madag.
salicifolia, *Hook. f. & Thoms. l. c.* iv. (1860) 124.— Reg. Himal.
salicifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 271 = I. Balsamina.
scabricula, *DC. Prod.* i. 687.—Reg. Himal.
scabriuscula, *Heyne, in Roxb. Fl. Ind. ed. Carey*, ii. 464.—Ind. or.
scapiflora, *Heyne, in Roxb. l. c.* 464.—Ind. or.
scapiflora, Hook. Bot. Mag. t. 3537 = scaulis.
scapiflora, Wight & Arn. Prod. 137, partim = Beddomei.
semiverticillata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 594.—Ind. or.
serrata, *Benth. in Wall. Cat.* n. 4771.—Reg. Himal.
setacea, Colebr. in Hook. Exot. Fl. ii. t. 187 = chinensis.
setosa, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 138.—Ind. or.
sinensis, Bunch.-Ham. ex Benth. in Wall. Cat. n. 4735 = aurantia.
sinensis, J. F. Gmel. Syst. 415 = chinensis.

IMPATIENS:—
spiritlis, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 138.—Reg. Himal.
stenantha, *Hook. f. Fl. Brit. Ind.* i. 478.—Reg. Himal.
Stocksii, *Hook. f. in Journ. Linn. Soc.* iv. (1860) 113.—Ind. or.
subcordata, *Arn. in Hook. Comp. Bot Mag.* i. (1835) 323.—Zeylan.
sulcata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 456.— Reg. Himal.
Sultani, *Hook. f. Bot. Mag.* t. 6643.—Afr. trop.
sumatrana, *Miq. Fl. Ind. Bat. Suppl* 396.—Sumatra.
Tangachee, *Bedd. in Madr. Journ. Sc.* Ser. II. xx. (1859) 68.—Ind. or.
tavagua, Steud. Nom. ed. II. i. 805, sphalm. = seq.
tavoyana, *Benth. in Wall. Cat.* n. 4778.—Burma.
tenella, *R. Br. in Salt, Abyss. App.* 64, nomen.— Abyssin.
tenella, *Heyne, in Wall. Cat.* n. 4746 A.—Ind. or.
tenella, Wall. l. c. 4746 B.= Kleinii.
tenuicula, Steud. Nom. ed. II. i. 805 = tenella.
tenuis, Bedd. in Madr. Journ. Sc. Ser. II. xx. (1859) 69. t. 7. f. 13 = modesta.
tenuifolia, Buch.-Ham. ex Hook. f. & Thoms. in Journ. Linn. Soc. iv. (1860) 126 = tripetala.
Textori, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 76.— Japon.
Teysmannii, *Miq. Fl. Ind. Bat. Suppl.* 396.—Sumatra.
Thomsoni, *Hook. f. in Journ. Linn. Soc.* iv. (1860) 128.—Reg. Himal.
Thomsoni, *Oliver, l. c.* xxi. (1885) 398.—Afr. trop.
tinctoria, *A. Rich. Tent. Fl. Abyss.* i. 120.—Abyss.
tingens, Edgew. in Trans. Linn. Soc. xx. (1846) 41 = racemosa.
tomentosa, *Heyne, in Wall. Cat.* n. 4751.—Ind. or.
travancorica, *Bedd. Ic. Pl. Ind. Or.* i. 29.—Ind. or.
trichocentra, *Baker, in Journ. Linn. Soc.* xx. (1883) 116.—Madag.
tricornis, Lindl. Bot. Reg. (1840) t. 9 = scabrida.
triflora, Linn. Sp. Pl. 938 = Hydrocera triflora.
trilobata, *Colebr. in Hook. Exot. Fl.* ii. t. 141.—Reg. Himal.
tripetala, *Roxb. Hort. Beng.* 18 ; *Fl. Ind.* i. 658.— Reg. Himal.
tropæolifolia, *Griff. ex Hook. f. Fl. Brit. Ind.* i. 472.— Reg. Himal.
truncata, *Thw. Enum. Pl. Zeyl.* 66.—Zeylan.
tuberculata, *Hook. f. & Thoms. in Journ. Linn. Soc.* iv. (1860) 155.—Reg. Himal.
umbellata, *Heyne, ex Roxb. Fl. Ind. ed. Carey*, ii. 464. —Ind. or.
umbrosa, Edgew. in Trans. Linn. Soc. xx. (1846) 39 = amphorata.
uncinata, *Wight, in Madr. Journ. Sc.* Ser. I. v. (1837) 11.—Ind. or.
urticifolia, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 457. —Reg. Himal.
verrucosa, Bedd. in Madr. Journ. Sc. Ser. II. xx. (1859) 69. t. 7. f. 9 = viridula.
verticillata, *Wight, l. c.* Ser. I. v. (1837) 15.—Ind. or.
violæflora, *Hook. f. Fl. Brit. Ind.* i. 457.—Burma.
viridiflora, *Wight, in Madr. Journ. Sc.* Ser. I. v. (1837) 9.—Ind. or.
viscida, *Wight, l. c.* 12.—Ind. or.
viscosa, *Bedd. l. c.* Ser. II. xx. (1859) 68. t. 7. f. 7.— Ind. or.
Walkeri, *Hook. ex Arn. in Hook. Comp. Bot. Mag.* i. (1835) 324.—Zeylan.
Walleriana, *Hook. f. in Oliver, Fl. Trop. Afr.* i. 302.— Afr. trop.
Wightiana, *Bedd. in Madr. Journ. Sc.* Ser. II. xx. (1859) 67.—Ind. or.
Zippelii, *Miq. Illustr. Ind. Archip.* 94.—N. Guin.

IMPERATA, Cyrill, Fl. Rar. Ic. ii. 98. t. 11 (1793), *GRAMINEÆ*, Benth. & Hook. f. iii. 1195.
SYLLEPIS, Foern. ex Benth. & Hook. f. Gen. iii. 1195 (1883).
Allang, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 295 = arundinacea.
arundinacea, *Cyrilli, Fl. Rar. Ic.* ii. 96. t. 11.—Reg. calid.
brasiliensis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 381.—Am. trop.
brevifolia, Vasey, Cat. Grass. U. St. 24 = Hookeri.
canaliculata, Steffan. in Bull. Torrey Bot. Club, ix. (1882) 86 = Hookeri.

IMPERATA:—
caudata, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1832) 331 = exaltata.
condensata, Steud. Syn. Pl. Gram. 481 = arundinacea.
cylindrica, Beauv. Agrost. 165 = arundinacea.
cylindroides, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 289 = Miscanthus sacchariflorus. !
exaltata, *Brongn. in Duperr. Voy. Coq. Bot.* 101.—Malaya; N. Hebrid.; Am. trop.
filifolia, Nees, ex Steud. Syn. Pl. Gram. 405 = arundinacea.
Hookeri, *Rupr. ex Anderss. in Oefvers. Vet. Acad. Stockh.* (1855) 160.—Am. bor.; Mexic.
jaculatoria, Roem. & Schult. Syst. ii. 289 = exaltata.
Klaga, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 298 = Saccharum spontaneum.
Kornigii, Beauv. Agrost. 165 = arundinacea.
ovata, Tratt. Tabul. t. 569; ex Roem. & Schult. Syst ii. 401 = Lagurus ovatus.
pedicellata, Steud. in Flora, xxix. (1846) 22 = arundinacea.
ramosa, Anderss. in Oefvers. Vet. Akad. Stockh. (1855) 158 = exaltata.
sacchariflora, Anderss. ex Benth. in Journ. Linn. Soc. xix. (1881) 65 = Miscanthus sacchariflorus.
sacchariflora, Maxim. Prim. Fl. Amur. 331 = Miscanthus sacchariflorus.
Sagi, Anderss. in Oefvers. Vet. Akad. Stockh. (1855) 159 = brasiliensis.
Sara, Schult. Mant. ii. 466 = Saccharum Sara.
Sieberi, Opiz, Natural. x. (1832) 190 = arundinacea.
Sieca, Beauv. ex Steud. Nom. ed. II. i. 805 = arundinacea.
spontanea, Beauv. Agrost. 8 = Saccharum spontaneum.
Thunbergii, Beauv. l. c. 165 = arundinacea.
tinctoria, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 289 (Miscanthi sp.).—Japon.

IMPERATIA, Moench, Meth. 60 (1794) = **Tunica**, Scop. (Caryophyll.).
filiformis, Moench, l. c. = T. Saxifraga.

IMPERATORIA, [Tourn.] Linn. Syst. ed. I (1735) = **Peucedanum**, Tourn. (Umbellif.).
Angelica, Borck. ex Gaertn. Mey. & Scherb. Fl. Wett. i. 378 = Angelica sylvestris.
angustifolia, Bell. in Mém. Acad. Turin, x, xi. (1802–3) 449 = P. angustifolium.
aragonica, Pourr. in Willk. & Lange, Prod. Fl. Hisp. iii. 44 = P. hispanicum.
aromatica, Salisb. Prod. 167 = P. Ostruthium
caucasica, Spreng. Prod. Umbellif. 17 = P. caucasicum.
Chabraei, Bess. ex DC. Prod. iv. 178 = P. Schottii.
Chabraei, Spreng. Sp. Umbellif. 64 = P. carvifolium.
dahurica, D. Dietr. Syn. Pl. ii. 969 = P. terebinthaceum.
decursiva, C. A. Mey. ex Glehn in Ind. Sem. Hort. Petrop. (1868) 87 (= *Tommasinia Szowitsii*).—Orient.
decursiva, Szow. ex Boiss. in Ann. Sc. Nat. Sér. III. i (1842) 302 (= *Tommasinia purpurascens*).—Oriens.
flavescens, Bess. Prim. Fl. Galic. i. 213 = Angelica sylvestris.
glauca, Bartl. in Bartl. & Wendl. f. Beitr. 98 = P. Schottii.
hispanica, Boiss. Elench. 48 = P. hispanicum.
laevigata, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 328 = P. latifolium.
Lowei, Coss. in Bull. Soc. Bot. Fr. xv. (1868) 99.—Ins. Mader.
major, S. F. Gray, Nat. Arr. Brit. Pl. ii. 519 = P. Ostruthium.
mexicana, Hort. Chels. ex G. Don, Gen. Syst. iii. 886. —Mexic.
montana, DC. Fl. Fr. v. 504 = Angelica sylvestris.
nodiflora, Lam. Fl. Fr. 417 = Trochiscanthes nodiflorus.
ostruthium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 198 = sq.
Ostruthium, Linn. Sp. Pl. 359 = P. Ostruthium.
palustris, Bess. Fl. Galic. i. 214 = Angelica palustris.
pimpinelloides, Sehlm. ex Boenn. Prod. Fl. Monast. 91 = P. carvifolium.
Rauuli, Hort. Vindob. ex Schult. Syst. vi. 602 = Angelica Razulii.
Segueri, Spreng. Sp. Umbellif. 66 = Ligusticum Seguieri.
sylvestris, Lam. Fl. Fr. iii. 417 = Angelica sylvestris.

IMPERATORIA:—
trilobata, Gilib. Fl. Lituan. ii. 84 = Peucedanum Ostruthium.
triternata, Viv. ex DC. Prod. iv. 183 = P. Ostruthium.
verticillaris, DC. Fl. Fr. iv. 287 = P. verticillare.
verticillata, Lam. Encyc. iii. 243 = prasc. !

IMPERIALIS, Adans. Fam. ii. 48 (1763) = **Fritillaria**, Tourn. (Liliac.).
comosa, Moench, Meth. 305 = F. Imperialis.
coronata, Dum.-Cours. Bot. Cult. ed. II. ii. 196 = F. Imperialis.
dagana, Turcz. in Bull. Soc. Nat. Mosc. (1857) n. i. 62 = F. dagana.
leucantha, Pisch. ex R. Grah. in Edinb. N. Phil Journ. (Apr.–June 1831) 190 = F. verticillata.
superba, Mirb. Hist. Pl. ed. II. (1804) viii. 274 = F. Imperialis.

IMPIA, Bluff & Fingerh. Comp. Fl. Germ. ii. 342 (1825) = **Filago**, Tourn. (Compos.).
germanica, Bluff & Fingerh. l. c. = F. germanica.
pyramidata, Bluff & Fingerh. l. c. 343 = F. germanica.

INCARVILLAEA, Orb. Dict. vii. 28 (1849) = seq.

INCARVILLEA, Juss. Gen. 138 (1789). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1049.
arguta, Royle, Illustr. Bot. Himal. 296 = Amphicome arguta.
chinensis, Poir. in Dict. Sc. Nat. xxiii. 52 = sinensis.
chinensis, Spreng. ex DC. Prod. ix. 237 = Tecoma grandiflora.
compacta, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 521.—China.
diffusa, Royle, Illustr. Bot. Himal. t. 72 = Amphicome arguta.
dubia, Wall. Cat. n. 4877 = Lysionotus serrata.
Emodi, Wall. l. c. n. 4877 = Amphicome arguta, A. Emodi.
foliosa, Steud. Nom. ed. II. i. 805 = Crotalaria medicaginea.
grandiflora, Poir. in Dict. Sc. Nat. xxiii. 53 = Tecoma grandiflora.
grandiflora, Spreng. Syst. ii. 836 = Aeschynanthus grandiflora.
Koempanni, Hort. Germ.—Cf. Gard. Chron. (1880) ii. 725 = Olgan.
olivaeifolia, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 118 = Clitria acuminata.
Olgae, *Regel, Gartenfl.* (1880) 3; *Pl. Nov. Fedsch.* 62. —Kokania.
parasitica, Roxb. Hort. Beng. 47; Fl. Ind. iii. 112 = Aeschynanthus grandiflora
sinensis, *Lam. Encyc.* iii. 243.—China.
tomentosa, Roxb. ex DC. Prod. ix. 237 = Paulownia imperialis.
tomentosa, Spreng. Syst. ii. 836 = Bignonia mollis.

INDIGASTRUM, Jaub. & Spach, Illustr. v. 101. t. 492 (1857) = **Indigofera**, Linn. (Legumin.).
deflexum, Jaub. & Spach, l. c. = Indigofera parviflora.
macrostachyum, Jaub. & Spach, l. c.—Orient.

INDIGOFERA, Linn. Hort. Cliff. 487 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 494.
ACANTHONOTUS, Benth. in Hook. Niger Fl. 298 (1849).
AMICARPUS, Benth. in Lindl. Veg. Kingd. 554 (1847).
BRISSONIA, Desv. [Journ. Bot. i. (1814) 78] in Ann. Sc. Nat. Sér. I. ix. (1826) 411.
ELIRANTHUS, Hochst. in Flora, xxix. (1846) 593.
HEMISPADON, Endl. l. c. xv. (1832) ii. 885.
INDIGASTRUM, Jaub. & Spach, Illustr. v. 101. t. 492 (1857).
ONOBROPIS, G. Don, Gen. Syst. iii. 214 (1832).
SPHAERIDIOPHORUM, Desv. Journ. Bot. iii. (1813) 195. t. 6.
TRICOELENDUS, Raf. Fl. Tellur. ii. 97 (1836).
abyssinica, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 97 = amorphoides.
acanthocarpa, Lindl. in Mitch. Three Exped. ii. 17 = Glycyrrhiza pauciloides.
acutifolia, *Schlecht. in Linnaea*, xii. (1838) 283.—Mexic.
adenocarpa, *E. Mey. Comm. Pl. Afr. Austr.* 105.—Afr. austr.

INDIGOFERA:—
adenophylla, R. Grah. in Wall. Cat. n. 5462 = trifoliata.
adenostachyoides, *A. Gray, Bot. U. St. Expl. Exped.* i. 405.—Austral.
adenensis, E. Mey. Comm. Pl. Afr. Austr. 99 = pollotes.
adscendens, *Eckl. & Zeyh. Enum.* 236.—Afr. austr.
aeruginis, Schweinf. Beitr. Fl. Aethiop. 11 = trigonelloides.
affinis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 184.—Afr. austr.
agowensis, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 81 = viridiflora.
alata, Schweinf. Beitr. Fl. Aethiop. 11 = amorphoides.
alba, *A. Gousult, in Rev. Hortic.* Sér. IV. iii. (1854) 361. t. 19.—China.
albicans, *R. Br. in Salt. Abyss. App.* 65, nomen.—Abyss.
albicans, *Span. in Linnaea*, xv. (1841) 190.—Ins. Timor.
alopecuroides, DC. Prod. ii. 231 = coriacea
alpina, Ekkl. & Zeyh. Enum 236 = stipularia.
alta, *Schweinf. Beitr. Fl. Aethiop.* 11.—Abyss.
alternans, *DC. Prod.* ii. 229.—Afr. austr.
amoena, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 68.—Afr. austr.
amoena, E. Mey. Comm. Pl. Afr. Austr. 106 = intermedia.
amorphoides, *Jaub. & Spach, Illustr.* v. 93.—Afr. trop.; As. occ.
anabaptista, *Steud. Nom.* ed. II. i. 805.—Afr. bor.; As. occ.
ancepe, Vahl, ex Poir. Encyc. Suppl. iii. 147 = endecaphylla.
angolensis, *D. Dietr. Syn. Pl.* iv. 1036.—Afr. trop.
angulata, Lindl. Bot. Reg. t. 991 = australis.
angulata, *Rottl. ex Spreng. Syst.* iii. 277.—Ind. or.
angulosa, *Edgew. in Journ. As. Soc. Beng.* xxi. (1858) 168.—Ind. or.
angustata, E. Mey. Comm. Pl. Afr. Austr. 94 = stenophylla.
angustifolia. Blanco, Fl. Filip. ed. I. 596 = hirsuta.
angustifolia, Curt. Bot Mag. t. 466 = stricta
angustifolia, *Linn. Mant.* ii. 272.—Afr. austr.
angustifolia, Perr. ex DC. Prod. ii. 224 = Tephrosia linearia.
angustifolia, Thunb. Fl. Cap. ed. Schult. 599 = brachycalyx.
Anil, *Linn. Mant.* ii. 272.—Am. austr.
Anil, Sm. ex Wight & Arn. Prod. 202 = paucifolia.
ajhylla, Benth. Hort. Kew. Brit. 309; ex Link, Enum. Hort. Berol. ii. 251 = filifolia.
arabica, *Jaub. & Spach, Illustr.* v. 89. t. 479.—Arab.
arborea, Roxb. Hort. Beng. 57; Fl. Ind. iii. 381 = pulchella.
arborescens, *Zuccagni, in Rorm. Collect.* 151.—Hab.?
arcuata, Willd. Sp. Pl. iii. 1393 = trita.
arenaria, E. Mey. Comm. Pl. Afr. Austr. 107 = alternans.
arenaria, *A. Rich. Tent. Fl. Abyss.* i. 188.—Abyss.
argentea, Blanco, Fl. Filip. ed. II. 415 = tinctoria.
argentea, Buch.-Ham. ex Roxb. Hort. Beng. 57; Fl. Ind. iii. 374 = pauciloba.
argentea, Burm. f. Fl. Ind. 171 = semitrijuga.
argentea, *Linn. Mant.* ii. 278.—Afr. bor.; As. occ.
argentea, Wall. Cat. n. 5455, partim = trita.
argyrea, *Eckl. & Zeyh. Enum.* 239.—Afr. austr.
argyroides, *E. Mey. Comm. Pl. Afr. Austr.* 106.—Afr. austr.
aristata, *Sperng. Syst.* iii. 278.—Afr. austr.
armata, Wall. Cat. n. 5458 = trita.
arrecta, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 188.—Afr. austr.
arrecta, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 184.—Abyss.
articulata, Gouan, Illustr. 49 = argentea.
ascendens, Walp. in Linnaea, xiii. (1839) 520 = adscendens.
aspalathifolia, Roxb. Hort. Beng. [98]; Fl. Ind. iii. 371 = aspalathoides.
aspalathoides, *Vahl, ex DC. Prod.* ii. 231.—Ind. or.
aspera, *Perr. ex DC. l. c.* 230.—Afr. trop.
asperifolia, *Bong. ex Benth. in Ann. Nat. Hist.* Ser. I. iii. (1839) 431.—Am. austr.
asperifolia, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 79 = trigonelloides.

INDIGOFERA :—
astragalina, *DC. Prod.* ii. 228.—Afr. trop.
athrophylla, Eckl. & Zeyh. Enum. 325 = denudata.
atricapa, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 190. —Afr. trop.
atropurpurea, *Buch.-Ham. ex Roxb. Hort. Beng.* 57; *Fl. Ind.* iii. 881.—Reg. Himal.; China.
auricoma, *E. Mey. Comm. Pl. Afr. Austr.* 107.—Afr. austr.
australis, *Willd. Sp. Pl.* iii. 1235.—Austral.
axillaris, E. Mey. in Linnaea, vii. (1832) 168 = Amphithalea virgata.
Bailleyi, *F. Muell. Fragm.* ix. 48.—Austral.
Baineii, *Baker*, in *Oliver, Fl. Trop. Afr.* ii. 71.—Afr. trop.
barbata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 410 = hirsuta.
Baukeana, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 230.—Afr. trop.
benguelensis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 87.—Afr. trop.
Benthamiana, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 219.—China.
Bernieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 999. —Madag.
Berteriana, *Spreng. Syst.* iii. 277.—Ins. Guadalup.
Bertolonii, Stend. Nom. ed. II. i. 807 = lateritia, *Bertol.*
biflora, *Roth, Nov. Pl. Sp.* 258.—Ind. or.
bifrons, *E. Mey. Comm. Pl. Afr. Austr.* 97.—Afr. austr.
bijuga, *Vog. ex Walp. in Linnaea*, xiii. (1839) 523.—Afr. austr.
bilabiata, *Loisel, ex Drapiez, Herb. Amat. Fl.* vi. (1838) t. 401.—Hab.?
Binderi, *Kotschy, in Sitzb. Acad. Wien, Math.-Nat.* li. *Abth.* ii. (1865) 364. t. 6 B.—Afr. trop.
Blanchetiana, *Benth. in Mart. Fl. Bras.* xv. i. 40.—Bras.
Boiviniana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 998.—Madag.
Bojeri, *Baker, in Journ. Linn. Soc.* xviii. (1881) 266. —Madag.
Bojeri, *Vatke, in Bremen, Abh.* vii. (1882) 245.—Madag.
bongensis, *Kotschy & Peyr. Pl. Tinn.* 8. n. 11. t. 4.—Afr. trop.
brachycarpa, R. Grah. in Wall. Cat. n. 5470 = argentea.
brachypoda, Stend. ex A. Rich. Tent. Fl. Abyss. i. 179 = stenophylla.
brachystachya, *E. Mey. in Linnaea*, vii. (1832) 168.—Afr. austr.
bracteata, *R. Grah. in Wall. Cat.* n. 5477.—Reg. Himal.
bracteola, *DC. Prod.* ii. 228.—Afr. trop.
Brandi, *Baker, in Oliver, Fl. Trop. Afr.* ii. 76.—Afr. trop.
brevidens, *Benth. in Mitch. Journ. Trop. Austral.* 385.—Austral.
Brunoniana, *R. Grah. in Wall. Cat.* n. 5491; *et in Pl. As. Rar.* iii. 48.—Burma.
bufalina, *Lour. Fl. Cochinch.* 458.—Cochinch.
Bungeana, *Walp. in Linnaea*, xiii. (1839) 523.—China.
Burchellii, *DC. Prod.* ii. 231.—Afr. austr.
Burchellii, E. Mey. Comm. Pl. Afr. Austr. 106 = argyraea.
Burkeana, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 197. —Afr. austr.
Burmanni, Boiss. Fl. Orient. ii. 187 = semitrijuga.
caerulea, Roxb. Hort. Beng. 57; Fl. Ind. iii. 877 = argentea.
cassia, Zipp. ex Span. in Linnaea, xv. (1841) 190 = timorensis.
caespitosa, Wight, in Wall. Cat. n. 5447 = enneaphylla.
calocaeura, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 229.—Burma.
cana, E. Mey. Comm. Pl. Afr. Austr. 104 = humifusa.
Canerori, *Baker, in Journ. Linn. Soc.* xv. (1876) 93. —Afr. trop.
campestris, *Bong. ex Benth. in Mart. Fl. Bras.* xv. i. 36.—Bras.
candicans, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 67. —Afr. austr.
candicans, E. Mey. Comm. Pl. Afr. Austr. 95 = heterophylla.
candicans, Sieber, ex DC. Prod. ii. 231 = filiformis.
Candolleana, *Meisn. in Hook. Lond. Journ. Bot.* ii. (1843) 35.—Afr. austr.
canescens, Lam. Encyc. iii. 251 = trita.
canescens, Spreng. Syst. iii. 274 = trifoliata.
capillaris, *Thunb. Prod. Pl. Cap.* 133.—Afr. austr.
capitata, R. Grah. in Wall. Cat. n. 5490 = Crotalaria medicaginea.
capitata, *Kotschy, in Sitzb. Acad. Wien, Math.-Nat. li. Abth.* ii. (1865) 365.—Afr. trop.
cardiophylla, *Harv. in Harv. & Sond. Fl. Cap.* ii. 175. —Afr. austr.
caroliniana, *Mill. Gard. Dict.* ed. VIII. n. 3 (Quid?).
carolina, Ind. hort.
caroliniana, *Walt. Fl. Carol.* 187.—Am. bor.
cassioides, Rottl. ex DC. Prod. ii. 225 = pulchella.
celebica, *Miq. Fl. Ind. Bat.* i. 1. 211.—Ins. Celebes
centrota, Eckl. & Zeyh. Enum. 238 = denudata.
Chamissoniana, D. Dietr. Syn. Pl. iv. 1041 = I. Anil.
chinensis, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843).—China.
cinerascens, DC. Prod. ii. 226 = argentea.
cinerascens, Eckl. & Zeyh. Enum. 241 = Zeyheri.
cinerea, Buckl. in. Proc. Acad. Sc. Philad. 1861 (1862) 451 = leptosepala.
cinerea, Willd. Sp. Pl. iii. 1235 = trita.
cinericolor, *F. Muell. Fragm.* v. 138.—Austral.
circinnata, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 202. —Afr. austr.
circinnata, *Roxb. ex Steud. Nom.* ed. II. i. 806 (Quid?). —Ind. or.
clitoriotides, *G. Don, Gen. Syst.* ii. 206.—Afr. trop.
coccinea, Broegelm. Beschr. Vorneegl. Pfl. 116 = Swainsonia galegifolia.
cocrinea, Lour. Fl. Cochinch. 457 = trita.
collina, Eckl. & Zeyh. Enum. 239 = argyraea.
colorata, Roxb. ex Wight & Arn. Prod. 202 = pauciflora.
coluteifolia, Jaub. & Spach, Illustr. v. 96. t. 487, 488 = emarginella.
complanata, *Reichb. ex Spreng. Syst.* iv. *Cur. Post.* 235. —Afr. austr.
complicata, *Eckl. & Zeyh. Enum.* 235.—Afr. austr.
compressa, *Lam. Encyc.* iii. 246.—Ins. Mascar.
concava, *Harv. in Harv. & Sond. Fl. Cap.* ii. 186.—Afr. austr.
concinna, *Baker, in Oliver, Fl. Trop. Afr.* ii. 80.—Afr. trop.
argentea, R. Grah. in Wall. Cat. n. 5471 = trifoliata.
congesta, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 70.—Afr. trop.
congoensis, *Baker, l. c.* 75.—Afr. trop.
consanguinea, Klotzsch, in Peters, Reise Mossamb. Bot. 50 = viscosa.
cordifolia, *Heyne, ex Roth, Nov. Pl. Sp.* 357.—Geront. trop.
coriacea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 68.—Afr. austr.
Cornevenlo, [Moç. & Sesse] ex DC. Prod. ii. 225 = I. Anil.
corniculata, *E. Mey. Comm. Pl. Afr. Austr.* 101.—Afr. austr.
corollillifolia, *A. Cunn. ex Benth. Fl. Austral.* ii. 201. —Austral.
corollillioides, *Taub. & Spach, Illustr.* v. 95. t. 487, 488 = Alysmin.
corollillioides, *Hurt. & Gal. in Bull. Acad. Brux.* x. (1843) 45.—Mexic.
costata, *Guill. & Perr. Fl. Senag. Tent.* i. 187.—Afr. trop.
costaricensis, *Benth. in Kjoeb. Vidensk. Meddel.* (1858) 5.—Costa Rica.
crassifolia, *Stend. Nom.* ed. II. i. 806.—Arab.
crotalarioides, *Baker, in Oliver, Fl. Trop. Afr.* ii. 85. —Afr. trop.
cryptoscelis, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 196. —Afr. austr.
cuneata, *Baker, in Journ. Linn. Soc.* xv. (1876) 93.—Afr. trop.
cuneifolia, *Eckl. & Zeyh. Enum.* 234.—Afr. austr.
cylindracea, *R. Grah. in Wall. Cat.* n. 5482.—Reg. Himal.
cylindrica, *DC. Prod.* ii. 226.—Afr. austr.
cytisoides, *Linn. Mant.* ii. 271, in nota; *Thunb. Prod. Pl. Cap.* 133.—Afr. austr.
daleoides, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 200. —Afr. austr.
decalbata, *Harv. l. c.* 170.—Afr. austr.
debilis, R. Grah. in Wall. Cat. n. 5466 = endecaphylla.
declinata, *E. Mey. Comm. Pl. Afr. Austr.* 104.—Afr. austr.
decora, *Lindl. in Journ. Hort. Soc.* i. (1846) 68.—China.
decumbens, *Hill, Hort. Kew.* 295.—China.
defexa, Hookst. ex A. Rich. Tent. Fl. Abyss. i. 178 = parviflora.
dendroides, *Zucc. in Abh. Akad. Muench.* iii. t. 571; *Coll.* ii. 367.—Afr. trop.
densiflora, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 44.—Mexic.
denudata, *Linn. f. Suppl.* 335; *Jacq. Hort. Schoenb.* ii. 57. t. 231.—Afr. austr.
denudata, *Thunb. Prod. Pl. Cap.* 132.—Afr. austr.
depressa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 175.—Afr. austr.
depressa, *Thunb. Prod. Fl. Cap.* 132.—Afr. austr.
desmodioides, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 158 = Pycnospora hedysaroides.
diffusa, R. Br. in Salt, Abyss. App. 65, nomen = Salitana.
diffusa, *Desv. Journ. Bot.* iii. (1814) 79.—Ins. Afr.
digitata, Eckl. & Zeyh. Enum. 239 = strigosa.
digitata, *Linn. f. Suppl.* 335; *Lam. Encyc.* iii. 251; *Thunb. Prod. Fl. Cap.* 133.—Afr. austr.
dillwynioides, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 178.—Afr. austr.
dimidiata, *Vog. ex Walp. in Linnaea*, xiii. (1839) 521.—Afr. austr.
diphylla, *Vent. Choix*, 800. t. 80; *Pers. Syn.* ii. 325.—Afr. occ.
discolor, E. Mey. Comm. Pl. Afr. Austr. 97 = procumbens.
disperma, *Linn. Syst. Nat.* ed. XII. iii. 232.—Ind. or.
distichora, *Baker, in Journ. Linn. Soc.* xv. (1876) 93. —Afr. trop.
distichia, *Eckl. & Zeyh. Enum.* 243.—Afr. austr.
divaricata, *Jacq. Hort. Schoenb.* iii. 61. t. 365.—Hab.?
diversifolia, *DC. Prod.* ii. 228.—Ins. Mascar.
dodecaphylla, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 18.—Afr. trop.
dominguensis, Spreng. ex DC. Prod. ii. 227 = microcarpa.
Dorycnium, Fenzl, in Flora, xxvii. (1844) 312 = pulchra.
Dossa, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 244.—Reg. Himal.
Dossa, Wall. Cat. n. 5481 = Gerardiana.
Dregeana, *E. Mey. Comm. Pl. Afr. Austr.* 100.—Afr. austr.
dubia, Stend. Nom. ed. II. i. 806 = elliptica.
dumosa, E. Mey. Comm. Pl. Afr. Austr. 93 = denudata.
echinata, *Willd. Sp. Pl.* iii. 1222.—Ind. or.
effusa, E. Mey. Comm. Pl. Afr. Austr. 107 = alternans.
efoliata, *F. Muell. in Melb. Chemist*, iii. (Nov. 1883) —Austral.
Ehrenbergiana, *Stend. Nom.* ed. II. i. 806.—Mexic.
elatior, *Carr. in Rev. Hortic.* (1866) 270.—Asa. trop.?
elegans, *Schum. & Thonn. Beskr. Guin. Pl.* 368.—Afr. trop.
elliptica, *E. Mey. Comm. Pl. Afr. Austr.* 99.—Afr. austr.
elliptica, Roxb. Hort. Beng. 57; Fl. Ind. iii. 880 = pulchella.
elongata, *G. Don, Gen. Syst.* ii. 206.—Afr. trop.
elongata, *Mickell, in Kjoeb. Vidensk. Meddel.* (1875) 64.—Bras.
emarginata, *G. Don, Gen. Syst.* ii. 209.—Afr. trop.
emarginata, Perr. in Ann. Mus. Hist. (1830); ex Guill. & Perr. Fl. Senag. Tent. i. 178 = tinctoria.
emarginella, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 184. —Abyss.
endecaphylla, *Jacq. Ic. Rar.* t. 570.—Geront. trop.
enneaphylla, Eckl. & Zeyh. Enum. 244 = alternans.
enneaphylla, *Linn. Mant.* ii. 272.—Asa. et Austral. trop.
enneaneis, E. Mey. Comm. Pl. Afr. Austr. 105 = disticha.
erecta, Eckl. & Zeyh. Enum. 239 = gracilis.
erecta, *Thunb. Prod. Pl. Cap.* 132.—Afr. austr.
erecta, Willd. ex Steud. Nom. ed. II. i. 806 = filiformis.
ericocarpa, *E. Mey. Comm. Pl. Afr. Austr.* 103.—Afr. austr.
erucoides, Meism. in Lehm. Pl. Preiss. i. 86 = australis.
erucoides, *A. Rich. Tent. Fl. Abyss.* i. 179.—Abyss.
erythrostoba, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 86 = paucifolia.

INDIGOFERA :—
ecythrogramma, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 78.—Afr. trop.
ecoslea, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 45.—Mexic.
exigua, Eckl. & Zeyh. Enum. 244 = alternans.
exigua, *E. Mey. Comm. Pl. Afr. Austr.* 108.—Afr. austr.
falcata, *E. Mey. l. c.* 98.—Afr. austr.
fastigiata, *E. Mey. l. c.* 102.—Afr. austr.
ferruginea, Schum. & Thonn. Beskr. Guin. Pl. 279 = hirsuta.
filicaulis, *Eckl. & Zeyh. Enum.* 239.—Afr. austr.
filifolia, *Thunb. Prod. Pl. Cap.* 132.—Afr. austr.
filiformis, *Thunb. l. c.* 133.—Afr. austr.
filipes, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 198.—Afr. austr.
Findlayensiana, Wall. Cat. n. 5488 = galegoides.
flabellata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 180.—Afr. austr.
flaccida, Koen. ex Roxb. Hort. Beng. [98]; Fl. Ind. iii. 375 = subulata.
flavicana, *Baker. in Oliver, Fl. Trop. Afr.* ii. 78.—Afr. trop.
flexuosa, Eckl. & Zeyh. Enum. 232 = denudata.
flexuosa, R. Grah. in Wall. Cat. n. 5473 = Tephrosia purpurea.
floribunda, Hort.—Cf. Gard. Chron. (1851) 11. 836 = Gerrardiana.
florida, E. Mey. Comm. Pl. Afr. Austr. 96 = cuneifolia.
foliolosa, R. Grah. in Wall. Cat. n. 5485 = Wightii.
follosa, *E. Mey. Comm. Pl. Afr. Austr.* 102.—Afr. austr.
follosa, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 217 = Crotalaria follosa.
fragrans, Retz. Obs. iv. 29 = pentaphylla.
frumentacea, Roxb. ex Wight & Arn. Prod. 196 = glandulosa.
frutescens, *Linn. f. Suppl.* 334 ; *Thunb. Prod. Pl. Cap.* 133.—Afr. austr.
frutticosa, Harv. & Sond. Fl. Cap. ii. 611 (in Ind.) = coriacea.
fruticulosa, Walp. in Linnæa, xiii. (1839) 536 = exigua.
fulcrata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 192.—Afr. austr.
fulgens, *Baker, in Oliver, Fl. Trop. Afr.* ii. 101.—Afr. trop.
fusca, G. Don, Gen. Syst. ii. 211 = hirsuta.
galegoides, *DC. Prod.* ii. 225.—As. trop. ; Malaya.
Garckeana, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 321.—Afr. trop.
geminata, *Baker, in Oliver, Fl. Trop. Afr.* ii. 81.—Afr. trop.
Gerardiana, *R. Grah. in Wall. Cat.* n. 5486.—Reg. Himal.
Gerrardiana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 595.—Afr. austr.
Gibsonii, J. Grah. Cat. Bomb. Pl. 46 = pulchella.
glabra, Linn. Sp. Pl. 751 = pentaphylla.
glandulifera, *Pagv, ex Steud. Nom.* ed. II. i. 807 (Quid ?).—Afr. austr.
glandulosa, Roxb. Hort. Beng. 57 ; Fl. Ind. iii. 372 = trifoliata.
glandulosa, Wendl. Bot. Beobacht. 55 = seq. ?
glandulosa, *Willd. Sp. Pl.* iii. 1227.—As. et Austral. trop.
glauca, Lam. Encyc. iii. 246 = argentea.
glauca, Perr. ex DC. Prod. ii. 228 = diphylla.
glaucescens, *Eckl. & Zeyh. Enum.* 242.—Afr. austr.
glaucovorea, R. Grah. in Wall. Cat. n. 5484 = pulchella.
glomerata, *E. Mey. in Linnæa*, vii. (1832) 166.—Afr. austr.
glutinosa, Perr. ex DC. Prod. ii. 227 = viscosa.
gnidioides, Vahl, ex DC. l. c. = secundiflora.
gonioides, *Hochst. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 86.—Afr. trop.
grandiflora, *Bong. ex Benth. in Ann. Nat. Hist.* iii. (1839) 481.—Bras.
grandiflora, *Spreng. Syst. iv. Cur. Post.* 285.—Afr. austr.
Grahamiana, Steud. Nom. ed. II. i. 807 = I. Dosua.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 997.—Madag.
grandifolia, *Carr. in Rev. Hortic. Sér.* IV. viii. (1866) 668.—Ind. or.

INDIGOFERA :—
grata, *E. Mey. Comm. Pl. Afr. Austr.* 108.—Afr. austr.
grevenolea, Schrad. Sert. Hannov. 20 = viscosa.
grisea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 80.—Afr. trop.
grisea, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 410 = lespedezioides.
Guatimala, Lunan, Hort. Jamaic. i. 220 = brachycarpa.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 267 = pilosa.
haitensis, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 410.—Ins. S. Doming.
Hamiltonii, R. Grah. in Wall. Cat. n. 5465 = atropurpurea.
haplophylla, *F. Muell. Fragm.* iii. 101.—Austral.
heliantala, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 101.—Reg. Himal.
hedranophylla, *Eckl. & Zeyh. Enum.* 232.—Afr. austr.
hedysantha, *Eckl. & Zeyh. l. c.* 243.—Afr. austr.
hedysaroides, Lam. Encyc. iii. 250 = trita.
headecapylla, Jacq. Coll. ii. 358 = endecaphylla.
heterantha, Wall. Cat. n. 5480 A, B, C, partim = Gerardiana.
heterantha, Wall. l. c. n. 5480 C, partim = I. Dosua
heterocarpa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 90.—Afr. trop.
heterophylla, *Presl, Bot. Bemerk.* 55.—Mexic.
heterophylla, Roxb. ex Wight & Arn. Prod. 200, 201 = pentaphylla, pentaphylla.
heterophylla, *Thunb. Prod. Pl. Cap.* 133.—Afr. austr.
heterotricha, *DC. Prod.* ii. 227.—Afr. austr.
Hildebrandtii, *Benth. ex Oliver, Fl. Trop. Afr.* ii. 85.—Afr. trop.
hilaria, *Eckl. & Zeyh. Enum.* 241.—Afr. austr.
hippocrepioides, Schlecht. in Linnæa, xii. (1838) 283 = Ionscacia angustifolia.
hirsuta, *Harv. in Harv. & Sond. Fl. Cap.* ii. 194.—Afr. austr.
hirsuta, Jacq. Ic. Rar. t. 859 = lateritia.
hirsuta, *Linn. Sp. Pl.* 751.—Geront. trop.
hirta, *Boj. Hort. Maurit.* 91.—Madag.
hirta, *E. Mey. Comm. Pl. Afr. Austr.* 101.—Afr. austr.
hispida, *Eckl. & Zeyh. Enum.* 240.—Afr. austr.
Hochstetteri, Baker, in Oliver, Fl. Trop. Afr. ii. 101 = malacostachya.
holoteuca, Benth. ex Harv. & Sond. Fl. Cap. ii. 200 = alternans.
Hookeriana, *Meisen. in Hook. Lond. Journ. Bot.* ii. (1843) 85.—Afr. austr.
Hoser, Forsk. Fl. Aegypt. Arab. 137 = lateritia.
Humboldtiana, *Spreng. Syst.* iii. 276.—Mexic.
humifusa, *Eckl. & Zeyh. Enum.* 244.—Afr. austr.
humilis, *H. B. & K. Nov. Gen. et Sp.* vi. 454.—Peruv.
humilis, *Zipp. ex Span. in Linnæa*, xv. (1841) 190, mantissa.
inamoena, Thw. Enum. Pl. Zeyl. 88 = Wightii.
incana, *Thunb. Prod. Pl. Cap.* 132.—Afr. austr.
indica, Lam. Encyc. iii. 245 = tinctoria.
indica, Mill. Gard. Dict. ed. VIII. n. 4 = hirsuta.
inhambanensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 48.—Afr. trop.
inquinans, *Willd. Sp. Pl.* iii. 1336.—Ins. S. Doming.
intermedia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 190.—Afr. austr.
intricata, *Boiss. Fl. Orient.* ii. 196.—Persia.
jamaicensis, *Spreng. Syst.* iii. 277.—Jamaic.
Jirahaiba, Buch.-Ham. in Wall. Cat. n. 5464 = pulchella.
juncea, Dec. Herb. Amat. t. 237 = filifolia.
Kirkii, *Oliver, in Hook. Ic. Pl.* t. 1416.—Afr. trop.
Klotzschii, Wight & Arn. Prod. 206 = endecaphylla.
Kotschyana, *Kotschy, in Sitzb. Acad. Wiss, Math.-Nat.* l. *Abth.* i. (1865) 363.—Afr. trop.
Kraussiana, *Meisen. in Hook. Lond. Journ. Bot.* ii. (1843) 83.—Afr. austr.
Lagascana, *DC. Prod.* ii. 229.—Mexic.
lasiantha, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 410.—Afr. austr.
lasiantha, F. Muell. in Rep. Gregory's Exped. 8 = brevidens.
lateritia, *Bertol. in Nov. Comm. Bonon.* ii. (1836) 218.—Afr. trop.
lateritia, Willd. Sp. Pl. iii. 1338 = viscosa.

INDIGOFERA :—
latifolia, *Michels, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. vii. 13.—Paraguay.
leptocarpa, *Eckl. & Zeyh. Enum.* 236.—Afr. austr.
leptocarpa, *Hochst. & Steud. ex Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 510.—Arab. ; Afr. trop.
leptoculis, Eckl. & Zeyh. Enum. 248 = angustifolia.
leptophylla, E. Mey. Comm. Pl. Afr. Austr. 99 = Zeyheri.
leptosepala, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 298.—Texas.
leptostachya, *DC. Prod.* ii. 225.—Reg. Himal.
Leschenaultii, DC. l. c. 228 = trita.
lespedezioides, *H. B. & K. Nov. Gen. et Sp.* vi. 457.—Mexic.
leucoclada, *Baker, in Journ. Linn Soc.* xviii. (1881) 267.—Madag.
Lindheimeriana, *Scheele, in Linnæa*, xxi. (1848) 464.—Am. bor. occ.
Lindleyana, *Spreng. ex Steud. Nom.* ed. II. i. 807.—Austral.
linearis, *DC. Prod.* ii. 228.—Afr. trop.
linearis, Guill. & Perr. Fl. Senceg. Tent. i. 184 = parviflora.
linifolia, *Retz. Obs.* iv. 29 ; vi. 33. t. 2.—Geront. trop.
longeracemosa, *Boiv. ex Baill. in Bull. Soc. Linn. Par.* i. (1882) 899.—Madag.
lotoides, *Lam. Encyc.* iii. 247.—Afr. austr.
lotoides, E. Mey. in Linnæa, vii. (1832) 168 = hispida.
lotoides, Schlecht. l. c. xii. (1838) 282 = Ehrenbergiana.
Lyalli, *Baker, in Journ. Linn. Soc.* xx. (1883) 128.—Madag.
machærocarpa, Fenzl, ex Baker, in Oliver, Fl. Trop. Afr. ii. 83 = parviflora.
macra, *E. Mey. Comm. Pl. Afr. Austr.* 105.—Afr. austr.
macrocalyx, *Guill. & Perr. Fl. Senceg. Tent.* i. 175. t. 46.—Afr. trop.
macrocarpa, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 409.—Peruv.
macrocarpa, *Lepr. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 73.—Afr. trop.
macrophylla, *Schum. & Thonn. Beskr. Guin. Pl.* 272.—Afr. trop.
macrostachya, Willd. ex Spreng. Syst. iii. 237 = Dalea Mutisii.
macrostachys, *Vent. Jard. Malm.* t. et p. 44.—China.
madagascariensis, *Schrank, ex Colla, Hort. Ripul. App.* ii. 250.—Madag.
madagascariensis, *Vatke, in Bremen, Abh.* vii. (1882) 345.—Madag.
malacostachya, *Benth. in Harv & Sond. Fl. Cap.* ii. 184.—Afr. austr.
marginata, Walp. Rep. i. 661, sphalm. = *emarginata*.
marginata, *R. Grah. in Wall. Cat.* n. 5467.—Ind. or.
marttima, *Baker, in Oliver, Fl. Trop. Afr.* ii. 89.—Afr. trop.
marmorata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 510.—Ins. Socotr.
mauritanica, Thunb. Prod. Pl. Cap. 133 = alopecuroides, coriacea.
medicaginea, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 86.—Afr. trop.
melanadenia, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 195.—Afr. austr.
melanotricha, Steud. ex A. Rich. Tent. Fl. Abyss. i. 181 = secundiflora.
mellitoides, Hance, in Journ. Bot. xvi. (1869) 168 = Astragalus tenuis.
melolobioides, Benth. ex Harv. & Sond. Fl. Cap. ii. 201 = patens.
mexicana, *Benth. Pl. Hartw.* 286.—Mexic.
mexicana, Linn. f. Suppl. 335 = Psoralea Mutisii.
Meyeriana, Eckl. & Zeyh. Enum. 235 = bifrons.
micrantha, Bunge, Enum. Pl. Chin. Bor. 16 = Bungeana.
micrantha, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 410 = I. Anil.
microcalyx, *E. Mey. Comm. Pl. Afr. Austr.* 104.—Afr. austr.
microcarpa, *Desv. Journ. Bot.* iii. (1814) 79.—Bras.
microphylla, Lam. Encyc. iii. 250 = sarmentosa.
microps, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 400.—Madag.
microstachya, *Presl, Bot. Bemerk.* 54.—Ind. or.
Mimosella, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 408.—Madag.

INDIGOFERA :—
mimosoides, *Baker, in Oliver, Fl. Trop. Afr.* ii. 98.—Afr. trop.
minuta, *Ortog. Hort. Madr. Dec.* 98.—Cuba.
minutifora, *Walp. Rep.* i. 666.—Afr. austr.
mollis, *Eckl. & Zeyh. Enum.* 237.—Afr. austr.
molucana, Dc. Prod. ii. 229 = trifoliata.
monophylla, *DC. l. c.* 232.—Austral.
monostachya, *Eckl. & Zeyh. Enum.* 257.—Afr. austr.
mouroundavensis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 897.—Madag.
mucronata, Lam. Encyc. iii. 247 = cytisoides.
mucronata, Spreng. ex DC. Prod. ii. 227 = subulata.
mucronata, Willd. ex Spreng. Syst. iii. 276 = Hamboldtiana.
multicaulis, DC. Prod. ii. 223 = trifoliata.
multijuga, *Baker, in Oliver, Fl. Trop. Afr.* ii. 83.—Afr. trop.
Mundiana, *Eckl. & Zeyh. Enum.* 243.—Afr. austr.
Mutivii, Spreng. Syst. iii. 274 = Psoralea Mutisii.
myosoroisis, *Rottl. ex DC. Prod.* ii. 222.—Ind. or.
nana, Eckl. & Zeyh. Enum. 242 = Zeyheri.
nephrocarpa, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1883) 509.—Ins. Socotr.
nigricans, *Vahl, ex Pers. Syn.* ii. 327.—Afr. trop.
nigritana, *Hook. f. in Hook. Niger Fl.* 294.—Afr. trop.
nigromontana, Eckl. & Zeyh. Enum. 233 = denudata.
nivea, E. Mey. Comm. Pl. Afr. Austr. 98 = glomerata.
nives, *Willd. ex Spreng. Syst.* iii. 273.—Afr. austr.
nuda, G. Don, in Lond. Hort. Brit. 301; et Gen. Syst. ii. 209 = filifolia ?
nudicaulis, *E. Mey. Comm. Pl. Afr. Austr.* 93.—Afr. austr.
nummularia, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 74.—Afr. trop.
obcordata, *Eckl. & Zeyh. Enum.* 232.—Afr. austr.
oblongifolia, *Forsk. Fl. Aegypt. Arab.* 137.—Arab.
obrajillensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 404. —Peruv.
oligophylla, *Klotzsch, in Peters, Reise Mossamb. Bot.* 49.—Afr. trop.
oligophylla, Lam. ex Baker, Fl. Maurit. 70 = tinctoria.
oligosperma, DC. Prod. ii. 228 = secundiflora.
onobrychioides, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 898.—Madag.
orixensis, Roxb. ex Wight & Arn. Prod. 201 = trifoliata.
ornithopodioides, *Cham. & Schlecht. in Linnaea*, v. (1830) 577.—Mexic.
ornithopodioides, Hochst. & Steud. ex Jaub. & Spach, Illustr. v. t. 480 = anabaptista.
ornithopodioides, Roxb. ex Wall. Cat. sub n. 5455 = trita.
ornithopodioides, Schum. & Thonn. Beskr. Guin. Pl. 373 = tinctoria.
oroboides, E. Mey. Comm. Pl. Afr. Austr. 94 = monostachya.
orthocarpa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 99.—Afr. trop.
orthocarpa, *Berg & Vathe, Ind. Sem. Hort. Berol.* (1876) *App.* 8.—Hab. ?
orthocarpa, *Presl, Bot. Bemerk.* 56.—Am. centr.
owata, *Thunb. Prod. Pl. Cap.* 135.—Afr. austr.
ovina, *Harv. in Harv. & Sond. Fl. Cap.* ii. 189.—Afr. austr.
oxalides, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 86.—Afr. trop.
oxyorpa, *Denc. Journ. Bot.* iii. (1814) 79.—Ind. occ.
oxycarpa, F. Muell. Fragm. iii. 103 = parviflora.
Oxytropis, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 193. —Afr. austr.
paludosa, Lepr. ex Guill. & Perr. Fl. Seneg. Tent. i. 186 = endecaphylla.
palustris, *Vahl, in Oestr. Bot. Zeitschr.* xxvii. (1878) 201.—Afr. trop.
pasiculata, *Pers. Syn.* ii. 325.—Afr. trop.
Parkeri, *Baker, in Journ. Linn. Soc. xx.* (1883) 126.—Madag.
parviflora, *Heyne, in Wall. Cat.* n. 5457.—Geront. trop.
parvula, *Delile, in Caill. Voy. Meroë*, 2. t. 84.—Afr. trop.
pascorum, *Benth. in Ann. Nat. Hist. Ser.* I. iii. (1839) 481.—Am. occ.
patens, *Eckl. & Zeyh. Enum.* 236.—Afr. austr.
pauciflora, *Eckl. & Zeyh. l. c.* 244.—Afr. austr.

INDIGOFERA :—
pauciflora, Heyne, ex Wight & Arn. Prod. Index, sphalm. = parviflora.
pauciflora, E. Mey. Comm. Pl. Afr. Austr. 99 = stricta.
pauciflolia, *Delile, Fl. Egypte*, 251.—As. et Afr. trop.
pauciflolia, Wight, in Wall. Cat. n. 5454 = prasc.
pectinata, *Baker, in Journ. Linn. Soc.* xx. (1883) 127. —Madag.
pedicellata, *Wight & Arn. Prod.* 200.—Ind. or.
pedunculata, *Hilarnb. & Bgl. ex Baker, in Journ. Bot.* xx. (1882) 67 —Madag.
pentaphylla, *Burch. ex Harv. & Sond. Fl. Cap.* ii. 178. —Afr. austr.
pentaphylla, *Murr. Syst. Veg.* ed. XIII. 564.—As. et Afr. trop.
peregrina, DC. Prod. ii. 224 = trifoliata.
Perrinana, *Spreng. Neue Entdeck.* ii. 161.—Am. austr.
Perrottetii, *DC. Prod.* ii. 228.—Afr. trop.
phyllanthoides, *Baker, in Oliver, Fl. Trop. Afr.* ii. 96. —Afr. trop.
pilosa, *Poir. Encyc. Suppl.* iii. 151.—Afr. trop.
pinifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 127. —Madag.
platyphylla, Miq. Fl. Ind. Bat. i. 1. 292 (= *Kiesera gracilis*).—Java.
platyphylla, E. Mey. Comm. Pl. Afr. Austr. 95 = complicata.
plumosa, *Spreng. Syst.* iii. 278.—Hab. ?
podophylla, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 168. —Afr. austr.
Pohliana, Benth. in Ann. Nat. Hist. Ser. I. iii. (1839) 481 = lespedecioides.
polioten, *Eckl. & Zeyh. Enum.* 242.—Afr. austr.
polycarpa, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 191. —Afr. austr.
polycarpa, *Willd. ex Spreng. Syst.* iii. 276.—Am. austr.
polygonoides, *Wendl. Bot. Beobacht.* 55.—Hab. ?
polyphylla, DC. Prod. ii. 227 = Gerardiana.
polyphylla, Heyne, ex Wight & Arn. Prod. 202 = myseroisis.
polyphylla, Rottl. ex Wight & Arn. l. c. = Wightii.
porrecta, *Eckl. & Zeyh. Enum.* 234.—Afr. austr.
praecana, *F. Muell. Rep. Burdek. Exp.* 10.—Austral.
Prieureana, *Guill. & Perr. Fl. Senag. Tent.* i. 187.—Afr. trop.
procera, *Schum. & Thonn. Beskr. Guin. Pl.* 365.—Afr. trop.
procumbens, Hort. ex Kranz, in Flora, xxvii. (1844) 555 = incana.
procumbens, *Linn. Mant.* ii. 271.—Afr. austr.
prostrata, Klein, ex Wight & Arn. Prod. 204 = Kleinii.
prostrata, Perr. ex DC. Prod. ii. 228 = Perrottetii.
prostrata, Roxb. Hort. Beng. [98]; Fl. Ind. iii. 373 = echinata.
prostrata, Willd. Sp. Pl. iii. 1236 = trifoliata.
pruinosa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 93.—Afr. trop
psilostachya, *Welw. ex Baker, l. c.* 84.—Afr. trop.
psoralioides, *Linn. f. Mant.* ii. 271.—Afr. austr.
pulchella, *Roxb. Hort. Beng.* 57; *Fl. Ind.* iii. 383.—Ker. Himal.
pulchra, *Willd. Sp. Pl.* iii. 1089.—Afr. trop.
punctata, Boiv. ex Baill. in Bull. Soc. Linn. Par. i. (1883) 898 = Boiviniana.
punctata, Eckl. & Zeyh. Enum. 241 = polioten.
punctata, Thunb. Prod. Pl. Cap. 138 = Zeyheri.
pungens, *E. Mey. Comm. Pl. Afr. Austr.* 98.—Afr. austr.
purpurascens, Roxb. Hort. Beng. 57; Fl. Ind. iii. 383 = pulchella.
purpurea, *Page, ex Steud. Nom.* ed. II. i. 606 (Quid ?). —Afr. austr.
pusilla, Lam. Encyc. ii. 248 = endecaphylla.
quadrangularis, R. Grah. in Wall. Cat. n. 5468 = Gerardiana.
quartiniana, *A. Rich. Tent. Fl. Abyss.* i. 193.—Abyss.
quinquefolia, *E. Mey. Comm. Pl. Afr. Austr.* 98.—Afr. austr.
racemosa, Llan. Fl. Ras. Afr. n. 87; Amoen. Acad. vi. 94 = paucifolia.
rariflola, Steud. Nom. ed. II. i. 606 = pauciflolia.
reclusa, Eckl. & Zeyh. Enum. 233 = denudata.
reflexa, E. Mey. Comm. Pl. Afr. Austr. 100 = glaucescens.

INDIGOFERA :—
retroflexa, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 899.—Madag.
retusa, R. Grah. in Wall. Cat. n. 5476 = argentea.
rhachodes, Walp. in Linnaea, xiii. (1839) 519 = *rechodes*.
rhodesiantha, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 305 = hirsuta.
rhynchocarpa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 92.—Afr. trop.
rhytidocarpa, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 209.—Afr. austr.
Richardiana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 898.—Madag.
rigescens, E. Mey. Comm. Pl. Afr. Austr. 94 = denudata.
rigida, Willd. Enum. Hort. Berol. 780 = trita.
Rothii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 77.—Afr. trop.
rotundifolia, *Yeume St. Hil. Mém. Indig.* 9. t. 2 (Quid ?).—Ind. or.
Roxburghii, Tausch, in Flora, xiv. (1831) 671 = linifolia.
rufescens, *E. Mey. Comm. Pl. Afr. Austr.* 103.—Afr. austr.
rufescens, Poir. Encyc. Suppl. iii. 148 = pulchra.
rugosa, *Benth. Pl. Australas.* ii. 199.—Austral.
rupestris, Eckl. & Zeyh. Enum. 241 = polioten.
sabulicola, *Benth. in Mart. Fl. Bras.* xv. i. 40.—Bras.
Saliana, *Steud. Nom.* ed. II. i. 808.—Abyssin.
sarmentosa, Herb. Holm. ex Harv. & Sond. Fl. Cap. ii. 77 = Dichilus gracilis.
sarmentosa, *Linn. f. Suppl.* 334; *Thunb. Prod. Pl. Cap.* 132.—Afr. austr.
saxicola, *F. Muell. ex Benth. Fl. Austral.* ii. 199.—Austral.
scabra, Roth, Nov. Pl. Sp. 359 = subulata.
Schimperi, *Jaub. & Spach, Illustr.* v. 94. t. 484.—Afr. trop.
Schimperiana, Hochst. in Flora, xxiv. (1841) 1. Intell. 81 = endecaphylla.
Schultziana, *F. Muell. Fragm.* vii. 105.—Austral.
scoparia, Vahl, ex DC. Prod. ii. 229 = senegalensis.
secunda, E. Mey. Comm. Pl. Afr. Austr. 102 = hedyantha.
secundiflora, *Poir. Encyc. Suppl.* iii. 148.—Afr. trop.
Sedgewickiana, *Vatke & Hildebr. in Oestr. Bot. Zeitschr.* xxviii. (1878) 202.—Afr. trop.
semitrijuga, *Forsk. Fl. Aegypt. Arab.* 137.—Afr. bor.; As. occ.
senegalensis, Blanco, Fl. Filip. ed. II. 612 = Tephrosia insolensis.
senegalensis, *Lam. Encyc.* ii. 248.—Afr. trop.
sericea, *Benth. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 76.—Afr. trop.
sericea, Linn. Mant. ii. 271 = Amphithalea densa.
sericea, Thunb. ex Harv. & Sond. Fl. Cap. ii. 22 = Amphithalia ericaefolia.
sessilifiora, *DC. Prod.* ii. 228.—Afr. trop.
sessilifolia, *DC. l. c.* 231.—Afr. austr.
setacea, E. Mey. Comm. Pl. Afr. Austr. 95 = gracilis.
setiolosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 196.—Afr. austr.
setifiora, *Baker, in Oliver, Fl. Trop. Afr.* ii. 90.—Afr. trop.
setulosa, *Bertol. in Nov. Comm. Bonon.* ii. (1836) 314. t. 8.—Afr. austr.
Sieberiana, *Schenle, in Linnaea*, xvii. (1843) 387.—Afr. austr.
simplicifolia, *Dennst. Schlüssw. Hort. Malab.* 35 (Quid ?).—Ind. or.
simplicifolia, Guill. & Perr. Fl. Seneg. Tent. i. 178 = macrocarpa.
simplicifolia, Heyne, in Wall. Cat. n. 5455 = argentea.
simplicifolia, *Lam. Encyc.* iii. 251.—Afr. trop.
somaliensis, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 201.—Afr. trop.
sordida, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 196.—Afr. austr.
Spachii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 93.—Afr. trop.
sparsa, *Baker, l. c.* 79.—Afr. trop.
sphaerocarpa, Hochst. ex Baker, l. c. 80 = suavcolens.
sphaerocarpa, *A. Gray, Pl. Wright.* ii. 37.—N. Mexic.
spicata, *Forsk. Fl. Aegypt. Arab.* 138.—Arab.

INDIGOFERA :—
spinescens, *E. Mey. Comm. Pl. Afr. Austr.* 98.—Afr. austr.
spinifera, Hochst. ex Steud. Nom. ed. II. i. 808 = spinosa.
spinosa, *Forsk. Fl. Aegypt. Arab.* 137.—Arab.
spiendens, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 19.—Afr. trop.
spolista, Hoffmgg. Verz. Pfl. Nachtr. ii. 138 = filifolia.
stachyoides, Lindl. Bot. Reg. (1843) t. 14 = I. Dosua.
stenophylla, *Eckl. & Zeyh. Enum.* 234.—Afr. austr.
stenophylla, *Guill. & Perr. Fl. Senag. Tent.* i. 188. t. 48.—Afr. trop.
stenosepala, *Baker, in Journ. Bot.* xx. (1882) 68.—Madag.
stipularis, Eckl. & Zeyh. Enum. 237 = dimidiata.
stipularis, *Link, Enum. Hort. Berol.* ii. 250.—Afr. austr.
stricta, *Linn. f. Suppl.* 334; *Jacq. Hort. Schoenbr.* t. 326.—Afr. austr.
stricta, Thunb. Prod. Pl Cap. 138 = Tephrosia stricta.
strigosa, *Spreng. Neue Entdeck.* iii. 84.—Afr. austr.
strobilifera, *Hochst. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 75.—Afr. trop.
suaveolens, *Jaub. & Spach, Illustr.* v. 98. t. 489.—Abyss.
subcorymbosa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 91.—Afr. trop.
subtilis, E. Mey. Comm. Pl. Afr. Austr. 98 = filicaulis
subulata, *Vahl, in Poir. Encyc. Suppl.* iii. 150.—Reg. trop.
subulifera, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 96.—Afr. trop.
suffruticosa, Mill. Gard. Dict. ed. VIII. n. 2 = I. Anil.
sulcata, *DC. Prod.* ii. 231.—Afr. austr.
sumatrana, Gaertn. Fruct. ii. 317. t. 148 = tinctoria.
sutherlandioides, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 101.—Afr. trop.
sylvatica, Sieber, ex Spreng. Syst. iii. 278 = australis.
tenella, Schum. & Thonn. Beskr. Guin. Pl. 367 = senegalensis.
tenella, *Vahl, ex DC. Prod.* ii. 229 = pusilla.
tenuicaulis, Klotzsch, in Peters, Reise Mossamb. Bot 50 = vicioides.
tenuifolia, Eckl. & Zeyh. Enum 248 = angustifolia.
tenuifolia, Lam. Encyc. iii. 294 = angustifolia, capillaris.
tenuifolia, *Rottl. ex Wight & Arn. Prod.* 200.—Ind. or.
tenuissima, *E. Mey. Comm. Pl. Afr. Austr.* 94.—Afr. austr.
tephroides, *H. B. & K. Nov. Gen. et Sp.* vi. 455.—Ecuador.
tephrosioides, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 398.—Madag.
terminalis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 70.—Afr. trop.
ternata, Roxb. ex Wight & Arn. Prod. 204 = trita.
tetragonoloba, L'Hérit. ex Steud. Nom. ed. I. 481 = Cyamopsis psoraloides.
tetragonoloba, *E. Mey. Comm. Pl. Afr. Austr.* 106.—Afr. austr.
tetrasperma, *Schum. & Thonn. Beskr. Guin. Pl.* 365.—Afr. trop.
tetrasperma, Vahl, ex Pers. Syn. ii. 325 = senegalensis.
tettensis, Klotzsch, in Peters, Reise Mossamb. Bot. 51 = Schimperi.
texana, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 451 = leptosepala.
Teysmanni, *Miq. Fl. Ind. Bat.* i. 1. 1068.—Malaya.
Theuschii, *O. Hoffm. in Linnaea*, xliii. (1880-82) 126.—Afr. trop.
Thibaudiana, *DC. Prod.* ii. 225.—Mexic.
Thonningii, Schum. & Thonn. Beskr. Guin. Pl. 366 = subulata.
thymoides, *Baker, in Journ. Linn. Soc.* xx. (1883) 126.—Madag.
timoriensis, DC. Prod. ii. 223 = trita.
tinctoria, Blanco, Fl. Filip. ed. I. 591 = I. Anil.
tinctoria, Chapm. Fl. S. U. St. 96 = sphaerocarpa.
tinctoria, Forsk. Fl. Aegypt. Arab. 138 = argentea.
tinctoria, Hook. Comp. Bot. Mag. i. (1835) 32 = leptosepala.
tinctoria, *Linn. Sp. Pl.* 751.—Reg. trop. (omit.).
tomentosa, *Eckl. & Zeyh. Enum.* 237.—Afr. austr.
tomentosa, R. Grah. in Wall. Cat. n. 5467 = I. Dosua.

INDIGOFERA :—
torulosa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 91.—Afr. trop.
torulosa, Hook. & Arn. Bot. Beech. Voy. 386 = subulata.
torulosa, *E. Mey. Comm. Pl. Afr. Austr.* 105.—Afr. austr.
trachyphylla, *Benth. in Hook. Ic. Pl.* t. 1854.—Afr. trop.
trifoliolata, *Boiss. Fl. Orient.* ii. 189.—Afr. trop.; As. occ.
trichopoda, *Lepr. ex Guill. & Perr. Fl. Senag. Tent* i. 177.—Afr. trop.
trifoliata, D. Don, Prod. Fl. Nep. 245 = multicaulis.
trifoliata, *Linn. Cent. Pl.* ii. 29; *Amoen. Acad.* iv. 327.—As. et Austral. trop.
trigonelloides, *Jaub. & Spach, Illustr.* v. 92. t. 482.—Afr. trop.; As. occ.
triquetra, *Dals. in Hook. Kew Journ.* ii. (1850) 86.—Ind. or.
triquetra, *E. Mey. Comm. Pl. Afr. Austr.* 95.—Afr. austr.
tristis, *E. Mey. l. c* 101.—Afr. austr.
trita, *Linn. f. Suppl.* 335.—As. et Austral. trop.
truxillensis, *H. B. & K. Nov. Gen. et Sp.* v. 456.—Peruv.
umbonata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 98.—Afr. trop.
umbraticola, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1878) 202.—Afr. trop.
uncinata, G. Don, Gen. Syst. ii 208 = I. Anil.
uncinata, Roxb. Hort. Beng. 57; Fl. Ind. iii. 382 = galegoides.
uniflora, *Buch.-Ham. ex Roxb. ll. cc.* 57; 374.—Ind. or.
uniflora, Heyne, ex Wall. Cat. sub n. 5444 = enneaphylla.
urostachya, Fenzl, ex Oliver, Fl. Trop. Afr. ii. 94 = secundiflora.
varia, *E. Mey. Comm. Pl. Afr. Austr.* 104.—Afr. austr.
velutina, *E. Mey. l. c.* 104.—Afr. austr.
venulosa, *Champ. ex Benth. in Hook. Kew Journ. Bot.* iv. (1852) 44.—China.
venusta, *Eckl. & Zeyh. Enum.* 235.—Afr. austr.
verrucosa, Eckl. & Zeyh. l. c. 242 = Zeyheri.
verrucosa, R. Grah. in Wall. Cat n. 5469 = pulchella.
vestita, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 96.—Ind. or.
vestita, *Harv. in Harv. & Sond. Fl. Cap.* ii. 182.—Afr. austr.
vicioides, *Jaub. & Spach, Illustr.* v. 91. t. 481.—Afr. trop.
villosa, Berg. ex Walp. Rep. i. 668 = glomerata.
viminea, *E. Mey. Comm. Pl. Afr. Austr.* 102.—Afr. austr.
violacea, Roxb. Hort. Beng. 57; Fl. Ind. iii. 380 = palchella.
virgata, DC. Prod. ii. 224 = trita.
virgata, Roxb. Hort. Beng. 57; Fl. Ind. iii. 388 = I. Gerardiana, I. Dosua.
viscidissima, *Baker, in Oliver, Fl. Trop. Afr.* ii. 100.—Afr. trop.
viscosa, *Lam. Encyc.* iii. 247.—Geront. trop.
vivea, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 64.—Bras.
vohemarensis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 400.—Madag.
volubilis, Wendl. Bot. Beobacht. 55 = Rhynchosia phaseoloides.
Welwitschii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 84.—Afr. trop.
Wightii, *R. Grah. in Wall. Cat.* n. 5458.—Ind or wplwyneuous, *S. Moore, in Journ. Bot.* xvi. (1878) 131.—Afr. austr.
Zeyheri, *Spreng. ex Eckl. & Zeyh. Enum.* 241.—Afr. austr.
Zollingeriana, Miq. Fl Ind. Bat. i. 1. 310 = galegoides.

INFANTEA, Remy, in C. Gay, Fl. Chil. iv. 257. t. 48 (1849) = Amblyopappus, Hook. (Compos.).
chilensis, Remy, l. c. 259 = A. pusillus.

INGA, Scop. Introd. 298 (1777). *LEGUMINOSAE*, Benth. & Hook. f. i. 599.
Dendaria, Rafin. Sylva Tellur. 119 (1838).
acrocephala, *Steud. in Flora*, xxvi. (1843) 759.—Guian.

INGA :—
acuminata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 600.—Ind. or.
Acus, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 139.—Hab. ?
acutangula, R. Grah. in Wall. Cat. n. 5271 = Pithecolobium angulatum.
acutifolia, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 614 = affinis.
adiantifolia, Kunth, Mimos. 66. t. 21 = Pithecolobium adiantifolium.
affinis, *DC. Prod.* ii. 433.—Bras.
affinis, Steud. in Flora, xxvi. (1843) 758 = setifera.
afheloides, Lodd. ex G. Don, in Loud. Hort. Brit. 605 = Calliandra hymenaeoides.
aggregata, *G. Don, Gen. Syst.* ii. 391.—Am. austr.
alba, *Wild. Sp. Pl.* iv. 1013.—Guian.
albicans, Walp. in Linnaea, xiv. (1840) 298 = capitata.
alternifolia, *G. Don, Gen. Syst.* ii. 395.—Mexic.
angulata, R. Grah. in Wall. Cat. n. 5271 = Pithecolobium angulatum.
angustifolia, *Willd. Sp. Pl.* iv. 1012.—Veneruela.
annularis, R. Grah. in Wall. Cat. n. 5269 = Pithecolobium bigeminum.
anomala, Kunth, Mimos. 70. t. 22 = Calliandra grandiflora.
Arrabidae, Steud. Nom. ed. II. i. 809 = affinis.
asplenioides, Steud. l. c. = Calliandra mertensioides.
axillaris, Miq. in Linnaea, xix. (1847) 180 = Buorgoni.
attenuata, R. Grah. in Wall. Cat. n. 5276 = Pithecolobium lobatum.
bahiensis, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 618.—Bras.
barbata, *Benth. l. c.* 606.—Bras.
bauhiniifolia, Poepp. & Endl. Nov. Gen. et Sp. iii. 80 = Calliandra amazonica.
Benthamiana, *Meisn. in Linnaea*, xxi. (1848) 258.—Guian.
Berteriana, DC. Mém. Lég. xii; Prod. ii. 434 = vera.
Berterii, *Spreng. Syst.* iii. 127.—Am. austr.
bigemina, Hook. & Arn. Bot. Beech. Voy. 182 = Pithecolobium lucidum.
bigemina, Willd. Sp. Pl. iv. 1007 = Pithecolobium bigeminum.
bigloboza, Willd. l. c. 1025 = Parkia africana.
Billbergiana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 585.—N. Granat.
Blanchetiana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 624.—Am. trop.
Bonplandiana, *H. B. & K. Nov. Gen. et Sp.* vi. 288.—N. Granat.
borbonica, Hassk. in Flora, xxv. (1842) ii. Beibl. 54 = Albizzia Lebbek.
brachypters, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 610.—N. Granat.
brachystachya, DC. Prod. ii. 440 = Pithecolobium luxorum.
bracteosa, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 609.—Guian.
brevipes, *Benth. in Hook. Journ. Bot.* ii. (1840) 144.—Guian.
bubalina, Jack, in Malay. Misc. ii. (1822) vii. 77 = Pithecolobium bubalinum.
bubalina, Wall. Cat. n. 5272 = Pithecolobium microcarpum.
bullata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 607.—Bras.
Buorgoni, *DC. Prod.* ii. 434 (Burgoni).—Am. trop.
byrsinocarpa, Hochst. in Flora, xxix. (1846) 597 = Millettia ferruginea.
caffra, *Meisn. in Hook. Lond. Journ. Bot.* ii. (1843) 102.—Afr. austr.
calocephala, Poepp. & Endl. Nov. Gen. et Sp. iii. 78 = macrophylla.
calycina, Benth. in Hook. Journ. Bot. iii. (1841) 612 = sessilis.
calycina, Salzm. ex Benth. in Trans. Linn. Soc. xxx. (1875) 611 = capitata.
Camatchili, *Perr. in Mém. Soc. Linn. Par.* iii. (1826) 192.—Ins. Philipp.
campanulata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 607.—Bras.
candida, H. B. & K. Nov. Gen. et Sp. vi. 299 = Pithecolobium excelsum.
canescens, Cham. & Schlecht. in Linnaea, v. (1830) 592 = Calliandra canescens.
capitata, *Desv. Journ. Bot.* iii. (1814) 71.—Guian.
capitata, Miq. ex Benth. in Trans. Linn. Soc. xxx. (1875) 582 = Pithecolobium lactum.

INGA:—
caripensis, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1009 = Calliandra tergemina.
carnosa, *G. Don, Gen. Syst. ii.* 391.—Peruv.
Catharctas, H. B. & K. Nov. Gen. et Sp. vi. 297 = Pithecolobium glomeratum.
Catharinæa, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 605 = striata.
cauliflora, Willd. Sp. Pl. iv. 1021 = Pithecolobium cauliflorum.
cauranensis, *Sagot, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 626.—Guiana.
chartacea, *Poepp. & Endl. Nov. Gen. et Sp.* iv. 79.—Bras.
ciliata, Poepp. & Endl. l. c. 78 = Poeppigiana.
ciliata, *Presl, Symb. Bot.* ii. 11. t. 58.—Bras.
cinerea, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1024 = Pithecolobium Saman.
cinnamomea, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 606.—Bras.
circinalis, Willd. Sp. Pl. iv. 1023 = Pithecolobium circinale.
cissigera, Moritz, ex Benth. in Trans. Linn. Soc. xxx. (1876) 630 = Cæsalpinia.
Clypearia, Jack, in Malay. Misc. ii. (1822) vii. 78 = Pithecolobium Clypearia.
Clypearia, Wall. Cat. n. 5270 B = Pithecolobium fasciculatum.
coccinea, G. Don, Gen. Syst. ii. 390 = Pithecolobium coccineum.
cochleata, Willd. Sp. Pl. iv. 1020 = Pithecolobium cochleatum.
cochlearpa, Mart. Reise Bras. i. 549 = Pithecolobium Auaremotense.
coerulescens, Walp. in Linnæa, xiv. (1840) 298 = ciliata.
cognata, Schlecht. l. c. xii. (1838) 560 = Pithecolobium cognatum.
comervynsiana, *Miq. Stirp. Surin. Sel.* 1.—Guian.
comosa, Willd. Sp. Pl. iv. 1026 = Calliandra comosa.
conferta, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 620.—Peruv.
consimilis, R. Grah. in Wall. Cat. n. 5268 = Pithecolobium contortum.
corcovadinæa, DC. Prod. ii. 441 = Pithecolobium umbellatum.
cordistipula, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 111.—Bras.
coriacea, Desv. Journ. Bot. iii. (1814) 71 = sertulifera.
coriacea, *G. Don, Gen. Syst.* ii. 390.—Mexic.
coriacea, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1010 = Calliandra emarginata.
cornigera, Kostel. Allg. Med.-Pharm. Flora, iv. 1855 = Acacia sphærocephala, spadicigera.
corozoana, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1017.—N. Granat.
corymbifera, Benth. in Hook. Journ. Bot. ii. (1840) 144 = nobilis.
crassifolia, Klotzsch, ex Benth. in Trans. Linn. Soc. xxx. (1875) 583 = Pithecolobium flamentosum.
Crevauxii, *Sagot, in Ann. Sc. Nat.* Sér. VI. xiii. (1882) 331.—Guiana.
Cumingiana, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 614 = Feuillei.
cylacarpa, Willd. Sp. Pl. iv. 1096 = Enterolobium cyclocarpum.
cylindrica, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 114.—Bras.
cyanostroides, *Sodd. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 306.—Ind.
darienesis, *Sam. Bot. Voy. Herald*, 117. t. 28.—Panama.
densiflora, *Benth. in Trans. Linn. Soc.* xxx. (1875) 617.—Am. trop.
diadema, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 114.—Bras.
dichotoma, *Mart. l. c.*—Bras.
dimidiata, Hook. & Arn. Bot Beech. Voy. 181 = Pithecolobium Clypearia.
dimidiata, Miq. Fl. Ind. Bat. i. 1. 38 = Albizia stipulata.
discolor, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1032 = Pithecolobium adiantifolium.
disticha, *Benth. in Hook. Journ. Bot.* ii. (1840) 148.—Guian.
divaricata, Bong. ex Benth. in Mart. Fl. Bras. xv. ii. 448 = Pithecolobium divaricatum.
dolabriformis, R. Grah. in Wall. Cat. n. 5279 = Xylia dolabriformis.

INGA:—
dominicensis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 618.—Ind. occ.
dubia, *R. Grah. in Wall. Cat.* n. 5274.—Ind. or.
dulcis, Mart. in Flora, xx. (1837) ii. Beibl. 113 = affinis.
dulcis, Willd. Sp. Pl. iv. 1005 = Pithecolobium dulce.
dumosa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 612.—Bras.
ebulis, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 113.—Bras.
elegans, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 321.—Mexic.
elliptica, *Blume, Cat. Gew. Buitenz.* 88.—Java.
emarginata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1009 = Calliandra emarginata.
erianthæ, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 614 = Lushnathiana.
eriocarpa, *Benth. l. c.* 615.—Mexic.
excelsa, *H. B. & K. Nov. Gen. et Sp.* vi. 299.—N. Granat.
excelsa, Kunth, Mimos. 57. t. 16 = Pithecolobium excelsum.
excelsa, Poepp. & Endl. Nov. Gen. et Sp. iii. 78 = marginata.
excelsa, Wight, ex Wall. Cat. n. 5286 = Pithecolobium umbellatum.
fasciculifera, Desv. in Ham. Prod. Pl. Ind. Occ. 61 = Parkia africana.
fagacifolia, Mart. ex Benth. in Trans. Linn. Soc. xxx. (1875) 619 = ciliata.
fagifolia, G. Don, Gen. Syst. ii. 391 = Ruiziana.
fagifolia, *Willd. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 607.—Am. trop.
falcifolia, Hassk. Cat. Hort. Bog. Alt. 291 = Pithecolobium montanum.
falciformis, DC. Prod. ii. 435 = Pithecolobium Vahlianum.
falciformis, Hassk. in Flora, xxv. (1842) ii. Beibl. 54 = Pithecolobium montanum.
fasciculata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 79.—Peruv.
fasciculata, Willd. Sp. Pl. iv. 1022 = Calliandra surinamensis.
fasciformis, Hassk. in Hoev. & De Vriese, Tijdschr. x (1843) 149 = falciformis.
fastigiata, Steud. Nom. ed. II. i. 809 = Albizzia fastigiata.
fastuosa, *Willd. Sp. Pl.* iv. 1014.—Venezuela.
felina, Stokes, Bot. Mat. Med. iii. 522 = Pithecolobium Unguis-cati.
Fendleriana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 630.—N. Granat.
ferruginea, Guill. & Perr. Fl. Seneg. Tent. i. 236 = Albizzia ferruginea.
ferruginea, Planch. in Fl. des Serres, Sér. I. viii. (1852-53) t. 778 = vulpina.
ferruginea, *Mart. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 625.—Bras.
Feuillei, *DC. Prod.* ii. 433.—Peruv.
filicina, Willd. Sp. Pl. iv. 1025 = Parkia filicina.
filipes, Vent. Choix, 88 = Pithecolobium filipes.
Finlaysoniana, R. Grah. in Wall. Cat. n. 5284 = Pithecolobium contortum.
flagelliformis, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 112.—Bras.
flexuosa, R. Grah. in Wall. Cat. n. 5286 = Pithecolobium umbellatum.
flexuosa, Schlecht. in Linnæa, xii. (1838) 559 = xalapensis.
floribunda, Benth. in Hook. Journ. Bot. ii. (1840) 148 = splendens.
foetida, Willd. Sp. Pl. iv. 1008 = Pitpadenia foetida.
foliosa, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 597 = Ruiziana.
Forfex, Kunth, Mimos. 52. t. 16 = Pithecolobium Unguis-cati.
forficata, *Hoffmgg. Vers. Pfl. Nachtr.* ii. 139.—Bras.
fragrans, Macfad. Fl. Jamaic. i. 209 = Pithecolobium fragrans.
fraxinea, Willd. Sp. Pl. iv. 1019 = alba.
fulgens, Kunth, Mimos. 56. t. 21 = spectabilis.
gallitica, Duchass. & Walp. in Linnæa, xxiii. (1850) 747 = ingoides.
geminata, Wight & Arn. Prod. 269 = Pithecolobium geminatum.

INGA:—
gladiata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 427 = Thibaudiana.
glandulosa, *Stend. Nom.* ed. I. 481.—Guian.
globosa, *Blume, Cat. Gew. Buitenz.* 88.—Java.
globuliflora, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 585.—Am. centr.
glomerata, DC. Prod. ii. 438 = Pithecolobium glomeratum.
gracilifolia, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 582.—Guian.
gracilis, Jungh. ex Miq. Fl. Ind. Bat. i. 1. 22 = Albizzia procera.
grandiflora, Wall. Cat. n. 5285 = Serianthes grandiflora.
guadalupensis, Desv. Journ. Bot. iii. (1814) 70 = Pithecolobium Unguis-cati.
guatemalensis, Hook. & Arn. Bot. Beech. Voy. 419 = Mimosa guatemalensis.
guayaquilensis, G. Don, Gen. Syst. ii. 391 = marginata.
Guilleminiana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 605.—Bras.
Haematoxylon, Hort. Calc. ex Hassk. Retzia, 217 = Calliandra haematocephala.
Harrisii, Lindl. Bot. Reg. (1839) t. 41 = Calliandra Harrisii.
Hayesii, *Benth. in Trans. Linn. Soc.* xxx. (1876) 617.—Panama.
heterophylla, *Willd. Sp. Pl.* iv. 1020.—Bras.
heteroptera, *Benth. in Hook. Lond. Journ. Bot.* iv (1845) 610.—N. Granat.
hirsuta, G. Don, Gen. Syst. ii. 395 = Calliandra hirsuta.
hispida, *Schott, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 625.—Bras.
Histrix, A. Rich. Ess. Fl. Cub. 471 = Pithecolobium Hystrix.
Houstoni, DC. Prod. ii. 442 = Calliandra Houstoni.
Humboldtiana, H. B. & K. Nov. Gen. et Sp. vi. 285 = nobilis.
hymenaeaefolia, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1006 = Pithecolobium hymenaeaefolium.
hymenaeoides, Desv. Journ. Bot. iii. (1814) 70 = Calliandra hymenaeoides.
inæqualis, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1019 = Pithecolobium inæquale.
ingoides, *Willd. l. c.* 1012.—Guian.
Inicuil, *Cham. & Schlecht. in Linnæa*, v. (1830) 392.—Mexic.
inidgni, *Kunth, Mimos.* 43. t. 13.—Ecuador.
Isombergiana, A. Rich. Tent. Fl. Abyss. i. 236 = Albizzia Julibrissin.
javana, DC. Prod. ii. 434 = Pithecolobium dulce.
Jiniculi, Schlecht. in Linnæa, xii. (1838) 559 = I. Inicuil.
Jiringa, Jack, in Malay. Misc. ii. (1822) vii. n. 78 = Pithecolobium lobatum
Jiringa, Wall Cat. n. 5288 = Pithecolobium fasciculatum.
jugladifolia, *Willd. Sp. Pl.* iv. 1018.—Venezuela.
Kaaringa, Voigt, Hort. Suburb. Calc. 258 = Pithecolobium lobatum.
Kamaburunae, Vahl, l. c. = Pithecolobium angulatum.
Kikium, Blume, ex Miq. Fl. Ind. Bat. i. 1. 23 = Albizzia procera.
Kœnigii, Wight & Arn. Prod. 269 = Pithecolobium geminatum.
Kouringa, Kostel. Allg. Med.-Pharm. Flora, iv. 1854 = Pithecolobium lobatum.
lanata, Poepp. & Endl. Nov. Gen. et Sp. iii. 80 = Pithecolobium lanatum.
lanigata, *Mart. & Gal. in Bull. Acad. Brux.* x. tt. (1843) 320.—Mexic.
lalleata, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1876) 628.—Peruv.
lamprophylla, C. Wright, ex Benth. l. c. 627 = vera †
lanceolata, *Benth. l. c.* 606.—Bras.
lanceolata, Blanco, Fl. Filip. ed. II. 270 = Pithecolobium dulce.
lanceolata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1005 = Pithecolobium ligustrinum.
lanuifloras, *Miq. in Linnæa*, xix. (1847) 181.—Guian.
latifolia, Willd. Sp. Pl. iv. 1020 = Pithecolobium latifolium.
lanuifolia, *Bong. ex Benth. in Hook. Journ. Bot.* ii. (1840) 24.—Bras.
lanuina, *Willd. Sp. Pl.* iv. 1016.—Am. austr.
lanuifora, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 617.—Peruv.

INGA:—
leiocalycina. *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 598.—Guian.
lentiscifolia, *Benth. in Trans. Linn. Soc.* xxx. (1875) 613.—Bras.
lentiscifolia, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 492.—Bras.
leptantha, *Benth. l. c.* 608.—Bras.
leptoloba, *Schlecht. in Linnaea*, xii. (1838) 560.—Mexic.
leptophylla, *Lag. Gen. et Sp. Nov.* 16.—Am. trop.
leptopus, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 580 = sertulifera.
leptostachya, Benth. in Trans. Linn. Soc. xxx. (1875) 608 = marginata.
leucantha, Presl, Bot. Bemerk. 65 = Pithecolobium dulce.
leucoxylon, Hassk. in Flora, xxv. (1842) ii. Beibl. 54 = Albizzia Lebbek.
lignosa, R. Grah. in Wall. Cat. n. 5278 = Xylia dolabriformis.
ligustrina, Willd. Sp. Pl. iv. 1007 = Pithecolobium ligustrinum.
Lindeniana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 608.—Am. trop.
lineata, *Benth. l. c.* 594.—Peruv.
lobata, R. Grah. in Wall. Cat. n. 8280 = Pithecolobium lobatum.
longiflora, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 600.—Am. trop.
longifolia, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1010 = Pithecolobium Vahlianum.
longipes, *Benth in Trans. Linn. Soc.* xxx. (1875) 617. —Peruv.
lucida, H. B. & K. Nov. Gen. et Sp. vi. 287 = spectabilis.
lucida, Wall. Cat. n. 5267 = Albizzia lucida.
lucidior, Steud. Nom. ed. II. i. 810 = Albizzia lucida.
Lushnathiana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 618.—Bras.
macradenia, Mart. ex Benth. in Trans. Linn. Soc. xxx. (1875) 615 = Thibaudiana.
macrophylla, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 132 = portobellensis.
macrophylla, Hook. Bot. Mag. t. 5076 = setifera.
macrophylla, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1015.—Venezuela.
macrostachya, DC. Prod. ii. 437 = Pithecolobium macrostachyum.
macrostachys, *Steud. Nom.* ed. I. 481.—Guian.
Magdalenae, Spreng. ex DC. Prod. ii. 472 = Desmanthus brachylobus.
malacophylla, A. Rich. Tent. Fl. Abyss. i. 235 = Albizzia pallida, ferruginea.
marginata, *H. B. & K. Nov. Gen. et Sp.* vi. 285.—Am. trop.
marginata, *Willd. Sp. Pl.* iv. 1015.—Am. trop.
marginata, Willd. l. c. (quoad syn.) = fagifolia.
maritima, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 601.—Bras.
Martiana, Spreng. ex DC. Prod. ii. 441 = Pithecolobium parvifolium.
martinicensis, *Presl, Symb. Bot.* i. 65. t. 42.—Ins. Martinic.
Matheuziana, *Benth. in Hook. Lond. Journ Bot.* iv. (1845) 594.—Peruv.
mayencis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 618.—Am. trop.
Melanocarpa, *Miq. Stirp. Surin. Sel.* 2.—Guian.
mellifera, Willd. Sp. Pl. iv. 1006 = Acacia mellifera.
membranacea, *Benth. in Trans. Linn. Soc.* xxx. (1875) 606.—Am. trop.
Meriana, Spitg. in Hoev. & De Vriese, Tijdschr. ix (1843) 113 = ingoides.
merinensioides, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 36 = Calliandra mertensioides.
micradenia, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 620 —Am. trop.
microcalyx, *Spruce, ex Benth. l. c.* 611.—Bras.
microphylla, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1004 = Pithecolobium Unguis-cati.
microphylla, Salzm. ex Benth. in Trans. Linn. Soc. xxx. (1875) 619 = ellisia.
Mocniana, G. Don, Gen. Syst. ii. 386 = vera.
mollis, *R. Grah. in Wall. Cat. n.* 5275.—Ind. or.
mollissima, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1023 = Calliandra mollissima.
mollissecula, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 427 (Calliandræ sp. ?).—Hab. ?

INGA:—
monilifermis, DC. Prod. ii. 440 = Pithecolobium monilliferum.
montana, Jungh. Reise Topog. 388 = Albizzia montana.
montana, Span. ex Miq. Fl. Ind. Bat. i. 1. 86 = Pithecolobium montanum.
Mucuna, Walp. Ann. ii. 459 = Lindeniana.
multicaulis, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 624.—Ecuador.
multiflora, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 598 = leiocalycina.
multijuga, *Benth. in Trans. Linn. Soc.* xxx. (1875) 615.—Am. trop.
myriantha, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 77. t. 229.—Am. trop.
nandinaefolia, DC. Prod. ii. 440 = Pithecolobium Anaremotomo.
Nefasia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 237 = Albizzia Abyssinica.
negrensis, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 621.—Bras.
Nemu, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1027 = Piptadenia peregrina.
nitens, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 319.—Mexic.
nitida, *Willd. Sp. Pl.* iv. 1018.—Bras.
nobilis, *Willd. Enum. Hort. Berol.* 1047.—Bras.
nodosa, Willd. Sp. Pl. iv. 1018 = Cassia bacillaris.
nuda, *Salzm. ex Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 607.—Am. trop.
nutans, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 114.—Bras.
obovalis, A. Rich. Ess. Fl. Cub. 472 = Pithecolobium obovale.
obtusa, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 489.—Bras.
obtusata, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 621.—Bras.
obtusifolia, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1022 = Calliandra purpurea.
odorata, G. Don, Gen. Syst. ii. 388 = marginata.
Oerstediana, *Benth. in Seem. Bot. Voy. Herald*, 117.—Costa Rica.
ornata, Kunth, Mimos. 46. t. 14 = Ingoides.
ornifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 291.—Ecuador.
packycarpa, Benth. Fl. Hartw. 172 = insignis.
panamensis, *Seem. Bot Voy. Herald*, 117.—Panama.
paniculata, Spreng. ex DC. Prod. ii. 437 = Pithecolobium pubescens.
panurensis, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 602.—Bras.
parviflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 189. —Bras.
parvifora, Sagot, ex Benth. in Trans. Linn. Soc. xxx. (1875) 610 = alba.
parvifora, Spruce, ex Benth. l. c. 602 = laterifiora.
parvifolia, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 209 = Calliandra bicolor.
patens, Hook. & Arn. Bot. Beech. Voy. 419 = Piptadenia patens.
pauciflora, Duchass. ex Walp. in Linnæa, xxiii. (1850) 746 = vera.
Pavoniana, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 598 = nobilis.
Pavoniana, *G. Don, Gen. Syst.* ii. 388.—Am. trop.
pectinata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1027 = Parkia pectinata.
pedicellaris, DC. Prod. ii. 441 = Pithecolobium pedicellare.
peduncularis, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 599 = stenoptera.
pedunculata, Mart. ex Benth. in Trans. Linn. Soc. xxx. (1875) 611 = capitata.
pendula, Willd. Sp. Pl. iv. 1025 = Parkia pendula.
penatula, Cham. & Schlecht. in Linnæa, v. (1830) 593 = Acacia pennatula.
Perrottetii, *Sagot, in Ann. Sc. Nat.* Sér. VI. xiii. (1882) 385.—Guian.
pexisiflora, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 587.—Guian.
pilosiuscula, *Desv. Journ. Bot.* iii. (1814) 71.—Guian.
Pisana, *G. Don, Gen. Syst.* ii. 383.—Peruv.
pistaciaefolia, Spreng. Syst. iii. 121 = Mimosa pistaciaefolia.

INGA:—
plantisiliqua, Mart. in Flora, xx. (1837) ii. Beibl. 111 = cordistipula.
platycarpa, Benth. in Hook. Journ. Bot. ii. (1845) 142 = setifera.
platyptera, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 602.—Bras.
plumifera, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 621.—Bras.
Poeppigiana, *Benth. in Hook. Lond. Journ. Bot.* iv (1845) 602 —Bras.
polystachya, Benth. l. c. 587 - cylindrica.
portobellensis, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 122.—Panama
Prieuri, *Sagot, in Ann. Sc Nat.* Sér. VI. xiii. (1882) 385.—Guian.
protracta, Steud. in Flora, xxvi (1843) 758 = heterophylla.
pruriens, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 78.—Peruv.
pterocarpa, DC. Prod. ii. 441 = Peltophorum ferrugineum.
puberula, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 589 = marginata.
pubescens, Bert. ex DC. Prod. ii. 437 = Pithecolobium pubescens.
pubigera, *Zipp. ex Span. in Linnæa*, xv. (1841) 198, nomen.—Ins. Timor.
pubiramea, *Steud. in Flora*, xxvi. (1843) 759.—Guian.
pulcherrima, Cerv. ex Sweet, Hort. Brit. ed. I. 483 = Calliandra Tweedii.
punctata, *Willd. Sp. Pl.* iv. 1016 (excl. syn.).—Venezuela.
pungens, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1004 = Pithecolobium dulce.
purpurascens, Hassk. in Flora, xxv. (1842) ii. Beibl. 103 = Albizzia stipulata.
purpurea, Willd. Sp. Pl. iv. 1021 = Calliandra purpurea.
pycnostachya, Benth. in Hook. Lond. Journ. Bot. iv (1845) 589 = marginata.
pyriformis, Jungh ex Miq. Fl. Ind. Bat. i. 1. 52 = Parkia Roxburghii.
Quarhniana, A. Rich. Tent Fl. Abyss. i. 235 = Albizzia Quartiniana.
quassiaefolia, Willd. Sp. Pl. iv. 1013 = altida.
quatemata, *Poepp. & Endl. Nov Gen. et Sp.* iii. 79.—Bras.
quinquenina, Hoffingg. ex Mart. in Flora, xx. (1837) ii. 116 = Pithecolobium cauliflorum.
ramiflora, Bong. ex Benth. in Trans. Linn. Soc. xxx (1875) 598 = Pithecolobium ramiflorum.
ramiflora, G. Don, Gen. Syst. iii. 392 = Pithecolobium glomeratum.
ramiflora, Steud. in Flora, xxvi. (1843) 759 = Pithecolobium cauliflorum.
reticulata, Spreng. Syst. iii. 130 = Feuillei.
rhoifolia, Hort. ex Planch. in Fl. des Serres, Sér. I. viii. (1859–53) sub t. 775 = vulpina.
rhoifolia, *Willd. Enum. Hort. Berol.* 1046.—Bras.
Riedeliana, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 595 = nobilis.
rosea, Steud. Nom. ed. I. 481 = Pithecolobium Unguis-cati.
rubiginosa, *DC. Prod.* ii. 434.—Guian.
ruiescens, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 586.—Am. sentr.
ruiinervia, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 612.—Bras.
rufescta, *Benth. l. c.* 621.—Am. trop.
Rusiana, *G. Don, Gen. Syst.* ii. 391.—Am. trop.
rutilans, Spruce, ex Benth. in Trans. Linn. Soc. xxx. (1875) 602 = umbellifera.
salutaris, H. B. & K. Nov. Gen. et Sp. vi. 304 = Pithecolobium Saman.
Salzmanniana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 606.—Bras.
Samea, Willd. Sp. Pl. iv. 1024 = Pithecolobium Saman.
sanguinea, Burch. ex Benth. in Trans. Linn. Soc. xxx (1875) 594 = Pithecolobium sanguineum.
sapida, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 590 = aggregata.
sapida, H. B. & K. Nov. Gen. et Sp vi. 286 = marginata.
sapindoides, *Willd. Sp. Pl.* iv. 1012.—Am. trop.
saponaria, Willd. l. c. 1008 = Albizzia saponaria *Sassa*, Willd. l. c. 1027 = Albizzia fastigiata.

INGA :—
scabriuscula, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 606.—Bras.
Schiedeana, Steud. Nom. ed. II. i. 810 = xalapensis.
schiolfolia, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 584.—Bras.
Sciadica, *Steud. in Flora*, xxvi. (1843) 758.—Guian.
Sellowiana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 583.—Bras.
semialata, Mart. in Flora, xx. (1837) ii. Beibl. 111 = marginata.
semicordata, Bertol. Fl. Guatim. 41 = Calliandra Seemanni.
senegalensis, DC. Prod. ii. 442 = Parkia africana.
sericantha, Miq. in Linnæa, xix. (1847) 132 = nobilis.
sericea, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 318.—Mexic.
sericocephala, A. Rich. Tent. Fl. Abyss. i. 236 = Albizzia amara.
sertulifera, *DC. Prod.* ii. 436.—Guian.
sessilis, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 114.—Bras.
setifera, *DC. Prod.* ii. 432—Guian.
setigera, Poepp. & Endl. Nov. Gen. et Sp. iii. 79 = rufiseta.
setosa, *G. Don, Gen. Syst.* ii. 388.—Am. trop.
sinemariensis, G. Don, l. c. = laurina ?
speciosa, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 320 = Calliandra Cumingii.
speciosa, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 630.—Bras.
spectabilis, *Willd. Sp. Pl.* iv. 1017.—Am. trop.
spinæfolia, Desv. in Journ. Bot. iii. (1814) 72 = Pithecolobium circinale.
spinifolia, Ham. Prod. Pl. Ind. Occ. 61 = præc. ?
splendens, Benth. in Trans. Linn. Soc. xxx. (1875) 612 = dumosa.
splendens, *Willd. Sp. Pl.* iv. 1017.—Bras.
Sprengelii, *G. Don, Gen. Syst.* ii. 392.—Am. austr.
Spruceana, Benth. in Hook. Kew Journ. ii. (1850) 289 = ulei.
sputa, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1011.—Venezuela.
stenocalyx, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 611.—Ecuador.
stenocarpa, Spruce, ex Benth. l. c. 603 = heterophylla.
stenoptera, *Benth. in Hook. Journ. Bot.* ii. (1840) 143. —Guiana.
stenostachya, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 437 = Pitpadenia psilostachya.
stipulacea, G. Don, Gen. Syst. ii. 391 = rufiseta.
stipularia, *DC. Mém. Lég.* 440.—Guian.
striata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 606.—Bras.
strigillosa, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 612.—Bras.
subfalcata, Zoll. & Mor. in Nat. Genootsch. Arch. Nederl. Ind. iii. (1846) 81 = Pithecolobium montanum.
subsnbnda, *Salzm. ex Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 613.—Bras.
suaveolens, Lem. Jard. Fleur. iii. (1853) t. 299–300 ; Misc. 84 = vulpina.
Tapinhuana, *Jacq. ex Heynh. Nom.* i. 417 (Quid ?).—Am. austr.
tarapotensis, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 609.—Peruv.
taxifolia, Kunth, Mimos. 64. t. 20 = Calliandra taxifolia.
tenerrima, Jungh. ex Benth. in Miq. Pl. Jungh. 270 = Acacia pennata.
tengerensis, Zoll. & Mor. in Nat. en Geneesk. Arch. Nederl. Indie, iii. (1846) 78, 81 = Pithecolobium Junghuhnianum.
tenuiflora, Salzm. ex Benth. in Hook. Lond. Journ. Bot. iv. (1845) 596 = Thibaudiana.
tenuifolia, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 587.—Bras.
tenuis, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 114.—Bras.
tergemina, Willd. Sp. Pl. iv. 1008 = Calliandra tergemina.
tetraphylla, G. Don, Gen. Syst. ii. 392 = Calliandra tetraphylla.
tetraphylla, Mart. in Flora, xx. (1837) ii. Beibl. 113 = fagifolia.
Thibaudiana, *DC. Prod.* ii. 434.—Guian.
thyrsoidea, Desv. Journ. Bot. iii. (1814) 71 = alba.
timoriana, DC. Prod. ii. 442 = Parkia Roxburghii.
tomentosa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 609.—Am. trop.
trapezifolia, DC. Prod. ii. 441 = Pithecolobium trapezifolium.
trapeziformis, Steud. in Flora, xxvi. (1843) 759 = Pithecolobium corymbosum.
trifora, G. Don, Gen. Syst. ii. 394 = Pithecolobium trifloram.
tubulifera, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 584.—Am. austr.
umbellata, G. Don, Gen. Syst. ii. 391 = heterophylla.
umbellata, Willd. Sp. Pl. iv. 1027 = Pithecolobium umbellatum.
umbellata, Zoll. in Flora, xxx. (1847) 706 = Pithecolobium monilliferum.
umbellifera, *Steud. ex DC. Prod.* ii. 432.—Am. austr.
umbraculiformis, Jungh. in Flora, xxx. (1847) 608 = Albizzia stipulata.
umbratica, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 77.—Bras.
umbrosa, Wall. Pl. As. Rar. ii. 23. t. 124 = Calliandra umbrosa.
uncinata, Spruce, ex Benth. in Trans. Linn. Soc. xxx. (1875) 630 = scabriuscula.
Unguis-cati, Willd. Sp. Pl. iv. 1006 = Pithecolobium Unguis-cati.
uniujuga, Poepp. & Endl. Nov. Gen. et Sp. iii. 79 = Macrolobium limbatum.
unguenda, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 302.—Uruguay.
Vachliana, DC. Prod. ii. 438 = Pithecolobium Vahlianum.
Velloziana, Mart. in Flora, xx. (1837) ii. Beibl. 114, 118 = maritima.
velutina, *Willd. Sp. Pl.* iv. 1014.—Bras.
venosa, *Trinch. Fl. Brit. W. Ind.* 711, nomen.—Ind. occ.
ventricosa, R. Grah. in Wall. Cat. n. 5266, partim = Serianthes grandiflora.
vera, H. B. & K. Nov. Gen. et Sp. vi. 289 = edulis.
vera, *Willd. Sp. Pl.* iv. 1010.—Am. trop.
verrucosa, *Presl, Bot. Bemerk.* 66.—Bras.
versicolor, Spruce, ex Benth. in Trans. Linn. Soc. xxx. (1875) 616 = setifera.
vestita, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 604.—Am. trop.
villosa, *Mart. & Gal. in Bull. Acad. Brux.* x. ii. (1843) 319.—Mexic.
villosissima, *Benth. in Trans. Linn. Soc.* xxx. (1875) 624.—Venezuela.
virescens, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 606.—Bras.
virgultosa, *Desv.* in *Ann. Sc. Nat.* Sér. I. ix. (1826) 426.—Guian.
vismiæfolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 79. —Bras.
vonapanfolia, Spruce, ex Benth. in Trans. Linn. Soc. xxx. (1875) 605 = heterophylla.
vulpina, *Mart. ex Benth. l. c.* 625.—Am. trop.
Wightiana, R. Grah. in Wall. Cat. n. 5281 = Pithecolobium bigeminum.
xalapensis, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 616.—Mexic.
xylocarpa, DC. Prod. ii. 439 = Xylia dolabriformis.
Zygia, DC. Mém. Lég. xii. 440. t. 65 = Albizzia Brownei.

INGARIA, Rafin. Sylva Tellur. 119 (1838) = præc.

INGENHOUSIA, Endl. Gen. 1415, in syn. (1841) = INGENHOUSSIA, Dennst. = **Vitis**, Tourn. (Ampelid.).

INGENHOUSIA, Spach, Hist. Vég. Phan. iii. 843 (1834) = **Ingenhousia**, Moç. & Sesse (Malvac.).

INGENHOUSIA, Steud. Nom. ed. II. i. 810 (1840) = INGENHOUZIA, Bert. = **Rhetinodendron**, Meisn. (Compos.).

INGENHOUSIA, Steud. Nom. ed. II. i. 810 (1840) = INGENHOUSSIA, E. Mey. = **Amphithalea**, Eckl. & Zeyh. (Legumin.).

INGENHOUSSIA, Dennst. Schlüssel. Hort. Malab. 33 (1818) = **Vitis**, Tourn. (Ampelid.).
umbellata, Dennst. l. c.—Ind. or.

INGENHOUSSIA, E. Mey. Comm. Pl. Afr. Austr. 20 (1835) = **Amphithalea**, Eckl. & Zeyh. (Legum.).
ericæfolia, E. Mey. l. c. 21 = A. ericæfolia.
micrantha, E. Mey. l. c. = A. micrantha.
rosea, E. Mey. l. c. 153 = A. densa.
rugosa, E. Mey. l. c. 22 = Coelidium ciliare.
spinosa, E. Mey. l. c. = Coelidium spinosum (in Addend.).
tortilis, E. Mey. l. c. = Coelidium Vogelii.
verticillata, E. Mey. l. c. 21 = Priestleya elliptica.
violacea, E. Mey. l. c. 12 = A. ericæfolia.

INGENHOUSSIA, Reichb. in Moessl. Handb. ed. II. p. lxiii. (1837) = **Ingenhousia**, Moç. & Sesse (Malvac.).

INGENHOUSZIA, Meisn. Gen. 97 (1837) = præc.

INGENHOUZIA, Bert. ex DC. Prod. vi. 447 (1837) = **Rhetinodendron**, Meisn. (Compos.).
thurifera, Bert. l. c. = R. Berteroi.

INGENHOUZIA, [Moç. & Sesse] ex DC. Prod. i. 474 (1824). *MALPACEÆ* [Benth. & Hook. f. i. 209, 982].
triloba, [*Moç. & Sesse*] l. c.—Mexic.

INGENHUSIA, Vell. Fl. Flum. Ic. viii. t. 93 (1827) = **Trichocline**, Cass. (Compos.).
radiata, Vell. l. c. = T. polymorpha.

INGENHUSIA, Vell. Fl. Flum. 351 (1825) = **Trichocline**, Cass. (Compos.).
radiata, Vell. l. c. = T. polymorpha.

INGOCARPUS, Forst. Char. Gen. 65. t. 33 (1776). *LEGUMINOSÆ*, Benth. & Hook. f. i. 552.
Bocoa, Aubl. Pl. Guian. Suppl 33. t. 391 (1775).
Etaballia, Benth. in Hook. Journ. Bot. ii. (1840) 99.
Inodaphnis, Miq. Fl. Ind. Bat. Suppl. 357 (1860). edulis, *Forst. Char. Gen.* 66.—Ins. Pacif.

INODAPHNIS, Miq. Fl. Ind. Bat. Suppl. 357 (1860) = **Inocarpus**, Forst. (Legumin.).
lanceolata, Miq. l. c. = Inocarpus edulis.

INOPSIDIUM, Walp. Rep. i. 175 (1842) = IONOPSIDIUM, Reichb. = **Cochlearia**, Tourn. (Crucif.).

INOPSIS, Steud. Nom. ed. I. 432 (1821) = **Ionopsis**, Kth. (Orchid.).

INTHYBUS, Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) i. 195 = INTYBUS, Fries = **Crepis**, Vaill. (Compos.).

INTKUSARIA, Rafin. Fl. Tellur. iv. 66 (1836) = **Asystasia**, Blume (Acanthac.).
vismiæfolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 79. —Bras.
vonapanfolia, Spruce, ex Benth. in Trans. Linn. Soc. xxx. (1875) 605 = heterophylla.
vulpina, *Mart. ex Benth. l. c.* 625.—Am. trop.
Wightiana, R. Grah. in Wall. Cat. n. 5281 = Pithecolobium bigeminum.
xalapensis, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 616.—Mexic.
xylocarpa, DC. Prod. ii. 439 = Xylia dolabriformis.
Zygia, DC. Mém. Lég. xii. 440. t. 65 = Albizzia Brownei.

INTRODUCTORIA, Rafin. Fl. Tellur. iv. 66 (1836) = **Asystasia**, Blume (Acanthac.).
bicolor, Rafin. l. c. = A. intrusa.

INTSIA, Thou. Gen. Nov. Madag. 22 (1806) = **Afzelia**, Sm. (Legumin.).
amboinensis, DC. Prod. ii. 509 = A. bijuga.
coriacea, Maingay, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 275 = A. coriacea.
madagascariensis, DC. Prod. ii. 509 = A. bijuga.
Moeleri, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 359.—N. Caled.
palembanica, Miq. Fl. Ind. Bat. Suppl. 289 = A. palembanica.
pubescens, Miq. l. c. 288.—Sumatra.

INTUTIS, Rafin. Sylva Tellur. 108 (1838) = **Capparis**, Linn.
amygdalina, Rafin. l. c.—C. odoratissima.
ferruginea, Rafin. l. c.—C. odoratissima.

INTYBELLIA, Cass. in Bull. Soc. Philom. (1821) 124 = **Pterothrix**, Cass. (Compos.).
glareosa, Schott & Kotschy, ex Boiss. Fl. Orient. iii. 858 = Lagoseris frigida.
purpurea, DC. Prod. vii. 160 = P. bifida.
rosea, Cass. in Dict. Sc. Nat. xxiii. 548 = P. bifida.
taurica, Less. Syn. Comp. 144 = P. bifida.

INTYBELLIA, Monn. Ess. Hieracc. 78 (1829) = **Crepis**, Vaill. (Compos.).
incarnata, Monn. l. c. 79 = C. incarnata.
praemorsa, Monn. l. c. = C. praemorsa.
pulchra, Monn. l. c. = Phoecosiun lampanoides.

INTYBUS, Fries, Novit. Fl. Suec. ed. II. 244 (1828) = **Crepis**, Vaill. (Compos.).
bulbosus, Fries, l. c. 245 = C. bulbosa.
incarnatus, Fries, l. c. in obs. = C. incarnata.
multicaulis, Fries, in Cat. Sem. Hort. Upsal. (1855); et in Bot. Notiser (1857), 181 = C. multicaulis.
praemorsus, Fries, Novit. Fl. Suec. ed. II. 245, in obs. = C. praemorsa.

INULA, Linn. Dias. Pan. 35 (1737); Amoen Acad. i. 409 (1749). *COMPOSITAE*, Benth. & Hook. f. ii. 330.
Amphiraphis, DC. Prod. v. 343 (1836).
Corvisartia, Mérat, Fl. Par. ed. I. 228 (1812); ed. II. ii. 261 (1818).
Cupularia, Crem. & Godr. Fl. Fr. ii. 180 (1850).
Diplogon, Rafin. Cat. 13 (1824).
Dumaldea, DC. Prod. v. 366 (1836).
Enula, Neck. Elem. i. 4 (1790).
Erithesis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 464 (1821).
Helenium, Linn. Syst. ed. I (1735); Adans. Fam. ii. 125 (1763).
Inulastrum, Sch. Bip. in Flora, xxv. (1842) i. Beibl. 134, 147, nomen; et in A. Rich. Tent. Fl. Abyss. i. 399 (1847).
Limbarda, Adans. Fam. ii. 125 (1763).
Panjopsis, Rafin. Fl. Tellur. ii. 49 (1836).
Schizogyne, Cass. in Dict. Sc. Nat. lvi. 23 (1828)
Ulina, Opiz, Seznam, 100 (1852).
acaulis, *Schott & Kotschy, ex Tchihat. Asie Min. Bot.* ii. 243.—As. Min.
acris, Bernh. Syst. Verz. Erf. 149 = Erigeron acris.
acuminata, *DC. Prod.* v. 472.—Reg. Himal.
adriatica ×, Borb. in Oest. Bot. Zeitschr. xxvii. (1877) 187 = squarrosa.
aestuans, Linn. Sp. Pl. ed. II. 1236 = Verbesina aestuans.
africana, *Lam. Encyc.* iii. 256.—Afr. austr.
alpina, Adam, in Web. & Mohr, Beitr. i. 68 = grandiflora.
ammophila, *DC. Prod.* v. 470.—China.
anatolica, *Boiss. Diagn.* Ser. I. xi. 6.—As. Min.
angustifolia, *DC. Prod.* v. 385.—Persia.
angustifolia, Wender. in Flora, xii. (1829) 1. Erg. 26 = ensifolia.
anthelmintica, Salisb. Prod. 201 = I. Helenium.
appendiculata, Wall. Cat. n. 2966 = Vicoa auriculata.
apula, Cranta, Inst. i. 327 = Pulicaria odora.
arabica, Bové, ex DC. Prod. v. 478 = Pulicaria deserto rum.
arabica, Linn. Mant. i. 114 - Pulicaria arabica.
Arbuscula, *Delile, in Ann. Sc. Nat.* Sér. II. xx. (1843) 93.—Abyss.
argentea, Pers. Syn. ii. 452 = Chrysopsis graminifolia.
aromatica, Linn. Pl. Afr. Rar. 23; Amoen. Acad. vi. 108 = Printzia aromatica.
Aschersoniana, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 179.—Balkan.
aspera, Polr. Encyc. Suppl. iii. 154 = salicina.
asperrima, Edgew. in Trans. Linn. Soc. xx. (1846) 69 = nervosa.
Aucheriana, *DC. Prod.* vii. 284.—As. Min.
aurantiaca, Spreng. Syst. iii. 524 = Dysodia appendiculata.
aurea, *Hassk. in Flora*, ii. (1868) 27.—Malaya.
auriculata, *Boiss. & Bal. Diagn.* Ser. II. iii. 13.—As. Min.
auriculata, Scher, in Verh. Siebenb. Ver. Naturw. x. (1859) 68 = salicina.
auriculata, Wall. Cat. n. 2965 = Vicoa auriculata.
auriculata, Hasskarl. ex Beck, in Denkschr. Acad. Wien, Math.-Nat. xliv. (1882) 225 = candida.
barbata, *Wall. Cat.* n. 2961.—Reg. Himal.
Barthiana, Schur, in Oestr. Bot. Zeitschr. xxi. (1871) 108 = stricta.
bifrons, *Linn. Sp. Pl.* ed. II. 1236.—Europ. austr.
bonhuica, Boiss. Fl. Orient. iii. 199 = graveolens.
brevanica, Bieb. Fl. Taur. Cauc. iii. 317, partim = dysenterica.
britannica, *Linn. Sp. Pl.* 882.—Europ.; As. bor.
Buhoniana, Hichw. Skizze, 147 = ensifolia.
Buhoniana, Hort. Prag. ex Tausch, in Syll. Ratisb. ii. (1828) 253 = stricta.
Buhoniana, Jacq. Fl. Austr. v. 26 = squarrosa.
canculus, Linn. Mant. ii. 471 = Printzia Bergii.
candyrina, Presl, Fl. Sic. p. xxix; Spreng. Syst. iii. 521 = montana.

INULA:—
campestris, Bess. Enum. Pl. Volh. 108 = I. Oculus-Christi.
canadensis, Bernh. Syst. Verz. Erf. 150 = Erigeron canadensis.
canariensis, *Mill. Gard. Dict.* ed. VIII. n. 12.—Ins. Canar.
candida, *Cass. Dict. Sc. Nat.* xxiii. (1826) 554.—Europ. austr.; As. Min.
candida, Ledeb. Fl. Ross. iii 502 = heterolepis.
caprusis, Spreng. Neue Entdeck. iii 43 = Pulicaria capensis.
Capps, *DC. Prod.* v. 469.—Reg. Himal.; Java; China.
caspica, *Ledeb. Ind. Sem. Hort. Dorpat.* (1822) 10.—Reg. Casp.
caucasica, Pers. Syn. ii. 450 = Baphthalmum speciosum.
cernua, Berg. Desc. Pl. Cay. 288 = Printzia Bergii.
chinensis, Rupr. ex Maxim. Prim. Fl. Amur. 149 = britannica.
chrysocomoides, Polr. Voy. Barb. ii. 239 = Pulicaria sicula.
cinerea, Lam. Encyc. iii. 259 = Vaillantii.
comosa, Lam. Fl. Fr. ii. 147 = britannica.
conferta flora, *A. Rich. Tent. Fl. Abyss.* i. 399.—Abyss.
Conyza, *DC. Prod.* v. 464.—Europ; As. occ.
conyzaea, Lam. Fl. Fr. ii. 149 = Pulicaria dysenterica
conyzoides, DC. Prod. vii. 283 (= *Varthemia montana*).—Syria.
conyzoides, Desf. Choix, 49 = Pulicaria calycina.
cordata, *Boiss. Diagn.* Ser. I. iv. 3.—Banat.; As. Min.
cordata, Freyn, ex Nym. Consp. 391 = salicina.
cordatofolia, Gillib. Fl. Lituan. i. 207 = Pulicaria dysenterica
cordato-germanica, Schur, Verh. Siebenb. Ver. Naturw. x. (1859) 64 = germanica.
coriacea, Schur, l. c. iv. (1858) 38 = salicina.
crasnfolia, Salisb. Prod. 201 = erithmoides.
cretica, Sch. Bip. ex Beck, in Denkschr. Acad. Wien, Math.-Nat. xliv. (1882) 225 = candida.
crispa, Pers. Syn. ii. 450 = Pulicaria crispa.
crispa, Sieber, ex DC. Prod. v. 479 - Pulicaria undulata.
crithmifolia, *Linn. Syst.* ed. X. 1219.—Europ.; Reg. Cauca.
crithmifolia, Willd. Sp. Pl. iii. 2101 = crithmoides.
crithmoides, *Linn. Sp. Pl.* 883.—Europ.; Reg. Mediterr.
crithmoides, Spreng. Syst. iii. 250 = multicaulis.
crocea, Krock, Fl. Siles. Suppl. ii. 313 = I Helenium ?
Castoi, *Borb. in Magyar Növenyt. Lapok*, vii. (1883) 41.—Hungar.
cuspidata, *C. B. Clarke, Comp. Ind.* 125.—Reg. Himal.
culchica, C. B. Clarke, l. c. 126 = grantioides.
cylindrica, Wtb. ex DC. Prod. v. 478 = Pulicaria vulgaris.
dentata, Sibth. & Sm. Fl. Graec. t. 874 = Pulicaria dysenterica.
dichotoma, Zuccagni, in Roem. Collect. 164 = britannica.
discoides, *Boiss. in Tchihat. Asie Min. Bot.* ii. 239. t. 20.—As. Min.
dioscorica, Neck. Delic. Gallo-Belg. ii. 356 = Pulicaria dysenterica.
divaricata, *Boiss. Fl. Orient.* iii. 200.—Persia; Scind.
dinaricata, Nutt. Gen. Am. ii. 152 = Aplopappus divaricatus.
dubia, Pourr. ex Timb. Précis Herbor. (1870) 176 = helenioides.
dubia, Thunb. Fl. Jap. 318 = Aster japonicus.
dumetorum, Salsm. ex DC. Prod. v. 479 = Pulicaria odora.
dysenterica, Georgi, Reise, i. 230 = britannica.
dysenterica, Linn. Sp. Pl. 882 = Pulicaria dysenterica.
enceloides, Hornem. ex Ledeb. Fl. Ross. ii. 506 = britannica.
ensifolia, *Linn. Sp. Pl.* 883.—Europ. austr.; Reg. Cauc.
ericoides, Torr. in Ann. Lyc. N. York, ii. (1828) 212 = Aster ericoides.
eriophora, DC. Prod. v. 470 = I. Cappa.
etrusca, Nym. Consp. 391, sphalm. = *hetrusca*.
eupatorioides, *DC. Prod.* v. 469.—Reg. Himal.; Burma.
falcata, Pursh, Fl. Am. Sept. ii. 532 = Chrysopsis villosa.
Falconeri, *Hook. f. Fl. Brit. Ind.* iii. 294.—Reg. Himal.
fasciculata, Gillib. Fl. Lituan. i. 209 = germanica.

INULA:—
flexuosa, *Boiss & Haussk. ex Boiss. Fl. Orient.* iii 200.—Persia; Scind
foetida, Dulac, Fl. Hautes-Pyr 516 = I. Conyza.
foetida, Linn. Syst ed. X. 1218 = Nidorella foliosa
foliosa, Schrad ex Nym Consp. 392 = squarrosa
fraglia, *Boiss. & Haussk ex Boiss Fl Orient* iii 197.—As. Min.
fruticosa, *Mill. Gard Dict* ed. VIII. n. 15.—Venezuela
fruticosa, Sch. Bip ex A. Rich. Tent. Fl. Abyss. i 396 = Arbuscula.
germanica, *Linn. Sp Pl.* 883.—Europ.; As Min
germanica, Vill Hist. Pl. Dauph iii. 219; Lam Encyc iii. 258 = squarrosa.
glabra, Bess. Hort. Crem., ex DC. Prod. v. 465 = bifrons.
glabra, Gillib. Fl Lituan. i 205 = salicina
glandulosa, Lam Encyc. iii. 259 = Chrysopsis gossypina ?
glandulosa, *Puschk. ex Willd. Sp. Pl.* iii. 2097.—Reg. Caucas.
glomerata, *Oliver & Hiern, in Oliver, Fl. Trop. Afr* iii. 359.—Afr. trop
glomeriflora, Lam Fl Fr. ii. 150 = bifrons.
glutinosa, Pers. Syn. ii. 450 = Grindelia glutinosa
gnaphalodes, Vent. Jard. Cels, t. 75 = Pulicaria gnaphalodes.
gossypina, Michx. Fl. Bor. Am. ii. 122 = Chrysopsis gossypina.
graminifolia, Michx. l. c = Chrysopsis graminifolia.
grandiflora, S. F. Gray, Nat Arr. Brit. Pl ii. 469 = I. Helenium.
grandiflora, *Willd. Sp. Pl* iii. 2096.—Reg. Himal et Caucas.
grandis, *Schrenk, Enum. Pl. Nov.* 50.—Songaria.
grantioides, *Boiss. Diagn.* Ser. II. iii. 14.—Ind. bor occ.
graveolens, *Desf. Fl Atlant.* ii. 275.—Reg. Mediterr.
Griffithii, *C. B. Clarke, Comp. Ind.* 126.—Reg. Himal
Hulleri, Durk. ex Beck, in Denkschr Acad Wien, Math. Nat. xliv (1882) 225 = Vaillantii.
Haussknechtii, Huter, in Oestr. Bot. Zeitschr. xiii (1863) 136 = hirta.
helenia, St. Lag in Ann. Soc. Bot. Lyon, vii. (1880) 128 = I. Helenium.
helenioides, *DC. Fl. Fr. Suppl.* 470.—Gall. austr.
Helenium, Asso, Syn. Stirp. Arag. 122 = helenioides
Helenium, Hook. f & Thoms. ex Fl. Brit Ind. iii. 292 = racemosa
Heleniun, *Linn Sp. Pl.* 881.—Europ.; As. bor.
helvetica, *Gaudin, Dec.* 17.—Helvet.
heterolepis, *Boiss. Diagn.* Ser. II. iii. 12.—Europ austr.
hetrusca, Moretti, Fl. Bras. Dec. iv. 14 = salicina.
hirsuta, Goeldem. Reise, ii. 72 = hirta.
hirsuta, Vitm. Summa, v. 84 = britannica.
hirta, *Linn. Sp. Pl.* 883.—Europ.; As bor.
hirta, Poll Hist. Pl. Palat. ii. 467 = britannica.
hispida, Schur, Enum. Pl. Transs. 314 = squarrosa.
Hookeri, *C. B. Clarke, Comp. Ind.* 122.—Reg. Himal
hybrida, Baumg. Enum. Stirp. Transs. iii. 132 = ensifolia.
incisa, Lam. Encyc. iii. 256 = Pulicaria undulata.
indica, Linn. Sp. Pl. ed II. 1236 = Vicoa auriculata.
indica, Russ. ex DC. Prod. v. 449 = Laggera pterodonta.
intermedia, Steud. Nom. ed. I. 482 = Chrysopsis gossypina ?
intermixta, *Kern. in Oestr. Bot. Zeitschr.* xxv (1875) 382.—Europ.
involucrata, Kalenic. in Bull. Soc. Nat. Mosc. xviii (1845) 1. 284 = hirta.
involucrata, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 171 = salicina.
japonica, *Thunb. Fl. Jap.* 318.—Japon.
Kalapani, *C. B. Clarke, Comp. Ind.* 123.—Reg. Himal.
laciniata, Lam. Fl. Fr. ii. 152 = Senecio uniflorus.
lanata, Banks, ex Pursh, Fl. Am. Sept. ii. 532 = Chrysopsis gossypina.
lanciflola, Wender. in Flora, xii. (1829) 1. Erg. 26 = ensifolia.
Langeana, Beck, in Denkschr. Acad Wien, Math.-Nat xliv. (1882) 225 = graveolens.
lanuginosa, Gillib. Fl. Lituan. i. 206 = I. Oculus-Christi latifolia, *DC. Prod.* v. 468.—Ind. or.
leptoclada, Webb, in Hook. Niger Fl. 137 = Vicoa auriculata.

INULA :—
limonifolia, *Boiss. Fl. Orient.* iii. 198.—Graec.
linariaefolia, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 134 = britannica.
limifolia, Wender. in Flora, xii. (1829) i. Erg. 26 = ensifolia.
litoralis, *Berb. ex Bech. in Denkschr. Acad. Wien, Math.-Nat.* xliv. (1882)311.—Europ.
lucens, Dulac. Fl. Hautes-Pyr. 516 = salicina.
macrocephala, Boiss. & Kotschy, ex Boiss. Fl. Orient. iii. 186 = Royleana?
macrolepis, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 330 = britannica.
macrophylla, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 488 = grandis.
macrophylla, *Sch. Bip. in Schweinf. & Aschers. Aufzachl.* 265, 307.—Abyss.
macrosperma, *Hook. f. Fl. Brit. Ind.* iii. 292.—Reg. Himal.
Mannii, *Benth. & Hook. f. Gen.* ii. 831.—Afr. trop.
Mariana, Linn. Sp. Pl. ed. II. 1240 = Chrysopsis Mariana.
media, Heb. Fl. Taur. Cauc. Suppl. 576 = germanica.
melanolepidea, Kalenic. in Bull. Soc. Nat. Mosc. xviii. (1845) i. 285 = hirta.
micrantha, Potr. Encyc. Suppl. iii. 153 = germanica.
micranthos, DC. Prod. v. 467 = britannica.
mollis, Bernh. ex Link, Enum. Hort. Berol. ii. 335 = suaveolens.
montana, Bal. & Bourg. ex Boiss. Fl. Orient. iii. 192 = Montbretiana.
montana, Hieb. Fl. Taur. Cauc. ii. 317 = I. Oculus-Christi.
montana, Bivona, Pl. Rar. Cent. i. 34 = Pulicaria calycina.
montana, *Linn. Sp. Pl.* 884.—Europ. austr.
montana, Lucé, Fl. Oail. 279 = salicina.
montana, Poll. Hist. Fl. Palat. ii. 469 = hirta.
Montbretiana, *DC. Prod.* vii. 284.—As. Min.
multicaulis, Boiss. Diagn. Ser. I. iv. 8 = anatolica.
multicaulis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 30.—Reg. Casp.
myrtifolia, Linn. Syst. ed. X. 1219 = squarrosa.
Neilreichii, A. Kern. Sched. Fl. Exsicc. i. 91; Beck, in Denkschr. Acad. Wien, Math.-Nat. xliv. (1882) 315 = sticta.
nervosa, *Wall. Cat.* n. 2960.—Reg. Himal.
nitida, Edgew. in Trans. Linn. Soc. xx. (1846) 68 = nervosa.
oblonga, DC. Prod. v. 469 = I. Cappa.
obovata, Nutt. ex Spreng. Syst. iii. 544 = Aster reticulatus.
obtusifolia, *Kern. in Ber. Naturw. Ver. Innsbruck*, i. (1870) 111.—Reg. Himal.
oboallata, Kit. in Linnaea, xxxii. (1863) 377 = hirta.
Oculus, Schrank, Baier. Fl. ii. 394 = I. Oculus-Christi.
Oculus-Christi, C. B. Clarke, Comp. Ind. 120 = obtusifolia.
Oculus-Christi, Lam. Encyc. iii. 254; Lapeyr. Hist. Abr. Fl. Pyr. 522 = helenioloides.
Oculus-Christi, *Linn. Sp. Pl.* 881.—Europ.; Oriena.
odora, Forsk. Fl. Aegypt. Arab. 150 = Pulicaria undulata.
odora, Linn. Sp. Pl. 881 = Pulicaria odora.
odora, Urv. in Mém. Soc. Linn. Par. i. (1822) 367 = montana.
Oettetiana, Reichb. in Moesal. Handb. ed. II. ii. 1506 = britannica.
orientalis, Lam. Encyc. iii. 255 = glandulosa.
orientalis, Urv. ex Boiss. Fl. Orient. iii. 193 = britannica.
orientalis, Willd. Sp. Pl. iii. 2100 = germanica.
oxylepis, Sch. Bip. ex Beck, in Denkschr. Acad. Wien, Math.-Nat. xliv. (1882) 325 = candida.
paludosa, DC. Prod. v. 480 = seq.
palustris, Link, Handb. i. 788 = Pulicaria paludosa.
parnassica, Boiss. & Heldr. Diag. Ser. II. iii. 11 = candida.
parviflora, Salisb. Prod. 201 = Pulicaria vulgaris.
phyllocoides, Willd. ex Spreng. Syst. iii. 536 = Tetramolopium phylloides.
pinifolia, Linn. Fl. Afr. Rar. 22; Amoen. Acad. vi. 402 = Senecio pinifolius.
planifolia, Link, ex DC. Prod. v. 479 = Pulicaria planifolia.
polycephala, Klatt, in Sitzb. Akad. Muench. (1878) 85 = capsidata.
polygonata, *DC. Prod.* v. 505.—Burma.

INULA :—
Portenschlagii, *Bech, in Denkschr. Acad. Wien, Math.-Nat.* xliv. (1882) 326.—Europ.
prenatica, Dum. Fl. Belg. 68 = germanica.
primulaefolia, *Lam. Encyc.* iii. 261.—Ins. S. Doming.
prostata, Gilib. Fl. Lituan. i. 205 = Pulicaria vulgaris.
provincialis, Linn. Sp. Pl. 884 = Senecio uniflorus.
Pseudo-Bubonium, Schur, Enum. Pl. Transs. 318 = salicina.
Pseudocappa, DC. Prod. v. 469 = I. Cappa.
Pseudogermanica, *Bech, in Denkschr. Acad. Wien, Math.-Nat.* xliv. (1882) 299.—Europ.
Pseudoglutinosa, Simkov. ex Beck, i. c. 303 = salicina.
Pulicaria, Brot. ex DC. Prod. v. 480 = Pulicaria paludosa.
Pulicaria, Linn. Sp. Pl. 882 = Pulicaria vulgaris.
Pulicaria, Urv. in Mém. Soc. Linn. Par. i. (1822) 367 = Pulicaria dysenterica.
pulicariaeformis, *DC. Prod.* vii. 284.—Persia.
punctata, Less. Syn. Comp. 159 = africana.
punctata, Muhl. Cat. 76 = Heterotheca grandiflora.
quadridentata, Lag. Nov. Gen. et Sp. 30 = graveolens.
quadrifida, Buch.-Ham. in Wall. Cat. n. 2965 = Pulicaria crispa.
racemosa, *Hook. f. Fl. Brit. Ind.* iii. 292.—Reg. Himal.
ramonssima, [Clairv.] Man. Herb. 244 = Vaillantii.
repanda, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 134 = britannica.
repens, Fisch. ex Treviran. Ind. Sem. Hort. Wratisl. (1819); Stev. ex Link, Enum. Hort. Berol. ii. 336 = Pulicaria dysenterica.
revoluta, Hoffmgg. & Link, Fl. Port. ii. 290 = Pulicaria revoluta.
rhizantha, *Tatarin. in Baer. & Helm. Beitr. Kenntn. Russ. Reich.* xx. 251.—Soongaria.
rhizocephala, *Schrenk, Enum. Pl. Nov.* 51.—Persia; Soongaria.
rhizocephaloides, *C. B. Clarke, Comp. Ind.* 124.—Reg. Himal.
rhodostemma, Boiss. Fl. Orient. iii. 172 = Lachnophyllum Lessmum.
rigida, Poell, Fl. Baden, iii. 1865 = hirta.
rigida, *Vill. Prosp.* 22 (Quid?).—Gallia.
Royleana, C. B. Clarke, Comp. Ind. 118 = racemosa.
Royleana, *DC. Prod.* v. 464.—Reg. Himal.
rubricaulis, *Benth. & Hook. f. Gen.* ii. 831.—Reg. Himal.
rupestris, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 169.—Afghan.
sallolifolia, *Gueldenst. Reise Russl.* ii. 78, 144, 146, 207, 237.—Kossia.
salicina, Bunge, Enum. Pl. Chin. Bor. 89. n. 214 = britannica.
salicina, C. B. Clarke, Comp. Ind. 123 = acuminata.
salicina, *Linn. Sp. Pl.* 882.—Europ.; As. bor.
salicina, Pall. Ind. Taur. [87]; ex Bieb. Fl. Taur.-Cauc. ii. 320 = squarrosa.
salsuloides, Sch. Bip. in Zoll. Syst. Verz. fnd. Archip. 122 = I. Cappa.
sarana, *Boiss. Fl. Orient.* iii. 199.—Cilicia.
Saturejae, Spreng. Syst. iii. 530 = Pectis saturejoides.
saturejoides, Mill. Gard. Dict. ed. VIII. n. 18 = Pectis saturejoides.
Savii, *Beck, in Denkschr. Acad. Wien, Math.-Nat.* xliv. (1882) 306.—Europ.
saxatilis, Lam. Fl. Fr. ii. 158 = viscosa.
scabra, Pursh, Fl. Am. Sept. ii. 581; Nutt. Gen. Am. ii. 131 = Heterotheca Lamarckii.
scabra, Wall. ex DC. Prod. v. 474 = Vicoa appendiculata.
schizogyne, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 199.—Ins. Canar.
Seidlitzii, *Boiss. Fl. Orient.* iii. 189.—Persia.
semiamplexicaulis, Rest. in Mém. Soc. Phys. Genève. vii. (1836) 169 = salicina.
semicordata, Borb.—Cf. Oestr. Bot. Zeitschr. xxix. (1879) 410 = rigida.
semihirta, Berb.—Cf. l. c. = rigida.
serica, Kit. ex Kanitz, in Verh. Zool. Bot. Ges. Wien, xiii. (1863) 692; St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = I. Oculus-Christi.
serrata, Gilib. Fl. Lituan. i. 207 = britannica.
serrata, Pers. Syn. ii. 450 = Grindelia inuloides.

INULA :—
setigera, *Beck. in Denkschr. Acad. Wien, Math.-Nat.* xliv. (1882) 331.—Europ.
shirensis, *Oliver, in Hook. fc. Pl.* t. 1399.—Afr. trop.
Simonsii, *C. B. Clarke, Comp. Ind* 121.—Reg. Himal.
somalensis, *Vatke, in Oestr. Boi. Zeitschr.* (1875) 326. —Afr. trop.
spathulata, Hort. Par. ex Pour. Encyc. Suppl. iii. 154 = sapera.
spiraeifolia, C. Koch, ex Boiss. Fl. Orient. iii. 188 = cordata.
spiraeifolia, Linn. Syst. ed. X. 1219 = squarrosa.
spuria, Kern. in Oestr. Bot. Zeitschr. (1871) 60 = rigida.
squarrosa, Bernh. ex DC. Prod. v. 464 = Conyza squarrosa.
squarrosa, Griseb. in Wiegm. Archiv, xviii. (1852) 337 = salicina.
squarrosa, Krock. Fl. Siles. ii. 11. 442 = britannica.
squarrosa, *Linn. Sp. Pl.* ed. II. 1240.—Europ.; Reg. Caucas.
Stoloschai, C. B. Clarke, Comp. Ind. 118 = Royleana.
stolonifera, J. Gay, ex DC. Prod. v. 849 = Aplopappus stolonifera.
stricta, *Tausch, in Syll. Ratisb.* ii. (1828) 253.—Hab.?
suaveolens, *Jacq. Hort. Vindob.* iii. 29. t. 51.—Europ. austr.
suaveolens, Pall. ex Bieb. Fl. Taur. Cauc. ii. 317 = I. Oculus Christi.
subauillaris, Lam. Encyc. iii. 259 = Heterotheca Lamarckii.
thapsoides, *Boiss. Fl. Orient.* iii. 190.—As. Min.; Persia.
thapsoides, *Spreng. Pugill.* i. 58.—Reg. Caucas.
Thomsoni, C. B. Clarke, Comp. Ind. 120 = obtusifolia.
transilvanica, Schur, Enum. Pl. Transs. 312 = germanica.
trichophylla, Nutt. Gen. Am. ii. 150 = Chrysopsis trichophylla.
Trixis, Linn. Syst. ed. X. 1219 = Trixis frutescens.
tuberosa, Lam. Encyc. iii. 260 = Jasonia tuberosa.
viligenea, Sibth. ex Sm. Fl. Brit. 892 = Pulicaria vulgaris.
undulata, Linn. Mant. i. 115 = Pulicaria undulata.
undulata, *Gilib. Fl. Lituan.* ii. 247.—Europ.
Vaillantii, Schur, Enum. Pl. Transs. 318 = britannica.
Vaillantii, *Vill. Hist. Fl. Dauph.* iii. 216.—Europ. austr.
valesiana, Tausch, ex Beck; Denkschr. Acad. Wien, Math.-Nat. xliv. (1882) 318 = hybrida.
verbascifolia, Potr. Encyc. Suppl. iii. 154 = thapsoides.
verrucosa, Klatt, in Sitzb. Akad. Muench. (1878) 86 = nervosa.
vestita, Wall. Cat. n. 2962 = Vicoa vestita.
villosa, Linn. Syst. ed. X. 1219 = montana.
villosa, Spreng. Syst. iii. 532 = Chrysopsis villosa.
villosa, Vahl, ex Homem. Hort. Hafn. ii. 822 = Pulicaria arabica.
vicidula, *Boiss. & Kotschy, in Boiss. Fl. Orient.* iii. 188.—Armenia.
viscosa, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 228.—Reg. Mediterr.
Vrabelyiana, Kern. in Oestr. Bot. Zeitschr. xviii. (1868) 297 = sticta.
vulgaris, Trevis. Prosp. Fl. Eugan. (1842) 29 = Pulicaria vulgaris.

INULASTER, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) 1. Intell. 42; in Flora, xxv. (1842) 1. Beibl. 134, 147, nomen; et in A. Rich. Tent. Fl. Abyss. i. 299 (1847) = Inula, Linn. (Compos.).
Kotschyi, Sch. Bip. ll. cc. 42; 134; 147 = Vicoa auriculata.
macrophyllus, Sch. Bip. in A. Rich. Tent. Fl. Abyss. i. 299 = I. macrophylla.

INVOLUCRARIA, Ser. in Mém. Soc. Phys. Genève. iii. 1. (1826) 25. t. 5 = Trichocoronatthes, Linn. (Cucurb.).
Anguina, M. Roem. Syn. Pepon. 97 = T. Anguina.
cordata, M. Roem. l. c. = T. cordata.
cucumerina, M. Roem. l. c. 98 = T. pubera.
globosa, M. Roem. l. c. 98 = T. globosa.
javanica, Miq. Fl. Ind. Bat. I. i. 678 = Thladiantha cordifolia.
Leptocarpa, Naud. in Huber, Cat. (1866) 11 = T. Leptocarpa.
Modecca, M. Roem. Syn. Pepon. 98 = T. pubera.
palmata, M. Roem. l. c. 98 = T. pubera.
trifoliata, M. Roem. l. c. 99 = T. trifoliata.

INVOLUCRARIA :—
Wallichiana, Ser. in Mém. Soc. Phys. Genèv. iii. t. (1825) 25 = Trichosanthes Wallichiana.
Wightiana, M. Roem. Syn. Pepon. 98 = T. pubera.

IOCASTE, E. Mey. ex DC. Prod. vii. 298 (1838) = **Phymaspermum**, Less. (Compos.).

ICACULON, Raf. New Fl. Am. ii. 53 (1836) = **Crotalaria**, Tourn. (Leguminos.)
cuneifolia, Raf. l. c. 55 = C. cuneifolia.
linearis, Raf. l. c. 54 = C. linearis.
longipes, Raf. l. c. = C. longipes, *Raf.*
lunulata, Raf. l. c. 55 = C. lunulata.
ovalis, Raf. l. c. = C. ovalis.
pilosa, Raf. l. c. 54 ~ C. pilosa, *Raf.*
pumila, Raf. l. c. 56 = C. pumila, *Kaufm.*
sagittalis, Raf. l. c. = C. sagittalis.

IOCHROMA, Benth. in Bot. Reg. xxx. (1845) t. 20. *SOLANACEAE*, Benth. & Hook. f. ii. 895.
CHARNSTHUS, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 336.
CLEOCHROMA, Miers, l. c. vii. 349 (1848).
DIPLURION et VALTETA, Raf. Sylva Tellur. 53 (1838) nomina, priora.
arborea, *Griseb. in Goett. Abh.* xix. (1874) 219.—Reg. Argent.
australe, *Griseb. l. c.* 218.—Reg. Argent.
Benthamiana, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 138.—Peruv.
calycina, *Benth. in Bot. Reg.* (1845) subt. 20.—Guian.
cocuinea, *Schrad in Fl. des Serres*, Sér. II. ii. (1857) 121. t. 1261.—Mexic.
corallofolia, *Miers, in Journ. Lond. Journ. Bot.* vii. (1848) 347.
fuchsioides, *Miers, l. c.* 345.—Peruv.
grandiflora, *Benth. in Bot. Reg.* (1845) sub t. 20.—Peruv.
lanceolata, *Miers, in Hook. Lond. Journ. Bot* vii. (1848) 347.—Ecuador.
longipes, *Miers, l. c.* 344.—Ecuador.
loxensis, *Miers, l. c.* 347.—Ecuador.
macrocalyx, *Miers, l. c. iv.* (1845) 339.—Peruv.
Schiechtendaliana, *Dun. in DC. Prod.* xiii. 1. 673.—Am. trop.
tetradynama, *Dun. l. c.* 490.—Am. austr.
tubulosa, *Benth. in Bot. Reg.* (1845) t. 20.—N. Granat.
umbrosa, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 346.—N. Granat.
Warscewiczii, *Regel, Gartenfl.* (1855) 245. t. 130.—Peruv.

IODANTHUS, Torr. & Gray, Fl. N. Am. i. 72 (1838). *CRUCIFERAE*, Benth. & Hook. f. i. 70.
hesperidoides, Torr. & Gray, in A. Gray, Man. U. St. Bot. ed. I. 33 = pinnatifidus.
pinnatifidus, *Steud. Nom.* ed. II. i. 812.—Am. bor.

IODES, Blume, Bijdr. 29 (1826) (Ioedes.). *OLACINEAE*, Benth. & Hook. f. i. 355.
ERYTHROSTAPHYLIS, Hance, in Journ. Bot. xi. (1873) 266.
africana, *Weiw. ex Oliver, Fl. Trop. Afr.* i. 358.—Afr. trop.
Brandisii, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 298.—Burma.
cirrhosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 261.—Java.
Hookeriana, *Baill. Adansonia*, x. (1871-72) 268.—Ind. or.
Horsfieldii, *Baill. l. c.* 267.—Java.
madagascariensis, *Baill. l. c.* ix. (1868-70) 146.—Madag.
oblonga, *Planch. ex Mast. in Hook. f. Fl. Brit. Ind.* i. 587.—Penang.
ovalis, *Blume, Bijdr.* 30.—Java.
Thomsoniana, *Baill. Adansonia*, x. (1871-72) 270.—Ind. or.
tomentella, Miq. Fl. Ind. Bat. i. 1. 796 = ovalis.

IODINA, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 178. *SANTALACEAE*, Benth. & Hook. f. iii. 228.
cuneifolia, *Miers, in Journ. Linn. Soc.* xvii. (1878) 84.—Reg. Argent.

IODINA :—
rhombifolia, *Hook. & Arn. ex Reissek, in Mart. Fl. Bras* xi. 1. 78.—Bras.
ruscifolia, Walp. Rep. i. 541 = rhombifolia.

ION, Medic. Malv. 102 (1787) = **Viola**, Tourn.
tricolor, Medic. l. c. = V. tricolor.

IONCOMELOS, Raf. Fl. Tellur. iii. 62 (1836) = **Solla**, Linn. (Liliac.).
purpureus, Raf. l. c. = S. Copani.

IONDRABA, Reichb. Consp. 183 (1828) = JONDRABA, Medic. = **Biscutella**, Linn. (Crucif.).

IONE, Lindl. Fol. Orch. (1853) = **Bulbophyllum**, Thou. (Orchid.).
bicolor, Lindl. l. c. 3 = B. bicolor †
candida, Lindl. l. c.—Reg. Himal.
cirrhata, Lindl. l. c. 1.—Reg. Himal.
fusco-purpurea, Lindl. l. c. 2 = B. fusco-purpureum.
khasiana, Lindl. l. c. = B. repens †
palacea, Lindl. l. c. = B. palaceum.
virens, Lindl. l. c. 1.—Reg. Himal.

IONIA, Pers. ex Steud. Nom. ed. I. 482 (1821) = Ionidium, Vent. (Violar.).

IONIDIOPSIS, Walp. Ann. i. 69 (1848) = JONIDIOPSIS, Presl = **Noisettia**, H. B. & K. (Violar.).

IONIDIUM, Vent. Jard. Malm. t. 27 (1803). *VIOLARIEAE*, Benth. & Hook. f. i. 117.
CALCHOLARIA, Loefl. It. 188 (1758) nomen, prius.
CUSELIUM, Raf. Cat. 18 (1824).
HYBANTHUS, Jacq. Enum. Pl. Carib. 2 (1760).
PIGEA, DC. Prod i. 307 (1824).
POMBALIA, Vand. Fasc. Pl 7 (1771).
SOLEA, Spreng. in Schrad. Journ. Bot. iv. (1800) 192.
VLAMINGIA, De Vriese, in Lehm. Pl. Preiss i. 398 (1845).
aethiopicum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 330.—Afr. trop.
album, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 477.—Bras.
angustifolium, H. B. & K. Nov. Gen. et Sp. v. 377 = oppositifolium.
anomalum, *H. B. & K. l. c.* 381. t. 500.—N. Granat.
atropurpureum, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 490.—Bras.
attenuatum, Willd. ex Roem. & Schult. Syst. v. 402 = oppositifolium.
Aubletii, Roem. & Schult. l. c. 395 = Corynostylis Hybanthus.
auranticum, *F. Muell. ex Benth. Fl. Austral.* i. 102.—Austral.
Australasicum, Behr, in Linnaea, xx. (1847) 629 = floribundum.
austro-caledonicum, *Virill. in Bull. Soc. Linn. Normand.* ix. (1865) 332.—N. Caled.
Balbisii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 556.—N. Granat.
Banksianum, Steud. Nom. ed. II. i. 813 = heterophyllum.
barselonense, Karst. Fl. Columb. i. 189 = oppositifolium.
bicolor, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 475.—Bras.
bigibbosum, *A. St. Hil. l. c.* 488.—Bras.
Botterii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 556.—Mexic.
brevicaule, *Mart. Spec. Mat. Med. Bras.* 15; *Reise Bras.* i. 553.—Bras.
brevifolium, *Benth. Fl. Austral.* i. 102.—Austral.
buxifolium, *Vent. Jard. Malm.* 27 in adn.—Madag.
caffrum, *Sond. in Linnaea*, xxiii. (1850) 18.—Afr. austr.
Calcolaria, Vent. Jard. Malm. sub 27 = I. lpecacuanha.
calceolarium, *Ging. in DC. Prod.* i. 311.—Mexic.
calycinum, *Steud. Nom.* ed. II. i. 813 —Austral.
capense, *Roem. & Schult. Syst.* v. 393.—Afr. austr.
chamaedrifolium, *Presl, Rel. Haenk.* ii. 96.—Chili.
chamaedryfolium, Poepp. ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 360 = glutinosum.
circaeoides, *H. B. & K. Nov. Gen. et Sp.* v. 379. t. 496. —Am. austr.
claytonioides, *Roem. & Schult. Syst.* v. 402.—Hab. ?
commune, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 469.—Bras.
concolor, *Benth. & Hook. f. ex S. Wats. Bibliog. Ind. N. Am. Bot.* 31.—Am. bor.

IONIDIUM :—
debilissimum, *F. Muell. Fragm.* xi. 4.—Austral.
diandrum, Roem. & Schult. Syst. v. 399 = Corynostylis diandra.
diffusum, Gill ex Hook. Bot. Misc. iii. (1833) 145 = Cynanchum diffusum.
elatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 556.—Mexic.
ennaspermum, Vent. Jard. Malm. sub 27; Ging. in DC. Prod. i. 308 = heterophyllum.
erectum, Ging. l. c. 311 = heterophyllum.
ericaeoides, Fritz. Ic. Ind. 886, sphalm. = circaeodes.
filiforme, *F. Muell. Vict. Pl.* i. 66, 324.—Austral.
floribundum, *Walp. Rep.* ii 767.—Austral.
frutescens, Blume, Bijdr. 58 = heterophyllum.
frutescens, *Ging. in DC. Prod.* i. 311.—Ind. or.
fruticulosum, *Benth. Bot. Voy. Sulph. T.* t. 3.—Calif Galeottii, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii (1854) ii. 339.—Mexic.
glaucum, Steud. Nom ed. II. i. 813 = calycinum.
glutinosum, *Vent. Jard. Malm.* sub 27.—Reg. Argent.
gracile, *Mog. & Sesse, in DC Prod* i. 309.—Mexic.
grandiflorum, *Turcz. in Bull. Soc. Nat Mosc.* xxvii. (1854) ii. 338.—Venezuela.
guaraniticum, A. St. Hil. in Mém. Mus. Par. xi. (1824) 474 = commune.
heterophyllum, *Vent. Jard. Malm.* 27, in adn.—Geront. trop
heterosepalum, *Eichl. in Mart. Fl. Bras.* xiii. t. 367.—Bras.
hexaspermum, Dalz. in Hook. Kew Journ. iv. (1852) 342 = suffruticosum.
Hieronymi, *Griseb. in Goett. Abh.* xxiv. (1879) 21.—Reg. Argent.
hirtum, Klotzsch, in Peters, Reise Mossamb. Bot. 148 = heterophyllum.
hispidulum, *Presl, Bot. Bemerk.* 13.—Hab. ?
Hybanthus, Roem. & Schult. Syst. v. 396 = Corynostylis Loeflingii.
ilicifolium, *Vieill. in Bull. Soc. Linn. Normand* ix. (1865) 334.—N. Caled.
indecorum, *A. St. Hil. in Mém. Mus. Par.* ix. (1822) 329.—Bras.
Ipecacuanha, *Vent. Jard Malm.* sub t. 27.—Bras.
Itoubou, Steud. Nom. ed. II. i. 813 = seq.
Itubu, H. B. & K. Nov. Gen. et Sp. v. 380 = I. Ipecacuanha.
Jacquinianum, *Roem. & Schult. Syst.* v. 397.—Cuba.
lanatum, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 482.—Bras.
lanceolatum, *Presl, Rel. Haenk.* v. 96.—Chili.
leptorhizum, DC. ex Ging. in DC. Prod. i. 306 = heterophyllum.
linarifolium, Ging. l. c. 309 = strictum.
linarioides, Presl, Bot. Bemerk. 12 = filiforme.
lineare, *Torr. in Ann. Lyc. N. York*, ii. (1828) 168.—Am. bor.
lineatum, *Ging. in DC. Prod.* i. 311.—Cuba.
lintfolium, *DC. l. c.* 309.—Madag.
lobelioides, *Schlecht. in Linnaea*, xii. (1838) 208.—Mexic.
Loeflingianum, Roem. & Schult. Syst. v. 396 = I. Ipecacuanha.
longifolium, *Mog. & Sesse, ex DC. Prod.* i. 311.—Mexic.
longifolium, Roem. & Schult. Syst. v. 398 = Noisettia longifolia.
Lorentzianum, *Eichl. ex Griseb. in Goett. Abh.* xix. (1874) 74.—Reg. Argent.
Maximiliani, *Eichl. in Mart. Fl. Bras.* xiii. t. 369.—Bras.
microphyllum, H. B. & K. Nov. Gen. et Sp. v. 374. t. 495 = glutinosum.
molle, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 338.—Java.
monopetalum, Roem. & Schult. Syst. i. 400 = filiforme.
multiflorum, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 340 = floribundum.
nanum, A. St. Hil. in Mém. Mus. Par. xi. (1824) 484 = album.
nanum, *Haro. in Haro. & Sond. Fl. Cap.* ii. 585.—Afr. austr.
occultum, *Polak. in Linnaea*, xli. (1877) 548.—Costa Rica.
oppositifolium, *Roem. & Schult. Syst.* v. 395.—Venezuela.
orchidiflorum, Roem. & Schult. l. c. 400 = Noisettia longifolia.
parietariaefolium, DC. Prod. i. 308 = oppositifolium.

IONIDIUM :—
parviflorum, Vent. Jard. Malm. 27, in Adn. = glutinosum.
phyllanthoides, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 125.—N. Granat.
Poiya, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 482.—Bras.
polygalaefolium, *Vent. Jard. Malm.* 27. t. 27 — Mexic.
procumbens, *Griseb Cat. Pl. Cub.* 11.—Cuba.
racemosum, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1825) 49.—Bras.
ramosissimum, *Thw. Enum. Pl. Zeyl.* 21.—Zeylan.
rhabdospermum, Hochst. in Flora, xxiv. (1841) i. Intell. 43 = heterophyllum ?
riparium, A. Gray, Pl. Wright. ii. 16 = lineare.
riparium, H. B. & K. Nov. Gen. et Sp. v. 378 = oppositifolium.
scariosum, A. St. Hil. in Mém. Mus. Par. xi. (1824) 478 = brevicaule.
setigerum, *A. St. Hil. l. c.* 470 —Bras.
Sprengelianum, Roem. & Schult. Syst. v. 401 = concolor.
Spruceì, *Eichl. in Mart. Fl. Bras.* xiii. t. 378.—Bras.
stipulaceum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 144 = lineare.
stipulare, Roem. & Schult. Syst. v. 892 = Viola stipularis.
strictum, *Vent. Jard. Malm.* 27, in Adn.—Ina. S. Doming.
suffruticosum, Ging. in DC Prod. i. 311 = heterophyllum.
sylvaticum, A. St. Hil. in Mém. Mus. Par. xi. (1824) 472 = commune.
tenuifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 337.—Bras.
thesiifolium, Roem. & Schult. Syst. v. 398 = heterophyllum.
thymifolium, *Presl, Rel. Haenk* ii. 91.—Afr. austr.
travancoricum, *Bedd. Ic Pl. Ind. Or.* i. 55.—Ind. or.
urticaefolium, Mart. Spec. Mat. Med. Bras. 16; Reise Bras. i. 353 = circaeoides.
verbenaceum, *H. B. & K. Nov. Gen. et Sp.* v. 370 t. 497.—Mexic.
Vernonii, *F. Muell. Pl. Vict.* i. 232.—Austral.
verticillatum, Roem. & Schult. Syst. v. 399 = polygalaefolium.
villosissimum, A. St. Hil. in Mém. Mus. Par. xi. (1824) 480.—Bras.
viscidulum, H. B. & K. Nov. Gen. et Sp. v. 377 = oppositifolium.
Wightiana, Wall. Cat. n. 4020 = heterophyllum.
Zippelii, *Oudem. in Mig. Ann. Mus. Bot. Lugd. Bat.* iii. 73.—Ins. Timor.

IONIRIS, Baker, in Gard. Chron. N. S. vi. (1876) 149, in obs. = JONIRIS, Klatt = **Iris**, Tourn.

IONOPSIDIUM, Reichb. Ic. Fl. Germ. vii. 26. t. 649 (1829) = **Cochlearia**, Tourn. (Crucif.).
acaule, Reichb. Ic. Fl. Germ. vii. 26 = C. acaulis.
albiflorum, Dur. in Duch. Rev. Bot. ii. (1846) 433.—Algeria.
Savianum, Ball, ex Caruel, in Atti Soc. Sc. Nat. vili. (1865) 439 = Bivonaea Saviana.

IONOPSIS, H. B. & K. Nov. Gen. et Sp. i. 348. t. 88 (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 567.
CYRTION, Spreng. Syst. iii. 679, 721 (1826).
IANTHA, Hook. Exot. Fl. t. 118 (1832).
brevifolia, *Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 26.—Mexic.
Burchellii, *Reichb. f. in Linnaea*, xli. (1877) 66.—Bras.
Gardneri, *Lindl. in Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 13, in obs.—Ind. occ.
pallidiflora, *Lindl. Bot. Reg.* sub t. 1904.—Ins. Trinit.
paniculata, *Lindl. l. c.*—Bras.
pulchella, *H. B. & K. Nov. Gen. et Sp. i.* 348. t. 88.—N. Granat.
satyrioides, Reichb. f. in Walp. Ann. vi. 688 = teniculata.
tenera, Lindl. Bot. Reg. t. 1904 = utriculariioides.
teres, *Lindl. l. c.* (1838) Misc. 96.—Guian.
teniculata, *Lindl. Gen. et Sp. Orch.* 193.—Jamaic.
utriculariioides, *Lindl. Coll. Bot.* t. 39 A.—Jamaic.
xonalis, Lindl. in Lindl. & Paxt. Flow. Gard. ii. (1851–52) 13. in obs. = utriculariioides.

IONOSMANTHUS, Jord. & Fourr. Ic. Pl. Europ. ii. 13 (1870) = **Ranunculus**, Tourn.
plantagineus, Jord. & Fourr. l. c. 14 = R. bupleurifolius ?
rhomboifolius, Jord. & Fourr. l. c. = R. bullatus.
semicalvus, Jord. & Fourr. l. c. = R. bullatus.

IONTHLASPI, Adans. Fam. ii. 423 (1763) = **Clypeola**, Linn. (Crucif.).
clypeolatum, Link, Handb. ii. 296 = C. Jonthlaspi.
eriophorum, Link, l. c. = C. eriocarpa.

IOSOTOMA, Griseb. Fl. Brit. W. Ind. 388 (1864), err. typ. = **Isotoma**, Lindl. (Campan.).

IOSTEPHANE, Benth. in Benth. & Hook. f. Gen. ii. 568 (1873). *COMPOSITAE*, Benth. & Hook. f. i. c. heterophylla, *Benth. l. c.*—Mexic.
trilobata, *Hemsl. Biol. Centr. Am. Bot.* ii. 169.—Mexic.

IOXYLON, Rafin. in Am. Monthly Mag. iv. (1818) 195 = TOXYLON, Rafin. = **Maclura**, Nutt. (Urtic.).

IOZOSMENE, Lindl. Introd. Nat. Syst. ed. II. 202 (1836) = IOZOSTE, Nees = **Actinodaphne**, Nees (Laurin.).

IOZOSTE, Nees, in Wall. Pl. As. Rar. ii. 68 (1831) = **Actinodaphne**, Nees (Laurin.).
acuminata, Plume, Mus. Bot. Lugd. Bat. i. 264 = A. acuminata.
chinensis, Blume, l. c. = A. chinensis.
lancifolia, Blume, l. c. = A. lancifolia.
pedunculata, Blume, l. c. = A. pedunculata.
rotundifolia, Nees, in Wall. Pl. As. Rar. ii. 68 = A. chinensis.

IPECACUANA, Rafin. Sylva Tellur. 147 (1838) = seq.

IPECACUANHA, Arruda, Diss. 44 (1810); ex A. St. Hil. Fl. U. St. Bras. t. 6 (1824) = **Psychotria**, Linn. (Rubiac.).
Brasica. Arruda, Diss. ex Koster, Trav. Braz. 497 = Ionidium Ipecacuanha.
fusca, Rafin Sylva Tellur. 147 = P. Ipecacuanha.
officinalis, Arruda, Diss. ex Koster, Trav. Braz. 497 = P. Ipecacuanha.
Prota, Arruda, l. c. = P. Ipecacuanha.

IPHEION, Rafin Fl. Tellur. ii. 12 (1836) = **Milla**, Cav. (Liliac.).
uniflorum, Rafin. l. c. = M. uniflora.

IPHIGENIA, Kunth, Enum. Pl. iv. 212 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 824.
APHOMA, Rafin. Fl. Tellur. ii. 31 (1836).
NEPHROCLAS, Salisb. Gen. Pl. Prael. 84 (1866).
caricina, Kunth, Enum. Pl. iv. 214 = indica.
guineensis, *Baker, in Journ. Linn. Soc.* xvii. (1879) 461.—Afr. trop.
indica, *A. Gray, in Kunth, Enum. Pl.* iv. 213.—Ind. or.
Novae-Zelandiae, *Baker, in Journ. Linn. Soc.* xvii. (1879) 451.—N. Zel.
pallida, *Baker, l. c.*—Ind. or.
racemosa, A. Gray, in Kunth, Enum. Pl. iv. 213 = indica.
tenuis, *Baker, in Journ. Linn. Soc.* xx. (1883) 275.—Madag.

IPHIONA, Cass. in Bull. Soc. Philom. (1817) 158; et in Dict. Sc. Nat. xxiii. 600 (1823). *COMPOSITAE*, Benth. & Hook. f. ii. 333.
BLANCHEA, Boiss. Fl. Orient. iii. 219, in syn. (1875).
CARPHOPAPPUS, Sch. Bip. in Walp. Rep. ii. 954 (1843).
KARTHEMIA, Sch. Bip. l. c. 953 (1843), sphalm.
VARTHEMIA, DC. Prod. v. 473 (1836).
acuminata, *Benth. & Hook. f. Gen.* ii. 333.—Afr. austr.
arabica, *Benth. & Hook. f. l. c.*—Arab.
baccharidifolia, *Benth. & Hook. f. l. c.*—Afr. austr.
dubia, Cass. in Bull. Soc. Philom. (1817) 158 = juniperifolia.
horrida, *Boiss. Diagn.* Ser. I. vi. 80.—Arab.

IPHIONA :—
lilicifolia, *Benth. & Hook. f. Gen.* ii. 333.—Madag.
iphionoides, *Benth. & Hook. f. l. c.*—Syria.
juniperifolia, *Cass. in Dict. Sc. Nat.* xxiii. 610.—Aegypt.; Arab.
microphylla, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1877) 196.—Afr. trop.
persica, *Benth. & Hook. f. Gen.* ii. 333.—Persia.
polygalifolia, *Benth. & Hook. f. l. c.*—Afr. austr.
punctata, Cass. in Dict. Sc. Nat. xxiii. 609 = Inula africana.
radiata, *Benth. in Henders. & Hume, Lahore*, 333.—Reg. Himal.
rotundifolia, *Benth. & Hook. f. Gen.* ii. 333; *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 360.—Afr. trop.
scabra, *DC. ex Decne. in Ann. Sc. Nat.* Sér. II. iii. (1834) 368.—Arab.; Aegypt.
spinosa, DC. ex Stend. Nom. ed. II. i. 815 = juniperifolia.

IPHISIA, Wight & Arn. Contrib. 32 (1834) = **Tylophora**, R. Br. (Asclep.).
Govani, Wight, l. c. = T. Govanii.
multiflora, Wight & Arn. l. c. = T. Iphisia.

IPNUM, Phil. Sert. Mendoc ii 53 (1871). *GRAMINEAE*, Benth. & Hook. f. iii. 1187.
mendocinum, *Phil. l. c.*—Chili.

IPO, Pers. Syn. ii. 566 (1807) = **Antiaris**, Lesch. (Urtic.).
toxicaria, Pers. l. c. = A. toxicaria.

IPOMAEA, Burm. f. Fl. Ind. 50 (1768) = **Ipomoea**, Linn. (Convolv.).

IPOMEA, All. Misc. Taur. v. 62 (1773) = praec.

IPOMERIA, Nutt. Gen. Am. i. 124 (1818) = **Gilia**, Ruiz & Pav. (Polem.).
aggregata, Nutt. l. c. = G. aggregata.
alba, Nutt. l. c. = G. laciniata.
coronopifolia, Nutt. l. c. = G. coronopifolia.
inconspicua, Nutt. l. c. = G. parviflora.

IPOMOEA, Linn. Syst. ed. I (1735). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 870.
ANISIA, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 431 (Conv. Or. 99).
APOPLEUMON, Rafin. Fl. Tellur. iv. 72 (1836).
BATATAS, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 434 (Conv. Or. 52.)
BOMBYCOSPERMUM, Presl, Rel. Haenk. ii. 127. t. 71 (1831).
BONANOX, Rafin. Fl. Tellur. iv. 77 (1836).
CALEOA, Cav. Ic. v. 51. t. 476 (1799).
CALONYCTION, Choisy, in Mém. Soc. Phys. Genèv vi. (1833) 441 (Conv. Or. 59).
CALYCANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 243. t. 40 (1861).
CAMONEA, Rafin. Fl. Tellur. iv. 81 (1836).
CAULOTILIS, Rafin. l. c. 75 (1836).
CLEISTERA, Rafin. l. c. 77 (1836).
CLITOCYAMOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128, 183.
COILADENA, Rafin. Fl. Tellur. ii. 12 (1836).
CONVOLVULOIDES, Moench, Meth. 451 (1794).
DECALOBA, Rafin. Fl. Tellur. iv. 76 (1836).
DIATREMA, Rafin. l. c. 75 (1836).
DISTIMAKE, Rafin. l. c. 82 (1836).
DOREMA, Rafin. l. c. 75 (1836).
ELYTHROSTAMNA, Boj. Hort. Maurit. 230 (1837).
EURYLOMA, Rafin. Fl. Tellur. iv. 75 (1836).
EXOGONIUM, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 448 (Conv. Or. 61).
FRAXIMA, Rafin. Fl. Tellur. iv. 83 (1836).
GYNOSIA, Rafin. l. c. 75 (1836).
GOMPHIPUS, Rafin. l. c. (1836).
KOLOPONIA, Rafin. l. c. 73 (1836).
LEPTOCALLIS, G. Don, Gen. Syst. iv. 260 (1837).
LORAKE, Rafin. Fl. Tellur. iv. 81 (1836).
MACROSTEMA, Pers. Syn. i. 185 (1805).
MARCELLIA, Mart. ex Choisy, in Mém. Soc. Phys. Genèv. x. (1841) 448.
MELASCUS, Rafin. Fl. Tellur. iv. 80 (1836).
MIMA, Cerv. in La Llave & Lex. Nov. Veg. Descr. fasc. i. 3 (1824).

IPOMOEA :—
MODESTA, Rafin. Fl. Tellur. iv. 75 (1836).
MORNOA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 5 (1824).
NEMANTHERA, Rafin. Fl. Tellur. iv. 80 (1836).
NEOTHOSIS, Rafin. l. c. 125 (1836).
NIL, Medic. in Staatsw. Vorles. Churpf. Phys. Oek. Gen. i. (1791) 210.
OPERCULINA, Silva Manso, Enum. Subst. Bras. 16 (1836).
ORNITHOSPERMA, Rafin. Fl. Ludov. 149 (1817).
PHARBITIS, Choisy, in Mém. Soc. Phys. Genève. vi. (1833) 438 (Conv. Or. 56).
PIPTOSTIGIA, Reichb. Consp. 118 (1841).
QUAMOCLIT, Moench, Meth. 463 (1794).
SKINNERIA, Choisy, in Mém. Soc. Phys. Genève. vi. (1833) 487 (Conv. Or. 105. t. 6).
SPIRANTHERA, Boj. Hort. Maurit. 226 (1837).
STOMADENA, Rafin. Fl. Tellur. ii. 12 (1836).
TERRIETRA, Rafin. l. c. iv. (1836).
TIRTALIA, Rafin. l. c. 71 (1836).
TREMASPERMA, Rafin. l. c. 124 (1836).
TRIVOLVULUS, Moç. & Sesse, ex Choisy, in DC. Prod. ix. 368 (1845).
TURBINA, Rafin. Fl. Tellur. iv. 81 (1836).
TURPITHUM, Rafin. l. c. 71 (1836).
abrupta, *R. Br. Prod.* 485.—Austral.
abscisa, Boj. ex Choisy, in DC. Prod. ix. 438, 392 = Breweria abscissa.
abutiloides, *G. Don, Gen. Syst.* iv. 278.—Ecuador.
abyssinica, *Schweinf. Beitr. Fl. Aethiop.* 94.—Abyss.
acanthocarpa, *Hochst. ex Choisy, in DC. Prod.* ix. 346, 392.—Afr. trop.
acetosaefolia, Roem. & Schult. Syst. iv. 246 - carnosa.
acetosaefolia, *Choisy, in DC. Prod.* ix. 362.—Ins. Maurit.
aculeata, Blume, Bijdr. 715 = I. Bona-nox.
acuminata, *Roem. & Schult. Syst.* iv. 228.—Ind. occ.
acuminata, Ruiz & Pav. Fl. Per. ii. t. 120. f. b = coccinea.
acutangula, Ruiz & Pav. l. c. 11 = coccinea.
acutiflora, *A. Rich. Tent. Fl. Abyss.* ii. 71.—Abyss.
adspersa, Mart. ex Meisn. in Mart. Fl. Bras. vii. 252 = polymorpha.
aegyptia, Lam. Sp. Pl. 162 = pentaphylla.
affinis, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 268.—Mexic.
affinis, Poepp. ex Choisy, in DC. Prod. ix. 392, 438 = Calystegia Soldanella.
afra, *Choisy, in DC. Prod.* ix. 340.—Afr. trop.
Afzelii, *Choisy, l. c.* 373.—Afr. trop.
aggregata, *Pov. Encyc. Suppl.* iv. 683.—Cochinch.
Altoni, *Lindl. Bot. Reg.* t. 1794.—Afr. austr.
alata, *R. Br. Prod.* 484.—Austral.
alatipes, *Hook. Bot. Mag.* t. 5380. – Venezuela.
alba, Garcke, in Linnaea, xxii. (1849) 67 = fastigiata.
alba, Linn. Sp. Pl. 161 = I. Bona-nox.
albiflora, *Mertc. Pl. Nouv. Am.* 114. t. 70.—Bras.
albivena, *Smeet, Hort. Brit.* ed. II. 372.—Afr. austr.
alternifolia, *Griseb. Cat. Pl. Cub.* 202.—Cuba.
altissima, Bert. ex G. Don, Gen. Syst. iv. 278 = racemosa.
altissima, *Mart. ex Choisy, in DC. Prod.* ix. 350.—Bras.
alulata, *Miq. in Linnaea*, xviii. (1844) 599.—Cuian.
ambigua, Endl. Prod. Fl. Norf. 33 = I. Bona-nox.
amoena, Blume, Bijdr. 718 = Argyreia Roxburghii.
amoena, *Choisy, in DC. Prod.* ix. 365.—Afr. trop.
amplexipafolia, G. Don, Gen. Syst. iv. 279 = quinquefolia ?
amplista, *Choisy, in DC. Prod.* ix. 361.—Mexic.
anceps, Roem. & Schult. Syst. iv. 231 = I. Turpethum.
angularis, Choisy, in Mém. Soc. Phys. Genève. vi. (1833) 454 = vitifolia.
angularis, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 197 = hederifolia.
angularis, Willd. in Nov. Act. Nat. Cur. iv. (1770) 197 = coccinea ?
angulata, *Mart. ex Choisy, in DC. Prod.* ix. 371.—Bras.
angulata, Ortog. Hort. Matr. Dec. 88 ; Ruiz & Pav. Fl. Per. ii. 11 = coccinea.
angustifolia, *Choisy, in DC. Prod.* ix. 366.—Bras.
angustifolia, *Jacq. Coll.* ii. 367.—Geront. trop.
angustifolia, *Parodi, Contrib.* 21.—Paraguay.
anisoloba, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 269.—Mexic.
aquatica, *Forsk. Fl. Aegypt. Arab.* 44.—Geront. trop.
arachnoides, *Boj. Hort. Maurit.* 225.—Afr. trop.
arachnosperma, *Welw. Apont.* 558.—Afr. trop.
arborescens, *Sweet, Hort. Brit.* ed. II. 378, nomen ; *G. Don, Gen. Syst.* iv. 267.—Mexic.
arenaria, *Roem. & Schult. Syst.* iv. 247.—Ind. occ.
arenaria, *Steud. Nom.* ed. II. i. 815.—Ins. occ.
argentea, *Mœsn. in Mart. Fl. Bras.* vii. 247.—Am. trop.
argentifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 131.—Ind. occ.
Argyreia, *Meisn. in Mart. Fl. Bras.* vii. 246.—Am. trop.
argyrocalyces, *Choisy, in DC. Prod.* ix. 357.—Afr. austr
argyrophylla, *Vatke, in Linnaea*, xliii. (1880-82) 510 —Afr. trop.
aristolochiaefolia, *G. Don, Gen. Syst.* iv. 277.—Venezuela.
aristulata, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11 (1845) 268.—Mexic.
armata, *Roem. & Schult. Syst.* iv. 214.—Mexic.
Arnottiana, Steud. Nom. ed. II. i. 815 = dissecta.
asarifolia, *Roem. & Schult. Syst.* iv. 231.—Afr. trop.
asclepiadea, Choisy, ex Steud. Nom. ed. II. i. 816 = Argyreia Guichenotii.
aspera, Vatke, in Linnaea, xliii. (1880-82) 508 = grandiflora.
aspersa, *Mart. ex Choisy, in DC. Prod.* ix. 365.—Bras.
atropurpurea, Choisy, in Mém. Soc. Phys. Genève. vi. (1833) 457 = Lettsomia atropurpurea.
atrosanguinea, Hook. Bot. Mag. t. 2170 = Argyreia cuneata.
sturensis, *G. Don, Gen. Syst.* iv. 266.—Venezuela.
auricoma, *A. Rich. Tent. Fl. Abyss.* ii. 65.—Abyss.
aurita, G. Don, Gen. Syst. iv. 282 = Convolvulus auritus.
auricularis, Ruiz. Fl. Ludov. 47 = hederacea.
Baaian, *Montr. in Mem. Acad. Lyon*, x. (1860) 238.—N. Caled.
baccifera, Kuas. ex Steud. Nom. ed. II. i. 815 = Evolvulus gangeticus, Argyreia tiliaefolia.
Bachi, *Choisy, in Mém. Soc. Phys. Genève.* viii. i. (1838) 60.—Afr. trop.
bahiensis, *Willd. ex Roem. & Schult. Syst.* iv. 189.—Am. trop.
barbata, Choisy, in Mém. Soc. Phys. Genève. vi. (1833) 458 = Lettsomia barbata.
barbata, Roth, Catalect. i. 37 = hederacea.
barbatisepala, *A. Gray, Syn. Fl. N. Am.* ii. 1. 212.—Texas.
barbigera, *Sweet, Brit. Flow. Gard.* t. t. 86.—Am. bor.
barleroides, *Benth. & Hook. f. Gen.* ii. 872.—Ind. or.
Batatas, *Poir. Encyc.* vi. 14.—Reg. trop. (cult.).
Batatilla, *G. Don, Gen. Syst.* iv. 275.—Venezuela.
batatoides, Benth. Pl. Hartw. 46 = mastilanica.
batatoides, *Choisy, in Mém. Soc. Phys. Genève.* viii. i. (1838) 58.—Bras.
Beladambœe, *Roem. & Schult. Syst.* iv. 238.—Ind. or.
bengalensis, Roth, ex Roem. & Schult. l. c. 229 = Rivea tiliaefolia.
beraviensis, *Vatke, in Linnaea*, xliii. (1880-82) 514.—Madag.
bicolor, *Lam. Tabl. Encyc.* i. 485.—Am. trop.
bicolor, Sweet, Hort. Brit. ed. I. 389 = Hewittia bicolor.
bidentata, G. Don, Gen. Syst. iv. 266 = angustifolia, Jacq.
bifida, Roem. & Schult. Syst. iv. 241 ; Roth, Nov. Pl. Sp. 115 = cymosa.
bifida, Wehmn. in Syll. Ratisb. ii. (1832) 17 = holosericea.
biflora, R. Br. Prod. 487 = stocta.
biflora, *Pers. Syn.* i. 183.—China.
bignonioides, *Sims, Bot. Mag.* t. 2645.—Guian.
biloba, *Forsk. Fl. Aegypt. Arab.* 44.—Amphig. trop.
bilobata, G. Don, in Sweet, Hort. Brit. ed. III. 483 = biloba.
bipedunculata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 216.—Reg. Himal.
Blanchetii, *Choisy, in DC. Prod.* ix. 357.—Bras.
Blancoi, *Choisy, l. c.* 389.—Ins. Philipp.
blanda, Sweet, Hort. Brit. ed. I. 388 = cymosa.
blepharocepala, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 72.—Abyss.
Blumii, Steud. Nom. ed. II. i. 815 = angustifolia, Jacq.
bogotensis, *G. Don, Gen. Syst.* iv. 273.—N. Granat.
bombycina, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 384.—Mexic.
Bona-nox, Heyne, ex Steud. Nom. ed. II. i. 815 = grandiflora.
bonariensis, *Hook. Bot. Mag.* t. 3665.—Reg. Argent.
Bouvetii, *Duchass. & Walp. in Linnaea*, xxiii. (1850) 752.—Guadeloup.
brachypoda, *Benth. Bot. Voy. Sulph.* 135.—Mexic.
bractesta, *Gav. Ic.* v. 51. t. 447.—Mexic.
bracteata, J. Grah. Cat. Pl. Bomb. 181 = Argyreia sericea.
bracteata, Heyne, ex Wall. Cat. n. 1419 = Argyreia bracteata.
bracteata, Roem. & Schult. Syst. iv. 227 ; Blume, Bijdr. 711 = Hewittia bicolor.
bractesta, Rudolph, ex Ledeb. in Schrad. Neues Journ. ii. (1807) 292 = Rudolphii.
bractesta, *Wight, Ic.* t. 1374 —Ind. or.
brasiliana, *Meisn. in Mart. Fl. Bras.* vii. 261.—Bras.
brasiliensis, G. F. W. Mey. Prim. Fl. Esseq. 97 = biloba.
brevifora, G. F. W. Mey. l. c. 100 = setifera.
brevipes, *Moç. & Sesse, ex Choisy, in DC. Prod.* ix. 349. —Mexic.
Brownei, Roem. & Schult. Syst. iv. 252 = obscura, *Ker-Gawl.*
Buaraiap, *Montr. in Mem. Acad. Lyon*, x. (1860) 238. —N. Caled.
Buchananii, Choisy, in Mém. Soc. Phys Genève. vi. (1833) 474 = salicifolia.
bufalina, *Choisy, l. c.* 452.—Cochinch.
bulbocastana, Moç. & Sesse, ex Choisy, in DC. Prod. ix. 344 = heterophylla, *Ortog.*
Burchellia, *Meisen. in Mart. Fl. Bras.* vii. 271.—Bras.
Burmanni, *Choisy, in DC. Prod.* ix. 350.—Am. austr.
Burmanni, Koen. ex Choisy, l. c. 350, 392 = Argyreia Roxburghii.
cærulea, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 296.—Porto Rico.
caerulea, Koen. ex Roxb. Fl. Ind. ed. Carey, ii. 91 = hederacea.
caerulescens, Roxb. Hort. Beng. 14 ; Fl. Ind. i. 500 = hederacea.
cæsia, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 140.—Bras.
cairica, Sweet, Hort. Brit. ed. I. 387 = palmata.
calantha, *Griseb. Cat. Pl. Cub.* 202.—Cuba.
caliginosa, Choisy, in Mém. Soc. Phys. Genève. vi. (1833) 461 = cymosa.
Calobra, *Hill & F. Muell. Fragm.* xi. 78.—Austral.
calonoura, *Meisen. in Mart. Fl. Bras.* vii. 281.—Bras.
calophylla, *Wright, ex Griseb. Cat. Pl. Cub.* 204.—Cuba.
calycina, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 201.—Ind. or. ; Afr. trop.
calycina, *Meisen. in Mart. Fl. Bras.* vii. 260.—Bras.
calystegioides, E. Mey. in Drège, Zwei Pfl. Docum. 196 = crassipes.
campanulata, Linn. ex Steud. Nom. ed. II. i. 815 = Argyreia tiliaefolia.
campanulata, *Linn. Sp. Pl.* 160.—Ind. or. ; Malaya.
campestris, *Meisen. in Mart. Fl. Bras.* vii. 254.—Bras.
cana, E. Mey. in Drège, Zwei Pfl. Docum. 195 = argyreioides.
candicans, Sweet, Hort. Brit. ed. I. 389 = fastigiata.
canescens, G. Don, Gen. Syst. iv. 278 = Jacquemontia violacea.
capillacea, G. Don, l. c. 267 = arnata.
capillata, Spaa. in Linnæa, xv. (1841) 540 = palmata ?
capillifolia, Bert. ex Choisy, in DC. Prod. ix. 388 = I. Carolina.
capitata, Choisy, l. c. 365 = guineensis.
capitata, Roem. & Schult. Syst. iv. 238 = Lettsomia strigosa.
capitellata, Choisy, in Mém. Soc. Phys. Genève. vi. (1833) 446 = I. Pes-tigridis.
capparoides, *Choisy, l. c.* viii. i. (1838) 59.—Bras.
cardiophylla, *A. Gray, Syn. Fl. N. Am.* ii. t. 213.—Texas.
cardiosepala, Hochst. ex Choisy, in DC. Prod. ix. 429, 392 = calycina, *Benth.*

IPOMOEA :—
cardiosepala, *Meisen. in Mart. Fl. Bras.* vii. 265.—Bras.
carinata, Endl. Prod. Fl. Norf. 58 = I. Bonc nox.
cariosepala, Klotzsch, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 201 = calycina, *Benth.*
carnea, Forst. f. Prod. 19 = denticulata.
carnea, *Jacq. Enum. Pl. Carib.* 18; *Select. Stirp. Am.* 26.—Am. austr.
carnosa, *R. Br. Prod.* 485.—Prop. reg.
Carolina, *Linn. Sp. Pl.* 160.—Ind. occ.
Carolina, Pursh, Fl. Am. Sept. i. 145 = commutata.
caroliniana, Poir. Encyc. vi. 11 = I. Carolina.
cataractae, *Endl. Prod. Fl. Norf.* 53.—Ins. Norfolc.
Catesbaei, G. F. W. Mey. Prim. Fl. Esseq. 113 = I. Batatas.
cathartica, *Poir. Encyc. Suppl.* iv. 633.—Am. trop.
Cavanillesii, *Roem. & Schult. Syst.* iv. 214 —Hab.?
chenopodifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 385. —Mexic.
chilensis, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1857) 1.—Chili.
Choisyi, *Montr. in Mém. Acad. Lyon*, x. (1860) 237. —N. Caled.
cholulensis, H. B. & K. Nov. Gen. et Sp. iii. 112 = coccinea.
chrysoides, [*Ker-Gawl.*] *in Bot. Reg.* t. 270.—As. et Austral. trop.
chrysorrhiza, *Hook. f. Handb. N. Zeal. Fl.* 760.—N. Zel.
chrysotricha, *Meisen. in Mart. Fl. Bras.* vii. 243 — Am. trop.
ciliolata, *Pers. Syn.* i. 183.—Am. bor.
ciliosa, Pursh, Fl. Am. Sept. i. 140 = pracc.
cincta, Koem. & Schult. Syst. iv. 254 = lracciata, *Cav.*
cinerascens, R. Br. Prod. 486 = incisa
cissoides, *Griseb. Fl. Brit. W. Ind.* 478.—Mexic.
Clappertoni, *R. Br. in Denh. & Clapp. Trav. App.* 240. —Afr. trop.
Clarkei, *Hook. f. Fl. Brit. Ind.* iv. 734.—Ind. or.
clavata, Rudolph, ex Ledeb. Dia. Pl. Dom. Dec. n. 5; ex Schrad. Neues Journ. il. (1807) 292 = triloba.
coccinea, *Linn. Sp. Pl.* 160.—Am. calid. (albi cult.).
coccinea, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 197 = bederifolia.
codonantha, *Benth. Pl. Hartw.* 120.—Ecuador.
coelestis, Boj. ex Choisy, in DC. Prod. ix. 439, 393 = Brweria spectabilis.
commutophylla, *A. Rich. Tent. Fl. Abyss.* ii. 65.— Abyssin.
commutata, *Roem. & Schult. Syst.* iv. 228.—Am. bor.
compressa, *Gust. Ind. Sem. Hort. Boccad.* (1826) 7.— Ind. or.
conferta, G. Don, Gen. Syst. iv. 267 = eriocarpa?
congesta, *R. Br. Prod.* 485.—Austral.
conica, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1854) 224.—Afr. trop.
contorquens, *Choisy, in DC. Prod.* ix. 345.—Bras.
contorta, *Choisy, l. c.* 350.—Afr. austr.
coptica, Roth, Nov. Pl. Sp. 110 = dissecta.
cordifolia, *Carey, ex Voigt, Hort. Suburb. Calc.* 353.— Ind. occ.?
cordifolia, *Tenore, Hort. Nap.* 58; *ex Choisy, in DC. Prod.* ix. 390, nomen.—Hab.?
cordigera, *Mart. ex Regel, Ind. Sem. Hort. Petrop.* (1860) 39.—Hab.?
cordofana, *Choisy, in DC. Prod.* ix. 350.—Afr. trop.
coriacea, *Choisy, l. c.* 358.—Bras.
corallinensis, *Choisy, l. c.* 361.—Mexic.
corymbosa, Roth, Nov. Pl. Sp. 109 = sidaefolia.
coscinosperma, *Hochst. ex Choisy, in DC. Prod.* ix. 354.—Afr. trop.
costata, *F. Muell. ex Benth. Fl. Austral.* iv. 419.— Austral.
costellata, *Torr. Bot. Mex. Bound.* 149.—N. Mexic.
crassifolia, Cav. Desc. 100 = triloba.
crassipes, *Hook. Bot. Mag.* t. 4068.—Afr. austr.
crotonifolia, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 160.—Bras.
Cruckshanksii, *Choisy, in DC. Prod.* ix. 339.—Ecuador.
cumanensis, G. Don, Gen. Syst. iv. 278 = ferruginea.
cuneifolia, *A. Gray, in Proc. Am. Acad.* xix. (1883) 90.—Arizona.
cuneifolia, *Meisen. in Mart. Fl. Bras.* vii. 245.— Bras.
curassavica, Balb. in Roem. & Schult. Syst. iv. 237 = sibirica?
cuspidata, *D. Don, Prod. Fl. Nep.* 98.—Reg. Himal.

IPOMOEA :—
cuspidata, Kuiz & Pav. Fl. Per. ii. 11 = bicolor.
cyamecilis, St. Leg. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = I. Quamoclit.
cyanantha, *Griseb. Fl. Brit. W. Ind.* 469.—Ind. occ.
Cymbalaria, Fenzl, in Flora, xxvii. (1844) 812 = reniformis.
cymosa, Baker, Fl. Maurit. 208 = denticulata.
cymosa, Lindl. Bot. Reg. (1848) t. 24 = sidaefolia.
cymosa, G. F. W. Mey. Prim. Fl. Esseq. 99 = fastigiata.
cymosa, *Roem. & Schult. Syst.* iv. 241.—As. et Austral. trop.
cyanchifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 208.—Reg. Himal.
cynanchifolia, *Meisen. in Mart. Fl. Bras.* vii. 274.— Bras.
dactylophylla, *Griseb. Cat. Pl. Cub.* 203.—Cuba.
dasysperma, *Jacq. Eclog.* i. 132 t. 89.—Ind. or.
daturaefiora, *Meissn. in Mart. Fl. Bras.* vii. 273.— Bras.
Davenporti, *F. Muell. Fragm.* vi. 97.—Austral.
dealbata, *Hemsl. Biol. Centr. Am. Bot.* ii. 386.—Mexic.
debilis, Boj. ex Choisy, in DC. Prod ix. 353 = desmophylla.
declinata, Mart. ex Meisen. in Mart. Fl. Bras. vii. 252 = polymorpha.
demeraranna, *Choisy, in DC. Prod.* ix. 361.—Guian.
dendroides, *Choisy, in Mem. Soc. Phys. Genèv.* viii. 1. (1838) 55.—Afr. trop.
dentata, Willd. ex Roem. & Schult. Syst. iv. 789 = chrysotricha.
denticulata, R. Br. Prod. 485 = angustifolia, *Jacq.*
denticulata, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 447.—As. et Austral. trop.
Deppeana, *G. Don, Gen. Syst.* iv. 276.—Mexic.
desmophylla, *Boj. ex Choisy, in DC. Prod.* ix. 353.— Madag.
Desrousseauxii, *Steud. Nom.* ed. II. i. 816.—Am. austr.
diantha, Roem. & Schult. Syst. iv. 254 = erecta.
dichotoma, *Choisy, in DC. Prod.* ix. 383.—Bras.
dichroantha, H. B. & K. Nov. Gen. et Sp. iii. 112 = coccinea.
dichroa, Choisy, in DC. Prod. ix. 364 = pilosa.
digitata, *Linn. Syst.* ed. X. 924.—Amphig. trop.
digitata, *Swartz. Contrib.* 28.—Paraguay
digitifolia, Sweet, Hort. Brit. ed. II. 372 = palmata.
Dillenii, Roem. & Schult. Syst. iv. 237 = hederacea.
dissecta, *G. Don, Gen. Syst.* iv. 270.—Bras.
discolor, Hort. ex Sims, Bot. Mag. t. 2645 = bignonioides.
discolor, Jacq. Hort. Schoenb. iii. 6 = purpurea.
dissecta, Purh, Fl. Am. Sept. i. 145 = sinuata.
dissecta, *Willd. Phytogr.* 5. t. 2.—Geront. trop.
distans, *Choisy, in DC. Prod.* ix. 378.—Am. trop.
diversifolia, Am. ex Steud. Nom. ed. II. i. 816 = dissecta.
diversifolia, *R. Br. Prod.* 487.—Austral.
diversifolia, F. Didr. in Kjoeb. Vidensk. Meddel. (1854) 223.—Afr. trop.
dubia, *Hemsl. Biol. Centr. Am. Bot.* ii. 386.—Mexic.
dubia, Roem. & Schult. Syst. iv. 216 = coccinea.
dumetorum, *Willd. ex Roem. & Schult. l. c.* 789.—N. Granat.
ebracteata, *Choisy, in DC. Prod.* ix. 377.—Am. trop.
echinocalyx, *Meisen. in Mart. Fl. Bras.* vii. 223.— Bras.
echioides, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 54.—Bras.
Edwardsii, *Regel, Ind. Sem. Hort. Petrop.* (1856) 28. —Ins. Cap. Virid.
elegans, *A. Distr. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 313.—Bras.
elegans, *Meisen. in Mart. Fl. Bras.* vii. 243.—Bras.
elliptica, Roth, Nov. Pl. Sp. 113 = Lettsomia elliptica.
elongata, *Choisy, in DC. Prod.* ix. 356.—Mexic.
emetica, *Choisy, l. c.* 376.—Mexic.
ennealoba, *Benoo. Pl. Ou.* ii. 68. t. 101.—Afr. trop.
erecta, *R. Br. Prod.* 487.—Austral.
ericoides, *Meisen. in Mart. Fl. Bras.* vii. 251.—Bras.
ericoides, *Meisen. l. c.* 286.—Bras.
eriocarpa, *R. Br. Prod.* 484.—Geront. trop.

IPOMOEA :—
eriocephala, Moric. Pl. Nouv. Am. 43. t. 29 = Jacquemontia eriocephala.
eriosperma, Beauv. Fl. Ow. ii. 73 = digitata.
eriosperma, *Bert. ex Spreng. Syst. i.* 598; *et Choisy, in DC. Prod.* ix. 388.—Guadalup.
eriosperma, Rafin. Fl. Tellur. iv. 74 = Desrousseauxii?
erythraea, Moç. & Sesse, ex Choisy, in DC. Prod. ix. 335, 398 = cocciues.
eustachiana, Jacq. Obs. 2. t. 36 = triloba.
evolvuloides, Moric. Pl. Nouv. Am. 47. t. 32 = Jacquemontia evolvuloides.
falklandica, *Griseb. Cat. Pl. Cub.* 206.—Cuba.
fastigiata, Chapm. Fl. S. U. St. 438 = cathartica.
fastigiata, Choisy, in DC. Prod. ix. 340 = Argyreia populifolia.
fastigiata, *Sweet, Hort. Brit.* ed. I. 283; ed. II. 372.— Am. trop.
ferruginea, *Roem. & Schult. Syst.* iv. 240.—Am. austr.
ficifolia, *Lindl. in Bot. Reg.* (1840) *Misc.* 90.—Reg. Argent.
filicaulis, Blume, Bijdr. 721 = angustifolia, *Jacq.*
filifolia, G. Don, Gen. Syst. iv. 268 = Cruckshanksii.
filiformis, *Jacq. Enum. Pl. Carib.* 13.—Ind. occ.
filiformis, Voigt, Hort. Suburb. Calc. 356 = angustifolia.
Preuss. Stat. (1879) t. 2.—Afr. trop.
delphinifolia, *Mart. & Gal. in Bull. Acad. Brux.* xii. ii. (1845) 365.—Mexic.
delphinioides, Choisy, in Mém. Soc. Phys. Genèv. viii. 1. (1838) 58 = polymorpha.
demeraranna, *Choisy, in DC. Prod.* ix. 361.—Guian.
dendroides, *Choisy, in Mem. Soc. Phys. Genèv.* viii. 1. (1838) 55.—Afr. trop.
denutata, Willd. ex Roem. & Schult. Syst. iv. 789 = chrysotricha.
denticulata, R. Br. Prod. 485 = angustifolia, *Jacq.*
denticulata, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 447.—As. et Austral. trop.
Deppeana, *G. Don, Gen. Syst.* iv. 276.—Mexic.
desmophylla, *Boj. ex Choisy, in DC. Prod.* ix. 353.— Madag.
Desrousseauxii, *Steud. Nom.* ed. II. i. 816.—Am. austr.
diantha, Roem. & Schult. Syst. iv. 254 = erecta.
dichotoma, *Choisy, in DC. Prod.* ix. 383.—Bras.
dichroantha, H. B. & K. Nov. Gen. et Sp. iii. 112 = coccinea.
dichroa, Choisy, in DC. Prod. ix. 364 = pilosa.
digitata, *Linn. Syst.* ed. X. 924.—Amphig. trop.
digitata, *Swartz. Contrib.* 28.—Paraguay
digitifolia, Sweet, Hort. Brit. ed. II. 372 = palmata.
Dillenii, Roem. & Schult. Syst. iv. 237 = hederacea.
dissecta, *G. Don, Gen. Syst.* iv. 270.—Bras.
discolor, Hort. ex Sims, Bot. Mag. t. 2645 = bignonioides.
discolor, Jacq. Hort. Schoenb. iii. 6 = purpurea.
dissecta, Purh, Fl. Am. Sept. i. 145 = sinuata.
dissecta, *Willd. Phytogr.* 5. t. 2.—Geront. trop.
distans, *Choisy, in DC. Prod.* ix. 378.—Am. trop.
diversifolia, Am. ex Steud. Nom. ed. II. i. 816 = dissecta.
diversifolia, *R. Br. Prod.* 487.—Austral.
diversifolia, F. Didr. in Kjoeb. Vidensk. Meddel. (1854) 223.—Afr. trop.
dubia, *Hemsl. Biol. Centr. Am. Bot.* ii. 386.—Mexic.
dubia, Roem. & Schult. Syst. iv. 216 = cocciues.
dumetorum, *Willd. ex Roem. & Schult. l. c.* 789.—N. Granat.
ebracteata, *Choisy, in DC. Prod.* ix. 377.—Am. trop.
echinocalyx, *Meisen. in Mart. Fl. Bras.* vii. 223.— Bras.
echioides, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 54.—Bras.
Edwardsii, *Regel, Ind. Sem. Hort. Petrop.* (1856) 28. —Ins. Cap. Virid.
elegans, *A. Distr. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 313.—Bras.
elegans, *Meisen. in Mart. Fl. Bras.* vii. 243.—Bras.
elliptica, Roth, Nov. Pl. Sp. 113 = Lettsomia elliptica.
elongata, *Choisy, in DC. Prod.* ix. 356.—Mexic.
emetica, *Choisy, l. c.* 376.—Mexic.
ennealoba, *Benoo. Pl. Ou.* ii. 68. t. 101.—Afr. trop.
erecta, *R. Br. Prod.* 487.—Austral.
ericoides, *Meisen. in Mart. Fl. Bras.* vii. 251.—Bras.
ericoides, *Meisen. l. c.* 286.—Bras.
eriocarpa, *R. Br. Prod.* 484.—Geront. trop.

IPOMOEA :—

glaucescens, G. Don, Gen. Syst. iv. 275 = dumetorum.
glaucifolia, *Linn. Sp. Pl.* 161.—Am. trop.
Glenieii, *Thw. ex C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* iv. 208.—Zeylan.
globosa, Meisn. in Mart. Fl. Bras. vii. 219 = pedunculatis.
gnaphalosperma, Hochst. ex Choisy, in DC. Prod. ix. 361 = rumicifolia.
Gomezii, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* iv. 211.—Burma.
gossypifolia, Stend. Nom. ed. II. i. 816 = seq.
gossypiifolia, Willd. Enum. Hort. Berol. 208 = digitata.
gossypioides, *Parodi, Contrib.* 15.—Paraguay.
goyazensis, *Gardn. in Hook. Ic. Pl.* t. 479.—Bras.
gracilis, *R. Br. Prod.* 486.—Austral.
graminea, *R. Br. l. c.* 485.—Austral.
graminiformis, *Meisn. in Mart. Fl. Bras.* vii. 250.—Bras.
graminicoides, Meisn. l. c. 417, Index = praec.
grandiflora, Jacq. Hort. Vind iii. t. 69 = latiflora.
grandiflora, *Lam. Tabl. Encyc* i. 467, *Roem. & Schult. Syst.* iv. 240.—Geront. trop.
grandiflora, *Parodi, Contrib.* 21.—Paraguay.
grandiflora, Roxb. Hort. Beng. 14; Fl. Ind. i. 497 = I. Bona-nox.
Grantii, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 116. —Afr. trop
granulata, *Parodi, Contrib.* 17.—Paraguay.
guineensis, *G. Don, Gen. Syst.* iv. 269.—Afr. trop.
guyanensis, *Choisy*, in *DC. Prod.* ix. 366.—Guian.
Habeliana, *Oliver, in Hook. Ic. Pl.* t. 1099.—Ins. Galapag.
Hanekeana, *Choisy, in DC. Prod.* ix. 358.—Am. trop
halophila, Miq. in Linnæa, xviii. (1844) 598 = carnosa.
halophyla, Poepp. ex Steud. Nom. ed. II. i. 817 = hilobæ.
Hamiltonii, G. Don, Gen. Syst. iv. 268 = triquetra.
Hamiltonii, Steud. Nom. ed. II. i. 817 = cuspidata.
Hardingii ×, Part. Mag. Bot. xi. (1844) 217.
Hartmanni, *Vatke & Kuntze*, in *Linnæa*, xliii. (1880–82) 514.—Afr. trop.
Hartwegi, *Benth. Pl. Hartw.* 15.—Mexic.
hartwegii, Meisn. in Mart. Fl. Bras. vii. 230 = pedunculatis.
hastata, *Linn. Mant.* ii. 204.—Java.
hastigera, H. B. & K. Nov. Gen. et Sp. iii. 111 = hederifolia.
havanensis, Choisy, in DC. Prod. ix. 368 = Convolvulus havanensis.
Haylocki, *Regel, Cat. Pl. Hort. Asiahon.* 77, nomen.—Hab.?
hederacea, *Jacq. Coll.* i. 124.—Amphig. trop.
hederifolia, *Linn. Syst. ed.* X. 925.—Ind. occ.
hepaticæfolia, Linn. Sp. Pl. 161 = I. Pes-tigridis.
hepaticifolia, Blanco, Fl. Filip. ed. II. 72 = angustifolia.
heptaphylla, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 527.—Ind. occ.
heptaphylla, Voigt, Hort. Suburb. Calc. 360 = palmata.
Hermanniæ, G. Don, Gen. Syst. iv. 276 = Convolvulus Hermanniæ.
heterophylla, *R. Br. Prod.* 487.—Austral.
heterophylla, G. Don, Gen. Syst. iv. 274 = sagittata.
heterophylla, *Orteg. Hort. Matr. Dec.* 9; *Willd. Enum. Hort. Berol.* 207.—Mexic.; Cuba.
heterophylla, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 81.—Bras.
heterophylla, Torr. Bot. Mex. Bound. 149 = Lindheimeri.
heterotricha, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1854) 290.—Afr. trop.
heterotricha, *Meisn. in Mart. Fl. Bras.* vii. 230.—Bras.
Heynei, Roem. & Schult. Syst. iv. 287 = cymosa.
Heynei, Wall. Cat. n. 1887 = sepiaria.
hirsuta, R. Br. Prod. 486 = quinata.
hirsuta, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 80 = Martiniana.
hirsutissima, *Gardn. in Hook. Ic. Pl.* sub t. 471.—Bras.
hirsutula, Hort. Vindob. ex Hornem. Hort. Hafn. Suppl. 28 = hederacea.
hirsutula, *Jacq. f. Eclog. Pl.* 65. t. 44.—Am. austr.
hirta, *Mart. & Gal. in Bull. Acad. Brux.* xii. ii. (1846) 394.—Mexic.
hirtiflora, *Mart. & Gal. l. c.* 267.—Mexic.

IPOMOEA:—

Hispaniolæ, *G. Don, Gen. Syst.* iv. 280.—Ins S. Doming.
hispida, *Parodi, Contrib.* 18.—Paraguay.
hispida, Roem. & Schult. Syst. iv. 238 = eriocarpa.
hispida, Zuccagni, in Roem. Collect. 127 = purpurea.
hispidula, *Ledeb. Ind. Sem Hort. Dorpat.* (1822) 11 (Quid?).—Hab.?
holosericea, *E. Mey. in Drège, Zwei Pfl. Docum.* 195, nomen.—Afr. austr.
holosericea, *Weinm. in Syll. Ratisb.* ii. (1828) 17.—Bras.
Hookeri, G. Don, Gen. Syst. iv. 274 = rubro-cærulea.
Hornei, *Baker, Fl. Maurit.* 207.—Ins. Seychell.
Horsfalliae, *Hook. Bot Mag* t. 3315.—Ind. or.
Horsfalliae, Hort.—Cf. Gard. Chron. (1885) i. 566 = ternata.
Horsfeldiana, Miq. Fl. Ind Bat ii. 611 = eriocarpa.
Hostmanni, *Meisn. in Mart. Fl. Bras.* vii. 230.—Am. trop.
Humboldtiana, Roem. & Schult. Syst. iv. 789 = hederifolia.
humilis, *G. Don, Gen. Syst.* iv. 267.—Afr. trop.
humilis, *Raffin. New Fl. Am.* iv. 57 (Quid?).—Am. bor.
hypargyræa, *Griseb. Cat. Pl. Cub.* 204.—Ind. occ.
igatimiana, *Parodi, Contrib.* 17.—Paraguay.
imbricata, *Parodi, l. c.* 16.—Paraguay.
imbricata, Roth, Nov. Pl. Sp. 112 = Lettsomia aggregata.
incanata, *Choisy, in DC. Prod.* ix. 360.—Venezuela.
incisa, *R. Br. Prod.* 486.—Austral.
incurva, G. Don, Gen. Syst. iv. 266 = Clapperton.
insignis, Andr. Bot. Rep. t. 686 = digitata
insignis, *Ker-Gawl. Bot. Reg.* 75; *Sims, Bot. Mag.* t. 1790.—Reg. trop.
insuavis, Blume, Bijdr. 716 = obscura.
insularis, Steud. Nom. ed. II. i. 817 = congesta.
intermedia, Schult. Obs. Bot. 37 = purpurea
involucrata, Beauv. Fl. Owar. ii. 52 t. 89 = pileata.
involucrata, F. G. Dietr. ex Choisy, in DC. Prod ix. 394, 344 = villosa.
involucrata, Hance, in Journ. Linn. Soc. xiii. (1872) 113 = pileata.
Jacquini, *Regel, Ind. Sem. Hort. Petrop.* (1856) 27.—Hab.?
Jalapa, *Cosz, in Journ. Am. Med. Sc.* v. 1829 (1830): *Hayne, Arzneigew.* xiii t. 37.—Mexic.
Jalapa, Pursh, Fl. Am. Sept. i. 146 = macrorrhiza, *Michx.*
Jalapa, Schiede & Deppe, ex G. Don, Gen. Syst. iv. 271 = I. Purga.
jalapoida, *Griseb. Cat. Pl. Cub.* 202.—Cuba.
jamaicensis, *G. Don, Gen. Syst.* iv. 278.—Mexic.
Jamesoni, *Choisy, in DC. Prod.* ix. 367.—Ecuador.
japonica, Roem. & Schult. Syst. iv. 245 = angustifolia, *Jacq.*!
jucunda, Thw. Enum. Pl. Zeyl. 211 = grandiflora, *Lam.*
juncea, Choisy, in DC. Prod. ix. 355 = sturensis.
Kadsuru, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 475.—Japon.
Karwinskianum, *Regel, Ind. Sem. Hort. Petrop.* (1857) 46.—Mexic.
katgurhiensis, *Hoffmgg. Vers. Pfl. Nachtr.* ii. 141.—Ind. or.
kentrocarpa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 70.—Abyss.
kentrocalyx, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* iv. 213.—Geront. trop.
Kinbergi, *Anderss. in Vet. Akad. Handl. Stockh.* 1858 (1855) 212.—Ins. Galapag.
kituiensis, *Vatke*, in *Linnæa*, xliii. (1880–82) 611.—Afr. trop.
Kleiniana, Roem. & Schult. Syst. iv. 789 = Argyreia bracteata.
Kotschyana, *Hochst. ex Choisy, in DC. Prod.* ix. 354. —Abyss.
Krusensternii, Ledeb. in Mém. Acad. Pétersb. iv. (1811) 401 = I. Bona-nox.
Kunthiana, *Meisn. in Mart. Fl. Bras.* vii. 253.—Am. trop.
lachnæa, *Spreng. Neue Entdech.* iii. 29.—Ins. S. Doming.
lachnosperma, *Choisy, in DC. Prod.* ix. 356.—Afr. trop.
lachnita, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 89.—Ins. Socotra.

IPOMOEA:—

laciniata, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* iv 200.—Ind. or.
lactea, *Wall. ex Voigt, Hort. Suburb. Calc.* 361 — Ind. or.
lactescens, *Benth. Pl. Hartw.* 120.—Ecuador.
lacunosa, *Linn. Sp. Pl.* 161.—Am. bor
laevigata, Soland ex Benth. Fl Austral. iv. 422 = denticulata.
laevigata, *Steud Nom.* ed II. i. 817 — Ins. Societ., Austral.
lancelata, G. Don, Gen Syst. iv. 282 = uniflora
Langsdorffii, *Choisy, in DC. Prod* ix 368 —Am. trop
lapidosa, *Vatke*, in *Linnæa*, xliii. (1880–82) 507.—Afr trop
laniociada, Choisy, in DC Prod ix. 357 = Jacquemontia Choisyana
latiflora, Landl. Bot. Reg. t. 889 = I. Bona-nox.
latiflora, *Roem. & Schult. Syst.* iv. 240 —Ind. or et occ.
latifolia, *Mart & Gal. in Bull. Acad. Brux.* xii. ii (1845) 266.—Mexic
laurifolia, Sweet, Hort. Brit. ed. I 288 = Lettsomia elliptica.
Leari, *Paxt. Mag. Bot.* vi (1839) 267 —Am. trop.
Ledebourii, *Choisy, in DC. Prod.* ix. 388.—Am. bor
Lemmoni, *A. Gray*, in *Proc Am. Acad.* xix. (1883) 90 —Arizona.
leptophylla, *Torr. in Frem. 1st Rep.* 94.—Am bor. occ.
leptotoma, *Torr. Bot. Mex. Bound.* 150.—Arizona.
leucantha, *Jacq. Coll.* ii. 280.—Am. austr.
ligulata, Boj. Hort. Maurit. 229 = eriocarpa.
lilacina, Blume, Bijdr 716 = Argyreia tiliæfolia?
lilacina, Kunze, ex Choisy, in DC. Prod. ix. 390 = Schlechtendahlii.
lilacina, *Schlecht. in Linnæa*, xx. (1847) 31.—Mexic.
lilacina, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii (1822) 81 = Bras.
Lindeni, *Mart. & Gal. in Bull. Acad. Brux* xii. ii. (1846) 264.—Mexic.
Lindheimeri, *A. Gray, Syn. Fl. N. Am.* ii. i. 210.—Mexic.
Lindleyi, Choisy, in DC. Prod. ix. 371 = fragrans.
linearifolia, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 204.—Ins. Cap Virid
linifolia, *Blume, Bijdr.* 721.—Malaya; Austral.
linoides, Choisy, in DC. Prod. ix. 354 = Jacquemontia linoides.
littoralis, *Blume, Bijdr.* 713.—Java.
littoralis, Boiss. Fl. Orient. iv. 113 = carnosa.
littoralis, Thw. Enum. Pl. Zeyl. 211 = denticulata.
livescens, *Meisn. in Mart. Fl. Bras.* vii. 224.—Am trop.
Llaveana, Melsm. l. c. 219 = I. Funis.
longepeduncularis, *Hemsl. Biol. Centr. Am. Bot.* ii 389.—Mexic.
longeranosa, *Choisy, in DC. Prod.* ix. 384.—Am. trop.
longicuspis, Meisn. in Mart. Fl. Bras. vii. 227 = bicolor.
longiflora, R. Br. Prod. i. 484 = grandiflora.
longiflora, Humb. & Bompl. ex Willd. Enum. Hort. Berol. 207 = I. Bona-nox.
longifolia, Benth. Pl. Hartw. 16 = carnosa.
longipes, *Garcke*, in *Linnæa*, xxii. (1849) 66.—Guian.
Landii, *Choisy*, in *Mém. Soc. Phys. Genèv.* viii. t. (1838) 56.—Bras.
lutea, *Hemsl. Diagn. Pl. Nov.* 34.—Guatemala.
lutoia, R. Br. Prod. 485 = obscura, *Ker-Gawl.*
lutoia, Jacq. Coll. ii. 286 = coccinea.
luxurians, *Moric. Pl. Nouv. Am.* 58. t. 39.—Bras.
macrantha, G. Don, Gen. Syst. iv. 267 = mansooides.
macrantha, Roem. & Schult. Syst. iv. 251 = grandiflora, *Lam.*
macrocalyx, Choisy, in DC. Prod. ix. 362 = glabra.
macrophylla, Choisy, l. c. 374.—Am. trop.
macropoda, Boj. Hort. Maurit. 229 = capitata.
macrorrhiza, *Michx. Fl. Bor. Am.* i. 141.—Am. bor.
macrorrhizos, *Roem. & Schult. Syst.* iv. 211.—Ins. S. Doming.
madagascariensis, *Choisy, in DC. Prod.* ix. 367.—Madag.
Mairei, *Choisy, l. c.* 374.—Mexic.
malabarica, Roem. & Schult. Syst. iv. 235 = Argyreia malabarica.

IPOMOEA :—
malvacoides, *Meisn. in Mart. Fl. Bras.* vii. 251.—Am. trop.
malvavacoides, *Meisn. l. c.* 384.—Am. trop.
mammosa, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 475.—Cochinch. ; Polynes. †
maragtnensis, Choisy, in DC. Prod. ix. 351 = contorquens.
Marcellii, *Meisn. in Mart. Fl. Bras.* vii. 257.—Am. trop.
marianensis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 468.—Ins. Marian.
maritima, R. Br. Prod. 486 = biloba.
Martii, *Meisn. in Mart. Fl. Bras.* vii. 235.—Am. trop.
martianicensis, *G. F. W. Mey. Prim. Fl. Esseq.* 98.—Amphig. trop.
Martiusana, *Steud. Nom.* ed. II. i. 817.—Bras.
matamoranga, G. Don, Gen. Syst. iv. 289 = Convolvulus matamoranga.
Matthioli, *Tausch, ex Steud. Nom.* ed. II. i. 817, nomen.—Hab. ?
mauranoidioides, *Meisn. in Mart. Fl. Bras.* vii. 275.—Am. trop.
mauritiana, Jacq. Coll. iv. 216 = digitata.
maxima, G. Don, in Sweet, Hort. Brit. ed. II. 372 = sepiaria.
Maximiliani, *Meisn. in Mart. Fl. Bras.* vii. 285.—Am. trop.
Mechoacan, Nutt. in Am. Journ. Sc. (1822) 289 = l. Jalapa.
megapotamica, *Choisy, in DC. Prod.* ix. 375.—Bras.
melanea, *Lindl. in Journ. Hort. Soc.* ii. (1847) 310.—Hab. ?
melanosticta, G. Don, Gen. Syst. iv. 271 = Argyreia tiliaefolia.
Mendesii, *Welw. Apont.* 584.—Afr. trop.
mestitlanica, *Choisy, in DC. Prod.* ix. 389.—Mexic.
mexicana, *A. Gray, Syn. Fl. N. Am.* ii. i. 210.—*N.* Mexic.
mexicana, Moç. & Sesse, ex Choisy, in DC. Prod. ix. 394, 835 = coccinea.
Meyeri, G. Don, Gen. Syst. iv. 275 = Convolvulus Meyeri.
micans, *Garcke, in Linnaea*, xxii. (1849) 56.—Bras.
Michauxii, Sweet, Hort. Brit. ed. I. 288 = I. Jalapa.
microdactyla, *Griseb. Cat. Pl. Cub.* 204.—Cuba.
microphylla, Roth, Nov. Pl. Sp. 113 = Convolvulus pluricaulis.
microsepala, *Benth. Bot. Voy. Sulph.* 136.—Mexic.
modesta, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 129, 131.—Malaya.
modesta, F. Muell. Fragm. ii. 22 = Porana sericea.
mojosensis, *Vatke, in Linnaea*, xliii. (1880–82) 515.—Afr. trop.
mollicoma, Miq. Stirp. Surin. Sel. 132 = umbellata.
mollis, *G. Don, Gen. Syst.* iv. 275.—Am. trop.
montesanana, *Vatke, in Linnaea*, xliii. (1880–82) 515.—Afr. trop.
monosperma, *Spreng. ex Choisy, in DC. Prod.* ix. 382.—Venezuela.
montana, Moric. Pl. Nouv. Am. 44. t. 30 = Jacquemontia montana.
montevidensis, G. Don, Gen. Syst. iv. 276 = Convolvulus montevidensis.
monticola, Mart. ex Meisn. in Mart. Fl. Bras. vii. 240 = patula.
Morelli, *Duchass. & Walp. in Linnaea*, xxiii. (1850) 752.—Panama.
Mueller, *Benth. Fl. Austral.* iv. 423.—Austral.
multiflora, Mill. ex Choisy, in DC. Prod. ix. 377 = umbellata.
multiflora, *Roem. & Schult. Syst.* iv. 234.—Jamaic.
multiflora, *Roxb. Hort. Beng.* 84 ; *Fl. Ind.* i. 499.—Ind. or.
multicaula, *Welw. Apont.* 589.—Afr. trop.
muricata, Cav. Ic. v. 52. t. 478 = armata.
muricata, *Jacq. Hort. Schoenb.* iii. 40. t. 293.—Reg. Himal. ; Japon.
muricata, Roxb. Hort. Beng. 14 ; Fl. Ind. i. 499 = grandiflora, *Lam.*
muricoides, *Roem. & Schult. Syst.* iv. 248.—Mexic.
mutabilis, [Ker-Gawl.] Bot. Reg. t. 20 = acuminata.
Nationis, *Nichols. Dict. Gard.* ii. 191.—Peruv.
nemorosa, G. Don, Gen. Syst. iv. 283 = pentaphylla.
nephrophylla, *Meisn. in Mart. Fl. Bras.* vii. 219.—Am. trop.
neriifolia, *Gardn. in Hook. Ic. Pl.* t. 471.—Bras.
novoinensis, G. Don, Gen. Syst. iv. 282 = Jacquemontia blanda.

IPOMOEA :—
Newmannii, *Lodd. Cat.* (1826) 19.—Hab. ?
nicoharica, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 141 = denticulata.
nigricans, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 180 = sinuata.
Nil, Meisn. in Fl. Bras. vii. 228. t. 79. f. 1 = mexicana, *A. Gray*.
Nil, Roth, Catalect. fasc. i. 36 = hederacea.
nitida, *Griseb. ex Lorentz, Veget. Nord.* 84, nomen ; *et in Goett. Abh.* xxiv. (1879) 264.—Reg. Argent.
noctiflora, Griff. Notul. iv. 386 = I. Bona-nox.
noctiluca, Herb. in Bot. Reg. t. 917, in nota = I. Bona-nox.
nutans, *Choisy, in DC. Prod.* ix. 368.—Mexic.
nyctaginea, *Choisy, l. c.* 369.—Am. trop.
Nyctelea, Linn. Sp. Pl. 160 = Ellisia Nyctelea.
Nyctylea, Hill, Veg. Syst. vii. 56 = præc.
nymphaeaefolia, Blume, Bijdr. 719 = peltata.
nymphaeaefolia, *Griseb. Cat. Pl. Cub.* 203.—Cuba.
oblonga, *Benth. Bot. Voy. Sulph.* 133.—Mexic.
oblongata, *E. Mey. ex Choisy, in DC. Prod.* ix. 368.—Afr. austr.
obscura, *Hassk. Pl. Jav. Rar.* 519.—Java.
obscura, [*Ker-Gawl.*] *Bot. Reg.* t. 389.—Ind. or. ; Austral.
obtusata, *Griseb. Cat. Pl. Cub.* 202.—Cuba.
obtusiloba, *Meisn. in Mart. Fl. Bras.* vii. 233.—Am. trop.
ochraces, *G. Don, Gen. Syst.* iv. 270.—Afr. trop.
ochroleuca, Span. in Linnaea, xv. (1841) 340 = obscura.
ocularis, Baril. l. c. xiv. (1840) 126 = obscura.
oenotheroides, Kañn. Fl. Tellur. iv. 74 = argyreioides ?
oenotheroides, *Regel, Cat. Pl. Hort. Askabov.* 77, nomen.—Hab. ?
officinalis, Polt. ex Choisy, in DC. Prod. ix. 342 = cathartica.
olanda, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 211, err. for *blanda*.
olefacea, *Welw. Apont.* 589.—Afr. trop.
oliganthe, Choisy, in DC. Prod. ix. 350 = fastigiata.
oligocephala, *Benth. Bot. Voy. Sulph.* 136.—Ecuador.
operculata, *Mart. Nov. Gen. Bras.* i. 147.—Bras.
orbicularis, Ell. Sketch, 257 = biloba.
orinabensis, *Ledenois. ex Steud. Nom.* ed. II. i. 818.—Hab. ?
Ortegae, Poir. Encyc. Suppl. iv. 688 = heterophylla, *Ortg.*
oxyensis, Roth, Nov. Pl. Sp. 117 = Lettsomia aggregata.
Ottoniana, Choisy, in DC. Prod. ix. 378 = Convolvulus Ottonis.
ovalifolia, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 449 = Jacquemontia sandwicensis, et Breweriae sp. (= *Bonamia Menziesii*).
ovata, *E. Mey. in Drège, Zwei Pfl. Docum.* 195.—Afr. austr.
owariensis, *Beauv. Fl. Owar.* ii. 40.—Afr. trop.
palmata, *Forsk. Fl. Aegypt. Arab.* 43.—Amphig. trop. ; Aegypt.
palmatisecta, *Baj. ex Choisy, in DC. Prod.* ix. 352.—Afr. trop.
palmato-partita, Benth. & Hook. f. Gen. ii. 874, sphalm. = sep.
palmato-pinnata, *Benth. & Hook. f. l. c.*—Bras.
pandurae folia, E. Mey. in Drège, Zwei Pfl. Docum. 195 = Hewittia bicolor.
pandurae formis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 476.—Cochinch.
paradoxa, G. F. W. Mey. Prim. Fl. Esseq. 100 = fastigiata.
paniculata, R. Br. Prod. 486 = digitata.
paniculata, Barm. f. Fl. Ind. 50 = Convolvulus malvaceus.
pannosa, R. Br. Prod. 487 = erecta.
Papirio, Ruiz & Pav. Fl. Per. ii. 11 = pubescens.
papposa, *Phil. Fl. Atac.* 95.—Chili.
parasitica, *G. Don, Gen. Syst.* iv. 275.—Venezuela.
pareirae folia, *G. Don, l. c.* 273.—N. Granat.
Parkeri, *Choisy, in DC. Prod.* ix. 351.—Guian.
parviflora, Pers. Syn. i. 138 = Convolvulus parviflorus.
parviflora, *Vahl, Symb. Bot.* iii. 84.—Ind. occ.
patula, *Choisy, in DC. Prod.* ix. 368.—Bras.
patuliflora, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 366.—Mexic.
Pavonii, *Choisy, in DC. Prod.* ix. 390.—Ecuador.
Peckoltii, *Meisn. in Mart. Fl. Bras.* vii. 266.—Am. trop.

IPOMOEA :—
pedata, G. Don, Gen. Syst. iv. 281 = Pavonii.
pedata, *Pait. ex Choisy, in DC. Prod.* ix. 894, 847.—Am. austr.
pedata, Steud. & Hochst. ex Choisy, l. c. 384 = pailophylla.
pedata, Volgt, Hort. Suburb. Calc. 260 = dasysperma.
pedatiscta, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 365.—Mexic.
pedicellaris, *Benth. Bot. Voy. Sulph.* 135.—Am. centr.
peduncularis, *Bertol. Fl. Guatim.* 8. t. 2.—Mexic.
peltata, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 452 (*Conv. Or.* 70).—Ins. Malay. et Pacif.
peltata, Heynе, ex Wall. Cat. nab. n. 1276 = pilleata.
pendula, R. Br. Prod. 486 = palmata.
pentadactylis, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 471 = quinata.
pentagona, Sweet, Hort. Brit. ed. II. 372 = cymosa.
pentaloba, Roem. & Schult. Syst. iv. Index, 828 = digitata.
pentaphylla, Cav. Ic. iii. 29 = Cavanillesii.
pentaphylla, *Jacq. Coll.* ii. 297 ; *Ic. Pl. Rar.* ii. 10. t. 319.—Amphig. trop.
perfoliata, G. Don, Gen. Syst. iv. 269 = involucrata.
Perrottetii, *Choisy, in DC. Prod.* ix. 378.—Afr. trop.
Perryana, *Duchass. & Walp. in Linnaea*, xxiii. (1850) 751.—Panama.
Pes-caprae, Roth, Nov. Pl. Sp. 109 = biloba.
Pes-tigridis, *Linn. Sp. Pl.* 162.—Geront trop.
petaloides, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 451.—Ind. or. ; Ins. Timor.
Petersiana, *Klotzsch, in Peters, Reise Mossamb. Bot.* 259.—Afr. trop.
petiolaris, G. Don, Gen. Syst. iv. 275 = I. Bona-nox.
philadelphica, Hort. ex Roem. & Schult. Syst. iv. 236 = leucantha.
philippinensis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 475.—Ins. Philipp. ; China.
phoenicea, Roxb. Hort. Beng. 14 ; Fl. Ind. i. 502 = coccinea.
phylloura, *Baker, in Journ. Linn. Soc.* xxi. (1885) 426.—Madag.
physatoides, Spreng. Nov. Prov. 24 = hederacea.
physanthifolia, *Regel, Cat. Pl. Hort. Askabov.* 77, nomen.—Hab. ?
pilesta, *Roxb. Fl. Ind. ed. Carey*, ii. 94.—Ind. or. ; China.
pilosa, Cav. Ic. iv. 11 = pentaphylla.
pilosa, *Sweet, Hort. Brit.* ed. I. 289.—Ind. or. ; Afr. trop.
pilosissima, *Mart. & Gal. in Bull. Acad. Brux.* xii. 11. (1845) 364.—Mexic.
pinifolia, *Meisn. in Mart. Fl. Bras.* vii. 250.—Am. trop.
pinnata, *Hockst. ex Choisy, in DC. Prod.* ix. 358.—Abyss.
pinnatifida, G. Don, Gen. Syst. iv. 280 = Convolvulus pinnatifidus.
platanifolia, Roem. & Schult. Syst. iv. 220 = fastigiata.
platensis, [Ker-Gawl.] Bot. Reg. t. 338 = digitata.
plebeia, *R. Br. Prod.* 484.—Austral.
Pohlii, *Choisy, in DC. Prod.* ix. 355.—Bras.
polyantha, *Miq. Fl. Ind. Bat.* ii. 613.—Ind. or. ; Java.
polyantha, Roem. & Schult. Syst. iv. 234 = umbellata, *G. F. W. Mey.*
polycarpa, G. Don, Gen. Syst. iv. 270 = Convolvulus micranthus.
polygonoides, *Schweinf. Beitr. Fl. Aethiop.* 95.—Afr. trop.
polymorpha, *Riedel, ex Meisn. in Mart. Fl. Bras.* vii. 262.—Bras.
polymorpha, Roem. & Schult. Syst. iv. 254 = heterophylla.
polyrrhiza, *Choisy, in DC. Prod.* ix. 356.—Am. trop.
polystricha, Sweet, Hort. Brit. ed. II. 373 = pentaphylla.
pomacea, Roxb. ex Steud. Nom. ed. II. i. 818 = Argyreia bracteata.
porsoaides, *C. B. Clarke, in Hook f. Fl. Brit. Ind.* iv. 208.—Reg. Himal.
portobellensis, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 189.—Panama.
portoricensis, G. Don, Gen. Syst. iv. 278 = cathartica.
potentilloides, *Meisn. in Mart. Fl. Bras.* vii. 236.—Am. trop.
praecox, *Wright, in Sauv. Fl. Cub.* 107.—Cuba.
prismatoflora, *G. Don, Gen. Syst. iv.* 270.—Afr. trop.

IPOMOEA :—
Primatosyphon, *Welw. Apont.* 585. n. 18.—Afr. trop.
procumbens, *Mart. & Choisy, in DC. Prod.* ix. 351.—Bras.
procurrens, *Meisen. in Mart. Fl. Bras.* vii. 254.—Am. trop.
prostrata, *Meisen. l. c.*—Am. trop.
proximum, *Hemsl. Biol. Centr. Am. Bot.* ii. 392.—Mexic.
Prunetii, Perr. ex Steud. Nom. ed. II. i. 818 (*Brunetii*, p 816. in syn.) = dissecta.
Pseudomurucoides, Bernh. ex Link, Enum. Hort. Berol. i. 200 = grandiflora.
psilophylla, *Steud. Nom.* ed. II. i. 818.—Arab.
pterocarpa, G. Don, Gen. Syst. iv. 282 = martinicensis.
pteroclea, *Choisy, in DC. Prod.* ix. 361.—Guiana.
pterodes, Seem. Bot. Voy. Herald, 171 = slattipes.
pterygocaulis, *Choisy, in DC. Prod.* ix. 381.—Abyss.
pubescens, *Hornem. Hort. Hafn.* i. 195.—Ind. occ.
pubescens, *Lem. Illustr.* i. 465. n. 2123.—Mexic.
pudibunda, Sweet, Hort. Brit. ed. I. 489 ; G. Don, Gen. Syst. iv. 276 = catharticus.
pulchella, G. Don, l. c. = fastigiata.
pulchella, Roth, Nov. Pl. Sp. 115 = palmata.
pulchra, *Blume, Bijdr.* 716.—Ins. Timor. ; Molucc.
pulneyensis, *Badd. in Trans. Linn. Soc.* xxv. (1865) 222.—Ind. or.
pumila, *Span. in Linnaea*, xv. (1841) 841.—Ins. Timor.
punctata, *Macfad. in Hook. Bot. Misc.* ii. (1831) 116. —Ind. occ.
punctata, Vers. Syn. i. 184 = hederacea.
punctata, *Wright, in Sauv. Fl. Cub.* 105.—Cuba.
puncticulata, *Benth. Bot. Voy. Sulph.* 136.—Mexic.
punices, *Choisy, in DC. Prod.* ix. 355.—Bras.
Purga, *Hayne. Arzneigew.* xii. tt. 33, 34.—Mexic.
purpurea, *Roth, Bot. Abh.* 27.—Am. calid.
Purshii, G. Don, in Sweet, Hort. Brit ed. III. 484 = macrorhiza.
quadrangularis, *Boj. ex Choisy, in DC. Prod.* ix. 357. —Madag.
Quamoclit, *Linn. Sp. Pl.* 159.—Amphig. trop.
quinata, *R. Br. Prod.* 486.—As. et Austral. trop.
quinquefolia, *Linn. Sp Pl.* 160 ; *Vriesb. in Mem. Am. Acad.* N. S. vii. (1863) 526.—Am. trop.
quinquelaba, Desf. Tabl. ed. I. 74 = palmata.
quinqueloba, *Roem. & Schult. Syst.* iv. 208.—Ins. St Crucis.
quinqueloba, Willd. ex Choisy, Conv. Or. 54 = paniculata.
quinquepartita, *Roem. & Schult. Syst.* iv. 247.—Ind. occ.
racemosa, *Poir. Encyc. Suppl.* iv. 638.—Ins. S. Doming.
racemosa, Roth, Nov. Pl. Sp. 115 = staphylina.
radiatifolia, *Kellogg, in Proc. Calif. Acad.* vii. (1877) 168.—Calif.
radicans, Bert. ex Choisy, in DC. Prod. ix. 387, partim = I. Carolina.
radicans, Blume, Bijdr. 712 = cymosa.
radicans, Choisy, in DC. Prod. ix. 387, partim = tenera.
Ramoni, *Choisy, l. c.* 380.—Cuba.
ramosissima, *Choisy, l. c.* 377.—Peruv.
reflexa, *Span. in Linnaea*, xv. (1841) 841.—Ins. Timor.
Regnellii, *Meisen. in Mart. Fl. Bras.* vii. 266.—Am. trop.
Reineweraltiana, Blume, Bijdr. 720 = Argyreia Guichenotii.
reniformis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 446.—As. et Afr. trop.
repanda, *Vacq. Enum. Pl. Carib.* 13.—Ind. or.
repens, Lam. Illustr. i. 467. n. 2134 ; Encyc. vi. 18 = I. Beladamboë.
repens, Roth, Nov. Pl. Sp. 110 = aquatica.
reptans, Poir. Encyc. Suppl. iii. 460 = aquatica.
rhynocrinica, *Dals. in Hook. Kew Journ.* iii. (1851) 179.—Ind. or.
Riedelianus, *Oliver, in Hook. Ic. Pl.* t. 1424.—Arch. Ind.
Riedelii, *Meisen. in Mart. Fl. Bras.* vii. 285.—Am. trop.
rigaria, G. Don, Gen. Syst. iv. 265 = Beclei.
Roget, *Choisy, in DC. Prod.* ix. 381.—Afr. trop.
roseo, *Choisy, l. c.* 384.—Bras.
roseo, *Martin. Cat. Pl.* 5 ; *ex Choisy, l. c.* 390 (Quid ?). —Hab. ?

IPOMOEA :—
Rothii, Roem. & Schult. Syst. iv. 241 = cymosa.
rotundifolia, G. Don, Gen. Syst. iv. 265 = bilobe.
Roxburghii, Steud. Nom. ed. II. i. 819 = I. Bona-nox.
Roxburghii, Sweet, Hort. Brit. ed. I. 389 = Argyreia Roxburghii.
rubens, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 468 = Lettsomia rubens.
rubra, Murr. Syst. Veg. ed. XIII. 171 = Gilia coronopifolia.
rubro-caerulea, *Hook. Bot. Mag.* t. 3397.—Mexic.
ruderaria, G. Don, Gen. Syst. iv. 267 = Convolvulus havanensis.
Rudolphii, *Roem. & Schult. Syst.* iv 227.—Ins. S. Doming.
rufescens, Walp. Ann. iii. 110, sphalm. = *racemosa*, Roth.
rugosa, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 446 = I. Beladamboe.
rumicifolia, *Choisy, l. c.* 447.—Ind. or. ; Afr. trop.
Rumphii, *Mig Fl. Ind. Bat.* ii. 606.—Ins. Molucc.
sagittaefolia, *Burm. f. Fl. Ind.* 50.—Java.
sagittaefolia, Hook. & Arn. Bot. Beech. Voy. 151 = Convolvulus luteolus.
sagittifaefolia, Ker-Gawl. Bot. Reg. t. 437 = sagittata.
sagittata, Moç. & Sesse, ex Choisy, in DC. Prod. ix. 376 = emetica.
sagittata, *Poir. Voy. Barb.* ii. 122 ; *Cav. Ic.* ii. 4. t. 107. —Am. bor. ; Ind. occ.
sagittata, Roxb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 201 = calycina, *Benth.*
sagittifera, G. Don, Gen. Syst. iv. 278 = umbellata.
salicifolia, Desr. ex Steud. Nom. ed. II. i. 819 = martinicensis.
salicifolia Herb. Madr ex Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 407 = Argyreia tiliaefolia.
sallcifolia, *Roxb. Hort. Beng.* 14 ; *Fl. Ind.* i. 498.—Ind. or.
Salsmanni, Choisy, in Mém. Soc. Phys. Genèv. viii. 1. (1838) 59 = bahiensis.
sanguinea, Vahl, Symb. Bot. iii. 33 = hederifolia.
scabra, *Forsk. Fl. Aegypt. Arab.* 44.—Arabia.
scabra, J. F Gmel. Syst. 345 = hederacea.
scabrida, Roem. & Schult. Syst. iv. 238 = hederacea.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xviii. (1882-83) 123.—Mexic.
Schiedeana, *Ham. in Lindl. Bot. Reg.* (1838) *Misc.* 19. —Mexic.
Schiedeana, Zucc. in Flors, xiv. (1831) 801 = I. Purga.
schimperi, *Kunze, Del. Sem. Hort. Lips.* (1845) 2. —Abyss. Argent.
Schlechtendalii, *Vis. Hort. Patav.* (1842) 141 ; *ex Choisy, in DC. Prod.* ix. 390.—Mexic.
Schomburgkii, *Choisy, l. c.* 354.—Bras.
Schrankii, Steud. Nom. ed. II. i. 819 = Illacina, *Schrank*.
Schultzeni, Weinm. ex Choisy, in DC. Prod. ix. 395, 435 = Hewittia bicolor.
scrobiculata, *G. Don, Gen. Syst.* iv. 281.—Ins. Trinit.
secunda, *G. Don, l. c.* 283.—Afr. trop.
Sellol, *Meisen. in Mart. Fl. Bras.* vii. 271.—Bras.
Sellowii, Pensy, ex G. Don, in Loud. Hort Brit. Suppl. i. 591 = bonariensis.
semidigyna, Sweet, ex Steud. Nom. ed. II. i. 819 = Bona-nox.
senegalensis, Lam. Illustr. i. 464 = palmata.
Senegambiae, *Choisy, in DC. Prod.* ix. 351.—Afr. trop.
sepiaria, *Korn. ex Roxb. Hort. Beng.* 14 ; *Fl. Ind.* i. 496.—As. et Austral. trop.
septemaba, Choisy, in DC. Prod. ix. 362 = glabra.
sericantha, *Griseb. Fl. Brit. W. Ind.* 471.—Am. trop.
sericantha, *Mig. Stirp. Surin. Sel.* 131.—Guian.
sericea, Benth. ex Meisen. in Mart. Fl. Bras. vii. 247 = argentea.
sericea, *Parodi, Contrib.* 18.—Paraguay.
sericea, Raño. Fl. Tellur. iv. 74 ; Blume, ex Choisy, in DC. Prod. ix. 395, 431 = Argyreia mollis.
sericea, Spreng. ex Choisy, l. c. 368 = sericantha, *Griseb.*
sericophylla, *Meisen. in Mart. Fl. Bras.* vii. 260.—Am. trop.
serotina, *Roem. & Schult. Syst.* iv. 215.—Mexic.
serpens, *Meisen. in Mart. Fl. Bras.* vii. 276.—Bras.
serpyllifolia, *G. Don, Gen. Syst.* iv. 267.—Mexic.
serrata, Choisy, in Mém. Soc. Phys. Genèv. viii. 1. (1838) 57 = Jacquemontia serrata.

IPOMOEA :—
Sescossiana, *Baill. in Bull. Soc. Linn. Par.* i. (1888) 885.—Mexic.
sessiliflora, Roth, Nov. Pl. Sp. 116 = eriocarpa.
setifera, *Poir. Encyc.* vi. 17.—Guian.
setosa, Blume, Bijdr. 714 = trichocalyx.
setosa, [*Ker-Gawl.*] *Bot. Reg.* t. 335.—Bras.
setulosa, Zoll. & Mor. Syst. Verz. Zoll. 51 = limifolia.
shirensis, *Oliver, in Hook. Ic. Pl.* t. 1474.—Afr. trop.
Shumardi, Torr. in Marcy, Rep. 191 = sep.
Shumardiana, Torr. Bot. Mex. Bound. 149 = longifolia.
sibirica, *Pers. Syn.* i. 183.—Reg. Himal. ; As. bor.
sidaefolia, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 459.—Mexic.
sidaefolia, Schrad. in Goett. Gel. Anz. i. (1821) 719 = rubata.
Silvana, *Choisy, in DC. Prod.* ix. 374.—Am. trop
simplex, *Thunb. Prod. Pl. Cap.* 36.—Afr. austr.
simulans, *Hanbury, in Journ. Linn. Soc.* xi. (1871) 281. t. 2.—Mexic.
sinensis, *Choisy, in Mém. Soc. Phys Genèv.* vi. (1833) 459.—China.
sinensis, *Fisch. Hort. Gorensk.* ed. II. 38 ; (*sp. dub. ex Choisy, in DC. Prod.* ix. 390) —China.
sinuata, *Ortog Hort. Matr. Dec.* 84.—Am calid.
solanifiora, *Thunb. Mus. Nat. Ups.* 168 (sp. dub.).—Hab. ?
solanifolia, Burm. f. Fl. Ind. 49 = obscura.
solanifolia, *Linn. Sp. Pl.* 161.—Ind. occ.
soldanellaefolia, *Schrank, in Syll. Ratisb.* i. (1824) 198.—Bras.
speciosa, Pers. Syn. i. 183 = Argyreia speciosa, *Sweet*.
spectabilis, Boj. ex Choisy, in Mém. Soc. Phys. Genèv viii. 1. (1838) 68 (Cav. Ker. 146) = Breweria spectabilis.
sphaerica, Hort. Calc. ex Choisy, l. c. vi. (1833) 465 = sinuata.
sphaerocephala, D. Don, Prod. Fl. Nep. 98 = Lettsomia atropurpurea.
sphaerocephala, Sweet, Hort. Brit. ed. I. 288 ; ed. II. 373 = eriocarpa.
sphaerostigma, Poir. ex Steud. Nom. ed. II. i. 819 = Jacquemontia hirsuta.
spicaeflora, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 54.—Bras.
spicata, H. B. & K. Nov. Gen. et Sp. iii. 112 = bracteata.
spiendens, Sims, Bot. Mag. t. 2628 = Argyreia splendens.
Spruceana, *Benth. ex Meisen. in Mart. Fl. Bras.* vii. 224.—Bras.
squamosa, *Choisy, in DC. Prod.* ix. 376.—Am. trop.
stachyoides, *Meisen. in Mart. Fl. Bras.* vii. 240.—Am trop.
stans, *Cav. Ic.* iii. 26. t. 25.—Mexic.
staphylina, *Roem. & Schult. Syst.* iv. 249.—Ind. or. ; Malaya.
stellata, *Parodi, Contrib.* 18.—Paraguay.
stenocolpa, Garcke, in Linnaea, xxii. (1849) 67 = fastigiata.
stenophylla, *Meisen. in Mart. Fl. Bras.* vii. 249.—Am trop.
stipulacea, Jacq. Hort. Schoenb. ii. 39. t. 199 = palmata.
stipulacea, Sweet, Hort. Brit. ed. I. 289 ; ed. II. 372 = sepiaria.
stipulata, Jacq. Hort. Schoenb. t. 199 = calrica.
St. Nicolai, *C. Bolle, in Bonplandia*, ix. (1861) 58.—Ins. Canar.
Stocksii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 204.—Ind. or.
Stocksii, C. B. Clarke, l. c. 207 = Clarkei.
stolonifera, J. F. Gmel. Syst. 345 = carnosa.
striata, *Pers. Syn.* i. 183.—Ind. or.
striata, Roth, Nov. Pl. Sp. 111 = sepiaria.
strigosa, Roth, l. c. 118 = Lettsomia setosa.
stylosa, Hort. Madr. ex Choisy, in DC. Prod. ix. 395, 435 = coccinea.
suaveolens, *Hemsl. Biol. Centr. Am. Bot.* ii. 394.—Mexic.
subdentata, Miq. Fl. Ind. Bat. ii. 614 = aquatica. dentiniata.
subincana, *Meisen. in Mart. Fl. Bras.* vii. 259.—Am. trop.
subpeltata, Desf. Cat. Hort. Par. ed. III. 397 = dissecta.
submollis, *Choisy, in DC. Prod.* ix. 376.—Guian.
subtrifiora, Zoll. & Mor. Syst. Verz. Zoll. 51 = chrysoides.

IPOMOEA :—
subtriloba, Ruiz & Pav. Fl. Per. ii. 12 = pubescens.
subtrilobata, Miq. Fl. Ind. Bat. ii. 618 = separata.
suffruticosa, *Burch. Trav. Afr.* ii. 226.—Afr. austr.
suffulta, *G. Don, Gen. Syst.* iv. 276.—Mexic.
sulcata, *Parodi, Contrib.* 19.—Paraguay.
sulphurea, G. Don, Gen. Syst. iv. 266 = corralinensis.
sulphurea, *Hochst. ex Choisy, in DC. Prod.* ix. 356.—Afr. trop.
superba, *G. Don, Gen. Syst.* iv. 275.—Mexic.
superba, *Ledeb. Ind. Sem. Hort. Dorpat.* (1822) 11 ; *ex Choisy, in DC. Prod.* ix. 390 (Quid ?).—Hab. ?
superba, *Schrank, ex Colla, Hort. Ripul. App.* ii. 350, 351 (Quid ?).—Am.
surinamensis, *Miq. in Linnaea*, xviii. 544.—Surinam.—Guian.
syringaefolia, Buek, Ind. ad DC. Prod. iii. 338 = Silvana.
syringaefolia, *Meissn. in Mart. Fl. Bras.* vii. 270.—Am. trop.
tamnifolia, Burm. f. Fl. Ind. 50 = I. Pes-tigridis.
tamnifolia, Linn. Sp. Pl. 161 = Jacquemontia tamnifolia.
tenera, *Meissn. in Mart. Fl. Bras.* vii. 239.—Am. trop.
tenuiloba, *Torr. Bot Mex. Bound.* 146.—Texas.
tenuirostris, *Choisy, in DC. Prod.* ix. 379.—Abyss.
tenuissima, *Choisy, l. c.* 376.—Ins. S. Doming.
tenuissima, *Choisy, l. c.* 378.—Afr. trop.
terminalis, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 54.—Bras.
ternata, *Jacq. Hort. Schoenb.* i. 16. t. 37.—Hab. ?
ternifolia, *Cav. Ic.* v. 69. t. 478. f. 1.—Mexic.
Thomsoniana, Mast. in Gard. Chron. (1888) ii. 818 = ternata.
Thouningii, G. Don, Gen. Syst. iv. 280 = dissecta.
Thurberi, *A. Gray, Syn. Fl. N. Am.* ii. 1. 212.—Arizona.
tigrina, Pers. Syn. i. 188 = I. Pes-tigridis.
tigrina, Stokes, Bot. Mat. Med. i. 330 = I. Pes-tigridis.
tiliacea, Choisy, in DC. Prod. ix. 375 = fastigiata.
tiliaefolia, Roem. & Schult. Syst. iv. 229 = Argyreia tiliaefolia.
timorensis, Blume, Bijdr. 711 = Hewittia bicolor.
tomentosa, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 55.—Bras.
trematosperma, Hochst. ex Choisy, in DC. Prod. ix. 367 = sessiliflora.
triantha, Roem. & Schult. Syst. iv. 248 = separia ?
trichocalyx, G. Don, Gen. Syst. iv. 275 = ochracea.
trichocalyx, *Steud. Nom.* ed. II. i. 819.—Java.
trichocarpa, Ell. Sketch. ii. 260 = commutata.
trichocephala, G. Don, Gen. Syst. iv. 269 = Jacquemontia tamnifolia.
trichosperma, *Blume, Bijdr.* 710.—Ind. or. ; Burma ; Malaya.
trichotosa, *Blume, l. c.* 717.—Java.
tricolor, *Cav. Ic.* iii. t. 1. 208.—Hab. ?
tridentata, *Roth, in Roem. Archiv*, i. ii. 88.—Ind. or. ; Malaya.
trifida, *G. Don, Gen. Syst.* iv. 280.—Am. trop.
trifida, *Forsk. Fl. Aegypt. Arab.* 44.—Arab.
trifiora, *Maria & Velasco, in La Naturaleza* (1870), 388.—Mexic.
trifoveata, Choisy, in Mém. Soc. Phys. Genèv. viii. 1. (1838) 53 = polymorpha.
triloba, *Linn. Sp. Pl.* 161.—Am. trop.
triloba, Thunb. Fl. Jap. 86 = bederacea.
triquetra, Roem. & Schult. Syst. iv. 231 = I. Turpethum.
Tuba, G. Don, Gen. Syst. iv. 271 = grandiflora.
tubata, *Nees, in Flora*, iv. (1821) 301.—Bras.
tuberculata, *Denr. ex Regel, Ind. Sem. Hort. Petrop.* (1856) 27.—Ins. Borbon.
tuberculata, [Ker-Gawl.] in Bot. Reg. t. 86 = dasysperma.
tuberculata, Roem. & Schult. Syst. iv. 208 = palmata.
tuberculosa, Desf. Tabl. ed. I. 74 = palmata.
tuberosa, *Linn. Sp. Pl.* 160.—Reg. trop.
tuberosa, Lour. Fl. Cochinch. 136 = palmata.
tuberosa, G. F. W. Mey. Prim. Fl. Esseq. 118 = digitata.
tuberosa, A. Rich. Tent. Fl. Abyss. ii. 67 = kentrocaulos.
tubiflora, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 204.—Ins. Galapag.
tubulosa, *Willd. ex Roem. & Schult. Syst.* iv. 789.—Am. austr.
turbinata, Lag. ex Choisy, in DC. Prod. ix. 396, 345 = muricata.
Turpethum, *R. Br. Prod.* 485.—As. et Austral. trop.
Tweediei, *Hook. Bot. Mag.* t. 8978.—Reg. Argent.
tyrianthina, *Lindl. in Bot. Reg.* (1836) *Misc.* 87.—Mexic.

IPOMOEA :—
ukambensis, *Vatke, in Linnaea*, xliii. (1880-82) 510.—Afr. trop.
umbellata, *Linn. Syst.* ed. X. 924.—Am. austr.
umbellata, *G. F. W. Mey. Prim. Fl. Essq.* 99.—Mexic.
umbelliflora, *Choisy, in DC. Prod.* ix. 389.—Am. trop.
umiflora, Roem. & Schult. Syst. iv. 247 = martiuncnsis.
urbica, Choisy, in DC. Prod. ix. 349 = asarifolia.
urcolata, *R. Br. Prod.* 485.—Australd.
uruguensis, *Meissn. in Mart. Fl. Bras.* vii. 272.—Am. trop.
utilis, *Choisy, in DC. Prod.* ix. 376.—Am. trop.
varia, Roth, Catal. fasc ii. 17 = pubescens, *Hornem.*
variabilis, *Choisy, in DC. Prod.* ix. 388.—Mexic.
variifolia, *Meissn. in Mart. Fl. Bras.* vii. 275.—Am. trop.
Vellosiana, Mart. in Flora, xxi. (1838) ii. Beibl. 64 = Jacquemontia grandiflora.
venitsua, *R. Br. Prod.* 485.—Austral.
venosa, *Roem. & Schult. Syst.* iv. 219.—Ins. Mascar.
ventricosa, *G. Don, Gen. Syst.* iv. 274 ; *Choisy, in DC. Prod.* ix. 360.—Ind. occ.
venustum, *Hemsl. Biol. Centr. Am. Bot.* ii. 395.—Mexic.
verbascoides, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 56.—Afr. trop.
vernetefiua, Meisan. in Mart. Fl. Bras. vii. 267 = pentaphylla.
vernicifolia, *Arn. in Madr. Journ. Sc.* Ser. I. v. (1837) 10.—Ind. or.
vernicipes, *C. A. Mey. in Fisch. Mey. & Ave Lall. Ind. Sem. Hort. Petrop.* ix. 76 ; *Choisy, in DC. Prod.* ix. 377.—Bras.
verrucosa, *Blume, Bijdr.* 718.—Java.
verrucosa, Hortg. Hort. Matr. Dec. 10 = dissecta.
verruculosa, Mart. ex Choisy, in DC. Prod. ix. 378 = verrucipes.
vernicolor, *Meissn. in Mart. Fl. Bras.* vii. 220.—Am. trop.
verticillata, *Forsk. Fl. Aegypt. Arab.* 44.—Arabia.
verticillata, Stend. Nom. ed. II. i. 820 = Jacquemontia hirsuta.
vesiculosa, Beauv. Fl. Owar. ii. 78. t. 106 = palmata ?
vespertina, *Colla, Hort. Rippul App* iii. 40.—Bras. ?
villosa, *Meissn. in Mart. Fl. Bras.* vii. 244.—Bras.
villosa, *Ruiz & Pav. Fl. Per.* ii. 12.—Peruv.
villosula, *Blume, ex Steud. Nom.* ed. II. i. 820.—Java.
violacea, Lin. Sp. Pl. 161.—Ins. St. Crucis.
virgata, *Meissn. in Mart. Fl. Bras.* vii. 241.—Am. trop.
viridis, *Choisy, in DC. Prod.* ix. 374.—Am. trop.
vitifolia, *Sweet, Hort. Brit.* ed. I. 289.—Ind. or. ; Malaya.
Waldeckii, *Regel, Cat. Pl. Hort. Absakow.* 77.—Hab. ?
Wallichii, Steud. Nom. ed. II. i. 820 = Leptatemon Wallichii.
Wallii, *Hemsl. Biol. Centr. Am. Bot.* ii. 396.—Mexic.
Warmingii, *Meissn. in Mart. Fl. Bras.* vii. 272.—Am. trop.
Weinmanni, Roem. & Schult. Syst. iv. 248 = Hewittia bicolor.
Wightii, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 470.—Ind. or.
Wildenowii, Roem. & Schult. Syst. iv. 211 = heterophylla.
Wrightii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 218.—Texas.
xanthantha, Kurz, For. Fl. Brit. Burma, ii. 219 = petaloides.
Yetira, *Parodi, Contrib.* 17.—Paraguay.
Yomae, Kurz, For. Fl. Brit. Burma, ii. 218 = trichosperma.
zebrina, *Porr. ex Choisy, in DC. Prod.* ix. 382.—Afr. trop.
zeylanica, Gaertn. Fruct. ii. 462 = Argyreia populifolia.
Zuccagnii, Roem. & Schult. Syst. iv. 230 = purpurea.

IPOMOPSIS, Michx. Fl. Bor. Am. i. 141 (1803) = **Gilia**, Ruiz & Pav. (Polemoniac.).
elegans, Lindl. Bot. Reg. t. 1881 = G. aggregata.
elegans, Michx. Fl. Bor. Am. i. 143 = G. coronopifolia.
inconspicua, Sm. Exot. Bot. i. 25 = G. inconspicua.
picta, Hort. Gall. ex G. Don, Gen. Syst. iv. 344 = G. coronopifolia.
tuberosa, Willd. ex Stend. Nom. ed. II. i. 820 = Ipomoea muricata.
verticillata, Willd. ex Roem. & Schult. Syst. iv. 793 = Ipomoea muricata.

IPSEA, Lindl. Gen. et Sp. Orch. 124 (1831) = **Pachystoma**, Blume (Orchid.).
speciosa, Lindl. l. c. = P. sessile.

IRASEKIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 229 (1821) = IRASEKIA, F. W. Schmidt = **Anagallis**, Linn. (Primul.).

IRAUPALOS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 87 = **Viburnum**, Linn. (Caprifol.).

IRENEIS, Moq. in DC. Prod. xiii. ii. 349 (1849) = **Iresine**, Linn. (Amarant.).
interrupta, Moq. l. c. = Iresine interrupta.

IREON, Burm. f. Prod. Fl. Cap. 6 (1768) = **Roridula**, Linn. (Droserac.).
verticillatum, Burm. f. l. c. = R. dentata.

IREON, Scop. Introd. 229 (1777) = **Prismatocarpus**, L'Hérit. (Campanul.).
ciliatum, Rafin. Sylva Tellur. 138 = P. subulatus.

IRESINE, P. Br. Hist. Jamaic. 358 (1756). *AMARANTACEAE*, Benth. & Hook. f. iii. 49.
CRUCITA, Linn. Sp. Pl. ed. II. 179 (1762).
CRUZETA, Loefl. It. Hisp. 208 (1758).
IRENEIS, Moq. in DC. Prod. xiii. ii. 349 (1849).
ROSEA, Mart. Nov. Gen. et Sp. ii. 98 (1826).
XERANDRA, Rafin. Fl. Tellur. iii. 48 (1836).
acuminata, *Moq. in DC. Prod.* xiii. ii. 345.—Mexic.
acutifolia, Sw. ex Moq. l. c. 343 = elatior.
aggregata, Moq. l. c. 340 = Philoxerus verniculatus.
amarantoides, *Moq. l. c.* 345.—Malaya.
angustifolia, *Euphrasén, Beskr. St. Barthel.* 165.—Ind. occ.
angustifolia, Hort. ex Moq. in DC. Prod. xiii. ii. 343 = elatior.
angustifolia, Rich. ex Moq. l. c. 348, 353 = interrupta.
argentiata, D. Dietr. Syn. Pl. i. 870 = Hebenthe rariflora ?
aurata, D. Dietr. l. c. = Alternanthera aurata
bonariensis, Miers, ex Seub. in Mart. Fl. Bras. v. i. 227 = celosioides.
Brownii, Moq. in DC. Prod. xiii. ii. 341 = Gomphrena diffusa.
canescens, Bonpl. ex Moq. l. c. 349, 352 = Hebenthe parviflora !
canescens, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 765.—Mexic.
cassiniaeformis, *Schau. in Linnaea*, xix. (1847) 709.—Mexic.
Celosia, Linn. Syst. ed. X. 1291 = seq.
celosioides, *Linn. Sp. Pl.* ed. II. 1456.—Am. bor.
celosioides, Sw. Obs. Bot. 376 = elatior.
conica, Moq. in DC. Prod. xiii. ii. 342 = Gomphrena conica.
crassifolia, *Moq. l. c.* 340.—Ind. occ.
Cunninghamii, Moq. l. c. 342 = Gomphrena Maitlandi.
diffusa, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 766 = celosioides.
dyscitcta, Spreng. Syst. iv. Cur. Post. 104 = Alternanthera aurata.
Edmonstonei, *Hook. f. in Trans. Linn. Soc.* xx. (1847 190.—Ins. Galapag.
elatior, *Rich. ex Willd. Sp. Pl.* iv. 766.—Am. trop.
elatior, Sieber, ex Moq. in DC. Prod. xiii. ii. 349, 353 = Alternanthera aurata.
elongata, Hook. & Arn. Bot. Beech. Voy. 308 = I. Hookeri.
elongata, *Humb. & Bonpl. in Willd. Sp. Pl.* iv. 765.—N. Granat.
erianthos, Poir. Encyc. Suppl. iii. 160 = Hebenthe virgata.
eriophora, *Poyr. in Linnaea*, xxx. (1859-60) 91.—Mexic.
eriophylla, Moq. in DC. Prod. xiii. ii. 347 = celosioides.
flavescens, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 766 = Alternanthera flavescens.
floribunda, Mart. & Gal. in Bull. Acad. Bruz. x. (1843) 347 = celosioides.
frutescens, *Moq. in DC. Prod.* xiii. ii. 344.—Mexic.
glabra, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 933, nomen.—Guian.
glomerata, Spreng. Neue Entdeck. ii. 110 = Pfaffia glauca.
gnaphalantha, *A. Rich. Fl. Cub. Fanerog.* ii. 177.—Cuba.

IRESINE:—
gossypina, *Nees, in Linnaea*, xix. (1847) 709.—Mexic.
gracilis, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 847.—Mexic.
grandiflora, Hook. Ic. Pl. t. 102 = Hebanthe deciplens.
havanensis, *H. B. & K. Nov. Gen. et Sp.* ii. 199.—Cuba.
herbacea, Desf. Tabl. ed. II. 53 = celosioides.
Herbstii, *Hook. in Gard. Chron.* (1864) 654, 1206.—Bras.
Hookeri, *Moq. in DC. Prod.* xiii. 11. 344.—Mexic.
interrupta, *Benth. Bot. Voy. Sulph.* 156.—Mexic.
javanica, Burm. f. Fl. Ind. 213 (err. typ. 312) = Aerva tomentosa.
lanceolata, *Moq. in DC. Prod.* xiii. 11. 347.—Hab.?
latifolia, *Benth. & Hook. f. Gen.* iii. 42.—Mexic.
latifolia, Moq. in DC. Prod. xiii. 11. 351 = Alternanthera pulverulenta.
Lindeni, *Van Houtte, in Fl. des Serres*, Sér. II. vii (1867-68) 41.—Hab.?
linearis, *Moq. in DC. Prod.* xiii. 11. 339.—Ind. occ.
luxuliflora, *Griseb. Fl. Brit. W. Ind.* 65.—Am. trop., Reg. Argent.
macrocephala, Moq. in DC. Prod. xiii. 11. 342 = Gomphrena conferta.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* ii. 200.—N. Granat.
obtusifolia, H. B. & K. l. c. 198 = Alternanthera flavescens.
paniculata, Poir. Encyc. Suppl. iii, 181 = elatior.
panniculata, Spreng. Syst. iv. Cur. Post. 108 = Hebanthe paniculata, *Mart.*
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent* v. (1878) 316; *Contrib.* 88.—Reg. Argent.
parvifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 198.—Cuba.
persica, Burm. f. Fl. Ind. 212 (err. typ. 312) = Aerva tomentosa.
polymorpha, Mart. Nov. Gen. et Sp Bras ii. 56. t. 153 = celosioides.
spicaligera, *Moq. in DC. Prod.* xiii. 11. 341.—Bras.
pulverulenta, D. Dietr. Syn. Pl. i. 870 = Alternanthera pulverulenta.
racemosa, Poir. Encyc. Suppl. iii. 180 = elatior.
reticulata, Mart.—Cf. Gard. Chron. (1874) i. 415 = Herbstii.
scandens, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 87; *Contrib.* 89.—Paraguay.
serpyllifolia, *Moq. in DC. Prod.* xiii. 11. 341.—Ins. S. Doming.
silvatica, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 320; *Contrib.* 88.—Reg. Argent.
spicata, Spreng. Syst. iv. Cur. Post. 104 = Hebanthe spicata.
spiciligera, *Seub. in Mart. Fl. Bras.* v. t. 228.—Bras.
surinamensis, *Moq. in DC. Prod.* xiii. 11. 339.—Guian.
tenella, Moq. l. c. 348 = Gomphrena tenella.
vermicularis, Moq. l. c. 340 = Philoxerus vermicularis.
Verschaffelti, *Lem. ex Van Houtte, Fl. des Serres*, xv. (1863-65) 101. t. 1501.—Bras.
verticillata, Spreng. Syst. i. 821 = celosioides.
virgata, Poepp. ex Moq. in DC. Prod. xiii. 11. 343 = elatior.
virgata, Spreng. Syst. iv. Cur. Post. 104 = Hebanthe virgata.
Wallisii, *Orty. in Regel, Gartenfl.* (1878) 80.—N. Granat.?

IREUM, Stend. Nom. ed. II. ii. 465, in syn. (1840) = IREON, Burm. f. = **Boridula**, Linn. (Droserac.).

IRIA, Hedw. f. Gen. 360 (1806) = **Fimbristylis**, Vahl (Cyperac.).
caribæa, Stend. Nom. ed. I. 437 = F. monostachya.
indica, Schult. Mant. ii. 94 = F. monostachya.

IRIARTEA, Ruiz & Pav. Prod. Fl. Per. 149. t. 32 (1794). *PALMAE*, Benth. & Hook. f. iii. 900.
DECKERIA, Kunt. in Linnæa, xxviii. (1856) 258.
afinis, *Karst. ex Linden, Illustr. Hortic.* xxviii. (1881) 81.—N. Granat.
altissima, *Klotzsch, ex Linden, l. c.* 81.—Venezuela.
andicola, Spreng. Syst. ii. 638 = Ceroxylon andicolum.
concuta, *H. Wendl. in Bonplandia*, viii. (1860) 102.—Bras.

IRIARTEA:—
costata, *Linden, Illustr. Hortic.* xxviii. (1881) 81.—N. Granat.
deltoidea, *Ruiz & Pav. Syst.* 298.—Peruv.
durissima, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 80 = Socratea durissima.
exorrhiza, Mart. Hist. Nat. Palm. ii. 36. t. 33, 34 = Socratea exorrhiza.
gigantea, *H. Wendl in Kerch. Palm.* (1878) 262. f. 155.—Hab.?
glaucescens, *Linden. Illustr. Hortic.* xxviii. (1881) 81—N. Granat.
Lamarckiana, Mart. Hist. Nat. Palm. iii. 190 = Dictyocaryum Lamarckianum.
leprosa, Zipp in Bijdr Wetensch. v. 178 = Drymophloeus olivaeformis.
monogyna, Zipp. l. c. 178 = Drymophloeus appendiculatus.
Orbignyana, Mart. Hist Nat. Palm. iii. 187 = Socratea Orbignyana.
phaeocarpa, *Mart. l. c.* 196.—Bolivia.
phlonotisa, Barb-Rodr. Enum. Palm. Nov. 13 = Socratea exorrhiza.
praemorsa, Klotzsch, in Linnæa, xx. (1847) 448 = Catoblastus praemorsus.
prurisens, Spruce, in Journ. Linn. Soc. xi. (1871) 136 = Iriartella setigera.
pubescens, Karst. in Linnæa, xxviii (1856) 262 = Catoblastus pubescens.
pygmæa, *Linden, Illustr. Hortic.* xxviii. (1881) 81.—Bras.
robusta, Hort. Versch. ex H. Wendl. in Kerch. Palm. 247 = deltoidea.
setigera, Mart. Hist. Nat. Palm. ii. 39. t. 37 = Iriartella setigera.
Spruciana, Barb-Rodr. Enum. Palm. Nov. 13 = Iriartella setigera.
ventricosa, *Mart. Hist. Nat. Palm.* ii. 37. t. 35, 36.—Bras.
xanthorhiza, *Klotzsch, ex Linden, Illustr. Hortic.* xxviii (1881) 81.—Venezuela.
zamonia, *Linden, l. c.*—Ecuador.

IRIARTELLA, H. Wendl. in Bonplandia, viii. (1860) 103. *PALMAE*, Benth. & Hook. f. iii. 901.
setigera, *H. Wendl. l. c.* 104.—Bras.

IRIDION, Roem. & Schult. Syst. v. 436, in syn. (1819) = IREON, Burm. f. = **Boridula**, Linn. (Droserac.).

IRIDOPSIS, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 270 = **Moraea**, Linn. (Irid.).
textilis, Welw. ex Baker, l. c. = M. textilis.

IRIDORCHIS, Blume, Orch. Archip. Ind. 90. t. 26 (1858) = **Cymbidium**, Sw. (Orchid.).
gigantea, Blume, l. c. 91 = C. giganteum.

IRIDROGALVIA, Pers. Syn. i. 399, in syn. (1805), sphalm. = ISIDROGALVIA, Ruiz & Pav. = **Tofieldia**, Huds. (Liliac.).

IRILLIUM, Kunth, Enum. Pl. v. 121 (1850), sphalm. = **Trillium**, Mill. (Liliac.).

IRINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2; Blume, Bijdr. 330 (1826) = **Pometia**, Forst (Sapindac.).
alnifolia, Blume, Rumphia, fil. 117.—Borneo.
Diplocordia, Blume, l. c. 118.—N. Guinea.
glabra, Blume, Bijdr. 330 = P. tomentosa.
integerrima, Blume, l. c. 331 = Meliosma sumatrana.
tomentosa, Blume, l. c. = P. tomentosa.

IRINE, Hassk. Pl. Jav. Rar. 238 (1848) = prsec.

IRIO, Linn. Syst. ed. I (1735) = **Claymbrium**, Tourn. (Crucif.).
ruderalis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 331 = S. Irio.

IRIPA, Adans. Fam. ii. 508 (1763) = **Cynometra**, Linn. (Legumin.).

IRIS, Tourn. ex Linn. Syst. ed. I (1735). *IRIDEAE*, Benth. & Hook. f. iii. 686.
BEVERNA, Adans. Fam. ii. (20) (1763).
IRIS, Medic. in SCHREB. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 357; Phil. Bot. ii. 97 (1791).

IRIS:—
CHAMOLETTA, Adans. Fam. ii. 60 (1763).
CORESANTHE, Alef. in Bot. Zeit. xxi. (1863) 398.
COSTIA, Willk. l. c. xvii. (1860) 181.
DIAPHANE, Salisb. in Trans. Hort. Soc. i. (1812) 305.
EVANSIA, Salisb. l. c. 308 (1812).
GATTENHOFIA, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 418.
GYNANDRIRIS, Parl. Nuov. Gen. e Sp. 49 (1854).
HERMODACTYLUS, Adans. Fam. ii. 60 (1763).
IONIRIS, Baker, in Gard. Chron. (1876) ii. 148, in obs.
Isis, Tratt. Arch. Gew. 668 (1812).
JONIRIS, Klatt, in Bot. Zeit xxx. (1872) 502.
JUNO, Tratt. in Roem. & Schult. Syst. i. 471, 474 (1817).
LIMNIRION, Opiz, Seznam, 59 (1852).
LIMNIRIS, Fuss. Fl. Transs. 636 (1866).
NEUBECKIA, Alef. in Bot. Zeit. xxi. (1863) 290, 297.
ONCOCYCLUS, Siemss. in Bot. Zeit. iv. (1846) 706.
PSEUDO-IRIS, Medic. in Act Acad. Theod. Palat. vi. Phys. (1790) 417.
SISYRINCHIUM, Adans. Fam. ii. 60 (1763).
SPATHULA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 163.
THELYSIA, Salisb. in Trans. Hort. Soc. i. (1812) 308.
XERIS, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 256.
XIPHION, Tourn. ex Medic. Phil. Bot. i. 173 (1789).
XURIS, Adans. Fam. ii. (20) (1763).
XYRIDION, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii (1869) 163; Klatt, in Bot. Zeit. xxx. (1872) 499.
acaulis, Pall. Rese, iii 213 = tenuifolia.
acoriformis, Bor. Fl. Cent. Fr. ed. III ii. 635 = I. Pseudacorus.
acoroides, Spach, Hist. Vég. Phan. xiii. 44 = I. Pseudacorus.
acuta, Willd. Enum. Hort. Berol. Suppl. 4 = sibirica.
acutiloba, *C. A. Mey. ex Regel, Gartenfl.* (1874) 223. t. 873.—Reg. Caucas.
acutiloba, *C. A. Mey. Vers. Pfl. Cauc.* 32.—Reg. Caucas.; Persia.
Adami, Willd. ex Link, Jahrb. i. iii. 72 = graminea.
aegyptia, Delile, in Laborde, Voy. Arab. Petr. 82 = I. Sisyrinchium.
aequiloba, Ledeb. Ind. Sem. Hort. Dorp. (1828) 4 = pumila.
Aitchisoni, *Baker, in Journ. Bot.* xiii. (1875) 108.—Afghan.
alata, *Pour. Voy. Barb* ii. 86.—Reg. Mediterr.
alba, Savi, Fl. Pis. i. 82 = florentina.
Alberti, *Regel, in Act. Hort. Petrop.* v. (1877) 260.—Turkestan.
albicans, Lange, in Kjoeb. Vidensk. Meddel. (1860) 76 = florentina.
alpina, Pall. ex Roem. & Schult. Syst. i. 476 = humilis.
amoena, [DC. in] Red. Liliac. t. 336 = hybrida.
angica, Hort. ex Steud. Nom. ed. I. 437 = siphioides.
angusta, Thunb. Diss. Irid. 19 = Moraea angusta.
angustifolia, Gilib. Exercit. Bot. ii. 498 = sibirica.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 13 = pumila.
aphylla, *Linn. Sp. Pl.* 38.—Europ.; Reg. Caucas.
arenaria, *Waldst. & Kit. Pl. Rar. Hung.* i. 57. t. 57.—Hungar.
atroviolacea, *Lange, in Bot. Tidsk.* xiii. (1882) 18.—Hab.?
attica, Boiss. & Heldr. in Boiss. Diagn. Ser. II. iv. 91 = pumila.
aurea, *Lindl. Bot. Reg.* (1847) t. 59.—Reg. Himal.
aurea, Link, Enum. Hort. Berol. i. 59 = Gueldenstaedtiana.
australis, Tod. Nuov. Gen. e Sp. 49 = pallida.
autumnalis, Tausch, in Flora, xvii. (1834) 523 = cenata?
balkana, Janka, in Magyar Tud. Akad. Math. Term. xii. (1874) 178. t. 14 = I. Chamaeiris.
Bartoni, *Foster, in Gard. Chron.* (1868) i. 275.—Afghan.
Bastardi, Bor. Fl. Cent. Fr. ed. III. ii. 635 = I. Pseudacorus.
bayonnensis, Darrach. ex Gren. & Godr. Fl. Fr. iii. 248 = graminea.

IRIS :—
Beecheyana, Herb. in Hook. & Arn. Bot. Beech. Voy. 395 = Douglasiana.
bicolor, Lindl. Bot. Reg. t. 1404 = Moraea bicolor.
bicolor, Mill. Gard. Dict. ed. VIII. n. 18 = sibirica.
biflora, Falk, Beitr. ii. n. 59 = glaucescens.
biflora, *Linn. Sp. Pl.* 38.—Lusitan. ; Tauris.
biflora, Pall. Reise, i. 171 = aphylla.
bifurca, Stev. ex Baker, in Journ. Linn. Soc. xvi. (1877) 146 = aphylla.
biglumis, Vahl, Enum. ii. 149 = ensata.
binata, Schur, in Oestr. Bot. Zeit. (1860) 354 = virescens.
biglorera, Host, Fl. Austr. i. 46 = aphylla.
bispathacea, Spach, Hist. Vég. Phan. xiii. 15 = tuberosa.
bituminosa, Linn f. Suppl. 98 = Moraea bituminosa. Blondowii, *Ledeb. Fl. Alt.* iv. 231.—Sibir. alt.
bohemica, F. W. Schmidt. Fl. Boem. iv. 5 = aphylla.
Boissieri, *Henrq. in Bol. Soc. Broter.* iii. (1885) 183. —Hispan.
Boltoniana, Hort. ex Regel, Ind. Sem. Hort. Petrop. (1860) 31 = pulchella.
Boltoniana, Schult. Mant. i. 306 = prismatica.
brachycarpa, Ledd. ex Stend. Nom ed. II. i. 321 = seq. ?
brachycuspis, Fisch. ex Sims, Bot. Mag. t. 2826 = setosa.
brachystigma, *Schrele, in Linnaea*, xxii. (1849) 848.—Texas.
bracteata, *S. Wats. in Proc. Am. Acad.* xx. (1885) 375.—Am. bor. occ.
breviculis, *Rafin. Fl. Ludov.* 20 (Quid ?).—Am. bor.
brevicuspis, Fisch. ex Schult. Mant. i. 306 = setosa.
brevicspa, *Opiz, Naturol.* x. (1825) 191 (Quid ?). —Europ.
Bungei, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 569.—Mongol.
caesia x, E. Berg, in Flora, xviii. (1835) 563.
caespitosa, *Berb. in Oestr. Bot. Zeitschr.* xxviii. (1878) 264, nomen.—Europ.
caespitosa, Pall. ex Link, Jahrb. i. iii. 71 = ruthenica.
capensis, Burm. f. Fl. Cap. Prod. 2 = Moraea edulis.
carinifolia, Pall. ex Link, Jahrb. i. iii. 73 = ensata.
Carolina, *Radius, ex Stend. Nom.* ed. II. i. 321.—Carolina.
catenulata, Baker, in Journ. Linn. Soc. xvi. (1877) 147 = Moraea catenulata.
caucasica, *Hoffm. in Comm. Soc. Phys. Mosq.* i. (1806) 40.—Reg. Caucas.
caurina, Herb. ex Hook. Fl. Bor. Am. ii. 206 = versicolor.
Cengialii, Ambrosi, Fl. Tirol. Austr. i. 649 = pallida.
cepafolia, Stokes, Bot. Comment. t. 233 = xiphioides. Chamaeiris *Bertol. Fl. Ital.* iii. 409.—Europ. austr.
chinensis, Curt. Bot. Mag. t. 378 = japonica.
ciliata, Linn. f. Suppl. 98 = Moraea ciliata.
clandestina, *Lehm. Sem. Hort. Hamb.* (1828) 16.—Bras.
Clusiana, Reichb. Ic. Fl. Germ. t. 238 = pumila.
Clusiana, Tausch, in Flora, xii. (1829) 1. Erg. 49 = aphylla.
coelestina, *Sweet, Hort. Brit.* ed. I. 394, nomen.—Am. bor.
convulsa, Spach, Hist. Vég. Phan. xiii. 50 = pumila.
collina, Salisb. Prod. 45 = I. Sisyrinchium.
Comitissae, *Andrs. ex Trautv. in Act. Hort. Petrop.* ix. (1886) 240, nomen.—Rossia.
compressa, Linn. f. Suppl. 98 = Moraea iridioides.
compressa, Moench, Meth. 539 = gramines.
compressa, Thunb. Diss. Irid. 12 = moraeoides.
concolor, Baker, in Gard. Chron. (1876) ii. 774 = sambucina.
convoluta, *Rafin. New Fl. Am.* ii. 93.—Am. bor.
coronaria, Salisb. Prod. 45 = I. Xiphium.
crassifolia, Ledd. Bot. Cab. t. 1861 = Moraea iridioides.
crocosmia, *Yanke, in Oestr. Bot. Zeitschr.* (1867) 376. —Graecia ; Syria.
crispa, Linn. f. Suppl. 98 = Moraea crispa.
cristata, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 70.—Am. bor.
crocea, Jacq. ex Baker, in Journ. Linn. Soc. xvi. (1877) 141 = aurea.
cuculiata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 74 = pallida.
cuprea, Pursh, Fl. Am. Sept. i. 50 = fulva.

IRIS :—
curtopetala, [Delar. in] Red. Lil. t. 340 = I. Pseudacorus.
daemonia, Kotschy, ex Baker, in Journ. Linn. Soc. xvi. (1877) 140 = spuria.
dahurica, *Herb. ex Klatt, in Bot. Zeit.* xxx. (1872) 514.—Dahuria.
Dandfordiae, *Boiss. Fl. Orient.* v. 124.—Cilicia.
darwasica, *Regel, in Act. Hort. Petrop.* viii. (1884) 679.—Bokhara.
decora, Wall. Fl. As. Rar. i. 77. t. 86 = nepalensis.
defexa, Knowles & Westc. Flor. Cab. ii. (1838) 19. t. 51 = germanica.
desertorum, Balb. Cat. Hort. Taur. App. i. 12 = Swertii.
desertorum, Gueldenst. Reise, i. 80 = spuria.
desertorum, Moench, Meth. 539 = Gueldenstaedtiana.
diantha, C. Koch, in Linnaea, xxi. (1848) 687 = aphylla.
dianthe, Schur, in Oestr. Bot. Zeitschr. x. (1860) 354 = virescens.
dichotoma, *Pall. Reise*, iii. 712.—Dahuria.
Dirrichii, Hort. Berol. ex C. Koch, Ind. Sem. Hort. Berol. (1853) 17 = spuria.
dilata, Bieb. Cent. i. Fl. Ross. t. 81 = Gueldenstaedtiana.
diversifolia, Steud. ex A. Rich. Tent. Fl. Abyss. ii. 305 = Moraea diversifolia.
Dominae, Spach, Hist. Vég. Phan. xiii. 84 = ensata.
Douglasiana, *Herb. in Hook. & Arn. Bot. Beech. Voy.* 395.—Am. bor. occ.
Douglasiana, Torr. Bot. Whipple Exped. 144 = macrosiphon.
dubia, Poir. Encyc. Suppl. iii. 184 = Gueldenstaedtiana.
Duetschii, *Buchl. in Am. Journ. Sc.* xlv. (1843) 176. —Am. bor.
edulis, Linn. f. Suppl. 98 = Moraea edulis.
elegans, *Pers. ex Steud. Nom.* ed. II. i. 83.—Hab. ?
elongata, Fisch. ex Baker, in Journ. Linn. Soc. xvi. (1877) 144 = glaucescens.
ensata, *Thunb. in Transc. Linn. Soc.* ii. (1794) 328.—Reg. Himal. ; As. bor.
erratica, Tod. Nuovi Gen. e Sp. 10 = lutescens.
Eulefeldi, *Regel, in Act. Hort. Petrop.* v. (1877) 633. —Turkestan.
extrafoliacea, Mikan, f. ex Roem. & Schult. Syst. i. 466 = aphylla.
falcata, Tausch, in Flora, vii. (1824) 83 = aphylla.
falcifolia, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 505.—Afghan. ; Turkest.
Falconeriana, Penny, ex Lond. Gard. Mag. v. (1829) 47 = germanica ?
Fieberi, Seidl, in Opiz, Natural. n. ix. (1825) 138 ; et in Oek. Tech. Fl. Bohem. i. 48 = aphylla.
filifolia, *Boiss. Voy. Bot. Espagne*, 602. t. 170.—Hispan.
filifolia, Bunge, in *Mém. Sav. Etr. Pétersb.* vii. (1847) 506 = longiscapa.
fimbriata, Vent. Jard. Cels, t. 9 = japonica.
flaccida, Spach, Hist. Vég. Phan. xiii. 41 = versicolor.
flavescens, [*Delile, in*] *Red. Lil.* t. 375.—Bom. ; Reg. Caucas.
flavissima, Besi. Enum. Pl. Volh. 40 = arenaria.
flavissima, *Pall. Reise*, iii. 715.—Sibiria.
fœtucea, Murr. in Nov. Comm. Gotting. vii. (1776) 80. t. 4 = sibirica.
florentina, Kunze, in Flora, xxix. (1846) 698 = albicans.
florentina, *Linn. Syst.* ed. X. 868.—Europ. austr.
florentina, Savi, Mat. Med. Tosc. 40. t. 48 = germanica.
floridana, *Rafin. New Fl. Am.* ii. 92 (Quid ?).—Am. bor.
foetida, Thunb. Diss. Irid. 16 = foetidissima.
foetidissima, *Linn. Sp. Pl.* 39.—Europ. ; Orient.
Fontanesii, *Gode. in Gren. & Godr. Fl. Fr.* iii. 345, adn.—Algeria.
fragrans, Lindl. Bot. Reg. (1840) t. 2 = ensata.
fragrans, Salisb. in Trans. Hort. Soc. i. (1812) 308 = biflora.
frigida, *Bunge, ex Bess. in Flora*, xvii. (1834) i. *Beibl.* 25, nomen.—Sibir. baical.
fugax, *Pers. Syn.* i. 54.—Afr. austr.
fugax, Tenore, Fl. Nap. i. 15 = I. Sisyrinchium.
fulgida, E. Berg, in Flora, xviii. (1835) 563 = pallida.
fulva, *Ker-Gawl. in Bot. Mag.* t. 1496.—Am. bor.
fumosa, *Boiss. & Hausskn. ex Baker, in Journ. Linn. Soc.* xvi. (1877) 134.—Syria.

IRIS :—
furcata, Bieb. Cent. Pl. Ross. ii. t. 51 = aphylla.
Gawleri, [Delar. in] Red. Lil. sub t. 310 = Gueldenstaedtiana.
geminiflora, *Bess. in Flora*, xv. (1832) ii. *Beibl.* 11.—Ross. austr.
georgica, Gueldenst. Reise, ii. 490 = iberica.
germanica, Falk, Beitr ii. 101 = glaucescens.
germanica, Forsk. ex Steud. Nom. ed. II. i. 821 = sumbucina.
germanica, *Linn. Sp. Pl.* 38.—Europ. ; Oriens.
germanica, Sibth. Fl. Graec. t. 40 = pallida.
gigantea, Carr. in Rev. Hortic. (1875) 356 = ochroleuca.
glauca, Salisb. in Trans. Hort. Soc. i. (1812) 302 = pallida.
glaucescens, *Bunge, ex Ledeb. Fl. Alt.* i. 55.—Sibir. alt.
glotiana, Nelder, ex E. Berg. in Flora, xvi. (1833) 17 = Swertii.
glumacea, *Rafin. New Fl. Am.* ii. 98 (Quid ?).—Am. bor.
Gmelini, Ledeb. in Denkschr. Bot. Ges. Regensb. iii. (1841) 46 = lacustris.
goniocarpa, *Baker, in Gard. Chron.* (1876) ii. 710.—Reg. Himal.
gracilipes, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 412.—Japon.
gracilis, E. Berg, in Flora, xviii. (1835) 563 = pumila.
gracilis, Bigel. Fl. Bost. 12 = prismatica.
gracilis, Licht. ex Roem. & Schult. Syst. i. 477 = Moraea tenuis.
gracilis, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 597.—China.
graminea, Georgi, Reise, i. 196 = ruthenica.
graminea, Griseb. Spicil. Fl. Rumel. ii. 370 = Sintenisii.
graminea, Heuff. ex Nym. Consp. 702 = I. Pseudocyperus.
gramines, *Linn. Sp. Pl.* 39.—Europ. austr.
gramines, Thunb. Fl. Jap. 24 = ensata.
graminifolia, Ledeb. Fl. Ross. iv. 106 = geminiflora.
graminifolia, *Pall. Reise Prov. Russ. Reich.* iii. 584, 668 (Quid ?).—Rossia.
grandiflora, Salisb. Prod. 48 = I. Susiana.
Grijii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 515.—China.
Gueldenstaedtiana, *Lepech. in Act. Acad. Petrop.* i. 1781 (1784) 292. t. 8.—Oriens.
Guldenstaedtii, J. F. Gmel. Syst. 115 = pracc.
Guldenstaedtii, Hoffmgg. Verz. Pfl. Nachtr. ii. 141 = pracc.
haematophylla, Fisch. ex Link, Enum. Hort. Berol. i. 60 = ensata.
haematophylla, Fisch. ex Sweet, Brit. Flow. Gard. t. 118 = sibirica.
halophila. Ker-Gawl. in Bot. Mag. t. 575 = spuria.
halophila, Pall. Reise, ii. 728 = Gueldenstaedtiana.
halophylla, Georgi, Beschr. Russ. Reich. III. iv. 671 = pracc.
Hartwegii, *Baker, in Gard. Chron.* (1876) ii. 822. —Calif.
Haynei, *Baker, l. c.* 710.—Palaestin.
Helena, C. Koch, in Linnaea, xxi. (1848) 689 = acutiloba.
Helenae, *Barbey, ex Boiss. Fl. Orient.* v. 182.—Aegypt.
heterophylla, Host. Fl. Austr. ex Spreng. Syst. iv. Cur. Post. 24 = Gueldenstaedtiana ?
hexagona, *Walt. Fl. Carol.* 66.—Am. bor.
Heylandiana, Boiss. & Kent. ex Baker, in Journ. Linn. Soc. xvi. (1877) 143 = iberica.
hirsuta, Licht. ex Roem. & Schult. Syst. i. 478 = Moraea papilionacea.
hirsuta, Hort. ex Steud. Nom. ed. I. 433 = I. Xiphium.
Histrio, *Reichb. f. in Bot. Zeit.* xxx. (1872) 488.—Palaest.
Hoeri, Penny, ex G. Don, in Loud. Hort. Brit. Suppl. i. 591 = tridentata.
hortensis, Tausch, in Flora, xi. (1828) 585 = pallida.
Humea, G. Don, in Loud. Encyc. Fl. Suppl. i. 1160 = germanica.
humilis, Ruoh. *Fl. Taur. Cauc.* i. 33.—Transsylv. ; Reg. Caucas.
humilis, Georgi, Bemerk. Reise Russ. Reich. i. 196 = arenaria.
humilis, Schur, Enum. Pl. Transs. 667 = ruthenica.
humilis, Sieber, ex Baker, in Journ. Linn. Soc. xvi. (1877) 138 = crenocola.

IRIS :—
hungarica, Waldst. & Kit. Pl. Rar. Hung. iii. 251 = aphylla.
hybrida, *Rets. Fasc. Obs. Bot.* (1774) 6.—Hab. †
iberica, *Hofm. in Comm. Soc. Phys. Mosq.* i. (1808) 41. —Orient.
iberica, *Stev. ex Bieb. Fl. Taur. Cauc.* i. 80.—Reg. Caucas.
illyrica, Tomm. ex Vis. Fl. Dalm. Suppl. ii. 53 = germanica.
imberbis, Klatt, in Linnaea, xxxiv. (1865–66) 571, in syn. = junces.
imberbis, Van Hasen, Cat. Arb. et Fl. 67; ex Ker. Gawl. in Bot. Mag. t. 1238 (= *Vieussensia fugax*). —Afr. austr.
imbricata, Lindl. Bot. Reg. (1845) t. 35 = flavescens.
involuta, Garsia, in Effem. Sc. Sicilia (1834), 286 = I. Sisyrinchium.
italica, Parl. Nuov. Gen. e Sp. 87 = I. Chamaeiris.
Itzhatsel, Hasak. Cat. Hort. Bog. Alt. 85 = lacvigata.
japonica, *Thunb. in Transa. Linn. Soc.* ii. (1794) 327. —Japon.
juncea, Brot. Fl. Lusit. i. 51 = filifolia.
juncea, *Pers. Voy. Barb.* ii. 85; *Desf. Fl. Atlant.* i. 39. —Mauritan.
juncifolia, *Pall. Reise Russ. Reich.* i. 94.—Rossia.
Junonia, Schott & Kotschy, ex Schott, in Oestr. Wochenbl. (1854) 309 = aphylla.
Kaempferi, Siebold, ex Lem. Illustr. Hortic. v. (1858) t. 157 = lacvigata.
kamaonensis, *Wall. Cat.* n. 5052.—Reg. Himal.
kashmiriana, *Baker, in Gard. Chron.* (1877) ii. 744.— Reg. Himal.
Kerneriana, [Aschers. & Sint.] ex Baker, in Gard. Chron. (1884) i. 795 = ochroleuca ?
Kolpakowskiana, *Regel, in Act. Hort. Petrop.* v. (1877) 268, 684.—Turkestan.
Korolkowi, *Regel, l. c.* ii. (1873) 432.—Turkestan.
kumaonensis, Auct. = kamaonensis.
lacera, Lam. Encyc. iii. 304 = Moraea polystachya.
lacinista, Hort. Hamb. ex E. Berg, in Flora, xvi. (1833) i. Beibl. 36 = lurida ?
lactea, Pall. Reise, iii. 713 = ensata ?
lacustris, *Nutt. Gen. Am.* i. 23.—Am. bor.
lacvigata, *Pisch. ex Besch. & Mey. Ind. Sem. Hort. Petrop.* v. 36.—Sibir.; Japon.
lamprophylla, Lange, in Bot. Tidsskr. xiii. (1882) 17 = graminea.
latifolia, Gilib. Exercit. ii. 497 = germanica.
Leichtlini, *Regel, in Act. Hort. Petrop.* viii. (1884) 680.—Bokhar.
lepida, Herb. in Flora, xxvii. (1853) 621 = squalens.
leucographa, *Kern. in Oestr. Bot. Zeitschr.* xiii. (1863) 213.—Europ.
Libani, Reut. ex Baker, in Gard. Chron. (1876) i. 359 = I. Histrio.
lilacina, Borb. ex Nym. Consp. 702 = Gueldenstaedtiana.
linbata, Bess. ex Stend. Nom. ed. I. 488 = variegata.
livida, *Booth, ex E. Berg, in Flora*, xvi. (1833) i. *Beibl* 30.—Hab. †
livida, Tratt. Arch. Pict. t. 129 (nigr. 176) = sauana.
longiflora, Herb. ex Baker, in Journ. Linn. Soc. xvi (1877) 144 = glaucescens.
longiflora, Ledeb. Ind. Sem. Hort. Dorpat. (1822) 11 = pumila.
longiflora, Vest, ex Schult. Mant. i. 304 = biflora.
longifolia, Baker, in Journ. Linn. Soc. xvi. (1877) 140, in syn. = I. Pseudacorus.
longifolia, Royle, Illustr. Bot. Himal. 372. t. 91. f. 2 = ensata.
longifolia, Schneev. Ic. t. 20; Andr. Bot. Rep. i. t. 45 = Moraea edulis.
longifolia, Spach, Hist. Veg. Phan. xiii. 14 = tuberosus.
longipetala, *Herb. in Hook. & Arn. Bot. Beech. Voy.* 398.—Am. bor. occ.
longiscapa, *Ledeb. Fl. Ross.* iv. 98.—As. bor.
longispatha, Fisch. ex Sims, Bot. Mag. t. 2628 = ensata.
lorea, Janka, in Termész. Füzetek, i. (1877) 245 = Ioetidissima.
Lortetii, *Barbey, ex Boiss. Fl. Orient.* v. 131.—Syria.
Ludwigi, *Maximo. in Bull. Acad. Pétersb.* xxvi. (1880) 508.—Mont. Altai.
lurida, Ker-Gawl. in Bot. Mag. t. 989 = squalens.
luridu, Reichb. ex Nym. Consp. 700 = germanica, squalens.
lutéola, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 68.— Europ.

IRIS :—
lusitanica, Ker-Gawl. in Bot. Mag. t. 679 = juncea.
lutea, Lam. Fl. Fr. iii. 496 = I. Pseudacorus.
lutescens, [Delaun. in] Red. Lil. t. 268 = I. Chamaeiris.
lutescens, Guss. Cat. Hort. Bocc. (1821) 103 = I. Pseudopumila.
lutescens, *Lam. Encyc* iii. 297,—Europ.: As. Min.
lutescens, Reichb. Ic Crit. x. t. 917 = virescens.
lutescens, Sprun. ex Nym. Consp. 701 = pumila.
Maackii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 542.—Mandchuria.
macrosiphon, *Torr. in Pacif. Rail. Rep.* iv. 144.— Calif.
maculata, *Baker, in Gard. Chron* (1876) ii. 517.— Mesopot.
Mandralisae, Tod. Nuov. Gen. e Sp. 42 = pallida.
mandshurica, *Maxim.* in *Bull. Acad. Pétersb.* xxvi. (1880) 530.—Manchuria.
maricoides, *Regel, in Act. Hort. Petrop.* viii. (1884) 676.—Bokhara.
maritima, Lam. Fl. Fr. iii. 497 = spuria.
maritima, Mill. Gard. Dict. ed. VIII. n. 11 = sibirica.
martinicensis, Jacq. Enum. Pl. Carib. 19 = Trumeia lurida.
Matthioli, *Tausch, in Flora*, xii. (1829) i. *Erg.* 48.— Bohem.
mauritanica, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. 270 = I. 244 = juncea.
meda, *Stapf, in Denkschr. Acad. Wien*, i. (1885) 20. —Persia.
melitta, *Yarsha, in Magyar. Tud. Akad. Math. Term.* xii. (1874) 172.—Thracia.
microptera, Vahl, Enum. ii. 142 = alata.
Milesii, *Baker, ex Foster, in Gard. Chron.* (1883) ii. 231.—Kaschmir. Himal.
minima, *Johann, ex Vest, in Schult. Mant.* i. 313.— Hab. †
minuta, *Franch. & Sav. Enum. Pl. Jap* ii. 42.— Japon.
minuta, Linn. f. Suppl. 98 = Moraea minuta.
missouriensis, *Nutt.* in *Journ. Acad. Philad.* vii. i. (1834) 58.—Am. bor.
Monnieri, [DC. in] Red. Lil. v. t. 236.—Creta; Rhodos.
monophylla, Boiss. & Heldr. in Atti Congr. Bot. Firenz. 1874 (1876) 284 = I. Sisyrinchium.
Moorcroftiana, Wall. Cat. n. 5092 = ensata.
moraeoides, Ker-Gawl. in Bot. Mag. sub t. 1407 = Moraea iridioides.
mutila, *Licht. ex Roem. & Schult. Syst.* i. 477.— Afr. austr.
Myleri, C. Koch, in C. Koch & Fint. Wochenschr. (1859) 206 = virescens.
nana, Pers. Syn. i. 53 = verna.
neglecta ×, Hornem. Hort. Hafn. i. 55.
neglecta, *Parl. Nuov. Gen. e Sp.* 41.—Hab. †
nepalensis, *D. Don, Prod. Fl. Nep.* 54.—Reg. Himal.
nepalensis, Wall. ex D. Don, in Trans. Linn. Soc. xvii. (1841) 310 (sp. dub.).—Nepal.
nepalensis, Wall. in Lindl. Bot. Reg. t. 818 = germanica.
Nertchinskia, Lodd. Bot. Cab. t. 1843 = sibirica.
nihilensis, Lange, in Bot. Tidsak. xiii. (1882) 17 = graminea.
Northiana, Pers. Syn. i. 53 = Marica Northiana.
notha, Bieb. Cent. Pl. Ross. t. 77 = spuria.
nudicaulis, Hook. f. in Bot. Mag. t. 5806 = biflora.
nudicaulis, Lam. Encyc. iii. 296 = aphylla.
ochroleuca, *Ker-Gawl.* in Gard. Gen. 48, ex parte = Monniori.
ochroleuca, *Linn. Mant.* 175.—As. Min.
odorata, Pers. Syn. i. 53 = cristata.
odoratissima, Jacq. Hort. Schoenb. i. t. 9 = pallida.
officinalis, Salisb. Prod. 43 = florentina.
olbiensis, Henon, in Ann. Soc. Agric. Lyon, viii. (1845) 462; ex Gren. & Godr. Fl. Fr. iii. 240 = I. Chamaeiris.
orchioides, *Carr. in Rev. Hortic.* (1880) 337.—Turkestan.
orientalis, Mill. Gard. Dict. ed. VIII. n. 9 = ochroleuca.
orientalis, Thunb. in Trans. Linn. Soc. ii. (1794) 328 = sibirica.
oxypetala, C. A. Mey. in Ind. Pl. Mus. Acad. Petrop. ii. (1834) 8; ex Ledeb. Fl. Ross. iv. 99 = ensata.
palaestina, *Boiss. Fl. Orient.* v. 130.—Syria; Mesopotam.
Pallasii, Fisch. ex Trev. Ind. Sem. Hort. Vrat. (1831) = ensata.

IRIS :—
pallida, *Lam. Encyc.* iii. 294.—Europ. austr.; Syria.
pallida, Sallab. Prod. 44 = Gueldenstaedtiana.
pallida, Tenore, Fl. Nap. iii. 36 = florentina.
pallido-caerulaea. Pers. Syn. i. 51 = pallida.
paludosa, Vers. i. c. 62 = I. Pseudacorus.
palustris, Moench, Meth. 528 = I. Pseudacorus.
pandurata, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 529.—China.
panormitana, Tod. Nuov. Gen. e Sp. 3 = I. Pseudopumila.
papilionacea, Linn. f. Suppl. 98 = Moraea papilionacea.
paradoxa, *Stev. in Mém. Soc. Nat. Mosc.* v. (1814) 355.—Reg. Caucas.
Pavonia, Linn. f. Suppl. 98 = Moraea Pavonia.
Pavonia, Linn. f. Suppl. 98 = Moraea trucsupis.
pedunculata, *Medic. in Act. Acad. Theod. Palat. vi. Phys.* (1790) 417 (Quid ?).—Europ. ?
persica, *Linn. Sp. Pl.* 40.—Cappadoc.; Persia.
picta, Mill. Gard. Dict. ed. VIII. n. 16 = versicolor.
picta, [Spreng.] Bot. Gart. Halle, 50 = Swertii ?
plicata, *Lam. Encyc.* iii. 296.—Hab. ?
plumaria, Thunb. Diss. Irid. 15 = Moraea plumaria.
Polakii, *Stapf, in Denkschr. Acad. Wien*, i. (1885) 20.—Persia.
polystachya, Thunb. Diss. Irid. 25 = Moraea polystachya.
pomeridiana, Fisch. ex Klatt, in Linnaea, xxxiv. (1865–66) 612 = dichotoma.
Potanini, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 528.—Dahuria.
praecox, Sallab. Prod. 45 = persica.
pratensis, Lam. Fl. Fr. iii. 496 = sibirica.
pramatica, *Parsh. Fl. Ag. Sept.* 30.—Am. bor.
Pseudacorus, *Linn. Sp. Pl.* 38.—Europ; As. or.
Pseudo-cyperus, *Schur, Enum. Pl. Transs.* 657.— Hungar.
Pseudo-graminea, Schur, l. c. 928, Index = graminea ?
Pseudo-pulcherrima, E. Berg, in Flora, xviii. (1835) 570 = pumila.
Pseudo-pumila, *Tineo, Cat. Hort. Panorm.* (1827) 388. —Sicil.
Pseudo-sibirica, Schur, Enum. Pl. Transs. 657 = sibirica.
pulchella, *Regel, Gartenfl.* (1859) 310.—Hab. ?
pulcherrima, E. Berg, in Flora, xvi. (1833) i. Beibl. 34 = pumila.
pumila, Bivona, Sic. Pl. ii. 32 = I. Pseudo-pumila.
pumila, Griseb. Spicil. Fl. Rumel. ii. 370, partim = rubromarginata.
pumila, *Linn. Sp. Pl.* 38.—Europ.; As. bor.
pumila, Sibth. & Sm. Fl. Graec. Prod. i. 27 = attica.
pumila, Vill. Hist. Pl. Dauph. ii. 234 = I. Chamaeiris.
punctata, Moench, Meth. 527 = susiana.
purpurascens, E. Berg, in Flora, xviii. (1835) 569 = pumila.
pyrenaica, Bubani, Sched. Crit. 8 = xiphioides.
pyrenaica, *Linn. Gen. Pl.* i. 4; *Amoen. Acad.* iv. 264. —Mont. Pyren.
ramosa, Thunb. Diss. Irid. 18 = Moraea ramosa.
ramosissima, Linn. f. Suppl. 99 = Moraea ramosa.
Redoutéana, Spach, Hist. Veg. Phan. xiii. 56 = lurida.
reflexa, E. Berg, in Flora, xvi. (1833) i. Beibl. 33 = aphylla.
Regeli, *Maxim. ex Regel, in Act. Hort. Petrop.* vi. (1879) 495.—Turkestan.
Reichenbachiana, Klatt, in Linnaea, xxxiv. (1865–66) 613 = spuria.
Reichenbachii, *Heuff. in Oestr. Bot. Zeitschr.* viii. (1858) 28.—Servia.
repanda, E. Berg, in Flora, xvii. (1835) 565 = japonica.
reticulata, *Bieb. Fl. Taur. Cauc.* i. 84.—As. Min.; Persia.
rigida, Sieber, ex Klatt, in Linnaea, xxxiv. (1865–66) 598 = aphylla.
Robinsoniana, F. Muell. Fragm. vii. 153 = Moraea Robinsoniana.
Rosenbachiana, *Regel, in Act. Hort. Petrop.* viii. (1884) 675.—Turkestan.
Rossii, *Baker, in Gard. Chron.* (1877) ii. 809.— Mandchuria.
rubescens, *Rafin. Fl. Ludov.* 20 (Quid ?).—Am. bor.
rubro-marginata, *Baker, in Gard. Chron.* (1875) 524.—Anatolia.
rupestris, Salisb. Prod. 43 = biflora.

IRIS:—
ruthenica, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. i. 117. —Sibir.
ruthenica, Ker-Gawl. Irid. Gen. 54, partim = humilis.
Saarii, Schott, ex Baker, in Gard. Chron. (1876) i. 788 = Sari.
salsa, Fall. Reise, ii. 461 = Gueldenstaedtiana.
Samarstanii, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 234 = I. Sisyrinchium.
sambucina, *Linn. Syst.* ed. X. 863.—Europ. austr.
sanguinea, Donn, Hort. Cant. ed. VI. 17; Hornem. Hort. Hafn. i. 58 = sibirica.
Sari, *Schott, ex Baker, in Gard. Chron.* (1876) 788 (Saarii); *et ex Boiss. Fl. Orient.* v. 131.—Cilicia.
sativa, Mill. Gard. Dict. ed. VIII. n. 15 = versicolor.
scariosa, Willd. ex Link, Jahrb. i. III. 71 = aphylla.
Schmidtii, Hort. ex Baker, in Journ. Linn. Soc. xvi. (1877) 145 = aphylla.
scorpioides, Desf. Fl. Atlant. i. 40. t. 6 = alata.
serbica, Pand. ex Nym. Consp. 701 = virescens
serotina, *Willd. in Willd. & Lange, Prod. Fl. Hisp.* i 141.—Hispan.
setacea. Thunb. Diss. Irid. 20 = Moraea setacea.
setifolia, Linn. f. Suppl. 99 = Moraea setacea.
setosa, *Pall. ex Link, Jahrb.* i. III. 71.—Sibir.
sibirica, Hook. & Arn. Bot. Beech. Voy. 117 = setosa.
sibirica, *Linn. Sp. Pl.* 39.—Europ.; As. bor.
sicula, Tod. Nuov. Gen. e Sp. 7 = pallida.
silvestris, *Vell. Fl. Flum.* 34; i. t. 82.—Bras.
sindjarensis, *Boiss & Haussk. ex Boiss. Fl. Orient.* v. 132.—Mesopot
Sintenisii, *Janka, in Termész. Füzetek*, i. (1877) 244.—As. Min.
Sisyrinchium, *Linn. Sp. Pl.* 40.—Reg. Mediterr.; Oriens; Afghan.
sogdiana, Bunge, in Mém. Sav. Etr Pétersb. vii. (1847) 507 = spuria.
songarica, *Schrenk, Enum. Pl. Nov.* i. 3.—Persia; Afghan.
sordida, Retz. Fasc. Obs. Bot. (1774) 6 = spuria ?
sordida, Soland. ex Salisb. in Trans. Hort. Soc. i. (1812) 308 = juncea.
sordida, *Willd. Enum. Hort. Berol. Suppl.* 4.—Hab. ?
spathacea, Jaume St. Hil. '19. t. 27'; ex Roem. & Schult. Syst. i. 465 = spuria.
spathacea, Thunb. Diss. Irid. 17 = Moraea spathacea.
spathulata, Cav. Desc. 298, in obs. = sibirica.
spathulata, Lam. Encyc. ii. 300 = spuria.
spathulata, Linn. f. Suppl. 99 = Moraea spathacea.
spathulata, Willd. Enum. Hort. Berol. Suppl. 4 = Gueldenstaedtiana.
spectabilis, Salisb. Prod. 43 = germanica.
spectabilis, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 98 = I. Xiphium.
specularis, *Hance, in Journ. Bot.* xiii. (1875) 196.—Hongkong.
spuria, Bertol. Fl. Ital. ii. 776 = foetidissima.
spuria, *Linn. Sp. Pl.* 39.—Europ.
spuria, Fall. Reise, iii. 713 = ensata.
spuria, Reichb. Ic. Pl. Crit. x. & f. 1235 = Gueldenstaedtiana.
squalens, E. Berg, in Flora, xvi. (1833) i. Beibl. 14 = variegata.
squalens, Georgi, Beschr. Russ. Reich. iv. 668 = germanica.
squalens, *Linn. Syst.* ed. X. 863.—Europ.; Caucas.; Persia.
squalens, Petagna, Inst. Bot. ii. 88 = sambucina.
squalens, Thunb. Fl. Jap. 33 = japonica.
Statellae, Tod. Nuov. Gen. e Sp. 5 = lutescens.
stenogyna, [Delar. in] Red. Lil. vi. t. 310 = Gueldenstaedtiana.
stenoloba, A. DC. ex Baker, in Journ. Linn. Soc. xvi. (1877) 148 = pumila.
Stocksii, *Boiss. Fl. Orient.* v. 133.—Afghan.
stolonifera, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 535.—Kokan.
stricta, Moench, Meth. 538 = sibirica.
stylosa, Desf. Fl. Atlant. i. 40. t. 5 = unguicularis.
suaveolens, *Boiss. & Reut. Diagn.* Ser. I. xiii. 15.—Orient.
suavis, Salisb. Prod. 44 = graminea.
subbarbata, Joo, in Verh. Siebenb. Ver. Naturw. ii. (1851) 98 = Gueldenstaedtiana.
subbiflora, Brot. Fl. Lusit. i. 50. t. 26 = biflora.
subbiflora, Fleber, ex Klatt, in Linnaea, xxxiv. (1865–66) 599 = aphylla.
sulcata, Wall. Cat. n. 5049 = nepalensis.

IRIS:—
sulphurea, C. Koch, in Linnaea, xxi. (1848) 687 = virescens.
superba, E. Berg, in Flora, xviii. (1835) 567 = squalens.
susiana, *Linn. Sp. Pl.* 38.—Persia.
Swertii, *Hort. ex Lam. Encyc.* iii. 296.—Hab. ?
sylvatica, Balb. ex Roem. & Schult. Syst. i. 476 = graminea.
syrtica, Viv. Fl. Lib. Spec. 3. t. 1. f. 1 = I. Sisyrinchium.
tangerica, *Hort. Ch^s. ex G. Don, in Loud. Hort. Brit.* 19 (Quid ?).—Marocc.
tardiflora, E. Berg, in Flora, xviii. (1835) 566 = squalens.
taurica, *Lodd. Bot. Cab.* t. 1506.—Tauria.
tectorum, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 380.—Japon.
tenax, *Dougl. in Bot. Reg.* t. 1218.—Am. bor. occ.
Tenoreana, Nym Consp. 962, Index = seq.
Tenorii, *Parl. Fl. Ital.* iii. 278.—Europ.
tenuifolia, *Pall. Reise*, iii. 714.—Rossia.
tenuis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 380.—Am. bor. occ
Tigridia, *Bunge, in Ledeb. Fl. Alt.* i. 60.—Mont. Altai.
tigrina, Jacquem. ex Baker, in Journ. Linn. Soc. xvi. (1877) 144 = kamaonensis.
Timaei, Tod. Nuov. Gen. e Sp. 46 = pallida.
tingitana, *Boiss. & Reut. Pugill. Pl. Nov.* 118 = Mauritania.
Tolmiana, Herb. in Hook. & Arn. Bot. Beech. Voy. 396 = missouriensis.
tomiolopha, Hance, in Journ. Bot. x. (1872) 229 = tectorum.
torta, *Tausch, in Flora*, xvii. (1834) 523.—Hab. ?
transilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 78 = pumila.
trantaguna, Brot. Fl. Lusit. i. 52 = alata.
tralata, Bot. Phyt. ii. 44. t. 95 = alata.
tricuspidata, Linn. f. Suppl. 98 = Moraea tricuspis ?
tricuspis, Thunb. Diss. Irid. 14 = Moraea edulis.
tricuspis, Willd. Sp. Pl. i. 281 = Moraea tricuspis.
tridentata, *Pursh, Fl. Am. Sept.* i. 30.—Am. bor.
tridentata, Sweet, Brit. Flow. Gard. t. 274 = tridentata.
trifoca, Balb. Misc. Bot. 6. t. 1 = ensata.
trigonocarpa, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1853) 17 = sibirica.
tripetala, Hook. Bot. Mag. t. 2886 = tridentata.
tripetala, Linn. f. Suppl. 92 = Moraea tripetala.
tripetala, *Walt Fl. Carol.* 66.—Am. bor.
tristis, Linn. f. Suppl. 97 = Moraea tristis.
tristis, Reichb. Ic. Fl. Germ. xix. 3 = pumila.
tuberosa, *Linn. Sp. Pl.* 40.—Europ. austr.
unguicularis, *Poir. Voy. Barb.* ii. 86.—Mauritan.
unifora, *Pall. ex Link, Jahrb.* i. III. 71.—Sibir.
unifolia, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 173 (Quid ?).—Afr. austr.
variabilis, Jacq. Coll. ii. 321 = I. Xiphium.
variegata, *Linn. Sp. Pl.* 38.—Europ.
Vartani, *Foster, in Gard. Chron.* (1885) i. 488.—Syria.
venticosa, *Pall. Reise*, iii. 712.—Dahuria.
venusta, Booth, ex E. Berg, in Flora, xvi. (1833) i. Beibl. 81 = lurida ?
verna, *Linn. Sp. Pl.* 39.—Am. bor.
verna, Pall. Reise, iii. 213 = ruthenica.
vernalis, Salisb. Prod. 44 = verna.
versicolor, *Linn. Sp. Pl.* 39.—Am. bor.
versicolor, Thunb. Fl. Jap. 34 = laevigata.
villosa, Ker-Gawl. in Bot. Mag. t. 571 = Moraea Pavonia.
violacea, Klatt, in Linnaea, xxxv. (1867–68) 884 = laevigata.
violacea, Savi, Bot. Etrusc. ii. 9 = germanica.
violacea, *Vahl, Enum. Hort. Brit.* ed. I. 238.—Europ. austr.
virescens, [*Delar. in*] *Red. Lil.* t. 295.—Europ.
virgata, *Raf. New Fl. Am.* ii. 92 (Quid ?).—Am. bor.
virginica, A. Gray, Man. Bot. N. U. St. ed. V. 516 = prismatica.
virginica, *Linn. Sp. Pl.* 39.—Am. bor.
virginica, Michx. Fl. Bor. Am. i. 22, partim = hexagona.
viscaria, Linn. f. Suppl. 98 = Moraea viscaria.
vulgaris, Pohl, Tent. Fl. Boh.; ex Schult. Mant. i. 204 = germanica.
Winkleri, *Regel, in Act. Hort. Petrop.* viii. (1884) 677.—Turkestan.

IRIS:—
xiphia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = I. Xiphium.
xiphioides, *Ehrh. Beitr.* viii. 140.—Europ. austr.
Xiphium, Desf. Fl. Atlant. i. 37 = Fontanesii.
Xiphium, Georgi, Reise, i. 196 (Quid ?).—Rossia.
Xiphium, Jacq. Coll. ii. 320 = xiphioides.
Xiphium, *Linn. Sp. Pl.* 40.—Hispan.
yedoensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 522.—Japon.
zelanica, Vigo, ex Parl Nuov. Gen. e Spec. 58 = I. Sisyrinchium.

IRIUM, Steud. Nom ed. I 440 (1821) = IRION, Burm. f. = **Boridula**, Linn. (Droserac.).

IRLBACHIA, Mart. Nov. Gen. et Sp. ii. 101. t. 179 (1827) = **Lisianthus**, Aubl. (Gentian.).
Bonplandiana, Fenzl, Nov. Stirp. Dec. (1839) 12; ex Griseb. in DC. Prod. ix. 72.—N. Granat.
cornuicensis, Griseb. Gen. et Sp. Gentian. 195 = L. corniculensis.
elegans, Mart. Nov. Gen. et Sp. ii. 102. t. 179 = L. paniculatus.
recurva, Prog. in Mart. Fl. Bras. vi. i. 229 = L. recurvus.
subcordata, Prog. l. c. = L. subcordatus.

IRMA, Bouton, ex A. DC. Prod. xv. i. 406 (1864) = **Begonia**, Tourn.

IRMISCHIA, Schlecht in Linnaea, xix. (1847) 738 = **Metastelma**, R. Br. (Asclep.).
floribunda, Schlecht l. c. 739.—Mexic.

IRON, P. Br. Hist. Jamaic. 179. t. 12. f. 3 (1765) = **Sauvagesia**, Linn. (Violar.).

IROUCANA, Aubl. Pl. Gui. i. 328. t. 127 (1775) = **Cassaria**, Jacq. (Samydac.).
guianensis, Aubl. l. c. 329. t. 127 = C. ramiflora.

IRSIOLA, P. Br. Hist. Jamaic. 147. t. 4. f. 2 (1756) = **Vitis**, Tourn. (Ampelid.).
sicyoides, Raf. Sylva Tellur. 86 = V. sicyoides.

IRULIA, Bedd. Forest. Man. Bot. 235 (1873) = **Melocanna**, Trin. (Gramin.).
travancorica, Bedd. l. c. = M. travancorica.

IRVINGIA, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 167. *SIMARUBEAE*, Benth. & Hook. f. i. 314.
Barteri, *Hook. f. l. c.*—Afr. trop.
malayana, *Oliver, ex A. Benn. in Hook. f. Fl. Brit. Ind.* i. 532.—Malacca.
Smithii, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 167.—Afr. trop.
tenuifolia, *Hook. f. l. c.*—Afr. trop.

IRVINGIA, F. Muell. Fragm. v. 17 (1865) = **Hedera**, Tourn. (Araliac.).
australiana, F. Muell. l. c. 18 = H. australiana.

ISABELIA, Rodrig. Orch. Nov. i. 75 (1877). *ORCHIDEAE.*
virginalis, *Rodrig. l. c.* 76.—Bras.

ISACANTHUS, Nees, in DC. Prod. xi. 276 (1847) = **Volkameria**, Linn.
Vogelii, Nees, l. c. 279 = S. Vogelii.

ISACHNE, R. Br. Prod. 196 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1100.
GRAYA, Nees, ex Steud. Syn. Pl. Gram. 119 (1854). SPHAEROCARYUM, Nees, ex Steud. Nom. ed. II. ii. 630 (1841).
adstans, *Miq. Fl. Ind. Bat.* iii. 461.—Malacca.
albens, *Trin. Ic. Gram.* t. 65.—Ind. or.; Java.
altissima, *Deboeuf, in Act. Linn. Soc. Bord.* xxx. (1875) 122.—China.
angusta, Nees, ex Steud. Syn. Pl. Gram. 96 = Panicum stenanthum.
arundinacea, Griseb. Fl. Brit. W. Ind. 558 = paniculata.
atrovirens, Trin. Gram. Panic. 251 = australis.
australis, *R. Br. Prod.* i. 196.—Austral.
dispar, *Trin. Ic. Gram.* t. 72.—Reg. Himal.

ISACHNE:—
disperma, Doell, in Mart. Fl. Bras. ii. 11. 274 = panicum.
dubia, Kunth, Rév. Gram. i. 42; Enum. Pl. i. 137 = panicea.
fimula, *Busse, in. Mig. Pl. Jungh.* 879.—Java.
Gardneri, *Benth. & Hook. f. Gen.* iii. 1100.—Zeylan.
geniculata, *Griff. Notul.* iii. 41.—Ind. or.
hispidula, Nees, ex Steud. Syn. Pl. Gram. 96 = Coelachne pulchella.
javana, *Nees, ex Steud. l. c.* in syn.—Java.
Kunthiana, *Nees, ex Steud. l. c.* 96.—Java.
Lamarckii, Kunth, Rév. Gram. i. 42 = mauritiana.
lersioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 533.—Cuba.
lipidota, Walp. Ann. i. 934 = australis.
mauritiana, *Kunth, Rév. Gram.* i. 343. t. 33.—Ins. Mascar.
Menoritana, R. Br. Prod. 196.= australis.
miliacea, *Roth, Nov. Pl. Sp.* 56.—Ind. or.
minutula, *Kunth, Rév. Gram.* ii. t. 117.—Ins. Mariscae.
monticola, *Busse, in Mig. Pl. Jungh.* 879.—Java.
muricata, Nees, ex Steud. Nom. ed. II. i. 323 = Panicum muricatum.
Myosotis, *Nees, in Hook. Kew Journ.* ii. (1850) 98.—Ins. Philipp.
Neesiana, Arn. ex Steud. Syn. Pl. Gram. 96 = Kunthiana.
nilagirica, Hochst. ex Benth. & Hook. f. Gen. iii. 1100 = Gardneri.
pangerangensis, *Zoll. & Mor. Syst. Verz. Zoll.* 102.—Java.
panicea, *Trin. Gram. Panic.* 253.—Am. trop.
perpusilla, Arn. ex Steud. Syn. Pl. Gram. 96 = Panicum perpusillum.
polygonoides, Doell, in Mart. Fl. Bras. ii. 11. 273 = Panicum polygonoides.
prostrata, Hort. Berol. ex Steud. Syn. Pl. Gram. 95 = Panicum horticolum.
pulchella, *Roth, Nov. Pl. Sp.* 58.—Ind. or.
pygmaea, *Griseb. Fl. Brit. W. Ind.* 553.—Ind. occ.
retracta, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 227. —Afr. occ.
rigens, *Trin. Gram. Panic.* 252.—Ind. occ.
rigida, *Nees, ex Steud. Syn. Pl. Gram.* 95.—Java.
simplicissecula, Wight & Arn. ex Thw. Enum. Pl. Zeyl. 378, in syn. = Coelachne brachiata.
stigmatosa, *Griff. Notul.* iii. 42.—Ind. or.
Streptostachys, Nees, ex Steud. Syn. Pl. Gram. 79 = Panicum Streptostachys.
trachyspermа, *Nees, ex Seem. Bot. Voy. Herald*, 224. —Panama.
tricrinata, *Roth, Nov. Pl. Sp.* 57.—Ind. or.
ventricosa, *Doell, in Mart. Fl. Bras.* ii. 11. 274.—Am. trop.
virgata, *Nees, ex Steud. Syn. Pl. Gram.* 96, in syn.—Ind. or.: Malaya.
Walkeri, *Arn. ex Steud. l. c.* 97.—Zeylan.

ISANDRA, F. Muell. Census, 140 (1882), nomen. *SOLANACEAE.*
Bancroftii, *F. Muell. l. c.*; *et in Wing, South. Sc. Record*, iii. (1883) 2.—Austral.

ISANDRA. Salisb. Gen. Pl. Fragm. 67 (1866) = **Thysanotus**, R. Br. (Liliac.).

ISANDRINA, Rafin. Sylva Tellur. 136 (1838) = **Cassia**, Tourn. (Legumin.).
arborescens, Rafin. l. c. = C. emarginata.

ISANTHERA, Nees, in Trans. Linn. Soc. xvii. (1834) 83. *GESNERACEAE*, Benth. & Hook. f. ii. 1016.
cronata, C. B. Clarke, in DC. Monog. Phan. v. 193 = discolor.
discolor, *Meisner. in Bull. Acad. Pétersb.* xix. (1874) 533.—Ins. Formosa.
exinia, *C. B. Clarke, in DC. Monog. Phan.* v. 193.—Java.
floribunda, Garda. in Calc. Journ. Nat. Hist. vi. (1846) 433 = parmollis.
parmollis, *Mart, in Trans. Linn. Soc.* xvii. (1834) 83. —Ind. or.

ISANTHINA, Reichb. ex Steud. Nom. ed. II. i. 400 (1840) = **Cummolinia**, Linn.

ISANTHUS, DC. Prod. vii. 63, in syn. (1838) = **Perasia**, Lag. (Compos.).

ISANTHUS, Michx. Fl. Bor. Am. ii. 3. t. 80 (1803). *LABIATAE*, Benth. & Hook. f. ii. 1220.
coeruleus, *Michx. Fl. Bor. Am.* ii. 4. t. 80.—Am. bor.
multiflorus, Rafin. New Fl. Am. ii. 35 = coeruleus.
pubescens, Rafin. l. c. = coeruleus.
pumilus, Rafin. l. c. 34 = coeruleus.

ISARTIA, Dun. Anal. Fam. 32 (1829) = **Isertia**, Schreb. (Rubiac.).

ISATIS, Tourn. ex Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 94.
CHAETOLOMA, Bunge, in Bot. Zeit. ii. (1844) 249.
GLASTUM, Rupp. Fl. Jen. ed. Hall 68 (1745).
SAMERARIA, Desv. Journ. Bot. iii. (1814) 161.
aegyptiaca, Crantz, Inst. ii. 94 = acq.
aegyptica, Linn. Sp. Pl. 671 = Cakile maritima.
agyptia, Forsk. Fl. Aegypt. Arab. 121 = praec.
alepplca, *Scop. Delic. Insub.* ii. 31. t. 16.—As. Min.; Syria.
alpina, *All. Fl. Pedem.* i. 259.—Italia.
alpina, Vill. Prosp. 38; Hist. Pl. Dauph. iii. 808 = lasiocarpa.
alyssifolia, *Boiss. Fl. Orient.* i. 378.—As. Min.
apiculata, Boiss. in Ann. Sc. Nat. Sér. II xvii. (1842) 199 = latisiliqua.
apiculata, Jord. ex Nym. Consp. 67 = tinctoria.
araratica, *Rupr. Fl. Cauc.* 182, in textu.—Armen.
armena, Bieb. Fl. Taur. Cauc ii. 91 = latisiliqua.
armena, *Ann. Sc. Fl.* 870.—Oriens.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 202.—As. Min.
baeuatica, Hort. Cremen. 74; ex Link, Enum. Hort. Berol. ii. 149 = tinctoria.
Besseri, *Trautv. in Mém. Sav. Etr. Pétersb.* iv. (1845) 813. t. 11.—As. Min.; Persia.
biscutellefolia, *Sieber & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 28.—Persia.
Boissieriana, *Reichb. f. in Journ. Bot.* xiv. (1876) 46. —Samarkand.
brachycarpa, *C. A. Mey. in Bichw. Casp. Cauc.* 17.—Armen.
Bungeana, *Seidl. Bot. Reise, Transkauk.* 87, 83. —Armenia.
callifera, *Boiss. & Bal. Diagn.* Ser. II. v. 46.—Cilicia.
campestris, Stev. in DC. Syst. ii. 571 = tinctoria.
campyloearpa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 208.—Persia.
canescens, DC. Fl. Fr. Suppl. 598 = tinctoria.
cappadocica, Desv. Journ. Bot. iii. (1814) 174 = latisiliqua.
caucasica, Kolenati, ex Rupr. Fl. Cauc. 182 = reticulata.
cochlearis, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 202.—Syria.
cordigera, Boiss. l. c. 198 = Besseri.
corymbosa, *Boiss. l. c.* 200.—As. Min.
costata, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 204.—Reg. Himal. et Altai
dalmatica, Mill. Gard. Dict. ed. VIII. n. 2 = tinctoria.
dasycarpa, Ledeb. Ind. Sem. Hort. Dorpat. (1834) 4 = praecox.
denticata, Pers. Syn. ii. 193 = alepplca.
Djerjaeana, *Cosa. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 628.—Algeria.
emarginata, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 134.—Soongaria.
frigida, *Boiss. & Ketzsky, Diagn.* Ser. II. ii. 547.—As. Min.
frutescens, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xiv. (1841) 356.—Soongaria.
Garcini, DC. Syst. ii. 565 = Fortunia Garcini.
glauca, *Auch. ex Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 201.—As. Min.; Persia.
glauca, Gilib. Fl. Lituan. ii. 66 = tinctoria.
glauca, Willd. ex DC. Prod. i. 210 = lusitanica.
heterocarpa, DC. Syst. ii. 569 = littoralis.
heterocarpa, C. A. May. in Ledeb. Fl. Alt. iii. 205 = lasiocarpa.
heterocarpa, Andrz. ex Bess. Enum. Pl. Volh. 82 = tinctoria.

ISATIS:—
heterocarpa, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 11.—Turkestan.
hirsuta, Pers. Syn. ii. 193; Link, Handb. ii. 384 = tinctoria.
hirtocalyx, *Fenzl. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 234.—Turkestan.
iberica, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 267. —Reg. Caucas.
icosia, *Boiss. & Heldr. Diagn.* Ser. I. viii. 48.—As. Min.
indigotica, *Fortune, ex Lindl. in Journ. Hort. Soc.* i. (1846) 276.—China.
intermedia, Turcz. ex Ledeb. Fl. Ross. i. 212 = oblongata.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 75.—Japon.
Kotschyana, *Boiss. & Hohen. Diagn.* Ser. I. viii. 48.—Persia.
laevigata, *Trautv. Del. Sem. Hort. Kiov.* (1840) 6.—Sibir.
lasiocarpa, *Fisch. ex Regel, Pl. Nov. Fedsch.* 10.—Turkestan.
lasiocarpa, Ledeb. Fl. Ross. i. 211 = tinctoria.
latisiliqua, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 267.—As. Min.; Persia.
leiocarpa, DC. Syst. ii. 567 = latisiliqua.
leuconeura, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 28.—Reg. Caucas.
littoralis, Bess. Enum. Pl. Volh. 71 = praecox.
littoralis, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 267.—Tauria.
Lockmannians, *Kotschy, ex Boiss. Fl. Orient.* i. 381.—Cilicia.
Lunaria, Crants, ex DC. Syst. ii. 285 = Ricotia Lunaria.
lusitanica, *Brot. Fl. Lusit.* i. 360.—Lusitan.
lusitanica, S. G. Gmel. It. iii. 806 = orientalis.
lusitanica, Linn. Sp. Pl. 670 = alepplca.
lusitanica, Moris, Fl. Sard. i. 114 = tinctoria.
maeotica, DC. Syst. ii. 571 = tinctoria.
maritima, Rupr. Fl. Cauc. 183, in textu = tinctoria.
meyacarpa, Fisch. ex Steud. Nom. ed. II. i. 624 = tinctoria.
membrances, *Trautv. in Mém. Sav. Etr. Pétersb.* iv. (1845) 316.—Hab.?
microcarpa, *Y. Gay, ex Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 201.—Arab.
minima, Bunge, Del. Sem. Hort. Dorpat. (1843) 7 = iberica.
minor, Moench, Meth. 292 = alepplca.
nummularia, Trautv. in Mém. Sav. Etr. Pétersb. iv. (1845) 312 = latisiliqua.
oblongata, *DC. Syst.* ii. 571.—Sibir.
orientalis, Willd. Hort. Berol. ii. 663 = alepplca.
pilosa, Schur, Enum. Pl. Transs. 73 = tinctoria.
pinnata, *Forsk. Fl. Aegypt. Arab.* 121.—Arab.
platycarpa, *Bunge, Del. Sem. Hort. Dorpat.* (1843) 7. —Persia.
platycarpa, Jaub. & Spach, Illustr. iii. 31. t. 23 = latisiliqua.
platyloba, Link, ex Steud. Nom. ed. I. 440 = lusitanica.
praecox, *Kit. ex Tratt. Arch.* ii. 40. t. 68; *DC. Syst.* ii. 568.—Europ.
psilocarpa, Ledeb. Fl. Ross. i. 210 = latisiliqua.
raphanifolia, *Boiss. Diagn.* Ser. I. vi. 19.—Persia.
reticulata, *C. A. Mey. in Beitr. Pfl. Russ. Reich.* vi. 53.—Reg. Caucas.
rostellata, Bertol. Fl. Ital. vi. 515 = tinctoria.
rugulosa, *Bunge, ex Boiss. Fl. Orient.* i. 384.—Persia.
sabulosa, *Stev. ex Ledeb. Fl. Ross.* i. 213.—Transcauc. *sibirica*, Trautv. in Mém. Sav. Etr. Pétersb. iv. (1845) 305 = lasiocarpa.
songarica, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 309 = iberica.
spathulata, Edgew. ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 179 = Dipterygium glaucum.
Steveniana, Trautv. in Mém. Sav. Etr. Pétersb. iv. (1845) 306 = psilocarpa.
Stockati, *Boiss. Fl. Orient.* i. 376.—Beluchist.
suberos, *Czern. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 310.—Bessarab.
subnudula, *Rupr. Fl. Cauc.* 181, in textu.—Reg. Caucas.
suffruticescens, Boiss. in Kotschy, ex Boiss. Fl. Orient. i. 379 = iberica.

ISATIS:—
taurica, Bieb. Fl. Taur. Cauc. iii. 423 = tinctoria.
tinctoria, *Linn. Sp. Pl.* 670.—Europ.; Orient.
tomentella, Boiss. & Bal. Diagn. Sér. II. v. 46 = tinctoria.
trachycarpa, *Trautv. in Act. Hort. Petrop.* ix. (1886) 870.—Turkest.
undulata, *Auch. ex Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 200.—As. Min.
velleroïdes, Boiss. & Noé, Diagn. Ser. II. i. 46 = Aucheri.
velutina, *Boiss. & Huet, i. c.* v. 45.—As. Min.
Villarsii, *Gaud. Syn. Fl. Helv.* 296.—Helvet.
violascens, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 166.—Turkest.; Persia.
virens, Cosa. in Bull. Soc. Bot. Fr. xxii. (1875) 54, nomer = tinctoria.

ISAURA, Comm. ex Poir. Encyc. Suppl. iii. 185 (1813) = **Stephanotis**, Thou. (Asclep.).
alliacea, Stend. Nom. ed. II. i. 824 = S. laxa.
allicia, Comm. ex Poir. Encyc. Suppl. iii. 185 = S. floribunda.

ISAUXIS, Reichb. Nom. 210 (1841) = **Vatica**, Linn. (Dipterocarp.).
cordifolia, Hook. f. Fl. Brit. Ind. i. 722 = V. cordifolia.
Roxburghiana, Thw. Enum. Pl. Zeyl. 37 = V. affinis.

ISCHAEMOPOGON, Griseb. Fl. Brit. W. Ind. 560 (1864) = **Ischaemum**, Linn. (Gramin.).
arcuatus, Griseb. l. c. (Spodiopogonis sp.).—Ind. occ.
latifolius, Griseb. l. c. = Ischaemum latifolium.

ISCHAEMUM, Linn. Gen. ed. II. 525 (1742). *GRAMINEAE*, Benth. & Hook. f. iii. 1192.
COLLADOA, Cav. Ic. v. 37. t. 460 (1799).
HOLOSARUM, Nees, in Edinb. N. Phil. Journ. xviii. (Jan. 1835) 185.
ISCHAEMOPOGON, Griseb. Fl. Brit. W. Ind. 560 (1864).
MEOSCHIUM, Beauv. Agrost. iii. t. 21 (1812).
SCHOENANTHUS, Adans. Fam. ii. 38 (1763).
SKHIMA, Forsk. Fl. Aegypt. Arab. 178 (1775).
acuminatum, Herb. Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 54 = Crypsis aculeata.
antephoroides, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 102.—Japon.
aristatum, *Linn. Sp. Pl.* 1049.—Ind. or.; China.
aristatum, Thunb. Prod. Pl. Cap. 21 = filiforme.
aristatum, Willd. Sp. Pl. iv. 939; Roxb. Hort. Beng. 8; Fl. Ind. i. 319 = ciliare.
arundinaceum, *F. Muell. ex Benth. Fl. Austral.* vii. 519.—Austral.
auritum, Nees, ex Steud. Syn. Pl. Gram. 361 = Rottboellia aurita.
australe, *R. Br. Prod.* 205.—Austral.
barbatum, Baker, Fl. Maurit. 447 = fasciculatum.
barbatum, Miq. in Ann. Mus. Bot. Lugd. Bat. iii. 198 = antephoroides.
barbatum, Retz. Obs. vi. 35 = aristatum.
binatum, *Bunge, ex De Vriese, Pl. Ind. Bat. Reinw.* 108.—Ins. Solor.
ciliare, *Retz, Obs.* vi. 36.—As. et Austral. trop.
Colladoa, Spreng. Syst. i. 298 = rugosum.
commutatum, *Hack. ex Trimen, Cat. Pl. Ceyl.* 107.—Zeyl.
conjugatum, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 321.—Ind. or.
corollatum, *Nees, ex Steud. Syn. Pl. Gram.* 369.—Reg. Himal.
crinitum, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 298 = Pogonatherum saccharoideum.
cuspidatum, Roxb. Hort. Beng. 8; Fl. Ind. i. 324 = Vossia procera.
decumbens, *Benth. Fl. Austral.* vii. 521.—Austral.
digitatum, *Brongn. in Duperr. Voy. Coq. Bot.* 70. t. 1a.—Ins. Molucc.
falcatum, *Nees, ex Steud. Syn. Pl. Gram.* 369.—China.
fasciculatum, *Brongn. in Duperr. Voy. Coq. Bot.* 73.—Ins. Maurit.
filiforme, *Willd Sp. Pl.* iv. 940.—Afr. austr.
fragile, *R. Br. Prod.* 205.—Austral.
graniculatum, Roxb. Hort. Beng. 8; Fl. Ind. i. 322 = aristatum.
gibbum, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 295 = aristatum.

ISCHAEMUM:—
glabratum, J. & C. Presl, Rel. Haenk. i. 338 = muticum.
glabrum, Walt. Fl. Carol. 249 = Tripsacum dactyloides.
guianense, *Kunth, Rév. Gram.* i. 168.—Guian.
hirsutum, Nees, ex Steud. Syn. Pl. Gram. 361 = Rottboellia hirsuta.
hirsutum, *Spreng. Syst.* i. 298.—Bras.
hirtiscabes, *Munro, ex Hitch. in Journ. Linn. Soc.* xviii. (1880) 111, nomen.—Afghan.
hispidum, H. B. & K. Nov. Gen. et Sp. i. 194 = Arundinella brasiliensis.
imberbe, Retz. Obs. vi. 35 = aristatum.
importunum, *Lour. Fl. Cochinch* 646.—Cochinch.
inscalptum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 472 = laxum.
intermedium, *Brongn. in Duperr. Voy. Coq. Bot.* 73. —Ins. Carolin.
involutum, Forst. f. Prod. 73 = Thuarea involuta.
involutum, Labill. ex Kunth, Enum. Pl. i. 512 = muticum.
Kotschyi, Hochst. in Flora, xxvii. (1844) 247 = laxum.
latifolium, *Kunth, Rév. Gram.* i. 166.—Am. trop.
latifolium, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 291 = Rottboellia latifolia.
laxum, *R. Br. Prod.* 205.—Geront. trop.
lecrioides, *Munro, in Proc. Am. Acad.* iv. (1864–65) 368.—China; Calif.
macrostachyum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 472 = laxum.
mastruccatum, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 298 = Rottboellia hirsuta.
meliocides, Koen. ex Willd. Sp. Pl. iv. 941 = Enteropogon meliocides.
Meyenianum, Steud. Nom. ed. II. i. 824 = aristatum.
minus, J. & C. Presl, Rel. Haenk. i. 339 = Urvillea laxa.
murinum, *Forst. f. Prod.* 73.—Ind. or.; Ins. Tanna.
muticum, *Linn. Sp. Pl.* 1049.—Ind. or.; Austral.; Ins. Pacif.
Myurus, Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 54 = Manisuris Myurus.
nervosum, Thw. Enum. Pl. Zeyl. 365 = laxum.
ophioides, *Munro, in Proc. Am. Acad.* iv. (1864) 368. —China.
palaceum, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 293 = Apocopis Royleana.
pallidum, Kunth, Enum. Pl. i. 515 = Arundinella pallida.
panicrum, Chauv. ex Steud. Syn. Pl. Gram. 375 = Isachne australis.
pectinatum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 296.—Ind. or.; Austral.
perveniens, Kunth, Enum. Pl. i. 515 (Arundinellae sp.). —Peruv.
pilosum, *Trimen, Cat. Pl. Ceyl.* 107.—Zeylan.
polystachyum, J. & C. Presl, Rel. Haenk. i. 338 = digitatum.
repens, Roxb. Hort. Beng. 83; Fl. Ind. i. 322 = muticum.
rottboellioides, R. Br. Prod. 206 = Rottboellia ophiuroides.
Royleanum, Miq. Fl. Ind. Bat. iii. 498 = rugosum.
rugosum, *Salisb. ic. Stirp. Rar.* t. 1.—As. trop.
rugosum, Trin. in Spreng. Neu. Entdeck. ii. 79; et in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 294 = segetum.
scaberrimum, Walt. Fl. Carol. 249 (Planta dubia).—Am. bor.
scrobiculatum, Wight & Arn. ex Steud. Syn. Pl. Gram. 378 = ciliare.
secundatum, Walt. Fl. Carol. 249 = Stenotaphrum glabrum.
segetum, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 294 = rugosum.
Sekima, Spreng. Syst. i. 298 = laxum.
semisagittatum, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 320. —Ind. or.
Sieboldii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 291.—China; Japon.
speciosum, Nees, ex Steud. Syn. Pl. Gram. 375 (Rottboellinae sp.).—Ind. or.
tenellum, Roxb. Hort. Beng. 83; Fl. Ind. i. 323 = ciliare.
timorense, Kunth, Rév. Gram. i. 169. t. 26 = Spodiopogon Dyrenis.
trilicoum, *R. Br. Prod.* 205.—Austral.

ISCHAEMUM:—
truncatiglume, *F. Muell. ex Benth. Fl. Austral.* vii. 518.—Austral.
Urvillaeanum, *Kunth, Rév. Gram.* i. 167.—Bras.
villosum, R. Br. Prod. 205 = australe.
vrylanicum, *Hack. ex Trimen, Cat. Pl. Ceyl.* 107.—Zeylan.
Zollingeri, Miq. Fl. Ind. Bat. iii. 499 = Andropogon Zollingeri.

ISCHALEON, Ehrh. Beitr. iv. 147 (1789) = **Microcala**, Hoffmgg. & Link (Gentian.).

ISCHARUM, Blume, Rumphia, i. 144. t. 29 (1885) = **Blarum**, Schott (Aroid.).
alexandrinum, Schott, Syn. Aroid. 8 = B. Olivieri.
angustatum, Hook. f. Bot. Mag. t. 6355 = B. angustatum.
Bovei, Schott, Syn. Aroid. 7 = B. Bovei.
Carsaani, Schott, Prod. Aroid. 67 = B. Bovei.
crispulum, Schott, l. c. 68 = B. Bovei.
disparum, Schott, Syn. Aroid. i. 7 = B. Bovei.
eximium, Schott & Kotschy, in Oestr. Bot. Wochenbl. (1854) 81 = B. eximium.
Frasianum, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 98 = B. Frasianum.
Haenseleri, Schott, Syn. Aroid. 8 = B. Haenseleri.
Hemaid, Schott, l. c. = B. Russellianum.
Kotschyi, Schott, l. c. 7 = B. Bovei.
Lehmanni, Schott, l. c. = Helicophyllum Lehmanni.
nobile, Schott, Prod. Aroid. 66 = B. Pyrami.
Olivieri, Schott, Syn. Aroid. 8 = B. Olivieri.
Pyrami, Schott, Prod. Aroid. 66 = B. Pyrami.
Sprunneri, Schott, Syn. Aroid. 7 = B. Sprunneri.

ISCHINA, Walp. Rep. vi. 520 (1847) = ISCHNIA, DC. = **Tamonea**, Aubl. (Verbenac.).

ISCHNANTHUS, Roem. & Schult. Syst. ii. 28, 497 (1817) = **Iohnanthus**, Beauv. (Gramin.).

ISCHNIA, DC. ex Meisn. Gen. 298 [206] (1840); et Prod. ix. 257 (1845) = **Tamonea**, Aubl. (Verbenac.).
verbenacea, DC. Prod. ix. 257.—Mexic.

ISCHNOSIPHON, Koern. in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 316. t. 10, 11. *SCITAMINEAE*, Benth. & Hook. f. iii. 649.
HYMENOCHARIS, Sallab. in Trans. Hort. Soc. i. (1812) 376, nomen prius.
affinis, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 82 = I. Parkeri.
Arouna, *Koern. in Nouv. Mém. Soc. Nat. Mosc.* xi. (1859) 348.—Am. trop.
bambusaceus, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 94; *Hovan. Prod. Scitam.* 11.—Peruv.
densiflorus, *Koern. l. c.* 88.—Bras.
gracilis, *Koern. l. c.* 94.—Am. trop.
guianensis, *Koern. l. c.* 87.—Guian.
laxus, *Koern. l. c.* 85.—Bras.
leucophaeus, *Koern. in Nouv. Mém. Soc. Nat. Mosc.* xi. (1859) t. 10, 11.—Bras.
obliquus, *Koern. l. c.* 341.—Guian.
orbiculatus, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 95.—Bras.
ovatus, *Koern. l. c.* 90.—Bras.
Parkeri, *Koern. l. c.* 81.—N. Granat.
plurispicatus, *Koern. l. c.* 83.—Bras.
polyphyllus, *Koern. l. c.* 93; *Hovan. Prod. Scitam.* 11.—Bras.
rotundifolius, *Koern. l. c.* 95; *Horan. l. c.*—Peruv.
Seubertianus, Koern. l. c. 85 = laxus.
spicatus, *Koern. l. c.* 83.—Guian.
surinamensis, *Koern. l. c.* 93.—Guian.

ISCHNURUS, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 98. *GRAMINEAE*.
pulchellus, *Balf. f. l. c.*—Ins. Socotr.

ISCHUROCHLOA, Buese, in Miq. Pl. Jungh. 389 (1851–55) = **Bambusa**, Schreb. (Gramin.).
floribunda, Buese, l. c. 390 = B. nana.
spinosa, Buese, l. c. = B. Blumeana.

ISCHYRANTHERA, Steud. ex Nand. in Ann. Sc. Nat. Sér. III. xvi. (1851) 103 = **Bellucia**, Neck. (Melast.).
laevigata, Steud. ex Nand. l. c. = B. Aubletii.

ISCHYROLEPIS, Steud. in Flora, xxiii. (1850) 229, nomen; Syn. Pl. Cyp. 249 (1855) = **Restio**, Linn. *subverticillata*, Steud. ll. cc. = R. subverticillatus.

ISEILEMA, Anderss. in Nov. Act. Soc. Sc. Upsal. Ser. III. ii. (1856) 280 = **Anthistiria**, Linn. (Gramin.). *argutum*, Anderss. l. c. 282 = A. argutus. *Mitchellii*, Anderss. l. c. = A. membranaceus. *prostrata*, Anderss. l. c. 281 = A. prostratus. *Wightii*, Anderss. l. c. = A. prostratus.

ISERTA, Batsch, Tabul. 235 (1802) = seq.

ISERTIA, Schreb. Gen. 284 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 85.
BRIGNOLIA, DC. Prod. v. 444 (1830).
BRUINSMANIA, Miq. in Linnaea, xvii. (1843) 72.
PHOSANTHUS, Rafin. in Ann. Sc. Gén. vi. (1820) 82.
breviflora, *Mart. ex Steud. Nom.* ed. II. i. 225 (Quid?). —Bras.
coccinea, Bartl. ex DC. Prod. iv. 457 = Hazenkeana.
coccinea, *Vahl, Eclog. Am.* iii. 27.—Guian.
commutata, Miq. Stirp. Surin. Sel. 179 = spicaeformis.
flava, *Miq. in Linnaea*, xviii. (1844) 613.—Guian. Hazenkeana, *DC. Prod.* iv. 457.—Mexic.
hypoleuca, *Benth. in Hook. Journ. Bot.* iii. (1841) 220. —Guian.
parviflora, *Vahl, Eclog. Am.* ii. 28. t. 15.—Ins. Trinit.
spicaeformis, *DC. Prod.* iv. 457.—Guian.

ISEXINA, Rafin. Sylva Tellur. 112 (1838) = **Cissome**, Linn. (Capparid.).
aurea, Rafin. l. c. = C. lutea, *E. Mey.*

ISGARUM, Rafin. Fl. Tellur. iii. 46 (1836) = **Salsola**? Linn. (‘certe non Salsolacea’ Moq. in DC. Prod. xiii. 1. 192).
didymum, Rafin. l. c.—Hab.?

ISIAS, De Not. in Mem. Acc. Sc. Torin. Ser. II. vi. (1844) 418 = **Serapias**, Linn. (Orchid.).
triloba, De Not. l. c. = S. neglecta.

ISICA, Moench, Meth. 504 (1794) = ISIKA, Adans. = **Lonicera**, Linn. (Caprifol.).

ISIDOREA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 384. t. 25. *RUBIACEAE*, Benth. & Hook. f. ii. 46.
amoena, *A. Rich. l. c.*—Ind. occ.

ISIDROGALVIA, Ruiz & Pav. Fl. Per. iii. 69. t. 302 (1802) = **Tofieldia**, Huds. (Lilac.).
borealis, Stend. Nom. ed. I. 859, in syn. = T. palustris.
calyculata, Ruiz & Pav. Fl. Per. iii. 69, in obs. = T. calyculata.
falcata, Ruiz & Pav. l. c. = T. falcata.
guyanensis, Klotzsch, in Schomb. Faun. & Fl. Gui. 1065, nomen.—Guian.
Moritziana, Klotzsch, ex Baker, in Journ. Linn. Soc. xvii. (1879) 489 = T. falcata.

ISIKA, Adans. Fam. ii. 501 (1763) = **Lonicera**, Linn. (Caprifol.).
alpigena, Borck. Forst. Bot. ii. tt. 1683-4 = L. alpigena.
cerulea, Medic. Phil. Bot. i. 136 = L. caerulea.
lucida, Moench, Meth. 504 = L. alpigena.

ISIPHIA, Rafin. Med. Fl. ii. 232 (1830) = **Aristolochia**, Linn.
glabra, Rafin. l. c. = A. Sipho.
tomentosa, Rafin. l. c. = A. tomentosa.
tripteris, Rafin. l. c. = A. tripteris.

ISIS, Trutt. Arch. Gew. n. 668 (1812) = **Iris**, Tourn.
fimbriata, Trutt. l. c. = Iris japonica.
fulva, Trutt. l. c. n. 669 = Iris fulva.
ruthenica, Trutt. l. c. 670 = Iris ruthenica.

ISMARIA, Rafin. Sylva Tellur. 117 (1838) = **Brickellia**, Ell. (Compos.).
glandulosa, Rafin. l. c.—Hab.?

ISMELIA, Cass. in Dict. Sc. Nat. xli. 40 (1826) = **Chrysanthemum**, Tourn. (Compos.).
Bromonetii, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276 = C. Bromonetii.
corinata, Sch. Bip. l. c. 271 = C. carinatum.
coronopifolia, Sch. Bip. l. c. 273. t. 92 = C. coronopifolium, *Moqf.*

ISMELIA :—
frutescens, Less. Syn. Comp. 255 = Phymaspermum carnosulum.
maderensis, D. Don, in Sweet, Brit. Flow. Gard. N. S. t. 342 = Chrysanthemum pinnatifidum.
nudosa, Less. Syn. Comp. 255 = C. nodosum.
versicolor, Cass. in Dict. Sc. Nat. xli. 40 = C. carinatum.

ISMENE, Salisb. in Trans. Hort. Soc. i. (1812) 342 = **Hymenocallis**, Salisb. (Amaryll.).
Amancaes, Herb. App. 46 = H. Amancaes.
Andreana. Baker, in Gard. Chron. (1884) t. 11 = H. Andreana.
calathiformis, M. Roem Syn. Ensat. 186 = H. calathina.
calathina, Herb. App. 46 = H. calathina.
crimifolia, Salisb. in Trans. Hort. Soc. i. (1812) 342 = H. Amancaes.
defixa, Herb. in Bot. Reg. (1839) Misc. 86.—Peruv.
integra, M. Roem. Syn. Ensat. 186 = H. Amancaes.
Knightii, Knowles & Westc. Flor. Cab. ii. (1838) 51 = H. lacera.
Macleana, Herb. ex Hook. in Bot. Mag. t. 3675 = H. Macleana.
narcissiflora, M. Roem. Syn. Ensat. 186 = H. lacera.
nutans, Herb. App. 46 = Pancratium nutans.
pedunculatum, Herb. Amaryll. 222. t. 35. f. 2 = H Macleana.
Tagliabuei, M. Roem. Syn. Ensat. 186 = H. calathina.
tenuifolia, Baker, in Hook. Bot. Mag. t. 6397 = H. quitoensis.
viridiflora, Lindl. Bot. Reg. (1841) t. 12 = H. Macleana.

ISNARDA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 176 = seq.

ISNARDIA, Linn. Gen. ed. I. 337 (1737) = **Ludwigia**, Linn. (Onagrar.).
alata, DC. Prod. iii. 61 = L. alata.
alternifolia, DC. l. c. 60 = L. alternifolia.
ascendens, Hall. ex Eaton & Wright, Man. ed. VIII. 283 = L. palustris.
aurantiaca, DC. Prod. iii. 61 = L. alternifolia.
capitata, DC. l. c. 60 = L. capitata.
cylindrica, DC. l. c. = L. cylindrica.
discolor, Klotzsch, in Peters, Reise Mossamb. Bot. 70. —Afr. trop.
hastata, Ruiz & Pav. Fl. Per. i. 66. t. 85. f. 6 = Ammania latifolia.
hirsuta, DC. Prod. iii. 60 = L. pilosa.
hirsuta, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 26 = L. hirtella.
hirsuta, Roem. & Schult. Syst. iii. 477 = L. pilosa, hirtella.
lanceolata, DC. Prod. iii. 61 = L. microcarpa.
linearis, DC. l. c. 60 = L. linearis.
microcarpa, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 26 = L. alata.
microcarpa, Poir. Encyc. Suppl. iii. 188 = L. microcarpa.
multiflora, Poir. l. c.; DC. Prod. iii. 60 = L. pilosa.
multiflora, Guill. & Perr. Fl. Seneg. Tent. i. 295 = L. parviflora.
nitida, Poir. Encyc. Suppl. iii. 188 = L. palustris.
palustris, Linn. Sp. Pl. 120 = L. palustris.
palustris, Lusé, Fl. Onil. 38 = Hippuris vulgaris.
pedunculosa, DC. Prod. iii. 60 = L. arcuata.
ramosa, DC. l. c. 61 = L. ramosa.
repens, DC. l. c. 60 = L. repens.
sphaerocarpa, DC. l. c. 61 = L. sphaerocarpa.
stricta, Griseb. Cat. Pl. Cub. 107.—Cuba.
subhastata, Ruiz & Pav. Fl. Per. i. 66 = Ammannia latifolia.
trifolia, Poir. Encyc. Suppl. iii. 188 = trifolia.
virgata, DC. Prod. iii. 60 = L. virgata.

ISOCARPHA, R. Br. in Trans. Linn. Soc. xii. (1816) 110. *COMPOSITAE*, Benth. & Hook. f. ii. 365.
DUNANTIA, DC. Prod. v. 626 (1836).
alternifolia, Cass. in Dict. Sc. Nat. xxiv. 18 = atriplicifolia.
Amalisa, Spreng. ex DC. Prod. v. 493 = Salmia Eupatoria.
angustata, *Griseb. Fl. Brit. W. Ind.* 376.—Ind. occ.
atriplicifolia, *R. Br. in Trans. Linn. Soc.* xii. (1816) 110.—Am. austr.
Bertorii, Balb. ex DC. Prod. v. 140 = Eupatorium Dales.

ISOCARPHA :—
Bilbergiana, *Less. in Linnaea*, vi. (1831) 405.—Am. austr.
divaricata, *Benth. Bot. Voy. Sulph.* 110. t. 41.—Ecuador.
echioides, Less. in Linnaea, v. (1830) 141. t. 2 = Ageratum echioides.
eupatorioides, Gardn in Hook. Lond. Journ. Bot. v. (1846) 456 = Trichogonia salviaefolia.
fastigiata, Gardn. l. c. 455 = Alomia fastigiata.
foliosa, Gardn. l. c. 457 = Alomia foliosa.
Kunthii, Cass. in Dict. Sc. Nat. xxiv. 19 = Splianthes leucantha.
oppositifolia, *R. Br. in Trans. Linn. Soc.* xii. (1816) 110.—Ind. occ.
Pyrethraria, Cass. in Dict. Sc. Nat. xxvi. 280 = Cotula Pyrethraria.
trichopala, Cass. l. c. in nota = oppositifolia.

ISOCHILOS, Spreng. Syst. iii. 754 (1826) = seq.

ISOCHILUS, R. Br. in Ait. Hort. Kew. ed. II. v. 209 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 527.
LEPTOTHRIUM, Kunth, in H. B. & K. Nov. Gen. et Sp. i. 340 (1815).
carnosiflorus, *Lindl. in Gard. Chron.* (1845); *ex Reichb. f. in Walp. Ann.* vi. 448.—Mexic.
cernuus, Lindl. in Ann. & Mag. Nat. Hist. Ser. I. x. (1842) 184 = Epidendrum Karwinskii.
crassiflorus, *Rich. & Gal. in Ann. Sc. Nat. Sér.* III. iii. (1845) 22.—Mexic.
dubius, A. Rich. & Gal. l. c. 23 = Scaphyglottis dubia.
elegans, Focke, in Tijdschr. Natuurk. Wetensch. iv. (1851) 68 = Epidendrum stenopetalum.
fusiformis, *Lindl. Gen. et Sp. Orch.* 113.—Ins. Trinit.
globosus, Lindl. l. c. 112 = Epidendrum globosum.
graminifolius, *H. B. & K. Nov. Gen. et Sp.* i. 340. t. 78.—N. Granat.
graminoides, Hook. Exot. Fl. t. 196 = Dichaea graminoides.
grandiflorum, *Lindl. Bot. Reg.* (1841) sub t. 1.—Peruv.
lancifolius, *Lindl. Gen. et Sp. Orch.* 113.—Ecuador.
latibracteatus, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 22.—Mexic.
leucanthus, *Rodrig. Orch. Noe.* i. 47.—Bras.
linearis, *R. Br. in Ait. Hort. Kew.* ed. II. v. 209.—Ind. occ.
linifolius, *Lindl. Gen. et Sp. Orch.* 113.—Peruv.
luridus, Lindl. Bot. Reg. (1839) Misc. 36 = Scaphyglottis dubia.
major, *Cham. & Schlecht. in Linnaea*, vi. (1831) 60.—Mexic.
proliferum, R. Br. in Ait. Hort. Kew. ed. II. v. 209 = Pomera prolifera.
ramosus, *Focke, in Tijdschr. Natuurk. Wetensch.* iv. (1851) 69.—Guian.
ramosus, Spreng. Syst. iii. 734 = Epidendrum ramosum.
tertifolius, Lindl. Gen. et Sp. Orch. 112 = Epidendrum tertifolium.

ISOCHORISTE, Miq. Fl. Ind. Bat. ii. 822 (1856). *ACANTHACEAE*, Benth. & Hook. f. ii. 1109.
africana, *S. Moore, in Journ. Bot.* xviii. (1880) 300.—Afr. trop.
javanica, *Miq. Fl. Ind. Bat.* ii. 822.—Java.

ISOCOMA, Nutt. in Trans. Am. Phil. Soc. N. S. vi. (1841) 393 = **Bigelovia**, Nutt. (Compos.).
vernonioides, Nutt. l. c. = B. veneta.

ISODECA, Rafin. Fl. Tellur. iii. 88 (1836) = **Lemaea**, Burm. (Labiat.).
flaccida, Rafin. l. c. = L. flaccida.

ISODENDRION, A. Gray, in Proc. Am. Acad. ii. (1852) 324. *VIOLARIEAE*, Benth. & Hook. f. i. 118.
lasiflorium, *A. Gray, l. c.*—Ins. Sandvic.
longifolium, *A. Gray, l. c.*—Ins. Sandvic.
pyrifolium, *A. Gray, l. c.*—Ins. Sandvic.

ISODESMIA, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 339. *LEGUMINOSAE*, Benth. & Hook. f i. 514.
Blanchetiana, *Benth. in Mart. Fl. Bras.* xv. i. 71.—Bras.
tomentosa, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 339.—Bras.

ISODON, Schrad. ex Benth. Lab. Gen. et Sp. 40 (1833) = **Plectranthus**, L'Hérit. (Labiat.).

plectranthoides, Schrad. ex Benth. l. c. 43 = P. rugosus.

ISOËTES *lacustris*, Weig. Obs. Bot. (1772) 86. t. 2. f. 7 = Juncus bufonius.

ISOËTOPSIS, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 174. *COMPOSITAE*, Benth. & Hook. f. ii. 462.

graminifolia, *Turcz. l. c.* 175.—Austral.

ISOGLOSSA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 155. *ACANTHACEAE*, Benth. & Hook. f. ii. 1111.

ECTINANTHUS, T. Anders. in Journ. Linn. Soc. vii. (1863) 45.

angusta, *Benth. & Hook. f. Gen.* ii. 1111; *Baker, in Journ. Linn. Soc.* xxi. (1885) 430.—Madag.

barleioides, *S. Moore, in Journ. Bot.* xv. (1877) 70.—Afr. trop.

ciliata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 155. t. 3.—Afr. austr.

gracillima, *Baker, in Journ. Linn. Soc.* xxi. (1885) 430.—Madag.

justicioides, *Baker, l. c.* xx. (1883) 221.—Madag.

Mellerî, *Baker, l. c.* xxi. (1885) 451.—Madag.

Rutenbergiana, *Vatke, in Bremen, Abh.* ix. (1885) 133. —Madag.

ISOLEPIS, R. Br. Prod. 221 (1810) = **Scirpus**, Linn. (Cyperac.).

acuulis, F. Muell. ex Benth. Fl. Austral. vii. 324 = S. humillimus.

acaulis, Phil. ex Boeck. in Linnaea, xxxvi. (1869–70) 440 = S. acaulis.

acicularis, Schlecht. Fl. Berol. i. 86 = Eleocharis acicularis.

acrostachya, A. Dietr. Sp. Pl. ii. 116 = Ficinia compositis.

acuminata, Schult. Mant. ii. 582 = S. Thouarsianus.

acuminata, A. Dietr. Sp. Pl. ii. 103 = Ficinia acuminata.

albescens, E. Desv. in C. Gay, Fl. Chil. vi. 188 = S. inundatus.

albescens, Miq. in Linnaea, xix. (1847) 226 (Rynchosporae sp.).—Bras.

alticaulos, Steud. Syn. Pl. Cyp. 91 = Eleocharis atropurpurea.

alpina, Hook. f. Fl. Tasm. ii. 86. t. 143 = S. Gunnii.

ambigua, Nees, in Linnaea, ix. (1834) 291 (Bulbostylidis sp.).—Am. trop.

ambigua, Steud. Syn. Pl. Cyp. 91 = Eleocharis Chaetaria.

ambigua, Zoll. Syst. Verz. Ind. Arch. ii. 62 = S. supinus.

anaschilienis, Steud. Syn. Pl. Cyp. 94 = S. inundatus.

ancepa, A. Dietr. Sp. Pl. ii. 125 = Fimbristylis complanata.

angularis, Schrad. ex Schult. Mant. ii. 69 = Fimbristylis quinquangularis.

antarctica, Nees, in Linnaea, vii. (1832) 505; viii. (1833) 33; x. (1836) 162 = S. diabolicus.

antarctica, Roem. & Schult. Syst. ii. 112 = S. antarcticus.

aphylla, Boeck. in Flora, xli. (1858) 417 = Eleocharis albida.

aquatilis, Kunth, Enum. Pl. ii. 215 = S. tenuissimus.

arcuata, Eckl. ex Kunth, l. c. 204 = Ficinia lateralis.

arenaria, Nees, in Linnaea, ix. (1834) 291 (Bulbostylidis sp.).—Am. trop.

aristata, Presl. Delic. Prag. 144 = S. pungens.

armerioides, Miq. Fl. Ind. Bat. iii. 210 (Fimbristylidis sp.).—Ins. Banca.

articulata, Nees, in Wight, Contrib. 108 = S. articulatus.

Ascolepis, A. Rich. Tent. Fl. Abyss. ii. 501 = Ascolepis eriocauloides.

asperiuscula, Kunth, Enum. Pl. ii. 212 = seq.

aspera, H. B. & K. Nov. Gen. et Sp. i. 221 (Bulbostylidis sp.).—Am. trop.

atacamensis, Phil. Fl. Atac. 53 = S. atacamensis.

atropurpurea, Nees, in Linnaea, vii. (1832) 495 = S. tenuis.

atropurpurea, Roem. & Schult. Syst. ii. 106 = Eleocharis atropurpurea.

aucklandica, Hook. f. Fl. Antarct. 88. t. 50 = S. aucklandicus.

ISOLEPIS :—

australis, Roem. & Schult. ex G. Don, in Loud. Hort. Brit. 21 = S. Holoschoenus.

autumnalis, J. & C. Presl, Rel. Haenk. i. 189 = Fimbristylis autumnalis.

barbata, R. Br. Prod. 232 (Bulbostylidis sp.).—Geront. trop.

basilaris, Hook. f. Handb. N. Zeal. Fl. 302 = S. basilaris.

bifolia, Steud. Syn. Pl. Cyp. 316 = S. Isolepis.

Bergiana, Schult. Mant. ii. 582 = S. antarcticus.

Berlandieri, Steud. Syn. Pl. Cyp. 108 (Bulbostylidis sp.).—Mexic

Beyrichii, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 88 (Bulbostylidis sp.).—Bras.

bicolor, Carmich. in Trans. Linn. Soc. xii. (1819) 503 = S. Thouarsianus.

bicolor, Nees, in Linnaea, ix. (1834) 291; x. (1836) 148 = S. venustulus.

bispicata, Schult. Mant. ii. 61 = Fimbristylis schoenoides.

bivalvis, Steud. Syn. Pl. Cyp. 101 (Fimbristylidis sp.). —Bras.

Buchetiert, Oliver, in Trans. Linn. Soc. xxix. (1875) 167 = Mariscus coloratus.

brachyphylla, J. & C. Presl, Rel. Haenk. i. 187 = S. cernuus.

brachyphylla, Schult. Mant. ii. 64 = Fimbristylis capillaris.

brevicalmis, Kunth, Enum. Pl. ii. 207 (Bulbostylidis sp.).—Afr austr.

brevifolia, J. & C. Presl, Rel. Haenk. i. 851 = S. cernuus.

brevis, Brongn. in Duperr. Voy. Coq. Bot. 180 = S. cernuus.

bufonia, H. B. & K. Nov. Gen. et Sp. i. 222 = Fimbristylis capillaris.

bulbosa, Nees, in Linnaea, vii. (1834) 507 = Ficinia bulbosa.

caespitula, Liebm. in Videnak. Skr. V. ii. (1851) 287 = S. microanthus.

capillaris, Roem. & Schult. Syst. ii. 118 = Fimbristylis barbata.

carinata, Hook & Arn. ex Torr. in Ann. Lyc. N. York, iii. (1836) 249 = S. carinatus.

Carmichaelii, A. Dietr. Sp. Pl. ii. 107 = S. sulcatus.

cartilaginea, R. Br. Prod. 232 = S. antarcticus.

cernua, Roem. & Schult. Syst. ii. 106 = S. cernuus.

chaetaria, Link, in Buch, Beschr. Canar. Ins. 188 = S. cernuus.

chlorostachya, Nees, in Linnaea, ix. (1834) 291 = S. cernuus.

chrysocarpa, Nees, l. c. vii. (1832) 499 = S. antarcticus.

ciliata, J. & C. Presl, Rel. Haenk. i. 188 (Bulbostylidis sp.).—Am. trop.

ciliatifolia, Torr. in Ann. Lyc. N. York, iii. (1836) 250 (Bulbostylidis sp.).—Am. bor.

cilicifolia, Steud. Syn. Pl. Cyp. 99 (Bulbostylidis sp.).— Guiana.

cinnamomoturnus, Roem. & Schult. Syst. ii. 119 = Fimbristylis cyperoides.

circinnata, Kunth, Enum. Pl. ii. 214 (Bulbostylidis sp.). —Bras.

coarctata, Torr. in Ann. Lyc. N. York, iii. (1836) 102 (Bulbostylidis sp.).—Am. bor.

cochleata, Steud. Syn. Pl. Cyp. 100 = Fimbristylis setacea.

collina, Kunth, Enum. Pl. ii. 208 (Bulbostylidis sp.).— Afr. austr.

communis, Nees, in Linnaea, x. (1836) 161 = Ficinia communis.

complanata, Roem. & Schult. Syst. ii. 119 = Fimbristylis complanata.

compressa, Nees, in Linnaea, x. (1836) 157 = S. Neesii.

Conceptionis, Steud. Syn. Pl. Cyp. 94 = S. inundatus.

congesta, Schrad. ex Schult. Mant. ii. 70 (Bulbostylidis sp.).—Bras.

consanguinea, Kunth, Enum. Pl. ii. 211 (Bulbostylidis sp.).—Bras.

consociata, Steud. Syn. Pl. Cyp. 318 = Fimbristylis exilis.

ISOLEPIS :—

conspera, Nees, in Endl. Prod. Fl. Norf. 23 = Scirpus inundatus.

controversa, Steud. Syn. Pl. Cyp. 92 = S. cernuus.

coronaria, Roem. & Schult. Syst. ii. 113 = Cyperus leucocephalus.

corymbosa, J. & C. Presl, Reliq. Haenk. i. 188 (Rhynchosporae sp.).—Panama.

corymbosa, Roem. & Schult. Syst. ii. 110 = S. corymbosus.

costata, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 499 = S. costatus.

crassiuscula, Hook. f. Fl. Tasm. ii. 86. t. 143 = S. crassiusculus.

crinita, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 82 (Bulbostylidis sp.).—Am. trop.

Cunninghii, Steud. Syn. Pl. Cyp. 101 = Carex setifolia.

curvata, Zoll. Syst. Verz. Ind. Arch. ii. 62 = S. fluitans.

curvifolia, Schrad. in Schult. Mant. ii. 70 = Fimbristylis diphylla.

curvula, Kunth, Enum. Pl. ii. 189 = S. fluitans.

cuspidata, A. Dietr. Sp. Pl. ii. 130 = Fuirena Wallichiana.

cyperoides, R. Br. Prod. 232 = S. cyperoides.

decipiens, Nees, in Linnaea, ix. (1834) 291 = S. paludicola.

deesa, Schult. Mant. ii. 70 (Bulbostylidis sp.).—Reg. Himal.

depauperata, Link, Jahrb. i. iii. 80 = Fimbristylis diphylla.

desertícola, Phil. Fl. Atac. 53 = S. semisubterraneus.

diabolica, Schrad. Anal. Cap. Cyp. 21; A. Dietr. Sp. Pl. ii. 116 = S. diabolicus.

dichotoma, H. B. & K. Nov. Gen. et Sp. i. 223 = Fimbristylis diphylla.

dichroa, Steud. Syn. Pl. Cyp. 91 = Eleocharis atropurpurea.

digitata, Nees, ex Schrad. Anal. Cap. Cyp. 20 = S. flaccifolius.

diocca, Kunth, Enum. Pl. ii. 199 = S. diocciua.

dipsacea, Roem. & Schult. Syst. ii. 119 = Fimbristylis dipsacea.

dissoluta, Kunth, Enum. Pl. ii. 219 = S. flaccifolius.

distichs, Boeck. in Flora, xli. (1858) 418.—Ins. S. Thomas.

divaricata, A. Dietr. Sp. Pl. ii. 131 = S. divaricatus.

Dregeana, Kunth, Enum. Pl. ii. 204 = S. Hystrix.

Drummondi, Torr. & Hook. ex Torr. in Ann. Lyc. N. York, iii. (1836) 250 = Fimbristylis anomala.

dubia, Kunth, Enum. Pl. ii. 216 = S. flaccifolius.

duna, Zoll. & Mor. Syst. Verz. Zoll. 97 = Fimbristylis sapcrrima.

Echinidium, Nees, in Linnaea, ix. (1834) 291 = S. inoonstulus.

echinocephala, Oliver, in Trans. Linn. Soc. xxix. (1875) 167 = S. cubensis.

eckinulata, Kunth, Enum. Pl. ii. 205 = Cyperus aristatus.

Echeniana, Nees, in Linnaea, vii. (1832) 506 = Ficinia paradoxa.

Echeniana, Schrad. Anal. Cap. Cyp. 15. t. 1. f. 3 = S. verruculus.

Echinis, A. Dietr. Sp. Pl. ii. 116 = Ficinia Eckloca.

Ekrenbergii, Otto, ex Sweet, Hort. Brit. ed. II. 550 (Quid ?).—Aegypt.

elachista, Schult. Mant. ii. 61 = S. dipsacea ?

Elliottii, A. Dietr. Sp. Pl. ii. 131 = Fimbristylis diphylla.

elongata, C. A. Mey. in Ledeb. Fl. Alt. i. 64 = S. pumilus.

erubescens, Steud. in Flora, xxv. (1842) 604 = S. cernuus.

erythrorhizus, Steud. Syn. Pl. Cyp. 96 = S. prolifer.

exiguus, Hook & Arn. Bot. Beech. Voy. 319 = Eleocharis acicularis.

exilis, H. B. & K. Nov. Gen. et Sp. i. 224 = Fimbristylis exilis.

exilis, Nees, in Linnaea, ix. (1834) 291 = S. tenuulus.

expallescens, Kunth, Enum. Pl. ii. 196 = S. expallescens.

falcata, Roem. & Schult. Syst. ii. 116 = Fimbristylis paucispicata.

fasciculata, Kunth, Enum. Pl. ii. 188 = S. Ludwigii.

fasciculata, Sieber, ex Kunth, l. c. 209 (Bulbostylidis sp.).—Afr. trop.

ferruginea, Schlecht. in Linnaea, vi. (1831) 87 = Fimbristylis ferruginea, *Hook. & Arn.*

ISOLEPIS :—

festucoides, Kunth, Enum. Pl. ii. 207 (Bulbostylidis sp.).—Madag.

fibrosa, Nees, in Linnæa, ix. (1834) 292 = Ficinia tenuifolia.

filiculoides, Steud. Syn. Pl. Cyp. 98 = Ficinia tenuifolia.

filamentosa, Roem. & Schult. Syst. ii. 113 (Bulbostylidis sp.).—Guinea.

filiformis, Roem. & Schult. l. c. 106 = Eleocharis tenuis.

fimbriata, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 88 (Bulbostylidis sp.).—Bras.

fistulosa, Delile, Fl. Aegypt. Illustr. 50 = Scirpus articulatus.

fluitans, R. Br. Prod. 221 = S. fluitans.

fornicularis, Schrad. ex Boeck. in Linnæa, xxxv. (1868–70) 752 (Bulbostylidis sp.).—Abyss.

fontana, Steud. Syn. Pl. Cyp. 97 = S. Holoschoenus.

Funchii, Steud. l. c. 91 (Bulbostylidis sp.).—Venezuela.

furcata, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 88 = S. cernuus.

fusca, Link, Jahrb. i. 111. 79 (Bulbostylidis sp.).—Am. austr.

fuscata, Meyen, Reise, i. 447, nomen.—Peruv.

fuscata, Nees, in Linnæa, ix. (1834) 291 = S. cernuus.

fuscescens, Steud. Syn. Pl. Cyp. 92 = S. nervosus.

fuscopurpurea, Steud. l. c. 99 = Eleocharis maculosa.

Gardneriana, Steud. l. c. 99 (Bulbostylidis sp.).—Bras.

Gaudichaudiana, Kunth, Enum. Pl. ii. 201 = S. inundatus.

Giraudyi, Steud. Syn. Pl. Cyp. 91 = S. fluitans.

glaucescens, Schrad. & Schult. Mant. ii. 68 = Fimbristylis squarrosa.

globosa, J. Buch. in Trans. N. Z. Inst. iii. (1871) 211 = S. prolifer.

globulifera, Steud. Syn. Pl. Cyp. 97 = S. Holoschoenus.

globulosa, Schult. Mant. ii. 68 = Fimbristylis globulosa.

glomerata, Schrad. ex Schult. Mant. ii. 67 = Fimbristylis spathacea.

gracilis, Nees, in Linnæa, x. (1835–36) 161 = Ficinia Poiretii.

gracilis, Nees, in Wight, Contrib. 109 (Bulbostylidis sp.).—Ind. or.

gracilis, Sweet, Hort. Brit. ed. I. 488 = S. nodosus.

grandispica, Steud. Syn. Pl. Cyp. 818 = S. maritimus.

Gunnii, Steud. l. c. 94 = S. Gunnii, inundatus.

Haenkei, J. & C. Presl, Rel. Haenk. 187 = Fimbristylis spathacea.

hamulosa, Kunth, Enum. Pl. ii. 204 = S. hamulosus.

Hartmanni, Steud. Syn. Pl. Cyp. 100 = Fimbristylis Vahlii.

hemisphærica, A. Dietr. Sp. Pl. ii. 109 = S. Isolepis.

heterolepis, Steud. Syn. Pl. Cyp. 93 = Carex Berteroana.

heteromorpha, Steud. l. c. 100 = Eleocharis nodulosa.

heterophylla, Steud. in Flora, xxv. (1842) 608 ; et l. c. 93 = S. cernuus.

hirta, H. B. & K. Nov. Gen. et Sp. i. 224 = Fimbristylis squarrosa.

hirtella, Schrad. ex Schult. Mant. ii. 70 (Bulbostylidis sp.).—Bras.

hispida, Steud. Syn. Pl. Cyp. 104 (Bulbostylidis sp.).—Am. austr.

Holoschœnus, Mellius, St. Hel. 343 = S. nodosus.

Holoschœnus, Roem. & Schult. Syst. ii. 115 = S. Holoschœnus.

Hookeri, Nees, ex Boeck. in Linnæa, xxxvi. (1869–70) 498 = S. corinutus.

Hookeriana, Boeck. in Flora, xli. (1858) 418 = S. nemoralis.

Humboldtii, Roem. & Schult. Syst. ii. 113 = S. micranthus.

humilis, Kunth, Enum. Pl. ii. 207 (Bulbostylidis sp.).—Afr. austr.

Hystrix, Kunth, l. c. 204 = S. Hystrix.

inclinata, Delile, Fl. Aegypt. Illustr. 50 = S. corymbosus.

inconspicua, Nees, in Linnæa, ix. (1834) 291 = S. inundatus.

incurvata, Nees, in Wight, Contrib. 108 = S. articulatus.

intricata, Steud. Syn. Pl. Cyp. 818 = S. fluitans.

inundata, R. Br. Prod. 222 = S. inundatus.

involucellata, Steud. Syn. Pl. Cyp. 101 (Bulbostylidis sp.).—Java.

junciformis, H. B. & K. Nov. Gen. et Sp. i. 229 (Bulbostylidis sp.).—Am. austr.

Junodiana, Mtz. Fl. Ind. Bot. iii. 813 = S. supinus.

Kunthii, Steud. Syn. Pl. Cyp. 93 = S. cernuus.

koilolepis, Steud. Syn. Pl. Cyp. 818 = Scirpus carinatus.

Krauss, Phil. in Linnæa, xxx. (1859–60) 208 = Schœnus nitens.

Kunthiana, Steud. Syn. Pl. Cyp. 95 = S. incomtulus.

kyllingioides, A. Rich. Tent. Fl. Abyss. ii. 502 = S. kyllingioides.

Labillardieri, Steud. Syn. Pl. Cyp. 96 (Bulbostylidis sp.).—N. Caled.

laciniata, Eckl. ex Kunth, Enum. Pl. ii. 255 = Ficinia Ecklonea.

lanata, H. B. & K. *Nov. Gen. et Sp. i.* 220. t. 68 (Bulbostylidis sp.).—Am. austr.

Langsdorffiana, Kunth, Enum. Pl. ii. 214 (Bulbostylidis sp.).—Bras. ; Peruv.

latifolia, A. Dietr. Sp. Pl. ii. 132 = Pleurostachys elegans.

lenticularis, R. Br. Prod. 222 = S. lenticularis.

lepida, Nees, in Linnæa, ix. (1834) 291, nomen.—Hab. ?

leptalea, Schult. Mant. ii. 62 = S. cernuus.

leptocaulis, Torr. in Pacif. Rail. Rep. iv. 153 = S. cernuus.

leptos, Steud. Syn. Pl. Cyp. 91 = S. nanus.

leptostachya, Kunth, Enum. Pl. ii. 190 = S. leptostachyus.

Lereschii, Shuttlw. ex Parl. Fl. Ital. ii. 68 = Eleocharis acicularis.

leucostachya, H. B. & K. Nov. Gen. et Sp. i. 220 (Bulbostylidis sp.).—Am. trop.

Lichtensteiniana, Kunth, Enum. Pl. ii. 207 (Bulbostylidis sp.).—Ins. S. Helena.

linearis, Nees, in Linnæa, x. (1835) 160 = Ficinia gracilis.

linearis, Roem. & Schult. Syst. ii. 117 = S. linearis.

longifolia, Nees, in Linnæa, ix. (1834) 291, nomen.—Hab. ?

longiflora, Steud. Syn. Pl. Cyp. 90 = Eleocharis acicularis.

longispica, Steud. l. c. 104 (Fimbristylidis sp.).—Hab. ?

Ludwigii, Kunth, Enum. Pl. ii. 189 = S. Ludwigii.

Maguirei, Nees, in Wight, Contrib. 107 = S. quinquefarius.

magellanica, Gandich. in Freyc. Voy. Bot. 414 = S. cernuus.

margaritifera, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 46 = Schœnus apogon.

marginata, A. Dietr. Sp. Pl. ii. 110 = Ficinia filiformis.

Martiana, Schrad. ex Nees, in Mart. Fl. Bras. i. t. 84 (Bulbostylidis sp.).—Bras.

Martii, Roem. & Schult. Syst. ii. 117 = Cladium germanicum.

maxima, A. Dietr. Sp. Pl. ii. 130 = S. grossus.

megapotamica, A. Dietr. l. c. 122 = Fycreus megapotamicus.

membranacea, Nees, in Linnæa, x. (1836) 159 = S. membranaceus.

mesantha, Nees, l. c. ix. (1834) 291 = S. supinus.

Meyeriana, Nees, l. c. ii. (1827) 291 = S. cernuus.

Michelianus, Roem. & Schult. Syst. ii. 114 = S. Michelianus.

micrantha, Roem. & Schult. l. c. 110 = S. micranthus.

microcarpa, Nees, in Linnæa, x. (1836) 150 = S. cernuus.

miliacea, J. & C. Presl, Rel. Haenk. i. 188 = Fimbristylis miliacea.

Minæ, Parl. Giorn. Bot. Ital. ii. i. (1846) 146 = S. cernuus.

minima, Schrad. in Goett. Gel. Anz. (1821) iii. 2068 = S. Isolepis.

modesta, Phil. in Linnæa, xxix. (1857–58) 79 = S. cernuus.

monandra, Link, Jahrb. i. iii. 78, in nota = S. setaceus.

monocephala, Steud. Syn. Pl. Cyp. 97 = Ficinia brevifolia.

monostachya, Spreng. Neue Entdeck. iii. 11 = Fimbristylis schœnoides.

mucronata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 178 = S. mucronatus.

Muehlenbergii, A. Dietr. Sp. Pl. ii. 106 (Bulbostylidis sp.).—Am. bor.

multicaulis, Schlecht. in Linnæa, xx. (1847) 562 = S. cernuus.

multiarrectus, Boeck. in Flora, xli. (1858) 418 = S. inundatus.

nana, Phil. in Linnæa, xxix. (1857–58) 79 = S. inundatus.

natans, A. Dietr. Sp. Pl. ii. 106 = S. rivularis.

nervosa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 499 = Scirpus nervosus.

nigrescens, Steud. Syn. Pl. Cyp. 91 = Eleocharis nigrescens.

nigricans, H. B. & K. Nov. Gen. et Sp. i. 220 = S. inundatus.

nitens, Roem. & Schult. Syst. ii. 117 = Rynchospora nitens.

nodosa, R. Br. Prod. 221 = S. nodosus.

nolata, Nees, in Lehm. Pl. Preiss. ii. 74 = S. antarcticus.

nudipes, Kunth, *Enum. Pl. ii.* 206 = *Eleocharis grandis.*

numidiana, Roem. & Schult. Syst. ii. 110 = S. cernuus.

obtusifolia, Roem. & Schult. l. c. 118 = Fimbristylis obtusifolia.

oligantha, A. Dietr. Sp. Pl. ii. 103 = Ficinia filiformis.

oligantha, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 197 = S. pumilus.

oryzetorum, Steud. Syn. Pl. Cyp. 96 = S. supinus.

Ottonis, Boeck. in Flora, xli. (1858) 419 (Fimbristylidis sp.).—Bras.

pallescens, A. Dietr. Sp. Pl. ii. 121 = Fimbristylis dichotoma.

pallida, Nees, in Linnæa, viii. (1833) 81. t. 3. f. 2 = S. cernuus.

paludicola, Kunth, Enum. Pl. ii. 198 = S. paludicolus.

palustris, Schrad. Anal. Cap. Cyp. 17. t. 1. f. 7 = S. nivalaris.

panamensis, J. & C. Presl, Rel. Haenk. i. 351 (Rynchospora sp. ?).—Panama.

paniculata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 78 = S. Holoschœnus.

panormitana, Parl. Giorn. Bot. Ital. ii. i. (1846) 146 = S. Holoschœnus.

paradoxa, Kunth, Enum. Pl. ii. 206 (Bulbostylidis sp.).—Venezuela.

paradoxa, Schrad. in Goett. Gel. Anz. (1821) iii. 2068 = Ficinia paradoxa.

paratnis, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 86 (Bulbostylidis sp.).—Bras.

pauciflora, Steud. Syn. Pl. Cyp. 104 (Bulbostylidis sp.).—Mexic.

pentagona, Schult. Mant. ii. 69 = Fimbristylis quinquangularis.

pentasticha, Boeck. in Flora, xlii. (1859) 446 = S. supinus.

perpusilla, Phil. in Anal. Univ. Chil. (1873) 565 = S. perpusillus.

Perrottetii, Steud. Syn. Pl. Cyp. 818 = Fimbristylis exilis.

phæocarpa, Nees, in Linnæa, x. (1836) 153 = S. antarcticus.

pholoides, Steud. Syn. Pl. Cyp. 98 = S. cernuus.

pilosa, Steud. l. c. 97 (Bulbostylidis sp.).—Austral.

planifolia, Spreng. Neue Entdeck. iii. 10 = S. paludicolus.

plebeia, Schrad. Anal. Cap. Cyp. 18. t. 1. f. 1, 6 = S. setaceus.

pleurocarpa, Hochst. ex Boiss. Fl. Orient. v. 379 = S. setaceus.

podocarpa, Boeck. in Flora, xliii. (1860) 179 (Fimbristylidis sp. ?).—Hab. ?

Pohlsana, Boeck. l. c. xli. (1858) 596 (Bulbostylidis sp.).—Bras.

Poiretii, Roem. & Schult. Syst. ii. 116 = S. Holoschœnus.

Poiretti, Steud. Syn. Pl. Cyp. 98 = Ficinia Poiretti.

polycolea, De Not. Ind. Sem. Hort. Genev. (1847) = S. supinus.

polyphylla, A. Rich. Tent. Fl. Abyss. ii. 503 = S. Steudneri.

prælongata, Nees, in Wight, Contrib. 108 = S. articulatus.

Preslii, A. Dietr. Sp. Pl. ii. 129.—Panama.

prolifera, R. Br. Prod. 222 = S. prolifer.

prolifera, Carmich. in Trans. Linn. Soc. xii. (1819) 503 = S. Thouarsianus.

prolifera, Hook. f. Fl. N. Zel. i. 271 ; et Fl. Tasm. ii. 87. t. 144 = S. inundatus, subcinctus, cernuus.

prolifera, Mellius, St. Hel. 344 = S. cernuus.

prolongata, Nees, in Wight, Contrib. 108 = S. articulatus.

propinqua, R. Br. Prod. 222 = S. inundatus.

propinqua, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 46 = S. antarcticus.

prostrata, Steud. Syn. Pl. Cyp. 95 = S. supinus.

Pseudo-junciformis, Boeck. in Flora, xl. (1857) 85 (Bulbostylidis sp.).—Mexic.

Pseudo-schœnus, A. Dietr. Sp. Pl. ii. 103 = Ficinia tristachya.

ISOLEPIS :—
psilocarpa. Kunth, Enum. Pl. ii. 193 = S. inundatus.
ptycholepis, Steud. Syn. Pl. Cyp. 93 = Scirpus verrucosulus.
puberula, Kunth, Enum. Pl. ii. 213 (Bulbostylidis sp.). —Madag.
pubescens, Roem. & Schult. Syst. ii. 118 = Fuirena pubescens.
publiculmis, Boeck. in Linnæa, xxxvii. (1871-73) 38 = Fimbristylis exilis.
pubigera, Reichb. ex Kunth, Enum. Pl. ii. 227 = Fimbristylis exilis.
pubigera, Schrad. Ind. Sem. Hort. Gotting. fasc. 1; ex Schult. Mant. ii. 476 = Fimbristylis dichotoma.
pulchella, Thw. Enum. Pl. Zeyl. 350 (Fimbristylidis sp.) —Zeylan.
pumila, Link, Hort. Berol. i. 286 = S. cernuus.
pumila, Roem. & Schult. Syst. ii. 106 = S. pumilus.
punctulata, Steud. Syn. Pl. Cyp. 92 = S. cernuus.
purpurascens, Steud. l. c. = S. cernuus.
pusilla, Kunth, Enum. Pl. ii. 190 = S. tenuis.
pygmæa, Kunth, Enum. Pl. ii. 191 = S. cernuus.
racemosa. Schult. Mant. ii. 532 = S. racemosus.
radiata, Roem. & Schult. Syst. ii. 113 = Ficinia radiata.
radicifera, Steud. Syn. Pl. Cyp. 318 (Bulbostylidis sp.)—Am. bor.
ramulosa, Steud. in Lechl. Berb. Am. Austr. 50 = S. inundatus.
rariflora, Schrad. ex Schult. Mant. ii. 61 = Fimbristylis schoenoides.
repens, Nees, in Linnæa, ix. (1834) 291 = Ficinia repens.
rigescens, Presl, in Oken, Isis, xxi. (1828) 268 (Fimbristylidis sp.).—Ins. Maurit.
riparia, Steud. in Lechl. Berb. Am. Austr. 56 = S. rigidus.
riparia, K. Br. Prod 222 = S. cernuus.
riparia, Schrad. Anal. Cap. Cyp. 19. t. 1. f. 5 = S. rivularis.
robustula, Steud. Syn. Pl. Cyp. 90 = S. capillifolius.
rumana, Roem. & Schult. ex G. Don, in Loud. Hort. Brit. 21 = S. Holoschœnus!
Royleï, Nees, in Wight, Contrib. 107 = S. quinquefarius.
rubescens, Presl, Delic. Prag. 157.—Bras.
rubicunda, Kunth, Enum. Pl. ii. 188 = S. Ludwigii.
rugosa, Boeck. in Linnæa, xxxvi. (1869–70) 702 = S. articulatus.
rupestris, Kunth, Enum. Pl. ii. 193 = S. cernuus.
Sariana, Schult. Mant. ii. 65 = S. cernuus.
Savii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 174 = S. cernuus.
scabra, J. & C. Presl, Rel. Haenk. 188 (Bulbostylidis sp.).—Peruv.
scariosa, Nees, in Linnæa, vii. (1832) 501 = Ficinia scariosa.
Schimperiana, Hochst. in Flora, xxiv. (1841) t. Intell. 21 (Bulbostylidis sp.).—Abyssin.
schœnoides, Kunth, Enum. Pl. ii. 208 (Bulbostylidis sp.). —Afr. austr.
Schomburghii, Steud. Syn. Pl. Cyp. 102 = Fimbristylis acutivalis.
Schraderi, A. Dietr. Sp. Pl. ii. 120 = Fimbristylis spathacea.
Schraderiana, Steud. Syn. Pl. Cyp. 103 (Bulbostylidis sp.).—Bras.
Schweinfurthiana, Oliver, in Trans. Linn. Soc. xxix. (1875) 167 (Bulbostylidis sp.).—Afr. trop.
Sellowiana, Kunth, Enum. Pl. ii. 208 (Fimbristylidis sp.).—Bras.
semipedunculata, Boeck. in Flora, xli. (1858) 417 = S. antarctious.
senegalensis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 500 = S. articulatus.
sepulcralis, Steud. Syn. Pl. Cyp. 94 = S. cernuus.
sesleriordes, Kunth, Enum. Pl. ii. 195 = S. antarcticus.
setacea, R. Br. Prod. 222 = S. setaceus.
setifolia. A. Rich. Tent. Fl. Abyss. ii. 498 = Elecocharis atropurpures.
setosa, Raoul, ex A. Rich. Voy. Astrol. Bot. 104 = S. cernuus.
sicula, Presl, Cyp. Gram. Sicul. 13 = S. cernuus.
Sieberi, Schrad. Anal. Cap. Cyp. 28, nota (Bulbostylidis sp.).—Afr. austr.
similima, Steud. Syn. Pl. Cyp. 95 = S. sapinus.
sororia, Kunth, Enum. Pl. ii. 192.—Afr. austr.
sphærocarpa, Nees, in Linnæa, x. (1836) 158 = S. antarcticus.
sphærophora, Ehrenb. ex Boeck. in Linnæa, xxxvi. (1869–70) 721 = S. Holoschœnus.

ISOLEPIS :—
spiralis, A. Rich. Voy. Astrol. Bot. t. 19 = S. frondosus.
squarrosa, Carmich. in Trans. Linn. Soc. xii. (1819) 508 = Scirpus Thouarsianus.
squarrosa, H. B. & K. Nov. Gen. et Sp. i. 221 = S. microanthus.
squarrosa, Roem. & Schult. Syst. ii. 111 = S. squarrosus.
stemphylla, Torr. in Ann. Lyc. N. York, iii. (1836) 288 (Bulbostylidis sp.).—Am. bor.
Steudelii, Schrad. Anal. Cap. Cyp. 19 = Ficinia Ecklonea.
striata, Kunth, Enum. Pl. ii. 189 = S. capillifolius.
striatella, Steud. Syn. Pl. Cyp. 92 = S. cernuus.
stricta, A. Dietr. Sp. Pl. ii. 131 = Fimbristylis spathacea.
striolata, Nees, ex Boeck. in Linnæa, xxxvi. (1869–70) 700 = S. sapinus.
subaculis, Steud. Syn. Pl. Cyp. 318 = Fimbristylis Vahlii.
subcuculiata, Bergg. in Minneskr. Fisiog. Sallsk. Lund, (1877) 23.—N. Zel.
subquarrosa, Schrad. ex Schult. Mant. ii. 64 = S. microanthus.
subtilis, Kunth, Enum. Pl. ii. 191 = S. cernuus.
subtilissima, Boeck. in Flora, xli. (1858) 416 = S. cernuus.
subtristachya, Hochst. ex Boeck. in Linnæa, xxxvi. (1869–70) 752 (Bulbostylidis sp.).—Abyss.
sulcata, Carmich. in Trans. Linn. Soc. xii. (1819) 508 = S. sulcatus.
supina, R. Br. Prod. 221 = S. supinus.
tenella, Kunth, Enum. Pl. ii. 213 (Fimbristylidis sp.). —Am. trop.
tenerrima, Fisch. & Mey. ex Ledeb. Fl. Ross. iv. 258 (Bulbostylidis sp.).—As. centr.
tenuifolia. A. Dietr. Sp. Pl. ii. 131 (Bulbostylidis sp.). —Am. trop.
tenuior, Steud. Syn. Pl. Cyp. 91 = S. Ludwigii.
tenuis, Presl, Delic. Prag. 144 = S. cernuus.
tenuis, Schrad. Anal. Cap. Cyp. 15 = S. tenuis.
tenuissima, D. Don, Prod. Fl. Nep. 40 (Bulbostylidis sp.) —Nepal.
tenuissima, Nees, in Linnæa, ix. (1834) 291 = S. tenuissimus.
terragona, Schult. Mant. ii. 69 = Fimbristylis millacea.
Thouarsii, Nees, in Linnæa, ix. (1834) 291 (Bulbostylidis sp.).—Hab.!
Thuarsii. A. Dietr. Sp. Pl. ii. 109 = S. Thouarsianus.
Thunbergii, Schrad. in Goett. Gel. Anz. (1821) iii. 2068 = S. Thunbergii.
trachycarpa, Boeck. in Linnæa, xxxvi. (1869–70) 501, in obs. = S. cernuus.
trachycarpa, Nees, l. c. x. (1836) 152 = S. cernuus.
trichoides, Schrad. in Schult. Mant. ii. 64 (Bulbostylidis sp.).—Bras.
trichoides, Steud. Syn. Pl. Cyp. 96 (Bulbostylidis sp.). —Java.
trifida, Nees, in Wight, Contrib. 108 (Bulbostylidis sp.).—Nepal.
trigyna, Kunze, ex Boeck. in Linnæa, xxxvi. (1869–70) 502 = S. cernuus.
tristachya, Phil. in Linnæa, xxx. (1859–60) 208.—Chili.
tristachya, Roem. & Schult. Syst. ii. 110 = Ficinia tristachya.
truncata, Nees, in Linnæa, vii. (1832) 508 = Ficinia truncata.
truncata, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 85 (Bulbostylidis sp.).—Bras.
turbinatocarpa, Kegel, in Act. Hort. Petrop. vii. (1880) 561 = Fimbristylis diphylla.
uliginosa, Steud. Syn. Pl. Cyp. 91 = Elecocharis acicularis.
uncinata, Roem. & Schult. Syst. ii. 111 = Fuirena uncinata.
unioides, Delile, Fl. Egypt. 153 = S. supinus.
Urvillei, Steud. Syn. Pl. Cyp. 94 = S. inundatus.
vaginata, Eckl. ex Steud. Nom. ed. II. i. 528 = Fimbristylis 'bulbosa' (Quid?).
Vahlii, H. B. & K. Nov. Gen. et Sp. i. 221 = Fimbristylis Vahlii.
Valdiviæ, Steud. Syn. Pl. Cyp. 94 = S. inundatus.
varia, Schrad. ex Nees, in Mart. Fl. Bras. ii. t. 75 = Fimbristylis diphylla.
variana, Steud. Syn. Pl. Cyp. 94 = S. inundatus.
variegata, Steud. l. c. 98 = Carex setifolia.
venustula, Kunth, Enum. Pl. ii. 192 = S. venustulus.
verruciflora, Maxim. Prim. Fl. Amur. 300 (Fimbristylidis dipsacea).

ISOLEPIS :—
verrucosula, Steud. Syn. Pl. Cyp. 93 = Scirpus verrucosulus.
Vincentii, Kunth, Enum. Pl. ii. 210 (Bulbostylidis sp.). —Am. trop.
Vincentii, Steud. Syn. Pl. Cyp. 102 = Fimbristylis Vahlii.
vivipara, Schrad. ex E. Desv. in C. Gay, Fl. Chil. vi. 169 = S. inundatus.
Wallichiana, Schult. Mant. ii. 538 (Bulbostylidis sp.). —Reg. Himal.
Wawri, Torr. in Ann. Lyc. N. York, iii. (1836) 354 (Bulbostylidis sp.).—Florida.
Wildenowii, Kunth, Enum. Pl. ii. 210 (Bulbostylidis sp.).—Afr. trop.
Wildenowii, Roem. & Schult. Syst. ii. 120 = Fimbristylis complanata.

ISOLOBA, Rafin. Fl. Tellur. iv. 58 (1836) = **Pinguicula**, Tourn. (Lentib.).
elatior, Rafin. l. c. 59 = P. elatior.
lusitanica, Rafin. l. c. = P. lusitanica.
lutea, Rafin. l. c. = P. lutea.
pumila, Rafin. l. c. = P. pumila.
recurva, Rafin. l. c. = P. lutea.

ISOLOBUS, A. DC. in Dec. Prod. vii. 352 (1839) = **Lobella**, Linn. (Campanulac.).
caespitosus, Hasak. in Bonplandia, vii. (1859) 180 = L. radicans.
campanuloides, A. DC. in DC. Prod. vii. 353 = L. radicans.
concolor, A. DC. l. c. 354 = Pratia erecta.
corymbosus, A. DC. l. c. 352 = L. corymbosa.
Cunninghamii, A. DC. l. c. 354 = Pratia erecta.
Echlonianus, Sond. in Harv. & Sond. Fl. Cap. iii. 536 = L. Ecklonianus.
jasionoides, A. DC. in DC. Prod. vii. 353 = L. corymbosa.
Kerii, A. DC. l. c. = L. radicans.
radicans, A. DC. l. c. = L. radicans.
Roxburghianus, A. DC. l. c. = L. radicans.
sparsiflorus, A. DC. l. c.—Afr. austr.

ISOLOMA, Decne. in Rev. Hortic. Sér. III. ii. (1848) 466. *GESNERACEAE*, Benth. & Hook. f. ii. 1001.
BRACHYLOMA, Hanst. in Linnæa, xxvi. (1853) 308.
CALYCOSTEMMA, Hanst. l. c. xxix. (1857–58) 506.
CRYPTOLOMA, Hanst. l. c. (1857–58).
GIESLERIA, Regel, in Flora, xxxii. (1849) 181.
KOHLERIA, Regel, l. c. xxxi. (1848) 250.
MOURONIA, Regel, l. c. 249 (1848).
PEARCEA, Regel, Gartenfl. xvi. (1867) 886.
SCIADOCALYX, Regel, l. c. ii. (1859) 257. t. 61, partim.
TYDAEA, Decne. in Rev. Hortic. Sér. III. ii. (1848) 468.
brevifloruu, *Lindl. & Paxt. Flow. Gard.* i. (1850-51) 63.—Panama.
cordifolium, *Hemsl. Biol. Centr. Am. Bot.* ii. 478.—Costa Rica.
costaricense, *Hemsl. l. c.*—Costa Rica.
Deppeanum, *Hemsl. l. c.*—Mexic.
elatius, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 465.—Mexic.
elegans, *Hemsl. Biol. Centr. Am. Bot.* ii. 478.—Guatemala.
erianthum, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 465. —N. Granat.
hirsutum, *Regel, in Bot. Zeit.* ix. (1851) 893.—Ind. occ.
hondense, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 465. —N. Granat.
hypocyttiforum, *Benth. & Hook. f. Gen.* ii. 1002.—Ecuador.
ignoratum, *Hemsl. Biol. Centr. Am. Bot.* ii. 478.—Mexic.
inæquale, *Benth. & Hook. f. Gen.* ii. 1002.—N. Granat.
incanum, *Hemsl. Biol. Centr. Am. Bot.* ii. 478.—Costa Rica.
incurvum, *Hemsl. l. c.*—Panama.
Kranerianum, *Lehm. in Hamb. Gartenz.* x. (1854) 456. —Am. centr.
lasianthum, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 465.—Mexic.
Linkianum, *Hemsl. Biol. Centr. Am. Bot.* ii. 478.—Guatemala.

ISOLOMA:—
longifolium, *Decne. in Rev. Hortic*, Sér. III. ii. (1848) 465.—Guatemala.
longipes, *Decne. l. c.*—N. Granat.
molle, *Decne. l. c.*—Venezuela.
ocellatum, *Benth. & Hook. f. Gen.* ii. 1002.—Panama.
petiolare, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 465.—Panama.
pictum, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 185.—N. Granat.
pictum, *Regel, Gartenfl.* iii. (1854) 110.—Panama.
pilosum, *Hemsl. Biol. Centr. Am. Bot.* ii. 479.—Costa Rica.
purpureum, *Benth. & Hook. f. Gen.* ii. 1003.—Peruv.
rhynchocarpum, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 465.—N. Granat.
rubricaule, *Regel*, in *Bot. Zeit.* ix. (1851) 893.—Venezula.
rupestre, *Hemsl. Biol. Centr. Am. Bot.* ii. 479.—Nicaragua.
Schiedeanum, *Hemsl. l. c.*—Mexic.
Seemannii, *Benth. & Hook. f. Gen.* ii. 1002.—Panama.
spicatum, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 465.—Mexic.
strictum, *Hemsl. Biol. Centr. Am. Bot.* ii. 479.—Costa Rica.
tetragonum, *Hemsl. l. c.* 480.—Costa Rica.
Trianaei, *Regel, Gartenfl.* (1854) 107.—N. Granat.
triflorum, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 465.—N. Granat.
tubiflorum, *Decne. l. c.*—Panama.
verticillatum, *Decne. l. c.*—Peruv.
vestitum, *Decne. l. c.*—N. Granat.

ISOLOPHUS, Spach, Hist. Vég. Phan. vii. 112 (1839) = **Polygala**, Linn.
sanatilis, Spach, l. c. 114 = P. rupestris.

ISOMERIA, D. Don, ex DC. Prod. v. 24 (1836) = **Vernonia**, Schreb. (Compos.).
chinensis, Wight, ex DC. l. c. = V. cinerea.
cinerea, Wight, ex DC. l. c. = V. cinerea.

ISOMERIS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 124 (1838). *CAPPARIDEÆ*, Benth. & Hook. f. i. 106. ℓ. ii. 657.
arborea, *Nutt. ex Torr. & Gray, l. c.*—Am. bor. occ.

ISONANDRA, Wight, Ic. ii. 4. tt. 859, 860 (1840); et tt. 1219, 1220. *SAPOTACEÆ*, Benth. & Hook. f. ii. 657.
acuminata, Drury, Useful Pl. Ind. 260 = Dichopsis elliptica.
acuminata, *Miq. Fl. Ind. Bat. Suppl.* 581.—Sumatra.
argentata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1858) 415.—Java.
Benjamina, *De Vriese, ex Miq. in Journ. Bot. Neerl.* i. (1861) 258.—Borneo.
calonevra, Kurs, For. Fl. Brit. Burma, ii. 119 = Bassia calonevra.
calophylla, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 88 = Bassia calonevra.
calophylla, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 36.—Java.
canaliculata, Thw. Enum. Pl. Zeyl. 177 = Dichopsis canaliculata.
Candolleana, *Wight, Ic.* t. 1220.—Ind. or.
dasyphylla, *Miq. Fl. Sumgh.* 201 (gen. dub.).—Java.
diplostemon, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 540.—Ind. or.
grandis, Thw. Enum. Pl. Zeyl. 176 = Dichopsis grandis.
Gutta, Hook. Lond. Journ. Bot. vi. (1847) 463 = Dichopsis Gutta.
hexandra, Griff. Notul. iv. 393 = Dichopsis hexandra.
laevifolia, Thw. Enum. Pl. Zeyl. 177 = Dichopsis laevifolia.
lampongo, *Miq. Fl. Ind. Bat. Suppl.* 581.—Sumatra.
lanceolata, Thw. Enum. Pl. Zeyl. 443 = Dichopsis lanceolata.
lanceolata, *Wight, Ic.* tt. 859, 860.—Ind. or.
macrophylla, *De Vriese, ex Miq. in Journ. Bot. Neerl.* i. (1861) 258.—Borneo.
microphylla, *De Vriese, ex Miq. l. c.* 260.—Borneo.
Mottleyana, Miq. l. c. 257 = Bassia Mottleyana.
obovata, Griff. Notul. iv. 393 = Dichopsis obovata.
pauciflora, Thw. Enum. Pl. Zeyl. 177 = Dichopsis pauciflora.
Perrottetiana, *A. DC. in DC. Prod.* viii. 168.—Ind. or.
polyandra, Wight, Ic. t. 1589 = Payena lucida.

ISONANDRA:—
polyantha, Kurs, For. Fl. Brit. Burma, ii. 119 = Dichopsis polyantha.
puberula, *Miq. Fl. Sumgh.* 201 (gen. dub.).—Sumatra.
pulchra, *Burck. in Ann. Jard. Buitenz. v.* (1885) 21.—Sumatra.
quercifolia, *De Vriese, ex Miq. in Journ. Bot. Neerl.* i. (1861) 260.—Borneo.
Richii, *A. Gray, in Proc. Am. Acad. v.* (1861–62) 327.—Ins. Amicor.
rostrata, *Miq. Fl. Ind. Bat. Suppl.* 581.—Ins. Banca.
rubiginosa, Thw. Enum. Pl. Zeyl. 177 = Dichopsis rubiginosa.
Stocksii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 539.—Ind. or.
sumatrana, *Burck. in Ann. Jard. Buitenz. v.* (1885) 21.—Sumatra.
villosa, *Wight, Ic* t. 860.—Ind. or.
Wightiana, *A. DC. in DC. Prod.* viii. 167.—Zeylan.
xanthocyma, *De Vriese, ex Miq. in Journ. Bot. Neerl.* i. (1861) 259.—Borneo.

ISONEMA, R. Br. in Mem. Wern. Soc. i. (1809) 63. *APOCYNACEÆ*, Benth. & Hook. f. ii. 712.
Smeathmanni, *Roem & Schult. Syst.* iv. 401.—Afr. trop.

ISONEMA, Cass. in Bull. Soc. Philom. (1817) 152 = **Vernonia**, Schreb. (Compos.).
ovata, Cass. in Dict. Sc. Nat. xxiv. 25 = V. chinensis.

ISOPAPPUS, Torr. & Gray, Fl. N. Am. ii. 329 (1840) = **Aplopappus**, Cass. (Compos.).
divaricatus, Torr. & Gray, l. c. = A. divaricatus.
Hookerianus, Torr. & Gray, l. c. = A. Hookerianus.

ISOPARA, Rafin. Atl. Journ. 144 (1832) = **Cleomella**, DC. (Capparⁱ.).
mexicana, Rafin. l. c. = C. mexicana.

ISOPETALUM, Sweet, Geran. t. 126 (1820) = **Pelargonium**, L'Hérit. (Geraniac.).
alchemilloides, Eckl. & Zeyh. Enum. 76 = P. alchemilloides.
aphanoides, Eckl. & Zeyh. l. c. = P. aphanoides.
articulatum, Eckl. & Zeyh. l. c. = P. articulatum.
Calyledonis, Sweet, Geran. t. 126 = P. Coryledonis.
dimidiatum, Eckl. & Zeyh. Enum. 76 = P. alchemilloides.
discipes, Colv. ex Sweet, Hort. Brit. ed. I. 85 = P. discipes.
elongatum, Eckl. & Zeyh. Enum. 76 = P. tabulare.
geranioides, Eckl. & Zeyh. l. c. = P. alchemilloides.
ranunculophyllum, Eckl. & Zeyh. l. c. = P. alchemilloides.

ISOPHYLLUM, Hoffm. Gen. Umbellif. 116 (1814) = **Bupleurum**, Tourn. (Umbellif.).
affine, Schur, Enum. Pl. Transs. 251 = B. affine.
australe, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 391 = B. Gerardi.
baldense, Hoffm Gen. Umbellif. 115 = B. baldense.
falcatum, Hoffm. l. c. = B. falcatum.
Gerardi, Hoffm. l. c. = B Gerardi.
glaucum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 391 = B. glaucum.
Jacquinianum, Fourr. l. c. = B. Gerardi.
junceum, Hoffm. Gen. Umbellif. 115 = B. juniceum.
petraeum, Hoffm. l. c. = B. petraeum.
tenuissimum, J. & C. Presl, Fl. Cech. 66 = B. tenuissimum.

ISOPHYLLUM, Spach, Hist. Vég. Phan. v. 432 (1836); et in Ann. Sc. Nat. Sér. II. v. (1836) 367 = **Ascyrum**, Linn. (Hypericin.).
Drummondii, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = Hypericum Isophyllum.

ISOPHYSIS, T. Moore, in Proc. Linn. Soc. ii. (1858) 212 = **Hewardia**, Hook. (Liliac.).
tasmanica, T. Moore, l. c. = H. tasmanica.

ISOPLESION, Rafin. Fl. Tellur. iv. 86 (1836) = **Behrtun**, Tourn. (Borag.).
giganteum, Rafin. l. c. = E. giganteum.
italicum, Rafin. l. c. = E. italicum.
pyrenaicum, Rafin. l. c. = E. italicum.
rubrum, Rafin. l. c. = E. rubrum.

ISOPLEXIS, Lindl. ex Benth. in Bot. Reg. sub t. 1770 (1835). *SCROPHULARINEÆ*, Benth. & Hook. f. ii. 961.
CALLIANASSA, Webb & Benth. Phyt. Canar. iii. 148. t. 188 (1836–50).
canariensis, *Steud. Nom.* ed. II. i. 329.—Ins. Canar.
Sceptrum, *Steud. l. c.*—Ins. Mader.

ISOPOGON, R. Br. ex Knight, Prot. 93 (1809); et R. Br. in Trans. Linn. Soc. x. (1810) 71. *PROTEACEÆ*, Benth. & Hook. f. iii. 179.
ATYLUS, Salisb. Parad. Lond. t. 67 (1807), partim.
adenanthoides, *Meisn. in Hook. Kew Journ.* vii. (1855) 69.—Austral.
anemonefolius, *Knight, Prot.* 93.—Austral.
anethifolius, *Knight, l. c.* 94.—Austral.
asper, *R. Br. Prot. Nov.* 8.—Austral.
attenuatus, *R. Br. in Trans. Linn. Soc.* x. (1810) 73.—Austral.
axillaris, *R. Br. l. c.* 74.—Austral.
Baxteri, *R. Br. Prot. Nov.* 9.—Austral.
buxifolius, *R. Br. in Trans. Linn. Soc.* x. (1810) 73.—Austral.
ceratophyllus, *R. Br. l. c.* 79.—Austral.
cornigerus, Lindl. Swan River App. 34 = teretifolius.
erithmifolius, *F. Muell. Fragm.* vi. 239.—Austral.
cuneatus, *R. Br. in Trans. Linn. Soc.* x. (1810) 73.—Austral.
divaricatus, Sweet, Hort. Brit. ed. II. 448 = anethifolius.
divergens, *R. Br. Prot. Nov.* 7.—Austral.
Drummondii, *Benth. Fl. Austral.* v. 344.—Austral.
formosus, *R. Br. in Trans. Linn. Soc.* x. (1810) 72.—Austral.
heterophyllus, *Meisn. in Lehm. Pl. Preiss.* i. 504.—Austral.
horridus, F. Muell. ex Meisn. in DC. Prod. xiv. 280 = ceratophyllus.
lanceolatus, Meisn. in Lehm. Pl. Preiss. i. 503 = cuneatus.
latifolius, *R. Br. Prot Nov* 8.—Austral.
linearis, *Meisn. in Hook. Kew Journ.* vii. (1855) 69.—Austral.
longifolius, *R. Br. in Trans. Linn. Soc.* x. (1810) 73.—Austral.
Loudoni, Baxter, ex R. Br. Prot. Nov. 8 = cuneatus.
pedunculatus, R. Br. l. c. 7 = Petrophila seminuda ?
petiolaris, *A. Cunn. ex R. Br. l. c.* 8.—Austral.
petrophiloides, R. Br. Prot. Nov. 7 = teretifolius.
petrophioloides, Meisn. in Lehm. Pl. Preiss. i. 503 = Drummondii.
polycephalus, *R. Br. in Trans. Linn. Soc.* x. (1810) 73.—Austral.
propinquus, *A. Cunn. ex G. Don, in Loud. Hort. Brit.* 97, nomen.—Austral.
Protea, Meisn. in Hook. Kew Journ. iv. (1852) 183 = latifolius.
roseus, *Lindl. Bot. Reg.* (1842) *Misc.* 39.—Austral.
scaber, Lindl. Swan River App. 34 = asper.
scaber, Meisn. in Lehm. Pl. Preiss. i. 506 = roseus.
scabriusculus, *Meisn. in Hook. Kew Journ.* iv. (1852) 183.—Austral.
spathulatus, R. Br. Prot. Nov. 8 = buxifolius.
spheraroccphalus, *Lindl. Swan River App.* 34.—Austral.
teretifolius, *R. Br. in Trans. Linn. Soc.* x. (1810) 71.—Austral.
tridens, *F. Muell. Fragm.* vi. 239.—Austral.
trilobus, *R. Br. in Trans. Linn. Soc.* x. (1810) 72.—Austral.
trilobus, Meisn. in Lehm. Pl. Preiss. i. 507 = tripartitus.
tripartitus, *R. Br. Prot. Nov.* 8.—Austral.
uncinatus, *R. Br. l. c.*—Austral.
villosus, *Meisn. in Hook. Kew Journ.* iv. (1852) 183.—Austral.

ISOPTERIS, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 197; et Abh. Acad. Berl. (1855) 181. t. 12 B (Isopteryx in textu) = **Begonia**, Tourn.
longirostris, Klotzsch, ll. cc. = B. longirostris.
umbellata, Klotzsch, in Monatsb. l. c. = B. umbellata.

ISOPTERIS, Wall. Cat. n. 7261 (1831) = **Trigonostemon**, Miq. (Polygal.).
penanginna, Wall. l. c. = T. hypoleucum.

ISOPTERYX, Klotzsch, in Abh. Akad. Berl. (1855) 131. t. 12 B = **Begonia**, Tourn.
umbellata, Klotzsch, l. c. 132 = B. umbellata.

ISOPYRUM, Adans. Fam. ii. 460 (1763) = **Anemone**, Linn. (Ranunc.).

ISOPYRUM, Linn. Gen. ed. II. 945 (1742). *RANUNCULACEAE*, Benth. & Hook. f. i. 8
ENNEION, Rafin. in Journ. Phys. sci. (1820) 70.
FONTANELLA, Kluk, ex Besa. Prim. Fl. Gallic. ii. 868 (1808).
GONTARELLA, Gilib. ex Steud. Nom. ed. I. 880 (1821).
LEPTOPYRUM, Reichb. Consp. 192 (1828).
OLPA, Adans. Fam. ii. 458 (1763).
adiantifolium, *Hook. f. & Thomas. Fl. Ind.* i. 42.—Reg. Himal.
adoxoides, *DC. Syst.* i. 894.—Japon. ; China.
adoxoides, Franch. & Sav. Enum. Pl. Jap. i. 11 = nipponicum.
album, Dulac, Fl. Hautes-Pyr. 221 = thalictroides.
anemonoides, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 135 = thalictroides.
aquilegioides, Bory & Chaub. Fl. Pélop. 35 = Thalictrum orientale.
aquilegioides, *Linn. Sp. Pl.* 557.—Europ. austr.
biternatum, *Torr. & Gray, Fl. N. Am.* i. *Suppl.* 660.—Am. bor.
caespitosum, *Boiss. & Hohen. Diagn.* Ser. I. viii. 7.—Persia.
capnoides, Flach. ex DC. Prod. i. 48 = adoxoides.
Clarkii, *Kellogg, in Proc. Calif. Acad.* vi. (1877) 181. —Calif.
dicarpon, Franch. & Sav. Enum. Pl. Jap. i. 11 ; ii. 271 = stoloniferum.
disarpon, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 7.—Japon.
dicarpon, S. Moore, in Journ. Bot. xvi. (1878) 193 = trachyspermum.
fumariaefolium, Saliab. in Trans. Linn. Soc. viii. (1807) 306 = seq.
fumarioides, *Linn. Sp. Pl.* 557.—Europ. ; As. bor. grandiflorum, *Fisch. ex DC. Prod.* i. 48.—Sibir. altaic. ; Reg. Himal.
Hallii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 374.—Am. bor. occ.
japonicum, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 181 = adoxoides.
microphyllum, Royle, Illustr. Bot. Himal. 54. t. 11. f. 4 = grandiflorum.
nipponicum, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 83.—Japon.
occidentale, *Hook. & Arn. Bot. Beech. Voy.* 816. —Calif.
Raddeanum, *C. Muell. in Walp. Ann.* vii. 28 ; *Maxim. in Bull. Acad. Pétersb.* xxix. (1884) 61.—Mandshur.
stipitatum, *A. Gray, in Proc. Am. Acad.* xii. (1877) 54.—Calif.
stipulaceum, *Franch. & Sav. Enum. Pl. Jap.* ii. 270.—Japon.
stoloniferum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 60.—Japon.
thalictrifolium, Gilib. Fl. Lituan. ii. 281 ; Saliab. in Trans. Linn. Soc. viii. (1807) 304 = thalictroides.
thalictroides, *Linn. Sp. Pl.* 557.—Europ. austr.
thalictroides, Short, Cat. Fl. Kentuck. i. 8 = biternatum.
trachyspermum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 60.—Japon.
uniflorum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 149.—Afghan.

ISORA, Adans. Fam. ii. 382 (1763) ; Schott & Endl. Melet. 31 (1832) = **Helicteres**, Linn. (Stercul.).
corylifolia, Schott & Endl. l. c. = H. isora.
grewioefolia, Schott & Endl. l. c. = H. spicata.

ISORIUM, Rafin. Fl. Tellur. ii. 61 (1836) = **Lobostemon**, Lehm. (Borag.).
formosum, Rafin. l. c. = L. formosa.

ISOSCHOENUS, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 49 = **Schoenus**, Linn. (Cyperac.).
acuminatus, Nees, l. c. = S. acuminatus.
Armeria, Nees, l. c. = S. Armeria.
barbatus, Nees, in Lehm. Pl. Preiss. ii. 80 = S. barbatus.
Drummondii, Steud. Syn. Pl. Cyp. 167 = S. flavus.
flavus, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 49 = S. flavus.

ISOSTIGMA, Less. in Linnaea, vi. (1831) 513. *COMPOSITAE*, Benth. & Hook. f. ii. 389.
brasiliense, *Benth. & Hook. f. Gen.* ii. 389.—Besa.
crithmifolium, *Less. in Linnaea*, vi. (1831) 515.—Bras.
crithmifolium, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 389 = brasiliense.
lobatum, *Baker*, in *Mart. Fl. Bras.* vi. iii. 239.—Bras.
microcephalum, Baker, l. c. = brasiliense.
pseudaxillaris, *Less. in Linnaea*, vi. (1831) 514.—Bras.
Riedelii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. iii. 240 = speciosum.
simplicifolium, *Less. in Linnaea*, vi. (1831) 513.—Bras.
speciosum, *Less. l. c.* 515.—Bras.

ISOSTOMA, D. Dietr. Syn. Pl. i. 569, 737 (1839), sphalm. = **Isotoma**, Lindl. (Campan.).

ISOSTYLIS, Spach, Hist. Vég. Phan. x. 402 (1840) = **Banksia**, Gaertn. (Proteac.).

ISOTOMA, Lindl. Bot. Reg. t. 964 (1826). *CAMPANULACEAE*, Benth. & Hook. f. ii. 588.
HIPPOBROMA, G. Don, Gen. Syst. iii. 717 (1834).
axillaris, *Lindl. Bot. Reg.* t. 964.—Austral.
Baueri, *Presl, Prod. Lobel.* 42.—Austral.
brevifiora, *Steud. Nom.* ed. II. i. 839.—Am. austr.
brevifolia, Presl, Prod. Lobel. 43 = Brownii.
Brownii, *Don. Gen. Syst.* iii. 718.—Austral.
fasciculata, Schlecht. in Lechl. Berb. Am. Austr. 58 = Rhizocephalum Candollei.
fluviatilis, *F. Muell. ex Benth. Fl. Austral.* iv. 186.—Austral.
Gulliverii, *F. Muell. Fragm.* x. 89.—Austral.
longiflora, *Presl, Prod. Lobel.* 42.—Ind. occ.
petraea, *F. Muell. in Linnaea*, xxv. (1852) 420.—Austral.
pusilla, *Benth. in Enum. Pl. Hueg.* 75.—Austral.
rhytidophora, F. Muell. Fragm. x. 40 = Lobelia rhytidosperma.
runcinata, *Hassk. in Bonplandia*, vii. (1859) 181.—Hab. ?
scapigera, *G. Don, Gen. Syst* iii. 716.—Austral.
senecioides, A. DC. in DC. Prod. vii. 412 = axillaris.

ISOTREMA, Rafin. in Am. Monthly Mag. (1819) 195 = **Aristolochia**, Linn.
Sipho, Rafin. l. c. = A. Sipho.

ISOTRIA, Rafin. ined. Hesp. Redes. N. York, v. (1808) 857 = **Pogonia**, Juss. (Orchid.).
medeoloides, Rafin. Fl. Tellur. iv. 47 = P. verticillata.
verticillata, Rafin. in Med. Repos. N. York, v. (1808) 857 = P. verticillata.

ISOTROPIS, Benth. in Enum. Pl. Hueg. 28 (1837). *LEGUMINOSAE*, Benth. & Hook. f. i. 468.
atropurpurea, *F. Muell. Fragm.* iii. 16.—Austral.
blidea, Benth. in Enum. Pl. Hueg. 29 = striata.
canescens, *F. Muell. Fragm.* x. 51.—Austral.
cuneifolia, *Benth. in Enum. Pl. Hueg.* 29, in nota.—Austral.
Drummondii, *Meisn. in Lehm. Pl. Preiss.* i. 81.—Austral.
filicaulis, *Benth. in Ann. Wien. Mus.* ii. (1838) 71.—Austral.
Forrestii, *F. Muell. in Wing, South Sc. Record*, ii.—(1882) 259.—Austral.
juncea, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 281.—Austral.
parviflora, *Benth. in Ann. Wien. Mus.* ii. (1838) 71. —Austral.
striata, *Benth. in Enum. Pl. Hueg.* 28.—Austral.
Wheeleri, *F. Muell. ex Benth. Fl. Austral.* ii. 40.—Austral.
Winneckeana, *F. Muell. Census, Suppl.* i. 8.—Austral.

ISOTYPUS, H. B. & K. Nov. Gen. et Sp. iv. 11. t. 307 (1820) = **Onoseris**, DC. (Compos.).
onoseroides, H. B. & K. l. c. = O. isotypus.
rosiflorus, Triana, ex Hérincq, L'Hortic. Franç. (1864) 137.—N. Granat.

ISQUIERDA, Willd. Sp. Pl. iv. 115 (1805) = IZQUIERDIA, Ruiz & Pav. = **Myrsine**), Linn.

ISQUIERDIA, Poir. Encyc. Suppl. iii. 190 = preec.

ISYPUS, Rafin. Fl. Tellur. iv. 73 (1836) = **Aesculus**, Linn. (Sapindac.).
ochraceus, Rafin. l. c. = Ae. glabra.

ITEA, Gronov. ex Linn. Gen. ed. II. 85 (1742). *SAXIFRAGEAE*, Benth. & Hook. f. i. 647.
DICONANGIA, Miich. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 208.
carolinana, Lam. Encyc. iii. 315 = Cyrilla racemiflora.
chinensis, *Hook. & Arn. Bot. Beech. Voy.* 189. t. 39.—Reg. Himal. ; China.
Cyrilla, Sw. Prod. Veg. Ind. Occ. 50 = Cyrilla racemiflora.
floribunda, Saliab. Prod. 142 = Cyrilla racemiflora.
fragrans, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 430 = Polyosma fragrans.
japonica, Hort. ex Presl, Symb. Bot. i. 40 = Sloanea monosperma.
japonica, *Oliver, in Journ. Linn. Soc.* lx. (1867) 164. —Japon.
javanica, *Blume, Bijdr.* 160.—Java.
macrophylla, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 419.—Reg. Himal.
nutans, *Royle, Illustr. Bot. Himal.* 226.—Reg. Himal.
padifolia, Saliab. Prod. 142 = virginica.
rosmarinifolia, Poir. Encyc. Suppl. iii. 190 = seq.
Rosmarinus, Roem. & Schult. Syst. v. 408 = Baeckea cochinchinensis.
spinosa, Andr. Bot. Rep. t. 314 = Bursaria spinosa.
umbellata, *Roxb. Hort. Beng.* 85 ; *Fl. Ind.* i. 682.—Ins. Malay.
virginica, *Linn. Sp. Pl.* 188.—Am. bor.

ITEADAPHNE, Blume, Mus. Bot. Lugd. Bat. i. 865 (1851). *LAURINEAE*, Benth. & Hook. f. iii. 162
confusa, *Blume*, l. c.—Malaya.

ITIA, Molina, Sagg. Chile, ed. II. 281 (1810) ; ex Roem. & Schult. Syst. v. 259, in obs. (1819) = **Lonicera**, Linn. (Caprifol.).

ITICANIA, Rafin. Sylva Tellur. 125 (1836) = **Loranthus**, Linn.
loniceroides, Rafin. l. c. = L. lonicerioides.

ITTNERA, C. C. Gmel. Fl. Bad. iii. 590. t. 4 (1808) = **Najas**, Linn.
major, C. C. Gmel. l. c. = N. major.
minor, C. C. Gmel. l. c. 592 = N. minor.
Najas, C. C. Gmel. l. c. 590 = N. major.

ITUTERION, Rafin. Sylva Tellur. 86 (1836) = **Witis**, Tourn. (Ampelid.).
arborea, Rafin. l. c. = V. arborea.

IUGA, Reichb. Consp. 157 (1828), sphalm. = **Inga**, Scop. (Legumin.).

IULOCROTON, Baill. Adansonia, iv. (1866) 367 = **Julocroton**, Mart. (Euphorb.).

IULUS, Saliab. Gen. Pl. Fragm. 92 (1866) = **Allium**, Tourn. (Liliac.).
arenosus, Saliab. l. c. = A. subhirsutum.
subhirsutus, Saliab. l. c. = A. subhirsutum.

IVA, Linn. Hort. Upsal. 285 (1748) ; Amoen. Acad. iii. 22 (1757). *COMPOSITAE*, Benth. & Hook. f. ii. 252.
CYCLACHAENA, Fresen. Ind. Sem. Hort. Frankf. (1836) 4 ; et in Linnaea, xii. (1838) Lit. 78.
DENIRA, Adans. Fam. ii. 118 (1763).
EUPHROSYNE, DC. Prod. v. 530 (1836).
ambrosiaefolia, *A. Gray, Syn. Fl. N. Am.* i. ii. 346.—Am. bor.
angustifolia, *Nutt. ex DC. Prod.* v. 529.—Am. bor. austr.
annua, *Linn. Sp. Pl.* 988.—Am. austr.
annua, Michx. Fl. Bor. Am. ii. 164 = ciliata.
asperifolia, *Less. in Linnaea*, v. (1830) 151.—Mexic.
axillaris, *Pursh, Fl. Am. Sept.* ii. 743.—Am. bor.
cheiranthifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 276.—Cuba.
ciliata, *Willd. Sp. Pl.* iii. 2286.—Am. bor.
declinata, *A. Gray, Pl. Wright.* i. 104.—N. Mexic.
foliolosa, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 346 = axillaris.

IVA :—
frutescens, *Linn. Sp. Pl.* 989.—Am. bor.
Hayesiana, *A. Gray, in Proc. Am. Acad.* xi. (1876) 78. —Calif.
imbricata, *Walt. Fl. Carol.* 232.—Am. bor.
integrifolia, Banks, ex Pursh, Fl. Am. Sept. ii. 580 = imbricata.
microcephala, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 346.—Am. bor.
monophylla, Walt. Fl. Carol. 232 = Ambrosia artemisiifolia.
nevadensis, *M. E. Jones, in Am. Natural.* xvii. (1883) 978.—Am. bor. occ.
paniculata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 347 = xanthifolia.
xanthifolia, *Nutt. Gen. Am.* ii. 135.—Am. bor.

IVESIA, Torr. & Gray, ex Torr. in Pacif. Rail. Rep. vi. 79 (1857) = **Potentilla**, Linn. (Rosac.).
Baileyi, S. Wats. in Bot. King's Exped. 90, 449.—Am. bor. occ.
depauperata, A. Gray, in Bot. Calif. i. 184 = P. depauperata.
Gordoni, A. Gray, in Pacif. Rail. Rep. vi. 79 = P. Gordoni.
gracilis, Torr. & Gray, l. c. 79. t. 11 = P. Newberryi.
Kingii, S. Wats. in Bot. King's Exped. 91, 448.—Am. bor. occ.
Lemmoni, S. Wats. in Proc. Am. Acad. xx. (1885) 368.—Arizona.
lycopodioides, A. Gray, l. c. vi. (1864–65) 530 = P. Gordoni.
Muirii, A. Gray, l. c. viii. (1873) 697 —Calif.
Pickeringii, Torr. in Bot. U. St. Expl. Exped. 288. t. 4 = P. Pickeringii.
pinnatifida, S. Wats. in Proc. Am. Acad. xx. (1885) 364.—Am. bor. occ.
pygmæa, A. Gray, l. c. vi. (1864–65) 531 = P. Gordoni.
santolinoides, A. Gray, l. c. = P. santolinoides.
tridentata, A. Gray, l. c. vii. (1868) 336.—Am. bor. occ.
unguiculata, A. Gray, l. c. 339.—Calif.
utahensis, S. Wats. l. c. xvii. (1881–82) 371.—Am. bor. occ.
Webberi, A. Gray, l. c. x. (1874) 71.—Calif.

IVIRA, Aubl. Pl. Gui. ii. 694. t. 279 (1775) = **Sterculia**, Linn.
pruriens, Aubl. l. c. 696 = S. Ivira.

IVONIA, Vell. Fl. Flum. 100 (1825) ; iii. t. 4 (1827) (Quid ?).
amplexicaulis, Vell. ll. oc.—Bras.

IXALUM, Forst. f. Prod. 92 (1786) = **Spinifex**, Linn. (Gramin.).
inerme, Forst. f. l. c. = S. hirsutus.

IXANTHUS, Griseb. Gen. et Sp. Gent. 129 (1839). *GENTIANEÆ*, Benth. & Hook. f. ii. 810.
viscosus, *Griseb. l. c.*—Ins. Canar.

IXAUCHENUS, Cass. in Dict. Sc. Nat. lvi. 176 (1828) = **Lagenophora**, Cass. (Compos.).
lyratus, Less. Syn. Comp. 193 = L. Billardieri.
sublyratus, Cass. in Dict. Sc. Nat. lvi. 176 = L. Billardieri.

IXERBA, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 249. *SAXIFRAGEÆ*, Benth. & Hook. i. 645.
brexioides, *A. Cunn. l. c.*—N. Zel.

IXERIS, Cass. in Bull. Soc. Philom. (1821) 178 ; et Dict. Sc. Nat. xxiv. 49 (1822) = **Lactuca**, Tourn. (Compos.).
albiflora, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 397 = L. albiflora.
debilis, A. Gray, l. c. = L. debilis.
floribunda, DC. Prod. vii. 151 = L. polycephala.
Gregorii, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 232 = L. Thunbergii.
integrifolia, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 398.—Ins. Bonin.
lyrata, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 190.—Japon.
monophlebia, Cass. in Dict. Sc. Nat. xxix. 389 (= *Microrhynchus pumilus*).—Ins. Falkland.

IXERIS :—
polycephala, Cass. in Dict. Sc. Nat. xxiv. 50 = Lactuca polycephala.
ramosissima, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 397 = L. denticulata.
repens, A. Gray, l. c. = L. repens.
sonchifolia, Hance, in Journ. Linn. Soc. xiii. (1873) 108 = L. denticulata ?
stolonifera, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 396 = L. stolonifera.
Thunbergii, *A. Gray*, l. c. 397 = L. Thunbergii.
versicolor, DC. Prod. vii. 151 = L. Fischeriana.

IXERRA, Pritz. Ic. Ind. i. 602 (1855), sphalm. = **Ixerba**, Cunn. (Saxifr.).

IXIA, Linn. Coroll. Gen. 1 (1787). *IRIDEÆ*, Benth. & Hook. f. iii. 704.
FREESIA, Eckl. Top. Verz. 80 (1827).
HYALIS, Salisb. in Trans. Hort. Soc. i. (1812) 317.
MORÆPHILA, Ker-Gawl. Irid. Gen. 103 (1827).
WURTHIA, Regel, in Bot. Zeit. ix. (1851) 656.
abbreviata, Houtt. Handleid. xii. 41. t. 78. f. 3 = maculata.
acuta, Bartr. Fl. N. Am. i. 76 = Nemastylis geminiflora.
acuta, Licht. ex Roem. & Schult. Syst. i. 353 = Hesperantha longituba.
africana, Hornem. Cat. Hort. Havn. 68 ; ex Roem. & Schult. Syst. i. 424 = Babiana rubrocyanea.
africana, Linn. Sp. Pl. 56 = Aristæa cyanea.
alba, *Eckl. Top. Verz.* 87.—Afr. austr.
alba, Linn. Syst. ed. X. 862 = Sparaxis bulbifera ?
alboflavens, Eckl. Top. Verz. 87 = maculata.
alopecuroides, Linn. f. Suppl. 92 = Watsonia spicata.
amœna, *Salisb. Prod.* 8.—Afr. austr. ?
americana, Aubl. Pl. Gui. i. 33 = Eleutherine plicata.
amœna, *Link, Enum. Hort. Berol.* i. 52.—Afr. austr.
amœna, Salisb. Prod. 8 = maculata.
anemonæflora, [DC. in] Red. Lil. t. 85 = Sparaxis bulbifera.
anemonæflora, *Jacq. Coll. Suppl.* 9.—Afr. austr.
angusta, Jacq. in Ker. ii. 7. t. 279 = Hesperantha angusta.
angustifolia, *Eckl. Top. Verz.* 25.—Afr. austr.
autumnalis*formis*, *Lam. Encyc.* iii. 334.—Afr. austr.
arcuata, *Burm. f. Fl. Cap. Prod.* 9.—Afr. austr.
aristata, *Ker-Gawl. in Bot. Mag.* t. 589.—Afr. austr.
aristata, Schneev. Ic. t. 87 = patens.
aristata, [Soland. in] Ait. Hort. Kew. ed. I. i. 57 = Sparaxis grandiflora.
aristata, Thunb. Diss. Ixia, 14 = leucantha.
aulica, [Soland. in] Ait. Hort. Kew. ed. I. i. 57 = scariosa.
aurantiaca, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 397 = Intes.
autumnalis, Salisb. Prod. 38 = Crocus autumnalis.
azurea, Banks, ex Ker-Gawl. in Bot. Mag. t. 598 = Geissorhiza Rochensis.
bulbifera, Linn. Cent. Pl. ii. 4 ; Amœn. Acad. iv. 300 = Sparaxis bulbifera.
bulbifera, Thunb. Diss. Ixia, n. 16, partim = Sparaxis grandiflora.
bulbocodioides, D. Delar. Desc. Pl. Nov. 19 = Romulea bulbocodioides.
Bulbocodium, Lam. Encyc. iii. 335 = Romulea rosea.
Bulbocodium, Linn. Sp. Pl. ed. II. 51 = Romulea Bulbocodium.
Bulbocodium, Sibth. & Sm. Fl. Græc. i. 36. t. 36 = Romulea Linaresii.
Bulbocodium, Thunb. Diss. Ixia, 6 = Romulea crocata.
bulbocoides, *D. Delar. Desc. Pl. Nov.* 19 ; *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
Burmanni, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* iv. 408.—Afr. austr.
caerulescens, *Eckl. Top. Verz.* 26.—Afr. austr.
campanulata, Banks, ex Roem. & Schult. Syst. i. 391 = speciosa.
campanulata, *Houtt. Handleid.* xii. 42. t. 78. f. 4.—Afr. austr.
campanulata, Lam. Illustr. i. 109 = Romulea hirsuta.

IXIA :—
campanuloides, *Eckl. Top. Verz.* 26, nomen.—Afr. austr.
cana, Eckl. l. c. = viridiflora.
candida, [Delile, in] Red. Lil. viii. t. 496 = leucantha.
capillaris, *Thunb. Diss. Ixia*, 12. t. 2. f. 2.—Afr. austr.
capitata, Andr. Bot. Rep. t. 50 = maculata.
carnosina, *Hort. ex Hoeven & De Vriese, Tijdschr.* viii. (1841) 184.—Hab. ?
cartilaginea, Lam. Encyc. iii. 340 = Watsonia marginata.
caryophyllacea, *Burm. f. Fl. Cap. Prod.* 1*.—Afr. austr.
cepacea, [DC. in] Red. Lil. ii. t. 96 = Micranthus fistulosus.
chinensis, Linn. Sp. Pl. 36 = Belamcanda punctata.
chloroleuca, Jacq. Coll. iv. 180 = Romulea chloroleuca.
ciliaris, Salisb. ex Ker-Gawl. in Bot. Mag. sub t. 549 = Geissorhiza ciliaris.
cinnamomea, Andr. Bot. Rep. t. 44 = Hesperantha falcata.
cinnamomea, Thunb. Diss. Ixia, 10 = Hesperantha cinnamomea.
coccinea, *Eckl. Top. Verz.* 24.—Afr. austr.
coccinea, Thunb. Fl. Cap. i. 241 = speciosa, patens.
coelestina, Bartr. Trav. ed. Carmin. 152. t. 3 = Nemastylis coelestina.
corruscrescens, Pers. Syn. i. 48 = Babiana villosa.
columellaris, [*Ker-Gawl. in*] *Bot. Mag.* t. 630.—Afr. austr.
Columnæ, Schult. Mant. i. 378 = Romulea Columnæ.
columnaris, Salisb. Prod. 36 = monadelpha.
conclina, *Salisb. l. c.*—Afr. austr.
concolor, Salisb. l. c. = patens.
conica, Salisb. l. c. = maculata.
conspicua, *Salisb. l. c.*—Afr. austr.
corymbosa, Linn. Cent. Pl. ii. 4 ; Amœn. Acad. iv. 300 = Lapeyrousia corymbosa.
crateroides, Ker-Gawl. in Bot. Mag. t. 594 = speciosa.
crispa, Linn. f. Suppl. 91 = Tritonia undulata.
crispifolia, Andr. Bot. Rep. t. 35 = Lapeyrousia corymbosa.
crocata, [Delar. in] Red. Lil. t. 335 = Tritonia miniata.
crocata, Linn. Sp. Pl. ed. II. 52 = Tritonia crocata.
crocea, Thunb. Fl. Cap. i. 218 = Romulea subulata.
cruciata, Jacq. Coll. Suppl. 16 = Romulea cruciata.
curta, Andr. Bot. Rep. t. 564 = monadelpha.
cyanea, Pers. Syn. i. 49 = Babiana stricta.
densiflora, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 397 = patens.
deusta, [Soland. in] Ait. Hort. Kew. ed. I. i. 60 = Tritonia deusta.
disticha, Lam. Encyc. iii. 333 = Witsenia maura.
dubia, Vent. Choix, 10 = lutea.
elliptica, *Thunb. Fl. Cap.* i. 244.—Afr. austr.
elongata, Vahl, Enum. Pl. ii. 51 = Romulea elongata.
emarginata, Lam. Encyc. iii. 342 = Lapeyrousia juncea.
enatifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. *Ixi.* 19 = Perdanthus chinensis.
erecta, Berg. Descr. Pl. Cap. 8 = polystachya.
erecta, Jacq. Hort. Schoenb. t. 18 = latea.
erosa, *Salisb. Prod.* 36.—Afr. austr.
europæa, Pers. Syn. i. 46 = Trichonema Bulbocodium.
excisa, Linn. f. Suppl. 93 = Lapeyrousia juncea.
excisa, Thunb. Diss. Ixia, 19 = Geissorhiza excisa.
Fabricii, D. Delar. Desc. Pl. Nov. 13 = Lapeyrousia anceps.
falcata, Thunb. Diss. Ixia, 19 = Hesperantha falcata.
fallax, *Salisb. Prod.* 34.—Afr. austr.
fastigiata, Lam. Encyc. iii. 337 = Lapeyrousia corymbosa.
fenestrata, Jacq. Coll. iii. 270 = Tritonia hyalina.
filifolia, [Delar. in] Red. Lil. t. 351. f. 2 = Romulea subulata.
filiformis, Vent. Jard. Cels, t. 48 = patens.
fimbriata, Lam. Encyc. iii. 339 = Sparaxis grandiflora.
fistulosa, Andr. Bot. Rep. t. 59 = Hesperantha radiata.
fistulosa, [Ker-Gawl. in] Bot. Mag. t. 583 = Micranthus fistulosus.
flabelliformis, D. Delar. Desc. Pl. Nov. 34 = Babiana plicata.
flabelliformis, Salisb. Prod. 37 = Babiana stricta.
flabellularis, Vahl, Enum. ii. 67 = Tritonia flava.
flaccida, Salisb. Prod. 35 = patens.
flava, Hornem. Hort. Hafn. Suppl. 5 = maculata.
flava, Lam. Illustr. i. 109 = Romulea bulbocodioides.

IXIA :—

flavescens, Eckl. Top. Vers. 26 = polystachya.
flavovirens, *Eckl. l. c.*—Afr. austr.
flexuosa, [Ker-Gawl.] in Bot. Mag. t. 127 = hybrida.
flexuosa, *Linn. Sp. Pl.* ed. II. 51.—Afr. austr.
fragrans, *Jacq. Coll. Suppl.* 9.—Afr. austr.
fruticosa, Linn. f. Suppl. 93 = Witsenia ramosa.
fruticosa, Thunb. Diss. Ixia, n. 1. t. 1. f. 3 = Aristea fruticosa.
fucata, Ker-Gawl. in Bot. Mag. t. 1879 = scariosa.
fugacissima, Linn. f. Suppl. 94 = Galaxia graminea.
fugax, Hornem. Hort. Hafn. 1. 50 = Romulea rosea.
fugax, *Salisb. Prod.* 34.—Afr. austr.
furva, Banks, ex Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 224 = Geissorhiza furva.
fuscocitrina, [DC. in] Red. Lil. t. 86 = maculata.
Galaxia, Linn. f. Suppl. 93 = Galaxia ovata.
Gawleri, Schrad. Neues Journ. iv. (1810) 67 = Lapeyrousia juncea.
geminata, Vahl, Enum. ii. 68 = Geissorhiza obtusata.
geminiflora, *Schrank*, *in Denkschr. Bot. Ges. Regensb.* ii. (1822) 188.—Afr. austr.
gibba, Salisb. Prod. 38 = Tritonia deusta.
gladiata, Linn. f. Suppl. 97 = Babiana gladiata.
gladiolaris, Lam. Encyc. iii. 341 = Tritonia securigera.
gracilis, Salisb. Prod. 37 = capillaris.
grandiflora, D. Delar. Desc. Fl. Nov. 23 = Sparaxis grandiflora.
grandiflora, Pers. Syn. i. 48 = monadelpha.
heterophylla, Willd. Sp. Pl. i. 199 = Lapeyrousia fasciculata.
hexandra, *Schrank*, *in Denkschr. Bot. Ges. Regensb.* ii. (1822) 188.—Afr. austr.
hirsuta, Lam. Ment. i. 37 = Diliztis corymbosa.
hirta, Thunb. Diss. Ixia, 9 = Geissorhiza hirta.
Hochstetteriana, A. Rich. Tent. Fl. Abys. ii. 309 = Hesperantha Pettitiana.
holosericea, Jacq. Hort. Schoenb. i. 9 = Sparaxis grandiflora.
humilis, Thunb. Diss. Ixia, 8 = Geissorhiza humilis.
hyalina, [DC. in] Red. Lil. t. 87 = Tritonia squalida.
hyalina, Linn. f. Suppl. 91 = Tritonia hyalina.
hyalina, Salisb. Prod. 38; Willd. Enum. Hort. Berol. 57 = Tritonia crocata.
hybrida, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 227.—Afr. austr.
imbricata, D. Delar. Desc. Pl. Nov. 17 = Geissorhiza imbricata.
incarnata, Jacq. Coll. Suppl. 12 = scariosa.
inflexa, D. Delar. Desc. Pl. Nov. 15 = Geissorhiza inflexa.
iridifolia, D. Delar. l. c. 24 = Tritonia crocata.
iridifolia, Link, Enum. Hort. Berol. i. 52 = Watsonia Meriana.
juncea, Link, l. c. 50 = Geissorhiza juncea.
lacera, Herb. Linn. ex Roem. & Schult. Syst. i. 396 = Sparaxis grandiflora.
lancea, Jacq. Ic. t. 281; Thunb. Diss. Ixia, 18 = Tritonia scariosa.
Lapeirousia, J. F. Gmel. Syst. 108 = Lapeyrousia compressa.
Larochei, Roem. & Schult. Syst. i. 879 = Geissorhiza Rochenesis.
latifolia, D. Delar. Pl. Nov. 22 = Geissorhiza latifolia.
laurea, Steud. Nom. ed. II. i. 779, 841, sphalm. = *lancea*.
leucantha, *Jacq. Coll. Suppl.* 11.—Afr. austr.
lilacina, *Eckl. Top. Vers.* 27, nomen.—Afr. austr.
Liliago, [DC. in] Red. Lil. t. 109 = Sparaxis grandiflora.
linearis, Jacq. Coll. iv. 188 = Hesperantha angusta.
linearis, *Linn. f. Suppl.* 93; *Thunb. Diss. Ixia*, 12.—Afr. austr.
lineata, *Salisb. Prod.* 35.—Afr. austr.
longiflora, Berg. Desc. Pl. Cap. 7 = Tritonia Rochensis.
longiflora, Lam. Encyc. iii. 343 (Watsoniæ sp. ?).—Afr. austr.
longiflora, [Soland. in] Ait. Hort. Kew. ed. I. i. 58 = paniculata.
longifolia, Jacq. Hort. Vindob. iii. 47 = Hexaglottis longifolia.
lutea, *Baker*, *in Journ. Linn. Soc.* xvi. (1877) 91.—Afr. austr.
lutea, Eckl. Tóp. Verz. 24 = curta.
maculata, *Linn. Sp. Pl.* ed. II. 1664; Mant. ii. 230.—Afr. austr.
magellanica, Lam. Illustr. i. 109; Encyc. iii. 334 = Tapeinia magellanica.

IXIA :—

marginata, [Soland. in] Ait. Hort. Kew. ed. I. i. 59 = Watsonia marginata.
Meriana, Link, Enum. Hort. Berol. i. 52 = Watsonia Meriana.
micrandra, *Baker*, *in Journ. Bot.* xiv. (1876) 367.—Afr. austr.
Millerî, Berg. Desc. Pl. Cap. 8 = maculata.
miniata, [DC. in] Red. Lil. t. 89 = Tritonia deusta.
miniata, Jacq. Hort. Schoenb. i. 10. t. 24 = Tritonia miniata.
minima, *Schrank*, *in Denkschr. Bot. Ges. Regensb.* ii. (1822) 179.—Afr. austr.
minima, Tenore, Fl. Nap. i. 13, in obs.; Syll. 24 = Romulea Columnæ.
minuta, Linn. f. Suppl. 92 = Pasithia hypoxidoides.
monadelpha, *D. Delar. Desc. Pl. Nov.* 22.—Afr. austr.
monadelphia, Burm. f. Fl. Cap. Prod. 1* = prasc. ?
monanthes, D. Delar. Desc. Pl. Nov. 21 = Sparaxis grandiflora.
monantha, Thunb. Fl. Cap. i. 226 = Geissorhiza Rochensis.
mutabilis, *Salisb. Prod.* 34.—Afr. austr.
neglecta, Schult. Mant. i. 279 = Romulea speciosa.
nervosa, Vent. Choix, sub t. 10 = maculata, polystachya.
nigro-albida, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 296 = maculata.
obtusata, Soland. ex [Ker-Gawl.] in Bot. Mag. t. 672 = Geissorhiza obtusata.
ochracea, *Eckl. Top. Vers.* 25.—Afr. austr.
ochroleuca, Sweet, Hort. Brit. ed. II. 509 = maculata.
ochroleuca, Vahl, Enum. ii. 60 = Romulea ochroleuca.
odorata, *Ker-Gawl. Irid. Gen.* 101.—Afr. austr.
ornithogaloides, Lichtst. ex Roem. & Schult. Syst. i. 374 = Trichonema ornithogaloides.
ovata, Burm. f. Fl. Cap. Prod. 1* = Geissorhiza excisa.
ovata, *Sweet, Hort. Brit.* ed. I. 299; *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 296.—Afr. austr.
pallens, Steud. Nom. ed. I. 458 = patens.
pallidiflorens, Eckl. Top. Vers. 25 = curta.
pallido-rosea, Eckl. l. c. 24 = flexuosa.
paniculata, *D. Delar. Desc. Pl. Nov.* 24, t. 1.—Afr. austr.
parviflora, Salisb. Prod. 38 = Romulea Columnæ.
patens, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 59.—Afr. austr.
patens, Willd. Enum. Hort. Berol. 56 = flexuosa.
pectinata, Vahl, Enum. ii. 62 = Tritonia pectinata.
pendula, Thunb. Diss. Ixia, 16 = Dierama ensifolium.
pentandra, Linn. f. Suppl. 92 = Tritonia scillaris.
Petitiana, A. Rich. Tent. Fl. Abyss. ii. 309 = Hesperantha Petitiana.
pholangioides, Roem. & Schult. Syst. i. 385 = Geissorhiza scillaris.
phlogiflora, [Delile, in] Red. Lil. viii. t. 432 = scariosa.
pigmæa, Burm. f. Fl. Cap. Prod. 1* = *pygmæa*.
pilosa, Linn. f. Suppl. 93 = Hesperantha pilosa.
planifolia, Mill. Ic. 160. t. 239. f. 2; ex Roem. & Schult. Syst. i. 400 = Tritonia crocata.
planiscapa, [Soland. in] Ait. Hort. Kew. ed. I. i. 59 = Micranthus plantagineus.
pliciata, Linn. Cent. Pl. ii. 4; Amœn. Acad. iv. 300 = Babiana stricta.
polystachya, [DC. in] Red. Lil. t. 126 = flexuosa.
polystachya, Jacq. Coll. iii. 269 = Tritonia scillaris.
polystachya, Ker-Gawl. in Bot. Mag. t. 629 = Watsonia petiana.
polystachya, *Linn. Sp. Pl.* ed. II. 51.—Afr. austr.
praxina, Soland. ex Baker, in Journ. Linn. Soc. xvi. (1877) 92 = viridiflora.
propinqua, *Salisb. Prod.* 38.—Afr. austr.
pudica, Banks, ex Ker-Gawl. in Bot. Mag. sub t. 575 = Romulea pudica.
pulcherrima, Eckl. Top. Vers. 24 = speciosa.
pulchra, Salisb. in Trans. Hort. Soc. i. (1812) 320 = viridiflora.
pumila, Forst. f. in Comm. Gotting. ix. (1789) t. 2 = Tapeinia magellanica.
punctata, Andr. Bot. Rep. t. 177 = Watsonia punctata.
punicea, *Eckl. Top. Vers.* 24.—Afr. austr.
punicea, Jacq. Coll. Suppl. 12; Ic. Pl. Rar. t. 287 = Babiana stricta.
punicea, Spreng. Neue Entdeck. i. 254 = patens.
purpurascens, Tenore, Fl. Nap. i. 13. t. 3 = Romulea purpurascens.
purpurea, Jacq. Coll. iii. 268 = Babiana stricta.
purpurea, Lam. Encyc. iii. 341 = Tritonia crocata.
pusilla, Andr. Bot. Rep. t. 945 = Geissorhiza secunda.
pygmæa, Roem. & Schult. Syst. i. 403 = Babiana nana.

IXIA :—

pyramidalis, Lam. Encyc. iii. 334 = Lapeyrousia anceps.
quadrangula, D. Delar. Desc. Pl. Nov. 16 = Geissorhiza quadrangula.
quadrangularis, Burm. f. Fl. Cap. Prod. 1* = prasc.
Quartiniana, A. Rich. Tent. Fl. Abyss. ii. 310 = Acidanthera bicolor.
radians, Thunb. in Hoffm. Phytog. Blastt. i. (1803) 3 = Geissorhiza Rochensis.
radiata, Jacq. Coll. Suppl. 15 = Hesperantha radiata.
ramiflora, Tenore, Syll. 25 = Romulea ramiflora.
ramosa, [Ker-Gawl. in] Bot. Mag. sub t. 549 = Geissorhiza secunda.
rapunculoides, [Delile in] Red. Lil. viii. t. 431 = scariosa.
recurva, [DC. in] l. c. t. 251. f. 1 = Romulea bulbocodioides.
recurva, Vahl, Enum. ii. 56 = Hesperantha radiata.
recurvifolia, Poir. Encyc. Suppl. iii. 201 = Trichonema recurvifolium.
reflexa, Andr. Bot. Rep. t. 14 = Tritonia scillaris.
reflexa, Thunb. Fl. Cap. ed. II. 55 = Romulea bulbocodioides.
reticularis, *Salisb. Prod.* 37.—Afr. austr.
reticulata, Thunb. Fl. Cap. ed Schult. 60 = Tritonia lineata.
retusa, Salisb. Prod. 35 = Tritonia scillaris.
Rochensis, [Ker-Gawl.] in Bot Mag. t. 598 = Geissorhiza Rochensis.
rosea, Linn. Syst. ed. XII. 75 = Romulea rosea.
rotata, [Ker-Gawl.] Andr. Recens. Pl. Nov. Bot. Rep. 3 = Tritonia scillaris.
rubrocyanea, Jacq. Coll. iii. 268 = Babiana stricta.
scariosa, *Thunb. Fl. Cap.* i. 248.—Afr. austr.
Scaptrum, Hort. ex Baker, in Journ. Linn. Soc. xvi. (1877) 158 = Watsonia marginata.
scillaris, Linn. Sp. Pl. ed. II. 52 = Tritonia scillaris.
scillaris, Mill. Gard. Dict. ed. VIII. n. 3 = Babiana stricta.
scillaris, Thunb. Diss. Ixia, 13 = Geissorhiza secunda.
secunda, *Berg. Desc. Pl. Cap.* 6.—Afr. austr.
secunda, D. Delar. Desc. Pl. Nov. 17 = Geissorhiza secunda.
serotina, [Masson] Salisb. Prod. 34 = polystachya.
setacea, Thunb. Diss. Ixia, 18 = Geissorhiza setacea.
similis, Salisb. Prod. 36 = Tritonia squalida.
socialis, *Salisb. l. c.* 35.—Afr. austr.
sordida, *Hornem. Hort. Hafn. Suppl.* 6.—Afr. austr.
Sparmannii, Schult. Mant. i. 285 (= *Meristostigma Sparmanni*).—Afr. austr.
sparsa, Mill. Gard. Dict. ed. VIII. n. 7 = Sparaxis grandiflora.
spathacea, [Soland.] ex Ker-Gawl. in Bot. Mag. sub t. 1468 = Gladiolus trichonemifolius.
speciosa, *Andr. Bot. Rep.* t. 156.—Afr. austr.
spectabilis, Salisb. Prod. 35 = viridiflora.
spicata, Andr. Bot. Rep. i. t. 29 = maculata.
spicata, Burm. f. Fl. Cap. Prod. 1* = Lapeyrousia juncea.
spicata, Willd. Sp. Pl. i. 200 = Watsonia spicata
Sprengeliana, Schult. Mant. i. 283 = patens.
squalida, [Soland. in] Ait. Hort. Kew. ed. I. i. 61 = Tritonia squalida.
squalida, Thunb. in Hoffm. Phytogr. Blastt. i. (1803) 4 = Tritonia lineata.
stellata, *Hill, Veg. Syst.* x. 3.—Afr. austr.
stellata, Hort. ex Roem. & Schult. Syst. i. 401 = Sparaxis tricolor.
stellata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 296.—Afr. austr.
striata, Vahl, Enum. ii. 65 = scariosa.
subluta, Lam. Encyc. iii. 335 = Romulea sublutea.
tardiflora, *Salisb. Prod.* 38.—Afr. austr.
tenella, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 296 = flexuosa.
tenuiflora, Vahl, Enum. ii. 66 = Tritonia longiflora.
tenuifolia, Vahl, l. c. 62 = linearis.
tereticlia, Banks, ex [Ker-Gawl.in] Bot. Mag. t. 528 = Watsonia spicata.
thrysiflora, D. Delar. Desc. Pl. Nov. 20 = flexuosa.
Thunbergii, Roem. & Schult. Syst. i. 391 = Tritonia lineata.
tortuosa, Licht. ex Roem. & Schult. l. c. 375 = Romulea tortuosa.
tricolor, Schneev. Ic. t. 20 = Sparaxis tricolor.
triticea, Burm. f. Fl. Cap. Prod. 1* = Micranthus plantagineus.
tubulosa, Burm. f. l. c. = Babiana tubiflora.

IXIA:—
umbellata, Burm. f. Fl. Cap. Prod. 2 = Dilatris corymbosa.
undulata, Burm. f. l. c. 1° = Tritonia undulata.
undulata, Salisb. Prod. 37 = crispa.
unifiora, Linn. Mant. i. 27 = Sparaxis grandiflora.
variegata, Banks, ex Schult. Mant. i. 225 = monadelpha.
venosa, *Link, Enum. Hort. Berol.* i. 51.—Afr. austr.
vernalis, Salisb. Prod. 32 = Crocus vernus.
verrucosa, B. Vog. in Trew, Pl. Rar. t. 24. f. 2 = Geissorhiza excisa.
villosa, [Soland. in] Ait. Hort. Kew. ed. I. i. 58 = Babiana stricta.
villosula, J. F. Gmel. Syst. 110 = Babiana stricta.
violacea, Banks, ex Ker-Gawl. in Bot. Mag. t. 598 = Geissorhiza Rochensis.
virgata, Willd. Sp. Pl. i. 203 = Hexaglottis virgata.
virginea, Soland. ex Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 225 = Hesperantha angusta.
viridiflora, *Lam. Encyc.* iii. 340.—Afr. austr.
viridis, Thunb. Fl. Cap. i. 242 = viridiflora.
vitellina, Eckl. Top. Verz. 25 = curta.
Xiphidium, Loefl. ex Roem. & Schult. Syst. i. 487 = Xiphidium floribundum.

IXIA, Mnhl. ex Spreng. Syst. i. 171 (1825) = **Hydrilla**, Rich. (Hydroch.).
aquatica, Mnhl. l. c. = H. verticillata.

IXIANTHES, Benth. in Hook. Comp. Bot. Mag. ii. (1836) 53. *SCROPHULARINEÆ*, Benth. & Hook f. ii. 937.
retusoides, *Benth. l. c.*—Afr. austr.

IXIANTHUS, Reichb. Nom. 116 (1841) = prec.

IXIAUCHENUS, Less. Syn. Comp. 198 (1832) = IXAUCHENUS, Cass. = **Lagenophora**, Cass. (Compos.).

IXIDIUM, Eichl. in Mart. Fl. Bras. v. II. 130. t. 31 (1868) = **Bremolepis**, Griseb. (Loranthac.).
Schottii, Eichl. l. c. 131.—Am. trop.
Wrightii, Eichl. l. c. 132.—Am. trop.

IXINA, Rafin. Act. Journ. 144 (1832) = **Krameria**, Linn. (Polygal.).

IXINE, Hill, Veg. Syst. iv. 17 (1762) = **Cnicus**, Tourn. (Compos.).
arvensis, Hill, Herb. Brit. i. 74 = C. arvensis.
Casabonæ, Hill, Hort. Kew. 51 = C. Casabonæ.
tuberosa, Hill, l. c. = C. tuberosus.

IXIOCHLAMYS, F. Muell. & Sond. in Linnæa, xxv. (1852) 466 = **Podocoma**, Cass. (Compos.).
cuneifolia, F. Muell. & Sond. l. c. = P. cuneifolia.

IXIOLÆNA, Benth. in Enum. Pl. Hueg. 66 (1837). *COMPOSITÆ*, Benth. & Hook. f. ii. 315.
brevicompta, *F. Muell. Fragm.* i. 53.—Austral.
chrysantha, Steetz, in Lehm. Preiss. i. 459 = Podolhica chrysantha.
leptolepis, *Benth. Fl. Austral.* iii. 597.—Austral.
supina, *F. Muell. in Trans. Vict. Inst.* i. (1855) 87.—Austral.
tomentosa, *Sond. & Muell. in Linnæa*, xxv. (1852) 504.—Austral.
viscosa, *Benth. in Enum. Pl. Hueg.* 66.—Austral.

IXIOLIRION, Fisch. ex Herb. App. 37 (1821). *AMARYLLIDEÆ*, Benth. & Hook. f. iii. 735.
KOLPAKOWSKIA, Regel, in Gartenfl. xxvii. (1878) 294. t. 953.
Kolpakowskianum, *Regel, in Act. Hort. Petrop.* vi. (1879) 491.—Turkestan.
Ledebouri, Fisch. & Mey. in Bong. & Mey. Suppl. Fl. Alt. n. 298 = montanum.
montanum, *Herb. App.* 37.—Syria; Persia.
montanum, Kunth, Enum. Pl. v. 817 = Pallasii.
Pallasii, *Fisch. & Mey. ex Ledeb. Fl. Ross.* iv. 116.—Reg. Casp.; Afghan.
tataricum, Herb. App. 37 = montanum.
tataricum, Schult. f. Syst. vii. 752 = Pallasii.

IXIONANTHES, Endl. Gen. 1055 (1840) = **Ixonanthes**, Jack (*Linnæa*).

IXIOSPORUM, F. Muell. Fragm. ii. 76 (1860) = **Citriobatea**, A. Cunn. (Plitosp.).
spinescens, F. Muell. l. c. = C. pauciflora.

IXODIA, R. Br. in Ait. Hort. Kew. ed. II. iv. 517 (1813); Bot. Mag. t. 1584 (1813). *COMPOSITÆ*, Benth. & Hook. f. ii. 318.
schilleoides, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 517.—Austral.

achilleoides, Schlecht. in Linnæa, xx. (1847) 593 = schilleoides.
Ammobium, Spreng. Syst. iv. Cur. Post. 300 = Ammobium alatum.
ptarmicoides, F. Muell. in Linnæa, xxv. (1852) 405 = schilleoides.

IXODIA, Soland. ex DC. Syst. ii. 87 (1821) = **Brassenia**, Schreb. (Nymph.).

IXONANTHES, Jack, in Malay. Misc. ii. (1822) VII. 51. *LINEÆ*, Benth. & Hook. f. i. 245.
BREWSTERA, M. Roem. Syn. i. Hesper. 132, 141 (1846).
EMMENANTHUS, Hook. & Arn. Bot. Beech. Voy. 317 (1836).
PIEROTIA, Blume, Mus. Bot. Lugd. Bat. i. 179 (1850).
chinensis, *Champ. in Proc. Linn. Soc.* ii. (1850) 100.—Hongkong.
cuneata, *Miq. Fl. Ind. Bat. Suppl.* 484.—Sumatra.
dodecandra, Griff. Pl. Cantor. 12 = Icosandra.
icosandra, *Jack, in Malay. Misc.* ii. (1822) VII. 53.—Malaya.
khasiana, *Hook. f. Fl. Brit. Ind.* i. 416.—Reg. Himal.
lucida, Blume, Mus. Bot. Lugd. Bat. i. 396 = icosandra.
obovata, *Hook. f. Fl. Brit. Ind.* i. 417.—Malayas.
petiolaria, *Blume, Mus. Bot. Lugd. Bat.* i. 396.—Malaya.
reticulata, *Jack, in Malay. Misc.* ii. (1822) VII. 51.—Malaya.

IXOPHORUS, Schlecht. in Linnæa, xxxi. (1861–62) 420 = **Setaria**, Beauv. (Gramin.).
Schiedeanus, Schlecht. l. c. 421 = S. Schiedeana.
unisetus, Schlecht. l. c. = S. uniseta.

IXORA, Linn. Syst. ed. I (1735). *RUBIACEÆ*, Benth. & Hook. f. ii. 113.
EMSETIA, Rafin. Sylva Tellur. 12 (1838).
EUMACHIA, DC. Prod. iv. 478 (1830).
PANCHEZIA, Montr. in Mém. Acad. Lyon, x. (1860) 223.
SCHETTI, Adans. Fam. ii. 146 (1763).
SIDERODENDRUM, Schreb. Gen. 71 (1789).
SIDEROXYLOIDES, Jacq. Select. Am. 19. t. 175. f. 9 (1763).
abyssinica, Oliver, in Trans. Linn. Soc. xxix. (1878) 87 = Pavetta abyssinica, Oliveriana.
acuminata, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 883.—Reg. Himal.
acuminata, Thw. Enum. Pl. Zeyl. 155 = Thwaitesii.
acuminatissima, *Muell. Arg. in Flora*, lvii. (1875) 454.—Bras.
acutifolia, Reinw. ex Miq. Fl. Ind. Bat. ii. 272 = Pavetta nigricans.
affinis, Wall. Cat. n. 6144 = nigricans.
africana, *Beauv. ex A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 117.—Afr. trop.
alba, *Linn. Sp. Pl.* 110.—Ind. or.
alba, Roxb. Hort. Beng. [10]; Fl. Ind. i. 879 = stricta.
Alba, Sm. ex Wight & Arn. Prod. 401 = Stylocoryne Webera.
alternifolia, Jacq. Enum. Pl. Carib. 12 = Cestrum nocturnum.
amabilis, *Hort. ex Flor. Mag.* x. (1871) tt. 549, 550.—Hab.?
amara, Steud. Nom. ed. II. i. 842, sphalm. = amoena.
amboinica, *DC. Prod.* iv. 487.—Ins. Amboin.
americana, Jacq. Hort. Schoenb. iii. 4. t. 257 = Bouvardia triphylla.
americana, Linn. Syst. ed. X. 898 = Bouvardia chlorantha, Famarca odoratissima.
amoena, *Wall. Cat.* n. 6121.—Burma; Malaya.
amplifolia, *A. Gray, in Proc. Am. Acad.* iv. (1860) 40.—Ins. Samoa.
angulata, Benth. ex Benth. & Hook. Gen. ii. 114 = Appunia angulata.
angustifolia, Lam. Encyc. iii. 45 = Pavetta angustifolia.
arborea, Lodd. Bot. Cab. t. 609 = coccinea.

IXORA:—
arborea, Roxb. ex Sm. in Rees, Cycl. xix. n. 5 = parviflora.
arborescens, Hask. in Retzia, i. 22 = grandifolia.
arenosa, Poir. Encyc. Suppl. iii. 309 = Pavetta arenosa.
arguta, R. Br. in Wall. Cat. n. 6137 = nigricans.
Arnottiana, Miq. ex Hook. f. Fl. Brit. Ind. iii. 142 = brachiata.
attenuata, Wall. Cat. n. 6164 = Chasalia curviflora.
bahiensis, *Benth. in Linnæa*, xxiii. (1850) 447.—Bras.
Bandhuca, Roxb. Hort. Beng. [10]; Fl. Ind. i. 876 = coccinea.
barbata, *Roxb. ll. cc.* [10]; 886.—Ins. Andam.
Beckleri, *Benth. Fl. Austral.* iii. 413.—Austral.
Benthamiana, *Muell. Arg. in Flora*, lvii. (1875) 455.—Bras.
Berseiana, *Baill. Adansonia*, xii. (1876–79) 218.—Madag.
blanda, Ker-Gawl. in Bot. Reg. t. 100 = chinensis.
Blumei, *Zoll. & Mor. Syst. Verz. Zoll.* 66.—Java.
brachiata, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 881.—Ind. or.
brachypoda, DC. Prod. iv. 488 = laxiflora.
brachyrhipha, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 165.—Afr. trop.
bracteolaris, *Muell. Arg. in Flora*, lvii. (1875) 458.—Bras.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 216.—Burma.
brevifiora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 165.—Afr. trop.
brevifolia, *Benth. in Linnæa*, xxiii. (1850) 448.—Bras.
brunnescens, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 217.—Ins. Andam.
Brunonis, *Wall. Cat.* n. 6136.—Burma; Malaya.
Burbidgei, *Hort. Veitch.—Cf. Gard. Chron.* (1881) ii. 231.—Hab.?
Burchellitana, *Muell. Arg. in Flora*, lvii. (1875) 454.—Bras.
buxifolia, Spreng. Syst. i. 409 = Rondeletia buxifolia.
buxina, *Baill. Adansonia*, xii. (1876–79) 294.—N. Caled.
cæfra, Poir. Encyc. Suppl. iii. 309 = Pavetta cæfra.
cæfra, Sond. in Linnæa, xxiii. (1850) 50 = Oxyanthus natalensis.
calycina, *Thw. Enum. Pl. Zeyl.* 155.—Zeylan.
Canjia, Wall. Cat. n. 6130 = undulata.
Carewi, *Horne, ex Baker, in Journ. Linn. Soc. xx.* (1883) 364.—Ins. Fiji.
carnea, *Benth. & Hook. f. Gen.* ii. 114.—Ins. Amicor.
Casei, *Hance, in Walp. Ann.* ii. 754.—China.
cauliflora, *Montr. in Mém. Acad. Lyon*, x. (1860) 224.—N. Caled.
chinensis, *Lam. Encyc.* iii. 344.—China.
coccinea, Comm. ex Lam. Encyc. ii. 355 = Embothrium coccineum.
coccinea, Lour. Bot. Mag. t. 169 = chinensis.
coccinea, *Linn. Sp. Pl.* 110.—As. trop.
Colebrookii, Voigt, Hort. Suburb. Calc. 391 = lanceolaria.
compactiflora, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 215.—Burma.
concinna, *R. Br. in Wall. Cat.* n. 6149.—Malaya.
congesta, *Roxb. Hort. Beng.* 83; *Fl. Ind.* i. 887.—Burma; Malaya.
Conjia, Buch.-Ham. ex Steud. Nom. ed. II. i. 842 = *Canjia*.
cordifolia, [Moç. & Sesse,] ex DC. Prod. iv. 486 = Bouvardia cordifolia.
coriacea, R. Br. in Wall. Cat. n. 6151 = Webera grandifolia.
corymbosa, Heyne, in Wall. l. c. n. 6155 = polyantha.
corymbosa, Mirb. in Hist. Nat. Pl. ed. II. xiv. 57 = Lonicera corymbosa.
crocata, Lindl. Bot. Reg. t. 782 = chinensis.
Cuminglana, *Vidal, Phan. Cuming. Philipp.* 183.—Ins. Philipp.
cuneifolia, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 880.—Ind. or.; Burma.
cyathocsperma, Wall. Cat. sub n. 6121 = amoena.
Cyphori × —Cf. Gard. Chron. (1874) ii. 49.
Dallachyana, F. Muell. Census, 75 = Webera Dallachiana.
decipiens, DC. Prod. iv. 488 = parviflora.
densa, R. Br. in Wall. Cat. n. 6150 = nigricans?
densiflora, *Muell. Arg. in Flora*, lvii. (1875) 455.—Bras.

IXORA:—
dimorphophylla, *Muell. Arg. in Flora*, lviii. (1875) 454.—Bras.
divaricifolia, *R. Br. in Wall. Cat.* n. 6146.—Burma.
dubia, Schult. Mant. iii. 121 = chinensis.
Duffii, T. Moore, Flor. & Pomol. (1878) 76 = macrothyrsa.
elliptica, R. Br. in Wall. Cat. n. 6158 = grandifolia.
elongata, *Heyne, ex Wall. Cat.* n. 6131.—Ind. or.
eminensis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 413.—Madag.
erianthe, *A. Grey, in Proc. Am. Acad.* iv. (1860) 40.—Bras.
erubescens, *Wall. Cat.* n. 6148.—Burma.
erythrocrypta, *Wall. ex Voigt, Hort. Suburb. Calc.* 393.—Hab.!
expandens, F. Muell. Census, 75 = Webera expandens.
fasciculata, Sw. Prod. Veg. Ind. Occ. 80 = Chomelia fasciculata.
ferrea, *Benth. in Linnaea*, xxiii. (1850) 447.—Ind. occ. Finlaysoniana, *Wall. Cat.* n. 6166.—Siam!
flammea, Salisb. Prod. 92 = stricta.
flava, *Lodd. ex G. Don, in Loud. Hort. Brit.* 49, nomen.—Ind. or.
floribunda, *Griseb. Cat. Pl. Cub.* 134.—Cuba.
foliosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 166.—Afr. trop.
formosa ×.—Cf. Gard. Chron. (1878) i. 10; (1881) ii. 336.
fragrans, A. Gray, in Proc. Am. Acad. iv. (1860) 39 = Cephaelis fragrans.
Francavillana, *Muell. Arg. in Flora*, lviii. (1875) 454.—Bras.
fulgens, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 373.—Burma; Malaya.
fulgens, Wall. Cat. n. 6512 B. = congesta.
Gardneriana, *Benth. in Linnaea*, xxiii. (1850) 448.—Bras.
glandulosa, Blanco, Fl. Filip. ed. I. 61 (= *Canthium confertum*).—Ins. Philipp.
glaucina, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 146.—Burma.
Glaziovi, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 460.—Bras.
gracilliflora, *Benth. in Linnaea*, xxiii. (1850) 448.—Guiana.
gracilis, *R. Br. ex Fawc. in Forbes, Nat. Wand. East. Archip.* 508.—Ins. Timor.
grandiflora, [Ker-Gawl.] in Bot. Reg. t. 154 = coccinea.
grandiflora, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 139 = Notoniana.
grandifolia, R. Br. in Wall. Cat. n. 6184 = Webera grandifolia.
grandifolia, *Muell. Arg. in Flora*, lviii. (1875) 454.—Bras.
grandifolia, *Zoll. & Mor. Syst. Verz. Zoll.* 65.—Malaya.
grandis, Miq. ex Hook. f. Fl. Brit. Ind. iii. 140 = polyantha.
Griffithii, Hook. Bot. Mag. t. 4325 = congesta.
guianensis, Spreng. Syst. i. 409 = Coffea guianensis.
guianensis, *Benth. in Hook. Niger Fl.* 414.—Afr. trop.
Helferi, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 216 = Webera Helferi.
heterodoxa, *Muell. Arg. in Flora*, lviii. (1875) 454.—Bras.
heterophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 68.—Bras.
incarnata, DC. Prod. iv. 486 = coccinea.
incarnata, Roxb. ex Sm. in Rees. Cycl. xix. n. 4 = chinensis.
indica, Baill. Hist. Pl. vii. 378 = Pavetta indica.
inundata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 166.—Afr. trop.
javanica, *DC. Prod.* iv. 487.—Java.
Johnsoni, *Hook. f. Fl. Brit. Ind.* iii. 136.—Ind. or.
Jookei, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 262.—Ins. Fiji.
jucunda, *Thw. Enum. Pl. Zeyl.* 155.—Zeylan.
Kingiana, *Hook. f. Fl. Brit. Ind.* iii. 140.—Malacca.
Klenderiana, F. Muell. Fragm. v. 16 = timorensis.
Korthalsiana, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 146.—Malaya.
Kurzeana, Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind. xxvii. (1864) 100 = grandifolia.
lanceolaris, *Colebr. in Roxb. Fl. Ind. ed. Carey*, i. 397.—Ind. or.
lanceolata, Lam. Encyc. iii. 346 = fulgens.
lanciflora, *Sm. in Rees, Cycl.* xix. n. 3.—Afr. trop.
lancifolia, Heyne, in Wall. Cat. n. 6148 = lanceolaris. Lobbii, *Loud. Encyc. Pl. Suppl.* ii. 1543.—Hab.?
longiflora, Hook. Bot. Mag. t. 4482, in syn. sphalm. = laxiflora.
longiflora, Poir. Encyc. Suppl. iii. 308 = Pavetta longiflora.
longifolia, G. Don, Gen. Syst. iii. 573 = Webera longifolia.
longifolia, Sm. in Rees, Cycl. xix. n. 3 = fulgens.
longifolia, Walp. Ann. ii. 756, sphalm. = Pavetta longiflora.
longipes, *Zoll. & Mor. Syst. Verz. Zoll.* 65.—Java.
Lourviri, G. Don, Gen. Syst. iii. 573 = alba!
lucida, *R. Br. in Wall. Cat.* n. 6135.—Penang.
macrophylla, *Bartl. ex DC. Prod.* iv. 487.—Ins. Philipp.
macrophylla, R. Br. in Wall. Cat. n. 6165 = Webera longifolia.
macrostephon, Kurz, in Journ. Bot. xiii. (1875) 327 = grandifolia.
macrothyrsa, [*Trijsm. & Binn.*].—*Cf. Gard. Chron.* (1884) ii. 257.—Malaya.
manila, Blanco, Fl. Filip. ed. I. 60 = Scyphiphora hydrophyllacea.
mauritiana, Boj. Hort. Maurit. 189 = Myonima multiflora.
maxima, *Sesm. Fl. Vit.* 134.—Ins. Fiji.
membranacea, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 458.—Bras.
memecylifolia, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 216 = erubescens.
merguensis, *Hook. f. Fl. Brit. Ind.* iii. 140.—Burma.
micrantha, Heyne, ex Wall. Cat. sub n. 6142 = brachiata.
microcrithe, Roem. & Schult. Syst. iii. 179 = Myonima multiflora.
minutiflora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 167.—Afr. trop.
modesta, *Hiern, l. c.* 167.—Afr. trop.
montana, *Lour. Fl. Cochinch.* 76.—Cochinch.
montana, Miq. ex Hook. f. Fl. Brit. Ind. iii. 40 = Wendlandia Notoniana.
multiflora, Sw. Prod. Veg. Ind. Occ. 80 = Myonima multiflora.
mutabilis, Reinw. ex Miq. Fl. Ind. Bat. ii. 268 = Pavetta indica.
nerifolia, *Yach, in Malay. Misc.* ii. (1822) vii. 89.—Sumatra.
nigricans, *R. Br. in Wall. Cat.* n. 6154.—Ind. or.; Malaya.
nigricans, Zoll. ex Miq. Fl. Ind. Bat. ii. 271 = Pavetta paludosa.
nitida, Schum. & Thonn. Beskr. Guin. Pl. 77 = Pavetta Baconis.
Nondana, *Wall. Cat.* n. 6132.—Ind. or.
novemnervia, *Lour. Fl. Cochinch.* 76.—Cochinch.
Nunnyapata, Roxb. ex Wight & Arn. Prod. 421 = Pavetta indica.
oblonga, *R. Br. in Wall. Cat.* n. 6147 A.—Burma; Malaya.
obovata, Heyne, in Roth, Nov. Pl. Sp. 90 = coccinea.
obscura, *Muell. Arg. in Flora*, lviii. (1875) 455.—Bras.
obtusata, Miq. ex Hook. f. Fl. Brit. Ind. iii. 142 = brachiata.
occidentalis, Forsk. Fl. Aegypt. Arab. p. cv = Pavetta longiflora.
occidentalis, *Linn. Syst. ed.* X. 893.—Ind. occ.
odoratа, *Roxb. Hort. Mag.* t. 4191.—Madag.
odorata, Spreng. Syst. i. 409 = Coffea odorata.
odoratissima, Klotzsch, in Abh. Acad. Berl. (1857) 227 = laxiflora.
opaca, *R. Br. in Wall. Cat.* n. 6141.—Burma; Malaya.
ovariensis, Poir. Encyc. Suppl. iii. 307 = Pavetta ovariensis.
oxyphylla, Wall. Cat. n. 6159 = subsessilis.
pallida, Reinw. ex Miq. Fl. Ind. Bat. ii. 269 = stricta.
paludosa, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 146.—Malaya.
paniculata, Lam. Encyc. iii. 344 = Pavetta indica.
paniculata, Retzi. in Wall. Cat. n. 6143 = oblonga!
parenensis, *Muell. Arg. in Flora*, lviii. (1875) 454.—Bras.
parasitica, Poir. Encyc. Suppl. iii. 309 = Pavetta parasitica.
parviflora, Lam. Illustr. n. 1473. t. 66. f. 2 = Myonima multiflora.
parviflora, *Vahl, Symb. Bot.* iii. 11. t. 52.—Ind. or.
pauciflora, DC. Prod. iv. 487 = Pavetta pauciflora.
Pavetta, Andr. Bot. Rep. t. 75 = parviflora.
Pavetta, Roxb. Hort. Beng. [10]; Fl. Ind. i. 385 = Pavetta indica.
pavetoides, Dalz. in Hook. Kew Journ. iii. (1851) 121 = elongata.
pelagica, *Seem. Fl. Vit.* 134.—Ins. Fiji.
pendula, *Yach, in Malay. Misc.* i. (1820) v. 11.—Malaya.
pentamera, *Benth. Fl. Austral.* iii. 416.—Austral.
pentandra, West, ex DC. Prod. iv. 486 = Guettarda parviflora.
podocarpa, *Benth. ex Benth. & Hook. Gen.* ii. 113.—Bras.
polyantha, *Wight, Ic.* t. 1066.—Ind. or.
propinqua, R. Br. in Wall. Cat. n. 6130 = coccinea.
pruinosa, *Baill. Adansonia*, xii. (1876–79) 217.—Ins. Mascar.
puberula, Wall. Cat. n. 6145 = cuneifolia.
pubescens, *Willd. ex Schult. Mant.* iii. 126.—Bras.
pulsiflora, *DC. Prod.* iv. 489.—Ind. or.
pudica, *Baker, Fl. Maurit.* 151.—Ins. Seychelles.
purpurea, Flech. ex Loud. Hort. Brit. 49 = coccinea.
quinquefida, *R. Br. ex Fawc. in Forbes, Nat. Wand. East. Archip.* 508.—Ins. Timor.
radiata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 163.—Afr. trop.
recurva, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 146.—Burma.
Richardiana, *Muell. Arg. in Flora*, lviii. (1875) 454.—Bras.
Riedeliana, *Muell. Arg. l. c.* 455.—Bras.
riparia, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 164.—Afr. trop.
rosacea, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 128.—Java.
rosea, Sims, in Bot. Mag. t. 2438 = chinensis.
rosea, Wall. in Roxb. Fl. Ind. ed. Carey, i. 398 = cuneifolia.
roxella, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 217 = grandifolia.
rufa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 62.—Bras.
rugosula, *Wall. Cat.* n. 6156.—Burma.
rugulosa, Steud. Nom. ed. II. i. 843 = praec.
salicifolia, DC. Prod. iv. 487 = fulgens.
samoënsis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 40.—Ins. Samoa.
Sandankwa, Hort. Bogor. ex Hassk. Retzia, i. 87 = Viburnum Sandankwa.
Schomburgkiana, *Benth. in Linnaea*, xxiii. (1850) 448.—Guian.
Schottiana, *Muell. Arg. in Flora*, lviii. (1875) 455.—Bras.
serices, *Baker, Fl. Maurit.* 151.—Ins. Seychelles.
sessiliflora, *Baill. Adansonia*, xii. (1876–79) 219.—Ins. Nossi-bé.
sessilifolia, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 216 = rugosula.
sessilifolia, Spreng. Syst. i. 409 = Faramea sessilifolia.
sessilis, Hemsl. Nom. i. 427, sphalm. = subsessilis.
siamensis, *Wall. Cat.* n. 6175.—Siam.
Seyanxii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 166.—Afr. trop.
speciosa, Willd. Enum. Hort. Berol. 157 = stricta.
spectabilis, *Wall. Cat.* n. 6138.—Burma.
spinosa, Lam. Encyc. iii. 344 = Chomelia spinosa.
splendida, *André, in Illustr. Hortic.* xxix. (1882) t. 463.—Hab.?
Spruceana, *Muell. Arg. in Flora*, lviii. (1875) 454.—Bras.
stipulata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 65.—Bras.
Storckii, *Seem. Fl. Vit.* 134.—Ins. Fiji.
stricta, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 379.—Ind. or.; Malaya.
stricta, Wight, ex Wall. Cat. sub n. 6161 = Notoniana.
subcapitata, *Wall. l. c.* n. 6130.—Ind. or.
subsessilis, *Wall. l. c.* n. 6129.—Reg. Himal.
sumatrensis, *Zoll. & Mor. Syst. Verz. Zoll.* 65.—Sumatra.
syringaefolia, *Muell. Arg. in Flora*, lviii. (1875) 455.—Bras.

IXORA:—
tamulifora, *Roxb. Hort. Beng.* 88; *Fl. Ind.* i. 387.—Ins. Molucc.
ternifolia, Cav. Ic. iv. t. 308 = Bouvardia triphylla.
ternifolia, Hook. f. ex Oliver, in Trans. Linn. Soc. xxix. (1873) 86 = Pavetta ternifolia.
Thomsoni, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 164.—Afr. trop.
Thwaitesiana, F. Muell. Fragm. ii. 132 = Randia densiflora.
Thwaitesii, *Hook. f. Fl. Brit. Ind.* iii. 138.—Zeylan.
thyrsiflora, Polr. Encyc. Suppl. iii. 208 = Pavetta cafra.
thyrsoides, *Muell. Arg. in Flora*, lviii. (1875) 455.—Bras.
timorensis, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 418.—Ins. Timor.
tinifolia, *Benth. in Linnaea*, xxiii. (1850) 447.—Ind. occ.?
tomentosa, Roxb. Hort. Beng. [10]; Fl. Ind. i. 386 = Pavetta indica.
tomentosa, Thw. Enum. Pl. Zeyl. 156 = Pavetta Gleniei.
triflora, *R. Br. ex Benth. Fl. Austral.* iii. 416.—Austral.
trifiora, Seem. Fl. Vit. 138 = Pavetta trifiora.
truncata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 459.—Bras.
undulata, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 385.—Ind. or.
velutina, Wall. Cat. n. 6174 = Pavetta indica.
venulosa, *Benth. in Linnaea*, xxiii. (1850) 446.—Bras.
vermicolor, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 135 = Hellcteres Isora.
verticillata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 59.—Bras.
villosa, Polr. Encyc. Suppl. iii. 208 = Pavetta villosa.
villosa, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 383.—Ind. or.; Malaya.
violacea, *Lour. Fl. Cochinch.* 76.—Cochinch.
vitiensis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 40.—Ins. Fiji.
Wallichii, Wight, ex Hook. f. Fl. Brit. Ind. iii. 138 = lanceolaria.
Wallichii, *Wight & Arn. Prod.* 428 = canellfolia!
Warmingii, *Muell. Arg. in Flora*, lviii. (1875) 455.—Bras.
weberaefolia, Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 133 = Webera Kursii.
Wightiana, Wall. Cat. n. 6161 = Notoniana.

IZQUIERDIA, Ruiz & Pav. Prod. 140 (1794) = **Myrsine** †, Linn.
aggregata, Ruiz & Pav. Syst. Veg. 273 (= *Samaria Izquierdia*).—Peruv.

J

JABOROSA, Juss. Gen. 125 (1789). *SOLANACEAE*, Benth. & Hook. f. ii. 868.
DORVSTIGMA, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 347.
LONCHESTIGMA, Dun. in DC. Prod. xiii. i. 476 (1848).
Bergii, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 263.—Reg. Argent.
bonariensis, j. F. Gmel. Syst. 880 = integrifolia.
casalcosna, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 347.—Chili.
crispa, *Benth. & Hook. f. Gen.* ii. 898.—Bolivia.
decorosa, *Miers, Trav. Chil.* ii. 531.—Chili.
integrifolia, *Lam. Encyc.* iii. 120.—Reg. Argent.
laciniata, *Miers, Trav. Chil.* ii. 531.—Parva.
longiflora, *Mog. & Sesse, ex Dun. in DC. Prod.* xiii. i. 481.—Mexic.
magellanica, *Benth. & Hook. f. Gen.* ii. 898.—Reg. Magell.
montevidensis, *Casar. Nov. Stirp. Bras. Dec.* 78.—Reg. Argent.
runcinata, Lam. Encyc. iii. 189 = Himenanthes runcinata.

JABUTAPITA, Adans. Fam. ii. 364 (1763) = **Gomphia**, Schreb. (Ochnac.).

JABROSA, Steud. Nom. ed. II. i. 795 (1840) = Jaborosa, Juss. (Solan.).

JACARANDA, Juss. Gen. 138 (1789). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1050.
KORDILLERIA, Ruiz & Pav. Syst. Veg. 279 (1798).
KORELLERIS, Aruda, in Koster, Trav. Bras. i. 500 (1816).
PTEROPODIUM, Dc. in Meisn. Gen. 300; Comm. 209 (1840).
RAPINESQUIA, Rafin. Sylva Tellur. 79 (1838).
acutifolia, *Humb. & Bonpl. Pl. Aequin.* i. 59. t. 17.—Peruv.
alba, Spreng. Syst. ii. 854 = Spathodia bracteosa.
Arrabidae, *Steud. Nom.* ed. II. i. 795.—Bras.
arvensis, *Steud. l. c.*—Bras.
bahaensis, R. Br. in Bot. Mag. t. 3327 = caerulea.
brasiliana, *Pers. Syn.* ii. 174.—Bras.
caerulea, *Juss. Gen.* 138; *Griseb. Fl. Brit. W. Ind.* 446.—Ind. occ.
Caroba, *DC. Prod.* ix. 232.—Bras.
Caroba, Hort. ex Lem. Illustr. Hortic. xi. (1864) sub t. 393 = digitalifiora, glozinisaefiora.
carolinensa, Pers. Syn. ii. 174 = caerulea.
Chelonia, *Griseb. in Goett. Abh.* xix. (1874) 223.—Reg. Argent.
Clausseniana, *Casar. Nov. Stirp. Bras. Dec.* vi. 58.—Bras.
coerulea, Auct. = caerulea.
Copaia, *D. Don, in Edinb. Phil. Journ.* (1828) 264 = Panama.
curialis, *DC. Prod.* ix. 232.—Bras.
cuspidifolia, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 51.—Bras.
decurrens, *Cham. in Linnaea*, vii. (1832) 544.—Bras.
densiconna, *Mart. ex A. DC. in DC. Prod.* ix. 232.—Bras.
digitalifiora, *Lem. Illustr. Hortic.* xi. (1864) t. 393.—Bras.
echinata, *Spreng. Syst.* iii. 834.—Am. austr.
elegans, Mart. ex DC. Prod. ix. 231 = oxyphylla.
elliptica, *Steud. Nom.* ed. II. i. 795; *Mart. ex DC. l. c.* 232.—Bras.
endotricha, *DC. Prod.* ix. 231.—Bras.
filicifolia, *D. Don, in Edinb. Phil. Journ.* (1828) 266.—Panama.
glozinisaefiora, Hort. ex Lem. Illustr. Hortic. xi. (1864) sub t. 393 = digitalifiora.
glozinisaefiora, *Lem. l. c.* iii. (1856) *Misc.* 44.—Bras.
hebephora, *DC. Prod.* ix. 231.—Bras.
intermedia, *Sond. in Linnaea*, xxii. (1849) 563.—Bras.
macrantha, *Cham. l. c.* vii. (1832) 552.—Bras.
micrantha, *Cham. l. c.* 554.—Bras.
mimosaefolia, D. Don, in Bot. Reg. t. 881 = ovalifolia.
nitida, *DC. Prod.* ix. 230.—Bras.
obovata, *Cham. in Linnaea*, vii. (1832) 549.—Bras.
obovata, Mart. ex DC. Prod. ix. 230 = subrhombea.
obtusifolia, *Humb. & Bonpl. Pl. Aequin.* i. 62. t. 18.—Venezuela.
orbiculata, Spreng. Syst. ii. 834 = Anemopaegma orbiculatum.
ovalifolia, *R. Br. in Bot. Mag.* t. 2327.—Am. austr.
oxyphylla, *Cham. in Linnaea*, vii. (1832) 546.—Bras.
paucifoliata, *Mart. ex DC. Prod.* ix. 230.—Bras.
paulistana, *Silva Manso, Enum. Subst. Bras.* 40.—Bras.
procera, *Spreng. Syst.* iii. 834.—Guiana.
pteroides, *Silva Manso, Enum. Subst. Bras.* 40.—Bras.
pubescula, *Cham. in Linnaea*, vii. (1832) 550.—Bras.
pubescens, Guill. in A. DC. in DC. Prod. ix. 231 = tomentosa.
pubescens, Lodd. ex G. Don, in Loud. Hort. Brit. 248 = ovalifolia.
quinquefolia, Steud. ex A. DC. in DC. Prod. ix. 233 = Zelyera montana.
racemosa, *Cham. in Linnaea*, vii. (1832) 547.—Bras.
rhombifolia, *G. F. W. Mey. Prim. Fl. Esseq.* 213.—Guian.
rufa, *Silva Manso, Enum. Subst. Bras.* 40.—Bras.
Sagreana, *DC. Prod.* ix. 230.—Ind. occ.
semiserrata, *Cham. in Linnaea*, vii. (1832) 551.—Bras.
spectabilis, *Mart. ex DC. Prod.* ix. 230.—Bras.
subrhombea, *DC. l. c.* 230.—Bras.

JACARANDA:—
subvolutina, Mart. ex DC. Prod. ix. 231 = tomentosa.
tomentosa, *R. Br. in Bot. Mag.* t. 2327.—Bras.
tuberculosa, *Steud. Nom.* ed. II. i. 795.—Bras.
undulata, *Steud. l. c.*—Bras.

JACARATIA, A. DC. Prod. xv. i. 419 (1864) = **Jaracatia**, Marg. (Passifl.).

JACEA, (Tourn.] Linn. Syst. ed. I (1735); Juss. Gen. 173 (1789) = **Centaurea**, Linn. (Compos.).
alata, Lam. Fl. Fr. ii. 53 = C. montana.
alba, Delarb. Fl. Auv. ed. II. 302 = C. amara.
barbata, Delarb. l. c. = C. phrygia †.
Cineraria, Delarb. l. c. 300 = C. Cineraria.
communis, Delarb. l. c. 301 = C. Jacea.
decumbens, Delarb. l. c. = C. amara.
graminifolia, Lam. Fl. Fr. iii. 638 = C. trinervia.
involucrata, Lam. l. c. ii. 54 = C. pullata.
nigra, S. F. Gray, Nat. Arr. Brit. Pl. ii. 443 = C. Jacea.
nigra, Hill, Herb. Brit. i. 28; Lam. Fl. Fr. ii. 52; Moench, Meth. 565 = C. nigra.
paniculata, Lam. l. c. 50 = C. paniculata.
plumosa, Lam. l. c. 51 = C. phrygia.
pratensis, Lam. l. c. 54; Cass. in Dict. Sc. Nat. xxiv. 89 = C. Jacea.
raguzina, Lam. ex Steud. Nom. ed. II. i. 795 = C. raguzina.
Scabiosa, Lam. Fl. Fr. ii. 51 = C. Scabiosa.
segetalis, Lam. ex Steud. Nom. ed. II. i. 795 = seq.
segetum, Lam. Fl. Fr. ii. 54 = C. Cyanus.
supina, Lam. l. c. 53 = C. amara.
tomentosa, Gilib. Fl. Lituan. i. 198 = C. Jacea.
vulgaris, Bernh. Syst. Verz. Erf. 158 = C. nigra.

JACKIA, Blume, Bijdr. 60 (1825) = **Xanthophyllum**, Roxb. (Polygal.).
excelsa, Blume, l. c. 62 = X. excelsum.
longifolia, Blume, l. c. 61 = X. longifolium.
vitellina, Blume, l. c. = X. vitellinum.

JACKIA, Spreng. Syst. iii. 85 (1826) = **Briolaena**, DC. (Sterculiac.).
spectabilis, Spreng. l. c. = E. spectabilis.

JACKIA, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 321 (1824). *RUBIACEAE*, Benth. & Hook. f. ii. 99.
ZUCCARINA, Spreng. Syst. iv. Cur. Post. 50 (1827).
ornata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 321.—Malaya.

JACKSONIA, R. Br. in Ait. Hort. Kew. ed. II. iii. 12 (1811). *LEGUMINOSAE*, Benth. & Hook. f. i. 469.
PITTOMERIS, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 257.
alata, *Benth. in Enum. Pl. Hueg.* 80.—Austral.
angulata, *Benth. Fl. Austral.* ii. 62.—Austral.
capitata, Benth. in Ann. Wien. Mus. ii. (1838) 74 = umbellata.
capitata, *Meisn. in Lehm. Pl. Preiss.* i. 45.—Austral.
carduacea, *Meisn. in Bot. Zeit.* xiii. (1855) 25.—Austral.
compressa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 260.—Austral.
cupulifera, Meisn. in Bot. Zeit. xiii. (1855) 27 = scoparia.
densiflora, *Benth. in Lindl. Swan River App.* 18.—Austral.
dilatata, *Benth. in Ann. Wien. Mus.* ii. (1838) 74.—Austral.
dumosa, Meisn. in Lehm. Pl. Preiss. i. 44 = furcellata.
fasciculata, Meisn. l. c. ii. 212 = hakeoides.
floribunda, *Endl. Nov. Stirp. Dec.* 87.—Austral.
floribunda, Meisn. in Lehm. Pl. Preiss. i. 43 = densiflora.
foliosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 260.—Austral.
furcellata, *DC. Prod.* ii. 107.—Austral.
gracilis, Meisn. in Lehm. Pl. Preiss. i. 44 = scrices.
grandiflora, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 259 = floribunda.
hakeoides, *Meisn. in Lehm. Pl. Preiss.* i. 45.—Austral.
horrida, *DC. Prod.* ii. 107.—Austral.
juncea, Turcz. in Bull. Soc. Nat Mosc. xxvi. (1853) i. 261 = umbellata.
Lehmanni, *Meisn. in Lehm. Pl. Preiss.* i. 46.—Austral.

JACKSONIA:—
macrocalyx, *Meisn. in Bot. Zeit.* xiii. (1855) 26.—Austral.
scaberrima, Benth. in Ann. Wien. Mus. ii. (1838) 74 = scoparia.
nematoclada, *F. Muell. Fragm.* v. 50.—Austral.
odontoclada, *F. Muell. ex Benth. Fl. Austral.* ii. 55.—Austral.
Piptomeris, *Benth. l. c.* 62.—Austral.
pteroclada, *F. Muell. Fragm.* v. 87.—Austral.
purpurascens, *F. Muell. l. c.* iv. 161.—Austral.
racemosa, *Meisn. in Lehm. Pl. Preiss.* ii. 212.—Austral.
ramosissima, *Benth. in Mitch. Journ. Trop. Austral.* 258.—Austral.
resticlea, *Meisn. in Lehm. Pl. Preiss.* i. 46.—Austral.
reticulata, DC. Prod. ii. 107 = Pultenæa reticulata.
rhadinoclada, *F. Muell. Fragm.* v. 88.—Austral.
scoparia, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 13.—Austral.
sericea, *Benth. in Enum. Pl. Hueg.* 31.—Austral.
spinosa, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 13.—Austral.
Stackhousii, *F. Muell. in Proc. Linn. Soc. N. S. W.* vi. (1881) 791.—Austral.
Sternbergiana, *Hueg. Bot. Arch.* t. 3.—Austral.
stricta, *Meisn. in Bot. Zeit.* xiii. (1855) 27.—Austral. thesioides, *A. Cunn. ex Benth. in Ann. Wien. Mus.* ii. (1838) 74.—Austral.
sulcina, Meisn. in Bot. Zeit. xiii. (1855) 26 = hakeoides.
umbellata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 261.—Austral.
velutina, *Benth. Fl. Austral.* ii. 60.—Austral.
vermicosa, *F. Muell. ex Benth. Fl. Austral.* ii. 58.—Austral.
virmicalis, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 75 = Mirbelia oxyclada.

JACKSONIA, Hort. ex Schlecht. in Linnæa, xxvii. (1854) 513 = Jasminum, Tourn. (Olesc.).
nitida, Hort. ex Schlecht. l. c. = Jasm. heterophyllum.

JACKSONIA, Raña. in Med. Repos. N. York, v. (1808) 352 = Cliscoma, Linn. (Capparⁱ.).
trifoliata, Raña. l. c. = C. gravcolens.

JACOBAEA, [Tourn.] Linn. Syst. ed. I (1735); Thunb. Fl. Cap. Prod. Pranf. (1800) = Senecio, Tourn. (Compos.).
abrotanifolia, Moench, Meth. 587 = S. abrotanifolius.
alpina, Moench, l. c. = S. alpinus.
angustifolia, Thunb. Prod. Fl. Cap. 156 = S. angustifolius.
anthoreæfolia, Presl, Delic. Prag. 92 = S. delphinifolius.
aquatica, Gaertn. Mey. & Scherb. Fl. Wett. iii. 210 = S. aquaticus.
aurea, Moench, Meth. Suppl. 239 = S. aureus.
auriculata, Schrank, in Denkschr. Akad. Muench. (1816–17) 194 = S. auriculatus.
Balsamita, Raña. Fl. Ludov. 140 = S. aureus.
bidentata, Thunb. Prod. Pl. Cap. 156 = S. paniculatus.
carniolica, Schrank, in Denkschr. Acad. Muench. 1813 (1814) 816 = S. carniolicus.
cervaria, Presl, Delic. Prag. 93 = S. sethnensis.
cordata, Opiz, Seznam, 54 = S. alpinus.
crassifolia, Raña. in Med. Repos. N. York, v. (1808) 354, nomen (Quid?).—Am. bor.
denticulata, Gilib. Fl. Lituan. i. 200 = S. sylvaticus.
Doria, Gaertn. Mey. & Scherb. Fl. Wett. iii. 214 = S. Doria.
elegans, Moench, Meth. 586 = S. elegans.
erratica, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = S. Jacobæa.
erucifolia, Gaertn. Mey. & Scherb. Fl. Wett. iii. 208 = S. erucifolius.
flaccuosa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = S. Jacobæa.
frondosa, Opiz, Seznam, 54 = S. nemorensis.
Fuchsii, C. A. Mey. Verz. Pfl. Cauc. 81 = S. nemorensis.
giustifolia, Schrank, in Syll. Ratisb. i. (1824) 45 = S. giustifolius, *Linn. f.*
gracilis, Raña. ex DC. Prod. vi. 426 = S. gracilis.
incana, Gilib. Fl. Lituan. i. 209 = S. vernalis.
incana, Raña. in Med. Repos. N. York, v. (1808) 354, nomen (Quid?).—Am. bor.

JACOBAEA:—
incisa, Presl, Delic. Prag. 94 = Senecio sethnensis.
Kustelcki, Opiz, Seznam, 54 = S. paludosus.
lanuginosa, Presl, Delic. Prag. 90 = S. nebrodensis.
macrophylla, Claus. in Goebel, Reise, ii. 238 = S. Doria.
macrophylla, C. A. Mey. Verz. Pfl. Cauc. 81 = S. macrophyllus.
mucronata, Thunb. Prod. Pl. Cap. 156 = S. mucronatus.
nebrodensis, Raña. Chlor. Astr. 6 = S. nebrodensis.
nemorensis, Moench, Meth. 587 = S. nemorensis.
nemorosa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = S. Jacobæa.
nivea, Thunb. Prod. Pl. Cap. 156; Nov. Gen. 171 = S. niveus.
opuntifolia, Raña. Fl. Ludov. 67 = S. opuntiæfolius.
Othonnæ, C. A. Mey. Verz. Pfl. Cauc. 81 = S. Othonnæ.
ovata, Gaertn. Mey. & Scherb. Fl. Wett. iii. 212 = S. Fuchsii.
paludosa, Gaertn. Mey. & Scherb. l. c. 211 = S. paludosus.
paniculata, Thunb. Prod. Pl. Cap. 156 = S. paniculatus.
persicifolia, Thunb. l. c. = S. persicifolius.
pseudanisifolia, Thunb. l. c. 157 = S. paniculatus.
purpurea, Thunb. l. c. = S. purpureus.
reclinata, Thunb. l. c. = S. paniculatus.
rosmarinifolia, Moench, Meth. Suppl. 239 = S. rosmarinifolius.
rubella, Moench, Meth. 586 = S. erucifolius.
saracenica, Moench, l. c. 588 = S. saracenicus.
scabra, Thunb. Nov. Gen. 178 = S. rigidus.
scabra, Thunb. Prod. Pl. Cap. 156 = S. erubescens, rigidus.
sicca, Gilib. Fl. Lituan. i. 199 = S. Jacobæa.
sinuata, Gilib. l. c. 201 = S. vernalis.
sinuata, Moench, Meth. Suppl. 239 = S. aegyptiacus.
squalida, C. A. Mey. Verz. Pfl. Cauc. 81 = S. vernalis.
suaveolens, Gilib. Fl. Lituan. i. 201 = S. erucifolius.
sylvatica, Moench, Meth. 586 = S. sylvaticus.
tenuifolia, J. & C. Presl, Fl. Čech. 175 = S. erucifolius.
tomentosa, Moench, Meth. 587 = S. Cineraria.
tomentosa, Thunb. Nov. Gen. 176; Prod. Pl. Cap. 157 = S. cinerascens.
vernæ, Raña. Chlor. Astr. 6 = S. vernalis?
vernalis, C. A. Mey. Verz. Pfl. Cauc. 81 = S. vernalis.
vestita, Thunb. Prod. Pl. Cap. 156 = S. vestitus.
viscosa, Gilib. Fl. Lituan. i. 200 = S. viscosus.
vulgaris, Claus. in Goebel, Reise, ii. 233 = S. vulgaris.
vulgaris, Gaertn. Fruct. ii. 445 = S. Jacobæa.

JACOBANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 408 = Senecio, Tourn. (Compos.).
crassifolius, Fourr. l. c. = S. crassifolius.
gallicus, Fourr. l. c. 404 = S. gallicus.

JACOREA, Thunb. Nov. Gen. xiii. 169 (1801) = JACOBAEA, Tourn. = Senecio, Tourn. (Compos.).

JACOBINIA, Moric. Pl. Nouv. Am. 156. t. 93 (1846). *ACANTHACEAE*, Benth. & Hook. f. ii. 1114.
CARDIACANTHUS, Nees, in DC. Prod. xi. 331 (1847).
CYRTANTHERA, Nees, in Mart. Fl. Bras. ix. 99. t. 14 (1847).
CYRTANTHERELLA, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 148. t. 3. ff. 10, 11.
DREJERA, Nees, in Mart. Fl. Bras. ix. 112. t. 17 (1847).
ETHESIA, Raña. Fl. Tellur. iv. 68 (1836), nomen prius.
LIEBNIA, C. Koch, in Linden, Prix-Cour. (1868) 2, cum. tab.
PACHYSTACHYS, Nees, in Mart. Fl. Bras. ix. 99 (1847).
SERICOSONIA, Linden & André, Illustr. Hortic. xxii. (1875) t. 198.
SERICOGRAPHIS, Nees, in Mart. Fl. Bras. ix. 107 (1847).
Aschenbornianæ, *Hemsl. Biol. Centr. Am. Bot.* ii. 520.—Mexic.
aurea, *Hemsl. Diagn. Pl. Nov.* 35.—Mexic.
aurea, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 58.—Bras.
barlerioides, *Nees, in DC. Prod.* xi. 383.—Ecuador.
calcicolella, *Grisek. in Gentt. Abh.* xxiv. (1879) 261.—Reg. Argent.

JACOBINIA:—
californica, *Nees, in DC. Prod.* xi. 739.—Calif.
canadicosa, Benth. & Hook. f. Gen. ii. 1115 = Dianthera candicana.
chrysostephana, *Benth. & Hook. f. l. c.*—Mexic.
ciliata, Nees, in DC. Prod. xi. 383 = Dianthera ciliata.
coccinea, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 84.—Bras.
Gleisbrechtiana, *Benth. & Hook. f. Gen.* ii. 1115.—Mexic.
Glaziovii, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 85.—Bras.
harpochlooides, *Benth. & Hook. f. Gen.* ii. 1114, in obs., 1115.—Cuba.
heterophylla, *Hemsl. Biol. Centr. Am. Bot.* ii. 520.—Mexic.
incana, *Hemsl. l. c.* 521.—Mexic.
lepida, *Moric. Pl. Nouv. Am.* 157. t. 93.—Bras.
Liboniana, *Benth. & Hook. f. Gen.* ii. 1115.—Bras.
Lindeni, *Nichols. Dict. Gard.* ii. 206.—Mexic.
longiflora, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 521.—Mexic.
macrantha, *Benth. & Hook. f. Gen.* ii. 1116.—Guatemala.
macrophylla, *Benth. & Hook. f. l. c.* 1115.—Mexic.
mexicana, *Seem. Bot. Voy. Herald*, 295. t. 66.—Mexic.
Mohintli, *Benth. & Hook. f. Gen.* ii. 1115; *Hemsl. Biol. Centr. Am. Bot.* ii. 521.—Mexic.
monticola, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 84.—Bras.
Necsiana, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 521.—Mexic.
neglecta, *A. Gray, Syn. Fl. Am.* ii. 1. 395.—Mexic.
nutans, *Hemsl. Biol. Centr. Am. Bot.* ii. 521.—Mexic.
paniculata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 153.—Mexic.
pauciflora, *Benth. & Hook. f. Gen.* ii. 1115.—Bras.
Pohliana, *Benth. & Hook. f. l. c.*—Bras.
polita, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 84.—Guatemala.
pubescens, *Nees, in DC. Prod.* xi. 382.—Peruv.
Sellowiana, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 85.—Bras.
sericea, *Nees, in DC. Prod.* xi. 382.—Peruv.
sericostomotecha, Nees, l. c. 481 = pubescens.
sulcata, *Nees, l. c.* 383.—Bras.
tinctoria, *Hemsl. Biol. Centr. Am. Bot.* ii. 522.—Guatemala.
virgata, *Hemsl. l. c.*—Mexic.

JACOSTA, DC. Prod. vi. 76 (1837) = Phymasperma, Less. (Compos.).
acicularis, E. Mey. ex Steud. Nom. ed. II. i. 796 = P. acicularc.

JACQUEMONTIA, Bélang. Voy. Ic. [t. 10] (1836) = Gamolepis, Less. (Compos.).
ericæfolia, *l. c.* = G. ericoides.

JACQUEMONTIA, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 476 (Conv. cr. 94). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 674.
abutiloides, *Benth. Bot. Voy. Sulph.* 84.—Calif.
acrocephala, *Meisn. in Mart. Fl. Bras.* vii. 206.—Bras.
agrestis, *Meisn. l. c.*—Bras.
asurea, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 476; vii. i. (1836) 93 = hirsuta.
Blanchetii, Choisy, ll. cc. 476; 64 = glanoscens.
Blanchetii, *Moric. Pl. Nouv. Am.* 41. t. 27.—Bras.
bracteosa, *Meisn. in Mart. Fl. Bras.* vii. 204.—Bras.
caerulea, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 476.—Hab.?
cenocarpa, Benth. in Lindl. Bot. Reg. (1847) t. 27 = violacea.
cenocarpa, *Benth. Pl. Hartw.* 236.—Mexic.
capitata, *G. Don, Gen. Syst.* iv. 268.—Ab. trop.
capitellata, *Choisy, in DC. Prod.* ix. 396.—Bras.
Choisyana, *Meisn. in Mart. Fl. Bras.* vii. 202.—Bras.
coelestis, *Hort. Van Houtt. ex Planch. in Fl. des Serres*, Sér. II. i. (1856) 121.—Am. trop.
cornuta, Anct. vide caerulea.
corymbulosa, *Meisn. in Mart. Fl. Bras.* vii. 204.—Bras.
corymbulosa, *Benth. Bot. Voy. Sulph.* 127.—Ecuador.
crassifolia, *Schaefe, in Linnaea*, xxi. (1848) 753.—Bras.

JACQUEMONTIA :—
erecta, *Choisy, in DC. Prod.* ix. 396.—Bras.
ericophala, *Meisen. in Mart. Fl. Bras.* vii. 303.—Bras.
evolvuloides, *Meisen. l. c.* 307.—Bras.
ferruginea, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 61.—Bras.
glaucescens, *Choisy, l. c.* 64.—Bras.
glomerata, *Parodi, Contrib.* 27.—Paraguay.
gracilis, *Choisy, in DC. Prod.* ix. 396.—Bras.
grandiflora, *Meisen. in Mart. Fl. Bras.* vii. 300.—Bras.
guyanensis, *Meisen. l. c.* 301.—Am. trop.
hastigera, Boj. Rapp. Desj. 33 ; Hort. Maurit. 229 = Ipomœa denticulata.
hirsuta, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 63.—Mexic.
hispida, *Schreele, in Linnaea*, xxi. (1848) 751.—Bras.
lactescens, *Seem. Bot. Voy. Herald*, 171.—Panama.
linarioides, *Meisen. in Mart. Fl. Bras.* vii. 308.—Bras.
linoides, *Meisen. l. c.*—Bras.
Martii, *Choisy, in DC. Prod.* ix. 398.—Bras.
menispermoides, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 63.—Bras.
micrantha, G. Don, Gen. Syst. iv. 283 = Convolvulus micranthus.
montana, *Meisen. in Mart. Fl. Bras.* vii. 304.—Bras.
nodiflora, G. Don, Gen. Syst. iv. 283 = confusa.
nummularia, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 62.—Am. centr.
ovata, *Œrsrin, ex Regel, Gartenfl.* (1860) 271. t. 300.—Chili.
parviflora, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 65.—Mexic.
pentantha. *G. Don, Gen. Syst.* iv. 283.—Am. trop.
Pringlei, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 228.—Texas.
prostrata, *Choisy, in DC. Prod.* ix. 399.—Bras.
pycnocephala, *Benth. Bot. Voy. Sulph.* 187.—Mexic.
racemosa, *Meisen. in Mart. Fl. Bras.* vii. 308.—Bras.
rufo-velutina, *Meisen. l. c.* 305.—Bras.
sandwicensis, *A. Gray, in Proc. Am. Acad.* v. (1861-62) 336.—Isn. Sandvic.
secunda, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 62.—Peruv.
serrata, *Meisen. in Mart. Fl. Bras.* vii. 304.—Bras.
sphærocephala, *Meisen. l. c.* 306.—Bras.
subsessilis, *Moric. Pl. Nouv. Am.* 42. t. 28.—Bras.
tamaifolia, *Griseb. Fl. Brit. W. Ind.* 474.—Am. bor. et trop.
umbellata, *Boj. Hort. Maurit.* 229 (Quid ?).—Ins. Comor.
velutina, *Choisy, in DC. Prod.* ix. 398.—Bras.
violacea, *Choisy, in Mém. Soc. Phys. Genèv.* viii. 1. (1838) 61.—Am. trop.

JACQUINA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 173 = seq.

JACQUINIA, Linn. Dias. Fl. Jam. App. (1759) ; Amœn. Acad. v. 388 (1760) (Jaquinia) ; Jacq. Enum. Pl. Carib. 2 (1760). *MYRSINEÆ*, Benth. & Hook. f. ii. 650.
BONBELLIA, Bert. in Colla, Hort. Ripul. App. iii. 21 (1826).
angustifolia, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 119.—Mexic.
arborea, *Vahl, Eclog. Am.* i. 26.—Am. trop.
aristata, *Jacq. Fragm.* 84.—Ind. occ.
armillaris, *Jacq. Enum. Pl. Carib.* 15.—Ind. occ
aurantiaca, *Ait. Hort. Kew.* ed. II. ii. 6.—Ins. Sandvic. *aurantiaca*, Bert. ex A. DC. in DC. Prod. viii. 151 = umbellata.
axillaris, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 121.—Mexic.
Berterii, *Griseb. Cat. Pl. Cub.* 168.—Cuba.
Berterii, *Spreng. Syst.* i. 668.—Ins. S. Doming.
caracasana, *H. B. & K. Nov. Gen. et Sp.* iii. 251.—Venezuela.
ferruginea, *Spreng. Syst.* i. 666.—Ins. S. Doming.
indica, Wight, ex DC. Prod. viii. 151 ; ix. 464 = Eryclhe pentamera.
linearis, *Jacq. Enum. Pl. Carib.* 15.—Ind. occ.
macrocarpa, *Cav. Ic.* v. 55. t. 458.—Am. trop.
macrocarpa, Cham. in Linnæa, vi. (1831) 394 = racemosa.

JACQUINIA :—
macrocarpa, Spreng. Syst. i. 668 = aurantiaca.
mexicana, *Regel, Ind. Sem. Hort. Petrop.* (1865) 48.—Mexic.
mucronata, Willd. ex Spreng. Syst. i. 668 = pubescens.
pubescens, *H. B. & K. Nov. Gen. et Sp.* iii. 251. t. 244.—Am. austr.
pungens, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 325.—Bras.
racemosa, *A. DC. in DC. Prod.* viii. 150.—Mexic.
revoluta, *Jacq. Fragm.* 84.—Venezuela.
ruscifolia, Hort. Tur. ex Steud. Nom. ed. II. i. 796 = Bontia daphnoides.
ruscifolia, *Jacq. Enum. Pl. Carib.* 15 ; *Linn. Sp. Pl.* ed. II. 272.—Ind. occ. ; Mexic.
ruscifolia, Spreng. Syst. i. 668 = linearis.
sarmentosa, *Planch. ex Linden, Pl. Nouv.* (1859) 8.—Mexic.
umbellata, *A. DC. in DC. Prod.* viii. 150.—Mexic.
venosa, Sw. Prod. Veg. Ind. Occ. 47 = Chione glabra.

JACQUINIA, Mutis, ex Linn. Mant. ii. 247 (1771) = **Prockia**, Linn. (Tiliac.).

JACQUINOTIA, Homb. & Jacquinot, ex Decne. Voy. Bot. Astrol. & Zél. Bot. Phan. Dicot. t. 82 (1853) = **Lebetanthus**, Endl. (Epacrid.).
Myrsinites, Homb. & Jacquin. ex Decne. l. c. = L. americanus.
prostrata, Hom'. & Jacquin. l. c. = L. americanus.
volubilis, Homb. & Jacquin. l. c. = L. americanus.

JACUANGA, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 341 = **Costus**, Linn. (Scitamin.).

JACULARIA, Rafin. in Lond. Gard. Mag. vii. (1832) 247 = **Cunninghamia**, R. Br. (Conif.).

JAEGERA, Gieske, Prael. Linn. 208, 286 (1792) = **Zingiber**, Adans. (Scitamin.).
montana, Gieske, l. c. = Z. montanum.

JAEGERIA, H. B. & K. Nov. Gen. et Sp. iv. 277. t. 400. *COMPOSITÆ*, Benth. & Hook. f. ii. 460.
MACELLA, C. Koch, Ind. Sem. Hort. Berol. (1855) App. 13.
abyssinica, Spreng. Syst. iii. 590 = Gnizotia abyssinica.
bellidioides, *Spreng. l. c.* 591.—Reg. Argent.
calendulacea, Spreng. l. c. 500 = Wedelia calendulacea.
gracilis, *Kook. f. in Trans. Linn. Soc. xx.* (1847) 218.—Ins. Galap.
hirta, *Less. Syn. Comp.* 228.—Am. austr.
macrocephala, *Less. l. c. ; et in Linnæa*, ix. (1834) 270.—Am. austr.
miniodes, *H. B. & K. Nov. Gen. et Sp.* iv. 277. t. 400.—Mexic.
parviflora, DC. Prod. v. 544 = hirta.
pedunculata, *Hook. & Arn. Bot. Beech. Voy.* 299.—Mexic.
prorepens, *Hook. f. in Trans. Linn. Soc. xx.* (1847) 214. Galap.
repens, DC. Prod. v. 544 = hirta.
uliginosa, Spreng. Syst. iii. 590 = Spilanthes Acmella.
urticæfolia, Spreng. l. c. = Sabazia urticæfolia.

JAEGERIERA, Kurz, in Journ. As. Soc. Beng. xxxix. ii. (1870) 280. t. 18. *GENTIANEÆ*, Benth. & Hook. f. ii. 816.
gentianoides, *Kurz, l. c.*—Reg. Himal.
latisepala, *C. B. Clarke, in Journ. Linn. Soc.* xiv. (1875) 441.—Reg. Himal.
microcarpa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 119.—Reg. Himal.

JAGERA, Blume, Rumphia, iii. 155 (1847). *SAPINDACEÆ*, Benth. & Hook. f. i. 408.
glabra, *Hassk. Hort. Bog. Desc.* 157.—Java.
javanica, Blume, Mus. Bot. Lugd. Bat. i. 227 = speciosa.
madagascariensis, Blume, Rumphia, iii. 155 = Garuga madagascariensis.
Pseudorhus, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 231 = Cupania Pseudorhus.
Roxburghii, *Blume, Rumphia*, iii. 155.—Ins. Molucc.
serrata, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 208 = Roxburghii.
speciosa, *Blume, Rumphia*, iii. 155.—Malaya ; N. Guin.

JALAMBICA, Rafin. New Fl. Am. iv. 71 (1836).—Cf. NEURELMIS, Rafin. (Compos.).
pumila, Rafin. l. c. 72.—' Cuba vel Florida.'

JALAMBICEA, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 12 (1825) = **Limnobium**, Rich. (Hydrocharid.).
repens, Cerv. l. c. = L. stoloniferum.

JALAPA, Tourn. ex Adans. Fam. ii. 265 (1763) = **Mirabilis**, Linn. (Nyctagin.).
congesta, Moench, Meth. 508 = M. Jalapa.
dichotoma, Crantz, Inst. ii. 266 = M. dichotoma.
longiflora, Crantz, l. c. = M. longiflora.
officinalis, Crantz, l. c. = M. Jalapa.
officinarum, Martyn, Cent. i. t. 1 = M. dichotoma.
undulata, Moench, Meth. Suppl. 196 = M. dichotoma.

JALAPPA, Burm. Thes. Zeyl. 124 (1737) = præc.

JALOMBICEA, Steud. Nom. ed. II. i. 796 (1840), sphalm. = JALAMBICEA, Cerv. = **Limnobium**, Rich. (Hydroch.).

JALTOMATA, Schlecht. Ind. Sem. Hort. Hal. (1838) n. 7 ; ex Linnæa, xiii. (1839) Litt. 98 = **Saracha**, Ruiz & Pav. (Solanac.).
edulis, Schlecht. ll. cc. = S. Jaltomata.

JALTONIA, Steud. Nom. ed. II. i. 796 (1840) = præc.

JAMBOLANA, Adans. Fam. ii. 508 (1763) = **Acronychia**, Forst. (Rutac.).

JAMBOLIFERA, Houtt. Handleid. ii. 272 (1774) = **Eugenia**, Mich. (Myrtac.).
coromandelica, Houtt. l. c. 275.—Ind. or.
pedunculata, Houtt. l. c. 273 ; Gaertn. Fruct. i. 178 = E. Jambolifera.

JAMBOLIFERA, Linn. Dias. Dess. 15 (1747) ; Amœn. Acad. i. 298 (1749) = **Acronychia**, Forst. (Rutac.).
arborea, Zoll. & Mor. Syst. Verz. Zoll. 14 = A. arborea.
chinensis, Spreng. Syst. iii. 216 = A. laurifolia.
oblongifolia, Steud. Nom. ed. II. i. 796 = A. laevis.
odorata, Lour. Fl. Cochinch. 231.—Cochinch.
pedunculata, Linn. Sp. Pl. 349 = A. laurifolia.
resinosa, Lour. Fl. Cochinch. 231 = A. laurifolia.

JAMBOS, Adans. Fam. ii. 88 (1763) = seq.

JAMBOSA, DC. Prod. iii. 286 (1828) = **Eugenia**, Mich. (Myrtac.).
acida, Hort.—Cf. Gard. Chron. (1876) i. 735.—Hab. ?
acuminata, Sweet, Hort. Brit. ed. II. 219 = E. acuminata.
acuminatissima, Hassk. Cat. Hort. Bog. Alt. 362 = E. acuminatissima.
acutata, Miq. Fl. Ind. Bat. i. 1. 482 = E. acutata.
aemula, Blume, Mus. Bot. Lugd. Bat. i. 99 = E. polypetala.
alba, G. Don, Gen. Syst. iii. 868 = E. javanica.
ambigua, Blume, Mus. Bot. Lugd. Bat. i. 96 = E. javanica.
amplericaulis, DC. Prod. iii. 287 = E. amplexicaulis.
anastomosans, Miq. Fl. Ind. Bat. Suppl. 309.—Sumatra.
appendiculata, Blume, Mus. Bot. Lugd. Bat. i. 361 = Memecylon appendiculatum.
aquea, DC. Prod. iii. 286 = E. aquea.
aromatica, Miq. Fl. Ind. Bat. i. 1. 486.—Java.
attenuata, Miq. l. c. 487.—Java.
auriculata, Blume, Mus. Bot. Lugd. Bat. i. 104.—Java !
australis, DC. Prod. iii. 287 = E. myrtifolia, *Sims*.
bifaria, Miq. Fl. Ind. Bat. i. 1. 492 = E. bifaria.
blanica, Miq. l. c. Suppl. 310.—Ins. Banca.
borneensis, Miq. l. c. i. 1. 484.—Malaya.
Brachenridgei, Bronga. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 161.—N. Caled.
bracteata, Miq. Fl. Ind. Bat. i. 1. 487 = E. zeylanica.
buxifolia, Miq. l. c. 1086.—Ins. Banca.
calophylla, Miq. l. c. 493.—Java.
canaliculata, DC. Prod. iii. 287 = E. polycephala.
celebica, Blume, Mus. Bot. Lugd. Bat. i. 107.—Ins. Celebes.

JAMBOSA :—
cerasiformis, Hassk. Cat. Hort. Bog. Alt. 262.—Java.
clavata, Korth. in Nederl. Kruidk. Arch. i. (1847) 201 = Eugenia rhododendrifolia.
coarctata, Blume, Mus. Bot. Lugd. Bat. i. 99 = E. macrocarpa.
Commersonii, Blume, l. c. 105.—Java.
condensata, Miq. Fl. Ind. Bat. i. i. 419.—Java.
confusa, Korth. in Nederl. Kruidk. Arch. i. (1847) 202 = E. marginifolia ?
confusa, Blume, Mus. Bot. Lugd. Bat. i. 101.—Sumatra.
cornifolia, Blume, l. c. 92.—Ins. Celebes.
corymbosa, Miq. Fl. Ind. Bat. i. i. 420.—Java.
costata, Miq. l. c. 415.—Java.
cuneata, Blume, Mus. Bot. Lugd. Bat. i. 105.—Java.
cuspidata, Blume, ex Miq. Fl. Ind. Bat. i. i. 425.—Java.
cylindrica, Thw. Enum. Pl. Zeyl. 115 = E. cylindrica.
cymifera, E. Mey. in Drège, Zwei Pfl. Docum. 194.—Afr. austr.
cymnifera, Presl, Bot. Bemerk. 70, sphalm. = præc.
densiflora, DC. Prod. iii. 287 = E. densiflora.
dentatica, DC. l. c. 288 = E. malaccensis.
eucalyptoides, F. Muell. Fragm. i. 226 = E. eucalyptoides.
firma, Blume, Mus. Bot. Lugd. Bat. i. 108 = E. grandis.
firma, Miq. Fl. Ind. Bat. i. i. 429.—Java.
formosa, G. Don, Gen. Syst. ii. 869 = E. formosa.
glabrata, DC. Prod. iii. 287 = E. ramosissima.
glandulifera, Miq. Fl. Ind. Bat. i. i. 438 = E. zeylanica.
glandulosa, Korth. in Nederl. Kruidk. Arch. i. (1847) 201.—Borneo.
gracilipes, C. Muell. in Walp. Ann. iv. 849 = E. gracilipes.
gracilis, Korth. in Nederl. Kruidk. Arch. i. (1847) 202.—Java.
grandis, Blume, Mus. Bot. Lugd. Bat. i. 108 = E. grandis.
Hardwickiæ, G. Don, Gen. Syst. ii. 869.—Ins. Maurit.
hemisphærica, Wight, Illustr. Ind. Bot. ii. 14 = E. hemisphærica.
hirta, Korth. in Nederl. Kruidk. Arch. i. (1847) 200.—Borneo.
Horsfieldii, Miq. Fl. Ind. Bat. i. i. 420.—Java.
hypericifolia, DC. Prod. iii. 287.—Java.
hypericifolia, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 100 = E. macrocarpa, *Roxb.*
inophylla, Miq. Fl. Ind. Bat. i. i. 433 = E. inophylla.
insignis, Blume, Mus. Bot. Lugd. Bat. i. 100.—Borneo.
Kiguruug, Blume, l. c. 107.—Java.
Klamph, Miq. Fl. Ind. Bat. i. i. 421.—Java.
Koenigii, Blume, Mus. Bot. Lugd. Bat. i. 100 = E. zeylanica.
Korthalsii, Blume, l. c. 101.—Malaya.
lacta, Blume, l. c. 104 = E. lacta.
laevis, Montr. in Mém. Acad. Lyon, x. (1860) 208.—N. Caled.
lanceolaria, Blume, Mus. Bot. Lugd. Bat. i. 101 = E. lanceolaria.
lanceolata, Korth. ex Miq. Fl. Ind. Bat. i. i. 426.—Java.
lanciflora, Miq. l. c. 427.—Borneo.
latifolia, Miq. l. c. 429.—Borneo.
laurifolia, DC. Prod. iii. 287 = E. bifaria.
leptostachya, Blume, Mus. Bot. Lugd. Bat. i. 99 = E. leptostemon.
leptostemon, Korth. in Nederl. Kruidk. Arch. i. (1847) 201 = E. leptostemon.
linearis, Korth. l. c. 199 = E. lacta.
linnata, DC. Prod. iii. 287 = E. linnata, *Duthie.*
litoralis, Blume, Mus. Bot. Lugd. Bat. i. 102.—Java.
longifolia, Bronga. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 181.—N. Caled.
macrocarpa, Sweet, Hort. Brit. ed. II. 212 = E. macrocarpa.
macrophylla, Boj. Hort. Maurit. 148 = E. Bojeri.
macrophylla, DC. Prod. iii. 286 = E. malaccensis.
macrophylla, Llanos, in Mem. Acad. Ciene. Madr. ii. (1859) 800.—Hab. ?
macrophylla, Zoll. ex Miq. Fl. Ind. Bat. i. i. 415 = *palymurea.*
madagascariensis, Blume, Mus. Bot. Lugd. Bat. i. 103 = E. aquea.

JAMBOSA :—
malaccensis, DC. Prod. iii. 286 = Eugenia malaccensis.
mappacea, Korth. in Nederl. Kruidk. Arch. i. (1847) 200 = E. formosa, *Wall.*
marginata, Miq. Fl. Ind. Bat. i. i. 428.—Java.
maritima, Miq. l. c. 435.—Java.
media, Korth. in Nederl. Kruidk. Arch. i. (1847) 199.—Borneo.
melanocarpa, Miq. Fl. Ind. Bat. i. i. 430.—Java.
melanoticta, Miq. l. c. 432.—Java.
melastomafolia, Blume, Mus. Bot. Lugd. Bat. i. 102.—Malaya.
memecyloides, Teijsm. & Binn. Cat. Hort. Bog. 394, nomen.—Ins. Banca.
Munronii, Wall. Rep. ii. 191 = E. Munronii.
myrtifolia, Heynh. Nom. i. 426 = E. myrtifolia, *Sims.*
nervifolia, Bronga. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 181.—N. Caled.
neurocalyx, C. Muell. in Walp. Ann. iv. 849 = E. neurocalyx.
nitida, Korth. in Nederl. Kruidk. Arch. i. (1847) 202 = E. nervosa.
obtusissima, DC. Prod. iii. 287 = E. javanica.
oenifolia, Blume, Mus. Bot. Lugd. Bat. i. 98.—N. Guinea.
owarinsis, DC. Prod. iii. 287 = E. owarinsis.
palembanica, Blume, Mus. Bot. Lugd. Bat. i. 93 = E. javanica.
pauciflora, Walp. Rep. ii. 192 = E. lacta.
peregrina, Blume, Mus. Bot. Lugd. Bat. i. 92.—Malaya.
perfoliata, Miq. Fl. Ind. Bat. Suppl. 809.—Ins. Banca.
polycephala, Miq. l. c. i. i. 439 = E. polycephala.
polyneura, Miq. l. c. 416.—Java.
polyneura, Walp. Rep. ii. 192 = E. polypetala.
Pseudo-malaccensis, Vieill. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 182.—N. Caled.
pterocarpa, Korth. in Nederl. Kruidk. Arch. i. (1847) 200.—Borneo.
pulchella, Miq. Fl. Ind. Bat. i. i. 422 = E. oblata.
puncticulata, Miq. Fl. Ind. Bat. Suppl. 810.—Sumatra.
purpurascens, DC. Prod. ii. 286 = E. malaccensis.
purpurea, Korth. in Nederl. Kruidk. Arch. i. (1847) 201.—Sumatra.
purpurea, Wight & Arn. Prod. 238 = E malaccensis.
quadrangularis, C. Muell. in Walp. Ann. iv. 849 = E. quadrangularis.
reticulata, Blume, Mus. Bot. Lugd. Bat. i. 92, in obs. = E. marginifolia.
rhyncophylla, Miq. Fl. Ind. Bat. i. i. 430.—Java.
rhytidocarpa, Zoll. in Miq. Choix Pl. Buitons. t. 4.—Malaya.
Richii, C. Muell. in Walp. Ann. iv. 849 = E. Richii.
rostrata, Miq. Fl. Ind. Bat. i. i. 436.—Sumatra.
rotundifolia, G. Don, Gen. Syst. ii. 869 = E. rotundifolia.
rubens, Blume, Mus. Bot. Lugd. Bat. i. 106 = E. rubens.
rubrinervis, Miq. Fl. Ind. Bat. i. i. 432 = E. rubrinervis.
saligna, Miq. l. c.—Java.
samarangensis, DC. Prod. iii. 286 = E. javanica.
sarawakensis, Miq. Fl. Ind. Bat. i. i. 422.—Java.
splendens, Miq. l. c. 435.—Java.
subsessilis, Miq. l. c. Suppl. 809.—Sumatra.
sulcata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 54.—Malaya.
sumatrana, Miq. Fl. Ind. Bat. i. i. 419.—Sumatra.
symphyocarpa, Korth. ex Miq. l. c. 436.—Sumatra.
syzygioides, Miq. l. c. 431.—Java.
tawaknensis, Korth. in Nederl. Kruidk. Arch. i. (1847) 202.—Borneo.
tenuicuspis, Miq. Fl. Ind. Bat. i. i. 431 = E. cymosa.
tenuifolia, Sweet, Hort. Brit. ed. II. 212 = E. formosa, *Wall.*
tenuiramis, Miq. Fl. Ind. Bat. i. i. 437.—Java.
tetraedra, Miq. Fl. Ind. Bat. Suppl. 810 = E. tetraedra.
tetragona, Blume, Mus. Bot. Lugd. Bat. i. 106.—Java.
tetraptera, Miq. Fl. Ind. Bat. Suppl. 811.—Ins. Banca.
tetrapyxidata, Miq. l. c. i. i. 433.—Java.
Teysmanni, Miq. l. c. 429 = E. ramosissima.
Theodelana, F. Muell. Fragm. i. 228 = E. myrtifolia, *Sims.*
timorensis, Blume, Mus. Bot. Lugd. Bat. i. 97 = E. javanica.
trifida, Miq. Fl. Ind. Bat. i. i. 435.—Sumatra.

JAMBOSA :—
urceolata, Korth. ex Miq. Fl. Ind. Bat. i. i. 418, partim = Eugenia grandis.
venosa, DC. Prod. iii. 286 = E. venosa.
virens, Miq. Fl. Ind. Bat. i. i. 426.—Java.
Vriesiana, Miq. l. c. 424.—Java.
vulgaris, DC. Prod. iii. 286 = E. Jambos.
Wightiana, Blume, Mus. Bot. Lugd. Bat. i. 106 = E. rubens.
Zollingeriana, Miq. Fl. Ind. Bat. i. i. 418.—Java.

JAMBUS, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 18 = JAMBOSA, DC. = **Eugenia**, Mich. (Myrtac.).
acidus, Noronha, l. c. (= *Jambosa hypericifolia*).—Java.
calycinus, Noronha, l. c. (Quid ?).—Java.
rostrus, Noronha, l. c. = E. Jambos.
ruber, Noronha, l. c. = E. aquea.

JAMESIA, Nees, in Nenwied, Reise N. Am. ii. 442 (1844) = **Stephanomeria**, Nutt. (Compos.).
pauciflora, Nees, l. c. = S. minor.

JAMESIA, Rafin. Atl. Journ. 145 (1832) = **Dalea**, Linn. (Legumin.).
obovata, Rafin. l. c. = D. Jamesii.

JAMESIA, *Torr. & Gray*, Fl. N. Am. i. 638 (1840).
SAXIFRAGEÆ, Benth. & Hook. f. i. 643.
americana, *Torr. & Gray*, l. c.—N. Mexic.

JANASIA, Rafin. Fl. Tellur. iv. 63 (1836) = **Phlogacanthus**, Nees (Acanthac.).
rosea, Rafin. l. c. = P. asperulus.
rubra, Rafin. l. c. = P. asperulus.

JANCAEA, Boiss. *Fl.* Orient. Nov. Dec. i. 4 (1875) = **Ramondia**, Rich. (Gesnerac.).
Heldreichii, Boiss. l. c. 5 = R. Heldreichii.

JANDINEA, Stend. in Flora, xxxiii. (1850) 229, sphalm. = JARDINEA, Steud. = **Thelepogon**, Roth (Gramin.).

JANGARACA, Rafin. in Ann. Sc. Gén. Phys. vi. (1820) 82, sphalm. = TANGARACA, Adams. = **Hamelia**, Jacq. (Kubiac.).

JANIA, Schult. f. Syst. vii. p. xcviii, 1528 (1830) = **Basomstra**, Salisb. (Liliac.).
Breyniana, Schult. f. l. c. 1528 = B. columellaris.

JANIPHA, H. B. & K. Nov. Gen. et Sp. ii. 106. t. 109 (1817) = **Manihot**, Adans. (Euphorb.).
aesculifolia, H. B. & K. l. c. 107. t. 109 = M. aesculifolia.
angustifolia, Pohl, ex Heynh. Nom. i. 428.—Bras.
anisophylla, Griseb. in Goett. Abh. xix. (1874) 95.—Reg. Argent.
arborea, Mart. ex Pohl, Pl. Bras. i. 62 = Jatropha obtusifolia.
foetida, H. B. & K. Nov. Gen. et Sp. ii. 106 = M. foetida.
Fuquilla, Mnell. Arg. in DC. Prod. xv. ii. 1073, in syn. = F. *Fuquilla*.
Lanjhingii, H. B. & K. Nov. Gen. et Sp. ii. 107 = M. carthagenensis.
Manihot, H. B. & K. l. c. 108 = M. utilissima.
obtusifolia, Mart. ex Baill. Adansonia, iv. (1863–64) 275 = Jatropha obtusifolia.
phyllacantha, Mart. ex Pohl, Ic. Pl. Bras. i. 62 = Jatropha phyllacantha.
urens, Poir. ex Pohl, l. c. 62 = Jatropha urens.
violacea, Griseb. in Goett. Abh. xxiv. (1879) 63.—Reg. Argent.
Yuquilla, H. B. & K. Nov. Gen. et Sp. ii. 107 = M. carthagenensis.

JANKAEA, Boiss. Fl. Orient. iv. 89 (1879) = **Ramondia**, Rich. (Gesnerac.).
Heldreichii, Boiss. l. c. 89 = R. Heldreichii.

JANRAIA, Adanss. Fam. ii. 76 (1763) = **Eugenia**, Linn. (Dicocor.).

JANSONIA, Kipp. in Proc. Linn. Soc. i. (1847) 330; et in Trans. Linn. Soc. xx. (1851) 394. t. 16. *LEGUMINOSÆ*, Benth. & Hook. f. i. 467.
CRYPTOSEMA, Meissn. in Lehm. Pl. Preiss. ii. 207 (1847).
formosa, *Kipp. ll. cc.*—Austral.

JANTHA, Stend. Nom. ed. II. i. 797 (1840) = IANTHA, Hook. = **Ionopsis**, H. B. & K. (Orchid.).

JANTHE, Griseb. Spicil. Fl. Rumel. ii. 40 (1844) = **Celsia**, Linn. (Scrophulac.).
bugulifolia, Griseb. l. c. = C. bugulifolia.

JANUSIA, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 260; et in Arch. Mus. Par. iii. (1843) 503. t. 2 (Monog. 249). *MALPIGHIACEÆ*, Benth. & Hook. f. i. 369.
amazonica, *Griseb. in Mart. Fl. Bras.* xii. i. 103.—Bras.
asiandra, *Griseb. l. c.*—Bras.
argentea, *Griseb. in Goett. Abh.* xxiv. (1879) 68.—Arg. Argent.
californica, *Benth. Bot. Voy. Sulph.* 8.—Calif.
caudata, *Griseb. in Mart. Fl. Bras.* xii. i. 103.—Bras.
discolor, A. Juss. in Arch. Mus. Par. iii. (1843) 504 = Camarea discolor.
gracilis, *A. Gray, Pl. Wright.* i. 37.—Texas.
guaranitica, *A. Juss. in Arch. Mus. Par.* iii. (1843) 504. t. 21.—Bras.
linearifolia, A. Juss. l. c. 503 = Camarea juncea.
muricata, *Griseb. in Mart. Fl. Bras.* xii. i. 103.—Bras.
sericea, A. Juss. in Arch. Mus. Par. iii. (1843) 504 = Camarea pulchella.

JAPOTAPITA, Endl. Gen. 1142, in syn. (1841) = JABOTAPITA, Adans. = **Gomphia**, Schreb. (Ochnac.).

JAQUINIA, Linn. Dies. Fl. Jamaic. App. (1759); Amoen. Acad. v. 398 (1760) = **Jacquinia**, Linn. (Myrsin.).

JAQUINOTIA, Walp. Ann. i. 481 (1848–49) = JACQUINOTIA, Hombr. = **Lobetanthus**, Endl. (Epacrid.).

JARACATIA, Marcg. ex Endl. Gen. 983 (1840). *PASSIFLORÆ*, Benth. & Hook. f. i. 815.
conica, *Kerber, in Eichl. Fahrb. Berl. ii.* (1888) 282.—Mexic.
dodecaphylla, *A. DC. in DC. Prod.* xv. i. 420.—Bras.
heptaphylla, *A. DC. l. c.*—Bras.
mexicana, *A. DC. l. c.*—Mexic.
spinosa, *A. DC. l. c.* 419.—Am. trop.

JARAPHA, Stend. Nom. ed. II. i. 797 (1840) = JARAVA, Ruiz & Pav. = **Stipa**, Linn. (Gramin.).

JARAPHAEA, Stend. Nom. ed. II. i. 796 (1840) = JARAVAEA, Scop. = **Microlicia**, D. Don (Melastom.).

JARAVA, Ruiz & Pav. Prod. Fl. Per. 2. t. 1 (1794) = **Stipa**, Linn. (Gramin.).
arundinacea, Willd. ex Stend. Nom. ed. II. i. 799 = S. Jarava.
Ichu, Ruiz & Pav. Fl. Per. i. 5 = S. Jarava.
ustata, Pers. Syn. i. 6 = S. Jarava.

JARAVAEA, Nock. Elem. ii. 118 (1790) = JARAVAEA, Scop. = **Microlicia**, D. Don (Melastom.).

JARDINEA, Stend. in Flora, xxiii. (1850) 229 (Jardinea); Syn. Pl. Gram. 350 (1855) = **Thalepogon**, Roth (Gramin.).
abyssinica, Stend. Syn. Pl. Gram. 350 = T. elegans.
gabonensis, Stend. l. c. (1855) 229.—Afr. trop.

JARDINIA, Benth. & Hook. f. Gen. iii. 1181 (1883) = prec.

JARDINIA, Sch. Bip. in Flora, xxxvi. (1853) 84, in syn. = **Brianæa**, Sch. Bip. (Compos.).
plumosa, Sch. Bip. l. c. = E. plumosa.

JASIONE, Linn. Syst. ed. I (1735). *CAMPANULACEÆ*, Benth. & Hook. f. ii. 554.
OVILLA, Adans. Fam. ii. 134 (1763).
acaulis, Vahl, Symb. ex Willd. & Lange, Prod. Fl. Hisp. ii. 353 = humilis.
amethystina, *Lag. & Rodr. in Anal. Clenc. Nat.* v. (1805) 271.—Hispan.
amethystina, Rent. ex Willd. & Lange, Prod. Fl. Hisp. ii. 354 = perennis.
armeria, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 553 = corymbosa.
atlantica, *Ball, in Journ. Bot.* xi. (1873) 272.—Marocc.
Amédéïi, *Regel, Gartenfl.* (1861) 356.—Austral.
bastica, Rodr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 281 = blepharodon.
blepharodon, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 72.—Hispan.
Bovei, *Boiss. & Reut. l. c.* 74.—Afr. bor.
caespitosa, *Koch, ex Roth, Enum. Pl. Phan. Germ.* i. 774.—Europ.
capensis, Berg in Nov. Act. Soc. Sc. Upsal. iii. (1780) 167. t. 10 = Astrantia ciliaris.
Cariensi, Bor. Fl. Cent. Fr. ii. 387 = montana.
carpetana, Boiss. & Reut. ex Boiss. Voy. Bot. Espagne, ii. 745 = perennis.
cossetana, *Pomel, Nouv. Mat. Fl. Atl.* 258.—Alger.
cornuta, *Ball, in Journ. Bot.* xi. (1873) 278.—Marocc.
corymbosa, *Poir. Encyc. Suppl.* iii. 181.—Hispan.
depressa, Huet, ex Nym. Consp. 487 = supina.
echinata, Boiss. & Reut. Pugill. Pl. Nov. (1852) 73 = montana.
fallax, Willk. in Flora, xxxv. (1852) 198 = humilis.
foliosa, *Cav. Ic. Rar.* ii. 88. t. 148. f. 1.—Hispan.
falcica, Willk. Enum. Pl. Nov. 44 = mariana.
glabra, Dur. ex Boiss. & Reut. Pugill. Pl. Nov. (1852) 73 = corymbosa.
glabra, *Valenz. in Oestr. Bot. Zeitschr.* xxxiv. (1884) 424.—Europ.
glabrata, Dum. Fl. Belg. 59 = montana.
Heldreichii, Boiss. & Orph. Diagn. Ser. II. vi. 120 = montana.
Heldreichii, Janka, ex Nym. Consp. 486 = Jankæ.
humilis, Costa, Fl. Catal. 161 = amethystina.
humilis, Cat. Fl. Madrit. 456 = humilis.
humilis, Jan, Elench. 8 = montana.
humilis, *Loisel, Not. Pl. Fr.* 49.—Europ.
Jankæ, *Heldr. Herb. Nachtr. Exp. Slav.* 48.—Banat.
laevis, Lam. Fl. Fr. ii. 8 = perennis.
litoralis, Fries, Summa Veg. 11 = montana.
lusitanica, *A. DC. Monog. Camp.* 105.—Lusitan.
montana, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 284.—Hispan.
maritima, Duf. ex A. DC. in DC. Prod. vii. 415 = montana.
megalocalyx, *Pomel, Nouv. Mat. Fl. Atl.* 258.—Alger.
montana, *Linn. Sp. Pl.* 928.—Europ.; Afr. bor.
montana, Schousb. Gew. Marok. 70 = corymbosa.
montana, Stend. Nom. ed. II. i. 797 = Alepidea ciliaris.
orbiculata, Griseb. Spicil. Fl. Rumel. ii. 253 = supina.
Pavonei, Clarion, Fl. Cent. Fr. ii. 287 = montana.
pencillata, *Boiss. in DC. Prod.* vii. 416.—Hispan.
perennis, Bertol. Lucubr. 22. n. 118 = montana.
perennis, *Lam. Encyc.* iii. 216.—Europ. austr.
pygmaea, Timb. in Bull. Soc. Sc. Phys. Nat. Toul. vi. (1888–84) 149 = perennis.
romularis, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 74.—Hispan.
scabiosa-capitata, Gilib. Fl. Lituan. i. 87 = montana.
sessiliflora, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 21 = humilis.
stricta, *Pomel, Nouv. Mat. Fl. Atl.* 1.—Alger.
supina, *Sieber, ex Spreng. Syst.* i. 810.—Bosnia; As. Min.!
tristis, Borry, in Ann. Gén. Sc. Phys. iii. (1820) 3 = amethystina.
umbellata, Gilib. Fl. Lituan. i. 58 = montana.
undulata, Lam. Fl. Fr. ii. 8 = humilis, montana.

JASMINANTHES, Blume, Mus. Bot. Lugd. Bat. i. 143. t. 28 (1850) = **Stephanostia**, Thou. (Asclepiad.).
suaveolens, Blume, l. c. 149.—Borneo.

JASMINARIA, Heist. Syst. 10 (1748) (Quid?).

JASMINIUM, Dum. Anal. Fam. 36 (1829) = **Jasminum**, Tourn. (Oleac.).

JASMINOIDES, Medic. in Phil. Bot. i. 184 (1789) = **Lycium**, Linn. (Solanac.).
afrum, Medic. l. c. = L. afrum.
flaccidum, Moench, Meth. 476 = L. barbarum.
linearifolium, Moench, Meth. Suppl. 177 = L. afrum.
rhombifolium, Moench, Meth. 476 = L. europæum.

JASMINUM, [Tourn.] Linn. Syst. ed. I (1735). *OLEACRÆ*, Benth. & Hook. f. ii. 674.
JACKSONIA, Hort. ex Schlecht. in Linnæa, xxvii. (1854) 512.
MOGORIUM, Juss. Gen. 106 (1789).
abyssinicum, *R. Br. in Salt, Abyss. App.* 63; *Hochst. ex DC. Prod.* viii. 311.—Abyssin.
acuminatisalmum, *Blume, Mus. Bot. Lugd. Bat.* i. 276. —Java.
acuminatum, R. Br. Prod. 521 = simplicifolium.
acuminatum, Heyne, ex Wall. Cat. n. 2881 = flexile.
acuminatum, Pers. Syn. i. 7 = trinerve
adenophyllum, *Wall. Cat.* n. 2876.—Reg. Himal.
adiantifolium, *Planch. Hort. Donat.* 64.—Hab.?
aemulum, *R. Br. Prod.* 521.—Austral.
affine, *Blume, Mus. Bot. Lugd. Bat.* i. 37.—Malaya.
affine, Royle, in Bot. Reg. (1845) t. 26 = officinale.
affine, Wight, Ic. t. 1255 = auriculatum.
ambiguum, *Blume, Mus. Bot. Lugd. Bat.* i. 278.—Malaya.
amoenum, *Blume, l. c.* 270.—Ins. Celebes.
amplexicaule, Buch.-Ham. in Wall. Cat. n. 2875 = undulatum.
anastomosans, *Wall. Cat.* n. 2868.—Reg. Himal.
angulare, Bunge, Enum. Pl. Chin. Bor. 49 = nudiflorum.
angulare, *Vahl, Symb. Bot.* iii. 1.—Afr. austr.
angustifolium, Ker-Gawl. in Bot. Reg. t. 521 = laurifolium.
angustifolium, Wall. Cat. n. 2873, pro parte = sessiliflorum.
angustifolium, *Willd. Sp. Pl.* i. 36; *Vahl, Enum.* i. 29.—Ind. or.
arborescens, Roj. Hort. Maurit. 203 = trifoliorum?
arborescens, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 95.—Ind. or.
arborescens, Buch.-Ham. ex Loud. Arboret. ii. 1249 = heterophyllum.
arborescens, Salisb. Mant. i. 90 = arborescens.
aristatum, Wall. Cat. n. 2858 = undulatum.
aristatum, Zipp. ex Span. in Linnæa, xv. (1841) 336 = lanciolium.
artense, *Montr. in Mém. Acad. Lyon*, x. (1860) 231.—N. Caled.
attenuatum, *Roxb. Hort. Beng.* 9, nomen; *DC. Prod.* viii. 309.—Reg. Himal.; Burma.
attenuatum, Wall. Cat. n. 2865 B = flexile.
aureum, D. Don, Prod. Fl. Nep. 106 = grandiflorum.
auriculatum, *Vahl, Symb. Bot.* iii. 1.—Ind. or.
australe, Pers. Syn. i. 8 = simplicifolium.
azoricum, Heyne, ex Wall. Cat. n. 2881 = flexile.
azoricum, Hook. & Arn. Bot. Beech. Voy. 66 = didymum.
azoricum, *Linn. Sp. Pl.* 7.—Ins. Fortunat.
bahiense, *DC. Prod.* viii. 311.—Bras.
Barrelieri, Webb & Berth. Phyt. Canar. iii. 185 = odoratissimum.
bicorollatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. 1. Art. iv. 19 = J. Sambac.
Bidevillii, Vis. in Act. Ist. Venet. Sc. Ser. III. v. (1856–59) 186 = lineare.
bifarium, *Wall. Cat.* n. 2866.—Malaya.
bignoniaceum, Wall. l. c. n. 2868 = humile.
Blancoi, Hassk. in Flora, xlvii. (1864) 49 = J. Sambac.
bracteatum, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 598 = Rottlerianum.
bracteatum, Roxb. Hort. Beng. 8; Fl. Ind. i. 94 = pubescens.
bracteatum, Wall. Cat. n. 2867 = bifarium.
brevilobum, *A. DC. in DC. Prod.* viii. 307.—Ind. or.
Burmannianum, *Blume, Mus. Bot. Lugd. Bat.* i. 279. —Zeylan.
calcareum, *F. Muell. Fragm.* i. 213.—Austral.
calophyllum, *Wall. Cat.* n. 2869.—Ind. or.
calycinum, *Wall. ex Voigt, Hort. Suburb. Calc.* 552, nomen.—Hab.?
campanulatum, Link, Jahrb. i. 111. 30 = tortuosum.
capense, Mill. Gard. Dict. ed. VIII. n. 7 = Gardenia florida.
capense, *Thunb. Prod. Fl. Cap.* 2.—Afr. austr.
carinatum, *Blume, Mus. Bot. Lugd. Bat.* i. 278.—Ins. Celebes.
catalonicum, Hort. ex DC. Prod. viii. 313 = grandiflorum.
candatum, *Wall. Cat.* n. 2854.—Reg. Himal.
chœnse, *Delile, in Rochet, Sec. Voy. Chan*, 343.—Afr. trop.
chrysanthemum, Roxb. Hort. Beng. 3; Fl. Ind. i. 99 = humile.
coarctatum, *Roxb. ll. cc.* 3; 92.—Burma.
collinum, Salisb. Prod. 12 = fruticans.

JASMINUM :—
confusum, DC. Prod. viii. 309 = simplicifolium.
congestum, Buch.-Ham. in Wall. Cat. n. 2874 = pubescens.
cordifolium, *Wall. Cat.* n. 2858.—Ind. or.
courtallense, Wight. Ic. t. 1252 = calophyllum.
crassifolium, *Blume, Bijdr.* 679.—Java.
crepidatum, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1802) 192 = rigidum.
Dallachii, F. Muell. Fragm. iv. 150 = didymum.
decussatum, *Wall. Cat.* n. 2860.—Burma.
dianthifolium, Vis. in Act. Ist. Venet. Sc. Ser. III. (1858–59) 137 = suaveissimum.
dichotomum, D. Don, Prod. Fl. Nep. 105 = glandulosum.
dichotomum, *Vahl, Symb. Bot.* i. 36.—Afr. trop.
didymum, *Forst. f. Prod.* 3.—Austral.; Ins. Pacif.
dispermum, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 99.—Reg. Himal.
distichum, *Blume, Mus. Bot. Lugd. Bat.* i. 275.—Sumatra.
divaricatum, R. Br. Prod. 521 = didymum.
elongatum, *Blume, Bijdr.* 678.—Java.
elongatum, Willd. Sp. Pl. i. 37; Roxb. ex DC. Prod. viii. 314 = Roxburghianum.
ensatum, *Blume, Mus. Bot. Lugd. Bat.* i. 277.—Ins. Celebes.
erectiflorum, A. DC. in DC. Prod. viii. 308 = cordifolium.
exsertum, Buch.-Ham. ex Wall. Cat. sub n. 2852 = Rottlerianum.
extensum, *Wall. Cat.* n. 2862.—Burma.
Finlaysonianum, *Wall. l. c.* n. 7126.—Malaya.
flavum, Sieber, ex DC. Prod. viii. 312 = revolutum.
flexile, Desf. ex DC. l. c. 310 = auriculatum.?
flexile, Jacq. Hort. Schoenb. iv. 46. t. 490 = tortuosum.
flexile, *Vahl, Symb. Bot.* iii. 1.—Ind. or.
floribundum, *R. Br. ex Preurn. in Mus. Senckenb.* ii. (1837) 168.—Abyssin.
floridum, *Bunge, Enum. Pl. Chin. Bor.* 42.—China.
fluminense, Vell. Fl. Flum. 10; t. t. 23 = bahlense.
Forsteni, *Blume, Mus. Bot. Lugd. Bat.* i. 277.—Malaya.
Forsteni, F. Muell. Fragm. vi. 86 = simulum.
fragrans, Salisb. Prod. 12 = J. Sambac.
fraternum, *Miq. Fl. Ind. Bat.* ii. 585.—Malaya.
frutescens, Gueldenst. Reisen, i. 422 = fruticans.
fruticans, *Linn. Sp. Pl.* 7.—Reg. Mediterr.; Orient.
fruticosum, Willd. in Mém. Acad. Berl. 1790–91 (1796) t. 2, sphalm. = praec.
fulvum, *Blume, Mus. Bot. Lugd. Bat.* i. 275.—Sumatra.
funale, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 405 = simplicifolium.
furostum, *Blume, Mus. Bot. Lugd. Bat.* i. 279.—Java.
Gardnerianum, Wight, Illustr. Ind. Bot. ii. 159 = trichotomum.
geminiflorum, Vent. Choix, t. 8 = simplicifolium.
glabellum, Herb. Madr. ex Wall. Cat. n. 2857 = malabaricum, arborescens.
glabriusculum, *Blume, Bijdr.* 679.—Java.
glaberum, Willd. ex Link, Jahrb. i. 111. 21 = bifarium?
glandulosum, *Wall. Cat.* n. 2849.—Reg. Himal.
glaucum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 9.—Afr. austr.
gracile, Andr. Bot. Rep. t. 137 = simplicifolium.
gracillimum, *Hook. f. in Gard. Chron.* (1881) i. 9.—Borneo.
grandiflorum, *Linn. Sp. Pl.* ed. II. 9.—Reg. Himal.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 593.—Malacca.
guineense, *G. Don, Gen. Syst.* iv. 61.—Afr. trop.
Hamiltonii, G. Don, l. c. = glandulosum.
Hasseltianum, *Blume, Mus. Bot. Lugd. Bat.* i. 278.—Java.
heterophyllum, Moench, Meth. 467 = fruticans.
heterophyllum, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 100.—Reg. Himal.
heterophyllum, *Blume, Mus. Bot. Lugd. Bat.* i. 272.—Ind. or.
Heynianum, Wall. Cat. n. 2871; G. Don, Gen. Syst. iv. 62 = J. Sambac.
hirsutum, Hoffmgg. Verz. Pfl. 69 = molle.
hirsutum, Wall. Cat. n. 2853 C = Rottlerianum.
hirsutum, Willd. Sp. Pl. i. 36 = pubescens, Gaertnerda speciosa.
hispanicum, Hort. ex DC. Prod. viii. 313 = grandiflorum.

JASMINUM :—
Horsfieldii, *Miq. Fl. Ind. Bat.* ii. 584.—Ins. Banca.
humile, Gueldenst. Reisen, i. 421 = fruticans.
humile, Heyne, ex Wall. Cat. n. 2867 = revolutum.
humile, *Linn. Sp. Pl.* 7.—As. trop.
insolorum, Jacquem. ex Decne. in Jacquem. Voy. Bot. 189 = humile.
insigne, *Blume, Mus. Bot. Lugd. Bat.* i. 278.—Sumatra.
intermedium, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 599 = trichotomum.
Jacquemontii, Jacquem. ex Decne. in Jacquem. Voy. Bot. t. 143 = humile.
Kitchingii, *Baker, in Journ. Linn. Soc.* xviii. (1881) 272.—Madag.
lactum, Wall. Cat. n. 2859 = scandens.
lanceolarium, Roxb. Hort. Beng. 8 = lanceolaria.
lanceolaria, *Roxb. l. c.*; *Fl. Ind.* i. 98.—Reg. Himal.
lanceolatum, Hort. Berol. ex Steud. Nom. ed. II. i. 798 = laurifolium.
lanceolatum, *Ruiz & Pav. Fl. Per.* i. 6. t. 7. f. a.—Peruv.
lanciflorum, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 404.—Ins. Timor.
latifolium, Buch.-Ham. ex Wall. Cat. sub n. 2852 = Rottlerianum.
latifolium, Roxb. Hort. Beng. 8; Fl. Ind. i. 95 = arborescens.
latipetalum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 599.—Burma.
laurifolium, Gamble, Darjeel. List, 55 = glandulosum.
laurifolium, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 92.—Reg. Himal.; Burma.
laurifolium, Wall. Cat. n. 2847 A = anastomosans.
Leschenaultii, A. DC. in DC. Prod. viii. 304 = bifarium.
ligustrifolium, Lam. Encyc. iii. 209 = glaucum.
ligustrifolium, Wall. ex Boj. Hort. Maurit. 204 = rigidum.
ligustrinum, *Blume, Mus. Bot. Lugd. Bat.* i. 276.—Malaya.
Lindleyanum, Blume, l. c. 272 = anastomosans?
lineare, *R. Br. Prod.* 521.—Austral.
lonchophyllum, Voigt, Hort. Suburb. Calc. 551 = lanceolaria.
Loudonianum, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 143.—Hab.?
lucidum, Banks, ex DC. Prod. viii. 309 = simplicifolium.
luteum, Gueldenst. Reisen, i. 422 = fruticans.
lusoniense, *Vidal, Phan. Cuming. Philipp.* 185.—Ins. Philipp.
macrophyllum, Hort. ex DC. Prod. viii. 312 = heterophyllum.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 594.—Penang.
malabaricum, *Wight, Ic.* t. 1250.—Ind. or.
mariannum, *DC. Prod.* viii. 307.—Ins. Marian.
mauritianum, Boj. Hort. Maurit. 204 = auriculatum.
Mesnyi, *Hance, in Journ. Bot.* xx. (1882) 87.—China.
microcalyx, *R. Br. in Sturt, Exped. Centr. Austral.* ii. App. 83.—Austral.
microcalyx, *Hance, in Journ. Bot.* xxi. (1883) 293.—China.
Mitchellii, Lindl. in Mitch. Trop. Austral. 365 = mixtinervium, *Blume, Mus. Bot. Lugd. Bat.* i. 273.—Java.
molle, R. Br. Prod. 521 = simplicifolium.
molle, Buch.-Ham. ex Wall. Cat. sub n. 2856 = arborescens.
multisinuosum, Wall. ex DC. Prod. viii. 307 = brevilobum.
montanum, Roth, Nov. Pl. Sp. 5 = arborescens.
mucronatum, Kelchb. ex Baker, Fl. Maurit. 290 = auriculatum.
mucronatum, Roth, Nov. Pl. Sp. 6 = tortuosum.
multiflorum, Andr. Bot. Rep. t. 496 = pubescens?
multiflorum, Roth, Nov. Pl. Sp. 8 = pubescens.
multipartitum, *Hochst. in Flora*, xxvii. (1844) 825.—Afr. austr.
myrtifolium, DC. Prod. viii. 305 = glaucum.
myrtophyllum, Zenk. Fl. Ind. 5. t. 7 = rigidum.
Nanqueulica, Buch.-Ham. in Wall. Cat. sub n. 2856 = arborescens.
nepalense, Sprong. Syst. iv. Cur. Post. 12 = glandulosum.
nervosum, *Lour. Fl. Cochinch.* 20.—Cochinch.

JASMINUM :—
nilagiricum, Hohen. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 600 = brevilobum.
nobile, *C. B. Clarke, l. c.* 597.—Burma.
nocti florum, *Afzel. Remed. Guin.* 25.—Afr. trop.
nudiflorum, *Lindl. in Trans. Hort. Soc.* i. (1846) 168.—China.
nummularioides, Blume, Mus. Bot. Lugd. Bat. i. 275 = bifarium.
oblongum, *Burm. f. Fl. Ind.* 6. t. 8. f. 2.—Ind. or.
odoratissimum, *Linn. Sp. Pl.* 7.—Ins. Fortunat.
odoratum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = J. Sambac.
odorum, Salisb. Prod. 12 = odoratissimum.
officinale, *Linn. Sp. Pl.* 7.—Ind. bor. occ.; China.
ovalifolium, Wight, Ic. t. 1256 = auriculatum.
ovatum, Wall. Cat. n. 2855 = flexile, caudatum.
paniculatum, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 99.—China.
parviflorum, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 405 = didymum.
pauciflorum, *Benth. in Hook. Niger Fl.* 448.—Afr. trop.
pedale, *Blume, Mus. Bot. Lugd. Bat.* i. 274.—Java.
pedatum, *Blume, l. c.* 276.—Java.
Perrottetianum, A. DC. in DC. Prod. viii. 304 = rigidum?
populifolium, *Blume, Mus. Bot. Lugd. Bat.* i. 276.—Ins. Philipp.
puberulum, *Baker, in Journ. Linn. Soc. xx.* (1883) 203.—Madag.
pubescens, Buch.-Ham. ex Wall. Cat. sub n. 2850 = J. Sambac.
pubescens, DC. Prod. viii. 313 = pubigerum.
pubescens, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 598 = Rottlerianum.
pubescens, *Willd. Sp. Pl.* i. 37.—As. trop.
pubigerum, *D. Don, Prod. Fl. Nep.* 106.—Reg. Himal.
punctatum, Buch.-Ham. in Wall. Cat. n. 2877 = arborescens.
purpureum, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 598 = J. Ritchiei.
quadrifolium, Buch.-Ham. ex Wall. Cat. n. 2868 = J. Sambac.
quinquefloruin, Heyne, ex Wall. l. c. n. 2878 = J. Sambac.
quinquenervum, Lamb. ex D. Don, Prod. Fl. Nep. 106 = dispermum.
quinquenervium, *Blume, Mus. Bot. Lugd. Bat.* i. 278.—Java.
racemosum, *F. Muell. Fragm.* i. 19.—Austral.
reticulatum, Wall. Cat. n. 2869 = coarctatum.
revolutum, Sims, Bot. Mag. t. 1731 = humile.
rigidum, Thw. Enum. Pl. Zeyl. 190 = Rottlerianum.
rigidum, *Zenk. Pl. Ind.* 5. t. 6.—Ind. or.
Ritchiei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 598.—Ind. or.
rivulare, Salisb. Ic. 15. t. 8 = glaucum.
Rottlerianum, *Wall. Cat.* n. 2865.—Ind. or.
rpzndiflorum, *Baif. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 77.—Hanoi.—Ins. Socotr.
Roxburghianum, *Wall. Cat.* n. 2870.—Ind. or.
rupestre, *Blume, Mus. Bot. Lugd. Bat.* i. 280.—N. Guinea.
Sambac, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 8.—As. trop.
Sanjurium, Buch.-Ham. ex DC. Prod. viii. 302 = quinquefolium.
scandens, Griff. Itin. Notes, 103 = undulatum.
scandens, *Vahl, Symb. Bot.* iii. 2.—Ind. or.
Schimperi, *Vatke, in Linnaea*, xl. (1876) 210.—Abyssin.
scutiflorum, *Vahl, Enum.* i. 29.—Ind. or.
Siebeldianum, Blume, Mus. Bot. Lugd. Bat. i. 280 = undiflorum.
silhetense, Blume, l. c. 271 = anastomosans.
simplicifolium, *Forst. f. Prod.* 3.—Austral.; Ins. Pacif.
smilacifolium, *Griff. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 600.—Malacca.
solenanthum, Dec. Prod. viii. 306 = bifarium?
stenopetalum, Lindl. Bot. Reg. sub t. 918 = anastomosans.
stenptopus, *E. Mey. ex DC. Prod.* viii. 307.—Afr. austr.
suaveolens, Salisb. Prod. 12 = asiaticum.
suaveissimum, *Lindl. in Mitch. Journ. Trop. Austral.* 365.—Austral.
subulosigatum, *Blume, Mus. Bot. Lugd. Bat.* i. 276.—Malaya.

JASMINUM:—
subglanduloum, *Kurz, in Journ. Bot.* xiii. (1875) 329. —Burma.
subpubescens, *Blume, Mus. Bot. Lugd. Bat.* i. 274.—Java.
subtriplinerve, *Blume, l. c.* 272.—Reg. Himal.
subulatum, Lindl. Bot. Reg. (1842) Misc. 57 = floridum.
sylvestre, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. J. Art. iv. 19 (Quid ?).—Java.
syriacum, Boiss. & Gaill. Diagn. Ser. II. vi. 120 = fruticans.
syringæfolium, *Wall. Cat.* n. 2861.—Burma; Assam.
tenuifolium, *Regel, Gartenfl.* (1886) 330.—Hab. ?
tetraphyx, Wight & Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 92 = rigidum.
tetraquetrum, *A. Gray, in Proc. Am Acad.* v. (1861–62) 382.—Ins. Fiji.
tettense, Klotzsch, in Peters, Reise Mossamb. Bot. 284 = auriculatum.
tortuosum, *Willd. Enum. Hort. Berol.* 10.—Afr. austr.
transcorvense, Heyne, ex Wall. Cat. sub n. 2881 = flexile.
trichotomum, *Heyne, ex Roth, Nov. Pl. Sp.* 6.—Ind. or.
trifoliatum, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 596 = rigidum.
trifjorum, Pers. Syn. i. 7 = angustifolium.
trifolitatum, Moench, Meth. 467 = azoricum.
trifolitatum, Pers. Syn. i. 7 = auriculatum.
trinerve, Roxb. Hort. Beng. 8; Fl. Ind. i. 93 = ane stomosans.
trinerve, *Vahl, Symb. Bot.* iii. 2—Java.
trinerve, Wall. Cat. n. 2845, partim = subtriplinerve.
tubiflorum, *Roxb. Hort. Beng.* [80]; *Fl. Ind.* i. 97.—Malaya.
undulatum, Hort. Heyne, ex Wall. Cat. n. 2871 = J. Sambac.
undulatum, *Ker-Gawl. in Bot. Reg.* t. 436.—Reg. Himal.; China.
undulatum, *Willd. Sp. Pl.* i. 36.—Malaya.
venustum, *Regel, Cat. Pl. Hort. Aksakow.* 75.—Hab. ?
villosum, Stokes, Bot. Comment. i. 18 = Roxburghianum.
viminale, Salisb. Prod. 12 = officinale.
vimineum, Willd. Sp. Pl. i. 36 = angustifolium.
volubile, *Jacq. Hort. Schoenb.* iii. 39.—Afr. austr.
vulcanicum, *Blume, Mus. Bot. Lugd. Bat.* i. 276.—Malaya.
vulgatius, Lam. Fl. Fr. ii. 806 = officinale.
Waltzianum, *Blume, Mus. Bot. Lugd. Bat.* i. 277.—Sambac.
Wallichianum, Lindl. Bot. Reg. t. 1409 = humile.
Wightii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 596.—Ind. or.
Zambac, Roxb. Hort. Beng. 3 = J. Sambac.
zanzibaricum, Boj. ex DC. Prod. vii. 810 = auriculatum.
zeylanicum, Burm. f. Fl. Ind. t. 39 = Tabernæmontana coronaria.
Zippelianum, *Blume, Mus. Bot. Lugd. Bat.* i. 279.—Ins. Amboin.

JASONIA, Cass. in Bull. Soc. Philom. (1815) 175; et in Dict. Sc. Nat. xxiv. 200 (1822). *COMPOSITÆ*, Benth. & Hook. f. ii. 382.
CHILIADENUS, Cass. in Dict. Sc. Nat. xxxiv. 34 (1825).
MYRIADENUS, Cass. in Bull. Soc. Philom. (1817) 188.
ORZINA, Bertol. in Ann. Stor. Nat. Bologn. I. ii. (1829) 862.
dichotoma, DC. Prod. v. 477 = Allagopappus dichotomus.
discoidea, Cass. in Dict. Sc. Nat. xxiv. 201; xxxix. 407 = Pulicaria sicula.
fœtida, Cass. in Bull. Soc. Philom. (1815) 175 = Nidorella foliosa.
glutinosa, DC. Prod. v. 476 = Inula viscosa.
hirrigata, *DC. l. c.* 477.—Ins. Teneriff.
longifolia, *Cass. in Bull. Soc. Philom.* (1815) 175.—Am. bor.
radiata, *Cass. in Dict. Sc. Nat.* xxiv. 201.—Europ. austr.
rupestris, *Pomel, Nouv. Mat. Fl. Atl.* 37.—Alger.
sanatilis, Cass. Fl. Sic. Syn. ii. 488 = glutinosa.
sicula, DC. ex Decne. in Ann. Sc. Nat. Bot. II. ii. (1834) 261 = Pulicaria sicula.
tuberosa, *DC. Prod.* v. 476.—Europ. austr.

JATEORHIZA, Miers, in Hook. Niger Fl. 212 (1849). *MENISPERMACEÆ*, Benth. & Hook. f. i. 34.
DISCIPHANIA, Eichl. in Mart. Fl. Bras. xiii. i. 168. t. 36 (1864).
Calumba, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiii. (1864) 134 = seq
Columba, *Miers, in Hook. Niger Fl.* 214, in nota.—Afr. trop.
lobata, Herb. Monac. ex Eichl. in Mart. Fl. Bras. xiii. i. 168 = Disciphania lobata.
Miersii, *Oliver, Fl. Trop. Afr.* i. 42.—Afr. trop.
palmata, Miers, in Hook. Niger Fl. 214. t. 18, in nota = J. Columba.
strugosa, *Miers, l. c.* 213.—Afr. trop.

JATROPHA, Scop. Introd. 248 (1777) = seq.

JATROPHA, Linn. Syst. ed. 7 (1785). *EUPHORBIACEÆ*, Benth. & Hook. f. iii. 290.
ADENOPIUM, Pohl, Ic. Pl. Bras. i. 12 t. 9 (1827).
CASTIGLIONIA, Ruiz & Pav. Prod. 139. t. 37 (1794).
CNIDOSCOLUS, Pohl, Ic. Pl. Bras i. 56. t. 49 (1827).
CURCAS, Adans. Fam. ii. 356 (1763).
JUSSIEVIA, Houst. Reisig. 6. t. 15 (1781).
LOUKRIRA, Cav. Ic. v. 17. t. 429 (1799).
MAZINNA, Spach, Hist. Vég. Phan. ii. 487 (1834).
MOLINNA, Orteg. Hort. Matr. Dec. 194. t. 13 (1797).
acerifolia, Salisb. Prod. 389 = J. Curcas.
aconitifolia, *Mill. Gard. Dict.* ed. VIII. n. 6.—Am. trop.
acuminata, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* iv. 76.—Arab.
acuminata, Desr. in Lam. Encyc. iv. 8 = hastata.
acuminata, Steud. Nom. ed. II. i. 799 = Manihot aesculifolia.
aethiopica, *Muell. Arg. in Flora*, xlvii. (1864) 485.—Afr. trop.
alamosana, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 207.—Mexic.
Andrieuxii, *Muell. Arg. l. c.* 208.—Mexic.
angustifolens, *Muell. Arg. in DC. Prod.* xv. ii. 1102.—Mexic.
angustifolia, *Griseb. in Goett. Nachr.* (1865) 171.—Cuba.
angustifolia, Steud. Nom. ed. II. i. 799 (nom. delend.).—Cuba.
anomala, Steud. l. c. = Manihot anomala.
antisiphilitica, *Spegz. in Anal. Soc. Cient. Argent.* xvi. (1883) 91.—Reg. Argent.
arcuata, Steud. Nom. ed. II. i. 799 = Manihot violacea.
australis, Ledd. ex G. Don, in Loud. Hort. Brit 891 = Sterculia diversifolia, *G. Don.*
Berlandieri, *Torr. Bot. Mex. Bound.* 193.—Mexic.
Berterii, *Spreng. Syst.* iii. 76.—N. Granat.
cajanifolia, Steud. Nom. ed. II. i. 799 = Manihot tripartita.
calyculata, Vahl, ex Steud. l. c. (nom. fals.).
canescens, *Muell. Arg. in DC. Prod.* xv. ii. 1079.—Calif.
capensis, *Sond. in Linnæa*, xxiii. (1850) 118.—Afr. austr.
cardiophylla, *Muell. Arg. in DC. Prod.* xv. ii. 1079.—Mexic.
caricæfolia, Steud. Nom. ed. II. i. 799 = Manihot caricæfolia.
carthagenensis, Jacq. Enum. Pl. Carib. 32 = Manihot carthagenensis.
ceroxylifolia, Jacq. Enum. Pl. Carib. 32 = Manihot violacea.
ciliata, *Cave. in Supl. Gas. Lit. Mexico* (Jul. 2, 1794) 4.—Mexic.
ciliata, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 209.—Peruv.
cinerea, *Muell. Arg. in DC. Prod.* xv. ii. 1079.—Mexic.
clavuligera, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 209.—Bolivia.
stenocarpifolia, Muell. Arg. in Mart. Fl. Bras. xi. ii. 476 = Manihot tripartita.
coccinea, *Hort. Cels, ex Link, Enum. Hort. Berol.* ii. 406.—Hab. ?
corvulus, Steud. Nom. ed. II. i. 799 = Manihot coeruliscens.
coeruliscens, Muell. Arg. in Mart. Fl. Bras. xi. ii. 469 = Manihot coeruliscens.
Condor, Wall. Cat. n. 2799 = J. Curcas.
cordata, *Muell. Arg. in DC. Prod.* xv. ii. 1078.—Mexic.

JATROPHA:—
crinita, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 207. —Afr. trop.
crotalariæformis, Steud. Nom. ed. II. i. 799 = Manihot crotalariæformis.
Curcas, *Linn. Sp. Pl.* 1006.—Amphig. trop.
Curcas, Wall. Cat. n. 7799 D = Wightiana.
dalechampiæformis, Steud. Nom. ed. II. i. 799 = Manihot tripartita.
diffusa, Steud. l. c. = Manihot palmata.
digitiformis, Steud. l. c. = Manihot digitiformis.
dioica, *Cerv. in Supl. Gas. Lit. Mexico* (Jul. 2, 1794) 4.—Mexic.
divaricata, *Sw. Prod. Veg. Ind. Occ.* 98.—Ind. occ.
divergens, Baill. Adansonia, iv. (1863–64) 268 = Pohliana.
divergens, Steud. Nom. ed. II. i. 799 = Manihot violacea.
diversifolia, *A. Rich. in Sagra, Fl. Cub. Fanerog.* ii. 207. —Cuba.
diversifolia, Steud. Nom. ed. II. i. 799 (nom. delend.).
dulcis, [J. F. Gmel.] Onomat. Bot. v. 7 = Manihot palmata.
edulis, *Cerv. in Supl. Gas. Lit. Mexico* (Jul. 2, 1794) 3.—Mexic.
elastica, Linn. f. Suppl. 422 = Hevea guyanensis.
elegans, Klotzsch, in Seem. Bot. Voy. Herald, 103 = gossypifolia.
elliptica, Muell. Arg. in Mart. Fl. Bras. xi. ii. 489 = opifera.
excisa, *Griseb. in Goett. Abh.* xix. (1874) 94.—Reg. Argent.
ferox, *Muell. Arg. in Mart. Fl. Bras* xi. ii. 497.—Bras *Paschers*, Steud. Nom. ed. II. i. 799 (nom. delend.).
flabellifolia, Steud. l. c. = Manihot palmata.
foetida, Steud. l. c. = Manihot foetida.
fragrans, *H. B. & K. Nov. Gen. et Sp.* ii. 105.—Cuba.
frutescens, Auct. ex Steud. Nom. ed. II. i. 799 = J. Janipha.
gallabatensis, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 661.—Afr. trop.
glandulifera, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 688. —As. et Afr. trop.
glandulosa, *Vahl, Symb. Bot.* i. 80.—Arab.
glauca, Griseb. in Goett. Nachr. (1865) 170 = angustifolia.
glauca, A. Rich. Tent. Fl. Abyss. ii. 250 = lobata.
glauca, Salisb. Prod. 388 = J. Manihot.
glauca, Vahl, Symb. Bot. i. 78 = glandulifera.
globosa, Gaertn. Fruct. ii. 122 = Toxicodendrum capense.
gossypifolia, *Linn. Sp. Pl.* 1008.—Am. et Afr. trop.
gracilis, Steud. Nom. ed. II. i. 799 = Manihot gracilis.
guaranitica, *Speg. in Anal. Soc. Cient. Argent.* xvi. (1883) 93.—Reg. Argent.
hamosa, *Muell. Arg. in DC. Prod.* xv. ii. 1097.—Bras.
hastata, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 158.—Manihot diversifolia.
hastata, *Jacq. Enum. Pl. Carib.* 32.—Cuba.
herbacea, Linn. Sp. Pl. 1007; Desr. in Lam. Encyc. iv. 16 = urens.
hemandiæfolia, *Vent. Jard. Malm.* 62.—Am. trop.
heterophylla, Steud. Nom. ed. II. i. 799 = Manihot heterophylla.
Heudelotii, Baill. Adansonia, i. (1860) 64 = Ricinodendron africanum.
hirsuta, *Hochst. in Flora*, xxviii. (1845) 82.—Afr. austr.
horrida, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 210. —Bras.
integerrima, Jacq. Enum. Pl. Carib. 32 = Manihot diversifolia.
Isabelli, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 489.—Bras.
Jacquini, Baill. Adansonia, iv. (1863–64) 268 = gossypifolia.
Janipha, Blanco, Fl. Filip. ed. I. 758 = multifida.
Janipha, Linn. Mant. i. 126 = Manihot carthagenensis.
Janipha, Lour. Fl. Cochinch. 585 = Manihot Loureiri.
Kuntiana, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 211.—Mexic.
Lacerti, *Silva Manso, Enum. Subst. Bras.* 8.—Bras.
laciniosa, Mart. Fl. Bras. xi. ii. 488 = Manihot laciniosa.
lagarithoides, *Sond. in Linnæa*, xxiii. (1850) 118.—Afr. austr.
lanciniosa, Steud. Nom. ed. II. i. 799 = Manihot laciniosa.
Liebmanni, *Muell. Arg. in Linnæa*, xxxiv. (1865–66) 212.—Mexic.

JATROPHA:—
lobata, *Muell. Arg. in DC. Prod.* xv. ii. 1065.—Arab.
longepetiolata, Steud. Nom. ed. II. i. 799 = Manihot longepetiolata.
Lourairii, Steud. l. c. = Manihot Lourairii.
luxurians, Baill. Adansonia, iv. (1863–64) 268 = Pohliana.
macrantha, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 209.—Peruv.
macrocarpa, *Griseb. in Goett. Abh.* xix. (1874) 94.—Reg. Argent.
macrorhiza, *Benth. Pl. Hartw.* 8.—Mexic.
Manihot, Linn. Sp. Pl. 1007 = Manihot utilissima.
Manihot, Vell. Fl. Flum. x. t. 80 = Manihot palmata.
Martiusii, *Baill. Adansonia*, iv. (1863–64) 268.—Bras.
mitis, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 301 = Manihot palmata.
mollissima, Baill. Adansonia, iv. (1863–64) 268 = Pohliana.
moluccana, Linn. Sp. Pl. 1006 = Aleurites triloba.
moluccana, Wall. Cat. n. 7799 = J. Curcas.
montana, Willd. Sp. Pl. iv. 563 = Baliospermum axillare.
multifida, *Linn. Sp. Pl.* 1006.—Amphig. trop.
mutabilis, *Baill. Adansonia*, iv. (1863–64) 267.—Bras.
nana, *Dais. et Gibs. Bomb. Fl.* 229.—Ind. or.
napaeifolia, Desr. in Lam. Encyc. iv. 15 = acanitifolia.
nalaleuia, *Muell. Arg. in Flora*, xlvii. (1864) 485.—Afr. austr.
neglecta, Houst. ex Baill. Etud. Gén. Euph. 304, nomen = urens.
neriifolia, *Muell. Arg. in Flora*, xlvii. (1864) 486.—Afr. trop.
nudicaulis, *Benth. Bot. Voy. Sulph.* 165.—Am. trop.
obtusifolia, *Muell. Arg. in DC. Prod.* xv. ii. 1097.—Bras.
octandra, *Cerv. in Supl. Gas. Lit. Mexico* (Jul. 2, 1794) 4.—Mexic.
officinalis, Mart. ex Pohl, Ic. Pl. Bras. i. 13. t. 9 = opifera.
officinalis, *Pohl, l. c.* 13.—Bras.
oligandra, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 502.—Bras.
olivacea, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 207.—Mexic.
opifera, *Mart. Reise Bras.* i. 548.—Am. trop.
orbicularis, Steud. Nom. ed. II. i. 799 = Manihot orbicularis.
osteocarpa, Schott, ex Pohl, Ic. Pl. Bras. i. 63 = urens.
palmata, Vell. Fl. Flum. x. t. 81 = Manihot palmata.
palmata, Willd. Sp. Pl. iv. 562 = acanitifolia.
pandurae folia, Andr. Bot. Rep. t. 267 = hastata.
Papaya, Medic. Beobacht. (1782) 194 = acanitifolia.
pauciflora, *Wright, ex Griseb. in Goett. Nachr.* (1865) 170.—Cuba.
paviaefolia, Steud. Nom. ed. II. i. 799 = Manihot paviaefolia.
pelargoniaefolia, Courb. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 150 = glandulosa.
peltata, *Cerv. in Supl. Gas. Lit. Mexico* (Jul. 2, 1794) 3; *H. B. & K. Nov. Gen. et Sp.* ii. 104.—Am. trop.
peltata, Steud. Nom. ed. II. i. 800 = Manihot peltata.
peltata, Wight, Ic. t. 1169 = Wightiana.
pentaphylla, Steud. Nom. ed. II. i. 800 = Manihot pentaphylla.
peruviana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 210.—Peruv.
phyllacantha, *Muell. Arg. in DC. Prod.* xv. ii. 1098.—Bras.
pilosa, Steud. Nom. ed. II. i. 800 = Manihot pilosa.
platyphylla, *Muell. Arg. in DC. Prod.* xv. ii. 1077.—Mexic.
podagrica, *Hook. Bot. Mag.* t. 4376.—Panama.
Pohliana, *Muell. Arg. in Mém. Soc. Phys. Genèv.* xvii. ii. (1864) 449.—Bras.
porrecta, Steud. Nom. ed. II. i. 800 = Manihot tripartita.
pronifolia, Steud. l. c. = Manihot pronifolia.
pruinosa, Steud. l. c. = Manihot pruinosa.
Pseudo-curcas, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 206.—Mexic.
pubescens, Steud. Nom. ed. II. i. 800 = Manihot pubescens.
pungens, Forsk. Fl. Aegypt. Arab. 168 = Tragia cordifolia.

JATROPHA:—
purpureo-costata, Steud. Nom. ed. II. i. 800 = Manihot purpureo-costata.
pusilla, Steud. l. c. = Manihot pusilla.
quinquefolia, Steud. l. c. = Manihot quinquefolia.
quinquefoliata, Steud. l. c. = prace.!
quinquelobata, *Cerv. in Supl. Gas. Lit. Mexico* (Jul. 2, 1794) 4.—Mexic.
quinquelobata, Steud. Nom. ed. II. i. 800 = Manihot quinquelobata.
quinquelobata, *Mill. Gard. Dict.* ed. VIII. n. 2.—Cuba.
reniformis, Steud. Nom. ed. II. i. 800 = Manihot reniformis.
ribifolia, Baill. Adansonia, iv. (1863–64) 268 = gossypifolia.
ricinifolia, Fenzl, ex Baill. l. c. i. (1860) 62 = lobata.
rotundifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 211.—Mexic.
Sabdariffa, *Schweinf. Britr. Fl. Aethiop.* 87.—Afr. trop.
sagittato-partita, Steud. Nom. ed. II. i. 800 = Manihot sagittato-partita.
salicifolia, Steud. l. c. = Manihot salicifolia.
silvestris, Vell. Fl. Flum. x. t. 83 = Manihot palmata.
sinuata, Steud. Nom. ed. II. i. 800 = Manihot sinuata.
sparsifolia, Steud. l. c. = Manihot sparsifolia.
spathulata, *Muell. Arg. in DC. Prod.* xv. ii. 1081.—Mexic.
spinosa, *Vahl, Symb. Bot.* i. 79.—Arab.
stephanifolia, Mill. Gard. Dict. ed. VIII. n. 9 = gossypifolia.
stimulosa, Michx. Fl. Bor. Am. ii. 216 = urens.
stipulata, Vell. Fl. Flum. x. t. 82 = Manihot utilissima.
stipulosa, Steud. Nom. ed. II. i. 800 = *stimulosa*.
sylvestris, Auct. = *silvestris*.
tenerrima, Steud. Nom. ed. II. i. 800 = Manihot gracilis.
tenuifolia, Steud. l. c. = Manihot gracilis.
texana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 211.—Texas.
tomentella, Steud. Nom. ed. II. i. 800 = Manihot tripartita.
tomentosa, Spreng. Syst. iii. 77 = Vitex cymosa.
tomentosa, Steud. Nom. ed. II. i. 800 = Manihot tomentosa.
triloba, *Cerv. in Supl. Gas. Lit. Mexico* (Jul. 2, 1794) 4.—Mexic.
tripartita, Spreng. Syst. iii. 76 = Manihot tripartita.
trip hylla, Steud. Nom. ed. II. i. 800 = Manihot triphylla.
tuberosa, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 219.—Mexic.
tupifolia, *Griseb. in Goett. Nachr.* (1865) 170.—Cuba.
unicostata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 94.—Ins. Socotr.
urens, H. B. & K. Nov. Gen. et Sp. ii. 106 = Knuthiana.
urens, *Linn. Sp. Pl.* 1007.—Am. trop.
varians, Steud. Nom. ed. II. i. 800 = Manihot varians.
variegata, *Vahl, Symb. Bot.* i. 79.—Arab.
villosa, Baill. Adansonia, iv. (1863–64) 268 = Pohliana.
villosa, Muell. Arg. in DC. Prod. xv. ii. 1085 = glandulosa.
villosa, Wight, Ic. t. 1159 = Wightiana.
viminea, *Rets. ex Steud. Nom.* ed. I. 426.—Hab.?
violacea, Steud. l. c. ed. II. i. 800 = Manihot violacea.
vitifolia, *Mill. Gard. Dict.* ed. VIII. n. 5.—Bras.
Weddelliana, *Baill. Adansonia*, iv. (1863–64) 267.—Paraguay.
Wightiana, *Muell. Arg. in DC. Prod.* xv. ii. 1080.—Ind. or.
Zeyheri, *Sond. in Linnaea*, xxiii. (1850) 117.—Afr. austr.

JATROPS, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 93 = **Maregravia**!, Plum. (Ternstroem.).
myristicoides, Rottb. l. c. (Quid?).—Guian.
umbellatus, Rottb. l. c. = M. umbellata!

JAUBERTIA, Guill. in Ann. Sc. Nat. Sér. II. xvi. (1841) 60 = **Guillonia**, A. Rich. (Rubiac.).
Auchers, Guill. l. c. = G. Auchers.

JAUBERTIA, Spach, ex Jaub. & Spach, Illustr. iii. 131. t. 289 (1849) = **Diptelocoma**, Pisch. (Compos.).
hookeriaeides, Spach, l. c. 133 = pusilla.

JAUMEA, Pers. Syn. ii. 397 (1807). *COMPOSITAE*, Benth. & Hook. f. ii. 397.
CHARTHYRSUS, Hook. & Arn. Bot. Beech. Voy. 298. t. 62 (1841).
COINOGYNE, Less. in Linnaea, vi. (1831) 520.
ESPEJOA, DC. Prod. v. 660 (1836).
HYPERICOPHYLLUM, Steetz, in Peters, Reise Mossamb. Bot. 498 t. 50 (1863).
KLEINIA, Juss. in Ann. Mus. Par. ii. (1803) 424. t. 61. f. 1.
camosa, *A. Gray, Bot. U. St. Expl. Exped.* 360.—Calif.
compositarum, *Benth. & Hook. f. ex Oliver, Fl. Trop. Afr.* iii. 395.—Afr. trop.
linearifolia, DC. Prod. v. 663 = seq.
linearis, *Pers. Syn.* ii. 397.—Bras.
mexicana, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 209.—Mexic.
pedunculata, *Benth. & Hook. f. ex Oliver, Fl. Trop. Afr.* iii. 395, in nota.—Mexic.

JEANNERETIA, Gaudich. Bot. Voy. Bonite, t. 25 (1844–52) = **Pandanus**, Linn.
littoralis, Gaudich. l. c. = P. littoralis.

JEFFERSONIA, Bart. in Trans. Am. Phil. Soc. iii. (1793) 334. *BERBERIDEAE*, Benth. & Hook. f. i. 44.
PLAGIORHEGMA, Maxim. Prim. Fl. Amur. 34. t. 1 (1859).
VINDICTA, Rafin. Fl. Tellur. ii. 51 (1836).
Bartonia, Michx. Fl. Bor. Am. i. 237 = binata.
binata, *Bart. in Trans. Am. Phil. Soc.* iii. (1793) 334.—Am. bor.
diphylla, Pers. Syn. i. 418 = binata.
dubia, *Benth. & Hook. f. ex Baker & Moore, in Journ. Linn. Soc.* xvii. (1879) 377.—Manchur.
lobata, Nutt. in Journ. Acad. Phil. vii. (1834) 99 = binata.
manchuriensis, Hance, in Journ. Bot. xvii. (1880) 258 = dubia.
odorata, Rafin. Med. Fl. ii. 11 = binata.

JEFFERSONIA, Brickell, in Med. Repos. N. York, i. (1800) 555 = **Gelsemium**, Juss. (Logan.).
sempervirens, Brickell, l. c. = G. sempervirens.

JEHLIA, Hort. Germ. ex Planch. in Fl. des Serres, Sér. I. vii. (1851–52) 177 = **Lopezia**, Cav. (Onagrar.).
fuchsioides, Hort. Germ. ex Planch. l. c. = L. macrophylla.

JELLIUM, Hort. ex Regel, Gartenfl. (1869) 102 = **Telanthera**, R. Br. (Amarant.).
tricolor, Hort. ex Regel, l. c. = T. Bettzichiana.

JENKINSIA, Griff. in Calc. Journ. Nat. Hist. iv. (1843) 231. t. 19 = **Miquelia**, Meisn. (Olacin.).
assamica, Griff. l. c. = M. Kleinii.

JENKINSIA, Wall. ex Volgt, Hort. Suburb. Calc. 544 (1845) = **Myriopteron**, Griff. (Asclepiad.).
cristata, Wall. l. c. = M. paniculatum.

JENKINSONIA, Sweet, Geran. t. 79 (1820) = **Pelargonium**, L'Hérit. (Geraniac.).
anemonefolia, Sweet, Hort. Brit. ed. I. 84 = P. myrrhifolium.
antidysenterica, Eckl. & Zeyh. Enum. 70 = P. antidysentericum.
bullata, Sweet, Hort. Brit. ed. I. 84 = P. myrrhifolium.
canariensis, Sweet, l. c. = P. canariense.
cancellifolia, Sweet, l. c. = P. cancellifolium.
coriandrifolia, Sweet, Geran. sub t. 99 = P. myrrhifolium.
dichotoma, Eckl. & Zeyh. Enum. 70 = P. dichotomum.
lacera, Sweet, Geran. sub t. 342; Hort. Brit. ed. I. 84 = P. myrrhifolium.
longicaule, Hoffmgg. Verz. Pfl. Nachtr. iii. 44 = P. longicaule.
multicaule, Hoffmgg. Verz. Pfl. 69 = P. multicaule.
myrrhifolia, Sweet, Hort. Brit. ed. I. 84 = P. myrrhifolium.
pendula, Sweet, Geran. t. 188 = P. pendulum.
quinata, Sweet, l. c. t. 79 = P. quinatum.
Synnoti, Sweet, l. c. t. 343 = P. Synnoti.
tetragona, Sweet, l. c. t. 99 = P. tetragonum.
tripartita, Hoffmgg. Verz. Pfl. 69 = P. tripartitum.

JENSOA, Rafin. Fl. Tellur. iv. 38 (1836) = **Cymbidium**, Sw. (Orchid.).
ensata, Rafin. l. c. = C. ensifolium.

JERDONIA, Wight, Ic. t. 1362 (1848). *GESNERACEAE*, Benth. & Hook. f. ii. 1020.
Indica, *Wight, l. c.*—Ind. or.

JERONIA, Prltz. Ic. Ind. i. 579 (1855), sphalm. = **Feronia**, Correa (Rutac.).

JESSENIA, Karst. in *Linnaea*, xxviii. (1856) 887. *PALMAE*, Benth. & Hook. f. iii. 897.
amazonum, *Drude, in Mart. Fl. Bras.* iii. ii. 474.—Bras.
oligocarpa. *Griseb. & H. Wendl. ex Griseb. Fl. Brit. W. Ind.* 516.—Ind. occ.
polycarpa, *Karst. in Linnaea*, xxviii. (1856) 888.—Bras.
repanda, *Engl. l. c.* xxxiii. (1864–65) 691.—N. Granat.

JESSIANA, H. Wendl. in Kerch. Palm. 247 (1878) = prsec.

JEZABEL, Banks, ex Saliab. Gen. Pl. Fragm. 5 (1866) = **Freycinetia**, Gaudich. (Pandan.).

JIMENSIA, Rafin. Fl. Tellur. iv. 38 (1836) = **Bletia**, Ruiz & Pav. (Orchid.).
nervosa, Rafin. l. c. = B. hyacinthina.

JIRASECKIA, Dam. Fl. Belg. 29 (1827) = seq.

JIRASEKIA, F. W. Schmidt, Neue a. Selt. Pfl. 49 (1793) = **Anagallis**, Linn. (Primulac.).
alpina, F. W. Schmidt, l. c. = A. tenella.
alternifolia, Reichb. ex Steud. Nom. ed. II. i. 802 = A. alternifolia.
crassifolia, Reichb. ex Steud. l. c. = A. crassifolia.
tenella, Reichb. Fl. Germ. Excurs. 408 = A. tenella.

JOACHIMEA, Benth. & Hook. f. Gen. iii. 1100 (1883), sphalm. = seq.

JOACHIMIA, Tenore, ex Roem. & Schult. Syst. ii. 695 (1817) = **Bookmannia**, Host (Gramin.).
phalaroides, Tenore, ex Roem. & Schult. l. c. = B. ercaeformis.

JOANNEA, Spreng. Anleit. ii. ii. 535 (1818) = **Chuquiraga**, Juss. (Compos.).
brasiliensis, Spreng. Syst. iii. 382 = C. Jussieui.
trplinervis, Spreng. ex Less. in Linnaea, v. (1830) 251 = C. paniculata.

JOANNESIA, Pers. Syn. ii. 388 (1807) = **Chuquiraga**, Juss. (Compos.).
insignis, Pers. l. c. = C. Jussieui.

JOANNESIA, Vell. Alograf. 199 (1798). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 290.
ANDA, A. Juss. in Dict. Sc. Nat. ii. 118 (1804); et Euph. Tent. 39. t. 12 (1824).
ANDICUS, Vell. Fl. Flum. 80 (1825).
Princeps, *Vell. Alograf.* 199.—Bras.

JOBAPHES, Phil. Fl. Atac. 27. t. 4 (1860) = **Hyalis**, D. Don (Compos.).
virgatus, Phil. l. c. 28.—Chili.

JOBINIA, Fourn. in Mart. Fl. Bras. vi. iv. 327 (1866). *ASCLEPIADEAE*.
bernardiflia, *Fourn. l. c.*—Bras.
Lindbergii, *Fourn. l. c.*—Bras.

JOCASTE, Kunth, Enum. Pl. v. 154 (1850) = **Smilacina**, Desf. (Liliac.).
purpurea, Kunth, l. c. = S. purpurea.

JOCASTE, Meisn. Gen. 211; Comm. 185 (1839) = **IOCASTE**, E. Mey. = **Phymaspermum**, Less. (Compos.).

JOCAVENA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 62, sphalm. = **Tocoyena**, Aubl. (Rubiac.).

JOHANNESIA, Endl. Gen. 1114 (1840) = **Joannesia**, Vell. (Euphorb.).

JOHANNIA, Willd. Sp. Pl. iii. 1705 (1803) = **Chuquiraga**, Juss. (Compos.).
brasiliensis, Steud. Nom. ed. II. i. 818 = C. glabra.
elegans, Willd. ex DC. Prod. vii. 9 = C. lanciflola.
imaginis, Willd. Sp. Pl. iii. 1705 = C. Jussieui.
lancifolia, Steud. Nom. ed. II. i. 813 = C. lanciflola.
microphylla, Steud. l. c. ed. I. 433 = C. Jussieui.

JOHNIA, Roxb. Hort. Beng. 6 (1814); Fl. Ind. i. 168 (1832) = **Salacia**, Linn. (Celastr.).
coromandeliana, Roxb. l. cc. 5; 169 = S. prinoides.
salaciodes, Roxb. l. cc. 5; 168 = S. Roxburghii.
serrata, Buch.-Ham. in Wall. Cat. sub n. 4219 = S. prinoides.

JOHNIA, Wight & Arn. Prod. 449 (1834) = **Glycine**, Linn. (Legumin.).
congesta, Dalz. & Gibs Bomb. Fl. Suppl. 28 = Dolichos bifloras.
Petitiana, A. Rich. Tent. Fl. Abyss. i. 210. t. 40 = G. Petitiana.
Wightii, Wight & Arn. Prod. 449 = G. javanica.
Willdenowia, Hook. f. in Hook. Niger Fl. 305 = G. hedysaroides.

JOHNSONIA, Adans. Fam. ii. 348 (1763) = **Cedrela**, P. Br. (Meliac.).

JOHNSONIA, R. Br. Prod. 287 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 795.
acaulis, *Endl. in Lehm. Pl. Preiss.* ii. 41.—Austral.
hirta, Lindl. Swan River App. 57 = pubescens.
hirta, F. Muell. Fragm. vii. 87, partim = pubescens.
longifolia, Endl. in Lehm. Pl. Preiss. n. 40 = pubescens.
lupulina, *R. Br. Prod.* 287.—Austral.
mucronata, Endl. in Lehm. Pl. Preiss. ii. 40 = pubescens.
pubescens, *Lindl. Swan River App.* 57.—Austral.
Sawalli, F. Muell. Fragm. vii. 85 = Stawellia dimorphantha.
teretifolia, Endl. in Lehm. Pl. Preiss. ii. 40 = lupulina.

JOHNSONIA, T. Dale, ex Mill. Gard. Dict. ed. VI. App. 75 (1752) = **Calliearpa**, Linn. (Verbenac.).
americana, Mill. l. c. ed. VIII. = C. americana.

JOHNSONIA, Neck. Elem. ii. 49 (1790) = **Lycium**, Linn. (Solanac.).

JOHRENIA, DC. Coll. Mém. v. 54 (1829); Prod. iv. 198 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 925.
DICHOROPETALUM, Fenzl, Pugill. Pl. Nov. Syr. 17 (1842).
PHLOIODICARPUS, Turcz. ex Bess. in Flora, xvii. (1834) i. Beibl. 14.
alpina, *Fenzl, in Russegg Reise*, ii. 961. t. 17.—Cilicia.
aurea, *Boiss. & Bal. Diagn.* Ser. II. vi. 81.—Cappadoc.
bertyas, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 1012—As. Min.
Candollei, *Boiss. in Ann. Sc. Nat., Sér.* III. i. (1844) 806.—Persia.
dichotoma, Boiss. l. c. 804 = fungosa.
dichotoma, *DC. Coll. Mém.* v. 54. t. 1.—Syria.
Engleri, *Diagl. in Flora*, lxvi. (1883) 212.—Bithynia.
fungosa, *Boiss. Diagn.* Ser. I. x. 83.—As. Min.
graeca, *Boiss. & Sprun. ex Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 805.—Graecia.
juncea, *Boiss. Diagn.* Ser. I. x. 85.—Syria.
Meyeri, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 807 = Pencedanum paucifolium.
persica, Boiss. l. c. 806 = Caudollel.
platycarpa, *Boiss. l. c.* 808.—Persia.
scoparia, Boiss. Diagn. Ser. I. vi. 61 = Pencedanum scoparium.
selinoides, *Boiss. & Bal. l. c.* Ser. II. v. 99.—Cilicia.
tenuissima, Boiss. l. c. ii. 90 = Pencedanum scoparium.
Tordyliam, Fenzl, Pugill. Pl. Nov. Syr. 18 = Coriandrum tordylioides.

JOINVILLEA, Gaudich Bot. Voy. Bonite, t. 39–40 (1844–52). *FLAGELLAREAE*, Benth. & Hook. f. iii. 861.
ascendens, *Gaudich. l. c.*—Ins. Sandvic.
elegans, *Gaudich. l. c.*—N. Caled.
Gaudichaudiana, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* vii (1861) 269.—Ins. Sandvic.

JOIRA, Steud Nom ed. II. ii. 639, in syn. (1841) = JVIRA, Aubl. = **Sterculia**, Linn.

JOLIFFIA, *Boj. ex Delile, in Mém.* Soc. Hist. Nat. Par. iii. (1827) 314, 318 t. 6 = **Telfairia**, Hook. (Cucurb.).
africana, Delile, l. c. = T. pedata.

JONCQUETIA, Schreb. Gen. 308 (1789) = **Tapiria**, Juss. (Anacard.).
paniculata, Willd. Sp. Pl. ii. 750 = T. guianensis.

JONDRABA, Medic. Pflanzeng. 27 (1792) = **Biscutella**, Linn. (Crucif.).
auriculata, Webb & Berth. Phyt. Canar. i. 94, in obs. = B. auriculata.
cichoriifolia, Webb & Berth. l. c. = B. cichoriifolia.
sulphurea, Medic. Pflanzeng. 29 = B. auriculata.

JONESIA, Roxb. in As. Res. iv. (1795) 355 = **Saraca**, Linn. (Legumin.).
Asoca, Roxb. l. c. = S. indica.
confusa, Hask. Retzia, i. 194.—Java.
declinata, Jack, in Malay. Misc. ii. (1822) vii. 74 = S. declinata.
javanica, Blume, ex Miq. Fl. Ind. Bat. i. 1. 85 = S. declinata.
macroptera, Miq. l. c. 1080 = S. macroptera.
minor, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 80 = S. indica.
monopetala, Hassk. Retzia, i. 199.—Java.
palembanica, Miq. Fl. Ind. Bat. Suppl. 291 = S. palembanica.
punnata, Willd. Sp. Pl. ii. 287 = S. indica.
scandens, Roxb. Hort. Beng. [37]; Fl. Ind. ii. 320 = S. triandra.
triandra, Roxb. Fl. Ind. ii. 320 = S. triandra.

JONGHEA, Lem. Jard. Fleur. ii. sub t. 180, 181 (1852) = **Billbergia**, Thunb. (Bromeliac.).
spendida, Lem. l. c. = B. thyrsoidea.

JONIA, Steud. Nom. ed. II. i. 813 (1840) = **Ionidium**, Vent. (Violar.).

JONIDIOPSIS, Presl, in Abh. Bochm. Ges. V. Folge, iii. (1845) 443 (Bot. Bemerk. 13) = **Noisettia**, H. B. & K. (Violar.).
fruticulosa, Presl, l. cc. = N. longifolia.

JONIRIS, Klatt, in Bot. Zeit. xxx. (1872) 502 = **Iris**, Tourn.
biglumis, *Doniana*, *fragrans*, *longifolia*, *longispatha*, *Pallasii*, *trifora*, Klatt, l. c. = I. ensata.
humilis, Klatt, l. c. = I. humilis.
ruthenica, Klatt, l. c. = I. ruthenica.
sungarica, Klatt, l. c. = I. sungarica.
stylosa, Klatt, l. c. = I. unguicularis.
tenax, Klatt, l. c. = I. tenax.
tenuifolia, Klatt, l. c. = I. tenuifolia.

JONQUILLA, Haw. Monog. Narciss. (1831) 7 = **Narcissus**, Tourn. (Amaryll.).
bicolor, Rafin. Fl. Tellur. iv. 21 = N. Pseudo-narcissus.
incomparabilis, Rafin. l. c. = N. Incomparabilis.
juncifolia, Foarr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 164 = N. juncifolia.
major, Haw. Monog. Narciss. 7 = N. Jonquilla.
media, Haw. l. c. = N. Jonquilla.
minor, Haw. l. c. = N. Jonquilla.
parvicorona, Haw. l. c. = N. Jonquilla ?
Tazetta, Rafin. Fl. Tellur. iv. 21 = N. Tazetta.
triflora, Rafin. l. c. = N. odorus.
viridiflora, Rafin. l. c. = N. viridiflorus.

JONQUILLIA, Endl. Ench. 108 (1841) = prsec.

JONTANEA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 52 = TONTANEA, Aubl. = Cocoocypselum, K. Br. (Rubiac.).

JONTHLASPI, DC. Syst. ii. 326 (1821) = IONTHLASPI, Adans. = **Clypeola**, Linn. (Crucif.).

JOOSIA, Karst. Fl. Columb. i. 9. t. 5 (1858) = **Ladenbergia**, Klotzsch (Rubiac.).
umbellifera, Karst. l. c.—N. Granat.

JORDANIA, Boiss. Diagn. Ser. I. viii. 93 (1849) = **Acanthophyllum**, C. A. Mey. (Caryophyll.).
minuartioides, Boiss. & Heldr. l. c. 94 (= *Gypsophila minuartioides*.—As. Min.
spergulaefolia, Boiss. l. c. (= *Gyps. spergulaefolia*).—As. Min.

JORENA, Adans. Fam. ii. 249 (1763) (Quid ?).

JOSEPHA, Benth. & Hook. f. Gen. iii. 516 (1883). sphalm. = **Josephia**, Wight (Orchid.).

JOSEPHA, Vell. Fl. Flum. 154 (1825); iv. t. 16 (1827) = **Bougainvillaea**, Comm. (Nyctagin.).
augusta, Vell. ll. cc.—B. spectabilis.

JOSEPHIA, Salisb. in Knight, Prot. 110 (1809) = **Dryandra**, R. Br. (Proteac.).
armata, Poir. Encyc. Suppl. iii. 175 = D. armata.
blechnifolia, Poir. l. c. 176 = D. pteridifolia.
cuneata, Poir. l. c. 176 = D. cuneata.
falcata, Poir. l. c. = D. falcata.
floribunda, Poir. l. c. = D. floribunda.
formosa, Poir. l. c. = D. formosa.
longifolia, Poir. l. c. = D. longifolia.
macronulata, Poir. l. c. = D. mucronulata.
nivea, Poir. l. c. = D. nivea.
obtusa, Poir. l. c. = D. obtusa.
plumosa, Poir. l. c. = D. plumosa.
pteridifolia, Poir. l. c. 178 = D. pteridifolia.
rachidifolia, Knight, Prot. 111 = D. nivea.
sessilis, Knight, l. c. 110 = D. floribunda.
tenuifolia, Poir. Encyc. Suppl. iii. 176 = D. tenuifolia.

JOSEPHIA, Stend. Nom. ed. II. i. 814 (1840) = JOSEPHA, Vell. = **Bougainvillaea**, Comm. (Nyctagin.).

JOSEPHIA, Wight, Ic. Pl. v. 19. t. 1742, 1743 (1851). *ORCHIDEAE*, Benth. & Hook. f. iii. 516.
lanceolata, *Wight, l. c.* t. 1742.—Ind. or.
latifolia, *Wight, l. c.* t. 1743.—Ind. or.

JOSEPHINA, Pers. Syn. ii. 174 (1807) = seq.

JOSEPHINIA, Vent. Jard. Malm. 67. t. 67 (1804). *PEDALINEAE*, Benth. & Hook. f. ii. 1060.
africana, *Vathe, in Linnaea*, xliii. (1880–82) 541.—Afr. trop.
celebica, Blume, Bijdr. 779 = imperatricia.
Eugenia, *F. Muell. in Hook. Kew Journ.* ix. (1857) 270. t. 11.—Austral.
grandiflora, *R. Br. Prod.* 520.—Austral.
imperatricia, *Vent. Jard. Malm.* 67. t. 67.—Austral. ; Ins. Celebes.

JOSSINIA, Comm. ex DC. Prod. iii. 287 (err. typ. 337) (1828) = **Eugenia**, Linn. (Myrtac.).
buxifolia, DC. l. c. 288 = E. coriacea.
casuinoides, DC. l. c. = E. casuinoides, *Lam.*
celebica, Blume, Mus. Bot. Lugd. Bat. i. 194.—Ins. Celebes.
cordifolia, Boj. Hort. Maurit. 141 = E. cotinifolia.
cotinifolia, DC. Prod. iii. 288 = E. cotinifolia.
elliptica, DC. l. c. 287 = E. cotinifolia.
ferruginea, Boj. Hort. Maurit. 141 = E. cotinifolia.
glomerata, Blume, Mus. Bot. Lugd. Bat. i. 192.—Ins. Mascar.
hastilis, Blume, l. c.—Ins. Mascar.
indica, Wight, Ic. t. 523; Illustr. ii. 12. t. 122 = E. Jossinia.
Lamarckii, Blume, Mus. Bot. Lugd. Bat. i. 191.—Ins. Maurit.
lanceolata, Boj. Hort. Maurit. 141 = E. cotinifolia.
litoralis, Blume, Mus. Bot. Lugd. Bat. i. 194.—N. Guinea.
lucida, DC. Prod. iii. 287 = E. cotinifolia.
mespiloides, DC. l. c. = E. cotinifolia.
orbiculata, DC. l. c. = E. cotinifolia.
pinnata, Heynh. Nom. i. 481.—Bras.
Reinwardtiana, Blume, Mus. Bot. Lugd. Bat. i. 190 = E. Reinwardtiana.

JOSSINIA :—
revoluta, Boj Hort. Maurit. 140 = Eugenia mespiloides.
Richardii, Blume, Mus. Bot. Lugd. Bat. i. 193.—Madag.
Sieberiana, Blume, l. c. 191.—Ins. Maurit.
tatifolia, Nadeaud, Enum. Pl. Tahiti, 79.—Ins. Tahiti.
terminalis, Boj. Hort. Maurit. 141.—Ins. Mascar.
tinifolia, DC. Prod. iii. 288 = E. cotinifolia.

JOUVEA, Fourn. in Bull. Soc. Bot. Belg. xv. (1876) 475. *GRAMINEAE*, Benth. & Hook. f. iii. 1178
straminea, *Fourn. l. c.*—Mexic.

JOVELLANA, Ruiz & Pav. Fl. Per. i. 12 (1798) = **Calceolaria**, Linn. (Scrophul.).
petiolaris, Cav. Ic. v. 20. t. 445 = C. petiolaris.
punctata, Ruiz & Pav. Fl. Per. i. 13. t. 18. f. a = C. punctata.
scapiflora, Ruiz & Pav. l. c. t. 18. f. b = C. plantaginea.
triandra, Cav. Ic. v. 22. t. 453 = C. triandra.
violacea, G. Don, Gen. Syst. iv. 608 = C. violacea.

JOVIBARBA, Opiz, Seznam, 54 (1852) = **Sempervivum**, Rupp. (Crassul.).
arenaria, Opiz, l. c. 55 = S. arenarium.
hirta, Opiz, l. c. 54 = S. hirtum.
sobolifera, Opiz, l. c. = S. globiferum.

JOXOCARPUS, Pritz. Ic. Ind. i. 586 (1855), sphalm. = **Toxocarpus**, Wight & Arn. (Asclep.).

JOXYLON, Stend. Nom. ed. II. i. 814 (1840), sphalm = TOXYLON, Rafin. = **Maohera**, Nutt. (Urtic.).

JRILLIUM, Rafin. in Journ. Phys. xci. 72 (1820), sphalm. = **Trillium**, Mill. (Liliac.).
sessile, Rafin. l. c. = T. sessile.

JUANIA, Drude, in Goett. Nachr. (1878) 40. *PALMAE*, Benth. & Hook. f. iii. 905.
australis, *Drude, ex Hook. f. Rep. Kew Gard.* (1884) *App.* 57.—Ins. J. Fernand.

JUANULLOA, Ruiz & Pav. Prod. Fl. Per. 27. t. 4 (1794). *SOLANACEAE*, Benth. & Hook. f. ii. 902.
Ectozoma, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iii. (1849) 166; et iv. (1849) 191.
Hydrocalyx, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 48.
Laurenia, Schlecht. in Linnaea, viii. (1833) 513.
Portaea, Tenore, in Atti 7ma Riun. Sc. Ital. 902 (1846).
Sarcophysa, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iii. (1849) 166; et iv. (1849) 190.
Ulloa, Pers. Syn. ii. 216 (1806).
aurantiaca, *Otto & Dietr. Allg. Gartens.* xii. (1844) 286.—Mexic.
elliptica, *Ruiz & Pav. ex Dun. in DC. Prod.* xiii. i. 580.—Mexic.
eximia, Hook. Bot. Mag. t. 5092 = Dyscochroma eximia.
grandiflora, *Dun. l. c.* 581.—Peruv.
Hookeriana, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 188 = aurantiaca.
mexicana, *Miers, l. c.*—Mexic.
salicophylla, *Dun. in DC. Prod.* xiii. i. 581.—Peruv.
panamensis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 190.—Panama.
parasitica, Hook. Bot. Mag. t. 4118 = aurantiaca.
parasitica, *Ruiz & Pav. Fl. Per.* ii. 47. t. 185.—Peruv.
speciosa, *Dun. in DC. Prod.* xiii. i. 582.—N. Granat.
umbellata, Dun. l. c. 580 = Iochroma calycina.

JUBAEA, H. B. & K. Nov. Gen. et Sp. i. 303. t. 96 (1815). *PALMAE*, Benth. & Hook. f. iii. 948.
Michococos, Phil. in Bot. Zeit. xvii. (1859) 205.
Molinara, Bert. in Mercr. Chil. (1829); ex Am. Journ. Sc. xx. (1831) 152.
spectabilis, *H. B. & K. Nov. Gen. et Sp.* i. 303. t. 96.—Chili.
Torallii, *H. Wendl. in Kerch. Palm.* 247.—Ecuador.
Toralyi, Linden, in Illustr. Hortic. xxviii. (1881) 81 = passc.

JUBELINA, A. Juss. in Deless. Ic. Sel. iii. 19. t. 82 (1837). *MALPIGHIACEAE*, Benth. & Hook. f. i. 360.
nicaragnensis, *Griseb. & Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 48.—Nicaragua.
riparia, *A. Juss. in Deless. Ic. Sel.* iii. 19. t. 82.—Guian.

JUBELINIA, Endl. Gen. Suppl. v. 76 (1850) = prec.

JUCHIA, Neck. Elem. i. 132 (1790) = **Lobelia**, Linn. (Campan.).

JUCHIA, M. Roem. Syn. Pepon. 11, 48 (1846) = **Melothria**, Linn. (Cucurbit.).
hastata, M. Roem. l. c. 48 = M. heterophylla.

JUCUNDA, Cham. in Linnaea, ix. (1834) 456 = **Miconia**, Ruiz & Pav. (Melastom.).
Bredemeyriana, Cham. l. c. 458 = M. laevigata.
Lhotskyana, Cham. l. c. 457 = M. stamines.
Martiana, Cham. l. c. 456 = M. stamines.
Olfersiana, Cham. l. c. 457 = M. jucunda.
Rudolphiana, Cham. l. c. 458 = M. laevigata.
Sellowiana, Cham. l. c. 456 = M. jucunda.
tomentosa, Benth. in Hook. Journ. Bot. ii. (1840) 302 = M. tomentosa.

JUDEOLA, Heist. Syst. 9 (1748). (Legum. gen. in-desc.).

JUERGENSENIA, Schlecht. in Bot. Zeit. vi. (1848) 435 = JURGENSENIA, Turcz. = **Befaria**, Mutis (Ericac.).

JUERGENSIA, Spreng. Syst. iii. 50 (1826) = **Alsodeia**, Thou. (Violar.).
angulfera, Spreng. l. c. = A. echinocarpa.

JUGA, Griseb. Fl. Brit. W. Ind. 711 (1864), err. typ. = **Inga**, Soop. (Legumin.).

JUGASTRUM, Miers, in Trans. Linn. Soc. xxx. (1874) 275 = **Leucythis**, Linn. (Myrtac.).
coriaceum, Miers, l. c. = L. coriacea.
depressum, Miers, l. c. 278.—Bras.
obtectum, Miers, l. c. 276.—Bras.
platypterum, Miers, l. c. 277.—Bras.
Poiteaui, Miers, l. c. 279 = L. Poiteaui.
subcinctum, Miers, l. c. 278.—Bras.

JUGLANS, Linn. Gen. ed. I. 291 (1737). *JUGLANDEAE*, Benth. & Hook. f. iii. 398.
Regia, Lond. ex C.DC. in Prod. xvi. ii. 136 (1864).
Wallia, Alef. in Bonplandia, ix. (1861) 285.
ailantifo'ia, Carr. in Rev. Hortic. (1878) 414 = Sieboldiana.
alba, Linn. Sp. Pl. 997 = Carya tomentosa.
alba, Mill. Card. Dict. ed. VIII. n. 4 = Carya alba.
alba-ovata, Marsh. Arbust. 69 = Carya alba.
amara, Michx. Hist. Arb. Am. Sept. i. 177. t. 18 = Carya amara.
amara, Mühl. Cat. 38 = Carya sulcata.
ambigua, Michx. l. c. 208 = Carya sulcata.
angustifolia, Ait. Hort. Kew. iii. 361 = Carya olivaeformis.
aquatica, Michx. Hist. Arb. Am. Sept. i. 182. t. 8 = Carya aquatica.
argeta, Will. Cat. n. 4944; ex C. DC. in DC. Prod. xvi. ii. 138 = regia.
australis, *Griseb. in Goett. Abh.* xxiv. (1879) 97.—Reg. Argent.
baccata, *Linn. Syst. ed.* X. 1272 (Quid ?).—Ind. occ.
californica, *S. Wats. in Proc. Am. Acad.* x. (1875) 349.—Calif.
Canirona, Lour. Fl. Cochinch. 578 = Aleurites triloba.
Cataspa, Lour. l. c. = Terminalia Catappa.
cathartica, Michx. Hist. Arb. Am. Sept. i. 185 = cinerea.
cinerea, *Linn. Syst. ed.* X. 1272.—Am. bor.
compressa, Gaertn. Fruct. ii. 91. t. 89 = Carya alba.
compressa, Mühl. in Gen. Meisurf. Pr. Berl. Neue Schr. iii. (1801) 390; Willd. Sp. Pl. iv. 458 = Carya sulcata.
cordiformis, *Maxim. in Bull. Acad. Pétersb.* xviii. (1872) 62.—Japon.

JUGLANS:—
cordensis, Siebold, in Verh. Batav. Gen. xii. (1830) 27 = Pterocarya rhoifolia.
cylindrica, Lam. Encyc. iv. 505 = Carya olivaeformis.
equatoriensis, Linden, Cat. n. 23 (1869) 26 = granatensis ?
filicifolia, Lodd. Cab. 1836 : ex Loud. Arboret. iii. 1424 = regia.
fraxinifolia, Lam. Encyc. iv. 502 = Pterocarya fraxinifolia.
glabra, J. F. Gmel. Syst. 755 = Carya alba.
glabra, Mill. Gard. Dict. ed. VIII. n. 5 ; Muhl. in Ges. Naturf. Fr. Berl. Neue Schr. iii. (1801) 891 = Carya porcina.
granatensis, *Linden*, Cat. n. 23 (1869) 26.—N. Granat.
heterophylla, Hort. ex Loud. Arboret. iii. 1424 = regia.
illinoënsis, Wangenh. Beitr. Forstw. Nord-Am. Holz. 84. t. 18 = Carya olivaeformis.
insularis, *Griseb. Cat. Pl. Cub.* 68.—Cuba.
intermedia, Carr. in Rev. Hortic. (1870–71) 494 = regia.
jamaicensis, *C. DC. in DC Prod.* xvi. ii. 138.—Jamaic.
japonica, Hort. ex Lavall. Arbor. Segrez. 1 = Sieboldiana.
japonica, *Siebold, ex Mig. Ann. Mus. Bot. Lugd. Bat.* iii. 103.—Japon.
Kurumi, Siebold, in Verh. Batav. Gen. xii. (1830) 27 = Platycarya strobilacea.
laciniosa, Michx. Hist. Arbr. Am. Sept. i. 199 = Carya sulcata.
longirostris, Carr. in Rev. Hortic. (1876) 53 = regia.
macrophylla, Carr. l. c. 415 = Sieboldiana.
mandshurica, *Maxim. in Bull. Phys.-Math. Acad. Petersb.* xv. (1856) 127.—Manchuria.
mandshurica, Miq. in Anu. Mus. Bot. Lugd. Bat. iii. 104 = Sieboldiana.
mollis, *Engelm. in Hemsl. Diagn. Pl. Nov.* 54.—Mexic.
mucronata, Michx. Fl. Bor. Am. ii. 192 = Carya sulcata.
myristicaeformis, Michx. Hist. Arb. Am. Sept. i. 211 = Carya myristicaeformis.
nigra, *Linn. Sp. Pl.* 997.—Am. bor.
nigra, Thunb. Fl. Jap. 195 = Sieboldiana.
obcordata, Lam. Encyc. iv. 504 = Carya porcina.
oblonga, Mill. Gard. Dict. ed. VIII. n. 8 = cinerea.
olivaeformis, Michx. Hist. Arb. Am. Sept. i. 173 = Carya olivaeformis.
ovata, Mill. Gard. Dict. ed. VIII. n. 6 = Carya alba.
Pecan, Walt. Fl. Carol. 236 = Carya olivaeformis.
Pitturesii, Morr. in Ann. Soc. Bot. Gand, iv. (1848) 179. t. 197 = nigra.
porcina, Michx. Hist. Arbr. Am. Sept. i. 206 = Carya porcina.
procera, Salisb. Prod. 392 = Carya alba.
pterocarpa, Buch.-Ham. ex Wall. Cat. sub n. 4940 = Engelhardtia spicata.
pterocarpa, Michx. Fl. Bor. Am. ii. 192 ; Willd. Sp. Pl. iv. 455 = Pterocarya fraxinifolia.
pterocera, Roxb. Hort. Beng. 68 ; Fl. Ind. iii. 681 = Engelhardtia spicata.
pubescens, Willd. Enum. Hort. Berol. Suppl. 64 = Carya sulcata.
pyriformis, *Liebm. in Kjoeb. Vidensk. Meddel.* (1850) 79.—Mexic.
pyriformis, Mnhl. Cat. 93 = Carya porcina.
regia, *Linn. Sp. Pl.* 997.—As. occ. ; Reg. Himal.
rubra, Gaertn. Fruct. ii. 51. t. 89 = Carya olivaeformis.
rupestris, *Engelm. in Torr. Bot. Stkgr. Rep.* 171. t. 15. —Am. bor. occ.
Sieboldiana, *Maxim. in Bull. Acad. Petersb.* xviii. (1873) 66.—Japon.
squamosa, Poir. in Lam. Encyc. iv. 504 ; Michx. Hist. Arbr. Am. Sept. i. 190 = Carya alba.
stenocarpa, *Maxim. Prim. Fl. Amur.* 78.—Reg. Amur.
sulcata, Willd. Berl. Baumz. 154 = Carya sulcata.
tomentosa, Lam. Encyc. iv. 504 = Carya alba.
tomentosa, Michx. Hist. Arbr. Am. Sept. i. 184 = Carya tomentosa.
villosa, Wall. Cat. n. 4945 = Engelhardtia Colebrookiana.

JUJUBA, Burm. Thes. Zeyl. 181. t. 61 (1737) = Ziziphus, Tourn. (Rhamn.).

JULIA, Steud. Nom. ed. II. i. 383, in syn. (1840) = JUNIA, Adans. = Clethra, Linn. (Vaccin.).

JULIANA, Reichb. Nom. 198 (1841) = seq.

JULIANIA, La Llave, in La Llave & Lex. Veg. Descr. fasc. ii. 4 (1825) = Choteya, Kunth (Rutac.).
caryophillata, La Llave, l. c. = C. tomata.

JULIANIA, Schlecht. in Linnaea, xvii. (1843) 746. *ANACARDIACEAE*, Benth. & Hook. f. i. 428.
HYPOTERVGIUM, Schlecht. l. c. (1843) 685.
adstringens, *Schlecht. l. c.* 686.—Mexic.
Jhuanii, *A. Gray, Bot. U. St. Expl. Exped.* i. 371.—Peruv.

JULIETA, Leschen. ex DC. Prod. vii. 764 (1839) = Lystnema, R. Br (Epacrid.).
unguiculata, Leschen. ex DC. l. c. 765 = L. ciliatum.

JULOCROTON, Mart. in Flora, xx. (1837) ii. Beibl. 119. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 296.
CENTRANDRA, Kunt. in Linnaea, xxviii. (1856) 440 ; et Fl. Columb. i. 117. t. 88 (1859).
CIECA, Adans. Fam. ii. 355 (1763).
HETEROCHLAMYS, Turcz. in Bull. Soc. Nat. Mosc. (1. (1848) 61.
Ackermannianus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 283.—Bras.
argenteus, *Didr. in Kjoeb. Vidensk. Meddel.* (1857) 184.—Am. centr.
conspurcatus, *Klotzsch, in Wiegm. Archiv.*, vii. (1841) 193.—Mexic.
decalobus, *Benth. & Hook. f. Gen.* iii. 296.—Guatemala.
dorstophyllus, *Muell. Arg. in DC. Prod.* xv. ii. 702.—Reg. Argent.
fuscescens, *Baill. Adansonia*, iv. (1863–64) 367.—Bras.
Gardneri, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 276. —Reg. Argent.
geraensis, *Baill. Adansonia*, iv. (1863–64) 370.—Bras.
boodensis, *Muell. Arg. in DC. Prod.* xv. ii. 704.—Am. trop.
humilis, *Didr. in Kjoeb. Vidensk. Meddel.* (1857) 183. —Bras.
lanatus, Klotzsch, ex Baill. Etud. Gén. Euph. 376 = fuscescens.
lanceolatus, *Klotzsch, in Wiegm. Archiv*, vii. (1841) 193.—Bras.
microcalyx, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 142.—Bras.
montevidensis, *Klotzsch, ex Baill. Etud. Gén. Euph.* 376.—Reg. Argent.
nervosus, *Baill. Adansonia*, iv. (1863–64) 369.—Bras.
nigricans, *Mart. ex Schlecht. in Linnaea*, xix. (1847) 345.—Bras.
peruvianus, *Muell. Arg. l. c.* xxxiv. (1865–66) 142.—Am. trop.
phagedaenicus, Schlecht. l. c. xix. (1847) 345 = seq.
phagedaenicus, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 119.—Bras.
pycnophyllus, *Schlecht. in Linnaea*, xix. (1847) 345. —Bras.
quinquenervius, Baill. Etud. Gén. Euph. 376 = argenteus.
Riedelianus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 278. —Bras.
rufescens, *Klotzsch, ex Baill. Etud. Gén. Euph.* 376, nomen.—Bras.
Salzmanni, *Baill. Adansonia*, iv. (1863–64) 369.—Bras.
serratus, *Muell. Arg. in Journ. Bot.* xii. (1874) 237.—Reg. Argent.
solanaceus, *Klotzsch, ex Baill. Etud. Gén. Euph.* 376 ; *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 279.—Bras.
stipularis, *Muell. Arg. l. c.* 277.—Bras.
subpannosus, *Muell. Arg. in Journ. Bot.* xii. (1874) 303.—Reg. Argent.
triqueter, *Baill. Adansonia*, iv. (1863–64) 368.—Bras.
triqueter, Didr. in Kjoeb. Vidensk. Meddel. (1857) 184 = phagedaenicus.
verbascifolius, *Klotzsch, ex Baill. Etud. Gén. Euph.* 376.—Bras.

JULOSTYLES, Benth. & Hook. f. Gen. i. 907 (1862) = seq.

JULOSTYLIS, Thw. Enum. Pl. Zeyl. 80 (1858). *MALVACEAE*, Benth. & Hook. f. i. 207.
angustifolia, *Thw. l. c.* 80.—Zeylan.

JUNAGO, Tourn. ex Moench, Meth. 644 (1794) = Triglochin, Linn. (Naiad.).
palustris, Moench, l. c. = T. palustre.

JUNCASTRUM, Heist. Syst. 12 (1748) = Juncus, Tourn.
acutum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 171 = J. acutus.
maritimum, Fourr. l. c. = J. maritimus.

JUNCELLA, F. Muell. 2nd Gen. Rep. 16 (1854) = Trithuria, Hook. f. (Centrolepid.).
tasmanica, F. Muell. l. c. = T. submersa.

JUNCINELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 171 = Juncus, Tourn.
capitata, Fourr. l. c. = J. capitatus.
pygmaea, Fourr. l. c. = J. pygmaeus.

JUNCOIDES, [Dill.] Adans. Fam. ii. 47 (1763) = Luzula, DC. (Juncac.).

JUNCUS, [Tourn.] Linn. Syst. ed. I (1735). *JUNCACEAE*, Benth. & Hook. f. iii. 67.
CEPHALOXYS, Desv. Journ. Bot. i. (1808) 321. t. 11.
JUNCASTRUM, Heist. Syst. 12 (1748).
JUNCINELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 171.
MARSIPPOSPERMUM, Desv. Journ. Bot. i. (1808) 328. t. 12. f. 1.
PHYLLOSCHOENUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 171.
TENAGRIA, Ehrh. Beitr. iv. 148 (1789).
aboriivus, Chapm. Fl. S. U. St. 495 = pelocarpus.
acuminatus, Balb. Elench. 87 = arcticus.
acuminatus, *Michx. Fl. Bor. Am.* i. 192.—Mexic.
acuminatus, Salzm. ex Steud. Syn. Pl. Cyp. 209 = Elliottii.
acutanguius, *Buchen. in Bremen, Abh.* iv. (1875) 480. —Afr. austr.
acutiflorus, Benth. Pl. Hartw. 341 = oxymeris.
acutiflorus, *Ehrh. Beitr.* vi. 92.—Europ. ; As. bor.
acutiflorus, Griseb. Spicil. Fl. Rumel. ii. 407 = stratus.
acutiflorus, Guss. Fl. Sic. Prod. i. 434 = Gussonii.
acutiflorus, Spreng. Neue Entdeck. ii. 107 = punctorius.
acutiflorus, Tenore, Fl. Nap. iii. 381 = lampocarpus.
acutus, *Linn. Sp. Pl.* 326.—Reg. temp.
acutus, Retz. Prod. Fl. Scand. ed. II. 79 (auctus) = maritimus.
acutus, Thuill. Fl. Par. 175 = glaucus.
acutus, Tsi in Comp. Fl. N. Midd. Stat. (1826) 156 = Roemerianus.
adscendens, Host, Gram. Austr. iii. 58. t. 87 = lampocarpus.
aemulans, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 88 = effusus.
affinis, R. Br. in Richards. App. Frankl. Journ. 11 = alpinus.
affinis, Gand. Agrost. Helvet. ii. 224 = supinus.
agrostophyllus, F. Muell. in Linnaea, xxvi. (1853–55) 345 = falcatus.
alatus, *Franch. & Sav. Enum. Pl. Jap.* ii. 584.—Japon.
albidus, Hoffm. Deutschl. Fl. ed. I. 126. t. 6 = Luzula angustifolia.
albidus, Willd. Sp. Pl. ii. 219 = Luzula nemorosa.
albus, *Ruffin. Fl. Ludov.* 17.—Am. bor.
alpestris, Gilem. ex Bab. in Journ. Linn. Soc. xi. (1871) 322 = lampocarpus.
alpestris, Hartm. Handb. Scand. Fl. ed. I. 141 = alpinus.
alpigenus, *C. Koch, in Linnaea*, xxi. (1848) 397.—Reg. Caucas.
alpinoplanus, Vill. Hist. Pl. Dauph. i. 318 = Luzula alpinoplanus.
alpinus, Pers. Syn. i. 384 = supinus.

JUNCUS :—
alpinus, *Vill. Hist. Pl. Dauph.* ii. 233.—Europ. ; Am. bor.
altus, *Buchen. in Bremen, Abh.* iv. (1875) 457.—Afr. austr.
ambiguus, Guss. Fl. Sic. Prod. i. 435 = J. Tenageia.
anceps, *Laharpe, in Mém. Soc. Hist. Nat. Par.* iii. (1827) 136.—Europ. ; Afr. bor.
andicolus, *Hook. Ic. Pl.* t. 714.—Reg. Andin.
Angelsii, Tenore, Viag. Abruz. 61 ; Syll. 178 = glaucus.
angustifolius, Wulf. in Jacq. Coll. iii. 56 = Luzula albida.
annuus, Krock. Fl. Siles. i. 566 = supinus
anonymus, *Steud. Syn. Pl. Cyp.* 304.—Afr. austr.
antarcticus, *Hook. Fl. Antarct.* 79. t. 46.– N. Zel.
Antonianus, Steud. in Lechl. Berb. Am. Austr. 56 = andicolus.
aquaticus, All. Fl. Pedem. ii. 217 = lampocarpus.
aquaticus, Brot. Fl. Lusit. i. 517 = acutiflorus ?
aquaticus, Sebast. & Mauri, Prod. Fl. Rom. (1818) 182 = obtusiflorus.
arcticus, Hook. Fl. Scot. 104 = baltícus.
arcticus, Lapeyr. Hist. Abr. Pl. Pyr. 103 = filiformis.
arcticus, *Willd. Sp. Pl.* ii. 206.—Reg. bor. et arct
arcuatus, Hook. in Scoresb. Journ. Greenl. 410 = Luzula hyperborea.
arcuatus, Wahlenb. Fl. Lapp. 67 = Luzula arctica.
aristatus, Link, Enum. Hort. Berol. i. 306 = tenuis.
aristatus, Pers. Syn. i. 385 = marginatus.
aristiflorus, [Clairv.] Man. Herb. 104 = lampocarpus.
aristulatus, Michx. Fl. Bor. Am. i. 192 = marginatus.
articulatus, DC. Fl. Fr. iii. 169 = obtusiflorus.
articulatus, Desf. Fl. Atlant. i. 318 = Fontanesii.
articulatus, Linn. Sp. Pl. 327 = lampocarpus.
asper, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 476.—Am. bor.
asper, Sauzé-Maill. Cat. Pl. Deux-Sèvres, 52 = striatus.
atlanticus, Labarpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 128 = heterophyllus.
atratus, Bess. Prim. Fl. Galic. i. 241 = acutiflorus.
atratus, Fries, Nov. Fl. Suec. Mant. iii. 23 = anceps.
atratus, *Krock. Fl. Siles.* i. 562.—Europ. ; Reg. Caucas.
atratus, Lam. Encyc. iii. 271 = biglumis.
atricephalus, Drejer, in Kröyers, Tidsskrift, ii. (1838) 182 = anceps.
atrofuscus, Ruprr. Betr. Pfl Russl. ii. 59 = Gerardi.
attenuatus, Vir. Fl. Cors. Sp. Nov. 5 = Gerardi.
auctus, Retz. Prod. Fl. Scand. ed. II. 79, sphalm. = maritimus.
aureus, Pourr. in Mém. Acad. Toul. iii. (1788) 321 = Luzula lutea.
austerus, *Buchen. in Bremen, Abh.* vi. (1879) 389. t. 8.—Chili.
australis, Desf. ex Steud. Nom. ed. I. 448 (Quid ?).—Hab. ?
australis, Hook. f. Fl. Tasm. ii. 66. t. 134 = paniciflorus.
Bachiti, *Hochst. ex Steud. Syn. Pl. Cyp.* 305.—Abyssin.
balticus, *Willd. in Ges. Naturf. Fr. Berl. Mag.* iii. (1809) 298.—Amphig. bor.
benghalensis, Kunth, Enum. Pl. iii. 360 = concinnus.
bicephalus, *Barcelo, Fl. Ins. Balear.* 472.—Ins. Majorc.
bicephalus, Viv. Fl. Cors. Sp. Nov. 5 = pygmaeus.
bicornis, Michx. Fl. Bor. Am. i. 191 = tenuis.
bifarus, Ell. Sketch, i. 407 = marginatus.
bifarus, Phil. in Anal. Univ. Chil. (1873) 552 = stipulatus.
bifolius, Hoppe, Taschenb. (1810) 163 = obtusiflorus.
biglumis, Jacq. Enum. Vind. 61, 237 = Jacquinii.
biglumis, Krock. Fl. Siles. i. 567 = triglumis.
biglumis, *Linn. Sp. Pl.* 325.—Reg. bor. et arct.
bogotensis, H. B. & K. Nov. Gen. et Sp. i. 235 = effusus.
Bolanderi, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 470.—Calif.
bottnicus, Wahlenb. Fl. Lapp. 32. t. 5 = Gerardi.
brachycarpus, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 467.—Am. bor.
brachyspathus, Maxim. Prim. Fl. Amur. 228 = filiformis.
bracteatus, *Buchen. in Engl. Bot. Jahrb.* vi. (1885) 230.—Reg. Himal.
brachestus, Stokes, Bot. Mat. Med. ii. 289 = Scirpus capsularis.
brachyocras, Kit. in Linnaea, xxxii. (1863) 323 = Luzula parviflora.

JUNCUS :—
brevifolius, T. Kirk, in Journ. Linn. Soc. xix. (1882) 289 = striatulus.
brevifolius, *Lieblm. in Kjoeb. Videnskab. Meddel.* (1850) 40.—Mexic.
brevifolius, Pers. Syn. i. 386 = Luzula nivea.
brevifolius, Roth. Dis. Junc. (1801) 35 = Luzula lactea.
brevirostris, Nees, in Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 482 = acutiflorus.
brevistylus, *Buchen. in Bremen, Abh.* iv. (1875) 433.—Afr. austr.
Breweri, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 475.—Calif.
Broteri, Steud. Syn. Pl. Cyp. 297 = maritimus.
Brownii, F. Muell. in Linnaea, xxvi. (1853–55) 245 ; Fine Gen. Report, 19 = revolutus.
brunneus, *Buchen. in Bremen, Abh.* vi. (1879) 403.—Reg. Andin.
Buchleyi, Engelm. in Trans. Acad. Sc. St. Louis, ii. (1868) 459 = leptocaulis.
bufonius, *Linn. Sp. Pl.* 328.—Amphig.
bulbosus, Ehrh. Beitr. vi. 81 = Gerardi.
bulbosus, Linn. Sp. Pl. ed. I. 327 = supinus.
bulbosus, Linn. f. c. ed. II. 466 = compressus.
bulbosus, Lour. Fl. Cochinch. 213 = Loureiroanus.
bupleuroides, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 135 = pygmaeus.
cacspiticus, *E. Mey. in Lehm. Pl. Preiss.* ii. 47.—Austral.
cæfer, Bertol. f. in Mem. Acc. Sc. Bolog. iii. (1851) 253 = maritimus.
campestris, Linn. Sp. Pl. 329 = Luzula campestris
canadensis, *Y. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par.* iii. (1827) 144.—Am. bor.
canaliculatus, *Engelm. in Coult. Bot. Gaz.* vii. (1882) 6.—Calif.
canalinervius, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 43 = marginatus.
canariensis, Willd. ex E. Mey. Syn. Luz. 29 = effusus.
candidus, Turcz. ex Ledeb. Fl. Koss. iv. 253 = trigluma.
capensis, *Thunb. Prod. Pl. Cap.* 66.—Afr. austr.
capillaceus, Hook. f. Fl. N. Zel. i. 264 = pusillus.
capillaceus, *Lam.* Encyc. iii. 266.—Am. austr.
capitatus, Schk. Handb. i. 302 = supinus.
capitatus, *Weigel, Obs. Bot.* 28 (1772).—Europ. ; Afr. bor.
capsularis, *Steud. Syn. Pl. Cyp.* 309.—Cochinch.
caricinus, Dur. Expl. Sc. Alger. t. 43. f. 1 = valvatus.
castaneus, [Clairv.] Man. Herb. 104 = Jacquinii.
castaneus, *Sm. Fl. Brit.* i. 358.—Reg. bor. et arct.
Castelli, Tineo, ex Guss. Fl. Sic. Syn. ii. 816 = Gussonli.
caudatus, Chapm. Fl. S. U. St. 495 = trigonocarpus.
Cephalotes, Hochst. ex Krauss, in Flora, xxviii. (1845) 349 = capensis.
Cephalotus, Thunb. Prod. Pl. Cap. 66 = Ionostaphyllus.
Chamaeschoenus, Wolff, ex Schult. f. Syst. vii. 210 = supinus.
Chamissonis, *Kunth, Enum. Pl.* iii. 343.—Reg. Andin.
chilensis, *C. Gay, Fl. Chil.* vi. 146.—Chili.
chinensis, Hort. ex Buchen. Kritisch. Verz. Junc. 18 = prismatocarpus.
chlorocephalus, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 485.—Calif.
chlorolicus, Schult. f. Syst. vii. 240 = tenuis.
chrysocoma, *Buchen. in Engl. Bot. Jahrb.* vi. (1885) 201.—Reg. Himal.
Clarkei, *Buchen. l. c.* 210.—Reg. Himal.
coarctatus, Hoppe, ex Schult. f. Syst. vii. 182 = glaucus.
coarctatus, Willd. ex Spreng. Syst. ii. 109 = tenuis.
coronus, Bieberst, in Trans. Linn. Soc. xii. (1819) 309 = Gerardi.
cognatus, Kunth, Enum. Pl. iii. 349 = tenuis ?
collinus, Steud. Syn. Pl. Cyp. 306 = Chamissonis.
communis, Steud. l. c. 300 = Domboyanus.
communis, E. Mey. Junc. 12 = *conglomeratus*, effusus.
communatus, Steud. Syn. Pl. Cyp. 301 = prismatocarpus.
compactus, Nees, ex Buchen. Kritisch. Verz. Junc. 14 = obtusiflorus.
complanatus, Schult. f. Syst. vii. 195 = mexicanus.
compressus, H. B. & K. Nov. Gen. et Sp. i. 235 = mexicanus.

JUNCUS :—
compressus, *Jacq. Enum. Stirp. Vind.* 60 ; 235.—Reg. temp.
compressus, Lindsay, ex Bab. in Journ. Linn. Soc. xi. (1870) 332 = Gerardi.
compressus, E. Mey. Junc. 16 ; et in Linnaea, iii. (1828) 393 = Lescurii.
compressus, Relh. Fl. Cantab. (1785) 142 = acutiflorus.
Conceptionis, Steud. Syn. Pl. Cyp. 295 = Lesueurii.
concinnus, *D. Don, Prod. Fl. Nep.* 44.—Reg. Himal.
confervaceus, St. Lag. in Ann. Soc. Bot. Lyon (1882) 749 = supinus.
congestus, Nees, in Heller, Suppl. Fl. Wirceb. (1815) 29 = obtusiflorus.
congestus, Schousb. ex Buchen. Kritisch. Verz. Junc. 15 = bufonius.
congestus, Thuill. Fl. Par. ed. II. (1799) 179 = Luzula campestris.
conglomeratus, Linn. Sp. Pl. 326 = effusus.
Conradi, Tuckerm. ex Torr. Fl. N. York, ii. 228 = pilocarpus.
consanguineus, Ziz, in Koch & Ziz, Cat. 8, 19 = Gerardi.
contractus, Hoppe, ex Schult. f. Syst. vii. 182 = glaucus.
Cooperi, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 486.—Calif.
corralensis, Phil. in Linnaea, xxxiii. (1864–65) 289 = stipulatus.
corrector, Steud. Syn. Pl. Cyp. 296 = pallidus.
crassifolius, Rosc, ex Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 140 = scirpoides.
cryptocarpus, Bebb, ex Engelm. in Trans. Acad. Sc. St. Louis, ii. (1868) 467 = brachycarpus.
cylindricus, M. A. Curt. in Am. Journ. Sc. xliv. (1843) 83 = marginatus.
cymosus, Lam. Encyc. iii. 267 = Ionostaphyllus.
cymosus, Schrank, Baier. Fl. i. 622 = Luzula vernalis.
cymosus, Spreng. Neue Entdeck. ii. 105 = capensis.
cyperinus, Willd. ex Spreng. Syst. ii. 106 = densiflorus.
cyperoides, *Laharpe, in Mém. Soc. Hist. Nat. Par.* iii (1827) 145.—Am. austr.
Czetzii, Schur, in Oestr. Bot. Zeitschr. xiii. (1863) 111 = castaneus.
Deangelisii, Tenore, Fl. Nap. iii. 330 = glaucus.
debilis, A. Gray, Man. Bot. U. St. ed. II. 480 = scirpinatus.
delicatulus, Steud. Cyp. Pl. Cyp. 304 = capensis.
densiflorus, Steud. l. c. 308 = planifolius.
densiflorus, *H. B. & K. Nov. Gen. et Sp.* i. 238.—Am. austr.
depauperatus, Tenore, Viag. Abruz. 62 ; Phil. Fl. Atac. 53 = glaucus.
deserticola, *Phil. l. c.* 52.—Chili.
diaphanus, *Buchen. in Bremen, Abh.* iv. (1875) 442.—Afr. austr.
diaphragmatus, Hochst. ex Nym. Consp. 747 = acutiflorus.
diaphragmaticus, Steud. Nom. ed. II. i. 831, sphalm. = *diaphramarius*.
diaphramarius, Brot. Fl. Lusit. i. 511 = glaucus.
dichotomus, *Ell. Sketch*, i. 406.—Am. bor. et austr.
dichotomus, Willd. ex Spreng. Syst. ii. 108 = microcephalus.
diffusissimus, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 9 = scuminatus.
diffusus, Hoppe, in Flora, i. (1819) 166 = effusus.
digynus ♀, Berb. in Magyar Növenyt. Lapok, ix. (1885) 59.—Europ.
dioicus, Steud. Syn. Pl. Cyp. 309 (an Junci sp. ?).—Austral.
diphyllus, Hoppe, Bot. Taschenb. (1810) 170, 169 = obtusiflorus.
discoriatus, Glith. Exercit. ii. 606 = bufonius.
disvaricatus, Wolff, Fl. Wirceb. 38 = obtusiflorus.
divergens, Koch, in Koch & Ziz, Cat. 8, 19 = obtusiflorus.
Domboyanus, *Y. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par.* iii. (1827) 133.—Am. austr.
Drogonus, *Kunth, Enum. Pl.* iii. 344.—Afr. austr.
Dregeanus, Presl, Bot. Bemerk. 117 = bufonius.
Drummondi, *E. Mey. in Lehm. Pl. Pruss.* iv. 326.—Ins. Koraginsk.
Drummondii, Steud. Syn. Pl. Cyp. 294 = pallidus.
dubius, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 480.—Calif.

JUNCUS:—
dubius, Wolff, ex Schult. f. Syst. vii. 197 = alpinus.
Duvalii, Loret, in Loret & Barr. Fl. Montp. 685, 848 = Tenoreanus.
ebracteatus, E. Mey. Syn. Junc. 28 = trinervis.
echinatus, Ell. Sketch, i. 410 = scirpoides.
echinoides, Brot. Fl. Lusit. i. 518 = valvatus.
echinuloides, Webb, It. Hisp. 6 = Fontanesii.
effusus, Auct. Am. ex Schult. f. Syst. vii. 247 = maritimus.
effusus, *Linn. Sp. Pl.* 326.—Amphig.
effusus, Pollich, Hist. Pl. Palat. i. 345 = glaucus.
effusus, Schum. ex Schult. f. Syst. vii. 185 = balticus.
elatior, Lange, in Kjoeb. Vidensk. Meddel. (1860) 67 = Gerardi.
elatus, Steud. Syn. Pl. Cyp. 295 = glaucus.
elegans, Ehrenb. ex Boiss. Fl. Orient. v. 859 = pyramidatus.
elegans, Royle, ex D. Don, in Trans. Linn. Soc. xviii (1840) 221 = concinnus.
Elliottii, *Chapm. Fl. S. U. St.* 494.—Am. bor.
elongatus, Vasey, ex Engelm. in Trans. Acad. Sc. St. Louis, ii. (1863) 459 = alpinus.
Engelmanni, *Buchen. Kritisch. Vers. Junc.* 18.—Am. bor.
enalifolius, *Wihstr. in Vet. Akad. Handl. Stockh.* (1823) *11.* 274.—Am. bor. occ.
equisetosus, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 365 = glaucus.
erectus, Bess. Prim. Fl. Galic. i. 241 = alpinus.
erectus, Pers. Syn. i. 386 = Lusula campestris.
ericetorum, Krock. Fl. Siles. i. 565. t. 50 = supinus.
ericetorum, Pollich, Hist. Pl. Palat. i. 351 = capitatus.
erythrocarpus, Chapm. ex Engelm. in Trans. Acad. Sc. St. Louis, ii. (1868) 478 = trigonocarpus.
exaltatus, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 16 = punctorius.
expansus, Jan, Catal. ex Schult. f. Syst. vii. 179 = effusus.
exsertus, *Buchen. in Bremen, Abh.* iv. (1875) 435.—Afr. austr.
falcatus, *E. Mey. Syn. Lusul.* 34.—Calif.
fasciculatis, Schrank, ex Steud. Nom. ed. I. 444, 445 = Luzula pilosa.
fasciculatus, Bertol. Fl. Ital. iv. 190 = bufonius.
fasciculatus, *Schousb. in E. Mey. Syn. Junc.* 28.—Marocc.
fasciculatus, Schrank, Baier. Fl. i. 616 = supinus.
fernandezianus, *Steud. Syn. Pl. Cyp.* 302.—Ins. J. Fernand.
filiformis, *Linn. Sp. Pl.* 326.—Reg. bor. temp.
filiformis, Walt. Fl. Carol. 124 = setaceus.
filipendulus, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 8 = leptocaulis.
Fischeri, Turcz. in Bull. Soc. Nat. Mosc. xxviii. (1855) t. 308 = alpinus.
flatulosus, Guss. Fl. Sic. Prod. 481 = effusus.
flaccidus, Steud. Syn. Pl. Cyp. 308 = capensis.
flaweczens, Host, Gram. Austr. iii. 62 = Luzula Hostii.
flexilis, *Turcz. ex Bess. in Flora*, xvii. (1834) t. *Beibl.* 92.—Sibir. boical.
floribundus, H. B. & K. Nov. Gen. et Sp. i. 237 = microcephalus.
floridanus, Raf. ex Engelm. in Trans. Acad. Sc. St. Louis, ii. (1866) 451 = Gerardi.
fluitans, Lam. Encyc. iii. 271 = supinus.
fluitans, Michx. Fl. Bor. Am. i. 191 = pelocarpos.
fluitans, Vest, ex Schult. f. Syst. vii. 201 = lampocarpus.
foliosus, Desf. Fl. Atlant. i. 315. t. 92 = bufonius.
foliosus, Hoppe, Bot. Taschenb. (1810) 159 = lampocarpus.
Fontanesii, Bory & Chaub. Nouv. Fl. Pélop. 25 = striatus.
Fontanesii, *Y. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par.* iii. (1827) 130.—Europ. austr.; Orient.
Forsteri, Sm. Fl. Brit. iii. 1305 = Luzula Forsteri.
fraternus, Kunth, Enum. Pl. iii. 340 = acuminatus.
fuscatus, Turcz. ex Ledeb. Fl. Ross. iv. 238 = triglumis
fuscoater, Schreb. in Schweigg. & Koerte, Fl. Erlang. 149 = alpinus.
Gayanus, Remy, Syn. Pl. Cyp. 300 = chilensis.
gemiculatus, Schrank, Baier. Fl. i. 618 = alpinus.
Gerardi, *Loisel. in Desv. Journ. Bot.* ii. (1808) 324.—Reg. bor. temp.
germanorum, Steud. Syn. Pl. Cyp. 305 = tenuis.
Gesneri, Sm. Eng. Fl. ii. 167 = tenuis.

JUNCUS:—
gibraltaricus, Salem. ex Buchen. Krit. Vers. Junc. 21 = striatus.
glabratus, Hoppe, ex Rostk. Junc. (1801) 27 = Luzula Desvauxii.
glabrescens, Schrank, Baier. Fl. i. 622 = Luzula vernalis.
glaucocarpus, Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 115 = glaucus.
glaucus, *Sibth. Fl. Oxon.* 113.—Amphig.
glaucus, Wahlenb. Fl. Lapp. 79 = balticus.
glomeratus, Thunb. Fl. Jap. 146 = effusus.
gracilis, *R. Br. Prod.* 259.—Austral.
gracilis, Lej. Fl. Spa, 166 = compressus
gracilis, Roth, Tent. Fl. Germ. i. 125 = capitatus.
gracilis, Sm. Engl. Bot. t. 2174 = tenuis.
gracilis, Walp. Ann. i. 878 = Kostkovia gracilis, *Hook. f.*
gracilus, Bory & Chaub. Nouv. Fl. Pélop. 23. t. 12. f. 1 = Luzula graeca.
graminifolius, Dum. Fl. Belg. 142 = supinus.
graminifolius, E. Mey. in Presl, Rel. Haenk. i. 144 = cyperoides.
grandiflorus, Linn. f. Suppl. 209 = Rostkovia grandiflora, *Hook. f.*
Greenei, *Buchen. & Tuckerm. in Am. Journ. Sc.* Ser. I. xlv. (1843) 87.—Am. bor.
Griesebachii, *Buchen. in Bremen, Abh.* iii. (1873) 295. —Reg. Himal.
Gunnii, *Hook. f.* Fl. Tasm. ii. 67 = pauciflorus.
Gussonianus, Tod. ex Parl. Fl. Ital. ii. 342 = seq
Gussonii, *Parl. Fl. Ital.* ii. 341.—Sicil.
Hallii, Mey. Syn. Junc. 10 = compressus.
Hallii, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 446.—Am. bor. occ.
Hancockii, *Hance, in Journ. Bot.* xvi. (1878) 111.—China.
Haussknechtii x, Ruhm. in Eichl. Jahrb. Berl. i. (1881) 257.—Europ.
Helderianus, Maras. ex Parl. Fl. Ital. ii. 315 = acutus.
helodes, Link, Enum. Hort. Berol. i. 305 = balticus.
heptogynius, Peterm. ex Koch, Syn. Fl. Germ. ed. II. 843 = tenuis.
heteranthus, Nutt. in Trans. Am. Phil. Soc. v. (1837) 158 = marginatus.
heterophyllus, *Duf. in Ann. Sc. Nat.* Sér. I. v. (1825) 38.—Europ. austr.
himalensis, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 60.—Reg. Himal.
Hochstetter, *Steud. Syn. Pl. Cyp.* 299 = striatus.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 60 = membranaceus.
Holoschoenus, *F. Br. Prod.* 259.—Austral. ; N. Zel.
homalocaulis, *F. Muell. First Gen. Report*, 19 ; *et in Benth. Fl. Australiæ* vii. 198.—Austral.
homalophyllus, Steud. Syn. Pl. Cyp. 308 = planifolius.
Hoockeriis, Steud. l. c. 307 = Rostkovia gracilis, *Hook. f. Hoppii*, Noé, ex Nym. Consp. 746 = acutus.
Hostii, Tausch, in Flora, xvii. (1834) *it.* 518 = trifidus.
hybridus, Brot. Fl. Lusit. i. 518 = pygmæus.
imbricatus, Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 149 = Chamissonis.
Inaequalis, *Buchen. in Bremen, Abh.* iv. (1875) 455.—Afr. austr.
inaequalis, Willd. ex Kunth, Enum. Pl. iii. 354 = bufonius.
inconspicuus, Urv. in Mém. Soc. Linn. Par. iv. (1826) 608 = scheuchzerioides.
indescriptus, *Steud. Syn. Pl. Cyp.* 304.—Afr. austr.
indicus, Royle, ex D. Don, in Proc. Linn. Soc. i. (1839) 10 ; et in Trans. Linn. Soc. xviii. (1840) 222 = prismatocarpus.
inflexus, Linn. Sp. Pl. 326 = glaucus !
inflexus, Retz. Prod. Fl. Scand. ed. II. 79 = balticus.
inflexus, With. Bot. Arr. ed. I. i. 211 = glaucus.
insularum, Viv. Fl. Cors. Sp. Nov. 5 = bufonius.
intermedius, Host, Gram. Austr. iii. 55. t. 99 = Luzula Desvauxii.
intermedius, Poir, Encyc. Suppl. iii. 158 = alpinus.
intermedius, Rohde, ex Buchen. Krit. Vers. Junc. 24 = acutiflorus.
intermedius, Thuill. Fl. Par. ed. II. 178 = Luzula campestris.
inundatus, Drejer, in Kroeyers, Nat. Tidsk. ii. (1838-39) 181 = balticus.
involucratus, T. Kirk, in Trans. N. Z. Inst. ix. (1877) 556 = enalifolius.
involucratus, *Steud. in Lechl. Berb. Am. Austr.* 58.—Forsv.

JUNCUS:—
iridifolius, Willd. ex Kunth, Enum. Pl iii. 337 = enalifolius.
isolepioides, Nees, in Linnaea, xx. (1847) 144 = inaequalis.
isthmiacus, Neck. Delic. Gallo-Belg. i. 168 = lampocarpus.
Jacquini, *Linn. Mant.* i. 68.—Europ.
Jacquini, Sm. in Linn. Fl. Lapp. ed. II 91 = arcticus.
Jacquini, Symons, Syn. Fl. Brit. 87 = castaneus.
Karelini, Steud. Syn. Pl. Cyp. 297 = acutus.
Kelloggii, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1858) 494.—Calif.
Kochii, F. Schultz, in Pollichia, xiii. (1855) 82 = supinus
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 102.—Persia.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* ii. 99.—Japon.
Kraussii, Hochst. in Flora, xxviii. (1845) 342 = spretus.
lactens, Link, in Schrad. Journ. ii. (1799) 316 = Luzula lactea.
laevis, Wallr. Sched. Crit. 142 = effusus.
lagenarius, J. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 130 = Fontanesii.
lampadocarpus, Davies, Welsh Botanol. 34 = lampocarpus.
lampocarpus, Baumg. Enum. Stirp. Transs. iii. 339 = Thomasii.
lampocarpus, *Ehrh. Calam.* n. 106; *ex Davies, in Trans. Linn. Soc.* x. (1810) 18.—Reg. bor. et austr.
lamprocarpus, Reichb. Fl. Germ. Excurs. 96 = prec.
lapponicus, Auct. ex Schult. f. Syst. vii. 177 = effusus.
latifolius, Wulf. in Jacq. Coll. iii. 59 = Luzula sylvatica
Lechleri, Steud. Syn. Pl. Cyp. 306 = Chamissonis.
Leersii, Marss. Fl. Neu-Vorpom. 451 = effusus.
Lehmanni, E. Mey. ex Hook. Fl. Bor. Am. ii. 192 = castaneus.
Leopoldii, Parl. in Giorn. Bot. Ital. ii. i. (1846) 324 = acutus.
leptocarpus, Buchen. in Goett. Nachr. (1869) 244 = glaucus.
leptocaulis, *Torr. & Gray, ex Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 454.—Am. bor. occ.
leptopetrus, *Buchen. in Engl. Bot. Jahrb.* vi. (1885) 203.—Reg. Himal.
Leschenaultii, J. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 137 = pumaticarpus.
Lesueurii, *Boland. in Proc. Calif. Acad.* ii. (1863) 179.—Calif.
leucanthus, *Royle, ex D. Don, in Proc. Linn. Soc.* i (1839) 10.—Reg. Himal.
leucomelas, *Royle, l. c.*—Reg. Himal.
limiger, Purt. Fl. Midl. Count. Engl. Suppl. 852. f. 9 = Luzula campestris.
Limbiwi, Siebold, in Verh. Bat. Gen. xii. (1830) 11 = effusus !
littoralis, C. A. Mey. Enum. Fl. Caucas. 24 = acutus.
lomatophloeus, Ehrh. Beitr. vi. 141 = Luzula albida.
lomatophyllus, *Spreng. Neue Entdeck.* ii. 108.—Afr. austr.
longicapsularis, Chev. Fl. Par. ii. 379 = lampocarpus.
longicornis, Bast. in Desv. Journ. Bot. iii. (1814) 21 = glaucus.
longifolius, Steud. Syn. Pl. Cyp. 299 = Dombeyanus.
longistylus, *Torr. in Bot. Mex. Bound.* 223.—Am. bor. occ.
Loureiranus, *Schult. f. Syst.* vii. 238.—Cochinch.
lucens, Burnham, in Trans. Linn. Soc. xix. (1846) 492 = effusus !
lucidus, Hochst. in Seub. Fl. Asor. 24. t. 4. f. 1 = tenuis.
Luetkei, Buchen. in Goett. Nachr. (1869) 243 = glaucus.
luteus, All. Fl. Pedem. ii. 216 = Luzula lutea.
lutosus, Chaix, in Vill. Hist. Pl. Dauph. i. 379 = Luzula lutea.
Lutzii-Astrei, Griseb. in Goett. Abh. xix. (1874) 268 = microcephalus.
Luzula, Krock. Fl. Siles. i. 560 = Luzula pilosa.
luzulaeformis, Gennari, Sp. Sard. (1857) 22 = bufonius.
luzulinus, Vill. Hist. Pl. Dauph. ii. 235 = Luzula flavescens.
luzuloides, Lam. Encyc. iii. 272 = Luzula nemorosa.
luzuloides, Steud. ex Schult. f. Syst. vii. 242 = capensis.
macer, S. F. Gray, Nat. Arr. Brit. Pl. ii. 164 = tenuis.
macrobotrys, Nees, in Linnaea, xx. (1847) 243 = acutus.
macrocephalus, Viv. Fl. Cors. Sp. Nov. 6 = lampocarpus
macrostemon, *Y. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par.* iii. (1827) 140 = scirpoides.
macrostigma, Colenso, in Trans. N. Z. Inst. xvii. (1885) 258 = pallidus.

JUNCUS:—
magellanicus, Lam. Encyc. iii. 366 = Rostkovia sphaerocarpa.
Maudeni, Buchen. in Bremen, Abh. iv. (1874) 121 = glaucus.
marginatus, *Roxb. Diss. Juncc.* 38.—Am. bor.
maritimus, *Lam. Encyc.* iii. 264.—Reg. temp.
maritimus, E. Mey, in Linnæa, vii. (1832) 180 = spretus.
maritimus, Moric. Fl. Venet. i. 179 = acutus.
maritimus, Bot. Hort. Mauric. 360 = effusus.
maximus, Retchard, Fl. Moeno-Francof ii. 192 = Luzula sylvatica.
maximus, Retz, Prod. Fl. Scand. ed. II. 82; Brot. Fl. Lusit. i. 515 = Luzula sylvatica.
maximus, Roth, Tent. Fl. Germ. i. 157, partim = Luzula albida.
megacephalus, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 132 = scirpoides.
megacephalus, Wood, Class-Book (1861) 734 = nodosus.
megalocos, Steud. Syn. Pl. Cyp. 298 = Dombeyanus.
melananthus, Reichb. Fl. Germ. Excurs. 96 = stratus.
melanocarpus, Michx. Fl. Bor. Am. i. 190 = Luzula parviflora.
melanocarpus, Phil. ex Buchen. Krit. Verz. Junc. 29 = stipulatus.
melanocephalus, Boiss. Fl. Orient. v. 360 = elipgenus.
melanocephalus, Frivald. in Flora, xix. (1836) ii. 487 = Thomasii.
membranaceus, *Royle, ex D. Don, in Proc. Linn. Soc.* i. (1839) 10; *et in Trans. Linn. Soc.* xviii. (1840) 320.—Reg. Himal.
Menziesii, R. Br. ex Hook. Fl. Bor. Am. ii. 192 = falcatus.
Mertensianus, *Bong. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 167.—Am. bor. occ.
Metsleri, F. Schultz, in Pollichia, xlii. (1855) 34 = compressus.
mexicanus, *Willd. ex Schult. f. Syst.* vii. 178.—Mexic.
micranthus, Desv. Obs. Pl. Ang. 82 = acutiflorus.
micranthus, Schrad. in E. Mey. Syn. Lus. 31 = scirpoides.
microcarpus, Nolte, ex Gaud. Fl. Helv. ii. 553 = alpinus.
microcephalus, *H. B. & K. Nov. Gen. et Sp.* i. 237.—Am. austr.
adilitaris, *Bigel. Fl. Bost.* ed. II. 189.—Am. bor.
Mimianni, Guill. ex Buchen. Krit. Verz. Junc. 30 = heterophyllus.
Minae, Strobl, ex Nym. Consp. 749 = pygmaeus.
minimus, *Buchen. in Bot. Zeit.* xxv. (1867) 145.—Reg. Himal.
minimus, E. Forst. in Symons, Syn. Pl. Brit. 197 = capitatus.
monanthos, Schrank, Naturh. Brief. ii. 208; ex ej. Baier. Fl. i. 930 = acq.†
monanthus, Jacq. Enum. Stirp. Vind. 61, 236 = trifidus.
montanus, Lam. Encyc. iii. 278 = Luzula sylvatica.
monticola, Steud. Syn. Pl. Cyp. 301 = prismatocarpus.
macrostiflorus, [Clairv.] Man. Herb. 104 = alpinus.
Muehlenbergii, Spreng. Syst. ii. 106 = pelocarpus.
Mueller, Trautv. in Act. Hort. Petrop. v. (1877) 119 = balticus.
multibractaetus, Tineo, in Guss. Suppl. Fl. Sic. Prod. 96 = acutus.
multiceps, Schult. f. Syst. vii. 198 = supinus.
multiceps, *Kunze, ex Kunth, Enum. Pl.* iii. 337.—Chili.
multiflorus, Bert. ex Buchen. in Bremen, Abh. vi. (1879) 379. t. 4 = Chamissonis.
multiflorus, Desf. Fl. Atlant. i. 213. t. 91 = subulatus.
multiflorus, Ehrh. [Calam]. 197; ex Hoffm. Fl. Deutschl. i. 169 = Luzula campestris.
multiflorus, Retz. Prod. Fl. Scand. ed. II. 82 = capensis.
mutabilis, Cav. Ic. iii. t. 296. f. 3 = capitatus.
mutabilis, Lam. Encyc. iii. 270 = supinus, pygmaeus, capitatus.
mutabilis, Savi, Fl. Pis. i. 364 = bufonius.
nanus, Dubois, Méthod Eprouv. Pl. Fr. 276 = pygmaeus.
nebrodensis, Tod. ex Parl. Fl. Ital. ii. 344 = Fontanesii.
Neesii, Heller, Fl. Wiroeb. Suppl. 27 = obtusiflorus.
nemorosus, Host, Gram. Austr. iii. t. 97. f. 2 = Luzula campestris.
nemorosus, Lam. Encyc. iii. 272 = Luzula vernalis.
nemorosus, Moench, Hass. 169 = Luzula sylvatica.
nemorosus, Pollich, Hist. Pl. Palat. i. 352 = Luzula albida.

JUNCUS:—
nemorosus, Sibth. Fl. Oxon. 114 = acutiflorus.
neozelandicus, *T. Wstz. in Proc. Am. Acad.* xiv. (1879) 303.—Am. bor. occ.
nigor, Wolff, ex Schult. f. Syst. vii. 208 = acutiflorus.
nigrescens, Schrank, ex Schult. f. l. c. 256 = Luzula spadicea.
nigricans, Drejer, ex Reichb. Ic. Fl. Germ. ix. 20 = ancepa.
nigricans, Willd. Sp. Pl. ii. 215 = Jacquini.
nigricans, Wolff, ex Schreb. in Schweigg. & Koerte, Fl. Erlang. 148 = acutiflorus.
nigritellus, D. Don, in Engl. Bot. Suppl. t. 2648 = lampocarpus.
nigritellus, Koch, Syn. Fl. Germ. ed. I. 780 = supinus.
nitidifolrus, Duf. in Ann. Sc. Nat. Sér. I. v. (1825) 86 = Gerardi.
nitidus, *Phil. Fl. Atac.* 53.—Chili.
nivalis, Kunth, Enum. Pl. iii. 310 = Luzula hyperborea.
niveus, Leers, Fl. Herborn. 91 = Luzula albida.
niveus, Linn. Syst. ed. X. 987; Sp. Pl. ed. II. 468 = Luzula nivea.
nodosus, *Linn. Sp. Pl.* ed. II. 466.—Am. bor.
nodosus, Schreb. ex Schult. f. Syst. vii. 215 = acuminatus, Dombeyanus.
nodulosus, Bory & Chaub. Exp. Sc. Morée, 105 = Luzula graeca.
nodulosus, Wahlenb. Fl. Upsal. 114 = alpinus.
Novae-Zandiae, *Hook. f. Fl. N. Zel.* i. 264.—N. Zel.
nutans, Vill. Fl. Delph. 34 = Luzula pilosa.
obtusatus, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 495.—Calif.
obtusatus, Kit. in Linnæa, xxxii. (1863) 381 = obtusiflorus.
obtusiflorus, *Ehrh. Beitr.* vi. 83.—Europ.; As. bor.
obtusiflorus, Ehdw. Skizze, 122 = alpinus.
obtusiflorus, Moench, Meth. 240 = lampocarpus.
ochraceus, *Buchen. in Bremen, Abh.* iii. (1872) 292.—Reg. Himal.
odoratus, Steud. Syn. Pl. Cyp. 304 = marginatus.
oliganthus, Phil. in Linnæa, xxix. (1857–58) 75 = stipulatus.
olympicus, Schott, Nyman & Kotschy, Analect. Bot. 3 = lampocarpus.
Oriabae, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 30 = mexicanus.
oxycarpus, E. Mey. in Drège, Zwei Pfl. Doesm. 194 = punctorius.
oxymeris, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1868) 468.—Calif.
pacificus, Engelm. l. c. 442, 490 = Lesueurii.
pallens, Pers. Syn. i. 384 = pygmaeus.
pallescens, Gaeseck. ex Lange, Consp. Fl. Groenl. t. 126 = Luzula campestris.
pallescens, Lam. Encyc. iii. 268 = acuminatus, Dombeyanus.
pallescens, E. Mey. ex Buchen. Krit. Verz. Junc. 34 = acuminatus.
pallescens, Schlecht. ex Buchen. in Bremen, Abh. vi. (1879) 402 = Sellowianus.
pallescens, Schrank, in Flora, ii. (1819) 445 = Luzula Hostii.
pallescens, Wahlenb. Fl. Lapp. 87 = Luzula campestris.
pallidiflorus, Beck, ex Buchen. Krit. Verz. Junc. 34; 64 = acutiflorus.
pallidus, *R. Br. Prod.* 258.—Austral.; N. Zel.
pallidus, Hoppe, ex E. Mey. Syn. Junc. 14 = glaucus.
pallidus, Willd. ex Kunth, Enum. Pl. iii. 348 = tenuis.
palustris, Gromow, ex Trautv. in Act. Hort. Petrop. ix. (1884) 388, nomen (Quid?)—Rossia.
paniculatus, Hoppe, ex Mert. & Koch, Deutschl. Fl. ii. 575, in obs. = glaucus.
paniculatus, Lucé, Topog. Nachr. Oesel, 107 = lampocarpus.
paucinodus, *Lang. Kurs, Phys. Fl. Ung. Nat.* (1846) 318.—Hungar.
papillosus, *Franch. & Sav. Enum. Pl. Jap.* ii. 98.—Japon.
paradoxus, Auct. ex Buchen. Krit. Verz. Junc. 34 = canadensis.
paradoxus, E. Mey. Syn. Junc. 30 = acuminatus.
Parryi, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 446.—Am. bor. occ.
parviflorus, Ehrh. Beitr. vi. 139 = Luzula parviflora.
parviflorus, Kit. in Linnæa, xxxii. (1863) 382 = compressus.

JUNCUS:—
parviflorus, Polr. Encyc. Suppl. iii. 160 = tenuis.
parviflorus, Retz. Prod. Fl. Scand. ed. II. 81 = Luzula spadicea.
parvulus, *E. Mey. & Buchen. in Bremen, Abh.* iv. (1875) 447.—Afr. austr.
patens, *E. Mey. Syn. Lusul.* 28.—Am. bor. occ.
pauciflorus, *R. Br. Prod.* 259.—As. or.; Austral.; N. Zel.
pauciflorus, T. Kirk, in Trans. N. Z. Inst. ix. (1877) 551 = antarcticus.
paucinodus, Moench, ex Schleich. Cat. Pl. Helv. 57 = arcticus.
pedatus, Jacq. ex Schult. f. Syst. vii. 264, in obs. = Luzula Hostii.
pedifornis, Chaix, in Vill. Hist. Pl. Dauph. i. 318; ii. 238 = Luzula pedifornis.
pelocarpus, A. Gray, Man. N. Am. Bot. ed. I. 507 = alpinus.
pelocarpus, *E. Mey. Syn. Lusul.* 30.—Am. bor.
persicus, Boiss. Diagn. Ser. I. vii. 101 = Gerardi.
phaeocephalus, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 484.—Calif.
phyzocaulroides, Baenitz, Prosp. Herb. (1878) 4 = capitatus.
pictus, Phil. in Linnæa, xxxiii. (1864–65) 268 = Lesueurii.
pilcus, *Steud. Syn. Pl. Cyp.* 305.—Afr. austr.
pilosus, Linn. Sp. Pl. 329 = Luzula vernalis.
pilosus, Schousb. Vextr. Marok. 181 = Luzula Forsteri.
pistilliferus, Jan, Cat. 333; ex Schult. f. Syst. vii. 268 = Luzula lutea.
planifolius, *R. Br. Prod.* 259.—Austral.; N. Zel.
platycaulis, C. Gay, Fl. Chil. vi. 146 = capillaceus.
platycaulis, H. B. & K. Nov. Gen. et Sp. i. 236 = Chamisonis.
plebeyus, R. Br. Prod. 259 = bufonius.
plumosus, Wall. ex Walp. Ann. iii. 652 = Luzula plumosa.
Pohlii, Steud. Syn. Pl. Cyp. 303 = densiflorus.
polycephalus, J. Gay, ex Laharpe, in Mém. Soc. Linn. Par. iii. (1818) 140 = densiflorus.
polycephalus, Hook. Fl. Bor. Am. ii. 190 = xiphioides.
polycephalus, Michx. Fl. Bor. Am. i. 192 = scirpoides.
polycephalus, Sm. Engl. Fl. ii. 177 = lampocarpus, supinus.
polytrichos, *E. Mey. & Buchen. in Bremen, Abh.* iv. (1875) 448.—Afr. austr.
Pondii, Wood, Class-Book (1861) 734 = acuminatus.
ponticus, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) n. 111. 94 = maritimus.
pratensis, Hoppe, ex Nym. Consp. 748 = lampocarpus.
pratensis, Wolff, ex Schult. f. Syst. vii. 206 = acutiflorus.
prismatocarpus, *R. Br. Prod.* 259.—As. or.; Austral.
procerus, *E. Mey. in Linnæa*, iii. (1828) 387.—Chili.
prolifer, H. B. & K. Nov. Gen. et Sp. i. 236 = bufonius.
prolifreus, Vahl, ex E. Mey. Syn. Junc. 37 = striatus.
proximus, Steud. Syn. Pl. Cyp. 306 = Dombeyanus.
punctorius, Lam. Encyc. iii. 269 = Dombeyanus.
punctorius, *Linn. f. Suppl.* 108; *Thunb. Prod. Pl. Cap.* 66.—Afr. austr.; As. occ.
pungens, Moris, ex Reichb. Fl. Germ. Excurs. 98 = acutus.
purpureus, Buch, in Abh. Akad. Berl. (1816) 362 = Luzula purpurea.
pusillus, *Buchen. in Bremen, Abh.* vi. (1879) 395.—Austral.; N. Zel.
pygmaeus, *Rich. in Thouill. Fl. Par.* ed. II. 178.—Europ.
pygmaeus, Savi, Bot. Etrusc. ii. 69 = bufonius.
Pylaei, Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 119 = effusus.
pyramidatus, *Laharpe, l. c.* 138.—Syria; Aegypt.
pyrenaeus, *Zenark. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* vi. (1828–34) 283.—Mont. Pyren.
Quartinianus, A. Rich. Tent. Fl. Abyss. ii. 329 = Fontanesii.
querioides, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 151 = bufonius.
radicans, Schleicht. in Linnæa, xviii. (1844) 442 = acuminatus.
radicans, Schur, Enum. Pl. Transs. 665 = lampocarpus.
Raddii, *Buchen. Krit. Verz. Junc.* 38.—Austral.
remotiflor, Nees, in Linnæa, xx. (1847) 268 = bufonius.

JUNCUS:—
variflorus, Hartm. Scand. Fl. ed. I. 141 = alpinus.
vaseuus, Wender. ex Schult. f. Syst. vii. 188 = glaucus.
vepens, J. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 198 = acutiflorus.
vepens, Maerkl. ex Schult. f. Syst. vii. 1660 = supinus.
repens, *Micks. Fl. Bor. Am.* i. 191.—Am. bor.
vepens, Nolte, Nov. Fl. Holsat. 88 = lampocarpus.
vepens, Req. in Guer. Desc. Vaucl. ed. II. 258 = Fontanesii.
Requienii, Parl. Fl. Ital. ii. 846 = alpinus.
retroflexus, Rafn, in Fl. Dan. t. 1872 = obtusiflorus.
revolutus, *R. Br. Prod.* 259.—Austral.
Richardsonianus, Schult. f. Syst. vii. 201 = alpinus.
rigidus, Desf. Fl. Atlant. i. 312 = maritimus.
rigidus, Wulf. ex Schult. f. Syst. vii. 273 = Luzula campestris.
riviularis, Poepp. ex Kunth, Enum. Pl. iii. 348 = cyperoides.
robustus, S. Wats. in Proc. Am. Acad. xiv. (1879) 302 = acutus.
Rochelianus, Schult. f. Syst. vii. 1658 = Thomasii.
Roemerianus, *Scheele, in Linnæa*, xxii. (1849) 348.—Am. bor. occ.
Rostkovii, E. Mey. in Presl, Rel. Haenk. i. 142 = phæocephalus.
Rostkovii, E. Mey. Syn. Junc. 36 = nodosus.
rostratus, *Buchen. in Bremen, Abh.* iv. (1875) 487.—Afr. austr.
rubellus, Hoppe, ex Schult. f. Syst. vii. 265 = Luzula nemorosa.
rubens, Lam. Encyc. iii. 366 = microcephalus.
rudis, Kunth, Enum. Pl. iii. 333 = microcephalus ?
rufus, Mielich. ex Huebn. in Flora, xxii. (1839) 490 = acutiflorus.
rugosus, Stend. Syn. Pl. Cyp 298 = acutiflorus.
rugulosus, Engelm. in Coult. Bot. Gea. vi. (1881) 224 = dubius.
rupestris, *Kunth, Enum. Pl.* iii. 344.—Afr. austr.
saginoides, Engelm. in Trans. Acad. Sc. St. Louis, ii. (1866) 436 = triformis.
salinus, Lange, in Willk. & Lange, Prod. Fl. Hisp. i. 180 = subulatus.
salsuginosus, Turcz. in Bull. Soc. Nat. Mosc. xxviii. (1855) i. 304 = Gerardi.
scabriusculus, *Kunth, Enum. Pl.* iii. 354.—Afr. austr.
schenchzeioides, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 100.—Chili; Ins. Antarct.
Schimperi, Hochst. in Flora, xxiv. (1841) i. Intell. 21 = punctorius.
Schlegrintweiti, Buchen. in Goett. Nachr. (1869) 255 = himalensis.
schoenoides, Mérat, Nouv. Fl. Par. ed. II. ii. 75 = acutiflorus.
Schrankianus, Moll, in Schrank & Moll, Naturh. Briefe, ii. 25 = trifidus.
scirpoides, *Lam. Encyc.* iii. 267.—Am. bor.
secundus, Poir. Encyc Suppl. iii. 160 = tenuis.
Sellowianus, *Kunth, Enum. Pl.* iii. 336.—Bras.
septangulus, Peterm. in Flora, xxvii. (1844) 361 = stratus.
serotinus, Schur, ex Buchen. Krit. Verz. Junc. 40 = stratus.
serratus, Linn. f. Suppl. 208 = Priesnium Palmita.
setaceus, Gerard, ex Buchen. Krit. Verz. Junc. 41 = Gerardi.
setaceus, *Roath. Sync.* 13. t. 1. f. 2; *Rafin. in Med. Repos. N. York*, v. (1808) 362.—Am. bor.
setaceus, Tenore, Fl. Neap. Prod. Suppl. 9 = capitatus.
setaceus, Van Hall, App. n. 12; ex ej. Fl. Belg. i. 388 = supinus.
setifolius, Ehrh. Beitr. vi. 58 = supinus.
siculus, Jan, Elench. i. 5.—Cf. Schult. f. Syst. vii. 205 = Fontanesii.
sieuleus, Tineo, in Guss. Fl. Sic. Syn. ii. 888 = ambulatus.
silvaticus, Auct. vide *sylvaticus*.
similis, Buchen. Krit. Verz. Junc 41 = cæspiticius.
simensis, J. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 187 = prismatocarpus.
singularis, *Steud. Syn. Pl. Cyp.* 306.—Afr. austr.
Smithii, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 444.—Am. bor.
Smithii, Kunth, Enum. Pl. iii. 349 = tenuis.
Soucheriensis, *Buchen. in Bremen, Abh.* iv. (1875) 476.—Afr. austr.

JUNCUS:—
sorantinus, Becker, in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 75 = compressus.
sorgnitius, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1843) 198 = Gerardi.
Sorrentonii, Parl. Fl. Ital. ii. 856 = pygmæus.
spadiceus, All. Fl. Pedem. ii. 216; Vill. Hist. Pl. Dauph. ii. 236 = Luzula spadicea.
spadiceus, Schreb. in Schweigg. & Koerte, Fl. Erlang 148 = acutiflorus.
spadiceus, Vill. Prosp. 18 = Luzula lutea.
spaniantkus, Steud. Syn. Pl. Cyp. 306 = capillaceus.
sparganiflolius, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* v. 856.—Syria.
sphacelatus, *Decne. in Jacquem. Voy. Bot.* 172. t. 172. Reg. Himal.
sphaerocarpus, *Nees, ex Funk, in Flora*, i. (1818) 521. —Europ.; Oriens.
sphaerocephalus, Salisb. ex Pons. in Mém. Soc. Linn. Par. iv. (1826) 561 = lampocarpus.
sphenostemon, *Buchen. in Engl. Bot. Jahrb.* vi. (1885) 233.—Reg. Himal.
spicatus, Lam. Encyc. iii. 274; Krock. Fl. Siles. i. 572. t. 52 = Luzula campestris.
spicatus, Linn. Sp. Pl. 329 = Luzula spicata.
spicatus, Walt. Fl. Carol. 124 = Luzula racemosa.
spicatus, Forsk. Fl. Aegypt. Arab. 75 = maritimus.
Sprengelii, *Nees, in Linnæa*, xx. (1847) 244 (excl. syn.).—Afr. austr.
Sprengelii, Willd. Prod. Fl. Berl. 125 = squarrosus.
spretus, *Schult. f. Syst* vii. 1656, in obs.—Afr. austr.
squamosus, Link, Enum. Hort. Berol. i. 805, spbalm. = squarrosus.
squarrosus, All. Fl. Pedem. ii. 215 = acutiflorus.
squarrosus, *Linn. Sp. Pl.* 327.—Europ.
stellatus, Desv. ex Spreng. Syst. ii 106 = striatus.
stellatus, Soland. ex Schult. f. Syst. vii. 222 = capitatus.
stellatus, Willd. ex Kunth, Enum. Pl. iii. 329 = Fontanesii.
stenophyllus, Steud. Syn. Pl. Cyp. 308 = capensis.
stipulatus, *Nees & Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 126.—Chili.
stoechasinus, Brot. Fl. Lusit. i. 516 = Luzula lactea.
stolonifer, Wohlleb. Suppl. Fl. Hal. 18 = lampocarpus.
striatus, Griseb. Spicil Fl. Rumel. ii. 407 = Thomasii.
striatus, *Schousb. ex E. Mey. Syn. Junc.* 27.—Europ.; Afr. bor.
strictus, Lucé, Topog. Nachr. Oesel, 107 = squarrosus.
stygius, *Linn. Syst.* ed. X. 967.—Europ.; Am. bor.
subglandulosus, Steud. Syn. Pl. Cyp. 308 = scabriusculus.
subincurvus, Steud. l. c. = repens.
submonocephalus, Steud. l. c. = Dregeanus.
subnodulosus, Schrank, Baier. Fl. i. 616 = obtusiflorus.
subpilosus, Gilib. Exercit. ii. 507 = Luzula campestris.
subtilis, E. Mey. Syn. Luzul. 31 = pelocarpus.
subulatus, *Forsk. Fl. Aegypt. Arab.* 75.—Reg. Mediterr.
subuliflorus, Drejer, Fl. Excurs. Hafn. 127 = effusus.
subuvrticillatus, Mahl. Desc. Uber. 209 = pelocarpus.
subverticillatus, Wulf. in Jacq. Coll. iii. 31 = supinus.
sudeticus, Krock. Fl. Siles. Suppl. ii. 34 = Luzula spicata.
sudeticus, Willd. Sp. Pl. 221 = Luzula campestris.
sulcatus, Hochst. ex Krauss, in Flora, xxviii. (1845) 343 = capensis.
supiniformis, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 441.—Am. bor. occ.
supinus, Bicheno, in Trans. Linn. Soc. xii. (1817) 317 = capitatus.
supinus, *Moench, Enum. Pl. Hass.* 167.—Europ.; Am. bor.
sylvaticus, Brot. Fl. Lusit. i. 517 = Thomasii.
sylvaticus, Desv. ex Schult. f. Syst. vii. 200 = lampocarpus.
sylvaticus, Host, Gram. Austr. iii. 57 = obtusiflorus.
sylvaticus, Huds. Fl. Angl. ed. I. 132 = Luzula sylvatica.
sylvaticus, Muhl. Cat. 36 = acuminatus, Dombeyanus.
sylvaticus, Parl. Fl. Palerm. i. 352 = Gussonii.
sylvaticus, Reichard, Fl. Moeno-Francof. ii. 151 = acutiflorus.
sylvaticus, Wahlenb. Fl. Lapp. 81 = alpinus.

JUNCUS:—
sylvaticus, Willk. & Lange, Prod. Fl. Hisp. i. 184 = Fontanesii, acutiflorus.
tasmanicus, Engelm. in Trans. Acad. Sc. St. Louis, ii. (1868) 458 = falcatus.
Tenagcia, *Ehrh. ex Linn. f. Suppl.* 208; *et Beitr.* i. 181.—Reg. Mediterr.
Tenagea, Host, Gram. Austr. iii. 60. t. 91 = sphaerocarpus.
tenagnius, Dulac, Fl. Hautes-Pyr. 103 (err. typ. 105) = præc.
tenax, *Buch, Beschr. Canar. Ins.* 191.—Ins. Canar.
tenax, Forst. f. Prod. 90 = pauciflorus ?
tenax, (Soland. in) Russell, Nat. Hist. Aleppo, ii. 251; Poir. Encyc. Suppl. iii. 156 = glaucus.
Tenegeia, Koth, Tent. Fl. Germ. ii. 409 = J. Tenagcia
tenellus, Geuns, Pl. Belg. Confed. Indig. Specil. 25 = capitatus.
tenuifolius, Steud. Syn. Pl. Cyp. ii 306 = Chamissonis.
tenuis, Sw. ex Schult. f. Syst. vii. 236 = compressus.
tenuis, *Willd. Sp. Pl.* ii. 214.—Europ.; Ind. occ.; Austral.
Thomasii, *Tenore, Ind. Sem. Hort. Neap.* (1827) 3.—Europ. austr.
Thomsoni, *Buchen. in Bot. Zeit.* xxv. (1867) 148.—Reg. Himal
thyrsiflorus, Vest, ex Schult. f. Syst. vii. 272 = Luzula spicata.
Tommasinii, Parl. Fl. Ital. ii. 315 = acutus.
transilvanicus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 77 = filiformis.
triandrus, Bosc, ex Schult. f. Syst. vii. 240 = marginatus.
triandrus, Gouan, Herbor. 26 = capitatus.
triandrus, Reichb. Ic. Fl. Germ. ix. t. 391 = pygmæus.
triandrus, Vill. Cat. Jard. Strasb. 81; ex E. Mey. Syn. Junc. 30 = supinus.
tricephalus, J. Gay, ex Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 132 = lampocarpus.
tricephalus, Wulf. ex Nym. Consp. 748 = supinus.
triceps, Rostk. Junc. 48 = castaneus.
trichocephalus, Cf. Bull. Férussac, xiii. (1828) 79 = lampocarpus.
trichodes, *Steud. Syn. Pl. Cyp.* 306.—Rossia.
trifidus, *Linn. Sp. Pl.* 326.—Reg. bor. et arct.
triformis, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 432.—Am. bor. occ.
triglumis, *Linn. Sp. Pl.* 328.—Reg. bor. et arct.
triglumis, Walt. Fl. Carol. 124 = marginatus.
trigonocarpus, *Steud. Syn. Pl. Cyp.* 308.—Am. bor.
trimervis, *Lisbon. in Kjoth. Vidensk. Meddel.* (1850) 41.—Mexic.
tristanianus, Hemsl. Bot. Voy. Challenger, i. 11. 154 = tenuis.
uliginosus, Kunth, Enum. Pl. iii. 334 = heterophyllus.
uliginosus, Roth, in Roem. & Usteri, Mag. i. 26tes St. (1787) 16 = supinus.
umbractatus, Griff. Notul. 329 = prismatocarpus.
uniglumis, Forst. ex Kunth, Enum. Pl. iii. 360 = Rostkovia grandiflora, *Hook. f.*
uruguensis, *Griseb. in Goett. Abh.* xxiv. (1879) 317.—Reg. Argent.
Urvillei, Steud. Syn. Pl. Cyp. 306 = capillaceus.
ustulatus, *Buchen. in Bremen, Abh.* vi. (1879) 401.—Bras.
ustulatus, Hoppe, Anleit. Grees. 80 = alpinus.
utriculosus, Auct. ex Schult. f. Syst. vii. 200 = lampocarpus.
vaccilans, Steud. Syn. Pl. Cyp. 305 = tenuis.
vaginatus, R. Br. Prod. 256 = pallidus.
vaginatus, Sieber, ex Schult. f. Syst. vii. 188 = paucifloras.
Vaillantii, Thuill. Fl. Par. i. 177 = J. Tenagcia.
Valdivius, Steud. Syn. Pl. Cyp. 296 = procerus.
valvatus, *Link, in Schrad. Journ.* ii. (1799) 316.—Lusit.; Alger.
variegatus, Carael, in Nuov. Giorn. Bot. ii. (1870) 276 = acutus.
Vaseyi, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 448.—Am. bor. occ.
vernalis, Bot. Fl. Lusit. i. 516 = Luzula Forsteri.
vernalis, Reichard, Fl. Moeno-Francof. ii. 152 = Luzula vernalis.
verticillatus, Pers. Syn. i. 384 = supinus.
viviparus, S. W. Conrad, in Journ. Acad. Philad. vi. (1832) 106 = pelocarpus.
viviparus, Relh. Fl. Cantab. 123 = supinus.

JUNCUS:—
vulgaris, Gueldenst. Reisen, i. 296 (Quid?).—Reg. Caucas.
Wallichianus, Laharpe, in Mém. Soc. Hist. Nat. Par. iii. (1827) 139 = prismatocarpus.
Woloszczakii, Hochst. ex Steud. Syn. Pl. Cyp. 304 = supinus.
Wulfeni, Micheli. ex Sauter, in Flora, xxxii. (1849) 666 = acutiflorus.
xanthocephs, Steud. Syn. Pl. Cyp. 308 = planifolius.
xiphioides, *E. Mey. Syn. Junc.* 50.—Am. bor. occ. ; Mexic.
zebrinus, Hort. Angl. ex André, Illustr. Hortic. xxvii. (1880) t. 393 = effusus.
zeylanicus, *E. Mey. Syn. Junc.* 66 (Quid?).—Zeylan.

JUNDZILLIA, Andrz. ex DC. Syst. ii. 529 (1821) = **Lepidium**, Tourn. (Crucif.).
Draba, Andrz. l. c. = L. Draba.

JUNGHANSIA, J. F. Gmel. Syst. 259 (1791) = **Curtisia**, Ait. (Cornac.).
fagrea, J. F. Gmel. l. c. = C. fagines.

JUNGHUHNIA, Miq. Fl. Ind. Bat. i. ii. 412 (1859) = **Codlaeanum**, Rumph. (Euphorb.).
glabra, Miq. l. c. = C. variegatum.

JUNGIA, Gaertn. Fruct. i. 175. t. 35 (1788) = **Baeckea**, Linn. (Myrtac.).
imbricata, Gaertn. l. c. = B. crenulata.
tenella, Gaertn. l. c. 176 = Micromyrtus microphylla.

JUNGIA, Heist. ex Moench, Meth. 378 (1794) = **Salvia**, Tourn. (Labiat.).
altissima, Moench, l. c. 379 = S. mexicana.
formosa, Moench, Meth. Suppl. 140 = S. leonuroides.
micrantha, Moench, l. c. 141 = S. tenella.

JUNGIA, Linn. f. Suppl. 58 (1781). *COMPOSITAE*. Benth. & Hook. f. ii. 502.
DUMERILIA, Lag. ex DC. in Ann. Mus. Par. xix. (1812) 71. t. 6 (15) et 7 (16).
MARTRASIA, Lag. Amen. Nat. i. 36 (1811).
PLEOCARPHUS, D. Don, in Trans. Linn. Soc. xvi. (1830) 228.
RHINACTINIA, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 139.
TRINACTS, Gaertn. Fruct. ii. 415 (1791).
affinis, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 460 = floribunda.
auriculata, *Steud. Nom.* ed. II. i. 834.—Hab. ?
axillaris, *Spreng. Syst. iv. Cur. Post.* 301.—Reg. Andin.
bullata, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 213.—Peruv.
crenata, *Steud. Nom.* ed. II. i. 834.—Hab. ?
ferruginea, *Linn. f. Suppl.* 390.—Reg. Andin.
floribunda, *Less. in Linnaea*, v. (1830) 88.—Bras.
floribunda, *Spruce, ex Benth. & Hook. f. Gen.* ii. 502. —Bras.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 537.—Bolivia.
paniculata, *A. Gray, in Proc. Am. Acad.* v. (1861) 148.—Peruv.
polita, *Griseb. in Goett. Abh.* xxiv. (1879) 215.—Reg. Argent.
pyramidalis, D. Don, in Trans. Linn. Soc. xvi. (1833) 299 = floribunda.
rugosa, *Less. in Linnaea*, v. (1830) 86.—Am. trop.
Sellowii, *Less. Syn. Comp.* 416.—Bras.
serices, *Steud. Nom.* ed. II. i. 834.—Hab. ?
spectabilis, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 297.—Peruv.
tomentosa, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. iii. 399 = floribunda.

JUNGIA, Loefl. ii. 199 (1758) = **Ayenia**, Linn. (Sterculiac.).

JUNIA, Adans. Fam. ii. 165 (1763) = **Clethra**, Linn. (Vaccin.).

JUNIPERUS, Tourn. ex Linn. Syst. ed. I (1735). *CONIFERAE*. Benth. & Hook. f. iii. 427.
ARCEUTHOS, Ant. & Kotschy, in Oestr. Bot. Wochenbl. (1854) 249.
OXYCEDRUS, Hort. ex Carr. Conif. ed. II. 18 (1867).
SABINA, Hall. in Rupp. Fl. Jen. ed. II. 336 (1745).
THUIARCARPUS, Trautv. Imag. Pl. Ross. 11. t. 6 (1844).

JUNIPERUS:—
abyssinica, Hort. ex C. Koch, Dendrol. ii. 11. 132 = procera.
acerosa, Griseb. Veg. Erde, 378, 572 = excelsa.
alba, Knight, ex Carr. Conif. ed. I. 58 = virginiana.
alpina, S. F. Gray, Nat. Arr. Brit. Pl. ii. 238 = communis.
alpina, Lodd. Cat. (1836) 48 = J. Sabina.
andina, Nutt. N. Am. Sylv. iii. 93. t. 110 = virginiana.
aquatica, Roxb. Hort. Beng. 73; Fl. Ind. iii. 838 = Taxodium heterophyllum.
arborea, *Budischef. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 314.—Manchur.
arborescens, Moench, Meth. 699 = virginiana.
argaea, Bal. ex Parl. in DC. Prod. xvi. ii. 480 = communis.
argentea, Hort. ex Gord. Pinet 118 = virginiana.
aromatica, Hort. ex Carr. Conif. ed. I. 127 = Cupressus Goveniana.
attica, Orph. in Heldr. Nutzpfl. Griech. 18 = macrocarpa.
australis, Endl. Syn. Conif. 26 = virginiana.
baccifornis, Carr. Conif. ed. I. 56 = phoenicea.
barbadensis, Linn. Sp. Pl. 1039 = bermudiana.
barbadensis, Thunb. Fl. Jap. 264 = chinensis.
Bedfordiana, Hort. ex Loud. Encyc. Trees, 1090 = virginiana.
bermudiana, *Linn. Sp. Pl.* 1039.—Ins. Bermud.
Biasoletti, Link, in Atti 7 ma. Riun. Scienz. Ital. Napoli, (1846) 878 = macrocarpa.
Biebersteiniana, Hort. ex C. Koch, Dendrol. ii. ii. 112 = J. Oxycedrus.
Bonatiana, Vis. in Mem. Ist. Venet. vi. (1856) 245 = thurifera.
borealis, Salisb. Prod. 397 = communis.
Bregeoni, *Regel, Ind. Sem. Hort. Petrop.* (1858) 23.—As. bor.
brevifolia, *Antoine, Cupress. Gatt.* 16. tt. 20–22.—Ins. Azor.
butanensis, Wender. Pfl. Bot. Gaert. Marb. 36 = recurva.
Cabiana, Vis. in Mem. Ist. Venet. vi. (1856) 246 = chinensis.
caesia, Carr. Conif. ed. I. 55 = Callitris arborea.
caesia, *Regel, Gartenfl.* (1857) 246.—Calif.
californica, *Carr. in Rev. Hortic. Sér.* IV. iii. (1854) 352.—Am. bor. occ.
canadensis, Lodd. Cat. (1836) 47 = communis.
canariensis, Knight, ex Gord. Pinet. 114 = J. Cedrus.
canescens, Wall. ex Gord. Pinet. 106 = recurva.
Cannartii, Hort. ex C. Koch, Dendrol. ii. ii. 140 = virginiana.
caperata, Lam. Encyc. ii. 626 = Callitris arborea.
carolinana, Mill. Gard. Dict ed. VIII. n. 4 = virginiana.
carolinisana, Duroi, Harbk. ed Pott, i. 497 = prasc.
cascacica, Fisch. ex Gord. Pinet. 98 = communis.
Cedrus, *Webb & Berth. Phyt. Canar.* iii. 277 t. 2. ff. 1–3.—Ins. Canar.
cernua, Roxb. Hort. Beng. 73; Fl. Ind. iii. 839 = chinensis.
cernuiformis, Kellogg, in Proc. Calif. Acad. ii. (1863) 57 = californica.
Chamberlayni, Carr. Man. Fl. iv. 313 = virginiana.
chinensis, Hort. ex Endl. Syn. Conif. 29 = J. Sabina.
chinensis, *Linn. Mant.* i. 127.—China.
chinensis, Roxb. Hort. Beng. 73; Fl. Ind. iii. 840 = Podocarpus macrophylla.
cinerascens, Hort. ex C. Koch, Dendrol. ii. ii. 141 = virginiana.
cinerea, Carr. Conif. ed. II. 35 = thurifera.
communis, *Linn. Sp. Pl.* 1040.—Reg. bor. temp. et arct.
communis, Thunb. Fl. Jap. 264 = rigida.
compressa, Hort. ex Carr. Conif. ed. I. 52 = communis.
conferta, *Parl. Nov. Conif.* 1.—Japon.
Cracovia, Hort. ex C. Koch, Dendrol. ii. ii. 115 = communis.
Cunninghamii, Hort. ex Carr. Conif. ed. I. 71 = Callitris quadrivalvis ?
daurica, Hort. ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 200 = communis.
davurica, *Pall. Fl. Ross.* i. ii. 12. t. 58.—Sibir.
deallata, Hort. ex Knight, Syn. Conif. 19 = occidentalis.
deallata, Lond. Encycl. Trees, 1090 = communis.
densa, Gord. Suppl. Pinet. 95 = recurva.
Deppeana, Steud. Nom. ed. II. i. 835 = mexicana.
depressa, Rafin. Med. Fl. ii. 12 = communis ?
diffusula, Gilib. Exercit. ii. 418 = communis.
dimorpha, Roxb. Hort. Beng. 73; Fl. Ind. iii. 839 = chinensis.

JUNIPERUS:—
dioica, Hort. ex Carr. Conif. ed. II. 45 = virginiana.
dimoricata, Hort. ex Carr. l. c. 51 = phoenicea.
drupacea, *Labill. Pl. Syr. Decad.* ii. 14. t. 8.—Graecia ; As. Min. ; Syria.
dumosa, Hort. ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 201 = J. Pseudo-sabina ?
duplicata, Goepp. Ind. Hort. Vratislav. (1871) = drupacea ?
echinoformis, Hort. ex Carr. Conif. ed. II. 18 = J. Oxycedrus.
echinoformis, Rinz, ex Bolse, in C. Koch, Wochenschr. xi. (1868) 284 = communis.
elata, Roxb. Hort. Beng. 73; Fl. Ind. iii. 838 = Dacrydium elatum.
elliptica, Hort. ex Carr. Conif. ed. I. 11 = macrocarpa.
elliptica, Hort. ex C. Koch, Dendrol. ii. ii. 118 = communis.
erioides, Hort. ex Carr. Conif. ed. II. 140 = Thuja orientalis.
erioides, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 495 = Thuja occidentalis.
erioides, Nois. ex Desf. Cat. Hort. Par. ed. III. 355 = Callitris calcarata.
excelsa, *Bieb. Fl. Taur. Cauc.* ii. 245.—As. Min.
excelsa, Griff. Notul. iv. 26 = macropods.
excelsa, Madden, ex Gord. Pinet. 107 = recurva.
excelsa, Pursh, Fl. Am. Sept. ii. 647 = occidentalis.
excelsa, Wall. Cat. n. 6041 = J. Pseudo-sabina.
excelsa, Webb, ex Parl. in DC. Prod. xvi. ii. 488 = chinensis.
excelsa, Willd. Sp. Pl. iv. 852 = J. Sabina.
fastigiata, Hort. ex Knight, Syn. Conif. 11 = communis.
flaccida, *Schlecht. in Linnaea*, xii. (1838) 495.—Mexic.
flagelliformis, *Hort. ex Loud. Encyc. Trees*, 1090 = chinensis.
foetida, Spach, Hist. Vég. Phan. xi. 314 = foetidissima.
foetidissima, Hort. ex Endl. Syn. Conif. 30 = communis.
foetidissima, *Willd. Sp. Pl.* iv. 853.—Graecia ; As. Min.
formosa, Hort. ex Carr. Conif. ed. II. 51 = phoenicea.
Fortunei, Hort. ex Carr. l. c. ed. I. 11 = macrocarpa.
Fortunei, Hort. ex Carr. l. c. ed. II. 32 = sphaerica.
fragrans, Knight, Syn. Conif. 18; Salisb. Prod. 397 = virginiana.
fragrans, Pasl, ex Gord. Pinet. Suppl. 37 = cassia.
gigantea, Kossl. Cat. 8 = mexicana.
glauca, Salisb. Prod. 397 = J. Oxycedrus.
glauca, Willd. Enum. Hort. Berol. Suppl. 67, nomen ; Cels, ex Link, Enum. Hort. Berol. ii. 455 = virginiana.
gossainthanea, Lodd. ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 202 = virginiana.
gracilis, Endl. Syn. Conif. 31 = virginiana.
gracilis, Hort. ex Carr. Conif. ed. I. 56 = Cupressus Comeyana.
grandiflolius, Link, in Buch, Beschr. Canar. Ins. 159 = J. Oxycedrus.
Hartwiesiana, *Spec. ex Koppen, Veh. Pfl. Acclimat. in Russl.* 6.—Hab. ?
hemisphaerica, Presl, Delic. Prag. 142 = communis.
Henryana, *R. Br. Camed. in Gard. Chron.* (1873) 8.—Am. bor. occ.
Hermanni, Pers. Syn. ii. 632 = virginiana.
heterocarpa, Timb. ex Nym. Conap. 676 = J. Oxycedrus.
hibernica, Lodd. ex Loud. Arboret. iv. 2490 = communis.
himalayensis, Hort. ex Carr. Man. Fl. iv. 314 = macropoda.
hispanica, Booth, ex Endl. Syn. Conif. 15 = communis.
hispanica, Mill. Gard. Dict. ed. VIII. n. 13 = thurifera.
Hochstetteri, *Antoine, Cupress. Gatt.* t. 83.—Abyssin.
horizontalis, Moench, Meth. 699.
hudsonica, Forb. Pinet. Woburn. 208 = J. Sabina.
humilis, Salisb. Prod. 397 = J. Sabina.
incurva, Buch.-Ham. ex Endl. Syn. Conif. 18 = recurva.
indica, Bertol. Mem. Bot. Bonn. 16 = J. Pseudo-sabina.
intermedia, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1850) 189 = communis.
interrupta, Wendl. ex Endl. Syn. Conif. 16 = communis.
isophylla, C. Koch, in Linnaea, xxii. (1849) 304 = excelsa.
japonica, Hort. ex Carr. Conif. ed. I. 32 = chinensis.
Lambertiana, Wall. ex Endl. Syn. Conif. 19 = recurva.

JUNIPERUS:—
Langoisiana, Hort. ex Gord. Pinet. Suppl. 38 = phoenicea.
Landoliana, Lawa. ex Gord. Pinet. 105 = excelsa.
Ittoralis, Maxim. in Bull. Acad. Pétersb. xii. (1868) 230 = conferta.
Lobelii, Guss. Fl. Sic. Syn. ii. 685 = macrocarpa.
Lorulaei, Hort. ex C. Koch, Dendrol. ii. 11. 185 = religiosa.
Lorulaei, Hort. ex Carr. Conif. ed. II. 36 = excelsa.
Lustitanica, Mill. Gard. Dict. ed. VIII. n. 11 = J. Sabina.
lycia, Linn. Sp. Pl. 1039 = phoenicea.
lycia, Pall. Fl. Ross. ii. 14. t. 56 = J. Sabina.
lycia, Siev. in Pall. N. Nord. Beitr. vii. 280 = J. Pseudosabina.
Macnabiana, Laws. ex Gord. Pinet. 64 = Cupressus Macnabiana.
macrocarpa, *Sibth. & Sm. Fl. Graec. Prod.* ii. 268.—Graec.; As. Min.
macrocarpa, Tenore, Syll. Fl. Neap. 482, partim = J. Oxycedrus.
macropoda, *Boiss. Fl. Orient.* v. 709.—Persia; Reg. Himal.
maiacocarpa, Carr. Conif. ed. II. 50 = phoenicea.
Marschalliana, Strev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 244; et xxx. (1857) i. 397 = J. Oxycedrus.
meldensis, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 495 = Thuya orientalis.
mexicana, *Schiede, in Linnaea*, v. (1830) 77.—Mexic.
mexicana, Spreng. Syst. iii. 909 = tetragona.
microphylla, *Antoine, Cupress. Gatt.* t. 31, 32.—Hab.?
montana, Hort. ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 200 = communis.
myosuros, Hort. ex Sénécl. Cat. (1854) 38 = phoenicea.
nana, Willd. Berl. Baums. 159 = communis.
neaboriensis, Laws. ex Gord. Pinet. 96 = macrocarpa.
nepalensis, Hort. ex Endl. Syn. Conif. 21 = chinensis.
nepalensis, Hort. ex Loud. Arboret. iv. 2499 = berm-udiana.
nepalensis, Rinz. ex Gord. Pinet. 106 = recurva.
nipponica, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 280.—Japon.
oblonga, Bieb. Fl. Taur. Cauc. ii. 496 = communis.
oblongata, Guss. Fl. Sic. Syn. ii. 685, in textu = macrocarpa, phoenicea.
occidentalis, *Hook. Fl. Bor. Am.* ii. 166.—Am. bor. occ.
occidentalis, Hort. ex Carr. Conif. ed. I. 54 = communis.
Olderiri, Carr. l. c. 57 = excelsa.
ophora, Kunze, in Flora, xxix. (1846) 687 = phoenicea.
oppositifolia, Moench, Meth. 699 = bermudiana.
oxycedrus, Sr. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = J. Oxycedrus.
Oxycedrus, Herb. Linn. ex Parl. in DC. Prod. xvi. 11. 477 = macrocarpa.
Oxycedrus, Hochst. Fl. Azor. 98 = J. Cedrus.
Oxycedrus, *Linn. Sp. Pl.* 1038.—Europ. austr.; Caucas.
pachyphlaea, *Torr. in Pacif. Rail. Rep.* iv. 142.—Mexic.
patens, *Raeb. Hort. Berg.* 78, nomen.—China.
pendula, Hort. ex Parl. in DC. Prod. xvi. 11. 470 = Cupressus torulosa.
Perbaani, Hort. ex Gord. Pinet. ed. IV. 144 = excelsa.
Philippiana, Wall. ex Carr. Conif. ed. I. 488 = Decrylium elatum.
phoenicea, *Linn. Sp. Pl.* 1040.—Reg. Mediterr.; Arab.
phoenicea, Pall. Fl. Ross. ii. 16. t. 57 = excelsa.
pojocarpos, C. Koch, in Linnaea, xxii. (1849) 303 = excelsa.
procera, *Hochst. ex Endl. Syn. Conif.* 26.—Afr. trop.
procumbens, Siebold, ex Lindl. in Journ. Hort. Soc. (1850) 253 = recurva.
procumbens, Siebold, in Ann. Soc. Hort. Pays-Bas (1844) 31 = japonica.
prostrata, Pers. Syn. ii. 632 = J. Sabina.
prostrata, *Risso, Hist. Nat. Europ. Merid.* ii. 459.—Europ. austr.
Pseudo-Sabina, *Fisch. & Mey. Sem. Hort. Petrop.* viii. 66.—Sconpar.; Reg. Himal.
pygmaea, C. Koch, in Linnaea, xxii. (1849) 303 = communis.
pyramidalis, Hort. ex Carr. Conif. ed. I. 38 = communis.
pyramidalis, Hort. ex Carr. l. c. ed. II. 48 = virginiana.
pyramidalis, Hort. ex Carr. l. c. ed. I. 51 = phoenicea.
pyramidalis, Salisb. Prod. 397 = bermudiana.

JUNIPERUS:—
pyriformis, A. Murr. in Gard. Chron (1855) 420 = occidentalis.
quadrangula, Miq. Cat. Hort. Bot. Amstel. 20 = Cupressus funebris.
racemosa, *Risso, Hist. Nat. Europ. Merid.* ii. 459.—Europ. austr.
recurva, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 55.—Reg. Himal.
Reevesiana, Hort. ex Endl. Syn. Conif. 31 = chinensis.
reflexa, Hort. ex Gord. Pinet. 98 = communis.
religiosa, Hort. ex Endl. Syn. Conif. 26 = excelsa.
religiosa, Royle, Illustr. Bot. Himal. 351 = recurva.
repanda, Hort. ex Carr. Conif. ed. I. 27 = recurva.
repens, Nutt. Gen. Am. ii. 245 = J. Sabina.
rhodocarpa, *Stev. in Bull. Soc. Nat. Mosc.* xxx. (1857) i. 397, in texto.—Reg. Caucas.
rigida, Gord. Pinet. 99 = taxifolia.
rigida, Nois. ex Desf. Cat. Hort. Paris, ed. III. 355 (= *Prenela rigida*).—Austral.
rigida, Pav. ex Carr. Conif. ed. I. 447 = Podocarpus glomeratus.
rigida, *Sieb. & Zucc. Fl. Jap.* ii. 109. t. 125.—Japon.
rigida, Wall. ex Carr. Conif. ed. II. 692 = Dacrydium elatum.
rigida, Wall. ex Endl. Syn. Conif. 19 = recurva
rufescens, Link, in Flora, xxiv. (1846) 579 = J. Oxycedrus.
Sabina, Gueldenst. Reisen, i. 98 = excelsa.
Sabina, *Linn. Sp. Pl.* 1039.—Europ.; As. bor.
Sabina, Pall. Reise, iii. 276 = J. Pseudo-Sabina.
Sabina, Sibth. & Sm. Fl. Graec. Prod. ii. 264 = foetidissima.
sabinoides, Endl. Syn. Conif. 24 = thurifera.
sabinoides, Grisch. Spicil. Fl. Rumel. ii. 352 = foetidissima.
sabinoides, Humb. ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 202 = mexicana.
saxatilis, Hort. ex Lindl. & Gord. l. c. 200 = communis.
Schottii, Hort. ex Gord. Pinet. Suppl. 36 = virginiana.
semiglobosa, *Regel, in Act. Hort. Petrop.* vi. (1879) 487.—Turkestan.
Sheppardii, Hort. ex Gord. Pinet. ed. II. 426 = sphaerica.
sibirica, Burgsd. Anleit. n. 272 = communis.
sinensis, J. F. Gmel. Syst. 1005 = chinensis.
sinensis, Hort. ex Carr. Conif. ed. II. 33 = sphaerica.
sphaerica, *Lindl. in Paxt. Flow. Gard.* i. (1850–51) 55. f. 35.—China.
sphaerocarpa, Antoine, Cupress. Gatt. II. t. 10 = macrocarpa.
squamata, Buch.-Ham. in Lamb. Pin. ed. I. ii. 17 = recurva.
squamata, Gord. Pinet. 110, ex parte = J. Pseudo-Sabina.
squamosa, Buch.-Ham. in Wall. Cat. n. 6043 = recurva.
stricta, Hort. ex Gord. l. c. ed. II. 144 = excelsa.
struthiacea, Knight, Syn. Conif. 13 = chinensis.
succica, Mill. Gard. Dict. ed. VIII. n. 3 = communis.
tamariscifolia, Hort. ex C. Koch, Dendrol. ii. 11. 125 = J. Sabina.
taurica, Hort. ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 200 = communis.
taurica, Strangways, ex Gord. Pinet. 100 = J. Oxycedrus.
taxifolia, *Hook. & Arn. Bot. Beech. Voy.* 271.—China; Japon.
tenella, Antoin. Cupress. Gatt. 20, 27–29 = J. Oxycedrus.
terminalis, Salisb. Prod. 397 = phoenicea.
tetragona, Moench, Meth. 699 = phoenicea.
tetragona, *Schlechtl. in Linnaea*, xii. (1838) 495.—Mexic.
Thunbergii, Hook. & Arn. Bot. Beech. Voy. 271 = chinensis.
thurifera, Bonpl. ex Endl. Syn. Conif. 68 = Cupressus thurifera.
thurifera, Hort. ex Carr. Conif. ed. II. 50 = phoenicea.
thurifera, Hort. Par. ex Endl. Syn. Conif. 25 = foetidissima.
thurifera, *Linn. Sp. Pl.* 1039.—Hispan.; Lusit.
turbinata, Guss. Fl. Sic. Syn. ii. 634 = phoenicea.
Ukdaensa, Gord. in Journ. Hort. Soc. v. (1850) 206 = Cupressus Lindleyi.
umbilicata, Godr. in Gren. & Godr. Fl. Fr. iii. 158, adnot. = macrocarpa.
uvifera, D. Don, ex Lamb. Pin. ed. II. ii. 116 = Libocedrus tetragona.

JUNIPERUS:—
uvifera, Loud. Encyc. Trees, 1058 = recurva.
vernata, Hort. ex Gord. Pinet. ed. II. 144 = excelsa.
virginiana, Endl. Syn. Conif. 27 = bermudiana.
virginiana, *Linn. Sp. Pl.* 1039.—Am. bor.
virginica, Thunb. Fl. Jap. 264 = chinensis.
Wallichiana, Hook. f. ex Parl. in DC. Prod. xvi. 11. 482 = J. Pseudo-Sabina.
Webbii, Carr. Conif. ed. I. 18 = J. Cedrus.
Whitleyana, Miq. Cat. Hort. Bot. Amstel. 21 = Cupressus sempervirens.
Willdommii, Antoine, Cupress. Gatt. 9. t. 7 = macrocarpa.
Wilhmanniana, Hort. ex Carr. Conif. ed. II. 18 = communis.
Wittmannianna, Fisch. ex Lindl. in Journ. Hort. Soc. v. (1850) 200 = J. Oxycedrus.
Wittmannianna, Stev. ex Parl. in DC. Prod. xvi. 11. 479 = communis.

JUNKIA, Rttg. in Marb. Schrift. ii. (1830) 128, sphalm. = **Funkia**, Spreng. (Liliac.).

JUNO, Tratt. Ausw. Gartenpfl. i. 135 (1817 ?) = **Iris**, Linn.
Aitchisoni, Klatt, in Abh. Naturf. Ges. Halle, xii. (1869) 362 = I. Aitchisoni.
Auckeri, Klatt, in Bot. Zeit. xxx. (1872) 498 = I. fumosa.
caucasica, Tratt. Ausw. Gartenpfl. i. 136 = I. caucasica.
Danfordiae, Klatt, in Abh. Naturf. Ges. Halle, xii. (1869) 362 = I. Danfordiae.
falcifolia, Klatt, in Bot. Zeit. xxx. (1872) 498 = I. palaestina.
persica, Tratt. Ausw. Gartenpfl. i. 136 = I. persica.
planifolia, Aschers. in Bot. Zeit. xxii. (1864) 114 = I. alata.
scorpioides, Tratt. Ausw. Gartenpfl. i. 135 = I. alata.
Stocksii, Klatt, in Abh. Naturf. Ges. Halle, xii. (1869) 363 = I. Stocksii.

JUNQUILLA, Fourn. in Ann. Soc. Linn. Lyon, N. S. xvi. (1869) 164 = JONQUILLA, Haw. = **Narcissus**, Tourn. (Amaryll.).

JUPICA, Rafin. Fl. Tellur. ii. 15 (1836) = **Kyris**, Linn.
caerulea, Rafin. l. c. = X. americana.

JURGENSENIA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 251 = **Befaria**, Mutis (Ericac.).
mexicana, Turcz. l. c. = B. discolor.

JURGENSIA, Benth. & Hook. f. Gen. i. 970 (1867) = JUERGENSIA, Spreng. = **Alkodsia**, Thou. (Violar.).

JURGENSIA, Rafin. Sylva Tellur. 146 (1838) = **Spermacoce**, Linn. (Rubiac.).
psylloides, Rafin. l. c. 147 = S. tenella.

JURINEA, Cass. in Bull. Soc. Philom. (1821) 140 = *COMPOSITAE*, Benth. & Hook. f. ii. 478.
ARGOPODON, Bohs. Diagn. Ser. I. vi. 123 (1845).
DERDERIA, Jaub. & Spach, Illustr. i. 129. t. 67 (1843).
DOLOMIAEA, DC. in Genil. Arch. Bot. ii. (1838) 330.
JURINELLA, Jaub. & Spach, Illustr. ii. 101. t. 183–185 (1846).
OUTREYA, Jaub. & Spach, l. c. i. 131. t. 68 (1843).
POLYTAXIS, Bunge, Del. Sem. Hort. Dorpat. (1842).
STECHMANNIA, DC. Prod. vi. 649 (1837).
TULAKRNIA, Rafin. Fl. Tellur. iv. 120 (1836).
Abramovi, *Regel & Herd. ex Regel, Fl. Nov. Fedsch.* 53.—Turkest.
adenocarpa, Schrenk, Enum. Fl. Nov. 46 = Saussurea Candolleana.
alata, *Cass. in Dict. Sc. Nat.* xxiv. 288.—Reg. Caucas.
albicaulis, *Bunge, in Flora*, xxiv. (1841) 156.—Sibir.
alpigena, *C. Koch, in Linnaea*, xxiv. (1851) 414.—As. Min.
ambigua, *DC. Prod.* vi. 675.—Rossia.
anatolica, *Boiss. Diagn.* Ser. I. iv. 21.—As. Min.
angustifolia, *Bunge, in Flora*, xxiv. (1841) 156.—Sibir.
arachnoidea, *Bunge, l. c.* 157.—Rossia; Reg. Caucas.
Auchericana, *DC. Prod.* vi. 674.—As. Min.
bellidioides, *Boiss. Fl. Orient.* iii. 583.—Reg. Caucas.
blanda, C. A. Mey. Vers. Pfl. Cauc. 67 = mollis.
Becconi, J. Gay, ex Nym. Consp. 416 = humilis.
Bocconi, *Guss. Fl. Sic. Syn.* ii. 448.—Sicil.; Hisp. mer.

JURINEA :—
bracteata, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 338.—Turkest.
brevicaulis, *Boiss. in Tchihat. Asie Min. Bot.* ii. 847. —Armen.
Bungei, *Boiss. Fl. Orient.* iii. 573.—Persia.
cadmea, *Boiss. Diagn.* Ser. I. iv. 93.—As. Min.
Capsil, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 393.—Turkestan.
cardiiformis, *Boiss. Diagn.* Ser. I. vi. 111.—Persia; Afghan.
catacnica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 580.—As. Min.
centaurcaps, *Benth. & Hook. f. Gen.* ii. 474.—Reg. Himal.
chaetocarpa, *Ledeb. ex DC. Prod.* vi. 676.—Sibir.
chaetolepis, Boiss. Diagn. Ser. II. iii. 50 = Tricholepis spartoides.
chia, Boiss. & Orph. ex Boiss. Fl. Orient. iii. 574 = anatolica.
consanguinea, DC. Prod. vi. 676 = anatolica.
cordata, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 581.—Persia.
cretacea, *Bunge, in Flora*, xxiv. (1841) 158.—Ross. austr.
cyanoides, *DC. Prod.* vi. 676.—Reg. Caucas.; Sibir.
depressa, *C. A. Mey. Verz. Pfl. Cauc.* 67.—As. Min. dumulosa, *Boiss. Fl. Orient.* iii. 569.—Persia.
dura, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 32 = anatolica.
elegans, *Stev. in DC. Prod.* vi. 675.—Reg. Caucas.
eriobasta, *DC. l. c.*—Persia.
erythrolepis, Boiss. Diagn. Ser. I. x. 98 = heterophylla.
Evermanni, *Bunge, in Flora*, xxiv. (1841) 155.— Rossia.
filicifolia, *Boiss. Fl. Orient.* iii. 569.—Reg. Caucas.
frigida, *Boiss. Diagn.* Ser. I. vi. 113.—Persia.
glycacantha, *DC. Prod.* vi. 677.—Graecia.
Gmelini, *C. A. Mey. ex Claus, Ind. Gorbel, Reise*, ii. 280.—Ross. med.
gnaphalioides, *Kotschy, in Sitzb. Akad. Muench.* (1878) 96.—Reg. Himal.
heterophylla, *Jaub. & Spach, Illustr.* ii. t. 181.— Persia.
heterophylla, Schur, Enum. Pl. Transs. 418 = mollis.
horrida, *Rupr. Sert. Tianschi.* 87.—Turkestan.
humilis, *DC. Prod.* vi. 677.—Reg. Mediterr.
integrifolia, Schur, Enum. Pl. Transs. 412 = mollis.
inuloides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 581.—Persia.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 322.—As. med.
lanipes, *Rupr. Sert. Tianschi.* 58.—Turkest.
lasiopoda, *Traute, in Act. Hort. Petrop.* viii. (1883) 510.—Turkest.
Ledebourii, Bunge, in Flora, xxiv. (1841) 157 = arachnoides.
leptoloba, *DC. Prod.* vi. 675.—Persia.
lineariloba, *DC. l. c.*—Reg. Caucas.; Sibir.
longifolia, DC. l. c. 674 = albicaulis.
lyrata, Trautv. Ind. Sem. Hort. Petrop. (1865) 87 = depressa.
macrocalathis, *C. Koch, in Linnaea*, xxiv. (1851) 415. —Banat.
macrocephala, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 378.—Reg. Himal.
macrocephala, *DC. Prod.* vi. 674.—Persia.
microcephala, *Boiss. Fl. Orient.* iii. 584.—Persia.
modesta, *Boiss. Diagn.* Ser. II. iii. 50.—Afghan.
mollis, *Reichb. Fl. Germ. Excurs.* 290.—Europ. austr. Monardi, *DC. Prod.* vi. 677.—Hispan.
mongolica, *Maxim. in Bull. Acad Pétersb.* xix. (1874) 519.—Mongol.
monoclona, Aschers. Fl. Brandenb. 852 = cynoides.
moschata, DC. Prod. vi. 677 = mollis.
multiloulis, *DC. l. c.* 674.—Persia.
Neumayriana, Vis. Ic. Fl. Dalm. t. 10. f. 2 = Amphoricarpos Neumayeri.
oblongifolia, Boiss. & Haussk. ex Boiss. Fl. Orient. iii. 578 = Ancherinia.
Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 51.—Turkest.
pinnata, *DC. Prod.* vi. 676.—Hispan.
pluricapa, C. Koch, in Linnaea, xxiv. (1851) 415 = arachnoides.
Pollichii, Nyman. Syll. 28, in syn. = cynoides.
polyclonos, *DC. Prod.* vi. 675.—Ross.; As. med.

JURINEA :—
proteoides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 579.—Persia.
pungens, *Boiss. Fl. Orient.* iii. 569.—Persia.
pyrenaica, Gren. & Godr. Fl Fr. ii. 270 = humilis.
radiata, *Boiss. Fl. Orient.* iii. 577.—Persia.
ramosissima, *DC. Prod.* vi. 677.—Persia; Afghan.
ramulosa, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 580.—Syria.
rhizantha, Fisch. & Mey. ex DC. Prod. vi. 677 = depressa.
robusta, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1844) 109.—Sconganria.
Rosgneri, C. Koch, in Linnaea, xxiv. (1851) 414 = arachnoides.
rosulata, *Klatt, in Sitzb. Akad. Muench.* (1878) 96.— Reg. Himal.
Ruprechti, *Boiss. Fl. Orient.* iii. 573.—Reg. Caucas.
salicifolia, *Boiss. l. c.* 580.—As. Min.
salicifolia, *Gruner, in Bull. Soc. Nat. Mosc.* xii. (1868) ii. 481.—Tauria.
sclerophylla, Boiss. & Heldr. ex eor. Diagn. Ser. I. x. 97 = Leuzea serratuloides.
sophiana, *Kalenicz. in Bull. Soc. Nat. Mosc.* xviii. (1845) i. 339.—Reg. Caucas.
sordida, Stev. l. c. xxix. (1856) ii. 401 = cynoides.
spectabilis, *Fisch. & Mey. in Ind. Sem. Hort. Petrop.* iv. 39.—Reg. Caucas.
squarrosa, *Fisch. & Mey. l. c.* xi. 73.—Reg. Caucas.
stoechadifolia, *DC. Prod.* vi. 674.—Ross. austr.
Stoechelinea, *Boiss. Diagn.* Ser. I. vi. 111.—Syria.
subacaulis, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 39 = depressa.
suffruticosa, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 166.—Orient.
tenuiloba, Bunge, in Flora, xxiv. (1841) 155 = cynoides.
tenuis, Bunge, l. c. 158 = Serratula tenuifolia.
thianschanica, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 322.—Turkestan.
tomentosa, Cass. in Dict. Sc. Nat. xxiv. 389 = cynoides.
transitoria, Schur, Enum. Pl. Transs. 413 = mollis. Trautvetteriana, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 52.—Turkestan.
Tricholepis, DC. Prod. vi. 678 (Tricholepidis sp.).— Reg. Himal.

JURINELLA, Jaub. & Spach, Illustr. ii. 101. t. 188–185 (1846) = Jurinea, Cass. (Compos.).
abrinthifolia, Jaub. & Spach, l. c. 108. t. 185 = Jurinea frigida.
Auchera, Jaub. & Spach, l. c. 101. t. 188 = Jurinea depressa.
Chamaecynara, Jaub. & Spach, l. c. 102. t. 184 = Jurinea depressa.

JUSSIA, Adans. Fam. ii. 85 (1763) = seq.

JUSSIAEA, Linn. Fl. Zeyl. 75 (1747) = seq.

JUSSIAEIA, Hill, Veg. Syst. xiii. 28 (1768) = **Jussieua**, Linn. (Onagrar.).

JUSSIEA, Linn. ex Sm. Laches. Lapp. i. 284 (1811) = **Potentilla**, Tourn. (Rosac.).

JUSSIENA, Reichb. Handb. 246 (1837) sphalm. = seq.

JUSSIEUA, Linn. Gen. ed. I. 196 (1737). *ONAGRARIEAE*, Benth. & Hook. f. i. 788.
CORYNOSTIGMA, Presl, Epim. Bot. 218 (1850).
CUBOSPERMUM, Lour. Fl. Cochinch. 275 (1790).
OLDENLANDIA, P. Br. Hist. Jamaic. 208 (1756).
acuminata, Sw. Fl. Ind. Occ. ii. 745 = linifolia.
adscendens, Linn. Mant. i. 69 = repens.
africana, DC. Prod. iii. 58.—Guinea.
alata, *G. Don, Gen. Syst.* ii. 693.—Bras.
alata, *Presl, Rel. Haenk.* ii. 84.—Mexic.
albiflora, Triana, ex Michell, in Mart. Fl. Bras. xiii. *11*. 156 = lithospermifolia.
alternifolia, E. Mey. ex Peters, Reise Mossamb. Bot. 69 = repens.
altissima, Perr. ex DC. Prod. iii. 55 = erecta.
ambigua, Miq. Stirp. Surin. Sel. 60 = decurrens.
amazonica, Spruce, ex Michell, in Mart. Fl. Bras. xiii. ii. 159 = inclinata.
amazonensis, *DC. Prod.* iii. 56.—Bras.
angustifolia, Blume, Bijdr. 1183 = Blumeana.

JUSSIEUA :—
angustifolia, Lam. Encyc. iii. 331 = suffruticosa.
Bertoriana, Steud. Nom. ed. II. i. 336 = repens.
Blumeana, *DC. Prod.* iii. 55.—Java.
bonariensis, *Michell, in Flora*, lvii. (1874) 303.—Am. trop.
Boydiana, Feathern. Rep. Louistan. Univ. (1870) 78 = repens.
brachycarpa, Lam. Encyc. iii. 331 = Ludwigia cylindrica.
brachycarpa, *Michell, in Flora*, lvii. (1874) 302.—Am. trop.
brachyphylla, *Michell, l. c.* 300.—Am. trop.
bracteata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 318 = elegans.
Burchellii, *Michell, in Flora*, lvii. (1874) 301.—Am. trop.
Burmanni, DC. Prod. iii. 57. suffruticosa.
calycina, Presl, Rel. Haenk. ii. 84 = octonervia.
Caparosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 258. —Bras.
caryophyllaea, Lam. Encyc. iii. 331 = Ludwigia parviflora.
Commersoni, Pohl, ex Michell, in Mart. Fl. Bras. xiii. ii. 155 = nervosa.
costata, Pohl, ex Michell, l. c. 151 = elegans.
costata, *Presl, Epim. Bot.* 217.—Ins. Philipp.
decumbens, Benth ex Wall. Cat. n. 6883 = suffruticosa.
decurrens, *DC. Prod.* iii. 56.—Am. bor.
densiflora, *Michell, in Flora*, lvii. (1874) 301.—Am. trop.
diffusa, Forsk. Fl. Aegypt.-Arab. 210 = repens.
distans, Pohl, ex Michell, in Mart. Fl. Bras. xiii. ii. 165 = affinis.
doclecandra, *DC. Prod.* iii. 58.—Guian.
edulis, Forsk. Fl. Aegypt. Arab. 210 = Corchorus Antichorus.
elegans, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 257.— Bras.
elevata, Pohl, ex Michell, in Mart. Fl. Bras. xiii. ii. 150 = Laruottenna.
elongata, Burch. ex Michell, l. c. 158 = Burchellii.
erecta, Blanco, Fl. Filip. ed. I. 366 = suffruticosa.
erecta, Linn. Amoen. Acad. viii. 256 = inclinata.
erecta, *Linn. Sp. Pl.* 388.—Am. trop.
erecta, Nutt. Gen. Am. I. 279 = decurrens.
erythracoulis, Mart. ex Michell, in Mart. Fl. Bras. xiii. ii. 170 = suffruticosa.
exaltata, Roxb. Hort. Beng. 83; Fl. Ind. ii. 401 = suffruticosa.
filiformis, *Michell, in Flora*, lvii. (1874) 300.—Am. trop.
floribunda, Griff. Notul. iv. 688 = repens.
fruticosa, *G. Don, Gen. Syst.* ii. 692.—Bras.
frutans, Hochst. in Flora, xxvii. (1844) 425 = repens.
furvistriis, Blume, Bijdr. 1183 = repens.
foliosa, *Wright, ex Griseb. Cat. Pl. Cub.* 107.—Cuba.
frutescens, *Vacq. f. Sem. Hort. Vind.* (1821); *ex DC. Prod.* iii. 56.—Hab.?
fruticosa, DC. l. c. 57 = suffruticosa.
grandiflora, Michx. Fl. Bor. Am. i. 267 = repens.
grandiflora, *Ruiz & Pav. Fl. Per.* iv. 382.—Peruv. Haenkeana, *Steud. Nom.* ed. II. i. 836.—Mexic.
helminthorhiza, Mart. in Flora, xxii. (1839) 1. Belbi. 61 = natans.
heterophylla, Salzm. ex Michell, in Mart. Fl. Bras. xiii. ii. 164 = pilosa.
hexamera, Miq. in Ann. Sc. Nat. Sér. III. i. (1844) 86 = affinis.
hexapetala, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 319 = repens.
hirsuta, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. centr.
hirsuta, Presl, Rel. Haenk. ii. 84 = Haenkeana.
hirsuta, Vell. Fl. Flum. 186; iv. t. 144 = mollis.
hirta, Lam. Encyc. iii. 331 = pubescens, scabra.
hirta, Vahl, Eclog. Am. ii. 51 = peruviana.
Hookeri, *Michell, in Flora*, lvii. (1874) 302.—Am. trop.
hyssopifolia, *G. Don, Gen. Syst.* ii. 693.—Afr. trop.
inclinata, Blanco, Fl. Filip. ed. I. 366 = repens.
inclinata, *Linn. f. Suppl.* 277.—Am. trop.
infusa, *Wright, in Journ. Linn. Soc.* x. (1869) 278.— Cuba.
Jungkuhniana, *Miq. Fl. Ind. Bat.* i. 1. 697.—Sumatra.
lanceolata, Cambess. in St. Hil. Fl. Bras. Mer. ii. 259 = myrtifolia.
lanceolata, Pohl, ex Michell, in Mart. Fl. Bras. xiii. ii. 156 = nervosa.
Laruotteana, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 256.—Bras.

JUSSIEUA:—
latifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 317.—Guian.
leptocarpa, Nutt. Gen. Am. i. 279 = pilosa.
ligustrifolia, H. B. & K. Nov. Gen. et Sp. vi. 100 = suffruticosa.
linearis, *Hooket. in Flora*, xxvii. (1844) 425.—Afr. austr.
linearis, *Willd. Sp. Pl.* ii. 875.—Afr. trop.
linifolia, *Vahl, Eclog. Am.* i. 32.—Am. et Afr. trop.
lithospermifolia, *Michell, in Flora*, lvii. (1874) 300.—Am. trop.
longifolia, *DC. in Mém. Soc. Phys. Genèv.* ii. (1824) 11. 141.—Bras.
longifolia, Reichb. Hort. Bot. i. 50. t. 67 = frutescens ?
longipes, Griff. Notul. iv. 689 = suffruticosa.
macrocarpa, H. B. & K. Nov. Gen. et Sp. vi. 102 = peruviana.
macropoda, *Presl, Rel. Haenk.* ii. 35.—Peruv.
madarapatana, Steud. Nom. ed. I. 447 = Ludwigia parviflora.
Maregravii, DC. Prod. iii. 58 = pilosa.
Martii, *Michell, in Flora*, lvii. (1874) 300.—Am. trop.
mauritiana, Presl, Bot. Bemerk. 70 = repens.
maypurensis, H. B. & K. Nov. Gen. et Sp. vi. 100 = nervosa.
meadiocrinervis, Gill. ex Hook. Bot. Misc. iii. (1833) 312 = repens.
micrantha, Kunze, Del. Sem. Hort. Lips. (1847); ex Linnæa, xxiv. (1851) 177 = linifolia.
microphylla, Mart. in Flora, xxii. (1839) i. Beibl. 61 = affinis.
Miquelii, Sagot, ex Michell, in Mart Fl. Bras. xiii. 11. 164 = pilosa.
mollis, H. B & K. Nov. Gen. et Sp. vi. 102 = peruviana.
montevidensis, Spreng. Syst. ii. 232 = repens.
multinervia, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 312.—Reg. Argent.
myrtifolia, *Cambess. in St Hil. Fl. Bras. Mer.* ii. 260. —Bras.
natans, *Humb. & Bonpl. Pl. Aequin.* i. 16. t. 8 B.—N. Granat.
nervosa, *Poir. Encyc. Suppl.* iii. 199.—Guian.
nodica, Hochst. ex Oliver, Fl. Trop. Afr. ii. 490 = linifolia.
oblonga, Gill. ex Hook. Bot. Misc. iii. (1833) 312 = repens.
occidentalis, Nutt. in Torr. & Gray, Fl. N. Am. i. 521 = octonervia.
octofila, DC. Prod. iii. 57 = suffruticosa.
octonervia, *Lam. Encyc.* iii. 332.—Am. trop.
octonalvis, Sw. Obs. Bot. 149; Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = suffruticosa.
Onagra, Mill. Card. Dict. ed. VIII. n. 4 = erecta.
oocarpa, Wright, ex Griseb. Cat. Pl. Cub. 108 = Oocarpon jussiæoides.
ovalifolia, *Sims, Bot. Mag.* t. 2580.—Madag.
palmicensis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 261 = nervosa.
palustris, G. F. W. Mey. Prim. Fl. Esseq. 173 = suffruticosa.
palustris, *Vell. Fl. Flum.* 186; iv. t. 143.—Bras.
paniculata, Poepp. ex Michell, in Mart. Fl. Bras. xiii. 11. 160 = decurrens.
parviflora, Cambess. in St. Hil. Fl. Bras. Mer. ii. 263 = suffruticosa.
parviflora, Salzm. ex Michell, in Mart. Fl. Bras. xiii. 11. 163 = linifolia.
patibilensis, H. B. & K. Nov. Gen. et Sp. vi. 97 = repens.
pauciflora, Pohl, ex Michell, in Mart. Fl. Bras. xiii. 11. 155 = nervosa.
pedunculosa, *Wright, ex Griseb. Cat. Pl. Cub.* 108.—Cuba.
peploides, H. B. & K. Nov. Gen. et Sp. vi. 97 = repens.
perotensiæfolia, Schlecht. in Linnæa, xii. (1838) 271 = suffruticosa.
peruviana, *Linn. Sp. Pl.* 888.—Am. trop.
pilosa, *H. B. & K. Nov. Gen. et Sp.* vi. 101.—Afr. trop.
pilosa, Pohl, ex Michell, in Mart. Fl. Bras. xiii. 11. 156 = brachyphylla.
Plumieriana, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 267.—Ins. Porto Rico.
polygonoides, Willd. ex Steud. Nom. ed. II. i. 586 = sq.
polygonoides, H. B. & K. Nov. Gen. et Sp. vi. 97 = repens.

JUSSIEUA:—
Potamogeton, *Michell, in Flora*, lvii. (1874) 301.—Am. trop.
Prieurea, *Guill. & Perr. Fl. Senegamb. Tent.* i. 294.—Afr. trop.
Pseudo-heterophylla, Salzm. ex Michell, in Mart. Fl. Bras. xiii. 11. 164 = pilosa.
ptrophora, Miq. in Ann. & Mag. Nat. Hist. Ser. I. xi. (1843) 19 = decurrens.
pulescens, *Linn. Sp. Pl.* ed. II. 555.—Am. calid.
pumila, Gill. ex Hook. Bot. Misc. iii. (1833) 312 = repens.
quadrangularis, *Michell, in Flora*, lvii. (1874) 302.—Am. trop.
racemosa, Rottl. ex DC. Prod. iii. 22 = Lumnitzera racemosa.
ramosa, *Jacq. f. ex Reichb. Ic. Bot. Exot.* 54. n. 75 (1827).—Bras.
ramulosa, DC. Prod. iii. 54 = repens.
repens, *Linn. Sp. Pl.* 888.—Amphig. trop.
rigida, Miq. Stirp. Surin. Sel. 58 = nervosa.
rigida, Poepp. ex Michell, in Mart. Fl. Bras. xiii. 11. 165 = affinis.
sagittata, Poir. Encyc. Suppl. iii. 198 = Ammannia latifolia.
Sagraeana, *A. Rich. Ess. Fl. Cub.* 534.—Cuba.
salicifolia, H. B. & K. Nov. Gen. et Sp. vi. 99 = suffruticosa.
scabra, Willd. Enum. Hort. Berol. i. 449 = suffruticosa
Schottii, *Michell, in Flora*, lvii. (1874) 302.—Bras.
sedoides, *Humb. & Bonpl. Pl. Aequin.* i. 13. t. 3 A. —Am. trop.
sericea, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 260.—Bras.
sessi(ar)flora, [Moç. & Sesse], ex DC. Prod. iii. 55 = erecta.
stenophylla, *Gill. in Hook. Bot. Misc.* iii. (1833) 312. —Arab. austr.
stolonifera, Guill. & Perr. Fl. Seneg. Tent. i. 292 = repens.
subacaulis, Pursh, Fl. Am. Sept. i. 304 = Oenothera heterantha.
suffruticosa, Griseb. ex Wright, in Journ. Linn. Soc. x. (1869) 279 = salicifolia.
suffruticosa, *Linn. Sp. Pl.* 888.—Amphig. trop.
surinamensis, *Miq. in Linnæa*, xviii. (1844) 870.—Guian.
Swartziana, DC. Prod. iii. 54 = repens.
tenella, *Burm. f. Fl. Ind.* 103. t. 35. f. 5; *Linn. Mant.* ii. 288.—Java.
tenuifolia, Nutt. in Ann. Journ. Sc. v. (1822) 294 = decurrens.
terminalis, Vell. Fl. Flum. 186; iv. t. 142 = anamomosana.
tetragona, Spreng. Syst. ii. 231 = suffruticosa.
tomentosa, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 254. —Bras.
torulosa, Am. in Ann. Sc. Nat. Sér. II. iii. (1835) 251, in texto = Oocarpon jussiæoides.
uruguayensis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 264.—Bras.
variabilis, G. F. W. Mey. Prim. Fl. Esseq. 174 = pilosa.
velutina, *G. Don, Gen. Syst.* ii. 695.—Afr. trop.
velutina, *Kunze, in Linnæa*, xx. (1847) 56.—Bras.
venosa, *Presl, Rel. Haenk.* ii. 35.—Mexic.
ventilifructus, Buch.-Ham. ex Wall. Cat. n. 6333 = villosa.
verticillata, Burch. ex Michell, in Mart. Fl. Bras. xiii. 11. 154 = inclinata.
villosa, Lam. Encyc. iii. 331 = suffruticosa.
villosissima, Pohl, ex Michell, in Mart. Fl. Bras. xiii. 11. 154 = sericea.
Weddelii, *M. Micl, in Flora*, lvii. (1874) 301.—Am. trop.

JUSSIEUAEA, [Rottl. ex] DC. Prod. iii. 22 (1828) = prsec.

JUSSIEUIA, Houst. Reliq. 8. t. 15 (1781) = Jatropha, Linn. (Euphorb.).
herbacea, Houst. l. c. = J. urens.

JUSSIEUIA, Thunb. Fl. Jap. 180 (1784) = Jussieua, Linn. (Onagrar.).

JUSSIEVA, Gleditsch, in Mem. Acad. Berol. v. (1751) 124 = prsec.

JUSSIEVIA, Linn. Syst. ed. II. 23 (1740) = prsec.

JUSTICA, Neck. Elem. i. 330 (1790) = seq.

JUSTICIA, Houst. ex Linn. Gen. ed. I. 4 (1737).
ACANTHACEÆ, Benth. & Hook. f. ii. 1108.
ADELODA, Rafin. Fl. Tellur. iv. 61 (1836).
ADNATODA, Nees, in DC. Prod. xi. 384 (1847), partim.
AMPHISCOPIA, Nees, in Wall. Pl. As. Rar. iii. 77, 112 (1832).
ANISOSTACHYA, Nees, in DC. Prod. xi. 366, 730 (1847).
Athlianthus, Endl. Gen. Suppl. ii. 63 (1843).
CAMPYLOSTEMON, E. Mey. in Drège, Zwei Pfl. Docum. 170 (1843), nomen.
DIPLANTHERA, Schrank, Pl. Rar. Hort. Monac. t. 62 (1819).
ECBOLIUM, Rtiv. ex Linn. Syst. ed. I (1735).
EMULARIA, Rafin. Fl. Tellur. iv. 63 (1836).
ETHESIA, Rafin. l. c. (1836).
GENDARUSSA, Nees, in Wall. Pl. As. Rar. iii. 76, 108 (1832).
GRUNSIA, Neck. Elem. i. 331 (1790).
HARNIERIA, Solms, in Sitzb. Ges. Naturf. Fr. Berl. (1864) 31.
HEMICHORISTE, Nees, in Wall. Pl. As. Rar. iii. 76, 103 (1832).
LEPTOSTACHYA, Nees, in DC. Prod. xi. 376 (1847).
MICROSPERMA, Rafin. Fl. Tellur. iv. 64 (1836).
MONECHMA, Hochst. in Flora, xxiv. (1841) 374.
PETALANTHERA, Rafin. Fl. Tellur. iii. 100 (1836).
RHAPHIDOSPERMA, G. Don, in Sweet, Hort. Brit. ed. III. 559 (1839).
RHAPHIDOSPORA, Nees, in Wall. Pl. As. Rar. iii. 115 (1832).
ROSLIMIA, Neck. Elem. i. 357 (1790).
ROSTELLARIA, Nees, in Wall. Pl. As. Rar. iii. 76, 100 (1832).
ROSTELLULARIA, Reichb. Handb. 190 (1837).
SAROTHECA, Nees, in Mart. Fl. Bras. ix. 113. t. 18 (1847).
SEMNERGIA, Neck. Elem. i. 366 (1790).
TYLOGLOSSA, Hochst. in Flora, xxv. (1842), i. Beibl. 144.
abnormis, Pohl, ex Nees, in DC. Prod. xi. 429, 482 = Dicliptera Pohliana.
acanthifolia, Willd. ex Nees, l. c. 302 = Aphelandra Acanthus.
acanthoides, Willd. l. c. = Barleria acanthoides.
acaulis, Linn. f. Suppl. 84 = Elytraria crenata.
acicularis, Sw. Prod. Veg. Ind. Occ. 13 = Anthacanthus jamaicensis.
acinoides, Hort. Kew. ex Nees, in DC. Prod. xi. 429, 498 = Peristrophe caulopsila.
Ackermanni, Mart. ex Nees, l. c. 429, 352 = Rhytiglossa cracaana.
acuminata, Vahl, Enum. i. 151 = Dicliptera acuminata.
acuminata, Wall. Cat. n. 2425 = Peristrophe acuminata, lers.
Adhatoda, Linn. Sp. Pl. 15 = Adhatoda Vasica.
Adhatoda, Mart. ex Nees, in DC. Prod. xi. 427 = Ecbolium Linnæanum.
adscendens, K. Br. Prod. 476 = procumbens.
ajugoides, Buch.-Ham. ex Nees, in DC. Prod. xi. 66 = Nelsonia tomentosa.
alata, Heyne, ex Steud. Nom. ed. II. i. 887 = diffusa.
alata, Vahl, Enum. i. 139 = Andrographis alata.
alba, Hort. Berol. ex Nees, in DC. Prod. xi. 731 = debilis.
alba, Roxb. Hort. Beng. 8; Fl. Ind. i. 116 = Eranthemum album.
albiflora, Blume, Bijdr. 787 = Peristrophe albiflora.
amabilis, Wall. Cat. Suppl. n. 71, 111 = Gymnostachyum venustum.
americana, Vahl, Symb. Bot. ii. 15 = Dianthera americana.
amœna, Mart. ex Nees, in DC. Prod. xi. 430 = Beloperone plumbaginifolia.
amygdalina, E. Mey. in Drège, Zwei Pfl. Docum. 196 = Adhatoda capensis.
Anagallis, Mart. ex Nees, in Mart. Fl. Bras. ix. 119 = Dianthera Anagallis.
anagalloides, *T. Anders. in Journ. Linn. Soc.* viii. (1864) 43.—Afr. austr.
andrographioides, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 584.—Burma.
androsæmifolia, Sieber, ex Nees, in DC. Prod. xi. 362 = Dianthera androsæmifolia.
angustifolia, Pohl, ex Nees, l. c. 339 = Dianthera angustifolia.

JUSTICIA :—
aciscanthus, *Schoenef. in Verh. Zool.-Bot. Ges. Wien*, xvii. (1868) 674.—Afr. trop.
annua, Stend. Nom. ed. 1. 374, 447 = Dicliptera scaurigularia.
Anselliana, T. Anders. in Journ. Linn. Soc. vii. (1864) 44 = Dianthera Anselliana.
antidota, *Sm. ex T. Anders. l. c.* 118 = J. Betonica.
antirrhina, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 89.—Bras.
appendiculata, Vahl, Enum. i. 159 = Beloperone appendiculata.
asperaa, Forsk. Fl. Aegypt. Arab. 6 = Barleria Prionitis.
arborea, Mill. Gard. Dict. ed. VIII. n. 7 = Aphelandra tetragona.
argyrostachya, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 510.—Burma.
aristata, Vahl, Symb. Bot. ii. 20 = Hypoëstes aristata.
aristata, Wall. Cat. n. 3481 = Monothecium aristatum.
armata, Sw. Prod. Veg. Ind. Occ. 13 = Asthacanthus armatus.
Arrebidea, *Steud. Nom.* ed. II. i. 867.—Bras.
arvensis, *Steud. l. c.*—Bras.
aspera, Perr. ex Nees, in DC. Prod. xi. 267 = Blepharis asperrima.
asperula, Lodd. Bot. Cab. t. 1681; Wall. Cat. n. 7171 = Phlogacanthus asperulus.
assamica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 597.—Reg. Himal.
assurgens, Linn. Syst. ed. X. 850 = Dicliptera assurgens.
Atherstonei, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 42.—Afr. austr.
Atkinsoni, *T. Anders. l. c.* ix. (1867) 509.—Reg. Himal.
atramentaria, Benth. Pl. Hartw. 69 = Jacobinia Mohlstili.
atropurpurea, *Dennst. Schluss. Hort. Malab.* 35.—Ind. or.
atrosanguinea, *Regel, Cat. Pl. Hort. Akakov.* 78.—Hab. ?
aurea, Schlecht. in Linnaea, vi. (1832) 398 = Jacobinia aurea.
auriculata, Wall. ex Nees, in DC. Prod. xi. 191 = Strobilanthes auriculatus.
australis, Gill. ex Nees, l. c. 395 = Adhatoda Gilliesii.
baphica, Spreng. Neue Entdeck. iii. 83 = Peristrophe tinctoria.
barbata, Bertol. in Nov. Comm. Acad. Bonon. iv. (1840) 405 (= *Rhytiglossa barbata*).—Guatemala.
barbata, Forsk. ex Nees, in DC. Prod. xi. 475 = Dicliptera Forsskalaea.
barleroides, Schott, ex Nees, in Mart. Fl. Bras. ix. 97 = Thyrsacanthus barleroides.
Barteri, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 39.—Afr. trop.
bengalensis, Spreng. Syst. i. 82 = Nelsonia campestris.
Betonica, T. Anders. in Journ. Linn. Soc. ix. (1867) 509 = trinervia.
Betonica, *Linn. Sp. Pl.* 15.—As. et Afr. trop.
betonicaefolia, Hort. Berol. ex Nees, in DC. Prod. xi. 480 = J. Betonica.
bibracteata, Wall. Cat. Suppl. n. 157 = Peristrophe tinctoria.
bicalyculata, Vahl, Symb. Bot. ii. 13 = Peristrophe bicalyculata.
bicolor, Sims, Bot. Mag. t. 1423 = Eranthemum bicolor.
bifora, Hort. ex Nees, in DC. Prod. xi. 430 = furcata.
bifora, Lam. Encyc. i. 629 = Asteotes trisulcus.
biflora, *Vahl, Symb. Bot.* ii. 9.—Ind. or.
bispinosa, Forsk. Fl. Aegypt. Arab. 6 = Barleria bispinosa.
bivalvis, Linn. Syst. ed. X. 850 = Dicliptera bivalvis.
bivalvis, Rovb. Hort. Beng. [80]; Fl. Ind. iii. 124 = Peristrophe commutata.
blepharophylla, Mart. ex Nees, in DC. Prod. xi. 459 = Chamaeranthemum Beyrichii.
blepharostegia, E. Mey. in Drège, Zwei Pfl. Docum. 195 = debilis.
Blumeana, *Klotzsch, ex Nees, in DC. Prod.* xi. 376.—Java.
boerhaaviaefolia, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 517.—Burma.
bonariensis, Gill. ex Nees, in DC. Prod. xi. 395 = Adhatoda Tweediana.
Bonneyana, *T. Muell. in Wing, South Sc. Record*, ii. (1882) 76.—Austral.
brachiata, Hook. ex Nees, in DC. Prod. xi. 475 = Dicliptera ciliaris.

JUSTICIA :—
brachiata, Pursh, Fl. Am. Sept. i. 18 = Dicliptera brachiata.
brachiata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Thyrsacanthus nitidus.
brachiata, Roth, ex Roem. & Schult. Syst. i. 167 = glauca.
brachiata, Salzm. ex Nees, in DC. Prod. xi. 478 = Dicliptera mucronifolia.
brachiata, Young, ex Nees, l. c. 346 = Gutesia lactevirens.
brachiata, Roem. & Schult. Syst. i. 167 = glauca.
brachionodes, Spreng. Syst. i. 82 = glauca.
brachystachya, *Thou. ex Schult. Mant.* i. 146.—Ins. Mauri.
bracteata, Pohl, ex Nees, in DC. Prod. xi. 367 = Herpetacanthus melancholicus.
bractelata, Jacq. Coll. iii. 258 = Thyrsacanthus bractelatus.
bractelata, Willd. Enum. Hort. Berol. 36 = Thyrsacanthus nitidus.
Brandisii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 514.—Burma.
brasiliana, Roth, Sp. Pl. Nov. 17 = Dianthera nodosa.
brasiliensis, Vell. ex Nees, in DC. Prod. xi. 430 = Amphilacopia Beyrichii.
brevicaulis, *S. Moore, in Journ. Bot.* xviii. (1880) 341. —Afr. trop.
brunelliaefolia, Lam. Tabl. Encyc. i. 40 = Nelsonia campestris.
Burchellii, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877-78) 80.—Bras.
burmanica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 597.—Burma.
caerulea, Auct. vide *cornulea*.
caerulea, Forsk. *Fl. Aegypt. Arab.* 5.—Arab.
calcarata, Hochst. in Flora, xxvi. (1843) 73 = neglecta.
calcarata, Wall. Pl. As. Rar. ii. 9. t. 113 = Rhinacanthus calcaratus.
callistachya, Cham. & Schlecht. in Linnaea, vi. (1831) 370 = Thyrsacanthus callistachyus.
callitricha, *Lodd. Bot. Cab.* t. 1921.—Bras.?
calonerva, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 98.—Burma.
calycotricha, Link & Otto, Ic. Pl. Sel. 113 t. 58 = Schaueria calycotricha.
calycotricha, Hook. Exot. Fl. t. 216 = Schaueria calycotricha.
campestris, *Griseb. in Goett. Abh.* xix. (1874) 235.—Reg. Argent.
campylostemon, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 44.—Afr. austr.
cancellata, Willd. ex Nees, in DC. Prod. xi. 506 = Hypoëstes cancellata.
canescens, Lam. Tabl. Encyc. i. 41 = Nelsonia campestris.
canescens, Wall. Cat. n. 3423 = Dicliptera Roxburghiana.
capensis, Eckl. ex Nees, in DC. Prod. xi. 498 = Peristrophe caulopelta.
capensis, E. Mey. ex Nees, l. c. 393 = Adhatoda incana.
capensis, *Thunb. Prod. Pl. Cap.* 104.—Afr. austr.
caracasana, Jacq. Coll. iv. 116 = Beloperone caracasana.
caracasana, Sieber, ex Nees, in DC. Prod. xi. 388 = Adhatoda Vasica.
caracasana, Willd. ex Nees, l. c. 340 = Dianthera secunda.
Caryana, Newm. ex Nees, l. c. 328 = Graptophyllum hortense.
Caryana, *Rovb. Hort. Beng.* 4, nomen.—Ind. or.
Caryana, Wall. Cat. n. 3456 = glabra.
carsipensis, H. B. & K. Nov. Gen. et Sp. ii. 233 = Dianthera secunda.
carnea, Hook. Bot. Mag. t. 3333 (= *Cyrtanthera magnifica*).—Bras.
carnea, Hook. ex Nees, in DC. Prod. xi. 320 (= *Packystachys asperula*).—Ins. Trist.
carthagrenensis, Benth. ex Nees, l. c. 409 = Adhatoda sphaerosperma.
carthagrenensis, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 56 (= *Adhatoda foliobrittana*).—Bras.
carthagrenensis, Willd. ex Nees, in DC. Prod. xi. 466 = Blechum Linnaei.
carthagrenensis, Jacq. Enum. Pl. Carib. 11 = Beloperone violacea.
carthaginensis, Vahl, Enum. i. 136 = Beloperone Lamarckiana.
caulopelta, E. Mey. in Drège, Zwei Pfl. Docum. 195 = Peristrophe caulopelta.

JUSTICIA :—
cavernarum, *F. Muell. Fragm.* vi. 91.—Austral.
Cephaladen, Herb. Berol. ex Link, Jahrb. i. 111. 48 = Lepidagathis hyalina.
ceylanica, *T. Anders. in Thw. Enum. Pl. Zeyl.* 233.—Zeylan.
Championi, *T. Anders. ex Benth. Fl. Hongk.* 264.—China.
charthagrinensis, Linn. Sp. Pl. ed. II. 1668, App. = Beloperone violacea ?
chellonida, *Rafin. Fl. Ludov.* 89 (Quid?).—Am. bor.
chinensis, Burm. f. Fl. Ind. 8. t. 4. f. 1 = Dicliptera Leonotis.
chinensis, Linn. Sp. Pl. 16 = Dicliptera chinensis.
chinensis, Mart. ex Nees, in DC. Prod. xi. 509 = Hypoëstes purpurea.
chinensis, Roxb. Hort. Beng. 4; Fl. Ind. i. 124 = Dicliptera Roxburghiana.
chinensis, Vahl, Symb. Bot. i. 4 = Dicliptera micrantha.
chinensis, Wall. Cat. n. 3466 = Dicliptera Leonotis, Roxburghiana, Hypoëstes trifora.
chlororptera, *Baker, in Journ. Linn. Soc.* xx. (1883 222.—Madag.
chrysotricha, *Benth. & Hook. f. Gen.* ii. 1109.—Am. trop.
chrysotrichoma, Pohl, ex Nees, in DC. Prod. xi. 408 = Adhatoda chrysotrichoma.
ciliaris, Linn. f. Suppl. 84 = Schwabea ciliaris.
ciliaris, Rusz. ex Nees, in DC. Prod. xi. 470 = Rungia parviflora.
ciliata, Jacq. Hort. Vindob. ii. 47 = Schwabea ciliaris.
ciliata, Pers. Syn. i. 23 = Beloperone ciliata.
cinerascens, Willd. ex Nees, in DC. Prod. xi. 382 = Jacobinia pubescens.
clandestina, Rusz. ex Wall. Cat. n. 2482 = Andrographis elongata.
clausula, Vahl, Enum. i. 146 = Dicliptera clavata.
cleomoides, *S. Moore, in Journ. Bot.* xviii. (1880) 313. —Afr. trop.
clinopodia, E. Mey. in Drège, Zwei Pfl. Docum. 195 = Dicliptera clinopodia.
coccinea, Aubl. Pl. Gui. i. 10 = Jacobinia coccinea.
coccinea, Cav. Ic. ii. 77 = Anisacanthus virgularis.
coccinea, Forsyth, ex Nees, in DC. Prod. xi. 341 = Dianthera secunda.
coccinea, Gosan, ex Nees, l. c. 398 = Adhatoda lithospermifolia.
coccinea, Hort. Berol. ex Nees, l. c. 431, 729 (= *Packystachys latior*).—Guian.
coccinea, Hort. Monsp. ex Nees, l. c. 415 = Adhatoda tetramera.
coccinea, Luschn. ex Nees, l. c. 420 = Beloperone plumbaginifolia.
coccinea, Willd. ex Nees, l. c. 420 (= *Packystachys asperula*).—Ind. occ.
cornuta, Poir. Encyc. Suppl. ii. 106 = Dicliptera pubescens.
cornuta, Steud. Nom. ed. II. i. 387 = Rungia parviflora.
collina, T. Anders. in Journ. Linn. Soc. ix. (1867) 515 = Dianthera collina.
colorata, Vahl, ex Nees, in DC. Prod. xi. 446 = Eranthemum nervosum.
comata, Lam. Encyc. iii. 632; Vahl, Symb. Bot. ii. 15 = Dianthera comata.
comata, Vell. ex Nees, in DC. Prod. xi. 432 = Adhatoda cuneata.
comosa, DC. ex Nees, l. c. 415 = Beloperone Rohrii.
comosa, *Vahl, Enum.* i. 155.—Am. austr.
confertiflora, Pohl, ex Nees, in DC. Prod. xi. 317 = Schaueria lachnostachys.
consanguinea, Nees, l. c. 473 = Strobilanthes exareolatus.
conspicua, Wall. Cat. n. 2460 = Barleria conspicua.
convena, E. Mey. in Drège, Zwei Pfl. Docum. 198 = Adhatoda protracta.
cordata, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 44.—Afr. trop.
cordifolia, Heyne, ex Wall. Cat. n. 2462 = Andrographis elongata.
corymbosa, Willd. ex Nees, in DC. Prod. xi. 500 = glabra.
corymbulosa, Bertol. in Nov. Comm. Acad. Bonon. iv. (1840) 404 (= *Rhytiglossa corymbulosa*).—Guatemala.
costata, Mart. ex Nees, in DC. Prod. xi. 57, 482 = Hygrophila longifolia.
cornuta, Pohl, ex Nees, l. c. 419 = Dianthera sodora.
cremulata, Pohl, ex Nees, l. c. 318 = Chaetothylax tomentosum.
crinita, DC. ex Nees, l. c. 444 = Sericospora crinita.

JUSTICIA :—
crinita, Thunb. Fl. Jap. 20 = Dicliptera Roxburghiana.
crispata, Pohl, ex Nees, in DC. Prod. xi. 356 = nummularia.
cristata, Jacq. Hort. Schoenb. iii. 38 = Aphelandra tetragona.
cristata, *Pohl, ex Nees, in DC. Prod.* xi. 356.—Bras.
cuneata, *Vahl, Symb. Bot.* ii. 10.—Afr. austr.
cuneata, Willd. ex Nees, in DC. Prod. xi. 392 = Adhatoda thymifolia.
cunefolia, Hook. ex Nees, l. c. 461 = Anthacanthus cuneatus.
cunefolia, *Nees & Mart. in Nov. Act. Nat. Cur.* i. (1823) 58.—Bras.
cuneiformis, E. Mey. ex Nees, in DC. Prod. xi. 398 = Adhatoda cuneata.
cuneiformis, Nees, in Mart. Fl. Bras. ix. 130 = cuneifolia.
curviflora, Wall. Pl. As. Rar. ii. 9. t. 119 = Phlogacanthus curviflorus.
cuspidata, Pers. Syn. i. 23 = Dicliptera peruviana.
cuspidata, Vahl, Symb. Bot. ii. 9 = Dicliptera micranthes.
cydoniifolia, *Griseb. Ind. Sem. Hort. Gotting.* (1878) 8.—Hab.!
cymosa, Herb. Madr. ex Nees, in DC. Prod. xi. 492 = glabra.
cynanchifolia, *R. Br. ex Nees, l. c.* 442.—Abyss.
Dahona, Buch.-Ham. ex Wall. Cat. sub n. 2442 = J. Gendarussa.
Dalzora, Blanco, Fl. Filip. ed. I. 14 = Peristrophe Dalzora.
dasycarpa, *Kurs, in Journ. As. Soc. Beng.* xlii. (1873) ii. 96.—Burma.
dasyclados, Mart. ex Nees, in DC. Prod. xi. 351 = Dianthera dasyclados.
debilis, *Lam. Encyc.* iii. 638 ; *Vahl, Symb. Bot.* ii. 15. —Arabia.
decussata, *Roxb. Hort. Beng.* 4 ; *Fl. Ind.* i. 127.—Burma.
dentata, Klein, ex Link, Jahrb. i. iii. 48 = Ecbolium Linneanum.
denticulata, Klein, ex Nees, in DC. Prod. xi. 428 = dentata.
depauperata, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 40.—Afr. trop.
dianthera, Vell. Fl. Flum. 13 ; i. t. 32 = Beloperone plumbaginifolia.
dichotoma, Blume, Bijdr. 788 = Dianthera dichotoma.
dichotoma, Herb. Madr. ex Wall. Cat. n. 2465 = glabra.
dichotoma, Roxb. in Gen. Nasurf. Fr. Neue Schr. iv. (1808) 221 = Rhinacanthus communis.
diffusa, Buch.-Ham. ex Wall. Cat. n. 2443. f. = J. Gendarussa.
diffusa, Sm. ex T. Anders. in Journ. Linn. Soc. vii. (1864) 117 = Lepidagathis hyalina.
diffusa, *Willd. Sp. Pl.* i. 87.—Geront. trop.
diversicta, Herb. Madr. ex Wall. Cat. n. 2492 = Daedalacanthus montanus.
divaricata, Roem. & Schult. Syst. i. 168 = capensis.
divaricata, Willd. ex Nees, in DC. Prod. xi. 391, 482 = Adhatoda divaricata.
divaricata, Zeyh. ex Nees, l. c. 386 = Dianthera ciliata.
Dohana, Buch.-Ham.—Cf. Wall. Cat. n. 2442. f. = J. Gendarussa.
dubia, *Forsk. Fl. Aegypt. Arab.* p. cii.—Arab. Fel.
dubia, Rafin. Atl. Journ. 146 (Quid !).—Am. bor.
Echolia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = Ecbolium Linneanum.
Echolium, Blanco, Fl. Filip. ed. I. 13 = Graptophyllum hortense.
Echolium, Linn. Sp. Pl. 15 = Ecbolium Linneanum.
Echolium, Wall. Cat. n. 2482 = dentata.
Echegaryi, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 61.—Reg. Argent.
echinoides, Crantz, Inst. i. 342 = sq.
echioides, Linn. Sp. Pl. 16 = Andrographis echioides.
Edgeworthii, T. Anders. in Journ. Linn. Soc. ix. (1867) 516 = pubigera.
elegans, Benav. Fl. Owar. 88 = Eranthemum elegans.
elegans, Pohl, ex Nees, in DC. Prod. xi. 382 = Sarothece elegans.
elegans, *Poir. Encyc. Suppl.* ii. 112.—Afr. trop.
elongata, Buch.-Ham. ex Wall. Cat. sub n. 2494 = Daedalacanthus strictus.
elongata, Vahl, Enum. i. 180 = Andrographis elongata.
emarginata, Nees, in DC. Prod. xi. 427 = Ecbolium Linneanum.
enaffroudy, Ell. Sketch, i. 11 = Dianthera americana.
enanthemoides, *F. Muell. Fragm.* vi. 90.—Austral.
eriandra, Salzm. ex Nees, in DC. Prod. xi. 387 = Gelasenoria nitida.
eustachiana, *Vacy. Enum. Pl. Carib.* 11.—Ind. occ.
eustachiana, Ker-Gawl. in Bot. Reg. t. 309 = Keriana.
eustachiana, Sims, Bot. Mag. t. 2076 = Chaetothylax eustachyana (in addend.).
eustachiana, Vahl, Symb. Bot. ii. 15 = Dicliptera Vahliana.
exserta, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 44.—Afr. trop.
falcata, Lam. Encyc. i. 629 = Dicliptera laevigata.
fasciata, *E. Mey. in Drège, Zwei Pfl. Docum.* 195.—Afr. austr.
fasciculata, Nees, in DC. Prod. xi. 457 = Eranthemum Hasenkeanum.
fasciculata, Russ. ex Wall. Cat. n. 2482 = Ecbolium Linneanum.
fastigiata, Linn. Illustr. i. 41 = Daedalacanthus montanus.
fastuosa, Linn. Mant. ii. 172 ; Vahl, Symb. Bot. i. 2 = Hypoëstes fastuosa.
filiformis, Ruiz, ex Nees, in DC. Prod. xi. 433, 211 = Ruellia pedunculosa.
fittonioides, *S. Moore, in Journ. Bot.* xvi. (1878) 184. —Afr. trop.
flaccida, *Kurs, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 80.—Burma.
flagellifera, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 535.—Burma.
flava, *Kurs, in Journ. As. Soc. Beng.* xlii. (1873) ii. 97.—Burma.
flava, Vahl, Symb. Bot. ii. 15 = Barleria flava.
flavicoma, Lindl. Bot. Reg. t. 1027 = Schaueria calycotricha.
flexibilis, Boj. ex Nees, in DC. Prod. xi. 505 = Hypoestes flexibilis.
fluminensis, *Steud. Nom.* ed. II. i. 838.—Bras.
foetida, Vell. Fl. Aegypt. Arab. 5 = Dicliptera bivalvis.
foliolosa, E. Mey. in Drège, Zwei Pfl. Docum. 195 = Adhatoda cuneata.
foliolosa, Willd. ex Nees, in DC. Prod. xi. 432 = Adhatoda cuneata.
foliosa, Jacq. ex Steud. Nom. ed. II. i. 838 = furcata.
formosa, Doan, Hort. Cantab. ed. III. 5 = Beloperone formosa.
formosa, Willd. Enum. Hort. Berol. i. 28 = Adhatoda tetramera.
Forskalei, Vahl, Symb. Bot. i. 2 ; ii. 13 = Hypoëstes Forskalei.
Forskohlii, R. Br. ex Steud. Nom. ed. II. i. 838 = praec.
Forsskohlii, Willd. ex Nees, in DC. Prod. xi. 509 = Hypoestes purpurea.
fragilis, *Dennst. Schlüssl. Hort. Malab.* 32.—Ind. or.
fragilis, *Wall. Cat.* n. 7174.—Burma.
frondosa, Vahl, Symb. Bot. ii. 8 = Dicliptera frondosa.
fruticosa, *Mill. Gard. Dict.* ed. VIII. n. 3.—Am. centr.
fulgida, Blume, Bijdr. 784 = Clinacanthus Burmanni.
fuliocoma, Cham. & Schlecht. in Linnaea, vi. (1831) 369 = Beloperone comosa.
furcata, *Jacq. Hort. Schoenb.* i. 2. t. 3.—Mexic.
furcata, Lam. Encyc. i. 631 = Ruellia paniculata.
Gandarussa, Linn. f. Suppl. 85 = J. Gendarussa.
gangetica, Linn. Cent. Pl. ii. 8 ; Amoen. Acad. iv. 299 = Asystasia coromandeliana.
Gendarusa, Blanco, Fl. Filip. t. 11 = Eranthemum bicolor.
Gendarussa, *Burm. f. Fl. Ind.* 10.—As. trop.
Gendarussa, Macrae, ex Nees, in DC. Prod. xi. 443 = Rhinacanthus communis.
geniculata, Sims, Bot. Mag. t. 2467 = Dianthera secunda.
geniculata, Vell. Fl. Flum. 13 ; i. t. 31 = Beloperone longipetiolata.
geniculata, Wall. Cat. n. 2443 = peploides.
genuflexa, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 87 (= *Rhytiglossa genuflexa*).—Bras.
Ghiesbreghtiana, Lem. in Fl. des Serres, Sér. I. iii. (1847) 243 verso = Jacobinia Ghiesbreghtiana.
Ghiesbreghtii, Hort. ex Lem. l. c. iv. (1848) 329 = praec.
Gilliesii, *Benth. & Hook. f. Gen.* ii. 1109.—Reg. Argent.
glabra, *Korn. ex Roxb. Hort. Beng.* 4 ; *Fl. Ind. ed. Carey*, i. 122.—Ind. or. ; Malaya.
glabrata, Rottl. ex Wall. Cat. n. 2455 C = glabra.
glandulifera, E. Mey. ex Nees, in DC. Prod. xi. 386 (= *Rhytiglossa origenioides*).—Afr. austr.
glandulosa, Pohl, ex Nees, l. c. 405 = Adhatoda glandulosa.
glandulosa, Roth, Nov. Pl. Sp. 13 = Andrographis Rothii.
glandulosus, Wall. Cat. n. 2485 = Andrographis glandulosa.
glauca, Heyne, ex Wall. l. c. sub n. 2494 = tranquebariensis.
glauca, *Rottl. in Ges. Naturf. Fr. Neue Schr.* iv. (1802) 219.—Ind. or.
glomerata, *Blume, Bijdr.* 786.—Java.
glomerata, Pohl, ex Nees, in DC. Prod. xi. 418 = Beloperone glomerata.
gonystachya, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 53 = Schaueria gonystachya.
gracilifrons, Wall. Cat. n. 2497 = Eranthemum gracilifrons.
gracilis, T. Anders. in Journ. Linn. Soc. ix. (1867) 512 = Rungia linifolia.
gracilis, Lyall, ex Nees, in DC. Prod. xi. 506 = Hypoëstes gracilis.
graminea, Koen. ex Roxb. Fl. Ind. ed. Carey, i. 134 = Rostellaria quinquenangularis.
graminea, Russ. ex Nees, in DC. Prod. xi. 376 = diffusa.
grandiflora, Dum.-Cours. Bot. Cult. ed. II. ii. 569 = Adhatoda tetramera.
grandifolia, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 510.—Burma.
Griffithii, T. Anders. l. c. 516 = Dianthera Griffithii.
grossa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 535.—Burma.
guttata, Wall. Pl. As. Rar. i. 24. t. 28 = Phlogacanthus guttatus, asperulus.
gymnostachya, Nees, in DC. Prod. xi. 426 = Ecbolium Linneanum.
harmatoder, Schlecht. in Linnaea, vii. (1832) 394 = Glockeria haematodes.
haplostachya, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 40 —Madag.
hedyotidifolia, Wall. Cat. n. 2450 = diffusa.
Helferi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 538.—Burma.
heterocarpa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 41.—Ind. occ. ; Abyss.
heterophylla, Schlecht. in Linnaea, v. (1830) 95 = Jacobinia heterophylla.
heterostegia, E. Mey. in Drège, Zwei Pfl. Docum. 195 = Dicliptera heterostegia.
hexangularis, Roem. & Schult. Syst. i. 168 in syn. = Dicliptera chinensis.
hirsuta, Hornem. ex Nees, in DC. Prod. xi. 433 = Strobilanthes hirta.
hirsuta, Jacq. Enum. Pl. Carib. 11. = Gonzalea spicata.
hirsuta, Pers. Syn. i. 23 = inficiens.
hirsuta, *Span. in Linnaea*, xv. (1841) 327.—Ins. Timor
hirsuta, Vahl, Symb. Bot. ii. 8 = Nelsonia campestris.
hirtella, Wall. Cat. n. 2448 = procumbens.
hispida, Heyne, ex Wall l. c. sub n. 2449 = simplex.
hispida, Willd. Sp. Pl. i. 84 = Eranthemum hispidum.
honamoriensis, Herb. Madr. ex Wall. Cat. n. 2491 A = Eranthemum malabaricum.
Hookeriana, *T. Anders. in Thw. Enum. Pl. Zeyl.* 233.—Zeylan.
humi/usa, Sw. Prod. Veg. Ind. Occ. 14 = Dianthera secunda.
humilis, Green, ex Nees, in DC. Prod. xi. 344 = Dianthera ovata.
humilis, Michx. Fl. Bor. Am. i. 8 = Dianthera ovata.
humuliflora, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 53 = Schaueria humuliflora.
hygrophiloides, *F. Muell. Fragm.* vi. 89.—Austral.
hypocraterjformis, Vahl, Enum. i. 165 = Eranthemum hypocrateriforme.
hysopifolia, Gouan, ex Nees, in DC. Prod. xi. 445 = Asiacantha virgata.
hysopifolia, *Linn. Sp. Pl.* 15.—Ins. Canar.
Hystrix, Lam. ex Nees, in DC. Prod. xi. 343 = Barleriola solanifolia.
imbricata, Pohl, ex Nees, l. c. 394 (= *Strobilorhachis prismatica*).—Bras.
imbricata, Vahl, Eclog. Am. i. 1 = Elytraria tridentata.
inaequalis, Benth. Pl. Hartw. 80 = Dianthera inaequalis
incana, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 42.—Afr. austr.
inconspicua, *Wall. Cat.* n. 2475.—Penang.
inficiens, *Vahl, Enum.* i. 161.—Peruv.
infracta, Vahl, l. c. 155 = Rungia parviflora.

JUSTICIA :—
infundibuliformis, Linn. Syst. ed. X. 850 = Crossandra undulaefolia.
lasiarla, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 40.—Afr. trop.
interrupta, E. Mey. in Drège, Zwei Pfl. Docum. 195 (= *Rhytiglossa ciliata*).—Afr. austr.
interrupta, Blanco, ex Nees, in DC. Prod. xi. 484 = Dicliptera interrupta.
interrupta, H. B. & K. Nov. Gen. et Sp. ii. 282 = Thyrsacanthus interruptus.
involucrata, Roxb. ex Koem. & Schult. Syst. ii. 141 = Hypoëstes involucrata, Peristrophe tinctoria.
jalapaefolia, Vahl, Enum. i. 180 = Peristrophe jalapaefolia.
japonica, Thunb. Fl. Jap. 80 = procumbens.
jasmnica, Lam. Encyc. ii. 638 = Peristrophe undulata.
juncea, R. Br. Prod. 476 = procumbens.
juncea, Koen. ex Roxb. Fl. Ind. ed Carey, i. 134 = quinqueangularis.
juncea, Rumph. ex Nees, in DC. Prod. xi. 476 = diffusa.
Kempeana, *F. Muell. Fragm.* xi. 101.—Austral.
Kerrana, *Sweet, Hort. Brit.* ed. 1. 398.—Ind. or.
khasiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 537.—Reg. Himal.
Kirkiana, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 39.—Afr. trop.
Klotzschiana, Hoffm. ex Nees, in DC. Prod. xi. 328 = Graptophyllum hortense.
Kursii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 537. —Burma.
ladanoides, DC. ex Nees, in DC. Prod. xi. 434 = Dicliptera Vahliana.
ladanoides, Lam. Tabl. Encyc. i. 43 = Adhatoda lithospermifolia.
laeta, Mart. ex Nees, in DC. Prod. xi. 350 = Dianthera laeta.
laeta, *S. Moore, in Journ. Bot.* xviii. (1880) 311.—Afr. trop.
laetevirens, Buckl. in Am. Journ. Sc. xlv. (1843) 176 = Gatesia laetevirens.
laetevirens, Vahl, Enum. i. 118 = Ecbolium Linneanum.
laevigata, Vahl, Symb. Bot. ii. 9 = Dicliptera javanica.
lamifolia, Koen. ex Roxb. Hort. Beng. 4; Fl. Ind. i. 134 = Nelsonia campestris.
lanosa, *Thunb. in Trans. Linn. Soc.* ii. (1794) 338.—Japon.
lanceata, Forsk. Fl. Aegypt. Arab. 6 = Barleria noctiflora.
lancolaria, Roxb. Hort. Beng. 4; Fl. Ind. i. 121 = Peristrophe lanceolaria.
lanceolata, Nees, in DC. Prod. xi. 240 = Barleria noctiflora.
lapathifolia, Vahl, Enum. i. 126 = Eranthemum lapathifolium.
latebrosa, Koen. ex Roxb. Hort. Beng. 80; Fl. Ind. i. 126 = Dicliptera parvibracteata.
latebrosa, Roxb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 551 = Dicliptera cuneata.
latebrosa, Russ. ex Wall. Cat. n. 2454 = Andrographis paniculata.
latevitia, Willd. ex Nees, in DC. Prod. xi. 420 = Beloperone plumbaginifolia.
latifolia, Vahl, Symb. Bot. ii. 4 = Eranthemum malabaricum.
lavandulaefolia, Pohl, ex Nees, in DC. Prod. xi. 348 (= *Rhytiglossa lavandulaefolia*).—Bras.
laxa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 43.—Afr. trop.
laxiflora, *Blume, ex Steud. Nom.* ed. II. i. 838 (Quid?). —Java.
Lazarus, *S. Moore, in Journ. Bot.* xvii. (1880) 313.—Afr. trop.
leptantha, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 44.—Malaog.
leptospermifolia, DC. ex Nees, in DC. Prod. xi. 435 = Dicliptera Vahliana.
leptostachya, Wall. ex Nees, l. c. = Gymnostachyum leptostachyum.
leucanthe, Blanch. ex Nees, l. c. 495, 452 = Eranthemum albiflorum.
leucantha, *Sebrank, in Syll. Ratisb.* i. (1824) 238.—Bras.
leucophlaea, Nees, in Mart. Fl. Bras. ix. 229 (= *Rhytiglossa leucophlaea*).—Bras.
ligularis, Salisb. ex Desf. Cat. Hort. Par. ed. III. 86 = quadrifida.
ligulata, DC. ex Nees, in DC. Prod. xi. 486 = Dicliptera resupinata.
ligulata, Lam. Encyc. i. 632 = Peristrophe bicalyculata.
ligustrina, Vahl, Enum. i. 118 = Ecbolium Linneanum.

JUSTICIA :—
Linaria, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 43.—Afr. trop.
linarifolia, Vahl, ex Nees, in DC. Prod. xi. 376 = diffusa.
Lindeni, T. Anders. in Journ. Agri-Hort. Soc. Ind. N. S. i. (1868) 280 = Beloperone violacea.
Lindeni, Houllet, in Rev. Hortic. (1870-71) 250 = Jacobinia Lindeni.
lineariifolia, Lam. Encyc. iii. 632; Tabl. Encyc. i. 41 = Dianthera americana.
linearis, Pers. Syn. i. 18 = lineolata.
lineata, Wight, ex Wall. Cat. n. 2486 = wynaadensis, Andrographis viscosula, A. lineata.
lineolata, *Ruiz & Pav. Fl. Per.* i. 9. t. 13. f. a.—Peruv.
linifolia, Heyne, in Wall. Cat. n. 2447 = Rungia linifolia.
lithospermifolia, Hort. Par. ex Nees, in DC. Prod. xi. 435 = Dicliptera resupinata.
lithospermifolia, Jacq. ex Nees, in DC. Prod. xi. 474 = Dicliptera resupinata.
lithospermifolia, Jacq. Hort. Schoenb. i. 3. t. 4 = Adhatoda lithospermifolia.
lithospermifolia, *Roth, Catalect. fasc.* iii. 1.—Hab.?
lithospermifolia, Vahl, Enum. i. 151 = Hygrophila guianensis.
livida, Nees, in DC. Prod. xi. 427 = Ecbolium Linneanum.
lobelioides, Wall. Cat n. 2464 = Andrographis lobelioides.
lobioides, *S. Moore, in Journ. Bot.* xviii. (1880) 310. —Afr. trop.
longepetiolata, Schott, ex Nees, in Mart. Fl. Bras. ix. 137 = Beloperone longepetiolata.
longifolia, Heyne, ex Wall. Cat. n. 2465 = Peristrophe montana.
longiflora, Vis. Sem. Hort. Patav. (1839).—Cf. Linnaea, xiv. (1840) Litt. 136 = Jacobinia longiflora.
longifolia, Balb. ex Nees, in DC. Prod. xi. 398 = Adhatoda reflexiflora.
longyfolia, Forst. f. Prod. 13 = Anthacanthus repandus.
longifolia, Heyne, ex Wall. Cat. n. 2467 = montana.
longifolia, Vahl, Symb. Bot. ii. 4 = Adhatoda longifolia.
longiracemosa, Hort. ex Lem. in Fl. des Serres, Sér. I. iii. (1847) 240 = Eranthemum coccineum.
longistaminea, Poepp. ex Nees, in DC. Prod. xi. 211 = Ruellia pedunculosa.
longistaminea, Ruiz & Pav. Fl. Per. i. 8 = Thyrsacanthus longistamineus.
longistylus, Gill. ex Nees, in DC. Prod. xi. 395 = Adhatoda Gilliesii.
Lourelriana, *Nees, l. c.* 428.—Cochinch.
lucens, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877-78) 80.—Bras.
lucida, Lam. Bot. Rep. t. 313 = Dianthera secunda.
lucida, Hort. Berol. ex Nees, in DC. Prod. xi. 435, 729 = Thyrsacanthus nitidus.
lupulina, Lam. ex Steud. Nom. ed. I. 111, in syn. = Ecbolium browei.
lupulina, E. Mey. in Drège, Zwei Pfl. Docum. 195 = Adhatoda lupulina.
lutea, Vahl, ex Nees, in Mart. Fl. Bras. ix. 104 = Schaueria sulfurea.
lutea, Ruiz & Pav. ex Schult. Mant. i. 140 (= *Pachystachys lutea*).—Bras.
lythroides, Fohl. ex Nees, in DC. Prod. xi. 314, 435 = Chaetothylax lythroides.
Macdonaldi, Hort. ex Regel, Cat. Pl. Hort. Akakov. 79 = Jacobinia aurea.
macilenta, E. Mey. in Drège, Zwei Pfl. Docum. 196 = Rhinacanthus communis.
macrantha, Benth. Fl. Hartw. 76 = Jacobinia macrantha.
macrophylla, Pohl, ex Nees, in Mart. Fl. Bras. ix. 105 = Schaueria macrophylla.
macrophylla, Spreng. Syst. i. 79 = Aphelandra squarrosa.
maculata, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 38.—Afr. trop.
maculata, *Ledd. Bot. Cab.* t. 696.—Hab.?
maculata, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 156.—Java.
maculosa, Boj. ex Nees, in DC. Prod. xi. 504 = Hypoëstes maculosa.
madagascarica, Burm. f. Fl. Ind. 9 = Calophanes littoralis.
magnifica, Pohl, ex Nees, in DC. Prod. xi. 339 (= *Cyrtanthera magnifica*).—Bras.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 534.—Ins. Penang.
major, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 39. —Abyss.
malabarica, [Soland. in] Ait. Hort. Kew. ed. I. i. 27 = Peristrophe bicalyculata.

JUSTICIA :—
Mannii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 44. —Afr. trop.
Marrubiastrum, Blanch. ex Nees, in Mart. Fl. Bras. ix. 22 = Hygrophila costata.
martinicensis, Jacq. Enum. Pl. Carib. 11 = Dicliptera martinicensis.
martinicensis, Sieber, ex Nees, in DC. Prod. xi. 466 = Blechum browei.
mauxia, Steud. Nom. ed. II. i. 888, sphalm. = *nauxia*.
matammensis, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 130.—Afr. trop.
mauritiana, Thou. ex Schult. Mant. i. 146 = glabra.
media, R. Br. Prod. 476 = procumbens.
megaphylla, Pohl, ex Nees, in Mart. Fl. Bras. ix. 14 = Clistax speciosus.
melgarinensis, Humb. & Boupl. ex Schult. Mant. i. 146 = Aphelandra melgarinensis.
membranacea, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 105 = polystachya.
mayuranensis, Klein, ex Nees, in DC. Prod. xi. 106 = Calophanes Nagchana.
micrantha, *Heyne, ex Wall. Cat.* n. 2449.—Ind. or.
micrantha, Pohl, ex Nees, in DC. Prod. xi. 408 = Adhatoda adenostachya.
micrantha, Wall. Cat. n. 2449, partim = procumbens.
microphylla, Lam. Illustr. i. 37 = Anthacanthus microphyllus.
microphylla, Pohl, ex Nees, in DC. Prod. xi. 424 = Beloperone involucrata.
minor, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 39. —Abyss.
mirabilioides, Lam. Tabl. Encyc. i. 39 = Beloperone Lamarckiana.
misera, Pohl, ex Nees, in DC. Prod. xi. 354 = Dianthera longiflora.
Mocti, Sesse, ex Nees, l. c. 486 = Jacobinia Mohintli.
Mohintli, Moç. & Sesse, ex Nees, l. c. 361 = Jacobinia Mohintli.
mollis, *E. Mey. in Drège, Zwei Pfl. Docum.* 196.—Afr. austr.
mollis, Pohl, ex Nees, in DC. Prod. xi. 486 = Dianthera rigida.
mollissima, Wall. Cat. n. 2445 = simplex.
monachinoides, *S. Moore, in Journ. Bot.* xviii. (1880) 311.—Afr. trop.
montana, Roxb. Corom. Pl. ii. 41. t. 176 = Daedalacanthus montanus.
montana, *Wall. Cat.* n. 2471.—Ind. or.
Moretiana, *Burm. f. Fl. Ind.* 10; *Vahl, Enum.* i. 162. —Zeylan.
mosaandes, *S. Moore, in Journ. Bot.* xviii. (1880) 313.—Afr. trop.
mucronata, Ruiz & Pav. Fl. Per. i. 8. t. 10. f. b = Aphelandra mucronata.
multiflora, Vahl, Enum. i. 160 = Dicliptera multiflora.
mycerensis, Klein, ex Schult. Mant. i. 146 = Calophanes Nagchana.
nauxia, Linn. Sp. Pl. 16 = Rhinacanthus communis.
natalensis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 38.—Afr. austr.
Necsiana, Mart. ex Nees, in DC. Prod. xi. 333 = Harpochilus Necsianus.
Necsiana, *Wall. Cat.* n. 7175.—Reg. Himal.
neglecta, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 40.—Afr. trop.
neilgherrensis, Auct. vide nilgherrensis.
nemorosa, Bert. ex Nees, in DC. Prod. xi. 468 = Tetramerium nervosum.
nemorosa, Sw. Prod. Veg. Ind. Occ. 14 = Beloperone nemorosa.
Nepeta, *S. Moore, in Journ. Bot.* xviii. (1880) 313.—Afr. trop.
nervosa, Vahl, Enum. i. 164 = Daedalacanthus nervosus.
nigriana, *Lour. Fl. Cochinch.* 24.—Cochinch.
nilgherrensis, Wall. Cat. n. 2480.—Ind. or.
nitida, Nees, in DC. Prod. xi. 328 = Graptophyllum hortense.
nitida, Jacq. Enum. Pl. Carib. 11 = Thyrsacanthus nitidus.
nitida, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 56 = Schaueria Martiniflori.
nodicaulis, Pohl, ex Nees, in DC. Prod. xi. 490 = Beloperone nodicaulis.
nodosa, Hook. Bot. Mag. t. 3914 = Dianthera nodosa.
nothe, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 537. —Ind. or.
nummularia, *Mart. ex Nees, in DC. Prod.* xi. 396.—Bras.

JUSTICIA :—

nummulariifolia, Vahl, Enum. i. 122 = Nelsonia campestris.

nummulariifolia, Wall. Cat. n. 2461 = Andrographis serpyllifolia.

nutans, Burm. f. Fl. Ind. 10. t. 5. f. 1 = Clinacanthus Burmanni.

oblongata, Link & Otto, Fl. Sel. 115. t. 54 = Beloperone plumbaginifolia.

ochroleuca, Blume, Bijdr. 786 = J. Betonica.

oxymoides, Lam. Encyc. i. 690 = Dicliptera micranthes.

odora, Lam. Encyc. i. 629 = Adhatoda odora.

odora, *Vahl, Symb. Bot.* ii. 11.—Arab.

odoratissima, Boj. ex Nees, in DC. Prod. xi. 443 = Rhinacanthus camosperma.

oleæfolia, Schlecht. ex Nees, l. c. 391 = Adhatoda capensis.

onygnatha, Pohl, ex Nees, l. c. 389 = Adhatoda cydoniæfolia.

opaca, Vahl, Enum. i. 123 = Barleria opaca.

orbiculata, T. Anders. in Journ. Linn. Soc. ix. (1867) 519, partim = simplex.

orbiculata, Wall. Cat. n. 2489 = diffusa, Eranthemum malaccense.

orchidea, E. Mey. ex Nees, in DC. Prod. xi. 394 = Adhatoda orchioides.

orchioides, Griff. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 544 = Sphinctacanthus Griffithii.

orchioides, *Linn. f. Suppl.* 85.—Afr. austr.

oreophila, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 556.—Ind. or.

origanoides, Griseb. Cat. Pl. Cub. 197 = Adhatoda origanoides.

origanoides, Vahl, Enum. i. 122 = Nelsonia tomentosa.

orixensis, Koen. ex Rozb. Hort. Beng. 80 ; Fl. Ind. i. 132 = glauca.

ovata, Dietz. ex Steud. Nom. ed. II. i. 338 = Dicliptera peruviana.

ovata, E. Mey. in Drège, Zwei Pfl. Docum. 196 (= *Rhytiglossa ovata*).—Afr. austr.

oxyphylla, DC. ex Nees, in DC. Prod. xi. 224 = Barleria micans.

pachyrrhiza, Mart. ex Nees, l. c. 456 = Eranthemum modestum.

pacifica, *Hemsl. Biol. Centr. Am. Bot.* ii. 516.—Mexic.

palæfera, Wall. Pl. As. Rar. i. 80. t. 92 = Eranthemum palæiferum.

palustris, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 88.—Abyss.

paniculata, Burm. f. Fl. Ind. 9 = Andrographis paniculata.

paniculata, Forsk. Fl. Aegypt. Arab. 4 = Hypoestes Forskålii.

parasitica, Lam. Tabl. Encyc. i. 49 = Agalmyla staminea.

parviflora, Ortog. Hort. Matr. Dec. 1 = Hemigraphis elegans !

parviflora, Retz, Obs. v. 9 = Rungia parviflora.

parvifolia, Hort. ex Nees, in DC. Prod. xi. 461 = Asthacanthus aciculatus.

parvifolia, Lam. Encyc. i. 627 = tranquebariensis.

patula, Lieht. ex Roem. & Schult. Syst. i. 164 = Adhatoda patula.

patula, E. Mey. ex Nees, in DC. Prod. xi. 392 = Adhatoda hyssopifolia.

pauciflora, *Griseb. in Gontt. Abh.* xxiv. (1879) 262.—Reg. Argent.

pauciflora, Vahl, in West, Bidr. Ste Croix, 267 ; Eclog. Am. i. 2 = sessilis.

pectinata, Linn. Cent. Pl. ii. 8 ; Amoen. Acad. iv. 299 = Rungia parviflora.

pectinata, Wight, ex Nees, in DC. Prod. xi. 366 = nilgherrensis.

pectoralis, Jacq. Enum. Pl. Carib. 11 = Diasthera pectoralis.

pedunculosa, Miehx. Fl. Bor. Am. i. 7 = Diasthera americana.

pentasticha, Buch.-Ham. ex Wall. Cat. n. 2370 = Nelsonia tomentosa.

pepleides, T. Anders. in Journ. Linn. Soc. ix. (1867) 511 = quinquenervularis.

pepleides, Boiss. Fl. Orient. iv. 825 = diffusa.

peripetaeflora, Jacq. Coll. Suppl. 5. t. 7. f. 2 = Adhatoda periploctifolia.

periplocifolia, Hort. Berol. ex Nees, in DC. Prod. xi. 398 = Adhatoda reflexiflora.

peruviana, Cav. Ic. i. 17. t. 26 = furcata.

peruviana, Lam. Encyc. iii. 685 = Dicliptera peruviana.

petiolaris, *E. Mey. in Drège, Zwei Pfl. Docum.* 196.—Afr. austr.

JUSTICIA :—

phænicea, Hort. Mus. Par. ex Dum. Cours. Bot. Cult. ed. II. ii. 566 = Beloperone plumbaginifolia.

picta, Linn. Sp. Pl. ed. II. 21 = Graptophyllum hortense.

picta, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 16 = Peristrophe salicifolia.

plicata, Vahl, Enum. i. 126 = Adhatoda plicata.

plumbaginea, Wall. Cat. n. 2487 = Asymasia coromandeliana.

plumbaginifolia, Jacq. Eclog. Am. 20. t. 2 = Beloperone plumbaginifolia.

polygonoides, H. B. & K. Nov. Gen. et Sp. ii. 282 (= *Rhytiglossa polygonoides*).—N. Granat.

polygonoides, Kozb. Hort. Beng. 3 ; Fl. Ind. i. 119 = Hygrophila polysperma.

polystachia, Vahl, Symb. Bot. ii. 7 = seq.

polystachya, *Lam. Tabl. Encyc.* i. 40.—Guian.

procumbens, T. Anders. ex Nees, in DC. Prod. xi. 358 = Dianthera pectoralis.

procumbens, Blume, Bijdr. 788 = Blumei.

procumbens, *Linn. Sp. Pl.* 15.—Ind. or.

procumbens, Linn. l. c. partim = micrantha, diffusa.

procumbens, Thib. ex Nees, in DC. Prod. xi. 363 = Lepidagathis alopecuroides.

procumbens, Wall. Cat. n. 2441 = quinquenervularis, simplex.

prolifera, E. Mey. ex Nees, in DC. Prod. xi. 337 (= *Rhytiglossa prolifera*).—*Afr. austr.*

prolixa, E. Mey. in Drège, Zwei Pfl. Docum. 196 (= *Rhytiglossa prolixa*).—Afr. austr.

prostrata, Schlecht. ex Nees, in DC. Prod. xi. 390 = Asthacanthus leptanthus.

protracta, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 41.—Afr. austr.

pruniflora, Pohl, ex Nees, in Mart. Fl. Bras. ix. 130 = nummularia.

Pseudo-betonica, Roth, Nov. Sp. Pl. 17 = J. Betonica.

psychotrioides, *Thou. ex Schult. Mant.* i. 146.—Ins. Mauric.

ptychostoma, Nees, in Wall. Pl. As. Rar. iii. 108 (sp. dub).—Ind. or.

ptychostoma, *Wall. Cat.* n. 2478.—Malaya.

pubescens, Lam. Tabl. Encyc. i. 40 = Gymnostachyum canescens.

pubescens, Ruiz, ex Nees, in DC. Prod. xi. 382 = Jacobinia pubescens.

pubescens, Scholt. Obs. Bot. 8 = furcata.

pubescens, Vahl, Symb. Bot. ii. 9 = Dicliptera pubescens.

pubigera, *Wall. Cat.* n. 2490.—Reg. Himal.

pulchella, Carey, ex Volgt, Hort. Suburb. Calc. 492 = Peristrophe pulchella.

pulchella, Rozb. Fl. Corom. ii. 41. t. 177 = Dasdalacanthus purpurascens.

pulchella, Wight, ex Wall. Cat. n. 2495 = Eranthemum montanum.

pulcherrima, *Jacq. Enum. Pl. Carib.* 11.—Ind. occ.

pulcherrima, Linn. f. Suppl. 84 = Aphelandra pulcherrima.

pulcra, Pohl, ex Nees, in DC. Prod. xi. 362 = Jacobinia pulcra.

pulegioides, E. Mey. in Drège, Zwei Pfl. Docum. 196 = seq.

Pulejum, E. Mey. ex Nees, in DC. Prod. xi. 390 = Adhatoda protracta.

pumila, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 197 = Asthacantha pumila.

punctata, Vahl, Symb. Bot. ii. 15 = Dianthera punctata.

punduana, *Wall. Cat.* n. 2458.—Reg. Himal.

purpurascens, Buch.-Ham. ex Wall. Pl. As. Rar. iii. 106 = Dasdalacanthus purpurascens.

purpurea, Forsyth, ex Nees, in DC. Prod. xi. 415 = Beloperone nemorosa.

purpurea, Linn. Sp. Pl. 16 = diffusa.

purpurea, Lour. Fl. Cochinch. 25 = Peristrophe tinctoria.

purpurea, Poit. ex Nees, in DC. Prod. xi. 490 = Dicliptera Vahliana.

purpurea, Vahl, Symb. Bot. ii. 13 = Hypoëstes purpurea.

pusilla, Pohl, ex Nees, in Mart. Fl. Bras. ix. 15 = Nelsonia campestris.

quadrangularis, Hook. Bot. Mag. t. 2845 = Phlogacanthus asperulus.

quadrifaria, T. Anders. in Journ. Linn. Soc. ix. (1867) 514. partim = Zollingeriana.

quadrifaria, *Wall. Cat.* n. 2479.—Reg. Himal. ; China.

quadrifida, Steud. Nom. ed. II. i. 338 = quadrifaria.

quadrifida, Vahl, Enum. i. 124 = Asthacantha virgulata.

JUSTICIA :—

quadrigona, Vahl, ex Link, Handb. i. 501 = Aphelandra cristata.

quinquenervularis, *Koen. ex Roxb. Hort. Beng.* 80 ; *Fl. Ind.* i. 138.—Ind. or.

quinquenervularis, Wall. Cat. n. 2443 = diffusa.

racemosa, *Ruiz & Pav. Fl. Per.* i. 9. t. 9. f. b.—Peruv.

racemulosa, Wikstr. in Vet. Acad. Handl. Stockh. (1825) 256 = Tetramerium racemulosum.

radicans, Vahl, Enum. i. 127 = Beloperone ciliata.

ramosa, Pohl, ex Nees, in DC. Prod. xi. 394 (= *Drejera ramosa*).—Bras.

ramosissima, Roxb. Hort. Beng 4 ; Fl. Ind. i. 129 = J. Betonica.

reflexiflora, Vahl, Enum. i. 127 = Adhatoda reflexiflora.

repanda, Forst. f. Prod. 8 = Asthacantha repandus.

repens, Deless. ex Nees, in DC. Prod. xi. 373 = Rouyeanna.

repens, Linn. Sp. Pl. 15 = Rungia repens.

repens, Nersud, ex T. Anders. in Journ. Linn. Soc. vii. (1864) 23 = Hypoëstes serpens.

repens, Pers. Syn. i. 23 = Dicliptera tomentosa.

repens, Vahl, ex Nees, in DC. Prod. xi. 470 = Rungia parviflora.

repens, Wall. Cat. n. 2459, B. partim = Rungia elegans.

reptans, Sw. Prod. Veg. Ind. Occ. 18 = Dianthera reptans.

resupinata, Hegewisch, ex Nees, in DC. Prod xi. 298 = Aphelandra Hasslariana.

resupinata, Vahl, Enum. i. 114 = Dicliptera resupinata.

reticulata, *Vahl, l. c.* 157.—Hab. ?

retorta, Vahl, l. c. 150 = Dicliptera parvibracteata.

retusa, *Vahl, l. c.* ii. 8.—Ins. S. Cruz.

rhodoptera, *Baker, in Journ. Linn. Soc. xx.* (1883) 221. —Madag.

rigida, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 87. —Ins. Socotr.

rivinæfolia, Vis. Sem. Hort. Patav. (1839).—Cf. Linnæa, xiv. (1840) Litt. 137 = Peristrophe speciosa.

rivinoides, Wall. Cat. n. 2468 ; et ex Nees, in DC. Prod. xi. 429 = Glossothigma spathulatum.

Rohrii, Vahl, Symb. Bot. ii. 6 = Beloperone Rohrii.

rosea, Vahl, Enum. i. 165 = Dasdalacanthus roseus.

rostrata, *Bertol. in Nov. Comm. Acad. Bonon.* iv. (1840) 406.—Guatemala.

Rottleriana, Wall. Cat. n. 2477 = Rhinacanthus communis.

rotundifolia, *E. Mey. in Drège, Zwei Pfl. Docum.* 196. —Afr. austr.

rotundifolia, Nees, in Wall. Pl. As. Rar. iii. 108 = Echolium Linneanum.

rotundifolia, Wall. Cat. n. 2489 = Dicliptera Roxburghiana, Rostellaria rotundifolia.

Roxburghiana, Blume, Bijdr. 786 = Peristrophe Blumeana.

Roxburghiana, Hort. Berol. ex Nees, in DC. Prod. xi. 723 = Nomaphila corymbosa.

Roxburghiana, Schult. Mant. i. 140 = Peristrophe tinctoria.

Rouyenna, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 587.—Zeylan.

rubicunda, Buch.-Ham. in Wall. Cat. n. 2480 = Eranthemum palæiferum.

rubra, Vahl, Eclog. Am. iii. 1 = Thymacanthus ruber.

rubra, Wight, ex Wall. Cat. n. 2468 = Peristrophe montana.

ruderalis, Rumx. ex Steud. Nom. ed. II. i. 339 = Peristrophe bicalyculata.

rupestris, Hochst. ex Nees, in DC. Prod. xi. 439 = debilis.

Ryani, Vahl, Eclog. Am. i. 1 = Beloperone cristachyana.

salicaria, Pohl, ex Nees, in DC. Prod. xi. 360 = Dianthera lasta.

salicifolia, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 514.—Reg. Himal.

salicifolia, *Blume, ex Steud. Nom.* ed. II. i. 339.—Java.

salicina, Vahl, Enum. i. 132 ; Jacq. ex Nees, in DC. Prod. xi. 439 = Jacobinia serices.

salicina, Vahl, ex Nees, in DC. Prod. xi. 410 = J. Gendarussa.

Salsola, *S. Moore, in Journ. Bot.* xviii. (1880) 340.—Afr. trop.

salsoloides, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 514.—Ind. or.

salviæfolia, *H. B. & K. Nov. Gen. et Sp.* ii. 288.—Mexic.

salviæfolia, Kunth, Syn. Pl. Aequin. ii. 81 = salviæfolia.

sanguinea, Humb. & Bonpl. ex Schult. Mant. i. 146 = Habracanthus sanguineus.

sanguinolentata, Vahl, Symb. Bot. ii. 10 = Gymnostachyum sanguinolentatum.

JUSTICIA :—
aculeva, Vahl, ex Nees, in DC. Prod. xi. 396 = Aphelandra pectinata.
scabrida, *J. Moore, in Journ. Bot.* xvlii. (1880) 310.—Afr. trop.
scandens, Michx. ex Nees, in DC. Prod. xi. 490 = Dicliptera Vahliana.
scandens, Vahl, Symb. Bot. ii. 7 = Rhinacanthus communis.
Schimperiana, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 86.—Abyss.
Schottii, Pohl, ex Nees, in Mart. Fl. Bras. ix. 107 = Schaueria Schottii.
scorpioides, *Griseb. in Goett. Abh.* xxiv. (1878) 262.—Reg. Argent.
scorpioides, Linn. Sp. Pl. ed. II. 21 = Tetranerium scorpioides.
secunda, Sieber, ex Steud. Nom. ed. II. i. 889 = Gonsales Coconcypselum.
secunda, Vahl, Symb. Bot. ii. 7 = Dianthera secunda.
secundiflora, Poepp. ex Nees, in DC. Prod. xi. 378 (= *Leptostachya Poeppigiana*).—Peruv.
secundiflora, Ruiz & Pav. ex Nees, l. c. 440 (= *Leptostachya secundiflora*).—Peruv.
secundiflora, *Vahl, Enum.* i. 158.—Peruv.
sericea, Ruiz & Pav. Fl. Per. I. 9. t. 9. f. b = Jacobinia sericea.
serpens, Vahl, Symb. Bot. ii. 9 = Hypoëstes serpens.
serpyllifolia, Rottl. ex Vahl, Enum. i. 169 = Andrographis serpyllifolia.
serrata Humb. ex Nees, in DC. Prod. xi. 394 = Barleria micans.
serrata, *Steud. Nom.* ed. II. i. 889 ; *et ex Nees, l. c.* 440. —Brs.
serrulata, Heyne, ex Wall. Cat. n. 2452 = Gymnostachyum cauescens.
sessilifolia, Herb. Madr. ex Wall. l. c. n. 2408 = Strobilanthes consanguineus.
sessiliflora, *Humb. & Bonpl. ex Schult. Mant.* i. 146. —Am. austr.
sessilis, Jacq. Enum. Pl. Carib. 11 = Dianthera sessilis.
setiloracta, Pohl, ex Nees, in Mart. Fl. Bras. ix. 129 (= *Rhytiglossa inaequilabia*).—Bras.
sexangularis, Cav. Ic. iii. 2. t. 208 = Dicliptera chinensis, resupinata.
sexangularis, Forsk. Fl. Aegypt. Arab. 5 = Dicliptera chinensis.
sexangularis, Gmelinest. ex Nees, in DC. Prod. xi. 490 = Dicliptera Vahliana.
sexangularis, Hort. Berol. ex Nees, in DC. Prod. xi. 440 = Dicliptera sexangula.
sexangularis, Linn. Sp. Pl. 16 = Dicliptera sexangularis.
sexangulata, Nees, in DC. Prod. xi. 474 = Dicliptera resupinata.
sibastica, Nees, l. c. 448 = Rhinacanthus communis.
sibaserrir, Nees, l. c. 404 = Peristrophe montana.
simplex, *D. Don, Prod. Fl. Nep.* 118.—As. et Afr. trop.
simensis, J. F. Gmel. Syst. 34 = Dicliptera Leonotis.
sinuata, Vahl, Symb. Bot. ii. 11 = Anthacanthus sinuatus.
somalensis, *Franch. in Révoil, Çomali*, 53.—Afr. trop.
spaciflora, Pohl, ex Nees, in DC. Prod. xi. 488 = Dicliptera spaciflora.
spartioides, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 48.—Afr. trop.
spathulata, Klotzsch, ex Nees, in DC. Prod. xi. 408 = Beloperone violacea.
spathulata, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 95 = Beloperone spathulata.
spathulata, *Thunb. Fl. Jap.* 23.—Japon.
speciosa, Roxb. Hort. Beng. 4; Fl. Ind. i. 122 = Peristrophe speciosa.
spergulaefolia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 48.—Afr. austr.
sphaerosperma, *Vahl, Symb. Bot.* ii. 8.—Ind. occ.
spicata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 18 (Rauthemi sp. ?).—Java.
spicata, *Ruiz & Pav. Fl. Per.* i. 8. t. 9. f. a.—Peruv.
spicata, Vell. ex Nees, in DC. Prod. xi. 440 (= *Pseuderanthemum Riedelianum*).—Bras.
spicigera, Schlecht. in Linnaea, vii. (1832) 895 = Jacobinia Meblenii.
spinosa, Jacq. Enum. Pl. Carib. 11; Linn. Sp. Pl. ed. II. 1668 = Anthacanthus spinosus.
spinulosa, Hochst. ex Nees, in DC. Prod. xi. 440 = Dicliptera microcalyx.

JUSTICIA :—
squamosa, Soland. ex Roem. & Schult. Syst. i. 139 = Rhytiglossa virgata.
squamosa, Thou. ex Schult. Mant. i. 146 = Isoella.
squarrosa, *Griseb. in Goett. Abh.* xix. (1874) 236.—Reg. Argent.
staminea, Vahl, Enum. i. 148 = Thyrsacanthus longistaminea.
staminea, *Humb. & Bonpl. ex Schult. Mant.* i. 146, non Am. austr.
stricta, Lam. ex Steud. Nom. ed. II. i. 889 = Andrographis paniculata.
stricta, Pohl, ex Nees, in DC. Prod. xi. 440, 360 = Dianthera rigida.
stricta, Vahl, Symb. Bot. ii. 6 = Nomaphila corymbosa.
strobilifera, *Linn. Tabl. Encyc.* i. 36.—Madag.
suaveolens, Willd. ex Nees, in DC. Prod. xi. 440, 358 (= *Rhytiglossa polygonoides*).—Ind. or.
subcaudata, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 190.—Afr. trop.
subfruticosa, E. Mey. in Drège, Zwei Pfl. Docum. 196 = Adhatoda tubulosa.
sulcata, Vahl, Symb. Bot. ii. 18 = Adhatoda sulcata.
sulfurea, Pohl, ex Nees, in Mart. Fl. Bras. ix. 104 = Schaueria sulfurea.
sulphurea, Pohl, ex Nees, in DC. Prod. xi. 318 = praec. sunatrana, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 142.—Ins. Nicobar.
superba, Hort. ex Nees, in DC. Prod. xi. 445 = Aniscanthus virgularis.
sylvatica, Vahl, Enum. i. 145 = Rhinacanthus communis.
sylvestris, Vahl, Cat. n. 2468 = Peristrophe montana.
syringifolia, *Vahl, Enum.* i. 117.—Madag.
tenuiseta, Hort. Liverp. ex Nees, in DC. Prod. xi. 419 = Dianthera nodosa.
tenuiseta, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 40. —Afr. trop.
tenuiflora, *Ruiz & Pav. Fl. Per.* i. 9. t. 12. f. a.—Peruv.
tenuiflora, Wall. Fl. As. Rar. iii. 116 = Andrographis tenuis.
tenuifolia, Spreng. Syst. i. 76 = tenuiflora.
tenuispica, *Steud. Nom.* ed. II. i. 889.—Bras.
tetragona, Hook. ex Nees, in DC. Prod. xi. 454 = Eranthemum tetragonum.
tetragona, Vahl, Symb. Bot. iii. 5 = Aphelandra tetragona.
thyrsiflora, Roxb. Hort. Beng. 3; Fl. Ind. i. 114 = Phlogacanthus thyrsiflorus.
thyrsoidea, Pohl, ex Nees, in DC. Prod. xi. 416 = Beloperone monticola.
thyrsoides, Roxb. ex Nees, l. c. 321 = Phlogacanthus thyrsiflorus.
tinctoria, Hoffm. ex Nees, l. c. 379 = Dianthera dichotoma.
tinctoria, Hort. Calc. ex Nees, l. c. 495 = Peristrophe speciosa.
tinctoria, *Lour. Fl. Cochinch.* 95.—Cochinch.
tinctoria, Roxb. Hort. Beng. 4; Fl. Ind. i. 128 = Peristrophe tinctoria.
tomentosa, Roxb. ll. ca. 4; 131 = Nelsonia campestris.
tomentosa, Vahl, Enum. i. 160 = Dicliptera tomentosa.
tranquebariensis, *Linn. f. Suppl.* 85.—Ind. or.
tranquebariensis, Roxb. Hort. Beng. 4; Fl. Ind. i. 133 = glauca.
triangularis, Heyne, ex Nees, in DC. Prod. xi. 399 = glauca.
tribracteata, Spreng. ex Nees, l. c. 781 = Beloperone leucopsicteiaris.
trichantha, Pohl, ex Nees, l. c. 424 = Beloperone involucrata.
trichocalyx, *Baker, in Journ. Linn. Soc. xxi.* (1885) 426.—Madag.
tridentata, E. Mey. in Drège, Zwei Pfl. Docum. 196 (= *Rostellularia tridentata*).—Afr. austr.
trifasciata, E. Mey. ex Nees, in DC. Prod. xi. 441 = Adhatoda fasciata.
trifloral, Forsk. Fl. Aegypt. Arab. 4. n. 10 = Hypoëstes triflora.
trifoliata, Roem. & Schult. Syst. i. 166 = Beloperone trifoliata.
trifoliolata, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 40.—Madag.
trinervia, *Vahl, Enum.* i. 156.—Ind. or.
trigonosa, Forsk. Fl. Aegypt. Arab. 6 = Barleria trispinosa.

FINIS TOMI PRIMI.

JUSTICIA :—
tristis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 88. —Afr. trop.
trivalva, Vahl, Symb. Bot. ii. 10 = Asistoles trivalvis.
trifiosa, *Baker, in Journ. Linn. Soc. xxi.* (1886) 429. —Madag.
tuberosa, Bertol. in Nov. Comm. Acad. Bonon. iv. (1840) 406 = Thyrsacanthus tuberosus.
tuberosa, Buch.-Ham. ex Wall. Cat. n. 2428 = Phlogacanthus tuberosus.
tuberosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 196.— Afr. austr.
tunicata, *Afz. Remed. Guin.* iii. 17.—Afr. trop.
Tweediana, *Benth. & Hook. f. Gen.* ii. 1109.—Bras.
umbellata, Hort. Madr. ex Nees, in DC. Prod. xi. 500 = glabra.
umbellata, Vahl, Enum. i. 111 = Dicliptera micranthes.
umbellata, Wall. Cat. n. 2467 = Dicliptera cuneata.
umbellata, Wight, ex Nees, in DC. Prod. xi. 488 = Dicliptera parvibractesta.
umbrosa, Benth. Pl. Hartw. 79 = Jacobinia aurea.
uncinulata, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 180.—Afr. trop.
undulata, Vahl, Symb. Bot. ii. 8 = Peristrophe undulata.
Vahliana, Schult. Mant. i. 138 = diffusa.
Vahlii, Dietr. ex Nees, in DC. Prod. xi. 441 = comosa.
Vahlii, Roth, Nov. Pl. Sp. 14 = diffusa.
Vandelli, *Roem. & Schult. Syst.* i. 166 — Bras.
variabilis, Zuccagni, in Roem. Collect. 116 = furcata.
variegata, *Acalt. Pl. Cat.* i. 13. t. 4.—Guinea.
vasculosa, *Wall. Cat.* n. 2469.—Reg. Himal ; Burma.
Velasqueziі, *Bertol. in Nov. Com. Acad. Bonon.* iv. (1840) 406.—Guatemala.
Vellozoi, Roem. & Schult. Syst. i. 166 = Beloperone hirsuta.
venosa, Pohl, ex Nees, in DC. Prod. xi. 358 = cristata.
venosa, Wight, ex Nees, l. c. 104 = Strobilanthes consanguineus.
ventricosa, *Wall. Cat.* n. 2426.—Burma ; China.
verbenaefolia, Wall. Pl. As. Rar. i. 58. t. 66 = Gymnostachyum venustum.
verticillaris, Linn. f. Suppl. 85 = Hypoëstes verticillaris.
verticillaris, Steud. Nom. ed. I. 450; ed. II. i. 840 = Hypoëstes aristata.
verticillata, Roxb. Hort. Beng. 80; Fl. Ind. i. 135 = Haplanthus verticillaris.
verticillata, Steud. Nom. ed. I. 450; ed. II. i. 840 = Dicliptera micranthes.
vestita, Schult. Mant. i. 145 = Nelsonia tomentosa.
villosa, Mart. ex Nees, in DC. Prod. xi. 390 = Ruellia villosa.
villosa, Pohl, ex Nees, l. c. 417 = Beloperone hirsuta.
vincoides, Lam. Encyc. i. 698 = Anthacanthus vincoides.
violacea, Neroniha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 18 = Graptophyllum hortense.
violacea, *Vahl, Symb. Bot.* ii. 15.—Arab.
virgata, T. Anders. in Journ. Linn. Soc. ix. (1867) 516 — Dianthera virgata.
virgularia, Salisb. Parad. Lond. t. 50 = Aniscanthus virgularis.
viridiflora, Buckl. ex Nees, in DC. Prod. xi. 346 = Gutzeia laetevirens.
viridis, Blanco, Fl. Filip. ed. I. 15 = Dicliptera viridis.
viridis, Forsk. Fl. Aegypt. Arab. 5 = Eebolium Linnaeanum.
viridis, Gill. ex Nees, in DC. Prod. xi. 395 = Adhatoda Tweediana.
vitellina, Roxb. Hort. Beng. 8; Fl. Ind. i. 115 = Phlogacanthus superbus.
volubilis, *Vand. ex Roem. & Schult. Syst.* i. 166.— Bras.
vulgaris, Bert. ex Schult. Mant. i. 135 = Adhatoda ophiorrhizoides.
Wallichii, T. Anders. in Journ. Linn. Soc. ix. (1867) 516 = Dianthera Wallichii.
Wenzingii, *Miers, in Ejusd. Viderunt. Addend.* (1877-78) 60.—Bras.
Wightiana, Wall. Cat. n. 2478 = Rungia Wightiana.
Wrightii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 320.—Am. bor. occ.
wynaadensis, *Heyne, in Wall. Cat.* n. 2474.—Ind. or.
xylocostoides, *Griseb. in Goett. Abh.* xix. (1874) 235.— Reg. Argent.
Zollingeriana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 530.—Burma ; Java.

DATE OF ISSUE

This book must be returned within 3,7,14 days of its issue. A fine of ONE ANNA per day will be charged if the book is overdue.

For Reference

NOT TO BE TAKEN FROM THIS ROOM